



(51) International Patent Classification:

A61K 8/37 (2006.01) *A61Q 5/10* (2006.01)
A61K 8/41 (2006.01) *A61K 8/19* (2006.01)

(21) International Application Number:

PCT/EP2013/060590

(22) International Filing Date:

23 May 2013 (23.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12179340.0 6 August 2012 (06.08.2012) EP

(71) **Applicant** (for AE, AG, AU, BB, BH, BN, BW, BZ, CA, CY, EG, GB, GD, GH, GM, IE, IL, IN, KE, KN, LC, LK, LS, MT, MW, MY, NA, NG, NZ, OM, PG, QA, RW, SC, SD, SG, SL, SZ, TT, TZ, UG, VC, ZA, ZM, ZW only): **UNILEVER PLC** [GB/GB]; a company registered in England and Wales under company no. 41424 of Unilever House, 100 Victoria Embankment, London Greater London EC4Y 0DY (GB).

(71) **Applicant** (for all designated States except AE, AG, AU, BB, BH, BN, BW, BZ, CA, CY, EG, GB, GD, GH, GM, IE, IL, IN, KE, KN, LC, LK, LS, MT, MW, MY, NA, NG, NZ, OM, PG, QA, RW, SC, SD, SG, SL, SZ, TT, TZ, UG, US, VC, ZA, ZM, ZW): **UNILEVER N.V.** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

(71) **Applicant** (for US only): **CONOPCO, INC., D/B/A UNILEVER** [US/US]; 800 Sylvan Avenue, AG West, S. Wing, Englewood Cliffs, New Jersey 07632 (US).

(72) **Inventors**: **INGER, Nicholas, John**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral Merseyside CH63 3JW (GB). **BATCHELOR, Stephen, Norman**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral Merseyside CH63 3JW (GB).

ton, Wirral Merseyside CH63 3JW (GB). **PELLETIER, Anne-Sophie**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral Merseyside CH63 3JW (GB). **RI-LEY, Robert, George**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral Merseyside CH63 3JW (GB).

(74) **Agent**: **TANSLEY, Sally, Elizabeth**; Unilever PLC, Unilever Patent Group, Colworth House, Sharnbrook, Bedfordshire MK44 1LQ (GB).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: HAIR COMPOSITION

(57) **Abstract**: A kit for colouring hair comprising: i) a first composition comprising a polyphenol ii) a second composition comprising a transition metal salt and/or complex; in which the first composition and/or the second composition further comprise a yellow and/or orange nitro dye.



- 1 -

HAIR COMPOSITION

The present invention relates to kits and methods for colouring hair.

5

Background

Two part colouring systems involving a metal ion and a polyphenol are described in EP0327345 (Beecham) and WO2010/135237 (Advance Cosmetic Technologies).

10

JP 04/164,017 (Lion) discloses a system for colouring hair consisting of shampoo containing a surfactant and gallic acid, tannic acid, salicylic acid, derivatives thereof, pyrogallol, catechol or haematin as a first agent. This is used in conjunction with a second agent containing a polyvalent metal salt and a cationic surfactant.

15

However; when metal/polyphenol based colouring systems are used to colour hair they can provide an unnatural blue shade, the effect is pronounced when using copper ion and/or iron ions.

20

The present system relates to a way of countering the blue hue.

Description of the Invention

25

Accordingly the present invention provides a kit for colouring hair comprising:

- i) a first composition comprising a polyphenol

- 2 -

ii) a second composition comprising an transition metal salt and/or complex;

in which the first composition and/or the second composition further comprise a yellow and/or orange nitro dye.

5

The invention further relates to a method of colouring hair comprising the steps of initially applying to the hair sequentially a composition comprising:

i) a first composition comprising a polyphenol

10

ii) a second composition comprising an transition metal salt and/or complex;

in which the first composition and/or the second composition further comprise a yellow and/or orange nitro dye.

15


The invention further relates to the use of a yellow and/or orange nitro dye to mitigate the blue hue on hair caused by a transition metal/polyphenol based colourant system

20

Detailed Description of the Invention

Yellow and Orange Nitro Dyes

25 The yellow and orange nitro dyes of the invention are substituted derivatives of nitrobenzene. The benzene ring is preferably substituted by one or two amino groups, preferably selected from -NH(CH₂CH₂OH) and NH(CH₂CH₂NH₂) and



- 3 -

The nitrobenzene ring may be further substituted by halides, alkyl and alkoxy groups, preferably selected from Cl, CF₃, OCH₂CH₂OH, OCH₃.

5 The dye preferably has a maximum optical absorption in the visible range (400-700nm) between 400 and 480nm in aqueous solution.

10 The yellow and orange dyes are preferably selected from HC Orange no 1, HC Orange no 2, HC Yellow no 4, HC Yellow no 2, HC Yellow no 9, HC Yellow no 10, HC Yellow no 13, more preferably , HC Yellow no 4, HC Yellow no 2, HC Yellow no 9, HC Yellow no 10, HC Yellow no 13, most preferably HC Yellow no 4, HC Yellow no 2. Yellow nitro dyes are particularly preferred.

The CAS-NO and COLIPA number of the preferred dyes are given below:

15

INCI Name	CAS-No	Colipa Number
HC ORANGE NO. 1	54381-08-7	B047
HC ORANGE NO. 2	85765-48-6	B067
HC YELLOW NO. 2	4926-55-0	B041
HC YELLOW NO.4	59820-43-8	B038
HC YELLOW NO. 9	141973-33-3	B069
HC YELLOW NO. 10	109023-83-8	B081
HC YELLOW NO. 13	10442-83-8	B102

Preferably the orange and yellow nitro dye is present at a level of the total composition from 0.01 to 1.5wt%, more preferably from 0.05 to 0.2wt%.

20

Preferably the yellow/orange nitro dye is present in the composition comprising the polyphenol and is absent from the composition comprising the transition metal. In a more preferred embodiment the yellow/orange nitro dye is present in a shampoo composition comprising polyphenol.

- 4 -

Polyphenol

The first composition of the kit comprises a polyphenol. The polyphenol of the invention comprises at least one benzene ring substituted with a first OH group
5 and a second OH or OCH₃ group, preferably OH. Preferably the first OH is ortho to the second OH group.

The OH containing ring may be fused to aromatic or non-aromatic rings. Polyphenols may be found in the classes of phenolic acid, anthocyanin,
10 anthocyanidin, flavanols, flavanones and gallates. For example flavan-3-ols, flavan-4-ols, flavan-3,4-ols.

It is highly preferable if the polyphenol is not an anthraquinone, thus the polyphenol is not derived or based on henna.

15

A preferred polyphenol contains a benzene ring substituted in the:
1 position by an X group, wherein X is a carbon containing organic group, preferably selected from an ester group, COOH, chromenes and 3,4 dihydro-chromenes. 3 position by a H, OH or OCH₃ group, preferably an OH group.
20 4 position by an OH group. 5 position by an OH or OCH₃ group, preferably an OH group.

Most preferably the 2 and 6 positions are substituted by H.

25 Preferably the polyphenol is a ortho-dihydroxybenzene derivative.

Preferably the polyphenol has a molecular weight of less than 1,000, more preferably less than 500.

- 5 -

The polyphenol of the invention is preferably selected from gallic acid, methyl gallate, ethyl gallate, propyl gallate or mixtures thereof, most preferably propyl gallate

- 5 Methyl gallate, ethyl gallate, propyl gallate are obtainable by esterification of gallic acid. Preferably the gallic acid is extracted from a natural source, preferably from hydrolysable tannins, preferably extracted from tara plants or gallnuts. Preferably, the polyphenols do not contain any active oxidase enzymes and have been heat treated to destroy any enzyme activity.

10

The polyphenol containing composition is preferably an aqueous polyphenol solution. The polyphenol composition preferably comprises from 0.01 wt% to 10.0 wt% of the total composition, more preferably from 0.2 wt% to 4.0 wt %, most preferably 0.5 wt% to 2.5 wt% of polyphenol.

15

To avoid oxidation of the polyphenols by air it is preferred that the composition is stored in an airtight container, preferably a bottle closed with an air tight cap.

20

The aqueous polyphenol solution contains water, preferably as the dominant ingredient. Auxiliary ingredients may be present for example to increase the viscosity, perfume and help solubilise the polyphenol. Solubilising ingredients include organic solvents and surfactants.

25

Preferably, the water used to formulate all compositions has a French hardness of from 0 to 36 degrees, more preferably 0 to 24 degrees, most preferably from 0 to 2 degrees.

30

Preferably, the water used to formulate all compositions contains less than 1ppm of chlorine based bleaching agents such as chlorine dioxide or hypochlorite. Most preferably less than 50ppb.

- 6 -

Preferably the composition comprising the polyphenol is a shampoo composition.

Preferably the pH of the composition comprising the polyphenol is in the range 5.9

5 to 9, most preferably 6 to 7.


Metal

The second composition of the kit comprises a transition metal. Preferably the
10 transition metal ion is selected from Cu(II), Fe(II) and Fe(III), more preferably Fe(II) and Fe(III). The metal ion may be present in the form of a salt or a complex.

Iron and copper containing compositions of the invention preferably comprise
from 0.01wt% to 5 wt% of metal, more preferably 0.05 to 4 wt% more preferably
15 from 0.4 to 1 wt%.

The level of metal ion in the formulation may be determined by quantitative elemental analysis.

20 For the avoidance of doubt, if the formulation contains 2wt% of the complex.



With molecular weight of 446.14, then it will contain $55.84/446.14 \times 2 = 0.25\text{wt\%}$ of
25 Fe(II) (2 decimal places).

The iron ion is preferably present as a complex with a ligand. The ligand is preferably gluconate and/or lactate. The ligand-metal complex may be pre-formed

- 7 -

before addition to the shampoo for example iron (II) gluconate. The ligand metal complex may be formed in the composition by the addition of an iron salt and the ligand for example, FeCl_2 and sodium lactate or tartrate. Preferably, the mole ratio of iron ion to ligand is from 1:1 to 1:4, more preferably from 1:1.5 to 1:2.5.

5

Preferably, the iron ion is in the form of a complex that is soluble in aqueous solution at the pH of the formulation. Most preferably the iron ion in the form of a salt and/or complex has a solubility in demineralised water at the pH of the formulation of greater than 0.1g/L.

10

The ability of a ligand to complex with a metal ion, M^{n+} , may be defined by its pM^{n+} value, wherein

$$pM^{n+} = -\log_{10}[M^{n+}]_{free}$$

15

and the ligand concentration is 10^{-5} mol/L and the total metal ion concentration is 10^{-6} mol/L and $[M^{n+}]_{free}$ is the molar concentration of uncomplexed metal ions.

Preferably the ligand used has a lower pM^{n+} , than gallic acid. Preferably the pM^{n+} are greater than 8 more preferably greater than 10. pM^{n+} values are most

20

preferably measured at pH 4 in de-mineralised water, with the chloride salt of the metal ion, most preferably ferric chloride.

Polyamino carboxylic acids are preferably present at weight % levels of less than 1/30th of the main ligand, preferably they are absent from the iron or copper

25

containing formulation. A polyamino carboxylic acid is a compound containing two or more amines connected through carbon atoms to two or more carboxylic acid groups. 2,2',2'',2'''-(Ethane-1,2-diyl dinitrilo)tetraacetic acid is a polyamino carboxylic acid. Diethylene triamine pentaacetic acid is a polyamino carboxylic acid. Ethylenediamine-*N,N'*-disuccinic acid is a polyamino carboxylic acid.

30

- 8 -

Particularly preferred are iron (III) gluconate, iron (II) gluconate and iron(III) lactate, iron(II) lactate, most preferably iron (III) gluconate.

Preferably, the iron is in a composition which further comprises a cationic surfactant, such as a conditioner composition.

5

Preferably pH of the second composition comprising the transition metal is in the range 5.5 to 8, most preferably 6 to 7.

Shampoo Composition

10

Shampoo compositions of the invention are generally aqueous, i.e. they have water or an aqueous solution or a lyotropic liquid crystalline phase as their major component. Suitably, the composition will comprise from 50 % to 98%, preferably from 60 % to 90% water by weight based on the total weight of the composition.

15

Anionic Cleansing Surfactant

Shampoo compositions according to the invention will generally comprise one or more anionic cleansing surfactants which are cosmetically acceptable and

20

suitable for topical application to the hair.

Examples of suitable anionic cleansing surfactants are the alkyl sulphates, alkyl ether sulphates, alkaryl sulphonates, alkanoyl isethionates, alkyl succinates, alkyl sulphosuccinates, alkyl ether sulphosuccinates, N-alkyl sarcosinates, alkyl

25

phosphates, alkyl ether phosphates, and alkyl ether carboxylic acids and salts thereof, especially sodium, magnesium, ammonium and mono-, di- and

triethanolamine salts. The alkyl and acyl groups generally contain from 8 to 18, preferably from 10 to 16 carbon atoms and may be unsaturated. The alkyl ether

sulphates, alkyl ether sulphosuccinates, alkyl ether phosphates and alkyl ether

30

carboxylic acids and salts thereof may contain from 1 to 20 ethylene oxide or propylene oxide units per molecule.

- 9 -

Typical anionic cleansing surfactants for use in shampoo compositions of the invention include sodium oleyl succinate, ammonium lauryl sulphosuccinate, sodium lauryl sulphate, sodium lauryl ether sulphate, sodium lauryl ether
5 sulphosuccinate, ammonium lauryl sulphate, ammonium lauryl ether sulphate, sodium dodecylbenzene sulphonate, triethanolamine dodecylbenzene sulphonate, sodium cocoyl isethionate, sodium lauryl isethionate, lauryl ether carboxylic acid and sodium N-lauryl sarcosinate.

10 Preferred anionic cleansing surfactants are sodium lauryl sulphate, sodium lauryl ether sulphate(n)EO, (where n is from 1 to 3), sodium lauryl ether sulphosuccinate(n)EO, (where n is from 1 to 3), ammonium lauryl sulphate, ammonium lauryl ether sulphate(n)EO, (where n is from 1 to 3), sodium cocoyl isethionate and lauryl ether carboxylic acid (n) EO (where n is from 10 to 20).

15

Mixtures of any of the foregoing anionic cleansing surfactants may also be suitable.

The total amount of anionic cleansing surfactant in shampoo compositions of the invention generally ranges from 0.5 % to 45%, preferably from 1.5 % to 35%,
20 more preferably from 5 % to 20 % by total weight anionic cleansing surfactant based on the total weight of the composition.

Further Ingredients

25

Optionally, a shampoo composition of the invention may contain further ingredients as described below to enhance performance and/or consumer acceptability.

30

- 10 -

Co-surfactant

The composition can include co-surfactants, to help impart aesthetic, physical or cleansing properties to the composition.

5

An example of a co-surfactant is a nonionic surfactant, which can be included in an amount ranging from 0.5 % to 8%, preferably from 2 % to 5% by weight based on the total weight of the composition.


10 For example, representative nonionic surfactants that can be included in shampoo compositions of the invention include condensation products of aliphatic (C₈ - C₁₈) primary or secondary linear or branched chain alcohols or phenols with alkylene oxides, usually ethylene oxide and generally having from 6 to 30 ethylene oxide groups.

15

Other representative nonionic surfactants include mono- or di-alkyl alkanolamides. Examples include coco mono- or di-ethanolamide and coco mono-isopropanolamide.

20 Further nonionic surfactants which can be included in shampoo compositions of the invention are the alkyl polyglycosides (APGs). Typically, APG is one which comprises an alkyl group connected (optionally via a bridging group) to a block of one or more glycosyl groups. Preferred APGs are defined by the following formula:

25



wherein R is a branched or straight chain alkyl group which may be saturated or unsaturated and G is a saccharide group.

30 R may represent a mean alkyl chain length of from about C₅ to about C₂₀. Preferably R represents a mean alkyl chain length of from about C₈ to about C₁₂.

- 11 -

Most preferably the value of R lies between about 9.5 and about 10.5. G may be selected from C₅ or C₆ monosaccharide residues, and is preferably a glucoside. G may be selected from the group comprising glucose, xylose, lactose, fructose, mannose and derivatives thereof. Preferably G is glucose.

5

The degree of polymerisation, n, may have a value of from about 1 to about 10 or more. Preferably, the value of n lies from about 1.1 to about 2. Most preferably the value of n lies from about 1.3 to about 1.5.

- 10 Suitable alkyl polyglycosides for use in the invention are commercially available and include for example those materials identified as: Oramix NS10 ex Seppic; Plantaren 1200 and Plantaren 2000 ex Henkel.

- 15 Other sugar-derived nonionic surfactants which can be included in compositions of the invention include the C₁₀-C₁₈ N-alkyl (C₁-C₆) polyhydroxy fatty acid amides, such as the C₁₂-C₁₈ N-methyl glucamides, as described for example in WO 92 06154 and US 5 194 639, and the N-alkoxy polyhydroxy fatty acid amides, such as C₁₀-C₁₈ N-(3-methoxypropyl) glucamide.

- 20 A preferred example of a co-surfactant is an amphoteric or zwitterionic surfactant, which can be included in an amount ranging from 0.5% to about 8%, preferably from 1% to 4% by weight based on the total weight of the composition.

- 25 Examples of amphoteric or zwitterionic surfactants include alkyl amine oxides, alkyl betaines, alkyl amidopropyl betaines, alkyl sulphobetaines (sultaines), alkyl glycinate, alkyl carboxyglycinates, alkyl amphoacetates, alkyl amphopropionates, alkylamphoglycinates, alkyl amidopropyl hydroxysultaines, acyl taurates and acyl glutamates, wherein the alkyl and acyl groups have from 8 to 19 carbon atoms. Typical amphoteric and zwitterionic surfactants for use in shampoos of the

- 12 -

invention include lauryl amine oxide, cocodimethyl sulphopropyl betaine, lauryl betaine, cocamidopropyl betaine and sodium cocoamphoacetate.

5 A particularly preferred amphoteric or zwitterionic surfactant is cocoamidopropyl betaine.

Mixtures of any of the foregoing amphoteric or zwitterionic surfactants may also be suitable. Preferred mixtures are those of cocoamidopropyl betaine with further amphoteric or zwitterionic surfactants as described above. A preferred further
10 amphoteric or zwitterionic surfactant is sodium cocoamphoacetate.

The total amount of surfactant (including any co-surfactant, and/or any emulsifier) in a shampoo composition of the invention is generally from 1% to 50%, preferably from 2 % to 40%, more preferably from 10 % to 25% by total weight surfactant
15 based on the total weight of the composition.

Cationic Polymers

Cationic polymers are preferred ingredients in a shampoo composition of the
20 invention for enhancing conditioning performance.

Suitable cationic polymers may be homopolymers which are cationically substituted or may be formed from two or more types of monomers. The weight average (M_w) molecular weight of the polymers will generally be between 100,000
25 and 2 million daltons. The polymers will have cationic nitrogen containing groups such as quaternary ammonium or protonated amino groups, or a mixture thereof. If the molecular weight of the polymer is too low, then the conditioning effect is poor. If too high, then there may be problems of high extensional viscosity leading to stringiness of the composition when it is poured.

30

- 13 -

The cationic nitrogen-containing group will generally be present as a substituent on a fraction of the total monomer units of the cationic polymer. Thus when the polymer is not a homopolymer it can contain spacer non-cationic monomer units. Such polymers are described in the CTFA Cosmetic Ingredient Directory, 3rd edition. The ratio of the cationic to non-cationic monomer units is selected to give polymers having a cationic charge density in the required range, which is generally from 0.2 to 3.0 meq/gm. The cationic charge density of the polymer is suitably determined via the Kjeldahl method as described in the US Pharmacopoeia under chemical tests for nitrogen determination.

Suitable cationic polymers include, for example, copolymers of vinyl monomers having cationic amine or quaternary ammonium functionalities with water soluble spacer monomers such as (meth)acrylamide, alkyl and dialkyl (meth)acrylamides, alkyl (meth)acrylate, vinyl caprolactone and vinyl pyrrolidine. The alkyl and dialkyl substituted monomers preferably have C1-C7 alkyl groups, more preferably C1-3 alkyl groups. Other suitable spacers include vinyl esters, vinyl alcohol, maleic anhydride, propylene glycol and ethylene glycol.

The cationic amines can be primary, secondary or tertiary amines, depending upon the particular species and the pH of the composition. In general secondary and tertiary amines, especially tertiary, are preferred.

Amine substituted vinyl monomers and amines can be polymerized in the amine form and then converted to ammonium by quaternization.

The cationic polymers can comprise mixtures of monomer units derived from amine- and/or quaternary ammonium-substituted monomer and/or compatible spacer monomers.

- 14 -

Suitable cationic polymers include, for example:

- cationic diallyl quaternary ammonium-containing polymers including, for example, dimethyldiallylammonium chloride homopolymer and copolymers of acrylamide and dimethyldiallylammonium chloride, referred to in the industry (CTFA) as Polyquaternium 6 and Polyquaternium 7, respectively;
- mineral acid salts of amino-alkyl esters of homo-and co-polymers of unsaturated carboxylic acids having from 3 to 5 carbon atoms, (as described in U.S. Patent 4,009,256);
- cationic polyacrylamides (as described in WO95/22311).

Other cationic polymers that can be used include cationic polysaccharide polymers, such as cationic cellulose derivatives, cationic starch derivatives, and cationic guar gum derivatives.

Cationic polysaccharide polymers suitable for use in compositions of the invention include monomers of the formula:



wherein: A is an anhydroglucose residual group, such as a starch or cellulose anhydroglucose residual. R is an alkylene, oxyalkylene, polyoxyalkylene, or hydroxyalkylene group, or combination thereof. R¹, R² and R³ independently represent alkyl, aryl, alkylaryl, arylalkyl, alkoxyalkyl, or alkoxyaryl groups, each group containing up to about 18 carbon atoms. The total number of carbon atoms for each cationic moiety (i.e., the sum of carbon atoms in R¹, R² and R³) is preferably about 20 or less, and X is an anionic counterion.

- 15 -

Another type of cationic cellulose includes the polymeric quaternary ammonium salts of hydroxyethyl cellulose reacted with lauryl dimethyl ammonium-substituted epoxide, referred to in the industry (CTFA) as Polyquaternium 24. These materials are available from the Amerchol Corporation, for instance under the tradename Polymer LM-200.

Other suitable cationic polysaccharide polymers include quaternary nitrogen-containing cellulose ethers (e.g. as described in U.S. Patent 3,962,418), and copolymers of etherified cellulose and starch (e.g. as described in U.S. Patent 3,958,581).

A particularly suitable type of cationic polysaccharide polymer that can be used is a cationic guar gum derivative, such a guar hydroxypropyltrimethylammonium chloride (commercially available from Rhodia in their JAGUAR trademark series). Examples of such materials are JAGUAR C13S, JAGUAR C14, JAGUAR C15, JAGUAR C17 and JAGUAR C16 Jaguar CHT and JAGUAR C162.

Mixtures of any of the above cationic polymers may be used.

Cationic polymer will generally be present in a shampoo composition of the invention at levels of from 0.01 % to 5%, preferably from 0.05 % to 1 %, more preferably from 0.08 % to 0.5% by total weight of cationic polymer based on the total weight of the composition.

Suspending Agent

Preferably, an aqueous shampoo composition of the invention further comprises a suspending agent. Suitable suspending agents are selected from polyacrylic acids, cross-linked polymers of acrylic acid, copolymers of acrylic acid with a hydrophobic monomer, copolymers of carboxylic acid-containing monomers and

- 16 -

acrylic esters, cross-linked copolymers of acrylic acid and acrylate esters, heteropolysaccharide gums and crystalline long chain acyl derivatives. The long chain acyl derivative is desirably selected from ethylene glycol stearate, alkanolamides of fatty acids having from 16 to 22 carbon atoms and mixtures thereof. Ethylene glycol distearate and polyethylene glycol 3 distearate are preferred long chain acyl derivatives, since these impart pearlescence to the composition. Polyacrylic acid is available commercially as Carbopol 420, Carbopol 488 or Carbopol 493. Polymers of acrylic acid cross-linked with a polyfunctional agent may also be used; they are available commercially as Carbopol 910, Carbopol 934, Carbopol 941 and Carbopol 980. An example of a suitable copolymer of a carboxylic acid containing monomer and acrylic acid esters is Carbopol 1342. All Carbopol (trademark) materials are available from Lubrizol.

Suitable cross-linked polymers of acrylic acid and acrylate esters are Pemulen TR1 or Pemulen TR2. A suitable heteropolysaccharide gum is xanthan gum, for example that available as Kelzan mu.

Mixtures of any of the above suspending agents may be used. Preferred is a mixture of cross-linked polymer of acrylic acid and crystalline long chain acyl derivative.

Suspending agent will generally be present in a shampoo composition of the invention at levels of from 0.1 % to 10%, preferably from 0.5 % to 6%, more preferably from 0.9 % to 4% by total weight of suspending agent based on the total weight of the composition.

Preferably the pH of shampoo is in the range from 5.9 to 9, most preferably from 6 to 7 at 20°C.

- 17 -

Conditioner Compositions

One part of the kit may be a conditioner for the treatment of hair (typically after shampooing) and subsequent rinsing. Such conditioner compositions preferably
5 comprise the iron salt/complex. Such conditioner compositions will typically comprise one or more conditioning surfactants which are cosmetically acceptable and suitable for topical application to the hair.

Suitable conditioning surfactants include those selected from cationic surfactants,
10 used singly or in admixture. Preferably, the cationic surfactants have the formula $N^+R^1R^2R^3R^4$ wherein R^1 , R^2 , R^3 and R^4 are independently (C_1 to C_{30}) alkyl or benzyl. Preferably, one, two or three of R^1 , R^2 , R^3 and R^4 are independently (C_4 to C_{30}) alkyl and the other R^1 , R^2 , R^3 and R^4 group or groups are (C_1 - C_6) alkyl or benzyl. More preferably, one or two of R^1 , R^2 , R^3 and R^4 are independently (C_6 to
15 C_{30}) alkyl and the other R^1 , R^2 , R^3 and R^4 groups are (C_1 - C_6) alkyl or benzyl groups. Optionally, the alkyl groups may comprise one or more ester (-OCO- or -COO-) and/or ether (-O-) linkages within the alkyl chain. Alkyl groups may optionally be substituted with one or more hydroxyl groups. Alkyl groups may be straight chain or branched and, for alkyl groups having 3 or more carbon atoms,
20 cyclic. The alkyl groups may be saturated or may contain one or more carbon-carbon double bonds (eg, oleyl). Alkyl groups are optionally ethoxylated on the alkyl chain with one or more ethyleneoxy groups.


Suitable cationic surfactants for use in conditioner compositions according to the
25 invention include cetyltrimethylammonium chloride, behenyltrimethylammonium chloride, cetylpyridinium chloride, tetramethylammonium chloride, tetraethylammonium chloride, octyltrimethylammonium chloride, dodecyltrimethylammonium chloride, hexadecyltrimethylammonium chloride, octyldimethylbenzylammonium chloride, decyldimethylbenzylammonium chloride,
30 stearyldimethylbenzylammonium chloride, didodecyldimethylammonium chloride, dioctadecyldimethylammonium chloride, tallowtrimethylammonium chloride,

- 18 -

dihydrogenated tallow dimethyl ammonium chloride (eg, Arquad 2HT/75 from Akzo Nobel), cocotrimethylammonium chloride, PEG-2-oleammonium chloride and the corresponding hydroxides thereof. Further suitable cationic surfactants include those materials having the CTFA designations Quaternium-5, Quaternium-31 and Quaternium-18. Mixtures of any of the foregoing materials may also be suitable. A particularly useful cationic surfactant for use in conditioners according to the invention is cetyltrimethylammonium chloride, available commercially, for example as GENAMIN CTAC, ex Hoechst Celanese. Another particularly useful cationic surfactant for use in conditioners according to the invention is behenyltrimethylammonium chloride, available commercially, for example as GENAMIN KDMP, ex Clariant.

Another example of a class of suitable cationic surfactants for use in the invention, either alone or together with one or more other cationic surfactants, is a combination of (i) and (ii) below:

(i) an amidoamine corresponding to the general formula (I):



in which R^1 is a hydrocarbyl chain having 10 or more carbon atoms, R^2 and R^3 are independently selected from hydrocarbyl chains of from 1 to 10 carbon atoms, and m is an integer from 1 to about 10; and

(ii) an acid.

As used herein, the term hydrocarbyl chain means an alkyl or alkenyl chain.

- 19 -

Preferred amidoamine compounds are those corresponding to formula (I) in which R^1 is a hydrocarbyl residue having from about 11 to about 24 carbon atoms, R^2 and R^3 are each independently hydrocarbyl residues, preferably alkyl groups, having from 1 to about 4 carbon atoms, and m is an integer from 1 to about 4.

5

Preferably, R^2 and R^3 are methyl or ethyl groups.

Preferably, m is 2 or 3, i.e. an ethylene or propylene group.

Preferred amidoamines useful herein include stearamido-propyldimethylamine, stearamidopropyldiethylamine, stearamidoethyldiethylamine,

10 stearamidoethyldimethylamine, palmitamidopropyldimethylamine, palmitamidopropyl-diethylamine, palmitamidoethyldiethylamine, palmitamidoethyldimethylamine, behenamidopropyldimethyl-amine, behenamidopropyldiethylamine, behenamidoethyldiethyl-amine, behenamidoethyldimethylamine, arachidamidopropyl-dimethylamine, 15 arachidamidopropyldiethylamine, arachid-amidoethyldiethylamine, arachidamidoethyldimethylamine, and mixtures thereof.

Particularly preferred amidoamines useful herein are

stearamidopropyldimethylamine, stearamidoethyldiethylamine, and mixtures 20 thereof.

Commercially available amidoamines useful herein include:

stearamidopropyldimethylamine with tradenames LEXAMINE S-13 available from Inolex (Philadelphia Pennsylvania, USA) and AMIDOAMINE MSP available from 25 Nikko (Tokyo, Japan), stearamidoethyldiethylamine with a tradename AMIDOAMINE S available from Nikko, behenamidopropyldimethylamine with a tradename INCROMINE BB available from Croda (North Humberside, England), and various amidoamines with tradenames SCHERCODINE series available from Scher (Clifton New Jersey, USA).

30

- 20 -

A protonating acid may be present. Acid may be any organic or mineral acid which is capable of protonating the amidoamine in the conditioner composition. Suitable acids useful herein include hydrochloric acid, acetic acid, tartaric acid, fumaric acid, lactic acid, malic acid, succinic acid, and mixtures thereof.

- 5 Preferably, the acid is selected from the group consisting of acetic acid, tartaric acid, hydrochloric acid, fumaric acid, lactic acid and mixtures thereof.

The primary role of the acid is to protonate the amidoamine in the hair treatment composition thus forming a tertiary amine salt (TAS) in situ in the hair treatment
10 composition. The TAS in effect is a non-permanent quaternary ammonium or pseudo-quaternary ammonium cationic surfactant.

Suitably, the acid is included in a sufficient amount to protonate more than 95 mole% (293 K) of the amidoamine present.
15

In conditioners of the invention, the level of cationic surfactant will generally range from 0.01 % to 10%, more preferably 0.05 % to 7.5%, most preferably 0.1 % to 5% by weight of the composition.

20 Conditioners of the invention will typically also incorporate a fatty alcohol. The combined use of fatty alcohols and cationic surfactants in conditioning compositions is believed to be especially advantageous, because this leads to the formation of a lamellar phase, in which the cationic surfactant is dispersed.

25 Representative fatty alcohols comprise from 8 to 22 carbon atoms, more preferably 16 to 22. Fatty alcohols are typically compounds containing straight chain alkyl groups. Examples of suitable fatty alcohols include cetyl alcohol, stearyl alcohol and mixtures thereof. The use of these materials is also advantageous in that they contribute to the overall conditioning properties of
30 compositions of the invention.

- 21 -

The level of fatty alcohol in conditioners of the invention will generally range from 0.01 % to 10%, preferably from 0.1 % to 8%, more preferably from 0.2 % to 7%, most preferably from 0.3% to 6% by weight of the composition. The weight ratio
5 of cationic surfactant to fatty alcohol is suitably from 1:1 to 1:10, preferably from 1:1.5 to 1:8, optimally from 1:2 to 1:5. If the weight ratio of cationic surfactant to fatty alcohol is too high, this can lead to eye irritancy from the composition. If it is too low, it can make the hair feel squeaky for some consumers.

- 10 Preferably the pH of conditioner is in the range from 5.5 to 8, more preferably 6 to 7 at 20°C.

Further Conditioning Agents

- 15 Compositions of the invention may comprise further conditioning agents to optimise wet and dry conditioning benefits.

The compositions of the invention can contain, emulsified droplets of a silicone conditioning agent, for enhancing conditioning performance. Silicone conditioning
20 agents may be present in the shampoo or conditioner.

- Suitable silicones include polydiorganosiloxanes, in particular polydimethylsiloxanes which have the CTFA designation dimethicone. Also suitable for use compositions of the invention (particularly shampoos and
25 conditioners) are polydimethyl siloxanes having hydroxyl end groups, which have the CTFA designation dimethiconol. Also suitable for use in compositions of the invention are silicone gums having a slight degree of cross-linking, as are described for example in WO 96/31188.

- 22 -

The viscosity of the emulsified silicone itself (not the emulsion or the final hair conditioning composition) is typically at least 10,000 cst at 25 °C the viscosity of the silicone itself is preferably at least 60,000 cst, most preferably at least 500,000 cst, ideally at least 1,000,000 cst. Preferably the viscosity does not
5 exceed 10^9 cst for ease of formulation.

Emulsified silicones for use in the shampoo compositions of the invention will typically have an average silicone droplet size in the composition of less than 30, preferably less than 20, more preferably less than 10 μm , ideally from 0.01 to
10 1 μm . Silicone emulsions having an average silicone droplet size of $\leq 0.15 \mu\text{m}$ are generally termed microemulsions.

Emulsified silicones for use in the conditioner compositions of the invention will typically have a size in the composition of less than 30, preferably less than 20,
15 more preferably less than 15. Preferably, the average silicone droplet is greater than 0.5 μm , more preferably greater than 1 μm , ideally from 2 to 8 μm .

Silicone particle size may be measured by means of a laser light scattering technique, for example using a 2600D Particle Sizer from Malvern Instruments.
20

Examples of suitable pre-formed emulsions include Xiameter MEM 1785 and microemulsion DC2-1865 available from Dow Corning. These are emulsions/microemulsions of dimethiconol. Cross-linked silicone gums are also available in a pre-emulsified form, which is advantageous for ease of formulation.
25

A further preferred class of silicones for inclusion in shampoos and conditioners of the invention are amino functional silicones. By "amino functional silicone" is meant a silicone containing at least one primary, secondary or tertiary amine group, or a quaternary ammonium group. Examples of suitable amino functional
30 silicones include: polysiloxanes having the CTFA designation "amodimethicone".

- 23 -

Specific examples of amino functional silicones suitable for use in the invention are the aminosilicone oils DC2-8220, DC2-8166 and DC2-8566 (all ex Dow Corning).

- 5 Suitable quaternary silicone polymers are described in EP-A-0 530 974. A preferred quaternary silicone polymer is K3474, ex Goldschmidt. Also suitable are emulsions of amino functional silicone oils with non ionic and/or cationic surfactant.
- 10 Pre-formed emulsions of amino functional silicone are also available from suppliers of silicone oils such as Dow Corning and General Electric. Specific examples include DC939 Cationic Emulsion and the non-ionic emulsions DC2-7224, DC2-8467, DC2-8177 and DC2-8154 (all ex Dow Corning).
- 15 With some shampoos it is preferred to use a combination of amino and non amino functional silicones.

The total amount of silicone is preferably from 0.01 wt% to 10 %wt of the total composition more preferably from 0.1 wt% to 5 wt%, most preferably 0.5 wt% to

20 3 wt% is a suitable level, especially for a shampoo composition.

Other Optional Ingredients

- A composition of the invention may contain other ingredients for enhancing
- 25 performance and/or consumer acceptability. Such ingredients include fragrance, dyes and pigments, pH adjusting agents, pearlescers or opacifiers, viscosity modifiers, preservatives, and natural hair nutrients such as botanicals, fruit extracts, sugar derivatives and amino acids.

- 24 -

Product Form

Compositions within the kit of the invention are typically “rinse-off” compositions to be applied to the hair and then rinsed away.

5

To avoid oxidation by air it is preferred that the compositions are stored in an air tight container such as a bottle closed with an air tight cap.

10 In one embodiment the kit preferably comprises a single dose (30 to 100ml) thio pre-treatment composition and multi-dose (150 to 300ml) shampoo and conditioner.

Method of use

15 Preferably, the compositions of the invention are applied to wet hair, necessitating the step of wetting the hair before application of the compositions of the invention.

20 Preferably the polyphenol is applied followed by the Fe-complex, preferably the polyphenol formulation is a shampoo and the iron complex is carried in a conditioner formulation.

A method preferably comprises the step of rinsing hair between application of the compositions of the invention.

25 Further conditioning and/or styling products may be applied as part of the colouring process.

The level of each composition applied to the head of hair is preferably from 5g to 100g.

30

- 25 -

Preferably, each composition comprising polyphenol or iron complex/salt remains on the hair for 5 to 600 seconds, more preferably 10 to 300 seconds.

5 Preferably, the water used to wet and rinse the hair has a French hardness of from 0 to 36 degrees, more preferably 0 to 24 degrees, most preferably from 0 to 2 degrees.

10 Preferably, the water used to wet and rinse the hair contains less than 1ppm of chlorine based bleaching agents such as chlorine dioxide or hypochlorite. Most preferably less than 50ppb

The invention will now be illustrated by the following non-limiting examples:

Examples

15 Shampoo formulation containing propyl gallate and yellow dye; and a conditioner formulation containing iron gluconate were prepared. The formulation details are given below.

- 26 -

Shampoo

Ingredient	wt%
Sodium Laureth Sulfate	17.0
Cocamidopropylbetaine	5.3
Carbomer	0.4
Hydrochloric acid	0.06
Mica	0.15
DMDM Hydantoin	0.1
Silicone	1.7
Guar Hydroxypropyltrimonium Chloride	0.2
Propyl gallate	2
Yellow dye	0.1
Sodium Chloride	0.5
NaOH	To pH = 6.5
water	to 100

The yellow dye was added last. A control shampoo was also made that did not contain yellow dye.

- 27 -






Conditioner

Ingredient	wt%
Behenyltrimonium Chloride	1.6
Stearamidopropyl Dimethylamine	0.4
Cetearyl Alcohol	4
Stearyl alc. And Cetearth 20	1.5
Fragrance	0.9
Silicone	1.5
DMDM Hydantoin	0.1
Methylchloroisothiazolinone Methylisothiazolinone	0.04
Iron Gluconate	4.0
Sodium hydroxide	To pH = 6.5
water	To 100

The shampoo and conditioner formulation were used to treat white Chinese hair
 switches. The hair was wet with water. 0.43g Shampoo per g hair was applied to
 5 the white hair and massaged into the hair for 30 seconds, the hair left for 3
 minutes, then rinsed for 30 seconds and the procedure repeated. 0.43g per g hair
 of the conditioner formulation was then applied and massaged into the hair for 30
 seconds, the hair left for 3 minutes, then rinsed for 30 seconds. The hair was
 10 dried and the colour measured using a reflectometer and expressed as the CIE
 L*a*b* values. The results are given in the table below.

The b* value is a measure of the colour on the yellow-blue axis, positive values
 indicate yellow and negative values blue. The initial b* of the switches was 25.6.

- 28 -

Dye	Structure	b*
Control (no dye)	-	1.6
Basic Orange 31 (CAS 97404-02-9)	 reference	- 0.5
HC Yellow no 7 (CAS 104226-21-3)	 reference	1.9
HC Orange no 1 (CAS 54381-08-7)		2.6
HC Yellow no 4 (59820-43-8)		3.6
HC Yellow no 2 (CAS 4926-55-0)		3.6

The control wash, without yellow dyes leads to a large drop in b*, due to the colouring effect of the propyl gallate, iron complex formed in the hair. Addition of nitro dyes, HC Orange no 1, HC Yellow no 4 and HC Yellow no 2 gives the largest increase in b* compared to the control and other yellow/orange dyes which excluded the nitro moiety.

10 The yellow nitro dyes provide the highest b* values.

- 29 -

CLAIMS

1. A kit for colouring hair comprising:
 - 5 i) a first composition comprising a polyphenol
 - (ii) a second composition comprising a transition metal salt and/or complex; in which the first composition and/or the second composition further comprise a yellow and/or orange nitro dye.
- 10 2. A kit according to claim 1 in which the nitro dye is present in the first composition i) comprising the polyphenol and not present in the second composition comprising the transition metal.
- 15 3. A kit according to any preceding claim in which the transition metal of the second composition ii) is selected from iron or copper.
4. A kit according to claim 3 in which the transition metal is iron
- 20 5. A kit as claimed in any preceding claim in which the iron is present as a complex of gluconate and/or lactate.
6. A kit according to any preceding claim in which iron is present as iron (II) or iron (III) gluconate.
- 25 7. A kit according to any preceding claim in which the nitro dye is selected from HC Orange no 1, HC Orange no 2, HC Yellow no 4, HC Yellow no 2, HC Yellow no 9, HC Yellow no 10, HC Yellow no 13, and mixtures thereof.
- 30 8. A kit according to any preceding claim in which the nitro dye is a yellow nitro dye.

- 30 -

9. A kit as claimed in any preceding claim in which the pH of the first composition is from 5.9 to 9 at 20° C.
- 5 10. A kit as claimed in any preceding claim in which the polyphenol is selected from selected from methyl gallate, ethyl gallate, propyl gallate and mixtures thereof.
- 10 11. A kit for colouring hair according to any preceding claim in which the iron containing composition further comprises a cationic conditioning surfactant.
12. A kit for colouring hair according to any preceding claim in which the polyphenol comprising composition further comprises an anionic surfactant.
- 15 13. A method of colouring hair comprising the steps of sequentially applying to the hair a kit for colouring hair comprising:
- i) a first composition comprising a polyphenol
- 20 (ii) a second composition comprising an transition metal salt and/or complex; in which the first composition and/or the second composition further comprise a yellow and/or orange nitro dye.
- 25 14. A method according to claim 12 which further comprises the step of rinsing the hair between application of the compositions.
15. A method according to claim 13 to 14 in which the polyphenol composition is applied to the hair before the iron comprising composition.
- 30 16. Use of a yellow and/or orange nitro dye to mitigate the blue hue on hair caused by a transition metal/polyphenol based colourant system.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/060590

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K8/37 A61K8/41 A61Q5/10 A61K8/19 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K A61Q		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP 2010 248103 A (SHISEI CHEMICAL KK) 4 November 2010 (2010-11-04) paragraphs [0006], [0009], [0010], [0012], [0013], [0019], [0023] - [0025], [0032], [0039] example 13	1-15
X	----- US 5 628 799 A (WENKE GOTTFRIED [US] ET AL) 13 May 1997 (1997-05-13) example 71 ----- -/--	1-4, 8, 9, 13
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search	Date of mailing of the international search report	
8 July 2013	15/07/2013	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Lenzen, Achim	

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2013/060590

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	Anonymous: "Green tea extract", Wikipedia, the free encyclopedia 7 March 2013 (2013-03-07), pages 1-4, XP055060866, Retrieved from the Internet: URL:http://en.wikipedia.org/wiki/Green_tea_extract [retrieved on 2013-04-24] page 2, paragraph 2 under "Biochemical properties" -----	1-16
A	Anonymous: "2-Amino-6-Chloro-4-Nitrophenol", Cosmetics Info page 2PP, XP002695924, Retrieved from the Internet: URL:http://www.cosmeticsinfo.org/ingredient_details.php?ingredient_id=1348 [retrieved on 2013-04-23] page 1, paragraph 1 -----	1-16
A	Anonymous: "Polyphenol", Wikipedia, the free encyclopedia 28 June 2012 (2012-06-28), pages 1-23, XP002679272, Retrieved from the Internet: URL:http://en.wikipedia.org/wiki/Polyphenol 1 [retrieved on 2012-07-05] page 2 -----	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2013/060590

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 2010248103	A	04-11-2010	NONE	

US 5628799	A	13-05-1997	NONE	
