



(51) International Patent Classification:

A01K 67/033 (2006.01) *A23K 50/00* (2016.01)
A01K 29/00 (2006.01) *A23K 50/90* (2016.01)
A01K 67/00 (2006.01)

(21) International Application Number:

PCT/US2017/050778

(22) International Filing Date:

08 September 2017 (08.09.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/385,125 08 September 2016 (08.09.2016) US

(71) Applicant: RUTGERS, THE STATE UNIVERSITY OF
NEW JERSEY [US/US]; Office of Corporate Liaison &
Technology Transfer, ASB III, 3 Rutgers Plaza, 3rd Floor,
New Brunswick, NJ 08901-8559 (US).

(72) Inventors: GAUGLER, Randy; 25 Quince Place, North
Brunswick, NJ 08902 (US). SUMAN, Devi S.; 9S, 9th Av-
enue, Apt. 3-K, Highland, NJ 08904 (US). CHANDEL,
Kshitij; 9S, 9th Avenue, Apt. 3-K, Highland Park, NJ
08094 (US). WANG, Yi; 3 Continental Court, South River,
NJ 08882 (US).

(74) Agent: RIGAUT, Kathleen D. et al.; Dann, Dorfman, Her-
rell and Skillman, 1601 Market Street, Suite 2400, Philadel-
phia, PA 19103-2307 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: A NON-MEMBRANNE FEEDING DEVICE AND DIET FORMULATION FOR MOSQUITO COLONY PRODUCTION




Fig. 1A

(57) Abstract: An artificial feeding system for maintenance and growth of hematophagous insects is disclosed to comprise a blood-free diet formulation and a feeding device. The blood-free diet formulation comprises effective amounts of a protein source and a carbohydrate source, wherein the blood-free diet formulation is suitable for egg production and colony maintenance of hematophagous insects. The feeding device comprises a diet reservoir for receiving the blood-free diet formulation and a feeding platform for hematophagous insects to feed the blood-free diet formulation from the diet reservoir.

A NON-MEMBRANE FEEDING DEVICE AND DIET FORMULATION FOR MOSQUITO COLONY PRODUCTION

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application claims priority from U.S. provisional patent application serial number 62/385,125, entitled "A non-membrane feeding device and diet formulation for mosquito colony production", filed September 08, 2016. The entire disclosure of this application is incorporated herein by reference.

FIELD OF THE INVENTION

10 This invention relates to the fields of mosquito maintenance and propagation. More specifically a non-membrane feeding device and diet formulation for mosquito colony production is disclosed.

BACKGROUND OF THE INVENTION

15 Numerous publications and patent documents, including both published applications and issued patents, are cited throughout the specification in order to describe the state of the art to which this invention pertains. The disclosure of each of these citations is incorporated herein by reference as though set forth in full.

20 Up to 700 million people are infected and more than a million die each year from mosquito-borne illnesses (Caraballo, Hector (May 2014)). Several diseases transmitted by mosquitoes are malaria, dengue, filariasis, West Nile virus, chikungunya, yellow fever, etc. The next generation of mosquito control technologies seem unlikely to require unilateral use of the toxic chemicals that have been dominate over the past 75 years. The new ecologically benign technologies currently being devised are increasingly focused on genetically modified or sterile mosquitoes, methods that require production of massive numbers of mosquitoes for field release.

25 Female mosquitoes generally require a blood meal to develop eggs for the next generation. Mosquitoes are divergent in host preference, which include humans, animals, birds, reptiles, and amphibians. They also vary in daily rhythms of blood feeding such as diurnal, nocturnal and crepuscular etc. In a laboratory setup, cultures are maintained mostly with a blood feeding system, where female mosquitoes are fed with vertebrate blood. This approach has certain

30 drawbacks. While the diet rich is in protein, diets based on vertebrate blood are expensive,

inconsistent in quality, and require pathogen-free certification. Moreover, such blood based diets are unstable during storage. Indeed, the blood must be maintained at 37 °C and most approaches require a membrane to mimic vertebrate skin for the mosquito to pierce. Laboratories maintaining mosquito colonies throughout the world are dependent on live animals or animal blood. Hence, production of large quantities of mosquitoes for mass-release in mosquito control trials places great pressure on animal care facilities and increases maintenance costs several-fold. The blood for these systems usually comes from live animals or birds, which are under constant surveillance and ethical regulations, requiring stringent animal care. Consequently, a laboratory maintaining only a small number of mosquito cages expends thousands of dollars for the blood feeding arrangements which rely on use of animals or artificial membrane feeding system. Understandably then, considerable research effort has been devoted to the development of *artificial* alternatives to blood. In some labs, researchers use blood components or synthetic blood-free diets based on the host-seeking and blood-feeding behavior of mosquitoes, in which case, the membrane feeding system replaces animal blood or live animals. While an artificial membrane feeding system has been developed and is available commercially (Hemotek), this system still requires the use of collected animal blood. There are huge variations in acceptability of this artificial membrane system for various mosquito species. Blood quality, age and composition also affect system performance (Phasomkusolsil et al., 2013). Often, quality blood for colony maintenance can be difficult to obtain. Some studies have used blood components or synthetic blood-free diet with the membrane feeding system to replace animal blood or live animals. These diets are based on host-seeking and blood feeding behavior of mosquitoes and contain multiple components including phagostimulants, protein sources, digestive activators, buffer solutions, attractants, etc. The performance of these diets depends entirely upon the membrane feeding system. Further, there are concerns regarding long term storage and cost effectiveness of such systems, as adenosine tri-phosphate (ATP), a widely used phagostimulant in the *artificial* alternatives to blood, used to encourage feeding, is expensive.

Clearly, a need exists for a commercially viable product that will replace costly, labor intensive and ethically vulnerable animal use for blood feeding and artificial membrane feeding systems.

SUMMARY OF THE INVENTION

In accordance with the present invention, an artificial feeding system for maintenance and growth of hematophagous insects is provided which exploits sugar feeding, rather than blood feeding behavior. An exemplary system comprises a diet reservoir for receiving a synthetic, blood-free diet formulation; at least one feeding platform consisting of a mosquito proof covering separating said reservoir from said platform, the system and diet formulation being suitable for egg production and colony maintenance. Another exemplary artificial feeding system comprises a blood-free diet formulation and a feeding device comprising a diet reservoir for receiving the blood-free diet formulation; and a feeding platform for hematophagous insects to sit and feed on the blood-free diet formulation from the diet reservoir. The blood-free diet formulation comprises effective amounts of a protein source and a carbohydrate source, wherein the blood-free diet formulation is suitable for egg production and colony maintenance of hematophagous insects.

In one embodiment, the system comprises a single feeding platform which is optionally a mesh. In another embodiment, the system comprises multiple feeding platforms, e.g., 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more platforms. In an embodiment, the feeding platform floats over the blood-free diet formulation in the diet reservoir. In another embodiment, at least a portion of the feeding platform is hydrophobic. Yet in another embodiment, the feeding platform is a mesh.

The system can be formed from polypropylene, polyethylene, polystyrene or any other inert plastic material or a metal, such as aluminum foil, or biodegradable materials such as wax paper and the like. The feeding platform can be made of any suitable materials, including but not limited to, polystyrene, aluminum foil, wax paper, sponge, resin, wood and cardboard coated with non-wetting agents.

The diet reservoir of the system can be any shape such as circular or polygonal e.g., square, rectangular or triangular.

In another embodiment of the invention a blood free diet formulation is provided comprising effective amounts of a protein source, a carbohydrate source and, optionally a lipid

source. In certain embodiments, the protein source is bovine serum albumin, egg albumin or a mixture thereof. The carbohydrate source can be glucose, sucrose, fructose, or mixtures thereof.

In certain embodiments, the lipid source is cholesterol. In a particularly preferred embodiment, the blood-free diet formulation lacks adenosine tri-phosphate, a phagostimulant and supports several different species of mosquitoes including for example, *Aedes*, *Anopheles* and *Culex* mosquitoes.

In another aspect of the invention, a method of maintaining and growing hematophagus insects is provided. An exemplary method comprises: (a) providing an artificial feeding system as disclosed hereinabove; (b) dissolving the blood-free diet formulation in water; and (c) feeding the hematophagus insects.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate particular embodiment(s) of the invention, and together with the written description, serve to explain certain principles of the invention.

Figs. 1A-1C shows acceptance of an exemplary blood-free diet of the present invention, by mosquito for feeding. Mosquito feeding status on new dietary system. (Fig. 1A) unfed, (Fig. 1B) half fed and (Fig. 1C) full fed. Arrow shows abdominal bulging after feeding on the exemplary blood-free diet of the present invention.

Figure 2 shows fecundity of *Aedes albopictus* after feeding on various diets: Blood-guinea pig, BS (Bovine serum albumin (BSA) & sugar), BCS (BSA, sugar & cholesterol), BAS (BSA, ATP & sugar), and BACS (BSA, ATP, sugar & cholesterol).

Figure 3 shows survival of *Aedes albopictus* after feeding on various diets: Blood-guinea pig, BS (Bovine serum albumin (BSA) & sugar), BCS (BSA, sugar & cholesterol), BAS (BSA, ATP & sugar), and BACS (BSA, ATP, sugar & cholesterol).

Figure 4 shows fecundity of *Aedes albopictus* as a function of the amount of BSA in diet.

Figure 5 shows survival of *Aedes albopictus* as a function of the amount of BSA in diet.

Figure 6 shows fecundity of *Aedes albopictus* at lower concentration of various diets (Egg albo-Egg albumin, ATP-ATP, BSA-Bovine serum albumin).

Figure 7 shows fecundity of *Ae. albopictus* females fed on diets containing phagostimulant (Adenosine tri phosphate) and without phagostimulant against. (Blood-guinea
5 pig, BS-Bovine serum albumin & sugar, BAS-Bovine serum albumin, ATP)

Figure 8 shows fecundity of *Culex pipiens molestus* after feeding on blood-free diet (BS-Bovine serum albumin & sugar) against control (sugar solution).

Figs. 9A-9C shows an exemplary artificial feeding system, in accordance with various embodiments of the present invention. (Fig. 9A) angular view, (Fig. 9B) lateral view, and (Fig.
10 9C) top view. Abbreviations: IP – inner feeding platform, MP – middle feeding platform, OP – outer feeding platform.

Figure 10 shows efficacy of blood-free diets containing 10% (W/V) and 20% (W/V) BSA with 5% (W/V) sugar to support multiple generations of *Aedes albopictus* mosquitoes in comparison to live Guinea pig feeding.

Figure 11 shows preference of blood-free diets prepared with distilled water and a PBS buffer to females *Aedes albopictus*.

Figure 12 shows *Aedes albopictus* egg production efficacy of blood-free diets (BSA 10% (W/V) + Sugar 5% (W/V)) and (BSA 20% (W/V) + Sugar 5% (W/V)) prepared in distilled water and PBS buffer.

Figure 13 shows efficacy of blood-free diet as gross and net production of egg per *Aedes albopictus* female after 24 h, 48 h and 72 h feeding durations.

Figs. 14A – 14C shows a bi-choice experimental setup developed for the estimation of feeding preferences using UV dyes (blue and red). Fig. 14A shows diet feeders with flat base, a diet reservoir for holding diet and a feeding platform; Fig. 14B shows centrifuge tubes
25 comprising exemplary diets with blue and red dyes, illuminated under UV light; and Fig. 14C shows a cup holding hungry mosquitoes and two diet feeders one with an exemplary diet comprising a red dye and the other with a blue dye.

Figs. 15A -15B shows detection of mosquito preference for diet and its components marked with blue and red dyes using the bi-choice experimental setup of Fig. 14 setup. Fig. 15A

shows mosquito fed on diets with dyes under normal daylight and Fig. 15B shows similar mosquitoes under UV light (Left side – fed on blue dye diet, Center – fed on red dye diet, Right side – fed on both blue and red dye diets). Choice experiment was tested for biases using the reciprocal exchange of sugar and BSA with blue and red dyes.

5 Figure 16 shows results of a bi-choice experiment for preference of diets: sugar (5% (W/V)), BSA (10% (W/V) in water) and mix diet (where females were fed both sugar and BSA) using UV dye as in Fig. 14 & 15.

 Figure 17 shows results of a bi-choice experiment for preference of diets: sugar only diet (5% (W/V)), BSA and sugar diet (BSA (10% (W/V)) + sugar (5% (W/V))) in water, and mix diet
10 (where females were fed both sugar only and BSA+sugar diets) using UV dye as in Fig. 14 & 15.

 Figure 18 shows feeding preference of a blood-free diet (BSA (20% (W/V)) + sugar (5% (W/V))) in comparison to a live animal blood feeding to *Anopheles quadrimaculatus*.

 Figure 19 shows egg laying efficacy of *Anopheles quadrimaculatus* fed on blood-free diet (BSA 20% (W/V) + Sugar 5% (W/V)) and on a live animal for blood.

15 Figure 20 shows preference of blood-free diet and a live animal blood feeding by diapausing *Aedes albopictus* females under short day photoperiod (8h) and lower temperature (21°C) conditions.

 Figure 21 shows efficacy of blood-free diet and a live animal blood feeding on *Aedes albopictus* diapause egg production under short day photoperiod (8h) and lower temperature
20 (21°C).

 Figure 22 shows an exemplary artificial feeding system, in accordance with various embodiments of the present invention.

 Figure 22 shows a method of maintaining and growing hematophagus insects, in accordance with various embodiments of the present invention.

25 **DETAILED DESCRIPTION OF THE INVENTION**

Reference will now be made in detail to various exemplary embodiments of the invention, examples of which are illustrated in the accompanying drawings. The following description is provided to assist the reader in understanding various embodiments and features of

embodiments of the invention, and should not be considered as a limitation on the invention as disclosed herein and claimed below.

The present invention relies on a different strategy and is dependent on the sugar feeding behavior of mosquitoes thereby providing alternative pathways to mosquito host-seeking and blood feeding behaviors. Based primarily on sugar feeding behavior, the present artificial feeding system does not require a membrane feeder, phagostimulant, ATP or host chemical cues *i.e.* lactic acid to get the protein diet described previously (Hosoi 1959, Kogon 1990, Accar et al. 2012, Pitts 2014, Gonzales et al. 2014). The artificial feeding system of the present invention is independent of blood feeding rhythms as sugar feeding does not have any specific pattern.

As used herein unless context dictates otherwise, the term “blood-free diet” includes food compositions which preferably exclude whole blood or plasma, or serum fluids of a person or animal. The “blood-free diet” of the present disclosure may include isolated blood components, such as bovine serum albumin or hemoglobin for example. The blood-free diets of the present invention support egg production and colony maintenance of hematophagous insects. In some embodiments, the hematophagous insects are sugar feeding insects.

While the following discussion is in the context of feeding of mosquitoes, the formulations described herein, as well as the methods of determining appropriate formulations, are expected to be applicable to other hematophagous insects as well (e.g., tsetse flies, lice, bed bugs, no-see-ums, fleas, sand flies, midges, snipe flies, horse flies, stable flies, or sheep flies).

Artificial Feeding System:

In an aspect of the present disclosure, the artificial feeding system, for maintenance and growth of hematophagous insects, includes a blood-free diet formulation and a feeding device. The blood-free diet formulation comprises effective amounts of a protein source and a carbohydrate source. In such embodiments, the blood-free diet formulation is suitable for egg production and colony maintenance of hematophagous insects. The feeding system includes a diet reservoir for receiving and holding the blood-free diet formulation and a feeding platform for hematophagous insects to feed the blood-free diet formulation from the diet reservoir.

The hematophagous insects are mosquito colonies chosen from *Aedes*, *Anopheles*, *Culex* mosquitoes.

Feeding device:

Figs. 9A-9C shows an exemplary feeding device (120), according to various embodiments of the present invention. The feeding device (120), as shown in Figs. 9A-9C has two parts, a diet reservoir (130) for holding the blood-free diet (not shown) and a feeding platform (140) for hematophagous insects, such as mosquitoes to sit on and feed on the blood-free diet formulation contained in the diet reservoir (130). The diet reservoir (130) is designed to provide a maximum reach of mosquitoes to diet by providing a feeding platform (140) for mosquitoes to sit in the center and on the outside edge of the periphery. In some embodiment, the feeding platform comprises a plurality of feeding platforms e.g., 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more platforms. In other embodiment, the plurality of feeding platforms divide the diet reservoir into a plurality of diet reservoirs separated by the platforms. In such an embodiment, the size of the diet reservoir can be increased or reduced depending on mosquito colony size, such as by using only a portion of the plurality of diet reservoirs. For example, Fig. 9A shows an exemplary feeding device comprising two diet reservoirs and three feeding platforms: inner platform (IP), middle platform (MP) and outer platform (OP), where the two diet reservoirs are separated by the middle feeding platform. In some embodiments, the feeding device may further comprise an optional mosquito-proof covering (not shown) separating the diet reservoir from the platform. In an embodiment, the covering is a mesh made of nylon, cotton, polyester fabric or metal screen. The mosquito-proof covering can be attached to the diet reservoir using hot glue or super glue. Alternatively, the optional mosquito-proof covering can be screwed or friction fit to the diet reservoir. In some embodiment, the feeding device is free of mosquito-proof covering, and comprises only a feeding platform and a diet reservoir, on which mosquitoes can successfully feed.

Figure 22 shows another exemplary artificial feeding system (200) comprising a blood-free diet formulation (210) and a feeding device (220), in accordance with various embodiments of the present invention. The feeding device (220) comprises a diet reservoir (230) for receiving the blood-free diet formulation and a feeding platform (240) that floats over the blood-free diet formulation in the diet reservoir. In some embodiments, at least a portion of the feeding platform is hydrophobic. In some embodiments, the feeding platform is in the form of a mesh. The feeding platform in the form of a mesh that floats, provides several advantages such as not only that the mesh can be used as a feeding platform that never gets wet, but it provides the

hematophagous insects an uninterrupted and/or continuous access to the blood-free diet through the mesh as the amount of blood-free diet decreases in the diet reservoir.

The feeding device (120), (220) can have any suitable shape, including but not limited to circular or polygonal. In some embodiments, the diet reservoir is circular in shape. In another
5 embodiment, the diet reservoir is square, triangular or rectangular in shape. In some other
embodiments, the diet reservoir is annular in shape. The feeding device can have any suitable size
depending upon the scale of feeding needed and the mosquito colony size, such as from 200 μ L
to 50 mL. In some embodiments, the diet reservoir can hold 200 μ L -2 mL of diet or from 200-
1000 μ L; or from 1-2 mL of diet which is sufficient to feed approximately 50-100 mosquitoes.
10 In other embodiments, the diet reservoir can hold from 10 to 50 mL of diet. Yet, in other
embodiment, multiple feeding devices can be used in large scale production as alternative of
larger sized device.

The diet reservoir of the feeding device can be made of any suitable material that can be
formed to hold water, including but not limited to, polypropylene, polyethylene, polystyrene or
15 any other inert plastic material or a metal, such as aluminum foil, or biodegradable materials
such as wax paper and the like. The feeding platform can be made of any suitable materials,
including but not limited to, polystyrene, aluminum foil, wax paper, sponge, resin, wood and
cardboard coated with non-wetting agents. In some embodiments, the floating feeding platform
is coated with non-wetting agents such as wax or any other hydrophobic material, to keep it from
20 wetting, and thereby prevent feeding mosquito from getting stuck. In certain embodiments, the
feeding platform is light enough to float on the liquid blood-free diet, so that it stays on top of the
diet for mosquitoes to use as a feeding platform. The pore size of the mesh forming the feeding
platform can be from 1x1 mm to 3x3 mm or from 1-3 mm in diameter or from $\sim$ 1.5-2 mm in
diameter, so that mosquito can stand on the mesh without getting stuck. In some embodiments,
25 the pore size of the mesh forming the feeding platform is 2 mm in size, which can be diameter
for circular pore or a side or diagonal for a polygonal pore.

Blood-free diet formulation:

In accordance with various embodiments, the artificial feeding system of the present
invention comprises a blood-free diet formulation in addition to a feeding device as disclosed
30 hereinabove. The blood-free diet formulation of the present invention comprises effective

amounts of a protein source and a carbohydrate source dispersed in a liquid, such as water (pH 7), 1 mM Phosphate Buffered Saline (PBS) (pH 7.2) or other appropriate physiological carriers suitable for egg production and colony maintenance of hematophagous insects.

The protein source present in the blood-free diet formulation can be selected from the group consisting of animal protein, milk protein, plant protein, insect protein, arthropod protein, and amino acids. In one embodiment, the protein source includes, bovine serum albumin, egg albumin, or a mixture thereof. Suitable carbohydrate sources include, but are not limited to glucose, sucrose, fructose, or mixtures thereof. The protein source can be present in an amount of 5-30%(W/V) or 10-20% (W/V) and the carbohydrate source can be present in an amount of 3-10% (W/V) or 4-8% (W/V). In a preferred embodiment, the blood-free diet formulation comprises glucose present in an amount of 5% (W/V). In another embodiment, the blood-free diet formulation comprises bovine serum albumin present in an amount of 10% (W/V).

In some embodiments, the blood-free diet formulation further comprises a lipid source, such as, for example, cholesterol. In another aspect, the blood-free diet formulation is free of a phagostimulant other than sugar. In certain embodiments, the blood-free diet formulation further comprises a traceable reagent, such as a UV dye.

Suitable examples of blood-free diet formulation, for egg development and survival of mosquitoes, include, but are not limited to:

Diet formulations:

- 20 Diet 1: Bovine Serum albumin (5 – 20%) (W/V)
 Sugar (5 – 10 %) (W/V)
 Dissolved in 1mM Phosphate Buffered Saline (pH 7.2)
- Diet 2: Bovine Serum albumin (5 – 10%) (W/V)
 Egg albumin (5 – 10%) (W/V)
 25 Sugar (5 – 10%) (W/V)
 Dissolved in 1mM Phosphate Buffered Saline (pH 7.2)
- Diet 3: Egg albumin (5 – 10%) (W/V)
 Sugar (5 – 10%) (W/V)
 Dissolved in 1mM Phosphate Buffered Saline (pH 7.2)

Diet 4: Bovine Serum albumin (5 – 20%) (W/V)

Sugar (5 – 10%) (W/V)

Cholesterol (1 – 2 mg/ml)

Dissolved in 1mM Phosphate Buffered Saline (pH 7.2)

5 Diet 5: Bovine Serum albumin (5 – 20%) (W/V)

Sugar (5 – 10%) (W/V)

Dissolved in Distilled water (pH 7)

As used herein the amounts shown in % (W/V) refer to the amount of solid component in
10 g dissolved in mL of liquid. In each formulation above, a range of 5-20% (W/V) means a range
between 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 and 20% (W/V). A range between 5-
10% (W/V) refers to any of 5, 6, 7, 8, 9, and 10% (W/V). Notably, additional components may
be added to the formulations above. These include, for example, animal proteins, milk proteins,
plant proteins, proteins obtained from insects or arthropods, amino acids and lipids.

15 In certain embodiments, the diet reservoir holding the diet composition can be placed in a
warming device having a temperature range between 30 to 35°C. The present inventors have
observed that warming the diet increases the success rate for the blood-free diet with respect to
egg laying and survival.

A method of maintaining and growing hematophagus insects is also provided and is
20 shown in Figure 23. The method comprises providing an artificial feeding system as disclosed
hereinabove; adding water and shaking it well to dissolve the blood-free diet composition in
water, followed by feeding the hematophagus insects. The feeding devices of the present
invention provide an easy way to both prepare diet and feed mosquitoes. Furthermore, the
feeding devices can have a variety of shapes and sizes, depending upon the colony size.

25 The device and diet composition of the present invention promotes both survival and egg
laying. The blood-free diet formulations, as disclosed hereinabove are suitable for production of
multiple generations of mosquitoes. The blood-free diet formulations of the present disclosure
are also suitable for rearing mosquitoes under diapausing conditions and production of viable
diapause eggs. The unique feeding mechanism utilizing sugar feeding behavior for mosquito egg
30 development renders the system independent of mosquito blood feeding schedules. Moreover,
the completely synthetic diet formulation is suitable for growth of a variety of different mosquito

species. Furthermore, the system does not require any electrical power for operation and can be used within a broad temperature range (25°C to 35°C). Finally, the blood-free diet formulation of the present disclosure can be stored for longer periods of time than blood under refrigeration.

We have developed a simple blood-free diet and feeding system offering the following
5 advances: (1) blood-free, (2) membrane-free, (3) ATP free, (4) room temperature feeding, (5) long-term storage stability even under field conditions, (6) inexpensive, (7) ease-of-handling, and (8) easy to scale-up. The key to our diet is the substitution of sugar for ATP to induce engorgement. There are two modalities in mosquito feeding: sugar and protein. Protein derived from blood supports egg production, whereas sugar derived from floral nectars supports
10 longevity. Sugar is also an extremely potent mosquito phagostimulant. The sugar in the artificial diet described herein serves to ‘fool’ the mosquito into concurrently imbibing the protein we provide. That is, the mosquito assumes a floral nectar rather than a host feeding mode, so neither a membrane nor heat to mimic a vertebrate host is needed. Thus, phagostimulants other than sugar and host attractants are not required for this maintenance free and reusable device. As
15 mentioned the system is non-species specific for mosquitoes and provides comparable fecundity and fertility similar to mosquitoes grown on a blood-based diet. Finally, the system supports both male and female development and survival.

The following examples describe illustrative methods of practicing the instant invention and are not intended to limit the scope of the invention in any way.

20

EXAMPLES

Materials Used

Bovine serum albumin (BSA) Lyophilized powder, Bio Reagent, suitable for cell culture, $\geq 96\%$
25 and albumin from chicken egg white powder, 62-88% (agarose gel electrophoresis) were obtained from Sigma Aldrich, St. Louis, MO.

General Method

A blood-free diet was prepared by dissolving a protein source (2-4 g) and a carbohydrate
30 source (1.0 g) in 20 mL amount of water (pH 7.0) or 1 mM Phosphate Buffer Saline solution

(PBS) (pH 7.2) by first mixing the components and shaking the mixture well. The resultant blood-free diet was ready to use once the foam developed during shaking dissipates.

A diet feeding system as shown in Fig. 9 was used in the following examples. Each experiment was conducted with a cohort of 10 or 15 females with three replicates, unless otherwise mentioned. Three types of mosquitos were used: *Aedes albopictus* females; *Culex pipiens molestus* females; and *Anopheles quadrimaculatus* females. The mosquitoes were provided with diet for 24 h and the diet was replenished periodically and observed for feeding at an interval of 30 minutes, unless otherwise mentioned

All the experiments on blood-free diet for attraction, feeding or oviposition were carried out under room temperature ($25^{\circ}\text{C}\pm 1$) and 16h light and 8h dark photoperiod except diapause egg experiments unless otherwise mentioned.

Experiment 1: Acceptance of proposed dietary system by mosquito for feeding:

Aedes albopictus females (5-7 days old) were put on fasting (by removing sugar food) for 24 h prior to acceptability test, as shown in Fig. 1A (unfed). Females were provided with exemplary artificial blood-free diet comprising BSA 10% (W/V) and Sugars 5% (W/V) in PBS for 2 hours and observed for feeding at an interval of 30 minutes. The experiment was conducted in a cohort of 10 females with three replicates. After feeding Females exhibited swollen abdomens. Fig. 1B shows a half fed mosquito and Fig. 1C) shows a full fed mosquito. All the females fed on exemplary blood-free diets. Arrow shows abdominal bulging after feeding on new dietary system.

Effect of Various Diet Formulations on Fecundity and Survival Rates of *Aedes albopictus* females

Experiment 2: Fecundity of *Aedes albopictus* females fed on various diet formulations.

Mosquitoes were fed on different diets containing bovine serum albumin, cholesterol, adenosine triphosphate, sugar and their combinations, as shown in Table 1 and on Guinea pig for blood feeding. Each treatment contains a group of 10 females per replicate and experiment was repeated three times.

Table 1

Sample		Composition
Control 1	Blood	blood fed on Guinea pig
Ex 2.1	BS	20% (W/V) BSA + 5% (W/V) Sugar in PBS
Ex. 2.1	BCS	20% (W/V) BSA + 0.2% (W/V) Cholesterol + 5% (W/V) Sugar in PBS
C.E. A.1	BAS	20% (W/V) BSA + 100 mM ATP + 5% (W/V) Sugar in PBS
C.E. A.2	BACS	20% (W/V) BSA + 100 mM ATP + 0.2% (W/V) Cholesterol + 5% (W/V) Sugar in PBS

Figure 2 shows that the exemplary diets, Ex. 2.1 (BS) and 2.2 (BCS), showed similar fecundity to blood feeding on the live animal (control) and other comparative diets, C.E. A.1 (BAS) and A.2 (BACS)) comprising phagostimulant ATP (Figure 2).

Experiment 3: Survival rate of females *Aedes albopictus* on various diet formulations:

Aedes albopictus females (5-7 days old) in a cohort of 15 were fed on different diet formulations, given in Table 1, after 24 h fasting. For control, fifteen females were blood fed on Guinea pig. Female survival recorded twice a week.

Figure 3 shows that the exemplary diet, Ex. 2.1 (BS) showed similar survival to blood feeding on the live animal (control 1) and other comparative diets, C.E. A.1 (BAS) and A.2 (BACS)) comprising phagostimulant ATP (Figure 2). However, the exemplary diet Ex. 2.2 (BCS) comprising cholesterol as lipid showed unexpectedly initially high mortality.

Effect of amount of protein source on Fecundity and Survival Rates of *Aedes albopictus* females

Exemplary diets comprising 2 g (10% (W/V)) or 4 g (20% (W/V)) of bovine serum albumin (BSA) as a protein source and 1 g of glucose as a sugar source, dissolved in 20 mL of PBS, were fed to 15 *Ae. albopictus* females for 24 hours.

Experiment 4: Figure 4 shows that there was no significant difference between the exemplary diets containing 10% (W/V) and 20% (W/V) concentration of bovine serum albumin (BSA).

Experiment 5: Figure 5 shows that there was no significant difference in adult survival between exemplary diets containing 10% (W/V) and 20% (W/V) concentrations of Bovine serum albumin.

5 **Effect of amount of protein source on Fecundity of *Aedes albopictus* females**

Experiment 6: Fecundity of *Aedes albopictus* females fed on various diet formulations compositions.

Aedes albopictus females (25 to 30) were fed on different diets, shown in Table 2 for 3 days after 24 h fasting. Individual females were kept into a 50 mL vial with water and oviposition paper for egg laying. After 4 days of oviposition, eggs were counted.

Table 2 summarizes various exemplary diets used:

Sample	Composition
Control 2	blood fed on Guinea pig
Ex 3.1	10% (W/V) Egg Albumin + 5% (W/V) Sugar in water
Ex. 3.2	10% (W/V) Egg Albumin + 100 mM ATP + 5% (W/V) Sugar in water
Ex. 3.3	5% (W/V) Egg Albumin + 5% (W/V) Sugar in water
Ex. 3.4	5% (W/V) Egg Albumin + 5% (W/V) BSA + 5% (W/V) Sugar in water
Ex. 3.5	5% (W/V) BSA + 5% (W/V) Sugar in water
Ex. 3.6	10% (W/V) BSA + 5% (W/V) Sugar in water

Figure 6 shows that there was no significant difference among blood, BSA (10% (W/V)) and egg albumin (5% (W/V)) + BSA (5% (W/V)). It should be noted that exemplary diet Ex. 3.1 comprising 10% (W/V) egg albumin performed better than similar diet, Ex. 3.2 comprising ATP. This indicates that protein source used in this diet (BSA) can be replaced by other animal proteins such as egg albumin, which provides added benefits of cost effectiveness and less stringent storage condition requirement. Furthermore, it also indicates that the sugar phagostimulant can also work with other protein sources and with mixtures of other protein sources and BSA.

Effect of Adenosine triphosphate (ATP) as phagostimulant in the Exemplary Diets on Fecundity of *Aedes albopictus* females

Experiment 7: Blood-free blood-free diet is effective in the absence of phagostimulant (Adenosine triphosphate (ATP)): Two diet formulations were prepared with and without ATP, summarized below in Table 3. We compared both diets with blood.

Table 3 summarizes various exemplary diets used:

Sample	Composition
Control 3	blood fed on Guinea pig
Ex 4.1	10% (W/V) BSA + 5% (W/V) Sugar in water
Ex. 4.2	10% (W/V) BSA + 100 mM ATP + 5% (W/V) Sugar in water

Figure 7 shows that blood-free diet Ex. 4.1 comprising BSA and sugar only works similar to blood control and another exemplary diet, Ex. 4.2 comprising ATP in addition to BSA and sugar, indicating that inclusion of ATP as a phagostimulant is not required in the diet formulations of the present invention.

Effect of exemplary diet on the Fecundity of *Culex pipiens molestus* against control (sugar solution)

Experiment 8: Fecundity of *Culex pipiens molestus* females fed on a blood-free diet containing BSA (20% (W/V)) & Sugar (5% (W/V)) in water. This mosquito is an autogenous, in that the first batch of eggs is laid without blood feeding. Females which have already laid first batch eggs were fed on blood-free diet and observed for oviposition. Twenty females were fed on diet formulation and observed for egg laying. Notably, eggs were also laid with blood-free diet. The control group did not lay eggs which show this species can be maintained for multiple gonotrophic cycles (Figure 8).

Efficacy of exemplary diets containing 10% (W/V) and 20% (W/V) BSA with 5% (W/V) sugar to support multiple generations of *Aedes albopictus* mosquitoes in comparison to live Guinea pig feeding

Experiment 9: Efficacy of blood-free diets containing 10% (W/V) and 20% (W/V) BSA with 5% (W/V) sugar to support multiple generations of *Aedes albopictus* mosquitoes in comparison to live Guinea pig feeding.

Initially, 50 females were taken in three groups (i.e. Blood, BSA (10% (W/V)) and BSA (20% (W/V))). One group was blood fed on Guinea pig, and the rest of the two groups were fed on exemplary diets having BSA (10% (W/V)) and 5% sugar or BSA (20% (W/V)) and 5% sugar, respectively. After feeding, fed and unfed females were sorted out and fed females were transferred to oviposition vial individually and observed for egg laying. Sugar solution (5% (W/V)) in cotton wicks was provided for survival. After 1 week of diet feeding, the females were observed for oviposition and number of eggs were counted for individual females.

Later, eggs were kept for hatching in 400 mL plastic cups and percent egg hatching rate was calculated. Hatched larvae were reared as described in Gaugler et al 2012. Brewer's yeast was provided as larval food. After pupation, pupae were transferred to 1 cu. feet mosquito cages for adult emergence and number of successful adult emerged were counted. For subsequent generations experiment, 50 females were taken and fed on blood-free dietary system and further assessed for oviposition, egg laying and larval development as described above for 10 generations. To compare the efficacy of the exemplary blood-free diets of the present invention, a control sample of blood feeding mosquitoes was taken from standard laboratory colony in each generation and were fed on Guinea pig.

Figure 10 shows that exemplary blood-free diets can support mosquito egg production for multiple generations in the lab colony and no differences were observed at the 10th generation with respect to egg production per female among various diets.

Preference of exemplary blood-free diets prepared with distilled water or a PBS buffer to females *Aedes albopictus*.

Experiment 10: Figure 11 shows that exemplary blood-free diet prepared with BSA (10% (W/V)) in distilled water was consumed by more *Aedes albopictus* females in comparison to the

diet prepared with BSA (10% (W/V)) in PBS buffer. This indicates that the exemplary blood-free diets of the present invention do not need PBS buffer and can be simplified further.

Effect of exemplary blood-free diets prepared with distilled water or a PBS buffer on fecundity of *Aedes albopictus* females

Experiment 11: Blood-free diet prepared in distilled water (BSA 10% (W/V) + Sugar 5% (W/V) and BSA 20% (W/V) + Sugar 5% (W/V)) produced more net and gross egg per *Aedes albopictus* female in comparison to diet prepared in PBS buffer (BSA 10% (W/V) + Sugar 5% (W/V) and BSA 20% (W/V) + Sugar 5% (W/V)).

During oviposition, few females died after laying eggs and it was unclear if they died after complete or partial egg laying. Hence, gross data shows eggs from all the females (live and dead), whereas, net data shows the egg number where females were alive at the time of egg counting.

Figure 12 shows that that the presence of water of PBS in the exemplary blood-free diets does affect the fecundity of *Aedes albopictus* females. Hence, the exemplary blood-free diets of the present invention do not require PBS buffer. Thus, the use of water in the exemplary blood-free diets of the present invention instead of PBS further simplifies the diet composition, making it cost effective and readily applicable to field conditions.

Effect of exemplary blood-free diets prepared with distilled water on fecundity, gross and net production of egg per *Aedes albopictus* female after 24 h, 48 h and 72 h feeding durations.

Experiment 12: Efficacy of blood-free diet as gross and net production of egg per *Aedes albopictus* female after 24 h, 48 h and 72 h feeding durations.

Females were provided with blood-free diet (BSA (10% (W/V)) and sugar (5% (W/V))) for different time intervals i.e. 24 h, 48 h and 72 h. After the specified duration of time, mosquitoes were provided with 5% (W/V) sugar solution in cotton wick. After 4 days, all females were transferred to oviposition vial individually and observed for egg laying.

Figure 13 shows that exemplary blood-free diet can produce similar eggs with less feeding time (24 h) in comparison to longer feeding time (72 h), suggesting that extended

feeding time such as 48 or 72 hours does not increase the egg production, and therefore, 24 hour feeding time is sufficient.

A Bi-choice Experimental Set Up

Experiment 13: A Bi-choice Experimental Set Up

5 Figs. 14A-14C show an experimental setup developed for the estimation of feeding preferences using UV dyes (blue and red): Fig. 14A shows an artificial feeding system (1400) comprising feeding devices (1420) with flat base, a diet reservoir (1420) for holding diet (1410) and a feeding platform (1440) ; Fig. 14B shows centrifuge tubes comprising exemplary diets with blue and red dyes, illuminated under UV light; and Fig. 14C shows a cup holding hungry
10 mosquitoes and two diet feeders one with an exemplary diet comprising a red dye and the other with a blue dye. The red and blue dyes were mixed in the exemplary blood-free diets at 0.01% (W/V) concentrations.

Experiment 14: Figure 15 shows detection of mosquito preference for diet and its components
15 marked with blue and red dyes using experimental setup, as shown in Figs 14A-14B. Fig. 15A shows mosquito fed on diets with dyes under normal daylight and Fig. 15B shows similar mosquitoes under UV light (Left side – fed on blue dye diet, Center – fed on red dye diet, Right side (referred to as mix) – fed on both blue and red dye diets). Choice experiment was tested for biases using the reciprocal exchange of sugar and BSA with blue and red dyes. We found no
20 biases in the system for any component.

A Bi-choice experiment for preference of sugar (5% (W/V)) against BSA (10% (W/V)) by *Aedes albopictus* mosquitoes

Experiment 15: A Bi-choice experimental set as described hereinabove was used to determine
25 preference of sugar (5% (W/V) in water) against BSA (10% (W/V) in water). Figure 16 shows that BSA is less preferred in comparison to sugar. Fifteen percent mosquitoes choose both sugar and BSA.

A Bi-choice experiment for preference of sugar (5% (W/V)) against exemplary blood-free diet comprising BSA and sugar by *Aedes albopictus* mosquitos

Experiment 16: A Bi-choice experimental set as described hereinabove was used to determine preference of sugar (5% (W/V)) against an exemplary diet comprising BSA (10% (W/V)) +
 5 sugar (5% (W/V)) in water.

Figure 17 shows a surprising result that BSA exhibits an additive effect in combination with sugar, as demonstrated by an increased preference of BSA+sugar by mosquitoes in comparison to sugar alone. Ten percent mosquitoes chose both sugar and BSA+sugar.

10 **Preference and efficacy of exemplary blood-free diet on fecundity of *Anopheles quadrimaculatus***

Experiment 17. Experimental setup was exactly the same as described hereinabove. Twenty females were provided proposed diet (BSA 20% (W/V) and sugar 5% (W/V)) for 24 hours and twenty females were fed on Guinea pig as control. After feeding, females were individually
 15 transferred to 450 mL plastic cup having smaller cups filled with 30 mL water for egg laying. Females were provided with 5% (W/V) sugar solution. After 7 days, individual females were observed for egg laying and eggs were counted.

Figure 18 shows that exemplary diet comprising BSA 20% (W/V) and Sugar 5% (W/V) in water is similar in preference to feeding on blood by *Anopheles quadrimaculatus*. Figure 19
 20 shows egg production efficacy of the exemplary diet in comparison to blood.

Preference and efficacy of exemplary blood-free diet on fecundity of *Aedes albopictus* for diapause egg production in short day photoperiod (8 h)

Experiment 18. Efficacy of blood-free diet (BSA 20% (W/V) + Sugar 5% (W/V)) for the
 25 production of *Aedes albopictus* diapause eggs at short day photoperiod (8h) and lower temperature (21°C) to produce diapause eggs in comparison to live animal blood feeding. Figure 20 shows that exemplary diet was readily accepted by mosquitoes reared under diapausing condition. Figure 21 shows that females lay diapause eggs under the same conditions and the egg production efficacy of the exemplary diet was similar in comparison to blood.

30 Pitts (2014) is perhaps the nearest diet to the exemplary blood-free diet, as disclosed hereinabove.

Pitts (2014) diet comprises BSA (200 mg/mL) and ATP (1 mM) in PBS and in common with nearly all artificial mosquito diets, include bovine serum albumin (BSA) as a blood-replacement protein. As with virtually all artificial mosquito diets, Pitt's system requires ATP as a phagostimulant, as well as a feeding device with heating, and a membrane, all of which are
 5 costly and complex to operate. Moreover, an 'odor blend of ammonium hydroxide, lactic acid, isovaleric acid, geranyl acetone, and butylamine was applied to the membrane as a host-seeking stimulant. This group of chemicals are complex to prepare, adding another obstacle to scale up. Despite this complexity, Pitts obtained good egg production from their feeding system, yet suboptimal egg hatch (71.8%) in comparison to exemplary blood-free diet described herein
 10 comprising BSA (200 mg/mL) and sugar (50 mg/mL) in water (volume made up to 1 mL) (98% egg hatch).

It will be apparent to those skilled in the art that various modifications and variations can be made in the practice of the present invention and in construction of this system without departing from the scope or spirit of the invention. Other embodiments of the invention will be
 15 apparent to those skilled in the art from consideration of the specification and practice of the invention. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

REFERENCES

- 20 Accar EB, Deane GF, Johanson E, Mullen ER, Myhrvold NP, Peterson NR, Tegreene CT, Whitmer C, Wood LL. 2012. Food composition for hematophagous insects. US Patent, US 8, 133, 524B1
- Allan SA, Bernier UR, Kline DL. 2010. Laboratory evaluation of lactic acid on attraction of *Culex* spp. (Diptera: Culicidae). *J. Vect. Ecol.* 35, 318-324.
- 25 Gonzales KK, Tsujimoto H, Hansen TA. 2014. Blood serum and BSA, but neither red blood cells nor hemoglobin can support vitellogenesis and egg production in the dengue vector *Aedes aegypti*. *PeerJ.* 3:e938.
- Hosoi T. 1959. Identification of blood components which induce gorging of the mosquito. *Journal of Insect Physiology.* 3, 191-218.
- 30 Kogan PH. 1990. Substitute blood meal for investigating and maintaining *Aedes aegypti* (Diptera: Culicidae). *J. Med. Entomol.* 27, 709-712.

Phasomkusolsil S, Tawong J, Monkanna N, Pantuwatana K, Damdangdee N, Khongtak W, Kertmanee Y, Evans BP, Schuster AL. 2013. Maintenance of mosquito vectors: effects of blood source on feeding, survival, fecundity, and egg hatching rates. *J. Vect. Ecol.* 38, 38-45.

Pitts RJ. 2014. A blood-free protein meal supporting oogenesis in the Asian tiger mosquito, *Aedes albopictus* (Skuse). *J. Insect Physiol.* 64, 1–6.

CLAIMS

What is claimed is:

1. An artificial feeding system for maintenance and growth of hematophagous insects comprising,

- 5 a) a diet reservoir for receiving a synthetic, blood-free diet formulation;
- b) at least one feeding platform; and
- c) an optional mosquito proof covering separating said diet reservoir from said platform, said system and diet formulation being suitable for egg production and colony maintenance, wherein said system lacks a membrane.

10 2. The system of claim 1 comprising multiple feeding platforms.

3. The system of claim 1 or 2, wherein said diet reservoir is 5mm or less in depth.

4. The system of any one of the preceding claims, wherein said covering is present and is a mesh having a pore size of about 2mm.

15 5. The system of any one of the preceding claims, wherein said diet reservoir is circular, square or triangular in shape.

6. The system of any one of the preceding claims, comprising said synthetic blood free diet formulation.

7. The system of claim 6, wherein said synthetic blood free diet formulation comprises a traceable reagent.

20 8. The system of claim 7, wherein said traceable reagent is a UV dye.

9. The system of claim 1, suitable for production of multiple generations of mosquito.

10. The system of claim 1 suitable for rearing mosquitoes under diapausing conditions and production of viable diapause eggs.

11. A synthetic blood-free diet formulation for maintenance of hematophagous insects comprising effective amounts of a protein source, a carbohydrate source and, optionally a lipid source, said diet being suitable for egg production and colony maintenance.

12. The diet formulation of claim 11, wherein said protein source is selected from the group consisting of animal protein, milk protein, plant protein, insect protein, arthropod protein, and amino acids.

13. The diet formulation of claim 11, wherein said protein source is bovine serum albumin and/or egg albumin.

14. The diet formulation of any one of the preceding claims 11-13, wherein said carbohydrate source comprises at least one of glucose, sucrose, fructose, or mixtures thereof.

15. The diet formulation of any one of the preceding claims 11-14, wherein said lipid is cholesterol.

16. The diet formulation of any one of the preceding claims 11-15, wherein said synthetic blood-free diet formulation lacks an ATP phagostimulant.

17. The diet formulation of any one of the preceding claims, for supporting a mosquito colony from Aedes, Anopheles and Culex mosquitoes.

18. The diet formulation of any one of the preceding claims, wherein the protein source is present in an amount of 5–30% (W/V) and the carbohydrate source is present in an amount of 3–10% (W/V).

19. The diet formulation of any one of the preceding claims 11-18, selected from the group consisting of:

Diet 1: Bovine Serum albumin (5 – 20%) (W/V)

Sugar (5 – 10 %) (W/V)

Dissolved in 1mM Phosphate Buffered Saline (pH 7.2);

Diet 2: Bovine Serum albumin (5 – 10%) (W/V)

Egg albumin (5 – 10%) (W/V)

Sugar (5 – 10%) (W/V)
dissolved in 1mM Phosphate Buffered Saline (pH 7.2);

Diet 3: Egg albumin (5 – 10%) (W/V)

5 Sugar (5 – 10%) (W/V)
 dissolved in 1mM Phosphate Buffered Saline (pH 7.2);

Diet 4: Bovine Serum albumin (5 – 20%) (W/V);

10 Sugar (5 – 10%) (W/V)
 Cholesterol (1 – 2 mg/ml)
 dissolved in 1mM Phosphate Buffered Saline (pH 7.2); and

Diet 5: Bovine Serum albumin (5 – 20%) (W/V)

15 Sugar (5 – 10 %) (W/V)
 dissolved in Distilled water.

21. The feeding system of claim 1, wherein at least a portion of the feeding platform is hydrophobic and floats.

20 22. The feeding system of claim 21, wherein the feeding platform is formed from a mesh.

23. An artificial feeding system for maintenance and growth of hematophagous insects as claimed in claim 1, further comprising a blood-free diet formulation comprising effective amounts of a protein source and a carbohydrate source, wherein the blood-free diet formulation
25 is suitable for egg production and colony maintenance of hematophagous insects.

24. The system of claim 23, wherein at least a portion of the feeding platform is hydrophobic and floats.

25. The system of any one of the preceding claims 21-24, wherein the blood-free diet formulation comprises the effective amounts of a protein source and a carbohydrate source dispersed in a liquid.

26. A method of maintaining and growing hematophagus insects, comprising:

- 5 (a) providing an artificial feeding system of claim 1 or claim 25;
(b) dissolving the blood-free diet formulation in water; and
(c) feeding the hematophagus insects.




Fig. 1A




Fig. 1B




Fig. 1C




Figure 2

2/15




Figure 3




Figure 4




Figure 5




Figure 6

4/15




Figure 7




Figure 8




Fig 9A




Fig. 9B




Fig. 9C




Figure 10




Figure 11

7/15




Figure 12




Figure 13

8/15




Fig. 14A




Fig. 14B

9/15




Fig. 14C

10/15




Fig. 15A




Fig. 15B

11/15




Figure 16




Figure 17

12/15




Figure 18




Figure 19

13/15




Figure 20




Figure 21




Figure 22

15/15




Figure 23

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2017/050778

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A01K 67/033; A01K 29/00; A01K 67/00; A23K 50/00; A23K 50/90 (2017.01)

CPC - A01K 67/0337; A01K 29/00; A01K 67/00; A01K 67/033; A01K 67/0333; A23K 50/00; A23K 50/90 (2017.08)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2012/0148705 A1 (ACAR et al) 14 June 2012 (14.06.2012) entire document	11-14
Y		1, 2, 9, 23, 25
Y	US 2007/0169209 A1 (HOFFMAN et al) 19 July 2007 (19.07.2007) entire document	1, 2, 9, 23, 25
A	US 5,133,289 A (GEORGI) 28 July 1992 (28.07.1992) entire document	1-3, 9-14, 21-25

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 October 2017

Date of mailing of the international search report

14 NOV 2017

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2017/050778

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-8, 15-19, 26
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.