

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

(10) **DK/EP 3540429 T3**

-
- (51) Int.Cl.: **G 01 N 33/00 (2006.01)** **G 01 N 33/84 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2021-02-15**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-11-11**
- (86) Europæisk ansøgning nr.: **19150428.1**
- (86) Europæisk indleveringsdag: **2014-09-02**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-09-18**
- (30) Prioritet: **2013-08-30 US 201361872149 P**
- (62) Stamansøgningsnr: **14841207.5**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **University of Maryland, College Park, Office of Technology Commercialization , 2130 Mitchell Bldg., College Park, MD 20742, USA**
Children's National Medical Center, 111 Michigan Avenue N.W., Washington, DC 20010, USA
The U.S.A. as represented by the Secretary, Department of Health and Human Services, Office of Technology Transfer , National Institutes of Health , 6011 Executive Boulevard, Suite 325 , MSC 7660, Bethesda, MD 20892-7660, USA
- (72) Opfinder: **AYYUP, Omar Bilal, 11429 Palatine Drive, Potomac, MD 20832, USA**
BEHRENS, Adam Michael, 3608 Toddsbury Lane, Olney, MD 20832, USA
KOFINAS, Peter, 5809 Nicholson Lane Apt. 610, North Bethesda, MD 20852, USA
SUMMAR, Marshall Lynn, 3131 Connecticut Avenue N.W. 2410, Washington, DC 20008, USA
CABRERA-LUQUE, Juan Manuel, 6400 Montrose Road, Rockville, MD 20852, USA
CUNNINGHAM, Gary, 3900 14th Street N.W. 201, Washington, DC 20011, USA
SIMEONOV, Anton, 7805 Green Twig Road, Bethesda, MD 20817, USA
MARUGAN, Juan, 16500 Apache Lane, Gaithersburg, MD 20878, USA
- (74) Fuldmægtig i Danmark: **Patrade A/S, Ceresbyen 75, 8000 Århus C, Danmark**
- (54) Benævnelse: **INDRETNING OG FREMGANGSMÅDER TIL ANVENDELSE AF INDRETNING TIL DETEKTION AF HYPERAMMONIÆMI**
- (56) Fremdragne publikationer:
MARTINA BAUMGARTNER ET AL: "Evaluation of flow injection analysis for determination of urea in sheep's and cow's milk", ACTA VETERINARIA HUNGARICA., vol. 50, no. 3, 1 July 2002 (2002-07-01), pages 263-271, XP055354034, HU ISSN: 0236-6290, DOI: 10.1556/AVet.50.2002.3.2
AYYUB OMAR B ET AL: "Simple and inexpensive quantification of ammonia in whole blood", MOLECULAR GENETICS AND METABOLISM, vol. 115, no. 2, 2015, pages 95-100, XP029130368, ISSN: 1096-7192, DOI: 10.1016/J.YMGME.2015.04.004
EGGENSTEIN C ET AL: "A DISPOSABLE BIOSENSOR FOR UREA DETERMINATION IN BLOOD BASED ON AN AMMONIUM-SENSITIVE TRANSDUCER", BIOSENSORS AND BIOELECTRONICS, ELSEVIER SCIENCE LTD. UK,

Fortsættes ...

**AMSTERDAM, NL, vol. 14, no. 1, 1 January 1999 (1999-01-01), pages 33-41, XP001020406, ISSN: 0956-5663, DOI:
10.1016/S0956-5663(98)00103-1**

DESCRIPTION

GOVERNMENT SUPPORT

[0001] This disclosure was made jointly by the NIH and with government support under HHSN268201200360P awarded by the NIH. The United States government has certain rights in the disclosure.

FIELD OF THE DISCLOSURE

[0002] The disclosure relates generally to devices that quantify and identify the presence or absence of ammonia or ammonium ion in a sample of bodily fluid, water, or other environmental sample. In some embodiments, the disclosure relates to diagnosing a subject with an hyperammonemia by detecting the presence, absence, or quantity of ammonia or ammonium ion in a sample of bodily fluid. In some embodiments, the device is a test strip only requiring a sample of whole bodily fluid for detection and/or quantification of ammonia or ammonium ion.

BACKGROUND OF THE DISCLOSURE

[0003] Elevated ammonia levels, oftentimes called hyperammonemia, is a potentially fatal symptom associated with a variety of diseases such as cirrhosis of the liver and urea cycle disorders found in neonatal infants. Left untreated, hyperammonemia can lead to cognitive developmental issues, seizures, other neurological problems, and death. The current testing methods include fluorometry and tandem mass spectroscopy performed by central laboratories, which could take multiple days to produce a reliable diagnosis. These methods involve large, cumbersome, and expensive machinery, which prevents testing of ammonia levels at the bedside or home once the disorder has been identified. Therefore, a system for a point of care testing device may be desired, as this may allow administration of treatment to occur more rapidly, in turn improving the neurological development of infants as well as making cirrhosis more manageable. Devices able to test for hyperammonemia may also be modified inexpensively to detect amino acid levels for applications in diagnosing and treating aminoacidopathies and other diseases.

[0004] Eggenstein C, et al. A disposable biosensor for urea determination in blood based on an ammonium-sensitive transducer. *Biosensors & Bioelectronics*. 1999 Jan;14(1):33-41 discloses a disposable biosensor for urea determination in blood based on an ammonium-sensitive transducer, using double matrix membrane technology.

SUMMARY OF THE INVENTION

[0005] The subject matter for which protection is sought is as defined in the claims.

[0006] The present invention generally relates to a test strip comprising:

a first and second vessel;

a fluid exchange opening positioned between the first and second vessel;

a conduit in fluid communication with the first vessel, the conduit configured to receive a fluid from a point external to the biosensor; and

a membrane comprising perfluorinated ionomers positioned at the fluid exchange opening;

wherein the membrane is capable of selectively transporting ammonia; and

wherein the second vessel comprises a hypohalite, an alkali buffer, and a phenolic compound suitable for performing a Berthelot reaction.

[0007] The present invention also provides the test strip of the invention, wherein the test strip is adapted for a portable device comprising a voltmeter and/or amperometer and a digital display such that, when the test strip is contacted to the device, the first and second electrodes become operably connected to a closed electrical circuit comprising the voltmeter and/or amperometer and the digital display, and, upon contact with a sample of bodily fluid, the at least one metabolic enzyme or functional fragment thereof catalyzes oxidation of an amino acid resulting in a current on the first electrode corresponding to a concentration value of amino acid in the sample of bodily fluid, such concentration value readable on the display of the portable device.

[0008] The present invention also provides the test strip of the invention, wherein the alkali buffer is used in a range from about 0.5 to about 1.0 mol/liter.

[0009] The present invention also provides the test strip of the invention, wherein the hypohalite is used in a range from about 50 to about 120 mmol/liter.

[0010] The present invention also provides the test strip of the invention, wherein the alkali buffer is one or a combination of: calcium acetate, calcium chloride, zinc acetate, zinc chloride, or any equivalent mono, di, or tri, -valent salt thereof.

[0011] The present invention also provides the test strip of the invention, wherein the phenolic reagent or indophenol reagent is used in a range from about 68 to about 70 mmol/liter

[0012] The present invention also provides the test strip of the invention, wherein the perfluorinated ionomer is Nafion.

[0013] The present invention also provides the test strip of the invention, wherein the perfluorinated ionomer has a thickness from about 0.001 mm to about 0.069 mm.

[0014] The present invention also provides the test strip of the invention, wherein the first vessel is no more than about 100 microliters in volume.

[0015] The present invention also provides a method of diagnosing a metabolic disease in a subject comprising:

1. (a) contacting a sample of bodily fluid to the test strip of the invention;
2. (b) quantifying one or more concentration values of ammonia in the sample;
3. (c) comparing the one or more concentration values of ammonia in the sample to a threshold value of ammonia concentration identified as being in a healthy range; and
4. (d) identifying the subject as having a metabolic disease if the one or more concentration values of ammonia in the sample exceed or fall below the threshold value.

[0016] The present invention also provides the method of diagnosing a metabolic disease in a subject wherein the metabolic disease is hyperammonemia.

[0017] The present invention also provides a method of determining patient responsiveness to a therapy comprising:

1. (a) contacting a sample of bodily fluid to the to the test strip of the invention;
2. (b) quantifying one or more ammonia or ammonium ion concentration values;
3. (c) comparing the one or more concentration values to one or more threshold values associated with a

metabolic disease.

[0018] The present invention also provides a method of determining patient responsiveness to a therapy, wherein the step of contacting comprises exposing the sample to the test strip for a sufficient period of time to allow modification of the indophenol reagent or indophenol related compound to become modified into a indophenol or indophenol reaction product.

[0019] The present invention also provides a method of determining patient responsiveness to a therapy, wherein the method does not comprise exposing the sample of bodily fluid to any external stimuli or reagent prior to contacting the sample to the at least one electrically conductive supports.

[0020] The present invention also provides a method of determining patient responsiveness to a therapy comprising, wherein the sample of bodily fluid is whole blood from a subject.

[0021] The present disclosure encompasses the recognition that hyperammonemia can be identified and/or characterized by identifying the levels or quantities of ammonia or ammonium ion in any sample, including a bodily fluid including human and non-human whole blood samples. In some embodiments, the present disclosure relates to identifying the quantity, presence, or absence of ammonia or ammonium ion in bodily fluids by contacting a bodily fluid to a device disclosed herein. In some embodiments, the methods disclosed herein do not comprise contacting the bodily fluid with any reagent or external stimuli prior to identifying or quantifying whether or how much one or more ammonia or ammonium ion are present in the bodily fluid.

SUMMARY OF OTHER ASPECTS OF THE DISCLOSURE

[0022] According to at least one exemplary aspect of the disclosure, a system, method, and apparatus for point of care hyperammonemia sensors may be disclosed. The system may utilize a phenol, 2-phenylphenol, ninhydrin, potassium tetraiodomercurate(II), nitroprusside, sodium hydroxide, similar reagents, catalysts, and buffers, or a combination thereof. The system may also utilize hyohalite, chloramine T, bleach, or similar chemical. Oftentimes called Berthelot's Reaction or an indophenol reaction, this reaction may determine ammonia levels in various mediums by changing color upon detection ammonia concentration. This may be useful for medical systems, such as in diagnosing hyperammonemia and various aminoacidopathies; for civil engineering systems, such as in determining ammonia levels of wastewater treatment plants; or for home based systems, such as ammonia detection in aquariums or pipes.

[0023] According to at least one exemplary aspect of the disclosure, an apparatus for point of care hyperammonemia sensors may be disclosed. The apparatus used may have a concavity, a fossa, or any other type of well as desired for the placement of the reagents and sample to be tested. Separating the sample and reagents may be a cation exchange membrane filter, such as Nafion or similar perfluorinated ionomers, to allow the passage of ammonia between the two sections of the well. Anion exchange membranes may also be used, as well as various polymeric hydrogels such as acrylamide, poly(ethylene glycol) diacrylate, poly(2-hydroxyethyl methacrylate), or poly(vinyl alcohol). Additionally, other exemplary aspects of the disclosure may include mechanisms for quantitative analysis of the color change by means of photodiodes and sensors or microfluidic devices that require smaller amounts of reagent and samples.

[0024] The present disclosure relates to a biosensor capable of measuring the total concentration of ammonia or ammonium ion in a sample with the use of a system comprising reagents for an indophenol or Berthelot reaction, such as hypochlorite, phenylphenol, a basic aqueous solution such as NaOH, and an alkali such as sodium acetate. In some aspects of the disclosure, the sensor or system comprises at least a first vessel comprising a basic buffer in aqueous or dried phase. In some aspects of the disclosure, comprises at least a first vessel comprises a gel or hydrogel that comprises at least one or a combination of: an indophenol reactant or reactants in dried or aqueous

phase, a basic buffer in aqueous or dried phase, an alkali solution in aqueous or dried phase, and/or an enzyme that oxidizes at least one amino acid substrate. The disclosure provides an ammonia or ammonium ion biosensor for measuring the total concentration of ammonia or ammonium ion. In some aspects of the disclosure, the detection or quantification of ammonia or ammonium ion is accomplished through colorimetric analysis whereby the reaction products of ammonia or ammonium ion are capable of emitting a wavelength in the visible spectrum of light. In some aspects of the disclosure, the system and/or biosensor comprises a diode configured to emit light in at least one vessel and a spectrophotometer configured to receive light emitted in a vessel containing indophenol or Berthelot reaction reactant products.

[0025] In some aspects of the disclosure, the system and/or biosensor also detects the absence, presence or quantity of amino acids in solution. In some aspects of the disclosure, the system and/or biosensor comprises at least a first electrically conductive surface (for measuring) and at least a second electrically conductive surface (counter electrode), wherein the first electrically conductive surface having one or more indophenol reaction reagents described herein or a combination of any one or more indophenol reaction reagents described herein and any one or combination of constituent factors, mediators, one or a plurality of enzymes, wherein, if the device comprise one or more enzymes, the one or more enzymes selectively utilize one or more amino acids as substrates. In those aspects of the disclosure with at least a first or second electrode, the one or plurality of enzymes produce reaction products by reacting with the specified amino acids as substrates, wherein the mediators transport electrons between the reaction products and the electrode measures amino acid concentrations, and wherein applied voltages at measuring between the first and second electrically conductive surfaces include such an applied voltage that, on a working curve representing the relationship between current value and applied voltage with respect to each of the one or plurality of specified amino acids, the distribution of current value at unchanged applied voltage as to individual amino acids. In other aspects of the disclosure in which the device, system, and/or biosensor comprise at least a first and/or second electrode, the first and/or second electrodes are positioned in, substantially adjacent to, or adjacent to at least one vessel in which an indophenol reagent described herein may react with one or more components of the reagents. In some aspects of the disclosure, ammonia or ammonium ion may be the reaction product of one of the enzymatic reactions in which the indophenol reaction, using a phenol or phenol related compound, can take place

[0026] According to at least one exemplary aspect of the disclosure, an apparatus, device, and/or system for point of care hyperammonemia sensors is disclosed. The apparatus comprises at least a first vessel, or a concavity, a fossa, or any other type of well as desired for the placement of the reagents and sample to be tested. The first vessel may be bifurcated by a membrane disclosed herein or the first vessel may be immediately adjacent to a second vessel in fluid communication with the second vessel via a fluid exchange opening. In some aspects of the disclosure, a membrane is positioned at the fluid exchange opening. In some aspects of the disclosure, the membrane is capable of transporting ions from the first vessel to the second vessel or vice versa. In some aspects of the disclosure, the membrane is a cation exchange membrane filter, such as Nafion® or similar membrane comprising perfluorinated ionomers. The membrane allows the passage of ammonia between the two vessels or between the two bifurcated sections of the at least first vessel. Anion exchange membranes may also be used, as well as various polymeric hydrogels such as acrylamide, poly(ethylene glycol) diacrylate, poly(2-hydroxyethylmethacrylate), or poly(vinyl alcohol).

[0027] Other exemplary aspects of the disclosure may include methods and mechanisms for quantitative analysis of ammonia or ammonium ion concentration in a sample by contacting a sample to a vessel comprising at least one indophenol reagent and/or a basic buffer, in either a solid or liquid phase, a section of the vessel exposed to at least a portion of a membrane disclosed herein. In some aspects of the disclosure, the method comprises detecting or quantitating the intensity of a color change within at least the first or second vessel before and after addition of a sample to the vessel or vessel. In some aspects of the disclosure, the method comprises contacting sample to at least a first vessel, a section or portion exposed or covered by at least one membrane disclosed herein, such first vessel also optionally comprising at least one indophenol reagent disclosed herein and/or a basic buffer, either in solid or liquid phase. In some aspects of the disclosure, if the at least first vessel comprises a buffer, the buffer may be an alkali solution such as sodium acetate or calcium acetate. In some aspects the disclosure relates to a method of contacting a sample to the device, biosensor or system disclosed herein

comprising at least a first and second vessel, said method comprises contacting or exposing the sample to the basic buffer in the at least first vessel, allowing ammonia from the sample to transfer to the second vessel comprising the indophenol reagents disclosed herein. In some aspects of the disclosure aspects the disclosure relates to a method of contacting a sample to the device, biosensor or system disclosed herein comprising at least a first and second vessel, said method comprises contacting or exposing the sample to the alkali solution in the at least first vessel, allowing ammonia from the sample to transfer to the second vessel comprising the indophenol reagents disclosed herein, the second vessel comprising one or a plurality of indophenol reactants, which after coming in contact with the ammonia produce an indophenol or indophenol related compound, In some aspects of the disclosure, the contents of the second vessel are exposed to light measure absorbance of light by indophenol compound or indophenol related compound at specific visible wavelengths of light, the absorbance is indicative of or proportionate to a quantity of ammonia or ammonium ion in the sample and whose absorbance is detected by an individual performing the test or by a device that measures wavelengths which is incorporated in the device, biosensor, or system disclosed herein. In some aspects of the disclosure, the method comprises comparing the absorbance of the color or wavelength to a standard which indicates the degree or severity of a hypereammonemia. In some aspects of the disclosure, the method comprises contacting a sample to a device, biosensor, or system disclosed comprising a diode, photodiode, and/or spectrophotometer or other device capable of measuring the absorbance of wavelength by the indophenol or indophenol related compounds produced as a product of an indophenol reaction within the device and exposed to a light. In some aspects of the disclosure, the device, biosensor, and/or system comprise a microfluidic circuit that comprises at least one conduit configured to receive the sample from a point external to the device, biosensor, and/or system, such microfluidic circuit comprises a conduit or series of conduits in fluid communication with at least the first and/or second vessel and the one or combination of: a spectrophotometer, diode, or other device capable of measuring the absorbance of specific wavelengths by the indophenol or indophenol related compound upon its exposure to light.

[0028] In some aspects of the disclosure, the disclosure relates to contacting or exposing a sample with an alkali buffer and/or a membrane disclosed herein within a vessel attached to an electrode able to measure the electron flow produced by indophenol or an indophenol related compound or redox transformation of the metabolite being analyzed. The concentration of ammonia and or ammonium ion and/or metabolite in blood correlates with the electron flow or current measurements on the circuit that comprise the at least one electrically conductive surfaces. The disclosure relates to the reduction to practice of this concept, showing how to select the metabolite, how to choose an immobilized enzyme, how to do the immobilization (what polymer, what additives, etc), how to attach the components to the electrode, how to make a measurement and how to develop a prototype. This disclosure is used to measure ammonia or ammonium ion and/or metabolites in blood of patients in real time. Aside from the sensor disclosed herein, there are no known sensors able to measure the proposed metabolites in real time.

[0029] The disclosure also relates to a device or system comprising at least one electrically conductive surface (such as an electrode) operably connected to a diode, a spectrophotometer, voltmeter and/or amperometer, the electrode comprises components that, when combined and in the presence of an ammonia, cause an indophenol reaction. The indophenol reaction product comprises a molecule that emits a visible or known wavelength after exposure to light. In some aspects of the disclosure, the device and system disclosed herein comprise a diode, such as a photodiode, which emits light into the vessel comprising the indophenol reaction product thereby exciting the reaction product and causing the reaction product to emit a visible or known wavelength. In some aspects of the disclosure, the device and system disclosed herein comprise a spectrophotometer that detects and/or quantitates the intensity of the visible or known wavelength of light emitted by the indophenol reaction product.

[0030] The disclosure also relates to a device and/or system that detects and quantifies amino acids. In some aspects of the disclosure, the device and/or systems comprise a vessel or well that comprises a metabolic enzyme disclosed herein or a functional fragment thereof. In some aspects of the disclosure, the enzyme or fragment thereof is immobilized to the vessel into which sample is initially placed in the device, biosensor, system, test strip, or cartridge. After contact with a sample, the enzyme or functional fragment thereof releases at least one or a series of electrons and ammonia, such that ammonia is free in solution and capable of moving between a first vessel to a second vessel through a membrane disclosed herein. In some aspects of the disclosure, the device comprises at least a first and second electrically conductive surface, wherein the first electrically conductive surface comprises a

hydrogel comprising an enzyme disclosed herein and the second electrically conductive surface does not comprise a hydrogel or an enzyme; wherein a voltmeter and/or amperometer are configured in a circuit such that the voltmeter can detect a voltage differential between the first and second electrodes in the presence of an amino acid and/or wherein the amperometer can detect an increased current in the first electrode as compared to the second electrode in the presence of an amino acid. The at least one or a series of electrodes are released after one or more enzymes within the hydrogel catalyzes the oxidation of the amino acid in a bodily sample in the presence of the one or more amino acids.

[0031] Hydrogel formulations are used to entrap one or more enzymes (that utilizes the metabolite/analyte as a specific substrate for its reaction) along with, in some aspects of the disclosure, a requisite cofactor in close proximity to the at least first electrode surface, with the hydrogel providing a simultaneous exclusion of interfering ions and macromolecules (contained within the patient's blood sample) from the electrode sensor. The coated electrode is contained within an electrochemical detection device capable of converting redox equivalents generated by the enzyme reaction into electron flow which in turn is measured as a current or voltage differential. Analyte concentration is derived using a calibration curve that correlates amperage or voltage differential to concentration of amino acid in the sample of bodily fluid. In some aspects of the disclosure, a small volume of whole blood is applied to or ammonia from the whole blood diffuses to a vessel exposed to the electrode and the result is reported within minutes of the application or contact to the electrode. Depending on the exact analyte, specific enzyme(s) and cofactor(s) are incorporated into the electrode in order to achieve analyte-specific reaction and response. For example, to detect elevated phenylalanine, the enzyme phenylalanine dehydrogenase is immobilized to the at least one electrically conductive surface optionally contained within a hydrogel.

[0032] The disclosure provides a method of sorting a mixture of samples of bodily fluid comprising: contacting a plurality of bodily fluid samples to a device or system disclosed herein. In some aspects of the disclosure, the method of sorting or cataloging a mixture of samples of bodily fluid further comprises the step of determining one or more concentrations of ammonia, ammonium ion, and/or amino acid in the bodily fluid sample, if in respect to the ammonia or ammonium ion concentration, based upon the presence or quantity of indophenol reaction products in one or more vessels or a current value or voltage differential value measured by the device; and, if in respect to determining one or more concentrations of amino acid in solution, based upon a current value or voltage differential value measured by the device. In some aspects of the disclosure, the method further comprises the step of comparing the one or more concentration of ammonia, ammonium ion, and/or amino acid in a sample of bodily fluid with one or more concentrations of ammonia, ammonium ion, and/or amino acid in sample of bodily fluid obtained from subject who does not have or is not suspected of having one or more aminoacidopathies or hyperammonemia, and cataloging, compiling, or identifying whether a sample of bodily fluid from a subject has an aminoacidopathy and/or hyperammonemia based upon their similarities or differences in concentration value to a sample of bodily fluid from a subject without an aminoacidopathy and/or hyperammonemia. The disclosure provides a method of diagnosing a subject with an hyperammonemia comprising: contacting at least one sample of bodily fluid from the subject to a device or system disclosed herein. In some aspects of the disclosure, the method of diagnosing further comprises the step of determining one or more concentrations of ammonia and/or ammonium ion in a bodily fluid sample based upon a current value, voltage differential value, or a presence or absence of a wavelength of light emitted by an indophenol reaction product, indophenol or an indophenol related compound. In each case, the device and or system disclosed herein detects and/or measures such values. In some aspects of the disclosure, the method further comprises the step of comparing the one or more concentration of ammonia, ammonium ion, and/or amino acid in the one or more samples from the subject with one or more concentrations of amino acids in sample of bodily fluid obtained from subject who does not have or is not suspected of having one or more aminoacidopathies and/or hyperammonemia, identifying whether a sample of bodily fluid from a subject has an aminoacidopathy and or hyperammonemia based upon their similarities or differences in concentration value to the sample of bodily fluid from a subject without an aminoacidopathy and/or hyperammonemia.

[0033] The disclosure also provides a method of monitoring the concentrations of ammonia or ammonium ion in subject over time in a sample of bodily fluid from a subject diagnosed or suspected as having hyperammonemia, the method comprising: contacting one or more samples of bodily fluid from a subject to a device or system disclosed herein and measuring the concentration of ammonia or ammonium ion of bodily fluid from the subject at

one time point and performing a repeating of the measurement at least once at a different time point. In some of the disclosure, the method of monitoring the concentrations of ammonia or ammonium ion in subject over time in a sample of bodily fluid from a subject diagnosed or suspected as having hyperammonemia further comprises the step of cataloguing the concentration values of the ammonia or ammonium ion over time. In some aspects of the disclosure, the method further comprises the step of comparing the one or more concentration of amino acids from the plurality of samples of bodily fluid from the subject over time and, optionally notifying a subject if the concentration of one or more ammonia or ammonium ion reaches or exceeds or drops below a threshold value that requires medical treatment or modification of diet.

[0034] In some aspects of the disclosure, samples of bodily fluid are isolated from a subject having been diagnosed with or suspected as having hyperammonemia. For example, in some aspects of the disclosure, a sample of bodily fluid such as a urine sample or a blood sample is isolated from the subject. The sample of bodily fluid is contacted to at least one electrode comprising at least one enzyme disclosed herein and the amino acid concentration in the sample of bodily fluid is measured based upon the magnitude of the voltage differential or current detected by the device comprising the at least one electrode. In further aspects of the disclosure, method of the present disclosure comprises contacting a sample of bodily fluid to at least one electrode comprising an immobilized enzyme disclosed herein, measuring the current or voltage difference between the at least one electrode and an electrically conductive surface that does not comprise an immobilized enzyme disclosed herein, determining the concentration of one or more amino acids in the sample of bodily fluid, and optionally, providing a readout of one or more concentration values to a subject from which the sample of bodily fluid was obtained. In further aspects of the disclosure, method of detecting ammonia or ammonium ion comprises contacting a sample of bodily fluid to at least one vessel comprising an hyohalite, an aqueous basic solution, and at least one compound comprising a phenyl group disclosed herein, measuring the current or voltage difference between the at least one electrode and an electrically conductive surface that does not comprise an immobilized enzyme disclosed herein, determining the concentration of ammonia or ammonium ion in the sample of bodily fluid, and optionally, providing a readout of one or more concentration values to a subject from which the sample of bodily fluid was obtained.

[0035] In some aspects of the disclosure, the present disclosure provides methods comprising contacting a sample of bodily fluid from a subject to an aqueous basic solution or a basic buffer in a dried or powdered phase, exposing the sample to hyohalite and at least one compound comprising a phenyl group in the presence (or absence --- to establish a control value) of a membrane comprising an ionomer, and optionally contacting a gel. In some aspects of the disclosure, the gel is a hydrogel comprising alginate. In some aspects of the disclosure, the present disclosure provides methods comprising detecting presence or level ammonia or ammonium ion in a sample of bodily fluid between cells in the sample. In some aspects of the disclosure, provided methods comprise determining that a particular set of detected interactions defines an threshold value (or control value) that is characteristic of an increased severity of hyperammonemia in that it distinguishes them from elevated or non-elevated amino acid levels in another sample of bodily fluid from the subject or from a sample of bodily fluid that is a reference or control sample such that, if the threshold value is reached, the device or system disclosed herein provides the subject with a signal or notification that treatment or diet modification should be sought. In some aspects of the disclosure, the step of detecting comprises detecting presence or level of ammonia or ammonium ion concentrations in a sample of bodily fluid that is characteristic of particular severity of disease in the sample in that it distinguishes them from a sample of bodily fluid that is a reference or control sample.

[0036] In some aspects of the disclosure, the step of detecting comprises detecting presence or level of ammonia or ammonium ion concentrations in a sample of bodily fluid that is characteristic of particular severity of disease in the sample in that it distinguishes them from a sample of bodily fluid that is a reference or control sample.

[0037] In some aspects of the disclosure, any of the methods disclosed herein do not comprise pre-treating the sample of bodily fluid prior to contacting the sample with the test strip, conduit, biosensor, and/or at least one electrically conductive surface. In some aspects of the disclosure, any of the methods disclosed herein do not comprise using at step of treating the sample with liquid chromatography, gas chromatography, and/or electrophoresis before, simultaneously with or after contacting the sample to the test strip, conduit, biosensor,

and/or at least one electrically conductive surface. In some aspects of the disclosure, any of the methods disclosed herein comprise contacting the sample to at least one electrode that does not comprise an enzyme obtained from an organism other than a bacteria or a plant.

[0038] The disclosure relates to methods of detecting the levels of ammonia or ammonium ion in whole blood by exposing a whole blood sample to one of the biosensors, systems, or devices disclosed herein. The disclosure also relates to manufacturing a biosensor disclosed herein by treating the membrane with one, two, three or more washes of an acidic solution prior to placement of the membrane at a fluid exchange opening or at a vessel. The disclosure relates to manufacturing a biosensor disclosed herein by treating the membrane with one, two, three or more washes of an acidic solution from about 0.1 M to about 1 M H_2SO_4 prior to placement of the membrane at a fluid exchange opening or at a vessel. The disclosure also relates to manufacturing a biosensor disclosed herein by treating the membrane with one, two, three or more washes of a hydrogen peroxide solution from about 0.1 M to about 1 M H_2O_2 prior to placement of the membrane at a fluid exchange opening or at a vessel. The disclosure relates to manufacturing a biosensor disclosed herein by treating the membrane with one, two, three or more washes of an acid solution and/or a hydrogen peroxide solution prior to placement of the membrane at a fluid exchange opening or at a vessel. The disclosure also relates to manufacturing a biosensor disclosed herein by treating the membrane with one, two, three or more washes of an acid solution from about 0.1 M to about 1 M H_2O_2 prior to placement of the membrane at a fluid exchange opening or at a vessel. The disclosure also relates to manufacturing a biosensor disclosed herein by treating the membrane with one, two, three or more of an acid solution comprising from about 0.1 M to about 1 M H_2SO_4 and with one, two, three or more washes of a hydrogen peroxide solution comprising from about 0.1 M to about 1 M H_2O_2 prior to placement of the membrane at a fluid exchange opening or at a vessel.

[0039] In some aspects of the disclosure, the present disclosure provides a system comprising one or more devices disclosed herein optionally in operable connection to a electronic storage medium that compiles ammonia or ammonium ion and/or amino acid concentration values of a subject. In some aspects of the disclosure, the electronic storage medium comprises compiled amino acid concentration values of a subject over time. In some aspects of the disclosure, the system comprises at least one electrically conductive surface that comprises an enzyme disclosed herein, a mediator, and optionally a gel or hydrogel. In some aspects of the disclosure, the system comprises an electronic circuit that is in operable connection to the at least one electrodes and a diode, spectrophotometer, voltmeter and/or amperometer. In the case of a diode and/or spectrophotometer, the diode or spectrophotometer detect and wavelength of light emitted from the at least one vessel. In the case of the voltmeter and/or amperometer, the voltmeter and/or amperometer measures the respective voltage and/or amperage of the circuit across the at least one electrode when the at least one electrode is in the presence of one or more amino acids and/or ammonia concentrations. In some aspects of the disclosure, system comprising one or more devices disclosed herein optionally in operable connection to a electronic storage medium that compiles amino acid concentration values of a subject determines one or a plurality of concentration values of ammonia concentration values and/or amino acids in a sample of bodily fluid when the sample of bodily fluid is in contact with the at least one electrode and under conditions and for a time sufficient for the indophenol reaction to take place or the one or more enzymes disclosed herein to oxidize its amino acid substrate, create a voltage differential or current change in the circuit and the device to display the concentration value on one or more displays. In some aspects of the disclosure, the device, system, and/or biosensor do not comprises one or more electrodes.

[0040] In some aspects of the disclosure, the disclosure provides for a method comprising steps of: contacting a sample comprising cells with an electrode. The disclosure further provides for a method comprising steps of: contacting a sample under conditions and for a time sufficient for a set of interactions to occur between ammonia in a sample and the membrane described herein. The disclosure further provides for a method comprising steps of: contacting a whole blood sample under conditions and for a time sufficient for a set of interactions to occur between ammonia in a sample and the membrane described herein. The disclosure further provides for a method comprising steps of: contacting a sample comprising bodily fluid under conditions and for a time sufficient for a set of interactions to occur between the ammonia in the sample and the one or plurality of indophenol reaction reagents described herein.


[0041] The disclosure relates to a biosensor comprising: at least one electrically conductive support, the electrically conductive support attached to a hydrogel, the hydrogel comprising at least one electron mediator, at least one reduction agent, and at least one metabolic enzyme or functional fragment thereof, wherein the hydrogel comprises alginate; and an amperometer and/or voltmeter operably connected to the at least one electrically conductive support or surface.

[0042] In some aspects of the disclosure, the biosensor comprises at least three electrically conductive supports. In some aspects of the disclosure, the at least one electrically conductive support is a silver and silver chloride wire. In some aspects of the disclosure, the at least one electrically conductive support comprises at least one or a combination of metabolic enzymes chosen from: leucine dehydrogenase, tyrosine dehydrogenase, phenylalanine dehydrogenase, leucine oxidoreductase, tyrosine monooxygenase, alanine dehydrogenase, or glutamate dehydrogenase; or functional fragments thereof. In some aspects of the disclosure, the biosensor comprises at least a first and a second electrically conductive support, wherein the first electrically conductive support is attached to a hydrogel, the hydrogel comprising at least one electron mediator, at least one reduction agent, and at least one metabolic enzyme or functional fragment thereof, wherein said first and second electrically conductive supports being operably connected to said voltmeter and/or amperometer to apply a voltage therebetween.

[0043] In some aspects of the disclosure, the at least one electrically conductive support comprises an electronegative or anionic chemical component. In some aspects of the disclosure, the at least one hydrogel comprises trehalose. In some aspects of the disclosure, the biosensor does not comprise one or more of the following: (i) uricase or a functional fragment thereof; (ii) a hydrogel comprising dextran or a derivative thereof; (iii) a bacterial cell; (iv) an electronic dipole configured for electrophoresis; and (v) 3, 4-DHB. In some aspects of the disclosure, the biosensor is at least 70% biologically active after about sixteen days in storage at 4 degrees Celsius. In some aspects of the disclosure, the biosensor is at least 70% biologically active after about thirty days in storage at 4 degrees Celsius. In some aspects of the disclosure, the biosensor is at least 80% biologically active after about thirty days in storage at 4 degrees Celsius. In some aspects of the disclosure, the biosensor is at least 90% biologically active after about thirty days in storage at 4 degrees Celsius. In some aspects of the disclosure, the biosensor is at least 95% biologically active after about thirty days in storage at 4 degrees Celsius. In some aspects of the disclosure, the biosensor is not functionally dependent upon exposure to UV light or addition of any stimulus external to the biosensor. In some aspects of the disclosure, the at least one enzyme or functional fragment thereof is derived from a bacterial species and is immobilized in the hydrogel. In some aspects of the disclosure, the at least one enzyme or functional fragment thereof is derived from a thermophilic bacterial species and is immobilized in the hydrogel. In some aspects of the disclosure, the at least one enzyme or functional fragment thereof comprises at least about 70% sequence identity to SEQ ID NO:1 or SEQ ID NO:2.

[0044] In some aspects of the disclosure, the disclosure relates to a biosensor, device, or system disclosed herein comprise a circuit comprising at least a first and second electrode in electric communication to at least one or a combination of a diode, photodiode, spectrophotometer, or other device capable of measuring the presence, absence, or intensity of light emitted by an amount of indophenol or indophenol relate compound exposed to light. In some embodiments the circuit comprises a wire. In some aspects of the disclosure, the wire comprises silver and silver chloride in operable connection to the voltmeter and/or amperometer.

[0045] In some aspects of the disclosure, the biosensor, device, and/or system disclosed herein comprises a membrane optionally comprising alginate comprises a block polymer with a formula



wherein m and n are any positive integer.

[0046] In some aspects of the disclosure, the biosensor the at least one electrically conductive support is not

covered by a membrane comprising cellulose or a derivative thereof. In some aspects of the disclosure, the at least one electron mediator is selected from: thionine, o-phenylenediamine, methylene blue, and toluidine blue. In some aspects of the disclosure, the at least one reduction agent is chosen from: NAD⁺ or FAD⁺.

[0047] The disclosure also relates to a biosensor comprising: at least one electrically conductive support, the electrically conductive support attached to at least one hydrogel, the hydrogel comprising at least one electron mediator, at least one reduction agent, and at least one metabolic enzyme or functional fragment thereof; wherein the at least one enzyme or functional fragment thereof is at least 70% homologous to a phenylalanine dehydrogenase from *Geobacillus thermoglucosidiasus*; and an amperometer and/or voltmeter operably connected to the at least one electrically conductive support. In some aspects of the disclosure, the enzyme or functional fragment thereof is at least 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% homologous to SEQ ID NO:1 or at least 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% homologous to a functional fragment of SEQ ID NO:1. In some aspects of the disclosure, the enzyme or functional fragment thereof is not derived from a species other than a bacterial cell. In some aspects of the disclosure, the enzyme or functional fragment thereof is at least 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% homologous to SEQ ID NO:2 or at least 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% homologous to a functional fragment of SEQ ID NO:2.

[0048] The disclosure relates to a system comprising a biosensor comprising at least a first and second vessel; a fluid exchange opening positioned between the at least first and second vessel; at least one conduit in fluid communication with the at least first vessel, the at least one conduit configured to receive a fluid from a point external to the biosensor; and a membrane positioned at the fluid exchange opening; wherein the membrane comprises an ionomer, and wherein the first vessel or the second vessel comprise, individually or in combination: hyohalite, an aqueous basic solution, and at least one compound comprising a phenyl group.

[0049] The disclosure also relates to a system comprising a biosensor disclosed herein optionally comprising an electric circuit comprising any one or combination of: a diode (such as a photodiode), a spectrophotometer, an amperometer and/or voltmeter operably connected to the at least one electrically conductive support or surface; wherein the biosensor is in operable connection to at least one computer storage memory. In some aspects of the disclosure, the system further comprises a sample of bodily fluid, such as whole blood. In some aspects of the disclosure, the system further comprises a digital display in operable connection to the at least one electrically conductive support (or surface) by an electrical circuit capable of carrying an electrical signal corresponding to a measurement of current and/or voltage differential from the diode, a spectrophotometer, voltmeter and/or amperometer to the digital display, wherein the digital display is configured to display one or more concentration values of ammonia or ammonium ion and/or an amino acid in a sample over time when the at least one electrically conductive support (or surface) is in contact with the sample for a time period sufficient for the indophenol reaction to take place.

[0050] In some aspects of the disclosure, the system further comprises a computer processor in operable connection with the at least one computer storage memory. In some aspects of the disclosure, the metabolic enzyme is a phenylalanine dehydrogenase immobilized within the hydrogel and wherein the alginate concentration of the hydrogel is from about 1% to about 3% weight to volume of the total volume attached and/or contacted to the at least one electrically conductive support.

[0051] The disclosure also relates to a kit comprising a biosensor comprising a diode, spectrophotometer, voltmeter and/or amperometer and a display configured in an electrical circuit such that, upon contact with at least one removable electrically conductive support, the circuit becomes closed such that the diode, spectrophotometer, voltmeter and/or amperometer are in operable communication with at least one electrically conductive support.

[0052] In some aspects of the disclosure, the kit comprises at least one of the following: a plurality of test strips comprising one or a plurality of vessels configured to receive a sample, such as whole blood, wherein the one or plurality of test strips further comprises at least one conduit in fluid communication with the at least first vessel. In some aspects of the disclosure, the kit comprises at least one of the following: a plurality of test strips comprising one or a plurality of vessels configured to receive a sample, such as whole blood, wherein the one or

plurality of test strips further comprises at least one conduit in fluid communication with the at least first vessel and, individually or in combination: hyohalite, an aqueous basic solution, and at least one compound comprising a phenyl group. In some aspects of the disclosure, the kit comprises at least one of the following: a plurality of test strips comprising one or a plurality of vessels configured to receive a sample, such as whole blood, wherein the one or plurality of test strips further comprises at least one conduit in fluid communication with the at least first vessel and a biosensor comprising a membrane disclosed herein. In some aspects of the disclosure, the kit comprises at least one of the following: a plurality of test strips comprising: one or a plurality of vessels configured to receive a sample, such as whole blood, wherein the one or plurality of test strips further comprises at least one conduit in fluid communication with the at least first vessel; and a biosensor comprising a membrane disclosed herein.

[0053] In some aspects of the disclosure, the membrane comprises a hydrogel layer. In some aspects of the disclosure, the hydrogel layer comprises alginate. In some aspects of the disclosure, the a control or reference sample of bodily fluid; a set of data comprising threshold values; and a set of instructions, wherein the set of instructions or the set of data optionally accessible remotely through an electronic medium. In some aspects of the disclosure, the kit comprises a solid support that comprises at least a first and a second electrode, wherein the first electrode comprises a hydrogel, the hydrogel comprising at least one electron mediator, at least one reduction agent, and at least one metabolic enzyme or functional fragment thereof; and wherein the second electrode is a control or reference electrode.

In some aspects of the disclosure, the kit comprises a test strip comprising a solid support attached to a first and a second electrode described herein.

[0054] The disclosure also relates to a method of determining or identifying a concentration of an ammonia or ammonium ion in a sample of bodily fluid comprising: (a) contacting a sample of bodily fluid to: (i) a biosensor comprising at least one electrically conductive support, the electrically conductive support attached to a vessel in fluid communication with a membrane disclosed herein and, optionally an amperometer and/or voltmeter operably connected to the at least one electrically conductive support; or (ii) a system comprising a biosensor comprising: at least one electrically conductive support attached to a vessel in fluid communication with a membrane disclosed herein and, optionally an amperometer and/or voltmeter operably connected to the at least one electrically conductive support; or (iii) a test strip disclosed herein; and/or (b) determining a quantity of ammonia or ammonium ion in the sample. In some aspects of the disclosure, the sample of bodily fluid comprises blood or serum from a subject. In some aspects of the disclosure, the sample consists of whole blood or consists essentially of whole blood.

[0055] The disclosure also relates to a method of quantifying a concentration of ammonia and/or an amino acid in a sample of bodily fluid comprising: (a) contacting a sample of bodily fluid to: (i) a biosensor comprising at least one electrically conductive support or surface, the electrically conductive support or surface attached to a hydrogel, the hydrogel comprising at least one electron mediator, at least one reduction agent, and at least one metabolic enzyme or functional fragment thereof, wherein the hydrogel comprises alginate; and an amperometer and/or voltmeter operably connected to the at least one electrically conductive support or surface; or (ii) a system comprising a biosensor comprising: at least one electrically conductive support or surface, the electrically conductive support or surface attached to at least one hydrogel, the hydrogel comprising at least one electron mediator, at least one reduction agent, and at least one metabolic enzyme or functional fragment thereof; wherein the at least one enzyme or functional fragment thereof is at least 70% homologous to a phenylalanine dehydrogenase from *Geobacillus thermoglucosidiasus*; and an amperometer and/or voltmeter operably connected to the at least one electrically conductive support; or (iii) a test strip disclosed herein; or (b) determining a quantity of amino acid in the sample. In some aspects of the disclosure, the method further comprises comparing a concentration value obtained by the quantifying or identifying steps to a threshold value associated with one or more metabolic diseases.

[0056] The disclosure further relates to a method comprising a step of contacting a biosensor, system, or test strip disclosed herein, wherein the step of contacting a sample of bodily fluid of a subject to any of the disclosed biosensors, systems, or test strips comprises contacting the sample for a sufficient period of time to allow ammonia

transport through the membrane and to expose the ammonia from the sample to reagents associated with an indophenol reaction. If amino acids are also being tested by the biosensors, systems, or test strips, such methods comprise contacting a sample for a sufficient period of time to allow oxidation of at least one amino acid in the sample of bodily fluid by the metabolic enzyme or functional fragment thereof. In some aspects of the disclosure, the method does not comprise exposing the sample of bodily fluid to any external stimuli or reagent prior to contacting the sample to the at least one electrically conductive supports. In some aspects of the disclosure, the method does not comprise exposing the sample of bodily fluid to iron ions and/or hydrozide ions prior to, simultaneously with, or after exposing the sample to the at least one electrode comprising a hydrogel. In some aspects of the disclosure, the method does not comprise exposing the sample to a non-porous carrier, such as glass beads, contained within the device, test strip or biosensor. In some aspects of the disclosure, the sample of bodily fluid contains whole blood or serum from a subject. In some aspects of the disclosure, the sample of bodily fluid does not contain urine. In some aspects of the disclosure, the sample of bodily fluid does not contain bodily fluid other than whole blood or blood serum.

[0057] The disclosure further relates to a method of diagnosing a metabolic disease in a subject comprising: (a) contacting a sample of bodily fluid to one or a combination of: (i) a biosensor comprising at least one electrically conductive support or surface, the electrically conductive support or surface attached to a vessel comprising an amount of indophenol or indophenol related compound; and, optionally an amperometer and/or voltmeter operably connected to the at least one electrically conductive support or surface; or (ii) a system comprising a biosensor comprising: at least one electrically conductive support or surface, the electrically conductive support or surface exposed to the at least first vessel or second vessel comprising the indophenol and/or indophenol related compounds; and an amperometer and/or voltmeter operably connected to the at least one electrically conductive support; or (iii) a test strip disclosed herein; (b) quantifying one or more concentration values of ammonia or ammonium ion in the sample; (c) comparing the one or more concentration values of ammonia or ammonium ion in the sample to a threshold value of ammonia or ammonium ion concentration identified as being in a healthy range or not within a range or concentration indicative of hyperammonemia; and (d) identifying the subject as having hyperammonemia or a metabolic disease related to hyperammonemia if the one or more concentration values of amino acids in the sample exceed or fall below the threshold value. In some aspects of the disclosure, the metabolic disease is a hyperammonemia related disorder.

[0058] The disclosure also relates to a method of determining patient responsiveness to a therapy comprising: (a) contacting a sample of bodily fluid to one or a combination of: (i) a biosensor comprising at least one electrically conductive support or surface, the electrically conductive support or surface attached to a vessel comprising an amount of indophenol or indophenol related compound; and, optionally an amperometer and/or voltmeter operably connected to the at least one electrically conductive support or surface; or (ii) a system comprising a biosensor comprising: at least one electrically conductive support or surface, the electrically conductive support or surface exposed to the at least first vessel or second vessel comprising the indophenol and/or indophenol related compounds; and an amperometer and/or voltmeter operably connected to the at least one electrically conductive support; or (iii) a test strip disclosed herein; (b) quantifying one or more concentration values of ammonia or ammonium ion in the sample; (c) comparing the one or more concentration values of ammonia or ammonium ion in the sample to a threshold value of ammonia or ammonium ion concentration identified as being in a healthy range or not within a range or concentration indicative of hyperammonemia; and (d) identifying the subject as having hyperammonemia or a metabolic disease related to hyperammonemia if the one or more concentration values of amino acids in the sample exceed or fall below the threshold value.

[0059] The disclosure also relates to a test strip comprising a solid support and at least a first and a second electrode attached to the solid support, wherein the first electrode comprises a membrane, the membrane comprising a perfluorinated ionomer. In some aspects of the disclosure, the test strip is adapted for a portable device comprising: a diode, spectrophotometer, voltmeter and/or amperometer; and a digital display such that, when the test strip is contacted to the device, the first and second electrodes become operably connected to a closed electrical circuit comprising diode, spectrophotometer, voltmeter and/or amperometer and the digital display, and, upon contact with light emitted from an indophenol or indophenol related compound, resulting in a current on the first electrode corresponding to a concentration value of amino acid in the sample of bodily fluid, such

concentration value readable on the display of the portable device. In some aspects of the disclosure, the test strip comprises the at least one or combination of indophenol reagents in solid or liquid phase optionally separated from but in fluid communication with a conduit, volume, or space the at least first vessel.

[0060] The disclosure also relates to a method of manufacturing any of the disclosed biosensors, test strips, systems disclosed herein that comprises at least one electrode, the method comprising: contacting the at least one electrode with a solution comprising at least one vessel, at least one conduit in fluid communication with the at least one vessel, and at least one indophenol reagent; subsequently contacting the at least one electrode with a basic buffer with a Na⁺, Ca⁺, Cl⁻, and/or acetate concentration at or below about 1 M.

BRIEF DESCRIPTION OF THE DRAWINGS

[0061] Advantages of embodiments of the present invention will be apparent from the following detailed description of the exemplary embodiments. The following detailed description should be considered in conjunction with the accompanying figures in which:

FIG. 1 is an exemplary view of a system having the ability to detect ammonia or ammonium ion levels in a given sample applied to a first and second vessel separated by a membrane positioned at an fluid exchange opening.

FIG. 2 is an exemplary view of a system comprising multiple vessels within which more than one indophenol reaction may be performed in parallel.

FIG. 3 shows exemplary reaction otherwise known as Berthelot's Reaction or an indophenol reaction.

FIG. 4 shows an exemplary embodiment of a microfluidic testing device.

FIG. 5 shows an exemplary flowchart for a method of quantitative point of care hyperammonemia sensing using embodiments of the disclosure.

FIG. 6 shows an exemplary embodiment of a blood test strip for use with an electronic testing device.

FIG. 7 shows an exemplary embodiment of a device comprising an electronic circuit comprising an electrode exposed to a vessel configured for performance of the indophenol reaction; an analog to digital convertor, a microchip in electronic communication with a display.

FIG. 8 shows the chemical composition of Nafion.

FIG. 9 shows experimental data demonstrating how the concentration of sodium salt yields high recovery and transfer of ammonia from a sample.

FIG. 10 shows experimental data demonstrating the differences in device performance when using sodium acetate versus sodium chloride as a basic buffer.

FIG. 11 depicts a photograph of the 3D printed modular pieces snapped together around Nafion to form the bisected well utilized for the sensing experiments

FIG. 12 depicts an indophenol reaction produces a linear curve with concentrations of ammonium chloride ranging from 0-750 μ M with a COD of 0.9939.

FIG. 13 depicts reagents for the indophenol reaction were stored at room temperature and used to generate an ammonia standard curve at regular intervals for 100 days. The response to 500 μ M ammonia began to degrade at day 75. The reagents of the indophenol reaction are stable at room temperature for up to 50 days before its response to different concentrations of ammonia begins to deteriorate.

FIG. 14 depicts 1mM concentrations of each of the 21 amino were tested using the indophenol reaction. The absorbance measured at 635nm for each amino acid after the indophenol reaction was calculated as percentage of the response from indophenol reaction with 1mM ammonium chloride. The radar graph displays the percent

response as compared to ammonium chloride. The highest response was threonine which produced an absorbance value that was just 7% of ammonia's response.

FIG. 15: The constructed sensor's response to a range of ammonia concentrations in 1X PBS. The COD is 0.9758 with n=5 samples.

FIG 16: Initial experiments of determining blood ammonia concentration demonstrated a limited response. Responses were hindered and would not exceed an absorbance of 0.35 indicating some degree of interference.

FIG 17: Concentrations of 2-10X hypochlorite were utilized in the analysis of 500mM ammonia in 1X PBS and whole sheep's blood. Increasing the concentration of hypochlorite utilized in the indophenol reaction reduced the negative interference small blood molecules had on the indophenol reaction. At concentrations higher than 3X, reaction itself began to degrade. A 3-fold increase in hypochlorite concentration was optimal.

FIG 18: The bisected well sensor was again used to extract ammonia in whole human blood. The extracted ammonia solutions were tested with the 3X hypochlorite-modified indophenol reaction and the absorbance measured at 635nm. In the range of 0-500 μ M the COD was 0.9573 with n=5 samples.

FIG 19: The sensor's response to blood ammonia concentrations ranging from 0-150 μ M. The relative standard deviation is ~10% with a COD of 0.9777 with n=5 samples.

FIG. 20: depicts a CAD sketch of the front face of the well plate of an embodiment.

FIG. 21: depicts a CAD sketch demonstrating the area of the well plate of an embodiment that the adhesive silicone should be attached to (black area) prior to completing the manufacturing of the device.

FIG. 22 depicts a photograph of the 3D-printed modular left and right sides pieces snapped together around Nafion® to divide the well into two sections.

FIG. 23 depicts an engineering drawing of 3D printed well.

FIG. 24 depicts a CAD sketch of the top piece of a disposable cartridge, with dimensions in mm. Channels 1 through 5 are labeled.

FIG. 25: CAD sketch of the bottom piece of the chip with channel 6 labeled.

FIG. 26: depicts the representations of a concentration profile at i) t=0 seconds (s) ; ii) t=13 s , and iii) t=24 s after a whole blood sample in 40 microliters is loaded into well number 6.

FIG. 27 depicts the top half of an embodiment comprising a microfluidic device used to quantify ammonia levels in whole blood.

FIG. 28 depicts the bottom half of an embodiment comprising a microfluidic device used to quantify ammonia levels in whole blood.

FIG. 29: is an exemplary view of a system having the ability to detect ammonia or ammonium ion levels and amino acids in a given sample applied to a first and second vessel separated by a membrane positioned at an fluid exchange opening, wherein the reaction is catalyzed by an enzyme.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0062] Various terms relating to the methods and other aspects of the present disclosure are used throughout the specification and claims. Such terms are to be given their ordinary meaning in the art unless otherwise indicated. Other specifically defined terms are to be construed in a manner consistent with the definition provided herein.

[0063] As used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise.

[0064] The term "about" as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of $\pm 20\%$, $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, or $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods.

[0065] As used herein, the terms "attach," "attachment," "adhere," "adhered," "adherent," or like terms generally refer to immobilizing or fixing, for example, a group, a compound or enzyme, to a surface, such as by physical absorption, chemical bonding, and like processes, or combinations thereof.

[0066] As used herein, the terms "biopsy" means a cell sample, collection of cells, or bodily fluid removed from a subject or patient for analysis. In some embodiments, the biopsy is a bone marrow biopsy, punch biopsy, endoscopic biopsy, needle biopsy, shave biopsy, incisional biopsy, excisional biopsy, or surgical resection.

[0067] As used herein, the terms "bodily fluid" means any fluid isolated from a subject including, but not necessarily limited to, blood sample, serum sample, a whole blood sample, urine sample, mucus sample, saliva sample, and sweat sample. The sample may be obtained from a subject by any means such as intravenous puncture, biopsy, swab, capillary draw, lancet, needle aspiration, collection by simple capture of excreted fluid.

[0068] As used herein the terms "electronic medium" mean any physical storage employing electronic technology for access, including a hard disk, ROM, EEPROM, RAM, flash memory, nonvolatile memory, or any substantially and functionally equivalent medium. In some embodiments, the software storage may be co-located with the processor implementing an embodiment of the disclosure, or at least a portion of the software storage may be remotely located but accessible when needed.

[0069] As used herein, the word "exemplary" means "serving as an example, instance or illustration." The embodiments described herein are not limiting, but rather are exemplary only. It should be understood that the described embodiment are not necessarily to be construed as preferred or advantageous over other embodiments. Moreover, the terms "embodiments of the invention", "embodiments" or "invention" do not require that all embodiments of the invention include the discussed feature, advantage or mode of operation. In addition, those skilled in the art may appreciate the wide variations in sizing scales that may be incorporated into the disclosed or related designs for use with samples many orders of magnitude larger or smaller than those disclosed.

[0070] As used herein, the term "aminoacidopathy" is meant to refer to those diseases and disorders characterized by dysfunction of a metabolic catalysis of amino acids that results in over production or under production of amino acids in the body of a subject.. Examples of aminoaciopathies are listed in the definition of a metabolic disease, terms that are used interchangeably in this application.

[0071] As used herein, "sequence identity" is determined by using the stand-alone executable BLAST engine program for blasting two sequences (bl2seq), which can be retrieved from the National Center for Biotechnology Information (NCBI) ftp site, using the default parameters (Tatusova and Madden, FEMS Microbiol Lett., 1999, 174, 247-250). To use the term "homologous to" is synonymous with a measured "sequence identity." In some embodiments, if an embodiment comprises a nucleic acid sequence or amino acid sequence with a percent sequence identity the term refers to a disclosed nucleic acid sequence or amino acid sequence possessing a homology to a disclosed sequence over its entire length.

[0072] The term "subject" is used throughout the specification to describe an animal from which a sample of bodily fluid is taken. In some embodiment, the animal is a human. For diagnosis of those conditions which are specific for a specific subject, such as a human being, the term "patient" may be interchangeably used. In some instances in the description of the present disclosure, the term "patient" will refer to human patients suffering from a particular disease or disorder. In some embodiments, the subject may be a human suspected of having or being identified as at risk to develop an aminoacidopathy. In some embodiments, the subject may be diagnosed as having at least

one aminoacidopathy. In some embodiments, the subject is suspected of having or has been diagnosed with hyperammonemia. In some embodiments, the subject may be a human suspected of having or being identified as at risk to develop hyperammonemia. In some embodiments, the subject may be a mammal which functions as a source of the isolated sample of bodily fluid. In some embodiments, the subject may be a non-human animal from which a sample of bodily fluid is isolated or provided. The term "mammal" encompasses both humans and non-humans and includes but is not limited to humans, non-human primates, canines, felines, murines, bovines, equines, and porcines.

[0073] As used herein, "conservative" amino acid substitutions may be defined as set out in Tables A, B, or C below. Metabolic enzymes include those amino acid sequences wherein conservative substitutions have been introduced by modification of polynucleotides encoding polypeptides disclosed herein. Amino acids can be classified according to physical properties and contribution to secondary and tertiary protein structure. A conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in Table A.

Table A -- Conservative Substitutions I

Side Chain Characteristics	Amino Acid
Aliphatic	
Non-polar	G A P I L V F
Polar - uncharged	C S T M N Q
Polar - charged	D E K R
Aromatic	H F W Y
Other	N Q D E

[0074] Alternately, conservative amino acids can be grouped as described in Lehninger, (Biochemistry, Second Edition; Worth Publishers, Inc. NY, N.Y. (1975), pp. 71-77) as set forth in Table B.

Table B -- Conservative Substitutions II

Side Chain Characteristic	Amino Acid
Non-polar (hydrophobic)	
Aliphatic:	A L I V P
Aromatic:	F W Y
Sulfur-containing:	M
Borderline:	G Y
Uncharged-polar	
Hydroxyl:	S T Y
Amides:	N Q
Sulfhydryl:	C
Borderline:	G Y
Positively Charged (Basic):	K R H
Negatively Charged (Acidic):	D E

[0075] Alternately, exemplary conservative substitutions are set out in Table C.

Table C -- Conservative Substitutions III

Original Residue	Exemplary Substitution
Ala (A)	Val Leu Ile Met
Arg (R)	Lys His
Asn (N)	Gln

Original Residue	Exemplary Substitution
Asp (D)	Glu
Cys (C)	Ser Thr
Gln (Q)	Asn
Glu (E)	Asp
Gly (G)	Ala Val Leu Pro
His (H)	Lys Arg
Ile (I)	Leu Val Met Ala Phe
Leu (L)	Ile Val Met Ala Phe
Lys (K)	Arg His
Met (M)	Leu Ile Val Ala
Phe (F)	Trp Tyr Ile
Pro (P)	Gly Ala Val Leu Ile
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr Phe Ile
Tyr (Y)	Trp Phe Thr Ser
Val (V)	Ile Leu Met Ala

[0076] It should be understood that the polypeptides comprising polypeptide sequences associated with the extracellular matrix described herein are intended to include polypeptides bearing one or more insertions, deletions, or substitutions, or any combination thereof, of amino acid residues as well as modifications other than insertions, deletions, or substitutions of amino acid residues.

[0077] As used herein, the term "prognosing" means determining the probable course and/or outcome of a disease.

[0078] As used herein, the term indophenol related compound - a small chemical compound that is a reaction product of an indophenol reaction. In some embodiment, it comprises at least one carbon atom in a 4, 5, 6-membered ring and emits a visible wavelength of light upon excitation of the small chemical compound by light emitted by from light source. In some embodiments, the small chemical compound is a product of the indophenol reaction and emits a wavelength of light visible to the human eye upon excitation of the chemical compound by light emitted from a light source. In some embodiments, the small chemical compound emits a wavelength from about 400 nm to about 600 nm when it is excited by light from a light source. In some embodiments, the biosensor, device, and/or system comprises a light source and at least one diode and/or spectrophotometer, or other device capable of measuring the light emitted by the indophenol or the indophenol related compound.

[0079] The term "vessel" as used herein is any chamber, indentation, container, receptacle, or space. In some embodiments, a vessel is a well capable of holding no more than about 1,000, 900, 800, 700, 600, 500, 400, 300, 200, 100, 50, 40, 30, 20, 15, 12, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 μ L of sample. bodily fluid.

[0080] The term "membrane" means any monomer or polymer in a solid phase. In some embodiments, the membrane comprises an ionomer. In some embodiments, the membrane is incapable of gas chromatography.

[0081] The terms "point of care" disclosed herein refer to a device, biosensor, system, test strip, or cartridge, either individually or configured to function with one or more additional components, capable of analyzing the presence, absence, or quantity of a reaction product, such as ammonia, and/or a sample component, such as an amino acid, within a time period no more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, or 30 minutes. In some

embodiments, the terms refer to a device, biosensor, system, test strip, or cartridge, either individually or configured to function with one or more additional components, capable of analyzing the presence, absence, or quantity of ammonia and/or an amino acid within a time period no more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, or 30 minutes, or capable analyzing the presence, absence, or quantity of ammonia and/or an amino acid at or substantially near the point from which the sample was taken. For instance, in some embodiments, the sample may be taken from a subject suspected of or previously diagnosed with hyperammonemia or a hyperammonemia-related disorder. Without sending and analyzing the ammonia content of a sample to a different location from the source of the sample, in some embodiments, the point of care device or biosensor or system is a point of care device which is capable of detecting the presence, absence, or quantity of ammonia or ammonium ion in a sample.


[0082] The term "fluid exchange opening" means any space or void through which a fluid may pass from one vessel to an adjacent vessel or another vessel in fluid communication with the one vessel.

[0083] The terms "individually comprise" in respect to a claimed element or elements mean that only one claimed element comprises each of the listed elements and not in combination with any other element named.

[0084] The terms "a compound comprising a phenol substituent" means any molecule comprising a phenyl group attached to a 4, 5, 6, or more-membered atomic ring comprising at least one carbon atom.

[0085] The term "ionomer" as used herein refers to any polymer comprising an ion. In some embodiments, the ionomer is a perfluorinated ionomer. In some embodiments, the ionomer comprises Formula I or a salt thereof.

Formula I:











Where $\text{X}_1=\text{F}-\text{O}-\text{CF}_2-\text{Y}$, F_2-SO_2 , or $\text{F}_2-\text{CF}_2-\text{CO}_2\text{CH}_3$

$\text{X}_2=\text{CF}_3$, or, if X_1 is F_2 , X_2 is null

Where $\text{Y}=\text{CF}_2-\text{SO}_2\text{F}$, $\text{CF}_2-\text{CF}-\text{SO}_2\text{F}$, or $\text{CF}_3-\text{CO}_2\text{CH}_3$

[0086] In some embodiments, the ionomer comprises one or a combination of:





[0088] As used herein, the term "functional fragment" means any portion of a disclosed polypeptide that is of a sufficient length to retain at least partial biological function that is similar to or substantially similar to the function of the wild-type polypeptide upon which the fragment is based. In some embodiments, a functional fragment of a polypeptide associated with the function of a metabolic enzyme is a polypeptide that comprises at least 70%, 75%, 80, 85, 90, 95, 96, 97, 98, or 99% sequence identity of any polypeptides disclosed herein and has sufficient length to retain at least partial binding affinity to one or a plurality of substrates that bind to the polypeptide. In some embodiments, the fragment is a fragment of any polypeptide disclosed herein and has a length of at least about 10, about 20, about 30, about 40, about 50, about 60, about 70, about 80, about 90, or about 100 contiguous amino acids. In some embodiments, the fragment is a fragment of any polypeptide disclosed herein and has a length of at least about 50 amino acids. In some embodiments, the fragment is a fragment of any polypeptide disclosed herein and has a length of at least about 100 amino acids. In some embodiments, the fragment is a fragment of any polypeptide disclosed herein and has a length of at least about 150 amino acids. In some embodiments, the fragment is a fragment of any polypeptide disclosed herein and has a length of at least about 200 amino acids. In some embodiments, the fragment is a fragment of any polypeptide disclosed herein and has a length of at least about 250 amino acids.

[0090] In some embodiments, the device, system, membrane, or vessel, may comprise any of the disclosed reagents or formula disclosed herein or any salt. Salts may be formed by reacting the free base, or a salt, enantiomer or racemate thereof, with one or more equivalents of the appropriate acid. In some embodiments, salts of the present disclosure refer to salts of the disclosed reagents or formula disclosed herein having at least one basic group or at least one basic radical. In some embodiments, salts of the present disclosure refer to salts of the disclosed reagents or formula disclosed herein having a free amino group, a free guanidino group, a pyrazinyl radical, or a pyridyl radical that forms acid addition salts. In some embodiments, salts of the present disclosure

refer to salts of the disclosed reagents or formula disclosed herein that are acid addition salts of the subject compounds with (for example) inorganic acids, such as hydrochloric acid, sulfuric acid or a phosphoric acid, or with suitable organic carboxylic or sulfonic acids, for example aliphatic mono-or di-carboxylic acids, such as trifluoroacetic acid, acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, fumaric acid, hydroxymaleic acid, malic acid, tartaric acid, citric acid or oxalic acid, or amino acids such as arginine or lysine, aromatic carboxylic acids, such as benzoic acid, 2-phenoxy-benzoic acid, 2-acetoxybenzoic acid, salicylic acid, 4-aminosalicylic acid, aromatic-aliphatic carboxylic acids, such as mandelic acid or cinnamic acid, heteroaromatic carboxylic acids, such as nicotinic acid or isonicotinic acid, aliphatic sulfonic acids, such as methane-, ethane- or 2-hydroxyethane-sulfonic acid, or aromatic sulfonic acids, for example benzene-, p-toluene- or naphthalene-2-sulfonic acid. When several basic groups are present mono-or poly-acid addition salts may be formed. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, for example, water, dioxane, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed in vacuo or by freeze drying. The reaction may also be a metathetical process or it may be carried out on an ion exchange resin. Salts according to the present disclosure may be found in their anhydrous form or as in hydrated crystalline form (i.e., complexed or crystallized with one or more molecules of water).

[0091] As used herein, the term "antibody" refers to any immunoglobulin, whether natural or wholly or partially synthetically produced. In some embodiments, an antibody is a complex comprised of 4 full-length polypeptide chains, each of which includes a variable region and a constant region, e.g., substantially of the structure of an antibody produced in nature by a B cell. In some embodiments, an antibody is a single chain. In some embodiments, an antibody is cameloid. In some embodiments, an antibody is an antibody fragment. In some embodiments, an antibody is chimeric. In some embodiments, an antibody is bi-specific. In some embodiments, an antibody is multi-specific. In some embodiments, an antibody is monoclonal. In some embodiments, an antibody is polyclonal. In some embodiments, an antibody is conjugated (*i.e.*, antibodies conjugated or fused to other proteins, radiolabels, cytotoxins). In some embodiments, an antibody is a human antibody. In some embodiments, an antibody is a mouse antibody. In some embodiments, an antibody is a rabbit antibody. In some embodiments, an antibody is a rat antibody. In some embodiments, an antibody is a donkey antibody. In some embodiments, the biosensor or system described herein comprises an antibody or plurality of antibodies.

[0092] *Characteristic:* As is used herein, the term "characteristic" refers to any detectable feature of a sample of bodily fluid that allows it to be distinguished from a comparable sample of bodily fluid. In some embodiments, a characteristic is an amount or identity of ammonia or ammonium ion in bodily fluid, in an environmental sample, or water sample. In some embodiments, a characteristic is an amount, sequence of, or modification of a amino acid. In some embodiments a characteristic is an amount of a carbohydrate. In some embodiments, a characteristic is an amount of a small molecule.

[0093] *Comparable:* As is used herein, the term "comparable" is used to refer to two entities that are sufficiently similar to permit comparison, but differing in at least one feature.

[0094] *Metabolic Enzyme:* As is used herein, the term "metabolic enzyme" means an enzyme responsible for catalysis of at least one step in the metabolic pathway of one or more amino acids. In some embodiments, the metabolic enzyme is phenylalanine dehydrogenase, glutamate dehydrogenase, respective functional fragments or a combination thereof or a fusion protein thereof.

[0095] As used herein the terms "metabolic disease" is any one of a group of disorders caused by a defect in an enzymatic step in the metabolic pathway of one or more amino acids or in a protein mediator necessary for transport of certain amino acids into or out of cells. In some embodiments, the metabolic disease is chosen from: Argininemia (ARG, arginase deficiency) Argininosuccinate acidemia (ASA, argininosuccinase) Citrullinemia type I (CIT-I, argininosuccinate synthetase) Citrullinemia type II (CIT-II, citrin deficiency) Defects of bipterin cofactor biosynthesis (BIOPT-BS) Defects of bipterin cofactor regeneration (BIOPT-RG) Homocystinuria (HCY, cystathionine beta synthase) Hyperphenylalaninemia (H-PHE) Hypermethioninemia (MET) Maple syrup urine disease (MSUD, branched-chain ketoacid dehydrogenase) Phenylketonuria (PKU, phenylalanine hydroxylase) Tyrosinemia type I (TYR-1, fumarylacetoacetate hydrolase), Tyrosinemia type II (TYR-II, tyrosine

aminotransferase), and Tyrosinemia type III (TYR-III, hydroxyphenylpyruvate dioxygenase) where the parenthetical phrases after each disease state represent an abbreviation for the disease accompanied by the enzyme that is generally defective in the subject suffering from the disease state.

[0096] *Polypeptide*: The term "polypeptide", as used herein, generally has its art-recognized meaning of a polymer of at least three amino acids. Those of ordinary skill in the art will appreciate that the term "polypeptide" is intended to be sufficiently general as to encompass not only polypeptides having the complete sequence recited herein, but also to encompass polypeptides that represent functional fragments (i.e., fragments retaining at least one activity) of such complete polypeptides. Moreover, those of ordinary skill in the art understand that protein sequences generally tolerate some substitution without destroying or significantly reducing activity. Thus, any polypeptide that retains activity and shares at least about 30-40% overall sequence identity, often greater than about 50%, 60%, 70%, 75%, 80%, or 85%, and further usually including at least one region of much higher identity, often greater than 90% or even 95%, 96%, 97%, 98%, or 99% in one or more highly conserved regions, usually encompassing at least 3-4 and often up to 20 or more amino acids, with another polypeptide of the same class, is encompassed within the relevant term "polypeptide" as used herein.

[0097] As used herein, the term "threshold value" is the concentration of ammonia or ammonium ion or amino acid in a sample of bodily fluid that indicates whether the amount of ammonia or ammonium ion or amino acid in the sample is considered abnormally high or low resulting in a diagnosis or suspected diagnosis of a particular disorder, such as a metabolic disease. For instance, in the case of a blood sample, known threshold values for certain aminoacidopathies are indicated in Table 1 below:

Table 1: Aminoacidopathies and their associated amino acid markers detectable in a sample

Disorder	Marker	Abnormal Range
ARG	Arginine	> 100 umol/L
ASA	Argininosuccinic acid	> 4.0 umol/L
	ASA/Arg	> 0.75
CIT-I and CIT-II	Citrulline	> 60 umol/L
	Cit/Tyr	> 1.0
	Cit/Arg	> 6.0
HCY and MET	Methionine	> 70 umol/L
	Met/Phe	> 1.2
MSUD	Leucine	> 250 umol/L
	Valine	> 250 umol/L
	Leu/Phe	> 4.0
	Val/Phe	> 3.5
PKU, H-PHE	Phenylalanine	> 130 umol/L
BIOPT-BS and BIOPT-RG	Phe/Tyr	> 2.0
TYR-I, TYR-II, and TYR-III	Tyrosine	> 250 umol/L

[0098] In some embodiments, information about a threshold value or reference sample of bodily fluid is obtained prior to or simultaneously with information about an experimental sample of bodily fluid. In some embodiments, information about a reference cell or cell type is historical. In some embodiments, information about a threshold value or reference sample of bodily fluid is stored for example in a computer-readable storage medium. In some embodiments, comparison of a particular concentration value with a threshold value or reference sample of bodily fluid differentiates the concentration values of ammonia in an experimental sample of bodily fluid with the threshold values thereby allowing a comparison that results in diagnosing a subject with one or more metabolic diseases or a change in severity of one or more metabolic diseases.

[0099] *Reference electrode*: As will be understood from context, a reference electrode or control electrode is an

electrically conductive support such as an electrode placed in a circuit with an at least one electrically conductive support comprising hydrogel and/or immobilized enzymes disclosed herein, to permit a relevant comparison of voltage difference between the reference or control electrode and the at least one electrically conductive support comprising hydrogel and/or immobilized enzymes disclosed herein.

[0100] *Sample:* As used herein, the term "sample" refers to a biological sample obtained or derived from a source of interest, as described herein. In some embodiments, a source of interest comprises an organism, such as an animal or human. In some embodiments, a biological sample comprises biological tissue or fluid. In some embodiments, a biological sample may be or comprise bone marrow; blood; blood cells; ascites; tissue or fine needle biopsy samples; cell-containing body fluids; free floating nucleic acids; sputum; saliva; urine; cerebrospinal fluid, peritoneal fluid; pleural fluid; feces; lymph; gynecological fluids; skin swabs; vaginal swabs; oral swabs; nasal swabs; washings or lavages such as a ductal lavages or broncheoalveolar lavages; aspirates; scrapings; bone marrow specimens; tissue biopsy specimens; surgical specimens; feces, other body fluids, secretions, and/or excretions; and/or cells therefrom, *etc.* In some embodiments, a biological sample is or comprises bodily fluid. In some embodiments, a sample is a "primary sample" obtained directly from a source of interest by any appropriate means. For example, in some embodiments, a primary biological sample is obtained by methods selected from the group consisting of biopsy (*e.g.*, fine needle aspiration or tissue biopsy), surgery, collection of body fluid (*e.g.*, blood, lymph, feces *etc.*), *etc.* In some embodiments, as will be clear from context, the term "sample" refers to a preparation that is obtained by processing (*e.g.*, by removing one or more components of and/or by adding one or more agents to) a primary sample. For example, filtering using a semi-permeable membrane. Such a "processed sample" may comprise, for example nucleic acids or proteins extracted from a sample or obtained by subjecting a primary sample to techniques such as amplification or reverse transcription of mRNA, isolation and/or purification of certain components, *etc.* in some embodiments, the methods disclosed herein do not comprise a processed sample.

[0101] The system, test strip, device, biosensor, and/or catridge comprises a concentration of any one or combination of the reagents disclosed on pages 78-84 of this disclosure. Notwithstanding any indication to the contrary, the invention always relates to a test strip.

Table 2

Enzyme	Gene Sequence	Accession Numbers
Phenylalanine Dehydrogenase	ATGGAAATCTTCGAGGAAATCAAACGGCGGGGACACGAGCAA	AEW06037.1
	ATTCTGTTCAATTATGATCGGGCTTCCG	YP_005257709.1
	GTTTGAAAGCAATTATCGCCATTCACAATACTACGTTGGGGC	AEH47572.1
	CGGCGTTGGGCGGGTGCCGAATGTTACC	YP_004587653.1
	GTATCAAACGGAAGAGGCGGCCCTCGAGGATGCGCTGCGGTT	YP_004581770.1
	GTCGGAAGGGATGACCTATAAAGCGGCC	AEH07849.1
	GCCGCCGGGCTCGATTTCGGCGGGGGCAAACGGTGATTATC	ACF96938.1
	GGGGATCCGATGAAAGACAAGTCCGAGG	YP_007466124.1
	CCCTGTTTCGTGCGCTCGGGCGTTTTATCGAGACCTTGAAAG	EZP75760.1
	GCCGTTACCTTACGGGAGAAGACGTAGG	AGT95551.1
	AACCAACGAAGAAGATTTTGTCTGGGCTCGTCGGGAAACCCG	EWG09095.1
	TTATGTTGTCTGGATTGCCGCCGGCTTAT	YP_008456272.1
	GGCGGGTCCGGCGATACGGGTGACAATACCGCGCGCGCGTC	EME23486.1
	ATTCAAGCGATGCGCGCCGCGTTGATGC	EJS99791.1
	ACCGGTACGGTTCGCCGGATCTCCAGGGCCGGCGGATTGCCG	EIT85807.1
	TCCAAGGGCTGGGCAAAGTAGGCTATCA	AAA22646.1
	TGTGGCGCGACGGGCCATCGAGGCCGGCGCTCGAGTGATTGC	EDL64419.1
	GGCCGATATCAATCCGCATGTAGTCGGC	
	CGAGTGGCGTCCGCTTGGGGGATTGAAGCCACCGATCCGTGG	
	GCTGTGGTGGAACCCCTGCGATATTT	
	TCGCCCCCTGTGCGTTGGGTAACGTCATTACGGAACGGACCG	
	TGTCCGCCCTCCAATGTCAGGTGGTGGC	
	CGGTTCGGCCAACAAATCAGCTGGCGGATGATCGACTGGCCGA	
	TGATTTAGCTGCCCGCGGCATTCTCTAT	
	GCGCCGATTTTATTGCGAATGCCGGCGGATTGATTTCAGGTG	
	GCGGATGAAATTCGGGGATATCATGAAG	
	AACGGGTCCGTCATCAAATAGACGGGATTTATGACGTCCTGC	
	TCGAGATTTTTCGGAAGGCGGACGCCTC	
	CGGCCGATCAACCGTGGCGGTTGCGGTAGACGAGGCGCGTCG	
	CCGTTTGGACACCATTCAGGCCATCCAC	

Enzyme	Gene Sequence	Accession Numbers
	CGCCTGTACGGATCATAG	EAR66050.1
		BAA08816.1
Phenylalanine Ammonia-Lyase	CTGCAGGTCAACGGATCATATTCTACACATATATAATGCACTCCAATTGACATAATACATAACGTGACATATGATACACTTATTAATATTAATTGTCACTTTACACTTCACATATTAAATACTCTCGTATGAATGCAA TTTGAAACATTTTAAATTAAATTGATTGATATATATTGAACAAAACCTAACAAAAATGCACCCTCTTGG TTCACAAAGAACTTTCTTCTATTTCTCACTTATTTCTGCTAGTGTCTTTCCTATTCAAAGCCATCATTT CCATCAACCTTCACAATACCATGTTTAAAAAGTCATTAAAAATCAATTTT TTAAATAGAAAAACAAGA AGATGGAAACCACTTGGTTGGTACTATATATTTAGTTGTTAAGTTTGACTCATACCGTGTATTGACCAAT ATAAATAAAATCTTATTTCAAATAAATTCAAAAGTTCAATAAATATATATTCGTTCATAACTTATAATAA AATTGATTACACATAGTCCTCCCCCATTCACTTTTACTGATCAATTATTTCTAAAATACTATTATTACTTT TACTTGTTACTTTTAATAAATTAAGAAAATATAATACTCCCTTCGTTTTTAAAAAATACCTAGTTTGAC TTGAAACGGAGTTTAAATAAAGAAAGAGACTTGTTAATCTTGTGATTCTAAATTAAAGCTATGTCAAAT GTACCAAAACGTCCTTTAATCTTGTGGTCTTAACATGTCACATGAAAAATTAAAGTCTTCAAAAAA GAAAGGGGCTCAATGTCATTCTTTTTTAAACAGACTAAAAAGAAATAAACTCATTCTTTTGAAACGGAG AGAGTAATCTTTCCACGTTTACTCATCAATATTAAATATTATTCTCTAGATCATCTATAAGATCTAA TAGTGGACATCAATTAATACCTATGTCACCTATTATTATTTTAATAATTGTATCAAGTCAAATAATAACA AGTAAAAAGGAGTACCTACTATTAACTCTCAACAACCACAATTTACTAGTTTTTTCTAGCAACCCCT CTCACATACTTCACCATTTACTGGTTTTTCTCTAGCAACCCCTCTCACA TATTTTGTCTTACCAACCATC ATTTGTCTCTATATATACTCACCACATGATAGATACATATATATACCA CAACCAAAACAAAAGGTTTT ATAAGTTCACAACATTTTTTATATACATACAAATAAACCTAACCATTTTCTCTTCACTAAAATTTCTTC ATTACAAATCTAACAATTTACTTGATCCAATGGCACCATCAATTGCACAA AATGGACACATTAATGGAGA AGTAGCTATGGATTGTGCAAGAAATCAATCAATGATCCATTGAATTGGG AAATGGCTGCTGATTCTTTA AGAGGCAGCCATTTGGATGAAGTGAAAAAGATGGTGGATGAATTTAGAAA GCCAATTGTGAACTTGGGG GTGAACTCTGTCAAGTTGCACAAGTTGCATCCATTGCAAAATGTTGATGAC AAAAGTAACGGGTAAAGT GGAACTTTCTGAAAGTGCAAGGGCTGGTGTGAAAGCTAGTAGTGATTGGG TTATGGATAGTATGAGTAAA GGTACAGATAGTTATGGTGTACTGCTGGATTGGAGCAACATCTCATAG AAGAACAAAAAATGGTGGTG CTCTTCAAAAAGAACTTATTAGGTAAACAACTATTTTTTTTCTGTTATAT ATACTAACAAATGTAAAGAAT TTAATATCTCTTTGTTATATATACTAACAAATGTAAAAAATTTAATATTTT	AAA34179.2
		ADR78835.1
		AAA99500.1
		AAC18871.1
		AAC18870.1
		AAA33805.1
		AIC66437.1
		AGY49231.1
		AEW43005.1
		AFP24940.1
		AER58180.1
		ADD12041.1
		AEE81750.1
		AAP59440.1
		AAP59439.1
		AAP59438.1
		ACG80829.1
		ACG80828.1
		ACG56648.1
		ACG56647.1
	TTTGTATATATACTAACAA TGTAAGAATTTAATATTTTTTTGTTATACATAGCTTATCGACTACTTAA GTGCTCCATTGATAAGATT TTTTTTTGTTTTACGCGAAGGGGATTCCGGATGAATTGAGTTAAAAATGTG ATCTTAATGAATTATCATAT TTTTTTGTAGGTTCTTGAATGCTGGAGTTTTTGGTAATGGAATAGAATCA TTTCACACATTGCCACATTC AGCAACAAGGGCAGCTATGCTTGTTAGGATCAACACTCTGCTTCAAGGCT ACTCTGGCATTAGATTTGAG ATCTTGGAAAGCAATCACTAAGTTGATCAATAGCAACATCACCCCGTGTTT CCCTCTCCGTCCACCATCA CTGCCTCGGGTGATCTCGTCCCTTTGTCTATATTGCTGGTTTGCTCACT GGCAGACCTAATCCAAGGC TGTTGGACCAATGGTGAGAACTTAATGCTGAGGAAGCTTCTGCGTGG CTGGTATTAGTGGTGGATTT TTCGAGTTGCAGCCTAAGGAAGGACTTGCACCTTGTGAATGGCAGCAGT TGGTTCTGCTATGGCATCAA TAGTCCTGTTTGAGTCCAATATCTTTGCTGTTATGCTGAAGTTTATCA GCGATTTTTACTGAAGTGAT GAACGGAAAGCCCGAATTCACTGACTATTTGACACACAAGTTGAAGCATC ACCCTGGTCAGATTGAGGCT GCTGCTATTATGGAACACATTTTGGATGGAAGCTCTTATGTGAAGGTAGC TCAGAAGCTCCATGAAATGG ATCCTCTTCAAAAACCAAAGCAAGATCGTTATGCTCTCCGAACATCTCCA CAATGGCTTGGACCTCAGAT TCAACGTCATTCCTGCTGCAACGTAACGATGCAACGACGACGACGATCAACGTCAC	

Enzyme	Gene Sequence	Accession Numbers
	TGAAATCATTCGTCCTGCAACCTAAGATGATCGAGAGGCGAGATTTACTCAG TGAACGACAATCCATTGATC GATGTTTCAAGAAACAAGGCCTTACATGGTGGCAACTTCCAAGGAACCCC TATTGGTGTCTCCATGGATA ATACAAGATTGGCCCTTGCATCAATTGGTAAATTGATGTTTGCCCAATTCTCAGAGCTTGTCAACGACTA TTACAACAACGGGTTGCCATCTAATCTGACAGCAGGAAGGAATCCAAGCTTGGACTATGGTTTCAAGGGC GCTGAAATCGCGATGGCTTCTTACTGCTCGGAACCTCAATTCTTGGCAAA TCCAGTGACTAACCATGTCT AAAGTGCTGAGCAACACAACCAAGATGTGAATTCCTTGGGCTTAATTTCA GCCAGGAAAACAGCTAAGGC TGTTGATATCTTGAAGATAATGTCATCAACCTATCTCGTGGCTCTTTGCC AAGCTATTGACTTACGACAT TTGGAGGAAAACCTTGAAGAGTGTTGTCAAGAACACAGTTAGCCAAGTAGCTAAGAGAACTTTGACAAATGG GTGCTAATGGTGAACCTCATCCAGCAAGATTCAGCGAAAAAGAATTGCTTCGAGTCGTGGATAGAGAATA CTTGTTTGCCATATGCTGATGATCCCTGCAGCTCCAACCTACCCTTTGATGC AGAAGCTGAGACAAGTCCTT GTTGATCAAGCAATGAAGAATGGTGAAAGTGAGAAGAATGTCAACAGCTCAATCTTCCAAAAGATTGGAG CTTTCGAGGACGAATTAATCGCTGTGTTGCCTAAAGAAGTTGAGAGTGTA AGAGCTGTTTTTGAAAGTGG CAACCCCTTAATTCGTAAACAGGATCACAGAATGCAGATCATATCCATTGT ACAGGTTGGTGAGAGAAGAA	
	CTTGGAACAGAATTGTTGACGGGTGAAAAAGTTCGATCACCTGGTGAGGA GATTGATAAAGTGTTTACAG CAATATGTAATGGACAGATTATTGATCCATTGTTGGAGTGCTGAAGAGCTGGAATGGTGCTCCTCTTCC AATCTGCTAAATGTGTTATTCTTTCAAGTTCCTTTTTTTGTACCTTTTAGTGAATTACTAGAATTATAATG ATGTTATGAACTTATATTAAAAAAAATATTTTTGACTATAAAATTTAGTTTGTTATTGAAATTAAAGG CTCAATCTGTGTTCTTTCCTTCTGTTATCTGAATATTATAAGAATTCAAGTAATCTTTTAGCTTTGTGAA CATGATGACATGCTTCTT	
Histidine Ammonia- Lyase	ATGATCACGCTTACCCCCGGCCACCTGACCCTCCCGCAACTGCGCCAGATCGCGCGCGAGCCCGTGCAAGC TGACGCTGGATCCGGCCAGCTTCGCGAAGATCGACGCGGGCGCGAAGGCCGTGTCCGACATCGCCGCGAA GGGCGAGCCGGCGTACGGCATCAACACGGGGCTTCGGTCGTCTGGCCAGCACGATATCCCGCACGATCAG CTCGAATTGCTGCAGAAGAACCTCGTGCTGTGCGATGCAGTCGGTGTCGGCGAGCCGATGGCGCGTTCTGT CGGTGCGTCTGCTGATCGCGCTGAAGCTGTGCGAGCCTCGGCCGCGGCCATTCGGGCATTGCGCGCGAAGT GATGGACGCGCTGATCAAGCTGTTCAACGCCGACGTGCTGCCGCTGATTCCGGTGAAAGGGCTCGGTGCGC GCATCGGGCGACCTCGCGCCGCTCGCGCACATGTGGCCGTGCTGCTCGGCGTCGGCGAAGTGTTTATTCT GCGGCGAGCGCGGAGCGCGGTGGACGGGTTGCGCGTCGCGGGCCTCGCGCCGCTGACGCTGCAGGCGAA GGAAGGCCTCGCGCTGCTGAACGGTACGCAGGCGTCGACGGCGCTCGCGCTCGACAACCTGTTGCGGATC GAAGACCTGTACCGCACGGCGCTCGTCGCGGCGCGCTGTGGTTCGATGCGCGGCCGGCTCGGTGAAGC CGTTGACGCGCGCATCCACGAACTGCGCGGCCATCGCGGCCAGATCGATGCGGCGGCCGCTATCGCGA GCTGCTCGAAGGCTCGGCGATCAACCTCTCGCATCGCGACTGCGGCAAGGTGAGGATCCGTACAGCCTG CGCTGCCAGCCGCAGGTGATGGGCGCGTGCTGGACCAGATGCGTCATGCGCCGACGTGCTGCTCGT AGGCGAACGCGGTATCGGACAACCCGCTGATCTTCCCGGATACCGGCGAA GTGCTGTGGGCGGCAATTT CCATGCGGAGCCCGTCGCGTTTCGCGGCCGACAACCTCGCGCTCGCGGCTGCGGAAATCGGCGCGCTGGCC GAGCGCCGCATCGCGCTGCTGATCGACGCGACGCTGTGCGGCGCTGCGGCCGCTGCGGCTGCGG GCGTGAAGGCTTCATGATTGCGCACGTGACGGCAGCTGCGCTCGCATCGGAGACAAGACGCTCGC GCATCCGGCGTCGGTTCGATTGCTGCGGACCTCGGCGAACCGGAAGACC	BAG44062.1 YP_005225923.1 CDF52938.1 ABR76232.1 AAL19728.1 AEW60321.1 AEW51583.1 ABQ54772.1 AAX64695.1 AAU27462.1 WP_021000087.1 YP_005185682.1 YP_001250118.1 EFC47317.1 AAH89809.1 BAH62483.1 XP_00268006

Enzyme	Gene Sequence	Accession Numbers
	ACGTGTCGATGGCGACGTTTC GCGGCACGCAAGCTGGCCGACATCGCCGACAACACGAAGCACATCCTCGC GATCGAACTGCTCGCGGCCG CGCAGGGCGTCGATCTGCGCGAGAACGAGACGAGCCCGAAGCTCGCGGAA	1.1
		AAO73411.1
		CAI79696.1
		CAI79696.1
Tyrosine Ammonia- Lyase	GTGATGAAGACGATTTCGCAG CAAGGTCGCGCATTACGAGCTCGACCACTACTTTGCGCCGGACATCGCCG TGATCGCGAAGCTCGTCGTC GAGCGCGCGTTTCGCGAAGCACTGCCCGTTGCGCTTCGCATCGGAGCAGTA A GTGACGCAGGTCGTTGGAACGTCAGGCTGATCGGCTCAGCAGCAGGGAGTA CCTGGCCCGGGTCGTGCGCA GCGCCGGGTGGGACGCCGGTCTCACCTCGTGCAACCGACGAGGAGATCGTC CGGATGGGCGCGAGCGCGC CACCATCGAGGAGTACCTGAAGTCCGACAAGCCCATCTACGGCCTGACGC AGGGCTTCGGTCCGCTGGTG CTGTTGACGCGCGACTCGGAGCTGGAGCAGGGCGGCTCGCTGATCTCGCA CCTGGGCACCGGCCAGGGCG CGCCACTGGCCCCGGAGGTGTGCGGGCTGATCCTCTGGCTGCGCATCCAG AACATGCGCAAGGGGTACTC GGCGGTCTCGCCGGTGTCTGGCAGAAGCTCGCCGACCTGTGGAACAAGG GGTTCACCCCGCGATCCCC CGGCACGGCACGCTCAGCGCGAGCGGCGACCTGCAACCGCTGGCGCACGC CGCGCTCGCCTTCACCGGTG TCGGCGAGGCGTGGACCCGGGACGCCGACGGCCGGTGGTCCACCGTGCCG GCCGTGGACGCGCTCGCCGC GCTGGGGGCGGAGCCGTTGACTGGCCGGTGCGCGAGGCGCTGGCGTTG TCAACGGGACCGGCGCGAGC CTCGCGGTGGCTGTGCTCAACCACCGGTCCGCCCTGCGGCTGGTCCGCGC CTGCGCCGTGCTCTCCGCGC GGCTGGCGACCTGCTGGGGGCCAATCCCGAGCACTACGACGTGGGGCAC GGTGTGCGCGCGGCGCAGGT CGGTGAGCTGACCGCGGCGAGTGGATCCGGCAGGGGCTGCCCGGGGCA TGGTGGCGGACGGCAGCCGC CCGCTCCAGGAGCCGTACAGCCTGCGGTGCGCGCCGCGAGGTGCTCGGCGC GGTGCTCGACCAGCTCGACG GCGCGGGCGACGTGCTGGCGCGGGAGGTGACGGCTGCCAGGACAACCCG ATCACCTACGAGGGCGAGCT GCTGCACGGCGGCAACTTCCACGCCATGCCGGTGGGTTTCGCCTCCGACC AGATCGGGTTGGCCATGCAC ATGGCCGCCTACCTGGCCGAGCGCCAGCTGGGTCTGCTGGTCAGCCCGGT GACCAACGGCGACCTGCCGC CCATGCTCACCCCGCGCGCCGGGCGCGGTGCCGGGCTGGCCGGGGTGCAG ATCAGCGCGACCTCGTTGCT CTCGCGGATCCGGCAGCTGGTGTTCGCCGCCCTCGCTGACCACCCTGCCGA CCAACGGCTGGAACCAGGAC CACGTGCCGATGGCGCTCAACGGGGCGAACTCGGTGTTGAGGCGTTGGA GCTCGGCTGGCTGACGGTCG GGTCGCTGGCGGTGGGCGTCGCGCAGCTCGCGGCCATGACCGGCCACGCC GCGGAGGGCGTCTGGGCGGA GCTGGCCGGGATCTGCCCGCCGCTGGACGCCGACCGCCGCTGGGCGCCG AGGTGCGCGCCGCGCGCGAC CTGCTGTCCGCGCACGCGGACCAACTGCTCGTCGACGAGGCAGACGGGAA GGATTTCGGATGA	YP_00703999 9.1
		Q8GMG0.1
		WP_0151032 37.1
		CCH33126.1
		AGZ04575.1
		GAK34477.1
		AIG26365.1
		WP_0308142 63.1
		WP_0305926 22.1
		WP_0305838 02.1
		WP_0302258 85.1
		WP_0301070 56.1
		WP_0102616 15.1
		WP_0090658 11.1
		WP_0290439 04.1
		WP_0290276 07.1
		WP_0290256 70.1
		WP_0290239 88.1
		WP_0290202 80.1
		WP_0286735 81.1
Glutamate Dehydrogenase	ATGTCAGCAAAGCAAGTCTCGAAAGATGAAGAAAAAGAAGCTCTTAACTT ATTTCTGTCTACCAAACAA TCATTAAAGGAAGCCCTTCGGAAGCTGGGTTATCCGGGAGATATGTATGAA CTCATGAAAGAGCCCGCAGAG	P39633.3
		KEG08275.1
		NP_00123385

Enzyme	Gene Sequence	Accession Numbers
	AATGCTCACTGTCCGCATTCCGGTCAAAATGGACAATGGGAGCGTCAAAG TGTTCAAGGCTACCGGTCA CAGCACAAATGATGCTGTCCGTCCGACAAAGGGGGCGTTCGCTTCCATCC AGAAGTTAATGAAGAGGAAG TAAAGGCATTATCCATTGGATGACGCTCAAATGCCGGATTGCCAATCTT CCTTACGGCGCGGGAAGGG CGGTATTATTTGTGATCCGCGGACAATGTCATTTGGAGAACTGGAAAGGC TGAGCAGGGGGTATGTCCGT GCCATCAGCCAGATCGTCCGTCCGACAAAGGATATTCCAGCTCCCGATGT GTACACCAATTCCGAGATTA TGGCGTGGATGATGGATGAGTACAGCCGGCTGCGGGAATTCGATTCTCCG GGCTTTATTACAGGTAAACC GCTTGTTTTGGGAGGATCGCAAGGACGGGAAACAGCGACGGCACAGGGCG TCACGATTTGTATTGAAGAG GCGGTGAAGAAAAAGGGATCAAGCTGCAAAACGCGCGCATCATCATACA GGGCTTTGGAAACGCGGGTA GCTTCCTGGCCAAATTCATGCACGATGCGGGCGCGAAGGTGATCGGGATT TCTGATGCCAATGGCGGGCT CTACAACCCAGACGGCCTTGATATCCCTTATTTGCTCGATAAACGGGACA GCTTTGGTATGGTCACCAAT TTATTTACTGACGTCATCACAAATGAGGAGCTGCTTGAAAAGGATTGCGA TATTTTAGTGCCTGCCGCGA TCTCCAATCAAATCACAGCCAAAAACGCACATAACATTTCAGGCGTCAATC GTCGTTGAAGCGGCGAAACGG CCCGACAACCATTGATGCCACTAAGATCCTGAATGAAAGAGGCGTGCTGC TTGTGCCGATATCCTAGCG AGTGCCGGCGGCGTCACGGTTTCTTATTTGAATGGGTGCAAAACAACCA AGGATATTATTGGTCGGAAG AAGAGGTTGCAGAAAACTGAGAAGCGTCATGGTCAGCTCGTTCGAAACA ATTTATCAAACAGCGGCAAC ACATAAAGTGGATATGCGTTTGGCGGCTTACATGACGGGCATCAGAAAT CGGCAGAAGCATCGCGTTTC CGCGATGGGTCTAA	0.1
		NP_00126803
		9.1
		AEW04907.1
		YP_00716125
		5.1
		YP_00525657
		9.1
		YP_00493265
		2.1
		YP_00444244
		4.1
		YP_00441234
		8.1
		YP_00441098
		6.1
		YP_00437273
		1.1
		YP_00436766
		7.1
		YP_00436636
		6.1
		YP_00434396
		8.1
		YP_00434335
		6.1
		YP_00426176
		6.1
		YP_00427038
		2.1
		YP_00409996
		1.1
		YP_00396781
		1.1
Glutamate Ammonia- Lyase	ATGTCCATCAAAGACGCTGTAAAACTGATTGAAGAAAGCGAAGCCCGCTT TGTCGATTTGCGCTTTACCG ATACCAAAGGCAAGCAGCACCACCTTACCGTGCCTGCGCGCATCGTGTTG GAAGACCCCGAAGAGTGGTT CGAAAACGGACAGGCGTTTGACGGTTCGTCCATCGGCGGCTGGAAAGGCA TTCAGGCTTCCGATATGCAG CTTCGCCCCGATCCCGCCACGGCGTTTATCGATCCTTTTTATGATGATGT TACCGTCGTCATTACCTGCG ACGTTATCGATCCCGCCGACGGTCAGGGTTACGACCGCGACCCGCGCTCC ATCGCACGCCGCGCGGAAGC CTATTTGAAATCTTCCGGTATCGGCGACACGGCATACTTCGGTCCCGAAC CCGAGTTTTTCGTCTTCGAC GGCGTAGAATTTGAAACCGATATGCACAAAACCCGTTACGAAATCACGTC	CBX22311.1
	CGAAAGCGGCGCATGGGCCA GCGGCCTGCATATGGACGGTCAAAACACCGGCCACCGCCCTGCCGTCAA GGCGGTTACGCGCCCGTCGC GCCGATTGACTGCGGTGAGGATTTGCGTTCGCGGATGGTAAACATTTTGG AAGGACTCGGCATCGAAGTC GAAGTGCACCACAGCGAAGTCGGTACCGGCAGCCAAATGGAAATCGGCAC	

Enzyme	Gene Sequence	Accession Numbers
Threonine Dehydrogenase	GCGCTTCGCCACCTTGGTCA AACGCGCCGACCAAACCCAAGACATGAAATATGTGATTCAAAATGTGCGC CACAACTTCGGCAAAACCGC CACCTTCATGCCCAAACCCATTATGGGCGACAACGGCAGCGGTATGCACG TTCACCAATCCATCTGGAAA GACGGTCAAAACCTGTTTCGCAGGCGACGGCTATGCCGGCTTGAGCGACAC CGCGCTCTACTACATCGGCG GCATCATCAAACACGCCAAAGCCCTGAACGCGATTACCAATCCGTCCACC AACTCCTACAAACGCCTTGT GCCGCACTTTGAAGCGCCGACCAAACCTGGCATATTCCGCCAAAAACCGTT CCGCTTCCATCCGTATTCCG TCTGTGAACAGCAGCAAGGCGCGCCGCATCGAAGCGCGTTTCCCCGACCC GACCGCCAACCCGTACTTGG CGTTCGCTGCCCTGCTGATGGCGGGTTTGGACGGCATTCAAAAACAAAATC CATCCGGGCGATCCTGCCGA TAAAAATCTCTACGACCTGCCGCCGGAAGAAGACGCGCTCGTCCCGACCG TTTGCGCTTCTTTAGAAGAA GCCCTCGCCGCGCTCAAAGCCGACCACGAATTCTCTTTACGCGGCGGGGT GTTCAGCAAAGACTGGATCG ACAGCTACATCGCCTTTAAAGAGGAAGATGTCCGCCGCATCCGTATGGCG CCGCATCCGCTGGAATTTGA AATGTATTACAGCCTGTAA	
	AGGAGGTGTTTTAATAATGAAAGGTTTTGCAATGCTCAGTATCGGTAAAG TCGGTTGGATTGAAAAAGAA AAGCCTACTCCCGGCCCTTTTGACGCTATTGTAAGACCTCTAGCTGTGGC CCCTTGCACTTCGGACGTTT ATACCGTTTTTTGAAGGTGCTATTGGCGAAAGACATAACATGATACTCGGT CACGAAGCTGTAGGTGAAGT AGTTGAAGTAGGTAGTGAGGTAAAAGATTTTAAACCTGGTGATCGCGTTG TGGTACCAGCTATTACCCCT GATTGGCGAACCTCTGAAGTGCAAAGAGGATATCACCAACACTCTGGTGG AATGCTGGCAGGCTGGAAAT TTTCGAATATAAAAGATGGTGTTTTTGGTGAATTTTTTCATGTGAACGAT GCTGATATGAATTTAGCACA TCTGCCTAAGGAAATTCCATTGGAAGCTGCAGTTATGATTCCCGATATGA TGACTACTGGCTTTCACGGA GCCGAAGTGGCAGATATAGAATTAGGTGCGACGGTAGCGGTTTTGGGTAT TGGCCCAGTAGGTCTTATGG CAGTCGCTGGTGCCAAATTGCGGGGTGCTGGAAGGATTATCGCAGTAGGC AGTAGACCAGTTTGTGTAGA TGCTGCAAATACTATGGAGCTACTGATATTGTAACTATAAAGATGGTC CTATCGACAGTCAGATTATG GATTTAACGGAAGGCAAAGGTGTTGATGCTGCCATCATCGCTGGAGGAAA TGTTGACATCATGGCTACAG	NP_622353.1 EPX86072.1 AFT82159.1 YP_00679615 8.1 EJZ15419.1 YP_00172763 0.1 ACA82186.1 AGZ44086.1 AEB44998.1 YP_00873713 9.1 EPX87740.1 YP_00440559 8.1 BAN60779.1 EPE39095.1 EPC57128.1 EME23086.1 ACI75705.1 ACI75704.1 ACI75703.1 ACI75702.1
	CAGTTAAGATTGTTAAACCTGGTGGCACCATCGCTAATGTAAATTACTTT GGCGAAGGAGATGTTTTGCC TGTTCCCTCGTCTTGAATGGGGTTGCGGCATGGCTCATAAACTATAAAAG GCGGGCTATGCCCCGGTGGA CGTCTAAGAATGGAAAGACTGATTGACCTTGTTGTTTATAAGCGTGTCTGA TCCTTCTAAGCTCGTCACTC ACGTTTTCCGGGATTTGACAATATTGAAAAAGCCTTTATGTTGATGAAA GACAAACCAAAGACCTAAT CAAACCTGTTGTAATATTAGCATAA	

Enzyme	Gene Sequence	Accession Numbers
Threonine Ammonia-Lyase	ATGGCTGACTCGCAACCCCTGTCCGGTACCCCGGAAGGTGCCGAATATTT AAGAGCGGTGCTGCGCGCGC CGGTCTACGAAGCGGCGCAGGTCACGCCGCTACAGAAAATGGAAAACTG TCGTGCGGTCTCGATAACGT GATTCTGGTGAAGCGCGAAGATCGCCAGCCAGTTCATAGCTTTAAGTTGC GCGGCGCATACGCCATGATG GCGGGCCTGACGGAAGAACAAGCACACGGCGTGATTACCGCTTCTGC AGGTAACCACGCGCAGGGCG TCGCGTTTTCTTCCGCACGGTTAGGCGTGAAGGCGCTGATCGTCATGCCA ACCGCCACCGCCGATATCAA AGTTGATGCGGTGCGCGGCTTTGGCGGCGAAGTGCTGCTTACGGGCGCAA ATTTCGATGAAGCGAAAGCG AAAGCGATCGAACTGTACAGCAGCAGGGTTTACCTGGGTACCGCCGTT CGATCATCCGATGGTGATCG CCGGGCAAGGCACGCTGGCGCTGGAAGTCTCCAGCAGGACGCCCATCTC GACCGCGTATTTGTACCGGT CGGCGGCGGCGGTCTGGCAGCGGGTGTGGCGGTGCTGATCAAACAAGTGA TGCCGCAAATCAAAGTAATC GCCGTGGAAGCGGAAGATTCCGCCTGCCTGAAAGCGGCGCTGGATGCGGG TCATCCCGTTGATCTGCCCC GCGTGGGGCTGTTTGCTGAAGGCGTCGCGGTAAAACGCATCGGCGATGAA ACCTTCCGTTTGTGCCAGGA GTATCTTGACGACATCATCACCGTCGATAGCGATGCCATCTGTGCGGCGA TGAAAGATCTGTTCTGAAGAT GTGCGCGCGGTGGCGGAACCTTCCGGCGCGCTGGCGCTGGCGGGGATGAA AAAATACATCGCCAGCACA ACATTCGCGGTGAACGGCTGGCGCATATTCTTTCCGGTGCTAACGTGAAC TTTCACGGTCTGCGCTACGT CTCGGAACGCTGCGAACTGGGCGAACAGCGTGAAGCGTTGTTGGCGGTGA CCATTCGGAAGAAAAAGGC AGCTTCCTCAAATTCTGCCAACTGCTTGGCGGGCGTTCGGTCACCGAGTT CAACTACCGTTTTTGCCGATG CCAAAAACGCCTGCATCTTTGTGCGCGTGCGCTTAAGCCGTGGCCTCGAA GAGCGCAAAGAAATTTTGA GATGCTCAACGACGGTGGCTACAGCGTGTTGATCTCTCCGACGACGAAA TGGCGAAGCTGCATGTGCGC TATATGGTTGGCGGGCGTCCATCGCATCCGTTGCAGGAACGCCTATACAG CTTCGAATTCCCGGAATCAC CGGGCGCGCTGCTGCGCTTCCTCAACACGCTGGGTACGCACTGGAACATC TCGCTGTTCCATTATCGCAG CCACGGTACCGACTACGGGCGCGTACTGGCGGCGTTTCGAGCTTGGCGATC	EGP22802.1
		AIL15845.1
		KFJ14411.1
		B22317
		ESE06785.1
		ESD87895.1
		ESD77040.1
		ESD56952.1
		ESD26867.1
		ESD18649.1
		ESC98561.1
		ESA95751.1
		ESA86931.1
		ESA78951.1
		ESA72735.1
		ESA67809.1
		ERL21545.1
		ERK40933.1
		ERJ97484.1
		ERH28800.1
	ATGAACCGGATTTTGAAACC CGGTTGAATGAACTGGGCTACGATTGCCACGACGAAACCAATAACCGGC GTTCAGGTTCTTTTGGCGG GTTAG	
Serine Dehydrogenase	ATGAGCGGTACCATCCTCATCACCGGCGCCACGTCCGGCTTCGGACAGGC CACGGCGCGGCGTTTCGTCA AGGAAGGCTGGAAGGTCATCGGCACAGGTCGGCGGGCGGAACGGCTGGAG GCGCTGGCGCAAGAACTCGG CTCCGCCTTTCACGGCGCTGCCTTCGATGTTACCGACGAAGATGCCACTA GAAAGGCACTTGCGGCTTTG CCGGAAGGTTTCCGGGACATCGATATTCTCGTCAACAATGCGGGGCTTGC GCTCGGCACCGCACCTGCAC CGCAGGTGCCGTGAAAGACTGGCAGACCATGGTGAACACCAACATCACC GGTCTTTTGAACATCACCCA CCATCTTTTGCCACGTTGATCGACCGCAAGGGCATTGTCATCAACCTTT CCTCGGTAGCTGCGCACTGG CCCTATGCGGGCGGCAATGTCTATGCCGGAACGAAAGCCTTCCTGCGGCA ATTCTCGCTCGGTCTGCGCT CCGACCTGCATGGCAAGGCGTGCGCGTCACCTCGATCGAACCGGGCATG TGCGAAACGGAAATTCACGCT TGTTGCGACCGGCGGCAATCAGGATGCCTCGGACAATCTTTACAAGGGCG TCAATCCGATCACGGCCGAG GATATCGCCAATACGATCCATTGGGTGCGCTCGCAGCCCAAACATATCAA CATCAACAGCCTCGAACTCA TGCCGGTCAACCAGTCCTTTGCCGGTTTCCAAGTGCATCGGGAAAGTTGA	ADY67207.1
		YP_004444298.1
		EAZ63492.1
		XP_001387515.1
		BAB07807.1
		EMS96834.1
		EKJ96295.1
		EHJ96027.1
		EHH03760.1
		WP_028707025.1
		NP_356536.1

Enzyme	Gene Sequence	Accession Numbers
		AEQ50417.1
		AAK89321.1
		YP_004898167.1
		YP_064393.1
		WP_003522480.1
		EGP55658.1
		EGL63994.1
		KFC62486.1
		WP_031354348.1
Serine Ammonia-Lyase	ATGATGACCAAAAACGAAATCCAAAAGTGGGTAAAGGAATTCCCGCTGCTTGAAACGATCATGGCGGCCG AAGAGGTATTTTGGCGCAATCCAAAATATCACGCGTTTGCGCAAGCTATT CGAACGATTCCTTTACGCGA ACGCGATGTCAAGGAGGCCGAAGAGCGATTGCGCCGCTTTGCCCCCTACA TCGCGAAAGTGTTCCTCCGAG ACGCGAACGGCCACGGTATCATCGAATCCCCTTTAGTGCGGATTCCGAA CATGAAACAGCGTTTGGA AGATGTTTCAGACCAACATCGAGGGGGATCTGTTGCTAAAATGCGACAGC CATCTTCCCATCTCCGGATC GATCAAGGCGAGAGGGGGAATCTACGAGGTTCTGAAACATGCGGAAGAAC TCGCTCTGGCAAACCATATG ATCACCATGGGGGATGACTATGCGGTCATGGCCAGCGAAGAATTCGGCA GTTCTTTTCCCGCTATTGCG TTGTCGTTGGTTCGACGGGAAATTTAGGCTTGAGTATCGGCATCATCGGG GCGCAGCTTGGGTTCCGCGT TACCGTTTATATGTCAGCCGATGCGAAACAATGGAAAAAGACTTGTTGC GAAGCAAAGGGGTTGCGGTC ATCGAACATCTCACCGACTACAACAAGGTGGTGGAAGAGGCGCGAAGACA GTCCGCCGAGGATCCAACGT CGTATTTTATCGATGATGAGAACTCGATCCATCTGTTTTTAGGCTATGCG GTGGCGGCGTTTCGGCTGAA AAAGCAATTAGAGGACATGAACATCACGGTTGATGAAAACCACCCGCTCT	KFL14920.1
		AIF56070.1
		KFI03369.1
		KFH36969.1
		KFH35774.1
		KFF56112.1
		WP_031409141.1
		KFC30598.1
		KEZ84476.1
		KEY95863.1
		KER46054.1
		WP_030024949.1
		KEK24273.1
		KEK22892.1
		KEK18491.1
		KEK12402.1
		WP_029761212.1
		WP_029758174.1
		WP_029714078.1
		WP_0295983
		16.1
	TTGTATATCTTCCTTGCGGC GTCGGCGGCGGTCCGGGCGGGTGACGTTTGGGCTGAAGCTCGTGTACGG CGATCATGTCCATTGCTTTT TCGCTGAGCCGACGCATTCGCCTTGCAATGTTGCTCGGCCTGATGACGGGA CAGCACGACCGCGTGTGGT GCAAGATTTTGGCTTCGACAATAAGACCGAAGCGGACGGGCTAGCGGTGG GGCGGCCGTCAAGGTTGGTG GGGAACATGCTTGAGAACGTCATCAGCGGCGTCTATACGGTGGACGATGC GACGCTTTACCGCTTGCTCG CGGCGATGGTGGAACGGAGGAAATCTATTTAGAGCCGTCCGCCTTGCGG GGGGTGGCGGGGCTGTTG GCTGTTTCGTGATTTGGCGGGGCAAACGTACGTAGAGGCAAACGGTTTGA	

Enzyme	Gene Sequence	Accession Numbers
	AAGAAAAGATGAAAAACGCC GTCCATATTGGCTGGGCGACAGGCGGAAGCATGGTGCTAAAGGATGTGAT GGAGGCCTATTATCGGGAAG GCGTGCGCATCGAAACGATGACAGGGAACGGTTTTTCTGAAGGACGATAA	
Leucine Dehydrogenase	ATGCTGATGTTCTGAAGAAATCCAGGCGCGCGGCCACGAGAGCGTCAAGCT GCTGCACCACGCCCCCAGCG GCCTGCGCGCCGTGCTCGCCGTGCACTCCACCGTGCTCGGCCCTGCCATT GCCGGCTGCCGCCTGATGCC CTGCACCGAGGAACGCGCCGTGCGCGACGCCCTCGCCCTCAGCGAGTCCG TCACGCTCAAGGCCGCCCTC GCGGGCCTGAACTACGGCGGGGGCGCGTGCGTCATGCTCCCCCGGAAGG CGGCGACATCGACGGGCACG CCCGCGAGGCGTGTTCGCGCGCTCGGCCGGCAGATCCGTTACCGCGGT GGCCGCGTCATCCTCACCGA GGACGTCGGCGTGACCGGCCGCGACATCGCCTTCGCCGCGCAGGAAACCG ACAGCACCATGGGCATGCAC ACCGACACGCCACCGTCAACCGGTACGGCGGTGTACCGGGCATCAAGGC CGCCGCGCGCGCCTACCTCG GCGGCGAGAGCATGCGCGCGGTGCGCGTCGCGCTGCTCGGCGCGGGCGCA GTCGGGCGCACCCCTCGCGCA GCACCTGCACCGCGAGGGCGCGCGCCTCACCGTCGCAGACCTGATGTCTG AGCGCGCGCAGGCCCTCGCG GACGACCTCGGCGAGCGCGTCAACCGTCGTGAGCGCCGTGACATCTTCGA CGTGCCGTGCGACGTATTCTG CGCCGTGCGCGTTCGGGCGACAGCATCAAAAGCGCCGACGTGCCCGGCTTG CAGTGCCGGGTGATCGCCGG CAGCGAACACCACCGCTCAGCCACAACGGCGAGACGCTCGTGCGCGAAG CGGGCATCACATACATCCCG GACTTCGCCATCAACAGCGCCGGCCTGATGAGCGCCGCGCAGAACCTCAG CATCGAAACGGCGGCGGAAC GCGTGTACGAGAGCGTCGCGCAGATCTGCGCGACCGCGCAGAAGTACGAG AAGCCGCCGCACGTCGTGCG CCGTAAACTCGCGCTGCGCCGCATCGAACTGATCGGCTCCATCAGCGGCC AGTACGCCGGCCAGTAA	YP_00416978 5.1
		ADV66120.1
		ADY26991.1
		AEW05136.1
		YP_00525680 8.1
		YP_00425660 8.1
		YP_00434624 5.1
		AEA45407.1
		YP_00410199 2.1
		YP_00410199 1.1
		YP_00382593 2.1
		ADU51265.1
		ADU51264.1
		ADL08309.1
		AFY88585.1
		YP_00405400 7.1
		YP_00709245 4.1
		YP_00382521 6.1
		ADR21899.1
		ADL07593.1
Aspartate Dehydrogenase	TCATGTGCCAACACGTATGTTATCACTTAAAATTTTTAGTAAAGTGACTG CTGAATATGCTGCCAAAACA CTTGTTTTTGATTTAATTACACACAGTGTTTTTTGTTATAGATTTAAA	ADP76847.1
		YP_00400360 9.1
		ADE37476.1
	CTCTCCAAAATCTCCTTTAA CATGGACTTCATGGATATTGTGTTCAACTTCAGGATCTGCAATTATCTTT ACATCCGCATCTATTCCAGA GGCTAGACTTAATGCCGCAGCAACGTTAATATTCAGTGAAATTTTTTAA CAGCTTCTGAGGATTTCCCT TTAAACACGACCTCCTTTTTTTTGGTCTTAACACCTAACGAAGTAGGTGA TTTTCTCGTTATAAGTTTTTA TTTCTTTTATCTTACCTAAGGATGCGGCTTTTACACCATCTAAACCAAT ATTGCACCGGAAGGTATGTA TATATTAGCTCCTGATTCTCTAGATTCCTTTATCAATCTTCTTCTAACTT TCTCATCTAATAGTGCACCC ACACTCATAATCAAACATCTATACCTCTACTAATTATATTGGGCACAAT TTCTTTTACTGCCTCTTGAG	AEH60264.1
		AEH50568.1
		YP_00461548 3.1
		YP_00465966 4.1
		YP_00354312 1.1
		YP_00389589

Enzyme	Gene Sequence	Accession Numbers
	AAGCAGATTCAATTATCAAATCAACTCCATTGAACATTTCTTCTACCTTT TTTACGGCAGTGCCATTTGT TAAATTTGCTAGCTTCTTAGCTTTTCTAAAATTTCTGTCATAAAAAATATT TTAATTTTATTTTTTTGATA TCTTGTTTTAAGACAAGGTAACTATTGTATTTGCAATTGCACCACATCC TATAATCCCACATCTCAT	1.1
		ADN37453.1
		ADV47603.1
		YP_00416310 1.1
		ADY50896.1
		ABX33598.1
		YP_00427271 8.1
		ADN60949.1
		ACL 18032.1
		ACL 16745.1
		ABX00971.1
		ADD08173.1
Aspartate Ammonia- Lyase	ATGTCCTCGCCTGCATCATCGCGCATCGAAAAAGACCTGCTTGGTGTTCT CGAAGTACCTGCCAACGCGT ATTACGGCATCCAGACCCTGCGAGCGGTGAACAACCTTTCACCTCTCCGGC GTGCCGCTTTCGCACTACCC GAAACTGGTAGTCGCGCTGGCCATGGTCAAGCAGGCGGCAGCGGATGCAA ACCATCAGCTCGGACACCTC AATGACGCCAAGCATGCGGCGATCAGCGAGGCCTGTGCCCCCTGATCCG CGGCGACTTCCACGATCAGT TCGTGGTCGACATGATCCAGGGCGGCGCTGGCACGTCGACCAACATGAAT GCCAACGAAGTCATCGCCAA CATCGCTCTGGAAACCATGGGTTCGAGAAAGGCGCATACAAACACCTGC ACCCCAACAACGATGTCAAC ATGGCGCAGTCGACCAACGACGCCTACCCACGGCGATCCGCTTGGGTCT GCTGCTGGGTCACGACGCTC TGCTCGCCAGCCTTTCAGCCTGATTCAGGCCTTCGCCGCCAAGGGCGAA GAATTCAACCATGTGCTGAA GATGGGCCCGCACCCAGTTGCAGGACGCCGTTCCAATGACCCTGGGTCAGG AATTCGCGCCTTCGCCACC ACCCTGACAGAAGACCTGAACCGCCTGCGCAGCCTGGCGCCAGAGCTGTT GACCGAAGTGAACCTCGGCG GAACCGCCATCGGCACCGGCATCAACGCCGACCCTGGCTATCAGAAGCTG GCAGTCGATCGTCTGGCACT CATCAGCGGCCAGCCTCTGGTGCCAGCAGCCGACCTGATCGAAGCGACCT CCGACATGGGCGCCTTCGTG TTGTTCTCGGGCATGCTCAAGCGTACTGCGGTCAAGCTGTCGAAAATCTG CAACGACCTGCGCCTGCTGT CCAGCGGCCCACGCACCGGCATCAACGAAATCAACCTGCCGGCACGTGAG CCAGGCAGCTCGATCATGCC CGGCAAGGTCAACCCGGTGATCCCGGAAGCGGTCAATCAGGTTGCCTTCG AAATCATCGGCAACGACCTG	ELS44542.1
		EXL32019.1
		EPF69098.1
		EDZ32290.1
		ACC77466.1
		ETO09916.1
		ETN58394.1
		AGZ94384.1
		EGU12843.1
		AGQ54567.1
		BAN21048.1
		ELU36465.1
		ELU36464.1
		EDS31003.1
		BAM20634.1
		ACO48312.1
		XP_00182883 3.2
		EAU92840.2
		XP_00184988 0.1
		XP_00165898 8.1
	TCGCTGACCATGGCAGCCGAAGGAGGACAATTGCAGCTCAACGTGATGGA GCCGCTGATCGCCTACAAGA TCTTCGACTCGATCCGCCTGCTGCAGCGCGCCATGGACATGCTGCGCGAG CACTGCATCGTCGGCATCAC AGCCAACGAACAGCGCTGCCGCGAGCTGGTCGAGCATTCGATCGGTCTGG TCACCGCCCTGAACCCTTAC ATCGGTTACGAGAACTCCACCCGTATCGCCCGCATCGCGCTGGAAACCGG CCGCGGCGTGCTGGAACCTGG TGCGTGAGGAAGGTCTGCTCGACGACGCCATGCTCGACGACATCCTGCGC CCGGAAAACATGATCGCTCC	

Enzyme	Gene Sequence	Accession Numbers
	GCGTCTGGCCCCCTTGAAGGCCTGA	
Valine Dehydrogenase	TCAGCGACCGCGGGCCTCGGCCATCCGCTGCTCGGCGATCCGGTCGGCCG CCGCGGCGGGCGGAATGCCG TCCGCCCTTCGCACGTGCGAATATTTCCAGCGTGGTGTCTGAAGATCTTCGT CGCCTTCGCCCTTGACCCGGT CGAAGTCGAACCCGTGCAGCTCGTCGGCGACCTGGATCACGCCGCCGGCG TTGACCACATAGTCGGGTGC GTAGAGGACCGACCGGTGCGCCAGGTCCTTCTCGACACCCGGGTGGGCCA GCTGTTTGTGGCCGCGCCG CACACCACCTTCGCCGTGAGCACCGGAACGGTCGCGTCGTTGAGCGCGCC GCCGAGCGCGCAGGGCGCGT AGATGTCGAGACCCCTCGGTGCGGATCAGCGTCTCGGTGTCCGCCACCACG GTGACCTCGGGGTGCAGATC GGTGATCCGGCGCACCGACTCCTCGCGTACGTGCGTGATCACGACCTCGG CCCGTCTGGAGAGCAGGTGC TCGACGAGGTGGTGGCCACCTTGCCGACCCCGCGACGCCGACCTTGCG GCCGCGCAGCGTCGGGTGCG CCACAGGTGCTGGGCCGAGGCCCGCATGCCCTGGAAGACACCGAACGCG GTGAGGACGGAGGAGTCGCC GGCGCCGCCGTTCTCGGGGAGCGGCCGGTGGTCCAGCGGCACTCCCTGG CGACGACGTCCATGTCGGCG ACGTAGGTGCCGACATCGCAGGCGGTGACGTACCGGCCGCCGAGCGAGGC GACGAACCGGCCGTAGGCCA GGAGGAGTTCCTCCGTCTTGATCTTCTCCGGGTCGCCGATGATGACGGCC TTGCCGCCACCGTGGTCGAG TCCGCCCAGGGCGTTCTTGACGACATCCCGCGCGACAGGTTGAGCGCGT CGCGGACGGCCTCGGCCTCG GTCGCGTACGGGTAGAAGCGGGTGCCGCCGAGGCCGGGGCCAGGGCGGT GGAGTGGAGGGCGATGACGG CCTTGAGGCCGGTGGCACGGTCCTGGCAGATCACGACTTGCTCGTGACCC CCCTGATCCGAGTGGAACAG GGTGTCAGGACGCCGTTAGTCACATCGGTCAC	YP_00793265 2.1
		AGK78767.1
		NP_628270.1
		YP_00197323 4.1
		AIJ14557.1
		YP_00752320 9.1
		WP_0156594 26.1
		CCK29082.1
		CAR62534.1
		AGT93561.1
		AEK45617.1
		ADI08852.1
		YP_00845428 2.1
		ESQ05180.1
		ESP98677.1
		YP_00553507 4.1
		YP_00496398 3.1
		EOD63988.1
		EME98953.1
		EME52779.1
Glycine Dehydrogenase	CTAGTTGTAAAAGTCGAGGGAGGCGCAACTGCACATGAGGTGACGATCTC CGTAAACCCCGTCAATGCGA CCACAGTCGGCCAGTACTTTTCAACGTACGAGTAAAGGATAAGGGGAATGC CGCCAAACGCCGATCATATG GTTTGTCCCATTTATCATCGGTGACACATCTTGCCGTGTGTGGTGCATTG TTCAAAACATTGTTATCCAC TGGTTGTTACCTTTTTCAATGGCGGCAATCTCACCTCGAATGGAAATTA	KEG12217.1
		ADH66904.1
		YP_00367941 0.1
		YP_00350749 1.1
		ADN74845.1
		ADD28471.1
		YP_00444520 3.1
	GTGCATCTGCCAAGCGATCC AACTCCCGCTTGGGTTCTGATTGCGTGGGTTCAATCATTAAAGTCCCGGG TACAGGAAACGCCAGTCTTG GCGAGTGAATCCGTAGTCCATCAACCGTTTGGCCACGTCTCTCGCCTCA ATATGAGCTGTCCTCTTGAA CCGTGGAAGATCAACGATAAACTCATGAGCGCAGTAGTTTTCTCCACCCA GGAAAAGAATCGTATAATGG TTCTCTAGGCGCTTCTTCAAGTAGTTTGCAATTCAAAACGGCGTACTCTGT ACAAGTTTTGAGCCCGTGTG ATCCAAGCATTAACATCAACATGTACGATATCGGAAGAATTGATGCTGAT CCGTACGCTGATTGTGAGAC TTGGCCGAATGGCTGTGAACCGCCAACTTTTTGGTTGAAAACAGAAATTTG GCAAAAAGGGGGCCAGATGT TGACGGACAGCTATAGGGCCCATTCGGGGCGCGCCACCACCATGGGGAAT TGAAAACGTCTTGTGGAGAT	YP_00391191 9.1
		ADQ81869.1
		AEH88507.1
		YP_00713821 9.1

Enzyme	Gene Sequence	Accession Numbers
	TAATGTGGCACACGTCGCCACCGATATATCCAGGGCCTGTATAGCCAACC ATGGCGTTAAGATTTGCCCC ATCAATGTAGCATTGTCCACCGTAGTAGTGCGCCATTGATGTAATGGATA AAATATCCTTGTCAAACAAG CCATACGTACTTGGATAATGTTATCATGATACACGACAACCTCTTTGCGTG TTTTTGGCAAGATTTCTCCA GGTCATTGATATCAACCCTGCCGTTAGACAAGCATTTCACCAAGACAATA TTCATTCTCGCCAATGTTGC CGAAGCTGGATTGCTACCATGCGCACTCTCTGGAATCAAACAGACGTTGC CGTGCTCCTTCCTTCATTGAT AGATGGTACGCACGAATAACACGAAGCCAGCGTATTACACCTTGGGCGCC ACTATTAGGCTGAAGCGATA CCGCATCCAGACCGGTAATTTCCCTTAACTTTTGCTCAAGATCTAGACAC AACGCACTGTACCCTCGCAC TTGGTCCACTGGGGCAAGGGGATGCACATTGGTGAATTCTGGCCAAGAGA GTGGTAACATAGCAGCGGCA GGGTAAAGCTTCATGGTGCAAGATCCCAACGGGACGCAACCATGCGTAA3 GCCGTAATCCTTTCGTTGTA GACGATGAATATAGCGCATCAGTTCACCTTTCACCTCTTGTACTTTTGAAAC CTTGACTGTTTCAGGAAATC AGACTTCCGCAACAGATCCAACGGTAGTACCGAATTTCTGATCGGCTAATTT TGGAAAGGGCTGCGACGACG GGAAGCTTCAACCCTGCAGCCTCCAAAAGTGACACAATGTGTCCATCCGT TGTTGCCTCATCCACAGAAA TGGAGACAGTCCCATTACTGTAATCAACAAAACATTAATACCCTTCTCA ACACATCGTGCTTGTAAATC CTCCGCTGTAATGCCTTTTAGGTTACAGTAACAGTGTGAAAAATGCAC TGTTTACCACAGAGTGTCTT ACTGATTCCATACCAACAGCGAGCACTTTCGCCTTGCCGTGTATCTCATT GGCAATCTCAATTAGACCAT CTGGACCATGGTAGCGCGCATAAAACCCACTCACCTTGGCCAATAACGCT TGTGCAGTACAGATATTTGA TGTGGCGCGCTCACGCTTAATATGTTGTTACGCTGTCTGCAGCGCCATGC GTATGGATGGCTCTCCGGCA GAATCCTTACTGACGCGGATCAGCGTCCCGGCATCAACCTCTTAAACT3 CTCCTTGACAGCAAAGAACG	YP_00461260 1.1
		YP_00417031 8.1
		YP_00416355 9.1
		YP_00404537 5.1
		YP_00478719 0.1
		YP_00714236 1.1
		YP_00706789 6.1
		YP_00710078 8.1
		YP_00477304 3.1
	CGGCGTGAGGACCTCCATATCCTAGTGGAACACCAAACGCTGGGAGGAT CCCACAACCACATCTGCATT CATTTCAACAGGTGGCTTGACAAGAACAAGCCATCAAGTCGGTCCCAC AGCAACTAATGACACCGTGC TTCCTTTGCATTCTCGAACAGTGGTGAGAAGTCATGAAGCATGCCCATCGC ATCTGGTGTTTGTACAAGGA TACCAAACAAGGAACTGTCAGTCCAGTCAATCAGATTCTGTGCGCCACG ACGACGTTTATCTTGAGCGG TTCGGCTCTTGTCTTAACCATCTCAATGCAGGATGGAAAAACAGTTTTTG ATACGAAGAACGTATTCCGC TTTCGTTGACCATGCTGAAAAGCAAGATGCATCGCCTCGGATGATGCTGT CGCTTGGTCAAGAAGAGATG CATTTGCCACATCCATCTTTGTCAAATCCATAACCATGGTTTGGAAATTC AAAAGGGACTCCAGACGTCC TTGTGCAATCTCAGCTTGGTATGGTGTGTAGGGTGTGTACCATCCAGGAT TTTCAATGACGTTGCGAAGT ATGACAGGAGGAGTAATGGACTCGTAGTACCCCTGACCAATCATGCTTTT TAGTACCTTGTTTCGCGCAC CAAGAGAGCGCACGAGTGCGAGAGCATCCATCTCACTCATAGCCGCCACC TCCGTCAAGGGTGGGCGTAC AATATCCCTTGGAAATAGCAGCCGTCATCAAATCAGAGAGACTCTCTTTTC CAACCGTTCGAAGCATCGAC ATTGTCTCAGCCGTTGTTGGACCAATATGGCGGTAAATATAGCTGTCCGT GGCAGTCCATCGAACAAATG TCACGCATGGCAAAGAGCCACGAAACAAACGACGGTACAT	
Alanine Dehydrogenase	ATGATCATTGGCCTGCCGAAAGAGATCAAAGTTAAGGAAAACCGCGTGGC ACTCACGCCCGGGGGCGTCG CCAGCCTCGTGCGCCGCGGCCACACCGTCATCGTGGAACGCAGCGCCGGC GTGGGCAGCGGCATCCAGGA CACCGAGTACGAGCAGGCCGGCGCGCAGCTCGGCAGCGCCGCCGAGGCGT GGGCCGCGCAGATGGTCGTG AAGGTCAAGGAGCCCATCAAGAGCGAATACGGGTACCTCCGCCCGGACCT GCTGCTGTTACGTACCTGC ACCTCGCTGCCGACCAGCCCCCTCACGGACGCCCTGCTGAGCGCCGGCACG ACCGCCGTTGCGTACGAGAC GGTGCAGCTCGACGACCGCAGCCTGCCGCTGCTCACGCCCATGAGTGAGG TCGCGGGCCGCTGAGCGTG CAGGCCGGCGCGTACCACCTGCAAAAGCCCATCGGCGGGCGCGGCGTGCT	YP_00417139 5.1
		ADV67730.1
		ADY25885.1
		ADV48359.1
		AFZ35471.1
		AFZ05172.1
		AEW05285.1

Enzyme	Gene Sequence	Accession Numbers
	GCTCGGCGGCGTGCCGGGGCG TGCAGGCGGGCCACGTCTGTCGTGATTGGCGGGCGGCGTCTGCGGCACGAAC GCCGCGAAAATGGCCATGGG CCTCGGCGCGAAGGTCACGGTGCTGGACGTGAACCACGGGCGCCTCTCGT ACCTCGACGACGTGTTCTTC GGGAAGCTCACCACCATGATGAGCAACGAGGCGAACATCCGCTCCATCCT GCCCCGAAGCGGACCTCGTGA TCGGCGGCGTGCTGATCCCCGGGGCGAAGGCGCCGCACCTTGTCACGCGC GACATGCTGGCGACCATGCA GGAAGGCAGCGTCATCGTCGACGTGGCGGTGGACCAGGGCGGATGCGTGG AGACCATTACGCGACGACG CACGACGATCCACGTACATCGTGGACGGCGTGATCCACTACGGCGTGGC	AEW04533.1
		AEM70054.1
		YP_00525695 7.1
		YP_00525620 5.1
		YP_00445049 2.1
		YP_00436810 3.1
		YP_00434043 2.1
		YP_00426160 9.1
		YP_00425550 2.1
		YP_00416385 7.1
		YP_00478747 6.1
		YP_00713243 7.1
		YP_00711358 8.1
Proline Dehydrogenase	GAACATGCCGGGCGCGGTGC CGCGCACCAGCACGTTCGCGCTCACGAACCAGACCATTGGGTACGTGCTG CAGCTCGCGGACAAGGGCGT GGAGGCACTCAGCGCCAGCAAGCCGCTGCTGCGTGGCCTGAACACCATCG GCGGGAAGCTGACGTACGCG GGCGTCGCGGAAGCGTTGCGCCTGACGTACACCGCGCCTGAAGTGGCGCT GGCGTAA	ADY26965.1
		ADI14996.1
		YP_00443747 0.1
		YP_00436897 4.1
		YP_00434574 4.1
		YP_00434068 4.1
		YP_00425658 2.1
		YP_00417068 0.1
		YP_00370553 9.1
		AEA44906.1
		AEB12864.1
		ADV67015.1
		AEE14339.1
		AEA34625.1

Enzyme	Gene Sequence	Accession Numbers
		EFH87253.1
		NP_868270.1
		ADQ16526.1
		AEL26370.1
		AFK04422.1
		AEM71761.1
Lysine Dehydrogenase	ATGAAAAACATTGTGGTTATCGGCGCGGGCAATATCGGTTCTGGCAATCGCCTGGATGCTGGCCGCATCAGGCGATTATCGCATCACGGTTGCCGATCGTTCAGCCGATCAGCTGGCCAATGTGCCGGCGCATGAACGGGTGACATCGTCGACATTACCGACCGTCCCGCTCTGGAAGCACTGCTAAAAGGCAAATTCGCCGTGCTCTCCGCCGCCCCACCGAATTCCACCTGACGGCGGGTATTGCCGAAGCGGGCGTTGCCGTCGGCACGCATTATCTCGATCTCACCGAAGACGTGGAATCCACCCGCAAGGTCAAGGCGCTGGCGGAAACGGCCGAAACCGCGCTCATTCCGCAATGCGGCCTCGCCCCGGCTTCATCTCCATCGTCGCTGCCGATCTCGCCGTCAAGTTCGACAAGCTGGACAGCGTGCGCATGCGCGTCGGCGCTCTGCCGCAATATCCGTCGAATGCGCTCAACTACAACC TCACCTGGAGTACCGACGGGCTGATCAACGAATATATCGAGCCCTGCGAA	BAH80102.1
		AAV93559.1
		YP_165503.1
		AIA01810.1
		AIA03878.1
		AIA03381.1
		AIA00889.1
		EXU92064.1
		EOT00338.1
		NP_882461.1
		EIJ80893.1
		AIA06975.1
		AIA05885.1
		EXU88317.1
		EOT04629.1
		AIA07859.1
	GGATTCTGTCGAAGGCCGCTCACCGCCGTTCCGGCCCTTGAGGAGCGCGAGGAGTTCTCGCTCGATGGCATCACCCTACGAGGCGTTCAACACCTCGGGCGGTCTCGGTACGCTTTCGCGACGCTGGAAGGCAAGGTGCGGACCATGAACTACCGCACTATCCGTTATCCCGGCCATGTGGCGATCATGAAGGCGCTTTTGAACGACCTCAACCTGCGCAACCGCCGCGATGTGCTGAAGGACCTGTTCGAAAACGCCCTGCCCGGCACCATGCAGGATGTGGTCATCGTCTTCGTCACCGTCTGCGGCACCCGCAACGGCCGCTTCCTGCAGGAAACCTATGCCAACAAAGTCTATGCCGGCCCGGTTTCCGGCCGGATGATGAGCGCCATCCAGATCACTACCGCCGCCGGCATCTGCACGGTTCTCGACCTGCTCGCGGAAGGCGCCCTGCCGAGAAGGGCTTCGTTGACAGGAGGAAGTGCGCGTGCCGAAGTTCCTCGAAAACCGGTTTGCCCGGTATTATGGCTCGCATGAGCCGCTGGCGCGGGTTGGGTGA	AIA04440.1
		AIA03522.1
		AIA02686.1

[0102] The disclosure relates to an ammonia or ammonium ion biosensor for measuring a total concentration of a ammonia in the blood. The ammonia biosensor comprises a measuring electrode which include as components, a mediator and an enzyme, which selectively act on the plurality of specific amino acids each serving as a substrate, and a counter electrode. In the amino-acid biosensor, the enzyme has a substrate affinity to each of the plurality of specific amino acids. The enzyme is operable to catalyze a reaction in each of the plurality of specific amino acids as a substrate so as to form a reaction product. The mediator is operable, during amino-acid concentration measurement, to carry electrons between the reaction product and the measuring electrode. Further, the amino-

acid biosensor is designed to apply a voltage between the measuring electrode and the counter electrode at a measurement point in such a manner that, in an analytical curve representing a relationship between an applied voltage and a current value in a specific concentration for each of the plurality of specific amino acids, the applied voltage is a voltage allowing the variety of the current values for the amino acids in the same concentration and at the same applied voltage.

[0103] In some aspects of the disclosure, the measuring electrode (at least a first electrode) further comprises a a hydrogel that comprises a coenzyme or reduction agent as a component. In some embodiments, the enzyme consists of a dehydrogenase. Further, the reaction product consists of a reduced coenzyme derived by reduction of the coenzyme, and the mediator is operable, during the amino-acid concentration measurement, to carry electrons from the reduced coenzyme to the measuring electrode.

[0104] In some aspects of the disclosure, a biosensor or system disclosed herein is used in conjunction with one or a combination of the following:

1. 1. a power source in electrical connection with the electrodes and capable of supplying an electrical potential difference between the electrodes sufficient to cause diffusion limited electro-oxidation of the reduced form of the mediator at the surface of the working electrode; and
2. 2. at least one meter, (such as a spectrophotometer, voltmeter and/or amperometer) in electrical connection with the electrodes and capable of measuring the diffusion limited current produced by of the reduced form of the mediator with the above-stated electrical potential difference is applied.

[0105] The meter will normally be adapted to apply an algorithm to the current measurement, whereby an ammonia or ammonium ion concentration is provided and visually displayed. Improvements in such power source, meter, and biosensor system are the subject of commonly assigned U.S. Pat. No. 4,963,814, issued Oct. 16, 1990; U.S. Pat. No. 4,999,632, issued Mar. 12, 1991; U.S. Pat. No. 4,999,582, issued Mar. 12, 1991; U.S. Pat. No. 5,243,516, issued Sep. 7, 1993; U.S. Pat. No. 5,352,351, issued Oct. 4, 1994; U.S. Pat. No. 5,366,609, issued Nov. 22, 1994; White et al., U.S. Pat. No. 5,405,511, issued Apr. 11, 1995; and White et al., U.S. Pat. No. 5,438,271, issued Aug. 1, 1995.

[0106] Ammonia or ammonium ion concentrations from a plurality of samples may be analyzed in parallel. For example, human and non-human body fluids such as whole blood, plasma, sera, lymph, bile, urine, semen, cerebrospinal fluid, spinal fluid, lacrimal fluid and stool specimens as well as other biological fluids readily apparent to one skilled in the art may be measured. Fluid preparations of tissues from humans and non-human animals can also be assayed, along with foods, water samples, fermentation products and environmental substances, which potentially contain environmental contaminants. In some aspects of the disclosure, human serum is assayed with the disclosed biosensor. In some aspects of the disclosure, the biosensor comprises or is configured to assay whole blood.

[0107] After reaction is complete, a power source (e.g., a battery) applies a potential difference between electrodes. When the potential difference is applied, the amount of oxidized form of the mediator at the auxiliary electrode and the potential difference must be sufficient to cause diffusion-limited electro-oxidation of the reduced form of the at least one mediator at the surface of the working electrode. In some aspects of the disclosure, the working electrode comprises a hydrogel disclosed herein. A current measuring meter (not shown) measures the diffusion-limited current generated by the oxidation of the reduced form of the mediator at the surface of the working electrode. The measured current may be accurately correlated to the concentration of ammonia or ammonium ion and/or one or more amino acids in sample when the following requirements are satisfied:

1. 1. The rate of the indophenol reaction based upon the concentration of indophenol reagents is governed by the rate of diffusion of the ammonia from the sample in a first vessel to the second vessel comprising a surface of the working electrode.

[0108] To manufacture biosensor a roll of metallized film is fed through guide rolls into an ablation/washing and drying station. A laser system capable of ablating bottom plate element 14 is known to those of ordinary skill in the art. Non-limiting examples of which include excimer lasers, with the pattern of ablation controlled by mirrors, lenses, and masks. A non-limiting example of such a system is the LPX-300 or LPX-200 both commercially available from LPKF Laser Electronic GmbH, of Garbsen, Germany.

[0109] In the laser ablator, the metallic layer of the metallized film is ablated in a pre-determined pattern, to form a ribbon of isolated electrode sets. The metallized film is further ablated, after the isolated electrode sets are formed to create recesses positioned adjacent the electrochemical area. The ribbon is then passed through more guide rolls, with a tension loop and through an optional inspection camera. The camera is used for quality control in order to check for defects.

[0110] Reagent is compounded and applied in a liquid form to the center of the electrochemical area at a dispensing and drying station. Reagent application techniques are well known to one of ordinary skill in the art as described in U.S. Pat. No. 5,762,770. It is appreciated that reagent may be applied to array in a liquid or other form and dried or semi-dried onto the center of the electrochemical area in accordance with this disclosure.

[0111] In addition, a roll of top plate element material is fed into an assembly station along with a roll of spacer material. Liners on either side of the spacer material are removed in that station and the top plate element or surface scaffold is applied to one side of the spacer material to form a top plate element/spacer subassembly. The top plate element/spacer subassembly is slit into the appropriate width for a row of biosensors. Next, a new release liner is added to the side of the spacer material opposite the cover and the subassembly is wound into a roll.

[0112] The ribbon of the reagent-coated bottom plate element is unwound and fed into a sensor assembly station along with the top plate element/spacer subassembly. The liner is removed from the spacer and the subassembly is placed on bottom plate element to cover reagent. Next, the assembled material is cut to form individual biosensors, which are sorted and packed into vials, each closed with a stopper, to give packaged sensor test strips.

[0113] Although ablating recesses is described herein, it is appreciated that the method of forming recesses in bottom plate element is also not limited. For example, the recesses may be formed by etching (e.g., using photolithographic methods) or otherwise removing a portion of the surface of top plate element. The nearest electrode edge is approximately about 10 μm to about 500 μm from the recess, or about 100 μm to about 400 μm from the recess, or from about 200 μm to about 300 μm from the recess. Biosensors that are formed with recesses in accordance with this disclosure yield a reagent profile with generally uniform thickness of chemistry. A generally uniform thickness of chemistry allows for more accurate sample analysis.


[0114] The processes and products described above include a disposable biosensor, especially for use individually as a diagnostic device or in combination with other components such as a pump system or spectrophotometer configured to diagnose hyperammonemia, abnormal function, or abnormally high or low amounts of ammonia in a sample.

Variations on the Indophenol Reaction

[0115] The disclosure relates to contacting a sample with one or a plurality of reagents in independently variable phases of dried, powdered or aqueous phases. The reaction has four major components: a compound comprising a phenyl group, a hypohalite, a catalyst and an alkali buffer. When these reagents are exposed to ammonia, an indophenol compound is produced that, when exposed to a light source at a particular wavelength, absorbs and/or emits a particular wavelength of light. In some embodiments, any of the methods disclosed herein may comprise a step of detecting the presence, absence, or quantity of ammonia or ammonium ion by measuring the absorbance of the contents of at least the first vessel or the second vessel.




Family of Phenols

[0116] Different compounds comprising a phenyl group can be used as long as the compound comprises a 4, 5, or 6-membered ring with at least one carbon atom and a unsubstituted 'para-position. '



Family of Hypohalites


[0117]




Hypobromite



Chloramine T



Sodium dichloroisocyanurate



Family of Catalysts/Coupling Agents


[0118]

Sodium Nitroprusside

Chromium

Iron

Manganese



Alkali Conditions

[0119] Any buffer capable of creating an alkali microenvironment for the reaction to take place with ammonia from a sample may be used. In some embodiments, a vessel comprising an alkali buffer with pH from about 8.5 to about 13 can be used in the biosensor, test strip, or system disclosed herein. Any compound that can create these alkali conditions can be used including sodium and potassium hydroxide, or sodium or potassium acetate. In some embodiments, the alkali buffer is in a powdered form, lyophilized, or aqueous solution in a vessel located within the biosensor or kit disclosed herein.

Electrode

[0120] In some embodiments, the biosensor, system or test strip disclosed herein comprise one or more electrodes. In some embodiments, the one or more electrodes transmit current variation generated by the reaction between the indophenol reagents and ammonia or ammonium ion from a sample and/or transmit current variation generated by a battery source to the light source or other equipment necessary to provide a readout of the levels of ammonia in a sample, for instance, in the case of a spectrophotometer to measure absorbance of a reactant vessel in the biosensor. In some embodiments, the electrodes comprise metal. In some embodiments, the electrodes comprise a carbon scaffold upon which a metal is deposited. In some embodiments, the electrodes comprise a carbon scaffold of carbon nanotubes.

[0121] Electrode structures which are suitable for the present disclosure and methods for the production of such structures have already been suggested in biosensor technology for other purposes. In this regard, reference is made to U.S. Pat. No. 6,645,359. Electrodes or Electrically conductive tracks are created or isolated on first surface. Tracks represent the electrodes of biosensor. As used herein, the phrase "electrode set" is a set of at least two electrodes, for example 2 to 200, or 3 to 20, electrodes. These electrodes may, for example, be a working (or measuring) electrode and an auxiliary electrode. In some embodiments, tracks cooperate to form an interdigitated electrode array positioned within the periphery of recesses and leads that extend from array and between recesses toward end.

[0122] Tracks are constructed from electrically conductive materials. Non-limiting examples of electrically-conductive materials include aluminum, carbon (such as graphite), cobalt, copper, gallium, gold, indium, iridium, iron, lead, magnesium, mercury (as an amalgam), nickel, niobium, osmium, palladium, platinum, rhenium, rhodium, selenium, silicon (such as highly doped polycrystalline silicon), silver, tantalum, tin, titanium, tungsten, uranium, vanadium, zinc, zirconium, mixtures thereof, and alloys, oxides, or metallic compounds of these elements. Preferably, tracks include gold, platinum, palladium, iridium, or alloys of these metals, since such noble metals and their alloys are unreactive in biological systems. In some embodiments, the track is a working electrode made of silver and/or silver chloride, and track is an auxiliary electrode that is also made of silver and/or silver chloride and is substantially the same size as the working electrode.

[0123] Tracks are isolated from the rest of the electrically conductive surface by laser ablation. Techniques for forming electrodes on a surface using laser ablation are known. Techniques for forming electrodes on a surface using laser ablation are known. See, for example, U.S. patent application Ser. No. 09/411,940, filed Oct. 4, 1999, and entitled "LASER DEFINED FEATURES FOR PATTERNED LAMINATES AND ELECTRODE". Tracks are

preferably created by removing the electrically conductive material from an area extending around the electrodes. Therefore, tracks are isolated from the rest of the electrically-conductive material on a surface by a gap having a width of about 5 μm to about 500 μm , preferably the gap has a width of about 100 μm to about 200 μm . Alternatively, it is appreciated that tracks may be created by laser ablation alone on bottom substrate. Further, tracks may be laminated, screen-printed, or formed by photolithography.

[0124] Multi-electrode arrangements are also possible in accordance with this disclosure. For example, it is contemplated that a biosensor may be formed that includes an additional electrically conductive track. In a three-electrode arrangement such as the arrangement depicted in Figure 4, the first track is a working electrode, the second is a counter electrode, and the third electrode is a reference electrode. It is also appreciated that an alternative three-electrode arrangement is possible where tracks are working electrodes and a third electrode is provided as an auxiliary or reference electrode. It is appreciated that the number of tracks, as well as the spacing between tracks in array may vary in accordance with this disclosure and that a number of arrays may be formed as will be appreciated by one of skill in the art. In some embodiments, the electrodes are embedded on or attached to a solid support, such as a test strip comprising a plastic and/or paper material.


[0125] Micro-electrode arrays are structures generally having two electrodes of very small dimensions, typically with each electrode having a common element and electrode elements or micro-electrodes. If "interdigitated" the arrays are arranged in an alternating, finger-like fashion (See, e.g., U.S. Pat. No. 5,670,031). These are a subclass of micro-electrodes in general. Interdigitated arrays of micro-electrodes, or IDAs, can exhibit desired performance characteristics; for example, due to their small dimensions, IDAs can exhibit excellent signal to noise ratios.

[0126] Interdigitated arrays have been disposed on non-flexible substrates such as silicon or glass substrates, using integrated circuit photolithography methods. IDAs have been used on non-flexible substrates because IDAs have been considered to offer superior performance properties when used at very small dimensions, e.g., with feature dimensions in the 1-3 micrometer range. At such small dimensions, the surface structure of a substrate (e.g., the flatness or roughness) becomes significant in the performance of the IDA. Because non-flexible substrates, especially silicon, can be processed to an exceptionally smooth, flat, surface, these have been used with IDAs. In some embodiments, the at least one electrode is a component of any IDA disclosed herein.


Membrane

[0127] In some embodiments, the membrane positioned at a fluid exchange opening comprises an ionomer. In some embodiments, the membrane comprises one or a combination of the following polymers:


1. 1.




2. 2.




3. 3.







4. 4.



5. 5.



6. 6.



wherein each of the variables p, q, r, s, t, u, v, w, x, y, and z are independently variable and are 0 or any positive integers; and wherein R is independently selected from an amine, hydroxy, hydroxyl, carbonyl, H, =O, -OH, -COOH, -N, -CH₃, -CH₂-X, halo, aryl, arylalkoxy, arylalkyl, alkynyl, alkenyl, alkylene, alkyl, alkyl-halo, arylamido, alkylheterocycle, alkylamino, alkylguanidino, alkanol, alkylcarboxy, cycloalkyl, heteroaryl, heteroarylalkyl, heteroarylalkoxy, or heterocyclyl; or any salt thereof.

[0128] In some embodiments, the R group is acidic or an electronegative substituent. In some embodiments, the variables p, q, r, s, t, u, v, w, x, y, z are independently variable and are 0 or positive integers from about 1 to about 200. In some embodiments, the variables p, q, r, s, t, u, v, w, x, y, z are independently variable and are 0 or positive integers from about 10 to about 100. In some embodiments, the variables p, q, r, s, t, u, v, w, x, y, z are independently variable and are 0 or positive integers from about 10 to about 100 across many species within a matrix of material comprising many species of polymer. A- represents the anionic or acidic groups that can include sulfonate, carboxylate, or other similar functional group. M+ represents the counter ion and may include H⁺, Li⁺, Na⁺, or similar cation. Letters (p-z) accompanied by parenthesis or brackets represent repeat units that can range from 0 to any integer value. Any polymer containing any combination of Carbon (C), Fluorine (F), Sulfur (S), Oxygen (O), Hydrogen (H), Nitrogen (N), Phosphorous (P), or any similar element, which may be used to create an ionic exchange membrane may also be utilized.

[0129] Ion exchange membranes can be constructed from polymers including perfluorinated ionomers (1&2), polyphosphazene based ionomers (3), polystyrene based ionomers (4), polystyrene based block-co-polymer

ionomers (5), and poly(arylene ether sulfone) based ionomers (6).

[0130] Total acid content for ionic exchange membranes may range from about 0.57 to about 3.5 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 4.0 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 3.0 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.9 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.8 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.7 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.6 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.5 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.4 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.3 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.2 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.1 meq/g. In some embodiments, the total acid content for ionic exchange is from about 0.57 to about 2.0 meq/g.

[0131] Membranes constructed from these ionomers may range in thickness from about .025 to about .69 mm in thickness. In some embodiments the membrane is from about .001 to about 069 mm in thickness. In some embodiments the membrane is from about .001 to about 068 mm in thickness. In some embodiments the membrane is from about .001 to about 067 mm in thickness.

[0132] In some embodiments the membrane is from about .001 to about 066 mm in thickness. In some embodiments the membrane is from about .001 to about 065 mm in thickness. In some embodiments the membrane is from about .001 to about 064 mm in thickness. In some embodiments the membrane is from about .001 to about 063 mm in thickness. In some embodiments the membrane is from about .001 to about 062 mm in thickness. In some embodiments the membrane is from about .001 to about 061 mm in thickness. In some embodiments the membrane is from about .001 to about 060 mm in thickness. In some embodiments the membrane is from about .001 to about 059 mm in thickness. In some embodiments the membrane is from about .001 to about 058 mm in thickness. In some embodiments the membrane is from about .001 to about 050 mm in thickness. In some embodiments the membrane is from about .001 to about 040 mm in thickness. In some embodiments the membrane is from about .001 to about 030 mm in thickness. In some embodiments the membrane is from about .001 to about 020 mm in thickness. In some embodiments the membrane is from about .001 to about 010 mm in thickness. In some embodiments the membrane is from about .025 to about 065 mm in thickness. In some embodiments the membrane is from about .025 to about 064 mm in thickness. In some embodiments the membrane is from about .025 to about 063 mm in thickness. In some embodiments the membrane is from about .025 to about 062 mm in thickness. In some embodiments the membrane is from about .025 to about 061 mm in thickness. In some embodiments the membrane is from about .025 to about 060 mm in thickness. In some embodiments the membrane is from about .025 to about 059 mm in thickness. In some embodiments the membrane is from about .025 to about 058 mm in thickness. In some embodiments the membrane is from about .025 to about 050 mm in thickness. In some embodiments the membrane is from about .025 to about 040 mm in thickness. In some embodiments the membrane is from about .025 to about 030 mm in thickness. In some embodiments the membrane is from about .025 to about 020 mm in thickness. In some embodiments the membrane is from about .025 to about 010 mm in thickness.

[0133] Higher total acid content and smaller membrane thickness leads to faster diffusion times. Membranes may be formed through extrusion casting, drop casting, hot pressing, or similar method.

Catridges and Disposable Devices

[0134] The biosensor, device, system, and or test strip may be or comprise a cartridge. In some embodiments, the cartridge is disposable after one use or can be used more than once per ammonia or ammonium ion detection event. In some embodiments, the cartridge comprises a plurality of microfluidic conduits in fluid communication with

a storage portion, a mixing portion and a readout portion of the cartridge. The storage portion comprises a plurality of compartments that store one or a combination of indophenol reagents either crystalized, dried, lyophilized or in solution. In some embodiments, the compartments may be partitioned from an adjacent conduit by plastic wall or other inert material. The mixing portion of the cartridge comprises a trunk-shaped conduit where one or more reagents being stored mix after they are released from the storage portion of the device. The reagents may mix with a sample and/or each other at different points in the microfluidic channels adjacent to the storage portion of the device. In some embodiments of the device the readout portion of the microfluidic conduits is adjacent to the mixing portion of the device. In some embodiments of the device, the cartridge comprises only a storage portion and a readout portion, wherein the readout portion comprises a microfluidic conduit configured to align to an instrument that measures the amount of ammonia or ammonium in a sample but also allows mixing of samples prior to any detection or quantification step takes place through the instrument. In some embodiments, the cartridge does not comprise an instrument for detection of the amount of ammonia or ammonium ion in a sample (spectrophotometer), but is configured to align the readout portion of the cartridge to a instrument capable of determining the amount of ammonia or ammonium ion in a sample. In some embodiments, the cartridge comprises an instrument for detection of the amount of ammonia or ammonium ion in a sample, such as a photodiode. In some embodiments, the cartridge comprises readout portion comprising microfluidic conduits for detection or quantification adjacent to the mixing portion of the device. In some embodiments, the cartridge comprises an instrument for detection of the amount of ammonia or ammonium ion in a sample, such as a photodiode, such instrument comprising a light source aligned to or with the readout portion of the device such that light from the light source may penetrate the readout portion and such instrument may detect the presence, absence or absorbance of wavelength of light in the readout portion.

[0135] In some embodiments, the cartridge comprises a microfluidic circuit comprising a storage portion in fluid communication with a mixing portion which is also in fluid communication with a readout portion. Fluid in such an embodiment is designed to flow from the storage portion to the mixing portion, and from the mixing portion to the readout portion of the cartridge. In some embodiments the storage portion comprises one compartment for each indophenol reagent. In some embodiments, the storage portion comprises a first compartment comprising a hypohalite (such as hypochlorite), a second compartment comprising an basic buffer (such as NaOH), and a third compartment comprising at least one indophenol reagent or indophenol related compound (such as 2-phenylphenol). In some embodiments, the storage portion comprises a fourth compartment comprising a catalyst or coupling reagent (such as Sodium Nitroprusside). In some embodiments, the storage portion comprises a fifth compartment comprising an alkali buffer (such as sodium acetate or calcium acetate or zinc acetate). In some embodiments, the cartridge comprises a fluid exchange opening between a microfluidic conduit the compartment comprising a an alkali buffer (such as sodium acetate or calcium acetate or zinc acetate). In some embodiments, a membrane disclosed herein is positioned over at least a portion of the fluid exchange opening such that when a sample comes in contact with the alkali buffer, ammonia can be transported across the membrane into the adjacent microfluidic conduit.

[0136] In some embodiments, the storage portion comprises a compartment optionally comprising an electrode. In some embodiments the compartment optionally comprising an electrode is adjacent to a compartment comprising the alkali buffer in solid or liquid phase, such compartment having an opening through which a sample may be deposited into the cartridge from a point exterior to the cartridge. In some embodiments, the cartridge comprises a sixth compartment comprising an opening and optionally comprising an electrode, such compartment having an opening through which a sample may be deposited into the cartridge from a point exterior to the cartridge. In some embodiments, the cartridge comprises a sixth compartment comprising an opening and optionally comprising an electrode, such compartment having an opening through which a sample may be deposited into the cartridge from a point exterior to the cartridge; wherein the cartridge further comprises a a compartment comprising an alkali buffer that is positioned at or substantially near the compartment comprising the opening, such that, upon inserting a sample into the compartment with an opening, the alkali buffer is transported to the compartment comprising the opening and mixes with the sample.

[0137] In some embodiments, a compartment has a volume of no more than about 100 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 100 microliters of

fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 90 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 80 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 70 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 60 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 50 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 40 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 30 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 20 microliters of fluid. In some embodiments, one or more compartments in the cartridge has a volume of no more than about 10 microliters of fluid.

[0138] FIGs. 24 through 28 depict an embodiment of the invention that is a cartridge. one half of the cartridge is depicted in FIG. 24 while the opposite facing half of the cartridge is depicted in FIG. 25. The two halves of the cartridge may be secured together by one or a plurality of screws, dowels or fasteners. The two halves of the cartridge may be made out of one or a plurality of inert materials such as a plastic and/or glass.

[0139] The cartridge half disclosed in FIG. 24 comprises a first, second, third, fourth and fifth storage compartment. FIG. 24 depicts a first, second, third, fourth and fifth compartment (labeled 1, 2, 3, 4, and 5 respectively) that define a volume immediately adjacent to, but partitioned from, a microfluidic conduit on a bottom half of the cartridge. The partition is delineated by the small solid dash bisecting the space between the compartment and the microfluidic conduit (labeled 10, 11, 12, 13, and 14 for each of the compartments 1, 2, 3, 4, and 5 respectively). In this embodiment, the first compartment comprises hypochlorite, the second compartment comprises an acidic buffer, the third compartment comprises a catalyst, the fourth compartment comprises an indophenol reagent (such as a phenolic compound), and the fifth compartment comprises an alkali buffer, which, if in aqueous solution, may be at a concentration from about 500 mM to about 1 M sodium acetate. The storage portion of the microfluidic circuit comprises the storage points 1, 2, 3, 4, 5, and 6. Any membrane disclosed herein may be placed at or near position 14 such that, upon introduction of a sample such as whole blood in the compartment 6 of FIG. 25, mixing of the reagents can occur. Fluid from compartment 5 is mixed with the sample and ammonia ions in solution may be transferred from 5 and 6 into the mixing portion of the device 20 across the membrane. The reagents in compartments 1 through 4 are also released such that after a period of about 4-5 seconds, all reagents have entered the mixing portion of the device 20. The upper branched portion of the mixing portion 20 mixes the indophenol reagents contained in the compartments 1, 2, 3 and 4 while the ammonia from the sample and buffer in 5 and 6 mix in the lower trunk of the cartridge. Once in use, FIG. 26 depicts the anticipated flow of fluid from each compartment to the mixing portion of the cartridge. Lighter shades of grey show the bolus of reagents from each compartment as they travel at 0 seconds (i); at 13 seconds (ii); and 24 seconds (iii) through the microfluidic circuit. After mixing is complete in the mixing portion of the device, all reagents mix in the portion of the mixing portion closest proximal to the readout portion 25 of the circuit. At the readout portion of the device, the cartridge may have an opening through which light may travel and expose the fluid to a certain measurable wavelength of light. An instrument such as a photodiode may be present near or adjacent to the readout portion of the device so that measurements of absorbance may be taken.


[0140] In some embodiments, the cartridge comprises at least one electrode that detects the presence or absence of ammonia or ammonium ion in a sample in a vessel configured to receive a sample from a point external to the cartridge. Once the electrode is activated by the presence of a sample, the storage portion of the cartridge opens and releases their contents such that a solution from each compartment is released into the mixing portion of the microfluidic conduits. The microfluidic conduits are of a length sufficient to mix all of the reagents from each compartment such that, by the time total fluid volume of reactants reach the readout portion of the cartridge, an indophenol reaction has taken place and an indophenol reaction product (such as indophenol or an indophenol related compound) have formed in the microfluidic conduits.

Table 3

Examples of Indophenol Reagent Concentration Ranges	
Reagent	Range
2-phenylphenol	From about 50 to about 70 mmol/liter
Sodium Nitroprusside	About 7 micromoles/liter
Sodium Hydroxide	From about 50 to about 500 mmol/liter
Sodium Hypochlorite	From about 50 to about 120 mmol/liter
Sodium/Calcium Acetate	From about 0.5 to about 1 mol/liter

Hydrogel

[0141] The biosensor comprises a hydrogel in some aspects of the disclosure. The hydrogel may be a cross-linked polymeric material that swells in water but does not dissolve. It is envisioned that the hydrogel may be capable of absorbing at least about 1 to about 10 times, and in one aspect at least about 100 times, its own weight of a liquid. The hydrogel chosen for use in the biosensor should depend directly on the method of functionalization. It is envisioned that the hydrogel may be biocompatible. In some aspects of the disclosure, the hydrogel comprises sodium alginate. In some aspects of the disclosure, the hydrogel comprises from about 0.1% to about 5% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.1% to about 4% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.1% to about 3% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.1% to about 2% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.1% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.1% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.2% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises sodium alginate. In some aspects of the disclosure, the hydrogel comprises from about 0.3% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.4% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.5% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.6% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.7% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.8% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 0.9% to about 1% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 1.0% to about 3.0% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 1.0% to about 2.0% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises from about 1.0% to about 1.5% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises about 1%, about 2%, or about 3% alginate weight/volume. In some aspects of the disclosure, the hydrogel comprises sodium alginate. The alginate may be any individual polymer of alginate used in bulk form or repitive pattern of monomers, G blocks, M blocks, and/or GM blocks. In some aspects of the disclosure the alignate comprises the formula:



where m and n are any positive integer. In some aspects of the disclosure m and n are indepedently variable and any positive integer from about 1 to about 1000. In some aspects of the disclosure, the hydrogel may be polymerized from acrylic monomers. The acrylic monomer may be one or a combination of the following: acrylamido-glycolic acid, acrylamido-methyl-propa-ne-sulfonic acid, acrylamido-ethylphosphate, diethyl-aminoethyl-acrylamide-, trimethyl-amino-propyl-methacrylamide, N-octylacrylamide, N-phenyl-acrylamide and tert-butyl-acrylamide. In aspects of the disclosure in which the device contains a cross-linking agent, exemplary cross-linking

agents may be N,N'-methylene-bis-acrylamide, N,N'-methylene-bismethacrylamide, diallyltartardiamide and poly(ethylene glycol)dimethacrylate. Examples of suitable hydrogels may also include silicon wafers, borosilicate glass substrates, 2-hydroxyethyl methacrylate (HEMA), N-Isopropylacrylamide (NIPAAm), and polyethylene glycol (PEG).

[0142] The hydrogel may include any number of molecules. For example, the hydrogel may include a polymerized monomer or hydrogel a cross linking agent and optionally a chemical or UV-light activated inducer agent. Examples of such monomers or dimers include vinyl acetates, vinyl pyrrolidones, vinyl ethers, olefins, styrenes, vinyl chlorides, ethylenes, acrylates, methacrylates, nitriles, acrylamides, maleates, epoxies, epoxides, lactones, ethylene oxides, ethylene glycols, ethyloxazolines, amino acids, saccharides, proteins, anhydrides, amides, carbonates, phenylene oxides, acetals, sulfones, phenylene sulfides, esters, fluoropolymers, imides, amide-imides, etherimides, ionomers, aryetherketones, amines, phenols, acids, benzenes, cinnamates, azoles, silanes, chlorides, and epoxides, N,N'-methylenebisacrylamide, methylenebismethacrylamide ethyleneglycol-dimethacrylate, N,N'-methylenebisacrylamide, polyethyleneglycoldiacrylate (PEGDA), polyethyleneglycoldimethacrylate (PEGDMA), polyethyleneglycoldiacrylate (PEGDA), polyethyleneglycoldimethacrylate (PEGDMA), poly(vinyliden fluoride) (PVdF) based polymer, a polyacrylonitrile (PAN) based polymer, a polymethylmethacrylate (PMMA) based polymer, a polyvinyl chloride (PVC) based polymer, and a mixture of the poly(vinyliden fluoride) (PVdF) based polymer, polyacrylonitrile (PAN) based polymer, polymethylmethacrylate (PMMA) based polymer, and polyvinyl chloride (PVC) based polymer, and mixtures of any two or more thereof. IN some aspects of the disclosure, the hydrogel does not comprise 3,4-dihydroxybenzoic acid (3, 4-DHB) or an analog thereof.

[0143] Cross linking agents and optionally the chemical or UV-light activated inducer agent may include N,N'-methylenebisacrylamide, methylenebismethacrylamide ethyleneglycol-dimethacrylate and agent N,N'-methylenebisacrylamide. Irgacure 2959 (Ciba); 2,2-dimethoxy-2-phenylacetophenone, 2-methoxy-2-phenylacetone, benzyl-dimethyl-ketal, ammonium sulfate, benzophenone, ethyl benzoin ether, isopropyl benzoin ether, .alpha.-methyl benzoin ether, benzoin phenyl ether, 2,2-diethoxy acetophenone, 1,1-dichloro acetophenone, 2-hydroxy-2-methyl-1-phenylpropane 1-on, 1-hydroxy cyclohexyl phenyl ketone, anthraquinone, 2-ethyl anthraquinone, 2-chloroanthraquinone, tioxantone, isopropyltioxantone, chloro tioxantone, 2,2-chlorobenzophenone, benzyl benzoate, and benzoyl benzoate, TEMED, and ammonium persulfate (APS). In some aspects of the disclosure, hydrogel comprises a protein, peptide, glycoprotein, proteoglycans, glycosaminoglycans, and/or carbohydrate that is secreted by cells into the extracellular environment. In some aspects of the disclosure, the secreted protein, peptide, glycoprotein, proteoglycans, glycosaminoglycans, and/or carbohydrate, or structures composed thereof.

[0144] In some aspects of the disclosure, the disclosure relates to a coated biosensor device comprising at least one coating, wherein the biosensor comprises a metabolic enzyme covalently bound or immobilized to the coating, wherein the metabolic enzyme shares at least 70% sequence identify to SEQ ID NO:1 or SEQ ID NO:2 or shares at least 70% sequence identify to functional fragments of SEQ ID NO:1 or SEQ ID NO:2. In some aspects of the disclosure, the disclosure relates to a coated biosensor device comprising at least one coating, wherein the biosensor comprises a metabolic enzyme covalently bound or immobilized within the coating, wherein the coating comprises a composition comprising a hydrogel matrix, said matrix comprising any one or combination of: alginate, trehalose, at least one electron mediator, and at lest one reduction agent. In some aspects of the disclosure, the disclosure relates to a coated biosensor device comprising at least one coating, wherein the biosensor comprises a metabolic enzyme covalently bound or immobilized to the coating, wherein the coating comprises a composition comprising a hydrogel matrix, said matrix comprising any one or combination of: poly(ethylene glycol) dimethacrylate with a molecular weight of about 1000 (PEGDMA-1000), 2-hydroxy-2 methyl propiophenone (HMPP) and at least one acrylate, wherein the acrylate is selected from the group consisting of methacrylic acid (MAA) and methyl methacrylate (MMA), wherein the ratio of PEGDMA:Acrylate is from about 10:90 mol % to about 70:30 mol %, and said HMPP is at a concentration of from about 0.2% to about 0.6%, total weight.

[0145] In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose or an analog thereof at a concentration from about 1 nM to about 999 mM. In some aspects of the disclosure, the hydrogel

solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 10 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 9 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 8 mM. In some emboidiments, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 7 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 6 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 5 mM. In some emboidiments, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 4 mM. In some emboidiments, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 3 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 2 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 1 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 10 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 100 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 200 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 300 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 400 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 500 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 600 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 700 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution prior to curing comprises trehalose at a concentration from about 800 μ M to about 1 mM. In some aspects of the disclosure, the hydrogel solution (prior to contacting with the electrode) comprises trehalose at a concentration from about 900 μ M to about 1 mM.

Enzymes

[0146] Any one or more metabolic enzymes may be chosen to used with the present disclosure. Metabolic enzymes that can be used individually or in combination with the biosensor, system or test strip disclosed herein include: any bacterial clone of phenylalanine dehydrogenase, histidine ammonia lyase, mistidine oxidase. pheylalanine lyase, glutamate dehydrogenase. In some embodiments the enzyme is chosen from any one or combination of enzymes disclosed below or their respective functional fragments that are at least 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% homoglous to the full-length enzyme or nucleic acid encoding such enzyme.

Organism	Enzyme	GenBank Accession No	SEQ ID NO
Thermoactinomyces intermedius	phenylalanine dehydrogenase	D00631.1	2
Solanum lycopersicum	phenylalanine ammonialvase	XM_004246602	7
Thermoactinomyces intermedius	phenylalanine dehydrogenase	DD421709.1	8
Caenorhabditis remanei	phenylalanine dehydro genase	XM_003102740	9
Arabidopsis thaliana	glutamate dehydrogenase	NM_121822.3	10
Spirochaeta africana	Hisitidine ammonia lyase	NC_017098.1	

SEQ ID NO:2

MRDVFEMMDRYGHEQVIFCRHPQTGLKAIILHNTTAGPALGGC
RMIPYASTDEALEDVLRSLKGMTYKCSLADVDFGGGKMVIIGDPKKDKSPELFRVIGR
FVGGLNGRFYTGTDMGTNPEDFVHAARESKSFAGLPKSYGGKGDTSIPTALGVFHGMRAT
ARFLWGTDQLKGRVVAIQGVGKVGERLLQLLVEVGAYCKIADIDSVRCEQLKEKYG
DKVQLVDVNRHKECDIFSPCAKGGVVNDTIDEFRCLAIVGSANNQLVEDRHHGALL
QERSIGVARDVYLVNACGLIQVADELFCETFERDYAKETFAVSNMAGDIEFTRAEVEMTTT

QKKSICYAPDYLVNAGGLIQVADELEGTHEEKVLAKTEAIYDMVLDIFHRAKNEINI I
CEAADRIVMERLKKLTDIRILLEDPRNSARR

SEQ ID NO:7

MASSIVQNGHVNGEAMDLCCKSINVNDPLNWEMAAESLRGSHLD
EVKKMVDEFRKPIVKLGGETLTVAQVASIANVDNKSNGVKVELSESARAGVKASSDWV
MDSMGKGTDSYGVTTGFGATSHRRTKNGGALQKELIRFLNAGVFGNGTESSHTLPHSA
TRAAMLVRINTLLQGYS GIRFEILEAITKLINSNITPCLPLRGTITASGDLVPLSYIA
GLLTGRPN SKAVGPNGEKLN AEEAFRVAGVTSGFFELQPKEGLALVNGTAVGSGMASM
VLFESNILAVMSEVLSAIFAEVMNGKPEFTDYLTHKLKHHPGQIEAAAIMEHILDGSS
YVKAAQKLHEMDPLQKPKQDRYALRTSPQWLGPQIEVIRAATKMIEREINSVNDNPLI
DVSRNKALHGGNFQGTPIGVSM DNTRLALASIGKLMFAQFSELVNDYYNNGLPSNLTA
GRNP SLDYGLKGAEIAMASYCSELQFLANPVTNHVQSAEQHNQDVNSLGLISARKTAE
AVDILKLMSSTYLVALCQAIDLRHLEENLRS AVKNTVSQVAKRTLTMGANGELHPARF
CEKELLRVVDREYVFAYADDP CSSTYPLMQKL RQVLVDHAMKNGESEKNVNSSIFQKI
VAFEDELKAVLPKEVESARAVVESGNPAIPNRITECRSYPLYRLVRQELGSELLTGEK
VRSPGEEIDKVFTAMCNGQIIDPLLECLKSWNGAPLPIC

SEQ ID NO:8

atgcgcgacg tgtttgaaat gatggaccgc tatggccacg agcaggatcat ttttgccgt
61 catccgcaaa ccggtctcaa agcgatcatc gccttgcata atacaaccgc ggggcccggt
121 ttgggtggat gccgatgat ccgatgct tegacggacg aagccttga ggatgtttg
181 cggttgtcca aaggcatgac ctataatgc agtctggcgg atgtggactt tggcggggga
241 aaaatggtta tcatcgcgga tccgaaaaa gataaatgc cggagtgtt tcgctgac
301 ggcggtttg tggcggggtt aaacggccgt ttctataccg gaaccgacat gggaaccaat
361 ccggaagatt ttgcatgc cgcagggaa tcgaaatctt ttgccggatt gccgaaatc
421 tacggcggaa agggggacac atccatccc accgcgctcg ggggtttca cggaatcgg
481 gccaccgccc ggttttatg ggggacggat cagctgaaag ggcgtgtgtt tgcaccaa
541 ggagtcggca aggtgggaga gcgctgttg cagctttgg tcgaagtggg ggctactgc
601 aaaattccg acatcgatc ggtgcgatgc gaacagctga aagaaaagta tggcgacaag
661 gtccaattgg tggatgtgaa ccgattcac aaggagagt gcgatattt ctgccttgc
721 gccaaaggcg gcgtggtcaa tgatgacac atgacgagt tccgttgct gccattgc
781 ggatccgcca acaaccaact ggtggaagac cggcatggg cactgctca aaaacggagc
841 attgttatg cacccgatta tctggtgaat gccggcgggc tgattcaagt ggctgatgaa
901 ctggaaggct tccatgaaga gagagtgc tcgcaaaaccg aagcgattta tgacatggtc
961 ctggatattt ttaccgggc gaaaaatgag aatattacca cttgtgaggc agcggaccgg
1021 atcgtgatgg agcgttgaa aaagtaacc gatattgcc ggatcttgtt ggaggatccc
1081 cgcaacagcg caaggagta a

SEQ ID NO:9

MDFKAKLLAEMAKKRKAVSGLEVKEGGAKFVRGADLESKRTQEY
EAKQEELAIKKRKADDEILQESTSRAKIVPEVPEAEFDEKTPMPEIHARLRQRGQPIL
LFGESLSVRKRLHQLEIEQPELNEG WENEMQTAMKFIGKEMDKAVVEGTADSATRHD
IALPQGYEEDNWKSIEHASTLLGVGDEM KRDCDIILSICRYILARWARDLND RPLDVK
KTAQGMHEAAHHKQTTMHLKSLMTSMEKYNVNNDIRHHLAKICRLLVIERNYLEANNA
YMEMAIGNAPWPVGVT RSGIHQRPGSAKAYVS NIAHVLNDETQRKYIQAFKRLMTKLQ
EYFPTDPSKSV E FVKKS V

SEQ ID NO:10

MNALAATNRNFKLAARLLGLDSKLEKSL LIPFREIKVECTIPKD
DGTLASFVGFRVQHDNARGPMKGGIRYHPEVDPDEVNALAQLMTWKTAVAKIPYGGAK
GGIGCDPSKLSISELERLTRVFTQKI HDLIGIHTDVPAPDMGTGPQTMAWILDEYSKF
HGYS PAVVTGKPIDL GGS LGRDAATGRGVMFGTEALLNEHGKTISGQRFVIQGF GNVG
SWAAKLISEKGGKIVAVSDITGA IKNKD GIDIPALLKHTKEHRGVKGF DGADPIDPNS
ILVEDCDILVPAALGGVINRENANEIKAKFIIEA ANHPTDPDADEILSKKG VVILPDI
YANS GGVTVS YFEWVQNIQGFMWEE EKVNDELKTYMTRSF KDLKEMCKTHSCDLRMGA
FTLGVNRVAQATILRGWGA

MNTVTNQWKAVDIFTQIRDHEQV VFCNDKNTGLKAIHAIHDTTL
GPALGGCRMYPYATVEDALFDVLR LSKGMTYKCLAADVDFGGGKAVIIGDPHKDKTPE
LFRAFQQFVESLNGRFYTG TDMGTTPDDFVHAMKETNCIVGVPEEYGGSGDSSVPTAL
GVYIGIQATNKVIWGSDELHGKTYAIQGLGKVGRKVAERLLKEGADLYVCDIHPTAIE
AIVSYAKKLGANVKVVQGTEIYRTDADIFVPCAFGNVNDNTIHVLKVKAIVGSANNQ
LLDVRH GQLLKEKGILYAPDYIVNAGGLIQVADELYGLNKERV LQKTKAIYSTLLHIY
SRAEADHITTIEAANRFCEERLQQRSRRNDFFTHRKQPKWDIRR
(SEQ ID NO:1).

Solid Support

[0147] There are many forms of ammonia- or ammonium ion-measuring devices; one common type is represented by hand-held electronic meters which receive blood samples via enzyme-based test strips. In using these systems, the patient may for example lance a finger or alternate body site to obtain a blood sample, the strip is inserted into a test strip opening in the meter housing, the sample is applied to the test strip and the electronics in the meter convert a current generated by the enzymatic reaction in the test strip to a amino acid concentration value.

[0148] Solid supports of the disclosure may be solid state but are a flexible substrate. According to the disclosure, the interdigitated array or at least one electrode is disposed proximal to, e.g., on, a flexible substrate. To act as a flexible substrate, a material must be flexible and also insulating, and is typically relatively thin. The substrate should be capable of adhering components of an IDA, or additional components of a sensor, to its surface. Such thin, insulative, flexible substrates are known in the art of flexible circuits and flex circuit photolithography. "Flexible substrates" according to the present disclosure can be contrasted to non-flexible substrates used in integrated circuit (IC) photolithography but not in flexible circuit photolithography. Examples of non-flexible substrates used in IC photolithography include silicon, aluminum oxide, and other ceramics. These non-flexible substrates are chosen to be processable to a very flat surface. Typical flexible substrates for use in the disclosure are constructed of thin plastic materials, e.g., polyester, especially high temperature polyester materials; polyethylene naphthalate (PEN); and polyimide, or mixtures of two or more of these. Polyimides are available commercially, for example under the trade name Kapton®, from I.E. duPont de Nemours and Company of Wilmington, Del. (duPont). Polyethylene naphthalate is commercially available as Kaladex®, also from duPont. A particularly preferred flexible substrate is 7 mil thick Kaladex® film.

[0149] Interdigitated arrays of the disclosure can be used in applications generally known to incorporate electrodes, especially applications known to involve interdigitated arrays of electrodes. Various applications are known in the arts of electronics and electrochemistry, including applications relating to process and flow monitoring or control, and chemical analytical methods. The arrays may be particularly useful as a component of an electrochemical sensor, where there is added value, benefit, or cost efficiency, to the use of a flexible substrate, or where there is value, benefit, or cost efficiency in having an interdigitated array of dimensions relatively larger than the dimensions of interdigitated arrays conventionally disposed on non-flexible substrates.

[0150] An interdigitated array of the disclosure can, for example, be included in an electrochemical sensor (sometimes referred to as a "biosensor" or simply "sensor") used in electrochemical detection methods. Electrochemical detection methods operate on principles of electricity and chemistry, or electrochemistry, e.g., on principles of relating the magnitude of a current flowing through a substance, the resistance of a substance, or a voltage across the substance given a known current, to the presence of a chemical species within the substance. Some of these methods can be referred to as potentiometric, chronoamperometric, or impedance, depending on how they are practiced, e.g., whether potential difference or electric current is controlled or measured. The methods and sensors, including sensors of the disclosure, can measure current flowing through a substance due directly or indirectly to the presence of a particular chemical compound (e.g., an analyte or an electroactive compound), such as a compound within blood, serum, interstitial fluid, or another bodily fluid, e.g., to identify levels of amino acids, blood urea, nitrogen, cholesterol, lactate, and the like. Adaptations of some electrochemical methods and electrochemical sensors, and features of their construction, electronics, and electrochemical operations, are described, for example, in U.S. Pat. Nos. 5,698,083, 5,670,031, 5,128,015, and 4,999,582.

[0151] Any of the above biosensor cartridges, devices, or methods comprise a volume of anticoagulant. In some aspects of the disclosure, the volume of the anticoagulant disclosed herein in a volume of about 10 microliters. In some aspects of the disclosure, the volume of the anticoagulant disclosed herein in a volume of about 20 microliters. In some aspects of the disclosure, the volume of the anticoagulant disclosed herein in a volume of about 30 microliters. In some aspects of the disclosure, the volume of the anticoagulant disclosed herein in a volume of about 40 microliters. In some aspects of the disclosure, the volume of the anticoagulant disclosed herein in a volume of about 50 microliters. In some aspects of the disclosure, the volume of the anticoagulant disclosed herein in a volume of about 100 microliters.

[0152] In some aspects of the disclosure, the methods disclosed herein comprise a step of mixing a sample comprising blood with an anticoagulant such as heparin, Acenocoumarol, phenprocoumon, Atromentin, Brodifacoum, Phenindione, Coumadin or the like. In some aspects of the disclosure the biosensor, catridge, device, or test strip comprise a mechanical shaker mechanism configured to shake one or more volumes within the at least one vessel, microfluidic conduit, or mixing portion of the biosensor, catridge, device, or test strip.

Methods

[0153] The disclosure relates to a method of diagnosing or prognosing a clinical outcome of a subject with hyperammonemia or a hyperammonia related disorder, comprising contacting a sensor, system, or test strip disclosed herein with a sample of bodily fluid obtained from a patient, and quantifying a level of ammonia or ammonium ion in the sample; and comparing the level of amino acid in the sample to a threshold value of what is considered normal level of amino acid level in the bodily fluid. In some embodiments, the method relates to to a method of diagnosing or prognosing a clinical outcome of a subject suspected of having or having been previously diagnosed with hyperpammonemia or a hyperammonemia-related disorder and/or at least one aminoacidopathy.

[0154] In some embodiments, the method relates to to a method of diagnosing or prognosing a clinical outcome of a subject suspected of having or having been previously diagnosed with at least one hyperpammonemia or a hyperammonemia-related disorder. The ranges of what ammonia or ammonium ion levels are considered normal for each age type are below in **Table 4**. If, after performing the quantification steps provided herein, the amount of ammonia or ammonium ion in the sample solution exceeds or falls below the ranges provided, diet regimen, exercise regimen, and/or medical treatment may be initiated or changed such that ammonia or ammonium ion levels are monitored until the subject's levels have stabilized or fall within what is considered a healthy range.

Table 4

Ammonia Ranges	
Case	Range
Newborn - Healthy	Less than 110 micromoles/liter
Newborn - Suspected Metabolic Disorder	Greater than 200 micromoles/liter
Older than Newborn - Healthy	50-80 micromoles/liter
Older than Newborn - Suspected Metabolic Disorder	Greater than 100 micromoles/liter
Hepatic Encephalopathy	Greater than 70 micromoles/liter

[0155] The disclosure relates to a method of detecting the presence or absence or quantity of ammonia or ammonium related disorder in bodily fluids. The disclosure also relates to a method of quantifying the concentration of ammonia or ammonium ion in bodily fluids of a subject. Quantification can occur at the point-of-care due to the quick enzymatic reaction readout caused by the generation of a detectable current within a circuit after exposure of a sample from a subject to one or a plurality of vessels comprising any one or combination of indophenol reagents disclosed herein. In some embodiments, the device or system described herein may be utilized to detect if a person has abnormally high or low levels of ammonia in the blood, after which an electronic message or display may then be provided to the user of the device or system or activated on a display by one or more processors or microchips that remotely or directly access one or more storage memories comprising one or rmore concentration values of ammonia or ammonium ion in sample of the subject. In some embodiments, multiple concentration values may be obtained either simultaneously or in series, compared or analyzed by the one or more processors operably connected to the device or system disclosed herein. In some embodiments, multiple concentration values of a subject over a time period may be compared or analyzed by the one or more processors operably connected to the device or system disclosed herein, after which a message comprising the concentration value and/or threshold values are displayed. In some embodiments, the message optionally includes a signal indicating that the subject should seek medical treatment or alter diet to control ammonia or ammonium ion levels in the subject.

[0156] The disclosure also relates to a method of diagnosing a subject with a liver dysfunction comprising:

1. (a) contacting a sample of bodily fluid from a subject to the to the biosensor, system or test strip disclosed herein;
2. (b) quantifying one or more concentration values of ammonia in the sample;
3. (c) comparing the one or more concentration values of ammonia in the sample to a threshold value of ammonia concentration identified as being in a healthy range; and
4. (d) identifying the subject as having a metabolic disease if the one or more concentration values of ammonia in the sample exceed or fall below the threshold value. In some embodiments, if the sample is blood or whole blood, the method comprises contacting the sample with an anticoagulant before or simultaneously with step (a).

[0157] The disclosure also relates to a method of diagnosing a subject with hyperammonemia comprising:

1. (a) contacting a sample of bodily fluid from a subject to the to the biosensor, system or test strip disclosed herein;
2. (b) quantifying one or more concentration values of ammonia in the sample;
3. (c) comparing the one or more concentration values of ammonia in the sample to a threshold value of ammonia concentration identified as being in a healthy range; and
4. (d) identifying the subject as having a metabolic disease if the one or more concentration values of ammonia in the sample exceed or fall below the threshold value. In some embodiments, if the sample is blood or whole blood, the method comprises contacting the sample with an anticoagulant before or simultaneously with step (a).

[0158] The disclosure also relates to a method of quantifying the amount of amino acid in sample comprising:

1. (a) contacting a sample of bodily fluid from a subject to the to the biosensor, system or test strip disclosed herein;
2. (b) quantifying one or more concentration values of ammonia in the sample;
3. (c) comparing the one or more concentration values of ammonia in the sample to a threshold value of ammonia concentration identified correlating to amino acid quantity; and
4. (d) identifying the amino acid levels if the one or more concentration values of ammonia in the sample exceed or fall below the threshold value. Any amino acid may be detected using the reference information from Figure 14 wherein the the biosensor, system or test strip disclosed herein comprises an enzyme disclosed herein or a functional fragment that has 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% sequence identity to any enzyme disclosed herein. One of ordinary skill in the art would know, for instance, that to detect the presence, absence, or quantity of amino acids listed on Table 5, one or more recombinant or synthetic enzymes disclosed herein or a functional fragment thereof that has 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% sequence identity to any sequence (either nucleic acid or encoded amino acid) disclosed herein.

[0159] In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 52 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 54 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 56 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 58 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 60 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 62 to about 70 mmol/liter. In some embodiments, the

phenolic reagent or indophenol reagent is used in a range from about 64 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 66 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 68 to about 70 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 68 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 66 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 64 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 62 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 60 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 58 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 56 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 54 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in a range from about 50 to about 52 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is used in concentration about 59 mmol/liter. In some embodiments, the phenolic reagent or indophenol reagent is 2-phenylphenol.

[0160] In some embodiments, the catalyst is used in a concentration of about 7 micromoles/liter. In some embodiments, the catalyst is sodium nitroprusside.

[illegible]

the basic buffer is used in a range from about 100 to about 120 mmol/liter. In some embodiments, the basic buffer is used in a concentration about 50 mmol/liter. In some embodiments, the basic buffer is sodium hydroxide. In some embodiments, the basic buffer is used in a concentration about 100 mmol/liter. In some embodiments, the basic buffer is sodium hydroxide. In some embodiments, the basic buffer is used in a concentration about 200 mmol/liter. In some embodiments, the basic buffer is sodium hydroxide. In some embodiments, the basic buffer is used in a concentration about 300 mmol/liter. In some embodiments, the basic buffer is sodium hydroxide. In some embodiments, the basic buffer is used in a concentration about 400 mmol/liter. In some embodiments, the basic buffer is sodium hydroxide. In some embodiments, the basic buffer is used in a concentration about 500 mmol/liter. In some embodiments, the basic buffer is sodium hydroxide.

[illegible]

some embodiments, the hypohalite is used in a range from about 50 to about 90 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 88 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 86 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 84 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 82 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 80 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 78 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 76 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 74 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 72 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 70 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 68 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 66 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 64 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 62 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 60 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 58 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 56 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 54 mmol/liter. In some embodiments, the hypohalite is used in a range from about 50 to about 52 mmol/liter. In some embodiments, the hypohalite is used in a concentration about 100 mmol/liter. In some embodiments, the hypohalite is sodium hypochlorite.

[illegible]

concentration about 0.9 mol/liter. In some embodiments, the alkali buffer is used in a concentration about 0.75 mol/liter.

[0164] The disclosure relates to a method of diagnosing liver dysfunction or hyperammonemia in a subject comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) detecting the presence, absence, or quantity of ammonia;
3. (c) correlating the quantity of ammonia to the levels of amino acid in the sample;
4. (d) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 100 micromoles/liter of sample.

[0165] The disclosure relates to a method of diagnosing liver dysfunction or hyperammonemia in a subject comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) detecting the presence, absence, or quantity of ammonia;
3. (c) correlating the quantity of ammonia to the levels of amino acid in the sample;
4. (d) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 90 micromoles/liter of sample.

[0166] The disclosure relates to a method of diagnosing liver dysfunction or hyperammonemia in a subject comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) detecting the presence, absence, or quantity of ammonia;
3. (c) correlating the quantity of ammonia to the levels of amino acid in the sample;
4. (d) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 80 micromoles/liter of sample.

[0167] The disclosure relates to a method of diagnosing liver dysfunction or hyperammonemia in a subject comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) detecting the presence, absence, or quantity of ammonia;
3. (c) correlating the quantity of ammonia to the levels of amino acid in the sample;
4. (d) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 70 micromoles/liter of sample.

[0168] Although not part of the invention, the disclosure provides a method of treating a subject with liver dysfunction or hyperammonemia comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 70 micromoles/liter of sample; and
3. (c) treating the subject by administering steroids, arginine supplements, sodium benzoate, phenylacetate, and/or a glucose solution.

[0169] Although not part of the invention, the disclosure provides a method of treating a subject with liver dysfunction or hyperammonemia comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 80 micromoles/liter of sample; and
3. (c) treating the subject by administering steroids, arginine supplements, sodium benzoate, phenylacetate, and/or a glucose solution.

[0170] Although not part of the invention, the disclosure provides a method of treating a subject with liver dysfunction or hyperammonemia comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 90 micromoles/liter of sample; and
3. (c) treating the subject by administering steroids, arginine supplements, sodium benzoate, phenylacetate, and/or a glucose solution.

[0171] Although not part of the invention, the disclosure provides a method of treating a subject with liver dysfunction or hyperammonemia comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) diagnosing the subject as having liver dysfunction or hyperammonemia if the ammonia levels are quantified as above about 100 micromoles/liter of sample; and
3. (c) treating the subject by administering steroids, arginine supplements, sodium benzoate, phenylacetate, and/or a glucose solution.

[0172] In any of the above methods, the method comprises detecting the ammonia or ammonium ion levels in whole blood, water, or a sample taken from a microenvironment such as a test solution reconstituted from a swab taken from a microenvironment.

[0173] The disclosure relates to a method of diagnosing a metabolic disorder in a subject comprising:

1. (a) contacting a sample of the subject to a system, cartridge, test strip, biosensor or device disclosed herein;
2. (b) detecting the presence, absence, or quantity of ammonia;
3. (c) correlating the quantity of ammonia to the levels of amino acid in the sample;
4. (d) diagnosing the subject as having a metabolic disorder if the amino acid levels are quantified as above those levels set forth in Table 1.

[0174] In some embodiments, any methods disclosed herein comprises taking multiple steps of detecting the presence, absence, or quantity of ammonia in a sample by performing 1, 2, 3, or more tests simultaneously or in series.

[0175] In some embodiments, the step of detecting the presence, absence, or quantity of ammonia comprises detecting the wavelength emitted or absorbed by a indophenol reaction product. In any of the above methods, the step of detecting the presence, absence, or quantity of ammonia comprises detecting the wavelength emitted or absorbed by a indophenol reaction product by looking at the visible light in one or more vessels. In some

embodiments, the step of detecting the presence, absence, or quantity of ammonia comprises detecting the wavelength absorbed by a indophenol reaction product wherein the wavelength from about 500nm to about 700nm.

[0176] In some embodiments, any of the above methods, the step of detecting the presence, absence, or quantity of ammonia comprises detecting the wavelength emitted or absorbed by a indophenol reaction product.

[0177] In some embodiments, any of the above methods do not comprise a step of converting liquid to a gas or any step involving gas chromatography.

[0178] In some embodiments, any of the above biosensor catridges, devices, or methods comprise mixing a volume of any of the reagents disclosed herein in a volume of from about 10 microliters to about 150 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise mixing a volume of any of the reagents disclosed herein in a volume of from about 10 microliters to about 100 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise a volume of any of the reagents disclosed herein in a volume of from about 10 microliters to about 150 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise a volume of any of the reagents disclosed herein in a volume of about 10 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise a volume of any of the reagents disclosed herein in a volume of about 20 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise a volume of any of the reagents disclosed herein in a volume of about 30 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise a volume of any of the reagents disclosed herein in a volume of about 40 microliters. In some embodiments, any of the above biosensor catridges, devices, or methods comprise a volume of any of the reagents disclosed herein in a volume of about 50 microliters.

[0179] The disclosure relates to a computer-implemented method of quantifying ammonia or ammonium ions and/or amino acid concentration in a sample.

[0180] The disclosure relates to a system comprising a processor that performs a computer-implemented method of quantifying amino acid concentration in a sample of a subject. In some aspects of the disclosure, the system comprises a processor optionally located at a remote location and accessible by internet connection, operably connected to a computer storage memory that stores subejct's concentration values over time. In some aspects of the disclosure, the subject or the subject's healthcare provider may accesses the internet to communicate with a server linked to the computer storage memory. Subject data reports may be generated and obtained by the subject after initiating a retrieve command through the processor. In some aspects of the disclosure, the system comprises a computer program-product that performs a function convert current signals generated by a biosensor disclosed herein to concentration of a particular amino acid and/or ammonia in a sample. The disclosure relates to a system including at least one processor and a computer readable memory, said computer readable memory having stored thereon program code for quantifying amino acid concentration in a sample of bodily fluid comprising: means for storing data associated with a subject; means for, responsive to receiving a level of current response from a biosensor or its computer storage memory, presenting a concentration value to a user as part of a user interface. In some aspects of the disclosure, the user is the subject or healthcare provider of the subject. In some aspects of the disclosure, the disclosure relates to a system that comprises at least one processor, a program storage, such as memory, for storing program code executable on the processor, and one or more input/output devices and/or interfaces, such as data communication and/or peripheral devices and/or interfaces. In some aspects of the disclosure, the user device and computer system or systems are communicably connected by a data communication network, such as a Local Area Network (LAN), the Internet, or the like, which may also be connected to a number of other client and/or server computer systems. The user device and client and/or server computer systems may further include appropriate operating system software.

[0181] The present disclosure relates generally to definition and/or use of concentration values that characterize a subject's modification of behavior. In some aspects of the disclosure, the concentration values corresponding to the concentration of amino acids in a sample of bodily fluid may characterize the degree to which a subject is advised

to modify a diet or seek medical treatment.

[0182] In some aspects of the disclosure, the present disclosure provides biosensors or test strips for use in diagnostic assays. In some embodiments the test strips are provided as part of a diagnostic or detection kit. In certain embodiments, kits for use in accordance with the present disclosure may include one or more reference samples; instructions (e.g., for processing samples, for performing tests, for interpreting results, etc.); media; and/or other reagents necessary for performing tests.

[0183] The disclosure provides a test strip comprising: a solid support, a at least a first vessel in fluid communication with at least one conduit, wherein the test strip comprises a hydrogel disclosed herein. In some embodiments, the solid support is a slide optionally coated with a polymer. In some embodiments, the solid support is coated with a polymer. In some embodiments, the polymer is polyacrylamide. In some embodiments, the solid support is a material chosen from: polystyrene (TCPS), glass, quartz, quartz glass, poly(ethylene terephthalate) (PET), polyethylene, polyvinyl difluoride (PVDF), polydimethylsiloxane (PDMS), polytetrafluoroethylene (PTFE), polymethylmethacrylate (PMMA), polycarbonate, polyolefin, ethylene vinyl acetate, polypropylene, polysulfone, polytetrafluoroethylene, silicones, poly(meth)acrylic acid, polyamides, polyvinyl chloride, polyvinylphenol, and copolymers and mixtures thereof. In some embodiments, the test strip is a paper product. In some embodiments, the at least one electrode is attached to the solid support.

[0184] The disclosure provides a software component or other non-transitory computer program product that is encoded on a computer-readable storage medium, and which optionally includes instructions (such as a programmed script or the like) that, when executed, cause operations related to the calculation of amino acid concentration values. In some aspects of the disclosure, the computer program product is encoded on a computer-readable storage medium that, when executed: quantifies one or more ammonia or ammonium ion concentration values; normalizes the one or more ammonia or ammonium ion concentration values over a control set of data; creates an amino acid profile or signature of a subject; and displays the profile or signature to a user of the computer program product. In some aspects of the disclosure, the computer program product is encoded on a computer-readable storage medium that, when executed: calculates one or more ammonia or ammonium ion concentration values, normalizes the one or more ammonia or ammonium ion concentration values, and creates an amino acid signature, wherein the computer program product optionally displays the amino acid signature and/or one or more ammonia or ammonium ion concentration values on a display operated by a user. In some aspects of the disclosure, the disclosure relates to a non-transitory computer program product encoded on a computer-readable storage medium comprising instructions for: quantifying one or more ammonia or ammonium ion concentration values; and displaying the one or more ammonia or ammonium ion concentration values to a user of the computer program product.

[0185] In some embodiments, the step of calculating one or more ammonia or ammonium ion concentration values comprises quantifying an average and standard deviation of counts on replicate trials of contacting the device or test strip with one or more samples of bodily fluids.

[0186] In some aspects of the disclosure, the one or more hydrogel coated electrodes are attached to a solid phase support. In some aspects of the disclosure, a solid phase support comprises any solid or semi-solid surface. In some aspects of the disclosure, a solid phase comprises any traditional laboratory material for growing or maintaining cells in culture including petri dishes, beakers, flasks, test tubes, microtitre plates, and/or culture slides. In some aspects of the disclosure, a solid phase comprises a glass slide, a plastic slide, a paper test strip, or combination thereof.

[0187] In some aspects of the disclosure, the one or more hydrogel coated electrodes are attached to discrete addressable sites on a solid phase support. In some aspects of the disclosure, a solid phase comprises polyamides, polyesters, polystyrene, polypropylene, polyacrylates, polyvinyl compounds (e.g. polyvinylchloride), polycarbonate, polytetrafluoroethylene (PTFE), nitrocellulose, cotton, polyglycolic acid (PGA), cellulose, dextran, gelatin, glass, fluoropolymers, fluorinated ethylene propylene, polyvinylidene, polydimethylsiloxane, polystyrene, silicon substrates (such as fused silica, polysilicon, or single silicon crystals) or combinations thereof.

[0188] The disclosure relates to a catalogue of medical records relating to a subject comprising test results from the one or plurality of methods described herein. Such catalogue, in some aspects of the disclosure, being stored on a computer readable medium being accessible remotely through a wireless internet connection.

[0189] As described above, certain aspects of the disclosure of the present disclosure may be used to distinguish between samples of bodily fluid obtained from a subject who does or is suspected of having an hyperammonemia and a subject who does not have a metabolic disease. This system is potentially useful, for example, when testing whole blood samples of a subject to determine whether disease is present. Diagnosing a patient using one or more ammonia or ammonium ion concentration values would include, for example, comparing one or more ammonia or ammonium ion concentration values of a sample from a subject with the measured reference values or threshold values of a subject.

Kits

[0190] In some aspects of the disclosure, kits in accordance with the present disclosure may be used to quantify amino acid concentration in samples of bodily fluid.

[0191] The disclosure further provides for a kit comprising one or a plurality of containers that comprise one or a plurality of the polypeptides or fragments disclosed herein. In some aspects of the disclosure, the kit comprises a test strip and/or a biosensor comprising a test strip, or any animal-based derivative of serum that enhances the culture or proliferation of cells. In some aspects of the disclosure, the kit comprises: a biosensor disclosed herein, any test strip disclosed herein, and a computer program product disclosed herein optionally comprising instructions to perform any one or more steps of any method disclosed herein. In some aspects of the disclosure, the kit does not comprise cell media. In some aspects of the disclosure, the kit comprises a solid support comprising a membrane disclosed herein and/or embedded with at least one electrode disclosed herein optionally comprising any one or combination of a hypohalite, an aqueous basic solution, and at least one compound comprising a phenyl group in one or a plurality of containers. In some aspects of the disclosure, the kit comprises a device to affix a hydrogel to a solid support.

[0192] The kit may contain two or more containers, packs, or dispensers together with instructions for preparation of an array. In some aspects of the disclosure, the kit comprises at least one container comprising the biosensor or system described herein and a second container comprising a solution for maintenance, use, and/or storage of the biosensor such as storage buffer. In some aspects of the disclosure, the kit comprises a composition comprising any molecule disclosed herein in solution or lyophilized or dried and accompanied by a rehydration mixture. In some aspects of the disclosure, the molecules and rehydration mixture may be in one or more additional containers. In some aspects of the disclosure, the kit comprises a composition comprising any one or combination of

[0193] The compositions included in the kit may be supplied in containers of any sort such that the shelf-life of the different components are preserved, and are not adsorbed or altered by the materials of the container. For example, suitable containers include simple bottles that may be fabricated from glass, organic polymers, such as polycarbonate, polystyrene, polypropylene, polyethylene, ceramic, metal or any other material typically employed to hold reagents or food; envelopes, that may consist of foil-lined interiors, such as aluminum or an alloy. Other containers include test tubes, vials, flasks, and syringes. The containers may have two compartments that are separated by a readily removable membrane that upon removal permits the components of the compositions to mix. Removable membranes may be glass, plastic, rubber, or other inert material.

[0194] The kit may contain a biosensor described herein and/or a test strip comprising a hypohalite, an aqueous basic solution, and at least one compound comprising a phenyl group. The kit may also contain a solid support such as a test strip comprising any membrane disclosed herein.

[0195] Kits may also be supplied with instructional materials. Instructions may be printed on paper or other

substrates, and/or may be supplied as an electronic-readable medium, such as a floppy disc, CD-ROM, DVD-ROM, zip disc, videotape, audio tape, or other readable memory storage device. Detailed instructions may not be physically associated with the kit; instead, a user may be directed to an internet web site specified by the manufacturer or distributor of the kit, or supplied as electronic mail.

[0196] The disclosure also provides a kit comprising: a biosensor comprising: a solid support and a plurality of electrodes, wherein at least one electrode comprises a hydrogel disclosed herein. In some aspects of the disclosure, the hydrogel comprises an immobilized metabolic enzyme or a functional fragment thereof; and optionally comprising at least one vessel comprising a hydrogel, an aqueous basic buffer, in liquid or solid phase, and at least one compound comprising a phenyl group. In some aspects of the disclosure, the kit further comprises at least one of the following: a sample, and a set of instructions, optionally accessible remotely through an electronic medium.

[0197] Generally referring to Figs. 1-7, a system, method, and apparatus for point of care hyperammonemia sensors may be described. In the exemplary embodiments described by the figures, samples may be tested for ammonia levels, amino acid levels, or other compound levels by being in contact with certain reagents to utilize an indophenol reaction. Color change in the reaction may be measured and correspond to certain concentrations of specific compounds and molecules by manual comparison to an extensive color-matching sheet or automated electronic analysis with the use of calibration curves.

[0198] Fig. 1 shows one exemplary embodiment of a system demonstrating the ability to detect ammonia levels in various samples. A well 100 may be made of plastic, wood, metal, composite materials, or a combination thereof. Additionally, well 100 may be comprised of synthetic compounds or polymers, such as silicone. Well 100 may further be divided into two or more sections, and may be separated by a membrane filter 105 interposed in or near the center of well 100. Membrane filter 105 may be made of a cation exchange filter such as Nafion, shown in Supplementary Fig. A, or similar perfluorinated ionomers to allow for only the passage of small positively charged and neutral molecules between sections. Therefore, membrane filter 105 may be selected to allow for the passage of various molecules or biological components based on charge, size, or similar characteristics. Other membrane filters may consequently be used for desired functionality, such as acrylamide, poly(ethylene glycol) diacrylate, poly(2-hydroxyethyl methacrylate), poly(vinyl alcohol), or other similar polymeric hydrogels. The selection of membrane filter 105 for a hyperammonemia sensor may depend on the membrane's ability to allow for the passage of molecules such as ammonia, and the ability to limit the passage of proteins, amino acids, and other molecules or compounds.

[0199] Still referring to Fig. 1, reagent section 101 may contain reagents such as phenol, 2-phenylphenol, sodium salicylate, other phenolic reagents or polymers, or a combination thereof. Further, reagent section 101 may also contain bleach, hypochlorite, chloramine T, a similar anion, or a combination thereof, catalysts such as nitroprusside, and a basic buffer such as sodium hydroxide or potassium hydroxide to maintain alkali conditions. Sample section 102 may contain serum, blood, plasma, or other liquid desired to be tested. Membrane filter 105 may only allow the passage of ammonia from section 102 to section 101. A chemical reaction, described in Fig. 3, may take place upon reception of ammonia or similar molecule into section 101, turning the reagents a blue color, as shown in section 103. Section 104 may describe the tested sample after the reaction takes place. Color sheets may be available for a qualitative comparison between colors representing specific ammonium concentrations.

[0200] In order for the cation exchange membrane, such as Nafion, to be useable for this application, a certain washing procedure and method may be disclosed. The membrane may be washed in a hydrogen peroxide aqueous solution, which may be at boiling temperatures. Additionally, the membrane may be washed in deionized water, ethylenediaminetetraacetic acid or other chelating agents, sulfuric acid, and other similar aqueous materials. The membrane may be exposed to extreme temperatures and pressures to further ensure washing.

[0201] Fig. 2 shows an exemplary embodiment of a device fitted with multiple wells. Wells 200 may be depressions or fossa in a mounting plate 203. Mounting plate 203 may be comprised of plastic, wood, metals, composite materials, or a combination thereof. Additionally, mounting plate 203 may be comprised of synthetic compounds or

polymers, such as silicone. As shown, mounting plate 203 carries three wells 200, yet those skilled in the art may appreciate the ability for a mounting plate 203 to carry substantially more or fewer wells as desired. Membrane filter 205 may be made of Nafion or similar membranes, and may be disposed of in any angle, such as a vertical placement as shown in Fig. 2, a horizontal placement, or a different angle as desired. Reagent section 201 may be filled with phenol, 2-phenylphenol, other phenolic reagents, or a combination thereof; bleach, hypochlorite, chloramine T, a similar anion, or a combination thereof; sodium hydroxide, potassium hydroxide, or a similar basic buffer to maintain alkali conditions; and one or more catalysts, such as nitroprusside. Sample section 202 may be filled with serum, blood, plasma, or similar material desired to be tested. The various wells 200 may be interconnected to facilitate the fluid flow between respective sections in order to test samples multiple times to further accuracy, or to test samples with multiple different membrane filters or reagents. Generally, if sample section 202 contains sufficient levels of ammonia, the ammonia may diffuse through membrane filter 205 and into reagent section 202, which may allow the reaction to be described in Fig. 3 to take place.

[0202] Fig. 3 shows exemplary reactions that may take place in a point of care hyperammonemia sensor, sometimes known as an indophenol reaction or Berthelot's Reaction. Reactions 300, 330, and 360 may take place upon diffusion of ammonia from one section of a well to another through a membrane filter, as described in Figs. 1-2. Anion 302 may be hypochlorite, as shown, bleach, calcium hypochlorite, sodium hypochlorite, or other similar anions. Anion 302 may then react with ammonia 301, and produce chloramine 303, or similar ammonia derivative. Chloramine 303 may then react with further reagents, such as phenol 331. A phenol-chloramine intermediate 333 may further react with additional phenol 331 molecules, producing indophenol 363 which may appear visibly blue in color. Phenol 331 may also be replaced with 2-phenylphenol for further efficacy, with other phenolic reagents such as sodium salicylate, with phenol polymers, or with a combination thereof. The color change in the reagent section of the well or depression may demonstrate the presence of ammonia in the sample section.

[0203] Fig. 4 shows a further exemplary embodiment of a testing device, comprising of a microfluidic. The microfluidic 400 may be suitable for home use in a similar fashion to blood glucose meters to provide ongoing, rapid, reliable testing for hyperammonemia, various aminoacidopathies, and other similar applications. The device 401 may be manufactured of plastic, wood, metal, composites, or a combination thereof, or a synthetic polymer or compound, such as silicone. A user may use a lancet to excrete a small amount of blood from the tip of a finger or other location on the body, and apply a small amount of blood, serum, plasma, or similar component at opening of a conduit channel 402. The sample may be transported through conduit channel 402 by capillary action and reach sample section 403. Sample section 403 may be separated from reagent section 404 by a cation exchange membrane 405, such as Nafion, whereby allowing ammonia to diffuse through membrane 405 into reagent section 404. Prior to the application of a blood sample, a squeezable reservoir 406 containing either dry or liquid bleach, hypochlorite, chloramine T, or similar anion may be manually or electronically stimulated, allowing for the flow of bleach into interposed reagent section 404. The bleach may be separate from reagents in reagent section 404 to ensure accurate and timely chemical reactions. Reagent section 404 may contain liquid or dry components of reagents disclosed in Figs. 1-3, such as phenol, 2-phenylphenol, other phenolic reagents, or a combination thereof; sodium hydroxide, potassium hydroxide, or a similar basic buffer to maintain alkali conditions; and may also contain one or more catalysts, such as nitroprusside. Upon the presence of a certain level of ammonium in the sample, the reagent section 404 may turn into a blue color, which may be compared to a separate or included color schematic for the user to identify.

[0204] Still referring to Fig. 4, the microfluidic 400 may be used multiple times or manufactured to be a single-use device. Additionally, changes may be implemented to the design and range of chemicals used to determine amino acid levels in samples. Those skilled in the art may also appreciate the ability for a device or similar device to conform to various biological or non-biological samples, such as saliva, urine, waste water, or perhaps various chemicals to be used in a laboratory or medical setting.

[0205] In addition to the qualitative methods of determining presence or levels of ammonia in applicable samples, a quantitative apparatus, system, and method may be disclosed.

[0206] Fig. 5 shows an exemplary flowchart of a sequence of events that may take place to accurately and

quantitatively identify the amount of ammonia in a sample, and is closely related to the exemplary apparatus and system disclosed in Figs. 6-7. Additionally, those skilled in the art may appreciate that quantitative analysis in this, or a similar fashion, be added to any of the apparatuses or systems disclosed in Figs. 1-4.

[0207] Fig. 5 therefore shows an exemplary flowchart of steps for a quantitative point of care hyperammonemia sensor. It may be appreciated that these steps may be interchangeable chronologically, may be altered significantly, or eliminated while receiving similar results. Block 501 may refer to a test strip of any size, similar to sizing of the testing strips of blood glucose meters. The insertion mechanism block 501 test strip may be manual or automated. Upon insertion into a device, block 502 may further disclose the initiation of a series of events that may take place under program control. A bleach reservoir may be opened, manually or automatically, into a reagent section within the device. The reagent section, sample section, or both may contain reagents necessary for an indophenol reaction, or reagents used for diagnosing aminoacidopathies or similar diseases and conditions. Block 503 may further disclose the application of a blood sample by way of lancet excretion. The blood sample may be substituted for other biological samples, which may then be transported through a conduit channel to a sample section separated by a reagent section by a cation exchange membrane, such as Nafion. Block 503 may further initiate a microchip under program control which may serve as a timing device, allowing for consistent timing between various steps. This microchip may direct a photodiode or photoresistor to remain inactive for a desired duration to allow for an adequate period of time for certain reactions to take place.

[0208] Still referring to Fig. 5, Block 504 may further disclose the diffusion of ammonia or similar compound from sample section to reagent section to initiate any reaction. After a determined period of time, the reagent section may turn blue in the presence of ammonia. The degree of coloration may be dependent on the amount of ammonia in the sample section, which will allow for accurate quantitative analysis. Block 505 may further disclose the initiation of a photodiode or photoresistor near the reagent section to measure the degree of coloration. The photodiode or photoresistor may change the current of the system based on the coloration, whereby block 506 may disclose the step of converting photodiode or photoresistor signal from an analog to a digital signal. Block 507 may further disclose the reception of a digital signal to a microchip under program control. Upon reception, a microchip of block 507 may utilize a predetermined calibration curve in order to correlate a signal to an accurate ammonia concentration value, as further disclosed in block 508. Block 509 may further disclose a transmission of data from the microchip to a display device, which may be either physically or wirelessly connected to microchip, for user accessibility. This method may include the use of fewer or significantly more microchips and controllers under additional program control. Further microchips may be useful for various tests, display mechanisms, data analysis, and both visual and auditory aesthetics. Microchips may also facilitate communication between an exemplary device and an at-home computer, cell phone, TV, or other common display and communication devices.

[0209] Fig. 6 shows an exemplary embodiment of a blood test strip for use with an electronic device further disclosed in Fig. 7. The testing strip may be large or small in nature, for use in either laboratory settings or personal home use. Conduit channel 601 may be the reception point of a sample to be tested. A blood droplet, excreted by lancet, may be placed on distal edge of conduit channel 601, where capillary action may transport sample into sample section 602. Sample section 602 may be U-shaped to increase surface area with a cation exchange membrane 604, such as Nafion. On the opposing side of membrane 604, a reagent section 603 may be filled with reagents commonly used with an indophenol or Berthelot's reaction. Bleach, or a similar anion, may be located in a separated reservoir either on the testing strip or within the electronic device in order to ensure the reactivity of certain reagents.

[0210] Fig. 7 shows an exemplary embodiment of a testing device under program control and a display device for the presentation of quantitative analysis. A blood test strip 701, such as a strip disclosed in Fig. 6, may be inserted into a port or aperture located on testing device 700, and a blood droplet 702 may be dispensed onto a conduit located distally on blood test strip 701. Upon insertion, an injection mechanism 770 may either automatically or manually add bleach or a similar anion to a reagent section 703. Bleach, chloramine T, or similar dry or liquid anion may be stored in reservoir 775, and may be refillable as desired. A photodiode or photoresistor 771 may remain inactive for a predetermined period of time until a fill sensor within microchip 773 directs the photodiode or

photoresistor to generate a signal corresponding to the coloration of reagent section 703. Photodiode or photoresistor 771 may then alter the current or voltage of the system with or without the means of an instrumentational amplifier and emit a signal sent to an analog-to-digital converter 772. Upon conversion to a digital signal, this may be sent to microchip 773 for analysis and further program control. Microchip 773 may compute signal and equate to a concentration of ammonium, or specific amino acids, within sample section 702 by means of pre-programmed calibration curves. Microchip 773 may then send data and information to display device 774 for user readability. Display device 774 may be wholly integrated into testing device 700, or may be connected to testing device 700 physically or wirelessly. Additionally, an alternate embodiment of testing device 700 may incorporate multiple microchips for further program control, and may be connected wirelessly or physically to an external display device, such as a computer, cell phone, TV, LCD screen, printer, or similar display and communication devices. Testing device 700 may also be in communication with devices at hospitals or laboratories for ease of information transfer to a user's doctor or medical facility.

[0211] Fig. 8 shows the chemical composition of Nafion. Other similar cation exchange membranes or perfluorinated ionomer membranes may also be used interchangeably.

References:

[0212] PCT Application Serial No. PCT/US2013/065548.

1. 1. J. Zschocke, G. F. Hoffmann, *Vademecum Metabolicum* (Milupa Metabolics, Friedrichsdorf, Germany, ed. 3rd, 2011).
2. 2. B. C. Lanpher, A. L. Gropman, K. A. Chapman, U. Lichter-Konecki, M. L. Summar, *Urea Cycle Disorders Overview* (NCBI Bookshelf, 2003).
3. 3. M. L. Summar, S. Koelker, D. Freedenberg, C. Le Mons, J. Haberle, H.-S. Lee, B. Kirmse, The incidence of urea cycle disorders., *Mol. Genet. Metab.* 110, 179-80 (2013).
4. 4. R. H. Singh, Nutritional management of patients with urea cycle disorders., *J. Inherit. Metab. Dis.* 30, 880-7 (2007).
5. 5. M. Msall, Neurological Outcome in Children with Inborn Errors of Urea Synthesis.pdf, *N. Engl. J. Med.* 310, 1500-1505 (1984).
6. 6. A. L. Gropman, M. L. Batshaw, Cognitive outcome in urea cycle disorders., *Mol. Genet. Metab.* 81 Suppl 1, S58-62 (2004).
7. 7. M. L. Batshaw, S. Brusilow, L. Waber, W. Blom, A. M. Brubakk, B. K. Burton, H. M. Cann, D. Kerr, P. Mamunes, R. Matalon, D. Myerberg, I. A. Schafer, Treatment of Inborn Errors of Urea Synthesis, *N. Engl. J. Med.* 306, 1387-1392 (1982).
8. 8. F. F. Poordad, Review article: the burden of hepatic encephalopathy., *Aliment. Pharmacol. Ther.* 25 Suppl 1, 3-9 (2007).
9. 9. R. F. Butterworth, J. F. Giguere, J. Michaud, J. Lavoie, G. P. Layrargues, Ammonia: key factor in the pathogenesis of hepatic encephalopathy, *Neurochem Pathol* 6, 1-12 (1987).
10. 10. R. F. Butterworth, Pathophysiology of hepatic encephalopathy: a new look at ammonia., *Metab. Brain Dis.* 17, 221-7 (2002).
11. 11. J. Stahl, Studies of the Blood Ammonia in Liver Disease, *Ann. Intern. Med.* 58 (1963).
12. 12. I. Eijgelshoven, S. Demirdas, T. A. Smith, J. M. T. van Loon, S. Latour, A. M. Bosch, The time consuming nature of phenylketonuria: A cross-sectional study investigating time burden and costs of phenylketonuria in the Netherlands, *Mol. Genet. Metab.* 109, 237-242 (2013).
13. 13. P. V. D. Burg, H. W. Mook, A simple and rapid method for the determination of ammonia in blood, *Clin. Chim. Acta* 8, 162-164 (1962).
14. 14. Y. Murawaki, K. Tanimoto, C. Hirayama, Y. Ikuta, N. Watabe, A simple and rapid microdiffusion method for blood ammonia using a reflectance meter and a reagent plate, and its clinical evaluation for liver diseases., *Clin. Chim. Acta* 144 (1984).
15. 15. R. J. Barsotti, Measurement of ammonia in blood, *J. Pediatr.* 138, S11-S20 (2001).
16. 16. J. Buttery, R. Ratnaike, B. Chamberlain, The measurement of erythro-cyte ammonia using the Hyland

- ammonia kit, J Clin Chem Clin Biochem 20 (1982).
17. 17. S. Dienst, An ion exchange method for plasma ammonia concentration, J. Lab. Clin. Med. 58 (1961).
 18. 18. J. Huizenga, C. Gips, Determination of blood ammonia using the Ammonia Checker, Ann Clin Biochem 20 (1983).
 19. 19. H. van Anken, M. Schiphorst, A kinetic determination of ammonia in plasma, Clin Chim Acta 56 (1974).
 20. 20. L. Rover Junior, J. C. Fernandes, G. de Oliveira Neto, L. T. Kubota, E. Katekawa, S. H. Serrano, Study of NADH stability using ultraviolet-visible spectrophotometric analysis and factorial design., Anal. Biochem. 260, 50-5 (1998).
 21. 21. M. Berthelot, B, Repert. Chim. Appl. , 254 (1859).
 22. 22. E. D. Rhine, G. K. Sims, R. L. Mulvaney, E. J. Pratt, Improving the Berthelot Reaction for Determining Ammonium in Soil Extracts and Water, Soil Sci. Soc. Am. J. 62 (1998).
 23. 23. T. T. Ngo, A. P. H. Phan, C. F. Yam, H. M. Lenhoff, Interference in Determination of Ammonia with the Hypochlorite-Alkali Phenol Method of Berthelot, , 46-49 (1981).

[0213] The foregoing description and accompanying figures illustrate the principles, preferred aspects of the disclosure and modes of operation of the disclosure. However, the disclosure should not be construed as being limited to the particular aspects of the disclosure or applications discussed above. Additional variations, modifications, and applications of the aspects of the disclosure discussed above will be appreciated by those skilled in the art. Additional variations and modifications may include, but are not limited to, the detection of a variety of different amino acids, such as phenylalanine, histidine, tyrosine, glutamate, threonine, serine, leucine, isoleucine, aspartate, valine, glycine, alanine, tryptophan, proline, lysine, arginine, or others. Detection of these amino acids may involve placing dehydrogenase enzymes or other ammonia lyase enzymes in the sample section of the well, along with the blood, serum, or plasma. Possible applications for the detection of the presence of ammonia or ammonium ion is to diagnose phenylketonuria or other aminoacidopathies.

[0214] Therefore, the above-described aspects of the disclosure should be regarded as illustrative rather than restrictive. Accordingly, it should be appreciated that variations to those aspects of the disclosure can be made by those skilled in the art without departing from the scope of the disclosure as defined by the following claims.

EXAMPLES

Example 1

[0215] The presented work demonstrates how the systematic investigation of previously known technologies yielded the fabrication of an effective blood ammonia sensor. The indophenol reaction, in tandem with a polyelectrolyte membrane, was explored as a means to quantify ammonia concentrations in whole blood.

[0216] The ammonia-indophenol standard curve was produced using a range of ammonium chloride concentrations in IX phosphate buffered saline (PBS) of 0 to 750 μM . The following concentrations were utilized in the indophenol reaction: 59mM 2-phenylphenol in ethanol, 7 μM sodium nitroprusside in water, 500mM sodium hydroxide in water, and 0.2-0.25% aqueous hypochlorite. These concentrations were mixed in a 1:1:1:0.5 ratio with an equal volume of the ammonium solution of interest and allowed to react at room temperature for 10 minutes. The absorbance of the resulting solution was measured at a wavelength of 635nm.

Example 2

Stability Studies

[0217] The reagents utilized in the indophenol reaction were investigated for long term stability. Aqueous solutions of hypochlorite, sodium nitroprusside, sodium hydroxide and a solution of 2-phenylphenol in ethanol were stored in separate 50mL falcon tubes, with limited exposure to light. At intervals of 3, 5, 7, 15, 21, 28, 35, 50, 75 and 100 days the hypochlorite, sodium nitroprusside, sodium hydroxide and 2-phenylphenol were utilized to develop a standard curve using ammonia concentrations ranging from 0-750 μ M. Significant deviations from the original standard curve indicated the degradation of the stored reagents. It should be noted that fresh ammonia samples were utilized at each test interval.

Response to Amino Acids

[0218] Primary amines can also undergo the indophenol reaction. Total amino acid concentrations in blood can be as high as 2.5mM, therefore the selectivity of 2-phenylphenol was determined in the indophenol reaction. 1mM solutions of each of the 21 amino acids was prepared in IX PBS. The same protocol utilized with the indophenol reagents for the ammonia standard curve was utilized with each amino acid solution. 10 minutes after the indophenol reagents and amino acid solution was mixed, its absorbance at 635nm was measured using a plate reader. The response was directly compared to the response seen from a 1mM solution of ammonium chloride and expressed as a percentage of the ammonium response.

Sensor Design

[0219] A bisected well containing blood in one section and a concentrated alkali solution in the other would provide a means for cation exchange of the whole blood to occur, yielding a strong recovery of the ammonium. A computer-aided design of the well that is both reusable and modular was 3D printed. As seen in Figure 11, two modular pieces were 3D printed from acrylonitrile-butadienestyrene thermoplastic. The pieces will snap together with the membrane in the middle, forming a Nafion bisected well. This design was chosen to provide a uniform platform for all future experiments involving this sensing mechanism. Silicone gasketing material, at a 1/64" thickness, was glued to the inner face of each well-half to ensure a water tight seal. The wells were then backfilled with polydimethylsiloxane to improve their mechanical properties.

[0220] Figure 11 Photograph of the 3D printed modular pieces snapped together around Nafion to form the bisected well utilized for the sensing experiments.

Sensor Response to Ammonia in Phosphate Buffered Saline and Whole Blood

[0221] The 3D printed wells were constructed with 1cm² pieces of Nafion membrane. In one bisection of the well a range of 0-500 μ M concentrations of ammonium chloride in IX PBS was added. In the opposing bisection a 1M alkali solution was added. Ion-exchange of ammonium was allowed to occur for 20 minutes. The alkali solution, now containing ammonia, was then extracted and utilized in the indophenol reaction. The absorbance of the resulting indophenol reaction was measured at 635nm after 10 minutes using a microplate reader.

[0222] Whole human blood was spiked using ammonium chloride to generate concentrations of ammonia of 25, 50, 75, 100, 150, 200, 250, 300, 400 and 500 μ M. This method of producing ammonia-spiked blood was verified utilizing a Siemens RXL to determine the true ammonia concentrations of the resulting whole blood. The ammonia-spiked whole blood was pipetted into the sensor in a protocol identical to the one used in the case of the ammonium in IX PBS. In one section of the well the ammonia-spiked blood was added. In the other section was the concentrated alkali solution. After 20 minutes the ion-exchange has taken place and the ammonia is extracted into the alkali solution. The ammonia-containing alkali solution was then mixed with the hypochlorite, sodium hydroxide, sodium nitroprusside, and 2-phenylphenol. The resulting indophenol reaction's absorbance was

measured at 635nm after 10 minutes.

Hypochlorite Concentrations Effect on Indophenol Response to Blood Ammonia

[0223] To reduce interference from reducing species in blood, higher concentrations of hypochlorite than conventionally utilized were employed in the indophenol reaction with ammonia extracted from whole sheep's blood. 1, 2, 3, 5, and 10X concentrations of hypochlorite were utilized and the resulting absorbance at 635nm was recorded.

Ammonia-Indophenol Standard Curve

[0224] Figure 12. The indophenol reaction produces a linear curve with concentrations of ammonium chloride ranging from 0-750 μ M with a COD of 0.9939.

[0225] The efficacy of the indophenol reaction was initially evaluated for its lower limit of quantification (LLoQ), resolution, range and, response time. The utilized reagents were optimized to produce a response from 25-1000 μ M ammonium chloride demonstrated in Figure 12. An LLoQ of 25 μ M was recorded with an average error of -15%, therefore the sensor's resolution in terms of concentration is higher at lower ammonium concentrations.

Stability Studies

[0226] One major advantage of using the indophenol reaction for determining ammonia concentrations is that it does not require any biological components such as enzymes, which are prone to stability issues. The shelf-life of the solutions used for the indophenol reaction was examined over the course of 100 days. The components of the indophenol reaction are not stable when mixed together, potentially due to the hypochlorite and the coupling agent's, sodium nitroprusside, reactivity. The response to the range of ammonium chloride concentrations was stable for up to 50 days. As seen in Figure 13, the response to 25, 150 and 500 μ M ammonium chloride does not change significantly until day 75.

Figure 13 The reagents for the indophenol reaction were stored at room temperature and used to generate an ammonia standard curve at regular intervals for 100 days. The response to 500 μ M ammonia began to degrade at day 75. The reagents of the indophenol reaction are stable at room temperature for up to 50 days before its response to different concentrations of ammonia begins to deteriorate.

Response to Amino Acids

[0227] The mechanism for the indophenol reaction is also applicable to other primary amine containing compounds. For whole blood applications this is problematic due to the presence of small amine containing molecules such as amino acids which would cause interference when measuring blood ammonia. The phenol compound utilized in the indophenol reaction, 2-phenylphenol, is thought to introduce some form of selectivity due the large phenyl group adding a degree of steric hindrance to the reaction. The selectivity of the reaction was tested with a large array of different amino acids. Since the response to the amino acids was so low, a concentration of 500 μ M was utilized for ammonia and 1mM for the amino acids. The absorbance values recorded for the amino acids were normalized with the ammonia acting as 100%. The radar graph in Figure 14 shows the response of each amino acid, the highest of which was threonine that was just 7% of the ammonia response.

[0228] Figure 14 1mM concentrations of each of the 21 amino were tested using the indophenol reaction. The absorbance measured at 635nm for each amino acid after the indophenol reaction was calculated as percentage of the response from indophenol reaction with 1mM ammonium chloride. The radar graph displays the percent

response as compared to ammonium chloride. The highest response was threonine which produced an absorbance value that was just 7% of ammonia's response.

Cation Exchange of Whole Blood

[0229] The other major source of interference for the indophenol reaction is proteins. Small quantities of proteins can completely disable the reaction from proceeding. In order to rapidly separate ammonium from whole blood while excluding any proteins, Nafion, a cation exchange membrane, was utilized. Nafion has had previous application in biosensors but almost entirely for protection of electrodes in electrochemical sensors. In this case Nafion is being operated as a cation exchange membrane rather than a nanoporous form of Teflon, as used in electrochemical sensors. Nafion is a fluorinated ionomer block copolymer. When cast into films, usually from solution in a hot press, the ionomeric block aggregate into long-range pores of the sulfones surrounded by a matrix of the fluoropolymer. The pores are highly negatively charged due to the sulfonic acids groups and are generally 1-4nm in size. These pores allow for the rapid diffusion of hydroxyl containing molecules and cations through the Nafion while inhibiting anions and completely preventing macromolecules. This would allow for the rapid diffusion of ammonia while reducing amino acid diffusion and completely eliminating proteins from passing and disabling the indophenol reaction.

[0230] The ion-exchange of ammonium through the use of the Nafion is the main mechanism of recovery of the analyte. Ammonium will diffuse across the membrane passively as well, but at a rate that is not sufficient for a beneficial point-of-care sensor. Alkali solutions of different ionic strength were tested for their effectiveness in exchanging with the ammonium from a PBS solution. It was expected that higher concentrations of salt would yield larger recoveries of ammonium. Bisected wells were prepared with Nafion membranes. A 500mM solution of ammonium chloride in PBS was placed on the 'analyte' side of the bisected well and solutions of a concentrated aqueous alkali in the opposing bisection. Distilled water resulted in a 10% recovery, in the control case, while the concentrated aqueous alkali resulted in a 75% recovery of the ammonium. The larger concentration of ammonium in the alkali solution versus the analyte is indicative of the ion exchange mechanism occurring, as the concentration would be equal if the mechanism was simply passive diffusion.

Sensor Response to Ammonia in PBS

[0231] The sensor was initially challenged with ammonium chloride solution in PBS, before introducing an environment as complex as whole blood. Concentrations ranging from 0-500 μM were analyzed. In healthy adults ammonia levels are generally 50-80 μM whereas concentrations greater than 100 μM are suspect. These numbers are higher in neonates in which case less than 110 μM is normal, up to 180 μM could be attributed to other illnesses and greater than 200 μM is cause for concern. In severe cases ammonia levels can be as high as 500 μM (1).

[0232] Figure 15 The constructed sensor's response to a range of ammonia concentrations in 1X PBS. The COD is 0.9758 with n=5 samples.

[0233] The sensor reliably extracted the ammonia in 20 minutes time. The extracted solution was then tested using the indophenol reaction and the developed color analyzed using a plate reader measuring absorbance at 635nm. This process produced the standard curve seen in Figure 15. The COD for detection in PBS was 0.97, with an error of 5-15%. In the range of 0-100mM, there was a resolution of ~ 30 μM ammonium. The sensor was efficacious over the entire clinically relevant range of ammonia levels in PBS.

Initial Sensor Response to Ammonia in Whole Blood

[0234] Initial studies in blood produced a non-linear relationship between ammonia concentration and absorbance. The response was limited at an absorbance of 0.35 at a blood ammonia concentration of 500 mM. These responses are markedly reduced from the same concentrations of ammonia in PBS. This suggests that the ammonia is either inhibited from diffusing across the Nafion membrane or small molecules from blood are interfering with the indophenol reaction. Negative interference of the indophenol reaction can also occur from the presence of certain small molecules. It has been previously reported that high concentrations of amines, thiols and reducing agents will disrupt the indophenol reaction, all of which are present in blood(23). Reducing agents will readily react with hypochlorite, an oxidizing agent, effectively disabling the indophenol reaction. To determine if this was the case, ammonia extract from whole sheep blood was exposed to a modified indophenol reaction using 2, 3, 5 and 10X more concentrated hypochlorite than conventionally utilized. As seen in Figure 17, increasing the hypochlorite concentration up to 3X improved the reaction's response to the ammonia extracted from the whole blood. At 5 and 10X concentrated hypochlorite, the reactions response to ammonia began to degrade, indicating that 3X hypochlorite is optimal for reducing the interference introduced by reducing agents found in blood.

Modified Sensor Response to Ammonia in Whole Blood

[0235] The 3X hypochlorite-modified indophenol reaction was examined in conjunction with the Nafion based separation technique for its effectiveness in distinguishing blood ammonia concentrations ranging from 25 to 500 μM , representing healthy to diseased levels. The resulting standard curve, in Figure 18, demonstrated the response in this range. There was a significant correlation, with a COD of 0.9573, between blood ammonia concentrations and the resulting absorbance at 635nm after the indophenol reaction.

[0236] Figure 18. The bisected well sensor was again used to extract ammonia in whole human blood. The extracted ammonia solutions were tested with the 3X hypochlorite-modified indophenol reaction and the absorbance measured at 635nm. In the range of 0-500 μM the COD was 0.9573 with n=5 samples.

[0237] In the range of 25-150 μM , where high resolution measurements are critical for examining treatment effectiveness, the COD was 0.9777, seen in Figure 19. The error in this range was -10%, giving a preliminary resolution of 15 μM . The relative standard deviation of 10% percent falls within the FDA guide for validation of a bioanalytical method which requires an relative standard deviation of 15% at n=5 samples. The LLoQ, 25 μM , is at least 3σ above the mean background reading of 0.04483 $\pm$ 0.00117 absorbance. Additionally, the sensor can reliably differentiate between 50 and 100 μM blood ammonia with a p=0.0001.

[0238] Figure 19 depicts the sensor's response to blood ammonia concentrations ranging from 0-150 μM . The relative standard deviation is -10% with a COD of 0.9777 with n=5 samples.

[0239] The investigated bioanalytical method for evaluating blood ammonia levels demonstrated a high degree of correlation between blood ammonia and sensor response. In the range of from about 25 to about 150mM, the relative standard deviation was approximately 10%. The sensor has about a twenty- minute response time, and the interference from other small molecules was greatly reduced. The components used are stable at room temperature for up 50 days and inexpensive.

[0240] Validation of the functionality of the biosensor will also be performed using the following experimental design. The evaluation of the efficacy of the phenylalanine sensor requires the construction of a three carbon electrode modified with an alginate hydrogel consisting of alginate, CaCl_2 , Toluidine Blue, Phenylalanine Dehydrogenase, and NAD(P) $^+$. The hydrogel will act as a filter to prevent interference from small molecules and proteins in whole blood. Initially 32 whole blood samples will be tested with phenylalanine concentrations ranging from about 35 μM to about 2000 μM . Specifically the following Phe concentrations will be tested on the enzyme electrode: 35, 100, 250, 500, 1000, 1250, 1500, 2000 μM . In this experiment 35 μM will represent a physiologically normal concentration and each other concentration above 100 μM will represent a variety of different diseased concentrations. These concentrations will be generated by doping whole blood of a concentration lower than 35 μM . Additionally, subject samples will be tested to ensure the sensor operates without issues form unforeseen

abnormalities with patient whole blood. All phenylalanine concentration will be verified by the use of high performance liquid chromatography (HPLC), which is the gold standard for determining phenylalanine levels in blood. The samples will not require preprocessing. The blood will be taken using sodium heparin vacuum tubes and then used unmodified outside of doping the blood with higher phenylalanine concentrations. The expected detection limit is 35 μ M with a range of 35-2000 μ M and a resolution of 20 μ M. Statistical Evaluations will be performed to assess the reliability of the concentration measurements. The concentration measurements will be analyzed using ANOVA single factor analysis to demonstrate differences between groups assuming a normal data distribution. Confidence intervals will be assessed and the sensitivity and the reproducibility of the method demonstrated. Concentration measurements ranging from normal physiological conditions to diseased conditions with a confidence 95% ($p < 0.05$) or higher will be deemed statistically significant. Statistical differences between diseased concentration levels and healthy, physiological concentration levels will first be demonstrated. Subsequent experimentation will be used to validate quantification of over the full range of discrete concentration values.

Example 3

Measuring Ammonia in Whole Blood with a Modular Well Plate

Materials

[0241] 2-phenylphenol, sodium nitroprusside, sodium hydroxide, sodium hypochlorite, sodium acetate, and ammonium chloride were purchased from Sigma-Aldrich. Nafion 111 was purchased from Ion-Power. 1/64" silicone gasket with adhesive backing was purchased from McMaster-Carr, Acrylonitrile Butadiene Styrene Resin.

Methods

Preparation of Sample

[0242]

- Prepare stock solutions of 59mM 2-phenylphenol in ethanol, 7 μ M sodium nitroprusside in water, 500mM sodium hydroxide in water and 0.6-0.75% sodium hypochlorite in water.
- Prepare a stock solution of 1M sodium acetate in water.
- 3D printed wells are produced from fused deposition modeling using acrylonitrile butadiene styrene based on the model seen in Figure 210.
- Attach a layer of the 1/64" silicone to the area of the well shaded in Figure 21.
- Remove the 25 μ m thick Nafion 111 from plastic backing and cut it into 1.5X1.5 cm squares.

Ammonia Exchange

[0243]

- Snap two of the modular 3D printed pieces around one of the Nafion Squares, creating a Bisected Well. The final configuration is shown in Figure 22.

- Pipette 45 μ l of the sodium acetate solution into section B of the well.
- Pipette 100 μ M of the ammonia containing sample (whole blood) into section A of the well.
- Wait 20 minutes for ion exchange to occur.

Collection of Data

[0244]

- Pipette 35 μ l of the sodium acetate/ammonia solution from section B into a 384 well plate.
- Pipette 10 μ l each of the 2-phenylphenol, sodium nitroprusside and sodium. hydroxide stock solutions into the same well as the sodium acetate.
- Add 5 microliters of the sodium hypochlorite solution to the well and mix thoroughly by pipette.
- Wait ten minutes for the indophenol reaction to proceed.
- Measure the absorbance of the well at 635nm using a 384 plate reader.
- Compare the measured absorbance to a standard curve to determine unknown ammonia concentration.

[0245] Figure 20 CAD sketch of the front face of the well plate.

[0246] Figure 21 CAD sketch demonstrating the area of the well plate that the adhesive silicone should be attached to (black area).

[0247] Figure 22 Photograph of the 3D printed modular pieces snapped together around Nafion to divide the well into two sections.

[0248] Figure 23 Engineering Drawing of 3D printed well.

Example 4

Measuring Ammonia in Whole Blood with a Fluidic Device

Materials

[0249] 2-phenylphenol, sodium nitroprusside, sodium hydroxide, sodium hypochlorite, sodium acetate, and ammonium chloride were purchased from Sigma-Aldrich. Nafion 111 was purchased from Ion-Power.

Methods

Preparation of Sample

[0250]

- Prepare stock solutions of 59mM 2-phenylphenol in ethanol, 7 μ M sodium nitroprusside in water, 500mM sodium hydroxide in water and 0.75% sodium hypochlorite in water.

- Prepare a stock solution of 1M sodium acetate in water.
- Molds are produced from a 3D printer for the 2 individual pieces of the device.
- Fill each mold with the PDMS elastomer and heat at 60°C for one hour.
- Remove each side of the device from the mold with a spatula.
- Remove the 25 µm thick Nafion 111 from plastic backing and cut it into a 1.5X1.5 cm square.
- Glue the square of Nafion over the well in the channel 6 using PDMS.
- Glue the top piece of the device to the bottom piece using PDMS, ensuring channel 6 lines up with channel 5.
- Heat the device to 60°C for one hour.

Ammonia Exchange

[0251]

- Insert a needle through the bottom of the device into well 6, and fill with 40 µl of blood.
- Fill channels 1-4 with 5 µl of 2-phenylphenol, sodium nitroprusside, sodium hydroxide and sodium hypochlorite respectively .
- Fill channel 5 with 20 µl of sodium acetate solution.
- Wait 20 minutes for ion exchange to occur.

Collection of Data

[0252]

- Apply a flow rate of 1mm/sec to channels 1 through 5 for 24 seconds.
- Wait ten minutes for the indophenol reaction to proceed.
- Insert the device into the custom photo-spectrometer to acquire absorbance data.
- Compare the measured absorbance to a standard curve to determine unknown ammonia concentration.

[0253] An embodiment depicting a method of detecting amino acids based upon ammonia levels is depicted in Figure 29. After addition of blood as depicted on the left handside of FIG. 29, ammonia levels are visualized by a color change of ammonia reaction with indophenol reagents in the left well of the device depicted on the righthand side. The left well is more grey (an indication of blue color) than the left well in the device depicte on the lefthand side (before the indophenol reaction takes place).

Example 5

Detection of Amino Acids in Whole Blood through Use of Ammonia Producing Enzymes with a Modular Well Plate

[0254] Dehydrogenases and ammonia-lyases generally affect amino acids by cleaving off the primary amine thereby generating ammonia. Using the ammonia dection modular well plate system described herein, the ammonia generated, measured via the color change produced, can then be correlated to the concentration of a given amino acid. This example tests for the presence of phenylaline using phenylalanine ammonia-lyase, although

other amino acids can be tested by using the appropriate enzyme(s) listed on Table 5.

Table 5

Phenylalnine	Histidine	Tyrosine	Glutamate	Threonine	Serine
Phenylalanine	Histidine	Tyrosine	Glutamate	Threonine	Serine
Ammonia	Ammonia	Ammonia	Dehydrogenase	Ammonia	Ammonia
Lyase,	Lyase	Lyase		Lyase,	Lyase, Serine
Phenylalanine				Threonine	Dehydrogenase
Dehydrogenase				Dehydrogenase	
Leucine	Isoleucine	Aspartate	Valine	Glycine	Alanine
Leucine	Isoleucine	Aspartate	Valine	Glycine	Alanine
Dehydrogenase	Dehydrogenase	Ammonia	Dehydrogenase	Dehydrogenase	Dehydrogenase
		Lyase,			
		Aspartate			
		Dehydrogenase			
Tryptophan	Proline	Lysine	Arginine		
Tryptophan	Proline	Lysine	Arginine		
Dehydrogenase	Dehydrogenase	Dehydrogenase	Dehydrogenase		

Materials

[0255] 2-phenylphenol, sodium nitroprusside, sodium hydroxide, sodium hypochlorite, sodium acetate, and ammonium chloride were purchased from Sigma-Aldrich. Nafion 111 was purchased from Ion-Power. 1/64" silicone gasket with adhesive backing was purchased from McMaster-Carr, Acrylonitrile Butadiene Styrene Resin, phenylalanine ammonia-lyase, sodium alginate from brown algae (optional), phosphate buffered saline (optional), 0.1M CaCl2 solution (optional).

Methods

Preparation of Sample

[0256]

- Prepare stock solutions of 59mM 2-phenylphenol in ethanol, 7 μM sodium nitroprusside in water, 500mM sodium hydroxide in water and 0.6-0.75% sodium hypochlorite in water.
- Prepare a stock solution of 1M sodium acetate in water.
- 3D printed wells are produced from fused deposition modeling using acrylonitrile butadiene styrene based on the model seen in Figure 210.
- Attach a layer of the 1/64" silicone to the area of the well shaded in Figure 21.
- Remove the 25 μm thick Nafion 111 from plastic backing and cut it into 1. 5X1.5 cm squares.

[0257] The enzyme can be added to the amino acid containg sample (whole blood) in two ways. Either by adding in the enzyme directly to the sample, or by immobilizing the enzyme in a gel placed inside the sample well, introducing greater enzyme stability.

Ammonia Exchange with free enzyme (option 1)

[0258]

- Snap two of the modular 3D printed pieces around one of the Nafion Squares, creating a Bisected Well. The final configuration is shown in Figure 22.
- Pipette 45 μ l of the sodium acetate solution into section B of the well.
- Pipette 100 μ M of the amino acid containing sample (whole blood) into section A of the well.
- Add 40 units of phenylalanine ammonia-lyase to the sample.
- Wait 20 minutes for ion exchange to occur.

Ammonia Exchange with gel-immobilized enzyme (option 2)

[0259]

- Snap two of the modular 3D printed pieces around one of the Nafion Squares, creating a Bisected Well. The final configuration is shown in Figure 22.
- Prepare a pre-gel solution of 40 units of phenylalanine ammonia-lyase, and 1% weight/volume sodium alginate from brown algae in 1 mL 1x phosphate buffered saline.
- Place 10 μ L of the pre-gel solution into section A of the well (where the whole blood will go).
- Spray the pre-gel solution in the well with 0.1M CaCl_2 solution using a Badger 200N airbrush at 7.5 psi for 1 second, depositing $\sim 5\mu\text{L}$ of the CaCl_2 solution. The gel will be allowed to cure for 30 minutes in a humid environment.
- Pipette 45 μ l of the sodium acetate solution into section B of the well.
- Pipette 100 μ M of the amino acid containing sample (whole blood) into section A of the well.
- Wait 20 minutes for ion exchange to occur.

Collection of Data

[0260]

- Pipette 35 μ l of the sodium acetate/ammonia solution from section B into a 384 well plate.
- Pipette 10 μ l each of the 2-phenylphenol, sodium nitroprusside and sodium. hydroxide stock solutions into the same well as the sodium acetate.
- Add 5 microliters of the sodium hypochlorite solution to the well and mix thoroughly by pipette.
- Wait ten minutes for the indophenol reaction to proceed.
- Measure the absorbance of the well at 635nm using a 384 plate reader.
- Compare the measured absorbance to a standard curve to determine unknown ammonia concentration.

Example 6

Detection of Amino Acids in Whole Blood through Use of Ammonia Producing Enzymes with a Fluidic Device

[0261] Dehydrogenases and ammonia-lyases generally affect amino acids by cleaving off the primary amine thereby generating ammonia. Using the ammonia detection modular well plate system described herein, the ammonia generated, measured via the color change produced, can then be correlated to the concentration of a given amino acid. This example tests for the presence of phenylalanine using phenylalanine ammonia-lyase, although other amino acids can be tested by using the appropriate enzyme(s) listed on Table 5.

Materials

[0262] 2-phenylphenol, sodium nitroprusside, sodium hydroxide, sodium hypochlorite, sodium acetate, and ammonium chloride were purchased from Sigma-Aldrich. Nafion 111 was purchased from Ion-Power, phenylalanine ammonia-lyase, sodium alginate from brown algae (optional), phosphate buffered saline (optional), 0.1M CaCl₂ solution (optional).

Methods

Preparation of Sample

[0263]

- Prepare stock solutions of 59mM 2-phenylphenol in ethanol, 7 μ M sodium nitroprusside in water, 500mM sodium hydroxide in water and 0.75% sodium hypochlorite in water.
- Prepare a stock solution of 1M sodium acetate in water.
- Molds are produced from a 3D printer for the 2 individual pieces of the device.
- Fill each mold with the PDMS elastomer and heat at 60°C for one hour.
- Remove each side of the device from the mold with a spatula.
- Remove the 25 μ m thick Nafion 111 from plastic backing and cut it into a 1.5X1.5 cm square.
- Glue the square of Nafion over the well in the channel 6 using PDMS.
- Glue the top piece of the device to the bottom piece using PDMS, ensuring channel 6 lines up with channel 5.
- Heat the device to 60°C for one hour.

[0264] The enzyme can be added to the amino acid containing sample (whole blood) in two ways. Either by adding in the enzyme directly to the sample, or by immobilizing the enzyme in a gel placed inside the sample well, introducing greater enzyme stability.

Ammonia Exchange with free enzyme (option 1)

[0265]

- Insert a needle through the bottom of the device into well 6, and fill with 40 μ l of blood.
- Add 40 units of phenylalanine ammonia-lyase to the blood sample.
- Fill channels 1-4 with 5 μ l of 2-phenylphenol, sodium nitroprusside, sodium hydroxide and sodium hypochlorite respectively.
- Fill channel 5 with 20 μ l of sodium acetate solution.
- Wait 20 minutes for ion exchange to occur.

Ammonia Exchange with gel-immobilized enzyme (option 2)**[0266]**

- Prepare a pre-gel solution of 40 units of phenylalanine ammonia-lyase, and 1% weight/volume sodium alginate from brown algae in 1 mL 1x phosphate buffered saline.
- Place 10 µL of the pre-gel solution into section A of the well (where the whole blood will go).
- Spray the pre-gel solution in the well with 0.1M CaCl₂ solution using a Badger 200N airbrush at 7.5 psi for 1 second, depositing ~5µL of the CaCl₂ solution. The gel will be allowed to cure for 30 minutes in a humid environment.
- Insert a needle through the bottom of the device into well 6, and fill with 40 µl of blood.
- Add 40 units of phenylalanine ammonia-lyase to the blood sample.
- Fill channels 1-4 with 5 µl of 2-phenylphenol, sodium nitroprusside, sodium hydroxide and sodium hypochlorite respectively .
- Fill channel 5 with 20 µl of sodium acetate solution.
- Wait 20 minutes for ion exchange to occur.

Collection of Data**[0267]**

- Apply a flow rate of 1 mm/sec to channels 1 through 5 for 24 seconds.
- Wait ten minutes for the indophenol reaction to proceed.
- Insert the device into the custom photo-spectrometer to acquire absorbance data.
- Compare the measured absorbance to a standard curve to determine unknown ammonia concentration.

Equivalents

[0268] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the disclosure described herein. The scope of the present disclosure is not intended to be limited to the above Description, but rather is as set forth in the following claims:

SEQUENCE LISTING**[0269]**

<110> The University of Maryland, College Park
Children's National Medical Center
Ayyub, Omar Bilal
Behrens, Adam Michael
Kofinas, Peter
Summar, Marshall Lynn
Cabrera-Luque, Juan Manuel
Cunningham, Gary

<120> DEVICE AND METHODS OF USING DEVICE FOR DETECTION OF HYPERAMMONEMIA

<130> P103171EP-WO/TCL

<140> EP14841207.5

<141> 2014-09-02

<150> PCT/US14/53756

<151> 2014-09-02

<150> US 61/872,149

<151> 2013-08-30

<160> 24

<170> PatentIn version 3.5

<210> 1

<211> 378

<212> PRT

<213> Artificial Sequence

<220>

<223> Dehydrogenase mutant 1

<400> 1

Met Asn Thr Val Thr Asn Gln Trp Lys Ala Val Asp Ile Phe Thr Gln
1 5 10 15

Ile Arg Asp His Glu Gln Val Val Phe Cys Asn Asp Lys Asn Thr Gly
20 25 30

Leu Lys Ala Ile Ile Ala Ile His Asp Thr Thr Leu Gly Pro Ala Leu
35 40 45

Gly Gly Cys Arg Met Tyr Pro Tyr Ala Thr Val Glu Asp Ala Leu Phe
50 55 60

Asp Val Leu Arg Leu Ser Lys Gly Met Thr Tyr Lys Cys Leu Ala Ala
65 70 75 80

Asp Val Asp Phe Gly Gly Gly Lys Ala Val Ile Ile Gly Asp Pro His
85 90 95

Lys Asp Lys Thr Pro Glu Leu Phe Arg Ala Phe Gly Gln Phe Val Glu
100 105 110

Ser Leu Asn Gly Arg Phe Tyr Thr Gly Thr Asp Met Gly Thr Thr Pro
115 120 125

Asp Asp Phe Val His Ala Met Lys Glu Thr Asn Cys Ile Val Gly Val
130 135 140

Pro Glu Glu Tyr Gly Gly Ser Gly Asp Ser Ser Val Pro Thr Ala Leu
145 150 155 160

Gly Val Ile Tyr Gly Ile Gln Ala Thr Asn Lys Val Ile Trp Gly Ser
165 170 175

Asp Glu Leu His Gly Lys Thr Tyr Ala Ile Gln Gly Leu Gly Lys Val
180 185 190

Gly Arg Lys Val Ala Glu Arg Leu Leu Lys Glu Gly Ala Asp Leu Tyr
195 200 205

Val Cys Asp Ile His Pro Thr Ala Ile Glu Ala Ile Val Ser Tyr Ala
210 215 220

Lys Lys Leu Gly Ala Asn Val Lys Val Val Gln Gly Thr Glu Ile Tyr
225 230 235 240

Arg Thr Asp Ala Asp Ile Phe Val Pro Cys Ala Phe Gly Asn Val Val
245 250 255

Asn Asp Asn Thr Ile His Val Leu Lys Val Lys Ala Ile Val Gly Ser
260 265 270

Ala Asn Asn Gln Leu Leu Asp Val Arg His Gly Gln Leu Leu Lys Glu
275 280 285

Lys Gly Ile Leu Tyr Ala Pro Asp Tyr Ile Val Asn Ala Gly Gly Leu
290 295 300

Ile Gln Val Ala Asp Glu Leu Tyr Gly Leu Asn Lys Glu Arg Val Leu
305 310 315 320

Gln Lys Thr Lys Ala Ile Tyr Ser Thr Leu Leu His Ile Tyr Ser Arg
325 330 335

Ala Glu Ala Asp His Ile Thr Thr Ile Glu Ala Ala Asn Arg Phe Cys
340 345 350

Glu Glu Arg Leu Gln Gln Arg Ser Arg Arg Asn Asp Phe Phe Thr His
355 360 365

Arg Lys Gln Pro Lys Trp Asp Ile Arg Arg
370 375

<210> 2

<211> 366

<212> PRT

<213> Artificial Sequence

<220>

<223> Mutant Enzyme 2

<400> 2

Met Arg Asp Val Phe Glu Met Met Asp Arg Tyr Gly His Glu Gln Val
1 5 10 15

Ile Phe Cys Arg His Pro Gln Thr Gly Leu Lys Ala Ile Ile Ala Leu
20 25 30

His Asn Thr Thr Ala Gly Pro Ala Leu Gly Gly Cys Arg Met Ile Pro
35 40 45

Tyr Ala Ser Thr Asp Glu Ala Leu Glu Asp Val Leu Arg Leu Ser Lys
50 55 60

Gly Met Thr Tyr Lys Cys Ser Leu Ala Asp Val Asp Phe Gly Gly Gly
65 70 75 80

Lys Met Val Ile Ile Gly Asp Pro Lys Lys Asp Lys Ser Pro Glu Leu
85 90 95

Phe Arg Val Ile Gly Arg Phe Val Gly Gly Leu Asn Gly Arg Phe Tyr
100 105 110

Thr Gly Thr Asp Met Gly Thr Asn Pro Glu Asp Phe Val His Ala Ala
115 120 125

Arg Glu Ser Lys Ser Phe Ala Gly Leu Pro Lys Ser Tyr Gly Gly Lys
130 135 140

150 155 160
 Gly Asp Thr Ser Ile Pro Thr Ala Leu Gly Val Phe His Gly Met Arg
 145 150 155 160
 Ala Thr Ala Arg Phe Leu Trp Gly Thr Asp Gln Leu Lys Gly Arg Val
 165 170 175
 Val Ala Ile Gln Gly Val Gly Lys Val Gly Glu Arg Leu Leu Gln Leu
 180 185 190
 Leu Val Glu Val Gly Ala Tyr Cys Lys Ile Ala Asp Ile Asp Ser Val
 195 200 205
 Arg Cys Glu Gln Leu Lys Glu Lys Tyr Gly Asp Lys Val Gln Leu Val
 210 215 220
 Asp Val Asn Arg Ile His Lys Glu Ser Cys Asp Ile Phe Ser Pro Cys
 225 230 235 240
 Ala Lys Gly Gly Val Val Asn Asp Asp Thr Ile Asp Glu Phe Arg Cys
 245 250 255
 Leu Ala Ile Val Gly Ser Ala Asn Asn Gln Leu Val Glu Asp Arg His
 260 265 270
 Gly Ala Leu Leu Gln Lys Arg Ser Ile Cys Tyr Ala Pro Asp Tyr Leu
 275 280 285
 Val Asn Ala Gly Gly Leu Ile Gln Val Ala Asp Glu Leu Glu Gly Phe
 290 295 300
 His Glu Glu Arg Val Leu Ala Lys Thr Glu Ala Ile Tyr Asp Met Val
 305 310 315 320
 Leu Asp Ile Phe His Arg Ala Lys Asn Glu Asn Ile Thr Thr Cys Glu
 325 330 335
 Ala Ala Asp Arg Ile Val Met Glu Arg Leu Lys Lys Leu Thr Asp Ile
 340 345 350
 Arg Arg Ile Leu Leu Glu Asp Pro Arg Asn Ser Ala Arg Arg
 355 360 365

<210> 3

<211> 1068

<212> DNA

<213> Artificial Sequence

<220>

<223> nucleic acid encoding enzyme 3

<400> 3

atggaaatct tcgaggaaat caaacggcgg ggacacgagc aaattctgtt caattatgat 60
 cgggcttccg gtttgaaagc aattatcgcc attcacaata ctacgttggg gccggcgttg 120
 ggcggtgcc gaatgttacc gtatcaaacg gaagaggcgg ccctcgagga tgcgctgcgg 180
 ttgtcggaag ggatgaccta taaagcggcc gccgccgggc tcgatttcgg cgggggcaaa 240
 acggtgatta tcggggatcc gatgaaagac aagtccgagg ccctgtttcg tgcgctcggg 300
 cgttttatcg agaccttgaa aggcggttac cttacgggag aagacgtagg aaccaacgaa 360
 gaagattttg tctgggctcg tcgggaaacc cgttatgttg tcggattgcc gccggcttat 420
 ggcggtccg gcgatacggg tgacaatacc gcgcgcggcg tcattcaagc gatgcgcgcc 480
 gcgttgatgc accggtacgg ttccgccgat ctccagggcc gccggattgc cgtccaaggg 540

ctgggcaaag taggctatca tgtggcgcga cgggcatcg aggcgggcgc tcgagtgatt	600
gcgcccgata tcaatccgca tgtagtcggc cgagtggcgt ccgcttgggg gattgaagcc	660
accgatccgt gggctgtggt ggaaaccccc tgcgatatct tcgccccctg tgcgttgggt	720
aacgtcatta cggaacggac cgtgtccgcc ctccaatgtc aggtggtggc cggttcggcc	780
aacaatcagc tggcggatga tcgactggcc gatgatttag ctgcccgcg cttctctat	840
gcgccggatt ttattgcgaa tgccggcgga ttgattcagg tggcggatga aattcgggga	900
tatcatgaag aacgggtccg tcatcaaata gacgggattt atgacgtcct gctcgagatt	960
tttcggaagg cggacgcctc cggccgatca accgtggcgg ttgcggtaga cgaggcgcgt	1020
cgcggtttgg acaccattca ggccatccac cgcctgtacg gatcatag	1068

<210> 4

<211> 3939

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid encoding enzyme 4

<400> 4

ctgcaggatca acggatcata ttctacacat atataatgca ctccaattga cataatacat	60
aacgtgacat atgatacatt tattaatatt aattgtcaca ttacacttc acatattaaa	120
atactctcgt atgaatgcaa ttgaaacat attttaaatt aattgattga tatatattga	180
acaaaacctc acaaaaatgc accctcttgg ttcacaaaga aactttcttc tattttctac	240
ttattttctgc tagtgtcttt cctattcaaa gccatcattt ccatcaacct tcacaatacc	300
atgtttaaaa agtcattaaa aatcaatttt taaatagaa aaaaacaaga agatggaaat	360
cacttggttg gtactatata tttagttggt aagtttgact cataccgtgt attgaccaat	420
ataaataaaa tcttatttca aataaattca aaagtccaat aaatatatat tcgttcataa	480
cttataataa aattgattat acatagtcct ccccatctca cttttactga tcaattattt	540
ctaaaatata ttattacttt tacttggtat ttttaataaa ttaagaaaat ataatactcc	600
cttcgttttt aaaaaatac ctagtttgac ttgaaacgga gtttaataaa agaaagaaga	660
cttgttaatc ttgtgattct aaattaaagt tatgtcaaat gtaccaaaat gtcctttaat	720
cttgtggtct taaacatgtc acatgaaaaa ttaaagtgtt tccaaaaaa gaaaggggtc	780
aatgtcattc ttttttaaac agactaaaaa agaaataaac tcattctttt tgaaacggag	840
agagtaattt ttccacggtt ttactcatta atattaaata ttattctcta gatcatccta	900
taagatctaa tagtggacat caattaatac ctatgtcact tattattatt ttaataattg	960
tatcaagtca aataataaca agtaaaaatg gactacctac tattaatctt caacaaccac	1020
aatttactag ttttttctca gcaacccctc ctcacatatt tcaccattta ctggtttttt	1080
cctagcaacc ccctctcaca tattttgttt accaaccatc atttggtcct ctatatatac	1140
tcaccacatg atagatacat atatatacca caacccaaaac aaaagggttt ataagttcac	1200
aacatttttt atatacatc aaataaactc taaccatttt ctcttcacta aaatttcttc	1260
attacaaatc taacaattta cttgatccaa tggcaccatc aattgcacaa aatggacata	1320
ttaatggaga agtagctatg gatttgtgca agaaatcaat caatgatcca ttgaattggg	1380
aaatggctgc tgattcttta agaggcagcc atttggtatg agtgaaaaag atgggtggatg	1440
aatttagaaa gccaatgtg aaacttgggg gtgaaacttt gtcagttgca caagttgcat	1500
ccattgcaaa tgttgatgac aaaagtaatg gggttaaagt ggaactttct gaaagtgcaa	1560
gggctggtgt gaaagctagt agtgattggg ttatggatag tatgagtaaa ggtacagata	1620
gttatgggtg tactgctgga ttggagcaa catctcatag aagaacaaaa aatggtggtg	1680
ctcttcaaaa agaacttatt agttaacaa actatttttt ttctttatat atactaacaa	1740

```

cccccaaaa agaacctatc aggcacaaca acctaccccc ccggtatata atactaaca 1740
tgtaaagaat ttaatatattt ttgtttatat atactaaca tgtaaaaaat ttaatatattt 1800
tttgtttatat atactaaca tgtaaagaat ttaatatattt ttgtttatata atagcttata 1860
gactacttaa gtgctccatt gataaagatt tttttttgtt ttacgcgaa ggggattcgg 1920
atgaattcag ttaaaatgtg atcttaatga attatgatat tttttttagt gttcttgaat 1980
gctggagttt ttggtaatgg aatagaatca ttccacacat tgccacattc agcaacaagg 2040
gcagctatgc ttgttaggat caacactctg cttcaaggct actctggcat tagatttgag 2100
atcttggaag caatcactaa gttgatcaat agcaacatca ccccggttt gcctctccgt 2160
ggcacgatca ctgcctcggg tgatctcgtc cttttgtcct atattgctgg ttgctcact 2220
ggcagacctt attccaaggc tgttggaacc aatggtgaga aacttaatgc tgaggaagct 2280
ttctgcgtgg ctggtattag tgggtgattt ttcgagttgc agcctaagga aggacttgca 2340
cttgtgaatg gcacagcagt tggttctgct atggcatcaa tagtctgtt tgagtccaat 2400
atctttgctg ttatgtctga agttttatca gcgattttta ctgaagtgat gaacggaaag 2460
cccgattca ctgactattt gacacacaag ttgaagcatc accctggtca gattgaggct 2520
gctgctatta tggaacacat ttggatgga agctcttatg tgaaggtagc tcagaagctc 2580

catgaaatgg atcctcttca aaaaccaaag caagatcgtt atgctctccg aacatctcca 2640
caatggcttg gacctcagat tgaagtcatt cgtgctgcaa ctaagatgat cgagagggag 2700
attaactcag tgaacgaca tccattgatc gatgtttcaa gaaacaaggc cttacatggt 2760
ggcaacttcc aaggaacccc tattggtgtc tccatggata atacaagatt ggcccttgca 2820
tcaattggta aattgatgtt tgcccaattc tcagagcttg tcaacgacta ttacaacaac 2880
gggttgccat ctaatctgac agcaggaagg aatccaagct tggactatgg tttcaagggc 2940
gctgaaatcg cgatggcttc ttactgctcg gaacttcaat tcttggcaaa tccagtgaat 3000
aaccatgtct aaagtgtga gcaacacaac caagatgtga attccttggg ctttaatttca 3060
gccaggaaaa cagctaaggc tgttgatata ttgaagataa tgtcatcaac ctatctcgtg 3120
gctctttgcc aagctattga cttacgacat ttggaggaaa acttgaagag tgttgtcaag 3180
aacacagtta gccaaagtagc taagagaact ttgacaatgg gtgctaattg tgaacttcat 3240
ccagcaagat tcagcgaaaa agaattgctt cgagtcgtgg atagagaata cttgtttgcc 3300
tatgctgatg atccctgcag ctccaactac cttttgatgc agaagctgag acaagtcctt 3360
gttgatcaag caatgaagaa tggtgaaagt gagaagaatg tcaacagctc aatcttccaa 3420
aagattggag ctttcgagga cgaattaatc gctgtgttgc ctaaagaagt tgagagtgt 3480
agagctgttt ttgaaagtgg caacccttta attcgtaca ggatcacaga atgcagatca 3540
tatccattgt acaggttggg gagagaagaa cttggaacag aattgttgac gggtgaaaaa 3600
gttcgatcac ctggtgagga gattgataaa gtgtttacag caatatgtaa tggacagatt 3660
attgatccat tgttgagtg tctgaagagc tggaaatggtg ctctcttcc aatctgctaa 3720
atgtgttatt ctttcaagtt ctttttttgt accttttagt gaattactag aattataatg 3780
atgttatgaa cttatattaa aaaaaatat ttttgactat aaaatttagt tttgttattg 3840
aaattaaagg ctcaatctgt gttctttcct tctgttatct gaatattata agaattcaag 3900
taatctttta gctttgtgaa catgatgaca tgctttctt 3939

```

<210> 5

<211> 1521

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 5

<400> 5

atgatcacgc ttacccccgg ccacctgacc ctcccgcaac tgcgccagat cgcgcgcgag	60
cccgtgcagc tgacgttga tccggccagc ttcgtgaaga tcgacgcggg cgcgaaggcc	120
gtgtccgaca tcgccgcgaa gggcgagccg gcgtacggca tcaacacggg cttcggtcgt	180
ctggccagca cgcataatcc gcacgatcag ctccaattgc tgcagaagaa cctcgtgctg	240
tgcgatgcag tcggtgtcgg cgagccgatg gcgcgttcgt cgggtgcgtct gctgatcgcg	300
ctgaagctgt cgagcctcgg ccgcggccat tcgggcatte gccgcgaagt gatggacggg	360
ctgatcaagc tgttcaacgc cgacgtgctg ccgctgatte cgggtgaaggg ctcggtcggc	420
gcacgcggcg acctcgcgcc gctcgcgcac atgtcggccg tgcgtgctcg cgtcggcgaa	480
gtgttcattc gcggcgagcg cgcgagcgcg gtggacgggt tgcgcgtcgc gggcctcgcg	540
ccgctgacgc tgcaggcgaa ggaaggcctc gcgctgctga accgtacgca ggcgtcgacg	600
gcgctcgcgc tcgacaacct gttcgcgac gaagacctgt accgcacggc gctcgtcgcc	660
ggcgcgctgt cggctgatgc ggcggccggc tcggtgaagc cgttcgacgc gcgcatccac	720
gaactgcgcg gccatcgcgg ccagatcgat gcggcgggccg cgtatcgca gctgctcgaa	780
ggctcggcga tcaacctctc gcacgcgcac tgcggcaagg tgcaggatcc gtacagcctg	840
cgtgccagc cgcagtgat gggcgctgc ctggaccaga tgcgtcatgc ggccgacgtg	900
ctgctcgtcg aggcgaacgc ggtatcggac aaccgctga tcttcccga taccggcgaa	960
gtgctgtcgg gcggcaattt ccatgcggag cccgtcgcgt tcggggccga caacctcgcg	1020
ctcgcggctg cggaaatcgg cgcgctggcc gagcgccga tcgcgctgct gatcgacgcg	1080
acgctgtcgg gcctgccgcc gttcctcgtg aaggatggcg gcgtgaactc gggcttcacg	1140
attgcgcacg tgacggcagc tgcgctcgca tcggagaaca agacgctcgc gcacccggcg	1200
tcggtcgatt cgtgccgac ctccggcaac caggaagacc acgtgtcgat ggcgacgttc	1260
gcggcacgca agctggccga catcgccgac aacacgaagc acatcctcgc gatcgaaactg	1320
ctcgcggccg cgcaggcgt cgatctcgc gagaacgaga cgagcccgaa gctcgcggaa	1380
gtgatgaaga cgattcgcag caaggtcgcg cattacgagc tcgaccacta ctttgcgcgcg	1440
gacatcgccg tgatcgcgaa gctcgtcgtc gagcgcgct tcgcgaagca ctgcccgttc	1500
gccttcgcat cggagcagta a	1521

<210> 6

<211> 1533

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid Sequence encoding enzyme 6

<400> 6

gtgacgcagg tcgtggaacg tcaggctgat cggctcagca gcaggagta cctggcccg	60
gtcgtgcgca gcgccgggtg ggacgccggt ctcacctcgt gcaccgacga ggagatcgtc	120
cggatgggcg cgagcgcgcg caccatcgag gagtacctga agtccgaaa gccatctac	180
ggcctgacgc agggcttcgg tccgctgggt ctgttcgacg ccgactcgga gctggagcag	240
ggcggctcgc tgatctcgca cctgggcacc ggccaggggc gccactggc cccggagggtg	300
tgcgggtga tcctctggct gcgcatccag aacatgcgca aggggtactc ggcggtctcg	360
ccggtgttct ggcagaagct cgcgcacctg tggaaacaagg ggttcacccc ggcgatcccc	420
cggcacggca cggtcagcgc gagcggcgac ctgcaaccgc tggcgacgc cgcgctcgcc	480
ttcaccgggt tcggcgaggc gtggacccgg gacgccgacg gccgggtggc caccgtgccg	540
gccgtggacg cgctcgccgc gctggggggc gagccgttcg actggccggt gcgcgaggcg	600

```

ctggcggttcg tcaacgggac cggcgcgagc ctgcggtgg ctgtgctcaa ccaccggtcc 660
gccctgcggc tggccgcgc ctgcgccgtg ctctccgcgc ggctggcgac cctgctgggg 720
gccaatcccg agcactacga cgtggggcac ggtgtcgcgc gcggccaggt cggtcagctg 780
accgcggcgg agtggatccg gcaggggctg ccccggggca tggcgcgcga cggcagccgc 840
ccgctccagg agccgtacag cctgcgggtg gcgccgcagg tgctcggcgc ggtgctcgac 900
cagctcgacg gcgcgggcga cgtgctggcg cgggaggtcg acggctgccg ggacaaccgg 960
atcacctacg agggcgagct gctgcacggc ggcaacttcc acgccatgcc ggtgggtttc 1020
gcctccgacc agatcgggtt ggccatgcac atggccgcct acctggccga gcgccagctg 1080
ggtctgctgg tcagcccggg gaccaacggc gacctgccgc ccattgctcac cccgcgcgcc 1140
gggcgcgggt cgggctggc cggggtgcag atcagcgcga cctcgttcgt ctgcggatc 1200
cggcagctgg tgttccccgc ctgcgtgacc acctgcccga ccaacggctg gaaccaggac 1260
cacgtgccga tggcgtcaa cggggcgaac tcggtgttcg aggcgttgga gctcggctgg 1320
ctgacggctg gtcgctggc ggtgggcgtc gcgcagctcg cggccatgac cggccacgcc 1380
gcggagggcg tctgggcgga gctggccggg atctgccgc cgctggacgc cgaccgccg 1440
ctgggcgccg aggtgcgcgc cgcgcgcgac ctgctgtccg cgcacgcgga ccaactgctc 1500
gtcgacgagg cagacgggaa ggatttcgga tga 1533

```

<210> 7

<211> 721

<212> PRT

<213> Solanum lycopersicum

<400> 7

```

Met Ala Ser Ser Ile Val Gln Asn Gly His Val Asn Gly Glu Ala Met
1           5           10          15

Asp Leu Cys Lys Lys Ser Ile Asn Val Asn Asp Pro Leu Asn Trp Glu
20          25          30

Met Ala Ala Glu Ser Leu Arg Gly Ser His Leu Asp Glu Val Lys Lys
35          40          45

Met Val Asp Glu Phe Arg Lys Pro Ile Val Lys Leu Gly Gly Glu Thr
50          55          60

Leu Thr Val Ala Gln Val Ala Ser Ile Ala Asn Val Asp Asn Lys Ser
65          70          75          80

Asn Gly Val Lys Val Glu Leu Ser Glu Ser Ala Arg Ala Gly Val Lys
85          90          95

Ala Ser Ser Asp Trp Val Met Asp Ser Met Gly Lys Gly Thr Asp Ser
100         105         110

Tyr Gly Val Thr Thr Gly Phe Gly Ala Thr Ser His Arg Arg Thr Lys
115         120         125

Asn Gly Gly Ala Leu Gln Lys Glu Leu Ile Arg Phe Leu Asn Ala Gly
130         135         140

Val Phe Gly Asn Gly Thr Glu Ser Ser His Thr Leu Pro His Ser Ala
145         150         155         160

Thr Arg Ala Ala Met Leu Val Arg Ile Asn Thr Leu Leu Gln Gly Tyr
165         170         175

Ser Gly Ile Arg Phe Glu Ile Leu Glu Ala Ile Thr Lys Leu Ile Asn
180         185         190

Ser Asn Ile Thr Pro Cys Leu Pro Leu Arg Gly Thr Ile Thr Ala Ser

```

195					200					205					
Gly 210	Asp	Leu	Val	Pro	Leu	Ser 215	Tyr	Ile	Ala	Gly	Leu 220	Leu	Thr	Gly	Arg
Pro 225	Asn	Ser	Lys	Ala	Val 230	Gly	Pro	Asn	Gly	Glu 235	Lys	Leu	Asn	Ala	Glu 240
Glu	Ala	Phe	Arg	Val 245	Ala	Gly	Val	Thr	Ser 250	Gly	Phe	Phe	Glu	Leu 255	Gln
Pro	Lys	Glu	Gly 260	Leu	Ala	Leu	Val	Asn 265	Gly	Thr	Ala	Val	Gly 270	Ser	Gly
Met	Ala	Ser 275	Met	Val	Leu	Phe	Glu 280	Ser	Asn	Ile	Leu	Ala 285	Val	Met	Ser
Glu 290	Val	Leu	Ser	Ala	Ile	Phe 295	Ala	Glu	Val	Met	Asn 300	Gly	Lys	Pro	Glu
Phe 305	Thr	Asp	Tyr	Leu	Thr 310	His	Lys	Leu	Lys	His 315	His	Pro	Gly	Gln	Ile 320
Glu	Ala	Ala	Ala	Ile 325	Met	Glu	His	Ile	Leu 330	Asp	Gly	Ser	Ser	Tyr 335	Val
Lys	Ala	Ala	Gln 340	Lys	Leu	His	Glu	Met 345	Asp	Pro	Leu	Gln	Lys 350	Pro	Lys
Gln	Asp 355	Arg	Tyr	Ala	Leu	Arg	Thr 360	Ser	Pro	Gln	Trp	Leu 365	Gly	Pro	Gln
Ile 370	Glu	Val	Ile	Arg	Ala	Ala 375	Thr	Lys	Met	Ile	Glu 380	Arg	Glu	Ile	Asn
Ser 385	Val	Asn	Asp	Asn	Pro 390	Leu	Ile	Asp	Val	Ser 395	Arg	Asn	Lys	Ala	Leu 400
His	Gly	Gly	Asn 405	Phe	Gln	Gly	Thr	Pro	Ile 410	Gly	Val	Ser	Met	Asp 415	Asn
Thr	Arg	Leu	Ala 420	Leu	Ala	Ser	Ile	Gly 425	Lys	Leu	Met	Phe	Ala	Gln	Phe
Ser	Glu	Leu 435	Val	Asn	Asp	Tyr	Tyr 440	Asn	Asn	Gly	Leu	Pro 445	Ser	Asn	Leu
Thr	Ala 450	Gly	Arg	Asn	Pro	Ser 455	Leu	Asp	Tyr	Gly	Leu 460	Lys	Gly	Ala	Glu
Ile 465	Ala	Met	Ala	Ser	Tyr 470	Cys	Ser	Glu	Leu	Gln 475	Phe	Leu	Ala	Asn	Pro 480
Val	Thr	Asn	His 485	Val	Gln	Ser	Ala	Glu	Gln 490	His	Asn	Gln	Asp	Val 495	Asn
Ser	Leu	Gly	Leu 500	Ile	Ser	Ala	Arg	Lys 505	Thr	Ala	Glu	Ala	Val 510	Asp	Ile
Leu	Lys	Leu 515	Met	Ser	Ser	Thr	Tyr 520	Leu	Val	Ala	Leu	Cys 525	Gln	Ala	Ile
Asp	Leu 530	Arg	His	Leu	Glu	Glu 535	Asn	Leu	Arg	Ser	Ala 540	Val	Lys	Asn	Thr
Val	Ser	Gln	Val	Ala	Lys	Arg	Thr	Leu	Thr	Met	Gly	Ala	Asn	Gly	Glu
545					550						555				560

Leu His Pro Ala Arg Phe Cys Glu Lys Glu Leu Leu Arg Val Val Asp
565 570 575

Arg Glu Tyr Val Phe Ala Tyr Ala Asp Asp Pro Cys Ser Ser Thr Tyr
580 585 590

Pro Leu Met Gln Lys Leu Arg Gln Val Leu Val Asp His Ala Met Lys
595 600 605

Asn Gly Glu Ser Glu Lys Asn Val Asn Ser Ser Ile Phe Gln Lys Ile
610 615 620

Val Ala Phe Glu Asp Glu Leu Lys Ala Val Leu Pro Lys Glu Val Glu
625 630 635 640

Ser Ala Arg Ala Val Val Glu Ser Gly Asn Pro Ala Ile Pro Asn Arg
645 650 655

Ile Thr Glu Cys Arg Ser Tyr Pro Leu Tyr Arg Leu Val Arg Gln Glu
660 665 670

Leu Gly Ser Glu Leu Leu Thr Gly Glu Lys Val Arg Ser Pro Gly Glu
675 680 685

Glu Ile Asp Lys Val Phe Thr Ala Met Cys Asn Gly Gln Ile Ile Asp
690 695 700

Pro Leu Leu Glu Cys Leu Lys Ser Trp Asn Gly Ala Pro Leu Pro Ile
705 710 715 720

Cys

<210> 8

<211> 1101

<212> DNA

<213> Thermoactinomyces intermedius

<400> 8

atgcgcgacg tgtttgaaat gatggaccgc tatggccacg agcaggtcat tttttgccgt	60
catccgcaaa ccggtctcaa agcgatcatc gccttgcata atacaaccgc ggggccggct	120
ttgggtggat gccgcatgat cccgtatgct tcgacggacg aagccttgga ggatgttttg	180
cggttgtcca aaggcatgac ctataaatgc agtctggcgg atgtggactt tggcggggga	240
aaaatggtta tcatcggcga tccgaaaaaa gataaatcgc cggagttggt tcgcgtgatc	300
ggcgtttttg tgggcggggt aaacggccgt ttctataccg gaaccgacat gggaaccaat	360
cgggaagatt ttgtccatgc cgccaggga tcgaaatctt ttgccggatt gccgaaatcg	420
tacggcggaa agggggacac atccattccc accgcgctcg ggggtgtttca cggaatgcgg	480
gccaccgccc ggtttttatg ggggacggat cagctgaaag ggcgtgtggt tgccatccaa	540
ggagtccgca aggtgggaga gcgcttggtg cagcttttgg tcgaagtggg ggcttactgc	600
aaaattgccg acatcgattc ggtgcgatgc gaacagctga aagaaaagta tggcgacaag	660
gtccaattgg tggatgtgaa ccggattcac aaggagagtt gcgatatttt ctgccttgc	720
gccaaaggcg gcgtggtcaa tgatgacacc attgacgagt tccgttgcct ggccattgtc	780
ggatccgcca acaaccaact ggtggaagac cggcatgggg cactgcttca aaaacggagc	840
at ttgttatg cacccgatta tctggtgaat gccggcgggc tgattcaagt ggctgatgaa	900
ctggaaggct tccatgaaga gagagtgctc gccaaaaccg aagcgattta tgacatggtc	960
ctggatattt ttcaccgggc gaaaaatgag aatattacca cttgtgaggc agcggaccgg	1020
atcgtgatgg agcgtttgaa aaagttaacc gatattcgcc ggatcttggt ggaggatccc	1080

cgcaacagcg caaggaggta a

1101

<210> 9

<211> 352

<212> PRT

<213> Caenorhabditis remanei

<400> 9

Met	Asp	Phe	Lys	Ala	Lys	Leu	Leu	Ala	Glu	Met	Ala	Lys	Lys	Arg	Lys
1			5					10						15	

Ala	Val	Ser	Gly	Leu	Glu	Val	Lys	Glu	Gly	Gly	Ala	Lys	Phe	Val	Arg
			20				25						30		

Gly	Ala	Asp	Leu	Glu	Ser	Lys	Arg	Thr	Gln	Glu	Tyr	Glu	Ala	Lys	Gln
	35					40					45				

Glu	Glu	Leu	Ala	Ile	Lys	Lys	Arg	Lys	Ala	Asp	Asp	Glu	Ile	Leu	Gln
	50				55					60					

Glu	Ser	Thr	Ser	Arg	Ala	Lys	Ile	Val	Pro	Glu	Val	Pro	Glu	Ala	Glu
65				70					75					80	

Phe	Asp	Glu	Lys	Thr	Pro	Met	Pro	Glu	Ile	His	Ala	Arg	Leu	Arg	Gln
			85					90						95	

Arg	Gly	Gln	Pro	Ile	Leu	Leu	Phe	Gly	Glu	Ser	Glu	Leu	Ser	Val	Arg
			100				105						110		

Lys	Arg	Leu	His	Gln	Leu	Glu	Ile	Glu	Gln	Pro	Glu	Leu	Asn	Glu	Gly
	115					120						125			

Trp	Glu	Asn	Glu	Met	Gln	Thr	Ala	Met	Lys	Phe	Ile	Gly	Lys	Glu	Met
	130				135						140				

Asp	Lys	Ala	Val	Val	Glu	Gly	Thr	Ala	Asp	Ser	Ala	Thr	Arg	His	Asp
145					150				155						160

Ile	Ala	Leu	Pro	Gln	Gly	Tyr	Glu	Glu	Asp	Asn	Trp	Lys	Ser	Ile	Glu
			165					170						175	

His	Ala	Ser	Thr	Leu	Leu	Gly	Val	Gly	Asp	Glu	Met	Lys	Arg	Asp	Cys
		180					185						190		

Asp	Ile	Ile	Leu	Ser	Ile	Cys	Arg	Tyr	Ile	Leu	Ala	Arg	Trp	Ala	Arg
	195					200						205			

Asp	Leu	Asn	Asp	Arg	Pro	Leu	Asp	Val	Lys	Lys	Thr	Ala	Gln	Gly	Met
	210				215						220				

His	Glu	Ala	Ala	His	His	Lys	Gln	Thr	Thr	Met	His	Leu	Lys	Ser	Leu
225				230						235					240

Met	Thr	Ser	Met	Glu	Lys	Tyr	Asn	Val	Asn	Asn	Asp	Ile	Arg	His	His
			245					250						255	

Leu	Ala	Lys	Ile	Cys	Arg	Leu	Leu	Val	Ile	Glu	Arg	Asn	Tyr	Leu	Glu
			260				265						270		

Ala	Asn	Asn	Ala	Tyr	Met	Glu	Met	Ala	Ile	Gly	Asn	Ala	Pro	Trp	Pro
	275					280						285			

Val	Gly	Val	Thr	Arg	Ser	Gly	Ile	His	Gln	Arg	Pro	Gly	Ser	Ala	Lys
	290					295					300				

Ala	Tyr	Val	Ser	Asn	Ile	Ala	His	Val	Leu	Asn	Asp	Glu	Thr	Gln	Arg
305				310						315					320

Lys Tyr Ile Gln Ala Phe Lys Arg Leu Met Thr Lys Leu Gln Glu Tyr
 325 330 335

Phe Pro Thr Asp Pro Ser Lys Ser Val Glu Phe Val Lys Lys Ser Val
 340 345 350

<210> 10

<211> 411

<212> PRT

<213> Arabidopsis thaliana

<400> 10

Met Asn Ala Leu Ala Ala Thr Asn Arg Asn Phe Lys Leu Ala Ala Arg
 1 5 10 15

Leu Leu Gly Leu Asp Ser Lys Leu Glu Lys Ser Leu Leu Ile Pro Phe
 20 25 30

Arg Glu Ile Lys Val Glu Cys Thr Ile Pro Lys Asp Asp Gly Thr Leu
 35 40 45

Ala Ser Phe Val Gly Phe Arg Val Gln His Asp Asn Ala Arg Gly Pro
 50 55 60

Met Lys Gly Gly Ile Arg Tyr His Pro Glu Val Asp Pro Asp Glu Val
 65 70 75 80

Asn Ala Leu Ala Gln Leu Met Thr Trp Lys Thr Ala Val Ala Lys Ile
 85 90 95

Pro Tyr Gly Gly Ala Lys Gly Gly Ile Gly Cys Asp Pro Ser Lys Leu
 100 105 110

Ser Ile Ser Glu Leu Glu Arg Leu Thr Arg Val Phe Thr Gln Lys Ile
 115 120 125

His Asp Leu Ile Gly Ile His Thr Asp Val Pro Ala Pro Asp Met Gly
 130 135 140

Thr Gly Pro Gln Thr Met Ala Trp Ile Leu Asp Glu Tyr Ser Lys Phe
 145 150 155 160

His Gly Tyr Ser Pro Ala Val Val Thr Gly Lys Pro Ile Asp Leu Gly
 165 170 175

Gly Ser Leu Gly Arg Asp Ala Ala Thr Gly Arg Gly Val Met Phe Gly
 180 185 190

Thr Glu Ala Leu Leu Asn Glu His Gly Lys Thr Ile Ser Gly Gln Arg
 195 200 205

Phe Val Ile Gln Gly Phe Gly Asn Val Gly Ser Trp Ala Ala Lys Leu
 210 215 220

Ile Ser Glu Lys Gly Gly Lys Ile Val Ala Val Ser Asp Ile Thr Gly
 225 230 235 240

Ala Ile Lys Asn Lys Asp Gly Ile Asp Ile Pro Ala Leu Leu Lys His
 245 250 255

Thr Lys Glu His Arg Gly Val Lys Gly Phe Asp Gly Ala Asp Pro Ile
 260 265 270

Asp Pro Asn Ser Ile Leu Val Glu Asp Cys Asp Ile Leu Val Pro Ala
 275 280 285

Ala Leu Gly Gly Val Ile Asn Arg Glu Asn Ala Asn Glu Ile Lys Ala
290 295 300

Lys Phe Ile Ile Glu Ala Ala Asn His Pro Thr Asp Pro Asp Ala Asp
305 310 315 320

Glu Ile Leu Ser Lys Lys Gly Val Val Ile Leu Pro Asp Ile Tyr Ala
325 330 335

Asn Ser Gly Gly Val Thr Val Ser Tyr Phe Glu Trp Val Gln Asn Ile
340 345 350

Gln Gly Phe Met Trp Glu Glu Glu Lys Val Asn Asp Glu Leu Lys Thr
355 360 365

Tyr Met Thr Arg Ser Phe Lys Asp Leu Lys Glu Met Cys Lys Thr His
370 375 380

Ser Cys Asp Leu Arg Met Gly Ala Phe Thr Leu Gly Val Asn Arg Val
385 390 395 400

Ala Gln Ala Thr Ile Leu Arg Gly Trp Gly Ala
405 410

<210> 11

<211> 1275

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid Sequence encoding enzyme 11

<400> 11

atgtcagcaa agcaagtctc gaaagatgaa gaaaaagaag ctcttaactt atttctgtct	60
acccaaacaa tcattaagga agcccttcgg aagctgggtt atccgggaga tatgtatgaa	120
ctcatgaaag agccgcagag aatgctcact gtccgcattc cggtcaaaat ggacaatggg	180
agcgtcaaag tggtcacagg ctaccggtca cagcacaatg atgctgtcgg tccgacaaag	240
gggggcggtc gcttccatcc agaagttaat gaagaggaag taaaggcatt atccatttgg	300
atgacgctca aatgcgggat tgccaatctt ccttacggcg gcgggaaggg cggtattatt	360
tgtgatccgc ggacaatgtc atttggagaa ctggaaaggc tgagcagggg gtatgtccgt	420
gccatcagcc agatcgtcgg tccgacaaag gatattccag ctcccgatgt gtacaccaat	480
tgcagatta tggcgtggat gatggatgag tacagccggc tgcgggaatt cgattctccg	540
ggctttatta caggtaaacc gcttgttttg ggaggatcgc aaggacggga aacagcgacg	600
gcacagggcg tcacgatttg tattgaagag gcggtgaaga aaaaagggat caagctgcaa	660
aacgcgcgca tcatcataca gggctttgga aacgcgggta gcttcctggc caaattcatg	720
cacgatgcgg gcgcgaaggat gatcgggatt tctgatgcca atggcgggct ctacaacca	780
gacggccttg atatccctta tttgctcgat aaacgggaca gctttggtat ggtcaccaat	840
ttatttactg acgtcatcac aaatgaggag ctgcttgaaa aggattgcga tatttttagtg	900
cctgccgcga tctccaatca aatcacagcc aaaaacgcac ataacattca ggcgtcaatc	960
gtcgttgaag cggcgaacgg cccgacaacc attgatgcca ctaagatcct gaatgaaaga	1020
ggcgtgctgc ttgtgccgga taccctagcg agtgccggcg gcgtcacggg ttcttatttt	1080
gaatgggtgc aaaacaacca aggatattat tggtcgggaag aagaggttgc agaaaaactg	1140
agaagcgtca tggtcagctc gttcgaaaca atttatcaaa cagcggcaac acataaagtg	1200
gatatgcgtt tggcggctta catgacgggc atcagaaaat cggcagaagc atcgcgtttc	1260
cgcgcatggg tctaa	1275

--- --

<210> 12

<211> 1419

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid Sequence encoding enzyme 12

<400> 12

```

atgtccatca aagacgctgt aaaactgatt gaagaaagcg aagcccgtt tgcgatttg      60
cgctttaccg ataccaaagg caagcagcac cactttaccg tgccctgcgcg catcgtgttg      120
gaagaccccg aagagtgggt cgaaaacgga caggcgtttg acggttcgtc catcggcggc      180
tggaaaggca ttcaggttc cgatatgcag ctcgcgcccg atcccgccac ggcgtttatc      240
gatacctttt atgatgatgt taccgtcgtc attacctgcg acgttatcga tcccgccgac      300
ggtcaggggt acgaccgcga cccgcgctcc atcgcacgcc gcgcgaagc ctatttgaaa      360
tcttcgggta tcggcgacac ggcatacttc ggtccgaac ccgagttttt cgtcttcgac      420
ggcgtagaat ttgaaaccga tatgcacaaa acccgttacg aaatcacgtc cgaaagcggc      480
gcatggggca gcggcctgca tatggacggt caaaacaccg gccaccgcc tgccgtcaaa      540

ggcggttacg cccccgtcgc gccgattgac tgcggtcagg atttgcgttc cgcgatggta      600
aacatttttg aaggactcgg catcgaagtc gaagtgcacc acagcgaagt cggtagccggc      660
agccaaatgg aaatcggcac gcgcttcgcc accttggtca aacgcgccga ccaaacccaa      720
gacatgaaat atgtgattca aaatgtcgcc cacaacttog gcaaaaccgc caccttcatg      780
cccaaaccga ttatgggcga caacggcagc ggtatgcacg ttcaccaatc catctggaaa      840
gacgggtcaa acctgttcgc aggcgacggc tatgccgggt tgagcgacac cgcgctctac      900
tacatcggcg gcatcatcaa acacgccaaa gccctgaacg cgattacca tccgtccacc      960
aactcctaca aacgccttgt gccgcacttt gaagcgccga ccaaactggc atattccgcc     1020
aaaaaccgtt ccgcttccat ccgtattccg tctgtgaaca gcagcaaggc gcgcgcgac     1080
gaagcgcgtt tccccgaccc gaccgccaac ccgtacttgg cgttcgctgc cctgctgat     1140
gcggggtttg acggcattca aaacaaaatc catccgggcg atcctgccga taaaaatct     1200
tacgacctgc cgcggaaga agacgcgctc gtcccgaccg tttgcgcttc tttagaagaa     1260
gccctcgccg cgtcaaagc cgaccacgaa ttctctttac gcggcgggcg gttcagcaaa     1320
gactggatcg acagctacat cgcctttaa gaggaagatg tccgccgcat ccgtatggcg     1380
cgcgatccgc tggaatttga aatgtattac agcctgtaa      1419

```

<210> 13

<211> 1075

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid Sequence encoding enzyme 13

<400> 13

```

aggaggtgtt ttaataatga aagggtttgc aatgctcagt atcggtaaag tcggttggat      60
tgaaaaagaa aagcctactc ccggcccttt tgacgctatt gtaagacctc tagctgtggc      120
cccttgcact tcggacgttc ataccgtttt tgaagggtgt attggcgaaa gacataacat      180
gatactcggc cacgaagctg taggtgaagt agttgaagta ggtagtgagg taaaagattt      240
taaacctggc gatcgcggtg tggtagcagc tattaccctt gattggcgaa cctctgaagt      300
gcaaagagga tatcaccaac actctgggtg aatgctggca ggctggaaat tttcgaatat      360

```

```

aaaagatggt gtttttgggtg aatTTTTTca tgtgaacgat gctgatatga atttagcaca 420
tctgcctaag gaaattccat tggaagctgc agttatgatt cccgatatga tgactactgg 480
ctttcacgga gccgaactgg cagatataga attaggtgcg acggtagcgg ttttgggtat 540
tggtccagta ggtcttatgg cagtcgctgg tgccaaattg cggggtgctg gaaggattat 600
cgcagtaggc agtagaccag tttgtgtaga tgctgcaaaa tactatggag ctactgatat 660

tgtaaaactat aaagatggtc ctatcgacag tcagattatg gatttaacgg aaggcaaagg 720
tgttgatgct gccatcatcg ctggaggaaa tgttgacatc atggctacag cagttaagat 780
tgttaaacct ggtggcacca tcgctaattg aaattacttt ggccaaggag atgttttgcc 840
tgttcctcgt cttgaatggg gttgcggcat ggctcataaa actataaaag gcgggctatg 900
ccccggtgga cgtctaagaa tggaaagact gattgacctt gttgtttata agcgtgtcga 960
tccttctaag ctcgtcactc acgttttccg gggatttgac aatattgaaa aagcctttat 1020
gttgatgaaa gacaaacca aagacctaat caaacctgtt gtaatattag cataa 1075

```

<210> 14

<211> 1545

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 14

<400> 14

```

atggctgact cgcaaccctt gtccgggtacc ccggaagggtg ccgaatattt aagagcgggtg 60
ctgcgcgcgc cggctctacga agcggcgcag gtcacgcgcg tacagaaaat ggaaaaactg 120
tcgtcgcgtc tcgataacgt gattctgggtg aagcgcgaag atcgccagcc agttcatagc 180
tttaagttgc gcggcgcata cgccatgatg gcgggcctga ccgaagaaca aaaagcacac 240
ggcgtgatta ccgcttctgc aggttaaccac gcgcaggggc tcgcgttttc ttccgcacgg 300
ttaggcgtga aggcgctgat cgtcatgcc aaccgccacc ccgatatcaa agttgatgcg 360
gtgcgcggct ttggcggcga agtgctgctt cacggcgcaa atttcgatga agcgaagcgc 420
aaagcgatcg aactgtcaca gcagcagggt ttacactggg taccgccgtt cgatcatccg 480
atggtgatcg ccgggcaagg cacgctggcg ctggaactgc tcagcagga cggccatctc 540
gaccgcgtat ttgtaccggt cggcggcggc ggtctggcag cgggtgtggc ggtgctgac 600
aaacaactga tgccgcaa atcaaagtaat gccgtggaag ccgaagattc cgcctgcctg 660
aaagcggcgc tggatgcggg tcatcccggt gatctgccc gcgtggggct gtttgcgtgaa 720
ggcgtcgcgg taaaacgcat cggcgatgaa accttccgtt tgtgccagga gtatcttgac 780
gacatcatca ccgtcgatag cgatgccatc tgtcggcgga tgaaagatct gttcgaagat 840
gtgcgcgcgg tggcggaacc ttccggcgcg ctggcgctgg cggggatgaa aaaatacatc 900
gccagcaca acattcgcgg tgaacggctg gcgcatatct tttccggtgc taacgtgaac 960
tttcacggtc tgcgtacgt ctcggaacgc tgcgaactgg gcgaacagcg tgaagcgttg 1020
ttggcgggtg ccattccgga agaaaaaggc agcttcctca aattctgcca actgcttggc 1080
gggcgttcgg tcaccgagtt caactaccgt tttgccgatg ccaaaaacgc ctgcatcttt 1140
gtcggcgtgc gcttaagccg tggcctcgaa gagcgcaaag aaattttgca gatgctcaac 1200

gacggtggct acagcgtggt tgatctctcc gacgacgaaa tggcgaagct gcatgtgcgc 1260
tatatgggtg gcgggcgtcc atcgcatccg ttgcaggaac gcctatacag cttcgaattc 1320
ccggaatcac cgggcgcgct gctgcgcttc ctcaacacgc tgggtacgca ctggaacatc 1380
tcgctgttcc attatcgag ccacgggtacc gactacgggc gcgtactggc ggcgttcgag 1440
cttggcgatc atgaaccgga ttttgaaacc cggttgaatg aactgggcta cgattgccac 1500

```

gacgaaacca ataaccgcgc gtccaggttc tttttggcgg gtttag 1545

<210> 15

<211> 750

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic acid sequence encoding enzyme 15

<400> 15

atgagcggta ccatacctcat caccggcgcc acgtccggct tcggacaggc caccggcggg	60
cgtttcgtca aggaaggctg gaaggctatc ggacacaggc ggccggcgga acggctggag	120
gcgctggcgc aagaactcgg ctccgccttt caccggcgtg ccttcgatgt taccgacgaa	180
gatgccacta gaaaggcact tgcggccttg ccggaagggt tccgggacat cgatattctc	240
gtcaacaatg cggggcttgc gtcggcacc gcacctgcac cgcagggtgc gctgaaagac	300
tggcagacca tggatgaacac caacatcacc ggtcttttga acatcaccca ccatactttg	360
cccacgttga tcgaccgcaa gggcattgtc atcaaccttt cctcggtagc tgcgcactgg	420
ccctatgcgg gcggcaatgt ctatgccgga acgaaagcct tcctgcggca attctcgctc	480
ggtctgcgct ccgacctgca tggcaagggc gtgcgcgtca cctcgatcga accgggcatg	540
tgcgaaacgg aattcacgct tgttcgcacc ggccgcaatc aggatgcctc ggacaatctt	600
tacaagggcg tcaatccgat caccggcgag gatatcgcca atacgatcca ttgggtcgcc	660
tgcagccca aacatatcaa catcaacagc ctggaactca tgcgggtcaa ccagtccttt	720
gccggtttcc aagtgcacg ggaaagttga	750

<210> 16

<211> 1380

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 16

<400> 16

atgatgacca aaaacgaaat ccaaaagtgg gtaaaggaat tcccgtgct tgaaacgac	60
atggcggccg aagaggtatt ttggcgcaat ccaaaatata acgcgtttgc gcaagctatt	120
cgaacgattc ctttacgcga acgcgatgtc aaggaggccg aagagcgatt gcgcgcgttt	180
gccccctaca tcgcgaaagt gtttcccgag acgcgaacgg cccacggtat catcgaatcc	240
ccttttagtgc ggattccgaa catgaaacag cgtttggaaa agatgtttca gaccaacatc	300
gaggggggac tgttgctaaa atgcgacagc catcttccca tctccggatc gatcaaggcg	360
agagggggaa tctacgaggt tctgaaacat gcggaagaac tcgctctggc aaaccatatg	420
atcaccatgg gggatgacta tgcggtcacg gccagcgaag aattccggca gttcttttcc	480
cgctattcgc ttgtcgttgg ttgcacggga aatttaggct tgagtatcgg catcatcggg	540
gcgcagcttg ggttccgcgt taccgttcat atgtcagccg atgcgaaaca atggaaaaaa	600
gacttgttgc gaagcaaagg ggttcgggtc atcgaacatc tcaccgacta caacaagggtg	660
gtggaagagg cgcgaagaca gtccgccgag gatccaaagt cgtattttat cgatgatgag	720
aactcgatcc atctgttttt aggcctatgcg gtggcgcggt ttcggctgaa aaagcaatta	780
gaggacatga acatcacggt tgatgaaaac caccgcctct ttgtatatct tccttgcggc	840
gtcggcgggc gtccggggcg ggtgacgttt gggctgaagc tcgtgtacgg cgatcatgtc	900
cattgctttt tcgctgagcc gaocgattcg ccttgcatgt tgctcggcct gatgacggga	960

cagcacgacc gcgtgtcggg gcaagatfff ggccctcgaca ataagaccga agcggacggg	1020
ctagcgggtgg ggcggccgct aagggttggtg gggaacatgc ttgagaacgt catcagcggc	1080
gtctatacgg tggacgatgc gacgctttac cgcttgctcg cggcgatggt ggaaacggag	1140
gaaatctatt tagagccgct cgccttggcg ggggtggcgg ggccgttctg gctgtttcgt	1200
gatttggcgg ggcaaacgta cgtagaggca aacggtttga aagaaaagat gaaaaacgcc	1260
gtccatattg gctgggcgac aggcggaagc atggtgctaa aggatgtgat ggaggcctat	1320
tatcgggaag gcgtgcgcat cgaaacgatg acagggaaacg gtttttctga aggacgataa	1380

<210> 17

<211> 1047

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 17

<400> 17

atgctgatgt tcaagaaat ccaggcgcgc ggccacgaga gcgtcacgct gctgcaccac	60
gccccagcg gcctgcgcgc cgtgctcgcc gtgcactcca ccgtgctcgg ccctgccatt	120
gcccggctgcc gcctgatgcc ctgcaccgag gaacgcgcgc tgccgcgacgc cctcgccctc	180
agcgagtccg tcacgctcaa ggccgcccctc gcgggcctga actacggcgg ggccgcgtgc	240
gtcatgctcc ccccggaagg cggcgacatc gacgggcacg ccccgagagg gctgttcgcg	300
gcgctcggcc ggcagatccg ttaccgcggt ggccgcgtca tctcaccga ggacgtcggc	360
gtgaccggcc gcgacatcgc cttcgccgcg caggaaaccg acagcaccat gggcatgcac	420
accgacacgc ccaccgtcac cgcgtacggc gtgtaccgcg gcatcaaggc cgcgcgcgc	480
gcctacctcg gcggcgagag catgcgcggc gtgcgcgtcg cgctgctcgg cgcgggcgca	540
gtcgggcgca ccctcgcgca gcacctgcac cgcgagggcg cgcgcctcac cgtcgcagac	600
ctgatgtctg agcgcgcgca ggccctcgcg gacgacctcg gcgagcgcgt caccgtcgtg	660
agcgcgcgtg acatcttcga cgtgccgtgc gacgtattcg cgcgtgctgc gttcgggcac	720
agcatcaaaa gcgcgcgacgt gcccgcgttg cagtgcgcgg tgatcgccgg cagcgaacac	780
cacccgctca gccacaacgg cgagacgctc gtgcgcgaag cgggcatcac atacatcccg	840
gacttcgcca tcaacagcgc cggcctgatg agcgcgcgc agaacctcag catcgaaacg	900
gcggcggaac gcgtgtacga gagcgtcgcg cagatctgcg cgaccgcgca gaagtacgag	960
aagccgcgcg acgtcgtcgc ccgtaaactc gcgctgcgcc gcacggaact gatcggctcc	1020
atcagcggcc agtacgcgg ccagtaa	1047

<210> 18

<211> 768

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 18

<400> 18

tcatgtgcca acacgtatgt tatcacttaa aatfttttagt aaagtgactg ctgaatatgc	60
tgccaaaaca cttgtttttg gatttaattc acacacagtg ttttttggtt tagattttaa	120
ctctccaaaa tctcctttta catggacttc atggatattg tgttcaactt caggatctgc	180
aattatcttt acatccgcat ctattccaga ggctagactt aatgccgcag caacgttaat	240
attcactgga aatfttttta cagcttctga ggatttccct ttaaacacga cctccttttt	300

tttgggtctta acacctaacg aagtaggtga ttttctcggt ataagtttta tttcttttat	360
ottacctaag gatgcgggtt ttacaccatc taaaccaatt attgcaccgg aaggtatgta	420
tatattagct cctgattctc tagattcctt tatcaatctt cttctaactt tctcatctaa	480
tagtgcaccc acactcataa tcaaaacatc tatacctcta ctaattatat tgggcacaa	540
ttcttttact gcctcttgag aagcagattc aattatcaaa tcaactccat tgaacatttc	600
ttctaccttt tttacggcag tgccatttgt taaatttggt agcttcttag cttttctaaa	660
atctctgtca taaaaatatt ttaattttat ttttttgata tcttggttta agacaagggt	720
aactattgta tttgcaattg caccacatcc tataatccca catctcat	768

<210> 19

<211> 1425

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 19

<400> 19

atgtcctcgc ctgcatcatc ggcacatcga aaagacctgc ttgggtgttct cgaagtacct	60
gccaacgcgt attacggcat ccagaccctg cgagcgggtga acaactttca cctctccggc	120
gtgccgcttt cgcactaccc gaaactggta gtcgcgctgg ccattggtcaa gcaggcggca	180
gcggatgcaa accatcagct cggacacctc aatgacgcca agcatgcggc gatcagcgag	240
gcctgtgccc gcctgatccg cggcgacttc cacgatcagt tcgtggtcga catgatccag	300
ggcggcgctg gcacgtcgac caacatgaat gccaacgaag tcatcgccaa catcgctctg	360
gaaaccatgg gtttcgagaa aggcgcatac aaacacctgc accccaacaa cgatgtcaac	420
atggcgcagt cgaccaacga cgcctacccc acggcgatcc gcttgggtct gctgctgggt	480
cacgacgctc tgctcgccag cctttccagc ctgattcagg ccttcgccgc caagggcgaa	540
gaattcaacc atgtgctgaa gatggggcgc acccagttgc aggacgccgt tccaatgacc	600
ctgggtcagg aattccgcgc cttcgccacc accctgacag aagacctgaa ccgcctgcgc	660
agcctggcgc cagagctgtt gaccgaagtg aacctcggcg gaaccgccat cggcacccgc	720
atcaacgccg accctggcta tcagaagctg gcagtcgac gtctggcact catcagcggc	780
cagcctctgg tgccagcagc cgacctgac gaagcgacct ccgacatggg cgccttcgtg	840
ttgttctcgg gcatgctcaa gcgactgcg gtcaagctgt cgaaaatctg caacgacctg	900
cgcctgctgt ccagcggccc acgcaccggc atcaacgaaa tcaacctgcc ggcacgtcag	960
ccaggcagct cgatcatgcc cggcaaggtc aacccggtga tcccggagc ggtcaatcag	1020
gttgcccttcg aaatcatcgg caacgacctg tcgctgacca tggcagccga aggaggacaa	1080
ttgcagctca acgtgatgga gccgctgac gcctacaaga tcttcgactc gatccgcctg	1140
ctgcagcgcg ccatggacat gctgcgcgag cactgcatcg tcggcatcac agccaacgaa	1200
cagcgtgcc gcgagctggt cgagcattcg atcggctctg tcaccgccct gaaccttac	1260
atcggttacg agaactccac ccgtatcgcc cgcacgcgc tggaaaccgg ccgcggcgtg	1320
ctggaactgg tgcgtgagga aggtctgctc gacgacgcca tgctcgacga catcctgcgc	1380
ccggaaaaca tgatcgctcc gcgtctggcc cccttgaagg cctga	1425

<210> 20

<211> 1083

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 20

<400> 20

tcagcgaccg cgggcctcgg ccatccgctg ctccggcgatc cggtcgggccg ccgcggcggg	60
cggaatgccg tccgccttcg cactgtcgaa tatttccagc gtggtgtcga agatcttcgt	120
cgccttcgcc ttgcaccggt cgaagtcgaa cccgtgcagc tcgtcggcga cctggatcac	180
gccgccggcg ttgaccacat agtcgggtgc gtagaggacc gaccggtcgg ccaggtcctt	240
ctcgacaccc gggtagggcca gctgggtgtt ggccgcgccg cacaccacct tcgccgtgag	300
caccggaacg gtgcgcgtcg tgagcgcgcc gccgagcgcg cagggcgcgt agatgtcgag	360
accctcggtg cggatcagcg tctcgggtgc cgcaccacg gtgacctcgg ggtgcagatc	420
ggtgatccgg cgcaccgact cctcgcgtac gtccgtgata acgacctcgg ccccgtagga	480
gagcaggtgc tcgacgaggt ggtggccac cttgccgacc ccggcgacgc cgaccttgcg	540
gccgcgcagc gtccgggtgc cccacaggtg ctgggccgag gcccgcatgc cctggaagac	600
accgaacgcg gtgaggacgg aggagtcgcc gccgcgccg ttctcggggg agcggccggg	660
ggtccagcgg cactccctgg cgacgacgtc catgtcggcg acgtagggtgc cgacatcgca	720
ggcggtagcg taccggccgc cgagcgaggc gacgaaccgg ccgtaggcca ggaggagtgc	780
ctccgtcttg atcttctccg ggtcgcgat gatgacggcc ttgccgccac cgtggtagag	840
tccggccagg gcgttcttgt acgacatccc gcgcgacagg ttcagcgcgt cggcgacggc	900
ctcggcctcg gtgcgtacg ggtagaagcg ggtgccgccg aggcggggg ccagggcggg	960
ggagtggagg gcgatgacgg ccttgaggcc ggtggcacgg tcctggcaga tcacgacttg	1020
ctcgtgacct cctgatccg agtggaacag ggtgtgcagg acgccgttag tcacatcggg	1080
cac	1083

<210> 21

<211> 2910

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 21

<400> 21

ctagttgtaa aagtcgaggg aggcgcaact gcacatgagg tgacgatctc cgtaaacccc	60
gtcaatgcga cccacagtcg gccagtactt ttcaacgtac gagtaaggat aagggaatgc	120
cgcctaacgc cgatcatatg gtttgtccca tttatcatcg gtgacacatc ttgccgtgtg	180
tggtgcattc ttcaaaacat tgttatccac tgggtgttca cctttttcaa tggcggcaat	240
ctcacctcga atggaaatta gtgcatctgc caagcgatcc aactcccgct tgggtttctga	300
ttcgggtggg tcaatcatta aagtcccggg tacaggaaac gccagtgttg gcgagtgaat	360
tccgtagtcc atcaaccgtt tggccacgtc ctccgcctca atatgagctg tcttcttgaa	420
ccgtcgaaga tcaacgataa actcatgagc gcagtagttt tctccacca gaaaagaat	480
cgtataatgg ttctctagcg gcttcttcaa gtagtttgca ttcaaaacgg cgtactctgt	540
acaagttttg agcccggtg atccaagcat taacatcaac atgtacgata tcggaagaat	600
tgatgctgat ccgtacgctg attgtgagac ttggccgaat ggctgtgaac cgccaacttt	660
ttggttgaaa acagaatttg gcaaaaaggg ggccagatgt tgacggacag ctatagggcc	720
cattccgggg ccgccaccac catggggaat tgaaaacgtc ttgtggagat taatgtggca	780
cacgtcgcca ccgatatac cagggcctgt atagccaacc atggcgtaa gatttgcccc	840
atcaatgtag cattgtccac cgtagtagtg cgccattgat gtaatggata aaatatcctt	900
gtcaacaag ccatacgtac ttggatatgt tatcatgata cagcacaact ctttgcgtg	960
tttttgcaa gatttctcca ggtcattgat atcaaccctg ccgttagaca agcatttcac	1020

caagacaata ttcattcctg ccaatgttgc cgaagctgga ttcgtaccat gcgcactctc 1080
 tggaaatcaaa cagacgttgc ggtgtccttc cttcattgat agatggtacg cacgaataac 1140
 acgaagccca gcgtattcac cttgggcgcc actattaggc tgaagcgata ccgcatccag 1200
 accggttaatt tcccttaact tttgctcaag atctagacac aacgcactgt accctcgac 1260
 ttggtccact ggggcaaggg gatgcacatt ggtgaattct ggccaagaga gtggtaacat 1320
 agcagcggca gggttaagct tcatggtgca agatcccaac gggacgcaac catgcgtaag 1380
 gccgtaatcc tttcgttgta gacgatgaat atagcgcata agttcacttt cactcttgta 1440
 cttttgaaac gttgagtgtt tcaggaaatc agacttccgc accagatcca acggtagtac 1500
 cgatttctga tcggctattt tggaaagggc tgcgacgacg ggaagcttca accctgcagc 1560
 ctccaaaagt gacacaatgt gtccatccgt tgttgctca tccacagaaa tggagacagt 1620
 ccattactg taatcaacaa aaacattaat acccttctca acacatcgtg tcttgtaatc 1680
 ctccgctgta atgcctttta ggttcacagt aacagtgtcg aaaaatgcac tgtttaccac 1740
 agagtgtcct actgattcca taccacagc gagcacttcc gccttgccgt gtatctcatt 1800
 ggcaatctca tttagaccat ctggaccatg gtaggcggca taaaaccac tcacgttggc 1860
 caataacgct tgtgcagtac agatatttga tgtggcgcgc tcacgcttaa tatgtgttc 1920
 acgtgtctgc agcgccatgc gtatggatgg ctctccggca gaatccttac tgacgccgat 1980
 cacacgtccc ggcacaaacc tcttaaaactg ctcttgaca gcaaagaacg cggcgtgagg 2040
 acctccatat cctagtggaa caccaaaacg ctgggaggat cccacaacca catctgcatt 2100
 catttcacca ggtggcttga caagaacaca agccatcaag tcgggtccac agcaactaat 2160
 gacaccgtgc ttctttgcat tctcgaacag tgggtgagaag tcatgaagca tgcccatcgc 2220
 atctggtgtt tgtacaagga taccaacaa ggaactgtca gtccagtcaa tcagattcgt 2280

 gtcgcccacg acgacgttta tcttgagcgg ttcggctctt gtcttaacca tctcaatgca 2340
 ggatggaaaa acagtttttg atacgaagaa cgtattccgc tttcgttgac catgctgaaa 2400
 agcaagatgc atcgctcgg atgatgctgt cgcttggtca agaagagatg catttgccac 2460
 atccatcttt gtcaaatcca taaccatggt ttggaaatc aaaagggact ccagacgtcc 2520
 ttgtgcaatc tcagcttgg atggtgtgta ggggtgtgtac catccaggat tttcaatgac 2580
 gttgcgaagt atgacaggag gagtaatgga ctcgtagtac ccctgaccaa tcatgctttt 2640
 tagtaccttg tttcgcgcac caagagagcg cagcagtgcg agagcatcca tctactcat 2700
 agccgccacc tccgtcaagg gtgggcgtac aatatccctt ggaatagcag ccgtcatcaa 2760
 atcagagaga ctctcttttc caaccgttcg aagcatcgac attgtctcag ccgttggtgg 2820
 accaatatgg cggttaatat agctgtccgt ggcagtccat cgaacaaatg tcacgcatgg 2880
 caaagagcca cgaaacaaac gacggtacat 2910

<210> 22

<211> 1107

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 22

<400> 22

atgatcattg gcctgccgaa agagatcaaa gttaaggaaa accgcgtggc actcacgcc 60
 gggggcgtcg ccagcctcgt gcgcgcggc cacaccgtca tcgtggaacg cagcgcggc 120
 gtgggcagcg gcatccagga caccgagtag gagcaggcgg gcgcgcagct cggcagcgcc 180
 gccgaggcgt gggccgcgca gatggtcgtg aaggtaagg agcccatcaa gagcgaatac 240
 gggtaacctc gcccggaact gctgctgttc acgtacctgc acctcgctgc ggaccagccc 300
 ctacggacg ccctgctgag cgcggcacg accgcggttgc cgtacgagac ggtgcagctc 360

```

gacgaccgca gcctgccgct gctcacgccc atgagtgagg tcgcggggccg cctgagcgtg      420
caggccggcg cgtaccacct gcaaaagccc atcggcgggc gggcgctgct gctcggcggc      480
gtgccgggcg tgcaggcggg ccacgtcgtc gtgattggcg gggcgctcgt cggcacgaac      540
gccgcgaaaa tggccatggg cctcggcgcg aaggtcacgg tgctggacgt gaaccacggg      600
cgctctcgt acctcgacga cgtgttcttc ggaagctca ccaccatgat gagcaacgag      660
gcgaacatcc gctccatcct gcccgaggcg gacctcgtga tcggcgggcg gctgatcccc      720
ggggcgagg cgccgcacct tgtcacgcgc gacatgctgg cgaccatgca ggaaggcagc      780
gtcatcgtcg acgtggcggt ggaccagggc ggatgcgtgg agaccattca cgcgacgacg      840
cacgacgatc ccacgtacat cgtggacggc gtgatccact acggcggtgg gaacatgccg      900
ggcgcggtgc cgcgcaccag cacgttcgcg ctcacgaacc agaccattgg gtacgtgctg      960
cagctcgcgg acaaggcggt ggaggcactc agcgcagca agccgctgct gcgtggcctg     1020
aacaccatcg gcgggaagct gacgtacgcg ggcgtcgcgg aagcgttcgg cctgacgtac     1080
accgcgcctg aagtggcgct ggcgtaa                                     1107

```

<210> 23

<211> 948

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 23

<400> 23

```

atggagccca ctatgagcca attcgaacag ctgtaccgcc aggtggccct cagtgtcgcc      60
ggcaaccgcg tcgtggaaaa agtcttgagc aagcagggct gggcgctggc gcagcgtttt     120
gtatcgggcg agacggcgca ggacgccatc aaggccatca agcggctgga agccagggc     180
atctccggca acctcgacct gctgggagag ttctgtaaca ccccggaacc cgccaatgcc     240
aacaccgaga tgattctggc gaccattgac caggtgcacg cggcgggcct cacgccctac     300
aacagcgtga aaatgtcggc gctgggcca aaggcagacc cgcgggacgg ccaggacctc     360
ggctacgtca acaccgccc cgtcgtggag cgggccaagc gctacggcgg ctctgtcaat     420
ctggacatgg aagaccacac ccgcgtggac tcgactctgc agattttccg ccgcctggtc     480
aaggagttcg gccaccagca tgtgggaacg gtgttcaggg cctacctgca ccgctcggaa     540
gacgaccgcc gcagcctgga cgacctgcgc cccaacctcc gcatggtgaa gggcgcctac     600
ctggagcccg cctccgtcgc cctgcagagc aaaaccgaca ttgacgccgc ctaccgccgc     660
ctggtctacg agcacctcaa ggccggcaac tactgcaacg tggccacca cgaccaccac     720
atcatctacg acgtgatgca ctttgcgctg gccacggcga tccctaagga ccagttcgaa     780
ttccagctgc tgtacggcat ccgcgaggac ctgcagcgcg aattggccga ggccggctac     840
acggtgcgct cgtacattcc tttcggcaag gactggtaag gctactactc gcgccgcac     900
gccgagcgcc cgcagaacgt gatgttcgtg ctgcgcggcc tgctgtaa                       948

```

<210> 24

<211> 1101

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic Acid sequence encoding enzyme 24

<400> 24

```

atgaaaaaca ttgtggttat cggcgcgggc aatatcggtt cggcaatcgc ctggatgctg      60

```

```

gccgcacacag gcgattatcg catcacgggt gccgatcggt cagccgatca gctggccaat      120
gtgccggcgc atgaacgggt cgacatcgtc gacattaccg accgtcccgc tctggaagca      180
ctgctaaaag gcaaattcgc cgtgctctcc gccgccccca ccgaattcca cctgacggcg      240
ggtattgccg aagcggccgt tgccgtcggc acgcattatc tcgatctcac cgaagacgtg      300
gaatccaccc gcaaggtcaa ggcgtggcg gaaacggccg aaaccgcgct cattccgcaa      360
tgccggcctcg ccccgggctt catctccatc gtcgctgccg atctcgccgt caagttcgac      420
aagctggaca gcgtgcgcat gcgcgtcggc gctctgccgc aatatccgtc caatgcgctc      480
aactacaacc tcacctggag taccgacggg ctgatcaacg aatatatcga gccctgcgaa      540
ggattcgctg aaggccgcct caccgccgtt ccggcccttg aggagcgcga ggagttctcg      600
ctcgatggca tcacctacga ggcgttcaac acctcggggc gtctcggtac gctttgcgcg      660
acgttggaag gcaaggtgcg gaccatgaac taccgcaacta tccgttatcc cggccatgtg      720
gcgatcatga aggcgctttt gaacgacctc aacctgcgca accgcgcgca tgtgctgaag      780
gacctgttcg aaaacgccct gcccggcacc atgcaggatg tggtcacgtt cttcgtcacc      840
gtctgcggca cccgcaacgg ccgcttcctg caggaaacct atgccaacaa ggtctatgcc      900
ggcccggttt ccggccggat gatgagcgcc atccagatca ctaccgccgc cggcatctgc      960
acggttctcg acctgctcgc ggaaggcgcc ctgccgcaga agggcttcgt tcgacaggag     1020
gaagtggcgc tgccgaagtt cctcgaaaac cggtttggcc ggtattatgg ctgcgatgag     1080
ccgctggcgc gggttgggtg a                                     1101

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US4963814A [0105]
- US4999632A [0105]
- US4999582A [0105] [0150]
- US5243516A [0105]
- US5352351A [0105]
- US5366609A [0105]
- US5405511A [0105]
- US5438271A [0105]
- US5762770A [0110]
- US6645359B [0121]
- US41194099 [0123]
- US5670031A [0125] [0150]
- US5698063A [0150]
- US5128015A [0150]
- US2013065548W [0212]
- EP14841207 [0269]
- US1453756W [0269]

- [US61672149B](#) [0269]

Non-patent literature cited in the description

- **EGGENSTEIN C et al.** A disposable biosensor for urea determination in blood based on an ammonium-sensitive transducer *Biosensors & Bioelectronics*, 1999, vol. 14, 133-41 [0004]
- **TATUSOVAMADDEN** *FEMS Microbiol Lett.*, 1999, vol. 174, 247-250 [0071]
- **LEHNINGER** *Biochemistry* *Worth Publishers, Inc.* 1975 000071-77 [0074]
- **J. ZSCHOCKE. F. HOFFMANN** *Vademecum Metabolicum* (Milupa Metabolics, Friedrichsdorf 2011 0000 [0212]
- **B. C. LANPHERA. L. GROPMANK. A. CHAPMANU. LICHTER-KONECKIM. L. SUMMAR** *Urea Cycle Disorders Overview* *NCBI Bookshelf* 2003 0000 [0212]
- **M. L. SUMMARS. KOELKERD. FREEDENBERGC. LE MONSJ. HABERLEH.-S. LEEB. KIRMSE** *The incidence of urea cycle disorders* *Mol. Genet. Metab.*, 2013, vol. 110, 179-80 [0212]
- **R. H. SINGH** *Nutritional management of patients with urea cycle disorders.* *J. Inherit. Metab. Dis.*, 2007, vol. 30, 880-7 [0212]
- **M. MSALL** *Neurological Outcome in Children with Inborn Errors of Urea Synthesis.* *pdfN. Engl. J. Med.*, 1984, vol. 310, 1500-1505 [0212]
- **A. L. GROPMANM. L. BATSHAW** *Cognitive outcome in urea cycle disorders* *Mol. Genet. Metab.*, 2004, vol. 81, 158-62 [0212]
- **M. L. BATSHAWS. BRUSILOWL. WABERW. BLOMA. M. BRUBAKKB. K. BURTONH. M. CANND. KERRP. MAMUNESR. MATALON** *Treatment of Inborn Errors of Urea Synthesis* *N. Engl. J. Med.*, 1982, vol. 306, 1387-1392 [0212]
- **F. F. POORDAD** *Review article: the burden of hepatic encephalopathy* *Aliment. Pharmacol. Ther.*, 2007, vol. 25, 13-9 [0212]
- **R. F. BUTTERWORTHJ. F. GIGUEREJ. MICHAUDJ. LAVOIEG. P. LAYRARGUES** *Ammonia: key factor in the pathogenesis of hepatic encephalopathy* *Neurochem Pathol*, 1987, vol. 6, 1-12 [0212]
- **R. F. BUTTERWORTH** *Pathophysiology of hepatic encephalopathy: a new look at ammonia.* *Metab. Brain Dis.*, 2002, vol. 17, 221-7 [0212]
- **J. STAHL** *Studies of the Blood Ammonia in Liver Disease* *Ann. Intern. Med.*, 1963, vol. 58, [0212]
- **I. EIJGELSHOVENS. DEMIRDAST. A. SMITHJ. M. T. VAN LOONS. LATOURA. M. BOSCH** *The time consuming nature of phenylketonuria: A cross-sectional study investigating time burden and costs of phenylketonuria in the Netherlands* *Mol. Genet. Metab.*, 2013, vol. 109, 237-242 [0212]
- **P. V. D. BURGH. W. MOOKA** *A simple and rapid method for the determination of ammonia in blood* *Clin. Chim. Acta*, 1962, vol. 8, 162-164 [0212]
- **Y. MURAWAKIK. TANIMOTOC. HIRAYAMAY. IKUTAN. WATABEA** *A simple and rapid microdiffusion method for blood ammonia using a reflectance meter and a reagent plate, and its clinical evaluation for liver diseases* *Clin. Chim. Acta*, 1984, vol. 1, 144- [0212]
- **R. J. BARSOTTI** *Measurement of ammonia in blood* *J. Pediatr*, 2001, vol. 138, S11-S20 [0212]
- **J. BUTTERYR. RATNAIKEB. CHAMBERLAIN** *The measurement of erythro-cyte ammonia using the Hyland ammonia kit* *J Clin Chem Clin Biochem*, 1982, vol. 20, [0212]
- **S. DIENST** *An ion exchange method for plasma ammonia concentration* *J. Lab. Clin. Med.*, 1961, vol. 58, [0212]
- **J. HUIZENGAC. GIPS** *Determination of blood ammonia using the Ammonia Checker* *Ann Clin Biochem*, 1983, vol. 20, [0212]
- **H. VAN ANKENM. SCHIPHORST** *A kinetic determination of ammonia in plasma* *Clin Chim Acta*, 1974, vol. 56, [0212]
- **L. ROVER JUNIORJ. C. FERNANDESG. DE OLIVEIRA NETOL. T. KUBOTAE. KATEKAWAS. H. SERRANO** *Study of NADH stability using ultraviolet-visible spectrophotometric analysis and factorial design.* *Anal. Biochem.*, 1998, vol. 260, 50-5 [0212]

- **M. BERTHELOT, B**Repert. Chim. Appl., vol. 254, 1859- [\[0212\]](#)
- **E. D. RHINEG. K. SIMSR. L. MULVANEYE. J. PRATT**Improving the Berthelot Reaction for Determining Ammonium in Soil Extracts and WaterSoil Sci. Soc. Am. J., 1998, vol. 62, [\[0212\]](#)
- **T. T. NGOA. P. H. PHANC. F. YAMH. M. LENHOFF**Interference in Determination of Ammonia with the Hypoehlorite-Alkali Phenol Method of Berthelot, 1981, 46-49 [\[0212\]](#)

PATENTKRAV

1. En teststrimmel (701) omfattende:

- 5 en første (602) og en anden beholder (603);
 en fluidudvekslingsåbning placeret mellem den første (602) og den anden be-
 holder (603);
 en ledning (601) i fluidkommunikation med den første beholder (602), led-
 ningen konfigureret til at modtage en fluid fra et punkt uden for biosensoren
10 og
 en membran (604) omfattende perfluorerede ionomerer placeret ved fluidud-
 vekslingsåbningen;
 hvor membranen (604) er i stand til selektivt at transportere ammoniak; og
 hvor den anden beholder omfatter en hypohalit, en alkalibuffer og en phenol-
15 forbindelse, der er egnet til at udføre en Berthelot-reaktion.

2. Teststrimmel (701) ifølge krav 1, hvor teststrimlen (701) er tilpasset til en bærbar
indretning (700) omfattende et voltmeter og/eller amperemeter og et digitalt display
(774) således, at når teststrimlen (701) er i kontakt med indretningen, bliver den første
20 og anden elektrode operativt forbundet til et lukket elektrisk kredsløb omfattende
 voltmeteret og/eller amperemeteret og det digitale display (774) og ved kontakt med
 en prøve af kropsfluid (702), det mindst ene metaboliske enzym eller funktionelle
 fragment deraf katalyserer oxidation af en aminosyre, hvilket resulterer i en strøm på
 den første elektrode svarende til en koncentrationsværdi af aminosyre i prøven af
25 kropsfluid (702), en sådan koncentrationsværdi kan aflæses på displayet (774) af den
 bærbare indretning (700).

3. Teststrimmel (701) ifølge et hvilket som helst af kravene 1 eller 2, hvor alkalibuffe-
ren anvendes i et interval fra ca. 0,5 til ca. 1,0 mol/liter.

30

4. Teststrimmel (701) ifølge et hvilket som helst af kravene 1 til 3, hvor hypohaliten
anvendes i et interval fra ca. 50 til ca. 120 mmol/liter.

5. Teststrimmel (701) ifølge et hvilket som helst af kravene 1 til 4, hvor alkalibufferen er en eller en kombination af: calciumacetat, calciumchlorid, zinkacetat, zinkchlorid eller en hvilken som helst ækvivalent mono, di eller tri, -valent salt deraf.
- 5 6. Teststrimmel (701) ifølge et hvilket som helst af kravene 1 til 4, hvor phenolreagenset eller indophenolreagenset anvendes i et interval fra ca. 68 til ca. 70 mmol/liter.
7. Teststrimmel (701) ifølge et hvilket som helst af kravene 1 - 6, hvor den perfluorerede ionomer er Nafion.
- 10 8. Teststrimmel (701) ifølge et hvilket som helst af kravene 1-7, hvor den perfluorerede ionomer har en tykkelse fra ca. 0,001 mm til ca. 0,069 mm.
- 15 9. Teststrimmel (701) ifølge et hvilket som helst af kravene 1 - 8, hvor den første beholder (602) ikke har mere end ca. 100 mikroliter i volumen.
10. En fremgangsmåde til diagnosticering af en metabolisk sygdom hos et individ omfattende:
- 20 (a) at bringe en prøve af kropsfluid (702) i kontakt med teststrimlen (701) ifølge et hvilket som helst af kravene 1-9;
- 25 (b) kvantificering af en eller flere koncentrationsværdier af ammoniak i prøven;
- 30 (c) sammenligning af en eller flere koncentrationsværdier af ammoniak i prøven med en tærskelværdi for ammoniakkoncentration identificeret som værende i et sundt interval; og
- 30 (d) identificering af individet som havende en metabolisk sygdom, hvis en eller flere koncentrationsværdier af ammoniak i prøven overstiger eller falder under tærskelværdien.
11. Fremgangsmåde ifølge krav 10, hvor den metaboliske sygdom er hyperammoniami.

12. Fremgangsmåde til bestemmelse af patientmodtagelighed overfor en terapi omfattende:

- 5 (a) at bringe en prøve af kropsfluid (702) i kontakt med teststrimlen (701) ifølge et hvilket som helst af kravene 1-9;
- (b) kvantificering af en eller flere koncentrationer af ammoniak eller ammoniumion;
- (c) sammenligning af en eller flere koncentrationsværdier med en eller flere tærskelværdier forbundet med en metabolisk sygdom.

10

13. Fremgangsmåde ifølge krav 12, hvor trinnet med kontakt omfatter eksponering af prøven (702) for teststrimlen (701) i et tilstrækkeligt tidsrum til, at modifikation af indophenolreagenset eller indophenol-relateret forbindelse kan modificeres til en indophenol eller indophenol-reaktionsprodukt.

15

14. Fremgangsmåde ifølge et hvilket som helst af kravene 12 - 13, hvor fremgangsmåden ikke omfatter eksponering af prøven af kropsfluid (702) for nogen eksterne stimuli eller reagenser før prøven bringes i kontakt med den mindst ene elektrisk ledende understøtning.

20

15. Fremgangsmåde ifølge et hvilket som helst af kravene 12 - 14, hvor prøven af kropsfluid (702) er fuldblod fra et individ.

DRAWINGS




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28



Catalyzed by the enzyme Phenylalanine Ammonia Lyase




FIG. 29