



(86) **Date de dépôt PCT/PCT Filing Date:** 2015/01/21
(87) **Date publication PCT/PCT Publication Date:** 2015/07/30
(45) **Date de délivrance/Issue Date:** 2024/04/30
(85) **Entrée phase nationale/National Entry:** 2016/07/13
(86) **N° demande PCT/PCT Application No.:** EP 2015/051081
(87) **N° publication PCT/PCT Publication No.:** 2015/110449
(30) **Priorité/Priority:** 2014/01/21 (EP14151960.3)

(51) **Cl.Int./Int.Cl. C12N 15/85** (2006.01),
A61K 45/00 (2006.01), **C12N 15/86** (2006.01)
(72) **Inventeurs/Inventors:**
VANDENDRIESSCHE, THIERRY, BE;
CHUAH, MARINEE, BE;
DE BLESER, PIETER, BE
(73) **Propriétaires/Owners:**
VRIJE UNIVERSITEIT BRUSSEL, BE;
VIB VZW, BE;
UNIVERSITEIT GENT, BE
(74) **Agent:** SMART & BIGGAR LP

(54) **Titre : ELEMENTS REGULATEURS D'ACIDE NUCLEIQUE EXPRIME DANS UN MUSCLE, METHODES ET UTILISATION ASSOCIEES**
(54) **Title: MUSCLE-SPECIFIC NUCLEIC ACID REGULATORY ELEMENTS AND METHODS AND USE THEREOF**

(57) **Abrégé/Abstract:**

The present invention relates to nucleic acid regulatory elements that are able to enhance muscle-specific expression of genes, in particular expression in cardiac muscle and/or skeletal muscle, methods employing these regulatory elements and uses of these elements. Expression cassettes and vectors containing these nucleic acid regulatory elements are also disclosed. The present invention is particularly useful for applications using gene therapy, more particularly muscle-directed gene therapy, and for vaccination purposes.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2015/110449 A1

(43) International Publication Date
30 July 2015 (30.07.2015)

(51) International Patent Classification:
C12N 15/864 (2006.01)

(21) International Application Number:
PCT/EP2015/051081

(22) International Filing Date:
21 January 2015 (21.01.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
14151960.3 21 January 2014 (21.01.2014) EP

(71) Applicants: **VRIJE UNIVERSITEIT BRUSSEL** [BE/BE]; Pleinlaan 2, B-1050 Brussel (BE). **VIB VZW** [BE/BE]; Rijnvischestraat 120, B-9052 Gent (BE). **UNIVERSITEIT GENT** [BE/BE]; Sint-Pietersnieuwstraat 25, B-9000 Gent (BE).

(72) Inventors: **VANDENDRIESSCHE, Thierry**; Tiensesteenweg 250, B-3360 Bierbeek (BE). **CHUAH, Marinee**; Tiensesteenweg 250, B-3360 Bierbeek (BE). **DE BLESER, Pieter**; Prosper Bosteelstraat 18, B-9255 Buggenhout (BE).

(74) Agents: **VANHALST, Koen Victor Rachel** et al.; De Clercq & Partners, E. Gevaertdreef 10a, B-9830 Sint-Martens-Latem (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: MUSCLE-SPECIFIC NUCLEIC ACID REGULATORY ELEMENTS AND METHODS AND USE THEREOF

(57) Abstract: The present invention relates to nucleic acid regulatory elements that are able to enhance muscle-specific expression of genes, in particular expression in cardiac muscle and/or skeletal muscle, methods employing these regulatory elements and uses of these elements. Expression cassettes and vectors containing these nucleic acid regulatory elements are also disclosed. The present invention is particularly useful for applications using gene therapy, more particularly muscle-directed gene therapy, and for vaccination purposes.



WO 2015/110449 A1

MUSCLE-SPECIFIC NUCLEIC ACID REGULATORY ELEMENTS AND METHODS AND USE THEREOF

FIELD

The present invention relates to nucleic acid regulatory elements that are able to enhance muscle-specific expression of genes, methods employing these regulatory elements and use thereof. The invention further encompasses expression cassettes, vectors and pharmaceutical compositions comprising these regulatory elements. The present invention is particularly useful for applications using gene therapy, more particularly muscle-directed gene therapy, and for vaccination purposes.

BACKGROUND

Muscle is an attractive target for gene therapy. Gene delivery to muscle can be used to augment expression of muscle structural proteins, such as dystrophin and sarcoglycans, e.g. to treat muscular dystrophy. In addition, muscle can be used as a therapeutic platform to express non-muscle secretory/regulatory pathway proteins for diabetes, atherosclerosis, hemophilia, cancer, etc.

Efforts to deliver transgenes to muscle have focused on vectors derived from adenoviruses, retroviruses, lentiviruses, and adeno-associated viruses (AAV), and plasmids. Adenoviral vectors have a relatively large cloning capacity can be produced at high titers and display relatively efficient transduction of muscle. Unfortunately, these vectors can elicit a robust cellular immune response against viral and some transgene proteins. Moreover, they can evoke a rapid activation of the innate immune system that can contribute to a dose-limiting and potentially dangerous inflammatory immune response. Adenoviral vectors do not integrate into the host genome, so their ability to persist for long periods of time is unclear. Retroviral and lentiviral vectors integrate stably into the target cell genome, potentially allowing persistent gene transfer. Whereas lentiviral vectors can transduce both dividing and non-dividing cells, conventional retroviral vectors derived from Moloney murine leukemia virus (MoMLV) can only transduce dividing cells. Consequently, lentiviral vectors can be used to transduce non-dividing skeletal muscle cells, whereas these are refractory to transduction by direct injection with retroviral vectors. Nevertheless, even lentiviral transduction of skeletal muscle is not very efficient. Naked plasmid DNA displays a remarkable ability to transfer genes to muscle. Plasmids display minimal immunogenicity and toxicity, and have an extremely large cloning capacity. The primary disadvantage of plasmids is their relatively poor transfection

efficiency under typical delivery protocols. Retention of plasmids is another important consideration.

Adeno-associated viral vector (AAV) is by far the most promising gene delivery vehicle for muscle-directed gene therapy. AAV's natural tropism to muscle cells, their long-term
5 persistent transgene expression, their multiple serotypes, as well as their minimal immune response have made AAV vectors well suited for muscle-directed gene therapy. AAV vector can be delivered into both skeletal muscle and cardiac muscle by means of local, regional, and systemic administrations.

There however remain concerns regarding the efficacy and safety of some gene delivery
10 approaches. The major limiting factors are: insufficient and/or transient transgene expression levels, and inappropriate expression of the transgene in unwanted cell types. In particular, it has been shown that inadvertent transgene expression in antigen-presenting cells (APCs), increases the risk of untoward immune responses against the gene-modified cells and/or the therapeutic transgene product that consequently curtails
15 long-term gene expression.

Conventional methods of vector design relied on haphazard trial-and-error approaches whereby transcriptional enhancers were combined with promoters to boost expression levels. Though this could sometimes be effective, it often resulted in non-productive combinations that resulted in either modest or no increased expression levels of the gene
20 of interest and/or loss of tissue specificity. Moreover, these conventional approaches did not take into account the importance of including evolutionary conserved regulatory motifs into the expression modules, which is particularly relevant for clinical translation.

A computational approach depending upon a modified distance difference matrix (DDM) – multidimensional scaling (MDS) strategy (De Bleser et al. 2007. Genome Biol 8, R83) has
25 proven to be useful for the *in silico* identification of clusters of evolutionary conserved transcription factor binding site (TFBS) motifs associated with robust tissue-specific expression in liver (WO 2009/130208) and heart (WO2011/051450).

There remains a need in the art for safe and efficient gene delivery to muscle.

SUMMARY

30 The present inventors have relied on a modified DDM-MDS strategy (De Bleser et al., 2007) combined with an enhanced screening strategy to identify evolutionarily conserved transcription factor binding site (TFBS) motifs associated with highly expressed muscle-specific genes defined herein as nucleic acid regulatory elements. As shown in the

experimental section, the inventors could identify nucleic acid regulatory elements that specifically enhanced gene expression in both heart and skeletal muscle, and skeletal muscle-specific nucleic acid regulatory elements. These regulatory elements were subsequently validated *in vivo* yielding efficient and tissue-specific gene expression. This approach hence, allows for the use of lower and thus safer vector doses, while maximizing therapeutic efficacy.

The invention therefore provides the following aspects:

Aspect 1: a nucleic acid regulatory element for enhancing muscle-specific gene expression comprising, consisting essentially of, or consisting of a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment comprises at least 20, preferably at least 25, more preferably at least 50, at least 100, at least 200 or at least 250, contiguous nucleotides from the sequence from which it is derived, and wherein said functional fragment comprises at least 1, preferably at least 5, more preferably at least 10 or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which it is derived.

Aspect 2: the nucleic acid regulatory element according to aspect 1 for enhancing cardiac and skeletal muscle-specific gene expression, comprising a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment comprise at least 20, preferably at least 25, more preferably at least 50, at least 100, at least 200 or at least 250, contiguous nucleotides from the sequence from which it is derived, and wherein said functional fragment comprises at least 1, preferably at least 5, more preferably at least 10 or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which it is derived.

Aspect 3: the nucleic acid regulatory element according to aspect 1 for enhancing skeletal muscle-specific gene expression comprising a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a sequence having at least 80%,

preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment comprise at least 20, preferably at least 25, more preferably at least 50, at least 100, at least 200 or at least 250, contiguous nucleotides from the sequence
 5 from which it is derived, and wherein said functional fragment comprises at least 1, preferably at least 5, more preferably at least 10 or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which it is derived.

Aspect 4: the nucleic acid regulatory element according to aspect 1, 2, or 3, comprising, consisting essentially of, or consisting of a sequence selected from the group consisting
 10 of: the nucleotide sequence from position 33 to 58 in SEQ ID NO:1; the nucleotide sequence from position 90 to 142 in SEQ ID NO:1; the nucleotide sequence from position 143 to 233 in SEQ ID NO:1; the nucleotide sequence from position 240 to 310 in SEQ ID NO:1; the nucleotide sequence from position 90 to 233 in SEQ ID NO:1; the nucleotide sequence from position 47 to 130 in SEQ ID NO:5; the nucleotide sequence from position
 15 252 to 293 in SEQ ID NO:5; the nucleotide sequence from position 330 to 450 in SEQ ID NO:5; the nucleotide sequence from position 10 to 180 in SEQ ID NO:10 (i.e. SEQ ID NO:37); the nucleotide sequence from position 190 to 240 in SEQ ID NO:10 (i.e. SEQ ID NO:38); the nucleotide sequence from position 241 to 300 in SEQ ID NO:10 (i.e. SEQ ID NO:39); the nucleotide sequence from position 241 to 360 in SEQ ID NO:10 (i.e. SEQ ID
 20 NO:41); the nucleotide sequence from position 380 to 420 in SEQ ID NO:10 (i.e. SEQ ID NO:40); or a sequence having at least 95% identity to any of said sequences.

Aspect 5: the nucleic acid regulatory element according to any one of aspects 1 to 4 comprising, consisting essentially of or consisting of a sequence selected from the group consisting of: the nucleotide sequence from position 33 to 310 in SEQ ID NO:1; the
 25 nucleotide sequence from position 47 to 450 in SEQ ID NO:5; the nucleotide sequence from position 10 to 420 in SEQ ID NO:10, or a sequence having at least 95% identity to any of said sequences.

Aspect 6: the nucleic acid regulatory element according to any one of aspects 1 to 5, comprising, consisting essentially of, or consisting of a sequence selected from the group
 30 consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

Aspect 7: the nucleic acid regulatory element according to any one of aspects 1, 2, or 4 to 6 for enhancing cardiac and skeletal muscle-specific gene expression comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

Aspect 8: the nucleic acid regulatory element according to any one of aspects 1, 3 to 6 for enhancing skeletal muscle-specific gene expression comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

Aspect 9: the nucleic acid regulatory element according to any one of aspects 1 to 8, comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, a functional fragment thereof comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

Aspect 10: the nucleic acid regulatory element according to any one of aspects 1, 2, 4 to 7, 9, comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, a functional fragment thereof comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, or SEQ ID NO:6, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

Aspect 11: the nucleic acid regulatory element according to any one of aspects 1, 3 to 6, 8, or 9, comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, a functional fragment thereof comprising SEQ

ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

- 5 Aspect 12: a nucleic acid regulatory element for enhancing muscle-specific gene expression comprising, consisting essentially of, or consisting of the complement of a sequence as defined in any one of aspects 1 to 11.

Aspect 13: a nucleic acid regulatory element for enhancing muscle-specific gene expression hybridizing under stringent conditions to the nucleic acid regulatory element
10 according to any one of aspects 1 to 12.

Aspect 14: the nucleic acid regulatory element according to any one of aspects 1 to 13, having a maximal length of 1000 nucleotides, preferably 800 nucleotides, more preferably 600 nucleotides, such as 550 nucleotides, 500 nucleotides, 450 nucleotides, 400 nucleotides, 350 nucleotides, or 300 nucleotides, still comprising the regulatory element
15 defined by any one of SEQ ID Nos: 1-13, or any one of the functional fragments thereof defined in any one of aspects 3 to 5.

Aspect 15: use of the nucleic acid regulatory element according to any one aspects 1 to 14 in a nucleic acid expression cassette, or a vector, more particularly for enhancing muscle-specific expression of said nucleic acid expression cassette or vector.

- 20 Aspect 16: a nucleic acid expression cassette comprising at least one, such as one, two, three, four, five or more, nucleic acid regulatory element according to any one of aspects 1 to 14, operably linked to a promoter.

Aspect 17: the nucleic acid expression cassette according to aspect 16, wherein the nucleic acid regulatory element is operably linked to a promoter and a transgene.

- 25 Aspect 18: the nucleic acid expression cassette according any one of aspects 16 or 17, wherein the promoter is a muscle-specific promoter, preferably the promoter from the desmin (DES) gene.

Aspect 19: the nucleic acid expression cassette according to any one of aspects 17 or 18, wherein the transgene encodes a therapeutic protein or an immunogenic protein.

- 30 Aspect 20: the nucleic acid expression cassette according to any one of aspects 17 to 19, wherein the transgene encodes a secretable protein or a structural protein, such as dystrophin or sarcoglycan.

Aspect 21: the nucleic acid expression cassette according to any one of aspects 16 to 20, further comprising an intron, preferably the Minute Virus of Mouse (MVM) intron.

Aspect 22: the nucleic acid expression cassette according to any one of aspects 16 to 21, further comprising a polyadenylation signal, preferably the Simian Virus 40 (SV40) polyadenylation signal.

Aspect 23: a vector comprising the nucleic acid regulatory element according to any one of aspects 1 to 14, or the nucleic acid expression cassette according to any one of aspects 16 to 22.

Aspect 24: the vector according to aspect 23, which is a viral vector, preferably an adeno-associated viral (AAV) vector, more preferably an AAV9 vector.

Aspect 25: the vector according to aspect 23, which is a non-viral vector, preferably a plasmid, a minicircle or a transposon-based vector, such as a PiggyBac-based vector or a Sleeping Beauty-based vector.

Aspect 26: a pharmaceutical composition comprising the nucleic acid expression cassette according to any one of aspects 16 to 22, or the vector according to any one of aspects 23 to 25, and a pharmaceutically acceptable carrier.

Aspect 27: the nucleic acid regulatory element according to any one of aspects 1 to 14, the nucleic acid expression cassette according to any one of aspects 16 to 22, the vector according to any one of aspects 23 to 25, or the pharmaceutical composition according to aspect 26 for use in medicine.

Aspect 28: the nucleic acid regulatory element according to any one of aspects 1 to 14, the nucleic acid expression cassette according to any one of aspects 16 to 22, the vector according to any one of aspects 23 to 25, or the pharmaceutical composition according to aspect 26 for use in gene therapy, preferably muscle-directed gene therapy.

Aspect 29: the nucleic acid regulatory element, the nucleic acid expression cassette, the vector, or the pharmaceutical composition according for use according to aspect 28, wherein the gene therapy is for muscular dystrophy (e.g. Duchenne muscular dystrophy (DMD) / Becker muscular dystrophy (BMD)), myotonic dystrophy, neuromuscular disease, motor neuron diseases (MND), such as e.g. Charcot-Marie-Tooth disease (CMT), spinal muscular atrophy (SMA), and amyotrophic lateral sclerosis (ALS), Emery-Dreifuss muscular dystrophy, facioscapulohumeral muscular dystrophy (FSHD), congenital muscular dystrophies, congenital myopathies, limb girdle muscular dystrophy, metabolic myopathies, muscle inflammatory diseases, myasthenia, mitochondrial myopathies,

anomalies of ionic channels, nuclear envelop diseases, cardiomyopathies, cardiac hypertrophy, heart failure, distal myopathies, cardiovascular diseases, hemophilia, including hemophilia A and B, and diabetes.

Aspect 30: the nucleic acid regulatory element according to any one of aspects 1 to 14, the nucleic acid expression cassette according to any one of aspects 16 to 22, the vector according to any one of aspects 23 to 25, or the pharmaceutical composition according to aspect 26 for use as a vaccine, preferably a prophylactic vaccine, or for use in vaccination therapy, preferably prophylactic vaccination.

Aspect 31: A method, preferably an in vitro or ex vivo method, for expressing a transgene product in muscle cells, preferably heart muscle cells and/or skeletal muscle cells, comprising:

- introducing the nucleic acid expression cassette according to any one of aspects 16 to 22, or the vector according to any one of aspects 23 to 25 into the muscle cells;
- expressing the transgene product in the muscle cells.

The present disclosure includes:

[1] A nucleic acid regulatory element for enhancing muscle-specific gene expression, comprising the sequence of SEQ ID NO:5, a sequence that is at least 95% identical to SEQ ID NO:5 over the entire length and that enhances muscle-specific gene expression, or consisting of a functional fragment of the sequence of SEQ ID NO:5, wherein said functional fragment comprises the sequence from positions 47 to 130 of SEQ ID NO:5; the sequence from positions 252 to 293 of SEQ ID NO:5; or the sequence from positions 330 to 450 of SEQ ID NO:5;

[2] The nucleic acid regulatory element according to [1], wherein the nucleic acid regulatory element enhances cardiac and skeletal muscle-specific gene expression;

[3] The nucleic acid regulatory element according to [1] or [2], wherein said functional fragment consists of the nucleotide sequence from positions 47 to 130 of SEQ ID NO:5; the sequence from positions 252 to 293 of SEQ ID NO:5; or the sequence from positions 330 to 450 of SEQ ID NO:5;

[4] The nucleic acid regulatory element according to any one of [1] to [3], having a maximal length of 600 nucleotides;

[5] The nucleic acid regulatory element according to any one of [1] to [3], having a maximal length of 550 nucleotides;

[6] The nucleic acid regulatory element according to [4] or [5], comprising the sequence of SEQ ID NO:5, or a sequence that is at least 95% identical to SEQ ID NO:5 over the entire length;

- [7] Use of the nucleic acid regulatory element according to any one of [1] to [6] in a nucleic acid expression cassette, or a vector;
- [8] A nucleic acid expression cassette comprising at least one nucleic acid regulatory element according to any one of [1] to [6], operably linked to a promoter and a transgene;
- [9] The nucleic acid expression cassette according to [8], wherein the promoter is a muscle-specific promoter;
- [10] The nucleic acid expression cassette according to [9], wherein the promoter is the promoter of the desmin (DES) gene;
- [11] The nucleic acid expression cassette according to any one of [8] to [10], wherein the transgene encodes a therapeutic protein, an immunogenic protein, or a structural protein;
- [12] The nucleic acid expression cassette according to any one of [8] to [10], wherein the transgene encodes dystrophin or a sarcoglycan;
- [13] A vector comprising the nucleic acid regulatory element according to any one of [1] to [6], or the nucleic acid expression cassette according to any one of [8] to [12];
- [14] The vector according to [13], which is a viral vector;
- [15] The vector according to [13], which is an adeno-associated viral vector;
- [16] A pharmaceutical composition comprising the nucleic acid expression cassette according to any one of [8] to [12], or the vector according to any one of [13] to [15], and a pharmaceutically acceptable carrier;
- [17] Use of the nucleic acid regulatory element according to any one of [1] to [6], the nucleic acid expression cassette according to any one of [8] to [12], the vector according to any one of [13] to [15], or the pharmaceutical composition according to [16], in gene therapy;
- [18] Use of the nucleic acid regulatory element according to any one of [1] to [6], the nucleic acid expression cassette according to any one of [8] to [12], the vector according to any one of [13] to [15], or the pharmaceutical composition according to [16], in vaccination therapy;
- [19] Use of the nucleic acid regulatory element according to any one of [1] to [6], the nucleic acid expression cassette according to any one of [8] to [12], the vector according to any one of [13] to [15], or the pharmaceutical composition according to [16], in prophylactic vaccination; and
- [20] An in vitro or ex vivo method for expressing a transgene product in muscle cells comprising: introducing the nucleic acid expression cassette according to any one of [8] to [12], or the vector according to any one of [13] to [15] into the muscle cells; and expressing the transgene product in the muscle cells.

BRIEF DESCRIPTION OF DRAWINGS

FIG 1: Flow diagram of the identification and validation of nucleic acid regulatory elements. A computational approach was used to identify the nucleic acid regulatory elements involving the following steps: (1) identification of tissue-specific genes that are highly expressed e.g. based on statistical analysis of micro-array expression data of normal human tissues; (2) extraction of the corresponding promoter sequences from publicly available databases; (3) identification of the regulatory modules and the transcription factor binding sites (TFBS) they contain (4) Next, the genomic context of the highly expressed genes was searched for evolutionary conserved clusters of TFBS (i.e. nucleic acid regulatory elements). The identified nucleic acid regulatory elements were *de novo* designed and validated *in vivo* by testing whether inclusion in a construct increases expression of a reporter gene.

FIG 2: Nucleotide sequences of the cardiac and skeletal muscle-specific regulatory elements CSk-SH1 (A, SEQ ID NO:1), CSk-SH2 (B, SEQ ID NO:2), CSk-SH3 (C, SEQ ID NO:3), CSk-SH4 (D, SEQ ID NO:4), CSk-SH5 (E, SEQ ID NO:5), and CSk-SH6 (F, SEQ ID NO:6).

FIG 3 shows a schematic representation of the AAV9sc-CSk-SH/Sk-SH-Des-Luc2 vectors disclosed herein. The expression cassette was packaged in a self-complementary (sc) adeno-associated virus, serotype 9 (AAV9). The cardiac and skeletal muscle-specific

desmin (Des) promoter regulates transcription of the luciferase (Luc2) transgene. The identified cardiac and skeletal muscle-specific (CSk-SH) or muscle-specific (Sk-SH) nucleic acid regulatory elements were cloned upstream of the Des promoter. The expression cassette further comprises the Minute Virus of Mouse (MVM) intron and a Simian virus 40 (SV40) polyadenylation signal (pA). The expression cassette is flanked by inverted terminal repeats (ITR) from adeno-associated virus, serotype 2 (AAV2).

FIG 4: Difference in luciferase expression **(A)** and Luc mRNA level **(B)** in heart and muscle tissue of mice that were intravenously injected with, from left to right, AAV9sc-CSk-SH1-Des-Luc2 (CSk-SH1), AAV9sc-CSk-SH2-Des-Luc2 (CSk-SH2), AAV9sc-CSk-SH3-Des-Luc2 (CSk-SH3), AAV9sc-CSk-SH4-Des-Luc2 (CSk-SH4), AAV9sc-CSk-SH5-Des-Luc2 (CSk-SH5), or AAV9sc-CSk-SH6-Des-Luc2 (CSk-SH6) vector compared to mice that were injected with the control vector AAV9sc-Des-Luc2 without nucleic acid regulatory element (control, no Csk-SH). Luciferase expression was measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissues at 9 weeks post-injection. Results were presented as mean $\pm$ standard error of the mean, *p<0.05; **p<0.001. Luc mRNA levels were measured by a quantitative RT-PCR method (qRT-PCR) at the end of the experiment from total RNA extracted from biopsies from the indicated tissues. Results were presented as mean $\pm$ standard error of the mean, *p<0.05; **p<0.001.

FIG 5: **(A)** Luciferase expression in selected tissues of mice that were intravenously injected with, from left to right, AAV9sc-Des-Luc2 (control, no CSk-SH), AAV9sc-CSk-SH1-Des-Luc2 (CSk-SH1), AAV9sc-CSk-SH2-Des-Luc2 (CSk-SH2), AAV9sc-CSk-SH3-Des-Luc2 (CSk-SH3), AAV9sc-CSk-SH4-Des-Luc2 (CSk-SH4), AAV9sc-CSk-SH5-Des-Luc2 (CSk-SH5), or AAV9sc-CSk-SH6-Des-Luc2 (CSk-SH6) vector. Luciferase expression was measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissue at 9 weeks post-injection. Results were presented as mean $\pm$ standard error of the mean. **(B)** Luc mRNA level in selected tissues of mice that were intravenously injected with, from left to right, AAV9sc-CSk-SH1-Des-Luc2 (CSk-SH1), AAV9sc-CSk-SH2-Des-Luc2 (CSk-SH2), AAV9sc-CSk-SH3-Des-Luc2 (CSk-SH3), AAV9sc-CSk-SH4-Des-Luc2 (CSk-SH4), AAV9sc-CSk-SH5-Des-Luc2 (CSk-SH5), or AAV9sc-CSk-SH6-Des-Luc2 (CSk-SH6) vector compared to mice that were injected with the control vector AAV9sc-Des-Luc2 (control, no Csk-SH). Luc mRNA levels were measured by a quantitative RT-PCR method (qRT-PCR) at the end of the experiment from total RNA extracted from

biopsies from the indicated tissues. Results were presented as mean $\pm$ standard error of the mean.

FIG 6: Transduction efficiency in different organs of mice injected with AAV9sc-CSk-SH1-Des-Luc2 (CSk-SH1) (A) or AAV9sc-CSk-SH5-Des-Luc2 (CSk-SH5) (B). AAV copy number relative to 100 ng of genomic DNA was determined for both constructs at a dose of 5×10^9 vg/mouse.

FIG 7: Nucleotide sequences of the muscle-specific regulatory elements Sk-SH1 (A, SEQ ID NO: 7), Sk-SH2 (B, SEQ ID NO: 8), Sk-SH3 (C, SEQ ID NO: 9), Sk-SH4 (D, SEQ ID NO: 10), Sk-SH5 (E, SEQ ID NO: 11), Sk-SH6 (F, SEQ ID NO: 12), Sk-SH7 (G, SEQ ID NO: 13).

FIG 8 shows difference in luciferase expression in heart and muscle tissue of mice that were intravenously injected with, from left to right, AAV9sc-Sk-SH1-Des-Luc2 (Sk-SH1, n=4), AAV9sc-Sk-SH2-Des-Luc2 (Sk-SH2, n=4), AAV9sc-Sk-SH3-Des-Luc2 (Sk-SH3, n=4), AAV9sc-Sk-SH4-Des-Luc2 (Sk-SH4, n=4), AAV9sc-Sk-SH5-Des-Luc2 (Sk-SH5, n=1), AAV9sc-Sk-SH6-Des-Luc2 (Sk-SH5, n=3) or AAV9sc-Sk-SH7-Des-Luc2 (Sk-SH6, n=4) vector compared to mice that were injected with the control vector AAV9sc-Des-Luc2 without nucleic acid regulatory element (control, no Sk-SH, n=5). Luciferase expression was measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissues at 7 weeks post-injection. Results were presented as mean $\pm$ standard error of the mean, *p<0.05; **p<0.001.

FIG 9 shows luciferase expression in selected tissues of mice that were intravenously injected with, from left to right, AAV9sc-Des-Luc2 (control, no Sk-SH, n=5), AAV9sc-Sk-SH1-Des-Luc2 (Sk-SH1, n=4), AAV9sc-Sk-SH2-Des-Luc2 (Sk-SH2, n=4), AAV9sc-Sk-SH3-Des-Luc2 (Sk-SH3, n=4), AAV9sc-Sk-SH4-Des-Luc2 (Sk-SH4, n=4), AAV9sc-Sk-SH5-Des-Luc2 (Sk-SH5, n=1), AAV9sc-Sk-SH6-Des-Luc2 (Sk-SH5, n=3), or AAV9sc-Sk-SH7-Des-Luc2 (Sk-SH6, n=4) vector. Luciferase expression was measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissue at 7 weeks post-injection. Results were presented as mean $\pm$ standard error of the mean, *p<0.05; **p<0.001.

FIG 10: (A) Schematic diagram of the AAVsc-CMV-Luc2-SV40pA plasmid construct with indication where the Cytomegalovirus promoter (CMVp) is cloned upstream of the Firefly Luciferase gene (Luc2). Abbreviations used are: ITR: viral inverted terminal repeat;

SV40pA: Simian Virus 40 polyadenylation site. **(B)** Luciferase expression in selected tissues of mice that were intravenously injected with, from left to right, AAV9sc-CMV-Luc2 (CMV, n=4), AAV9sc-Sk-SH4-Des-Luc2 (Sk-SH4, n=2), AAV9sc-CSk-SH1-Des-Luc2 (CSk-SH1, n=4), or AAV9sc-CSk-SH5-Des-Luc2 (CSk-SH5, n=4) vector. Luciferase expression was measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissue. Results were presented as mean $\pm$ standard error of the mean. **(C)** Difference in luciferase expression in heart and muscle tissue of mice that were intravenously injected with, from left to right, AAV9sc-Sk-SH4-Des-Luc2 (Sk-SH4, n=2), AAV9sc-CSk-SH1-Des-Luc2 (CSk-SH1, n=4), or AAV9sc-CSk-SH5-Des-Luc2 (CSk-SH5, n=4) vector compared to mice that were injected with vector AAV9sc-CMV-Luc2 (CMV, n=4). Luciferase expression was measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissues at 7 weeks post-injection. Results were presented as mean $\pm$ standard error of the mean.

FIG 11 shows functional fragments and the indicated transcription binding sites (TFBS) of CSk-SH1 **(A)**, CSk-SH-5 **(B)**, and Sk-SH4 **(C)** mapped on a schematic representation of respectively, SEQ ID NO:1 (A), SEQ ID NO:5 (B), and SEQ ID NO:10 (C).

FIG 12: Difference in Luc mRNA level in heart and muscle tissue of mice that were intravenously injected with, from left to right, AAV9sc-Sk-SH1-Des-Luc2 (Sk-SH1), AAV9sc-Sk-SH2-Des-Luc2 (Sk-SH2), AAV9sc-Sk-SH3-Des-Luc2 (Sk-SH3), AAV9sc-Sk-SH4-Des-Luc2 (Sk-SH4), AAV9sc-Sk-SH6-Des-Luc2 (Sk-SH6), or AAV9sc-Sk-SH7-Des-Luc2 (Sk-SH7) vector compared to mice that were injected with the control vector AAV9sc-Des-Luc2 without nucleic acid regulatory element (control, no Sk-SH). Luc mRNA levels were measured by a quantitative RT-PCR method (qRT-PCR) from total RNA extracted from biopsies from the indicated tissues. Results were presented as mean $\pm$ standard error of the mean, *p<0.05; **p<0.001.

FIG 13: Transduction efficiency in different organs of mice injected with AAV9sc-Sk-SH4-Des-MVM-Luc-pA vector **(A)** or AAV9sc-Sk-SH1-Des-MVM-Luc-pA vector **(B)**. AAV copy number relative to 100 ng of genomic DNA was determined for both vectors (n=3).

FIG 14 shows luciferase expression in heart and muscle tissue of mice that were intravenously injected with, from left to right, AAV9sc-Des-Luc (control, no Sk-SH), AAV9sc-Sk-SH4-Des-Luc (Sk-SH4), AAV9sc-Sk-SH4^a-Des-Luc (Sk-SH4^a), AAV9sc-Sk-SH4^b-Des-Luc (Sk-SH4^b), AAV9sc-Sk-SH4^c-Des-Luc (Sk-SH4^d), AAV9sc-Sk-SH4^d-Des-Luc (Sk-SH4^e), or AAV9sc-Sk-SH4^e-Des-Luc (Sk-SH4^e) vector. Luciferase expression was

measured as total flux, expressed in photons per second per centimetre squared per steradian (photons/sec/cm²/sr), released by luciferase activity in the selected tissue 5 weeks post-injection. Results were presented as mean $\pm$ standard error of the mean. The fold difference in luciferase expression in mice injected with Sk-SH4 and Sk-SH4^b compared to mice that were injected with the control vector AAV9sc-Des-Luc2 without nucleic acid regulatory element is indicated.

FIG 15: Chromatin immunoprecipitation (CHIP) assay for heart (**A, C**) and muscle (**B, D**) tissue of mice injected with AAV9sc-Sk-SH4-Des-Luc (**A, B**) or AAV9sc-CSk-SH5-Des-Luc (**C, D**) (5x10⁹vg/mouse). Antibodies specific for the transcription factors CEBP, SRF and MEF2 were used. PCR primers were designed to amplify a region of Sk-SH4 that binds CEBP and SRF, or a region of CSk-SH5 that bind CEBP and MEF2, and as negative control (-) an un-transcribed region on chromosome 17 was used. Binding events for 10³ cells were determined for each of the corresponding primer pairs. Results are presented as mean $\pm$ standard error of the mean. Significant difference compared to the negative control is indicated (t-test, * P $\leq$ 0.05).

FIG 16 shows schematic representations of single-stranded (ss) AAV plasmid constructs disclosed herein, which comprise the microdystrophin1 (MD1) (**A**, AAVss-SkSH4-Des-MVM-MD1) or follistatin (FST) (**C**, AAVssSkSH4-Des-MVM-FST-2A-Luc2) transgene regulated by the Desmin promoter operably linked to the muscle-specific nucleic acid regulatory element SkSH4 cloned upstream of the Desmin promoter. The follistatin gene was linked to the Luc2 reporter gene via a 2A peptide. The expression cassettes further comprise the Minute Virus of Mouse (MVM) intron and the 49bp synthetic Proudfoot polyadenylation site (pA). The expression cassettes are flanked by inverted terminal repeats (ITR). (**B**) Nucleotide sequence of the AAVss- SkSH4-Des-MVM-MD1 plasmid construct (SEQ ID NO:44). (**D**) Nucleotide sequence of the AAVss-SkSH4-Des-MVM-FST-2A-Luc2 plasmid construct (SEQ ID NO:45).

FIG 17: Treadmill test for MDX-SCID mice injected with, from left to right, PBS (control), AAVss-Sk-SH4-Des-MVM-FST-2A-Luc, AAVss-Sk-SH4-Des-MVM-MD1, or the combination of AAVss-Sk-SH4-Des-MVM-FST-2A-Luc and AAVss-Sk-SH4-Des-MVM-MD1. Results are expressed as the calculated distance covered by each group of mice when run on a treadmill machine.

FIG 18: (**A**) Hematoxylin/eosin staining of gastrocnemius muscle tissues of MDX/SCID mice injected with PBS (a, control), AAVss-Sk-SH4-Des-MVM-FST-2A-Luc (b), AAVss-Sk-SH4-Des-MVM-MD1 (c) or the combination of AAVss-Sk-SH4-Des-MVM-MD1 and AAVss-Sk-SH4-Des-MVM-FST-2A-Luc (d). (**B**) Quantification of centrally nucleated cells

in wild-type C57Bl/6 mice, untreated MDX/SCID mice (control), and MDX/SCID mice injected with AAVss-Sk-SH4-Des-MVM-MD1 (MD1), AAVss-Sk-SH4-Des-MVM-FST-2A-Luc (FST) or both (FST+MD1) therapeutic vectors. Statistical analysis was performed on H&E-stained transversally transected myofibers of gastronemius muscle embedded in paraffin. *** $\leq 0,0001$ ** $\leq 0,001$ * $\leq 0,05$

FIG 19: Microdystrophin1 (MD1) (A,B) and follistatin (FST) (C) mRNA levels in heart and muscle (gastrocnemius and quadriceps) tissues from mice that were intravenously injected with the AAVss-Sk-SH4-Des-MVM-MD1 vector (A), the combination of AAVss-Sk-SH4-Des-MVM-MD1 and AAVss-Sk-SH4-Des-MVM-FST-2A-Luc (B), or AAVss-Sk-SH4-Des-MVM-FST (C) relative to expression of endogenous housekeeping gene (GAPDH: Glyceraldehyde-3-phosphate dehydrogenase). Results are presented as relative expression of MD1 or FST ($\Delta\Delta CT$).

FIG 20 shows schematic representations of self-complementary (sc) AAV constructs disclosed herein, which comprise the luciferase (Luc) transgene regulated by (a) the cardiac and skeletal muscle-specific desmin (Desmin) promoter, (b) the SPc5-12 promoter, (c) the cytomegalovirus (CMV) promoter, (d) the Desmin promoter operably linked to a muscle-specific (Sk-SH) nucleic acid regulatory element cloned upstream of the Desmin promoter, or (e) the SPc5-12 promoter operably linked to a muscle-specific (Sk-SH) nucleic acid regulatory element cloned upstream of the SPc5-12 promoter. Expression cassettes (a), (b), (d) and (e) further comprise the Minute Virus of Mouse (MVM) intron and a polyadenylation signal (pA). The expression cassettes are flanked by inverted terminal repeats (ITR).

FIG. 21: Difference in Luc mRNA levels in selected tissues of CB17/lcrTac/Prkdcscid mice that were intravenously injected with, from left to right, AAVsc-CMV-Luc, AAVsc-Des-MVM-Luc, AAVsc-Sk-SH4-Des-MVM-Luc, AAVsc-SPc5-12-MVM-Luc or AAVsc-Sk-SH4-SPc5-12-MVM-Luc vector as schematically shown in Figure 20, compared to mice that were injected with AAVsc-CMV-Luc vector. Luc mRNA levels were measured by qRT-PCR from total RNA extracted from biopsies of the indicated tissues. Results were presented as mean $\pm$ standard error of the mean, * $p < 0.05$; ** $p < 0.001$.

DESCRIPTION

As used herein, the singular forms “a”, “an”, and “the” include both singular and plural referents unless the context clearly dictates otherwise.

The terms “comprising”, “comprises” and “comprised of” as used herein are synonymous with “including”, “includes” or “containing”, “contains”, and are inclusive or open-ended and

do not exclude additional, non-recited members, elements or method steps. The terms also encompass “consisting of” and “consisting essentially of”, which enjoy well-established meanings in patent terminology.

5 The recitation of numerical ranges by endpoints includes all numbers and fractions subsumed within the respective ranges, as well as the recited endpoints.

The terms “about” or “approximately” as used herein when referring to a measurable value such as a parameter, an amount, a temporal duration, and the like, are meant to encompass variations of and from the specified value, such as variations of +/-10% or less, preferably +/-5% or less, more preferably +/-1% or less, and still more preferably +/-
10 0.1% or less of and from the specified value, insofar such variations are appropriate to perform in the disclosed invention. It is to be understood that the value to which the modifier “about” refers is itself also specifically, and preferably, disclosed.

Whereas the terms “one or more” or “at least one”, such as one or more members or at least one member of a group of members, is clear per se, by means of further
15 exemplification, the term encompasses inter alia a reference to any one of said members, or to any two or more of said members, such as, e.g., any ≥ 3 , ≥ 4 , ≥ 5 , ≥ 6 or ≥ 7 etc. of said members, and up to all said members. In another example, “one or more” or “at least one” may refer to 1, 2, 3, 4, 5, 6, 7 or more.

The discussion of the background to the invention herein is included to explain the context
20 of the invention. This is not to be taken as an admission that any of the material referred to was published, known, or part of the common general knowledge in any country as of the priority date of any of the claims.

Throughout this disclosure, various publications, patents and published patent specifications are referenced by an identifying citation.
25

Unless otherwise defined, all terms used in disclosing the invention, including technical and scientific terms, have the meaning as commonly understood by one of ordinary skill in
30 the art to which this invention belongs. By means of further guidance, term definitions are included to better appreciate the teaching of the invention. When specific terms are defined in connection with a particular aspect of the invention or a particular embodiment of the invention, such connotation is meant to apply throughout this specification, i.e., also in the context of other aspects or embodiments of the invention, unless otherwise defined.

In the following passages, different aspects or embodiments of the invention are defined in more detail. Each aspect or embodiment so defined may be combined with any other aspect(s) or embodiment(s) unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous.

Reference throughout this specification to "one embodiment", "an embodiment" means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present invention. Thus, appearances of the phrases "in one embodiment" or "in an embodiment" in various places throughout this specification are not necessarily all referring to the same embodiment, but may. Furthermore, the particular features, structures or characteristics may be combined in any suitable manner, as would be apparent to a person skilled in the art from this disclosure, in one or more embodiments. Furthermore, while some embodiments described herein include some but not other features included in other embodiments, combinations of features of different embodiments are meant to be within the scope of the invention, and form different embodiments, as would be understood by those in the art. For example, in the appended claims, any of the claimed embodiments can be used in any combination.

For general methods relating to the invention, reference is made *inter alia* to well-known textbooks, including, e.g., "Molecular Cloning: A Laboratory Manual, 2nd Ed." (Sambrook *et al.*, 1989), "Current Protocols in Molecular Biology" (Ausubel *et al.*, 1987).

In an aspect, the invention relates to a nucleic acid regulatory element for enhancing muscle-specific gene expression comprising, consisting essentially of (i.e., the regulatory element may for instance additionally comprise sequences used for cloning purposes, but the indicated sequences make up the essential part of the regulatory element, e.g. they do not form part of a larger regulatory region such as a promoter), or consisting of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, or a functional fragment thereof (i.e. a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ

ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, or of a sequence having high percentage sequence identity to any of said sequences).

A 'regulatory element' as used herein refers to a transcriptional control element, in particular a non-coding cis-acting transcriptional control element, capable of regulating and/or controlling transcription of a gene, in particular tissue-specific transcription of a gene. Regulatory elements comprise at least one transcription factor binding site (TFBS), more in particular at least one binding site for a tissue-specific transcription factor, most particularly at least one binding site for a muscle-specific transcription factor. Typically, regulatory elements as used herein increase or enhance promoter-driven gene expression when compared to the transcription of the gene from the promoter alone, without the regulatory elements. Thus, regulatory elements particularly comprise enhancer sequences, although it is to be understood that the regulatory elements enhancing transcription are not limited to typical far upstream enhancer sequences, but may occur at any distance of the gene they regulate. Indeed, it is known in the art that sequences regulating transcription may be situated either upstream (e.g. in the promoter region) or downstream (e.g. in the 3'UTR) of the gene they regulate *in vivo*, and may be located in the immediate vicinity of the gene or further away. Of note, although regulatory elements as disclosed herein typically comprise naturally occurring sequences, combinations of (parts of) such regulatory elements or several copies of a regulatory element, i.e. regulatory elements comprising non-naturally occurring sequences, are themselves also envisaged as regulatory element. Regulatory elements as used herein may comprise part of a larger sequence involved in transcriptional control, e.g. part of a promoter sequence. However, regulatory elements alone are typically not sufficient to initiate transcription, but require a promoter to this end. The regulatory elements disclosed herein are provided as nucleic acid molecules, i.e. isolated nucleic acids, or isolated nucleic acid molecules. Said nucleic acid regulatory element hence have a sequence which is only a small part of the naturally occurring genomic sequence and hence is not naturally occurring as such, but is isolated therefrom.

The term "nucleic acid" as used herein typically refers to an oligomer or polymer (preferably a linear polymer) of any length composed essentially of nucleotides. A nucleotide unit commonly includes a heterocyclic base, a sugar group, and at least one, e.g. one, two, or three, phosphate groups, including modified or substituted phosphate groups. Heterocyclic bases may include *inter alia* purine and pyrimidine bases such as adenine (A), guanine (G), cytosine (C), thymine (T) and uracil (U) which are widespread in naturally-occurring nucleic acids, other naturally-occurring bases (e.g., xanthine, inosine,

hypoxanthine) as well as chemically or biochemically modified (e.g., methylated), non-natural or derivatised bases. Sugar groups may include *inter alia* pentose (pentofuranose) groups such as preferably ribose and/or 2-deoxyribose common in naturally-occurring nucleic acids, or arabinose, 2-deoxyarabinose, threose or hexose sugar groups, as well as modified or substituted sugar groups. Nucleic acids as intended herein may include naturally occurring nucleotides, modified nucleotides or mixtures thereof. A modified nucleotide may include a modified heterocyclic base, a modified sugar moiety, a modified phosphate group or a combination thereof. Modifications of phosphate groups or sugars may be introduced to improve stability, resistance to enzymatic degradation, or some other useful property. The term "nucleic acid" further preferably encompasses DNA, RNA and DNA/RNA hybrid molecules, specifically including hnRNA, pre-mRNA, mRNA, cDNA, genomic DNA, amplification products, oligonucleotides, and synthetic (e.g., chemically synthesised) DNA, RNA or DNA/RNA hybrids. A nucleic acid can be naturally occurring, e.g., present in or isolated from nature; or can be non-naturally occurring, e.g., recombinant, i.e., produced by recombinant DNA technology, and/or partly or entirely, chemically or biochemically synthesised. A "nucleic acid" can be double-stranded, partly double stranded, or single-stranded. Where single-stranded, the nucleic acid can be the sense strand or the antisense strand. In addition, nucleic acid can be circular or linear.

As used herein "transcription factor binding site", "transcription factor binding sequence" or "TFBS" refers to a sequence of a nucleic acid region to which transcription factors bind. Non-limiting examples of TFBS include binding sites for transcription factor 3, also known as TCF3 or E2A; binding sites for nuclear factor I, also known as NF1; binding sites for CCAAT-enhancer-binding protein, also known as C/EBP; binding sites for myogenic differentiation, also known as MyoD; binding sites for sterol regulatory element-binding protein, also known as SREBP; binding sites for leukemia/lymphoma-related factor, also known as LRF; binding sites for protein 53, also known as p53; binding sites for hepatocyte nuclear factor 3-alpha, also known as HNF3a; binding sites for hepatocyte nuclear factor 3-beta, also known as HNF3b; binding sites for hepatocyte nuclear factor 4, also known as HNF4; binding sites for myocyte-specific enhancer factor 2A, also known as MEF2A or RSRFC4; binding sites for peroxisome proliferator-activated receptor, also known as PPAR; binding sites for serum response factor, also known as SRF; binding sites for transcription activator-like protein 1b, also known as Tal1_b. Transcription factor binding sites may be found in databases such as Transfac®.

Sequences disclosed herein may be part of sequences of regulatory elements capable of controlling transcription of muscle-specific genes *in vivo*, in particular controlling the

following genes: desmin also known as DES, CSM1 or CSM2; actinin, alpha 2 also known as ACTN2 or CMD1AA; filamin-C (FLNC) also known as actin-binding-like protein (ABLP), filamin-2 (FLN2), ABP-280, ABP280A, ABPA, ABPL, MFM5, or MPD4; sarcoplasmic/endoplasmic reticulum calcium ATPase 1 also known as ATP2A1, ATP2A, or SERCA1; troponin I type 1 (slow skeletal muscle) also known as TNNI1, SSTNI, or TNN1; myosin light chain phosphorylatable fast skeletal muscle (MYLPF); myosin-1 also known as MYH1, MYHSA1, MYHa; MyHC-2X/D, or MyHC-2x; tropomyosin alpha-3 chain also known as TPM3, CFTD, NEM1, OK/SW-cl.5, TM-5, TM3, TM30, TM30nm, TM5, TPMsk3, TRK, hTM5, or hscp30; and ankyrin repeat domain-containing protein 2 also known as ANKRD2, or ARPP. Accordingly, in embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from Des regulatory elements, i.e. regulatory elements that control expression of the Des gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:1, SEQ ID NO: 17, SEQ ID NO:2, SEQ ID NO:18, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from ACTN2 regulatory elements, i.e. regulatory elements that control expression of the ACTN2 gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:3, SEQ ID NO:19, SEQ ID NO:4, SEQ ID NO:20, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from FLNC regulatory elements, i.e. regulatory elements that control expression of the FLNC gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:5, SEQ ID NO:21, SEQ ID NO:6, SEQ ID NO:22, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from ATP2A1 regulatory elements, i.e. regulatory elements that control expression of the ATP2A1 gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:7, SEQ ID NO:23, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from TNNI1 regulatory elements, i.e. regulatory elements that control expression of the TNNI1 gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:8, SEQ ID NO:24, SEQ ID NO:9, SEQ ID NO:25, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from MYLPF regulatory elements, i.e. regulatory elements that control expression of the MYLPF gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:10, SEQ ID NO:26, or functional fragments thereof, such as regulatory elements comprising or consisting of SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40 or SEQ ID NO:41. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from MYH1 regulatory elements, i.e. regulatory elements that control expression of the MYH1 gene *in vivo*, e.g. regulatory elements comprising SEQ ID

NO:11, SEQ ID NO:27, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from TPM3 regulatory elements, i.e. regulatory elements that control expression of the TPM3 gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:12, SEQ ID NO:28, or functional fragments thereof. In embodiments, the nucleic acid regulatory elements disclosed herein comprise a sequence from ANKRD2 regulatory elements, i.e. regulatory elements that control expression of the ANKRD2 gene *in vivo*, e.g. regulatory elements comprising SEQ ID NO:13, SEQ ID NO:29, or functional fragments thereof.

As used herein, the terms "identity" and "identical" and the like refer to the sequence similarity between two polymeric molecules, e.g., between two nucleic acid molecules, e.g., two DNA molecules. Sequence alignments and determination of sequence identity can be done, e.g., using the Basic Local Alignment Search Tool (BLAST) originally described by Altschul et al. 1990 (J Mol Biol 215: 403-10), such as the "Blast 2 sequences" algorithm described by Tatusova and Madden 1999 (FEMS Microbiol Lett 174: 247-250). Typically, the percentage sequence identity is calculated over the entire length of the sequence. As used herein, the term "substantially identical" denotes at least 90%, preferably at least 95%, such as 95%, 96%, 97%, 98% or 99%, sequence identity.

The term 'functional fragment' as used in the application refers to fragments of the regulatory element sequences disclosed herein that retain the capability of regulating muscle-specific expression, i.e. they can still confer tissue specificity and they are capable of regulating expression of a (trans)gene in the same way (although possibly not to the same extent) as the sequence from which they are derived. Functional fragments may preferably comprise at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100, at least 120, at least 150, at least 200, at least 250, at least 300, at least 350, or at least 400 contiguous nucleotides from the sequence from which they are derived. Also preferably, functional fragments may comprise at least 1, more preferably at least 2, at least 3, or at least 4, even more preferably at least 5, at least 10, or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which they are derived.

"Muscle-specific expression" as used in the application, refers to the preferential or predominant expression of a (trans)gene (as RNA and/or polypeptide) in muscles or muscle tissue, as compared to other (i.e. non-muscle) tissues. According to particular embodiments, at least 50%, more particularly at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99% or 100% of the (trans)gene expression occurs within muscle. According to a particular embodiment,

muscle-specific expression entails that there is no 'leakage' of expressed gene product to other organs or tissue than muscle, such as lung, liver, brain, kidney and/or spleen.

As used herein "cardiac and skeletal muscle-specific expression" refers to the preferential or predominant expression of a (trans)gene in heart, in particular heart muscle, and skeletal muscle. According to particular embodiments, at least 50%, more particularly at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99% or 100% of the (trans)gene expression occurs within heart and skeletal muscle. Thus, according to particular embodiments, less than 10%, less than 5%, less than 2% or even less than 1% of the (trans)gene expression occurs in an organ or tissue other than heart and skeletal muscle.

As used herein "skeletal muscle-specific expression" refers to the preferential or predominant expression of a (trans)gene in skeletal muscle. According to particular embodiments, at least 50%, more particularly at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99% or 100% of the (trans)gene expression occurs within skeletal muscle. Thus, according to particular embodiments, less than 10%, less than 5%, less than 2% or even less than 1% of the (trans)gene expression occurs in an organ or tissue other than skeletal muscle.

The same applies *mutatis mutandis* for myocyte-specific and myoblast-specific expression, which may be considered as a particular form of muscle-specific expression. Throughout the application, where muscle-specific is mentioned in the context of expression, myocyte-specific and myoblast-specific expression are also explicitly envisaged. Similarly, where cardiac and skeletal muscle-specific expression is used in the application, cardiomyocyte and skeletal myocyte-specific expression and cardiac myoblast and skeletal myoblast-specific expression is also explicitly envisaged. Similarly, where skeletal muscle-specific expression is used in the application, skeletal myocyte-specific and skeletal myoblast-specific expression is also explicitly envisaged.

As used herein, the terms "heart muscle" or "cardiac muscle" refer to the autonomically regulated, striated muscle type found in the heart.

As used herein, the term "skeletal muscle" refers to the voluntarily controlled, striated muscle type that is attached to the skeleton. Non-limiting examples of skeletal muscle include the biceps, the triceps, the quadriceps, the tibialis interior, and the gastrocnemius muscle.

The term "myocyte," as used herein, refers to a cell that has been differentiated from a progenitor myoblast such that it is capable of expressing muscle-specific phenotype under

appropriate conditions. Terminally differentiated myocytes fuse with one another to form myotubes, a major constituent of muscle fibers. The term "myocyte" also refers to myocytes that are de-differentiated. The term includes cells *in vivo* and cells cultured *ex vivo* regardless of whether such cells are primary or passaged.

- 5 The term "myoblast" as used herein, refers to an embryonic cell in the mesoderm that differentiates to give rise to a muscle cell or myocyte. The term includes cells *in vivo* and cells cultured *ex vivo* regardless of whether such cells are primary or passaged.

In embodiments, the invention relates to a nucleic acid regulatory element for enhancing muscle-specific gene expression comprising, consisting essentially of, or consisting of a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment comprises at least 20, preferably at least 25, more preferably at least 50, at least 100, at least 200 or at least 250, contiguous nucleotides from the sequence from which it is derived, and wherein said functional fragment comprises at least 1, preferably at least 5, more preferably at least 10 or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which it is derived.

In further embodiments, the invention relates to a nucleic acid regulatory element for enhancing cardiac and skeletal muscle-specific gene expression comprising a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment comprise at least 20, preferably at least 25, more preferably at least 50, at least 100, at least 200 or at least 250, contiguous nucleotides from the sequence from which it is derived, and wherein said functional fragment comprises at least 1, preferably at least 5, more preferably at least 10 or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which it is derived.

In yet further embodiments, the invention relates to a nucleic acid regulatory element for enhancing cardiac and skeletal muscle-specific gene expression comprising a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:1 and SEQ ID

- NO:5, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment consists of the nucleotide sequence from position 33 to 58 in SEQ ID NO:1; the nucleotide sequence from position 90 to 142 in SEQ ID NO:1; the nucleotide sequence from position 143 to 233 in SEQ ID NO:1; the nucleotide sequence from position 240 to 310 in SEQ ID NO:1; the nucleotide sequence from position 90 to 233 in SEQ ID NO:1; the nucleotide sequence from position 47 to 130 in SEQ ID NO:5; the nucleotide sequence from position 252 to 293 in SEQ ID NO:5; or the nucleotide sequence from position 330 to 450 in SEQ ID NO:5.
- 10 In further embodiments, the invention relates to a nucleic acid regulatory element for enhancing skeletal muscle-specific gene expression comprising a functional fragment of a sequence selected from the group consisting of: SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, wherein said functional fragment comprise at least 20, preferably at least 25, more preferably at least 50, at least 100, at least 200 or at least 250, contiguous nucleotides from the sequence from which it is derived, and wherein said functional fragment comprises at least 1, preferably at least 5, more preferably at least 10 or at least 15, of the transcription factor binding sites (TFBS) that are present in the sequence from which it is derived.

- In yet further embodiments, the invention relates to a nucleic acid regulatory element for enhancing skeletal muscle-specific gene expression comprising a functional fragment of SEQ ID NO:10 or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to SEQ ID NO:10, wherein said functional fragment consists of the nucleotide sequence from position 10 to 180 in SEQ ID NO:10; the nucleotide sequence from position 190 to 240 in SEQ ID NO:10; the nucleotide sequence from position 241 to 300 in SEQ ID NO:10; the nucleotide sequence from position 241 to 360 in SEQ ID NO:10; or the nucleotide sequence from position 380 to 420 in SEQ ID NO:10.

- In certain embodiments, the nucleic acid regulatory elements of the invention comprise or consist of a sequence selected from the group consisting of: the nucleotide sequence from position 33 to 58 in SEQ ID NO:1; the nucleotide sequence from position 90 to 142 in SEQ ID NO:1; the nucleotide sequence from position 143 to 233 in SEQ ID NO:1; the nucleotide sequence from position 240 to 310 in SEQ ID NO:1; the nucleotide sequence

from position 90 to 233 in SEQ ID NO:1; the nucleotide sequence from position 47 to 130 in SEQ ID NO:5; the nucleotide sequence from position 252 to 293 in SEQ ID NO:5; the nucleotide sequence from position 330 to 450 in SEQ ID NO:5; the nucleotide sequence from position 10 to 180 in SEQ ID NO:10; the nucleotide sequence from position 190 to 240 in SEQ ID NO:10; the nucleotide sequence from position 241 to 300 in SEQ ID NO:10; the nucleotide sequence from position 241 to 360 in SEQ ID NO:10; the nucleotide sequence from position 380 to 420 in SEQ ID NO:10; or a sequence having at least 95% identity to any of said sequences.

In certain embodiments, the nucleic acid regulatory elements of the invention comprise or consist of a sequence selected from the group consisting of: the nucleotide sequence from position 33 to 310 in SEQ ID NO:1; the nucleotide sequence from position 47 to 450 in SEQ ID NO:5; the nucleotide sequence from position 10 to 420 in SEQ ID NO:10, or a sequence having at least 95% identity to any of said sequences.

In further embodiments, the invention provides a nucleic acid regulatory element for enhancing muscle-specific gene expression comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

In yet further embodiments, the invention provides for a nucleic acid regulatory element for enhancing cardiac and skeletal muscle-specific gene expression comprising a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

In yet further embodiments, the invention provides for a nucleic acid regulatory element for enhancing skeletal muscle-specific gene expression comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

In further embodiments, the nucleic acid regulatory elements for enhancing muscle-specific gene expression, comprise, consist essentially of, or consist of a sequence selected from the group consisting of: SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, a functional fragment thereof comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

In further embodiments, the nucleic acid regulatory elements for enhancing cardiac and skeletal muscle-specific gene expression comprise, consist essentially of, or consist of a sequence selected from the group consisting of: SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, a functional fragment thereof comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, or SEQ ID NO:6, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

In further embodiments, the nucleic acid regulatory elements for enhancing skeletal muscle-specific gene expression comprise, consist essentially of, or consist of a sequence selected from the group consisting of: SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, a functional fragment thereof comprising SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13, or a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences.

It is also possible to make nucleic acid regulatory elements that comprise an artificial sequence by combining two or more identical or different sequences disclosed herein or functional fragments thereof. Accordingly, in certain embodiments a nucleic acid regulatory element for enhancing muscle-specific gene expression is provided comprising at least two sequences selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, , SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a sequence having at least 90%, preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, or a functional fragment thereof.

Also disclosed herein is a nucleic acid regulatory element for enhancing muscle-specific gene expression, in particular cardiac and skeletal muscle-specific gene expression, comprising at least two sequences selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, a sequence
 5 having at least 90%, preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, or a fragment thereof.

Also disclosed herein is a nucleic acid regulatory element for enhancing muscle-specific gene expression, in particular skeletal muscle-specific gene expression, comprising at least two sequences selected from the group consisting of: SEQ ID NO:7, SEQ ID NO:8,
 10 SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a sequence having at least 90%, preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, or a fragment thereof.

For example, disclosed herein is a nucleic acid regulatory element comprising, consisting essentially of, or consisting of SEQ ID NO:1 and SEQ ID NO:5; a nucleic acid regulatory element comprising, consisting essentially of, or consisting of SEQ ID NO:1 and SEQ ID
 15 NO:10; a nucleic acid regulatory element comprising, consisting essentially of, or consisting of SEQ ID NO:1, SEQ ID NO:5, and SEQ ID NO:10; a nucleic acid regulatory element comprising, consisting essentially of, or consisting of 2, 3, 4, or 5 repeats, e.g. tandem repeats, of SEQ ID NO:1; a nucleic acid regulatory element comprising, consisting
 20 essentially of, or consisting of 2, 3, 4, or 5 repeats, e.g. tandem repeats, of SEQ ID NO:5; or a nucleic acid regulatory element comprising, consisting essentially of, or consisting of 2, 3, 4, or 5 repeats, e.g. tandem repeats, of SEQ ID NO:10.

Particular examples of nucleic acid regulatory elements that comprise an artificial sequence include the regulatory elements that are obtained by rearranging the
 25 transcription factor binding sites (TFBS) that are present in the sequences disclosed herein. Said rearrangement may encompass changing the order of the TFBSs and/or changing the position of one or more TFBSs relative to the other TFBSs and/or changing the copy number of one or more of the TFBSs. For example, also disclosed herein is a nucleic acid regulatory element for enhancing muscle-specific gene expression, in
 30 particular cardiac and skeletal muscle-specific gene expression, comprising binding sites for E2A, HNH1, NF1, C/EBP, LRF, MyoD, and SREBP; or for E2A, NF1, p53, C/EBP, LRF, and SREBP; or for E2A, HNH1, HNF3a, HNF3b, NF1, C/EBP, LRF, MyoD, and SREBP; or for E2A, HNF3a, NF1, C/EBP, LRF, MyoD, and SREBP; or for E2A, HNF3a, NF1, CEBP, LRF, MyoD, and SREBP; or for HNF4, NF1, RSRFC4, C/EBP, LRF, and MyoD, or
 35 NF1, PPAR, p53, C/EBP, LRF, and MyoD. For example, also disclosed herein is a nucleic

acid regulatory element for enhancing muscle-specific gene expression, in particular skeletal muscle-specific gene expression, comprising binding sites for E2A, NF1, SRFC, p53, C/EBP, LRF, and MyoD; or for E2A, NF1, C/EBP, LRF, MyoD, and SREBP; or for E2A, HNF3a, C/EBP, LRF, MyoD, SEREBP, and Tal1_b; or for E2A, SRF, p53, C/EBP, LRF, MyoD, and SREBP; or for HNF4, NF1, RSRFC4, C/EBP, LRF, and SREBP; or for E2A, HNF3a, HNF3b, NF1, SRF, C/EBP, LRF, MyoD, and SREBP; or for E2A, CEBP, and MyoD. In further examples, these nucleic acid regulatory elements comprise at least two, such as 2, 3, 4, or more copies of one or more of the recited TFBSs.

In case the regulatory element is provided as a single stranded nucleic acid, e.g. when using a single-stranded AAV vector, the complement strand is considered equivalent to the disclosed sequences. Hence, also disclosed herein is a nucleic acid regulatory element for enhancing muscle-specific gene expression comprising, consisting essentially of, or consisting of the complement of a sequence described herein, in particular a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, or a functional fragment thereof.

Also disclosed herein is a nucleic acid regulatory element for enhancing muscle-specific gene expression hybridizing under stringent conditions to a nucleic acid regulatory element described herein, in particular to the nucleic acid regulatory element comprising, consisting essentially of, or consisting of a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, a sequence having at least 90%, preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity to any of these sequences, a functional fragment thereof, or to its complement. Said nucleic acid regulatory elements do not need to be of equal length as the sequence they hybridize to. In preferred embodiments, the size of said hybridizing nucleic acid regulatory element does not differ more than 25% in length, in particular 20% in length, more in particular 15% in length, most in particular 10% in length from the sequence it hybridizes to.

The expression 'hybridize under stringent conditions', refers to the ability of a nucleic acid molecule to hybridize to a target nucleic acid molecule under defined conditions of temperature and salt concentration. Typically, stringent hybridization conditions are no

more than 25°C to 30°C (for example, 20°C, 15°C, 10°C or 5°C) below the melting temperature (T_m) of the native duplex. Methods of calculating T_m are well known in the art. By way of non-limiting example, representative salt and temperature conditions for achieving stringent hybridization are: 1x SSC, 0.5% SDS at 65°C. The abbreviation SSC
5 refers to a buffer used in nucleic acid hybridization solutions. One liter of the 20x (twenty times concentrate) stock SSC buffer solution (pH 7.0) contains 175.3 g sodium chloride and 88.2 g sodium citrate. A representative time period for achieving hybridization is 12 hours.

Preferably the regulatory elements as described herein are fully functional while being
10 only of limited length. This allows their use in vectors or nucleic acid expression cassettes without unduly restricting their payload capacity. Accordingly, in embodiments, the regulatory element disclosed herein is a nucleic acid of 1500 nucleotides or less, 1000 nucleotides or less, 900 nucleotides or less, 800 nucleotides or less, 700 nucleotides or less, more preferably 600 nucleotides or less, such as 550 nucleotides or less, 500
15 nucleotides or less, 450 nucleotides or less, 400 nucleotides or less, 350 nucleotides or less, or 300 nucleotides or less (i.e. the nucleic acid regulatory element has a maximal length of 1500 nucleotides, 1000 nucleotides, 900 nucleotides, 800 nucleotides, 700 nucleotides, preferably 600 nucleotides, such as 550 nucleotides, 500 nucleotides, 450 nucleotides, 400 nucleotides, 350 nucleotides, or 300 nucleotides).

20 However, it is to be understood that the disclosed nucleic acid regulatory elements retain regulatory activity (i.e. with regard to specificity and/or activity of transcription) and thus they particularly have a minimum length of 20 nucleotides, 25 nucleotides, 30 nucleotides, 35 nucleotides, 40 nucleotides, 45 nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, 250 nucleotides, 300 nucleotides, 350 nucleotides or 400
25 nucleotides.

In certain embodiments, the invention provides for a nucleic acid regulatory element of 1000 nucleotides or less, preferably 600 nucleotides or less, such as 550 nucleotides or less, 500 nucleotides or less, 450 nucleotides or less, 400 nucleotides or less, 350 nucleotides or less, or 300 nucleotides or less, for enhancing muscle-specific gene
30 expression comprising a sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, a sequence having at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, such as 95%, 96%, 97%, 98%, or 99%, identity
35 to any of these sequences, or a functional fragment thereof.

The nucleic acid regulatory elements disclosed herein may be used in a nucleic acid expression cassette. Accordingly, in an aspect the invention provides for the use of the nucleic acid regulatory elements as described herein in a nucleic acid expression cassette.

- 5 In an aspect the invention provides a nucleic acid expression cassette comprising a nucleic acid regulatory element as described herein, operably linked to a promoter. In embodiments, the nucleic acid expression cassette does not contain a transgene. Such nucleic acid expression cassette may be used to drive expression of an endogenous gene. In preferred embodiments, the nucleic acid expression cassette comprises a nucleic acid regulatory element as described herein, operably linked to a promoter and a transgene.

As used herein, the term 'nucleic acid expression cassette' refers to nucleic acid molecules that include one or more transcriptional control elements (such as, but not limited to promoters, enhancers and/or regulatory elements, polyadenylation sequences, and introns) that direct (trans)gene expression in one or more desired cell types, tissues or organs. Typically, they will also contain a transgene, although it is also envisaged that a nucleic acid expression cassette directs expression of an endogenous gene in a cell into which the nucleic acid cassette is inserted.

The term 'operably linked' as used herein refers to the arrangement of various nucleic acid molecule elements relative to each such that the elements are functionally connected and are able to interact with each other. Such elements may include, without limitation, a promoter, an enhancer and/or a regulatory element, a polyadenylation sequence, one or more introns and/or exons, and a coding sequence of a gene of interest to be expressed (i.e., the transgene). The nucleic acid sequence elements, when properly oriented or operably linked, act together to modulate the activity of one another, and ultimately may affect the level of expression of the transgene. By modulate is meant increasing, decreasing, or maintaining the level of activity of a particular element. The position of each element relative to other elements may be expressed in terms of the 5' terminus and the 3' terminus of each element, and the distance between any particular elements may be referenced by the number of intervening nucleotides, or base pairs, between the elements. As understood by the skilled person, operably linked implies functional activity, and is not necessarily related to a natural positional link. Indeed, when used in nucleic acid expression cassettes, the regulatory elements will typically be located immediately upstream of the promoter (although this is generally the case, it should definitely not be interpreted as a limitation or exclusion of positions within the nucleic acid expression

cassette), but this needs not be the case *in vivo*. E.g., a regulatory element sequence naturally occurring downstream of a gene whose transcription it affects is able to function in the same way when located upstream of the promoter. Hence, according to a specific embodiment, the regulatory or enhancing effect of the regulatory element is position-independent.

In particular embodiments, the nucleic acid expression cassette comprises one nucleic acid regulatory element as described herein. In alternative embodiments, the nucleic acid expression cassette comprises two or more, such as, e.g., 2, 3, 4, 5, 6, 7, 8, 9, or 10, nucleic acid regulatory elements as described herein, i.e. they are combined modularly to enhance their regulatory (and/or enhancing) effect. In further embodiments, at least two of the two or more nucleic acid regulatory elements are identical or substantially identical. In yet further embodiments, all of the two or more regulatory elements are identical or substantially identical. The copies of the identical or substantially identical nucleic acid regulatory elements may be provided as tandem repeats in the nucleic acid expression cassette. In alternative further embodiments, at least two of the two or more nucleic acid regulatory elements are different from each other. The nucleic acid expression cassette may also comprise a combination of identical and substantially identical nucleic acid regulatory elements and non-identical nucleic acid regulatory elements.

For example, the nucleic acid expression cassette may comprise a nucleic acid regulatory element comprising SEQ ID NO:1, and a nucleic acid regulatory element comprising SEQ ID NO:5; or the nucleic acid expression cassette may comprise a nucleic acid regulatory element comprising SEQ ID NO:1 and a nucleic acid regulatory element comprising SEQ ID NO:10; or the nucleic acid expression cassette may comprise a nucleic acid regulatory element comprising SEQ ID NO:1, a nucleic acid regulatory element comprising SEQ ID NO:5, and a nucleic acid regulatory element comprising SEQ ID NO:1; or the nucleic acid regulatory element may comprise 2, 3, 4, or 5 nucleic acid regulatory elements comprising SEQ ID NO:1; or the nucleic acid regulatory element may comprise 2, 3, 4, or 5 nucleic acid regulatory elements comprising SEQ ID NO:5; or the nucleic acid regulatory element may comprise 2, 3, 4, or 5 nucleic acid regulatory elements comprising SEQ ID NO:10.

As used in the application, the term 'promoter' refers to nucleic acid sequences that regulate, either directly or indirectly, the transcription of corresponding nucleic acid coding sequences to which they are operably linked (e.g. a transgene or endogenous gene). A promoter may function alone to regulate transcription or may act in concert with one or more other regulatory sequences (e.g. enhancers or silencers, or regulatory elements). In the context of the present application, a promoter is typically operably linked to a

regulatory element as disclosed herein to regulate transcription of a (trans)gene. When a regulatory element as described herein is operably linked to both a promoter and a transgene, the regulatory element can (1) confer a significant degree of muscle-specific, in particular cardiac and skeletal muscle-specific or skeletal muscle-specific, expression *in vivo* (and/or in myoblasts, myocytes, or muscle-derived cell lines, in particular cardiac and skeletal or skeletal myoblasts, cardiac and skeletal or skeletal myocytes, or cardiac and skeletal muscle or skeletal muscle-derived cell lines *in vitro*) of the transgene, and/or (2) can increase the level of expression of the transgene in muscle, in particular cardiac and skeletal muscle or skeletal muscle (and/or in myoblasts, myocytes, or muscle-derived cell lines, in particular cardiac and skeletal or skeletal myoblasts, cardiac and skeletal or skeletal myocytes, or cardiac and skeletal muscle or skeletal muscle-derived cell lines *in vitro*).

The promoter may be homologous (i.e. from the same species as the animal, in particular mammal, to be transfected with the nucleic acid expression cassette) or heterologous (i.e. from a source other than the species of the animal, in particular mammal, to be transfected with the expression cassette). As such, the source of the promoter may be any virus, any unicellular prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, or may even be a synthetic promoter (i.e. having a non-naturally occurring sequence), provided that the promoter is functional in combination with the regulatory elements described herein. In preferred embodiments, the promoter is a mammalian promoter, in particular a murine or human promoter.

The promoter may be an inducible or constitutive promoter.

The enrichment in muscle-specific TFBS in the nucleic acid regulatory elements disclosed herein in principle allows the regulatory elements to direct muscle-specific expression even from a promoter that itself is not muscle-specific. Hence, the regulatory elements disclosed herein can be used in nucleic acid expression cassettes in conjunction with their natural promoter, as well as with another promoter. Preferably, the nucleic acid expression cassettes disclosed herein comprise a muscle-specific promoter. This to increase muscle-specificity and/or avoid leakage of expression in other tissues. Non-limiting examples of muscle-specific promoters, include the desmin (DES) promoter, the alpha 2 actinin (ACTN2) promoter, the filamin-C (FLNC) promoter, the sarcoplasmic/endoplasmic reticulum calcium ATPase 1 (ATP2A1) promoter, the troponin I type 1 (TNNI1) promoter, the myosin-1 (MYH1) promoter, the phosphorylatable, fast skeletal muscle myosin light chain (MYLPP) promoter, the alpha-3 chain tropomyosin (TPM3) promoter, the ankyrin repeat domain-containing protein 2 (ANKRD2) promoter the myosin heavy-chain (MHC)

promoter, the myosin light-chain (MLC) promoter, the muscle creatine kinase (MCK) promoter, synthetic muscle promoters as described in Li et al. (1999. Nat Biotechnol. 17:241-245), such as the SPc5-12 promoter, the muscle creatine kinase (MCK) promoter, the dMCK promoter and the tMCK promoter consisting of respectively, a double or triple
5 tandem of the MCK enhancer to the MCK basal promoter as described in Wang et al. (2008. Gene Ther. 15:1489-1499).

In particularly preferred embodiments, the promoter is a mammalian muscle-specific promoter, in particular a murine or human muscle-specific promoter.

In preferred embodiments, the promoter is from the desmin gene, in particular the murine
10 or human desmin gene, such as the promoter as defined in SEQ ID NO: 16. For example, the murine desmin promoter is commercially available as pDRIVE-mDesmin (Invivogen). The desmin promoter is expressed in both cardiac muscle and skeletal muscle.

In embodiments, the promoter is a skeletal muscle-specific promoter, in particular a muscle creatine kinase (MCK) promoter, more particularly the double MCK promoter or
15 triple MCK promoter consisting of a double or triple tandem of MCK enhancer and MCK basal promoter as described in Wang et al. (2008. Gene Ther. 15:1489-1499).

Furthermore, the promoter does not need to be the promoter of the transgene in the nucleic acid expression cassette, although it is possible that the transgene is transcribed from its own promoter.

20 To minimize the length of the nucleic acid expression cassette, the regulatory elements may be linked to minimal promoters, or shortened versions of the promoters described herein. A 'minimal promoter' (also referred to as basal promoter or core promoter) as used herein is part of a full-size promoter still capable of driving expression, but lacking at least part of the sequence that contributes to regulating (e.g. tissue-specific) expression. This
25 definition covers both promoters from which (tissue-specific) regulatory elements have been deleted- that are capable of driving expression of a gene but have lost their ability to express that gene in a tissue-specific fashion and promoters from which (tissue-specific) regulatory elements have been deleted that are capable of driving (possibly decreased) expression of a gene but have not necessarily lost their ability to express that gene in a
30 tissue-specific fashion. Preferably, the promoter contained in the nucleic acid expression cassette disclosed herein is 1000 nucleotides or less in length, 900 nucleotides or less, 800 nucleotides or less, 700 nucleotides or less, 600 nucleotides or less, 500 nucleotides or less, 400 nucleotides or less, 300 nucleotides or less, or 250 nucleotides or less.

The term 'transgene' as used herein refers to particular nucleic acid sequences encoding a polypeptide or a portion of a polypeptide to be expressed in a cell into which the nucleic acid sequence is introduced. However, it is also possible that transgenes are expressed as RNA, typically to control (e.g. lower) the amount of a particular polypeptide in a cell into which the nucleic acid sequence is inserted. These RNA molecules include but are not limited to molecules that exert their function through RNA interference (shRNA, RNAi), micro-RNA regulation (miR) (which can be used to control expression of specific genes), catalytic RNA, antisense RNA, RNA aptamers, etc. How the nucleic acid sequence is introduced into a cell is not essential to the invention, it may for instance be through integration in the genome or as an episomal plasmid. Of note, expression of the transgene may be restricted to a subset of the cells into which the nucleic acid sequence is introduced. The term 'transgene' is meant to include (1) a nucleic acid sequence that is not naturally found in the cell (i.e., a heterologous nucleic acid sequence); (2) a nucleic acid sequence that is a mutant form of a nucleic acid sequence naturally found in the cell into which it has been introduced; (3) a nucleic acid sequence that serves to add additional copies of the same (i.e., homologous) or a similar nucleic acid sequence naturally occurring in the cell into which it has been introduced ; or (4) a silent naturally occurring or homologous nucleic acid sequence whose expression is induced in the cell into which it has been introduced.

The transgene may be homologous or heterologous to the promoter (and/or to the animal, in particular mammal, in which it is introduced, e.g. in cases where the nucleic acid expression cassette is used for gene therapy).

The transgene may be a full length cDNA or genomic DNA sequence, or any fragment, subunit or mutant thereof that has at least some biological activity. In particular, the transgene may be a minigene, i.e. a gene sequence lacking part, most or all of its intronic sequences. The transgene thus optionally may contain intron sequences. Optionally, the transgene may be a hybrid nucleic acid sequence, i.e., one constructed from homologous and/or heterologous cDNA and/or genomic DNA fragments. By 'mutant form' is meant a nucleic acid sequence that contains one or more nucleotides that are different from the wild-type or naturally occurring sequence, i.e., the mutant nucleic acid sequence contains one or more nucleotide substitutions, deletions, and/or insertions. The nucleotide substitution, deletion, and/or insertion can give rise to a gene product (i.e. e., protein or nucleic acid) that is different in its amino acid/nucleic acid sequence from the wild type amino acid/nucleic acid sequence. Preparation of such mutants is well known in the art. In

some cases, the transgene may also include a sequence encoding a leader peptide or signal sequence such that the transgene product will be secreted from the cell.

The transgene that may be contained in the nucleic acid expression cassettes described herein typically encodes a gene product such as RNA or a polypeptide (protein).

- 5 In embodiments, the transgene encodes a therapeutic protein. The therapeutic protein may be a secretable protein. Non-limiting examples of secretable proteins, in particular secretable therapeutic proteins, include clotting factors, such as factor VIII or factor IX, insulin, erythropoietin, lipoprotein lipase, antibodies or nanobodies, growth factors, cytokines, chemokines, plasma factors, etc. The therapeutic protein may also be a structural protein. Non-limiting examples of structural proteins, in particular structural therapeutic proteins, include dystrophin and sarcoglycans. In embodiments, the transgene comprises microdystrophin 1 (*MD1*) gene or follistatin (*FST*) gene.
- 10

In embodiments, the transgene encodes an immunogenic protein. Non-limiting examples of immunogenic proteins include epitopes and antigens derived from a pathogen.

- 15 As used herein, the term "immunogenic" refers to a substance or composition capable of eliciting an immune response.

Other sequences may be incorporated in the nucleic acid expression cassette disclosed herein as well, typically to further increase or stabilize the expression of the transgene product (e.g. introns and/or polyadenylation sequences).

- 20 Any intron can be utilized in the expression cassettes described herein. The term "intron" encompasses any portion of a whole intron that is large enough to be recognized and spliced by the nuclear splicing apparatus. Typically, short, functional, intron sequences are preferred in order to keep the size of the expression cassette as small as possible which facilitates the construction and manipulation of the expression cassette. In some
- 25 embodiments, the intron is obtained from a gene that encodes the protein that is encoded by the coding sequence within the expression cassette. The intron can be located 5' to the coding sequence, 3' to the coding sequence, or within the coding sequence. An advantage of locating the intron 5' to the coding sequence is to minimize the chance of the intron interfering with the function of the polyadenylation signal. In embodiments, the
- 30 nucleic acid expression cassette disclosed herein further comprises an intron. Non-limiting examples of suitable introns are Minute Virus of Mice (MVM) intron, beta-globin intron (betaIVS-II), factor IX (FIX) intron A, Simian virus 40 (SV40) small-t intron, and beta-actin intron.

Preferably, the intron is MVM intron.

Any polyadenylation signal that directs the synthesis of a polyA tail is useful in the expression cassettes described herein, examples of those are well known to one of skill in the art. Exemplary polyadenylation signals include, but are not limited to, polyA sequences derived from the Simian virus 40 (SV40) late gene, the bovine growth hormone (BGH) polyadenylation signal, the minimal rabbit β -globin (mRBG) gene, and the synthetic polyA s(SPA) site as described in Levitt et al. (1989, Genes Dev 3:1019-1025) (SEQ ID NO:46).

Preferably, the polyadenylation signal is derived from SV40 (i.e. SV40 pA).

In particular embodiments, the invention provides a nucleic acid expression cassette comprising a nucleic acid regulatory element consisting of SEQ ID NO:10 or a sequence having 95% identity to said sequence, operably linked to a promoter, preferably a promoter selected from the group consisting of the promoter from the desmin gene or the SPc5-12 promoter, and a transgene, preferably a transgene encoding a luciferase. In further embodiments, the nucleic acid expression cassette further comprises an MVM intron. In yet further embodiments the nucleic acid expression cassette further comprises a polyadenylation signal, preferably a polyadenylation signal derived from SV40.

In particular embodiments, the invention provides a nucleic acid expression cassette comprising a nucleic acid regulatory element consisting of SEQ ID NO:10 or a sequence having 95% identity to said sequence, operably linked to a promoter, preferably the promoter from the desmin gene, and a transgene, preferably a transgene encoding microdystrophin 1 or follistatin. In further embodiments, the nucleic acid expression cassette further comprises an MVM intron. In yet further embodiments, the nucleic acid expression cassette further comprises a polyadenylation signal, preferably wherein the polyadenylation signal has SEQ ID NO:46.

The nucleic acid regulatory element and the nucleic acid expression cassette disclosed herein may be used as such, or typically, they may be part of a nucleic acid vector. Accordingly, a further aspect relates to the use of a nucleic acid regulatory element as described herein or a nucleic acid expression cassette as described herein in a vector, in particular a nucleic acid vector.

In an aspect, the invention also provides a vector comprising a nucleic acid regulatory element as disclosed herein. In further embodiments, the vector comprises a nucleic acid expression cassette as disclosed herein.

The term 'vector' as used in the application refers to nucleic acid molecules, e.g. double-stranded DNA, which may have inserted into it another nucleic acid molecule (the insert nucleic acid molecule) such as, but not limited to, a cDNA molecule. The vector is used to

transport the insert nucleic acid molecule into a suitable host cell. A vector may contain the necessary elements that permit transcribing the insert nucleic acid molecule, and, optionally, translating the transcript into a polypeptide. The insert nucleic acid molecule may be derived from the host cell, or may be derived from a different cell or organism.

5 Once in the host cell, the vector can replicate independently of, or coincidental with, the host chromosomal DNA, and several copies of the vector and its inserted nucleic acid molecule may be generated. The vectors can be episomal vectors (i.e., that do not integrate into the genome of a host cell), or can be vectors that integrate into the host cell genome. The term 'vector' may thus also be defined as a gene delivery vehicle that

10 facilitates gene transfer into a target cell. This definition includes both non-viral and viral vectors. Non-viral vectors include but are not limited to cationic lipids, liposomes, nanoparticles, PEG, PEI, plasmid vectors (e.g. pUC vectors, bluescript vectors (pBS) and pBR322 or derivatives thereof that are devoid of bacterial sequences (minicircles)) transposons-based vectors (e.g. PiggyBac (PB) vectors or Sleeping Beauty (SB) vectors),

15 etc. Viral vectors are derived from viruses and include but are not limited to retroviral, lentiviral, adeno-associated viral, adenoviral, herpes viral, hepatitis viral vectors or the like. Typically, but not necessarily, viral vectors are replication-deficient as they have lost the ability to propagate in a given cell since viral genes essential for replication have been eliminated from the viral vector. However, some viral vectors can also be adapted to

20 replicate specifically in a given cell, such as e.g. a cancer cell, and are typically used to trigger the (cancer) cell-specific (onco)lysis. Virosomes are a non-limiting example of a vector that comprises both viral and non-viral elements, in particular they combine liposomes with an inactivated HIV or influenza virus (Yamada et al., 2003). Another example encompasses viral vectors mixed with cationic lipids.

25 In preferred embodiments, the vector is a viral vector, such as a retroviral, lentiviral, adenoviral, or adeno-associated viral (AAV) vector, more preferably an AAV vector. AAV vectors are preferably used as self-complementary, double-stranded AAV vectors (scAAV) in order to overcome one of the limiting steps in AAV transduction (i.e. single-stranded to double-stranded AAV conversion) (McCarty, 2001, 2003; Nathwani et al, 2002, 2006,

30 2011; Wu et al., 2008), although the use of single-stranded AAV vectors (ssAAV) are also encompassed herein.

AAV serotype 9 (AAV9) is ideally suited to achieve efficient transduction in heart and skeletal muscle. Accordingly, in particularly preferred embodiments, the vector is an AAV9 vector, more particularly a self-complementary AAV9 vector (scAAV9).

In other embodiments, the vector is a non-viral vector, preferably a plasmid, a minicircle, or a transposon-based vector, such as a Sleeping Beauty(SB)-based vector or piggyBac(PB)-based vector.

In yet other embodiments, the vector comprises viral and non-viral elements.

- 5 In particular embodiments, the invention provides a vector comprising a nucleic acid expression cassette comprising a nucleic acid regulatory element consisting of SEQ ID NO:10, a promoter, preferably the promoter from the desmin gene, an MVM intron, a transgene, preferably a transgene encoding microdystrophin 1, and a polyadenylation signal, preferably the polyadenylation signal having SEQ ID NO:46. In
10 particular embodiments, said vector has SEQ ID NO: 44.

- In particular embodiments, the invention provides a vector comprising a nucleic acid expression cassette comprising a nucleic acid regulatory element consisting of SEQ ID NO:10, a promoter, preferably the promoter from the desmin gene, an MVM intron, a transgene, preferably a transgene encoding follistatin, and a polyadenylation signal,
15 preferably the polyadenylation signal having SEQ ID NO:46. In particular embodiments, said vector has SEQ ID NO: 45.

- The nucleic acid expression cassettes and vectors disclosed herein may be used, for example, to express proteins that are normally expressed and utilized in muscle (i.e. structural proteins), or to express proteins that are expressed in muscle and that are then
20 exported to the blood stream for transport to other portions of the body (i.e. secretable proteins). For example, the expression cassettes and vectors disclosed herein may be used to express a therapeutic amount of a gene product (such as a polypeptide, in particular a therapeutic protein, or RNA) for therapeutic purposes, in particular for gene therapy. Typically, the gene product is encoded by the transgene within the expression
25 cassette or vector, although in principle it is also possible to increase expression of an endogenous gene for therapeutic purposes. In an alternative example, the expression cassettes and vectors disclosed herein may be used to express an immunological amount of a gene product (such as a polypeptide, in particular an immunogenic protein, or RNA) for vaccination purposes.

- 30 The nucleic acid expression cassettes and vectors as taught herein may be formulated in a pharmaceutical composition with a pharmaceutically acceptable excipient, i.e., one or more pharmaceutically acceptable carrier substances and/or additives, e.g., buffers, carriers, excipients, stabilisers, etc. The pharmaceutical composition may be provided in the form of a kit.

The term “pharmaceutically acceptable” as used herein is consistent with the art and means compatible with the other ingredients of the pharmaceutical composition and not deleterious to the recipient thereof.

Accordingly, a further aspect of the invention relates to a pharmaceutical composition comprising a nucleic acid expression cassette or a vector described herein.

The use of nucleic acid regulatory elements described herein for the manufacture of these pharmaceutical compositions is also disclosed herein.

In embodiments, the pharmaceutical composition may be a vaccine. The vaccine may further comprise one or more adjuvants for enhancing the immune response. Suitable adjuvants include, for example, but without limitation, saponin, mineral gels such as aluminium hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, bacilli Calmette-Guerin (BCG), *Corynebacterium parvum*, and the synthetic adjuvant QS-21. Optionally, the vaccine may further comprise one or more immunostimulatory molecules. Non-limiting examples of immunostimulatory molecules include various cytokines, lymphokines and chemokines with immunostimulatory, immunopotentiating, and pro-inflammatory activities, such as interleukins (e.g., IL-1, IL-2, IL-3, IL-4, IL-12, IL-13); growth factors (e.g., granulocyte-macrophage (GM)-colony stimulating factor (CSF)); and other immunostimulatory molecules, such as macrophage inflammatory factor, Flt3 ligand, B7.1; B7.2, etc.

In a further aspect, the invention relates to the nucleic acid regulatory elements, the nucleic acid expression cassettes, the vectors, or the pharmaceutical compositions described herein for use in medicine.

As used herein, the terms “treat” or “treatment” refer to both therapeutic treatment and prophylactic or preventative measures. Beneficial or desired clinical results include, but are not limited to, prevention of an undesired clinical state or disorder, reducing the incidence of a disorder, alleviation of symptoms associated with a disorder, diminishment of extent of a disorder, stabilized (i.e., not worsening) state of a disorder, delay or slowing of progression of a disorder, amelioration or palliation of the state of a disorder, remission (whether partial or total), whether detectable or undetectable, or combinations thereof. “Treatment” can also mean prolonging survival as compared to expected survival if not receiving treatment.

As used herein, the terms “therapeutic treatment” or “therapy” and the like, refer to treatments wherein the object is to bring a subjects body or an element thereof from an undesired physiological change or disorder to a desired state, such as a less severe or

unpleasant state (e.g., amelioration or palliation), or back to its normal, healthy state (e.g., restoring the health, the physical integrity and the physical well-being of a subject), to keep it at said undesired physiological change or disorder (e.g., stabilization, or not worsening), or to prevent or slow down progression to a more severe or worse state compared to said undesired physiological change or disorder.

As used herein the terms "prevention", "preventive treatment" or "prophylactic treatment" and the like encompass preventing the onset of a disease or disorder, including reducing the severity of a disease or disorder or symptoms associated therewith prior to affliction with said disease or disorder. Such prevention or reduction prior to affliction refers to administration of the nucleic acid regulatory elements, the nucleic acid expression cassettes, the vectors, or the pharmaceutical compositions described herein to a patient that is not at the time of administration afflicted with clear symptoms of the disease or disorder. "Preventing" also encompasses preventing the recurrence or relapse-prevention of a disease or disorder for instance after a period of improvement. In embodiments, the nucleic acid regulatory elements, the nucleic acid expression cassettes, the vectors, or the pharmaceutical compositions described herein may be for use in gene therapy, in particular muscle-directed gene therapy, more particularly heart and skeletal muscle-directed gene therapy or skeletal muscle-directed gene therapy.

Also disclosed herein is the use of the nucleic acid regulatory elements, the nucleic acid expression cassettes, the vectors, or the pharmaceutical compositions described herein for the manufacture of a medicament for gene therapy, in particular muscle-directed gene therapy, more particularly heart and skeletal muscle-directed gene therapy or skeletal muscle-directed gene therapy.

Also disclosed herein is a method for gene therapy, in particular muscle-directed gene therapy, more particularly heart and skeletal muscle-directed gene therapy or skeletal muscle-directed gene therapy, in a subject in need of said gene therapy comprising:

- introducing in the subject, in particular in muscle of the subject, more particularly in heart muscle or skeletal muscle of the subject, a nucleic acid expression cassette, a vector or a pharmaceutical composition described herein, wherein the nucleic acid expression cassette, the vector or the pharmaceutical composition comprises a nucleic acid regulatory element described herein operably linked to a promoter and a transgene; and

- expressing a therapeutically effective amount of the transgene product in the subject, in particular in muscle of the subject, more particularly in heart and skeletal muscle or in skeletal muscle of the subject.

The transgene product may be a polypeptide, in particular a structural protein such as, e.g., dystrophin or a sarcoglycan, or a secretable protein such as, e.g., a clotting factor, e.g., factor IX or factor VIII, a cytokine, a growth factor, an antibody or nanobody, a chemokine, a plasma factor, insulin, erythropoietin, lipoprotein lipase. In particular embodiments, the transgene product is follistatin or microdystrophin, in particular microdystrophin 1. Alternatively, the transgene product may be RNA, such as siRNA.

- Exemplary diseases and disorders that may benefit from gene therapy using the nucleic acid regulatory elements, the nucleic acid expression cassettes, the vectors, or the pharmaceutical compositions described herein include muscular dystrophy (e.g. Duchenne muscular dystrophy (DMD) / Becker muscular dystrophy (BMD)), myotonic dystrophy, neuromuscular disease, motor neuron diseases (MND), such as e.g. Charcot-Marie-Tooth disease (CMT), spinal muscular atrophy (SMA), and amyotrophic lateral sclerosis (ALS), Emery-Dreifuss muscular dystrophy, facioscapulohumeral muscular dystrophy (FSHD), congenital muscular dystrophies, congenital myopathies, limb girdle muscular dystrophy, metabolic myopathies, muscle inflammatory diseases, myasthenia, mitochondrial myopathies, anomalies of ionic channels, nuclear envelop diseases, cardiomyopathies, cardiac hypertrophy, heart failure, distal myopathies, cardiovascular diseases, hemophilia, including hemophilia A and B, and diabetes.

Gene therapy protocols have been extensively described in the art. These include, but are not limited to, intramuscular injection of plasmid (naked or in liposomes), hydrodynamic gene delivery in various tissues, including muscle, interstitial injection, instillation in airways, application to endothelium, intra-hepatic parenchyme, and intravenous or intra-arterial administration. Various devices have been developed for enhancing the availability of DNA to the target cell. A simple approach is to contact the target cell physically with catheters or implantable materials containing DNA. Another approach is to utilize needle-free, jet injection devices which project a column of liquid directly into the target tissue under high pressure. These delivery paradigms can also be used to deliver vectors. Another approach to targeted gene delivery is the use of molecular conjugates, which consist of protein or synthetic ligands to which a nucleic acid-or DNA-binding agent has been attached for the specific targeting of nucleic acids to cells (Cristiano et al., 1993). In embodiments, the nucleic acid regulatory elements, the nucleic acid expression cassettes,

the vectors, or the pharmaceutical compositions described herein may be for use as a vaccine, more particularly for use as a prophylactic vaccine.

Also disclosed herein is the use of the nucleic acid regulatory elements, the nucleic acid expression cassettes, the vectors, or the pharmaceutical compositions described herein
5 for the manufacture of a vaccine, in particular for the manufacture of a prophylactic vaccine.

Also disclosed herein is a method of vaccination, in particular prophylactic vaccination, of a subject in need of said vaccination comprising:

- 10 - introducing in the subject, in particular in muscle of the subject, more particularly in heart muscle or skeletal muscle of the subject, a nucleic acid expression cassette, a vector or a pharmaceutical composition described herein, wherein the nucleic acid expression cassette, the vector or the pharmaceutical composition comprises a nucleic acid regulatory element described herein operably linked to a promoter and a transgene; and
- 15 - expressing an immunologically effective amount of the transgene product in the subject, in particular in muscle of the subject, more particularly in heart and skeletal muscle or in skeletal muscle of the subject.

As used herein, a phrase such as "a subject in need of treatment" includes subjects that would benefit from treatment of a recited disease or disorder. Such subjects may include,
20 without limitation, those that have been diagnosed with said disease or disorder, those prone to contract or develop said disease or disorder and/or those in whom said disease or disorder is to be prevented.

The terms "subject" and "patient" are used interchangeably herein and refer to animals, preferably vertebrates, more preferably mammals, and specifically include human patients
25 and non-human mammals. "Mammalian" subjects include, but are not limited to, humans, domestic animals, commercial animals, farm animals, zoo animals, sport animals, pet and experimental animals such as dogs, cats, guinea pigs, rabbits, rats, mice, horses, cattle, cows; primates such as apes, monkeys, orang-utans, and chimpanzees; canids such as dogs and wolves; felids such as cats, lions, and tigers; equids such as horses, donkeys,
30 and zebras; food animals such as cows, pigs, and sheep; ungulates such as deer and giraffes; rodents such as mice, rats, hamsters and guinea pigs; and so on. Preferred patients or subjects are human subjects.

A 'therapeutic amount' or 'therapeutically effective amount' as used herein refers to the amount of gene product effective to treat a disease or disorder in a subject, i.e., to obtain

a desired local or systemic effect. The term thus refers to the quantity of gene product that elicits the biological or medicinal response in a tissue, system, animal, or human that is being sought by a researcher, veterinarian, medical doctor or other clinician. Such amount will typically depend on the gene product and the severity of the disease, but can be decided by the skilled person, possibly through routine experimentation.

An "immunologically effective amount" as used herein refers to the amount of (trans)gene product effective to enhance the immune response of a subject against a subsequent exposure to the immunogen encoded by the (trans)gene. Levels of induced immunity can be determined, e.g. by measuring amounts of neutralizing secretory and/or serum antibodies, e.g., by plaque neutralization, complement fixation, enzyme-linked immunosorbent, or microneutralization assay.

Typically, the amount of (trans)gene product expressed when using an expression cassette or vector as described herein (i.e., with at least one muscle-specific nucleic acid regulatory element) are higher than when an identical expression cassette or vector is used but without a nucleic acid regulatory element therein. More particularly, the expression is at least double as high, at least five times as high, at least ten times as high, at least 20 times as high, at least 30 times as high, at least 40 times as high, at least 50 times as high, or even at least 60 times as high as when compared to the same nucleic acid expression cassette or vector without nucleic acid regulatory element. Moreover, the higher expression remains specific to muscle, in particular both heart and skeletal muscle or skeletal muscle alone. Furthermore, the expression cassettes and vectors described herein direct the expression of a therapeutic amount of the gene product for an extended period. Typically, therapeutic expression is envisaged to last at least 20 days, at least 50 days, at least 100 days, at least 200 days, and in some instances 300 days or more. Expression of the gene product (e.g. polypeptide) can be measured by any art-recognized means, such as by antibody-based assays, e.g. a Western Blot or an ELISA assay, for instance to evaluate whether therapeutic expression of the gene product is achieved. Expression of the gene product may also be measured in a bioassay that detects an enzymatic or biological activity of the gene product. Also disclosed herein is the use of the nucleic acid regulatory elements, the nucleic acid expression cassettes, or the vectors disclosed herein for transfecting or transducing muscle cells, preferably heart muscle cells and/or skeletal muscle cells.

Further disclosed herein is the use of the nucleic acid expression cassettes or the vectors disclosed herein for expressing a transgene product in muscle cells, preferably heart muscle cells and/or skeletal muscle cells, wherein the nucleic acid expression cassette or

the vector comprises a nucleic acid regulatory element disclosed herein operably linked to a promoter and a transgene.

Further disclosed herein is a method for expressing a transgene product in muscle cells, preferably heart muscle cells and/or skeletal muscle cells, comprising:

- 5 - transfecting or transducing the muscle cells with a nucleic acid expression cassette or a vector disclosed herein, wherein the nucleic acid expression cassette or the vector comprises a nucleic acid regulatory element disclosed herein operably linked to a promoter and a transgene; and
 - expressing the transgene product in the muscle cells.
- 10 Non-viral transfection or viral vector-mediated transduction of muscle cells may be performed by *in vitro*, *ex vivo* or *in vivo* procedures. The *in vitro* approach requires the *in vitro* transfection or transduction of muscle cells, e.g. muscle cells previously harvested from a subject, muscle cell lines or muscle cells differentiated from e.g. induced pluripotent stem cells or embryonic cells. The *ex vivo* approach requires harvesting of the
- 15 muscle cells from a subject, *in vitro* transfection or transduction, and optionally re-introduction of the transfected muscle cells into the subject. The *in vivo* approach requires the administration of the nucleic acid expression cassette or the vector disclosed herein into a subject. In preferred embodiments, the transfection of the muscle cells is performed *in vitro* or *ex vivo*.
- 20 It is understood by the skilled person that the use of the nucleic acid regulatory elements, the nucleic acid expression cassettes and vectors disclosed herein has implications beyond gene therapy, e.g. coaxed differentiation of stem cells into myogenic cells, transgenic models for over-expression of proteins in muscle, etc.

The invention is further explained by the following non-limiting examples

25 **EXAMPLES**

Example 1: Identification of cardiac and skeletal muscle-specific regulatory elements

Experimental procedures

- 30 Genes highly expressed both in heart and muscle, but showing only minimal expression in other tissues, were identified using the Specificity Measure (SPM) method described in Xiao et al. (2010. Bioinformatics 26:1273-1275). As data set we used the U133A/GNF1H Gene Atlas (GSE1133) of the human protein-encoding transcriptomes (Su et al. 2004. Proc. Natl. Acad. Sci. USA 101:6062-6067). This resulted in a short list of 5 genes:

- filamin-c gene (*FLNC*), actinin, alpha 2 gene (*ACTN2*), myosin regulatory light chain 2, ventricular/cardiac muscle isoform gene (*MYL2*), desmin gene (*DES*) and telethonin gene (*TCAP*). Next, the genomic context of these genes was searched for cross-species conserved regions enriched for transcription factor binding sites (TFBS) associated with
- 5 high expression in muscle and heart. Additional filtering was done by selecting the putative nucleic acid regulatory elements that either overlapped or contained regions bearing reproducible biochemical features associated with transcription regulation and/or a open chromatin structure as defined in the ENCODE project (The ENCODE Project Consortium. 2012. Nature 489:57-74).
- 10 For the heart-specific regulatory elements we used a high-density gene expression database of 18,927 unique genes based on microarrays that were derived from 158 normal human samples from 19 different organs of 30 different individuals (Son, S. et al. 2005. Database of mRNA gene expression profiles of multiple human organs. Genome Res. 15:443–450). This was used to identify a set of most highly expressed (i.e. 'over-
- 15 expressed') genes in the heart compared to any of the other tissues. A two-tailed t-test was used for each pairwise comparison. Conversely, a set of 'under-expressed' genes was identified, corresponding to those genes that exhibited the lowest expression in these respective organs compared to any of the other tissues. This analysis resulted in a set of 43 over-expressed genes and a collection of 37 under-expressed heart-specific genes.
- 20 Next, the Reference Sequence (RefSeq) identifiers (IDs) lists of these 'over-expressed' and 'under-expressed' heart-specific genes were used to extract the corresponding promoter sequences upstream the reported transcription start sites (TSSs) by 1000 bases (NCBI36/hg18 genome assembly), using the transcription start location data stored in the refGene table of the UCSC Genome Browser database.
- 25 This resulted in two sets of heart-specific promoter sequences corresponding to promoters of 'over-expressed' or 'under-expressed' genes. In order to make a non-redundant set of representative promoter sequences, the promoter sequences were filtered using 'uclust'.
- The two sets of non-redundant tissue-specific promoter sequences corresponding to
- 30 promoters of 'over-expressed' or 'under-expressed' genes were used as input for the DDM/MDS method, described in detail in De Bleser et al. (2007. Genome Biol 8:R83). The observed differential behaviour might be explained by the presence of one or more TFBS elements characteristic of the promoters of the up-regulated or down-regulated group of genes. These 'differential' TFBS elements could be found using following
- 35 procedure. First, a library of TFBS positional

weight matrices (PWMs) (TRANSFAC® 2010.3) was used to predict TFBS on every promoter sequence. For the muscle-specific promoters we used the Find Individual Motif Occurrences (fimo) application with a P-value cut-off of 10^{-3} . The number of predicted TFBS elements per PWM per promoter was collected in the form of a matrix in which each

5 row corresponds to a promoter sequence, while the columns corresponded to the used PWM. Two TFBS were considered correlated if their corresponding columns in the matrix were similar what could be measured using a distance function. With this approach, distance matrices summarizing all TFBS associations were constructed for the TFBS in both sets of promoters. Finally, by calculating the distance difference matrix (DDM) and

10 performing multidimensional scaling on this DDM to visualize its content in two dimensions, TFBS could be distinguished that did not contribute to the observed differential gene expression as they were mapped near the origin of the DDM-MDS plot from 'deviating' TFBS that are likely responsible for the observed differential gene expression. As the MDS procedure plots TFBS that are strongly associated closer

15 together than less associated ones, it was able to highlight interactions between TFBS in the promoter datasets. This procedure resulted in a list of TFBS associated with high tissue-specific expression for heart-specific promoters.

For the muscle-specific regulatory elements, a list of muscle-specific genes was extracted from the Tissue-specific Gene Expression and Regulation (TiGER) database. This was

20 used to identify a set of the most highly expressed (i.e. 'over-expressed') genes in muscle tissue. Conversely, a set of 'under-expressed' genes was identified, corresponding to those genes that exhibited the lowest expression in muscle. Next, the Reference Sequence (RefSeq) identifiers (IDs) lists of these 'over-expressed' and 'under-expressed' muscle-specific genes were used to extract the corresponding promoter sequences

25 upstream the reported transcription start sites (TSSs) by 1000 bases (NCBI36/hg18 genome assembly), using the transcription start location data stored in the refGene table of the UCSC Genome Browser database. This resulted in two sets of muscle-specific promoter sequences corresponding to promoters of 'over-expressed' or 'under-expressed' genes. In order to make a non-redundant set of

30 representative promoter sequences, the promoter sequences were filtered using 'uclust'.

The two sets of non-redundant tissue-specific promoter sequences corresponding to promoters of 'over-expressed' or 'under-expressed' genes were used as input for the DDM/MDS method, described in detail in De Bleser et al. (2007. *Genome Biol* 8:R83).

35 The observed differential behaviour

might be explained by the presence of one or more TFBS elements characteristic of the promoters of the up-regulated or down-regulated group of genes. These 'differential' TFBS elements could be found using following procedure. First, a library of TFBS positional weight matrices (PWMs) (TRANSFAC® 2010.3) was used to predict TFBS on every promoter sequence. For the muscle-specific promoters we used the Find Individual Motif Occurrences (fimo) application with a P-value cut-off of 10^{-3} . The number of predicted TFBS elements per PWM per promoter was collected in the form of a matrix in which each row corresponds to a promoter sequence, while the columns corresponded to the used PWM. Two TFBS were considered correlated if their corresponding columns in the matrix were similar what could be measured using a distance function. With this approach, distance matrices summarizing all TFBS associations were constructed for the TFBS in both sets of promoters. Finally, by calculating the distance difference matrix (DDM) and performing multidimensional scaling on this DDM to visualize its content in two dimensions, TFBS could be distinguished that did not contribute to the observed differential gene expression as they were mapped near the origin of the DDM-MDS plot from 'deviating' TFBS that are likely responsible for the observed differential gene expression. As the MDS procedure plots TFBS that are strongly associated closer together than less associated ones, it was able to highlight interactions between TFBS in the promoter datasets. This procedure resulted in a list of TFBS associated with high tissue-specific expression for muscle-specific promoters.

Results

This computational approach led to the identification of 6 cardiac and skeletal muscle-specific regulatory sequences, summarized in Table 1.

Table 1: Cardiac and skeletal muscle-specific regulatory elements. Bp: base pairs.

Sequence	Name	Gene regulated by sequence	Size (bp)	Conserved TFBS present
SEQ ID NO: 1	CSk-SH1	Des	381	E2A, HNH1, NF1, CEBP, LRF, MyoD, SREBP
SEQ ID NO: 2	CSk-SH2	Des	435	E2A, NF1, p53, CEBP, LRF, SREBP
SEQ ID NO: 3	CSk-SH3	ACTN2	551	E2A, HNH1, HNF3a, HNF3b, NF1, CEBP, LRF,

				MyoD, SREBP
SEQ ID NO: 4	CSk-SH4	ACTN2	430	E2A, HNF3a, NF1, CEBP, LRF, MyoD, SREBP
SEQ ID NO: 5	CSk-SH5	FLNC	454	HNF4, NF1, RSRFC4, CEBP, LRF, MyoD
SEQ ID NO: 6	CSk-SH6	FLNC	453	NF1, PPAR, p53, CEBP, LRF, MyoD

Example 2: *In vivo* validation of cardiac and skeletal muscle-specific regulatory elements via AAV vectors

Experimental procedures

Generation of the AAV plasmid constructs (pAAV-CSk-SH-Des-Luc2)

- 5 The cardiac and skeletal muscle-specific regulatory elements (CSk-SH1 to CSk-SH6) were synthesized by conventional oligonucleotide synthesis and flanked with Acc65I and MluI restriction sites. The different CSk-SHs were cloned upstream of the Desmin (Des) promoter that drives expression of a Firefly Luciferase (Luc2) reporter gene in the context of an adeno-associated viral vector (AAV) backbone (designated as pAAV-Des-Luc2),
- 10 schematically represented in Figure 3. The corresponding AAV plasmid constructs were designated as pAAV-CSk-SH-Des-Luc2. The plasmids also contained a Minute Virus of Mouse (MVM) intron and a Simian Virus 40 (SV40) polyadenylation site (pA).

AAV vector production (AAV9sc.CSk-SH1-6.Des.Luc2) and purification

- AAV9 vectors were produced by calcium phosphate (Invitrogen Corp, Carlsbad, CA) co-
- 15 transfection of 293T human embryonic kidney cells with the pAAV plasmid of interest, an adenoviral helper plasmid and a chimeric packaging construct that delivers the AAV2 *rep* gene together with the AAV9 *cap* gene, as described in Vandendriessche et al. (2007. J Thromb Haemost 5:16-24).

- Briefly, two days post transfection, cells were harvested and vector particles were purified
- 20 using isopycnic centrifugation methods. Harvested cells were lysed by successive freeze/thaw cycles and sonication, treated with benzonase (Novagen, Madison, WI) and deoxycholic acid (Sigma-Aldrich, St. Louis, MO) and subsequently subjected to 3 successive rounds of cesium chloride (Invitrogen Corp, Carlsbad, CA) density gradient ultracentrifugation. Fractions containing the AAV vector were collected, concentrated in
- 25 1mM MgCl₂ in Dulbecco's phosphate buffered saline (PBS) (Gibco, BRL) and stored at -80°C.

Vector titers (in viral genomes (vg)/ml) were determined by quantitative real-time PCR using SYBR Green mix (which included SYBR Green dye, Taqman polymerase, ROX and dNTP's all in one) and luciferase specific primers on an ABI 7500 Real-Time PCR System (Applied Biosystem, Foster city, CA, USA). The forward and reverse primers used were 5'-
 5 CCCACCGTCGTATTCTGTGAG-3' (SEQ ID NO: 14) and 5'-
 TCAGGGCGATGGTTTTGTCCC-3' (SEQ ID NO: 15), respectively.

Typically, for all vectors titers in the range of $1.5\text{--}6.1 \times 10^{11}$ vg/ml were achieved from a small production batch of 20 petri dishes of producer cells. If higher number of petri dishes such as 60 dishes of producer cells were used, a higher titer typically in the range of $10^{12}\text{--}10^{13}$ gc/ml of AAV particles were achieved. Known copy numbers ($10^2\text{--}10^7$) of the
 10 respective vector plasmids used to generate the corresponding AAV vectors, carrying the appropriate cDNAs were used to generate the standard curves.

Animal studies

All animal procedures were approved by the institutional animal ethics committee of the
 15 Free University of Brussels (VUB) (Brussels, Belgium). All mice were housed under specific pathogen-free conditions; food and water were provided *ad libitum*.

Two-days old CB.17/lcrTac/Prkdcscid mice were intravenously injected into the periorbital vein with 50 μ l of concentrated vectors (5×10^9 vg/mouse) containing the different CSk-SH (i.e. CSk-SH1 to CSk-SH6) regulatory elements or AAV9-Des-Luc2 control vector (5×10^9
 20 vg/mouse) as summarized in Table 2.

Table 2: Experimental design for the injection of cardiac and skeletal muscle-specific regulatory elements. Vector	Mouse number (n)	Dose	Titre (gc/ml)	Volume vector (μ l)	Volume PBS (μ l)	Total volume (μ l)
AAV9sc. Des.Luc2	3	5×10^9	5.7×10^{11}	8.7	41.3	50
AAV9sc.CSk-	4	5×10^9	6.1×10^{11}	8.2	41.8	50

SH1.Des.Luc2						
AAV9sc.CSk-SH2.Des.Luc2	2	5×10^9	6.0×10^{11}	8.3	41.7	50
AAV9sc.CSk-SH3.Des.Luc2	5	5×10^9	1.5×10^{11}	33.3	16.7	50
AAV9sc.CSk-SH4.Des.Luc2	3	5×10^9	5.2×10^{11}	9.6	40.4	50
AAV9sc.CSk-SH5.Des.Luc2	4	5×10^9	1.7×10^{11}	39.4	20.6	50
AAV9sc.CSk-SH6.Des.Luc2	4	5×10^9	3.1×10^{11}	16.1	33.9	50

Mice were imaged between 5 and 8 weeks post-injection once per week using a biospace *In Vivo* photo Imaging System (IVIS). The CCD was cooled to -120°C and the field of view (FOV) set at 25 cm height of the sample shelf. The charged coupled device (CCD) camera operates by converting photons that strike the CCD pixel into electrons at wavelengths between 400-1000 nm, allowing detection of visible imaged infrared light through by anesthetizing with 2% isoflurane and oxygen. D-luciferin substrate was injected intravenously, at a dose of 150 $\mu\text{g/g}$ of body weight. Mice were euthanized 9 weeks post injection and intact organs were harvested and imaged using a biospace *In Vivo* photo Imaging System (IVIS).

mRNA analysis

Total RNA was extracted from different organs of the mice by a silica-membrane based purification kit according to the manufacturer's instructions (Invitrogen Corp, Carlsbad, CA, USA). Subsequently, 50 ng of total RNA from each sample was subjected to reverse transcription (RT) using a cDNA synthesis kit (Invitrogen Corp, Carlsbad, CA, USA). Next, a cDNA amount corresponding to 10 ng of total RNA was amplified by quantitative(q) PCR on an ABI 7700 (Applied Biosystems, Foster City, CA, USA), using 5'-CCCACCGTCGTATTCGTGAG-3' (SEQ ID NO: 14) as a forward primer and 5'-TCAGGGCGATGGTTTTGTCCC-3' (SEQ ID NO: 15) as reverse primer (amplicon 217 bp). The qPCR standards consisted of serially diluted AAV9-CSK-SH-Des-Luc2 plasmids of known quantity. The Luc2 mRNA levels were normalized to mRNA levels of the endogenous murine glyceraldehyde-3-phosphate dehydrogenase (mGAPDH) gene, using

5'-TGTGTCCGTCGTGGATCTGA-3' (SEQ ID NO:31) as forward primer and 5'-GCCTGCTTCACCACCTTCTTGA-3' (SEQ ID NO:32) as the reverse primer (amplicon 82 bp). RNA samples were amplified with and without reverse transcriptase to exclude DNA amplification. The size of the amplified PCR fragments was verified on a 1.8% agarose gel.

Transduction efficiency and vector biodistribution

Transduction efficiency of the viral vectors and biodistribution were evaluated by quantifying Luc2 transgene copy numbers in the different organs and tissues as described previously (Pacak, C.A., et al. 2008. Genet Vaccines Ther 6: 13). Briefly, genomic DNA was extracted from 30 mg of each tissue according to DNeasy Blood & Tissue Kit protocol (Qiagen, Chatsworth, CA, USA) and 100 ng of genomic DNA from each sample was subjected to qPCR, using the Luc2-specific forward primer 5'-CCCACCGTCGTATTCGTGAG-3' (SEQ ID NO: 14) and reverse primer 5'-TCAGGGCGATGGTTTTGTCCC-3' (SEQ ID NO: 15) (amplicon 217 bp). Known copy numbers (10^2 - 10^7) of the corresponding plasmid pAAV-CSk-SH-Des-Luc2 were used to generate the standard curve. The results were expressed as mean AAV copy number/100 ng of genomic DNA.

Results

To assess the effect of the *in silico* identified cardiac and skeletal muscle-specific regulatory elements (CSk-SH) *in vivo*, adeno-associated vectors were generated that expressed the luciferase gene *luc2* from a chimeric promoter. This promoter was composed of the muscle-specific desmin (Des) promoter linked to the cardiac and skeletal muscle-specific regulatory elements CSk-SH1-6. The vectors were intravenously injected in mice and whole body images were taken from the mice at 5 and 6 weeks post-injection to examine luciferase expression level.

All mice injected with a vector comprising a cardiac and skeletal muscle-specific regulatory element (CSk-SH1-6) showed increased luciferase activity compared to control mice that were injected with a corresponding AAV9 vector without a regulatory element, indicating that all of the regulatory elements tested increased luciferase expression (data not shown). We observed very robust and enhanced luciferase activity in mice that were injected with AAV9 vector comprising the regulatory element CSk-SH1, CSk-SH3 or CSk-SH5 at 5 and 6 weeks post-injection as compared to luciferase activity of the control mice.

We also determined luciferase activity in the individual organs. Nine weeks post injection, the mice were euthanized and individual organs were analyzed to evaluate luciferase expression and the biodistribution pattern. We observed significantly increased levels of luciferase activity in the heart, the biceps, the triceps, the quadriceps, the tibialis interior, and the gastrocnemius muscle of mice injected with AAV vector comprising a regulatory element CSk-SH1-6 as compared to control mice. Up to 48-fold augmentation of luciferase activity was measured in the heart of mice injected with AAV vector comprising the CSk-SH5 regulatory element as compared to control mice (Fig. 4A). This regulatory element was also the most robust in muscle, in particular gastrocnemius muscle, tibialis interior, quadriceps, biceps and triceps. Hence, the most potent regulatory element was CSk-SH5. The second most robust regulatory element was CSk-SH1. A 25-fold augmentation of luciferase activity was measured in the heart of mice injected with AAV vector comprising CSk-SH1 when compared to control mice. Robust luciferase activity was also measured in other muscle of mice injected with CSk-SH1. The third robust regulatory element was CSk-SH3, followed by CSk-SH6, CSk-SH4 and CSk-SH2. Furthermore the expression was specific for cardiac and skeletal muscle as luciferase expression was absent or limited in all other organs (Fig. 5A), despite transduction of the vector into the other organs (Fig. 6A-B). The level of luciferase activity measured in the organs was consistent with the luciferase mRNA levels measured in the respective organs (Fig. 4B, 5B). These *in vivo* data show that the nucleic acid regulatory elements CSk-SH, in particular CSk-SH1, CSk-SH3 and CSk-SH5, more particularly CSk-SH1 and CSk-SH5, can enhance heart and skeletal muscle-specific luciferase expression.

Example 3: Identification of skeletal muscle-specific regulatory elements

Experimental procedures

First, a list of muscle-specific genes was extracted from the Tissue-specific Gene Expression and Regulation (TiGER) database. This was used to identify a set of the most highly expressed (i.e. 'over-expressed') genes in muscle tissue. Conversely, a set of 'under-expressed' genes was identified, corresponding to those genes that exhibited the lowest expression in muscle. Next, the Reference Sequence (RefSeq) identifiers (IDs) lists of these 'over-expressed' and 'under-expressed' muscle-specific genes were used to extract the corresponding promoter sequences upstream the reported transcription start sites (TSSs) by 1000 bases (NCBI36/hg18 genome assembly), using the transcription start location data stored in the refGene table of the UCSC Genome Browser database. This resulted in two sets of muscle-specific promoter sequences corresponding to promoters of 'over-expressed' or 'under-expressed' genes.

In order to make a non-redundant set of representative promoter sequences, the promoter sequences were filtered using 'uclust'.

The two sets of non-redundant tissue-specific promoter sequences corresponding to promoters of 'over-expressed' or 'under-expressed' genes were used as input for the DDM/MDS method, described in detail in De Bleser et al. (2007. *Genome Biol* 8:R83). The observed differential behaviour might be explained by the presence of one or more TFBS elements characteristic of the promoters of the up-regulated or down-regulated group of genes. These 'differential' TFBS elements could be found using following procedure. First, a library of TFBS positional weight matrices (PWMs) (TRANSFAC® 2010.3) was used to predict TFBS on every promoter sequence. For the muscle-specific promoters we used the Find Individual Motif Occurrences (fimo) application with a P-value cut-off of 10^{-3} . The number of predicted TFBS elements per PWM per promoter was collected in the form of a matrix in which each row corresponds to a promoter sequence, while the columns corresponded to the used PWM. Two TFBS were considered correlated if their corresponding columns in the matrix were similar what could be measured using a distance function. With this approach, distance matrices summarizing all TFBS associations were constructed for the TFBS in both sets of promoters. Finally, by calculating the distance difference matrix (DDM) and performing multidimensional scaling on this DDM to visualize its content in two dimensions, TFBS could be distinguished that did not contribute to the observed differential gene expression as they were mapped near the origin of the DDM-MDS plot from 'deviating' TFBS that are likely responsible for the observed differential gene expression. As the MDS procedure plots TFBS that are strongly associated closer together than less associated ones, it was able to highlight interactions between TFBS in the promoter datasets. This procedure resulted in a list of TFBS associated with high tissue-specific expression for muscle-specific promoters.

Next, the genomic context of the tissue-specific over-expressed genes was searched for cross-species conserved regions for the TFBS associated with high tissue-specific gene expression. For that purpose, the sequences of all conserved sequence elements in the NCBI36/hg18 genome assembly were downloaded based on the information stored in the phastConsElements44way table of the UCSC Genome Browser database.

The predicted conserved sequence elements are assigned a log-odds score equal to its log probability under the conserved model minus its log probability under the non-conserved model. This allows to restrict the search for putative regulatory elements that coincide with the most conserved sequence elements. The conserved sequence

elements were scanned for TFBS associated with high tissue-specific expression using the fimo application, as described above. Using internally developed Perl scripts, this led to the identification of highly conserved sequence elements containing clusters of TFBS associated with high tissue-specific expression (i.e. nucleic acid regulatory elements).

- 5 Additional filtering was done by selecting the putative nucleic acid regulatory elements that either overlapped or contained regions bearing reproducible biochemical features associated with transcription regulation and/or a open chromatin structure as defined in the ENCODE project (The ENCODE Project Consortium. 2012. Nature 489:57-74).

Results

- 10 This computational approach led to the identification of 6 muscle-specific regulatory sequences, summarized in Table 3.

Table 3: Muscle-specific regulatory elements. Bp: base pairs.

Sequence	Name	Gene regulated by sequence	Size (bp)	Conserved TFBS present
SEQ ID NO: 7	Sk-SH1	ATP2A1	495	E2A, NF1, SRFC, p53, CEBP, LRF, MyoD
SEQ ID NO: 8	Sk-SH2	TNNI1	344	E2A, NF1, CEBP, LRF, MyoD, SREBP
SEQ ID NO: 9	Sk-SH3	TNNI1	430	E2A, HNF3a, CEBP, LRF, MyoD, SREBP, Tal1_b
SEQ ID NO: 10	Sk-SH4	MYLPP	435	E2A, SRF, p53, CEBP, LRF, MyoD, SREBP
SEQ ID NO: 11	Sk-SH5	MYH1	474	HNF4, NF1, RSRFC4, CEBP, LRF, SREBP
SEQ ID NO: 12	Sk-SH6	TPM3	519	E2A, HNF3a, HNF3b, NF1, SRF, CEBP, LRF, MyoD, SREBP
SEQ ID NO: 13	Sk-SH7	ANKRD2	372	E2A, CEBP, MyoD

Example 4: *In vivo* validation of skeletal muscle-specific regulatory elements via AAV vectors comprising a desmin promoter

- 15 Experimental procedures

- AAV plasmid constructs comprising a skeletal muscle-specific regulatory element (pAAV-Sk-SH-Des-Luc2) were generated according to the protocol described in Example 2. Briefly, the skeletal muscle-specific regulatory elements were synthesized by conventional oligonucleotide synthesis and flanked with Acc65I and MluI restriction sites (Sk-SH1, 2, 3, 4, 5 and 7) or BsiWI and MluI restriction sites (Sk-SH6). The different Sk-SHs were cloned upstream of the Desmin (Des) promoter that drives expression of a Firefly Luciferase (Luc2) reporter gene in the context of the AAV vector backbone pAAV-Des-Luc2. The plasmids also contained a Minute Virus of Mouse (MVM) intron and a Simian Virus 40 (SV40) polyadenylation site (pA).
- AAV vector production and purification and animal studies were carried out as described in Example 2. The experimental design is summarized in Table 4.

Table 4: Experimental design for the injection of skeletal muscle-specific regulatory elements.

Vector	Mouse number (n)	Dose	Titre (gc/ml)	Volume vector (μl)	Volume PBS (μl)	Total volume (μl)
AAV9sc. Des.Luc2	5	5×10^9	5.7×10^{11}	8.7	41.3	50
AAV9sc. Sk-SH1.Des.Luc2	4	5×10^9	1.7×10^{11}	29.4	20.6	50
AAV9sc. Sk-SH2.Des.Luc2	4	5×10^9	1.8×10^{11}	27.7	22.3	50
AAV9sc. Sk-SH3.Des.Luc2	4	5×10^9	1.9×10^{11}	26.3	23.7	50
AAV9sc. Sk-SH4.Des.Luc2	4	5×10^9	1.6×10^{11}	31.3	18.7	50
AAV9sc. Sk-SH5.Des.Luc2	1	5×10^9	8.1×10^{11}	62	3	65
AAV9sc. Sk-SH6.Des.Luc2	3	5×10^9	1.4×10^{11}	35.7	14.3	50
AAV9sc. Sk-SH7.Des.Luc2	4	5×10^9	1.8×10^{11}	27.7	22.3	50

mRNA analysis and transduction efficiency and vector biodistribution were assessed as described in Example 2.

Results

Adeno-associated vectors were generated that expressed the luciferase gene *luc2* from a chimeric promoter composed of the desmin (Des) promoter linked to the skeletal muscle-specific regulatory elements Sk-SH1-7. The vectors were intravenously injected in mice and whole body images were taken from the mice at 5 and 6 weeks post-injection to examine luciferase expression level.

All mice injected with a vector comprising a skeletal muscle-specific regulatory element (Sk-SH1-7) showed increased luciferase activity compared to control mice that were injected with a corresponding AAV9 vector without a regulatory element, indicating that all of the regulatory elements tested increased luciferase expression from the desmin promoter (data not shown). We observed very robust and enhanced luciferase activity in mice that were injected with AAV9 vector comprising the regulatory element Sk-SH1 or Sk-SH4 at 5 and 6 weeks post-injection as compared to luciferase activity of the control mice.

7 weeks post injection, the mice were euthanized and individual organs were analyzed to evaluate luciferase expression and the biodistribution pattern (Fig. 8 and 9). Up to 200 to 400-fold augmentation of luciferase activity was measured in skeletal muscle, in particular gastrocnemius muscle, tibialis interior, quadriceps, biceps and triceps, of mice injected with AAV vector comprising the Sk-SH4 regulatory element as compared to control mice (Fig. 8). There was still a 36-fold augmentation of luciferase activity measured in the heart of said mice (Fig. 8), due to the use of the desmin promoter, which is specific for both, heart and skeletal muscle. Luciferase expression was absent or limited in all other organs than cardiac and skeletal muscle tissue (Fig. 9). The second most robust regulatory element was Sk-SH1, which also specifically increased luciferase expression from the desmin promoter in skeletal muscle, and to a lesser extent in heart muscle.

The level of luciferase activity measured in the heart and different skeletal muscles was consistent with the luciferase mRNA levels measured in the respective organs (Fig. 12). Notwithstanding the transduction of the vector into different organs as shown in Figure 13, luciferase expression was absent or limited in all other organs than cardiac and skeletal muscle tissue (Fig. 9).

These *in vivo* data show that the nucleic acid regulatory elements Sk-SH, in particular Sk-SH1 and Sk-SH4, can enhance heart and skeletal muscle-specific luciferase expression.

The nucleic acid regulatory element Sk-SH4 is by far the most robust element that led to the highest level of luciferase expression in the heart and skeletal muscle as compared to the other 5 regulatory elements that were identified. The highest activity that we measured was a 400-fold upregulation of luciferase activity in the tibialis.

5 **Example 5: In vivo comparison of muscle-specific regulatory elements and CMV promoter**

Experimental procedures

AAV plasmid constructs comprising a muscle-specific regulatory element (pAAV-Sk-SH4-Des-Luc2, pAAV-CSk-SH1-Des-Luc2, pAAV-CSk-SH5-Des-Luc2) were generated
10 according to the protocol described in Example 2. Briefly, the muscle-specific regulatory elements were synthesized by conventional oligonucleotide synthesis and cloned upstream of the Desmin (Des) promoter that drives expression of a Firefly Luciferase (Luc2) reporter gene in the context of the AAV vector backbone pAAV-Des-Luc2. The plasmids also contained a Minute Virus of Mouse (MVM) intron and a Simian Virus 40
15 (SV40) polyadenylation site (pA).

A pAAVsc-CMV-Luc2-SV40pA plasmid construct was generated, wherein the Cytomegalovirus (CMV) promoter (SEQ ID NO: 30) is cloned upstream the Firefly Luciferase (Luc2) reporter gene instead of the desmin promoter in the context of the AAV vector backbone pAAV-Des-Luc2. The plasmid also contained a Simian Virus 40 (SV40)
20 polyadenylation site (pA). A schematic representation of pAAVsc-CMV-Luc2-SV40pA is shown in Figure 10A. For the cloning of the AAVsc-CMV-Luc2-SV40pA, the fragment 5'-Acc65I-CMV-HindIII-3' was synthesised and cloned into an AAVsc-Luc2-SV40pA vector which was restricted with the same pair of enzymes (i.e. Acc65I and HindIII).

AAV vector production and purification were carried out as described in Example 2.

25 6 weeks old male CB17 SC SCID mice having an average weight of 17.4 ± 1.6 g were intravenously injected into the periorbital vein with 1×10^{10} vg/mouse of concentrated vectors. Mice were imaged 4 weeks post-injection as described in Example 2. Mice were euthanized 5 weeks post-injection and intact organs were harvested and imaged as described in Example 2.

30 Results

Luciferase activity was compared in mice that were injected with a vector that expressed the luciferase gene *luc2* from the desmin (Des) promoter operably linked to a muscle-specific regulatory element: SK-SH4, CSk-SH1 or CSk-SH5, versus mice that were

injected with a vector that expressed the luciferase gene *luc2* from the Cytomegalovirus (CMV) promoter. The CMV promoter is considered one of the most powerful promoters known to allow robust gene expression in the heart. No muscle-specific regulatory element was present in the latter vectors.

- 5 Figures 10A and 10B show that the vectors comprising a muscle-specific regulatory element of the invention operably linked to the desmin promoter allow for much better expression than the vector comprising the CMV promoter.

Example 6: *In vivo* validation of skeletal muscle-specific regulatory elements via AAV vectors comprising a skeletal muscle-specific promoter

10 Experimental procedures

AAV plasmids are constructed as described in Example 4, wherein the desmin promoter is replaced by the muscle-specific promoter described in Wang et al. (2008. Gene Ther. 15:1489-1499).

- 15 AAV vector production and purification and animal studies are carried out as described in Example 2.

Results

- When using a skeletal muscle-specific promoter instead of the desmin promoter, the increase in luciferase expression is confined to skeletal muscle only, showing that the nucleic acid regulatory elements Sk-SH1-7, in particular Sk-SH4 and Sk-SH1 enhance
20 skeletal muscle-specific luciferase expression.

Example 7: *In vivo* validation of functional fragments of the identified muscle-specific regulatory elements

Experimental procedures

- The identified nucleic acid regulatory elements CSk-SH1 (SEQ ID NO: 1; 381bp), CSk-SH5 (SEQ ID NO:5; 454bp) and Sk-SH4 (SEQ ID NO:10; 435bp) were split into smaller
25 functional fragments. The transcription factor binding sites (TFBS) of the regulatory elements were mapped on the respective sequence (Fig. 11A-C), and functional fragments were generated by randomly clustering regions with TFBS. The following functional fragments were generated:

30 Functional fragments of SEQ ID NO:1:

Cluster 1A: the nucleotide sequence from position 33 to 58 in SEQ ID NO:1

Cluster 1B: the nucleotide sequence from position 90 to 142 in SEQ ID NO:1

Cluster 1C: the nucleotide sequence from position 143 to 233 in SEQ ID NO:1

Cluster 1D: the nucleotide sequence from position 240 to 310 in SEQ ID NO:1

Cluster 1E: the nucleotide sequence from position 90 to 233 in SEQ ID NO:1

Functional fragments of SEQ ID NO:5:

Cluster 5A: the nucleotide sequence from position 47 to 130 in SEQ ID NO:5

5 Cluster 5B: the nucleotide sequence from position 252 to 293 in SEQ ID NO:5

Cluster 5C: the nucleotide sequence from position 330 to 450 in SEQ ID NO:5

Functional fragments of SEQ ID NO:10:

Cluster 4A: the nucleotide sequence from position 10 to 180 in SEQ ID NO:10 (SEQ ID NO: 37);

10 Cluster 4B: the nucleotide sequence from position 190 to 240 in SEQ ID NO:10 (SEQ ID NO: 38);

Cluster 4C: the nucleotide sequence from position 241 to 300 in SEQ ID NO:10 (SEQ ID NO: 39);

15 Cluster 4E: the nucleotide sequence from position 241 to 360 in SEQ ID NO:10 (SEQ ID NO:41)

Cluster 4D: the nucleotide sequence from position 380 to 420 in SEQ ID NO:10 (SEQ ID NO: 40)

The functional fragments of Sk-SH4 are summarized in Table 5.

Table 5: Functional fragments of the muscle-specific regulatory element Sk-SH4. Bp: base
20 pairs.

Sequence	Name	Gene regulated by sequence	Size (bp)	Conserved TFBS present
SEQ ID NO: 37	Sk-SH4 ^a	MYLPF	171	CEBP, E2A, LRF, SRFb, SRFc
SEQ ID NO: 38	Sk-SH4 ^b	MYLPF	51	CEBP, E2A, LRF, SRFb
SEQ ID NO: 39	Sk-SH4 ^c	MYLPF	60	E2A, LRF, SRFb, MyoD
SEQ ID NO: 40	Sk-SH4 ^d	MYLPF	41	LRF, SRFb, p53, MyoD
SEQ ID NO: 41	Sk-SH4 ^e	MYLPF	120	CEBP, E2A, LRF, SRFb

The functional fragments of Sk-SH4 were synthesized by conventional oligonucleotide synthesis and cloned into the pAAV9sc-Des-Luc2 vector as described in Example 2. AAV vector production and purification were carried out as described in Example 2. 6 weeks old CB17/lcrTac/Prkdc scid adult mice with an average weight of about 17-18 g per mouse

were intravenously injected with the AAV vectors at a dose of 1×10^{10} vg per mouse. After 2 and 4 weeks post injection, the mice were subjected to imaging using the same biospace *In Vivo* photo Imaging System as described in Example 2.

Results

- 5 The results shown in Fig. 14 show that all 5 fragments of the Sk-SH4 regulatory element were capable of augmenting the expression of the luciferase gene driven from the desmin promoter when compared to the reference construct without any regulatory element. The fragment Sk-SH4^b showed the highest luciferase expression compared to the other fragments. However, none of the fragments could achieve higher or equal luciferase
10 expression when compared to the expression induced by the full length Sk-SH4 fragment.

Example 8: *In vivo* validation of modules of muscle-specific regulatory elements

- The muscle-specific regulatory elements are further validated by making different combinations of the regulatory elements and/or by using several copies of the same regulatory element (e.g. CSk-SH1 combined with CSk-SH5; CSk-SH1 combined with Sk-SH4; CSk-SH1 combined with CSk-SH5 and SkSH4; CSk-SH1 repeated 3x or 4x; CSk-SH5 repeated 3x or 4x; Sk-SH4 repeated 3x or 4x). These constructs are incorporated
15 upstream of the Des promoter in accordance with the protocol described in Example 2. AAV vector production and purification and animal studies are carried out as described in Example 2.

- 20 Further combinations are made with functional fragments of the identified muscle-specific regulatory elements as generated in Example 7.

Example 9: Binding of transcription factors to muscle-specific regulatory elements

Experimental procedures

- 4 weeks old mice were intravenously injected in the periorbital vein with AAV vectors containing a Sk-SH4 and CSk-SH5 regulatory element, i.e. AAV9sc-Sk-SH4-Des-Luc and
25 AAV9sc-CSk-SH5-Des-Luc, at a dose of 5×10^9 vg/per mouse.

- Heart and muscle tissues harvested from the mice, were submersed in phosphate buffered saline (PBS) containing 1% formaldehyde, cut into small pieces and incubated at room temperature for 15 minutes. Fixation was stopped by the addition of 0.125 M glycine
30 (final concentration). The tissue pieces were then treated with a Tissue-Tearer and finally spun down and washed twice in PBS. Chromatin was isolated by the addition of lysis buffer, followed by disruption with a Dounce homogenizer. Lysates were sonicated and the DNA sheared to an average length of 300-500 bp. Genomic DNA (Input) was prepared

by treating aliquots of chromatin with RNase, proteinase K and heat for de-crosslinking, followed by ethanol precipitation. Pellets were re-suspended and the resulting DNA was quantified on a NanoDrop spectrophotometer. Extrapolation to the original chromatin volume allowed quantitation of the total chromatin yield. An aliquot of chromatin (30 µg) was pre-cleared with protein A agarose beads (Invitrogen, catalogue number 15918-014). Genomic DNA regions of interest were isolated using 4 µg of antibody against MEF2 (Santa Cruz, sc-313), SRF (Santa Cruz, sc-335) and CEBP (Santa Cruz, sc-150). Complexes were washed, eluted from the beads with SDS buffer, and subjected to RNase and proteinase K treatment. Crosslinks were reversed by incubation overnight at 65°C, and ChIP DNA was purified by phenol-chloroform extraction and ethanol precipitation. Quantitative PCR (QPCR) reactions were carried out in triplicate on specific genomic regions using SYBR Green Supermix (Bio-Rad). The resulting signals were normalized for primer efficiency by carrying out QPCR for each primer pair using Input DNA. Primer sequences for Sk-SH4 were: 5'-GTCCCTCACTCCCAACTCAG-3' (SEQ ID NO: 33; forward) and 5'-GAGGAGAAGGAGATCAGACACTG-3' (SEQ ID NO: 34; reverse); for CSk-SH5: 5'-TAGCTGGGCCTTTCCTTCTC-3' (SEQ ID NO: 35; forward) and 5'-CGTCTCCCTAGCAGCAACAG-3' (SEQ ID NO: 36; reverse). Negative control primers were purchased from Active Motif (Carlsbad, CA, USA) (#71012) and are specific for non-transcribed gene sequences on chromosome 17.

20 Results

The muscle-specific regulatory element Sk-SH4 contains several putative transcription factor binding sites (TFBS), including E2A, SRF, p53, CEBP, LRF, MyoD and SREBP. Using a chromatin immuno-precipitation (CHIP) assay, binding of CEBP and SRF on the Sk-SH4 element was confirmed in the heart and skeletal muscle from mice that were injected with AAV9sc-Sk-SH4-Des-Luc vector (Fig. 15 A,B). Similarly, binding of CEBP and MEF2 element on the CSk-SH5 regulatory element was shown in the heart and skeletal muscle from mice injected with AAV9sc-CSk-SH5-Des-Luc (Fig. 15 C,D).

Example 10: Therapeutic evaluation of muscle-specific regulatory elements via AAV vectors comprising a desmin promoter

30 The effect of the muscle-specific regulatory element Sk-SH4 on the expression and therapeutic efficacy of two therapeutic genes, in particular micro-dystrophin and follistatin, which have therapeutic potential for the treatment of muscle disease such as Duchenne muscular dystrophy (DMD) by gene therapy, was evaluated. MDX-SCID mice replicate the disease manifestations of Duchenne muscular dystrophy in patients and are therefore well

suited to assess therapeutic efficacy of the AAVss-Sk-SH4-Des-MVM-MD1 and AAVss-Sk-SH4-Des-MVM-FST-2A-Luc vectors.

Experimental procedures

Cloning follistatin and MD1 genes into AAV

- 5 The microdystrophin 1 (MD1) (Koo et al., 2011) and follistatin (FST) (Kota et al., 2009) genes were cloned and driven from desmin promoter, which was operably linked to the muscle-specific regulatory element Sk-SH4. The MD1 gene was flanked by MluI and XhoI restriction sites at the 5' and 3' ends, while the FST gene was flanked by MluI and SalI restriction sites at the 5' and 3' ends, respectively and were synthesized by conventional
10 oligonucleotide synthesis. The Sk-SH4 regulatory element, operably linked to the desmin promoter, was cloned upstream of the MVM intron in the context of a single stranded adeno-associated viral vector (AAVss) backbone. The vector also contained a 49bp synthetic proudfoot polyadenylation site (Levitt N et al, 1989). The follistatin gene was linked to a luciferase reporter gene via a 2A polypeptide. The generated constructs were
15 designated as pAAVss-Sk-SH4-Des-MVM-MD1 and pAAVss-Sk-CRM4-Des-MVM-FST-2A-Luc, respectively, and are schematically shown in Figure 16.

AAV vector production and purification were carried out as described in Example 2.

Treadmill test for phenotypic correction of MDX-SCID mice

- 4 weeks old MDX-SCID mice (bred in house) were injected intravenously in a total volume
20 of 100 μ l with a dose of 2×10^{10} vg per mouse of the AAVss-Sk-SH4-Des-MVM-MD1 and AAVss-Sk-SH4-Des-MVM-FST-2A-Luc vectors, individually or in combination. 8 weeks post-injection, the treated and control (injected with PBS) MDX-SCID mice were subjected to the treadmill test. The treadmill tests were performed using the Exer-3/6 open treadmill (Columbus instruments, USA). The inclination of the belt was adjusted to 10° uphill before
25 performing the test. The initial speed was set at 10 m/min and thereafter the speed was increased by 1 m every minute. The test was terminated at a point when the mice sat for $\geq$ 5 seconds on the pulse grill. At that point the distance covered by the mice was recorded and the total distance covered by the mice during the course of the test was calculated by using the formula, distance= $((N+n)/2) \times (N-n+1)$ where N is the time (in min) at the point of
30 termination of the test and n is the time (in minutes) at the start of the test.

mRNA analysis

mRNA analysis was performed as described in Example 2. To quantify the mRNA of the microdystrophin and follistatin genes, a qPCR-based method was used using the primers

5'-GTGCCCTACTACATCAA-3' (SEQ ID NO:42) as forward primer and 5'-AGGTTGTGCTGGTCCA-3' (SEQ ID NO:43) as reverse primer (amplicon 206 bp) for the microdystrophin, and the Luc2-specific forward primer 5'- CCCACCGTCGTATTTCGTGAG-3' (SEQ ID NO: 14) and reverse primer 5'- TCAGGGCGATGGTTTTGTCCC-3' (SEQ ID NO: 15) (amplicon 217 bp) for the follistatin gene.

Results

The results of the treadmill test clearly showed that the mice treated by gene therapy with either MD1 or FST or the combination of both, outperformed the untreated control MDX-SCID mice (Fig. 17). The MDX-SCID mice injected with either the AAVss-Sk-SH4-Des-MVM-MD1 or the AAVss-Sk-SH4-Des-MVM-FST-2A-Luc vector covered twice the distance covered by the control mice. When both vectors were co-injected into the MDX-SCID mice, a distance of 4 km was covered, which is 4 times more than the distance covered by the control MDX-SCID mice injected with PBS. These results clearly demonstrate that therapeutic efficacy can be achieved using these vectors.

The significant physiologic effect of the gene therapy using the AAVss-Sk-SH4-Des-MVM-MD1 or the AAVss-Sk-SH4-Des-MVM-FST-2A-Luc vector or their combined use was confirmed by histologic examination. Hematoxylin/eosin-stained muscle sections were scored for % centrally nucleated cells for the different experimental groups (Fig. 18). The muscle sections of C57B6 wild-type mice showed predominantly peripheral localization of the nucleus in the myofibers (data not shown) and a minimal percentage (<2%) of the nuclei was centrally located (Fig. 18B). In untreated, control, MDX/SCID mice, the nuclei were predominantly (about 50% of the nuclei) centrally located in the transversally transected myofibers (Fig. 18Aa, Fig. 18B). MDX/SCID mice injected with AAVss-Sk-SH4-Des-MVM-MD1 (AAV9-MD1) (Fig. 18Ac, Fig. 18B) or AAVss-Sk-SH4-Des-MVM-FST-2A-Luc (AAV9-FST) (Fig. 18Ab, Fig. 18B), all showed a decreased number of myofibers that are centrally nucleated. Co-administration of both, AAVss-Sk-SH4-Des-MVM-MD1 and AAVss-Sk-SH4-Des-MVM-FST-2A-Luc vectors, resulted in an even more profound shift in nuclear localization towards the periphery of the myofibers, and a further reduction in the percentage of centrally located nuclei (Fig. 18Ad, Fig. 18B). These results are indicative of a reduced regenerative stimulus and phenotypic correction after the gene therapy. Consequently, the shift of nuclear localization from a central to a more perihelal location in the myofibers is consistent with the phenotypic correction as shown by the improved muscle strength and mobility in the treadmill assay .

Using quantitative RT-PCR, the expression of the micro-dystrophin (MD1) and the follistatin (FST) genes in biopsies from heart and muscle tissues, in particular

gastrocnemius and quadriceps, of the mice injected with the pAAVss-Sk-SH4-Des-MVM-MD1 construct, the pAAVss-Sk-SH4-Des-MVM-FST-2A-Luc construct or the combination of both constructs was assessed (Fig. 19).

Example 11: *In vivo* comparison of muscle-specific regulatory elements with CMV and SPc5-12 promoters

Experimental procedures

AAV vector production and purification were carried out as described in Example 2.

The AAVsc-Des-MVM-Luc vector is a self-complementary AAV vector containing a luciferase (Luc) transgene driven from the desmin promoter. The scAAV vector backbone also contained a Minute Virus of Mouse (MVM) intron and a Simian Virus 40 (SV40) polyadenylation site (pA).

The AAVsc-SPc5-12-MVM-Luc vector had the same configuration as the AAVsc-Des-MVM-Luc vector, but the desmin promoter was replaced by the SPc5-12 promoter.

AAV vectors comprising the muscle-specific regulatory element Sk-SH4 were generated according to the protocol described in Example 2. Briefly, the muscle-specific regulatory element Sk-SH4 was synthesized by conventional oligonucleotide synthesis and cloned upstream of the desmin or the SPc5-12 promoter in the context of the AAV vector backbone of AAVsc-Des-MVM-Luc or AAVsc-SPc5-12-MVM-Luc, respectively.

The AAVsc-CMV-Luc vector was generated by cloning the Cytomegalovirus (CMV) promoter (SEQ ID NO: 30) upstream the luciferase reporter gene instead of the desmin promoter in the context of the AAV vector backbone of AAVsc-Des-Luc, which also contained a polyadenylation site (pA).

A schematic representation of the different vectors is shown in Figure 20.

Adult CB17/IcrTac/Prkdcscid mice were intravenously injected with the different vectors (n=3) at a dose of 1×10^{10} vg/mouse.

mRNA analysis was performed as described in Example 3.

Results

Luciferase mRNA levels in mice injected with a vector that expressed the luciferase gene from the desmin promoter operably linked to the muscle-specific regulatory element Sk-SH4 were compared versus mice that were injected with a vector that expressed the luciferase gene from the Cytomegalovirus (CMV) or the SPc5-12 promoters, which were not operably linked to a muscle-specific regulatory element. The CMV and SPc5-12

promoters are considered the most powerful promoters known to allow robust gene expression in the heart and skeletal muscles. Figure 21 shows that the vector comprising the Sk-SH4 muscle-specific regulatory element of the invention operably linked to the desmin promoter allows for much better expression than the vectors comprising the CMV or the SPc 5-12 promoter without muscle-specific regulatory element.


Similarly, when the Sk-SH4 muscle-specific regulatory element was operably linked to the SPc5-12 promoter, increased luciferase mRNA levels were observed in the heart and all skeletal muscle types tested (Fig. 21). However, the mRNA level induced from this construct was lower than when the Sk-SH4 muscle-specific regulatory element of the invention is operably linked to the desmin promoter.

CLAIMS:

1. A nucleic acid regulatory element for enhancing muscle-specific gene expression, comprising the sequence of SEQ ID NO:5, a sequence that is at least 95% identical to SEQ ID NO:5 over the entire length and that enhances muscle-specific gene expression, or
5 consisting of a functional fragment of the sequence of SEQ ID NO:5, wherein said functional fragment comprises the sequence from positions 47 to 130 of SEQ ID NO:5; the sequence from positions 252 to 293 of SEQ ID NO:5; or the sequence from positions 330 to 450 of SEQ ID NO:5.
2. The nucleic acid regulatory element according to claim 1, wherein the nucleic acid regulatory element enhances cardiac and skeletal muscle-specific gene expression.
- 10 3. The nucleic acid regulatory element according to claim 1 or 2, wherein said functional fragment consists of the nucleotide sequence from positions 47 to 130 of SEQ ID NO:5; the sequence from positions 252 to 293 of SEQ ID NO:5; or the sequence from positions 330 to 450 of SEQ ID NO:5.
4. The nucleic acid regulatory element according to any one of claims 1 to 3, having
15 a maximal length of 600 nucleotides.
5. The nucleic acid regulatory element according to any one of claims 1 to 3, having a maximal length of 550 nucleotides.
6. The nucleic acid regulatory element according to claim 4 or 5, comprising the sequence of SEQ ID NO:5, or a sequence that is at least 95% identical to SEQ ID NO:5 over
20 the entire length.
7. Use of the nucleic acid regulatory element according to any one of claims 1 to 6 in a nucleic acid expression cassette, or a vector.
8. A nucleic acid expression cassette comprising at least one nucleic acid regulatory element according to any one of claims 1 to 6, operably linked to a promoter and a transgene.
- 25 9. The nucleic acid expression cassette according to claim 8, wherein the promoter is a muscle-specific promoter.
10. The nucleic acid expression cassette according to claim 9, wherein the promoter is the promoter of the desmin (DES) gene.
11. The nucleic acid expression cassette according to any one of claims 8 to 10, wherein the
30 transgene encodes a therapeutic protein, an immunogenic protein, or a structural protein.
12. The nucleic acid expression cassette according to any one of claims 8 to 10, wherein the transgene encodes dystrophin or a sarcoglycan.

13. A vector comprising the nucleic acid regulatory element according to any one of claims 1 to 6, or the nucleic acid expression cassette according to any one of claims 8 to 12.
14. The vector according to claim 13, which is a viral vector.
15. The vector according to claim 13, which is an adeno-associated viral vector.
- 5 16. A pharmaceutical composition comprising the nucleic acid expression cassette according to any one of claims 8 to 12, or the vector according to any one of claims 13 to 15, and a pharmaceutically acceptable carrier.
- 10 17. Use of the nucleic acid regulatory element according to any one of claims 1 to 6, the nucleic acid expression cassette according to any one of claims 8 to 12, the vector according to any one of claims 13 to 15, or the pharmaceutical composition according to claim 16, in gene therapy.
- 15 18. Use of the nucleic acid regulatory element according to any one of claims 1 to 6, the nucleic acid expression cassette according to any one of claims 8 to 12, the vector according to any one of claims 13 to 15, or the pharmaceutical composition according to claim 16, in vaccination therapy.
19. Use of the nucleic acid regulatory element according to any one of claims 1 to 6, the nucleic acid expression cassette according to any one of claims 8 to 12, the vector according to any one of claims 13 to 15, or the pharmaceutical composition according to claim 16, in prophylactic vaccination.
- 20 20. An in vitro or ex vivo method for expressing a transgene product in muscle cells comprising:
- introducing the nucleic acid expression cassette according to any one of claims 8 to 12, or the vector according to any one of claims 13 to 15 into the muscle cells; and
 - expressing the transgene product in the muscle cells.

FIG. 1



2/38

FIG. 2

A

SEQ ID NO: 1

GTTCTCCTCTATAAATACCCGCTCTGGTATTTGGGGTTGGCAGCTGTTGCTGCCAGGGAGA
TGGTTGGGTTGACATGCGGCTCCTGACAAAACACAAACCCCTGGTGTGTGTGGGCGTGGGT
GGTGTGAGTAGGGGGATGAATCAGGGAGGGGGCGGGGACCCAGGGGGCAGGAGCCACACA
AAGTCTGTGCGGGGGTGGGAGCGCACATAGCAATTGGAACTGAAAGCTTATCAGACCCTT
TCTGGAAATCAGCCCCTGTTTATAAACTTGAGGCCCCACCCTCGACAGTACCGGGGAGGA
AGAGGGCCTGCACTAGTCCAGAGGGAACTGAGGCTCAGGGCTAGCTCGCCCATAGACATA
CATGGCAGGCAGGCT

B

SEQ ID NO: 2

CGTCTATAAATTCCAGGGAAGGTCTCTGATTGGCCCTGCTCATTCCCAGGCCCATTCTTG
ACCCAGTCACTGAAGTCAGGGAGATGCAGTAATAAGACTGGCTGGAATCAGGGTCTTTAGG
GGTGGAGGGATGGGGAGGAGGCACAGCATGTTCATCAAAATAAGGAAATTGCAAAAGAAAGC
TTGCAGGCTACTTTGAATGACAATGAGAAAGACGGTGCTGCCTGAGTGTGTTAAGGATCCA
CATGGTCTCCAAAATCCTCCAGGAGCATAACAGTCTAGTCTGGGAGATGAGACACAAAAATA
ACCAGAACACACAGCTTGCACTGACTCGAGGGCTGGATAAGAATATCTGGAACCTCCCCA
TCTATTTTCAGAAGCTTGTCTCTTGATGAAAATTAGACACTTAATGGGAAAGGGCTTTGAA
AAGAGTGC

C

SEQ ID NO: 3

AAAAAATAAAAAATAAAAAATAAAAAATAAAGTGGATGGCTTAGAGTATCTTTTCTGTT
TAGACCTGACTAAAGCTTAGACATAATTGTTAGTTTAGGCTCTCAGGGTAAATTTATTAC
TGTAATCCAAAAATCCCTTCTTCTTTTTTTTTTTTTTTTTTTGGTCCTTGAATTAAAT
GCTGTCACCTCCTTCTTGAAAGGAGAACTATTAGTCAGATTTGAAAATCCTCTTTATCAC
CCAGGAAAATCATTTTTATGGACACTTTGTCTTTCTGTAGTCTGACTTAGAAGCAGCCTGT
TTTTGATAGGTTGAAGTTTTTCATCTTGAACACAAACCCTGTTTGTGTGTCCCCTACTCCCC
AGTTTGATGTGCCAGGCACTTTGTTCTCAAGCCAGCAGCTGTTGTGGGATGAGGGGACAT
TTTGCATGCTTAGCCAGCAGCTGCCAGAAACATTTCTAATCTGGTTTTTGGCAGGAAATAGG
GCACAAGTGGAAGCCAAGTTAAAAGAAGCTGGAAAAATAAACAGAATAACTTTAGATGTCA
CT

3/38

FIG. 2

D

SEQ ID NO: 4

TGCACTGTAGACTATTTATGTTTGCATATTTTTCAGGATTGTGTACAGTAAACCTTAAGTG
AATACAAAGGATCCAATTGTCCTGTAAGACTCACTTCAATTACAGATTGTGCTCAGTATTA
AACTTTGCTAGTACTTTCAGGATGCAGTCAATCAAGAGGCAGGGAAAGGTCTGTCAGCATC
CACAGCCTCCTTTTCAAATTGGACACTTAGTCTCTGGTCCGAATAATAGGTGCCCCCAGTG
TCACCTCACACAATGGGGTATACACAGTTTTTAAAAATTTATTTTCAGCTGGGCACAGTAG
CTCACACCTAAAATCCCAGCACTTTGGGAGGCTGAGGCAGGTGGATCACCTGAGGTCAGGA
GTTCAAGACCAGCCTGGCCAACGTGGCAAAATCCTGTCTCTACTAAAAATACAAAAATTAA
CCG

E


SEQ ID NO: 5

CAGTTTACTCACCAGGGATTTCAGAGGCAGCACTGCTGAACCCTGAGCCCTTGGCACATCAG
GTTGGCTGTCAGAAGTCGGCCTTTGTACATACACAGTTCCCTTGTGAGGCCCAGCTGCGTG
TCCTAGGAGCGGGGCCTCTCTCCACAGCAGAGCTCAGCCTCTCAAGTGTATGGACAGCACG
GGTGCCTGATGGGTGGATTTAGCCATGAGTTGAAGGTGGCTTGGGGAGAATGAGAGTTCTA
GAGATAGGGAGAAGGGGTTGCCAATAGGAGAGTGGAATTCCTGAGCACCTCGTCACAGGCA
GCCGACAGAACATGAGCCGCAGGGCCCAGGCTATTTATACCTCGCCTGTCACTATCAGGGT
CCCCACAGCTCCCCCCCACCTCCAGCCACACACAGCAGGTCCTTTTGCTCTTTCTGGTCCCT
TCTCTACTCCTCCCCCTCCCTACCTAA

F

SEQ ID NO: 6


CCCTTTCCCTGGCAGGATCTGCCCTGTGGCCCAAATGGGCATGTTGCCCAGGGGGCTCCCT
GGCACTATGGGGGAAGAGTCTCTCCTTCCCCTCTTATCATCTCAGTTGAGTCAGACTTG
GGAGGGGGGATACACAGTGTGAGTCACTGGGTACCCTTTTCCTGAGCTCAGCTTCATACCGA
GGCGATGAGGCCAAACGGGCTGGTGACAGGGACACTGAGTCAGGGGCAGGGGCCCCGGTCT
TACTCCTGGGCCTCTGGATTTGGGCCCTACATGAGGCTTTTCTATCTGTAAAGTCAAGCAA
TGGCTGGGAGGCACACACAACCCCCCGCCCCCGCAGGCTTCTCCTTCATTGGCCCCGGGC
AAGGTCCCTGCTTCCCTCTCAGGCCGTCTCTGCACAAGCACACACACTTCCCTTCCCTGTCC
ACAGGTGGACAATGCCCTGGGCTAGG

FIG. 3

5/38

FIG 4

A



6/38

FIG 4

B





FIG 5


A



8/38

FIG. 5


B




9/38

FIG. 6

A



B



10/38

FIG. 7**A**

SEQ ID NO: 7

CTCACTCCCCGCCAGGCAGCAAGGAGCCCACACCCTCATGCCCCCTCAGCTTCAGCCCCCA
CCTCCAGGAGGCCCTACCCACGCTCATGACCTTGCTATTCTGGGCCTTGCTGTCTGTAGGG
AGATGGACAGGAGACAGCTGGGCTTCCAGGCCACCCAGGCGGGGGGCTAGCCGAGGGAAGC
CTGCTGGCTCTCCTGCTTGCTCTAATTTCTGGGGCTCCCCAAACCTTGGCCTCAGGAGACT
GGGGATAGGACCGGCCTTGAAAGTGGGGGAAGCTTTGGAGAGCCGGGTGCTGGGTTCCTAG
TGAGATGGCCAGTGAAGGCTGTGGTGCCCCGAGGTAAGCAGGGCCTGATCCCCCTCCTAATC
TTCCAGCAGCAACTGGTGCTCTGAGGCTCCCCCTCCCCCAGCCCTGCCAGCCTTCAGGGAC
CTGCCTTCCAAAGATGGGCAGGGGAGGGGGACGAGGACACCCACCCACTCCTCAGACCAGC
ATGTCTT

B

SEQ ID NO: 8

CCCTCCAGATGGGTTTCCTGGAATCTAGATTTCCCAGGTTCCAAAGGACACCCGAGTCTCA
TGCCTGGAACCTCAGTGAGACTAATTCACCTCTCCTCTGCCCTAATCTTCATCTCCAGCCAG
AAGCCAACAGATCCCAGGGGACTGGAGCCACAGGGGCTGCACCTGTTTACCGGGTATTTTTT
AGGATGGTTGATGAACACATAATACCCACCCTATAGTCAGAGAAAGACAATGCCTGCTATG
TTAATCCTGTGGCTATTATAGTCTGTCATCTCATGGGTTGGGGCAGGACACTGACCCTCTC
AGAGGCCAGAGAGAGGCCTCGCAAGCAGGAGGTTAGGGA

C

SEQ ID NO: 9

ATGGAGACAATCCATGAATTCCTGAGATGCTTGGCTGGTATTAGATTTTATGGGCAGCTGC
TTATTCTTAGGGCTCTGCTTCTCCAAAGACACTGAGGAAGTCCAAAGGAAACACCAGCTGG
CGAAGAGCCACCTCCAGGCCCATCTGTCCATCATCAGCCTCCAGGAATGCCAGTGTCCAGA
GGGCACCAGGTCTGCGTCTGTCTCCCTGGGATGTGCCTTGTCCTTGGTGGGCATTTGGCAG
TGATCATGCCTCCCTGTCTCCCTCAGAGATCCAACTGTCCCCATTGTGGGGCCCTACCTTC
CAAGGCCGTTTACACCTCCTGCCAAGCTCCGGGGCCTGCCCCAGCCTGCCTCACTGACA
AATGCCAGACCAAGGGGTCCCACGTCAGGCAAGAGGCCTCAGCCTGTGCTCTGACACCCCT
CAG

D

SEQ ID NO: 10

TTCTGAGTCCTCTAAGGTCCCTCACTCCCAACTCAGCCCCATGTCCTGTCAATTCCCCTC
AGTGTCTGATCTCCTTCTCCTCACCTTTCCCATCTCCCGTTTGACCCAAGCTTCCTGAGCT
CTCCTCCCATTCCCCTTTTTGGAGTCCTCCTCCTCTCCCAGAACCAGTAATAAGTGGGCT
CCTCCCTGGCCTGGACCCCCGTGGTAACCCTATAAGGCGAGGCAGCTGCTGTCTGAGGCAG
GGAGGGGCTGGTGTGGGAGGCTAAGGGCAGCTGCTAAGTTTAGGGTGGCTCCTTCTCTCTT
CTTAGAGACAACAGGTGGCTGGGGCCTCAGTGCCCAGAAAAGAAAATGTCTTAGAGGTATC
GGCATGGGCCTGGAGGAGGGGGGACAGGGCAGGGGGAGGCATCTTCCTCAGGACATCGGGT
CCTAGAGG

11/38

FIG. 7

E

SEQ ID NO: 11

GACTAGGAATAAATCACATATCCTCAATCCCTGGACAACTTGTTTACTTCTAGTGTTAGTT
TTTTCTTAAAAAAAAAATTGAAATCATTCTGAGGCTGGAATACTTTGGACATGCCCAGCAG
TCCTGGCAGTTCCACAGAAGCATTACCTCATGACTGGAGTGGGTAAAGCATACTGTGGG
CTATGGATAAGACTGACATTAACCACAAGCATGTTTGGCAGCAGACTGGTGCTTTACAAGC
TCCATGTTTCAGCAGGAGCTGCAAAGTGTTCTCCAAACCAATATTTGTCATTCTTGGATTC
TATTTAGGAGGTCCTGTTACTCACATGTTTCAATATCAGCAGAAGCCAGTTTCCCTGTGGT
ACCGAAGTGGATCCTGATGAATTTACCCTTGTAAGTAAAAAAATGATGTTATACCCAAAG
CTTGAAGTACGTAGTGGGGATGCCACTGAAATAATTCAGACATGCTT

F

SEQ ID NO: 12

GTGCTCATAGCTCCACCTTTTGTTTCCTAATATGGTCTTTCCAGCTCCCTCCACCCCATCAT
TGTTCTCCTGGGGGAACACAGGGTGAGACGCTTTGATGAACTGACATCACCAGCAAAAAA
ATATCTAGCAACAGCTGAGGCTGATTTTAGACAATGGAAAGTGGGGAGGGAAGAGGTTCT
CCCTGACCCTGAACTTTCCACTCATTCTGGGCAGCTCTATGGATGTTTTAAAGAAGAGG
AAGAGGGGAGGGAAGAACATTGAAATAGAGAAGTGACTTTGGCAATTCTAGGTGGCAGT
TTGCATCCAGGGGGTCCTGGTTGCCTTTTCAGCTTCCCGTTTCACTCTCCCCCAGACTGTGT
TGAATGCTGGTCAAACCTCCGTTAGTTGAGTTTTAGCTTTTGATTCTGGTATTCAAGGAGC
TTGGGCACAGGGAAGAGGGGAGGTCATCATGATCCTTAACAATTCTCCAGATCCCCAGA
TCAAATTGCTGTGCTATTCTGGGAGTCTCCG

G

SEQ ID NO: 13

ATCGTGTGTCAGAGGTTTGTGTCAGCTTCCCAGCAAGGGAACCAGAAAGGAAAAGGAACCG
GTTCCCTCATGCTTCCTAGGGGAATGCATGCATATCTGAAGAGAAGGGAATCTTATATAAGG
CTGTTTAGCTAAGGGCAGCCACCAGCCAGGTGAGCCTTACAGAAGCACAGGGCTGGGTGTC
TGCAGTTCCCTAGCAGATTAACCTGGGTCACAGTGACTCAGAGCTCCAGCATGCGAGTTCC
AGGTGTGGAAGTACAGATCTGCTTTTGCTCCACTTGGGAGTATTTTTCCTTCT
TAGTGAGCATGGGCAGCCTCCTGGCCAGGGAAGTCTGGCACTGTCTGGGCCTGACAGGGAA
ACCCTG

FIG. 8






FIG. 9



14/38

FIG. 10


A.



15/38

FIG.10

B



16/38

FIG.10

C




FIG.11

A




FIG. 11

B






FIG. 11

C



20/38


FIG. 12




21/38

FIG. 13

A



B



22/38

FIG. 14





FIG. 15

A



B






FIG. 15

C




D



25/38

FIG. 16

A



26/38

FIG. 16**B**

SEQ ID NO:44


CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCCGGGCGTCG
GGCGACCTTTGGTCGCCCCGGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAAC
TCCATCACTAGGGGTTCCTCGTACGTTCTGAGTCTCTAAGGTCCCTCACTCCCAACTCAG
CCCCATGTCTGTCAATTCCCCTCAGTGTCTGATCTCCTTCTCCTCACCTTTCCCATCTC
CCGTTTGACCCAAGCTTCCTGAGCTCTCCTCCCATTCCCCTTTTTGGAGTCTCCTCCTCT
CCCAGAACCCAGTAATAAGTGGGCTCCTCCCTGGCCTGGACCCCCGTGGTAACCTATAAG
GCGAGGCAGCTGCTGTCTGAGGCAGGGAGGGGCTGGTGTGGGAGGCTAAGGGCAGCTGCTA
AGTTTAGGGTGGCTCCTTCTCTCTTCTTAGAGACAACAGGTGGCTGGGGCCTCAGTGCCCA
GAAAAGAAAATGTCTTAGAGGTATCGGCATGGGCCTGGAGGAGGGGGGACAGGGCAGGGGG
AGGCATCTTCTCAGGACATCGGGTCCTAGAGGGACCTTGCTTCCTAGCTGGGCCTTTCTCT
TCTCCTCTATAAATACCAGCTCTGGTATTTGCGCTTGGCAGCTGTTGCTGCTAGGGAGACG
GCTGGCTTGACATGCATCTCCTGACAAAACACAAACCCGTGGTGTGAGTGGGTGTGGGCGG
TGTGAGTAGGGGGATGAATCAGAGAGGGGGCGAGGGAGACAGGGGCGCAGGAGTCAGGCAA
AGGCGATGCGGGGGTGC GACTACACGCAGTTGGAAACAGTCGTCAGAAGATTCTGGAAACT
ATCTTGCTGGCTATAAACTTGAGGGAAGCAGAAGGCCAACATTCTCCCAAGGGAAACTGA
GGCTCAGAGTTAAACCCAGGTATCAGTGATATGCATGTGCCCCGGCCAGGGTCACTCTCT
GACTAACCGGTACCTACCCTACAGGCCTACCTAGAGACTCTTTTGAAAGGATGGTAGAGAC
CTGTCCGGGCTTTGCCCACAGTCGTTGGAAACCTCAGCATTTTCTAGGCAACTTGTGCGAA
TAAACACTTCGGGGGTCTTCTTGTTCATTCCAATAACCTAAACCTCTCCTCGGAGAAA
ATAGGGGGCCTCAAACAAACGAAATTCTCTAGCCCGCTTTCCCCAGGATAAGGCAGGCATC
CAAATGGAAAAAAGGGGCCGGCCGGGGGTCTCCTGTCAGCTCCTTGCCCTGTGAAACCCA
GCAGGCCTGCCTGTCTTCTGTCCTCTTGGGGCTGTCCAGGGGCGCAGGCCTCTTGCGGGGG
AGCTGGCCTCCCCGCCCCCTCGCCTGTGGCCGCCCTTTTCTGTCAGGACAGAGGGATCCT
GCAGCTGTCAGGGGAGGGGCGCCGGGGGGTGTGTCAGGAGGGCTACAAATAGTGCAGACA
GCTAAGGGGCTCCGTCACCCATCTTCACATCCACTCCAGCCGGCTGCCCGCCCGCTGCCTC
CTCTGTGCGTCCGCCAGCCAGCCTCGTCCACGCCGCCACCTCTAGAAAGAGGTAAGGGTT
TAAGGGATGGTTGGTTGGTGGGGTATTAATGTTTAATTACCTGGAGCACCTGCCTGAAATC
ACTTTTTTTTTCAGGTTGGACGCGTGCCACCATGCTGTGGTGGGAGGAAGTGGAGGACTGCTA
CGAGAGAGAGGACGTGCAGAAGAAAACCTTCACCAAGTGGGTGAACGCCCAGTTCAGCAAG
TTCGGCAAGCAGCACATCGAGAACCTGTTTCAGCGACCTGCAGGATGGCAGGAGACTGCTGG
ATCTGCTGGAGGGACTGACCGGCCAGAAGCTGCCCAAGGAGAAGGGCAGCACCAGAGTGCA
CGCCCTGAACAACGTGAACAAGGCCCTGAGAGTGCTGCAGAACAACAACGTGGACCTGGTG
AATATCGGCAGCACCGACATCGTGGACGGCAACCACAAGCTGACCCTGGGCCTGATCTGGA
ACATCATCCTGCACTGGCAGGTGAAGAACGTGATGAAGAACATCATGGCCGGCCTGCAGCA
GACCAACAGCGAGAAGATCCTGCTGAGCTGGGTGAGGCAGAGCACCAGAACTACCCCCAG
GTGAACGTGATCAACTTCACCACCTCCTGGAGCGACGGCCTGGCCCTGAACGCCCTGATCC
ACAGCCACAGACCCGACCTGTTGACTGGAACAGCGTGGTGTGTCAGCAGAGCGCCACCCA
GAGACTGGAGCACGCCTTCAACATCGCCAGATACCAGCTGGGCATCGAGAAGCTGCTGGAC
CCCGAGGACGTGGACACCACCTACCCCGACAAGAAAAGCATCCTGATGTATATTACCTCTC
TGTTTTTCAGGTGCTGCCCCAGCAGGTGTCCATCGAGGCCATCCAGGAAGTGGAATGCTGCC
CAGGCCCCCCCCACCGTGTCCCTGGCCCAGGGCTATGAGAGAACCAGCAGCCCCAAGCCCAGA
TTCAAGAGCACCGTGTCCCTGGCCCAGGGCTATGAGAGAACCAGCAGCCCCAAGCCCAGAT
TCAAGAGCTACGCCTACACCCAGGCCGCCTACGTGACCACCTCCGACCCCCACCAGAAGCCC
CTTCCCCAGCCAGCACCTGGAGGCCCGGAGGACAAGAGCTTCGGCAGCAGCCTGATGGAG

AGCGAAGTGAACCTGGACAGATACCAGACCGCCCTGGAGGAAGTGCTGTCTTGGCTGCTGT
CCGCCGAGGACACCCTGCAGGCCCAGGGCGAGATCAGCAACGACGTGGAAGTGGTGAAGGA
CCAGTTCCACACCCACGAGGGCTACATGATGGATCTGACCGCCCACCAGGGCAGAGTGGGC
AATATCCTGCAGCTGGGCAGCAAGCTGATCGGCACCGGCAAGCTGAGCGAGGACGAGGAGA
CCGAAGTGCAGGAGCAGATGAACCTGCTGAACAGCAGATGGGAGTGCCTGAGAGTGGCCAG
CATGGAGAAGCAGAGCAACCTGCACCGCGTGCTGATGGACCTGCAGAACCAGAAGCTGAAG
GAGCTGAACGACTGGCTGACCAAGACCGAGGAGCGGACCAGAAAAGATGGAGGAGGAGCCCC
TGGGCCCCGACCTGGAGGACCTGAAGAGACAGGTGCAGCAGCACAAAGTGCTGCAGGAGGA
CCTGGAACAGGAGCAGGTGCGCGTGAACAGCCTGACCCACATGGTGGTCGTGGTGGACGAG
AGCAGCGGCGACCACGCCACAGCCGCCCTGGAAGAGCAGCTGAAAGTGCTGGGCGACAGAT
GGGCCAACATCTGCCGGTGGACCGAGGACAGATGGGTGCTGCTGCAGGACATCCTGCTGAA
GTGGCAGAGACTGACAGAGGAGCAGTGCCTGTTTAGCGCCTGGCTGAGCGAGAAGGAGGAC
GCCGTGAACAAGATCCACACCACCGGCTTCAAGGACCAGAACGAGATGCTGAGCAGCCTGC
AGAAGCTGGCCGTGCTGAAGGCCGATCTGGAGAAGAAAAAGCAGAGCATGGGCAAGCTGTA
CTCCCTGAAGCAGGACCTGCTGTCCACCCTGAAGAACAAGAGCGTGACCCAGAAAACCGAG
GCCTGGCTGGACAATTTGCCCCGGTGCTGGGACAATCTGGTGCAGAACTGGAGAAGAGCA
CCGCCCAGATCAGCCAGGCCGTGACCACCACCCAGCCCAGCCTGACACAGACCACCGTGAT
GGAGACCGTGACCACAGTGACCACCAGGGAGCAGATCCTGGTGAAGCACGCCCAGGAGGAG
CTGCCCCCTCCCCCCCCCTCAGAAGAAGCGGCAGATCACAGTGGACACCCTGGAGAGACTGC
AGGAGCTGCAGGAAGCCACCGACGAGCTGGACCTGAAGCTGAGACAGGCCGAAGTGATCAA
GGGCAGCTGGCAGCCTGTGGGCGATCTGCTGATCGACAGCCTGCAGGACCACCTGGAGAAA
GTGAAGGCCCTGCGGGGCGAGATCGCCCCCTGAAGGAGAATGTGAGCCACGTGAACGACC
TGGCCAGACAGCTGACCACCCTGGGCATCCAGCTGAGCCCCTACAATCTGAGCACCCCTGGA
AGATCTGAACACCCGGTGGAAACTGCTGCAGGTGGCCGTGGAGGATAGAGTGAGGCAGCTG
CACGAGGCCACAGAGACTTCGGCCCTGCCTCCAGCACTTCCTGAGCACCAAGCGTGACAGG
GCCCCCTGGGAGAGAGCCATCTCCCCCAACAAAGTGCCCTACTACATCAACCACGAGACCCA
GACCACCTGCTGGGACCACCCTAAGATGACCGAGCTGTACCAGAGCCTGGCCGACCTGAAC
AATGTGCGGTTTCAGCGCTACAGAACCGCCATGAAGCTGCGGAGACTGCAGAAGGCCCTGT
GCCTGGACCTGCTGAGCCTGAGCGCCGCCTGCGACGCCCTGGACCAGCACAACTGAAGCA
GAACGACCAGCCCATGGACATTCTGCAGATCATCAACTGCCTGACCACCATCTACGATCGG
CTGGAGCAGGAGCACAAACCTGGTGAACGTGCCCTGTGCGTGGACATGTGCCTGAATT
GGCTGCTGAACGTGTACGACACCGGCAGGACCGGCAGAATCAGAGTGCTGTCTTCAAGAC
CGGCATCATCAGCCTGTGCAAGGCCACCTGGAGGATAAGTACCGCTACCTGTTCAAGCAG
GTGGCCAGCAGCACCGGCTTCTGCGATCAGAGGAGACTGGGCCTGCTGCTGCACGATAGCA
TCCAGATCCCTAGGCAGCTGGGCGAAGTGGCCAGCTTTGGCGGCAGCAACATCGAGCCCTC
TGTGAGGAGCTGCTTCCAGTTCGCCAACAACAAGCCCGAGATCGAGGCCGCCCTGTTCCCTG
GATTGGATGAGGCTGGAGCCCCAGAGCATGGTGTGGCTGCCTGTGCTGCACAGAGTGGCCG
CCGCCGAGACCGCCAAGCACCAGGCCAAGTGCAACATCTGCAAGGAGTGCCCCATCATCGG
CTTCCGGTACAGGAGCCTGAAGCACTTCAACTACGACATCTGCCAGAGCTGCTTTTTTCAGC
GGCAGAGTGGCCAAGGGCCACAAGATGCACTACCCCATGGTGGAGTACTGCACCCCCACCA
CCTCCGGCGAGGATGTGAGAGACTTCGCCAAAGTGCTGAAGAATAAGTTCCGGACCAAGCG
GTACTTTGCCAAGCACCCCAAGGATGGGCTACCTGCCCCGTGCAGACCGTGCTGGAGGGCGAC
AACATGGAGACCGACACCATGTGATGATGACTCGAGAATAAAAGATCTTTATTTTCATTAG
ATCTGTGTGTTGGTTTTTTGTGTGAGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCTCT
GCGCGCTCGCTCGCTCACTGAGGCCGGGCGACCAAAGGTCGCCCGACGCCCGGGCTTTGCC
CGGGCGGCCTCAGTGAGCGAGCGAGCGCGCAGCTGCCTGCAGGGGCGCCTGATGCGGTATT
TTCTCCTTACGCATCTGTGCGGTATTTACACCCGCATACGTCAAAGCAACCATAGTACGCG
CCCTGTAGCGGCGCATTAAGCGCGGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACAC
TTGCCAGCGCCCTAGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCCTTTCTCGCCACGTTGCG

CGGCTTTCCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTA
CGGCACCTCGACCCCCAAAAAATTGATTTGGGTGATGGTTCACGTAGTGGGCCATCGCCCT
GATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGA CTCTTGTT
CCAAACTGGAACAACACTCAACCCTATCTCGGGCTATTCTTTTGATTTATAAGGGATTTTG
CCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTTTA
ACAAAATATTAACGTTTACAATTTTATGGTGCACCTCTCAGTACAATCTGCTCTGATGCCGC
ATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGCTCTG
CTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCTCAGAGGT
TTTCACCGTCATCACCGAAACGCGCGAGACGAAAGGGCCTCGTGATACGCCTATTTTTATA
GGTTAATGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTTCGGGGAAATGTG
CGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGAC
AATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTT
CCGTGTCGCCCTTATTCCTTTTTTTCGGGCATTTTGCCTTCCTGTTTTTGTCTACCCAGAA
ACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCGAAC
TGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTTCGCCCCGAAGAACGTTTTTCCAATGAT
GAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCCGGGCAAGAG
CAACTCGGTTCGCCGCATACACTATTCTCAGAATGACTTG GTTGAGTACTACCAGTCACAG
AAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAG
TGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCT
TTTTTGCACAACATGGGGGATCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAATG
AAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGCG
CAAATATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATG
GAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTG
CTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGA
TGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGAGTCAGGCAACTATGGATGAA
CGAAATAGACAGATCGCTGAGATAGGTGCCCTCACTGATTAAGCATTGGTA ACTGTCTCAGACC
AAGTTTACTCATATATACTTTAGATTGATTTAAAACTTCATTTTTTAATTTAAAAGGATCTA
GGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTTCGTTCCAC
TGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCG
TAATCTGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTTGTTTGCCGGATCA
AGAGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGATACCAAATACT
GTCCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACAT
ACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTAC
CGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGCTGAACGGGGGGT
TCGTGCACACAGCCCAGCTTGAGAGCAACGACCTACACCGAACTGAGATACCTACAGCGTG
AGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGG
CAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTAT
AGTCCTGTGCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTTGTGATGCTCGTCAGGGG
GGCGGAGCCTATGAAAAACGCCAGCAACGCGGCCCTTTTTACGGTTCTCTGGCCTTTTGCTG
GCCTTTTGCTCACATGT

FIG. 16

C



30/38

FIG. 16**D**

SEQ ID NO:45

CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCCGGGCGTCG
GGCGACCTTTGGTCGCCCCGGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAAC
TCCATCACTAGGGGTTCCTCGTACGTTCTGAGTCCTCTAAGGTCCCTCACTCCCAACTCAG
CCCCATGTCTGTCAATTCCCACCTCAGTGTCTGATCTCCTTCTCCTCACCTTTCCCATCTC
CCGTTTGACCCAAGCTTCCTGAGCTCTCCTCCCATTCCCCTTTTTGGAGTCCTCCTCCTCT
CCCAGAACCCAGTAATAAGTGGGCTCCTCCCTGGCCTGGACCCCCGTGGTAACCTATAAG
GCGAGGCAGCTGCTGTCTGAGGCAGGGAGGGGCTGGTGTGGGAGGCTAAGGGCAGCTGCTA
AGTTTAGGGTGGCTCCTTCTCTCTTCTTAGAGACAACAGGTGGCTGGGGCCTCAGTGCCCA
GAAAAGAAAATGTCTTAGAGGTATCGGCATGGGCCTGGAGGAGGGGGGACAGGGCAGGGGG
AGGCATCTTCTCAGGACATCGGGTCCTAGAGGGACCTTGCTTCCTAGCTGGGCCTTTCTCT
TCTCCTCTATAAATACCAGCTCTGGTATTTGCGCTTGGCAGCTGTTGCTGCTAGGGAGACG
GCTGGCTTGACATGCATCTCCTGACAAAACACAAACCCGTGGTGTGAGTGGGTGTGGGCGG
TGTGAGTAGGGGGATGAATCAGAGAGGGGGCGAGGGAGACAGGGGCGCAGGAGTCAGGCAA
AGGCGATGCGGGGGTGC GACTACACGCAGTTGGAAACAGTCGTCAGAAGATTCTGGAAACT
ATCTTGCTGGCTATAAACTTGAGGGAAGCAGAAGGCCAACATTCTCCCAAGGGAAACTGA
GGCTCAGAGTTAAAACCCAGGTATCAGTGATATGCATGTGCCCCGGCCAGGGTCACTCTCT
GACTAACCGGTACCTACCCTACAGGCCTACCTAGAGACTCTTTTGAAAGGATGGTAGAGAC
CTGTCCGGGCTTTGCCCACAGTCGTTGGAAACCTCAGCATTTTCTAGGCAACTTGTGCGAA
TAAACACTTCGGGGGTCTTCTTGTTCATTCCAATAACCTAAAACCTCTCCTCGGAGAAA
ATAGGGGGCCTCAAACAAACGAAATTCTCTAGCCCGCTTTCCCCAGGATAAGGCAGGCATC
CAAATGGAAAAAAGGGGCCGGCGGGGGTCTCCTGTCAGCTCCTTGCCCTGTGAAACCCA
GCAGGCCTGCCTGTCTTCTGTCCTCTTGGGGCTGTCCAGGGGCGCAGGCCTCTTGCGGGGG
AGCTGGCCTCCCCGCCCCCTCGCCTGTGGCCGCCCTTTTCTGTCAGGACAGAGGGATCCT
GCAGCTGTCAGGGGAGGGGCGCCGGGGGGTGTGTCAGGAGGGCTACAAATAGTGCAGACA
GCTAAGGGGCTCCGTCACCCATCTTCACATCCACTCCAGCCGGCTGCCCGCCCGCTGCCTC
CTCTGTGCGTCCGCCAGCCAGCCTCGTCCACGCCGCCACCTCTAGAAAGAGGTAAGGGTT
TAAGGGATGGTTGGTTGGTGGGGTATTAATGTTTAATTACCTGGAGCACCTGCCTGAAATC
ACTTTTTTTTTCAGGTTGGACGCGTATGGTCCGCGCGAGGCACCAGCCGGGTGGGCTTTGCCT
CCTGCTGCTGCTGCTCTGCCAGTTCATGGAGGACCGCAGTGCCAGGCTGGGAACTGCTGG
CTCCGTCAAGCGAAGAACGGCCGCTGCCAGGTCTGTACAAGACCGAACTGAGCAAGGAGG
AGTGCTGCAGCACCGGCCGGCTGAGCACCTCGTGGACCGAGGAGGACGTGAATGACAACAC
ACTCTTCAAGTGGATGATTTTCAACGGGGGCGCCCCCAACTGCATCCCCTGTAAAGAAACG
TGTGAGAACGTGGACTGTGGACCTGGGAAAAAATGCCGAATGAACAAGAAGAACAAACCCC
GCTGCGTCTGCGCCCCGGATTGTTCCAACATCACCTGGAAGGGTCCAGTCTGCGGGCTGGA
TGGGAAAACCTACCGCAATGAATGTGCACTCCTAAAGGCAAGATGTAAAGAGCAGCCAGAA
CTGGAAGTCCAGTACCAAGGCAGATGTAAAAAGACTTGTGCGGATGTTTTCTGTCCAGGCA
GCTCCACATGTGTGGTGGACCAGACCAATAATGCCTACTGTGTGACCTGTAATCGGATTTG
CCCAGAGCCTGCTTCTCTGAGCAATATCTCTGTGGGAATGATGGAGTCACCTACTCCAGT
GCCTGCCACCTGAGAAAGGCTACCTGCCTGCTGGGCAGATCTATTGGATTAGCCTATGAGG
GAAAGTGTATCAAAGCAAAGTCCTGTGAAGATATCCAGTGCCTGGTGGGAAAAAATGTTT
ATGGGATTTCAAGGTTGGGAGAGGCCGGTGTTCCTCTGTGATGAGCTGTGCCCTGACAGT
AAGTCGGATGAGCCTGTCTGTGCCAGTGACAATGCCACTTATGCCAGCGAGTGTGCCATGA
AGGAAGCTGCCTGCTCCTCAGGTGTGCTACTGGAAGTAAAGCACTCCGGATCTTGCAACTC
CATTTTCGGAAGACACCGAGGAAGAGGAGGAAGATGAAGACCAGGACTACAGCTTTCCTATA

TCTTCTATTCTAGAGTGGGTTCGAGAGGTCCGGCGGCGGAGAGGGCAGAGGAAGTCTTCTAA
CATGCGGTGACGTGGAGGAGAATCCCGGCCCAATGGAAGATGCCAAAAACATTAAGAAGGG
CCCAGCGCCATTCTACCCACTCGAAGACGGGACCGCCGGCGAGCAGCTGCACAAAGCCATG
AAGCGCTACGCCCTGGTGCCCGGCACCATCGCCTTTACCGACGCACATATCGAGGTGGACA
TTACCTACGCCGAGTACTTCGAGATGAGCGTTCGGCTGGCAGAAGCTATGAAGCGCTATGG
GCTGAATAACAAACCATCGGATCGTGGTGTGCAGCGAGAATAGCTTGCAGTTCTTCATGCC
GTGTTGGGTGCCCTGTTTCATCGGTGTGGCTGTGGCCCCAGCTAACGACATCTACAACGAGC
GCGAGCTGCTGAACAGCATGGGCATCAGCCAGCCCACCGTCGTATTTCGTGAGCAAGAAAGG
GCTGCAAAAGATCCTCAACGTGCAAAAGAAGCTACCGATCATACAAAAGATCATCATCATG
GATAGCAAGACCGACTACCAGGGCTTCCAAAGCATGTACACCTTCGTGACTTCCCATTTCG
CACCCGGCTTCAACGAGTACGACTTCGTGCCCCGAGAGCTTCGACCGGGACAAAACCATCGC
CCTGATCATGAACAGTAGTGGCAGTACCGGATTGCCCAAGGGCGTAGCCCTACCGCACCGC
ACCGCTTGTGTCCGATTTCAGTCATGCCCCGCGACCCCATCTTCGGCAACCAGATCATCCCCG
ACACCGCTATCCTCAGCGTGGTGCCATTTACACCGGCTTCGGCATGTTTCACACGCTGGG
CTACTTGATCTGCGGCTTTTCGGGTGCTGCTCATGTACCGCTTCGAGGAGGAGCTATTCTTG
CGCAGCTTGCAAGACTATAAGATTCAATCTGCCCTGCTGGTGCCACACTATTTAGCTTCT
TCGCTAAGAGCACTCTCATCGACAAGTACGACCTAAGCAACTTGCACGAGATCGCCAGCGG
CGGGGCGCCGCTCAGCAAGGAGGTAGGTGAGGCCGTGGCCAAACGCTTCCACCTACCAGGC
ATCCGCCAGGGCTACGGCCTGACAGAAACAACCAGCGCCATTCTGATCACCCCCGAAGGGG
ACGACAAGCCTGGCGCAGTAGGCAAGGTGGTGCCCTTCTTCGAGGCTAAGGTGGTGGACTT
GGACACCGGTAAGACACTGGGTGTGAACCAGCGCGGCGAGCTGTGCGTCCGTGGCCCCATG
ATCATGAGCGGCTACGTTAACAACCCCGAGGCTACAAACGCTCTCATCGACAAGGACGGCT
GGCTGCACAGCGGCGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTGGAACG
GCTGAAGAGCCTGATCAAATACAAGGGCTACCAGGTAGCCCCAGCCGAACCTGGAGAGCATC
CTGCTGCAACACCCCCAACATCTTCGACGCGGGGTCGCCGGCCTGCCCGACGACGATGCCG
GCGAGCTGCCCGCCGCGAGTCGTGCTGCTGGAACACGGTAAAACCATGACCGAGAAGGAGAT
CGTGGACTATGTGGCCAGCCAGGTTACAACCGCCAAGAAGCTGCGCGGTGGTGTGTGTTT
GTGGACGAGGTGCCTAAAGGACTGACCGGCAAGTTGGACGCCCCGAAGATCCGCGAGATT
TCATTAAGGCCAAGAAGGGCGGCAAGATCGCCGTGTAAAATAAAAGATCTTTATTTTCATT
AGATCTGTGTGTTGGTTTTTGTGTGAGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCT
CTGCGCGCTCGCTCGCTCACTGAGGCCGGGCGACCAAAGGTGCCCCGACGCCCGGGCTTTG
CCCGGGCGGCCTCAGTGAGCGAGCGAGCGCGCAGCTGCCTGCAGGGGCGCCTGATGCGGTA
TTTTCTCCTTACGCATCTGTGCGGTATTTACACCCGCATACGTCAAAGCAACCATAGTACG
CGCCCTGTAGCGGCGCATTAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTAC
ACTTGCCAGCGCCCTAGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCTTTCTCGCCACGTT
GCCGGCTTTCCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTT
TACGGCACCTCGACCCCCAAAAAATGATTTGGGTGATGGTTACGTAGTGGGCCATCGCC
CTGATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTG
TTCCAAACTGGAACAACACTCAACCCTATCTCGGGCTATTCTTTTGATTTATAAGGGATTT
TGCCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTT
TAACAAAAATATTAACGTTTACAATTTTATGGTGCACCTCTCAGTACAATCTGCTCTGATGCC
GCATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTC
TGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCAGAG
GTTTTACCCGTCATCACCGAAACGCGCGAGACGAAAGGGCCTCGTGATACGCCTATTTTTA
TAGGTTAATGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATG
TGCGCGGAACCCCTATTTGTTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAG
ACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACAT
TTCCGTGTCGCCCTTATTCCTTTTTTTCGGGCATTTTGCCTTCTGTTTTTGTCTACCCAG
AAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGA

ACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATG
ATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCCGGGCAAG
AGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCAC
AGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATG
AGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCG
CTTTTTTGCACAACATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAA
TGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTG
CGCAAACCTATTAAGTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGA
TGGAGGCGGATAAAGTTGCAGGACCACCTTCTGCGCTCGGCCCTTCGGGCTGGCTGGTTTAT
TGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCA
GATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATG
AACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAAGCATTGGTAACTGTCAGA
CCAAGTTTACTCATATATACTTTAGATTGATTTAAACTTCATTTTTAATTTAAAAGGATC
TAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCTGTTCC
ACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCG
CGTAATCTGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGAT
CAAGAGCTACCAACTCTTTTTCCGAAGGTAACTGGCTTCAGCAGAGCGCAGATACCAAATA
CTGTCCTTCTAGTGCTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTAC
ATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTT
ACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTCTGGGCTGAACGGGGG
GTTTCGTGCACACAGCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCG
TGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGC
GGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTT
ATAGTCCTGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGG
GGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGC
TGGCCTTTTGCTCACATGT

33/38

FIG. 17






FIG. 18

A




B




35/38

FIG. 19

A




B



36/38

FIG. 19


C




37/38

FIG. 20


a




b




c



d



e



38/38

FIG. 21

