



US 20070128184A1

(19) **United States**

(12) **Patent Application Publication**

Podack et al.

(10) **Pub. No.: US 2007/0128184 A1**

(43) **Pub. Date:**

Jun. 7, 2007

(54) **IMMUNOMODULATING TUMOR NECROSIS FACTOR RECEPTOR 25 (TNFR25) AGONISTS, ANTAGONISTS AND IMMUNOTOXINS**

(76) Inventors: **Eckhard Podack**, Miami, FL (US); **Vadim Deyev**, Miami, FL (US); **Robert Levy**, Cooper City, FL (US)

Correspondence Address:

MCDERMOTT WILL & EMERY LLP
600 13TH STREET, N.W.
WASHINGTON, DC 20005-3096 (US)

(21) Appl. No.: **11/512,412**

(22) Filed: **Aug. 30, 2006**

Related U.S. Application Data

(60) Provisional application No. 60/712,084, filed on Aug. 30, 2005.

Publication Classification

(51) **Int. Cl.**

A61K 39/395 (2006.01)

C07K 16/46 (2006.01)

C07K 16/28 (2006.01)

(52) **U.S. Cl. 424/143.1; 530/388.22; 424/178.1;**
530/391.1

(57)

ABSTRACT

It is an object of the invention to provide novel compositions and methods utilizing immunomodulating agents that can either stimulate or indirectly augment the immune system or in other cases have an immunosuppressive effect. TNFR25 agonists disclosed herein have an anti-inflammatory and healing effect. They can be used, among other things, to treat disease caused by asthma and chronic inflammation such as for example inflammatory bowel diseases including ulcerative colitis and Crohn's Disease. TNFR25 antagonists disclosed herein are capable of inhibiting CD8 T cell-mediated cellular immune responses and can for example, mitigate organ or tissue rejection following a tissue transplantation. TNFR25 agonists disclosed herein represent biological response modifiers that alter the interaction between the body's cellular immune defenses and cancer cells to boost, direct, or restore the body's ability to fight the cancer when given with tumor vaccines. TNFR25 specific immunotoxins disclosed herein are also capable of increasing the effectiveness of a chemotherapeutic regimen by depleting a cancer patient of naturally occurring immunosuppressive cells.




Figure 1a**Figure 1b****Figure 1c**




Figure 2a**Figure 2b****Figure 2c**


Figure 3a




Figure 3b**Figure 3c****Figure 3d**




Figure 3e**Figure 3f****Figure 3g**


Figure 4a





Figure 4b**Figure 4c****Figure 4d****Figure 4e**






Figure 5a**Figure 5b****Figure 5c****Figure 5d****Figure 5e**





Figure 6a**Figure 6b****Figure 6c****Figure 6d**





Figure 6e**Figure 6f****Figure 6g****Figure 6h**







Figure 7a**Figure 7b****Figure 7c**

Figure 7d**Figure 7e**

control IgG anti TL1A, L4G6



Figure 7f**Figure 7g**




Figure 7h**Figure 7i****Figure 7j**


Figure 8


Figure 9

Figure 10




Figure 11

Enhanced OT-I Expansion by TNFR25
Costimulation





Figure 12


Figure 13

Figure 14

Requirement of TNF25 for recovery from Colitis





Figure 15


Figure 16

Figure 17

SEQ ID NO: 1 - Mouse TNFER25 cDNA; NM_033042

1	gggagttgtt	ctggatggcg	cggggggcggg	cgggcagcag	ctactctagt	ctaggaacat
61	aggggctgag	ctgggtgggg	aagccccggg	ttacgcgacc	gaccagagcc	gcactcacaa
121	gggcccaggc	ggtacacacc	gcaatggagg	cacggctgtc	gcggggctgc	gtggtggagc
181	ctctgttccct	accactgctg	ctgctgtcgc	tgctgtctgt	gctgtcttgg	ggcaggggcc
241	agggcggcat	gtctggcagg	tgtactgtg	ccagtggatc	ccagaagagg	tatggccctgt
301	tttggttcag	gggctgccc	aaggggacact	acatgaaggc	ccctcgccca	gaaccctgtg
361	gcaactccac	ctgccttccc	tgtccctcgg	acaccttctt	gaccagagac	aaccactta
421	agactgactg	tacccgtcgc	caagtctgtg	atgaagaggc	ccttcaagtg	acccttgaga
481	actgctcgcc	aaagtccggac	acccactgtg	gctgcccggc	aggctgggt	gttactgtct
541	ccaccgtgcc	atgtgggaaa	agctcacctt	tctttgtgt	cccatgggg	gctacgacac
601	cagtccatga	ggctccaacc	ccccggccct	gcctgcctgg	cttctatata	cgtggcaatg
661	actgcacgtc	ctgccccacg	ggcttcagca	gcgtttgccc	taaggcttgc	actgtgtct
721	gtggctggaa	gcagatgttt	tgggtccagg	tgcttctagg	agtcgcgttc	ctttttgggg
781	ctatcctgat	ctgtcataat	tgtcgtatggc	agccttgtaa	ggccgtggtc	actgcagaca
841	cagctggac	ggagccccctg	gcctcaccac	agactgccc	tctctcagcc	tcagacagcg
901	cccacacccct	cttggcacct	ccaagcaaga	ctggggaaat	ctgttacact	gtccagttgg
961	taggcaacaa	ctggaccctt	ggcttacccc	agactcagga	gttggcttgc	ggacaggcct
1021	cacaacccctg	ggatcagctg	ccaaacacaaa	cttttggaaac	tctcttgca	tctccgtct
1081	cgccagcgcc	ccctgcgggc	tctccggctg	ctgtgttcc	gcctggcccg	cagcttacg
1141	atgtatgg	tgcggtcccc	gcacgaagg	ggaggagtt	ctgtgcacg	ctggggctgc
1201	gggaagcgga	aattgaagcc	gtggagggtt	aaatctgcgg	cttccgagac	cagcgtatg
1261	agatgctcaa	gctgtggcgt	cagcagcagc	ctgcaggcc	cggtgcccac	tatgcggctc
1321	tggagcgc	gggtctggaa	ggctgtccgg	aggacctcg	cagccgcctg	cagcgtggcc
1381	cgtgtatgg	ggtccatcat	ccacttttgc	accctagtga	cccttgaagg	agccttaagt
1441	attgttactt	atgcgtgttag	acattttatg	tcaattacta	acccctgc	gtggccttgc
1501	gtagcaggc	tggctgcctc	acttttgc	atctgcagca	cggagcttct	gctaaggaa
1561	gctgtatgg	gaaataccag	aaggggccaa	gtgattggtt	gctcagctgt	taatttagccc
1621	gagtttggac	tttgtattaa	atttcataag	aaaagcagct	gcttg	

SEQ ID NO: 2 - Human TNFER25 cDNA; NM_148965

1	cgggccctgc	gggcgcgggg	ctgaaggcgg	aaccacgacg	ggcagagagc	acggagccgg
61	gaagccctg	ggcgcggcgtc	ggagggtat	ggagcagcgg	ccgcggggct	gcgcggcggt
121	ggcggccggcg	ctcttcctgg	tgctgtggg	ggcccgggcc	cagggcggca	ctcgtagccc
181	caggtgtgac	tgtgcgggt	acttccacaa	gaagattgtt	ctgttttgtt	gcagaggctg
241	cccagccggg	cactacctg	aggccccctg	cacggagccc	tgccggcaact	ccacctgcct
301	tgtgtgtccc	caagacaccc	tcttggcctg	ggagaaccac	cataattctg	aatgtccccg
361	ctgccaggcc	tgtgtatggc	aggccctccca	ggtggcgctg	gagaactgtt	cagcagtggc
421	cgacacccgc	tgtggctgt	agccaggctg	gtttgtggag	tgccaggctca	gccaatgtgt
481	cagcaggta	cccttctact	gccaaccatg	cctagactgc	ggggccctgc	accgccacac
541	acggctactc	tgttcccgca	gagatactg	ctgtgggacc	tgcctgcctg	gttcttatga
601	acatggcgat	ggctgcgtgt	cctgccccac	gcaacccccc	tcccttgca	gagcaccctg
661	gggagctgtc	cagagcgtc	tgccgcgtc	tgtggcttgg	ggcagatgtt	gtgtgttctg
721	gttccagggt	ctcttggctg	gccttgggtt	cccccttcgt	cttggggcca	ccctgaccta
781	cacataccgc	cactgtgtgc	ctcacaagcc	cctgggtact	gcagatgtt	ctgggatgg
841	ggctctgacc	ccaccaccgg	ccacccatct	gtcacccttg	gacagcccc	acacccttct
901	agcacccct	gacagcgtg	agaagatctg	caccgtccag	ttggtgggtt	acagctggac
961	ccctggctac	cccgagaccc	aggaggcgct	ctgcccgcag	gtgacatgtt	cctgggacca
1021	gttgcggc	agagcttgc	gccccgtgc	tgcgtccaca	ctctcgccag	agtccccagc
1081	cggctcgcc	gccatgtatgc	tgcagccggg	cccgagctc	tacgacgtt	tggacgcgg
1141	cccaaggccgg	cgtgttgcagg	atttcgtgc	cacgtgggg	ctgcgcgagg	cagagatcg
1201	agccgtggag	gtggagatcg	gccgttccc	agaccagcag	tacgagatgc	tcaagcgctg
1261	gcccacgc	cagcccgccgg	gcctcgagc	cgtttacgc	gccctggagc	gcatggggct

Figure 17 Continue

1321	ggacggctgc	gtgaaagact	tgcgccagccg	cctgcagcgc	ggcccggtac	acggcgccca
1381	cttgcacact	aggcgtctg	gtggcccttg	cagaagccct	aagtacgtt	acttatgcgt
1441	gtagacattt	tatgtcaactt	attaagccgc	tggcacggcc	ctgcgttagca	gcaccagccg
1501	gcccccacccc	tgctcgcccc	tatcgctcca	gccaaggcga	agaagcacga	acgaatgtcg
1561	agaggggggtg	aagacatttc	tcaacttctc	ggccggagtt	tggctgagat	cgcggattta
1621	aatctgtgaa	aaaaaaacaaa	acaaaaacaaa	aaaaaaaaaa	aaaaaa	

SEQ ID NO: 3 - Mouse TL1A cDNA; AF520786

1	atggcagagg	agctggggtt	gggcttcgga	gaaggagttcc	cagtggaaagt	gctggccggaa
61	ggctgttagac	acaggccaga	ggccaggggcc	gggcttagctg	ccaggagcaa	agcctgcctg
121	gctctcacct	gtgcctgtt	gtcatttccc	atcctcgag	gacttagcac	cctccctaattg
181	gtggccagc	tccgggtccc	cgaaaagac	tgtatgttc	gggcccataac	agaagagaga
241	tctgagccctt	caccacagca	agtttactca	cctcccaagag	gcaagccgag	agcacacctg
301	acaattaaga	aacaaacccc	agcaccacat	ctgaaaaatc	agctctctgc	tctacacttgg
361	gaacatgacc	tagggatggc	cttcaccaag	aacgggatga	agtacatcaa	caaatccctg
421	gtgatcccaag	agtcaaggaga	ctatttcatc	tactcccaga	tcacattccg	agggaccacaca
481	tctgtgtgtg	gtgacatcg	tccggggaga	cgaccacaaaca	agccagactc	catcaactgtg
541	gtcatcacca	aggttagcaga	cagctaccct	gagcctgccc	gcctactaaac	agggtccaaag
601	tctgtgtgtg	aaataagcaa	caactggttc	cagttcccttct	accttggggc	catgttctcc
661	ttggaagaag	gggacagact	aatggtaaac	gtcagtgaca	tctccttgggt	ggattacaca
721	aaagaagata	aaactttctt	tggagctttc	ttgctataa		

SEQ ID NO: 4 - Mouse DN TNFR25 cDNA

atggagg cacggctgct gcggggctgc gtgggtggagc
ctctgttcct accactgctg ctgctgctgc tctgtctgct gctgcttggt ggccaggggcc
agggcggcat gtctggcagg tgtgactgtg ccagttagtc ccagaaggagg tatggcccgat
tttgttgcag gggctgccc aagggacact acatgaaggc cccctgcgcga aaccctgtg
gcaactccac ctgccttccc tgccttcgg acacccctt gaccagagac aaccacttta
agactgactg taccggctgc caagtctgtg atgaagaggg ccttcaagtg acccttgaga
actgctcggc aaagtggac acccaactgtg gctgcccagtc aggctggtgt gttgactgct
ccaccgtgcc atgtggggaaa agctcacctt tctcttgggt cccatgcgggg gctacgacac
cagttccatga ggctccaaacc ccccgccct gcctgcctgg cttctatata cgtggcaatg
actgcacgtc ctggcccaacg ggcttcagca gcgtttgccc taaggcttgc actgctgtct
gtggctggaa gcagatgttt tgggtccagg tgcttctagg agtcgcgttc cttttgggg
ctatccatgtat ctgtgcataat

SEQ ID NO: 5 - Human DN TNFR25 cDNA (extracellular + transmembrane only)

atggagcagcgg ccgcggggct gcgcggcggt
ggcggcggcg ctccctcctgg tgctgctggg ggcccgccg cagggcgca ctcgtagccc
cagggtgtac tgtgcgggtg acttccacaa gaagatttgt ctgttttgtt gcagaggctg
cccagcgggg cactaccta agggcccttg cacggagccc tgccgcaact ccacactgcct
tgtgtgtccc caagacacct tcttggcctg ggagaaccac cataattctg aatgtcccg
ctgccaggcc tgtgtatgagc aggcctccca ggtggcgctg gagaactgtt cagcagtggc
cgacacccgc tgtggctgta agccaggctg gtttggag tgccaggctca gccaatgtgt
cagcagttca cccttctact gccaaccatg cctagactgc ggggcccctgc accgcccacac
acggctactc tgttcccgcgca gagatactga ctgtgggacc tgccctgcctg gcttctatg
acatggcgat ggctgcgtgt cctgccccac gccacccccc tcccttgca gggcaccctg
gggagctgtc cagagcgctg tgccgctgtc tgtggctgga ggcagagtag gtgtgttctg
ggtccaggtg ctccctggctg gccttgttgt ccccttcctg ctggggcca ccctgaccta
caca

Figure 17 Continue

SEQ ID NO: 6 - Human DN TNFR25 cDNA (extracellular only)

```

at ggagcagcgg ccgcggggct gcgcggcgg
ggcggcggcg ctccctcgg tgctgctggg ggcggggcc cagggcggca ctcgtagccc
cagggtgtac tgtccgggtg acttccacaa gaagattggt ctgtttgtt gcagaggctg
cccagcgggg cactacactga aggccccttg cacggagccc tgccgcaact ccacctgcct
tgtgtgtccc caagacacct tcttggctg ggagaaccac cataattctg aatgtccccg
ctgccaggcc tgtgtatgac aggcctccca ggtggcgctg gagaactgtt cagcagtggc
cgacaccgcg tggctgtta agccaggctg gtttggag tgccaggtca gccaatgtgt
cagcagtctca cccttctact gccaaccatg cctagactgc gggccctgc accgccccac
acggctactc tgccccgca gagatactga ctgtgggacc tgccctgcctg gcttctatga
acatggcgat ggctgcgtgt cctgccccac gccacccccc tcccttgcag gaggcaccctg
gggagctgtc cagagcgtgtc tgccgctgtc tgtggctgga ggcagagtag gtgtgttctg
g

```

SEQ ID NO: 7 - Human TL1A; cDNA; AF520785

```

1 gagagggaaa agggaaaggag gagactgagt gattaagtca cccactgtga agagctggc
61 ttcttatttaa tggggggctct ctctgcccag gagtccagg tgcctccagg aacccggcgg
121 gcatggccga ggatctggga ctgagctttg gggaaacacgc cagtgtggaa atgtgccag
181 agcacggcag ctgcaggccc aaggccagga gcagcagcgc acgctggct ctcacctgt
241 gcctgggttt gctcccccctc cttgcaggac tcaccacata cctgctgtc aacccggcgtcc
301 gggcccaggag agaggccctgt gtgcagttcc aggccttaaa aggacaggag tttgcaccc
361 cacatcagca agtttatgca cctcttagag cagacggaga taagccaagg gcacacctga
421 cagttgtgag acaaactccc acacagcact tttttttatca gttcccaact ctgcactggg
481 aacatgaact aggcctggcc ttccaccaaga accgaatgaa ctataccaac aaattccctgc
541 tgatcccaga gtcgggagac tacttcattt actcccaagg cacattccgt gggatgaccc
601 ctgagtgca gtaaatcaga caagcaggcc gacccaaacaa gcccacactcc atcaactgtgg
661 tcatcacccaa ggttaacagac agtacccctg agccaaacccca gtcctccatg gggaccaagt
721 ctgtatgcga agtaggttagc aactgttcc agcccatcta cctcggagcc atttctctt
781 tgcaagaagg ggacaagcta atggtaacg tcagtgcacat ctctttggg gattacacaa
841 aagaagataa aacccctttt ggagccctt tactatagga ggagagcaaa tatcattata
901 tgaaaagtccct ctgccaccga gttccctaatt ttctttgtt aaatgttaatt ataaaccagg
961 gttttcttgg ggccgggagt agggggcatt ccacaggac aacggtttag ctatgaaatt
1021 tggggcccaa aatttcacac ttcatgtgc ttactgatga ggtactaac tggaaaaggc
1081 tgaagagagc aaatataattt ttaagatggg ttggaggatt ggcgagttc taaatattaa
1141 gacactgatc actaaatgaa tggatgatct actcgggtca ggattgaaag agaaatattt
1201 caacacctcc ctgtatatac atggtcacca gtggtccagt tattgttcaa ttgtatcata
1261 aatttgcctt aattcaggag ctttgaagga agtccaaagg aacccggcgtcc ttttgcgttt
1321 aacttcaga ggcacccatcc ttcaccaatt ttccacata ctttgcgtttt tggccctaaaa
1381 aaaatgaaaa gagagtttgt atgttcatg aatgttccaca cagaaggagt tggggggat
1441 gtcatctaca gcatatgaga aaagctaccc ttctttgtat tatgtacaca gatatctaaa
1501 taaggaagta tgagtttccat atgtatatac aaaatacaac agttgcgttgc attcagtaga
1561 gttttcttgc ccacccatcc ttttttttttgc ttttttttttgc cccagaagac actatgaaaa
1621 acaagacaga ctccactcaa aatttatatg aacaccacta gatactccct gatcaaacat
1681 cagtcacat actctaaaga ataactccaa gtcttggccca ggcgcgtgg ctcacacctg
1741 taatcccaac actttgggag gccaaggatgg ttggatcatc taaggccggg agtcaagac
1801 cagccgtacc aacgtggaga aaccccatct ctactaaaaa tacaaaattt gcccggcggt
1861 gtacgcgtatc gctgtatcc tggctactca ggaggccggag gcagaagaat tgcttgcact
1921 ggggaggcag aggttgcgtt gagcccatgt cgccgttgc cactccagcc tgggttaacaa
1981 gagcaaaact ctgtccaaaa aaaaaaaaaa aaaaaaaaaa

```

SEQ ID NO: 16 - mDN-TNFR25 cDNA

```

ATGGAGGCAC GGCTGCTGCG GGGCTGCGTG GTGGAGCCTC TGTTCTTACCC ACTGCTGCTG
CTGCTGCTGC TGCTGCTTGG TGGCCAGGGC CAGGGCGGCA TGTCTGGCAG GTGTGACTGT
GCCAGTGAGT CCCAGAAGAG GTATGGCCCG TTTTGTGCA GGGGCTGCC AAAGGGACAC

```

Figure 17 Continue

TACATGAAGG	CCCCCTGCGC	AGAACCCCTGT	GGCAACTCCA	CCTGCCTTCC	CTGTCCCTCG
GACACCTTCT	TGACCAAGAGA	CAACCACTTT	AAGACTGACT	GTACCCGCTG	CCAAGTCTGT
GATGAAGAGG	CCCTTCAAGT	GACCCCTTGAG	AACTGCTCGG	CAAAGTCGGA	CACCCACTGT
GGCTGCCAGT	CAGGCTGGTG	TGTTGACTGC	TCCACCGAGC	CATGTGGGAA	AAGCTCACCT
TTCTCTTGTG	TCCCATGCGG	GGCTACAAACA	CCAGTCCATG	AGGCTCCAAC	CCCCCGGGCC
TGCCTGCCGT	GCTTCTATAT	ACGTGGCAAT	GACTGCACGT	CCTGCCCCAC	GGGCTTCAGC
AGCGTTTGCC	CTAAGGCTTG	CACTGCTGTC	TGTGGCTGGA	AGCAGATGTT	TTGGGTCCAG
GTGCTTCTAG	GAGTCGCGTT	CCTTTTGGG	GCTATCCTGA	TCTGTGCATA	TGGCTCTGGA
AGCGGGAGCG	TCGACGATCT	TTACGACGAT	GATAAAATAGT	AA	

IMMUNOMODULATING TUMOR NECROSIS FACTOR RECEPTOR 25 (TNFR25) AGONISTS, ANTAGONISTS AND IMMUNOTOXINS**PRIORITY**

[0001] This application claims priority to U.S. Ser. No. 60/712,084, filed Aug. 30, 2005, which is incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION**[0002] 1. Field of the Invention**

[0003] This invention relates Tumor Necrosis Factor Receptor Super Family 25 (TNFR25) agonists, immunotoxins, antagonists and their use in treating cancer, inflammation and effecting immunosuppression, respectively.

[0004] 2. Background

[0005] Many disorders of the human immune system fall into two broad categories: those characterized by an attenuated immune response and those characterized by overzealous immune responses. Immunodeficiency is characterized by an attenuated response. There are congenital (inborn) and acquired forms of immune deficiency. Chronic granulomatous disease, in which phagocytes have trouble destroying pathogens, is an example of the former. AIDS ("Acquired Immune Deficiency Syndrome"), an infectious disease, caused by the HIV virus that destroys CD4+ T cells, is an example of the latter. An additional disease that may be characterized by an attenuated immune response is cancer. In contrast to healthy individuals, cancer patients' immune systems are no longer capable of effectively recognizing and/or destroying tumor cells.

[0006] Despite high hopes, there are no medications to date that directly increase the activity of the immune system. However, biological therapies have recently been used to recruit the immune system, either directly or indirectly, to fight diseases such as cancer. Monoclonal (MAb) antibodies are now frequently used as a biologic therapy. For example, monoclonal antibodies may react with specific types of cancer cells, and have direct or indirect antitumor effects.

[0007] Tumor vaccines may be employed therapeutically or for prophylaxis after primary therapy. Anti-tumor vaccines may need to induce cellular immunity in the form of tumor-specific cytotoxic T cells of the CD4 or CD8 phenotype. It is thought that effective anti-tumor immunity requires the generation and maintenance for long periods of times of such cytotoxic cells. In addition evidence indicates that the innate arm of the immune system must be activated in order to generate effective anti-tumor vaccines. Vaccines that enhance or generate humoral responses produce antibodies that can be detected over a relatively long period. To be effective, these antibodies need to be capable of targeting cell surface antigens in live cell assays. Maintaining specific cellular immune responses to antigen epitopes (adaptive immunity) may require more frequent immunizations, although memory cells can sustain the ability to respond and rechallenge the immunizing epitope. As such, it would be substantial benefit to have access to therapies that would be capable of boosting cancer specific cellular immune responses to tumor vaccines.

[0008] On the other end of the scale, an overactive immune system figures in a number of other disorders,

particularly autoimmune disorders such as lupus erythematosus, type I diabetes (sometimes called "juvenile onset diabetes"), multiple sclerosis, psoriasis, rheumatoid arthritis and inflammatory bowel diseases such as Crohn's Disease and ulcerative colitis (UC). In these, the immune system fails to properly distinguish between self and non-self and attacks a part of the patient's own body. Other examples of overzealous immune responses in disease include hypersensitivities such as allergies and asthma.

[0009] Suppression of the immune system is often used to control autoimmune disorders or inflammation when this causes excessive tissue damage. Immunosuppressive medication intentionally induces an immunodeficiency in order to prevent rejection of transplanted organs. Commonly used immunosuppressants include glucocorticoids, azathioprine, methotrexate, cyclosporin, cyclophosphamide and mercaptopurine. In organ transplants, selective T cell inhibition prevents organ rejection, and cyclosporin, tacrolimus, mycophenolate mofetil and various others are used.

[0010] T lymphocytes play a central role in regulating immune responses. Helper T cells express the CD4 surface marker and provide help to B cells for antibody production and help CD8 T cells to develop cytotoxic activity. Other CD4 T cells inhibit antibody production and cytotoxicity. T cells regulate the equilibrium between attack of infected or tumorigenic cells and tolerance to the body's cells. A dysregulated immune attack can lead to autoimmunity, while diminished immune responsiveness results in chronic infection and cancer.

[0011] Tumor Necrosis Factor Receptor 25 (TNFR25) also interchangeably referred to herein as Death receptor 3 (DR3), as discussed herein, is a regulator of T cell function. Death receptor 3 (DR3) (Chinaiyan et al., *Science* 274:990, 1996) is a member of the TNF-receptor family. It is also known as TRAMP (Bodmer et al., *Immunity* 6:79, 1997), wsl-1 (Kitson et al., *Nature* 384:372, 1996), Apo-3 (Marsters et al., *Curr Biol* 6:1669, 1996), and LARD (Screaton et al., *Proc Natl Acad Sci USA* 94:4615, 1997) and contains a typical death domain. Transfection of 293 cells with human DR3 (hDR3) induced apoptosis and activated NF- κ B. The cognate ligand for DR3 has recently been identified as TL1A (Migone et al., *Immunity* 16:479, 2002) and has been shown to have costimulatory activity for DR3 on T cells through the induction of NF- κ B and suppression of apoptosis by expression cIAP2 (Wen et al., *J Biol Chem* 25:25, 2003). TL1A also binds to the decoy receptor 3 (DcR3/FR-6), indicating that fine-tuning of biological TL1A accessibility is of critical importance. Multiple spliced forms of human DR3 mRNA have been observed, indicating regulation at the post transcriptional level (Screaton et al., *Proc Natl Acad Sci USA* 94:4615, 1997).

[0012] Many TNF-receptor family members have the ability to induce cell death by apoptosis or induce costimulatory signals for T cell function. The regulation of these opposing pathways has recently been clarified for TNF-R1, the prototypic death domain-containing receptor that can cause apoptosis or proliferation of receptor positive T cells (Micheau and Tschopp. *Cell* 114:181, 2003). NF- κ B activation by a signaling complex composed of TNF-R1 via TRADD, TRAF2 and RIP induces FLIPL association with a second signaling complex composed of TNFR1, TRADD and FADD, preventing caspase 8 activation as long as the

NF- κ B signaling persists. DR3 has been shown to be able to induce apoptosis in transfected cells and to induce NF- κ B and all three MAP-kinase pathways (Chinnaiyan et al., *Science* 274:990, 1996; Bodmer et al., *Immunity* 6:79, 1997; Kitson et al., *Nature* 384:372, 1996; Marsters et al., *Curr Biol* 6:1669, 1996; Scream et al., *Proc Natl Acad Sci USA* 94:4615, 1997; Wen et al., *J Biol Chem* 25:25, 2003). Blocking of NF- κ B, but not of MAP-kinase and inhibition of protein synthesis resulted in DR3-mediated cell death, indicating that NF- κ B signals mediate anti-apoptotic effects through the synthesis of anti-apoptotic proteins.

[0013] Expression of human DR3 mRNA is pronounced in lymphoid tissues, mainly in the spleen, lymph nodes, thymus, and small intestine, indicating an important role for DR3 in lymphocytes. Murine DR3 has been deleted by homologous recombination in embryonic stem cells (Wang et al., *Mol Cell Biol* 21:3451, 2001). DR3-/-mice show diminished negative selection by anti-CD3 in the thymus but normal negative selection by superantigens and unimpaired positive selection of thymocytes. Mature peripheral T cells were unaffected by DR3 deficiency. Despite a significant amount of preliminary research, the physiological function of DR3 remains poorly characterized.

[0014] All scientific publications including patent documents cited herein are incorporated by reference in their entirety for all purposes.

SUMMARY OF THE INVENTION

[0015] One aspect of the invention relates to an antibody that binds a Tumor Necrosis Factor Receptor 25 (TNFR25) antigen and that can act as a TNFR25 agonist. In one embodiment, the antibody is capable of increasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody. In a further embodiment, the antibody is purified monoclonal antibody 4C12.

[0016] Another aspect of this invention relates to a TNFR25-specific toxin comprising a toxic agent linked to polypeptide that binds the TNFR25 receptor. In one embodiment of this aspect, the toxin-comprising portion includes the monoclonal antibody 4C12 or an immunospecific portion of 4C12. In another embodiment, the toxic agent is selected from a radioactive isotope, ricin, abrin, diphtheria toxin, *Pseudomonas* exotoxin, or metal ion. In a further embodiment, the polypeptide that binds the TNFR25 receptor is the TL1A protein or a fragment or variant thereof.

[0017] In another aspect of the invention, the TNFR25-specific toxin is used in a method of treating cancer in a patient. Specifically, the method includes depleting a patient of CD4+/CD25+T regulatory cells (Tregs) by providing the patient with the TNFR25-specific toxin and also providing the patient with a chemotherapeutic agent.

[0018] Yet another aspect of this invention relates to a method of activating TNFR25 receptor expressed on a cell comprising contacting the cell with a TNFR25 agonist. The agonist may be selected from a monoclonal antibody 4C12; an antibody that binds TNFR25 and that can increase OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody; a soluble TL1A protein; an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25 agonist antibody; an expression vector with an expression cassette

capable of driving the transgenic expression of a soluble TL1A; or an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25. This method further includes observing an increase in TNFR25 receptor signaling.

[0019] An additional aspect of this invention relates to an antibody that is capable of acting as a TNFR25 antagonist. In one embodiment, the antibody binds a TL1A and is capable of decreasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody. In a further embodiment, the antibody is the purified monoclonal antibody L4G6.

[0020] A further aspect of the invention relates to a method of inhibiting TNFR25 receptor signaling in a cell. The method includes contacting the cell with a TNFR25 antagonist. The method further includes observing a decrease in TNFR25 receptor signaling.

[0021] Another aspect of this invention relates to a tumor vaccine comprising a tumor antigen and a TNFR25 agonist as a biological response modifier. A further embodiment of this vaccine also includes an adjuvant.

[0022] In yet another aspect of this invention, after isolating a tumor specific antigen, a vaccine comprising a tumor specific antigen and a TNFR25 agonist, is used to immunize a patient against the tumor.

[0023] A further aspect of this invention relates to a method of treating and/or preventing gut inflammation comprising providing a patient in need thereof with an effective amount of a therapeutic composition comprising a TNFR25 agonist.

[0024] A further aspect of this invention relates to a therapeutic composition for the facilitation of an organ transplant comprising a TNFR25 antagonist and an immunosuppressant. In one embodiment of this aspect, the immunosuppressant is glucocorticoid, azathioprine, methotrexate, cyclosporin, cyclophosphamide, mercaptopurine, tacrolimus or mycophenolate mofetil.

[0025] In another aspect of the invention, a TNFR25 antagonist composition is used in a method of transplanting a tissue from a donor into a host. This method includes the steps of obtaining tissue from a donor; providing a host with a TNFR25 antagonist composition; and transplanting the tissue into the host.

[0026] Another aspect of this invention relates to a method of inhibiting the clonal expansion of a population of cognate CD8 T cells. This method includes exposing the CD8 T cells to their cognate antigen and exposing the CD8 T cells to a TNFR25 antagonist. In a further embodiment, the cognate antigen is associated with tissue to be transplanted from a donor into a host.

[0027] Another aspect of the invention relates to an isolated TNFR25 antagonist comprising a polypeptide encoded by a nucleic acid comprising sequence that hybridizes under stringent conditions to SEQ ID NOs: 4, 5, 6 and/or 16, and wherein the sequence encodes an amino acid sequence capable of binding a TL1A protein. In one embodiment, the sequence hybridizes under stringent conditions to SEQ ID NO: 4. In another embodiment, the sequence hybridizes under stringent conditions to SEQ ID NO: 5. In a further embodiment, the sequence hybridizes under stringent con-

ditions to SEQ ID NO: 6. In a further embodiment, the sequence hybridizes under stringent conditions to SEQ ID NO: 16. In yet another embodiment, the TL1A is human or mouse TL1A.

[0028] Another aspect of the invention relates to a method of treating and/or preventing lung inflammation comprising providing a patient in need thereof a with an effective amount of a therapeutic composition comprising the TNFR25 antagonist comprising a polypeptide encoded by a nucleic acid comprising sequence that hybridizes under stringent conditions to SEQ ID NOs: 4, 5, 6 and/or 16, and wherein the sequence encodes an amino acid sequence capable of binding a TL1A protein.

[0029] Another aspect of the invention relates to a method of transplanting a tissue from a donor into a host comprising obtaining the tissue from the donor; providing the host with the TNFR25 antagonist comprising a polypeptide encoded by a nucleic acid comprising sequence that hybridizes under stringent conditions to SEQ ID NOs: 4, 5, 6 and/or 16, and wherein the sequence encodes an amino acid sequence capable of binding a TL1A protein; and transplanting the tissue into the host.

[0030] Another aspect of the invention relates to a composition comprising a polypeptide encoded by a sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein the sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein; and a toxic agent. In one embodiment, the toxic agent is selected from the group consisting of a radioactive isotope, ricin, abrin, diphtheria toxin, *Pseudomonas* exotoxin, and metal ion.

[0031] Another aspect of the invention relates to a method of treating cancer in a patient comprising depleting a patient of CD4+/CD25+T regulatory cells (Tregs) by providing the patient with a composition comprising the toxin of claim 39; and providing a patient with a chemotherapeutic agent.

[0032] A further aspect of the invention relates to a method of treating and/or preventing gut inflammation comprising providing a patient in need thereof a with an effective amount of a composition comprising a polypeptide encoded by a sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein the sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein. In one embodiment, the inflammation is a result of irritable bowel syndrome. In another embodiment, the gut inflammation is a result of Crohn's disease.

[0033] A further aspect of the invention relates to a tumor vaccine comprising a tumor antigen and a polypeptide encoded by a sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein the sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein, as a biological response modifier.

[0034] Still another aspect of the invention relates to an expression vector comprising a nucleic acid sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and that encodes an amino acid sequence capable of binding a TNFR25 receptor protein.

[0035] Yet a further aspect of the invention relates to an expression vector comprising a nucleic acid that hybridizes

under stringent conditions to SEQ ID NOs: 4, 5, 6 and/or 16, and wherein the sequence encodes an amino acid sequence capable of binding a TL1A protein

[0036] Additional advantages of the present invention will become readily apparent to those skilled in this art from the following detailed description, wherein only the preferred embodiment of the invention is shown and described, simply by way of illustration of the best mode contemplated of carrying out the invention. As will be realized, the invention is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the invention. The present invention may be practiced without some or all of these specific details. In other instances, well known process operations have not been described in detail, in order not to unnecessarily obscure the present invention. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

BRIEF DESCRIPTION OF THE DRAWINGS

[0037] FIG. 1A depicts the expression of murine TNFR25 in lymph node cells. MFI is indicated by black numbers for isotype control antibody as primary antibody and in shaded for anti-TNFR25. Detection of TNFR25 required a triple sandwich of primary hamster anti-TNFR25 monoclonal antibody, followed by goat anti-hamster biotin and PE-labeled Strept-Avidin. 1B depicts the expression of TNFR25 on activated lymphocytes. Activation of splenocytes was done with immobilized anti-CD3 (5 µg/ml) and anti-CD28 (1 µg/ml) or LPS (1 µg/ml) for 24 hours. Cells were gated for CD4 or CD8 or B220 positive and 7-AAD negative cells (subscript a stands for activated CD4, CD8 or B cells). In the histograms, MFI for expression of TNFR25 on resting splenocytes and on activated splenocytes is shown. 1C depicts the expression of TNFR25 on thymocytes. Thymocytes were gated on CD4/CD8 double negative, double positive or single positive cells and evaluated for anti-TNFR25 fluorescence.

[0038] FIG. 2A depicts splice forms of murine TNFR25. Splice forms were obtained by RT-PCR of mRNA obtained from resting murine splenocytes and murine cell lines. CRD: cysteine-rich domain; TM: transmembrane domain; DD: death domain. Asterisks: in frame stop codon. In splice forms 1-4 the intron between exon 2 and 3 is not spliced out and contains a premature stop codon. Splice forms 1-4 are likely to be nonfunctional proteins. Murine Δ5 and Δ6 (corresponding to human Δ6 and Δ6,7) lack a complete transmembrane domain and are predicted to be secreted forms that could act as soluble decoy receptor for TL1A. Δ5,6 (potentially corresponding to human A3) and FL splice forms are studied as transgenes in this report. Δ5,6,8 and Δ5,6,9 (no human homologues) are predicted to be membrane anchored but lack the death domain and may have altered signaling properties or may act as dominant negative splice forms. The preferred DN TNFR25 disclosed herein was truncated after the TM domain. 2B depicts activation-induced alternative splicing of TNFR25. Both mouse and human TNFR25 are spliced after activation. The human splicing is shown here because splice forms of TNFR25 are separated better in size after gel electrophoresis than murine forms. Splice forms were confirmed by sequencing. PBL were isolated by Ficoll-Hypaque gradient centrifugation. Five million cells were used in each sample and mRNAs

were extracted and converted to cDNA using the Invitrogen cDNA synthesis kit. Activation of PBL by PHA (5 µg/ml), immobilized anti-CD3 (5 µg/ml) and soluble anti-CD28 (1 µg/ml), or PMA (10 ng/ml) and ionomycin (400 ng/ml) as indicated. The cells were harvested at the indicated time points and RT-PCRs were performed; β-actin was used as the internal control. 2C depicts activation-induced splicing is PKC-dependent and protein synthesis-independent. Freshly isolated PBL cells were stimulated with PMA (10 ng/ml) alone, or ionomycin (400 ng/ml) alone or in combination. PBL were pretreated with H7 (50 µM), or cycloheximide (10 µg/ml) for half an hour, then PMA and ionomycin were added into the cell culture. The cells were harvested after 12 hours for RT-PCR analysis.

[0039] FIG. 3 illustrates expression and function of TNFR25-transgenes under the CD2 promoter and enhancer. 3A depicts expression of TNFR25 in transgenic mice compared to B6 wild-type mice and isotype control. Inguinal lymph node cells were gated on CD4, CD8, B220, CD11c positive cells, or NK1.1 positive and CD3 negative cells or NK1.1/CD3 double positive cells. The corresponding MFI is indicated. The expression profile for FL TNFR25, Δ5,6 TNFR25 and DN TNFR25 were identical. 3B depicts a Western blot of alternatively spliced TNFR25 transfected tumor cells. 50 µg protein from P815 lysates transfected with three splice forms of TNFR25 were loaded and blotted with the anti TNFR25 antibody 1OD1. Lane 1: DN TNFR25; lane 2: Δ5,6 TNFR25; lane 3: FL TNFR25. The antibody does not detect Δ5,6 TNFR25 in western blots. 3C illustrates reduced cellularity of the CD4 and CD8 positive cells in FL TNFR25-tg lymph node cells and thymocytes compared to w.t. littermates (n=5)* p<0.05; **p<0.01; ***p<0.001. 3D depicts impaired activation-induced proliferation of FL and Δ5,6 TNFR25-transgenic cells. Proliferation of purified CD4 or CD8 cells was measured after 3 days of stimulation by thymidine uptake during the last 6 hours. Cells were activated in microtiter plates with immobilized anti CD3 (2 µg/ml) with or without soluble anti CD28 (1 µg/ml) or PMA (10 ng/ml) and ionomycin (400 ng/ml). Recombinant mouse IL-2 was used at 1000 U/ml. 3E illustrates FL TNFR25 and Δ5,6 TNFR25 activate NF-κB NF-κB activation was measured in EL4 cells transfected with FL TNFR25 (upper panel) or with A5,6 TNFR25 (lower panel) in response to TNFR25 triggering. Cells were treated with the agonistic TNFR25 antibody 4C12 (5 µg/ml) for 50 min; soluble TL1A was given for 25, 50, or 75 min as indicated in the form of 25% supernatants from TL1A transfected EL4 cells; membrane bound TL1A (MTL1A) was given for 50 min by adding TL1A transfected EL4-cells directly to TNFR25 expressing EL4. Controls received EL4 (untransfected) supernatants for 50 min. Nuclear extracts were prepared and analyzed by EMSA; the arrow indicates activated NF-κB. 3F depicts primary Th2 biased cytokine production by w.t., FL TNFR25 and Δ5,6 TNFR25-transgenic CD4 T cells. CD4 T cells from spleens were purified by negative selection and activated with immobilized anti-CD3 (2 µg/ml) and soluble anti-CD28 (1 µg/ml) for 3 days. Supernatants were collected for cytokine ELISA assays. The figure is representative of three independent experiments. n.s.: not significant; * p<0.05; **p<0.01; ***p<0.001. 3G illustrates that TNFR25 can costimulate Th1 or Th2 cytokine production. Cytokine production of restimulated FL TNFR25-tg CD4 T cells was determined under non-polarizing (Th neutral), Th1, or Th2 polarizing conditions. CD4 cells were activated with immo-

bilized anti CD3 (2 µg/ml) and soluble anti CD28 (1 µg/ml) alone (Th neutral), or combined with IL-12 (5 ng/ml) and anti-IL-4 (20 µg/ml) for Th1 polarization, or combined with IL-4 (10 ng/ml), anti-IFN-γ (10 µg/ml), and anti-IL-12 (10 µg/ml) for Th2 polarization for 4 days. The cells were harvested, washed and replated on anti CD3 for 24 hours and the supernatants collected for cytokine ELISA analysis.

[0040] FIG. 4 depicts diminished proliferation of TNFR25-transgenic cells is not due to apoptosis, lack of IL-2 or IL-2 receptor expression. 4A illustrates normal upregulation of IL-2-Ra (CD25) in transgenic CD4 and CD8 cells upon activation. After 72-hour activation with immobilized anti-CD3 and soluble anti-CD28, splenocytes were harvested, washed and stained with anti-CD25-FITC, anti-CD8-PE, and anti-CD4-CY. Upper panels wild type, lower panels Δ5,6 TNFR25-tg cells. 4B demonstrates that transgenic cells do not undergo increased apoptosis upon activation. CD4 cells were activated with immobilized anti-mouse CD3 (2 µg/ml) with soluble anti-mCD28 (1 µg/ml) for three days and stained with Annexin-V-PE and 7-AAD. Annexin-V positive and 7-AAD negative cells represented the apoptotic cells. 4C illustrates Reduced IL-2 production by Δ5,6 TNFR25 transgenic T cells. T cells were purified by negative selection and activated with immobilized anti-CD3 and soluble anti-CD28 for 3 days. Supernatants were analyzed for IL-2 production by ELISA assay; ** p=0.0078. 4D illustrates that dominant negative TNFR25-transgenic cells are not Th2 polarized in primary response. CD4 T cells from the spleen were purified by negative selection and activated with immobilized anti-CD3 (2 µg/ml) and soluble anti-CD28 (1 µg/ml) for 3 days. The supernatants were collected for cytokine ELISA assays. The figure is representative of three independent experiments. 4E depicts the same antibody-isotype response of DN TNFR25-transgenic and w.t. mice. Mice were immunized intraperitoneally with 100 µg DNP-KLH in sterile PBS and DNP-specific IgG1 and IgG2a antibodies in serum were evaluated by ELISA three weeks after immunization. High-binding 96-well plates were coated with DNP-BSA at 0.8 µg/ml to detect anti-DNP specific antibodies. Figure represents one of three independent experiments.

[0041] FIG. 5 illustrates that TNFR25 transgenic mice develop a Th2-biased response after in vivo challenge. 5A demonstrates that an antibody-isotype is Th2 biased in immunized TNFR25-tg mice. Mice were immunized intraperitoneally with 100 µg DNP-KLH in sterile PBS without adjuvant. DNP-specific IgG1 and IgG2a antibodies in serum were evaluated by ELISA one week after immunization. High-binding 96-well plates were coated with DNP-BSA at 0.8 µg/ml to detect anti-DNP specific antibodies. Data represent one of three independent experiments. ** p<0.01; n.s.: not significant 5B illustrates TNFR25 signals regulate eosinophilia in broncho-alveolar fluid (BALF). Immunization and airway challenge of w.t. and Δ5,6 TNFR25-tg mice was done as described in Examples 10-15. Airways were lavaged and the differential cell count obtained from Wright-Giemsa stained cytopsins preparations. *: p<0.05; n.s.: not significant 5C depicts increased lung inflammation in TNFR25-tg mice. Lung histology of mice i.p. sensitized and airway challenged with ovalbumin. After bronchial lavage, lungs were removed and fixed in 10% neutral buffered formalin. The lungs were then embedded in paraffin, cut 5 µm thick, and stained with H&E (left panels) or periodic acid-Schiff (PAS, right panels) to detect mucus production.

Upper two panels wild type B6 mice; lower panels Δ5,6 TNFR25-tg mice. 5D illustrates that TNFR25 signals control serum IgE levels. Mice were bled on day 0 and three days after the aerosol challenge (day 15). Serum was separated and analyzed by ELISA for ovalbumin specific IgE by sandwich ELISA. Since there is no standard protein available for OVA-specific IgE, results are presented in O.D. units. The figure represents one of three independent experiments. **: p<0.01. 5E depicts increased Th2 cytokine production by bronchial lymph node cells in TNFR25-tg mice. Bronchial lymph nodes were harvested one day after aerosol challenge (day 13) and cells were restimulated in vitro with 100 µg/ml ovalbumin for 4 days. Supernatants were then analyzed for cytokine production by ELISA. This figure is the representative of two independent experiments. * p<0.05; **p<0.01; ***p<0.001

[0042] FIG. 6 illustrates that dominant negative TNFR25 interferes with Th2 polarization and lung inflammation. 6A illustrates that DN TNFR25 transgene blocks cytokine costimulation by endogenous TNFR25. W.t. and DN TNFR25-tg CD4 cells were stimulated for three days with anti CD3, or with the agonistic anti TNFR25 antibody 4C12 and anti CD3 combined and the supernatants analyzed for cytokines by ELISA. 6B illustrates that DN TNFR25 blocks costimulation of proliferation by endogenous TNFR25. In proliferation assays w.t. and DN TNFR25-tg CD4 cells were activated with anti CD3, with 4C12, or with the combination of anti CD3 and 4C12 for 3 days and thymidine incorporation measured during the final 8 hours. 6C illustrates that DN TNFR25-tg CD4 T cells produce diminished Th2 cytokines in secondary activation. W.t. and transgenic CD4 T cells were purified by negative selection and activated with immobilized anti-CD3 (2 µg/ml) and soluble anti-CD28 (1 µg/ml) for 3 days. Cells were then harvested, washed, replated and restimulated with immobilized anti-CD3 (1 µg/ml) for 2 days. The supernatants were collected for cytokine ELISA assay. n.s.—not significant; *** p<0.001. 6D depicts that DN TNFR25-tg CD4 T cells resist Th2 polarization in vitro. W.t. and DN TNFR25-tg CD4 cells were purified and activated for four days with anti CD3 and anti CD28 under neutral (ThN) or Th2 polarizing conditions (by adding IL-4 and blocking IFN-γ with antibody). Cells were harvested, washed and replated on anti CD3 for 24 h, supernatants harvested and cytokines analyzed by ELISA. 6E illustrates diminished cellular exudation in BALF in DN TNFR25-tg mice compared to w.t. mice. Mice were primed and then aerosol challenged with ovalbumin according to the standard protocol; n=5; *: p<0.05. 6F depicts suppression of lung inflammation in DN TNFR25-tg mice after immunization and airway challenge. Upper panel: Absence of perivascular infiltrates in DN TNFR25-tg mice after antigen aerosol exposure. Lower panel: Absence of mucus over production and goblet cell hyperplasia in DN TNFR25-tg mice. Lung inflammation was induced as in FIG. 5 by ovalbumin immunization and subsequent aerosol exposure. W.t. control mice had typical inflammation as shown in FIG. 5. 6G illustrates suppression of ovalbumin specific IgE production in DN TNFR25-tg mice. Ovalbumin specific IgE in serum was determined by ELISA. ** p<0.01. 6H depicts suppression of Th2 but not Th1 cytokine production by DN TNFR25 in bronchial lymph nodes. Bronchial lymph nodes were harvested one day after aerosol challenge and cells were restimulated with 1001 µg/ml ovalbumin for 4 days. Supernatants were then analyzed for cytokine production by

ELISA. This figure is the representative of two independent experiments. n.d.: not detected; n.s.: not significant; *: p<0.05

[0043] FIG. 7 illustrates that TL1A blocking antibodies abrogate lung inflammation and diminish Th2 cytokine production in the lung. 7A-D depict that anti TL1A antibody L4G6 is a functional antagonist of TL1A. a. P815 transfectants with FL TNFR25, b. with Δ5,6 TNFR25 and c. with TL1A stained by flow cytometry with the appropriate antibodies. d. L4G6 blocks TL1A mediated cytotoxicity of TNFR25 transfected cells. Serially diluted soluble TL1A harvested from supernatants of P815-TL1A transfected cells were mixed with 51Cr-labeled P815-TNFR25 target cells. Different anti-murine TL1A monoclonal antibodies were added into the assay and 51Cr release was analyzed five hours later. 7E illustrates TL1A blocking antibody L4G6 suppresses mucus production and lung inflammation in vivo in wild type C57B16 mice. Schema: Schedule of ovalbumin priming, aerosol challenge and administration of blocking antibody (L4G6) or control antibody. Lung histology after PAS staining with control IgG (left) and L4G6-IgG (right). Notice the lack of mucus production in L4G6 treated animals (arrows point to mucus in mice treated with control IgG). 7F demonstrates diminished cellular exudation in BALF in L4G6 treated mice compared to control IgG treated animals. *: p<0.05; **: p<0.01; n.s. not significant. 7G illustrates diminished IL-5 and IL-13 production by ovalbumin restimulated bronchial lymph node cells after TL1A blockade with L4G6. **p<0.01; *** p<0.001. Experimental details as in FIG. 5. 7H depicts TL1A expression after aerosol challenge on a subpopulation of CD11c positive cells (arrow) in the lung. Bronchial lymph nodes were harvested before and after ova-aerosol challenge, and CD11c cells analyzed for TL1A expression. All other bronchial lymph node cells were TL1A negative. 7I illustrates a lack of TL1A expression on lymphocytes from bronchial lymph node cells of ovalbumin immunized mice after airway challenge. Single cell suspensions were stained in a triple sandwich with anti-TL1A as primary monoclonal antibody. Cells were gated using the respective labeled antibody as population marker and the TL1A histogram displayed. Anti-TL1A; isotype control. 7J illustrates that TL1A is expressed on activated T lymphocytes in vitro. Splenocytes were activated for 24 h with plate bound anti-CD3 or with LPS and then stained with the anti-TL1A triple sandwich as in FIG. 1 and with the population marker as indicated. B cells are TL1A negative even after LPS activation. Gating on the population marker, TL1A expression on activated cells is shown.

[0044] FIG. 8 depicts that TNFR25 signals are required in NKT cells for induction of lung inflammation. NKT deficient Jα18 knock out mice (Cui, J. et al. Science 278, 1623-6 (1997)) were primed with ovalbumin and alum as in the standard protocol in material and methods. On day 11 the mice received by i.v. adoptive transfer 3.1 million partially purified w.t. NKT cells or DN TNFR25-tg NKT cells or PBS as indicated. The mice were aerosolized on day 12 with ovalbumin and on day 14 analyzed. W.t. mice served as positive controls for induction of lung inflammation. Jα18 k.o. mice were immunized and ovalbumin aerosolized without adoptive cell transfer as negative controls. The data are for four mice in each group in two independent experiments. Jα are NKT deficient mice (Cui, J. et al. Science 278, 1623-6 (1997)).

[0045] FIG. 9 illustrates a model for TL1A/TNFR25 mediated triggering of NKT cells and Th2 polarized CD4 cells in the lung. The model depicts the potential interaction between TL1A and TNFR25 that may contribute to IL-13 production and induction of AHR. Evidence for the need of IL-13 production by NKT cells has been provided previously (Akbari, O. et al. Nat Med 9, 582-8 (2003)). The present communication shows the need for TNFR25 signals on NKT cells for lung inflammation, the expression of TL1A by CD11c+ cells and the costimulation of CD4 Th2 effectors by TNFR25. Since CD4 effectors can express TL1A it is possible that TL1A/TNFR25 interaction between CD4 and NKT cells provides the molecular link for their synergy in asthma. In addition other lung associated cells may express TL1A and help triggering lung inflammation.

[0046] FIG. 10 depicts the results of an experiment in which mice received 1 million-OT-I-gfp i.v. Two days later they were immunized i.p. with the indicated dose of ovalbumin in PBS (blue curve); with EG7-gp96-Ig or 3T3-ova-gp96-Ig—the amount of gp96-Ig (in ng) secreted within 24 h by the number of injected cells is indicated on the abscissa; with 3T3-ova—the amount of ovalbumin secreted within 24 h by the number of injected cells is indicated on the abscissa. After 5 days OT-I expansion was determined by flow cytometry in the peritoneal cavity (PEC) and in the spleen. Frequency of OT-I among CD8 cells was up to 50% in the PEC and up to 8% in the spleen with 3T3-ova-gp96-Ig immunization.

[0047] FIG. 11A depicts the results of an experiment in which anti-TNFR25 mAb 4C12 acted agonistically and killed TNFR25 transfected EL4 cells, but not w.t. EL4. In 11B, 50 µg of agonistic 4C12 or antagonistic anti-TL1A L4G6 or control IgG was given i.p. 24 h and 72 h after EG7-gp96-1 g immunization. 4C12 caused 8-10 fold increase of OT-I expansion and doubling of peritoneal exudate cells (n=4 mice per group). Anti TL1A inhibits CD8 expansion.

[0048] FIG. 12 depicts cross priming of CD8 cells by heat shock protein gp96. Vaccine cells (allogeneic or syngeneic) were transfected with gp96-Ig and ovalbumin whereupon they secreted gp96-Ig chaperoning ovalbumin peptides. Gp96 was detected by CD91 and TLR2/4 on DC resulting in their activation and engulfment of gp96_Ig with its bound peptides. Activated DC up-regulate B7 (independent of CD40-L) and cross-resent gp96-bound peptides via Kb (in B6 mice). OT-I are gfp expressing, TCR transgenic CD8 cells adoptively transferred to B6 mice specific for Kb-ova. Their expansion can be easily measured by green fluorescent protein.

[0049] FIG. 13 depicts FoxP3 positive CD4+ CD25+Tregs expressing TNFR25. CD4 cells were purified by depleting B cells CD8 cells and monocytes and then positively selected by magnetic sorting for CD25 and then activated.

[0050] FIG. 14 depicts a lack of recovery of DN-TNFR25-tg mice from DSS induced colitis. 5 mice in each group received 2% DSS in drinking water for 8 days and were then restored to normal water.

[0051] FIG. 15 shows that TNFR25 signals abolish Treg inhibition.

[0052] FIG. 16 shows Gfp-OT-I locating to the mucosa after gp96-Ig immunization i.p. Mice received 1 million

gfp-OT-I i.v. by adoptive transfer. After 2 days the mice received 4 million EG7-gp96-Ig (left) or 4 million EG7 (right panels) i.p. as stimulus. Four days later the frequency of gfp-OT-I was analyzed in IEL, Peyer's patches and LPL in addition to the usual analysis of cells in the PEC, in the spleen and in lymph nodes.

[0053] FIG. 17 shows SEQ ID NOs: 1-7 and SEQ ID NO: 16 and selected public database accession numbers.

DETAILED DESCRIPTION OF THE INVENTION

[0054] It is an object of the invention to provide novel compositions and methods utilizing immunomodulating agents that can either stimulate or indirectly augment the immune system or in other cases, have an immunosuppressive effect. TNFR25 agonists disclosed herein represent biological response modifiers that alter the interaction between the body's cellular immune defenses and cancer cells to boost, direct, or restore the body's ability to fight the cancer when given with tumor vaccines. TNFR25-specific toxic agents disclosed herein are capable of increasing the effectiveness of a chemotherapeutic regimen by depleting a cancer patient of naturally occurring immunosuppressive cells. TNFR25 agonists disclosed herein can also have a healing effect. They can be used, among other things, to treat disease that caused by chronic inflammation such as inflammatory bowel disease. TNFR25 antagonists disclosed herein are capable of inhibiting CD8 T cell-mediated cellular immune responses and can for example, to treat asthma and mitigate organ or tissue rejection following a tissue transplantation.

[0055] 1. Definitions

[0056] In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

[0057] An "antigen" includes any substance that may be specifically bound by an antibody molecule. Thus, the term "antigen" encompasses biologic molecules including, but not limited to, simple intermediary metabolites, sugars, lipids, autoacids, and hormones, as well as macromolecules such as complex carbohydrates, phospholipids, nucleic acids and proteins.

[0058] An "antigenic composition" may comprise an antigen (e.g., a peptide or polypeptide), a nucleic acid encoding an antigen (e.g., an antigen expression vector), or a cell expressing or presenting an antigen. See U.S. Pub. No. 2003/0185840, which is hereby incorporated by reference in its entirety.

[0059] An "immunogen" is a macromolecular antigen that is capable of initiating lymphocyte activation resulting in an antigen-specific immune response. An immunogen therefore includes any molecule which contains one or more epitopes that will stimulate a host's immune system to initiate a secretory, humoral and/or cellular antigen-specific response.

[0060] The term "antibody" encompasses polyclonal and monoclonal antibody preparations. Antibodies of the invention may be prepared in any mammal, including mice, rats, rabbits, goats and humans. The antibody may be a member of one of the following immunoglobulin classes: IgG, IgM, IgA, IgD, or IgE, and the subclasses thereof, and preferably is an IgG1 antibody.

[0061] The term antibody also refers to functional equivalents of the antibodies described in this specification. Functional equivalents have binding characteristics comparable to those of the antibodies, and include, for example, chimerized, hybrid, humanized and single chain antibodies as well as fragments thereof. Methods of producing such functional equivalents are disclosed in PCT Application WO 93/21319, European Patent Application No. 239,400; PCT Application WO 89/09622; European Patent Application 338,745; and European Patent Application EP 332,424. Functional equivalents include polypeptides with amino acid sequences substantially the same as the amino acid sequence of the variable or hypervariable regions of the antibodies of the invention. "Substantially the same" amino acid sequence is defined herein as a sequence with at least 70%, preferably at least about 80%, and more preferably at least about 90% homology to another amino acid sequence, as determined by the FASTA search method in accordance with Pearson and Lipman, Proc. Natl. Acad. Sci. USA 85, 2444-2448 (1988).

[0062] As used herein, the term "monoclonal antibody" refers to an antibody composition having a homogeneous antibody population. The term is not limited regarding the species or source of the antibody, nor is it intended to be limited by the manner in which it is made. The term encompasses whole immunoglobulins.

[0063] Methods of making polyclonal and monoclonal antibodies are known in the art. Polyclonal antibodies are generated by immunizing a suitable animal, such as a mouse, rat, rabbit, sheep or goat, with an antigen of interest. In order to enhance immunogenicity, the antigen can be linked to a carrier prior to immunization. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lipid aggregates (such as oil droplets or liposomes), and inactive virus particles. Such carriers are well known to those of ordinary skill in the art. Furthermore, the antigen may be conjugated to a bacterial toxoid, such as toxoid from diphtheria, tetanus, cholera, etc., in order to enhance the immunogenicity thereof.

[0064] Rabbits, sheep and goats are preferred for the preparation of polyclonal sera when large volumes of sera are desired. These animals are good design choices also because of the availability of labeled anti-rabbit, anti-sheep and anti-goat antibodies. Immunization is generally performed by mixing or emulsifying the antigen in saline, preferably in an adjuvant such as Freund's complete adjuvant ("FCA"), and injecting the mixture or emulsion parenterally (generally subcutaneously or intramuscularly). The animal is generally boosted 2-6 weeks later with one or more injections of the antigen in saline, preferably using Freund's incomplete adjuvant ("FIA"). Antibodies may also be generated by in vitro immunization, using methods known in the art. Polyclonal antisera is then obtained from the immunized animal.

[0065] Monoclonal antibodies are generally prepared using the method of Kohler and Milstein, Nature (1975) 256:495-497, or a modification thereof or Campbell in "Monoclonal Antibody Technology, The Production and Characterization of Rodent and Human Hybridomas" in Burdon et al., Eds., Laboratory Techniques in Biochemistry

and Molecular Biology, Volume 13, Elsevier Science Publishers, Amsterdam (1985); as well as by the recombinant DNA method described by Huse et al in Science 246, 1275-1281 (1989).

[0066] Typically, a mouse or rat is immunized as described above. However, rather than bleeding the animal to extract serum, the spleen (and optionally several large lymph nodes) is removed and dissociated into single cells. If desired, the spleen cells may be screened (after removal of non-specifically adherent cells) by applying a cell suspension to a plate or well coated with the antigen. B-cells, expressing membrane-bound immunoglobulin specific for the antigen, will bind to the plate, and are not rinsed away with the rest of the suspension. Resulting B-cells, or all dissociated spleen cells, are then induced to fuse with myeloma cells to form hybridomas, and are cultured in a selective medium (e.g., hypoxanthine, aminopterin, thymidine medium, "HAT"). The resulting hybridomas are plated by limiting dilution, and are assayed for the production of antibodies which bind specifically to the immunizing antigen (and which do not bind to unrelated antigens). The selected monoclonal antibody-secreting hybridomas are then cultured either in vitro (e.g., in tissue culture bottles or hollow fiber reactors), or in vivo (e.g., as ascites in mice).

[0067] The "antigen-binding site," or "binding portion" refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable ("V") regions of the heavy ("H") and light ("L") chains. Three highly divergent stretches within the V regions of the heavy and light chains are referred to as "hypervariable regions" which are interposed between more conserved flanking stretches known as "framework regions," or "FRs". Thus the term "FR" refers to amino acid sequences which are naturally found between and adjacent to hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as "complementarity-determining regions," or "CDRs."

[0068] As used herein, the terms "immuno-specific," "immunological binding" and "immunological binding properties" refer to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and on geometric parameters that equally influence the rate in both directions. Thus, both the "on rate constant" (K_{on}) and the "off rate constant" (K_{off}) can be determined by calculation of the concentrations and the actual rates of association and dissociation. The ratio of K_{off}/K_{on} enables cancellation of all

parameters not related to affinity, and is thus equal to the dissociation constant K_d . See, generally, Davies et al. (1990) Annual Rev. Biochem. 59:439-473.

[0069] A number of therapeutically useful molecules are known in the art which comprise antigen-binding sites that are capable of exhibiting immunological binding properties of an antibody molecule. The proteolytic enzyme papain preferentially cleaves IgG molecules to yield several fragments, two of which (the "F(ab)" fragments) each comprise a covalent heterodimer that includes an intact antigen-binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the "F(ab')2" fragment which comprises both antigen-binding sites. An "Fv" fragment can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent VH:VL heterodimer including an antigen-binding site which retains much of the antigen recognition and binding capabilities of the native antibody molecule. Inbar et al. (1972) Proc. Nat. Acad. Sci. USA 69:2659-2662; Hochman et al. (1976) Biochem 15:2706-2710; and Ehrlich et al. (1980) Biochem 19:4091-4096.

[0070] A single chain Fv ("sFv") polypeptide is a covalently linked VH:VL heterodimer which is expressed from a gene fusion including VH-and VL-encoding genes linked by a peptide-encoding linker. Huston et al. (1988) Proc. Nat. Acad. Sci. USA 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated—but chemically separated—light and heavy polypeptide chains from an antibody V region into an sFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

[0071] "Cognate antigen" as used herein refers to an antigen for which a CD8 T cell receptor (TCR) is immuno-specific. Such antigen, which are generally derived from e.g. pathogen, transplanted tissues (alloantigen) and tumor cells, are recognized by the immune system as non-self. Binding of the cognate antigen to its CD8 T cell receptor results in the clonal expansion of that T cell. A growing population of cognate CD8 T cells is then in a position to mount a cellular immunological response against the source of the offending cognate antigen.

[0072] Antibodies of the invention can also be used to make "immunotoxins." The hybrid molecule combines the specificity of an antibody or antigen with the toxicity of the toxin. As such, immunotoxin molecules have an antigen binding portion and a toxic agent portion. Immunotoxins are preferably specific for a cell surface molecule, e.g., TNFR25, and facilitate the delivery of a toxic agent to a cell expressing the aforementioned cell surface molecule.

[0073] Preferably, the "toxic agents" have a cytostatic and/or cytotoxic effect on the cell to which it is delivered. Preferred toxic agents are, for example, radioactive isotopes, fluorescers, chemiluminescers, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, dyes, and metal ions. Toxic agents include but are not limited to Iodine-131, Indium-111, and Technetium-99m, Technetium-99m,

Indium-111, Yttrium-90, doxorubicin (Yang et al. (1988) Proc. Natl. Acad. Sci. USA 85:1189-1193), daunorubicin (Diener et al. (1985) Science 231:148-150; Dillman et al. (1988) Cancer Res. 48:6097-6102), methotrexate (Uadia et al. (1985) J Nati Cancer Inst. 74:29-35; Deguchi et al. (1986) Cancer Res. 46:3751-3755), and chlorambucil (Smyth et al. (1986) J. Immunol. 137:3361-3366). Other toxic agents include ricin, abrin, diphtheria toxin and *Pseudomonas* exotoxin, or an enzymatically active portion (A chains) thereof. See, e.g., U.S. Pat. No. 4,753,894 to Frankel et al.; Nevelle, et al. (1982) Immunol Rev 62:75-91; Ross et al. (1980) European J Biochem 104; Vitteta et al. (1982) Immunol Rev 62:158-183; Raso et al. (1982) Cancer Res 42:457-464, and Trowbridge et al. (1981) Nature 294:171-173. Also included are enzymatically active toxins and fragments thereof such as modeccin, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), *momordica charantia* inhibitor, curcin, crotin, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, and enomycin.

[0074] "Toxic agents" can also be chemotherapeutic agents. The term "chemotherapeutic agent" as used herein generally relates to compounds administered to stop, slow or reverse unwanted proliferation of cells. Preferably such agents have an anti-proliferative effect.

[0075] Chemotherapeutic agents may be anti-metabolites such as 5-FU, for example. 5-FU-based chemotherapy comprises administration of 5-FU, its derivatives, alone or with other chemotherapeutics, such as leucovorin or with a DPD inhibitor such as uracil, 5-ethynyluracil, bromovinyluracil, thymine, benzyloxybenzyluracil (BBU) or 5-chloro-2,4-dihydroxypyridine. Furthermore, it has been found that co-administration of a 5'-deoxy-cytidine derivative of the formula (I) with 5-FU or a derivative thereof significantly improves delivery of a chemotherapeutic agent selectively to tumor tissues as compared with the combination of 5-FU or a derivative thereof with a DPD inhibitor 5-ethynyluracil.

[0076] Alternatively, genotoxic agents are those that form persistent genomic lesions and are preferred for use as chemotherapeutic agents in the clinical management of unwanted cellular proliferation. The rate of cellular repair of genotoxin-induced DNA damage, as well as the rate of cell growth via the cell division cycle, affects the outcome of genotoxin therapy. A general class of genotoxic compounds that are used for treating many cancers are DNA alkylating agents and DNA intercalating agents. Psoralens are genotoxic compounds known to be useful in the photochemotherapeutic treatment of cutaneous diseases such as psoriasis, vitiligo, fungal infections and cutaneous T cell lymphoma. Harrison's Principles of Internal Medicine, Part 2 Cardinal Manifestations of Disease, Ch. 60 (12th ed. 1991). Another general class of genotoxic compounds, members of which can alkylate or intercalate into DNA, includes synthetically and naturally sourced antibiotics. Of particular interest herein are antineoplastic antibiotics, which include but are not limited to the following classes of compounds represented by: amsacrine; actinomycin A, C, D (alternatively known as dactinomycin) or F (alternatively KS4); azaserine; bleomycin; carminomycin (carubicin); daunomycin (daunorubicin), or 14-hydroxydaunomycin (adriamycin or doxorubicin); mitomycin A, B or C; mitoxantrone; plicamycin (mithramycin); and the like. Still

another general class of genotoxic agents that are commonly used and that alkylate DNA, are those that include the haloethylnitrosoureas, especially the chloroethylnitrosoureas. Representative members of this broad class include carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine and streptozotocin. Haloethylnitrosoureas first agents can be analogs or derivatives of any of the foregoing representative compounds.

[0077] Tumors currently manageable by platinum coordination compounds such as cisplatin or oxaliplatin include testicular, endometrial, cervical, gastric, squamous cell, adrenocortical and small cell lung carcinomas along with medulloblastomas and neuroblastomas. Other cytotoxic anti-cancer therapeutic agents include, for example, BEP (bleomycin, etoposide, cisplatin) for testicular cancer, MVAC (methotrexate, vinblastine, doxorubicin, cisplatin) for bladder cancer, MVP (mitomycin C, vinblastine, cisplatin) for non-small cell lung cancer treatment.

[0078] Yet another general class of genotoxic agents, members of which alkylate DNA, includes the sulfur and nitrogen mustards. These compounds damage DNA primarily by forming covalent adducts at the N7 atom of guanine. Representative members of this broad class include chlorambucil, cyclophosphamide, ifosfamide, melphalan, mechlorethamine, novembicin, trofosfamide and the like. Oligonucleotides or analogs thereof that interact covalently or noncovalently with specific sequences in the genome of selected cells can also be used as genotoxic agents, if it is desired to select one or more predefined genomic targets as the locus of a genomic lesion.

[0079] Another class of agents, members of which alkylate DNA, include the ethylenimines and methylmelamines. These classes include altretamine (hexamethylmelamine), triethylenephosphoramide (TEPA), triethylenethiophosphoramide (ThioTEPA) and triethylenemelamine, for example.

[0080] Additional classes of DNA alkylating agents include the alkyl sulfonates, represented by busulfan; the azinidines, represented by benzodepa; and others, represented by, e.g., mitoguazone, mitoxantrone and procarbazine. Each of these classes includes analogs and derivatives of the respective representative compounds.

[0081] In one embodiment, chemotherapeutic agents are inhibitors of receptor tyrosine kinases such as EGFR and HER2-neu and are employed as selective inhibitors of the growth of proliferative cells. For example, erbstatin, an EGF receptor tyrosine kinase inhibitor, reduces the growth of EGFR expressing human carcinoma cells. Various derivatives of styrene are also stated to possess tyrosine kinase inhibitory properties and to be of use as anti-tumour agents. Two such styrene derivatives are Class I RTK inhibitors whose effectiveness have been demonstrated by attenuating the growth of human squamous cell carcinoma injected into nude mice. Certain 4-anilinoquinazoline derivatives are useful as inhibitors of receptor tyrosine kinases. The very tight structure-activity relationships shown by these compounds indicates a clearly-defined binding mode, where the quinazoline ring binds in the adenine pocket and the anilino ring binds in an adjacent, unique lipophilic pocket. Three 4-anilinoquinazoline analogues (two reversible and one irreversible inhibitor) have been evaluated clinically as anticancer drugs. Additionally, the monoclonal antibody trasta-

zumab (Herceptin™) for the treatment of HER2-neu overexpressing metastatic breast cancers. Scheurle, et al., Anticancer Res 20:2091-2096, 2000.

[0082] An "expression vector" is any genetic element, e.g., a plasmid, chromosome, virus, behaving either as an autonomous unit of polynucleotide replication within a cell. (i.e., capable of replication under its own control) or being rendered capable of replication by insertion into a host cell chromosome, having attached to it another polynucleotide segment, so as to bring about the replication and/or expression of the attached segment. Suitable vectors include, but are not limited to viruses, plasmids and cosmids.

[0083] Expression vectors contain an "expression cassette" which includes a polynucleotide sequence to be transcribed operably linked to polynucleotide sequences which are necessary to effect ligation or insertion of the vector into a desired host cell and to effect the transcription of the polynucleotide to be expressed. Such sequences include promoter sequences to effect transcription, enhancer sequences to increase transcription, ribosomal binding site sequences and transcription and translation termination sequences. Alternatively, expression vectors may be capable of directly expressing nucleic acid sequence products encoded therein without ligation or integration of the vector into host cell DNA sequences.

[0084] The term "operably linked" refers to the linkage of a DNA segment to another DNA segment in such a way as to allow the segments to function in their intended manners. A DNA sequence encoding a gene product is operably linked to a regulatory sequence when it is ligated to the regulatory sequence, such as, for example, promoters, enhancers and/or silencers, in a manner which allows modulation of transcription of the DNA sequence, directly or indirectly. For example, a DNA sequence is operably linked to a promoter when it is ligated to the promoter downstream with respect to the transcription initiation site of the promoter, in the correct reading frame with respect to the transcription initiation site and allows transcription elongation to proceed through the DNA sequence. An enhancer or silencer is operably linked to a DNA sequence coding for a gene product when it is ligated to the DNA sequence in such a manner as to increase or decrease, respectively, the transcription of the DNA sequence. Enhancers and silencers may be located upstream, downstream or embedded within the coding regions of the DNA sequence. A DNA for a signal sequence is operably linked to DNA coding for a polypeptide if the signal sequence is expressed as a preprotein that participates in the secretion of the polypeptide. Linkage of DNA sequences to regulatory sequences is typically accomplished by ligation at suitable restriction sites or via adapters or linkers inserted in the sequence using restriction endonucleases known to one of skill in the art.

[0085] "Small molecule" as used herein refers to molecular weight synthetic compounds designed to interact with a specific protein known to be involved in a given disease condition. Libraries of such compounds may be readily synthesized by combinatorial chemistry for example, and screened for TNFR25 agonist/antagonist activity using conventional techniques.

[0086] The terms "TNFR-SF25", "TNFR25" or "DR3" are all used interchangeably herein for a member of the TNF receptor family whose complete biological function was

previously not known. See U.S. Pat. No. 6,713,061, and Borysenko, et al., *Biochem Biophys Res Commun.* 2005 Mar. 18;328(3):794-9, Sheikh, et al., *Curr. Cancer Drug Targets.* 2004 February;4(1):97-104, which are incorporated by reference in their entirety. However, the inventors, however, have made a number of important discoveries. The cDNA sequence encoding mouse TNFR25 is shown as SEQ ID NO: 1. The cDNA encoding human TNFR25 is shown as SEQ ID NO: 2.

[0087] Unlike that of any other member of the TNF-R family, DR3 expression was found to be controlled by alternative mRNA splicing. Resting T cells express little or no DR3 protein, but contained high levels of randomly spliced DR3 mRNA. Upon T cell activation via the T cell receptor, protein kinase C (PKC) is activated. PKC activation in turn mediates correct splicing of full-length DR3 and surface expression of the protein. This unique regulation of DR3 expression allows for rapid DR3 protein expression on T cells and enables environmental regulation of DR3 expression via influencing PKC levels responsible for DR3 splicing and expression.

[0088] DR3 is also involved in co-stimulating T cell polarization and in stimulating the production of IL-13 and IL-10 in Th2 polarized cells. This is an important observation because among other things, IL-10 production plays a critical role in suppressing inflammatory bowel disease.

[0089] Transgenic expression of TNFR25 in T cells mediates TH2 polarization of cytokine and antibody production upon T cell activation and antigen exposure. In addition transgenic TNFR25 partially inhibits TCR driven proliferation of CD4 and CD8 cells and reduced total T cell numbers in lymphoid organs without inducing apoptosis. CD8 cells were more affected by TNFR25 than CD4 cells. As such, TNFR25 signals are important in effector responses to pathogens by shaping the ensuing polarization towards TH2 or towards a mixed TH1/TH2 response.

[0090] TNFR25 transgenic mice are highly susceptible to antigen induced airway inflammation in an asthma model in mice and produced increased quantities of IL-13 and eosinophils in the lung upon antigen exposure by inhalation. Transgenic mice expressing a dominant negative form of TNFR25 showed increased resistance to airway hyperreactivity when compared to wild type mice.

[0091] "TL1A" is referred to herein as a TNF-like factor that acts as a costimulator of IFN-gamma secretion through binding to the death domain-containing receptor, DR3. TL1A, like TNF, is also presumed to circulate as a homotrimeric soluble form. As such, "soluble TL1A" as used herein, refers to homotrimeric TL1A. The term is not limited to any species specific form. However, the cDNA sequence for the human TL1A monomer is provided as SEQ ID NO: 7 and that of mouse as SEQ ID NO: 3.

[0092] TL1A has been suggested to play a role in inflammatory bowel disease (IBD) by functioning as a Th1-polarizing cytokine. It has been shown that the amount of TL1A protein and the number of TL1A-positive cells correlate with the severity of inflammation, most significantly in Crohn's Disease (CD). It has also been shown that the addition of recombinant human TL1A to cultures of PHA-stimulated lamina propria mononuclear from CD patients significantly augmented IFN-gamma production by 4-fold,

whereas a minimal effect was observed in control patients. Additionally, a blocking anti TL1A antibody was able to ameliorate asthma in wild type mice indicating that TNFR25 and TL1A are involved in the pathogenesis of asthma.

[0093] The receptors of the human body work by being stimulated or inhibited by natural (such as hormones, cytokines and neurotransmitters) or synthetic (such as drugs, e.g., antibodies or small molecules) agonists.

[0094] "TNFR25 agonist" is referred to herein as a substance that binds to the TNFR25 receptor and triggers a response in the cell on which the TNFR25 receptor is expressed similar to a response that would be observed by exposing the cell to a natural TNFR25 ligand, e.g., TL1A. An agonist is the opposite of an antagonist in the sense that while an antagonist may also bind to the receptor, it fails to activate the receptor and actually completely or partially blocks it from activation by endogenous or exogenous agonists. A partial agonist activates a receptor but does not cause as much of a physiological change as does a full agonist.

[0095] Soluble TL1A might be given in therapeutic form to a patient to increase the activation of the TNFR25 receptor in a given cell population as a TNFR25 agonist. Alternatively, another example of a TNFR25 agonist is an antibody that is capable of binding and activating TNFR25. For example, the monoclonal antibody 4C12 binds and activates TNFR25 signaling. In another embodiment and in the context of this invention, a TNFR25 agonist may be derived from an expression vector with an expression cassette capable of ectopically driving the transgenic expression of a TNFR25 agonist antibody and/or TL1A protein at a chosen location or at a chosen time. In yet another embodiment, a TNFR25 agonist leading to an increase in TNFR25 signaling in a tissue is provided by an expression vector with an expression cassette capable of driving the transgenic expression of TNFR25 itself. This would be useful in a situation where increased TNFR25 signaling is desired in a tissue in which there is an excess of exogenous or endogenous TNFR25 agonist(s) relative to receptor.

[0096] "TNFR25 antagonist" is referred to herein as a substance that inhibits the normal physiological function of a TNFR25 receptor. Such agents work by interfering in the binding of endogenous receptor agonists/ligands such as TL1A, with TNFR25 receptor. An example of a TNFR25 antagonist is a dominant negative TNFR25 receptor. Preferably, the TNFR25 antagonist used herein is an antibody specific to TL1A which interferes with TL1A's ability to activate the TNFR25 receptor. Most preferably, that antibody is the monoclonal antibody L4G6. In another embodiment, the TNFR25 antagonist is a fusion protein of the extracellular portion of TNFR25 or an alternative splice form of TNFR25 with The Fc portion of immunoglobulin or any other suitable fusionpartner. In another embodiment, the TNFR25 antagonist is a soluble form of TNFR25 made by truncation above the transmembrane binding domain, either as alternative splice form or as an artificial construct. In another embodiment, the TNFR25 antagonist is an antibody that specifically binds TNFR25 and interferes with its binding to its natural ligand(s). In another embodiment, TNFR25 antagonist may be an expression vector with an expression cassette capable of driving the transgenic expression of an antisense mRNA, RNAi or ribozyme that is capable of

knocking down endogenous TNFR25 and/or TL1A mRNA transcription and/or translation at a chosen location or at a chosen time. In yet another embodiment, one may decrease TNFR25 signaling in a tissue by providing an expression vector with an expression cassette capable of driving the transgenic expression of a dominant negative TNFR25.

[0097] TNFR25 antagonists or agonists may be in the form of aptamers. "Aptamers" are DNA or RNA molecules that have been selected from random pools based on their ability to bind other molecules. In one embodiment, aptamers specifically bind TNFR25 to block binding of its natural ligand, e.g., TL1A, or which bind TL1A itself, and prevent it from binding TNFR25. In another embodiment, aptamers specifically bind the TNFR25 receptor and activate it.

[0098] "Dominant negative" or "DN" as used herein refers to an exogenously provided structural variant of TNFR25 that acts to block endogenous TNFR25. For example, a molar excess of a DN will out compete endogenous TNFR25 for binding of the TNFR25 ligand, e.g., TL1A. Preferably the DN is the same as the wild type TNFR25 except that it is missing the intracellular domain. Alternatively, the DN is the same as the wild type TNFR25 except that it is missing the transmembrane and the intracellular domain. The coding sequence for the mouse DN TNFR25 is provided in SEQ ID NO: 4. The coding sequence for the human DN TNFR25 is provided in SEQ ID NO: 5. The coding sequence for the human DN TNFR25 containing only the extracellular domain is provided in SEQ ID NO: 6.

[0099] The invention also involves the coding sequences that are substantially identical to SEQ ID NOS: 3-7. The skilled artisan will also appreciate that oligonucleotide sequences substantially identical to SEQ ID NOS: 3-7 may differ from SEQ ID NOS: 3-7, respectively, with respect to the identity of at least one nucleotide base. However, all oligonucleotides sequences substantially identical to SEQ ID NOS: 3-7 will hybridize under stringent conditions (as defined herein) to all or a portion of the complements of SEQ ID NOS: 3-6 (i.e., target sequences), respectively. The terms "hybridize(s) specifically" or "specifically hybridize(s)" refer to complementary hybridization between an oligonucleotide (e.g., a primer or labeled probe) and a target sequence. The term specifically embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired priming for the PCR polymerases or detection of hybridization signal.

[0100] Under stringent hybridization conditions, only highly complementary, i.e., substantially identical nucleic acid sequences, hybridize. Preferably, such conditions prevent hybridization of nucleic acids having 3 or more mismatches out of 20 contiguous nucleotides, more preferably 2 or more mismatches out of 20 contiguous nucleotides, most preferably one or more mismatch out of 20 contiguous nucleotides. The hybridizing portion of the hybridizing nucleic acid is at least about 90%, preferably at least about 95%, or most preferably about at least about 98%, identical to the sequence of a target sequence, or its complement.

[0101] Hybridization of a nucleic acid to a nucleic acid sample under stringent conditions is defined below. Nucleic acid duplex or hybrid stability is expressed as a melting temperature (T_m), which is the temperature at which the probe dissociates from the target DNA. This melting temperature is used to define the required stringency conditions.

If sequences are to be identified that are substantially identical to the probe, rather than identical, then it is useful to first establish the lowest temperature at which only homologous hybridization occurs with a particular concentration of salt (e.g. SSC or SSPE). Then assuming that 1% mismatching results in a 1° C. decrease in T_m , the temperature of the final wash in the hybridization reaction is reduced accordingly (for example, if sequences having >95% identity with the probe are sought, the final wash temperature is decrease by 5° C.). In practice, the change in T_m can be between 0.5° C. and 1.5° C. per 1% mismatch.

[0102] Stringent conditions involve hybridizing at 68° C. in 5×SSC/5× Denhart's solution/1.0% SDS, and washing in 0.2×SSC/0.1% SDS at room temperature. Moderately stringent conditions include washing in 3×SSC at 42° C. The parameters of salt concentration and temperature may be varied to achieve optimal level of identity between the primer and the target nucleic acid. Additional guidance regarding such conditions is readily available in the art, for example, Sambrook, Fischer and Maniatis, Molecular Cloning, a laboratory manual, (2nd ed.), Cold Spring Harbor Laboratory Press, New York, (1989) and F. M. Ausubel et al eds., Current Protocols in Molecular Biology, John Wiley and Sons (1994).

[0103] "Immunosuppressants" as the term is used herein are important drugs necessary for the treatment of inflammatory diseases and facilitation of organ transplants. For example, Cyclosporin A (CsA) has considerable immunosuppressive activity. It has revolutionized organ transplantation and is commonly used in the treatment of autoimmune diseases. For a recent review of the use of CsA and its mechanisms of action, see Wenger et al; Cyclosporine Chemistry, Structure-activity relationships and Mode of Action, Progress in Clinical Biochemistry and Medicine, Vol. 2, 176 (1986). However, CsA is a powerful medication that can have severe side effects such as renal failure, bone marrow suppression and infertility.

[0104] Corticosteroids are also used in inflammatory conditions for their anti-inflammatory effects. They have a rapid onset of action, and profoundly affect many parts of the immune system as well as most other body systems. Corticosteroids are a cornerstone of treating most types of vasculitis, and are often used in combination with other immunosuppressive medications. However, long term use of corticosteroid can also have severe side effects including hypertension, weight gain, acne, and a swollen face. Other immunosuppressives include azathioprine, methotrexate, cyclophosphamide, mercaptopurine, tacrolimus and mycophenolate mofetil.

[0105] As a general matter, the agonists and antagonists may be provided to a patient in either compositions that are to be swallowed, injected, inhaled or provide by way of suppositories. Alternatively, the compositions may be formulating into ear or eye drops.

[0106] The term "asthma" as used herein includes any asthmatic condition marked by recurrent attacks of paroxysmal dyspnea (i.e., "reversible obstructive airway passage disease") with wheezing due to spasmotic contraction of the bronchi (so called "bronchospasm"). Asthmatic conditions which may be treated or even prevented in accordance with this invention include allergic asthma and bronchial allergy characterized by manifestations in sensitized persons pro-

voked by a variety of factors including exercise, especially vigorous exercise ("exercise-induced bronchospasm"), irritant particles (pollen, dust, cotton, cat dander) as well as mild to moderate asthma, chronic asthma, severe chronic asthma, severe and unstable asthma, nocturnal asthma, and psychologic stresses. The methods of this invention may be particularly useful in preventing the onset of asthma in mammals e.g., humans afflicted with reversible obstructive disease of the lower airway passages and lungs as well as exercise-induced bronchospasm. In methods for treating asthma disclosed herein, the preferred method of delivering the inventive antagonists is through inhalation

[0107] There are several different types of devices which use generally different mechanisms and methodologies to produce aerosols for inhalation. The most commonly used device is a metered dose inhaler (MDI) which comprises a drug formulation container with the formulation including a low boiling point propellant. The formulation is held in the container under pressure and a metered dose of formulation is released as an aerosol when the valve on the container is opened. The low boiling point propellant quickly evaporates or "flashes" when the formulation is exposed to atmospheric pressure outside the container. The particles of formulation containing the drug without the propellant are inhaled into the patient's lungs and thereafter migrate into the patient's circulatory system. There are a number of different types of MDI devices. Devices of this type are disclosed in U.S. Pat. No. 5,404,871 issued Apr. 11, 1995 and U.S. Pat. No. 5,364,838 issued Nov. 15, 1994.

[0108] Another type of device is the dry powder inhaler (DPI) device. As indicated by the name such devices use formulations of dry powder which powder is blown into an aerosolized cloud via a burst of gas. Typical DPI devices are shown in U.S. Pat. No. 5,775,320 issued Jul. 7, 1998 and U.S. Pat. No. 5,740,794 issued Apr. 21, 1998.

[0109] Yet another type of aerosol delivery device forces a formulation through a porous membrane. Formulation moving through the pores breaks up to form small particles which are inhaled by the patient. Devices of this type are shown in U.S. Pat. No. 5,554,646 issued Aug. 13, 1996 and U.S. Pat. No. 5,522,385 issued Jun. 4, 1996.

[0110] With respect to inhalable compositions, suitable carrier materials may be in the form of an amorphous powder, a crystalline powder, or a combination of amorphous and crystalline powders. Suitable materials include carbohydrates, e.g., monosaccharides such as fructose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as lactose, trehalose, cellobiose, and the like; cyclodextrins, such as 2-hydroxypropyl-beta-cyclodextrin; and polysaccharides, such as raffinose, maltodextrins, dextrans, and the like; (b) amino acids, such as glycine, arginine, aspartic acid, glutamic acid, cysteine, lysine, and the like; (c) organic salts prepared from organic acids and bases, such as sodium citrate, sodium ascorbate, magnesium gluconate, sodium gluconate, tromethamine hydrochloride, and the like; (d) peptides and proteins, such as aspartame, human serum albumin, gelatin, and the like; (e) alditols, such as mannitol, xylitol, and the like. A preferred group of carriers includes lactose, trehalose, raffinose, maltodextrins, glycine, sodium citrate, tromethamine hydrochloride, human serum albumin, and mannitol.

[0111] Such carrier materials may be combined with the inventive agonists or antagonists prior to spray drying, i.e.,

by adding the carrier material to the buffer solution which is prepared for spray drying. In that way, the carrier material will be formed simultaneously with and as part of the inventive agonist or antagonist particles. Typically, when the carrier is formed by spray drying together with the inventive agonists or antagonists, the inventive agonists or antagonists will be present in each individual particle at a weight percent in the range from 5% to 95%, preferably from 20% to 80%. The remainder of the particle will primarily be carrier material (typically being from 5% to 95%, usually being from 20% to 80% by weight), but will also include buffer(s) and may include other components as described above. The presence of carrier material in the particles which are delivered to the alveolar region of the lung (i.e., those in the requisite size range below 10 μm) has been found not to significantly interfere with systemic absorption of inventive agonists or antagonists.

[0112] Alternatively, the carriers may be separately prepared in a dry powder form and combined with the dry powder inventive agonists or antagonists by blending. The separately prepared powder carriers will usually be crystalline (to avoid water absorption), but might in some cases be amorphous or mixtures of crystalline and amorphous. The size of the carrier particles may be selected to improve the flowability of the inventive agonists or antagonists powder, typically being in the range from 25 μm to 100 μm . Carrier particles in this size range will generally not penetrate into the alveolar region of the lung and will often separate from the inventive agonists or antagonists in the delivery device prior to inhalation. Thus, the particles which penetrate into the alveolar region of the lung will consist essentially of inventive agonists or antagonists and buffer. A preferred carrier material is crystalline mannitol having a size in the above-stated range.

[0113] The dry powder compositions of the present invention are preferably aerosolized by dispersion in a flowing air or other physiologically acceptable gas stream in a conventional manner. One system suitable for such dispersion is described in copending application Ser. No. 07/910,048, which has been published as WO 93/00951, the full disclosures of which are incorporated herein by reference.

[0114] Sterile injectable solutions are prepared by incorporating the compositions and formulations of the invention in the required amount in the appropriate solvent, such as sodium phosphate-buffered saline, followed by filter sterilization. As used herein, "a physiologically acceptable carrier" includes any and all solvents, dispersion media, antibacterial and antifungal agents that are non-toxic to humans, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. The media or agent must be compatible with maintenance of proper conformation of the inventive protein chains, and its use in the therapeutic compositions. Supplementary active ingredients can also be incorporated into the compositions. A description of exemplary pharmaceutically acceptable carriers and diluents, as well as pharmaceutical formulations, can be found in Remington's Pharmaceutical Sciences, a standard text in this field, and in USP/NF.

[0115] The dosage and mode of administration of claimed compositions should be adjusted according to the identity, formulation, route of administration and other relevant characteristics pertaining, as is known, or readily discernable in the art.

[0116] 2. TNFR25 Agonists as Direct Potentiators of T Cell Polarization and Lung Inflammation

[0117] TNFR25 and its ligand TL1A were discovered to be direct potentiators of T cell polarization and lung inflammation. The important role of TL1A and TNFR25 in lung disease is supported by the exaggerated lung inflammation observed in agonistic TNFR25-transgenic mice. Over-expression of TNFR25 results in a strong Th2 bias of CD4 cells upon activation. This bias in turn is likely to be responsible for increased lung inflammation in the ovalbumin model.

[0118] The inventors observed that in a murine model of asthma, blockade of TNFR25 signaling results in reduced Th2 cytokine production which inhibits lung inflammation. TNFR25 on CD4 cells triggers IL-13 secretion on both Th1 and Th2 polarized cells and TNFR25 signals are needed in NKT-cells that initiate and promote lung inflammation. Chronic IL-13 production, triggered by TNFR25, also is responsible for the sequelae of chronic lung inflammation including but not restricted to chronic asthma, for airway remodeling and fibrosis. Blockade of TNFR25 signaling on NKT cells by a dominant negative TNFR25 mutant in adoptive transfer experiments abrogates lung inflammation. Blockade of TNFR25 signals such as this would be useful in the treatment of acute and chronic asthma and other lung disorders.

[0119] In the chain of events leading to lung inflammation, IL-13 production by NKT cells is an early step (Akbari, O. et al. Nat Med 9, 582-8 (2003)). The critical role for NKT cells in asthma is supported by this study since the adoptive transfer experiments indicated that one of the principal molecular switches used by NKT cells to induce lung inflammation is TNFR25. TNFR25 has the additional ability to modulate incipient effector responses by antigen specific CD4 memory cells through PKC mediated TNFR25 splicing in TCR activated cells. TNFR25 thus acts in the very early phase of the initiation of the memory response in the lung. Therefore, blockade of TNFR25 signaling by anti-TL1A or by other procedures interrupts the cascade of events leading to acute lung inflammation which is thought to be responsible for asthma attacks.

[0120] Blockade of TNFR25 was achieved in wild type mice genetically as well as by antibody blockade with the two independent methods giving similar results. Interfering with TNFR25 signals in vivo resulted in diminished IL-13, IL-5 and IL-4 production by antigen restimulated draining bronchial lymph nodes and in suppression of lung inflammation. Importantly, anti TL1A antibody blockade during the phase of airway antigen challenge of primed and Th2 polarized mice was effective in inhibiting lung inflammation indicating a direct role of TNFR25 in the effector phase in the lung.

[0121] 3. Constitutive TNFR25 Expression on NKT Cells and Inducible Expression on Activated T Cells.

[0122] To study the biological functions of TNFR25 and its cognate ligand TL1A, hamster anti-mouse monoclonal antibodies were generated by standard protocols. To reliably detect the low level of TNFR25 expression by flow cytometry on primary cells it was necessary to develop a triple layer sandwich assay. Without activation TNFR25, was detected at low levels on naïve CD4 T cells and at even lower levels also on CD8 T cells, but not on B cells (FIG.

1a). In addition a subpopulation of CD11c+ cells expressed TNFR25. NKT cells constitutively expressed relatively high levels of TNFR25, while only a small fraction of CD3 negative NK11+ cells showed TNFR25 expression (FIG. 1a). In the thymus single positive CD4 and CD8 cells expressed TNFR25 similar to peripheral T cells. CD4, CD8 double positive and double negative thymocytes did not express TNFR25 (FIG. 1c).

[0123] Upon activation of peripheral T cells with anti-CD3 and anti-CD28, TNFR25 expression was upregulated on both CD4+ and CD8+ cells (FIG. 1b). LPS activation of B cells on the other hand did not result in TNFR25 expression. Murine TNFR25 mRNA is constitutively expressed in T cells but randomly spliced similar to human TNFR25 (Screaton, G. R. et al. Proc Natl Acad Sci USA 94, 4615-9. (1997)) (FIG. 2). Increased TNFR25 protein expression on activated T cells was associated with activation induced splicing of full length TNFR25 from randomly spliced TNFR25 mRNA. Activation induced splicing of TNFR25 in T cells was blocked by the chemical inhibitor H7 indicating a role for PKC in splicing (FIG. 2).

[0124] The preferential expression of TNFR25 on NKT and activated T cells and its activation induced splicing to full length TNFR25 and rapid increase in surface expression raised the question of the biological function of TNFR25 in the immune system. The ablation of the TNFR25 gene in mice did not reveal a definitive phenotype for TNFR25 deficiency, except for a mild defect in negative selection in the thymus. Hence, the biological effects of T cell-expressed TNFR25-transgenes driven by the CD2 promoter and enhancer were analyzed.

[0125] Full length TNFR25 (FL TNFR25, in FIG. 2) and an alternative splice product of TNFR25 lacking exon 5 and 6 (Δ 5,6 TNFR25, FIG. 2) were used for transgenic expression. Δ 5,6 TNFR25 lacks exon 5 and 6 encoding part of the fourth cysteine rich, extracellular domain. It is anchored in the membrane, however, and has a complete intracellular signaling domain like FL TNFR25, and binds TL1A and an agonistic anti TNFR25 antibody (4C12). In addition a dominant negative mutant, DN TNFR25, truncated immediately after the transmembrane domain was expressed as transgene and will be described below.

[0126] 4. Transgenic Over-Expression of TNFR25 Promotes Th2 Polarization of CD4 Cells and Mediates Increased Lung Inflammation in the Ovalbumin Model of AHR.

[0127] Four independent founders for each TNFR25 transgene were obtained and analyzed. The CD2 promoter and enhancer supported position independent transgene expression in all founders Δ 5,6 TNFR25, FL TNFR25 and DN TNFR25 revealed high level expression in resting T cells, NKT cells, NK cells and in a subpopulation of CD11c+ cells. B cells did not express the transgene (FIG. 3a). The authenticity of antibody detection by flow cytometry was verified by Western blots of transfected tumor cells (FIG. 3b) and identical Western blot bands were detected on transgenic splenocytes while the endogenous molecule was below the level of detection by Western blot. The Δ 5,6-splice form was not detected by the antibody 10D1 in Western blots (FIG. 3b) indicating that it binds to exon 5 or 6. Transgenic over-expression of FL TNFR25 was associated with diminished numbers of T cells in primary and secondary lymphoid

organs compared to non-transgenic litter mates (FIG. 3c); the effect of the Δ5,6 transgene on cellularity was modest in the thymus and not significant secondary lymphoid organs. The reduced number of T cells in transgenic mice was accompanied by diminished proliferation in response to anti CD3 and CD28 stimulation when comparing equal numbers of purified transgenic with non-transgenic littermate CD4 and CD8 cells (FIG. 3d). Diminished proliferation was seen at all time points from 24 to 72 hours. However, stimulation of TNFR25-tg CD4 or CD8 cells with the phorbol ester PMA and the Ca-ionophore ionomycin restored normal proliferation indicating that transgenic cells did not have an intrinsic defect in their ability to proliferate. CD3/CD28 activated TNFR25-tg T cells upregulated CD25 normally but produced only about one half the amount of IL-2 (FIG. 4) compared to littermate controls. Exogenous addition of excess IL-2 did not restore proliferation (FIG. 3d). TNFR25-tg T cells did not undergo increased apoptosis as measured by annexin V staining (FIG. 4) indicating that the proliferative deficit is not due to TNFR25 signals for cell death.

[0128] FL and Δ5,6 TNFR25 transfected EL4 cells were used to compare the signaling properties of the splice variants triggered with soluble TL1A, membrane bound TL1A (EL4-TL1A) or an agonistic anti TNFR25 antibody (4C12). All three ligands rapidly induced NF-κB activation within 25 min as detected by EMSA (FIG. 3e).

[0129] After primary activation with plate bound anti-CD3 and soluble anti-CD28 FL and Δ5,6 TNFR25 transgenic cells produced significantly increased quantities of Th2 cytokines including IL-4, IL-5, IL-13 and IL-10 when compared to non-transgenic littermates (FIG. 3f). IFN-γ was diminished when FL TNFR25-tg CD4 cells were activated but not with Δ5,6 TNFR25 transgenes, indicating a subtle difference in the function of the splice variants. Although proliferation of TNFR25-transgenic CD4 cells was diminished, increased Th2 cytokine production was detectable already within 24 hours of activation and continued to increase in the following days, indicating Th2 bias existed prior to activation.

[0130] Next it was determined whether the Th2 bias of FL TNFR25-transgenic CD4 cells could be overruled under Th1 polarizing conditions. Under Th neutral conditions TNFR25-tg CD4 cells produced as much IL-4 as w.t. CD4 cells under Th2 polarizing conditions (FIG. 3g). Incubation of FL TNFR25-tg cells under Th2-polarizing conditions had no additional effect on IL-4 or IL-13 production indicating that the cells were already maximally Th2 polarized during primary activation under Th-neutral conditions. However, FL TNFR25-tg CD4 cells could be polarized to Th1 by including in the culture antibodies to IL-4 and adding exogenous IL-12. Under these conditions FL TNFR25-tg cells produced IFN-γ at higher levels than Th11-polarized wild type cells (FIG. 3g) indicating that the TNFR25 transgene can also costimulate Th1 cytokines. Th1 polarized FL TNFR25-tg CD4 cells, unlike w.t. Th1 cells, also produced IL-13 but only minimal amounts of IL-4. Transgenic over expression of TNFR25 while spontaneously biased towards Th2, nonetheless can costimulate either Th1 or Th2 type cytokine production under appropriate polarizing conditions. In addition TNFR25 signals costimulate IL-13 production under either Th1 or Th2 polarizing conditions.

[0131] The spontaneous Th2 bias of TNFR25-tg mice in vivo by immunization and analysis of antibody isotype

production was evaluated. In vivo studies were carried out with Δ5,6 TNFR25-transgenic mice. Δ5,6-TNFR25 shows the same signaling properties as FL TNFR25 with regard to NF-κB induction (FIG. 3e) and induction of apoptosis and generates a similar Th2 biased cytokine profile. However, unlike the FL-transgenic mice, Δ5,6 transgenic mice have normal CD4 T cell cellularity in lymph nodes (FIG. 3c) and spleens and therefore may more accurately represent TNFR25 function for in vivo experiments.

[0132] DNP-KLH immunized TNFR25-tg mice generated increased ratios of antigen specific IgG1/IgG2a antibody compared to non-transgenic litter mates, indicative of an increased Th2 type antibody ratio in vivo (FIG. 5a). Without immunization, IgG1 and IgG2a levels of w.t. and transgenic mice were identical, evidencing that activation is necessary to reveal the Th2 bias.

[0133] The Th2 bias of TNFR25-tg CD4 cells and their increased IL-13 production suggested that TNFR25 overexpression may predispose to increased allergic lung inflammation characteristic for asthma, as IL-13 is the signature cytokine of that condition (Elias, J. A. et al. Am J Respir Cell Mol Biol 28, 401-4 (2003)). This hypothesis was tested using the classical ovalbumin model for experimental lung inflammation in mice. TNFR25-tg B6 mice and w.t. controls were primed i.p. with ovalbumin and alum on day 0 and 5. On day 12 they were airway challenged with aerosolized ovalbumin and analyzed one to three days later. TNFR25-tg mice contained dramatically increased numbers of eosinophils in the broncho-alveolar fluid (BALF, FIG. 5b), associated with increased ovalbumin specific IgE levels in serum (FIG. 5d) and elevated Th2 cytokine production by ovalbumin restimulation of bronchial lymph node cells (FIG. 5e). IFN-γ production was diminished in transgenic bronchial lymph node cells compared to w.t. cells. Histopathological analysis of the lungs of TNFR25-tg mice showed massively increased perivascular lung infiltration by eosinophils, increased bronchial mucus production and goblet cell hyperplasia stained with PAS (FIG. 5c) consistent with exacerbation of lung inflammation by TNFR25 over-expression on T cells.

[0134] 5. Genetic or Antibody Blockade of TNFR25 During Airway Challenge of Primed Mice Blocks Lung Inflammation.

[0135] A dominant negative mutant of TNFR25, DN TNFR25 was made to block TNFR25 signaling during airway challenge, and expressed the construct as transgene under the CD2 promoter and enhancer. The DN TNFR25-transgene lacks the entire intracellular signaling domain but is identical to full length TNFR25 in its transmembrane and extracellular domain.

[0136] Transgenic DN TNFR25 was expressed at identical levels as the agonistic TNFR25 transgenes (FIG. 3a) as determined by flow cytometry. Using surface fluorescence intensity as measure for the number of expressed molecules, a three to four fold molar excess of transgenic TNFR25 expression over endogenous TNFR25 was determined. This level of overexpression of DN TNFR25 silenced the activity of endogenous TNFR25. Primary anti CD3 activation of both w.t. and DN-transgenic CD4 cells stimulates both Th1 and Th2 cytokine secretion (FIG. 4, panel d) Triggering of TNFR25 with an agonistic antibody (4C12) during primary anti CD3 activation of w.t cells costimulates both Th1 and

Th2 cytokine production but this costimulatory effect of TNFR25 is blocked by the DN TNFR25 transgene (FIG. 6a), indicating the transgene blocks the function of the endogenous gene. Similarly, the agonistic effect of 4C12 costimulates proliferation of w.t. CD4 cells which was blocked in DN TNFR25-tg CD4 cells (FIG. 6b) indicating that endogenous TNFR25 is silenced. Importantly, the expression of DN TNFR25 on CD4 cells greatly diminished the normally observed, upregulated Th2 cytokine production upon secondary activation of cells that had been primed under Th-neutral conditions (FIG. 6c), indicating that TNFR25 signals strongly promote or are required for Th2 polarization and that these signals are blocked by the DN-transgene. More importantly, DN TNFR25-tg cells could not be Th2 polarized even under Th2 polarizing conditions which were provided by the addition of IL-4 and blocking antibodies to IFN- γ and IL-12 (FIG. 6d). Th1 polarization of DNTNFR25-tg CD4 cells on the other hand was not affected even under non-polarizing (Th neutral, ThN) conditions. Thus although TNFR25 costimulates the production of both Th1 and Th2 cytokines in primary activation of CD4 cells, TNFR25 signals appear to be necessary to promote Th2 polarization in vitro. Th1 polarization is independent of TNFR25, but TNFR25 signals in Th1 cells promote IL-13 production (FIG. 3g). The absence of TNFR25 signals in DN TNFR25-tg mice *in vivo* does not affect the normal IgG1 to IgG2a antibody ratio found in w.t. mice following immunization of DN TNFR25-tg mice with DNP-KLH (FIG. 4).

[0137] Next, it was determined whether DN TNFR25-tg mice had an altered response in the lung inflammation model. Compared to w.t. mice, DN TNFR25-tg mice upon airway challenge exhibited significantly diminished eosinophilic infiltration in the BALF, absent lung inflammation and diminished IgE production in serum when examined by histopathology and PAS staining (FIG. 6e-g). Restimulating bronchial draining lymph nodes from airway challenged DN TNFR25 tg mice with ovalbumin showed diminished Th2 cytokine production but normal IFN- γ production (FIG. 6h). The data suggest that TNFR25 plays a critical role in pulmonary immune responses.

[0138] To validate the genetic data of TNFR25 blockade in w.t. mice, monoclonal antibodies to murine TL1A were developed. Using TNFR25 and TL1A transfected cells (FIG. 7 a,b), TL1A blocking antibodies that abrogated TNFR25 signaling were identified. TL1A transfected cells express TL1A on their surface (FIG. 7c) and secrete TL1A into the supernatant similar to other TNF superfamily members. TL1A containing supernatants caused rapid 51Cr release from FL TNFR25 or Δ 5,6 TNFR25 transfected tumor cells through apoptosis as reported previously (Chinnaiyan, A. M. et al. *Science* 274, 990-2. (1996); Kitson, J. et al. *Nature* 384, 372-5. (1996); Screamton, G. R. et al. *Proc Natl Acad Sci USA* 94, 4615-9. (1997); Bodmer, J. L. et al. *Immunity* 6, 79-88. (1997); Marsters, S. A. et al. *Curr Biol* 6, 1669-76. (1996); Tan, K. B. et al. *Gene* 204, 35-46 (1997)). (FIG. 7d). One of the anti-TL1A antibodies, L4G6, completely blocked TL1A mediated lysis of TNFR25-transfected cells, while several other anti-TL1A antibodies either had no effect or mediated only incomplete inhibition of lysis (FIG. 7d). L4G6 antibody therefore was selected for use of in vivo blockade of TL1A to block TNFR25 signaling in genetically unmodified wild type mice. Mice were immunized twice with ovalbumin in alum as usual. One day prior to airway challenge and for the next three days thereafter 50 μ g L4G6 was injected i.p. and

then the mice were analyzed (FIG. 7e). Controls received the same amount and schedule of hamster IgG. L4G6 administered in this way during and after the period of aerosol challenge phase inhibited eosinophil exudation into BALF, blocked excessive mucus production and diminished the Th2 cytokine production of bronchial lymph node cells upon rechallenge with ovalbumin *in vitro* (FIG. 7e-g).

[0139] Blockade of lung inflammation by anti-TL1A during aerosol challenge is evidence of TL1A expression in the airways. It was found that modest levels of TL1A are expressed on a subpopulation of CD11c+ cells in bronchial draining lymph nodes after airway challenge but not prior to airway challenge (FIG. 7h). All other cell populations in bronchial lymph nodes were TL1A negative before and after aerosol challenge (FIG. 7i). Inguinal lymph nodes did not express TL1A on CD11c+ cells or any other cell type at any time before or after ovalbumin priming or airway challenge. TL1A expression, however, can be induced *in vitro* within 24 h on purified CD4+ and CD8+ spleen or lymph node cells by activation with anti-CD3 and anti-CD28 (FIG. 7j). LPS activated, proliferating B cells do not express TL1A (FIG. 7j).

[0140] 6. DN TNFR25-Transgenic NKT Cells Fail to Support Lung Inflammation in Antigen Primed and Aerosol Challenged NKT Deficient Mice.

[0141] It has been shown that adoptive transfer of w.t. NKT cells to NKT deficient mice restores lung inflammation and airway hyper reactivity in the ovalbumin model and that IL-13 production by NKT cells is required (Akbari, O. et al. *Nat Med* 9, 582-8 (2003); Lisbonne, M. et al. *J Immunol* 171, 1637-41 (2003); Meyer, E. H. et al. *Proc Natl Acad Sci U S A* 103, 2782-7 (2006)). NKT cells have also been implicated in the pathophysiology of asthma patients (Sen, Y. et al. *J Immunol* 175, 4914-26 (2005). Akbari, O. et al. *N Engl J Med* 354, 1117-29 (2006)). To determine whether TNFR25, which is constitutively expressed on NKT cells (FIG. 1a), is involved in triggering lung inflammation, transferred wild type and DN TNFR25-tg NKT cells were adoptively transferred into ovalbumin primed NKT-deficient mice (Cui, J. et al. *Science* 278, 1623-6 (1997)) ($\text{Ja}18$ k.o.) (FIG. 8). While adoptively transferred w.t. NKT cells restored lung inflammation upon airway antigen challenge, the same number of DN TNFR25-tg NKT cells was unable to do so. The data demonstrate that TNFR25 signals in NKT cells are critical for triggering lung inflammation during airway antigen exposure of sensitized mice.

[0142] 7. TNFR25 Agonists as Direct Potentiators of Anti-Tumor Immune Responses

[0143] Mature dendritic cells carry out an important process referred to as "cross-presentation" that enables them to effectively prime T-cytotoxic cells that are specific for tumor-specific peptides. Tumor antigens are degraded and are presented on MHC Class I proteins to circulating CD8 T cells.

[0144] To demonstrate that TNFR25 agonists are effective tumor vaccine BRMs, mice were injected with EG7 tumor cells and OT-I cells. OT-I cells were then observed for clonal expansion. EG7 are EL4 mouse ascites lymphoma lymphoblast cells that have been genetically altered to express ovalbumin, the major protein constituent of chick egg white. Mice were inoculated with EG7 cells and received an

adoptive transfer of ovalbumin-specific T-cell receptor transgenic cells (OT-I). OT-I cells act as indicator cells in vivo by responding to the tumor specific ovalbumin antigen. Theoretically, the mouse's dendritic cells present the tumor-specific ovalbumin to and activate the ovalbumin-specific (OT-I) T cells. However, under these circumstances and as is often seen in human tumor vaccine trials, OT-I cells react with energy to the EG7 tumor cells.

[0145] In contrast, co-transfected EG7 cells with a construct that encodes a secreted version of the heat shock protein gp96 (gp-96-Ig) provided for a tumor secreting gp961g and containing ovalbumin as surrogate antigen. Exposure of the CD8 OT-I cells to the secreted chaperone heat shock protein in combination with the tumor specific ovalbumin resulted in an expansion of OT-I from an initial frequency of 0.5% to over 50% of all CD8 cells following primary injection and one boost with gp961g secreting. Therefore, cross priming of CD8 cells by gp96-Ig-ovalbumin compared to cross-priming by intact ovalbumin protein is enhanced 10,000 to 1,000,000 fold. See Example 19 and FIGS. 10 and 12. See also Am J Reprod Immunol. 2002 October;48(4):220-5.

[0146] When OT-I CD8 cells were cross-primed by gp96-Ig-ovalbumin, in the presence of the TNFR25 agonist antibody 4C12, OT-I CD8 expansion increased by an additional 10-fold over a control antibody. However, when OT-I CD8 cells were cross-primed by gp96-Ig-ovalbumin, in the presence of the TL1A blocking antibody L4G6, OT-I CD8 expansion was decreased by an 10-fold over a control antibody. See FIG. 11.

[0147] As such, TNFR25 agonists are effective biological response modifiers for tumor vaccines because they boost T cell activation and the cellular immune response to a tumor specific antigen, whereas TNFR25 antagonists blocked or inhibited T cell activation. Therefore, another aspect of the invention relates to methods and therapeutic agents that increase the effectiveness of a tumor vaccine.

[0148] Tumor vaccines attempt to the use of elements of the body's natural immune system to fight cancer. Tumor vaccines contain one or more tumor specific antigens and may contain an adjuvant and biological response modifiers. A tumor specific antigen is a polypeptide that is substantially limited to expression in or on tumor cells and which can be used to stimulate an immune response intended to target those tumor cells. Different types of vaccines are used to treat different types of cancer. For an antigenic composition to be useful as a vaccine, an antigenic composition must induce an immune response to the antigen in a cell or tissue. As used herein, an "antigenic composition" may comprise an antigen (e.g., a peptide or polypeptide), a nucleic acid encoding an antigen (e.g., an antigen expression vector), or a cell expressing or presenting an antigen. See U.S. Pub. No. 2003/0185840, which is hereby incorporated by reference in its entirety.

[0149] Biologic response modifiers (BRM), which have been shown to upregulate T cell immunity or downregulate suppressor cell activity. Such BRMs include, but are not limited to, Cimetidine (CIM; 1200 mg/d) (Smith/Kline, PA); low-dose Cyclophosphamide (CYP; 300 mg/m²) (Johnson/Mead, NJ), cytokines such as g-interferon, IL-2, or IL-12 or genes encoding proteins involved in immune helper functions, such as B-7.

[0150] In one embodiment of this aspect of the invention, the tumor vaccine composition includes a tumor antigenic composition and a TNFR25 agonist. In another embodiment, the TNFR25 agonist is the antibody 4C12. In the preferred embodiment, a TNFR25 agonist is added to a tumor vaccine as a biological response modifier. Even more preferably, the TNFR25 agonist is the antibody 4C12. In another embodiment the tumor vaccine includes an adjuvant.

[0151] Tumor vaccine adjuvants may include IL-1, IL-2, IL-4, IL-7, IL-12, gamma-interferon, GM-CSF, BCG, aluminum hydroxide, MDP compounds, such as thur-MDP and nor-MDP, CGP (MTP-PE), lipid A, and monophosphoryl lipid A (MPL). RIBI, which contains three components extracted from bacteria, MPL, trehalose dimycolate (TDM) and cell wall skeleton (CWS) in a 2% squalene/Tween 80 emulsion is also contemplated. MHC antigens may even be used. Exemplary, often preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvants and aluminum hydroxide adjuvant.

[0152] The preparation of vaccines which contain peptide sequences as active ingredients is generally well understood in the art, as exemplified by U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all incorporated herein by reference. Typically, such vaccines are prepared as injectables. Either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccines.

[0153] 8. TNFR25 Immunotoxins as Indirect Potentiators of Innate Anti-Tumor Immune Responses

[0154] Many scientists and corporations work on CD4+/CD25+T regulatory cells (Tregs) because of their immense potential impact on the treatment of many diseases. Although many people are exposed to the same environmental allergens and sensitization to allergens is common, only a fraction of people develop allergic diseases such as asthma. The reason for this is unclear at present but could be related to the presence of efficient regulatory Tregs that suppress airway inflammation in healthy allergen exposed individuals. As such, Tregs are known contribute to the maintenance of peripheral tolerance against self and non-self. However, Tregs have also been documented to impede the body's ability to fight cancer. In those cases, Tregs interfere with the body's tumor-killing immune cells. As such, Tregs function as dedicated suppressor cells and may play a role in preventing tumors, e.g., in squamous cell carcinoma of the head and neck, from being recognized by the immune system. See Br J Cancer, 2005 Mar. 14;92(5):913-20.

[0155] The Inventors have observed that Tregs have properties that suggest that they are activated. Other TNF Receptors have been reported to be expressed on Tregs. GITR is

expressed by activated Tregs. Its ligation has been found to abolish the inhibitory activity of Tregs (Nocentini et al., Eur J. Immunol. 2005 April;35(4):1016-22). TNFR25 has many properties that could make it a versatile regulator of Tregs cells. a) TNFR25 protein expression is rapidly up-regulated by PKC induced mRNA splicing; b) Several functional splice versions including a decoy receptor and a dominant negative form allow fine tuning of regulation; c) TL1A expression appears to be highly regulated on Crohn's Disease; in addition TL1A expression on activated lymphocytes allows sensing of lymphocyte density. Finally, they exert their regulatory function at least in part by secreting IL-10 and IL-13, a cytokine triggered by TNFR25 signals. The inventors are the first to note that FoxP3 expressing, CD4+/CD25+ cultured Tregs express inordinately high TNFR25 levels. As such, the Inventors conclude that TNFR25 modulates Treg function and that TNFR25 can be used as a molecular tag to deplete Tregs *in vivo*. See Example 20. Moreover, TNFR25 agonists abolish Treg Inhibition. See FIG. 15.

[0156] Therefore, another aspect of the invention relates to methods and therapeutic agents that are useful in increasing the potency of anti-cancer therapies by depleting a patient of CD4+/CD25+T regulatory cells (Tregs). In one embodiment of this aspect of the invention, an immunotoxin used to deplete Tregs. In this embodiment, the immunotoxin has an antigen binding portion that is specific for TNFR25; and is conjugated to a toxic agent. In an alternate embodiment, the patient is provided with soluble TL1A conjugated to a toxic agent. Yet another embodiment relates to a chemotherapeutic composition having a chemotherapeutic agent and a TNFR25-specific immunotoxin. Still another embodiment relates to a chemotherapeutic composition having a chemotherapeutic agent and a TL1A conjugated to a toxic agent.

[0157] Another aspect of the invention relates to methods and therapeutic agents that are useful in increasing the potency of anti-cancer therapies by providing a patient with TNFR25 agonists reduce inhibition mediated by CD4+/CD25+T regulatory cells (Tregs).

[0158] 9. TNFR25 Agonists as Anti-Inflammatory Agents

[0159] Using a dominant negative form of TNFR25 (DN-TNFR25) lacking the intracellular death domain and an alternatively spliced form of TNFR25 (delta 5,6-TNFR25) lacking exon 5 and 6 encoding the fourth extra-cellular cysteine rich domain, the Inventors found that TNFR25 function is required to restore homeostatic balance after a mucosal insult. Specifically, the inventors transgenically expressed a dominant negative form of TNFR25 (DN-TNFR25) under the CD2 promoter in mice. The mice were given dextran sodium sulfate (DSS) to induce colitis as a model for human Crohn's disease. Wild type C57B1/6 mice developed colitis and diarrhea and lost weight after 5 days of drinking water with 2% DSS. However, if restored to normal water, wild type mice recovered within a week and regained weight, whereas DN-TNFR25-tg expressing mice acquired disease in a similar manner to wild type mice but diarrhea appeared more severe. Additionally, restoration to normal water did not result in their recovery. Instead all DN-TNFR-tg expressing mice continued to loose weight and died within the second week. Mice expressing functional transgenes of TNFR25 (full length TNFR25 or a splice version Δ5,6-TNFR25) recovered in a manner similar to wild type mice. See Examples 21 and 22 and FIG. 14.

[0160] Current treatment of Crohn's disease uses anti inflammatory agents, immunosuppressives, and TNF inhibitors. All these are symptomatic. Stimulating TNFR25 signaling moves one step up in the pathogenetic chain of events. Activated TNFR25 stimulates IL-10 and IL-13 production, which, in turn, stimulates TGF-beta production resulting in restoration of homeostasis. Such a treatment is curative or approaching cure.

[0161] Therefore, another aspect of the invention relates to a method of treating inflammatory bowel disease in a patient by administering to a patient a therapeutic amount of a TNFR25 agonist. Another embodiment relates to a method of treating inflammatory bowel disease in a patient by raising IL-10 levels in the mucosal areas in the intestine. In another embodiment, the TNFR25 agonist is the antibody 4C12. In a further embodiment, the TNFR25 agonist is the a soluble form of TL1A. In the preferred embodiment, the inflammatory bowel disease is Crohn's disease.

[0162] Given this anti-inflammatory activity of TNFR25 agonists, it is another aspect of the invention to provide methods and therapeutic agents to a patient requiring reduction of inflammation. In one embodiment, a patient is provided a composition containing a TNFR25 agonist to decrease inflammation and promote healing. In one embodiment, a patient is provided with a composition containing the TNFR25 agonist 4C12. Such embodiments are useful for alleviating the symptoms of disorders mediated by chronic inflammatory responses at the cellular level, including cardiovascular diseases (e.g., atherosclerosis), autoimmune diseases including systemic lupus erythematosus (SLE), multiple sclerosis (MS), diabetes (especially type I diabetes), ankylosing spondulitis, arthritis (particularly rheumatoid arthritis), asthma and allergy, bone resorative disorders, ophthalmological disorders including retinopathies, and fibrotic diseases.

[0163] 10. TNFR25 Antagonists as Immunosuppressives

[0164] Many organs and tissues are now routinely transplanted from one human to another. Except for the rare cases where the donor and recipient are monozygotic "identical" twins, such grafts are called allografts. Tissue matching for the transplantation of tissues from one individual to another is critical because a tissue recipient will mount a strong humoral and cellular immune reaction against all non-self proteins. Tissue typing involves identifying MHC antigens on both donor and recipient cells and using donor cells with as many MHC alleles identical to those of the recipient as possible. Matching MHC Class I (especially HLA-B) and Class II HLA-DR alleles is more important for successful transplantation than matching other MHC antigens; and matching MHC is more important than matching minor histocompatibility antigens.

[0165] HLA matching improves graft survival but does not prevent rejection, even in MHC-identical siblings (except for identical twins). Allogeneic MHC is recognized by either CD8 T cells (Class I) or CD4 T cells (Class II); up to 10% of T cells can recognize a given allogeneic MHC because it resembles self MHC+ foreign peptide.

[0166] Improved success in transplantation is due to growing technical expertise, increasing availability of transplant centers to do HLA matching and minimize organ delivery time, and the increased availability of immunosuppressive

drugs (cyclosporin and tacrolimus) that block T cell activation to alloantigens. Still problematic are shortages of organs, the ability of existing disease to destroy the transplanted organ (diabetes and HBV infection are two examples), side effects of immunosuppressive drugs and high cost.

[0167] Given the major side effects associated with current immunosuppressives that block T cell activation and the observation that TNFR25 antagonists such as TL1A blocking antibody L4G6, make effective inhibitors of cognate CD8 T cell clonal expansion (see FIG. 11, Example 19), another aspect of the invention involves the use of TNFR25 antagonists for the facilitation of tissue transplantation to prevent tissue rejection. In one embodiment, TNFR25 antagonists are provided to a transplant recipient to suppress the clonal expansion of CD8 T cells that carry alloantigen-specific T cell receptors (TCRs) and to relieve suppression of T-reg by TNFR25. In another embodiment, TNFR25 antagonists are provided to a transplant recipient in combination with an immunosuppressive agent.

EXAMPLES

Example 1

Media and Reagents

[0168] Cells were cultured in Iscove's Modified Dulbecco's Minimal Essential Medium (Invitrogen) supplemented with 10% heat-inactivated FBS (Invitrogen), 10 µg/ml gentamycin (Invitrogen), and 50 µM β-mercapto-ethanol (Bio-Rad). Monoclonal anti-mouse CD3 and anti-human CD3 were purified from culture supernatants of the 2C11 and the OKT3 cell lines, respectively (ATCC, Manassas, Va.). Monoclonal anti-mouse CD28 and anti-human CD28 were purchased from eBioscience (San Diego, Calif.). ConA, PHA, and LPS were purchased from Sigma (St. Louis, Mo.). Recombinant murine IL-2 was from BioSource International (Camarillo, Calif.). PMA, ionomycin, H7, and cycloheximide were purchased from Calbiochem (San Diego, Calif.).

[0169] Directly conjugated monoclonal antibodies, including FITC-CD4, Cychrome-CD4, PE-CD8a, Cychrome-CD8a, FITC-B220, PE-B220, FITC-CD25, PE-CD11c, PEDX5, FITC-CD3, PE-NK1.1, PE-Annexin V and 7-AAD were purchased from BD/PharMingen (San Diego, Calif.). Hamster IgG control was purchased from eBioscience.

Example 2

Generation of Monoclonal Antibodies Against mTNFR25 and mTL1A

[0170] Armenian hamsters were immunized intraperitoneally three times biweekly with 50 µg of mTNFR25-Ig or mTL1A-MBP (maltose binding protein) in Freund's adjuvant. Three days prior to the fusion, hamsters were boosted with 50 µg of the respective proteins intravenously. Hamster splenocytes were fused with the murine myeloma SP20 with PEG and then plated in methylcellulose-based medium for two weeks using ClonaCell-HY kit (StemCell Technologies Inc., BC, Canada). One thousand colonies were picked and analyzed by ELISA in plates coated with the immunizing fusion protein or control protein-Ig fusion protein. Supernatants from positive clones were tested for the ability to

detect mTNFR25 isoforms in transfected cells by flow cytometry and western blotting. Antibodies were purified from a Nutridoma-SP (Roche, Indianapolis, Ind.) supernatant on a protein G column, dialyzed into PBS and filter sterilized.

Example 3

Flow Cytometric Analysis for the Expression of mTNFR25 and mTL1A

[0171] Single cell suspensions were prepared from lymphoid organs indicated in the individual experiment. Prior to staining, cells were treated with purified anti-mouse CD16/CD32 (Fc-γIII/II receptor; BD) and purified human IgG (Jackson ImmunoResearch, West Grove, Pa.) to block non-specific binding to FcRs. Cells were stained with Armenian hamster anti-mouse TNFR25 or anti-mouse TL1A for 30 minutes at 4° C. Cells were washed in FACS buffer (PBS containing 0.5% BSA and 2 mM EDTA) and then stained with biotin-labeled goat anti-Armenian hamster IgG (Jackson ImmunoResearch) for 30 minutes at 4° C. Cells were washed and then stained with Streptavidin-PE or Streptavidin-Cychrome (BD) for 30 minutes at 4° C. Cells were washed and then stained with directly conjugated cell surface markers for distinct cell populations. Samples were analyzed using a Becton Dickinson FACS LSR instrument and CELLQuest™ software.

Example 4

RT-PCR

[0172] Identification of splice forms of mTNFR25: Messenger RNA was extracted from murine cell lines or tissues with the Micro Fast-Track kit (Invitrogen, Carlsbad, Calif.) and cDNA was reverse transcribed using the Superscript II kit (Invitrogen). RT-PCR products were subcloned into the PCR II vector using the TOPO cloning kit (Invitrogen) and were confirmed as splice forms of mTNFR25 by DNA sequencing. For splicing analysis of murine TNFR25 the following primers were used. Upstream primer, exon 2: CAG TGA GTC CCA GAA GAG GT (SEQ ID NO: 8); downstream primer, exon 7: GGA TAG CCC CAA AAA GGA AC (SEQ ID NO: 9); upstream primer, exon 7: TCC TTT TTG GGG CTA TCC TG (SEQ ID NO: 10); downstream primer, exon 10: GGT ATT TCT CCA TGA CGC TT (SEQ ID NO: 11).

[0173] Activation-induced alternative splicing of human TNFR25 was analyzed because in the mouse the PCR products of different splice forms were more difficult to distinguish on agarose gels but appeared similar to human splice forms. The following PCR primers were used. Upstream exon 4: TTC ACC CTT CTA CTG CCA AC (SEQ ID NO: 12); downstream, exon 7: TAA CCA GGG GCT TGT GAG GC (SEQ ID NO: 13). Human peripheral blood mononuclear cells were isolated from healthy donors by Ficoll Hypaque density gradient centrifugation. Five million cells per sample were activated with PHA (5 µg/ml), or immobilized anti-CD3 (OKT3, 5 µg/ml) and anti-CD28 (1 µg/ml), or PMA (10 ng/ml) and ionomycin (400 ng/ml). The cells were harvested at the indicated time points and mRNA was extracted and converted to cDNA using the Invitrogen kit. Human β-actin was used as internal control. Quantitation of PCR products was done with the aid of Molecular Analyst software (BioRad).

Example 5

Generation of Transgenic Mice

[0174] Murine TNFR25 constructs were cloned under the human CD2 promoter and local control region (gift from Dr. A. Singer, NIH) using the restriction endonuclease sites EcoR I and Sal I. Three mTNFR25 constructs were generated by PCR using a proofreading enzyme. The constructs were the full length molecule of murine TNFR25 (FL TNFR25), the TNFR25 splice variant lacking the 5th and 6th exon (Δ 5,6 TNFR25), and the dominant negative version of TNFR25 (DN TNFR25, as 1-234) terminating at the end of the transmembrane domain and lacking the entire intracellular domain. The sequence of the PCR products was confirmed by sequencing. Microinjections of DNA into the fertilized eggs were performed by the transgenic facility at the University of Miami, School of Medicine. Potential founders were screened by PCR of DNA from tail biopsies. The primer pair was located upstream and downstream of the cloning sites, therefore the same primer pair was used for all the mTNFR25 transgenes. The upstream primer is 5' CGC TCT TGC TCT CTG TGT ATG 3' (SEQ ID NO: 14) and the downstream primer is 5' CTG CCA GCC CTC TTC CAT C3' (SEQ ID NO: 15). Transgenic mice were bred into the C57BL/6J background by serially mating hemizygous transgenic animals with wild-type C57BL/6J (Jackson Laboratories, Bar Harbor, Me.). All mice were used at 6-12 weeks of age and were maintained in pathogen-free facilities. The University of Miami Animal Care and Use Committee approved all animal use procedures.

Example 6

Nuclear Extract Preparation and Electrophoretic Mobility Shift Assays for NF κ B Activation

[0175] One hundred and seven of EL4- Δ 5,6 TNFR25 or EL4-FL TNFR25 cells were treated with soluble or membrane bound TL1A or with the TNFR25 agonistic antibody 4C12 as indicated in the figure legends and then collected by centrifugation at 800 g for 5 min. Nuclear extracts were isolated using a minipreparation protocol and subjected to EMSA as described (Harhaj, E. W. et al. *Virology* 333, 145-58 (2005). Nuclear extracts (6 μ g) were incubated at room temperature for 20 min with a 32P-labeled high-affinity κ B probe, followed by resolving the DNA-protein complexes on native 5% polyacrylamide gels.

Example 7

T Cell Proliferation Assay

[0176] Splenocytes were plated in triplicate at 1×10^5 cells/well in 96-well flat-bottom plates. Cells were activated with immobilized anti-CD3 (2 μ g/ml) with or without soluble anti-CD28 (1 μ g/ml), or with ConA (5 μ g/ml) or with PMA (1 μ g/ml) and ionomycin (400 ng/ml). For T cell proliferation, purified CD4 T cells at 1×10^5 cells/well or CD8 T cells at 5×10^4 cells/well were stimulated with coated anti-CD3 (2 μ g/ml) and soluble anti-CD28 (1 μ g/ml). Recombinant mIL-2 was added to the culture at 1000 U/ml in indicated experiments. Cells were cultured for 72 hr and pulsed for the last 6 hours of incubation with 1 μ Ci/well of 3H-thymidine (Perkin Elmer, Boston, Mass.), and thymidine incorporation was quantitated using a scintillation counter.

[0177] Murine CD4 or CD8 or T cells were purified from splenocytes and/or lymph nodes by negative selection using SpinSep kit (StemCell Technology Inc.) according to the manufacturer's protocol. The purity was routinely around 90%-96% examined by FACS analysis.

Example 8

Immunization of Mice with DNP-KLH, Antibody Isotype Determination, and Cytokine ELISA

[0178] Adult (6-10 wk old) transgenic and wt mice were immunized intraperitoneally with 100 μ g DNP-conjugated keyhole limpet hemocyanin (DNP-KLH) (CalBiochem). One week and three weeks after immunization, mice were bled and serum was separated for analysis of anti-DNP specific IgG1, IgE, and IgG2a antibodies by ELISA according to manufacturer's protocol (BD). Sera from individual animals were absorbed to 96-well plates coated with 0.8 μ g/ml DNP-albumin (DNP-BSA) (CalBiochem) and the isotype of bound antibody determine by ELISA.

[0179] To examine the cytokine production in the supernatants of cell cultures, sandwich ELISAs were performed per the manufacturer's instructions. Antibody pairs from BD were used for IL-2, IFN- γ , and IL-4 analysis. Reagents for IL-13 ELISA were purchased from R&D Systems (Minneapolis, Minn.) and reagents for IL-5 and IL-10 ELISA were purchased from eBioscience.

Example 9

In vitro polarization CD4 T cells into Th1 or Th2 cells

[0180] CD4 T cells were purified by negative selection as mentioned above. CD4 T cells were activated with immobilized anti-mCD3 (2 μ g/ml) and soluble anti-mCD28 (1 μ g/ml) alone, or in the presence of IL-12 (10 ng/ml) and anti-mIL-4 (20 μ g/ml) for Th1 differentiation, or with IL-4 (10 ng/ml), anti-mIFN- γ (10 μ g/ml), and anti-mIL-12 (10 μ g/ml) for 7 days. The cells were harvested, washed and replated at 1×10^5 cells per well and restimulated with immobilized anti-CD3. After 24 hours the supernatants were harvested and evaluated for cytokine production by ELISA.

Example 10

Immunization Protocols for the Murine Model of Allergic Asthma

[0181] DN TNFR25-tg (encoded by SEQ ID NO: 16), Δ 5,6 TNFR25-tg, and FL TNFR25-tg mice generated as described before and backcrossed at least seven generations into the C57BL/6J background were compared with wild-type C57BL/6J mice purchased from National Cancer Institution (Frederick, Md.). Mice were sensitized by intraperitoneal injection of 66 μ g ovalbumin (crystallized chicken egg albumin, grade V; Sigma) absorbed to 6.6 mg aluminum potassium sulfate (alum; Sigma) in 200 μ l PBS on day 0. On day 5, mice were boosted intraperitoneally with the same dose of ovalbumin in alum. On day 12, mice were aerosol challenged with 0.5% ovalbumin in PBS for one hour using an Ultrasonic Nebulizer (MABIS Healthcare Inc., Lake Forest, Ill.). Mice were assessed for allergic inflammation of the lungs three days after the single aerosol exposure. Mice were sacrificed by inhalation of CO₂. After cannulation of

the trachea the lung was lavaged 4 times with 1 ml of PBS. Cells recovered from the BAL fluid were counted and used for cytopsin preparations (50,000 cells or fewer/slide). >200 cells were counted for each cytopsin slide stained with Wright-Giemsa stain (Sigma) to determine differential cell counts for macrophages, eosinophils, lymphocytes, and neutrophils.

Example 11

Lung Histology

[0182] Lungs were removed from mice after the bronchial lavage procedure and fixed in 10% neutral buffered formalin. Samples were submitted to the Histopathology Core of the Sylvester Cancer Center at the University of Miami School of Medicine where specimens were embedded, sectioned, and stained with haematoxylin and eosin. Sections were also stained with periodic acid-Schiff (PAS) to determine mucus production.

Example 12

ELISA for Serum Total IgE and Ovalbumin-Specific Ig

[0183] Mice were bled before sensitization (day 0), 3 days after aerosol challenge (day 15) and in some experiments one day before aerosol challenge (day 11). The total IgE level was quantitated by ELISA according to the manufacturer's protocol (BD). Ovalbumin specific IgE was determined in Sandwich ELISA by first coating plate with 0.01% OVA in PBS, followed by loading diluted serum samples and then the secondary biotin-labeled anti-IgE antibody (BD).

Example 13

In vitro Restimulation of Bronchial Lymph Node Cells and Cytokine Production

[0184] One day or three days after aerosol challenge, bronchial lymph nodes were harvested and single cell suspensions were prepared. Cells were seeded into round-bottom 96-well plates at 1×10^6 cells/well and cultured with 100 μ g/ml ovalbumin for 4 days. Then supernatants were collected for cytokine ELISA assays as described.

Example 14

Cytotoxicity Assay

[0185] Serially diluted soluble mTL1A supernatants harvested from P815-TL1A transfected cells were added to ^{51}Cr -labeled P815-TNFR25 transfected target cells. To test for TL1A blocking activity different anti-TL1A monoclonal antibodies were added into the culture and Cr release determined after 4 hours in triplicate samples. Spontaneous release was calculated from the wells that contained only ^{51}Cr -labeled target cells. 100% release (positive control) was calculated from the wells that contained 51Cr-labeled target cells and 1% SDS. The percentage of cytotoxicity activity was calculated as the following: (mean readout of sample—mean readout of spontaneous release)/mean readout of positive control. Similar data were also obtained with EL4-transfetsants.

Example 15

Blocking of Lung Inflammation by Antagonistic Anti-mTL1A Antibody

[0186] Mice were sensitized intraperitoneally with ovalbumin in alum on day 0 and day 5 followed by aerosol challenge with 0.5% ovalbumin in PBS for one hour on day 12. Mice were given L4G6 or an equivalent amount of the control hamster IgG (Jackson Immuno Research) by intraperitoneal injection of 50 μ g/mouse each day from day 11 to 14. Allergic lung inflammation was evaluated on day 15.

Example 16

Adoptive Transfer of NKT Cells

[0187] J α 18 k.o. mice (Cui, J. et al. Science 278, 1623-6 (1997)) were a gift from Michael Lotze (U. Pittsburgh) with kind permission from M. Taniguchi (Ciba University, Japan). NKT cells from w.t. and DN TNFR25-tg mice were isolated from pooled spleen cells from 10 mice by positive selection using the EasySep mouse Pan NK Positive Selection Kit (StemCell Technologies, Vancouver, Canada) according to the manufacturer's instructions.

Example 17

Statistical Analyses

[0188] Statistical analyses using a two-tailed Student's t test were performed with the GraphPad Prism Software (San Diego, Calif.); $p < 0.05$ is considered significant. Data in the text are presented as the mean \pm SEM.

Example 18

Generation of DR3 and TL1A Antibodies

[0189] A DR3-Ig fusion protein was generated, purified and used to immunize hamsters. Hybridoma supernatants were obtained and screened by ELISA using the DR3-Ig fusion protein as a screening agent. The nature of the hybridomas was verified by flow cytometry of DR3 transfected tumor cells, by Western blots, and by functional studies. All of the antibodies detected full-length and alternatively spliced DR3 on transfected cells by FACS, one of the antibodies detected DR3 in Western blots, and the antibody (4C12) displayed agonistic activity, mediating DR3 signaling in the absence of TL1A.

[0190] TL1A monoclonal antibodies were obtained by immunizing hamsters with a TL1A-maltose-binding-protein fusion. The TL1A antibodies detected transfected TL1A by flow cytometry. The antibody (L4G6) displayed antagonistic activity, blocking TL1A binding to DR3.

Example 19

Signaling Through TNFR25 Enhances CD8 Cross-Priming

[0191] A novel heat shock protein gp96 based system that mediates strong, antigen specific CD8-CTL expansion in vivo was recently described in Strbo et al., Am J Reprod Immunol. 2002 October;48(4):220-5. In this model system released gp96-Ig (engineered to be secreted) activates dendritic cells and provides chaperoned peptides for cross

presentation to and cross priming of CD8 cells (FIG. 12). The system is very useful because it is independent of CD4 help. Secreted gp96 provides the activation signal for DC through CD91 and TLR2/4 which is otherwise provided by CD40-L on CD4 cells. Accordingly, CD8 (OT-I) expansion in this system works well in CD40-L deficient mice. This has been employed to study mucosal immunity and to determine the role of TNFR25 in CD8 expansion.

[0192] EG7-gp96 is a cell line derived from the EL4 lymphoma by transfection with ovalbumin and gp96-Ig. The cells secrete gp96-Ig associated with ovalbumin peptides. Ovalbumin-peptides chaperoned by gp96-Ig enhance cross-priming of CD8 cells (FIG. 10) by about 10,000 fold when compared to ovalbumin alone. 100 ng ova-gp96-Ig expand OT-I in B6 mice from a frequency of 0.5% among CD8 cells post transfer to 20% in the spleen after EG7-gp96-Ig immunization.

[0193] In order to determine the effect of TNFR25 signals on CD8 expansion TCR transgenic OT-I model as described above, were used together with EG7-gp96-Ig mediated stimulation. To determine the effect of TNFR25 signals the mice received an agonistic anti TNFR25 antibody (4C12), a TNFR25 binding but not agonistic antibody (L4G6) or a control IgG 24 h and 72 h after of EG7-gp96-Ig immunization. OT-I expansion was monitored in the peritoneal cavity on day 5 after immunization. 4C 12 caused a increased recruitment of cells into the peritoneal cavity by EG7-gp96-Ig resulting in a doubling of the cell number. In addition 4C12 specifically caused an over 8-fold increase in the expansion of OT-I. The L4G6 anti TNFR25 antibody did not induce increased recruitment of cells to the peritoneal cavity and inhibited OT-I expansion.

[0194] These data show that agonistic anti TNFR25 antibodies costimulate CD8 cells and/or inhibit suppressive effects of Tregs via TNFR25. Costimulation of naïve T cells by TNFR25 results in increased proliferation and Th2 polarization upon secondary activation. In addition signaling of TNFR25 on T regulatory cells results in their temporary inhibition of suppression. The combined effect then is responsible for the increased CD8 expansion and cell recruitment seen in FIG. 11.

Example 20

CD4+ CD25+T Regulatory Cells Express High Levels on TNFR25

[0195] In order to determine the expression of TNFR25 CD4+ CD25 Tregs were purified from spleens by negative selection of CD4 cells followed magnetic sorting with anti CD25. The cells were cultured with anti CD3, anti CD28 beads at a bead to cell ratio of 3:1 and 2000 u/ml human IL-2 was added. Under these conditions the cells will begin to proliferate after 3-4 days and keep expanding for about 3 weeks. The cultured cells were analyzed by FACS analysis for CD4 and CD25, by intracellular cytofluorimetry for FoxP3 expression and for surface analysis of TNFR25. FIG. 13 shows that Tregs obtained in this way are essentially pure and express both FoxP3 and TNFR25.

Example 21

TNFR25-Blockade Causes Lethality of Dextran Sodium Sulfate Colitis

[0196] The dextran sodium sulfate (DSS) model has widely been used as a colitis model resembling in some

aspects Crohn's disease. The initial insult is the damaging effect of DSS on the permeability barrier normally provided by the gut epithelium. This effect of DSS allows access of the normal gut flora to sites in the mucosal immune system that set off an inflammatory immunological reaction resembling Crohn's colitis. Wild type (w.t.) B6 mice during an 8 day course of exposure to drinking water containing 2% DSS develop diarrhea and loose weight. Upon restoration of normal water, B6 mice recover and regain their normal weight. TNFR25 also influences the course of disease in Colitis. In the present experiment, w.t. and transgenic were exposed to mice to 2% DSS water for seven days. As shown in FIG. 14, DN-TNFR25-tg mice developed disease similar to wild type mice, however when normal water was restored, DN-TNFR25-tg mice did not recover as w.t. mice did. Instead DN-TNFR25-tg mice continued to loose weight and died between day 12 and 16. The Δ5,6-TNFR25-tg mice resembled w.t. mice although the death of one mouse could also suggest a disturbed immune response. Two conclusions were reached: In DNTNFR25-tg mice, the ensuing immune response is much stronger than in w.t. mice leading to lethality and the restoration of normal health and homeostasis in w.t. mice is dependent on normally functioning T regulatory (Treg) cells. Treg function is disturbed in DNTNFR25-tg mice. The latter is likely since it is known that Treg function is extraordinarily important to maintain homeostasis in the mucosal immune system by maintaining the correct balance between tolerance to nutrients and normal gut flora and immune response to gut pathogens.

Example 22

Immunization with EG7-gp96-Ig Induced the OT-I Cells to Migrate to Mucosal Sites Peyer's Patches, Lamina Propria Lymphocytes (LPL) and Intraepithelial Lymphocytes (IEL)

[0197] As shown in FIG. 16 (right column), EG7-cells are unable to cause clonal expansion of OT-I or migration to mucosal sites. EG7-cells secreting gp96-Ig on the other hand cause clonal expansion of OT-I in spleen, lymph nodes and peritoneal cavity (not shown) and their migration to mucosal sites (FIG. 16). In Peyer's patches 8% of the cells are CD8+ and 6.7% of the CD8-cells are OT-I; in IEL 61% of the cells are CD8+ and 9% of the CD8 are OT-I. In LPL 29% of the CD8 cells are OT-I. The OT-Icells migrating to IEL after immunization are αEb7+ and α4β7+ but remain CD8αβ and TCRαβ, unlike resident CD8 IEL the majority of which are CD8αα and 50% TCRγδ.

[0198] In this disclosure there are described only the preferred embodiments of the invention and but a few examples of its versatility. It is to be understood that the invention is capable of use in various other combinations and environments and is capable of changes or modifications within the scope of the inventive concept as expressed herein. Thus, for example, those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such equivalents are considered to be within the scope of this invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1
<211> LENGTH: 1665
<212> TYPE: DNA
<213> ORGANISM: Mouse

<400> SEQUENCE: 1

gggagttgtt	ctggatggcg	cggggcgccc	cgggcagcag	ctactctagt	ctaggaacat	60
aggggctgag	ctgggtgggg	aaggccccggg	ttacgcgacc	gaccagagcc	gcactcacaa	120
gggcccaggc	ggtacacacc	gcaatggagg	cacggctgct	gcggggctgc	gtgggtggagc	180
ctctgttcct	accactgtcg	ctgctgtcg	tgctgtcgct	gctgttgggt	ggccaggggcc	240
agggcggcat	gtctggcagg	tgtgactgtg	ccagttagtc	ccagaagagg	tatggcccg	300
tttgttgcag	gggctgccc	aaggcacact	acatgaaggc	cccctgcgca	gaaccctgtg	360
gcaactccac	ctgccttccc	tgtccctcgg	acaccttctt	gaccagagac	aaccacttta	420
agactgactg	tacccgctgc	caagtctgtg	atgaagaggc	ccttcaagtg	acccttgaga	480
actgctcggc	aaagtccggac	accactgtg	gctgccatgc	aggctgggt	gttactgtct	540
ccaccgtgcc	atgtggaaa	agctcacctt	tctcttgggt	cccatgcggg	gctacgacac	600
cagtccatga	ggctccaaacc	ccccggccct	gcctgcctgg	cttctatata	cgtggcaatg	660
actgcacgtc	ctgccccacg	ggcttcagca	gcgtttgc	taaggcttgc	actgtgtct	720
gtggctggaa	gcagatgttt	tgggtccagg	tgcttctagg	agtgcgcgtt	ctttttgggg	780
ctatcctgat	ctgtgcata	tgtcgatggc	agccttgtaa	ggccgtggtc	actgcagaca	840
cagctggac	ggagccccgt	gcctcaccac	agactgccc	tctctcagcc	tcagacagcg	900
cccacccct	cttggcacct	ccaagcgt	ctggaaaat	ctgttaccact	gtccagggtgg	960
taggcaacaa	ctggacccct	ggcttatccc	agactcagga	ggtggcttgc	ggacaggccc	1020
cacaaccctg	ggatcagctg	ccaaacagaa	ctcttggaa	tcctctggca	tctccgtct	1080
cgtccagcgcc	ccctgcgggc	tctccggctg	ctgtgtccca	gcctggcccg	cagctctacg	1140
atgtgatgga	tgcggtccca	gcacgaagg	ggaaggagg	cgtgcgcacg	ctggggctgc	1200
ggaaagcgg	aattgaagcc	gtggagggtgg	aaatctgcg	cttccggagac	cagcgtatg	1260
agatgctcaa	gcgcgtggcgt	cagcagcgc	ctgcaggccat	cggtgccatc	tatgcggctc	1320
tggagcgc	gggtctggaa	ggctgtgc	aggacctgcg	cagccgcctg	cagcgtggcc	1380
cgtatggaa	ggtccatcat	ccactttgac	acccttagtga	cccttgaagg	agccttaagt	1440
attgttacit	atgcgtgtag	acatttatg	tcaattacta	acccctgcc	gtggtctgc	1500
gttagcaggc	tggctgcctc	acttttgc	atctgcagca	cggagcttcc	gctaaaggaa	1560
gcgtcatgga	gaaataccag	aaggggccaa	gtgattgggt	gctcagctgt	taattagccc	1620
gagtttggac	ttggtattaa	atttcataag	aaaagcagct	gcttg		1665

<210> SEQ ID NO 2
<211> LENGTH: 1665
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 2

```

cggggccctgc gggcgcgggg ctgaaggcgg aaccacgacg ggcagagagc acggagccgg 60
gaagcccccgtc ggcccccgtc ggagggctat ggagcagcgg ccgcggggct gcgcggcggt 120
ggcggccggcg ctccctcctgg tgctgctggg ggcccgggcc cagggcggca ctctgtagccc 180
cagggtgtgac tgtgcccgtg acttccacaa gaagatttgt ctgttttgtt gcagaggctg 240
cccagcgggg cactacctga aggccccttg cacggagccc tgccggcaact ccacctgcct 300
tgtgtgtccc caagacacact tcttggcctg ggagaaccac cataattctg aatgtgccc 360
ctgccaggcc tgtgtatgac aggcctccca ggtggcgctg gagaacttgtt cagcagtggc 420
cgacacccgc tggcggctgt a gccaggctg gtttgtggag tgccaggctca gccaatgtgt 480
cagcagttca cccttctact gccaaccatg cctagactgc ggggcccctgc accgcccacac 540
acggctactc tggcccgcg a gataactga ctgtgggacc tgccctgcctg gcttctatga 600
acatggcgat ggctgcgtgt cctgccccac gccacccccc tcccttgcag gacgaccctg 660
gggagctgtc cagagcgctg tgccgctgtc tgtggctgga ggcagagtag gtgtgttctg 720
gttccagggtg ctccctggctg gcttgtgtt ccccttctg cttggggccca ccctgaccta 780
cacataccgc cactgctggc ctcacaagcc cctggttact gcagatgaag ctgggatgga 840
ggctctgacc ccaccacccgg ccacccatct gtcacccatg gacagcgccc acacccttct 900
agcaccccttct gacagcgtg agaagatctg caccgtccag ttgggtggta acagctggac 960
ccctggctac cccgagaccc aggaggcgctg ctgcccgcag gtgacatgtt cctgggacca 1020
gttggccacggc agagctcttg gccccgtgc tgcccccaca ctctcgccag agtccccagc 1080
cggctcgcca gccatgatgc tgccgggggg cccgcagctc tacgacgtga tggacgcgg 1140
ccccagcgccgg cgcttggaaagg agttcgctgc caccgtgggg ctgcgcgagg cagagatcga 1200
agccgtggag gtggagatcg gccgcttccg agaccagcag tacgagatgc tcaagcgctg 1260
gcccgcagcag cagcccgccgg gcccgcggc cgtttacgcg gcccctggagc gcatggggct 1320
ggacggctgc gtggaaagact tgccgcggcc cctgcacgcgc gcccgtgc acggcgccca 1380
cttgcacactt aggccgtctg gtggcccttg cagaagccct aagtacggtt acttatgcgt 1440
gttagacattt tatgtcaattt attaaggccg tggcacggcc ctgcgtacca gacccaggcc 1500
gccccacccccc tgctcgcccc tatcgctcca gccaaggcga agaagcacga acgaatgtcg 1560
agaggggggtg aagacatttc tcaacttctc ggcggagtt tggctgagat cgccgtattt 1620
aatctgtgaa agaaaacaaa acaaaaacaaa aaaaaaaaaa aaaaa 1665

```

<210> SEQ ID NO 3

<211> LENGTH: 759

<212> TYPE: DNA

<213> ORGANISM: Mouse

<400> SEQUENCE: 3

```

atggcagagg agctggggtt gggcttcggg gaaggaggccc cagtggaaatg gctgcggaa 60
ggctgttagac acaggccaga ggccaggccc gggctagctg ccaggagcaa agcctgcctg 120
gctctcacct gctgcctgtt gtcatatccc atcctcgccag gacttagcac cctccataatg 180
gctggccaggc tccgggtccc cggaaaagac tgtatgttc gggccataac agaagagaga 240
tctgagccctt caccacagca agtttactca cctcccagag gcaagccgag agcacacctg 300

```

-continued

acaattaaga aacaaacccc	360
agcaccacat ctgaaaaatc	
agctctctgc tctacactgg	
gaacatgacc tagggatggc	420
cttcaccaag aacgggatga	
agtacatcaa caaatccctg	
gtgatcccag agtcaggaga	480
ctatttcatc tactccaga	
tcacattccg agggaccaca	
tctgtgtgtg gtgacatcg	540
tcgggggaga cgaccaaaca	
agccagactc catcaactgt	
gtcatcacca aggtacgaga	600
cagctaccct gagcctgccc	
gcctactaac agggtccaag	
tctgtgtgtg aaataagcaa	660
caactggttc cagtcctct	
accttggggc catgttctcc	
ttggaagaag gggacagact	720
aatggtaaac gtcagtaca	
tctccttggt ggattacaca	
aaagaagata aaactttctt	759
tggagctttc ttgctataa	

<210> SEQ ID NO 4

<211> LENGTH: 657

<212> TYPE: DNA

<213> ORGANISM: Mouse

<400> SEQUENCE: 4

atggaggcac ggctgtcg	60
ggctgtcggt gtggaggc	
tc ttccatacc actgtgt	
ctgtgtgtc tggtgtgt	120
gttttttttcc cagggcagg	
ggccatgtc tggcagggt	
gactgtgcca gtgagtc	180
cccaaaagaa gaagaggta	
tttttttttggt ggccgtttt	
gttgcagggg ctgccc	
ggccaaaggg ctgccc	
ggccaaaggg ctgccc	240
ggccaaaggg ctgccc	
ccctcgac ctttgc	
ccctcgac ctttgc	300
ccctcgac ctttgc	
ccctcgac ctttgc	
ccctcgac ctttgc	360
ccctcgac ctttgc	
ccctcgac ctttgc	
ccctcgac ctttgc	420
ccctcgac ctttgc	
ccctcgac ctttgc	480
ccctcgac ctttgc	
ccctcgac ctttgc	540
ccctcgac ctttgc	
ccctcgac ctttgc	600
ccctcgac ctttgc	
ccctcgac ctttgc	657

<210> SEQ ID NO 5

<211> LENGTH: 696

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

atggagcagc ggccgcgggg	60
ctgcgcggcg gtggcggcg	
cgctccctct ggtgtgt	
ggggcccgcc cccaggccgg	120
cactcgtagc cccagggtgt	
actgtgcgg tgacttccac	
aagaagattt gtcgttttg	180
ttgcagaggc tgcccagccg	
ggcactacct gaaggccct	
tgcacggagc cctgcggcaa	240
ctccacctgc cttgtgtgtc	
cccaagacac cttttggcc	
tgggagaacc accataattc	300
tgaatgtgc cgctgcagg	
cctgtgtatga gcaggctcc	
cagggtggcgc tggagaactg	360
ttcagcagtg gcccacacc	
gctgtggctg taaggccaggc	
tggtttgtgg agtgcagg	420
cagccaatgt gtcagcagg	
tacccttcta ctgccaacca	
tgcctagact gcggggccct	480
gcaccgcac acacggctac	
tctgttcccg cagagatact	
gactgtggaa cctgcctgcc	540
tggcttctat gaacatggcg	
atggctgtcg gtccctgccc	
acggccacccc cgtcccttgc	600
aggagcacc tggggagctg	
tccagagcgc tgcgcctgt	
tctgtggctg gaggcagagt	660
agggtgttcc tgggtccagg	
tgctccttggc tggccttgg	

-continued

gtccccctcc	tgcttggggc	caccctgacc	tacaca	696	
<210> SEQ_ID NO 6					
<211> LENGTH: 633					
<212> TYPE: DNA					
<213> ORGANISM: Homo sapiens					
<400> SEQUENCE: 6					
atggagcagc	ggccgcgggg	ctgcgcggcg	gtggcggcgg	cgctcctcct	60
ggggcccccggg	cccagggcgg	cactcgttagc	cccaggtgtg	actgtgccgg	120
aagaagattt	gtctgttttg	ttgcagggc	tgcccaggg	ggcactaccc	180
tgcacggagc	cctgcggcaa	ctccacactgc	cttgtgtgtc	cccaagacac	240
tgggagaacc	accataattt	tgaatgtgcc	cgctgccagg	cctgtgtatga	300
caggtggcgc	tggagaactt	ttcagcagt	gccgacaccc	gctgtggctg	360
tggtttgtgg	agtgcgcagg	cagccatgt	gtcagcagtt	cacccttcta	420
tgccttagact	ggggggccct	gcaccggccac	acacggctac	tctgttcccg	480
gactgtggga	cctgcctgcc	tggcttctat	gaacatggcg	atggctgcgt	540
acgccaccccc	cgtcccttgc	aggagcaccc	tggggagctg	tccagagcgc	600
tctgtggctg	gaggcagagt	aggtgtgttc	tgg		633
<210> SEQ_ID NO 7					
<211> LENGTH: 2016					
<212> TYPE: DNA					
<213> ORGANISM: Homo sapiens					
<400> SEQUENCE: 7					
gagagggaaa	agggaaggag	gagactgagt	gattaagtca	cccactgtga	60
ttctattnaa	tgggggcctct	ctctgcccag	gagtcaagg	tgcctccagg	120
gcatggccga	ggatctggga	ctgagctttt	gggaaacagc	cagtgtggaa	180
agcacggcag	ctgcaggccc	aaggccagga	gcagcagcgc	acgctggct	240
gcctgggtt	gctcccttc	cttgcaggac	tcaccacata	cctgcctgtc	300
ggggccaggg	agaggcctgt	gtcagttcc	aggctctaaa	aggacaggag	360
cacatcagca	agtttatgca	cctcttagag	cagacggaga	taagccaagg	420
cagttgttag	acaaaactccc	acacagcact	ttaaaaatca	gttcccagct	480
acatgaact	aggcctggcc	ttcaccaaga	accgaatgaa	ctataccaac	540
tgatcccaga	gtcggggagac	tacttcattt	actcccaggt	cacattccgt	600
ctgagtgca	tgaatcaga	caagcaggcc	gaccaaaca	gccagactcc	660
tcatcacca	ggtaacagac	agctaccctg	agccaaacca	gctcctcatg	720
ctgtatgcg	atgttttttt	aaactgttcc	agcccatcta	cctcgagcc	780
tgcagaagg	ggacaagcta	atggtaac	tcagtgtat	ctctttgtt	840
aagaagataa	aaccttcttt	ggagccttct	tactatagga	ggagagcaaa	900
tgaaagtcc	ctgccaccga	gttcctaattt	ttctttgttc	aatgttaatt	960
gttttcttgg	ggccgggagt	agggggcattt	ccacaggac	aacggtttag	1020
tggggcccaa	aatttcacac	ttcatgtgcc	ttactgtatga	gagtactaac	1080

-continued

tgaagagagc aaatatatta ttaagatggg ttggaggatt ggcgagttc taaatattaa	1140
gacactgatc actaaatgaa tggatgatct actcgggtca ggattgaaag agaaatattt	1200
caacacctcc ctgctataca atggtcacca gtggccagt tattgttcaa tttgatcata	1260
aatttgcctc aattcaggag ctttgaagga agtccaagga aagctctaga aaacagtata	1320
aacttcaga ggcaaaatcc ttcaccaatt tttccacata ctttcatgcc ttgcctaaaa	1380
aaaatgaaaa gagagtttgt atgtctcatg aatgttccaca cagaaggagt tggtttcat	1440
gtcatctaca gcatatgaga aaagctacct ttctttgtat tatgtacaca gatatctaaa	1500
taaggaagta tgagtttac atgtatatac aaaataacaac agttgcttgtt attcagtaga	1560
gttttcttgc ccaccttattt tgtgctgggt tctaccttaa cccagaagac actatgaaaa	1620
acaagacaga ctccactcaa aatttatatg aacaccacta gatacttccat gatcaaacat	1680
cagtcaacat actctaaaga ataactccaa gtcttggcca ggcgcagtgg ctcacacctg	1740
taatcccaac actttggag gccaagggtgg gtggatcatc taaggccggg agttcaagac	1800
caggctgacc aacgtggaga aaccccatct ctactaaaaa tacaaaattt gccggcggt	1860
gtagcgcattg gctgtatcc tggctactca ggaggccgag gcagaagaat tgcttgaact	1920
ggggaggcag aggttgcgtt gagcccagat cgcgcattt cactccagcc tggtaacaa	1980
gagcaaaact ctgtccaaaa aaaaaaaaaa aaaaaa	2016

<210> SEQ_ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized
<400> SEQUENCE: 8

cagtgagtcc cagaagaggt	20
-----------------------	----

<210> SEQ_ID NO 9
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized
<400> SEQUENCE: 9

ggatagcccc aaaaaggaac	20
-----------------------	----

<210> SEQ_ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized
<400> SEQUENCE: 10

tccttttgg ggctatcctg	20
----------------------	----

<210> SEQ_ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized

-continued

<400> SEQUENCE: 11
ggtatttctc catgacgctt 20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized

<400> SEQUENCE: 12
ttcaccccttc tactgccaac 20

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized

<400> SEQUENCE: 13
taaccagggg ctttgaggc 20

<210> SEQ ID NO 14
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized

<400> SEQUENCE: 14
cgctcttgct ctctgtgtat g 21

<210> SEQ ID NO 15
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized

<400> SEQUENCE: 15
ctgcccagcccc tcttccatc 19

<210> SEQ ID NO 16
<211> LENGTH: 702
<212> TYPE: DNA
<213> ORGANISM: Mouse

<400> SEQUENCE: 16
atggaggcac ggctgctgac gggctgcgtg gtggagccctc tggccatacc actgctgctg 60
ctgtctgtgc tgctgcttgg tggccaggcc caggccggca tggctggcag gtgtgactgt 120
gccagtgtgtt cccagaagag gtatggcccg ttttggca ggggctgccc aaaggacac 180
tacatgaagg ccccccgtgc agaaccctgt ggcaactcca cctgccttcc ctgtccctcg 240
gacaccccttct tgaccagaga caaccacttt aagactgact gtacccgctg ccaagtctgt 300
gatgaagagg cccttcaagt gacccttgg aactgctcgg caaagtccgaa cacccactgt 360
ggctgccagt caggctgggtg tggtgactgc tccaccgagc catgtggaa aagctcacct 420
ttctcttgcgt tccccatgcgg ggctacaaca ccagtccatg aggctccaaac ccccccggccc 480

-continued

tgccctgcctg gcttctatat acgtggcaat gactgcacgt cctgccccac gggcttcagc	540
agcggttgcc ctaaggcttg cactgctgtc tgtggctgga agcagatgtt ttgggtccag	600
gtgcttctag gagtcgcgtt ccttttggg gctatcctga tctgtgcata tggctctgga	660
agcgggagcg tcgacgatct ttacgacgat gataaaatgt aa	702

What is claimed is:

1. An isolated antibody that binds a Tumor Necrosis Factor Receptor 25 (TNFR25) antigen wherein said antibody is capable of acting as a TNFR25 agonist.
2. The antibody of claim 1, wherein said antibody is capable of increasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody.
3. The antibody of claim 1, wherein the Tumor Necrosis Factor Receptor 25 (TNFR25) antigen is human or mouse.
4. A Tumor Necrosis Factor Receptor 25 (TNFR25)-specific toxin comprising a toxic agent linked to a polypeptide that binds the TNFR25 receptor.
5. The toxin of claim 4 wherein said polypeptide comprises monoclonal antibody 4C12 or an immunospecific portion thereof.
6. The toxin of claim 4 wherein said toxic agent is selected from the group consisting of a radioactive isotope, ricin, abrin, diphtheria toxin, *Pseudomonas* exotoxin, and metal ion.
7. The toxin of claim 4 wherein said polypeptide comprises a TL1A protein of a TNFR25-binding portion thereof.
8. A method of activating Tumor Necrosis Factor Receptor 25 (TNFR25) receptor expressed on a cell comprising:
 - a. contacting the cell with an TNFR25 agonist, wherein said agonist is selected from the group consisting of:
 - i. a monoclonal antibody 4C12;
 - ii. an antibody that binds to a Tumor Necrosis Factor Receptor 25 (TNFR25), wherein said antibody is capable of increasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody;
 - iii. a soluble TL1A;
 - iv. an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25 agonist antibody;
 - v. an expression vector with an expression cassette capable of driving the transgenic expression of a soluble TL1A; and
 - vi. an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25;
 - b. observing an increase in TNFR25 receptor signaling.
9. An isolated antibody that binds a Tumor Necrosis Factor Receptor 25 (TNFR25) antigen wherein said antibody is capable of acting as a TNFR25 antagonist.
10. The antibody of claim 9, wherein said antibody binds a TL1A antigen, wherein said antibody is capable of decreasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody.

11. The antibody of claim 9, wherein said Tumor Necrosis Factor Receptor 25 (TNFR25) antigen is human or mouse.

12. A method of inhibiting Tumor Necrosis Factor Receptor 25 (TNFR25) receptor signaling in a cell comprising:

- a. contacting the cell with a TNFR25 antagonist, wherein said antagonist is selected from the group consisting of:
 - i. a monoclonal antibody L4G6
 - ii. an antibody that binds to a Tumor Necrosis Factor Receptor 25 (TNFR25) wherein said antibody is capable of decreasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody;
 - iii. an antibody specific for soluble TL1A;
 - iv. a fusion protein comprising an extracellular portion of TNFR25;
 - v. an expression vector with an expression cassette capable of driving the transgenic expression of the extracellular portion of TNFR25;
 - vi. an expression vector with an expression cassette capable of driving the transgenic expression of an antibody specific for soluble TL1A; and
 - vi. an expression vector with an expression cassette capable of driving the transgenic expression of L4G6;
- b. observing an decrease in TNFR25 receptor signaling.
- 13.** A tumor vaccine comprising a tumor antigen and a TNFR25 agonist as a biological response modifier.
- 14.** The tumor vaccine of claim 12 wherein said TNFR25 agonist is selected from the group consisting of consisting of:
 - a. a monoclonal antibody 4C12;
 - b. an antibody that binds to a Tumor Necrosis Factor Receptor 25 (TNFR25), wherein said antibody is capable of increasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody;
 - c. a soluble TL1A;
 - d. an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25 agonist antibody;
 - e. an expression vector with an expression cassette capable of driving the transgenic expression of a soluble TL1A; and

f. an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25.

15. The tumor vaccine of claim 14 further comprising an adjuvant.

16. A method of immunizing a patient against a tumor comprising isolating a tumor specific antigen from said tumor; and administering to the patient a tumor vaccine of claim 14.

17. A method of treating cancer in a patient comprising depleting a patient of CD4+/CD25+T regulatory cells (Tregs) by providing the patient with a composition comprising the toxin of claim 4; and providing a patient with a chemotherapeutic agent.

18. A method of treating and/or preventing gut inflammation comprising providing a patient in need thereof with an effective amount of a therapeutic composition comprising a TNFR25 agonist.

19. The method of claim 18, wherein said gut inflammation condition is selected from the group consisting of inflammatory bowel disease, Crohn's disease, and ulcerative colitis.

20. The method of claim 18, wherein said TNFR25 agonist is selected from the group consisting of

a. soluble TL1A

b. a monoclonal antibody 4C12

c. an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25 agonist antibody;

d. an expression vector with an expression cassette capable of driving the transgenic expression of a soluble TL1A; and

e. an expression vector with an expression cassette capable of driving the transgenic expression of a TNFR25.

21. A therapeutic composition for the facilitation of an organ transplant comprising a TNFR25 antagonist and an immunosuppressant.

22. The composition of claim 21 wherein the immunosuppressant is selected from the group consisting of glucocorticoid, azathioprine, methotrexate, cyclosporin, cyclophosphamide, mercaptopurine, tacrolimus and mycophenolate mofetil.

23. The composition of claim 21 wherein said TNFR25 antagonist is selected from the group consisting of:

a. a monoclonal antibody L4G6

b. an antibody that binds to a Tumor Necrosis Factor Receptor 25 (TNFR25) is capable of decreasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody;

c. an antibody specific for soluble TL1A;

d. a fusion protein comprising an extracellular portion of TNFR25;

e. an expression vector with an expression cassette capable of driving the transgenic expression of the extracellular portion of TNFR25;

f. an expression vector with an expression cassette capable of driving the transgenic expression of an antibody specific for soluble TL1A; and

g. an expression vector with an expression cassette capable of driving the transgenic expression of L4G6.

24. A method of transplanting a tissue from a donor into a host comprising:

a. obtaining the tissue from the donor;

b. providing the host with the composition of claim 21;

c. transplanting the tissue into the host.

25. A method of inhibiting the clonal expansion of a population of cognate CD8 T cells comprising,

a. exposing the CD8 T cells to their cognate antigen; and

b. exposing the CD8 T cells to a TNFR25 antagonist.

26. The method of claim 25 wherein said TNFR25 antagonist is selected from the group consisting of:

a. a monoclonal antibody L4G6

b. an antibody that binds to a Tumor Necrosis Factor Receptor 25 (TNFR25) wherein said antibody is capable of decreasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody;

c. an antibody specific for soluble TL1A;

d. a fusion protein comprising an extracellular portion of TNFR25;

e. an expression vector with an expression cassette capable of driving the transgenic expression of the extracellular portion of TNFR25;

f. an expression vector with an expression cassette capable of driving the transgenic expression of an antibody specific for soluble TL1A; and

g. an expression vector with an expression cassette capable of driving the transgenic expression of L4G6.

27. The method of claim 26 wherein cognate antigen is associated with tissue to be transplanted from a donor into a host.

28. A method of treating and/or preventing lung inflammation comprising providing a patient in need thereof with an effective amount of a therapeutic composition comprising a TNFR25 antagonist.

29. The method of claim 28, wherein said inflammation condition is a result of asthma.

30. The method of claim 28 wherein said TNFR25 antagonist is selected from the group consisting of:

a. a monoclonal antibody L4G6

b. an antibody that binds to a Tumor Necrosis Factor Receptor 25 (TNFR25) wherein said antibody is capable of decreasing OT-I CD8 cell expansion when cross-primed by gp96-Ig-ovalbumin relative to a control antibody;

c. an antibody specific for soluble TL1A;

d. a fusion protein comprising an extracellular portion of TNFR25;

e. an expression vector with an expression cassette capable of driving the transgenic expression of the extracellular portion of TNFR25;

- f. an expression vector with an expression cassette capable of driving the transgenic expression of an antibody specific for soluble TL1A; and
- g. an expression vector with an expression cassette capable of driving the transgenic expression of L4G6.

31. An isolated TNFR25 antagonist comprising a polypeptide encoded by a nucleic acid comprising a sequence that hybridizes under stringent conditions to SEQ ID NOs: 4, 5, 6 and/or 16, and wherein said sequence encodes an amino acid sequence capable of binding a TL1A protein.

32. The isolated TNFR25 antagonist of claim 31, wherein said sequence hybridizes under stringent conditions to SEQ ID NO: 4.

33. The isolated TNFR25 antagonist of claim 31, wherein said sequence hybridizes under stringent conditions to SEQ ID NO: 5.

34. The isolated TNFR25 antagonist of claim 31, wherein said sequence hybridizes under stringent conditions to SEQ ID NO: 6.

35. The isolated TNFR25 antagonist of claim 31, wherein said TL1A is human TL1A.

36. The isolated TNFR25 antagonist of claim 31, wherein said TL1A is mouse TL1A.

37. A method of treating and/or preventing lung inflammation comprising providing a patient in need thereof with an effective amount of a therapeutic composition comprising the TNFR25 antagonist of claim 31.

38. A method of transplanting a tissue from a donor into a host comprising:

- a. obtaining the tissue from the donor;
- b. providing the host with the TNFR25 antagonist of claim 31;
- c. transplanting the tissue into the host.

39. A composition comprising a polypeptide encoded by a sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein said sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein; and a toxic agent.

40. The composition of claim 39 wherein said toxic agent is selected from the group consisting of a radioactive isotope, ricin, abrin, diphtheria toxin, *Pseudomonas* exotoxin, and metal ion.

41. A method of treating cancer in a patient comprising depleting a patient of CD4+/CD25+T regulatory cells (Tregs) by providing the patient with a composition comprising the toxin of claim 39; and providing a patient with a chemotherapeutic agent.

42. A method of treating and/or preventing gut inflammation comprising providing a patient in need thereof with an effective amount of a composition comprising a polypeptide encoded by a sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein said sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein.

43. The method of claim 42 wherein said gut inflammation associated with irritable bowel syndrome.

44. The method of claim 42 wherein said gut inflammation associated with Crohn's disease.

45. A tumor vaccine comprising a tumor antigen and a polypeptide encoded by a sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein said sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein, as a biological response modifier.

46. An expression vector comprising a nucleic acid sequence that hybridizes under stringent conditions to SEQ ID NOs: 3 and/or 7, and wherein said sequence encodes an amino acid sequence capable of binding a TNFR25 receptor protein.

47. An expression vector comprising a nucleic acid sequence that hybridizes under stringent conditions to SEQ ID NOs: 4, 5, 6 and/or 16, and wherein said sequence encodes an amino acid sequence capable of binding a TL1A protein

* * * * *