

(43) International Publication Date  
25 November 2010 (25.11.2010)(10) International Publication Number  
**WO 2010/135521 A2**(51) International Patent Classification:  
*C07K 16/10* (2006.01)(21) International Application Number:  
PCT/US2010/035559(22) International Filing Date:  
20 May 2010 (20.05.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/180,027 20 May 2009 (20.05.2009) US  
61/234,145 14 August 2009 (14.08.2009) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 61/180,027 (CIP)  
Filed on 20 May 2009 (20.05.2009)  
US 61/234,145 (CIP)  
Filed on 14 August 2009 (14.08.2009)(71) Applicant (for all designated States except US): **THER-ACLONE SCIENCES, INC.** [US/US]; Seattle Life Sciences Building, 1124 Columbia Street, Suite 300, Seattle, WA 98104 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GRANDEA, Andres, G.** [US/US]; 20024 Palatine Avenue N., Shoreline, WA 98133 (US). **KING, Gordon** [US/US]; 15716 1st Avenue N.W., Shoreline, WA 98177 (US). **COX,****Thomas, C.** [US/US]; 7504 230th Avenue NE, Redmond, WA 98053 (US). **OLSEN, Ole** [CA/US]; 5309 117th Street SE, Everett WA 98208 (GB). **MITCHAM, Jennifer** [US/US]; 20330 NE 85th Street, Redmond, WA 98053 (US). **MOYLE, Matthew** [US/US]; 15737 NE 134th Street, Redmond, WA 98052 (US). **HAMMOND, Phil** [US/US]; 7733 30th Avenue N.W., Seattle, WA 98117 (US).(74) Agents: **ELRIFI, Ivor, R.** et al.; Nintz Levin Cohn Ferris Glovsky And Popeo, P.C., One Financial Center, Boston, MA 02111 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK,

[Continued on next page]

(54) Title: COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF INFLUENZA




Fig. 10

(57) Abstract: The present invention provides novel human anti- influenza antibodies and related compositions and methods. These antibodies are used in the diagnosis and treatment of influenza infection.



---

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG). **Published:**

— *without international search report and to be republished  
upon receipt of that report (Rule 48.2(g))*

## **COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF INFLUENZA**

### **RELATED APPLICATIONS**

[01] This application claims the benefit of provisional applications USSN 61/180,027, filed May 20, 2009 and USSN 61/234,145 filed August 14, 2009, the contents of which are each herein incorporated by reference in their entirety.

### **FIELD OF THE INVENTION**

[02] The present invention relates generally to therapy, diagnosis and monitoring of influenza infection. The invention is more specifically related to methods of identifying influenza matrix 2 protein-specific antibodies and their manufacture and use. Such antibodies are useful in pharmaceutical compositions for the prevention and treatment of influenza, and for the diagnosis and monitoring of influenza infection.

### **BACKGROUND OF THE INVENTION**

[03] Influenza virus infects 5-20% of the population and results in 30,000-50,000 deaths each year in the U.S. Although the influenza vaccine is the primary method of infection prevention, four antiviral drugs are also available in the U.S.: amantadine, rimantadine, oseltamivir and zanamivir. As of December 2005, only oseltamivir (TAMIFLU™) is recommended for treatment of influenza A due to the increasing resistance of the virus to amantadine and rimantidine resulting from an amino acid substitution in the M2 protein of the virus.

[04] Disease caused by influenza A viral infections is typified by its cyclical nature. Antigenic drift and shift allow for different A strains to emerge every year. Added to that, the threat of highly pathogenic strains entering into the general population has stressed the need for novel therapies for flu infections. The predominant fraction of neutralizing antibodies is directed to the polymorphic regions of the hemagglutinin and neuraminidase proteins. Thus, such a neutralizing MAb would presumably target only one or a few strains. A recent focus has been on the relatively invariant matrix 2 (M2) protein. Potentially, a neutralizing MAb to M2 would be an adequate therapy for all influenza A strains.

[05] The M2 protein is found in a homotetramer that forms an ion channel and is thought to aid in the uncoating of the virus upon entering the cell. After infection, M2 can be found in

abundance at the cell surface. It is subsequently incorporated into the virion coat, where it only comprises about 2% of total coat protein. The M2 extracellular domain (M2e) is short, with the aminoterminal 2-24 amino acids displayed outside of the cell. Anti-M2 MAbs to date have been directed towards this linear sequence. Thus, they may not exhibit desired binding properties to cellularly expressed M2, including conformational determinants on native M2.

[06] Therefore, a long-felt need exists in the art for new antibodies that bind to the cell-expressed M2 and conformational determinants on the native M2.

### SUMMARY OF THE INVENTION

[07] The present invention provides fully human monoclonal antibodies specifically directed against M2e. Optionally, the antibody is isolated from a B-cell from a human donor. Exemplary monoclonal antibodies include 8i10, 21B15 23K12, 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, and 3242\_P05 described herein. Alternatively, the monoclonal antibody is an antibody that binds to the same epitope as 8i10, 21B15 23K12, 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, or 3242\_P05. The antibodies respectively referred to herein are huM2e antibodies. The huM2e antibody has one or more of the following characteristics: a) binds to an epitope in the extracellular domain of the matrix 2 ectodomain (M2e) polypeptide of an influenza virus; b) binds to influenza A infected cells; or c) binds to influenza A virus.

[08] The epitope that huM2e antibody binds to is a non-linear epitope of a M2 polypeptide. Preferably, the epitope includes the amino terminal region of the M2e polypeptide. More preferably the epitope wholly or partially includes the amino acid sequence SLLTEV (SEQ ID NO: 42). Most preferably, the epitope includes the amino acid at position 2, 5 and 6 of the M2e polypeptide when numbered in accordance with SEQ ID NO: 1. The amino acid at position 2 is a serine; at position 5 is a threonine; and at position 6 is a glutamic acid.

[09] A huM2e antibody contains a heavy chain variable having the amino acid sequence of SEQ ID NOS: 44 or 50 and a light chain variable having the amino acid sequence of SEQ ID NOS: 46 or 52. Preferably, the three heavy chain CDRs include an amino acid sequence at least 90%, 92%, 95%, 97% 98%, 99% or more identical to the amino acid sequence of NYYWS (SEQ ID NO: 72), FIYYGGNTKYNPSLKS (SEQ ID NO: 74), ASCSGGYCILD (SEQ ID NO: 76), SNYMS (SEQ ID NO: 103), VIYSGGSTYYADSVK (SEQ ID NO: 105),

CLSRMRGYGLDV (SEQ ID NO: 107) (as determined by the Kabat method) or ASCSGGYCILD (SEQ ID NO: 76), CLSRMRGYGLDV (SEQ ID NO: 107), GSSISN (SEQ ID NO: 109), FIYYGGNTK (SEQ ID NO: 110), GSSISN (SEQ ID NO: 111), GFTVSSN (SEQ ID NO: 112), VIYSGGSTY (SEQ ID NO: 113) (as determined by the Chothia method) and a light chain with three CDRs that include an amino acid sequence at least 90%, 92%, 95%, 97% 98%, 99% or more identical to the amino acid sequence of RASQNIYKYLN (SEQ ID NO: 59), AASGLQS (SEQ ID NO: 61), QQSYSPPLT (SEQ ID NO: 63), RTSQSISSYLN (SEQ ID NO: 92), AASSLQSGVPSRF (SEQ ID NO: 94), QQSYSMMPA (SEQ ID NO: 96) (as determined by the Kabat method) or RASQNIYKYLN (SEQ ID NO: 59), AASGLQS (SEQ ID NO: 61), QQSYSPPLT (SEQ ID NO: 63), RTSQSISSYLN (SEQ ID NO: 92), AASSLQSGVPSRF (SEQ ID NO: 94), QQSYSMMPA (SEQ ID NO: 96) (as determined by the Chothia method). The antibody binds M2e.

**[10]** An isolated anti-matrix 2 ectodomain (M2e) antibody, or antigen-binding fragment thereof, comprising a heavy chain variable (VH) domain and a light chain variable (VL) domain, wherein the VH domain and the VL domain each comprise three complementarity determining regions 1 to 3 (CDR1-3), and wherein each CDR includes the following amino acid sequences: VH CDR1: SEQ ID NOs: 179, 187, 196, 204, 212, 224, 230, 235, 242, 248, or 254; VH CDR2: SEQ ID NOs: 180, 188, 195, 197, 205, 213, 218, 225, 231, 236, 243, 249, 246, or 256; VH CDR3 SEQ ID NOs: 181, 189, 198, 206, 214, 219, 226, 232, 237, 244, or 250; VL CDR1: SEQ ID NOs: 184, 192, 199, 215, 220, 233, or 238; VL CDR2: SEQ ID NOs: 61, 185, 193, 200, 207, 211, 216, 227, 239, or 241; and VL CDR3: SEQ ID NOs: 63, 186, 194, 201, 208, 221, 228, 234, 240, 245, or 251.

**[11]** Alternatively, or in addition, an isolated anti-matrix 2 ectodomain (M2e) antibody, or antigen-binding fragment thereof, comprising a heavy chain variable (VH) domain and a light chain variable (VL) domain, wherein the VH domain and the VL domain each comprise three complementarity determining regions 1 to 3 (CDR1-3), and wherein each CDR includes the following amino acid sequences: VH CDR1: SEQ ID NOs: 182, 190, 202, 209, 222, 229, 247, 252, 257, 258, or 260; VH CDR2: SEQ ID NOs: 183, 191, 203, 210, 217, 223, 230, 246, 253, 259, or 261; VH CDR3 SEQ ID NOs: 181, 189, 195, 198, 206, 214, 219, 226, 232, 237, 244, or 250; VL CDR1: SEQ ID NOs: 184, 192, 199, 215, 220, 233, or 238; VL CDR2: SEQ ID NOs: 61, 185, 193, 200, 207, 211, 216, 227, 239, or 241; and VL CDR3: SEQ ID NOs: 63, 186, 194, 201, 208, 221, 228, 234, 240, 245, or 251.

**[12]** The heavy chain of an M2e antibody is derived from a germ line V (variable) gene such as, for example, the IgHV4 or the IgHV3 germline gene.

[13] The M2e antibodies of the invention include a variable heavy chain ( $V_H$ ) region encoded by a human IgHV4 or the IgHV3 germline gene sequence. An IgHV4 germline gene sequence is shown, *e.g.*, in Accession numbers L10088, M29812, M95114, X56360 and M95117. An IgHV3 germline gene sequence is shown, *e.g.*, in Accession numbers X92218, X70208, Z27504, M99679 and AB019437. The M2e antibodies of the invention include a  $V_H$  region that is encoded by a nucleic acid sequence that is at least 80% homologous to the IgHV4 or the IgHV3 germline gene sequence. Preferably, the nucleic acid sequence is at least 90%, 95%, 96%, 97% homologous to the IgHV4 or the IgHV3 germline gene sequence, and more preferably, at least 98%, 99% homologous to the IgHV4 or the IgHV3 germline gene sequence. The  $V_H$  region of the M2e antibody is at least 80% homologous to the amino acid sequence of the  $V_H$  region encoded by the IgHV4 or the IgHV3  $V_H$  germline gene sequence. Preferably, the amino acid sequence of  $V_H$  region of the M2e antibody is at least 90%, 95%, 96%, 97% homologous to the amino acid sequence encoded by the IgHV4 or the IgHV3 germline gene sequence, and more preferably, at least 98%, 99% homologous to the sequence encoded by the IgHV4 or the IgHV3 germline gene sequence.

[14] The M2e antibodies of the invention also include a variable light chain ( $V_L$ ) region encoded by a human IgKV1 germline gene sequence. A human IgKV1  $V_L$  germline gene sequence is shown, *e.g.*, Accession numbers X59315, X59312, X59318, J00248, and Y14865. Alternatively, the M2e antibodies include a  $V_L$  region that is encoded by a nucleic acid sequence that is at least 80% homologous to the IgKV1 germline gene sequence. Preferably, the nucleic acid sequence is at least 90%, 95%, 96%, 97% homologous to the IgKV1 germline gene sequence, and more preferably, at least 98%, 99% homologous to the IgKV1 germline gene sequence. The  $V_L$  region of the M2e antibody is at least 80% homologous to the amino acid sequence of the  $V_L$  region encoded the IgKV1 germline gene sequence. Preferably, the amino acid sequence of  $V_L$  region of the M2e antibody is at least 90%, 95%, 96%, 97% homologous to the amino acid sequence encoded by the IgKV1 germline gene sequence, and more preferably, at least 98%, 99% homologous to the sequence encoded by the IgKV1 germline gene sequence.

[15] In another aspect the invention provides a composition including an huM2e antibody according to the invention. The composition is optionally a pharmaceutical composition including any one of the M2e antibodies described herein and a pharmaceutical carrier. In various aspects the composition further includes an anti-viral drug, a viral entry inhibitor or a viral attachment inhibitor. The anti-viral drug is for example a neuraminidase inhibitor, a HA inhibitor, a sialic acid inhibitor or an M2 ion channel inhibitor. The M2 ion channel inhibitor

is for example amantadine or rimantadine. The neuraminidase inhibitor is for example zanamivir, or oseltamivir phosphate. In a further aspect the composition further includes a second anti-influenza A antibody.

[16] In a further aspect the huM2e antibodies according to the invention are operably-linked to a therapeutic agent or a detectable label.

[17] Additionally, the invention provides methods for stimulating an immune response, treating, preventing or alleviating a symptom of an influenza viral infection by administering an huM2e antibody to a subject

[18] Optionally, the subject is further administered with a second agent such as, but not limited to, an influenza virus antibody, an anti-viral drug such as a neuraminidase inhibitor, a HA inhibitor, a sialic acid inhibitor or an M2 ion channel inhibitor, a viral entry inhibitor or a viral attachment inhibitor. The M2 ion channel inhibitor is for example amantadine or rimantadine. The neuraminidase inhibitor for example zanamivir, or oseltamivir phosphate. The subject is suffering from or is predisposed to developing an influenza virus infection, such as, for example, an autoimmune disease or an inflammatory disorder.

[19] In another aspect, the invention provides methods of administering the huM2e antibody of the invention to a subject prior to, and/or after exposure to an influenza virus. For example, the huM2e antibody of the invention is used to treat or prevent rejection influenza infection. The huM2e antibody is administered at a dose sufficient to promote viral clearance or eliminate influenza A infected cells.

[20] Also included in the invention is a method for determining the presence of an influenza virus infection in a patient, by contacting a biological sample obtained from the patient with a huM2e antibody; detecting an amount of the antibody that binds to the biological sample; and comparing the amount of antibody that binds to the biological sample to a control value.

[21] The invention further provides a diagnostic kit comprising a huM2e antibody.

[22] Other features and advantages of the invention will be apparent from and are encompassed by the following detailed description and claims.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

[23] Figure 1 shows the binding of three antibodies of the present invention and control hu14C2 antibody to 293-HEK cells transfected with an M2 expression construct or control vector, in the presence or absence of free M2 peptide.

- [24] Figures 2A and B are graphs showing human monoclonal antibody binding to influenza A/Puerto Rico/8/32.
- [25] Figure 3A is a chart showing amino acid sequences of extracellular domains of M2 variants.
- [26] Figures 3B and C are bar charts showing binding of human monoclonal anti-influenza antibody binding to M2 variants shown in Figure 3A.
- [27] Figures 4A and B are bar charts showing binding of human monoclonal anti-influenza antibody binding to M2 peptides subjected to alanine scanning mutagenesis.
- [28] Figure 5 is a series of bar charts showing binding of MAbs 8i10 and 23K12 to M2 protein representing influenza strain A/HK/483/1997 sequence that was stably expressed in the CHO cell line DG44.
- [29] Figure 6A is a chart showing cross reactivity binding of anti-M2 antibodies to variant M2 peptides.
- [30] Figure 6B is a chart showing binding activity of M2 antibodies to truncated M2 peptides.
- [31] Figure 7 is a graph showing survival of influenza infected mice treated with human anti-influenza monoclonal antibodies.
- [32] Figure 8 is an illustration showing the anti-M2 antibodies bind a highly conserved region in the N-Terminus of M2e.
- [33] Figure 9 is a graph showing anti-M2 rHMAb clones from crude supernatant bound to influenza on ELISA, whereas the control anti-M2e mAb 14C2 did not readily bind virus.
- [34] Figure 10 is a series of photographs showing anti-M2 rHMAbs bound to cells infected with influenza. MDCK cells were or were not infected with influenza A/PR/8/32 and Ab binding from crude supernatant was tested 24 hours later. Data were gathered from the FMAT plate scanner.
- [35] Figure 11 is a graph showing anti-M2 rHMAb clones from crude supernatant bound to cells transfected with the influenza subtypes H3N2, HK483, and VN1203 M2 proteins. Plasmids encoding full length M2 cDNAs corresponding to influenza strains H3N2, HK483, and VN1203, as well as a mock plasmid control, were transiently transfected into 293 cells. The 14C2, 8i10, 23K12, and 21B15 mABs were tested for binding to the transfectants, and were detected with an AF647-conjugated anti-human IgG secondary antibody. Shown are the mean fluorescence intensities of the specific mAB bound after FACS analysis.

**DETAILED DESCRIPTION**

[36] The present invention provides fully human monoclonal antibodies specific against the extracellular domain of the matrix 2 (M2) polypeptide. The antibodies are respectively referred to herein as huM2e antibodies.

[37] M2 is a 96 amino acid transmembrane protein present as a homotetramer on the surface of influenza virus and virally infected cells. M2 contains a 23 amino acid ectodomain (M2e) that is highly conserved across influenza A strains. Few amino acid changes have occurred since the 1918 pandemic strain thus M2e is an attractive target for influenza therapies. In prior studies, monoclonal antibodies specific to the M2 ectodomain (M2e) were derived upon immunizations with a peptide corresponding to the linear sequence of M2e. In contrast, the present invention provides a novel process whereby full-length M2 is expressed in cell lines, which allows for the identification of human antibodies that bound this cell-expressed M2e. The huM2e antibodies have been shown to bind conformational determinants on the M2-transfected cells, as well as native M2, either on influenza infected cells, or on the virus itself. The huM2e antibodies did not bind the linear M2e peptide, but they do bind several natural M2 variants, also expressed upon cDNA transfection into cell lines. Thus, this invention has allowed for the identification and production of human monoclonal antibodies that exhibit novel specificity for a very broad range of influenza A virus strains. These antibodies may be used diagnostically to identify influenza A infection and therapeutically to treat influenza A infection.

[38] The huM2e antibodies of the invention have one or more of the following characteristics: the huM2e antibody binds a) to an epitope in the extracellular domain of the matrix 2 (M2) polypeptide of an influenza virus; b) binds to influenza A infected cells; and/or c) binds to influenza A virus (i.e., virions). The huM2e antibodies of the invention eliminate influenza infected cells through immune effector mechanisms, such as ADCC, and promote direct viral clearance by binding to influenza virions. The huM2e antibodies of the invention bind to the amino-terminal region of the M2e polypeptide. Preferably, the huM2e antibodies of the invention bind to the amino-terminal region of the M2e polypeptide wherein the N-terminal methionine residue is absent. Exemplary M2e sequences include those sequences listed on Table I below

[39] Table I

| Type | Name                     | Subtype | M2E Sequence             | SEQ ID NO    |
|------|--------------------------|---------|--------------------------|--------------|
| A    | BREVIG<br>MISSION.1.1918 | H1N1    | MSLLTEVETPTRNEWGCRCNDSSD | SEQ ID NO: 1 |
| A    | FORT MONMOUTH.1.1947     | H1N1    | MSLLTEVETPTKNEWECRCNDSSD | SEQ ID NO: 2 |
| A    | .SINGAPORE.02.2005       | H3N2    | MSLLTEVETPIRNEWECRCNDSSD | SEQ ID NO: 3 |

|   |                                   |         |                           |               |
|---|-----------------------------------|---------|---------------------------|---------------|
| A | WISCONSIN.10.98                   | H1N1    | MSLLTEVETPIRNGWECKCNDSSD  | SEQ ID NO: 4  |
| A | WISCONSIN.301.1976                | H1N1    | MSLLTEVETPIRSEWGCRCNDSSD  | SEQ ID NO: 5  |
| A | PANAMA.1.66                       | H2N2    | MSFLPEVETPIRNEWGCRCNDSSD  | SEQ ID NO: 6  |
| A | NEW YORK.321.1999                 | H3N2    | MSLLTEVETPIRNEWGCRCNDSSN  | SEQ ID NO: 7  |
| A | CARACAS.1.71                      | H3N2    | MSLLTEVETPIRKEWGCRCNDSSD  | SEQ ID NO: 8  |
| A | TAIWAN.3.71                       | H3N2    | MSFLTEVETPIRNEWGCRCNDSSD  | SEQ ID NO: 9  |
| A | WUHAN.359.95                      | H3N2    | MSLPTEVETPIRSEWGCRCNDSSD  | SEQ ID NO: 10 |
| A | HONG KONG.1144.99                 | H3N2    | MSLLPEVETPIRNEWGCRCNDSSD  | SEQ ID NO: 11 |
| A | HONG KONG.1180.99                 | H3N2    | MSLLPEVETPIRNGWGCRCNDSSD  | SEQ ID NO: 12 |
| A | HONG KONG.1774.99                 | H3N2    | MSLLTEVETPTRNGWECRCGSSD   | SEQ ID NO: 13 |
| A | NEW YORK.217.02                   | H1N2    | MSLLTEVETPIRNEWYRCNDSSD   | SEQ ID NO: 14 |
| A | NEW YORK.300.2003                 | H1N2    | MSLLTEVETPIRNEWYRCSDSSD   | SEQ ID NO: 15 |
| A | SWINE.SPAIN.54008.2004            | H3N2    | MSLLTEVETPTRNGWECRYSDSSD  | SEQ ID NO: 16 |
| A | GUANGZHOU.333.99                  | H9N2    | MSFLTEVETLTRNGWECRCSDSSD  | SEQ ID NO: 17 |
| A | HONG KONG.1073.99                 | H9N2    | MSLLTEVETLTRNGWECKCRDSSD  | SEQ ID NO: 18 |
| A | HONG KONG.1.68                    | H3N2    | MSLLTEVETPIRNEWGCRCNDSSD  | SEQ ID NO: 19 |
| A | SWINE.HONG KONG.126.1982          | H3N2    | MSLLTEVETPIRSEWGCRCNDSGD  | SEQ ID NO: 20 |
| A | NEW YORK.703.1995                 | H3N2    | MSLLTEVETPIRNEWECRCNGSSD  | SEQ ID NO: 21 |
| A | SWINE.QUEBEC.192.81               | H1N1    | MSLPTEVETPIRNEWGCRCNDSSD  | SEQ ID NO: 22 |
| A | PUERTO RICO.8.34                  | H1N1    | MSLLTEVETPIRNEWGCRCNGSSD  | SEQ ID NO: 23 |
| A | HONG KONG.485.97                  | H5N1    | MSLLTEVDTLTRNGWGCRCSDSSD  | SEQ ID NO: 24 |
| A | HONG KONG.542.97                  | H5N1    | MSLLTEVETLTKNWGCRCSDSSD   | SEQ ID NO: 25 |
| A | SILKY CHICKEN.SHANTOU.1826.2004   | H9N2    | MSLLTEVETPTRNGWECKCSDSSD  | SEQ ID NO: 26 |
| A | CHICKEN.TAIWAN.0305.04            | H6N1    | MSLLTEVETHTRNGWECKCSDSSD  | SEQ ID NO: 27 |
| A | QUAIL.ARKANSAS.16309-7.94         | H7N3NSA | MSLLTEVKTPTRNGWECKCSDSSD  | SEQ ID NO: 28 |
| A | HONG KONG.486.97                  | H5N1    | MSLLTEVETLTRNGWGCRCSDSSD  | SEQ ID NO: 29 |
| A | CHICKEN.PENNSYLVANIA.13552-1.98   | H7N2NSB | MSLLTEVETPTRDGWECKCSDSSD  | SEQ ID NO: 30 |
| A | CHICKEN.HEILONGJIANG.48.01        | H9N2    | MSLLTEVETPTRNGWGCRCSDSSD  | SEQ ID NO: 31 |
| A | SWINE.KOREA.S5.2005               | H1N2    | MSLLTEVETPTRNGWECKCNDSSD  | SEQ ID NO: 32 |
| A | HONG KONG.1073.99                 | H9N2    | MSLLTEVETLTRNGWECKCSDSSD  | SEQ ID NO: 33 |
| A | WISCONSIN.3523.88                 | H1N1    | MSLLTEVETPIRNEWGCKCNDSSD  | SEQ ID NO: 34 |
| A | X-31 VACCINE STRAIN               | H3N2    | MSFLTEVETPIRNEWGCRCNGSSD  | SEQ ID NO: 35 |
| A | CHICKEN.ROSTOCK.8.1934            | H7N1    | MSLLTEVETPTRNGWECRCNDSSD  | SEQ ID NO: 36 |
| A | ENVIRONMENT.NEW YORK.16326-1.2005 | H7N2    | MSLLTEVETPIRKGWECNCSDSSD  | SEQ ID NO: 37 |
| A | INDONESIA.560H.2006               | H5N1    | MSLLTEVETPTRNEWECRCSDSSD  | SEQ ID NO: 38 |
| A | CHICKEN.HONG KONG.SF1.03          | H9N2    | MSLLTGVETHTRNGWGCCKCSDSSD | SEQ ID NO: 39 |
| A | CHICKEN.HONGKONG.YU427.03         | H9N2    | MSLLPEVETHTRNGWGCRCSDSSD  | SEQ ID NO: 40 |

[40] In one embodiment, the huM2e antibodies of the invention bind to a M2e that wholly or partially includes the amino acid residues from position 2 to position 7 of M2e when numbered in accordance with SEQ ID NO: 1. For example, the huM2e antibodies of the invention bind wholly or partially to the amino acid sequence SLLTEVET (SEQ ID NO: 41)

Most preferably, the huM2e antibodies of the invention bind wholly or partially to the amino acid sequence SLLTEV (SEQ ID NO: 42). Preferably, the huM2e antibodies of the invention bind to non-linear epitope of the M2e protein. For example, the huM2e antibodies bind to an epitope comprising position 2, 5, and 6 of the M2e polypeptide when numbered in accordance to SEQ ID NO: 1 where the amino acid at a) position 2 is a serine; b) position 5 is a threonine; and c) position 6 is a glutamic acid. Exemplary huM2e monoclonal antibodies that binds to this epitope are the 8I10, 21B15 or 23K12 antibodies described herein.

[41] The 8I10 antibody includes a heavy chain variable region (SEQ ID NO: 44) encoded by the nucleic acid sequence shown below in SEQ ID NO: 43, and a light chain variable region (SEQ ID NO: 46) encoded by the nucleic acid sequence shown in SEQ ID NO: 45.

[42] The amino acids encompassing the CDRs as defined by Chothia, C. et al. (1989, Nature, 342: 877-883) are underlined and those defined by Kabat E.A. et al. (1991, Sequences of Proteins of Immunological Interest, 5<sup>th</sup> edit., NIH Publication no. 91-3242 U.S.

Department of Health and Human Services.) are highlighted in bold in the sequences below.

[43] The heavy chain CDRs of the 8I10 antibody have the following sequences per Kabat definition: NYYWS (SEQ ID NO: 72), FIYYGGNTKYNPSLKS (SEQ ID NO: 74) and ASCSGGYCILD (SEQ ID NO: 76). The light chain CDRs of the 8I10 antibody have the following sequences per Kabat definition: RASQNIYKYLN (SEQ ID NO: 59), AASGLQS (SEQ ID NO: 61) and QQSYSPPLT (SEQ ID NO: 63).

[44] The heavy chain CDRs of the 8I10 antibody have the following sequences per Chothia definition: GSSISN (SEQ ID NO: 109), FIYYGGNTK (SEQ ID NO: 110) and ASCSGGYCILD (SEQ ID NO: 76). The light chain CDRs of the 8I10 antibody have the following sequences per Chothia definition: RASQNIYKYLN (SEQ ID NO: 59), AASGLQS (SEQ ID NO: 61) and QQSYSPPLT (SEQ ID NO: 63).

#### >8I10 VH nucleotide sequence: (SEQ ID NO: 43)

CAGGTGCAATTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCAC  
CTGCACTGTCTCTGGTTCGTCCATCAGTAATTACTACTGGAGCTGGATCCGGCAGTCCCCAG  
GGAAGGGACTGGAGTGGATTGGGTTTATCTATTACGGTGGAAACACCAAGTACAATCCCTCC  
CTCAAGAGCCGCGTCACCATATCACAAGACACTTCCAAGAGTCAGGTCTCCCTGACGATGAG  
CTCTGTGACCGCTGCGGAATCGGCCGTCTATTTCTGTGCGAGAGCGTCTTGTAGTGGTGGTT  
ACTGTATCCTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCG

#### >8I10 VH amino acid sequence: (SEQ ID NO: 44)

Kabat Bold, Chothia underlined

|   |   |   |   |          |          |          |          |          |          |          |          |          |          |   |   |   |   |   |   |   |
|---|---|---|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|---|---|---|---|---|---|---|
| Q | V | Q | L | Q        | E        | S        | G        | P        | G        | L        | V        | K        | P        | S | E | T | L | S | L | T |
| C | T | V | S | <u>G</u> | <u>S</u> | <u>S</u> | <u>I</u> | <u>S</u> | <b>N</b> | <b>Y</b> | <b>Y</b> | <b>W</b> | <b>S</b> | W | I | R | Q | S | P | G |

|          |          |          |   |   |   |   |          |          |          |          |          |          |          |          |          |          |          |          |          |          |
|----------|----------|----------|---|---|---|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| K        | G        | L        | E | W | I | G | <b>F</b> | <b>I</b> | <b>Y</b> | <b>Y</b> | <b>G</b> | <b>G</b> | <b>N</b> | <b>T</b> | <b>K</b> | <b>Y</b> | <b>N</b> | <b>P</b> | <b>S</b> | <b>L</b> |
| <b>K</b> | <b>S</b> | R        | V | T | I | S | <u>Q</u> | D        | T        | S        | K        | S        | <u>Q</u> | V        | S        | L        | T        | M        | S        | S        |
| V        | T        | A        | A | E | S | A | V        | Y        | F        | C        | A        | R        | <b>A</b> | <b>S</b> | <b>C</b> | <b>S</b> | <b>G</b> | <b>G</b> | <b>Y</b> | <b>C</b> |
| <b>I</b> | <b>L</b> | <b>D</b> | Y | W | G | Q | G        | T        | L        | V        | T        | V        | S        |          |          |          |          |          |          |          |

**>8I10 VL nucleotide sequence: (SEQ ID NO: 45)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
 CACTTGCCGGGCGAGTCAGAACATTTACAAGTATTTAAATTGGTATCAGCAGAGACCAGGGA  
 AAGCCCCTAAGGGCCTGATCTCTGCTGCATCCGGGTTGCAAAGTGGGGTCCCATCAAGGTTT  
 AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCACCAGTCTGCAACCTGAAGATTT  
 TGCAACTTACTACTGTCAACAGAGTTACAGTCCCCCTCTCACTTTCGGCGGAGGGACCAGGG  
 TGGAGATCAAAC

**>8I10 VL amino acid sequence: (SEQ ID NO: 46)**

Kabat Bold, Chothia underlined

|   |   |          |          |          |          |          |          |          |          |          |          |          |          |   |   |   |   |   |   |   |
|---|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|---|---|---|---|---|---|---|
| D | I | Q        | M        | T        | Q        | S        | P        | S        | S        | L        | S        | A        | S        | V | G | D | R | V | T | I |
| T | C | <b>R</b> | <b>A</b> | <b>S</b> | <u>Q</u> | <b>N</b> | <b>I</b> | <b>Y</b> | <b>K</b> | <b>Y</b> | <b>L</b> | <b>N</b> | W        | Y | Q | Q | R | P | G | K |
| A | P | K        | G        | L        | I        | S        | <b>A</b> | <b>A</b> | <b>S</b> | <b>G</b> | <b>L</b> | <u>Q</u> | <b>S</b> | G | V | P | S | R | F | S |
| G | S | G        | S        | G        | T        | D        | F        | T        | L        | T        | I        | T        | S        | L | Q | P | E | D | F | A |
| T | Y | Y        | C        | <u>Q</u> | <u>Q</u> | <b>S</b> | <b>Y</b> | <b>S</b> | <b>P</b> | <b>P</b> | <b>L</b> | <b>T</b> | F        | G | G | G | T | R | V | E |
| I | K |          |          |          |          |          |          |          |          |          |          |          |          |   |   |   |   |   |   |   |

[45] The 21B15 antibody includes antibody includes a heavy chain variable region (SEQ ID NO: 44) encoded by the nucleic acid sequence shown below in SEQ ID NO: 47, and a light chain variable region (SEQ ID NO: 46) encoded by the nucleic acid sequence shown in SEQ ID NO: 48.

[46] The amino acids encompassing the CDRs as defined by Chothia et al. 1989, are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[47] The heavy chain CDRs of the 21B15 antibody have the following sequences per Kabat definition: NYYWS (SEQ ID NO: 72), FIYYGGNTKYNPSLKS (SEQ ID NO: 74) and ASCSGGYCILD (SEQ ID NO: 76). The light chain CDRs of the 21B15 antibody have the following sequences per Kabat definition: RASQNIYKYLN (SEQ ID NO: 59), AASGLQS (SEQ ID NO: 61) and QQSYSPPLT (SEQ ID NO: 63).

[48] The heavy chain CDRs of the 21B15 antibody have the following sequences per Chothia definition: GSSISN (SEQ ID NO: 111), FIYYGGNTK (SEQ ID NO: 110) and ASCSGGYCILD (SEQ ID NO: 76). The light chain CDRs of the 21B15 antibody have the following sequences per Chothia definition: RASQNIYKYLN (SEQ ID NO: 59), AASGLQS (SEQ ID NO: 61) and QQSYSPPLT (SEQ ID NO: 63).

**>21B15 VH nucleotide sequence: (SEQ ID NO: 47)**

CAGGTGCAATTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCAC  
 CTGCACTGTCTCTGGTTCGTCCATCAGTAATTACTACTGGAGCTGGATCCGGCAGTCCCCAG  
 GGAAGGGACTGGAGTGGATTGGGTTTATCTATTACGGTGGAACACCAAGTACAATCCCTCC  
 CTCAAGAGCCGCGTCACCATATCACAAGACACTTCCAAGAGTCAGGTCTCCCTGACGATGAG  
 CTCTGTGACCGCTGCGGAATCGGCCGTCTATTTCTGTGCGAGAGCGTCTTGTAAGTGGTGGTT  
 ACTGTATCCTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCG

**>21B15 VH amino acid sequence: (SEQ ID NO: 44)**

Kabat Bold, Chothia underlined

|          |          |          |   |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |
|----------|----------|----------|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| Q        | V        | Q        | L | Q        | E        | S        | G        | P        | G        | L        | V        | K        | P        | S        | E        | T        | L        | S        | L        | T        |
| C        | T        | V        | S | <u>G</u> | S        | S        | I        | S        | <b>N</b> | <b>Y</b> | <b>Y</b> | <b>W</b> | <b>S</b> | W        | I        | R        | Q        | S        | P        | G        |
| K        | G        | L        | E | <u>W</u> | <u>I</u> | <u>G</u> | <b>F</b> | <b>I</b> | <b>Y</b> | <b>Y</b> | <b>G</b> | <b>G</b> | <b>N</b> | <b>T</b> | <b>K</b> | <b>Y</b> | <b>N</b> | <b>P</b> | <b>S</b> | <b>L</b> |
| <b>K</b> | <b>S</b> | R        | V | T        | I        | S        | Q        | D        | T        | S        | K        | S        | Q        | V        | S        | L        | T        | M        | S        | S        |
| V        | T        | A        | A | E        | S        | A        | V        | Y        | F        | C        | A        | R        | <b>A</b> | <b>S</b> | <b>C</b> | <b>S</b> | <b>G</b> | <b>G</b> | <b>Y</b> | <b>C</b> |
| <b>I</b> | <b>L</b> | <b>D</b> | Y | W        | G        | Q        | G        | T        | L        | V        | T        | V        | S        |          |          |          |          |          |          |          |

**>21B15 VL nucleotide sequence: (SEQ ID NO: 48)**

GACATCCAGGTGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
 CACTTGCCGCGCGAGTCAGAACATTTACAAGTATTTAAATTGGTATCAGCAGAGACCAGGGA  
 AAGCCCCTAAGGGCCTGATCTCTGCTGCATCCGGGTTGCAAAGTGGGGTCCCATCAAGGTTTC  
 AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCACCAGTCTGCAACCTGAAGATTT  
 TGCAACTTACTACTGTCAACAGAGTTACAGTCCCCCTCTCACTTTCGGCGGAGGGACCAGGG  
 TGGATATCAAAC

**>21B15 VL amino acid sequence: (SEQ ID NO: 46)**

Kabat Bold, Chothia underlined

|   |   |          |          |          |          |          |          |          |          |          |          |          |          |   |   |   |   |   |   |   |
|---|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|---|---|---|---|---|---|---|
| D | I | Q        | V        | T        | Q        | S        | P        | S        | S        | L        | S        | A        | S        | V | G | D | R | V | T | I |
| T | C | <b>R</b> | <b>A</b> | <b>S</b> | <b>Q</b> | <b>N</b> | <b>I</b> | <b>Y</b> | <b>K</b> | <b>Y</b> | <b>L</b> | <b>N</b> | W        | Y | Q | Q | R | P | G | K |
| A | P | K        | G        | L        | I        | S        | <b>A</b> | <b>A</b> | <b>S</b> | <b>G</b> | <b>L</b> | <b>Q</b> | <b>S</b> | G | V | P | S | R | F | S |
| G | S | G        | S        | G        | T        | D        | F        | T        | L        | T        | I        | T        | S        | L | Q | P | E | D | F | A |
| T | Y | Y        | C        | <b>Q</b> | <b>Q</b> | <b>S</b> | <b>Y</b> | <b>S</b> | <b>P</b> | <b>P</b> | <b>L</b> | <b>T</b> | F        | G | G | G | T | R | V | D |
| I | K |          |          |          |          |          |          |          |          |          |          |          |          |   |   |   |   |   |   |   |

[49] The 23K12 antibody includes antibody includes a heavy chain variable region (SEQ ID NO: 50) encoded by the nucleic acid sequence shown below in SEQ ID NO: 49, and a light chain variable region (SEQ ID NO: 52) encoded by the nucleic acid sequence shown in SEQ ID NO: 51.

[50] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[51] The heavy chain CDRs of the 23K12 antibody have the following sequences per Kabat definition: SNYMS (SEQ ID NO: 103), VIYSGGSTYYADSVK (SEQ ID NO: 105) and CLSRMRGYGLDV (SEQ ID NO: 107). The light chain CDRs of the 23K12 antibody have the following sequences per Kabat definition: RTSQSISSYLN (SEQ ID NO: 92), AASSLQSGVPSRF (SEQ ID NO: 94) and QQSYSMMPA (SEQ ID NO: 96).

[52] The heavy chain CDRs of the 23K12 antibody have the following sequences per Chothia definition: GFTVSSN (SEQ ID NO: 112), VIYSGGSTY (SEQ ID NO: 113) and CLSRMRGYGLDV (SEQ ID NO: 107). The light chain CDRs of the 23K12 antibody have the following sequences per Chothia definition: RTSQSISSYLN (SEQ ID NO: 92), AASSLQSGVPSRF (SEQ ID NO: 94) and QQSYSMMPA (SEQ ID NO: 96).

**>23K12 VH nucleotide sequence: (SEQ ID NO: 49)**

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTCCAGCCTGGGGGGTCCCTGAGAATCTC  
CTGTGCAGCCTCTGGATTCACCGTCAGTAGCAACTACATGAGTTGGGTCCGCCAGGCTCCAG  
GGAAGGGGCTGGAGTGGGTCTCAGTTATTTATAGTGGTGGTAGCACATACTACGCAGACTCC  
GTGAAGGGCAGATTCTCCTTCTCCAGAGACAACTCCAAGAACACAGTGTTCCTTCAAATGAA  
CAGCCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGCGAGATGTCTGAGCAGGATGCGGG  
GTTACGGTTTAGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCG

**>23K12 VH amino acid sequence: (SEQ ID NO: 50)**

Kabat Bold, Chothia underlined

|          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| E        | V        | Q        | L        | V        | E        | S        | G        | G        | G        | L        | V        | Q        | P        | G        | G        | S        | L        | R        | I        | S        |
| C        | A        | A        | S        | <u>G</u> | <u>F</u> | <u>T</u> | <u>V</u> | <u>S</u> | <b>S</b> | <b>N</b> | <b>Y</b> | <b>M</b> | <b>S</b> | W        | V        | R        | Q        | A        | P        | G        |
| K        | G        | L        | E        | W        | V        | S        | <b>V</b> | <b>I</b> | <b>Y</b> | <b>S</b> | <b>G</b> | <b>G</b> | <b>S</b> | <b>T</b> | <b>Y</b> | <b>Y</b> | <b>A</b> | <b>D</b> | <b>S</b> | <b>V</b> |
| <b>K</b> | G        | R        | F        | S        | F        | S        | R        | D        | N        | S        | K        | N        | T        | V        | F        | L        | Q        | M        | N        | S        |
| L        | R        | A        | E        | D        | T        | A        | V        | Y        | Y        | C        | A        | R        | <b>C</b> | <b>L</b> | <b>S</b> | <b>R</b> | <b>M</b> | <b>R</b> | <b>G</b> | <b>Y</b> |
| <b>G</b> | <b>L</b> | <b>D</b> | <b>V</b> | W        | G        | Q        | G        | T        | T        | V        | T        | V        | S        |          |          |          |          |          |          |          |

**>23K12 VL nucleotide sequence: (SEQ ID NO: 51)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGACAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGGA  
AAGCCCCTAAACTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCGGTCTGCAACCTGAAGATTT  
TGCAACCTACTACTGTCAACAGAGTTACAGTATGCCTGCCTTTGGCCAGGGGACCAAGCTGG  
AGATCAAA

**>23K12 VL amino acid sequence: (SEQ ID NO: 52)**

Kabat Bold, Chothia underlined

|   |   |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |          |   |
|---|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|---|
| D | I | Q        | M        | T        | Q        | S        | P        | S        | S        | L        | S        | A        | S        | V        | G        | D        | R        | V        | T        | I |
| T | C | <b>R</b> | <b>T</b> | <b>S</b> | <b>Q</b> | <b>S</b> | <b>I</b> | <b>S</b> | <b>S</b> | <b>Y</b> | <b>L</b> | <b>N</b> | W        | Y        | Q        | Q        | K        | P        | G        | K |
| A | P | K        | L        | L        | I        | Y        | <b>A</b> | <b>A</b> | <b>S</b> | <b>S</b> | <b>L</b> | <b>Q</b> | <b>S</b> | <b>G</b> | <b>V</b> | <b>P</b> | <b>S</b> | <b>R</b> | <b>F</b> | S |
| G | S | G        | S        | G        | T        | D        | F        | T        | L        | T        | I        | S        | G        | L        | Q        | P        | E        | D        | F        | A |
| T | Y | Y        | C        | <u>Q</u> | <u>Q</u> | <b>S</b> | <b>Y</b> | <b>S</b> | <b>M</b> | <b>P</b> | <b>A</b> | F        | G        | Q        | G        | T        | K        | L        | E        | I |

[53] The 3241\_G23 antibody (also referred to herein as G23) includes antibody includes a heavy chain variable region (SEQ ID NO: 116) encoded by the nucleic acid sequence shown below in SEQ ID NO: 115, and a light chain variable region (SEQ ID NO: 118) encoded by the nucleic acid sequence shown in SEQ ID NO: 117.

[54] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[55] The heavy chain CDRs of the G23 antibody have the following sequences per Kabat definition: GGGYSWN (SEQ ID NO: 179), FMFHSGSPRYNPTLKS (SEQ ID NO: 180) and VGQMDKYYAMDV (SEQ ID NO: 181). The light chain CDRs of the G23 antibody have the following sequences per Kabat definition: RASQSIGAYVN (SEQ ID NO: 184), GASNLQS (SEQ ID NO: 185) and QQTYSTPIT (SEQ ID NO: 186).

[56] The heavy chain CDRs of the G23 antibody have the following sequences per Chothia definition: GGPVSGGG (SEQ ID NO: 182), FMFHSGSPR (SEQ ID NO: 183) and VGQMDKYYAMDV (SEQ ID NO: 181). The light chain CDRs of the G23 antibody have the following sequences per Chothia definition: RASQSIGAYVN (SEQ ID NO: 184), GASNLQS (SEQ ID NO: 185) and QQTYSTPIT (SEQ ID NO: 186).

**>3241\_G23 VH nucleotide sequence (SEQ ID NO: 115)**

CAGGTGCAGCTGCAGCAGTCGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTCAC  
TTGCACTGTCTCTGGTGGCCCCGTCAGCGGTGGTGGTTACTCCTGGAACCTGGATCCGCCAAC  
GCCCAGGACAGGGCCTGGAGTGGGTGGGTTCATGTTTCACAGTGGGAGTCCCCGCTACAAT  
CCGACCCTCAAGAGTCGAATTACCATCTCAGTCGACACGTCTAAGAACCTGGTCTCCCTGAA  
GCTGAGCTCTGTGACGGCCGCGGACACGGCCGTGTATTTTGTGCGCGAGTGGGGCAGATGG  
ACAAGTACTATGCCATGGACGTCTGGGGCCAAGGACCACGGTCACCGTCTCGAGC

**>3241\_G23 VH amino acid sequence (SEQ ID NO: 116)**

Kabat Bold, Chothia underlined

QVQLQQSGPGLVKPSQTL~~SLTCTV~~SGGPVS**GGGYSWN**WIRQRPQGQLEWVG**FMFHSGSPRYN**  
**PTLKS**RITISVDTSKNLVSLKLSSVTAADTAVYFCARVGQMDKYYAMDVWGQGT~~TVTVSS~~

**>3241\_G23 VL nucleotide sequence (SEQ ID NO: 117)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTTCTCTGTCTGGAGACAGAGTCACCAT  
 CACTTGCCGGGCAAGTCAGAGCATTGGCGCCTATGTAAATTGGTATCAACAGAAAGCAGGGA  
 AAGCCCCCAGGTCCTGATCTTTGGTGCTTCCAATTTACAAAGCGGGGTCCCATCAAGGTTTC  
 AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGACTT  
 TGCAACTTACTTCTGTCAACAGACTTACAGTACCCCGATCACCTTCGGCCAAGGGACACGAC  
 TGGAGATTAAACG

**>3241\_G23 VL amino acid sequence (SEQ ID NO: 118)**

DIQMTQSPSSLSSSVGDRVTITC**RASQSIGAYVN**WYQQKAGKAPQVLI**F****GASNLQS**GVPSRF  
 SGSGSGTDFTLTISSLQPEDFATYFC**QQTYSTPIT**FGQGRLEIK

[57] The 3244\_I10 antibody (also referred to herein as I10) includes antibody includes a heavy chain variable region (SEQ ID NO: 120) encoded by the nucleic acid sequence shown below in SEQ ID NO: 119, and a light chain variable region (SEQ ID NO: 122) encoded by the nucleic acid sequence shown in SEQ ID NO: 121.

[58] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991, are highlighted in bold in the sequences below.

[59] The heavy chain CDRs of the I10 antibody have the following sequences per Kabat definition: SDYWS (SEQ ID NO: 187), FFYNGGSTKYNPSLKS (SEQ ID NO: 188) and HDAKFSGSYVAS (SEQ ID NO: 189). The light chain CDRs of the I10 antibody have the following sequences per Kabat definition: RASQSISTYLN (SEQ ID NO: 192), GATNLQS (SEQ ID NO: 193) and QQSYNTPLI (SEQ ID NO: 194).

[60] The heavy chain CDRs of the I10 antibody have the following sequences per Chothia definition: GGSITS (SEQ ID NO: 190), FFYNGGSTK (SEQ ID NO: 191) and HDAKFSGSYVAS (SEQ ID NO: 189). The light chain CDRs of the I10 antibody have the following sequences per Chothia definition: RASQSISTYLN (SEQ ID NO: 192), GATNLQS (SEQ ID NO: 193) and QQSYNTPLI (SEQ ID NO: 194).

**>3244\_I10 VH nucleotide sequence (SEQ ID NO: 119)**

CAGGTCCAGCTGCAGGAGTCGGGCCCAGGACTGCTGAAGCCTTCGGACACCCTGGCCCTCAC  
 TTGCACTGTCTCTGGTGGCTCCATCACCAGTGACTIONTGGAGCTGGATCCGGCAACCCCCAG  
 GGAGGGGACTGGACTGGATCGGATTCTTCTATAACGGCGGAAGCACCAAGTACAATCCCTCC  
 CTCAAGAGTCGAGTCACCATTTTCAGCGGACACGTCCAAGAACCAGTTGTCCCTGAAATTGAC  
 CTCTGTGACCGCCGCAGACACGGGCGTGTATTATTGTGCGAGACATGATGCCAAATTTAGTG  
 GGAGCTACTACGTTGCCTCCTGGGGCCAGGGAACCCGAGTCACCGTCTCGAGC

**>3244\_I10 VH amino acid sequence (SEQ ID NO: 120)**

Kabat Bold, Chothia underlined

QVQLQESGPGLLKPSDTLALTCTVSGGSIT**SDYWS**WIRQPPGRGLDWIG**FFYNGGSTKYNPS**  
**LKSRVTISADTSKNQLSLKLT**SVTAADTGVYYCAR**HDAKFSGSYVAS**WGQGTRVTVSS

**>3244\_I10 VL nucleotide sequence (SEQ ID NO: 121)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
 CTCTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTAAATTGGTATCAGCAGCAACCTGGGA  
 AAGCCCCTAAGGTCCTCATTTTTTGGTGCAACCAACTTGCAAAGTGGGGTCCCATCTCGCTTC  
 AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTT  
 TGCAACTTACTACTGTCAACAGAGTTACAATACCCCCCTCATTTTTTGGCCAGGGGACCAAGC  
 TGGAGATCAAACG

**>3244\_I10 VL amino acid sequence (SEQ ID NO: 122)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITISC**RASQSISTYLN**WYQQQPGKAPKVLIF**GATNLQS**GVPSRF  
 SGSGSGTDFTLTISLQPEDFATYYC**QQSYNTPLI**FGQGTKLEIK

[61] The 3243\_J07 antibody (also referred to herein as J07) includes antibody includes a heavy chain variable region (SEQ ID NO: 124) encoded by the nucleic acid sequence shown below in SEQ ID NO: 123, and a light chain variable region (SEQ ID NO: 126) encoded by the nucleic acid sequence shown in SEQ ID NO: 125.

[62] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[63] The heavy chain CDRs of the J07 antibody have the following sequences per Kabat definition: SDYWS (SEQ ID NO: 187), FFYNGGSTKYNPSLKS (SEQ ID NO: 188) and HDVKFSGSYVAS (SEQ ID NO: 195). The light chain CDRs of the J07 antibody have the following sequences per Kabat definition: RASQSISTYLN (SEQ ID NO: 192), GATNLQS (SEQ ID NO: 193) and QQSYNTPLI (SEQ ID NO: 194).

[64] The heavy chain CDRs of the J07 antibody have the following sequences per Chothia definition: GGSITS (SEQ ID NO: 190), FFYNGGSTK (SEQ ID NO: 191) and HDVKFSGSYVAS (SEQ ID NO: 195). The light chain CDRs of the J07 antibody have the following sequences per Chothia definition: RASQSISTYLN (SEQ ID NO: 192), GATNLQS (SEQ ID NO: 193) and QQSYNTPLI (SEQ ID NO: 194).

**>3243\_J07 VH nucleotide sequence (SEQ ID NO: 123)**

CAGGTCCAGCTGCAGGAGTCGGGCCCAGGACTGCTGAAGCCTTCGGACACCCTGGCCCTCAC  
 TTGCACTGTCTCTGGTGGCTCCATCACCAGTGACTACTGGAGCTGGATCCGGCAACCCCCAG

GGAGGGGACTGGACTGGATCGGATTCTTCTATAACGGCGGGAGCACCAAGTACAATCCCTCC  
CTCAAGAGTCGAGTCACCATATCAGCGGACACGTCCAAGAACCAGTTGTCCCTGAAATTGAC  
CTCTGTGACCGCCGCAGACACGGGCGTGTATTATTGTGCGAGACATGATGTCAAATTTAGTG  
GGAGCTACTACGTTGCCTCCTGGGGCCAGGGAACCCGAGTCACCGTCTCGAGC

**>3243\_J07 VH amino acid sequence (SEQ ID NO: 124)**

Kabat Bold, Chothia underlined

QVQLQESGPGLLKPSDTLALTCTVSGGSIT**SDYWS**WIRQPPGRGLDWIG**FFYNGGSTKYNPS**  
**LKSRVT**ISADTSKNQLSLKLTSTVAADTGVYYCAR**HDVKFSGSYVAS**WGQGTRVTVSS

**>3243\_J07 VL nucleotide sequence (SEQ ID NO: 125)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
CTCTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTAAATTGGTATCAGCAGCAACCTGGGA  
AAGCCCCTAAGGTCCTGATCTCTGGTGCAACCAACTTGCAAAGTGGGGTCCCATCTCGCTTC  
AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTT  
TGCAACTTACTACTGTCAACAGAGTTACAATACCCCCCTCATTTTGGCCAGGGGACCAAGC  
TGGAGATCAAACG

**>3243\_J07 VL amino acid sequence (SEQ ID NO: 126)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITISC**RASQSISTYLN**WYQQQPGKAPKVLIS**GATNLQS**GVPSRF  
SGSGSGTDFTLTISLQPEDFATYYC**QQSYNTPLI**FGQGTKLEIK

[65] The 3259\_J21 antibody (also referred to herein as J21) includes antibody includes a heavy chain variable region (SEQ ID NO: 128) encoded by the nucleic acid sequence shown below in SEQ ID NO: 127, and a light chain variable region (SEQ ID NO: 130) encoded by the nucleic acid sequence shown in SEQ ID NO: 129.

[66] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[67] The heavy chain CDRs of the J21 antibody have the following sequences per Kabat definition: SYNWI (SEQ ID NO: 196), HIYDYGRTFYNSSLQS (SEQ ID NO: 197) and PLGILHYYAMD (SEQ ID NO: 198). The light chain CDRs of the J21 antibody have the following sequences per Kabat definition: RASQSIDKFLN (SEQ ID NO: 199), GASNLHS (SEQ ID NO: 200) and QQSFSVPA (SEQ ID NO: 201).

[68] The heavy chain CDRs of the J21 antibody have the following sequences per Chothia definition: GGSISS (SEQ ID NO: 202), HIYDYGRTF (SEQ ID NO: 203) and PLGILHYYAMD (SEQ ID NO: 198). The light chain CDRs of the J21 antibody have the

following sequences per Chothia definition: RASQSIDKFLN (SEQ ID NO: 199), GASNLHS (SEQ ID NO: 200) and QQSFSVPA (SEQ ID NO: 201).

**>3259\_J21 VH nucleotide sequence (SEQ ID NO: 127)**

CAGGTGCAGCTGCAGGAGTCGGGCCCCACGAGTGGTGAGGCCTTCGGAGACCCTGTCCCTCAC  
CTGCACTGTCTCGGGGGGCTCCATCAGTTCTTACAACCTGGATTTGGATCCGGCAGCCCCCTG  
GGAAGGGACTGGAGTGGATTGGGCACATATATGACTATGGGAGGACCTTCTACAACCTCCTCC  
CTCCAGAGTCGACCTACCATATCTGTAGACGCGTCCAAGAATCAGCTCTCCCTGCGATTGAC  
CTCTGTGACCGCCTCAGACACGGCCGTCTATTACTGTGCGAGACCTCTCGGTATACTCCACT  
ACTACGCGATGGACCTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3259\_J21 VH amino acid sequence (SEQ ID NO: 128)**

Kabat Bold, Chothia underlined

QVQLQESGPRVVRPSETLSLTCTVSGGSIS**SYNWI**WIRQPPGKLEWIG**HIYDYGRTFYNSS**  
**LQSRPTISVDASKNQLSRLTSVTASDTAVYYCAR**PLGILHYYAMDLWGQGTTVTVSS

**>3259\_J21 VL nucleotide sequence (SEQ ID NO: 129)**

GACATCCAGATGACCCAGTCTCCATTATCCGTGTCTGTATCTGTTCGGGGACAGGGTCACCAT  
CGCTTGCCGGGCAAGTCAGAGTATTGACAAGTTTTTAAATTGGTATCAGCAGAAACCAGGGA  
AAGCCCCTAAACTCCTGATCTATGGTGCCCTCAATTTGCACAGTGGGGCCCCATCAAGGTTT  
AGTGCCAGTGGGTCTGGGACAGACTTCACTCTAACAATCACCAATATACAGACTGAAGATTT  
CGCAACTTACCTCTGTCAACAGAGTTTCAGTGTCCCCGCTTTCGGCGGAGGGACCAAGGTTG  
AGATCAAACG

**>3259\_J21 VL amino acid sequence (SEQ ID NO: 130)**

Kabat Bold, Chothia underlined

DIQMTQSPLSVSVSGDRVTIAC**RASQSIDKFLN**WYQQKPGKAPKLLIY**GASNLHS**GAPSRF  
SASGSGTDFTLTITNIQTEDFATYLC**QQSFSVPA**FGGGTKVEIK

[69] The 3245\_O19 antibody (also referred to herein as O19) includes a heavy chain variable region (SEQ ID NO: 132) encoded by the nucleic acid sequence shown below in SEQ ID NO: 131, and a light chain variable region (SEQ ID NO: 134) encoded by the nucleic acid sequence shown in SEQ ID NO: 133.

[70] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[71] The heavy chain CDRs of the O19 antibody have the following sequences per Kabat definition: STYMN (SEQ ID NO: 204), VFYSETRTYADSVKG (SEQ ID NO: 205) and VQRLSYGMDV (SEQ ID NO: 206). The light chain CDRs of the O19 antibody have the following sequences per Kabat definition: RASQSISTYLN (SEQ ID NO: 192), GASTLQS (SEQ ID NO: 207) and QQTYSIPL (SEQ ID NO: 208).

[72] The heavy chain CDRs of the O19 antibody have the following sequences per Chothia definition: GLSVSS (SEQ ID NO: 209), VFYSETRTY (SEQ ID NO: 210) and VQRLSYGMDV (SEQ ID NO: 206). The light chain CDRs of the O19 antibody have the following sequences per Chothia definition: RASQSISTYLN (SEQ ID NO: 192), GASTLQS (SEQ ID NO: 207) and QQTYSIPL (SEQ ID NO: 208).

**>3245\_O19 VH nucleotide sequence (SEQ ID NO: 131)**

GAGGTGCAACTGGTGGAGTCTGGAGGGGGCTTGGTCCAGCCTGGGGGGTCCCTGAGACTCTC  
CTGTACGGCCTCTGGGTAAAGTGTGTCAGTTCCACCTACATGAACTGGGTCCGCCAGGCTCCAG  
GGAAGGGGCTGGAATGGGTCTCAGTTTTTTATAGTGAGACCAGGACGTACTACGCAGACTCC  
GTGAAGGGCCGATTACCGTCTCCAGACACAATTCCAACAACACGCTCTATCTTCAGATGAA  
CAGCCTGAGAGTTGAAGACACGGCCGTGTATTATTGTGCGAGAGTCCAGAGATTGTCGTACG  
GTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3245\_O19 VH amino acid sequence (SEQ ID NO: 132)**

Kabat Bold, Chothia underlined

EVQLVESGGGLVQPGGSLRLSCTASGLSVS**STYMN**WVRQAPGKGLEWVSVFYSETRTY**YADS**  
**VKGR**FTVSRHNSNNTLYLQMNSLRVEDTAVYYCARVQRLSYGMDVWGQGTITVTVSS

**>3245\_O19 VL nucleotide sequence (SEQ ID NO: 133)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTTGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTAAATTGGTATCAGAAGAGACCAGGGA  
AAGCCCCTAAACTCCTGGTCTATGGTGCATCCACTTTGCAGAGTGGGGTCCCATCAAGGTTT  
AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCGCCAGTCTGCAACCTGAAGATTC  
TGCAACTTACTACTGTCAACAGACTTACAGTATCCCCCTCTTCGGCCAGGGGACACGGCTGG  
AGATTAAACG

**>3245\_O19 VL amino acid sequence (SEQ ID NO: 134)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGRVTITCRASQSISTYLNWYQKRPKAPKLLVYGASTLQSGVPSRF  
SGSGSGTDFTLTIASLQPEDSATYYCQQTYSIPLFGQGTGLEIK

[73] The 3244\_H04 antibody (also referred to herein as H04) includes antibody includes a heavy chain variable region (SEQ ID NO: 136) encoded by the nucleic acid sequence shown below in SEQ ID NO: 135, and a light chain variable region (SEQ ID NO: 138) encoded by the nucleic acid sequence shown in SEQ ID NO: 137.

[74] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[75] The heavy chain CDRs of the H04 antibody have the following sequences per Kabat definition: STYMN (SEQ ID NO: 204), VFYSETRTYADSVKG (SEQ ID NO: 205) and

VQRLSYGMDV (SEQ ID NO: 206). The light chain CDRs of the H04 antibody have the following sequences per Kabat definition: RASQSISTYLN (SEQ ID NO: 192), GASSLQS (SEQ ID NO: 211) and QQTYSIPL (SEQ ID NO: 208).

[76] The heavy chain CDRs of the H04 antibody have the following sequences per Chothia definition: GLSVSS (SEQ ID NO: 209), VFYSETRTY (SEQ ID NO: 210) and VQRLSYGMDV (SEQ ID NO: 206). The light chain CDRs of the H04 antibody have the following sequences per Chothia definition: RASQSISTYLN (SEQ ID NO: 192), GASSLQS (SEQ ID NO: 211) and QQTYSIPL (SEQ ID NO: 208).

**>3244\_H04 VH nucleotide sequence (SEQ ID NO: 135)**

GAGGTGCAGCTGGTGGGAATCTGGAGGGGGCTTGGTCCAGCCTGGGGGGTCCCTGAGACTCTC  
CTGTACAGCCTCTGGGTAAAGCGTCAGTTCCACCTACATGAACTGGGTCCGCCAGGCTCCAG  
GGAAGGGGCTGGAATGGGTCTCAGTTTTTTATAGTGAAACCAGGACGTATTACGCAGACTCC  
GTGAAGGGCCGATTCACCGTCTCCAGACACAATTCCAACAACACGCTGTATCTTCAAATGAA  
CAGCCTGAGAGCTGAAGACACGGCCGTGTATTATTGTGCGAGAGTCCAGAGACTGTCATACG  
GTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3244\_H04 VH amino acid sequence (SEQ ID NO: 136)**

Kabat Bold, Chothia underlined

EVQLVESGGGLVQPGGSLRLSCTASGLSVS**STYMN**WVRQAPGKGLEWVS**VFYSETRTYADS**  
**VKGRFT**VSRLHNSNNTLYLQMNSLRAEDTAVYYCAR**VQRLSYGMDV**WGQGTTVTVSS

**>3244\_H04 VL nucleotide sequence (SEQ ID NO: 137)**

GACATCCAGATGACCCAGTCTCCATCGTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTAAATTGGTATCAGAAGAGACCAGGGA  
AAGCCCCTAAACTCCTGGTCTATGGTGCATCCAGTTTGCAGAGTGGGGTCCCATCAAGGTTT  
AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCGCCAGTCTGCAACCTGAAGATTC  
TGCAGTTTATTACTGTCAACAGACTTACAGTATCCCCCTCTTCGGCCAGGGGACACGACTGG  
AGATTAAACG

**>3244\_H04 VL amino acid sequence (SEQ ID NO: 138)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITTC**RASQSISTYLN**WYQKRPGKAPKLLVY**GASSLQS**GVPSRF  
SGSGSGTDFTLTIASLQPEDSAVYYC**QQTYSIPL**FGQGTTRLEIK

[77] The 3136\_G05 antibody (also referred to herein as G05) includes antibody includes a heavy chain variable region (SEQ ID NO: 140) encoded by the nucleic acid sequence shown below in SEQ ID NO: 139, and a light chain variable region (SEQ ID NO: 142) encoded by the nucleic acid sequence shown in SEQ ID NO: 141.

[78] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[79] The heavy chain CDRs of the G05 antibody have the following sequences per Kabat definition: SDFWS (SEQ ID NO: 212), YVYNRGSTKYSPSLKS (SEQ ID NO: 213) and NGRSSTSWGIDV (SEQ ID NO: 214). The light chain CDRs of the G05 antibody have the following sequences per Kabat definition: RASQSISTYLH (SEQ ID NO: 215), AASSLQS (SEQ ID NO: 216) and QQSYSPPLT (SEQ ID NO: 63).

[80] The heavy chain CDRs of the G05 antibody have the following sequences per Chothia definition: GGSISS (SEQ ID NO: 202), YVYNRGSTK (SEQ ID NO: 217) and NGRSSTSWGIDV (SEQ ID NO: 214). The light chain CDRs of the G05 antibody have the following sequences per Chothia definition: RASQSISTYLH (SEQ ID NO: 215), AASSLQS (SEQ ID NO: 216) and QQSYSPPLT (SEQ ID NO: 63).

**>3136\_G05 VH nucleotide sequence (SEQ ID NO: 139)**

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCCTCGGAGACCCTGTCCCTCAC  
CTGCAGTGTCTCTGGTGGCTCCATTAGTAGTGATTTCTGGAGTTGGATCCGACAGCCCCCAG  
GGAAGGGACTGGAGTGGATTGGGTATGTCTATAACAGAGGGAGCACTAAGTACAGTCCCTCC  
CTCAAGAGTCGAGTCACCATATCAGCAGACATGTCCAAGAACCAGTTTTCCCTGAATATGAG  
TTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGAAAAATGGTCAAGTAGCACCA  
GTTGGGGCATCGACGTCTGGGGCAAAGGGACCACGGTCACCGTCTCGAGC

**>3136\_G05 VH amino acid sequence (SEQ ID NO: 140)**

Kabat Bold, Chothia underlined

QVQLQESGPGLVKPSETLSLTCSVSGGSISSDFWSWIRQPPGKGLEWIGYVYNRGSTKYSPS  
LKSRVTISADMSKNQFSLNMSSVTAADTAVYYCAKNGRSSTSWGIDVWGKGTTTVTSS

**>3136\_G05 VL nucleotide sequence (SEQ ID NO: 141)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTGGGAGACAGACTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTACATTGGTATCAGCAGAAACCAGGGA  
AAGCCCCTAACTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGGCAGTAGATCAGGAACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGATGACTT  
TGCAACTTACTACTGTCAACAGAGTTACAGTCCCCCCTCACTTTCGGCCCTGGGACCAAAG  
TGGATATGAAACG

**>3136\_G05 VL amino acid sequence (SEQ ID NO: 142)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRLTITCRASQSISTYLHWYQQKPGKAPKLLIYAASSLQSGVPSRF  
SGSRSGTDFTLTISLQPDFFATYYCQQSYSPPLTFGPGTKVDMK

[81] The 3252\_C13 antibody (also referred to herein as C13) includes antibody includes a heavy chain variable region (SEQ ID NO: 144) encoded by the nucleic acid sequence shown below in SEQ ID NO: 143, and a light chain variable region (SEQ ID NO: 146) encoded by the nucleic acid sequence shown in SEQ ID NO: 145.

[82] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[83] The heavy chain CDRs of the C13 antibody have the following sequences per Kabat definition: SDYWS (SEQ ID NO: 187), YIYNRGSTKYTPSLKS (SEQ ID NO: 218) and HVGGHTYGIDY (SEQ ID NO: 219). The light chain CDRs of the C13 antibody have the following sequences per Kabat definition: RASQSIISNYLN (SEQ ID NO: 220), AASSLQS (SEQ ID NO: 216) and QQSYNTPIT (SEQ ID NO: 221).

[84] The heavy chain CDRs of the C13 antibody have the following sequences per Chothia definition: GASISS (SEQ ID NO: 222), YIYNRGSTK (SEQ ID NO: 223) and HVGGHTYGIDY (SEQ ID NO: 219). The light chain CDRs of the C13 antibody have the following sequences per Chothia definition: RASQSIISNYLN (SEQ ID NO: 220), AASSLQS (SEQ ID NO: 216) and QQSYNTPIT (SEQ ID NO: 221).

**>3252\_C13 VH nucleotide sequence (SEQ ID NO: 143)**

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCAC  
CTGCACTGTCTCTGGTGCTCCATCAGTAGTGACTACTGGAGCTGGATCCGGCTGCCCCCAG  
GGAAGGGACTGGAGTGGATTGGGTATATCTATAATAGAGGGAGTACCAAGTACACCCCTCC  
CTGAAGAGTCGAGTCACCATATCACTAGACACGGCCGAGAACCAGTTCTCCCTGAGGCTGAG  
GTCGGTGACCGCCGCAGACACGGCCATCTATTACTGTGCGAGACATGTAGGTGGCCACACCT  
ATGGAATTGATTACTGGGGCCAGGGAACCCTGGTCACCGTCTCGAGC

**>3252\_C13 VH amino acid sequence (SEQ ID NO: 144)**

Kabat Bold, Chothia underlined

QVQLQESGPGLVKPSSETLSLTCTVSGASIS**SDYWS**WIRLPPGKGLEWIG**YIYNRGSTKYTPS**  
**LKSRVTISLDTAENQFSLRLRSVTAADTAIYYCAR**HVGGHTYGIDYWGQGTLLTVSS

**>3252\_C13 VL nucleotide sequence (SEQ ID NO: 145)**

GACATCCAGATGACCCAGTCTCCATCGTCCCTGTCTGCCTCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCAACTATTTAAATTGGTATCAACACAAACCTGGGG  
AAGCCCCCAAGCTCCTGAACTATGCTGCGTCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGCCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTTCAACCTGAAGATTT  
TGCCACTTACTACTGTCAACAGAGTTACAATACTCCGATCACCTTCGGCCAAGGGACACGAC  
TGGAATTAACG

**>3252\_C13 VL amino acid sequence (SEQ ID NO: 146)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITTC**RASQ****SISNYLN**WYQHKPGEAPKLLNY**AASSLQS**GVPSRF  
SASGSGTDFTLTISSLQPEDFATYYC**QQSYNTPIT**FGQGTRLEIK

[85] The 3259\_J06 antibody (also referred to herein as J06) includes antibody includes a heavy chain variable region (SEQ ID NO: 148) encoded by the nucleic acid sequence shown below in SEQ ID NO: 147, and a light chain variable region (SEQ ID NO: 150) encoded by the nucleic acid sequence shown in SEQ ID NO: 149.

[86] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[87] The heavy chain CDRs of the J06 antibody have the following sequences per Kabat definition: SDYWS (SEQ ID NO: 187), YIYNRGSTKYTPSLKS (SEQ ID NO: 218) and HVGGHTYGIDY (SEQ ID NO: 219). The light chain CDRs of the J06 antibody have the following sequences per Kabat definition: **RASQ****SISNYLN** (SEQ ID NO: 220), **AASSLQS** (SEQ ID NO: 216) and **QQSYNTPIT** (SEQ ID NO: 221).

[88] The heavy chain CDRs of the J06 antibody have the following sequences per Chothia definition: GASISS (SEQ ID NO: 222), YIYNRGSTK (SEQ ID NO: 223) and HVGGHTYGIDY (SEQ ID NO: 219). The light chain CDRs of the J06 antibody have the following sequences per Chothia definition: **RASQ****SISNYLN** (SEQ ID NO: 220), **AASSLQS** (SEQ ID NO: 216) and **QQSYNTPIT** (SEQ ID NO: 221).

**>3255\_J06 VH nucleotide sequence (SEQ ID NO: 147)**

CAGGTGCAGCTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCAC  
CTGCACTGTCTCTGGTGCCTCCATCAGTAGTGACTACTGGAGCTGGATCCGGCTGCCCCCAG  
GGAAGGGACTGGAGTGGATTGGGTATATCTATAATAGAGGGAGTACCAAGTACACCCCCTCC  
CTGAAGAGTCGAGTCACCATATCACTAGACACGGCCGAGAACCAGTTCTCCCTGAGGCTGAG  
GTCGGTGACCGCCGACACACGGCCGTCTATTACTGTGCGAGACATGTGGGTGGCCACACCT  
ATGGAATTGATTACTGGGGCCAGGGAACCCTGGTCACCGTCTCGAGC

**>3255\_J06 VH amino acid sequence (SEQ ID NO: 148)**

Kabat Bold, Chothia underlined

QVQLQESGPGLVKPSETLSLTCTVSGASIS**SDYWS**WIRLPPGKGLEWIG**YIYNRGSTKYTPS**  
**LKSRVTISLDTAENQFSLRLRSVTAADTAVYYCAR****HVGGHTYGIDY**WGQGLTVTVSS

**>3255\_J06 VL nucleotide sequence (SEQ ID NO: 149)**

GACATCCAGATGACCCAGTCTCCATCGTCCCTGTCTGCCTCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCAACTATTTAAATTGGTATCAACACAAACCTGGGG  
AAGCCCCCAAGCTCCTGAACTATGCTGCGTCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGCCAGTGGATCTGGGACAGATTTCACTCTCAGCATCAGCGGTCTTCAACCTGAAGATTT

TGCCACTTACTACTGTCAACAGAGCTACAATACTCCGATCACCTTCGGCCCAGGGACACGAC  
TGGAAATTAAACG

**>3255\_J06 VL amino acid sequence (SEQ ID NO: 150)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITTCRASQSISNYLNWYQHKPGEAPKLLNYAASSLQSGVPSRF  
SASGSGTDFTLISGLQPEDFATYYCQQSYNTPITFGPGTRLEIK

[89] The 3410\_I23 antibody (also referred to herein as I23) includes antibody includes a heavy chain variable region (SEQ ID NO: 152) encoded by the nucleic acid sequence shown below in SEQ ID NO: 151, and a light chain variable region (SEQ ID NO: 154) encoded by the nucleic acid sequence shown in SEQ ID NO: 153.

[90] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[91] The heavy chain CDRs of the I23 antibody have the following sequences per Kabat definition: SYSWS (SEQ ID NO: 224), YLYYSGSTKYNPSLKS (SEQ ID NO: 225) and TGSESTTGYGMDV (SEQ ID NO: 226). The light chain CDRs of the I23 antibody have the following sequences per Kabat definition: RASQSISTYLN (SEQ ID NO: 192), AASSLHS (SEQ ID NO: 227) and QQSYSPIT (SEQ ID NO: 228).

[92] The heavy chain CDRs of the I23 antibody have the following sequences per Chothia definition: GDSISS (SEQ ID NO: 229), YLYYSGSTKS (SEQ ID NO: 230) and TGSESTTGYGMDV (SEQ ID NO: 226). The light chain CDRs of the I23 antibody have the following sequences per Chothia definition: RASQSISTYLN (SEQ ID NO: 192), AASSLHS (SEQ ID NO: 227) and QQSYSPIT (SEQ ID NO: 228).

**>3420\_I23 VH nucleotide sequence (SEQ ID NO: 151)**

CAGGTGCAGCTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCGTCAC  
CTGCAAAGTCTCTGGTGACTCCATCAGTAGTTATTCCTGGAGCTGGATCCGGCAGCCCCCAG  
GGAAGGGACTGGAGTGGGTTGGCTATTTGTATTATAGTGGGAGCACCAAGTACAACCCCTCC  
CTCAAGAGTCGAACCACCATATCAGTAGACACGTCCACGAACCAGTTGTCCCTGAAGTTGAG  
TTTTGTGACCGCCGCGGACACGGCCGTGTATTTCTGTGCGAGAACCGGCTCGGAATCTACTA  
CCGGCTACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3420\_I23 VH amino acid sequence (SEQ ID NO: 152)**

Kabat Bold, Chothia underlined

QVQLQESGPGLVKPSETLSVTCKVSGDSIS**SYSWS**WIRQPPGKLEWVGYLYYSGSTKYNPS  
**LKS**RTTISVDTSTNQSLKLSFVTAADTAVYFCARTGSESTTGYGMDVWGQGTQTVTVSS

**>3420\_I23 VL nucleotide sequence (SEQ ID NO: 153)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
 CACTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTAAATTGGTATCAGCAGAAACCAGGGA  
 AAGCCCCTAAGCTCCTGATCTATGCTGCATCCAGTTTGCACAGTGGGGTCCCATCAAGGTTT  
 AGTGGCAGTGGATCTGGGACAGATTTGCTCTCACCATCAGCAGTCTGCAACCTGAAGATTT  
 TGCAACTTACTACTGTCAACAGAGTTACAGTCCCCCGATCACCTTCGGCCAAGGGACACGAC  
 TGGAGATTAAACG

**>3420\_I23 VL amino acid sequence (SEQ ID NO: 154)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITTC**RASQSISTYLN**WYQQKPGKAPKLLIYAASSLHSGVPSRF  
 SGSGSGTDFALTISSLQPEDFATYYCQQSYSPPITFGQGRLEIK

[93] The 3139\_P23 antibody (also referred to herein as P23) includes antibody includes a heavy chain variable region (SEQ ID NO: 156 ) encoded by the nucleic acid sequence shown below in SEQ ID NO:155, and a light chain variable region (SEQ ID NO:158) encoded by the nucleic acid sequence shown in SEQ ID NO:157.

[94] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[95] The heavy chain CDRs of the P23 antibody have the following sequences per Kabat definition: NSFVG (SEQ ID NO: 230), YVYNSGNTKYNPSLKS (SEQ ID NO: 231) and HDDASHGYSIS (SEQ ID NO: 232). The light chain CDRs of the P23 antibody have the following sequences per Kabat definition: RASQTISTYLN (SEQ ID NO: 233), AASGLQS (SEQ ID NO: 61) and QQSYNTPLT (SEQ ID NO: 234).

[96] The heavy chain CDRs of the P23 antibody have the following sequences per Chothia definition: GGSISN (SEQ ID NO: 258), YVYNSGNTK (SEQ ID NO: 259) and HDDASHGYSIS (SEQ ID NO: 232). The light chain CDRs of the P23 antibody have the following sequences per Chothia definition: RASQTISTYLN (SEQ ID NO: 233), AASGLQS (SEQ ID NO: 61) and QQSYNTPLT (SEQ ID NO: 234).

**>3139\_P23 VH nucleotide sequence (SEQ ID NO: 155)**

CAGGTGCAGCTGCAGGAGTCGGGCCCAAGACTGGTGAAGCCTTCGGAGAGCCTGTCCCTCAC  
 CTGCACTGTCTCTGGTGGCTCCATTAGTAATTCCTTCTGGGGCTGGATCCGGCAGCCCCCAG  
 GGGAGGGACTGGAGTGGATTGGTTATGTCTATAACAGTGGCAACACCAAGTACAATCCCTCC  
 CTAAGAGTCGAGTCACCATTTGCGCGACACGTCCAAGAGTCAACTCTACATGAAGCTGAG  
 GTCTGTGACCGCCGCTGACACGGCGTGTACTACTGTGCGAGGCATGACGACGCAAGTCATG  
 GCTACAGCATCTCCTGGGGCCACGGAACCCTGGTCACCGTCTCGAGC

**>3139\_P23 VH amino acid sequence (SEQ ID NO: 156)**

Kabat Bold, Chothia underlined

QVQLQESGPRLVKPSESLSLTCTVSGGSIS**NSFWG**WIRQPPGEGLEWIG**YVYNSGNTKYNPS**  
**LKSRVTISRDT**SKSQLYMKLRSVTAADTAVYYCAR**HDDASHGYSIS**WGHTLVTVSS

**>3139\_P23 VL nucleotide sequence (SEQ ID NO: 157)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGGGACAGAGTCACCAT  
 CACTTGCCGGGCAAGTCAGACCATTAGTACTTATTTAAATTGGTATCAACAGAAATCAGGGA  
 AAGCCCCTAAGCTCCTGATCTATGCTGCATCCGGTTTGCAAAGTGGAGTCCCATCAAGGTTT  
 AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTTCAACCTGAAGATTT  
 TGCAACTTACTTCTGTCAACAGAGTTACAATACTCCCTGACGTTTCGGCCAAGGGACCAAGG  
 TGGAAATCAAA

**>3139\_P23 VL amino acid sequence (SEQ ID NO: 158)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITITC**RASQTISTYLN**WYQQKSGKAPKLLIY**AASGLQS**GVPSRF  
 SGSGSGTDFTLTISLQPEDFATYFC**QQSYNTPLT**FGQGTKVEIK

[97] The 3248\_P18 antibody (also referred to herein as P18) includes antibody includes a heavy chain variable region (SEQ ID NO:160 ) encoded by the nucleic acid sequence shown below in SEQ ID NO:159 , and a light chain variable region (SEQ ID NO:162) encoded by the nucleic acid sequence shown in SEQ ID NO:161.

[98] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[99] The heavy chain CDRs of the P18 antibody have the following sequences per Kabat definition: AYHWS (SEQ ID NO: 235), HIFDSGSTYYNPSLKS (SEQ ID NO: 236) and PLGSRYYYGMDV (SEQ ID NO: 237). The light chain CDRs of the P18 antibody have the following sequences per Kabat definition: RASQSIERYLN (SEQ ID NO: 238), GASTLQN (SEQ ID NO: 239) and QQSYSVPA (SEQ ID NO: 240).

[100] The heavy chain CDRs of the P18 antibody have the following sequences per Chothia definition: GGSISA (SEQ ID NO: 260), HIFDSGSTY (SEQ ID NO: 261) and PLGSRYYYGMDV (SEQ ID NO: 237). The light chain CDRs of the P18 antibody have the following sequences per Chothia definition: RASQSIERYLN (SEQ ID NO: 238), GASTLQN (SEQ ID NO: 239) and QQSYSVPA (SEQ ID NO: 240).

**>3248\_P18 VH nucleotide sequence (SEQ ID NO: 159)**

CAGGTGCAACTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCAC  
 CTGCACTGTCTCGGGTGGCTCCATCAGTGCTTACCACTGGAGCTGGATCCGCCAGCCCCCAG  
 GGAAGGGACTGGAGTGGATTGGGCACATCTTTGACAGTGGGAGCACTTACTACAACCCCTCC

CTTAAGAGTCGAGTCACCATATCACTAGACGCGTCCAAGAACCAGCTCTCCCTGAGATTGAC  
CTCTGTGACCGCCTCAGACACGGCCATATATTACTGTGCGAGACCTCTCGGGAGTCGGTACT  
ATTACGGAATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3248\_P18 VH amino acid sequence (SEQ ID NO: 160)**

Kabat Bold, Chothia underlined

QVQLQESGPGGLVKPSETLSLTCTVSGGSIS**AYHWS**WIRQPPGKGLEWIG**HIFDSGSTYYNPS**  
**LKSRVTISLDASKNQLSRLTSVTASDTAIYCAR**PLGSRYYYGMDVWGQGTTVTVSS

**>3248\_P18 VL nucleotide sequence (SEQ ID NO: 161)**

GACATCCAGATGACCCAGTCTCCGTCCTCCCTGTCTGCATCTGTCTCGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGTATTAGCAGGTATTTAAATTGGTATCAGCAGAAACCAGGGA  
AAGCCCCTAAGCTCCTGATCTATGGTGCCTCCACTTTGCAAAATGGGGCCCCATCAAGGTTTC  
AGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTACAACCTGAAGATTC  
CGCAACTTACCTCTGTCAACAGAGTTACAGTGTCCCTGCTTTCGGCGGAGGAACCAAGGTGG  
AGGTCAAA

**>3248\_P18 VL amino acid sequence (SEQ ID NO: 162)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITITCRASQ**SISRYLN**WYQQKPGKAPKLLIYGASTLQNGAPSRF  
SGSGSGTDFTLTISSSLQPEDSATYLCQQSY**SVPA**FGGGTKVEVK

[101] The 3253\_P10 antibody (also referred to herein as P10) includes antibody includes a heavy chain variable region (SEQ ID NO: 164) encoded by the nucleic acid sequence shown below in SEQ ID NO: 163, and a light chain variable region (SEQ ID NO: 166) encoded by the nucleic acid sequence shown in SEQ ID NO: 165.

[102] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[103] The heavy chain CDRs of the P10 antibody have the following sequences per Kabat definition: SDYWS (SEQ ID NO: 187), FFYNGGSTKYNPSLKS (SEQ ID NO: 188) and HDAKFSGSYVAS (SEQ ID NO: 189). The light chain CDRs of the P10 antibody have the following sequences per Kabat definition: RASQSISTYLN (SEQ ID NO: 192), GATDLQS (SEQ ID NO: 241) and QQSYNTPLI (SEQ ID NO: 194).

[104] The heavy chain CDRs of the P10 antibody have the following sequences per Chothia definition: GGSITS (SEQ ID NO: 190), FFYNGGSTK (SEQ ID NO: 191) and HDAKFSGSYVAS (SEQ ID NO: 189). The light chain CDRs of the P10 antibody have the following sequences per Chothia definition: RASQSISTYLN (SEQ ID NO: 192), GATDLQS (SEQ ID NO: 241) and QQSYNTPLI (SEQ ID NO: 194).

**>3253\_P10 VH nucleotide sequence (SEQ ID NO: 163)**

CAGGTCCAGCTGCAGGAGTCGGGCCAGGACTGCTGAAGCCTTCGGACACCCCTGGCCCTCAC  
 TTGCACTGTCTCTGGTGGCTCCATCACCAGTGACTACTGGAGCTGGATCCGGCAACCCCCAG  
 GGAGGGGACTGGACTGGATCGGATTCTTCTATAACGGCGGGAGCACCAAGTACAATCCCTCC  
 CTAAGAGTCGAGTCACCATATCAGCGGACACGTCCAAGAACCAGTTGTCCCTGAAATTGAC  
 CTCTGTGACCGCCGCAGACACGGGCGTGTATTATTGTGCGAGACATGATGCCAAATTTAGTG  
 GGAGCTACTACGTTGCCTCCTGGGGCCAGGGAACCCGAGTCACCGTCTCGAGC

**>3253\_P10 VH amino acid sequence (SEQ ID NO: 164)**

Kabat Bold, Chothia underlined

QVQLQESGPGLLKPSDTLALTCTVSGGSIT**SDYWS**WIRQPPGRGLDWIG**FFYNGGSTKYNPS**  
**LKSRVT**ISADTSKNQLSLKLTSVTAADTGVYYCAR**HDAKFSGSYYVAS**WGQGTRVTVSS

**>3253\_P10 VL nucleotide sequence (SEQ ID NO: 165)**

GACATCCAGATGACCCAGTCTCCCTCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
 CTCTTGCCGGGCAAGTCAGAGCATTAGCACCTATTTAAATTGGTATCAGCAGCAACCTGGGA  
 AAGCCCCTAAGGTCCTGATCTCTGGTGAACCGACTTGCAAAGTGGGGTCCCATCTCGCTTC  
 AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTT  
 TGCAACTTACTACTGTCAACAGAGTTACAATACCCCCCTCATTTTTGGCCAGGGGACCAAGC  
 TGGAGATCAAA

**>3253\_P10 VL amino acid sequence (SEQ ID NO: 166)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITIS**RASQSIS**TYLNWYQQQPGKAPKVLIS**GATDLQS**GVPSRF  
 SGSGSGTDFTLTISSSLQPEDFATYYC**QQSYNTPLI**FGQGTKLEIK

[105] The 3260\_D19 antibody (also referred to herein as D19) includes antibody includes a heavy chain variable region (SEQ ID NO: 168) encoded by the nucleic acid sequence shown below in SEQ ID NO: 167, and a light chain variable region (SEQ ID NO: 170) encoded by the nucleic acid sequence shown in SEQ ID NO: 169.

[106] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[107] The heavy chain CDRs of the D19 antibody have the following sequences per Kabat definition: DNYIN (SEQ ID NO: 242), VFYSADRTSYADSVKG (SEQ ID NO: 243) and VQKSYYGMDV (SEQ ID NO: 244). The light chain CDRs of the D19 antibody have the following sequences per Kabat definition: RASQSISRYLN (SEQ ID NO: 238), GASSLQS (SEQ ID NO: 211) and QQTFSIPL (SEQ ID NO: 245).

[108] The heavy chain CDRs of the D19 antibody have the following sequences per Chothia definition: GFSVSD (SEQ ID NO: 247), VFYSADRTS (SEQ ID NO: 246) and VQKSYYGMDV (SEQ ID NO: 244). The light chain CDRs of the D19 antibody have the

following sequences per Chothia definition: RASQSIERYLN (SEQ ID NO: 238), GASSLQS (SEQ ID NO: 211) and QQTFSIPL (SEQ ID NO: 245).

**>3260\_D19 VH nucleotide sequence (SEQ ID NO: 167)**

GACATGCAGCTGGTGGAGTCTGGAGGAGGCTTGGTCCCGCCGGGGGGGTCCCTGAGACTCTC  
CTGCGCAGCCTCTGGGTTTTCCGTCAGTGACAACTACATAAACTGGGTCCGCCAGGCTCCAG  
GGAAGGGGCTGGACTGGGTCTCAGTCTTTTATAGTGCTGATAGAACATCCTACGCAGACTCC  
GTGAAGGGCCGATTACCGTCTCCAGCCACGATTCCAAGAACACAGTGTACCTTCAAATGAA  
CAGTCTGAGAGCTGAGGACACGGCCGTTTATTACTGTGCGAGAGTTCAGAAGTCCTATTACG  
GTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3260\_D19 VH amino acid sequence (SEQ ID NO: 168)**

Kabat Bold, Chothia underlined

DMQLVESGGGLVPPGGSLRLSCAASGFSVS**DN**YINWVRQAPGKGLDWVSVFYSADRTSYADS  
**VKGR**FTVSSHDSKNTVYLQMNSLRAEDTAVYYCARVQKSYYGMDVWGQGTITVTVSS

**>3260\_D19 VL nucleotide sequence (SEQ ID NO: 169)**

GGCATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCAGATATTTAAATTGGTATCTGCAGAAACCAGGGA  
AAGCCCCTAAGCTCCTGATCTCTGGTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGGCACTGGGTCTGGGACAGAATTCAGTCTCACCATCAGCAGTTTGCAACCTGAAGATTT  
TGCAACTTACTACTGTCAACAGACTTTCAGTATCCCTCTTTTTTGGCCAGGGGACCAAGGTGG  
AGATCAAA

**>3260\_D19 VL amino acid sequence (SEQ ID NO: 170)**

Kabat Bold, Chothia underlined

GIQMTQSPSSLSASVGDRVITITCRASQSIERYLNWYLQKPGKAPKLLISGASSLQSGVPSRF  
SGTGSGETFTLTISLQPEDFATYYCQQTFSIPLFGQGTKVEIK

[109] The 3362\_B11 antibody (also referred to herein as B11) includes antibody includes a heavy chain variable region (SEQ ID NO: 172) encoded by the nucleic acid sequence shown below in SEQ ID NO: 171, and a light chain variable region (SEQ ID NO: 174) encoded by the nucleic acid sequence shown in SEQ ID NO: 173.

[110] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[111] The heavy chain CDRs of the B11 antibody have the following sequences per Kabat definition: SGAYYWT (SEQ ID NO: 248), YIYYSGNTYYNPSLKS (SEQ ID NO: 249) and AASTSVLGYGMDV (SEQ ID NO: 250). The light chain CDRs of the B11 antibody have

the following sequences per Kabat definition: RASQISIRYLN (SEQ ID NO: 238), AASSLQS (SEQ ID NO: 216) and QQSYSTPLT (SEQ ID NO: 251).

[112] The heavy chain CDRs of the B11 antibody have the following sequences per Chothia definition: GDSITSGA (SEQ ID NO: 252), YIYYSGNTY (SEQ ID NO: 253) and AASTSVLGYGMDV (SEQ ID NO: 250). The light chain CDRs of the B11 antibody have the following sequences per Chothia definition: RASQISIRYLN (SEQ ID NO: 238), AASSLQS (SEQ ID NO: 216) and QQSYSTPLT (SEQ ID NO: 251).

**>3362\_B11 VH nucleotide sequence (SEQ ID NO: 171)**

CAGGTGCAGCTGCAGGCGTCGGGCCCAGGACTGGTGAAGCCTTCAGAGACCCTGTCCCTCAC  
CTGCACTGTCTCTGGTGAAGTCCATCACCAGTGGTGTCTTACTACTGGACCTGGATCCGCCAGC  
ACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAACACCTACTACAAC  
CCGTCCCTCAAGAGTCGAGTTACCATATCACTAGACACGTCTAAGAACCAGTTCTCCCTGAA  
GGTGAAGTCTGTGACTGCCGCGGACACGGCCGTATATTACTGTGCGCGAGCTGCTTCGACTT  
CAGTGCTAGGATACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3362\_B11 VH amino acid sequence (SEQ ID NO: 172)**

Kabat Bold, Chothia underlined

QVQLQASGPGLVKPSETLSLTCTVSGDSIT**SGAYYWT**WIRQHPGKGLEWIG**YIYYSGNTYYN**  
**PSLKS**RVTIISLDTSKNQFSLKVN SVTAADTAVYYCAR**AASTSVLGYGMDV**WGQGTITVTVSS

**>3362\_B11 VL nucleotide sequence (SEQ ID NO: 174)**

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCAGATATTTAAATTGGTATCAGCAGGAACCAGGGA  
AGGCCCTAAGCTCCTGGTCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATAAGCAGTCTTCAACCTGAAGATTT  
TGCAACTTACTACTGTCAACAGAGTTATAGTACCCCCCTCACCTTCGGCCAAGGGACACGAC  
TGGAGATTAAA

**>3362\_B11 VH amino acid sequence (SEQ ID NO: 175)**

Kabat Bold, Chothia underlined

DIQMTQSPSSLSASVGDRVITITC**RASQISIRYLN**WYQQEPGKAPKLLVY**AASSLQS**GVPSRF  
SGSGSGTDFTLTISSLQPEDFATYYC**QQSYSTPLT**FGQGRLEIK

[113] The 3242\_P05 antibody (also referred to herein as P05) includes antibody includes a heavy chain variable region (SEQ ID NO: 176) encoded by the nucleic acid sequence shown below in SEQ ID NO: 175, and a light chain variable region (SEQ ID NO: 178) encoded by the nucleic acid sequence shown in SEQ ID NO 177.

[114] The amino acids encompassing the CDRs as defined by Chothia et al., 1989 are underlined and those defined by Kabat et al., 1991 are highlighted in bold in the sequences below.

[115] The heavy chain CDRs of the P05 antibody have the following sequences per Kabat definition: VSDNYIN (SEQ ID NO: 254), VFYSADRTSYAD (SEQ ID NO: 256) and VQKSYYGMDV (SEQ ID NO: 244). The light chain CDRs of the P05 antibody have the following sequences per Kabat definition: RASQSIERYLN (SEQ ID NO: 238), GASSLQS (SEQ ID NO: 211) and QQTFSIPL (SEQ ID NO: 245).

[116] The heavy chain CDRs of the P05 antibody have the following sequences per Chothia definition: SGFSV (SEQ ID NO: 257), VFYSADRTS (SEQ ID NO: 246) and VQKSYYGMDV (SEQ ID NO: 244). The light chain CDRs of the P05 antibody have the following sequences per Chothia definition: The light chain CDRs of the P05 antibody have the following sequences per Kabat definition: RASQSIERYLN (SEQ ID NO: 238), GASSLQS (SEQ ID NO: 211) and QQTFSIPL (SEQ ID NO: 245).

**>3242\_P05 VH nucleotide sequence (SEQ ID NO: 175)**

GACATGCAGCTGGTGGAGTCTGGAGGAGGCTTGGTCCCGCCGGGGGGGTCCCTGAGACTCTC  
CTGCGCAGCCTCTGGGTTTTCCGTCAGTGACAACTACATAAACTGGGTCCGCCAGGCTCCAG  
GGAAGGGGCTGGACTGGGTCTCAGTCTTTTATAGTGCTGATAGAACATCCTACGCAGACTCC  
GTGAAGGGCCGATTACCGTCTCCAGCCACGATTCCAAGAACACAGTGTACCTTCAAATGAA  
CAGTCTGAGAGCTGAGGACACGGCCGTTTATTACTGTGCGAGAGTTCAGAAGTCCTATTACG  
GTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCGAGC

**>3242\_P05 VH amino acid sequence (SEQ ID NO: 176)**

Kabat Bold, Chothia underlined

DMQLVESGGGLVPPGGSLRLSCAASGFS**VSDNYIN**WVRQAPGKGLDWVS**VFYSADRTSYADS**  
**VKGRFTVSSHDSKNTVYLQMNSLR**AEDTAVYYCAR**VQKSYYGMDV**WGQGT

**>3242\_P05 VL nucleotide sequence (SEQ ID NO: 177)**

GGCATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCAT  
CACTTGCCGGGCAAGTCAGAGCATTAGCAGATATTTAAATTGGTATCTGCAGAAACCAGGGA  
AAGCCCCTAAGCTCCTGATCTCTGGTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTT  
AGTGGCACTGGGTCTGGGACAGAATCACTCTCACCATCAGCAGTTTGCAACCTGAAGATTT  
TGCAACTTACTACTGTCAACAGACTTTCAGTATCCCTCTTTTGGCCAGGGGACCAAGGTGG  
AGATCAAA

**>3242\_P05 VL amino acid sequence (SEQ ID NO: 178)**

Kabat Bold, Chothia underlined

GIQMTQSPSSLSASVGRVTITC**RASQSIERYLN**WYLQKPGKAPKLLIS**GASSLQS**GVPSRF  
SGTGS

[117] HuM2e antibodies of the invention also include antibodies that include a heavy chain variable amino acid sequence that is at least 90%, 92%, 95%, 97% 98%, 99% or more identical the amino acid sequence of SEQ ID NO: 44 or 49. and/or a light chain variable amino acid that is at least 90%, 92%, 95%, 97% 98%, 99% or more identical the amino acid sequence of SEQ ID NO: 46 or 52.

[118] Alternatively, the monoclonal antibody is an antibody that binds to the same epitope as 8I10, 21B15, 23K12, 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, or 3242\_P05.

[119] The heavy chain of a M2e antibody is derived from a germ line V (variable) gene such as, for example, the IgHV4 or the IgHV3 germline gene.

[120] The M2e antibodies of the invention include a variable heavy chain ( $V_H$ ) region encoded by a human IgHV4 or the IgHV3 germline gene sequence. An IgHV4 germline gene sequence is shown, *e.g.*, in Accession numbers L10088, M29812, M95114, X56360 and M95117. An IgHV3 germline gene sequence is shown, *e.g.*, in Accession numbers X92218, X70208, Z27504, M99679 and AB019437. The M2e antibodies of the invention include a  $V_H$  region that is encoded by a nucleic acid sequence that is at least 80% homologous to the IgHV4 or the IgHV3 germline gene sequence. Preferably, the nucleic acid sequence is at least 90%, 95%, 96%, 97% homologous to the IgHV4 or the IgHV3 germline gene sequence, and more preferably, at least 98%, 99% homologous to the IgHV4 or the IgHV3 germline gene sequence. The  $V_H$  region of the M2e antibody is at least 80% homologous to the amino acid sequence of the  $V_H$  region encoded by the IgHV4 or the IgHV3  $V_H$  germline gene sequence. Preferably, the amino acid sequence of  $V_H$  region of the M2e antibody is at least 90%, 95%, 96%, 97% homologous to the amino acid sequence encoded by the IgHV4 or the IgHV3 germline gene sequence, and more preferably, at least 98%, 99% homologous to the sequence encoded by the IgHV4 or the IgHV3 germline gene sequence.

[121] The M2e antibodies of the invention also include a variable light chain ( $V_L$ ) region encoded by a human IgKV1 germline gene sequence. A human IgKV1  $V_L$  germline gene sequence is shown, *e.g.*, Accession numbers X59315, X59312, X59318, J00248, and Y14865. Alternatively, the M2e antibodies include a  $V_L$  region that is encoded by a nucleic acid sequence that is at least 80% homologous to the IgKV1 germline gene sequence. Preferably, the nucleic acid sequence is at least 90%, 95%, 96%, 97% homologous to the IgKV1 germline gene sequence, and more preferably, at least 98%, 99% homologous to the IgKV1 germline gene sequence. The  $V_L$  region of the M2e antibody is at least 80%

homologous to the amino acid sequence of the V<sub>L</sub> region encoded the IgKV1 germline gene sequence. Preferably, the amino acid sequence of V<sub>L</sub> region of the M2c antibody is at least 90%, 95%, 96%, 97% homologous to the amino acid sequence encoded by the IgKV1 germline gene sequence, and more preferably, at least 98%, 99% homologous to the sequence encoded by the IgKV1 germline gene sequence.

[122] Unless otherwise defined, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures utilized in connection with, and techniques of, cell and tissue culture, molecular biology, and protein and oligo- or polynucleotide chemistry and hybridization described herein are those well known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (*e.g.*, electroporation, lipofection). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The practice of the present invention will employ, unless indicated specifically to the contrary, conventional methods of virology, immunology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, many of which are described below for the purpose of illustration. Such techniques are explained fully in the literature. See, *e.g.*, Sambrook, *et al.* Molecular Cloning: A Laboratory Manual (2nd Edition, 1989); Maniatis *et al.* Molecular Cloning: A Laboratory Manual (1982); DNA Cloning: A Practical Approach, vol. I & II (D. Glover, ed.); Oligonucleotide Synthesis (N. Gait, ed., 1984); Nucleic Acid Hybridization (B. Hames & S. Higgins, eds., 1985); Transcription and Translation (B. Hames & S. Higgins, eds., 1984); Animal Cell Culture (R. Freshney, ed., 1986); Perbal, A Practical Guide to Molecular Cloning (1984).

[123] The nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[124] The following definitions are useful in understanding the present invention:

[125] The term "antibody" (Ab) as used herein includes monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific antibodies), and antibody fragments, so

long as they exhibit the desired biological activity. The term “immunoglobulin” (Ig) is used interchangeably with “antibody” herein.

[126] An “isolated antibody” is one that has been separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody is purified: (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator; or (3) to homogeneity by SDS-PAGE under reducing or non-reducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[127] The basic four-chain antibody unit is a heterotetrameric glycoprotein composed of two identical light (L) chains and two identical heavy (H) chains. An IgM antibody consists of five of the basic heterotetramer units along with an additional polypeptide called a J chain, and therefore, contains ten antigen binding sites, while secreted IgA antibodies can polymerize to form polyvalent assemblages comprising 2-5 of the basic 4-chain units along with J chain. In the case of IgGs, the 4-chain unit is generally about 150,000 daltons. Each L chain is linked to an H chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. Each H and L chain also has regularly spaced intrachain disulfide bridges. Each H chain has at the N-terminus, a variable domain ( $V_H$ ) followed by three constant domains ( $C_H$ ) for each of the  $\alpha$  and  $\gamma$  chains and four  $C_H$  domains for  $\mu$  and  $\epsilon$  isotypes. Each L chain has at the N-terminus, a variable domain ( $V_L$ ) followed by a constant domain ( $C_L$ ) at its other end. The  $V_L$  is aligned with the  $V_H$  and the  $C_L$  is aligned with the first constant domain of the heavy chain ( $C_{H1}$ ). Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains. The pairing of a  $V_H$  and  $V_L$  together forms a single antigen-binding site. For the structure and properties of the different classes of antibodies, see, *e.g.*, Basic and Clinical Immunology, 8th edition, Daniel P. Stites, Abba I. Terr and Tristram G. Parslow (eds.), Appleton & Lange, Norwalk, Conn., 1994, page 71, and Chapter 6.

[128] The L chain from any vertebrate species can be assigned to one of two clearly distinct types, called kappa ( $\kappa$ ) and lambda ( $\lambda$ ), based on the amino acid sequences of their constant

domains ( $C_L$ ). Depending on the amino acid sequence of the constant domain of their heavy chains ( $C_H$ ), immunoglobulins can be assigned to different classes or isotypes. There are five classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, having heavy chains designated alpha ( $\alpha$ ), delta ( $\delta$ ), epsilon ( $\epsilon$ ), gamma ( $\gamma$ ) and mu ( $\mu$ ), respectively. The  $\gamma$  and  $\alpha$  classes are further divided into subclasses on the basis of relatively minor differences in  $C_H$  sequence and function, *e.g.*, humans express the following subclasses: IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2.

[129] The term “variable” refers to the fact that certain segments of the V domains differ extensively in sequence among antibodies. The V domain mediates antigen binding and defines specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the 110-amino acid span of the variable domains. Instead, the V regions consist of relatively invariant stretches called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called “hypervariable regions” that are each 9-12 amino acids long. The variable domains of native heavy and light chains each comprise four FRs, largely adopting a  $\beta$ -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the  $\beta$ -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC).

[130] The term “hypervariable region” when used herein refers to the amino acid residues of an antibody that are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a “complementarity determining region” or “CDR” (*e.g.*, around about residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the  $V_L$ , and around about 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the  $V_H$  when numbered in accordance with the Kabat numbering system; Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)); and/or those residues from a “hypervariable loop” (*e.g.*, residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the  $V_L$ , and 26-32 (H1), 52-56 (H2) and 95-101 (H3) in the  $V_H$  when numbered in accordance with the Chothia numbering system; Chothia and Lesk, J. Mol. Biol. 196:901-917 (1987)); and/or those residues from a “hypervariable loop”/CDR (*e.g.*, residues 27-38 (L1), 56-65 (L2)

and 105-120 (L3) in the V<sub>L</sub>, and 27-38 (H1), 56-65 (H2) and 105-120 (H3) in the V<sub>H</sub> when numbered in accordance with the IMGT numbering system; Lefranc, M.P. et al. Nucl. Acids Res. 27:209-212 (1999), Ruiz, M. et al. Nucl. Acids Res. 28:219-221 (2000)). Optionally the antibody has symmetrical insertions at one or more of the following points 28, 36 (L1), 63, 74-75 (L2) and 123 (L3) in the V<sub>L</sub>, and 28, 36 (H1), 63, 74-75 (H2) and 123 (H3) in the V<sub>H</sub> when numbered in accordance with AHO; Honneger, A. and Plunkthun, A. J. Mol. Biol. 309:657-670 (2001)).

**[131]** By “germline nucleic acid residue” is meant the nucleic acid residue that naturally occurs in a germline gene encoding a constant or variable region. “Germline gene” is the DNA found in a germ cell (*i.e.*, a cell destined to become an egg or in the sperm). A “germline mutation” refers to a heritable change in a particular DNA that has occurred in a germ cell or the zygote at the single-cell stage, and when transmitted to offspring, such a mutation is incorporated in every cell of the body. A germline mutation is in contrast to a somatic mutation which is acquired in a single body cell. In some cases, nucleotides in a germline DNA sequence encoding for a variable region are mutated (*i.e.*, a somatic mutation) and replaced with a different nucleotide.

**[132]** The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations that include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier “monoclonal” is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies useful in the present invention may be prepared by the hybridoma methodology first described by Kohler et al., Nature, 256:495 (1975), or may be made using recombinant DNA methods in bacterial, eukaryotic animal or plant cells (see, *e.g.*, U.S. Pat. No. 4,816,567). The “monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352:624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222:581-597 (1991), for example.

**[133]** The monoclonal antibodies herein include “chimeric” antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in

antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see U.S. Pat. No. 4,816,567; and Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81:6851-6855 (1984)). The present invention provides variable domain antigen-binding sequences derived from human antibodies. Accordingly, chimeric antibodies of primary interest herein include antibodies having one or more human antigen binding sequences (*e.g.*, CDRs) and containing one or more sequences derived from a non-human antibody, *e.g.*, an FR or C region sequence. In addition, chimeric antibodies of primary interest herein include those comprising a human variable domain antigen binding sequence of one antibody class or subclass and another sequence, *e.g.*, FR or C region sequence, derived from another antibody class or subclass. Chimeric antibodies of interest herein also include those containing variable domain antigen-binding sequences related to those described herein or derived from a different species, such as a non-human primate (*e.g.*, Old World Monkey, Ape, etc). Chimeric antibodies also include primatized and humanized antibodies.

[134] Furthermore, chimeric antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. For further details, see Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol. 2:593-596 (1992).

[135] A “humanized antibody” is generally considered to be a human antibody that has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as “import” residues, which are typically taken from an “import” variable domain. Humanization is traditionally performed following the method of Winter and co-workers (Jones *et al.*, Nature, 321:522-525 (1986); Reichmann *et al.*, Nature, 332:323-327 (1988); Verhoeyen *et al.*, Science, 239:1534-1536 (1988)), by substituting import hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such “humanized” antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species.

[136] A “human antibody” is an antibody containing only sequences present in an antibody naturally produced by a human. However, as used herein, human antibodies may comprise

residues or modifications not found in a naturally occurring human antibody, including those modifications and variant sequences described herein. These are typically made to further refine or enhance antibody performance.

[137] An “intact” antibody is one that comprises an antigen-binding site as well as a C<sub>L</sub> and at least heavy chain constant domains, C<sub>H</sub> 1, C<sub>H</sub> 2 and C<sub>H</sub> 3. The constant domains may be native sequence constant domains (*e.g.*, human native sequence constant domains) or amino acid sequence variant thereof. Preferably, the intact antibody has one or more effector functions.

[138] An “antibody fragment” comprises a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (see U.S. Pat. No. 5,641,870; Zapata *et al.*, Protein Eng. 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[139] The phrase “functional fragment or analog” of an antibody is a compound having qualitative biological activity in common with a full-length antibody. For example, a functional fragment or analog of an anti-IgE antibody is one that can bind to an IgE immunoglobulin in such a manner so as to prevent or substantially reduce the ability of such molecule from having the ability to bind to the high affinity receptor, Fc<sub>ε</sub>RI.

[140] Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, and a residual “Fc” fragment, a designation reflecting the ability to crystallize readily. The Fab fragment consists of an entire L chain along with the variable region domain of the H chain (V<sub>H</sub>), and the first constant domain of one heavy chain (C<sub>H</sub> 1). Each Fab fragment is monovalent with respect to antigen binding, *i.e.*, it has a single antigen-binding site. Pepsin treatment of an antibody yields a single large F(ab')<sub>2</sub> fragment that roughly corresponds to two disulfide linked Fab fragments having divalent antigen-binding activity and is still capable of cross-linking antigen. Fab' fragments differ from Fab fragments by having additional few residues at the carboxy terminus of the C<sub>H</sub>1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[141] The “Fc” fragment comprises the carboxy-terminal portions of both H chains held together by disulfides. The effector functions of antibodies are determined by sequences in

the Fc region, which region is also the part recognized by Fc receptors (FcR) found on certain types of cells.

[142] “Fv” is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (three loops each from the H and L chain) that contribute the amino acid residues for antigen binding and confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[143] “Single-chain Fv” also abbreviated as “sFv” or “scFv” are antibody fragments that comprise the V<sub>H</sub> and V<sub>L</sub> antibody domains connected into a single polypeptide chain. Preferably, the sFv polypeptide further comprises a polypeptide linker between the V<sub>H</sub> and V<sub>L</sub> domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994); Borrebaeck 1995, *infra*.

[144] The term “diabodies” refers to small antibody fragments prepared by constructing sFv fragments (see preceding paragraph) with short linkers (about 5-10 residues) between the V<sub>H</sub> and V<sub>L</sub> domains such that inter-chain but not intra-chain pairing of the V domains is achieved, resulting in a bivalent fragment, *i.e.*, fragment having two antigen-binding sites. Bispecific diabodies are heterodimers of two “crossover” sFv fragments in which the V<sub>H</sub> and V<sub>L</sub> domains of the two antibodies are present on different polypeptide chains. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

[145] As used herein, an antibody that “internalizes” is one that is taken up by (*i.e.*, enters) the cell upon binding to an antigen on a mammalian cell (*e.g.*, a cell surface polypeptide or receptor). The internalizing antibody will of course include antibody fragments, human or chimeric antibody, and antibody conjugates. For certain therapeutic applications, internalization *in vivo* is contemplated. The number of antibody molecules internalized will be sufficient or adequate to kill a cell or inhibit its growth, especially an infected cell. Depending on the potency of the antibody or antibody conjugate, in some instances, the uptake of a single antibody molecule into the cell is sufficient to kill the target cell to which the antibody binds. For example, certain toxins are highly potent in killing such that

internalization of one molecule of the toxin conjugated to the antibody is sufficient to kill the infected cell.

[146] As used herein, an antibody is said to be “immunospecific,” “specific for” or to “specifically bind” an antigen if it reacts at a detectable level with the antigen, preferably with an affinity constant,  $K_a$ , of greater than or equal to about  $10^4 \text{ M}^{-1}$ , or greater than or equal to about  $10^5 \text{ M}^{-1}$ , greater than or equal to about  $10^6 \text{ M}^{-1}$ , greater than or equal to about  $10^7 \text{ M}^{-1}$ , or greater than or equal to  $10^8 \text{ M}^{-1}$ . Affinity of an antibody for its cognate antigen is also commonly expressed as a dissociation constant  $K_D$ , and in certain embodiments, HuM2e antibody specifically binds to M2e if it binds with a  $K_D$  of less than or equal to  $10^{-4} \text{ M}$ , less than or equal to about  $10^{-5} \text{ M}$ , less than or equal to about  $10^{-6} \text{ M}$ , less than or equal to  $10^{-7} \text{ M}$ , or less than or equal to  $10^{-8} \text{ M}$ . Affinities of antibodies can be readily determined using conventional techniques, for example, those described by Scatchard *et al.* (*Ann. N.Y. Acad. Sci. USA* 51:660 (1949)).

[147] Binding properties of an antibody to antigens, cells or tissues thereof may generally be determined and assessed using immunodetection methods including, for example, immunofluorescence-based assays, such as immuno-histochemistry (IHC) and/or fluorescence-activated cell sorting (FACS).

[148] An antibody having a “biological characteristic” of a designated antibody is one that possesses one or more of the biological characteristics of that antibody which distinguish it from other antibodies. For example, in certain embodiments, an antibody with a biological characteristic of a designated antibody will bind the same epitope as that bound by the designated antibody and/or have a common effector function as the designated antibody.

[149] The term “antagonist” antibody is used in the broadest sense, and includes an antibody that partially or fully blocks, inhibits, or neutralizes a biological activity of an epitope, polypeptide, or cell that it specifically binds. Methods for identifying antagonist antibodies may comprise contacting a polypeptide or cell specifically bound by a candidate antagonist antibody with the candidate antagonist antibody and measuring a detectable change in one or more biological activities normally associated with the polypeptide or cell.

[150] An “antibody that inhibits the growth of infected cells” or a “growth inhibitory” antibody is one that binds to and results in measurable growth inhibition of infected cells expressing or capable of expressing an M2e epitope bound by an antibody. Preferred growth inhibitory antibodies inhibit growth of infected cells by greater than 20%, preferably from about 20% to about 50%, and even more preferably, by greater than 50% (*e.g.*, from about 50% to about 100%) as compared to the appropriate control, the control typically being

infected cells not treated with the antibody being tested. Growth inhibition can be measured at an antibody concentration of about 0.1 to 30  $\mu\text{g/ml}$  or about 0.5 nM to 200 nM in cell culture, where the growth inhibition is determined 1-10 days after exposure of the infected cells to the antibody. Growth inhibition of infected cells *in vivo* can be determined in various ways known in the art. The antibody is growth inhibitory *in vivo* if administration of the antibody at about 1  $\mu\text{g/kg}$  to about 100 mg/kg body weight results in reduction the percent of infected cells or total number of infected cells within about 5 days to 3 months from the first administration of the antibody, preferably within about 5 to 30 days.

[151] An antibody that “induces apoptosis” is one which induces programmed cell death as determined by binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies). Preferably the cell is an infected cell. Various methods are available for evaluating the cellular events associated with apoptosis. For example, phosphatidyl serine (PS) translocation can be measured by annexin binding; DNA fragmentation can be evaluated through DNA laddering; and nuclear/chromatin condensation along with DNA fragmentation can be evaluated by any increase in hypodiploid cells. Preferably, the antibody that induces apoptosis is one that results in about 2 to 50 fold, preferably about 5 to 50 fold, and most preferably about 10 to 50 fold, induction of annexin binding relative to untreated cell in an annexin binding assay.

[152] Antibody “effector functions” refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (*e.g.*, B cell receptor); and B cell activation.

[153] “Antibody-dependent cell-mediated cytotoxicity” or “ADCC” refers to a form of cytotoxicity in which secreted Ig bound to Fc receptors (FcRs) present on certain cytotoxic cells (*e.g.*, Natural Killer (NK) cells, neutrophils, and macrophages) enable these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. The antibodies “arm” the cytotoxic cells and are required for such killing. The primary cells for mediating ADCC, NK cells, express Fc $\gamma$ RIII only, whereas monocytes express Fc $\gamma$ RI, Fc $\gamma$ RII and Fc $\gamma$ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as

that described in U.S. Pat. No. 5,500,362 or U.S. Pat. No. 5,821,337 may be performed.

Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, *e.g.*, in a animal model such as that disclosed in Clynes *et al.*, PNAS (USA) 95:652-656 (1998).

[154] “Fc receptor” or “FcR” describes a receptor that binds to the Fc region of an antibody. In certain embodiments, the FcR is a native sequence human FcR. Moreover, a preferred FcR is one that binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors. FcγRII receptors include FcγRIIA (an “activating receptor”) and FcγRIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (*see* review M. in Daeron, Annu. Rev. Immunol. 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, Annu. Rev. Immunol 9:457-92 (1991); Capel *et al.*, Immunomethods 4:25-34 (1994); and de Haas *et al.*, J. Lab. Clin. Med. 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, J. Immunol. 117:587 (1976) and Kim *et al.*, J. Immunol. 24:249 (1994)).

[155] “Human effector cells” are leukocytes that express one or more FcRs and perform effector functions. Preferably, the cells express at least FcγRIII and perform ADCC effector function. Examples of human leukocytes that mediate ADCC include PBMC, NK cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source, *e.g.*, from blood.

[156] “Complement dependent cytotoxicity” or “CDC” refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system (C1q) to antibodies (of the appropriate subclass) that are bound to their cognate antigen. To assess complement activation, a CDC assay, *e.g.*, as described in Gazzano-Santoro *et al.*, J. Immunol. Methods 202:163 (1996), may be performed.

[157] The terms “influenza A” and “Influenzavirus A” refer to a genus of the Orthomyxoviridae family of viruses. Influenzavirus A includes only one species: influenza A

virus which causes influenza in birds, humans, pigs, and horses. Strains of all subtypes of influenza A virus have been isolated from wild birds, although disease is uncommon. Some isolates of influenza A virus cause severe disease both in domestic poultry and, rarely, in humans.

[158] A “mammal” for purposes of treating an infection, refers to any mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

[159] “Treating” or “treatment” or “alleviation” refers to both therapeutic treatment and prophylactic or preventative measures; wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. A subject or mammal is successfully “treated” for an infection if, after receiving a therapeutic amount of an antibody according to the methods of the present invention, the patient shows observable and/or measurable reduction in or absence of one or more of the following: reduction in the number of infected cells or absence of the infected cells; reduction in the percent of total cells that are infected; and/or relief to some extent, one or more of the symptoms associated with the specific infection; reduced morbidity and mortality, and improvement in quality of life issues. The above parameters for assessing successful treatment and improvement in the disease are readily measurable by routine procedures familiar to a physician.

[160] The term “therapeutically effective amount” refers to an amount of an antibody or a drug effective to “treat” a disease or disorder in a subject or mammal. See preceding definition of “treating.”

[161] “Chronic” administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. “Intermittent” administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

[162] Administration “in combination with” one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

[163] “Carriers” as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers that are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low

molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN<sup>TM</sup> polyethylene glycol (PEG), and PLURONICS<sup>TM</sup>.

[164] The term “cytotoxic agent” as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.*, At<sup>211</sup>, I<sup>131</sup>, I<sup>125</sup>, Y<sup>90</sup>, Re<sup>186</sup>, Re<sup>188</sup>, Sm<sup>153</sup>, Bi<sup>212</sup>, P<sup>32</sup> and radioactive isotopes of Lu), chemotherapeutic agents *e.g.*, methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents, enzymes and fragments thereof such as nucleolytic enzymes, antibiotics, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof, and the various antitumor or anticancer agents disclosed below. Other cytotoxic agents are described below.

[165] A “growth inhibitory agent” when used herein refers to a compound or composition which inhibits growth of a cell, either *in vitro* or *in vivo*. Examples of growth inhibitory agents include agents that block cell cycle progression, such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vinca alkaloids (vincristine, vinorelbine and vinblastine), taxanes, and topoisomerase II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled “Cell cycle regulation, oncogenes, and antineoplastic drugs” by Murakami *et al.* (W B Saunders: Philadelphia, 1995), especially p. 13. The taxanes (paclitaxel and docetaxel) are anticancer drugs both derived from the yew tree. Docetaxel (TAXOTERE<sup>TM</sup>, Rhone-Poulenc Rorer), derived from the European yew, is a semisynthetic analogue of paclitaxel (TAXOL®, Bristol-Myers Squibb). Paclitaxel and docetaxel promote the assembly of microtubules from tubulin dimers and stabilize microtubules by preventing depolymerization, which results in the inhibition of mitosis in cells.

[166] “Label” as used herein refers to a detectable compound or composition that is conjugated directly or indirectly to the antibody so as to generate a “labeled” antibody. The label may be detectable by itself (*e.g.*, radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable.

[167] The term “epitope tagged” as used herein refers to a chimeric polypeptide comprising a polypeptide fused to a “tag polypeptide.” The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide is also preferably fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

[168] A “small molecule” is defined herein to have a molecular weight below about 500 Daltons.

[169] The terms “nucleic acid” and “polynucleotide” are used interchangeably herein to refer to single- or double-stranded RNA, DNA, or mixed polymers. Polynucleotides may include genomic sequences, extra-genomic and plasmid sequences, and smaller engineered gene segments that express, or may be adapted to express polypeptides.

[170] An “isolated nucleic acid” is a nucleic acid that is substantially separated from other genome DNA sequences as well as proteins or complexes such as ribosomes and polymerases, which naturally accompany a native sequence. The term embraces a nucleic acid sequence that has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogues or analogues biologically synthesized by heterologous systems. A substantially pure nucleic acid includes isolated forms of the nucleic acid. Of course, this refers to the nucleic acid as originally isolated and does not exclude genes or sequences later added to the isolated nucleic acid by the hand of man.

[171] The term “polypeptide” is used in its conventional meaning, *i.e.*, as a sequence of amino acids. The polypeptides are not limited to a specific length of the product. Peptides, oligopeptides, and proteins are included within the definition of polypeptide, and such terms may be used interchangeably herein unless specifically indicated otherwise. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like, as well as other modifications

known in the art, both naturally occurring and non-naturally occurring. A polypeptide may be an entire protein, or a subsequence thereof. Particular polypeptides of interest in the context of this invention are amino acid subsequences comprising CDRs and being capable of binding an antigen or Influenza A-infected cell.

[172] An “isolated polypeptide” is one that has been identified and separated and/or recovered from a component of its natural environment. In preferred embodiments, the isolated polypeptide will be purified (1) to greater than 95% by weight of polypeptide as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or non-reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes the polypeptide *in situ* within recombinant cells since at least one component of the polypeptide's natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

[173] A “native sequence” polynucleotide is one that has the same nucleotide sequence as a polynucleotide derived from nature. A “native sequence” polypeptide is one that has the same amino acid sequence as a polypeptide (*e.g.*, antibody) derived from nature (*e.g.*, from any species). Such native sequence polynucleotides and polypeptides can be isolated from nature or can be produced by recombinant or synthetic means.

[174] A polynucleotide “variant,” as the term is used herein, is a polynucleotide that typically differs from a polynucleotide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the polynucleotide sequences of the invention and evaluating one or more biological activities of the encoded polypeptide as described herein and/or using any of a number of techniques well known in the art.

[175] A polypeptide “variant,” as the term is used herein, is a polypeptide that typically differs from a polypeptide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the above polypeptide sequences of the invention and evaluating one or more biological activities of the polypeptide as described herein and/or using any of a number of techniques well known in the art.

[176] Modifications may be made in the structure of the polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a variant or

derivative polypeptide with desirable characteristics. When it is desired to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, variant or portion of a polypeptide of the invention, one skilled in the art will typically change one or more of the codons of the encoding DNA sequence.

[177] For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of its ability to bind other polypeptides (*e.g.*, antigens) or cells. Since it is the binding capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences that encode said peptides without appreciable loss of their biological utility or activity.

[178] In many instances, a polypeptide variant will contain one or more conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged.

[179] In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

[180] It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within  $\pm 2$  is preferred, those

within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent 4,554,101 states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

[181] As detailed in U. S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate ( $+3.0 \pm 1$ ); glutamate ( $+3.0 \pm 1$ ); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine ( $-0.4$ ); proline ( $-0.5 \pm 1$ ); alanine ( $-0.5$ ); histidine ( $-0.5$ ); cysteine ( $-1.0$ ); methionine ( $-1.3$ ); valine ( $-1.5$ ); leucine ( $-1.8$ ); isoleucine ( $-1.8$ ); tyrosine ( $-2.3$ ); phenylalanine ( $-2.5$ ); tryptophan ( $-3.4$ ). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within  $\pm 2$  is preferred, those within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred.

[182] As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

[183] Amino acid substitutions may further be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by substitution, deletion or addition of five amino acids or fewer. Variants may also (or alternatively) be modified by, for example, the deletion

or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydrophobic nature of the polypeptide.

[184] Polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

[185] When comparing polynucleotide and polypeptide sequences, two sequences are said to be “identical” if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A “comparison window” as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

[186] Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) *Proc. Natl. Acad., Sci. USA* 80:726-730.

[187] Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the

search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

[188] One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information.

[189] In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

[190] For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment.

[191] In one approach, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is

calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residues occur in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

[192] “Homology” refers to the percentage of residues in the polynucleotide or polypeptide sequence variant that are identical to the non-variant sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology. In particular embodiments, polynucleotide and polypeptide variants have at least 70%, at least 75%, at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% polynucleotide or polypeptide homology with a polynucleotide or polypeptide described herein.

[193] “Vector” includes shuttle and expression vectors. Typically, the plasmid construct will also include an origin of replication (e.g., the ColE1 origin of replication) and a selectable marker (e.g., ampicillin or tetracycline resistance), for replication and selection, respectively, of the plasmids in bacteria. An “expression vector” refers to a vector that contains the necessary control sequences or regulatory elements for expression of the antibodies including antibody fragment of the invention, in bacterial or eukaryotic cells. Suitable vectors are disclosed below.

[194] As used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural references unless the content clearly dictates otherwise.

[195] The present invention includes HuM2e antibodies comprising a polypeptide of the present invention, including those polypeptides encoded by a polynucleotide sequence set forth in Example 1 and amino acid sequences set forth in Example 1 and 2, and fragments and variants thereof. In one embodiment, the antibody is an antibody designated herein as 8i10, 21B15, 23K12, 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, or 3242\_P05. These antibodies preferentially bind to or specifically bind to influenza A infected cells as compared to uninfected control cells of the same cell type.

[196] In particular embodiments, the antibodies of the present invention bind to the M2 protein. In certain embodiments, the present invention provides HuM2e antibodies that bind to epitopes within M2e that are only present in the native conformation, *i.e.*, as expressed in cells. In particular embodiments, these antibodies fail to specifically bind to an isolated M2e polypeptide, e.g., the 23 amino acid residue M2e fragment. It is understood that these antibodies recognize non-linear (*i.e.* conformational) epitope(s) of the M2 peptide.

[197] These specific conformational epitopes within the M2 protein, and particularly within M2e, may be used as vaccines to prevent the development of influenza infection within a subject.

[198] As will be understood by the skilled artisan, general description of antibodies herein and methods of preparing and using the same also apply to individual antibody polypeptide constituents and antibody fragments.

[199] The antibodies of the present invention may be polyclonal or monoclonal antibodies. However, in preferred embodiments, they are monoclonal. In particular embodiments, antibodies of the present invention are fully human antibodies. Methods of producing polyclonal and monoclonal antibodies are known in the art and described generally, *e.g.*, in U.S. Patent No. 6,824,780. Typically, the antibodies of the present invention are produced recombinantly, using vectors and methods available in the art, as described further below. Human antibodies may also be generated by *in vitro* activated B cells (see U.S. Pat. Nos. 5,567,610 and 5,229,275).

[200] Human antibodies may also be produced in transgenic animals (*e.g.*, mice) that are capable of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region ( $J_H$ ) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array into such germ-line mutant mice results in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits *et al.*, Proc. Natl. Acad. Sci. USA, 90:2551 (1993); Jakobovits *et al.*, Nature, 362:255-258 (1993); Bruggemann *et al.*, Year in Immuno., 7:33 (1993); U.S. Pat. Nos. 5,545,806, 5,569,825, 5,591,669 (all of GenPharm); U.S. Pat. No. 5,545,807; and WO 97/17852. Such animals may be genetically engineered to produce human antibodies comprising a polypeptide of the present invention.

[201] In certain embodiments, antibodies of the present invention are chimeric antibodies that comprise sequences derived from both human and non-human sources. In particular embodiments, these chimeric antibodies are humanized or primatized<sup>TM</sup>. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[202] In the context of the present invention, chimeric antibodies also include fully human antibodies wherein the human hypervariable region or one or more CDRs are retained, but

one or more other regions of sequence have been replaced by corresponding sequences from a non-human animal.

[203] The choice of non-human sequences, both light and heavy, to be used in making the chimeric antibodies is important to reduce antigenicity and human anti-non-human antibody responses when the antibody is intended for human therapeutic use. It is further important that chimeric antibodies retain high binding affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, chimeric antibodies are prepared by a process of analysis of the parental sequences and various conceptual chimeric products using three-dimensional models of the parental human and non-human sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

[204] As noted above, antibodies (or immunoglobulins) can be divided into five different classes, based on differences in the amino acid sequences in the constant region of the heavy chains. All immunoglobulins within a given class have very similar heavy chain constant regions. These differences can be detected by sequence studies or more commonly by serological means (*i.e.* by the use of antibodies directed to these differences). Antibodies, or fragments thereof, of the present invention may be any class, and may, therefore, have a gamma, mu, alpha, delta, or epsilon heavy chain. A gamma chain may be gamma 1, gamma 2, gamma 3, or gamma 4; and an alpha chain may be alpha 1 or alpha 2.

[205] In a preferred embodiment, an antibody of the present invention, or fragment thereof, is an IgG. IgG is considered the most versatile immunoglobulin, because it is capable of carrying out all of the functions of immunoglobulin molecules. IgG is the major Ig in serum, and the only class of Ig that crosses the placenta. IgG also fixes complement, although the IgG4 subclass does not. Macrophages, monocytes, PMN's and some lymphocytes have Fc receptors for the Fc region of IgG. Not all subclasses bind equally well: IgG2 and IgG4 do not bind to Fc receptors. A consequence of binding to the Fc receptors on PMN's, monocytes

and macrophages is that the cell can now internalize the antigen better. IgG is an opsonin that enhances phagocytosis. Binding of IgG to Fc receptors on other types of cells results in the activation of other functions. Antibodies of the present invention may be of any IgG subclass.

[206] In another preferred embodiment, an antibody, or fragment thereof, of the present invention is an IgE. IgE is the least common serum Ig since it binds very tightly to Fc receptors on basophils and mast cells even before interacting with antigen. As a consequence of its binding to basophils and mast cells, IgE is involved in allergic reactions. Binding of the allergen to the IgE on the cells results in the release of various pharmacological mediators that result in allergic symptoms. IgE also plays a role in parasitic helminth diseases. Eosinophils have Fc receptors for IgE and binding of eosinophils to IgE-coated helminths results in killing of the parasite. IgE does not fix complement.

[207] In various embodiments, antibodies of the present invention, and fragments thereof, comprise a variable light chain that is either kappa or lambda. The lambda chain may be any of subtype, including, *e.g.*, lambda 1, lambda 2, lambda 3, and lambda 4.

[208] As noted above, the present invention further provides antibody fragments comprising a polypeptide of the present invention. In certain circumstances there are advantages of using antibody fragments, rather than whole antibodies. For example, the smaller size of the fragments allows for rapid clearance, and may lead to improved access to certain tissues, such as solid tumors. Examples of antibody fragments include: Fab, Fab', F(ab')<sub>2</sub> and Fv fragments; diabodies; linear antibodies; single-chain antibodies; and multispecific antibodies formed from antibody fragments.

[209] Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (*see, e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992); and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. Fab, Fv and ScFv antibody fragments can all be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of these fragments. Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')<sub>2</sub> fragments (Carter *et al.*, *Bio/Technology* 10:163-167 (1992)). According to another approach, F(ab')<sub>2</sub> fragments can be isolated directly from recombinant host cell culture. Fab and F(ab')<sub>2</sub> fragment with increased *in vivo* half-life comprising a salvage receptor binding epitope residues are described in U.S. Pat. No. 5,869,046. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner.

[210] In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; U.S. Pat. Nos. 5,571,894; and 5,587,458. Fv and sFv are the only species with intact combining sites that are devoid of constant regions. Thus, they are suitable for reduced nonspecific binding during *in vivo* use. sFv fusion proteins may be constructed to yield fusion of an effector protein at either the amino or the carboxy terminus of an sFv. See Antibody Engineering, ed. Borrebaeck, *supra*. The antibody fragment may also be a “linear antibody”, *e.g.*, as described in U.S. Pat. No. 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

[211] In certain embodiments, antibodies of the present invention are bispecific or multi-specific. Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of a single antigen. Other such antibodies may combine a first antigen binding site with a binding site for a second antigen. Alternatively, an anti-M2e arm may be combined with an arm that binds to a triggering molecule on a leukocyte, such as a T-cell receptor molecule (*e.g.*, CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16), so as to focus and localize cellular defense mechanisms to the infected cell. Bispecific antibodies may also be used to localize cytotoxic agents to infected cells. These antibodies possess an M2e-binding arm and an arm that binds the cytotoxic agent (*e.g.*, saporin, anti-interferon- $\alpha$ , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments (*e.g.*, F(ab')<sub>2</sub> bispecific antibodies). WO 96/16673 describes a bispecific anti-ErbB2/anti-FcγRIII antibody and U.S. Pat. No. 5,837,234 discloses a bispecific anti-ErbB2/anti-FcγRI antibody. A bispecific anti-ErbB2/Fc $\alpha$  antibody is shown in WO98/02463. U.S. Pat. No. 5,821,337 teaches a bispecific anti-ErbB2/anti-CD3 antibody.

[212] Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, Nature, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, EMBO J., 10:3655-3659 (1991).

[213] According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. Preferably, the fusion is with an Ig heavy chain constant domain, comprising at least part of the hinge, C<sub>H</sub>2, and C<sub>H</sub>3 regions. It is preferred to have the first heavy-chain constant region (C<sub>H</sub>1) containing the site necessary for light chain bonding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host cell. This provides for greater flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yield of the desired bispecific antibody. It is, however, possible to insert the coding sequences for two or all three polypeptide chains into a single expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios have no significant affect on the yield of the desired chain combination.

[214] In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

[215] According to another approach described in U.S. Pat. No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers that are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C<sub>H</sub> 3 domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (*e.g.*, tyrosine or tryptophan). Compensatory “cavities” of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.*, alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[216] Bispecific antibodies include cross-linked or “heteroconjugate” antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Pat. No. 4,676,980, along with a number of cross-linking techniques.

[217] Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, Science, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')<sub>2</sub> fragments. These fragments are reduced in the presence of the dithiol complexing agent, sodium arsenite, to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[218] Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, J. Exp. Med., 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')<sub>2</sub> molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[219] Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, J. Immunol., 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers.

The “diabody” technology described by Hollinger *et al.*, Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a  $V_H$  connected to a  $V_L$  by a linker that is too short to allow pairing between the two domains on the same chain. Accordingly, the  $V_H$  and  $V_L$  domains of one fragment are forced to pair with the complementary  $V_L$  and  $V_H$  domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, J. Immunol., 152:5368 (1994).

[220] Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.*, J. Immunol. 147: 60 (1991). A multivalent antibody may be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The antibodies of the present invention can be multivalent antibodies with three or more antigen binding sites (*e.g.*, tetravalent antibodies), which can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody. The multivalent antibody can comprise a dimerization domain and three or more antigen binding sites. The preferred dimerization domain comprises (or consists of) an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fc region. The preferred multivalent antibody herein comprises (or consists of) three to about eight, but preferably four, antigen binding sites. The multivalent antibody comprises at least one polypeptide chain (and preferably two polypeptide chains), wherein the polypeptide chain(s) comprise two or more variable domains. For instance, the polypeptide chain(s) may comprise  $VD1-(X1)_n-VD2-(X2)_n-Fc$ , wherein  $VD1$  is a first variable domain,  $VD2$  is a second variable domain,  $Fc$  is one polypeptide chain of an Fc region,  $X1$  and  $X2$  represent an amino acid or polypeptide, and  $n$  is 0 or 1. For instance, the polypeptide chain(s) may comprise:  $VH-CH1-flexible\ linker-VH-CH1-Fc$  region chain; or  $VH-CH1-VH-CH1-Fc$  region chain. The multivalent antibody herein preferably further comprises at least two (and preferably four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a  $C_L$  domain.

[221] Antibodies of the present invention further include single chain antibodies.

[222] In particular embodiments, antibodies of the present invention are internalizing antibodies.

[223] Amino acid sequence modification(s) of the antibodies described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the antibody may be prepared by introducing appropriate nucleotide changes into a polynucleotide that encodes the antibody, or a chain thereof, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution may be made to arrive at the final antibody, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antibody, such as changing the number or position of glycosylation sites. Any of the variations and modifications described above for polypeptides of the present invention may be included in antibodies of the present invention.

[224] A useful method for identification of certain residues or regions of an antibody that are preferred locations for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells in *Science*, 244:1081-1085 (1989). Here, a residue or group of target residues are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with PSCA antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed anti- antibody variants are screened for the desired activity.

[225] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue or the antibody fused to a cytotoxic polypeptide. Other insertional variants of an antibody include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.*, for ADEPT) or a polypeptide that increases the serum half-life of the antibody.

[226] Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antibody molecule replaced by a different residue. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but

FR alterations are also contemplated. Conservative and non-conservative substitutions are contemplated.

[227] Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

[228] Any cysteine residue not involved in maintaining the proper conformation of the antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

[229] One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody. Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (*e.g.*, 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (*e.g.*, binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and an antigen or infected cell. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

[230] Another type of amino acid variant of the antibody alters the original glycosylation pattern of the antibody. By altering is meant deleting one or more carbohydrate moieties found in the antibody, and/or adding one or more glycosylation sites that are not present in the antibody.

[231] Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-aceylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[232] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[233] The antibody of the invention is modified with respect to effector function, *e.g.*, so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antibody. This may be achieved by introducing one or more amino acid substitutions in an Fc region of the antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, J. Exp Med. 176:1191-1195 (1992) and Shopes, B. J. Immunol. 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-infection activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.*, Cancer Research 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.*, Anti-Cancer Drug Design 3:219-230 (1989).

[234] To increase the serum half-life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Pat. No. 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub>, or IgG<sub>4</sub>) that is responsible for increasing the in vivo serum half-life of the IgG molecule.

[235] Antibodies of the present invention may also be modified to include an epitope tag or label, *e.g.*, for use in purification or diagnostic applications. The invention also pertains to therapy with immunoconjugates comprising an antibody conjugated to an anti-cancer agent such as a cytotoxic agent or a growth inhibitory agent. Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above.

[236] Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, maytansinoids, a trichothene, and CC1065, and the derivatives of these toxins that have toxin activity, are also contemplated herein.

[237] In one preferred embodiment, an antibody (full length or fragments) of the invention is conjugated to one or more maytansinoid molecules. Maytansinoids are mitototic inhibitors that act by inhibiting tubulin polymerization. Maytansine was first isolated from the east African shrub *Maytenus serrata* (U.S. Pat. No. 3,896,111). Subsequently, it was discovered that certain microbes also produce maytansinoids, such as maytansinol and C-3 maytansinol esters (U.S. Pat. No. 4,151,042). Synthetic maytansinol and derivatives and analogues thereof are disclosed, for example, in U.S. Pat. Nos. 4,137,230; 4,248,870; 4,256,746; 4,260,608; 4,265,814; 4,294,757; 4,307,016; 4,308,268; 4,308,269; 4,309,428; 4,313,946; 4,315,929; 4,317,821; 4,322,348; 4,331,598; 4,361,650; 4,364,866; 4,424,219; 4,450,254; 4,362,663; and 4,371,533.

[238] In an attempt to improve their therapeutic index, maytansine and maytansinoids have been conjugated to antibodies specifically binding to tumor cell antigens. Immunoconjugates containing maytansinoids and their therapeutic use are disclosed, for example, in U.S. Pat. Nos. 5,208,020, 5,416,064 and European Patent EP 0 425 235 B1. Liu *et al.*, Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996) described immunoconjugates comprising a maytansinoid designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic towards cultured colon cancer cells, and showed antitumor activity in an *in vivo* tumor growth assay.

[239] Antibody-maytansinoid conjugates are prepared by chemically linking an antibody to a maytansinoid molecule without significantly diminishing the biological activity of either the antibody or the maytansinoid molecule. An average of 3-4 maytansinoid molecules conjugated per antibody molecule has shown efficacy in enhancing cytotoxicity of target cells without negatively affecting the function or solubility of the antibody, although even one molecule of toxin/antibody would be expected to enhance cytotoxicity over the use of naked antibody. Maytansinoids are well known in the art and can be synthesized by known techniques or isolated from natural sources. Suitable maytansinoids are disclosed, for

example, in U.S. Pat. No. 5,208,020 and in the other patents and nonpatent publications referred to hereinabove. Preferred maytansinoids are maytansinol and maytansinol analogues modified in the aromatic ring or at other positions of the maytansinol molecule, such as various maytansinol esters.

[240] There are many linking groups known in the art for making antibody conjugates, including, for example, those disclosed in U.S. Pat. No. 5,208,020 or EP Patent 0 425 235 B1, and Chari *et al.*, Cancer Research 52: 127-131 (1992). The linking groups include disulfide groups, thioether groups, acid labile groups, photolabile groups, peptidase labile groups, or esterase labile groups, as disclosed in the above-identified patents, disulfide and thioether groups being preferred.

[241] Immunoconjugates may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl)hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). Particularly preferred coupling agents include N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP) (Carlsson *et al.*, Biochem. J. 173:723-737 [1978]) and N-succinimidyl-4-(2-pyridylthio) pentanoate (SPP) to provide for a disulfide linkage. For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026. The linker may be a "cleavable linker" facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker, Cancer Research 52: 127-131 (1992); U.S. Pat. No. 5,208,020) may be used.

[242] Another immunoconjugate of interest comprises an antibody conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics is capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. Pat. Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, 5,877,296 (all to American Cyanamid Company). Another drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily

cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

[243] Examples of other agents that can be conjugated to the antibodies of the invention include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. Pat. Nos. 5,053,394, 5,770,710, as well as esperamicins (U.S. Pat. No. 5,877,296).

[244] Enzymatically active toxins and fragments thereof that can be used include, *e.g.*, diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232.

[245] The present invention further includes an immunoconjugate formed between an antibody and a compound with nucleolytic activity (*e.g.*, a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

[246] For selective destruction of infected cells, the antibody includes a highly radioactive atom. A variety of radioactive isotopes are available for the production of radioconjugated anti-PSCA antibodies. Examples include At<sup>211</sup>, I<sup>131</sup>, I<sup>125</sup>, Y<sup>90</sup>, Re<sup>186</sup>, Re<sup>188</sup>, Sm<sup>153</sup>, Bi<sup>212</sup>, P<sup>32</sup>, Pb<sup>212</sup> and radioactive isotopes of Lu. When the conjugate is used for diagnosis, it may comprise a radioactive atom for scintigraphic studies, for example tc<sup>99m</sup> or I<sup>123</sup>, or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, MRI), such as iodine-123, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

[247] The radio- or other label is incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as tc<sup>99m</sup> or I<sup>123</sup>, Re<sup>186</sup>, Re<sup>188</sup> and In<sup>111</sup> can be attached via a cysteine residue in the peptide. Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker *et al.* (1978) Biochem. Biophys. Res. Commun. 80: 49-57) can be used to incorporate iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.

[248] Alternatively, a fusion protein comprising the antibody and cytotoxic agent is made, *e.g.*, by recombinant techniques or peptide synthesis. The length of DNA may comprise

respective regions encoding the two portions of the conjugate either adjacent one another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

[249] The antibodies of the present invention are also used in antibody dependent enzyme mediated prodrug therapy (ADET) by conjugating the antibody to a prodrug-activating enzyme which converts a prodrug (*e.g.*, a peptidyl chemotherapeutic agent, see WO81/01145) to an active anti-cancer drug (*see, e.g.*, WO 88/07378 and U.S. Pat. No. 4,975,278).

[250] The enzyme component of the immunoconjugate useful for ADEPT includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form. Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratia protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as  $\beta$ -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs;  $\beta$ -lactamase useful for converting drugs derivatized with  $\beta$ -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as “abzymes”, can be used to convert the prodrugs of the invention into free active drugs (*see, e.g.*, Massey, *Nature* 328: 457-458 (1987)). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a infected cell population.

[251] The enzymes of this invention can be covalently bound to the antibodies by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using recombinant DNA techniques well known in the art (*see, e.g.*, Neuberger *et al.*, *Nature*, 312: 604-608 (1984)).

[252] Other modifications of the antibody are contemplated herein. For example, the antibody may be linked to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene

glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol. The antibody also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate)microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980).

[253] The antibodies disclosed herein are also formulated as immunoliposomes. A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant that is useful for delivery of a drug to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82:3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77:4030 (1980); U.S. Pat. Nos. 4,485,045 and 4,544,545; and WO97/38731 published Oct. 23, 1997. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556.

[254] Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem. 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon *et al.*, J. National Cancer Inst. 81(19)1484 (1989).

[255] Antibodies of the present invention, or fragments thereof, may possess any of a variety of biological or functional characteristics. In certain embodiments, these antibodies are Influenza A specific or M2 protein specific antibodies, indicating that they specifically bind to or preferentially bind to Influenza A or the M2 protein thereof, respectively, as compared to a normal control cell. In particular embodiments, the antibodies are HuM2e antibodies, indicating that they specifically bind to a M2e protein, preferably to an epitope of the M2e domain that is only present when the M2 protein is expressed in cells or present on a virus, as compared to a normal control cell.

[256] In particular embodiments, an antibody of the present invention is an antagonist antibody, which partially or fully blocks or inhibits a biological activity of a polypeptide or cell to which it specifically or preferentially binds. In other embodiments, an antibody of the present invention is a growth inhibitory antibody, which partially or fully blocks or inhibits the growth of an infected cell to which it binds. In another embodiment, an antibody of the present invention induces apoptosis. In yet another embodiment, an antibody of the present invention induces or promotes antibody-dependent cell-mediated cytotoxicity or complement dependent cytotoxicity.

#### Methods of Identifying and Producing Antibodies Specific for Influenza Virus

[257] The present invention provides novel methods for the identification of HuM2e antibodies, as exemplified in Example 4. These methods may be readily adapted to identify antibodies specific for other polypeptides expressed on the cell surface by infectious agents, or even polypeptides expressed on the surface of an infectious agent itself.

[258] In general, the methods include obtaining serum samples from patients that have been infected with or vaccinated against an infectious agent. These serum samples are then screened to identify those that contain antibodies specific for a particular polypeptide associated with the infectious agent, such as, e.g., a polypeptide specifically expressed on the surface of cells infected with the infectious agent, but not uninfected cells. In particular embodiments, the serum samples are screened by contacting the samples with a cell that has been transfected with an expression vector that expresses the polypeptide expressed on the surface of infected cells.

[259] Once a patient is identified as having serum containing an antibody specific for the infectious agent polypeptide of interest is identified, mononuclear and/or B cells obtained from the same patient are used to identify a cell or clone thereof that produces the antibody, using any of the methods described herein or available in the art. Once a B cell that produces the antibody is identified, cDNAs encoding the variable regions or fragments thereof of the antibody may be cloned using standard RT-PCR vectors and primers specific for conserved antibody sequences, and subcloned in to expression vectors used for the recombinant production of monoclonal antibodies specific for the infectious agent polypeptide of interest.

[260] In one embodiment, the present invention provides a method of identifying an antibody that specifically binds influenza A-infected cells, comprising: contacting an Influenza A virus or a cell expressing the M2 protein with a biological sample obtained from a patient having been infected by Influenza A; determining an amount of antibody in the

biological sample that binds to the cell; and comparing the amount determined with a control value, wherein if the value determined is at least two-fold greater than the control value, an antibody that specifically binds influenza A-infected cells is indicated.

In various embodiments, the cells expressing an M2 protein are cells infected with an Influenza A virus or cells that have been transfected with a polynucleotide that expressed the M2 protein. Alternatively, the cells may express a portion of the M2 protein that includes the M2e domain and enough additional M2 sequence that the protein remains associated with the cell and the M2e domain is presented on the cell surface in the same manner as when present within full length M2 protein. Methods of preparing an M2 expression vector and transfecting an appropriate cell, including those described herein, may be readily accomplished, in view of the M2 sequence being publicly available. See, for example, the Influenza Sequence Database (ISD) ([flu.lanl.gov](http://flu.lanl.gov) on the World Wide Web, described in Macken et al., 2001, "The value of a database in surveillance and vaccine selection" in *Options for the Control of Influenza IV*. A.D.M.E., Osterhaus & Hampson (Eds.), Elsevier Science, Amsterdam, pp. 103-106) and the Microbial Sequencing Center (MSC) at The Institute for Genomic Research (TIGR) ([tigr.org/msc/infl\\_a\\_virus.shtml](http://tigr.org/msc/infl_a_virus.shtml) on the World Wide Web).

[261] The M2e-expressing cells or virus described above are used to screen the biological sample obtained from a patient infected with influenza A for the presence of antibodies that preferentially bind to the cell expressing the M2 polypeptide using standard biological techniques. For example, in certain embodiments, the antibodies may be labeled, and the presence of label associated with the cell detected, e.g., using FMAT or FACs analysis. In particular embodiments, the biological sample is blood, serum, plasma, bronchial lavage, or saliva. Methods of the present invention may be practiced using high throughput techniques.

[262] Identified human antibodies may then be characterized further. For example the particular conformational epitopes within the M2e protein that are necessary or sufficient for binding of the antibody may be determined, e.g., using site-directed mutagenesis of expressed M2e polypeptides. These methods may be readily adapted to identify human antibodies that bind any protein expressed on a cell surface. Furthermore, these methods may be adapted to determine binding of the antibody to the virus itself, as opposed to a cell expressing recombinant M2e or infected with the virus.

[263] Polynucleotide sequences encoding the antibodies, variable regions thereof, or antigen-binding fragments thereof may be subcloned into expression vectors for the recombinant production of HuM2e antibodies. In one embodiment, this is accomplished by

obtaining mononuclear cells from the patient from the serum containing the identified HuM2e antibody was obtained; producing B cell clones from the mononuclear cells; inducing the B cells to become antibody-producing plasma cells; and screening the supernatants produced by the plasma cells to determine if it contains the HuM2e antibody. Once a B cell clone that produces an HuM2e antibody is identified, reverse-transcription polymerase chain reaction (RT-PCR) is performed to clone the DNAs encoding the variable regions or portions thereof of the HuM2e antibody. These sequences are then subcloned into expression vectors suitable for the recombinant production of human HuM2e antibodies. The binding specificity may be confirmed by determining the recombinant antibody's ability to bind cells expressing M2e polypeptide.

[264] In particular embodiments of the methods described herein, B cells isolated from peripheral blood or lymph nodes are sorted, *e.g.*, based on their being CD19 positive, and plated, *e.g.*, as low as a single cell specificity per well, *e.g.*, in 96, 384, or 1536 well configurations. The cells are induced to differentiate into antibody-producing cells, *e.g.*, plasma cells, and the culture supernatants are harvested and tested for binding to cells expressing the infectious agent polypeptide on their surface using, *e.g.*, FMAT or FACS analysis. Positive wells are then subjected to whole well RT-PCR to amplify heavy and light chain variable regions of the IgG molecule expressed by the clonal daughter plasma cells. The resulting PCR products encoding the heavy and light chain variable regions, or portions thereof, are subcloned into human antibody expression vectors for recombinant expression. The resulting recombinant antibodies are then tested to confirm their original binding specificity and may be further tested for pan-specificity across various strains of isolates of the infectious agent.

[265] Thus, in one embodiment, a method of identifying HuM2e antibodies is practiced as follows. First, full length or approximately full length M2 cDNAs are transfected into a cell line for expression of M2 protein. Secondly, individual human plasma or sera samples are tested for antibodies that bind the cell-expressed M2. And lastly, MAbs derived from plasma- or serum-positive individuals are characterized for binding to the same cell-expressed M2. Further definition of the fine specificities of the MAbs can be performed at this point.

[266] These methods may be practiced to identify a variety of different HuM2e antibodies, including antibodies specific for (a) epitopes in a linear M2e peptide, (b) common epitopes in multiple variants of M2e, (c) conformational determinants of an M2 homotetramer, and (d) common conformational determinants of multiple variants of the M2 homotetramer. The last

category is particularly desirable, as this specificity is perhaps specific for all A strains of influenza.

[267] Polynucleotides that encode the HuM2e antibodies or portions thereof of the present invention may be isolated from cells expressing HuM2e antibodies, according to methods available in the art and described herein, including amplification by polymerase chain reaction using primers specific for conserved regions of human antibody polypeptides. For example, light chain and heavy chain variable regions may be cloned from the B cell according to molecular biology techniques described in WO 92/02551; U.S. Patent No. 5,627,052; or Babcook et al., *Proc. Natl. Acad. Sci. USA* 93:7843-48 (1996). In certain embodiments, polynucleotides encoding all or a region of both the heavy and light chain variable regions of the IgG molecule expressed by the clonal daughter plasma cells expressing the HuM2e antibody are subcloned and sequenced. The sequence of the encoded polypeptide may be readily determined from the polynucleotide sequence.

Isolated polynucleotides encoding a polypeptide of the present invention may be subcloned into an expression vector to recombinantly produce antibodies and polypeptides of the present invention, using procedures known in the art and described herein.

[268] Binding properties of an antibody (or fragment thereof) to M2e or infected cells or tissues may generally be determined and assessed using immunodetection methods including, for example, immunofluorescence-based assays, such as immuno-histochemistry (IHC) and/or fluorescence-activated cell sorting (FACS). Immunoassay methods may include controls and procedures to determine whether antibodies bind specifically to M2e from one or more specific strains of Influenza A, and do not recognize or cross-react with normal control cells.

[269] Following pre-screening of serum to identify patients that produce antibodies to an infectious agent or polypeptide thereof, e.g., M2, the methods of the present invention typically include the isolation or purification of B cells from a biological sample previously obtained from a patient or subject. The patient or subject may be currently or previously diagnosed with or suspect or having a particular disease or infection, or the patient or subject may be considered free of a particular disease or infection. Typically, the patient or subject is a mammal and, in particular embodiments, a human. The biological sample may be any sample that contains B cells, including but not limited to, lymph node or lymph node tissue, pleural effusions, peripheral blood, ascites, tumor tissue, or cerebrospinal fluid (CSF). In various embodiments, B cells are isolated from different types of biological samples, such as a biological sample affected by a particular disease or infection. However, it is understood

that any biological sample comprising B cells may be used for any of the embodiments of the present invention.

[270] Once isolated, the B cells are induced to produce antibodies, *e.g.*, by culturing the B cells under conditions that support B cell proliferation or development into a plasmacyte, plasmablast, or plasma cell. The antibodies are then screened, typically using high throughput techniques, to identify an antibody that specifically binds to a target antigen, *e.g.*, a particular tissue, cell, infectious agent, or polypeptide. In certain embodiments, the specific antigen, *e.g.*, cell surface polypeptide bound by the antibody is not known, while in other embodiments, the antigen specifically bound by the antibody is known.

[271] According to the present invention, B cells may be isolated from a biological sample, *e.g.*, a tumor, tissue, peripheral blood or lymph node sample, by any means known and available in the art. B cells are typically sorted by FACS based on the presence on their surface of a B cell-specific marker, *e.g.*, CD19, CD138, and/or surface IgG. However, other methods known in the art may be employed, such as, *e.g.*, column purification using CD19 magnetic beads or IgG-specific magnetic beads, followed by elution from the column. However, magnetic isolation of B cells utilizing any marker may result in loss of certain B cells. Therefore, in certain embodiments, the isolated cells are not sorted but, instead, phicol-purified mononuclear cells isolated from tumor are directly plated to the appropriate or desired number of specificities per well.

[272] In order to identify B cells that produce an infectious agent-specific antibody, the B cells are typically plated at low density (*e.g.*, a single cell specificity per well, 1-10 cells per well, 10-100 cells per well, 1-100 cells per well, less than 10 cells per well, or less than 100 cells per well) in multi-well or microtitre plates, *e.g.*, in 96, 384, or 1536 well configurations. When the B cells are initially plated at a density greater than one cell per well, then the methods of the present invention may include the step of subsequently diluting cells in a well identified as producing an antigen-specific antibody, until a single cell specificity per well is achieved, thereby facilitating the identification of the B cell that produces the antigen-specific antibody. Cell supernatants or a portion thereof and/or cells may be frozen and stored for future testing and later recovery of antibody polynucleotides.

[273] In certain embodiments, the B cells are cultured under conditions that favor the production of antibodies by the B cells. For example, the B cells may be cultured under conditions favorable for B cell proliferation and differentiation to yield antibody-producing plasmablast, plasmacytes, or plasma cells. In particular embodiments, the B cells are cultured in the presence of a B cell mitogen, such as lipopolysaccharide (LPS) or CD40

ligand. In one specific embodiment, B cells are differentiated to antibody-producing cells by culturing them with feed cells and/or other B cell activators, such as CD40 ligand.

[274] Cell culture supernatants or antibodies obtained therefrom may be tested for their ability to bind to a target antigen, using routine methods available in the art, including those described herein. In particular embodiments, culture supernatants are tested for the presence of antibodies that bind to a target antigen using high-throughput methods. For example, B cells may be cultured in multi-well microtitre dishes, such that robotic plate handlers may be used to simultaneously sample multiple cell supernatants and test for the presence of antibodies that bind to a target antigen. In particular embodiments, antigens are bound to beads, *e.g.*, paramagnetic or latex beads) to facilitate the capture of antibody/antigen complexes. In other embodiments, antigens and antibodies are fluorescently labeled (with different labels) and FACS analysis is performed to identify the presence of antibodies that bind to target antigen. In one embodiment, antibody binding is determined using FMAT™ analysis and instrumentation (Applied Biosystems, Foster City, CA). FMAT™ is a fluorescence macro-confocal platform for high-throughput screening, which mix-and-read, non-radioactive assays using live cells or beads.

[275] In the context of comparing the binding of an antibody to a particular target antigen (*e.g.*, a biological sample such as infected tissue or cells, or infectious agents) as compared to a control sample (*e.g.*, a biological sample such as uninfected cells, or a different infectious agent), in various embodiments, the antibody is considered to preferentially bind a particular target antigen if at least two-fold, at least three-fold, at least five-fold, or at least ten-fold more antibody binds to the particular target antigen as compared to the amount that binds a control sample.

[276] Polynucleotides encoding antibody chains, variable regions thereof, or fragments thereof, may be isolated from cells utilizing any means available in the art. In one embodiment, polynucleotides are isolated using polymerase chain reaction (PCR), *e.g.*, reverse transcription-PCR (RT-PCR) using oligonucleotide primers that specifically bind to heavy or light chain encoding polynucleotide sequences or complements thereof using routine procedures available in the art. In one embodiment, positive wells are subjected to whole well RT-PCR to amplify the heavy and light chain variable regions of the IgG molecule expressed by the clonal daughter plasma cells. These PCR products may be sequenced.

[277] The resulting PCR products encoding the heavy and light chain variable regions or portions thereof are then subcloned into human antibody expression vectors and

recombinantly expressed according to routine procedures in the art (*see, e.g.*, US Patent No. 7,112,439). The nucleic acid molecules encoding a tumor-specific antibody or fragment thereof, as described herein, may be propagated and expressed according to any of a variety of well-known procedures for nucleic acid excision, ligation, transformation, and transfection. Thus, in certain embodiments expression of an antibody fragment may be preferred in a prokaryotic host cell, such as *Escherichia coli* (*see, e.g.*, Pluckthun et al., *Methods Enzymol.* 178:497-515 (1989)). In certain other embodiments, expression of the antibody or an antigen-binding fragment thereof may be preferred in a eukaryotic host cell, including yeast (*e.g.*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Pichia pastoris*); animal cells (including mammalian cells); or plant cells. Examples of suitable animal cells include, but are not limited to, myeloma, COS, CHO, or hybridoma cells. Examples of plant cells include tobacco, corn, soybean, and rice cells. By methods known to those having ordinary skill in the art and based on the present disclosure, a nucleic acid vector may be designed for expressing foreign sequences in a particular host system, and then polynucleotide sequences encoding the tumor-specific antibody (or fragment thereof) may be inserted. The regulatory elements will vary according to the particular host.

[278] One or more replicable expression vectors containing a polynucleotide encoding a variable and/or constant region may be prepared and used to transform an appropriate cell line, for example, a non-producing myeloma cell line, such as a mouse NSO line or a bacteria, such as *E.coli*, in which production of the antibody will occur. In order to obtain efficient transcription and translation, the polynucleotide sequence in each vector should include appropriate regulatory sequences, particularly a promoter and leader sequence operatively linked to the variable domain sequence. Particular methods for producing antibodies in this way are generally well known and routinely used. For example, molecular biology procedures are described by Sambrook *et al.* (*Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, New York, 1989; *see also* Sambrook et al., 3rd ed., Cold Spring Harbor Laboratory, New York, (2001)). While not required, in certain embodiments, regions of polynucleotides encoding the recombinant antibodies may be sequenced. DNA sequencing can be performed as described in Sanger *et al.* (*Proc. Natl. Acad. Sci. USA* 74:5463 (1977)) and the Amersham International plc sequencing handbook and including improvements thereto.

[279] In particular embodiments, the resulting recombinant antibodies or fragments thereof are then tested to confirm their original specificity and may be further tested for pan-specificity, *e.g.*, with related infectious agents. In particular embodiments, an antibody identified or produced according to methods described herein is tested for cell killing via

antibody dependent cellular cytotoxicity (ADCC) or apoptosis, and/or well as its ability to internalize.

### Polynucleotides

[280] The present invention, in other aspects, provides polynucleotide compositions. In preferred embodiments, these polynucleotides encode a polypeptide of the invention, *e.g.*, a region of a variable chain of an antibody that binds to Influenza A, M2, or M2e.

Polynucleotides of the invention are single-stranded (coding or antisense) or double-stranded DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include, but are not limited to, HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Alternatively, or in addition, coding or non-coding sequences are present within a polynucleotide of the present invention. Also alternatively, or in addition, a polynucleotide is linked to other molecules and/or support materials of the invention. Polynucleotides of the invention are used, *e.g.*, in hybridization assays to detect the presence of an Influenza A antibody in a biological sample, and in the recombinant production of polypeptides of the invention.

[281] Therefore, according to another aspect of the present invention, polynucleotide compositions are provided that include some or all of a polynucleotide sequence set forth in Example 1, complements of a polynucleotide sequence set forth in Example 1, and degenerate variants of a polynucleotide sequence set forth in Example 1. In certain preferred embodiments, the polynucleotide sequences set forth herein encode polypeptides capable of preferentially binding a Influenza A-infected cell as compared to a normal control uninfected cell, including a polypeptide having a sequence set forth in Examples 1 or 2. Furthermore, the invention includes all polynucleotides that encode any polypeptide of the present invention.

[282] In other related embodiments, the invention provides polynucleotide variants having substantial identity to the sequences set forth in Figure 1, for example those comprising at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide sequence of this invention, as determined using the methods described herein, (*e.g.*, BLAST analysis using standard parameters). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like.

[283] Typically, polynucleotide variants contain one or more substitutions, additions, deletions and/or insertions, preferably such that the immunogenic binding properties of the polypeptide encoded by the variant polynucleotide is not substantially diminished relative to a polypeptide encoded by a polynucleotide sequence specifically set forth herein.

In additional embodiments, the present invention provides polynucleotide fragments comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise at least about 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all intermediate lengths there between. As used herein, the term “intermediate lengths” is meant to describe any length between the quoted values, such as 16, 17, 18, 19, *etc.*; 21, 22, 23, *etc.*; 30, 31, 32, *etc.*; 50, 51, 52, 53, *etc.*; 100, 101, 102, 103, *etc.*; 150, 151, 152, 153, *etc.*; including all integers through 200-500; 500-1,000, and the like.

[284] In another embodiment of the invention, polynucleotide compositions are provided that are capable of hybridizing under moderate to high stringency conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-60°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS. One skilled in the art will understand that the stringency of hybridization can be readily manipulated, such as by altering the salt content of the hybridization solution and/or the temperature at which the hybridization is performed. For example, in another embodiment, suitable highly stringent hybridization conditions include those described above, with the exception that the temperature of hybridization is increased, *e.g.*, to 60-65°C or 65-70°C.

[285] In preferred embodiments, the polypeptide encoded by the polynucleotide variant or fragment has the same binding specificity (*i.e.*, specifically or preferentially binds to the same epitope or Influenza A strain) as the polypeptide encoded by the native polynucleotide. In certain preferred embodiments, the polynucleotides described above, *e.g.*, polynucleotide variants, fragments and hybridizing sequences, encode polypeptides that have a level of binding activity of at least about 50%, preferably at least about 70%, and more preferably at least about 90% of that for a polypeptide sequence specifically set forth herein.

[286] The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. A nucleic acid fragment of almost any length is employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative polynucleotide segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are included in many implementations of this invention.

[287] It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are multiple nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that encode a polypeptide of the present invention but which vary due to differences in codon usage are specifically contemplated by the invention. Further, alleles of the genes including the polynucleotide sequences provided herein are within the scope of the invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

[288] In certain embodiments of the present invention, mutagenesis of the disclosed polynucleotide sequences is performed in order to alter one or more properties of the encoded polypeptide, such as its binding specificity or binding strength. Techniques for mutagenesis are well-known in the art, and are widely used to create variants of both polypeptides and polynucleotides. A mutagenesis approach, such as site-specific mutagenesis, is employed for the preparation of variants and/or derivatives of the polypeptides described herein. By this approach, specific modifications in a polypeptide sequence are made through mutagenesis of the underlying polynucleotides that encode them. These techniques provides a straightforward approach to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the polynucleotide.

Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences include the nucleotide sequence of the desired mutation, as well as

a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations are employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

[289] In other embodiments of the present invention, the polynucleotide sequences provided herein are used as probes or primers for nucleic acid hybridization, *e.g.*, as PCR primers. The ability of such nucleic acid probes to specifically hybridize to a sequence of interest enable them to detect the presence of complementary sequences in a given sample. However, other uses are also encompassed by the invention, such as the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructions. As such, nucleic acid segments of the invention that include a sequence region of at least about 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence disclosed herein is particularly useful. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) including full length sequences, and all lengths in between, are also used in certain embodiments.

[290] Polynucleotide molecules having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide sequence disclosed herein, are particularly contemplated as hybridization probes for use in, *e.g.*, Southern and Northern blotting, and/or primers for use in, *e.g.*, polymerase chain reaction (PCR). The total size of fragment, as well as the size of the complementary stretch(es), ultimately depends on the intended use or application of the particular nucleic acid segment. Smaller fragments are generally used in hybridization embodiments, wherein the length of the contiguous complementary region may be varied, such as between about 15 and about 100 nucleotides, but larger contiguous complementarity stretches may be used, according to the length complementary sequences one wishes to detect.

[291] The use of a hybridization probe of about 15-25 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 12 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. Nucleic acid

molecules having gene-complementary stretches of 15 to 25 contiguous nucleotides, or even longer where desired, are generally preferred.

[292] Hybridization probes are selected from any portion of any of the sequences disclosed herein. All that is required is to review the sequences set forth herein, or to any continuous portion of the sequences, from about 15-25 nucleotides in length up to and including the full length sequence, that one wishes to utilize as a probe or primer. The choice of probe and primer sequences is governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence.

[293] Polynucleotide of the present invention, or fragments or variants thereof, are readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments are obtained by application of nucleic acid reproduction technology, such as the PCR<sup>TM</sup> technology of U. S. Patent 4,683,202, by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

#### Vectors, Host Cells and Recombinant Methods

[294] The invention provides vectors and host cells comprising a nucleic acid of the present invention, as well as recombinant techniques for the production of a polypeptide of the present invention. Vectors of the invention include those capable of replication in any type of cell or organism, including, *e.g.*, plasmids, phage, cosmids, and mini chromosomes. In various embodiments, vectors comprising a polynucleotide of the present invention are vectors suitable for propagation or replication of the polynucleotide, or vectors suitable for expressing a polypeptide of the present invention. Such vectors are known in the art and commercially available.

[295] Polynucleotides of the present invention are synthesized, whole or in parts that are then combined, and inserted into a vector using routine molecular and cell biology techniques, including, *e.g.*, subcloning the polynucleotide into a linearized vector using appropriate restriction sites and restriction enzymes. Polynucleotides of the present invention are amplified by polymerase chain reaction using oligonucleotide primers complementary to each strand of the polynucleotide. These primers also include restriction enzyme cleavage sites to facilitate subcloning into a vector. The replicable vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, and one or more marker or selectable genes.

[296] In order to express a polypeptide of the present invention, the nucleotide sequences encoding the polypeptide, or functional equivalents, are inserted into an appropriate expression vector, *i.e.*, a vector that contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods well known to those skilled in the art are used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Such techniques are described, for example, in Sambrook, J., et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y.

[297] A variety of expression vector/host systems are utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (*e.g.*, baculovirus); plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids); or animal cell systems. Within one embodiment, the variable regions of a gene expressing a monoclonal antibody of interest are amplified from a hybridoma cell using nucleotide primers. These primers are synthesized by one of ordinary skill in the art, or may be purchased from commercially available sources (*see, e.g.*, Stratagene (La Jolla, California), which sells primers for amplifying mouse and human variable regions. The primers are used to amplify heavy or light chain variable regions, which are then inserted into vectors such as ImmunoZAP™ H or ImmunoZAP™ L (Stratagene), respectively. These vectors are then introduced into *E. coli*, yeast, or mammalian-based systems for expression. Large amounts of a single-chain protein containing a fusion of the V<sub>H</sub> and V<sub>L</sub> domains are produced using these methods (*see Bird et al., Science* 242:423-426 (1988)).

[298] The “control elements” or “regulatory sequences” present in an expression vector are those non-translated regions of the vector, *e.g.*, enhancers, promoters, 5' and 3' untranslated regions, that interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, are used.

[299] Examples of promoters suitable for use with prokaryotic hosts include the phoA promoter,  $\beta$ -lactamase and lactose promoter systems, alkaline phosphatase promoter, a tryptophan (trp) promoter system, and hybrid promoters such as the tac promoter. However, other known bacterial promoters are suitable. Promoters for use in bacterial systems also usually contain a Shine-Dalgarno sequence operably linked to the DNA encoding the polypeptide. Inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSORT1 plasmid (Gibco BRL, Gaithersburg, MD) and the like are used.

[300] A variety of promoter sequences are known for eukaryotes and any are used according to the present invention. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

[301] In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. Polypeptide expression from vectors in mammalian host cells are controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (*e.g.*, Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus (CMV), a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker. One example of a suitable expression vector is pcDNA-3.1 (Invitrogen, Carlsbad, CA), which includes a CMV promoter.

[302] A number of viral-based expression systems are available for mammalian expression of polypeptides. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus that is capable of expressing the polypeptide in infected host cells

(Logan, J. and Shenk, T. (1984) Proc. Natl. Acad. Sci. 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

[303] In bacterial systems, any of a number of expression vectors are selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are desired, vectors that direct high level expression of fusion proteins that are readily purified are used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase, so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) J. Biol. Chem. 264:5503-5509); and the like. pGEX Vectors (Promega, Madison, WI) are also used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems are designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

[304] In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH are used. Examples of other suitable promoter sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. For reviews, see Ausubel *et al.* (*supra*) and Grant *et al.* (1987) Methods Enzymol. 153:516-544. Other yeast promoters that are inducible promoters having the additional advantage of transcription controlled by growth conditions include the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Yeast enhancers also are advantageously used with yeast promoters.

[305] In cases where plant expression vectors are used, the expression of sequences encoding polypeptides are driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV are used alone or in combination

with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters are used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J., et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, e.g., Hobbs, S. or Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

[306] An insect system is also used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding the polypeptide are cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence renders the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses are then used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae, in which the polypeptide of interest is expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91 :3224-3227).

[307] Specific initiation signals are also used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon are provided. Furthermore, the initiation codon is in the correct reading frame to ensure correct translation of the inserted polynucleotide. Exogenous translational elements and initiation codons are of various origins, both natural and synthetic.

[308] Transcription of a DNA encoding a polypeptide of the invention is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are known, including, e.g., those identified in genes encoding globin, elastase, albumin,  $\alpha$ -fetoprotein, and insulin. Typically, however, an enhancer from a eukaryotic cell virus is used. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, *Nature* 297:17-18 (1982) on

enhancing elements for activation of eukaryotic promoters. The enhancer is spliced into the vector at a position 5' or 3' to the polypeptide-encoding sequence, but is preferably located at a site 5' from the promoter.

[309] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) typically also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-PSCA antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein.

[310] Suitable host cells for cloning or expressing the DNA in the vectors herein are the prokaryote, yeast, plant or higher eukaryote cells described above. Examples of suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, *Enterobacteriaceae* such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 Apr. 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

[311] *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and used herein, such as *Schizosaccharomyces pombe*; *Kluyveromyces* hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris*. (EP 183,070); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[312] In certain embodiments, a host cell strain is chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation,

carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing that cleaves a “prepro” form of the protein is also used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, COS, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, are chosen to ensure the correct modification and processing of the foreign protein.

[313] Methods and reagents specifically adapted for the expression of antibodies or fragments thereof are also known and available in the art, including those described, *e.g.*, in U.S. Patent Nos. 4816567 and 6331415. In various embodiments, antibody heavy and light chains, or fragments thereof, are expressed from the same or separate expression vectors. In one embodiment, both chains are expressed in the same cell, thereby facilitating the formation of a functional antibody or fragment thereof.

[314] Full length antibody, antibody fragments, and antibody fusion proteins are produced in bacteria, in particular when glycosylation and Fc effector function are not needed, such as when the therapeutic antibody is conjugated to a cytotoxic agent (*e.g.*, a toxin) and the immunoconjugate by itself shows effectiveness in infected cell destruction. For expression of antibody fragments and polypeptides in bacteria, see, *e.g.*, U.S. Pat. Nos. 5,648,237, 5,789,199, and 5,840,523, which describes translation initiation region (TIR) and signal sequences for optimizing expression and secretion. After expression, the antibody is isolated from the *E. coli* cell paste in a soluble fraction and can be purified through, *e.g.*, a protein A or G column depending on the isotype. Final purification can be carried out using a process similar to that used for purifying antibody expressed *e.g.*, in CHO cells.

[315] Suitable host cells for the expression of glycosylated polypeptides and antibodies are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, *e.g.*, the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses are used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco are also utilized as hosts.

[316] Methods of propagation of antibody polypeptides and fragments thereof in vertebrate cells in culture (tissue culture) are encompassed by the invention. Examples of mammalian

host cell lines used in the methods of the invention are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TR1 cells (Mather *et al.*, Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[317] Host cells are transformed with the above-described expression or cloning vectors for polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

[318] For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines that stably express a polynucleotide of interest are transformed using expression vectors that contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells are allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced sequences. Resistant clones of stably transformed cells are proliferated using tissue culture techniques appropriate to the cell type.

[319] A plurality of selection systems are used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. *et al.* (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. *et al.* (1990) *Cell* 22:817-23) genes that are employed in *tk*<sup>-</sup> or *aprt*<sup>-</sup> cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance is used as the basis for selection; for example, *dhfr*, which confers resistance to methotrexate (Wigler, M. *et al.* (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); *npt*, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. *et al.* (1981) *J. Mol. Biol.* 150:1-14); and *als* or *pat*, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*). Additional

selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, and *hisD* allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). The use of visible markers has gained popularity with such markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. *et al.* (1995) *Methods Mol. Biol.* 55:121-131).

[320] Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression is confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences are identified by the absence of marker gene function. Alternatively, a marker gene is placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells that contain and express a desired polynucleotide sequence are identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include, for example, membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

[321] A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Nonlimiting examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide is preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. *et al.* (1990; *Serological Methods, a Laboratory Manual*, APS Press, St Paul, Minn.) and Maddox, D. E. *et al.* (1983; *J. Exp. Med.* 158:1211-1216).

[322] Various labels and conjugation techniques are known by those skilled in the art and are used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof are cloned into a vector for the

production of an mRNA probe. Such vectors are known in the art, are commercially available, and are used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures are conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which are used include, but are not limited to, radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

[323] The polypeptide produced by a recombinant cell is secreted or contained intracellularly depending on the sequence and/or the vector used. Expression vectors containing polynucleotides of the invention are designed to contain signal sequences that direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane.

[324] In certain embodiments, a polypeptide of the invention is produced as a fusion polypeptide further including a polypeptide domain that facilitates purification of soluble proteins. Such purification-facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Amgen, Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the encoded polypeptide are used to facilitate purification. An exemplary expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. *et al.* (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors used for producing fusion proteins is provided in Kroll, D. J. *et al.* (1993; *DNA Cell Biol.* 12:441-453).

[325] In certain embodiments, a polypeptide of the present invention is fused with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. The heterologous signal sequence selected preferably is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. For prokaryotic host cells, the signal sequence is selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or

heat-stable enterotoxin II leaders. For yeast secretion, the signal sequence is selected from, e.g., the yeast invertase leader,  $\alpha$  factor leader (including *Saccharomyces* and *Kluyveromyces*  $\alpha$  factor leaders), or acid phosphatase leader, the *C. albicans* glucoamylase leader, or the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

[326] When using recombinant techniques, the polypeptide or antibody is produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the polypeptide or antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Carter *et al.*, Bio/Technology 10:163-167 (1992) describe a procedure for isolating antibodies that are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min. Cell debris is removed by centrifugation. Where the polypeptide or antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Optionally, a protease inhibitor such as PMSF is included in any of the foregoing steps to inhibit proteolysis and antibiotics is included to prevent the growth of adventitious contaminants.

[327] The polypeptide or antibody composition prepared from the cells are purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the polypeptide or antibody. Protein A is used to purify antibodies or fragments thereof that are based on human  $\gamma_1$ ,  $\gamma_2$ , or  $\gamma_4$  heavy chains (Lindmark *et al.*, J. Immunol. Meth. 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human  $\gamma_3$  (Guss *et al.*, EMBO J. 5:1567-1575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the polypeptide or antibody comprises a C<sub>H</sub> 3 domain, the Bakerbond ABX™ resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™

chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the polypeptide or antibody to be recovered.

Following any preliminary purification step(s), the mixture comprising the polypeptide or antibody of interest and contaminants are subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (*e.g.*, from about 0-0.25M salt).

#### Pharmaceutical Compositions

[328] The invention further includes pharmaceutical formulations including a polypeptide, antibody, or modulator of the present invention, at a desired degree of purity, and a pharmaceutically acceptable carrier, excipient, or stabilizer (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)). In certain embodiments, pharmaceutical formulations are prepared to enhance the stability of the polypeptide or antibody during storage, *e.g.*, in the form of lyophilized formulations or aqueous solutions.

Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include, *e.g.*, buffers such as acetate, Tris, phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; tonicifiers such as trehalose and sodium chloride; sugars such as sucrose, mannitol, trehalose or sorbitol; surfactant such as polysorbate; salt-forming counter-ions such as sodium; metal complexes (*e.g.* Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG). In certain embodiments, the therapeutic formulation preferably comprises the polypeptide or antibody at a concentration of between 5-200 mg/ml, preferably between 10-100 mg/ml.

[329] The formulations herein also contain one or more additional therapeutic agents suitable for the treatment of the particular indication, *e.g.*, infection being treated, or to

prevent undesired side-effects. Preferably, the additional therapeutic agent has an activity complementary to the polypeptide or antibody of the present invention, and the two do not adversely affect each other. For example, in addition to the polypeptide or antibody of the invention, an additional or second antibody, anti-viral agent, anti-infective agent and/or cardioprotectant is added to the formulation. Such molecules are suitably present in the pharmaceutical formulation in amounts that are effective for the purpose intended.

[330] The active ingredients, *e.g.*, polypeptides and antibodies of the invention and other therapeutic agents, are also entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and polymethylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

[331] Sustained-release preparations are prepared. Suitable examples of sustained-release preparations include, but are not limited to, semi-permeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Nonlimiting examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and  $\gamma$  ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

[332] Formulations to be used for *in vivo* administration are preferably sterile. This is readily accomplished by filtration through sterile filtration membranes.

### Diagnostic Uses

[333] Antibodies and fragments thereof, and therapeutic compositions, of the invention specifically bind or preferentially bind to infected cells or tissue, as compared to normal control cells and tissue. Thus, these influenza A antibodies are used to detect infected cells or tissues in a patient, biological sample, or cell population, using any of a variety of diagnostic and prognostic methods, including those described herein. The ability of an anti-M2e specific antibody to detect infected cells depends upon its binding specificity, which is readily

determined by testing its ability to bind to infected cells or tissues obtained from different patients, and/or from patients infected with different strains of Influenza A.

Diagnostic methods generally involve contacting a biological sample obtained from a patient, such as, *e.g.*, blood, serum, saliva, urine, sputum, a cell swab sample, or a tissue biopsy, with an Influenza A, *e.g.*, HuM2e antibody and determining whether the antibody preferentially binds to the sample as compared to a control sample or predetermined cut-off value, thereby indicating the presence of infected cells. In particular embodiments, at least two-fold, three-fold, or five-fold more HuM2e antibody binds to an infected cell as compared to an appropriate control normal cell or tissue sample. A pre-determined cut-off value is determined, *e.g.*, by averaging the amount of HuM2e antibody that binds to several different appropriate control samples under the same conditions used to perform the diagnostic assay of the biological sample being tested.

[334] Bound antibody is detected using procedures described herein and known in the art. In certain embodiments, diagnostic methods of the invention are practiced using HuM2e antibodies that are conjugated to a detectable label, *e.g.*, a fluorophore, to facilitate detection of bound antibody. However, they are also practiced using methods of secondary detection of the HuM2e antibody. These include, for example, RIA, ELISA, precipitation, agglutination, complement fixation and immuno-fluorescence.

[335] In certain procedures, the HuM2e antibodies are labeled. The label is detected directly. Exemplary labels that are detected directly include, but are not limited to, radiolabels and fluorochromes. Alternatively, or in addition, labels are moieties, such as enzymes, that must be reacted or derivatized to be detected. Nonlimiting examples of isotope labels are  $^{99}\text{Tc}$ ,  $^{14}\text{C}$ ,  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^3\text{H}$ ,  $^{32}\text{P}$  and  $^{35}\text{S}$ . Fluorescent materials that are used include, but are not limited to, for example, fluorescein and its derivatives, rhodamine and its derivatives, auramine, dansyl, umbelliferone, luciferin, 2,3-dihydrophthalazinediones, horseradish peroxidase, alkaline phosphatase, lysozyme, and glucose-6-phosphate dehydrogenase.

[336] An enzyme label is detected by any of the currently utilized colorimetric, spectrophotometric, fluorospectro-photometric or gasometric techniques. Many enzymes which are used in these procedures are known and utilized by the methods of the invention. Nonlimiting examples are peroxidase, alkaline phosphatase,  $\beta$ -glucuronidase,  $\beta$ -D-glucosidase,  $\beta$ -D-galactosidase, urease, glucose oxidase plus peroxidase, galactose oxidase plus peroxidase and acid phosphatase.

[337] The antibodies are tagged with such labels by known methods. For instance, coupling agents such as aldehydes, carbodiimides, dimaleimide, imidates, succinimides, bis-diazotized

benzadine and the like are used to tag the antibodies with the above-described fluorescent, chemiluminescent, and enzyme labels. An enzyme is typically combined with an antibody using bridging molecules such as carbodiimides, periodate, diisocyanates, glutaraldehyde and the like. Various labeling techniques are described in Morrison, *Methods in Enzymology* 32b, 103 (1974), Syvanen *et al.*, *J. Biol. Chem.* 284, 3762 (1973) and Bolton and Hunter, *Biochem J.* 133, 529(1973).

[338] HuM2e antibodies of the present invention are capable of differentiating between patients with and patients without an Influenza A infection, and determining whether or not a patient has an infection, using the representative assays provided herein. According to one method, a biological sample is obtained from a patient suspected of having or known to have an Influenza A infection. In preferred embodiments, the biological sample includes cells from the patient. The sample is contacted with an HuM2e antibody, *e.g.*, for a time and under conditions sufficient to allow the HuM2e antibody to bind to infected cells present in the sample. For instance, the sample is contacted with an HuM2e antibody for 10 seconds, 30 seconds, 1 minute, 5 minutes, 10 minutes, 30 minutes, 1 hour, 6 hours, 12 hours, 24 hours, 3 days or any point in between. The amount of bound HuM2e antibody is determined and compared to a control value, which may be, *e.g.*, a pre-determined value or a value determined from normal tissue sample. An increased amount of antibody bound to the patient sample as compared to the control sample is indicative of the presence of infected cells in the patient sample.

[339] In a related method, a biological sample obtained from a patient is contacted with an HuM2e antibody for a time and under conditions sufficient to allow the antibody to bind to infected cells. Bound antibody is then detected, and the presence of bound antibody indicates that the sample contains infected cells. This embodiment is particularly useful when the HuM2e antibody does not bind normal cells at a detectable level.

[340] Different HuM2e antibodies possess different binding and specificity characteristics. Depending upon these characteristics, particular HuM2e antibodies are used to detect the presence of one or more strains of Influenza A. For example, certain antibodies bind specifically to only one or several strains of Influenza virus, whereas others bind to all or a majority of different strains of Influenza virus. Antibodies specific for only one strain of Influenza A are used to identify the strain of an infection.

[341] In certain embodiments, antibodies that bind to an infected cell preferably generate a signal indicating the presence of an infection in at least about 20% of patients with the infection being detected, more preferably at least about 30% of patients. Alternatively, or in

addition, the antibody generates a negative signal indicating the absence of the infection in at least about 90% of individuals without the infection being detected. Each antibody satisfies the above criteria; however, antibodies of the present invention are used in combination to improve sensitivity.

[342] The present invention also includes kits useful in performing diagnostic and prognostic assays using the antibodies of the present invention. Kits of the invention include a suitable container comprising a HuM2e antibody of the invention in either labeled or unlabeled form. In addition, when the antibody is supplied in a labeled form suitable for an indirect binding assay, the kit further includes reagents for performing the appropriate indirect assay. For example, the kit includes one or more suitable containers including enzyme substrates or derivatizing agents, depending on the nature of the label. Control samples and/or instructions are also included.

#### Therapeutic/ Prophylactic Uses

[343] Passive immunization has proven to be an effective and safe strategy for the prevention and treatment of viral diseases. (See Keller et al., Clin. Microbiol. Rev. 13:602-14 (2000); Casadevall, Nat. Biotechnol. 20:114 (2002); Shibata et al., Nat. Med. 5:204-10 (1999); and Igarashi et al., Nat. Med. 5:211-16 (1999), each of which are incorporated herein by reference)). Passive immunization using human monoclonal antibodies provide an immediate treatment strategy for emergency prophylaxis and treatment of influenza

[344] HuM2e antibodies and fragments thereof, and therapeutic compositions, of the invention specifically bind or preferentially bind to infected cells, as compared to normal control uninfected cells and tissue. Thus, these HuM2e antibodies are used to selectively target infected cells or tissues in a patient, biological sample, or cell population. In light of the infection-specific binding properties of these antibodies, the present invention provides methods of regulating (*e.g.*, inhibiting) the growth of infected cells, methods of killing infected cells, and methods of inducing apoptosis of infected cells. These methods include contacting an infected cell with an HuM2e antibody of the invention. These methods are practiced *in vitro*, *ex vivo*, and *in vivo*.

[345] In various embodiments, antibodies of the invention are intrinsically therapeutically active. Alternatively, or in addition, antibodies of the invention are conjugated to a cytotoxic agent or growth inhibitory agent, *e.g.*, a radioisotope or toxin, that is used in treating infected cells bound or contacted by the antibody.

[346] In one embodiment, the invention provides methods of treating or preventing infection in a patient, including the steps of providing an HuM2e antibody of the invention to a patient diagnosed with, at risk of developing, or suspected of having an Influenza A infection. The methods of the invention are used in the first-line treatment of the infection, follow-on treatment, or in the treatment of a relapsed or refractory infection. Treatment with an antibody of the invention is a stand alone treatment. Alternatively, treatment with an antibody of the invention is one component or phase of a combination therapy regime, in which one or more additional therapeutic agents are also used to treat the patient.

[347] Subjects at risk for an influenza virus -related diseases or disorders include patients who have come into contact with an infected person or who have been exposed to the influenza virus in some other way. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the influenza virus -related disease or disorder, such that a disease or disorder is prevented or, alternatively, delayed in its progression.

[348] In various aspects, the huM2e is administered substantially contemporaneously with or following infection of the subject, i.e., therapeutic treatment. In another aspect, the antibody provides a therapeutic benefit. In various aspects, a therapeutic benefit includes reducing or decreasing progression, severity, frequency, duration or probability of one or more symptoms or complications of influenza infection, virus titer, virus replication or an amount of a viral protein of one or more influenza strains. In still another aspect, a therapeutic benefit includes hastening or accelerating a subject's recovery from influenza infection.

[349] Methods for preventing an increase in influenza virus titer, virus replication, virus proliferation or an amount of an influenza viral protein in a subject are further provided. In one embodiment, a method includes administering to the subject an amount of a huM2e antibody effective to prevent an increase in influenza virus titer, virus replication or an amount of an influenza viral protein of one or more influenza strains or isolates in the subject.

[350] Methods for protecting a subject from infection or decreasing susceptibility of a subject to infection by one or more influenza strains/isolates or subtypes, i.e., prophylactic methods, are additionally provided. In one embodiment, a method includes administering to the subject an amount of huM2e antibody that specifically binds influenza M2 effective to protect the subject from infection, or effective to decrease susceptibility of the subject to infection, by one or more influenza strains/isolates or subtypes.

[351] Optionally, the subject is further administered with a second agent such as, but not limited to, an influenza virus antibody, an anti-viral drug such as a neuraminidase inhibitor, a HA inhibitor, a sialic acid inhibitor or an M2 ion channel inhibitor, a viral entry inhibitor or a viral attachment inhibitor. The M2 ion channel inhibitor is for example amantadine or rimantadine. The neuraminidase inhibitor for example zanamivir, or oseltamivir phosphate.

[352] Symptoms or complications of influenza infection that can be reduced or decreased include, for example, chills, fever, cough, sore throat, nasal congestion, sinus congestion, nasal infection, sinus infection, body ache, head ache, fatigue, pneumonia, bronchitis, ear infection, ear ache or death.

[353] For *in vivo* treatment of human and non-human patients, the patient is usually administered or provided a pharmaceutical formulation including a HuM2e antibody of the invention. When used for *in vivo* therapy, the antibodies of the invention are administered to the patient in therapeutically effective amounts (*i.e.*, amounts that eliminate or reduce the patient's viral burden). The antibodies are administered to a human patient, in accord with known methods, such as intravenous administration, *e.g.*, as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. The antibodies may be administered parenterally, when possible, at the target cell site, or intravenously. Intravenous or subcutaneous administration of the antibody is preferred in certain embodiments. Therapeutic compositions of the invention are administered to a patient or subject systemically, parenterally, or locally.

[354] For parenteral administration, the antibodies are formulated in a unit dosage injectable form (solution, suspension, emulsion) in association with a pharmaceutically acceptable, parenteral vehicle. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils and ethyl oleate are also used. Liposomes are used as carriers. The vehicle contains minor amounts of additives such as substances that enhance isotonicity and chemical stability, *e.g.*, buffers and preservatives. The antibodies are typically formulated in such vehicles at concentrations of about 1 mg/ml to 10 mg/ml.

[355] The dose and dosage regimen depends upon a variety of factors readily determined by a physician, such as the nature of the infection and the characteristics of the particular cytotoxic agent or growth inhibitory agent conjugated to the antibody (when used), *e.g.*, its therapeutic index, the patient, and the patient's history. Generally, a therapeutically effective amount of an antibody is administered to a patient. In particular embodiments, the amount of

antibody administered is in the range of about 0.01 mg/kg to about 100 mg/kg of patient body weight, or more preferably, in the range of about 0.1 mg/kg to about 40 mg/kg of patient body weight. Depending on the type and severity of the infection, about 0.1 mg/kg to about 40 mg/kg body weight (*e.g.*, about 0.1- 40 mg/kg/dose) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. In alternative embodiments, the amount of antibody administered is in the range of 0.01 mg/kg to 0.1 mg/kg, 0.1 mg/kg to 0.10 mg/kg, 0.10 mg/kg to 1 mg/kg, 1 mg/kg to 10 mg/kg, 10 mg/kg to 20 mg/kg, 20 mg/kg to 30 mg/kg, 30 mg/kg to 40 mg/kg, 40 mg/kg to 50 mg/kg, 50 mg/kg to 60 mg/kg, 60 mg/kg to 70 mg/kg, 70 mg/kg to 80 mg/kg, 80 mg/kg to 90 mg/kg, or 90 mg/kg to 100 mg/kg of patient body weight. In other aspects, the amount of antibody administered is in the range of 0.01 mg/kg to 100 mg/kg, 0.1 mg/kg to 60 mg/kg, 10 mg/kg to 40 mg/kg, 20 mg/kg to 30 mg/kg of patient body weight or any range in between. The progress of this therapy is readily monitored by conventional methods and assays and based on criteria known to the physician or other persons of skill in the art.

[356] In one particular embodiment, an immunoconjugate including the antibody conjugated with a cytotoxic agent is administered to the patient. Preferably, the immunoconjugate is internalized by the cell, resulting in increased therapeutic efficacy of the immunoconjugate in killing the cell to which it binds. In one embodiment, the cytotoxic agent targets or interferes with the nucleic acid in the infected cell. Examples of such cytotoxic agents are described above and include, but are not limited to, maytansinoids, calicheamicins, ribonucleases and DNA endonucleases.

[357] Other therapeutic regimens are combined with the administration of the HuM2e antibody of the present invention. The combined administration includes co-administration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities. Preferably such combined therapy results in a synergistic therapeutic effect.

[358] In certain embodiments, it is desirable to combine administration of an antibody of the invention with another antibody directed against another antigen associated with the infectious agent.

[359] Aside from administration of the antibody protein to the patient, the invention provides methods of administration of the antibody by gene therapy. Such administration of nucleic acid encoding the antibody is encompassed by the expression "administering a

therapeutically effective amount of an antibody". See, for example, PCT Patent Application Publication WO96/07321 concerning the use of gene therapy to generate intracellular antibodies.

[360] In another embodiment, anti-M2e antibodies of the invention are used to determine the structure of bound antigen, e.g., conformational epitopes, the structure of which is then used to develop a vaccine having or mimicking this structure, e.g., through chemical modeling and SAR methods. Such a vaccine could then be used to prevent Influenza A infection.

[361] All of the above U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet are incorporated herein by reference, in their entirety.

## EXAMPLES

### Example 1: Screening and Characterization of M2e-specific Antibodies Present in Human Plasma Using Cells Expressing Recombinant M2e Protein

[362] Fully human monoclonal antibodies specific for M2 and capable of binding to influenza A infected cells and the influenza virus itself were identified in patient serum, as described below.

#### *Expression of M2 in Cell Lines*

[363] An expression construct containing the M2 full length cDNA, corresponding to the derived M2 sequence found in Influenza subtype H3N2, was transfected into 293 cells.

[364] The M2 cDNA is encoded by the following polynucleotide sequence and SEQ ID NO: 53:

```
ATGAGTCTTCTAACCGAGGTCGAAACGCCTATCAGAAACGAATGGGGGTGCAGATGCAACGATTC
AAGTGATCCTCTTGTTGTTGCCGCAAGTATCATTGGGATCCTGCACTTGATATTGTGGATTCTTG
ATCGTCTTTTTTTCAAATGCATTTATCGTCTCTTTAAACACGGTCTGAAAAGAGGGCCTTCTACG
GAAGGAGTACCAGAGTCTATGAGGGAAGAATATCGAAAGGAACAGCAGAGTGCTGTGGATGCTGA
CGATAGTCATTTTGTCAACATAGAGCTGGAG
```

[365] The cell surface expression of M2 was confirmed using the anti-M2e peptide specific MAb 14C2. Two other variants of M2, from A/Hong Kong/483/1997 (HK483) and A/Vietnam/1203/2004 (VN1203), were used for subsequent analyses, and their expression was determined using M2e-specific monoclonal antibodies of the present invention, since 14C2 binding may be abrogated by the various amino acid substitutions in M2e.

#### *Screening of Antibodies in Peripheral Blood*

[366] Over 120 individual plasma samples were tested for antibodies that bound M2. None of them exhibited specific binding to the M2e peptide. However, 10% of the plasma samples contained antibodies that bound specifically to the 293-M2 H3N2 cell line. This indicates that the antibodies could be categorized as binding to conformational determinants of an M2 homotetramer, and binding to conformational determinants of multiple variants of the M2 homotetramer; they could not be specific for the linear M2e peptide.

*Characterization of Anti-M2 MAbs*

[367] The human MAbs identified through this process proved to bind to conformational epitopes on the M2 homotetramer. They bound to the original 293-M2 transfectant, as well as to the two other cell-expressed M2 variants. The 14C2 MAb, in addition to binding the M2e peptide, proved to be more sensitive to the M2 variant sequences. Moreover, 14C2 does not readily bind influenza virions, while the conformation specific anti-M2 MAbs did.

[368] These results demonstrate that the methods of the invention provide for the identification of M2 MAbs from normal human immune responses to influenza without a need for specific immunization of M2. If used for immunotherapy, these fully human MAbs have the potential to be better tolerated by patients than humanized mouse antibodies. Additionally, and in contrast to 14C2 and the Gemini Biosciences MAbs, which bind to linear M2e peptide, the MAbs of the invention bind to conformational epitopes of M2, and are specific not only for cells infected with A strain influenza, but also for the virus itself. Another advantage of the MAbs of the invention is that they each bind all of the M2 variants yet tested, indicating that they are not restricted to a specific linear amino acid sequence.

Example 2: Identification of M2-Specific Antibodies

[369] Mononuclear or B cells expressing three of the MAbs identified in human serum as described in Example 1 were diluted into clonal populations and induced to produce antibodies. Antibody containing supernatants were screened for binding to 293 FT cells stably transfected with the full length M2E protein from influenza strain Influenza subtype H3N2. Supernatants which showed positive staining/binding were re-screened again on 293 FT cells stably transfected with the full length M2E protein from influenza strain Influenza subtype H3N2 and on vector alone transfected cells as a control.

[370] The variable regions of the antibodies were then rescue cloned from the B cell wells whose supernatants showed positive binding. Transient transfections were performed in 293 FT cells to reconstitute and produce these antibodies. Reconstituted antibody supernatants were screened for binding to 293 FT cells stably transfected with the full length M2E protein

as detailed above to identify the rescued anti-M2E antibodies. Three different antibodies were identified: 8i10, 21B15 and 23K12. A fourth additional antibody clone was isolated by the rescue screens, 4C2. However, it was not unique and had the exact same sequence as clone 8i10 even though it came from a different donor than clone 8i10.

[371] The sequences of the kappa and gamma variable regions of these antibodies are provided below.

*Clone 8i10:*

[372] The Kappa LC variable region of the anti M2 clone 8i10 was cloned as Hind III to BsiW1 fragment (see below), and is encoded by the following polynucleotide sequences, and SEQ ID NO: 54 (top) and SEQ ID NO: 55 (bottom):

HindIII

AAGCTTCCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAGGT G  
TTCGAAGGTGGTACCTGTACTCCCAGGAGCGAGTCGAGGACCCGAGGACGATGAGACCGAGGCTCCA C  
CCAGATGTGACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC A  
GGTCTACACTGTAGGTCTACTGGGTCAGAGGTAGGAGGGACAGACGTAGACATCCTCTGTCTCAGTGG T  
TCACTTGCCGGGCGAGTCAGAACATTTACAAGTATTTAAATTGGTATCAGCAGAGACCAGGGAAAGCC C  
AGTGAACGGCCCGCTCAGTCTTGTAATGTTCAATAATTTAACCATAGTCGTCTCTGGTCCCTTTTCGG G  
CTAAGGGCCTGATCTCTGCTGCATCCGGGTTGCAAAGTGCGGTCCCATCAAGGTTTCAGTGGCAGTGGA T  
GATTCCCGGACTAGAGACGACGTAGGCCCAACGTTTCACCCAGGGTAGTTCCAAGTCACCGTCACCT A  
CTGGGACAGATTTCACTCTCACCATCACCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAA C  
GACCCTGTCTAAAGTGAGAGTGGTAGTGGTCAGACGTTGGACTTCTAAAACGTTGAATGATGACAGTT G

BsiWI

AGAGTTACAGTCCCCCTCTCACTTTCGGCGGAGGGACCAGGGTGGAGATCAAACGTACG  
TCTCAATGTCAGGGGGAGAGTGAAAGCCGCCTCCCTGGTCCACCTCTAGTTTGCATGC

[373] The translation of the 8i10 Kappa LC variable region is as follows, polynucleotide sequence (above, SEQ ID NO: 54, top) and amino acid sequence (below, corresponding to SEQ ID NO: 56):

HindIII

AAGCTTCCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAGGT G  
M D M R V L A Q L L G L L L L W L R G  
CCAGATGTGACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC A  
A R C D I Q M T Q S P S S L S A S V G D R V T  
TCACTTGCCGGGCGAGTCAGAACATTTACAAGTATTTAAATTGGTATCAGCAGAGACCAGGGAAAGCC C  
I T C R A S Q N I Y K Y L N W Y Q Q R P G K A  
CTAAGGGCCTGATCTCTGCTGCATCCGGGTTGCAAAGTGCGGTCCCATCAAGGTTTCAGTGGCAGTGGA T  
P K G L I S A A S G L Q S G V P S R F S G S G  
CTGGGACAGATTTCACTCTCACCATCACCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAA C  
S G T D F T L T I T S L Q P E D F A T Y Y C Q

BsiWI

AGAGTTACAGTCCCCCTCTCACTTTCGGCGGAGGGACCAGGGTGGAGATCAAACGTACG  
Q S Y S P P L T F G G G T R V E I K R T

[374] The amino acid sequence of the 8i10 Kappa LC variable region is as follows, with specific domains identified below (CDR sequences defined according to Kabat methods):

|                                  |                                |
|----------------------------------|--------------------------------|
| MDMRVLAQLLGLLLLWLRGARC           | VK leader (SEQ ID NO: 57)      |
| DIQMTQSPSSLASVGDRVITC            | FR1 (SEQ ID NO: 58)            |
| RASQNIYKYLN                      | CDR1 (SEQ ID NO: 59)           |
| WYQQRPGKAPKGLIS                  | FR2 (SEQ ID NO: 60)            |
| AASGLQS                          | CDR2 (SEQ ID NO: 61)           |
| GVPSRFSGSGSGTDFTLTITSLQPEDFATYYC | FR3 (SEQ ID NO: 62)            |
| QQSYSPPLT                        | CDR3 (SEQ ID NO: 63)           |
| FGGGTRVEIK                       | FR4 (SEQ ID NO: 255)           |
| RT                               | Start of Kappa constant region |

[375] The following is an example of the Kappa LC variable region of 8i10 cloned into the expression vector pcDNA3.1 which already contained the Kappa LC constant region (upper polynucleotide sequence corresponds to SEQ ID NO: 65, lower polynucleotide sequence corresponds to SEQ ID NO: 66, amino acid sequence corresponds to SEQ ID NO: 56 shown above). Bases in black represents pcDNA3.1 vector sequences, blue bases represent the cloned antibody sequences. The antibodies described herein have also been cloned into the expression vector pCEP4.

**[376]** The 8i10 Gamma HC variable region was cloned as a Hind III to Xho I fragment, and is encoded the following polynucleotide sequences, and SEQ ID NO: 67 (top) and SEQ ID NO: 68 (bottom).

100

[377] The translation of the 8i10 Gamma HC is as follows, polynucleotide sequence (above, SEQ ID NO: 67, top) and amino acid sequence (below, corresponding to SEQ ID NO: 69):

HindIII

```

AAGCTTCCACCATGAAACACCTGTGGTTCTTCCTTCTCCTGGTGGCAGCTCCCAGCTGGGT C
      M   K   H   L   W   F   F   L   L   L   V   A   A   P   S   W   V
CTGTCCCAGGTGCAATTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTG
  L   S   Q   V   Q   L   Q   E   S   G   P   G   L   V   K   P   S   E   T   L
TCCCTCACCTGCACTGTCTCTGGTTCGTCCATCAGTAATTACTACTGGAGCTGGATCCGG
  S   L   T   C   T   V   S   G   S   S   I   S   N   Y   Y   W   S   W   I   R
CAGTCCCCAGGGAAGGGACTGGAGTGGATTGGGTTTATCTATTACGGTGGAAACACCAAG
  Q   S   P   G   K   G   L   E   W   I   G   F   I   Y   Y   G   G   N   T   K
TACAATCCCTCCCTCAAGAGCCGCGTCACCATATCACAAGACACTTCCAAGAGTCAGGTC
  Y   N   P   S   L   K   S   R   V   T   I   S   Q   D   T   S   K   S   Q   V
TCCCTGACGATGAGCTCTGTGACCGCTGCGGAATCGGCCGTCTATTTCTGTGCGAGAGCG
  S   L   T   M   S   S   V   T   A   A   E   S   A   V   Y   F   C   A   R   A
TCTTGTAGTGGTGGTTACTGTATCCTTGACTACTGGGGCCAGGGAACCCTGGTCAACGTC
  S   C   S   G   G   Y   C   I   L   D   Y   W   G   Q   G   T   L   V   T   V
TCGAG
  S

```

XhoI

[378] The amino acid sequence of the 8i10 Gamma HC is as follows with specific domains identified below (CDR sequences defined according to Kabat methods):

|                                  |                           |
|----------------------------------|---------------------------|
| MKHLWFFLLLVAAPSWVLS              | VH leader (SEQ ID NO: 70) |
| QVQLQESGPGLVKPSETLSLTCTVSGSSIS   | FR1 (SEQ ID NO: 71)       |
| NYYWS                            | CDR1 (SEQ ID NO: 72)      |
| WIRQSPGKGLEWIG                   | FR2 (SEQ ID NO: 73)       |
| FIYYGGNTKYNPSLKS                 | CDR2 (SEQ ID NO: 74)      |
| RVTISQDTSKSQVSLTMSSVTAAESAVYFCAR | FR3 (SEQ ID NO: 75)       |
| ASCSGGYCILD                      | CDR3 (SEQ ID NO: 76)      |
| YWGQGTLVTVS                      | FR4 (SEQ ID NO: 77)       |

[379] The following is an example of the Gamma HC variable region of 8i10 cloned into the expression vector pcDNA3.1 which already contained the Gamma HC constant region (upper polynucleotide sequence corresponds to SEQ ID NO: 78, lower polynucleotide sequence corresponds to SEQ ID NO: 79, amino acid sequence corresponds to SEQ ID NO: 69 shown above). Bases in black represents pcDNA3.1 vector sequences, blue bases represent the cloned antibody sequences.

PmeI (900)  
NheI (894) HindIII (910)

TGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCCAGCTGGCTAGCGTTTAAACCTTAAGCTTCCACCATGAAACACCTGTGGTT  
ACCGAATAGCTTTAATTATGCTGAGTGATATCCCTCTGGGTTCGACCGATCGCAAATTTGAATTCGAGAAGGTGCTACTTTGTGGACACCAA  
CTTCCTTCTCCTGGTGGCAGCTCCCAGCTGGGTCTGTCCCAGGTGCAATTGCAGGAGTCGGGGCCCA  
GAAGGAAGAGGACCACCGTCGAGGGTCGACCCAGGACAGGGTCCACGTTAACGTCCTCAGCCCCGGGT  
GGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCACCTGCACTGTCTCTGGTTCTGCCATCAGTAATT  
CCTGACCACTTCGGAAGCCTCTGGGACAGGGAGTGGACGTGACAGAGACCAAGCAGGTAGTCATTAA  
ACTACTGGAGCTGGATCCGGCAGTCCCCAGGGAAGGGACTGGAGTGGATTGGGTTTATCTATTACGG  
TGATGACCTCGACCTAGGCCGTGAGGGTCCCTTCCCTGACCTCACCTAACCCAAATAGATAATGCC  
TGGAAACACCAAGTACAATCCCTCCCTCAAGAGCCGCGTCACCATATCACAAGACACTTCCAAGAGT  
ACCTTTGTGGTTCATGTTAGGGAGGGAGTTCTCGGCGCAGTGGTATAGTGTCTGTGAAGGTTCTCA  
CAGGTCTCCCTGACGATGAGCTCTGTGACCGCTGCGGAATCGGCCGTCTATTTCTGTGCGAGAGCGT  
GTCCAGAGGGACTGCTACTCGAGACACTGGCGACGCCTTAGCCGGCAGATAAAGACACGCTCTCGCA  
CTTGTAGTGGTGGTTACTGTATCCTTGACTACTGGGGCCAGGGAACCCCTGGTCACCGTCTCGAGAGCCCTCCA  
GAACATCACCACCAATGACATAGGAACTGATGACCCCGGTCCCTTGGGACCAGTGGCAGCTCTCGGAGGT  
S C S G G Y C I L D Y W G Q G T L V T V S R A S  
CCAAGGGCCCATCGGTCTTCCCTTGGCACCCTCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCCTGGTCAAGGACTACTTCCCGGAACCGGTGA C  
GGTTCCCGGGTAGCCAGAAGGGGACCGTGGGAGGAGTTCTCGTGGAGACCCCGTGTGCGCGGACCCGACGACAGTTCCTGATGAAGGGGCTTGGCCACT G  
T K G P S V F P L A P S S K S T S G G T A A L G C L V K D Y F P E P V T  
GGTGTCTGGAACCTCAGGCGCCCTGACAGCGCGGTGCACACCTTCCCGGCTGTCTACAGTCTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTC C  
GCACAGCACCTTGAAGTCCGCGGGACTGGTCCCGCACCTGTGGAAGGGGCCGACAGGATGTCAGGAGTCTCGAGATGAGGGAGTCTCGCACCCTGGCAGCGGAG G  
V S W N S G A L T S G V H T F P A V L Q S S G L Y S L S S V V T V P S  
AGCAGCTTGGGCAACCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAATCTTGTGACAAAACCTCACACA T  
TCGTCGAACCCGTTGGTCTGGATGTAGACGTTCGCACTTAGTGTTCGGGTCTGTGTGGTTCACCTGTCTCTCAACTCGGGTTTAGAACACTGTCTTTGAGTGTGT A  
S S L G T Q T Y I C N V N H K P S N T K V D K R V E P K S C D K T H T  
GCCCCCGTGGCCAGCACCTGAACCTCGGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACAT G  
CGGGTGGCAACGGTCTGGACTTGGAGGACCCCTTGGCAGTCAAGAGGAGGAGGGGGTTCGGGTTCTGTGGGAGTACTAGGGGCGCTGGGGAATCCAGTGTA C  
C P P C P A P E L L G G P S V F L G P P K P K D T L M I S R T E V T C  
CGTGGTGGTGGACGTGAGCCACGAAGACCCCTGAGGTCAAGTCAACTGGTACCTGGACGGCGTGGAGGTGCATAATGCCAAGACAAGCCGCGGGAGGAGCAGTA C  
GCACCCACCTGCACTCGGTGCTTCTGGGACTCCAGTTCAGTTGACCATGCACTGCGCGCACTCCACGTATTACGTTCTGTCTCGGCGCCCTCTCTGTCAT G  
V V V D V S H E D P E V K F N W Y V D G V E V H N A K T K P R E E Q Y  
AACAGCACGTACCTGTGGTTCAGGTCCTCACCCTCTGCAACAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAAACAAAGCCCTCCAGCCCCC A  
TTGTCGTGATGGCACACAGTTCGAGGAGTGGCAGGAGTGGTCTGACCGACTTACCGTTCTCTCATGTTACGTTCCAGAGGTGTTTTCGGGAGGGTCCGGGG T  
N S T Y R V V S V L T V L H Q D W L N G K E Y K C K V S N K A L P A P  
TCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCGGAGAACCACAGGTGTACACCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGACGCTGACCT G  
AGCTCTTTTGGTAGAGTTTTCGGTTTCCCGTCGGGGCTCTTGGTGTCCACATGTGGGACGGGGTATGGGCCCTCTCTACTGTTCTTGGTCCAGTCCGACTGCA C  
I E K T I S K A K G Q P R E G T C V Y T L P P S R E E M T K N Q V S L T C  
CCTGGTCAAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGACGCCGAGAACAACTACAAGACACGCCCTCCCGTGTGGACTCCGACGG C  
GGACCACTTTCCGAAGATAGGGTCTGTAGCGGCACCTCACCTCTCTGTTACCCGTCGGGCTCTTGTGATGTTCTGGTGGGAGGGACGACCTGAGGCTGCC G  
L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V L D S D G  
TCCTTCTCTCTATAGCAAGCTCACCGTGACAAAGAGCAGGTGGCAGCAGGGGAACGCTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACG C  
AGGAAGAAGGAGATATCGTTGAGTGGCACCTGTCTCTGTCACCGTCTGTCGAGAGAGTACGAGGCACTACGTACTCCGAGACGTGTTGGTGTATGTGC G  
S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T  
Apal (2339)  
DraIII (2338)  
XbaI (2333) PmeI (2345)  
AGAAGAGCCTCTCCCTGCTCTCCGGTAAATGAGTTCTAGA GGGCCCGTTTAAACCCGCTGATCAGCCTCGACTGTGCCCTCTAGTTGCCAGCCATCTGTTGTTTG C  
TCTTCTCGGAGAGGGACAGAGGCCATTACTCAAGATCTCCCGGGCAAAATTTGGGCGACTAGTCGGAGCTGACACGGAAGATCAACGGTTCGGTAGACAACAAAC G  
Q K S L S L S P G K

[380] The framework 4 (FR4) region of the Gamma HC normally ends with two serines (SS), so that the full framework 4 region should be WGQGLTVTVSS (SEQ ID NO: 80). The

accepting Xho 1 site and one additional base downstream of the Xho1 site in the vector, in which the Gamma HC constant region that the Gamma HC variable regions are cloned, supplies the last bases, which encode this final amino acid of framework 4. However, the original vector did not adjust for the silent mutation made when the Xho1 site (CTCGAG, SEQ ID NO: 81) was created and contained an “A” nucleotide downstream of the Xho1 site, which caused an amino acid change at the end of framework 4: a serine to arginine (S to R) substitution present in all the working Gamma HC clones. Thus, the full framework 4 region reads WGQGTTLVTVSR (SEQ ID NO: 82). Future constructs are being created wherein the base downstream of the Xho 1 site is a “C” nucleotide. Thus, the creation of the Xho 1 site used for cloning of the Gamma HC variable region sequences in alternative embodiments is a silent mutation and restores the framework 4 amino acid sequence to its proper WGQGTTLVTVSS (SEQ ID NO: 80). This is true for all M2 Gamma HC clones described herein.

*Clone 21B15:*

[381] The Kappa LC variable region of the anti M2 clone 21B15 was cloned as Hind III to BsiW1 fragment, and is encoded by the following polynucleotide sequences and SEQ ID NO: 83 and SEQ ID NO: 84:

HindIII

```
AAGCTTCCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAGGTGC
TTCGAAGGTGGTACCTGTACTCCCAGGAGCGAGTCGAGGACCCCGAGGACGATGAGACCGAGGCTCCACG
CAGATGTGACATCCAGGTGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATC
GTCTACACTGTAGGTCCACTGGGTCAGAGGTAGGAGGGACAGACGTAGACATCCTCTGTCTCAGTGGTAG
ACTTGCCGCGCGAGTCAGAACATTTACAAGTATTTAAATTGGTATCAGCAGAGACCAGGGAAAGCCCCTA
TGAACGGCGCGCTCAGTCTTGTAATGTTTCATAAATTTAACCATAGTCGTCTCTGGTCCCTTTCGGGGAT
AGGGCCTGATCTCTGCTGCATCCGGGTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTGGCAGTGGATCTGG
TCCCGGACTAGAGACGACGTAGGCCCAACGTTTACCCCCAGGGTAGTTCCAAGTCAACGTCACCTAGAC C
GACAGATTTCACTCTCACCATCACCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCAACAGAGT
CTGTCTAAAGTGAGAGTGGTAGTGGTCAGACGTTGGACTTCTAAAACGTTGAATGATGACAGTTGTCTCA
```

BsiW1

```
TACAGTCCCCCTCTCACTTTCGGCGGAGGGACCAGGGTGGATATCAAACGTACG
ATGTCAGGGGGAGAGTGAAAGCCGCTCCCTGGTCCCACCTATAGTTTGCATGC
```

[382] The translation of the 21B15 Kappa LC variable region is as follows, polynucleotide sequence (above, SEQ ID NO: 83, top) and amino acid sequence (below, corresponding to SEQ ID NO: 56):

AAGCTTCCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAGGT  
M D M R V L A Q L L G L L L L W L R G  
GCCAGATGTGACATCCAGGTGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCAC C  
A R C D I Q V T Q S P S S L S A S V G D R V T  
ATCACTTGCCGCGCGAGTCAGAACATTTACAAGTATTTAAATTGGTATCAGCAGAGACCAGGGAAAGC C  
I T C R A S Q N I Y K Y L N W Y Q Q R P G K A  
CCTAAGGGCCTGATCTCTGCTGCATCCGGGTTGCAAAGTGCGGTCCCATCAAGGTTCAGTGGCAGTGG A  
P K G L I S A A S G L Q S G V P S R F S G S G  
TCTGGGACAGATTTCACTCTCACCATCACCAGTCTGCAACCTGAAGATTTTGCAACTTACTACTGTCA A  
S G T D F T L T I T S L Q P E D F A T Y Y C Q  
BsiWI  
CAGAGTTACAGTCCCCCTCTCACTTTTCGGCGGAGGGACCAGGGTGGATATCAAACGTACG  
Q S Y S P P L T F G G G T R V D I K R T

R T Start of Kappa constant region

**[385]** The 21B15 Gamma HC variable region was cloned as a Hind III to Xho I fragment, and is encoded by the following polynucleotide sequences and SEQ ID NO: 85 (top), and SEQ ID NO: 86 (bottom):

HindIII

AAGCTTCCACCATGAAACACCTGTGGTTCTTCCTTCTCCTGGTGGCAGCTCCCAGCTGGGTC C  
 TTCGAAGGTGGTACTTTGTGGACACCAAGAAGGAAGAGGACCACCGTCGAGGGTTCGACCCAG G  
 TGTCCCAGGTGCAATTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCC C  
 ACAGGGTCCACGTAAACGTCCTCAGCCCGGGTCTGACCACTTCGGAAGCCTCTGGGACAGG G  
 TCACCTGCACTGTCTCTGGTTTCGTCCATCAGTAATTACTACTGGAGCTGGATCCGGCAGTCC C  
 AGTGGACGTGACAGAGACCAAGCAGGTAGTCATTAATGATGACCTCGACCTAGGCCGTCAGG G  
 CAGGGAAGGGACTGGAGTGGATTGGGTTTATCTATTACGGTGGAAACACCAAGTACAATCCC T  
 GTCCCTTCCCTGACCTCACCTAACCCAAATAGATAATGCCACCTTTGTGGTTTCATGTTAGGG A  
 CCCTCAAGAGCCGCGTCACCATATCACAAGACACTTCCAAGAGTCAGGTCTCCCTGACGATG A  
 GGGAGTTCTCGGCGCAGTGGTATAGTGTTCTGTGAAGGTTCTCAGTCCAGAGGGACTGCTAC T  
 GCTCTGTGACCGCTGCGGAATCGGCCGTCTATTTCTGTGCGAGAGCGTCTTGTAAGTGGTGGT T  
 CGAGACACTGGCGACGCCTTAGCCGGCAGATAAAGACACGCTCTCGCAGAACATCACCACCA A

XhoI

ACTGTATCCTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCGAG  
 TGACATAGGAACTGATGACCCCGGTCCCTTGGGACCAGTGGCAGAGCTC

[386] The translation of the 21B15 Gamma HC is as follows, polynucleotide sequence

(above, SEQ ID NO: 87, top) and amino acid sequence (below, corresponding to SEQ ID

NO: 69):

HindIII

AAGCTTCCACCATGAAACACCTGTGGTTCTTCCTTCTCCTGGTGGCAGCTCCCAGCTGGGTC  
 M K H L W F F L L L V A A P S W V  
 CTGTCCCAGGTGCAATTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCC C  
 L S Q V Q L Q E S G P G L V K P S E T L S  
 CTCACCTGCACTGTCTCTGGTTTCGTCCATCAGTAATTACTACTGGAGCTGGATCCGGCAGTCC C  
 L T C T V S G S S I S N Y Y W S W I R Q S  
 CCAGGGAAGGGACTGGAGTGGATTGGGTTTATCTATTACGGTGGAAACACCAAGTACAATCC C  
 P G K G L E W I G F I Y Y G G N T K Y N P  
 TCCCTCAAGAGCCGCGTCACCATATCACAAGACACTTCCAAGAGTCAGGTCTCCCTGACGAT G  
 S L K S R V T I S Q D T S K S Q V S L T M  
 AGCTCTGTGACCGCTGCGGAATCGGCCGTCTATTTCTGTGCGAGAGCGTCTTGTAAGTGGTGG T  
 S S V T A A E S A V Y F C A R A S C S G G  
 TACTGTATCCTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCGAG  
 Y C I L D Y W G Q G T L V T V S

XhoI

[387] The amino acid sequence of the 21B15 Gamma HC is as follows, with specific domains identified below (CDR sequences defined according to Kabat methods):

MKHLWFFLLLVAAPSWVLS

VH leader (SEQ ID NO: 70)

QVQLQESGPGLVKPSETLSLTCTVSGSSIS

FR1 (SEQ ID NO: 71)

NYYWS

CDR1 (SEQ ID NO: 72)

WIRQSPGKGLEWIG

FR2 (SEQ ID NO: 73)

FIYYGGNTKYNPSLKS

CDR2 (SEQ ID NO: 74)

RVTISQDTSKSQVSLTMSSVTAAESAVYFCAR

FR3 (SEQ ID NO: 75)

ASCSGGYCILD

CDR3 (SEQ ID NO: 76)

YWGQGTLVTVS

FR4 (SEQ ID NO: 77)

*Clone 23K12:*

[388] The Kappa LC variable region of the anti M2 clone 23K12 was cloned as Hind III to BsiWI fragment (see below), and is encoded by the following polynucleotide sequences SEQ ID NO: 88 (top) and SEQ ID NO: 89 (below).

HindIII

AAGCTTCCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAG G  
 TTCGAAGGTGGTACCTGTACTCCCAGGAGCGAGTCGAGGACCCCGAGGACGATGAGACCGAGGCTC C  
 TGCCAGATGTGACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGT C  
 ACGGTCTACACTGTAGGTCTACTGGGTGAGAGGTAGGAGGGACAGACGTAGACATCCTCTGTCTCA G  
 ACCATCACTTGCCGGACAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGG A  
 TGGTAGTGAACGGCCTGTTTCAGTCTCGTAATCGTCGATAAATTTAACCATAGTCGTCTTTGGTCCC T  
 AAGCCCCCTAAACTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTG G  
 TTCGGGGATTGAGGACTAGATACGACGTAGGTCAAACGTTTCACCCAGGGTAGTTCCAAGTCAC C  
 CAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCGGTCTGCAACCTGAAGATTTTGCAACCTA C  
 GTCACCTAGACCCTGTCTAAAGTGAGAGTGGTAGTCGCCAGACGTTGGACTTCTAAAACGTTGGAT G  
 TACTGTCAACAGAGTTACAGTATGCCTGCCTTTGGCCAGGGGACCAAGCTGGAGATCAAACGTACG  
 ATGACAGTTGTCTCAATGTCATACGGACGGAAACCGGTCCCCTGGTTCGACCTCTAGTTTGCATGC

BsiWI

[389] The translation of the 23K12 Kappa LC variable region is as follows, polynucleotide sequence (above, SEQ ID NO: 90, top) and amino acid sequence (below, corresponding to SEQ ID NO: 91).

HindIII

AAGCTTCCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAG G  
 M D M R V L A Q L L G L L L L W L R G  
 TGCCAGATGTGACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGT C  
 A R C D I Q M T Q S P S S L S A S V G D R V  
 ACCATCACTTGCCGGACAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCAGGG A  
 T I T C R T S Q S I S S Y L N W Y Q Q K P G  
 AAGCCCCCTAAACTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGTTTCAGTG G  
 K A P K L L I Y A A S S L Q S G V P S R F S G  
 CAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCGGTCTGCAACCTGAAGATTTTGCAACCTA C  
 S G S G T D F T L T I S G L Q P E D F A T Y  
 TACTGTCAACAGAGTTACAGTATGCCTGCCTTTGGCCAGGGGACCAAGCTGGAGATCAAACGTACG  
 Y C Q Q S Y S M P A F G Q G T K L E I K R T

BsiWI

[390] The amino acid sequence of the 23K12 Kappa LC variable region is as follows, with specific domains identified below (CDR sequences defined according to Kabat methods):

|                            |                                   |
|----------------------------|-----------------------------------|
| MDMRVLAQLLGLLLLWLRGARC     | VK leader (SEQ ID NO: 57)         |
| DIQMTQSPSSLSASVGDRVTITC    | FR1 (SEQ ID NO: 58)               |
| RTSQSISSYLN                | CDR1 (SEQ ID NO: 92)              |
| WYQQKPGKAPKLLIY            | FR2 (SEQ ID NO: 93)               |
| AASSLQSGVPSRF              | CDR2 (SEQ ID NO: 94)              |
| SGSGSGTDFTLTISGLQPEDFATYYC | FR3 (SEQ ID NO: 95)               |
| QQSYMPA                    | CDR3 (SEQ ID NO: 96)              |
| FGQGTKLEIK                 | FR4 (SEQ ID NO: 114)              |
| RT                         | Start of Kappa LC constant region |

[391] The 23K12 Gamma HC variable region was cloned as a Hind III to Xho I fragment, and is encoded by the following polynucleotide sequences and SEQ ID NO: 97 (top) and SEQ ID NO: 98 (bottom).

HindIII

AAGCTTCCACCATGGAGTTGGGGCTGTGCTGGGTTTTCTTGTGCTATTTTAAAAGGTGTCCAGT  
 TTCGAAGGTGGTACCTCAACCCCGACACGACCCAAAAGGAACAACGATAAAATTTTCCACAGGTCA  
 GTGAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTCCAGCCTGGGGGGTCCCTGAGAATCTCC T  
 CACTCCACGTCGACCACCTCAGACCCCTCCGAACCAGGTCCGACCCCCCAGGGACTCTTAGAGGA  
 GTGCAGCCTCTGGATTACCGTCAGTAGCAACTACATGAGTTGGGTCCGCCAGGCTCCAGGGAAG G  
 CACGTCGGAGACCTAAGTGGCAGTCATCGTTGATGTACTCAACCCAGGCGGTCCGAGGTCCCTTC C  
 GGCTGGAGTGGGTCTCAGTTATTTATAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGC A  
 CCGACCTCACCCAGAGTCAATAAATATCACCACCATCGTGTATGATGCGTCTGAGGCACTTCCCG T  
 GATTCTCCTTCTCCAGAGACAACCTCCAAGAACACAGTGTTTCTTCAAATGAACAGCCTGAGAGCC G  
 CTAAGAGGAAGAGGTCTCTGTTGAGGTTCTTGTGTACAAAGAAGTTTACTTGTGCGGACTCTCGG C  
 AGGACACGGCTGTGTATTACTGTGCGAGATGTCTGAGCAGGATGCGGGGTTACGGTTTAGACGTC T  
 TCCTGTGCCGACACATAATGACACGCTCTACAGACTCGTCCTACGCCCCAATGCCAAATCTGCAG A

XhoI

GGGGCCAAGGGACACGGTCACCGTCTCGAG  
 CCGCGGTTCCCTGGTGCCAGTGGCAGAGCTC

[392] The translation of the 23K12 Gamma HC variable region is as follows, polynucleotide sequence (above, SEQ ID NO: 99, top), and amino acid sequence (below, corresponding to SEQ ID NO: 100):

HindIII  
AAGCTTCCACCATGGAGTTGGGGCTGTGCTGGGTTTTCTTGTGTGCTATTTTAAAAGGTGTCCAG  
M E L G L C W V F L V A I L K G V Q  
TGTGAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTCCAGCCTGGGGGGTCCCTGAGAATCTC C  
C E V Q L V E S G G G L V Q P G G S L R I S  
TGTGCAGCCTCTGGATTACCGTCAGTAGCAACTACATGAGTTGGGTCCGCCAGGCTCCAGGGAAG  
C A A S G F T V S S N Y M S W V R Q A P G K  
GGGCTGGAGTGGGTCTCAGTTATTTATAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGG C  
G L E W V S V I Y S G G S T Y Y A D S V K G  
AGATTCTCCTTCTCCAGAGACAACCTCCAAGAACACAGTGTTTCTTCAAATGAACAGCCTGAGAGC C  
R F S F S R D N S K N T V F L Q M N S L R A  
GAGGACACGGCTGTGTATTACTGTGCGAGATGTCTGAGCAGGATGCGGGGTTACGGTTTAGACGT C  
E D T A V Y Y C A R C L S R M R G Y G L D V  
XhoI  
TGGGGCCAAGGGACCACGGTCACCGTCTCGAG  
W G Q G T T V T V S

[393] The amino acid sequence of the 23K12 Gamma HC variable region is as follows, with specific domains identified below (CDR sequences defined according to Kabat methods):

|                                   |                            |
|-----------------------------------|----------------------------|
| MELGLCWVFLVAILKGVQC               | VH leader (SEQ ID NO: 101) |
| EVQLVESGGGLVQPGGSLRISCAASGFTVS    | FR1 (SEQ ID NO: 102)       |
| SNYMS                             | CDR1 (SEQ ID NO: 103)      |
| WVRQAPGKGLEWVS                    | FR2 (SEQ ID NO: 104)       |
| VIYSGGSTYYADSVK                   | CDR2 (SEQ ID NO: 105)      |
| GRFSFSRDNSKNTVFLQMNSLRAEDTAVYYCAR | FR3 (SEQ ID NO: 106)       |
| CLSRMRGYGLDV                      | CDR3 (SEQ ID NO: 107)      |
| WGQGTTVTVS                        | FR4 (SEQ ID NO: 108)       |

#### Example 3: Identification of Conserved Antibody Variable Regions

[394] The amino acid sequences of the three antibody Kappa LC and Gamma HC variable regions were aligned to identify conserved regions and residues, as shown below.

**[395]** Amino acid sequence alignment of the Kappa LC variable regions of the three clones:

|                              |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |
|------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Translation of mp 73 21B15_F | A | S | M | M | V | L | A | L | L | G | L | L | L | L | L | L | G | A | C | I | V | S | S | P | L |
| Translation of mp 147 8i10 k | A | S | M | M | V | L | A | L | L | G | L | L | L | L | L | L | G | A | C | I | M | S | S | P | L |
| Translation of mp 137 23K12  | A | S | M | M | V | L | A | L | L | G | L | L | L | L | L | L | G | A | C | I | M | S | S | P | L |

|                              |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |
|------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Translation of mp 73 21B15_F | A | V | G | V | I | C | A | S | I | L | P | S | G | A | G | L |
| Translation of mp 147 8i10 k | A | V | G | V | I | C | A | S | I | L | P | S | G | A | G | L |
| Translation of mp 137 23K12  | A | V | G | V | I | C | A | S | I | L | P | S | G | A | G | L |

|                              |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |
|------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Translation of mp 73 21B15_F | I | A | A | S | G | L | G | V | F | S | G | G | G | I | T | L | I | S | L | L |
| Translation of mp 147 8i10 k | I | A | A | S | G | L | G | V | F | S | G | G | G | I | T | L | I | S | L | L |
| Translation of mp 137 23K12  | I | A | A | S | G | L | G | V | F | S | G | G | G | I | T | L | I | S | L | L |

|                              |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |
|------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Translation of mp 73 21B15_F | A | T | C | S | S | L | M | P | A | G | G | I | V | I | T |
| Translation of mp 147 8i10 k | A | T | C | S | S | L | M | P | A | G | G | I | V | I | T |
| Translation of mp 137 23K12  | A | T | C | S | S | L | M | P | A | G | G | I | V | I | T |

**[396]** Amino acid sequence alignment of the Gamma HC variable regions of the three clones:

|                               |    |   |   |   |  |   |   |   |   |   |   |   |   |   |   |   |   |   |    |   |   |   |   |   |   |   |  |  |  |  |  |  |  |  |  |  |
|-------------------------------|----|---|---|---|--|---|---|---|---|---|---|---|---|---|---|---|---|---|----|---|---|---|---|---|---|---|--|--|--|--|--|--|--|--|--|--|
|                               | 10 |   |   |   |  |   |   |   |   |   |   |   |   |   |   |   |   |   | 20 |   |   |   |   |   |   |   |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 81 21B15_F2 | A  | S | T | M |  | L | G | L | C | V | L | L | L | V | A | A | F | S | V  | L | S | V | L | A | S | V |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 145 23K12_F | A  | S | T | M |  | L | G | L | C | V | V | L | L | V | A | I | L | G | V  | L | C | V | L | V | S | V |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 153 8i10 ga | A  | S | T | M |  | L | G | L | C | V | L | L | L | V | A | A | F | S | V  | L | S | V | L | A | S | V |  |  |  |  |  |  |  |  |  |  |

|                               |    |   |   |   |   |  |   |   |   |   |   |   |   |   |   |   |   |   |    |   |   |   |   |   |   |   |   |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|-------------------------------|----|---|---|---|---|--|---|---|---|---|---|---|---|---|---|---|---|---|----|---|---|---|---|---|---|---|---|--|--|--|--|--|--|--|--|--|----|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
|                               | 30 |   |   |   |   |  |   |   |   |   |   |   |   |   |   |   |   |   | 40 |   |   |   |   |   |   |   |   |  |  |  |  |  |  |  |  |  | 50 |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 81 21B15_F2 | G  | P | G | L | V |  | P | S | T | L | S | L | T | C | T | V | S | G | S  | T | I | S | M | S | V | S | M |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 145 23K12_F | G  | G | G | L | V |  | P | G | G | S | L | I | S | C | A | A | G | T | T  | V | S | S | M | M | S | M |   |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 153 8i10 ga | G  | P | G | L | V |  | P | S | T | L | S | L | T | C | T | V | S | G | S  | T | I | S | M | S | V | S | M |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

|                               |    |  |   |   |   |  |   |   |   |   |   |   |   |   |   |   |   |  |    |   |   |   |  |  |  |   |  |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|-------------------------------|----|--|---|---|---|--|---|---|---|---|---|---|---|---|---|---|---|--|----|---|---|---|--|--|--|---|--|--|--|--|--|--|--|--|--|--|----|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
|                               | 60 |  |   |   |   |  |   |   |   |   |   |   |   |   |   |   |   |  | 70 |   |   |   |  |  |  |   |  |  |  |  |  |  |  |  |  |  | 80 |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 81 21B15_F2 | I  |  | S | T | G |  | G | L | M | I | G | T | I | Y | G | G | T |  | Y  | T | P | L |  |  |  |   |  |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 145 23K12_F | V  |  | A | P | G |  | G | L | M | V | V | I | Y | G | G | T |   |  | A  | S | V |   |  |  |  | G |  |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 153 8i10 ga | I  |  | S | T | G |  | G | L | M | I | G | T | I | Y | G | G | T |  |    | P | L |   |  |  |  | S |  |  |  |  |  |  |  |  |  |  |    |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

|                               |    |   |   |   |   |   |   |  |  |   |   |   |   |   |   |   |   |   |     |   |   |  |   |   |   |   |   |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|-------------------------------|----|---|---|---|---|---|---|--|--|---|---|---|---|---|---|---|---|---|-----|---|---|--|---|---|---|---|---|--|--|--|--|--|--|--|--|--|-----|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
|                               | 90 |   |   |   |   |   |   |  |  |   |   |   |   |   |   |   |   |   | 100 |   |   |  |   |   |   |   |   |  |  |  |  |  |  |  |  |  | 110 |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 81 21B15_F2 |    | V | T | I | S | T | S |  |  | S | V | S | L | T | M | S | S | V | T   | A | A |  |   | A | V | T | I |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 145 23K12_F |    | T | T | S |   | T | S |  |  | T | V | F | L | S | M | S | L |   | A   |   |   |  | T | A | V | T | I |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 153 8i10 ga |    | V | T | I | S | T | S |  |  | S | V | S | L | T | M | S | S | V | T   | A | A |  |   | A | V | T | I |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

|                               |     |   |  |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |     |   |   |   |   |   |  |  |  |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|-------------------------------|-----|---|--|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|-----|---|---|---|---|---|--|--|--|--|--|--|--|--|--|--|--|--|-----|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
|                               | 120 |   |  |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   | 130 |   |   |   |   |   |  |  |  |  |  |  |  |  |  |  |  |  | 140 |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 81 21B15_F2 | C   | A |  | A |   | C | S | G | G |   | C | I | L |   | M | G |   | G | T   | L | V | T | V | S |  |  |  |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 145 23K12_F | C   | A |  | C | L |   | M | G |   | G | L |   | V | M | G |   | G | T | T   | V | T | V | S |   |  |  |  |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Translation of mp 153 8i10 ga | C   | A |  | A |   | C | S | G | G |   | C | I | L |   | M | G |   | G | T   | L | V | T | V | S |  |  |  |  |  |  |  |  |  |  |  |  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

**[397]** Clones 8i10 and 21B15 came from two different donors, yet they have the same exact Gamma HC and differ in the Kappa LC by only one amino acid at position 4 in the framework 1 region (amino acids M versus V, see above), (excluding the D versus E wobble position in framework 4 of the Kappa LC).

**[398]** Sequence comparisons of the variable regions of the antibodies revealed that the heavy chain of clone 8i10 was derived from germline sequence IgHV4 and that the light chain was derived from the germline sequence IgKV1.

**[399]** Sequence comparisons of the variable regions of the antibodies revealed that the heavy chain of clone 21B15 was derived from germline sequence IgHV4 and that the light chain was derived from the germline sequence IgKV1.

[400] Sequence comparisons of the variable regions of the antibodies revealed that the heavy chain of clone 23K12 was derived from germline sequence IgHV3 and that the light chain was derived from the germline sequence IgKV1.

Example 4: Production and characterization of M2 Antibodies

[401] The antibodies described above were produced in milligram quantities by larger scale transient transfections in 293 PEAK cells. Crude un-purified antibody supernatants were used to examine antibody binding to influenza A/Puerto Rico/8/1932 (PR8) virus on ELISA plates, and were compared to the binding of the control antibody 14C2, which was also produced by larger scale transient transfection. The anti-M2 recombinant human monoclonal antibodies bound to influenza while the control antibody did not (Figure 9).

[402] Binding was also tested on MDCK cells infected with the PR8 virus (Figure 10). The control antibody 14C2 and the three anti M2E clones: 8I10, 21B15 and 23K12, all showed specific binding to the M2 protein expressed on the surface of PR8-infected cells. No binding was observed on uninfected cells.

[403] The antibodies were purified over protein A columns from the supernatants. FACs analysis was performed using purified antibodies at a concentration of 1 ug per ml to examine the binding of the antibodies to transiently transfected 293 PEAK cells expressing the M2 proteins on the cell surface. Binding was measured testing binding to mock transfected cells and cells transiently transfected with the Influenza subtype H3N2, A/Vietnam/1203/2004 (VN1203), or A/Hong Kong/483/1997 HK483 M2 proteins. As a positive control the antibody 14C2 was used. Unstained and secondary antibody alone controls helped determined background. Specific staining for cells transfected with the M2 protein was observed for all three clones. Furthermore, all three clones bound to the high path strains A/Vietnam/1203/2004 and A/Hong Kong/483/1997 M2 proteins very well, whereas the positive control 14C2 which bound well to H3N2 M2 protein, bound much weaker to the A/Vietnam/1203/2004 M2 protein and did not bind the A/Hong Kong/483/1997 M2 protein. See Figure 11.

[404] Antibodies 21B15, 23K12, and 8I10 bound to the surface of 293-HEK cells stably expressing the M2 protein, but not to vector transfected cells (see Figure 1). In addition, binding of these antibodies was not competed by the presence of 5 mg/ml 24-mer M2 peptide, whereas the binding of the control chimeric mouse V-region/human IgG1 kappa 14C2 antibody (hu14C2) generated against the linear M2 peptide was completely inhibited by the M2 peptide (see Figure 1). These data confirm that these antibodies bind to

conformational epitopes present in M2e expressed on the cell or virus surface, as opposed to the linear M2e peptide.

Example 5: Viral Binding of human anti-influenza monoclonal antibodies

[405] UV-inactivated influenza A virus (A/PR/8/34) (Applied Biotechnologies) was plated in 384-well MaxiSorp plates (Nunc) at 1.2 µg/ml in PBS, with 25 µl/well, and was incubated at 4°C overnight. The plates were then washed three times with PBS, and blocked with 1% Nonfat dry milk in PBS, 50 µl/well, and then were incubated at room temp for 1 hr. After a second wash with PBS, MAbs were added at the indicated concentrations in triplicate, and the plates were incubated at room temp for 1 hour. After another wash with PBS, to each well was added 25 µl of a 1/5000 dilution of horseradish peroxidase (HRP) conjugated goat anti-human IgG Fc (Pierce) in PBS/1% Milk, and the plates were left at room temp for 1 hr. After the final PBS wash, the HRP substrate 1-Step™ Ultra-TMB-ELISA (Pierce) was added at 25 µl/well, and the reaction proceeded in the dark at room temp. The assay was stopped with 25 µl/well 1N H<sub>2</sub>SO<sub>4</sub>, and light absorbance at 450 nm (A<sub>450</sub>) was read on a SpectroMax Plus plate reader. Data are normalized to the absorbance of MAb 8I10 binding at 10 µg/ml. Results are shown in Figures 2A and 2B.

Example 6: Binding of Human Anti-Influenza Monoclonal Antibodies to Full-Length M2 Variants

[406] M2 variants (including those with a high pathology phenotype in vivo) were selected for analysis. See Figure 3A for sequences.

[407] M2 cDNA constructs were transiently transfected in HEK293 cells and analyzed as follows: To analyze the transient transfectants by FACS, cells on 10 cm tissue culture plates were treated with 0.5 ml Cell Dissociation Buffer (Invitrogen), and harvested. Cells were washed in PBS containing 1% FBS, 0.2% NaN<sub>3</sub> (FACS buffer), and resuspended in 0.6 ml FACS buffer supplemented with 100 µg/ml rabbit IgG. Each transfectant was mixed with the indicated MAbs at 1 µg/ml in 0.2 ml FACS buffer, with 5 x 10<sup>5</sup> to 10<sup>6</sup> cells per sample. Cells were washed three times with FACS buffer, and each sample was resuspended in 0.1 ml containing 1 µg/ml alexafluor (AF) 647-anti human IgG H&L (Invitrogen). Cells were again washed and flow cytometry was performed on a FACSCanto device (Becton-Dickenson). The data is expressed as a percentage of the mean fluorescence of the M2-D20 transient transfectant. Data for variant binding are representative of 2 experiments. Data for alanine

mutants are average readouts from 3 separate experiments with standard error. Results are shown in Figure 3B and 3C.

#### Example 7: Alanine Scanning Mutagenesis to Evaluate M2 Binding

[408] To evaluate the antibody binding sites, alanine was substituted at individual amino acid positions as indicated by site-directed mutagenesis.

[409] M2 cDNA constructs were transiently transfected in HEK293 cells and analyzed as described above in Example 6. Results are shown in Figure 4A and 4B. Figure 8 shows that the epitope is in a highly conserved region of the amino terminus of the M2 polypeptide. As shown in Figures 4A, 4B and Figure 8, the epitope includes the serine at position 2, the threonine at position 5 and the glutamic acid at position 6 of the M2 polypeptide.

#### Example 8: Epitope Blocking

[410] To determine whether the MAbs 8I10 and 23K12 bind to the same site, M2 protein representing influenza strain A/HK/483/1997 sequence was stably expressed in the CHO (Chinese Hamster Ovary) cell line DG44. Cells were treated with Cell Dissociation Buffer (Invitrogen), and harvested. Cells were washed in PBS containing 1% FBS, 0.2% NaN<sub>3</sub> (FACS buffer), and resuspended at 10<sup>7</sup> cells/ml in FACS buffer supplemented with 100 µg/ml rabbit IgG. The cells were pre-bound by either MAb (or the 2N9 control) at 10 µg/ml for 1 hr at 4 °C, and were then washed with FACS buffer. Directly conjugated AF647-8I10 or -23K12 (labeled with the AlexaFluor® 647 Protein Labeling kit (Invitrogen)) was then used to stain the three pre-blocked cell samples at 1 µg/ml for 10<sup>6</sup> cells per sample. Flow cytometric analyses proceeded as before with the FACSCanto. Data are average readouts from 3 separate experiments with standard error. Results are shown in Figure 5.

#### Example 9: Binding of human anti-influenza monoclonal antibodies to M2 Variants and Truncated M2 Peptides

[411] The cross reactivity of mAbs 8i10 and 23K12 to other M2 peptide variants was assessed by ELISA. Peptide sequences are shown in Figures 6A and 6B. Additionally, a similar ELISA assay was used to determine binding activity to M2 truncated peptides.

[412] In brief, each flat bottom 384 well plate (Nunc) was coated with a concentration of 2 µg/mL peptide and 25 µL/ well of PBS buffer overnight at 4°C. Plates were washed three times and blocked with 1% Milk/PBS for one hour at room temperature. After washing three times, MAb titers were added and incubated for one hour at room temperature. Diluted HRP

conjugated goat anti-human immunoglobulin FC specific (Pierce) was added to each well after washing three times. Plates were incubated for one hour at room temperature and washed three times. 1-Step™ Ultra-TMB-ELISA (Pierce) was added at 25 µl/well, and the reaction proceeded in the dark at room temp. The assay was stopped with 25 µl/well 1N H<sub>2</sub>SO<sub>4</sub>, and light absorbance at 450 nm (A450) was read on a SpectroMax Plus plate reader. Results are shown in Figures 6A and 6B.

Example 10: In Vivo Evaluation of the Ability of Human Anti-Influenza Monoclonal Antibodies to Protect From Lethal Viral Challenge

[413] The ability of antibodies, 23K12 and 8I10, to protect mice from lethal viral challenge with a high path avian influenza strain was tested.

[414] Female BALB/c mice were randomized into 5 groups of 10. One day prior (Day -1 (minus one)) and two days post infection (Day +2 (plus two)), 200 ug of antibody was given via 200 µl intra-peritoneal injection. On Day 0 (zero), an approximate LD90 (lethal dose 90) of A/Vietnam/1203/04 influenza virus, in a volume of 30 µl was given intra-nasally. Survival rate was observed from Day 1 through Day 28 post-infection. Results are shown in Figure 7.

Example 11 : Characterization of M2 Antibodies 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, and 3242\_P05

[415] FACS

[416] Full length M2 cDNA (A/Hong Kong/483/97) were synthesized (Blue Heron Technology) and cloned into the plasmid vector pcDNA3.1 which was then transfected into CHO cells with Lipofectamine (Invitrogen) to create a stable pool of CHO-HK M2-expressing cells. For the panel of anti-M2 Mabs, 20 µl samples of supernatant from transient transfections from each of the IgG heavy and light chain combinations was used to stain the CHO-HK M2 stable cell line. Bound anti-M2 mabs were visualized on viable cells with Alexafluor 647-conjugated goat anti-Human IgG H&L antibody (Invitrogen). Flow cytometry was performed with a FACSCanto, and analysis on the accompanying FACSDiva software (Becton Dickenson).

[417] ELISA

[418] Purified Influenza A (A/Puerto Rico/8/34) inactivated by β-propiolactone (Advanced Biotechnologies, Inc.) was biotinylated (EZ-Link Sulfo-NHS-LC-Biotin, Pierce) and adsorbed for 16 hours at 4°C to 384-well plates in 25 µl PBS that were pre-coated with

neutravidin (Pierce). Plates were blocked with BSA in PBS, samples of supernatant from transient transfections from each of the IgG heavy and light chain combinations were added at a final dilution of 1:5, followed by HRP-conjugated goat anti-human Fc antibody (Pierce), and developed with TMB substrate (ThermoFisher).

[419] The results of this analyses are shown below in Table 2

| Transfection no. | BCC well ID | Sequence ID   |               | FACS         | Virus                        |
|------------------|-------------|---------------|---------------|--------------|------------------------------|
|                  |             | Gamma         | Light         | M2-HK<br>MFI | ELISA<br>OD A <sub>450</sub> |
| 322              | 3241_G23    | G4_005        | K1_004        | 1697         | 3.02                         |
| 352              | 3244_I10    | G4_007        | K2_006        | 434          | 3.01                         |
| 339              | 3243_J07    | G4_007        | K1_007        | 131          | 2.94                         |
| 336              | 3259_J21    | G4_005        | K2_005        | 1673         | 2.40                         |
| 348              | 3245_O19    | G3_004        | K1_001        | 919          | 3.51                         |
| 345              | 3244_H04    | G3_003        | K1_006        | 963          | 3.31                         |
| 346              |             | Pos Cont (HC) | Pos Cont (LC) | 754          | 2.69                         |
| 347              |             | Neg Cont (HC) | Neg Cont (LC) | 11           | 0.15                         |
| 374              | 3136_G05    | G4_007        | K1_007        | 109          | ND                           |
| 386              | 3252_C13    | G4_013        | K1_002        | 449          | ND                           |
| 390              | 3255_J06    | G4_013        | K2_007        | 442          | ND                           |
| 400              | 3420_I23    | G4_004        | K1_003        | 112          | ND                           |
| 432              | 3139_P23    | G4_016        | K1_007a       | 110          | 1.02                         |
| 412              | 3248_P18    | G4_009        | K1_006        | 967          | 0.56                         |
| 413              | 3253_P10    | G4_007        | K1_004        | 43           | 0.50                         |
| 434              | 3260_D19    | G3_004a       | K2_001        | 846          | 2.46                         |
| 439              | 3362_B11    | G4_010a       | K1_007        | 218          | 1.83                         |
| 408              | 3242_P05    | G3_005        | K2_004        | 596          | 0.50                         |
| 451              |             | Pos Cont (HC) | Pos Cont (LC) | 1083         | 1.87                         |
| 452              |             | Neg Cont (HC) | Neg Cont (LC) | 6            | 0.05                         |

Positive control: supernatant from transient transfection with the IgG heavy and light chain combination of mAb 8I10

Negative control: supernatant from transient transfection with the IgG heavy and light chain combination of mAb 2N9

MFI= mean fluorescence intensity

## OTHER EMBODIMENTS

[420] Although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

[421] While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

[422] The patent and scientific literature referred to herein establishes the knowledge that is available to those with skill in the art. All United States patents and published or unpublished United States patent applications cited herein are incorporated by reference. All published foreign patents and patent applications cited herein are hereby incorporated by reference. Genbank and NCBI submissions indicated by accession number cited herein are hereby incorporated by reference. All other published references, documents, manuscripts and scientific literature cited herein are hereby incorporated by reference.

[423] While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

## CLAIMS

What is claimed is:

1. A human monoclonal antibody that specifically binds to an epitope in the extracellular domain of the matrix 2 ectodomain (M2e) polypeptide of an influenza virus wherein said monoclonal antibody is 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, or 3242\_P05.
2. An isolated anti-matrix 2 ectodomain (M2e) antibody, or antigen-binding fragment thereof, comprising a heavy chain variable (VH) domain and a light chain variable (VL) domain, wherein the VH domain and the VL domain each comprise three complementarity determining regions 1 to 3 (CDR1-3), and wherein each CDR comprises the following amino acid sequences:  
VH CDR1: SEQ ID NOs: 179, 187, 196, 204, 212, 224, 230, 235, 242, 248, or 254;  
VH CDR2: SEQ ID NOs: 180, 188, 195, 197, 205, 213, 218, 225, 231, 236, 243, 249, 246, or 256;  
VH CDR3 SEQ ID NOs: 181, 189, 198, 206, 214, 219, 226, 232, 237, 244, or 250;  
VL CDR1: SEQ ID NOs: 184, 192, 199, 215, 220, 233, or 238;  
VL CDR2: SEQ ID NOs: 61, 185, 193, 200, 207, 211, 216, 227, 239, or 241; and  
VL CDR3: SEQ ID NOs: 63, 186, 194, 201, 208, 221, 228, 234, 240, 245, or 251.
3. An isolated anti-matrix 2 ectodomain (M2e) antibody, or antigen-binding fragment thereof, comprising a heavy chain variable (VH) domain and a light chain variable (VL) domain, wherein the VH domain and the VL domain each comprise three complementarity determining regions 1 to 3 (CDR1-3), and wherein each CDR comprises the following amino acid sequences:  
VH CDR1: SEQ ID NOs: 182, 190, 202, 209, 222, 229, 247, 252, 257, 258, or 260;  
VH CDR2: SEQ ID NOs: 183, 191, 203, 210, 217, 223, 230, 246, 253, 259, or 261;  
VH CDR3 SEQ ID NOs: 181, 189, 195, 198, 206, 214, 219, 226, 232, 237, 244, or 250;  
VL CDR1: SEQ ID NOs: 184, 192, 199, 215, 220, 233, or 238;  
VL CDR2: SEQ ID NOs: 61, 185, 193, 200, 207, 211, 216, 227, 239, or 241; and

VL CDR3: SEQ ID NOs: 63, 186, 194, 201, 208, 221, 228, 234, 240, 245, or 251.

4. A pharmaceutical composition comprising any one of the antibodies according to any of claims 1-3 and a pharmaceutical carrier.

5. An antibody that binds the same epitope as monoclonal antibody 3241\_G23, 3244\_I10, 3243\_J07, 3259\_J21, 3245\_O19, 3244\_H04, 3136\_G05, 3252\_C13, 3255\_J06, 3420\_I23, 3139\_P23, 3248\_P18, 3253\_P10, 3260\_D19, 3362\_B11, or 3242\_P05.

6. The composition of claim 4, further comprising an anti-viral drug, a viral entry inhibitor or a viral attachment inhibitor.

7. The composition of claim 6, wherein said anti-viral drug is a neuraminidase inhibitor, a HA inhibitor, a sialic acid inhibitor or an M2 ion channel inhibitor.

8. The composition of claim 7, wherein said M2 ion channel inhibitor is amantadine or rimantadine.

9. The composition of claim 7, wherein said neuraminidase inhibitor zanamivir, or oseltamivir phosphate.

10. The composition of claim 4, further comprising a second anti-influenza A antibody.

11. A method for stimulating an immune response in a subject, comprising administering to the patient the composition of claim 4.

12. A method for the treatment or prevention of an influenza virus infection in a subject, comprising administering to the subject the composition of claim 4.

13. The method of claim 12, wherein the method further comprises administering an anti-viral drug, a viral entry inhibitor or a viral attachment inhibitor.
14. The method of claim 13, wherein said anti-viral drug is a neuraminidase inhibitor, a HA inhibitor, a sialic acid inhibitor or an M2 ion channel.
15. The method of claim 14, wherein said M2 ion channel inhibitor is amantadine or rimantadine.
16. The method of claim 12, wherein said neuraminidase inhibitor zanamivir, or oseltamivir phosphate.
17. The method of claim 12, wherein said composition is administered prior to or after exposure to influenza virus.
18. The method of claim 12, wherein said composition is administered at a dose sufficient to promote viral clearance or eliminate influenza A infected cells.
19. A method for determining the presence of a influenza virus infection in a patient, comprising the steps of:
  - (a) contacting a biological sample obtained from the patient with the antibody according any one of claims 1-3;
  - (b) detecting an amount of the antibody that binds to the biological sample; and
  - (c) comparing the amount of antibody that binds to the biological sample to a control value, and therefrom determining the presence of the influenza virus in the patient.
20. A diagnostic kit comprising the antibody according to any one of claims 1-3.

1/17




Fig. 1

2/17




Fig. 2A

3/17

VIRAL BINDING (with Z3G1)  
INFLUENZA A/PR/8/32 ELISA




Fig. 2B

4/17

## FULL-LENGTH M2 VARIANT BINDING

## AMINO ACID SEQUENCES OF EXTRACELLULAR DOMAINS OF M2 VARIANTS.

|    |                                     | 10 |   |   |   |   |   |   |   |   |   | 20 |   |   |   |   |   |   |   |   |   |   |   |   |   |
|----|-------------------------------------|----|---|---|---|---|---|---|---|---|---|----|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1  | A.Brevig.Mission.1.1918.H1N1        | M  | S | L | L | T | E | V | E | T | P | T  | R | N | E | W | G | C | R | C | N | D | S | S | D |
| 2  | A.Fort.Monmouth.1.1947.H1N1         | M  | S | L | L | T | E | V | E | T | P | T  | K | N | E | W | E | C | R | C | N | D | S | S | D |
| 3  | A.Singapore.02.2005.H3N2            | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | E | C | R | C | N | D | S | S | D |
| 4  | A.Wisconsin.10.98.H1N1              | M  | S | L | L | T | E | V | E | T | P | I  | K | N | G | W | E | C | K | C | N | D | S | S | D |
| 5  | A.Wisconsin.301.1976.H1N1           | M  | S | L | L | T | E | V | E | T | P | I  | R | S | E | W | G | C | R | C | N | D | S | S | D |
| 6  | A.Panama.1.66.H2N2                  | M  | S | F | L | P | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | D | S | S | D |
| 7  | A.New York.321.1999.H3N2            | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | D | S | S | N |
| 8  | A.Caracas.1.71.H3N2                 | M  | S | L | L | T | E | V | E | T | P | I  | R | K | E | W | G | C | R | C | N | D | S | S | D |
| 9  | A.Taiwan.3.71.H3N2                  | M  | S | F | L | T | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | D | S | S | D |
| 10 | A.Wuhan.359.95.H3N2                 | M  | S | L | P | T | E | V | E | T | P | I  | R | S | E | W | G | C | R | C | N | D | S | S | D |
| 11 | A.Hong Kong.1144.99.H3N2            | M  | S | L | L | P | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | D | S | S | D |
| 12 | A.Hong Kong.1180.99.H3N2            | M  | S | L | L | P | E | V | E | T | P | I  | R | N | G | W | G | C | R | C | N | D | S | S | D |
| 13 | A.Hong Kong.1774.99.H3N2            | M  | S | L | L | T | E | V | E | T | P | T  | R | N | G | W | E | C | R | C | S | G | S | S | D |
| 14 | A.New York.217.02.H1N2              | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | E | Y | R | C | N | D | S | S | D |
| 15 | A.New York.300.2003.H1N2            | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | E | Y | R | C | S | D | S | S | D |
| 16 | A.swine.Spain.54008.2004.H3N2       | M  | S | L | L | T | E | V | E | T | P | T  | R | N | G | W | E | C | R | Y | S | D | S | S | D |
| 17 | A.Guangzhou.333.99.H9N2             | M  | S | F | L | T | E | V | E | T | L | T  | R | N | G | W | E | C | R | C | S | D | S | S | D |
| 18 | A.Hong Kong.1073.99.H9N2            | M  | S | L | L | T | E | V | E | T | L | T  | R | N | G | W | E | C | K | C | R | D | S | S | D |
| 19 | A.Hong Kong.1.68.H3N2               | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | D | S | S | D |
| 20 | A.swine.Hong Kong.126.1982.H3N2     | M  | S | L | L | T | E | V | E | T | P | I  | R | S | E | W | G | C | R | C | N | D | S | G | D |
| 21 | A.New York.703.1995.H3N2            | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | E | C | R | C | N | G | S | S | D |
| 22 | A.swine.Quebec.192.81.H1N1          | M  | S | L | P | T | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | D | S | S | D |
| 23 | A.Puerto Rico.8.34.H1N1             | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | G | S | S | D |
| 24 | A.Hong Kong.485.97.H5N1             | M  | S | L | L | T | E | V | D | T | L | T  | R | N | G | W | G | C | R | C | S | D | S | S | D |
| 25 | A.Hong Kong.542.97.H5N1             | M  | S | L | L | T | E | V | E | T | L | T  | K | N | G | W | G | C | R | C | S | D | S | S | D |
| 26 | A.silky chicken.Shantou.1826.2004.H | M  | S | L | L | T | E | V | E | T | P | T  | R | N | G | W | E | C | K | C | S | D | S | S | D |
| 27 | A.chicken.Taiwan.0305.04.H6N1       | M  | S | L | L | T | E | V | E | T | H | T  | R | N | G | W | E | C | K | C | S | D | S | S | D |
| 28 | A.Quail.Arkansas.16309-7.94.H7N3    | M  | S | L | L | T | E | V | K | T | P | T  | R | N | G | W | E | C | K | C | S | D | S | S | D |
| 29 | A.Hong Kong.486.97.H5N1             | M  | S | L | L | T | E | V | E | T | L | T  | R | N | G | W | G | C | R | C | S | D | S | S | D |
| 30 | A.Chicken.Pennsylvania.13552-1.98   | M  | S | L | L | T | E | V | E | T | P | T  | R | D | G | W | E | C | K | C | S | D | S | S | D |
| 31 | A.chicken.Heilongjiang.48.01.H9N2   | M  | S | L | L | T | E | V | E | T | P | T  | R | N | G | W | G | C | R | C | S | D | S | S | D |
| 32 | A.swine.Korea.S5.2005.H1N2          | M  | S | L | L | T | E | V | E | T | P | T  | R | N | G | W | E | C | K | C | N | D | S | S | D |
| 33 | A.Hong Kong.1073.99.H9N2            | M  | S | L | L | T | E | V | E | T | L | T  | R | N | G | W | E | C | K | C | S | D | S | S | D |
| 34 | A.Wisconsin.3523.88.H1N1            | M  | S | L | L | T | E | V | E | T | P | I  | R | N | E | W | G | C | K | C | N | D | S | S | D |
| 35 | A.X-31 Vaccine strain H3N2          | M  | S | F | L | T | E | V | E | T | P | I  | R | N | E | W | G | C | R | C | N | G | S | S | D |
| 36 | A.Chicken.Rostock.8.1934.H7N1       | M  | S | L | L | T | E | V | E | T | P | T  | R | N | G | W | E | C | R | C | N | D | S | S | D |
| 37 | A.environment.New York.16326-1.2    | M  | S | L | L | T | E | V | E | T | P | I  | R | K | G | W | E | C | N | C | S | D | S | S | D |
| 38 | A.Indonesia.560H.2006.H5N1          | M  | S | L | L | T | E | V | E | T | P | T  | R | N | E | W | E | C | R | C | S | D | S | S | D |
| 39 | A.Chicken.Hong Kong.SF1.03.H9N2     | M  | S | L | L | T | G | V | E | T | H | T  | R | N | G | W | G | C | K | C | S | D | S | S | D |
| 40 | A.chicken.Hong Kong.YU427.03.H9N    | M  | S | L | L | P | E | V | E | T | H | T  | R | N | G | W | G | C | R | C | S | D | S | S | D |

EXTRACELLULAR SEQUENCE OF D20 IS IDENTICAL  
TO #19, HK483 TO #29, AND VN1203 TO #38.

Fig. 3A




Fig. 3B-1

6/17





Fig. 3B-2

7/17



8/17




Fig. 3C-2

9/17




Fig. 4A

10/17




Fig. 4B

11/17




Fig. 5

12/17

## CROSS REACTIVITY BINDING OF ANTI-M2 ANTIBODIES TO VARIANT M2 PEPTIDES

| seqNo | Name     | Size  | Description             | ELISA (OD 450) |      |       |     |
|-------|----------|-------|-------------------------|----------------|------|-------|-----|
|       |          |       |                         | 14C2           | 8i10 | 23K12 | 2N9 |
| 1     | M2       | 23 aa | SLLTEVETPIRNEWGCRCNDSSD | +              | -    | -     | -   |
| 2     | M2SG     | 23 aa | SLLTEVETPIRSEWGCRCNDSSD | +              | -    | -     | -   |
| 3     | M2EG     | 23 aa | SLLTEVETPIRNEWECRCNGSSD | +              | -    | -     | -   |
| 4     | M2P      | 23 aa | SLPTEVETPIRNEWGCRCNDSSD | +              | -    | -     | -   |
| 5     | M2G      | 23 aa | SLLTEVETPIRNEWGCRCNGSSD | +              | -    | -     | -   |
| 6     | M2DLTGS  | 23 aa | SLLTEVDLTRNGWGCRCSDSSD  | -              | -    | +     | -   |
| 7     | M2KNS    | 23 aa | SLLTEVETPIRKEWGCNCSDSSD | +              | -    | -     | -   |
| 8     | M2LGS    | 23 aa | SLLTEVETLIRNGWGCRCSDSSD | -              | -    | -     | -   |
| 9     | M2LTGKS  | 23 aa | SLLTEVETLTNGWGCRCSDSSD  | -              | -    | -     | -   |
| 10    | M2SY     | 23 aa | SLLTEVETPIRSEWGCRCNDSSD | +              | -    | -     | -   |
| 11    | M2TGEKS  | 23 aa | SLLTEVETPTRNGWECKCSDSSD | +              | -    | -     | -   |
| 12    | M2HTGEKS | 23 aa | SLLTEVETHTRNGWECKCSDSSD | -              | -    | -     | -   |
| 13    | M2KTGEKS | 23 aa | SLLTEVKTPTNGWECKCSDSSD  | -              | -    | -     | -   |
| 14    | M2LTGS   | 23 aa | SLLTEVETLTRNGWGCRCSDSSD | -              | -    | +     | -   |
| 15    | M2TDGEKS | 23 aa | SLLTEVETPTRDGWECKCSDSSD | +              | -    | -     | -   |
| 16    | M2TGS    | 23 aa | SLLTEVETPTRNGWGCRCSDSSD | +              | -    | W     | -   |
| 17    | M2TGEK   | 23 aa | SLLTEVETPTRNGWECKCNDSSD | +              | -    | -     | -   |
| 18    | M2LTGEKS | 23 aa | SLLTEVETLTRNGWECKCSDSSD | -              | -    | W     | -   |
| 19    | M2K      | 23 aa | SLLTEVETPIRNEWGCKCNDSSD | +              | W    | +     | -   |
| 20    | M2FG     | 23 aa | SFLTEVETPIRNEWGCRCNGSSD | +              | W    | -     | -   |
| 21    | M2TGE    | 23 aa | SLLTEVETPTRNGWECCNDSSD  | +              | -    | -     | -   |
| 22    | M2KGENS  | 23 aa | SLLTEVETPIRKGWECNCSDSSD | +              | -    | -     | -   |
| 23    | M2TES    | 23 aa | SLLTEVETPTRNEWECRCSDSSD | +              | -    | -     | -   |
| 24    | M2GHTGKS | 23 aa | SLLTGVEHTRNGWGCKCSDSSD  | -              | -    | -     | -   |
| 25    | M2PHTGS  | 23 aa | SLLPEVETHTRNGWGCRCSDSSD | -              | -    | -     | -   |

PERCENTAGE COMPARED RELATIVE TO BINDING TO WILD-TYPE PEPTIDE (Seq 1)

NOTE: mAbs WERE TESTED AT 5 µg/mL

>25 %      -      NO BINDING  
 25 - 40 %      W      WEAK BINDING  
 > 40 %      +      POSITIVE BINDING

Fig. 6A

13/17

## BINDING ACTIVITY OF M2 ANTIBODIES TO TRUNCATED M2 PEPTIDES

| seqNo | Name  | Size  | Description              | 14C2 | 8i10 | 23K12 | 2N9  |
|-------|-------|-------|--------------------------|------|------|-------|------|
| 1     | M2    | 23 aa | SLLTEVETPIRNEWGCRCNDSSD  | 3.85 | 0.11 | 0.22  | 0.06 |
| 26    | M16   | 16 aa | LLTEVETPIRNEWGCR         | 3.94 | 0.09 | 0.21  | 0.09 |
| 27    | M15   | 15 aa | LTEVETPIRNEWGCR          | 3.95 | 0.09 | 0.21  | 0.09 |
| 28    | M12   | 12 aa | VETPIRNEWGCR             | 0.15 | 0.09 | 0.20  | 0.09 |
| 29    | CM17  | 17 aa | ETPIRNEWGCRCNDSSD        | 0.19 | 0.11 | 0.34  | 0.11 |
| 30    | CM16  | 16 aa | TPIRNEWGCRCNDSSD         | 0.23 | 0.13 | 0.35  | 0.12 |
| 31    | CM15  | 15 aa | PIRNEWGCRCNDSSD          | 0.19 | 0.12 | 0.34  | 0.11 |
| 32    | CM14  | 14 aa | IRNEWGCRCNDSSD           | 0.23 | 0.14 | 0.36  | 0.13 |
| 33    | CM13  | 13 aa | RNEWGCRCNDSSD            | 0.22 | 0.14 | 0.34  | 0.13 |
| 34    | CM12  | 12 aa | NEWGCRCNDSSD             | 0.27 | 0.14 | 0.39  | 0.14 |
| 35    | NM17  | 17 aa | SLLTEVETPIRNEWGCR        | 3.99 | 0.26 | 0.58  | 0.10 |
| 36    | NM16  | 16 aa | SLLTEVETPIRNEWGC         | 3.90 | 0.29 | 0.62  | 0.09 |
| 37    | NM15  | 15 aa | SLLTEVETPIRNEWG          | 3.97 | 0.12 | 0.30  | 0.11 |
| 38    | NM14  | 14 aa | SLLTEVETPIRNEW           | 3.97 | 0.11 | 0.24  | 0.09 |
| 39    | NM13  | 13 aa | SLLTEVETPIRNE            | 0.18 | 0.11 | 0.25  | 0.10 |
| 40    | NM12  | 12 aa | SLLTEVETPIRN             | 0.20 | 0.10 | 0.24  | 0.09 |
| 41    | NM11  | 11 aa | SLLTEVETPIR              | 0.21 | 0.13 | 0.30  | 0.12 |
| 42    | NM10  | 10 aa | SLLTEVETPI               | 0.17 | 0.10 | 0.24  | 0.10 |
| 43    | NM8   | 8 aa  | SLLTEVET                 | 0.15 | 0.10 | 0.20  | 0.09 |
| 44    | NM7   | 7 aa  | SLLTEVE                  | 0.14 | 0.10 | 0.20  | 0.08 |
| 45    | NM9   | 9 aa  | SLLTEVETP                | 0.21 | 0.12 | 0.30  | 0.19 |
| 46    | M2e   | 24 aa | MSLLTEVETPIRNEWGCRCNDSSD | 3.98 | 0.13 | 0.43  | 0.10 |
| CMV   | HVIR1 |       |                          | 0.16 | 0.11 | 0.21  | 3.99 |

NOTE: mAbs WERE TESTED AT 5 µg/mL

Fig. 6B




Fig. 7

15/17




Fig. 8




Fig. 9




Fig. 10




Fig. 11