

(19)



INTELLECTUAL PROPERTY
OFFICE OF SINGAPORE

(11) Publication number:

SG 177982 A1

(43) Publication date:

28.02.2012

(51) Int. Cl:

;

(12)

Patent Application

(21) Application number: 2012002978

(71) Applicant:

ABBOTT LABORATORIES 100 ABBOTT
PARK ROAD, ABBOTT PARK, IL
60064-3500 IN US

(22) Date of filing: 16.01.2008

US 60/880,767 16.01.2007

(72) Inventor:

VALDES, JOAQUIN MARIO 1290
KASTING LANE, MUNDELEIN, ILLINOIS

(30) Priority: US 60/925,960 24.04.2007

US 60/961,764 24.07.2007

US 60/997,012 28.09.2007

60060 US
CHARTASH, ELLIOT KEITH 4789
CRESTPARK LANE, MARIETTA, GA

30068 US
BARCHUK, WILLIAM T. 3923 RIVIERA
DRIVE, UNIT A, SAN DIEGO, CA 92109
US

PAULSON, SUSAN K. 1490 WOOD
AVENUE, DOWNTON GROVE, IL 60515
US

KIMBALL, ALEXANDRA B. 4
MONMOUTH STREET, BROOKLINE, MA
02445 US

(54) Title:

METHODS FOR TREATING PSORIASIS

(57) Abstract:

METHODS FOR TREATING PSORIASIS ABSTRACT This invention provides a method of treating psoriasis in a subject by administering to a subject an antibody capable of binding to the p40 subunit of IL-12 and/or IL23. Figure 1

METHODS FOR TREATING PSORIASIS

ABSTRACT

This invention provides a method of treating psoriasis in a subject by administering to a subject an antibody capable of binding to the p40 subunit of IL-12 and/or IL23.

Figure 1

METHODS FOR TREATING PSORIASIS

RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 5 60/880,767, filed on January 16, 2007; U.S. Provisional Application No. 60/904,022, filed on February 27, 2007; U.S. Provisional Application No. 60/925,960, filed on April 24, 2007; U.S. Provisional Application No. 60/961,764, filed July 24, 2007; and U.S. Provisional Application No. 60/997,012, filed on September 28, 2007, the entire contents of each of which are incorporated herein by reference.

10

BACKGROUND OF THE INVENTION

Psoriasis is a T cell-mediated inflammatory disease that is considered to be one of the most common autoimmune diseases, affecting approximately 2% to 3% of adults, though the global prevalence varies widely (Stern R.S., et al., *J Investig Dermatol Symp Proc* 2004, 9: 136–39; Davidson A and Diamond B. *N Engl J Med* 2001, 345: 340–50; Langley R.G.B., et al., *Ann Rheum Dis* 2005, 64(Suppl II): ii18–23). Psoriasis has a major impact on quality of life (de Korte J, et al., *J Investig Dermatol Symp Proc* 2004, 9: 140–7; Krueger G, et al., *Arch Dermatol* 2001, 137: 280–4; Finlay AY and Coles EC, *Br J Dermatol* 1995, 132: 236–44) and is associated with a number of psychological and 15 psychosocial problems (Kimball AB, et al., *Am J Clin Dermatol* 2005, 6: 383–92; Russo PA, et al., *Australas J Dermatol* 2004, 45: 155–9). Many traditional psoriasis therapies have toxic adverse effects; therefore, their long-term use is limited (Lebwohl M. and Ali S., *J Am Acad Dermatol* 2001, 45: 487–98; Lebwohl M. and Ali S., *J Am Acad Dermatol* 2001, 45: 649–61). In addition, many patients with psoriasis are dissatisfied 20 with traditional therapies (Stern RS, et al., *J Investig Dermatol Symp Proc* 2004, 9: 136–39; Finlay AY and Ortonne JP, *J Cutan Med Surg* 2004, 8: 310–20); thus, there is a clear need for therapies that are safer and easier to use and that can be prescribed on a long-term basis.

Interleukin-12 (IL-12) and the related cytokine IL-23 are members of the IL-12 30 superfamily of cytokines that share a common p40 subunit (Anderson EJR, et al., *Springer Semin Immunopathol* 2006, 27: 425–42). Both cytokines contribute to the development of the type 1 T helper cell (Th1) immune response in psoriasis, but each has a unique role (Rosmarin D and Strober BE, *J Drugs Dermatol* 2005, 4: 318–25; Hong

K, *et al.*, *J Immunol* 1999, 162: 7480–91; Yawalkar N, *et al.*, *J Invest Dermatol* 1998, 111: 1053–57). IL-12 primarily stimulates differentiation of Th1 cells and subsequent secretion of interferon-gamma, whereas IL-23 preferentially stimulates differentiation of naïve T cells into effector T helper cells (Th17) that secrete IL-17, a proinflammatory mediator Rosmarin D and Strober BE, *J Drugs Dermatol* 2005, 4: 318–25; Harrington Le, *et al.*, *Nature Immunol* 2005, 6: 1123–32; Park H, *et al.* *Nature Immunol* 2005, 6: 1132–41). The overexpression of IL-12 p40 and IL-23 p40 messenger RNA in psoriatic skin lesions suggests that the inhibition of IL-12 and IL-23 with a neutralizing antibody to the IL-12/23 p40 subunit protein may offer an effective therapeutic approach for the treatment of psoriasis (Yawalkar N, *et al.*, *J Invest Dermatol* 1998, 111: 1053–57; Lee E, *et al.*, *J Exp Med* 2004, 199: 125–30; Shaker OG, *et al.*, *Clin Biochem* 2006, 39: 119–25; Piskin G, *et al.*, *J Immunol* 2006, 176: 1908–15). Such therapeutic approaches for the treatment of psoriasis are clearly needed in the art.

15 SUMMARY OF THE INVENTION

The present invention provides methods and compositions for treating psoriasis, *e.g.*, chronic psoriasis, using an antibody, or antigen-binding portion thereof, that binds human IL-12 and/or human IL-23.

In one aspect, the invention provides a method of treating psoriasis in a subject comprising administering to a subject a biweekly, weekly or single dose of an antibody, or antigen-binding portion thereof, directed against human IL-12 and/or human IL-23.

In one embodiment, the subject maintains a response to the biweekly, weekly or single dose of the antibody, or antigen-binding portion thereof, for an extended period, *e.g.*, for at least about 12 weeks or for at least about 24 weeks..

25 In another embodiment, the subject maintains at least a PASI 75 response for an extended period following a biweekly, weekly or single dose of an antibody, or antigen-binding portion thereof, directed against human IL-12 and human IL-23 to the subject.

In yet another embodiment, the subject maintains at least a PASI 90 response for an extended period following a biweekly, weekly or single dose of an antibody, or antigen-binding portion thereof, directed against human IL-12 and human IL-23 to the subject.

30 In yet a further embodiment, the subject maintains at least a PASI 100 response for an extended period following a biweekly, weekly or single dose of an antibody, or antigen-binding portion thereof, directed against human IL-12 and human IL-23 to the subject.

In one embodiment, the dose of the antibody directed against human IL-12 and/or human IL-23 is about 200 mg or about 100 mg.

In one embodiment, the psoriasis is plaque psoriasis, *e.g.*, chronic plaque psoriasis. In another embodiment, the psoriasis is chronic psoriasis, *e.g.*, chronic plaque psoriasis. In yet another embodiment, the psoriasis is moderate to severe psoriasis, *e.g.*, moderate to severe plaque psoriasis, moderate to severe chronic psoriasis or moderate to severe chronic plaque psoriasis.

In one embodiment, the antibody, or antigen-binding portion thereof, is administered via subcutaneous administration.

10 In another aspect, the invention provides a method of treating psoriasis in a subject comprising the steps of: (i) selecting a subject who is suffering from chronic psoriasis; and (ii) administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23; thereby treating chronic psoriasis in the subject.

15 In one embodiment, the subject has had a clinical diagnosis of psoriasis for at least 6 months. In another embodiment, the subject has had stable plaque psoriasis for at least 2 months.

In yet another aspect, the invention provides a method of treating psoriasis in a subject comprising the steps of: (i) selecting a subject who has not had a condition selected from the group consisting of previous exposure to systemic or biologic anti-IL-12 therapy; nonplaque psoriasis; inability to discontinue topical psoriasis therapies at least 2 weeks before treatment; ultraviolet B light phototherapy at least 2 weeks before treatment; psoralen-ultraviolet-light phototherapy at least 4 weeks before treatment; systemic therapies at least 4 weeks before treatment; biologic therapies at least 12 weeks before treatment; required intake of oral or injectable corticosteroids during treatment; an exacerbation of asthma requiring hospitalization in the 10 years prior to screening; an infection or risk factors for severe infection; a history of malignancies other than successfully treated basal cell carcinoma, *e.g.*, with a history of squamous cell carcinoma, or cervical carcinoma *in situ*; and a history of major immunologic reaction, *e.g.*, serum sickness or anaphylactoid reaction, to an immunoglobulin G-containing agent, *e.g.*, intravenous gamma globulin, a fusion protein, or monoclonal antibody; and (ii) administering to the subject an antibody, or antigen-binding portion thereof, which is

capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23; thereby treating psoriasis in the subject.

In yet another aspect, the invention provides a method of treating psoriasis in a subject comprising the steps of: (i) selecting a subject who has not had vaccination with 5 a live viral agent within 1 month; and (ii) administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23; thereby treating psoriasis in the subject.

In a still further aspect, the invention provides a method of treating psoriasis in a subject comprising the steps of: (i) administering an antibody, or antigen-binding 10 portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23 to the subject; (ii) monitoring the subject for a clinically significant abnormal laboratory result selected from the group consisting of aspartate transaminase or alanine transaminase >5 times the upper limit of normal; serum total bilirubin >3 times the upper limit of normal; serum creatinine >3 times the upper limit of normal; 15 creatine phosphokinase >5 times the upper limit of normal; hemoglobin <8 g/dL; white blood cell count <2 × 10⁹/L; and platelet count <75 × 10⁹/L; and (iii) discontinuing administration of the antibody, or antigen-binding portion thereof, in a subject in which the clinically significant abnormal laboratory result is detected; thereby treating psoriasis in the subject.

20 In one embodiment, the antibody, or antigen-binding portion thereof, is administered biweekly. In another embodiment, the antibody, or antigen-binding portion thereof, is administered weekly or in a single dose.

In one embodiment the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 25 mg, 170 mg, 180 mg, 190 mg, 200 mg, 210 mg, or 220 mg.

In one embodiment, the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to the p35 subunit of IL-12. In another embodiment, the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to a p19 subunit of IL-23. In yet another embodiment, the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to a 30 p35 subunit of IL-12 and when the p40 subunit is bound to a p19 subunit of IL-23.

In one embodiment, the psoriasis is chronic psoriasis, *e.g.*, chronic plaque psoriasis, *e.g.*, moderate to severe chronic plaque psoriasis.

In another aspect, the invention provides a method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 90 response for an extended period following initial administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

In one embodiment, the extended period is at least about 12, 13, 14, 15, 16, 17, 10 18, 19, 20, 21, 22, 23 or 24 weeks.

In one embodiment, the antibody, or antigen-binding portion thereof, is administered biweekly. In another embodiment, the antibody, or antigen-binding portion thereof, is administered weekly. In yet another embodiment, the antibody is administered in a single dose.

15 In one embodiment the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170 mg, 180 mg, 190 mg, 200 mg, 210 mg, or 220 mg.

In one embodiment, the psoriasis is chronic psoriasis, *e.g.*, chronic plaque psoriasis, *e.g.*, moderate to severe chronic plaque psoriasis.

20 In yet another aspect, the invention provides a method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23 to the subject, wherein the subject maintains a clear or minimal PGA rating for an extended period following initial administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

25 In one embodiment, the extended period is at least about 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 or 24 weeks.

In one embodiment, the antibody, or antigen-binding portion thereof, is administered biweekly. In another embodiment, the antibody, or antigen-binding portion thereof, is administered weekly. In yet another embodiment, the antibody is administered in a single dose.

In one embodiment the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170 mg, 180 mg, 190 mg, 200 mg, 210 mg, or 220 mg.

5 In one embodiment, the psoriasis is chronic psoriasis, *e.g.*, chronic plaque psoriasis, *e.g.*, moderate to severe chronic plaque psoriasis.

In a still further aspect, the invention provides a method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23 to the subject, wherein the subject exhibits an improved PASI score by about 8 weeks 10 following initial administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

In one embodiment, the subject exhibits an improved PASI score by about 7 weeks, about 6 weeks, about 5 weeks, about 4 weeks, about 3 weeks, about 2 weeks or about 1 week following initial administration of the antibody, or antigen binding portion 15 thereof.

In one embodiment, the antibody, or antigen-binding portion thereof, is administered biweekly. In another embodiment, the antibody, or antigen-binding portion thereof, is administered weekly. In yet another embodiment, the antibody is administered in a single dose.

20 In one embodiment the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170 mg, 180 mg, 190 mg, 200 mg, 210 mg, or 220 mg.

In one embodiment, the psoriasis is chronic psoriasis, *e.g.*, chronic plaque psoriasis, *e.g.*, moderate to severe chronic plaque psoriasis.

25 In another aspect, the invention provides a method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 50 response for an extended period following discontinuation of administration of the antibody, or antigen-binding portion 30 thereof, thereby treating psoriasis in the subject.

In a related aspect, the invention provides a method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-

23, wherein the subject maintains at least a PASI 75 response for an extended period following discontinuation of administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

In yet another related aspect, the invention provides a method of treating
5 psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 90 response for an extended period following discontinuation of administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

10 In one embodiment, the extended period following discontinuation of administration of the antibody is at least about 12 weeks.

In one embodiment, the antibody is administered for at least about 12 weeks.

15 In one embodiment, the antibody, or antigen-binding portion thereof, is administered biweekly. In another embodiment, the antibody, or antigen-binding portion thereof, is administered weekly. In another embodiment, the antibody, or antigen-binding portion thereof, is administered in a single dose.

In one embodiment the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170 mg, 180 mg, 190 mg, 200 mg, 210 mg, or 220 mg.

20 In one embodiment, the psoriasis is chronic psoriasis, *e.g.*, chronic plaque psoriasis, *e.g.*, moderate to severe chronic plaque psoriasis.

In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention is capable of binding to an epitope of the p40 subunit of IL-12.

25 In another embodiment, the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to the p35 subunit of IL-12. In yet another embodiment, the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to a p19 subunit. In one embodiment, the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to the p35 subunit of IL-12 and when the p40 subunit is bound to a p19 subunit.

In one embodiment, the antibody, or antigen binding portion thereof, binds to an epitope of the p40 subunit of IL-12 to which an antibody selected from the group consisting of Y61 and J695 binds.

In another embodiment, the antibody is further capable of binding to a first 5 heterodimer and is also capable of binding to a second heterodimer, wherein the first heterodimer comprises the p40 subunit of IL-12 and the p35 subunit of IL-12, and wherein the second heterodimer comprises the p40 subunit of IL-12 and a p19 subunit.

In a further embodiment, the antibody neutralizes the activity of the first heterodimer. In another embodiment, the antibody neutralizes the activity of the second 10 heterodimer. In yet another embodiment, the antibody neutralizes the activity of the first heterodimer and the second heterodimer.

In a further embodiment, the antibody, or antigen binding portion thereof, used in the methods of the invention inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-9} M or less, or which inhibits human $IFN\gamma$ 15 production with an IC_{50} of 1×10^{-10} M or less.

In one embodiment, the antibody, or antigen binding portion thereof, used in the methods of the invention dissociates from the p40 subunit of IL-12 with a K_d of 1×10^{-10} M or less or a k_{off} rate constant of $1 \times 10^{-3} s^{-1}$ or less, as determined by surface plasmon resonance.

20 In one embodiment, the isolated antibody, or antigen binding portion thereof, used in the methods of the invention is a chimeric antibody, a humanized antibody or a human antibody.

In another embodiment, the antibody, or antigen binding portion thereof, used in the methods of the invention has a heavy chain CDR3 comprising the amino acid 25 sequence of SEQ ID NO: 25 and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26;

In a further embodiment, the antibody, or antigen binding portion thereof, used in the methods of the invention has a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 27 and a light chain CDR2 comprising the amino acid 30 sequence of SEQ ID NO: 28.

In one embodiment, the antibody, or antigen binding portion thereof, used in the methods of the invention has a heavy chain CDR1 comprising the amino acid sequence

of SEQ ID NO: 29 and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 30.

In another embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention is capable of binding to an interleukin comprising a p40 subunit. In one embodiment, the interleukin comprises a p40 subunit and a p35 subunit, *e.g.*, the interleukin is IL-12. In another embodiment, the interleukin comprises a p40 subunit and a p19 subunit. In yet another embodiment, the antibody, or antigen binding portion thereof, neutralizes the activity of the interleukin.

In one embodiment, the antibody, or antigen binding portion thereof, binds to an epitope of the p40 subunit.

In one embodiment, the antibody, or antigen-binding portion thereof, is administered to a subject in a pharmaceutical composition comprising the antibody, or antigen binding portion thereof, and a pharmaceutically acceptable carrier. The pharmaceutical composition may also comprise an additional agent, such as a therapeutic agent, *e.g.*, budesonide, epidermal growth factor, corticosteroids, cyclosporin, sulfasalazine, aminosalicylates, 6-mercaptopurine, azathioprine, metronidazole, lipoxygenase inhibitors, mesalamine, olsalazine, balsalazide, antioxidants, thromboxane inhibitors, IL-1 receptor antagonists, anti-IL-1 β monoclonal antibodies, anti-IL-6 monoclonal antibodies, growth factors, elastase inhibitors, pyridinyl-imidazole compounds, antibodies or agonists of TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, IL-18, EMAP-II, GM-CSF, FGF, and PDGF, antibodies of CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands, methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, ibuprofen, corticosteroids, prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, IRAK, NIK, IKK, p38, MAP kinase inhibitors, IL-1 β converting enzyme inhibitors, TNF α converting enzyme inhibitors, T-cell signalling inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors, soluble p55 TNF receptor, soluble p75 TNF receptor, sIL-1RI, sIL-1RII, sIL-6R, antiinflammatory cytokines, IL-4, IL-10, IL-11, IL-13 and TGF β .

In another embodiment, the therapeutic agent in the pharmaceutical composition administered to the subject may be selected from the group consisting of anti-TNF

antibodies and antibody fragments thereof, TNFR-Ig constructs, TACE inhibitors, PDE4 inhibitors, corticosteroids, budesonide, dexamethasone, sulfasalazine, 5-aminosalicylic acid, olsalazine, IL-1 β converting enzyme inhibitors, IL-1ra, tyrosine kinase inhibitors, 6-mercaptopurines and IL-11.

5 In another embodiment, the therapeutic agent may be selected from the group consisting of corticosteroids, prednisolone, methylprednisolone, azathioprine, cyclophosphamide, cyclosporine, methotrexate, 4-aminopyridine, tizanidine, interferon- β 1a, interferon- β 1b, Copolymer 1, hyperbaric oxygen, intravenous immunoglobulin, clabribine, antibodies or agonists of TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, 10 IL-18, EMAP-II, GM-CSF, FGF, PDGF, antibodies to CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD80, CD86, CD90 or their ligands, methotrexate, cyclosporine, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, ibuprofen, corticosteroids, prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, IRAK, NIK, 15 IKK, p38 or MAP kinase inhibitors, IL-1 β converting enzyme inhibitors, TACE inhibitors, T-cell signalling inhibitors, kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors, soluble p55 TNF receptor, soluble p75 TNF receptor, sIL-1RI, sIL-1RII, sIL-6R, sIL-13R, anti-P7s, p-selectin glycoprotein ligand 20 (PSGL), antiinflammatory cytokines, IL-4, IL-10, IL-13 and TGF β .

In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention binds to human IL-12 and/or human IL-23 and dissociates from human IL-12 and/or human IL-23, respectively, with a K_d of 1×10^{-10} M or less and a k_{off} rate constant of 1×10^{-3} s $^{-1}$ or less, as determined by surface plasmon resonance. In one embodiment, the antibody, or antigen-binding portion thereof, dissociates from human IL-12 and/or human IL-23 with a k_{off} rate constant of 1×10^{-4} s $^{-1}$ or less. In another embodiment, the antibody, or antigen-binding portion thereof, dissociates from human IL-12 and/or human IL-23 with a k_{off} rate constant of 1×10^{-5} s $^{-1}$ or less.

30 In another embodiment, the antibody, or antigen-binding portion thereof, binds to human IL-12 and/or human IL-23 and dissociates from human IL-12 and/or human IL-23, respectively, with a k_{off} rate constant of 1×10^{-2} s $^{-1}$ or less, as determined by surface plasmon resonance. In yet another embodiment, the antibody, or antigen-binding

portion thereof, dissociates from human IL-12 and/or human IL-23 with a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less. In a still further another embodiment, the antibody, or antigen-binding portion thereof, dissociates from human IL-12 and/or human IL-23 with a k_{off} rate constant of $1 \times 10^{-4} \text{ s}^{-1}$ or less. In another embodiment, the antibody, or antigen-binding portion thereof, dissociates from human IL-12 and/or human IL-23 with a k_{off} rate constant of $1 \times 10^{-5} \text{ s}^{-1}$ or less.

In still another embodiment, the antibody, or antigen-binding portion thereof, binds to human IL-12 and/or human IL-23 and dissociates from human IL-12 and/or human IL-23, respectively, with a K_d of $1.34 \times 10^{-10} \text{ M}$ or less. In yet another embodiment, the antibody, or antigen-binding portion thereof, binds to human IL-12 and/or human IL-23 and dissociates from human IL-12 and/or human IL-23, respectively, with a K_d of $9.74 \times 10^{-11} \text{ M}$ or less. In one embodiment, the antibody, or antigen-binding portion thereof, is a recombinant antibody, or antigen-binding portion thereof.

In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention is a neutralizing antibody, *e.g.*, neutralizes the activity of human IL-12 and/or human IL-23. In one embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-9} \text{ M}$ or less. In another embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-10} \text{ M}$ or less. In still another embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-11} \text{ M}$ or less. In yet another embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* phytohemagglutinin blast proliferation assay (PHA assay) with an IC_{50} of $1 \times 10^{-7} \text{ M}$ or less. In still another embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-8} \text{ M}$ or less. In one embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits human IFN γ production with an IC_{50} of $1 \times 10^{-10} \text{ M}$ or less. In still another embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits human IFN γ production with an IC_{50} of $1 \times 10^{-11} \text{ M}$ or less. In yet a

further embodiment, the neutralizing antibody, or antigen-binding portion thereof, inhibits human IFN γ production with an IC₅₀ of 5 x 10⁻¹² M or less.

In one embodiment, the antibody, or an antigen-binding portion thereof, used in the methods of the invention

- 5 a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less;
- b) has a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 25; and
- c) has a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26. In one embodiment, the antibody further has a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 27; and a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 28. In still another embodiment, the antibody, or antigen-binding portion thereof, further has a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 29; and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 30. In still another embodiment, the antibody, or antigen-binding portion thereof, further inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻¹⁰ M or less. In still another embodiment, the antibody, or antigen-binding portion thereof, further inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻¹¹ M or less.

20 In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention has a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 31, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 32.

25 In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention comprises a heavy chain constant region selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant regions. In one embodiment, the antibody heavy chain constant region is IgG1. In another embodiment, the antibody is a Fab fragment, F(ab')₂ fragment, or a single chain Fv fragment.

30 In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention dissociates from human IL-12 and/or human IL-23 with a K_d of 1 x 10⁻¹⁰ M or less and binds to an epitope on the p40 subunit of human IL-12 and/or human IL-23.

In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention is a human antibody, or antigen-binding portion thereof, which

- a) dissociates from human IL-12 with a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, as determined by surface plasmon resonance;
- 5 b) has a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 25; and
- c) has a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26.

In another embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention dissociates from human IL-12 with a k_{off} rate constant of $1 \times 10^{-4} \text{ s}^{-1}$ or less. In a further embodiment, the human antibody, or antigen-binding portion thereof, dissociates from human IL-12 with a k_{off} rate constant of $1 \times 10^{-5} \text{ s}^{-1}$ or less.

In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention is a human antibody, or antigen-binding portion thereof, that binds to human IL-12 and comprises:

a light chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 26; and

20 a heavy chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 25.

In one embodiment, the antibody, or antigen-binding portion thereof, has a light chain variable region (LCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 26, and has a heavy chain variable region (HCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 25. In another embodiment, the antibody, or antigen-binding portion thereof, comprises an LCVR further having a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 28 and an HCVR further comprising a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 27. In yet another embodiment, the LCVR further has CDRI domain comprising the amino acid sequence of SEQ ID NO: 30 and the HCVR has a CDRI domain comprising the amino acid sequence of SEQ ID NO: 29.

In one embodiment, the antibody, or antigen-binding portion thereof, binds human IL-12 and/or human IL-23 and is the antibody J695 (also referred to as ABT-874), or an antigen binding portion thereof.

In one embodiment, the antibody, or antigen-binding portion thereof, binds to human IL-12 and/or human IL-23 and dissociates from human IL-12 with a K_d of 1.34×10^{-10} M or less, and neutralizes human IL-12 and/or human IL-23. In one embodiment, the antibody, or antigen-binding portion thereof, dissociates from human IL-12 and/or human IL-23 with a K_d of 9.74×10^{-11} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-7} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-8} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-9} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-10} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-11} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits human IFN γ production with an IC_{50} of 1×10^{-10} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits human IFN γ production with an IC_{50} of 1×10^{-11} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits human IFN γ production with an IC_{50} of 5×10^{-12} M or less.

In one embodiment, the antibody, or antigen-binding portion thereof, used in the methods of the invention inhibits IL-12 and/or IL-23 binding to its receptor in an IL-12 or IL-23 receptor binding assay (RBA), respectively, with an IC_{50} of 1×10^{-9} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits IL-12 and/or IL-23 binding to its receptor in an IL-12 or IL-23 receptor binding assay (RBA), respectively, with an IC_{50} of 1×10^{-10} M or less. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits IL-12 and/or IL-23 binding to its receptor in an IL-12 or IL-23 receptor binding assay (RBA), respectively, with an IC_{50} of 1×10^{-11} M or less.

30

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the patient disposition of the trial. (The term “eow” refers to every other week dosing.)

Figure 2 shows the percentage of patients with at least a 75% improvement in the psoriasis area and severity index (PASI 75) during the 12-week portion of the trial. By week 8, with the exception of the 200 mg × 1 group, the percentage of patients who had a PASI 75 response was statistically significantly greater ($p<0.001$) in each ABT-874 treatment group for each comparison with placebo based on an analysis of variance of observed data for the intention-to-treat population. (The term "eow" refers to every other week dosing.)

Figure 3 shows the mean percentage improvement in psoriasis area and severity index (PASI) scores from baseline. The data show that $*p<0.001$ for each ABT-874 treatment group compared with placebo at all time points (except 100 mg eow at week 1, $p=0.023$) based on an analysis of variance of observed data for the intention-to-treat population. (The term "eow" refers to every other week dosing.)

Figures 4A-C show the percentage of patients who maintained a PASI 50, PASI 75 and PASI 90 response, respectively, at week 24 of the trial, *i.e.*, at 12 weeks following discontinuation of administration of the antibody.

Figure 4D shows the percentage of patients maintaining a PASI 75 response over time during the 24 week period of the trial.

DETAILED DESCRIPTION OF THE INVENTION

In order that the present invention may be more readily understood, certain terms are first defined.

The term "activity enhancing amino acid residue" includes an amino acid residue which improves the activity of the antibody. It should be understood that the activity enhancing amino acid residue may replace an amino acid residue at a contact, hypermutation or preferred selective mutagenesis position and, further, more than one activity enhancing amino acid residue can be present within one or more CDRs. An activity enhancing amino acid residue include, an amino acid residue that improves the binding specificity/affinity of an antibody, for example anti-human IL-12 antibody binding to human IL-12. The activity enhancing amino acid residue is also intended to include an amino acid residue that improves the neutralization potency of an antibody, for example, the human IL-12 antibody which inhibits human IL-12.

The term "antibody" includes an immunoglobulin molecule comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by

disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

10 In one embodiment, the antibody used in the compositions and methods of the invention is the antibody described in U.S. Patent No. 6,914,128, incorporated by reference herein. In another embodiment, the antibody used in the compositions and methods of the invention is the antibody ABT-874 (also referred to as J695; Abbott Laboratories).

15 The term "antigen-binding portion" of an antibody (or "antibody portion") includes fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., hIL-12). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883) . Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific

antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see e.g., Holliger, P., *et al.* (1993) *Proc. 5 Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994) *Structure* 2:1121-1123). Still further, an antibody or antigen-binding portion thereof may be part of a larger immunoadhesion molecules, formed by covalent or non-covalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a 10 tetrameric scFv molecule (Kipriyanov, S.M., *et al.* (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov, S.M., *et al.* (1994) *Mol. Immunol.* 31:1047-1058). Antibody portions, such as Fab and F(ab')₂ 15 fragments, can be prepared from whole antibodies using conventional techniques, such as papain or pepsin digestion, respectively, of whole antibodies. Moreover, antibodies, antibody portions and immunoadhesion molecules can be obtained using standard recombinant DNA techniques, as described herein. Preferred antigen binding portions 20 are complete domains or pairs of complete domains.

The term “backmutation “ refers to a process in which some or all of the 25 somatically mutated amino acids of a human antibody are replaced with the corresponding germline residues from a homologous germline antibody sequence. The heavy and light chain sequences of the human antibody of the invention are aligned separately with the germline sequences in the VBASE database to identify the sequences with the highest homology. Differences in the human antibody of the invention are returned to the germline sequence by mutating defined nucleotide positions encoding 30 such different amino acid. The role of each amino acid thus identified as candidate for backmutation should be investigated for a direct or indirect role in antigen binding and any amino acid found after mutation to affect any desirable characteristic of the human antibody should not be included in the final human antibody; as an example, activity enhancing amino acids identified by the selective mutagenesis approach will not be subject to backmutation. To minimize the number of amino acids subject to backmutation those amino acid positions found to be different from the closest germline sequence but identical to the corresponding amino acid in a second germline sequence

can remain, provided that the second germline sequence is identical and colinear to the sequence of the human antibody of the invention for at least 10, preferably 12 amino acids, on both sides of the amino acid in question. Backmutation may occur at any stage of antibody optimization; preferably, backmutation occurs directly before or after the 5 selective mutagenesis approach. More preferably, backmutation occurs directly before the selective mutagenesis approach.

The phrase "human interleukin 12" (abbreviated herein as hIL-12, or IL-12), as used herein, includes a human cytokine that is secreted primarily by macrophages and dendritic cells. The term includes a heterodimeric protein comprising a 35 kD subunit 10 (p35) and a 40 kD subunit (p40) which are both linked together with a disulfide bridge. The heterodimeric protein is referred to as a "p70 subunit". The structure of human IL-12 is described further in, for example, Kobayashi, *et al.* (1989) *J. Exp Med.* 170:827-845; Seder, *et al.* (1993) *Proc. Natl. Acad. Sci.* 90:10188-10192; Ling, *et al.* (1995) *J. Exp Med.* 154:116-127; Podlaski, *et al.* (1992) *Arch. Biochem. Biophys.* 294:230-237. 15 The term human IL-12 is intended to include recombinant human IL-12 (rh IL-12), which can be prepared by standard recombinant expression methods.

The terms "Kabat numbering", "Kabat definitions and "Kabat labeling" are used interchangeably herein. These terms, which are recognized in the art, refer to a system of numbering amino acid residues which are more variable (*i.e.* hypervariable) than 20 other amino acid residues in the heavy and light chain variable regions of an antibody, or an antigen binding portion thereof (Kabat *et al.* (1971) *Ann. NY Acad. Sci.* 190:382-391 and , Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the hypervariable region ranges from amino 25 acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

The Kabat numbering is used herein to indicate the positions of amino acid 30 modifications made in antibodies of the invention. For example, the Y61 anti-IL-12 antibody can be mutated from serine (S) to glutamic acid (E) at position 31 of the heavy chain CDR1 (H31S → E), or glycine (G) can be mutated to tyrosine (Y) at position 94 of the light chain CDR3 (L94G → Y).

The term "human antibody" includes antibodies having variable and constant regions corresponding to human germline immunoglobulin sequences as described by Kabat *et al.* (See Kabat, *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (*e.g.*, mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. The mutations preferably are introduced using the "selective mutagenesis approach" described herein. The human antibody can have at least one position replaced with an amino acid residue, *e.g.*, an activity enhancing amino acid residue which is not encoded by the human germline immunoglobulin sequence. The human antibody can have up to twenty positions replaced with amino acid residues which are not part of the human germline immunoglobulin sequence. In other embodiments, up to ten, up to five, up to three or up to two positions are replaced. In a preferred embodiment, these replacements are within the CDR regions as described in detail below. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The phrase "recombinant human antibody" includes human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further in Section II, below), antibodies isolated from a recombinant, combinatorial human antibody library (described further in Section III, below), antibodies isolated from an animal (*e.g.*, a mouse) that is transgenic for human immunoglobulin genes (see *e.g.*, Taylor, L.D., *et al.* (1992) *Nucl. Acids Res.* 20:6287-6295) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences (See Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic

for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*. In certain 5 embodiments, however, such recombinant antibodies are the result of selective mutagenesis approach or backmutation or both.

An "isolated antibody" includes an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds hIL-12 is substantially free of antibodies that specifically bind 10 antigens other than hIL-12). An isolated antibody that specifically binds hIL-12 may bind IL-12 molecules from other species (discussed in further detail below). Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

A "neutralizing antibody" (or an "antibody that neutralized hIL-12 activity") 15 includes an antibody whose binding to hIL-12 results in inhibition of the biological activity of hIL-12. This inhibition of the biological activity of hIL-12 can be assessed by measuring one or more indicators of hIL-12 biological activity, such as inhibition of human phytohemagglutinin blast proliferation in a phytohemagglutinin blast proliferation assay (PHA), or inhibition of receptor binding in a human IL-12 receptor 20 binding assay (see Example 3-Interferon-gamma Induction Assay of US Patent No. 6,914,128). These indicators of hIL-12 biological activity can be assessed by one or more of several standard *in vitro* or *in vivo* assays known in the art (see Example 3 of US Patent No. 6,914,128).

The term "activity" includes activities such as the binding specificity/affinity of 25 an antibody for an antigen, for example, an anti-hIL-12 antibody that binds to an IL-12 antigen and/or the neutralizing potency of an antibody, for example, an anti-hIL-12 antibody whose binding to hIL-12 inhibits the biological activity of hIL-12, e.g. inhibition of PHA blast proliferation or inhibition of receptor binding in a human IL-12 receptor binding assay (see Example 3 of US Patent No. 6,914,128).

30 The phrase "surface plasmon resonance" includes an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIACore system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, NJ). For further

descriptions, see Example 5 of US Patent No. 6,914,128 and Jönsson, U., *et al.* (1993) *Ann. Biol. Clin.* 51:19-26; Jönsson, U., *et al.* (1991) *Biotechniques* 11:620-627; Johnsson, B., *et al.* (1995) *J. Mol. Recognit.* 8:125-131; and Johnnson, B., *et al.* (1991) *Anal. Biochem.* 198:268-277.

5 The term "K_{off}", as used herein, is intended to refer to the off rate constant for dissociation of an antibody from the antibody/antigen complex.

The term "K_d", as used herein, is intended to refer to the dissociation constant of a particular antibody-antigen interaction.

10 The phrase "nucleic acid molecule" includes DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

15 The phrase "isolated nucleic acid molecule", as used herein in reference to nucleic acids encoding antibodies or antibody portions (e.g., VH, VL, CDR3) that bind hIL-12 including "isolated antibodies"), includes a nucleic acid molecule in which the nucleotide sequences encoding the antibody or antibody portion are free of other 20 nucleotide sequences encoding antibodies or antibody portions that bind antigens other than hIL-12, which other sequences may naturally flank the nucleic acid in human genomic DNA. Thus, for example, an isolated nucleic acid of the invention encoding a VH region of an anti-IL-12 antibody contains no other sequences encoding other VH regions that bind antigens other than IL-12. The phrase "isolated nucleic acid molecule" is also intended to include sequences encoding bivalent, bispecific antibodies, such as 25 diabodies in which VH and VL regions contain no other sequences other than the sequences of the diabody.

25 The term "vector" includes a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of 30 autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes

to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used 5 interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The phrase "recombinant host cell" (or simply "host cell") includes a cell into 10 which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term 15 "host cell" as used herein.

The term "modifying", as used herein, is intended to refer to changing one or more amino acids in the antibodies or antigen-binding portions thereof. The change can be produced by adding, substituting or deleting an amino acid at one or more positions. The change can be produced using known techniques, such as PCR mutagenesis.

20 The phrase "contact position" includes an amino acid position of in the CDR1, CDR2 or CDR3 of the heavy chain variable region or the light chain variable region of an antibody which is occupied by an amino acid that contacts antigen in one of the twenty-six known antibody-antigen structures. If a CDR amino acid in any of the 26 known solved structures of antibody-antigen complexes contacts the antigen, then that 25 amino acid can be considered to occupy a contact position. Contact positions have a higher probability of being occupied by an amino acid which contact antigen than non-contact positions. Preferably a contact position is a CDR position which contains an amino acid that contacts antigen in greater than 3 of the 26 structures (>11.5 %). Most preferably a contact position is a CDR position which contains an amino acid that 30 contacts antigen in greater than 8 of the 25 structures (>32%).

The term "hypermutation position" includes an amino acid residue that occupies position in the CDR1, CDR2 or CDR3 region of the heavy chain variable region or the light chain variable region of an antibody that is considered to have a high frequency or

probability for somatic hypermutation during *in vivo* affinity maturation of the antibody. "High frequency or probability for somatic hypermutation" includes frequencies or probabilities of a 5 to about 40% chance that the residue will undergo somatic hypermutation during *in vivo* affinity maturation of the antibody. It should be
5 understood that all ranges within this stated range are also intended to be part of this invention, e.g., 5 to about 30%, e.g., 5 to about 15%, e.g., 15 to about 30%.

10 The term "preferred selective mutagenesis position" includes an amino acid residue that occupies a position in the CDR1, CDR2 or CDR3 region of the heavy chain variable region or the light chain variable region which can be considered to be both a contact and a hypermutation position.

15 The phrase "selective mutagenesis approach" includes a method of improving the activity of an antibody by selecting and individually mutating CDR amino acids at at least one preferred selective mutagenesis position, hypermutation, and/or contact position. A "selectively mutated" human antibody is an antibody which contains a
20 mutation at a position selected using a selective mutagenesis approach. In another embodiment, the selective mutagenesis approach is intended to provide a method of preferentially mutating selected individual amino acid residues in the CDR1, CDR2 or CDR3 of the heavy chain variable region (hereinafter H1, H2, and H3, respectively), or the CDR1, CDR2 or CDR3 of the light chain variable region (hereinafter referred to as
25 L1, L2, and L3, respectively) of an antibody. Amino acid residues may be selected from preferred selective mutagenesis positions, contact positions., or hypermutation positions. Individual amino acids are selected based on their position in the light or heavy chain variable region. It should be understood that a hypermutation position can also be a contact position. In an embodiment, the selective mutagenesis approach is a "targeted approach". The language "targeted approach" is intended to include a method of preferentially mutating selected individual amino acid residues in the CDR1, CDR2 or CDR3 of the heavy chain variable region or the CDR1, CDR2 or CDR3 of the light chain variable region of an antibody in a targeted manner, e.g., a "Group-wise targeted approach" or "CDR-wise targeted approach". In the "Group-wise targeted approach",
30 individual amino acid residues in particular groups are targeted for selective mutations including groups I (including L3 and H3), II (including H2 and L1) and III (including L2 and H1), the groups being listed in order of preference for targeting. In the "CDR-wise targeted approach", individual amino acid residues in particular CDRs are targeted

for selective mutations with the order of preference for targeting as follows: H3, L3, H2, L1, H1 and L2. The selected amino acid residue is mutated, *e.g.*, to at least two other amino acid residues, and the effect of the mutation on the activity of the antibody is determined. Activity is measured as a change in the binding specificity/affinity of the antibody, and/or neutralization potency of the antibody. It should be understood that the selective mutagenesis approach can be used for the optimization of any antibody derived from any source including phage display, transgenic animals with human IgG germline genes, human antibodies isolated from human B-cells. Preferably, the selective mutagenesis approach is used on antibodies which can not be optimized further using phage display technology. It should be understood that antibodies from any source including phage display, transgenic animals with human IgG germline genes, human antibodies isolated from human B-cells can be subject to backmutation prior to or after the selective mutagenesis approach.

The term "activity enhancing amino acid residue" includes an amino acid residue which improves the activity of the antibody. It should be understood that the activity enhancing amino acid residue may replace an amino acid residue at a preferred selective mutagenesis position, contact position, or a hypermutation position and, further, more than one activity enhancing amino acid residue can be present within one or more CDRs. An activity enhancing amino acid residue include, an amino acid residue that improves the binding specificity/affinity of an antibody, for example anti-human IL-12 antibody binding to human IL-12. The activity enhancing amino acid residue is also intended to include an amino acid residue that improves the neutralization potency of an antibody, for example, the human IL-12 antibody which inhibits human IL-12.

The term "dosing", as used herein, refers to the administration of a substance (*e.g.*, an anti-IL-12, anti-IL-23 antibody) to achieve a therapeutic objective (*e.g.*, the treatment of rheumatoid arthritis).

The terms "biweekly dosing regimen", "biweekly dosing", and "biweekly administration", as used herein, refer to the time course of administering a substance (*e.g.*, an anti-IL-12, anti-IL-23 antibody) to a subject to achieve a therapeutic objective, wherein the time course is every other week (eow). The biweekly dosing regimen is not intended to include a weekly dosing regimen. Preferably, the substance is administered every 9-19 days, more preferably, every 11-17 days, even more preferably, every 13-15 days, and most preferably, every 14 days.

The term “combination” as in the phrase “a first agent in combination with a second agent” includes co-administration of a first agent and a second agent, which for example may be dissolved or intermixed in the same pharmaceutically acceptable carrier, or administration of a first agent, followed by the second agent, or

5 administration of the second agent, followed by the first agent. The present invention, therefore, includes methods of combination therapeutic treatment and combination pharmaceutical compositions.

The term “concomitant” as in the phrase “concomitant therapeutic treatment” includes administering an agent in the presence of a second agent. A concomitant therapeutic treatment method includes methods in which the first, second, third, or additional agents are co-administered. A concomitant therapeutic treatment method also includes methods in which the first or additional agents are administered in the presence of a second or additional agents, wherein the second or additional agents, for example, may have been previously administered. A concomitant therapeutic treatment method

10 may be executed step-wise by different actors. For example, one actor may administer to a subject a first agent and a second actor may administer to the subject a second agent, and the administering steps may be executed at the same time, or nearly the same time, or at distant times, so long as the first agent (and additional agents) are after administration in the presence of the second agent (and additional agents). The actor

15 and the subject may be the same entity (e.g., human).

20

The term “combination therapy”, as used herein, refers to the administration of two or more therapeutic substances, e.g., an anti-IL-12, anti-IL-23 antibody and another drug. The other drug(s) may be administered concomitant with, prior to, or following the administration of an anti-IL-12, anti-IL-23 antibody.

25 The term "kit" as used herein refers to a packaged product comprising components with which to administer the anti-IL-12, anti-IL-23 antibody of the invention for treatment of a IL-12 related disorder. The kit preferably comprises a box or container that holds the components of the kit. The box or container is affixed with a label or a Food and Drug Administration approved protocol. The box or container holds

30 components of the invention which are preferably contained within plastic, polyethylene, polypropylene, ethylene, or propylene vessels. The vessels can be capped-tubes or bottles. The kit can also include instructions for administering an anti-IL-12, anti-IL-23 antibody.

Various aspects of the invention are described in further detail in the following subsections.

I. Human Antibodies that Bind Human IL-12

5 This invention provides methods and compositions for using human antibodies, or antigen-binding portions thereof, that bind to human IL-12 for the treatment of psoriasis. The invention also includes methods and compositions for using an antibody which binds both IL-12 and IL-23. Preferably, the human antibodies used in the invention are recombinant, neutralizing human anti-hIL-12 antibodies.

10 In one embodiment, the antibody used in the invention is the antibody ABT-874 (see US Patent No. 6,914,128). ABT-874 is a fully human antibody against interleukin 12 (IL-12) and IL-23. It binds with great affinity to the p40 subunit common to both IL-12 and IL-23, validated targets in the treatment of psoriasis (Ps).

15 Antibodies that bind to human IL-12 can be selected, for example, by screening one or more human V_L and V_H cDNA libraries with hIL-12, such as by phage display techniques as described in Example 1 of US Patent No. 6,914,128. Screening of human V_L and V_H cDNA libraries initially identified a series of anti-IL-12 antibodies of which one antibody, referred to herein as "Joe 9" (or "Joe 9 wild type"), was selected for further development. Joe 9 is a relatively low affinity human IL-12 antibody (e.g., a 20 K_{off} of about 0.1 sec^{-1}), yet is useful for specifically binding and detecting hIL-12. The affinity of the Joe 9 antibody was improved by conducting mutagenesis of the heavy and light chain CDRs, producing a panel of light and heavy chain variable regions that were "mixed and matched" and further mutated, leading to numerous additional anti-hIL-12 antibodies with increased affinity for hIL-12 (see Example 1, table 2 (see Appendix A) 25 of US Patent No. 6,914,128 and the sequence alignments of Figures 1A-D of US Patent No. 6,914,128).

30 Of these antibodies, the human anti-hIL-12 antibody referred to herein as Y61 demonstrated a significant improvement in binding affinity (e.g., a K_{off} of about $2 \times 10^{-4} \text{ sec}^{-1}$). The Y61 anti-hIL-12 antibody was selected for further affinity maturation by individually mutating specific amino acids residues within the heavy and light chain CDRs. Amino acids residues of Y61 were selected for site-specific mutation (selective mutagenesis approach) based on the amino acid residue occupying a preferred selective mutagenesis position, contact and/or a hypermutation position. A summary of the

substitutions at selected positions in the heavy and light chain CDRs is shown in Figures 2A-2H of US Patent No. 6,914,128. A preferred recombinant neutralizing antibody of the invention, referred to herein as J695 (also referred to as ABT-874 (Abbott Laboratories), resulted from a Gly to Tyr substitution at position 50 of the light chain CDR2 of Y61, and a Gly to Tyr substitution at position 94 of the light chain CDR3 of Y61.

5 Amino acid sequence alignments of the heavy and light chain variable regions of a panel of anti-hIL-12 antibodies used in the invention, on the lineage from Joe 9 wild type to J695, are shown in Figures 1A-1D of US Patent No. 6,914,128. These sequence 10 alignments allowed for the identification of consensus sequences for preferred heavy and light chain variable regions of antibodies of the invention that bind hIL-12, as well as consensus sequences for the CDR3, CDR2, and CDR1, on the lineage from Joe 9 to J695. Moreover, the Y61 mutagenesis analysis summarized in Figures 2A-2H allowed 15 for the identification of consensus sequences for heavy and light chain variable regions that bind hIL-12, as well as consensus sequences for the CDR3, CDR2, and CDR1 that bind hIL-12 on the lineage from Y61 to J695 that encompasses sequences with 20 modifications from Y61 yet that retain good hIL-12 binding characteristics. Preferred CDR, VH and VL sequences of the invention (including consensus sequences) as identified by sequence identifiers in the attached Sequence Listing, are summarized below.

SEQ ID NO:	ANTIBODY CHAIN	REGION	SEQUENCE
1	Consensus Joe 9 to J695	CDR H3	(H/S) - G - S - (H/Y) - D - (N/T/Y)
2	Consensus Joe 9 to J695	CDR L3	Q - (S/T) - Y - (D/E) - (S/R/K) - (S/G/Y) - (L/F/T/S) - (R/S/T/W/H) - (G/P) - (S/T/A/L) - (R/S/M/T/L) - (V/I/T/M/L)
3	Consensus Joe 9 to J695	CDR H2	F - I - R - Y - D - G - S - N - K - Y - Y - A - D - S - V - K - G
4	Consensus Joe 9 to J695	CDR L2	(G/Y) - N - (D/S) - (Q/N) - R - P - S
5	Consensus	CDR H1	F - T - F - S - (S/E) - Y - G - M - H

	Joe 9 to J695		
6	Consensus Joe 9 to J695	CDR L1	(S/T) - G - (G/S) - (R/S) - S - N - I - (G/V) - (S/A) - (N/G/Y) - (T/D) - V - (K/H)
7	Consensus Joe 9 to J695	VH	(full VH sequence; see sequence listing)
8	Consensus Joe 9 to J695	VL	(full VL sequence; see sequence listing)
9	Consensus Y61 to J695	CDR H3	H - (G/V/C/H) - (S/T) - (H/T/V/R/I) - (D/S) - (N/K/A/T/S/F/W/H)
10	Consensus Y61 to J695	CDR L3	Q - S - Y - (D/S) - (Xaa) - (G/D/Q/L/F/R/H/N/Y) - T - H - P - A - L - L
11	Consensus Y61 to J695	CDR H2	(F/T/Y) - I - (R/A) - Y - (D/S/E/A) - (G/R) - S - (Xaa) - K - (Y/E) - Y - A - D - S - V - K - G
12	Consensus Y61 to J695	CDR L2	(G/Y/S/T/N/Q) - N - D - Q - R - P - S
13	Consensus Y61 to J695	CDR H1	F - T - F - (Xaa) - (Xaa) - (Y/H) - (G/M/A/N/S) - M - H
14	Consensus Y61 to J695	CDR L1	S - G - G - R - S - N - I - G - (S/C/R/N/D/T) - (N/M/I) - (T/Y/D/H/K/P) - V - K
15	Consensus Y61 to J695	VH	(full VH sequence; see sequence listing)
16	Consensus Y61 to J695	VL	(full VL sequence; see sequence listing)
17	Y61	CDR H3	H - G - S - H - D - N
18	Y61	CDR L3	Q - S - Y - D - R - G - T - H - P - A - L - L
19	Y61	CDR H2	F - I - R - Y - D - G - S - N - K - Y - Y - A - D - S - V - K - G
20	Y61	CDR L2	G - N - D - Q - R - P - S
21	Y61	CDR H1	F - T - F - S - S - Y - G - M - H
22	Y61	CDR L1	S - G - G - R - S - N - I - G - S - N - T - V - K
23	Y61	VH	(full VH sequence; see sequence listing)

24	Y61	VL	(full VL sequence; see sequence listing)
25	J695	CDR H3	H-G-S-H-D-N
26	J695	CDR L3	Q-S-Y-D-R-Y-T-H-P-A-L-L
27	J695	CDR H2	F-I-R-Y-D-G-S-N-K-Y-Y-A-D-S-V-K-G
28	J695	CDR L2	Y-N-D-Q-R-P-S
29	J695	CDR H1	F-T-F-S-S-Y-G-M-H
30	J695	CDR L1	S-G-S-R-S-N-I-G-S-N-T-V-K
31	J695	VH	(full VH sequence; see sequence listing)
32	J695	VL	(full VL sequence; see sequence listing)

Antibodies produced from affinity maturation of Joe 9 wild type were functionally characterized by surface plasmon resonance analysis to determine the K_d and K_{off} rate. A series of antibodies were produced having a K_{off} rate within the range 5 of about 0.1 s^{-1} to about $1 \times 10^{-5}\text{ s}^{-1}$, and more preferably a K_{off} of about $1 \times 10^{-4}\text{ s}^{-1}$ to $1 \times 10^{-5}\text{ s}^{-1}$ or less. Antibodies were also characterized *in vitro* for their ability to inhibit phytohemagglutinin (PHA) blast proliferation, as described in Example 3 of US Patent 10 No. 6,914,128. A series of antibodies were produced having an IC_{50} value in the range of about $1 \times 10^{-6}\text{ M}$ to about $1 \times 10^{-11}\text{ M}$, more preferably about $1 \times 10^{-10}\text{ M}$ to $1 \times 10^{-11}\text{ M}$ or less.

Accordingly, in one aspect, the invention provides methods and compositions for using an isolated human antibody, or antigen-binding portion thereof, that binds to human IL-12 and dissociates from human IL-12 with a K_{off} rate constant of 0.1 s^{-1} or less, as determined by surface plasmon resonance, or which inhibits phytohemagglutinin 15 blast proliferation in an *in vitro* phytohemagglutinin blast proliferation assay (PHA assay) with an IC_{50} of $1 \times 10^{-6}\text{ M}$ or less. In preferred embodiments, the isolated human IL-12 antibody, or an antigen-binding portion thereof, dissociates from human IL-12 with a K_{off} rate constant of $1 \times 10^{-2}\text{ s}^{-1}$ or less, or inhibits phytohemagglutinin blast 20 proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-7}\text{ M}$ or less. In more preferred embodiments, the isolated human IL-12 antibody, or an antigen-binding

portion thereof, dissociates from human IL-12 with a K_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, or inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-8} \text{ M}$ or less. In more preferred embodiments, the isolated human IL-12 antibody, or an antigen-binding portion thereof, dissociates from human IL-12 with a K_{off} rate constant of $1 \times 10^{-4} \text{ s}^{-1}$ or less, or inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-9} \text{ M}$ or less. In more preferred embodiments, the isolated human IL-12 antibody, or an antigen-binding portion thereof, dissociates from human IL-12 with a K_{off} rate constant of $1 \times 10^{-5} \text{ s}^{-1}$ or less, or inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-10} \text{ M}$ or less. In even more preferred embodiments, the isolated human IL-12 antibody, or an antigen-binding portion thereof, dissociates from human IL-12 with a K_{off} rate constant of $1 \times 10^{-5} \text{ s}^{-1}$ or less, or inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of $1 \times 10^{-11} \text{ M}$ or less.

The dissociation rate constant (K_{off}) of an IL-12 antibody can be determined by surface plasmon resonance (see Example 5 of US Patent No. 6,914,128). Generally, surface plasmon resonance analysis measures real-time binding interactions between ligand (recombinant human IL-12 immobilized on a biosensor matrix) and analyte (antibodies in solution) by surface plasmon resonance (SPR) using the BiAcore system (Pharmacia Biosensor, Piscataway, NJ). Surface plasmon analysis can also be performed by immobilizing the analyte (antibodies on a biosensor matrix) and presenting the ligand (recombinant IL-12 in solution). Neutralization activity of IL-12 antibodies, or antigen binding portions thereof, can be assessed using one or more of several suitable *in vitro* assays (see Example 3 of US Patent No. 6,914,128).

It is well known in the art that antibody heavy and light chain CDRs play an important role in the binding specificity/affinity of an antibody for an antigen. Accordingly, the invention encompasses human antibodies having light and heavy chain CDRs of Joe 9, as well as other antibodies having CDRs that have been modified to improve the binding specificity/affinity of the antibody. As demonstrated in Example 1 of US Patent No. 6,914,128, a series of modifications to the light and heavy chain CDRs results in affinity maturation of human anti-hIL-12 antibodies. The heavy and light chain variable region amino acid sequence alignments of a series of human antibodies ranging from Joe 9 wild type to J695 that bind human IL-12 is shown in Figures 1A-1D

of US Patent No. 6,914,128. Consensus sequence motifs for the CDRs of antibodies can be determined from the sequence alignment. For example, a consensus motif for the VH CDR3 of the lineage from Joe 9 to J695 comprises the amino acid sequence: (H/S)-G-S-(H/Y)-D-(N/T/Y) (SEQ ID NO: 1), which encompasses amino acids from position 95 to

5 102 of the consensus HCVR shown in SEQ ID NO: 7. A consensus motif for the VL CDR3 comprises the amino acid sequence: Q-(S/T)-Y-(D/E)-(S/R/K)-(S/G/Y)-(L/F/T/S)-(R/S/T/W/H)-(G/P)-(S/T/A/L)-(R/S/M/T/L-V/I/T/M/L) (SEQ ID NO: 2), which encompasses amino acids from position 89 to 97 of the consensus LCVR shown in SEQ ID NO: 8.

10 Accordingly, in another aspect, the invention provides methods and compositions comprising an isolated human antibody, or an antigen-binding portion thereof, which has the following characteristics:

- a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁶ M or less;
- 15 b) has a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 1; and
- c) has a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 2.

20 In a preferred embodiment, the antibody further comprises a VH CDR2 comprising the amino acid sequence: F-I-R-Y-D-G-S-N-K-Y-Y-A-D-S-V-K-G (SEQ ID NO: 3) (which encompasses amino acids from position 50 to 65 of the consensus HCVR comprising the amino acid sequence SEQ ID NO: 7) and further comprises a VL CDR2 comprising the amino acid sequence: (G/Y)-N-(D/S)-(Q/N)-R-P-S (SEQ ID NO: 4) (which encompasses amino acids from position 50 to 56 of the consensus LCVR 25 comprising the amino acid sequence SEQ ID NO: 8).

25 In another preferred embodiment, the antibody further comprises a VH CDR1 comprising the amino acid sequence: F-T-F-S-(S/E)-Y-G-M-H (SEQ ID NO: 5) (which encompasses amino acids from position 27 to 35 of the consensus HCVR comprising the amino acid sequence SEQ ID NO: 7) and further comprises a VL CDR1 comprising the 30 amino acid sequence: (S/T)-G-(G/S)-(R/S)-S-N-I-(G/V)-(S/A)-(N/G/Y)-(T/D)-V-(K/H) (SEQ ID NO: 6) (which encompasses amino acids from position 24 to 34 of the consensus LCVR comprising the amino acid sequence SEQ ID NO: 8).

In yet another preferred embodiment, the antibody used in the invention comprises a HCVR comprising the amino acid sequence of SEQ ID NO: 7 and a LCVR comprising the amino acid sequence of SEQ ID NO: 8.

Additional consensus motifs can be determined based on the mutational analysis 5 performed on Y61 that led to the J695 antibody (summarized in Figures 2A-2H of US Patent No. 6,914,128). As demonstrated by the graphs shown in Figures 2A-2H of US Patent No. 6,914,128, certain residues of the heavy and light chain CDRs of Y61 were amenable to substitution without significantly impairing the hIL-12 binding properties of the antibody. For example, individual substitutions at position 30 in CDR H1 with 10 twelve different amino acid residues did not significantly reduce the K_{off} rate of the antibody, indicating that this position is amenable to substitution with a variety of different amino acid residues. Thus, based on the mutational analysis (i.e., positions within Y61 that were amenable to substitution by other amino acid residues) consensus motifs were determined. The consensus motifs for the heavy and light chain CDR3s are 15 shown in SEQ ID NOS: 9 and 10, respectively, consensus motifs for the heavy and light chain CDR2s are shown in SEQ ID NOS: 11 and 12, respectively, and consensus motifs for the heavy and light chain CDR1s are shown in SEQ ID NOS: 13 and 14, respectively. Consensus motifs for the VH and VL regions are shown in SEQ ID NOS: 16 and 17, respectively.

20 Accordingly, in one aspect, the invention includes an isolated human antibody, or an antigen-binding portion thereof, which has the following characteristics:

- a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC_{50} of 1×10^{-9} M or less;
- b) has a heavy chain CDR3 comprising the amino acid sequence of SEQ ID 25 NO: 9; and
- c) has a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 10.

In a preferred embodiment, the antibody further comprises a VH CDR2 comprising the amino acid sequence of SEQ ID NO: 11 and further comprises a VL 30 CDR2 comprising the amino acid sequence of SEQ ID NO: 12.

In another preferred embodiment, the antibody further comprises a VH CDR1 comprising the amino acid sequence of SEQ ID NO: 13 and further comprises a VL CDR1 comprising the amino acid sequence of SEQ ID NO: 14.

In yet another preferred embodiment, the antibody used in the invention comprises a HCVR comprising the amino acid sequence of SEQ ID NO: 15 and a LCVR comprising the amino acid sequence of SEQ ID NO: 16.

A preferred antibody used in the invention, the human anti-hIL-12 antibody Y61, 5 can be produced by affinity maturation of Joe 9 wild type by PCR mutagenesis of the CDR3 (as described in Example 1 of US Patent No. 6,914,128). Y61 had an improved specificity/binding affinity determined by surface plasmon resonance and by *in vitro* neutralization assays. The heavy and light chain CDR3s of Y61 are shown in SEQ ID NOs: 17 and 18, respectively, the heavy and light chain CDR2s of Y61 are shown in 10 SEQ ID NOs: 19 and 20, respectively, and the heavy and light chain CDR1s of Y61 are shown in SEQ ID NOs: 21 and 22, respectively. The VH of Y61 has the amino acid sequence of SEQ ID NO: 23 and the VL of Y61 has the amino acid sequence of SEQ ID NO: 24 (these sequences are also shown in Figures 1A-1D of US Patent No. 6,914,128, aligned with Joe9).

15 Accordingly, in another aspect, the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which

- a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less;
- b) has a heavy chain CDR3 comprising the amino acid sequence of SEQ ID 20 NO: 17; and
- c) has a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 18.

In a preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the methods and compositions of the invention has a heavy 25 chain CDR2 comprising the amino acid sequence of SEQ ID NO: 19 and a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 20.

In another preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the methods and compositions of the invention, has a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 21 and a light 30 chain CDR1 comprising the amino acid sequence of SEQ ID NO: 22.

In yet another preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the methods and compositions of the invention comprising a the heavy chain variable region comprising the amino acid sequence of

SEQ ID NO: 23, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 24.

In certain embodiments, the full length antibody comprises a heavy chain constant region, such as IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant regions, 5 and any allotypic variant therein as described in Kabat (Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Preferably, the antibody heavy chain constant region is an IgG1 heavy chain constant region. Alternatively, the antibody portion can be an Fab fragment, an F(ab')₂ fragment or a single chain Fv 10 fragment.

Modifications of individual residues of Y6I led to the production of a panel of antibodies shown in Figures 2A-2H of US Patent No. 6,914,128. The specificity/binding affinity of each antibody was determined by surface plasmon resonance and/or by *in vitro* neutralization assays.

15 Accordingly, in another aspect, the invention features an isolated human antibody, or an antigen-binding portion thereof, which

- a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less;
- b) has a heavy chain CDR3 comprising the amino acid sequence selected 20 from the group consisting of SEQ ID NO: 404-SEQ ID NO: 469; and
- c) has a light chain CDR3 comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 534-SEQ ID NO: 579.

In preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the methods and compositions of the invention has a heavy 25 chain CDR2 comprising the amino acid sequence selected from the group consisting of SEQ ID NO:335-SEQ ID NO: 403; and a light chain CDR2 comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 506-SEQ ID NO: 533.

In another preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, has a heavy chain CDR1 comprising the amino acid sequence 30 selected from the group consisting of SEQ ID NO: 288-SEQ ID NO: 334; and a light chain CDR1 comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 470-SEQ ID NO: 505.

In yet another preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, comprising a the heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 23, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 24.

5 In certain embodiments, the full length antibody comprising a heavy chain constant region such as IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant regions and any allotypic variant therein as described in Kabat (, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Preferably, the antibody
10 heavy chain constant region is an IgG1 heavy chain constant region. Alternatively, the antibody portion can be a Fab fragment, an F(ab')₂ fragment or a single chain Fv fragment.

A particularly preferred recombinant, neutralizing antibody, J695, which may be used in the invention was produced by site-directed mutagenesis of contact and
15 hypermutation amino acids residues of antibody Y61 (see Example 2 of US Patent No. 6,914,128 and section III below). J695 differs from Y61 by a Gly to Tyr substitution in Y61 at position 50 of the light chain CDR2 and by a Gly to Tyr substitution at position 94 of the light chain CDR3. The heavy and light chain CDR3s of J695 are shown in SEQ ID NOs: 25 and 26, respectively, the heavy and light chain CDR2s of J695 are
20 shown in SEQ ID NOs: 27 and 28, respectively, and the heavy and light chain CDR1s of J695 are shown in SEQ ID NOs: 29 and 30, respectively. The VH of J695 has the amino acid sequence of SEQ ID NO: 31 and the VL of J695 has the amino acid sequence of SEQ ID NO: 32 (these sequences are also shown in Figures 1A-ID of US Patent No. 6,914,128, aligned with Joe9).

25 Accordingly, in another aspect, the invention features an isolated human antibody, or an antigen-binding portion thereof, which a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less; b) has a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 25; and c) has a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26.

30 In preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the invention has a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 27, and a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 28.

In another preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the invention has a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 29, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 30.

5 In yet another preferred embodiment, the isolated human antibody, or an antigen-binding portion thereof, used in the invention has a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 31, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 32.

In certain embodiments, the full length antibody comprises a heavy chain 10 constant region, such as IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant regions and any allotypic variant therein as described in Kabat (, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Preferably, the antibody heavy chain constant region is an IgG1 heavy chain constant region. Alternatively, the 15 antibody portion can be an Fab fragment, an F(ab')₂ fragment or a single chain Fv fragment.

Additional mutations in the preferred consensus sequences for CDR3, CDR2, and CDR1 of antibodies on the lineage from Joe 9 to J695, or from the lineage Y61 to J695, can be made to provide additional anti-IL-12 antibodies of the invention. Such 20 methods of modification can be performed using standard molecular biology techniques, such as by PCR mutagenesis, targeting individual contact or hypermutation amino acid residues in the light chain and/or heavy chain CDRs-, followed by kinetic and functional analysis of the modified antibodies as described herein (e.g., neutralization assays described in Example 3 of US Patent No. 6,914,128, and by BIACore analysis, as 25 described in Example 5 of US Patent No. 6,914,128).

Accordingly, in another aspect the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which

- a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁶ M or less;
- 30 b) comprises a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 1, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 3 and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 5, or a mutant thereof having one or more amino acid substitutions at a preferred selective

mutagenesis position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 1, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 3, and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 5; and

5 sequence of SEQ ID NO: 5; and

c) comprises a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 2, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO:

4, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 6, or a mutant thereof having one or more amino acid substitutions at a preferred selective

10 mutagenesis position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a light chain CDR3 comprising

the amino acid sequence of SEQ ID NO: 2, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 4, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 6.

15 In another aspect the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which

a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less;

b) comprises a heavy chain CDR3 comprising the amino acid sequence of
20 SEQ ID NO: 9, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID

NO: 11 and a heavy chain CDRI comprising the amino acid sequence of SEQ ID NO: 13, or a mutant thereof having one or more amino acid substitutions at a preferred selective mutagenesis position, contact position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a

25 heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 9, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 11, and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 13; and

c) comprises a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 10, a light chain CDR2 comprising the amino acid sequence of SEQ ID

30 NO: 12, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 14, or a mutant thereof having one or more amino acid substitutions at a preferred selective mutagenesis position, contact position or a hypermutation position, wherein

light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 10, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 12, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 14.

An ordinarily skilled artisan will also appreciate that additional mutations to the 5 CDR regions of an antibody, for example in Y61 or in J695, can be made to provide additional anti-IL-12 antibodies of the invention. Such methods of modification can be performed using standard molecular biology techniques, as described above. The functional and kinetic analysis of the modified antibodies can be performed as described in Example 3 of US Patent No. 6,914,128 and Example 5 of US Patent No. 6,914,128, 10 respectively. Modifications of individual residues of Y61 that led to the identification of J695 are shown in Figures 2A-2H of US Patent No. 6,914,128 and are described in Example 2 of US Patent No. 6,914,128.

Accordingly, in another aspect the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which

- 15 a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less;
- b) comprises a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 17, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 19 and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 21, or a mutant thereof having one or more amino acid substitutions at a preferred 20 selective mutagenesis position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 17, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 19, and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 21; and
- 25 c) comprises a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 18, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 20, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 22, or a mutant thereof having one or more amino acid substitutions at a preferred 30 selective mutagenesis position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 18, a light chain CDR2 comprising

the amino acid sequence of SEQ ID NO: 20, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 22.

In another aspect the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which

- 5 a) inhibits phytohemagglutinin blast proliferation in an *in vitro* PHA assay with an IC₅₀ of 1 x 10⁻⁹ M or less;
- 10 b) comprises a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 25, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 27 and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 29, or a mutant thereof having one or more amino acid substitutions at a preferred selective mutagenesis position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 25, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 27, and a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 29; and
- 15 c) comprises a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 28, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 30, or a mutant thereof having one or more amino acid substitutions at a preferred selective mutagenesis position or a hypermutation position, wherein said mutant has a k_{off} rate no more than 10-fold higher than the antibody comprising a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 28, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 30.

25 In yet another embodiment, the invention provides use of an isolated human antibodies, or antigen-binding portions thereof, that neutralize the activity of human IL-12, and at least one additional primate IL-12 selected from the group consisting of baboon IL-12, marmoset IL-12, chimpanzee IL-12, cynomolgus IL-12 and rhesus IL-12, but which do not neutralize the activity of the mouse IL-12.

30

II Selection of Recombinant Human Antibodies

Recombinant human antibodies which may be used in the invention can be isolated by screening of a recombinant combinatorial antibody library, preferably a scFv

phage display library, prepared using human VL and VH cDNAs prepared from mRNA derived from human lymphocytes. Methods for identifying antibodies which may be used in the methods and compositions of the invention are described in US Patent No. 6,914,128, incorporated by reference herein. Methodologies for preparing and screening 5 such libraries are known in the art. In addition to commercially available kits for generating phage display libraries (e.g., the Pharmacia *Recombinant Phage Antibody System*, catalog no. 27-9400-01; and the Stratagene *SurfZAP*™ phage display kit, catalog no. 240612), examples of methods and reagents particularly amenable for use in generating and screening antibody display libraries can be found in, for example, Kang 10 *et al.* PCT Publication No. WO 92/18619; Winter *et al.* PCT Publication No. WO 92/20791; Breitling *et al.* PCT Publication No. WO 93/01288; McCafferty *et al.* PCT Publication No. WO 92/01047; Garrard *et al.* PCT Publication No. WO 92/09690; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; McCafferty *et al.*, 15 *Nature* (1990) 348:552-554; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982.

20 The antibody libraries used in this method are preferably scFv libraries prepared from human VL and VH cDNAs. The scFv antibody libraries are preferably screened using recombinant human IL-12 as the antigen to select human heavy and light chain sequences having a binding activity toward IL-12. To select for antibodies specific for the p35 subunit of IL-12 or the p70 heterodimer, screening assays were performed in the 25 presence of excess free p40 subunit. Subunit preferences can be determined, for example by, micro-Friguet titration, as described in Example 1 of US Patent No. 6,914,128.

Once initial human VL and VH segments are selected, "mix and match" experiments, in which different pairs of the selected VL and VH segments are screened 30 for IL-12 binding, are performed to select preferred VL/VH pair combinations (see Example 1 of US Patent No. 6,914,128). Additionally, to further improve the affinity and/or lower the off rate constant for hIL-12 binding, the VL and VH segments of the preferred VL/VH pair(s) can be randomly mutated, preferably within the CDR3 region

of VH and/or VL, in a process analogous to the *in vivo* somatic mutation process responsible for affinity maturation of antibodies during a natural immune response. This *in vitro* affinity maturation can be accomplished by amplifying VH and VL regions using PCR primers complimentary to the VH CDR3 or VL CDR3, respectively, which 5 primers have been "spiked" with a random mixture of the four nucleotide bases at certain positions such that the resultant PCR products encode VH and VL segments into which random mutations have been introduced into the VH and/or VL CDR3 regions. These randomly mutated VH and VL segments can be reselected and rescreened for 10 binding to hIL-12 and sequences that exhibit high affinity and a low off rate for IL-12 binding can be selected. table 2 (see Appendix A of US Patent No. 6,914,128) shows 15 antibodies that displayed altered binding specificity/affinity produced as a result of *in vitro* affinity maturation.

Following selection, isolation and screening of an anti-hIL-12 antibody of the 15 invention from a recombinant immunoglobulin display library, nucleic acid encoding the selected antibody can be recovered from the phage particle(s) (e.g., from the phage genome) and subcloned into other expression vectors by standard recombinant DNA techniques. If desired, the nucleic acid can be further manipulated to create other 20 antibody forms of the invention (e.g., linked to nucleic acid encoding additional immunoglobulin domains, such as additional constant regions). To express a recombinant human antibody isolated by screening of a combinatorial library, the DNA 25 encoding the antibody is cloned into a recombinant expression vector and introduced into a mammalian host cells, as described in further detail in Section IV below.

Methods for selecting human IL-12 binding antibodies by phage display 25 technology, and affinity maturation of selected antibodies by random or site-directed mutagenesis of CDR regions are described in further detail in Example 1 of US Patent No. 6,914,128.

As described in Example 1 of US Patent No. 6,914,128, screening of human VL 30 and VH cDNA libraries identified a series of anti-IL-12 antibodies, of which the Joe 9 antibody was selected for further development. A comparison of the heavy chain variable region of Joe 9 with the heavy chain germline sequences selected from the VBASE database, revealed that Joe 9 was similar to the COS-3 germline sequence. COS-3 belongs to the V_H3 family of germline sequences.

The $V_{H}3$ family is part of the human VH germline repertoire which is grouped into seven families, $V_{H}1-V_{H}7$, based on nucleotide sequence homology (Tomlinson *et al.* (1992) *J. Mol. Biol.*, 227, 776-798 and Cook *et al.* (1995) *Immunology Today*, 16, 237-242). The $V_{H}3$ family contains the highest number of members and makes the 5 largest contribution to the germline repertoire. For any given human $V_{H}3$ - germline antibody sequence, the amino acid sequence identity within the entire $V_{H}3$ family is high (See e.g., Tomlinson *et al.* (1992) *J. Mol. Biol.*, 227, 776-798 and Cook *et al.* (1995) *Immunology Today*, 16, 237-242). The range of amino acid sequence identity between any two germline VH sequences of the $V_{H}3$ family varies from 69-98 residues 10 out of approximately 100 VH residues, (*i.e.*, 69-98% amino acid sequence homology between any two germline VH sequences). For most pairs of germline sequences there is at least 80 or more identical amino acid residues, (*i.e.*, at least 80% amino acid sequence homology). The high degree of amino acid sequence homology between the $V_{H}3$ family members results in certain amino acid residues being present at key sites in 15 the CDR and framework regions of the VH chain. These amino acid residues confer structural features upon the CDRs.

Studies of antibody structures have shown that CDR conformations can be grouped into families of canonical CDR structures based on the key amino acid residues that occupy certain positions in the CDR and framework regions. Consequently, there 20 are similar local CDR conformations in different antibodies that have canonical structures with identical key amino acid residues (Chothia *et al.* (1987) *J. Mol. Biol.*, 196, 901-917 and Chothia *et al.* (1989) *Nature*, 342, 877-883). Within the $V_{H}3$ family there is a conservation of amino acid residue identity at the key sites for the CDR1 and CDR2 canonical structures (Chothia *et al.* (1992) *J. Mol. Biol.*, 227, 799-817).

25 The COS-3 germline VH gene, is a member of the $V_{H}3$ family and is a variant of the 3-30 (DP-49) germline VH allele. COS-3, differs from Joe9 VH amino acid sequences at only 5 positions. The high degree of amino acid sequence homology between Joe9 VH and COS-3, and between Joe9 VH and the other $V_{H}3$ family members also confers a high degree of CDR structural homology (Chothia *et al.* (1992) *J. Mol. Biol.*, 227, 799-817; Chothia *et al.* (1987) *J. Mol. Biol.*, 196, 901-917 and Chothia *et al.* (1989) *Nature*, 342, 877-883). 30

The skilled artisan will appreciate that based on the high amino acid sequence and canonical structural similarity to Joe 9, other V_{H3} family members could also be used to generate antibodies that bind to human IL-12. This can be performed, for example, by selecting an appropriate VL by chain-shuffling techniques (Winter *et al.* 5 (1994) *Annual Rev. Immunol.*, 12, 433-55), or by the grafting of CDRs from a rodent or other human antibody including CDRs from antibodies of this invention onto a V_{H3} family framework.

The human V lambda germline repertoire is grouped into 10 families based on nucleotide sequence homology (Williams *et al.* (1996) *J. Mol. Biol.*, 264, 220-232). A 10 comparison of the light chain variable region of Joe 9 with the light chain germline sequences selected from the VBASE database, revealed that Joe 9 was similar to the DPL8 lambda germline. The Joe9 VL differs from DPL8 sequence at only four framework positions, and is highly homologous to the framework sequences of the other $V_{\lambda}1$ family members. Based on the high amino acid sequence homology and canonical 15 structural similarity to Joe 9, other $V_{\lambda}1$ family members may also be used to generate antibodies that bind to human IL-12. This can be performed, for example, by selecting an appropriate VH by chain-shuffling techniques (Winter *et al. Supra*, or by the grafting of CDRs from a rodent or other human antibody including CDRs from antibodies of this invention onto a $V_{\lambda}1$ family framework.

20 The methods of the invention are intended to include recombinant antibodies that bind to hIL-12, comprising a heavy chain variable region derived from a member of the V_{H3} family of germline sequences, and a light chain variable region derived from a member of the $V_{\lambda}1$ family of germline sequences. Moreover, the skilled artisan will appreciate that any member of the V_{H3} family heavy chain sequence can be combined 25 with any member of the $V_{\lambda}1$ family light chain sequence.

Those skilled in the art will also appreciate that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the germline may exist within a population (e.g., the human population). Such genetic polymorphism in the germline sequences may exist among individuals within a population due to natural allelic 30 variation. Such natural allelic variations can typically result in 1-5 % variance in the nucleotide sequence of the a gene. Any and all such nucleotide variations and resulting

amino acid polymorphisms in germline sequences that are the result of natural allelic variation are intended to be within the scope of the invention.

Accordingly, in one aspect, the invention features an isolated human antibody, or an antigen-binding portion thereof, which has the following characteristics:

- 5 a) that binds to human IL-12 and dissociates from human IL-12 with a k_{off} rate constant of 0.1 s^{-1} or less, as determined by surface plasmon resonance, or which inhibits phytohemagglutinin blast proliferation in an *in vitro* phytohemagglutinin blast proliferation assay (PHA assay) with an IC_{50} of $1 \times 10^{-6}\text{M}$ or less.
- 10 b) has a heavy chain variable region comprising an amino acid sequence selected from a member of the V_H3 germline family, wherein the heavy chain variable region has a mutation at a contact or hypermutation position with an activity enhancing amino acid residue.
- 15 c) has a light chain variable region comprising an amino acid sequence selected from a member of the $V_{\lambda}1$ germline family, wherein the light chain variable region has a mutation at a preferred selective mutagenesis position, contact or hypermutation position with an activity enhancing amino acid residue.

In a preferred embodiment, the isolated human antibody, or antigen binding has mutation in the heavy chain CDR3. In another preferred embodiment, the isolated human antibody, or antigen binding has mutation in the light chain CDR3. In another preferred embodiment, the isolated human antibody, or antigen binding has mutation in the heavy chain CDR2. In another preferred embodiment, the isolated human antibody, or antigen binding has mutation in the light chain CDR2. In another preferred embodiment, the isolated human antibody, or antigen binding has mutation in the heavy chain CDR1. In another preferred embodiment, the isolated human antibody, or antigen binding has mutation in the light chain CDR1.

An ordinarily skilled artisan will appreciate that based on the high amino acid sequence similarity between members of the V_H3 germline family, or between members of the light chain $V_{\lambda}1$ germline family, that mutations to the germlines sequences can provide additional antibodies that bind to human IL-12. table 1 of US Patent No. 30 6,914,128 (see also Appendix A of US Patent No. 6,914,128) shows the germline sequences of the V_H3 family members and demonstrates the significant sequence homology within the family members. Also shown in table 1 of US Patent No.

6,914,128 are the germline sequences for $V_{\lambda}1$ family members. The heavy and light chain sequences of Joe 9 are provided as a comparison. Mutations to the germline sequences of V_H3 or $V_{\lambda}1$ family members may be made, for example, at the same amino acid positions as those made in the antibodies of the invention (e.g. mutations in 5 Joe 9). The modifications can be performed using standard molecular biology techniques, such as by PCR mutagenesis, targeting individual amino acid residues in the germline sequences, followed by kinetic and functional analysis of the modified antibodies as described herein (e.g., neutralization assays described in Example 3 of US Patent No. 6,914,128, and by BIACore analysis, as described in Example 5 of US Patent 10 No. 6,914,128).

Accordingly, in one aspect, the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which has the following characteristics:

- a) has a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 595-667, wherein the heavy chain 15 variable region has a mutation at a preferred selective mutagenesis position, contact or hypermutation position with an activity enhancing amino acid residue.
- b) has a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 669-675, wherein the light chain variable region has a mutation at a preferred selective mutagenesis position, contact or 20 hypermutation position with an activity enhancing amino acid residue.

An ordinarily skilled artisan will appreciate that based on the high amino acid sequence similarity between Joe 9 and COS-3 heavy chain germline sequence, and between Joe 9 and DPL8 lambda germline sequence, that other mutations to the CDR regions of these germlines sequences can provide additional antibodies that bind to 25 human IL-12. Such methods of modification can be performed using standard molecular biology techniques as described above.

Accordingly, in one aspect, the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which has the following characteristics:

- a) that binds to human IL-12 and dissociates from human IL-12 with a 30 k_{off} rate constant of 0.1s^{-1} or less, as determined by surface plasmon resonance, or which inhibits phytohemagglutinin blast proliferation in an *in vitro* phytohemagglutinin blast proliferation assay (PHA assay) with an IC_{50} of $1 \times 10^{-6}\text{M}$ or less.

b) has a heavy chain variable region comprising the COS-3 germline amino acid sequence, wherein the heavy chain variable region has a mutation at a preferred selective mutagenesis position, contact or hypermutation position with an activity enhancing amino acid residue.

5 c) has a light chain variable region comprising the DPL8 germline amino acid sequence, wherein the light chain variable region has a mutation at a preferred selective mutagenesis position, contact or hypermutation position with an activity enhancing amino acid residue.

Due to certain amino acid residues occupying key sites in the CDR and
10 framework regions in the light and heavy chain variable region, structural features are conferred at these regions. In particular, the CDR2 and CDR1 regions are subject to canonical structural classifications. Since there is a high degree of amino acids sequence homology between family members, these canonical features are present between family members. The skilled artisan will appreciate that modifications at the
15 amino acid residues that confer these canonical structures would produce additional antibodies that bind to IL-12. The modifications can be performed using standard molecular biology techniques as described above.

Accordingly, in another aspect, the invention features use of an isolated human antibody, or an antigen-binding portion thereof, which has the following characteristics:

20 a) that binds to human IL-12 and dissociates from human IL-12 with a k_{off} rate constant of 0.1 s^{-1} or less, as determined by surface plasmon resonance, or which inhibits phytohemagglutinin blast proliferation in an *in vitro* phytohemagglutinin blast proliferation assay (PHA assay) with an IC_{50} of $1 \times 10^{-6}\text{M}$ or less.

25 b) has a heavy chain variable region comprising an amino acid sequence selected from a member of the V_H3 germline family, wherein the heavy chain variable region comprises a CDR2 that is structurally similar to CDR2s from other V_H3 germline family members, and a CDR1 that is structurally similar to CDR1s from other V_H3 germline family members, and wherein the heavy chain variable region has a mutation at a preferred selective mutagenesis position, contact or hypermutation position with an
30 activity enhancing amino acid residue;

c) has a light chain variable region comprising an amino acid sequence selected from a member of the $V_{\lambda}1$ germline family, wherein the light chain variable region comprises a CDR2 that is structurally similar to CDR2s from other $V_{\lambda}1$ germline

family members, and a CDR1 that is structurally similar to CDR1s from other $V\lambda 1$ germline family members, and wherein the light chain variable region has a mutation at a preferred selective mutagenesis position, contact or hypermutation position with an activity enhancing amino acid residue.

5 Recombinant human antibodies used in the invention have variable and constant regions which are homologous to human germline immunoglobulin sequences selected from the VBASE database. Mutations to the recombinant human antibodies (e.g., by random mutagenesis or PCR mutagenesis) result in amino acids that are not encoded by human germline immunoglobulin sequences. Also, libraries of recombinant antibodies
10 which were derived from human donors will contain antibody sequences that differ from their corresponding germline sequences due to the normal process of somatic mutation that occurs during B-cell development. It should be noted that if the "germline" sequences obtained by PCR amplification encode amino acid differences in the framework regions from the true germline configuration (i.e., differences in the
15 amplified sequence as compared to the true germline sequence), it may be desirable to change these amino acid differences back to the true germline sequences (i.e., "backmutation" of framework residues to the germline configuration). Thus, the present invention can optionally include a backmutation step. To do this, the amino acid sequences of heavy and light chain encoded by the germline (as found as example in
20 VBASE database) are first compared to the mutated immunoglobulin heavy and light chain framework amino acid sequences to identify amino acid residues in the mutated immunoglobulin framework sequence that differ from the closest germline sequences. Then, the appropriate nucleotides of the mutated immunoglobulin sequence are mutated back to correspond to the germline sequence, using the genetic code to determine which
25 nucleotide changes should be made. Mutagenesis of the mutated immunoglobulin framework sequence is carried out by standard methods, such as PCR-mediated mutagenesis (in which the mutated nucleotides are incorporated into the PCR primers such that the PCR product contains the mutations) or site-directed mutagenesis. The role of each amino acid identified as candidate for backmutation should be investigated for a
30 direct or indirect role in antigen binding and any amino acid found after mutation to affect any desirable characteristic of the human antibody should not be included in the final human antibody; as an example, activity enhancing amino acids identified by the selective mutagenesis approach will not be subject to backmutation. Assays to determine

the characteristics of the antibody resulting from mutagenesis can include ELISA, competitive ELISA, *in vitro* and *in vivo* neutralization assays and/or (see e.g. Example 3 of US Patent No. 6,914,128) immunohistochemistry with tissue sections from various sources (including human, primate and/or other species).

5 To minimize the number of amino acids subject to backmutation those amino acid positions found to be different from the closest germline sequence but identical to the corresponding amino acid in a second germline sequence can remain, provided that the second germline sequence is identical and colinear to the sequence of the human antibody of the invention for at least 10, preferably 12 amino acids, on both sides of the 10 amino acid in question. This would assure that any peptide epitope presented to the immune system by professional antigen presenting cells in a subject treated with the human antibody of the invention would not be foreign but identical to a self-antigen, i.e. the immunoglobulin encoded by that second germline sequence. Backmutation may occur at any stage of antibody optimization; preferably, backmutation occurs directly 15 before or after the selective mutagenesis approach. More preferably, backmutation occurs directly before the selective mutagenesis approach.

III. Modifications to Preferred Selective Mutagenesis Positions, Contact and/or Hypermutation Positions

20 Typically, selection of antibodies with improved affinities can be carried out using phage display methods, as described in section 11 above and in US Patent No. 6,914,128, incorporated by reference herein. This can be accomplished by randomly mutating combinations of CDR residues and generating large libraries containing 25 antibodies of different sequences. However, for these selection methods to work, the antibody-antigen reaction must tend to equilibrium to allow, over time, preferential binding of higher affinity antibodies to the antigen. Selection conditions that would allow equilibrium to be established could not be determined (presumably due to additional non-specific interactions between the antigen and phage particle) when phage display methods were used to improve the affinity of selected anti-IL-12 antibodies, 30 upon attaining a certain level of affinity achieved (*i.e.*, that of antibody Y61). Accordingly, antibodies with even higher affinities could not be selected by phage display methods. Thus, for at least certain antibodies or antigens, phage display methods are limiting in their ability to select antibodies with a highly improved binding

specificity/affinity. Accordingly, a method termed Selective Mutagenesis Approach which does not require phage display affinity maturation of antibodies, was established to overcome this limitation and is provided by the invention. Although this Selective Mutagenesis Approach was developed to overcome limitations using the phage display system, it should be noted that this method can also be used with the phage display system. Moreover, the selective mutagenesis approach can be used to improve the activity of any antibody.

To improve the activity (e.g., affinity or neutralizing activity) of an antibody, ideally one would like to mutate every CDR position in both the heavy and light chains to every other possible amino acid residue. However, since there are, on average, 70 CDR positions within an antibody, such an approach would be very time consuming and labor intensive. Accordingly, the method of the invention allows one to improve the activity of the antibody by mutating only certain selected residues within the heavy and/or light chain CDRs. Furthermore, the method of the invention allows improvement in activity of the antibody without affecting other desirable properties of the antibody.

Determining which amino acid residues of an antibody variable region are in contact with an antigen cannot be accurately predicted based on primary sequence or their positions within the variable region. Nevertheless, alignments of sequences from antibodies with different specificities conducted by Kabat *et al.* have identified the CDRs as local regions within the variable regions which differ significantly among antibodies (Kabat *et al.* (1971) *Ann. NY Acad. Sci.* 190:382-393, , Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Structural studies have shown that the antigen binding surface is formed by amino acid residues present in the CDRs. Other amino acid residues outside the CDR are also known to play structural roles or be directly involved in antigen binding. Therefore, for each antigen-antibody pair, amino acid residues within and outside of the CDRs may be important.

The sequence alignment studies by Tomlison *et al* identified a number of positions in the heavy and light chain CDR1 and CDR2, and in a portion of the kappa chain CDR3 which are frequent sites of somatic mutation. (Tomlison *et al* (1996) *J. Mol. Biol.* 256: 813-817). In particular, positions H31, H31B, H33, H33B, H52B, H56, H58, L30, L31, L31A, L50, L53, L91, L92, L93 and L94 were identified as frequent sites for somatic mutation. However, this analysis excludes the important heavy chain

CDR3 regions, and sections of the light chain CDR3 which are known to lie in the center of an antibody binding site, and potentially provide important interactions with an antigen. Furthermore, Tomlison *et al.* propose that somatic diversity alone does not necessarily predict a role of a specific amino acid in antigen binding, and suggest 5 conserved amino acid residues that contact the antigen, and diverse amino acid residues which do not contact the antigen. This conclusion is further supported by mutational studies on the role of somatic mutations to antibody affinity (*Sharon, (1990), PNAS, 87:4814-7*). Nineteen somatic mutations in a high-affinity anti-p-azophenylarsonate (Ars) antibody were simultaneously replaced with their corresponding germline 10 residues, generating a germline version of the anti-Ars antibody which had a two-hundred fold loss in activity. The full affinity of the anti-Ars antibody could be recovered by restoring only three of the nineteen somatic mutations, demonstrating that many somatic mutations may be permitted that do not contribute to antigen binding 15 activity.

15 The result can be explained in part by the nature of antibody diversity itself. Immature B-cells may produce initially low affinity antibodies that recognize a number of self or non-self antigens. Moreover, antibodies may undergo in the course of affinity maturation sequence variations that may cause self-reactivity. Hypermutation of such 20 low affinity antibodies may serve to abolish self-reactivity ("negative selection") and increase affinity for the foreign antigen. Therefore, the analysis of primary and structural data of a large number of antibodies does not provide a method of predicting either (1) the role of somatic hyper-mutation sites in the affinity maturation process versus the process of decreasing affinity towards unwanted antigens, or (2) how a given amino acid contributes to the properties of a specific antigen-antibody pair.

25 Other attempts to address the role of specific amino acid residues in antigen recognition were made by analyzing a number of crystal structures of antigen-antibody complexes (MacCallum *et al.* (1996) *J. Mol. Biol.* 262: 732-745). The potential role of positions located within and outside the CDRs was indicated. Positions in CDRs involved in antigen binding in more than 10 of 26 analyzed structures included H31, 30 H33, H50, H52, H53, H54, H56, H58, H95, H96, H97, H98 and H100 in the heavy chain and L30A, L32, L91, L92, L93, L94, L96 in the light chain. However, the authors noted that prediction of antigen contacts using these and other structural data may over

and under predict contact positions, leading to the speculation that a different strategy may have to be applied to different antigens.

Pini *et al.* describe randomizing multiple residues in antibody CDR sequences in a large phage display library to rapidly increase antibody affinity (Pini *et al.* (1998) *J. Biol Chem.* 273: 21769-21776). However, the high affinity antibodies discussed by Pini *et al.* had mutations in a total of eight positions, and a reductionary analysis of which changes are absolutely required to improve affinity of the antibody becomes impractical because of the large number of possible combinations to be tested for the smallest number of amino acids required.

Furthermore, randomizing multiple residues may not necessarily preserve other desired properties of the antibody. Desirable properties or characteristics of an antibody are art-recognized and include for example, preservation of non-cross reactivity, e.g., with other proteins or human tissues and preservation of antibody sequences that are close to human germline immunoglobulin sequences improvement of neutralization potency. Other desirable properties or characteristics include ability to preserve species cross reactivity, ability to preserve epitope specificity and ability to preserve high expression levels of protein in mammalian cells. The desirable properties or characteristics can be observed or measured using art-recognized techniques including but not limited to ELISA, competitive ELISA, *in vitro* and *in vivo* neutralization assays (see e.g. Example 3 of US Patent No. 6,914,128), immunohistochemistry with tissue sections from different sources including human, primate or other sources as the need may be, and studies to expression in mammalian cells using transient expression or stable expression.

In addition, the method of Pini *et al* may introduce more changes than the minimal number actually required to improve affinity and may lead to the antibodies triggering anti-human-antibody (HAMA) formation in human subjects. Further, as discussed elsewhere, the phage display as demonstrated here, or other related method including ribosome display may not work appropriately upon reaching certain affinities between antibody and antigen and the conditions required to reach equilibrium may not be established in a reasonable time frame because of additional interactions including interactions with other phage or ribosome components and the antigen.

The ordinarily skilled artisan may glean interesting scientific information on the origin of antibody diversity from the teachings of the references discussed above. The

present invention, however, provides a method for increasing antibody affinity of a specific antigen-antibody pair while preserving other relevant features or desirable characteristics of the antibody. This is especially important when considering the desirability of imparting a multitude of different characteristics on a specific antibody

5 including antigen binding.

If the starting antibody has desirable properties or characteristics which need to be retained, a selective mutagenesis approach can be the best strategy for preserving these desirable properties while improving the activity of the antibody. For example, in the mutagenesis of Y61, the aim was to increase affinity for hIL-12, and to improve the

10 neutralization potency of the antibody while preserving desired properties. Desired properties of Y61 included (1) preservation of non-cross reactivity with other proteins or human tissues, (2) preservation of fine epitope specificity, i.e. recognizing a p40 epitope preferably in the context of the p70 (p40/p35) heterodimer, thereby preventing binding interference from free soluble p40; and (3) generation of an antibody with heavy and

15 light chain amino acid sequences that were as close as possible to their respective germline immunoglobulin sequences.

In one embodiment, the method of the invention provides a selective mutagenesis approach as a strategy for preserving the desirable properties or characteristics of the antibody while improving the affinity and/or neutralization potency. The term "selective mutagenesis approach" is as defined above and includes a method of individually mutating selected amino acid residues. The amino acid residues to be mutated may first be selected from preferred selective mutagenesis positions, then from contact positions, and then from hypermutation positions. The individual selected position can be mutated to at least two other amino acid residue and the effect of the

20 mutation both on the desired properties of the antibody, and improvement in antibody activity is determined.

The Selective Mutagenesis approach comprises the steps of:

selecting candidate positions in the order 1) preferred selective mutagenesis positions; 2) contact positions; 3) hypermutation positions and ranking the positions

25 based on the location of the position within the heavy and light chain variable regions of an antibody (CDR3 preferred over CDR2 preferred over CDR1);

individually mutating candidate preferred selective mutagenesis positions, hypermutation and/or contact positions in the order of ranking, to all possible other

amino acid residues and analyzing the effect of the individual mutations on the activity of the antibody in order to determine activity enhancing amino acid residues;

if necessary, making stepwise combinations of the individual activity enhancing amino acid residues and analyzing the effect of the various combinations on the activity of the antibodies; selecting mutant antibodies with activity enhancing amino acid residues and ranking the mutant antibodies based on the location and identity of the amino acid substitutions with regard to their immunogenic potential. Highest ranking is given to mutant antibodies that comprise an amino acid sequence which nearly identical to a variable region sequence that is described in a germline database, or has an amino acid sequence that is comparable to other human antibodies. Lower ranking is given to mutant antibodies containing an amino acid substitution that is rarely encountered in either germline sequences or the sequences of other human antibodies. The lowest ranking is given to mutant antibodies with an amino acid substitution that has not been encountered in a germline sequence or the sequence of another human antibody. As set forth above, mutant antibodies comprising at least one activity enhancing amino acid residue located in CDR3 is preferred over CDR2 which is preferred over CDR1. The CDRs of the heavy chain variable regions are preferred over those of the light chain variable region.

The mutant antibodies can also be studied for improvement in activity, *e.g.* when compared to their corresponding parental antibody. The improvement in activity of the mutant antibody can be determined for example, by neutralization assays, or binding specificity/affinity by surface plasmon resonance analysis (see Example 3 of US Patent No. 6,914,128). Preferably, the improvement in activity can be at least 2-20 fold higher than the parental antibody. The improvement in activity can be at least " x_1 " to " x_2 " fold higher than the parental antibody wherein " x_1 " and " x_2 " are integers between and including 2 to 20, including ranges within the state range, *e.g.* 2-15, *e.g.* 5-10.

The mutant antibodies with the activity enhancing amino acid residue also can be studied to determine whether at least one other desirable property has been retained after mutation. For example, with anti-hIL-12 antibodies testing for, (1) preservation of non-cross reactivity with other proteins or human tissues, (2) preservation of epitope recognition, *i.e.* recognizing a p40 epitope preferably in the context of the p70 (p40/p35) heterodimer, thereby preventing binding interference from free soluble p40; and (3) generation of antibodies with heavy and light chain amino acid sequences that were as

close as possible to their respective germline immunoglobulin sequences, and determining which would be least likely to elicit a human immune response based on the number of differences from the germline sequence. The same observations can be made on an antibody having more than one activity enhancing amino acid residues, e.g. at 5 least two or at least three activity enhancing amino acid residues, to determine whether retention of the desirable property or characteristic has occurred.

An example of the use of a "selective mutagenesis approach", in the mutagenesis of Y61 is described below. The individual mutations H31S→E, L50→Y, or L94G→Y each improved neutralization activity of the antibody. However, when combination 10 clones were tested, the activity of the combined clone H31S→E + L50→Y + L94G→Y was no better than L50→Y + L94G→Y (J695). Therefore, changing the germline amino acid residue Ser to Glu at position 31 of CDR1 was unnecessary for the improved activity of J695 over Y61. The selective mutagenesis approach therefore, identified the minimal number of changes that contributed to the final activity, thereby reducing the 15 immunogenic potential of the final antibody and preserving other desired properties of the antibody.

Isolated DNA encoding the VH and VL produced by the selected mutagenesis approach can be converted into full length antibody chain genes, to Fab fragment genes as to a scFV gene, as described in section IV. For expression of VH and VL regions 20 produced by the selected mutagenesis approach, expression vectors encoding the heavy and light chain can be transfected into variety host cells as described in detail in section IV. Preferred host cells include either prokaryotic host cells, for example, *E coli*, or eukaryotic host cells, for example, yeast cells, e.g., *S. cerevisiae*. Most preferred eukaryotic host cells are mammalian host cells, described in detail in section IV.

25 The selective mutagenesis approach provides a method of producing antibodies with improved activities without prior affinity maturation of the antibody by other means. The selective mutagenesis approach provides a method of producing antibodies with improved affinities which have been subject to back mutations. The selective mutagenesis approach also provides a method of improving the activity of affinity 30 matured antibodies.

The skilled artisan will recognize that the selective mutagenesis approach can be used in standard antibody manipulation techniques known in the art. Examples include, but are not limited to, CDR grafted antibodies, chimeric antibodies, scFV fragments,

Fab fragments of a full length antibodies and human antibodies from other sources, e.g., transgenic mice.

Rapid large scale mutational analysis of antibodies include *in vitro* transcription and translation using ribosome display technology (see e.g., Hanes *et al.*, (1997) *Proc.*

5 *Natl. Acad. Sci.* 94: 4937-4942; Dall Acqua *et al.*, (1998) *Curr. Opin. Struc. Biol.* 8: 443-450; He *et al.*, (1997) *Nucleic Acid Res.* 25: 5132-5134), and U.S. Patent Nos. 5,643,768 and 5,658,754 issued to Kawasaki. The selective mutagenesis approach also provides a method of producing antibodies with improved activities that can be selected using ribosomal display techniques.

10 In the methods of the invention, antibodies or antigen binding portions thereof are further modified by altering individual positions in the CDRs of the HCVR and/or LCVR. Although these modifications can be made in phage-displayed antibodies, the method is advantageous in that it can be performed with antibodies that are expressed in other types of host systems, such as bacterial, yeast or mammalian cell expression systems. The individual positions within the CDRs selected for modification are based 15 on the positions being a contact and/or hypermutation position.

Preferred contact positions and hypermutation positions as defined herein are shown in table 3 of US Patent No. 6,914,128 (see Appendix A of US Patent No. 6,914,128) and their modification in accordance with the method of the invention is 20 described in detail in Example 2 of US Patent No. 6,914,128. Preferred contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96. Preferred hypermutation positions are selected from the group consisting of H30, H31, H31B, H32, H52, H56, 25 H58, L30, L31, L32, L53 and L93. More preferred amino acid residues (referred to as “preferred selective mutagenesis positions”) are both contact and hypermutation positions and are selected from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94. Particularly preferred contact positions are selected from the group consisting of L50 and L94.

30 Preferred activity enhancing amino acid residues replace amino acid residues located at positions selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94, and L96. More preferred activity

enhancing amino acid residues replace amino acid residues located at positions H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94.

Particularly, preferred activity enhancing amino acid residues replace amino acid residues located at positions selected from the group consisting of L50 and L94.

5 In general, the method of the invention involves selecting a particular preferred selective mutagenesis position, contact and/or hypermutation position within a CDR of the heavy or light chain of a parent antibody of interest, or antigen binding portion thereof, randomly mutagenizing that individual position (e.g., by genetic means using a mutagenic oligonucleotide to generate a "mini-library" of modified antibodies), or

10 mutating a position to specific desired amino acids, to identify activity enhancing amino acid residues expressing, and purifying the modified antibodies (e.g., in a non-phage display host system), measuring the activity of the modified antibodies for antigen (e.g., by measuring k_{off} rates by BIAcore analysis), repeating these steps for other CDR positions, as necessary, and combining individual mutations shown to have improved

15 activity and testing whether the combination(s) generate an antibody with even greater activity (e.g., affinity or neutralizing potency) than the parent antibody, or antigen-binding portion thereof.

Accordingly, in one embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

20 a) providing a parent antibody or antigen-binding portion thereof;

 b) selecting in order a 1) preferred selective mutagenesis position, 2) contact position, or 3) hypermutation position within a complementarity determining region (CDR) for mutation, thereby identifying a selected preferred selective mutagenesis position, contact or hypermutation position;

25 c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof;

 d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof;

30 e) optionally, repeating steps a) through d) for at least one other preferred selective mutagenesis position, contact or hypermutation position;

- f) combining, in the parent antibody, or antigen-binding portion thereof, individual mutations shown to have improved activity, to form combination antibodies, or antigen-binding portions thereof; and
- g) evaluating the activity of the combination antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof; until an antibody, or antigen-binding portion thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained. Preferably, the selected antibody or antibodies have an improved activity without loss or with retention of at least one desirable characteristic or property of the parental antibody as described above. The desirable characteristic or property can be measured or observed by the ordinarily skilled artisan using art-recognized techniques.

Preferred contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96. Preferred hypermutation positions are selected from the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93. More preferred selective mutagenesis positions are selected from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93 and L94. Particularly preferred contact positions are selected from the group consisting of L50 and L94.

In another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a parent antibody or antigen-binding portion thereof;
- b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation;
- c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof;
- d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof,
- 30 thereby identifying an activity enhancing amino acid residue;
- e) optionally, repeating steps a) through d) for at least one other preferred selective mutagenesis position, contact or hypermutation position;

- f) combining, in the parent antibody, or antigen-binding portion thereof, two individual activity enhancing amino acid residues shown to have improved activity, to form combination antibodies, or antigen-binding portions thereof; and
 - g) evaluating the activity of the combination antibodies, or antigen-binding portions thereof with two activity enhancing amino acid residues, relative to the parent antibody or antigen-binding portion thereof;
- 5 until an antibody, or antigen-binding portion thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferred contact positions are selected from the group consisting of H30, H31, 10 H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96. Preferred hypermutation positions are selected from the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93. More preferred selective mutagenesis positions are selected from the group consisting of H30, H31, H31B, H32, 15 H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93 and L94. Particularly preferred contact positions are selected from the group consisting of L50 and L94.

In another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a parent antibody or antigen-binding portion thereof;
- 20 b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation;
- c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof;
- 25 d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof, thereby identifying an activity enhancing amino acid residue;
- e) optionally, repeating steps a) through d) for at least one other preferred selective mutagenesis position, contact or hypermutation position;
- 30 f) combining, in the parent antibody, or antigen-binding portion thereof, three individual activity enhancing amino acid residues shown to have improved activity, to form combination antibodies, or antigen-binding portions thereof; and

g) evaluating the activity of the combination antibodies, or antigen-binding portions thereof with two activity enhancing amino acid residues, relative to the parent antibody or antigen-binding portion thereof;
until an antibody, or antigen-binding portion thereof, with an improved activity, relative
5 to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferably, the activity enhancing amino acid residue replaces amino acid residues located at positions selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96.

10 Following mutagenesis of individual selected positions, mutated clones can be sequenced to identify which amino acid residues have been introduced into the selected position in each clone. A small number of clones (e.g., about 24) can be selected for sequencing, which statistically should yield 10-15 unique antibodies, whereas larger numbers of clones (e.g., greater than 60) can be sequenced to ensure that antibodies with
15 every possible substitution at the selected position are identified.

In one embodiment, contact and/or hypermutation positions within the CDR3 regions of the heavy and/or light chains are first selected for mutagenesis. However, for antibodies that have already been affinity matured *in vitro* by random mutagenesis of the CDR3 regions via phage display selection, it may be preferably to first select contact
20 and/or hypermutation positions within CDR1 or CDR2 of the heavy and/or light chain.

In a more preferred embodiment, preferred selective mutagenesis positions within the CDR3 regions of the heavy and/or light chains are first selected for mutagenesis. However, for antibodies that have already been affinity matured *in vitro* by random mutagenesis of the CDR3 regions via phage display selection, it may be
25 preferably to first select preferred selective mutagenesis positions within CDR1 or CDR2 of the heavy and/or light chain.

In another preferred embodiment, the optimization of a selected antibody by the selective mutagenesis approach is done sequentially as follows: preferred selective mutagenesis positions selected from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94 are mutated first to at least 2 other amino acids each (preferably 5-14 other amino acids) and the resulting antibodies are characterized for increased affinity, neutralization potency (and possibly also for at least one other retained characteristic or property discussed elsewhere). If a

mutation of a single preferred selective mutagenesis position does not increase the affinity or neutralization potency at all or sufficiently and if even the combination of multiple activity enhancing amino acids replacing amino acids in preferred selective mutagenesis positions does not result in an combination antibody which meets the target

5 activity (including affinity and/or neutralization potency), additional amino acid residues will be selected for selective mutagenesis from the group consisting of H35, H50, H53, H54, H95, H96, H97, H98, L30A and L96 are mutated to at least 2 other amino acids each (preferably 5-14 other amino acids) and the resulting antibodies are characterized for increased affinity, neutralization potency (and possibly also for at least one other

10 retained characteristic or property discussed elsewhere).

If a mutation of a single amino acid residue selected from the group consisting of H35, H50, H53, H54, H95, H96, H97, H98, L30A and L96 does not increase the activity (including affinity and/or neutralization potency) at all or not sufficiently and if even the combination of multiple activity enhancing amino acids replacing amino acids in those

15 positions does not result in an combination antibody which meets the targeted activity (including affinity and/or target neutralization potency), additional amino acid residues will be selected for selective mutagenesis from the group consisting of H33B, H52B, L31A and are mutated to at least 2 other amino acids each (preferably 5-14 other amino acids) and the resulting antibodies are characterized for increased affinity, neutralization

20 potency (and possibly also for at least one other retained characteristic or property discussed elsewhere).

It should be understood that the sequential selective mutagenesis approach may end at any of the steps outline above as soon as an antibody with the desired activity (including affinity and neutralization potency) has been identified. If mutagenesis of the

25 preselected positions has identified activity enhancing amino acids residues but the combination antibody still do not meet the targets set for activity (including affinity and neutralization potency) and/or if the identified activity enhancing amino acids also affect other desired characteristics and are therefore not acceptable, the remaining CDR residues may be subjected to mutagenesis (see section IV).

30 The method of the invention can be used to improve activity of an antibody, or antigen binding portion thereof, to reach a predetermined target activity (e.g. a predetermined affinity and/or neutralization potency, and/or a desired property or characteristic).

Accordingly, the invention provides a method of improving the activity of an antibody, or antigen-binding portion thereof, to attain a predetermined target activity, comprising:

- a) providing a parent antibody a antigen-binding portion thereof;
- 5 b) selecting a preferred selective mutagenesis position selected from group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94.
- c) individually mutating the selected preferred selective mutagenesis position to at least two other amino acid residues to hereby create a first panel of mutated 10 antibodies, or antigen binding portions thereof;
- d) evaluating the activity of the first panel of mutated antibodies, or antigen binding portions thereof to determine if mutation of a single selective mutagenesis position produces an antibody or antigen binding portion thereof with the predetermined target activity or a partial target activity;
- 15 e) combining in a stepwise fashion, in the parent antibody, or antigen binding portion thereof, individual mutations shown to have an improved activity, to form combination antibodies, or antigen binding portions thereof.
- f) evaluating the activity of the combination antibodies, or antigen binding portions thereof to determine if the combination antibodies, or antigen binding portions 20 thereof have the predetermined target activity or a partial target activity.
- g) if steps d) or f) do not result in an antibody or antigen binding portion thereof having the predetermined target activity, or result an antibody with only a partial activity, additional amino acid residues selected from the group consisting of H35, H50, H53, H54, H95, H96, H97, H98, L30A and L96 are mutated to at least two other amino 25 acid residues to thereby create a second panel of mutated antibodies or antigen-binding portions thereof;
- h) evaluating the activity of the second panel of mutated antibodies or antigen binding portions thereof, to determine if mutation of a single amino acid residue selected from the group consisting of H35, H50, H53, H54, H95, H96, H97, H98, L30A and L96 results an antibody or antigen binding portion thereof, having the 30 predetermined target activity or a partial activity;

- i) combining in stepwise fashion in the parent antibody, or antigen-binding portion thereof, individual mutations of step g) shown to have an improved activity, to form combination antibodies, or antigen binding portions thereof;
 - j) evaluating the activity of the combination antibodies or antigen binding portions thereof, to determine if the combination antibodies, or antigen binding portions thereof have the predetermined target activity or a partial target activity;
 - k) if steps h) or j) do not result in an antibody or antigen binding portion thereof having the predetermined target activity, or result in an antibody with only a partial activity, additional amino acid residues selected from the group consisting of H33B,
- 10 H52B and L31A are mutated to at least two other amino acid residues to thereby create a third panel of mutated antibodies or antigen binding portions thereof;
- l) evaluating the activity of the third panel of mutated antibodies or antigen binding portions thereof, to determine if a mutation of a single amino acid residue selected from the group consisting of H33B, H52B and L31A resulted in an antibody or
 - 15 antigen binding portion thereof, having the predetermined target activity or a partial activity;
 - m) combining in a stepwise fashion in the parent antibody, or antigen binding portion thereof, individual mutation of step k) shown to have an improved activity, to form combination antibodies, or antigen binding portions, thereof;
- 20 n) evaluating the activity of the combination antibodies or antigen-binding portions thereof, to determine if the combination antibodies, or antigen binding portions thereof have the predetermined target activity to thereby produce an antibody or antigen binding portion thereof with a predetermined target activity.

A number of mutagenesis methods can be used, including PCR assembly, Kunkel (dut-ung-) and thiophosphate (Amersham Sculptor kit) oligonucleotide-directed mutagenesis.

A wide variety of host expression systems can be used to express the mutated antibodies, including bacterial, yeast, baculoviral and mammalian expression systems (as well as phage display expression systems). An example of a suitable bacterial expression vector is pUC119(Sfi). Other antibody expression systems are known in the art and/or are described below in section IV.

The modified antibodies, or antigen binding portions thereof, produced by the method of the invention can be identified without the reliance on phage display methods

for selection. Accordingly, the method of the invention is particularly advantageous for improving the activity of a recombinant parent antibody or antigen-binding portion thereof, that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in the phage-display system.

5 Accordingly, in another embodiment, the invention provides a method for improving the affinity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a recombinant parent antibody or antigen-binding portion thereof; that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;
- 10 b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation, thereby identifying a selected contact or hypermutation position;
- 15 c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof, and expressing said panel in a non-phage display system;
- 20 d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof;
- 25 e) optionally repeating steps b) through d) for at least one other preferred selective mutagenesis position, contact or hypermutation position;
- 30 f) combining, in the parent antibody, or antigen-binding portion thereof, individual mutations shown to have improved activity, to form combination antibodies, or antigen-binding portions thereof; and
- 35 g) evaluating the activity of the combination antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof; until an antibody, or antigen-binding portion thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferred contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96. Preferred hypermutation positions are selected from the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93. More preferred preferred selective mutagenesis positions are selected from the group consisting of H30, H31, H31B, H32,

H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93 and L94. Particularly preferred contact positions are selected from the group consisting of L50 and L94.

With available methods it is not possible or it is extremely laborious to derive an antibody with increased binding affinity and neutralization potency while retaining other properties or characteristics of the antibodies as discussed above. The method of this invention, however, can readily identify such antibodies. The antibodies subjected to the method of this invention can come from any source.

Therefore, in another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- 10 a) providing a recombinant parent antibody or antigen-binding portion thereof ;
 b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation, thereby identifying a selected preferred selective mutagenesis position, contact or hypermutation position;
- 15 c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof and expressing said panel in an appropriate expression system;
- 20 d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof, thereby identifying an activity enhancing amino acid residue;
- 25 e) evaluating the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof for at least one other property or characteristics, wherein the property or characteristic is one that needs to be retained in the antibody;
until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

In a preferred embodiment, the contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope

recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In another preferred embodiment, the hypermutation positions are selected from
5 the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody
10 with a close to germline immunoglobulin sequence.

In a more preferred embodiment the residues for selective mutagenesis are selected from the preferred selective mutagenesis positions from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with
15 other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In a more preferred embodiment, the contact positions are selected from the
20 group consisting of L50 and L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

25 If therefore, the affinity of an antibody for a specific antigen should be improved, but where the phage display (or related system including ribosome display) method is no longer applicable, and other desirable properties or characteristics should be retained, the method of the invention can be used. Accordingly, in another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:
30

a) providing a recombinant parent antibody or antigen-binding portion thereof; that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;

- b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation, thereby identifying a selected preferred selective mutagenesis position, contact or hypermutation position;
 - 5 c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof, and expressing said panel in a non-phage display system;
 - 10 d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;
 - 15 e) evaluating the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof for at least one other property or characteristic, wherein the property or characteristic is one that needs to be retained, until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.
 - 20 f) optionally, repeating steps a) through e) for at least one other preferred selective mutagenesis position, contact or hypermutation position;
 - 25 g) combining, in the parent antibody, or antigen-binding portion thereof, at least two individual activity enhancing amino acid residues shown to have improved activity and at least one retained property or characteristic, to form combination antibodies, or antigen-binding portions thereof; and
 - 30 h) evaluating the activity of the combination antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof; until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained other property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

In a preferred embodiment, the contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope

recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In another preferred embodiment, the hypermutation positions are selected from 5 the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody 10 with a close to germline immunoglobulin sequence.

In a more preferred embodiment the residues for selective mutagenesis are selected from the preferred selective mutagenesis positions from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with 15 other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In a more preferred embodiment, the contact positions are selected from the 20 group consisting of L50 and L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

25 In another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

a) providing a recombinant parent antibody or antigen-binding portion thereof; that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;

30 b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation, thereby identifying a selected contact or hypermutation position;

- c) individually mutating said selected preferred selective mutagenesis position, contact or hypermutation position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof, and expressing said panel in a non-phage display system;
- 5 d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;
- 10 e) evaluating the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof for at least one other property or characteristic, wherein the property or characteristic is one that needs to be retained, until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

In a preferred embodiment, the contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In another preferred embodiment, the hypermutation positions are selected from the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In a more preferred embodiment the residues for selective mutagenesis are selected from the preferred selective mutagenesis positions from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing

p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In a more preferred embodiment, the contact positions are selected from the 5 group consisting of L50 and L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

10 In another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

a) providing a recombinant parent antibody or antigen-binding portion thereof; that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;

15 b) selecting a preferred selective mutagenesis position, contact or hypermutation position within a complementarity determining region (CDR) for mutation, thereby identifying a selected contact or hypermutation position;

c) individually mutating said selected preferred selective mutagenesis positions, contact or hypermutation position to at least two other amino acid residues to thereby 20 create a panel of mutated antibodies, or antigen-binding portions thereof, and expressing said panel in a non-phage display system;

d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;

25 e) evaluating the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof for at least one other property or characteristic, wherein the property or characteristic is one that needs to be retained, until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained characteristic, relative to the parent antibody, or 30 antigen-binding portion thereof, is obtained.

f) optionally, repeating steps a) through e) for at least one other preferred selective mutagenesis position, contact or hypermutation position;

g) combining, in the parent antibody, or antigen-binding portion thereof, at least two individual activity enhancing amino acid residues shown to have improved activity and at least one retained other characteristic, to form combination antibodies, or antigen-binding portions thereof; and

5 h) evaluating the activity of the combination antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof; until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

10 In a preferred embodiment, the contact positions are selected from the group consisting of H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope

15 recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

20 In another preferred embodiment, the hypermutation positions are selected from the group consisting of H30, H31, H31B, H32, H52, H56, H58, L30, L31, L32, L53 and L93 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

25 In a more preferred embodiment the residues for selective mutagenesis are selected from the preferred selective mutagenesis positions from the group consisting of H30, H31, H31B, H32, H33, H52, H56, H58, L30, L31, L32, L50, L91, L92, L93, L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

In a more preferred embodiment, the contact positions are selected from the group consisting of L50 and L94 and the other characteristic is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 5 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

IV. Modifications of other CDR residues

Ultimately, all CDR residues in a given antibody-antigen pair identified by any 10 means to be required as activity enhancing amino acid residues and/or required directly or indirectly for binding to the antigen and/or for retaining other desirable properties or characteristics of the antibody. Such CDR residues are referred to as “preferred selective mutagenesis positions”. It should be noted that in specific circumstances that 15 preferred selective mutagenesis residues can be identified also by other means including co-crystallization of antibody and antigen and molecular modeling.

If the preferred attempts to identify activity enhancing amino acids focussing on the preferred selective mutagenesis positions, contact or hypermutation positions described above are exhausted, or if additional improvements are required, the remaining CDR residues may be modified as described below. It should be understood 20 that the antibody could already be modified in any one or more contact or hypermutation positions according to the embodiments discussed above but may require further improvements. Therefore, in another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- 25 a) providing a parent antibody or antigen-binding portion thereof;
- b) selecting an amino acid residue within a complementarity determining region (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- c) individually mutating said selected position e.g., to at least two other amino 30 acid residues to thereby create a mutated antibody or a panel of mutated antibodies, or antigen-binding portions thereof;
- d) evaluating the activity of the mutated antibody or the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or

antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;

- e) evaluating the mutated antibody or the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof, for changes in at least one other property or characteristic until an antibody, or antigen-binding portion thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

5

Preferably, the other characteristic or property is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope 10 recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence

If mutagenesis of a single residue is not sufficient other residues can be included; therefore, in another embodiment, the invention provides a method for improving the 15 activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a parent antibody or antigen-binding portion thereof;
- b) selecting an amino acid residue within a complementarity determining region (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, 20 L55, L91, L92, L93, L94 and L96;
- c) individually mutating said selected position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof;
- d) evaluating the activity of the panel of mutated antibodies, or antigen-binding 25 portions thereof, relative to the parent antibody or antigen-binding portion thereof, thereby identifying an activity enhancing amino acid residue;
- e) repeating steps b) through d) for at least one other CDR position which is neither the position selected under b) nor a position at H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, 30 L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- f) combining, in the parent antibody, or antigen-binding portion thereof, at least two individual activity enhancing amino acid residues shown to have improved activity, to form combination antibodies, or antigen-binding portions thereof; and

g) evaluating the activity of the combination antibodies, or antigen-binding portions thereof with two activity enhancing amino acid residues, relative to the parent antibody or antigen-binding portion thereof until an antibody, or antigen-binding portion thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

If the preferred attempts to identify activity enhancing amino acids focussing on the contact or hypermutation positions described above are exhausted, or if additional improvements are required, and the antibody in question can not further be optimized by mutagenesis and phage display (or related ribosome display) methods the remaining 10 CDR residues may be modified as described below. It should be understood that the antibody could already be modified in any one or more preferred selective mutagenesis position, contact or hypermutation positions according to the embodiments discussed above but may require further improvements.

Therefore, in another embodiment, the invention provides a method for 15 improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a recombinant parent antibody or antigen-binding portion thereof; that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;
- b) selecting a selecting an amino acid residue within a complementarity 20 determining region (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and;
- c) individually mutating said selected contact or hypermutation position to at 25 least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof, and expressing said panel in a non-phage display system;
- d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;
- e) evaluating the panel of mutated antibodies, or antigen-binding portions 30 thereof, relative to the parent antibody or antigen-binding portion thereof, for changes in at least one other property or characteristic, until an antibody, or antigen-binding portion

thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferably, the other characteristic or property is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope 5 recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence.

If a single mutagenesis is not sufficient to increase the affinity of the antibody other residues may be included in the mutagenesis. Therefore, in another embodiment, 10 the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a parent antibody or antigen-binding portion thereof that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;
- 15 b) selecting an amino acid residue within a complementarity determining region (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- 20 c) individually mutating said selected position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof and expression in a non-phage display system;
- d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;
- 25 e) repeating steps b) through d) for at least one other position which is neither the position selected under b) nor a position at H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 ;
- 30 g) combining, in the parent antibody, or antigen-binding portion thereof, at least two individual activity enhancing amino acid residues shown to have improved activity, to form combination antibodies, or antigen-binding portions thereof; and

h) evaluating the activity and other property or characteristic of the combination antibodies, or antigen-binding portions thereof with two activity enhancing amino acid residues, relative to the parent antibody or antigen-binding portion thereof; until an antibody, or antigen-binding portion thereof, with an improved activity, relative 5 to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferably, the other characteristic or property is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope 10 recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence

The preferred attempts to identify activity enhancing amino acids focussing on the preferred selective mutagenesis positions, contact or hypermutation positions described may be exhausted, or additional improvements may be required, and it is important to retain other properties or characteristics of the antibody.

15 Therefore, in another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, without affecting other characteristics, comprising:

- a) providing a parent antibody or antigen-binding portion thereof;
- b) selecting an amino acid residue within a complementarity determining region 20 (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- c) individually mutating said selected position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions 25 thereof;
- d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;
- e) evaluating the panel of mutated antibodies, or antigen-binding portions 30 thereof, relative to the parent antibody or antigen-binding portion thereof, for changes in at least one other property or characteristic until an antibody, or antigen-binding portion thereof, with an improved activity and retained other property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferably, the other characteristic or property is selected from 1) preservation of non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce 5 an antibody with a close to germline immunoglobulin sequence

If mutagenesis of a single residue is not sufficient other residues can be included; therefore, in another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a parent antibody or antigen-binding portion thereof;
- 10 b) selecting an amino acid residue within a complementarity determining region (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- 15 c) individually mutating said selected position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof;
- d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof, thereby identifying an activity enhancing amino acid residue;
- 20 e.) evaluating the panel of mutated antibodies or antigen-binding portions thereof, relative to the parent antibody or antigen-portion thereof, for changes in at least one other characteristic or property;
- e) repeating steps b) through e) for at least one other CDR position which is neither the position selected under b) nor a position at H30, H31, H31B, H32, H33, H35, 25 H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- f) combining, in the parent antibody, or antigen-binding portion thereof, at least two individual activity enhancing amino acid residues shown to have improved activity and not affecting at least one other property or characteristic, to form combination 30 antibodies, or antigen-binding portions thereof; and
- g) evaluating the activity and the retention of at least one other property or characteristic of the combination antibodies, or antigen-binding portions thereof with two activity enhancing amino acid residues, relative to the parent antibody or antigen-

binding portion thereof until an antibody, or antigen-binding portion thereof, with an improved activity and at least one retained other property or characteristic, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

Mutagenesis of the preferred selective mutagenesis position, contact and

5 hypermutation residues may not have increased the affinity of the antibody sufficiently, and mutagenesis and the phage display method (or related ribosome display method) may no longer be useful and at least one other characteristic or property of the antibody should be retained.

Therefore, in another embodiment the invention provides a method to improve
10 the affinity of an antibody or antigen-binding portion thereof, comprising:

a) providing a parent antibody or antigen-binding portion thereof that was obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;

b) selecting an amino acid residue within a complementarity determining region
15 (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;

c) individually mutating said selected position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions
20 thereof and expression in a non-phage display system;

d) evaluating the activity of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof thereby identifying an activity enhancing amino acid residue;

e) evaluating the panel of mutated antibodies, or antigen-binding portions
25 thereof, relative to the parent antibody or antigen-binding portion thereof, for changes in at least one other property or characteristic until an antibody, or antigen-binding portion thereof, with an improved activity, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

Preferably, the other characteristic or property is selected from 1) preservation of
30 non-crossreactivity with other proteins or human tissues, 2) preservation of epitope recognition, i.e. recognizing p40 epitope preferably in the context of the p70 p40/p35 heterodimer preventing binding interference from free, soluble p40 and/or 3) to produce an antibody with a close to germline immunoglobulin sequence

If mutagenesis of a single residue is not sufficient other residues can be included; therefore, in another embodiment, the invention provides a method for improving the activity of an antibody, or antigen-binding portion thereof, comprising:

- a) providing a parent antibody or antigen-binding portion thereof that was
- 5 obtained by selection in a phage-display system but whose activity cannot be further improved by mutagenesis in said phage-display system;
- b) selecting an amino acid residue within a complementarity determining region (CDR) for mutation other than H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53,
- 10 L55, L91, L92, L93, L94 and L96;
- c) individually mutating said selected position to at least two other amino acid residues to thereby create a panel of mutated antibodies, or antigen-binding portions thereof and expression in a non-phage display system;
- d) evaluating the activity and retention of at least one other property or
- 15 characteristic of the panel of mutated antibodies, or antigen-binding portions thereof, relative to the parent antibody or antigen-binding portion thereof, thereby identifying an activity enhancing amino acid residue;
- e) repeating steps b) through d) for at least one other CDR position which is neither the position selected under b) nor a position at H30, H31, H31B, H32, H33, H35, H50, H52, H52A, H53, H54, H56, H58, H95, H96, H97, H98, H101, L30, L31, L32, L34, L50, L52, L53, L55, L91, L92, L93, L94 and L96;
- f) combining, in the parent antibody, or antigen-binding portion thereof, at least two individual activity enhancing amino acid residues shown to have improved activity and not to affect at least one other property or characteristic, to form combination
- 25 antibodies, or antigen-binding portions thereof; and
- g) evaluating the activity and retention of at least one property or characteristic of the combination antibodies, or antigen-binding portions thereof with two activity enhancing amino acid residues, relative to the parent antibody or antigen-binding portion thereof until an antibody, or antigen-binding portion thereof, with an improved
- 30 activity and at least one other retained characteristic or property, relative to the parent antibody, or antigen-binding portion thereof, is obtained.

V. Expression of Antibodies

An antibody, or antibody portion, of the invention can be prepared by recombinant expression of immunoglobulin light and heavy chain genes in a host cell. To express an antibody recombinantly, a host cell is transfected with one or more recombinant expression vectors carrying DNA fragments encoding the immunoglobulin light and heavy chains of the antibody such that the light and heavy chains are expressed in the host cell and, preferably, secreted into the medium in which the host cells are cultured, from which medium the antibodies can be recovered. Standard recombinant DNA methodologies are used to obtain antibody heavy and light chain genes, incorporate these genes into recombinant expression vectors and introduce the vectors into host cells, such as those described in Sambrook, Fritsch and Maniatis (eds), *Molecular Cloning; A Laboratory Manual, Second Edition*, Cold Spring Harbor, N.Y., (1989), Ausubel, F.M. et al. (eds.) *Current Protocols in Molecular Biology*, Greene Publishing Associates, (1989) and in U.S. Patent No. 4,816,397 by Boss et al.

To obtain a DNA fragment encoding the heavy chain variable region of Joe 9 wt or a Joe 9 wt-related antibody, antibodies specific for human IL-12 were screened from human libraries and mutated, as described in section II. Once DNA fragments encoding Joe 9 wt or Joe 9 wt-related VH and VL segments are obtained, mutagenesis of these sequences is carried out by standard methods, such as PCR site directed mutagenesis (PCR-mediated mutagenesis in which the mutated nucleotides are incorporated into the PCR primers such that the PCR product contains the mutations) or other site-directed mutagenesis methods. Human IL-12 antibodies that displayed a level of activity and binding specificity/affinity that was desirable, for example J695, were further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term "operatively linked", as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (CH1, CH2 and CH3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat, E.A.,

et al. (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or 5 IgD constant region and any allotypic variant therein as described in Kabat (, Kabat, E.A., et al. (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242), but most preferably is an IgG1 or IgG4 constant region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding 10 only the heavy chain CH1 constant region.

The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see e.g., 15 Kabat, E.A., et al. (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region, but most preferably is a lambda constant region.

20 To create a scFv gene, the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly₄-Ser)₃, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see e.g., Bird et al. (1988) *Science* 242:423-426; Huston et al. (1988) *Proc. Natl. 25 Acad. Sci. USA* 85:5879-5883; McCafferty et al., *Nature* (1990) 348:552-554).

To express the antibodies, or antibody portions of the invention, DNAs encoding partial or full-length light and heavy chains, obtained as described above, are inserted 30 into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. In this context, the term "operatively linked" is intended to mean that an antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host

cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector or, more typically, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the antibody gene 5 fragment and vector, or blunt end ligation if no restriction sites are present). Prior to insertion of the J695 or J695-related light or heavy chain sequences, the expression vector may already carry antibody constant region sequences. For example, one approach to converting the J695 or J695-related VH and VL sequences to full-length antibody genes is to insert them into expression vectors already encoding heavy chain 10 constant and light chain constant regions, respectively, such that the VH segment is operatively linked to the CH segment(s) within the vector and the VL segment is operatively linked to the CL segment within the vector. Additionally or alternatively, the recombinant expression vector can encode a signal peptide that facilitates secretion 15 of the antibody chain from a host cell. The antibody chain gene can be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

In addition to the antibody chain genes, the recombinant expression vectors of the invention carry regulatory sequences that control the expression of the antibody 20 chain genes in a host cell. The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). It 25 will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as 30 promoters and/or enhancers derived from cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)) and polyoma. For further description of viral regulatory elements, and sequences thereof, see e.g., U.S.

Patent No. 5,168,062 by Stinski, U.S. Patent No. 4,510,245 by Bell *et al.* and U.S. Patent No. 4,968,615 by Schaffner *et al.*, U.S. Patent No. 5,464,758 by Bujard *et al.* and U.S. Patent No. 5,654,168 by Bujard *et al.*

In addition to the antibody chain genes and regulatory sequences, the 5 recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see e.g., U.S. Patents Nos. 4,399,216, 10 4,634,665 and 5,179,017, all by Axel *et al.*). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr⁻ host cells with methotrexate selection/amplification) and the *neo* gene (for G418 selection).

For expression of the light and heavy chains, the expression vector(s) encoding 15 the heavy and light chains is transfected into a host cell by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, e.g., electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like. Although it is theoretically possible to express the 20 antibodies of the invention in either prokaryotic or eukaryotic host cells, expression of antibodies in eukaryotic cells, and most preferably mammalian host cells, is the most preferred because such eukaryotic cells, and in particular mammalian cells, are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active antibody. Preferred mammalian host cells for expressing the 25 recombinant antibodies of the invention include Chinese Hamster Ovary (CHO cells) (including dhfr⁻ CHO cells, described in Urlaub and Chasin, (1980) *Proc. Natl. Acad. Sci. USA* 77:4216-4220, used with a DHFR selectable marker, e.g., as described in R.J. Kaufman and P.A. Sharp (1982) *Mol. Biol.* 159:601-621), NS0 myeloma cells, COS cells and SP2 cells. When recombinant expression vectors encoding antibody genes are 30 introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the

host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods.

Host cells can also be used to produce portions of intact antibodies, such as Fab fragments or scFv molecules. It will be understood that variations on the above 5 procedure are within the scope of the present invention. For example, it may be desirable to transfet a host cell with DNA encoding either the light chain or the heavy chain (but not both) of an antibody of this invention. Recombinant DNA technology may also be used to remove some or all of the DNA encoding either or both of the light and heavy chains that is not necessary for binding to hIL-12. The molecules expressed 10 from such truncated DNA molecules are also encompassed by the antibodies of the invention. In addition, bifunctional antibodies may be produced in which one heavy and one light chain are an antibody of the invention and the other heavy and light chain are specific for an antigen other than hIL-12 by crosslinking an antibody of the invention to a second antibody by standard chemical crosslinking methods.

15 In a preferred system for recombinant expression of an antibody, or antigen-binding portion thereof, of the invention, a recombinant expression vector encoding both the antibody heavy chain and the antibody light chain is introduced into dhfr- CHO cells by calcium phosphate-mediated transfection. Within the recombinant expression vector, the antibody heavy and light chain genes are each operatively linked to 20 enhancer/promoter regulatory elements (e.g., derived from SV40, CMV, adenovirus and the like, such as a CMV enhancer/AdMLP promoter regulatory element or an SV40 enhancer/AdMLP promoter regulatory element) to drive high levels of transcription of the genes. The recombinant expression vector also carries a DHFR gene, which allows for selection of CHO cells that have been transfected with the vector using methotrexate 25 selection/amplification. The selected transformant host cells are culture to allow for expression of the antibody heavy and light chains and intact antibody is recovered from the culture medium. Standard molecular biology techniques are used to prepare the recombinant expression vector, transfet the host cells, select for transformants, culture the host cells and recover the antibody from the culture medium. Antibodies or antigen- 30 binding portions thereof of the invention can be expressed in an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see e.g., Taylor, L.D. *et al.* (1992) Nucl. Acids Res. 20: 6287-6295). Plant cells can also be modified to create transgenic plants that express the antibody or antigen binding portion thereof, of the invention.

In view of the foregoing, another aspect of the invention pertains to nucleic acid, vector and host cell compositions that can be used for recombinant expression of the antibodies and antibody portions of the invention. Preferably, the invention features isolated nucleic acids that encode CDRs of J695, or the full heavy and/or light chain 5 variable region of J695. Accordingly, in one embodiment, the invention features an isolated nucleic acid encoding an antibody heavy chain variable region that encodes the J695 heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 25. Preferably, the nucleic acid encoding the antibody heavy chain variable region further encodes a J695 heavy chain CDR2 which comprises the amino acid sequence of SEQ 10 ID NO: 27. More preferably, the nucleic acid encoding the antibody heavy chain variable region further encodes a J695 heavy chain CDR1 which comprises the amino acid sequence of SEQ ID NO: 29. Even more preferably, the isolated nucleic acid encodes an antibody heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 31 (the full VH region of J695).

15 In other embodiments, the invention features an isolated nucleic acid encoding an antibody light chain variable region that encodes the J695 light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 26. Preferably, the nucleic acid encoding the antibody light chain variable region further encodes a J695 light chain CDR2 which comprises the amino acid sequence of SEQ ID NO: 28. More preferably, 20 the nucleic acid encoding the antibody light chain variable region further encodes a J695 light chain CDR1 which comprises the amino acid sequence of SEQ ID NO: 30. Even more preferably, the isolated nucleic acid encodes an antibody light chain variable region comprising the amino acid sequence of SEQ ID NO: 32 (the full VL region of J695).

25 The invention also provides recombinant expression vectors encoding both an antibody heavy chain and an antibody light chain. For example, in one embodiment, the invention provides a recombinant expression vector encoding:

- a) an antibody heavy chain having a variable region comprising the amino acid sequence of SEQ ID NO: 31; and
- 30 b) an antibody light chain having a variable region comprising the amino acid sequence of SEQ ID NO: 32.

The invention also provides host cells into which one or more of the recombinant expression vectors of the invention have been introduced. Preferably, the host cell is a

mammalian host cell, more preferably the host cell is a CHO cell, an NS0 cell or a COS cell. Still further the invention provides a method of synthesizing a recombinant human antibody of the invention by culturing a host cell of the invention in a suitable culture medium until a recombinant human antibody of the invention is synthesized. The 5 method can further comprise isolating the recombinant human antibody from the culture medium.

VI. Pharmaceutical Compositions and Pharmaceutical Administration

The antibodies and antibody-portions of the invention can be incorporated into 10 pharmaceutical compositions suitable for administration to a subject. Typically, the pharmaceutical composition comprises an antibody or antibody portion of the invention and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are 15 physiologically compatible. Examples of pharmaceutically acceptable carriers include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Pharmaceutically acceptable carriers may further comprise 20 minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody or antibody portion.

The antibodies and antibody-portions of the invention can be incorporated into a pharmaceutical composition suitable for parenteral administration. Preferably, the 25 antibody or antibody-portions will be prepared as an injectable solution containing 0.1-250 mg/ml antibody. The injectable solution can be composed of either a liquid or lyophilized dosage form in a flint or amber vial, ampule or pre-filled syringe. The buffer can be L-histidine (1-50 mM), optimally 5-10mM, at pH 5.0 to 7.0 (optimally pH 6.0). Other suitable buffers include but are not limited to, sodium succinate, sodium 30 citrate, sodium phosphate or potassium phosphate. Sodium chloride can be used to modify the toxicity of the solution at a concentration of 0-300 mM (optimally 150 mM for a liquid dosage form). Cryoprotectants can be included for a lyophilized dosage form, principally 0-10% sucrose (optimally 0.5-1.0%). Other suitable cryoprotectants

include trehalose and lactose. Bulking agents can be included for a lyophilized dosage form, principally 1-10% mannitol (optimally 2-4%). Stabilizers can be used in both liquid and lyophilized dosage forms, principally 1-50 mM L-Methionine (optimally 5-10 mM). Other suitable bulking agents include glycine, arginine, can be included as 0-5 0.05% polysorbate-80 (optimally 0.005-0.01%). Additional surfactants include but are not limited to polysorbate 20 and BRIJ surfactants.

In a preferred embodiment, the pharmaceutical composition includes the antibody at a dosage of about 100 mg - 200 mg dose.

The compositions of this invention may be in a variety of forms. These include, 10 for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for 15 passive immunization of humans with other antibodies. The preferred mode of administration is parenteral (e.g., intravenous, subcutaneous, intraperitoneal, intramuscular). In a preferred embodiment, the antibody is administered by subcutaneous injection.

Therapeutic compositions typically must be sterile and stable under the conditions 20 of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the active compound (*i.e.*, antibody or antibody portion) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed 25 by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile, lyophilized powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and spray-drying that yields a powder of the active 30 ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions

can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

The antibodies and antibody-portions of the present invention can be administered by a variety of methods known in the art, although for many therapeutic applications, the 5 preferred route/mode of administration is subcutaneous injection, intravenous injection or infusion. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal 10 patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, e.g., *Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel 15 Dekker, Inc., New York, 1978.

In certain embodiments, an antibody or antibody portion of the invention may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's 20 diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the invention by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its 25 inactivation.

Supplementary active compounds can also be incorporated into the compositions. In certain embodiments, an antibody or antibody portion of the invention is coformulated with and/or coadministered with one or more additional therapeutic agents that are useful for treating disorders in which IL-12 activity is detrimental. For 30 example, an anti-hIL-12 antibody or antibody portion of the invention may be coformulated and/or coadministered with one or more additional antibodies that bind other targets (e.g., antibodies that bind other cytokines or that bind cell surface molecules). Furthermore, one or more antibodies of the invention may be used in

combination with two or more of the foregoing therapeutic agents. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies. It will be appreciated by the skilled practitioner that when the

5 antibodies of the invention are used as part of a combination therapy, a lower dosage of antibody may be desirable than when the antibody alone is administered to a subject (e.g., a synergistic therapeutic effect may be achieved through the use of combination therapy which, in turn, permits use of a lower dose of the antibody to achieve the desired therapeutic effect).

10 Interleukin 12 plays a critical role in the pathology associated with a variety of diseases involving immune and inflammatory elements. These diseases include, but are not limited to, rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin

15 dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, atopic dermatitis, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis,

20 Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure,

25 myocardial infarction, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, spondyloarthropathy,

30 atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic

mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female

5 infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus

10 associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial

15 lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated hypoglycemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation,

20 osteoarthritis, primary sclerosing cholangitis, idiopathic leucopenia, autoimmune neutropenia, renal disease NOS, glomerulonephritides, microscopic vasulitis of the kidneys, lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), insulin-dependent diabetes mellitus, sympathetic ophthalmia, pulmonary hypertension secondary to connective

25 tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Takayasu's disease/arteritis, autoimmune thrombocytopenia, idiopathic thrombocytopenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary

30 myxoedema, phacogenic uveitis, primary vasculitis and vitiligo. The human antibodies, and antibody portions of the invention can be used to treat autoimmune diseases, in particular those associated with inflammation, including, rheumatoid spondylitis, allergy, autoimmune diabetes, autoimmune uveitis.

Preferably, the antibodies of the invention or antigen-binding portions thereof, are used to treat rheumatoid arthritis, Crohn's disease, multiple sclerosis, insulin dependent diabetes mellitus and psoriasis, as described in more detail in section VII.

A human antibody, or antibody portion, of the invention also can be

- 5 administered with one or more additional therapeutic agents useful in the treatment of autoimmune and inflammatory diseases.

Antibodies of the invention, or antigen binding portions thereof can be used alone or in combination to treat such diseases. It should be understood that the IL-12 antibodies of the invention or antigen binding portion thereof can be used alone or in

- 10 combination with an additional agent, e.g., a therapeutic agent, said additional agent being selected by the skilled artisan for its intended purpose. For example, the additional agent can be a therapeutic agent art-recognized as being useful to treat the disease or condition being treated by the antibody of the present invention. The additional agent also can be an agent which imparts a beneficial attribute to the
- 15 therapeutic composition e.g., an agent which effects the viscosity of the composition.

It should further be understood that the combinations which are to be included within this invention are those combinations useful for their intended purpose. The agents set forth below are illustrative for purposes and not intended to be limited. The combinations which are part of this invention can be the antibodies of the present

- 20 invention and at least one additional agent selected from the lists below. The combination can also include more than one additional agent, e.g., two or three additional agents if the combination is such that the formed composition can perform its intended function. Furthermore, additional agents described herein used in combination with an IL-12 antibody, are not limited to the disorder to which they are attributed for
- 25 treatment.

Preferred combinations are non-steroidal anti-inflammatory drug(s) also referred to as NSAIDS which include drugs like ibuprofen. Other preferred combinations are corticosteroids including prednisolone; the well known side-effects of steroid use can be reduced or even eliminated by tapering the steroid dose required when treating patients

- 30 in combination with the anti-IL-12 antibodies of this invention. Non-limiting examples of therapeutic agents for rheumatoid arthritis with which an antibody, or antibody portion, of the invention can be combined include the following: cytokine suppressive anti-inflammatory drug(s) (CSAIDs); antibodies to or antagonists of other human

cytokines or growth factors, for example, TNF (including adalimumab / HUMIRA), LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, 5 CD30, CD40, CD45, CD69, CD80 (B7.1), CD86 (B7.2), CD90, or their ligands including CD154 (gp39 or CD40L).

Preferred combinations of therapeutic agents may interfere at different points in the autoimmune and subsequent inflammatory cascade; preferred examples include TNF antagonists like chimeric, humanized or human TNF antibodies, D2E7, (U.S. application 10 serial number 08/599,226 filed February 9, 1996), cA2 (RemicadeTM), CDP 571, anti-TNF antibody fragments (*e.g.*, CDP870), and soluble p55 or p75 TNF receptors, derivatives thereof, (p75TNFR1gG (EnbrelTM) or p55TNFR1gG (Lenercept), soluble 15 IL-13 receptor (sIL-13), and also TNF α converting enzyme (TACE) inhibitors; similarly IL-1 inhibitors (*e.g.*, Interleukin-1-converting enzyme inhibitors, such as Vx740, or IL-1RA etc.) may be effective for the same reason. Other preferred combinations include 20 Interleukin 11, anti-P7s and p-selectin glycoprotein ligand (PSGL). Yet another preferred combination are other key players of the autoimmune response which may act parallel to, dependent on or in concert with IL-12 function; especially preferred are IL-18 antagonists including IL-18 antibodies or soluble IL-18 receptors, or IL-18 binding 25 proteins. It has been shown that IL-12 and IL-18 have overlapping but distinct functions and a combination of antagonists to both may be most effective. Yet another preferred combination are non-depleting anti-CD4 inhibitors. Yet other preferred combinations include antagonists of the co-stimulatory pathway CD80 (B7.1) or CD86 (B7.2) including antibodies, soluble receptors or antagonistic ligands.

25 Anti-IL12 antibodies, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, 6-MP, azathioprine sulphasalazine, mesalazine, olsalazine chloroquine/hydroxychloroquine, pencillamine, aurothiomalate (intramuscular and oral), azathioprine, cochicine, corticosteroids (oral, inhaled and local injection), beta-2 adrenoreceptor agonists (salbutamol, terbutaline, salmeterol), 30 xanthines (theophylline, aminophylline), cromoglycate, nedocromil, ketotifen, ipratropium and oxitropium, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement

inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (e.g. IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors (e.g., Vx740), anti-P7s, p-selectin glycoprotein ligand (PSGL), TNF α converting enzyme (TACE) inhibitors, T-cell signalling inhibitors 5 such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g. soluble p55 or p75 TNF receptors and the derivatives p75TNFR IgG (EnbrelTM) and p55TNFR IgG (Lenercept), sIL-1RI, sIL-1RII, sIL-6R, soluble IL-13 receptor (sIL-13)) and antiinflammatory cytokines (e.g. IL-4, IL-10, IL-10, IL-11, IL-13 and TGF β). Preferred combinations include methotrexate or leflunomide and in moderate or severe rheumatoid arthritis cases, cyclosporine.

Non-limiting examples of therapeutic agents for inflammatory bowel disease with which an anti-IL-12 antibody, or antibody portion, can be combined include the following: budesonide; epidermal growth factor; corticosteroids; cyclosporin, 15 sulfasalazine; aminosalicylates; 6-mercaptopurine; azathioprine; metronidazole; lipoxygenase inhibitors; mesalamine; olsalazine; balsalazide; antioxidants; thromboxane inhibitors; IL-1 receptor antagonists; anti-IL-1 β monoclonal antibodies; anti-IL-6 monoclonal antibodies; growth factors; elastase inhibitors; pyridinyl-imidazole compounds; antibodies to or antagonists of other human cytokines or growth factors, 20 for example, TNF (including adalimumab / HUMIRA), LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands. The antibodies of the invention, or antigen binding portions 25 thereof, may also be combined with agents, such as methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (e.g. IRAK, NIK, 30 IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors (e.g., Vx740), anti-P7s, p-selectin glycoprotein ligand (PSGL), TNF α converting enzyme inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme

inhibitors, soluble cytokine receptors and derivatives thereof (e.g. soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R, soluble IL-13 receptor (sIL-13)) and antiinflammatory cytokines (e.g. IL-4, IL-10, IL-11, IL-13 and TGF β).

Preferred examples of therapeutic agents for Crohn's disease in which an antibody or an antigen binding portion can be combined include the following: TNF antagonists, for example, anti-TNF antibodies, D2E7 (adalimumab / HUMIRA), cA2 (RemicadeTM), CDP 571, anti-TNF antibody fragments (e.g., CDP870), TNFR-Ig constructs(p75TNFR IgG (EnbrelTM) and p55TNFR IgG (Lenercept)) , anti-P7s, p-selectin glycoprotein ligand (PSGL), soluble IL-13 receptor (sIL-13), and PDE4 inhibitors.

Antibodies of the invention or antigen binding portions thereof, can be combined with corticosteroids, for example, budesonide and dexamethasone. Antibodies may also be combined with agents such as sulfasalazine, 5-aminosalicylic acid and olsalazine, and agents which interfere with synthesis or action of proinflammatory cytokines such as IL-1, for example, IL-1 β converting enzyme inhibitors (e.g., Vx740) and IL-1ra.

Antibodies or antigen binding portion thereof may also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors 6-mercaptopurines. Antibodies or antigen binding portions thereof, can be combined with IL-11.

Non-limiting examples of therapeutic agents for multiple sclerosis with which an antibody, or antibody portion, can be combined include the following: corticosteroids; prednisolone; methylprednisolone; azathioprine; cyclophosphamide; cyclosporine; methotrexate; 4-aminopyridine; tizanidine; interferon- β 1a (Avonex; Biogen); interferon- β 1b (Betaseron; Chiron/Berlex); Copolymer 1 (Cop-1; Copaxone; Teva Pharmaceutical Industries, Inc.); hyperbaric oxygen; intravenous immunoglobulin; clabribine; antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD80, CD86, CD90 or their ligands. The antibodies of the invention, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, cyclosporine, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by

proinflammatory cytokines such as TNF α or IL-1 (e.g. IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors (e.g., Vx740), anti-P7s, p-selectin glycoprotein ligand (PSGL), TACE inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, 5 angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g. soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R, soluble IL-13 receptor (sIL-13)) and antiinflammatory cytokines (e.g. IL-4, IL-10, IL-13 and TGF β).

Preferred examples of therapeutic agents for multiple sclerosis in which the antibody or antigen binding portion thereof can be combined to include interferon- β , for 10 example, IFN β 1a and IFN β 1b; copaxone, corticosteroids, IL-1 inhibitors, TNF inhibitors, and antibodies to CD40 ligand and CD80.

An antibody, antibody portion, may be used in combination with other agents to treat skin conditions. For example, an antibody, antibody portion, or other IL-12 inhibitor of the invention is combined with PUVA therapy. PUVA is a combination of 15 psoralen (P) and long-wave ultraviolet radiation (UVA) that is used to treat many different skin conditions. The antibodies, antibody portions, or other IL-12 inhibitors of the invention can also be combined with pimecrolimus. In another embodiment, the antibodies of the invention are used to treat psoriasis, wherein the antibodies are administered in combination with tacrolimus. In a further embodiment, tacrolimus and 20 IL-12 inhibitors are administered in combination with methotrexate and/or cyclosporine. In still another embodiment, the IL-12 inhibitor of the invention is administered with excimer laser treatment for treating psoriasis.

The pharmaceutical compositions of the invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of an antibody or antibody portion of the invention. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the antibody or antibody portion may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the 25 individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired 30 amount.

prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

Dosage regimens may be adjusted to provide the optimum desired response (*e.g.*, 5 a therapeutic or prophylactic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used 10 herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active 15 compound and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

In one embodiment, the IL-12 antibody, or antigen-binding portion thereof, is administered on a biweekly dosing regimen, including, for example, a biweekly dosage 20 ranging from about 50 to 300 mg, a dosage ranging from about 100 mg to about 200 mg, and a dosage from about 125 to about 175 mg. Alternatively, the IL-12 antibody may be administered as a one time dose, including, for example, a dose of about 200 mg dose, a dose of about 100 mg. In another embodiment, the IL-12 antibody is administered on a weekly dosing regimen, including, for example, a dose ranging from about 50 to 300 25 mg, a dosage ranging from about 100 mg to about 200 mg, and a dosage from about 125 to about 175 mg. It should be noted that doses within the specified ranges are also included herein, *e.g.*, 85 mg, 97 mg, etc.

In another embodiment, a human IL-12 antibody, or antigen-binding portion thereof, is administered as a single dose to a subject having a disorder in which IL-12 30 activity is detrimental, *e.g.*, psoriasis, which results in treatment. A response to the IL-12 antibody, or antigen-binding portion thereof, may be maintained for an extended period in a subject. Maintenance of a response may be monitored in accordance with the disorder being treated. For example, maintenance of a response with an IL-12

antibody, or antigen-binding portion thereof, for treating psoriasis may be determined by the subject's PASI 75 response over time.

It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, 5 specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

10

VII. Uses of the Invention

The invention provides a method for inhibiting IL-12 activity in a subject suffering from a disorder in which IL-12 activity is detrimental. In one embodiment, the invention provides a method treating psoriasis comprising administering a single 15 dose of an IL-12 antibody, or antigen-binding portion thereof.

IL-12 has been implicated in the pathophysiology of a wide variety of disorders (Windhagen *et al.*, (1995) *J. Exp. Med.* 182: 1985-1996; Morita *et al.* (1998) *Arthritis and Rheumatism*. 41: 306-314; Bucht *et al.*, (1996) *Clin. Exp. Immunol.* 103: 347-367; Fais *et al.* (1994) *J. Interferon Res.* 14:235-238; Parronchi *et al.*, (1997) *Am. J. Path.* 20 150:823-832; Monteleone *et al.*, (1997) *Gastroenterology*. 112:1169-1178, and Berrebi *et al.*, (1998) *Am. J. Path* 152:667-672; Parronchi *et al* (1997) *Am. J. Path.* 150:823-832). The invention provides methods for inhibiting IL-12 activity in a subject suffering from such a disorder, which method comprises administering to the subject an antibody or antibody portion of the invention such that IL-12 activity in the subject is inhibited. 25 Preferably, the IL-12 is human IL-12 and the subject is a human subject. Alternatively, the subject can be a mammal expressing a IL-12 with which an antibody of the invention cross-reacts. Still further the subject can be a mammal into which has been introduced hIL-12 (e.g., by administration of hIL-12 or by expression of an hIL-12 transgene). An antibody of the invention can be administered to a human subject for therapeutic 30 purposes (discussed further below). Moreover, an antibody of the invention can be administered to a non-human mammal expressing a IL-12 with which the antibody cross-reacts for veterinary purposes or as an animal model of human disease. Regarding the latter, such animal models may be useful for evaluating the therapeutic efficacy of

antibodies of the invention (e.g., testing of dosages and time courses of administration).

As used herein, the phrase "a disorder in which IL-12 activity is detrimental" is intended to include diseases and other disorders in which the presence of IL-12 in a subject suffering from the disorder has been shown to be or is suspected of being either 5 responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which IL-12 activity is detrimental is a disorder in which inhibition of IL-12 activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of IL-12 in a biological fluid of a subject 10 suffering from the disorder (e.g., an increase in the concentration of IL-12 in serum, plasma, synovial fluid, *etc.* of the subject), which can be detected, for example, using an anti-IL-12 antibody as described above. There are numerous examples of disorders in which IL-12 activity is detrimental. In one embodiment, the antibodies or antigen binding portions thereof, can be used in therapy to treat the diseases or disorders 15 described herein. In another embodiment, the antibodies or antigen binding portions thereof, can be used for the manufacture of a medicine for treating the diseases or disorders described herein. The use of the antibodies and antibody portions of the invention in the treatment of a few non-limiting specific disorders is discussed further below:

20

A. Rheumatoid Arthritis:

Interleukin-12 has been implicated in playing a role in inflammatory diseases such as rheumatoid arthritis. Inducible IL-12p40 message has been detected in synovia from rheumatoid arthritis patients and IL-12 has been shown to be present in the 25 synovial fluids from patients with rheumatoid arthritis (see *e.g.*, Morita *et al.*, (1998) *Arthritis and Rheumatism* 41: 306-314). IL-12 positive cells have been found to be present in the sublining layer of the rheumatoid arthritis synovium. The human antibodies, and antibody portions of the invention can be used to treat, for example, rheumatoid arthritis, juvenile rheumatoid arthritis, Lyme arthritis, rheumatoid 30 spondylitis, osteoarthritis and gouty arthritis. Typically, the antibody, or antibody portion, is administered systemically, although for certain disorders, local administration of the antibody or antibody portion may be beneficial. An antibody, or antibody portion,

of the invention also can be administered with one or more additional therapeutic agents useful in the treatment of autoimmune diseases.

In the collagen induced arthritis (CIA) murine model for rheumatoid arthritis, treatment of mice with an anti-IL-12 mAb (rat anti-mouse IL-12 monoclonal antibody, 5 C17.15) prior to arthritis profoundly suppressed the onset, and reduced the incidence and severity of disease. Treatment with the anti-IL-12 mAb early after onset of arthritis reduced severity, but later treatment of the mice with the anti-IL-12 mAb after the onset of disease had minimal effect on disease severity.

10 B. *Crohn's Disease*

Interleukin-12 also plays a role in the inflammatory bowel disease, Crohn's disease. Increased expression of IFN- γ and IL-12 occurs in the intestinal mucosa of patients with Crohn's disease (see e.g., Fais *et al.*, (1994) *J. Interferon Res.* 14: 235-238; Parronchi *et al.*, (1997) *Amer. J. Pathol.* 150: 823-832; Monteleone *et al.*, (1997) 15 *Gastroenterology* 112: 1169-1178; Berrebi *et al.*, (1998) *Amer. J. Pathol.* 152: 667-672). Anti-IL-12 antibodies have been shown to suppress disease in mouse models of colitis, e.g., TNBS induced colitis IL-2 knockout mice, and recently in IL-10 knock-out mice. Accordingly, the antibodies, and antibody portions, of the invention, can be used in the treatment of inflammatory bowel diseases.

20

C. *Multiple Sclerosis*

Interleukin-12 has been implicated as a key mediator of multiple sclerosis. Expression of the inducible IL-12 p40 message or IL-12 itself can be demonstrated in lesions of patients with multiple sclerosis (Windhagen *et al.*, (1995) *J. Exp. Med.* 182: 25 1985-1996, Drulovic *et al.*, (1997) *J. Neurol. Sci.* 147: 145-150). Chronic progressive patients with multiple sclerosis have elevated circulating levels of IL-12. Investigations with T-cells and antigen presenting cells (APCs) from patients with multiple sclerosis revealed a self-perpetuating series of immune interactions as the basis of progressive multiple sclerosis leading to a Th1-type immune response. Increased secretion of IFN- γ 30 from the T cells led to increased IL-12 production by APCs, which perpetuated the cycle leading to a chronic state of a Th1-type immune activation and disease (Balashov *et al.*, (1997) *Proc. Natl. Acad. Sci.* 94: 599-603). The role of IL-12 in multiple sclerosis has been investigated using mouse and rat experimental allergic encephalomyelitis (EAE)

models of multiple sclerosis. In a relapsing-remitting EAE model of multiple sclerosis in mice, pretreatment with anti-IL-12 mAb delayed paralysis and reduced clinical scores. Treatment with anti-IL-12 mAb at the peak of paralysis or during the subsequent remission period reduced clinical scores. Accordingly, the antibodies or antigen binding portions thereof of the invention may serve to alleviate symptoms associated with multiple sclerosis in humans.

5 **D. *Insulin-Dependent Diabetes Mellitus***

Interleukin-12 has been implicated as an important mediator of insulin-dependent diabetes mellitus (IDDM). IDDM was induced in NOD mice by administration of IL-12, and anti-IL-12 antibodies were protective in an adoptive transfer model of IDDM. Early onset IDDM patients often experience a so-called "honeymoon period" during which some residual islet cell function is maintained. These residual islet cells produce insulin and regulate blood glucose levels better than administered insulin. Treatment of these early onset patients with an anti-IL-12 antibody may prevent further destruction of islet cells, thereby maintaining an endogenous source of insulin.

10 **E. *Psoriasis***

Interleukin-12 (IL-12) and the related cytokine IL-23 have been implicated as key mediators in psoriasis. Psoriasis involves acute and chronic skin lesions that are associated with a TH1-type cytokine expression profile (Hamid et al. (1996) J. Allergy Clin. Immunol. 1:225-231; Turka et al. (1995) Mol. Med. 1:690-699). Both IL-12 and IL-23 contribute to the development of the type 1 T helper cell (Th1) immune response in psoriasis. Moreover, the IL-12 p40 and IL-23 p40 messenger RNA is overexpressed in psoriatic skin lesions. Accordingly, the antibodies or antigen binding portions thereof of the invention may serve to alleviate chronic skin disorders such psoriasis.

In one embodiment, the invention provides a method for treating psoriasis. Treatment for psoriasis often includes a topical corticosteroids, vitamin D analogs, and topical or oral retinoids, or combinations thereof. In one embodiment, an IL-12 and/or IL-23 antibody is administered in combination with or the presence of one of these common treatments. Additional therapeutic agents which can be combined with the IL-12 and/or IL-23 antibody for treatment of psoriasis are described in more detail below.

The diagnosis of psoriasis is usually based on the appearance of the skin. Additionally a skin biopsy, or scraping and culture of skin patches may be needed to rule out other skin disorders. An x-ray may be used to check for psoriatic arthritis if joint pain is present and persistent.

5 Improvements in psoriasis in a subject can be monitored by the subject's Psoriasis Area and Severity Index Score (PASI). The method for determining the PASI has been described in Fredriksson and Pettersson (1978) *Dermatologica* 157:238 and Marks *et al.* (1989) *Arch Dermatol* 125:235. Briefly, the index is based on evaluation of four anatomic sites, including the head, upper extremities, trunk, and lower extremities, 10 for erythema, induration, and desquamation using a 5 point scale (0= no symptoms; 1=slight; 2= moderate; 3=marked; 4=very marked). Based on the extent of lesions in a given anatomic site, the area affected is assigned a numerical value (0=0; 1 = < 10%; 2 = 10-29%; 3 = 30-49%; 4 = 50-69%; 5 = 70-89%; 6 = 90-100%). The PASI score is then calculated, wherein the possible range of PASI score is 0.0 to 72.0 with the highest score 15 representing complete erythroderma of the severest degree.

In one embodiment of the invention, an IL-12 and/or IL-23 antibody is used for the treatment of psoriasis, including chronic plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, pemphigus vulgaris, erythrodermic psoriasis, psoriasis associated with inflammatory bowel disease (IBD), and psoriasis associated with 20 rheumatoid arthritis (RA). In another embodiment, an IL-12 and/or IL-23 antibody, such as J695 / ABT-874, is used to treat subjects who have psoriasis in combination with PsA. Specific types of psoriasis included in the treatment methods of the invention are described in detail below:

25 a. *Chronic plaque psoriasis*

Chronic plaque psoriasis (also referred to as psoriasis vulgaris) is the most common form of psoriasis. Chronic plaque psoriasis is characterized by raised reddened patches of skin, ranging from coin-sized to much larger. In chronic plaque psoriasis, the plaques may be single or multiple, they may vary in size from a few millimeters to several centimeters. The plaques are usually red with a scaly surface, and reflect light 30 when gently scratched, creating a "silvery" effect. Lesions (which are often symmetrical) from chronic plaque psoriasis occur all over body, but with predilection for extensor surfaces, including the knees, elbows, lumbosacral regions, scalp, and nails. Occasionally chronic plaque psoriasis can occur on the penis, vulva and flexures, but

scaling is usually absent. Diagnosis of patients with chronic plaque psoriasis is usually based on the clinical features described above. In particular, the distribution, color and typical silvery scaling of the lesion in chronic plaque psoriasis are characteristic of chronic plaque psoriasis.

5 b. *Guttate psoriasis*

Guttate psoriasis refers to a form of psoriasis with characteristic water drop shaped scaly plaques. Flares of guttate psoriasis generally follow an infection, most notably a streptococcal throat infection. Diagnosis of guttate psoriasis is usually based on the appearance of the skin, and the fact that there is often a history of recent sore

10 throat.

c. *Inverse psoriasis*

Inverse psoriasis is a form of psoriasis in which the patient has smooth, usually moist areas of skin that are red and inflamed, which is unlike the scaling associated with plaque psoriasis. Inverse psoriasis is also referred to as intertiginous psoriasis or

15 flexural psoriasis. Inverse psoriasis occurs mostly in the armpits, groin, under the breasts and in other skin folds around the genitals and buttocks, and, as a result of the locations of presentation, rubbing and sweating can irritate the affected areas.

d. *Pustular psoriasis*

Pustular psoriasis, also referred to as palmar plantar psoriasis, is a form of

20 psoriasis that causes pus-filled blisters that vary in size and location, but often occur on the hands and feet. The blisters may be localized, or spread over large areas of the body. Pustular psoriasis can be both tender and painful, can cause fevers.

e. *Other psoriasis disorders*

Other examples of psoriatic disorders which can be treated with the IL-12 and/or

25 IL-23 antibody include erythrodermic psoriasis, vulgaris, psoriasis associated with IBD, and psoriasis associated with arthritis, including rheumatoid arthritis.

The present invention is further illustrated by the following examples which should not be construed as limiting in any way. The contents of all cited references,

30 including literature references, issued patents, and published patent applications, as cited throughout this application are hereby expressly incorporated herein by reference. It should further be understood that the contents of all the tables attached hereto (see

Appendix A of US Patent No. 6,914,128) as well as the entire contents of U.S. Patent No. 6,914,128 are incorporated herein by reference.

EXAMPLES

5

Example 1: Efficacy of the Fully Human IL-12/IL-23 Monoclonal Antibody, ABT-874, In the Treatment of Moderate to Severe Plaque Psoriasis

ABT-874 is a fully human antibody against interleukin-12 (IL-12) and IL-23. It binds with great affinity to the p40 subunit common to both IL-12 and IL-23, both 10 validated targets in the treatment of psoriasis (Ps).

The objective of the following study was to evaluate the efficacy of subcutaneous injections of ABT-874 in the treatment of patients with moderate to severe plaque Ps.

Adult patients with Ps affecting $\geq 10\%$ body surface area (BSA) and a Psoriasis 15 Area and Severity Index (PASI) score ≥ 12 at baseline were eligible for this 12-week, double-blind, placebo-controlled study. Patients were randomized to 1 of 6 arms: 1) 100-mg ABT-874 every other week (eow) for 12 weeks; 2) one 200-mg ABT-874 dose at Week 0; 3) 200-mg ABT-874 every week for 4 weeks; 4) 200-mg ABT-874 eow for 12 weeks; 5) 200-mg ABT-874 every week for 12 weeks; or 6) placebo. Primary 20 endpoint was a \geq PASI75 response at Week 12. Other efficacy assessments included the PASI50 and Physician's Global Assessment (PGA). Patients who met the primary endpoint entered a 36-week blinded/retreatment phase and were monitored for time to loss of response.

A total of 180 patients enrolled in the study, 30 in each arm. Baseline 25 characteristics were similar between arms and indicative of moderate to severe Ps (all mean values except % male): age, 46 yrs, 74% male; 21 yrs duration of Ps; PASI 19; and 25% BSA affected. At Week 12, the percentages of patients achieving \geq PASI75 were statistically significantly greater for patients in each of the 5 ABT-874 arms vs. placebo (93%, 63%, 90%, 93%, 90%, vs. 3%, respectively, $p<0.001$, ITT). In addition, 30 the percentages of patients achieving \geq PASI50 were statistically significantly greater for patients in each of the 5 ABT-874 arms vs. placebo (100%, 77%, 97%, 97%, and 100%, vs. 17%, $p<0.001$). The mean percentage decreases (improvements) in PASI at Week 12 were 90%, 70%, 92%, 92%, and 90%, respectively, in the ABT-874 arms, and 26% for

placebo. Similarly, the percentages of patients with a PGA of Clear/Minimal were 83%, 50%, 73%, 87% and 87%, respectively, in the ABT-874 arms, and 3% for placebo.

In conclusion, ABT-874 was significantly more efficacious than placebo in the treatment of moderate to severe plaque psoriasis.

5

Example 2: Safety and Efficacy of the Fully Human IL-12/23 Monoclonal Antibody, ABT-874, in the Treatment of Moderate to Severe Plaque Psoriasis

ABT-874 is a fully human antibody against interleukin 12 (IL-12) and IL-23. It binds with great affinity to the p40 subunit common to both IL-12 and IL-23, validated 10 targets in the treatment of psoriasis (Ps). The objective of this Phase II study was to investigate the efficacy and safety of subcutaneous injections of ABT-874 in the treatment of moderate to severe plaque Ps.

Adults with Ps affecting $\geq 10\%$ body surface area (BSA) and a PASI score ≥ 12 were eligible for this 12-wk, double-blind, placebo-controlled study. Patients were 15 randomized to 1 of 6 arms: 1) 100-mg ABT-874 every other week (eow) for 12 wks; 2) one 200-mg ABT-874 dose at Wk 0; 3) 200-mg ABT-874 every wk for 4 wks; 4) 200-mg ABT-874 eow for 12 wks; 5) 200-mg ABT-874 every wk for 12 wks; or 6) placebo. The primary endpoint was a \geq PASI75 response at Wk 12. Patients who met the primary endpoint entered a 36-wk blinded/retreatment phase and were monitored for time to loss 20 of response. All patients were evaluated for safety through Wk 54.

180 patients enrolled, 30 in each arm. Baseline characteristics were similar between arms (mean values presented except % male): age, 46 yrs, 74% male; 21 yrs duration of Ps; PASI=19; and 25% BSA affected. At Wk 12, the %s of patients with \geq PASI75 were statistically significantly greater in each of the 5 ABT-874 arms vs. 25 placebo (93%, 63%, 90%, 93%, 90%, vs. 3%, respectively, $p<0.001$, ITT). During the 12-wk, DB phase, infectious AEs for the ABT-874 groups ranged from 23–43% and for the placebo group was 23%, with the most common being nasopharyngitis (7–17% for ABT-874; 3% for placebo). There were no statistically significant differences between arms. No serious infectious AEs were reported, and no deaths occurred.

30 In conclusion, ABT-874 was significantly more efficacious than placebo in the treatment of moderate to severe plaque Ps, and appears to have a favorable safety profile.

Example 3: Maintenance of Response with the Fully Human IL-12/-23 Monoclonal Antibody, ABT-874, in the Treatment of Moderate to Severe Plaque Psoriasis

The efficacy and safety of ABT-874 was evaluated in a 12-week, Phase II, randomized controlled trial and 36-week follow-up phase. The objective of the

5 following example was to analyze maintenance of response following discontinuation of therapy during the second 12 weeks of this Phase II study of subcutaneous injections of ABT-874 in the treatment of moderate to severe plaque Ps.

Adults with Ps affecting $\geq 10\%$ body surface area (BSA) and a PASI score ≥ 12 were eligible for this 12-week, double-blind, placebo-controlled study. Patients were
10 randomized to 1 of 6 arms:

- 1) 100-mg ABT-874 every other week (eow) for 12 wks;
- 2) one 200-mg ABT-874 dose at Wk 0;
- 3) 200-mg ABT-874 every wk for 4 wks;
- 4) 200-mg ABT-874 eow for 12 wks;
- 15 5) 200-mg ABT-874 every wk for 12 wks; or
- 6) placebo.

The primary endpoint was a \geq PASI75 response at Week 12. Patients who met the primary endpoint entered a 36-week blinded/retreatment phase. Treatment with study drug was discontinued, and patients were monitored for time to loss of response (a
20 decrease in PASI score, any time during the 36-week follow-up period, to $<$ PASI 50). Maintenance of PASI response was evaluated through Week 24.

A total of 180 patients enrolled, 30 in each arm. Baseline characteristics were similar between arms (mean values presented except % male): age, 46 years, 74% male; 21 years duration of Ps; PASI=19; and 25% BSA affected.

25 At Week 12, the percentages of patients with \geq PASI75 were statistically significantly greater in each of the 5 ABT-874 arms vs. placebo (Table I). At Week 24, substantial percentages of PASI 75 responders in the active treatments arms had maintained at least a PASI 50 response.

30 Table 1: 24-Week Efficacy of ABT-874

		Maintenance of PASI Response: Wk 24 vs. Wk 12
	\geq PASI75 at Wk 12	
100 mg eow for 12 wks	28/30 (93%)*	24/28 (86%)
200 mg, one dose	19/30 (63%)*	15/19 (79%)
200-mg every wk for 4 wks	27/30 (90%)*	23/27 (85%)
200-mg eow for 12 wks	28/30 (93%)*	26/28 (93%)
200-mg every wk for 12 wks	27/30 (90%)*	26/27 (96%)
Placebo	1/30 (3%)	—

*p<0.001 vs. placebo, NRI.

In conclusion, ABT-874 was significantly more efficacious than placebo in the treatment of moderate to severe plaque Ps. Substantial percentages of PASI 75

5 responders maintained these responses at Week 24, following discontinuation of active therapy.

Example 4: Safety and Efficacy of ABT-874, a Fully Human IL-12/23 Monoclonal Antibody, in the Treatment of Moderate to Severe Chronic Plaque Psoriasis

10

The objective of the following example was to demonstrate the efficacy and safety of a range of doses of a human IL-12/23 monoclonal antibody (ABT-874) compared with placebo in the treatment of patients with clinically stable moderate to severe chronic plaque psoriasis.

15

I. Materials and Methods

A. Study design: The following study was a 12-week, multicentre, randomised, double-blind, phase II, placebo-controlled trial that was conducted at 24 centres in the United States (16 sites) and Canada (8 sites). ABT-874 (Abbott Laboratories, Abbott Park, IL) is a human monoclonal antibody with genetically engineered complementarity-determining regions that have high affinity for the IL-12/23 p40 subunit protein. Patients

were randomised in a 1:1:1:1:1:1 ratio to receive 1 of 6 treatments: 200 mg of ABT-874, 1 dose at week 0 (200 mg \times 1); 100 mg of ABT-874 every other week (eow) for 12 weeks (100 mg eow); 200 mg of ABT-874 weekly for the first 4 weeks (200 mg \times 4); 200 mg of ABT-874 eow for 12 weeks (200 mg eow); 200 mg of ABT-874 weekly for 5 weeks (200 mg weekly); or placebo. After week 12, all patients who achieved at least a 75% reduction in psoriasis area and severity index (PASI 75) response continued into a 36-week blinded observation/retreatment phase.

B. Patients: Patients were \geq 18 years of age and had a clinical diagnosis of 10 psoriasis for at least 6 months (determined by patient interview and confirmation of diagnosis through physical examination by the investigator), stable plaque psoriasis for at least 2 months before screening and at baseline visits as determined by subject interview, moderate to severe plaque psoriasis defined by \geq 10% body surface area (BSA) involvement at the baseline visit, a PASI score of \geq 12 at the baseline visit, and a 15 physician's global assessment (PGA) of at least moderate disease at the baseline visit.

Patients were ineligible if they had previous exposure to systemic or biologic anti-IL-12 therapy; nonplaque psoriasis; inability to discontinue the following therapies before the baseline visit: topical psoriasis therapies at least 2 weeks before, ultraviolet B light phototherapy at least 2 weeks before, psoralen-ultraviolet-light phototherapy at 20 least 4 weeks before, systemic therapies at least 4 weeks before, and biologic therapies at least 12 weeks before; required intake of oral or injectable corticosteroids during the study (inhaled corticosteroids for stable medical conditions were allowed); an exacerbation of asthma requiring hospitalization in the 10 years prior to screening; an 25 infection or risk factors for severe infection; a history of malignancies other than successfully treated basal cell carcinoma (patients with a history of squamous cell carcinoma were excluded) or cervical carcinoma *in situ*; or a history of major immunologic reaction (eg, serum sickness or anaphylactoid reaction) to an immunoglobulin G-containing agent (eg, intravenous gamma globulin, a fusion protein, or monoclonal antibody).

30 Patients were allowed to continue treatment with medicated shampoos that did not contain corticosteroids, bland (without beta- or alpha-hydroxy acids) emollients, or Class VI or VII low-potency topical corticosteroids on their palms, soles, face, inframammary area, and groin area during the course of the study. Application of these

topical psoriasis therapies was not to occur within 24 hours of a study visit. Vaccination with a live viral agent was not allowed within 1 month prior to dosing with ABT-874, during the study, or for 1 month after the last dose of study drug was administered.

Occurrence of any of the following clinically significant abnormal laboratory results led to immediate withdrawal of a patient from the study: aspartate transaminase or alanine transaminase >5 times the upper limit of normal; serum total bilirubin >3 times the upper limit of normal; serum creatinine >3 times the upper limit of normal; creatine phosphokinase >5 times the upper limit of normal; hemoglobin <8 g/dL; white blood cell count <2 × 10⁹/L; or platelet count <75 × 10⁹/L.

10

C. Efficacy assessments: The primary efficacy assessment was the percentage of patients achieving a PASI 75 response at week 12, defined as at least a 75% reduction in PASI score relative to the baseline score. PASI is a measure of the severity of psoriatic lesions (in terms of erythema, induration, and desquamation) and the extent of BSA involvement. The PASI score ranges from 0 (no psoriasis) to 72 (severe disease) (Fredriksson T, Pettersson U. *Dermatologica* 1978; **157**: 238–44). Other efficacy measures included the percentage of patients who achieved at least PASI 75 at weeks 1, 2, 4, and 8; the percentage of patients who achieved at least PASI 50 or PASI 90 at weeks 1, 2, 4, 8, and 12; and the percentage of patients who attained a PGA of clear or minimal at week 12 and at weeks 1, 2, 4, and 8. The PGA measures the severity of disease on a 6-point scale, which ranges from 0 (no disease, or clear) to 5 (very severe) (Ko H-S. Clinical trial design in psoriasis. Presented at: 49th Meeting of the Dermatologic and Ophthalmologic Advisory Committee; March 20, 1998; Bethesda, MD).

25

D. Safety assessments: Adverse events, laboratory data, and vital signs were assessed throughout the study. Patients were closely monitored for signs of infection, malignancy, and immunologic reaction. Treatment-emergent AEs were defined as those events that occurred between week 0 and the earlier of 45 days after the last nonmissing study drug dose or 1 day prior to the first retreatment dose (for those patients continuing on to the 36-week trial).

E. Statistical analysis: The sample size was calculated using nQuery Advisor® 4.0 (Statistical Solutions, Saugus, MA). With the assumption that 15% of the patients in the placebo group would achieve a PASI 75 response at week 12, the study designers determined that a sample size of 26 in each dosage group would be adequate to detect at 5 least a 45% difference from a treated group using the Fisher exact test with 90% power at a 0.05 2-sided significance level. The study was designed to enroll approximately 180 patients, with 30 patients in each group.

The intention-to-treat population included all patients who were randomised at week 0 and received at least 1 injection of study drug; this population was used for the 10 efficacy analyses. All tests were performed at $\alpha=0.05$. Nonresponder imputation was used for all efficacy analyses; any patient with a missing PASI or PGA score at any visit was considered a nonresponder at that visit. To assess the impact of the missing data, sensitivity analyses of week-12 data were completed using the last-observation-carried-forward method. The primary analysis of PASI 75 response at week 12 was performed 15 using the following sequential order to adjust for multiplicity: 200 mg weekly versus placebo, 200 mg eow versus placebo, 100 mg eow versus placebo, 200 mg \times 4 versus placebo, and 200 mg \times 1 versus placebo. The treatment difference between each ABT-874 treatment group and the placebo group for mean percentage change in PASI score was assessed using analysis of variance, with baseline PASI score and treatment group 20 as factors. The safety analyses were conducted using the safety population, which included all patients who received at least 1 injection of study drug.

II. Results

A. Patients: A total of 180 patients were enrolled and randomised to 1 of the 6 25 treatment groups (Figure 1). The majority of patients (76.7% of placebo-treated patients and 98% of all ABT-874 treatment group patients) completed the 12-week portion of the study.

Patients were well balanced across treatment groups with respect to demographic characteristics and disease activity (table 1). Patients were predominantly male (74.4%) 30 and white (92.2%). Mean BSA involvement was 25% and mean PASI score was 18.8.

B. Efficacy: The percentage of patients achieving the primary endpoint of PASI 75 response at week 12 was statistically significantly greater ($p<0.001$) in all of the

ABT-874 treatment groups (200 mg × 1: 63.3%, 19 of 30; 100 mg eow: 93.3%, 28 of 30; 200 mg × 4: 90.0%, 27 of 30; 200 mg eow: 93.3%, 28 of 30; 200 mg weekly: 90.0%, 27 of 30) compared with placebo (3.3%, 1 of 30). For the relatively short duration of this trial, PASI 75 responses in all ABT-874 treatment groups were similar with the

5 exception of the 200 mg × 1 treatment group (Figure 2).

A subgroup analysis by demographics (gender, age, race, and weight), baseline disease characteristics (history of psoriatic arthritis, BSA, and PASI score), and baseline therapy for psoriasis within 12 months of receiving study treatment (systemic biologic and nonbiologic, topical, and phototherapy) demonstrated that ABT-874–treated patients
10 within the various subgroups consistently achieved high levels of PASI 75 response at week 12.

Nearly 100% of the higher ABT-874 dosage groups attained at least a PASI 50 response by week 12 (200 mg × 1: 76.7%, 23 of 30; 100 mg eow: 100.0%, 30 of 30; 200 mg × 4: 96.7%, 29 of 30; 200 mg eow: 96.7%, 29 of 30; 200 mg weekly: 100.0%, 30 of 30; placebo: 16.7%, 5 of 30; p<0.001 for each comparison with placebo). The percentage of patients achieving at least a PASI 90 response at week 12 was statistically significantly greater (p<0.001) in all but 1 (200 mg × 1) of the ABT-874 treatment groups when compared with placebo, as follows: 200 mg × 1: 16.7%, 5 of 30; 100 mg eow: 53.3%, 16 of 30; 200 mg × 4: 63.3%, 19 of 30; 200 mg eow: 76.6%, 23 of 30; 200
20 mg weekly: 53.3%, 16 of 30; and placebo: 0%, 0 of 30. In addition, by week 12, significantly more (p<0.001) patients in all ABT-874 treatment groups had attained a clear or minimal PGA rating compared with patients in the placebo group, as follows: 200 mg × 1: 50.0%, 15 of 30; 100 mg eow: 83.3%, 25 of 30; 200 mg × 4: 73.3%, 22 of 30; 200 mg eow: 86.7%, 26 of 30; 200 mg weekly: 86.7%, 26 of 30; versus placebo:
25 3.3%, 1 of 30.

The percentage of patients achieving the primary endpoint of PASI 100 response at week 12 was statistically significantly greater (p<0.001) in the following ABT-874 treatment groups (200 mg eow: 46.7%, 14 of 30; 200 mg weekly: 36.7%, 11 of 30) compared with placebo (0%, 0 of 30).

30 Response to ABT-874 was rapid. The mean percentage improvement in PASI scores from baseline increased over time for all ABT-874 treatment groups (Figure 3) and were statistically significantly greater for each ABT-874 treatment group compared

with placebo at each time point ($p<0.001$, except for the 100 mg eow group at week 1, $p=0.023$).

C. Safety: ABT-874 therapy was generally well tolerated (table 2). One (0.7%) patient treated with ABT-874 discontinued the study owing to a localised skin discolouration; 2 (6.7%) patients treated with placebo discontinued the study, 1 for psoriatic arthropathy and 1 for ovarian cancer. Two (1.1%) patients experienced serious adverse effects (AEs); 1 placebo-treated patient was diagnosed with ovarian cancer on day 37, and 1 ABT-874-treated patient (200 mg \times 1) was diagnosed with costochondritis on day 10. No patients experienced myocardial or cerebral infarctions, and there were no deaths.

Patients receiving any dose of ABT-874 were significantly ($p=0.033$) more likely than patients receiving placebo to experience an AE at least possibly related to study drug (ABT-874: 36.0%, 54 of 150; placebo: 10.0%, 3 of 30; table 2); most of these AEs were related to the injection site (injection-site reaction, erythema, pruritus, or irritation).

Most AEs were mild (mild AEs occurred in 46.0% [69 of 150] of ABT-874-treated patients and 30.0% [9 of 30] placebo-treated patients). The most common AE was injection-site reaction, occurring in 16.7% (25 of 150) of patients treated with any dose of ABT-874 (no reported injection-site reactions for placebo-treated patients; $p=0.028$; table 3). There were no statistically significant differences between the incidences of other AEs in the ABT-874-treated patients compared with placebo-treated patients. The next most frequently reported AEs were nasopharyngitis and upper respiratory tract infection.

Infectious AEs were reported by 32.8% (59 of 180) of all patients (placebo: 23.3%, 7 of 30; all ABT-874-treated patients: 34.7%, 52 of 150). The most common infectious AEs reported for any ABT-874 treatment group were nasopharyngitis (12.0%, 18 of 150), upper respiratory tract infection (10.7%, 16 of 150), and bronchitis and viral infection (both 2.7%, 4 of 150). No serious infectious AEs were reported.

Two patients reported malignancies during the study. One placebo-treated patient was diagnosed with ovarian cancer, which was ongoing as of day 129. One ABT-874-treated patient (200 mg \times 4) was diagnosed with a non-melanoma skin cancer

(squamous cell carcinoma) that was removed on day 133. The medical history for this patient included removal of a benign skin growth in March 2005.

There were no clinically significant hematology, chemistry (including blood glucose concentrations), or vital sign changes compared with placebo.

Table 1: Baseline demographics and clinical characteristics

Characteristic	Treatment Group						All	
	200 mg		100 mg		200 mg			
	Placebo N=30	× 1 eow	× 4	eow	weekly	ABT-874 N=150		
Age, y	49±14.4	52±12.0	45±13.8	43±13.8	44±16.0	46±14.0	46±14.1	
Male, No. (%)	22 (73.3)	23 (76.7)	22 (73.3)	21 (70.0)	23 (76.7)	23 (76.7)	112 (74.7)	
White, No. (%)	28 (93.3)	25 (83.3)	28 (93.3)	27 (90.0)	30 (100.0)	28 (93.3)	138 (92.0)	
Weight, kg	89±17.6	94±21.2	94±17.9	92±27.8	93±24.1	95±18.0	94±21.9	
Duration of psoriasis, y	21±12.4	20±13.2	24±14.6	22±14.2	18±11.5	18±10.9	21±13.0	
PASI score	16±2.9	18±6.7	20±6.3	20±7.6	20±6.2	19±6.3	19±6.6	
BSA affected, %	21±9.2	24±13.6	28±15.7	24±13.0	29±16.8	23±12.6	26±14.5	
PGA, No. (%)								
Mild	1 (3.3)	0	0	0	0	0	0	
Moderate	20 (66.7)	19 (63.3)	17 (56.7)	13 (43.3)	15 (50.0)	17 (56.7)	81 (54.0)	

Severe	9 (30.0)	11 (36.7)	12 (40.0)	14 (46.7)	13 (43.3)	11 (36.7)	61 (40.7)
History of PsA, No. (%)	9 (30.0)	7 (23.3)	12 (40.0)	9 (30.0)	6 (20.0)	9 (30.0)	43 (28.7)
Previous psoriasis treatment,*							
No. (%)							
Topical therapy	19 (63.3)	21 (70.0)	26 (86.7)	15 (50.0)	21 (70.0)	23 (76.7)	106 (70.7)
Phototherapy	1 (3.3)	6 (20.0)	4 (13.3)	4 (13.3)	3 (10.0)	5 (16.7)	22 (14.7)
Systemic nonbiologic	6 (20.0)	4 (13.3)	7 (23.3)	5 (16.7)	6 (20.0)	8 (26.7)	30 (20.0)
Systemic biologic	3 (10.0)	3 (10.0)	7 (23.3)	6 (20.0)	4 (13.3)	7 (23.3)	27 (18.0)

Values are mean±SD unless otherwise noted. *Within past 12 months prior to study treatment. BSA=body surface area; eow=every other week; PASI=psoriasis area and severity index; PGA=physician's global assessment; PsA=psoriatic arthritis

Table 2: Clinical treatment-emergent adverse events summary

Event	Treatment Group					All
	200 mg	100 mg	200 mg	200 mg	200 mg	
Placebo	× 1	eow	× 4	eow	weekly	ABT-874
N=30	N=30	N=30	N=30	N=30	N=30	N=150
					No. (%)	
Any AE	18 (60.0)	18 (60.0)	22 (73.3)	21 (70.0)	21 (70.0)	19 (63.3)
Any AE at least possibly drug-related*	3 (10.0)	9 (30.0)	12 (40.0)	14 (46.7)	11 (36.7)	8 (26.7)
Any severe AE	3 (10.0)	1 (3.3)	0	0	0	1 (3.3)
Any serious AE†	1 (3.3)	1 (3.3)	0	0	0	0
Any AE leading to discontinuation of study drug	2 (6.7)	1 (3.3)	0	0	0	1 (0.7)
Any AE at least possibly drug-related* and serious	0	0	0	0	0	0
Any infectious AE	7 (23.3)	7 (23.3)	9 (30.0)	13 (43.3)	13 (43.3)	10 (33.3)
						52 (34.7)

Any serious infectious AE	0	0	0	0	0	0
Any malignant neoplasms	1 (3.3)	0	0	1 (3.3)	0	0
Deaths	0	0	0	0	0	0

* As assessed by the investigator. † Serious adverse events included the following: any event that resulted in death; any event that was life-threatening; any event that resulted in admission to the hospital for any length of time; any event that occurred while the patient was hospitalised and resulted in prolongation of hospital stay; any event that resulted in persistent or significant disability/incapacity; or any important medical event that required medical or surgical intervention to prevent serious outcome. AE=adverse event; eow=every other week.

Table 3: Treatment-emergent adverse events with an incidence $\geq 5\%$ in any treatment group by descending frequency of patients treated with any dosage of ABT-874

Event	Treatment Group					All N=150
	200 mg x N=30	100 mg N=30	200 mg x N=30	200 mg weekly N=30	200 mg eow	
	No. (%)					
Injection-site reaction	0	2 (6.7)	7 (23.3)	5 (16.7)	7 (23.3)	4 (13.3) 25 (16.7)
Nasopharyngitis	1 (3.3)	4 (13.3)	4 (13.3)	3 (10.0)	2 (6.7)	5 (16.7) 18 (12.0)
Upper respiratory tract infection	2 (6.7)	2 (6.7)	4 (13.3)	3 (10.0)	5 (16.7)	2 (6.7) 16 (10.7)
Headache	2 (6.7)	5 (16.7)	0	1 (3.3)	3 (10.0)	2 (6.7) 11 (7.3)
Injection site pruritus	0	0	1 (3.3)	2 (6.7)	2 (6.7)	2 (6.7) 7 (4.7)
Injection site erythema	0	0	0	4 (13.3)	2 (6.7)	1 (3.3) 7 (4.7)
Injection site irritation	0	1 (3.3)	3 (10.0)	2 (6.7)	0	0 6 (4.0)
Fatigue	0	2 (6.7)	2 (6.7)	0	0	1 (3.3) 5 (3.3)
Pain in extremity	0	1 (3.3)	0	0	1 (3.3)	2 (6.7) 4 (2.7)

Arthralgia	0	2 (6.7)	0	0	0	2 (6.7)	4 (2.7)
Viral infection	0	0	0	2 (6.7)	1 (3.3)	1 (3.3)	4 (2.7)
Bronchitis	0	1 (3.3)	0	1 (3.3)	2 (6.7)	0	4 (2.7)
Nausea	1 (3.3)	0	3 (10.0)	0	0	0	3 (2.0)
Otitis externa	0	0	0	0	2 (6.7)	0	2 (1.3)
Vomiting	1 (3.3)	0	0	2 (6.7)	0	0	2 (1.3)
Urinary tract infection	2 (6.7)	1 (3.3)	0	1 (3.3)	0	0	2 (1.3)
Herpes simplex	0	0	2 (6.7)	0	0	0	2 (1.3)
Limb injury	0	2 (6.7)	0	0	0	0	2 (1.3)
Pruritus	2 (6.7)	0	0	0	0	0	0

*As assessed by the investigator.

III. Conclusion

The phase II, multicentre, randomised, double-blind, placebo-controlled trial described in this Example demonstrated statistically and clinically significant efficacy of ABT-874 in the treatment of moderate to severe chronic plaque psoriasis. With the exception of the ABT-874 200 mg × 1 treatment group, 90% or more of patients in all ABT-874 treatment groups achieved PASI 75 or greater by week 12, compared with 3.3% of placebo-treated patients. Even in the group that received only 1 dose of study drug (200 mg × 1), a majority (63.3%) of patients had achieved at least PASI 75 by week 12. In addition, almost 100% of patients treated with ABT-874 reached PASI 50 or greater, which is considered to be a clinically significant improvement (Carlin CS, Feldman SR, Krueger JG, Menter A, Krueger GG. *J Am Acad Dermatol* 2004; **50**: 859–66) by week 12. The results for other secondary endpoints, such as PASI 90 and PGA of clear or minimal, were consistent with and supported the primary efficacy analysis.

Response to ABT-874 was rapid. Statistically significant separation between placebo- and ABT-874–treated patients occurred as early as week 1 for the mean percentage improvement in PASI scores. Improvement was sustained for the 12-week duration of the trial, even for patients in the ABT-874 200 mg × 1 and 200 mg × 4 dosage groups.

ABT-874 was well tolerated, and most AEs were mild. Although ABT-874–treated patients were significantly more likely to experience an AE at least possibly related to study drug, most of these were injection site–related AEs (injection-site reaction, erythema, pruritus, or irritation). There was no apparent association between an increased dose of ABT-874 and an increased incidence of AEs. Of note, there were no myocardial or cerebral infarctions.

Immunologic-related events are of particular interest for patients receiving anti–IL-12/23 antibodies. The most frequently reported infectious AEs were nasopharyngitis, upper respiratory tract infection, bronchitis, and viral infection. There were no serious infectious AEs reported for the duration of this trial. Of the 2 malignancies diagnosed during the study, ovarian cancer was diagnosed in a placebo-treated patient, and non-melanoma skin cancer was diagnosed in an ABT-874–treated patient who had a history of a benign skin growth.

In summary, ABT-874 demonstrated statistically and clinically significant benefit for the treatment of patients with moderate to severe chronic plaque psoriasis, and was well tolerated.

5 **Example 5: Maintenance of Response with the Fully Human IL-12/-23 Monoclonal Antibody, ABT-874, in the Treatment of Moderate to Severe Plaque Psoriasis**

The efficacy and safety of ABT-874 was evaluated in a 12-week, Phase II, randomized controlled trial and 36-week follow-up phase. The objective of the following example was to analyze maintenance of response following discontinuation of 10 therapy during the second 12 weeks of this Phase II study of subcutaneous injections of ABT-874 in the treatment of moderate to severe plaque Ps.

Adults with Ps affecting $\geq 10\%$ body surface area (BSA) and a PASI score ≥ 12 were eligible for this 12-week, double-blind, placebo-controlled study. Patients were randomized to 1 of 6 arms:

15 1) 100-mg ABT-874 every other week (eow) for 12 wks;
2) one 200-mg ABT-874 dose at Wk 0;
3) 200-mg ABT-874 every wk for 4 wks;
4) 200-mg ABT-874 eow for 12 wks;
5) 200-mg ABT-874 every wk for 12 wks; or
20 6) placebo.

The primary endpoint was a \geq PASI 75 response at Week 12. Patients who met the primary endpoint entered a 36-week blinded/retreatment phase. Treatment with study drug was discontinued, and patients were monitored for PASI score at various times during the 36-week follow-up period, including PASI 50, PASI 75 and PASI 90 25 responses. Maintenance of PASI response was evaluated through Week 24.

A total of 180 patients enrolled, 30 in each arm. Baseline characteristics were similar between arms (mean values presented except % male): age, 46 years, 74% male; 21 years duration of Ps; PASI=19; and 25% BSA affected.

At Week 12, the percentages of patients with \geq PASI 75 were statistically 30 significantly greater in each of the 5 ABT-874 arms vs. placebo (Table 4). At Week 24, substantial percentages of PASI 75 responders in the active treatments arms had maintained at least a PASI score of \geq PASI 50. Further, substantial percentages of PASI 75 responders in the active treatments arms had also maintained at least a PASI score of

\geq PASI 75, as well as a PASI score of \geq PASI 90 (Table 4 and Figures 4A-C). The percentage of patients maintaining a PASI 75 response over time during the 24 week period is depicted in Figure 4D.

5 Table 4: 24-Week Efficacy of ABT-874

	\geq PASI 75 at Wk 12	Maintenance of \geq PASI 50 Response: Wk 24 vs. Wk 12	Maintenance of \geq PASI 75 Response: Wk 24 vs. Wk 12	Maintenance of \geq PASI 90 Response: Wk 24 vs. Wk 12
100 mg eow for 12 wks	93%*	71%	60%	33%
200 mg, one dose	63%*	68%	23%	7%
200-mg every wk for 4 wks	90%*	82%	60%	23%
200-mg eow for 12 wks	93%*	89%	73%	53%
200-mg every wk for 12 wks	90%*	85%	83%	57%
Placebo	3%	—	7%	7%

*p<0.001 vs. placebo, NRI.

In conclusion, ABT-874 was significantly more efficacious than placebo in the treatment of moderate to severe plaque Ps. Substantial percentages of PASI 75
10 responders maintained a response of \geq PASI 50, \geq PASI 75, and \geq PASI 90 at Week 24, following discontinuation of active therapy.

EQUIVALENTS

15 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS

1. A method of treating psoriasis in a subject comprising the steps of:
(i) selecting a subject who is suffering from chronic psoriasis; and
5 (ii) administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23; thereby treating chronic psoriasis in the subject.

2. The method of claim 1, wherein said subject has had a clinical diagnosis
10 of psoriasis for at least 6 months.

3. The method of claim 1, wherein said subject has had stable plaque psoriasis for at least 2 months.

15 4. A method of treating psoriasis in a subject comprising the steps of:
(i) selecting a subject who has not had a condition selected from the group consisting of previous exposure to systemic or biologic anti-IL-12 therapy; nonplaque psoriasis; inability to discontinue topical psoriasis therapies at least 2 weeks before treatment; ultraviolet B light phototherapy at least 2 weeks before treatment; psoralen-
20 ultraviolet-light phototherapy at least 4 weeks before treatment; systemic therapies at least 4 weeks before treatment; biologic therapies at least 12 weeks before treatment; required intake of oral or injectable corticosteroids during treatment; an exacerbation of asthma requiring hospitalization in the 10 years prior to screening; an infection or risk factors for severe infection; a history of malignancies other than successfully treated
25 basal cell carcinoma; and a history of major immunologic reaction to an immunoglobulin G-containing agent; and
(ii) administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23; thereby treating psoriasis in the subject.

30 5. A method of treating psoriasis in a subject comprising the steps of:
(i) selecting a subject who has not had vaccination with a live viral agent within 1 month; and
35 (ii) administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23; thereby treating psoriasis in the subject.

6. A method of treating psoriasis in a subject comprising the steps of:

(i) administering an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23 to the subject,

(ii) monitoring the subject for a clinically significant abnormal laboratory result selected from the group consisting of aspartate transaminase or alanine transaminase >5 times the upper limit of normal; serum total bilirubin >3 times the upper limit of normal; serum creatinine >3 times the upper limit of normal; creatine phosphokinase >5 times the upper limit of normal; hemoglobin <8 g/dL; white blood cell count $<2 \times 10^9/L$; and platelet count $<75 \times 10^9/L$;

(iii) discontinuing administration of the antibody, or antigen-binding portion thereof, in a subject in which the clinically significant abnormal laboratory result is detected;

thereby treating psoriasis in the subject.

7. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is administered biweekly.

8. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is administered weekly.

9. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 200 mg.

10. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg.

11. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to the p35 subunit of IL-12.

12. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to a p19 subunit of IL-23.

13. The method of any one of claims 1-6, wherein the antibody, or antigen-binding portion thereof, is capable of binding to the epitope of the p40 subunit when the p40 subunit is bound to the p35 subunit of IL-12 and when the p40 subunit is bound to a p19 subunit of IL-23.

14. The method of any one of claims 1-3, wherein the chronic psoriasis is chronic plaque psoriasis.

5 15. The method of any one of claims 4-6, wherein the psoriasis is chronic psoriasis.

16. The method of claim 15, wherein the chronic psoriasis is chronic plaque psoriasis.

10

17. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 90 response for an extended period following initial administration of the 15 antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

18. The method of claim 17, wherein the extended period is at least about 12 weeks.

20 19. The method of claim 17, wherein the antibody, or antigen-binding portion thereof, is administered biweekly.

20. The method of claim 17, wherein the antibody, or antigen-binding portion thereof, is administered weekly.

25

21. The method of claim 17, wherein the antibody is administered in a single dose.

22. The method of claim 17, wherein the antibody, or antigen-binding 30 portion thereof, is administered in a dose of about 200 mg.

23. The method of claim 17, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg.

35

24. The method of claim 17, wherein the psoriasis is chronic psoriasis.

25. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23 to the subject, wherein the subject maintains a clear or minimal PGA rating for an extended period following initial 5 administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

26. The method of claim 25, wherein the extended period is at least about 12 weeks.

10

27. The method of claim 25, wherein the antibody, or antigen-binding portion thereof, is administered biweekly.

15

28. The method of claim 25, wherein the antibody, or antigen-binding portion thereof, is administered weekly.

29. The method of claim 25, wherein the antibody, or antigen-binding portion thereof, is administered in a single dose.

20

30. The method of claim 25, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 200 mg.

31. The method of claim 25, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg.

25

32. The method of claim 25, wherein the psoriasis is chronic psoriasis.

30

33. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23 to the subject, wherein the subject exhibits an improved PASI score by about 8 weeks following initial administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

35

34. The method of claim 33, wherein the subject exhibits an improved PASI score by about 4 weeks following initial administration of the antibody, or antigen binding portion thereof.

35. The method of claim 33, wherein the subject exhibits an improved PASI score by about 2 weeks following initial administration of the antibody, or antigen binding portion thereof.

5 36. The method of claim 33, wherein the subject exhibits an improved PASI score by about 1 week following initial administration of the antibody, or antigen binding portion thereof.

10 37. The method of claim 33, wherein the antibody, or antigen-binding portion thereof, is administered biweekly.

38. The method of claim 33, wherein the antibody, or antigen-binding portion thereof, is administered weekly.

15 39. The method of claim 33, wherein the antibody, or antigen-binding portion thereof, is administered in a single dose.

40. The method of claim 33, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 200 mg.

20 41. The method of claim 33, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg.

42. The method of claim 33, wherein the psoriasis is chronic psoriasis.

25 43. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 100 response for an extended period following initial administration of the 30 antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

44. The method of claim 43, wherein the extended period is at least about 12 weeks.

35 45. The method of claim 43, wherein the antibody, or antigen-binding portion thereof, is administered biweekly.

46. The method of claim 43, wherein the antibody, or antigen-binding portion thereof, is administered weekly.

47. The method of claim 43, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 200 mg.

48. The method of claim 43, wherein the psoriasis is chronic psoriasis.

49. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 50 response for an extended period following discontinuation of administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

15

50. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 75 response for an extended period following discontinuation of administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

51. A method of treating psoriasis in a subject comprising administering to the subject an antibody, or antigen-binding portion thereof, which is capable of binding to an epitope of the p40 subunit of IL-12 and/or IL-23, wherein the subject maintains at least a PASI 90 response for an extended period following discontinuation of administration of the antibody, or antigen-binding portion thereof, thereby treating psoriasis in the subject.

30 52. The method of any one of claims 49-51, wherein the extended period is at least about 12 weeks.

53. The method of any one of claims 49-51, wherein the antibody is administered for at least about 12 weeks.

35

54. The method of any one of claims 49-51, wherein the antibody, or antigen-binding portion thereof, is administered biweekly.

55. The method of any one of claims 49-51, wherein the antibody, or antigen-binding portion thereof, is administered weekly.

56. The method of any one of claims 49-51, wherein the antibody, or 5 antigen-binding portion thereof, is administered in a single dose.

57. The method of any one of claims 49-51, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 200 mg.

10 58. The method of any one of claims 49-51, wherein the antibody, or antigen-binding portion thereof, is administered in a dose of about 100 mg.

59. The method of any one of claims 49-51, wherein the psoriasis is chronic psoriasis.

15 60. A method of treating psoriasis in a subject comprising administering an antibody directed against human IL-12 and human IL-23 to the subject on a biweekly dosing regimen, such that psoriasis is treated.

SEQUENCE LISTING

<110> Abbott Laboratories, et al.

5 <120> Methods for Treating Psoriasis

<130> BBI-276PC

10 <140> Not yet available

<141> Concurrently Herewith

<150> 60/880767

<151> 2007-01-16

15 <150> 60/904022

<151> 2007-02-27

<150> 60/925960

<151> 2007-04-24

20 <150> 60/961764

<151> 2007-07-24

<150> 60/997012

25 <151> 2007-09-28

<160> 675

<170> PatentIn Ver. 2.0

30 <210> 1

<211> 6

<212> PRT

<213> Homo sapiens

35 <220>

<223> Xaa at position 1 could be either His or Ser

<220>

40 <223> Xaa at position 4 could be either Tyr or His

<220>

<223> Xaa at position 6 could be either Tyr, Asn or Thr

45 <400> 1

Xaa Gly Ser Xaa Asp Xaa

1 5

50 <210> 2

<211> 12

<212> PRT

<213> Homo sapiens

55 <220>

<223> Xaa at position 2 could be either Ser or Thr

<220>

<223> Xaa at position 4 could be either Asp or Glu

60 <220>

<223> Xaa at position 5 could be either Ser, Arg or Lys

<220>

65 <223> Xaa at position 6 could be either Ser, Gly or Tyr

<220>

<223> Xaa at position 7 could be either Leu, Phe, Thr or

Ser
 5 <220>
 <223> Xaa at position 8 could be either Arg, Ser, Thr,
 Trp or His
 10 <220>
 <223> Xaa at position 9 could be either Gly or Pro
 15 <220>
 <223> Xaa at position 10 could be either Ser, Thr, Ala
 or Leu
 20 <220>
 <223> Xaa at position 11 could be either Arg, Ser, Met,
 Thr or Leu
 25 <220> 2
 Gln Xaa Tyr Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10
 30 <210> 3
 <211> 17
 <212> PRT
 <213> Homo sapiens
 35 <210> 3
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 40 <210> 4
 <211> 7
 <212> PRT
 <213> Homo sapiens
 45 <220>
 <223> Xaa at position 1 could be either Gly or Tyr
 50 <220>
 <223> Xaa at position 3 could be either Asp or Ser
 55 <220>
 <223> Xaa at position 4 could be either Gln or Asn
 <400> 4
 Xaa Asn Xaa Xaa Arg Pro Ser
 1 5
 60 <210> 5
 <211> 9
 <212> PRT
 <213> Homo sapiens
 65 <220>
 <223> Xaa represents either Ser or Glu
 <400> 5
 Phe Thr Phe Ser Xaa Tyr Gly Met His
 1 5

5 <210> 6
 <211> 13
 <212> PRT
 5 <213> Homo sapiens

10 <220>
 <223> Xaa at position 1 could be either Ser or Thr

15 10 <220>
 <223> Xaa at position 3 could be either Ser or Gly

15 <220>
 <223> Xaa at position 4 could be either Arg or Ser

15 <220>
 <223> Xaa at position 8 could be either Gly or Val

20 20 <220>
 <223> Xaa at position 9 could be either Ser or Ala

20 <220>
 <223> Xaa at position 10 could be either Asn, Gly or Tyr

25 25 <220>
 <223> Xaa at position 11 could be either Thr or Asp

30 30 <220>
 <223> Xaa at position 13 could be either Lys or His

30 <400> 6
 Xaa Gly Xaa Xaa Ser Asn Ile Xaa Xaa Xaa Val Xaa
 1 5 10

35 35 <210> 7
 <211> 115
 <212> PRT
 <213> Homo sapiens

40 40 <220>
 <223> Xaa at position 6 could be either Gln or Glu

45 45 <220>
 <223> Xaa at position 16 could be either Arg or Gly

45 <220>
 <223> Xaa at position 31 could be either Ser or Glu

50 50 <220>
 <223> Xaa at position 84 could be either Lys or Asn

50 <220>
 <223> Xaa at position 97 could be either Thr, Ala or Lys

55 55 <220>
 <223> Xaa at position 98 could be either Thr or Lys

60 60 <220>
 <223> Xaa at position 99 could be either Ser or His

60 <220>
 <223> Xaa at position 102 could be either Tyr or His

65 65 <220>
 <223> Xaa at position 104 could be either Tyr, Asn or
 Thr

65 <400> 7

Gln Val Gln Leu Val Xaa Ser Gly Gly Gly Val Val Gln Pro Gly Xaa
 1 5 10 15

5 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Xaa Tyr
 20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

10 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Asx
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

15 Leu Gln Met Xaa Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

20 Xaa Xaa Xaa Gly Ser Xaa Asp Xaa Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

25 <210> 8
 <211> 112
 <212> PRT
 <213> Homo sapiens

30 <220>
 <223> Xaa at position 1 could be either Ser or Gln

35 <220>
 <223> Xaa at position 2 could be either Tyr or Ser

<220>
 <223> Xaa at position 13 could be either Thr or Ala

40 <220>
 <223> Xaa at position 23 and 91 could be either Ser or Thr

<220>
 <223> Xaa at position 25 could be either Gly or Ser

<220>
 <223> Xaa at position 26 could be either Arg or Ser

50 <220>
 <223> Xaa at position 30 could be either Gly or Val

<220>
 <223> Xaa at position 31 could be either Ser or Ala

55 <220>
 <223> Xaa at position 35 could be either Lys or His

<220>
 <223> Xaa at position 51 could be either Gly or Lys

<220>
 <223> Xaa at position 54 could be either Gln or Asn

65 <220>
 <223> Xaa at position 79 could be either Val or Leu

<220>

<223> Xaa at position 93 could be either Asp or Glu
 <220>
 <223> Xaa at position 94 could be either Ser, Arg or Lys
 5 <220>
 <223> Xaa at position 95 could be either Ser, Gly or Tyr
 <220>
 10 <223> Xaa at position 96 could be either Leu, Phe, Thr or Ser
 <220>
 <223> Xaa at position 97 could be either Arg, Ser, Thr, 15 Trp or His
 <220>
 <223> Xaa at position 98 could be either Gly or Pro
 20 <220>
 <223> Xaa at position 99 could be either Ser, Thr, Ala or Leu
 <220>
 25 <223> Xaa at position 100 could be either Arg, Ser, Met, Thr or Leu
 <220>
 <223> Xaa at position 101 could be either Val, Ile, Thr, 30 Met or Leu
 <220>
 <223> Xaa at position 32 could be either Asn, Gly or Tyr
 35 <220>
 <223> Xaa at position 33 could be either Thr or Asp
 <220>
 <223> Xaa at position 53 could be either Asp or Ser
 40 <400> 8
 Xaa Xaa Val Leu Thr Gln Pro Pro Ser Val Ser Gly Xaa Pro Gly Gln
 1 5 10 15
 45 Arg Val Thr Ile Ser Cys Xaa Gly Xaa Xaa Ser Asn Ile Xaa Xaa Xaa
 20 25 30
 Xaa Val Xaa Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45
 50 Ile Tyr Xaa Asn Xaa Xaa Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Xaa Gln
 55 65 70 75 80
 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Xaa Tyr Xaa Xaa Xaa Xaa
 85 90 95
 60 Xaa Xaa Xaa Xaa Xaa Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110
 <210> 9
 <211> 6
 <212> PRT
 65 <213> Homo sapiens

```

<220>
<223> Xaa at position 2 could be either Gly, Val, Cys or
      His

5  <220>
<223> Xaa at position 3 could be either Ser or Thr

<220>
<223> Xaa at position 4 could be either His, Thr, Val, Arg, or Ile
10
<220>
<223> Xaa at position 5 could be either Asp or Ser

<220>
15  <223> Xaa at position 6 could be either Asn, Lys, Ala,
      Thr, Ser, Phe, Trp, or His

<400> 9
His Xaa Xaa Xaa Xaa Xaa
20      1           5

<210> 10
<211> 12
<212> PRT
25  <213> Homo sapiens

<220>
<223> Xaa at position 4 could be either Asp or Ser

30  <220>
<223> Xaa at position 5 represents any amino acid

<220>
<223> Xaa at position 6 could be either Gly, Asp, Gln,
35  Leu, Phe, Arg, His, Asn or Tyr

<400> 10
Gln Ser Tyr Xaa Xaa Xaa Thr His Pro Ala Leu Leu
      1           5           10
40
<210> 11
<211> 17
<212> PRT
<213> Homo sapiens

45
<220>
<223> Xaa at position 1 could be either Phe, Thr or Tyr

<220>
50  <223> Xaa at position 3 could be either Arg or Ala

<220>
<223> Xaa at position 5 could be either Asp, Ser, Glu or
      Ala
55
<220>
<223> Xaa at position 6 could be either Gly or Arg

<220>
60  <223> Xaa at position 8 represents any amino acid

<220>
<223> Xaa at position 10 could be either Tyr or Glu

65  <400> 11
Xaa Ile Xaa Tyr Xaa Xaa Ser Xaa Lys Xaa Tyr Ala Asp Ser Val Lys
      1           5           10           15

```

Gly

5 <210> 12
 <211> 7
 <212> PRT
 <213> Homo sapiens

10 <220>
 <223> Xaa at position 1 could be either Gly, Tyr, Ser,
 Thr, Asn or Gln

15 <400> 12
 Xaa Asn Asp Gln Arg Pro Ser
 1 5

20 <210> 13
 <211> 9
 <212> PRT
 <213> Homo sapiens

25 <220>
 <223> Xaa at position 4 and 5 represents any amino acid

25 <220>
 <223> Xaa at position 6 could be either Tyr or His

30 <220>
 <223> Xaa at position 7 could be either Gly, Met, Ala,
 Asn or Ser

35 <400> 13
 Phe Thr Phe Xaa Xaa Xaa Xaa Met His
 1 5

40 <210> 14
 <211> 13
 <212> PRT
 <213> Homo sapiens

45 <220>
 <223> Xaa at position 9 could be either Ser, Cys, Arg,
 Asn, Asp or Thr

50 <220>
 <223> Xaa at position 10 could be either Asn, Met or Ile

50 <220>
 <223> Xaa at position 11 could be either Thr, Tyr, Asp,
 His, Lys or Pro

55 <400> 14
 Ser Gly Gly Arg Ser Asn Ile Gly Xaa Xaa Xaa Val Lys
 1 5 10

60 <210> 15
 <211> 114
 <212> PRT
 <213> Homo sapiens

65 <220>
 <223> Xaa at position 30 could be Ser or Glu

65 <220>
 <223> Xaa at position 83 could be Lys or Asn

65 <220>

<223> Xaa at position 5 could be either Gln or Glu
 <400> 15
 Gln Val Gln Val Xaa Ser Gly Gly Gly Val Val Gln Pro Gly Arg Ser
 5 1 5 10 15
 Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Xaa Tyr Gly
 20 25 30
 10 Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala
 35 40 45
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 50 55 60
 15 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu
 65 70 75 80
 Gln Met Xaa Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Lys
 20 25 30 35 40 45 50 55 60 65 70 75 80 85 90 95 98 99
 Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr Val
 100 105 110
 25 Ser Ser

 <210> 16
 <211> 112
 30 <212> PRT
 <213> Homo sapiens

 <220>
 <223> Xaa at position 1 could be either Ser or Gln
 35
 <220>
 <223> Xaa at position 2 could be Tyr or Ser

 <220>
 40 <223> Xaa at position 13 could be either Thr or Ala

 <220>
 <223> Xaa at position 25 could be either Gly or Ser

 45 <220>
 <223> Xaa at position 51 and 95 could be either Gly or
 Tyr

 <220>
 50 <223> Xaa at position 79 could be either Val or Leu

 <400> 16
 Xaa Xaa Val Leu Thr Gln Pro Pro Ser Val Ser Gly Xaa Pro Gly Gln
 1 5 10 15
 55 Arg Val Thr Ile Ser Cys Ser Gly Xaa Arg Ser Asn Ile Gly Ser Asn
 20 25 30
 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 60 35 40 45
 Ile Tyr Xaa Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 65 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Xaa Gln
 65 70 75 80
 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Xaa Thr

	85	90	95
	His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly		
	100	105	110
5	<210> 17		
	<211> 6		
	<212> PRT		
	<213> Homo sapiens		
10	<400> 17		
	His Gly Ser His Asp Asn		
	1	5	
15	<210> 18		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
20	<400> 18		
	Gln Ser Tyr Asp Arg Gly Thr His Pro Ala Leu Leu		
	1	5	10
25	<210> 19		
	<211> 17		
	<212> PRT		
	<213> Homo sapiens		
30	<400> 19		
	Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys		
	1	5	10
	15		
35	Gly		
	<210> 20		
	<211> 7		
40	<212> PRT		
	<213> Homo sapiens		
	<400> 20		
	Gly Asn Asp Gln Arg Pro Ser		
45	1	5	
	<210> 21		
	<211> 9		
	<212> PRT		
50	<213> Homo sapiens		
	<400> 21		
	Phe Thr Phe Ser Ser Tyr Gly Met His		
	1	5	
55	<210> 22		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
60	<400> 22		
	Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Thr Val Lys		
	1	5	10
65	<210> 23		
	<211> 115		
	<212> PRT		
	<213> Homo sapiens		

<400> 23
 Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 5 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 10 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60
 15 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 20 Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110
 Val Ser Ser
 25 115

 <210> 24
 <211> 112
 30 <212> PRT
 <213> Homo sapiens

 <400> 24
 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
 35 1 5 10 15
 Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Trp Ile Gly Ser Asn
 20 25 30
 40 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45
 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 45 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
 65 70 75 80
 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr
 50 85 90 95
 His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110
 55
 <210> 25
 <211> 6
 <212> PRT
 <213> Homo sapiens
 60
 <400> 25
 His Gly Ser His Asp Asn
 1 5
 65
 <210> 26
 <211> 12
 <212> PRT

<213> Homo sapiens
 <400> 26
 Gln Ser Tyr Asp Arg Tyr Thr His Pro Ala Leu Leu
 5 1 5 10

<210> 27
 <211> 17
 10 <212> PRT
 <213> Homo sapiens

<400> 27
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 15 1 5 10 15

Gly

20 <210> 28
 <211> 7
 <212> PRT
 <213> Homo sapiens

25 <400> 28
 Tyr Asn Asp Gln Arg Pro Ser
 1 5

30 <210> 29
 <211> 9
 <212> PRT
 <213> Homo sapiens

<400> 29
 35 Phe Thr Phe Ser Ser Tyr Gly Met His
 1 5

<210> 30
 <211> 13
 40 <212> PRT
 <213> Homo sapiens

<400> 30
 Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn Thr Val Lys
 45 1 5 10

<210> 31
 <211> 115
 50 <212> PRT
 <213> Homo sapiens

<400> 31
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 55 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

60 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

65 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

5 Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

10 <210> 32
 <211> 112
 <212> PRT
 <213> Homo sapiens

15 <400> 32
 Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15

20 Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn
 20 25 30

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

25 Ile Tyr Tyr Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
 65 70 75 80

30 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Tyr Thr
 85 90 95

35 His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

<210> 33
 <211> 115
 <212> PRT
 40 <213> Homo sapiens

<400> 33
 Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

45 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

50 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

55 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

60 Thr Thr Ser Gly Ser Tyr Asp Tyr Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

<210> 34

<211> 112
 <212> PRT
 <213> Homo sapiens

5 <400> 34
 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
 1 5 10 15

10 Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
 20 25 30

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

15 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
 65 70 75 80

20 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu
 85 90 95

25 Arg Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

<210> 35
 <211> 115
 30 <212> PRT
 <213> Homo sapiens

<400> 35
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

40 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

45 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

50 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys Ser Gly Ser Tyr Asp Tyr Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

55 Val Ser Ser
 115

<210> 36
 60 <211> 112
 <212> PRT
 <213> Homo sapiens

<220>
 65 <223> Xaa at position 32 represents either Gly or Tyr

<400> 36
 Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln

1	5	10	15	
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Xaa				
	20	25	30	
5	Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu			
	35	40	45	
10	Ile Tyr Gly Asn Ser Asn Arg Pro Ser Gly Val Pro Asp Arg Phe Ser			
	50	55	60	
	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln			
	65	70	75	80
15	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu			
	85	90	95	
	Ser Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly			
	100	105	110	
20	<210> 37			
	<211> 115			
	<212> PRT			
25	<213> Homo sapiens			
	<400> 37			
	Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg			
	1	5	10	15
30	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr			
	20	25	30	
	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val			
35	35	40	45	
	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val			
	50	55	60	
40	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr			
	65	70	75	80
	Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys			
	85	90	95	
45	Thr Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr			
	100	105	110	
	Val Ser Ser			
50	115			
	<210> 38			
	<211> 112			
55	<212> PRT			
	<213> Homo sapiens			
	<400> 38			
60	Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln			
	1	5	10	15
	Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn			
	20	25	30	
65	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu			
	35	40	45	
	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser			

	50	55	60
	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln		
5	65 70 75 80		
	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu		
	85 90 95		
10	Arg Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly		
	100 105 110		
	<210> 39		
	<211> 115		
15	<212> PRT		
	<213> Homo sapiens		
	<400> 39		
20	Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg		
	1 5 10 15		
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr		
	20 25 30		
25	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
	35 40 45		
	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50 55 60		
30	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65 70 75 80		
	Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
35	85 90 95		
	Thr Thr Ser Gly Ser Tyr Asp Tyr Trp Gly Gln Gly Thr Met Val Thr		
	100 105 110		
40	Val Ser Ser		
	115		
	<210> 40		
45	<211> 112		
	<212> PRT		
	<213> Homo sapiens		
	<400> 40		
50	Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln		
	1 5 10 15		
	Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn		
	20 25 30		
55	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35 40 45		
	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
60	50 55 60		
	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln		
	65 70 75 80		
65	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Phe		
	85 90 95		
	Thr Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly		

	100	105	110
5	<210> 41 <211> 115 <212> PRT <213> Homo sapiens		
10	<400> 41 Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg 1 5 10 15		
15	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30		
20	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45		
25	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val 50 55 60		
30	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr 65 70 75 80		
35	Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95		
40	Thr Thr Ser Gly Ser Tyr Asp Tyr Trp Gly Gln Gly Thr Met Val Thr 100 105 110		
45	Val Ser Ser 115		
50	<210> 42 <211> 112 <212> PRT <213> Homo sapiens		
55	<400> 42 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln 1 5 10 15		
60	Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn 20 25 30		
65	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu 35 40 45		
70	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser 50 55 60		
75	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln 65 70 75 80		
80	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu 85 90 95		
85	Trp Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly 100 105 110		
90	<210> 43 <211> 115 <212> PRT		

<213> Homo sapiens

<400> 43

5 Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

10 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

15 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

20 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Thr Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
100 105 110

25 Val Ser Ser
115

<210> 44

30 <211> 112

<212> PRT

<213> Homo sapiens

<400> 44

35 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
20 25 30

40 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

45 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
65 70 75 80

50 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Phe
85 90 95

55 Thr Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
100 105 110

55

60 <210> 45

<211> 115

<212> PRT

<213> Homo sapiens

65 <400> 45

Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

5 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

10 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

15 Thr Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

20 Val Ser Ser
 115

25 <210> 46
 <211> 112
 <212> PRT
 <213> Homo sapiens

30 <400> 46
 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
 20 25 30

35 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

40 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
 65 70 75 80

45 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu
 85 90 95

Trp Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

50

55 <210> 47
 <211> 115
 <212> PRT
 <213> Homo sapiens

60 <400> 47
 Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

65 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

15 <210> 48
 <211> 112
 <212> PRT
 20 <213> Homo sapiens

<400> 48
 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
 1 5 10 15

25 Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn
 20 25 30

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 30 35 40 45

Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

35 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
 65 70 75 80

Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Thr Tyr Asp Lys Gly Phe
 85 90 95

40 Thr Gly Ser Ser Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

45

<210> 49
 <211> 115
 50 <212> PRT
 <213> Homo sapiens

<400> 49
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 55 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

60 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

65 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

5 Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

10 <210> 50
 <211> 112
 <212> PRT
 <213> Homo sapiens

15 <400> 50
 Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15

20 Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn
 20 25 30

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

25 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

30 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
 65 70 75 80

Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Thr Tyr Asp Lys Gly Phe
 85 90 95

35 Thr Gly Ser Ser Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

40 <210> 51
 <211> 115
 <212> PRT
 45 <213> Homo sapiens

<400> 51
 Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

55 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

65 Thr Thr His Gly Ser His Asp Thr Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

5 <210> 52
 <211> 112
 <212> PRT
 <213> Homo sapiens

10 <400> 52
 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
 1 5 10 15

15 Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
 20 25 30

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

20 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
 65 70 75 80

25 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu
 85 90 95

30 Trp Gly Thr Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

35

<210> 53
 <211> 115
 <212> PRT
 <213> Homo sapiens

40 <400> 53
 Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

45 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

50 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

55 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

60 Thr Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

Val Ser Ser
 115

65

<210> 54
 <211> 112

<212> PRT
<213> Homo sapiens

<400> 54
5 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Val Ser Asn
20 25 30

10 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

15 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
65 70 75 80

20 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Phe
85 90 95

Thr Gly Ser Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
100 105 110

25

30 <210> 55
<211> 115
<212> PRT
<213> Homo sapiens

35 <400> 55
Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

45 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

50 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

55 Thr Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
100 105 110

Val Ser Ser
115

60

<210> 56
<211> 112
<212> PRT
<213> Homo sapiens

65 <400> 56
Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
1 5 10 15

	Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Val Ser Asn
	20 25 30
5	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
	35 40 45
	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
	50 55 60
10	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
	65 70 75 80
15	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Phe
	85 90 95
	Thr Gly Ala Arg Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
	100 105 110
20	
25	<210> 57
	<211> 115
	<212> PRT
	<213> Homo sapiens
	<400> 57
30	Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
	1 5 10 15
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
	20 25 30
35	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
	35 40 45
40	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
	50 55 60
	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
	65 70 75 80
45	Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
	85 90 95
	Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
	100 105 110
50	Val Ser Ser
	115
55	<210> 58
	<211> 112
	<212> PRT
	<213> Homo sapiens
60	<400> 58
	Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
	1 5 10 15
65	Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
	20 25 30
	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
	35 40 45

	Ile	Tyr	Gly	Asn	Asp	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
	50					55					60					
5	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Thr	Gly	Val	Gln
	65					70					75					80
	Ala	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Gln	Thr	Tyr	Asp	Lys	Gly	Phe
						85				90					95	
10	Thr	Gly	Ser	Ser	Val	Phe	Gly	Thr	Gly	Thr	Lys	Val	Thr	Val	Leu	Gly
					100				105					110		
15																
	<210>	59														
20	<211>	115														
	<212>	PRT														
	<213>	Homo sapiens														
	<400>	59														
25	Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
	1				5						10			15		
	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr
					20				25					30		
30	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
					35				40					45		
	Ala	Phe	Ile	Arg	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala	Asp	Ser	Val
					50				55					60		
35	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
					65				70			75			80	
	Leu	Gln	Met	Lys	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
40						85				90					95	
	Lys	Thr	His	Gly	Ser	His	Asp	Asn	Trp	Gly	Gln	Gly	Thr	Met	Val	Thr
						100				105					110	
45	Val	Ser	Ser													
			115													
	<210>	60														
50	<211>	112														
	<212>	PRT														
	<213>	Homo sapiens														
	<400>	60														
55	Ser	Tyr	Val	Leu	Thr	Gln	Pro	Pro	Ser	Val	Ser	Gly	Thr	Pro	Gly	Gln
	1				5					10				15		
	Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Gly	Arg	Ser	Asn	Ile	Gly	Ser	Asn
					20				25					30		
60	Thr	Val	Lys	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	Leu
					35				40					45		
	Ile	Tyr	Gly	Asn	Asp	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
65					50				55					60		
	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Thr	Gly	Val	Gln
					65				70			75			80	

	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Glu Arg Gly Phe		
	85	90	95
5	Thr Gly Ser Met Val Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly		
	100	105	110
10			
	<210> 61		
	<211> 115		
	<212> PRT		
15	<213> Homo sapiens		
	<400> 61		
	Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg		
	1	5	10
20	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr		
	20	25	30
25	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
	35	40	45
	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50	55	60
30	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
	Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90	95
35	Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr		
	100	105	110
	Val Ser Ser		
40	115		
	<210> 62		
	<211> 112		
45	<212> PRT		
	<213> Homo sapiens		
	<400> 62		
	Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln		
50	1	5	10
	Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn		
	20	25	30
55	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35	40	45
	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
	50	55	60
60	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln		
	65	70	75
	80		
65	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr		
	85	90	95
	His Pro Leu Thr Ile Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly		
	100	105	110

5 <210> 63
 <211> 115
 <212> PRT
 <213> Homo sapiens
 10 <400> 63
 Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 15 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 20 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60
 25 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 30 Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110
 Val Ser Ser
 115
 35 <210> 64
 <211> 112
 <212> PRT
 40 <213> Homo sapiens
 <400> 64
 Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
 1 5 10 15
 45 Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
 20 25 30
 50 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45
 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 55 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
 65 70 75 80
 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Ser
 85 90 95
 60 His Pro Ala Leu Thr Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110
 65 <210> 65

<211> 115
<212> PRT
<213> Homo sapiens

5 <400> 65
Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
10 20 25 30
Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
15 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
20 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
25 Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
100 105 110
Val Ser Ser
115
30
<210> 66
<211> 112
<212> PRT
<213> Homo sapiens
35 <400> 66
Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln
1 5 10 15
40 Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn
20 25 30
Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45
45 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60
50 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln
65 70 75 80
Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr
85 90 95
55 His Pro Leu Thr Met Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
100 105 110
60
<210> 67
<211> 115
<212> PRT
65 <213> Homo sapiens
<400> 67
Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg

	1	5	10	15
5	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr	20	25	30
10	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val	35	40	45
15	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val	50	55	60
20	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr	65	70	75
25	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	85	90	95
30	Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr	100	105	110
35	Val Ser Ser	115		
40	<210> 68			
45	<211> 112			
50	<212> PRT			
55	<213> Homo sapiens			
60	<400> 68			
65	Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln	1	5	10
70	Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn	20	25	30
75	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu	35	40	45
80	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser	50	55	60
85	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln	65	70	75
90	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr	85	90	95
95	His Pro Leu Thr Met Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly	100	105	110
100	<210> 69			
105	<211> 115			
110	<212> PRT			
115	<213> Homo sapiens			
120	<400> 69			
125	Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg	1	5	10
130	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr	20	25	30
135	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val			

	35	40	45
	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50	55	60
5	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
10	85	90	95
	Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr		
	100	105	110
15	Val Ser Ser		
	115		
	<210> 70		
20	<211> 112		
	<212> PRT		
	<213> Homo sapiens		
	<400> 70		
25	Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln		
	1	5	10
	Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn		
	20	25	30
30	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35	40	45
	Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
35	50	55	60
	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln		
	65	70	75
	80		
40	Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr		
	85	90	95
	His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly		
	100	105	110
45			
	<210> 71		
50	<211> 115		
	<212> PRT		
	<213> Homo sapiens		
55	<400> 71		
	Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg		
	1	5	10
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Glu Tyr		
60	20	25	30
	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
	35	40	45
65	Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50	55	60
	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		

65	70	75	80
Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys			
85 90 95			
5	Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr		
100 105 110			
10	Val Ser Ser		
115			
<210> 72			
<211> 112			
15	<212> PRT		
<213> Homo sapiens			
<400> 72			
Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln			
20	1	5	10
15			
Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn			
20 25 30			
25	Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
35 40 45			
Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser			
50 55 60			
30	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln		
65 70 75 80			
Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr			
35	85 90 95		
His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly			
100 105 110			
40			
<210> 73			
45	<211> 115		
<212> PRT			
<213> Homo sapiens			
<400> 73			
Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg			
50	1	5	10
15			
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr			
20 25 30			
55	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
35 40 45			
Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val			
60	50	55	60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr			
65 70 75 80			
65	Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
85 90 95			
Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr			

100

105

110

Val Ser Ser
115

5

<210> 74
<211> 112
<212> PRT
10 <213> Homo sapiens

<400> 74
Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

15 Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn
20 25 30

20 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

Ile Tyr Tyr Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50 55 60

25 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
65 70 75 80

Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Gly Thr
85 90 95

30 His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
100 105 110

35

<210> 75
<211> 115
40 <212> PRT
<213> Homo sapiens

<400> 75
Gln Val Gln Leu Val Gln Ser Gly Gly Val Val Gln Pro Gly Arg
45 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

50 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

55 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

60 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
100 105 110

65 Val Ser Ser
115

<210> 76
 <211> 112
 <212> PRT
 <213> Homo sapiens
 5
 <400> 76
 Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15

10 Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn
 20 25 30

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

15 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
 20 65 70 75 80

Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Tyr Thr
 85 90 95

25 His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

30

<210> 77
 <211> 6
 <212> PRT
 35 <213> Homo sapiens

<400> 77
 Ser Gly Ser Tyr Asp Tyr
 1 5

40

<210> 78
 <211> 6
 <212> PRT
 45 <213> Homo sapiens

<400> 78
 His Gly Ser His Asp Asn
 1 5

50

<210> 79
 <211> 6
 <212> PRT
 55 <213> Homo sapiens

<400> 79
 His Gly Ser Tyr Asp Tyr
 1 5

60

<210> 80
 <211> 6
 <212> PRT
 65 <213> Homo sapiens

<400> 80
 Arg Arg Arg Ser Asn Tyr

1 5

5 <210> 81
<211> 6
<212> PRT
<213> Homo sapiens

10 <400> 81
Ser Gly Ser Ile Asp Tyr
1 5

15 <210> 82
<211> 6
<212> PRT
<213> Homo sapiens

20 <400> 82
His Gly Ser His Asp Asp
1 5

25 <210> 83
<211> 6
<212> PRT
<213> Homo sapiens

30 <400> 83
His Gly Ser His Asp Asn
1 5

35 <210> 84
<211> 12
<212> PRT
<213> Homo sapiens

40 <400> 84
Thr Thr His Gly Ser His Asp Asn Trp Gly Gln Gly
1 5 10

45 <210> 85
<211> 12
<212> PRT
<213> Homo sapiens

50 <400> 85
Ala Lys His Gly Ser His Asp Asn Trp Gly Gln Gly
1 5 10

55 <210> 86
<211> 12
<212> PRT
<213> Homo sapiens

60 <400> 86
Thr Thr His Gly Ser His Asp Asn Trp Ser Gln Gly
1 5 10

65 <210> 87
<211> 12
<212> PRT
<213> Homo sapiens

<400> 87
Thr Thr His Gly Ser His Asp Thr Trp Gly Gln Gly
1 5 10

5
<210> 88
<211> 12
<212> PRT
<213> Homo sapiens

10
<400> 88
Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly
1 5 10

15
<210> 89
<211> 12
<212> PRT
<213> Homo sapiens

20
<400> 89
Lys Thr His Gly Ser His Asp Asn Trp Gly His Gly
1 5 10

25
<210> 90
<211> 12
<212> PRT
<213> Homo sapiens

30
<400> 90
Thr Thr His Gly Ser His Asp Asn Trp Ser Gln Gly
1 5 10

35
<210> 91
<211> 12
<212> PRT
<213> Homo sapiens

40
<400> 91
Thr Thr His Arg Ser His Asn Asn Trp Gly Gln Gly
1 5 10

45
<210> 92
<211> 8
<212> PRT
<213> Homo sapiens

50
<400> 92
Thr Thr His Gly Ser His Asp Asn
1 5

55
<210> 93
<211> 8
<212> PRT
<213> Homo sapiens

60
<400> 93
Thr Thr His Gly Ser His Asp Thr
1 5

65
<210> 94
<211> 8
<212> PRT

<213> Homo sapiens

<400> 94
5 Thr Lys His Gly Ser His Asp Asn
1 5

<210> 95
<211> 8
<212> PRT
10 <213> Homo sapiens

<400> 95
15 Thr Thr Gln Gly Arg His Asp Asn
1 5

<210> 96
20 <211> 8
<212> PRT
<213> Homo sapiens

<400> 96
25 Lys Thr Arg Gly Arg His Asp Asn
1 5

<210> 97
30 <211> 8
<212> PRT
<213> Homo sapiens

<400> 97
35 Thr Thr His Gly Ser His Asp Lys
1 5

<210> 98
40 <211> 8
<212> PRT
<213> Homo sapiens

<400> 98
45 Thr Thr His Gly Ser His Asp Asp
1 5

<210> 99
50 <211> 8
<212> PRT
<213> Homo sapiens

<400> 99
55 Lys Thr His Gly Ser His Asp Asn
1 5

<210> 100
60 <211> 8
<212> PRT
<213> Homo sapiens

<400> 100
65 Lys Thr His Gly Ser His Asp Asn
1 5

5 <210> 101
 <211> 8
 <212> PRT
 <213> Homo sapiens
5 <400> 101
 Thr Thr His Gly Ser His Asp Asn
 1 5
10 <210> 102
 <211> 8
 <212> PRT
 <213> Homo sapiens
15 <400> 102
 Thr Thr Ser Gly Ser Tyr Asp Tyr
 1 5
20 <210> 103
 <211> 8
 <212> PRT
 <213> Homo sapiens
25 <400> 103
 Thr Thr His Gly Ser His Asp Asn
 1 5
30 <210> 104
 <211> 8
 <212> PRT
 <213> Homo sapiens
35 <400> 104
 Thr Thr His Gly Ser Gln Asp Asn
 1 5
40 <210> 105
 <211> 8
 <212> PRT
 <213> Homo sapiens
45 <400> 105
 Ala Thr His Gly Ser Gln Asp Asn
 1 5
50 <210> 106
 <211> 6
 <212> PRT
 <213> Homo sapiens
55 <400> 106
 His Gly Ser Gln Asp Thr
 1 5
60 <210> 107
 <211> 6
 <212> PRT
 <213> Homo sapiens
65 <400> 107
 Ser Gly Ser Tyr Asp Tyr
 1 5

```

      <210> 108
      <211> 6
  5  <212> PRT
      <213> Homo sapiens

      <400> 108
      His Gly Ser Gln Asp Asn
 10     1             5

      <210> 109
      <211> 9
 15  <212> PRT
      <213> Homo sapiens

      <400> 109
      Cys Lys Thr His Gly Ser His Asp Asn
 20     1             5

      <210> 110
      <211> 12
 25  <212> PRT
      <213> Homo sapiens

      <400> 110
      Gln Ser Tyr Asp Ser Ser Leu Arg Gly Ser Arg Val
 30     1             5             10

      <210> 111
      <211> 12
 35  <212> PRT
      <213> Homo sapiens

      <400> 111
      Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Arg Val
 40     1             5             10

      <210> 112
      <211> 12
 45  <212> PRT
      <213> Homo sapiens

      <400> 112
      Gln Ser Tyr Asp Ser Ser Leu Arg Gly Ser Arg Val
 50     1             5             10

      <210> 113
      <211> 12
 55  <212> PRT
      <213> Homo sapiens

      <400> 113
      Gln Ser Tyr Asp Ser Ser Leu Thr Gly Ser Arg Val
 60     1             5             10

      <210> 114
      <211> 12
 65  <212> PRT
      <213> Homo sapiens

      <400> 114

```

Gln Ser Tyr Asp Ser Ser Leu Trp Gly Ser Arg Val
1 5 10

5 <210> 115
<211> 12
<212> PRT
<213> Homo sapiens

10 <400> 115
Gln Thr Tyr Asp Ile Ser Glu Ser Gly Ser Arg Val
1 5 10

15 <210> 116
<211> 12
<212> PRT
<213> Homo sapiens

20 <400> 116
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Arg Val
1 5 10

25 <210> 117
<211> 12
<212> PRT
<213> Homo sapiens

30 <400> 117
Gln Thr Tyr Asp Arg Gly Phe Thr Gly Ser Arg Val
1 5 10

35 <210> 118
<211> 12
<212> PRT
<213> Homo sapiens

40 <400> 118
Gln Thr Tyr Asp Lys Gly Phe Thr Gly Ser Ser Val
1 5 10

45 <210> 119
<211> 12
<212> PRT
<213> Homo sapiens

50 <400> 119
Gln Ser Tyr Asp Arg Arg Phe Thr Gly Ser Arg Val
1 5 10

55 <210> 120
<211> 12
<212> PRT
<213> Homo sapiens

60 <400> 120
Gln Ser Tyr Asp Trp Asn Phe Thr Gly Ser Arg Val
1 5 10

65 <210> 121
<211> 12
<212> PRT
<213> Homo sapiens

<400> 121
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Arg Val
1 5 10
5
<210> 122
<211> 12
<212> PRT
10 <213> Homo sapiens

<400> 122
Gln Ser Tyr Asp Asn Gly Phe Thr Gly Ser Arg Val
1 5 10
15
<210> 123
<211> 12
<212> PRT
20 <213> Homo sapiens

<400> 123
Gln Ser Tyr Asp Asn Ala Val Thr Ala Ser Lys Val
1 5 10
25
<210> 124
<211> 12
<212> PRT
30 <213> Homo sapiens

<400> 124
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Arg Val
1 5 10
35
<210> 125
<211> 12
<212> PRT
40 <213> Homo sapiens

<400> 125
Gln Ser Tyr Asp Ser Ser Leu Trp Gly Thr Arg Val
1 5 10
45
<210> 126
<211> 12
<212> PRT
50 <213> Homo sapiens

<400> 126
Gln Ser Tyr Asp Arg Asp Phe Thr Gly Ser Arg Val
1 5 10
55
<210> 127
<211> 12
<212> PRT
60 <213> Homo sapiens

<400> 127
Gln Ser Tyr Glu Arg Gly Phe Thr Gly Ser Met Val
1 5 10
65
<210> 128
<211> 12

<212> PRT
<213> Homo sapiens

5 <400> 128
Gln Ser Tyr Asp Asn Gly Phe Thr Gly Ala Arg Val
1 5 10

10 <210> 129
<211> 12
<212> PRT
<213> Homo sapiens

15 <400> 129
Gln Ser Tyr Asp Arg Arg Phe Thr Gly Ser Arg Val
1 5 10

20 <210> 130
<211> 12
<212> PRT
<213> Homo sapiens

25 <400> 130
Gln Thr Tyr Asp Lys Gly Phe Thr Gly Ser Ser Val
1 5 10

30 <210> 131
<211> 12
<212> PRT
<213> Homo sapiens

35 <400> 131
Gln Ser Tyr Asp Arg Asp Phe Thr Gly Thr Arg Val
1 5 10

40 <210> 132
<211> 12
<212> PRT
<213> Homo sapiens

45 <400> 132
Gln Ser Tyr Asp Arg Gly Phe Tyr Gly Ser Met Val
1 5 10

50 <210> 133
<211> 12
<212> PRT
<213> Homo sapiens

55 <400> 133
Gln Thr Tyr Asp Lys Gly Phe Thr Gly Ser Ser Val
1 5 10

60 <210> 134
<211> 12
<212> PRT
<213> Homo sapiens

65 <400> 134
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ala Arg Val
1 5 10

5 <210> 135
<211> 12
<212> PRT
<213> Homo sapiens
<400> 135
Gln Ser Tyr Glu Arg Gly Phe Thr Gly Ala Arg Val
1 5 10
10 <210> 136
<211> 13
<212> PRT
<213> Homo sapiens
15 <400> 136
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Arg Val Phe
1 5 10
20 <210> 137
<211> 13
<212> PRT
<213> Homo sapiens
25 <400> 137
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Phe Lys Val Phe
1 5 10
30 <210> 138
<211> 13
<212> PRT
<213> Homo sapiens
35 <400> 138
Gln Ser Tyr Asp Arg Gly Phe Val Ser Ala Tyr Val Phe
1 5 10
40 <210> 139
<211> 13
<212> PRT
<213> Homo sapiens
45 <400> 139
Gln Ser Tyr Asp Arg Gly Leu Thr Val Thr Lys Val Phe
1 5 10
50 <210> 140
<211> 13
<212> PRT
<213> Homo sapiens
55 <400> 140
Gln Ser Tyr Asp Arg Gly Tyr Thr Ala Ser Arg Val Phe
1 5 10
60 <210> 141
<211> 13
<212> PRT
<213> Homo sapiens
65 <400> 141
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Lys Val Phe
1 5 10

5 <210> 142
 <211> 13
 <212> PRT
 <213> Homo sapiens

10 <400> 142
 Gln Ser Tyr Asp Arg Gly Leu Thr Gly Phe Arg Val Phe
 1 5 10

15 <210> 143
 <211> 13
 <212> PRT
 <213> Homo sapiens

20 <400> 143
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Tyr Lys Val Phe
 1 5 10

25 <210> 144
 <211> 13
 <212> PRT
 <213> Homo sapiens

30 <400> 144
 Gln Ser Tyr Asp Arg Gly Leu Thr Gly Tyr Arg Leu Phe
 1 5 10

35 <210> 145
 <211> 13
 <212> PRT
 <213> Homo sapiens

40 <400> 145
 Gln Ser Tyr Asp Arg Gly Phe Thr Asp Tyr Lys Val Phe
 1 5 10

45 <210> 146
 <211> 13
 <212> PRT
 <213> Homo sapiens

50 <400> 146
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Pro Arg Leu Phe
 1 5 10

55 <210> 147
 <211> 13
 <212> PRT
 <213> Homo sapiens

60 <400> 147
 Gln Ser Tyr Asp Arg Gly Leu Thr Gly Ser Arg Val Phe
 1 5 10

65 <210> 148
 <211> 13
 <212> PRT
 <213> Homo sapiens

 <400> 148

Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ala Arg Val Trp
1 5 10

5 <210> 149
<211> 13
<212> PRT
<213> Homo sapiens

10 <400> 149
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Tyr Arg Val Phe
1 5 10

15 <210> 150
<211> 13
<212> PRT
<213> Homo sapiens

20 <400> 150
Gln Ser Tyr Asp Arg Gly Phe Thr Gly Pro Arg Val Phe
1 5 10

25 <210> 151
<211> 13
<212> PRT
<213> Homo sapiens

30 <400> 151
Gln Ser Tyr Asp Arg Gly Met Thr Ser Ser Arg Val Phe
1 5 10

35 <210> 152
<211> 13
<212> PRT
<213> Homo sapiens

40 <400> 152
Gln Ser Tyr Asp Arg Asp Ser Thr Gly Ser Arg Val Phe
1 5 10

45 <210> 153
<211> 13
<212> PRT
<213> Homo sapiens

50 <400> 153
Gln Ser Tyr Asp Ser Ser Leu Arg Gly Ser Arg Val Phe
1 5 10

55 <210> 154
<211> 13
<212> PRT
<213> Homo sapiens

60 <400> 154
His Ser Tyr Asp Ser Asp Phe Thr Gly Ser Arg Val Phe
1 5 10

65 <210> 155
<211> 13
<212> PRT
<213> Homo sapiens

<400> 155
His Ser Ser Glu Ser Gly Phe Thr Gly Ser Arg Val Phe
1 5 10
5

<210> 156
<211> 13
<212> PRT
10 <213> Homo sapiens

<400> 156
His Ser Tyr Asp Asn Arg Phe Thr Gly Ser Arg Val Phe
1 5 10
15

<210> 157
<211> 13
<212> PRT
20 <213> Homo sapiens

<400> 157
His Ser Tyr Asp Ser Arg Phe Thr Gly Ser Arg Val Phe
1 5 10
25

<210> 158
<211> 13
<212> PRT
30 <213> Homo sapiens

<400> 158
Gln Ser Tyr Asp Ser Glu Phe Thr Gly Ser Arg Val Phe
1 5 10
35

<210> 159
<211> 13
<212> PRT
40 <213> Homo sapiens

<400> 159
Gln Ser Tyr Asp Thr Gly Phe Thr Gly Ser Arg Val Phe
1 5 10
45

<210> 160
<211> 13
<212> PRT
50 <213> Homo sapiens

<400> 160
His Ser Tyr Asp Ser Gly Phe Thr Gly Ser Arg Val Phe
1 5 10
55

<210> 161
<211> 13
<212> PRT
60 <213> Homo sapiens

<400> 161
Gln Ser Tyr Asp Thr Gly Phe Thr Gly Ser Arg Val Phe
1 5 10
65

<210> 162
<211> 13

<212> PRT
<213> Homo sapiens

5 <400> 162
His Ser Tyr Asp Thr Lys Phe Thr Gly Ser Arg Val Phe
1 5 10

10 <210> 163
<211> 13
<212> PRT
<213> Homo sapiens

15 <400> 163
His Ser Ser Asp Ser Gly Phe Thr Gly Ser Arg Val Phe
1 5 10

20 <210> 164
<211> 13
<212> PRT
<213> Homo sapiens

25 <400> 164
Gln Ser Tyr Asp Ser Asp Phe Thr Gly Ser Arg Val Phe
1 5 10

30 <210> 165
<211> 13
<212> PRT
<213> Homo sapiens

35 <400> 165
His Ser Tyr Glu Ser Gly Phe Thr Gly Ser Arg Val Phe
1 5 10

40 <210> 166
<211> 13
<212> PRT
<213> Homo sapiens

45 <400> 166
Gln Ser Tyr Asp Ala Pro Trp Ser Gly Ser Arg Val Phe
1 5 10

50 <210> 167
<211> 13
<212> PRT
<213> Homo sapiens

55 <400> 167
Gln Ser Tyr Asp Ser Asp Phe Thr Gly Ser Lys Val Phe
1 5 10

60 <210> 168
<211> 13
<212> PRT
<213> Homo sapiens

65 <400> 168
His Thr Asn Asp Ser Gly Phe Thr Gly Ser Arg Val Phe
1 5 10

<210> 169
 <211> 13
 <212> PRT
 <213> Homo sapiens
 5
 <400> 169
 His Ser Tyr Asp Thr Arg Phe Thr Gly Ser Arg Val Phe
 1 5 10

10
 <210> 170
 <211> 13
 <212> PRT
 <213> Homo sapiens
 15
 <400> 170
 Gln Ser Tyr Asp Met Arg Phe Thr Gly Ser Arg Val Phe
 1 5 10

20
 <210> 171
 <211> 13
 <212> PRT
 <213> Homo sapiens
 25
 <400> 171
 His Ser Ser Asp Ser Asp Ser Thr Gly Ser Arg Val Phe
 1 5 10

30
 <210> 172
 <211> 13
 <212> PRT
 <213> Homo sapiens
 35
 <400> 172
 Gln Ser Tyr Asn Thr Asp Phe Thr Gly Ser Arg Val Phe
 1 5 10

40
 <210> 173
 <211> 13
 <212> PRT
 <213> Homo sapiens
 45
 <400> 173
 Gln Ser Tyr Asp Ser Gly Phe Thr Gly Ser Arg Val Phe
 1 5 10

50
 <210> 174
 <211> 13
 <212> PRT
 <213> Homo sapiens
 55
 <400> 174
 His Ser Tyr Asp Met Gly Phe Thr Gly Ser Arg Val Phe
 1 5 10

60
 <210> 175
 <211> 13
 <212> PRT
 <213> Homo sapiens
 65
 <400> 175
 His Ser Tyr Asp Asn Gly Phe Thr Gly Ser Arg Val Phe
 1 5 10

5 <210> 176
<211> 13
<212> PRT
<213> Homo sapiens

10 <400> 176
His Ser His Asp Arg Asp Phe Thr Gly Ser Arg Val Phe
1 5 10

15 <210> 177
<211> 12
<212> PRT
<213> Homo sapiens

20 <400> 177
Gln Ser Tyr Asp Ser Ser Leu Arg Gly Ser Arg Val
1 5 10

25 <210> 178
<211> 13
<212> PRT
<213> Homo sapiens

30 <400> 178
Gln Ser Tyr Asp Arg Gly Ile His Gly Ser Arg Val Phe
1 5 10

35 <210> 179
<211> 13
<212> PRT
<213> Homo sapiens

40 <400> 179
Gln Ser Tyr Asp Ser Gly Phe Pro Gly Ser Arg Val Phe
1 5 10

45 <210> 180
<211> 13
<212> PRT
<213> Homo sapiens

50 <400> 180
Gln Ser Tyr Asp Ile Gly Ser Thr Gly Ser Arg Val Phe
1 5 10

55 <210> 181
<211> 13
<212> PRT
<213> Homo sapiens

60 <400> 181
Gln Ser Tyr Asp Ser Gly Leu Thr Gly Ser Arg Val Phe
1 5 10

65 <210> 182
<211> 13
<212> PRT
<213> Homo sapiens

<400> 182

Gln Ser Tyr Asp Ile Gly Met Thr Gly Ser Arg Val Phe
1 5 10

5 <210> 183
<211> 13
<212> PRT
<213> Homo sapiens

10 <400> 183
Gln Ser Tyr Asp Ile Gly Leu Thr Gly Ser Arg Val Phe
1 5 10

15 <210> 184
<211> 13
<212> PRT
<213> Homo sapiens

20 <400> 184
Gln Ser Tyr Asp Ser Gly Val Thr Gly Ser Arg Val Phe
1 5 10

25 <210> 185
<211> 13
<212> PRT
<213> Homo sapiens

30 <400> 185
Gln Ser Tyr Asp Arg Gly Leu Thr Ala Ser Arg Val Phe
1 5 10

35 <210> 186
<211> 13
<212> PRT
<213> Homo sapiens

40 <400> 186
Gln Ser Tyr Asp Thr Gly Leu Thr Gly Ser Arg Val Phe
1 5 10

45 <210> 187
<211> 13
<212> PRT
<213> Homo sapiens

50 <400> 187
Gln Ser Tyr Asp Thr Ala Leu Thr Gly Ser Arg Val Phe
1 5 10

55 <210> 188
<211> 13
<212> PRT
<213> Homo sapiens

60 <400> 188
Gln Ser Tyr Asp Ile Arg Phe Thr Gly Ser Arg Val Phe
1 5 10

65 <210> 189
<211> 13
<212> PRT
<213> Homo sapiens

<400> 189
 Gln Ser Tyr Asp Ile Arg Ser Thr Gly Ser Arg Val Phe
 1 5 10
 5

<210> 190
 <211> 13
 <212> PRT
 10 <213> Homo sapiens

<400> 190
 Gln Ser Tyr Asp Asn Arg Leu Thr Gly Ser Arg Val Phe
 1 5 10
 15

<210> 191
 <211> 13
 <212> PRT
 20 <213> Homo sapiens

<400> 191
 Gln Ser Tyr Glu Thr Ser Phe Thr Gly Ser Arg Val Phe
 1 5 10
 25

<210> 192
 <211> 13
 <212> PRT
 30 <213> Homo sapiens

<400> 192
 Gln Ser Tyr Asp Ser Ser Ser Thr Gly Ser Arg Val Phe
 1 5 10
 35

<210> 193
 <211> 13
 <212> PRT
 40 <213> Homo sapiens

<400> 193
 Gln Ser Tyr Asp Ser Gly Phe Thr Ala Ser Arg Val Phe
 1 5 10
 45

<210> 194
 <211> 13
 <212> PRT
 50 <213> Homo sapiens

<400> 194
 Gln Thr Tyr Asp Lys Gly Phe Thr Gly Ser Ser Val Phe
 1 5 10
 55

<210> 195
 <211> 13
 <212> PRT
 60 <213> Homo sapiens

<400> 195
 Gln Ser Tyr Asp Asn Gly Phe Thr Gly Ser Arg Val Phe
 1 5 10
 65

<210> 196
 <211> 13

<212> PRT
<213> Homo sapiens

5 <400> 196
Gln Ser Tyr Asp Thr Gly Phe Thr Lys Ser Arg Val Phe
1 5 10

10 <210> 197
<211> 13
<212> PRT
<213> Homo sapiens

15 <400> 197
Gln Ser Tyr Asp Ser Asp Val Thr Gly Ser Arg Val Phe
1 5 10

20 <210> 198
<211> 13
<212> PRT
<213> Homo sapiens

25 <400> 198
Gln Ser Tyr Asp Ala Gly Phe Thr Gly Ser Arg Val Phe
1 5 10

30 <210> 199
<211> 12
<212> PRT
<213> Homo sapiens

35 <400> 199
Gln Ser Tyr Asp Arg Gly Thr His Pro Ser Met Leu
1 5 10

40 <210> 200
<211> 12
<212> PRT
<213> Homo sapiens

45 <400> 200
Gln Ser Tyr Asp Arg Gly Thr Thr Pro Arg Pro Met
1 5 10

50 <210> 201
<211> 12
<212> PRT
<213> Homo sapiens

55 <400> 201
Gln Ser Tyr Asp Arg Gly Arg Asn Pro Ala Leu Thr
1 5 10

60 <210> 202
<211> 12
<212> PRT
<213> Homo sapiens

65 <400> 202
Gln Ser Tyr Asp Arg Gly Thr His Pro Trp Leu His
1 5 10

<210> 203
<211> 12
<212> PRT
<213> Homo sapiens
5
<400> 203
Gln Ser Tyr Asp Arg Gly Asn Ser Pro Ala Thr Val
1 5 10

10
<210> 204
<211> 12
<212> PRT
<213> Homo sapiens
15
<400> 204
Gln Ser Tyr Asp Arg Gly Thr Phe Pro Ser Pro Gln
1 5 10

20
<210> 205
<211> 12
<212> PRT
<213> Homo sapiens
25
<400> 205
Gln Ser Tyr Asp Arg Gly Leu Asn Pro Ser Ala Thr
1 5 10

30
<210> 206
<211> 12
<212> PRT
<213> Homo sapiens
35
<400> 206
Gln Ser Tyr Asp Arg Gly Lys Ser Asn Lys Met Leu
1 5 10

40
<210> 207
<211> 12
<212> PRT
<213> Homo sapiens
45
<400> 207
Gln Ser Tyr Asp Arg Gly His Thr Ala His Leu Tyr
1 5 10

50
<210> 208
<211> 12
<212> PRT
<213> Homo sapiens
55
<400> 208
Gln Ser Tyr Asp Arg Gly Gln Thr Pro Ser Ile Thr
1 5 10

60
<210> 209
<211> 12
<212> PRT
<213> Homo sapiens
65
<400> 209
Gln Ser Tyr Asp Arg Gly Tyr Pro Arg Asn Ile Leu
1 5 10

5 <210> 210
 <211> 12
 <212> PRT
 <213> Homo sapiens

10 <400> 210
 Gln Ser Tyr Asp Arg Gly Ile Thr Pro Gly Leu Ala
 1 5 10

15 <210> 211
 <211> 12
 <212> PRT
 <213> Homo sapiens

20 <400> 211
 Gln Ser Tyr Asp Arg Gly Gln Pro His Ala Val Leu
 1 5 10

25 <210> 212
 <211> 12
 <212> PRT
 <213> Homo sapiens

30 <400> 212
 Gln Ser Tyr Asp Arg Gly Asn Ser Pro Ile Pro Thr
 1 5 10

35 <210> 213
 <211> 12
 <212> PRT
 <213> Homo sapiens

40 <400> 213
 Gln Ser Tyr Asp Arg Gly Thr Pro Asn Asn Ser Phe
 1 5 10

45 <210> 214
 <211> 12
 <212> PRT
 <213> Homo sapiens

50 <400> 214
 Gln Ser Tyr Asp Ser Gly Val Asp Pro Gly Pro Tyr
 1 5 10

55 <210> 215
 <211> 12
 <212> PRT
 <213> Homo sapiens

60 <400> 215
 Gln Ser Tyr Asp Arg Gly Arg Pro Arg His Ala Leu
 1 5 10

65 <210> 216
 <211> 12
 <212> PRT
 <213> Homo sapiens

 <400> 216

Gln Ser Tyr Asp Arg Gly Pro Tyr His Pro Ile Arg
1 5 10

5 <210> 217
<211> 12
<212> PRT
<213> Homo sapiens

10 <400> 217
Gln Ser Tyr Asp Arg Gly Pro His Thr Gln Pro Thr
1 5 10

15 <210> 218
<211> 12
<212> PRT
<213> Homo sapiens

20 <400> 218
Gln Ser Tyr Asp Arg Gly His Asn Asn Phe Ser Pro
1 5 10

25 <210> 219
<211> 12
<212> PRT
<213> Homo sapiens

30 <400> 219
Gln Ser Tyr Asp Arg Gly Pro Thr His Leu Pro His
1 5 10

35 <210> 220
<211> 12
<212> PRT
<213> Homo sapiens

40 <400> 220
Gln Ser Tyr Asp Arg Gly Thr Pro Ser Tyr Pro Thr
1 5 10

45 <210> 221
<211> 12
<212> PRT
<213> Homo sapiens

50 <400> 221
Gln Ser Tyr Asp Ser Gly Thr Ser Asn Leu Leu Pro
1 5 10

55 <210> 222
<211> 12
<212> PRT
<213> Homo sapiens

60 <400> 222
Gln Ser Tyr Asp Arg Gly Asp Ser Asn His Asp Leu
1 5 10

65 <210> 223
<211> 12
<212> PRT
<213> Homo sapiens

<400> 223
Gln Ser Tyr Asp Arg Gly Leu Pro Arg Leu Thr His
1 5 10
5

<210> 224
<211> 12
<212> PRT
10 <213> Homo sapiens

<400> 224
Gln Ser Tyr Asp Arg Gly Ile Pro Thr Ser Tyr Leu
1 5 10
15

<210> 225
<211> 12
<212> PRT
20 <213> Homo sapiens

<400> 225
Gln Ser Tyr Asp Arg Gly Leu Arg Val Gln Ala Pro
1 5 10
25

<210> 226
<211> 12
<212> PRT
30 <213> Homo sapiens

<400> 226
Gln Ser Tyr Asp Arg Gly Leu Ser Asp Ser Pro Leu
1 5 10
35

<210> 227
<211> 12
<212> PRT
40 <213> Homo sapiens

<400> 227
Gln Ser Tyr Asp Ser Gly Ser Leu Arg Arg Ile Leu
1 5 10
45

<210> 228
<211> 12
<212> PRT
50 <213> Homo sapiens

<400> 228
Gln Ser Tyr Asp Arg Gly Pro Ala Arg Thr Ser Pro
1 5 10
55

<210> 229
<211> 12
<212> PRT
60 <213> Homo sapiens

<400> 229
Gln Ser Tyr Asp Arg Gly Arg Ala Ala His Pro Gln
1 5 10
65

<210> 230
<211> 12

<212> PRT
<213> Homo sapiens

5 <400> 230
1 Gln Ser Tyr Asp Arg Gly Thr Gln Pro Ala Asx Ile
5 10

10 <210> 231
<211> 12
<212> PRT
<213> Homo sapiens

15 <400> 231
1 Gln Ser Tyr Asp Arg Gly Thr His Pro Thr Met Ile
5 10

20 <210> 232
<211> 12
<212> PRT
<213> Homo sapiens

25 <400> 232
1 Gln Ser Tyr Asp Arg Gly Arg Ile Pro Ala Asx Thr
5 10

30 <210> 233
<211> 12
<212> PRT
<213> Homo sapiens

35 <400> 233
1 Gln Ser Tyr Asp Arg Gly Thr His Pro Val Pro Ala
5 10

40 <210> 234
<211> 12
<212> PRT
<213> Homo sapiens

45 <400> 234
1 Gln Ser Tyr Asp Arg Gly Ser Asx Pro Ile Pro Ala
5 10

50 <210> 235
<211> 12
<212> PRT
<213> Homo sapiens

55 <400> 235
1 Gln Ser Tyr Asp Arg Gly Thr His Pro Val Pro Ala
5 10

60 <210> 236
<211> 12
<212> PRT
<213> Homo sapiens

65 <400> 236
1 Gln Ser Tyr Asp Arg Gly Thr His Pro Thr Met Tyr
5 10

5 <210> 237
<211> 12
<212> PRT
<213> Homo sapiens
<400> 237
Gln Ser Tyr Asp Arg Gly His His Tyr Thr Thr Phe
1 5 10

10 <210> 238
<211> 12
<212> PRT
<213> Homo sapiens
<400> 238
Gln Ser Tyr Asp Arg Gly Ser His Pro Ala Ala Glu
1 5 10

15 <210> 239
<211> 12
<212> PRT
<213> Homo sapiens
<400> 239
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Ser Ile Glu
1 5 10

20 <210> 240
<211> 12
<212> PRT
<213> Homo sapiens
<400> 240
Gln Ser Tyr Asp Arg Gly Ser Ser Pro Ala Ile Met
1 5 10

25 <210> 241
<211> 12
<212> PRT
<213> Homo sapiens
<400> 241
Gln Ser Tyr Asp Arg Gly Ile Trp Pro Asn Leu Asn
1 5 10

30 <210> 242
<211> 12
<212> PRT
<213> Homo sapiens
<400> 242
Gln Ser Tyr Asp Arg Gly Thr His Pro Asn Leu Asn
1 5 10

35 <210> 243
<211> 12
<212> PRT
<213> Homo sapiens
<400> 243
Gln Ser Tyr Asp Arg Gly Thr His Pro Ser Ile Ser
1 5 10

40 <210> 244
<211> 12
<212> PRT
<213> Homo sapiens
<400> 244
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Asn Leu Asn
1 5 10

45 <210> 245
<211> 12
<212> PRT
<213> Homo sapiens
<400> 245
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Asn Leu Asn
1 5 10

50 <210> 246
<211> 12
<212> PRT
<213> Homo sapiens
<400> 246
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Asn Leu Asn
1 5 10

55 <210> 247
<211> 12
<212> PRT
<213> Homo sapiens
<400> 247
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Asn Leu Asn
1 5 10

60 <210> 248
<211> 12
<212> PRT
<213> Homo sapiens
<400> 248
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Asn Leu Asn
1 5 10

65 <210> 249
<211> 12
<212> PRT
<213> Homo sapiens
<400> 249
Gln Ser Tyr Asp Arg Gly Thr Ile Pro Asn Leu Asn
1 5 10

5 <210> 244
 <211> 12
 <212> PRT
 <213> Homo sapiens

10 <400> 244
 Gln Ser Tyr Asp Arg Gly Ser Ala Pro Met Ile Asn
 1 5 10

15 <210> 245
 <211> 12
 <212> PRT
 <213> Homo sapiens

20 <400> 245
 Gln Ser Tyr Asp Arg Gly His His Pro Ala Met Ser
 1 5 10

25 <210> 246
 <211> 12
 <212> PRT
 <213> Homo sapiens

30 <400> 246
 Gln Ser Tyr Asp Arg Gly Thr His Pro Ser Ile Thr
 1 5 10

35 <210> 247
 <211> 12
 <212> PRT
 <213> Homo sapiens

40 <400> 247
 Gln Ser Tyr Asp Arg Gly Thr Asp Pro Ala Ile Val
 1 5 10

45 <210> 248
 <211> 12
 <212> PRT
 <213> Homo sapiens

50 <400> 248
 Gln Ser Tyr Asp Arg Gly Thr His Pro Ala Leu Leu
 1 5 10

55 <210> 249
 <211> 12
 <212> PRT
 <213> Homo sapiens

60 <400> 249
 Gln Ser Tyr Asp Arg Gly Ser His Pro Ala Leu Thr
 1 5 10

65 <210> 250
 <211> 12
 <212> PRT
 <213> Homo sapiens

 <400> 250

Gln Ser Tyr Asp Arg Gly Thr Thr Pro Ala Pro Glu
1 5 10

5 <210> 251
<211> 12
<212> PRT
<213> Homo sapiens

10 <400> 251
Gln Ser Tyr Asp Arg Gly Ser His Pro Thr Leu Ile
1 5 10

15 <210> 252
<211> 12
<212> PRT
<213> Homo sapiens

20 <400> 252
Gln Ser Tyr Asp Arg Gly Thr His Pro Ser Met Leu
1 5 10

25 <210> 253
<211> 12
<212> PRT
<213> Homo sapiens

30 <400> 253
Gln Ser Tyr Asp Arg Gly Thr Thr Pro Arg Pro Met
1 5 10

35 <210> 254
<211> 12
<212> PRT
<213> Homo sapiens

40 <400> 254
Gln Ser Tyr Asp Arg Gly Arg Leu Pro Ala Gln Thr
1 5 10

45 <210> 255
<211> 12
<212> PRT
<213> Homo sapiens

50 <400> 255
Gln Ser Tyr Asp Arg Gly Thr His Pro Leu Thr Ile
1 5 10

55 <210> 256
<211> 12
<212> PRT
<213> Homo sapiens

60 <400> 256
Gln Ser Tyr Asp Arg Gly Gln Thr Pro Ser Ile Thr
1 5 10

65 <210> 257
<211> 12
<212> PRT
<213> Homo sapiens

<400> 257
Gln Ser Tyr Asp Arg Gly Thr His Phe Gln Met Tyr
1 5 10
5

<210> 258
<211> 12
<212> PRT
10 <213> Homo sapiens

<400> 258
Gln Ser Tyr Asp Arg Gly Arg Asn Pro Ala Leu Thr
1 5 10
15

<210> 259
<211> 12
<212> PRT
20 <213> Homo sapiens

<400> 259
Gln Ser Tyr Asp Arg Gly Thr His Pro Leu Thr Met
1 5 10
25

<210> 260
<211> 12
<212> PRT
30 <213> Homo sapiens

<400> 260
Gln Ser Tyr Asp Arg Gly Thr His Pro Leu Thr Met
1 5 10
35

<210> 261
<211> 12
<212> PRT
40 <213> Homo sapiens

<400> 261
Gln Ser Tyr Asp Ser Gly Tyr Thr Gly Ser Arg Val
1 5 10
45

<210> 262
<211> 12
<212> PRT
50 <213> Homo sapiens

<400> 262
Gln Ser Tyr Asp Ser Gly Phe Thr Gly Ser Arg Val
1 5 10
55

<210> 263
<211> 12
<212> PRT
60 <213> Homo sapiens

<400> 263
Gln Ser Tyr Asp Ser Arg Phe Thr Gly Ser Arg Val
1 5 10
65

<210> 264
<211> 12

<212> PRT
<213> Homo sapiens

5 <400> 264
Gln Ser Tyr Pro Asp Gly Thr Pro Ala Ser Arg Val
1 5 10

10 <210> 265
<211> 12
<212> PRT
<213> Homo sapiens

15 <400> 265
Gln Ser Tyr Ser Thr His Met Pro Ile Ser Arg Val
1 5 10

20 <210> 266
<211> 12
<212> PRT
<213> Homo sapiens

25 <400> 266
Gln Ser Tyr Asp Ser Gly Ser Thr Gly Ser Arg Val
1 5 10

30 <210> 267
<211> 12
<212> PRT
<213> Homo sapiens

35 <400> 267
Gln Ser Tyr Pro Asn Ser Tyr Pro Ile Ser Arg Val
1 5 10

40 <210> 268
<211> 10
<212> PRT
<213> Homo sapiens

45 <400> 268
Gln Ser Tyr Ile Arg Ala Pro Gln Gln Val
1 5 10

50 <210> 269
<211> 12
<212> PRT
<213> Homo sapiens

55 <400> 269
Gln Ser Tyr Leu Lys Ser Arg Ala Phe Ser Arg Val
1 5 10

60 <210> 270
<211> 12
<212> PRT
<213> Homo sapiens

65 <400> 270
Gln Ser Tyr Asp Ser Arg Phe Thr Gly Ser Arg Val
1 5 10

5 <210> 271
 <211> 12
 <212> PRT
 <213> Homo sapiens
10 <400> 271
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Met Val
 1 5 10
15 <210> 272
 <211> 12
 <212> PRT
 <213> Homo sapiens
20 <400> 272
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Met Val
 1 5 10
25 <210> 273
 <211> 12
 <212> PRT
 <213> Homo sapiens
30 <400> 273
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Phe Asp Gly
 1 5 10
35 <210> 274
 <211> 12
 <212> PRT
 <213> Homo sapiens
40 <400> 274
 Gln Ser Tyr Asp Arg Gly Thr Ala Pro Ala Leu Ser
 1 5 10
45 <210> 275
 <211> 12
 <212> PRT
 <213> Homo sapiens
50 <400> 275
 Gln Ser Tyr Asp Arg Gly Ser Tyr Pro Ala Leu Arg
 1 5 10
55 <210> 276
 <211> 12
 <212> PRT
 <213> Homo sapiens
60 <400> 276
 Gln Ser Tyr Asp Arg Gly Asn Trp Pro Asn Ser Asn
 1 5 10
65 <210> 277
 <211> 12
 <212> PRT
 <213> Homo sapiens
70 <400> 277
 Gln Ser Tyr Asp Arg Gly Thr Ala Pro Ser Leu Leu
 1 5 10

5 <210> 278
 <211> 12
 <212> PRT
 <213> Homo sapiens

10 <400> 278
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Met Val
 1 5 10

15 <210> 279
 <211> 12
 <212> PRT
 <213> Homo sapiens

20 <400> 279
 Gln Ser Tyr Asp Arg Gly Thr Thr Pro Arg Ile Arg
 1 5 10

25 <210> 280
 <211> 12
 <212> PRT
 <213> Homo sapiens

30 <400> 280
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Met Val
 1 5 10

35 <210> 281
 <211> 12
 <212> PRT
 <213> Homo sapiens

40 <400> 281
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Met Val
 1 5 10

45 <210> 282
 <211> 12
 <212> PRT
 <213> Homo sapiens

50 <400> 282
 Gln Ser Tyr Asp Arg Gly Met Ile Pro Ala Leu Thr
 1 5 10

55 <210> 283
 <211> 12
 <212> PRT
 <213> Homo sapiens

60 <400> 283
 Gln Ser Tyr Asp Arg Asn Thr His Pro Ala Leu Leu
 1 5 10

65 <210> 284
 <211> 12
 <212> PRT
 <213> Homo sapiens

 <400> 284

Gln Ser Tyr Asp Arg Phe Thr His Pro Ala Leu Leu
1 5 10

5 <210> 285
<211> 12
<212> PRT
<213> Homo sapiens

10 <400> 285
Gln Ser Tyr Asp Arg Tyr Thr His Pro Ala Leu Leu
1 5 10

15 <210> 286
<211> 12
<212> PRT
<213> Homo sapiens

20 <400> 286
Gln Ser Tyr Asp Arg Gly Thr His Pro Ala Leu Leu
1 5 10

25 <210> 287
<211> 12
<212> PRT
<213> Homo sapiens

30 <400> 287
Gln Ser Tyr Asp Arg Tyr Thr His Pro Ala Leu Leu
1 5 10

35 <210> 288
<211> 9
<212> PRT
<213> Homo sapiens

40 <400> 288
Phe Thr Phe Glu Ser Tyr Gly Met His
1 5

45 <210> 289
<211> 9
<212> PRT
<213> Homo sapiens

50 <400> 289
Phe Thr Phe Ser Ser Tyr Gly Met His
1 5

55 <210> 290
<211> 9
<212> PRT
<213> Homo sapiens

60 <400> 290
Phe Thr Phe Tyr Ser Tyr Gly Met His
1 5

65 <210> 291
<211> 9
<212> PRT
<213> Homo sapiens

<400> 291
Phe Thr Phe His Ser Tyr Gly Met His
1 5
5

<210> 292
<211> 9
<212> PRT
10 <213> Homo sapiens

<400> 292
Phe Thr Phe Lys Ser Tyr Gly Met His
1 5
15

<210> 293
<211> 9
<212> PRT
20 <213> Homo sapiens

<400> 293
Phe Thr Phe Arg Ser Tyr Gly Met His
1 5
25

<210> 294
<211> 9
<212> PRT
30 <213> Homo sapiens

<400> 294
Phe Thr Phe Asn Ser Tyr Gly Met His
1 5
35

<210> 295
<211> 9
<212> PRT
40 <213> Homo sapiens

<400> 295
Phe Thr Phe Thr Ser Tyr Gly Met His
1 5
45

<210> 296
<211> 9
<212> PRT
50 <213> Homo sapiens

<400> 296
Phe Thr Phe Gly Ser Tyr Gly Met His
1 5
55

<210> 297
<211> 9
<212> PRT
60 <213> Homo sapiens

<400> 297
Phe Thr Phe Val Ser Tyr Gly Met His
1 5
65

<210> 298
<211> 9

<212> PRT
<213> Homo sapiens

5 <400> 298
Phe Thr Phe Ile Ser Tyr Gly Met His
1 5

10 <210> 299
<211> 9
<212> PRT
<213> Homo sapiens

15 <400> 299
Phe Thr Phe Trp Ser Tyr Gly Met His
1 5

20 <210> 300
<211> 9
<212> PRT
<213> Homo sapiens

25 <400> 300
Phe Thr Phe Ser Glu Tyr Gly Met His
1 5

30 <210> 301
<211> 9
<212> PRT
<213> Homo sapiens

35 <400> 301
Phe Thr Phe Ser Cys Tyr Gly Met His
1 5

40 <210> 302
<211> 9
<212> PRT
<213> Homo sapiens

45 <400> 302
Phe Thr Phe Ser Ser Tyr Gly Met His
1 5

50 <210> 303
<211> 9
<212> PRT
<213> Homo sapiens

55 <400> 303
Phe Thr Phe Ser Tyr Tyr Gly Met His
1 5

60 <210> 304
<211> 9
<212> PRT
<213> Homo sapiens

65 <400> 304
Phe Thr Phe Ser His Tyr Gly Met His
1 5

<210> 305
<211> 9
<212> PRT
<213> Homo sapiens
5
<400> 305
Phe Thr Phe Ser Arg Tyr Gly Met His
1 5
10
<210> 306
<211> 9
<212> PRT
<213> Homo sapiens
15
<400> 306
Phe Thr Phe Ser Asn Tyr Gly Met His
1 5
20
<210> 307
<211> 9
<212> PRT
<213> Homo sapiens
25
<400> 307
Phe Thr Phe Ser Gln Tyr Gly Met His
1 5
30
<210> 308
<211> 9
<212> PRT
<213> Homo sapiens
35
<400> 308
Phe Thr Phe Ser Thr Tyr Gly Met His
1 5
40
<210> 309
<211> 9
<212> PRT
<213> Homo sapiens
45
<400> 309
Phe Thr Phe Ser Ala Tyr Gly Met His
1 5
50
<210> 310
<211> 9
<212> PRT
<213> Homo sapiens
55
<400> 310
Phe Thr Phe Ser Ile Tyr Gly Met His
1 5
60
<210> 311
<211> 9
<212> PRT
<213> Homo sapiens
65
<400> 311
Phe Thr Phe Ser Ser Glu Gly Met His
1 5

5 <210> 312
 <211> 9
 <212> PRT
 <213> Homo sapiens

10 <400> 312
 Phe Thr Phe Ser Ser Cys Gly Met His
 1 5

15 <210> 313
 <211> 9
 <212> PRT
 <213> Homo sapiens

20 <400> 313
 Phe Thr Phe Ser Ser Ser Gly Met His
 1 5

25 <210> 314
 <211> 9
 <212> PRT
 <213> Homo sapiens

30 <400> 314
 Phe Thr Phe Ser Ser Tyr Gly Met His
 1 5

35 <210> 315
 <211> 9
 <212> PRT
 <213> Homo sapiens

40 <400> 315
 Phe Thr Phe Ser Ser His Gly Met His
 1 5

45 <210> 316
 <211> 9
 <212> PRT
 <213> Homo sapiens

50 <400> 316
 Phe Thr Phe Ser Ser Arg Gly Met His
 1 5

55 <210> 317
 <211> 9
 <212> PRT
 <213> Homo sapiens

60 <400> 317
 Phe Thr Phe Ser Ser Asn Gly Met His
 1 5

65 <210> 318
 <211> 9
 <212> PRT
 <213> Homo sapiens

 <400> 318

Phe Thr Phe Ser Ser Thr Gly Met His
1 5

5 <210> 319
<211> 9
<212> PRT
<213> Homo sapiens

10 <400> 319
Phe Thr Phe Ser Ser Ala Gly Met His
1 5

15 <210> 320
<211> 9
<212> PRT
<213> Homo sapiens

20 <400> 320
Phe Thr Phe Ser Ser Val Gly Met His
1 5

25 <210> 321
<211> 9
<212> PRT
<213> Homo sapiens

30 <400> 321
Phe Thr Phe Ser Ser Leu Gly Met His
1 5

35 <210> 322
<211> 9
<212> PRT
<213> Homo sapiens

40 <400> 322
Phe Thr Phe Ser Ser Ile Gly Met His
1 5

45 <210> 323
<211> 9
<212> PRT
<213> Homo sapiens

50 <400> 323
Phe Thr Phe Ser Ser Tyr Asp Met His
1 5

55 <210> 324
<211> 9
<212> PRT
<213> Homo sapiens

60 <400> 324
Phe Thr Phe Ser Ser Tyr Glu Met His
1 5

65 <210> 325
<211> 9
<212> PRT
<213> Homo sapiens

<400> 325
Phe Thr Phe Ser Ser Tyr Cys Met His
1 5
5

<210> 326
<211> 9
<212> PRT
10 <213> Homo sapiens

<400> 326
Phe Thr Phe Ser Ser Tyr Ser Met His
1 5
15

<210> 327
<211> 9
<212> PRT
20 <213> Homo sapiens

<400> 327
Phe Thr Phe Ser Ser Tyr Tyr Met His
1 5
25

<210> 328
<211> 9
<212> PRT
30 <213> Homo sapiens

<400> 328
Phe Thr Phe Ser Ser Tyr Asn Met His
1 5
35

<210> 329
<211> 9
<212> PRT
40 <213> Homo sapiens

<400> 329
Phe Thr Phe Ser Ser Tyr Gly Met His
1 5
45

<210> 330
<211> 9
<212> PRT
50 <213> Homo sapiens

<400> 330
Phe Thr Phe Ser Ser Tyr Ala Met His
1 5
55

<210> 331
<211> 9
<212> PRT
60 <213> Homo sapiens

<400> 331
Phe Thr Phe Ser Ser Tyr Val Met His
1 5
65

<210> 332
<211> 9

<212> PRT
<213> Homo sapiens

5 <400> 332
Phe Thr Phe Ser Ser Tyr Met Met His
1 5

10 <210> 333
<211> 9
<212> PRT
<213> Homo sapiens

15 <400> 333
Phe Thr Phe Ser Ser Tyr Ile Met His
1 5

20 <210> 334
<211> 9
<212> PRT
<213> Homo sapiens

25 <400> 334
Phe Thr Phe Ser Ser Tyr Pro Met His
1 5

30 <210> 335
<211> 17
<212> PRT
<213> Homo sapiens

35 <400> 335
Glu Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

40

<210> 336
<211> 17
<212> PRT
45 <213> Homo sapiens

<400> 336
Cys Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

50 Gly

55 <210> 337
<211> 17
<212> PRT
<213> Homo sapiens

60 <400> 337
Tyr Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

65 <210> 338

<211> 17
<212> PRT
<213> Homo sapiens

5 <400> 338
His Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly
10

<210> 339
<211> 17
15 <212> PRT
<213> Homo sapiens

<400> 339
Lys Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
20 1 5 10 15

Gly

25 <210> 340
<211> 17
<212> PRT
<213> Homo sapiens

30 <400> 340
Asn Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly
35

40 <210> 341
<211> 17
<212> PRT
<213> Homo sapiens

45 <400> 341
Gln Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

50 <210> 342
<211> 17
<212> PRT
55 <213> Homo sapiens

<400> 342
Thr Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

60 <210> 343
<211> 17
<212> PRT
<213> Homo sapiens

1 5 10 15

Gly

5

<210> 349
<211> 17
<212> PRT

10 <213> Homo sapiens

<400> 349
Phe Ile Lys Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

15 Gly

20 <210> 350
<211> 17
<212> PRT
<213> Homo sapiens

25 <400> 350
Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

30

<210> 351
<211> 17
35 <212> PRT
<213> Homo sapiens

<400> 351
Phe Ile Gln Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
40 1 5 10 15

Gly

45

<210> 352
<211> 17
<212> PRT
<213> Homo sapiens

50 <400> 352
Phe Ile Thr Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

55 Gly

60 <210> 353
<211> 17
<212> PRT
<213> Homo sapiens

65 <400> 353
Phe Ile Gly Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

5 <210> 354
 <211> 17
 <212> PRT
 <213> Homo sapiens

10 <400> 354
 Phe Ile Ala Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

15

20 <210> 355
 <211> 17
 <212> PRT
 <213> Homo sapiens

25 <400> 355
 Phe Ile Val Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

30 <210> 356
 <211> 17
 <212> PRT
 <213> Homo sapiens

35 <400> 356
 Phe Ile Leu Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

40

45 <210> 357
 <211> 17
 <212> PRT
 <213> Homo sapiens

50 <400> 357
 Phe Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

55

60 <210> 358
 <211> 17
 <212> PRT
 <213> Homo sapiens

65 <400> 358
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

5 <210> 359
<211> 17
<212> PRT
<213> Homo sapiens
<400> 359
Phe Ile Arg Tyr Glu Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
10 Gly

15 <210> 360
<211> 17
<212> PRT
<213> Homo sapiens
<400> 360
Phe Ile Arg Tyr Ser Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
20 Gly

25

30 <210> 361
<211> 17
<212> PRT
<213> Homo sapiens
<400> 361
Phe Ile Arg Tyr Tyr Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
35 Gly

40 <210> 362
<211> 17
<212> PRT
<213> Homo sapiens
<400> 362
Phe Ile Arg Tyr Lys Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
45 Gly

50

55 <210> 363
<211> 17
<212> PRT
<213> Homo sapiens
<400> 363
Phe Ile Arg Tyr Arg Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
60 Gly

65 <210> 364
<211> 17
<212> PRT

<213> Homo sapiens

<400> 364
Phe Ile Arg Tyr Asn Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
5 1 5 10 15

Gly

10

<210> 365
<211> 17
<212> PRT
<213> Homo sapiens

15

<400> 365
Phe Ile Arg Tyr Gln Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

25

<210> 366
<211> 17
<212> PRT
<213> Homo sapiens

30

<400> 366
Phe Ile Arg Tyr Thr Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

35

<210> 367
<211> 17
<212> PRT
<213> Homo sapiens

40

<400> 367
Phe Ile Arg Tyr Ala Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

45

50

<210> 368
<211> 17
<212> PRT
<213> Homo sapiens

55

<400> 368
Phe Ile Arg Tyr Val Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

60

<210> 369
<211> 17
<212> PRT
<213> Homo sapiens

<400> 369

Phe Ile Arg Tyr Leu Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

5 Gly

10 <210> 370
<211> 17
<212> PRT
<213> Homo sapiens

15 <400> 370
Phe Ile Arg Tyr Ile Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

20 <210> 371
<211> 17
<212> PRT
<213> Homo sapiens

25 <400> 371
Phe Ile Arg Tyr Phe Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

30 Gly

35 <210> 372
<211> 17
<212> PRT
<213> Homo sapiens

40 <400> 372
Phe Ile Arg Tyr Asp Asp Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

45 <210> 373
<211> 17
<212> PRT
<213> Homo sapiens

50 <400> 373
Phe Ile Arg Tyr Asp Glu Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

55 Gly

60 <210> 374
<211> 17
<212> PRT
<213> Homo sapiens

65 <400> 374
Phe Ile Arg Tyr Asp Ser Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

5 <210> 375
<211> 17
<212> PRT
<213> Homo sapiens
10 <400> 375
Phe Ile Arg Tyr Asp Tyr Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

15 Gly

20 <210> 376
<211> 17
<212> PRT
<213> Homo sapiens
25 <400> 376
Phe Ile Arg Tyr Asp Lys Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

30 <210> 377
<211> 17
<212> PRT
<213> Homo sapiens
35 <400> 377
Phe Ile Arg Tyr Asp Arg Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

40 Gly

45 <210> 378
<211> 17
<212> PRT
<213> Homo sapiens
50 <400> 378
Phe Ile Arg Tyr Asp Asn Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

55
<210> 379
<211> 17
<212> PRT
60 <213> Homo sapiens
<400> 379
Phe Ile Arg Tyr Asp Gln Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
65 Gly

5 <210> 380
 <211> 17
 <212> PRT
 <213> Homo sapiens
 10 <400> 380
 Phe Ile Arg Tyr Asp Thr Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 15 <210> 381
 <211> 17
 <212> PRT
 <213> Homo sapiens
 20 <400> 381
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 25
 30 <210> 382
 <211> 17
 <212> PRT
 <213> Homo sapiens
 35 <400> 382
 Phe Ile Arg Tyr Asp Val Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 40
 45 <210> 383
 <211> 17
 <212> PRT
 <213> Homo sapiens
 <400> 383
 Phe Ile Arg Tyr Asp Phe Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 55 <210> 384
 <211> 17
 <212> PRT
 <213> Homo sapiens
 <400> 384
 60 Phe Ile Arg Tyr Asp Gly Ser Ser Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 65
 <210> 385
 <211> 17

<212> PRT
<213> Homo sapiens

5 <400> 385
1 Phe Ile Arg Tyr Asp Gly Ser Tyr Lys Tyr Tyr Ala Asp Ser Val Lys
5 10 15

Gly
10

<210> 386
<211> 17
<212> PRT
15 <213> Homo sapiens

<400> 386
1 Phe Ile Arg Tyr Asp Gly Ser His Lys Tyr Tyr Ala Asp Ser Val Lys
5 10 15

20 Gly

25 <210> 387
<211> 17
<212> PRT
<213> Homo sapiens

30 <400> 387
1 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
5 10 15

35 Gly

40 <210> 388
<211> 17
<212> PRT
<213> Homo sapiens

<400> 388
1 Phe Ile Arg Tyr Asp Gly Ser Thr Lys Tyr Tyr Ala Asp Ser Val Lys
5 10 15

45 Gly

50 <210> 389
<211> 17
<212> PRT
<213> Homo sapiens

55 <400> 389
1 Phe Ile Arg Tyr Asp Gly Ser Gly Lys Tyr Tyr Ala Asp Ser Val Lys
5 10 15

60 Gly

65 <210> 390
<211> 17
<212> PRT
<213> Homo sapiens

<400> 390
 Phe Ile Arg Tyr Asp Gly Ser Met Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

5 Gly

10 <210> 391
 <211> 17
 <212> PRT
 <213> Homo sapiens

15 <400> 391
 Phe Ile Arg Tyr Asp Gly Ser Leu Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

20

<210> 392
 <211> 17
 <212> PRT
 25 <213> Homo sapiens

<400> 392
 Phe Ile Arg Tyr Asp Gly Ser Ile Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

30 Gly

35 <210> 393
 <211> 17
 <212> PRT
 <213> Homo sapiens

40 <400> 393
 Phe Ile Arg Tyr Asp Gly Ser Pro Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

45

50 <210> 394
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 394
 Phe Ile Arg Tyr Asp Gly Ser Phe Lys Tyr Tyr Ala Asp Ser Val Lys
 55 1 5 10 15

Gly

60

<210> 395
 <211> 17
 <212> PRT
 <213> Homo sapiens

65 <400> 395
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Glu Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

5

<210> 396

<211> 17

<212> PRT

<213> Homo sapiens

10

<400> 396

Phe Ile Arg Tyr Asp Gly Ser Asn Lys Ser Tyr Ala Asp Ser Val Lys
1 5 10 15

15 Gly

20 <210> 397

<211> 17

<212> PRT

<213> Homo sapiens

25 <400> 397

Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

30

<210> 398

<211> 17

<212> PRT

35 <213> Homo sapiens

<400> 398

Phe Ile Arg Tyr Asp Gly Ser Asn Lys Asn Tyr Ala Asp Ser Val Lys
1 5 10 15

40

Gly

45 <210> 399

<211> 17

<212> PRT

<213> Homo sapiens

50 <400> 399

Phe Ile Arg Tyr Asp Gly Ser Asn Lys Val Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

55

<210> 400

<211> 17

60 <212> PRT

<213> Homo sapiens

<400> 400

Phe Ile Arg Tyr Asp Gly Ser Asn Lys Leu Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

5 <210> 401
 <211> 17
 <212> PRT
 <213> Homo sapiens

10 <400> 401
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Ile Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

15 <210> 402
 <211> 17
 <212> PRT
 <213> Homo sapiens

20 <400> 402
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Pro Tyr Ala Asp Ser Val Lys
 1 5 10 15

25 Gly

30 <210> 403
 <211> 17
 <212> PRT
 <213> Homo sapiens

35 <400> 403
 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Phe Tyr Ala Asp Ser Val Lys
 1 5 10 15

 Gly

40

45 <210> 404
 <211> 6
 <212> PRT
 <213> Homo sapiens

50 <400> 404
 Glu Gly Ser His Asp Asn
 1 5

55 <210> 405
 <211> 6
 <212> PRT
 <213> Homo sapiens

60 <400> 405
 Ser Gly Ser His Asp Asn
 1 5

65 <210> 406
 <211> 6
 <212> PRT
 <213> Homo sapiens

 <400> 406
 His Gly Ser His Asp Asn

5 <210> 407
<211> 6
<212> PRT
<213> Homo sapiens

10 <400> 407
Lys Gly Ser His Asp Asn
1 5

15 <210> 408
<211> 6
<212> PRT
<213> Homo sapiens

20 <400> 408
Gln Gly Ser His Asp Asn
1 5

25 <210> 409
<211> 6
<212> PRT
<213> Homo sapiens

30 <400> 409
Thr Gly Ser His Asp Asn
1 5

35 <210> 410
<211> 6
<212> PRT
<213> Homo sapiens

40 <400> 410
Ala Gly Ser His Asp Asn
1 5

45 <210> 411
<211> 6
<212> PRT
<213> Homo sapiens

50 <400> 411
Leu Gly Ser His Asp Asn
1 5

55 <210> 412
<211> 6
<212> PRT
<213> Homo sapiens

60 <400> 412
Pro Gly Ser His Asp Asn
1 5

65 <210> 413
<211> 6
<212> PRT
<213> Homo sapiens

<400> 413
Phe Gly Ser His Asp Asn
1 5

5
<210> 414
<211> 6
<212> PRT
<213> Homo sapiens

10
<400> 414
His Asp Ser His Asp Asn
1 5

15
<210> 415
<211> 6
<212> PRT
<213> Homo sapiens

20
<400> 415
His Cys Ser His Asp Asn
1 5

25
<210> 416
<211> 6
<212> PRT
<213> Homo sapiens

30
<400> 416
His His Ser His Asp Asn
1 5

35
<210> 417
<211> 6
<212> PRT
<213> Homo sapiens

40
<400> 417
His Arg Ser His Asp Asn
1 5

45
<210> 418
<211> 6
<212> PRT
<213> Homo sapiens

50
<400> 418
His Thr Ser His Asp Asn
1 5

55
<210> 419
<211> 6
<212> PRT
<213> Homo sapiens

60
<400> 419
His Gly Ser His Asp Asn
1 5

65
<210> 420
<211> 6
<212> PRT

<213> Homo sapiens

<400> 420
His Val Ser His Asp Asn
5 1 5

<210> 421
<211> 6
10 <212> PRT
<213> Homo sapiens

<400> 421
His Met Ser His Asp Asn
15 1 5

<210> 422
<211> 6
20 <212> PRT
<213> Homo sapiens

<400> 422
His Leu Ser His Asp Asn
25 1 5

<210> 423
<211> 6
30 <212> PRT
<213> Homo sapiens

<400> 423
His Ile Ser His Asp Asn
35 1 5

<210> 424
<211> 6
40 <212> PRT
<213> Homo sapiens

<400> 424
His Pro Ser His Asp Asn
45 1 5

<210> 425
<211> 6
50 <212> PRT
<213> Homo sapiens

<400> 425
His Trp Ser His Asp Asn
55 1 5

<210> 426
<211> 6
60 <212> PRT
<213> Homo sapiens

<400> 426
His Gly Asp His Asp Asn
65 1 5

<210> 427

<211> 6
<212> PRT
<213> Homo sapiens

5 <400> 427
His Gly Ser His Asp Asn
1 5

10 <210> 428
<211> 6
<212> PRT
<213> Homo sapiens

15 <400> 428
His Gly Tyr His Asp Asn
1 5

20 <210> 429
<211> 6
<212> PRT
<213> Homo sapiens

25 <400> 429
His Gly His His Asp Asn
1 5

30 <210> 430
<211> 6
<212> PRT
<213> Homo sapiens

35 <400> 430
His Gly Arg His Asp Asn
1 5

40 <210> 431
<211> 6
<212> PRT
<213> Homo sapiens

45 <400> 431
His Gly Asn His Asp Asn
1 5

50 <210> 432
<211> 6
<212> PRT
<213> Homo sapiens

55 <400> 432
His Gly Thr His Asp Asn
1 5

60 <210> 433
<211> 6
<212> PRT
<213> Homo sapiens

65 <400> 433
His Gly Gly His Asp Asn
1 5

<210> 434
<211> 6
<212> PRT
5 <213> Homo sapiens

<400> 434
His Gly Ala His Asp Asn
1 5
10

<210> 435
<211> 6
<212> PRT
15 <213> Homo sapiens

<400> 435
His Gly Ile His Asp Asn
1 5
20

<210> 436
<211> 6
<212> PRT
25 <213> Homo sapiens

<400> 436
His Gly Pro His Asp Asn
1 5
30

<210> 437
<211> 6
<212> PRT
35 <213> Homo sapiens

<400> 437
His Gly Trp His Asp Asn
1 5
40

<210> 438
<211> 6
<212> PRT
45 <213> Homo sapiens

<400> 438
His Gly Phe His Asp Asn
1 5
50

<210> 439
<211> 6
<212> PRT
55 <213> Homo sapiens

<400> 439
His Gly Ser His Asp Asn
1 5
60

<210> 440
<211> 6
<212> PRT
65 <213> Homo sapiens

<400> 440
His Gly Ser Arg Asp Asn

1 5

5 <210> 441
<211> 6
<212> PRT
<213> Homo sapiens

10 <400> 441
His Gly Ser Thr Asp Asn
1 5

15 <210> 442
<211> 6
<212> PRT
<213> Homo sapiens

20 <400> 442
His Gly Ser Ala Asp Asn
1 5

25 <210> 443
<211> 6
<212> PRT
<213> Homo sapiens

30 <400> 443
His Gly Ser Val Asp Asn
1 5

35 <210> 444
<211> 6
<212> PRT
<213> Homo sapiens

40 <400> 444
His Gly Ser Leu Asp Asn
1 5

45 <210> 445
<211> 6
<212> PRT
<213> Homo sapiens

50 <400> 445
His Gly Ser Ile Asp Asn
1 5

55 <210> 446
<211> 6
<212> PRT
<213> Homo sapiens

60 <400> 446
His Gly Ser Phe Asp Asn
1 5

65 <210> 447
<211> 6
<212> PRT
<213> Homo sapiens

<400> 447
His Gly Ser His Asp Asn
1 5

5
<210> 448
<211> 6
<212> PRT
<213> Homo sapiens

10
<400> 448
His Gly Ser His Ser Asn
1 5

15
<210> 449
<211> 6
<212> PRT
<213> Homo sapiens

20
<400> 449
His Gly Ser His Tyr Asn
1 5

25
<210> 450
<211> 6
<212> PRT
<213> Homo sapiens

30
<400> 450
His Gly Ser His His Asn
1 5

35
<210> 451
<211> 6
<212> PRT
<213> Homo sapiens

40
<400> 451
His Gly Ser His Arg Asn
1 5

45
<210> 452
<211> 6
<212> PRT
<213> Homo sapiens

50
<400> 452
His Gly Ser His Asn Asn
1 5

55
<210> 453
<211> 6
<212> PRT
<213> Homo sapiens

60
<400> 453
His Gly Ser His Gly Asn
1 5

65
<210> 454
<211> 6
<212> PRT

<213> Homo sapiens

<400> 454
His Gly Ser His Ala Asn
5 1 5

<210> 455
<211> 6
10 <212> PRT
<213> Homo sapiens

<400> 455
His Gly Ser His Val Asn
15 1 5

<210> 456
<211> 6
20 <212> PRT
<213> Homo sapiens

<400> 456
His Gly Ser His Ile Asn
25 1 5

<210> 457
<211> 6
30 <212> PRT
<213> Homo sapiens

<400> 457
His Gly Ser His Asp Ser
35 1 5

<210> 458
<211> 6
40 <212> PRT
<213> Homo sapiens

<400> 458
His Gly Ser His Asp His
45 1 5

<210> 459
<211> 6
50 <212> PRT
<213> Homo sapiens

<400> 459
His Gly Ser His Asp Lys
55 1 5

<210> 460
<211> 6
60 <212> PRT
<213> Homo sapiens

<400> 460
His Gly Ser His Asp Arg
65 1 5

<210> 461

<211> 6
<212> PRT
<213> Homo sapiens

5 <400> 461
His Gly Ser His Asp Asn
1 5

10 <210> 462
<211> 6
<212> PRT
<213> Homo sapiens

15 <400> 462
His Gly Ser His Asp Thr
1 5

20 <210> 463
<211> 6
<212> PRT
<213> Homo sapiens

25 <400> 463
His Gly Ser His Asp Gly
1 5

30 <210> 464
<211> 6
<212> PRT
<213> Homo sapiens

35 <400> 464
His Gly Ser His Asp Ala
1 5

40 <210> 465
<211> 6
<212> PRT
<213> Homo sapiens

45 <400> 465
His Gly Ser His Asp Leu
1 5

50 <210> 466
<211> 6
<212> PRT
<213> Homo sapiens

55 <400> 466
His Gly Ser His Asp Ile
1 5

60 <210> 467
<211> 6
<212> PRT
<213> Homo sapiens

65 <400> 467
His Gly Ser His Asp Pro
1 5

5 <210> 468
 <211> 6
 <212> PRT
 5 <213> Homo sapiens

 10 <400> 468
 His Gly Ser His Asp Trp
 1 5
 10

 15 <210> 469
 <211> 6
 <212> PRT
 15 <213> Homo sapiens

 20 <400> 469
 His Gly Ser His Asp Phe
 1 5
 20

 25 <210> 470
 <211> 13
 <212> PRT
 25 <213> Homo sapiens

 30 <400> 470
 Ser Gly Gly Arg Ser Asn Ile Gly Asp Asn Thr Val Lys
 1 5 10
 30

 35 <210> 471
 <211> 13
 <212> PRT
 35 <213> Homo sapiens

 40 <400> 471
 Ser Gly Gly Arg Ser Asn Ile Gly Cys Asn Thr Val Lys
 1 5 10
 40

 45 <210> 472
 <211> 13
 <212> PRT
 45 <213> Homo sapiens

 50 <400> 472
 Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Thr Val Lys
 1 5 10
 50

 55 <210> 473
 <211> 13
 <212> PRT
 55 <213> Homo sapiens

 60 <400> 473
 Ser Gly Gly Arg Ser Asn Ile Gly Tyr Asn Thr Val Lys
 1 5 10
 60

 65 <210> 474
 <211> 13
 <212> PRT
 65 <213> Homo sapiens

 70 <400> 474
 Ser Gly Gly Arg Ser Asn Ile Gly Lys Asn Thr Val Lys

	1	5	10
5	<210> 475		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
10	<400> 475		
	Ser Gly Gly Arg Ser Asn Ile Gly Arg Asn Thr Val Lys		
	1	5	10
15	<210> 476		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
20	<400> 476		
	Ser Gly Gly Arg Ser Asn Ile Gly Asn Asn Thr Val Lys		
	1	5	10
25	<210> 477		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
30	<400> 477		
	Ser Gly Gly Arg Ser Asn Ile Gly Thr Asn Thr Val Lys		
	1	5	10
35	<210> 478		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
40	<400> 478		
	Ser Gly Gly Arg Ser Asn Ile Gly Pro Asn Thr Val Lys		
	1	5	10
45	<210> 479		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
50	<400> 479		
	Ser Gly Gly Arg Ser Asn Ile Gly Ser Asp Thr Val Lys		
	1	5	10
55	<210> 480		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		
60	<400> 480		
	Ser Gly Gly Arg Ser Asn Ile Gly Ser Glu Thr Val Lys		
	1	5	10
65	<210> 481		
	<211> 13		
	<212> PRT		
	<213> Homo sapiens		

<400> 481
Ser Gly Gly Arg Ser Asn Ile Gly Ser Ser Thr Val Lys
1 5 10

5
<210> 482
<211> 13
<212> PRT
<213> Homo sapiens

10
<400> 482
Ser Gly Gly Arg Ser Asn Ile Gly Ser Tyr Thr Val Lys
1 5 10

15
<210> 483
<211> 13
<212> PRT
<213> Homo sapiens

20
<400> 483
Ser Gly Gly Arg Ser Asn Ile Gly Ser His Thr Val Lys
1 5 10

25
<210> 484
<211> 13
<212> PRT
<213> Homo sapiens

30
<400> 484
Ser Gly Gly Arg Ser Asn Ile Gly Ser Lys Thr Val Lys
1 5 10

35
<210> 485
<211> 13
<212> PRT
<213> Homo sapiens

40
<400> 485
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Thr Val Lys
1 5 10

45
<210> 486
<211> 13
<212> PRT
<213> Homo sapiens

50
<400> 486
Ser Gly Gly Arg Ser Asn Ile Gly Ser Gln Thr Val Lys
1 5 10

55
<210> 487
<211> 13
<212> PRT
<213> Homo sapiens

60
<400> 487
Ser Gly Gly Arg Ser Asn Ile Gly Ser Thr Thr Val Lys
1 5 10

65
<210> 488
<211> 13
<212> PRT

<213> Homo sapiens

<400> 488
5 Ser Gly Gly Arg Ser Asn Ile Gly Ser Gly Thr Val Lys
1 5 10

10 <210> 489
<211> 13
<212> PRT
<213> Homo sapiens

<400> 489
15 Ser Gly Gly Arg Ser Asn Ile Gly Ser Met Thr Val Lys
1 5 10

20 <210> 490
<211> 13
<212> PRT
<213> Homo sapiens

<400> 490
25 Ser Gly Gly Arg Ser Asn Ile Gly Ser Ile Thr Val Lys
1 5 10

30 <210> 491
<211> 13
<212> PRT
<213> Homo sapiens

<400> 491
35 Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Asp Val Lys
1 5 10

40 <210> 492
<211> 13
<212> PRT
<213> Homo sapiens

<400> 492
45 Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Cys Val Lys
1 5 10

50 <210> 493
<211> 13
<212> PRT
<213> Homo sapiens

<400> 493
55 Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Ser Val Lys
1 5 10

60 <210> 494
<211> 13
<212> PRT
<213> Homo sapiens

<400> 494
65 Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Tyr Val Lys
1 5 10

<210> 495

<211> 13
<212> PRT
<213> Homo sapiens

5 <400> 495
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn His Val Lys
1 5 10

10 <210> 496
<211> 13
<212> PRT
<213> Homo sapiens

15 <400> 496
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Lys Val Lys
1 5 10

20 <210> 497
<211> 13
<212> PRT
<213> Homo sapiens

25 <400> 497
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Arg Val Lys
1 5 10

30 <210> 498
<211> 13
<212> PRT
<213> Homo sapiens

35 <400> 498
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Asn Val Lys
1 5 10

40 <210> 499
<211> 13
<212> PRT
<213> Homo sapiens

45 <400> 499
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Gln Val Lys
1 5 10

50 <210> 500
<211> 13
<212> PRT
<213> Homo sapiens

55 <400> 500
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Thr Val Lys
1 5 10

60 <210> 501
<211> 13
<212> PRT
<213> Homo sapiens

65 <400> 501
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Ala Val Lys
1 5 10

<210> 502
<211> 13
<212> PRT
5 <213> Homo sapiens

<400> 502
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Val Val Lys
1 5 10
10

<210> 503
<211> 13
<212> PRT
15 <213> Homo sapiens

<400> 503
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Leu Val Lys
1 5 10
20

<210> 504
<211> 13
<212> PRT
25 <213> Homo sapiens

<400> 504
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Ile Val Lys
1 5 10
30

<210> 505
<211> 13
<212> PRT
35 <213> Homo sapiens

<400> 505
Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn Pro Val Lys
1 5 10
40

<210> 506
<211> 7
<212> PRT
45 <213> Homo sapiens

<400> 506
Asp Asn Asp Gln Arg Pro Ser
1 5
50

<210> 507
<211> 7
<212> PRT
55 <213> Homo sapiens

<400> 507
Glu Asn Asp Gln Arg Pro Ser
1 5
60

<210> 508
<211> 7
<212> PRT
65 <213> Homo sapiens

<400> 508
Cys Asn Asp Gln Arg Pro Ser

1 5

5 <210> 509
<211> 7
<212> PRT
<213> Homo sapiens

10 <400> 509
Ser Asn Asp Gln Arg Pro Ser
1 5

15 <210> 510
<211> 7
<212> PRT
<213> Homo sapiens

20 <400> 510
Tyr Asn Asp Gln Arg Pro Ser
1 5

25 <210> 511
<211> 7
<212> PRT
<213> Homo sapiens

30 <400> 511
His Asn Asp Gln Arg Pro Ser
1 5

35 <210> 512
<211> 7
<212> PRT
<213> Homo sapiens

40 <400> 512
Lys Asn Asp Gln Arg Pro Ser
1 5

45 <210> 513
<211> 7
<212> PRT
<213> Homo sapiens

50 <400> 513
Arg Asn Asp Gln Arg Pro Ser
1 5

55 <210> 514
<211> 7
<212> PRT
<213> Homo sapiens

60 <400> 514
Asn Asn Asp Gln Arg Pro Ser
1 5

65 <210> 515
<211> 7
<212> PRT
<213> Homo sapiens

<400> 515
Gln Asn Asp Gln Arg Pro Ser
1 5

5
<210> 516
<211> 7
<212> PRT
<213> Homo sapiens

10
<400> 516
Thr Asn Asp Gln Arg Pro Ser
1 5

15
<210> 517
<211> 7
<212> PRT
<213> Homo sapiens

20
<400> 517
Gly Asn Asp Gln Arg Pro Ser
1 5

25
<210> 518
<211> 7
<212> PRT
<213> Homo sapiens

30
<400> 518
Ala Asn Asp Gln Arg Pro Ser
1 5

35
<210> 519
<211> 7
<212> PRT
<213> Homo sapiens

40
<400> 519
Val Asn Asp Gln Arg Pro Ser
1 5

45
<210> 520
<211> 7
<212> PRT
<213> Homo sapiens

50
<400> 520
Met Asn Asp Gln Arg Pro Ser
1 5

55
<210> 521
<211> 7
<212> PRT
<213> Homo sapiens

60
<400> 521
Leu Asn Asp Gln Arg Pro Ser
1 5

65
<210> 522
<211> 7
<212> PRT

<213> Homo sapiens

<400> 522
5 Ile Asn Asp Gln Arg Pro Ser
1 5

<210> 523
<211> 7
10 <212> PRT
<213> Homo sapiens

<400> 523
15 Pro Asn Asp Gln Arg Pro Ser
1 5

<210> 524
<211> 7
20 <212> PRT
<213> Homo sapiens

<400> 524
25 Trp Asn Asp Gln Arg Pro Ser
1 5

<210> 525
<211> 7
30 <212> PRT
<213> Homo sapiens

<400> 525
35 Phe Asn Asp Gln Arg Pro Ser
1 5

<210> 526
<211> 7
40 <212> PRT
<213> Homo sapiens

<400> 526
45 Gly Asn Asp Ser Arg Pro Ser
1 5

<210> 527
<211> 7
50 <212> PRT
<213> Homo sapiens

<400> 527
55 Gly Asn Asp Tyr Arg Pro Ser
1 5

<210> 528
<211> 7
60 <212> PRT
<213> Homo sapiens

<400> 528
65 Gly Asn Asp Arg Arg Pro Ser
1 5

<210> 529

<211> 7
<212> PRT
<213> Homo sapiens

5 <400> 529
Gly Asn Asp Gln Arg Pro Ser
1 5

10 <210> 530
<211> 7
<212> PRT
<213> Homo sapiens

15 <400> 530
Gly Asn Asp Thr Arg Pro Ser
1 5

20 <210> 531
<211> 7
<212> PRT
<213> Homo sapiens

25 <400> 531
Gly Asn Asp Ala Arg Pro Ser
1 5

30 <210> 532
<211> 7
<212> PRT
<213> Homo sapiens

35 <400> 532
Gly Asn Asp Ile Arg Pro Ser
1 5

40 <210> 533
<211> 7
<212> PRT
<213> Homo sapiens

45 <400> 533
Gly Asn Asp Pro Arg Pro Ser
1 5

50 <210> 534
<211> 12
<212> PRT
<213> Homo sapiens

55 <400> 534
Gln Ser Tyr Asp Arg Gly Thr His Pro Ala Leu Leu
1 5 10

60 <210> 535
<211> 12
<212> PRT
<213> Homo sapiens

65 <400> 535
Gln Ser Tyr Cys Arg Gly Thr His Pro Ala Leu Leu
1 5 10

<210> 536
<211> 12
<212> PRT
5 <213> Homo sapiens

<400> 536
Gln Ser Tyr Ser Arg Gly Thr His Pro Ala Leu Leu
1 5 10
10

<210> 537
<211> 12
<212> PRT
15 <213> Homo sapiens

<400> 537
Gln Ser Tyr Tyr Arg Gly Thr His Pro Ala Leu Leu
1 5 10
20

<210> 538
<211> 12
<212> PRT
25 <213> Homo sapiens

<400> 538
Gln Ser Tyr Asn Arg Gly Thr His Pro Ala Leu Leu
1 5 10
30

<210> 539
<211> 12
<212> PRT
35 <213> Homo sapiens

<400> 539
Gln Ser Tyr Gln Arg Gly Thr His Pro Ala Leu Leu
1 5 10
40

<210> 540
<211> 12
<212> PRT
45 <213> Homo sapiens

<400> 540
Gln Ser Tyr Thr Arg Gly Thr His Pro Ala Leu Leu
1 5 10
50

<210> 541
<211> 12
<212> PRT
55 <213> Homo sapiens

<400> 541
Gln Ser Tyr Gly Arg Gly Thr His Pro Ala Leu Leu
1 5 10
60

<210> 542
<211> 12
<212> PRT
65 <213> Homo sapiens

<400> 542
Gln Ser Tyr Ala Arg Gly Thr His Pro Ala Leu Leu

	1	5	10
5	<210> 543		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
10	<400> 543		
	Gln Ser Tyr Leu Arg Gly Thr His Pro Ala Leu Leu		
	1	5	10
15	<210> 544		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
20	<400> 544		
	Gln Ser Tyr Ile Arg Gly Thr His Pro Ala Leu Leu		
	1	5	10
25	<210> 545		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
30	<400> 545		
	Gln Ser Tyr Trp Arg Gly Thr His Pro Ala Leu Leu		
	1	5	10
35	<210> 546		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
40	<400> 546		
	Gln Ser Tyr Phe Arg Gly Thr His Pro Ala Leu Leu		
	1	5	10
45	<210> 547		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
50	<400> 547		
	Gln Ser Tyr Asp Asp Gly Thr His Pro Ala Leu Leu		
	1	5	10
55	<210> 548		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		
60	<400> 548		
	Gln Ser Tyr Asp Cys Gly Thr His Pro Ala Leu Leu		
	1	5	10
65	<210> 549		
	<211> 12		
	<212> PRT		
	<213> Homo sapiens		

<400> 549
Gln Ser Tyr Asp Ser Gly Thr His Pro Ala Leu Leu
1 5 10

5
<210> 550
<211> 12
<212> PRT
<213> Homo sapiens

10
<400> 550
Gln Ser Tyr Asp Tyr Gly Thr His Pro Ala Leu Leu
1 5 10

15
<210> 551
<211> 12
<212> PRT
<213> Homo sapiens

20
<400> 551
Gln Ser Tyr Asp Arg Gly Thr His Pro Ala Leu Leu
1 5 10

25
<210> 552
<211> 12
<212> PRT
<213> Homo sapiens

30
<400> 552
Gln Ser Tyr Asp Asn Gly Thr His Pro Ala Leu Leu
1 5 10

35
<210> 553
<211> 12
<212> PRT
<213> Homo sapiens

40
<400> 553
Gln Ser Tyr Asp Gln Gly Thr His Pro Ala Leu Leu
1 5 10

45
<210> 554
<211> 12
<212> PRT
<213> Homo sapiens

50
<400> 554
Gln Ser Tyr Asp Thr Gly Thr His Pro Ala Leu Leu
1 5 10

55
<210> 555
<211> 12
<212> PRT
<213> Homo sapiens

60
<400> 555
Gln Ser Tyr Asp Gly Gly Thr His Pro Ala Leu Leu
1 5 10

65
<210> 556
<211> 12
<212> PRT

<213> Homo sapiens

<400> 556
5 Gln Ser Tyr Asp Ala Gly Thr His Pro Ala Leu Leu
1 5 10

10 <210> 557
<211> 12
<212> PRT
<213> Homo sapiens

<400> 557
15 Gln Ser Tyr Asp Val Gly Thr His Pro Ala Leu Leu
1 5 10

20 <210> 558
<211> 12
<212> PRT
<213> Homo sapiens

<400> 558
25 Gln Ser Tyr Asp Met Gly Thr His Pro Ala Leu Leu
1 5 10

30 <210> 559
<211> 12
<212> PRT
<213> Homo sapiens

<400> 559
35 Gln Ser Tyr Asp Leu Gly Thr His Pro Ala Leu Leu
1 5 10

40 <210> 560
<211> 12
<212> PRT
<213> Homo sapiens

<400> 560
45 Gln Ser Tyr Asp Ile Gly Thr His Pro Ala Leu Leu
1 5 10

50 <210> 561
<211> 12
<212> PRT
<213> Homo sapiens

<400> 561
55 Gln Ser Tyr Asp Pro Gly Thr His Pro Ala Leu Leu
1 5 10

60 <210> 562
<211> 12
<212> PRT
<213> Homo sapiens

<400> 562
65 Gln Ser Tyr Asp Trp Gly Thr His Pro Ala Leu Leu
1 5 10

<210> 563

<211> 12
<212> PRT
<213> Homo sapiens

5 <400> 563
Gln Ser Tyr Asp Arg Asp Thr His Pro Ala Leu Leu
1 5 10

10 <210> 564
<211> 12
<212> PRT
<213> Homo sapiens

15 <400> 564
Gln Ser Tyr Asp Arg Cys Thr His Pro Ala Leu Leu
1 5 10

20 <210> 565
<211> 12
<212> PRT
<213> Homo sapiens

25 <400> 565
Gln Ser Tyr Asp Arg Ser Thr His Pro Ala Leu Leu
1 5 10

30 <210> 566
<211> 12
<212> PRT
<213> Homo sapiens

35 <400> 566
Gln Ser Tyr Asp Arg Tyr Thr His Pro Ala Leu Leu
1 5 10

40 <210> 567
<211> 12
<212> PRT
<213> Homo sapiens

45 <400> 567
Gln Ser Tyr Asp Arg His Thr His Pro Ala Leu Leu
1 5 10

50 <210> 568
<211> 12
<212> PRT
<213> Homo sapiens

55 <400> 568
Gln Ser Tyr Asp Arg Arg Thr His Pro Ala Leu Leu
1 5 10

60 <210> 569
<211> 12
<212> PRT
<213> Homo sapiens

65 <400> 569
Gln Ser Tyr Asp Arg Asn Thr His Pro Ala Leu Leu
1 5 10

<210> 570
<211> 12
<212> PRT
5 <213> Homo sapiens

<400> 570
Gln Ser Tyr Asp Arg Gln Thr His Pro Ala Leu Leu
1 5 10
10

<210> 571
<211> 12
<212> PRT
15 <213> Homo sapiens

<400> 571
Gln Ser Tyr Asp Arg Thr Thr His Pro Ala Leu Leu
1 5 10
20

<210> 572
<211> 12
<212> PRT
25 <213> Homo sapiens

<400> 572
Gln Ser Tyr Asp Arg Gly Thr His Pro Ala Leu Leu
1 5 10
30

<210> 573
<211> 12
<212> PRT
35 <213> Homo sapiens

<400> 573
Gln Ser Tyr Asp Arg Ala Thr His Pro Ala Leu Leu
1 5 10
40

<210> 574
<211> 12
<212> PRT
45 <213> Homo sapiens

<400> 574
Gln Ser Tyr Asp Arg Val Thr His Pro Ala Leu Leu
1 5 10
50

<210> 575
<211> 12
<212> PRT
55 <213> Homo sapiens

<400> 575
Gln Ser Tyr Asp Arg Leu Thr His Pro Ala Leu Leu
1 5 10
60

<210> 576
<211> 12
<212> PRT
65 <213> Homo sapiens

<400> 576
Gln Ser Tyr Asp Arg Ile Thr His Pro Ala Leu Leu

	1	5	10		
5	<210> 577				
	<211> 12				
	<212> PRT				
	<213> Homo sapiens				
10	Gln Ser Tyr Asp Arg Pro Thr His Pro Ala Leu Leu				
	1	5	10		
15	<210> 578				
	<211> 12				
	<212> PRT				
	<213> Homo sapiens				
20	<400> 578				
	Gln Ser Tyr Asp Arg Trp Thr His Pro Ala Leu Leu				
	1	5	10		
25	<210> 579				
	<211> 12				
	<212> PRT				
	<213> Homo sapiens				
30	<400> 579				
	Gln Ser Tyr Asp Arg Phe Thr His Pro Ala Leu Leu				
	1	5	10		
35	<210> 580				
	<211> 48				
	<212> DNA				
	<213> synthetic construct				
40	<223> nucleotides at positions 16 to 34 can be substituted with any nucleotide such that the randomized nucleotides represent 12% of the sequence				
	<400> 580				
45	tgtcccttgg ccccagttagt catagctccc actggtcgta cagtaata				48
	<210> 581				
	<211> 35				
	<212> DNA				
	<213> synthetic construct				
50	<400> 581				
	gacacctcga tcagcggata acaatttcac acagg				35
55	<210> 582				
	<211> 15				
	<212> DNA				
	<213> synthetic construct				
60	<400> 582				
	tggggccaag ggaca				15
	<210> 583				
	<211> 45				
	<212> DNA				
65	<213> synthetic construct				
	<400> 583				
	attcgtccta taccgttcta ctttgcgtc tttccagacg ttagt				45

5 <210> 584
 <211> 18
 <212> DNA
 <213> synthetic construct
 10 <400> 584
 attcgtccta taccgttc 18

15 <210> 585
 <211> 66
 <212> DNA
 <213> synthetic construct
 <223> nucleotides from position 28 to 42 can be
 substituted with any nucleotide such that the
 randomized nucleotides represent 12% of the
 sequence

20 <400> 585
 ggtcccgatt ccgaagaccc tcgaacccct caggctgctg tcatatgact ggcagtaata 60
 gtcagc 66

25 <210> 586
 <211> 15
 <212> DNA
 <213> synthetic construct
 30 <400> 586
 tggggccaag ggaca 15

35 <210> 587
 <211> 24
 <212> DNA
 <213> synthetic construct
 40 <400> 587
 tgaagagacg gtgaccattg tccc 24

45 <210> 588
 <211> 16
 <212> DNA
 <213> synthetic construct
 50 <400> 588
 gacacctcga tcagcg 16

55 <210> 589
 <211> 48
 <212> DNA
 <213> synthetic construct
 60 <400> 589
 gagtcattct cgacttgcgg ccgcacctag gacggtcagc ttggccc 48

65 <210> 590
 <211> 12
 <212> PRT
 <213> Homo sapiens
 66 <400> 590
 Gln Ser Tyr Asp Arg Gly Phe Thr Gly Ser Met Val
 1 5 10

71 <210> 591
 <211> 12
 <212> PRT
 <213> Homo sapiens

5 <220>
 <223> Xaa is encoded by a randomized codon of sequence
 NNS with N being any nucleotide and S being either
 deoxycytosine or deoxyguanidine
 10 <400> 591
 Xaa Xaa Xaa Xaa Xaa Phe Thr Gly Ser Met Val
 1 5 10
 10
 15 <210> 592
 <211> 12
 <212> PRT
 <213> Homo sapiens
 20 <220>
 <223> Xaa is encoded by a randomized codon of sequence
 NNS with N being any nucleotide and S being either
 deoxycytosine or deoxyguanidine
 25 <400> 592
 Gln Ser Tyr Xaa Xaa Xaa Xaa Xaa Ser Met Val
 1 5 10
 25
 30 <210> 593
 <211> 12
 <212> PRT
 <213> Homo sapiens
 35 <220>
 <223> Xaa is encoded by a randomized codon of sequence
 NNS with N being any nucleotide and S being either
 deoxycytosine or deoxyguanidine
 40 <400> 593
 Gln Ser Tyr Asp Arg Gly Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10
 40
 45 <210> 594
 <211> 100
 <212> PRT
 <213> Homo sapiens
 50 <400> 594
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 50
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp His
 20 25 30
 55 Tyr Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 55
 Gly Arg Thr Arg Asn Lys Ala Asn Ser Tyr Thr Thr Glu Tyr Ala Ala
 50 55 60
 60 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Ser
 65 70 75 80
 65
 Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95
 65
 Tyr Cys Ala Arg
 100

<210> 595
 <211> 100
 <212> PRT
 5 <213> Homo sapiens

 <400> 595
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 10 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp His
 20 25 30

 Tyr Met Ser Trp Val Arg Gln Ala Gln Gly Lys Gly Leu Glu Leu Val
 15 35 40 45

 Gly Leu Ile Arg Asn Lys Ala Asn Ser Tyr Thr Thr Glu Tyr Ala Ala
 50 55 60

 20 Ser Val Lys Gly Arg Leu Thr Ile Ser Arg Glu Asp Ser Lys Asn Thr
 65 70 75 80

 Leu Tyr Leu Gln Met Ser Ser Leu Lys Thr Glu Asp Leu Ala Val Tyr
 85 90 95
 25 Tyr Cys Ala Arg
 100

 30 <210> 596
 <211> 100
 <212> PRT
 <213> Homo sapiens

 35 <400> 596
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

 40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp His
 20 25 30

 Tyr Met Ser Trp Val Arg Gln Ala Gln Gly Lys Gly Leu Glu Leu Val
 35 40 45

 45 Gly Leu Ile Arg Asn Lys Ala Asn Ser Tyr Thr Thr Glu Tyr Ala Ala
 50 55 60

 Ser Val Lys Gly Arg Leu Thr Ile Ser Arg Glu Asp Ser Lys Asn Thr
 65 70 75 80
 50 Met Tyr Leu Gln Met Ser Asn Leu Lys Thr Glu Asp Leu Ala Val Tyr
 85 90 95

 55 Tyr Cys Ala Arg
 100

 <210> 597
 <211> 100
 60 <212> PRT
 <213> Homo sapiens

 <400> 597
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 65 1 5 10 15

 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp His
 20 25 30

Tyr Met Ser Trp Val Arg Gln Ala Gln Gly Lys Gly Leu Glu Leu Val
 35 40 45

5 Gly Leu Ile Arg Asn Lys Ala Asn Ser Tyr Thr Thr Glu Tyr Ala Ala
 50 55 60

Ser Val Lys Gly Arg Leu Thr Ile Ser Arg Glu Asp Ser Lys Asn Thr
 65 70 75 80

10 Leu Tyr Leu Gln Met Ser Ser Leu Lys Thr Glu Asp Leu Ala Val Tyr
 85 90 95

15 Tyr Cys Ala Arg
 100

<210> 598
 <211> 98
 20 <212> PRT
 <213> Homo sapiens

<400> 598
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Arg
 25 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30

30 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Gly Ile Ser Trp Asn Ser Gly Ser Ile Gly Tyr Ala Asp Ser Val
 50 55 60

35 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

40 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr Tyr Cys
 85 90 95

Ala Lys

45 <210> 599
 <211> 98
 <212> PRT
 <213> Homo sapiens

50 <400> 599
 Glu Val Gln Leu Val Glu Ser Gly Gly Val Val Arg Pro Gly Gly
 1 5 10 15

55 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30

Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

60 Ser Gly Ile Asn Trp Asn Gly Gly Ser Thr Gly Tyr Ala Asp Ser Val
 50 55 60

65 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr His Cys
 85 90 95

Ala Arg

5 <210> 600
 <211> 98
 <212> PRT
 <213> Homo sapiens
 10 <400> 600
 Glu Val Gln Leu Val Glu Ser Gly Gly Val Val Val Gln Pro Gly Gly
 1 5 10 15
 15 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Thr Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 20 Ser Leu Ile Ser Trp Asp Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 25 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Thr Glu Asp Thr Ala Leu Tyr Tyr Cys
 85 90 95
 30 Ala Lys

```

35  <210> 601
    <211> 98
    <212> PRT
    <213> Homo sapiens

40  <400> 601
    Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
    1           5           10          15

    Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
    20          25          30

45  Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
    35          40          45

    Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
    50          55          60

    Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
    65          70          75          80

55  Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
    85          90          95

```

Ala Arg

60 <210> 602
<211> 98
<212> PRT
65 <213> Homo sapiens

<400> 602
Gln Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly

	1	5	10	15
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr			
	20	25		30
5	Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val			
	35	40	45	
	Ser Tyr Ile Ser Ser Ser Ser Tyr Thr Asn Tyr Ala Asp Ser Val			
10	50	55	60	
	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr			
	65	70	75	80
15	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys			
	85	90	95	
	Ala Arg			
20				
	<210> 603			
	<211> 100			
	<212> PRT			
25	<213> Homo sapiens			
	<400> 603			
	Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly			
	1	5	10	15
30	Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Ser			
	20	25	30	
	Ala Met His Trp Val Arg Gln Ala Ser Gly Lys Gly Leu Glu Trp Val			
35	35	40	45	
	Gly Arg Ile Arg Ser Lys Ala Asn Ser Tyr Ala Thr Ala Tyr Ala Ala			
	50	55	60	
40	Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr			
	65	70	75	80
	Ala Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr			
	85	90	95	
45	Tyr Cys Thr Arg			
	100			
50	<210> 604			
	<211> 100			
	<212> PRT			
	<213> Homo sapiens			
55	<400> 604			
	Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Lys Pro Gly Gly			
	1	5	10	15
60	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala			
	20	25	30	
	Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val			
	35	40	45	
65	Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala			
	50	55	60	
	Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr			

65	70	75	80
Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr			
85 90 95			
5	Tyr Cys Thr Thr		
	100		
10	<210> 605		
	<211> 100		
	<212> PRT		
	<213> Homo sapiens		
15	<400> 605		
	Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly		
	1	5	10
	15		
20	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala		
	20	25	30
	Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
	35	40	45
25	Gly Arg Ile Glu Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala		
	50	55	60
	Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr		
	65	70	75
30	80		
	Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr		
	85	90	95
35	Tyr Cys Thr Thr		
	100		
40	<210> 606		
	<211> 100		
	<212> PRT		
	<213> Homo sapiens		
	<400> 606		
45	Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly		
	1	5	10
	15		
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala		
	20	25	30
50			
	Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
	35	40	45
	Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala		
	50	55	60
55			
	Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr		
	65	70	75
	80		
60	Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr		
	85	90	95
	Tyr Cys Thr Thr		
	100		
65	<210> 607		
	<211> 100		
	<212> PRT		

<213> Homo sapiens

<400> 607

5 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

10 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asn Tyr Ala Ala
50 55 60

15 Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
65 70 75 80

20 Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
85 90 95

Tyr Cys Thr Thr
100

25

<210> 608

<211> 100

<212> PRT

<213> Homo sapiens

30

<400> 608

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

Trp Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

40 Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
50 55 60

45 Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
85 90 95

50 Tyr Cys Thr Thr
100

<210> 609

55 <211> 100

<212> PRT

<213> Homo sapiens

<400> 609

60 Glu Val Gln Leu Val Glu Ser Gly Gly Ala Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

65 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50 55 60

5 Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
 65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95

10 Tyr Cys Thr Thr
 100

15 <210> 610
 <211> 98
 <212> PRT
 <213> Homo sapiens

20 <400> 610
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Pro Ala Ser Gly Phe Thr Phe Ser Asn His
 20 25 30

25 Tyr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

30 Ser Tyr Ile Ser Gly Asp Ser Gly Tyr Thr Asn Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Asn Asn Ser Pro Tyr
 65 70 75 80

35 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Val Lys

40 <210> 611
 <211> 98
 <212> PRT
 <213> Homo sapiens

45 <400> 611
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn His
 20 25 30

Tyr Thr Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

55 Ser Tyr Ser Ser Gly Asn Ser Gly Tyr Thr Asn Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

65 Val Lys

<210> 612

<211> 98
<212> PRT
<213> Homo sapiens

5 <400> 612
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ser
10 20 25 30
Asp Met Asn Trp Val His Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
15 Ser Gly Val Ser Trp Asn Gly Ser Arg Thr His Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Ile Ile Ser Arg Asp Asn Ser Arg Asn Thr Leu Tyr
65 70 75 80
20 Leu Gln Thr Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Val Arg
25
<210> 613
<211> 98
<212> PRT
30 <213> Homo sapiens
<400> 613
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ser
20 25 30
Asp Met Asn Trp Ala Arg Lys Ala Pro Gly Lys Gly Leu Glu Trp Val
40 35 40 45
Ser Gly Val Ser Trp Asn Gly Ser Arg Thr His Tyr Val Asp Ser Val
50 55 60
45 Lys Arg Arg Phe Ile Ile Ser Arg Asp Asn Ser Arg Asn Ser Leu Tyr
65 70 75 80
Leu Gln Lys Asn Arg Arg Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys
85 90 95
50 Val Arg

<210> 614
55 <211> 98
<212> PRT
<213> Homo sapiens
<400> 614
60 Thr Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Glu Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ser
20 25 30
65 Asp Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Val Ser Trp Asn Gly Ser Arg Thr His Tyr Ala Asp Ser Val
50 55 60

5 Lys Gly Arg Phe Ile Ile Ser Arg Asp Asn Ser Arg Asn Phe Leu Tyr
65 70 75 80

Gln Gln Met Asn Ser Leu Arg Pro Glu Asp Met Ala Val Tyr Tyr Cys
85 90 95

10 Val Arg

15 <210> 615

<211> 97

<212> PRT

<213> Homo sapiens

20 <400> 615

Glu Val His Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ala Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30

25

Asp Met His Trp Val Arg Gln Ala Thr Gly Lys Gly Leu Glu Trp Val
35 40 45

30 Ser Ala Asn Gly Thr Ala Gly Asp Thr Tyr Tyr Pro Gly Ser Val Lys
50 55 60

Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Ser Leu Tyr Leu
65 70 75 80

35 Gln Met Asn Ser Leu Arg Ala Gly Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

Arg

40

<210> 616

<211> 97

<212> PRT

<213> Homo sapiens

45

<400> 616

Glu Val Gln Leu Val Glu Thr Gly Gly Leu Ile Gln Pro Gly Gly
1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Val Ser Ser Asn
20 25 30

Tyr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

55

Ser Val Ile Tyr Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
50 55 60

60

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu
65 70 75 80

Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

65 Arg

<210> 617

<211> 97
<212> PRT
<213> Homo sapiens

5 <400> 617
Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Ser Tyr
10 20 25 30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
15 Ser Ala Ile Gly Thr Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50 55 60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu
65 70 75 80
20 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala
85 90 95

Arg

25
30 <210> 618
<211> 97
<212> PRT
<213> Homo sapiens
<400> 618
Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Ser Tyr
35 20 25 30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
40 35 40 45
Ser Ala Ile Gly Thr Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50 55 60
45 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu
65 70 75 80
Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala
85 90 95
50

Arg

55 <210> 619
<211> 98
<212> PRT
<213> Homo sapiens
<400> 619
60 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
65 20 25 30
Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Lys

<210> 620
 <211> 98
 15 <212> PRT
 <213> Homo sapiens

<400> 620
 20 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ser Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

25 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Val
 35 40 45

Ser Ala Ile Ser Ser Asn Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

30 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

35 Val Gln Met Ser Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Val Lys

40

<210> 621
 <211> 98
 <212> PRT
 <213> Homo sapiens

45

<400> 621
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ser Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Val
 35 40 45

55 Ser Ala Ile Ser Ser Asn Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Val Gln Met Ser Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

65 Val Lys

<210> 622

<211> 98
<212> PRT
<213> Homo sapiens

5 <400> 622
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

10 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Val
 35 40 45

15 Ser Ala Ile Ser Ser Asn Gly Gly Ser Thr Tyr Tyr Ala Asn Ser Val
 50 55 60

20 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
Leu Gln Met Gly Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys
 85 90 95

25 Ala Arg

30 <210> 623
<211> 98
<212> PRT
<213> Homo sapiens

<400> 623
Glu Val Gln Leu Leu Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
35 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

40 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

50 Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Gly Asp Ser Val
 50 55 60
45 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

55 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys

55 <210> 624
<211> 98
<212> PRT
<213> Homo sapiens

60 <400> 624
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

65 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Thr Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Arg

15 <210> 625
 <211> 98
 <212> PRT
 <213> Homo sapiens

20 <400> 625
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 25 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

30 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

35 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

40 Ala Arg

<210> 626
 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 626
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 55 35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Ser Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

65 Ala Arg

5 <210> 627
 <211> 98
 <212> PRT
 <213> Homo sapiens

10 <400> 627
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 15 35 40 45
 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60
 20 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 25 Ala Arg

30 <210> 628
 <211> 98
 <212> PRT
 <213> Homo sapiens

35 <400> 628
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 40 35 40 45
 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 45 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 50 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg

55 <210> 629
 <211> 98
 <212> PRT
 <213> Homo sapiens

60 <400> 629
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 65 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

	35	40	45
	Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50	55	60
5	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
			80
10	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
		85	90
			95
	Ala Arg		
15	<210> 630		
	<211> 98		
	<212> PRT		
	<213> Homo sapiens		
20	<400> 630		
	Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg		
	1	5	10
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr		
25	20	25	30
	Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
	35	40	45
30	Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50	55	60
	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
			80
35	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90	95
	Ala Arg		
40	<210> 631		
	<211> 98		
	<212> PRT		
45	<213> Homo sapiens		
	<400> 631		
	Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg		
	1	5	10
50	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr		
	20	25	30
	Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
55	35	40	45
	Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val		
	50	55	60
60	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
			80
	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90	95
65	Ala Arg		

5 <210> 632
 <211> 98
 <212> PRT
 <213> Homo sapiens

10 <400> 632
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

15 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

20 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

25 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

30 Ala Arg

35 <210> 633
 <211> 98
 <212> PRT
 <213> Homo sapiens

40 <400> 633
 Gln Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ser Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

45 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Val
 35 40 45

50 Ser Ala Ile Ser Ser Asn Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

55 Val Gln Met Ser Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Val Lys

60 <210> 634
 <211> 98
 <212> PRT
 <213> Homo sapiens

65 <400> 634
 Gln Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ser Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Val
 35 40 45

Ser Ala Ile Ser Ser Asn Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Arg

<210> 635

15 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 635

20 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

25 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 30 50 55 60

Lys Gly Arg Phe Ala Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

35 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

40 <210> 636
 <211> 98
 <212> PRT
 <213> Homo sapiens

45 <400> 636

Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

55 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

65 Ala Arg

<210> 637

<211> 98
<212> PRT
<213> Homo sapiens

5 <400> 637
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

10 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

15 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys

25

<210> 638

<211> 97

<212> PRT

30 <213> Homo sapiens

<400> 638

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

40 Asp Met His Trp Val Arg Gln Ala Thr Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ala Ile Gly Thr Ala Gly Asp Thr Tyr Tyr Pro Gly Ser Val Lys
50 55 60

45 Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Ser Leu Tyr Leu
65 70 75 80

Gln Met Asn Ser Leu Arg Ala Gly Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

50

Arg

55 <210> 639
<211> 98
<212> PRT
<213> Homo sapiens

60 <400> 639
Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

65 Glu Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Arg

<210> 640
 <211> 98
 15 <212> PRT
 <213> Homo sapiens

<400> 640
 20 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

25 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

30 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Leu Arg Ala Arg Leu Cys Ile Thr Val
 35 85 90 95

Arg Glu

<210> 641
 40 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 641
 45 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

50 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 55 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

60 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

65 <210> 642
 <211> 98
 <212> PRT

<213> Homo sapiens

<400> 642

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

10 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

15 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg

25 <210> 643

<211> 98

<212> PRT

<213> Homo sapiens

30 <400> 643

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

40 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

45 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg

50

<210> 644

<211> 98

55 <212> PRT

<213> Homo sapiens

<400> 644

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
60 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

65 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val

	50	55	60														
	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	
	65				70						75					80	
5	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys																
					85						90					95	
	Ala Arg																
10																	
	<210> 645																
	<211> 98																
15	<212> PRT																
	<213> Homo sapiens																
	<400> 645																
20	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg		
	1			5						10					15		
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr																
					20					25					30		
25	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
	35					40						45					
	Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val																
30		50			55						60						
	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Arg	Leu	Tyr	
	65				70						75					80	
35	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys																
					85						90					95	
	Ala Arg																
40	<210> 646																
	<211> 98																
	<212> PRT																
	<213> Homo sapiens																
45	<400> 646																
50	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg		
	1			5						10					15		
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr																
					20					25					30		
	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val																
		35			40						45						
55	Ala	Val	Ile	Ser	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala	Asp	Ser	Val	
	50			55						60							
	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	
	65				70						75					80	
60	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys																
					85						90					95	
	Ala Arg																
65	<210> 647																
	<211> 98																

<212> PRT

<213> Homo sapiens

<400> 647

5 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

10 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

15 Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg

25

<210> 648

<211> 98

<212> PRT

30 <213> Homo sapiens

<400> 648

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

40 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

45 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Gly Thr Ala Val Tyr Tyr Cys
85 90 95

50 Ala Arg

55 <210> 649

<211> 98

<212> PRT

<213> Homo sapiens

60 <400> 649

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Gly
1 5 10 15

65 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Lys

15 <210> 650
 <211> 98
 <212> PRT
 <213> Homo sapiens

20 <400> 650
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 25 20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

30 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

35 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

40 Ala Lys

45 <210> 651
 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 651
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 50 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

55 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

65 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

5 <210> 652
 <211> 98
 <212> PRT
 <213> Homo sapiens

10 <400> 652
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

15 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

 Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

20 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

25 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

 Ala Lys

30 <210> 653
 <211> 95
 <212> PRT
 <213> Homo sapiens

35 <400> 653
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

45 Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

50 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Arg Lys
 85 90 95

55 <210> 654
 <211> 98
 <212> PRT
 <213> Homo sapiens

60 <400> 654
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

65 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Arg

<210> 655
 15 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 655
 20 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

25 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Ala
 30 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Thr Asn Thr Leu Phe
 65 70 75 80

35 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

40

<210> 656
 <211> 98
 <212> PRT
 <213> Homo sapiens

45

<400> 656
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

55

Ser Tyr Ile Ser Ser Ser Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60

60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

65 Ala Arg

<210> 657

<211> 98
<212> PRT
<213> Homo sapiens

5 <400> 657
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
10 20 25 30
Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
15 35 40 45
15 Ser Ser Ile Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
20 65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
25 85 90 95
Ala Arg

25

<210> 658
<211> 97
30 <212> PRT
<213> Homo sapiens

<400> 658
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
35 1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
40 20 25 30
Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
45 35 40 45
Ser Ser Ile Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val Lys
50 50 55 60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu
55 65 70 75 80
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
50 85 90 95
Arg

55 <210> 659
<211> 98
<212> PRT
<213> Homo sapiens

60 <400> 659
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
65 1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
65 20 25 30
Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
70 35 40 45

Ser Ser Ile Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

10 Ala Arg

15 <210> 660
 <211> 98
 <212> PRT
 <213> Homo sapiens

20 <400> 660
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

25 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Leu Glu Trp Val
 35 40 45

30 Ser Tyr Ile Ser Ser Ser Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

35 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

40 Ala Arg

45 <210> 661
 <211> 97
 <212> PRT
 <213> Homo sapiens

<400> 661
 Glu Asp Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

50 Ser Leu Arg Pro Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ser Tyr
 20 25 30

Val Leu His Trp Val Arg Arg Ala Pro Gly Lys Gly Pro Glu Trp Val
 55 35 40 45

Ser Ala Ile Gly Thr Gly Gly Asp Thr Tyr Tyr Ala Asp Ser Val Met
 50 55 60

60 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Ser Leu Tyr Leu
 65 70 75 80

Gln Met Asn Ser Leu Ile Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala
 85 90 95

65 Arg

5 <210> 662
 <211> 98
 <212> PRT
 <213> Homo sapiens

10 <400> 662
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

15 Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Val Trp Val
 35 40 45
 Ser Arg Ile Asn Ser Asp Gly Ser Ser Thr Ser Tyr Ala Asp Ser Val
 50 55 60

20 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

25 Ala Arg

30 <210> 663
 <211> 98
 <212> PRT
 <213> Homo sapiens

35 <400> 663
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Val Trp Val
 35 40 45

45 Ser Arg Ile Asn Ser Asp Gly Ser Ser Thr Ser Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

50 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

55 <210> 664
 <211> 98
 <212> PRT
 <213> Homo sapiens

60 <400> 664
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

65 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

5 Ala Asn Ile Lys Gln Asp Gly Ser Glu Lys Tyr Tyr Val Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

10 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

15

<210> 665
 <211> 98
 <212> PRT
 20 <213> Homo sapiens

<400> 665
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

25 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Val Trp Val
 30 35 40 45

Ser Arg Ile Asn Ser Asp Gly Ser Ser Thr Ser Tyr Ala Asp Ser Met
 50 55 60

35 Lys Gly Gln Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys
 85 90 95

40 Thr Arg

45 <210> 666
 <211> 98
 <212> PRT
 <213> Homo sapiens

50 <400> 666
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

55 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

60 Ala Asn Ile Lys Gln Asp Gly Ser Glu Lys Tyr Tyr Val Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

65 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

5 <210> 667
<211> 98
<212> PRT
<213> Homo sapiens

10 <400> 667
Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

15 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

20 Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

25 Leu Gln Met Lys Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

30 Thr Thr

35 <210> 668
<211> 98
<212> PRT
<213> Homo sapiens

<400> 668
Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Ala Ala Pro Gly Gln
1 5 10 15

40 Lys Val Thr Ile Ser Cys Ser Gly Ser Ser Asn Ile Gly Asn Asn
20 25 30

45 Tyr Val Ser Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35 40 45

Ile Tyr Asp Asn Asn Lys Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser
50 55 60

50 Gly Ser Lys Ser Gly Thr Ser Ala Thr Leu Gly Ile Thr Gly Leu Gln
65 70 75 80

Thr Gly Asp Glu Ala Asp Tyr Tyr Cys Gly Thr Trp Asp Ser Ser Leu
85 90 95

55 Ser Ala

60 <210> 669
<211> 98
<212> PRT
<213> Homo sapiens

<400> 669
65 Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Ala Ala Pro Gly Gln
1 5 10 15

Lys Val Thr Ile Ser Cys Ser Gly Ser Ser Asp Met Gly Asn Tyr

	20	25	30
5	Ala Val Ser Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35	40	45
Ile Tyr Glu Asn Asn Lys Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser			
50	55	60	
10	Gly Ser Lys Ser Gly Thr Ser Ala Thr Leu Gly Ile Thr Gly Leu Trp		
	65	70	75
			80
	Pro Glu Asp Glu Ala Asp Tyr Tyr Cys Leu Ala Trp Asp Thr Ser Pro		
	85	90	95
15	Arg Ala		
20	<210> 670		
	<211> 98		
	<212> PRT		
	<213> Homo sapiens		
25	<400> 670		
	Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln		
	1	5	10
			15
	Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn		
	20	25	30
30	Thr Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35	40	45
	Ile Tyr Ser Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
	50	55	60
35	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln		
	65	70	75
			80
40	Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu		
	85	90	95
	Asn Gly		
45	<210> 671		
	<211> 98		
	<212> PRT		
	<213> Homo sapiens		
50	<400> 671		
	Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln		
	1	5	10
			15
55	Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn		
	20	25	30
	Tyr Val Tyr Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu		
	35	40	45
60	Ile Tyr Arg Asn Asn Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser		
	50	55	60
	Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Arg		
	65	70	75
			80
	Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu		
	85	90	95

Ser Gly

5 <210> 672
<211> 98
<212> PRT
<213> Homo sapiens
10 <400> 672
Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Glu Ala Pro Arg Gln
1 5 10 15
15 Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Asn Asn
20 25 30
Ala Val Asn Trp Tyr Gln Gln Leu Pro Gly Lys Ala Pro Lys Leu Leu
35 40 45
20 Ile Tyr Tyr Asp Asp Leu Leu Pro Ser Gly Val Ser Asp Arg Phe Ser
50 55 60
25 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln
65 70 75 80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu
85 90 95
30 Asn Gly

35 <210> 673
<211> 99
<212> PRT
<213> Homo sapiens
40 <400> 673
Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30
45 Tyr Val Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45
50 Leu Ile Tyr Gly Asn Ser Asn Arg Pro Ser Gly Val Pro Asp Gln Phe
50 55 60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu
65 70 75 80
55 Gln Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Lys Ala Trp Asp Asn Ser
85 90 95
60 Leu Asn Ala

65 <210> 674
<211> 99
<212> PRT
<213> Homo sapiens
70 <400> 674
Gln Ser Val Val Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln

1	5	10	15
Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly			
20		25	30
Tyr Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu			
35		40	45
Leu Ile Tyr Gly Asn Ser Asn Arg Pro Ser Gly Val Pro Asp Arg Phe			
50		55	60
Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu			
65		70	75
80			
15 Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser			
85		90	95
Leu Ser Gly			
20			
<210> 675			
<211> 98			
<212> PRT			
25 <213> Homo sapiens			
<400> 675			
Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Gly Thr Pro Gly Gln			
1		5	10
15			
30 Arg Val Thr Ile Ser Cys Ser Gly Gly Arg Ser Asn Ile Gly Ser Asn			
20		25	30
35 Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu			
35		40	45
Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser			
50		55	60
40 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Val Gln			
65		70	75
80			
45 Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser Leu			
85		90	95
50 Arg Gly			

50




Fig. 1




Fig. 2





Fig. 3

Percentage of Week-12 PASI 75 Responders Who Maintained \geq PASI 50 Response at Week 24



* 1 of 1 patients in the placebo group who was a PASI 75 responder at Week 12 maintained a PASI 50 response at Week 24 following discontinuation of placebo at Week 12.

Fig. 4A






Fig. 4B

PASI 90 at Week 24 Following
Therapy Discontinuation at Week 12



[TT-NRI]

Fig. 4C



NRI.
* $p<0.001$ for all active treatment
arms vs. placebo.

Fig. 4D