

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29012 A2

(51) International Patent Classification⁷: **C07D 243/00**

(74) Agent: **WAECHTER, Dieter**; 124 Grenzacherstrasse,
CH-4070 Basle (CH).

(21) International Application Number: PCT/EP00/09554

(22) International Filing Date:
29 September 2000 (29.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99120519.6 15 October 1999 (15.10.1999) EP

(71) Applicant: **F. HOFFMANN-LA ROCHE AG** [CH/CH];
124 Grenzacherstrasse, CH-4070 Basle (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


(72) Inventors: **ADAM, Geo**; 8 Untere Staltenstrasse, 79650 Schopfheim (DE). **ALANINE, Alexander**; 5, rue d'Altkirch, F-68400 Riedisheim (FR). **GOETSCHI, Erwin**; 23 Landhofweg, CH-4153 Reinach (CH). **MUTEL, Vincent**; 15, place des Marechaux, F-68100 Mulhouse (FR). **WOLTERING, Thomas, Johannes**; 13 Riedlistrasse, 79576 Weil am Rhein (DE).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: BENZODIAZEPINE DERIVATIVES



(57) Abstract: The present invention relates to compounds of general formula (I) wherein R³ is phenyl; pyridine; thiophenyl or thiazolyl, which are optionally substituted. The compounds according to the present invention can be used for treating or preventing acute and/or chronic neurological disorders such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.

Benzodiazepine derivatives

The present invention relates to compounds of general formula I



wherein

X is a single bond or an ethynediyl group, wherein,

5 in case X is a single bond,

R¹ is hydrogen; halogen; nitro; lower alkyl; halo-lower alkyl; alkoxycarbonyl; lower cycloalkyl, optionally substituted with oxygen; benzoyl, optionally substituted with lower alkyl, halo-lower alkyl, or halogen; phenyl, optionally substituted with halogen, hydroxy, lower alkyl, halo-lower alkyl, lower cycloalkyl, or lower alkoxy, halo-lower alkoxy, or
 10 cyano; styrenyl; phenylethyl; naphthyl; biphenyl; benzofuranyl; or 5 or 6 membered heterocyclic ring, optionally substituted with oxo, benzyloxy, benzoyl, methanesulfonyl, benzenesulfonyl, or acetyl;

 in case X is an ethynediyl group,

R¹ is hydrogen; lower alkyl, optionally substituted with hydroxy; halo-lower alkyl; lower
 15 cycloalkyl, optionally substituted with hydroxy, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, or halogen; lower cycloalkenyl, optionally substituted with lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, halogen, or oxo; lower alkenyl; phenyl, optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower cycloalkyl, lower alkoxy, halo; 5 or 6 membered heterocyclic ring, optionally substituted with lower alkyl,
 20 halogen, oxo, benzyloxy, benzoyl, methanesulfonyl, benzenesulfonyl, acetyl, pivaloyl, tert. butoxycarbonyl, or tert. butylcarbonyl; or benzofuranyl;

R³ is phenyl; pyridine; thiophenyl or thiazolyl, which are optionally substituted with halogen, cyano, nitro, azido, hydroxy, carboxy, morpholine-4-carbonyl, carbamoyl, thiocarbamoyl, N-hydroxycarbamoyl, trimethylsilyl-ethynyl, or from lower alkyl, lower alkynyl, lower alkoxy, halo-lower alkyl, 4-lower alkyl-piperazine-1-carbonyl, lower alkylaminocarbonyl, which are optionally substituted by amino, lower alkylamino, acylamino, oxo, hydroxy; lower alkoxy, lower alkylthio, or carboxy which is optionally esterified or amidated; or a 5-membered aromatic heterocycle which is optionally substituted by amino, lower alkylamino, acylamino, oxo, hydroxy, lower alkoxy, lower alkylthio, carboxy which is optionally esterified with lower alkyl or amidated with lower alkylamino which is eventually substituted by hydroxy, or lower alkyl which is optionally substituted by halogen, hydroxy, amino, lower alkylamino, acylamino, or amidino, which is optionally substituted by lower alkyl, -C(NRR')=NR" (where R, R' and R" are hydrogen or lower alkyl), hydroxy, lower alkoxy, lower alkylthio, acyloxy, lower alkylsulfinyl, lower alkylsulfonyl, lower alkoxy-lower alkylsulfanyl, lower alkylsulfanyl, cycloalkylsulfinyl, cycloalkylsulfonyl, hydroxyimino, or lower alkoxyimino, which is optionally esterified or amidated, lower alkenyl, oxo, cyano, carbamoyloxy, or sulfamoyl which is optionally substituted by lower alkyl,

with the proviso that, if X is a single bond and R³ is pyridinyl, R¹ is not hydrogen, or methyl;

and their pharmaceutically acceptable acid addition salts.

It has surprisingly been found that the compounds of general formula I are metabotropic glutamate receptor antagonists. Compounds of formula I are distinguished by valuable therapeutic properties.

In the central nervous system (CNS) the transmission of stimuli takes place by the interaction of a neurotransmitter, which is sent out by a neuron, with a neuroreceptor.

L-glutamic acid, the most commonly occurring neurotransmitter in the CNS, plays a critical role in a large number of physiological processes. The glutamate-dependent stimulus receptors are divided into two main groups. The first main group forms ligand-controlled ion channels. The metabotropic glutamate receptors (mGluR) form the second main group and, furthermore, belong to the family of G-protein-coupled receptors.

At present, eight different members of these mGluR are known and of these some even have sub-types. On the basis of structural parameters, the different influences on the synthesis of secondary metabolites and the different affinity to low-molecular weight

chemical compounds, these eight receptors can be subdivided into three sub-groups: mGluR1 and mGluR5 belong to group I, mGluR2 and mGluR3 belong to group II and mGluR4, mGluR6, mGluR7 and mGluR8 belong to group III.

Ligands of metabotropic glutamate receptors belonging to the group II can be used for the
5 treatment or prevention of acute and/or chronic neurological disorders such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.

Other treatable indications in this connection are restricted brain function caused by bypass operations or transplants, poor blood supply to the brain, spinal cord injuries, head injuries, hypoxia caused by pregnancy, cardiac arrest and hypoglycaemia. Further treatable
10 indications are chronic and acute pain, Huntington's chorea, amyotrophic lateral sclerosis (ALS), dementia caused by AIDS, eye injuries, retinopathy, idiopathic parkinsonism or parkinsonism caused by medicaments as well as conditions which lead to glutamate-deficiency functions, such as e.g. muscle spasms, convulsions, migraine, urinary incontinence, nicotine addiction, opiate addiction, anxiety, vomiting, dyskinesia and
15 depressions.

Objects of the present invention are compounds of formula I and their pharmaceutically acceptable salts per se and as pharmaceutically active substances, their manufacture, medicaments based on one or more compounds in accordance with the invention and their production, as well as the use of the compounds in accordance with the invention in
20 the control or prevention of illnesses of the aforementioned kind, and, respectively, for the production of corresponding medicaments.

Preferred compounds of formula I are those in which R³ is phenyl substituted in meta position by cyano; halogen; or imidazolyl, which is optionally substituted by lower alkyl; or 1,3-thiazolyl which is optionally substituted by hydroxy-lower alkyl, carboxy, or -CO-NH-
25 (CH₂)₂OH; 1,3-oxazolyl; 1,2,3-triazolyl; 1,2,4-triazolyl which is optionally substituted with lower alkyl; tetrazolyl; or isoxazolyl, which is optionally substituted by lower alkyl;

The following are examples of such compounds:

3-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;
4-(3-Chloro-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
30 4-(3-Imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;
8-(4-Fluoro-phenylethynyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-
benzo[b][1,4]diazepin-2-one;

- 8-(4-Fluoro-phenylethynyl)-4-(3-[1,2,4]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(4-Fluoro-phenyl)-4-[3-(4-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 5 8-(4-Fluoro-2-methyl-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(4-Fluoro-2-hydroxy-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(4-Fluoro-phenylethynyl)-4-(3-tetrazol-1-yl-phenyl)-1,3-dihydro-
- 10 benzo[b][1,4]diazepin-2-one;
- 8-(2-Fluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(4-Fluoro-phenyl)-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 15 8-(2,4-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(2,3-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(2-Fluoro-phenyl)-4-[3-(4-hydroxymethyl-thiazol-2-yl)-phenyl]-1,3-dihydro-
- 20 benzo[b][1,4]diazepin-2-one;
- 2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid;
- 2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-4-methyl-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide; and
- 25 4-[3-(4,5-Dimethyl-4H-[1,2,4]triazol-3-yl)-phenyl]-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;

Compounds of formula I, in which R³ is thiophenyl, preferably thiophen-2-yl, optionally substituted with halogen, cyano; or pyridinyl, preferably pyridin-4-yl, optionally substituted, preferably in the 2-position, with halogen, or cyano; are also preferred.

- 30 The following are examples of such compounds:

- 8-(4-Fluoro-phenylethynyl)-4-(2-imidazol-1-yl-pyridin-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 2-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile;
- 35 4-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-pyridine-2-

carbonitrile; and

4-[7-(2,4-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile.

All tautomeric forms of the compounds of invention are also embraced herewith.

- 5 The term "lower alkyl" used in the present description denotes straight-chain or branched saturated hydrocarbon residues with 1-7 carbon atoms, preferably with 1-4 carbon atoms, such as methyl, ethyl, n-propyl, i-propyl and the like.

The term "lower alkynyl" used in the present description denotes straight-chain or branched unsaturated hydrocarbon residues with 2-7 carbon atoms, preferably with 2-4
10 carbon atoms, such as ethynyl, n-propynyl, and the like.

The term "lower cycloalkyl" used in the present description denotes cyclic saturated hydrocarbon residues with 3-5 carbon atoms, preferably with 3 carbon atoms, such as cyclopropyl.


The term "lower alkoxy" denotes a lower alkyl residue in the sense of the foregoing
15 definition bonded via an oxygen atom.

The term "halogen" embraces fluorine, chlorine, bromine and iodine.


The expression "5 or 6 membered heterocyclic ring" embraces thiophene, furane, thiazole, pyridine, partially hydrated pyridine, for example 2-pyridone, partially hydrogenated pyridine, for example tetrahydropyridine, five-membered aromatic heterocycle containing
20 up to 4 heteroatoms, selected from O, S, N, embracing imidazol-1-yl, imidazol-2-yl, imidazol-4-yl; pyrazol-1-yl, pyrazol-3-yl, pyrazol-4-yl; 1,3-thiazol-2-yl, 1,3-thiazol-4-yl, 1,3-thiazol-5-yl; 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 1,3-oxazol-5-yl, 1,2-oxazol-3-yl, 1,2-oxazol-4-yl, 1,2-oxazol-5-yl; 1,2,3-triazol-1-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-1-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-2-yl; 1,2,4-oxadiazol-2-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,3-oxadiazol-4-yl, 1,2,3-oxadiazol-5-yl; 1,2,4-thiadiazol-2-yl, 1,2,4-thiadiazol-3-yl,
25 1,2,4-thiadiazol-5-yl, 1,2,3-thiadiazol-4-yl, 1,2,3-thiadiazol-5-yl; tetrazol-1-yl, tetrazol-2-yl, tetrazol-5-yl;

The compounds of general formula I and their pharmaceutically acceptable salts can be manufactured according to the following methods:

- 6 -

Scheme A

Compounds of general formula I, in which R^1 , R^3 and X are as described above can be prepared according to scheme A, by, for example, cleaving the BOC protecting group in compounds of general formula II, and concomitant cyclization of the deprotected compound. The deprotection-cyclization step can be carried out by treating the compounds of general formula II with a bronsted acid, like for example trifluoroacetic acid (TFA), in an inert solvent, like for example dichloromethane (DCM). The reaction is preferably carried out at temperatures between 0 °C and 50 °C. It may be advantageous to use also anisole or 1,3-dimethoxybenzene as a carbocation scavenger in the reaction mixture. Any other suitable amino protecting group, such as e.g. Fmoc or benzyloxycarbonyl (Z), can be alternatively used instead of the BOC group.


Scheme B

Compounds of general formula II, in which R^1 , R^3 and X are as described above, can be prepared according to scheme B by for example reacting a compound of general formula III with a dioxinone (general formula IV) in an inert solvent like for example toluene or xylene at elevated temperatures, preferably between 80 °C and 160 °C.


Alternatively, compounds of general formula II can also be prepared by for example reaction of a compound of general formula III with a β -ketoester (general formula IVa), in which R^3 is as described above using the same conditions as described for the reaction with the dioxinones.

- 7 -

Scheme C



According to scheme C, compounds of general formula III in which R^1 is as described above for compounds where X is a single bond can be prepared by different routes from the iodo-compound IX, depending on the nature of R^1 . As shown in scheme C, the key steps are coupling reactions of Suzuki- and Stille-type in presence or absence of carbonmonoxide. The exact conditions for the respective compounds can be found in the experimental part.

Scheme D


GP A, method a: diphosgene, EtOAc, 77 °C; then t-BuOH

GP A, method b: Boc_2O , Cs_2CO_3 , 2-butanone, 52 °C

GP A, method c: i) Boc_2O , DMAP, THF; ii) TFA, DCM, 0 °C

According to scheme D, the key intermediate iodide IX can be prepared from commercially available 2-nitroaniline by a standard iodination-protection sequence.

5

Scheme E

According to scheme E, compounds of general formula IIIa in which R^1 is as described above for compounds where X is an ethynediyl group can be prepared by different routes from the iodo-compound IX, depending on the nature of R^1 . As shown in Scheme E, the transformation can for example be carried out

10

a) by directly attaching the R^1 -alkynediyl-substituent via a Sonogashira-type coupling followed by the reduction of the nitro group or

b) by two stepwise Sonogashira -type couplings, in which first trimethylsilyl-acetylene is coupled to iodide IX to yield, after deprotection with sodium hydroxide in methanol, the intermediate XII which then can be transformed via a second Sonogashira-type coupling


15

- 9 -

with the appropriate reactant R^1 -I, R^1 -Br or R^1 -OSO₂CF₃ and reduction of the nitro group to the desired compounds.

The exact conditions for the respective compounds can be found in the experimental part.

Scheme F




5

According to Scheme F, the dioxinones and β -keto esters building blocks with the general formula IV and IVa can be prepared by methods known to someone skilled in the art from the corresponding carboxylic acid derivatives R^3 -COR, i.e. free acids, methyl or ethyl esters and acid chlorides. The exact conditions for the corresponding compounds can be found


10 in the experimental part.

Another synthetic route to prepare compounds of general formula Ic, in which R^1 and X have the meaning as described above and R^3 is a phenyl-carboxamide of general formula $C(O)NR^4R^5$, in which R^4 and R^5 are hydrogen, lower alkyl or R^4 and R^5 together form a morpholino-residue or a N-methyl-piperazine is outlined in scheme G:

Scheme G

The exact conditions for the respective compounds can be found in the experimental part.

Still another way to prepare compounds of general formula I is the reaction of 4-aryl-8-iodo-1,3-dihydro-benzo[b][1,4]diazepin-2-ones (General formula Id, Synthetic Scheme H) with alkynes of general formula $\text{R}^1\text{-C}\equiv\text{C-}$, in which R^1 has the meaning as described above, in a Sonogashira-coupling.

Scheme H

The exact conditions for the respective compounds can be found in the experimental part.

The pharmaceutically acceptable salts can be manufactured readily according to methods known per se and taking into consideration the nature of the compound to be converted into a salt. Inorganic or organic acids such as, for example, hydrochloric acid, hydrobromic acid, sulphuric acid, nitric acid, phosphoric acid or citric acid, formic acid, fumaric acid, maleic acid, acetic acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like are suitable for the formation of pharmaceutically acceptable salts of basic compounds of formula I.

The compounds of formula I and their pharmaceutically acceptable salts are metabotropic glutamate receptor antagonists and can be used for the treatment or prevention of acute and/or chronic neurological disorders, such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits. Other treatable indications are restricted brain function caused by bypass operations or transplants, poor blood supply to the brain, spinal cord injuries, head injuries, hypoxia caused by pregnancy, cardiac arrest and

hypoglycaemia. Further treatable indications are acute and chronic pain, Huntington's chorea, ALS, dementia caused by AIDS, eye injuries, retinopathy, idiopathic parkinsonism or parkinsonism caused by medicaments as well as conditions which lead to glutamate-deficient functions, such as e.g. muscle spasms, convulsions, migraine, urinary
5 incontinence, nicotine addiction, psychoses, opiate addiction, anxiety, vomiting, dyskinesia and depression.

The compounds of formula I and pharmaceutically acceptable salts thereof can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées,
10 hard and soft gelatin capsules, solutions, emulsions or suspensions. However, the administration can also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formula I and pharmaceutically acceptable salts thereof can be processed with pharmaceutically inert, inorganic or organic carriers for the production of
15 pharmaceutical preparations. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts and the like can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatin capsules. Suitable carriers for soft gelatin capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like; depending on the nature of the active substance no carriers are, however, usually required in the case
20 of soft gelatin capsules. Suitable carriers for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar, glucose and the like. Adjuvants, such as alcohols, polyols, glycerol, vegetable oils and the like, can be used for aqueous injection solutions of water-soluble salts of compounds of formula I, but as a rule are not necessary. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats,
25 semi-liquid or liquid polyols and the like.

In addition, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

30 As mentioned earlier, medicaments containing one or more compounds of formula I or a pharmaceutically acceptable salt thereof and a therapeutically inert excipient are also an object of the present invention, as is a process for the production of such medicaments which comprises bringing one or more compounds of formula I or pharmaceutically acceptable salts thereof and, if desired, one or more other therapeutically valuable

substances into a galenical dosage form together with one or more therapeutically inert carriers.

The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, the effective dosage for oral or parenteral
5 administration is between 0.01-20 mg/kg/day, with a dosage of 0.1-10 mg/ kg/day being preferred for all of the indications described. The daily dosage for an adult human being weighing 70 kg accordingly lies between 0.7-1400 mg per day, preferably between 7 and 700 mg per day.

The present invention relates also to the use of compounds of formula I and of
10 pharmaceutically acceptable salts thereof for the production of medicaments, especially for the control or prevention of acute and/or chronic neurological disorders of the aforementioned kind.

The compounds of the present invention are group II mGlu receptor antagonists as determined using the assay described by Cartmell et al. (Br. J. Pharmacol. 1998, 123(3),
15 497-504).

The compounds show activities, as measured in the assay described below, of 50 μ M or less, typically 3 μ M or less, and ideally of 0.5 μ M or less. In the table below are described some specific pKi values of preferred compounds

Compound	Ki mGlu2 (μ M)
3-(7-Iodo-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile	0.017
4-(3-Chloro-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.040
4-(3-Imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.006
3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile	0.055

8-(4-Fluoro-phenylethynyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.004
8-(4-Fluoro-phenylethynyl)-4-(3-[1,2,4]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.049
8-(4-Fluoro-phenylethynyl)-4-(2-imidazol-1-yl-pyridin-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.004
8-(4-Fluoro-phenyl)-4-[3-(4-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.016
8-(4-Fluoro-2-methyl-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.050
8-(4-Fluoro-2-hydroxy-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.170
2-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile	0.250
8-(4-Fluoro-phenylethynyl)-4-(3-tetrazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.036
4-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-pyridine-2-carbonitrile	0.039
4-[7-(2,4-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile	0.026
8-(2-Fluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.008
8-(4-Fluoro-phenyl)-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.289
8-(2,4-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.010
8-(2-Fluoro-phenyl)-4-(2-imidazol-1-yl-thiazol-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.298

8-(2,3-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.013
8-(2-Fluoro-phenyl)-4-[3-(4-hydroxymethyl-thiazol-2-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one.	0.016
2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid.	0.016
2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-4-methyl-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	0.049
4-[3-(4,5-Dimethyl-4H-[1,2,4]triazol-3-yl)-phenyl]-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.108

[³H]-LY354740 binding on mGlu2 transfected CHO cell membranes

Transfection and cell culture

- cDNA encoding the rat mGlu2 receptor protein in pBluescript II was obtained from Prof. S. Nakanishi (Kyoto, Japan), and subcloned into the eukaryotic expression vector pcDNA I-amp from Invitrogen (NV Leek, The Netherlands). This vector construct (pcD1mGR2) was co-transfected with a psvNeo plasmid encoding the gene for neomycin resistance, into CHO cells by a modified calcium phosphate method described by Chen & Okayama (1988). The cells were maintained in Dulbecco's Modified Eagle medium with reduced L-glutamine (2mM final concentration) and 10% dialysed foetal calf serum from Gibco BRL (Basel, Switzerland). Selection was made in the presence of G-418 (1000 ug/ml final). Clones were identified by reverse transcription of 5µg total RNA, followed by PCR using mGlu2 receptor specific primers 5'-atcactgcttggttctggcactg-3' and 5'-agcatcactgtgggtggcataggagc-3' in 60mM Tris HCl (pH 10), 15mM (NH₄)₂SO₄, 2 mM MgCl₂, 25 units/ml Taq Polymerase with 30 cycles annealing at 60°C for 1 min., extension at 72°C for 30s, and 1 min. 95°C denaturation.

Membrane preparation

Cells, cultured as above, were harvested and washed three times with cold PBS and frozen at -80°C. The pellet was resuspended in cold 20 mM HEPES-NaOH buffer containing 10

mM EDTA (pH 7.4), and homogenised with a polytron (Kinematica, AG, Littau, Switzerland) for 10s at 10 000rpm. After centrifugation for 30 min. at 4°C, the pellet was washed once with the same buffer, and once with cold 20 mM HEPES-NaOH buffer containing 0.1 mM EDTA, (pH 7.4). Protein content was measured using the Pierce
5 method (Socochim, Lausanne, Switzerland) using bovine serum albumin as standard.

[³H]-LY354740 binding

After thawing, the membranes were resuspended in cold 50mM Tris-HCl buffer containing 2 mM MgCl₂ and 2 mM CaCl₂, (pH 7) (binding buffer). The final concentration of the membranes in the assays was 25 µg protein/ml. Inhibition experiments were performed
10 with membranes incubated with 10 nM [³H]-LY354740 at room temperature, for 1 hour, in presence of various concentrations of the compound to be tested. Following the incubations, membranes were filtered onto Whatmann GF/C glass fiber filters and washed 5 times with cold binding buffer. Non specific binding was measured in the presence of 10 µM DCG IV (TOCRIS No. 0975). After transfer of the filters into plastic vials containing
15 10 ml of Ultima-gold scintillation fluid (Packard, Zürich, Switzerland), the radioactivity was measured by liquid scintillation in a Tri-Carb 2500 TR counter (Packard, Zürich, Switzerland).

Data analysis.

The inhibition curves were fitted with a four parameter logistic equation giving IC₅₀
20 values, and Hill coefficients.

EXAMPLES

General procedure A (Synthetic Scheme D):

Preparation of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters from 4-iodo-2-nitroanilines.

25 Method a

To a solution of diphosgene (4.1 mL, 34.1 mmol) in EtOAc (40 mL) at 0 °C was added a solution of the 4-iodo-2-nitroaniline (45.5 mmol) in EtOAc (200-500 mL), and the mixture was heated to reflux for 18 h. The solvent was removed in vacuum to leave a brown solid, which was triturated with hot hexane (200 mL). The solid material was
30 filtered off and the filtrate was concentrated under reduced pressure to leave the pure 4-iodo-2-nitrophenylisocyanate as a yellow solid. This material was refluxed in a mixture of excess tert.-BuOH in CH₂Cl₂ for 2.5 h. Removal of the solvent left an orange solid which

was purified by silica gel column chromatography with hexane/EtOAc to give the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester as a yellow solid.

Method b

To a mixture of the 4-iodo-2-nitroaniline (142 mmol) and cesium carbonate (55.5 g, 170 mmol) in 2-butanone (740 mL) was dropwise added a solution of Boc₂O (37.8 g, 173 mmol) in 2-butanone (170 mL) and the resulting mixture was stirred at 52 °C for 26 h. The solvent was removed in vacuum, the residue was treated with a mixture of H₂O (240 mL) and MeOH (240 mL) and extracted with hexane (3 × 500 mL). The combined hexane layer was washed with brine (200 mL) and all aqueous layers were reextracted with hexane (300 mL). All combined hexane layers were dried over MgSO₄, filtered and the solvent was removed in vacuum to give an orange solid, which was purified by silica gel column chromatography with hexane/EtOAc to give the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester as a yellow solid.

Method c

To a solution of the 4-iodo-2-nitroaniline (550 mmol) and DMAP (1.22 g, 10 mmol) in THF (1000 mL) at 23 °C was dropwise added within 70 min a solution of Boc₂O (246 g, 1128 mmol) in THF (500 mL) and stirring was continued at 23 °C for 75 min. The entire mixture was evaporated to dryness and dried at HV to leave a dark brown solid (253.59 g). This material was dissolved in DCM (1100 mL), cooled to 0 °C and TFA (84 mL, 1100 mmol) was added dropwise. The mixture was stirred at 0 °C for 2 h, poured into icecold sat. NaHCO₃-sol., extracted with DCM, washed with brine and dried over MgSO₄. Removal of the solvent in vacuum left a dark brown solid (199.71 g) which was coated on silica gel and purified by silica gel column chromatography with hexane/EtOAc to give the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester as a yellow solid.

25 Example A1

(4-Iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

Prepared the isocyanate from 4-iodo-2-nitroaniline (12.0 g, 45.5 mmol; prepared from 2-nitroaniline according to Wilson, J. Gerald; Hunt, Frederick C. *Aust. J. Chem.* 1983, 36, 2317-25; CAS-No. [20691-72-9]) with diphosgene (4.1 mL, 34.1 mmol) in EtOAc (250 mL), followed by treatment with tert.-BuOH (12 mL) in CH₂Cl₂ (60 mL) according to the general procedure A (method a). Obtained as a yellow solid (8.23 g, 82 %).

MS (EI) 390 (M^+); mp 92-94 °C.

Example A2

(4-Isopropyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

- Prepared from 4-isopropyl-2-nitroaniline (CAS-No. [63649-64-9]) by reaction with Boc_2O and cat. DMAP in THF, followed by treatment with TFA in CH_2Cl_2 according to the general procedure A (method c). Obtained as a yellow oil (14.1 g).

MS (EI) 280 (M^+).

Example A3

(4-Cyclopropyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

- Prepared from 4-cyclopropyl-2-nitro-phenylamine prepared from nitration of N-(4-cyclopropyl-phenyl)-acetamide (CAS-No. [63649-64-9] with 65% HNO_3 in Ac_2O and subsequent saponification with 6N NaOH in refluxing dioxane) by reaction with Boc_2O and cat. DMAP in THF, followed by treatment with TFA in CH_2Cl_2 according to the general procedure A (method c). Obtained as an orange liquid (2.33 g).
- MS (EI) 278 (M^+).

General procedure B (Synthetic Scheme C)

Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by direct Suzuki-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with arylboronic acids.

- A mixture of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (3.0 mmol), the arylboronic acid (4.5 mmol) and $PdCl_2(PPh_3)_2$ (2 mol%) was refluxed in 1,4-dioxane (25 mL) and 2M Na_2CO_3 -sol. (7.5 mL) [or alternatively with 1M $NaHCO_3$ -sol. (7.5 mL), LiCl (6.0 mmol) and $(Ph_3P)_4Pd$ (3 mol%) in DME (30 mL); also possible with Et_3N (9.0 mmol), $Pd(OAc)_2$ (3 mol%), PPh_3 (6 mol%) in DMF (10 mL) at 100 °C] until tlc indicated complete conversion of the iodide. The mixture was transferred into a separating funnel, H_2O (25 mL) was added and the product was extracted with ether or EtOAc (3 x 30 mL). The combined organic layers were washed with brine (50 mL) and dried over Na_2SO_4 . Removal of the solvent left a brown residue, which was purified by silica gel column chromatography with cyclohexane/ether or cyclohexane/EtOAc to give the title compound.

Example B1

(4'-Methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 4-methoxyphenylboronic acid according to the general procedure B. Obtained as a yellow solid (637 mg).

MS (ISN) 343 [(M-H)⁻]; mp 107-109 °C.

Example B2

(2-Nitro-4-thiophen-3-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 3-thiopheneboronic acid according to the general procedure B. Obtained as a yellow solid (326 mg).

MS (ISN) 319 [(M-H)⁻].

Example B3

(4-Furan-2-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and furan-2-boronic acid according to the general procedure B. Obtained as an orange solid (282 mg).

MS (EI) 304 (M⁺); mp 169-172 °C.

Example B4

(4'-Ethyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 4-ethylbenzene boronic acid according to the general procedure B. Obtained as an orange solid (689 mg).

MS (EI) 343 [(M+H)⁺]; mp 94-99 °C.

Example B5

(3-Nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (994 mg, 2.73 mmol) and phenyl boronic acid (576 mg, 3.00 mmol) according to the general procedure B. Obtained as a bright yellow solid (800 mg).

MS (EI) 314 (M^+); mp 119-121 °C.

5 Example B6

(4-Furan-3-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and furan-3-boronic acid according to the general procedure B. Obtained as an orange solid (855 mg).

10 MS (ISP) 322 [$(M+NH_4)^+$] and 327 [$(M+Na)^+$]; mp 105-110 °C.

Example B7

(4-Naphthalen-1-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 1-naphthylboronic acid according to the general procedure B. Obtained as a yellow foam
15 (1.0 g).

MS (ISN) 363 [$(M-H)^-$]; mp 60-66 °C.

Example B8

(3'-Methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and
20 3-methoxyphenylboronic acid according to the general procedure B. Obtained as an orange solid (818 mg).

MS (ISP) 345 [$(M+H)^+$], 362 [$(M+NH_4)^+$] and 367 [$(M+Na)^+$]; mp 104-107 °C.

Example B9

(3-Nitro-4'-trifluoromethoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 4-(trifluoromethoxy)benzene boronic acid according to the general procedure B. Obtained as an orange solid (569 mg).

MS (ISN) 397 [(M-H)⁻]; mp 145-147 °C.

5 Example B10

(2'-Fluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 2-fluorobenzene boronic acid according to the general procedure B. Obtained as a yellow solid (1.48 g).

10 MS (ISN) 331 [(M-H)⁻]; mp 131-133 °C.

Example B11

(3'-Fluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 3-fluorobenzene boronic acid according to the general procedure B. Obtained as a yellow
15 solid (3.87 g).

MS (ISN) 331 [(M-H)⁻]; mp 93-96 °C.

Example B12

(4'-Fluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and
20 4-fluorobenzene boronic acid according to the general procedure B. Obtained as a yellow solid (1.08 g).

MS (ISN) 331 [(M-H)⁻]; mp 155-167 °C.

Example B13

(4'-Fluoro-2'-methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

25 Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 4-fluoro-2-methylbenzene boronic acid [CAS-no. 139911-29-8; prepared from 2-bromo-

5-fluorotoluene by reaction with n-BuLi at -78°C followed by treatment with $\text{B}(\text{OMe})_3$ and subsequent hydrolysis] according to the general procedure B. Obtained as a yellow solid (1.71 g).

MS (EI) 346 (M^+).

5 Example B14

(4'-Fluoro-2'-methoxymethoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 4-fluoro-2-methoxymethoxybenzene boronic acid [prepared from 1-bromo-4-fluoro-2-(methoxymethoxy)benzene (CAS-no. 162269-78-5) by reaction with n-BuLi at -78°C followed by treatment with $\text{B}(\text{OMe})_3$ and subsequent hydrolysis] according to the general procedure B. Obtained as a yellow solid (0.96 g).

MS (EI) 392 (M^+).

Example B15

(2',4'-Difluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

15 Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 2,4-difluorobenzene boronic acid according to the general procedure B. Obtained as a yellow solid (3.26 g).

MS (ISN) 349 [$(\text{M}-\text{H})^-$].

Example B16

20 (2'-Fluoro-6'-methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 2-fluoro-6-methoxybenzene boronic acid according to the general procedure B. Obtained as a yellow solid (0.95 g).

MS (ISN) 361 [$(\text{M}-\text{H})^-$]; mp $65-68^{\circ}\text{C}$.

25 Example B17

(2',5'-Difluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and 2,5-difluorobenzene boronic acid according to the general procedure B. Obtained as a yellow solid (2.85 g).

MS (ISN) 349 [(M-H)⁻]; mp 104 °C.

5 Example B18

(4-Benzofuran-2-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) and benzo[b]furan-2-boronic acid according to the general procedure B. Obtained as an orange solid (711 mg).

10 MS (EI) 354 (M⁺); mp 175-177 °C.

Example B19

(2',3'-Difluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (3.64 g, 10 mmol) and 2,3-difluorobenzene boronic acid (2.35 g, 14.9 mmol) according to the
15 general procedure B. Obtained as a yellow solid (3.22 g).

MS (ISN) 349 [(M-H)⁻]; mp 93 °C.

General procedure C (Synthetic Scheme C)

Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by Suzuki-coupling
20 of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with bis(pinacolato)diboron and subsequent reaction with aryl halides.

A mixture of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (2.0 mmol), bis(pinacolato)diboron (2.2 mmol), KOAc (6.0 mmol) and PdCl₂(PPh₃)₂ (3 mol%) in 1,4-dioxane (25 mL) was stirred at 100 °C until tlc indicated complete conversion of the iodide
25 [cf. *Tetr. Lett.* 1997, 38, 3841-3844]. After addition of the aryl halide (4.0 mmol), PdCl₂(PPh₃)₂ (3 mol%) and 2M Na₂CO₃-sol. (7.5 mL) the mixture was stirred at 100 °C until tlc indicated complete conversion of the intermediate boronic ester. The mixture was transferred into a separating funnel, H₂O (30 mL) was added and the product was

extracted with ether or EtOAc (3 x 50 mL). The combined organic layers were washed with brine (100 mL) and dried over Na₂SO₄. Removal of the solvent left a brown residue, which was purified by silica gel column chromatography with cyclohexane/ether or cyclohexane/EtOAc to give the title compound.

5 Example C1

(2'-Methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1), bis(pinacolato)diboron and 2-iodoanisole according to the general procedure C. Obtained as a yellow solid (735 mg).

10 MS (EI) 344 (M⁺)

Example C2

(4'-Chloro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1), bis(pinacolato)diboron and 1-chloro-4-iodobenzene according to the general procedure C.

15 Obtained as a yellow solid (779 mg).

MS (ISN) 347 [(M-H)⁺]; mp 150-155 °C (dec.).

Example C3

(2-Nitro-4-thiophen-2-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),

20 bis(pinacolato)diboron and 2-iodothiophene according to the general procedure C. Obtained as a yellow solid (91 mg).

MS (ISN) 319 [(M-H)⁺].

Example C4

(4'-Methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

25 Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1), bis(pinacolato)diboron and 4-iodotoluene according to the general procedure C. Obtained as an orange solid (542 mg).

MS (ISP) 346 [(M+NH₄)⁺] and 367 [(M+Na)⁺]; mp 105-108 °C.

Example C5

(2-Nitro-4-pyridin-2-yl-phenyl)-carbamic acid tert.-butyl ester.

- Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
5 bis(pinacolato)diboron and 2-bromopyridine according to the general procedure C.
Obtained as a yellow solid (407 mg).

MS (ISP) 316 [(M+H)⁺].

Example C6

(3'-Methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

- 10 Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
bis(pinacolato)diboron and 3-iodotoluene according to the general procedure C.
Obtained as a yellow solid (524 mg).

MS (ISP) 346 (M⁺) and 674 [(2M+NH₄)⁺]; mp 83-85 °C.

Example C7

- 15 (3',4'-Dichloro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
bis(pinacolato)diboron and 3,4-dichloro-iodobenzene according to the general procedure
C. Obtained as a yellow solid (540 mg).

MS (EI) 382 (M⁺).

- 20 Example C8

(2'-Chloro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
bis(pinacolato)diboron and 1-chloro-2-iodobenzene according to the general procedure C.
Obtained as a yellow oil (886 mg).

- 25 MS (ISN) 347 [(M-H)⁻].

Example C9

(2'-Methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
bis(pinacolato)diboron and 2-iodotoluene according to the general procedure C. Obtained
5 as a yellow oil (755 mg).

MS (ISN) 327 [(M-H)⁺].

Example C10

(2-Nitro-4-pyridin-3-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
10 bis(pinacolato)diboron and 3-bromopyridine according to the general procedure C.
Obtained as a yellow solid (587 mg).

MS (ISP) 316 [(M+H)⁺]; mp 107-109 °C.

Example C11

(2-Nitro-4-pyridin-4-yl-phenyl)-carbamic acid tert.-butyl ester.

15 Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
bis(pinacolato)diboron and 4-bromopyridine according to the general procedure C.
Obtained as a yellow solid (379 mg).

MS (ISP) 316 [(M+H)⁺].

Example C12

20 (3''-Nitro-[1,1';4',1'']terphenyl-4''-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1),
bis(pinacolato)diboron and 4-bromobiphenyl according to the general procedure C.
Obtained as a yellow solid (1.29 g).

MS (EI) 390 (M⁺).

25 Example C13

(4'-Cyano-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1), bis(pinacolato)diboron and 4-bromobenzonitrile according to the general procedure C. Obtained as a yellow solid (1.38 g).

MS (ISN) 338 [(M-H)⁻].

5 General procedure D (Synthetic Scheme C)

Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by carbonylative Suzuki-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with aryl boronic acids

- 10 A mixture of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (1.0 mmol), aryl boronic acid (1.1 mmol), K₂CO₃ (3.0 mmol) and PdCl₂(PPh₃)₂ (3 mol%) in anisole (6 mL) was stirred at 80 °C under a CO-atmosphere until tlc indicated complete conversion of the iodide [cf. *Tetr. Lett.* 1993, 34, 7595-7598]. The mixture was transferred into a separating funnel, H₂O (30 mL) was added and the product was extracted with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (50 mL) and dried over Na₂SO₄.
- 15 Removal of the solvent left a yellow residue, which was purified by silica gel column chromatography with or hexane/EtOAc to give the title compound.

Example D1

(4-Benzoyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester

- 20 Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (364 mg, 1.0 mmol) and phenylboronic acid (134 mg, 1.1 mmol) according to the general procedure D. Obtained as a yellow solid (242 mg).

MS (EI) 342 (M⁺).

Example D2

(2-Nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester

- 25 A solution of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (3.64 g, 10 mmol), hexabutylstannane (7.5 mL, 15 mmol) and (Ph₃P)₄Pd (116 mg, 0.1 mmol) in toluene (20 mL) was heated to 60° C for 5 days [cf. *Bull. Chem. Soc. Jpn.* 1983, 56, 3855-3856]. The reaction mixture was diluted with toluene (150 mL), washed with aqueous KF-sol. (2 x 50 mL), brine and dried over MgSO₄. Removal of the solvent in vacuum left a
- 30 brown oil, which was purified by silica gel column chromatography with hexane/ether 49:1

to give the (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester as a yellow liquid (3.8 g, 72 %).

MS (EI) 467, 469, 471 [all (M-butyl)⁺].

General procedure E (Synthetic Scheme C)

- 5 Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters or (4-{alkenyl-, cycloalkenyl- or heterocycloalkenyl}-2-nitro-phenyl)-carbamic acid tert.-butyl esters by Stille-coupling of (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester with aryl halides or vinyl triflates or Stille-coupling of (4-iodo-2-nitrophenyl)-carbamic acid tert.-butyl ester with trialkylarylstannanes.
- 10 A mixture of (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester (Example D2) (525 mg, 1.0 mmol), aryl halide or vinyl triflate (0.95-6.0 mmol), anhydrous LiCl (126 mg, 3.0mmol) and Pd(PPh₃)₄ (5 mol%) in DME (3 mL) was stirred at 100 °C under argon atmosphere until tlc indicated complete consumption of the stannane. The reaction was cooled to 23 °C, stirred with sat. aqueous KF-sol. (5 mL) for 45 min, filtered
- 15 through celite, washed with ether and the filtrate was dried over MgSO₄. Removal of the solvent in vacuum left a brown oil, which was purified by silica gel column chromatography with hexane/EtOAc to give the title compound.

Example E1

- 4-(4-tert.-Butoxycarbonylamino-3-nitro-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic
- 20 acid tert.-butyl ester.

- Prepared from (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester (Example D2) (3.18 g, 6.06 mmol) and 4-trifluoromethylsulfonyloxy-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (1.99 g, 6.0 mmol) [prepared from Boc-4-piperidone by treatment with LDA in THF at -78 °C followed by reaction with N-phenyl-
- 25 bis(trifluoromethanesulfonimide) according to Wustrow et al. *Synthesis* 1991, 1993] according to the general procedure E. Obtained as an orange solid (1.304 g, 52 %).

MS (ISP) 420 [(M+H)⁺], 442 [(M+Na)⁺] and 458 [(M+K)⁺]; mp 85-87 °C.

Example E2

(2-Nitro-4-thiazol-2-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester (Example D2) (1.0 g, 1.9 mmol) and 2-bromothiazole (0.56 mL, 6.27 mmol) according to the general procedure E. Obtained as a yellow solid (160 mg).

MS (EI) 321 (M^+).

5 Example E3

[4-(6-Benzyloxy-pyridin-3-yl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (728 mg, 2.0 mmol) and 2-(phenylmethoxy)-5-(trimethylstannyl)-pyridine (766 mg, 2.2 mmol) [CAS-No. [188881-22-3], WO 9709311] according to the general procedure E. Obtained as
10 a yellow solid (876 mg).

MS (ISP) 422 [$(M+H)^+$]; mp 119-122 °C.

General procedure F(Synthetic Scheme E):

Preparation of (4-alkynyl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by Sonogashira-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with acetylenic
15 compounds; also Sonogashira-coupling of (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl esters with aryl halides; and Sonogashira-coupling of 8-iodo-4-aryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones with acetylenic compounds.

A mixture of the halide (3.0-4.5 mmol), acetylenic compound (3.0-4.5 mmol), Et_3N (13.5 mmol), $PdCl_2(PPh_3)_2$ (5 mol%) and PPh_3 (2.5 mol%) in THF (12 mL) [with very insoluble
20 material DMF (up to 12 mL) could be added] was stirred for 20 min at 23 °C while being purged with Argon. CuI (1.2 mol%) was added and stirring was continued at 60 °C under Argon atmosphere until tlc indicated complete conversion of the minor component [cf. J. Org. Chem. 1998, 63, 8551]. The mixture was transferred into a separating funnel, 5% citric acid (50 mL) was added and the product was extracted with EtOAc (2 x 100 mL). The
25 combined organic layers were washed with sat. $NaHCO_3$ -sol. (50 mL) and brine (50 mL), followed by drying over $MgSO_4$. Removal of the solvent left a yellow residue, which was purified by silica gel column chromatography with hexane/EtOAc and/or triturated with hexane or aqueous EtOH to give the title compound.

Example F1

30 (2-Nitro-4-trimethylsilanylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (3.64 g, 10 mmol) and trimethylsilylacetylene (4.2 mL, 30 mmol) according to the general procedure F. Obtained as a green oil (3.6 g).

MS (EI) 334 (M^+).

5 Example F2

(4-Ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

To a solution of (2-nitro-4-trimethylsilanylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example F1) (3.54 g, 10.6 mmol) in MeOH (10 mL) and THF (20 mL) at 0 °C was added 1N NaOH (13 mL) and stirring was continued at 23 °C for 30 min. Poured into cold 5%
10 citric acid, extracted with EtOAc (300 mL), washed with sat. NaHCO_3 -sol. and brine, dried over MgSO_4 . Removal of the solvent in vacuum left a dark brown oil, which was purified by silica gel column chromatography with hexane/EtOAc 19:1. Obtained as a yellow solid (2.4 g).

MS (EI) 262 (M^+); mp 102 °C.

15 Example F3

(2-Nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (5.64 g, 15 mmol) and phenylacetylene (2.47 mL, 22.5 mmol) according to the general procedure F. Obtained as a yellow solid (4.96 g).

20 MS (EI) 338 (M^+); mp 146-148 °C.

Example F4

(2-Nitro-4-p-tolylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (728 mg, 2.0 mmol) and 4-tolylacetylene (349 mg, 3.0 mmol) according to the general
25 procedure F. Obtained as a yellow solid (632 mg).

MS (EI) 352 (M^+); mp 164-165 °C.

Example F5

[4-(2-Chloro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (1.82 g, 5.0 mmol) and 2-chlorophenylacetylene (1.02 g, 7.5 mmol) according to the general
5 procedure F. Obtained as a yellow solid (1.8 g).

MS (ISN) 371 [(M-H)-] and 373 [(M+2-H)-]; mp 152-155 °C.

Example F6

[4-(4-Chloro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (728 mg, 2.0 mmol) and 4-chlorophenylacetylene (416 mg, 3.0 mmol) according to the general
10 procedure F. Obtained as a yellow solid (658 mg).

MS (EI) 372 (M⁺) and 374 [(M+2)⁺]; mp 213-218 °C.

Example F7

(2-Nitro-4-thiazol-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester.

15 Prepared from (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F2) (262 mg, 1.00 mmol) and 2-bromothiazole (0.14 mL, 1.50 mmol) according to the general procedure F. Obtained as a yellow solid (215 mg).

MS (ISN) 344 [(M-H)⁻]; mp 137 °C.

Example F8

20 (2-Nitro-4-pyridin-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F2) (262 mg, 1.0 mmol) and 2-bromopyridine (0.15 mL, 1.6 mmol) according to the general procedure F. Obtained as a yellow solid (293 mg).

MS (ISP) 340 [(M+H)⁺]; mp 142-144 °C.

25 Example F9

[4-(4-Fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F2) (525 mg, 2.0 mmol) and 1-fluoro-4-iodobenzene (0.35 mL, 3 mmol) according to the general procedure F. Obtained as a yellow solid (793 mg).

MS (ISN) 355 [(M-H)⁻]; mp 157-158 °C.

5 Example F10

[4-(2-Fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F2) (525 mg, 2.0 mmol) and 2-fluoro-1-iodobenzene (0.35 mL, 3 mmol) according to the general procedure F. Obtained as a yellow solid (759 mg).

10 MS (ISN) 355 [(M-H)⁻]; mp 140-142 °C.

Example F11

[4-(2,4-Difluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F2) (525 mg, 2.0 mmol) and 2,4-difluoro-1-iodobenzene (0.36 mL, 3 mmol) according to the
15 general procedure F. Obtained as a yellow solid (807 mg).

MS (ISN) 373 [(M-H)⁻]; mp 134-136 °C.

Example F12

[2-Nitro-4-(4-trifluoromethoxy-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F2)
20 (701 mg, 2.67 mmol) and 1-iodo-4-(trifluoromethoxy)benzene (1.0 g, 3.47 mmol) according to the general procedure F. Obtained as a yellow solid (1.10 g).

MS (ISN) 421 [(M-H)⁻]; mp 129-131 °C.

General procedure G (Synthetic Scheme E)

Preparation of the (2-amino-phenyl)-carbamic acid tert.-butyl esters by reduction of (2-
25 nitro-phenyl)-carbamic acid tert.-butyl esters; Also preparation of 4-aryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones by reduction and concomitant cyclization of 3-aryl-N-(2-nitro-phenyl)-3-oxo-propionamides.

Method a: Catalytic hydrogenation

A mixture of the nitro compound (1.0 mmol) in MeOH or EtOH and THF (1:1 to 1:9 ca. 20 mL) and 10% Palladium on carbon (20 mg) or Raney-Ni (20 mg) was stirred vigorously at 23 °C under hydrogen atmosphere until tlc indicated complete conversion. The catalyst
5 was filtered off, washed thoroughly with MeOH or EtOH and THF (1:1), the solvent was removed in vacuum to give the title compound, which was generally pure enough for further transformations.

Method b: Reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$

A mixture of the nitro compound (1.0 mmol) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (5.0 mmol) in EtOH (30
10 mL) was stirred at 70-80 °C under Argon atmosphere until tlc indicated complete conversion [cf. *Tetr. Lett.* 1984, 25, 839]. The reaction mixture was brought to pH 8 by addition of sat. NaHCO_3 -sol. and extracted with EtOAc (2 x 100 mL). The combined organic layer were washed with brine and dried over Na_2SO_4 . Removal of the solvent left a yellow solid, which - if necessary - can be purified by silica gel column chromatography.

15 Method c: Reduction with Zn and NH_4Cl

To a mixture of the nitro compound (1.0 mmol) in EtOH/THF/sat. NH_4Cl -sol. (1:1:1, 30 mL) was added Zinc dust (3.0 mmol) and the mixture was stirred at 70 °C under Argon atmosphere until tlc indicated complete conversion. Aqueous workup as described in method b.

20 Method d: Reduction with Fe and HOAc

To a mixture of the nitro compound (1.0 mmol) in THF/ H_2O (4:1, 10-50 mL) was added Fe powder (6.0 mmol), followed by HOAc (10-12 drops) and the mixture was stirred at 70 °C under Argon atmosphere until tlc indicated complete conversion. Aqueous workup as described in method b.

25 Example G1

(2-Amino-4-iodo-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A1) (2.18 g, 6.0 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (6.77 g, 30 mmol) according to the general procedure G (method b). Obtained as a brown-yellow solid (2.0 g).

30 MS (EI) 334 (M^+); mp 127-130 °C.

Example G2

(2-Amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example F3) (403 mg, 1.19 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1.34 g, 5.96 mmol) according to the general procedure G (method b). Obtained as an orange solid (237 mg).

MS (ISP) 309 $[(\text{M}+\text{H})^+]$; mp 177-178 °C.

Example G3

(2-Amino-4-p-tolyethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (2-nitro-4-p-tolyethynyl-phenyl)-carbamic acid tert.-butyl ester (Example F4) (640 mg, 1.82 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.0 g, 9.1 mmol) according to the general procedure G (method b). Obtained as a beige solid (569 mg).

MS (ISP) 323 $[(\text{M}+\text{H})^+]$; mp 175 °C.

Example G4

[2-Amino-4-(2-chloro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (4-(2-chloro-phenylethynyl)-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F5) (1.61 g, 4.3 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (5.3 g, 23.5 mmol) according to the general procedure G (method b). Obtained as a light brown solid (1.1 g).

MS (EI) 342 (M^+) and 344 $[(\text{M}+2)^+]$; mp 120-122 °C.

Example G5

(3-Amino-4'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B1) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (77 mg).

MS (ISP) 315 $[(\text{M}+\text{H})^+]$ and 337 $[(\text{M}+\text{Na})^+]$.

Example G6

(2-Amino-4-thiophen-3-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-thiophen-3-yl-phenyl)-carbamic acid tert.-butyl ester (Example B2) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (278 mg).

MS (ISP) 291 [(M+H)⁺].

5 Example G7

(2-Amino-4-furan-2-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-furan-2-yl-2-nitrophenyl)-carbamic acid tert.-butyl ester (Example B3) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a grey powder (212 mg).

10 MS (EI) 274 (M⁺).

Example G8

(3-Amino-4'-ethyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-ethyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B4) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (311 mg).

15 MS (ISP) 313 [(M+H)⁺] and 335 [(M+Na)⁺].

Example G9

(3-Amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B5) (100 mg, 0.32 mmol) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a white solid (85 mg).

20 MS (ISP) 285 [(M+H)⁺]; mp 137 °C.

Example G10

(3-Amino-2'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2'-methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C1) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (169 mg).

MS (ISP) 315 [(M+H)⁺].

5 Example G11

(2-Amino-4-furan-3-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-furan-3-yl-2-nitrophenyl)-carbamic acid tert.-butyl ester (Example B6) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a light brown solid. (744 mg).

10 MS (ISP) 275 [(M+H)⁺] and 297 [(M+Na)⁺]; mp 158-161 °C (dec.).

Example G12

(3-Amino-4'-chloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-chloro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C2) by reduction with Zn/NH₄Cl according to the general procedure G (method c).

15 Obtained as a green solid (162 mg).

MS (EI) 318 (M⁺).

Example G13

[2-Amino-4-(4-chloro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

20 Prepared from (4-(4-chloro-phenylethynyl)-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example F6) (606 mg, 1.63 mmol) by reduction with SnCl₂·2H₂O (1.83 g, 8.1 mmol) according to the general procedure G (method b). Obtained as a beige solid (406 mg).

MS (ISP) 343 [(M+H)⁺] and 345 [(M+2+H)⁺]; mp 170-173 °C.

Example G14

(2-Amino-4-naphthalen-1-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-naphthalen-1-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B7) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (226 mg).

MS (EI) 334 (M^+).

5 Example G15

(2-Amino-4-thiazol-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (2-nitro-4-thiazol-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example F7) (205 mg, 0.59 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (668 mg, 2.95 mmol) according to the general procedure G (method b). Obtained as a yellow solid (108 mg).

10 MS (ISP) 316 [$(M+H)^+$]; mp 182 °C.

Example G16

(3-Amino-4'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C4) by catalytic hydrogenation with Pd/C according to the general procedure G (method
15 a). Obtained as a white solid (125 mg).

MS (ISP) 299 [$(M+H)^+$], 321 [$(M+Na)^+$] and 337 [$(M+K)^+$].

Example G17

(2-Amino-4-pyridin-2-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-pyridin-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example
20 C5) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a light brown solid (339 mg).

MS (ISP) 286 [$(M+H)^+$] and 308 [$(M+Na)^+$].

Example G18

(2-Amino-4-thiophen-2-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-thiophen-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example C3) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a beige powder (151 mg).

MS (ISP) 291 [(M+H)⁺].

5 Example G19

(2-Amino-4-pyridin-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-pyridin-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example F8) (262 mg, 0.772 mmol) by reduction with SnCl₂·2H₂O (871 mg, 3.86 mmol) according to the general procedure G (method b). Obtained as a light brown solid (130
10 mg).

MS (EI) 309 (M⁺); mp 178 °C.

Example G20

(3-Amino-3'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (3'-methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
15 C6) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white powder (212 mg).

MS (EI) 298 (M⁺).

Example G21

(3-Amino-3',4'-dichloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

20 Prepared from (3',4'-dichloro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C7) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (485 mg).

MS (ISP) 353 (M⁺); mp 168-171 °C.

Example G22

25 (3-Amino-2'-chloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2'-chloro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C8) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (180 mg).

MS (ISP) 319 [(M+H)⁺], 341 [(M+Na)⁺] and 357 [(M+K)⁺].

5 Example G23

(3-Amino-2'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2'-methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C9) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white powder (281 mg).

10 MS (ISP) 299 [(M+H)⁺], 321 [(M+Na)⁺] and 337 [(M+K)⁺].

Example G24

(2-Amino-4-benzoyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-benzoyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example D1) (280 mg, 0.82 mmol) by catalytic hydrogenation with Raney-Ni according to the general
15 procedure G (method a). Obtained as a yellow foam (269 mg).

MS (ISP) 313 [(M+H)⁺].

Example G25

(3-Amino-3'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (3'-methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester
20 (Example B8) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (165 mg).

MS (ISP) 315 [(M+H)⁺].

Example G26

(2-Amino-4-pyridin-3-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-pyridin-3-yl-phenyl)-carbamic acid tert.-butyl ester (Example C10) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a grey powder (200 mg).

MS (ISP) 286 [(M+H)⁺].

5 Example G27

(3-Amino-4'-trifluoromethoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (3-nitro-4'-trifluoromethoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B9) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a white solid (371 mg).

10 MS (ISP) 369 [(M+H)⁺] and 391 [(M+Na)⁺].

Example G28

(2-Amino-4-pyridin-4-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (2-nitro-4-pyridin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example C11) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a light grey powder (305 mg).

MS (ISP) 286 [(M+H)⁺].

Example G29

4-(3-Amino-4-tert.-butoxycarbonylamino-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester.

20 Prepared from 4-(4-tert.-butoxycarbonylamino-3-nitro-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (Example E1) (256 mg, 0.61 mmol) by reduction with Zn/NH₄Cl according to the general procedure G (method c). Obtained as an orange foam (75 mg).

MS (EI) 389 (M⁺).

25 Example G30

[2-Amino-4-(6-benzyloxy-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl ester

Prepared from [4-(6-benzyloxy-pyridin-3-yl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example E3) (768 mg, 1.82 mmol) by reduction with Zn/NH₄Cl according to the general procedure G (method c). Obtained as a light brown solid (678 mg).

MS (ISP) 392 [(M+H)⁺]; mp 176-177 °C.

5 Example G31

[2-Amino-4-(6-oxo-1,6-dihydro-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(6-benzyloxy-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl ester (Example G30) (160 mg, 0.409 mmol) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as an off-white solid (136 mg).

10 MS (ISP) 302 [(M+H)⁺]; mp 120-124 °C (dec.).

Example G32

(3''-Amino-[1,1';4',1'']terphenyl-4''-yl)-carbamic acid tert.-butyl ester.

Prepared from (3''-nitro-[1,1';4',1'']terphenyl-4''-yl)-carbamic acid tert.-butyl ester (Example C12) (160 mg, 0.409 mmol) by catalytic hydrogenation with Pd/C according to
15 the general procedure G (method a). Obtained as a beige solid (175 mg).

MS (ISP) 361 [(M+H)⁺]; mp 206-207 °C.

Example G33

[2-Amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [4-(4-fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester
20 (Example F9) (10.0 g, 28.1 mmol) by reduction with SnCl₂·2H₂O (31.7 g, 140.5 mmol) according to the general procedure G (method b). Obtained as a yellow solid (7.08 g).

MS (ISP) 327 [(M+H)⁺]; mp 180 °C.

Example G34

[2-Amino-4-(2-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [4-(2-fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example F10) (713 mg, 2 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.26 g, 10 mmol) according to the general procedure G (method b). Obtained as an orange solid (757 mg).

MS (ISP) 327 $[(\text{M}+\text{H})^+]$; mp 137-139 °C.

5 Example G35

[2-Amino-4-(2,4-difluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [4-(2,4-difluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example F11) (750 mg, 2 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.26 g, 10 mmol) according to the general procedure G (method b). Obtained as a brown solid (688 mg).

10 MS (ISP) 345 $[(\text{M}+\text{H})^+]$; mp 113-116 °C.

Example G36

[2-Amino-4-(4-trifluoromethoxy-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [2-nitro-4-(4-trifluoromethoxy-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example F12) (1.09 g, 2.58 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.91 g, 12.9 mmol) according to the general procedure G (method b). Obtained as a light brown solid (690 mg).

MS (ISP) 393 $[(\text{M}+\text{H})^+]$; mp 167-169 °C.

Example G37

(3-Amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

20 Prepared from (2'-fluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B10) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a yellow solid (1.31 g).

MS (ISP) 303 $[(\text{M}+\text{H})^+]$; mp 100-103 °C.

Example G38

25 (3-Amino-3'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (3'-fluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B11) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a brown solid (3.40 g).

MS (ISP) 303 [(M+H)⁺]; mp 125-128 °C.

5 Example G39

(3-Amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-fluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B12) by catalytic hydrogenation with Raney-Nickel according to the general procedure G (method a). Obtained as a light yellow solid (3.40 g).

10 MS (ISP) 303 [(M+H)⁺].

Example G40

(3-Amino-4'-cyano-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-cyano-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example C13) by reduction with SnCl₂·2H₂O according to the general procedure G (method b).

15 Obtained as a yellow solid (360 mg).

MS (ISP) 310 [(M+H)⁺]; mp 195 °C (dec.).

Example G41

(3-Amino-3'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-fluoro-2'-methyl-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B13) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a light yellow solid (1.51 g).

20

MS (ISP) 317 [(M+H)⁺]; mp 143 °C.

Example G42

(3-Amino-4'-fluoro-2'-methoxymethoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (4'-fluoro-2'-methoxymethoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B14) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a light purple foam (577 mg).

MS (ISP) 363 [(M+H)⁺].

5 Example G43

(3-Amino-2',4'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2',4'-difluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B15) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a light yellow solid (1.77 g).

10 MS (ISP) 321 [(M+H)⁺]; mp 120 °C (dec.).

Example G44

(3-Amino-2'-fluoro-6'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2'-fluoro-6'-methoxy-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B16) by catalytic hydrogenation with Pd/C according to the general
15 procedure G (method a). Obtained as a light purple foam (577 mg).

MS (ISP) 333 [(M+H)⁺]; mp 165-167 °C.

Example G45

(3-Amino-2',5'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2',5'-difluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester
20 (Example B17) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a light yellow gum (2.18 g).

MS (ISN) 319 [(M-H)⁻].

Example G46

(2-Amino-4-benzofuran-2-yl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-benzofuran-2-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B18) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure G (method b). Obtained as an orange solid (579 mg).

MS (ISN) 323 $[(\text{M}-\text{H})^-]$; mp 165 °C.

5 Example G47

(2-Amino-4-isopropyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-isopropyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A2) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a white solid (12.59 g).

10 MS (ISP) 251 $[(\text{M}+\text{H})^+]$; mp 100-101 °C.

Example G48

(2-Amino-4-cyclopropyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (4-cyclopropyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example A3) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure G (method b).

15 Obtained as a dark solid (1.96 g).

MS (ISP) 249 $[(\text{M}+\text{H})^+]$.

Example G49

(3-Amino-2',3'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester.

Prepared from (2',3'-difluoro-3-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example B19) (3.14 g, 8.96 mmol) by catalytic hydrogenation with Pd/C according to the general procedure G (method a). Obtained as a green solid (2.91 g).

20 MS (ISP) 321 $[(\text{M}+\text{H})^+]$; mp 78 °C.

General procedure H (Synthetic Scheme F)

Method a) Preparation of ethyl or tert.-butyl 3-aryl-3-oxo-propionates

25 The ethyl or tert.-butyl 3-aryl-3-oxo-propionates were prepared from the aryl acid chlorides and ethyl or tert.-butyl malonate potassium salt [CAS-no. 6148-64-7 and 75486-

33-8] with Et_3N and MgCl_2 in CH_3CN at $0\text{ }^\circ\text{C}$ to $23\text{ }^\circ\text{C}$ according to *Synthesis* 1993, 290. If the free aryl carboxylic acid was employed in this reaction, it was activated by treatment with ethyl chloroformate and Et_3N in $\text{THF}/\text{CH}_3\text{CN}$ at $0\text{ }^\circ\text{C}$ prior to reaction with the malonate salt.

5 Method b) Preparation of tert.-butyl 3-aryl-3-oxo-propionates

The tert.-butyl 3-aryl-3-oxo-propionates were alternatively prepared from the methyl or ethyl aryl esters by treatment with lithium tert.-butyl acetate [prepared by treatment of tert.-butyl acetate with lithium diisopropylamide in THF at $-78\text{ }^\circ\text{C}$] in the presence of lithium tert.-butoxide according to *Synthesis* 1985, 45. If the product contained residual
10 starting material after workup, thus could be removed by selective saponification with LiOH in $\text{THF}/\text{MeOH}/\text{H}_2\text{O}$ at $23\text{ }^\circ\text{C}$.

Method c) Preparation of 3-aryl-3-oxo-propionic acids

The 3-aryl-3-oxo-propionic acids were prepared from the aryl acid chlorides and bis(trimethylsilyl)malonate with Et_3N and LiBr in CH_3CN at $0\text{ }^\circ\text{C}$ according to *Synth.*
15 *Commun.* 1985, 15, 1039 (method c1) or with $n\text{-BuLi}$ in ether at $-60\text{ }^\circ\text{C}$ to $0\text{ }^\circ\text{C}$ according to *Synthesis* 1979, 787 (method c2).

Example H1

3-Oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionic acid ethyl ester.

Prepared from 3-[1,2,4]triazol-4-yl-benzoic acid [prepared by reaction of 3-aminobenzoic
20 acid with hydrazine hydrate and triethyl orthoformate in acetic acid at $120\text{ }^\circ\text{C}$] by activation with ethyl chloroformate/ Et_3N and reaction with ethyl malonate potassium salt with Et_3N and MgCl_2 in CH_3CN according to general procedure H (method a). Obtained as a white solid (5.74 g).

MS (EI) 259 (M^+).

25 Example H2

3-Oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester.

Prepared from 3-[1,2,3]triazol-1-yl-benzoic acid [prepared by refluxing of methyl 3-azidobenzoate [CAS-No. 93066-93-4] in trimethylsilylacetylene, followed by saponification with aqueous NaOH in refluxing EtOH] by activation with ethyl chloroformate/ Et_3N and

reaction with ethyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure H (method a). Obtained as a light yellow solid (2.22 g).

MS (EI) 259 (M⁺); mp 72-74 °C.

Example H3

5 3-(3-Cyano-phenyl)-3-oxo-propionic acid tert-butyl ester.

Prepared from methyl 3-cyanobenzoate [CAS-No. 13531-48-1] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a light brown oily semisolid.

MS (EI) 245 (M⁺).

10 Example H4

3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionic acid tert.-butyl ester.

Prepared from methyl 3-(1H-imidazol-1-yl)benzoate [prepared from 3-(1H-imidazol-1-yl)benzoic acid (*J. Med. Chem.* 1987, 30, 1342; CAS-No. [108035-47-8] by refluxing in conc. H₂SO₄/MeOH] by treatment with lithium tert.-butyl acetate according to general
15 procedure H (method b). Obtained as an orange-brown oil.

MS (ISP) 287 [(M+H)⁺].

Example H5

3-(2-Imidazol-1-yl-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester.

Prepared from 2-imidazol-1-yl-isonicotinoyl chloride hydrochloride [prepared by reaction
20 of tert.-butyl 2-chloroisonicotinoate with imidazole and NaH in DMF at 80 °C, treatment with formic acid at 50 °C and reaction with thionylchloride in toluene at 100 °C] and tert.-butyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure H (method a). Obtained as a brown solid (10.8 g).

MS (EI) 287 (M⁺); mp 80 °C (dec.).

25 Example H6

3-Oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionic acid tert.-butyl ester.

Prepared from methyl 3-[1,2,4]triazol-1-yl-benzoate [CAS-No. 167626-27-9] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as an orange liquid (2.41 g).

MS (EI) 287 (M^+).

5 Example H7

3-[3-(4-Methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from methyl 3-(4-methyl-imidazol-1-yl)-benzoate [prepared the corresponding acid from 3-isothiocyanatobenzoic acid and 2-aminopropionaldehyde dimethyl acetal according to *J. Med. Chem.* 1987, 30, 1342, followed by refluxing in conc. $H_2SO_4/MeOH$] by treatment with lithium tert.-butyl acetate according to general procedure H (method b).
10 Obtained as a yellow-brown oil (10.69 g).

MS (EI) 300 (M^+).

Example H8

3-[3-(2-Methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

15 Prepared from ethyl 3-(2-methyl-imidazol-1-yl)-benzoate [prepared by reaction of ethyl 3-aminobenzoate with ethyl acetimidate hydrochloride in EtOH at 0 °C, direct treatment with aminoacetaldehyde diethyl acetal in EtOH at 23 °C, followed by addition of conc. H_2SO_4 and refluxing.] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a brown oil (9.66 g).

20 MS (ISN) 299 [$(M-H)^+$].

Example H9

3-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from ethyl 3-(2,4-dimethyl-imidazol-1-yl)-benzoate [prepared by reaction of ethyl 3-aminobenzoate with ethyl acetimidate hydrochloride in EtOH at 0 °C, direct
25 treatment with 2-aminopropionaldehyde dimethyl acetal in EtOH at 23 °C, followed by addition of conc. H_2SO_4 and refluxing.] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a yellow-brown oil (6.00 g).

MS (ISN) 313 [$(M-H)^+$].

Example H10

3-(2-Cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester.

Prepared from 2-cyano-isonicotinic acid ethyl ester [CAS-No. 58481-14-4] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as
5 a light brown solid (7.70 g).

MS (ISN) 245 [(M-H)⁻].

Example H11

3-Oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionic acid tert.-butyl ester.

Prepared from methyl 3-[1,2,4]triazol-4-yl-benzoate [prepared by reaction of 3-aminobenzoic acid with hydrazine hydrate and triethyl orthoformate in acetic acid at 120
10 °C, followed by esterification with conc. H₂SO₄ in refluxing MeOH] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a light yellow gum (870 mg).

MS (ISN) 286 [(M-H)⁻].

15 Example H12

3-[3-(2-Methoxymethylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from ethyl 3-(2-methoxymethylsulfanyl-imidazol-1-yl)-benzoate [prepared by esterification of 3-(2-methoxymethylsulfanyl-imidazol-1-yl)-benzoic acid [CAS-No.
20 108035-46-7] with conc. H₂SO₄ in EtOH, followed by treatment with chloromethylmethyl ether and NaH in THF/DMF] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as an orange oil (1.82 g).

MS (EI) 362 (M⁺).

Example H13

25 3-[3-(2-Methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from ethyl 3-(2-methylsulfanyl-imidazol-1-yl)-benzoate [prepared by esterification of 3-(2-methoxymethylsulfanyl-imidazol-1-yl)-benzoic acid [CAS-No. 108035-46-7] with conc. H₂SO₄ in EtOH, followed by treatment methyl iodide and NaH in

THF/DMF] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a light brown oil (4.41 g).

MS (ISP) 333 [(M+H)⁺].

Example H14

- 5 3-[3-(3-Methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from ethyl 3-(3-methyl-isoxazol-5-yl)-benzoate [prepared by reaction of ethyl 3-ethynylbenzoate [CAS-No. 178742-95-5] with a mixture of NCS, acetaldoxime, Et₃N and cat. amount of pyridine in CHCl₃ at 50 °C according to *Tetrahedron* 1984, 40, 2985-2988] by treatment with lithium tert.-butyl acetate according to general procedure H (method b).

- 10 Obtained as a yellow solid (2.54 g).

MS (ISP) 302 [(M+H)⁺]; mp 50-56 °C.

Example H15

3-Oxo-3-(3-tetrazol-1-yl-phenyl)-propionic acid ethyl ester.

- 15 Prepared from 3-tetrazol-1-yl-benzoic acid [CAS-No. 204196-80-5] by activation with ethyl chloroformate/Et₃N and reaction with ethyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure H (method a). Obtained as a light yellow solid (211 mg).

MS (EI) 260 (M⁺).

Example H16

- 20 3-(3-Chloro-thiophen-2-yl)-3-oxo-propionic acid ethyl ester

Prepared from 3-chloro-2-thiophenecarbonyl chloride [CAS-No. 86427-02-3] by reaction with ethyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure H (method a). Obtained as a brown oil (6.84 g).

MS (EI) 232 (M⁺) and 234 [(M+2)⁺].

- 25 Example H17

3-(5-Cyano-thiophen-2-yl)-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 5-cyano-2-thiophenecarboxylate [CAS-No. 67808-35-9] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a yellow solid (6.66 g).

MS (EI) 251 (M^+); mp 78 °C.

5 Example H18

3-(5-Cyano-2-fluoro-phenyl)-3-oxo-propionic acid ethyl ester

Prepared from 5-cyano-2-fluoro-benzoyl chloride [prepared from the corresponding acid [CAS-No. 146328-87-2] by treatment with SOCl_2 , cat. DMF in toluene at 80 °C] by reaction with ethyl malonate potassium salt with Et_3N and MgCl_2 in CH_3CN according to
10 general procedure H (method a). Obtained as a light yellow solid (3.85 g).

MS (EI) 235 (M^+); mp 55-60 °C.

Example H19

3-(2-Imidazol-1-yl-thiazol-4-yl)-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 2-imidazol-1-yl-thiazole-4-carboxylate [CAS-No. 256420-32-3] by
15 treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as an orange oil (12.0 g).

Example H20

3-[2-(4-Methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 2-(4-methyl-imidazol-1-yl)-thiazole-4-carboxylate [prepared from
20 ethyl 2-amino-4-thiazolecarboxylate (CAS-No. [256420-32-3]) by the following synthetic sequence: 1.) NaH , 2-isothiocyanato-1,1-dimethoxy-propane, DMF, 23 °C; 2.) aq. H_2SO_4 , reflux; 3.) EtOH , conc. H_2SO_4 , 23 °C; 4.) 30% H_2O_2 , HOAc , 23 °C] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a brown oil (8.73 g).

25 MS (EI) 307 (M^+).

Example H21

3-[3-(1-Methyl-1H-imidazol-2-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 3-(1-methyl-1H-imidazol-2-yl)benzoate [CAS-No. 168422-44-4] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Obtained as a light yellow liquid (1.26 g).

MS (ISP) 301.3 [(M+H)⁺].

5 General procedure J (Synthetic Scheme F)

Preparation of 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones

Method a)

The 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones were prepared from 3-aryl-3-oxo-propionic acids and catalytic amount of conc. H₂SO₄ or trifluoroacetic acid (TFA) in isopropenyl acetate at 23 °C according to *Chem. Pharm. Bull.* 1983, 31, 1896. The final products were
10 purified by silica gel column chromatography with hexane/EtOAc.

Method b)

The 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones were prepared from the tert.-butyl 3-aryl-3-oxo-propionates by treatment with trifluoroacetic anhydride (TFAA) in a mixture of TFA
15 and acetone at 23 °C according to *Tetrahedron Lett.* 1998, 39, 2253. The final products were if necessary purified by silica gel column chromatography with hexane/EtOAc.

Example J1

2,2-Dimethyl-6-thiophen-2-yl-[1,3]dioxin-4-one.

The 3-oxo-3-thiophen-2-yl-propionic acid was prepared from thiophene-2-carbonyl chloride (5.3 mL, 50 mmol) and bis(trimethylsilyl)malonate (25.6 mL, 100 mmol) with n-BuLi (1.6M in hexane, 62.5 mL) in ether at -60 °C to 0 °C according to the general
20 procedure H (method c2). The crude material (7.88 g) was transformed into the title compound by stirring in isopropenyl acetate and TFA according to the general procedure J (method a). Obtained as a yellow solid (4.09 g).

25 MS (EI) 210 (M⁺); mp 42 °C (dec.).

Example J2

6-(3-Chloro-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-chloro-thiophen-2-yl)-3-oxo-propionic acid was prepared from 3-chloro-thiophene-2-carbonyl chloride (7.82 g, 43.2 mmol) and bis(trimethylsilyl)malonate (11.6 mL, 45.4 mmol) with Et₃N (12.65 mL, 90.7 mmol) and LiBr (3.53 g, 47.5 mmol) in CH₃CN at 0 °C according to general procedure H (method c1). The crude material (5.69 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as an orange solid (2.3 g).

MS (EI) 244 (M⁺) and 246 [(M+2)⁺]; mp 88-89 °C (dec.).

Example J3

6-(3-Cyano-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one.

10 The 3-(3-cyano-thiophen-2-yl)-3-oxo-propionic acid was prepared from 3-cyano-thiophene-2-carbonyl chloride (24.33 g, 140.6 mmol) and bis(trimethylsilyl)malonate (38.0 mL, 147.7 mmol) with Et₃N (41 mL, 295.4 mmol) and LiBr (13.5 g, 154.7 mmol) in CH₃CN at 0 °C according to general procedure H (method c1). The crude material (24.8 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as an orange solid (5.6 g).

MS (EI) 235 (M⁺); mp 116-120 °C (dec.).

Example J4

3-(2,2-Dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile.

20 The 3-(3-cyano-phenyl)-3-oxo-propionic acid was prepared from 3-cyanobenzoyl chloride (828 mg, 5 mmol) and bis(trimethylsilyl)malonate (2.56 mL, 10 mmol) with n-BuLi (1.6M in hexane, 6.25 mL) in ether at -60 °C to 0 °C according to general procedure H (method c2). The crude material (1.04 g) was transformed into the title compound by stirring in isopropenyl acetate and TFA according to general procedure J (method a). Obtained as a light yellow solid (0.8 g).

25 MS (EI) 229 (M⁺); mp 138 °C (dec.).

Example J5

2,2-Dimethyl-6-(3-trifluoromethyl-phenyl)-[1,3]dioxin-4-one.

The 3-oxo-3-(3-trifluoromethyl-phenyl)-propionic acid was prepared from 3-trifluoromethylbenzoyl chloride (10 mL, 67.6 mmol) and bis(trimethylsilyl)malonate (18.2

mL, 71 mmol) with Et₃N (20 mL, 142 mmol) and LiBr (6.46 g, 74.4 mmol) in CH₃CN at 0 °C according to general procedure H (method c1). The crude material (7.0 g of the obtained 15.4 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as a light
5 yellow solid (5.3 g).

MS (EI) 272 (M⁺); mp 77-78 °C (dec.).

Example J6

6-(3-Chloro-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-chloro-phenyl)-3-oxo-propionic acid was prepared from 3-chlorobenzoyl
10 chloride (11 mL, 85.7 mmol) and bis(trimethylsilyl)malonate (23.0 mL, 90.0 mmol) with Et₃N (25 mL, 180 mmol) and LiBr (8.19 g, 94.3 mmol) in CH₃CN at 0 °C according to general procedure H (method c1). The crude material (17.1 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as a yellow-brown solid (8.0 g).

15 MS (EI) 238 (M⁺) and 240 [(M+2)⁺]; mp 87-88 °C (dec.).

Example J7

6-(3-Iodo-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-iodo-phenyl)-3-oxo-propionic acid was prepared from 3-iodobenzoyl chloride (21.0 g, 78.8 mmol) and bis(trimethylsilyl)malonate (21.0 mL, 82.8 mmol) with Et₃N (23
20 mL, 165.5 mmol) and LiBr (7.54 g, 86.7 mmol) in CH₃CN at 0 °C according to general procedure H (method c1). The crude material (21.9 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as a yellow solid (9.6 g).

MS (EI) 330 (M⁺); mp 79-80 °C (dec.).

25 Example J8

2,2-Dimethyl-6-(3-trifluoromethoxy-phenyl)-[1,3]dioxin-4-one.

The 3-oxo-3-(3-trifluoromethoxy-phenyl)-propionic acid was prepared from 3-trifluoromethoxybenzoyl chloride and bis(trimethylsilyl)malonate with Et₃N and LiBr in CH₃CN at 0 °C according to general procedure H (method c1). The crude material was

transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as an orange solid (2.27 g).

MS (EI) 288 (M⁺); mp 49-54 °C (dec.).

Example J9

5 2,2-Dimethyl-6-pyridin-4-yl-[1,3]dioxin-4-one.

Prepared from 3-oxo-3-pyridin-4-yl-propionic acid [prepared from 4-acetylpyridine, magnesium methylcarbonate and CO₂ in DMF at 120 °C according to *Journal of Antibiotics* 1978, 31, 1245] by treatment with acetone, TFA and TFAA according to general procedure J (method b). Obtained as a white solid (1.3 g).

10 MS (EI) 205 (M⁺)

Example J10

6-(3-Imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-imidazol-1-yl-phenyl)-3-oxo-propionic acid was prepared from 3-(1H-imidazol-1-yl)benzoyl chloride hydrochloride [prepared by treatment of 3-(1H-imidazol-1-yl)benzoic acid (*J. Med. Chem.* 1987, 30, 1342; CAS-No. [108035-47-8] with SOCl₂) and
15 bis(trimethylsilyl)malonate with Et₃N and LiBr in CH₃CN at 0 °C according to general procedure H (method c1). The crude material was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as an orange semisolid (617 g).

20 MS (EI) 270 (M⁺).

Example J11

2,2-Dimethyl-6-(3-methoxy-phenyl)-[1,3]dioxin-4-one.

The 3-(3-methoxy-phenyl)-3-oxo-propionic acid was prepared from 3-methoxybenzoyl chloride (10.3 g, 60.4 mmol) and bis(trimethylsilyl)malonate (16.2 mL, 63.4 mmol) with
25 Et₃N (17.7 mL, 127 mmol) and LiBr (5.77 g, 66.4 mmol) in CH₃CN at 0 °C according to general procedure H (method c1). The crude material (6.38 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure J (method a). Obtained as a yellow oil (640 mg).

MS (ISP) 235 [(M+H)⁺] and 252 [(M+NH₄)⁺].

Example J12

2,2-Dimethyl-6-(3-nitro-phenyl)-[1,3]dioxin-4-one.

The 3-(3-nitro-phenyl)-3-oxo-propionic acid tert.-butyl ester was prepared from 3-nitrobenzoyl chloride (2.71 g, 14.6 mmol) and tert.-butyl malonate potassium salt (6.0 g, 30.0 mmol) with Et₃N (4.5 mL, 32.2 mmol) and MgCl₂ (3.48 g, 36.52 mmol) in CH₃CN according to general procedure H (method a). The crude material (3.88 g) was transformed into the title compound by stirring in TFA/acetone with TFAA according to general procedure J (method b). Obtained as a yellow solid (2.76 g).

10 MS (EI) 249 (M⁺); mp 110-117 °C.

Example J13

2,2-Dimethyl-6-(3-[1,2,4]triazol-1-yl-phenyl)-[1,3]dioxin-4-one.

The 3-oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionic acid tert.-butyl ester was prepared from 3-[1,2,4]triazol-1-yl-benzoic acid methyl ester [CAS-No. 167626-27-9] by treatment with lithium tert.-butyl acetate according to general procedure H (method b). Prepared from (Example H6) by stirring in TFA/acetone with TFAA according to general procedure J (method b). Obtained as a yellow solid (539 mg).

MS (EI) 271 (M⁺).

Example J14

20 6-(2-Imidazol-1-yl-pyridin-4-yl)-2,2-dimethyl-[1,3]dioxin-4-one.

Prepared from 3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H5) by stirring in TFA/acetone with TFAA according to general procedure J (method b). Obtained as a brown solid (10.8 g).

MS (EI) 271 (M⁺); mp 151 °C (dec.).

25 Example J15

2,2-Dimethyl-6-[3-(2-methyl-imidazol-1-yl)-phenyl]-[1,3]dioxin-4-one.

Prepared from 3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H8) by stirring in TFA/acetone with TFAA according to general procedure J (method b). Obtained as a beige solid (2.13 g).

MS (EI) 284 (M^+); mp 122 °C.

5 Example J16

4-(2,2-Dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-pyridine-2-carbonitrile.

Prepared from 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H10) by stirring in TFA/acetone with TFAA according to general procedure J (method b). Obtained as a brown solid (3.30 g).

10 MS (EI) 230 (M^+); mp 132 °C (dec.).

General procedure K (Synthetic Scheme B)

Preparation of {2-[3-aryl-3-oxo-propionylamino]-4-aryl-phenyl}-carbamic acid tert.-butyl ester by reaction of (2-amino-4-aryl-phenyl)-carbamic acid tert.-butyl esters with ethyl or
15 tert.-butyl 3-aryl-3-oxo-propionates or 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones; also 3-aryl-N-(2-nitro-4-aryl-phenyl)-3-oxo-propionamides by reaction of 2-nitro-4-aryl-phenylamines with 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones.

A mixture of the (2-amino-4-aryl-phenyl)-carbamic acid tert.-butyl ester or 2-nitro-4-aryl-phenylamine (1.0 mmol) and the ethyl or tert.-butyl 3-aryl-3-oxo-propionate or 6-aryl-
20 2,2-dimethyl-[1,3]dioxin-4-one (0.8-1.5 mmol) was refluxed in toluene (4-8 mL) until thin layer chromatography indicated complete consumption of the minor component. The solution was allowed to cool to 23 °C, whereupon the product generally crystallized (in cases where crystallization failed to appear it was induced by addition of hexane or the entire reaction mixture was directly subjected to chromatography). The solid was filtered
25 off, washed with ether or mixtures of ether/hexane and dried in vacuum to give the {2-[3-aryl-3-oxo-propionylamino]-4-aryl-phenyl}-carbamic acid tert.-butyl esters or 3-aryl-N-(2-nitro-4-aryl-phenyl)-3-oxo-propionamides, which was used directly in the following step or - if necessary - was purified by recrystallization or by silica gel column chromatography.

Example K1

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-iodo-phenyl}-carbamic acid tert.-butyl ester.

Prepared from (2-amino-4-iodo-phenyl)-carbamic acid tert.-butyl ester (Example G1)
5 (900 mg, 2.7 mmol) and 3-(3-cyano-phenyl)-3-oxo-propionic acid ethyl ester (880 mg, 4.1 mmol) according to the general procedure K. Obtained as a yellow solid (1.2 g).

MS (ISP) 506 [(M+H)⁺] and 528 [(M+Na)⁺]; mp 182-183 °C.

Example K2

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid
10 tert.-butyl ester.

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) (214 mg, 0.7 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (250 mg, 1.1 mmol) according to the general procedure K. Obtained as a light yellow solid (260 mg).

15 MS (ISP) 480 [(M+H)⁺], 497 [(M+NH₄)⁺] and 502 [(M+Na)⁺]; mp 168-170 °C (dec.).

Example K3

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4'-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester.

Prepared from (3-amino-4'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester
20 (Example G5) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a light brown solid (207 mg).

MS (ISP) 486 [(M+H)⁺], 508 [(M+Na)⁺] and 524 [(M+K)⁺]

Example K4

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-thiophen-3-yl-phenyl}-carbamic acid
25 tert.-butyl ester.

Prepared from (2-amino-4-thiophen-3-yl-phenyl)-carbamic acid tert.-butyl ester (Example G6) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4)

according to the general procedure K. Obtained as a brown solid (104 mg) and used crude in the next step (Example 7).

Example K5

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-furan-2-yl-phenyl}-carbamic acid
5 tert.-butyl ester.

Prepared from (2-amino-4-furan-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example G7) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a beige powder (271 mg).

MS (ISP) 446 [(M+H)⁺], 468 [(M+Na)⁺] and 484 [(M+K)⁺]

10 Example K6

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl
ester.

Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (250
mg, 0.88 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example
15 J4) (243 mg, 1.06 mmol) according to the general procedure K. Obtained as a white solid
(324 mg).

MS (ISP) 456 [(M+H)⁺]; mp 168 °C (dec.)

Example K7

{2-[3-(3-Iodo-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid
20 tert.-butyl ester.

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester
(Example G2) (1.0 g, 3.24 mmol) and 6-(3-iodo-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one
(Example J7) (1.78 g, 3.57 mmol) according to the general procedure K. Obtained as a light
yellow solid (1.9 g).

25 MS (ISP) 581 [(M+H)⁺] and 603 [(M+Na)⁺]; mp 193-195 °C (dec.)

Example K8

{3-[3-(3-Azido-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl
ester.

Prepared (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (569 mg, 2 mmol) and 3-(3-azido-phenyl)-3-oxo-propionic acid ethyl ester (700 mg, 3 mmol; prepared from 3-azido-benzoyl chloride (*Bioorg. Chem* 1986, 134) using the procedure described in *Synthesis*, 1993, 290, method A; MS (EI) 233 (M^+)) according to the general
5 procedure K. Obtained as an orange solid (367 mg).

MS (ISP) 472 [$(M+H)^+$] and 494 [$(M+Na)^+$]

Example K8

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2',3'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (3-amino-2',3'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G49) (160 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (115 mg, 0.5 mmol) according to the general procedure K. Obtained as a white solid (53 mg).

MS (ISP) 492 [$(M+H)^+$]; mp 118 °C.

15 Example K9

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-furan-3-yl-phenyl}-carbamic acid tert.-butyl ester.

- Prepared from (2-amino-4-furan-3-yl-phenyl)-carbamic acid tert.-butyl ester (Example G11) and 3-(3-cyano-phenyl)-3-oxo-propionic acid ethyl ester (*Pol. J. Chem.* 1978, 25)
20 according to the general procedure K. Obtained as an orange solid (460 mg).

MS (ISP) 446 [$(M+H)^+$], 463 [$(M+NH_4)^+$] and 468 [$(M+Na)^+$]

Example K10

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-naphthalen-1-yl-phenyl}-carbamic acid tert.-butyl ester.

- 25 Prepared from (2-amino-4-naphthalen-1-yl-phenyl)-carbamic acid tert.-butyl ester (Example G14) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as an off-white solid (92 mg) and used crude in the next step (Example 20).

Example K11

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-thiazol-2-ylethynyl-phenyl}-carbamic acid tert.-butyl ester.

- Prepared from (2-amino-4-thiazol-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester
5 (Example G11) (89 mg, 0.28 mmol) and 3-(3-cyano-phenyl)-3-oxo-propionic acid ethyl ester (*Pol. J. Chem.* 1978, 25) (74 mg, 0.34 mmol) according to the general procedure K. Obtained as a light yellow solid (129 mg).

MS (ISP) 487 [(M+H)⁺]; mp 131 °C

Example K12

- 10 {2-[3-(3-Chloro-thiophen-2-yl)-3-oxo-propionylamino]-4-pyridin-2-yl-phenyl}-carbamic acid tert.-butyl ester.

Prepared from (2-amino-4-pyridin-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example G17) and 6-(3-chloro-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J2) according to the general procedure K. Obtained as an amorphous brown solid (145 mg).

- 15 MS (ISP) 472 [(M+H)⁺] and 494 [(M+Na)⁺]

Example K13

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-pyridin-2-yl-phenyl}-carbamic acid tert.-butyl ester.

- Prepared from (2-amino-4-pyridin-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example
20 G17) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a light brown solid (90 mg).

MS (ISP) 457 [(M+H)⁺] and 479 [(M+Na)⁺].

Example K14

- [3-(3-Oxo-3-thiophen-3-yl-propionylamino)-biphenyl-4-yl]-carbamic acid tert.-butyl
25 ester

Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (144 mg, 0.51 mmol) and 3-oxo-3-thiophen-3-yl-propionic acid ethyl ester (FR 7191887)

(151 mg, 0.76 mmol) according to the general procedure K. Obtained as a yellow foam (181 mg).

MS (ISN) 435 [(M+H)⁺].

Example K15

- 5 {3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-3'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester.

Prepared from (3-amino-3'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G20) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a viscous orange oil (290 mg).

- 10 MS (ISP) 470 [(M+H)⁺], 492 [(M+Na)⁺] and 508 [(M+K)⁺].

Example K16

{3-[3-(3-Chloro-thiophen-2-yl)-3-oxo-propionylamino]-2'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester.

- 15 Prepared from (3-amino-2'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G23) and 6-(3-chloro-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J2) according to the general procedure K. Obtained as a viscous brown oil (154 mg).

MS (ISP) 485 [(M+H)⁺], 507 [(M+Na)⁺] and 523 [(M+K)⁺].

Example K17

- 20 {4-Benzoyl-2-[3-(3-chloro-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester.

Prepared from (2-amino-4-benzoyl-phenyl)-carbamic acid tert.-butyl ester (Example G24) (205 mg, 0.66 mmol) and 6-(3-chloro-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J6) (174 mg, 0.73 mmol) according to the general procedure K. Obtained as a brown foam (232 mg).

- 25 MS (ISP) 493 [(M+H)⁺].

Example K18

{4-Benzoyl-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester.

- Prepared from (2-amino-4-benzoyl-phenyl)-carbamic acid tert.-butyl ester (Example G24) (375 mg, 1.20 mmol) and 3-(3-cyano-phenyl)-3-oxo-propionic acid ethyl ester (*Pol. J. Chem.* 1978, 25) (313 mg, 1.44 mmol) according to the general procedure K. Obtained as a light yellow solid (170 mg).

MS (ISP) 484 [(M+H)⁺], 501 [(M+NH₄)⁺] and 506 [(M+Na)⁺]; mp 168 °C (dec.)

Example K19

- [4-Benzoyl-2-(3-oxo-3-thiophen-2-yl-propionylamino)-phenyl]-carbamic acid tert.-butyl ester.

- Prepared from (2-amino-4-benzoyl-phenyl)-carbamic acid tert.-butyl ester (Example G24) (259 mg, 0.5 mmol) and 2,2-dimethyl-6-thiophen-2-yl-[1,3]dioxin-4-one (Example J1) (135 mg, 0.55 mmol) according to the general procedure K. Obtained as a light yellow solid (60 mg).

MS (ISP) 465 [(M+H)⁺]

Example K20

{3-[3-(3-Chloro-thiophen-2-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester.

- Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (100 mg, 0.35 mmol) and 6-(3-chloro-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J2) (95 mg, 0.39 mmol) according to the general procedure K. Obtained as a white solid (127 mg).

MS (ISP) 471 [(M+H)⁺]; mp 165 °C.

- Example K21

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-3'-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester.

Prepared from (3-amino-3'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G25) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a beige solid (238 mg).

MS (ISP) 486 [(M+H)⁺], 508 [(M+Na)⁺] and 524 [(M+K)⁺].

5 Example K22

[5-(6-Oxo-1,6-dihydro-pyridin-3-yl)-2-(3-oxo-3-thiophen-2-yl-propionylamino)-phenyl]-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(6-oxo-1,6-dihydro-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl ester (Example G31) (200 mg, 0.664 mmol) and 2,2-dimethyl-6-thiophen-2-yl-[1,3]dioxin-4-one (Example J1) (140 mg, 0.665 mmol) according to the general procedure K. Obtained as a beige solid (235 mg).

MS (ISP) 454 [(M+H)⁺]

Example K23

15 {4-(6-Benzoyloxy-pyridin-3-yl)-2-[3-oxo-3-(3-trifluoromethyl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester.

Prepared from [2-amino-4-(6-benzyloxy-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl ester (Example G30) (203 mg, 0.52 mmol) and 2,2-dimethyl-6-(3-trifluoromethyl-phenyl)-[1,3]dioxin-4-one (Example J5) (150 mg, 0.55 mmol) according to the general procedure K. Obtained as an off-white solid (213 mg).

20 MS (ISP) 606 [(M+H)⁺]; mp 190 °C (dec.).

Example K24

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4'-trifluoromethoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester.

25 Prepared from (3-amino-4'-trifluoromethoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G27) and 3-(3-cyano-phenyl)-3-oxo-propionic acid ethyl ester (*Pol. J. Chem.* 1978, 25) according to the general procedure K. Obtained as a brown semisolid (94 mg).

MS (ISP) 540 [(M+H)⁺], 557 [(M+NH₄)⁺] and 562 [(M+Na)⁺].

Example K25

4-{4-tert.-Butoxycarbonylamino-3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester

Prepared from 4-(3-amino-4-tert.-butoxycarbonylamino-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (Example G29) (544 mg, 1.4 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (336 mg, 1.5 mmol) according to the general procedure K. Obtained as an orange solid (722 mg).

MS (ISP) 561 [(M+H)⁺]; mp 75-79 °C (dec.).

Example K26

10 {4-(6-Benzyloxy-pyridin-3-yl)-2-[3-(3-chloro-thiophen-2-yl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(6-benzyloxy-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl ester (Example G30) (216 mg, 0.55 mmol) and 6-(3-chloro-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J2) (142 mg, 0.58 mmol) according to the general procedure K. Obtained as a beige solid (172 mg).

MS (ISP) 578 [(M+H)⁺]; mp 158-159 °C (dec.).

Example K27

{2-[3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

20 Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) (154 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) (135 mg, 0.5 mmol) according to the general procedure K. Obtained as an orange oil (179 mg).

MS (ISN) 519 [(M-H)⁻].

25 Example K28

{4-Benzofuran-2-yl-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-benzofuran-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example G46) (324 mg, 1.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (252 mg, 1.1 mmol) according to the general procedure K. Obtained as a yellow solid (299 mg).

- 5 MS (ISP) 496 [(M+H)⁺]; mp 115 °C.

Example K29

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-[1,1';4',1'']terphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (3''-amino-[1,1';4',1'']terphenyl-4''-yl)-carbamic acid tert.-butyl ester (Example G32) (159mg, 0.44mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (111 mg, 0.49 mmol) according to the general procedure K. Obtained as an off-white solid (156 mg).

MS (ISN) 530 [(M-H)⁻]; mp 214-216 °C.

Example K30

- 15 [2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-(4-trifluoromethoxy-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester

- 20 Prepared from [2-amino-4-(4-trifluoromethoxy-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G36) (196 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (126 mg, 0.55 mmol) according to the general procedure K. Obtained as an off-white solid (175 mg).

MS (ISP) 564 [(M+H)⁺]; mp 152-154 °C.

Example K31

{2-[3-(3-Nitro-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

- 25 Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) (1.1 g, 4.0 mmol) and 2,2-dimethyl-6-(3-nitro-phenyl)-[1,3]dioxin-4-one (Example J12) (1.23 g, 4.4 mmol) according to the general procedure K. Obtained as a light yellow solid (989 mg).

MS (ISN) 498 [(M-H)⁻]; mp 177-179 °C.

Example K32

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 5 Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a yellow solid (257 mg).

MS (ISP) 474 [(M+H)⁺]; mp 177-179 °C.

Example K33

- 10 {4-(4-Fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G33) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) according to the general procedure K. Obtained as an orange oil (207 mg).

- 15 MS (ISN) 537 [(M-H)⁻].

Example K34

{3-[3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 20 Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (142 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) (170 mg, 0.63 mmol) according to the general procedure K. Obtained as a light yellow foam (248 mg).

MS (ISN) 495 [(M-H)⁻].

Example K35

- 25 {4'-Fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) according to the general procedure K. Obtained as a light yellow solid (489 mg).

MS (ISN) 513 [(M-H)⁺].

5 Example K36

(3-Amino-4'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G5) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) according to the general procedure K. Obtained as a yellow liquid (195 mg).

10 Example K37

{4-(4-Fluoro-phenylethynyl)-2-[3-oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G33) and 2,2-dimethyl-6-(3-[1,2,4]triazol-1-yl-phenyl)-[1,3]dioxin-4-one (Example J13) according to the general procedure K. Obtained as a yellow solid (230 mg).

mp 131-139 °C.

Example K38

{4-(2-Fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

20 Prepared from [2-amino-4-(2-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G34) (245 mg, 0.75 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) (300 mg, 1.1 mmol) according to the general procedure K. Obtained as a yellow-brown foam (211 mg).

MS (ISN) 537 [(M-H)⁺].

25 Example K39

{4'-Fluoro-3-[3-oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 2,2-dimethyl-6-(3-[1,2,4]triazol-1-yl-phenyl)-[1,3]dioxin-4-one (Example J13) according to the general procedure K. Obtained as an orange foam (233 mg).

Example K40

- 5 {4'-Cyano-3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-cyano-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G40) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a light red solid (185 mg).

10 Example K41

{4'-Cyano-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-4'-cyano-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G40) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) according to the general procedure K. Obtained as a yellow solid (228 mg).

Example K42

{4'-Fluoro-3-[3-(3-nitro-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 2,2-dimethyl-6-(3-nitro-phenyl)-[1,3]dioxin-4-one (Example J12) according to the general procedure K. Obtained as a light yellow solid (1.01 g).

Example K43

{4-(4-Fluoro-phenylethynyl)-2-[3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

- 25 Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G33) and 6-(2-imidazol-1-yl-pyridin-4-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J14) according to the general procedure K. Obtained as a brown solid (284 mg).

MS (ISP) 540 [(M+H)⁺]; mp 169 °C.

Example K44

{4'-Fluoro-3-[3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-
carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
5 G39) and 6-(2-imidazol-1-yl-pyridin-4-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J14)
according to the general procedure K. Obtained as a brown solid (361 mg).

MS (ISP) 516 [(M+H)⁺]; mp 124 °C (dec.).

Example K45

{2'-Fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-
10 carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
G37) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10)
according to the general procedure K. Obtained as a brown solid (352 mg).

MS (ISP) 515 [(M+H)⁺]; mp 50-58 °C.

15 Example K46

{4-(6-Benzylloxy-pyridin-3-yl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-
phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(6-benzylloxy-pyridin-3-yl)-phenyl]-carbamic acid tert.-butyl
ester (Example G30) (196 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-
20 [1,3]dioxin-4-one (Example J10) (229 mg, 0.85 mmol) according to the general procedure
K. Obtained as a yellow solid (209 mg).

MS (ISN) 602 [(M-H)⁻]; mp 79-83 °C.

Example K47

{4'-Fluoro-3-[3-[3-(4-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-
25 yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
G39) (362 mg, 1.2 mmol) and 3-[3-(4-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic

acid tert.-butyl ester (Example H7) (300 mg, 1.0 mmol) according to the general procedure K. Obtained as a light yellow solid (392 mg).

MS (ISP) 529 [(M+H)⁺]; mp 124 °C.

Example K48

- 5 {2'-Fluoro-3-[3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) and 6-(2-imidazol-1-yl-pyridin-4-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J14) according to the general procedure K. Obtained as an orange solid (103 mg).

- 10 MS (ISP) 516 [(M+H)⁺].

Example K49

(4-(4-Fluoro-phenylethynyl)-2-{3-[3-(4-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-phenyl)-carbamic acid tert.-butyl ester

- 15 Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G33) (392 mg, 1.2 mmol) and 3-[3-(4-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H7) (300 mg, 1.0 mmol) according to the general procedure K. Obtained as a yellow solid (407 mg).

MS (ISP) 553 [(M+H)⁺]; mp 166 °C.

Example K50

- 20 {4'-Fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-3'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G41) (158 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) (186 mg, 0.69 mmol) according to the general procedure K. Obtained as an
25 orange oil (168 mg).

MS (ISP) 529 [(M+H)⁺].

Example K51

{4'-Fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2'-methoxymethoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-2'-methoxymethoxy-biphenyl-4-yl)-carbamic acid
5 tert.-butyl ester (Example G42) (181 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) (135 mg, 0.5 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (221 mg).

MS (ISN) 573 [(M-H)⁻].

Example K52

10 (2'-Fluoro-3-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (362 mg, 1.2 mmol) and 3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic
acid tert.-butyl ester (Example H8) (300 mg, 1.0 mmol) according to the general procedure
15 K. Obtained as a yellow amorphous substance (312 mg).

MS (ISP) 529 [(M+H)⁺].

Example K53

(4'-Fluoro-3-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

20 Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) (362 mg, 1.2 mmol) and 3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H8) (300 mg, 1.0 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (302 mg).

MS (ISP) 529 [(M+H)⁺].

25 Example K54

{3-[3-(3-Cyano-thiophen-2-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) (302 mg, 1.0 mmol) and 6-(3-cyano-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J3) (250 mg, 1.06 mmol) according to the general procedure K. Obtained as a light yellow solid (251 mg).

- 5 MS (ISP) 480 [(M+H)⁺]; mp 156-157 °C.

Example K55

{3-[3-(3-Cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (302 mg, 1.0 mmol) and 6-(3-cyano-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J3) (280 mg, 1.19 mmol) according to the general procedure K. Obtained as a light yellow solid (446 mg).

MS (ISP) 480 [(M+H)⁺]; mp 63-66 °C.

Example K56

- 15 {4'-Fluoro-3-[3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionic acid ethyl ester (Example H15) according to the general procedure K. Obtained as a light yellow solid (159 mg).

- 20 MS (ISN) 515 [(M-H)⁻].

Example K57

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 25 Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a light yellow foam (239 mg).

MS (ISP) 474 [(M+H)⁺].

Example K58

{2'-Fluoro-3-[3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) and 3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionic acid ethyl ester (Example H15) according to the general procedure K. Obtained as a light red solid (129 mg).

MS (ISP) 517 [(M+H)⁺].

Example K59

(3-{3-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) (227 mg, 0.75 mmol) and 3-[3-(2,4-dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H9) (157 mg, 0.5 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (127 mg).

MS (ISP) 543 [(M+H)⁺].

Example K60

(2-{3-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) (231 mg, 0.75 mmol) and 3-[3-(2,4-dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H9) (157 mg, 0.5 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (140 mg).

MS (ISP) 549 [(M+H)⁺].

Example K61

{3-[3-(3-Chloro-thiophen-2-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) (360 mg, 1.2 mmol) and 3-(3-chloro-thiophen-2-yl)-3-oxo-propionic acid ethyl ester

(Example H16) (350 mg, 1.5 mmol) according to the general procedure K. Obtained as a white-yellow solid (353 mg).

MS (ISP) 489 [(M+H)⁺] and 491 [(M+2+H)⁺]; mp 168-169 °C.

Example K62

- 5 {2'-Fluoro-3-[3-(3-nitro-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) and 2,2-dimethyl-6-(3-nitro-phenyl)-[1,3]dioxin-4-one (Example J12) according to the general procedure K. Obtained as a yellow solid (113 mg).

- 10 MS (ISN) 492 [(M-H)⁻]; mp 167 °C.

Example K63

{3-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 15 Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H10) according to the general procedure K. Obtained as a red solid (118 mg).

Example K64

{3-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 20 Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H10) according to the general procedure K. Obtained as a white solid (151 mg).

mp 190 °C (dec.).

Example K65

- 25 {3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-3'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-3'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G38) (151 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (175 mg, 0.76 mmol) according to the general procedure K. Obtained as an orange solid (141 mg).

- 5 MS (ISP) 474 [(M+H)⁺]; mp 148-150 °C.

Example K66

{3'-Fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (3-amino-3'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G38) (151 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10) (135 mg, 0.5 mmol) according to the general procedure K. Obtained as an orange solid (150 mg).

MS (ISP) 515 [(M+H)⁺]; mp 70-83 °C.

Example K67

- 15 {2-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H10) according to the general procedure K. Obtained as a white solid (174 mg).

- 20 mp 189 °C (dec.).

Example K68

{2'-Fluoro-3-[3-oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 25 Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (302 mg, 1.0 mmol) and 3-oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionic acid tert.-butyl ester (Example H11) (345 mg, 1.2 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (207 mg).

MS (ISP) 516 [(M+H)⁺].

Example K69

{3-[3-(5-Cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (302 mg, 1.0 mmol) and 3-(5-cyano-thiophen-2-yl)-3-oxo-propionic acid tert.-butyl ester (Example H17) (276 mg, 1.1 mmol) according to the general procedure K. Obtained as a yellow solid (451 mg).

MS (ISP) 480 [(M+H)⁺]; mp 201 °C.

Example K70

- 10 (3-[3-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (333 mg, 1.1 mmol) and 3-[3-(2,4-dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H9) (314 mg, 1.0 mmol) according to the general procedure K. Obtained as a light yellow solid (374 mg).

MS (ISP) 543 [(M+H)⁺]; mp 145 °C.

Example K71

(2'-Fluoro-3-{3-[3-(2-methoxymethylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 20 Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (303 mg, 1.0 mmol) and 3-[3-(2-methoxymethylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H12) (512 mg, 1.41 mmol) according to the general procedure K. Obtained as a light yellow solid (552 mg).

MS (ISN) 589 [(M-H)⁻]; mp 83-86 °C.

- 25 Example K72

{3-[3-(5-Cyano-2-fluoro-phenyl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (302 mg, 1.0 mmol) and 3-(5-cyano-2-fluoro-phenyl)-3-oxo-propionic acid ethyl ester (Example H18) (362 mg, 1.5 mmol) according to the general procedure K. Obtained as a yellow-brown solid (352 mg).

- 5 MS (ISP) 492 [(M+H)⁺]; mp 170 °C.

Example K73

{3-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-2',4'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (3-amino-2',4'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G43) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H10) according to the general procedure K. Obtained as a light brown solid (207 mg).

MS (ISN) 491 [(M-H)⁻]; mp 160-161 °C.

Example K74

- 15 {2-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-4-isopropyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-isopropyl-phenyl)-carbamic acid tert.-butyl ester (Example G47) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H10) according to the general procedure K. Obtained as a light brown solid (183 mg).

- 20 MS (ISN) 421 [(M-H)⁻]; mp 163-165 °C.

Example K75

(2'-Fluoro-3-{3-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 25 Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (151 mg, 0.5 mmol) and 3-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionic acid tert.-butyl ester (Example H20) (154 mg, 0.5 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (157 mg).

MS (ISN) 534 [(M-H)⁻].

Example K76

(4'-Fluoro-3-[3-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
5 G39) (151 mg, 0.5 mmol) and 3-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionic acid tert.-butyl ester (Example H20) (154 mg, 0.5 mmol) according to the general procedure K. Obtained as a yellow amorphous substance (225 mg).

MS (ISN) 534 [(M-H)⁻].

Example K77

- 10 {2'-Fluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
G37) (152 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl
ester (Example H2) (200 mg, 0.77 mmol) according to the general procedure K. Obtained
15 as a yellow oil (191 mg).

MS (ISP) 516 [(M+H)⁺].

Example K78

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-cyclopropyl-phenyl}-carbamic acid
tert.-butyl ester

- 20 Prepared from (2-amino-4-cyclopropyl-phenyl)-carbamic acid tert.-butyl ester (Example G48) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a light brown solid (92 mg).

MS (EI) 419 (M⁺).

Example K79

- 25 {2-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-4-cyclopropyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-cyclopropyl-phenyl)-carbamic acid tert.-butyl ester (Example G48) and 4-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-pyridine-2-carbonitrile (Example J16) according to the general procedure K. Obtained as a light brown solid (148 mg).

MS (ISP) 421 [(M+H)⁺].

5 Example K80

{4-Cyclopropyl-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-cyclopropyl-phenyl)-carbamic acid tert.-butyl ester (Example G48) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J10)

10 according to the general procedure K. Obtained as a light yellow solid (79 mg).

MS (ISP) 461 [(M+H)⁺].

Example K81

{4'-Fluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

15 Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) (152 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example H2) (200 mg, 0.77 mmol) according to the general procedure K. Obtained as a yellow oil (107 mg).

MS (ISP) 516 [(M+H)⁺].

20 Example K82

{4-(4-Fluoro-phenylethynyl)-2-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G33) (163 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-

25 propionic acid ethyl ester (Example H2) (181 mg, 0.7 mmol) according to the general procedure K. Obtained as a yellow oil (107 mg).

MS (ISP) 540 [(M+H)⁺].

Example K83

{4'-Fluoro-3-[3-(2-imidazol-1-yl-thiazol-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
5 G39) (151 mg, 0.5 mmol) and 3-(2-imidazol-1-yl-thiazol-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H19) (235 mg, 0.8 mmol) according to the general procedure K. Obtained as an orange oil (162 mg).

MS (ISP) 522 [(M+H)⁺].

Example K84

- 10 (4'-Fluoro-3-[3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
G39) (151 mg, 0.5 mmol) and 3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionic
acid tert.-butyl ester (Example H14) (190 mg, 0.63 mmol) according to the general
15 procedure K. Obtained as an off-white solid (86 mg).

MS (ISP) 530 [(M+H)⁺]; mp 100-101 °C.

Example K85

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2',4'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 20 Prepared from (3-amino-2',4'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G43) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as an orange oil (95 mg).

MS (ISP) 492 [(M+H)⁺].

Example K86

- 25 {2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-isopropyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-isopropyl-phenyl)-carbamic acid tert.-butyl ester (Example G47) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. Obtained as a light red solid (100 mg).

MS (ISP) 422 [(M+H)⁺]; mp 179-180 °C.

5 Example K87

(4'-Fluoro-3-{3-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) and 3-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H13) according to the general procedure K. Obtained as a light yellow oil (181 mg).

MS (ISP) 561 [(M+H)⁺].

Example K88

15 (4-Isopropyl-2-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-phenyl)-carbamic acid tert.-butyl ester

Prepared from (2-amino-4-isopropyl-phenyl)-carbamic acid tert.-butyl ester (Example G47) and 2,2-dimethyl-6-[3-(2-methyl-imidazol-1-yl)-phenyl]-[1,3]dioxin-4-one (Example J15) according to the general procedure K. Obtained as a yellow oil (186 mg).

MS (ISP) 477 [(M+H)⁺].

20 Example K89

(2',4'-Difluoro-3-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (3-amino-2',4'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G43) and 2,2-dimethyl-6-[3-(2-methyl-imidazol-1-yl)-phenyl]-[1,3]dioxin-4-one (Example J15) according to the general procedure K. Obtained as a yellow oil (145 mg).

MS (ISP) 547 [(M+H)⁺].

Example K90

{2',4'-Difluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (3-amino-2',4'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester
5 (Example G43) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example H2) according to the general procedure K. Obtained as a light yellow oil (164 mg).

MS (ISP) 534 [(M+H)⁺].

Example K91

- 10 {2'-Fluoro-3-[3-(2-imidazol-1-yl-thiazol-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (151 mg, 0.5 mmol) and 3-(2-imidazol-1-yl-thiazol-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example H19) (235 mg, 0.8 mmol) according to the general procedure K.

- 15 Obtained as a white solid (98 mg).

MS (ISP) 522 [(M+H)⁺]; mp 115-130 °C.

Example K92

{3-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2',5'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 20 Prepared from (3-amino-2',5'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G45) (160 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (115 mg, 0.5 mmol) according to the general procedure K. Obtained as an amorphous white substance (110 mg).

MS (ISN) 490 [(M-H)⁻].

- 25 Example K93

{2',5'-Difluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2',5'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G45) (160 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example H2) (130 mg, 0.5 mmol) according to the general procedure K. Obtained as an amorphous off-white substance (129 mg).

5 MS (ISN) 532 [(M-H)⁻].

Example K94

(2'-Fluoro-3-{3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (3-amino-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G37) (151 mg, 0.5 mmol) and 3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H14) (190 mg, 0.63 mmol) according to the general procedure K. Obtained as a white solid (213 mg).

MS (ISN) 528 [(M-H)⁻]; mp 158-160 °C.

Example K95

15 {2',3'-Difluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (3-amino-2',3'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G49) (160 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example H2) (130 mg, 0.5 mmol) according to the general procedure K.

20 Obtained as an amorphous yellow substance (166 mg).

MS (ISP) 534 [(M+H)⁺].

Example K96

[2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-(2,4-difluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester

25 Prepared from [2-amino-4-(2,4-difluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G35) (172 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (126 mg, 0.55 mmol) according to the general procedure K. Obtained as an orange oil (198 mg).

MS (ISP) 516 [(M+H)⁺].

Example K97

[2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-(2-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester

- 5 Prepared from [2-amino-4-(2-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G34) (163 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (126 mg, 0.55 mmol) according to the general procedure K. Obtained as a brown solid (143 mg).

MS (ISN) 496 [(M-H)⁻]; mp 216-217 °C.

10 Example K98

[2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester

- Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G33) (245 mg, 0.75 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (190 mg, 0.825 mmol) according to the general procedure K. Obtained as a yellow oil (318 mg).

MS (ISN) 496 [(M-H)⁻].

General procedure M (Synthetic Scheme B)

- Preparation of 4,8-diaryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones, 4-aryl-8-aroylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones or 4-aryl-8-arylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones

- A suspension of the {2-[3-aryl-3-oxo-propionylamino]-4-aryl-phenyl}-carbamic acid tert.-butyl ester or {2-[3-aryl-3-oxo-propionylamino]-4-arylethynyl-phenyl}-carbamic acid tert.-butyl ester (1.0 mmol) in CH₂Cl₂ (5 mL) [anisole or 1,3-dimethoxybenzene (5 to 25 mmol) can be added if necessary] was treated with TFA (0.5-5.0 mL) at 0 °C and stirring was continued at 23 °C until tlc indicated complete consumption of the starting material. The solvent was removed in vacuum, the residue treated with little ether, whereupon it crystallized. The solid was stirred with sat. NaHCO₃-sol., filtered, washed with H₂O and ether or mixtures of ether/hexane and was dried to give the title compound, which if necessary can be purified by crystallization from THF/CH₂Cl₂/ether/hexane.

Example 1

3-(7-Iodo-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-iodo-phenyl}-carbamic acid tert.-butyl ester (Example K1) (1.15 g, 2.3 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (880 mg).

MS (EI) 387 (M⁺); mp 198-200 °C (dec.)

Example 2

3-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example K2) (230 mg, 0.48 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (135 mg).

MS (EI) 361 (M⁺); mp 245 °C (dec.)

Example 3

3-(4-Oxo-7-p-tolyethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- Prepared from (2-amino-4-p-tolyethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G3) (161 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (183 mg, 0.6 mmol) according to the general procedure K. Obtained as a light yellow solid (228 mg). This material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (86 mg).

MS (EI) 375 (M⁺); mp 236-239 °C (dec.).

Example 4

3-[7-(2-Chloro-phenylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from [2-amino-4-(2-chloro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G4) (172 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (230 mg, 0.75 mmol, 75% pure) according to the general procedure K. Obtained as a yellow solid (321 mg). This material was deprotected and

cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (148 mg).

MS (EI) 395 (M^+) and 397 [$(\text{M}+2)^+$]; mp 239-240 °C (dec.).

Example 5

5 4-(3-Chloro-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) and 6-(3-chloro-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J6) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as an
10 orange solid (155 mg).

MS (EI): 370 (M^+)

Example 6

3-[7-(4-Methoxy-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

15 Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4'-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K3) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow powder (122 mg).

MS (ISP) 368 [$(\text{M}+\text{H})^+$]; mp 236-237 °C (dec.).

Example 7

20 3-(4-Oxo-7-thiophen-3-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-thiophen-3-yl-phenyl}-carbamic acid tert.-butyl ester (Example K4) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a beige solid (54 mg).

MS (EI) 343 (M^+); mp 238-243 °C (dec.).

25 Example 8

3-(7-Furan-2-yl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-furan-2-yl-phenyl}-carbamic acid tert.-butyl ester (Example K5) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a brown powder (73 mg).

MS (EI) 327 (M⁺); mp 205-210 °C (dec.).

5 Example 9

3-[7-(4-Ethyl-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from (3-amino-4'-ethyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G8) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by
10 treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a beige solid (67 mg).

MS (EI) 365 (M⁺); mp 225-229 °C (dec.).

Example 10

3-(4-Oxo-7-phenyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

15 Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K6) (268 mg, 0.59 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a white solid (188 mg).

MS (EI) 337 (M⁺); mp 238-240 °C (dec.).

Example 11

20 4-(3-Iodo-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2-[3-(3-iodo-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example K7) (871 mg, 1.5 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (542 mg).

MS (EI) 462 (M⁺); mp 227-229 °C (dec.).

25 Example 12

8-Phenylethynyl-4-pyridin-4-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) and 2,2-dimethyl-6-pyridin-4-yl-[1,3]dioxin-4-one (Example J9) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a brown solid (50 mg).

MS (EI) 337 (M⁺); mp 198-200 °C (dec.).

Example 13

3-[7-(2-Methoxy-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

10 Prepared from (3-amino-2'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G10) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light green powder (30 mg).

15 MS (EI) 367 (M⁺)

Example 14

3-(7-Furan-3-yl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-furan-3-yl-phenyl}-carbamic acid tert.-butyl ester (Example K9) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow powder (73 mg).

MS (EI) 327 (M⁺); mp 228-233 °C (dec.).

Example 15

3-[7-(4-Chloro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from (3-amino-4'-chloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G12) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow powder (79 mg).

MS (EI) 371 (M^+); mp 244-250 °C (dec.).

Example 16

3-[7-(4-Chloro-phenylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 5 Prepared from [2-amino-4-(4-chloro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example G13) (187 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (184 mg, 0.6 mmol) according to the general procedure K. The obtained material (234 mg) was deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow powder (93 mg).
- 10 MS (EI) 395 (M^+) and 397 [$(M+2)^+$]; mp 237-240 °C (dec.).

Example 17

8-Phenyl-4-thiophen-2-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (69 mg, 0.243 mmol) and 2,2-dimethyl-6-thiophen-2-yl-[1,3]dioxin-4-one (Example J1) (54 mg, 0.257 mmol) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a brown solid (46 mg).
- 15

MS (ISP) 319 [$(M+H)^+$].

Example 18

- 20 8-Phenylethynyl-4-(3-trifluoromethyl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) and 2,2-dimethyl-6-(3-trifluoromethoxy-phenyl)-[1,3]dioxin-4-one (Example J8) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M.

- 25 Obtained as a yellow solid (157 mg).

MS (EI) 404 (M^+).

Example 19

4-(3-Chloro-phenyl)-8-phenyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (284 mg, 1.0 mmol) and 6-(3-Chloro-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example J6) (358 mg, 1.2 mmol) according to the general procedure K. The obtained material (339 mg) was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general
5 procedure M. Obtained as a yellow solid (188 mg).

MS (EI) 346 (M⁺) and 348 [(M+2)⁺]; mp 208 °C (dec.).

Example 20

3-(7-Naphthalen-1-yl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-naphthalen-1-yl-phenyl}-carbamic acid tert.-butyl ester (Example K10) by treatment with TFA in CH₂Cl₂
10 according to the general procedure M. Obtained as a light yellow powder (41 mg).

MS (ISP) 388 [(M+H)⁺]; mp 240-245 °C (dec.)

Example 21

3-(4-Oxo-7-thiazol-2-ylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

15 Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-thiazol-2-ylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example K11) (119 mg, 0.24 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (36 mg).

MS (EI) 368 (M⁺); mp 230 °C (dec.).

20 Example 22

3-(4-Oxo-7-p-tolyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from (3-amino-4'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G16) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and
25 cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a white solid (76 mg).

MS (ISP) 352 [(M+H)⁺]; mp 242-245 °C (dec.).

Example 23

4-(3-Chloro-thiophen-2-yl)-8-pyridin-2-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2-[3-(3-chloro-thiophen-2-yl)-3-oxo-propionylamino]-4-pyridin-2-yl-phenyl}-carbamic acid tert.-butyl ester (Example K12) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown powder (51 mg).

MS (EI) 353 (M⁺); mp 220-225 °C (dec.).

Example 24

3-(4-Oxo-7-thiophen-2-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from (2-amino-4-thiophen-2-yl-phenyl)-carbamic acid tert.-butyl ester (Example G18) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (108 mg).

MS (EI) 343 (M⁺); mp >250 °C (dec.).

15 Example 25

8-Iodo-4-thiophen-2-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-amino-4-iodo-phenyl)-carbamic acid tert.-butyl ester (Example G1) and 2,2-dimethyl-6-thiophen-2-yl-[1,3]dioxin-4-one (Example J1) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (30 mg).

MS (EI) 368 (M⁺); mp >260 °C.

Example 26

3-(4-Oxo-7-pyridin-2-ylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from (2-amino-4-pyridin-2-ylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G19) (124 mg, 0.4 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) (147 mg, 0.48 mmol) according to the general procedure K. The obtained material (199 mg) was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (124 mg).

MS (EI) 362 (M^+); mp 229-231 °C (dec.).

Example 27

4-(3-Methoxy-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

- 5 (Example G2) and 2,2-dimethyl-6-(3-methoxy-phenyl)-[1,3]dioxin-4-one (Example J11) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (131 mg).

MS (EI) 366 (M^+).

10 Example 28

3-(4-Oxo-7-pyridin-2-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-pyridin-2-yl-phenyl}-carbamic acid tert.-butyl ester (Example K13) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a brown powder (29 mg).

- 15 MS (EI) 338 (M^+); mp 243-244 °C (dec.).

Example 29

8-Phenyl-4-thiophen-3-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from [3-(3-oxo-3-thiophen-3-yl-propionylamino)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example K14) (110 mg, 0.25 mmol) by treatment with TFA in CH_2Cl_2

- 20 according to the general procedure M. Obtained as a light yellow solid (53 mg).

MS (EI) 318 (M^+); mp 233 °C (dec.).

Example 30

3-(4-Oxo-7-m-tolyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-3'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K15) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow powder (145 mg).

25

MS (ISP) 352 [(M+H)⁺]; mp 248-251 °C (dec.).

Example 31

3-[7-(3,4-Dichloro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 5 Prepared from (3-amino-3',4'-dichloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G21) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (150 mg).
- 10 MS (EI) 405 (M⁺); mp 252-255 °C (dec.).

Example 32

8-Phenyl-4-(3-trifluoromethyl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from (3-amino-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G9) (284 mg, 1.0 mmol) and 2,2-dimethyl-6-(3-trifluoromethyl-phenyl)-[1,3]dioxin-4-one
- 15 (Example J5) (408 mg, 1.2 mmol) according to the general procedure K. The obtained material (392 mg) was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (238 mg).

MS (EI) 380 (M⁺); mp 210 °C (dec.).

Example 33

- 20 3-[7-(2-Chloro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from (3-amino-2'-chloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G22) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a
- 25 white powder (43 mg).

MS (ISP) 372 [(M+H)⁺]; mp 240-244 °C (dec.).

Example 34

4-(3-Chloro-thiophen-2-yl)-8-o-tolyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {3-[3-(3-chloro-thiophen-2-yl)-3-oxo-propionylamino]-2'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K16) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow powder (60 mg).

MS (ISP) 366 (M⁺) and 368 [(M+2)⁺]; mp 247-248 °C (dec.).

5 Example 35

8-Benzoyl-4-(3-chloro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {4-benzoyl-2-[3-(3-chloro-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K17) (193 mg, 0.39 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (93 mg).

10 MS (EI) 374 (M⁺) and 376 [(M+2)⁺]; mp 199-202 °C (dec.).

Example 36

3-(7-Benzoyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {4-benzoyl-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K18) (263 mg, 0.54 mmol) by treatment with TFA
15 in CH₂Cl₂ according to the general procedure M. Obtained as an orange solid (77 mg).

MS (EI) 365 (M⁺); mp 207 °C (dec.).

Example 37

8-Benzoyl-4-thiophen-2-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from [4-benzoyl-2-(3-oxo-3-thiophen-2-yl-propionylamino)-phenyl]-carbamic
20 acid tert.-butyl ester (Example K19) (54 mg, 0.12 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (8 mg).

MS (EI) 346 (M⁺).

Example 38

4-(3-Chloro-thiophen-2-yl)-8-phenyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {3-[3-(3-chloro-thiophen-2-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K20) (100 mg, 0.21 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (93 mg).

MS (EI) 352 (M⁺) and 354 [(M+2)⁺]; mp 228 °C (dec.).

5 Example 39

3-[7-(3-Methoxy-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-3'-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K21) by treatment with TFA in CH₂Cl₂
10 according to the general procedure M. Obtained as a light yellow powder (118 mg).

MS (ISP) 368 [(M+H)⁺]; mp 240-243 °C (dec.).

Example 40

3-(4-Oxo-7-pyridin-3-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from (2-amino-4-pyridin-3-yl-phenyl)-carbamic acid tert.-butyl ester (Example
15 G26) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a beige powder (161 mg).

MS (EI) 338 (M⁺); mp 210-214 °C (dec.).

20 Example 41

8-(6-Oxo-1,6-dihydro-pyridin-3-yl)-4-thiophen-2-yl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from [5-(6-oxo-1,6-dihydro-pyridin-3-yl)-2-(3-oxo-3-thiophen-2-yl-propionylamino)-phenyl]-carbamic acid tert.-butyl ester (Example K22) (127 mg, 0.28
25 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (57 mg).

MS (ISP) 336 [(M+H)⁺]; mp >250 °C.

Example 42

8-(6-Benzyloxy-pyridin-3-yl)-4-(3-trifluoromethyl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4-(6-benzyloxy-pyridin-3-yl)-2-[3-oxo-3-(3-trifluoromethyl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K 23) (198 mg, 0.33 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (49 mg).

MS (ISP) 488 [(M+H)⁺]; mp 195 °C (dec.).

Example 43

- 10 3-[4-Oxo-7-(4-trifluoromethoxy-phenyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4'-trifluoromethoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K24) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow powder (23 mg).

- 15 MS (EI) 421 (M⁺); mp 237-241 °C (dec.).

Example 44

3-(4-Oxo-7-pyridin-4-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- Prepared from (2-amino-4-pyridin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example G28) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a red-brown solid (25 mg).

MS (EI) 338 (M⁺); mp >200 °C (dec.).

Example 45

- 25 3-[7-(1-Benzenesulfonyl-1,2,3,6-tetrahydro-pyridin-4-yl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 3-[4-oxo-7-(1,2,3,6-tetrahydro-pyridin-4-yl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile [prepared from 4-{4-tert.-butoxycarbonylamino-

3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (Example K25) (690 mg, 1.23 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a brown solid (309 mg)] (56 mg, 0.164 mmol) and benzenesulfonyl chloride (0.164 mmol) in THF (2 mL) at
5 23 °C. Obtained as a brown solid (40 mg).

MS (ISP) 483 [(M+H)⁺]; mp 92-95 °C (dec.).

Example 46

3-[7-(1-Methanesulfonyl-1,2,3,6-tetrahydro-pyridin-4-yl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

10 Prepared from 3-[4-oxo-7-(1,2,3,6-tetrahydro-pyridin-4-yl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile [prepared from 4-{4-tert.-butoxycarbonylamino-3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (Example K25) (690 mg, 1.23 mmol) by treatment with
15 TFA in CH₂Cl₂ according to the general procedure M. Obtained as a brown solid (309 mg)] (60 mg, 0.175 mmol), methanesulfonyl chloride (0.014 mL, 0.175 mmol) and Et₃N (0.025 mL, 0.175 mmol) in THF (3 mL) at 23 °C. Obtained as an orange solid (35 mg).

MS (ISP) 421 [(M+H)⁺]; mp 211-213 °C (dec.).

Example 47

3-[7-(1-Acetyl-1,2,3,6-tetrahydro-pyridin-4-yl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile
20

Prepared from 3-[4-oxo-7-(1,2,3,6-tetrahydro-pyridin-4-yl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile [prepared from 4-{4-tert.-butoxycarbonylamino-3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (Example K25) (690 mg, 1.23 mmol) by treatment with
25 TFA in CH₂Cl₂ according to the general procedure M. Obtained as a brown solid (309 mg)] (54 mg, 0.158 mmol) and Ac₂O (0.017 mL, 0.173 mmol) in THF (3 mL) at 23 °C. Obtained as an orange solid (65 mg).

MS (EI) 384 (M⁺); mp 220-222 °C (dec.).

Example 48

8-(6-Benzyloxy-pyridin-3-yl)-4-(3-chloro-thiophen-2-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4-(6-benzyloxy-pyridin-3-yl)-2-[3-(3-chloro-thiophen-2-yl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K26) (172 mg, 0.3 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (38 mg).

MS (ISP) 460 [(M+H)⁺]; mp 213-215 °C (dec.).

Example 49

- 10 3-[7-(4-Methoxy-phenylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 3-(7-iodo-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile (Example 1) (290 mg, 0.75 mmol) and 4-methoxy-phenylacetylene (350 mg, 1.57 mmol) according to the general procedure F. Obtained as a light yellow solid (172 mg).

- 15 MS (EI) 391 (M⁺); mp 234-235 °C (dec.).

Example 50

3-(4-Oxo-7-thiophen-2-ylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- Prepared from 3-(7-iodo-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile (Example 50) (185 mg, 0.48 mmol) and 2-ethynylthiophene (78 mg, 0.72 mmol) [prepared from 2-thiophenecarboxaldehyde according to *J. Org. Chem.* 1982, 47, 2201-2204] according to the general procedure F. Obtained as a light yellow solid (146 mg).

MS (EI) 367 (M⁺); mp 235-238 °C (dec.).

Example 51

- 25 3-[7-(2,3-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2',3'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K8) (35 mg, 0.07 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a grey solid (23 mg).

MS (ISP) 374 [(M+H)⁺]; mp 240 °C.

5 Example 52

4-(3-Imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example K27) (170 mg, 0.33 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a brown solid (64
10 mg).

MS (EI) 402 (M⁺); mp 193-196 °C.

Example 53

3-(7-Benzofuran-2-yl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {4-benzofuran-2-yl-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K28) (255 mg, 0.51 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (260
15 mg).

MS (EI) 377 (M⁺); mp >250 °C.

Example 54

20 3-(7-Biphenyl-4-yl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-[1,1';4',1'']terphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K29) (135 mg, 0.25 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (100
mg).

25 MS (EI) 413 (M⁺); mp 251-253 °C.

Example 55

3-[4-Oxo-7-(4-trifluoromethoxy-phenylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-(4-trifluoromethoxy-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example K30) (170 mg, 0.3 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a beige solid (112 mg).

MS (EI) 445 (M⁺); mp 239-241 °C.

Example 56

10 4-(3-Nitro-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2-[3-(3-nitro-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example K31) (970 mg, 1.94 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (631 mg).

MS (EI) 381 (M⁺); mp 228-229 °C.

15 Example 57

3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K32) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as an orange solid (141 mg).

20 MS (EI) 355 (M⁺).

Example 58

8-(4-Fluoro-phenylethynyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K33) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a beige solid (89 mg).

MS (EI) 420 (M⁺); mp 216-218 °C.

Example 59

4-(3-Imidazol-1-yl-phenyl)-8-phenyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K34) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (125 mg).

MS (EI) 378 (M⁺); mp 201-205 °C.

Example 60

8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4'-fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K35) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as an orange solid (261 mg).

MS (EI) 396 (M⁺).

Example 61

- 4-(3-Imidazol-1-yl-phenyl)-8-(4-methoxy-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (3-amino-4'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K36) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (100 mg).

- MS (EI) 408 (M⁺); mp 207-208 °C.

Example 62

8-(4-Fluoro-phenylethynyl)-4-(3-[1,2,4]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4-(4-fluoro-phenylethynyl)-2-[3-oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K37) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (156 mg).

MS (EI) 421 (M⁺); mp 217-218 °C.

Example 63

8-(2-Fluoro-phenylethynyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {4-(2-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K38) by treatment with
5 TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (99 mg).

MS (EI) 420 (M⁺); mp 201-203 °C.

Example 64

8-(4-Fluoro-phenyl)-4-(3-[1,2,4]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-
10 2-one

Prepared from {4'-fluoro-3-[3-oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K39) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (115 mg).

MS (EI) 397 (M⁺); mp 241-245 °C.

15 Example 65

3-[7-(4-Cyano-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {4'-cyano-3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K40) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (102 mg).

20 MS (EI) 362 (M⁺); mp 240 °C (dec.).

Example 66

4-[2-(3-Imidazol-1-yl-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-7-yl]-benzonitrile

Prepared from {4'-cyano-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert-butyl ester (Example K41) by treatment with TFA in
25 CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (128 mg).

MS (EI) 403 (M⁺); mp 209-213 °C (dec.).

Example 67

8-(4-Fluoro-phenyl)-4-(3-nitro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {4'-fluoro-3-[3-(3-nitro-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-
carbamic acid tert.-butyl ester (Example K42) by treatment with TFA in CH₂Cl₂ according
5 to the general procedure M. Obtained as a light yellow solid (719 mg).

MS (ISP) 376 [(M+H)⁺]; mp 220-226 °C (dec.).

Example 68

8-(4-Fluoro-phenylethynyl)-4-(2-imidazol-1-yl-pyridin-4-yl)-1,3-dihydro-
benzo[b][1,4]diazepin-2-one

10 Prepared from {4-(4-fluoro-phenylethynyl)-2-[3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-
propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K43) by treatment with
TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (189
mg).

MS (ISP) 422 [(M+H)⁺]; mp 206 °C (dec.).

15 Example 69

8-(4-Fluoro-phenyl)-4-(2-imidazol-1-yl-pyridin-4-yl)-1,3-dihydro-
benzo[b][1,4]diazepin-2-one

Prepared from {4'-fluoro-3-[3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionylamino]-
biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K43) by treatment with TFA in
20 CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (209 mg).

MS (ISP) 398 [(M+H)⁺]; mp 213 °C (dec.).

Example 70

4-(3-Amino-phenyl)-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from 8-(4-fluoro-phenyl)-4-(3-nitro-phenyl)-1,3-dihydro-
25 benzo[b][1,4]diazepin-2-one (Example 67) by catalytic hydrogenation with Raney-Ni
according to the general procedure G (method a). Obtained as a light yellow solid (16 mg).

MS (ISP) 346 [(M+H)⁺]; mp 210-217 °C (dec.).

Example 71

8-(2-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-
5 biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K45) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (176 mg).

MS (ISP) 397 $[(\text{M}+\text{H})^+]$; mp 179-182 °C.

Example 72

10 N-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-acetamide

Prepared from 4-(3-amino-phenyl)-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 70) by treatment with Ac_2O in HOAc at 80 °C. Obtained as a light yellow solid (9 mg).

MS (EI) 387 (M^+).

15 Example 73

N-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-methanesulfonamide

Prepared from 4-(3-amino-phenyl)-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 70) by treatment with methanesulfonyl chloride
20 and Et_3N in THF at 23 °C. Obtained as a light yellow solid (24 mg).

MS (ISP) 424 $[(\text{M}+\text{H})^+]$; mp 232-235 °C.

Example 74

8-(6-Benzyloxy-pyridin-3-yl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

25 Prepared from {4-(6-benzyloxy-pyridin-3-yl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K46) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (75 mg).

MS (ISP) 486 [(M+H)⁺]; mp 225-231 °C.

Example 75

4-(3-Imidazol-1-yl-phenyl)-8-(6-oxo-1,6-dihydro-pyridin-3-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 5 Prepared from 8-(6-benzyloxy-pyridin-3-yl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 74) by catalytic hydrogenation with Raney-Ni according to the general procedure G (method a). Obtained as a light yellow solid (17 mg).

MS (ISP) 396 [(M+H)⁺].

Example 76

- 10 8-(4-Fluoro-phenyl)-4-[3-(4-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (4'-fluoro-3-{3-[3-(4-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K47) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light
15 yellow solid (196 mg).

MS (EI) 410 (M⁺); mp 215 °C.

Example 77

8-(2-Fluoro-phenyl)-4-(2-imidazol-1-yl-pyridin-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 20 Prepared from {2'-fluoro-3-[3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K48) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (32 mg).

MS (EI) 397 (M⁺).

Example 78

- 25 8-(4-Fluoro-phenylethynyl)-4-[3-(4-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (4-(4-fluoro-phenylethynyl)-2-{3-[3-(4-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-phenyl)-carbamic acid tert.-butyl ester (Example K49) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (122 mg).

5 MS (EI) 434 (M⁺); mp 210 °C.

Example 79

8-(4-Fluoro-2-methyl-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

10 Prepared from {4'-fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K50) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a beige solid (82 mg).

MS (EI) 410 (M⁺); mp 170-172 °C.

Example 80

15 8-(4-Fluoro-2-hydroxy-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {4'-fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2'-methoxymethoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K51) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (13 mg).

20 MS (ISP) 413 [(M+H)⁺]; mp 220 °C.

Example 81

8-(2-Fluoro-phenyl)-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

25 Prepared from (2'-fluoro-3-[3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K52) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (174 mg).

MS (ISP) 411 [(M+H)⁺]; mp 187 °C.

Example 82

8-(4-Fluoro-phenyl)-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from (4'-fluoro-3-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K53) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (178 mg).

MS (ISP) 411 [(M+H)⁺]; mp 227 °C.

Example 83

- 10 2-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile

Prepared from {3-[3-(3-cyano-thiophen-2-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K54) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as an orange solid (209 mg).

- 15 MS (EI) 361 (M⁺); mp 239-240 °C.

Example 84

2-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile

- Prepared from {3-[3-(3-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K55) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (258 mg).

MS (EI) 361 (M⁺); mp 238-240 °C.

Example 85

8-(4-Fluoro-phenyl)-4-(3-tetrazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 25 Prepared from {4'-fluoro-3-[3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K56) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (54 mg).

MS (EI) 398 (M^+).

Example 86

3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-
5 carbamic acid tert.-butyl ester (Example K57) by treatment with TFA in CH_2Cl_2 according
to the general procedure M. Obtained as a light yellow solid (128 mg).

MS (EI) 355 (M^+).

Example 87

8-(4-Fluoro-phenylethynyl)-4-(3-tetrazol-1-yl-phenyl)-1,3-dihydro-
10 benzo[b][1,4]diazepin-2-one

Prepared from [2-amino-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl
ester (Example G33) and 3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionic acid ethyl ester
(Example H15) according to the general procedure K. The obtained material was
deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general
15 procedure M. Obtained as a beige powder (5.5 mg).

MS (EI) 422 (M^+).

Example 88

8-(2-Fluoro-phenyl)-4-(3-tetrazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-oxo-3-(3-tetrazol-1-yl-phenyl)-propionylamino]-biphenyl-
20 4-yl}-carbamic acid tert.-butyl ester (Example K58) by treatment with TFA in CH_2Cl_2
according to the general procedure M. Obtained as a light yellow solid (31 mg).

MS (EI) 398 (M^+); mp 179-180 °C.

Example 89

4-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-8-(4-fluoro-phenyl)-1,3-dihydro-
25 benzo[b][1,4]diazepin-2-one

Prepared from (3-{3-[3-(2,4-dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K59) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (54 mg).

MS (EI) 424 (M⁺); mp 188 °C.

5 Example 90

4-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-{3-[3-(2,4-dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K60) by treatment with
10 TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (57 mg).

MS (EI) 430 (M⁺); mp 134 °C (dec.).

Example 91

4-(3-Chloro-thiophen-2-yl)-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

15 Prepared from {3-[3-(3-chloro-thiophen-2-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K61) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (232 mg).

MS (EI) 370 (M⁺); mp 255-256 °C.

Example 92

20 3-[7-(2-Fluoro-6-methoxy-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from (3-amino-2'-fluoro-6'-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G44) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example J4) according to the general procedure K. The obtained material was deprotected
25 and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a white solid (47 mg).

MS (EI) 385 (M⁺).

Example 93

8-(2-Fluoro-phenyl)-4-(3-nitro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-(3-nitro-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-
carbamic acid tert.-butyl ester (Example K62) by treatment with TFA in CH₂Cl₂ according
5 to the general procedure M. Obtained as a light yellow solid (62 mg).

MS (EI) 375 (M⁺).

Example 94

4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-
carbonitrile

10 Prepared from {3-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-
4-yl}-carbamic acid tert.-butyl ester (Example K63) by treatment with TFA in CH₂Cl₂
according to the general procedure M. Obtained as an orange solid (23 mg).

MS (EI) 356 (M⁺).

Example 95

15 4-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-
carbonitrile

Prepared from {3-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-
4-yl}-carbamic acid tert.-butyl ester (Example K64) by treatment with TFA in CH₂Cl₂
according to the general procedure M. Obtained as a green solid (40 mg).

20 MS (EI) 356 (M⁺).

Example 96

3-[7-(3-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-3'-fluoro-biphenyl-4-yl}-
carbamic acid tert.-butyl ester (Example K65) by treatment with TFA in CH₂Cl₂ according
25 to the general procedure M. Obtained as a yellow solid (57 mg).

MS (EI) 355 (M⁺); mp 232-235 °C.

Example 97

8-(3-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {3'-fluoro-3-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K66) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (69 mg).

MS (EI) 396 (M⁺); mp 200-201 °C.

Example 98

4-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-pyridine-2-carbonitrile

Prepared from {2-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example K67) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (10 mg).

MS (EI) 362 (M⁺).

Example 99

8-(2-Fluoro-phenyl)-4-(3-[1,2,4]triazol-4-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K68) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (79 mg).

MS (EI) 397 (M⁺); mp 198 °C.

Example 100

5-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiophene-2-carbonitrile

Prepared from {3-[3-(5-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K69) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (263 mg).

MS (EI) 361 (M^+); mp >250 °C.

Example 101

4-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-8-(2-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 5 Prepared from (3-{3-[3-(2,4-dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K70) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a yellow solid (92 mg).

MS (EI) 424 (M^+); mp 92 °C.

Example 102

- 10 8-(2-Fluoro-phenyl)-4-[3-(2-methoxymethylsulfanyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2'-fluoro-3-{3-[3-(2-methoxymethylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K71) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light
15 yellow solid (100 mg).

MS (EI) 472 (M^+); mp 189-190 °C.

Example 103

4-Fluoro-3-[7-(2-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 20 Prepared from {3-[3-(5-cyano-2-fluoro-phenyl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K72) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a yellow solid (92 mg).

MS (EI) 373 (M^+); mp 239 °C.

Example 104

- 25 4-[7-(2,4-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile

Prepared from {3-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-2',4'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K73) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (24 mg).

MS (ISP) 375 $[(\text{M}+\text{H})^+]$; mp 250-253 °C.

5 Example 105

4-(7-Isopropyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-pyridine-2-carbonitrile

Prepared from {2-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-4-isopropyl-phenyl}-carbamic acid tert.-butyl ester (Example K74) by treatment with TFA in CH_2Cl_2 according
10 to the general procedure M. Obtained as a yellow solid (33 mg).

MS (ISP) 305 $[(\text{M}+\text{H})^+]$; mp 173-178 °C.

Example 106

8-(2-Fluoro-phenyl)-4-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

15 Prepared from (2'-fluoro-3-{3-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K75) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a yellow solid (75 mg).

MS (ISP) 418 $[(\text{M}+\text{H})^+]$; mp 229 °C.

20 Example 107

8-(4-Fluoro-phenyl)-4-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (4'-fluoro-3-{3-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K76) by
25 treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a yellow solid (101 mg).

MS (ISP) 418 $[(\text{M}+\text{H})^+]$; mp 229 °C.

Example 108

3-(4-Oxo-7-phenethyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from 3-(4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile (Example 2) by catalytic hydrogenation with Pd/C according to the general
5 procedure G (method a). Obtained as a light yellow solid (156 mg).

MS (EI) 365 (M^+); mp 212-214 °C.

Example 109

3-(4-Oxo-7-styryl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from 3-(4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-
10 benzonitrile (Example 2) by catalytic hydrogenation with Lindlar-catalyst in the presence of 10 eq. cyclohexene according to the general procedure G (method a). Obtained as a light yellow solid (159 mg).

MS (EI) 363 (M^+); mp 185-188 °C.

Example 110

15 8-(2-Fluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K77) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (100 mg).

20 MS (EI) 397 (M^+); mp 209-212 °C.

Example 111

3-(7-Cyclopropyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-cyclopropyl-phenyl}-carbamic acid tert.-butyl ester (Example K78) by treatment with TFA in CH_2Cl_2 according
25 to the general procedure M. Obtained as a light yellow solid (44 mg).

MS (ISP) 302 [$(M+H)^+$]; mp 204-209 °C.

Example 112

4-(7-Cyclopropyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-pyridine-2-carbonitrile

- Prepared from {2-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-4-cyclopropyl-phenyl}-carbamic acid tert.-butyl ester (Example K79) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (88 mg).

MS (ISP) 303 [(M+H)⁺]; mp 227-230 °C.

Example 113

8-Cyclopropyl-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4-cyclopropyl-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K80) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (41 mg).

MS (ISP) 303 [(M+H)⁺]; mp 198-200 °C.

Example 114

- 8-(4-Fluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K81) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (29 mg).

- MS (EI) 397 (M⁺); mp 218-220 °C.

Example 115

8-(4-Fluoro-phenylethynyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4-(4-fluoro-phenylethynyl)-2-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example K82) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (53 mg).

MS (EI) 421 (M^+); mp 246-248 °C.

Example 116

8-(4-Fluoro-phenyl)-4-(2-imidazol-1-yl-thiazol-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 5 Prepared from {4'-fluoro-3-[3-(2-imidazol-1-yl-thiazol-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K83) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light brown solid (64 mg).

MS (ISP) 404 [$(M+H)^+$]; mp 214 °C.

Example 117

- 10 8-(4-Fluoro-phenyl)-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from (4'-fluoro-3-{3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K84) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light
15 yellow solid (33 mg).

MS (EI) 411 (M^+); mp >250 °C.

Example 118

3-[7-(2,4-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 20 Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2',4'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K85) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (37 mg).

MS (ISP) 374 [$(M+H)^+$].

Example 119

- 25 3-(7-Isopropyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-isopropyl-phenyl}-carbamic acid tert.-butyl ester (Example K86) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as an off-white solid (25 mg).

MS (ISP) 304 [(M+H)⁺].

5 Example 120

8-(4-Fluoro-phenyl)-4-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (4'-fluoro-3-{3-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K87) by
10 treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (104 mg).

MS (ISP) 443 [(M+H)⁺].

Example 121

8-Isopropyl-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-
15 one

Prepared from (4-isopropyl-2-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-phenyl)-carbamic acid tert.-butyl ester (Example K88) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (94 mg).

20 MS (ISP) 359 [(M+H)⁺].

Example 122

8-(2,4-Difluoro-phenyl)-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2',4'-difluoro-3-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K89) by
25 treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (71 mg).

MS (ISP) 429 [(M+H)⁺].

Example 123

8-(2,4-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2',4'-difluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K90) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light brown solid (78 mg).

MS (ISP) 416 [(M+H)⁺].

Example 124

8-(2-Fluoro-phenyl)-4-(2-imidazol-1-yl-thiazol-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-3-[3-(2-imidazol-1-yl-thiazol-4-yl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K91) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (46 mg).

MS (ISP) 404 [(M+H)⁺]; mp 194-197 °C.

Example 125

3-[7-(2,5-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {3-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2',5'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K92) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (49 mg).

MS (EI) 373 (M⁺); mp 247 °C.

Example 126

8-(2,5-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2',5'-difluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K93) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (67 mg).

MS (EI) 415 (M^+); mp 231 °C.

Example 127

8-(2-Fluoro-phenyl)-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 5 Prepared from (2'-fluoro-3-{3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example K94) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (100 mg).

MS (EI) 411 (M^+); mp 222-224 °C.

10 Example 128

7-Methyl-4-[3-(1-methyl-1H-imidazol-2-yl)-phenyl]-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from (2-amino-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example G2) (620 mg, 2.0 mmol) and 3-[3-(1-methyl-1H-imidazol-2-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example H21) (1.26 g, 4.2 mmol) according to the general procedure K. The obtained material was deprotected and cyclized by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a light yellow solid (81 mg).

MS (ISP) 417.2 [$(M+H)^+$]; mp 202-205 °C.

Example 129

- 20 8-(2,3-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {2',3'-difluoro-3-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example K95) (118 mg, 0.22 mmol) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a yellow solid (74 mg).

mp 230 °C.

Example 130

3-[7-(2,4-Difluoro-phenylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-(2,4-difluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example K96) by treatment with
5 TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (87 mg).

MS (EI) 397 (M⁺); mp 240-246 °C.

Example 131

10 3-[7-(2-Fluoro-phenylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-(2-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example K97) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a yellow solid (82 mg).

15 MS (EI) 379 (M⁺); mp 213-214 °C.

Example 132

8-Phenylethynyl-4-(3-trimethylsilanylethynyl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (231 mg, 0.5 mmol) and trimethylsilylacetylene (0.4 mL, 1.0 mmol)
20 according to the general procedure F. Obtained as a light brown solid (102 mg).

MS (EI) 432 (M⁺); mp 169-171 °C (dec.).

Example 133

4-(3-Ethynyl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

25 Prepared from 8-phenylethynyl-4-(3-trimethylsilanylethynyl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 132) (63 mg, 0.146 mmol) by treatment with 1N NaOH (3drops) in MeOH (3 mL) and THF (0.5 mL) at 23 °C for 2 h. Obtained as a brown solid (31 mg).

MS (ISP) 361 [(M+H)⁺]; mp >250 °C (dec.).

Example 134

3-[7-(3-Hydroxy-3-methyl-but-1-ynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 5 Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (194 mg, 0.5 mmol) and 2-methyl-3-butyn-2-ol (105 mg, 1.25 mmol) according to the general procedure F. Obtained as a yellow solid (147 mg).

MS (EI) 343 (M⁺); mp 243 °C.

Example 135

- 10 3-[7-(1-Hydroxy-cyclohexylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (308 mg, 0.80 mmol) and 1-ethynyl-1-cyclohexanol (248 mg, 2.00 mmol) according to the general procedure F. Obtained as a beige solid (262 mg).

- 15 MS (ISP) 384 [(M+H)⁺]; mp 196 °C.

Example 136

3-(7-Cyclohex-1-enylethynyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- 20 Prepared from 3-[7-(1-hydroxy-cyclohexylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 135) (50 mg, 0.13 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (22 mg).

MS (EI) 365 (M⁺); mp 228 °C.

Example 137

- 25 3-[7-(3-Methyl-but-3-en-1-ynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 3-[7-(3-hydroxy-3-methyl-but-1-ynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 134) (50 mg, 0.15 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a sand solid (33 mg).

- 5 MS (ISP) 326 [(M+H)⁺]; mp 231 °C.

Example 138

3-[7-(3,6-Dihydro-2H-pyran-4-ylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from 3-[7-(4-hydroxy-tetrahydro-pyran-4-ylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile [prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (387 mg, 1.0 mmol) and 4-ethynyl-tetrahydro-pyran-4-ol [CAS-No. 57385-16-7] (315 mg, 2.5 mmol) according to the general procedure F. Obtained as a yellow solid (296 mg)] (50 mg, 0.13 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as a light yellow solid (16 mg).
- 10
15

MS (EI) 367 (M⁺); mp 238 °C.

Example 139

4-[2-(3-Cyano-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-7-ylethynyl]-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester

- Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (1.16 g, 3.0 mmol) and 4-ethynyl-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester [CAS-No. 177984-28-0] (0.93 g, 4.5 mmol) according to the general procedure F. Obtained as a yellow solid (620 mg).
- 20

MS (ISP) 467 [(M+H)⁺]; mp 239 °C.

- 25 Example 140

3-[4-Oxo-7-(1,2,3,6-tetrahydro-pyridin-4-ylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 4-[2-(3-cyano-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-7-ylethynyl]-3,6-dihydro-2H-pyridine-1-carboxylic acid tert.-butyl ester (Example 139) (573

mg, 1.13 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure M. Obtained as an off-white solid (434 mg).

MS (ISP) 367 [(M+H)⁺]; mp 184 °C.

Example 141

- 5 3-[7-(1-Acetyl-1,2,3,6-tetrahydro-pyridin-4-ylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 3-[4-oxo-7-(1,2,3,6-tetrahydro-pyridin-4-ylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 140) (55 mg, 0.15 mmol) by treatment with Ac₂O (0.017 mL, 0.165 mmol) in THF (3 mL) at 23 °C for 1 h. Obtained as a sand
10 solid (59 mg).

MS (ISP) 409 [(M+H)⁺]; mp 203 °C.

Example 142

3-[7-(1-Methanesulfonyl-1,2,3,6-tetrahydro-pyridin-4-ylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 15 Prepared from 3-[4-oxo-7-(1,2,3,6-tetrahydro-pyridin-4-ylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 140) (55 mg, 0.15 mmol) by treatment with methanesulfonyl chloride (0.012 mL, 0.15 mmol) and Et₃N (0.021 mL, 0.15 mmol) in THF (3 mL) at 23 °C for 23 h. Obtained as a yellow solid (38 mg).

MS (EI) 444 (M⁺); mp 235 °C.

- 20 Example 143

3-{7-[1-(2,2-Dimethyl-propionyl)-1,2,3,6-tetrahydro-pyridin-4-ylethynyl]-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl}-benzonitrile

- Prepared from 3-[4-oxo-7-(1,2,3,6-tetrahydro-pyridin-4-ylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 140) (55 mg, 0.15 mmol) by treatment
25 with pivaloyl chloride (0.02 mL, 0.165 mmol) and Et₃N (0.021 mL, 0.15 mmol) in THF (3 mL) at 23 °C for 19 h. Obtained as a sand solid (26 mg).

MS (ISP) 451 [(M+H)⁺]; mp 219 °C.

Example 144

3-[4-Oxo-7-(3-oxo-cyclohex-1-enylethynyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (500 mg, 1.29 mmol) and 3-ethynyl-cyclohex-2-enone [CAS-No. 124267-26-1] (310 mg, 2.58 mmol) according to the general procedure F. Obtained as a yellow solid (346 mg).

MS (EI) 379 (M^+); mp 110 °C.

Example 145

3-[7-(4-Fluoro-phenylethynyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example K98) (307 mg, 0.62 mmol) by treatment with TFA in CH_2Cl_2 according to the general procedure M. Obtained as a eggshell solid (146 mg).

MS (EI) 379 (M^+); mp 245-246 °C.

The following examples relate to the palladium-catalyzed carbonylation of the 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one in the presence of secondary amines leads directly to the corresponding amides as shown in synthetic scheme H.

General procedure N

Preparation of 4-[3-(Amino-4-carbonyl)-phenyl]-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones by Pd-catalyzed carbonylative amination of 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

A solution of 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (1.0 mmol), the secondary amine (5.0 mmol), PPh_3 (6 mol%) or dppp (3 mol%), $Pd(OAc)_2$ (3 mol%) and Et_3N (2.0 mmol) in DMF (4 mL) was stirred at 23 °C under CO-atmosphere until tlc indicated complete consumption of the iodide. After

dilution with EtOAc, washing with sat. NaHCO_3 -sol. and brine, the organic phase was dried over Na_2SO_4 . Removal of the solvent left a brown oil, which was purified by silica gel column chromatography with hexane/EtOAc to give the title compound.

Example 146

- 5 4-[3-(Morpholine-4-carbonyl)-phenyl]-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (471 mg, 1.02 mmol) and morpholine (0.44 mL, 5.09 mmol) according to the general procedure N. Obtained as a light yellow solid (324 mg).

- 10 MS (ISP) 450 $[(\text{M}+\text{H})^+]$; mp 142-145 °C (dec.).

Example 147

4-[3-(4-Methyl-piperazine-1-carbonyl)-phenyl]-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 15 Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (231 mg, 0.5 mmol) and N-methylpiperazine (0.28 mL, 2.5 mmol) according to the general procedure N. Obtained as a light yellow solid (65 mg).

MS (ISP) 463 $[(\text{M}+\text{H})^+]$; mp 168-172 °C (dec.).

Example 148

3-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzamide

- 20 Prepared from 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 11) (231 mg, 0.5 mmol) and hexamethyldisilazane (0.52 mL, 2.5 mmol) according to the general procedure N. Obtained as a yellow solid (104 mg).

MS (EI) 379 (M^+); mp >250 °C (dec.).

Example 149

- 25 3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamide

Preparation of 4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamides (according to Pen-Yuan Lin et al. *Synthesis* 1992, 1219]: To a solution of hexamethyldisilthiane (1.45 g, 8.0 mmol) in 1,3-dimethyl-2-imidazolidinone (8 mL) was added at 23 °C sodium methoxide (0.31 g, 6.8 mmol). The mixture was stirred for 5 min. and the blue solution formed was then added to a solution of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 57) (1.42 g, 4 mmol) in 1,3-dimethyl-2-imidazolidinone (12 mL). The mixture was stirred for 3 h at 23 °C and then poured into water (100 mL). Stirring was continued for 0.5 h and the precipitate was isolated by filtration and triturated with ethyl acetate (50 mL) to give the title compound as light yellow solid (1.38 g).

MS (ISN) 387.9 [(M-H)⁻]; mp 267-270 °C (dec.).

Example 150

3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamide

3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 86) (0.53 g) was subjected in an analogous manner to the procedure described in Example 149 to give, after trituration of the crude product with ethyl acetate, the title compound as light yellow solid (0.46 g).

MS (ISN) 387.9 [(M-H)⁻]; mp 232-234 °C (dec.).

Example 151

4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbothioic acid amide

(Example 94) (0.36 g) was subjected in an analogous manner the procedure described in Example 149 to give, after trituration of the crude product with acetone, the title compound as light yellow solid (0.33 g).

MS (ISN) 388.9 [(M-H)⁻]; mp 280-283 °C (dec.).

Example 152

8-(4-Fluoro-phenyl)-4-[3-(4-methyl-thiazol-2-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

To a suspension of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamide (Example 149) (97 mg, (0.25 mmol) in ethanol (5 mL) was added chloroacetone (0.1 mL, 1.25 mmol) and the mixture was heated at reflux. After 1 h and after 5 h of heating, more chloroacetone (2 times 0.1 mL, 1.25 mmol) was added. After 20 h the mixture was cooled to 0 °C and stirred for 0.5 h, and the title compound was isolated by filtration as light yellow solid (61 mg).

MS (ISP) 428.4 [(M+H)⁺]; mp 259-261 °C (dec.).

Example 153

2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid ethyl ester

A mixture of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamide (Example 149) (39 mg, 0.1 mmol) and ethyl bromopyruvate (29 mg, 0.15 mmol) in ethanol (0.5 mL) was heated at reflux for 7 h. The solution was cooled to 0 °C, stirred for 0.5 h, and the title compound was isolated by filtration as light yellow solid (29 mg).

MS (ISN) 484.2 [(M-H)⁻]; mp 236-238 °C (dec.).

Example 154

2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid ethyl ester

3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamide (Example 150) (0.24 g) was subjected in an analogous manner to the procedure described in Example 153. The reaction mixture was evaporated in vacuum and the residue was triturated with ethyl acetate to give the title compound as light yellow solid (0.29 g).

MS (ISP) 503.4 [(M+NH₄)⁺]; mp 172-175 °C.

Example 155

2-{4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridin-2-yl}-thiazole-4-carboxylic acid ethyl ester

- 4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbothioic acid amide (Example 151) (160 mg) was subjected in an analogous manner to the procedure described in Example 153. The cooled reaction solution was diluted with ethyl acetate, washed with NaHCO₃-sol. and with brine, dried over Na₂SO₄ and evaporated
5 in vacuum. The crude product was chromatographed on silica gel using ethyl acetate/hexane (2:1) as eluent, and the purified product was triturated with dichloromethane/diethyl ether to give the title compound as light yellow solid (40 mg).

MS (ISN) 485.1 [(M-H)⁻]; mp 238-240 °C (dec.).

Example 156

- 10 8-(4-Fluoro-phenyl)-4-[3-(4-hydroxymethyl-thiazol-2-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one.

- To a solution of 2-{3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid ethyl ester (Example 153) (110 mg, 0.23 mmol) in THF (7 mL) was added a 3.5 M solution of sodium dihydrido-bis(2-
15 methoxyethoxy)aluminate in toluene (0.66 mL, 2.3 mmol). The reaction mixture was stirred at 23 °C for 15 min. and then poured into ice-cold 10% aq. acetic acid. The mixture was extracted with ethyl acetate and the organic layer was washed successively with H₂O, sat. Na₂CO₃-sol. and brine, dried over Na₂SO₄ and evaporated in vacuum. The residue was triturated with ethyl acetate to give the title compound as light yellow solid (48 mg).

- 20 MS (ISN) 442.1 [(M-H)⁻]; mp 235-239 °C (dec.).

Example 157

8-(2-Fluoro-phenyl)-4-[3-(4-hydroxymethyl-thiazol-2-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one.

- 2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-
25 thiazole-4-carboxylic acid ethyl ester (Example 154) (97 mg) was subjected in an analogous manner to the procedure described in Example 156 to give the title compound as light yellow solid (30 mg).

MS (ISN) 442.1 [(M-H)⁻]; mp 238-244 °C (dec.).

Example 158

8-(4-Fluoro-phenyl)-4-[2-(4-hydroxymethyl-thiazol-2-yl)-pyridin-4-yl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one.

- 2-{4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridin-2-yl}-thiazole-4-carboxylic acid ethyl ester (Example 155) (84 mg) was subjected in an analogous manner to the procedure described in Example 156. The crude product was chromatographed on silica gel using ethyl acetate/hexane (2:1) and ethyl acetate as eluent, and the purified product was crystallized from ethyl acetate/hexane to give the title compound as light yellow solid (16 mg).
- 10 MS (ISP) 445.2 [(M+H)⁺]; mp 225-229 °C.

Example 159

2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid.

- To a solution of 2-{3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid ethyl ester (Example 153) (49 mg, 0.1 mmol) in a mixture of MeOH (1 mL) and DMSO (1.5 mL) was added 2N KOH (1 mL, 2 mmol). The mixture was stirred at 23 °C for 0.5 h and then partitioned between ethyl acetate and H₂O. The pH of the aqueous phase was set to 3 by the addition of 3N HCl and then stirred in the ice-bath for 0.5 h. The precipitate was collected by filtration, triturated with EtOH/Et₂O (1/1) and dried, to give the title compound as light yellow solid (25 mg).
- 15 20

MS (ISN) 456.4 [(M-H)⁻]; mp 286-290 °C (dec.).

Example 160

2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid.

- 25 2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid ethyl ester (Example 154) (97 mg) was subjected in an analogous manner to the procedure described in Example 159 to give the title compound as light yellow solid (60 mg).

MS (ISN) 456.2 [(M-H)⁻]; mp 261-263 °C (dec.).

Example 161

2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid (2-hydroxy-ethyl)-amide

A solution of 2-{3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid ethyl ester (Example 153) (49 mg, 0.1 mmol) in 2-amino-ethanol was heated to 40 °C for 40 min. The cooled solution was partitioned between ethyl acetate and H₂O and the organic layer was dried over Na₂SO₄ and evaporated in vacuum. The residue was triturated with ethyl acetate to give the title compound as light yellow solid (21 mg).

MS (ISP) 518.3 [(M+NH₄)⁺]; mp 238-245 °C (dec.).

Example 162

2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-4-methyl-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide

A mixture 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiobenzamide (Example 149) (78 mg, 0.2 mmol) and ethyl 4-chloro-3-oxo-butanoate (164 mg, 1.0 mmol) in ethanol (4 mL) was heated at reflux for 24 h. The solution was cooled to 0 °C and stirred for 0.5 h. The crystals were isolated by filtration and subsequently suspended in 2-amino-ethanol (4 mL). The mixture was stirred at 70 °C for 4 h, cooled to 23 °C, and partitioned between ethyl acetate and H₂O. The organic layer was dried over Na₂SO₄ and evaporated in vacuum and the residue was triturated with ethyl acetate to give the title compound as light yellow solid (37 mg).

MS (ISP) 515.3 [(M+H)⁺]; mp 248-251 °C (dec.).

Example 163

3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid

To a suspension of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 57) (355 mg, 1 mmol) in EtOH (2 mL) was added 2N KOH (5 mL). The mixture was heated at reflux for 3 h, then cooled to 23 °C and partitioned between ethyl acetate and H₂O. The pH of the aqueous phase was set to 3 by the addition of 3N HCl, a precipitate being formed. The mixture was stirred in the ice-bath for 0.5 h.

The crystalline solid was collected by filtration, triturated with ethyl acetate and dried, to give the title compound as light yellow solid (218 mg).

MS (ISN) 373.3 [(M-H)⁻]; mp 290-296 °C (dec.).*

Example 164

5 4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carboxylic acid

4-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile (Example 94) (356 mg) was subjected in an analogous manner to the procedure described in Example 163. The reaction mixture was partitioned between ethyl acetate and H₂O, and the pH of the aqueous phase was set to 3 by the addition of 3N HCl, a precipitate being formed. The mixture was extracted with ethyl acetate and the organic layer was washed with brine, dried over Na₂SO₄, and evaporated in vacuum. The residue was triturated with EtOH to give the title compound as yellow solid (190 mg).

MS (ISN) 374.2 [(M-H)⁻]; mp 211-213 °C (dec.).

15 Example 165

8-(4-Fluoro-phenyl)-4-[3-(5-methyl-[1,3,4]oxadiazol-2-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

A mixture of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid hydrazide (Starting material 8a-2) (58 mg, 0.15 mmol) and ethyl acetimidate hydrochloride (74 mg, 0.6 mmol) in pyridine (0.6 mL) was heated to 100 °C for 2 h. The mixture was cooled to 23 °C, diluted with H₂O (30 mL) and stirred for 20 min. The precipitate was collected by filtration, and the crude product was purified by chromatography on silica gel using ethyl acetate/hexane (2:1) as eluent. The purified product was crystallized from Et₂O/hexane to give the title compound as light yellow solid (28 mg).

MS (ISP) 430.3 [(M+NH₄)⁺]; mp 254-256 °C (dec.).

Starting material 8a-1

3-[7-(3-Fluoromethyl-phenyl)-8-methyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid allyl ester

To a solution of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid (Example 163) (749 mg, 2 mmol) and 1,1,3,3-tetramethylguanidine (230 mg, 2 mmol) in DMSO (24 mL) was added allyl bromide (242 mg, 2 mmol) and the mixture was stirred at 23 °C for 2.5 h. H₂O (100 mL) was added and stirring was continued for 20 min. The precipitate was collected by filtration and triturated with ethyl acetate/Et₂O (1:1) to give the title compound as light yellow solid (705 mg).

MS (ISP) 415.2 [(M+H)⁺]; mp 226-228 °C.

Starting material 8a-2

3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid hydrazide

To a suspension of 3-[7-(3-fluoromethyl-phenyl)-8-methyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid allyl ester (Starting material 8a-1) (207 mg, 0.5 mmol) in MeOH (3.5 mL) was added hydrazine hydrate (0.25 mL, 5 mmol). The mixture was stirred at 60 °C for 15 h, then cooled to 23 °C and diluted with H₂O (35 mL). The precipitate was collected by filtration and dried, and the crude product was purified by chromatography on silica gel using ethyl acetate/hexane (1:1) as eluent. The purified product was crystallized from Et₂O/hexane to give the title compound as light yellow solid (64 mg).

MS (ISN) 387.0 [(M-H)⁻]; mp 256-258 °C (dec.).

Example 166

4-[3-(4,5-Dimethyl-4H-[1,2,4]triazol-3-yl)-phenyl]-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

A mixture of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid hydrazide (Starting material 8a-2) (234 mg, 0.6 mmol) and ethyl acetimidate hydrochloride (185 mg, 1.5 mmol) in EtOH (12 mL) was heated at reflux for 0.5 h. The mixture was cooled and partitioned between ethyl acetate and H₂O. The organic layer was dried over Na₂SO₄ and evaporated in vacuum to give an amorphous residue (261 mg). A sample of this material (92 mg) was dissolved in a 33% solution (3 mL) of methylamine in EtOH. After being stirred for 0.5 h at 23 °C, the reaction mixture was evaporated in vacuum and the residue was heated in 1,4-dioxane (5 mL) to 100 °C for 2 h. The solvent was evaporated in vacuum and the crude product was chromatographed on silica gel using

ethyl acetate and ethyl acetate/methanol (4:1) as eluent. The purified product was triturated with ethyl acetate to give the title compound as light yellow solid (26 mg).

MS (ISP) 426.4 (M+H)⁺; mp 238-240 °C (dec.).

Example 167

- 5 3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-N-hydroxy-benzamidine

To a stirred suspension of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile (Example 57) (355 mg, 1 mmol) in EtOH (60 mL) was added at 23 °C over 6 h 50% aqueous hydroxylamine solution (3.48 mL, 60
10 mmol) in 6 portions. The mixture was cooled in the ice-bath, diluted with H₂O (100 mL) and stirred for 1 h. The precipitate was collected by filtration, washed with water and Et₂O, and triturated with ethyl acetate to give the title compound as light yellow solid (336 mg).

MS (ISP) 389.2 [(M+H)⁺]; mp 257-259 °C (dec.).

Example 168

- 15 8-(4-Fluoro-phenyl)-4-{3-[5-(3-hydroxy-propyl)-[1,2,4]oxadiazol-3-yl]-phenyl}-1,3-dihydro-benzo[b][1,4]diazepin-2-one

To a stirred suspension of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-N-hydroxy-benzamidine (Example 167) (78 mg, 0.2 mmol) in EtOH (3 mL) was added at 23 °C sodium hydride (55% dispersion in oil, 32 mg, 0.8
20 mmol) and the mixture was stirred at 23 °C for 5 min. Butyrolactone (103 mg, 1.2 mmol) was added and the mixture was heated to 70 °C for 3 h. The reaction mixture was cooled to 23 °C and partitioned between ethyl acetate and H₂O. The organic layer was dried over Na₂SO₄, evaporated in vacuum, and the residue was crystallized from dichloromethane to give the title compound as light yellow solid (60 mg).

- 25 MS (ISP) 457.3 [(M+H)⁺]; mp 220-222 °C.

Example 169

8-(4-Fluoro-phenyl)-4-[3-(3-methyl-[1,2,4]oxadiazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

To a solution of 3-[7-(4-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzoic acid (Example 163) (262 mg, 0.7 mmol) and 1-hydroxybenzotriazole (130 mg, 0.84 mmol) in CH₃CN/DMF (7 mL/10 mL), cooled to 0 °C, was added EDC (161 mg, 0.84 mmol) and the mixture was stirred at 0 °C for 1 h. N-hydroxy-acetamidine (62 mg, 0.84 mmol) was added and stirring was continued for 4 h at 23 °C. The reaction mixture was partitioned between ethyl acetate and H₂O and the organic layer was dried over Na₂SO₄ and evaporated in vacuum. The residue was heated in toluene (30 mL) for 20 h at reflux. The mixture was cooled, evaporated in vacuum and the residue was chromatographed on silica gel using ethyl acetate/hexane (2:1) as eluent. The purified product was triturated with ethyl acetate to give the title compound as light yellow solid (49 mg).

MS (ISN) 411.2 [(M-H)⁻]; mp 266-268 °C.

Example 170

8-(4-Fluoro-phenyl)-4-(2-hydroxy-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

A mixture of 4'-fluoro-biphenyl-3,4-diamine (prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) by treatment with TFA) (121 mg, 0.6 mmol) and 4-hydroxycoumarin (97 mmol, 0.6 mmol) in xylene (8 mL) was heated to 130 °C for 6 h. A yellow precipitate started to form after ca. 1 h. The mixture was cooled to 23 °C, diluted with ethyl acetate (8 mL) and hexane (8 mL) and stirring was continued for 15 min. The precipitate was collected by filtration and triturated with ethyl acetate/hexane (1:1) to give the title compound as a mixture with the isomeric 7-(4-fluoro-phenyl)-4-(2-hydroxy-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one as yellow solid (117 mg).

MS (ISP) 347.3 [(M+H)⁺].

Example 171

4-(3-Chloro-2,4-dihydroxy-phenyl)-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

By subjecting 4'-fluoro-biphenyl-3,4-diamine (prepared from (3-amino-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example G39) by treatment with TFA) (121 mg, 0.6 mmol) and 8-chloro-4,8-dihydroxycoumarin (128 mmol, 0.6 mmol) in an analogous manner to the procedure described in Example 170, the title compound was obtained as a mixture with the isomeric 4-(3-chloro-2,4-dihydroxy-phenyl)-7-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one as light yellow solid (103 mg).

- 135 -

MS (ISN) 395.1 [(M-H)⁻].Example I

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
5 Active ingredient	100
Powdered. lactose	95
White corn starch	35
Polyvinylpyrrolidone	8
Na carboxymethylstarch	10
10 Magnesium stearate	<u>2</u>
Tablet weight	250

Example II

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
15 Active ingredient	200
Powdered. lactose	100
White corn starch	64
Polyvinylpyrrolidone	12
20 Na carboxymethylstarch	20
Magnesium stearate	<u>4</u>
Tablet weight	400

Example III

Capsules of the following composition are produced:


- 136 -

	<u>mg/Capsule</u>
Active ingredient	50
Crystalline. lactose	60
Microcrystalline cellulose	34
5 Talc	5
Magnesium stearate	1
Capsule fill weight	150

The active ingredient having a suitable particle size, the crystalline lactose and the
10 microcrystalline cellulose are homogeneously mixed with one another, sieved and
thereafter talc and magnesium stearate are admixed. The final mixture is filled into hard
gelatin capsules of suitable size.

Claims

1. Compounds of general formula I



5 wherein

X is a single bond or an ethynediyl group, wherein,

in case X is a single bond,

R¹ is hydrogen; halogen; nitro; lower alkyl; halo-lower alkyl; alkoxy-carbonyl; lower
cycloalkyl, optionally substituted with oxygen; benzoyl, optionally substituted with lower
10 alkyl, halo-lower alkyl, or halogen; phenyl, optionally substituted with halogen, hydroxy,
lower alkyl, halo-lower alkyl, lower cycloalkyl, or lower alkoxy, halo-lower alkoxy, or
cyano; styrenyl; phenylethyl; naphthyl; biphenyl; benzofuranyl; or 5 or 6 membered
heterocyclic ring, optionally substituted with oxo, benzyloxy, benzoyl, methanesulfonyl,
benzenesulfonyl, or acetyl;

15 in case X is an ethynediyl group,

R¹ is hydrogen; lower alkyl, optionally substituted with hydroxy; halo-lower alkyl; lower
cycloalkyl, optionally substituted with hydroxy, lower alkyl, halo-lower alkyl, lower alkoxy,
halo-lower alkoxy, or halogen; lower cycloalkenyl, optionally substituted with lower alkyl,
halo-lower alkyl, lower alkoxy, halo-lower alkoxy, halogen, or oxo; lower alkenyl; phenyl,
20 optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower cycloalkyl, lower
alkoxy, halo; 5 or 6 membered heterocyclic ring, optionally substituted with lower alkyl,
halogen, oxo, benzyloxy, benzoyl, methanesulfonyl, benzenesulfonyl, acetyl, pivaloyl, tert.
butoxycarbonyl, or tert. butylcarbonyl; or benzofuranyl;

R³ is phenyl; pyridine; thiophenyl or thiazolyl, which are optionally substituted by halogen,
25 cyano, nitro, azido, hydroxy, carboxy, morpholine-4-carbonyl, carbamoyl, thiocarbamoyl,
N-hydroxycarbamoyl, trimethylsilyl-ethynyl, or from lower alkyl, lower alkynyl, lower

alkoxy, halo-lower alkyl, 4-lower alkyl-piperazine-1-carbonyl, lower alkylaminocarbonyl, which are optionally substituted by amino, lower alkylamino, acylamino, oxo, hydroxy; lower alkoxy, lower alkylthio, or carboxy which is optionally esterified or amidated; or a 5-membered aromatic heterocycle which is optionally substituted by amino, lower alkylamino, acylamino, oxo, hydroxy, lower alkoxy, lower alkylthio, carboxy which is optionally esterified with lower alkyl or amidated with lower alkylamino which is eventually substituted by hydroxy, or lower alkyl which is optionally substituted by halogen, hydroxy, amino, lower alkylamino, acylamino, or amidino, which is optionally substituted by lower alkyl, -C(NRR')=NR" (where R, R' and R" are hydrogen or lower alkyl), hydroxy, lower alkoxy, lower alkylthio, acyloxy, lower alkylsulfinyl, lower alkylsulfonyl, lower alkoxy-lower alkylsulfanyl, lower alkylsulfanyl, cycloalkylsulfinyl, cycloalkylsulfonyl, hydroxyimino, or lower alkoxyimino, which is optionally esterified or amidated, lower alkenyl, oxo, cyano, carbamoyloxy, or sulfamoyl which is optionally substituted by lower alkyl,

with the proviso that, if X is a single bond and R³ is pyridinyl, R¹ is not hydrogen, or methyl;

and their pharmaceutically acceptable acid addition salts.

2. Compounds according to claim 1, wherein R³ is phenyl substituted in meta position by cyano; halogen; or imidazolyl, which is optionally substituted by lower alkyl; or 1,3-thiazolyl which is optionally substituted by hydroxy-lower alkyl, carboxy, or -CO-NH-(CH₂)₂OH; 1,3-oxazolyl; 1,2,3-triazolyl; 1,2,4-triazolyl which is optionally substituted with lower alkyl; tetrazolyl; or isoxazolyl, which is optionally substituted by lower alkyl;


3. Compounds according to claim 2, being

3-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;
 4-(3-Chloro-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 4-(3-Imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;
 8-(4-Fluoro-phenylethynyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 8-(4-Fluoro-phenylethynyl)-4-(3-[1,2,4]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;

- 8-(4-Fluoro-phenyl)-4-[3-(4-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 8-(4-Fluoro-2-methyl-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 5 8-(4-Fluoro-2-hydroxy-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 8-(4-Fluoro-phenylethynyl)-4-(3-tetrazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 8-(2-Fluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 10 8-(4-Fluoro-phenyl)-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 8-(2,4-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 15 8-(2,3-Difluoro-phenyl)-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 8-(2-Fluoro-phenyl)-4-[3-(4-hydroxymethyl-thiazol-2-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 2-{3-[7-(2-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-thiazole-4-carboxylic acid;
 20 2-{3-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-phenyl}-4-methyl-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide; and
 4-[3-(4,5-Dimethyl-4H-[1,2,4]triazol-3-yl)-phenyl]-8-(4-fluoro-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one.
- 25 4. Compounds of formula I, in which R³ is thiophenyl, optionally substituted with halogen, cyano; or pyridinyl, optionally substituted with halogen, or cyano.
5. Compounds according to claim 4, being
- 8-(4-Fluoro-phenylethynyl)-4-(2-imidazol-1-yl-pyridin-4-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
 30 2-[7-(4-Fluoro-phenyl)-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile;
 4-(4-Oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-pyridine-2-carbonitrile; and
 4-[7-(2,4-Difluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile.
 35


- 140 -

6. A medicament containing one or more compounds of any one of claims 1 to 5 and pharmaceutically acceptable excipients.
7. A medicament according to claim 7 for the treatment or prevention of acute and/or chronic neurological disorders including psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.
8. A process for preparing compounds of formula I as defined in claim 1, which process comprises reacting a compound of formula III



III


with a compound of formula IV or IVa



IV

IVa

wherein R is ethyl or buthyl, to a compound of formula II



II

- which subsequently undergoes deprotection of the amino group and cyclization, to obtain a compound of formula I of claim 1.
9. A compound according to any one of claims 1-5, whenever prepared by a process as claimed in claim 9 or by an equivalent method.

- 141 -

10. The use of a compound according to any one of claims 1-5 for the treatment of acute and/or chronic neurological disorders including psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.
11. The use of one or more compounds according to claims 1 to 5 and/or one or more of
5 their pharmaceutically acceptable acid addition salts for the manufacture of medicaments for the treatment or prevention of diseases.
12. The use according to claim 11 for the treatment or prevention of acute and/or chronic neurological disorders including psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.
- 10 13. The invention as hereinbefore described.
