

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199948993 B2
(10) Patent No. 760168

(54) Title
Novel cyclosporins

(51)⁷ International Patent Classification(s)
C07K 007/64 A61K 038/13

(21) Application No: **199948993**

(22) Application Date: **1999.06.10**

(87) WIPO No: **WO99/65933**

(30) Priority Data

(31) Number	(32) Date	(33) Country
98110761	1998.06.12	EP

(43) Publication Date : **2000.01.05**

(43) Publication Journal Date : **2000.03.16**

(44) Accepted Journal Date : **2003.05.08**

(71) Applicant(s)
C-Chem AG

(72) Inventor(s)
Ernst Ellmerer-Mueller; Dagmar Brossner; Najib Maslouh; Horst-Dieter Ambrosi; Gerhard Jas; Gunter Fischer

(74) Agent/Attorney
GRIFFITH HACK,GPO Box 4164,SYDNEY NSW 2001

48993/99



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07K 7/64, A61K 38/13	A1	(11) International Publication Number: WO 99/65933 (43) International Publication Date: 23 December 1999 (23.12.99)
(21) International Application Number: PCT/EP99/04012 (22) International Filing Date: 10 June 1999 (10.06.99) (30) Priority Data: 98110761.8 12 June 1998 (12.06.98) EP (71) Applicant (for all designated States except US): C-CHEM AG [CH/CH]; Hauptstrasse 67, CH-4102 Binningen (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): ELLMERER-MUELLER, Ernst [AT/AT]; Institut für Organische Chemie, Leopold-Franzens-Universität Innsbruck, Innrain 52A, A-6020 Innsbruck (AT). BRÖSSNER, Dagmar [DE/AT]; Conradstrasse 6, A-6020 Innsbruck (AT). MASLOUH, Najib [AT/AT]; Holzhammerstrasse 11, A-6020 Innsbruck (AT). AMBROSI, Horst-Dieter [DE/DE]; Schwarzmeer- strasse 20, D-10319 Berlin (DE). JAS, Gerhard [DE/DE]; Falkenhausenweg 15, D-12249 Berlin (DE). <i>Fischer, Gunter</i> (74) Agent: KUNZ, Ekkehard; Agrolinz Melamin GmbH, Patentabteilung, St.-Peter-Strasse 25, A-4021 Linz (AT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i> <i>Otto Kanning Strasse 11</i> <i>D-06120 Halle/Saale</i> <i>Germany</i>
(54) Title: NOVEL CYCLOSPORINS (I) (57) Abstract Compounds of formula (I) and their pharmaceutically acceptable salts wherein the letters A to L have the meaning indicated in the description and processes for their preparation and their uses.		



Novel cyclosporins

The present invention relates to novel cyclosporins, processes for their production, their use as pharmaceuticals and pharmaceutical compositions comprising them. Furthermore, this invention discloses a novel general method for the exchange of substituents at the sarcosine residue of the cyclosporin macrocycle.

Cyclosporin A is well known for its immunosuppressive and antiinflammatory properties but many biological properties have been described in addition.

EP 0 194 972 describes cyclosporin derivatives with substituents on the sarcosine in position 3 of the macrocycle, the introduction of such substituents, as well as the immunosuppressive, antiinflammatory and antiparasitic activity of these cyclosporin derivatives. EP 0 484 281 describes cyclosporin derivatives with reduced immunosuppressive potency and activity against HIV.

The present invention discloses novel cyclosporins which can be used for the treatment of infectious diseases, of chronic inflammatory and autoimmune diseases, to prevent cardiac hypertrophy, to treat and prevent ischemia and reperfusion injury, to treat neurodegenerative diseases, and to induce processes of tissue regeneration.

A second embodiment of the present invention is a novel method to prepare cyclosporins with substituents at the sarcosine in position 3 of the macrocycle. EP 0 194 972 describes the introduction of certain substituents at the sarcosine. The method described in EP 0 194 972 involves treatment of a cyclosporin with strong base to generate a polyanion and subsequent reaction of this polyanion with electrophiles, such as disulfides, alkyl halides or other suitable alkylating agents. Halogens or sources of positive halogen can also be used, as well as aldehydes. There is no example in the prior art which describes the exchange of such a substituent by another. The present invention discloses such a method. In this novel method, a suitable substituent is first introduced into a cyclosporin polyanion and the resulting product is isolated. The substituent is subsequently activated to become a leaving group and replaced by the desired novel substituent. This novel method


allows the introduction of a wide variety of substituents into the sarcosine residue of the cyclosporin macrocycle.

The cyclosporin nomenclature and numbering systems used hereafter are those used by J. Kallen et al., "Cyclosporins: Recent Developments in Biosynthesis, Pharmacology and Biology, and Clinical Applications", Biotechnology, second edition, H.-J. Rehm and G. Reed, ed., 1997, p535-591 and are shown below:

Position numbering	Letter in Formula I	Amino acid in cyclosporin A
1	A	N-Methyl-butenyl-threonine (MeBmt)
2	B	α -aminobutyric acid (Abu)
3	C	Sarcosine (Sar)
4	D	N-Methyl-leucine (MeLeu)
5	E	Valine (Val)
6	F	N-Methyl-leucine (MeLeu)
7	G	Alanine (Ala)
8	H	(D)-Alanine ((D)-Ala)
9	I	N-Methyl-leucine (MeLeu)
10	K	N-Methyl-leucine (MeLeu)
11	L	N-Methylvaline (MeVal)




In a first aspect, the present invention provides compounds of the formula I



Formula I

and pharmaceutically acceptable salts thereof, wherein the letters A to L represent residues of the following amino acids:

A (L)-alpha-N-methylamino-beta-hydroxy acid of the formula II




Formula II

wherein R₁ is (E)-2-butenyl-1,

B alpha-amino-butyric acid, alpha-amino-valerianic acid (norvaline), threonine, or valine,

C substituted sarcosine of the formula III



Formula III

in which X is O-R₃, in which R₃ is

- (a) hydrogen, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 1-pentyl, 2-pentyl, 3-pentyl, isopentyl, tert-pentyl, neopentyl, hexyl or an isomer thereof, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, bicyclo[2.1.1]hexyl, norbornyl, bicyclo[2.2.2]octyl, vinyl, allyl, butenyl, pentenyl, pentadienyl, hexenyl, hexadienyl, formyl, acetyl,



propionyl, butyryl, pivaloyl, benzoyl or alkoxycarbonyl,
which can be substituted by hydroxy, methoxy, ethoxy, propoxy,
isopropoxy, halogen, carboxy or carbamido, or

- (b) amino, methylamino, ethylamino, n-propylamino, isopropylamino, n-
butylamino, isobutylamino, sec-butylamino, tert-butylamino, 1-
pentylamino, 2-pentylamino, 3-pentylamino, isopentylamino, tert-
pentylamino, neopentylamino, hexylamino or an isomer thereof,
N,N-dimethylamino, N-methyl-N-ethylamino, N,N-diethylamino, N-propyl-
N-methylamino, N-methyl-N-isopropylamino, dipropylamino,
diisopropylamino, N-butyl-N-methylamino or an isomer thereof, N-butyl-N-
ethylamino or an isomer thereof, N-butyl-N-propylamino or an isomer
thereof, N,N-dibutylamino or an isomer thereof, azetidine, pyrrolidine,
piperidine, morpholine, piperazine, N'-alkylpiperazine,
azabicyclo[2.1.1]hexane, azanorbornane, azabicyclo[2.2.2]octane, N-
formylamino, N-acetylamino, N-tert-butoxycarbonyl-amino, N-
benzyloxycarbonyl-amino, N-benzoyl-amino or N-phthaloyl, or

- (c) a residue of the formula VI and Y⁻ is an anion



- D is N-methyl-leucine, N-methyl-valine, or N-methyl-isoleucine,
E is valine,
F is N-methyl-leucine,
25 G is alanine,
H is glycine, (D)-alanine, (D)-serine, O-hydroxyethyl-(D)-serine,
I,K are N-methyl-leucine, and
L is N-methyl-valine.

- 30 In a second aspect, the present invention provides a process for the synthesis of
compounds of the formula I, characterised in, that in a compound of formula I, wherein
C is the substituted sarcosine of the formula III in which X is -SR₃, the -SR₃ is
exchanged by -OR₃.

- 35 In a third aspect, the present invention provides a process for the synthesis of
compounds of the formula I, characterised in, that the substituent OR₃ of formula III is
substituted by another substituent OR₃.




In a fourth aspect, the present invention provides a method of treating infectious diseases caused by bacterial, fungal or viral pathogens, or by protozoan parasites or helminthes, in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

In a fifth aspect, the present invention provides a method of treating chronic inflammation or autoimmune diseases in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

In a sixth aspect, the present invention provides a method of treating pathological conditions caused by trauma or reperfusion injury in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

In a seventh aspect, the present invention provides a method of inducing tissue regenerative processes in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

Also described herein are compounds of the formula IA




Formula IA

and their pharmaceutically acceptable salts wherein the letters A to L represent residues of the following amino acids:

A (L)-alpha-N-methylamino-beta-hydroxy acid of the formula II,






Formula II

wherein R1 is n-butyl or butenyl-1,

- B alpha-amino-butyric acid, alpha-amino-valerianic acid (norvaline), threonine, or valine,
- C substituted sarcosine of the formula III




Formula III

in which X is

S-(O)_n-R₂, in which n has the value zero, one or two and R₂ is hydrogen, unsubstituted or substituted, unbranched or branched, acyclic, monocyclic or polycyclic, saturated or unsaturated lower alkyl, substituted or unsubstituted aryl or heteroaryl, or

X is O-R₃, in which R₃ is hydrogen, unsubstituted or substituted, unbranched or branched, saturated or unsaturated, acyclic, monocyclic or polycyclic lower alkyl, or acyl, or

X is sulfonium groups of the formula IV,




Formula IV

in which R₄ and R₅ are independently selected from lower alkyl, aryl, or heteroaryl and Y⁻ is an anion, or



X is a group of the formula V,



in which R6 and R7 are independently selected from lower alkyl or aryl or form together a ring and R8 is hydrogen or substituted or unsubstituted lower alkyl and Y⁻ is an anion, or

C is a residue of the formula VI and Y⁻ is an anion,



- D N-methyl-leucine, gamma-hydroxy-N-methyl-leucine, N-methyl-valine, or N-methyl-isoleucine,
- E valine,
- F N-methyl-leucine,
- G alanine,
- H glycine, (D)-alanine, (D)-serine, O-hydroxyethyl-(D)-serine,
- I,K N-methyl-leucine, and
- L N-methyl-valine.

The 1-butenyl rest in A has preferably the trans configuration.

Examples for lower alkyl are methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 1-pentyl, 2-pentyl, 3-pentyl, isopentyl, tert-pentyl, neopentyl, hexyl and its isomers.

Examples for monocyclic lower alkyl groups are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl.

Examples for polycyclic lower alkyl groups are bicyclo[2.1.1]hexyl, norbornyl, bicyclo[2.2.2]octyl.

Examples for unsaturated lower alkyl are vinyl, allyl, butenyl, pentenyl, pentadienyl, hexenyl, hexadienyl.

Examples for substituents in these radicals are hydroxy, methoxy, ethoxy, propoxy, isopropoxy, amino, monoalkylamino, dialkylamino, acylamino, halogen, acyl, carboxy, carbamido.

Examples for monoalkylamino are methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, sec-butylamino, tert-butylamino, 1-pentylamino, 2-pentylamino, 3-pentylamino, isopentylamino, tert-pentylamino, neopentylamino, hexylamino and its isomers.

Examples for dialkylamino are N,N-dimethylamino, N-methyl-N-ethylamino, N,N-diethylamino, N-propyl-N-methylamino, N-methyl-N-isopropylamino, dipropylamino, diisopropylamino, N-butyl-N-methylamino and its isomers, N-butyl-N-ethylamino and its isomers, N-butyl-N-propylamino and its isomers, N,N-dibutylamino and its isomers.

The alkyl groups of dialkylamino may also form a ring together. Examples are azetidine, pyrrolidine, piperidine, morpholine, piperazine, N'-alkylpiperazine, azabicyclo[2.1.1]hexane, azanorbornane, azabicyclo[2.2.2]octane.

Examples for acyl are formyl, acetyl, propionyl, butyryl, pivaloyl, benzoyl, alkoxycarbonyl.

Examples for acylamino are N-formylamino, N-acetylamino, N-tert-butoxycarbonylamino, N-benzyloxycarbonylamino, N-benzoylamino, N-phthaloyl.


Examples for substituted aryl are tolyl, chlorophenyl, methoxyphenyl, aminophenyl, dimethylaminophenyl, 1-naphthyl, 2-naphthyl.

Examples for heteroaryl are thiazole, oxazole, imidazole, pyridine, pyrazole, pyrimidine, pyrazine, triazine, benzthiazole, benzoxazole, benzimidazole.

Examples of sulfonium groups are dimethylsulfonium, S-methyl-S-phenylsulfonium, S-methyl-S-allylsulfonium, S-methyl-S-carboxamidomethyl-sulfonium, S-dodecyl-S-methylsulfonium.

Examples for groups of the formula V are N-methyl-pyridinium-2-ylthio, N-methyl-pyridinium-4-ylthio, N-methyl-triazinylthio, N-methyl-benzthiazol-2-ylthio, pyridinium-2-ylthio toluenesulfonate, pyridinium-4-ylthio toluenesulfonate, pyridinium-2-ylthio methanesulfonate.

Compounds of the formula IA in which C is a sarcosine substituted by S-R₂ are prepared by forming polyanions from cyclosporins in which C is sarcosine and



reacting these polyanions with appropriate sulfur electrophiles like disulfides, thiolsulfonates, sulfonyl halides, or disulfide-derived sulfonium salts. The polyanions are in turn prepared by treating the cyclosporins in an appropriate solvent at low temperature with an excess of a strong base. Examples for strong bases are alkali amides like lithium amide, sodium amide, lithium diisopropylamide or lithium hexamethyl disilazide. Examples for inert solvents used for these reactions are tetrahydrofuran, dioxane, diethylether, methyl tert-butylether, or liquid ammonia.

Compounds of the formula IA in which C is sarcosine substituted by S-R₂ are furthermore prepared by exchanging the substituent X in compounds of formula I in which C is sarcosine substituted by X and where X is a residue of formula IV or formula V by thiols HS-R₂, in which R₂ is lower alkyl, aryl, or acyl.

A compound of the formula IA in which C is sarcosine substituted by S-R₂ in which R₂ is hydrogen is prepared by treating compounds of formula IA in which C is sarcosine substituted by S-R₂ and in which R₂ is acyl in an appropriate solvent with ammonia, hydrazine, hydroxylamine, or organic derivatives thereof, such as methylamine, benzylamine, methylhydrazine, or dimethylhydrazine. Appropriate solvents for these reactions are alcohols, such as methanol or ethanol, ethers such as diethylether, tetrahydrofuran, or dioxane, or inert aprotic solvents, such as dimethyl formamide. The resulting thiol can be isolated but is more conveniently alkylated directly by adding to these reactions an alkylating agent such as an alkyl halide or esters of alcohols with sulfuric acid, or organic sulfonic acids such as 4-toluene sulfonic acid, methane sulfonic acid, or trifluoromethane sulfonic acid. The present invention makes it therefore possible to produce compounds of the formula IA in which C is sarcosine substituted by S-R₂ in which R₂ is substituted or unsubstituted, branched or unbranched, saturated or unsaturated, acyclic, monocyclic, or polycyclic lower alkyl.

Compounds of the formula IA in which C is sarcosine substituted by SO-R₂ and by SO₂-R₂ are prepared by treating compounds of formula IA in which C is sarcosine substituted by S-R₂ with an appropriate oxidant in an inert solvent. Examples for such oxidants are hydrogen peroxide, sodium chlorate, sodium periodate, peroxyacetic acid, meta-chloroperoxybenzoic acid, or potassium persulfate. Solvents for these reactions are for example mixtures of water with organic solvents such as




tetrahydrofurane, dioxane or acetic acid, or anhydrous organic solvents such as dichloromethane, chloroform, tetrachloroethane, tetrahydrofurane, or dioxane.

Compounds of the formula I in which C is a sarcosine substituted by O-R3 can be prepared by exchanging the substituent S-R2 in compounds of formula I in which C is sarcosine substituted by S-R2. This exchange reaction is effected by metal salts which have an affinity for sulfur, such as mercuric acetate, silver acetate, copper acetate and others, but can also be effected by the presence of Bronsted or Lewis acids. The acidic nature of Bronsted acids is due to their capacity to act as proton donors. Such acids are for example sulfuric acid, toluene sulfonic or camphersulfonic acid, hydrochloric acid, but also acetic acid, formic acid and other organic carboxylic acids. Lewis acids are compounds having affinity for free electron pairs and forming coordination complexes with groups having free electron pairs. Examples for Lewis acids are boron trifluoride, titanium tetrachloride, aluminum chloride, or zinc chloride. Such Bronsted or Lewis acids or metal salts convert S-R2 or O-R3 substituents at the sarcosine position of the cyclosporin macrocycle into leaving groups, forming an intermediary cation of the formula VI which can then further react with nucleophiles present in the reaction mixture to form the desired products.

The present invention makes it therefore also possible to exchange one given O-R3 substituent for another O-R3' substituent by using the reaction conditions as described above.

A compound of formula I in which the amino acid residue of C is the cation of Formula VI is a common intermediate for these exchange reactions. This type of cation is well known to experts in the field and is analogous to the commonly accepted intermediate in the Mannich reaction. In the case of cyclosporins, however, such an intermediate has never been described and is new. Mannich reactions are used to introduce aminoalkyl residues into a wide variety of nucleophiles, such as enols, phenols, enamines, heterocycles such as indole, pyrrol, or furane. Other nucleophiles reacting with such cations are allyl and vinylsilanes and -stannanes as well as acetylenes. Therefore, cyclosporins in which the amino acid residue of C is the cation of Formula VI are an especially preferred embodiment of the present invention.



The compounds of the present invention act on enzymes called cyclophilins and inhibit their catalytic activity. Cyclophilins occur in a wide variety of different organisms, including human, yeast, bacteria, protozoa, metazoa, insects, plants, or viruses. In the case of infectious organisms, inhibition of the cyclophilin catalytic activity by compounds of the present invention often results in an inhibitory effect on the organism. Furthermore, in humans the catalytic activity of cyclophilins plays a role in many different disease situations. Inhibition of this catalytic activity is often associated to a therapeutic effect. Therefore, the compounds of the present invention can be used for the treatment of infections including that by HIV as well as fungal pathogens, protozoan and metazoan parasites. Furthermore, the compounds of the present invention can be used to treat chronic inflammatory and autoimmune diseases including but not limited to rheumatoid arthritis, psoriasis, and uveitis. In addition, the compounds of the present invention can be used to treat neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, and neuropathies.

Another use of the compounds of the present invention is protection against tissue damage associated to ischemia and reperfusion such as paralytic damage after spinal cord or head injuries or cardiac damage after myocardial infarct.

Furthermore, the compounds of the present invention induce regenerative processes such as that of hair, liver, gingiva, or nerve tissue damaged or lost due to injury or other underlying pathologies, such as damage of the optical nerve in glaucoma.

The compounds of the present invention can, together with pharmaceutically acceptable additives and/or excipients be administered either orally in the form of capsules, tablets, or drink solutions or parenterally in form of acute injections or infusions. They can also be applied locally in form of solutions, eye drops, or as gels and ointments. For topical and parenteral applications it is of special advantage that, unlike cyclosporin A, many of the compounds of the present invention have basic substituents which enable the formation of salts with physiologically acceptable acids.

The daily administered dosage depends on the structure of the medicament, the disease to be treated, and the type of formulation and is from about 1 mg to about 200 mg per kg body weight.

Examples

1. 3-(Pyridyl-2-thio)-cyclosporin:

Liquid ammonia (200 ml) is condensed in a flask under argon. Freshly cut sodium (1.78 g, 75 mmol) is added in 4 portions, followed by one crystal of ferric nitrate. The mixture is vigorously stirred for 15 minutes, at which time a dark grey color persists. A solution of cyclosporin A (5 g, 4 mmol) in tert.-butylmethyl ether (90 ml) is added over 15 minutes and the mixture is stirred at minus 40° Celsius for 1.5 hrs. Then, a solution of 2,2'-dipyridyl disulfide (7.2 g, 33 mmol) in 70 ml tert.-butylmethyl ether is added and stirring continued for 2 hrs at minus 35° minus 40°. Solid ammonium chloride (5.56 g, 104 mmol) is added, stirring continued at minus 35° for 10 minutes. After warming to room temperature, the mixture is stirred for another 2 hrs, filtered, the residue washed 3 times with tert.-butylmethyl ether, and filtrate and washings combined, extracted with 2N sodium hydroxide, brine and dried over sodium sulfate. Chromatography of the residue obtained upon evaporation on silica, eluting with diethyl ether/methanol = 96/4 yielded the title compound as two isomers.

The ¹H-NMR chemical shift of the proton at the sarcosine residue [$\delta(H3)$] for isomer A of the title compound is at 6.36 ppm, for isomer B at 7.13 ppm. The corresponding values in the ¹³C-NMR [$\delta(C3)$] are 58.81 and 58.45 ppm.

Analogous to example 1 the following disulfides were used to prepare the following products:

Example	Disulfide	Product
2	4,4'-dipyridyldisulfid	3-(Pyridyl-4-thio)-cyclosporin A
3	2,2'-dipyridyldisulfid	3-(Pyridyl-2-thio)-4-(γ -hydroxy-methylleucine)-cyclosporin
4	2,2'-dipyridyldisulfid	3-(Pyridyl-2-thio)-4-methylvaline-cyclosporin



5	2,2'-dipyridyldisulfid	3-(Pyridyl-2-thio)-2-norvaline-cyclosporin
6	2,2'-dipyridyldisulfid	3-(Pyridyl-2-thio)-2-valine-cyclosporin
7	2,2'-dipyridyldisulfid	3-(Pyridyl-2-thio)-8-(D)-serine-cyclosporin
8	diphenyldisulfid	3-(Phenylthio)-cyclosporin A
9	2,2'-dimethylaminoethylthio-	3-(2-dimethylaminoethylthio)-cyclosporin
10	2,2'-mercaptobenzthiazolyl-disulfide	3-(Mercaptobenzthiazol-2-ylthio)-cyclosporin

The products of these examples have the following physiochemical data:

Example	Product	δ (H3) (CDCl ₃)	δ (C3) (CDCl ₃)	Calc Mass	High Resolution MS
2	3-(Pyridyl-4-thio)-cyclosporin A				
	isomer A	6.36	60.19	1311.84	1311.8850
	isomer B	7.13	58.46		
3	3-(Pyridyl-2-thio)-4-(γ -hydroxy-methylleucine)-cyclosporin	7.12	58.58	1327.84	1327.7922
4	3-(Pyridyl-2-thio)-4-methylvaline-cyclosporin	7.12	58.45	1297.83	1297.7507
5	3-(Pyridyl-2-thio)-2-norvaline-cyclosporin	7.12	58.77	1325.86	1325.8316
6	3-(Pyridyl-2-thio)-2-valine-cyclosporin				
	isomer A	7.08	58.81	1325.86	1325.8517
	isomer B	7.33	58.45		

7	3-(Pyridyl-2-thio)-8-(D)-serine-cyclosporin	7.11	58.12		
	Isomer A	7.60	57.63	1327.84	1327.8436
	Isomer B				
8	3-(Phenylthio)-cyclosporin A	6.18	63.62	1310.85	1310.8732
9	3-(2-dimethylaminoethylthio)-cyclosporin	5.77	59.32	1305.89	1305.8762
10	3-(Mercptobenzthiazol-2-ylthio)-cyclosporin,				
	Isomer A	7.0	63.00	1367.86	1366.8120
	Isomer B	7.56	60.90		

11. 3'-Acetoxy-3-phenylthio-cyclosporin

To a stirred solution of 700 mg the product of example 8 and 300 mg of 4-dimethylamino pyridine in 3 ml pyridine was added 0.5 ml acetic anhydride. The solution was stirred at ambient temperature for 36 hrs and then diluted with 20 ml cold water. The mixture was extracted three times with ethyl acetate, the organic extracts were combined and washed successively with 1N sulfuric acid, water, brine and dried over sodium sulfate. The residue obtained from evaporation to dryness was chromatographed on silicagel using ethyl acetate as eluent. The title product was obtained as yellow foam.

12. 3',3-Diacetoxy-cyclosporin

The product of example 11 (100 mg) was dissolved in 2 ml acetic acid, mercuric acetate (100 mg) was added and the mixture was heated to 50° for 3 hrs. After evaporation to dryness the residue was taken up in ethyl acetate and washed with sodium bicarbonate solution. The organic phase was dried over sodium sulfate and evaporated to dryness. Chromatography of the residue on silica gel yielded 45 mg of the title compound. The characteristic

NMR signals of this compound in deuteriochloroform are at 6.80 ppm (sarcosine H) and 74.53 ppm (sarcosine C).

13. 3-Acetoxy-cyclosporin: To a stirred solution of 500 mg of the product of example 8 in 3 ml acetic acid was added silver acetate (100 mg) and the mixture was warmed to 50° for 20 hrs. After evaporation to dryness the residue was taken up in ethyl acetate and washed with sodium bicarbonate solution. The organic phase was dried over sodium sulfate and evaporated to dryness. Chromatography of the residue on silica gel yielded 45 mg of the title compound. The characteristic NMR signals of this compound in deuteriochloroform are at 6.82 ppm (sarcosine H) and 74.40 ppm (sarcosine C).

14. 3-Methoxycyclosporin

The product of example 1 (131 mg) was dissolved in 2 ml methanol, camphersulfonic acid (25 mg) was added and the mixture heated to 50° for 5 hrs. After addition of aqueous ammonia (0.5 ml) and evaporation to dryness the residue was taken up in ethyl acetate and washed with sodium bicarbonate solution. The organic phase was dried over sodium sulfate and evaporated to dryness. Chromatography of the residue on silica gel yielded 45 mg of the title compound. The characteristic NMR signals of this compound in deuteriochloroform are at 5.84 ppm (sarcosine H) and 83.30 ppm (sarcosine C).

This compound is also prepared under identical conditions using the product of example 12 as starting material in place of the product of example 1.

Analogous to example 14 the following alcohols were used to prepare the following products:

Example	Alcohol	Product	$\delta(\text{H3})$ (CDCl ₃)	$\delta(\text{C3})$ (CDCl ₃)
15	Ethanol	3-Ethoxycyclosporin	5.93	81.58
16	Isopropyl alcohol	3-Isopropyloxycyclosporin	6.02	79.50
17.	[D-Sar-(tert.-butoxy) ³]-cyclosporin: Following the procedure of example 14 but using the product of example 10 instead of the product of example 1 and using tert.-butanol instead of methanol, the title compound is obtained. The characteristic NMR signals of this compound in deuteriochloroform are at 6.09 ppm (sarcosine H) and 76.00 ppm (sarcosine C). The calculated mass is 1274.74, the mass determined by high resolution MS is 1273.898884.			
18.	[D-Sar-(allyloxy) ³]-cyclosporin: Following the procedure of example 17 and using allyl alcohol in place of tert.-butanol, the title compound is obtained. The characteristic NMR signals of this compound in deuteriochloroform are at 5.96 ppm (sarcosine H) and 80.80 ppm (sarcosine C). The calculated mass is 1258.69, the mass determined by high resolution MS is 1257.867583.			
19.	[D-Sar-(hydroxy) ³]-cyclosporin: The product of example 1 (130 mg) was dissolved in 2 ml tetrahydrofurane, 1N sulfuric acid (1 ml) was added and the mixture heated to 50° for 5 hrs. After addition of aqueous ammonia (0.5 ml), ethyl acetate was added, the mixture was vigorously shaken and, after separation of the organic layer it was washed with sodium bicarbonate solution. The organic phase was dried over sodium sulfate and evaporated to dryness. Chromatography of the residue on silica gel yielded 55 mg of the title compound. The characteristic NMR signal of this compound in			

- deuteriochloroform is at 6.96 ppm (sarcosine H). The calculated mass is 1218.63, the mass determined by high resolution MS is 1217.836248.
20. [D-Sar-(acetylthio)³]-cyclosporin: The product of example 1 (500 mg) was dissolved in 2 ml tetrahydrofurane and 2 ml thioacetic acid. Camphersulfonic acid (200 mg) was added and the mixture heated to 50° for 5 hrs. After addition of aqueous ammonia (1 ml), ethyl acetate was added, the mixture was vigorously shaken and, after separation of the organic layer it was washed with sodium bicarbonate solution. The organic phase was dried over sodium sulfate and evaporated to dryness. Chromatography of the residue on silica gel yielded 255 mg of the title compound. The characteristic NMR signals of this compound in deuteriochloroform are at 6.50 ppm (sarcosine H) and 57.60 ppm (sarcosine C). The calculated mass is 1276.73, the mass determined by high resolution MS is 1276.824005.
21. [D-Sar-(methylthio)³]-cyclosporin: Under argon atmosphere, the product of example 19 (120 mg) was dissolved in 1 ml ethanol. N,N-dimethylhydrazine (60 mg) was added and the mixture stirred at room temperature for 5 hrs. After addition of methyl iodide (0.5 ml) stirring was continued over night. The mixture was concentrated to dryness and the residue partitioned between ethyl acetate and 1N sulfuric acid. The organic layer was washed with sodium bicarbonate solution and brine, dried over anhydrous sodium sulfate and evaporated to dryness. Chromatography of the residue on silica gel yielded 75 mg of the title compound. The compound has the following optical rotation: $[\alpha]^{20}_D = -215^\circ$, c = 1, CHCl₃. A value of $[\alpha]^{20}_D = -215^\circ$, c = 1, CHCl₃, has been reported for this compound.
22. [D-Sar-(2-(N-Boc-3-aminopropoxy))³]-cyclosporin: The product of example 10 (140 mg) was dissolved in 5 ml dry tetrahydrofurane, 175 mg N-Boc-3-aminopropanol and Camphersulfonic acid (30 mg) were added and the mixture heated to 50° for 12 hrs. The mixture was evaporated to dryness, the residue taken up in ethyl acetate and saturated sodium bicarbonate solution. The layers were separated, the organic phase was dried over sodium sulfate and evaporated to dryness. Repeated chromatography of the residue on silica

gel yielded 75 mg of the title compound and 23 mg [D-Sar-(2-(3-aminopropoxy))³]-cyclosporin. The characteristic NMR signal of the title compound in deuteriochloroform is at 5.97 ppm (sarcosine H), that of [D-Sar-(2-(3-aminopropoxy))³]-cyclosporin at 5.91 ppm.

According to the procedure described in example 22 and using the following alcohols the following products are obtained:

Example	Alcohol	Product
23	N-Boc-2-aminoethanol	[D-Sar-(2-(N-Boc-3-aminoethoxy)) ³]-cyclosporin
24		[D-Sar-(2-(3-aminoethoxy)) ³]-cyclosporin
25	N-Boc-5-aminopentanol	[D-Sar-(2-(N-Boc-3-aminopentoxy)) ³]-cyclosporin
26		[D-Sar-(2-(3-aminopentoxy)) ³]-cyclosporin
27	N,N-dimethyl-aminoethanol	[D-Sar-(2-(N,N-dimethylaminoethoxy)) ³]-cyclosporin
28	N,N-diethylaminoethanol	[D-Sar-(2-(N,N-diethylaminoethoxy)) ³]-cyclosporin
29	N-methyl-N-(1,1-dimethyl-ethyl)-aminoethanol	[D-Sar-(2-(N-methyl-N-(1,1-dimethyl-ethyl)-2-aminoethoxy)) ³]-cyclosporin
30	N-phthaloyl-aminoethanol	[D-Sar-(2-(N-phthalimido-2-aminoethoxy)) ³]-cyclosporin
31	N-Boc-2-amino-3-phenylpropanol	[D-Sar-(2-(N-Boc-2-amino-3-phenyl-propoxy)) ³]-cyclosporin
32	N-Boc-2-hydroxymethyl-pyrrolidine	[D-Sar-(2-(N-Boc-pyrrolidine-2-yl)-methoxy) ³]-cyclosporin
33		[D-Sar-(2-(pyrrolidine-2-yl)-methoxy) ³]-cyclosporin
34	N-methyl-4-hydroxypiperidine	[D-Sar-(2-(N-methyl-piperidine-4-yl)-oxy) ³]-cyclosporin
35	N-methyl-3-hydroxyazetidine	[D-Sar-(2-(N-methyl-azetidine-3-yl)-oxy) ³]-cyclosporin

The characteristic NMR signals of these compounds are as follows:

Example	Product	δ (H3) (CDCl ₃)	δ (C3) (CDCl ₃)
23	[D-Sar-(2-(N-Boc-3-aminoethoxy)) ³]-cyclosporin	5.96	82.01
24	[D-Sar-(2-(3-aminoethoxy)) ³]-cyclosporin	5.92	81.53
25	[D-Sar-(2-(N-Boc-3-aminopentoxy)) ³]-cyclosporin	5.95	81.76
26	[D-Sar-(2-(3-aminopentoxy)) ³]-cyclosporin	5.94	81.73
27	[D-Sar-(2-(N,N-dimethylaminoethoxy)) ³]-cyclosporin	5.89	80.63
28	[D-Sar-(2-(N,N-diethylaminoethoxy)) ³]-cyclosporin	5.88	80.54
29	[D-Sar-(2-(N-methyl-N-(1,1-dimethyl-ethyl)-2-aminoethoxy)) ³]-cyclosporin	5.84	80.13
30	[D-Sar-(2-(N-phthalimido-2-aminoethoxy)) ³]-cyclosporin	5.99	82.12
31	[D-Sar-(2-(N-Boc-2-amino-3-phenyl-propoxy)) ³]-cyclosporin	5.96	81.89
32	[D-Sar-(2-(N-Boc-pyrrolidine-2-yl)-methoxy) ³]-cyclosporin	5.87	79.46
33	[D-Sar-(2-(pyrrolidine-2-yl)-methoxy) ³]-cyclosporin	5.85	79.32
34	[D-Sar-(2-(N-methyl-piperidine-4-yl)-oxy) ³]-cyclosporin	5.85	79.32
35	[D-Sar-(2-(N-methyl-azetidine-3-yl)-oxy) ³]-cyclosporin	6.03	78.50

Using the conditions of example 1 the following compounds can be obtained:

3-(2-Hydroxyethyl)thio-cyclosporin
 3-(3-Hydroxypropyl)thio-cyclosporin
 3-(2-Aminoethyl)thio-cyclosporin
 3-(2-Methylaminoethyl)thio-cyclosporin
 3-(2-Ethylaminoethyl)thio-cyclosporin
 3-(2-Ethyl-N-methylaminoethyl)thio-cyclosporin

3-(2-Diethylaminoethyl)thio-cyclosporin
3-(2-n-Propylaminoethyl)thio-cyclosporin
3-(2-Isopropylaminoethyl)thio-cyclosporin
3-(2-Cyclopropylaminoethyl)thio-cyclosporin
3-(2-n-Propyl-methylaminoethyl)thio-cyclosporin
3-(2-n-Propyl-ethylaminoethyl)thio-cyclosporin
3-(2-Methyl-isopropylaminoethyl)thio-cyclosporin
3-(2-Methylcyclopropylaminoethyl)thio-cyclosporin
3-(2-Ethyl-isopropylaminoethyl)thio-cyclosporin
3-(2-Diisopropylaminoethyl)thio-cyclosporin
3-(2-n-Propyl-isopropylaminoethyl)thio-cyclosporin
3-(2-n-Butylaminoethyl)thio-cyclosporin
3-(2-sec-Butylaminoethyl)thio-cyclosporin
3-(2-Isobutylaminoethyl)thio-cyclosporin
3-(2-tert-Butylaminoethyl)thio-cyclosporin
3-(2-n-Butyl-methylaminoethyl)thio-cyclosporin
3-(2-n-Butyl-ethylaminoethyl)thio-cyclosporin
3-(2-n-Butyl-isopropylaminoethyl)thio-cyclosporin
3-(2-sec-Butyl-methylaminoethyl)thio-cyclosporin
3-(2-sec-Butyl-ethylaminoethyl)thio-cyclosporin
3-(2-sec-Butyl-isopropylaminoethyl)thio-cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-cyclosporin
3-(2-Azetidinoethyl)thio-cyclosporin
3-(2-pyrrolidinoethyl)thio-cyclosporin
3-(2-piperidinoethyl)thio-cyclosporin
3-(2-morpholinoethyl)thio-cyclosporin
3-(2-piperazinoethyl)thio-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-cyclosporin
3-(3-azetidiny)thio-cyclosporin

3-(N-methyl-3-azetidiny)thio-cyclosporin
3-(N-methyl-3-azetidiny)thio-cyclosporin
3-(N-isopropyl-3-azetidiny)thio-cyclosporin
3-(3-pyrrolidiny)thio-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-cyclosporin
3-(N-isopropyl-3-pyrrolidiny)thio-cyclosporin
3-(4-piperidiny)thio-cyclosporin
3-(N-methyl-4-piperidiny)thio-cyclosporin
3-(N-methyl-4-piperidiny)thio-cyclosporin
3-(N-isopropyl-4-piperidiny)thio-cyclosporin

3-(2-Hydroxyethyl)thio-2-valine-cyclosporin
3-(3-Hydroxypropyl)thio-2-valine-cyclosporin
3-(2-Aminoethyl)thio-2-valine-cyclosporin
3-(2-Methylaminoethyl)thio-2-valine-cyclosporin
3-(2-Ethylaminoethyl)thio-2-valine-cyclosporin
3-(2-Ethyl-N-methylaminoethyl)thio-2-valine-cyclosporin
3-(2-Diethylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Propylaminoethyl)thio-2-valine-cyclosporin
3-(2-Isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-Cyclopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Propyl-methylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Propyl-ethylaminoethyl)thio-2-valine-cyclosporin
3-(2-Methyl-isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-Methylcyclopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-Ethyl-isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-Diisopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Propyl-isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Butylaminoethyl)thio-2-valine-cyclosporin
3-(2-sec-Butylaminoethyl)thio-2-valine-cyclosporin
3-(2-Isobutylaminoethyl)thio-2-valine-cyclosporin

3-(2-tert-Butylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Butyl-methylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Butyl-ethylaminoethyl)thio-2-valine-cyclosporin
3-(2-n-Butyl-isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-sec-Butyl-methylaminoethyl)thio-2-valine-cyclosporin
3-(2-sec-Butyl-ethylaminoethyl)thio-2-valine-cyclosporin
3-(2-sec-Butyl-isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-2-valine-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-2-valine-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-2-valine-cyclosporin
3-(2-Azetidinoethyl)thio-2-valine-cyclosporin
3-(2-pyrrolidinoethyl)thio-2-valine-cyclosporin
3-(2-piperidinoethyl)thio-2-valine-cyclosporin
3-(2-morpholinoethyl)thio-2-valine-cyclosporin
3-(2-piperazinoethyl)thio-2-valine-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-2-valine-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-2-valine-cyclosporin
3-(3-azetidiny)thio-2-valine-cyclosporin
3-(N-methyl-3-azetidiny)thio-2-valine-cyclosporin
3-(N-methyl-3-azetidiny)thio-2-valine-cyclosporin
3-(N-isopropyl-3-azetidiny)thio-2-valine-cyclosporin
3-(3-pyrrolidiny)thio-2-valine-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-2-valine-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-2-valine-cyclosporin
3-(N-isopropyl-3-pyrrolidiny)thio-2-valine-cyclosporin
3-(4-piperidiny)thio-2-valine-cyclosporin
3-(N-methyl-4-piperidiny)thio-2-valine-cyclosporin
3-(N-methyl-4-piperidiny)thio-2-valine-cyclosporin
3-(N-isopropyl-4-piperidiny)thio-2-valine-cyclosporin

3-(2-Hydroxyethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(3-Hydroxypropyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin

3-(2-Aminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Methylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Ethylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Ethyl-N-methylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Diethylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Propylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Cyclopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Propyl-methylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Propyl-ethylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Methyl-isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Methylcyclopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-
cyclosporin
3-(2-Ethyl-isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Diisopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Propyl-isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-
cyclosporin
3-(2-n-Butylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-sec-Butylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-Isobutylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-tert-Butylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Butyl-methylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Butyl-ethylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-n-Butyl-isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-sec-Butyl-methylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-sec-Butyl-ethylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-sec-Butyl-isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-
cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-4-(gamma-hydroxy-methylleucine)-
cyclosporin

3-(2-Azetidinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-pyrrolidinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-piperidinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-morpholinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-piperazinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-azetidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-methyl-3-azetidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-methyl-3-azetidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-isopropyl-3-azetidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-pyrrolidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-isopropyl-3-pyrrolidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(4-piperidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-methyl-4-piperidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-methyl-4-piperidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(N-isopropyl-4-piperidiny)thio-4-(gamma-hydroxy-methyllleucine)-cyclosporin

3-(2-Hydroxyethyl)thio-2-norvaline-cyclosporin
3-(3-Hydroxypropyl)thio-2-norvaline-cyclosporin
3-(2-Aminoethyl)thio-2-norvaline-cyclosporin
3-(2-Methylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Ethylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Ethyl-N-methylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Diethylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Propylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Cyclopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Propyl-methylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Propyl-ethylaminoethyl)thio-2-norvaline-cyclosporin

3-(2-Methyl-isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Methylcyclopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Ethyl-isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Diisopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Propyl-isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Butylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-sec-Butylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Isobutylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-tert-Butylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Butyl-methylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Butyl-ethylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-n-Butyl-isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-sec-Butyl-methylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-sec-Butyl-ethylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-sec-Butyl-isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-2-norvaline-cyclosporin
3-(2-Azetidinoethyl)thio-2-norvaline-cyclosporin
3-(2-pyrrolidinoethyl)thio-2-norvaline-cyclosporin
3-(2-piperidinoethyl)thio-2-norvaline-cyclosporin
3-(2-morpholinoethyl)thio-2-norvaline-cyclosporin
3-(2-piperazinoethyl)thio-2-norvaline-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-2-norvaline-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-2-norvaline-cyclosporin
3-(3-azetidiny)thio-2-norvaline-cyclosporin
3-(N-methyl-3-azetidiny)thio-2-norvaline-cyclosporin
3-(N-methyl-3-azetidiny)thio-2-norvaline-cyclosporin
3-(N-isopropyl-3-azetidiny)thio-2-norvaline-cyclosporin
3-(3-pyrrolidiny)thio-2-norvaline-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-2-norvaline-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-2-norvaline-cyclosporin

3-(N-isopropyl-3-pyrrolidinyl)thio-2-norvaline-cyclosporin

3-(4-piperidinyl)thio-2-norvaline-cyclosporin

3-(N-methyl-4-piperidinyl)thio-2-norvaline-cyclosporin

3-(N-methyl-4-piperidinyl)thio-2-norvaline-cyclosporin

3-(N-isopropyl-4-piperidinyl)thio-2-norvaline-cyclosporin

3-(2-Hydroxyethyl)thio-4-methylvaline-cyclosporin

3-(3-Hydroxypropyl)thio-4-methylvaline-cyclosporin

3-(2-Aminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Methylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Ethylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Ethyl-N-methylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Diethylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Propylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Isopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Cyclopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Propyl-methylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Propyl-ethylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Methyl-isopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Methylcyclopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Ethyl-isopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Diisopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Propyl-isopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Butylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-sec-Butylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-Isobutylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-tert-Butylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Butyl-methylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Butyl-ethylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-n-Butyl-isopropylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-sec-Butyl-methylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-sec-Butyl-ethylaminoethyl)thio-4-methylvaline-cyclosporin

3-(2-sec-Butyl-isopropylaminoethyl)thio-4-methylvaline-cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-4-methylvaline-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-4-methylvaline-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-4-methylvaline-cyclosporin
3-(2-Azetidinoethyl)thio-4-methylvaline-cyclosporin
3-(2-pyrrolidinoethyl)thio-4-methylvaline-cyclosporin
3-(2-piperidinoethyl)thio-4-methylvaline-cyclosporin
3-(2-morpholinoethyl)thio-4-methylvaline-cyclosporin
3-(2-piperazinoethyl)thio-4-methylvaline-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-4-methylvaline-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-4-methylvaline-cyclosporin
3-(3-azetidiny)thio-4-methylvaline-cyclosporin
3-(N-methyl-3-azetidiny)thio-4-methylvaline-cyclosporin
3-(N-methyl-3-azetidiny)thio-4-methylvaline-cyclosporin
3-(N-isopropyl-3-azetidiny)thio-4-methylvaline-cyclosporin
3-(3-pyrrolidiny)thio-4-methylvaline-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-4-methylvaline-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-4-methylvaline-cyclosporin
3-(N-isopropyl-3-pyrrolidiny)thio-4-methylvaline-cyclosporin
3-(4-piperidiny)thio-4-methylvaline-cyclosporin
3-(N-methyl-4-piperidiny)thio-4-methylvaline-cyclosporin
3-(N-methyl-4-piperidiny)thio-4-methylvaline-cyclosporin
3-(N-isopropyl-4-piperidiny)thio-4-methylvaline-cyclosporin

3-(2-Hydroxyethyl)thio-4-methylisoleucine-cyclosporin
3-(3-Hydroxypropyl)thio-4-methylisoleucine-cyclosporin
3-(2-Aminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Methylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Ethylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Ethyl-N-methylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Diethylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Propylaminoethyl)thio-4-methylisoleucine-cyclosporin

3-(2-Isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Cyclopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Propyl-methylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Propyl-ethylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Methyl-isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Methylcyclopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Ethyl-isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Diisopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Propyl-isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Butylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-sec-Butylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Isobutylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-tert-Butylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Butyl-methylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Butyl-ethylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-n-Butyl-isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-sec-Butyl-methylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-sec-Butyl-ethylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-sec-Butyl-isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-Azetidinoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-pyrrolidinoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-piperidinoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-morpholinoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-piperazinoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-4-methylisoleucine-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-4-methylisoleucine-cyclosporin
3-(3-azetidiny)thio-4-methylisoleucine-cyclosporin
3-(N-methyl-3-azetidiny)thio-4-methylisoleucine-cyclosporin
3-(N-methyl-3-azetidiny)thio-4-methylisoleucine-cyclosporin

3-(N-isopropyl-3-azetidiny)thio-4-methylisoleucine-cyclosporin
3-(3-pyrrolidiny)thio-4-methylisoleucine-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-4-methylisoleucine-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-4-methylisoleucine-cyclosporin
3-(N-isopropyl-3-pyrrolidiny)thio-4-methylisoleucine-cyclosporin
3-(4-piperidiny)thio-4-methylisoleucine-cyclosporin
3-(N-methyl-4-piperidiny)thio-4-methylisoleucine-cyclosporin
3-(N-methyl-4-piperidiny)thio-4-methylisoleucine-cyclosporin
3-(N-isopropyl-4-piperidiny)thio-4-methylisoleucine-cyclosporin

3-(2-Hydroxyethyl)thio-8-(D)-serine-cyclosporin
3-(3-Hydroxypropyl)thio-8-(D)-serine-cyclosporin
3-(2-Aminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Methylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Ethylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Ethyl-N-methylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Diethylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Propylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Cyclopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Propyl-methylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Propyl-ethylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Methyl-isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Methylcyclopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Ethyl-isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Diisopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Propyl-isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Butylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-sec-Butylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Isobutylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-tert-Butylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Butyl-methylaminoethyl)thio-8-(D)-serine-cyclosporin

3-(2-n-Butyl-ethylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-n-Butyl-isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-sec-Butyl-methylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-sec-Butyl-ethylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-sec-Butyl-isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-tert-Butyl-methylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-tert-Butyl-ethylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-tert-Butyl-isopropylaminoethyl)thio-8-(D)-serine-cyclosporin
3-(2-Azetidinoethyl)thio-8-(D)-serine-cyclosporin
3-(2-pyrrolidinoethyl)thio-8-(D)-serine-cyclosporin
3-(2-piperidinoethyl)thio-8-(D)-serine-cyclosporin
3-(2-morpholinoethyl)thio-8-(D)-serine-cyclosporin
3-(2-piperazinoethyl)thio-8-(D)-serine-cyclosporin
3-(2-N-methylpiperazinoethyl)thio-8-(D)-serine-cyclosporin
3-(2-N-tert-butylpiperazinoethyl)thio-8-(D)-serine-cyclosporin
3-(3-azetidiny)thio-8-(D)-serine-cyclosporin
3-(N-methyl-3-azetidiny)thio-8-(D)-serine-cyclosporin
3-(N-methyl-3-azetidiny)thio-8-(D)-serine-cyclosporin
3-(N-isopropyl-3-azetidiny)thio-8-(D)-serine-cyclosporin
3-(3-pyrrolidiny)thio-8-(D)-serine-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-8-(D)-serine-cyclosporin
3-(N-methyl-3-pyrrolidiny)thio-8-(D)-serine-cyclosporin
3-(N-isopropyl-3-pyrrolidiny)thio-8-(D)-serine-cyclosporin
3-(4-piperidiny)thio-8-(D)-serine-cyclosporin
3-(N-methyl-4-piperidiny)thio-8-(D)-serine-cyclosporin
3-(N-methyl-4-piperidiny)thio-8-(D)-serine-cyclosporin
3-(N-isopropyl-4-piperidiny)thio-8-(D)-serine-cyclosporin

3-(2-Hydroxyethoxy)-cyclosporin
3-(3-Hydroxypropoxy)-cyclosporin
3-(2-Chloroethoxy)-cyclosporin
3-(2-Bromoethoxy)-cyclosporin

3-(2-Hydroxyethoxy)-cyclosporin
3-(3-Chloropropoxy)-cyclosporin
3-(3-Bromopropoxy)-cyclosporin
3-(3-Hydroxypropoxy)-cyclosporin
3-(3-Chlorobutoxy)-cyclosporin
3-(3-Bromobutoxy)-cyclosporin
3-(2-Aminoethoxy)-cyclosporin
3-(2-Methylaminoethoxy)-cyclosporin
3-(2-Dimethylaminoethoxy)-cyclosporin
3-(2-Ethylaminoethoxy)-cyclosporin
3-(2-Ethyl-N-methylaminoethoxy)-cyclosporin
3-(2-Diethylaminoethoxy)-cyclosporin
3-(2-n-Propylaminoethoxy)-cyclosporin
3-(2-Isopropylaminoethoxy)-cyclosporin
3-(2-Cyclopropylaminoethoxy)-cyclosporin
3-(2-n-Propyl-methylaminoethoxy)-cyclosporin
3-(2-n-Propyl-ethylaminoethoxy)-cyclosporin
3-(2-Methyl-isopropylaminoethoxy)-cyclosporin
3-(2-Cyclopropylaminoethoxy)-cyclosporin
3-(2-Ethyl-isopropylaminoethoxy)-cyclosporin
3-(2-Diisopropylaminoethoxy)-cyclosporin
3-(2-n-Propyl-isopropylaminoethoxy)-cyclosporin
3-(2-n-Butylaminoethoxy)-cyclosporin
3-(2-sec-Butylaminoethoxy)-cyclosporin
3-(2-Isobutylaminoethoxy)-cyclosporin
3-(2-tert-Butylaminoethoxy)-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-cyclosporin

3-(2-tert-Butyl-methylaminoethoxy)-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-cyclosporin
3-(2-Azetidinoethoxy)-cyclosporin
3-(2-pyrrolidinoethoxy)-cyclosporin
3-(2-piperidinoethoxy)-cyclosporin
3-(2-morpholinoethoxy)-cyclosporin
3-(2-piperazinoethoxy)-cyclosporin
3-(2-N-methylpiperazinoethoxy)-cyclosporin
3-(2-N-methylpiperazinoethoxy)-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-cyclosporin
3-(3-azetidinyloxy)-cyclosporin
3-(N-methyl-3-azetidinyloxy)-cyclosporin
3-(N-isopropyl-3-azetidinyloxy)-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-cyclosporin
3-(3-pyrrolidinyloxy)-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-cyclosporin
3-(N-tert-butyl-3-pyrrolidinyloxy)-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-cyclosporin
3-(4-piperidinyloxy)-cyclosporin
3-(N-methyl-4-piperidinyloxy)-cyclosporin
3-(N-isopropyl-4-piperidinyloxy)-cyclosporin
3-(N-tert-butyl-4-piperidinyloxy)-cyclosporin

3-(2-Hydroxyethoxy)-2-valine-cyclosporin
3-(3-Hydroxypropoxy)-2-valine-cyclosporin
3-(2-Chloroethoxy)-2-valine-cyclosporin
3-(2-Bromoethoxy)-2-valine-cyclosporin
3-(2-Hydroxyethoxy)-2-valine-cyclosporin
3-(3-Chloropropoxy)-2-valine-cyclosporin
3-(3-Bromopropoxy)-2-valine-cyclosporin
3-(3-Hydroxypropoxy)-2-valine-cyclosporin

3-(3-Chlorobutoxy)-2-valine-cyclosporin
3-(3-Bromobutoxy)-2-valine-cyclosporin
3-(2-Aminoethoxy)-2-valine-cyclosporin
3-(2-Methylaminoethoxy)-2-valine-cyclosporin
3-(2-Dimethylaminoethoxy)-2-valine-cyclosporin
3-(2-Ethylaminoethoxy)-2-valine-cyclosporin
3-(2-Ethyl-N-methylaminoethoxy)-2-valine-cyclosporin
3-(2-Diethylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Propylaminoethoxy)-2-valine-cyclosporin
3-(2-Isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-Cyclopropylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Propyl-methylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Propyl-ethylaminoethoxy)-2-valine-cyclosporin
3-(2-Methyl-isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-Cyclopropylaminoethoxy)-2-valine-cyclosporin
3-(2-Ethyl-isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-Diisopropylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Propyl-isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Butylaminoethoxy)-2-valine-cyclosporin
3-(2-sec-Butylaminoethoxy)-2-valine-cyclosporin
3-(2-Isobutylaminoethoxy)-2-valine-cyclosporin
3-(2-tert-Butylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-2-valine-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-2-valine-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-2-valine-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-tert-Butyl-methylaminoethoxy)-2-valine-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-2-valine-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-2-valine-cyclosporin
3-(2-Azetidinoethoxy)-2-valine-cyclosporin
3-(2-pyrrolidinoethoxy)-2-valine-cyclosporin

3-(2-piperidinoethoxy)-2-valine-cyclosporin
3-(2-morpholinoethoxy)-2-valine-cyclosporin
3-(2-piperazinoethoxy)-2-valine-cyclosporin
3-(2-N-methylpiperazinoethoxy)-2-valine-cyclosporin
3-(2-N-methylpiperazinoethoxy)-2-valine-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-2-valine-cyclosporin
3-(3-azetidinyloxy)-2-valine-cyclosporin
3-(N-methyl-3-azetidinyloxy)-2-valine-cyclosporin
3-(N-isopropyl-3-azetidinyloxy)-2-valine-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-2-valine-cyclosporin
3-(3-pyrrolidinyloxy)-2-valine-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-2-valine-cyclosporin
3-(N-tert-butyl-3-pyrrolidinyloxy)-2-valine-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-2-valine-cyclosporin
3-(4-piperidinyloxy)-2-valine-cyclosporin
3-(N-methyl-4-piperidinyloxy)-2-valine-cyclosporin
3-(N-isopropyl-4-piperidinyloxy)-2-valine-cyclosporin
3-(N-tert-butyl-4-piperidinyloxy)-2-valine-cyclosporin

3-(2-Hydroxyethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-Hydroxypropoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Chloroethoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Bromoethoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Hydroxyethoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-Chloropropoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-Bromopropoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-Hydroxypropoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-Chlorobutoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(3-Bromobutoxy)- 4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Aminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Methylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Dimethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin

3-(2-Ethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Ethyl-N-methylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Diethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Propylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Cyclopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Propyl-methylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Propyl-ethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Methyl-isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Cyclopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Ethyl-isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Diisopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Propyl-isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Butylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-sec-Butylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Isobutylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-tert-Butylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-tert-Butyl-methylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-Azetidinoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-pyrrolidinoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-piperidinoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-morpholinoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-piperazinoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin
3-(2-N-methylpiperazinoethoxy)-4-(gamma-hydroxy-methyllleucine)-cyclosporin

3-(2-N-methylpiperazinoethoxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(3-azetidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-methyl-3-azetidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-isopropyl-3-azetidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(3-pyrrolidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-tert-butyl-3-pyrrolidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(4-piperidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-methyl-4-piperidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-isopropyl-4-piperidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin
3-(N-tert-butyl-4-piperidinyloxy)-4-(gamma-hydroxy-methylleucine)-cyclosporin

3-(2-Hydroxyethoxy)-2-norvaline-cyclosporin
3-(3-Hydroxypropoxy)-2-norvaline-cyclosporin
3-(2-Chloroethoxy)-2-norvaline-cyclosporin
3-(2-Bromoethoxy)-2-norvaline-cyclosporin
3-(2-Hydroxyethoxy)-2-norvaline-cyclosporin
3-(3-Chloropropoxy)-2-norvaline-cyclosporin
3-(3-Bromopropoxy)-2-norvaline-cyclosporin
3-(3-Hydroxypropoxy)-2-norvaline-cyclosporin
3-(3-Chlorobutoxy)-2-norvaline-cyclosporin
3-(3-Bromobutoxy)-2-norvaline-cyclosporin
3-(2-Aminoethoxy)-2-norvaline-cyclosporin
3-(2-Methylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Dimethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Ethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Ethyl-N-methylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Diethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Propylaminoethoxy)-2-norvaline-cyclosporin

3-(2-Isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Cyclopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Propyl-methylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Propyl-ethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Methyl-isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Cyclopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Ethyl-isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Diisopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Propyl-isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Butylaminoethoxy)-2-norvaline-cyclosporin
3-(2-sec-Butylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Isobutylaminoethoxy)-2-norvaline-cyclosporin
3-(2-tert-Butylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-2-norvaline-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-tert-Butyl-methylaminoethoxy)-2-norvaline-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-2-norvaline-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-2-norvaline-cyclosporin
3-(2-Azetidinoethoxy)-2-norvaline-cyclosporin
3-(2-pyrrolidinoethoxy)-2-norvaline-cyclosporin
3-(2-piperidinoethoxy)-2-norvaline-cyclosporin
3-(2-morpholinoethoxy)-2-norvaline-cyclosporin
3-(2-piperazinoethoxy)-2-norvaline-cyclosporin
3-(2-N-methylpiperazinoethoxy)-2-norvaline-cyclosporin
3-(2-N-methylpiperazinoethoxy)-2-norvaline-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-2-norvaline-cyclosporin
3-(3-azetidinyloxy)-2-norvaline-cyclosporin
3-(N-methyl-3-azetidinyloxy)-2-norvaline-cyclosporin

3-(N-isopropyl-3-azetidinyloxy)-2-norvaline-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-2-norvaline-cyclosporin
3-(3-pyrrolidinyloxy)-2-norvaline-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-2-norvaline-cyclosporin
3-(N-tert-butyl-3-pyrrolidinyloxy)-2-norvaline-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-2-norvaline-cyclosporin
3-(4-piperidinyloxy)-2-norvaline-cyclosporin
3-(N-methyl-4-piperidinyloxy)-2-norvaline-cyclosporin
3-(N-isopropyl-4-piperidinyloxy)-2-norvaline-cyclosporin
3-(N-tert-butyl-4-piperidinyloxy)-2-norvaline-cyclosporin

3-(2-Hydroxyethoxy)-4-methylvaline-cyclosporin
3-(3-Hydroxypropoxy)-4-methylvaline-cyclosporin
3-(2-Chloroethoxy)-4-methylvaline-cyclosporin
3-(2-Bromoethoxy)-4-methylvaline-cyclosporin
3-(2-Hydroxyethoxy)-4-methylvaline-cyclosporin
3-(3-Chloropropoxy)-4-methylvaline-cyclosporin
3-(3-Bromopropoxy)-4-methylvaline-cyclosporin
3-(3-Hydroxypropoxy)-4-methylvaline-cyclosporin
3-(3-Chlorobutoxy)-4-methylvaline-cyclosporin
3-(3-Bromobutoxy)-4-methylvaline-cyclosporin
3-(2-Aminoethoxy)-4-methylvaline-cyclosporin
3-(2-Methylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Dimethylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Ethylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Ethyl-N-methylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Diethylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Propylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Cyclopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Propyl-methylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Propyl-ethylaminoethoxy)-4-methylvaline-cyclosporin

3-(2-Methyl-isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Cyclopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Ethyl-isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Diisopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Propyl-isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Butylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-sec-Butylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Isobutylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-tert-Butylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-tert-Butyl-methylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-4-methylvaline-cyclosporin
3-(2-Azetidinoethoxy)-4-methylvaline-cyclosporin
3-(2-pyrrolidinoethoxy)-4-methylvaline-cyclosporin
3-(2-piperidinoethoxy)-4-methylvaline-cyclosporin
3-(2-morpholinoethoxy)-4-methylvaline-cyclosporin
3-(2-piperazinoethoxy)-4-methylvaline-cyclosporin
3-(2-N-methylpiperazinoethoxy)-4-methylvaline-cyclosporin
3-(2-N-methylpiperazinoethoxy)-4-methylvaline-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-4-methylvaline-cyclosporin
3-(3-azetidinyloxy)-4-methylvaline-cyclosporin
3-(N-methyl-3-azetidinyloxy)-4-methylvaline-cyclosporin
3-(N-isopropyl-3-azetidinyloxy)-4-methylvaline-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-4-methylvaline-cyclosporin
3-(3-pyrrolidinyloxy)-4-methylvaline-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-4-methylvaline-cyclosporin

3-(N-tert-butyl-3-pyrrolidinyloxy)-4-methylvaline-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-4-methylvaline-cyclosporin
3-(4-piperidinyloxy)-4-methylvaline-cyclosporin
3-(N-methyl-4-piperidinyloxy)-4-methylvaline-cyclosporin
3-(N-isopropyl-4-piperidinyloxy)-4-methylvaline-cyclosporin
3-(N-tert-butyl-4-piperidinyloxy)-4-methylvaline-cyclosporin

3-(2-Hydroxyethoxy)-4-methylisoleucine-cyclosporin
3-(3-Hydroxypropoxy)-4-methylisoleucine-cyclosporin
3-(2-Chloroethoxy)-4-methylisoleucine-cyclosporin
3-(2-Bromoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Hydroxyethoxy)-4-methylisoleucine-cyclosporin
3-(3-Chloropropoxy)-4-methylisoleucine-cyclosporin
3-(3-Bromopropoxy)-4-methylisoleucine-cyclosporin
3-(3-Hydroxypropoxy)-4-methylisoleucine-cyclosporin
3-(3-Chlorobutoxy)-4-methylisoleucine-cyclosporin
3-(3-Bromobutoxy)-4-methylisoleucine-cyclosporin
3-(2-Aminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Methylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Dimethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Ethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Ethyl-N-methylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Diethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Propylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Cyclopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Propyl-methylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Propyl-ethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Methyl-isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Cyclopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Ethyl-isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Diisopropylaminoethoxy)-4-methylisoleucine-cyclosporin

3-(2-n-Propyl-isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Butylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-sec-Butylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Isobutylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-tert-Butylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-tert-Butyl-methylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-4-methylisoleucine-cyclosporin
3-(2-Azetidinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-pyrrolidinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-piperidinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-morpholinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-piperazinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-N-methylpiperazinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-N-methylpiperazinoethoxy)-4-methylisoleucine-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-4-methylisoleucine-cyclosporin
3-(3-azetidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-methyl-3-azetidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-isopropyl-3-azetidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-4-methylisoleucine-cyclosporin
3-(3-pyrrolidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-tert-butyl-3-pyrrolidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-4-methylisoleucine-cyclosporin
3-(4-piperidinyloxy)-4-methylisoleucine-cyclosporin
3-(N-methyl-4-piperidinyloxy)-4-methylisoleucine-cyclosporin

3-(N-isopropyl-4-piperidinyloxy)-4-methylisoleucine-cyclosporin

3-(N-tert-butyl-4-piperidinyloxy)-4-methylisoleucine-cyclosporin

3-(2-Hydroxyethoxy)-8-(D)-serine-cyclosporin

3-(3-Hydroxypropoxy)-8-(D)-serine-cyclosporin

3-(2-Chloroethoxy)-8-(D)-serine-cyclosporin

3-(2-Bromoethoxy)-8-(D)-serine-cyclosporin

3-(2-Hydroxyethoxy)-8-(D)-serine-cyclosporin

3-(3-Chloropropoxy)-8-(D)-serine-cyclosporin

3-(3-Bromopropoxy)-8-(D)-serine-cyclosporin

3-(3-Hydroxypropoxy)-8-(D)-serine-cyclosporin

3-(3-Chlorobutoxy)-8-(D)-serine-cyclosporin

3-(3-Bromobutoxy)-8-(D)-serine-cyclosporin

3-(2-Aminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Methylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Dimethylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Ethylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Ethyl-N-methylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Diethylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-n-Propylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Isopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Cyclopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-n-Propyl-methylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-n-Propyl-ethylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Methyl-isopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Cyclopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Ethyl-isopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Diisopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-n-Propyl-isopropylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-n-Butylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-sec-Butylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-Isobutylaminoethoxy)-8-(D)-serine-cyclosporin

3-(2-tert-Butylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-n-Butyl-methylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-n-Butyl-ethylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-n-Butyl-isopropylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-sec-Butyl-methylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-sec-Butyl-ethylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-sec-Butyl-isopropylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-tert-Butyl-methylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-tert-Butyl-ethylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-tert-Butyl-isopropylaminoethoxy)-8-(D)-serine-cyclosporin
3-(2-Azetidinoethoxy)-8-(D)-serine-cyclosporin
3-(2-pyrrolidinoethoxy)-8-(D)-serine-cyclosporin
3-(2-piperidinoethoxy)-8-(D)-serine-cyclosporin
3-(2-morpholinoethoxy)-8-(D)-serine-cyclosporin
3-(2-piperazinoethoxy)-8-(D)-serine-cyclosporin
3-(2-N-methylpiperazinoethoxy)-8-(D)-serine-cyclosporin
3-(2-N-methylpiperazinoethoxy)-8-(D)-serine-cyclosporin
3-(2-N-isopropylpiperazinoethoxy)-8-(D)-serine-cyclosporin
3-(3-azetidinyloxy)-8-(D)-serine-cyclosporin
3-(N-methyl-3-azetidinyloxy)-8-(D)-serine-cyclosporin
3-(N-isopropyl-3-azetidinyloxy)-8-(D)-serine-cyclosporin
3-(N-tert-butyl-3-azetidinyloxy)-8-(D)-serine-cyclosporin
3-(3-pyrrolidinyloxy)-8-(D)-serine-cyclosporin
3-(N-methyl-3-pyrrolidinyloxy)-8-(D)-serine-cyclosporin
3-(N-tert-butyl-3-pyrrolidinyloxy)-8-(D)-serine-cyclosporin
3-(N-isopropyl-3-pyrrolidinyloxy)-8-(D)-serine-cyclosporin
3-(4-piperidinyloxy)-8-(D)-serine-cyclosporin
3-(N-methyl-4-piperidinyloxy)-8-(D)-serine-cyclosporin
3-(N-isopropyl-4-piperidinyloxy)-8-(D)-serine-cyclosporin
3-(N-tert-butyl-4-piperidinyloxy)-8-(D)-serine-cyclosporin

36. Test for cyclophilin binding:

To measure the affinity and inhibition of test compounds for cyclophilin, the test described by H. Fliri in "Antibiotics and Antiviral compounds", 1993, Verlag Chemie, K.Krohn, H.A. Kirst, and H. Maag, eds., p 229-240 was used. The activity of the test compounds shown in the table is expressed as relative IC₅₀, i.e. IC₅₀ (cyclosporin A)/IC₅₀ (test compound):

Product of example	Relative IC ₅₀ (ng/ml)
1	0.2
2	0.2
3	0.2
4	0.2
5	0.2
6	0.2
7	0.2
8	0.4
9	1
12	1.5
13	1.5
14	2
15	2

37 Test for inhibition of cyclophilin :

The test described by G. Fischer et al., Biomed. Biochim. Acta, Vol. 43, 1101 - 1111 (1984) was used. Specifically, 5 μ L of a 1 μ M solution of recombinant human cyclophilin 18 was dissolved, at 10,0 $\pm$ 0.3°C in 1250 μ L HEPES buffer ; α -chymotrypsin was added to give a final concentration of 30 μ M. Test compounds were dissolved in DMSO as 10 μ M solutions and added to the enzyme solution to give final concentrations of 0.1, 0.5, 1.0, 5.0, 10.0, 50.0, 100.0, 500.0, and 1000 nM concentrations. After 5 minutes preincubation, Succ-Ala-Phe-Pro-Phe-4-nitroanilide was added to give a final concentration of 50 μ M. The hydrolysis of the substrate was measured

spectrophotometrically at 390 - 450 nM. The following IC⁵⁰ values were obtained :

Product of Example	Name	IC50 [nM]
	Cyclosporin A	12.43 ± 0.40
14	3-Methoxy-cyclosporin	11.57 ± 1.59
15	3-Ethoxy-cyclosporin	11.13 ± 1.03
16	3-Isopropoxy-cyclosporin	20.39 ± 1.26
17	3-tert-Butoxy-cyclosporin	18.94 ± 1.03
18	3-Allyloxy-cyclosporin	10.15 ± 2.21

38. Inhibition of HIV replication

The inhibitory effect of test compounds on the HIV-induced cytopathic effect on the MT4 T cell line as described by R. Pauwels in J. Virol. Methods, 1988, Vol. 20, pp 309-321 was used. The activity of test compounds is shown in the table:

Product of example	IC50 (ng/ml)
1	450
2	1000
3	1000
4	1000
5	1000
6	1000
7	1000
8	500
9	30
12	150
13	150
14	100

39. Activity against *Toxoplasma gondii*:

Using microtiter plates, *T. gondii* tachyzoites, strain RH were co-incubated with fibroblasts (MRC5, Bio-Merieux) for 72 hrs in the presence and absence of test compounds. *T. gondii* was quantified using an ELISA. The activities of the test compounds are as follows:

Product of example	Activity against <i>T. gondii</i> (IC ₅₀ , µg/ml)
9	0.1
13	1
14	0.6
15	0.6



40 Antiinflammatory activity

The antiinflammatory activity of test compounds was determined by the adjuvant arthritis test described by Pearson, "Arthr. Rheum", 1959, Vol. 2., 440. Test compounds were administered orally as solutions in olive oil and tween 80. Their activity is expressed as ED₅₀ in mg/kg, i.e. the dose effective in reducing the joint swelling by 50%, and is as shown below.

Product of example	ED ₅₀ (mg/kg)
1	25
2	25
3	25
4	25
5	25
6	25
7	25
8	25
9	15


12	15
13	15
14	10
15	15

It is to be understood that a reference herein to a prior art document does not constitute an admission that the document forms part of the common general knowledge in the art in Australia.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:


1. A compound of formula I



Formula I

or a pharmaceutically acceptable salt thereof, wherein the letters A to L represent residues of the following amino acids:

- A (L)-alpha-N-methylamino-beta-hydroxy acid of the formula II,




Formula II

wherein R1 is (E)-2-butenyl-1,

- B alpha-amino-butyric acid, alpha-amino-valerianic acid (norvaline), threonine, or valine,

- C substituted sarcosine of the formula III



Formula III

in which X is O-R3, in which R3 is

- (a) hydrogen, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 1-pentyl, 2-pentyl, 3-pentyl, isopentyl, tert-pentyl, neopentyl, hexyl or an isomer thereof,
cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
bicyclo[2.1.1]hexyl, norbornyl, bicyclo[2.2.2]octyl,




vinyl, allyl, butenyl, pentenyl, pentadienyl, hexenyl, hexadienyl, formyl, acetyl, propionyl, butyryl, pivaloyl, benzoyl or alkoxycarbonyl, which can be substituted by hydroxy, methoxy, ethoxy, propoxy, isopropoxy, halogen, carboxy or carbamido, or

- 5 (b) amino, methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, sec-butylamino, tert-butylamino, 1-pentylamino, 2-pentylamino, 3-pentylamino, isopentylamino, tert-pentylamino, neopentylamino, hexylamino or an isomer thereof, N,N-dimethylamino, N-methyl-N-ethylamino, N,N-diethylamino, N-propyl-N-methylamino, N-methyl-N-isopropylamino, dipropylamino, 10 diisopropylamino, N-butyl-N-methylamino or an isomer thereof, N-butyl-N-ethylamino or an isomer thereof, N-butyl-N-propylamino or an isomer thereof, N,N-dibutylamino or an isomer thereof, azetidine, pyrrolidine, piperidine, morpholine, piperazine, N'-alkylpiperazine, 15 azabicyclo[2.1.1]hexane, azanorbornane, azabicyclo[2.2.2]octane, N-formylamino, N-acetylamino, N-tert-butoxycarbonyl-amino, N-benzyloxycarbonyl-amino, N-benzoyl-amino or N-phthaloyl, or

(c) residue of the formula VI and Y⁻ is an anion

20




Formula VI

- D is N-methyl-leucine, N-methyl-valine, or N-methyl-isoleucine,
E is valine,
25 F is N-methyl-leucine,
G is alanine,
H is glycine, (D)-alanine, (D)-serine, O-hydroxyethyl-(D)-serine,
I,K are N-methyl-leucine, and
L is N-methyl-valine.

30

2. A process for the synthesis of compounds of the formula I, characterized in, that in a compound of formula I, wherein C is a substituted sarcosine of the formula III in which X is -SR₃, the -SR₃ is exchanged by -OR₃.



3. A process for the synthesis of compounds of the formula I, characterized in, that the substituent OR3 of formula III is substituted by another substituent OR3.
4. Use of compounds of formula I and its pharmaceutically acceptable salts as pharmaceuticals and for the manufacture of pharmaceutical compositions.
5. Use of compounds of formula I and its pharmaceutically acceptable salts as inhibitors of cyclophilins.
6. Use of compounds of formula I and its pharmaceutically acceptable salts for the treatment of infectious diseases caused by bacterial, fungal, viral pathogens as well as by protozoan parasites and helminthes.
7. Use of compounds of formula I and its pharmaceutically acceptable salts for the treatment of chronic inflammatory and autoimmune diseases.
8. Use of compounds of formula I and its pharmaceutically acceptable salts for the treatment of pathologic conditions caused by trauma and reperfusion injury.
9. Use of compounds of formula I and its pharmaceutically acceptable salts to induce tissue regenerative processes.
10. A method of treating infectious diseases caused by bacterial, fungal or viral pathogens, or by protozoan parasites or helminthes, in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.
11. A method of treating chronic inflammation or autoimmune diseases in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.
12. A method of treating pathological conditions caused by trauma or reperfusion injury in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.



13. A method of inducing tissue regenerative processes in a patient comprising administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

14. A compound of formula I or a pharmaceutically acceptable salt thereof substantially as herein described with reference to the Examples.

Dated this 3rd day of January 2001

C-CHEM AG

By their Patent Attorneys

GRIFFITH HACK

