

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 July 2009 (16.07.2009)

PCT

(10) International Publication Number
WO 2009/089062 A2

(51) International Patent Classification:
C07K 16/28 (2006.01) *A61K 38/17* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BI, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2009/000162

(22) International Filing Date: 9 January 2009 (09.01.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/010,734 11 January 2008 (11.01.2008) US

(71) Applicant (for all designated States except US): **SYNOVEX CORPORATION** [US/US]; 200 Boston Avenue, Suite 1100, Medford, MA 02155 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MCARTHUR, James, G.** [US/US]; 94 Upland Road, Concord, MA 01742 (US).

(74) Agents: **BROOK, David, E.** et al.; Hamilton, Brook, Smith & Reynolds, P.C., 530 Virginia Road, P.O. Box 9133, Concord, MA 01742-9133 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description

WO 2009/089062 A2

(54) Title: CADHERIN-11 ANTAGONISTS AND METHODS FOR THE TREATMENT OF INFLAMMATORY JOINT DISORDERS

(57) Abstract: The present invention relates to Cadherin-11 antagonists and compositions comprising Cadherin-11 antagonists. The invention also relates to methods for treating inflammatory joint disorders, such as rheumatoid arthritis, in a mammalian subject by administering a therapeutically effective amount of a Cadherin-11 antagonist.

CADHERIN-11 ANTAGONISTS AND METHODS FOR THE TREATMENT OF INFLAMMATORY JOINT DISORDERS

RELATED APPLICATION(S)

This application claims the benefit of U.S. Provisional Application No. 5 61/010,734, filed on January 11, 2008. The entire teachings of the above application are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Patients with advanced chronic joint inflammation suffer from severe joint deterioration including bone and cartilage destruction, resulting in long-term pain, 10 deformity, loss of joint function, reduced mobility and shortened life expectancy. Joint inflammation is associated with an increased number of cells and inflammatory substances in the joint, which cause irritation, wearing down of cartilage and swelling of the joint lining. Several different autoimmune disorders are known to trigger inappropriate or misdirected inflammation in a joint, resulting in chronic 15 inflammation in the joints of individuals who suffer from these disorders. Common inflammatory joint disorders include rheumatoid arthritis, psoriatic arthritis, Reiter's syndrome and ankylosing spondylitis.

Rheumatoid arthritis (RA) is the most common form of inflammatory arthritis and is estimated to affect approximately 1 percent of the U.S. population, or 20 about 2.1 million Americans. RA is a chronic disease that is characterized by inflammation of the lining, or synovium, of the joints, and can lead to significant bone and cartilage damage over time. RA is more common in women than in men and as many as 3% of women may develop rheumatoid arthritis in their lifetime. Currently, the cause of RA is unknown.

25 RA can lead to long-term joint damage, resulting in chronic pain, loss of function and disability. In addition, recent research indicates that people with RA,

particularly those whose disease is not well controlled, may have a higher risk for heart disease and stroke. Thus, RA is a major national health burden and there is an urgent need to develop new agents for the prevention and treatment of rheumatoid arthritis, and other inflammatory joint disorders.

5

SUMMARY OF THE INVENTION

The present invention encompasses, in one embodiment, a Cadherin-11 antagonist that specifically binds an extracellular 1 (EC1) domain of a mammalian Cadherin-11 protein, and inhibits aggregation of cells that express the mammalian Cadherin-11. In a particular embodiment, the Cadherin-11 antagonist is an antibody or an antibody fragment. In another embodiment, the Cadherin-11 antagonist is a fusion protein that comprises the EC1 domain of a Cadherin-11 protein (e.g., SEQ ID NO:3).

In an additional embodiment, the invention relates to methods of treating an inflammatory joint disorder in a mammalian subject (e.g., a human). The method comprises administering to the mammalian subject a therapeutically effective amount of a Cadherin-11 antagonist of the invention, thereby resulting in a desired therapeutic effect in the mammal. In a particular embodiment, the methods of the invention can be used to treat rheumatoid arthritis.

In another embodiment, the invention encompasses a pharmaceutical composition comprising a Cadherin-11 antagonist of the invention and a pharmaceutically-acceptable carrier. In one embodiment, the pharmaceutical composition further comprises a second agent, such as a disease-modifying anti-rheumatic drug or an anti-inflammatory agent.

25

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

FIG. 1A is a Western blot showing detection of a Cadherin-11-EC1-5-Fc fusion protein using anti-Cad-11 antibodies 23C6, 13C2 and 27F3 (see solid arrows). These antibodies did not recognize the Cadherin-11-EC1-Fc and Cadherin-11-EC1/2-Fc fusion proteins that were also present on the membrane (see open arrows for positions of the Cadherin-11-EC1-Fc and Cadherin-11-EC1/2-Fc proteins on the blot).

FIG. 1B is a graph depicting the binding of public Cadherin-11 antibodies 13C2, 23C6 and 5F82 to human Cad-11-EC1-5-Fc fusion protein, but not the Cad-11-EC1-Fc fusion protein, as determined by ELISA. In contrast, the EC1 antibody 10 H1M1 binds both Cad-11-EC1-Fc fusion protein and the Cad-11-EC1-5-Fc fusion protein.

FIG. 2 is an amino acid sequence alignment of the first 34 amino acids of the EC1 domains of human Cad-11 (SEQ ID NO:3), MN-Cad (SEQ ID NO:4), and Cad-8 (SEQ ID NO:5) that are involved in cadherin binding. Donor sequences containing residues that extend into the pocket of a cadherin counter-receptor are indicated by underlining of the left half of sequence and residues of the pocket sequence are indicated by the underlining of the right half of sequence in SEQ ID NO:3.

FIG. 3 is a graph depicting the binding of a Cadherin-11-binding Fab to a 20 human Cad-11 EC1 domain peptide as well as the Cad-11-EC1-Fc fusion protein, but not the Cad-8 or MN-Cad EC1 domain peptides, as determined by ELISA. Clone 7 demonstrated significant binding to the Cad-11 EC1 domain peptide and fusion protein, but not the MN-Cad or Cad-8 EC1 domain peptides.

FIG. 4 is a graph depicting data from an *in vitro* Cad-11 cell aggregation 25 assay. Cad-11 antagonists added to the media, such as a Fab made from the anti-Cad-11 antibody 13C2, or varying concentrations of an anti-Cadherin-11 EC1 Fab directed to the first 35 amino acids of the EC1 domain of Cadherin-11 (designated EC1 Fab clone 7), block Cad-11 mediated 431-D-11 cell aggregation. The anti-Cadherin-11 EC1 Fab (clone 7) inhibited aggregation of A-431-D-11 epidermoid 30 carcinoma cells at all concentrations tested in a range of 0.3 μ g/ml to 10 μ g/ml. In contrast, the Fab made from the 13C2 anti-Cadherin-11 antibody only inhibited cell aggregation at a concentration of 10 μ g/ml.

FIG. 5 is a graph depicting data from a second *in vitro* Cad-11 cell aggregation assay. Percent aggregation of 431-D-11 cells is shown at 40 min. after addition of either SME media (designated control), varying concentrations of a fusion protein comprising the EC1 domain of Cad-11 fused to the human IgG2 hinge, CH2 and CH3 domains (designated Cad-11-EC1-Fc), varying concentrations of an anti-Cadherin-11 EC1 Fab directed to the first 35 amino acids of the EC1 domain of Cadherin-11 (designated Cad-11 EC1 Fab) or varying concentrations of a control anti-green fluorescent protein (anti-GFP) Fab (designated GFP fAb). The anti-Cadherin-11 EC1 Fab (clone 7) inhibited aggregation of Cad-11 expressing 431-D-11 cells at concentrations of 3 μ g/ml, 1 μ g/ml and 0.1 μ g/ml. The EC1-Fc fusion protein inhibited aggregation of 431-D-11 cells at concentrations of 3 μ g/ml. In contrast, the anti-GFP Fab failed to inhibit cell aggregation significantly at any of the test concentrations.

FIG. 6 is a graph depicting inhibition of Cad-11 mediated cell aggregation by various anti-Cadherin-11 EC1 Fabs that have binding specificity for Cad-11 alone (EC1 fAb clone 7 and clone 4), Cad-11 and Cad-8 (EC1 fAb clone 6), or Cad-11 and MN-Cad (EC1 fAb clone 5), using an *in vitro* cell aggregation assay. All Fabs tested inhibited 431-D-11 cell aggregation relative to the control (D-11 SME; left bar).

FIG. 7 is a graph depicting inhibition of Cad-11 mediated cell aggregation by anti-Cad-11 Fabs that have binding specificity for Cad-11 alone (EC1 fAb clone 7), or Cad-11 and MN-Cad (EC1 fAb clone 8), using an *in vitro* cell aggregation assay. The specificity of the Fabs tested is shown in parentheses next to each Fab designation. Both cadherin-specific Fabs inhibited cell aggregation (middle and right bars) relative to a control Fab that was specific for GFP (left bar).

FIG. 8 shows the nucleotide (DNA) sequence (SEQ ID NO:6) of the human Cad-11-EC1-hIgG2-Fc1 fusion protein (Cad-11-EC1-Fc). The sequence of the human Cadherin-11 extracellular domain is shown in italics, the BglII site is underlined, and the sequence encoding the human IgG2-Fc1 region is shown in bold lettering.

FIG. 9 shows the amino acid sequence (SEQ ID NO:7) of the human Cad-11-EC1-hIgG2-Fc1 fusion protein (Cad-11-EC1-Fc). The sequence of the human

Cadherin-11 extracellular domain is shown in italics, the sequence encoded by the BglII site is underlined, and the sequence of the human IgG₂-Fc1 region is shown in bold lettering.

FIG. 10 is an image of an SDS polyacrylamide gel that has been stained with Coomassie Blue, which shows the predominant intense bands corresponding to the monomeric form of the purified Cad-11-EC1-hIgG₂-Fc1 (middle lane) and Cad-11-EC1/2-hIgG₂-Fc1 (right lane) fusion proteins, respectively, following purification from cell culture medium using a protein A column. Molecular weight standards are shown in the left lane.

FIG. 11 is a Western blot showing the detection of human Cad-11-EC1-hIgG₂-Fc1 (middle lane) and Cad-11-EC1/2-hIgG₂-Fc1 (right lane) fusion proteins using an anti-human IgG antibody conjugated to horse radish peroxidase (HRP). The predominant observed band in each lane corresponds to the locations of the monomeric forms of the fusion proteins. The locations of the dimeric forms of the fusion proteins are also visible (see less intense higher molecular weight bands), due to incomplete reducing conditions. Molecular weight standards are shown in the left lane.

FIG. 12 is a graph depicting that a Cad-11-EC1-Fc fusion protein and a mouse anti-Cad-11 antibody, 13C2, inhibit the invasion of Cad-11 expressing human fibroblast-like synoviocytes into a matrigel plug at the indicated concentrations compared to untreated cells, labeled *Invasion*. Data is pooled from two independent experiments.

FIGS. 13A and B are graphs depicting data from two *in vitro* Cad-11 cell aggregation assays. Percent aggregation of Cad-11 expressing 431-D-11 cells is shown at 40 min. after addition of either SME media (designated control) or a Cad-11 fusion protein. FIG. 13A shows the inhibition of aggregation in the presence of varying concentrations of a fusion protein comprising the 5 extracellular domains of Cad-11 fused to the human IgG2 hinge, CH2 and CH3 domains (designated Cad-11-EC1-5-Fc). FIG. 13B shows the inhibition of aggregation of varying concentrations of a fusion protein comprising either the N-terminal extracellular domain (EC1 domain) of Cad-11 fused to the human IgG2 hinge, CH2 and CH3 domains (designated Cad-11-EC1-Fc) or Cad-11-EC1-5-Fc.

FIGS. 14A-C show the human Cadherin-11 cDNA sequence (SEQ ID NO:1; see Genbank Accession No. NM001797).

FIG. 15 shows the human Cadherin-11 protein sequence (SEQ ID NO:2; see Genbank Accession No. NP001788).

5 FIG. 16 is a graph depicting the level of binding of antibodies in media from peptide 4 hybridomas (HL), or control hybridoma media (Media), to proteins containing the EC1-2 domains of Cad-11, Cad-8 or MN-Cad, as determined by ELISA.

10 FIGS. 17A-C are representative graphs depicting the intensity of cell staining (MFI; mean fluorescence intensity) as a measure of binding of H14 antibody to Cad-11-expressing 431-D-11 cells.

FIGS. 17D-F are representative graphs depicting the absence of 431-D cell staining (MFI; mean fluorescence intensity) relative to FIGS. 17A-C, indicating a lack of binding of H14 antibody to the Cad-11 negative cells.

15 FIGS. 17G-I are representative graphs depicting the intensity of cell staining (MFI; mean fluorescence intensity) as a measure of binding of H1M1 antibody to Cad-11-expressing 431-D-11 cells.

FIG. 18A is a graph depicting the binding of H14 antibody to Cad-11-expressing cells, and the absence of H14 binding to Cad-11 negative control cells, at 20 varying concentrations of antibody, as measured by the intensity of cell staining (MFI; mean fluorescence intensity).

FIG. 18B is a graph depicting the binding of H1M1 antibody to Cad-11-expressing cells, and the absence of binding of H1M1 to Cad-11 negative control cells, at varying concentrations of antibody, as measured by the intensity of cell staining (MFI; mean fluorescence intensity).

25 FIG. 19A is a graph depicting the degree of binding of the H14 anti-Cad-11 antibody to Cad-11 and Cad-8 EC1 domain peptides at various antibody concentrations, as determined by ELISA.

FIG. 19B is a graph depicting the absence of binding of the H14 anti-Cad-11 antibody to Cad7, MN Cad, Cad9, Cad18, Cad20 or Cad24 EC1 domain peptides at 30 various antibody concentrations, as determined by ELISA.

FIG. 20 is a graph depicting the binding of the H1M1 anti-Cad-11 antibody to Cad-11 and Cad-8 EC1 domain peptides at varying antibody concentrations, as determined by ELISA.

5 FIG. 21A is a graph depicting the degree of binding of the H1M1 anti-Cad-11 antibody to various Cad-11 EC1 domain peptide immunogens (PEP1, PEP2, PEP3 and PEP4), as well as the Cad-11 EC1 domain fusion protein (EFL) and human IgG control (Fc block), as determined by ELISA.

10 FIG. 21B is a graph depicting the degree of binding of the H14 anti-Cad-11 antibody to various Cad-11 EC1 domain peptide immunogens (PEP1, PEP2, PEP3 and PEP4), as well as the Cad-11 EC1 domain fusion protein (EFL) and human IgG control (Fc block), as determined by ELISA.

15 FIG. 22 is a schematic diagram depicting the sequence of the first 37 amino acids of the EC1 domain of human Cadherin-11 and the portions of this sequence encompassed by each of Peptides 1-4. Amino acid residues shared by Peptides 2 and 4 that are upstream of Peptide 3 are highlighted in the boxed region. Amino acids directly involved in Cad-11 to Cad-11 binding are underlined.

FIG. 23A is a photograph showing a large mass of aggregated Cad-11-expressing cells that were treated with a control isotype antibody.

20 FIG. 23B is a photograph showing small clumps of H1M1-treated Cad-11-expressing cells that did not progress to form the large masses observed in FIG. 23A.

FIG. 23C is a photograph showing that untreated parental Cad-11 negative cells remain as groups of single or double cells.

25 FIG. 24A is a photograph depicting a culture of Cad-11 expressing cells with large masses of aggregated cells.

FIG. 24B is a photograph depicting a culture of Cad-11 expressing cells with predominantly single cells with small and infrequent cell clusters relative to those shown in FIG. 24A following treatment with the H14 Cad-11 EC1 domain antibody.

30 FIG. 25 is a graph depicting inhibition of arthritis-associated joint swelling in mice treated with increasing dosages of H1M1 anti-Cad-11 antibody relative to untreated control mice.

FIG. 26 is a graph depicting inhibition of arthritis-associated joint swelling in mice treated with 0.3 mg of either H14 or H1M1 anti-Cad-11 antibodies every second day relative to untreated control mice.

5 FIG. 27 is a graph showing that treatment with 0.3 mg of either H1M1 or H14 antibody delayed the development of arthritis in a mouse model compared to an untreated control.

10 FIG. 28 is a graph depicting the degree of binding of antibody-containing media from peptide 3 hybridomas (HL), or control hybridoma media (Media), to the EC1-2 domains of Cad-11, Cad-8, and MN-Cadherin, or a Cad-11 EC1-Fc fusion protein.

FIG. 29 is a graph depicting the degree of binding of anti-Cad-11 antibodies from the peptide 3 hybridomas to cells expressing human Cad-11 protein (see arrow) and non-Cad-11-expressing control cells that expressed Neos.

15 DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, the terms “Cadherin-11,” “Cad-11,” and “OB-Cadherin” refer to a naturally occurring or endogenous Cadherin-11 (*e.g.*, mammalian, for example human) protein, and to proteins having an amino acid sequence that is the same as that of naturally occurring or endogenous Cadherin-11 protein (*e.g.*, recombinant proteins, synthetic proteins). Accordingly, the terms “Cadherin-11,” “Cad-11,” and “OB-Cadherin,” which are used interchangeably herein, include polymorphic or allelic variants and other isoforms of a Cadherin-11 protein (*e.g.*, mammalian, human) produced by, *e.g.*, alternative splicing or other cellular processes, that occur naturally in mammals (*e.g.*, humans, non-human primates). Preferably, the Cadherin-11 protein is a human protein that has the amino acid sequence of SEQ ID NO:2 (See, Genbank Accession No. NP001788 and FIG. 15).

As defined herein, a “Cadherin-11 antagonist” is an agent (*e.g.*, antibody, fusion protein, peptide, peptidomimetic, small molecule, nucleic acid) that 30 specifically binds an EC1 domain of a Cadherin-11 protein and inhibits (*e.g.*, reduces, prevents) one or more Cadherin-11-mediated activities in a cell. Cadherin-

11-mediated activities include, but are not limited to, binding of a Cadherin-11 protein to one or more other Cadherin-11 proteins in a homotypic fashion, aggregation of cells that express Cadherin-11, induction of enzyme (e.g., collagenase, serine proteases, MMP1, MMP3, MMP13) expression or activity, and 5 induction of cytokines or growth factors (e.g., IL-6, IL-8 or RANKL or TRANCE). In one embodiment, the Cadherin-11 antagonist can inhibit the binding of a Cadherin-11 protein to one or more other Cadherin-11 proteins by, for example, blocking the interaction between the donor sequences in the EC1 domain of a Cad-11 protein (e.g., a Cad-11 protein expressed on the surface of a cell) with the pocket 10 sequence in the EC1 domain of one or more other Cad-11 proteins (e.g., one or more Cad-11 proteins expressed on the surface of another cell).

As used herein, a Cadherin-11 antagonist that “specifically binds” an EC1 domain of a Cadherin-11 protein refers to a Cadherin-11 antagonist that binds (e.g., under physiological conditions) an EC1 domain of a Cadherin-11 protein with an 15 affinity (e.g., a binding affinity) that is at least about 5 fold, preferably at least about 10 fold, greater than the affinity with which the Cadherin-11 antagonist binds an EC1 domain of another cadherin protein (e.g., MN-Cadherin, Cadherin-8). In a particular embodiment, the Cadherin-11 antagonist that specifically binds an EC1 domain of a Cadherin-11 protein binds an epitope present in SEQ ID NO:3, the N-terminal portion of the EC1 domain of human Cadherin-11, with an affinity that is at 20 least about 5 fold, preferably at least about 10 fold, greater than the affinity with which the Cadherin-11 antagonist binds an epitope present in SEQ ID NO:4, the N-terminal portion of the EC1 domain of human MN-Cadherin, and the affinity with which the Cadherin-11 antagonist binds an epitope present in SEQ ID NO:5, the N-terminal portion of the EC1 domain of human Cadherin-8.

As used herein, the term “antibody” is intended to encompass both whole antibodies and antibody fragments (e.g., antigen-binding fragments of antibodies, for example, Fv, Fc, Fd, Fab, Fab', F(ab'), and dAb fragments). “Antibody” refers to both polyclonal and monoclonal antibodies and includes naturally-occurring and 30 engineered antibodies. Thus, the term “antibody” includes, for example, human, chimeric, humanized, primatized, veneered, single chain, and domain antibodies

(dAbs). (See *e.g.*, Harlow *et al.*, *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988).

5 The term “epitope” refers to a unit of structure conventionally bound by an immunoglobulin V_H/V_L pair. An epitope defines the minimum binding site for an antibody, and thus represent the target of specificity of an antibody.

The term “fusion protein” refers to a naturally occurring, synthetic, semi-synthetic or recombinant single protein molecule that comprises all or a portion of two or more heterologous polypeptides

10 The term “polypeptide” refers to a polymer of amino acids, and not to a specific length; thus, peptides, oligopeptides and proteins are included within the definition of a polypeptide.

15 As used herein, the term “peptide” refers to a compound consisting of from about 2 to about 100 amino acid residues wherein the amino group of one amino acid is linked to the carboxyl group of another amino acid by a peptide bond. Such peptides are typically less than about 100 amino acid residues in length and preferably are about 10, about 20, about 30, about 40 or about 50 residues.

20 As used herein, the term “peptidomimetic” refers to molecules which are not peptides or proteins, but which mimic aspects of their structures. Peptidomimetic antagonists can be prepared by conventional chemical methods (see *e.g.*, Damewood J.R. “Peptide Mimetic Design with the Aid of Computational Chemistry” in *Reviews in Computational Biology*, 2007, Vol. 9, pp.1-80, John Wiley and Sons, Inc., New York, 1996; Kazmierski W.K., “*Methods of Molecular Medicine: Peptidomimetic Protocols*,” Humana Press, New Jersey, 1999).

25 As defined herein, “therapy” is the administration of a particular therapeutic or prophylactic agent to a subject (*e.g.*, a mammal, a human), which results in a desired therapeutic or prophylactic benefit to the subject.

30 As defined herein a “treatment regimen” is a regimen in which one or more therapeutic or prophylactic agents are administered to a mammalian subject at a particular dose (*e.g.*, level, amount, quantity) and on a particular schedule or at particular intervals (*e.g.*, minutes, days, weeks, months).

As defined herein, a “therapeutically effective amount” is an amount sufficient to achieve the desired therapeutic or prophylactic effect under the

conditions of administration, such as an amount sufficient to inhibit (*i.e.*, reduce, prevent) inflammation in a joint (*e.g.*, by inhibiting the aggregation of cells, for example synoviocytes, that express Cadherin-11). The effectiveness of a therapy (*e.g.*, the reduction of inflammation in a joint and/or prevention of inflammation in a 5 joint) can be

determined by suitable methods (*e.g.*, imaging methods, such as MRI, NMR, CT).

Cadherins

Cadherins belong to a large family of Ca^{2+} -dependent adhesion molecules 10 that mediate cell adhesion by binding to other cadherins in a homotypic manner (MJ Wheelock and KR Johnson, *Ann. Rev. Cell Dev. Biol.* 19: 207-235 (2003). Classical cadherins are single-pass transmembrane proteins that contain five extracellular cadherin (EC) domains, each approximately 110 amino acids in length, a transmembrane region and a conserved cytoplasmic domain. Cadherins are divided 15 into either type I or type II cadherins based on the degree of homology between the EC domains. Type II cadherins include human cadherins-5, -6, -8, -11, and -12, and MN-cadherin. The relative importance of the role of each of the extracellular domains in mediating inter-cellular binding is unclear.

20 Cadherin-11 activity in synoviocytes

Cadherin-11 mediates synoviocyte to synoviocyte binding in the synovial lining of articulated joints (Valencia *et al.*, *J. Exp. Med.* 200(12):1673-1679 (2004); Kiener and Brenner, *Arthritis Res Ther.* 7(2):49-54 (2005)). A fusion protein that comprised all five extracellular cadherin domains of human Cadherin-11, fused to 25 the hinge-CH₂-CH₃ domain of human IgG₂, inhibited synoviocyte lining formation *in vitro* (Kiener *et al.*, *Am. J. Pathol.* 168 (2006)). In addition, antagonistic anti-Cadherin-11 antibodies and a fusion protein that comprised EC1-5 of murine Cadherin-11, fused to the hinge-CH₂-CH₃ domains of murine IgG2a, inhibited 30 inflammation and joint swelling in murine models of rheumatoid arthritis (Lee *et al.*, *Science* 315:1006-1010 (2007)).

Cadherin-11 antagonists

A Cadherin-11 antagonist of the invention can be any agent that specifically binds an EC1 domain of a Cadherin-11 protein and inhibits (e.g., reduces, prevents) one or more Cadherin-11-mediated activities in a cell. Cadherin-11-mediated activities include, but are not limited to, aggregation of cells that express Cadherin-11 on the cell surface, and expression or secretion of factors such as, for example, collagenase, serine proteases, MMP1, MMP3, IL-6, IL-8 or RANKL/TRANCE. The agent can be an antibody, a fusion protein, a peptide, a peptidomimetic, a small molecule, or a nucleic acid, among others.

10

Cadherin-11 antibodies

As described herein, antibodies that bind an epitope within an N-terminal portion of the EC1 domain of human Cadherin-11 that comprises the donor sequences and cadherin-binding pocket of Cad-11 (e.g., SEQ ID NO:3), block Cadherin-11 activity *in vitro* more effectively than antibodies that bind to epitopes in other regions of this protein (See Examples 1 and 2).

Accordingly, in one embodiment, the invention provides an antibody or antigen-binding fragment thereof that binds (e.g., specifically binds) an epitope that is present in the N-terminal portion of the EC1 domain of a Cadherin-11 protein that comprises the donor sequences and cadherin-binding pocket of Cad-11. The term "antibody" is intended to encompass all types of polyclonal and monoclonal antibodies (e.g., human, chimeric, humanized, primatized, veneered, single chain, domain antibodies (dAbs)) and antigen-binding fragments of antibodies (e.g., Fv, Fc, Fd, Fab, Fab', F(ab'), dAb). (See e.g., Harlow *et al.*, *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988). In a particular embodiment, the Cad-11 EC1 domain-specific antibody is a human antibody or humanized antibody. Cad-11 EC1 domain-specific antibodies can also be directly or indirectly linked to a cytotoxic agent.

Other antibodies or antibody fragments that specifically bind to an N-terminal portion of the EC1 domain of a Cad-11 protein and inhibit the activity of the Cad-11 protein can also be produced, constructed, engineered and/or isolated by conventional methods or other suitable techniques. For example, antibodies which

are specific for the EC1 domain of a Cadherin-11 protein can be raised against an appropriate immunogen, such as a recombinant mammalian (e.g., human) Cadherin-11 EC1 domain peptide (e.g., SEQ ID NO:3) or a portion thereof (including synthetic molecules, e.g., synthetic peptides). A variety of methods have been

5 described (see e.g., Kohler *et al.*, *Nature*, 256: 495-497 (1975) and *Eur. J. Immunol.* 6: 511-519 (1976); Milstein *et al.*, *Nature* 266: 550-552 (1977); Koprowski *et al.*, U.S. Patent No. 4,172,124; Harlow, E. and D. Lane, 1988, *Antibodies: A*

10 *Laboratory Manual*, (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY); *Current Protocols In Molecular Biology*, Vol. 2 (Supplement 27, Summer '94), Ausubel, F.M. *et al.*, Eds., (John Wiley & Sons: New York, NY), Chapter 11, (1991)). Antibodies can also be raised by immunizing a suitable host (e.g., mouse) with cells that express the EC1 domain of Cadherin-11 (e.g., cancer cells/cell lines) or cells engineered to express the EC1 domain of Cadherin-11 (e.g., transfected cells). (See e.g., Chuntharapai *et al.*, *J. Immunol.*, 152:1783-1789 (1994);

15 Chuntharapai *et al.* U.S. Patent No. 5,440, 021). For the production of monoclonal antibodies, a hybridoma can be produced by fusing a suitable immortal cell line (e.g., a myeloma cell line such as SP2/0 or P3X63Ag8.653) with antibody producing cells. The antibody producing cells can be obtained from the peripheral blood, or preferably, the spleen or lymph nodes, of humans or other suitable animals

20 immunized with the antigen of interest. The fused cells (hybridomas) can be isolated using selective culture conditions, and cloned by limited dilution. Cells which produce antibodies with the desired specificity can be selected by a suitable assay (e.g., ELISA).

Antibody fragments can be produced by enzymatic cleavage or by recombinant techniques. For example, papain or pepsin cleavage can generate Fab or F(ab')₂ fragments, respectively. Other proteases with the requisite substrate specificity can also be used to generate Fab or F(ab')₂ fragments. Antibodies can also be produced in a variety of truncated forms using antibody genes in which one or more stop codons has been introduced upstream of the natural stop site. For example, a chimeric gene encoding a F(ab')₂ heavy chain portion can be designed to include DNA sequences encoding the CH₁ domain and hinge region of the heavy chain. Single chain antibodies, and human, chimeric, humanized or primatized

(CDR-grafted), or veneered antibodies, as well as chimeric, CDR-grafted or veneered single chain antibodies, comprising portions derived from different species, and the like are also encompassed by the present invention and the term "antibody". The various portions of these antibodies can be joined together 5 chemically by conventional techniques, or can be prepared as a contiguous protein using genetic engineering techniques. For example, nucleic acids encoding a chimeric or humanized chain can be expressed to produce a contiguous protein. See, e.g., Cabilly *et al.*, U.S. Patent No. 4,816,567; Cabilly *et al.*, European Patent No. 0,125,023 B1; Boss *et al.*, U.S. Patent No. 4,816,397; Boss *et al.*, European Patent 10 No. 0,120,694 B1; Neuberger, M.S. *et al.*, WO 86/01533; Neuberger, M.S. *et al.*, European Patent No. 0,194,276 B1; Winter, U.S. Patent No. 5,225,539; Winter, European Patent No. 0,239,400 B1; Queen *et al.*, European Patent No. 0 451 216 B1; and Padlan, E.A. *et al.*, EP 0 519 596 A1. See also, Newman, R. *et al.*, *BioTechnology*, 10: 1455-1460 (1992), regarding primatized antibody, and Ladner *et 15 al.*, U.S. Patent No. 4,946,778 and Bird, R.E. *et al.*, *Science*, 242: 423-426 (1988)) regarding single chain antibodies.

Humanized antibodies can be produced using synthetic or recombinant DNA technology using standard methods or other suitable techniques. Nucleic acid (e.g., cDNA) sequences coding for humanized variable regions can also be constructed 20 using PCR mutagenesis methods to alter DNA sequences encoding a human or humanized chain, such as a DNA template from a previously humanized variable region (see e.g., Kamman, M., *et al.*, *Nucl. Acids Res.*, 17: 5404 (1989)); Sato, K., *et al.*, *Cancer Research*, 53: 851-856 (1993); Daugherty, B.L. *et al.*, *Nucleic Acids Res.*, 19(9): 2471-2476 (1991); and Lewis, A.P. and J.S. Crowe, *Gene*, 101: 297-302 (1991)). Using these or other suitable methods, variants can also be readily 25 produced. In one embodiment, cloned variable regions (e.g., dAbs) can be mutated, and sequences encoding variants with the desired specificity can be selected (e.g., from a phage library; see e.g., Krebber *et al.*, U.S. 5,514,548; Hoogenboom *et al.*, WO 93/06213, published April 1, 1993).

30 Other suitable methods of producing or isolating antibodies of the requisite specificity can be used, including, for example, methods which select a recombinant antibody or antibody-binding fragment (e.g., dAbs) from a library (e.g., a phage

display library), or which rely upon immunization of transgenic animals (e.g., mice). Transgenic animals capable of producing a repertoire of human antibodies are well-known in the art (e.g., Xenomouse[®] (Abgenix, Fremont, CA)) and can be produced using suitable methods (see e.g., Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90: 5 2551-2555 (1993); Jakobovits *et al.*, *Nature*, 362: 255-258 (1993); Lonberg *et al.*, U.S. Patent No. 5,545,806; Surani *et al.*, U.S. Patent No. 5,545,807; Lonberg *et al.*, WO 97/13852).

The invention encompasses, in one embodiment, a Cad-11 antibody that binds to an epitope that is present in the first about 37 amino acids of the EC1 domain of human Cad-11 (SEQ ID NO: 13). In a particular embodiment, the invention relates to a Cad-11 antibody that binds to an epitope that is present in SEQ ID NO:10. In a further embodiment, the invention relates to a Cad-11 antibody that binds to an epitope that comprises SEQ ID NO:11. In another embodiment, the invention relates to a Cad-11 antibody that binds to an epitope that is present in SEQ 10 ID NO:12.

In one embodiment, the invention relates to a Cad-11 antibody produced by hybridoma H1M1 (ATCC accession number _____), having been deposited on January 8, 2009 at the American Type Culture Collection (ATCC), P.O. Box 1549, Manassas, Virginia 20108, United States of America. In another embodiment, the invention provides a Cad-11 antibody produced by hybridoma H14 (ATCC 20 accession number _____), having been deposited on January 8, 2009 at the American Type Culture Collection (ATCC), P.O. Box 1549, Manassas, Virginia 20108, United States of America.

The invention also encompasses antibodies that specifically compete with a Cad-11 antibody produced by hybridoma H1M1 and/or a Cad-11 antibody produced by hybridoma H14 for binding to a human Cad-11 protein or an EC1-domain containing portion thereof (e.g., SEQ ID NO:3, 10, 12, 13). In a particular embodiment, an antibody that specifically competes with a Cad-11 antibody produced by hybridoma H1M1 and/or hybridoma H14 blocks (e.g., inhibits, diminishes, prevents) the binding of a Cad-11 antibody produced by hybridoma H1M1 and/or hybridoma H14 to a human Cad-11 protein or EC1-domain containing portion thereof (e.g., SEQ ID NO:3, 10, 12, 13).

In addition, the invention encompasses antibodies having a binding affinity for a human Cad-11 protein or EC1-domain containing portion thereof (e.g., SEQ ID NO:3, 10, 12, 13) that is at least as great as the binding affinity of a Cad-11 antibody produced by hybridoma H1M1 and/or a Cad-11 antibody produced by hybridoma 5 H14 for a human Cad-11 protein or EC1-domain containing portion thereof.

Cadherin-11 fusion proteins

In addition, immunoglobulin fusion proteins that contain only the EC1 10 domain of human Cad-11 (e.g., the EC1 domain of human Cad-11 fused to a portion of human IgG) inhibited Cad-11 activity *in vitro* more effectively than a fusion protein that included a larger portion of the EC region of Cad-11, which contained all 5 EC domains.

Cadherin-11 antagonists also encompass chimeric, or fusion, proteins that 15 comprise at least about the N-terminal 35 amino acids of the EC1 domain of human Cad-11 (SEQ ID NO:2) operatively linked to all or a portion of a heterologous protein. “Operatively linked” indicates that the portion of the Cad-11 EC1 domain and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the protein. For example, the fusion 20 protein can be a GST-fusion protein in which the protein sequences are fused to the C-terminus of a GST sequence. Other types of fusion proteins include, but are not limited to, enzymatic fusion proteins, for example, β -galactosidase fusion proteins, yeast two-hybrid GAL fusion proteins, poly-His fusions, FLAG-tagged fusion proteins, GFP fusion proteins, and immunoglobulin (Ig) fusion proteins. Such 25 fusion protein can facilitate purification (e.g., of a recombinant fusion protein). In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence. Therefore, in another embodiment, the fusion protein contains a heterologous signal sequence at its N-terminus.

30 EP-A-O 464 533 discloses fusion proteins comprising various portions of immunoglobulin constant regions. The Fc is useful in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (see, for example,

EP-A 0232 262). In drug discovery, for example, human proteins have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists (Bennett *et al.*, *Journal of Molecular Recognition* 8:52-58 (1995); Johanson *et al.*, *J. Biol. Chem.*, 270(16):9459-9471 (1995)). Thus, this invention 5 also encompasses soluble fusion proteins containing a protein Cad-11 antagonist of the invention and various portions of the constant regions of heavy and/or light chains of immunoglobulins of various subclasses (e.g., IgG, IgM, IgA, IgE). Advantages of immunoglobulin fusion proteins of the present invention include one or more of the following: (1) increased avidity for multivalent ligands due to the 10 resulting bivalence of dimeric fusion proteins, (2) longer serum half-life, (3) the ability to activate effector cells via the Fc domain, (4) ease of purification (for example, by protein A chromatography), (5) affinity for Cad-11 and (6) the ability to block Cad-11 mediated activity.

Accordingly, in particular embodiments, the Cad-11 antagonist is a fusion 15 protein that comprises a portion of the extracellular region of a Cadherin-11 protein that includes an N-terminal portion of the EC1 domain (amino acids 54-90 of SEQ ID NO:2), operatively linked to all, or a portion of, a mammalian immunoglobulin protein. In a particular embodiment, the immunoglobulin fusion proteins of the invention do not comprise a portion of the extracellular region of Cadherin-11 that 20 includes all five EC domains that are contained within amino acids 1-609 of SEQ ID NO:2. In certain embodiments, the portion of the human Cadherin-11 extracellular region can include, for example, amino acids 1-160, amino acids 1-259 or amino acids 1-269 of SEQ ID NO:2. In a particular embodiment, the fusion protein lacks the leader and pro-region of human Cadherin-11 (amino acids 1-53 of SEQ ID 25 NO:2) and uses a heterologous leader sequence. The immunoglobulin portion can be from any vertebrate source, such as murine, but preferably, is a human immunoglobulin protein. In one embodiment, the mammalian immunoglobulin protein is a human IgG₂ protein or a portion thereof, such as the hinge-CH₂-CH₃ portion of human IgG₂.

30 A chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for different protein sequences (e.g., a Cad-11 EC1 domain peptide and a mammalian

immunoglobulin) are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of nucleic acid fragments can be carried out using anchor primers 5 that give rise to complementary overhangs between two consecutive nucleic acid fragments that can subsequently be annealed and re-amplified to generate a chimeric nucleic acid sequence (see Ausubel *et al.*, Current Protocols in Molecular Biology, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST moiety, an Fc moiety). A nucleic acid molecule 10 encoding protein Cad-11 antagonist can be cloned into such an expression vector that the fusion moiety (*e.g.*, immunoglobulin) is linked in-frame to the protein.

The immunoglobulin fusion proteins of the invention can be provided as monomers, dimers, tetramers or other multimers (*e.g.*, polymers). For example, 15 variable domains of the immunoglobulin portion of the fusion protein may be linked together to form multivalent ligands by, for example, provision of a hinge region at the C-terminus of each V domain and disulphide bonding between cysteines in the hinge regions; or provision of heavy chains each with a cysteine at the C-terminus of the domain, the cysteines being disulphide bonded together; or production of V-CH & V-CL to produce a Fab format; or use of peptide linkers (for example Gly₄Ser 20 linkers) to produce dimers, trimers and further multimers. For example, such ligands can be linked to an antibody Fc region comprising one or both of C_H2 and C_H3 domains, and optionally a hinge region. For example, vectors encoding ligands linked as a single nucleotide sequence to an Fc region may be used to prepare such ligands (*e.g.*, by expression).

25 The immunoglobulin fusion proteins of the invention can be conjugated to other moieties including, but not limited to, multimers of polyethelene glycol (PEG) or its derivatives (*e.g.*, poly methyl ethylene glycol), radionuclides, cytotoxic agents and drugs, and subsequently used for *in vivo* therapy. Examples of radionuclides include ²¹²Bi, ¹³¹I, ¹⁸⁶Re, and ⁹⁰Y, among others. The radionuclides exert their 30 cytotoxic effect by locally irradiating the cells, leading to various intracellular lesions, as is known in the art of radiotherapy. Cytotoxic drugs that can be conjugated to the fusion proteins include, but are not limited to, daunorubicin,

doxorubicin, methotrexate, and Mitomycin C. Cytotoxic drugs interfere with critical cellular processes including DNA, RNA, and protein synthesis. For a fuller exposition of these classes of drugs, which are known in the art, and their mechanisms of action, see Goodman, A.G., *et al.*, *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, 8th Ed., Macmillan Publishing Col, 1990. 5 Katzung, ed., *Basic and Clinical Pharmacology*, Fifth Edition, p 768-769, 808-809, 896, Appleton and Lange, Norwalk, Conn.

As used herein, the term "immunoglobulin fusion protein" includes fragments of the immunoglobulin fusion proteins of the invention. Such fragments 10 are intended to be within the scope of this invention. For example, once the molecules are isolated, they can be cleaved with protease to generate fragments that remain capable of binding the EC1 domain of human Cad-11.

Peptide antagonists

15 The Cadherin-11 antagonist of the invention can also be a peptide that binds to the EC1 domain of a Cadherin-11 protein. The peptide can comprise any suitable L-and/or D-amino acid, for example, common α -amino acids (e.g., alanine, glycine, valine), non- α -amino acids (e.g., β -alanine, 4-aminobutyric acid, 6-aminocaproic acid, sarcosine, statine), and unusual amino acids (e.g., citrulline, homocitrulline, 20 homoserine, norleucine, norvaline, ornithine). The amino, carboxyl and/or other functional groups on a peptide can be free (e.g., unmodified) or protected with a suitable protecting group. Suitable protecting groups for amino and carboxyl groups, and methods for adding or removing protecting groups are known in the art and are disclosed in, for example, Green and Wuts, *"Protecting Groups in Organic 25 Synthesis"*, John Wiley and Sons, 1991. The functional groups of a peptide can also be derivatized (e.g., alkylated) using art-known methods.

The peptide Cad-11 antagonist can comprise one or more modifications (e.g., amino acid linkers, acylation, acetylation, amidation, methylation, terminal 30 modifiers (e.g., cyclizing modifications)), if desired. The peptide can also contain chemical modifications (e.g., N-methyl- α -amino group substitution). In addition, the peptide antagonist can be an analog of a known and/or naturally-occurring peptide, for example, a peptide analog having conservative amino acid residue

substitution(s). These modifications can improve various properties of the peptide (e.g., solubility, binding), including its Cadherin-11 antagonist activity.

Cad-11 antagonists that are peptides can be linear, branched or cyclic, e.g., a peptide having a heteroatom ring structure that includes several amide bonds. In a 5 particular embodiment, the peptide is a cyclic peptide. Such peptides can be produced by one of skill in the art using standard techniques. For example, a peptide can be derived or removed from a native protein by enzymatic or chemical cleavage, or can be synthesized by suitable methods, for example, solid phase peptide synthesis (e.g., Merrifield-type synthesis) (see, e.g., Bodanszky *et al.* "Peptide 10 Synthesis," John Wiley & Sons, Second Edition, 1976). Peptides that are Cadherin-11 antagonists can also be produced, for example, using recombinant DNA methodologies or other suitable methods (see, e.g., Sambrook J. and Russell D.W., *Molecular Cloning: A Laboratory Manual*, 3rd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).

15 Peptides can be synthesized and assembled into libraries comprising a few to many discrete molecular species. Such libraries can be prepared using methods of combinatorial chemistry, and can be screened using any suitable method to determine if the library comprises peptides with a desired biological activity. Such peptide antagonists can then be isolated using suitable methods.

20

Peptidomimetic antagonists

Cadherin-11 antagonists can also be peptidomimetics. For example, 25 polysaccharides can be prepared that have the same functional groups as peptides. Peptidomimetics can be designed, for example, by establishing the three dimensional structure of a peptide agent in the environment in which it is bound or will bind to a target molecule. The peptidomimetic comprises at least two components, the binding moiety or moieties and the backbone or supporting structure.

The binding moieties are the chemical atoms or groups which will react or 30 form a complex (e.g., through hydrophobic or ionic interactions) with a target molecule, for example, with amino acids in the EC1 domain of Cad-11. For example, the binding moieties in a peptidomimetic can be the same as those in a

peptide or protein antagonist. The binding moieties can be an atom or chemical group which reacts with the receptor in the same or similar manner as the binding moiety in the peptide antagonist. For example, computational chemistry can be used to design peptide mimetics of the donor sequences of the EC1 domain of a

5 Cadherin-11 protein, for instance, which can bind to the pocket sequence in the EC1 domain of Cad-11 proteins. Examples of binding moieties suitable for use in designing a peptidomimetic for a basic amino acid in a peptide include nitrogen containing groups, such as amines, ammoniums, guanidines and amides or phosphoniums. Examples of binding moieties suitable for use in designing a

10 peptidomimetic for an acidic amino acid include, for example, carboxyl, lower alkyl carboxylic acid ester, sulfonic acid, a lower alkyl sulfonic acid ester or a phosphorous acid or ester thereof.

The supporting structure is the chemical entity that, when bound to the binding moiety or moieties, provides the three dimensional configuration of the

15 peptidomimetic. The supporting structure can be organic or inorganic. Examples of organic supporting structures include polysaccharides, polymers or oligomers of organic synthetic polymers (such as, polyvinyl alcohol or polylactide). It is preferred that the supporting structure possess substantially the same size and dimensions as the peptide backbone or supporting structure. This can be determined

20 by calculating or measuring the size of the atoms and bonds of the peptide and peptidomimetic. In one embodiment, the nitrogen of the peptide bond can be substituted with oxygen or sulfur, for example, forming a polyester backbone. In another embodiment, the carbonyl can be substituted with a sulfonyl group or sulfinyl group, thereby forming a polyamide (e.g., a polysulfonamide). Reverse amides of the peptide can be made (e.g., substituting one or more CONH-groups for a NHCO-group). In yet another embodiment, the peptide backbone can be substituted with a polysilane backbone.

These compounds can be manufactured by known methods. For example, a polyester peptidomimetic can be prepared by substituting a hydroxyl group for the

30 corresponding α -amino group on amino acids, thereby preparing a hydroxyacid and sequentially esterifying the hydroxyacids, optionally blocking the basic and acidic side chains to minimize side reactions. Determining an appropriate chemical

synthesis route can generally be readily identified upon determining the chemical structure.

Peptidomimetics can be synthesized and assembled into libraries comprising a few to many discrete molecular species. Such libraries can be prepared using well-known methods of combinatorial chemistry, and can be screened to determine if the library comprises one or more peptidomimetics which have the desired activity. Such peptidomimetic antagonists can then be isolated by suitable methods.

Small molecule antagonists

10 Cadherin-11 antagonists can also be small molecules. Examples of small molecules include organic compounds, organometallic compounds, inorganic compounds, and salts of organic, organometallic or inorganic compounds. Atoms in a small molecule are typically linked together via covalent and/or ionic bonds. The arrangement of atoms in a small organic molecule may represent a chain (*e.g.* a carbon-carbon chain or a carbon-heteroatom chain), or may represent a ring containing carbon atoms, *e.g.* benzene or a polycyclic system, or a combination of carbon and heteroatoms, *i.e.*, heterocycles such as a pyrimidine or quinazoline. Although small molecules can have any molecular weight, they generally include molecules that are less than about 5,000 daltons. For example, such small molecules can be less than about 1000 daltons and, preferably, are less than about 750 daltons or, more preferably, are less than about 500 daltons. Small molecules and other non-peptidic Cadherin-11 antagonists can be found in nature (*e.g.*, identified, isolated, purified) and/or produced synthetically (*e.g.*, by traditional organic synthesis, bio-mediated synthesis, or a combination thereof). See *e.g.* Ganesan, *Drug Discov.* 25 Today 7(1): 47-55 (January 2002); Lou, *Drug Discov. Today*, 6(24): 1288-1294 (December 2001). Examples of naturally occurring small molecules include, but are not limited to, hormones, neurotransmitters, nucleotides, amino acids, sugars, lipids, and their derivatives.

30 A small molecule Cadherin-11 antagonist according to the present invention, and physiologically acceptable salts thereof, can inhibit the homotypic binding of a Cadherin-11 protein (*e.g.*, by directly competing with a donor sequence in the EC1 domain of a Cad-11 protein for binding to the binding pocket of another Cadherin-

11, by directly competing with the binding pocket in the EC1 domain of a Cad-11 protein for binding to a donor sequence of another Cadherin-11).

Nucleic acid antagonists

5 Cad-11 antagonists of the invention can also be nucleic acid molecules (*e.g.*, oligonucleotides) that bind to the EC1 domain of a human Cadherin-11. Suitable nucleic acid Cad-11 antagonists include aptamers, which are capable of binding to a particular molecule of interest (*e.g.*, the EC1 domain of human Cadherin-11) with high affinity and specificity through interactions other than classic Watson-Crick 10 base pairing (Tuerk and Gold, *Science* 249:505 (1990); Ellington and Szostak, *Nature* 346:818 (1990)).

Aptamers, like peptides generated by phage display or monoclonal antibodies (MAbs), are capable of specifically binding to selected targets and, through binding, block their targets' ability to function. Created by an *in vitro* 15 selection process from pools of random sequence oligonucleotides, aptamers have been generated for over 100 proteins including growth factors, transcription factors, enzymes, immunoglobulins, and receptors. A typical aptamer is 10-15 kDa in size (30-45 nucleotides), binds its target with sub-nanomolar affinity, and discriminates against closely related targets (*e.g.*, will typically not bind other proteins from the 20 same gene family). A series of structural studies have shown that aptamers are capable of using the same types of binding interactions (hydrogen bonding, electrostatic complementarity, hydrophobic contacts, steric exclusion, etc.) that drive affinity and specificity in antibody-antigen complexes.

An aptamer that binds to a target of interest (*e.g.*, an EC1 domain of a human 25 Cad-11 protein) can be generated and identified using a standard process known as "Systematic Evolution of Ligands by Exponential Enrichment" (SELEX), described in, *e.g.*, U.S. Pat. Nos. 5,475,096 and U.S. Pat. No. 5,270,163.

Identification of Cadherin-11 antagonists

30 Agents having Cadherin-11 binding specificity, including small molecules, can be identified in a screen, for example, a high-throughput screen of chemical compounds and/or libraries (*e.g.*, chemical, peptide, nucleic acid libraries).

Antibodies that specifically bind the EC1 domain of human Cadherin-11 can be identified, for example, by screening commercially available combinatorial antibody libraries (Dyax Corp., MorphoSys AG). Suitable combinatorial antibody libraries and standard methods of screening these libraries are described in Hoet *et al.*, *Nature Biotechnology* 23(3):344-348 (2005) and Rauchenberger *et al.*, *J. Biol. Chem.* 278(40):38194-38205 (2003), the contents of which are incorporated herein by reference. Such libraries or collections of molecules can also be prepared using well-known chemical methods.

Alternatively murine antibodies that specifically bind the EC1 domain of human Cadherin-11 can be identified, for example, by immunizing mice with EC1 protein domains or EC1 peptides along with an adjuvant to break tolerance to the antigen. These antibodies can be screened for the desired specificity and activity and then humanized using known techniques to create suitable agents for the treatment of human disease.

Compounds or small molecules can be identified from numerous available libraries of chemical compounds from, for example, the Chemical Repository of the National Cancer Institute and the Molecular Libraries Small Molecules Repository (PubChem), as well as libraries of the Institute of Chemistry and Cell Biology at Harvard University and other libraries that are available from commercial sources (e.g., Chembridge, Peakdale, CEREP, MayBridge, Bionet). Such libraries or collections of molecules can also be prepared using well-known chemical methods, such as well-known methods of combinatorial chemistry. The libraries can be screened to identify compounds that bind and inhibit Cadherin-11.

Identified compounds can serve as lead compounds for further diversification using well-known methods of medicinal chemistry. For example, a collection of compounds that are structural variants of the lead can be prepared and screened for Cadherin-11 binding and/or inhibitory activity. This can result in the development of a structure activity relationship that links the structure of the compounds to biological activity. Compounds that have suitable binding and inhibitory activity can be developed further for *in vivo* use.

Agents that bind Cadherin-11 can be evaluated further for Cadherin-11 antagonist activity. For example, a composition comprising a Cadherin-

11 protein can be used in a screen or binding assay to detect and/or identify agents that bind and antagonize the Cadherin-11 protein. Compositions suitable for use include, for example, cells that naturally express a Cadherin-11 protein (e.g., a synoviocyte), extracts of such cells, and recombinant Cadherin-11 protein.

5 An agent that binds a Cadherin-11 protein can be identified in a competitive binding assay, for example, in which the ability of a test agent to inhibit the binding of Cadherin-11 to a reference agent is assessed. The reference agent can be a full-length Cad-11 protein or a portion thereof that comprises the EC1 domain. The reference agent can be labeled with a suitable label (e.g., radioisotope, epitope label, 10 affinity label (e.g., biotin and avidin or streptavidin), spin label, enzyme, fluorescent group, chemiluminescent group, dye, metal (e.g., gold, silver), magnetic bead) and the amount of labeled reference agent required to saturate the Cadherin-11 protein in the assay can be determined. The specificity of the formation of the complex between the Cadherin-11 protein and the test agent can be determined using a 15 suitable control (e.g., unlabeled agent, label alone).

The capacity of a test agent to inhibit formation of a complex between the reference agent and a Cadherin-11 protein can be determined as the concentration of test agent required for 50% inhibition (IC₅₀ value) of specific binding of labeled reference agent. Specific binding is preferably defined as the total binding (e.g., 20 total label in complex) minus the non-specific binding. Non-specific binding is preferably defined as the amount of label still detected in complexes formed in the presence of excess unlabeled reference agent. Reference agents suitable for use in the method include molecules and compounds which specifically bind to Cadherin-11, e.g., an antibody that binds Cadherin-11.

25 An agent that antagonizes a Cadherin-11 protein can be identified by screening for agents that have an ability to antagonize (reduce, prevent, inhibit) one or more activities of Cadherin-11, such as, for example, a binding activity (e.g., homotypic Cad-11 binding). Such activities can be assessed using an appropriate *in vitro* or *in vivo* assay. Exemplary assays for Cadherin-11 activity have been described 30 previously (Patel, SD, *et al.*, *Cell* 124: 1255-1268 (2006); Lee *et al.*, *Science* 315:1006-1010 (2007)).

Once a Cadherin-11 antagonist is identified, the ability of the Cadherin-11

antagonist to interfere with (e.g., reduce, inhibit, prevent) one or more biological functions or properties associated with Cadherin-11 activity in a cell can be assessed, for example, using a cell-based assay designed to measure a particular biological function or property associated with Cadherin-11. Biological functions and properties that are known to be associated with Cadherin-11 expression and/or activity include, but are not limited to, cell adhesion, cell migration, cell invasion, cell sorting, cell condensation, cell rearrangement, maintenance of tissue integrity and architecture, contact inhibition of cell proliferation and malignant transformation of cancer (e.g., tumor) cells (Kiener and Brenner, *Arthritis Res Ther.* 7(2):49-54 (2005)). In addition Cad-11 antagonists are shown herein to inhibit production of active MMPs by synoviocytes. Suitable assays for assessing one or more biological functions of cadherins are known to those of skill in the art (see, e.g., Patel, SD, *et al.*, *Cell* 124: 1255-1268 (2006)) and include, for example, the cell aggregation assay described herein (see Exemplification, Materials and Methods section).

Methods of Therapy

Without wishing to be bound by any one theory, it is believed that the first about 35 amino acids (e.g., about 33 to about 37 amino acids) of the EC1 domain of Cad-11 are required for homotypic cadherin binding and that agents that specifically bind to this region of Cad-11 can effectively inhibit binding between Cad-11 molecules. Accordingly, such agents are useful in the treatment and prevention of inflammatory joint disorders (e.g., rheumatoid arthritis) associated with Cad-11 expression and activity in synoviocytes and other cell types in inflamed joints. Thus, one aspect of the present invention relates to a method for treating an inflammatory joint disorder in a mammalian subject comprising administering to the subject a therapeutically effective amount of a Cadherin-11 antagonist that binds a human Cadherin-11 EC1 domain peptide (SEQ ID NO:3).

Using the methods of the invention, an inflammatory joint disorder in a mammal (e.g., a human) can be treated by administering a Cadherin-11 antagonist of the invention (e.g., antibodies, fusion proteins, small molecules, nucleic acids, peptides, peptidomimetics) in an amount that is sufficient to provide a therapeutic

benefit, for example, by inhibiting the aggregation of cells, or inhibiting the migration of cells, or inhibiting expression of active proteases or inflammatory molecules by cells, that express Cadherin-11 in an articulated joint (e.g., synoviocytes).

5 Accordingly, one aspect of the invention relates to a method for treating an inflammatory joint disorder in a mammalian subject comprising administering to the subject a therapeutically effective amount of a Cadherin-11 antagonist of the invention. The inflammatory joint disorder can be any disorder that is associated with or characterized by Cadherin-11 expression in cells (e.g., synoviocytes) of an 10 articulated joint. Examples of inflammatory joint disorders that can be treated by the present invention include, but are not limited to, rheumatoid arthritis, psoriatic arthritis, Reiter's syndrome and ankylosing spondylitis. In a particular embodiment, the inflammatory joint disorder is rheumatoid arthritis.

In one aspect, a therapeutically effective amount of a Cadherin-11 15 antagonist is administered to a patient in need thereof. The amount of the Cadherin-11 antagonist to be administered (e.g., a therapeutically effective amount) can be determined by a clinician using the guidance provided herein and other methods known in the art and is dependent on several factors including, for example, the particular agent chosen, the subject's age, sensitivity, tolerance to drugs and overall 20 well-being. For example, suitable dosages for Cad-11 antagonists that are antibodies can be from about 0.01 mg/kg to about 300 mg/kg body weight per treatment and preferably from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 1 mg/kg to about 10 mg/kg body weight per treatment. Suitable dosages for a small molecule Cad-11 antagonist can be from about 0.001 25 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.01 mg/kg to about 1 mg/kg body weight per treatment. Suitable dosages for Cadherin-11 antagonists that are proteins or peptides (linear, cyclic, mimetic), will result in a plasma concentration of the peptide from about 0.1 μ g/mL to about 200 μ g/mL. Determining the dosage for a 30 particular agent, patient and cancer is well within the abilities of one skilled in the art. Preferably, the dosage does not cause, or produces minimal, adverse side effects

(*e.g.*, immunogenic response, nausea, dizziness, gastric upset, hyperviscosity syndromes, congestive heart failure, stroke, pulmonary edema).

A therapeutically effective amount of a Cadherin-11 antagonist can be administered alone, or in combination with one or more other therapeutic agents (5 *e.g.*, anti-inflammatory agents). Suitable anti-inflammatory agents that are useful for treating inflammatory joint disorders, particularly RA, which can be administered in combination with Cad-11 antagonists of the invention, include, but are not limited to, (i) non-steroidal anti-inflammatory drugs (NSAIDs; *e.g.*, detopropfen, diclofenac, diflunisal, etodolac, fenoprofen, flurbiprofen, ibuprofen, 10 indomethacin, ketoprofen, meclofenamate, mefenamic acid, meloxicam, nabumeone, naproxen sodium, oxaprozin, piroxicam, sulindac, tolmetin, celecoxib, rofecoxib, aspirin, choline salicylate, salsalate, and sodium and magnesium salicylate); (ii) steroids (*e.g.*, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone); (iii) DMARDs, *i.e.*, 15 disease modifying antirheumatic drugs (*e.g.*, cyclosporine, azathioprine, methotrexate, leflunomide, cyclophosphamide, hydroxychloroquine, sulfasalazine, D-penicillamine, minocycline, and gold); or (iv) recombinant proteins (*e.g.*, ENBREL® (etanercept, a soluble TNF receptor), REMICADE® (infliximab, a chimeric monoclonal anti-TNF antibody), ORENCLIA® (abatacept, a soluble 20 CTLA4 receptor), ACTEMRA® (Tocilizumab, a monoclonal antibody to the IL-6 receptor), and RITUXAN® (rituximab, a monoclonal antibody to CD20)).

Thus, a Cadherin-11 antagonist can be administered as part of a combination therapy (*e.g.*, with one or more other therapeutic agents). The Cad-11 antagonist can be administered before, after or concurrently with one or more other therapeutic 25 agents. In some embodiments, the Cadherin-11 antagonist and other therapeutic agent can be co-administered simultaneously (*e.g.*, concurrently) as either separate formulations or as a joint formulation. Alternatively, the agents can be administered sequentially, as separate compositions, within an appropriate time frame as determined by the skilled clinician (*e.g.*, a time sufficient to allow an overlap of the 30 pharmaceutical effects of the therapies). The Cadherin-11 antagonist and one or more other therapeutic agents can be administered in a single dose or in multiple doses, in an order and on a schedule suitable to achieve a desired therapeutic effect

(*e.g.*, a reduction in and/or inhibition of joint inflammation). Suitable dosages and regimens of administration can be determined by a clinician and are dependent on the agent(s) chosen, pharmaceutical formulation and route of administration, various patient factors and other considerations.

5 The effectiveness of a therapy (*e.g.*, the reduction or elimination of joint inflammation and/or the prevention or inhibition of joint inflammation) can be determined by any suitable method (*e.g.*, imaging (MRI, NMR)).

10 According to the methods of the invention, a therapeutically effective amount of a Cad-11 antagonist is administered to a mammalian subject to treat an inflammatory joint disorder. The term “mammalian subject” is defined herein to include mammals such as primates (*e.g.*, humans) cows, sheep, goats, horses, dogs 15 cats, rabbits, guinea pigs, rats, mice or other bovine, ovine, equine, canine feline, rodent and murine species.

Agents that are Cad-11 antagonists can be administered to a mammalian 15 subject by a variety of routes. For example, the agent can be administered by any suitable parenteral or nonparenteral route, including, for example, topically (*e.g.*, cream, ointment), or nasally (*e.g.*, solution, suspension). Parenteral administration can include, for example, intraarticular, intramuscular, intravenous, intraventricular, intraarterial, intrathecal, subcutaneous, or intraperitoneal administration. The agent 20 can also be administered orally (*e.g.*, in capsules, suspensions, tablets or dietary), transdermally, intradermally, topically, by inhalation (*e.g.*, intrabronchial, intranasal, oral inhalation or intranasal drops), transmucosally or rectally. Administration can be local or systemic as appropriate, and more than one route can be used 25 concurrently, if desired. Localized administration of a Cad-11 antagonist can be achieved by intraarticular injection (*e.g.*, direct injection of the agent into a joint). The preferred mode of administration can vary depending upon the particular agent chosen. However, systemic intravenous or subcutaneous administration is generally preferred for antibodies.

Delivery can also be by injection into the brain or body cavity of a patient or 30 by use of a timed release or sustained release matrix delivery systems, or by onsite delivery using micelles, gels and liposomes. Nebulizing devices, powder inhalers, and aerosolized solutions are representative of methods that may be used to

administer such preparations to the respiratory tract. Delivery can be *in vitro*, *in vivo*, or *ex vivo*.

Agents that are proteins (*e.g.*, fusion protein) can be administered via *in vivo* expression of recombinant protein. *In vivo* expression can be accomplished by 5 somatic cell expression according to suitable methods (see, *e.g.*, U.S. Patent No. 5,399,346). Further, a nucleic acid encoding the protein can also be incorporated into retroviral, adenoviral or other suitable vectors (preferably, a replication deficient infectious vector) for delivery, or can be introduced into a transfected or transformed host cell capable of expressing the protein for delivery. In the latter 10 embodiment, the cells can be implanted (alone or in a barrier device), injected or otherwise introduced in an amount effective to express the protein in a therapeutically effective amount.

Nucleic acid-based Cadherin-11 antagonists (*e.g.*, aptamers) can be introduced into a mammalian subject of interest in a number of ways. For instance, 15 nucleic acids may be expressed endogenously from expression vectors or PCR products in host cells or packaged into synthetic or engineered compositions (*e.g.*, liposomes, polymers, nanoparticles) that can then be introduced directly into the bloodstream of a mammalian subject (by, *e.g.*, injection, infusion). Anti-Cadherin-11 nucleic acids or nucleic acid expression vectors (*e.g.*, retroviral, adenoviral, 20 adeno-associated and herpes simplex viral vectors, engineered vectors, non-viral-mediated vectors) can also be introduced into a mammalian subject directly using established gene therapy strategies and protocols (see *e.g.*, Tochilin V.P. *Annu Rev Biomed Eng* 8:343-375, 2006; Recombinant DNA and Gene Transfer, Office of Biotechnology Activities, National Institutes of Health Guidelines).

25 Agents that are Cadherin-11 antagonists (*e.g.*, small molecules) can be administered to a mammalian subject as part of a pharmaceutical or physiological composition, for example, as part of a pharmaceutical composition comprising a Cadherin-11 antagonist and a pharmaceutically acceptable carrier. Formulations or compositions comprising a Cadherin-11 antagonist or compositions comprising a 30 Cadherin-11 antagonist and one or more other therapeutic agents (*e.g.*, an anti-inflammatory agent) will vary according to the route of administration selected (*e.g.*, solution, emulsion or capsule). Suitable pharmaceutical carriers can contain inert

ingredients which do not interact with the Cadherin-11 antagonist. Standard pharmaceutical formulation techniques can be employed, such as those described in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA. Suitable pharmaceutical carriers for parenteral administration include, for example, 5 sterile water, physiological saline, bacteriostatic saline (saline containing about 0.9% mg/ml benzyl alcohol), phosphate-buffered saline, Hank's solution, Ringer's lactate and the like. Formulations can also include small amounts of substances that enhance the effectiveness of the active ingredient (*e.g.*, emulsifying agents, solubilizing agents, pH buffering agents, wetting agents). Methods of encapsulation 10 compositions (such as in a coating of hard gelatin or cyclodextran) are known in the art. For inhalation, the agent can be solubilized and loaded into a suitable dispenser for administration (*e.g.*, an atomizer or nebulizer or pressurized aerosol dispenser).

The pharmaceutical agent can be administered as a neutral compound or as a salt or ester. Pharmaceutically acceptable salts include those formed with free 15 amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic or tartaric acids, and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc. Salts of compounds containing an amine or other basic group can be obtained, for example, by reacting 20 with a suitable organic or inorganic acid, such as hydrogen chloride, hydrogen bromide, acetic acid, perchloric acid and the like. Compounds with a quaternary ammonium group also contain a counteranion such as chloride, bromide, iodide, acetate, perchlorate and the like. Salts of compounds containing a carboxylic acid or other acidic functional group can be prepared by reacting with a suitable base, for 25 example, a hydroxide base. Salts of acidic functional groups contain a countercation such as sodium or potassium.

The present invention will now be illustrated by the following Examples, which are not intended to be limiting in any way.

Exemplification

Example 1: Identification of Fabs having binding specificity for an epitope within the N-terminal 35 amino acids of the EC1 domain of human Cadherin-11.

5

Materials and Methods

Western blotting

Proteins were separated by SDS-PAGE and transferred to a nitrocellulose (NC) membrane using standard methods. Briefly NC membrane was rinsed with tris buffered-saline-tween (TBST) (8.8 g/L of NaCl, 0.2g/L of KCl, 3g/L of Tris base, 500ul/L of Tween-20, pH to 7.4). The membrane was blocked with 4% BSA dissolved in TBST for hour at 22°C. The NC membrane was rinsed 3X for 5 min each with TBST. Mouse anti-human Cad-11 antibody was diluted to 0.5 µg/ml in TBST and the NC was incubated for 1 hour at 22°C. The NC membrane was rinsed 3X for 5 minutes each in TBST. Goat anti-mouse Ig antibody conjugated with horse radish peroxidase (HRP) was diluted to 1 µg/ml in TBST and the NC membrane was incubated in secondary solution for a minimum time of 1 hour @ room temperature (RT) at 22°C. The NC membrane was rinsed 3X for 5 min each in TBST. Signal was developed using standard HRP method.

20

ELISA

The antigen (either 5 µg/ml or 50 µg of Cad-11-EC-1-Fc or 5 µg/ml of Cadherin peptide) was diluted in a buffer and used to coat the plates overnight at 4°C. The plates were washed and then blocked with 1.5 % BSA, 5% low fat milk powder in PBS Dilution buffer: 1.5 % BSA, 2.5% low fat milk powder, 0.1% Tween-20 in PBS. The plates were then incubated with bacterial lysate containing the anti-Cad-11 human fAbs or purified anti-Cad-11 human fAbs for 1 hr. After washing, the secondary antibody (Cy5-conjugated a-hu-Fab diluted 1/100) was applied for 25 min. The plates were then washed and the resulting fluorescence read.

30

Results

Three sets of previously reported Cadherin-11-specific antibodies were tested for an ability to bind to the EC1 domain of human Cadherin-11. These antibodies included antibodies that were raised against a mouse Cadherin-11-Fc fusion protein 5 immunogen in Cadherin-11 knock-out or deficient mice (Lee *et al.*, *Science* 315:1006-1010 (2007)), antibodies that were raised in Cadherin-11 wild type mice against a human Cadherin-11-Fc fusion protein immunogen that had been produced in CHO cells (Valencia *et al.*, *J. Exp. Med.* 200(12):1673-1679 (2004)), and antibodies that were raised in Cadherin-11 wild type mice against a bacterially- 10 produced protein containing the EC1-3 domains of human Cadherin-11. These antibodies were tested by western analysis for an ability to bind fusion proteins that contained only the EC1 domain of human Cad-11(Cadherin-11-EC1-Fc), the EC1 and EC2 domains of Cad-11 (Cad11-EC1/2-Fc) or all 5 EC domains of Cad-11 (Cadherin-11-EC1-5-Fc). None of the antibodies tested recognized the EC1-Fc or 15 the EC1-2-Fc fusion proteins on a Western blot (FIG. 1A). However, antibodies from each of the three sets tested recognized the human Cad-11-Fc fusion protein that included extracellular domains 1 through 5 (FIG. 1B). These results indicate that the tested antibodies did not bind to the EC1 or EC2 domains of human Cad-11, but recognized epitopes elsewhere in the extracellular region of this protein.

20 The available published anti-Cad11 antibodies that bind Cad11 expressing cells, 13C2, 23C6, 5F82 (Lifespan Science) and 283416 (R&D Systems), as well as the Cad11 EC1-binding antibody H1M1, and the control antibody, MOPC, were tested by ELISA for the ability to bind fusion proteins that contained only the EC1 domain 25 of human Cad-11 (Cadherin-11-EC1-Fc) or all 5 EC domains of Cad-11 (Cadherin-11-EC1-5-Fc). None of the available published anti-Cad11 antibodies tested recognized the EC1-Fc (FIG. 1B, open bars) (data for 283416 is not shown here). However, the Cad11 EC1 binding H1M1 antibody bound both the Cadherin-11-EC1-Fc and Cadherin-11-EC1-5-Fc (FIG. 1B closed bar). The control MOPC 30 antibody bound neither fusion protein. These results indicate that the available published anti-Cad11 antibodies do not bind to the EC1 domain of human Cad-11, but recognized epitopes elsewhere in the extracellular region of this protein.

To create an antibody specific for an epitope within the N-terminal 35 amino acids of the EC1 domain of human Cadherin-11, a phage display library (MorphoSys AG) 5 encoding human Fabs was screened. Candidate Fabs were identified using two selection criteria - a positive selection for binding to a peptide that included the first 35 amino acids of the human Cadherin-11 EC1 domain, and a negative selection for binding to corresponding peptides from the EC1 domains of two closely related and highly homologous cadherins, Cadherin-8 and MN-Cadherin (FIG. 2). ELISA was 10 used to assess binding.

Two screens were conducted. In the first screen, 96 Fab clones that bound the Cadherin-11 EC1 peptide were identified by ELISA. Seven (7) candidate Fabs bound the Cad-11 EC-1 peptide; however, only two of these bound to both EC-1 15 peptide and the EC1-2-Fc fusion protein. One of these two Fabs also bound to MN-Cad peptide. Accordingly, only one of the seven Fab clones specifically bound the EC1-Fc fusion protein, but did not bind to both MN-Cad and Cadherin-8 EC1 domain peptides. In a second screen, similar results were observed, as only 1 of 96 Fabs (clone F9) showing specificity for the Cadherin-11 EC1 peptide and EC1-2-Fc 20 fusion protein, failed to bind MN-Cad and Cadherin-8 EC1 domain peptides (FIG. 3). The majority of the Cad-11 EC1 domain-binding Fabs tested showed cross reactivity with the MN-Cad peptide, which contains an EEY CAR sequence that overlaps with the EEY CAR sequence of Cad-11.

25

Example 2: A Fab that binds the EC1 domain of Cadherin-11 inhibits Cad-11 mediated cell aggregation in an *in vitro* assay

Materials and Methods

30 *In vitro* Cadherin-11 aggregation assay

431-D cells grow in suspension and do not normally express any cadherins and do not aggregate. 431-D-11 cells have been genetically modified to express Cad-11.

When 431-D-11 cells are incubated in media alone and they begin to aggregate over 40 min and the clumps of cells settle to the bottom of well and the remaining non-aggregated cells in suspension can be measured and the percentage of aggregated 431-D-11 calculated. For the aggregation assay, 431-D-11 cells (D-11 cells) were 5 grown to sub-confluence in a flask and then were removed from the flask using 0.05% Trypsin plus 0.53 mM EDTA. Approximately 2×10^6 431-D-11 cells were added to 2 ml of SME media (Dulbecco's Modified Eagle's Medium-high glucose, 0.1M Hepes pH 7.4 and 5U/ml DNase) and were preincubated for 15 min on ice, either in the absence or presence of a test agent (e.g., antibody, Fab, fusion protein). 10 After pre-incubation with the test agent, the cells were transferred to a round bottom well on a 24-well plate and incubated at 37°C while rotating at 130 rpm on a rotary shaker. As cells aggregate they sink to the bottom of the well. At 0 min and 40 min, 200 μ l from the middle of the sample were removed from the well and mixed with 25 μ l of 8% glutaraldehyde to fix the cells. 200 μ l of the fixed sample of cells were 15 added to 9.8 ml of Coulter Counter isotonic saline solution and counted using a Coulter Counter set at the 8 μ m to 24 μ m threshold. 3 cell counts per sample were recorded. The percentage of cell decrease or aggregated cells at 40 min. compared to the percentage at the 0 min. time point was calculated.

20 Results

A candidate Fab (clone F9) having binding specificity for an epitope within the N-terminal 35 amino acids of the Cadherin-11 EC1 domain, which does not bind the EC1 domains of MN-Cad or Cad-8, was tested for an ability to inhibit Cad-11 25 mediated cell aggregation using an *in vitro* Cadherin-11 cell aggregation assay. The candidate Fab significantly inhibited Cadherin-11 mediated aggregation of cells at concentrations of 1 μ g/ml or lower (FIGS. 4 and 5). In contrast, a Fab made from the 13C2 antibody that binds to an epitope in the extracellular region of Cad-11 outside the EC1/2 domains inhibited Cadherin-11 aggregation only at a concentration of 10 μ g/ml, suggesting that the F9 Fab inhibits Cad-11 activity more 30 effectively at lower concentrations than antibodies which bind to other portions of the extracellular domain of Cad-11.

Cad-11 mediated cell aggregation was also inhibited by various anti-Cadherin-11-EC1 domain Fabs that were specific for either Cad-11 alone, Cad-11 and Cad-8, Cad-11 and MN-Cad, or Cad-11, Cad-8 and MN-Cad (FIGS. 6 and 7). All cadherin-EC1-domain specific Fabs that were tested inhibited cell aggregation *in vitro* relative to the control samples (e.g., SME medium (FIG. 6), a Fab specific for GFP (FIG. 7).

Example 3: Generation of Cadherin-11/Immunoglobulin Fusion Proteins Containing the EC1 domain of human Cadherin-11.

10 The Cadherin-11 EC1 region was prepared from a vector encoding the full length human Cadherin-11 cDNA (human Cad-11 cloned into the Not1 and Kpn-1 sites of the Invitrogen pCEP4® vector) using polymerase chain reaction (PCR) performed under standard conditions using the following oligonucleotide primers to introduce EcoR1 and BglII sites (see underlined sequences in forward and reverse primers, 15 respectively) into the amplified product:

Forward Primer:

tttttttttgaattcatgaaggagaactactgtttacaagc (SEQ ID NO:8)
EcoR1

20

Reverse Primer:

tttttttttagatctctggaccttgacaatgaattccgacgg (SEQ ID NO:9)
BglII

25 The amplified product was digested with restriction enzymes EcoR1 and BglII, and the digestion product was isolated and ligated into the pFUSE-hIgG2e1-Fc1 vector (InvivoGen) using the corresponding EcoR1 and BglII sites. TOP10 competent bacteria (Invitrogen) were transformed as described by the manufacturer with the ligation product and bacteria with the Cadherin-11-EC1-Fc plasmid were selected 30 with zeomycin. Cadherin-11-EC1-Fc plasmid was isolated, sequenced and then used to transiently transfect 293F cells. Conditioned media was collected and the Cadherin-11-EC1-Fc fusion protein (SEQ ID NO:9) was purified using tangential flow filtration followed by isolation on a 50/50 mix protein A/protein G column equilibrated in 20 mM HEPES pH 7.4, 137mM NaCl, 3mM KCl and 1mM CaCl2.

The purified Cadherin-11-EC1-Fc fusion protein was eluted from the column using 0.1 M Glycine (pH 3) and 1 mM CaCl₂ and into tubes containing 200 µl of 1M Tris pH 7.4, and 1mM CaCl₂. The eluates with the Cad-11 fusion protein were then dialyzed against 20 mM Hepes 5 (pH 7.4), 137 mM NaCl, 3 mM KCl and 1 mM CaCl₂. The size of the protein was confirmed by SDS PAGE (FIG. 10) and identity was confirmed by Western analysis using an antibody that recognizes the human Fc region (FIG. 11) and N-terminal sequencing (not shown). Cad-11-EC1-2-Fc was produced using techniques and conditions similar 10 those described above.

Example 4: A Cad-11-EC1-Fc immunoglobulin fusion protein inhibits Cad-11 mediated cell aggregation in an *in vitro* assay

15 Materials and methods

Cell invasion/migration into a matrigel plug

FLS migratory activity was assessed in Matrigel ECM-coated transwells in FLS media (Dulbecco's Modified Eagle's Medium-high glucose [Sigma #D7777], 10% Fetal Bovine Serum [Benchmark #100-106], 1% Pencillin-Streptomycin [Gibco 20 315140-122], 1% L-Glutamine [Gibco #25030], 0.5% Gentamicin [Gibco #15710-064]. Human FLS cell suspensions in FLS medium containing 1×10^4 cells were added to the control insert or matrigel coated insert set in the well of a 24-well plate containing 0.750 mL of FLS medium. The plates were then incubated in a humidified tissue culture 25 incubator at 37°C, 5% CO₂ atmosphere for 22 hours.

To calculate the number of cells that migrated, non-invading cells were removed 30 from the upper surface of the membrane of control inserts by wiping with a cotton swab. A second wipe using a cotton swap wetted with FLS medium is repeated. Control inserts were then fixed and stained using a differential staining kit [Fisher #122-911]. Inserts are allowed to dry and cells are counted in 4 quadrants of the

control insert using a microscope with a 10x objective. Triplicate inserts are counted and the totals averaged.

To calculate the number of cells that invaded the matrigel inserts, non-invading cells
5 were gently removed from the surface of the matrigel insert by wiping with a cotton swab. A second wipe using a cotton swap wetted with FLS medium is repeated. Inserts were then fixed and stained using a differential staining kit [Fisher #122-911]. Inserts are allowed to dry and cells are counted in 4 quadrants of the control
10 insert using a microscope with a 10x objective. Triplicate inserts are counted and the totals averaged.

Results

Cad-11-EC1-Fc significantly inhibited cell aggregation at a concentration of 3 µg/ml, while the full length Cad-11-EC1-5-Fc protein containing all 5 EC domains
15 of human Cad-11 inhibited Cad-11 mediated aggregation at a concentration of 100 µg/ml (FIG. 13). These data show that the Cad-11-EC1-Fc immunoglobulin fusion protein effectively inhibits Cad-11 mediated cell aggregation in an *in vitro* assay.

In addition, the ability of the Cad-11-EC1-Fc immunoglobulin fusion protein to
20 inhibit the invasion of human fibroblast like synoviocytes (FLS) into a matrigel plug was tested *in vitro*. Invasion of the FLS into matrigel is a complex process that involves the expression of MMP1, MMP-3, MMP-13, serine proteases, and other proteins by the FLS to degrade the matrigel as well as the migration of the FLS into matrigel. In a separate assay we saw no inhibition of migration of FLS through a
25 normal fiber insert. This suggests the impact of the EC-Fc or 13C2 mAb is to inhibit the degradation of the matrigel (a surrogate for joint cartilage). Both the Cad-11-EC1-Fc and murine anti-Cad-11 mAb 13C2 inhibited FLS invasion into a matrigel plug in two independent experiments.

Example 5: Generation of antibodies against an EC1 domain peptide of human Cadherin-11

Materials and Methods

5 Balb/c mice were immunized bi-weekly in the foot pad nine times over a one month period with 0.01 mg of a peptide corresponding to the first 33 amino acids of the human Cad-11 EC1 domain (GWWWN QFFVI EEYTG PDPVL VGRLH SDIDS GDG (SEQ ID NO:10)), covalently linked to BSA. This peptide is referred to herein as Peptide 4. Spleens from the immunized mice were harvested and fused

10 with a murine fusion partner, P3X63-Ag8.653, to create antibody-producing hybridomas. The hybridomas were expanded and subcloned at either 10, 3 or 0.5 cells/well, and the anti-Cad-11 antibody-containing media from the hybridomas were screened in an ELISA for the ability to specifically bind Cad-11 EC1-2 domain-containing protein produced in bacteria. Anti-Cad-11 antibody-containing

15 media from these Peptide 4 hybridomas were screened concurrently for the absence of binding to proteins encompassing the EC1-2 domains of human Cad-8 and MN-Cadherin. 96-well EIA plates were coated overnight at 4°C with 0.05 ml of 0.0 to 0.3 mg/ml of each of the EC1-2 Cad proteins and then washed several times with saline buffer. Plates were then blocked with 0.25 ml of casein-PBS buffer and

20 washed several times with saline buffer. Hybridoma media containing the anti-Cad-11 antibody were incubated neat in each well for 1 hr at 22°C and then washed twice with PBS-Tween (0.05%). 100 µl of a 1/1000 dilution of a goat anti-mouse IgG secondary antibody were added to each well, incubated for 30 min at 22°C, and then washed twice with PBS-Tween (0.05%). 100 µl/well of room temperature TMB (3, 25 3', 5, 5'-tetramethylbenzidine) reagent was added to each well and color was allowed to develop for 5 min at 22°C. The reaction was stopped with 100 µl of room temperature 2N sulfuric acid and the plate was read at 450 nm on a Wallac 1420 microplate reader.

30 The specificity of H1M1 and H14 anti-Cad-11 antibodies was tested further using an ELISA against the first 33 amino acids of the EC1 domains of Cad-11, Cad-7, Cad-8, Cad-20, Cad-24, Cad-9, Cad-18, and MN-Cad. Peptides corresponding to the

region of Cad-7, Cad-8, Cad-20, Cad-24, Cad-9, Cad-18, MN-Cad that overlapped with the G1-G33 region of the Cad-11 EC1 domain were synthesized and conjugated to biotin. 100 μ l of a 30 ng/ml solution of each of these peptides in PBS-Tween (0.05%) were incubated in each well of a 96-well Netravidin plate for 2-3 hrs 5 at 4°C and then washed twice with PBS-Tween (0.05%). Various concentrations of the anti-Cad-11 antibody were incubated in each well for 1 hr at 22°C and then washed twice with PBS-Tween (0.05%). 100 μ l of a 1/1000 dilution of a goat anti-mouse IgG secondary antibody were added to each well, incubated for 30 min at 22°C, and then washed twice with PBS-Tween (0.05%). 100 μ l/well of room 10 temperature TMB (3, 3', 5, 5'-tetramethylbenzidine) reagent was added to each well and color was allowed to develop for 5 min at 22°C. The reaction was stopped with 100 μ l of room temperature 2N sulfuric acid and the plate was read at 450 nm on a Wallac 1420 microplate reader.

15 Media from wells containing positive anti-Cad-11 antibody hybridomas were tested for the ability to bind to Cad-11 expressing cells. Frozen Cad-11-expressing 431D cells were thawed and washed twice in Hanks Balanced Saline Solution (HBSS) containing Ca^{2+} (0.137 M NaCl, 5.4 mM, KCl 0.25, mM Na_2HPO_4 , 0.44 mM KH_2PO_4 , 1.3 mM CaCl_2 , 1.0 mM MgSO_4 and 4.2 mM NaHCO_3) and then 20 resuspended at 10^6 cells/ml in HBSS containing Ca^{2+} . 10^5 cells/well were stained with either a 50% or 16% anti-Cad-11 antibody media for 45 min on ice, washed twice in HBSS containing Ca^{2+} , stained with a secondary goat anti-mouse IgG antibody conjugated with phytoerytherin (Jackson ImmunoResearch, West Grove, PA) a concentration of 1% for 45 min on ice and then washed again twice in HBSS 25 containing Ca^{2+} . Cells were then resuspended in 400 μ l of HBSS containing Ca^{2+} and 1% formaldehyde and subsequently analyzed on a FACScalibur (Becton Dickenson, Franklin Lakes, NJ) for PE positive cells.

Results

30 Anti-Cad-11 antibody-containing media from the Peptide 4 hybridomas bound to the Cad-11 EC1-2 protein (FIG. 16, HL vs CAD11), but not proteins containing the EC1-2 domains of Cad-8 and MN-Cad (FIG. 16, HL vs CAD8 and HL vs MNCAD,

respectively). Control hybridoma media did not bind any of the cadherin proteins tested (FIG. 16, Media vs CAD11, Media vs CAD8, and Media vs MNCAD). These data demonstrate the presence of Cad-11 specific antibodies to Peptide 4 in the hybridomas.

5

Two Peptide 4 hybridomas, referred to herein as H1M1 and H14, bound to cells expressing human Cad-11 protein (FIGS. 17A-C and 17G-I), but not to non-Cad-11 control 431-D cells (FIGS. 17D-17F). The hybridoma cell line referred to as H1M1 has the A.T.C.C. designation number _____, having been deposited on January 8, 10 2009. The hybridoma cell line referred to as H14 has the A.T.C.C. designation number _____, having been deposited on January 9, 2009. These hybridomas contain anti-Cad-11 antibodies that recognize both Peptide 4 and Cad-11-expressing cells *in vitro*. The binding of these antibodies to Cad-11-expressing cells was shown to titrate with the amount of Peptide 4 antibody that was used, as shown in the plots 15 of the titration of H1M1 (FIG. 18A) and H14 (FIG. 18B) versus the intensity of cell staining from the mean fluorescence intensity (MFI).

The H1M1 and H14 Peptide 4 anti-Cad-11 antibodies demonstrated >100-fold higher binding to Cad-11 than to any of the other cadherins tested, which included 20 Cad-7, Cad-8, Cad-20, Cad-24, Cad-9, Cad-18, and MN-Cad. In most cases, no binding of H1M1 and H14 anti-Cad-11 antibodies to the other cadherins was observed. The anti-Cad-11 antibody H14 showed strong binding to Cad-11 (FIG. 19A), with 468-fold lower binding to Cad-8 (FIG. 19A), and virtually no binding to Cad-7, MN-Cad, Cad-9, Cad-18, Cad-20 or Cad-24 (FIG. 19B). Similarly, the anti- 25 Cad-11 antibody H1M1 showed strong binding to Cad-11 (FIG. 20), with 365-fold lower binding to Cad-8 (FIG. 20), and no binding to Cad-7, MN-Cad, Cad-9, Cad-18, Cad-20 or Cad-24 (data not shown).

Example 6: The anti-Cad-11 EC1 domain antibodies H1M1 and H14 bind epitopes in the Cad-11 EC1 domain that include the amino acid sequence GPDP

Materials and Methods

5 To determine the epitope within the Cad-11 EC1 domain that the Peptide 4 Cad-11 EC1 antibodies H1M1 and H14 bind, four different peptides spanning the first 37 amino acids of the EC1 region (see FIG. 22) were immobilized in an ELISA format and the ability of the H1M1 and H14 antibodies to bind each of the four peptides was determined. 96-well Reactibind plates were coated overnight at 4°C with 0.3
10 ng/well of Peptide 1 (amino acids G1-P18 of the Cad-11 EC1 domain), 0.3 ng/well of Peptide 2 (amino acids G15-N34 of the Cad-11 EC1 domain), 0.3 ng/well of Peptide 3 (amino acids V19-Y37 of the Cad-11 EC1 domain), 0.3 ng/well of the immunogen Peptide 4 (amino acids G1-G33 of the Cad-11 EC1 domain), 20 ng of a fusion protein including the entire EC1 domain (EFL), or 20 ng of control human Ig
15 (Fc-block). The wells were washed twice with PBS-Tween (0.05%), blocked with casein in dH₂O for 3 hrs at 22°C and then washed again twice with PBS-Tween (0.05%). Various dilutions of the different Peptide 4 CAD-11 EC1 domain antibodies were transferred to the peptide- or protein-coated wells, incubated for 45 min at 22°C and then washed twice with PBS-Tween (0.05%). 100 µl of a 1/1000
20 dilution of goat anti-mouse IgG secondary antibody (Jackson ImmunoResearch, West Grove, PA) were added to each well, incubated for 30 min at 22°C and then washed twice with PBS-Tween (0.05%). 100 µl/well of room temperature TMB reagent was added to each well and color was allowed to develop for 5 min at 22°C. The reaction was stopped with 100 µl of room temperature 2 N sulfuric acid and the
25 plate was read at a wavelength of 450 nm on a Wallac 1420 microplate reader.

Results

The Peptide 4 anti-Cad-11 antibodies H1M1 at 1:11 (FIG. 21A) and H14 at 1:23 (FIG. 21B) both bound the Peptide 4 (PEP4) immunogen, as well as the EC1 domain fusion protein (EFL), in the ELISA as indicated by elevated OD450 plate readings relative to the control. Neither of these antibodies bound to the human IgG control

(Fc block). In addition, both antibodies bound Peptide 2 (PEP2), but not Peptide 1 (PEP1) or Peptide 3 (PEP3), in the ELISA (FIGS. 21A and 21B).

These results suggest that the anti-Cad-11 EC1 domain antibodies H1M1 and H14 bind a common epitope in Peptides 2 and 4 that is not present in the overlapping Peptide 3. Amino acids shared by Peptides 2 and 4 that are upstream of Peptide 3 are highlighted in the boxed region shown in FIG. 22. These four amino acids, GPDP (SEQ ID NO:11), beginning at G15 of the Cad-11 EC1 domain, are likely part of the epitope recognized by the H1M1 and H14 antibodies.

10

Example 7: The anti-Cad-11 EC1 domain antibodies H1M1 and H14 inhibit aggregation of Cad-11-expressing cells *in vitro*

Materials and Methods

15 To assess the ability of the Cad-11 antibodies to inhibit Cad-11 mediated cell aggregation, 30 μ g/ml of the H1M1 Peptide 4 antibody was cultured with 75,000 Cad-11 expressing A-431-D epidermoid carcinoma cells in 0.5 ml of DMEM-high glucose, 20mM Hepes pH 7.4, 10% FCS and 10U/ml DNase in a 24-well round bottom polypropylene plate. The 24-well plates were placed on a rotating platform 20 at approximately 60 rpm and incubated with 5% CO₂ overnight at 37°C. The next day, cell aggregation was assessed after photographing the plates at 100x (for H1M1 experiment) or 40x (for H14 experiment) magnification.

Results

25 In the presence of a control isotype antibody (30 μ g/ml), the Cad-11-expressing cells formed large masses (FIG. 23A), while the parental Cad-11 negative cells remain as single or double cell groups (FIG. 23C). The H1M1-treated Cad-11 cells remained as small clumps of cells (FIG. 23B) that did not progress to form the large masses obtained using the control antibody.

30

Using the same assay, the anti-Cad-11 antibody H14 was also shown to inhibit Cad-11-mediated aggregation. While the parental Cad-11-expressing cells formed large

clusters of aggregated cells (FIG. 24A), the H14 antibody (FIG. 24B) inhibited aggregation at a concentration of 30 μ g/ml, as cell clusters were small and infrequent. These results indicate that the anti-Cad-11 antibodies H1M1 and H14 inhibit Cad-11-mediated cell aggregation *in vitro*.

5

Example 8: The anti-Cad-11 EC1 domain antibodies, H1M1 and H14, inhibit arthritis-associated joint swelling *in vivo* in a murine model of rheumatoid arthritis

Materials and Methods

10 Study 1 – Six-week-old male C57/Bl6 mice were injected with 150 μ l of KBN sera on day 0 and day 2. KBN sera-treated mice received either saline injections (FIG. 25, unfilled triangles) or were treated with different doses of the H1M1 anti-Cad-11 EC1 antibody. Treatment regimens included dosing on day 0 with 0.5 mg of antibody/mouse and every second day (q2d) thereafter with 0.1 mg of antibody/mouse (0.5 mg+0.1 mg) (FIG. 25, filled triangles); dosing on day 0 with 0.5 mg of antibody/mouse (0.5 mg) (FIG. 25, diamonds); dosing every second day (q2d) with 0.1 mg of antibody/mouse (0.1 mg+0.1 mg) (FIG. 25, squares); or dosing every second day (q2d) with 0.3 mg of antibody/mouse (0.3 mg+0.3 mg) (FIG. 25, circles). The control group consisted of 5 mice and the treatment group consisted of 20 7 mice. Arthritis-associated joint swelling was determined by caliper measurements taken every second day.

25 Study 2 – Six-week-old male C57/Bl6 mice were injected with 150 μ l of KBN sera on day 0 and day 2, and then were treated with either saline every second day (q2d) (FIG. 26, triangles), or one of the anti-Cad-11 antibodies, H1M1 (FIG. 26, squares) or H14 (FIG. 26, circles), at 0.3mg/dose q2d. The control group consisted of 5 mice and the treatment group consisted of 7 mice. Arthritis-associated joint swelling was determined by caliper measurements taken every second day.

30 Results

Study 1 – The H1M1 anti-Cad-11 antibody inhibited joint swelling relative to the control mice. The greatest inhibition of arthritis-associated joint swelling was

observed by dosing KBN-treated mice with 0.3 mg of H1M1 antibody every second day (FIG. 25, circles).

5 Study 2 - Both of the anti-Cad-11 antibodies inhibited joint swelling relative to the control. In this study, the H14 antibody significantly delayed the onset of arthritis compared to the control animals (FIG. 27). All mice in the control group developed arthritis by day 3, while the H14-treated mice required 6 days before all of the animals developed arthritis.

10 These studies indicate that antibodies against the EC1 domain of human Cad-11 can inhibit the development and severity of arthritis *in vivo*.

Example 9: Generation of antibodies against another EC1 domain peptide of human Cadherin-11

15

Materials and Methods

Balb/c mice were immunized bi-weekly in the foot pad nine times over a 1 month period with 0.01 mg of peptide V19-Y37 (VL VGRLH SDIDS GDGNI KY (SEQ ID NO:12)), corresponding to 19 amino acids of the human Cad-11 EC1 domain, 20 covalently linked to BSA. This peptide is referred to herein as Peptide 3. Spleens from the immunized mice were harvested and fused with a murine fusion partner P3X63-Ag8.653, to create antibody-producing hybridomas. These hybridomas were expanded and the anti-Cad-11 antibody-containing media from the hybridomas were screened for the ability to bind to a protein corresponding to the EC1-2 domain of 25 Cad-11, which was produced in bacteria. The anti-Cad-11 antibody-containing media from these Peptide 3 hybridomas were screened concurrently for the absence of binding to proteins encompassing the EC1-2 domains of Cad-8 and MN-Cadherin. 96-well EIA plates were coated overnight at 4°C with 0.05 ml of 0 to 300 mg/ml of one of each of the EC1-2 Cad proteins, or CHO cell produced EC1-Fc 30 fusion protein, and then washed several times with saline buffer. Plates were then blocked using 0.25 ml of casein-PBS buffer and subsequently washed several times with saline buffer. Hybridoma media containing the Peptide 3 anti-Cad-11

antibodies were incubated neat in each well for 1 hr at 22°C and then washed twice with PBS-Tween (0.05%). 100 μ l of a 1/1000 dilution of a goat anti-mouse IgG secondary antibody were added to each well, incubated for 30 min at 22°C, and then washed twice with PBS-Tween (0.05%). 100 μ l/well of room temperature TMB (3, 5 3', 5, 5'-tetramethylbenzidine) reagent was added to each well and color was allowed to develop for 5 min at 22°C. The reaction was stopped with 100 μ l of room temperature 2N sulfuric acid and the plate was read at 450 nm on a Wallac 1420 microplate reader.

10 Media from the Peptide 3 hybridomas were also tested for the ability to bind to human Cad-11 protein expressed on cells. Frozen Cad-11-expressing 431D cells were thawed and washed twice in HBSS with Ca^{2+} and then resuspended at 10^6 cells/ml in HBSS containing Ca^{2+} . 10^5 cells/well were stained with either a 50% or 16% anti-Cad-11 antibody media for 45 min on ice, washed twice in HBSS

15 containing Ca^{2+} , and then stained with a secondary goat anti-mouse IgG antibody conjugated with phytoerytherin at a concentration of 1% for 45 on ice and then washed again twice in HBSS containing Ca^{2+} . Cells were then resuspended in 400 μ l of HBSS containing Ca^{2+} and 1% formaldehyde and subsequently analyzed on a FACScalibur for PE positive cells.

20

Results

Anti-Cad-11 antibody-containing media from the Peptide 3 hybridomas bound to the Cad-11 EC1-2 protein and the EC1-Fc fusion protein (FIG. 28, HL vs CAD11 and HL vs Cad11-EC1, respectively), but did not bind proteins containing the EC1-2 domains of Cad-8 and MN-Cad (FIG. 28, HL vs CAD8 and HL vs MNCAD, respectively). Control hybridoma media did not bind any of the cadherin proteins (FIG. 28, Media vs CAD11, Media vs CAD8, and Media vs MNCAD).

25 Anti-Cad-11 antibodies from the Peptide 3 hybridomas also bound to cells expressing human Cad-11 protein (FIG. 29, see arrow), but not to non-Cad-11-expressing control cells that expressed Neos. This result confirmed the presence of

anti-Cad-11 antibodies in the hybridomas that recognize both Peptide 3 and Cad-11-expressing cells *in vitro*.

5 The relevant teachings of all patents, published applications and references cited herein are incorporated by reference in their entirety.

While this invention has been particularly shown and described with references to example embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without 10 departing from the scope of the invention encompassed by the appended claims.

INDICATIONS RELATING TO DEPOSITED MICROORGANISM OR OTHER BIOLOGICAL MATERIAL
(PCT/Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 15, line 17 and 21; page 41, line 9 and 11; page 54, line 12, 14, 16, and 19

B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet

Name of depositary institution

AMERICAN TYPE CULTURE COLLECTION

Address of depositary institution (*including postal code and country*)

American Type Culture Collection
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

Date of deposit: 08 January 2009

Accession Number: Not Available

C. ADDITIONAL INDICATIONS (*leave blank if not applicable*) This information is continued on an additional sheet

In respect of those designations for which a European patent is sought, the Applicant hereby informs the International Bureau that the Applicant wishes that, until the publication of the mention of the grant of a European patent or for 20 years from the date of filing if the application is refused or withdrawn or deemed to be withdrawn, the biological material deposited with the American Type Culture Collection under Accession No. shall be made available as provided in Rule 28(3) EPC only by the issue of a sample to an expert nominated by the requester (Rule 28(4) EPC).

D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (*if the indications are not for all designated States*)

E. SEPARATE FURNISHING OF INDICATIONS (*leave blank if not applicable*)

The indications listed below will be submitted to the International Bureau later (*specify the general nature of the indications e.g., "Accession Number of Deposit"*)

For receiving Office use only

This sheet was received with the International application

Authorized officer

For International Bureau use only

This sheet was received by the International Bureau on:

Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL
(Additional Sheet)**

C. ADDITIONAL INDICATIONS (Continued)

In respect of the designation of Australia in the subject PCT application, and in accordance with Regulation 3.25(3) of the Australian Patents Regulations, the Applicant hereby gives notice that the furnishing of a sample of the biological material deposited with the American Type Culture Collection under Accession No. N/A shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention and who is nominated in a request for the furnishing of a sample.

In respect of the designation of Canada in the subject PCT application, the Applicant hereby informs the International Bureau that the Applicant wishes that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the biological material deposited with the American Type Culture Collection under Accession No. N/A and referred to in the application to an independent expert nominated by the Commissioner.

CLAIMS

What is claimed is:

1. A Cadherin-11 antagonist that specifically binds an EC1 domain of a mammalian Cadherin-11 protein, wherein the Cadherin-11 antagonist inhibits aggregation of cells that express said mammalian Cadherin-11 protein.
2. The Cadherin-11 antagonist of Claim 1, wherein the Cadherin-11 antagonist binds SEQ ID NO:3.
3. The Cadherin-11 antagonist of Claim 1, wherein the Cadherin-11 antagonist is selected from the group consisting of an antibody, a fusion protein, a peptide, a peptidomimetic, a nucleic acid, and a small molecule.
4. An isolated antibody that specifically binds an EC1 domain of a mammalian Cadherin-11 protein, wherein the antibody inhibits aggregation of cells that express said mammalian Cadherin-11 protein.
5. The isolated antibody of Claim 4, wherein the antibody binds an epitope that is present in SEQ ID NO:3.
6. The isolated antibody of Claim 5, wherein the epitope does not include the amino acid sequence EEY.
7. The isolated antibody of Claim 4, wherein the antibody is selected from the group consisting of a monoclonal antibody, a polyclonal antibody, a humanized antibody, a chimeric antibody, a single chain antibody, and an antibody fragment.
8. The isolated antibody of Claim 7, wherein the antibody is an antibody fragment.

9. The isolated antibody of Claim 8, wherein the antibody fragment is selected from the group consisting of an Fab, an Fab', an F(ab')₂ and an scFv.
- 5 10. A fusion protein comprising at least a portion of a mammalian immunoglobulin protein and a portion of a human Cadherin-11 extracellular region that includes amino acids 54-90 of SEQ ID NO:2, wherein the portion of the human Cadherin-11 extracellular region does not include the entire human Cadherin-11 extracellular region consisting of amino acids 1-609 of SEQ ID NO:2.
- 10 11. The fusion protein of Claim 10, wherein the fusion protein comprises amino acids 1-150 of SEQ ID NO:2.
- 15 12. The fusion protein of Claim 10, wherein the fusion protein comprises amino acids 1-160 of SEQ ID NO:2.
13. The fusion protein of Claim 10, wherein the fusion protein comprises amino acids 1-259 of SEQ ID NO:2.
- 20 14. The fusion protein of Claim 10, wherein the fusion protein comprises amino acids 1-269 of SEQ ID NO:2.
15. The fusion protein of Claim 10, wherein the fusion protein lacks amino acids 25 1-53 of SEQ ID NO:2.
16. The fusion protein of Claim 10, wherein the mammalian immunoglobulin protein is a human IgG2 protein.
- 30 17. The fusion protein of Claim 16, wherein the fusion protein comprises the hinge-CH₂-CH₃ portion of human IgG₂.

18. The fusion protein of Claim 10, wherein the fusion protein is a monomer, a dimer or a tetramer.
19. A method of treating an inflammatory joint disorder in a mammalian subject in need thereof, comprising administering to the subject a therapeutically effective amount of a Cadherin-11 antagonist that specifically binds an EC1 domain of a mammalian Cadherin-11 protein, wherein the Cadherin-11 antagonist inhibits aggregation of cells that express said mammalian Cadherin-11 protein in one or more joints of said subject.
20. The method of Claim 19, wherein the Cadherin-11 antagonist binds SEQ ID NO:3.
21. The method of Claim 19, wherein the inflammatory joint disorder is selected from the group consisting of rheumatoid arthritis, osteoarthritis, psoriatic arthritis, Reiter's syndrome and ankylosing spondylitis.
22. The method of Claim 19, wherein the inflammatory joint disorder is rheumatoid arthritis.
23. The method of Claim 19, wherein the Cadherin-11 antagonist is an isolated antibody.
24. The method of Claim 19, wherein the Cadherin-11 antagonist is a fusion protein comprising at least a portion of a mammalian immunoglobulin protein and a portion of a human Cadherin-11 extracellular region that includes amino acids 54-90 of SEQ ID NO:2, wherein the portion of the human Cadherin-11 extracellular region does not include the entire human Cadherin-11 extracellular region consisting of amino acids 1-609 of SEQ ID NO:2.
25. The method of Claim 19, wherein the mammalian subject is a human.

26. The method of Claim 19, wherein the Cadherin-11 antagonist is administered systemically.
- 5 27. The method of Claim 19, wherein the Cadherin-11 antagonist is administered intravenously.
28. The method of Claim 19, wherein the Cadherin-11 antagonist is administered by direct injection into a joint.
- 10 29. The method of Claim 19, wherein the Cadherin-11 antagonist inhibits migration, adhesion, invasion into cartilage, or intercellular signaling of cells that express said mammalian Cadherin-11 protein in one or more joints of said subject.
- 15 30. The method of Claim 19, wherein the Cadherin-11 antagonist inhibits induction of expression or activity of an enzyme selected from the group consisting of a collagenase, a serine protease, and a matrix metalloproteinase in cells that express said mammalian Cadherin-11 protein in one or more joints of said subject.
- 20 31. The method of Claim 19, wherein the Cadherin-11 antagonist inhibits induction of expression or activity of a cytokine or growth factor selected from the group consisting of a IL-6, IL-8, RANKL and TRANCE in cells that express said mammalian Cadherin-11 protein in one or more joints of said subject.
- 25 32. The method of Claim 19, wherein the Cadherin-11 antagonist is administered in combination with a disease-modifying anti-rheumatic drug.
- 30 33. The method of Claim 32, wherein the disease-modifying anti-rheumatic drug is methotrexate.

34. The method of Claim 19, wherein the Cadherin-11 antagonist is administered in combination with an anti-inflammatory agent.

5 35. The method of Claim 34, wherein the anti-inflammatory agent is an NSAID or a steroid.

36. The method of Claim 34, wherein the anti-inflammatory agent is a disease modifying antirheumatic drug or a recombinant protein.

10 37. A pharmaceutical composition comprising a Cadherin-11 antagonist that specifically binds an EC1 domain of a mammalian Cadherin-11 protein and inhibits aggregation of cells that express said mammalian Cadherin-11, and a pharmaceutically-acceptable carrier.

15 38. The pharmaceutical composition of Claim 37, wherein the Cadherin-11 antagonist binds SEQ ID NO:3.

39. The pharmaceutical composition of Claim 37, wherein the Cadherin-11 antagonist is an isolated antibody.

20 40. The pharmaceutical composition of Claim 37, wherein the Cadherin-11 antagonist is a fusion protein comprising at least a portion of a mammalian immunoglobulin protein and a portion of a human Cadherin-11 extracellular region that includes amino acids 54-90 of SEQ ID NO:2, wherein the portion of the human Cadherin-11 extracellular region does not include the entire human Cadherin-11 extracellular region consisting of amino acids 1-609 of SEQ ID NO:2.

25 41. The pharmaceutical composition of Claim 37, further comprising a disease-modifying anti-rheumatic drug.

42. The pharmaceutical composition of Claim 41, wherein the disease-modifying anti-rheumatic drug is methotrexate.
43. The pharmaceutical composition of Claim 37, further comprising an anti-inflammatory agent.
44. The pharmaceutical composition of Claim 43, wherein the anti-inflammatory agent is an NSAID or a steroid.
- 10 45. The pharmaceutical composition of Claim 43, wherein the anti-inflammatory agent is a disease modifying antirheumatic drug or a recombinant protein.
46. The isolated antibody of Claim 4, wherein the antibody binds an epitope that is present in SEQ ID NO:10.
- 15 47. The isolated antibody of Claim 4, wherein the antibody binds an epitope that comprises SEQ ID NO:11.
48. The isolated antibody of Claim 4, wherein the antibody binds an epitope that is present in SEQ ID NO:12.
- 20 49. The method of Claim 23, wherein the isolated antibody binds an epitope that comprises SEQ ID NO:11.
- 25 50. The method of Claim 23, wherein the isolated antibody binds an epitope that is present in at least one sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:12, and SEQ ID NO:13.
- 30 51. The pharmaceutical composition of Claim 39, wherein the isolated antibody binds an epitope that comprises SEQ ID NO:11.

52. The pharmaceutical composition of Claim 39, wherein the isolated antibody binds an epitope that is present in at least one sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:12, and SEQ ID NO:13.
- 5 53. An isolated nucleic acid encoding the antibody of Claim 4.
54. The isolated nucleic acid of Claim 53, wherein said nucleic acid is present in a vector.
- 10 55. An isolated cell expressing the antibody of Claim 4.
56. A cell of hybridoma H1M1 (ATCC accession number _____).
57. A cell of hybridoma H14 (ATCC accession number _____).
- 15 58. An antibody produced by hybridoma H1M1 (ATCC accession number - _____).
59. An antibody produced by hybridoma H14 (ATCC accession number - _____).
- 20 60. Use of a Cadherin-11 antagonist that specifically binds an EC1 domain of a mammalian Cadherin-11 protein for the treatment of an inflammatory joint disorder in a mammalian subject in need thereof.
- 25 61. Use of a Cadherin-11 antagonist that specifically binds an EC1 domain of a mammalian Cadherin-11 protein in the manufacture of a medicament for the treatment of an inflammatory joint disorder in a mammalian subject in need thereof.




FIG. 1A




FIG. 1B

3/32

Human Cad-11	<u>G</u> <u>W</u> <u>W</u> <u>W</u> <u>N</u> <u>Q</u> <u>FF</u> <u>V</u> <u>I</u>	EE <u>Y</u> <u>T</u> <u>G</u> <u>P</u> <u>D</u> <u>P</u> <u>V</u> <u>L</u> <u>V</u> <u>G</u> <u>R</u> <u>L</u> <u>H</u> <u>S</u> <u>D</u> <u>I</u> <u>D</u> <u>S</u> <u>G</u> <u>D</u> <u>G</u> <u>N</u>	(SEQ ID NO:3)
Human Cad-8	<u>G</u> <u>W</u> <u>W</u> <u>W</u> <u>N</u> <u>Q</u> <u>M</u> <u>F</u> <u>V</u> <u>L</u>	EE <u>F</u> <u>S</u> <u>G</u> <u>P</u> <u>E</u> <u>P</u> <u>I</u> <u>L</u> <u>V</u> <u>G</u> <u>R</u> <u>L</u> <u>H</u> <u>T</u> <u>D</u> <u>L</u> <u>D</u> <u>P</u> <u>G</u> <u>S</u> <u>K</u> <u>K</u>	(SEQ ID NO:4)
Human MN-Cad	<u>S</u> <u>W</u> <u>W</u> <u>W</u> <u>N</u> <u>Q</u> <u>F</u> <u>F</u> <u>V</u> <u>L</u>	EE <u>Y</u> <u>T</u> <u>G</u> <u>T</u> <u>D</u> <u>P</u> <u>L</u> <u>Y</u> <u>V</u> <u>G</u> <u>K</u> <u>L</u> <u>H</u> <u>S</u> <u>D</u> <u>M</u> <u>D</u> <u>R</u> <u>G</u> <u>D</u> <u>G</u> <u>S</u>	(SEQ ID NO:5)

FIG. 2





FIG. 3

FIG. 4






FIG. 5

7/32





FIG. 6



ATGAGGAGAACTACTGTTTACAAGCCCCCTGGCTGGCATGGCCATGCCACAGGCCATGCCCTGGC
CCCAGAGGGGGCACCTGGCCCTCCATGGCACCCATGAGAAGGGCAAGGAGGGCAGG
TGCTACAGGGCTCCAAGCGTGGCTGGTGTGGTGTGGATAGGGAGTACACCGGGCTGA
CCCCGTGCTTGTGGCAGGGCTTCAATTGAGTATTGACTCTGGTGTGGATGGAAACATTAAATACATTCTCAGG
GGAAGGGAGCTGGAAACCAATTGGTGTGGATGGACAAATCAGGGAAACATTGCCACCAAGACGGTTGGATC
GAGAAGAGAGCCCCAGTACACGTTGGATGGCTCAGGGTGGAGTGGAGTGGCCACCCACCTGTGGC
CACCGTGGAAATTCAATTGTCAGGGTCCAGAGATCTGTGGAGTGGCCACCCACCTGTGGC
AGGACCCCTCAGTCTCCCTCTCCCCAAAACCCAAAGGACACCCCTGATGATCTCCAGAACCCCTGAGGTC
ACGTGCGTGGTGGACGTCAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGG
GGAGGGTGCATAATGCCAAGACAAAGGCCACGGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGG
TCCCTACCGGCTGGCACCCAGGACTGGCTGAACGGCAAGGAGTACAAGTGCAGGTCAACTGGTACGGC
TCCAGCCCCCATCGAGAAAACCATCTCCAAACCAAGGGCAGCCCCGAGAACCCACAGGGTGTACAC
CTGCCCCATCCCCGGAGGAGATGACCAAGAACCCAGGTCAAGCTGCCCTGGTCAAAGGGCTCTAC
CCAGGGACATGCCGTGGAGTGGAGGAGCAATGGCAGGGAGAACAAACTACAAGACCAACACCTCC
CATGCTGGACTCCGACGGCTCCCTACAGCAAGCTCAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGG
GGGGAAACGTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAAACCACACAGAAGAGGAGGAGGAGGAGG
CCTGTCTGGTAAATGAGTGCCACGGCTAGCTGG (SEQ ID NO:6)

FIG. 8

MKENYCLQAALVCLGMLCHSHAFAPERRGHLRPSFHGHHEKGKEGQVLQRSSKR
GWWWNQFFVIEEYTGDPVLVGRLLHSDIDSGDGNIKYILSGEGAGTIVDDKSGNI
HATKTLDREREERAQYTLMAQAVDRDTNRPLEPPSEFIVKVQ
RSVECPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFN
WYVDGMEVHNNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGL
PAPIEKTIKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTPPMLSDGSSFFLYSKLTVDKSRWQQGNVFSCSVMMHEALHNHY
TQKSLSLSPGKVPRLA (SEQ ID NO:7)

FIG. 9

hCad11 EC1/2-hIgG2

hCad11 EC1-hIgG2



FIG. 10

12/32





FIG. 11






FIG. 13B




FIG. 13A

1 agatgccgcg ggggcccgtc gcagccgcg ctgacttgt aatgggaccg ggactggggc
61 cgggactgac accgcagcgc ttgcctgcg ccagggactg gcccgcgaa gggtgcgtcc
121 accctcaagg gccccagaaa tcactgtgtt ttcaagctcag cggccctgtg acattccttc
181 gtgttgtcat ttgttgagtg accaatcaga tgggtggagt gtgttacaga aattggcagc
241 aagtatccaa tgggtgaaga agaagctaac tggggacgtg ggcagccctg acgtgatgag
301 ctcaccaggc agagacattc catccaaaga gaggtctgcg tgacgcgtcc gggaggccac
361 cctcagcaag accaccgtac agttggtgga aggggtgaca gctgcattct cctgtgccta
421 ccacgtaacc aaaaatgaag gagaactact gtttacaagc cgcctggtg tgcctggca
481 tgctgtgcca cagccatgcc tttgccccag agcgggggg gcacctgcgg ccctccctcc
541 atgggcacca tgagaagggc aaggagggc aggtgctaca gcgcctcaag cgtggctgg
601 tctggAACCA gttcttcgtt atagaggagt acaccggcc tgacccctgt cttgtggca
661 ggcttcattc agatattgac tctggtgatg ggaacattaa atacattctc tcaggggaag
721 gagctggAAC CATTTCGTG attgtatgaca aatcaggaa cattcatgcc accaagacgt
781 tggatcgaga agagagagcc cagtacacgt tgatggctca ggcgggtggac agggacacca
841 atcggccact ggagccaccg tcggattca ttgtcaaggt ccaggacatt aatgacaacc
901 ctccggagtt cctgcacgag acctatcatg ccaacgtgcc tgagaggc aatgtggaa
961 cgtcagtaat ccaggtgaca gcttcagatg cagatgaccc cacttatgg aatagcgc
1021 agtttagtgt a cgtatcctc gaaggacaac cctatTTTC ggtggaaagca cagacaggta
1081 tcatcagaac agccctaccc aacatggaca gggaggccaa ggaggagtac cacgtggta
1141 tccaggccaa ggacatgggt ggacatatgg gcggactctc agggacaacc aaagtgacga
1201 tcacactgac cgtatgtcaat gacaacccac caaagtTTTC gcagagcgtt taccagatgt
1261 ctgtgtcaga agcagccgtc cctggggagg aagttaggaag agtggaaagct aaagatccag
1321 acattggaga aaatggctt a gtcacataca atattgttga tggagatggt atggaaatcgt
1381 ttgaaatcac aacggactat gaaacacagg aggggggtgat aaagctgaaa aagcctgttag
1441 attttggaaac caaaagagcc tatacgTTGA aggttagaggc agccaaacgtg cacatcgacc
1501 cgaagtttat cagcaatggc ctttcaagg acactgtgac cgtcaagatc tcagtagaa
1561 atgtgtatgt gccccctatg ttcttggccc caagttacat ccacgaagtc caagaaaatg
1621 cagctgctgg caccgtggtt gggagagtgc atgccaaaga ccctgtatgtc gccaacagcc

16/32

1681 cgataaggta ttccatcgat cgtcacactg acctcgacag atttttact attaatccag
1741 aggatggttt tattaaaact acaaaaacctc tggatagaga gggaaacagcc tggctcaaca
1801 tcactgtctt tgcagcagaa atccacaatc ggcacatcagga agccaaagtc ccagtggcca
1861 ttagggtcct tgatgtcaac gataatgctc ccaagttgc tgccccttat gaaggttca
1921 tctgtgagag tgatcagacc aagccacttt ccaaccagcc aattgttaca attagtgcag
1981 atgacaagga tgacacggcc aatggaccaa gatttatctt cagectaccc cctgaaatca
2041 ttcacaatcc aaatttcaca gtcagagaca accgagataa cacagcagggc gtgtacgccc
2101 ggcgtggagg gttcagtcgg cagaagcagg acttgtacct tctgcccata gtgatcagcg
2161 atggcggcat cccgcccattg agtagcacca acaccctcac catcaaagtc tgcgggtgcg
2221 acgtgaacgg ggcactgctc tccatgcacg cagaggccta cattctgaac gccggcctga
2281 gcacaggcgc cctgatcgcc atcctcgcc tgcacat tctcctggtc attgttagtat
2341 tgggggtgac cctgagaagg caaaaagaaaag aaccactcat tgcgttttag gaagaagatg
2401 tccgtgagaa catcattact tatgtatgt aaggggggtgg ggaagaagac acagaagcct
2461 ttgatattgc caccctccag aatcctgtatg gtatcaatgg atttatcccc cgcaaagaca
2521 tcaaaccctga gtatcagtac atgcctagac ctgggctccg gccagcgc aacagcgtgg
2581 atgtcgatga cttcatcaac acgagaatac aggaggcaga caatgacccc acggctcctc
2641 cttatgactc cattcaaatac tacggttatg aaggcagggg ctcagtgcc gggccctga
2701 gctccctaga gtcggccacc acagattcag acttgacta tgattatcta cagaactggg
2761 gacctcggtt taagaaacta gcagattgt atgggtccaa agacacttt gatgacgatt
2821 ctttacaata acgatacataa tttggccta agaactgtgt ctggcggttca caagaatcta
2881 gaagatgtgt aaacaggtat tttttaaat caagggaaagg ctcatttaaa acaggcaaag
2941 ttttacagag aggatacatt taataaaaact gcgaggacat caaagtggta aatactgtga
3001 aatacccttt ctcacaaaaa ggcaaataattt gaagttgtt atcaacttcg ctgggggggg
3061 aaaacacttg gcatacataa tatttaatgt aaggagaagt ctaacgctga actgacaatg
3121 aaggggaaatt gtttatgtgt tatgaacatc caagtcttc ttctttttt agttgtcaaa
3181 gaagcttcca caaaattaga aaggacaaca gttctgagct gtaatttcgc cttaaactct
3241 ggacactcta tatgtatgtc atttttaaac ttgaaatata taatattcag ccagcttaaa
3301 cccatataat gtatgtacaa tacaatgtac aattatgtct cttgagcattc aatcttggta
3361 ctgctgattc ttgttaatct ttttgcgttactt actttcatct taaactaata cgtgccat

FIG. 14B

17/32

3421 ataactgtct tgtttcagtg agagacgcc tatttctatg tcattttaa tgtatctatt
3481 tgtacaattt taaagttctt attttagtat acgtataaat atcagtattc tgacatgtaa
3541 gaaaatgtta cggcatcaca ctatatttt atgaacattt tactgttgct ttaatatgag
3601 cttcaatata agaagcaatc ttgaaataa aaaaagattt tttttaaaa aaaa (SEQ
ID NO:1)

FIG. 14C

1 mkenyclqaa lvclgmlchs hafaperrgh lrpsfhghhe kgkegqvqlqr skrgvwvnqf
61 fviieeytgpd pvlvgrlhds idsgdgniky ilsgegagti fviddksgni hatkldree
121 raqytlmaqa vdrdtnrple ppsefivkvq dindnppbefl hetyhanvpe rsnvgtsviq
181 vtasdaddpt ygnsaklvys ilegqpyfsv eaqtgiirta lpnmdreake eyhvvqakd
241 mgghmgglsg ttkvtiltd vndnppkfpq svyqmsvsea avpgeevgrv kakdpdigen
301 glvtnivdg dgmefiitt dyetqegvik lkkpvdfetk rayslkveaa nvhidpkfis
361 ngpfkdtvtv kisvedadep pmflapsyih evqenaaagt vvgrvhakdp daanspirys
421 idrhtdldrf ftinpedgfi kttkpldree tawlnitvfa aeihrhqeа kpvairvld
481 vndnapkfaa pyegficesd qtkplsnqpi vtisaddkdd tangprfifs lppeihnpn
541 ftvrdnrdnt agvyarrggf srqkqdlyll pivisdggip pmsstntlti kvcgcvnnga
601 llscnaeayi lnaglstgal iailacivil lvivvlfvtl rrqkkepliv feeedvreni
661 ityddeggge edteafdiat lqnpdgingf iprkdkpey qymprpglrlp apnsvdvddf
721 intriqeadn dptappydsi qiygyegrsgs vagslssles attdsldyd ylqnwgprfk
781 kladlygskd tfddds (SEQ ID NO: 2)













FIG. 16

FIG. 17C**FIG. 17B****FIG. 17A**

H14
Cad11+ cells

FIG. 17D

H14
Cad11- cells

FIG. 17F**FIG. 17E****FIG. 17I****FIG. 17H****FIG. 17G**

H1M1
Cad11+ cells

21/32




FIG. 18A

FIG. 18B

22/32

FIG. 19A





FIG. 19B



23/32





FIG. 20

24/32







FIG. 22

26/32



27/32




FIG. 25




FIG. 26




FIG. 27





FIG. 28



32/32




FIG. 29