



(11) **EP 3 498 308 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
27.11.2024 Bulletin 2024/48

(51) International Patent Classification (IPC):
A61K 51/04 ^(2006.01) **A61P 35/04** ^(2006.01)
A61P 35/00 ^(2006.01) **A61K 103/40** ^(2006.01)
A61K 103/00 ^(2006.01)

(21) Application number: **18212268.9**

(52) Cooperative Patent Classification (CPC):
A61K 51/0402; A61K 51/0497; A61P 35/00;
A61P 35/04

(22) Date of filing: **13.12.2018**

(54) **COMPLEX COMPRISING A PSMA-TARGETING COMPOUND LINKED TO A LEAD OR THORIUM RADIONUCLIDE**

KOMPLEX BEINHALTEND EINE AUF PSMA GERICHTETE VERBINDUNG, DIE AN EIN BLEI- ODER THORIUM-RADIONUKLID GEBUNDEN IST

COMPLEXE COMPRENANT UN COMPOSÉ CIBLANT PSMA LIÉ À UN RADIONUCLÉIDE DE PLOMB OU DE THORIUM

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **13.12.2017 EP 17206887**

(43) Date of publication of application:
19.06.2019 Bulletin 2019/25

(73) Proprietor: **Sciencons AS**
0484 Oslo (NO)

(72) Inventor: **LARSEN, Roy Hartvig**
N-0598 Oslo (NO)

(74) Representative: **Mewburn Ellis LLP**
Aurora Building
Counterslip
Bristol BS1 6BX (GB)

(56) References cited:

- **CLEMENS KRATOCHWIL ET AL: "Targeted [alpha]-Therapy of Metastatic Castration-Resistant Prostate Cancer with 225 Ac-PSMA-617: Dosimetry Estimate and Empiric Dose Finding", THE JOURNAL OF NUCLEAR MEDICINE, vol. 58, no. 10, 13 April 2017 (2017-04-13), US, pages 1624 - 1631, XP055565538, ISSN: 0161-5505, DOI: 10.2967/jnumed.117.191395**
- **MIKE SATHEKGE ET AL: "213Bi-PSMA-617 targeted alpha-radionuclide therapy in metastatic castration-resistant prostate cancer", EUROPEAN JOURNAL OF NUCLEAR MEDICINE AND MOLECULAR IMAGING, vol. 44, no. 6, 2 March 2017 (2017-03-02), DE, pages 1099 - 1100, XP055565545, ISSN: 1619-7070, DOI: 10.1007/s00259-017-3657-9**
- **JAMES M. KELLY ET AL: "Assessment of PSMA targeting ligands bearing novel chelates with application to theranostics: Stability and complexation kinetics of 68 Ga 3+ , 111 In 3+ , 177 Lu 3+ and 225 Ac 3+", NUCLEAR MEDICINE AND BIOLOGY., vol. 55, 1 December 2017 (2017-12-01), US, pages 38 - 46, XP055565558, ISSN: 0969-8051, DOI: 10.1016/j.nucmedbio.2017.10.001**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

FIELD OF THE INVENTION

[0001] The present invention relates to complexes comprising a PSMA targeting compound linked to a radionuclide ^{212}Pb . These compounds, and pharmaceutical compositions comprising them, can be used for medical applications. These applications include the treatment of prostate cancer, and the complexes allow for dual targeting of cancers. Peptide and peptidomimetic PSMA targeting urea derivatives conjugated to chelators for complexing ^{212}Pb for radio-therapeutic use. These can be used in single and dual targeting.

BACKGROUND OF THE INVENTION

[0002] Prostate cancer is among the most frequent causes of cancer related mortality in men. There is a great demand for new and effective treatment, especially in hormone refractory late stage disease. Skeletal metastases are a frequent problem in late stage disease and therefore the alpha-particle emitter ^{223}Ra (Xofigo) was introduced as a bone specific therapy for late stage prostate cancer patients with skeletal metastases.

[0003] Although, as a bone-seeker, ^{223}Ra shows significant clinical benefit for patients its activity is limited to the bone metastases and is not targeting soft tissue metastases.

[0004] Several carrier molecules for radioligand targeting of prostate specific membrane antigen (PSMA) exists. Lutetium-177 labeled PSMA-617 (^{177}Lu -PSMA-617) is the compound in most advanced clinical development stage for use in radionuclide therapy.

[0005] This molecule works in a suitable manner and give relevant tumor to normal tissue ratios for longer lived (i.e. a half-life of a few days) radionuclides, including ^{177}Lu and ^{225}Ac , but at early times points (typically a few hours after injection) shows high uptake in kidneys. With shorter lived radionuclides like ^{212}Pb (half-life of 10.6 hours), the initial kidney uptake represents a potential toxicity problem.

[0006] It is therefore advantageous to use a PSMA-ligand with less kidney uptake, but this should not compromise the tumor uptake. The PSMA ligand molecules are made up of (1) a PSMA-binding region, (2) a linker region and (3) a chelator, whereby the linker region connects the (1) and (3). The linker region also is used to adjust molecular size and polarity etc to affect the in vivo distribution properties. The PSMA-binding region (motif) used in PSMA-617 is a structure that can be found in several molecules of this class, developed by several different inventors and researchers, including PSMA-11 and PSMA I&T as well as ^{131}I and ^{211}At labelled PSMA binding ligands.

[0007] Kratochwil et al. (Journal of nuclear medicine, vol. 58, no. 10, 13 April 2017, pages 1624-1631) describes ^{225}Ac -PSMA-617 for use in the treatment of metastatic castration-resistant prostate cancer.

[0008] Sathekge et al. (European Journal of Nuclear medicine and molecular imaging, vol. 44, no. 6, 2 March 2017, pages 1099-1100) describes ^{213}Bi -PSMA-617 for use in the treatment of metastatic castration-resistant prostate cancer.

[0009] Kelly et al. (Nuclear medicine and biology, vol. 55, 1 December 2017, pages 38-46) describes assessment of PSMA targeting ligands with DOA and complexation of e.g. $^{68}\text{Ga}^{3+}$.


[0010] New compounds that contain a PSMA region are warranted because at current all ligands in testing have challenges, including a relatively low radiobiological effectiveness (RBE) and suboptimal biodistribution.

[0011] There is also a need for an improved alpha emitter that can target both the bone metastases and the soft tissue metastases.

[0012] Described herein are compounds that address these challenges.

SUMMARY OF THE INVENTION

[0013] A first aspect of the invention is a complex comprising a compound complexed with ^{212}Pb wherein the compound comprises a PSMA unit and chelating unit, wherein the PSMA unit is carbon-backbone linked to the chelating unit, and wherein the compound is of the formula:



p-SCN-Bn-TCMC-PSMA ligand 1

[0014] A second aspect of the invention is a pharmaceutical composition comprising the complex according to the first aspect, and a diluent, carrier, surfactant, and/or excipient.

[0015] In one embodiment, the pharmaceutical composition further comprises ^{224}Ra .

[0016] In one embodiment, the activity ratio (MBq) between ^{212}Pb to ^{224}Ra is between 0.5 and 2, such as 0.8 - 1.5, or such as 0.8 - 1.3, or preferably such as 0.9 - 1.15.

[0017] In one embodiment, the pharmaceutical composition is dosaged with a radioactivity of 100 kBq to 100 MBq per dose.

[0018] In one embodiment, the pharmaceutical composition is dosaged with a radioactivity of 1 MBq to 500 MBq per patient.

[0019] In one embodiment, the pharmaceutical composition is a solution having a volume of 500 μL to 100 mL.

[0020] A third aspect of the invention is the pharmaceutical composition according to the first aspect for use as a medicament.

[0021] A fourth aspect of the invention is the pharmaceutical composition according to the first aspect for use in the treatment of PSMA-expressing disease including soft tissue- and skeletal disease; wherein the skeletal disease is selected from the group consisting of skeletal metastases from cancers to the breast, prostate, kidneys, lung, bone, or multiple myeloma.

[0022] A fifth aspect of the invention is the pharmaceutical composition according to the first aspect for use as a medicament in the treatment of cancer.

[0023] In one embodiment, the cancer is prostate cancer.


[0024] In one embodiment, the cancer is PSMA-expressing prostate cancer.

[0025] In one embodiment, the cancer is advanced metastatic prostate cancer.

[0026] In one embodiment of the third, fourth, or fifth aspect, the pharmaceutical composition is administered at a dose in the range 50-150 kBq per kg of bodyweight.

BRIEF DESCRIPTION

[0027] Described herein are complexes comprising a compound complexed with ^{212}Pb , wherein the compound comprises a PSMA unit and chelating unit, wherein the PSMA unit is carbon-backbone linked to the chelating unit, and wherein the compound is of the formula:



p-SCN-Bn-TCMC-PSMA ligand 1.

[0028] In one embodiment is the linker TCMC or a TCMC derivative.

[0029] Also described herein is a radiopharmaceutical composition for use as a medicament.

[0030] Also described herein is a radiopharmaceutical composition for use in the treatment of soft tissue and or skeletal

disease.

[0031] In one embodiment the skeletal disease is selected from the group consisting of skeletal metastases from cancers to the breast, prostate, kidneys, lung, bone, or multiple myeloma, or non-cancerous diseases causing undesired calcification including ankylosing spondylitis.

FIGURES

[0032]

Figure 1 shows biodistribution of ^{212}Pb 2 hours after injecting ^{212}Pb -labeled p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 respectively, in the presence of ^{224}Ra .

Figure 2 shows survival of mice with C4-2 PSMA positive xenograft after treatment with saline, 52 MBq ^{177}Lu -OSMA-617 and 0.32 MBq ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1.

DETAILED DESCRIPTION

[0033] The references to the methods of treatment in this description are to be interpreted as references to compounds, complexes, pharmaceutical compositions, and medicaments for use in those methods.

Some abbreviations used:

[0034] Peptide mimetic- also termed peptidomimetic, is a small protein-like chain designed to mimic a peptide. They typically arise either from modification of an existing peptide, or by designing similar systems that mimic peptides, such as peptoids and β -peptides. Irrespective of the approach, the altered chemical structure is designed to advantageously adjust the molecular properties such as, stability or biological activity. This can have a role in the development of drug-like compounds from existing peptides. These modifications involve changes to the peptide that will not occur naturally (such as altered backbones and the incorporation of nonnatural amino acids). Based on their similarity with the precursor peptide, peptidomimetics can be grouped into four classes (A - D) where A features the most and D the least similarities. Classes A and B involve peptide-like scaffolds, while classes C and D include small molecules.

PSMA- Prostate-specific membrane antigen. Synonyms PSMA, Prostate Specific Cancer Antigen, PSM, FGCP, FOLH, GCP2, mGCP, GCP II, NAALAD1, NAALadase, FOLH1, Glutamate carboxypeptidase 2, Glutamate carboxypeptidase II, Membrane glutamate carboxypeptidase, N-acetylated-alpha-linked acidic dipeptidase I, Pteroylpolygamma-glutamate carboxypeptidase, Folylpoly-gamma-glutamate carboxypeptidase, Folate hydrolase 1, Prostate-specific membrane antigen, Cell growth-inhibiting protein 27

DOTMP - 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetra(methylene phosphonic acid)

EDTMP - ethylenediamine tetra(methylene phosphonic acid)

EDTA - ethylenediaminetetraacetic acid

p-SCN-Bn-DOTA - 2-(4-isothiocyanatobenzyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid

DOTA- 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid and also used for benzyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (e.g. conjugated to monoclonal antibody)

p-SCN-Bn-TCMC - 2-(4-isothiocyanatobenzyl)-1, 4, 7, 10-tetraaza-1, 4, 7, 10-tetra-(2-carbamonyl methyl)-cyclododecane

TCMC - 1, 4, 7, 10-tetraaza-1, 4, 7, 10-tetra-(2-carbamonyl methyl)-cyclododecane and also used for benzyl-1, 4, 7, 10-tetraaza-1, 4, 7, 10-tetra-(2-carbamonyl methyl)-cyclododecane (e.g. conjugated to monoclonal antibody)

mAb -monoclonal antibody.

HOPO - Me-3,2-HOPO, octadentate hydroxypyridinone for complexing ^{227}Th , 4-((4-(3-(bis(2-(3-hydroxy-1-methyl-2-oxo-1,2-dihydropyridine-4-carboxamido)ethyl)amino)-2-((bis(2-(3-hydroxy-1-methyl-2-oxo-1,2-dihydropyridine-

4-carboxamido)ethyl)amino)methyl)propyl)phenyl)amino)-4-oxobutanoic acid and derivatives.

Ligand 1 - p-SCN-Bn-TCMC-PSMA ligand 1, p-SCN-Bn-TCMC-PSMA etc.

Ligand 2 - p-SCN-Bn-DOTA-PSMA ligand 2, p-SCN-Bn-DOTA-PSMA etc

[0035] The same abbreviations are in the following used for acids, salts or partly or fully dissociated versions of the chelators.


[0036] Described herein are single and dual targeting property solutions based on a small molecular urea derivative as a prostate cancer cellular targeting agent for carrying ^{212}Pb .

[0037] This may be used with purified ^{212}Pb or in a dual targeting solution whereby ^{224}Ra will act as a skeletal treatment and ^{212}Pb -urea-derivative will act as a systemic therapy against cells expressing PSMA antigen which is associated mainly with advanced metastatic prostate cancer, and to some extent also other types of cancer.

[0038] It is known in the field that urea based compounds conjugated to a chelator group can facilitate the targeting of radionuclides to PSMA expressing cells. Radionuclides that have been evaluated for radiotherapeutic purposes with PSMA targeting includes ^{177}Lu , ^{211}At , ^{213}Bi , and ^{225}Ac .

[0039] The invention is in the field of radiolabeled therapy agents. Radiolabelled derivatives of urea based prostate-specific membrane antigen (PSMA) inhibitors are described herein.

[0040] Thus, described herein is a complex comprising a compound that is a PSMA unit and chelating unit, wherein the PSMA unit is carbon-backbone linked to the chelating unit, and wherein the compound is of the formula:



p-SCN-Bn-TCMC-PSMA ligand 1 linked to ^{212}Pb .

Linker

[0041] Also described herein but not according to the invention is a complex wherein the compound X is linked to a radionuclide, such as ^{177}Lu , ^{213}Bi , ^{225}Ac , ^{212}Pb or ^{227}Th , by a chelating moiety Z.

[0042] In one embodiment is the radionuclide ^{212}Pb .

[0043] Also described herein is the radionuclide ^{227}Th .

[0044] Also described herein is the radionuclide ^{177}Lu .

[0045] Also described herein is the radionuclide ^{213}Bi or ^{212}Bi .

[0046] Also described herein is the radionuclide ^{225}Ac .

[0047] It is to be understood that the complexing agent, or linker, or chelating moiety Z, according to the disclosure may also cover derivatives of the above-mentioned compounds (such as derivatives of EDTMP, DOTA, for example p-SCN-Bn-DOTA and TCMC, for example p-SCN-Bn-TCMC). It is of course to be understood that such derivatives must maintain the capability to complex ^{212}Pb with a higher stability constant than to ^{224}Ra .

[0048] The chelating moiety Z may be selected from the group consisting of acyclic chelators, cyclic chelators, cryptands, crown ethers, porphyrins or cyclic or noncyclic polyphosphonates, DOTMP, EDTMP and bisphosphonate derivatives, DOTA, a DOTA derivative such as p-SCN-Bn-DOTA, pamidronate conjugated to DOTA, TCMC, a TCMC derivative such as p-SCN-Bz-TCMC, pamidronate conjugated to TCMC, antibody-conjugated-DOTA, antibody-conjugated-TCMC, HBED-CC, NOTA, NODAGA, TRAP, NOPO, PCTA, DFO, DTPA, CHX-DTPA, AAZTA, DEDPA, and oxo-Do3A.

[0049] Also described herein but not according to the invention is the linker DOTA or a DOTA derivative, such as p-SCN-Bn-DOTA. Also described herein is the linker TCMC or a TCMC derivative, such as p-SCN-Bn-TCMC. The complexing agent may be linked via the carbon backbone allowing all "binding arms" of the chelator molecule interact with the metal. Alternatively, one of the arms may be used as a linker.

[0050] Also described herein but not according to the invention are suitable chelators including DOTA derivatives such

as p-isothiocyanatobenzyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (p-SCN-Bz-DOTA) and DOTA-NHS-ester.

[0051] Thus, for p-SCN-Bn-DOTA or p-SCN-Bn-TCMC can the complexing agent be linked via the carbon backbone (C-backbone) to the rest of the compound.

[0052] In a special embodiment an urea derivative based PSMA targeting complex is labeled with ^{212}Pb or ^{227}Th in a mixture of bone-seeking ^{224}Ra or ^{223}Ra , respectively, for dual targeting by (1) PSMA cellular targeting and (2) targeting of skeletal metastases related bone synthesis by radium cations.

[0053] Thus, described herein is a PSMA targeting urea derivative comprising a TCMC group, such as p-SCN-Bn-TCMC, for chelating ^{212}Pb .

[0054] Also described herein but not according to the invention is a PSMA targeting urea derivative comprising HOPO for chelating ^{227}Th .

[0055] Also described herein but not according to the invention is a PSMA targeting urea derivative comprising DOTA, such as p-SCN-Bn-TCMC, labeled with either ^{212}Pb or ^{227}Th .

[0056] In an embodiment, the complexing agent does not complex ^{224}Ra or substantially complex ^{224}Ra in the pharmaceutical composition.

[0057] In a further embodiment, the complexing agent complexes with a higher stability constant to ^{212}Pb than to ^{224}Ra .

[0058] In an embodiment, the stability constant for ^{212}Pb is at least twice the affinity for ^{224}Ra , such as at least four times higher, such as at least 8 times higher or such as at least 10 times higher.

[0059] As described herein but not according to the invention the complexing agent may be selected from the group consisting of, ligand-conjugated-DOTA, such as ligand-conjugated-p-SCN-Bn-DOTA, or ligand-conjugated-TCMC, such as ligand-conjugated-p-SCN-Bn-TCMC.

[0060] The ligand may be an antibody or polypeptide.

[0061] In a further embodiment, the amount of ^{224}Ra and ^{212}Pb is in radioactive equilibrium.

[0062] In yet a further embodiment, the activity ratio (in MBq) between ^{212}Pb to ^{224}Ra is between 0.5 and 2, such as 0.8 - 1.5, or such as 0.8 - 1.3, or preferably such as 0.9 - 1.15.

[0063] In the present context, the term "radioactive equilibrium" relates to the ratio in MBq between two radionuclides being the same or substantially the same over time. The term "activity ratio" e.g. between ^{212}Pb and ^{224}Ra relates to the ratio of MBq of ^{212}Pb to ^{224}Ra . In figure 5 is a table (table 2) showing the development of this activity ratio over time. It can be seen that after two days a radioactive equilibrium of 1.1 has been established for the activity ratio between ^{212}Pb to ^{224}Ra (7.3 divided by 6.8). Thus, in figure 5, it can also be seen that the radioactive equilibrium between ^{212}Pb and ^{224}Ra is reached after about 2 days.

[0064] In the present context, the terms "complexing agent", "scavenger", "linker", "chelating moiety Z", and "chelating agent" are used interchangeably. The terms relate to agents capable of forming complexes with ^{212}Pb , preferably by chelation and with a significant strength as measured in test systems while radium is not significantly affected by the presence of the complex as measured in the test systems.

[0065] Test systems include *in vivo* biodistribution and *in vitro* cation exchanger or size retention and centrifuge concentration cartridge for chelate-antibody binding of radionuclide. Alternatively, thin layer chromatography may be used as a test system.

[0066] In the present context "scavenging" (or complexing) is defined as at least 50% bound according to thin layer chromatography (TLC), centrifuge concentration separation or biodistribution profiles.

[0067] This means, as an example, at least 50% less blood uptake of ^{212}Pb with a small molecular chelator. With an antibody-conjugated chelator, where blood uptake is not a reliable indicator, at least 50% bound according to TLC analyses.

[0068] In one embodiment is at least 60% bound.

[0069] In another embodiment is at least 70% bound.

[0070] In another embodiment is at least 80% bound.

[0071] In another embodiment is at least 85% bound.

[0072] In another embodiment is at least 90% bound.

[0073] The compound or compounds may also be capable of complexing more radionuclides than ^{212}Pb .

[0074] In one embodiment, the compound and/or complex is at a concentration of 1 ng/mL to 1 g/mL.

[0075] In another embodiment, the compound and/or complex is at a concentration of 100 ng to 10 mg/mL.

[0076] The complex can comprise one, two, three, four, five or more compounds.

[0077] In one embodiment is the solution in a volume of 100 μL to 1000 mL, such as 500 μL to 100 mL, 1 mL to 10 mL.

[0078] In one embodiment is the radioactivity of the solution 1 kBq to 1 GBq, such as 10 kBq to 100 MBq, such as 100 kBq to 10 MBq.

[0079] In another embodiment is the radioactivity of the solution 100 kBq to 100 MBq.

[0080] Described herein but not according to the invention is a complexing agent conjugated to a compound selected from the group consisting of peptides and peptide mimetic urea derivatives with affinity for PSMA.

[0081] In some instances, the complexing agent is the chelator TCMC, such as p-SCN-Bn-TCMC, or DOTA, such as p-SCN-Bn-DOTA, conjugated to a compound selected from the group consisting of a monoclonal antibody, a polyclonal antibody, an antibody fragment, a synthetic protein, a peptide, a hormone or hormone derivative or a vitamin or a vitamin derivative.

[0082] For dosing purposes, a pure ^{212}Pb solution may be used. Alternatively, ^{224}Ra in mixture with ^{212}Pb may be used, the latter for dual targeting purposes, i.e., ^{224}Ra targets bone metastases and ^{212}Pb targets systemic cancer by its urea derivative carrier. If ^{224}Ra comprising solutions are administered they may have been stored for some time, e.g., 1 day or more preferably at least two days, such as 1-2 days or 1-3 days, to reach equilibrium between ^{224}Ra and $^{212}\text{Pb}/^{212}\text{Bi}$. This will ensure ^{212}Pb to ^{224}Ra activity ratios between 0.83 and 1.14. This can, e.g., be accomplished by the manufacturer by simply retaining the product for a day or so before shipment.

[0083] Alternatively, ^{212}Pb may be added to ^{224}Ra solutions to obtain a specific radionuclide ratio. E.g. if the soft tissue tumor burden is much higher than the skeletal tumor burden, a pure ^{212}Pb preparation or a solution with a high ^{212}Pb to ^{224}Ra ratio may be used.

[0084] The non-overlapping side effect profile of cationic radium and alpha emitters conjugated to PSMA binding urea-derivatives makes a mixture of ^{224}Ra cation and ^{212}Pb -PSMA targeting agent for dual targeting particularly attractive. This is because more modest dosing of each compound can be used to produce antitumor activity, since, at least for the skeletal component, the two different compounds will target the lesions independently.

[0085] It is important to keep radium as mainly un-complexed, or weakly complexed cation as this ensures a maximum uptake in bone and bone metastases and also ensures favourable excretion of eliminated product mainly through the intestines.

[0086] By adding complexing agent to a solution of radium the radioactive daughter can be made bone- or tumor-seeking and increase the therapeutic potential of the radium solution instead of being a health hazard. It should be a complexing agent that does not negatively affect the bone-seeking properties of radium, though. For example, can a TCMC-labeled urea derivative scavenge ^{212}Pb produced in the radium solution during transport and storage between the production site and the hospitals whereby the product is going to be administered.

[0087] Although it is possible to reduce the susceptibility by adding radiolytic inhibitors, tumor targeting peptides and peptide mimetics are often more susceptible to radiolysis and should probably be supplied in a kit format whereby they are added a few hours to a few minutes before administration of ^{224}Ra solutions with relatively long shelf-lives.

[0088] It is known in the field that calixarenes and EDTA to some extent can complex radium and also complex lead and bismuth. However, in the current work we found chelators that would leave radium mainly uncomplexed or weakly complexed, as determined by *in vivo* biodistribution measurements, while being able to rapidly and with relevant stability, complex the longest living daughter ^{212}Pb . The selective complexation can be used to make at least lead bone- or tumor-seeking while maintaining the favourable properties of radium in terms of treating sclerotic diseases, like skeletal metastases. The ^{212}Pb complex that targets bone or tumor cells generates the alpha emitter ^{212}Bi from the decay of ^{212}Pb . Thus, the beta emitter ^{212}Pb is used as an indirect alpha source for irradiating the targeted cells or tissue. Other potential chelates which could be suitable for ^{224}Ra daughter nuclide scavenging besides TCMC and DOTA includes but are not limited to porphyrins, DTPA and DTPA derivatives and also carboxyl linked DOTA.

[0089] Lead-212 is by far the longest living of the progenies from ^{224}Ra and this is the most important to complex, as it is an *in vivo* generator for the short-lived alpha-emitter ^{212}Bi . If a ^{212}Pb -chelate is taken up in bone or in tumor cells ^{212}Bi will also likely be retained in the target. In a ^{224}Ra solution in equilibrium with progenies there will be more than 10 times of ^{212}Pb vs. ^{212}Bi atoms. Thus, the amount of radiation generated from the ^{212}Bi atoms in these solutions are modest and probably not of a toxicologically importance compared with ^{224}Ra and ^{212}Bi decay series. The amount of ^{212}Bi is comparable to that of the ^{211}Pb which indirectly produces an alpha particle in the ^{223}Ra series and this has not been of a significant problem for the registration and clinical use of ^{223}Ra in equilibrium with progenies.

[0090] If, however, a high degree of chelation also of ^{212}Bi in an injectate should be needed, it may at least in some instances be necessary to add a stabilizing agent like NaI or HI since bismuth in aqueous solutions tends to exist in a state less suitable for chelation.

[0091] When comparing with current approved alpha-pharmaceutical for treatment of skeletal metastases, i.e., ^{223}Ra , the novel solutions described herein could give, in one of the embodiment, a product with improved properties for treatment of skeletal metastases since the daughter nuclide can be made targetable to circulating cancer cells and, to some extent, also soft tissue metastases. This may prevent recurrence from cancer recolonization of the skeleton due to CTC's.

[0092] Another aspect is that the shorter half-life of ^{224}Ra vs. ^{223}Ra may actually be of some benefit as the radium is embedded in the bone matrix. Because of the high density of the bone mineral the range of alpha-particles is strongly reduced in bone vs. soft tissues. Especially in rapid mineralizing areas like osseous cancer metastases, the embedment process may be of significance when using a volume-seeking alpha-pharmaceutical.

[0093] Therefore, ^{224}Ra could improve the tumor dose since, on average, it will be less embedded at the time of decay.

[0094] Diseases by which the novel ^{212}Pb solutions with or without ^{224}Ra may be used include but are not limited to

primary and metastatic cancers, autoimmune diseases and arteriosclerosis. The product may be administered intravenously or locally, including intraperitoneally, or in limb perfusion settings.

[0095] The chelators used in the novel solutions may be acyclic as well as cyclic chelators and cryptands, crown ethers, porphyrins or cyclic- or noncyclic polyphosphonates including DOTMP and EDTMP. Also a bisphosphonate, e.g., pamidronate, conjugated to DOTA, TCMC or similar may be used as scavenger in the ^{224}Ra solution.


[0096] One may argue that the amount of ^{212}Pb in therapeutic ^{224}Ra solution may be moderate to modest (i.e., at equilibrium about 1.1 times that of ^{224}Ra). If one assumes similar dosing of ^{224}Ra as is done with ^{223}Ra in patients but correct for the half-life difference, roughly 150 kBq per kg of bodyweight would be the administered dose.

[0097] At equilibrium this would translate into a ^{212}Pb -antibody conjugate dosage of 11.5 MBq in 5 litres of blood in a 70-kg patient (if ^{212}Pb is quantitatively chelated). The number of circulating tumor cells is typically less than 10 cells per ml, thus in 5 l blood there are less than 50 000 tumor cell in total. If only 1 in 100 000 of the injected ^{212}Pb -antibody conjugate molecules binds to the tumor cells this would mean at least 0.0023 Bq per cell, equivalent to approximately 127 ^{212}Pb atoms bound per cell, which would be highly destructive as it has been reported that a mean of 25 cell bound ^{212}Pb per cell would kill 90% of a cell population.

Compounds

[0098] Also described herein but not according to the invention is a compound X linked to a chelating moiety Z is defined by the formula I:

Formula I



or a pharmaceutically acceptable salt thereof,

wherein

W is a PSMA-targeting ligand;

A⁴ is a bond or a divalent linking moiety comprising 1 to 10 carbon atoms in a chain, a ring, or a combination thereof, wherein at least one carbon atom is optionally replaced with O, -NR³-, or -C(O)-;

G is C=O, C=S, C-NH₂, or C-NR³;

R¹ is hydrogen or a carboxylic acid protecting group;

R³ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, heterocycloalkyl, aryl, alkylaryl, and heteroaryl.

R¹¹, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each independently hydrogen, alkyl, alkoxyl, or

R¹⁷ and R¹⁸ are each independently hydrogen, alkyl, aryl, or alkylaryl;

R¹⁹ is selected from the group consisting of alkyl, alkoxyl, halide, haloalkyl, and

CN;

m is an integer from 1 to 6; and

o is an integer from 0 to 4, wherein when o is greater than 1, each R¹⁹ is the same or different.

[0099] Also described herein but not according to the invention is a complex, or a pharmaceutically acceptable salt thereof, wherein

A⁴ is a bond, (CH₂)_n, -HC(O)-, -(OCH₂CH₂)_n-, -(HCH₂CH₂)_n-, -H(CO)CH₂-, -HC(O)CH₂(OCH₂CH₂)_n-, or -HC(O)CH₂(HCH₂CH₂)_n-; and


L is a bond, (CH₂)_n, -(OCH₂CH₂)_n-, -(HCH₂CH₂)_n-, or -C(O)(CH₂)_n-;

wherein n is independently 1, 2, or 3.

[0100] Also described herein but not according to the invention is a complex, or a pharmaceutically acceptable salt thereof, wherein


A⁴ is a bond,
 -(OCH₂CH₂)_n-, or -HC(=O)CH₂(OCH₂CH₂)_n-; and
 L is a bond, or -(OCH₂CH₂)_n-;
 wherein n is independently 1 or 2.

[0101] Also described herein but not according to the invention is a complex, or a pharmaceutically acceptable salt thereof, wherein W has the structure:




wherein R²⁰ and R²¹ are each independently an amino acid residue linked via an amino group thereof to the adjacent -C(O)- group.

[0102] Also described herein but not according to the invention is a complex, or a pharmaceutically acceptable salt thereof, wherein W has the structure:




wherein R is hydrogen or a carboxylic acid protecting group.

[0103] Also described herein but not according to the invention is a complex, or a pharmaceutically acceptable salt thereof, having the structure:



or a pharmaceutically acceptable salt thereof,
 wherein R¹⁷ is aryl.

[0104] Also described herein but not according to the invention is a complex, or a pharmaceutically acceptable salt thereof, wherein the complex is PSMA-617:




[0105] With a radionuclide such as ^{212}Pb linked/chelated to four N.

[0106] Also described herein but not according to the invention is a complex, wherein the DOTA unit, such as p-SCN-Bz-DOTA, is substituted with a TCMC unit, such as p-SCN-Bz-TCMC.


[0107] The linker in PSMA-617 may also be covalently linked to from a C-atom in the backbone instead of link to N as can be seen in the above figure.

[0108] Thus, the urea derivative may have backbone C-linked or N-linked DOTA or TCMC conjugation.

[0109] For a backbone-C linked p-SCN-Bn-DOTA or p-SCN-Bn-TCMC conjugation is the compound:




[0110] Wherein Z is:



[0111] And wherein X is: -OH or NH₂.


[0112] X is -OH for p-SCN-Bn-DOTA and X is NH₂ for p-SCN-Bn-TCMC.

[0113] This means that the formula for backbone-C linked p-SCN-Bn-DOTA or p-SCN-Bn-TCMC will be:



wherein X is: -OH or NH₂.


[0114] Backbone-C linked p-SCN-Bn-DOTA is:



p-SCN-Bn-DOTA-PSMA ligand 2

[0115] Thus, described herein but not according to the invention is a compound which is p-SCN-Bn-DOTA-PSMA ligand 2.

[0116] Backbone-C linked p-SCN-Bn-TCMC is:



p-SCN-Bn-TCMC-PSMA ligand 1

[0117] Thus, described herein is a compound which is p-SCN-Bn-TCMA-PSMA ligand 1.

[0118] The p-SCN-Bn-TCMC-PSMA ligand (ligand 1) and p-SCN-Bn-DOTA-PSMA ligand (ligand 2) (as shown in the examples) is linked via the carbon backbone (C-backbone) to the rest of the compound and has an extended linker region including a isothiocyanato-benzyl linker and also uses a carbon substituted chelator with all 4 chelator arms free as opposed to PSMA-617 which has a shorter linker region and uses one of the chelator arms as linker attachment. It is shown in the later examples herein that these differences cause a significant different biodistribution of the radiolabelled product, making it more suitable for targeting of ^{212}Pb to PSMA-expressing tumors, as it reduces kidney exposure compared to PSMA-617.

[0119] The compounds p-SCN-Bn-TCMC-PSMA ligand 1 and p-SCN-Bn-DOTA-PSMA ligand 2 may be in the form of a trifluoroacetic acid salt.

[0120] CTT1401 and CTT1403 and derivatives (Choy et al, 2017) with TCMC or DOTA variants may also be used with ^{212}Pb in complexes.

[0121] By preparing a PSMA binding Urea derivative according to the present disclosure with a readily accessible amino group or another group suitable for conjugation, p-SCN-Bn-TCMC can be conjugated to the PSMA binding compound. Subsequent purification may be needed before radiolabelling with ^{212}Pb with or without ^{224}Ra in solution.

[0122] Thus, described herein is the linker p-SCN-Bn-TCMC. In some instances of a complex described herein the linker-chelator is p-SCN-Bn-TCMC. In some instances of a complex described herein the linker is p-SCN-Bn-DOTA. In other words -p-SCN-Bn- is the end part of the linker region which is attached to the TCMC or DOTA chelator group via the carbon backbone.

[0123] Human serum albumin can be used to prolong the half-life of drugs. Thus, the ^{212}Pb -labeled PSMA binding urea derivative described herein may further comprise a group that can associate with albumin to increase the circulation half-life of the radiolabeled product.

[0124] Also described herein is a complex described herein, wherein ^{212}Pb -labeled PSMA binding urea derivative described herein further comprises human serum albumin that has been directly conjugated to the complex, or is associated to the complex, for example through liposomes.

Pharmaceutical compositions

[0125] Also described herein is a solution comprising the compound and/or the complex described herein. The solution can also be a pharmaceutical composition.

[0126] Usually is an important element of a pharmaceutical composition a buffer solution, which to a substantial degree maintain the chemical integrity of the radioligand and is being physiologically acceptable for infusion into patients.

[0127] The pharmaceutical composition may comprise one or more pharmaceutically acceptable carriers and/or adjuvants.

[0128] Acceptable pharmaceutical carriers include but are not limited to non-toxic buffers, fillers, isotonic solutions, etc. More specifically, the pharmaceutical carrier can be but are not limited to normal saline (0.9 %), half-normal saline, Ringer's lactate, 5 % Dextrose, 3.3 % Dextrose/0.3 % Saline. The physiologically acceptable carrier can contain an anti-radiolytic stabilizer, e.g., ascorbic acid, which protect the integrity of the radiopharmaceutical during storage and shipment.

[0129] Described herein is a pharmaceutical composition comprising the complex described herein, and a diluent, carrier, surfactant, and/or excipient.

[0130] Also described herein is a single agent ^{212}Pb -labeled ligand. This would be the compound described herein complexed with ^{212}Pb and without any further radionuclides, such as ^{224}Ra .

[0131] Also described herein is a dual targeting solution containing ^{212}Pb -labeled ligand and cationic or weakly complexed ^{224}Ra . This would be the compound described herein complexed with ^{212}Pb with ^{224}Ra present as cationic or weakly complexed ^{224}Ra .

[0132] Also described herein is a pharmaceutical composition comprising the compound and/or the complex described herein, further comprising ^{224}Ra . ^{224}Ra can be cationic. The addition of ^{224}Ra allows for dual targeting for example when present together with ^{212}Pb .

[0133] Also described herein is a pharmaceutical composition comprising the complex described herein, wherein the radioactivity is 100 kBq to 100 MBq per dose.

[0134] Also described herein is a pharmaceutical composition comprising the complex described herein, wherein the amount of ^{224}Ra and ^{212}Pb is in radioactive equilibrium.

[0135] Also described herein is a pharmaceutical composition comprising the complex described herein, wherein the activity ratio (MBq) between ^{212}Pb to ^{224}Ra is between 0.5 and 2, such as 0.8 - 1.5, or such as 0.8 - 1.3, or preferably such as 0.9 - 1.15.

Kits

[0136] The solution should be made physiologically suitable for injections either at a centralized production site or be made up by a kit system of typically 2-4 vials whereby being physiologically suitable for injection after combination of the kit vials.

[0137] Accordingly, described herein is a kit comprising a first vial comprising a radiopharmaceutical composition described herein, and a second vial comprising a neutralizing solution to adjust pH and/or isotonicity of the radiopharmaceutical composition prior to administration to a patient.

[0138] For, e.g., a monoclonal antibody, it is usually advisable to keep the self-dose of the alpha particle producing radiopharmaceutical solution below 0.5 kGy to avoid reduced binding properties due to radiolysis. Thus, a kit system whereby chelator conjugated ligand is added to the ^{224}Ra (including daughters) solution a few hours to 10 minutes before injection is advised for concentrated solutions intended for remote shipping, depending of the radiolytic resistance of the radioligand that is generated.

[0139] Described herein is a kit comprising a first vial comprising a ^{224}Ra solution; a second vial comprising a complexing agent which is p-SCN-Bn-TCMC-PSMA ligand 1, wherein the complexing agent is capable of complexing a daughter nuclide of ^{224}Ra , such as ^{212}Pb , and wherein the complexing agent does not complex ^{224}Ra in the pharmaceutical solution; and optionally, instructions for mixing the first vial and the second vial, thereby forming a pharmaceutical composition ready to be administered to a patient 1 minute to 12 hours after mixing.

[0140] In one embodiment the kit is for use as a medicament.

[0141] In a specific embodiment, the term " ^{224}Ra solution" is to be understood as ^{224}Ra is free in the solution and not coupled to e.g. a surface such as a resin.

[0142] In an embodiment, the kit comprises a third vial comprising a neutralizing solution to adjust pH and/or isotonicity of the radiopharmaceutical solution prior to administration to a patient.

[0143] In yet a preferred embodiment, the amount of ^{224}Ra and ^{212}Pb is in radioactive equilibrium in the first vial.

[0144] In yet another preferred embodiment the activity ratio (MBq) between ^{212}Pb to ^{224}Ra in the first vial is between 0.5 and 2, such as 0.8 - 1.5, or such as 0.8 - 1.3, or such as 0.9 - 1.15.

[0145] In yet another embodiment the first vial has a radioactivity in the range 100 kBq to 100 MBq.

[0146] In one embodiment, the chelator conjugated ligand is added to the ^{224}Ra (including daughters) solution 30 min to 5 hour before injection, such as 1-3 hours before injection.

[0147] In one embodiment, the chelator conjugated ligand is added to the ^{224}Ra (including daughters) solution 1 min to 20 min before injection.

[0148] In one embodiment, the chelator conjugated ligand is added to the ^{224}Ra (including daughters) solution 1 min to 10 min before injection.

[0149] A kit with a chelate labelled protein or peptide in one vial and a ^{224}Ra solution in another vial whereby the content of the two are mixed 12 hours to 1 minute before administration is also described herein. In an embodiment, the mixing takes place a few hours (such as 5) to 30 minutes before administration to a patient as to bind ^{212}Pb and or ^{212}Bi to the chelate.

[0150] In one embodiment, the content of the two are mixed 30 min to 1 hour before injection.

[0151] In one embodiment, the content of the two are mixed 1 min to 20 min before injection.

[0152] In embodiment, the content of the two are mixed 1 min to 10 min before injection.

[0153] Optionally, a third vial containing a liquid used for dilution and isotonicity adjustment before administration of the radiopharmaceutical solution could be used. This third vial may contain EDTMP, which could chelate ^{212}Bi , if needed.

Medical applications

[0154] Also described herein is the radiopharmaceutical composition described herein for use as a medicament.

[0155] In one embodiment is the disease cancer.

[0156] Also described herein is the radiopharmaceutical composition described herein for use in the treatment of soft tissue and/or skeletal disease. The treatment is focused on PSMA-expressing disease including soft tissue- and skeletal disease.

[0157] In one embodiment is the skeletal disease selected from the group consisting of soft tissue and or skeletal metastases from cancers to the breast, prostate, kidneys, lung, bone, or multiple myeloma.

[0158] In one embodiment is the cancer prostate cancer. The cancer can also be breast cancer. The cancer can be kidney cancer. The cancer can also be lung cancer. The cancer can also be bone cancer. The cancer can also be multiple myeloma. The cancer can be metastases from these types of cancer.

[0159] In one embodiment is the solution administered at a dose in the range 50-150 kBq per kg of bodyweight, such as 50-100 kBq per kg of bodyweight.

[0160] Also described herein is a method of treatment of malignant or non-malignant disease by administration of a

radiopharmaceutical composition described herein to an individual in need thereof.

[0161] This may be used with purified ^{212}Pb -labeled ligand or in a dual targeting solution whereby ^{224}Ra will act as a skeletal treatment, and ^{212}Pb -urea-derivative will act as a systemic therapy against cells the expressing PSMA antigen which is associated with advanced metastatic prostate cancer.

[0162] Thus, the complexes and the solutions described herein can be used in the treatment of metastatic prostate cancer.

[0163] Also described herein is a pharmaceutical solution with dual targeting properties whereby ^{212}Pb is complexed by a urea based PSMA targeting agent as disclosed herein, and cationic ^{224}Ra is targeting bone metastases through calcium mimetic bone incorporation.

Methods for preparation

[0164] Also described herein is a method for providing a radiopharmaceutical composition described herein, the method comprising providing a first solution wherein the amount of ^{224}Ra and ^{212}Pb is in radioactive equilibrium; providing a second solution comprising a complexing agent which is p-SCN-Bn-TCMC-PSMA ligand 1, wherein the complexing agent is capable of complexing a daughter nuclide of ^{224}Ra , such as ^{212}Pb , and wherein the complexing agent does not complex ^{224}Ra ; and mixing the first composition and the second composition, thereby providing a pharmaceutical composition described herein.

PSMA derivatives

[0165] PSMA (also named Prostate Specific Cancer Antigen, PSM, FGCP, FOLH, GCP2, mGCP, GCPII, NAALAD1, NAALAdase, FOLH1, Glutamate carboxypeptidase 2, Glutamate carboxypeptidase II, Membrane glutamate carboxypeptidase, N-acetylated-alpha-linked acidic dipeptidase I, Pteroylpoly-gamma-glutamate carboxypeptidase, Folylpoly-gamma-glutamate carboxypeptidase, Folate hydrolase 1, Prostate-specific membrane antigen, Cell growth-inhibiting protein 27) is a prostate epithelial cell membrane antigen of type II transmembrane protein and consist of a short NH₂-terminal cytoplasmic domain, hydrophobic transmembrane region, and a large extracellular domain. PSMA is an enzyme that in humans is encoded by the FOLH1 (folate hydrolase 1) gene. Human GCPII contains 750 amino acids and weighs approximately 84 kDa.

[0166] Expression of PSMA is restricted to a few healthy tissues such as lacrimal and salivary glands, proximal renal tubules, epididymis, ovary, the luminal side of the ileum-jejunum and astrocytes within the central nervous system (CNS); healthy prostate gland expresses comparatively little PSMA, which is confined within the apical epithelium of secretory ducts. In these non-malignant tissues, uptake of PSMA-targeted probes may be limited by an intact blood-brain barrier, a healthy proximal small bowel lumen, and truncated cytoplasmic expression of PSMA within normal prostate. PSMA is most associated with high grade androgen independent, metastatic disease, although PSMA is expressed in most primary prostate tumors regardless of androgen status.

[0167] Small molecular PSMA inhibitors are zinc binding compounds and can be classified into three types 1) phosphonate-, phosphate-, and phosphoramidate compounds; 2) thiols; and 3) ureas. The urea derivatives seem to have particularly interesting properties as carrier for radionuclides for diagnosis and therapy.

[0168] Described herein is the use of ^{212}Pb -urea derivatives. It may be combined with androgen deprivation therapy to enhance PSMA expression for better uptake of radioligand (Bakht et al, 2017).

Activity level

[0169] If a ^{212}Pb -labeled urea derivative is used alone, the activity level would typically be between 1 MBq and 500 MBq per patient, more typically 10 -100 MBq per patient.

[0170] If ^{224}Ra in equilibrium with ^{212}Pb complexed with a PSMA binding urea derivative the dosing would typically be between 0.1 MBq and 100 MBq per patient, more typically between 1 and 20 MBq per patient.

[0171] Also described herein is a urea derivative based PSMA targeting complex described herein, labeled with ^{227}Th , which produces bone-seeking ^{223}Ra for dual targeting.

[0172] The ^{227}Th may be pure or containing various amounts of ^{223}Ra , eg. 10%, 50%, 100% or 250% of ^{223}Ra compared to ^{227}Th .

General

[0173] It should be understood that any feature and/or aspect discussed above in connections with the compounds described herein apply by analogy to the methods described herein.

[0174] The following figures and examples are provided below to illustrate the present invention. They are intended

to be illustrative and are not to be construed as limiting in any way.

EXAMPLES

[0175] In the following examples the xenograft model used is a tumor model with intermediary level of PSMA-ligand uptake of typically 10-15% of injected dose per gram (%ID/g) in mouse xenografts with ^{177}Lu -PSMA-617 as opposed to the PC3 PIP model used by other researchers which shows an uptake of typically 30-40% ID/g in mouse xenografts.

[0176] *Example 1. Novel PSMA-binding chelator ligands compared with PSMA-617.*

[0177] Background: Several carrier molecules for radioligand targeting of prostate specific membrane antigen (PSMA) exists. Lutetium-177 labeled PSMA-617 (^{177}Lu -PSMA-617) is the compound in most advanced clinical development stage for use in radionuclide therapy. This molecule works well and give relevant tumor to normal tissue ratios for longer lived (i.e., a half-life of a few days) radionuclides, including ^{177}Lu and ^{225}Ac , but at early times points (typically a few hours after injection) shows high uptake in kidneys. With shorter lived radionuclides like ^{212}Pb (half-life of 10.6 hours), the initial kidney uptake represents a potential toxicity problem. It is therefore advantageous to use a PSMA-ligand with less kidney uptake, but this should not compromise the tumor uptake. The PSMA ligand molecules are made up of (1) a PSMA-binding region, (2) a linker region and (3) a chelator, whereby the linker region connects the (1) and (3). The linker region also is used to adjust molecular size and polarity to affect the in vivo distribution properties. The PSMA-binding region (motif) used in PSMA-617 is a structure that can be found in several molecules of this class developed by several different inventors and researchers including PSMA-11 and PSMA I&T as well as ^{131}I - and ^{211}At -labelled PSMA binding ligands.

[0178] As can be seen, the p-SCN-Bn-TCMC-PSMA ligand 1 has an extended linker region including a isothiocyanato-benzyl linker and also uses a carbon substituted chelator with all 4 chelator arms free as opposed to PSMA-617 which has a shorter linker region and uses one of the chelator arms as linker attachment. It is shown in later examples herein that these differences cause a significant different biodistribution of the radiolabelled product, making p-SCN-Bn-TCMC-PSMA ligand 1 it more suitable for targeting of ^{212}Pb to PSMA-expressing tumors, as it reduces kidney exposure compared to PSMA-617.


[0179] Materials and methods: The PSMA-ligand precursors for radiolabeling were synthesized by a subcontractor commercial synthesis laboratory.

[0180] PSMA-617 was synthesized according to procedures described in the literature. The p-SCN-Bn-TCMC precursors were synthesized according to procedures described in the literature. In the final synthesis step the TCMC-PSMA ligand was synthesized by conjugating p-SCN-Bn-TCMC to amino group of a PSMA binding ligand intermediate. PSMA-617 and TCMC-Bn-PSMA-ligand 1 were both purified by HPLC to a purity of > 98% and dried and stored as the trifluoroacetic acid salts. Structures and molecule weights were determined with ^1H -NMR and MS analysis.


[0181] The p-SCN-Bn-TCMC-PSMA ligand 1 trifluoroacetic acid salt has the chemical formula $\text{C}_{65}\text{H}_{86}\text{F}_{12}\text{N}_{14}\text{O}_{21}\text{S}$ and a molecular weight of 1659.52 g/mol.

[0182] Chemical Name (IUPAC): (((1S)-1-carboxy-5-((2S)-3-(naphthalen-2-yl)-2-((1R,4S)-4-((3-(4-((1,4,7,10-tetrazakis(2-amino-2-oxoethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)thioureido)methyl)cyclohexane-1-carboxamido)propanamido)pentyl)carbamoyl)-L-glutamic acid; trifluoroacetic acid (1:4)

[0183] The PSMA-617 trifluoroacetic acid salt has the chemical formula $\text{C}_{57}\text{H}_{75}\text{F}_{12}\text{N}_9\text{O}_{24}$ and a molecular weight of 1498.25 g/mol.



p-SCN-Bn-TCMC-PSMA ligand 1 trifluoroacetic acid salt.



PSMA-617 trifluoroacetic acid salt

[0184] As can be seen, the p-SCN-Bn-TCMC-PSMA ligand 1 has an extended linker region including a benzyl linker and also uses a carbon substituted chelator with all 4 chelator arms free as opposed to PSMA-617 which has a shorter linker region and uses one of the chelator arms as linker attachment. It is shown in the following examples that these chemical differences cause a significant different biodistribution of the radiolabelled product, making it more suitable for targeting of ^{212}Pb to PSMA-expressing tumors, as it reduces kidney exposure compared to PSMA-617.


[0185] In conclusion, a novel molecule with same PSMA-binding region, but with different linker region and different chelation properties compared with PSMA-617 is described.

Reference Example 2

[0186] A carbon substituted p-SCN-Bn-DOTA-PSMA ligand 2 for radiolabeling with ^{212}Pb , ^{212}Bi , ^{213}Bi , ^{225}Ac , ^{227}Th , ^{177}Lu , etc.

[0187] Compared to PSMA-617 the ligand has larger size and a p-SCN-Bn-DOTA group, i.e., both a different linker region and a different DOTA-chelator version where all four chelator arms are free to cause chelation of radionuclides as compared to PSMA-617.

[0188] The p-SCN-Bn-DOTA-PSMA ligand 2 is the DOTA analogue of p-SCN-Bn-TCMC-PSMA ligand 1 described in example 1 and has carbon linker to the chelator backbone leaving the chelator groups free to interact with the radiolabel and is therefore expected to improve chelate stability after radionuclide labelling compared with radiolabelled PSMA-617. Due to the additional lipophilic benzyl unit in the linker region and larger size of molecule, less kidney uptake is expected compared with PSMA-617. This molecule can be synthesized in the same fashion as p-SCN-Bn-TCMC-PSMA ligand 1 by using a DOTA based precursor instead of a TCMC-based precursor in the last step of synthesis.



p-SCN-Bn-DOTA-PSMA ligand 2 trifluoroacetic acid salt.

[0189] The p-SCN-Bn-DOTA-PSMA ligand 2 trifluoroacetic acid salt has the chemical formula $\text{C}_{65}\text{H}_{82}\text{F}_{12}\text{N}_{10}\text{O}_{25}\text{S}$ and a molecular weight of 1663.46 g/mol.

[0190] With the carbons substituted attachment to the DOTA, this molecule has properties very suitable for radiolabeling with ^{212}Pb , ^{212}Bi , ^{213}Bi , ^{225}Ac , ^{227}Th , and also for positron emission tomography (PET) compatible radionuclides like,

⁶⁸Ga, as well.

[0191] Thus, a novel reference PSMA-binding molecule is described, p-SCN-Bn-DOTA-PSMA ligand 2, suitable for radioligand imaging and therapy.

[0192] In conclusion, carbon substituted p-SCN-Bn-DOTA-PSMA ligand 2 is described with different properties in terms of size and chelation properties compared with PSMA-617.

Example 3. Radionuclides tested.

[0193] Lutetium-177 was purchased as ready to use ¹⁷⁷LuCl₃ dissolved in diluted HCl. Lead-212 was obtained from ²²⁴Ra based solutions. Radium-224 was made from ²²⁸Th bound to Actinide resin (Eichrom Technologies, LLC) by eluting a column containing actinide resin with immobilized ²²⁸Th with 1 M HCl. The eluate was purified on a second Ac-resin column and the eluate evaporated to dryness using an evaporation vial with a cap with gas inlet and outlet placed in a heater block at approximately 110 °C and a gentle stream of nitrogen gas to evaporate of the solvent. When the evaporation vial was empty from solvent it was added 0.1 M HCl to dissolve the residue, typically 200-400 µl.

Example 4. Radiolabeling of PSMA-binding ligands.

[0194] In general, ¹⁷⁷Lu and ²²⁴Ra/²¹²Pb solutions were adjusted with HCl with 10% 5 M ammonium acetate to the desired volume and pH 5-6. The PSMA-binding ligands were dissolved in 0.1 M HCl with 0.5 M ammonium acetate to a concentration of 1 mg/ml. Typically a concentration of 20 microgram per 1 ml radioactive solution were used. The reaction mixture was incubated on a shaker for 15-30 minutes, typically a labelling was evaluated by thin layer chromatography. Lead-212 labeling was performed at room temperature or 37 °C with the p-SCN-Bn-TCMC-PSMA ligand 1, while ¹⁷⁷Lu-labeling and ²¹²Pb-labeling of PSMA-617 was performed at 90 °C. Typical radiolabeling yields were in the range 90-100% with the compounds and the radionuclides tested, providing a concentration of 1 µg per 20 µl or higher were used.

[0195] In conclusion, radiolabeling of both ligands worked well. The novel p-SCN-Bn-TCMC-PSMA ligand 1 could be radiolabelled with ²¹²Pb at room temperature as opposed to ¹⁷⁷Lu-PSMA-617 that had to be labeled at elevated temperature.

Example 5: Thin layer chromatography analyses

[0196] Thin layer chromatography (TLC) was performed using chromatography strips (model # 150-772, Biodex Medical Systems Inc, Shirley, NY, USA). A formulation buffer (FB) consisting of 7.5% human serum albumin and 5 mM EDTA in DPBS and adjusted to approximately pH 7 with NaOH was mixed with the antibody conjugates in ratio 2:1 for at least 5 minutes before application to the strips to determine free radionuclide. A small beaker with about 0.5 ml of 0.9% NaCl was used to place strips with a sample spot in. To the strip was typically added 1-4 µl of sample at approximately 10% above the bottom of the strip. After the solvent front had moved to about 20% from the top of the strip, the strip was cut in half and each half was placed in a 5 ml test tube for counting. In this system radiolabeled ligand stays at the bottom half while radionuclide complexed with EDTA migrates to the upper half. It was verified that free cations of both ²¹²Pb and ¹⁷⁷Lu would complex with EDTA and moved to the top.

[0197] In conclusion, a TLC system was used allowing a rapid determination of radiolabeling yield that would distinguish effectively between radioligand and free radionuclide.

Example 6. Separation of ²¹²Pb from ²²⁴Ra solutions

[0198] Radiolabeled ligand can be used as a component in ²²⁴Ra dual targeting solutions whereby ²²⁴Ra targets skeletal disease and ligand targets systemic metastatic disease. Alternatively, ²²⁴Ra generator solutions can be used for producing ²¹²Pb-labeled ligand by labelling in situ, i.e., ²¹²Pb is complexed by the ligand in the presence of ²²⁴Ra. To 40 µl Radium-224/²¹²Pb solutions in 0.5 M ammonium acetate, pH 5-6 was added 2 µl (1 µg/µl) of either p-SCN-Bn-TCMC-PSMA ligand 1 or PSMA-617 and reacted as described to generate the ²¹²Pb-labeled ligands. To purify ²¹²Pb-PSMA ligand the product was added about 10 µl of a formulation buffer consisting of 7% bovine serum albumin, 10 mM EDTA, and 10 mg/ml ascorbic acid. Thereafter the reaction mixture was added to a Sephadex G-10 column PD MiniTrap G-10 (GE Healthcare Life Sciences) and eluted with 0.9% NaCl. The eluate containing the ²¹²Pb, typically the fractions eluted after application of 0.7-1.5 ml, was collected and analysed on a gamma counter and TLC and radioligand binding assay was performed. The product purification procedure had high radiochemical yield (typically >80%) and had a high radiochemical purity with ²²⁴Ra amounting to less than 0.4% compared with ²¹²Pb-labeled ligand, typically.

[0199] In conclusion, p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 could be radiolabelled with ²¹²Pb in the presence of ²²⁴Ra yielding a dual targeting solution for combined bone metastases targeting with ²²⁴Ra and systemic tumor

cell targeting with ^{212}Pb -labeled PSMA ligand.

[0200] It was also found that using Sephadex G-10 gel filtration column, both ^{212}Pb -labeled PSMA-ligands tested could be separated from ^{224}Ra in the solution with a recovery above 80% and breakthrough of ^{224}Ra of less than 0.4 % yielding a highly purified ^{212}Pb -labeled PSMA radioligand for stand-alone PSMA targeting.

Example 7. Stability testing of ^{212}Pb -labeled ligands in vitro.

[0201] Radiolabeled ligand in $^{224}\text{Ra}/^{212}\text{Pb}$ solution was mixed 1:1 with PBS or bovine serum albumin and incubated at 37 °C for up to 48 hours. TLC analyses were performed at 1 h, 4 h, 24 h and 48 h of incubation.

[0202] The data are shown in table 1 and indicates that ^{212}Pb continuously, after it is generated from ^{224}Ra , react with the ligand and maintain a high percentage of radiochemical purity even after 48 hours. Thus, the PSMA-ligands are compatible with the use of a ^{224}Ra solution for in situ production of radioligand that is suitable for centralized production and shipment to the end user.

Table 1. Radiochemical purity of ^{212}Pb -TCMC-PSMA ligand 1 and ^{212}Pb -PSMA-617 in PBS and FBS

Time since start of incubation	^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1		^{212}Pb -PSMA-617	
	PBS	FBS	PBS	FBS
1h	93.26%	93.50%	91.62%	92.13%
4h	95.04%	94.18%	94.33%	94.37%
24h	95.00%	93.54%	93.26%	94.54%
48h	95.11%	90.81%	93.17%	95.65%
Conclusion: The data indicate that p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 labelled with ^{212}Pb are stable in ^{224}Ra solutions for a prolonged period of time indicating compatibility with centralized production and shipment to the end user of ready to use product. Such solutions could be used for treatment against cancer.				

Example 8. Prostate cancer cell binding of p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 labeled with ^{212}Pb .

[0203] the cell binding fractions measured by adding about 1 ng of radioligand to 0.2 ml of C4-2 cells (5×10^7 cells per ml) in a 5 ml test tube and incubate for 1 h before measuring applied activity, wash cells 3 times with 0.5 ml 0.5% bovine serum albumin in DPBS and thereafter recount the washed cell pellet. From multiple experiments it was found that the % bound was in the range of 40-53% after subtracting for nonspecific binding. Nonspecific binding was measured by blocking the cells with excess 10 $\mu\text{g}/\text{ml}$ of unlabelled ligand before adding radioligand. No significant difference in cell-binding was found between radiolabelled p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617. In conclusion, the radioligand of radiolabelled p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 had similar cell binding properties in vitro indicating that p-SCN-Bn-TCMC-PSMA ligand 1 has relevant antigen binding ability.

Example 9. Radiolytic stability of p-SCN-Bn-TCMC-PSMA ligand 1 assessed by radioligand binding ability with prostate cancer cells and TLC analyses.

[0204] Table 2 shows the cell binding fractions measured by adding about 1 ng of radioligand to 0.2 ml of C4-2 cells (5×10^7 cells per ml) in a 5 ml test tube and incubate for 1 h before measuring applied activity, wash cells 3 times with 0.5 ml 0.5% bovine serum albumin in DPBS and thereafter recount the washed cell pellet. Initial activity in the solution was about 5 MBq/ml causing an absorbed radiation dose to the solution of about 1.8 Gy after 24 h and 3.3 Gy after 48 hours. The data (Table 2) indicate a slight decline at the longest exposure time but in general a relatively strong radiolytic resistance of the ligand was observed, which is compatible with centralized production and shipment to the end user ^{224}Ra -based generator solution with or without removal of ^{224}Ra prior to the administration of product.

Table 2. Binding ability of ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 when kept in ^{224}Ra solution.

Time since start of incubation	Cell binding fraction	TLC measured RCP
0.5h		93.81%
1h		92.84%
4h	52.3%	94.75%

(continued)

Time since start of incubation	Cell binding fraction	TLC measured RCP
24h	48.16%	94.85%
48h	37.18%	94.85%
Conclusion: The data indicate that ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 is stable in ^{224}Ra solutions for a prolonged period of time and the PSMA-ligand is capable of complexing ^{212}Pb as it is generated from ^{224}Ra during storage, and provided that the absorbed radiation dose is kept below about 2 kGy such solutions could be used for treatment against cancer.		

Example 10. Biodistribution of radiolabelled ligands in nude mice with C4-2 PSMA-positive xenografts.

[0205] The biodistribution of $^{224}\text{Ra}/^{212}\text{Pb}$ solutions with p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 labeled with ^{212}Pb and ^{177}Lu were compared after intravenous injection in nude mice with C4-2 xenografts at various time points after injection. Each group usually consisted of three mice. The ^{212}Pb labelling was above 92% for the products. The molar concentration of ligand was significantly lower for the ^{177}Lu -PSMA-617 since much higher level of radionuclide was used with ^{177}Lu vs. ^{212}Pb . Approximately 16 kBq of $^{224}\text{Ra}/^{212}\text{Pb}$ was injected in each animal, i.e., approximately 0.2 nmol of ligand per mouse. Animals were given anaesthesia and sacrificed by cervical dislocation followed by dissection and harvesting of tissue-, blood- and urine samples. The samples were weighed and counted on a gamma counter.

Example 11. Comparison of tumor binding and kidney uptake of p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 labelled with ^{212}Pb in mice.

[0206] The biodistribution of $^{224}\text{Ra}/^{212}\text{Pb}$ solutions with p-SCN-Bn-TCMC-PSMA ligand 1 and PSMA-617 labeled with ^{212}Pb were compared after intravenous injection in nude mice with C4-2 xenografts at 4 h after injection. Three mice in each group. The ^{212}Pb labelling was above 92% for both products. The molar concentration was the same for both products, i.e., 12.5 nmol per MBq. Approximately 16 kBq of $^{224}\text{Ra}/^{212}\text{Pb}$ was injected in each animal, i.e., approximately 0.2 nmol of ligand per mouse. Animals were given anaesthesia and sacrificed by cervical dislocation followed by dissection and harvesting of tissue-, blood- and urine samples. The samples were weighed and counted on a gamma counter. Results: Four hours after injection, the ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 to PSMA-617 ratios were as follows: tumor 1.35; kidneys 0.20; blood 1.21; liver 2.67; spleen 0.71. It was confirmed by counting samples after 3 days storage that the ^{224}Ra biodistribution was not significantly altered by either of the PSMA-directed ligands. Discussion: ^{212}Pb p-SCN-Bn-TCMC-PSMA ligand 1 showed a significant different biodistribution compared to ^{212}Pb -PSMA-617 and especially noteworthy is the very low, and favourable ratio for uptake in the kidneys as kidneys are expected to be the main dose limiting normal tissue related to PSMA radioligand therapy with relatively short-lived alpha-emitters (Figure 1). In conclusion, ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 shows very promising early time point biodistribution compared with PSMA-617 which is important when using shorter lived radionuclides like ^{212}Pb .

Example 12. Comparison of tumor binding and kidney uptake of ^{212}Pb -labeled p-SCN-Bn-TCMC-PSMA ligand 1 and ^{177}Lu -labeled PSMA-617 in mice.

[0207] Methods: Tumor and kidney uptake of ^{212}Pb -labeled p-SCN-Bn-TCMC-PSMA ligand 1 and ^{177}Lu -labeled PSMA-617 was compared at 1 hours and 4 hours after administration of the products by intravenous injection into nude mice with C4-2 xenografts as described in Example 9. Results: The tumors and the kidneys were the tissues taking up the largest amounts of radioactivity. As examples the tumor and kidney uptake of ^{212}Pb -p-SCN-Bn-TCMC-PSMA were on average 13.9 and 8.1 percent of injected dose per gram (%ID/g) respectively at 4 hours after injection. The tumor and kidney uptake of ^{177}Lu -PSMA-617 were on average 13.6 and 17.4 %ID/g respectively 4 hours after injection. It is noteworthy that the molar amount of ligand injected was much lower for PSMA-617, which is known to reduce kidney uptake, but still the novel ^{212}Pb -labeled compound showed less kidney uptake. The tumor-to-kidney ratios at the 4-hour timepoint were as follows: ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1, 1.7; ^{177}Lu -PSMA-617, 0.8.

[0208] The average tumor-to-kidney ratios determined at the 1-hour point after administration were for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1, 0.40 and for ^{177}Lu -PSMA-617, 0.17. In conclusion, despite a higher molar ligand concentration for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1, it showed better tumor to kidney ratios than ^{177}Lu -PSMA-617 indicating that it may be well suited for ^{212}Pb based alpha emitter radioligand therapy.

Example 13. Biodistribution of single targeting solution containing ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 in mice with PSMA-positive xenografts.

[0209] Using a $^{212}\text{Pb}/^{224}\text{Ra}$ solution for reaction with p-SCN-Bn-TCMC-PSMA ligand 1 as described, purified ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 was purified using a Sephadex G-10 gel filtration column and about 30 kBq and 300 ng of the purified radioligand product was injected per animal. The data are shown in Table 3. As can be seen the kidney activity is reduced relatively quickly, while the tumor uptake shows good retention. The tumor-to-tissue ratios (Table 4) indicates suitability for radioligand targeting with ^{212}Pb .

[0210] In conclusion, ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 shows relevant targeting properties for use in radioligand therapy against PSMA-expressing prostate cancer.

Table 3. Biodistribution for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 at various timepoints post injection

Organ	% ID/g			
	1 h	2h	4 h	8 h
Blood	1,77	0,54	0,46	0,13
Urine	253,36	250,85	32,12	6,76
Testes	4,41	0,56	0,38	0,12
Prostate	13,95	3,80	1,48	-0,10
Salivary gland	0,72	0,32	0,34	0,15
Tumor	26,44	15,87	14,14	14,65
Skin	1,70	0,47	0,42	0,23
Kidneys	63,96	25,41	9,19	4,07
Liver	2,39	1,33	1,55	1,25
Spleen	1,24	0,44	0,54	0,36
Small intestine	0,34	0,23	0,30	0,08
Large intestine	0,25	0,39	0,18	0,17
Stomach	0,16	0,22	0,10	0,09
Lungs	1,19	0,40	0,75	0,17
Heart	0,78	0,20	0,52	0,09
Bladder	36,86	6,85	3,53	0,26
Femur	0,87	0,40	0,57	0,74
Muscle	0,62	0,22	0,23	0,04
Brain	0,61	0,04	0,07	0,02
Skull	0,77	0,38	0,37	0,63

Table 4. Tumor to tissue ratios for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 at various timepoints

	1h	2h	4 h	8 h
Tumor/blood	14.9	29.4	30.7	112.7
Tumor/kidneys	0.41	0.62	1.54	3.60
Tumor/muscle	42.6	72.1	61.5	366.2
Tumor/Femur	30.4	39.7	24.8	19.8

Example 14 - Dosage

[0211] The radiation energy produced for the two nuclides is mainly from alpha particles and therefore only the alpha particles are considered in the following estimate.

[0212] The ^{212}Pb and short-lived daughters produce on average 7.8 MeV alpha radiation per atom of ^{212}Pb . The half-life of ^{212}Pb is 10.6 h.

[0213] The ^{213}Bi and short-lived daughters produce on average 8.4 MeV of alpha energy per atom of ^{213}Bi . The half-life of ^{213}Bi is 46 min.

[0214] It is assumed an equivalent dose for alpha particles of 5 Sv/Gy.

[0215] Thus 1 Bq of ^{212}Pb produce an equivalent alpha dose of $1 \times (10.6 \times 60/46) \times 7.8/8.4 = 12.6$ Bq of ^{213}Bi when decaying completely.

[0216] It has been reported that salivary glands, kidneys and red marrow are the dose limiting tissues for ^{213}Bi and ^{225}Ac complexed to PSMA-617 (Kratochwil, et al, 2018).

[0217] Based on the imaging of PSMA-617 labeled with radionuclides suitable for positron emission tomography detection, it is assumed a maximum percentage uptake to be at the 30 minutes time point, that 70 % ^{213}Bi atoms relative to 90% of ^{212}Pb would reach and decay in tumors (i.e., the relative decay fraction).

[0218] The relative decay fraction for the dose limiting tissues is assumed to be 70% for all the tissues with ^{213}Bi -PSMA-617. For ^{212}Pb -PSMA-617 taken up it is assumed that 50% would decay in the salivary glands, 30% in kidneys and 20% in bone marrow.

[0219] By correcting for energy per Bq and the relative decay fraction for ^{212}Pb - and ^{213}Bi -labeled PSMA-617 the dose estimate for ^{212}Pb -PSMA-617 would be as presented in Table 5 together with previously published data for ^{213}Bi - and ^{225}Ac -labeled PSMA-617. Also, by using mouse data comparison of ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 and ^{212}Pb -PSMA-617 and assuming similar tissue uptake ratio in man dosimetry estimate for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 is presented in Table 5.

Table 5 Dose assessment assuming similar stability and affinity of products regardless of radionuclide.

Organ	^{225}Ac -PSMA-617 ^a Sv/MBq	^{213}Bi -PSMA-617 ^b Sv/GBq	^{212}Pb -PSMA-617 Sv/100 MBq	^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 Sv/100 MBq ^c
Salivary glands	2.33	8.1	5.67	5.67
Kidneys	0.74	8.1	3.40	1.70
Red marrow	0.05	0.52	0.19	0.19
Tumors	5.7	6.3	10.2	10.2

^aFrom Kratochwil et al 2017. ^bFrom Kratochwil et al 2018. ^cAssuming same uptake as for ^{212}Pb -PSMA-617 except for kidneys which is assumed to be reduced by 50%.

[0220] It has been reported that both ^{225}Ac - and ^{213}Bi -labeled PSMA-617 has been used clinically and shown considerable antitumor activity. Based on the estimates in this example it is indicated that ^{212}Pb -labeled urea based PSMA inhibitor is a very promising therapeutic tool against PSMA expressing cancer.

Example 15 Dosimetry estimate for dual targeting ^{224}Ra cation and ^{212}Pb -labeled PSMA targeting urea derivative.

[0221] Using ^{224}Ra in equilibrium with ^{212}Pb -labeled PSMA binding urea derivative.

[0222] For a dosage of 150 kBq of ^{224}Ra the total activity administered would be 10.5 MBq in a 70 kg person. By using published dosimetry for cationic ^{223}Ra in prostate cancer patient and correcting for half-life difference between ^{224}Ra and ^{223}Ra , assuming same biodistribution, and considering the different residence times in the various tissues it was found that ^{224}Ra would give per MBq administered 0.006 Gy to kidneys, 0.029 Gy to salivary glands, 0.26 Gy to red marrow and an estimated 5 times the red marrow uptake to tumors i.e., 1.3 Gy.

[0223] An equivalent dose of 5 Sv/Gy for alpha particle dose is assumed and the data is translated to Sv per injected dosage (10.5 MBq/patient) in Table 6.

Table 6 Dose assessment of ^{224}Ra cation + ^{212}Pb -labeled PSMA binding urea derivative*.

Organ	$^{224}\text{Ra}^{\text{a}}$ Sv/ 10.5 MBq	^{212}Pb -p-SCN-Bn-TCMC- PSMA ligand 1 Sv/10.5 MBq ^c	^{224}Ra cation + ^{212}Pb -p-SCN-Bn- TCMC-PSMA ligand 1 Sv/10.5 MBq
Salivary glands	37	0.60	0.97
Kidneys	0.39	0.36	1.7
Red marrow ^b	2.2	0.02	2.22
Tumors (bone metastases/soft tissue metastases)	10.5/0.09	1.07/1.07	11.6/1.16
*Assuming 1:1 ratio between ^{224}Ra and ^{212}Pb . ^a From Lassmann et al, 2002. ^b It should be noted that the red marrow is mainly irradiated from bone surface deposits of Ra, and due to short range of alpha-particles from the bone surfaces, substantial areas of red bone marrow are outside of reach from bone surface alpha particles. Equivalent dose of 5 for alpha-particle dose is assumed.			

Example 16. Dosimetry estimated for ^{227}Th -labeled PSMA binding urea derivative

[0224] In this example the ^{177}Lu is assumed labeled to p-SCN-Bn-DOTA-PSMA ligand 2 or a HOPO derived version of this molecule. The current estimates are based on adaptations of data from studies of ^{223}Ra in patients and ^{225}Ac -PSMA-617 in patients with prostate cancer (Chittenden et al., 2015, Kratochwil et al, 2017, 2018).

[0225] An equivalent dose of 5 Sv per Gy is assumed for the alpha particle radiation.

[0226] It is assumed that the ^{223}Ra generated in tissues would decay locally. It is assumed that 40% of the ^{223}Ra generated from whole body circulation of ^{227}Th is retained in the skeleton. Only the alpha particle dose is considered as this constitute 95% or more of the total radiation energy produced.

[0227] In tissue with a longer residence times for the radiolabeled PSMA urea derivative (salivary gland, kidneys and tumors) the cumulated activity of ^{223}Ra produced from ^{227}Th is assumed to be 20% of the ^{227}Th and for red bone marrow 5%.

[0228] The 20% and 5% radium generated from ^{227}Th in the various tissues are assumed to be in equilibrium with alpha-emitting progeny radionuclides, so each radium decay in effect produces 26.4 MeV of alpha radiation while ^{227}Th produces one alpha of 5.9 MeV.

[0229] Since residence times are lower than the half-lives of ^{225}Ac and ^{227}Th , the cumulated activity for ^{227}Th is assumed to be only 10% higher than for ^{225}Ac (not considering ^{225}Ac progenies) in all organs. It is assumed a total of 27.7 MeV of alpha dose per atom of ^{225}Ac from decay of the mother nuclide and alpha emitting progenies.

[0230] It is also assumed that the red marrow dose and skeletal tumor dose is increased by a factor of 2 vs. the red marrow dose, due to skeletal uptake of ^{223}Ra generated during systemic circulation etc of the ^{227}Th -product.

[0231] The data are presented in Table 7. The data indicate favorable tumor to tissue ratios for ^{227}Th -labeled PSMA binding urea derivative.

Table 7 Dosimetry estimated of equivalent doses for critical organs and tumors for ^{227}Th -labeled PSMA binding urea derivative (MBq/patient).

Organ	^{227}Th (Sv/MBq)	^{223}Ra (co-localized with ^{227}Th)	^{227}Th + ^{223}Ra
Salivary glands	0.55	0.49	1.04
Kidneys	0.17	0.15	0.32
Red marrow	0.012	0.003	0.030*
Tumors (soft tissue/bone met.)	1.33/1.33	1.19/1.19	2.52/2.67*
*Includes added skeletal dose of ^{223}Ra generated from ^{227}Th during circulation phase.			

[0232] The data in Table 7 represent ^{227}Th -labeled PSMA binding urea derivative (e.g. PSMA-617) which is purified

from ^{223}Ra prior to injection. Alternatively, it is possible to use the ^{227}Th -product solution with a presence of various amounts of ^{223}Ra to increase dose to the bone metastases, e.g., if the bone disease is very dominating compared with soft tissue disease.

[0233] It is assumed that the products described herein can be used in single treatment or in repeated treatment fashion.

[0234] In conclusion, the dosimetric estimates for PSMA-targeting urea derivatives labeled with ^{212}Pb or ^{227}Th indicates promising tumor to tissue ratios indicating that clinical benefit of use may be possible.

Example 17. Comparative therapy experiment with ^{177}Lu -PSMA-617 and ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 in nude mice with C4-2 Xenografts.

[0235] Male nude mice were inoculated in the flanks with C4-2 PSMA positive human prostate cells and 2 weeks later when tumors were 5-7 mm in diameters. Groups of 8 animals each received saline, 52 MBq of ^{177}Lu -PSMA-617 or 320 kBq of ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1. Animals were sacrificed when tumor size reached 20 mm due to animal welfare requirements. Tumor dosimetry was calculated based on the following assumptions: for ^{177}Lu -PSMA-617 effective half-life in tumor of 3 days, 10% of injected dose per gram decays in tumor and 80% of radiation from decays in tumors is absorbed in tumors and 0.15 MeV radiation energy per decay. For ^{212}Pb the effective half-life was assumed to be 10.6 h, 10% of injected dose per gram decays in tumor and 100% of radiation in a tumor is absorbed in the tumor, 100% retention of ^{212}Pb and daughters in the tumors and 8 MeV radiation energy per decay. The tumor dosimetry for the injected activities gave on average 35.9 Gy to tumors for ^{177}Lu -PSMA-617 and 2.06 Gy to tumors for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 groups respectively.

[0236] Results: At day 30 after treatment the following the data showed 0%, 12.5% and 75% survival in the saline, ^{177}Lu -PSMA-617 and ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 groups respectively (Figure 2). The median survival was 15 days, 20 days and >30 days (not reached) in the saline, ^{177}Lu -PSMA-617 and ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 groups, respectively.

[0237] In conclusion, the data indicates a strong tumor growth delay with ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 vs ^{177}Lu -PSMA-617 even though the radiation dose estimates show a 17 times higher radiation dose, in terms of Gy, delivered with ^{177}Lu -PSMA-617. Thus, the radiobiological effectiveness (RBE) for ^{212}Pb -p-SCN-Bn-TCMC-PSMA ligand 1 vs ^{177}Lu -PSMA-617 was at least 17. This high radiobiological effectiveness for therapeutic levels of alpha emitter was highly unexpected as usually an RBE of 2-5 is expected for alpha emitters vs beta emitters.

References

[0238]

An efficient chelator for complexation of thorium-227. Ramdahl T, Bonge-Hansen HT, Ryan OB, Larsen S, Herstad G, Sandberg M, Bjerke RM, Grant D, Brevik EM, Cuthbertson AS. *Bioorg Med Chem Lett*. 2016 Sep 1; 26(17):4318-21.

In Vitro and In Vivo Efficacy of a Novel CD33-Targeted Thorium-227 Conjugate for the Treatment of Acute Myeloid Leukemia. Hagemann UB, Wickstroem K, Wang E, Shea AO, Sponheim K, Karlsson J, Bjerke RM, Ryan OB, Cuthbertson AS. *Mol Cancer Ther*. 2016 Oct; 15(10):2422-2431.

Chittenden SJ., A phase 1, open-label study of the biodistribution, pharmacokinetics, and dosimetry of ^{223}Ra -dichloride in patients with hormone-refractory prostate cancer and skeletal metastases. *J Nucl Med* 56: 1304-1309 (2015).

Kratochwil C et al., Targeted α -therapy of metastatic castration resistant prostate cancer with ^{225}Ac -PSMA-617: Dosimetry estimate and empirical dose finding *J Nucl Med* 58: 1624-1631 (2017).

Kratochwil C et al., Targeted α -therapy of mCRPC: Dosimetry estimate of ^{213}Bi -PSMA-617. *Eur J Nucl Med Mol Imaging* 45:31-37 (2018)


Huang SS, Heston WDW. Should Low Molecular Weight PSMA Targeted Ligands Get Bigger and Use Albumin Ligands for PSMA Targeting? *Theranostics* 7: (7) 1940-1941 (2017)

Choy CJ et al., ^{177}Lu -Labeled Phosphoramidate-Based PSMA Inhibitors: The Effect of an Albumin Binder on Biodistribution and Therapeutic Efficacy in Prostate Tumor-Bearing Mice. *Theranostics*. 2017;7(7):1928-1939. doi:10.7150/thno.18719.

Lassmann, Met al., Therapy of ankylosing spondylitis with ^{224}Ra -radium chloride: Dosimetry and risk considerations. Radiat Environ Biophys 41: 173-178 (2002).

Claims

1. A complex comprising a compound complexed with ^{212}Pb wherein the compound comprises a PSMA unit and chelating unit, wherein the PSMA unit is carbon-backbone linked to the chelating unit, and wherein the compound is of the formula:




p-SCN-Bn-TCMC-PSMA ligand 1.

2. A pharmaceutical composition comprising the complex according to claim 1, and a diluent, carrier, surfactant, and/or excipient.
3. The pharmaceutical composition according to claim 2, further comprising ^{224}Ra .
4. The pharmaceutical composition according to claim 3, wherein the activity ratio (MBq) between ^{212}Pb to ^{224}Ra is between 0.5 and 2, such as 0.8 - 1.5, or such as 0.8 - 1.3, or preferably such as 0.9 - 1.15.
5. The pharmaceutical composition according to any one of claims 2-4, wherein the composition is dosaged with a radioactivity of 100 kBq to 100 MBq per dose.
6. The pharmaceutical composition according to claim 2, wherein the composition is dosaged with a radioactivity of 1 MBq to 500 MBq per patient.
7. The pharmaceutical composition according to any one of claims 2-6, wherein the pharmaceutical composition is a solution having a volume of 500 μL to 100 mL.
8. The pharmaceutical composition according to any one of claims 2-7 for use as a medicament.
9. The pharmaceutical composition according to any one of claims 2-7, for use in the treatment of PSMA-expressing disease including soft tissue- and skeletal disease; wherein the skeletal disease is selected from the group consisting of skeletal metastases from cancers to the breast, prostate, kidneys, lung, bone, or multiple myeloma.
10. The pharmaceutical composition according to any one of claims 2-7, for use as a medicament in the treatment of cancer.
11. The pharmaceutical composition for use according to claim 10, wherein the cancer is prostate cancer.
12. The pharmaceutical composition for use according to claim 10, wherein the cancer is PSMA-expressing prostate cancer.
13. The pharmaceutical composition for use according to claim 10, wherein the cancer is advanced metastatic prostate cancer.

14. The pharmaceutical composition for use according to any one of claims 8-13, wherein the composition is administered at a dose in the range 50-150 kBq per kg of bodyweight.

Patentansprüche

1. Komplex, der eine mit ^{212}Pb komplexierte Verbindung umfasst, wobei die Verbindung eine PSMA-Einheit und eine chelatierende Einheit umfasst, wobei die PSMA-Einheit ein an die chelatierende Einheit gebundenes Kohlenstoff-Gerüst ist und wobei die Verbindung der folgenden Formel entspricht:



p-SCN-Bn-TCMC-PSMA-Ligand 1.


2. Pharmazeutische Zusammensetzung, die einen Komplex nach Anspruch 1 und ein Verdünnungsmittel, einen Träger, ein Tensid und/oder einen Hilfsstoff umfasst.
3. Pharmazeutische Zusammensetzung nach Anspruch 2, die weiters ^{224}Ra umfasst.
4. Pharmazeutische Zusammensetzung nach Anspruch 3, wobei das Aktivitätsverhältnis (MBq) zwischen ^{212}Pb und ^{224}Ra zwischen 0,5 und 2 beträgt, wie z.B. 0,8-1,5 oder z.B. 0,8-1,3 oder vorzugsweise z.B. 0,9-1,15.
5. Pharmazeutische Zusammensetzung nach einem der Ansprüche 2 bis 4, wobei die Zusammensetzung mit einer Radioaktivität von 100 kBq bis 100 MBq pro Dosis dosiert ist.
6. Pharmazeutische Zusammensetzung nach Anspruch 2, wobei die Zusammensetzung mit einer Radioaktivität von 1 MBq bis 500 MBq pro Patient dosiert ist.
7. Pharmazeutische Zusammensetzung nach einem der Ansprüche 2 bis 6, wobei die pharmazeutische Zusammensetzung eine Lösung mit einem Volumen von 500 μl bis 100 ml ist.
8. Pharmazeutische Zusammensetzung nach einem der Ansprüche 2 bis 7 zur Verwendung als Medikament.
9. Pharmazeutische Lösung nach einem der Ansprüche 2 bis 7 zur Verwendung bei der Behandlung einer PSMA exprimierenden Erkrankung, einschließlich von Weichgewebe- und Skeletterkrankungen; wobei die Skeletterkrankung aus der aus Skelettmetastasen bei Brust-, Prostata-, Nieren-, Lungen-, Knochenkrebs oder multiplem Myelom bestehenden Gruppe ausgewählt ist.
10. Pharmazeutische Zusammensetzung nach einem der Ansprüche 2 bis 7 zur Verwendung als Medikament zur Behandlung von Krebs.
11. Pharmazeutische Zusammensetzung zur Verwendung nach Anspruch 10, wobei es sich bei dem Krebs um Prostatakrebs handelt.
12. Pharmazeutische Zusammensetzung zur Verwendung nach Anspruch 10, wobei es sich bei dem Krebs um PSMA exprimierenden Prostatakrebs handelt.
13. Pharmazeutische Zusammensetzung zur Verwendung nach Anspruch 10, wobei es sich bei dem Krebs um fortge-

schrittenen metastasierenden Prostatakrebs handelt.

14. Pharmazeutische Zusammensetzung zur Verwendung nach einem der Ansprüche 8 bis 13, wobei die Zusammensetzung in einer Dosis im Bereich von 50 bis 150 kBq pro kg Körpergewicht verabreicht wird.

Revendications

1. Complexe comprenant un composé complexé avec du ^{212}Pb , dans lequel le composé comprend une unité de PSMA et une unité chélatante, dans lequel l'unité de PSMA est un squelette carboné lié à l'unité chélatante, et dans lequel le composé répond à la formule :



p-SCN-Bn-TCMC-PSMA ligand 1.

2. Composition pharmaceutique comprenant le complexe selon la revendication 1, et un diluant, un support, un tensioactif et/ou un excipient.
3. Composition pharmaceutique selon la revendication 2, comprenant en outre du ^{224}Ra .
4. Composition pharmaceutique selon la revendication 3, dans laquelle le rapport d'activité (MBq) entre ^{212}Pb et ^{224}Ra est compris entre 0,5 et 2, comme 0,8 à 1,5, ou comme 0,8 à 1,3, ou de préférence comme 0,9 à 1,15.
5. Composition pharmaceutique selon l'une quelconque des revendications 2 à 4, dans laquelle la composition est dosée avec une radioactivité de 100 kBq à 100 MBq par dose.
6. Composition pharmaceutique selon la revendication 2, dans laquelle la composition est dosée avec une radioactivité de 1 MBq à 500 MBq par patient.
7. Composition pharmaceutique selon l'une quelconque des revendications 2 à 6, dans laquelle la composition pharmaceutique est une solution présentant un volume de 500 μL à 100 mL.
8. Composition pharmaceutique selon l'une quelconque des revendications 2 à 7, à utiliser comme médicament.
9. Composition pharmaceutique selon l'une quelconque des revendications 2 à 7, à utiliser dans le traitement d'une maladie exprimant le PSMA, y compris une maladie des tissus mous et du squelette ; dans laquelle la maladie du squelette est choisie dans le groupe comprenant des métastases de squelette provenant de cancers du sein, de la prostate, des reins, des poumons, des os, ou d'un myélome multiple.
10. Composition pharmaceutique selon l'une quelconque des revendications 2 à 7, à utiliser comme médicament dans le traitement du cancer.
11. Composition pharmaceutique à utiliser selon la revendication 10, dans laquelle le cancer est le cancer de la prostate.
12. Composition pharmaceutique à utiliser selon la revendication 10, dans laquelle le cancer est un cancer de la prostate exprimant le PSMA.

EP 3 498 308 B1

13. Composition pharmaceutique à utiliser selon la revendication 10, dans laquelle le cancer est un cancer de la prostate métastatique avancé.
14. Composition pharmaceutique à utiliser selon l'une quelconque des revendications 8 à 13, dans laquelle la composition est administrée à une dose comprise entre 50 et 150 kBq par kg de poids de corps.

5

10

15

20

25

30

35

40

45

50

55

Figure 1





Figure 2



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **KRATOCHWIL et al.** *Journal of nuclear medicine*, 13 April 2017, vol. 58, 1624-1631 [0007]
- **SATHEKGE et al.** *European Journal of Nuclear medicine and molecular imaging*, 02 March 2017, vol. 44, 1099-1100 [0008]
- **KELLY et al.** *Nuclear medicine and biology*, 01 December 2017, vol. 55, 38-46 [0009]
- **RAMDAHL T ; BONGE-HANSEN HT ; RYAN OB ; LARSEN S ; HERSTAD G ; SANDBERG M ; BJERKE RM ; GRANT D ; BREVIK EM ; CUTHBERTSON AS.** An efficient chelator for complexation of thorium-227. *Bioorg Med Chem Lett.*, 01 September 2016, vol. 26 (17), 4318-21 [0238]
- **HAGEMANN UB ; WICKSTROEM K ; WANG E ; SHEA AO ; SPONHEIM K ; KARLSSON J ; BJERKE RM ; RYAN OB ; CUTHBERTSON AS.** In Vitro and In Vivo Efficacy of a Novel CD33-Targeted Thorium-227 Conjugate for the Treatment of Acute Myeloid Leukemia. *Mol Cancer Ther.*, October 2016, vol. 15 (10), 2422-2431 [0238]
- **CHITTENDEN SJ.** A phase 1, open-label study of the biodistribution, pharmacokinetics, and dosimetry of ²²³Ra -dichloride in patients with hormone-refractory prostate cancer and skeletal metastases. *J Nucl Med*, 2015, vol. 56, 1304-1309 [0238]
- **KRATOCHWIL C et al.** Targeted α -therapy of metastatic castration resistant prostate cancer with ²²⁵Ac-PSMA-617: Dosimetry estimate and empirical dose finding. *J Nucl Med*, 2017, vol. 58, 1624-1631 [0238]
- **KRATOCHWIL C et al.** Targeted α -therapy of mCRPC: Dosimetry estimate of ²¹³Bismuth-PSMA-617. *Eur J Nucl Med Mol Imaging*, 2018, vol. 45, 31-37 [0238]
- **HUANG SS ; HESTON WDW.** Should Low Molecular Weight PSMA Targeted Ligands Get Bigger and Use Albumin Ligands for PSMA Targeting. *Theranostics*, 2017, vol. 7 (7), 1940-1941 [0238]
- **CHOY CJ et al.** ¹⁷⁷Lu-Labeled Phosphoramidate-Based PSMA Inhibitors: The Effect of an Albumin Binder on Biodistribution and Therapeutic Efficacy in Prostate Tumor-Bearing Mice. *Theranostics.*, 2017, vol. 7 (7), 1928-1939 [0238]
- **LASSMANN, M et al.** Therapy of ankylosing spondylitis with ²²⁴Ra-radium chloride: Dosimetry and risk considerations. *Radiat Environ Biophys*, 2002, vol. 41, 173-178 [0238]