

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2012338571 B2

(54) Title
Polymyxin derivatives

(51) International Patent Classification(s)
C07K 7/62 (2006.01) **A61K 38/12** (2006.01)

(21) Application No: **2012338571** (22) Date of Filing: **2012.11.16**

(87) WIPO No: **WO13/072695**

(30) Priority Data

(31) Number
61/561,361 (32) Date
2011.11.18 (33) Country
US

(43) Publication Date: **2013.05.23**
(44) Accepted Journal Date: **2016.06.16**

(71) Applicant(s)
Novacta Biosystems Limited;Cantab Anti-Infectives Limited

(72) Inventor(s)
Saadi, Mona;Duperchy, Esther;Brown, Pamela;Dawson, Michael John;Wadman, Sjoerd Nicolaas

(74) Agent / Attorney
FB Rice, Level 23 44 Market Street, Sydney, NSW, 2000

(56) Related Art
WO 2008/017734 A1

**(43) International Publication Date
23 May 2013 (23.05.2013)**

(51) International Patent Classification:

(21) International Application Number: PCT/GB2012/052844

(22) International Filing Date:
16 November 2012 (16.11.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 61/561,361 18 November 2011 (18.11.2011) US

(71) **Applicants:** NOVACTA BIOSYSTEMS LIMITED [GB/GB]; Biopark Hertfordshire, Broadwater Road, Welwyn Garden City, Hertfordshire AL7 3AX (GB). CAN-TAB ANTI-INFECTIVES LIMITED [GB/GB]; 4th Floor, Reading Bridge House, George Street, Reading, Berkshire RG1 8LS (GB).

(72) **Inventors:** **SAADI, Mona**; Cantab Biopharmaceuticals Limited, Biopark Hertfordshire, Broadwater Road, Welwyn Garden City, Hertfordshire AL7 3AX (GB). **DUPERCHY, Esther**; Cantab Biopharmaceuticals Limited, Biopark Hertfordshire, Broadwater Road, Welwyn Garden City, Hertfordshire AL7 3AX (GB). **BROWN, Pamela**; Cantab Biopharmaceuticals Limited, Biopark Hertfordshire, Broadwater Road, Welwyn Garden City, Hertfordshire AL7 3AX (GB). **DAWSON, Michael John**; Cantab Biopharmaceut-

icals Limited, Biopark Hertfordshire, Broadwater Road, Welwyn Garden City, Hertfordshire AL7 3AX (GB). **WADMAN, Sjoerd Nicolaas**; Rebgasse 4, 79618 Rheinfelden (DE).


(74) **Agents:** **WILLS, Jonathan** et al.; Mewburn Ellis LLP, 33 Gutter Lane, London, Greater London EC2V 8AS (GB).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DF, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: POLYMYXIN DERIVATIVES



(57) Abstract: Provided are compounds, the use of the said compounds in treatment, for example treatment of microbial infections, particularly by Gram negative bacteria. The compounds are polymyxin-based and are represented by the formula (I): and pharmaceutically acceptable salts thereof, where X is $-\text{NHC}(\text{O})-$, $-\text{C}(\text{O})-$, $-\text{OC}(\text{O})-$, $-\text{CH}_2-$ or $-\text{SO}_2-$; R^5 represents C_{10-12} alkyl(C_{4-6} heterocyclyl), or C_{2-12} , $\text{>} \text{alkyl}$ or C_{0-12} alkyl(C_{3-8} cycloalkyl), and the alkyl or cycloalkyl bears one, two or three hydroxyl groups, or a $-\text{NF}^6\text{R}^7$ group, or one $-\text{NR}^8\text{R}^7$ group and one or two hydroxyl groups; and R^1 to R^4 and R^6 to R^8 are as defined in the description.



Declarations under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

Published:

— *with international search report (Art. 21(3))*

POLYMYXIN DERIVATIVES

The present case claims the priority and benefit of US 61/561361 filed on 18 November 2011 (18/11/2011), the contents of which are incorporated by reference herein in their entirety.

The present disclosure relates to novel compounds, pharmaceutical compositions comprising 5 said compounds and the use of the said compounds and pharmaceutical compositions for treatment, for example treatment of microbial infections, particularly by Gram negative bacteria.

In susceptible individuals, certain Gram negative bacteria can cause serious complications and infections, such as pneumonia, urinary tract infections, wound infections, ear infections, 10 eye infections, intra-abdominal infections, oral bacterial overgrowth and sepsis. The treatment of serious bacterial infections in clinical practice can be complicated by antibiotic resistance. Recent years have seen a rise in infections by Gram negative bacteria which are resistant to many types of antimicrobials including broad spectrum antibiotics such as aminoglycosides, 15 cephalosporins and even carbapenems. There is therefore a need to identify new antimicrobials that are effective against Gram negative bacteria, in particular against multidrug resistant Gram negative bacteria.


Polymyxins are a class of antibiotics produced by the Gram positive bacterium *Bacillus polymyxa*. First identified in the late 1940s, polymyxins, particularly polymyxin B and polymyxin E (colistin) were used in the treatment of Gram negative infections. However, these 20 antibiotics exhibited side effects such as nephrotoxicity. Consequently, their use in therapy is limited to treatment of last resort.

WO 2008/017734 tries to address this toxicity problem by providing polymyxin derivatives carrying at least two but no more than three positive charges. These compounds are said to be effective antibacterial agents with reduced toxicity. It is hypothesised in the disclosure that 25 the reduced number of positive charges decreases the affinity of the compound for isolated rat kidney tissue which in turn may lead to a reduction in nephrotoxicity.

Surprisingly, the present inventors have found that certain alternative polymyxin type compounds including some with 4 or more charges have suitable antibacterial activity whilst also apparently exhibiting less toxicity, especially nephrotoxicity.

Summary of the Invention

Thus there is provided a compound of the formula (I):



wherein:

- X represents an $-\text{NHC(O)-}$, $-\text{C(O)-}$, $-\text{OC(O)-}$, $-\text{CH}_2-$ or $-\text{SO}_2-$; and
- 5 R^1 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a phenylalanine, leucine or valine residue;
- R^2 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a leucine, iso-leucine, phenylalanine, threonine, valine or nor-valine residue;
- 10 R^3 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a threonine or leucine residue;
- R^4 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents α,γ -diaminobutyric acid or a serine residue;
- R^5 represents
- 15 C_{0-12} alkyl(C_{4-6} heterocyclyl), or
 C_{2-12} alkyl or C_{0-12} alkyl(C_{3-8} cycloalkyl) wherein the alkyl or cycloalkyl bears:
 - i) one, two or three hydroxyl groups, or
 - ii) a $-\text{NR}^6\text{R}^7$ group, or
 - iii) one $-\text{NR}^6\text{R}^7$ group and one or two hydroxyl groups;
- 20 R^6 represents hydrogen or C_{1-4} alkyl; and
- R^7 represents hydrogen or C_{1-4} alkyl,
- R^8 represents hydrogen or methyl, or
 a pharmaceutically acceptable salt thereof.

The compounds of formula (I) are characterised in that the peptide part of the molecule contains only nine amino acids whereas natural polymyxins comprise 10 amino acids.

Detailed Description

Surprisingly the compounds of formula (I) seem to have lower toxicity than the parent 5 polymyxin compounds whilst retaining useful antibacterial activity.

It is known that polymyxin nonapeptide missing the acyl chain has reduced toxicity but lacks useful antibacterial activity. However, a study of chain lengths of simple acyl polymyxin B nonapeptide derivatives (K.Okimura *et. al*, *Bull. Chem. Soc. Jpn.*, 2007, **80**, 543) suggested the 10 importance of chain length for antibacterial activity, with an optimum of around eight carbon atoms, and demonstrated that the acetyl derivative had very poor activity against *E.coli* and *Salmonella typhimurium*. This was consistent with conclusions from the acyl decapeptide series (P.C de Visser *et al*, *J. Peptide Res.*, 2003, **61**, 298), where the pentanoyl and butanoyl 15 analogues showed a marked drop-off in activity.

We have surprisingly found good antibacterial activity together with reduced toxicity in 15 polymyxin B nonapeptides according to the invention, including those substituted acyl nonapeptides with short acyl chains, especially those bearing an amino substituent.

It is suspected that the toxicity of polymyxin type compounds results from a detergent-like 20 interaction with membranes of eukaryotic cells. In addition, nephrotoxicity of polymyxin type compounds may result from the fact that they are retained in kidney cells and thus accumulate rather than being excreted from the body. Whilst not wishing to be bound by theory it is 25 hypothesised that the compounds of the present invention have a group R⁵ which comprises a substituent which disrupts the hydrophobicity of the alkyl component thereof. The inventors believe that this disruption changes the balance of hydrophobic and hydrophilic nature of the molecules which means they are less well suited for aligning themselves in bilipid-layers which form membranes. In turn this inability to align in the membrane may result in lower residency time therein and thus may result in lower toxicity.

Polymyxin nonapeptide as employed herein is intended to refer to amino acids 2-10 of polymyxin B or polymyxin E.

An amino acid residue (for example a leucine residue, etc.) as employed herein is intended to 30 refer to an amino acid that has lost a water molecule and forms a bond with another entity (such as another amino acid) through the carbonyl end thereof and also forms a bond through the nitrogen end thereof to another entity (such as another amino acid). The bonds formed may for example be amide bonds.

Alkyl as used herein refers to straight chain or branched chain alkyl, such as, without limitation, methyl, ethyl, n-propyl, iso-propyl, butyl, n-butyl and tert-butyl. In one embodiment alkyl refers to straight chain alkyl.

Alkyl in the context of a linker molecule (i.e. substitute alkyl) clearly extends to alkylene fragments, including branched and straight chain versions thereof. Branches may terminate in alkyl radical such as -CH₃.

Heterocyclyl as employed herein is a saturated carbocyclic ring comprising at least one nitrogen ring atom, for example 1 or 2 nitrogen ring atoms, such as only 1 nitrogen ring atom and optionally containing a further ring heteroatom selected from oxygen and sulfur.

10 Examples of C₄₋₆ heterocyclyl groups include azetidine, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl. In one embodiment the heterocyclyl is linked to the remainder of the molecule through nitrogen. In the term "C₄₋₆ heterocyclyl", the expression C₄₋₆ represents the total number of ring atoms, including carbon and heteroatoms.

15 In one embodiment R¹ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents a phenylalanine residue, for example a D-phenylalanine or a leucine residue, such as a D-leucine residue.

In one embodiment R² together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents a leucine residue.

20 In one embodiment R³ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents a threonine residue.

In one embodiment R⁴ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents α,γ -diaminobutyric acid (Dab) or a serine residue, for example L-Dab or D-Ser.

In one embodiment X represents -C(=O).

25 In one embodiment R⁵ represents azetidine, pyrrolidinyl or piperidinyl.

In one embodiment the R⁵ C₂₋₁₂ alkyl component is C₂ alkyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, C₆ alkyl, C₇ alkyl, C₈ alkyl, C₉ alkyl, C₁₀ alkyl, C₁₁ alkyl or C₁₂ alkyl.

In one embodiment R⁵ C₂₋₁₂ alkyl component is C₃₋₁₀ alkyl, for example C₄₋₈ alkyl.

In one embodiment R⁶ is C₃₋₈ cycloalkyl, for example C₅ cycloalkyl or C₆ cycloalkyl.

30 In one embodiment R⁶ bears one substituent.

In one embodiment R^5 bears two substituents.

In one embodiment R^5 bears three substituents.

In one embodiment R^5 bears one, two or three hydroxyl groups, for example one hydroxyl group.

5 In one embodiment R^5 bears one amine group, for example a C_{2-12} alkyl bearing one amine, such as C_{2-4} alkyl bearing one amine.

In one embodiment R^5 bears one, two or three hydroxyl groups, such as one hydroxyl.

In one embodiment R^5 bears one amine group and one hydroxyl group.

In one embodiment R^5 bears one amine group and two hydroxyl groups.

10 In one embodiment wherein R^5 bears one or more hydroxyls then the alkyl chain is C_{5-12} .

In one embodiment R^5 does not bear more than one amine group.

In one embodiment wherein R^5 bears more than one substituent, the substituents are not located on the same carbon atom.

15 In one embodiment at least one R^5 substituent (such as one substituent) is located on C_2 alkyl, C_3 alkyl, C_4 alkyl, C_5 alkyl, C_6 alkyl, C_7 alkyl, C_8 alkyl, C_9 alkyl, C_{10} alkyl, C_{11} alkyl or C_{12} alkyl.

In one embodiment at least one R^5 substituent (such as one substituent) is on a terminal carbon of a straight alkyl chain or an alkyl branch, for example a straight alkyl chain.

20 When the substituent is on the terminal carbon of a straight alkyl chain (or indeed the terminal carbon of a branch) the remaining part of the alkyl chain (or indeed the alkyl linking part of the branch) will form an alkylene link. Thus the term alkyl as used herein is in fact a generic term which covers the situation wherein part or all of the alkyl moiety is in fact an alkylene moiety.

Terminal carbon as employed herein is intended to refer to carbon that would be a $-CH_3$ if it bore no substituents.

25 In one embodiment at least one (such as only one) substituent is not on a terminal carbon, i.e. $-CH(\text{substituent})-$.

In one embodiment R^6 is hydrogen.

In one embodiment R^6 is C_{1-4} alkyl, such as C_1 alkyl, C_2 alkyl, C_3 alkyl or C_4 alkyl, for example methyl.

In one embodiment R⁷ is hydrogen.

In one embodiment R⁷ is C₁₋₄ alkyl such as C₁ alkyl, C₂ alkyl, C₃ alkyl or C₄ alkyl, for example methyl.

In one embodiment both R⁶ and R⁷ represent methyl.


5 In one embodiment R⁶ represents H and R⁷ represents methyl.

In one embodiment R⁵ is selected from -CH(OH)(CH₂)₅CH₃, -CH₂NH₂, -CH₂CH₂NH₂, -CH₂CH₂CH₂NH₂, -(CH₂)₅NH₂, -(CH₂)₇NH₂, -CH₂CH₂NHCH₃, -CH₂CH₂N(CH₃)₂, and -(CH₂)₇OH.

In one embodiment R⁸ is methyl.

10 In one embodiment R⁸ is hydrogen.

In one embodiment, the compound is of formula (Ia):



or a pharmaceutically acceptable salt thereof.

Where R¹ (together with associated groups) represents phenylalanine, R² (together with associated groups) represents leucine, R³ (together with associated groups) represents threonine, R⁴ (together with associated groups) represents α,γ-diaminobutyric acid; and R⁸ represents methyl (and together with the associated groups represents threonine), the compound of formula (Ia) is a polymyxin nonapeptide having amino acids 2-10 of polymyxin B.

Where R¹ (together with associated groups) represents leucine, R² (together with associated groups) represents leucine, R³ (together with associated groups) represents threonine, R⁴ (together with associated groups) represents α,γ -diaminobutyric acid; and R⁸ represents methyl (and together with the associated groups represents threonine), the compound of formula (Ia) is a polymyxin nonapeptide having amino acids 2-10 of polymyxin E.

5 In one embodiment a compound of formula (I) has three positive charges.

In one embodiment a compound of formula (I) has four or five positive charges, such as four.

In one embodiment a compound of formula (I) has five positive charges.

In one embodiment a compound of formula (I) has six positive charges.

10 In one embodiment the compound is selected from:

2-Hydroxyoctanoyl polymyxin B nonapeptide;

2-Aminoethanoyl polymyxin B nonapeptide;

3-Aminopropanoyl polymyxin B nonapeptide;

3-(N,N-dimethylamino)-propanoyl polymyxin B nonapeptide;

15 4-Aminobutanoyl polymyxin B nonapeptide;

6-Aminohexanoyl polymyxin B nonapeptide;

8-Hydroxyoctanoyl polymyxin B nonapeptide;

8-Aminoctanoyl polymyxin B nonapeptide;

3-(N-methylamino) propanoyl polymyxin B nonapeptide;

20 2-Amino cyclopentane carbonyl polymyxin B nonapeptide;

3-Aminopropanoyl colistin (polymyxin E) nonapeptide;

3-Pyrrolidine-3-carbonyl polymyxin B nonapeptide;

3-Amino-3-cyclohexanepropanoyl polymyxin B nonapeptide, or
a pharmaceutically acceptable salt thereof.

25

Additionally or alternatively, the compound is selected from:

5-Aminopentanoyl polymyxin B nonapeptide

Hydroxyacetyl polymyxin B nonapeptide

3-Hydroxyoctanoyl polymyxin B nonapeptide

30 4-(N,N-dimethylamino)-butanoyl polymyxin B nonapeptide

7-Aminoheptanoyl polymyxin B nonapeptide

4-Morpholinylbutanoyl polymyxin B nonapeptide

6-Hydroxyhexanoyl polymyxin B nonapeptide

3-Hydroxybutanoyl polymyxin B nonapeptide
4-(*N*-methylamino)-butanoyl polymyxin B nonapeptide,
trans-4-aminocyclohexanecarbonyl polymyxin B nonapeptide,
4-Aminobutanoyl polymyxin E nonapeptide,
5 2-Hydroxyoctanoyl polymyxin E nonapeptide,
cis-4-Aminocyclohexane carbonyl polymyxin B nonapeptide,
4-Amino-4-methyl pentanoyl polymyxin B nonapeptide
4-Amino-5-methylhexanoyl polymyxin B nonapeptide, including for example 4-(*R*)-Amino-
5-methylhexanoyl polymyxin B nonapeptide
10 3-(1-Pyrrolidin-2-yl)-propionyl polymyxin B nonapeptide, including for example
3-(*S*)-(1-Pyrrolidin-2-yl)-propionyl polymyxin B nonapeptide
4-Aminopentanoyl polymyxin B nonapeptide, including for example 4-(*S*)-Aminopentanoyl
polymyxin B nonapeptide
trans-4-Hydroxycyclohexanecarbonyl polymyxin B nonapeptide,
15 3-Hydroxypropanoyl polymyxin B nonapeptide
(2-Hydroxy-2-cyclohexyl)ethanoyl polymyxin B nonapeptide
2-Amino octanoyl polymyxin B nonapeptide, or
a pharmaceutically acceptable salt thereof.

20 Examples of salts of compound of formula (I) include all pharmaceutically acceptable salts, such as, without limitation, acid addition salts of strong mineral acids such as HCl and HBr salts and addition salts of strong organic acids such as a methanesulfonic acid salt. Further examples of salts include sulphates and acetates such as trifluoroacetate or trichloroacetate.
In one embodiment the compounds of the present disclosure are provided as a sulphate salt.

25 A compound of the disclosure can also be formulated as prodrug. Prodrugs can include an antibacterial compound herein described in which one or more amino groups are protected with a group which can be cleaved *in vivo*, to liberate the biologically active compound. In one embodiment the prodrug is an "amine prodrug". Examples of amine prodrugs include sulphomethyl, as described in e.g., Bergen *et al*, *Antimicrob. Agents and Chemotherapy*, 2006,
30 50, 1953 or HSO₃-FMOC, as described in e.g. Schechter *et al*, *J. Med Chem* 2002, 45(19)
4264, and salts thereof. Further examples of amine prodrugs are given by Krise and Oliyai in *Biotechnology: Pharmaceutical Aspects*, 2007, 5(2), 101-131.
In one embodiment the compounds of the invention are provided as a prodrug.

35 The disclosure herein extends to solvates of compounds of formula (I). Examples of solvates include hydrates.

The compounds of the disclosure include those where the atom specified is replaced by a naturally occurring or non-naturally occurring isotope. In one embodiment the isotope is a stable isotope. Thus the compounds of the disclosure include, for example deuterium containing compounds and the like.

5 The present invention provides compounds having amino acids 2-10 of polymyxin B, or a variant thereof as described below, wherein the N terminal of the amino acid corresponding to residue 2 in polymyxin B, is modified with a group R^5 -X-. The variables R^5 and X are as defined above. In the compounds of the invention, residue 1 of polymyxin B is absent.

A variant of the compound is a compound in which one or more, for example, from 1 to 5, such 10 as 1, 2, 3 or 4 amino acids are substituted by another amino acid. The amino acid is at a position selected from positions 2, 3, 6, 7 or 10 (referring to the numbering of residues used in polymyxin B). The substitution may be for another amino acid or for a stereoisomer.

At position 2, the variant may have a D-Ser substitution.

At position 3, the variant may have a Ser substitution.

15 At position 6, the variant may have a Leu or Val substitution.

A position 7, the variant may have a Ile, Phe, Thr, Val or Nva (norvaline) substitution.


At position 10, the variant may have a Leu substitution.

A polymyxin E compound may be regarded as a polymyxin B compound having a Leu substitution at position 6.

20 For convenience, the compounds of the invention are represented by the formula (I) where the amino acids at positions 2, 3, 6, 7 or 10 are determined by the nature of the groups R^8 , R^4 , R^1 , R^2 and R^3 respectively. Compounds of the invention, which include the variants described above, are biologically active.

Compounds of formula (I) can be prepared by conventional peptide synthesis, using methods 25 known to those skilled in the art. Suitable methods include solution-phase synthesis such as described by Yamada *et al*, *J. Peptide Res.* **64**, 2004, 43-50, or by solid-phase synthesis such as described by de Visser *et al*, *J. Peptide Res.* **61**, 2003, 298-306, and Vaara *et al*, *Antimicrob. Agents and Chemotherapy*, **52**, 2008, 3229-3236. These methods include a suitable protection strategy, and methods for the cyclisation step. Alternatively, compounds 30 may be prepared from readily available polymyxins, for example by removal of the N-terminal amino acid of the polymyxin (residue 1). Such a method is described herein for the preparation of compounds based on residues 2-10 of polymyxins B and E.

The invention also provides a method of preparing certain compound of formula (I) by reacting a compound of formula (II):



or a protected derivative thereof wherein:


R^1 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a phenylalanine, leucine or valine residue;

R^2 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a leucine, iso-leucine, phenylalanine, threonine, valine or nor-valine residue;

R^3 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a threonine or leucine residue;

R^4 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents α,γ -diaminobutyric acid or a serine residue;

with a compound of formula (III):



or a protected derivative thereof

wherein

R^5 is defined above for compounds of formula (I);

X^1 represents group which after coupling to compounds of formula (II) is converted or can be converted into $-NHC(O)-$, $-C(O)-$, $-OC(O)-$, $-CH_2-$ or $-SO_2-$; and

L represents a leaving group,

m represents 0 or 1, or

a pharmaceutically acceptable salt thereof,

optionally followed by deprotection to provide a compound of formula (I).


Generally compounds of formula (II) will be employed in a form where all the free amines, which are not desired to participate in the proposed reaction, are protected by a suitable

5 protecting group for example *tert*-butyloxycarbonyl (BOC), 9-fluorenylmethoxycarbonyl (FMOC), or other suitable amine protecting group such as those described in "Protective groups in Organic Synthesis" by Theodora W. Green and Peter G. M. Wuts, Wiley, New York, 1999.

10 After the requisite chemical reactions deprotection to provide a compound of formula (I) can be carried out using standard methods such as those described in "Protective groups in Organic Synthesis" by Theodora W. Green and Peter G. M. Wuts, Wiley, New York, 1999.

In compounds of formula (I) wherein X represents $-\text{NHC}(=\text{O})-$, can be synthesized employing


15 a compound of formula (III) which corresponds to an isocyanate such as:



wherein R^5 is defined above.

20 The reaction may be performed in a suitable solvent such as dichloromethane, optionally in the presence of base such as triethylamine or *N*-ethyldiisopropylamine (DIPEA).

Alternatively compounds of formula (I) wherein X represents $-\text{NHC}(=\text{O})-$, can be synthesized employing a compound of Formula (IIIb):



wherein R^5 is defined above,

in the presence of base, as described in Gallon *et al*, *J.Org. Chem.*, 2005, **70**, 6960.

30 In compounds of formula I wherein X represents $-\text{C}(=\text{O})-$, $-\text{OC}(=\text{O})-$, or $-\text{SO}_2-$ can be synthesized employing a compound of formula (III) wherein R^5 is as hereinbefore described, X^1 represents $-\text{C}(=\text{O})-$, $-\text{OC}(=\text{O})-$, or $-\text{SO}_2-$ and L represents a leaving group, for example Cl or Br.


The reaction may be performed in a suitable solvent, such as a polar aprotic solvent such as dichloromethane.

Compounds of formula (I) wherein X represent -C(=O)-, can be prepared employing a
5 compound of formula (IIc):



wherein R^5 is defined above, for example in the presence of a coupling agent such as HATU, (O-(7-azabenzotriazol-1-yl)-N,N,N',N'- tetramethyluronium hexafluorophosphate), HBTU ((2-(1
10 H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), DCC (dicyclohexyl carbodiimide), or PYBOP (benzotriazole-1-yl-oxy-tri-pyrrolidino-phosphonium hexafluorophosphate), under basic conditions in a polar solvent.

Compounds of formula (I) wherein X represents -CH₂- can be prepared employing an
aldehyde of formula (IIId):



wherein R^5 is as hereinbefore described, for example in the presence of a reducing agent such as sodium triacetoxyborohydride, sodium cyanoborohydride, or polymer-supported cyanoborohydride in a solvent such as methanol, dichloromethane, DMF, using conditions such as described in March's Advanced Organic Chemistry, Wiley, 2001.

20 In one aspect, the invention provides a pharmaceutical composition comprising a compound of formula (I), for example a therapeutically effective amount thereof and a pharmaceutically acceptable excipient, diluent and/or carrier (including combinations thereof).

The routes for administration (delivery) include, but are not limited to, one or more of: oral (e.g. as a dry powder/ free flowing particulate formulation, tablet, capsule, or as an ingestible
25 solution or suspension) buccal, sublingual.

The compositions of the disclosure include those in a form especially formulated for parenteral, oral, buccal, rectal, topical, implant, ophthalmic, nasal, rectal or genito-urinary use. In one aspect of the invention, the agents are delivered orally, hence, the agent is in a form that is suitable for oral delivery.

30 In some instances it may be possible to deliver the compounds of the disclosure by a topical, parenteral (e. g. by an injectable form) or transdermal route, including mucosal (e. g. as a nasal spray or aerosol for inhalation), nasal, gastrointestinal, intraspinal, intraperitoneal, intramuscular, intravenous, intrauterine, intraocular, intradermal, intracranial, intratracheal,

intravaginal, intracerebroventricular, intracerebral, subcutaneous, ophthalmic (including intravitreal or intracameral).

There may be different composition/formulation requirements depending on the different delivery systems or different routes of administration. By way of example, the pharmaceutical

5 composition of the present disclosure may be formulated to be delivered using a mini-pump or by a mucosal route, for example, as a nasal spray or aerosol for inhalation or ingestible solution, or parenterally in which the composition is formulated in an injectable form, for delivery by, for example, an intravenous, intramuscular or subcutaneous route. Alternatively, the formulation may be designed to be delivered by both routes. Where appropriate, the

10 pharmaceutical compositions can be administered by inhalation, in the form of a suppository or pessary, topically in the form of a lotion, solution, cream, ointment or dusting powder, by use of a skin patch, orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents, or they can be injected

15 parenterally, for example intravenously, intramuscularly or subcutaneously.

For parenteral administration, the compositions may be best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or saccharides, in particular a monosaccharide, to make the solution isotonic with blood.

Examples of parenteral administration include one or more of: intravenously, intraarterially, 20 intraperitoneally, intrathecally, intraventricularly, intraurethrally, intrasternally, intracranially, intramuscularly or subcutaneously administering the agent, and/or by using infusion techniques.

In one embodiment the formulation of compounds of the invention is provided as a liposomal formulation. Liposomes can vary in size from low micrometer range to tens of micrometers,

25 unilamellar liposomes are typically in the lower size range with various targeting ligands attached to their surface allowing for their surface-attachment and accumulation in pathological areas for treatment of disease. Liposomes are artificially prepared vesicles made of lipid bilayer

In one embodiment the formulation is adapted for delivery by infusion or slow injection.

30 In one embodiment the formulation is adapted for delivery by bolus injection.

For buccal or sublingual administration the compositions may be administered in the form of tablets or lozenges which can be formulated in a conventional manner.

The compounds of the disclosure can be administered (e.g. orally or topically) in the form of tablets, capsules, ovules, elixirs, solutions or suspensions, which may contain flavouring or colouring agents, for immediate-, delayed-, modified-, sustained-, pulsed- or controlled-release applications.

5 The compounds of the disclosure may also be presented for human or veterinary use in a form suitable for oral or buccal administration, for example in the form of solutions, gels, syrups, mouth washes or suspensions, or a dry powder for constitution with water or other suitable vehicle before use, optionally with flavouring and colouring agents.

10 Solid compositions such as tablets, capsules, lozenges, pastilles, pills, powder, pastes, granules, bullets or premix preparations may also be used. Solid and liquid compositions for oral use may be prepared according to methods well known in the art. Such compositions may also contain one or more pharmaceutically acceptable carriers and excipients which may be in solid or liquid form.

15 The tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, calcium sulphate, dibasic calcium phosphate and glycine, mannitol, pregelatinised starch, corn starch, potato starch, disintegrants such as sodium starch glycollate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia.

20 Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

25 Solid compositions of a similar type may also be administered in gelatin or HPMC (hydroxypropyl methylcellulose) capsules. Suitable excipients in this regard include microcrystalline cellulose, lactose, calcium carbonate, calcium sulphate, dibasic calcium phosphate and, mannitol, pregelatinised starch, corn starch, potato starch or high molecular weight polyethylene glycols.

For aqueous suspensions and/or elixirs, the agent may be combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

30 Capsules may be filled with a powder (of medicament alone or as blend with selected filler(s)) or alternatively a liquid, each comprising one or more salts of the present disclosure and optionally a carrier. Where the capsule is filled with a powder the salts of the present

disclosure and/or the carrier may be milled or micronised to provide material with an appropriate particle size.

Alternatively, the tablet or a capsule, as appropriate, may be filled into another capsule (preferably a HPMC capsule such as Capsugel®) to provide either a tablet in capsule or

5 capsule in capsule configuration, which when administered to a patient yields controlled dissolution in the gastrointestinal tract thereby providing a similar effect to an enteric coating.

Thus in one aspect the disclosure provides a solid dose formulation of a salt of the present disclosure, for example where the formulation has an enteric coating.

In another aspect the disclosure provides a solid dose formulation comprising a protective

10 capsule as outer layer, for example as a tablet in a capsule or a capsule in a capsule. The enteric coating may provide an improved stability profile over uncoated formulations.

The compounds of the disclosure may also be administered orally, in veterinary medicine, in the form of a liquid drench such as a solution, suspension or dispersion of the active ingredient together with a pharmaceutically acceptable carrier or excipient.

15 The compounds of the invention may also, for example, be formulated as suppositories e.g. containing conventional suppository bases for use in human or veterinary medicine or as pessaries e. g. containing conventional pessary bases.

In one embodiment the formulation is provided as a formulation for topical administration including inhalation.

20 Suitable inhalable preparations include inhalable powders, metering aerosols containing propellant gases or inhalable solutions free from propellant gases. Inhalable powders according to the disclosure containing the active substance may consist solely of the abovementioned active substances or of a mixture of the abovementioned active substances with physiologically acceptable excipient.

25 These inhalable powders may include monosaccharides (e.g. glucose or arabinose), disaccharides (e.g. lactose, saccharose, maltose), oligo- and polysaccharides (e.g. dextrans), polyalcohols (e.g. sorbitol, mannitol, xylitol), salts (e.g. sodium chloride, calcium carbonate) or mixtures of these with one another. Mono- or disaccharides are preferably used, the use of lactose or glucose, particularly but not exclusively in the form of their hydrates.

30 Particles for deposition in the lung require a particle size less than 10 microns, such as 1-9 microns suitably from 0.1 to 5 µm, particularly preferably from 1 to 5 µm.

The propellant gases which can be used to prepare the inhalable aerosols are known from the prior art. Suitable propellant gases are selected from among hydrocarbons such as n-propane, n-butane or isobutane and halohydrocarbons such as chlorinated and/or fluorinated derivatives of methane, ethane, propane, butane, cyclopropane or cyclobutane. The above-

5 mentioned propellant gases may be used on their own or in mixtures thereof.

Particularly suitable propellant gases are halogenated alkane derivatives selected from among TG11, TG 12, TG 134a and TG227. Of the abovementioned halogenated hydrocarbons, TG134a (1,1,1,2-tetrafluoroethane) and TG227 (1,1,1,2,3,3,3- heptafluoro propane) and mixtures thereof are suitable for use in formulations of the present invention.

10 The propellant-gas-containing inhalable aerosols may also contain other ingredients such as co-solvents, stabilisers, surface-active agents (surfactants), antioxidants, lubricants and means for adjusting the pH. All these ingredients are known in the art.

The propellant-gas-containing inhalable aerosols according to the invention may contain up to 5 % by weight of active substance. Aerosols according to the disclosure may contain, for 15 example, 0.002 to 5 % by weight, 0.01 to 3 % by weight, 0.015 to 2 % by weight, 0.1 to 2 % by weight, 0.5 to 2 % by weight or 0.5 to 1 % by weight of active.

The salts of the disclosure may also be used in combination with other therapeutic agents. The disclosure thus provides, in a further aspect, a combination comprising a salt of the present disclosure together with a further therapeutic agent. The combination may, for example be a 20 combination of a salt of the compound of formula (I) and an antibiotic, such as vancomycin, fosfomycin, rifamycin, a beta-lactam (such as a cephalosporin or carbapenem), an aminoglycoside, a macrolide, a tetracycline, a lipopeptide, an oxazolidinone and/or an anti-inflammatory such as a steroid. The combination may be provided as a co-formulation or simply packaged together as separate formulations, for simultaneous or sequential delivery.

25 In one embodiment there is provided salts of the present disclosure in combination with a further therapeutic agent.

It is to be understood that not all of the compounds/salts of the combination need be administered by the same route. Thus, if the therapy comprises more than one active component, then those components may be administered by different routes.

30 The individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations by any convenient route.

When administration is sequential, either the salt of the disclosure or the second therapeutic agent may be administered first. When administration is simultaneous, the combination may be administered either in the same or a different pharmaceutical composition.

The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above together with a pharmaceutically acceptable carrier or excipient comprise a further aspect of the disclosure.

When combined in the same formulation it will be appreciated that the two compounds/salts must be stable and compatible with each other and the other components of the formulation.

When formulated separately they may be provided in any convenient formulation, in such manner as are known for such compounds in the art.

The compositions may contain from 0.01-99% of the active material. For topical administration, for example, the composition will generally contain from 0.01-10%, more such as 0.01-1 % of the active material.

When a salt of the disclosure is used in combination with a second therapeutic agent active against the same disease state the dose of each compound/salt may be the same or differ from that employed when the compound/salt is used alone. Appropriate doses will be readily appreciated by those skilled in the art. It will also be appreciated that the amount of a salt of the disclosure required for use in treatment will vary with the nature of the condition being treated and the age and the condition of the patient and will be ultimately at the discretion of the attendant physician or veterinarian.

Typically, a physician will determine the actual dosage which will be most suitable for an individual subject. The specific dose level and frequency of dosage for any particular individual may be varied and will depend upon a variety of factors including the activity of the specific salt employed, the metabolic stability and length of action of that salt, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the individual undergoing therapy.

For oral and parenteral administration to humans, the daily dosage level of the agent may be in single or divided doses. For systemic administration the daily dose as employed for adult human treatment will range from 2-100 mg/Kg body weight, such as 5-60 mg/Kg body weight, which may be administered in 1 to 4 daily doses, for example, depending on the route of administration and the condition of the patient. When the composition comprises dosage units, each unit will preferably contain 100 mg to 1 g of active ingredient. The duration of treatment will be dictated by the rate of response rather than by arbitrary numbers of days.

In one embodiment the treatment regime is continued for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 or more days.

As described above, the salts of the present disclosure may be employed in the treatment or prophylaxis of humans and/or animals.

5 There is further provided by the present disclosure a process of preparing a pharmaceutical composition, which process comprises mixing a salt of the disclosure or a pharmaceutically acceptable derivative thereof, together with a pharmaceutically acceptable excipient, diluent and/or carrier.

In another aspect, the invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof or a composition comprising the same for use in therapy, and in 10 particular, in the treatment infection such as bacterial infection, such as a Gram negative bacterial infection.

In one embodiment the compounds and compositions of the disclosure are useful in the treatment of pneumonia, urinary tract infections, wound infections, ear infections, eye 15 infections, intra-abdominal infections, bacterial overgrowth and/or sepsis.

In one embodiment the compounds are useful in the treatment of infections by bacteria which are multidrug resistant.

Examples of Gram negative bacteria include, but are not limited to, *Escherichia* spp., *Klebsiella* spp., *Enterobacter* spp., *Salmonella* spp., *Shigella* spp., *Citrobacter* spp.,

20 *Morganella morganii*, *Yersinia pseudotuberculosis* and other Enterobacteriaceae, *Pseudomonas* spp., *Acinetobacter* spp., *Moraxella*, *Helicobacter*, *Stenotrophomonas*, *Bdellovibrio*, acetic acid bacteria, *Legionella* and alpha-proteobacteria such as *Wolbachia* and numerous others. Other notable groups of Gram-negative bacteria include the cyanobacteria, spirochaetes, green sulfur and green non-sulfur bacteria.

25 Medically relevant Gram-negative cocci include three organisms, which cause a sexually transmitted disease (*Neisseria gonorrhoeae*), a meningitis (*Neisseria meningitidis*), and respiratory symptoms (*Moraxella catarrhalis*).

Medically relevant Gram-negative bacilli include a multitude of species. Some of them primarily cause respiratory problems (*Hemophilus influenzae*, *Klebsiella pneumoniae*,

30 *Legionella pneumophila*, *Pseudomonas aeruginosa*), primarily urinary problems (*Escherichia coli*, *Enterobacter cloacae*), and primarily gastrointestinal problems (*Helicobacter pylori*, *Salmonella enterica*).

Gram-negative bacteria associated with nosocomial infections include *Acinetobacter baumannii*, which causes bacteremia, secondary meningitis, and ventilator-associated pneumonia in intensive-care units of hospital establishments.

In one embodiment the compounds and compositions of the present invention are useful in the treatment of infection of one or more of the following Gram negative bacteria: *E. coli*, *S. enterica*, *Klebsiella*; *K. pneumoniae*, *K. oxytoca*; *Enterobacter*; *E. cloacae*, *E. aerogenes*, *E. agglomerans*, *Acinetobacter*; *A. calcoaceticus*, *A. baumannii*; *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia*, *Providencia stuartii*, *Proteus*; *P. mirabilis*, *P. vulgaris*.

In one embodiment compounds of formula (I) or pharmaceutically acceptable salts thereof or compositions comprising the same are useful for the treatment of *Pseudomonas* infections including *P. aeruginosa* infection, for example skin and soft tissue infections, gastrointestinal infection, urinary tract infection, pneumonia and sepsis.

In one embodiment compounds of formula (I), or pharmaceutically acceptable salts thereof, or compositions comprising the same are useful for the treatment of *Acinetobacter* infections including *A. baumanii* infection, for pneumonia, urinary tract infection and sepsis.

In one embodiment compounds of formula (I), or pharmaceutically acceptable salts thereof, or compositions comprising the same are useful for the treatment of *Klebsiella* infections including *K. pneumoniae* infection, for pneumonia, urinary tract infection, meningitis and sepsis.

In one embodiment compounds of formula (I), or pharmaceutically acceptable salts thereof, or compositions comprising the same are useful for the treatment of *E. coli* infection including *E. coli* infections, for bacteremia, cholecystitis, cholangitis, urinary tract infection, neonatal meningitis and pneumoniae

In one embodiment the compounds of formula (I) or pharmaceutically acceptable salts thereof or compositions comprising the same may be useful for long term treatment.

In one aspect there is provided a compound of formula (I) or a composition comprising the same for the manufacture of a medicament for one or more of the indications defined above.

In one aspect there is provided a method of treatment comprising the step of administering a therapeutically effective amount of a compound of formula (I) or a pharmaceutical acceptable salt thereof or a composition comprising the same to a patient (human or animal) in need thereof, for example for the treatment of an infection as described herein.

Where technically appropriate embodiments may be combined and thus the disclosure extends to all permutations/combinations of the embodiments provided herein.

Preferences given for compounds of formula (I) may equally apply to other compounds of the invention, disclosed herein, as technically appropriate.

Abbreviation	Meaning
PMBN	Polymyxin B nonapeptide
Thr	Threonine
Ser	Serine
DSer	D-serine
Leu	Leucine
Ile	Isoleucine
Phe	Phenylalanine
DPhe	D-phenylalanine
Val	Valine
Dab	α,γ -Diaminobutyric acid
DIPEA	<i>N,N</i> -diisopropylethylamine

5

Examples

Intermediate 1. Polymyxin B nonapeptide

A mixture of Polymyxin B (20 g), immobilised papain (185 ELU/g), potassium phosphate buffer (25 mM; pH 7, 1.25 L), potassium chloride (30 mM), EDTA (10 mM) and cysteine

10 (1 mM) was incubated at 37°C for 18h with gentle agitation. The progress of the reaction was monitored by LC-MS using the conditions outlined in Table 1. The immobilized papain was removed by filtration and the filtrate was concentrated *in vacuo* to leave a solid residue which was re-suspended in 10% aqueous methanol and left at room temperature overnight. The supernatant was decanted and concentrated *in vacuo*. Polymyxin B nonapeptide was purified 15 from the residue by SPE on C18 silica, eluting with 0-10% aqueous methanol. Evaporation of the appropriate fractions gave the product as a white solid. *m/z* 482, [M+2H]²⁺

Table 1. LC-MS conditions

Micromass Platform LC

Column: Zorbax 5 μ C18 (2) 150 x 4.6 mm


20 Mobile Phase A: 10% Acetonitrile in 90% Water, 0.15 %TFA or 0.1% formic

Mobile Phase B: 90% Acetonitrile in 10% Water, 0.15 %TFA or 0.1% formic

Flow rate: 1 mL/min
 Gradient: Time 0 min 100% A 0% B
 Time 10 min 0% A 100% B
 Time 11 min 0% A 100% B
 5 Time 11.2 min 100% A 0% B
 Time 15 min 100% A 0% B
 Cycle time 15 min
 Injection volume: 20 μ L
 Detection: 210 nm

10

Intermediate 2. tetra-(Boc) polymyxin B nonapeptide



15

20 Molecular Weight = 1363.63
 Exact Mass = 1362
 Molecular Formula = C63H106N14O19


Selective BOC protection of the free γ -amino groups on the Dab residues of polymyxin B nonapeptide was carried out using the procedure of H. O'Dowd *et al*, *Tetrahedron Lett.*, 2007, 48, 2003-2005. Polymyxin B Nonapeptide (**Intermediate 1** 7.5 g, 7.78 mmol) was suspended in water (65 mL) with sonication. Dioxane (65 mL) and triethylamine (65 mL) were added and the mixture was cooled in ice for 10 min prior to the addition of 1-(Boc-oxyimino)-2-phenyl acetonitrile (Boc-ON) (7.67 g; 31.15 mmol). The progress of the reaction was followed by LC-MS and reached completion after 30 minutes, whereupon the mixture was quenched by addition of 20% methanolic ammonia (50 mL). The liquid phase was decanted and the residual 25 solid was purified by chromatography (eluent 0-20% methanol in dichloromethane) on silica gel to afford tetra-(Boc) polymyxin B nonapeptide as a white solid (2.5 g, 24 %). TLC, R_f 0.2 (10% methanol in dichloromethane). m/z 1362.8[MH] $^+$.

Intermediate 3. Colistin (Polymyxin E) nonapeptide

Colistin (polymyxin E, 5 g) was treated with immobilised papain (185 ELU/g), potassium phosphate buffer (25 mM; pH 7, 1.25 L), potassium chloride (30 mM), EDTA (10 mM) and 5 cysteine (1 mM) at 37°C for 32 h with gentle agitation to produce colistin (polymyxin E) nonapeptide. The progress of the reaction was monitored by LC-MS using the conditions outlined in Intermediate 1, Table 1. The immobilized papain was removed by filtration and the filtrate was concentrated *in vacuo* to leave a solid residue which was re-suspended in 10% aqueous methanol and left at room temperature overnight. The supernatant was decanted and 10 concentrated *in vacuo*. Colistin (Polymyxin E) nonapeptide was purified from the residue by SPE on C18 silica (10 gm), eluting with 0-25% aqueous methanol. Evaporation of the appropriate fractions gave the product as a white solid. *m/z* 465.32 [M+2H]²⁺.

Intermediate 4 . tetra-(Boc) colistin (polymyxin E) nonapeptide

15




20

Molecular Weight = 1329.61
 Exact Mass = 1328
 Molecular Formula = C₆₀H₁₀₈N₁₄O₁₉

Colistin (Polymyxin E) Nonapeptide (2.5 g, 2.69 mmol) was suspended in water (35 mL) with sonication. Dioxane (35 mL) and triethylamine (35 mL) were added and the mixture was cooled in ice for 10 min prior to the addition of 1-(Boc-oxyimino)-2-phenyl acetonitrile (Boc-ON) 25 (2.65 g; 10.76 mmol). The progress of the reaction was followed by LC-MS and reached completion after 10 minutes, whereupon the mixture was quenched by addition of 20% methanolic ammonia (25 mL). The liquid phase was decanted and the residual solid was re-dissolved in water and extracted sequentially with dichloromethane and iso-butanol. Based on LC-MS analysis, the decanted liquid and both dichloromethane and iso-butanol extracts 30 were pooled together followed by concentration *in vacuo* to give yellow gum which was loaded


onto flash chromatography (Si 60A- 35-70). The column was eluted with 0-20% methanol (containing 2% ammonia) in dichloromethane. The column fractions eluted with 7-10% methanol (containing 2% ammonia) in dichloromethane afforded tetra-(Boc) colistin (polymyxin E) nonapeptide as a white solid (1.18 g, 33 %), *m/z* 1329.7 [M+H]⁺.

5 **Example 1. [2(R,S)-2-Hydroxyoctanoyl] polymyxin B nonapeptide, trifluoroacetate.**



Molecular Weight = 1105.36
Exact Mass = 1104
Molecular Formula = C51H88N14O13

15 **a) [2-(R,S)-2-Hydroxyoctanoyl] [tetra-(Boc)]-polymyxin B nonapeptide**



25
Molecular Weight = 1505.83
Exact Mass = 1504
Molecular Formula = C71H120N14O21


2-Hydroxyoctanoic acid (1.16 g, 7.34 mmol) was dissolved in dichloromethane (2 mL). *N,N*-Dilisopropylethylamine (1.19 mL, 7.34 mmol) and 2-(1*H*-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate (HATU) (2.79 g, 7.34 mmol) were then added to the reaction mixture. After 30 min stirring at room temperature compound of **intermediate 2** (2.0 g, 1.47 mmol) was added. After 16 h the completion of the reaction was confirmed by LC-MS and the reaction mixture was evaporated to dryness and purified using column chromatography on silica gel (eluent 0-10% methanol in dichloromethane). The appropriate

fractions were concentrated to leave [2(R,S)-2-hydroxyoctanoyl] [tetra-(Boc)]-Polymyxin B nonapeptide as a colourless oil (1.28 g, 58 %). TLC, R_f 0.6 (10% MeOH in dichloromethane). m/z 1527.5, $[M+Na]^+$.

5 b) Title compound: [2(R,S)-2-Hydroxyoctanoyl] polymyxin B nonapeptide, trifluoroacetate

2-Hydroxyoctanoyl [tetra-(Boc)]-polymyxin B nonapeptide 1.28 g, 0.85 mmol) was dissolved in dichloromethane (2 mL). Trifluoroacetic acid (3.9 mL, 51.02 mmol) was added and the mixture was stirred at room temperature for 16 h, after which time LC-MS confirmed completion of the 10 reaction. The reaction mixture was concentrated *in vacuo* to leave [2(R,S)-2-hydroxyoctanoyl] Polymyxin B nonapeptide, trifluoroacetate as a colourless oil (1.3 g, 93 %). TLC, R_f baseline (10% MeOH in dichloromethane). m/z 1104.8 $[MH]^+$.

Example 2. [2(R,S)-2-Hydroxyoctanoyl] polymyxin B nonapeptide, sulphate salt.



Molecular Weight = 1105.36
Exact Mass = 1104
Molecular Formula = C51H88N14O13

15

To the compound of Example 1 (1.3 g) was added water (1 mL) and the mixture was sonicated for 5 min. To the resulting suspension was added 1M NaHCO₃ (20 mL) until the mixture reached pH 9. The mixture was then passed through a 10 g C18 SPE column, eluting sequentially with 0, 40, 50, 60, 70, 80 and 100 % aqueous methanol. LC-MS analysis of each 20 fraction showed that the desired product eluted in the 60, 70 and 80% aqueous methanol fractions. These fractions were pooled and evaporated to leave a white solid (0.5 g), to which was added 0.1 M H₂SO₄ (30 mL) until pH 7 was reached. *tert*-Butanol (10 mL) was added and the mixture was stirred for 16 h at room temperature and subsequently freeze-dried to leave [2(R,S)-2-hydroxyoctanoyl] Polymyxin B nonapeptide, sulphate salt as a white solid (0.52 g).


Analysis by HPLC according to the conditions outlined in Table 2 gave a retention time of 5.93 minutes. m/z 1104.9 [MH]⁺.

Table 2. Analytical HPLC conditions

Column:	Zorbax 5 μ C18 (2) 150 x 4.6 mm			
5 Mobile Phase A:	10% Acetonitrile in 90% Water, 0.15 %TFA or 0.1% Formic acid			
Mobile Phase B:	90% Acetonitrile in 10% Water, 0.15 %TFA or 0.1% Formic acid			
Flow rate:	1mL/min			
Gradient:	Time 0 min	100%	A	0% B
	Time 10 min	0%	A	100% B
10	Time 11 min	0%	A	100% B
	Time 11.2 min	100%	A	0% B
	Cycle time 15 min			
Injection volume:	20 μ L			
Detection:	210 nm			

15


Example 3. 2-Aminoethanoyl polymyxin B nonapeptide, sulphate salt.



2-Aminoethanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 2-(tert-butoxycarbonylamino)-ethanoic acid following the


20 sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 4.99 min; m/z 1020.8 [MH]⁺.

Example 4. 3-Aminopropanoyl polymyxin B nonapeptide, sulphate salt.




3-Aminopropanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 3-(tert-butoxycarbonylamino)-propanoic acid following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 4.97 min; *m/z* 1034.42, $[\text{MH}]^+$.

Example 5. 3-(*N,N*-dimethylamino)-propanoyl polymyxin B nonapeptide, sulphate salt




10 3-(*N,N*-dimethylamino)-propanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 3-(*N,N*-dimethylamino)propanoic acid following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 5.01 min; *m/z* 531.92, $[\text{M}+2\text{H}]^{2+}$.


Example 6. 4-Aminobutanoyl polymyxin B nonapeptide, sulphate salt.

4-Aminobutanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 4-(tert-butoxycarbonylamino)-butanoic acid following the

5 sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 4.97min; *m/z* 524.91 [M+2H]²⁺.


Example 7. 6-Aminohexanoyl polymyxin B nonapeptide, sulphate salt.

10 6-Aminohexanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 6-(tert-butoxycarbonylamino)-hexanoic acid following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 4.97 min; *m/z* 1077.15 [MH]⁺.

Example 8. 8-Hydroxyoctanoyl polymyxin B nonapeptide, sulphate salt.

8-Hydroxyoctanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 8-hydroxyoctanoic acid following the sequence of reactions


5 described for Examples 1 and 2. Retention time (HPLC) 5.29 min; *m/z* 1104.87, [M]⁺.

Example 9. 8-Aminoctanoyl polymyxin B nonapeptide, sulphate salt.

8-Aminoctanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 8-(tert-butoxycarbonylamino)-octanoic acid following the

10 sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 5.02 min; *m/z* 1105.2, [MH]⁺.


Example 10. 3-(N-methylamino)propanoyl polymyxin B nonapeptide, sulphate salt.



3-(N-Methylamino)propanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and N-[(1,1-dimethylethoxy)carbonyl]-N-methyl β -Alanine,

5 following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 5.0 min; m/z 525.05, $[M+2H]^{2+}$.

Example 11. (1R,S/2R,S)-2-Aminocyclopentanecarbonyl polymyxin B nonapeptide, sulphate salt




10

(1R,S/2R,S)-2-Aminocyclopentanecarbonyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and *cis*-2-(tert-butoxycarbonylamino)-cyclopentane carboxylic acid following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 5.07 min; m/z 1074.87, $[MH]^{+}$.


15

Example 12. 3-Aminopropanoyl colistin (polymyxin E) nonapeptide, sulphate salt




Following the sequential reactions described for Examples 1 and 2, 3-Aminopropanoyl colistin (polymyxin E) nonapeptide, sulphate salt was prepared from tetra-(Boc) colistin (polymyxin E) nonapeptide (Intermediate 4) and Boc- β -alanine. Retention (HPLC) time of 4.98 minutes. *m/z* 501, $[\text{M}+2\text{H}]^{2+}$.

Example 13. [3-(R,S)-Pyrrolidine-3-carbonyl] polymyxin B nonapeptide, sulphate salt



[3(R,S)-Pyrrolidine-3-carbonyl] polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 3-(N-tert-butoxycarbonyl)-pyrrolidinecarboxylic acid following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 4.91 min; *m/z* 1060.58 $[\text{MH}]^+$.

Example 14. [3(R,S)-3-Amino-3-cyclohexanepropanoyl] Polymyxin B nonapeptide, sulphate salt



[3(R,S)-3-Amino-3-cyclohexanepropanoyl polymyxin B nonapeptide, sulphate salt was prepared from tetra-(Boc) polymyxin B nonapeptide and 3-(tert-butoxycarbonylamino)-3-cyclohexanepropanoic acid following the sequence of reactions described for Examples 1 and 2. Retention time (HPLC) 5.24 min; *m/z* 1116.78, [MH]⁺.

Additional Examples 15-35

Additional compounds of Examples 15-35 were prepared using the methods of preparation set out for Examples 1 and 2 above. Thus, a compound having a substituent at the polymyxin B nonapeptide N terminal was prepared from tetra-(Boc) polymyxin B nonapeptide (intermediate 2) and an appropriate carboxylic acid in the presence of coupling agents (e.g. HATU) and base (e.g. DIPEA) (as set out in Example 1a), followed by treatment with acid (e.g. TFA) (as set out in Example 1a), and an appropriate work up (as set out in Example 2). Similarly, a compound having a substituent at the polymyxin E nonapeptide N terminal was prepared from tetra-(Boc) colistin (polymyxin E) nonapeptide (Intermediate 4) and an appropriate carboxylic acid in the presence of coupling agents (e.g. HATU) and base (e.g. DIPEA) (as set out in Example 1b), followed by treatment with acid (e.g. TFA) (as set out in Example 1b), and conversion to the sulphate salt (as set out in Example 2).



The additional compounds of Examples 15-35 are presented in Table 3 below.

The recorded retention times and masses given in the Table were obtained using the LC-MS conditions described above in Table 2.

The compounds were isolated as the sulphate salt forms of the compounds shown.

Table 3

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
15		5-Aminopentanoyl polymyxin B nonapeptide	C ₄₈ H ₈₃ N ₁₅ O ₁₂ 1061.63	5.09	1062.7 (M ⁺)
16		Hydroxyacetyl polymyxin B nonapeptide	C ₄₅ H ₇₆ N ₁₄ O ₁₃ 1020.57	5.00	1021.1 (M ⁺)

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
17	 <p>3-(R,S)-3-Hydroxyoctanoyl polymyxin B nonapeptide</p>		$C_{51}H_{88}N_{14}O_{13}$ 1104.67	6.04	1126.6 ($M+Na^+$)
18	 <p>4-(N,N-dimethylamino)-butanoyl polymyxin B nonapeptide,</p>		$C_{49}H_{85}N_{15}O_{12}$ 1075.65	4.92	1076 (MH^+)



Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
19		7-Aminoheptanoyl polymyxin B nonapeptide	C ₅₀ H ₈₇ N ₁₅ O ₁₂ 1090.34	4.751	1091.76 (M ⁺)
20		4-Morpholinylbutyroyl polymyxin B nonapeptide	C ₅₁ H ₈₇ N ₁₅ O ₁₃ 1117.66	5.08	1116.9 (M ⁺)

- 35 -



Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
21		6-Hydroxyhexanoyl polymyxin B nonapeptide	C ₄₉ H ₈₄ N ₁₄ O ₁₃ 1076.63	5.01	539 (M+2H) ²⁺
22		3-(R,S)-3-Hydroxybutanoyl polymyxin B nonapeptide	C ₄₇ H ₈₀ N ₁₄ O ₁₃ 1048.60	4.83	525.3 (M+2H) ²⁺

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
23		4-(N-methylamino)-butanoyl polymyxin B nonapeptide	C48H83N15O12 1061.63	4.92	1062.4 (MH ⁺)
24		trans-4-aminocyclohexanecarbonyl polymyxin B nonapeptide	C50H85N15O12 1087.65	4.95	1087.1 (M ⁺)

- 37 -

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
25		4-aminobutanoyl polymyxin E nonapeptide	C ₄₄ H ₈₃ N ₁₅ O ₁₂ 1013.63	4.94	1036.0 (MNa ⁺)
26		2(R,S)-2-Hydroxyoctanoyl polymyxin E nonapeptide	C ₄₈ H ₉₀ N ₁₄ O ₁₃ 1070.68	5.92	1071.3 (MH ⁺)

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
27	<p><i>cis</i>-4-aminocyclohexane carbonyl polymyxin B nonapeptide</p>		$C_{50}H_{85}N_{15}O_{12}$ 1087.65	5.27	1087.0 (M^+)
28	<p>4-Amino-4-methyl pentanoyl polymyxin B nonapeptide</p>		$C_{49}H_{85}N_{15}O_{12}$ 1075.65	5.08	1076.3 (MH^+)

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
29	 <p>4-(R)-Amino-5-methylhexanoyl polymyxin B nonapeptide</p>	<p>C₅₀H₈₇N₁₅O₁₂ 1089.67</p>	5.16	1089.6 (M ⁺)	
30	 <p>3-(S)-(1-pyrrolidin-2-yl)propionyl polymyxin B nonapeptide</p>	<p>C₅₀H₈₅N₁₅O₁₂ 1087.65</p>	5.10	1087 (M ⁺)	


- 40 -

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
31		4-(S)-aminopentanoyl polymyxin B nonapeptide	C48H83N15O12 1061.63	5.07	1062.1 (MH ⁺)
32		trans-4-hydroxycyclohexanecarbonyl polymyxin B nonapeptide	C50H84N14O13 1088.63	5.13	1088.7 (M ⁺)

- 41 -

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
33		3-Hydroxypropanoyl polymyxinB nonapeptide	C46H78N14O13 1034.59	5.19	1034.3 (M ⁺)
34		2-(R,S)-(2-Hydroxy-2-cyclohexyl)ethanoyl polymyxin B nonapeptide	C51H86N14O13 1103.65	5.80, 6.01	1103.9 (MH ⁺)

- 42 -

Ex.	Compound	Name	Formula and Predicted Mass	Retention Time (min)	m/z
35	 <p>2-(R,S)-2-Amino octanoyl polymyxin B nonapeptide</p> <p>C₅₁H₈₉N₁₅O₁₂ 1103.68</p> <p>5.42, 5.79</p> <p>1104.94</p>				

Antibacterial activity

To evaluate the potency and spectrum of the compounds, susceptibility testing was performed against four strains of each of the four Gram negative pathogens, *Escherichia coli*,

5 *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Acinetobacter baumannii*.

The day before testing, 3 to 4 colonies were picked from fresh Mueller-Hinton Agar (MHA) plate and transferred into 10 mL of Cation adjusted MHB (CaMHB). Cultures were incubated at 37°C 250 rpm for 18-20 hours before being diluted 100-fold in fresh CaMHB. The

10 subcultures were grown further until the OD₆₀₀ reached 0.2-0.3 corresponding to 10⁵-10⁶ CFU/ml. The actively growing cultures were diluted 100-fold in fresh medium and used for the inoculum.

MIC testing was performed by two-fold serial antibiotic dilutions in CaMHB in sterile 96-well

15 microtitre plates in a total volume of 170 µL (150 µL broth containing the antimicrobial agent, 20 µL inoculum). The assays were performed in duplicate. Plates were incubated aerobically without shaking for 18-20 hours at 37°C with the MIC defined as the lowest concentration of drug that prevented visible growth.

20 Table 4 shows the MIC (micrograms/mL) of Examples 2 to 14 compared to Polymyxin B (PMB),

Additional studies on antibacterial activity

25 Table 4A shows the MIC values obtained for compounds of Additional Examples 15 to 35 plus Examples 2, 6 and 14. Data was obtained under similar conditions to Table 4 except that different batches of cation-adjusted Muller-Hinton broth were used. The MIC values for these compounds are compared with those values obtained for Polymyxin B, Colistin Sulphate, CB-182,804 and NAB739 (as the TFA salt). CB-182,804 is a polymyxin decapeptide derivative with an aryl urea substituent at the N-terminus, which has been claimed to have lower toxicity than Polymyxin B (compound 5 in WO 2010/075416. See page 37). NAB739 has been described previously by Vaara et al.

Table 4. MIC Data for Compounds 2 to 14 (micrograms/mL)

STRAIN	PMB	Eg2	Eg3	Eg4	Eg5	Eg6	Eg7	Eg8	Eg9	Eg10	Eg11	Eg12	Eg13	Eg14
<i>E. coli</i> ATCC25922	0.5	1	2	2	1	0.5	2	2	0.5	2	4	0.5	0.5	0.5
<i>E. coli</i> ATCC700928	0.25	4	0.06	0.5	0.125	ND	0.5	1	0.125	0.125	ND	0.25	0.25	0.25
<i>E. coli</i> NCTC9001	0.25	0.25	4	0.25	ND	1	8	2	2	1	0.5	ND	ND	ND
<i>E. coli</i> NCTC12900	0.125	0.25	8	1	2	2	4	ND	1	2	2	ND	2	0.5
<i>P. aeruginosa</i> ATCC27853	0.5	0.5	0.5	0.125	0.25	0.25	0.5	0.5	0.06	0.125	0.5	0.25	0.5	0.5
<i>P. aeruginosa</i> ATCC 9721	0.5	2	0.125	0.25	0.25	0.125	ND	ND	0.5	0.125	0.25	ND	0.125	0.25
<i>P. aeruginosa</i> ATCC10145	0.5	1	0.25	0.5	0.5	0.25	ND	2	2	0.125	0.25	ND	0.5	0.25
<i>P. aeruginosa</i> ATCCCFM-9027	0.25	0.25	0.125	0.25	0.5	0.125	0.25	0.25	0.5	0.125	0.125	ND	2	0.25
<i>K. pneumoniae</i> ATCC 4352	0.5	0.5	1	0.125	0.25	0.125	0.5	ND	0.5	0.125	0.5	0.5	0.5	1
<i>K. pneumoniae</i> ATCCBAA-1706	0.25	0.5	16	0.25	0.5	0.25	0.06	ND	0.5	0.25	0.25	ND	1	0.25
<i>K. pneumoniae</i> NCTC7427	0.25	4	0.5	0.06	2	0.125	0.25	1	0.5	0.06	0.25	ND	0.25	0.25
<i>K. pneumoniae</i> NCTC8172	0.5	4	>32	2	>32	ND	>32	ND	>32	>32	2	ND	4	1
<i>A. baumannii</i> ATCC19606	0.5	2	2	1	1	0.25	ND	ND	16	0.125	2	32	0.5	0.5
<i>A. baumannii</i> ATCCBAA-747	0.25	2	>32	1	16	ND	8	ND	>32	4	1	ND	4	0.5
<i>A. baumannii</i> NCTC13423	0.25	1	16	1	16	ND	0.5	ND	>32	2	0.5	ND	2	0.06
<i>A. baumannii</i> NCTC7844	0.25	2	16	4	16	2	16	16	>32	4	4	ND	4	0.5

ND: not determined

Table 4A. MIC Data for Additional Examples 15 to 35 plus Examples 2, 6 and 14 (micrograms/mL)

STRAIN	PMB	Collistin Sulphate	CB108 804	NAB739 TFA salt	2	6	14	15	16	17	18	19	20
<i>E.coli</i> ATCC25922	1	1	2	2	2	2	2	2	32	8	0.5	1	4
<i>E.coli</i> NCTC9001	0.25	2	ND	ND	2	8	4	2	ND	ND	ND	ND	ND
<i>E.coli</i> NCTC12900	0.25	0.5	ND	ND	8	8	4	2	ND	ND	4	ND	ND
<i>E.coli</i> ATCC700928	0.25	0.25	ND	ND	0.5	2	2	1	ND	ND	0.25	ND	ND
<i>P.aeruginosa</i> ATCC27853	0.5	0.5	1	1	0.5	0.25	0.25	0.25	0.25	2	0.125	0.25	1
<i>P.aeruginosa</i> ATCC10145	1	0.5	ND	ND	1	0.5	0.5	1	ND	ND	0.25	ND	ND
<i>P.aeruginosa</i> ATCC9721	0.125	0.5	ND	ND	0.5	0.125	0.25	0.125	ND	ND	ND	ND	ND
<i>P.aeruginosa</i> ATCC9027	0.25	0.5	ND	ND	0.5	0.25	0.125	0.125	ND	ND	2	ND	ND
<i>K.pneumoniae</i> ATCC4352	0.25	0.125	ND	2	1	1	0.5	0.5	8	8	0.5	2	>32
<i>K.pneumoniae</i> NCTC7427	0.25	0.25	ND	ND	0.5	0.125	0.5	0.25	ND	ND	0.125	ND	ND
<i>K.pneumoniae</i> NCTC8172	1	0.5	ND	ND	4	16	>32	4	ND	ND	>32	ND	ND
<i>K.pneumoniae</i> ATCCBAA-1706	ND	ND	ND	ND	ND	ND	0.5	ND	ND	ND	8	ND	ND
<i>A.baumannii</i> ATCC19606	0.125	ND	0.5	1	2	4	ND	1	32	4	1	16	>32
<i>A.baumannii</i> ATCCBAA-747	0.125	0.25	ND	ND	2	ND	0.5	8	ND	ND	16	ND	ND
<i>A.baumannii</i> NCTC13423	<0.06	<0.06	ND	ND	1	ND	0.125	4	ND	ND	4	ND	ND
<i>A.baumannii</i> NCTC7844	0.25	0.125	ND	ND	4	2	0.5	8	ND	ND	8	ND	ND

- 46 -

STRAIN	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
<i>E.coli</i> ATCC25922	32	4	0.5	2	8	8	2	4	8	16	>32	8	>32	0.5	ND
<i>E.coli</i> NCTC9001	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
<i>E.coli</i> NCTC12900	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
<i>E.coli</i> ATCC700928	ND	ND	ND	ND	ND	ND	ND	ND	2	ND	ND	ND	ND	ND	ND
<i>P.aeruginosa</i> ATCC27853	1	0.5	0.25	0.125	4	4	0.125	<0.06	0.125	0.125	0.25	0.125	0.5	0.125	0.25
<i>P.aeruginosa</i> ATCC10145	ND	ND	ND	ND	ND	ND	ND	ND	0.5	ND	ND	ND	ND	0.5	ND
<i>P.aeruginosa</i> ATCC9721	ND	ND	ND	ND	ND	ND	ND	ND	0.25	ND	ND	ND	ND	0.25	ND
<i>P.aeruginosa</i> AATCCRM 9027	ND	ND	ND	ND	ND	ND	ND	ND	0.5	ND	ND	ND	ND	0.5	ND
<i>K.pneumoniae</i> ATCC4352	32	8	0.5	0.125	4	8	0.125	0.25	0.5	1	0.5	4	4	1	1
<i>K.pneumoniae</i> NCTC7427	ND	ND	ND	ND	ND	ND	ND	ND	0.5	ND	ND	ND	ND	0.5	ND
<i>K.pneumoniae</i> NCTC8172	ND	ND	ND	ND	ND	ND	ND	ND	>32	ND	ND	ND	ND	>32	ND
<i>K.pneumoniae</i> ATCCBAA-1706	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
<i>A.baumannii</i> ATCC19606	>32	>32	8	2	16	16	0.5	4	2	8	8	16	>32	1	2
<i>A.baumannii</i> ATCCBAA-747	ND	ND	ND	ND	ND	ND	ND	ND	1	ND	ND	ND	ND	1	ND
<i>A.baumannii</i> NCTC13423	ND	ND	ND	ND	ND	ND	ND	ND	0.5	ND	ND	ND	ND	0.25	ND
<i>A.baumannii</i> NCTC7844			ND	ND	ND	ND	ND	ND	2	ND	ND	ND	ND	2	ND

ND: not determined

The *in vitro* antibacterial activities of the compounds of examples 2 and 6 were evaluated against a panel of 500 Gram-negative bacterial isolates alongside Colistin. The panel consisted of 100 clinical isolates of each of *A. baumannii*, *E. coli*, *K. pneumoniae* and *P. aeruginosa*. The panel represented the current epidemiology in Europe and the USA and included a number of relevant strains with defined resistant phenotypes to current clinically-used antibacterial agents. These resistant strains included 20 *A. baumannii*, 22 *E. coli*, 25 *K. pneumoniae* and 20 *P. aeruginosa* strains.

The results of the study are summarised in Table 4B. All compounds were tested up to a maximum concentration of 64 μ g/mL with the exception of colistin, which was evaluated up to a maximum concentration of 16 μ g/mL.

Table 4B - Summary of the MIC values (μ g/mL) of Examples 2 and 6, and Colistin against a panel of 400 Gram-negative clinical isolates and 100 Gram negative strains of defined resistant phenotypes

Example	Organism	Range Clinical isolates	Range Resistant strains	MIC ₅₀ Clinical isolates	MIC ₅₀ Resistant strains	MIC ₉₀ Clinical isolates	MIC ₉₀ Resistant strains
Colistin	<i>A. baumannii</i>	0.5-4	1-4	2	2	2	2
	<i>E. coli</i>	0.5-4	0.25-2	2	1	2	1
	<i>K. pneumoniae</i>	1-16	1->16	2	2	4	8
	<i>P. aeruginosa</i>	0.25-4	0.5-2	2	2	4	2
2	<i>A. baumannii</i>	1-32	2-4	4	4	8	4
	<i>E. coli</i>	1-16	0.5-4	2	1	4	2
	<i>K. pneumoniae</i>	2->64	2->64	8	8	32	32
	<i>P. aeruginosa</i>	1-8	1-4	2	2	4	4
6	<i>A. baumannii</i>	0.5->64	1-4	2	2	8	4
	<i>E. coli</i>	0.12->64	0.06-4	2	0.5	8	4
	<i>K. pneumoniae</i>	0.5->64	1->64	>64	>64	>64	>64
	<i>P. aeruginosa</i>	0.06-64	0.12-16	0.25	0.5	16	8

In vivo efficacy against *E. coli* thigh infection in mice

The *in vivo* efficacy of 8 compounds of the invention (Examples 2, 4, 5, 6, 7, 8, 10, and 11) was evaluated in a mouse thigh infection model of *E. coli*. The results are summarized in Table 5.

Groups of 5 female specific-pathogen-free CD-1 mice weighing 22 ± 2 g were used. The animals were made neutropenic by intraperitoneal administration of cyclophosphamide on days -4 (150 mg/kg) and -1 (100 mg/kg). On Day 0, animals were inoculated intramuscularly with 10^5 CFU/mouse of *Escherichia coli* isolate ATCC25922 into the right thigh. At 1 h, the

5 CFU count was determined from 5 mice and the remaining mice (five per group) were treated with a subcutaneous injection of the drug at + 1 and 6 hr post-infection. In each study, there were two dose groups per test compound, 1.5 and 5 mg/kg BID, respectively. Examples 2, 4, 5, 6, 7, 8, 10, 11 and polymyxin B were prepared in Normal Saline at 2 mg/mL and the solution was adjusted to pH 6-7 by addition of 0.1M H₂SO₄ or 4.2% NaHCO₃ as required. Twenty-four

10 hours after infection, the mice were euthanized humanely. The muscle of the right thigh of each animal was harvested, homogenized, serially diluted and plated on Brain Heart Infusion agar + 0.5% charcoal (w/v) for CFU determination. Decrease of the total CFU of right thigh as compared to control counts at 24 hrs post-infection was determined for each dose group.

15 The compounds 2 and 6 at 10 mg/kg/day demonstrated an efficacy comparable to that of polymyxin B with over $3 \log_{10}$ reduction in bacterial counts.

Table 5: *In vivo* Efficacy Versus *E. coli* ATCC25922 Thigh Infections in Neutropenic Mice

Example No	Total daily dosage (mg/kg)	Mean \log_{10} CFU reduction vs. control
Polymyxin B	3	2.5 ^a
	10	4.2 ^a
2	3	0.98 ^b
	10	4.48 ^b
4	3	0 ^b
	10	0.82 ^b
5	3	0.52
	10	0.51
6	3	0.72 ^b
	10	3.38 ^b
7	3	1.09
	10	2.15
8	3	0.53
	10	0.82
10	3	0.17
	10	0.56
11	3	1.19
	10	1.85

^amean values of 5 independent studies; ^bmean value of 2 independent studies.

Additional studies on the *in vivo* efficacy against *E. coli* thigh infection in mice

The *in vivo* efficacy of the compound of Example 14 was evaluated in a mouse thigh infection model of *E. coli*, using the methods described in the examples above. The result is

5 summarized in Table 5A in comparison with Polymyxin B.

Table 5A: *In vivo* Efficacy Versus *E. coli* ATCC25922 Thigh Infections in Neutropenic Mice

Example No	Total daily dosage (mg/kg)	Mean \log_{10} CFU reduction vs. control
Polymyxin B	3	3.75
	10	4.87
14	3	0
	10	4.05

10 Compounds 14 at 10 mg/kg/day demonstrated an efficacy comparable to that of polymyxin B with over 3 \log_{10} reduction in bacterial counts.

Additional studies on the *in vivo* efficacy against *Klebsiella pneumoniae* thigh infection in mice

15 Using the same procedure as described above, the *in vivo* efficacy of three compounds of the invention (Examples 2, 6, and 14) was evaluated in a mouse thigh infection model of *Klebsiella pneumoniae* ATCC10031, using Colistin (Polymyxin E) as comparator. The results are summarized in Table 5B. The compounds 2, 6 and 14 at 10 mg/kg/day demonstrated an efficacy comparable to that of Colistin with approx. 2 \log_{10} reduction in bacterial counts.

Table 5B: *In vivo* efficacy versus *K. pneumoniae* ATCC10031 thigh infections in neutropenic mice.

Example	Total daily dosage (mg/Kg)	Mean \log_{10} CFU reduction vs. control
Colistin	10	2.60
2	10	2.22
6	10	1.92
14	10	2.30

Pharmacokinetic and urinary clearance studies

The pharmacokinetics and urinary clearance of 3 compounds (Examples 2, 4 and 6) of the invention and polymyxin B were evaluated in rats.

5 Drug solutions were prepared at 4 mg/mL in Normal Saline and the pH adjusted to 6-7 by adding the appropriate volume of 0.1 M H₂SO₄ or 4.2% NaHCO₃. The solutions were filter-sterilized and stored at -80°C before use. On the day of the experiment, drug solutions were diluted to 1 mg/mL with sterile Normal Saline.

10 Groups of 3 male Sprague Dawley rats were acclimatised for a minimum of 4 days before the study. Rats were anesthetized using isofluorane and a cannula was inserted into the jugular vein. One day after surgery, rats were dosed with an intravenous bolus injection of the solution at 1 mg/kg through the cannula, followed by washing with Normal Saline. Blood was collected

15 manually through the cannula prior to administration of the compound and at 0.08, 0.25, 0.5, 1, 3, 6, 8 and 24 h thereafter. Plasma was harvested by centrifugation immediately after blood collection. Twenty-four hour urine samples were collected prior to and after administration of the compound in 0-4 h, 4-6 h, and 6- 24 h intervals. Plasma and urine samples were frozen at -20°C.

20 Determination of the plasma and urine concentrations of the drug was performed by Liquid Chromatography Mass Spectrometry (LC-MS/MS). Before analysis the plasma and urine samples were prepared as follows. Plasma samples were thawed on the day of analysis and mixed with 3 volumes of acetonitrile containing 0.1% (v/v) formic acid and 100 ng/mL of

25 internal standard. After centrifugation, supernatants were transferred to a 96-well plate and diluted 1:1 with water ready for analysis by LC-MS/MS. Urine samples were purified by solid-phase extraction (SPE) on Oasis HLB cartridges (Waters, UK) eluting with 100% methanol. An aliquot was transferred to a 96 well plate and diluted 1:1 with water before addition of internal standard solution, ready for analysis by LC-MS/MS.

30 Column: Kinetex 2.6 µm XB-C18 50 x 2.1 mm
Mobile Phase A: Water + 0.15% TFA or 0.1% formic acid
Mobile Phase B: Acetonitrile + 0.15% TFA or 0.1% formic acid
Flow rate: 0.5 mL/min
Gradient:

35 Time 0 min 95% A 5% B
Time 1.20 min 5% A 95 % B
Time 1.50 min 5 % A 95 % B
Time 1.51 min 95 % A 5 % B

Time 3.00 min 95 % A 5% B
 Cycle time 4.5 min
 Injection volume: 20 μ L

5 The pharmacokinetic parameters were determined by non-compartmental analysis using WinNonLin v5.3. The urinary recovery was recorded as the percentage of intact drug recovered in the urine for the first 24 h after injection.

Table 6 Pharmacokinetics of polymyxin B and PMBN derivatives

Example No	t _{1/2} (hr)	C _{max} (ng/mL)	AUC _{0-inf} (ng.hr/mL)	Cl (mL/hr/kg)	Vd (mL/kg)	Urinary recovery (% dose)
Polymyxin B	1.94	1455	2372	429	1120	0.3
2	1.34	2400	4009	251	488	0.5
4	1.33	2524	2033	492	690	3.9
6	0.56	3581	2619	386	289	8.9

10

Interestingly, all compounds show higher urinary recovery than polymyxin B. Previous studies reported that polymyxin E (colistin) undergoes extensive renal tubular reabsorption (Li *et al.*, *Antimicrob. Agents and Chemotherapy*, 2003, 47(5); Yousef *et al.*, *Antimicrob. Agents and Chemotherapy*, 2011, 55(9)). Whilst not wishing to be bound by theory, higher urinary

15 clearance of the compounds could reflect a decreased renal tubular reabsorption which could in turn reduce their nephrotoxicity potential.

In vitro renal cell toxicity assay

The renal cell toxicity of the compounds was assessed in an *in vitro* assay using the HK-2 cell line, an immortalized proximal tubule cell line derived from a normal human kidney. The 20 endpoint to describe the toxicity of the compounds was the reduction of resazurin correlating with the metabolic activity of the cells.

Cells were cultured in 150 cm² flasks in 25 mL supplemented KSF (with 5 ng/mL EGF and 50 μ g/mL BPE). Cells were maintained at 70% confluence with a maximum of 25 passages.

25 Day 1: Media was removed and cells were washed with 10ml DPBS. Six ml of a 0.25% trypsin solution with EDTA was then added to the flask and the cells returned to the incubator. After 1 to 2 minutes incubation, 14 ml media was added to the flask to inactivate the trypsin. The cell suspension was transferred to a centrifuge tube and the cells pelleted at 1000 rpm for

6 minutes. The cell pellet was then resuspended in fresh media supplemented with EGF and BPE. The cell number was counted and cells were diluted to 46875 cells/mL in fresh medium supplemented with EGF and BPE. 7500 cells were dispensed in each well in a volume of 160 μ l and incubated at 37°C for 24 h.

5

Day 2: Test compounds were prepared directly into the media. Nine point concentrations were prepared from 1000 μ g/mL to 1.95 μ g/mL in two-fold dilutions in fresh medium. The microtiter plates were removed from the incubator and the media replaced with 100 μ l of the dilutions of the compound solution. Every set of concentration was done in triplicate, and positive and 10 negative controls were added to each plate. The plates were then incubated for 24h at 37°C with 5% CO₂ in a humidified atmosphere.

Day 3: The reagent containing the resazurin (CellTiter-Blue, Promega) was diluted in PBS (1:4) and added at 20% (v/v) to each well. The plates were then incubated at 37°C for 2h 15 before the fluorescent reduction product was detected.

Media only background values were subtracted before the data was analysed using GraphPad Prism. Compound concentration values were plotted as log values to enable a dose-response curve to be fitted and IC₅₀ values determined (Table 7).

20

Table 7: IC₅₀ Data for Polymyxin B and Examples 2-14

Example No	IC ₅₀ HK-2 cells (μ g/mL) ^a
Polymyxin B	11 ^b
2	87
3	166
4	82
5	250
6	154
7	138
8	497
9	104
10	127
11	310
12	>500
13	158
14	60

^a Mean values of up to 6 independent studies; ^b Mean value of 16 independent studies.

Additional studies on the *in vitro* renal cell toxicity assay

5 The renal cell toxicity of the additional example compounds was assessed in an *in vitro* assay using the HK-2 cell as described in the example above. The IC₅₀ values for these compounds are set out in Table 7A below. For comparison, the renal cell toxicity Colistin, and CB182,804 (compound 5 in WO2010/075416) and NAB739 were also assessed..

Table 7A: IC₅₀ Data for Colistin and Examples 15-35

Example No	IC ₅₀ HK-2 cells (µg/mL) ^a
Colistin	28 ^a
CB182,804	22
NAB739 TFA salt	176
15	133
16	1000 ^c
17	84
18	>500
19	157 ^c
22	500 ^c
23	173
24	101
25	277
26	128
27	118
28	108
29	82
30	133
31	93
32	500
33	1000 ^c
34	86
35	82

^a Mean values of up to 6 independent studies; ^csolubility issues noted at top concentration

Additional studies on *In vivo* nephrotoxicity

A model of nephrotoxicity of polymyxins (adapted from Yousef et al., *Antimicrob. Agents Chemother.*, 2011, 55 (9): 4044-4049) was established in rats. The compounds of examples 2, 6, and 14 were examined in the model and compared to Colistin (in its sulphate form).

5 After one week acclimatisation, male Sprague-Dawley rats were surgically prepared with a jugular cannula and were housed individually, as required, either in pre-assigned housing cages or metabolic cages. Colistin and the example compounds were prepared in saline. Compounds were introduced *via* the jugular canula twice a day 7 hours apart for seven days. Each dose was increased progressively for three days up to the top dose that was then 10 administered until termination of the study. Twenty-four hour urine collection (on ice) was performed at pre-dose and on days 4 and 7. The dose regimen is set out in Table 8 below.

Table 8 - Dose regimen used in the *in vivo* nephrotoxicity study. Doses are indicated in mg drug base/kg.

Dose regimens	Day 1		Day 2		Day 3		Day 4 to Day 7 or Day 10	
	a.m.	p.m.	a.m.	p.m.	a.m.	p.m.	a.m.	p.m.
2 mg/kg bid	0.25	0.5	0.625	0.625	0.875	1.375	2	2
8 mg/kg bid	1	2	2.5	2.5	3.5	5.5	8	8

15 The activity in urine of the N-acetyl-beta-D-glucosaminidase (NAG) was determined spectrophotometrically using the NAG assay kit from Roche Applied Science. Biomarkers of kidney injury were determined using the Kidney Injury Panel II from the Multi-Spot® Assay System (Meso Scale Discovery).

20 Examples 2, 6, and 14 dosed using the 8 mg/kg regimen showed significantly reduced levels of the renal biomarkers NAG, albumin and cystatin C compared to Colistin at the same dose regimen (see Figures 1 to 3). The response was similar to that elicited by Colistin at a maximum concentration of 2mg/kg.

25 Figure 1 shows the concentration of NAG (ng/24 h) at days 0, 4 and 7 for compounds 2, 6, and 14 and Colistin. The left-hand graph shows from left to right Colistin (2 mg/kg BID), Colistin (8 mg/kg BID), compound 2 (8 mg/kg BID) and 6 (8 mg/kg BID). The right-hand graph shows Colistin (2 mg/kg BID), Colistin (8 mg/kg BID) and compound 14 (8 mg/kg BID).

Figure 2 shows the concentration of albumin (ng/24 h) at days 0, 4 and 7 for compounds 2, 6, and 14 and Colistin. The left-hand graph shows from left to right Colistin (2 mg/kg BID),


Colistin (8 mg/kg BID), compound 2 (8 mg/kg BID) and 6 (8 mg/kg BID). The right-hand graph shows Colistin (2 mg/kg BID), Colistin (8 mg/kg BID) and compound 14 (8 mg/kg BID).

Figure 3 shows the concentration of cystatin C (ng/24 h) at days 0, 4 and 7 for compounds 2, 6, and 14 and Colistin. The left-hand graph shows from left to right Colistin (2 mg/kg BID),

5 Colistin (8 mg/kg BID), compound 2 (8 mg/kg BID) and 6 (8 mg/kg BID). The right-hand graph shows Colistin (2 mg/kg BID), Colistin (8 mg/kg BID) and compound 14 (8 mg/kg BID).

Claims:

1. A compound of the formula (I):



wherein:

X represents an -C(O)-, -NHC(O)-, -OC(O)-, -CH₂- or -SO₂-; and

R¹ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a phenylalanine, leucine or valine residue;

R² together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a leucine, iso-leucine, phenylalanine, threonine, valine or nor-valine residue;

R³ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents a threonine or leucine residue;

R⁴ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached, represents α,γ -diaminobutyric acid or a serine residue;

R⁵ represents a group (a) to (g), where:

(a) is C₀₋₁₂ alkyl(C₃₋₈ cycloalkyl) and C₀₋₁₂ alkyl is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, butyl, n-butyl and tert-butyl, wherein the alkyl or the cycloalkyl is substituted with (i) one, two or three hydroxyl groups, or (ii) one -NR⁶R⁷ group, or (iii) one -NR⁶R⁷ group and one or two hydroxyl groups;

2012338571 16 Feb 2016

- (b) is C_{2-12} alkyl, wherein the alkyl is substituted with (i) one, two or three hydroxyl groups, or (ii) one $-NR^6R^7$ group at a terminal of the alkyl chain, or (iii) one $-NR^6R^7$ group and two hydroxyl groups;
- (c) is C_{0-12} alkyl(C_{4-6} heterocyclyl);
- (d) is C_{3-8} cycloalkyl, and the cycloalkyl is substituted with (i) one, two or three hydroxyl groups, or (iii) one $-NR^6R^7$ group and one or two hydroxyl groups;
- (e) is C_{3-12} alkyl, such as C_{3-10} alkyl, wherein the alkyl is substituted with one $-NR^6R^7$ group and one hydroxyl group;
- (f) is C_{6-12} alkyl substituted with one $-NR^6R^7$ group; and
- (g) is C_5 cycloalkyl substituted with one $-NR^6R^7$ group,

R^6 represents hydrogen or C_{1-4} alkyl; and

R^7 represents hydrogen or C_{1-4} alkyl,

R^8 represents methyl or hydrogen, or

a prodrug thereof, and/or

a pharmaceutically acceptable salt thereof.

2. A compound of formula (I) according to claim 1 where R^1 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents a phenylalanine residue.

3. A compound of formula (I) according to claim 2 where the phenylalanine is a D-phenylalanine residue.

4. A compound of formula (I) according to claim 1 where R^2 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents a leucine residue.

5. A compound of formula (I) according to claim 1 where R^3 together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents a threonine residue.

6. A compound of formula (I) according to claim 1 where R⁴ together with the carbonyl group and nitrogen alpha to the carbon to which it is attached represents an α,γ -diaminobutyric acid residue.
7. A compound of formula (I) according to claim 1 wherein R⁶ is hydrogen or methyl.
8. A compound of formula (I) according to claim 1 wherein R⁷ is hydrogen or methyl.
9. A compound of formula (I) according to claim 1 where X is -C(O)-.
10. A compound according to claim 1 selected from the group consisting of:
 - 2-Hydroxyoctanoyl polymyxin B nonapeptide;
 - 3-Aminopropanoyl polymyxin B nonapeptide;
 - 3-(*N,N*-Dimethylamino)-propanoyl polymyxin B nonapeptide;
 - 4-Aminobutanoyl polymyxin B nonapeptide;
 - 6-Aminohexanoyl polymyxin B nonapeptide;
 - 8-Hydroxyoctanoyl polymyxin B nonapeptide;
 - 8-Aminooctanoyl polymyxin B nonapeptide;
 - 3-(*N*-Methylamino)propanoyl polymyxin B nonapeptide;
 - 2-Amino cyclopentane carbonyl polymyxin B nonapeptide;
 - 3-Aminopropanoyl colistin (polymyxin E) nonapeptide;
 - 3-Pyrrolidine-3-carbonyl polymyxin B nonapeptide;
 - 3-Amino-3-cyclohexanepropanoyl polymyxinB nonapeptide;
 - 5-Aminopentanoyl polymyxin B nonapeptide;
 - 3-Hydroxyoctanoyl polymyxin B nonapeptide;
 - 4-(*N,N*-dimethylamino)-butanoyl polymyxin B nonapeptide;
 - 7-Aminoheptanoyl polymyxin B nonapeptide;
 - 4-Morpholinylbutanoyl polymyxin B nonapeptide;
 - 6-Hydroxyhexanoyl polymyxin B nonapeptide;
 - 3-Hydroxybutanoyl polymyxin B nonapeptide;
 - 4-(*N*-methylamino)-butanoyl polymyxin B nonapeptide;
 - 4-Aminobutanoyl polymyxin E nonapeptide;
 - 2-Hydroxyoctanoyl polymyxin E nonapeptide;
 - 4-Amino-5-methylhexanoyl polymyxin B nonapeptide;
 - 3-(1-Pyrrolidin-2-yl)-propionyl polymyxin B nonapeptide;

trans-4-Hydroxycyclohexanecarbonyl polymyxin B nonapeptide;
3-Hydroxypropanoyl polymyxin B nonapeptide;
(2-Hydroxy-2-cyclohexyl)ethanoyl polymyxin B nonapeptide;
2-Amino octanoyl polymyxin B nonapeptide; and
a pharmaceutically acceptable salt thereof.

11. A pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in claim 1 together with a pharmaceutically acceptable carrier.
12. A compound according to claim 1 wherein R^5 is C_{0-12} alkyl(C_{3-8} cycloalkyl) and C_{0-12} alkyl is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, butyl, n-butyl and tert-butyl, wherein the alkyl or the cycloalkyl is substituted with (i) one, two or three hydroxyl groups, or (ii) one $-NR^6R^7$ group, or (iii) one $-NR^6R^7$ group and one or two hydroxyl groups.
13. A compound according to claim 1 wherein R^5 is C_{0-12} alkyl(C_{3-8} cycloalkyl) wherein the alkyl or the cycloalkyl is substituted with one $-NR^6R^7$ group.
14. A compound according to claim 1 wherein the C_{3-8} cycloalkyl is C_5 or C_6 cycloalkyl.
15. A compound according to claim 1 wherein R^5 is C_{2-12} alkyl, wherein the alkyl is substituted with (i) one, two or three hydroxyl groups, or (ii) a $-NR^6R^7$ group at a terminal of the alkyl chain, or (iii) one $-NR^6R^7$ group and two hydroxyl groups.
16. A compound according to claim 15 wherein R^5 is C_{2-12} alkyl substituted with a $-NR^6R^7$ group at a terminal of the alkyl chain.
17. A compound according to claim 15, wherein R^5 is C_{2-12} alkyl is substituted with one, two or three hydroxyl groups.
18. A compound according to claim 1, wherein R^5 is C_{0-12} alkyl(C_{4-6} heterocyclil).
19. A compound according to claim 1, wherein R^5 is C_{3-8} cycloalkyl, and the cycloalkyl is substituted with (i) one, two or three hydroxyl groups, or (iii) one $-NR^6R^7$ group and one or two hydroxyl groups.

20. A compound according to claim 1, wherein R⁵ is C₃₋₁₂ alkyl wherein the alkyl is substituted with one -NR⁶R⁷ group and one hydroxyl group.
21. A compound according to claim 1, wherein R⁵ is C₆₋₁₂ alkyl substituted with one -NR⁶R⁷ group.
22. The compound of claim 1, wherein R⁵ is C₅ cycloalkyl substituted with one -NR⁶R⁷ group.
23. A compound of formula (I) according to claim 1 wherein R⁸ is methyl.
24. A method of treating a Gram negative bacterial infection comprising administering to a subject in need thereof a therapeutically effective amount of a compound according to claim 1.
25. A method according to claim 24 where the bacterium is multidrug resistant.
26. Use of a therapeutically effective amount of a compound according to claim 1 in the preparation of a medicament for the treatment of Gram negative bacterial infections.
27. The use according to claim 26 where the bacterium is multidrug resistant.

Figure 1




Figure 2




Figure 3

