




(43) International Publication Date
30 December 2015 (30.12.2015)

- (51) International Patent Classification:
B01L 3/14 (2006.01) *C40B 60/12* (2006.01)
C40B 30/06 (2006.01)
- (21) International Application Number:
PCT/US2015/037997
- (22) International Filing Date:
26 June 2015 (26.06.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/017,579 26 June 2014 (26.06.2014) US
- (71) Applicant: THE TRUSTEES OF THE UNIVERSITY
OF PENNSYLVANIA [US/US]; 3160 Chestnut Street,
Suite 200, Philadelphia, PA 19104 (US).
- (72) Inventor: FANG-YEN, Christopher; 224 S 23rd St.,
Philadelphia, PA 19103 (US).
- (74) Agents: RAGUSA, Paul, A. et al.; Baker Botts LLP, 30
Rockefeller Plaza, New York, NY 10012-4498 (US).

- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

- (54) Title: MICRO-FABRICATED MULTI-WELL CHIPS FOR LONG TERM AUTOMATED IMAGING OF C. ELEGANS
GROWTH AND BEHAVIOR



- (57) Abstract: A composition, including a
substrate having a planar array of depressions
each defined by concave walls and a moat dis-
posed around each depression of said array of
depressions.

FIG. 1a



Published:

— *with international search report (Art. 21(3))*

MICRO-FABRICATED MULTI-WELL CHIPS FOR LONG TERM AUTOMATED IMAGING OF *C. ELEGANS* GROWTH AND BEHAVIOR

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to U.S. Provisional Patent Application
Serial No. 62/017,579, filed June 26, 2014, the contents of which are incorporated by
reference herein in their entirety.

FIELD OF THE INVENTION

10 The presently disclosed subject matter relates to scalable compositions
for the cultivation and imaging of *C. elegans* and other organisms, as well as methods
of using the same.

BACKGROUND

 Aging is manifested in gradual changes in an adult organism which can
cause a reduction of function and an increase in mortality rate. Certain studies of
15 model organisms have identified highly conserved processes and pathways which
influence aging, including dietary restriction, insulin/insulin-like signaling, and the
cytoprotective DAF-16/FOXO pathway. The model roundworm *Caenorhabditis*
elegans (*C. elegans*) has been used to study aging biology, due to its short lifespan
and genetic manipulability. However, certain studies of aging in *C. elegans* have
20 several limitations. First, they can focus on measurement of lifespan, an incomplete
description that ignores potentially important aspects of the aging process. Second,
they can rely on visual observation of worms on agar plates, a laborious and
potentially-subjective process not suitable for high-throughput analysis. Third, they
can be limited to studies at the population level, with individual trajectories
25 unavailable.

 To address these limitations, certain automated or semi-automated
techniques for analysis of aging in *C. elegans* have been developed. The study of
worm aging in microtiter plates under liquid culture conditions permits an increase in
throughput, but also changes behavior, physiology, and lifespan. Constraining the
30 animal to nanodroplets can result in severely compromised survival. Another method
employs microfluidic chambers, which can have limited scalability and can usually be

limited to specific developmental stages. One technique can employ flatbed scanners to image tens of thousands of worms on standard agar plates. However, due to a low frame rate (~1 / hr), it cannot track young adult animals or easily quantify behavior. Another study described a system for long-term multi-well imaging in glass wells, which can allow for longitudinal imaging but is not easily scalable to large numbers of worms.

In view of the foregoing, there remains a need in the art for scalable compositions for the cultivation and imaging of *C. elegans*, as well as other organisms, which can allow for longitudinal imaging of large numbers of organisms. In addition to applications in aging, such methods can be used for monitoring nematode development, growth, behavior, fluorescence, morphological changes, and other aspects.

SUMMARY

In certain embodiments, the present disclosure provides compositions including: a substrate having a planar array of depressions, e.g., a well, each defined by concave walls; and a moat disposed around each depression of said array of depressions. In certain embodiments, the depressions have a 3 mm diameter, a 3 mm depth, and a center-to-center spacing of 4.5 mm. In certain embodiments, the moat disposed around each depression is 0.5 mm wide and 3 mm deep. In certain embodiments, the composition is made of at least one of polyurethane, polycarbonate, polyvinyl, polystyrene, polyvinylchloride (PVC), polypropylene, cyclic olefin copolymer (COC), and polydimethylsiloxane (PDMS).

In certain embodiments, the substrate having a planar array of depressions is adapted for insertion into a tray to form a microwell assembly. In certain embodiments, the substrate is of uniform thickness. In certain embodiments, the composition is made entirely of polyurethane, polycarbonate, polyvinyl, polystyrene, polyvinylchloride (PVC), polypropylene, or cyclic olefin copolymer (COC). In certain embodiments, the composition has an overall shape and dimension generally conforming to the ANSI/SLAS microplate standard. In certain embodiments, the composition comprises at least one of a 48-well microplate having an 8×6 array of wells; a 96-well microplate having an 8×12 array of wells; a 240-well

microplate having a 12 x 20 array of wells; and a 384-well microplate having a 16x24 array of wells.

In certain embodiments, the present disclosure relates methods of cultivating and imaging an organism, including by providing a composition that includes: (A) a substrate having a planar array of depressions each defined by concave walls; and (B) a moat disposed around each depression of said array of depressions. An organism can be introduced into a depression of said substrate, in addition to solid or liquid culture medium, any necessary food, and other reagents. The organism can be cultivated and imaged over a period of time, for example, up to several months. In certain embodiments, the organism is selected from the group consisting of *C. elegans*, other nematodes, zebrafish larvae, and *Drosophila* larvae.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 illustrates an exemplary design, fabrication, experimental setup, and data analysis according to the disclosed subject matter. Fig. 1a shows a schematic of a 16-well example embodiment of the disclosed subject matter (top view). Fig. 1b shows a fabrication process (cross section). Fig. 1c shows darkfield imaging with a CCD or CMOS camera inside a light-shielded imaging rig. A robotic plate handler moves each plate from a stack to the imaging rig for 10 minute-imaging periods. Fig. 1d shows an image of a 240-well example embodiment of the disclosed subject matter (containing L4 *C. elegans*). A machine vision algorithm measures activity of each worm. Distance between well centers is 4.5 mm. Fig. 1e shows an example of a well shape, with a cross section through center of well, where the well is symmetric around central axis. Fig. 1f shows a device having a convex rounded surface on the bottom of the device.

Fig. 2 shows that an example embodiment of the disclosed subject matter generates behavioral profiles and automates lifespan analysis. Figs. 2a-b illustrate activity profiles for N2 and Hawaiian (CB4856) strains at 25°C. White indicates high activity and black indicates zero activity. Each row denotes the activity profile of a single worm. Worms are sorted by time of death. Time of death for each individual is marked by a white 'x'. Fig. 1c shows an activity trace of a single N2 worm. Time of death is marked as the final moment of non-zero activity (arrow). Fig. 2d shows an individual Hawaiian worm activity trace. Time of death is marked

as the final moment of non-zero activity (arrow). Fig. 2e shows average survivor activity (solid curves) and surviving fraction (dashed curves) plotted over time for N2 and Hawaiian. Fig. 2f shows survival curves for worms cultivated under standard laboratory conditions (dashed lines) and those grown on an example embodiment of the disclosed subject matter (solid lines). Worms were scored as dead by hand for manual assays and were automatically scored as dead by machine vision for the example embodiment of the disclosed subject matter. Fig. 2g shows survival data for worms cultivated with RNAi on an example embodiment of the disclosed subject matter. N2 and mutants (solid curves) were treated with an empty vector. N2 worms were fed either *daf-2* or *daf-16* RNAi (dotted curves). Temperature = 20°C.

Fig. 3 illustrates lifespan prediction by behavioral profiles. Figs. 3a-d show activity profiles for N2 and three short-lived strains at 20°C. Time of death for each individual is marked by a white 'x'. Fig. 3e shows survivor average activity (solid curves) and surviving fraction (dashed curves) for N2 and *daf-16 (mu86)*. Average *daf-16* activity is higher than N2 just before their respective deaths. Fig. 3f shows survival curves for N2 and short-lived strains. Fig. 3g shows healthspan curves for N2 short-lived strains. Figs. 3h-k show lifespan versus healthspan for individual worms fitted to a line (dashed black line). Figs. 3l-o shows lifespan versus early life activity decay rate. Activity decay rate is the negative slope of individual activity traces during days 5-13 for N2 and days 2-6 for short-lived strains

Fig. 4 illustrates behavioral profiles in long-lived mutant strains. Figs. 4a-d show activity profiles for N2 and three long-lived strains. White x's mark death. Fig. 4e shows an example *age-1* mutant activity curve. Quiescence and revival are evident. Fig. 4f shows the fraction of quiescence for the same worm as Fig. 4e. The fraction of quiescence increases greatly and then declines before the worm dies. Fig. 4g shows an example *daf-2 (e1370)* activity curve. The precipitous decline in activity is evident on about day 10. Fig. 4h shows the fraction of quiescence for the same worm as in Fig. 4g. The fraction of quiescence increases from 0 to about 1 on about day 10 and stays high until the worm dies on day 50. Fig. 4i shows survivor average activity (solid curves) and survival (dashed curves) for N2, *age-1(hx546)*, and *daf-2(e1370)*. *age-1* revival is visible as a 'tail' to the right of the initial decline. *daf-2(e1370)* decline together and do not revive. Fig. 4j shows survival curves for long-

lived mutants. Fig. 4k shows locomotor healthspan curves for long-lived mutants. Fig. 4l shows the day of initial decline for long-lived mutants. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Error bars are s.e.m. Fig. 4m shows the day of initial decline normalized by lifespan for each individual. ***, $p < 0.001$. Error bars are s.e.m.

5 Fig. 5a illustrates a 3D rendering of well and moat geometry. Fig. 5b illustrates a rendering of 3D printed master for a 240-well example embodiment of the disclosed subject matter including alignment tabs. Fig. 5c illustrates a demolded PDMS device. Scale bar: 1 cm. Fig. 5d shows an example embodiment of the disclosed subject matter mounted in an OmniTray

10 Fig. 6 shows a comparison of image quality between standard 384-well plate and an example embodiment of the disclosed subject matter. Fig. 6a shows a bright field image of L4 worms crawling on agar in a polystyrene 384-well plate (Corning Costar). Worms are obscured by edges of wells. Darkfield images resulted in extremely poor image quality (not shown). Scale bar: 2 mm. Fig. 6b shows a darkfield
15 image of L4 worms on agar in an example embodiment of the disclosed subject matter. Well design enables more uniform illumination and easy identification of worms.

Fig. 7 illustrates that the well geometry of an example embodiment of the disclosed subject matter reduces burrowing and aversive moat prevents escape.
20 Fig. 7a shows the fraction of animals burrowing under the agar as a function of time in the example embodiment of the disclosed subject matter (red squares, $n=96$) and in 384-well microplate (blue circles, $n=80$). Fig. 7b shows the fraction of worms escaping wells as a function of time. A moat containing 100 mM CuSO_4 (red squares, $n=96$) is superior to a moat containing NGM buffer (blue circles, $N=96$) for
25 preventing animals from leaving their wells. NGM buffer is NGM agar without agar, peptone, or cholesterol.

Fig. 8 illustrates activity quantification. Consecutive images (left and center) are subtracted to yield a map of pixel gray-scale intensity changes. The absolute value of these subtracted grayscale values are shown in the difference image
30 (right). The number of pixels whose intensity changed above a threshold was summed up and the resultant value reported as 'activity'.

Fig. 9 illustrates automatic labeling of death time. Fig. 9a shows activity measurements throughout life of an individual N2 worm at 25° C. Fig. 9b illustrates that time of death was identified as the final moment the activity was greater than zero.

5 Fig. 10 illustrates the normalized healthspan of long-lived strains. Individual healthspan was divided by lifespan and strain averages are shown. Error bars are s.e.m. * $p < 0.05$; *** $p < 0.001$

Fig. 11 illustrates monitoring lethargus quiescence during development. Fig. 11a shows an activity trace for a single worm from L3 until adulthood. Fig. 11b shows the fraction of frames the worm in Fig. 11a was quiescent in a ten-minute moving window. Fig. 11c shows the fraction of quiescence heat map for a single 48-well example embodiment of the disclosed subject matter initially filled with L3 worms. Warmer colors indicate more quiescence and cooler colors indicate less quiescence. The fraction of quiescence traces for each animal were aligned for visualization. The L3 lethargus and L4 lethargus are visible as two bands of high quiescence.

Fig. 12 shows long-term fluorescent reporter tracking with an example embodiment of the disclosed subject matter. *mlt-10* is expressed in the hypodermis only during molting. Using a *mlt-10::gfp* reporter total worm fluorescence was tracked from the L3 stage to early adulthood. Heat map denotes fluorescence intensity in arbitrary units.

Fig. 13 illustrates measuring responses to aversive light stimulus. Violet light from an LED elicits an increase in speed in N2 worms. The light-sensing defective strain *lite-1* exhibits an attenuated response to light. Young adult animals were imaged sequentially for 30 seconds under bright field illumination. Violet light stimulation lasted 10 seconds and began 10 seconds after imaging commenced. Speed traces for each animal represent an average of 40 trials. Strain averages are denoted in thick lines, and individual traces are denoted by thin black lines for N2 and dashed blue lines for *lite-1*. N=6 animals for each strain.

30 Fig. 14 shows images of individual wells of example embodiments of the disclosed subject. Fig. 14a shows a brightfield image of a single well with adult worm and Fig. 14b shows a darkfield image of the same well.

DETAILED DESCRIPTION OF EMBODIMENTS

The present disclosure provides compositions and methods for highly scalable methods of cultivating and imaging organisms, including, but not limited to *C. elegans*.

5 A. Construction

In certain embodiments the present disclosure provides highly scalable compositions for the cultivation and imaging of *C. elegans*, other nematodes, zebrafish larvae, *Drosophila* larvae, among other organisms. In certain embodiments, the present disclosure provides compositions including microfabricated multi-well
10 arrays, including, but not limited to, certain embodiments identified here as the “WorMotel” (“WM”).

By conforming to the ANSI standard microplate format, certain embodiments of the of the present disclosure leverage existing automation technology such as robotic plate handlers, chemical library screening, and other tools. With
15 respect to embodiments relating to the cultivation and imaging of *C. elegans*, such existing automation technology also includes automated worm sorters.

In certain embodiments, the compositions, *e.g.*, WorMotels, are constructed by manually molding polydimethylsiloxane (PDMS) from a 3D-printed master. However, mass production using injection-molded transparent polymers such
20 as polyurethane, polycarbonate, polyvinyl, polystyrene, polyvinylchloride (PVC), polypropylene is also possible. In certain embodiments, the compositions are fabricated from any transparent material including PDMS, polystyrene, polycarbonate, and cyclic olefin. In certain embodiments, differences in thickness throughout the composition are acceptable. In certain embodiments uniform thickness throughout the
25 device is desirable. If uniform thickness throughout the device is desired, the composition can be produced via injection molding to achieve the desired uniform thickness.

With reference to Fig. 1, for the purpose of illustration and not limitation, in certain embodiments, the compositions of the present disclosure will
30 include a substrate 101 including an array of wells 102. In certain embodiments, the wells are arranged in relatively close proximity in a rectangular array or matrix. Common sizes for such arrays include a 4×4 matrix (16 wells; as illustrated in Fig.

1a), an 8×6 matrix (48 wells), an 8×12 matrix (96 wells), a 12×20 matrix (240 wells), and a 16×24 (384 wells), although larger arrays can also be used that include matrices of hundreds or even thousands of wells. The size and depth of each well can be varied according to the requirements of the particular assay being developed. For example, there is a tradeoff between size of the experiments and the total number of organisms that can be simultaneously studied. In certain embodiments, the wells have a diameter of about 2 mm to about 20 mm, e.g., 3 mm diameter, and a depth of about 2 mm to about 20 mm, e.g., a 3 mm depth, and a center-to-center spacing of about 2 mm to about 20 mm, e.g., 4.5 mm.

In certain embodiments the substrate including an array of wells will be a microplate. In certain embodiments the substrate including an array of wells will take the form of an insert that can be combined with a tray to provide a microwell assembly.

Conventional 96-well or 384-well microplates are not well-suited for imaging and cultivation due to three problems. First, the vertical walls of each well make it difficult to image the organisms when they are close to the edge of the wells. Second, in instances where the wells contain agar, organisms tend to crawl between the agar and well edges, making them difficult or impossible to image clearly. Third, in certain instances, organisms can climb over the walls of the wells, mixing with other organisms in other wells.

In certain embodiments, the well geometry can improve cultivation and imaging of one organism per well. For example, but not by way of limitation, a rounded concave well geometry (Fig. 1e) can reduce interference at the edge of the agar surface, and can prevent organisms from burrowing under the agar. In certain embodiments, the well array is a square array of a raised square boxes each containing a concave dimple. In certain embodiments, the space between the individual wells is a moat. In certain embodiments, the square box can be replaced with a cylinder containing the concave dimple. In certain of such embodiments, the moats are shaped like the space outside an array of circles.

As noted above, in certain embodiments, a moat surrounds each well. Such moats can function to prevent organisms from escaping from their individual wells. In certain embodiments, the moat disposed around each

depression is between about 0.3 mm and about 1 mm wide, e.g., 0.5 mm wide, and about 2 mm to about 20 mm deep, e.g., 3 mm deep. In certain embodiments, the moat is filled with an aversive compound. In certain embodiments the aversive compound will be: copper (e.g., as CuSO_4) or 2 % sodium dodecyl sulfate (SDS). In certain
5 embodiments, a moat containing 100 mM copper sulfate is employed.

In certain embodiments the composition is sealed. For example, but not by way of limitation, a composition pursuant to the instant disclosure can include a microplate or microwell assembly covered by a lid. In certain embodiments, the microplate or microwell assembly is wrapped with a sealing film. In certain
10 embodiments, the sealing film is gas permeable. In certain embodiments, Parafilm[®] is employed as the sealing film; however other gas permeable seals and tapes can also be used in the context of the subject matter disclosed herein.

To maintain humidity inside the microplates or microwell assemblies, water absorbing sodium polyacrylate crystals can be employed. Sterile distilled water
15 can be added to the crystals in a ratio of 30:1 (water:crystals) by weight. Approximately 15 g of hydrated crystals can then be added around the microplates or microwell assemblies.

B. Methods of Use

In certain embodiments, the present disclosure provides methods of
20 using the highly scalable compositions for cultivating and imaging organisms. In certain embodiments, the present disclosure provides for the use of such compositions for the long-term (>60 days) cultivation of organisms. For example, but not by way of limitation, the present disclosure provides methods of cultivating and imaging organisms, e.g., *C. elegans*, in studies of aging biology. The compositions of the
25 present disclosure, e.g., the WorMotel, are well-suited for other long-term assays including the quantification of behavior during development and the monitoring of fluorescent reporters.

The methods described herein can be a powerful and scalable platform for *C. elegans* aging research as well as research into aging of other organisms. By
30 tracking individually isolated organisms over their entire lifespans, the methods disclosed herein can enable longitudinal analysis of behavioral aging phenotypes. For example, as discussed herein, this analysis has uncovered previously undescribed

collapse-and-revival phenotypes in *C. elegans age-1* mutants. As disclosed herein, the instant methods can be used in understanding the genetic pathways and neural circuits governing aging quiescence in these and other strains.

5 In certain embodiments, detailed behavioral phenotypes, such as body posture, body bends, turning, and foraging can also be recorded by appropriately changing the camera's field of view and frame rate and employing existing tracking software.

Depending on the context, imaging can be carried out in a large range of spatial and temporal resolutions and over durations ranging from minutes to
10 months.

The techniques described herein can be used for performing high-throughput screening for genetic or pharmaceutical modulators of aging in *C. elegans* and other organisms. For example, but not by way of limitation, the WorMotel can be designed to work with standard automation tools such as the COPAS worm sorter,
15 liquid handlers, and robotic plate handlers. Furthermore, our approach is compatible with established RNAi protocols.

By employing near-standard culture conditions, the methods of the present disclosure can be directly compared with a large body of literature from aging studies performed on agar plates. Additionally, the organisms, *e.g.*, worms, can
20 remain experimentally accessible throughout their lifespans, and both the organisms and plates can be manipulated freely while maintaining the identities of the organisms under study.

In certain embodiments relating to assays of *C. elegans*, each well is filled with approximately 15 μ L of NGM agar and seeded with bacteria. A single
25 worm can be added to each well, either manually or using a COPAS Biosort (Union Biometrica) worm sorter. The rounded, aspheric geometry of each well can reduce interference from optical scattering at the edge of each well and to prevent worms from burrowing under the agar surface.

After loading the wells, the arrays can be imaged at 0.1 frames per
30 second or other desired temporal resolution under LED dark field illumination with an appropriate imaging system, *e.g.*, a 5 megapixel CMOS camera (Fig. 1c). In certain embodiments there is a trade-off between number of organisms and image resolution;

for example, but not by way of limitation, for a 240-well array the pixel resolution can be 36 μm when using a 5 megapixel CMOS camera. A plate handler robot with a microplate stacker can be used to serially image tens of thousands of organisms. Finally, a machine vision algorithm can quantify movement in each well between
 5 frames and records organism shape properties such as, in applications relating to *C. elegans*, centroid, area, and posture (Fig. 1d).

When the device is fabricated using a relatively high refractive index material, such as polystyrene ($n=1.55$) there can be high unwanted optical scattering from well due to a lensing effect of the rounded interface between the agar and
 10 concave well. This scattering can obscure the imaging of the organisms.

To address this problem, and with reference to Fig. 1f, a convex rounded surface can be provided on the bottom of the device, which is otherwise flat. The curvature of this surface can be set such that optical system consisting of the two refractive interfaces (agar-to-well-material and well-material-to-air) has a combined
 15 optical power of zero. This calculation can be performed using paraxial ray matrices or through an optical design software. For $n=1.55$, bottom of well radius of curvature $R_1=0.5$ mm, and a thickness t at the optical axis (center of well) of $t = 1\text{mm}$, the calculation gives an optimal radius of curvature of approximately $R_2 = 2.0$ mm as shown in Fig 1f.

20 EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the presently disclosed subject matter, and are not intended to limit the scope of what the inventors regard as their presently disclosed subject matter. It is understood that
 25 various other embodiments may be practiced, given the general description provided above.

A. A Scalable Platform For Long-Term Imaging Of Worm Lifespan And Behavior

Each WorMotel can consist of a transparent polydimethylsiloxane (PDMS) substrate containing a rectangular array of up to 240 wells (Fig. 1a, Fig. 5),
 30 each 3-mm in diameter. To fabricate the WorMotels a novel method of molding from a 3D-printed master (Fig. 1b) was developed. Each well can be filled with

approximately 15 μ L of NGM agar and seeded with bacteria, which the animal feeds on. A single worm can be added to each well, either manually or using a COPAS Biosort (Union Biometrica) worm sorter. The rounded, aspheric geometry of each well can be designed to reduce interference from optical scattering at the edge of each well and to reduce worms burrowing under the agar surface (Fig. 6). A narrow liquid-filled moat surrounding each well prevents animals from escaping from their wells. After loading the devices, the arrays can be imaged at 0.1 frames per second or other desired temporal resolution under bright field or dark field illumination with a CCD or CMOS camera (Fig. 1c). There is a trade-off between number of animals and image resolution; for a 240-well array and 5 megapixel camera the pixel resolution is 36 μ m. A plate handler robot with a microplate stacker can be used to serially image hundreds of plates and therefore tens of thousands of worms. Finally, a machine vision algorithm can quantify movement in each well between frames and records worm shape properties such as centroid, area, and posture (Fig. 1d).

Conventional 96-well or 384-well microplates are not well suited for worm imaging and cultivation on agar media due to three problems. First, the vertical walls of each well can make it difficult to image the worms when they are close to the edge of the wells (Fig. 6). Second, worms tend to crawl between the agar and well edges, again making them difficult to image clearly. Third, under humid conditions worms can climb over the walls of the wells, mixing with other worms in other wells.

The WorMotel uses well geometry for improved worm cultivation and imaging of one worm per well. A rounded concave well geometry can reduce interference at the edge of the agar surface, and can inhibit worms from burrowing under the agar. Less than 5% of young adult worms in the WorMotel burrow beneath the agar after 24 hours in comparison with over 80% of worms in a conventional 384-well microplate (Fig. 7). To prevent worms from leaving their designated wells, moats consisting of NGM liquid were tested, but after 3 days about 20% of worms escaped into the liquid. By contrast, only 5% worms escaped after 3 days when moat containing 100 mM copper sulfate was used (Fig. 7). Thus, the WorMotel can longitudinally monitor up to 240 worms under near standard laboratory conditions with an attrition rate of about 5% over the course of early adulthood.

(1) The WorMotel Can Generate Life-Long Behavior Profiles And Enable Automated Lifespan Analysis

To evaluate if the WorMotel could support worms throughout their lifespans, two strains were monitored: the reference strains N2 (“Bristol”) and CB4856 (“Hawaiian”). Image data from the WM aging experiments was used to continuously track behavioral history of each animal throughout its adult lifespan. Activity level, defined as the number of pixels that had changed by more than a certain threshold between temporally-adjacent images was plotted (Fig. 2a-b, Fig. 8). The time of death was determined as the final time at which activity greater than zero occurred between images acquired one minute apart (Fig. 2c-d, Fig. 9). Activity profiles were averaged across individuals to generate behavioral healthspan curves and compare them to survival data (Fig. 2e).

To determine whether the WorMotel accurately reproduced the lifespans of worms grown under standard laboratory conditions used in most aging experiments, the survival curves of worms on standard agar plates to those grown in WorMotels were compared. The lifespans of N2 were measured alongside the short-lived strain *daf-16* (*mu86*) and the long-lived strain *daf-2* (*e1370*). For worms grown on standard plates, lifespan assays were carried out using standard methods. Mean lifespan of *daf-16* animals (WM: 7.74 ± 0.29 days, n=61; Manual: 6.98 ± 0.15 days, n=117) was shorter than that of N2 (WM: 12.3 ± 0.33 days, n=123; Manual: 12.15 ± 0.51 days, n=94) while *daf-2* showed a longer lifespan (WM: 33.5 ± 1.94 days, n=46, Manual: 30.77 ± 1.95 days, n=52) than N2. No significant difference between survival curves acquired from worms grown on standard plates and those grown on the WorMotel were found (Fig. 2f). These results show the WorMotel can reproduce expected lifespans for wild type as well as both short and long-lived strains and can monitor individual animals for at least 60 days.

Next the compatibility of the WorMotel with methods for RNA interference was tested. Using RNAi by feeding, the lifespan of N2 in which either DAF-16 or DAF-2 was knocked down was measured. As a control, lifespans of N2, *daf-16*(*mu86*), and *daf-2*(*e1370*) fed with *E. coli* carrying an RNAi plasmid without a gene insert (“empty vector”) were also measured. *daf-16*(RNAi) and *daf-16*(*mu86*) animals had a lifespan shorter than the empty vector control, and *daf-2*(RNAi) and *daf-*

2(*e1370*)) animals lived longer than controls (Fig. 2g). These results show that the WorMotel platform is suitable for RNAi-based screening.

5 (2) Locomotion Profiles Can Predict Individual Worm Survival

In order to characterize the utility of the locomotor activity profiles produced by the WorMotel, the three short-lived mutants *daf-16* (*mu86*), *mir-71* (*n4115*), and *aak-2* (*ok524*) alongside N2 were monitored. Animals of these three mutant genotypes exhibited declines in their locomotion and died earlier than N2 animals (Fig. 3a-d, f, g) (N2 301; *daf-16* 302; *mir-71* 303; *aak-2* 304). The average survivor activity over time was plotted, however, short-lived mutants seemed to be healthier for a greater fraction of their lives compared to N2 (Fig. 3e). Each individual's healthspan was defined as the last time point at which its locomotion was greater than 50% of its maximum value. The fraction alive was plotted alongside the fraction healthy over time (Fig. 3f-g). Healthspan was well-correlated with lifespan for individuals within each strain (Fig. 3e-g). In all four strains, the curve depicting the fraction of healthy worms was shifted to the left of the curve depicting the fraction of live worms. As suggested by Figures 3e-g, the magnitude of this shift was greater for N2 than for the three short-lived mutants. These data suggest that while N2 animals live longer than these short-lived mutants, its fraction of life spent in good health might not be similarly prolonged.

The ability to track single animal trajectories throughout the adult life of the animal allowed an analysis of plotting individual lifespans as a function of their individual healthspan. The slope of a linear fit of these data was smaller for N2 animals ($m=0.74$) than for short-lived mutants (*daf-16*: $m=0.92$; *aak-2*: $m=1.09$; *mir-71*: $m=0.96$), suggesting that lifespan depends less strongly on healthspan in N2. Finally, the degree to which healthspan predicted lifespan, as reflected by the R^2 values of the least-squared linear regression fit, were larger for short-lived mutants (*daf-16* $R^2 = 0.71$; *aak-2* $R^2 = 0.69$; *mir-71* $R^2 = 0.34$) than for N2 ($R^2 = 0.12$). Choosing a more lenient threshold for healthy activity, for example, 25% of maximum activity rather than 50%, was more predictive of lifespan for all these strains (N2 $R^2 = 0.55$; *daf-16* $R^2 = 0.86$; *aak-2* $R^2 = 0.87$; *mir-71* $R^2 = 0.61$). This data shows that

behavioral data from the WorMotel reflects the health of individual animals at specific time points during aging.

The rate of decay in locomotion during days 3 through 7 can be negatively correlated with lifespan. However, it was unclear whether the locomotion profiles produced by the WM could be used in this way to predict individual life spans. Activity decay rate was calculated as the negative slope of a line fit to the activity of individual worms early in life, defined as the time until 25% of each strain's animals had died, which was day 6 for the short-lived strains and day 13 for N2. We then plotted lifespan against the activity decay rate (Fig. 3l-o). Lifespan was indeed negatively correlated with early life activity decay rate (*N2*: $R = 0.70$, $p < 10^{-8}$; *daf-16*: $R = 0.61$, $p < 10^{-5}$; *aak-2*: $R = 0.42$, $p < 0.005$; *mir-71*: $R = 0.69$, $p < 10^{-8}$). These results indicate that activity measurements from the WM reflect the health trajectories of individual animals and can be used to identify healthy animals early in life.

(3) Monitoring Of Individual Behavioral Profiles Can Reveal Unexpected Aging Phenotypes

Some long-lived mutant strains exhibited striking differences in their behavioral aging profiles (Fig. 4a-d). Behavior of a mutant in the gene *age-1*, whose gene product encodes the *C. elegans* homolog of mammalian phosphatidylinositol-3-OH kinase (PI(3)K) catalytic subunit, was examined. Like other mutants in the insulin/insulin-like (IIS) signaling pathway, *age-1(hx546)* animals have extended lifespan. *age-1* worms show a decline in activity on around day 8 (day of initial decline = 8.5 ± 0.3 days) followed by a quiescent period of about 3 days (quiescent duration = 2.8 ± 0.5 days, Fig. 4e-f, l) during which worms exhibit essentially no movement. This quiescent period is followed by a revival during which animals resume activity approximately half the amplitude of the worm's peak activity from early adulthood (peak revival activity = $44 \pm 4\%$ of peak early life activity, Fig. 4e). Revivals occur in 72% (34/47) worms tested. This non-monotonic behavioral profile in *age-1* worms supports the notion that long-term behavioral imaging can uncover novel phenotypes associated with aging.

Whether this mid-life quiescence in *age-1* was common to all loss-of-function mutants in the IIS pathway was tested. The behavioral profile of mutants for

the insulin receptor homolog *daf-2*, in which adult quiescence has previously been reported, was tested. *daf-2 (e1370)* worms exhibit a precipitous drop in activity around day 10 (Fig. 4d, g, h, l) and after that remain largely quiescent except for brief low-level bursts of activity until they die. By plotting survivor average activity along with survival it can be observed that the behavioral profiles of *age-1* and *daf-2* are different: the asynchronous revivals of *age-1* animals lead to an extended period of activity during days 10-20, which was absent in *daf-2* animals (Fig. 4i). Therefore, while *age-1* and *daf-2* worms both display more quiescence than N2, the collapse-and-revival behavior of *age-1* mutants appears to be distinct from the pattern of quiescence in *daf-2* animals.

In order to determine the relationship between healthspan and lifespan, survival for N2, *age-1*, and multiple alleles of *daf-2*, was plotted alongside the healthy fraction as determined by animals moving at greater than 50% of their maximum activity level. After normalizing individual healthspan by lifespan, only *age-1(hx546)* and *daf-2(e1370)* exhibit significant differences from N2 (Fig. 10). In order to further characterize the quiescence in these strains, the day of initial decline, the first time period of detectable prolonged quiescence, was quantified. All tested insulin-signaling mutants initially decline well before N2 (Fig. 4l) while most retain the ability to move normally (Fig. 4k). When the day of initial decline was normalized by the lifespan of individual animals, N2 first becomes quiescent on average at an age about 60% of its lifespan while *age-1* and *daf-2* mutants first become quiescent at an age between 30% and 40% of their lifespans (Fig. 4m). It is possible that prolonged quiescence can play a causal role in the long lifespan of insulin-signaling mutants. The WorMotel can play an important role in illuminating the role of such long-term phenotypes in extended longevity.

In addition to the applications to aging described here, the WorMotel method can be suited for other experiments in which longitudinal monitoring is desired. The disclosed subject matter has shown experiments in which the WorMotel was used to measurement of quiescence during development (Fig. 11), long-term monitoring of fluorescent reporters (Fig. 12) and monitoring responses to aversive stimuli (Fig. 13).

B. Construction & Use of WorMotel

(1) WorMotel Design and Fabrication

Using MATLAB scripts, a chip containing a rectangular array of either 48 or 240 rounded wells with 3 mm diameter, 3 mm depth, and center-to-center spacing of 4.5 mm (Fig. 5a) was designed. Each well was surrounded by a 0.5 mm wide and 3 mm deep 'moat'. A master corresponding to the negative of this shape in the material VeroBlack was fabricated using an Objet30 photopolymer 3D printer (Stratasys). (Fig. 5b).

To mold the PDMS devices, Sylgard 184 PDMS (Dow Corning) was mixed according to the manufacturer's instructions, bubbles were removed by vacuum degassing for 15 min, and then an appropriate amount of PDMS was poured into the masters. Devices were cured overnight at 40°C then removed from molds using a spatula. The first device cast from each master often did not de-mold cleanly, but subsequent casts were successful (Fig. 5c).

(2) WorMotel Preparation

To prepare devices for experiments, the chips were first treated with oxygen or air plasma for 3 minutes using a plasma cleaner (PE-50, Plasma Etch Inc.). This treatment can render PDMS temporarily hydrophilic, which can improve the ease of filling of wells and moats. The medium was based on standard NGM media except low-gelling temperature agarose (Gelling temp. 26-30°C, Research Products International) was substituted for agar to facilitate filling by pipette and streptomycin (200 ng/mL) was added to the medium to minimize bacterial contamination.

For aging experiments 5-fluoro-2'-deoxyuridine (FUdR) was added to prevent progeny growth. An FUdR stock solution of 10 mg/ml was prepared and added to molten agar at a concentration of 2.5 μ L per mL just prior to filling. This yielded a final concentration of FUdR of 100 μ M.

After the agar cooled and gelled, a suspension of the *Escherichiae coli* bacterial strain DA837, which is a streptomycin resistant derivative of OP50 (Brenner, S. The Genetics of *C. elegans*. *Genetics* **77**, 71–94 (1974)) was added to each well using a 200 μ L pipette.

A moat solution of 100 mM CuSO_4 was prepared, which was approximately in osmotic equilibrium with the agar medium. The moat solution was added to the moats by pipette.

For aging experiments, late L4 worms were added to the WorMotel
5 either manually with a platinum wire pick or automatically using a COPAS Biosort machine (Union Biometrica).

PDMS devices were placed inside either a 90 mm diameter petri dish for 48-well WorMotels or an OmniTray microplate (Nunc Thermo Scientific) for 240-well. 240-well WorMotels contained alignment tabs to keep devices in alignment
10 with respect to the OmniTray (Fig. 5d). To maintain humidity inside the dishes, water absorbing sodium polyacrylate crystals were used. Sterile distilled water was added to the crystals in a ratio of 30:1 (water:crystals) by weight. Approximately 15 g of hydrated crystals were added around the WorMotel. Lids were placed on all dishes. For aging experiments, a single layer of fully-stretched Parafilm (Bemis Co.) was
15 wrapped around the sides of the plate to further reduce water loss while permitting gas exchange.

(3) Image Acquisition

Images were captured with an Imaging Source DMK 72AUC02 camera (2592 x 1944 pixels) equipped with a Fujinon lens (HF12.5SA-1, 1:1.4/12.5 mm). IC
20 Capture software (The Imaging Source) was used to acquire time lapse images. All experiments were carried out under dark-field illumination using four 4.7" long red LED strips (Oznum) positioned approximately 2" below the WorMotel. Images were saved and processed by a 64-bit computer with a 3.40 GHz Intel Core i3 processor and 4 GB of RAM. Images were analyzed using custom MATLAB software.

25 Different spatial resolutions can be attained by adjusting the camera's field of view and thus by modulating the number of wells viewed at once. Imaging 6 wells at once gives approximately 5 μm resolution, imaging 12 wells gives 7 μm resolution, imaging 48 wells gives 15 μm resolution, and imaging 240 wells gives 36 μm resolution.

30 (4) Image Processing

Temporally adjacent images were subtracted to generate maps of pixel value intensity change (Fig. 8). A Gaussian smoothing filter with width 3-5 pixels

was applied to the resulting image in order to reduce camera noise. A binary threshold was then applied to the filtered intensity change image in order to score whether or not movement occurred at each pixel: those pixels which changed by much more than the noise level of the camera. All pixels in which movement occurred were counted and the resulting value was called the ‘activity’ between the two frames.

(5) Removal of camera artifacts

Approximately 0.17% of frames, in which camera errors resulted in shifted or distorted images, were censored. These events were automatically identified based on activity spikes above threshold occurring simultaneously across all regions of interest.

(6) Automatic death quantification

Time of death was quantified as the final time point during which the activity was greater than zero (Fig. 9).

(7) Healthspan calculation

Activity profiles were normalized to the maximum for each individual worm. The healthspan was defined as the final time point at which the individual’s activity value was greater than or equal to 50% of its maximum value.

(8) Lifespan prediction by activity decay rate

Early life was defined as day 2 until the 25th percentile of lifespan for each strain tested. This time point corresponded to day 13 for N2, day 6 for *daf-16*, day 6 for *aak-2*, and day 6 for *mir-71*. For each individual, the normalized activity during early life was fit by a linear function. The negative slope of the fit line was taken as the early life activity decay rate. Lifespan was then plotted against early life activity decay rate. A line was fit to the data and the R^2 value was generated for each strain (Fig. 3l-o).

(9) Fraction of quiescence and day of initial decline calculation

The fraction of quiescence was calculated by tallying the fraction of frames in a two-hour window during which the activity was less than a specified threshold (Fig. 4). The day of initial decline was calculated for each individual by measuring the first time point at which the fraction of quiescence rose above 0.5.

(10) Automated Imaging of Multiple Plates

A plate handler robot (Caliper Twister II) was used to automatically image multiple plates per day by sequentially lifting each plate from a stack and positioning it inside a dark field imaging rig for 10 minutes (Fig. 1c). The Twister II has a capacity of 400 plates, and can monitor 96,000 individual worms. Multiple OmniTrays containing prepared WorMotels were placed in a 'home' stack. The robot lifted each plate and positioned it in a darkfield imaging rig for 10 minutes at which point the plate was returned to an 'away' stack. Once all plates in the 'home' stack had been imaged, the plates in the 'away' stack were returned to the 'home' stack and the loop began again. The camera acquired images continuously throughout this plate rotation process. Plates were uniquely identified in each image via a printed barcode located at the top of each plate.

Although the foregoing presently disclosed subject matter has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the presently disclosed subject matter. The disclosures of all patent and scientific literature cited herein are expressly incorporated in their entirety by reference.

WHAT IS CLAIMED IS:

1. A composition, comprising:
 - a substrate having a planar array of depressions each defined by concave walls;
 - and
 - a moat disposed around each depression of said array of depressions.
2. The composition of claim 1, wherein the depressions have a 3 mm diameter, a 3 mm depth, and a center-to-center spacing of 4.5 mm.
3. The composition of claim 2, wherein the moat disposed around each depression is 0.5 mm wide and 3 mm deep.
4. The composition of claim 1, wherein the composition is made of at least one of polyeurathene, polycarbonate, polyvinyl, polystyrene, polyvinylchloride (PVC), polypropylene, polydimethylsiloxane (PDMS), and cyclic olefin copolymer (COC).
5. The composition of claim 1, wherein said substrate having a planar array of depressions is designed to insert into a tray to form a microwell assembly.
6. The composition of claim 1, wherein the substrate is of uniform thickness.
7. The composition of claim 6, wherein the plate is made entirely of polyeurathene, polycarbonate, polyvinyl, polystyrene, polyvinylchloride (PVC), polypropylene, or cyclic olefin copolymer (COC).
8. The composition of claim 7, wherein the plate is made of polystyrene.
9. The composition of claim 1, wherein the plate has an overall shape and dimension generally conforming to a standard ANSI/SLAS microplate.
10. The composition of claim 9, wherein the standard microplate comprises at least one of a 48-well microplate having a 8×6 array of wells; a 96-well microplate having a 8×12 array of wells; a 240-well microplate having a 12×20 array of wells; and a 384-well microplate having a 16×24 array of wells.
11. A method of cultivating and imaging an organism, comprising:
 - providing a composition comprising:
 - a substrate having a planar array of depressions each defined by concave walls; and
 - a moat disposed around each depression of said array of depressions

introducing an organism into a depression of said substrate; and
cultivating and imaging said organism.

12. The method of claim 11, wherein the organism is selected from the group consisting of *C. elegans*, other nematodes, zebrafish larvae, and *Drosophila* larvae.




FIG. 1a




FIG. 1b

2/25





FIG. 1c

Raw



Background Subtracted




FIG. 1d

Well shape

All dimensions in mm

3/25




FIG. 1e




FIG. 1f




FIG. 2a




FIG. 2b




FIG. 2c




FIG. 2d

5/25




FIG. 2e




FIG. 2f




FIG. 2g




FIG. 3a




FIG. 3b




FIG. 3c




FIG. 3d

7/25




FIG. 3e




FIG. 3f




FIG. 3g

8/25




FIG. 3h




FIG. 3i




FIG. 3j




FIG. 3k

9/25




FIG. 3l




FIG. 3m




FIG. 3n




FIG. 3o

10/25




FIG. 4a




FIG. 4b




FIG. 4c




FIG. 4d

11/25




FIG. 4e




FIG. 4f




FIG. 4g




FIG. 4h

12/25




FIG. 4i




FIG. 4j




FIG. 4k




FIG. 4l




FIG. 4m

14/25




FIG. 5b




FIG. 5d




FIG. 5a




FIG. 5c

15/25




FIG. 6a




FIG. 6b

16/25




FIG. 7a




FIG. 7b

17/25




FIG. 8

18/25




FIG. 9a




FIG. 9b

19/25




FIG. 10

20/25




FIG. 11a




FIG. 11b

21/25




FIG. 11c

22/25




FIG. 12

23/25




FIG. 13

24/25




FIG. 14A




FIG. 14B

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US15/37997

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - B01L 3/14; C40B 30/06, 60/12 (2015.01)

CPC - B01L 3/5085, 3/50855; C40B 30/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - B01L 3/14; C40B 30/06, 60/12 (2015.01)

CPC - B01L 3/5085, 3/50855, 2300/0829, 3/50857; C40B 30/06

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatSeer (US, EP, WO, DE, GB, FR, KR, ES, CA); Google Scholar; ProQuest; PubMed. Keywords: microplate, microtiter, microwell, depression, wells, concave, curve, moat, channel, trough, trench, cultivate, culture, imaging, picture, camera, wide, width, deep, depth

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y --- A	US 7,998,435 B2 (REED, MT) 16 August 2011; figures 22a, 26, 32, 150-151; column 9, lines 28-38; page 10, lines 11-19; column 14, lines 33-35; column 15, lines 34-43; column 20, lines 60-67; column 21 lines 1-3; column 39, lines 3-22	1-2, 4-12 ----- 3
Y	EP 0 353 591 B1 (ABBOTT LABORATORIES) 17 April 1996; figures 5, 8; page 10, lines 11-19	1-2, 4-12
Y	US 2007/0237683 A1 (HO, WZ et al.) 11 October 2007; abstract; paragraphs [0031], [0035]	2, 5
Y	US 6,051,191 A (IRELAND, RM) 18 April 2000; column 3, lines 31-54; column 5, lines 54-61; column 6, lines 62-63	8, 10
Y	(Yanik, FY et al.) Technologies for Micromanipulating, Imaging, and Phenotyping Small Invertebrates and Vertebrates. Annual Review of Biomedical Engineering. August 2011, Vol. 13, pages 185-217; page 193, figure 4; page 186, second paragraph; page 193, fifth-sixth paragraphs; page 203, first paragraph	11-12
Y	WO 1998/046981 A1 (LJL BIOSYSTEMS) 22 October 1998; page 7, lines 2-5	2
A	US 2004/0228770 A1 (GHANDHI, K et al.) 18 November 2004; figure 4A; paragraph [0054]	3
A	WO 2004/018623 A2 (CLINICAL MICROARRAYS, INC., et al.) 04 March 2004; figure 24A; page 42, lines 4-6; page 43, lines 12-13	3
A	US 6,908,594 B1 (SCHAEVITZ, SB et al.) 21 June 2005; figure 1A; column 3, lines 10-14, 49-55; column 4, lines 34-45	3

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

31 August 2015 (31.08.2015)

Date of mailing of the international search report

30 SEP 2015

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774