



US 20040242516A1

(19) **United States**

(12) **Patent Application Publication**
Crooke et al.

(10) **Pub. No.: US 2004/0242516 A1**

(43) **Pub. Date: Dec. 2, 2004**

(54) **ANTISENSE MODULATION OF
APOLIPOPROTEIN(A) EXPRESSION**

(76) Inventors: **Rosanne M Crooke**, Carlsbad, CA
(US); **Mark J Graham**, San Clemente,
CA (US)

Correspondence Address:
Mary E Bak
Howson and Howson
Spring House Corporate Center
Box 457
Spring House, PA 19477 (US)

(21) Appl. No.: **10/485,113**

(22) PCT Filed: **Aug. 5, 2002**

(86) PCT No.: **PCT/US02/24920**

Related U.S. Application Data

(63) Continuation of application No. 09/923,515, filed on
Aug. 7, 2001.

Publication Classification

(51) **Int. Cl.⁷** **A61K 48/00**; C07H 21/04

(52) **U.S. Cl.** **514/44**; 536/23.5

(57) **ABSTRACT**

Antisense compounds, compositions and methods are provided for modulating the expression of apolipoprotein(a). The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding apolipoprotein(a). Methods of using these compounds for modulation of apolipoprotein(a) expression and for treatment of diseases associated with expression of apolipoprotein(a) are provided.

ANTISENSE MODULATION OF APOLIPOPROTEIN(A) EXPRESSION

FIELD OF THE INVENTION

[0001] The present invention provides compositions and methods for modulating the expression of apolipoprotein(a). In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding apolipoprotein(a). Such compounds have been shown to modulate the expression of apolipoprotein(a).

BACKGROUND OF THE INVENTION

[0002] Lipoproteins are globular, micelle-like particles that consist of a non-polar core of acylglycerols and cholesteryl esters, surrounded by an amphiphilic coating consisting of protein, phospholipid and cholesterol. Lipoproteins have been classified into five broad categories on the basis of their functional and physical properties: chylomicrons (which transport dietary lipids from intestine to tissues), very low density lipoproteins (VLDL), intermediate density lipoproteins (IDL), low density lipoproteins (LDL), (all of which transport triacylglycerols and cholesterol from the liver to tissues), and high density lipoproteins (HDL) (which transport endogenous cholesterol from tissues to the liver).

[0003] Lipoprotein particles undergo continuous metabolic processing and have variable properties and compositions. Lipoprotein densities increase without decreasing particle diameter because the density of their outer coatings is less than that of the inner core. The protein components of lipoproteins are known as apolipoproteins. At least nine apolipoproteins are distributed in significant amounts among the various human lipoproteins.

[0004] Lipoprotein(a) (also known as Lp(a)) is a cholesterol rich particle of the pro-atherogenic LDL class. Because Lp(a) is found only in Old World primates and European hedgehogs, it has been suggested that it does not play an essential role in lipid and lipoprotein metabolism. Most studies have shown that high concentrations of Lp(a) are strongly associated with increased risk of cardiovascular disease (Rainwater and Kammerer, *J. Exp. Zool.*, 1998, 282, 54-61). These observations have stimulated numerous studies in humans and other primates to investigate the factors that control Lp(a) concentrations and physiological properties (Rainwater and Kammerer, *J. Exp. Zool.*, 1998, 282, 54-61).

[0005] Lp(a) contains two disulfide-linked distinct proteins, apolipoprotein(a) (or ApoA) and apolipoprotein B (or ApoB) (Rainwater and Kammerer, *J. Exp. Zool.*, 1998, 282, 54-61). Apolipoprotein(a) is a unique apolipoprotein encoded by the LPA gene which has been shown to exclusively control the physiological concentrations of Lp(a) (Rainwater and Kammerer, *J. Exp. Zool.*, 1998, 282, 54-61). It varies in size due to interallelic differences in the number of tandemly repeated Kringle 4-encoding 5.5 kb sequences in the LPA gene (Rainwater and Kammerer, *J. Exp. Zool.*, 1998, 282, 54-61).

[0006] Cloning of human apolipoprotein(a) in 1987 revealed homology to human plasminogen (McLean et al., *Nature*, 1987, 330, 132-137). The gene locus LPA encoding apolipoprotein(a) was localized to chromosome 6q26-27, in

close proximity to the homologous gene for plasminogen (Frank et al., *Hum. Genet.*, 1988, 79, 352-356).

[0007] Transgenic mice expressing human apolipoprotein(a) were found to be more susceptible than control mice to the development of lipid-staining lesions in the aorta and, consequently, apolipoprotein(a) is co-localized with lipid deposition in the artery walls (Lawn et al., *Nature*, 1992, 360, 670-672). As an extension of these studies, it was established that the major in vivo action of apolipoprotein(a) is inhibition of the conversion of plasminogen to plasmin which causes decreased activation of latent transforming growth factor-beta. Because transforming growth factor-beta is a negative regulator of smooth muscle cell migration and proliferation, inhibition of plasminogen activation indicates a possible mechanism for apolipoprotein(a) induction of atherosclerotic lesions (Grainger et al., *Nature*, 1994, 370, 460-462).

[0008] Elevated plasma levels of Lp(a), caused by increased expression of apolipoprotein(a), are associated with increased risk for atherosclerosis and its manifestations, which include hypercholesterolemia (Seed et al., *N. Engl. J. Med.* 1990, 322, 1494-1499), myocardial infarction (Sandkamp et al., *Clin. Chem.*, 1990, 36, 20-23), and thrombosis (Nowak-Gottl et al., *Pediatrics*, 1997, 99, E11).

[0009] Moreover, the plasma concentration of Lp(a) is strongly influenced by heritable factors and is refractory to most drug and dietary manipulation (Katan and Beynen, *Am. J. Epidemiol.*, 1987, 125, 387-399; Vessby et al., *Atherosclerosis*, 1982, 44, 61-71). Pharmacologic therapy of elevated Lp(a) levels has been only modestly successful and apheresis remains the most effective therapeutic modality (Hajjar and Nachman, *Annu. Rev. Med.*, 1996, 47, 423-442).

[0010] Morishita et al. have reported the use of ribozyme oligonucleotides against apolipoprotein(a) for inhibition of apolipoprotein(a) expression in HepG2 cells (Morishita et al., *Circulation*, 1998, 98, 1898-1904).

[0011] Disclosed and claimed in U.S. Pat. No. 5,721,138 are nucleotide sequences encoding the human apolipoprotein(a) gene 5'-regulatory region and isolated nucleotide sequences comprising at least thirty consecutive complementary nucleotides from human apolipoprotein(a) from nucleotide position -208 to -1448 (Lawn, 1998).

[0012] To date, investigative and therapeutic strategies aimed at inhibiting apolipoprotein(a) function have involved the previously cited use of Lp(a) apheresis and ribozyme oligonucleotides. Consequently, there remains a long-felt need for additional agents capable of effectively inhibiting apolipoprotein(a) function.

[0013] Antisense technology is emerging as an effective means of reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic and research applications involving modulation of apolipoprotein(a) expression.

[0014] The present invention provides compositions and methods for modulating apolipoprotein(a) expression.

SUMMARY OF THE INVENTION

[0015] The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding apolipoprotein(a), and which modu-

late the expression of apolipoprotein(a). Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of apolipoprotein(a) in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of apolipoprotein(a), by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

DETAILED DESCRIPTION OF THE INVENTION

[0016] The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding apolipoprotein(a), ultimately modulating the amount of apolipoprotein(a) produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding apolipoprotein(a). As used herein, the terms "target nucleic acid" and "nucleic acid encoding apolipoprotein(a)" encompass DNA encoding apolipoprotein(a), RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of apolipoprotein(a). In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

[0017] It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multi step process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding apolipoprotein(a). The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules;

5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding apolipoprotein(a), regardless of the sequence(s) of such codons.

[0018] It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

[0019] The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

[0020] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an

overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

[0021] Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

[0022] In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

[0023] Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

[0024] Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

[0025] For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with other antisense compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

[0026] Expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds which affect expression patterns.

[0027] Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, 2000, 480, 17-24; Celis, et al., *FEBS Lett.*, 2000, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., *Drug Discov. Today*, 2000, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, 1999, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2000, 97, 1976-81), protein arrays and proteomics (Celis, et al., *FEBS Lett.*, 2000, 480, 2-16; Jungblut, et al., *Electrophoresis*, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, 2000, 480, 2-16; Larsson, et al., *J. Biotechnol.*, 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., *Anal. Biochem.*, 2000, 286, 91-98; Larson, et al., *Cytometry*, 2000, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, 2000, 3, 316-21), comparative genomic hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*, 1998, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, *Eur. J. Cancer*, 1999, 35, 1895-904) and mass spectrometry methods (reviewed in (To, *Comb. Chem. High Throughput Screen*, 2000, 3, 235-41).

[0028] The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

[0029] In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

[0030] While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

[0031] As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

[0032] Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides. Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thiono-alkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2' -5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

[0033] Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

[0034] Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbone; and others having mixed N, O, S and CH₂ component parts.

[0035] Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

[0036] In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

[0037] Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular —CH₂—NH—O—CH₂—, —CH₂—N(CH₃)—O—CH₂— [known as a methylene (methylimino) or MMI backbone], —CH₂—O—N(CH₃)—CH₂—, —CH₂—N(CH₃)—N(CH₃)—CH₂— and —O—N(CH₃)—CH₂—CH₂—

[wherein the native phosphodiester backbone is represented as $—O—P—O—CH_2—$] of the above referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above referenced U.S. Pat. No. 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

[0038] Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O—, S—, or N-alkyl; O—, S—, or N-alkenyl; O—, S— or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C_1 to C_{10} alkyl or C_2 to C_{10} alkenyl and alkynyl. Particularly preferred are $O[(CH_2)_nO]_mCH_3$, $O(CH_2)_nOCH_3$, $O(CH_2)_nNH_2$, $O(CH_2)_nCH_3$, $O(CH_2)_nONH_2$, and $O(CH_2)_nON[(CH_2)_mCH_3]_2$, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C_1 to C_{10} lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH_3 , OCN, Cl, Br, CN, CF_3 , OCF_3 , $SOCH_3$, SO_2CH_3 , ONO_2 , NO_2 , N_3 , NH_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy ($2'-O—CH_2CH_2OCH_3$, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoxyethoxy, i.e., a $O(CH_2)_2ON(CH_3)_2$ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylamino-ethoxyethoxy (also known in the art as 2'-O-dimethylamino-ethoxyethyl or 2'-DMAEOE), i.e., $2'-O—CH_2—O—CH_2—N(CH_2)_2$, also described in examples hereinbelow.

[0039] A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene ($—CH_2—$)_n group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

[0040] Other preferred modifications include 2'-methoxy ($2'-O—CH_3$), 2'-aminopropoxy ($2'-OCH_2CH_2CH_2NH_2$), 2'-allyl ($2'-CH_2—CH=CH_2$), 2'-O-allyl ($2'-O—CH_2—CH=CH_2$) and 2'-fluoro ($2'-F$). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,

747; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

[0041] Oligonucleotides may also include nucleobase (often referred to in the art simply as “base”) modifications or substitutions. As used herein, “unmodified” or “natural” nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ($—C≡C—CH_3$) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine(1H-pyrimido [5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido [5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrimido[3', 2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J. I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S. T. and Lebleu, B. ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyl-adenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2° C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0042] Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos. 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,645,985; 5,830,653;

5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and U.S. Pat. No. 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

[0043] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, poly-ethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterol, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International patent application PCT/US92/09196, filed Oct. 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxysterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937). Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folic acid, a benzothiadiazide, chlorothiazide, a diazepam, indo-methicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in U.S. patent application Ser. No. 09/334,130 (filed Jun. 15, 1999) which is incorporated herein by reference in its entirety.

[0044] Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

[0045] It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

[0046] Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

[0047] The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis.

Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

[0048] The antisense compounds of the invention are synthesized *in vitro* and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the *in vivo* synthesis of antisense molecules.

[0049] The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

[0050] The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

[0051] The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO-93/24510 to Gosselin et al., published Dec. 9, 1993 or in WO 94/26764 and U.S. Pat. No. 5,770,713 to Imbach et al.

[0052] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

[0053] Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma. Sci.*, 1977, 66, 1-19). The base addition

salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

[0054] For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

[0055] The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal,

preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of apolipoprotein(a) is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

[0056] The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding apolipoprotein(a), enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding apolipoprotein(a) can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of apolipoprotein(a) in a sample may also be prepared.

[0057] The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

[0058] Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid,

capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C₁₋₁₀ alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in U.S. patent application Ser. No. 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

[0059] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodehydrofusidate. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pullulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyrrolidone, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in detail in U.S. applications Ser. No. 08/886,829 (filed Jul. 1, 1997), Ser. No. 09/108,673 (filed Jul. 1, 1998), Ser. No. 09/256,515 (filed Feb. 23, 1999), Ser. No. 09/082,624 (filed May 21, 1998) and Ser. No. 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

[0060] Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[0061] Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

[0062] The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0063] The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0064] In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

[0065] Emulsions

[0066] The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of droplets usually exceeding $0.1 \mu\text{m}$ in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed

as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

[0067] Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

[0068] Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: non-ionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

[0069] Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous

preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

[0070] A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

[0071] Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

[0072] Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

[0073] The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

[0074] In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be

defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 1985, p. 271).

[0075] The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

[0076] Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

[0077] Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

[0078] Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories—surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

[0079] Liposomes

[0080] There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term “liposome” means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

[0081] Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages in vivo.

[0082] In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable

transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

[0083] Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

[0084] Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

[0085] Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

[0086] Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

[0087] Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

[0088] Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

[0089] One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome

compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

[0090] Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

[0091] Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

[0092] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{M1}, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G_{M1}, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Pat. No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G_{M1} or a galactocerebroside sulfate ester. U.S. Pat. No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

[0093] Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C₁₂15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Pat. Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Pat. Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Pat. No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Pat. No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Pat. No. 5,540,935 (Miyazaki et al.) and U.S. Pat. No. 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

[0094] A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Pat. No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Pat. No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

[0095] Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

[0096] Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophilic/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[0097] If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

[0098] If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

[0099] If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

[0100] If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

[0101] The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[0102] Penetration Enhancers

[0103] In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

[0104] Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty

acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

[0105] Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

[0106] Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C₁₋₁₀ alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and diglycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

[0107] Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), gluchoic acid (sodium gluchocholate), glycholic acid (sodium glychocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium-taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, Pa., 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

[0108] Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as

compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

[0109] Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

[0110] Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Pat. No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

[0111] Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

[0112] Carriers

[0113] Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity per se) but is recognized as a nucleic acid by in vivo processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in

hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4' isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

[0114] Excipients

[0115] In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

[0116] Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

[0117] Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

[0118] Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

[0119] Other Components

[0120] The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preserva-

tives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

[0121] Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0122] Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylecyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphamide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

[0123] In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

[0124] The formulation of therapeutic compositions and their subsequent administration is believed to be within the

skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC_{50} s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

[0125] While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

EXAMPLES

Example 1

Nucleoside Phosphoramidites for Oligonucleotide Synthesis Deoxy and 2'-alkoxy amidites

[0126] 2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham Mass. or Glen Research, Inc. Sterling Va.). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Pat. No. 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

[0127] Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods (Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203) using commercially available phosphoramidites (Glen Research, Sterling Va. or ChemGenes, Needham Mass.).

2'-Fluoro amidites

2'-Fluorodeoxyadenosine amidites

[0128] 2'-fluoro oligonucleotides were synthesized as described previously (Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841) and U.S. Pat. No. 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoro-adenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S_N2 -displacement of a 2'-beta-trityl group. Thus

N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-dihydro-2H-pyridin-2-ylidene (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

2'-Fluorodeoxyguanosine

[0129] The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylsilyloxane (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

2'-Fluorouridine

[0130] Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

2'-Fluorodeoxycytidine

[0131] 2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

2'-O-(2-Methoxyethyl) modified amidites

[0132] 2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

[0133] 5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60° C. at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or

it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4° C.)

2'-O-Methoxyethyl-5-methyluridine

[0134] 2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160° C. After heating for 48 hours at 155-160° C., the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH₂Cl₂/acetone/MeOH (20:5:3) containing 0.5% Et₃NH. The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

[0135] 2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH₃CN (200 mL). The residue was dissolved in CHCl₃ (1.5 L) and extracted with 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl. The organic phase was dried over Na₂SO₄, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et₃NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

[0136] 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL) of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35° C. The residue was dissolved in CHCl₃ (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl₃. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

[0137] A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH₃CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH₃CN (1 L), cooled to -5° C. and stirred for 0.5 h using an overhead stirrer. POCl₃ was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10° C., and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1×300 mL of NaHCO₃ and 2×300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

[0138] A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH₄OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2×200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH₃ gas was added and the vessel heated to 100° C. for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

[0139] 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl₃ (700 mL) and extracted with saturated NaHCO₃ (2×300 mL) and saturated NaCl (2×300 mL), dried over MgSO₄ and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et₃NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

N4-Benzoyl-2'-N-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

[0140] N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH₂Cl₂ (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The

reaction mixture was extracted with saturated NaHCO₃ (1×300 mL) and saturated NaCl (3×300 mL). The aqueous washes were back-extracted with CH₂Cl₂ (300 mL), and the extracts were combined, dried over MgSO₄ and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

2'-O-(Aminoxyethyl) nucleoside amidites and
2'-O-(dimethylaminoxyethyl) nucleoside amidites

2'-(Dimethylaminoxyethoxy) nucleoside amidites

[0141] 2'-(Dimethylaminoxyethoxy) nucleoside amidites (also known in the art as 2'-O-(dimethylaminoxyethyl) nucleoside amidites) are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutryl in the case of guanosine.

5-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine

[0142] O²-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0 g, 0.416 mmol), dimethylaminopyridine (0.66 g, 0.013 eq, 0.0054 mmol) were dissolved in dry pyridine (500 mL) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8 g, 119.0 mL, 1.1 eq, 0.458 mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (Rf 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2×1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600 mL) and the solution was cooled to -10° C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3×200 mL) and dried (40° C., 1 mm Hg, 24 h) to 149 g (74.8%) of white solid. TLC and NMR were consistent with pure product.

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine

[0143] In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160° C. was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1 mm Hg) in a warm water bath (40-100° C.) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low

boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2 kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84 g, 50%), contaminated starting material (17.4 g) and pure reusable starting material 20 g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine

[0144] 5'-O-tert-Butylidiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20 g, 36.98 mmol) was mixed with triphenylphosphine (11.63 g, 44.36 mmol) and N-hydroxyphthalimide (7.24 g, 44.36 mmol). It was then dried over P₂O₅ under high vacuum for two days at 40° C. The reaction mixture was flushed with argon and dry THF (369.8 mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98 mL, 44.36 mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine

[0145] 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine (3.1 g, 4.5 mmol) was dissolved in dry CH₂Cl₂ (4.5 mL) and methylhydrazine (300 mL 4.64 mmol) was added dropwise at -10° C. to 0° C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH₂Cl₂ and the combined organic phase was washed with water, brine and dried over anhydrous Na₂SO₄. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5 mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine as white foam: (1.95 g, 78%).

5'-O-tert-Butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine

[0146] 5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77 g, 3.12 mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6 mL). Sodium cyanoborohydride (0.39 g, 6.13 mmol) was added to this solution at 10° C. under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10° C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH₂Cl₂). Aqueous NaHCO₃ solution (5%, 10 mL) was added and extracted with ethyl acetate (2x20 mL). Ethyl

acetate phase was dried over anhydrous Na₂SO₄, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6 mL). Formaldehyde (20% w/w, 30 mL, 3.37 mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10° C. in an ice bath, sodium cyanoborohydride (0.39 g, 6.13 mmol) was added and reaction mixture stirred at 10° C. for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO₃ (25 mL) solution was added and extracted with ethyl acetate (2x25 mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH₂Cl₂ to get 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6 g, 80%).

2'-O-(dimethylaminoxyethyl)-5-methyluridine

[0147] Triethylamine trihydrofluoride (3.91 mL, 24.0 mmol) was dissolved in dry THF and triethylamine (1.67 mL, 12 mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40 g, 2.4 mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH₂Cl₂). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH₂Cl₂ to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766 mg, 92.5%).

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine

[0148] 2'-O-(dimethylaminoxyethyl)-5-methyluridine (750 mg, 2.17 mmol) was dried over P₂O₅ under high vacuum overnight at 40° C. It was then co-evaporated with anhydrous pyridine (20 mL). The residue obtained was dissolved in pyridine (11 mL) under argon atmosphere. 4-dimethylaminopyridine (26.5 mg, 2.60 mmol), 4,4'-dimethoxytrityl chloride (880 mg, 2.60 mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH₂Cl₂ (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13 g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

[0149] 5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08 g, 1.67 mmol) was co-evaporated with toluene (20 mL). To the residue N,N-diisopropylamine tetrazonide (0.29 g, 1.67 mmol) was added and dried over P₂O₅ under high vacuum overnight at 40° C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4 mL) and 2-cyanoethyl-N,N,N¹,N¹-tetraisopropylphosphoramidite (2.12 mL, 6.08 mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70 mL) and washed with 5% aqueous NaHCO₃ (40 mL). Ethyl acetate

layer was dried over anhydrous Na_2SO_4 and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04 g, 74.9%).

2'-(Aminoxyethoxy) nucleoside amidites

[0150] 2'-(Aminoxyethoxy) nucleoside amidites (also known in the art as 2'-O-(aminoxyethyl) nucleoside amidites) are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

N2-isobutyryl-6-O-diphenylcarbonyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

[0151] The 2'-O-aminoxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbonyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbonyl-2'-O-(2-hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphorylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbonyl-2'-O-([2-phthalimidoxy]ethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites

[0152] 2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O- $\text{CH}_2\text{—O—CH}_2\text{—N}(\text{CH}_2)_2$, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

[0153] 2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. $\text{O}^2\text{-},2\text{'-anhydro-5-methyluridine}$ (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155° C. for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3×200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The resi-

due is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

[0154] To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH_2Cl_2 (2×200 mL). The combined CH_2Cl_2 layers are washed with saturated NaHCO_3 solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using $\text{MeOH}:\text{CH}_2\text{Cl}_2:\text{Et}_3\text{N}$ (20:1, v/v, with 1% triethylamine) gives the title compound.

5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite

[0155] Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH_2Cl_2 (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

Example 2

Oligonucleotide synthesis

[0156] Unsubstituted and substituted phosphodiester ($\text{P}=\text{O}$) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

[0157] Phosphorothioates ($\text{P}=\text{S}$) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55° C. (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Pat. No. 5,508,270, herein incorporated by reference.

[0158] Alkyl phosphonate oligonucleotides are prepared as described in U.S. Pat. No. 4,469,863, herein incorporated by reference.

[0159] 3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Pat. Nos. 5,610,289 or 5,625,050, herein incorporated by reference.

[0160] Phosphoramidite oligonucleotides are prepared as described in U.S. Pat. No. 5,256,775 or U.S. Pat. No. 5,366,878, herein incorporated by reference.

[0161] Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

[0162] 3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Pat. No. 5,476,925, herein incorporated by reference.

[0163] Phosphotriester oligonucleotides are prepared as described in U.S. Pat. No. 5,023,243, herein incorporated by reference.

[0164] Borano phosphate oligonucleotides are prepared as described in U.S. Pat. Nos. 5,130,302 and 5,177,198, both herein incorporated by reference.

Example 3

Oligonucleoside synthesis

[0165] Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedi-methylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligo-nucleosides, also identified as amide-4 linked oligonucleo-sides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Pat. Nos. 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

[0166] Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Pat. Nos. 5,264,562 and 5,264,564, herein incorporated by reference.

[0167] Ethylene oxide linked oligonucleosides are prepared as described in U.S. Pat. No. 5,223,618, herein incorporated by reference.

Example 4

PNA Synthesis

[0168] Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Pat. Nos. 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

Example 5

Synthesis of Chimeric Oligonucleotides

[0169] Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3'"wing" segments of linked nucleosides and a second "open end" type wherein the "gap"

segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

[2'-O-Me]—[2'-deoxy—[2'-O-Me]Chimeric Phosphorothioate Oligonucleotides

[0170] Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligo-nucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3-O-phosphor-amidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to ½ volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[2'-O-(2-Methoxyethyl)]—[2'-deoxy]—[2'-O-(Methoxyethyl)]Chimeric Phosphorothioate. Oligonucleotides

[0171] [2'-O-(2-methoxyethyl)]—[2'-deoxy]—[2'-O-(methoxy-ethyl)]chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

[2'-O-(2-Methoxyethyl)Phosphodiester]—[2'-deoxy Phosphorothioate]—[2'-O-(2-Methoxyethyl). Phosphodiester]Chimeric Oligonucleotides

[0172] [2'-O-(2-methoxyethyl phosphodiester)]—[2'-deoxy phosphorothioate]—[2'-O-(methoxyethyl)phosphodiester]chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

[0173] Other chimeric oligonucleotides, chimeric oligo-nucleo-sides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to U.S. Pat. No. 5,623,065, herein incorporated by reference.

Example 6

Oligonucleotide Isolation

[0174] After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55° C. for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by ³¹P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* 1991, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

Example 7

Oligonucleotide Synthesis—96 Well Plate Format

[0175] Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, Calif., or Pharmacia, Piscataway, N.J.). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

[0176] Oligonucleotides were cleaved from support and deprotected with concentrated NH₄OH at elevated temperature (55-60° C.) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

Example 8

Oligonucleotide Analysis—96 Well Plate Format

[0177] The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

Example 9

Cell Culture and Oligonucleotide Treatment

[0178] The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of

cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 7 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

[0179] T-24 Cells:

[0180] The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, Va.). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, Md.) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, Md.), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, Md.). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

[0181] For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

[0182] A549 cells:

[0183] The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, Va.). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, Md.) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, Md.), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, Md.). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

[0184] NHDF Cells:

[0185] Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville Md.). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville Md.) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

[0186] HEK Cells:

[0187] Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville, Md.). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville Md.) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

[0188] HepG2 Cells:

[0189] The human hepatoblastoma cell line HepG2 was obtained from the American Type Culture Collection (Manassas, Va.). HepG2 cells were routinely cultured in Eagle's MEM supplemented with 10% fetal calf serum,

non-essential amino acids, and 1 mM sodium pyruvate (Gibco/Life Technologies, Gaithersburg, Md.). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

[0190] For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

[0191] AML12 Cells:

[0192] AML12 (alpha mouse liver 12) cell line was established from hepatocytes from a mouse (CD1 strain, line MT42) transgenic for human TGF alpha. Cells are cultured in a 1:1 mixture of Dulbecco's modified Eagle's medium and Ham's F12 medium with 0.005 mg/ml insulin, 0.005 mg/ml transferrin, 5 ng/ml selenium, and 40 ng/ml dexamethasone, and 90% 10% fetal bovine serum. For subculturing, spent medium is removed and fresh media of 0.25% trypsin, 0.03% EDTA solution is added. Fresh trypsin solution (1 to 2 ml) is added and the culture is left to sit at room temperature until the cells detach.

[0193] Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

[0194] For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

Primary Mouse Hepatocytes:

[0195] Primary mouse hepatocytes were prepared from CD-1 mice purchased from Charles River Labs (Wilmington, Mass.) and were routinely cultured in Hepatocyte Attachment Media (Gibco) supplemented with 10Fetal Bovine Serum (Gibco/Life Technologies, Gaithersburg, Md.), 250 nM dexamethasone (Sigma), and 10 nM bovine insulin (Sigma). Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 10000 cells/well for use in RT-PCR analysis.

[0196] For Northern blotting or other analyses, cells are plated onto 100 mm or other standard tissue culture plates coated with rat tail collagen (200 ug/mL) (Becton Dickinson) and treated similarly using appropriate volumes of medium and oligonucleotide.

[0197] Treatment with Antisense Compounds:

[0198] When cells reach 80% confluency, they are treated with oligonucleotide. For cells grown in 96-well plates, wells are washed once with 200 μ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130 μ L of OPTI-MEM™-1 containing 3.75 μ g/mL LIPO-FECTIN™ (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium is replaced with fresh medium. Cells are harvested 16-24 hours after oligonucleotide treatment.

[0199] The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells

are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, TCCGTCATCGCTCCT-CAGGG, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, ATGCATTCTGCCCCAAGGA, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

Example 10

Analysis of oligonucleotide Inhibition of apolipoprotein(a) Expression

[0200] Antisense modulation of apolipoprotein(a) expression can be assayed in a variety of ways known in the art. For example, apolipoprotein(a) mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, Calif. and used according to manufacturer's instructions.

[0201] Protein levels of apolipoprotein(a) can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to apolipoprotein(a) can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, Mich.), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal-antisera are taught in, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

[0202] Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F. M. et al.,

Current Protocols in Molecular Biology, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

Example 11

Poly(A)+ mRNA isolation

[0203] Poly(A)+ mRNA can be isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium is removed from the cells and each well is washed with 200 μ L cold PBS. 60 μ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) is added to each well, the plate is gently agitated and then incubated at room temperature for five minutes. 55 μ L of lysate is transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine Calif.). Plates are incubated for 60 minutes at room temperature, washed 3 times with 200 μ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate is blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60 μ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70° C. is added to each well, the plate is incubated on a 90° C. hot plate for 5 minutes, and the eluate is then transferred to a fresh 96-well plate.

[0204] Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

Example 12

Total RNA Isolation

[0205] Total RNA can be isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia Calif.) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium is removed from the cells and each well is washed with 200 μ L cold PBS. 100 μ L Buffer RLT is added to each well and the plate vigorously agitated for 20 seconds. 100 μ L of 70% ethanol is then added to each well and the contents mixed by pipetting three times up and down. The samples are then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum is applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE is then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum is applied for an additional 10 minutes. The plate is then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate is then re-attached to the QIAVAC™ manifold fitted with a collec-

tion tube rack containing 1.2 mL collection tubes. RNA is then eluted by pipetting 60 μ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step is repeated with an additional 60 μ L water.

[0206] The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia Calif.). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

Example 13

Real-Time Quantitative PCR Analysis of apolipoprotein(a) mRNA Levels

[0207] Quantitation of apolipoprotein(a) mRNA levels can be determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, Calif.) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, Calif. or PE-Applied Biosystems, Foster City, Calif.) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, Calif. or PE-Applied Biosystems, Foster City, Calif.) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISMS™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

[0208] Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both

(multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

[0209] PCR reagents are obtained from PE-Applied Biosystems, Foster City, Calif. RT-PCR reactions are carried out by adding 25 μ L PCR cocktail (1 \times TAQMANTM buffer A, 5.5 mM MgCl₂, 300 μ M each of dATP, dCTP and dGTP, 600 μ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLI-TAQ GOLDTM, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 μ L total RNA solution. The RT reaction is carried out by incubation for 30 minutes at 48° C. Following a 10 minute incubation at 95° C. to activate the AMPLI-TAQ GOLDTM, 40 cycles of a two-step PCR protocol are carried out: 95° C. for 15 seconds (denaturation) followed by 60° C. for 1.5 minutes (annealing/extension).

[0210] Gene target quantities obtained by real time RT-PCR can be normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreenTM (Molecular Probes, Inc. Eugene, Oreg.). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreenTM RNA quantification reagent from Molecular Probes (Eugene, Oreg.). Methods of RNA quantification by RiboGreenTM are taught in Jones, L. J., et al, *Analytical Biochemistry*, 1998, 265, 368-374.

[0211] In this assay, 175 μ L of RiboGreenTM working reagent (RiboGreenTM reagent diluted 1:2865 in 10 mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25 μ L purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480 nm and emission at 520 nm.

[0212] Probes and primers to human apolipoprotein(a) are designed to hybridize to a human apolipoprotein(a) sequence, using published sequence information (GenBank accession number NM_005577, incorporated herein as SEQ ID NO: 3).

[0213] For human GAPDH the standard PCR primers are: forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 4) reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 5) and the PCR probe is: 5' JOE-CAAGCT-TCCCGTTCTCAGCC-TAMRA 3' (SEQ ID NO: 6) where JOE (PE-Applied Biosystems, Foster City, Calif.) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, Calif.) is the quencher dye.

Example 14

Northern Blot Analysis of apolipoprotein(a) mRNA Levels

[0214] Eighteen hours after antisense treatment, cell monolayers are washed twice with cold PBS and lysed in 1 mL RNAzolTM (TEL-TEST "B" Inc., Friendswood, Tex.). Total RNA is prepared following manufacturer's recom-

mended protocols. Twenty micrograms of total RNA is fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, Ohio). RNA is transferred from the gel to HYBONDTM-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, N.J.) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, Tex.). RNA transfer is confirmed by UV visualization. Membranes are fixed by UV cross-linking using a STRATALINKERTM UV Crosslinker 2400 (Stratagene, Inc, La Jolla, Calif.) and then probed using QUICKHYBTM hybridization solution (Stratagene, La Jolla, Calif.) using manufacturer's recommendations for stringent conditions.

[0215] To detect human apolipoprotein(a), a human apolipoprotein(a) specific probe is prepared by PCR using a forward and a reverse primer. To normalize for variations in loading and transfer efficiency, membranes are stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, Calif.).

[0216] Hybridized membranes are visualized and quantitated using a PHOSPHORIMAGERTM and IMAGEQUANTTM Software V3.3 (Molecular Dynamics, Sunnyvale, Calif.). Data are normalized to GAPDH levels in untreated controls.

Example 15

Chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap targeting human apolipoprotein(a)

[0217] In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human apolipoprotein(a) RNA, using published sequence (GenBank accession number NM_005577, incorporated herein as SEQ ID NO: 3). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3'directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines.

TABLE 1

Chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap targeting human apolipoprotein(a)					
ISIS #	REGION	TARGET		SEQUENCE	SEQ ID NO
		SEQ ID NO	TARGET SITE		
144367	Coding	3	174	GGCAGGTCCCTCCTGTGACA	7
144368	Coding	3	352	TCTGCGTCTGAGCATTGCGT	8
144369	Coding	3	522	AAGCTTGGCAGGTTCTTCCT	9
144370	Coding	3	1743	TCGGAGCGCGACGGCAGTC	10

TABLE 1-continued

Chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap targeting human apolipoprotein(a)					
ISIS #	REGION	TARGET		SEQUENCE	SEQ ID NO
		SEQ ID	TARGET SITE		
144371	Coding	3	2768	CGGAGGCGCGACGGCAGTCC	11
144372	Coding	3	2910	GGCAGGTCTTCTCTGTGACA	12
144373	Coding	3	3371	ATAACAATAAGGAGCTGCCA	13
144374	Coding	3	4972	GACCAAGCTTGGCAGGTTCT	14
144375	Coding	3	5080	TAACAATAAGGAGCTGCCAC	15
144376	Coding	3	5315	TGACCAAGCTTGGCAGGTTTC	16
144377	Coding	3	5825	TTCTGCGTCTGAGCATTGCG	17
144378	Coding	3	6447	AACAATAAGGAGCTGCCACA	18
144379	Coding	3	7155	ACCTGACACCGGGATCCCTC	19
144380	Coding	3	7185	CTGAGCATTGCGTCAGGTTG	20
144381	Coding	3	8463	AGTAGTTCATGATCAAGCCA	21
144382	Coding	3	8915	GACGGCAGTCCCTTCTGCGT	22
144383	Coding	3	9066	GGCAGGTCTTCCAGTGACA	23
144384	Coding	3	10787	TGACCAAGCTTGGCAAGTTC	24
144385	Coding	3	11238	TATAACACCAAGGACTAATC	25
144386	Coding	3	11261	CCATCTGACATTGGGATCCA	26
144387	Coding	3	11461	TGTGGTGTGATAGAGGACCA	27
144388	Coding	3	11823	ATGGGATCCTCCGATGCCAA	28
144389	Coding	3	11894	ACACCAAGGGCGAATCTCAG	29
144390	Coding	3	11957	TTCTGTCACTGGACATCGTG	30
144391	Coding	3	12255	CACACGGATCGGTTGTGTAA	31
144392	Coding	3	12461	ACATGTCCTTCTGTGACAG	32

TABLE 1-continued

Chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap targeting human apolipoprotein(a)					
ISIS #	REGION	TARGET		SEQUENCE	SEQ ID NO
		SEQ ID	TARGET SITE		
144393	Coding	3	12699	CAGAAGGAGGCCCTAGGCTT	33
144394	Coding	3	13354	CTGGCGGTGACCATGTAGTC	34
144395	3'UTR	3	13711	TCTAAGTAGGTTGATGCTTC	35
144396	3'UTR	3	13731	TCCTTACCCACGTTTCAGCT	36
144397	3'UTR	3	13780	GGAACAGTGTCTTCGTTTGA	37
144398	3'UTR	3	13801	GTTTGGCATAGCTGGTAGCT	38
144399	3'UTR	3	13841	ACCTTAAAAGCTTATACACA	39
144400	3'UTR	3	13861	ATACAGAATTTGTCAGTCAG	40
144401	3'UTR	3	13881	GTCATAGCTATGACACCTTA	41

Example 16

Western Blot Analysis of apolipoprotein(a) Protein Levels

[0218] Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to apolipoprotein(a) is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized, using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale Calif.).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 41

<210> SEQ ID NO 1

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 1

tccgtcatcg ctcctcaggg

-continued

```

<210> SEQ ID NO 2
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 2

atgcattctg cccccaagga                                20

<210> SEQ ID NO 3
<211> LENGTH: 13938
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (46)...(13692)

<400> SEQUENCE: 3

ctgggattgg gacacacttt ctggacactg ctggccagtc ccaaa atg gaa cat aag    57
                                     Met Glu His Lys
                                     1

gaa gtg gtt ctt cta ctt ctt tta ttt ctg aaa tca gca gca cct gag    105
Glu Val Val Leu Leu Leu Leu Phe Leu Lys Ser Ala Ala Pro Glu
  5                10                15                20

caa agc cat gtg gtc cag gat tgc tac cat ggt gat gga cag agt tat    153
Gln Ser His Val Val Gln Asp Cys Tyr His Gly Asp Gly Gln Ser Tyr
                25                30                35

cga ggc acg tac tcc acc act gtc aca gga agg acc tgc caa gct tgg    201
Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp
                40                45                50

tca tct atg aca cca cat caa cat aat agg acc aca gaa aac tac cca    249
Ser Ser Met Thr Pro His Gln His Asn Arg Thr Thr Glu Asn Tyr Pro
                55                60                65

aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca    297
Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala
  70                75                80

gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc    345
Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys
  85                90                95                100

aac ctg acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg    393
Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro
                105                110                115

act gtt acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg    441
Thr Val Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro
                120                125                130

act gag caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag    489
Thr Glu Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln
                135                140                145

agt tat cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa    537
Ser Tyr Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln
                150                155                160

gct tgg tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac    585
Ala Trp Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr
                165                170                175                180

tac cca aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct    633
Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala
                185                190                195

gtg gca gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag    681
Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu

```


-continued

805	810	815	820	
cct ggg gtg cag gag tgc tac cat ggt aat gga cag agt tat cga ggc				2553
Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly	825	830	835	
aca tac tcc acc act gtc aca gga aga acc tgc caa gct tgg tca tct				2601
Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser	840	845	850	
atg aca cca cac tcg cat agt cgg acc cca gaa tac tac cca aat gct				2649
Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala	855	860	865	
ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca gct cct				2697
Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Ala Pro	870	875	880	
tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg				2745
Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu	885	890	895	900
acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg act gtt				2793
Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val	905	910	915	
acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg act gag				2841
Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu	920	925	930	
caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag agt tat				2889
Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr	935	940	945	
cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa gct tgg				2937
Arg Gly Thr Tyr Ser Thr Val Thr Gly Arg Thr Cys Gln Ala Trp	950	955	960	
tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac tac cca				2985
Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro	965	970	975	980
aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca				3033
Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala	985	990	995	
gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc				3081
Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys	1000	1005	1010	
aac ctg acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg				3129
Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro	1015	1020	1025	
act gtt acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg				3177
Thr Val Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro	1030	1035	1040	
act gag caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag				3225
Thr Glu Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln	1045	1050	1055	1060
agt tat cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa				3273
Ser Tyr Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln	1065	1070	1075	
gct tgg tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac				3321
Ala Trp Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr	1080	1085	1090	
tac cca aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct				3369
Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala	1095	1100	1105	
gtg gca gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag				3417
Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu				

-continued

1110	1115	1120	
tac tgc aac ctg acg caa tgc tca gac gca gaa ggg act gcc gtc gcg Tyr Cys Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala 1125	1130	1135	1140 3465
cct ccg act gtt acc ccg gtt cca agc cta gag gct cct tcc gaa caa Pro Pro Thr Val Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln 1145	1150	1155	3513
gca ccg act gag caa agg cct ggg gtg cag gag tgc tac cat ggt aat Ala Pro Thr Glu Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn 1160	1165	1170	3561
gga cag agt tat cga ggc aca tac tcc acc act gtc aca gga aga acc Gly Gln Ser Tyr Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr 1175	1180	1185	3609
tgc caa gct tgg tca tct atg aca cca cac tcg cat agt cgg acc cca Cys Gln Ala Trp Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro 1190	1195	1200	3657
gaa tac tac cca aat gct ggc ttg atc atg aac tac tgc agg aat cca Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro 1205	1210	1215	3705
gat gct gtg gca gct cct tat tgt tat acg agg gat ccc ggt gtc agg Asp Ala Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg 1225	1230	1235	3753
tgg gag tac tgc aac ctg acg caa tgc tca gac gca gaa ggg act gcc Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala 1240	1245	1250	3801
gtc gcg cct ccg act gtt acc ccg gtt cca agc cta gag gct cct tcc Val Ala Pro Pro Thr Val Thr Pro Val Pro Ser Leu Glu Ala Pro Ser 1255	1260	1265	3849
gaa caa gca ccg act gag caa agg cct ggg gtg cag gag tgc tac cat Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly Val Gln Glu Cys Tyr His 1270	1275	1280	3897
ggt aat gga cag agt tat cga ggc aca tac tcc acc act gtc aca gga Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly 1285	1290	1295	3945
aga acc tgc caa gct tgg tca tct atg aca cca cac tcg cat agt cgg Arg Thr Cys Gln Ala Trp Ser Ser Met Thr Pro His Ser His Ser Arg 1305	1310	1315	3993
acc cca gaa tac tac cca aat gct ggc ttg atc atg aac tac tgc agg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg 1320	1325	1330	4041
aat cca gat gct gtg gca gct cct tat tgt tat acg agg gat ccc ggt Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly 1335	1340	1345	4089
gtc agg tgg gag tac tgc aac ctg acg caa tgc tca gac gca gaa ggg Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly 1350	1355	1360	4137
act gcc gtc gcg cct ccg act gtt acc ccg gtt cca agc cta gag gct Thr Ala Val Ala Pro Pro Thr Val Thr Pro Val Pro Ser Leu Glu Ala 1365	1370	1375	4185
cct tcc gaa caa gca ccg act gag caa agg cct ggg gtg cag gag tgc Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly Val Gln Glu Cys 1385	1390	1395	4233
tac cat ggt aat gga cag agt tat cga ggc aca tac tcc acc act gtc Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr Ser Thr Val 1400	1405	1410	4281
aca gga aga acc tgc caa gct tgg tca tct atg aca cca cac tcg cat Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr Pro His Ser His 1415			4329

-continued

1415	1420	1425	
agt cgg acc cca gaa tac tac cca aat gct ggc ttg atc atg aac tac Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr 1430 1435 1440			4377
tgc agg aat cca gat gct gtg gca gct cct tat tgt tat acg agg gat Cys Arg Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp 1445 1450 1455 1460			4425
ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa tgc tca gac gca Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser Asp Ala 1465 1470 1475			4473
gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg gtt cca agc cta Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro Val Pro Ser Leu 1480 1485 1490			4521
gag gct cct tcc gaa caa gca ccg act gag caa agg cct ggg gtg cag Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly Val Gln 1495 1500 1505			4569
gag tgc tac cat ggt aat gga cag agt tat cga ggc aca tac tcc acc Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr Ser Thr 1510 1515 1520			4617
act gtc aca gga aga acc tgc caa gct tgg tca tct atg aca cca cac Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr Pro His 1525 1530 1535 1540			4665
tcg cat agt cgg acc cca gaa tac tac cca aat gct ggc ttg atc atg Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met 1545 1550 1555			4713
aac tac tgc agg aat cca gat gct gtg gca gct cct tat tgt tat acg Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Pro Tyr Cys Tyr Thr 1560 1565 1570			4761
agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa tgc tca Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser 1575 1580 1585			4809
gac gca gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg gtt cca Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro Val Pro 1590 1595 1600			4857
agc cta gag gct cct tcc gaa caa gca ccg act gag caa agg cct ggg Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly 1605 1610 1615 1620			4905
gtg cag gag tgc tac cat ggt aat gga cag agt tat cga ggc aca tac Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr 1625 1630 1635			4953
tcc acc act gtc aca gga aga acc tgc caa gct tgg tca tct atg aca Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr 1640 1645 1650			5001
cca cac tcg cat agt cgg acc cca gaa tac tac cca aat gct ggc ttg Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu 1655 1660 1665			5049
atc atg aac tac tgc agg aat cca gat gct gtg gca gct cct tat tgt Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys 1670 1675 1680			5097
tat acg agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln 1685 1690 1695 1700			5145
tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro 1705 1710 1715			5193
gtt cca agc cta gag gct cct tcc gaa caa gca ccg act gag caa agg Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg 1720 1725 1730 1735			5241

-continued

	1720	1725	1730	
cct ggg gtg cag gag tgc tac cat ggt aat gga cag agt tat cga ggc				5289
Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly	1735	1740	1745	
aca tac tcc acc act gtc aca gga aga acc tgc caa gct tgg tca tct				5337
Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser	1750	1755	1760	
atg aca cca cac tcg cat agt cgg acc cca gaa tac tac cca aat gct				5385
Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala	1765	1770	1775	1780
ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca gct cct				5433
Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Ala Pro	1785	1790	1795	
tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg				5481
Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu	1800	1805	1810	
acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg act gtt				5529
Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val	1815	1820	1825	
acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg act gag				5577
Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu	1830	1835	1840	
caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag agt tat				5625
Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr	1845	1850	1855	1860
cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa gct tgg				5673
Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp	1865	1870	1875	
tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac tac cca				5721
Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro	1880	1885	1890	
aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca				5769
Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala	1895	1900	1905	
gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc				5817
Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys	1910	1915	1920	
aac ctg acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg				5865
Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro	1925	1930	1935	1940
act gtt acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg				5913
Thr Val Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro	1945	1950	1955	
act gag caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag				5961
Thr Glu Gln Arg Pro Gly Val Gln Cys Tyr His Gly Asn Gly Gln	1960	1965	1970	
agt tat cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa				6009
Ser Tyr Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln	1975	1980	1985	
gct tgg tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac				6057
Ala Trp Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr	1990	1995	2000	
tac cca aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct				6105
Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala	2005	2010	2015	2020
gtg gca gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag				6153
Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu				

-continued

2325	2330	2335	2340
agt cgg acc cca gaa tac tac cca aat gct ggc ttg atc atg aac tac Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr 2345 2350 2355			7113
tgc agg aat cca gat gct gtg gca gct cct tat tgt tat acg agg gat Cys Arg Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp 2360 2365 2370			7161
ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa tgc tca gac gca Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser Asp Ala 2375 2380 2385			7209
gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg gtt cca agc cta Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro Val Pro Ser Leu 2390 2395 2400			7257
gag gct cct tcc gaa caa gca ccg act gag caa agg cct ggg gtg cag Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly Val Gln 2405 2410 2415 2420			7305
gag tgc tac cat ggt aat gga cag agt tat cga ggc aca tac tcc acc Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr Ser Thr 2425 2430 2435			7353
act gtc aca gga aga acc tgc caa gct tgg tca tct atg aca cca cac Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr Pro His 2440 2445 2450			7401
tcg cat agt cgg acc cca gaa tac tac cca aat gct ggc ttg atc atg Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met 2455 2460 2465			7449
aac tac tgc agg aat cca gat gct gtg gca gct cct tat tgt tat acg Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys Tyr Thr 2470 2475 2480			7497
agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa tgc tca Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser 2485 2490 2495 2500			7545
gac gca gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg gtt cca Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro Val Pro 2505 2510 2515			7593
agc cta gag gct cct tcc gaa caa gca ccg act gag caa agg cct ggg Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly 2520 2525 2530			7641
gtg cag gag tgc tac cat ggt aat gga cag agt tat cga ggc aca tac Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr 2535 2540 2545			7689
tcc acc act gtc aca gga aga acc tgc caa gct tgg tca tct atg aca Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr 2550 2555 2560			7737
cca cac tcg cat agt cgg acc cca gaa tac tac cca aat gct ggc ttg Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu 2565 2570 2575 2580			7785
atc atg aac tac tgc agg aat cca gat gct gtg gca gct cct tat tgt Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys 2585 2590 2595			7833
tat acg agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln 2600 2605 2610			7881
tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro 2615 2620 2625			7929
gtt cca agc cta gag gct cct tcc gaa caa gca ccg act gag cag agg Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg			7977

-continued

2630	2635	2640	
cct ggg gtg cag gag tgc tac cac ggt aat gga cag agt tat cga ggc			8025
Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly			
2645	2650	2655	2660
aca tac tcc acc act gtc act gga aga acc tgc caa gct tgg tca tct			8073
Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser			
	2665	2670	2675
atg aca cca cac tcg cat agt cgg acc cca gaa tac tac cca aat gct			8121
Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala			
	2680	2685	2690
ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca gct cct			8169
Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala Ala Pro			
	2695	2700	2705
tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc aac ctg			8217
Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu			
	2710	2715	2720
acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg act gtt			8265
Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Val			
	2725	2730	2740
acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg act gag			8313
Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu			
	2745	2750	2755
caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag agt tat			8361
Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr			
	2760	2765	2770
cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa gct tgg			8409
Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ala Trp			
	2775	2780	2785
tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac tac cca			8457
Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro			
	2790	2795	2800
aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct gtg gca			8505
Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala Val Ala			
	2805	2810	2815
gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag tac tgc			8553
Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu Tyr Cys			
	2825	2830	2835
aac ctg acg caa tgc tca gac gca gaa ggg act gcc gtc gcg cct ccg			8601
Asn Leu Thr Gln Cys Ser Asp Ala Glu Gly Thr Ala Val Ala Pro Pro			
	2840	2845	2850
act gtt acc ccg gtt cca agc cta gag gct cct tcc gaa caa gca ccg			8649
Thr Val Thr Pro Val Pro Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro			
	2855	2860	2865
act gag caa agg cct ggg gtg cag gag tgc tac cat ggt aat gga cag			8697
Thr Glu Gln Arg Pro Gly Val Gln Glu Cys Tyr His Gly Asn Gly Gln			
	2870	2875	2880
agt tat cga ggc aca tac tcc acc act gtc aca gga aga acc tgc caa			8745
Ser Tyr Arg Gly Thr Tyr Ser Thr Thr Val Thr Gly Arg Thr Cys Gln			
	2885	2890	2895
gct tgg tca tct atg aca cca cac tcg cat agt cgg acc cca gaa tac			8793
Ala Trp Ser Ser Met Thr Pro His Ser His Ser Arg Thr Pro Glu Tyr			
	2905	2910	2915
tac cca aat gct ggc ttg atc atg aac tac tgc agg aat cca gat gct			8841
Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr Cys Arg Asn Pro Asp Ala			
	2920	2925	2930
gtg gca gct cct tat tgt tat acg agg gat ccc ggt gtc agg tgg gag			8889
Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp Pro Gly Val Arg Trp Glu			

-continued

2935	2940	2945	
tac tgc aac ctg acg Tyr Cys Asn 2950	caa tgc tca gac gca Gln Cys Ser Asp 2955	gaa ggg act gcc gtc gcg Glu Gly Thr Ala Val Ala 2960	8937
cct ccg act gtt acc Pro Pro Thr 2965	ccg gtt cca agc cta Pro Val Pro Ser 2970	gag gct cct tcc gaa caa Glu Ala Pro Ser Glu Gln 2975 2980	8985
gca ccg act gag cag Ala Pro Thr 2985	agg cct ggg gtg Arg Pro Gly Val 2985	cag gag tgc tac cac ggt aat Gln Glu Cys Tyr His Gly Asn 2990 2995	9033
gga cag agt tat cga Gly Gln Ser 3000	ggc aca tac tcc acc Gly Thr Tyr Ser Thr 3005	act gtc act gga aga acc Thr Val Thr Gly Arg Thr 3010	9081
tgc caa gct tgg tca Cys Gln Ala 3015	tct atg aca cca cac Ser Ser Met Thr Pro 3020	tcg cat agt cgg acc cca Ser His Ser Arg Thr Pro 3025	9129
gaa tac tac cca aat Glu Tyr Tyr 3030	gct ggc ttg atc atg Ala Gly Leu Ile Met 3035	aac tac tgc agg aat cca Tyr Cys Arg Asn Pro 3040	9177
gat gct gtg gca gct Asp Ala Val 3045	cct tat tgt tat acg Pro Tyr Cys Tyr Thr 3050	agg gat ccc ggt gtc agg Arg Asp Pro Gly Val Arg 3055 3060	9225
tgg gag tac tgc aac Trp Glu Tyr 3065	ctg acg caa tgc tca Leu Thr Gln Cys Ser 3070	gac gca gaa ggg act gcc Asp Ala Glu Gly Thr Ala 3075	9273
gtc gcg cct ccg act Val Ala Pro 3080	gtt acc ccg gtt cca Pro Thr Val Thr Pro 3085	agc cta gag gct cct tcc Ser Leu Glu Ala Pro Ser 3090	9321
gaa caa gca ccg act Glu Gln Ala 3095	gag cag agg cct ggg Glu Gln Arg Pro Gly 3100	gtg cag gag tgc tac cac Val Gln Glu Cys Tyr His 3105	9369
ggt aat gga cag agt Gly Asn Gly 3110	gat gct cct tat tgt Gln Ser Tyr Arg Gly 3115	atc atg aac tac tgc atc atg aac tac tgc 3120	9417
aga acc tgc caa gct Arg Thr Cys 3125	tgg tca tct atg aca Trp Ser Ser Met Thr 3130	cca cac tcg cat agt cgg Pro His Ser His Ser Arg 3135 3140	9465
acc cca gaa tac tac Thr Pro Glu Tyr 3145	cca aat gct ggc ttg Pro Asn Ala Gly Leu 3150	atc atg aac tac tgc agg Ile Met Asn Tyr Cys Arg 3155	9513
aat cca gat gct gtg Asn Pro Asp 3160	gca gct cct tat tgt Ala Val Ala Ala Pro 3165	atc atg aac tac tgc Tyr Cys Tyr Thr Arg 3170	9561
gtc agg tgg gag tac Val Arg Trp Glu 3175	tgc aac ctg acg caa Leu Thr Gln Cys Ser 3180	tca gac gca gaa ggg Ser Asp Ala Glu Gly 3185	9609
act gcc gtc gcg cct Thr Ala Val 3190	ccg act gtt acc ccg Pro Pro Thr Val Thr 3195	ggt cca agc cta gag gct Pro Val Pro Ser Leu Ala 3200	9657
cct tcc gaa caa gca Pro Ser Glu Gln 3205	ccg act gag cag agg Pro Thr Glu Gln Arg 3210	cct ggg gtg cag gag tgc Pro Gly Val Gln Glu Cys 3215 3220	9705
tac cac ggt aat gga Tyr His Gly Asn 3225	cag agt tat cga ggc Gln Ser Tyr Arg Gly 3230	aca tac tcc acc act gtc Thr Tyr Ser Thr Thr Val 3235	9753
act gga aga acc tgc Thr Gly Arg Thr 3240	caa gct tgg tca tct Gln Ala Trp Ser Ser 3245	atg aca cca cac tcg cat Met Thr Pro His Ser His 3250	9801

-continued

3240	3245	3250	
agt cgg acc cca gaa tac tac cca aat gct ggc ttg atc atg aac tac Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met Asn Tyr 3255 3260 3265			9849
tgc agg aat cca gat gct gtg gca gct cct tat tgt tat acg agg gat Cys Arg Asn Pro Asp Ala Val Ala Ala Pro Tyr Cys Tyr Thr Arg Asp 3270 3275 3280			9897
ccc ggt gtc agg tgg gag tac tgc aac ctg acg caa tgc tca gac gca Pro Gly Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser Asp Ala 3285 3290 3295 3300			9945
gaa ggg act gcc gtc gcg cct ccg act gtt acc ccg gtt cca agc cta Glu Gly Thr Ala Val Ala Pro Pro Thr Val Thr Pro Val Pro Ser Leu 3305 3310 3315			9993
gag gct cct tcc gaa caa gca ccg act gag cag agg cct ggg gtg cag Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly Val Gln 3320 3325 3330			10041
gag tgc tac cac ggt aat gga cag agt tat cga ggc aca tac tcc acc Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Tyr Ser Thr 3335 3340 3345			10089
act gtc act gga aga acc tgc caa gct tgg tca tct atg aca cca cac Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr Pro His 3350 3355 3360			10137
tcg cat agt cgg acc cca gaa tac tac cca aat gct ggc ttg atc atg Ser His Ser Arg Thr Pro Glu Tyr Tyr Pro Asn Ala Gly Leu Ile Met 3365 3370 3375 3380			10185
aac tac tgc agg aat cca gat cct gtg gca gcc cct tat tgt tat acg Asn Tyr Cys Arg Asn Pro Asp Pro Val Ala Ala Pro Tyr Cys Tyr Thr 3385 3390 3395			10233
agg gat ccc agt gtc agg tgg gag tac tgc aac ctg aca caa tgc tca Arg Asp Pro Ser Val Arg Trp Glu Tyr Cys Asn Leu Thr Gln Cys Ser 3400 3405 3410			10281
gac gca gaa ggg act gcc gtc gcg cct cca act att acc ccg att cca Asp Ala Glu Gly Thr Ala Val Ala Pro Pro Thr Ile Thr Pro Ile Pro 3415 3420 3425			10329
agc cta gag gct cct tct gaa caa gca cca act gag caa agg cct ggg Ser Leu Glu Ala Pro Ser Glu Gln Ala Pro Thr Glu Gln Arg Pro Gly 3430 3435 3440			10377
gtg cag gag tgc tac cac gga aat gga cag agt tat caa ggc aca tac Val Gln Glu Cys Tyr His Gly Asn Gly Gln Ser Tyr Gln Gly Thr Tyr 3445 3450 3455 3460			10425
ttc att act gtc aca gga aga acc tgc caa gct tgg tca tct atg aca Phe Ile Thr Val Thr Gly Arg Thr Cys Gln Ala Trp Ser Ser Met Thr 3465 3470 3475			10473
cca cac tcg cat agt cgg acc cca gca tac tac cca aat gct ggc ttg Pro His Ser His Ser Arg Thr Pro Ala Tyr Tyr Pro Asn Ala Gly Leu 3480 3485 3490			10521
atc aag aac tac tgc cga aat cca gat cct gtg gca gcc cct tgg tgt Ile Lys Asn Tyr Cys Arg Asn Pro Asp Pro Val Ala Ala Pro Trp Cys 3495 3500 3505			10569
tat aca aca gat ccc agt gtc agg tgg gag tac tgc aac ctg aca cga Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr Cys Asn Leu Thr Arg 3510 3515 3520			10617
tgc tca gat gca gaa tgg act gcc ttc gtc cct ccg aat gtt att ctg Cys Ser Asp Ala Glu Trp Thr Ala Phe Val Pro Pro Asn Val Ile Leu 3525 3530 3535 3540			10665
gct cca agc cta gag gct ttt ttt gaa caa gca ctg act gag gaa acc Ala Pro Ser Leu Glu Ala Phe Phe Glu Gln Ala Leu Thr Glu Glu Thr			10713

-continued

3845	3850	3855	3860	
gtg atg gaa tca act ctc ctc aca act ccc acg gtg gtc cca gtt cca				11673
Val Met Glu Ser Thr Leu Leu Thr Thr Pro Thr Val Val Pro Val Pro				
	3865	3870	3875	
agc aca gag ctt cct tct gaa gaa gca cca act gaa aac agc act ggg				11721
Ser Thr Glu Leu Pro Ser Glu Glu Ala Pro Thr Glu Asn Ser Thr Gly				
	3880	3885	3890	
gtc cag gac tgc tac cga ggt gat gga cag agt tat cga ggc aca ctc				11769
Val Gln Asp Cys Tyr Arg Gly Asp Gly Gln Ser Tyr Arg Gly Thr Leu				
	3895	3900	3905	
tcc acc act atc aca gga aga aca tgt cag tct tgg tcg tct atg aca				11817
Ser Thr Thr Ile Thr Gly Arg Thr Cys Gln Ser Trp Ser Ser Met Thr				
	3910	3915	3920	
cca cat tgg cat cgg agg atc cca tta tac tat cca aat gct ggc ctg				11865
Pro His Trp His Arg Arg Ile Pro Leu Tyr Tyr Pro Asn Ala Gly Leu				
	3925	3930	3935	3940
acc agg aac tac tgc agg aat cca gat gct gag att cgc cct tgg tgt				11913
Thr Arg Asn Tyr Cys Arg Asn Pro Asp Ala Glu Ile Arg Pro Trp Cys				
	3945	3950	3955	
tac acc atg gat ccc agt gtc agg tgg gag tac tgc aac ctg aca cga				11961
Tyr Thr Met Asp Pro Ser Val Arg Trp Glu Tyr Cys Asn Leu Thr Arg				
	3960	3965	3970	
tgt cca gtg aca gaa tcg agt gtc ctc aca act ccc aca gtg gcc ccg				12009
Cys Pro Val Thr Glu Ser Ser Val Leu Thr Thr Pro Thr Val Ala Pro				
	3975	3980	3985	
gtt cca agc aca gag gct cct tct gaa caa gca cca cct gag aaa agc				12057
Val Pro Ser Thr Glu Ala Pro Ser Glu Gln Ala Pro Pro Glu Lys Ser				
	3990	3995	4000	
cct gtg gtc cag gat tgc tac cat ggt gat gga cgg agt tat cga ggc				12105
Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Arg Ser Tyr Arg Gly				
	4005	4010	4015	4020
ata tcc tcc acc act gtc aca gga agg acc tgt caa tct tgg tca tct				12153
Ile Ser Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ser Trp Ser Ser				
	4025	4030	4035	
atg ata cca cac tgg cat cag agg acc cca gaa aac tac cca aat gct				12201
Met Ile Pro His Trp His Gln Arg Thr Pro Glu Asn Tyr Pro Asn Ala				
	4040	4045	4050	
ggc ctg acc gag aac tac tgc agg aat cca gat tct ggg aaa caa ccc				12249
Gly Leu Thr Glu Asn Tyr Cys Arg Asn Pro Asp Ser Gly Lys Gln Pro				
	4055	4060	4065	
tgg tgt tac aca acc gat ccg tgt gtg agg tgg gag tac tgc aat ctg				12297
Trp Cys Tyr Thr Thr Asp Pro Cys Val Arg Trp Glu Tyr Cys Asn Leu				
	4070	4075	4080	
aca caa tgc tca gaa aca gaa tca ggt gtc cta gag act ccc act gtt				12345
Thr Gln Cys Ser Glu Thr Glu Ser Gly Val Leu Glu Thr Pro Thr Val				
	4085	4090	4095	4100
gtt cca gtt cca agc atg gag gct cat tct gaa gca gca cca act gag				12393
Val Pro Val Pro Ser Met Glu Ala His Ser Glu Ala Ala Pro Thr Glu				
	4105	4110	4115	
caa acc cct gtg gtc cgg cag tgc tac cat ggt aat ggc cag agt tat				12441
Gln Thr Pro Val Val Arg Gln Cys Tyr His Gly Asn Gly Gln Ser Tyr				
	4120	4125	4130	
cga ggc aca ttc tcc acc act gtc aca gga agg aca tgt caa tct tgg				12489
Arg Gly Thr Phe Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ser Trp				
	4135	4140	4145	
tca tcc atg aca cca cac cgg cat cag agg acc cca gaa aac tac cca				12537
Ser Ser Met Thr Pro His Arg His Gln Arg Thr Pro Glu Asn Tyr Pro				

-continued

4150	4155	4160	
aat gat ggc ctg aca atg aac tac tgc agg aat cca gat gcc gat aca Asn Asp Gly Leu Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp Thr 4165	4170	4175	12585 4180
ggc cct tgg tgt ttt acc atg gac ccc agc atc agg tgg gag tac tgc Gly Pro Trp Cys Phe Thr Met Asp Pro Ser Ile Arg Trp Glu Tyr Cys 4185	4190	4195	12633
aac ctg acg cga tgc tca gac aca gaa ggg act gtg gtc gct cct ccg Asn Leu Thr Arg Cys Ser Asp Thr Glu Gly Thr Val Val Ala Pro Pro 4200	4205	4210	12681
act gtc atc cag gtt cca agc cta ggg cct cct tct gaa caa gac tgt Thr Val Ile Gln Val Pro Ser Leu Gly Pro Pro Ser Glu Gln Asp Cys 4215	4220	4225	12729
atg ttt ggg aat ggg aaa gga tac cgg ggc aag aag gca acc act gtt Met Phe Gly Asn Gly Lys Gly Tyr Arg Gly Lys Lys Ala Thr Thr Val 4230	4235	4240	12777
act ggg acg cca tgc cag gaa tgg gct gcc cag gag ccc cat aga cac Thr Gly Thr Pro Cys Gln Glu Trp Ala Ala Gln Glu Pro His Arg His 4245	4250	4255	12825 4260
agc acg ttc att cca ggg aca aat aaa tgg gca ggt ctg gaa aaa aat Ser Thr Phe Ile Pro Gly Thr Asn Lys Trp Ala Gly Leu Glu Lys Asn 4265	4270	4275	12873
tac tgc cgt aac cct gat ggt gac atc aat ggt ccc tgg tgc tac aca Tyr Cys Arg Asn Pro Asp Gly Asp Ile Asn Gly Pro Trp Cys Tyr Thr 4280	4285	4290	12921
atg aat cca aga aaa ctt ttt gac tac tgt gat atc cct ctc tgt gca Met Asn Pro Arg Lys Leu Phe Asp Tyr Cys Asp Ile Pro Leu Cys Ala 4295	4300	4305	12969
tcc tct tca ttt gat tgt ggg aag cct caa gtg gag ccg aag aaa tgt Ser Ser Ser Phe Asp Cys Gly Lys Pro Gln Val Glu Pro Lys Lys Cys 4310	4315	4320	13017
cct gga agc att gta ggg ggg tgt gtg gcc cac cca cat tcc tgg ccc Pro Gly Ser Ile Val Gly Gly Cys Val Ala His Pro His Ser Trp Pro 4325	4330	4335	13065 4340
tgg caa gtc agt ctc aga aca agg ttt gga aag cac ttc tgt gga ggc Trp Gln Val Ser Leu Arg Thr Arg Phe Gly Lys His Phe Cys Gly Gly 4345	4350	4355	13113
acc tta ata tcc cca gag tgg gtg ctg act gct gct cac tgc ttg aag Thr Leu Ile Ser Pro Glu Trp Val Leu Thr Ala Ala His Cys Leu Lys 4360	4365	4370	13161
aag tcc tca agg cct tca tcc tac aag gtc atc ctg ggt gca cac caa Lys Ser Ser Arg Pro Ser Ser Tyr Lys Val Ile Leu Gly Ala His Gln 4375	4380	4385	13209
gaa gtg aac ctc gaa tct cat gtt cag gaa ata gaa gtg tct agg ctg Glu Val Asn Leu Glu Ser His Val Gln Glu Ile Glu Val Ser Arg Leu 4390	4395	4400	13257
ttc ttg gag ccc aca caa gca gat att gcc ttg cta aag cta agc agg Phe Leu Glu Pro Thr Gln Ala Asp Ile Ala Leu Leu Lys Leu Ser Arg 4405	4410	4415	13305 4420
cct gcc gtc atc act gac aaa gta atg cca gct tgt ctg cca tcc cca Pro Ala Val Ile Thr Asp Lys Val Met Pro Ala Cys Leu Pro Ser Pro 4425	4430	4435	13353
gac tac atg gtc acc gcc agg act gaa tgt tac atc act ggc tgg gga Asp Tyr Met Val Thr Ala Arg Thr Glu Cys Tyr Ile Thr Gly Trp Gly 4440	4445	4450	13401
gaa acc caa ggt acc ttt ggg act ggc ctt ctc aag gaa gcc cag ctc Glu Thr Gln Gly Thr Phe Gly Thr Gly Leu Leu Lys Glu Ala Gln Leu 4455			13449

-continued

4455	4460	4465	
ctt gtt att gag aat gaa gtg tgc aat cac tat aag tat att tgt gct			13497
Leu Val Ile Glu Asn Glu Val Cys Asn His Tyr Lys Tyr Ile Cys Ala			
4470	4475	4480	
gag cat ttg gcc aga ggc act gac agt tgc cag ggt gac agt gga ggg			13545
Glu His Leu Ala Arg Gly Thr Asp Ser Cys Gln Gly Asp Ser Gly Gly			
4485	4490	4495	4500
cct ctg gtt tgc ttc gag aag gac aaa tac att tta caa gga gtc act			13593
Pro Leu Val Cys Phe Glu Lys Asp Lys Tyr Ile Leu Gln Gly Val Thr			
4505	4510	4515	
tct tgg ggt ctt ggc tgt gca cgc ccc aat aag cct ggt gtc tat gct			13641
Ser Trp Gly Leu Gly Cys Ala Arg Pro Asn Lys Pro Gly Val Tyr Ala			
4520	4525	4530	
cgt gtt tca agg ttt gtt act tgg att gag gga atg atg aga aat aat			13689
Arg Val Ser Arg Phe Val Thr Trp Ile Glu Gly Met Met Arg Asn Asn			
4535	4540	4545	
taa ttggacggga gacagagtga agcatcaacc tacttagaag ctgaaacgtg			13742
ggtaaggatt tagcatgctg gaaataatag acagcaatca aacgaagaca ctgttcccag			13802
ctaccagcta tgccaaacct tggcattttt ggtatttttg tgtataagct tttaaggctt			13862
gactgacaaa ttctgtatta aggtgtcata gctatgacat ttgttaaaaa taaactctgc			13922
acttattttg atttga			13938
<210> SEQ ID NO 4			
<211> LENGTH: 19			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: PCR Primer			
<400> SEQUENCE: 4			
gaaggtgaag gtcggagtc			19
<210> SEQ ID NO 5			
<211> LENGTH: 20			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: PCR Primer			
<400> SEQUENCE: 5			
gaagatggtg atgggatttc			20
<210> SEQ ID NO 6			
<211> LENGTH: 20			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: PCR Probe			
<400> SEQUENCE: 6			
caagcttccc gttctcagcc			20
<210> SEQ ID NO 7			
<211> LENGTH: 20			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Antisense Oligonucleotide			

-continued

<400> SEQUENCE: 7

ggcaggtcct tcctgtgaca

20

<210> SEQ ID NO 8

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 8

tctgcgtctg agcattgcgt

20

<210> SEQ ID NO 9

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 9

aagcttggca ggttcttcct

20

<210> SEQ ID NO 10

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 10

tcggaggcgc gacggcagtc

20

<210> SEQ ID NO 11

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 11

cggaggcgcg acggcagtcc

20

<210> SEQ ID NO 12

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 12

ggcaggttct tcctgtgaca

20

<210> SEQ ID NO 13

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 13

ataacaataa ggagctgcca

20

-continued

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 14

gaccaagctt ggcaggttct 20

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 15

taacaataag gagctgccac 20

<210> SEQ ID NO 16
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 16

tgaccaagct tggcaggttc 20

<210> SEQ ID NO 17
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 17

ttctgcgtct gagcattgcg 20

<210> SEQ ID NO 18
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 18

aacaataagg agctgccaca 20

<210> SEQ ID NO 19
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 19

acctgacacc gggatccctc 20

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 20
ctgagcattg cgtcaggttg 20

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 21
agtagttcat gatcaagcca 20

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 22
gacggcagtc cttctgctg 20

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 23
ggcaggttct tccagtgaca 20

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 24
tgaccaagct tggcaagttc 20

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 25
tataacacca aggactaatc 20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

-continued

<400> SEQUENCE: 26
ccatctgaca ttgggatcca 20

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 27
tgtggtgtca tagaggacca 20

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 28
atgggacacct ccgatgcaa 20

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 29
acaccaaggg cgaatctcag 20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 30
ttctgtcact ggacatcgtg 20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 31
cacacggatc ggttgtgtaa 20

<210> SEQ ID NO 32
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 32
acatgtcctt cctgtgacag 20

-continued

<210> SEQ ID NO 33
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 33

cagaaggagg ccctaggctt 20

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 34

ctggcgggtga ccatgtagtc 20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 35

tctaagtagg ttgatgcttc 20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 36

tccttaccca cgtttcagct 20

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 37

ggaacagtgt cttcgtttga 20

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 38

gtttgcata gctgtagct 20

<210> SEQ ID NO 39
<211> LENGTH: 20
<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 39

accttaaaag cttatacaca                20

<210> SEQ ID NO 40
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 40

atacagaatt tgtcagtcag                20

<210> SEQ ID NO 41
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense Oligonucleotide

<400> SEQUENCE: 41

gtcatagcta tgacacctta                20

```

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding human apolipoprotein(a), wherein said compound specifically hybridizes with a nucleic acid molecule encoding human apolipoprotein(a) and inhibits the expression of human apolipoprotein(a).

2. The compound of claim 1 which is an antisense oligonucleotide.

3. (Canceled).

4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.

6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.

10. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding human apolipoprotein(a).

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of human apolipoprotein(a) in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of human apolipoprotein(a) is inhibited.

16. A method of treating a human having a disease or condition associated with human apolipoprotein(a) comprising administering to said human a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of human apolipoprotein(a) is inhibited.

17. The method of claim 16 wherein the condition involves abnormal lipid metabolism.

18. The method of claim 16 wherein the condition involves abnormal cholesterol metabolism.

19. The method of claim 16 wherein the condition is atherosclerosis.

20. The method of claim 16 wherein the disease is cardiovascular disease.

* * * * *