

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(22) Date de dépôt/Filing Date: 2008/12/22

(41) Mise à la disp. pub./Open to Public Insp.: 2009/07/09

(45) Date de délivrance/Issue Date: 2020/08/25

(62) Demande originale/Original Application: 2 703 997

(30) Priorités/Priorities: 2007/12/26 (US61/016,793);
2008/02/25 (US61/031,353); 2008/04/18 (US61/046,353);
2008/05/02 (US61/050,172); 2008/07/10 (US61/079,779);
2008/09/22 (US61/099,178)

(51) Cl.Int./Int.Cl. *C07K 16/00* (2006.01),
C07K 16/22 (2006.01), *C07K 16/24* (2006.01),
C07K 16/28 (2006.01), *C07K 16/42* (2006.01),
C12N 15/13 (2006.01)

(72) **Inventeurs/Inventors:**
CHAMBERLAIN, AARON, US;
DAHIYAT, BASSIL, US;
DESJARLAIS, JOHN RUDOLPH, US;
KARKI, SHER BAHADUR, US;
LAZAR, GREGORY ALAN, US

(73) Propriétaire/Owner:
XENCOR, INC., US

(74) Agent: SMART & BIGGAR LLP

(54) Titre : VARIANTS DE FC AVEC UNE LIAISON ALTEREE A FCRN

(54) Title: FC VARIANTS WITH ALTERED BINDING TO FCRN

```

E1 Index
E1 Index 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138
lgp1 A S T K O P S V F P L A P S S K S T S O
lgp2 A S T K O P S V F P L A P S S K S T S O
lgp3 A S T K O P S V F P L A P S S K S T S O
lgp4 A S T K O P S V F P L A P S S K S T S E S

Z1 Index
Z1 Index 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159
lgp1 T A A L O G C L V K D Y F F P E P V T V S W N
lgp2 T A A L O G C L V K D Y F F P E P V T V S W N
lgp3 T A A L O G C L V K D Y F F P E P V T V S W N
lgp4 T A A L O G C L V K D Y F F P E P V T V S W N

E1 Index
E1 Index 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180
lgp1 S G A L T S G V H T F P A V L Q S S O L Y
lgp2 S G A L T S G V H T F P A V L Q S S O L Y
lgp3 S G A L T S G V H T F P A V L Q S S O L Y
lgp4 S G A L T S G V H T F P A V L Q S S O L Y

E1 Index
E1 Index 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201
lgp1 S S L S S V V T V P S S S L G T K T Y C N
lgp2 S S L S S V V T V P S S S L G T K T Y C N
lgp3 S S L S S V V T V P S S S L G T K T Y C N
lgp4 S L S S V V T V P S S S L G T K T Y C N

E1 Index
E1 Index 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220
lgp1 Y N H K P S N T K V D K R V E P K S C
lgp2 Y V H K P S N T K V D K R V E R K C Y G
lgp3 Y N H K P S N T K V D K R V E P K S C
lgp4 Y V H K P S N T K V D K R V E R K C Y G

Hinge
Fc >
E1 Index 221 222 223 224 225 226 227 228
lgp1 R T H T C P P
lgp2 R T H T C P P
lgp3 L G D G T P P C P R C P E P K S C D I T P P
lgp4 L G D G T P P C P S

E1 Index
E1 Index 229 230 231 232 233 234 235 236
lgp1 C P A P E L L A
lgp2 C P A P E L V I A
lgp3 C P A P E L L A
lgp4 E P A S K S C D I T P P C P R

E1 Index
E1 Index 237 238 239 240 241 242 243 244
lgp1 C P A P E L L A
lgp2 C P A P E L V I A
lgp3 C P A P E L L A
lgp4 E P A S K S C D I T P P C P R

```

(continued)

812	index	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257
g0	O	P	S	S	F	L	F	F	F	K	P	K	P	K	D	T	L	M	I	S	R	T
g1	O	P	S	S	F	L	F	F	F	K	P	K	P	K	D	T	L	M	I	S	R	T
g2	O	P	S	S	F	L	F	F	F	K	P	K	P	K	D	T	L	M	I	S	R	T
g3	O	P	S	S	F	L	F	F	F	K	P	K	P	K	D	T	L	M	I	S	R	T
g4	O	P	S	S	F	L	F	F	F	K	P	K	P	K	D	T	L	M	I	S	R	T
813	index	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278
g0	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	E	V	K	F	N	W
g1	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	E	V	K	F	N	W
g2	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	E	V	K	F	N	W
g3	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	E	V	K	F	N	W
g4	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	E	V	K	F	N	W
814	index	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299
g0	V	D	O	Q	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	N	T	S	T
g1	V	D	O	Q	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	N	T	S	T
g2	V	D	O	Q	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	N	T	S	T
g3	V	D	O	Q	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	N	T	S	T
g4	V	D	O	Q	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	N	T	S	T
815	index	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320
g0	U	R	V	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
g1	U	R	V	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
g2	U	R	V	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
g3	U	R	V	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
g4	U	R	V	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
816	index	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341
g0	C	K	C	K	S	S	N	N	K	G	L	P	A	P	I	E	K	T	S	K	A	K
g1	C	K	C	K	S	S	N	N	K	G	L	P	A	P	I	E	K	T	S	K	A	K
g2	C	K	C	K	S	S	N	N	K	G	L	P	A	P	I	E	K	T	S	K	A	K
g3	C	K	C	K	S	S	N	N	K	G	L	P	A	P	I	E	K	T	S	K	A	K
g4	C	K	C	K	S	S	N	N	K	G	L	P	A	P	I	E	K	T	S	K	A	K
817	index	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362
g0	Q	P	R	E	P	Q	V	V	T	L	P	P	S	S	S	S	S	S	S	S	S	S
g1	Q	P	R	E	P	Q	V	V	T	L	P	P	S	S	S	S	S	S	S	S	S	S
g2	Q	P	R	E	P	Q	V	V	T	L	P	P	S	S	S	S	S	S	S	S	S	S
g3	Q	P	R	E	P	Q	V	V	T	L	P	P	S	S	S	S	S	S	S	S	S	S
g4	Q	P	R	E	P	Q	V	V	T	L	P	P	S	S	S	S	S	S	S	S	S	S
818	index	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383
g0	O	V	S	L	T	C	L	V	K	O	F	F	Y	P	S	D	I	A	V	E	W	E
g1	O	V	S	L	T	C	L	V	K	O	F	F	Y	P	S	D	I	A	V	E	W	E
g2	O	V	S	L	T	C	L	V	K	O	F	F	Y	P	S	D	I	A	V	E	W	E
g3	O	V	S	L	T	C	L	V	K	O	F	F	Y	P	S	D	I	A	V	E	W	E
g4	O	V	S	L	T	C	L	V	K	O	F	F	Y	P	S	D	I	A	V	E	W	E
819	index	384	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403
g0	S	N	G	Q	Q	P	P	P	N	N	N	T	T	P	P	L	P	P	L	P	P	P
g1	S	N	G	Q	Q	P	P	P	N	N	N	T	T	P	P	L	P	P	L	P	P	P
g2	S	N	G	Q	Q	P	P	P	N	N	N	T	T	P	P	L	P	P	L	P	P	P
g3	S	N	G	Q	Q	P	P	P	N	N	N	T	T	P	P	L	P	P	L	P	P	P
g4	S	N	G	Q	Q	P	P	P	N	N	N	T	T	P	P	L	P	P	L	P	P	P
820	index	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424
g0	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	N	V	F	S	S	S
g1	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	N	V	F	S	S	S
g2	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	N	V	F	S	S	S
g3	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	N	V	F	S	S	S
g4	F	F	L	Y	S	K	L	T	V	D	K	S	R	W	Q	Q	N	V	F	S	S	S
821	index	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445
g0	C	S	V	M	H	E	L	N	H	N	K	Y	T	Q	K	S	L	S	L	S	P	P
g1	C	S	V	M	H	E	L	N	H	N	K	Y	T	Q	K	S	L	S	L	S	P	P
g2	C	S	V	M	H	E	L	N	H	N	K	Y	T	Q	K	S	L	S	L	S	P	P
g3	C	S	V	M	H	E	L	N	H	N	K	Y	T	Q	K	S	L	S	L	S	P	P
g4	C	S	V	M	H	E	L	N	H	N	K	Y	T	Q	K	S	L	S	L	S	P	P
822	index	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466
g0	O	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K
g1	O	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K
g2	O	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K
g3	O	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K
g4	O	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K	K

(57) **Abrégé/Abstract:**

The present application relates to a variant Fc region comprising at least one modification relative to a wild-type human Fc region, where the modification selected from the group consisting of 434S, 252Y/428L, 252Y/434S, and 428L/434S, and the numbering is according to the EU index.

52620-185D1

ABSTRACT

The present application relates to a variant Fc region comprising at least one modification relative to a wild-type human Fc region, where the modification selected from the group consisting of 434S, 252Y/428L, 252Y/434S, and 428L/434S, and the numbering is according to the EU index.

52620-185D1

Fc VARIANTS WITH ALTERED BINDING TO FcRn

[1] This application is a divisional application of patent application 2703997, which was filed on December 22, 2008 and claims priority to USSN 61/016,793, filed December 26, 2007; USSN 61/031,353, filed February 25, 2008; USSN 61/046,353, filed April 18, 2008; USSN 61/050,172, filed May 2, 2008; USSN 61/079,779, filed July 10, 2008; and USSN 61/099,178, filed September 22, 2008.

FIELD OF THE INVENTION

[2] The present application relates to optimized IgG immunoglobulin variants, engineering methods for their generation, and their application, particularly for therapeutic purposes.

BACKGROUND OF THE INVENTION

[3] Antibodies are immunological proteins that each binds a specific antigen. In most mammals, including humans and mice, antibodies are constructed from paired heavy and light polypeptide chains. Each chain is made up of individual immunoglobulin (Ig) domains, and thus the generic term immunoglobulin is used for such proteins. Each chain is made up of two distinct regions, referred to as the variable and constant regions. The light and heavy chain variable regions show significant sequence diversity between antibodies, and are responsible for binding the target antigen. The constant regions show less sequence diversity, and are responsible for binding a number of natural proteins to elicit important biochemical events. In humans there are five different classes of antibodies including IgA (which includes subclasses IgA1 and IgA2), IgD, IgE, IgG (which includes subclasses IgG1, IgG2, IgG3, and IgG4), and IgM. The distinguishing feature between these antibody classes is their constant regions, although subtler differences may exist in the V region. IgG antibodies are tetrameric proteins composed of two heavy chains and two light chains. The IgG heavy chain is composed of four immunoglobulin domains linked from N- to C-terminus in the order VH-CH1-CH2-CH3, referring to the heavy chain variable domain, heavy chain constant domain 1, heavy chain constant domain 2, and heavy chain constant domain 3 respectively (also referred to as VH-C γ 1-C γ 2-C γ 3, referring to the heavy chain variable domain, constant gamma 1 domain, constant gamma 2 domain, and constant gamma 3 domain respectively). The IgG light chain is composed of two immunoglobulin domains linked from N- to C-terminus in the order VL-CL, referring to the light chain variable domain and the light chain constant domain respectively.

[4] In IgG, a site on Fc between the C γ 2 and C γ 3 domains mediates interaction with the neonatal receptor FcRn. Binding to FcRn recycles endocytosed antibody from the endosome back to the bloodstream (Raghavan et al., 1996, Annu Rev Cell Dev Biol 12:181-220; Ghetie et al., 2000, Annu Rev Immunol 18:739-766). This process, coupled with preclusion of kidney filtration due to the large size of the full-length molecule, resulting in favorable antibody serum half-lives ranging from one to three weeks. Binding of Fc to FcRn also plays a key role in antibody transport. The binding site on Fc for FcRn is also the site at which the

52620-185

bacterial proteins A and G bind. The tight binding by these proteins is typically exploited as a means to purify antibodies by employing protein A or protein G affinity chromatography during protein purification. Thus the fidelity of this region on Fc is important for both the clinical properties of antibodies and their purification. Available structures of the rat Fc/FcRn complex (Burmeister et al., 1994, *Nature*, 372:379-383; Martin et al., 2001, *Mol Cell* 7:867-877), and of the complexes of Fc with proteins A and G (Deisenhofer, 1981, *Biochemistry* 20:2361-2370; Sauer-Eriksson et al., 1995, *Structure* 3:265-278; Tashiro et al., 1995, *Curr Opin Struct Biol* 5:471-481), provide insight into the interaction of Fc with these proteins. The FcRn receptor is also responsible for the transfer of IgG to the neonatal gut and to the lumen of the intestinal epithelia in adults (Ghetie and Ward, *Annu. Rev. Immunol.*, 2000, 18:739-766; Yoshida et al., *Immunity*, 2004, 20(6):769-783).

[5] Studies of rat and human Fc domains have demonstrated the importance of some Fc residues to the binding of FcRn. The rat and human sequences have about 64% sequence identity in the Fc regions (residues 237-443 in the numbering of EU index). See figures 3, 4, and 5 for the rat/human alignments of Fc, FcRn heavy chain, and FcRn light chain (beta-2-microglobulin). A model of the human Fc/FcRn complex has been built from the existing structure of the rat Fc/FcRn complex (Martin et al., 2001, *Mol Cell* 7:867-877). The rat and human sequences share some residues that are critical for FcRn binding, such as H310 and H435 (Medesan et al., 1997 *J. Immunol.* 158(5):221-7; Shields et al., 2001, *J. Biol. Chem.* 276(9):6591-6604). In many positions, however, the human and rat proteins have different amino acids, giving the residues in the human sequence different environments, and possibly a different identities, than in the rat sequence. This variability limits the ability to transfer characteristics from one homolog to the other homolog.

[6] In the murine Fc, random mutation and phage display selection at the sites, T252, T254, and T256 lead to a triple mutant, T252L/T254S/T256F, that has a 3.5-fold increase in FcRn affinity and a 1.5-fold increase in serum half-life (Ghetie et al., 1997, *Nat. Biotech.* 15(7): 637-640). Disruption of the Fc/FcRn interaction by mutations at positions 253, 310 and 435 also lead to decreased *in vivo* half-life (Medesan et al *J. Immunol.* 1997 158(5)2211-7).

[7] Mutational studies in human Fc γ have been done on some of the residues that are important for binding to FcRn and have demonstrated an increased serum half-life. In human Fc γ 1, Hinton et al. mutated three residues individually to the other 19 common amino acids. Hinton et al., found that some point mutants a double mutant increased the FcRn binding affinity (Hinton et al., 2004, *J. Biol. Chem.* 279(8): 6213-6216; Hinton et al. *Journal of Immunology* 2006, 176:346-356). Two mutations had increased half-lives in monkeys. Shields et al. mutated residues, almost exclusively to Ala, and studied their binding to FcRn and

52620-185

the FcγR's (Shields et al., 2001, *J. Biol. Chem.*, 276(9):6591-6604).

[8] Dall'Acqua et al. used phage display to select for Fc mutations that bound FcRn with increased affinity (Dall'Acqua et al. 2002, *J. Immunol.* 169:5171-5180).

The DNA sequences selected for were primarily double and triple mutants. The reference expressed the proteins encoded by many of their selected sequences and found some that bound to FcRn more tightly than the wild-type Fc.

[9] The administration of antibodies and Fc fusion proteins as therapeutics requires injections with a prescribed frequency relating to the clearance and half-life characteristics of the protein. Longer *in vivo* half-lives allow more seldom injections or lower dosing, which is clearly advantageous. Although the past mutations in the Fc domain have lead to some proteins with increased FcRn binding affinity and *in vivo* half-lives, these mutations have not identified the optimal mutations and enhanced *in vivo* half-life.

[10] One feature of the Fc region is the conserved N-linked glycosylation that occurs at N297. This carbohydrate, or oligosaccharide as it is sometimes called, plays a critical structural and functional role for the antibody, and is one of the principle reasons that antibodies must be produced using mammalian expression systems. Umaña et al., 1999, *Nat Biotechnol* 17:176-180; Davies et al., 2001, *Biotechnol Bioeng* 74:288-294; Mimura et al., 2001, *J Biol Chem* 276:45539-45547.; Radaev et al., 2001, *J Biol Chem* 276:16478-16483; Shields et al., 2001, *J Biol Chem* 276:6591-6604; Shields et al., 2002, *J Biol Chem* 277:26733-26740; Simmons et al., 2002, *J Immunol Methods* 263:133-147; Radaev et al., 2001, *J Biol Chem* 276:16469-16477; and Krapp et al., 2003, *J Mol Biol* 325:979-989).

[11] Antibodies have been developed for therapeutic use. Representative publications related to such therapies include Chamow et al., 1996, *Trends Biotechnol* 14:52-60; Ashkenazi et al., 1997, *Curr Opin Immunol* 9:195-200, Cragg et al., 1999, *Curr Opin Immunol* 11:541-547; Glennie et al., 2000, *Immunol Today* 21:403-410, McLaughlin et al., 1998, *J Clin Oncol* 16:2825-2833, and Cobleigh et al., 1999, *J Clin Oncol* 17:2639-2648. Currently for anticancer therapy, any small improvement in mortality rate defines success. Certain IgG variants disclosed herein enhance the capacity of antibodies to limit further growth or destroy at least partially, targeted cancer cells.

[12] Anti-tumor potency of antibodies is via enhancement of their ability to mediate cytotoxic effector functions such as ADCC, ADCP, and CDC. Examples include Clynes et al., 1998, *Proc Natl Acad Sci U S A* 95:652-656; Clynes et al., 2000, *Nat Med* 6:443-446 and Cartron et al., 2002, *Blood* 99:754-758).

[13] Human IgG1 is the most commonly used antibody for therapeutic purposes, and the majority of engineering studies have been constructed in this context. The different isotypes of the IgG class however, including IgG1, IgG2, IgG3, and IgG4, have unique physical, biological, and

clinical properties. There is a need in the art to design improved IgG1, IgG2, IgG3, and IgG4 variants. There is a further need to design such variants to improve binding to FcRn and/or increase *in vivo* half-life as compared to native IgG polypeptides. Additionally, there is a need to combine variants with improved pharmacokinetic properties with variants comprising modifications to improve efficacy through altered FcγR binding. The present application meets these and other needs.

SUMMARY OF THE INVENTION

[14] The present application is directed to Fc variants of a parent polypeptide including at least one modification in the Fc region of the polypeptide. In various embodiments, the variant polypeptides exhibit altered binding to FcRn as compared to a parent polypeptide. In certain variations, the modification can be selected from the group consisting of: 428L, 434M and 434S, where the numbering is according to the EU Index in Kabat et al.

[15] In another embodiment, the Fc variant includes at least two modifications selected from the group consisting of: 252Y/428L, 428L/434H, 428L/434F, 428L/434Y, 428L/434A, 428L/434M, and 428L/434S.

[16] In another embodiment, the Fc variant includes at least one modification selected from the group consisting of: M428L/N434S, V308F/M428L/N434S.

[17] In another embodiment, the Fc variant includes at least one modification selected from the group consisting of: 259I/434S, 308F/434S, 308F/428L/434S, 259I/308F/434S, 307Q/308F/434S, 250I/308F/434S, and 308F/319L/434S.

[18] In another embodiment, the Fc variant includes at least one modification selected from the group consisting of:

[19] In another embodiment, the invention includes a method of treating a patient in need of said treatment comprising administering an effective amount of an Fc variant described herein.

[20] In another embodiment, the invention includes a method of increasing the half-life of an antibody or immunoadhesin by modifying an Fc according to the modifications described herein.

[21] In another variant, the invention includes Fc variant with enhanced FcRn binding with additional Fc variants that modulate effector function.

[21a] The present disclosure includes:

- a polypeptide comprising an Fc variant of a human IgG Fc polypeptide, wherein said Fc variant comprises a serine at position 434 and an isoleucine or a valine at position 311, wherein numbering is according to the EU index; and
- a method of producing the polypeptide of the invention as described herein, said method comprising providing a cell comprising a nucleic acid encoding said polypeptide, wherein said nucleic acid is operably linked with control or regulatory sequences for expression of said polypeptide, and wherein said cell is cultured under conditions suitable for expression of said polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

[22] Figure 1. Sequence alignments of human IgG constant heavy chains. Gray indicates differences from IgG1, and boxed residues indicate common allotypic variations in the human population.

[23] Figure 2. (SEQ ID NO: 1-6) Amino acid sequences of constant regions used in the invention.

[24] Figure 3. (SEQ ID NO: 7-12) Amino acid sequences of exemplary variant constant regions.

[25] Figure 4. (SEQ ID NO: 13-22) Amino acid sequences of VH and VL variable regions used in the invention.

- [26] Figure 5. (SEQ ID NO: 23-29) Amino acid sequences of exemplary variant antibodies.
- [27] Figure 6. Relative VEGF binding by WT and select variant IgG1 anti-VEGF antibodies. The plot shows the Biacore response units (RU) at the end of the association phase for binding of antibody analyte to immobilized VEGF antigen. Anti-Her2 IgG1 antibody was used as a negative control.
- [28] Figure 7. Biacore sensorgrams of WT and variant IgG1 antibodies to immobilized human FcRn at low (6.0) and high (7.4) pH.
- [29] Figure 8. FcRn binding affinities of WT and select variant IgG1 antibodies to human FcRn at pH 6.0 as determined by Biacore. The graph shows a plot of the pseudo-affinity constant (K_a^*), on a log scale.
- [30] Figure 9. Relative binding of variant IgG1 anti-VEGF antibodies to human FcRn as determined by Biacore. The table shows the fold of the K_a^* of each variant relative to human WT (native) IgG1. n indicates the number of time each variant was tested, and Mean and SD indicate the average and standard deviation respectively for each variant over n binding experiments. Fold FcRn was calculated for all variants relative to WT IgG1 within each respective binding experiment. NB indicates no binding was detected. ND indicates that binding was not determined for that particular variant. NF indicates no fit was possible from the binding data.
- [31] Figure 10. Relative binding of variant IgG2 and IgG1/2 anti-VEGF antibodies to human FcRn as determined by Biacore. The table is as described in Figure 9.
- [32] Figure 11. Analysis of additive and synergistic substitution combinations. Figure 11a shows a plot of the experimentally determined fold binding to human FcRn by each variant versus the predicted fold FcRn binding as determined by the product of the single variants. Variant data points are labeled, and the line represents perfect additivity. Figure 11b shows the difference between experimental and predicted fold for each combination variant. Figure 11c shows the synergy of each variant combination. % synergy is calculated as the $100 \times [(\text{experimental fold/predicted fold}) - 1]$.
- [33] Figure 12. Relative binding of variant anti-TNF, -CD25, -EGFR, and -IgE antibodies to human FcRn as determined by Biacore. The table is as described in Figure 9.
- [34] Figure 13. In vivo pharmacokinetics of WT and variant antibodies in mFcRn-/- hFcRn+ mice. The graphs plot the serum concentration of antibody versus time after a single intravenous dose. Figure 13a shows data from one of the 4 studies carried out with IgG1 antibodies (Study 3), and Figure 13b shows data from a study carried out with IgG2 antibodies (Study 5).
- [35] Figure 14. Fitted PK parameters from all in vivo PK studies carried out in mFcRn-/- hFcRn+ mice with variant and WT antibodies. n represents the number of mice per group, with Mean and standard deviation (SD) data provided for PK parameters. Half-Life represents the beta phase that characterizes elimination of antibody from serum. Cmax is the maximal observed serum concentration, AUC is the area under the concentration time curve, and clearance is the clearance

of antibody from serum. Fold half-life is calculated as the half-life of variant antibody over that of the WT IgG1 or IgG2 parent within each study.

[36] Figure 15. Correlation between half-life of IgG1 (Figure 15a) and IgG2 (Figure 15b) variant antibodies in mFcRn^{-/-} hFcRn⁺ mice and fold FcRn binding relative to WT IgG1. Data on the y-axis are from Figure 14, and data on the x-axis are from Figures 9 and 10. Select variants are labeled, and variant data from repeat experiments are circled. Figure 15c shows both IgG1 and IgG2 correlation data, with the black and gray lines representing fits of the IgG1 and IgG2 data respectively.

[37] Figure 16. (SEQ ID NO: 30-35) Amino acid sequences of variant and parent anti-TNF Fc immunoadhesins used in the invention.

[38] Figure 17. Binding of anti-TNF immunoadhesins to TNF antigen as determined by Biacore.

[39] Figure 18. Relative binding of variant Fc immunoadhesins to human FcRn as determined by Biacore. The table shows the fold of the K_a^* of each variant relative to human WT (native) IgG1. n indicates the number of time each variant was tested, and Mean and SD indicate the average and standard deviation respectively for each variant over n binding experiments. Fold FcRn was calculated for all variants relative to the respective IgG parent within each respective binding experiment.

[40] Figure 19. In vivo pharmacokinetics of parent and variant Fc immunoadhesins in mFcRn^{-/-} hFcRn⁺ mice. The graphs plot the serum concentration of Fc fusion versus time after a single intravenous dose.

[41] Figure 20. Fitted PK parameters from the Fc fusion in vivo PK study in mFcRn^{-/-} hFcRn⁺ mice. Parameters are as described in Figure 14. % increase in half-life is calculated as 100 times the half-life of variant Fc fusion over that of the WT IgG1 or IgG2 parent.

[42] Figure 21. Relative binding of variant IgG1 anti-VEGF antibodies to cynomolgus monkey and human FcRn as determined by Biacore. Figure 21a shows the data in tabular form. Description of the figure is as in Figure 9, and data for binding to human FcRn are taken from Figure 9. Figure 21b shows a plot of the data.

[43] Figure 22. In vivo pharmacokinetics of WT and variant antibodies in cynomolgus monkeys. The graphs plot the serum concentration of antibody versus time after a single intravenous dose.

[44] Figure 23. Fitted PK parameters from the in vivo PK study in cynomolgus monkeys with variant and WT antibodies. Parameters are as described in Figure 14.

DETAILED DESCRIPTION OF THE INVENTION

[45] The present invention discloses the generation of novel variants of Fc domains, including those found in antibodies, Fc fusions, and immuno-adhesions, which have an increased binding to the FcRn receptor. As noted herein, binding to FcRn results in longer serum retention *in vivo*.

[46] In order to increase the retention of the Fc proteins *in vivo*, the increase in binding affinity must be at around pH 6 while maintaining lower affinity at around pH 7.4. Although still under

examination, Fc regions are believed to have longer half-lives *in vivo*, because binding to FcRn at pH 6 in an endosome sequesters the Fc (Ghetie and Ward, 1997 Immunol Today. 18(12): 592-598). The endosomal compartment then recycles the Fc to the cell surface. Once the compartment opens to the extracellular space, the higher pH, ~7.4, induces the release of Fc back into the blood. In mice, Dall'Acqua et al. showed that Fc mutants with increased FcRn binding at pH 6 and pH 7.4 actually had reduced serum concentrations and the same half life as wild-type Fc (Dall'Acqua et al. 2002, J. Immunol.

169:5171-5180). The increased affinity of Fc for FcRn at pH 7.4 is thought to forbid the release of the Fc back into the blood. Therefore, the Fc mutations that will increase Fc's half-life *in vivo* will ideally increase FcRn binding at the lower pH while still allowing release of Fc at higher pH. The amino acid histidine changes its charge state in the pH range of 6.0 to 7.4. Therefore, it is not surprising to find His residues at important positions in the Fc/FcRn complex (Figure 6.)

[47] An additional aspect of the invention is the increase in FcRn binding over wild type specifically at lower pH, about pH 6.0, to facilitate Fc/FcRn binding in the endosome. Also disclosed are Fc variants with altered FcRn binding and altered binding to another class of Fc receptors, the FcγR's (sometimes written FcγgammaR's) as differential binding to FcγRs, particularly increased binding to FcγRIIIb and decreased binding to FcγRIIb, has been shown to result in increased efficacy.

[48] *Definitions*

[49] In order that the application may be more completely understood, several definitions are set forth below. Such definitions are meant to encompass grammatical equivalents.

[50] By "ADCC" or "antibody dependent cell-mediated cytotoxicity" as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause lysis of the target cell.

[51] By "ADCP" or "antibody dependent cell-mediated phagocytosis" as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause phagocytosis of the target cell.

[52] By "modification" herein is meant an amino acid substitution, insertion, and/or deletion in a polypeptide sequence or an alteration to a moiety chemically linked to a protein. For example, a modification may be an altered carbohydrate or PEG structure attached to a protein. By "amino acid modification" herein is meant an amino acid substitution, insertion, and/or deletion in a polypeptide sequence.

[53] By "amino acid substitution" or "substitution" herein is meant the replacement of an amino acid at a particular position in a parent polypeptide sequence with another amino acid. For example, the substitution E272Y refers to a variant polypeptide, in this case an Fc variant, in which the glutamic acid at position 272 is replaced with tyrosine.

[54] By "amino acid insertion" or "insertion" as used herein is meant the addition of an amino acid sequence at a particular position in a parent polypeptide sequence. For example, -233E or ^233E designates an insertion of glutamic acid after position 233 and before position 234. Additionally, -233ADE or ^233ADE designates an insertion of AlaAspGlu after position 233 and before position 234.

[55] By "amino acid deletion" or "deletion" as used herein is meant the removal of an amino acid sequence at a particular position in a parent polypeptide sequence. For example, E233- or E233# designates a deletion of glutamic acid at position 233. Additionally, EDA233- or EDA233# designates a deletion of the sequence GluAspAla that begins at position 233.

[56] By "variant protein" or "protein variant", or "variant" as used herein is meant a protein that differs from that of a parent protein by virtue of at least one amino acid modification. Protein variant may refer to the protein itself, a composition comprising the protein, or the amino sequence that encodes it. Preferably, the protein variant has at least one amino acid modification compared to the parent protein, e.g. from about one to about seventy amino acid modifications, and preferably from about one to about five amino acid modifications compared to the parent. The protein variant sequence herein will preferably possess at least about 80% homology with a parent protein sequence, and most preferably at least about 90% homology, more preferably at least about 95% homology. Variant protein can refer to the variant protein itself, compositions comprising the protein variant, or the DNA sequence that encodes it. Accordingly, by "antibody variant" or "variant antibody" as used herein is meant an antibody that differs from a parent antibody by virtue of at least one amino acid modification, "IgG variant" or "variant IgG" as used herein is meant an antibody that differs from a parent IgG by virtue of at least one amino acid modification, and "immunoglobulin variant" or "variant immunoglobulin" as used herein is meant an immunoglobulin sequence that differs from that of a parent immunoglobulin sequence by virtue of at least one amino acid modification. "Fc variant" or "variant Fc" as used herein is meant a protein comprising a modification in an Fc domain. The Fc variants of the present invention are defined according to the amino acid modifications that compose them. Thus, for example, N434S or 434S is an Fc variant with the substitution serine at position 434 relative to the parent Fc polypeptide, wherein the numbering is according to the EU index. Likewise, M428L/N434S defines an Fc variant with the substitutions M428L and N434S. A relative to the parent Fc polypeptide. The identity of the WT amino acid may be unspecified, in which case the aforementioned variant is referred to as 428L/434S. It is noted that the order in which substitutions are provided is arbitrary, that is to say that, for example, 428L/434S is the same Fc variant as M428L/N434S, and so on. For all positions discussed in the present invention, numbering is according to the EU index. The EU index or EU index as in Kabat or EU numbering scheme refers to the numbering of the EU antibody (Edelman et al., 1969, Proc Natl Acad Sci USA 63:78-85.) The modification can be an addition, deletion, or substitution. Substitutions can include

naturally occurring amino acids and non-naturally occurring amino acids. Variants may comprise non-natural amino acids. Examples include US 6,586,207; WO 98/48032; WO 03/073238; US2004-0214988A1; WO 05/35727A2; WO 05/74524A2; J. W. Chin et al., (2002), Journal of the American Chemical Society 124:9026-9027; J. W. Chin, & P. G. Schultz, (2002), ChemBioChem 11:1135-1137; J. W. Chin, et al., (2002), PICAS United States of America 99:11020-11024; and, L. Wang, & P. G. Schultz, (2002), Chem. 1-10.

[57] As used herein, "protein" herein is meant at least two covalently attached amino acids, which includes proteins, polypeptides, oligopeptides and peptides. The peptidyl group may comprise naturally occurring amino acids and peptide bonds, or synthetic peptidomimetic structures, i.e. "analogs", such as peptoids (see Simon et al., PNAS USA 89(20):9367 (1992), entirely incorporated by reference). The amino acids may either be naturally occurring or non-naturally occurring; as will be appreciated by those in the art. For example, homo-phenylalanine, citrulline, and noreleucine are considered amino acids for the purposes of the invention, and both D- and L- (R or S) configured amino acids may be utilized. The variants of the present invention may comprise modifications that include the use of unnatural amino acids incorporated using, for example, the technologies developed by Schultz and colleagues, including but not limited to methods described by Cropp & Shultz, 2004, Trends Genet. 20(12):625-30, Anderson et al., 2004, Proc Natl Acad Sci USA 101(2):7566-71, Zhang et al., 2003, 303(5656):371-3, and Chin et al., 2003, Science 301(5635):964-7. In addition, polypeptides may include synthetic derivatization of one or more side chains or termini, glycosylation, PEGylation, circular permutation, cyclization, linkers to other molecules, fusion to proteins or protein domains, and addition of peptide tags or labels.

[58] By "residue" as used herein is meant a position in a protein and its associated amino acid identity. For example, Asparagine 297 (also referred to as Asn297 or N297) is a residue at position 297 in the human antibody IgG1.

[59] By "Fab" or "Fab region" as used herein is meant the polypeptide that comprises the VH, CH1, VL, and CL immunoglobulin domains. Fab may refer to this region in isolation, or this region in the context of a full length antibody, antibody fragment or Fab fusion protein. By "Fv" or "Fv fragment" or "Fv region" as used herein is meant a polypeptide that comprises the VL and VH domains of a single antibody.

[60] By "IgG subclass modification" as used herein is meant an amino acid modification that converts one amino acid of one IgG isotype to the corresponding amino acid in a different, aligned IgG isotype. For example, because IgG1 comprises a tyrosine and IgG2 a phenylalanine at EU position 296, a F296Y substitution in IgG2 is considered an IgG subclass modification.

[61] By "non-naturally occurring modification" as used herein is meant an amino acid modification that is not isotypic. For example, because none of the IgGs comprise a serine at

position 434, the substitution 434S in IgG1, IgG2, IgG3, or IgG4 is considered a non-naturally occurring modification.

[62] By “amino acid” and “amino acid identity” as used herein is meant one of the 20 naturally occurring amino acids or any non-natural analogues that may be present at a specific, defined position.

[63] By “effector function” as used herein is meant a biochemical event that results from the interaction of an antibody Fc region with an Fc receptor or ligand. Effector functions include but are not limited to ADCC, ADCP, and CDC.

[64] By “IgG Fc ligand” as used herein is meant a molecule, preferably a polypeptide, from any organism that binds to the Fc region of an IgG antibody to form an Fc / Fc ligand complex. Fc ligands include but are not limited to FcγRs, FcγRs, FcγRs, FcRn, C1q, C3, mannan binding lectin, mannose receptor, *staphylococcal* protein A, *streptococcal* protein G, and viral FcγR. Fc ligands also include Fc receptor homologs (FcRH), which are a family of Fc receptors that are homologous to the FcγRs (Davis *et al.*, 2002, *Immunological Reviews* 190:123-136). Fc ligands may include undiscovered molecules that bind Fc. Particular IgG Fc ligands are FcRn and Fc gamma receptors. By “Fc ligand” as used herein is meant a molecule, preferably a polypeptide, from any organism that binds to the Fc region of an antibody to form an Fc / Fc ligand complex.

[65] By “Fc gamma receptor”, “FcγR” or “FcγgammaR” as used herein is meant any member of the family of proteins that bind the IgG antibody Fc region and is encoded by an FcγR gene. In humans this family includes but is not limited to FcγRI (CD64), including isoforms FcγRIa, FcγRIb, and FcγRIc; FcγRII (CD32), including isoforms FcγRIIa (including allotypes H131 and R131), FcγRIIb (including FcγRIIb-1 and FcγRIIb-2), and FcγRIIc; and FcγRIII (CD16), including isoforms FcγRIIIa (including allotypes V158 and F158) and FcγRIIIb (including allotypes FcγRIIIb-NA1 and FcγRIIIb-NA2) (Jefferis *et al.*, 2002, *Immunol Lett* 82:57-65), as well as any undiscovered human FcγRs or FcγR isoforms or allotypes. An FcγR may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. Mouse FcγRs include but are not limited to FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16), and FcγRIII-2 (CD16-2), as well as any undiscovered mouse FcγRs or FcγR isoforms or allotypes.

[66] By “FcRn” or “neonatal Fc Receptor” as used herein is meant a protein that binds the IgG antibody Fc region and is encoded at least in part by an FcRn gene. The FcRn may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. As is known in the art, the functional FcRn protein comprises two polypeptides, often referred to as the heavy chain and light chain. The light chain is beta-2-microglobulin and the heavy chain is encoded by the FcRn gene. Unless other wise noted herein, FcRn or an FcRn protein refers to the complex of

FcRn heavy chain with beta-2-microglobulin. Sequences of particular interest of FcRn are shown in the Figures, particularly the human species.

[67] By "parent polypeptide" as used herein is meant an unmodified polypeptide that is subsequently modified to generate a variant. The parent polypeptide may be a naturally occurring polypeptide, or a variant or engineered version of a naturally occurring polypeptide. Parent polypeptide may refer to the polypeptide itself, compositions that comprise the parent polypeptide, or the amino acid sequence that encodes it. Accordingly, by "parent immunoglobulin" as used herein is meant an unmodified immunoglobulin polypeptide that is modified to generate a variant, and by "parent antibody" as used herein is meant an unmodified antibody that is modified to generate a variant antibody. It should be noted that "parent antibody" includes known commercial, recombinantly produced antibodies as outlined below.

[68] By "position" as used herein is meant a location in the sequence of a protein. Positions may be numbered sequentially, or according to an established format, for example the EU index for antibody numbering.

[69] By "target antigen" as used herein is meant the molecule that is bound specifically by the variable region of a given antibody. A target antigen may be a protein, carbohydrate, lipid, or other chemical compound.

[70] By "target cell" as used herein is meant a cell that expresses a target antigen.

[71] By "variable region" as used herein is meant the region of an immunoglobulin that comprises one or more Ig domains substantially encoded by any of the V κ , V λ , and/or V H genes that make up the kappa, lambda, and heavy chain immunoglobulin genetic loci respectively.

[72] By "wild type or WT" herein is meant an amino acid sequence or a nucleotide sequence that is found in nature, including allelic variations. A WT protein has an amino acid sequence or a nucleotide sequence that has not been intentionally modified.

[73] The present invention is directed to antibodies that exhibit increased binding to FcRn relative to a wild-type antibody. For example, in some instances, increased binding results in cellular recycling of the antibody and hence increased half-life. In addition, antibodies exhibiting increased binding to FcRn and altered binding to other Fc receptors, eg. Fc γ Rs, find use in the present invention.

Antibodies

[74] The present application is directed to antibodies that include amino acid modifications that modulate binding to FcRn. Of particular interest are antibodies that minimally comprise an Fc region, or functional variant thereof, that displays increased binding affinity to FcRn at lowered pH, and do not exhibit substantially altered binding at higher pH.

[75] Traditional antibody structural units typically comprise a tetramer. Each tetramer is typically composed of two identical pairs of polypeptide chains, each pair having one "light" (typically having a molecular weight of about 25 kDa) and one "heavy" chain (typically having a molecular

weight of about 50-70 kDa). Human light chains are classified as kappa and lambda light chains. Heavy chains are classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to IgG1, IgG2, IgG3, and IgG4. IgM has subclasses, including, but not limited to, IgM1 and IgM2. Thus, "isotype" as used herein is meant any of the subclasses of immunoglobulins defined by the chemical and antigenic characteristics of their constant regions. The known human immunoglobulin isotypes are IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgM1, IgM2, IgD, and IgE.

[76] The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. In the variable region, three loops are gathered for each of the V domains of the heavy chain and light chain to form an antigen-binding site. Each of the loops is referred to as a complementarity-determining region (hereinafter referred to as a "CDR"), in which the variation in the amino acid sequence is most significant.

[77] The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Kabat et al. collected numerous primary sequences of the variable regions of heavy chains and light chains. Based on the degree of conservation of the sequences, they classified individual primary sequences into the CDR and the framework and made a list thereof (see SEQUENCES OF IMMUNOLOGICAL INTEREST, 5th edition, NIH publication, No. 91-3242, E.A. Kabat et al.).

[78] In the IgG subclass of immunoglobulins, there are several immunoglobulin domains in the heavy chain. By "immunoglobulin (Ig) domain" herein is meant a region of an immunoglobulin having a distinct tertiary structure. Of interest in the present invention are the heavy chain domains, including, the constant heavy (CH) domains and the hinge domains. In the context of IgG antibodies, the IgG isotypes each have three CH regions. Accordingly, "CH" domains in the context of IgG are as follows: "CH1" refers to positions 118-220 according to the EU index as in Kabat. "CH2" refers to positions 237-340 according to the EU index as in Kabat, and "CH3" refers to positions 341-447 according to the EU index as in Kabat.

[79] Another type of Ig domain of the heavy chain is the hinge region. By "hinge" or "hinge region" or "antibody hinge region" or "immunoglobulin hinge region" herein is meant the flexible polypeptide comprising the amino acids between the first and second constant domains of an antibody. Structurally, the IgG CH1 domain ends at EU position 220, and the IgG CH2 domain begins at residue EU position 237. Thus for IgG the antibody hinge is herein defined to include positions 221 (D221 in IgG1) to 236 (G236 in IgG1), wherein the numbering is according to the EU index as in Kabat. In some embodiments, for example in the context of an Fc region, the lower hinge is included, with the "lower hinge" generally referring to positions 226 or 230.

[80] Of particular interest in the present invention are the Fc regions. By "Fc" or "Fc region", as used herein is meant the polypeptide comprising the constant region of an antibody excluding the first constant region immunoglobulin domain and in some cases, part of the hinge. Thus Fc refers

to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, as illustrated in Figure 1, Fc comprises immunoglobulin domains Cgamma2 and Cgamma3 (Cg2 and Cg3) and the lower hinge region between Cgamma1 (Cg1) and Cgamma2 (Cg2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in Kabat. Fc may refer to this region in isolation, or this region in the context of an Fc polypeptide, as described below. By "Fc polypeptide" as used herein is meant a polypeptide that comprises all or part of an Fc region. Fc polypeptides include antibodies, Fc fusions, isolated Fcs, and Fc fragments.

[81] In some embodiments, the antibodies are full length. By "full length antibody" herein is meant the structure that constitutes the natural biological form of an antibody, including variable and constant regions, including one or more modifications as outlined herein.

[82] Alternatively, the antibodies can be a variety of structures, including, but not limited to, antibody fragments, monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies (sometimes referred to herein as "antibody mimetics"), chimeric antibodies, humanized antibodies, antibody fusions (sometimes referred to as "antibody conjugates"), and fragments of each, respectively.

[83] *Antibody Fragments*

[84] In one embodiment, the antibody is an antibody fragment. Of particular interest are antibodies that comprise Fc regions, Fc fusions, and the constant region of the heavy chain (CH1-hinge-CH2-CH3), again also including constant heavy region fusions.

[85] Specific antibody fragments include, but are not limited to, (i) the Fab fragment consisting of VL, VH, CL and CH1 domains, (ii) the Fd fragment consisting of the VH and CH1 domains, (iii) the Fv fragment consisting of the VL and VH domains of a single antibody; (iv) the dAb fragment (Ward et al., 1989, Nature 341:544-546, entirely incorporated by reference) which consists of a single variable, (v) isolated CDR regions, (vi) F(ab')₂ fragments, a bivalent fragment comprising two linked Fab fragments (vii) single chain Fv molecules (scFv), wherein a VH domain and a VL domain are linked by a peptide linker which allows the two domains to associate to form an antigen binding site (Bird et al., 1988, Science 242:423-426, Huston et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:5879-5883), (viii) bispecific single chain Fv (WO 03/11161, hereby incorporated by reference) and (ix) "diabodies" or "triabodies", multivalent or multispecific fragments constructed by gene fusion (Tomlinson et. al., 2000, Methods Enzymol. 326:461-479; WO94/13804; Holliger et al., 1993, Proc. Natl. Acad. Sci. U.S.A. 90:6444-6448). The antibody fragments may be modified. For example, the molecules may be stabilized by the incorporation of disulphide bridges linking the VH and VL domains (Reiter et al., 1996, Nature Biotech. 14:1239-1245).

[86] *Chimeric and Humanized Antibodies*

[87] In some embodiments, the scaffold components can be a mixture from different species. As such, if the protein is an antibody, such antibody may be a chimeric antibody and/or a humanized antibody. In general, both "chimeric antibodies" and "humanized antibodies" refer to antibodies that combine regions from more than one species. For example, "chimeric antibodies" traditionally comprise variable region(s) from a mouse (or rat, in some cases) and the constant region(s) from a human. "Humanized antibodies" generally refer to non-human antibodies that have had the variable-domain framework regions swapped for sequences found in human antibodies. Generally, in a humanized antibody, the entire antibody, except the CDRs, is encoded by a polynucleotide of human origin or is identical to such an antibody except within its CDRs. The CDRs, some or all of which are encoded by nucleic acids originating in a non-human organism, are grafted into the beta-sheet framework of a human antibody variable region to create an antibody, the specificity of which is determined by the engrafted CDRs. The creation of such antibodies is described in, e.g., WO 92/11018; Jones, 1986, *Nature* 321:522-525, Verhoeyen et al., 1988, *Science* 239:1534-1536. "Backmutation" of selected acceptor framework residues to the corresponding donor residues is often required to regain affinity that is lost in the initial grafted construct (US 5530101; US 5585089; US 5693761; US 5693762; US 6180370; US 5859205; US 5821337; US 6054297; US 6407213). The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin, and thus will typically comprise a human Fc region. Humanized antibodies can also be generated using mice with a genetically engineered immune system. Roque et al., 2004, *Biotechnol. Prog.* 20:639-654, entirely incorporated by reference. A variety of techniques and methods for humanizing and reshaping non-human antibodies are well known in the art (See Tsurushita & Vasquez, 2004, *Humanization of Monoclonal Antibodies*, *Molecular Biology of B Cells*, 533-545, Elsevier Science (USA), and references cited therein). Humanization methods include but are not limited to methods described in Jones et al., 1986, *Nature* 321:522-525; Riechmann et al., 1988, *Nature* 332:323-329; Verhoeyen et al., 1988, *Science*, 239:1534-1536; Queen et al., 1989, *Proc Natl Acad Sci USA* 86:10029-33; He et al., 1998, *J. Immunol.* 160: 1029-1035; Carter et al., 1992, *Proc Natl Acad Sci USA* 89:4285-9, Presta et al., 1997, *Cancer Res.* 57(20):4593-9; Gorman et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:4181-4185; O'Connor et al., 1998, *Protein Eng* 11:321-8. Humanization or other methods of reducing the immunogenicity of nonhuman antibody variable regions may include resurfacing methods, as described for example in Roguska et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:696-973. In one embodiment, the parent antibody has been affinity matured, as is known in the art. Structure-based methods may be employed for humanization and affinity maturation, for example as described in USSN 11/004,590. Selection

based methods may be employed to humanize and/or affinity mature antibody variable regions, including but not limited to methods described in Wu et al., 1999, J. Mol. Biol. 294:151-162; Baca et al., 1997, J. Biol. Chem. 272(16):10678-10684; Rosok et al., 1996, J. Biol. Chem. 271(37): 22611-22618; Rader et al., 1998, Proc. Natl. Acad. Sci. USA 95: 8910-8915; Krauss et al., 2003, Protein Engineering 16(10):753-759. Other humanization methods may involve the grafting of only parts of the CDRs, including but not limited to methods described in USSN 09/810,510; Tan et al., 2002, J. Immunol. 169:1119-1125; De Pascalis et al., 2002, J. Immunol. 169:3076-3084.

[88] *Bispecific Antibodies*

[89] In one embodiment, the antibodies of the invention multispecific antibody, and notably a bispecific antibody, also sometimes referred to as "diabodies". These are antibodies that bind to two (or more) different antigens. Diabodies can be manufactured in a variety of ways known in the art (Holliger and Winter, 1993, Current Opinion Biotechnol. 4:446-449, entirely incorporated by reference), e.g., prepared chemically or from hybrid hybridomas.

[90] *Minibodies*

[91] In one embodiment, the antibody is a minibody. Minibodies are minimized antibody-like proteins comprising a scFv joined to a CH3 domain. Hu et al., 1996, Cancer Res. 56:3055-3061, entirely incorporated by reference. In some cases, the scFv can be joined to the Fc region, and may include some or the entire hinge region.

[92] *Antibody Fusions*

[93] In one embodiment, the antibodies of the invention are antibody fusion proteins (sometimes referred to herein as an "antibody conjugate"). One type of antibody fusions comprises Fc fusions, which join the Fc region with a conjugate partner. By "Fc fusion" as used herein is meant a protein wherein one or more polypeptides is operably linked to an Fc region. Fc fusion is herein meant to be synonymous with the terms "immunoadhesin", "Ig fusion", "Ig chimera", and "receptor globulin" (sometimes with dashes) as used in the prior art (Chamow et al., 1996, Trends Biotechnol 14:52-60; Ashkenazi et al., 1997, Curr Opin Immunol 9:195-200).

An Fc fusion combines the Fc region of an immunoglobulin with a fusion partner, which in general can be any protein or small molecule. Virtually any protein or small molecule may be linked to Fc to generate an Fc fusion. Protein fusion partners may include, but are not limited to, the variable region of any antibody, the target-binding region of a receptor, an adhesion molecule, a ligand, an enzyme, a cytokine, a chemokine, or some other protein or protein domain. Small molecule fusion partners may include any therapeutic agent that directs the Fc fusion to a therapeutic target. Such targets may be any molecule, preferably an extracellular receptor, which is implicated in disease. Thus, the IgG variants can be linked to one or more fusion partners. In one alternate embodiment, the IgG variant is conjugated or operably linked to another therapeutic compound. The therapeutic compound may be a cytotoxic agent, a chemotherapeutic

agent, a toxin, a radioisotope, a cytokine, or other therapeutically active agent. The IgG may be linked to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol.

[94] In addition to Fc fusions, antibody fusions include the fusion of the constant region of the heavy chain with one or more fusion partners (again including the variable region of any antibody), while other antibody fusions are substantially or completely full length antibodies with fusion partners. In one embodiment, a role of the fusion partner is to mediate target binding, and thus it is functionally analogous to the variable regions of an antibody (and in fact can be). Virtually any protein or small molecule may be linked to Fc to generate an Fc fusion (or antibody fusion). Protein fusion partners may include, but are not limited to, the target-binding region of a receptor, an adhesion molecule, a ligand, an enzyme, a cytokine, a chemokine, or some other protein or protein domain. Small molecule fusion partners may include any therapeutic agent that directs the Fc fusion to a therapeutic target. Such targets may be any molecule, preferably an extracellular receptor, which is implicated in disease.

[95] The conjugate partner can be proteinaceous or non-proteinaceous; the latter generally being generated using functional groups on the antibody and on the conjugate partner. For example linkers are known in the art; for example, homo-or hetero-bifunctional linkers as are well known (see, 1994 Pierce Chemical Company catalog, technical section on cross-linkers, pages 155-200).

[96] Suitable conjugates include, but are not limited to, labels as described below, drugs and cytotoxic agents including, but not limited to, cytotoxic drugs (e.g., chemotherapeutic agents) or toxins or active fragments of such toxins. Suitable toxins and their corresponding fragments include diphtheria A chain, exotoxin A chain, ricin A chain, abrin A chain, curcin, crotin, phenomycin, enomycin and the like. Cytotoxic agents also include radiochemicals made by conjugating radioisotopes to antibodies, or binding of a radionuclide to a chelating agent that has been covalently attached to the antibody. Additional embodiments utilize calicheamicin, auristatins, geldanamycin, maytansine, and duocarmycins and analogs; for the latter, see U.S. 2003/0050331A1.

[97] *Covalent modifications of Antibodies*

[98] Covalent modifications of antibodies are included within the scope of this invention, and are generally, but not always, done post-translationally. For example, several types of covalent modifications of the antibody are introduced into the molecule by reacting specific amino acid residues of the antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

[99] Cysteinyll residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyll residues may also be derivatized by reaction with

bromotrifluoroacetone, α -bromo- β -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole and the like.

[100] Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

[101] Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimide; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

[102] Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

[103] The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

[104] Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($\text{R}'\text{---N}=\text{C}=\text{N}\text{---R}'$), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

[105] Derivatization with bifunctional agents is useful for crosslinking antibodies to a water-insoluble support matrix or surface for use in a variety of methods, in addition to methods described below. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis (succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated

carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440, are employed for protein immobilization.

[106] Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention. Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86[1983]), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[107] *Glycosylation*

[108] Another type of covalent modification is glycosylation. In another embodiment, the IgG variants disclosed herein can be modified to include one or more engineered glycoforms. By "engineered glycoform" as used herein is meant a carbohydrate composition that is covalently attached to an IgG, wherein said carbohydrate composition differs chemically from that of a parent IgG. Engineered glycoforms may be useful for a variety of purposes, including but not limited to enhancing or reducing effector function. Engineered glycoforms may be generated by a variety of methods known in the art (Umaña et al., 1999, *Nat Biotechnol* 17:176-180; Davies et al., 2001, *Biotechnol Bioeng* 74:288-294; Shields et al., 2002, *J Biol Chem* 277:26733-26740; Shinkawa et al., 2003, *J Biol Chem* 278:3466-3473; US 6,602,684; USSN 10/277,370; USSN 10/113,929; PCT WO 00/61739A1; PCT WO 01/29246A1; PCT WO 02/31140A1; PCT WO 02/30954A1; (Potelligent® technology [Biowa, Inc., Princeton, NJ]; GlycoMAb® glycosylation engineering technology [Glycart Biotechnology AG, Zürich, Switzerland]). Many of these techniques are based on controlling the level of fucosylated and/or bisecting oligosaccharides that are covalently attached to the Fc region, for example by expressing an IgG in various organisms or cell lines, engineered or otherwise (for example Lec-13 CHO cells or rat hybridoma YB2/0 cells), by regulating enzymes involved in the glycosylation pathway (for example FUT8 [α 1,6-fucosyltransferase] and/or β 1-4- N-acetylglucosaminyltransferase III [GnTIII]), or by modifying carbohydrate(s) after the IgG has been expressed. Engineered glycoform typically refers to the different carbohydrate or oligosaccharide; thus an IgG variant, for example an antibody or Fc fusion, can include an engineered glycoform. Alternatively, engineered glycoform may refer to the IgG variant that comprises the different carbohydrate or oligosaccharide. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

[109] Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[110] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the antibody amino acid sequence is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[111] Another means of increasing the number of carbohydrate moieties on the antibody is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330 and in Aplin and Wriston, 1981, CRC Crit. Rev. Biochem., pp. 259-306.

[112] Removal of carbohydrate moieties present on the starting antibody may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, Arch. Biochem. Biophys. 259:52 and by Edge et al., 1981, Anal. Biochem. 118:131. Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, Meth. Enzymol. 138:350.

Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, J. Biol. Chem. 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

[113] Another type of covalent modification of the antibody comprises linking the antibody to various nonproteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in, for example, 2005-2006 PEG Catalog from Nektar Therapeutics (available at the Nektar website) US Patents 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody to facilitate the addition of polymers such as PEG. See for example, U.S. Publication No. 2005/0114037A1.

[114] *Labeled Antibodies*

[115] In some embodiments, the covalent modification of the antibodies of the invention comprises the addition of one or more labels. In some cases, these are considered antibody fusions. The term "labelling group" means any detectable label. In some embodiments, the labelling group is coupled to the antibody via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used in performing the present invention.

[116] In general, labels fall into a variety of classes, depending on the assay in which they are to be detected: a) isotopic labels, which may be radioactive or heavy isotopes; b) magnetic labels (e.g., magnetic particles); c) redox active moieties; d) optical dyes; enzymatic groups (e.g. horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase); e) biotinylated groups; and f) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.). In some embodiments, the labelling group is coupled to the antibody via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used in performing the present invention.

[117] Specific labels include optical dyes, including, but not limited to, chromophores, phosphors and fluorophores, with the latter being specific in many instances. Fluorophores can be either "small molecule" fluors, or proteinaceous fluors.

[118] By "fluorescent label" is meant any molecule that may be detected via its inherent fluorescent properties. Suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade BlueJ, Texas Red, IAEDANS, EDANS, BODIPY FL, LC Red 640, Cy 5, Cy 5.5, LC Red 705, Oregon green, the Alexa-Fluor dyes (Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 546, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 660, Alexa Fluor 680), Cascade Blue, Cascade Yellow and R-phycoerythrin (PE) (Molecular Probes, Eugene, OR), FITC, Rhodamine, and Texas Red (Pierce, Rockford, IL), Cy5, Cy5.5, Cy7 (Amersham Life Science, Pittsburgh, PA). Suitable optical dyes, including

fluorophores, are described in Molecular Probes Handbook by Richard P. Haugland, entirely incorporated by reference.

[119] Suitable proteinaceous fluorescent labels also include, but are not limited to, green fluorescent protein, including a Renilla, Ptilosarcus, or Aequorea species of GFP (Chalfie et al., 1994, Science 263:802-805), EGFP (Clontech Laboratories, Inc., Genbank Accession Number U55762), blue fluorescent protein (BFP, Quantum Biotechnologies, Inc. 1801 de Maisonneuve Blvd. West, 8th Floor, Montreal, Quebec, Canada H3H 1J9; Stauber, 1998, Biotechniques 24:462-471; Heim et al., 1996, Curr. Biol. 6:178-182), enhanced yellow fluorescent protein (EYFP, Clontech Laboratories, Inc.), luciferase (Ichiki et al., 1993, J. Immunol. 150:5408-5417), β galactosidase (Nolan et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:2603-2607) and Renilla (WO92/15673, WO95/07463, WO98/14605, WO98/26277, WO99/49019, U.S. Patent Nos. 5292658, 5418155, 5683888, 5741668, 5777079, 5804387, 5874304, 5876995, 5925558).

[120] *IgG Variants*

[121] In one embodiment, the invention provides variant IgG proteins. At a minimum, IgG variants comprise an antibody fragment comprising the CH2-CH3 region of the heavy chain. In addition, suitable IgG variants comprise Fc domains (e.g. including the lower hinge region), as well as IgG variants comprising the constant region of the heavy chain (CH1-hinge-CH2-CH3) also being useful in the present invention, all of which can be fused to fusion partners.

[122] An IgG variant includes one or more amino acid modifications relative to a parent IgG polypeptide, in some cases relative to the wild type IgG. The IgG variant can have one or more optimized properties. An IgG variant differs in amino acid sequence from its parent IgG by virtue of at least one amino acid modification. Thus IgG variants have at least one amino acid modification compared to the parent. Alternatively, the IgG variants may have more than one amino acid modification as compared to the parent, for example from about one to fifty amino acid modifications, preferably from about one to ten amino acid modifications, and most preferably from about one to about five amino acid modifications compared to the parent.

[123] Thus the sequences of the IgG variants and those of the parent Fc polypeptide are substantially homologous. For example, the variant IgG variant sequences herein will possess about 80% homology with the parent IgG variant sequence, preferably at least about 90% homology, and most preferably at least about 95% homology. Modifications may be made genetically using molecular biology, or may be made enzymatically or chemically.

[124] *Target Antigens for Antibodies*

[125] Virtually any antigen may be targeted by the IgG variants, including but not limited to proteins, subunits, domains, motifs, and/or epitopes belonging to the following list of target antigens, which includes both soluble factors such as cytokines and membrane-bound factors, including transmembrane receptors: 17-IA, 4-1BB, 4Dc, 6-keto-PGF1a, 8-iso-PGF2a, 8-oxo-dG,

A1 Adenosine Receptor, A33, ACE, ACE-2, Activin, Activin A, Activin AB, Activin B, Activin C, Activin RIA, Activin RIA ALK-2, Activin RIB ALK-4, Activin RIIA, Activin RIIB, ADAM, ADAM10, ADAM12, ADAM15, ADAM17/TACE, ADAM8, ADAM9, ADAMTS, ADAMTS4, ADAMTS5, Addressins, aFGF, ALCAM, ALK, ALK-1, ALK-7, alpha-1-antitrypsin, alpha-V/beta-1 antagonist, ANG, Ang, APAF-1, APE, APJ, APP, APRIL, AR, ARC, ART, Artemin, anti-Id, ASPARTIC, Atrial natriuretic factor, av/b3 integrin, Axl, b2M, B7-1, B7-2, B7-H, B-lymphocyte Stimulator (BlyS), BACE, BACE-1, Bad, BAFF, BAFF-R, Bag-1, BAK, Bax, BCA-1, BCAM, Bcl, BCMA, BDNF, b-ECGF, bFGF, BID, Bik, BIM, BLC, BL-CAM, BLK, BMP, BMP-2 BMP-2a, BMP-3 Osteogenin, BMP-4 BMP-2b, BMP-5, BMP-6 Vgr-1, BMP-7 (OP-1), BMP-8 (BMP-8a, OP-2), BMPR, BMPR-IA (ALK-3), BMPR-IB (ALK-6), BRK-2, RPK-1, BMPR-II (BRK-3), BMPs, b-NGF, BOK, Bombesin, Bone-derived neurotrophic factor, BPDE, BPDE-DNA, BTC, complement factor 3 (C3), C3a, C4, C5, C5a, C10, CA125, CAD-8, Calcitonin, cAMP, carcinoembryonic antigen (CEA), carcinoma-associated antigen, Cathepsin A, Cathepsin B, Cathepsin C/DPPI, Cathepsin D, Cathepsin E, Cathepsin H, Cathepsin L, Cathepsin O, Cathepsin S, Cathepsin V, Cathepsin X/Z/P, CBL, CCL, CCK2, CCL, CCL1, CCL11, CCL12, CCL13, CCL14, CCL15, CCL16, CCL17, CCL18, CCL19, CCL2, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCL26, CCL27, CCL28, CCL3, CCL4, CCL5, CCL6, CCL7, CCL8, CCL9/10, CCR, CCR1, CCR10, CCR10, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CD1, CD2, CD3, CD3E, CD4, CD5, CD6, CD7, CD8, CD10, CD11a, CD11b, CD11c, CD13, CD14, CD15, CD16, CD18, CD19, CD20, CD21, CD22, CD23, CD25, CD27L, CD28, CD29, CD30, CD30L, CD32, CD33 (p67 proteins), CD34, CD38, CD40, CD40L, CD44, CD45, CD46, CD49a, CD52, CD54, CD55, CD56, CD61, CD64, CD66e, CD74, CD80 (B7-1), CD89, CD95, CD123, CD137, CD138, CD140a, CD146, CD147, CD148, CD152, CD164, CEACAM5, CFTR, cGMP, CINC, *Clostridium botulinum* toxin, *Clostridium perfringens* toxin, Ckb8-1, CLC, CMV, CMV UL, CNTF, CNTN-1, COX, C-Ret, CRG-2, CT-1, CTACK, CTGF, CTLA-4, CX3CL1, CX3CR1, CXCL, CXCL1, CXCL2, CXCL3, CXCL4, CXCL5, CXCL6, CXCL7, CXCL8, CXCL9, CXCL10, CXCL11, CXCL12, CXCL13, CXCL14, CXCL15, CXCL16, CXCR, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, cytokeratin tumor-associated antigen, DAN, DCC, DcR3, DC-SIGN, Decay accelerating factor, des(1-3)-IGF-I (brain IGF-1), Dhh, digoxin, DNAM-1, Dnase, Dpp, DPPIV/CD26, Dlk, ECAD, EDA, EDA-A1, EDA-A2, EDAR, EGF, EGFR (ErbB-1), EMA, EMMPRIN, ENA, endothelin receptor, Enkephalinase, eNOS, Eot, eotaxin1, EpCAM, Ephrin B2/ EphB4, EPO, ERCC, E-selectin, ET-1, Factor IIa, Factor VII, Factor VIIIc, Factor IX, fibroblast activation protein (FAP), Fas, FcR1, FEN-1, Ferritin, FGF, FGF-19, FGF-2, FGF3, FGF-8, FGFR, FGFR-3, Fibrin, FL, FLIP, Flt-3, Flt-4, Follicle stimulating hormone, Fractalkine, FZD1, FZD2, FZD3, FZD4, FZD5, FZD6, FZD7, FZD8, FZD9, FZD10, G250, Gas 6, GCP-2, GCSF, GD2, GD3, GDF, GDF-1, GDF-3 (Vgr-2), GDF-5 (BMP-14, CDMP-1), GDF-6 (BMP-13, CDMP-2), GDF-7 (BMP-12, CDMP-3), GDF-8 (Myostatin), GDF-9, GDF-15 (MIC-1), GDNF, GDNF, GFAP, GFRa-1, GFR-alpha1, GFR-alpha2, GFR-alpha3, GITR, Glucagon, Glut 4, glycoprotein IIb/IIIa (GP IIb/IIIa),

GM-CSF, gp130, gp72, GRO, Growth hormone releasing factor, Hapten (NP-cap or NIP-cap), HB-EGF, HCC, HCMV gB envelope glycoprotein, HCMV gH envelope glycoprotein, HCMV UL, Hemopoietic growth factor (HGF), Hep B gp120, heparanase, Her2, Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), herpes simplex virus (HSV) gB glycoprotein, HSV gD glycoprotein, HGFA, High molecular weight melanoma-associated antigen (HMW-MAA), HIV gp120, HIV IIIB gp120 V3 loop, HLA, HLA-DR, HM1.24, HMFG PEM, HRG, Hrk, human cardiac myosin, human cytomegalovirus (HCMV), human growth hormone (HGH), HVEM, I-309, IAP, ICAM, ICAM-1, ICAM-3, ICE, ICOS, IFNg, Ig, IgA receptor, IgE, IGF, IGF binding proteins, IGF-1R, IGFBP, IGF-I, IGF-II, IL, IL-1, IL-1R, IL-2, IL-2R, IL-4, IL-4R, IL-5, IL-5R, IL-6, IL-6R, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-18R, IL-23, interferon (INF)-alpha, INF-beta, INF-gamma, Inhibin, iNOS, Insulin A-chain, Insulin B-chain, Insulin-like growth factor 1, integrin alpha2, integrin alpha3, integrin alpha4, integrin alpha4/beta1, integrin alpha4/beta7, integrin alpha5 (alphaV), integrin alpha5/beta1, integrin alpha5/beta3, integrin alpha6, integrin beta1, integrin beta2, interferon gamma, IP-10, I-TAC, JE, Kallikrein 2, Kallikrein 5, Kallikrein 6, , Kallikrein 11, Kallikrein 12, Kallikrein 14, Kallikrein 15, Kallikrein L1, Kallikrein L2, Kallikrein L3, Kallikrein L4, KC, KDR, Keratinocyte Growth Factor (KGF), laminin 5, LAMP, LAP, LAP (TGF- 1), Latent TGF-1, Latent TGF-1 bp1, LBP, LDGF, LECT2, Lefty, Lewis-Y antigen, Lewis-Y related antigen, LFA-1, LFA-3, Lfo, LIF, LIGHT, lipoproteins, LIX, LKN, Lptn, L-Selectin, LT-a, LT-b, LTB4, LTBP-1, Lung surfactant, Luteinizing hormone, Lymphotoxin Beta Receptor, Mac-1, MAdCAM, MAG, MAP2, MARC, MCAM, MCAM, MCK-2, MCP, M-CSF, MDC, Mer, METALLOPROTEASES, MGDF receptor, MGMT, MHC (HLA-DR), MIF, MIG, MIP, MIP-1-alpha, MK, MMAC1, MMP, MMP-1, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-2, MMP-24, MMP-3, MMP-7, MMP-8, MMP-9, MPIF, Mpo, MSK, MSP, mucin (Muc1), MUC18, Muellierian-inhibitin substance, Mug, MuSK, NAIP, NAP, NCAD, N-Cadherin, NCA 90, NCAM, NCAM, Neprilysin, Neurotrophin-3, -4, or -6, Neurturin, Neuronal growth factor (NGF), NGFR, NGF-beta, nNOS, NO, NOS, Npn, NRG-3, NT, NTN, OB, OGG1, OPG, OPN, OSM, OX40L, OX40R, p150, p95, PADPr, Parathyroid hormone, PARC, PARP, PBR, PBSF, PCAD, P-Cadherin, PCNA, PDGF, PDGF, PDK-1, PECAM, PEM, PF4, PGE, PGF, PGI2, PGJ2, PIN, PLA2, placental alkaline phosphatase (PLAP), PIGF, PLP, PP14, Proinsulin, Prorelaxin, Protein C, PS, PSA, PSCA, prostate specific membrane antigen (PSMA), PTEN, PTHrp, Ptk, PTN, R51, RANK, RANKL, RANTES, RANTES, Relaxin A-chain, Relaxin B-chain, renin, respiratory syncytial virus (RSV) F, RSV Fgp, Ret, Rheumatoid factors, RLIP76, RPA2, RSK, S100, SCF/KL, SDF-1, SERINE, Serum albumin, sFRP-3, Shh, SIGIRR, SK-1, SLAM, SLPI, SMAC, SMDF, SMOH, SOD, SPARC, Stat, STEAP, STEAP-II, TACE, TACI, TAG-72 (tumor-associated glycoprotein-72), TARC, TCA-3, T-cell receptors (e.g., T-cell receptor alpha/beta), TdT, TECK, TEM1, TEM5, TEM7, TEM8, TERT, testicular PLAP-like alkaline phosphatase, TfR, TGF, TGF-alpha, TGF-beta, TGF-beta Pan Specific, TGF-beta RI (ALK-5), TGF-beta RII, TGF-beta RIIB, TGF-beta RIII, TGF-beta1, TGF-beta2, TGF-beta3, TGF-

beta4, TGF-beta5, Thrombin, Thymus Ck-1, Thyroid stimulating hormone, Tie, TIMP, TIQ, Tissue Factor, TMEFF2, Tmpo, TMPRSS2, TNF, TNF-alpha, TNF-alpha beta, TNF-beta2, TNFc, TNF-RI, TNF-RII, TNFRSF10A (TRAIL R1 Apo-2, DR4), TNFRSF10B (TRAIL R2 DR5, KILLER, TRICK-2A, TRICK-B), TNFRSF10C (TRAIL R3 DcR1, LIT, TRID), TNFRSF10D (TRAIL R4 DcR2, TRUNDD), TNFRSF11A (RANK ODF R, TRANCE R), TNFRSF11B (OPG OCIF, TR1), TNFRSF12 (TWEAK R FN14), TNFRSF13B (TACI), TNFRSF13C (BAFF R), TNFRSF14 (HVEM ATAR, HveA, LIGHT R, TR2), TNFRSF16 (NGFR p75NTR), TNFRSF17 (BCMA), TNFRSF18 (GITR AITR), TNFRSF19 (TROY TAJ, TRADE), TNFRSF19L (RELT), TNFRSF1A (TNF RI CD120a, p55-60), TNFRSF1B (TNF RII CD120b, p75-80), TNFRSF26 (TNFRH3), TNFRSF3 (LTbR TNF RIII, TNFC R), TNFRSF4 (OX40 ACT35, TXGP1 R), TNFRSF5 (CD40 p50), TNFRSF6 (Fas Apo-1, APT1, CD95), TNFRSF6B (DcR3 M68, TR6), TNFRSF7 (CD27), TNFRSF8 (CD30), TNFRSF9 (4-1BB CD137, ILA), TNFRSF21 (DR6), TNFRSF22 (DcTRAIL R2 TNFRH2), TNFRST23 (DcTRAIL R1 TNFRH1), TNFRSF25 (DR3 Apo-3, LARD, TR-3, TRAMP, WSL-1), TNFSF10 (TRAIL Apo-2 Ligand, TL2), TNFSF11 (TRANCE/RANK Ligand ODF, OPG Ligand), TNFSF12 (TWEAK Apo-3 Ligand, DR3 Ligand), TNFSF13 (APRIL TALL2), TNFSF13B (BAFF BLYS, TALL1, THANK, TNFSF20), TNFSF14 (LIGHT HVEM Ligand, LTg), TNFSF15 (TL1A/VEGI), TNFSF18 (GITR Ligand AITR Ligand, TL6), TNFSF1A (TNF-a Conectin, DIF, TNFSF2), TNFSF1B (TNF-b LTa, TNFSF1), TNFSF3 (LTb TNFC, p33), TNFSF4 (OX40 Ligand gp34, TXGP1), TNFSF5 (CD40 Ligand CD154, gp39, HIGM1, IMD3, TRAP), TNFSF6 (Fas Ligand Apo-1 Ligand, APT1 Ligand), TNFSF7 (CD27 Ligand CD70), TNFSF8 (CD30 Ligand CD153), TNFSF9 (4-1BB Ligand CD137 Ligand), TP-1, t-PA, Tpo, TRAIL, TRAIL R, TRAIL-R1, TRAIL-R2, TRANCE, transferring receptor, TRF, Trk, TROP-2, TSG, TSLP, tumor-associated antigen CA 125, tumor-associated antigen expressing Lewis Y related carbohydrate, TWEAK, TXB2, Ung, uPAR, uPAR-1, Urokinase, VCAM, VCAM-1, VECAD, VE-Cadherin, VE-cadherin-2, VEGFR-1 (flt-1), VEGF, VEGFR, VEGFR-3 (flt-4), VEGI, VIM, Viral antigens, VLA, VLA-1, VLA-4, VNR integrin, von Willebrands factor, WIF-1, WNT1, WNT2, WNT2B/13, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, WNT16, XCL1, XCL2, XCR1, XCR1, XEDAR, XIAP, XPD, and receptors for hormones and growth factors.

[126] One skilled in the art will appreciate that the aforementioned list of targets refers not only to specific proteins and biomolecules, but the biochemical pathway or pathways that comprise them. For example, reference to CTLA-4 as a target antigen implies that the ligands and receptors that make up the T cell co-stimulatory pathway, including CTLA-4, B7-1, B7-2, CD28, and any other undiscovered ligands or receptors that bind these proteins, are also targets. Thus target as used herein refers not only to a specific biomolecule, but the set of proteins that interact with said target and the members of the biochemical pathway to which said target belongs. One skilled in the art will further appreciate that any of the aforementioned target antigens, the ligands or receptors that bind them, or other members of their corresponding biochemical pathway, may be operably linked

to the Fc variants of the present invention in order to generate an Fc fusion. Thus for example, an Fc fusion that targets EGFR could be constructed by operably linking an Fc variant to EGF, TGF- β , or any other ligand, discovered or undiscovered, that binds EGFR. Accordingly, an Fc variant of the present invention could be operably linked to EGFR in order to generate an Fc fusion that binds EGF, TGF- β , or any other ligand, discovered or undiscovered, that binds EGFR. Thus virtually any polypeptide, whether a ligand, receptor, or some other protein or protein domain, including but not limited to the aforementioned targets and the proteins that compose their corresponding biochemical pathways, may be operably linked to the Fc variants of the present invention to develop an Fc fusion.

[127] The choice of suitable antigen depends on the desired application. For anti-cancer treatment it is desirable to have a target whose expression is restricted to the cancerous cells. Some targets that have proven especially amenable to antibody therapy are those with signaling functions. Other therapeutic antibodies exert their effects by blocking signaling of the receptor by inhibiting the binding between a receptor and its cognate ligand. Another mechanism of action of therapeutic antibodies is to cause receptor down regulation. Other antibodies do not work by signaling through their target antigen. In some cases, antibodies directed against infectious disease agents are used.

[128] In one embodiment, the Fc variants of the present invention are incorporated into an antibody against a cytokine. Alternatively, the Fc variants are fused or conjugated to a cytokine. By "cytokine" as used herein is meant a generic term for proteins released by one cell population that act on another cell as intercellular mediators. For example, as described in Penichet et al., 2001, J Immunol Methods 248:91-101, cytokines may be fused to antibody to provide an array of desirable properties. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF- β ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12; IL-15, a tumor necrosis factor such as TNF- α or TNF- β ; C5a; and other polypeptide factors including LIF and kit ligand

(KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture, and biologically active equivalents of the native sequence cytokines.

[129] Cytokines and soluble targets, such as TNF superfamily members, are preferred targets for use with the variants of the present invention. For example, anti-VEGF, anti-CTLA-4, and anti-TNF antibodies, or fragments thereof, are particularly good antibodies for the use of Fc variants that increase the FcRn binding. Therapeutics against these targets are frequently involved in the treatment of autoimmune diseases and require multiple injections over long time periods. Therefore, longer serum half-lives and less frequent treatments, brought about from the variants of the present invention, are particularly preferred.

[130] A number of antibodies and Fc fusions that are approved for use, in clinical trials, or in development may benefit from the Fc variants of the present invention. These antibodies and Fc fusions are herein referred to as "clinical products and candidates". Thus in a preferred embodiment, the Fc polypeptides of the present invention may find use in a range of clinical products and candidates. For example, a number of antibodies that target CD20 may benefit from the Fc polypeptides of the present invention. For example the Fc polypeptides of the present invention may find use in an antibody that is substantially similar to rituximab (Rituxan®, IDEC/Genentech/Roche) (see for example US 5,736,137), a chimeric anti-CD20 antibody approved to treat Non-Hodgkin's lymphoma; HuMax-CD20, an anti-CD20 currently being developed by Genmab, an anti-CD20 antibody described in US 5,500,362, AME-133 (Applied Molecular Evolution), hA20 (Immunomedics, Inc.), HumaLYM (Intracel), and PRO70769 (PCT/US2003/040426, entitled "Immunoglobulin Variants and Uses Thereof"). A number of antibodies that target members of the family of epidermal growth factor receptors, including EGFR (ErbB-1), Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), may benefit from the Fc polypeptides of the present invention. For example the Fc polypeptides of the present invention may find use in an antibody that is substantially similar to trastuzumab (Herceptin®, Genentech) (see for example US 5,677,171), a humanized anti-Her2/neu antibody approved to treat breast cancer; pertuzumab (rhuMab-2C4, Omnitarg™), currently being developed by Genentech; an anti-Her2 antibody described in US 4,753,894; cetuximab (Erbix®, Imclone) (US 4,943,533; PCT WO 96/40210), a chimeric anti-EGFR antibody in clinical trials for a variety of cancers; ABX-EGF (US 6,235,883), currently being developed by Abgenix-Immunex-Amgen; HuMax-EGFr (USSN 10/172,317), currently being developed by Genmab; 425, EMD55900, EMD62000, and EMD72000 (Merck KGaA) (US 5,558,864; Murthy et al. 1987, Arch Biochem Biophys. 252(2):549-60; Rodeck et al., 1987, J Cell Biochem. 35(4):315-20; Kettleborough et al., 1991, Protein Eng. 4(7):773-83); ICR62 (Institute of Cancer Research) (PCT WO 95/20045; Modjtahedi et al., 1993, J. Cell Biophys. 1993, 22(1-3):129-46; Modjtahedi et al., 1993, Br J Cancer. 1993, 67(2):247-53; Modjtahedi et al, 1996, Br J Cancer, 73(2):228-35; Modjtahedi et al, 2003, Int J Cancer, 105(2):273-80); TheraCIM hR3 (YM Biosciences, Canada and Centro de Immunologia Molecular, Cuba (US 5,891,996; US 6,

506,883; Mateo et al, 1997, Immunotechnology, 3(1):71-81); mAb-806 (Ludwig Institute for Cancer Research, Memorial Sloan-Kettering) (Jungbluth et al. 2003, Proc Natl Acad Sci U S A. 100(2):639-44); KSB-102 (KS Biomedix); MR1-1 (IVAX, National Cancer Institute) (PCT WO 0162931A2); and SC100 (Scancell) (PCT WO 01/88138). In another preferred embodiment, the Fc polypeptides of the present invention may find use in alemtuzumab (Campath®, Millenium), a humanized monoclonal antibody currently approved for treatment of B-cell chronic lymphocytic leukemia. The Fc polypeptides of the present invention may find use in a variety of antibodies or Fc fusions that are substantially similar to other clinical products and candidates, including but not limited to muromonab-CD3 (Orthoclone OKT3®), an anti-CD3 antibody developed by Ortho Biotech/Johnson & Johnson, ibritumomab tiuxetan (Zevalin®), an anti-CD20 antibody developed by IDEC/Schering AG, gemtuzumab ozogamicin (Mylotarg®), an anti-CD33 (p67 protein) antibody developed by Celltech/Wyeth, alefacept (Amevive®), an anti-LFA-3 Fc fusion developed by Biogen, abciximab (ReoPro®), developed by Centocor/Lilly, basiliximab (Simulect®), developed by Novartis, palivizumab (Synagis®), developed by MedImmune, infliximab (Remicade®), an anti-TNFalpha antibody developed by Centocor, adalimumab (Humira®), an anti-TNFalpha antibody developed by Abbott, Humicade™, an anti-TNFalpha antibody developed by Celltech, etanercept (Enbrel®), an anti-TNFalpha Fc fusion developed by Immunex/Amgen, ABX-CBL, an anti-CD147 antibody being developed by Abgenix, ABX-IL8, an anti-IL8 antibody being developed by Abgenix, ABX-MA1, an anti-MUC18 antibody being developed by Abgenix, Pemtumomab (R1549, 90Y-muHMFG1), an anti-MUC1 In development by Antisoma, Therex (R1550), an anti-MUC1 antibody being developed by Antisoma, AngioMab (AS1405), being developed by Antisoma, HuBC-1, being developed by Antisoma, Thioplatin (AS1407) being developed by Antisoma, Antegren® (natalizumab), an anti-alpha-4-beta-1 (VLA-4) and alpha-4-beta-7 antibody being developed by Biogen, VLA-1 mAb, an anti-VLA-1 integrin antibody being developed by Biogen, LTBR mAb, an anti-lymphotoxin beta receptor (LTBR) antibody being developed by Biogen, CAT-152, an anti-TGF-β2 antibody being developed by Cambridge Antibody Technology, J695, an anti-IL-12 antibody being developed by Cambridge Antibody Technology and Abbott, CAT-192, an anti-TGFβ1 antibody being developed by Cambridge Antibody Technology and Genzyme, CAT-213, an anti-Eotaxin1 antibody being developed by Cambridge Antibody Technology, LymphoStat-B™ an anti-Blys antibody being developed by Cambridge Antibody Technology and Human Genome Sciences Inc., TRAIL-R1mAb, an anti-TRAIL-R1 antibody being developed by Cambridge Antibody Technology and Human Genome Sciences, Inc., Avastin™ (bevacizumab, rhuMab-VEGF), an anti-VEGF antibody being developed by Genentech, an anti-HER receptor family antibody being developed by Genentech, Anti-Tissue Factor (ATF), an anti-Tissue Factor antibody being developed by Genentech, Xolair™ (Omalizumab), an anti-IgE antibody being developed by Genentech, Raptiva™ (Efalizumab), an anti-CD11a antibody being developed by Genentech and Xoma, MLN-02 Antibody (formerly LDP-02), being developed by Genentech and Millenium

Pharmaceuticals, HuMax CD4, an anti-CD4 antibody being developed by Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Genmab and Amgen, HuMax-Inflam, being developed by Genmab and Medarex, HuMax-Cancer, an anti-Heparanase I antibody being developed by Genmab and Medarex and Oxford GcoSciences, HuMax-Lymphoma, being developed by Genmab and Amgen, HuMax-TAC, being developed by Genmab, IDEC-131, and anti-CD40L antibody being developed by IDEC Pharmaceuticals, IDEC-151 (Clenoliximab), an anti-CD4 antibody being developed by IDEC Pharmaceuticals, IDEC-114, an anti-CD80 antibody being developed by IDEC Pharmaceuticals, IDEC-152, an anti-CD23 being developed by IDEC Pharmaceuticals, anti-macrophage migration factor (MIF) antibodies being developed by IDEC Pharmaceuticals, BEC2, an anti-idiotypic antibody being developed by Imclone, IMC-1C11, an anti-KDR antibody being developed by Imclone, DC101, an anti-flk-1 antibody being developed by Imclone, anti-VE cadherin antibodies being developed by Imclone, CEA-Cide™ (labetuzumab), an anti-carcinoembryonic antigen (CEA) antibody being developed by Immunomedics, LymphoCide™ (Epratuzumab), an anti-CD22 antibody being developed by Immunomedics, AFP-Cide, being developed by Immunomedics, MyelomaCide, being developed by Immunomedics, LkoCide, being developed by Immunomedics, ProstaCide, being developed by Immunomedics, MDX-010, an anti-CTLA4 antibody being developed by Medarex, MDX-060, an anti-CD30 antibody being developed by Medarex, MDX-070 being developed by Medarex, MDX-018 being developed by Medarex, Osidem™ (IDM-1), and anti-Her2 antibody being developed by Medarex and Immuno-Designed Molecules, HuMax™-CD4, an anti-CD4 antibody being developed by Medarex and Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Medarex and Genmab, CNTO 148, an anti-TNF α antibody being developed by Medarex and Centocor/J&J, CNTO 1275, an anti-cytokine antibody being developed by Centocor/J&J, MOR101 and MOR102, anti-intercellular adhesion molecule-1 (ICAM-1) (CD54) antibodies being developed by MorphoSys, MOR201, an anti-fibroblast growth factor receptor 3 (FGFR-3) antibody being developed by MorphoSys, Nuvion® (visilizumab), an anti-CD3 antibody being developed by Protein Design Labs, HuZAF™, an anti-gamma interferon antibody being developed by Protein Design Labs, Anti- α 5 β 1 Integrin, being developed by Protein Design Labs, anti-IL-12, being developed by Protein Design Labs, ING-1, an anti-Ep-CAM antibody being developed by Xoma, and MLN01, an anti-Beta2 integrin antibody being developed by Xoma.

[131] The Fc polypeptides of the present invention may be incorporated into the aforementioned clinical candidates and products, or into antibodies and Fc fusions that are substantially similar to them. The Fc polypeptides of the present invention may be incorporated into versions of the aforementioned clinical candidates and products that are humanized, affinity matured, engineered, or modified in some other way.

[132] In one embodiment, the Fc polypeptides of the present invention are used for the treatment of autoimmune, inflammatory, or transplant indications. Target antigens and clinical products and candidates that are relevant for such diseases include but are not limited to anti- $\alpha 4\beta 7$ integrin antibodies such as LDP-02, anti-beta2 integrin antibodies such as LDP-01, anti-complement (C5) antibodies such as 5G1.1, anti-CD2 antibodies such as BTI-322, MEDI-507, anti-CD3 antibodies such as OKT3, SMART anti-CD3, anti-CD4 antibodies such as IDEC-151, MDX-CD4, OKT4A, anti-CD11a antibodies, anti-CD14 antibodies such as IC14, anti-CD18 antibodies, anti-CD23 antibodies such as IDEC 152, anti-CD25 antibodies such as Zenapax, anti-CD40L antibodies such as 5c8, Antova, IDEC-131, anti-CD64 antibodies such as MDX-33, anti-CD80 antibodies such as IDEC-114, anti-CD147 antibodies such as ABX-CBL, anti-E-selectin antibodies such as CDP850, anti-gpIIb/IIIa antibodies such as ReoPro/Abcixima, anti-ICAM-3 antibodies such as ICM3, anti-ICE antibodies such as VX-740, anti-FcR1 antibodies such as MDX-33, anti-IgE antibodies such as rhuMab-E25, anti-IL-4 antibodies such as SB-240683, anti-IL-5 antibodies such as SB-240563, SCH55700, anti-IL-8 antibodies such as ABX-IL8, anti-interferon gamma antibodies, anti-TNF (TNF, TNFa, TNFa, TNF-alpha) antibodies such as CDP571, CDP870, D2E7, Infliximab, MAK-195F, and anti-VLA-4 antibodies such as Antegren.

[133] Fc variants of the present invention such as those with increased binding to FcRn may be utilized in TNF inhibitor molecules to provide enhanced properties. Useful TNF inhibitor molecules include any molecule that inhibits the action of TNF-alpha in a mammal. Suitable examples include the Fc fusion Enbrel® (etanercept) and the antibodies Humira® (adalimumab) and Remicade® (infliximab). Monoclonal antibodies (such as Remicade and Humira) engineered using the Fc variants of the present invention to increase FcRn binding, may translate to better efficacy through an increased half-life.

[134] In some embodiments, antibodies against infectious diseases are used. Antibodies against eukaryotic cells include antibodies targeting yeast cells, including but not limited to *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Kluyveromyces fragilis* and *K. lactis*, *Pichia guillerimondii* and *P. pastoris*, *Schizosaccharomyces pombe*, *plasmodium falciparum*, and *Yarrowia lipolytica*.

[135] Antibodies against additional fungal cells are also useful, including target antigens associated with *Candida* strains including *Candida glabrata*, *Candida albicans*, *C. krusei*, *C. lusitaniae* and *C. maltosa*, as well as species of *Aspergillus*, *Cryptococcus*, *Histoplasma*, *Coccidioides*, *Blastomyces*, and *Penicillium*, among others

[136] Antibodies directed against target antigens associated with protozoa include, but are not limited to, antibodies associated with *Trypanosoma*, *Leishmania* species including *Leishmania donovani*; , *Plasmodium spp.*, *Pneumocystis carinii*, *Cryptosporidium parvum*, *Giardia lamblia*, *Entamoeba histolytica*, and *Cyclospora cayetanensis*.

[137] Antibodies against prokaryotic antigens are also useful, including antibodies against suitable bacteria such as pathogenic and non-pathogenic prokaryotes including but not limited to *Bacillus*, including *Bacillus anthracis*; *Vibrio*, e.g. *V. cholerae*; *Escherichia*, e.g. Enterotoxigenic *E. coli*, *Shigella*, e.g. *S. dysenteriae*; *Salmonella*, e.g. *S. typhi*; *Mycobacterium* e.g. *M. tuberculosis*, *M. leprae*; *Clostridium*, e.g. *C. botulinum*, *C. tetani*, *C. difficile*, *C. perfringens*; *Corynebacterium*, e.g. *C. diphtheriae*; *Streptococcus*, *S. pyogenes*, *S. pneumoniae*; *Staphylococcus*, e.g. *S. aureus*; *Haemophilus*, e.g. *H. influenzae*; *Neisseria*, e.g. *N. meningitidis*, *N. gonorrhoeae*; *Yersinia*, e.g. *Y. lambia*, *Y. pestis*, *Pseudomonas*, e.g. *P. aeruginosa*, *P. putida*; *Chlamydia*, e.g. *C. trachomatis*; *Bordetella*, e.g. *B. pertussis*; *Treponema*, e.g. *T. palladium*; *B. anthracis*, *Y. pestis*, *Brucella spp.*, *F. tularensis*, *B. mallei*, *B. pseudomallei*, *B. mallei*, *B. pseudomallei*, *C. botulinum*, *Salmonella spp.*, *SEB V. cholerae toxin B*, *E. coli O157:H7*, *Listeria spp.*, *Trichosporon beigellii*, *Rhodotorula species*, *Hansenula anomala*, *Enterobacter sp.*, *Klebsiella sp.*, *Listeria sp.*, *Mycoplasma sp.* and the like.

[138] In some aspects, the antibodies are directed against viral infections; these viruses include, but are not limited to, including orthomyxoviruses, (e.g. influenza virus), paramyxoviruses (e.g. respiratory syncytial virus, mumps virus, measles virus), adenoviruses, rhinoviruses, coronaviruses, reoviruses, togaviruses (e.g. rubella virus), parvoviruses, poxviruses (e.g. variola virus, vaccinia virus), enteroviruses (e.g. poliovirus, coxsackievirus), hepatitis viruses (including A, B and C), herpesviruses (e.g. Herpes simplex virus, varicella-zoster virus, cytomegalovirus, Epstein-Barr virus), rotaviruses, Norwalk viruses, hantavirus, arenavirus, rhabdovirus (e.g. rabies virus), retroviruses (including HIV, HTLV-I and -II), papovaviruses (e.g. papillomavirus), polyomaviruses, and picornaviruses, and the like.

[139] *Optimized IgG Variant Properties*

[140] The present application also provides IgG variants that are optimized for a variety of therapeutically relevant properties. An IgG variant that is engineered or predicted to display one or more optimized properties is herein referred to as an "optimized IgG variant". The most preferred properties that may be optimized include but are not limited to enhanced or reduced affinity for FcRn and increased or decreased *in vivo* half-life. Suitable embodiments include antibodies that exhibit increased binding affinity to FcRn at lowered pH, such as the pH associated with endosomes, e.g. pH 6.0, while maintaining the reduced affinity at higher pH, such as 7.4., to allow increased uptake into endosomes but normal release rates. Similarly, these antibodies with modulated FcRn binding may optionally have other desirable properties, such as modulated FcγR binding, such as outlined in U.S.S.N.s U.S.S.N.s 11/174,287, 11/124,640, 10/822,231, 10/672,280, 10/379,392, and the patent application entitled IgG Immunoglobulin variants with optimized effector function filed on October 21, 2005 having application no. 11/256,060. That is, optimized properties also include but are not limited to enhanced or reduced affinity for an FcγR. In one optional embodiment, the IgG variants are optimized to possess enhanced affinity for a human

activating FcγR, preferably FcγRIIIa in addition to the FcRn binding profile. In yet another optional alternate embodiment, the IgG variants are optimized to possess reduced affinity for the human inhibitory receptor FcγRIIb. That is, particular embodiments embrace the use of antibodies that show increased binding to FcRn, and increased binding to FcγRIIIa. Other embodiments utilize use of antibodies that show increased binding to FcRn, and increased binding to FcγRIIIa. These embodiments are anticipated to provide IgG polypeptides with enhanced therapeutic properties in humans, for example enhanced effector function and greater anti-cancer potency. In an alternate embodiment, the IgG variants are optimized to have increased or reduced affinity for FcRn and increased or decreased affinity for a human FcγR, including but not limited to FcγRI, FcγRIIa, FcγRIIb, FcγRIIc, FcγRIIIa, and FcγRIIIb including their allelic variations. These embodiments are anticipated to provide IgG polypeptides with enhanced therapeutic properties in humans, for example increased serum half-life and reduced effector function. In other embodiments, IgG variants provide enhanced affinity for FcRn and enhanced affinity for one or more FcγRs, yet reduced affinity for one or more other FcγRs. For example, an IgG variant may have enhanced binding to FcRn and FcγRIIIa, yet reduced binding to FcγRIIb. Alternately, an IgG variant may have reduced binding to FcRn and to FcγR's. In another embodiment, an IgG variant may have reduced affinity for FcRn and enhanced affinity for FcγRIIb, yet reduced affinity to one or more activating FcγRs. In yet another embodiment, an IgG variant may have increased serum half-life and reduced effector functions.

[141] Preferred embodiments comprise optimization of binding to a human FcRn and FcγR, however in alternate embodiments the IgG variants possess enhanced or reduced affinity for FcRn and FcγR from nonhuman organisms, including but not limited to rodents and non-human primates. IgG variants that are optimized for binding to a nonhuman FcRn may find use in experimentation. For example, mouse models are available for a variety of diseases that enable testing of properties such as efficacy, toxicity, and pharmacokinetics for a given drug candidate. As is known in the art, cancer cells can be grafted or injected into mice to mimic a human cancer, a process referred to as xenografting. Testing of IgG variants that comprise IgG variants that are optimized for FcRn may provide valuable information with regard to the clearance characteristics of the protein, its mechanism of clearance, and the like. The IgG variants may also be optimized for enhanced functionality and/or solution properties in aglycosylated form. The Fc ligands include but are not limited to FcRn, FcγRs, C1q, and proteins A and G, and may be from any source including but not limited to human, mouse, rat, rabbit, or monkey, preferably human. In an alternately preferred embodiment, the IgG variants are optimized to be more stable and/or more soluble than the aglycosylated form of the parent IgG variant.

[142] IgG variants can include modifications that modulate interaction with Fc ligands other than FcRn and FcγRs, including but not limited to complement proteins, and Fc receptor homologs

(FcRHs). FcRHs include but are not limited to FcRH1, FcRH2, FcRH3, FcRH4, FcRH5, and FcRH6 (Davis et al., 2002, *Immunol. Reviews* 190:123-136).

[143] Preferably, the Fc ligand specificity of the IgG variant will determine its therapeutic utility. The utility of a given IgG variant for therapeutic purposes will depend on the epitope or form of the target antigen and the disease or indication being treated. For most targets and indications, enhanced FcRn binding may be preferable as the enhanced FcRn binding may result in an increase in serum half-life. Longer serum half-lives allow less frequent dosing or lower dosing of the therapeutic. This is particularly preferable when the therapeutic agent is given in response to an indication that requires repeated administration. For some targets and indications, decreased FcRn affinity may be preferable. This may be particularly preferable when a variant Fc with increased clearance or decreased serum half-life is desired, for example in Fc polypeptides used as imaging agents or radio-therapeutics.

[144] IgG variants may be used that comprise IgG variants that provide enhanced affinity for FcRn with enhanced activating FcγRs and/or reduced affinity for inhibitory FcγRs. For some targets and indications, it may be further beneficial to utilize IgG variants that provide differential selectivity for different activating FcγRs; for example, in some cases enhanced binding to FcγRIIIa and FcγRIIIa may be desired, but not FcγRI, whereas in other cases, enhanced binding only to FcγRIIIa may be preferred. For certain targets and indications, it may be preferable to utilize IgG variants that alter FcRn binding and enhance both FcγR-mediated and complement-mediated effector functions, whereas for other cases it may be advantageous to utilize IgG variants that enhance FcRn binding, or serum half-life, and either FcγR-mediated or complement-mediated effector functions. For some targets or cancer indications, it may be advantageous to reduce or ablate one or more effector functions, for example by knocking out binding to C1q, one or more FcγR's, FcRn, or one or more other Fc ligands. For other targets and indications, it may be preferable to utilize IgG variants that provide enhanced binding to the inhibitory FcγRIIb, yet WT level, reduced, or ablated binding to activating FcγRs. This may be particularly useful, for example, when the goal of an IgG variant is to inhibit inflammation or auto-immune disease, or modulate the immune system in some way. Because auto-immune diseases are generally long-lasting and treatment is given in repeated dosing, their treatment with Fc variants with increased half-life from increased FcRn is particularly preferred.

[145] Modification may be made to improve the IgG stability, solubility, function, or clinical use. In a preferred embodiment, the IgG variants can include modifications to reduce immunogenicity in humans. In a most preferred embodiment, the immunogenicity of an IgG variant is reduced using a method described in USSN 11/004,590, entirely incorporated by reference. In alternate embodiments, the IgG variants are humanized (Clark, 2000, *Immunol Today* 21:397-402).

[146] The IgG variants can include modifications that reduce immunogenicity. Modifications to reduce immunogenicity can include modifications that reduce binding of processed peptides derived from the parent sequence to MHC proteins. For example, amino acid modifications would be engineered such that there are no or a minimal number of immune epitopes that are predicted to bind, with high affinity, to any prevalent MHC alleles. Several methods of identifying MHC-binding epitopes in protein sequences are known in the art and may be used to score epitopes in an IgG variant. See for example WO 98/52976; WO 02/079232; WO 00/3317; USSN 09/903,378; USSN 10/039,170; USSN 60/222,697; USSN 10/754,296; PCT WO 01/21823; and PCT WO 02/00165; Mallios, 1999, *Bioinformatics* 15: 432-439; Mallios, 2001, *Bioinformatics* 17: 942-948; Sturniolo *et al.*, 1999, *Nature Biotech.* 17: 555-561; WO 98/59244; WO 02/069232; WO 02/77187; Marshall *et al.*, 1995, *J. Immunol.* 154: 5927-5933; and Hammer *et al.*, 1994, *J. Exp. Med.* 180:2353-2358. Sequence-based information can be used to determine a binding score for a given peptide – MHC interaction (see for example Mallios, 1999, *Bioinformatics* 15: 432-439; Mallios, 2001, *Bioinformatics* 17: p942-948; Sturniolo *et al.*, 1999, *Nature Biotech.* 17: 555-561).

[147] *Engineering IgG Variants*

[148] Variants of the present invention may be designed by various means. The variants, as described herein, may be insertions, deletions, substitutions, other modifications, or combinations of these and other changes. A particularly novel embodiment of the present invention is the design of insertions and deletions that either improve or reduce the binding of an Fc polypeptide to an Fc ligand. As disclosed herein, insertions or deletions may be made that increase or decrease the affinity of the Fc polypeptide for FcRn. Insertions and deletions may be designed by rational approaches or by approaches that include the use of random components, such as random or semi-random library creation or screening. In an alternative embodiment, substitutions are disclosed that increase or decrease the affinity of the Fc polypeptide for FcRn.

[149] *Backbone Modifications: Insertions and deletions*

[150] Variant Fc polypeptides may be created by substituting a variant amino acid in place of the parent amino acid at a position in the Fc polypeptide. By substituting one or more amino acids for variant amino acids in the Fc polypeptide, the side chains at those positions are altered. Most useful substitutions modify the Fc properties by altering the Fc side chains. The substituted side chains may interact directly or indirectly with an Fc binding partner that is associated with an Fc function or property. The at least one substitution alters the covalent structure of one or more side chains of the parent Fc polypeptide.

[151] Alternatively, variant Fc polypeptides may be created that change the covalent structure of the backbone of the parent Fc polypeptide. The backbone atoms in proteins are the peptide nitrogen, the alpha carbon, the carbonyl or peptide carbon and the carbonyl oxygen. Changing the covalent structure of the backbone provides additional methods of altering the properties of the Fc

polypeptides. The covalent structure of the Fc backbone may be altered by the addition of atoms into the backbone, e.g. by inserting one or more amino acids, or the subtraction of atoms from the backbone, e.g. by deleting one or more amino acids. The covalent structure of the backbone may also be altered by changing individual atoms of the backbone to other atoms (Deechongkit et al., J Am Chem Soc. 2004. 126(51)16762-71). As is known in the art and is illustrated herein, insertions or deletions of amino acids in Fc polypeptides may be done by inserting or deleting the corresponding nucleotides in the DNA encoding the Fc polypeptide. Alternatively, as is known in the art, insertions or deletions of amino acids may be done during synthesis of Fc polypeptides.

[152] The design of insertions or deletions of amino acids that alter the interaction of the Fc polypeptide with one or more binding partners (e.g. FcγR's, FcRn, C1q) may be done by considering the structure of the complex of the Fc polypeptide and its binding partner. In a less preferred embodiment, the design may be done by considering the structure of the Fc polypeptide and information about the Fc region involved in binding the binding partner. This information may be obtained by mutagenesis experiments, phage display experiments, homology comparisons, computer modeling or other means.

[153] Preferred positions in the amino acid sequence for insertions or deletions that affect the Fc binding interactions, but do not affect the overall structure, stability, expression or use of the Fc polypeptide, are in loops that are involved in the Fc/Fc-binding partner interactions. To alter FcRn binding to the Fc polypeptide, positions 244-257, 279-284, 307-317, 383-390, and 428-435 are preferred loop locations for insertions or deletions (numbering from EU index of Kabat et al., Burmeister et al., 1994, Nature, 372:379-383; Martin et al., 2001, Mol Cell 7:867-877).

To alter the Fcγ receptor binding to the Fc polypeptide, positions 229-239, 266-273, 294-299, and 324-331 are preferred loop locations for insertions or deletions (numbering from EU index of Kabat et al., PDB code 1E4K.pdb Sonderrmann et al. Nature. 2000 406:267). Loops are regions of the polypeptide not involved in alpha helical or beta sheet structure. Loop positions are positions that are not in either alpha helical or beta sheet structures (van Holde, Johnson and Ho. Principles of Physical Biochemistry. Prentice Hall, New Jersey 1998, Chapter 1 pp2-67). Loop positions are preferred because the backbone atoms are typically more flexible and less likely involved in hydrogen bonds compared to the backbone atoms of alpha helices and beta sheets. Therefore, the lengthening or shortening of a loop due to an insertion or deletion of one or more amino acids is less likely to lead to large, disruptive changes to the Fc polypeptide, including stability, expression or other problems.

[154] Insertions and deletions may be used to alter the length of the polypeptide. For example, in loop regions, altering the loop length results in altered flexibility and conformational entropy of the loop. Insertions in a loop will generally increase the conformational entropy of the loop, which may

be defined as Boltzman's constant multiplied by the natural logarithm of the number of possible conformations (van Holde, Johnson and Ho. Principles of Physical Biochemistry. Prentice Hall, New Jersey 1998, pp78). By inserting at least one amino acid into a polypeptide, the total number of conformations available to the polypeptide increases. These additional conformations may be beneficial for forming favorable Fc/Fc-binding partner interactions because the Fc polypeptide may use one of the additional conformations in binding the Fc-binding protein. In this case, the insertion may lead to stronger Fc/Fc-binding partner interactions. If the additional conformations are not used in the binding interface, then the insertion may lead to weaker Fc/Fc-binding partner interactions, because the additional conformations would compete with the binding-competent conformation. Similarly, deletion of a polypeptide segment may also lead to either stronger or weaker Fc/Fc binding-partner interactions. If deletion of a segment, which reduces the possible number of backbone conformations, removes the binding-competent conformation, then the deletion may lead to weaker Fc/Fc-binding partner interactions. If the deletion does not remove the binding-competent conformation, then the deletion may lead to stronger Fc/Fc-binding partner interactions because the deletion may remove conformations that compete with the binding-competent conformation.

[155] Insertions and deletions may be used to alter the positions and orientations of the amino acids in the Fc polypeptide. Because insertions and deletions cause a change in the covalent structure of the backbone, they necessarily cause a change in the positions of the backbone atoms. Figure 7 compares the backbone positions at some loop segments, marked L1 to L4, in three different backbones. The reference backbone structure contains four loop segments, whereas the deletion backbone lacks segment L1 and the insertion segment comprises an additional segment before, ie, N-terminal to, segment L1. Deletions and insertions cause the largest change in the backbone structure near the site of the insertion or deletion. By deleting a segment near the N-terminal end of the loop, e.g. segment L1, the loop shortens and the remaining segments shift their position closer to the loop N-terminus. This has the effect of moving the L2 segment toward the prior location of the L1 segment and toward the loop N-terminus. This change in position of the L2 segment toward the L1 segment may strengthen the binding of the Fc/Fc-binding partner complex and is preferred when there is prior information suggesting that the amino acid or amino acids located in L2 make favorable interactions with the Fc-binding partner, when located in L1. For example, if L2 contains alanine and tyrosine and substitution of two L1 amino acids for alanine and tyrosine previously lead to an Fc variant with increased binding, then deletion of L1 may create an Fc variant with increased affinity for the Fc-binding partner.

[156] Similarly, an insertion of a polypeptide segment into an Fc polypeptide at the N-terminal side of a loop causes the positions of the loop segments to be shifted toward the C-terminal side of the loop. In Figure 7, an insertion of one or more amino acids before, i.e. N-terminally to, segment L1 alters the backbone conformation including a shift of the L1 segment toward the C-terminal end

of the loop. This type of insertion is preferred when the amino acids located in segment L1 are known to make favorable interactions when located in the L2 positions, as the insertion may lead to stronger Fc/Fc-binding partner interactions. If weaker Fc/Fc-binding partner interactions are desired, then the insertion may be used to shift unfavorable amino acid into a new position. The inserted, deleted and reference segments (L1 to L4 in Figure 7) may be one or more than one amino acid in the Fc polypeptide.

[157] Alternatively, insertions or deletions may be used at the C-terminal end of loops in a manner analogous to the insertions or deletions at the N-terminal end of loops. Insertions at the loop C-terminus may lead to a movement of the positions N-terminal of the insertion toward the loop N-terminus. Deletions at the loop C-terminus may lead to a movement of the positions N-terminal of the deletion toward the loop C-terminus. The choice of using an insertion or deletion at the N-terminal or C-terminal end of the loop is based on the amino acids located in the loop, the desire for increased or decreased Fc/Fc-binding partner affinity, and the positional shift desired.

[158] Insertions or deletions may be used in any region of an Fc polypeptide, including the loops, the alpha helical, and the beta sheet regions. Preferred locations for insertions and deletions include loop regions, which are those that are not alpha helical or beta sheet regions. Loops are preferred because they generally accept alterations in the backbone better than alpha helices or beta sheets. The particularly preferred locations for insertions or deletions that result in stronger protein/protein interactions are at the N-terminal or C-terminal edges of a loop. If the loop side chains are involved in the Fc/Fc-binding partner interactions, then insertions or deletion at the edges are less likely to lead to strongly detrimental changes in the binding interactions. Deletions within the exact center of the loop are more likely to remove important residues in the Fc/Fc-binding partner interface and insertions within the exact center of the loop are more likely to create unfavorable interactions in the Fc/Fc-binding partner interface. The number of residues deleted or inserted may be determined by the size of the backbone change desired with insertions or deletions of 15 or less residues being preferred, insertions or deletions of 10 or less residues being more preferred, and insertions or deletions of 5 or less residues being most preferred.

[159] Once the position and size of an Fc deletion variant is designed, the entire polypeptide sequence is completely determined and the polypeptide may be constructed by methods known in the art.

[160] Fc insertion variants, however, have the additional step of designing the sequence of the at least one amino acid to be inserted. Insertions of polar residues, including Ser, Thr, Asn, Gln, Ala, Gly, His, are preferred at positions expected to be exposed in the Fc polypeptide. The smaller amino acids, including Ser, Thr, and Ala, are particularly preferred as the small size is less likely to sterically interfere with the Fc/Fc-binding partner interactions. Ser and Thr also have the capability to hydrogen bond with atoms on the Fc-binding partner.

[161] Insertions also have the added flexibility that the inserted polypeptide may be designed to make favorable interactions with the Fc-binding partner as would be desired when stronger Fc/Fc-binding partner binding is desired. The length of the backbone insertion may be determined by modeling the variant backbone with a simple, generic sequence to be inserted. For example, polyserine, polyglycine or polyalanine insertions of different lengths may be constructed and modeled. Modeling may be done by a variety of methods, including homology modeling based on known three-dimensional structures of homologues comprising the insertion, and by computer modeling including MODELLER (M.A. Marti-Renom et al. Annu. Rev. Biophys. Biomol. Struct. 29, 291-325, 2000) and ROSETTA (Kuhlman et al. (2003). Science 302, 1364-8).

Typically, various backbone conformations are initially generated and the final backbone structure may be determined after the identities of the side chain are established. The side chains may be designed by PDA® algorithms (US 6,188,965; 6,269,312; 6,403,312; 6,801,861; 6,804,611; 6,792,356, 6,950,754, and USSN 09/782,004; 09/927,790; 10/101,499; 10/666,307; 10/666,311; 10/218,102).

[162] Insertions and deletions may be made to alter the binding of Fc polypeptides to FcγR in an analogous manner to the described method to alter FcγR-binding properties. Fc domains bind to the FcγR at the position indicated in Figure 1. Structures of the Fc/FcγR complex, including PDB codes 1T89 and 1IIS (Radaev S et al. J. Biol. Chem. 276, p. 16469-16477), demonstrate the interacting residues and loops between the two structures. Mutagenesis results such as those found in US11/124620 and US6737056) all have utility in determining appropriate shifts of backbone positioning.

[163] Insertions and deletions may be designed in any polypeptide besides Fc polypeptides by the methods described herein. For example, insertions or deletions in the TNF superfamily member, APRIL, may be designed with the aid of its three-dimensional structure (PDB code 1XU1.pdb, Hymowitz, et al. (2005) J. Biol. Chem. 280:7218, entirely incorporated by reference). Insertions or deletions may be designed to increase APRIL binding to its receptor, TACI. The loop residues preferred as insertion or deletion sites are residues Ser118-Val124, Asp164-Phe167, Pro192-Ala198, Pro221-Lys226. These loops interact with TACI in the APRIL/TACI complex and mediate binding.

[164] *Polypeptides incorporating variants*

[165] The IgG variants can be based on human IgG sequences, and thus human IgG sequences are used as the "base" sequences against which other sequences are compared, including but not limited to sequences from other organisms, for example rodent and primate sequences. IgG variants may also comprise sequences from other immunoglobulin classes such as IgA, IgE, IgD, IgM, and the like. It is contemplated that, although the IgG variants are engineered in the context of one parent IgG, the variants may be engineered in or "transferred" to the context of another,

second parent IgG. This is done by determining the "equivalent" or "corresponding" residues and substitutions between the first and second IgG, typically based on sequence or structural homology between the sequences of the IgGs. In order to establish homology, the amino acid sequence of a first IgG outlined herein is directly compared to the sequence of a second IgG. After aligning the sequences, using one or more of the homology alignment programs known in the art (for example using conserved residues as between species), allowing for necessary insertions and deletions in order to maintain alignment (i.e., avoiding the elimination of conserved residues through arbitrary deletion and insertion), the residues equivalent to particular amino acids in the primary sequence of the first IgG variant are defined. Alignment of conserved residues preferably should conserve 100% of such residues. However, alignment of greater than 75% or as little as 50% of conserved residues is also adequate to define equivalent residues. Equivalent residues may also be defined by determining structural homology between a first and second IgG that is at the level of tertiary structure for IgGs whose structures have been determined. In this case, equivalent residues are defined as those for which the atomic coordinates of two or more of the main chain atoms of a particular amino acid residue of the parent or precursor (N on N, CA on CA, C on C and O on O) are within 0.13 nm and preferably 0.1 nm after alignment. Alignment is achieved after the best model has been oriented and positioned to give the maximum overlap of atomic coordinates of non-hydrogen protein atoms of the proteins. Regardless of how equivalent or corresponding residues are determined, and regardless of the identity of the parent IgG in which the IgGs are made, what is meant to be conveyed is that the IgG variants discovered by can be engineered into any second parent IgG that has significant sequence or structural homology with the IgG variant. Thus for example, if a variant antibody is generated wherein the parent antibody is human IgG1, by using the methods described above or other methods for determining equivalent residues, the variant antibody may be engineered in another IgG1 parent antibody that binds a different antigen, a human IgG2 parent antibody, a human IgA parent antibody, a mouse IgG2a or IgG2b parent antibody, and the like. Again, as described above, the context of the parent IgG variant does not affect the ability to transfer the IgG variants to other parent IgGs.

[166] Methods for engineering, producing, and screening IgG variants are provided. The described methods are not meant to constrain to any particular application or theory of operation. Rather, the provided methods are meant to illustrate generally that one or more IgG variants may be engineered, produced, and screened experimentally to obtain IgG variants with optimized effector function. A variety of methods are described for designing, producing, and testing antibody and protein variants in USSN 10/754,296, and USSN 10/672,280.

[167] A variety of protein engineering methods may be used to design IgG variants with optimized effector function. In one embodiment, a structure-based engineering method may be used, wherein available structural information is used to guide substitutions, insertions or

deletions. In a preferred embodiment, a computational screening method may be used, wherein substitutions are designed based on their energetic fitness in computational calculations. See for example USSN 10/754,296 and USSN 10/672,280, and references cited therein.

[168] An alignment of sequences may be used to guide substitutions at the identified positions. One skilled in the art will appreciate that the use of sequence information may curb the introduction of substitutions that are potentially deleterious to protein structure. The source of the sequences may vary widely, and include one or more of the known databases, including but not limited to the Kabat database (Northwestern University); Johnson & Wu, 2001, *Nucleic Acids Res.* 29:205-206; Johnson & Wu, 2000, *Nucleic Acids Res.* 28:214-218), the IMGT database (IMGT, the international ImMunoGeneTics information system®; ; Lefranc *et al.*, 1999, *Nucleic Acids Res.* 27:209-212; Ruiz *et al.*, 2000 *Nucleic Acids Res.* 28:219-221; Lefranc *et al.*, 2001, *Nucleic Acids Res.* 29:207-209; Lefranc *et al.*, 2003, *Nucleic Acids Res.* 31:307-310), and VBASE. Antibody sequence information can be obtained, compiled, and/or generated from sequence alignments of germline sequences or sequences of naturally occurring antibodies from any organism, including but not limited to mammals. One skilled in the art will appreciate that the use of sequences that are human or substantially human may further have the advantage of being less immunogenic when administered to a human. Other databases which are more general nucleic acid or protein databases, i.e. not particular to antibodies, include but are not limited to SwissProt, GenBank Entrez, and EMBL Nucleotide Sequence Database. Aligned sequences can include VH, VL, CH, and/or CL sequences. There are numerous sequence-based alignment programs and methods known in the art, and all of these find use in the generation of sequence alignments.

[169] Alternatively, random or semi-random mutagenesis methods may be used to make amino acid modifications at the desired positions. In these cases positions are chosen randomly, or amino acid changes are made using simplistic rules. For example all residues may be mutated to alanine, referred to as alanine scanning. Such methods may be coupled with more sophisticated engineering approaches that employ selection methods to screen higher levels of sequence diversity. As is well known in the art, there are a variety of selection technologies that may be used for such approaches, including, for example, display technologies such as phage display, ribosome display, cell surface display, and the like, as described below.

[170] Methods for production and screening of IgG variants are well known in the art. General methods for antibody molecular biology, expression, purification, and screening are described in Antibody Engineering, edited by Duebel & Kontermann, Springer-Verlag, Heidelberg, 2001; and Hayhurst & Georgiou, 2001, *Curr Opin Chem Biol* 5:683-689; Maynard & Georgiou, 2000, *Annu Rev Biomed Eng* 2:339-76. Also see the methods described in USSN 10/754,296; USSN 10/672,280; and USSN 10/822,231; and 11/124,620.

Preferred variants of the present invention include those found in Figure 8. Alternatively preferred variants of the present invention include those found in Figure 9. Additionally alternatively preferred variants of the present invention include those found in Figure 10. These variants have shown increased binding to the Fc receptor, FcRn, as illustrated in the examples.

[171] Making IgG Variants

[172] The IgG variants can be made by any method known in the art. In one embodiment, the IgG variant sequences are used to create nucleic acids that encode the member sequences, and that may then be cloned into host cells, expressed and assayed, if desired. These practices are carried out using well-known procedures, and a variety of methods that may find use in are described in *Molecular Cloning - A Laboratory Manual*, 3rd Ed. (Maniatis, Cold Spring Harbor Laboratory Press, New York, 2001), and *Current Protocols in Molecular Biology* (John Wiley & Sons). The nucleic acids that encode the IgG variants may be incorporated into an expression vector in order to express the protein. Expression vectors typically include a protein operably linked, that is, placed in a functional relationship, with control or regulatory sequences, selectable markers, any fusion partners, and/or additional elements. The IgG variants may be produced by culturing a host cell transformed with nucleic acid, preferably an expression vector, containing nucleic acid encoding the IgG variants, under the appropriate conditions to induce or cause expression of the protein. A wide variety of appropriate host cells may be used, including but not limited to mammalian cells, bacteria, insect cells, and yeast. For example, a variety of cell lines that may find use are described in the ATCC cell line catalog, available from the American Type Culture Collection. The methods of introducing exogenous nucleic acid into host cells are well known in the art, and will vary with the host cell used.

[173] In a preferred embodiment, IgG variants are purified or isolated after expression. Antibodies may be isolated or purified in a variety of ways known to those skilled in the art. Standard purification methods include chromatographic techniques, electrophoretic, immunological, precipitation, dialysis, filtration, concentration, and chromatofocusing techniques. As is well known in the art, a variety of natural proteins bind antibodies, for example bacterial proteins A, G, and L, and these proteins may find use in purification. Often, purification may be enabled by a particular fusion partner. For example, proteins may be purified using glutathione resin if a GST fusion is employed, Ni²⁺ affinity chromatography if a His-tag is employed, or immobilized anti-flag antibody if a flag-tag is used. For general guidance in suitable purification techniques, see *Antibody Purification: Principles and Practice*, 3rd Ed., Scopes, Springer-Verlag, NY, 1994.

[174] Screening IgG Variants

[175] Fc variants may be screened using a variety of methods, including but not limited to those that use *in vitro* assays, *in vivo* and cell-based assays, and selection technologies. Automation and

high-throughput screening technologies may be utilized in the screening procedures. Screening may employ the use of a fusion partner or label, for example an immune label, isotopic label, or small molecule label such as a fluorescent or colorimetric dye.

[176] In a preferred embodiment, the functional and/or biophysical properties of Fc variants are screened in an *in vitro* assay. In a preferred embodiment, the protein is screened for functionality, for example its ability to catalyze a reaction or its binding affinity to its target.

[177] As is known in the art, subsets of screening methods are those that select for favorable members of a library. The methods are herein referred to as "selection methods", and these methods find use in the present invention for screening Fc variants. When protein libraries are screened using a selection method, only those members of a library that are favorable, that is which meet some selection criteria, are propagated, isolated, and/or observed. A variety of selection methods are known in the art that may find use in the present invention for screening protein libraries. Other selection methods that may find use in the present invention include methods that do not rely on display, such as *in vivo* methods. A subset of selection methods referred to as "directed evolution" methods are those that include the mating or breeding of favorable sequences during selection, sometimes with the incorporation of new mutations.

[178] In a preferred embodiment, Fc variants are screened using one or more cell-based or *in vivo* assays. For such assays, purified or unpurified proteins are typically added exogenously such that cells are exposed to individual variants or pools of variants belonging to a library. These assays are typically, but not always, based on the function of the Fc polypeptide; that is, the ability of the Fc polypeptide to bind to its target and mediate some biochemical event, for example effector function, ligand/receptor binding inhibition, apoptosis, and the like. Such assays often involve monitoring the response of cells to the IgG, for example cell survival, cell death, change in cellular morphology, or transcriptional activation such as cellular expression of a natural gene or reporter gene. For example, such assays may measure the ability of Fc variants to elicit ADCC, ADCP, or CDC. For some assays additional cells or components, that is in addition to the target cells, may need to be added, for example serum complement, or effector cells such as peripheral blood monocytes (PBMCs), NK cells, macrophages, and the like. Such additional cells may be from any organism, preferably humans, mice, rat, rabbit, and monkey. Antibodies may cause apoptosis of certain cell lines expressing the target, or they may mediate attack on target cells by immune cells which have been added to the assay. Methods for monitoring cell death or viability are known in the art, and include the use of dyes, immunochemical, cytochemical, and radioactive reagents. Transcriptional activation may also serve as a method for assaying function in cell-based assays. Alternatively, cell-based screens are performed using cells that have been transformed or transfected with nucleic acids encoding the variants. That is, Fc variants are not added exogenously to the cells.

[179] The biological properties of the IgG variants may be characterized in cell, tissue, and whole organism experiments. As is known in the art, drugs are often tested in animals, including but not limited to mice, rats, rabbits, dogs, cats, pigs, and monkeys, in order to measure a drug's efficacy for treatment against a disease or disease model, or to measure a drug's pharmacokinetics, toxicity, and other properties. The animals may be referred to as disease models. Therapeutics are often tested in mice, including but not limited to nude mice, SCID mice, xenograft mice, and transgenic mice (including knockins and knockouts). Such experimentation may provide meaningful data for determination of the potential of the protein to be used as a therapeutic. Any organism, preferably mammals, may be used for testing. For example because of their genetic similarity to humans, monkeys can be suitable therapeutic models, and thus may be used to test the efficacy, toxicity, pharmacokinetics, or other property of the IgGs. Tests of the in humans are ultimately required for approval as drugs, and thus of course these experiments are contemplated. Thus the IgGs may be tested in humans to determine their therapeutic efficacy, toxicity, immunogenicity, pharmacokinetics, and/or other clinical properties.

[180] *Methods of Using IgG Variants*

[181] The IgG variants may find use in a wide range of products. In one embodiment the IgG variant is a therapeutic, a diagnostic, or a research reagent, preferably a therapeutic. The IgG variant may find use in an antibody composition that is monoclonal or polyclonal. In a preferred embodiment, the IgG variants are used to kill target cells that bear the target antigen, for example cancer cells. In an alternate embodiment, the IgG variants are used to block, antagonize or agonize the target antigen, for example for antagonizing a cytokine or cytokine receptor. In an alternately preferred embodiment, the IgG variants are used to block, antagonize or agonize the target antigen and kill the target cells that bear the target antigen.

[182] The IgG variants may be used for various therapeutic purposes. In a preferred embodiment, an antibody comprising the IgG variant is administered to a patient to treat an antibody-related disorder. A "patient" for the purposes includes humans and other animals, preferably mammals and most preferably humans. By "antibody related disorder" or "antibody responsive disorder" or "condition" or "disease" herein are meant a disorder that may be ameliorated by the administration of a pharmaceutical composition comprising an IgG variant. Antibody related disorders include but are not limited to autoimmune diseases, immunological diseases, infectious diseases, inflammatory diseases, neurological diseases, and oncological and neoplastic diseases including cancer. By "cancer" and "cancerous" herein refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to carcinoma, lymphoma, blastoma, sarcoma (including liposarcoma), neuroendocrine tumors, mesothelioma, schwannoma, meningioma, adenocarcinoma, melanoma, and leukemia and lymphoid malignancies.

[183] In one embodiment, an IgG variant is the only therapeutically active agent administered to a patient. Alternatively, the IgG variant is administered in combination with one or more other therapeutic agents, including but not limited to cytotoxic agents, chemotherapeutic agents, cytokines, growth inhibitory agents, anti-hormonal agents, kinase inhibitors, anti-angiogenic agents, cardioprotectants, or other therapeutic agents. The IgG variants may be administered concomitantly with one or more other therapeutic regimens. For example, an IgG variant may be administered to the patient along with chemotherapy, radiation therapy, or both chemotherapy and radiation therapy. In one embodiment, the IgG variant may be administered in conjunction with one or more antibodies, which may or may not be an IgG variant. In accordance with another embodiment, the IgG variant and one or more other anti-cancer therapies are employed to treat cancer cells *ex vivo*. It is contemplated that such *ex vivo* treatment may be useful in bone marrow transplantation and particularly, autologous bone marrow transplantation. It is of course contemplated that the IgG variants can be employed in combination with still other therapeutic techniques such as surgery.

[184] A variety of other therapeutic agents may find use for administration with the IgG variants. In one embodiment, the IgG is administered with an anti-angiogenic agent. By "anti-angiogenic agent" as used herein is meant a compound that blocks, or interferes to some degree, the development of blood vessels. The anti-angiogenic factor may, for instance, be a small molecule or a protein, for example an antibody, Fc fusion, or cytokine, that binds to a growth factor or growth factor receptor involved in promoting angiogenesis. The preferred anti-angiogenic factor herein is an antibody that binds to Vascular Endothelial Growth Factor (VEGF). In an alternate embodiment, the IgG is administered with a therapeutic agent that induces or enhances adaptive immune response, for example an antibody that targets CTLA-4. In an alternate embodiment, the IgG is administered with a tyrosine kinase inhibitor. By "tyrosine kinase inhibitor" as used herein is meant a molecule that inhibits to some extent tyrosine kinase activity of a tyrosine kinase. In an alternate embodiment, the IgG variants are administered with a cytokine.

[185] Pharmaceutical compositions are contemplated wherein an IgG variant and one or more therapeutically active agents are formulated. Formulations of the IgG variants are prepared for storage by mixing the IgG having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed., 1980), in the form of lyophilized formulations or aqueous solutions. The formulations to be used for *in vivo* administration are preferably sterile. This is readily accomplished by filtration through sterile filtration membranes or other methods. The IgG variants and other therapeutically active agents disclosed herein may also be formulated as immunoliposomes, and/or entrapped in microcapsules.

[186] The concentration of the therapeutically active IgG variant in the formulation may vary from about 0.1 to 100% by weight. In a preferred embodiment, the concentration of the IgG is in the

range of 0.003 to 1.0 molar. In order to treat a patient, a therapeutically effective dose of the IgG variant may be administered. By "therapeutically effective dose" herein is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. Dosages may range from 0.01 to 100 mg/kg of body weight or greater, for example 0.01, 0.1, 1.0, 10, or 50 mg/kg of body weight, with 1 to 10mg/kg being preferred. As is known in the art, adjustments for protein degradation, systemic versus localized delivery, and rate of new protease synthesis, as well as the age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

[187] Administration of the pharmaceutical composition comprising an IgG variant, preferably in the form of a sterile aqueous solution, may be done in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, parenterally, intranasally, intraotically, intraocularly, rectally, vaginally, transdermally, topically (e.g., gels, salves, lotions, creams, etc.), intraperitoneally, intramuscularly, intrapulmonary (e.g., AERx® inhalable technology commercially available from Aradigm, or Inhance® pulmonary delivery system commercially available from Nektar Therapeutics, etc.). Therapeutic described herein may be administered with other therapeutics concomitantly, i.e., the therapeutics described herein may be co-administered with other therapies or therapeutics, including for example, small molecules, other biologicals, radiation therapy, surgery, etc.

EXAMPLES

[188] Examples are provided below to illustrate the present invention. These examples are not meant to constrain the present invention to any particular application or theory of operation. For all positions discussed in the present invention, numbering is according to the EU index as in Kabat (Kabat *et al.*, 1991, *Sequences of Proteins of Immunological Interest*, 5th Ed., United States Public Health Service, National Institutes of Health, Bethesda). Those skilled in the art of antibodies will appreciate that this convention consists of nonsequential numbering in specific regions of an immunoglobulin sequence, enabling a normalized reference to conserved positions in immunoglobulin families. Accordingly, the positions of any given immunoglobulin as defined by the EU index will not necessarily correspond to its sequential sequence.

EXAMPLE 1: DNA construction, expression, and purification of Fc variants

[189] Amino acid modifications were engineered in the Fc region of IgG antibodies to improve their affinity for the neonatal Fc receptor FcRn. Variants were screened in the context of a number of different human IgG constant chains (Figure 2), including IgG1, IgG2, and a hybrid IgG sequences that contains the CH1 and upper hinge of IgG1 and the Fc region of IgG2. It will be appreciated by those skilled in the art that because of the different interactions of the IgG1 and

IgG2 Fc region with Fc γ R_s and complement, these different parent Fc regions will have different Fc γ R- and complement-mediated effector function properties. Exemplary sequences of Fc variants in the context of these parent IgG constant chains are shown in Figure 3.

[190] Fc variants were engineered in the context of an antibody targeting vascular endothelial factor (VEGF). The heavy and light chain variable regions (VH and VL) are those of a humanized version of the antibody A4.6.1, also referred to as bevacizumab (Avastin®), which is approved for the treatment of a variety of cancers. The amino acid sequences of the VH and VL regions of this antibody are shown in Figure 4.

[191] Genes encoding the heavy and light chains of the anti-VEGF antibodies were constructed in the mammalian expression vector pTT5. Human IgG1 and IgG2 constant chain genes were obtained from IMAGE clones and subcloned into the pTT5 vector. The IgG1/2 gene was constructed using PCR mutagenesis. VH and VL genes encoding the anti-VEGF antibodies were synthesized commercially (Blue Heron Biotechnologies, Bothell WA), and subcloned into the vectors encoding the appropriate CL, IgG1, IgG2, and IgG1/2 constant chains. Amino acid modifications were constructed using site-directed mutagenesis using the QuikChange® site-directed mutagenesis methods (Stratagene, La Jolla CA). All DNA was sequenced to confirm the fidelity of the sequences.

[192] Plasmids containing heavy chain gene (VH-C γ 1-C γ 2-C γ 3) were co-transfected with plasmid containing light chain gene (VL-C κ) into 293E cells using lipofectamine (Invitrogen, Carlsbad CA) and grown in FreeStyle 293 media (Invitrogen, Carlsbad CA). After 5 days of growth, the antibodies were purified from the culture supernatant by protein A affinity using the MabSelect resin (GE Healthcare). Antibody concentrations were determined by bicinchoninic acid (BCA) assay (Pierce).

EXAMPLE 2. Fc variant antibodies maintain binding to antigen

[193] The fidelity of the expressed variant antibodies was confirmed by demonstrating that they maintained specificity for antigen. VEGF binding was monitored using surface plasmon resonance (SPR, Biacore), performed using a Biacore 3000 instrument. Recombinant VEGF (VEGF-165, PeproTech, Rocky Hill, NJ) was adhered to a CM5 chip surface by coupling with N-hydroxysuccinimide/N-ethyl-N'-(-3-dimethylamino-propyl) carbodiimide (NHS/EDC) using standard methods. WT and variant antibodies were injected as analytes, and response, measured in resonance units (RU), was acquired. The dissociation phase was too slow to measure true equilibrium constants, and so relative binding was determined by measuring RU's at the end of the association phase, which should be proportional to the protein concentration (which is held constant in the experiment) and the association rate constant. The data (Figure 6) show that the variant anti-VEGF antibodies maintain binding to antigen, in contrast to the negative control anti-Her2 antibody which does not bind VEGF.

EXAMPLE 3. Measurement of binding to human FcRn

[194] Binding of variant antibodies to human FcRn was measured at pH 6.0, the pH at which it is naturally bound in endosomes. Vectors encoding beta 2 microglobulin and His-tagged alpha chain genes of FcRn were constructed, co-transfected into 293T cells, and purified using nickel chromatography. Antibody affinity for human FcRn (hFcRn) at pH 6.0 was measured on a Biacore 3000 instrument by coupling human FcRn to a CM5 chip surface using standard NHS/EDC chemistry. WT and variant antibodies were used in the mobile phase at 25-100 nM concentration and response was measured in resonance units. Association and dissociation phases at pH 6.0 were acquired, followed by an injection of pH 7.4 buffer to measure release of antibody from receptor at the higher pH. A cycle with antibody and buffer only provided baseline response, which was subtracted from each sample sensorgram.

[195] Figure 7 shows Biacore sensorgrams for binding of native IgG1 and select Fc variant antibodies to human FcRn at the two relevant pH's. The data show that wild-type and variant antibodies bind readily to FcRn chip at pH 6.0 and dissociate slowly at that pH, as they would in the endosome, yet release rapidly at pH 7.4, as they would upon recycling of endosome to the membrane and exposure to the higher pH of serum.

[196] The FcRn association/dissociation curves did not fit to a simple Langmuir model, possibly due to the antibody and receptor multi-valency or chip heterogeneity. Pseudo-Ka values (referred to as Ka*) were determined by fitting to a conformational change model with the change in refractive index (RI) fixed at 0 RU. These values for select variant antibodies are plotted in Figure 8. The relative affinity of each variant as compared to its parent IgG was calculated according to the equation $\text{Fold} = (\text{WT Ka}^* / \text{Variant Ka}^*)$. The relative binding data for all Fc variants in an IgG1 Fc region are presented in Figure 9, and binding data for variants in antibodies with an IgG2 Fc region (constant chains IgG1 and IgG1/2) are presented in Figure 10. For many variants, the binding experiment was repeated multiple times (n), for which folds were calculated with reference to the WT IgG parent within each particular binding experiment. Averaging of these data provided a mean and standard deviation, as presented in Figures 9 and 10.

[197] Figures 9 and 10 show that a number of engineered variants bind with greater affinity to human FcRn binding at pH 6.0 relative to WT IgG1. Improvements were heavily dependent on the identity of the substitution at a given position. For example, using 2-fold as a criteria for improved binding, a number of mutations at position 434 in IgG2 increased affinity (A, S, Y, F, and W), some were neutral (within 2-fold of WT IgG2) (G, H, M, and T), and a number of substitutions reduced affinity (< 0.5 fold) (D, E, K, P, R, and V). Greater binding in the context of IgG1 did not necessarily translate to greater binding in IgG2 (for example 434T was improved in binding in IgG1 but not IgG2). Moreover, improvements provided by single variants were not always additive upon combination. Figure 11a demonstrates this graphically by plotting the experimental fold FcRn binding by select double substitution variants versus the product of the fold FcRn binding by the individual single variants that compose them. The straight line represents perfect additivity, i.e. the

value that would be expected or predicted from the product of the single substitutions. A number of double variants fall on or close to this line (259I/319I, 259I/428L, 319I/428L, and 308F/428L). Several variants are less than additive (319I/308F, 252Y/428L, and 428L/434M). For these variants, particularly in the case of the latter two (252Y/428L and 428L/434M), the affinity improvements of the single substitutions would seem to be incompatible with each other when combined. Surprisingly, the FcRn affinity improvements of variants 259I/308F and 428L/434S were greater than would be expected from the affinities of their respective single substitutions. These particular single substitutions had unexpected synergistic improvements when combined. The difference between experimental affinities and those predicted from the affinities of the single variants are plotted in Figure 11b, with variants grouped according to their composite single variants (259I, 308F, and 319I on the left, and combinations with 482L on the right). Synergy can be quantitated by calculating the fold of the experimental value relative to the predicted value, followed by normalization to 1 and conversion to a percentage (% synergy = $100 \times [(\text{experimental fold/predicted fold}) - 1]$). This analysis is plotted in Figure 11b, with variants grouped according to their composite single variants. This graph highlights again the synergy of some of the variants, particularly 259I/308F and 428L/434S. Figures 11b and 11c also emphasize the nonpredictive nature of combining many of the best single substitutions from the screen. For example, whereas combination of 428L with 434S and 259I provided synergistic binding improvements, 252Y or 434M had a negative impact when combined with 428L. The dramatic difference between combining 428L with 434S versus 434M further highlights the importance of the particular amino acid identity of the substitution at a given position.

EXAMPLE 4. Testing of variants in other antibody contexts

[198] Select variants were constructed in the context of antibodies targeting other antigens, including TNF (TNF α), CD25 (TAC), EGFR, and IgE. Figure 4 provides the amino acid sequences of the VH and VL regions of antibodies targeting these antigens that were used in the invention. The WT and Fc variant anti-TNF antibodies contain the variable region of the fully human antibody adalimumab (Humira®), currently approved for the treatment of rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA), ankylosing spondylitis (AS), and Crohn's disease (CD). The WT and Fc variant anti-CD25 antibodies are humanized versions of the antibody anti-TAC (Junghans et al., 1990, Cancer Research 50:1495-1502), referred to as H1.8/L1 anti-TAC. The WT and Fc variant anti-EGFR antibodies are humanized versions of the murine antibody C225, referred to as H4.42/L3.32 C225. Finally, the WT and Fc variant anti-IgE antibodies contain the variable region of the humanized antibody omalizumab (Xolair®), which is approved for the treatment of allergic asthma.

[199] WT and variant antibodies were constructed, expressed, and purified as described above. Antibodies were tested for binding to human FcRn at pH 6.0 by Biacore as described above. The relative binding data of the variant anti-TNF, -CD25, -EGFR, and -IgE antibodies to human FcRn

are provided in Figure 12. As can be seen, the variants improve FcRn affinity in the context of antibodies targeting a variety of antigens.

EXAMPLE 5. Pharmacokinetic experiments in human FcRn knock-in mice

[200] To test the ability of select variants to improve half-life in vivo, pharmacokinetic experiments were performed in B6 mice that are homozygous knock-outs for murine FcRn and heterozygous knock-ins of human FcRn (mFcRn^{-/-}, hFcRn^{+/+}) (Petkova et al., 2006, Int Immunol 18(12):1759-69, entirely incorporated by reference), herein referred to as hFcRn or hFcRn^{+/+} mice. A single, intravenous tail vein injection of anti-VEGF antibody (2 mg/kg) was given to groups of 4-7 female mice randomized by body weight (20-30g range). Blood (~50ul) was drawn from the orbital plexus at each time point, processed to serum, and stored at -80°C until analysis. Study durations were 28 or 49 days. Animals were not harmed during these studies.

[201] Antibody concentrations were determined using two ELISA assays. In the first two studies (referred to as Study 1 and Study 2), goat anti-human Fc antibody (Jackson Immuno research) was adhered to the plate, wells were washed with PBST (phosphate buffered saline with 0.05% Tween) and blocked with 3% BSA in PBST. Serum or calibration standards were then added, followed by PBST washing, addition of europium labeled anti-human IgG (Perkin Elmer), and further PBST washing. The time resolved fluorescence signal was collected. For Studies 3-5, serum concentration was detected using a similar ELISA, but recombinant VEGF (VEGF-165, PeproTech, Rocky Hill, NJ) was used as capture reagent and detection was carried out with biotinylated anti-human kappa antibody and europium-labeled streptavidin. PK parameters were determined for individual mice with a non-compartmental model using WinNonLin (Pharsight Inc, Mountain View CA). Nominal times and dose were used with uniform weighing of points. The time points used (lambda Z ranges) were from 4 days to the end of the study, although all time points were used for the faster clearing mutants, P257N and P257L.

[202] Five antibody PK studies in mFcRn^{-/-} hFcRn^{+/+} mice were carried out. Figure 13 shows serum concentration data for WT and variant IgG1 (Study 3) and IgG2 (Study 5) antibodies respectively. Fitted PK parameters from all in vivo PK studies carried out in mFcRn^{-/-} hFcRn^{+/+} mice are provided in Figure 14. PK data include half-life, which represents the beta phase that characterizes elimination of antibody from serum, C_{max}, which represents the maximal observed serum concentration, AUC, which represents the area under the concentration time curve, and clearance, which represents the clearance of antibody from serum. Also provided for each variant is the calculated fold improvement or reduction in half-life relative to the IgG1 or IgG2 parent antibody [Fold half-life = half-life(variant) / half-life (WT)].

[203] The data show that a number of the engineered Fc variant antibodies with enhanced FcRn affinity at pH 6.0 extend half-life in vivo. Figure 15a shows a plot of the in vivo half-life versus the fold FcRn binding for the IgG1 antibodies, with select variants labeled. Results from repeat experiments (circled in the figure) indicate that data from the in vivo model are reproducible. The

best single variants include 308F and 434S, the best double variants include 259I/308F, 308F/428L, 308F/434S, and 428L/434S, and the best triple variant is 259I/308F/428L. There is a general correlation between affinity for FcRn and in vivo half-life, but it is not completely predictive. Notably, variants 257L and 257N, which improved FcRn binding by 3.4 and 3.5 –fold respectively, reduced in vivo half-life by 0.6 and 0.3 respectively. The plot also highlights again the importance of the amino acid identity of substitution at a given position – whereas 308F/434S provided substantial half-life improvement, 308F/434M was barely better than WT IgG1.

[204] Figure 15b shows a plot of the in vivo half-life versus fold FcRn binding for the IgG2 variant antibodies with the variants labeled. When the IgG2 in vivo data were compared with the IgG1 in vivo data (Figure 15c), a surprising result was observed. The variants provided a substantially greater improvement to in vivo half-life in the context of an IgG2 Fc region than they do an IgG1 Fc region. The longest single variant and double variant half-lives from all antibodies in all 5 studies were 12.2 and 16.5, provided by 434S IgG2 and 428L/434S IgG2 respectively. The dramatic improvement in half-lives for the IgG2 variants relative to IgG1 were despite the fact that fold-improvements by the variants in IgG2 were comparable or even lower than they were in IgG1 (434S IgG1 fold = 3.8, 434S IgG2 fold = 4.9, 428L/434S IgG1 fold = 17.3, 428L/434S IgG2 fold = 14.8). Thus unexpectedly, the IgG2 antibody may be the best application for the Fc variants for improving in vivo half-life in mammals.

EXAMPLE 6. Variant immunoadhesins

[205] The Fc variants of the invention were also evaluated for their capacity to improve the half-life of immunoadhesins (also referred to as Fc fusions). Select Fc variants were engineered into the anti-TNF immunoadhesion etanercept (Enbrel®). Etanercept is a fusion of human TNF receptor 2 (TNF RII) and the Fc region of human IgG1, and is clinically approved for the treatment of rheumatoid arthritis, juvenile idiopathic arthritis, ankylosing spondylitis, psoriatic arthritis, and psoriasis. An IgG2 Fc region version of this Fc fusion was also constructed, and select Fc variants were constructed in this context as well. The amino acid sequences of the anti-TNF immunoadhesins characterized in the invention are provided in Figure 16. Genes were constructed using recursive PCR and subcloned into the pTT5 vector, and Fc variants were constructed using QuikChange® mutagenesis methods. Immunoadhesins were expressed in 293E cells and purified as described above.

[206] The binding specificity of the purified immunoadhesins was confirmed by testing binding to recombinant TNF by Biacore. Immunoadhesins were captured onto an immobilized Protein A/G (Pierce) CM5 biosensor chip (Biacore), generated using standard primary amine coupling. Immunoadhesins were immobilized on the Protein A/G surface, and recombinant TNF in serial dilutions was injected over antibody bound surface, followed by a dissociation phase. After each cycle, the surface was regenerated with buffer. Data were processed by zeroing time and response before the injection of receptor and by subtracting from a reference channel to account

for changes due to injections. Kinetic data were fit to a 1:1 binding model (Langmuir). Equilibrium association constants (K_a 's) obtained from these fits are provided in Figure 17. The results show that the variant immunoadhesins retained affinity for TNF, comparable to commercial enbrel.

[207] Variant immunoadhesins were tested for binding to human FcRn at pH 6.0 using Biacore as described above. The results (Figure 18) indicate that, similar as in the context of antibodies, the variants improve binding to FcRn relative to their IgG1 and IgG2 parent immunoadhesin proteins.

[208] The half-lives of the variant immunoadhesins were tested in the mFcRn^{-/-} hFcRn⁺ mice as described above. 12 mice per group were injected at 2 mg/kg of variant and parent IgG1 immunoadhesin. Serum concentration was detected using an ELISA similar to that described above, except that goat anti-human TNF RII antibody was used as capture reagent; detection was carried out with biotinylated anti-human kappa antibody and europium-labeled streptavidin. Figure 19 shows serum concentration data for WT IgG1 Fc and variant Fc immunoadhesins. Fitted PK parameters, as described above, from the PK study are provided in Figure 20. Also provided for each variant is the calculated % increase in half-life, calculated as 100 times the half-life of variant Fc fusion over that of the WT IgG1 Fc parent. The results indicate that the variants extend in vivo half-life in the context of the immunoadhesin.

EXAMPLE 7. Pharmacokinetic experiment in nonhuman primates

[209] The PK properties of biologics in non-human primates are well-established to be predictive of their properties in humans. A PK study was carried out in cynomolgus monkeys (*macaca fascicularis*) in order to evaluate the capacity of the variant anti-VEGF antibodies to improve serum half-life in non-human primates.

[210] In preparation for a PK study in cynomolgus monkeys, binding of the variant antibodies to cynomolgus (cyno) FcRn (cFcRn) at pH 6.0 was measured. cFcRn was constructed, expressed, and purified as described above for human FcRn. Binding of variant anti-VEGF antibodies to cFcRn was measured using Biacore as described above. The data are provided in Figure 21. The results show that the variants improve affinity for cyno FcRn similarly as they do for human FcRn. Dissociation at the higher pH (7.4) was also very rapid (data not shown), similar to as was observed for binding to human FcRn. These results are not surprising given the high sequence homology of the human and cyno receptors (FcRn alpha chain 96%, beta-2-microglobulin 91%).

[211] The PK of the variants were studied in vivo in non-human primates. Male cynomolgus monkeys (*macaca fascicularis*, also called crab-eating Macaque) weighing 2.3-5.1kg were randomized by weight and divided into 5 groups with 3 monkeys per group. The monkeys were given a single, 1 hour peripheral vein infusion of 4 mg/kg antibody. Blood samples (1 ml) were drawn from a separate vein from 5 minutes to 90 days after completion of the infusion, processed to serum and stored at -70°C. Animals were not harmed during these studies.

[212] Antibody concentrations were determined using the VEGF capture method as described above. PK parameters were determined by fitting the concentrations versus time to a non-compartmental model as was done in the mouse PK studies. However, time points from day 10 to day 90 were used for PK parameter determinations. The PK results are plotted in Figure 22, and the fitted parameters are provided in Figure 23. The results show that the variants enhanced the in vivo half-life of antibody up to 3.2-fold. In the best case (the 428L/434S variant) half-life was extended from 9.7 days to 31.1 days. The PK results obtained in cynomolgus monkeys are consistent with those obtained in mFcRn^{-/-} hFcRn⁺ mice, validating the hFcRn mouse model as a system for assessing the in vivo PK properties of the variants, and supporting the conclusions from those studies.

[213] Whereas particular embodiments of the invention have been described above for purposes of illustration, it will be appreciated by those skilled in the art that numerous variations of the details may be made without departing from the invention as described in the appended claims.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 52620-185 Seq 30-MAR-10 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110>	XENCOR, INC. LAZAR, GREGORY CHAMBERLAIN, AARON KEITH DAHIYAT, BASSIL I. DESJARLAIS, JOHN R. KARKI, SHER BAHADUR
<120>	FC VARIANTS WITH ALTERED BINDING TO FcRn
<130>	5026-WO03
<140>	PCT/US2008/088053
<141>	2008-12-22
<150>	61/016,793
<151>	2007-12-26
<150>	61/ 031,353
<151>	2008-02-25

<150> 61/046,353
 <151> 2008-04-18

<150> 61/050,172
 <151> 2008-05-02

<150> 61/079,779
 <151> 2008-07-10

<150> 61/099,178
 <151> 2008-09-22

<160> 35

<170> PatentIn version 3.5

<210> 1
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 1
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 1 5 10 15
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 20 25 30
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 35 40 45
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 50 55 60
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 65 70 75 80
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 85 90 95
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 100 105

<210> 2
 <211> 330
 <212> PRT
 <213> Homo sapiens

<400> 2
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160
 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175
 Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190
 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
 225 230 235 240
 Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320
 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325 330

<210> 3
 <211> 326
 <212> PRT
 <213> Homo sapiens

<400> 3
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15
 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr
 65 70 75 80
 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro
 100 105 110
 Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
 115 120 125
 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 130 135 140
 Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
 145 150 155 160
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn
 165 170 175
 Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
 180 185 190
 Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
 195 200 205
 Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
 210 215 220
 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
 225 230 235 240
 Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
 245 250 255

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
 260 265 270
 Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
 275 280 285
 Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
 290 295 300
 Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 305 310 315 320
 Ser Leu Ser Pro Gly Lys
 325

<210> 4
 <211> 377
 <212> PRT
 <213> Homo sapiens

<400> 4
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Thr Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Arg Val Glu Leu Lys Thr Pro Leu Gly Asp Thr Thr His Thr Cys Pro
 100 105 110
 Arg Cys Pro Glu Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg
 115 120 125
 Cys Pro Glu Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg Cys
 130 135 140
 Pro Glu Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg Cys Pro
 145 150 155 160
 Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
 165 170 175
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
 180 185 190
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Lys Trp Tyr
 195 200 205
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
 210 215 220
 Gln Tyr Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Leu His
 225 230 235 240
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 245 250 255
 Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln
 260 265 270
 Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
 275 280 285
 Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
 290 295 300
 Ser Asp Ile Ala Val Glu Trp Glu Ser Ser Gly Gln Pro Glu Asn Asn
 305 310 315 320
 Tyr Asn Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
 325 330 335
 Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Ile
 340 345 350

Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln
 355 360 365
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 370 375

<210> 5
 <211> 327
 <212> PRT
 <213> Homo sapiens

<400> 5
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15
 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
 65 70 75 80
 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro
 100 105 110
 Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125
 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 130 135 140
 Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
 145 150 155 160
 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
 165 170 175
 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 180 185 190
 Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
 195 200 205
 Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 210 215 220
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
 225 230 235 240
 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 245 250 255
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270
 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285
 Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320
 Leu Ser Leu Ser Leu Gly Lys
 325

<210> 6
 <211> 329
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> IgG1/2 constant heavy chain (CH1-hinge-CH2-CH3))

<400> 6

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
 115 120 125
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
 130 135 140
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
 145 150 155 160
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
 165 170 175
 Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His
 180 185 190
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 195 200 205
 Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln
 210 215 220
 Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
 225 230 235 240
 Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
 245 250 255
 Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
 260 265 270
 Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
 275 280 285
 Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
 290 295 300
 Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
 305 310 315 320
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325

<210> 7

<211> 330

<212> PRT

<213> Artificial Sequence

<220>

<223> IgG1 259I/308F

<400> 7

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80

```

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85      90      95
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
      100      105      110
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
      115      120      125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Ile Thr Cys
      130      135      140
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
      145      150      155      160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      165      170      175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Phe Leu
      180      185      190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      195      200      205
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
      210      215      220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
      225      230      235      240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      245      250      255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      260      265      270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
      275      280      285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      290      295      300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      305      310      315      320
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
      325      330

```

```

<210> 8
<211> 330
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> IgG1 434S/428L

```

```

<400> 8
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      20      25      30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      35      40      45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      50      55      60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
      65      70      75      80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85      90      95
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
      100      105      110
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
      115      120      125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      130      135      140
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
      145      150      155      160

```

```

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      165      170      175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
      180      185      190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      195      200      205
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
      210      215      220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
      225      230      235      240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      245      250      255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      260      265      270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
      275      280      285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      290      295      300
Val Phe Ser Cys Ser Val Leu His Glu Ala Leu His Ser His Tyr Thr
      305      310      315      320
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
      325      330

```

```

<210> 9
<211> 326
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> IgG2 434S

```

```

<400> 9
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1      5      10      15
Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      20      25      30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      35      40      45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      50      55      60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr
      65      70      75      80
Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85      90      95
Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro
      100      105      110
Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
      115      120      125
Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
      130      135      140
Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
      145      150      155      160
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn
      165      170      175
Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
      180      185      190
Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
      195      200      205
Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
      210      215      220
Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
      225      230      235      240

```

```

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
      245                      250                255
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
      260                      265                270
Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
      275                      280                285
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
      290                      295                300
Ser Val Met His Glu Ala Leu His Ser His Tyr Thr Gln Lys Ser Leu
      305                      310                315                320
Ser Leu Ser Pro Gly Lys
      325

```

```

<210> 10
<211> 326
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> IgG2 434S/428L

```

```

<400> 10
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1      5                      10                15
Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      20                      25                30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      35                      40                45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      50                      55                60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr
      65                      70                75                80
Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85                      90                95
Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro
      100                     105                110
Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
      115                     120                125
Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
      130                     135                140
Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
      145                     150                155                160
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn
      165                     170                175
Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
      180                     185                190
Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
      195                     200                205
Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
      210                     215                220
Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
      225                     230                235                240
Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
      245                     250                255
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
      260                     265                270
Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
      275                     280                285
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
      290                     295                300

```

Ser Val Leu His Glu Ala Leu His Ser His Tyr Thr Gln Lys Ser Leu
 305 310 315 320
 Ser Leu Ser Pro Gly Lys
 325

<210> 11
 <211> 329
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> IgG1/2 434S

<400> 11
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
 115 120 125
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
 130 135 140
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
 145 150 155 160
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
 165 170 175
 Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His
 180 185 190
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 195 200 205
 Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln
 210 215 220
 Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
 225 230 235 240
 Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
 245 250 255
 Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
 260 265 270
 Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
 275 280 285
 Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
 290 295 300
 Phe Ser Cys Ser Val Met His Glu Ala Leu His Ser His Tyr Thr Gln
 305 310 315 320
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325

<210> 12
 <211> 329
 <212> PRT
 <213> Artificial Sequence

<220>

<223> IgG1/2 434S/428L

<400> 12

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20      25      30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35      40      45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50      55      60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65      70      75      80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85      90      95
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100     105     110
Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
115     120     125
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
130     135     140
Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
145     150     155     160
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
165     170     175
Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His
180     185     190
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
195     200     205
Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln
210     215     220
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
225     230     235     240
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
245     250     255
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
260     265     270
Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
275     280     285
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
290     295     300
Phe Ser Cys Ser Val Leu His Glu Ala Leu His Ser His Tyr Thr Gln
305     310     315     320
Lys Ser Leu Ser Leu Ser Pro Gly Lys
325

```

<210> 13

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> Anti-VEGF VH

<400> 13

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1      5      10      15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
20      25      30
Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35      40      45

```

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 50 55 60
 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
 100 105 110
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 14
 <211> 107
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-VEGF VL

<400> 14
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
 20 25 30
 Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile
 35 40 45
 Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 15
 <211> 121
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF VH

<400> 15
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60
 Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 16
 <211> 107
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF VL

<400> 16
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ala Ala Ser Thr Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Val Ala Thr Tyr Tyr Cys Gln Arg Tyr Asn Arg Ala Pro Tyr
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 17
 <211> 116
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-CD25

<400> 17
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30
 Arg Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Trp Ile Asn Pro Ser Thr Gly Tyr Thr Glu Tyr Asn Gln Lys Phe
 50 55 60
 Gln Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Ile Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Gly Gly Val Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val
 100 105 110
 Thr Val Ser Ser
 115

<210> 18
 <211> 106
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-CD25 VL

<400> 18
 Gln Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Ser Ser Ile Ser Tyr Met
 20 25 30
 His Trp Phe Gln Gln Lys Pro Gly Gln Ser Pro Gln Leu Leu Ile Tyr
 35 40 45
 Thr Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60
 Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
 65 70 75 80
 Asp Phe Ala Val Tyr Tyr Cys His Gln Arg Ser Thr Tyr Pro Leu Thr
 85 90 95
 Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 19
 <211> 119
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-EGFR VH

<400> 19
 Gln Val Gln Leu Gln Gln Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Tyr
 20 25 30
 Gly Val His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met
 35 40 45
 Gly Ile Ile Trp Ser Gly Gly Ser Thr Asp Tyr Ser Thr Ser Leu Lys
 50 55 60
 Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val Val Leu
 65 70 75 80
 Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala
 85 90 95
 Arg Ala Leu Thr Tyr Tyr Asp Tyr Glu Phe Ala Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ser
 115

<210> 20
 <211> 107
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-EGFR VL

<400> 20
 Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Asn
 20 25 30
 Leu His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
 35 40 45
 Lys Tyr Ala Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala
 65 70 75 80
 Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Asn Asn Asn Trp Pro Thr
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 21
 <211> 121
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-IgE VH

<400> 21
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Tyr Ser Ile Thr Ser Gly
 20 25 30
 Tyr Ser Trp Asn Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
 35 40 45
 Val Ala Ser Ile Thr Tyr Asp Gly Ser Thr Asn Tyr Asn Pro Ser Val
 50 55 60
 Lys Gly Arg Ile Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr Phe Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Ser His Tyr Phe Gly His Trp His Phe Ala Val Trp Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 22
 <211> 111
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-IgE VL

<400> 22
 Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Val Asp Tyr Asp
 20 25 30
 Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 35 40 45
 Lys Leu Leu Ile Tyr Ala Ala Ser Tyr Leu Glu Ser Gly Val Pro Ser
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 65 70 75 80
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser His
 85 90 95
 Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105 110

<210> 23
 <211> 214
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF light chain

<400> 23
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

```

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Tyr
      20      25      30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
      35      40      45
Tyr Ala Ala Ser Thr Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
      50      55      60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
      65      70      75      80
Glu Asp Val Ala Thr Tyr Tyr Cys Gln Arg Tyr Asn Arg Ala Pro Tyr
      85      90      95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
      100      105      110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
      115      120      125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
      130      135      140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
      145      150      155      160
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
      165      170      175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
      180      185      190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
      195      200      205
Phe Asn Arg Gly Glu Cys
      210

```

<210> 24

<211> 451

<212> PRT

<213> Artificial Sequence

<220>

<223> Anti-TNF heavy chain IgG1 259I/308F

<400> 24

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1      5      10      15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
      20      25      30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
      35      40      45
Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
      50      55      60
Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
      65      70      75      80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
      85      90      95
Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
      100      105      110
Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
      115      120      125
Val Phe Pro Leu Ala Pro Ser Lys Ser Thr Ser Gly Gly Thr Ala
      130      135      140
Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
      145      150      155      160
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
      165      170      175
Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
      180      185      190
Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
      195      200      205

```

```

Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
210 215 220
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
225 230 235 240
Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
245 250 255
Ile Ser Arg Thr Pro Glu Ile Thr Cys Val Val Val Asp Val Ser His
260 265 270
Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
275 280 285
His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
290 295 300
Arg Val Val Ser Val Leu Thr Phe Leu His Gln Asp Trp Leu Asn Gly
305 310 315 320
Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
325 330 335
Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
340 345 350
Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser
355 360 365
Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
370 375 380
Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
385 390 395 400
Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
405 410 415
Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
420 425 430
His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
435 440 445
Pro Gly Lys
450

```

```

<210> 25
<211> 451
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Anti-TNF heavy chain IgG1 434S/428L

```

```

<400> 25
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
20 25 30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
50 55 60
Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
100 105 110
Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
115 120 125
Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
130 135 140
Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
145 150 155 160

```

```

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
      165      170      175
Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
      180      185      190
Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
      195      200      205
Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
      210      215      220
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
      225      230      235      240
Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
      245      250      255
Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
      260      265      270
Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
      275      280      285
His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
      290      295      300
Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
      305      310      315      320
Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
      325      330      335
Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
      340      345      350
Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser
      355      360      365
Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
      370      375      380
Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
      385      390      395      400
Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
      405      410      415
Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Leu
      420      425      430
His Glu Ala Leu His Ser His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
      435      440      445
Pro Gly Lys
      450

```

<210> 26

<211> 447

<212> PRT

<213> Artificial Sequence

<220>

<223> Anti-TNF heavy chain IgG2 434S

<400> 26

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1      5      10      15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
      20      25      30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
      35      40      45
Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
      50      55      60
Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
      65      70      75      80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
      85      90      95
Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
      100      105      110

```

Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
 115 120 125
 Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala
 130 135 140
 Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
 145 150 155 160
 Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
 165 170 175
 Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
 180 185 190
 Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His
 195 200 205
 Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys
 210 215 220
 Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 260 265 270
 Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Ser His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435 440 445

<210> 27
 <211> 447
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF heavy chain IgG2 434S/428L

<400> 27
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60
 Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

```

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
      85          90          95
Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
      100        105        110
Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
      115        120        125
Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala
      130        135        140
Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
      145        150        155        160
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
      165        170        175
Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
      180        185        190
Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His
      195        200        205
Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys
      210        215        220
Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val
      225        230        235        240
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
      245        250        255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
      260        265        270
Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
      275        280        285
Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser
      290        295        300
Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
      305        310        315        320
Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile
      325        330        335
Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
      340        345        350
Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
      355        360        365
Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
      370        375        380
Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser
      385        390        395        400
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
      405        410        415
Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Leu His Glu Ala Leu
      420        425        430
His Ser His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
      435        440        445

```

```

<210> 28
<211> 450
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Anti-TNF heavy chain IgG1/2 434S

```

```

<400> 28
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1      5      10      15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
20     25     30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35     40     45

```

Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60
 Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
 115 120 125
 Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
 130 135 140
 Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
 145 150 155 160
 Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
 165 170 175
 Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
 180 185 190
 Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
 195 200 205
 Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
 210 215 220
 Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
 225 230 235 240
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 245 250 255
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
 Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
 290 295 300
 Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
 305 310 315 320
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
 325 330 335
 Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 340 345 350
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 355 360 365
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met
 385 390 395 400
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 420 425 430
 Glu Ala Leu His Ser His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 435 440 445
 Gly Lys
 450

<210> 29

<211> 450

<212> PRT

<213> Artificial Sequence

<220>

<223> Anti-TNF heavy chain IgG1/2 434S/428L

<400> 29
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60
 Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
 115 120 125
 Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
 130 135 140
 Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
 145 150 155 160
 Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
 165 170 175
 Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
 180 185 190
 Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
 195 200 205
 Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
 210 215 220
 Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
 225 230 235 240
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 245 250 255
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
 Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
 290 295 300
 Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
 305 310 315 320
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
 325 330 335
 Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 340 345 350
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 355 360 365
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met
 385 390 395 400
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Leu His
 420 425 430
 Glu Ala Leu His Ser His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 435 440 445
 Gly Lys
 450

<210> 30

<211> 489

<212> PRT

<213> Artificial Sequence

<220>

<223> Anti-TNF Fc fusion with IgG1 Fc

<400> 30

```

Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
1      5      10      15
Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
      20      25      30
Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
      35      40      45
Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
      50      55      60
Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
      65      70      75      80
Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
      85      90      95
Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
      100     105     110
Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
      115     120     125
Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
      130     135     140
Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
      145     150     155     160
Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
      165     170     175
Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
      180     185     190
Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
      195     200     205
Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
      210     215     220
Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
      225     230     235     240
Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
      245     250     255
Asp Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
      260     265     270
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
      275     280     285
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
      290     295     300
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
      305     310     315     320
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
      325     330     335
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
      340     345     350
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
      355     360     365
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
      370     375     380
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
      385     390     395     400
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
      405     410     415
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
      420     425     430
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
      435     440     445

```

Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
 450 455 460
 Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
 465 470 475 480
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 485

<210> 31
 <211> 488
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF Fc fusion with IgG2 Fc

<400> 31
 Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15
 Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30
 Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45
 Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
 50 55 60
 Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
 65 70 75 80
 Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
 85 90 95
 Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
 100 105 110
 Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
 115 120 125
 Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
 130 135 140
 Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
 145 150 155 160
 Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
 165 170 175
 Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
 180 185 190
 Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
 195 200 205
 Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
 210 215 220
 Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
 225 230 235 240
 Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
 245 250 255
 Asp Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
 260 265 270
 Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
 275 280 285
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
 290 295 300
 Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val
 305 310 315 320
 Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
 325 330 335
 Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln
 340 345 350
 Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly
 355 360 365

```

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro
 370          375          380
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
385          390          395          400
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
          405          410          415
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
          420          425          430
Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
          435          440          445
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
          450          455          460
Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
465          470          475          480
Ser Leu Ser Leu Ser Pro Gly Lys
          485

```

```

<210> 32
<211> 489
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Anti-TNF Fc fusion with IgG1 259I/308F Fc

```

```

<400> 32
Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
1          5          10          15
Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
          20          25          30
Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
          35          40          45
Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
          50          55          60
Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
65          70          75          80
Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
          85          90          95
Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
          100          105          110
Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
          115          120          125
Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
          130          135          140
Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
145          150          155          160
Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
          165          170          175
Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
          180          185          190
Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
          195          200          205
Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
          210          215          220
Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
225          230          235          240
Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
          245          250          255
Asp Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
          260          265          270
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
          275          280          285

```

```

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Ile Thr Cys Val
290                295                300
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
305                310                315                320
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
325                330                335
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Phe Leu His
340                345                350
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
355                360                365
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
370                375                380
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
385                390                395                400
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
405                410                415
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
420                425                430
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
435                440                445
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
450                455                460
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
465                470                475                480
Lys Ser Leu Ser Leu Ser Pro Gly Lys
485

```

```

<210> 33
<211> 489
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Anti-TNF Fc fusion with IgG1 428L/434S Fc

```

```

<400> 33
Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
1                5                10                15
Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
20                25                30
Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
35                40                45
Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
50                55                60
Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
65                70                75                80
Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
85                90                95
Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
100               105               110
Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
115               120               125
Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
130               135               140
Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
145               150               155               160
Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
165               170               175
Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
180               185               190
Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
195               200               205

```

Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
 210 215 220
 Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
 225 230 235 240
 Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
 245 250 255
 Asp Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
 260 265 270
 Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
 275 280 285
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
 290 295 300
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
 305 310 315 320
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
 325 330 335
 Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
 340 345 350
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 355 360 365
 Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
 370 375 380
 Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
 385 390 395 400
 Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
 405 410 415
 Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
 420 425 430
 Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
 435 440 445
 Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
 450 455 460
 Phe Ser Cys Ser Val Leu His Glu Ala Leu His Ser His Tyr Thr Gln
 465 470 475 480
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 485

<210> 34
 <211> 488
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF Fc fusion with IgG2 434S Fc

<400> 34
 Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15
 Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30
 Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45
 Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
 50 55 60
 Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
 65 70 75 80
 Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
 85 90 95
 Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
 100 105 110
 Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
 115 120 125

```

Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
130          135          140
Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
145          150          155          160
Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
          165          170          175
Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
          180          185          190
Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
          195          200          205
Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
          210          215          220
Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
225          230          235          240
Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
          245          250          255
Asp Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
          260          265          270
Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
          275          280          285
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
          290          295          300
Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val
305          310          315          320
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
          325          330          335
Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln
          340          345          350
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly
          355          360          365
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro
          370          375          380
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
385          390          395          400
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
          405          410          415
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
          420          425          430
Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
          435          440          445
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
          450          455          460
Ser Cys Ser Val Met His Glu Ala Leu His Ser His Tyr Thr Gln Lys
465          470          475          480
Ser Leu Ser Leu Ser Pro Gly Lys
          485

```

<210> 35
 <211> 488
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Anti-TNF Fc fusion with IgG2 428L/434S Fc

<400> 35
 Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15
 Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30
 Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45

```

Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
50 55 60
Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
65 70 75 80
Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
85 90 95
Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
100 105 110
Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
115 120 125
Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
130 135 140
Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
145 150 155 160
Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
165 170 175
Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
180 185 190
Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
195 200 205
Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
210 215 220
Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
225 230 235 240
Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
245 250 255
Asp Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
260 265 270
Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
275 280 285
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
290 295 300
Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val
305 310 315 320
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
325 330 335
Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln
340 345 350
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly
355 360 365
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro
370 375 380
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
385 390 395 400
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
405 410 415
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
420 425 430
Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
435 440 445
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
450 455 460
Ser Cys Ser Val Leu His Glu Ala Leu His Ser His Tyr Thr Gln Lys
465 470 475 480
Ser Leu Ser Leu Ser Pro Gly Lys
485

```

CLAIMS:

1. A polypeptide comprising an Fc variant of a human IgG Fc polypeptide, wherein said Fc variant comprises a serine at position 434 and an isoleucine or a valine at position 311, wherein numbering is according to the EU index.
2. The polypeptide according to claim 1, wherein said polypeptide comprising an Fc variant has specificity for a target molecule that is selected from the group consisting of a cytokine, a soluble protein factor, and a protein expressed on cancer cells.
3. The polypeptide according to claim 1 or 2, wherein said polypeptide is an antibody.
4. The polypeptide according to claim 3, wherein said antibody is selected from the group consisting of a chimeric antibody, a humanized antibody, or a human antibody.
5. The polypeptide according to any one of claims 1-4, wherein said polypeptide is an Fc fusion.
6. A method of producing the polypeptide according to any one of claims 1-5, said method comprising providing a cell comprising a nucleic acid encoding said polypeptide, wherein said nucleic acid is operably linked with control or regulatory sequences for expression of said polypeptide, and wherein said cell is cultured under conditions suitable for expression of said polypeptide.
7. The method according to claim 6, wherein said nucleic acid is part of an expression vector.
8. A host cell comprising a nucleic acid encoding the polypeptide according to any one of claims 1-5.
9. An expression vector, wherein said expression vector encodes the polypeptide according to any one of claims 1-5.

Figure 1

CH1																							
EU Index	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138		
IgG1	A	S	T	K	G	P	S	V	F	P	L	A	P	S	S	K	S	T	S	G	G		
IgG2	A	S	T	K	G	P	S	V	F	P	L	A	P	C	S	R	S	T	S	E	S		
IgG3	A	S	T	K	G	P	S	V	F	P	L	A	P	C	S	R	S	T	S	G	G		
IgG4	A	S	T	K	G	P	S	V	F	P	L	A	P	C	S	R	S	T	S	E	S		
EU Index	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159		
IgG1	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N		
IgG2	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N		
IgG3	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N		
IgG4	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N		
EU Index	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180		
IgG1	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y		
IgG2	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y		
IgG3	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y		
IgG4	S	G	A	L	T	S	G	V	H	T	F	P	A	V	L	Q	S	S	G	L	Y		
EU Index	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201		
IgG1	S	L	S	S	V	V	T	V	P	S	S	S	L	G	T	Q	T	Y	I	C	N		
IgG2	S	L	S	S	V	V	T	V	P	S	S	N	F	G	T	Q	T	Y	T	C	N		
IgG3	S	L	S	S	V	V	T	V	P	S	S	S	L	G	T	Q	T	Y	T	C	N		
IgG4	S	L	S	S	V	V	T	V	P	S	S	S	L	G	T	K	T	Y	T	C	N		
EU Index	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220				
IgG1	V	N	H	K	P	S	N	T	K	V	D	K	K/R	V	E	P	K	S	C				
IgG2	V	D	H	K	P	S	N	T	K	V	D	K	T	V	E	R	K	C	C				
IgG3	V	N	H	K	P	S	N	T	K	V	D	K	R	V	E	L	K	T	P				
IgG4	V	D	H	K	P	S	N	T	K	V	D	K	R	V	E	S	K	Y	Q				
Hinge																			Fc		>		
EU Index	221				222	223	224	225	226	227	228												
IgG1	D				K	T	H	T	C	P	P												
IgG2					V		E		C	P	P												
IgG3	L	G	D	T	T	H	T	C	P	R	C	P	E	P	K	S	C	D	T	P	P		
IgG4							P	P	C	P	S												
EU Index																							
IgG1																							
IgG2																							
IgG3	P	C	P	R	C	P	E	P	K	S	C	D	T	P	P	P	C	P	R	C	P		
IgG4																							
EU Index													229	230	231	232	233	234	235	236			
IgG1													C	P	A	P	E	L	L	G			
IgG2													C	P	A	P	P	V	A				
IgG3	E	P	K	S	C	D	T	P	P	P	C	P	R	C	P	A	P	E	L	L	G		
IgG4													C	P	A	P	E	F	L	G			

Figure 2

Kappa constant light chain (Cκ) (SEQ ID NO:1)

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNFPYPREAKVQWKVDNALQSGNSQESVTEQDSKDS
 TYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

IgG1 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:2)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPK
 KDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ
 DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI
 AVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKS
 LSLSPGK

IgG2 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:3)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSNFGTQTYTCNVDHKPSNTKVDKVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTL
 MISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVHQQDWLN
 GKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE
 WESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSL
 SPGK

IgG3 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:4)

ASTKGPSVFPLAPCSRSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYTCNVNHKPSNTKVDKRVELKTPLGDTTHTCPRCPEPKSCDTPPPCPRC
 PEKSCDTPPPCPRCPEPKSCDTPPPCPRCAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVD
 VSHEDPEVQFNWYVDGVEVHNAKTKPREEQYNSTFRVSVLTVLHQDWLNGKEYKCKVSNKA
 LPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYN
 TTPPMLDSDGSFFLYSKLTVDKSRWQQGNIFCFSVMHEALHNRFQKSLSLSPGK

IgG4 constant heavy chain (CH1-hinge-CH2-CH3) (SEQ ID NO:5)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYTCNVNHKPSNTKVDKRVESKYGPPCPCPAPEFLGGPSVFLFPPKPKDT
 LMISRTPEVTCVVDVSDQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWL
 NGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVE
 WESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFCFSVMHEALHNHYTQKSLSL
 SLGK

IgG1/2 constant heavy chain (CH1-hinge-CH2-CH3)) (SEQ ID NO:6)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPK
 DTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVHQQD
 WLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA
 VEWESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKS
 LSLSPGK

Figure 3

IgG1 259I/308F (SEQ ID NO:7)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKP
 KDTLMISRTPEITCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTFHQQD
 WLNKGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA
 VEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHLEALHNHYTQKSL
 SLSPGK

IgG1 434S/428L (SEQ ID NO:8)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKP
 KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
 DWLNKGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI
 AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSL
 SLSPGK

IgG2 434S (SEQ ID NO:9)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPPCAPPVAGPSVFLFPPKPKDTL
 MISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLN
 GKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE
 WESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSL
 SPGK

IgG2 434S/428L (SEQ ID NO:10)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPPCAPPVAGPSVFLFPPKPKDTL
 MISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLN
 GKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE
 WESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSL
 SPGK

IgG1/2 434S (SEQ ID NO:11)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPK
 DTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQD
 WLNKGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA
 VEWESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSL
 SLSPGK

IgG1/2 434S/428L (SEQ ID NO:12)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPK
 DTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQD
 WLNKGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA
 VEWESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSL
 SLSPGK

Figure 4

Anti-VEGF VH (SEQ ID NO:13)

EVQLVESGGGLVQPGGSLRLSCAASGYTFTNYGMNWVRQAPGKGLEWVGWINTYTGEPTYAA
DFKRRFTFSLDTSKSTAYLQMNSLRAEDTAVYYCAKYPHYYGSSHWFYFDVWGQGTLVTVSS

Anti-VEGF VL (SEQ ID NO:14)

DIQMTQSPSSLSASVGDRVTITCSASQDISNYLNWYQQKPGKAPKVLIIYFTSSLHSGVPSRFSGS
GSGTDFTLTISLQPEDFATYYCQYSTVPWTFGQGTKVEIK

Anti-TNF VH (SEQ ID NO:15)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTLVTVSS

Anti-TNF VL (SEQ ID NO:16)

DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGS
GSGTDFTLTISLQPEDVATYYCQRYNRAPYTFGQGTKVEIK

Anti-CD25 VH (SEQ ID NO:17)

QVQLVQSGAEVKKPGASVKVSKASGYTFTSYRMHWVRQAPGQGLEWMGWINPSTGYTEYNQ
KFQGRVTITADKSISTAYMELSRLRSDDTAVYYCARGGGVFDYWGQGTLVTVSS

Anti-CD25 VL (SEQ ID NO:18)

QIVLTQSPATLSLSPGERATLSCRASSISYMHWFQQKPGQSPQLLIYTTNLSGVPARFSGSG
SGTDYTLTISLQPEDFAVYYCHQRSTYPLTFGSGTKLEIK

Anti-EGFR VH (SEQ ID NO:19)

QVQLQQSGPGLVKPSQTLSTCTVSGFSLSNYGVHWVRQAPGKGLEWMGIWSSGSTDYSTSL
KSRLTISKDTSKSQVLTMTNMDPVDATYYCARALTYDYEFAYWGQGTLVTVSS

Anti-EGFR VL (SEQ ID NO:20)

DIQLTQSPSSLSASVGDRVTITCRASQSISSNLHWYQQKPDQSPKLLIKYASESISGVPSRFSGSG
SGTDFTLTISLQAEDVAVYYCQQNNNWPTTFGQGTKLEIK

Anti-IgE VH (SEQ ID NO:21)

EVQLVESGGGLVQPGGSLRLSCAVSGYSITSGYSWNWVRQAPGKGLEWVASITYDGSTNYNPSV
KGRITISRDDSKNTFYLMNSLRAEDTAVYYCARGSHYFGHWHFAVWGQGTLVTVSS

Anti-IgE VL (SEQ ID NO:22)

DIQLTQSPSSLSASVGDRVTITCRASQSDYDGD SYMNWYQQKPGKAPKLLIYAASYLESVPSR
FSGSGSGTDFTLTISLQPEDFATYYCQQSHEDPYTFGQGTKVEIK

Figure 5

Anti-TNF light chain (SEQ ID NO:23)

DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGS
 GSGTDFLTITSLQPEDVATYYCQRYNRAPYTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS
 VVCLLNNFYPRKAVQWVKVDNALQSGNSQESVTEQDSKSTYSLSSTLTLSKADYEKHKVYACE
 VTHQGLSPVTKSFNRGEC

Anti-TNF heavy chain IgG1 259I/308F (SEQ ID NO:24)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
 VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGLTVTVSSASTKGPS
 VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
 SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISR
 TPEITCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTLHQQDWLNGKEY
 KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESN
 GQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK

Anti-TNF heavy chain IgG1 434S/428L (SEQ ID NO:25)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
 VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGLTVTVSSASTKGPS
 VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
 SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISR
 TPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQQDWLNGKEY
 YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESN
 GQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK

Anti-TNF heavy chain IgG2 434S (SEQ ID NO:26)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
 VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGLTVTVSSASTKGPS
 VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
 SSNFGTQTYTCNVNHHKPSNTKVDKTKVERKCCVECPAPPVAGPSVFLFPPKPKDTLMISRTPE
 VTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQQDWLNGKEYKC
 KVSNGKLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ
 PENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMEALHSHYTQKSLSLSPGK

Anti-TNF heavy chain IgG2 434S/428L (SEQ ID NO:27)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
 VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGLTVTVSSASTKGPS
 VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
 SSNFGTQTYTCNVNHHKPSNTKVDKTKVERKCCVECPAPPVAGPSVFLFPPKPKDTLMISRTPE
 VTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQQDWLNGKEYKC
 KVSNGKLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ
 PENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK

Anti-TNF heavy chain IgG1/2 434S (SEQ ID NO:28)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
 VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGLTVTVSSASTKGPS
 VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
 SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRT
 PEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQQDWLNGKEY
 KCKVSNKALPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESN
 GQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMEALHSHYTQKSLSLSPGK

Figure 5 (continued)

Anti-TNF heavy chain IgG1/2 434S/428L (SEQ ID NO:29)

EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWWSAITWNSGHIDYADS
 VEGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTTLTVSSASTKGPS
 VFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
 SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRT
 PEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEY
 KCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESN
 GQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK

Figure 6




Figure 7




Figure 8




Figure 9

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG1	Anti-VEGF		1.00	
T250I	IgG1	Anti-VEGF	1	1.31	
M252Q	IgG1	Anti-VEGF	5	2.36	3.07
M252Y	IgG1	Anti-VEGF	2	4.82	0.62
M252E	IgG1	Anti-VEGF	1	NB	
M252P	IgG1	Anti-VEGF	1	0.03	
M252R	IgG1	Anti-VEGF	1	0.02	
M252D	IgG1	Anti-VEGF	1	NB	
M252K	IgG1	Anti-VEGF	1	0.64	
R255Y	IgG1	Anti-VEGF	1	1.24	
P257L	IgG1	Anti-VEGF	1	3.39	
P257N	IgG1	Anti-VEGF	1	3.53	
V259I	IgG1	Anti-VEGF	2	1.14	0.19
V259F	IgG1	Anti-VEGF	1	1.44	
V259G	IgG1	Anti-VEGF		ND	
V259H	IgG1	Anti-VEGF		ND	
V259M	IgG1	Anti-VEGF	1	0.68	
V259N	IgG1	Anti-VEGF	1	1.87	
V259S	IgG1	Anti-VEGF		ND	
V259T	IgG1	Anti-VEGF		ND	
V259W	IgG1	Anti-VEGF		ND	
V259Y	IgG1	Anti-VEGF		ND	
T307Q	IgG1	Anti-VEGF	1	3.41	
T307N	IgG1	Anti-VEGF	1	0.83	
T307G	IgG1	Anti-VEGF	1	1.72	
V308F	IgG1	Anti-VEGF	6	4.96	1.06
Y319L	IgG1	Anti-VEGF	1	1.43	
Y319I	IgG1	Anti-VEGF	2	1.48	0.53
Y319A	IgG1	Anti-VEGF	1	2.66	
Y319G	IgG1	Anti-VEGF		ND	
Y319H	IgG1	Anti-VEGF	1	1.81	
Y319M	IgG1	Anti-VEGF	1	1.95	
Y319N	IgG1	Anti-VEGF		ND	
Y319Q	IgG1	Anti-VEGF		ND	
Y319S	IgG1	Anti-VEGF		ND	
Y319T	IgG1	Anti-VEGF		ND	
Y319V	IgG1	Anti-VEGF	1	2.03	

Figure 9 (continued)

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
Y319W	IgG1	Anti-VEGF	1	1.51	
E380A	IgG1	Anti-VEGF		ND	
E380Q	IgG1	Anti-VEGF	1	2.01	
E380R	IgG1	Anti-VEGF	1	3.41	
E380S	IgG1	Anti-VEGF	1	2.55	
E380V	IgG1	Anti-VEGF		ND	
M428L	IgG1	Anti-VEGF	3	3.00	0.37
N434A	IgG1	Anti-VEGF	4	2.15	0.54
N434S	IgG1	Anti-VEGF	6	3.77	0.79
N434M	IgG1	Anti-VEGF	3	4.83	1.23
N434T	IgG1	Anti-VEGF	1	3.32	
N434V	IgG1	Anti-VEGF		ND	
N434Y	IgG1	Anti-VEGF		ND	
P245G/V308F	IgG1	Anti-VEGF	1	2.89	
T250I/V308F	IgG1	Anti-VEGF	2	6.75	0.10
T250Q/M428L	IgG1	Anti-VEGF	3	6.30	0.82
M252Y/V259I	IgG1	Anti-VEGF	1	3.56	
M252Q/V259I	IgG1	Anti-VEGF	1	0.14	
M252Q/T307Q	IgG1	Anti-VEGF	1	0.43	
M252Q/V308F	IgG1	Anti-VEGF	1	1.04	
M252Y/V308F	IgG1	Anti-VEGF	2	9.63	0.32
M252Q/Y319I	IgG1	Anti-VEGF	1	0.21	
M252Y/Y319I	IgG1	Anti-VEGF	1	2.37	
M252Q/M428L	IgG1	Anti-VEGF	1	0.35	
M252Y/M428L	IgG1	Anti-VEGF	1	5.32	
M252Y/N434A	IgG1	Anti-VEGF	1	11.94	
M252Y/N434S	IgG1	Anti-VEGF	2	11.46	2.73
M252Q/N434M	IgG1	Anti-VEGF	1	0.20	
M252Q/N434S	IgG1	Anti-VEGF	1	0.48	
M252Y/N434M	IgG1	Anti-VEGF	1	2.79	
M252Y/N434Y	IgG1	Anti-VEGF	1	61.60	
S254T/V308F	IgG1	Anti-VEGF	1	3.56	
T256V/V308F	IgG1	Anti-VEGF	2	5.04	1.94
T256E/N434Y	IgG1	Anti-VEGF	1	35.53	
V259I/V308F	IgG1	Anti-VEGF	12	9.79	2.72
V259A/V308F	IgG1	Anti-VEGF	1	4.81	
V259L/V308F	IgG1	Anti-VEGF		ND	

Figure 9 (continued)

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
V259I/Y319I	IgG1	Anti-VEGF	1	2.00	
V259I/M428L	IgG1	Anti-VEGF	1	4.53	
V259I/N434S	IgG1	Anti-VEGF	1	5.67	
V259I/N434M	IgG1	Anti-VEGF	1	0.87	
V259I/N434Y	IgG1	Anti-VEGF	1	24.81	
L306I/V308F	IgG1	Anti-VEGF	1	7.92	
T307P/V308F	IgG1	Anti-VEGF	1	3.17	
T307Q/V308F	IgG1	Anti-VEGF	2	5.57	0.69
T307S/V308F	IgG1	Anti-VEGF	1	5.22	
T307Q/M428L	IgG1	Anti-VEGF	1	7.25	
T307Q/N434S	IgG1	Anti-VEGF	1	14.18	
T307Q/N434M	IgG1	Anti-VEGF	1	1.99	
T307Q/N434Y	IgG1	Anti-VEGF	1	50.43	
V308F/L309Y	IgG1	Anti-VEGF	1	5.37	
V308F/L309N	IgG1	Anti-VEGF	1	1.00	
V308F/Q311P	IgG1	Anti-VEGF		ND	
V308F/Y319L	IgG1	Anti-VEGF	2	5.84	0.30
V308F/Y319F	IgG1	Anti-VEGF	1	3.10	
V308F/M428L	IgG1	Anti-VEGF	4	13.71	2.13
V308F/N434M	IgG1	Anti-VEGF	2	4.20	0.78
V308F/N434S	IgG1	Anti-VEGF	2	14.63	0.28
V308F/N434Y	IgG1	Anti-VEGF	1	41.55	
Q311A/N434Y	IgG1	Anti-VEGF	1	27.42	
Y319I/V308F	IgG1	Anti-VEGF	3	4.93	0.82
Y319I/M428L	IgG1	Anti-VEGF	1	4.70	
Y319I/N434M	IgG1	Anti-VEGF	1	4.22	
Y319I/N434S	IgG1	Anti-VEGF	1	4.42	
Y319I/N434Y	IgG1	Anti-VEGF		ND	
M428L/N434A	IgG1	Anti-VEGF	1	12.74	
M428L/N434S	IgG1	Anti-VEGF	14	17.28	6.70
M428L/N434M	IgG1	Anti-VEGF	1	1.88	
M428L/N434Y	IgG1	Anti-VEGF	1	39.54	
N434S/Y436F	IgG1	Anti-VEGF	1	4.43	
N434S/Y436M	IgG1	Anti-VEGF	1	2.58	
T250I/V259I/V308F	IgG1	Anti-VEGF	1	11.29	
T250Q/V308F/M428L	IgG1	Anti-VEGF	1	17.08	
T250I/V308F/N434S	IgG1	Anti-VEGF	1	12.26	

Figure 9 (continued)

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
M252Y/S254T/T256E	IgG1	Anti-VEGF	10	11.37	2.90
M252Q/V259I/N434S	IgG1	Anti-VEGF	1	0.65	
M252Y/V308F/M428L	IgG1	Anti-VEGF	1	10.62	
M252Q/V308F/N434S	IgG1	Anti-VEGF	1	2.37	
M252Y/V308F/N434M	IgG1	Anti-VEGF	1	11.09	
M252Y/V308F/N434S	IgG1	Anti-VEGF	1	12.13	
M252Q/M428L/N434S	IgG1	Anti-VEGF	1	1.90	
V259I/T307Q/V308F	IgG1	Anti-VEGF	2	7.40	1.54
V259I/V308F/Y319L	IgG1	Anti-VEGF	1	3.53	
V259I/V308F/Y319I	IgG1	Anti-VEGF	1	4.10	
V259I/V308F/M428L	IgG1	Anti-VEGF	13	19.94	10.46
V259I/V308F/N434S	IgG1	Anti-VEGF	3	10.86	3.38
V259I/V308F/N434M	IgG1	Anti-VEGF	1	7.06	
V259I/Y319I/N434S	IgG1	Anti-VEGF	1	7.90	
V259I/M428L/N434S	IgG1	Anti-VEGF	1	9.57	
V259I/M428L/N434S	IgG1	Anti-VEGF	1	11.91	
T307Q/V308F/Y319L	IgG1	Anti-VEGF	1	4.24	
T307Q/V308F/N434S	IgG1	Anti-VEGF	1	4.04	
V308F/Y319I/M428L	IgG1	Anti-VEGF	1	10.20	
V308F/Y319L/M428L	IgG1	Anti-VEGF	1	NF	
V308F/Y319L/N434S	IgG1	Anti-VEGF	2	12.39	5.63
V308F/Y319I/N434M	IgG1	Anti-VEGF	1	3.78	
V308F/Y319I/N434S	IgG1	Anti-VEGF	1	NF	
V308F/M428L/N434S	IgG1	Anti-VEGF	5	16.68	4.89
V308F/M428L/N434M	IgG1	Anti-VEGF	1	8.14	
Y319I/M428L/N434S	IgG1	Anti-VEGF	1	13.90	
M252Y/S254T/T256E/N434A	IgG1	Anti-VEGF	1	11.37	
M252Y/S254T/T256E/N434S	IgG1	Anti-VEGF	2	10.29	2.99
M252Y/S254T/T256E/V308F	IgG1	Anti-VEGF	2	9.84	4.04
M252Y/S254T/T256E/M428L	IgG1	Anti-VEGF	2	11.43	6.01
M252Y/S254T/T256E/T307Q	IgG1	Anti-VEGF	1	13.50	
M252Y/S254T/T256E/N434M	IgG1	Anti-VEGF	1	3.62	
V259I/V308F/M428L/N434S	IgG1	Anti-VEGF	1	24.22	
V259I/V308F/M428L/N434S	IgG1	Anti-VEGF	2	14.23	0.75
V308F/Y319I/M428L/N434S	IgG1	Anti-VEGF	1	10.37	
M252Y/S254T/T256E/M428L/N434S	IgG1	Anti-VEGF	1	7.21	

Figure 10

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG2	Anti-VEGF		1.00	
V259I	IgG2	Anti-VEGF	1	1.02	
V308F	IgG2	Anti-VEGF	2	3.66	2.07
M428L	IgG2	Anti-VEGF	1	4.19	
N434A	IgG2	Anti-VEGF	1	3.83	
N434S	IgG2	Anti-VEGF	2	4.88	1.05
N434G	IgG2	Anti-VEGF	1	1.60	
N434H	IgG2	Anti-VEGF	1	1.45	
N434M	IgG2	Anti-VEGF	1	1.51	
N434T	IgG2	Anti-VEGF	1	1.22	
N434Y	IgG2	Anti-VEGF	1	21.39	
N434D	IgG2	Anti-VEGF	1	NB	
N434E	IgG2	Anti-VEGF	1	NB	
N434F	IgG2	Anti-VEGF	1	16.79	
N434K	IgG2	Anti-VEGF	1	NB	
N434P	IgG2	Anti-VEGF	1	0.26	
N434R	IgG2	Anti-VEGF	1	0.44	
N434V	IgG2	Anti-VEGF	1	0.48	
N434W	IgG2	Anti-VEGF	1	39.81	
T250Q/M428L	IgG2	Anti-VEGF	1	7.62	
V259I/V308F	IgG2	Anti-VEGF	1	6.58	
V308F/M428L	IgG2	Anti-VEGF	1	16.13	
V308F/N434S	IgG2	Anti-VEGF	1	16.36	
M428L/N434S	IgG2	Anti-VEGF	1	14.80	
M252Y/S254T/T256E	IgG2	Anti-VEGF	1	8.34	
V259I/V308F/M428L	IgG2	Anti-VEGF	1	15.29	

Figure 10 (continued)

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG1/2	Anti-VEGF		1.00	
N434S	IgG1/2	Anti-VEGF	1	5.28	
L251A/N434S	IgG1/2	Anti-VEGF	1	0.65	
L251F/N434S	IgG1/2	Anti-VEGF	1	1.95	
L251I/N434S	IgG1/2	Anti-VEGF	1	1.28	
L251M/N434S	IgG1/2	Anti-VEGF	1	2.83	
L251N/N434S	IgG1/2	Anti-VEGF	1	1.34	
L251V/N434S	IgG1/2	Anti-VEGF	1	2.31	
R255H/N434S	IgG1/2	Anti-VEGF	1	2.23	
R255Q/N434S	IgG1/2	Anti-VEGF	1	1.53	
T307I/N434S	IgG1/2	Anti-VEGF	1	4.09	
T307V/N434S	IgG1/2	Anti-VEGF	1	5.25	
Q311A/N434S	IgG1/2	Anti-VEGF	1	4.35	
Q311D/N434S	IgG1/2	Anti-VEGF	1	1.39	
Q311E/N434S	IgG1/2	Anti-VEGF	1	1.94	
Q311F/N434S	IgG1/2	Anti-VEGF	1	2.95	
Q311G/N434S	IgG1/2	Anti-VEGF	1	1.26	
Q311I/N434S	IgG1/2	Anti-VEGF	1	15.81	
Q311K/N434S	IgG1/2	Anti-VEGF	1	4.57	
Q311L/N434S	IgG1/2	Anti-VEGF	1	9.00	
Q311M/N434S	IgG1/2	Anti-VEGF	1	9.12	
Q311N/N434S	IgG1/2	Anti-VEGF	1	2.27	
Q311P/N434S	IgG1/2	Anti-VEGF	1	0.56	
Q311R/N434S	IgG1/2	Anti-VEGF	1	5.96	
Q311T/N434S	IgG1/2	Anti-VEGF	1	5.17	
Q311V/N434S	IgG1/2	Anti-VEGF	1	14.17	
Q311W/N434S	IgG1/2	Anti-VEGF	1	2.97	
Q311Y/N434S	IgG1/2	Anti-VEGF	1	2.65	
L314F/N434S	IgG1/2	Anti-VEGF	1	3.35	
L314I/N434S	IgG1/2	Anti-VEGF	1	3.54	
L314M/N434S	IgG1/2	Anti-VEGF	1	3.05	
L314Q/N434S	IgG1/2	Anti-VEGF	1	1.75	
L314V/N434S	IgG1/2	Anti-VEGF	1	2.34	
H429N/N434S	IgG1/2	Anti-VEGF	1	1.78	
H429V/N434S	IgG1/2	Anti-VEGF	1	0.69	
E430A/N434S	IgG1/2	Anti-VEGF	1	3.24	
E430D/N434S	IgG1/2	Anti-VEGF	1	1.79	

Figure 10 (continued)

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
E430F/N434S	IgG1/2	Anti-VEGF	1	3.79	
E430I/N434S	IgG1/2	Anti-VEGF	1	5.27	
E430L/N434S	IgG1/2	Anti-VEGF	1	4.40	
E430Q/N434S	IgG1/2	Anti-VEGF	1	4.20	
E430S/N434S	IgG1/2	Anti-VEGF	1	2.58	
E430T/N434S	IgG1/2	Anti-VEGF	1	2.37	
A431F/N434S	IgG1/2	Anti-VEGF	1	5.21	
A431G/N434S	IgG1/2	Anti-VEGF	1	4.31	
A431H/N434S	IgG1/2	Anti-VEGF	1	6.29	
A431N/N434S	IgG1/2	Anti-VEGF	1	5.06	
A431S/N434S	IgG1/2	Anti-VEGF	1	3.18	
A431T/N434S	IgG1/2	Anti-VEGF	1	2.46	
A431V/N434S	IgG1/2	Anti-VEGF	1	4.64	
L432A/N434S	IgG1/2	Anti-VEGF	1	4.83	
L432F/N434S	IgG1/2	Anti-VEGF	1	1.14	
L432G/N434S	IgG1/2	Anti-VEGF	1	1.71	
L432H/N434S	IgG1/2	Anti-VEGF	1	1.69	
L432I/N434S	IgG1/2	Anti-VEGF	1	1.96	
L432N/N434S	IgG1/2	Anti-VEGF	1	1.43	
L432Q/N434S	IgG1/2	Anti-VEGF	1	1.59	
L432V/N434S	IgG1/2	Anti-VEGF	1	2.24	
H433A/N434S	IgG1/2	Anti-VEGF	1	2.41	
H433D/N434S	IgG1/2	Anti-VEGF	1	0.75	
H433E/N434S	IgG1/2	Anti-VEGF	1	0.73	
H433F/N434S	IgG1/2	Anti-VEGF	1	0.76	
H433I/N434S	IgG1/2	Anti-VEGF	1	0.94	
H433K/N434S	IgG1/2	Anti-VEGF	1	3.65	
H433L/N434S	IgG1/2	Anti-VEGF	1	1.14	
H433M/N434S	IgG1/2	Anti-VEGF	1	1.91	
H433P/N434S	IgG1/2	Anti-VEGF	1	6.33	
H433S/N434S	IgG1/2	Anti-VEGF	1	3.68	
H433T/N434S	IgG1/2	Anti-VEGF	1	0.88	
H433V/N434S	IgG1/2	Anti-VEGF	1	2.41	
H433Y/N434S	IgG1/2	Anti-VEGF	1	0.56	
N434S/Y436A	IgG1/2	Anti-VEGF	1	0.03	
N434S/Y436D	IgG1/2	Anti-VEGF	1	1.79	
N434S/Y436E	IgG1/2	Anti-VEGF	1	4.36	

Figure 10 (continued)

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
N434S/Y436G	IgG1/2	Anti-VEGF	1	4.24	
N434S/Y436H	IgG1/2	Anti-VEGF	1	7.63	
N434S/Y436I	IgG1/2	Anti-VEGF	1	30.57	
N434S/Y436K	IgG1/2	Anti-VEGF	1	9.00	
N434S/Y436L	IgG1/2	Anti-VEGF	1	12.81	
N434S/Y436N	IgG1/2	Anti-VEGF	1	0.39	
N434S/Y436Q	IgG1/2	Anti-VEGF	1	2.14	
N434S/Y436R	IgG1/2	Anti-VEGF	1	5.60	
N434S/Y436S	IgG1/2	Anti-VEGF	1	6.67	
N434S/Y436T	IgG1/2	Anti-VEGF	1	4.71	
N434S/Y436V	IgG1/2	Anti-VEGF	1	17.46	
N434S/Y436W	IgG1/2	Anti-VEGF	1	19.47	
N434S/T437I	IgG1/2	Anti-VEGF	1	2.96	
G236R/L328R	IgG1	Anti-VEGF	1	1.25	
V308F/I332E	IgG1	Anti-VEGF	1	4.15	
I332E/N434S	IgG1	Anti-VEGF	1	3.91	
G236A/V308F/I332E	IgG1/2 ELLG 327A	Anti-VEGF	1	1.82	
G236A/I332E/N434S	IgG1/2 ELLG 327A	Anti-VEGF	1	3.87	

Figure 11a




Figure 11b




Figure 11c




Figure 12a

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG1	Anti-TNF		1.00	
V259I/V308F	IgG1	Anti-TNF	3	13.25	5.09
M428L/N434S	IgG1	Anti-TNF	2	19.76	4.42
V259I/V308F/M428L	IgG1	Anti-TNF	2	49.85	5.09
G236R/L328R	IgG1	Anti-TNF	1	0.99	
WT	IgG2	Anti-TNF		1.00	
N434S	IgG2	Anti-TNF	2	3.12	0.79
M428L/N434S	IgG2	Anti-TNF	3	19.99	8.69
L328R	IgG2	Anti-TNF	1	0.97	
L328R/M428L/N434S	IgG2	Anti-TNF	1	19.67	
WT	IgG1/2	Anti-TNF		1.00	
N434S	IgG1/2	Anti-TNF	2	7.22	0.16
M428L/N434S	IgG1/2	Anti-TNF	3	25.39	5.13
L328R	IgG1/2	Anti-TNF	1	1.87	
L328R/M428L/N434S	IgG1/2	Anti-TNF	1	39.90	

Figure 12b

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG1	Anti-CD25		1.00	
V259I/V308F	IgG1	Anti-CD25	1	3.49	
V259I/V308F/M428L	IgG1	Anti-CD25	1	19.97	
WT	IgG1/2	Anti-CD25		1.00	
N434S	IgG1/2	Anti-CD25	1	5.16	
M428L/N434S	IgG1/2	Anti-CD25	1	28.00	

Figure 12c

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG1	Anti-EGFR		1.00	
V259I/V308F	IgG1	Anti-EGFR	3	14.23	8.51
V259I/V308F/M428L	IgG1	Anti-EGFR	2	32.96	20.16
WT	IgG1/2	Anti-EGFR		1.00	
N434S	IgG1/2	Anti-EGFR	1	1.61	
N434S/M428L	IgG1/2	Anti-EGFR	1	5.93	

Figure 12d

Variant	IgG	Fv	n	Fold FcRn	
				Mean	SD
WT	IgG1	Anti-IgE		1.00	
M428L	IgG1	Anti-IgE	1	3.06	
N434S	IgG1	Anti-IgE	1	4.38	
V259I/V308F	IgG1	Anti-IgE	1	8.34	
V308F/Y319I	IgG1	Anti-IgE	1	5.39	
V308F/M428L	IgG1	Anti-IgE	1	14.25	
M428L/N434S	IgG1	Anti-IgE	1	12.63	
M252Y/S254T/T256E	IgG1	Anti-IgE	1	9.56	
V259I/V308F/M428L	IgG1	Anti-IgE	1	20.27	
V259I/V308F/M428L/N434S	IgG1	Anti-IgE	1	27.29	

Figure 13a




Figure 13b




Figure 14

Fc variant	IgG	n	Half-Life			Cmax		AUC		Clearance	
			(day)			(ug/mL)		(day*ug/mL)		(mL/day/kg)	
			Mean	SD	Fold	Mean	SD	Mean	SD	Mean	SD
Study 1											
WT (Native IgG1)	IgG1	6	3	0.5	1.0	28.5	7.3	69.8	8.4	29.1	4.2
P257L	IgG1	6	1.9	0.4	0.6	18.1	8.4	38	11.3	59.5	28.8
P257N	IgG1	6	0.9	0.1	0.3	30.1	4.7	38.2	6.1	53.4	8.8
V308F	IgG1	6	4.9	0.3	1.6	30.1	9.4	129.2	11	15.6	1.3
T250Q/M428L	IgG1	6	8.4	2.8	2.8	22.3	3.8	186	52.5	12	5.4
M252Y/S254T/T256E	IgG1	4	10.9	1.2	3.6	34.6	10.1	330.8	57.3	6.2	1.1
N434A	IgG1	6	5.5	2.1	1.8	24.5	6.2	143.1	26.9	14.3	2.2
Study 2											
WT (Native IgG1)	IgG1	7	3.9	0.52	1.0	36	5.6	92	13	22	2.9
V308F	IgG1	7	6.8	0.99	1.7	28	8.7	152	34.6	14	3.4
M252T/S254T/T256E	IgG1	7	11.3	1.2	2.9	25	7	241	64.2	8.9	2.6
Study 3											
WT (Native IgG1)	IgG1	6	2.8	0.3	1.0	27	6	69	10	29.4	4.6
V308F	IgG1	6	5.9	0.4	2.1	33	3	173	33	11.8	1.9
M252T/S254T/T256E	IgG1	6	10.4	1.5	3.7	34	6	317	67	6.6	1.5
N434S	IgG1	6	7.7	1.5	2.8	33	11	228	75	10	4.6
V259I/V308F	IgG1	6	9.2	1.5	3.3	36	5	262	47	7.9	1.4
V308F/M428L	IgG1	6	11.2	1.1	4.0	38	6	307	50	6.7	1.1
M252Y/V308F	IgG1	6	5.7	0.4	2.0	40	6	140	24	14.7	2.6
M428L/N434S	IgG1	6	12	2.9	4.3	37	9	400	112	5.5	2
V308F/N434S	IgG1	6	8.6	0.3	3.1	28	3	195	25	10.4	1.3
V259I/V308F/M428L	IgG1	6	13.3	2.7	4.8	37	6	383	68	5.3	0.8

Figure 14 (continued)

Fc variant	IgG	n	Half-Life			Cmax		AUC		Clearance	
			(day)			(ug/mL)		(day*ug/mL)		(mL/day/kg)	
			Mean	SD	Fold	Mean	SD	Mean	SD	Mean	SD
Study 4											
WT (Native IgG1)	IgG1	6	2.9	0.4	1.0	32.5	3.1	72.7	5.8	27.6	2.3
M252Y/S254T/T256E	IgG1	6	11.3	1.8	3.9	40.6	5.8	376.5	60.7	5.4	0.8
V308F/N434M	IgG1	6	4.9	0.8	1.7	38.6	5.4	153.6	21.3	13.2	1.9
V259I/V308F	IgG1	6	7.5	0.8	2.6	37.9	3.8	234.5	23.1	8.6	0.9
V308F	IgG1	6	6.1	0.4	2.1	37.8	1.9	172.2	16.7	11.7	1.2
V308F/Y319I	IgG1	6	6	0.5	2.1	38.1	6.5	206.5	71	10.5	3.1
M428L/N434S	IgG1	6	11.8	0.6	4.1	42	4.6	392.4	52.2	5.2	0.7
V308F/M428L/N434S	IgG1	6	9	0.4	3.1	31.8	7.9	204.6	34.2	10	1.8
V259I/V308F/N434S	IgG1	6	8.5	1.8	2.9	33.4	7.3	254.9	29	7.9	0.9
V259I/V308F/M428L	IgG1	6	10.9	0.6	3.8	35	7.5	295.1	54.5	7	1.3
Study 5											
WT (Native IgG2)	IgG2	6	5.9	0.9	1.0	43.8	5	212.3	33.8	9.6	1.5
N434S	IgG2	6	12.2	1.3	2.1	39.1	6.4	381.6	91.6	5.5	1.2
V259I/V308F	IgG2	6	10.2	0.6	1.7	42.3	1.3	342	21.2	5.9	0.4
M428L/N434S	IgG2	6	16.5	1.1	2.8	48	5.8	603.3	72.9	3.4	0.4

Figure 15a




Figure 15b




Figure 15c




Figure 16

Anti-TNF Fc fusion with IgG1 Fc (SEQ ID NO:30)

MAPVAVWAALAVGLELWAAHALPAQVAFPTYAPEPGSTCRLREYYDQTAQMCCSKCSPGQHA
 KVFC TKTS DTVCDSCEDSTYTQLWNWVPECLSCGSRCS SDQVETQACTREQNRICRPGWYC
 ALSKQEGCRLCAPLRKCRPGFGVARPGTETSDVVKPCAPGTFSNTTSSTDICRPHQICNVVAIP
 GNASMDAVCTSTSPTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTS FLLPMGPSPPAEGST
 GDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY
 VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS
 KLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Anti-TNF Fc fusion with IgG2 Fc (SEQ ID NO:31)

MAPVAVWAALAVGLELWAAHALPAQVAFPTYAPEPGSTCRLREYYDQTAQMCCSKCSPGQHA
 KVFC TKTS DTVCDSCEDSTYTQLWNWVPECLSCGSRCS SDQVETQACTREQNRICRPGWYC
 ALSKQEGCRLCAPLRKCRPGFGVARPGTETSDVVKPCAPGTFSNTTSSTDICRPHQICNVVAIP
 GNASMDAVCTSTSPTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTS FLLPMGPSPPAEGST
 GDEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWY
 VDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLY
 SKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Figure 16 (continued)**Anti-TNF Fc fusion with IgG1 259I/308F Fc (SEQ ID NO:32)**

MAPVAVWAALAVGLELWAAHALPAQVAFTPYAPEPGSTCRLREYYDQTAQMCCSKCSPGQHA
 KVFCTKTSDTVCDSCEDSTYTQLWNWVPECLSCGSRCSDDQVETQACTREQNRICRCRPGWYC
 ALSKQEGCRLCAPLRKCRPGFGVARPGTETSDVVKPCAPGTFSNTTSSTDICRPHQICNVVAIP
 GNASMDAVCTSTSPTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTSFLPMGSPSPAEGST
 GDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEITCVVVDVSHEDPEVKFNWY
 VDGVEVHNAKTKPREEQYNSTYRVVSVLTLFHDQDLNGKEYKCKVSNKALPAPIEKTISKAKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS
 KLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK

Anti-TNF Fc fusion with IgG1 428L/434S Fc (SEQ ID NO:33)

MAPVAVWAALAVGLELWAAHALPAQVAFTPYAPEPGSTCRLREYYDQTAQMCCSKCSPGQHA
 KVFCTKTSDTVCDSCEDSTYTQLWNWVPECLSCGSRCSDDQVETQACTREQNRICRCRPGWYC
 ALSKQEGCRLCAPLRKCRPGFGVARPGTETSDVVKPCAPGTFSNTTSSTDICRPHQICNVVAIP
 GNASMDAVCTSTSPTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTSFLPMGSPSPAEGST
 GDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY
 VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS
 KLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK

Anti-TNF Fc fusion with IgG2 434S Fc (SEQ ID NO:34)

MAPVAVWAALAVGLELWAAHALPAQVAFTPYAPEPGSTCRLREYYDQTAQMCCSKCSPGQHA
 KVFCTKTSDTVCDSCEDSTYTQLWNWVPECLSCGSRCSDDQVETQACTREQNRICRCRPGWYC
 ALSKQEGCRLCAPLRKCRPGFGVARPGTETSDVVKPCAPGTFSNTTSSTDICRPHQICNVVAIP
 GNASMDAVCTSTSPTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTSFLPMGSPSPAEGST
 GDEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWY
 VDGVEVHNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLY
 SKLTVDKSRWQQGNVFSCSVMEALHSHYTQKSLSLSPGK

Anti-TNF Fc fusion with IgG2 428L/434S Fc (SEQ ID NO:35)

MAPVAVWAALAVGLELWAAHALPAQVAFTPYAPEPGSTCRLREYYDQTAQMCCSKCSPGQHA
 KVFCTKTSDTVCDSCEDSTYTQLWNWVPECLSCGSRCSDDQVETQACTREQNRICRCRPGWYC
 ALSKQEGCRLCAPLRKCRPGFGVARPGTETSDVVKPCAPGTFSNTTSSTDICRPHQICNVVAIP
 GNASMDAVCTSTSPTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTSFLPMGSPSPAEGST
 GDEPKSCDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWY
 VDGVEVHNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLY
 SKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK

Figure 17




Figure 18

Variant	Fc region	Fusion partner	n	Fold FcRn	
				Mean	SD
WT	IgG1	TNFR		1.00	
V259I/V308F	IgG1	TNFR	4	4.86	2.22
M428L/N434S	IgG1	TNFR	2	7.24	1.57
WT	IgG2	TNFR		1.00	
N434S	IgG2	TNFR	3	6.93	2.02
M428L/N434S	IgG2	TNFR	3	11.78	4.33

Figure 19




Figure 20

Fc variant	Fc	n	Half-Life		Cmax	AUC	Clearance
			(day)	Increase (%)	(ng/mL)	(day*ng/mL)	(mL/day/kg)
WT	IgG1	12	0.60	n/a	22253.3	1324.7	1177.7
V259I/V308F	IgG1	12	0.81	136	12175.6	2291.5	741.9
WT	IgG2	12	0.65	n/a	14580.2	3686.3	423.2
N434S	IgG2	12	0.77	119	12086.8	2531.3	655.8
M428L/N434S	IgG2	12	1.01	156	30676.7	4405.2	354.1

Figure 21a

	Fold cyno FcRn			Fold human FcRn		
	n	Mean	SD	n	Mean	SD
WT IgG1		1.00			1.00	
M252Y/S254T/T256E	2	11.95	2.23	10	11.37	2.90
V259I/V308F	2	8.60	1.86	12	9.79	2.72
M428L/N434S	3	12.18	1.23	14	17.28	6.70
V259I/V308F/M428L	4	12.43	0.57	13	19.94	10.46

Figure 21b




Figure 22




Figure 23

Fc variant	IgG	n	Half-Life			Cmax		AUC		Clearance	
			(day)			(ug/mL)		Day*ug/mL		mL/day/kg	
			Mean	SD	Fold	Mean	SD	Mean	SD	Mean	SD
WT (Native IgG1)	IgG1	3	9.7		1.0	101.2		822.9		4.9	
M252Y/S254T/T256E	IgG1	3	24.2	1.6	2.5	111.2	13.9	1919	210.1	2.1	0.2
V259I/V308F	IgG1	3	16.2	6.4	1.7	102.8	3.2	1352.6	367	3.1	0.9
M428L/N434S	IgG1	3	31.1	7.9	3.2	126	17	2660.9	791	1.6	0.6
V259I/V308F/M428L	IgG1	3	25.1	5.9	2.6	124.6	14.3	2301.6	922.5	1.9	0.8

CH1

EU Index	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138
IgG1	A	S	T	K	Q	P	S	V	F	P	L	A	P	S	S	K	S	T	S	G	Q
IgG2	A	S	T	K	Q	P	S	V	F	P	L	A	P	C	S	R	S	T	S	E	S
IgG3	A	S	T	K	Q	P	S	V	F	P	L	A	P	C	S	R	S	T	S	Q	Q
IgG4	A	S	T	K	Q	P	S	V	F	P	L	A	P	C	S	R	S	T	S	E	S

EU Index	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159
IgG1	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N
IgG2	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N
IgG3	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N
IgG4	T	A	A	L	G	C	L	V	K	D	Y	F	P	E	P	V	T	V	S	W	N

EU Index	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
IgG1	S	Q	A	L	T	S	Q	V	H	T	F	P	A	V	L	Q	S	S	Q	L	Y
IgG2	S	Q	A	L	T	S	Q	V	H	T	F	P	A	V	L	Q	S	S	Q	L	Y
IgG3	S	Q	A	L	T	S	Q	V	H	T	F	P	A	V	L	Q	S	S	Q	L	Y
IgG4	S	Q	A	L	T	S	Q	V	H	T	F	P	A	V	L	Q	S	S	Q	L	Y

EU Index	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201
IgG1	S	L	S	S	V	T	V	P	S	S	S	L	G	T	Q	T	Y	I	C	N	
IgG2	S	L	S	S	V	T	V	P	S	S	N	F	G	T	Q	T	Y	T	C	N	
IgG3	S	L	S	S	V	T	V	P	S	S	S	L	G	T	Q	T	Y	T	C	N	
IgG4	S	L	S	S	V	T	V	P	S	S	S	L	G	T	K	T	Y	T	C	N	

EU Index	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220
IgG1	V	N	H	K	P	S	N	T	K	V	D	K	KR	V	E	P	K	S	C
IgG2	V	D	H	K	P	S	N	T	K	V	D	K	T	V	E	R	K	C	C
IgG3	V	N	H	K	P	S	N	T	K	V	D	K	R	V	E	L	K	T	P
IgG4	V	D	H	K	P	S	N	T	K	V	D	K	R	V	E	S	K	Y	G

Hinge

Fc >

EU Index	221	222	223	224	225	226	227	228
IgG1	D	K	T	H	T	C	P	P
IgG2		V		E		C	P	P
IgG3	L	G	D	T	T	H	T	C
IgG4				P	P	C	P	S

EU Index

EU Index	229	230	231	232	233	234	235	236
IgG1	P	C	P	R	C	P	E	P
IgG2								
IgG3	P	C	P	R	C	P	E	P
IgG4								

EU Index

EU Index	229	230	231	232	233	234	235	236
IgG1	C	P	A	P	E	L	L	G
IgG2	C	P	A	P	P	V	A	
IgG3	E	P	K	S	C	D	T	F
IgG4	C	P	A	P	E	F	L	Q