




(86) Date de dépôt PCT/PCT Filing Date: 2005/04/20
(87) Date publication PCT/PCT Publication Date: 2005/11/03
(85) Entrée phase nationale/National Entry: 2007/10/22
(86) N° demande PCT/PCT Application No.: EP 2005/004325
(87) N° publication PCT/PCT Publication No.: 2005/103713
(30) Priorité/Priority: 2004/04/20 (US60/563,661)

(51) Cl.Int./Int.Cl. *G01N 33/68* (2006.01),
G01N 33/566 (2006.01)
(71) Demandeur/Applicant:
GALAPAGOS N.V., BE
(72) Inventeurs/Inventors:
MERCHIEERS, PASCAL GERARD, BE;
SPITTAELS, KOENRAAD FREDERIK FLORENTINA,
BE;
HOFFMANN, MARCEL, NL
(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : METHODES, COMPOSITIONS ET ANALYSE DE COMPOSES DESTINEES A INHIBER LA PRODUCTION
D'UNE PROTEINE BETA-AMYLOIDE
(54) Title: METHODS, COMPOSITIONS AND COMPOUND ASSAYS FOR INHIBITING AMYLOID-BETA PROTEIN
PRODUCTION



(57) **Abrégé/Abstract:**

A method for identifying compounds that inhibit amyloid-beta precursor protein processing in cells, comprising contacting a test compound with a GPCR polypeptide, (eg. GPR3, GPR6, GPR12) or fragment thereof, and measuring a compound-GPCR property related to the production of amyloid-beta peptide. Cellular assays of the method measure indicators including second messenger and/or amyloid beta peptide levels. Therapeutic methods, and pharmaceutical compositions including effective amyloid-beta precursor processing-inhibiting amounts of GPCR expression inhibitors, are useful for treating conditions involving cognitive impairment such as Alzheimers Disease.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 November 2005 (03.11.2005)

PCT

(10) International Publication Number
WO 2005/103713 A3

(51) International Patent Classification:

G01N 33/68 (2006.01) *G01N 33/566* (2006.01)

(21) International Application Number:

PCT/EP2005/004325

(22) International Filing Date: 20 April 2005 (20.04.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/563,661 20 April 2004 (20.04.2004) US

(71) Applicant (for all designated States except US): **GALA-PAGOS N.V.** [BE/BE]; Generaal De Wittelaan L11 A3, B-2800 Mechelen (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MERCHIEERS, Pascal, Gerard** [BE/BE]; Loopstraat 23, B-2460 Tielen (BE). **SPITTAELS, Koenraad, Frederik, Florentina** [BE/BE]; Essendries 10, B-2870 Puurs (BE). **HOFFMANN, Marcel** [NL/NL]; Valeriuslaan 12, NL-1422 HT Uithoorn (NL).

(74) Agent: **HOOVELD, Arjen, Jan, Winfried**; Arnold & Siedsma, Sweelinckplein 1, NL-2517 GK Den Haag (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report

(88) Date of publication of the international search report:

8 September 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS, COMPOSITIONS AND COMPOUND ASSAYS FOR INHIBITING AMYLOID-BETA PROTEIN PRODUCTION



(57) Abstract: A method for identifying compounds that inhibit amyloid-beta precursor protein processing in cells, comprising contacting a test compound with a GPCR polypeptide, (eg. GPR3, GPR6, GPR12) or fragment thereof, and measuring a compound-GPCR property related to the production of amyloid-beta peptide. Cellular assays of the method measure indicators including second messenger and/or amyloid beta peptide levels. Therapeutic methods, and pharmaceutical compositions including effective amyloid-beta precursor processing-inhibiting amounts of GPCR expression inhibitors, are useful for treating conditions involving cognitive impairment such as Alzheimers Disease.

WO 2005/103713 A3

5

10

METHODS, COMPOSITIONS AND COMPOUND ASSAYS FOR INHIBITING
AMYLOID-BETA PROTEIN PRODUCTION

15

Field of the Invention

This invention relates to the field of mammalian neuronal cell disorders, and in particular, to methods for identifying effective compounds, and therapies and compositions using such compounds, useful for the prevention and treatment of diseases associated with progressive loss of intellectual capacities in humans.

20

The neurological disorder that is most widely known for its progressive loss of intellectual capacities is Alzheimer's disease (AD). Worldwide, about 20 million people suffer from Alzheimer's disease. AD is clinically characterized by the initial loss of memory, followed by disorientation, impairment of judgment and reasoning, which is commonly referred to as cognitive impairment, and ultimately by full dementia. AD patients finally

25 lapse into a severely debilitated, immobile state between four and twelve years after onset of the disease.

The key pathological evidence for AD is the presence of extracellular amyloid plaques and intracellular tau tangles in the brain, which are associated with neuronal degeneration (Ritchie and Lovestone (2002)). The extracellular amyloid plaques are believed

to result from an increase in the insoluble amyloid beta peptide 1-42 produced by the metabolism of amyloid-beta precursor protein (APP). Following secretion, these amyloid beta 1-42 peptides form amyloid fibrils more readily than the amyloid beta 1-40 peptides, which are predominantly produced in healthy people. It appears that the amyloid beta peptide is on top of the neurotoxic cascade: experiments show that amyloid beta fibrils, when injected into the brains of P301L tau transgenic mice, enhance the formation of neurofibrillary tangles (Gotz et al. (2001)). In fact, a variety of amyloid beta peptides have been identified as amyloid beta peptides 1-42, 1-40, 1-39, 1-38, 1-37, which can be found in plaques and are often seen in cerebral spinal fluid.

The amyloid beta peptides are generated (or processed) from the membrane anchored APP, after cleavage by beta secretase and gamma secretase at position 1 and 40 or 42, respectively (Figure 1A)(Annaert and De Strooper (2002)). In addition, high activity of beta secretase results in a shift of the cleavage at position 1 to position 11. Cleavage of amyloid-beta precursor protein by alpha secretase activity at position 17 and gamma secretase activity at 40 or 42 generates the non-pathological p3 peptide. Beta secretase was identified as the membrane anchored aspartyl protease BACE, while gamma secretase is a protein complex comprising presenilin 1 (PS1) or presenilin 2 (PS2), nicastrin, Anterior Pharynx Defective 1 (APH1) and Presenilin Enhancer 2 (PEN2). Of these proteins, the presenilins are widely thought to constitute the catalytic activity of the gamma secretase, while the other components play a role in the maturation and localization of the complex. The identity of the alpha secretase is still illustrious, although some results point towards the proteases ADAM 10 and TACE, which could have redundant functions.

A small fraction of AD cases (mostly early onset AD) are caused by autosomal dominant mutations in the genes encoding presenilin 1 and 2 (PS1; PS2) and the amyloid-beta precursor protein (APP), and it has been shown that mutations in APP, PS1 and PS2 alter the metabolism of amyloid-beta precursor protein leading to such increased levels of amyloid beta 1-42 produced in the brain. Although no mutations in PS1, PS2 and amyloid-beta precursor protein have been identified in late onset AD patients, the pathological characteristics are highly similar to the early onset AD patients. These increased levels of amyloid beta peptide could originate progressively with age from disturbed amyloid-beta precursor protein processing (e.g. high cholesterol levels enhance amyloid beta peptide production) or from decreased amyloid beta peptide catabolism. Therefore, it is generally accepted that AD in late onset AD patients is also caused by aberrant increased amyloid

5 Reported Developments

10 Therapies aimed at decreasing the levels of amyloid beta peptides in the brain, are increasingly being investigated and focus on the perturbed amyloid-beta precursor protein processing involving the beta- or gamma secretase enzymes.

The present invention is based on the discovery that certain known polypeptides are factors in the up-regulation and/or induction of amyloid beta precursor processing in neuronal cells, and that the inhibition of the function of such polypeptides are effective in reducing levels of amyloid beta peptides.

Summary of the Invention

The present invention relates to the relationship between the function of the G-protein coupled receptor(s) ("GPCR(s)") and amyloid-beta precursor protein processing in mammalian cells.

One aspect of the present invention is a method for identifying a compound that inhibits the processing of amyloid-beta precursor protein in a mammalian cell, comprising

- (a) contacting a compound with a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333; and
- 25 (b) measuring a compound-polypeptide property related to the production of amyloid-beta protein. .

Aspects of the present method include the *in vitro* assay of compounds using polypeptide domains of a GPCR, and cellular assays wherein GPCR inhibition is followed by observing indicators of efficacy, including second messenger levels and/or amyloid beta peptide levels.

Another aspect of the invention is a method of treatment or prevention of a condition involving cognitive impairment, or a susceptibility to the condition, in a subject suffering or susceptible thereto, by administering a pharmaceutical composition comprising an effective amyloid-beta precursor processing-inhibiting amount of a GPCR antagonist or inverse agonist.

A further aspect of the present invention is a pharmaceutical composition for use in said method wherein said inhibitor comprises a polynucleotide selected from the group of an antisense polynucleotide, a ribozyme, and a small interfering RNA (siRNA), wherein said agent comprises a nucleic acid sequence complementary to, or engineered from, a naturally occurring polynucleotide sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6.

Another further aspect of the present invention is a pharmaceutical composition comprising a therapeutically effective amyloid-beta precursor processing-inhibiting amount of a GPCR antagonist or inverse agonist or its pharmaceutically acceptable salt, hydrate, solvate, or prodrug thereof in admixture with a pharmaceutically acceptable carrier. The present polynucleotides and GPCR antagonist and inverse agonist compounds are also useful for the manufacturing of a medicament for the treatment of Alzheimer's disease.

Brief Description of the Drawings

Figure 1A: APP processing: The membrane anchored amyloid precursor protein (APP) is processed by two pathways: the amyloidogenic and non amyloidogenic pathway. In the latter pathway, APP is cleaved first by alpha secretase and then by gamma secretase, yielding the p3 peptides (17-40 or 17-42). The amyloidogenic pathway generates the pathogenic amyloid beta peptides (A beta) after cleavage by beta- and gamma-secretase respectively. The numbers depicted are the positions of the amino acids comprising the A beta sequences.

Figure 1B: Pictorial representation of transmembrane structure of GPCR proteins.

Figure 2: Evaluation of the APP processing assay: Positive (PS1G384L; PS1L392V and BACE1) and negative (eGFP, LacZ and empty) control viruses are infected in Hek293APPwt at random MOI, mimicking a screening. A and B: Transduction is performed respectively with 1 and 0.2 µl of virus and amyloid beta 1-42 levels are performed. Data are represented as relative light units and correlate to pM of amyloid beta 1-42.

Figure 3: Involvement of GPR3 in APP processing: HEK293 APPwt cells are transduced with Ad5/GPR3 and with negative control viruses (Ad5/empty, Ad5/LacZ, Ad5/eGFP and Ad5/luciferase) at different MOIs (2, 10, 50, 250). Resulting amyloid beta 1-42, 1-40, 11-42, x-42 and 1-x peptides were measured with the appropriate ELISAs. Data are represented in pM or as relative light units (rlu), which correlates to pM of amyloid beta.

Figure 4: Transfection with GPR3 siRNA reduces Amyloid beta 1-42: HEK293 APPwt cl29 cells are transfected with siRNA of GPR3, eGFP, Luciferase and BACE and amyloid beta 1-42 levels are determined. Cells are transfected and 24 hours after transfection, medium is refreshed and cells are allowed to accumulate amyloid beta for 24 hours (48 hours post transfection (p.t.)). Amyloid beta is determined by means of the amyloid beta 1-42 ELISA as described above. Data are presented in pM of amyloid beta. RNA levels of GPR3 are determined from these samples.

Figure.5: ClustalW protein sequence alignment of GPR3, GPR6 and GPR12.

Figure 6: Graph of amyloid beta peptide levels in neurons transfected with a variety of protein expression viruses at different MOI. The graph shows that increased levels of GPR3 overexpression in primary neurons result in a corresponding dose dependent increase of amyloid beta 1-42 levels compared to the negative controls.

Detailed Description

The following terms are intended to have the meanings presented therewith below and are useful in understanding the description of and intended scope of the present invention.

Definitions:

The term “agonist” refers to a ligand that activates the intracellular response of the receptor to which the agonist binds.

The term “amyloid beta peptide” means amyloid beta peptides processed from the amyloid beta precursor protein (APP). The most common peptides include amyloid beta peptides 1-40, 1-42, 11-40 and 11-42. Other species less prevalent amyloid beta peptides are described as y-42, whereby y ranges from 2-17, and 1-x whereby x ranges from 24-39 and 41.

The term "antagonist" means a moiety that bind competitively to the receptor at the same site as the agonists but which do not activate the intracellular response initiated by the active form of the receptor, and can thereby inhibit the intracellular responses by agonists. Antagonists do not diminish the baseline intracellular response in the absence of an agonist or
5 partial agonist.

The term "carrier" means a non-toxic material used in the formulation of pharmaceutical compositions to provide a medium, bulk and/or useable form to a pharmaceutical composition. A carrier may comprise one or more of such materials such as an excipient, stabilizer, or an aqueous pH buffered solution. Examples of physiologically
10 acceptable carriers include aqueous or solid buffer ingredients including phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other
15 carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counter ions such as sodium; and/or nonionic surfactants such as TWEEN.TM., polyethylene glycol (PEG), and PLURONICS.TM..

The term "compound" is used herein in the context of a "test compound" or a "drug
20 candidate compound" described in connection with the assays of the present invention. As such, these compounds comprise organic or inorganic compounds, derived synthetically or from natural sources. The compounds include inorganic or organic compounds such as polynucleotides, lipids or hormone analogs that are characterized by relatively low molecular weights. Other biopolymeric organic test compounds include peptides comprising from
25 about 2 to about 40 amino acids and larger polypeptides comprising from about 40 to about 500 amino acids, such as antibodies or antibody conjugates.

The term "constitutive receptor activation" means stabilization of a receptor in the active state by means other than binding of the receptor with its endogenous ligand or a chemical equivalent thereof.

30 The term "contact" or "contacting" means bringing at least two moieties together, whether in an in vitro system or an in vivo system.

The term "condition" or "disease" means the overt presentation of symptoms (i.e., illness) or the manifestation of abnormal clinical indicators (e.g., biochemical indicators), resulting from defects in one amyloid beta protein precursor processing. Alternatively, the term "disease" refers to a genetic or environmental risk of or propensity for developing such symptoms or abnormal clinical indicators.

The term "endogenous" shall mean a material that a mammal naturally produces. Endogenous in reference to, for example and not limitation, the term "receptor" shall mean that which is naturally produced by a mammal (for example, and not limitation, a human) or a virus. In contrast, the term non-endogenous in this context shall mean that which is not naturally produced by a mammal (for example, and not limitation, a human) or a virus. For example, and not limitation, a receptor which is not constitutively active in its endogenous form, but when manipulated becomes constitutively active, is most preferably referred to herein as a "non-endogenous, constitutively activated receptor." Both terms can be utilized to describe both "in vivo" and "in vitro" systems. For example, and not a limitation, in a screening approach, the endogenous or non-endogenous receptor may be in reference to an in vitro screening system. As a further example and not limitation, where the genome of a mammal has been manipulated to include a non-endogenous constitutively activated receptor, screening of a candidate compound by means of an in vivo system is viable.

The term "expression" comprises both endogenous expression and overexpression by transduction.

The term "expressible nucleic acid" means a nucleic acid coding for a proteinaceous molecule, an RNA molecule, or a DNA molecule.

The term "hybridization" means any process by which a strand of nucleic acid binds with a complementary strand through base pairing. The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C.sub.0t or R.sub.0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed). The term "stringent conditions" refers to conditions that permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt concentration, the concentration

of organic solvent, e.g., formamide, temperature, and other conditions well known in the art. In particular, reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature can increase stringency.

5 The term "inhibit" or "inhibiting", in relationship to the term "response" means that a response is decreased or prevented in the presence of a compound as opposed to in the absence of the compound.

10 The term "inverse agonist" mean a moiety that binds the endogenous form of the receptor, and which inhibits the baseline intracellular response initiated by the active endogenous form of the receptor below the normal base level of activity that is observed in the absence of the endogenous ligand, or agonists, or decrease GTP binding to membranes. Preferably, the baseline intracellular response is decreased in the presence of the inverse agonist by at least 30%, more preferably by at least 50%, and most preferably by at least 75%, as compared with the baseline response in the absence of the inverse agonist.

15 The term "ligand" means an endogenous, naturally occurring molecule specific for an endogenous, naturally occurring receptor.

The term "pharmaceutically acceptable prodrugs" as used herein means the prodrugs of the compounds useful in the present invention, which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients with undue toxicity, irritation, allergic response commensurate with a reasonable benefit/risk ratio, and effective for their intended use of the compounds of the invention. The term "prodrug" means a compound that is transformed in vivo to yield an effective compound useful in the present invention or a pharmaceutically acceptable salt, hydrate or solvate thereof. The transformation may occur by various mechanisms, such as through hydrolysis in blood. The compounds bearing metabolically cleavable groups have the advantage that they may exhibit improved bioavailability as a result of enhanced solubility and/or rate of absorption conferred upon the parent compound by virtue of the presence of the metabolically cleavable group, thus, such compounds act as pro-drugs. A thorough discussion is provided in Design of Prodrugs, H. Bundgaard, ed., Elsevier (1985); Methods in Enzymology; K. Widder et al, Ed., Academic Press, 42, 309-396 (1985); A Textbook of Drug Design and Development, Krogsgaard-Larsen and H. Bandaged, ed., Chapter 5; "Design and Applications of Prodrugs" 113-191 (1991); Advanced Drug Delivery Reviews, H. Bundgard, 8, 1-38, (1992); J. Pharm. Sci., 77, 285 (1988); Chem. Pharm. Bull., N. Nakeya et al, 32, 692

20
25
30

(1984); Pro-drugs as Novel Delivery Systems, T. Higuchi and V. Stella, 14 A.C.S. Symposium Series, and Bioreversible Carriers in Drug Design, E.B. Roche, ed., American Pharmaceutical Association and Pergamon Press, 1987, which are incorporated herein by reference. An example of the prodrugs is an ester prodrug. "Ester prodrug" means a compound that is convertible *in vivo* by metabolic means (e.g., by hydrolysis) to an inhibitor compound according to the present invention. For example an ester prodrug of a compound containing a carboxy group may be convertible by hydrolysis *in vivo* to the corresponding carboxy group.

The term "pharmaceutically acceptable salts" refers to the non-toxic, inorganic and organic acid addition salts, and base addition salts, of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of compounds useful in the present invention.

The term "polynucleotide" means a polynucleic acid, in single or double stranded form, and in the sense or antisense orientation, complementary polynucleic acids that hybridize to a particular polynucleic acid under stringent conditions, and polynucleotides that are homologous in at least about 60 percent of its base pairs, and more preferably 70 percent of its base pairs are in common, most preferably 90 per cent, and in a special embodiment 100 percent of its base pairs. The polynucleotides include polyribonucleic acids, polydeoxyribonucleic acids, and synthetic analogues thereof. The polynucleotides are described by sequences that vary in length, that range from about 10 to about 5000 bases, preferably about 100 to about 4000 bases, more preferably about 250 to about 2500 bases. A preferred polynucleotide embodiment comprises from about 10 to about 30 bases in length. A special embodiment of polynucleotide is the polyribonucleotide of from about 10 to about 22 nucleotides, more commonly described as small interfering RNAs (siRNAs). Another special embodiment are nucleic acids with modified backbones such as peptide nucleic acid (PNA), polysiloxane, and 2'-O-(2-methoxy)ethylphosphorothioate, or including non-naturally occurring nucleic acid residues, or one or more nucleic acid substituents, such as methyl-, thio-, sulphate, benzoyl-, phenyl-, amino-, propyl-, chloro-, and methanocarbanucleosides, or a reporter molecule to facilitate its detection.

The term "polypeptide" relates to proteins, proteinaceous molecules, fractions of proteins (such as kinases, proteases, GPCRs), peptides and oligopeptides.

The term "solvate" means a physical association of a compound useful in this invention with one or more solvent molecules. This physical association includes hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid.

5 "Solvate" encompasses both solution-phase and isolable solvates. Representative solvates include hydrates, ethanolates and methanolates.

The term "subject" includes humans and other mammals.

The term "effective amount" or "therapeutically effective amount" means that amount of a compound or agent that will elicit the biological or medical response of a subject that is
10 being sought by a medical doctor or other clinician. In particular, with regard to treating an neuronal disorder, the term "effective amount" is intended to mean that effective amyloid-beta precursor processing inhibiting amount of a compound or agent that will bring about a biologically meaningful decrease in the levels of amyloid beta peptide in the subject's brain tissue.

15 The term "treating" means an intervention performed with the intention of preventing the development or altering the pathology of, and thereby alleviating a disorder, disease or condition, including one or more symptoms of such disorder or condition. Accordingly, "treating" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treating include those already with the disorder as well as those in which the
20 disorder is to be prevented. The related term "treatment," as used herein, refers to the act of treating a disorder, symptom, disease or condition, as the term "treating" is defined above.

The background of the present inventors' discovery is described briefly below.

Background of the G-Protein Couple Receptors:

25 In 1994, Marchese and co-workers cloned the GPR3 gene (Marchese et al., 1994) and one year later, it was found that a single exon encoded this receptor protein of 330 amino acids, also called adenylate cyclase constitutive activator (ACCA). Based on the amino acid sequence, GPR3 can be classified in the same sub-family as GPR6 and GPR12: GPR3 and GPR6 exhibit 58% identity, and GPR3 and GPR12 57% (Figure 5).

30 G protein-coupled receptors (GPCR) share a common structural motif. All these receptors have seven sequences of between 22 to 24 hydrophobic amino acids that form seven alpha helices, each of which spans the membrane forming 7 transmembrane domains, an extracellular N-terminus and an intracellular C-terminus. The transmembrane helices are

joined by strands of amino acids having a larger loop between the fourth and fifth transmembrane helix on the extracellular side of the membrane. Another larger loop, composed primarily of hydrophilic amino acids, joins transmembrane helices five and six on the intracellular side of the membrane. See Figure 1B.

5 Under physiological conditions, GPCRs exist in the cell membrane in equilibrium between two different states or conformations: an "inactive" state and an "active" state. A receptor in an inactive state is unable to link to the intracellular transduction pathway to produce a biological response. Changing the receptor conformation to the active state allows linkage to the transduction pathway and produces a biological response. A receptor may be
10 stabilized in an active state by an endogenous ligand or an exogenous agonist ligand. Recent discoveries, including but not exclusively limited to, modifications to the amino acid sequence of the receptor, provide alternative mechanisms other than ligands to stabilize the active state conformation. These approaches effectively stabilize the receptor in an active state by simulating the effect of a ligand binding to the receptor. Stabilization by such
15 ligand-independent approaches is termed "constitutive receptor activation."

The major signal transduction cascades activated by GPCRs are initiated by the activation of heterotrimeric G-proteins, built from three different proteins; the G_α , G_β and G_γ subunits. It is believed that the loop joining helices five and six, as well as the carboxy terminus, interact with the G protein. Uhlenbrock and colleagues (2002) showed that GPR3,
20 GPR6 and GPR12 all confer constitutive activation of $G(a)(s)$ and $G(a)(i/o)$, and, recently, sphingosine-1-phosphate (S1P) and dihydrosphingosine 1-phosphate (DHS1P) have been identified as bioactive lipid ligands for GPR3, GPR6 and GPR12 (Uhlenbrock et al., 2002). The GPR3, GPR6 and GPR12 expression profile is also similar: they are all primary expressed in brain tissue.

25 The signal transduction cascade starts with the activation of the receptor by an agonist. Transformational changes in the receptor are then translated down to the G-protein. The G-protein dissociates into the G_α subunit and the $G_{\beta\gamma}$ subunit. Both subunits dissociate from the receptor and are both capable of initiating different cellular responses. Best known are the cellular effects that are initiated by the G_α subunit. It is for this reason that G-proteins
30 are categorized by their G_α subunit. The G-proteins are divided into four groups: G_s , $G_{i/o}$, G_q and $G_{12/13}$. Each of these G-proteins is capable of activating an effector protein, which results in changes in second messenger levels in the cell. The changes in second messenger level are

the triggers that make the cell respond to the extracellular signal in a specific manner. The activity of a GPCR can be measured by measuring the activity level of the second messenger.

The two most important second messengers in the cell are cAMP and Ca^{2+} . The α -subunit of the G_s class of G-proteins is able to activate adenylyl cyclase, resulting in an increased turnover from ATP to cAMP. The α -subunit of $G_{i/o}$ G-proteins does exactly the opposite and inhibits adenylyl cyclase activity resulting in a decrease of cellular cAMP levels. Together, these two classes of G-proteins regulate the second messenger cAMP. Ca^{2+} is regulated by the α -subunit of the G_q class of G-proteins. Through the activation of phospholipase C phosphatidylinositol 4,5-bisphosphate (PIP2) from the cell membrane are hydrolyzed to inositol 1,4,5-trisphosphate and 1,2-diacylglycerol, both these molecules act as second messengers. Inositol 1,4,5-trisphosphate binds specific receptors in the endoplasmatic reticulum, resulting in the opening of Ca^{2+} channels and release of Ca^{2+} in the cytoplasm.

No clear functions have been assigned to the GPCRs. The expression level of GPR3 and of GPR12 is increased in human umbilical vein endothelial cells after shear stress (Uhlenbrock et al., 2003). Since sphingosine-1-phosphate is a ligand for GPR3 and GPR12, the above data suggest a role for both GPCRs in sphingosine-1-phosphate-mediated intracellular signaling in human endothelial cells. As the expression of GPR3 and GPR6 is also differentially regulated in rodent obesity models, both GPCRs (+GPR12) are considered as putative drug targets in appetite, hunger and satiety control. GPR12, on the other hand, seems to be involved in the differentiation and maturation of post mitotic neurons (Ignatov et al., 2003).

References: Annaert, W. and B. De Strooper (2002). "A cell biological perspective on Alzheimer's disease." *Annu Rev Cell Dev Biol* 18: 25-51.

Gotz, J., F. Chen, et al. (2001). "Formation of neurofibrillary tangles in P3011 tau transgenic mice induced by Abeta 42 fibrils." *Science* 293(5534): 1491-5.

Ignatov, A.; Lintzel, J.; Hermans-Borgmeyer, I.; Kreienkamp, H-J., Joost, P.; Thomsen, S.; Methner, A. And Schaller, H.C. (2003). Role of the G-protein-coupled receptor GPR12 as high-affinity receptor for sphingosylphosphorylcholine and its expression and function in brain development. *J. Neurosci.* 23, 3: 907-914.

Lipinski, C. A., Lombardo, F., Dominy, B. W., and Feeney, P. J. Adv. Drug. Deliv. Rev., 23, 3-25, 1997

Marchese, A.; Docherty, JM.; Nguyen, T.; Heiber, M.; Cheng, R.; Heng, HH.; Tsui, LC.; Shi, X.; George SR. and O'Dowd, BF. (1994). Cloning of human genes encoding novel G protein-coupled receptors. Genomics, 23, 3: 609-618.
Marinissen, M. J. and J. S. Gutkind (2001). "G-protein-coupled receptors and signaling networks: emerging paradigms." Trends Pharmacol Sci 22(7): 368-76.

Ritchie, K. and S. Lovestone (2002). "The dementias." Lancet 360(9347): 1759-66.

Uhlenbrock, K.; Gassenhuber, H. And Kostenis, E. (2002). Sphingosine-1-phosphate is a ligand of the human GPR3, GPR6 and GPR12 family of constitutively active G protein-coupled receptors. Cell Signal, 14, 11: 941-953.

Uhlenbrock, K.; Huber, J.; Ardati, A.; Bush, AE. And Kostenis, E. (2003). Fluid shear stress differentially regulates GPR3, GPR6 and GPR12 expression in human umbilical vein endothelial cells. Cell Physiol. Biochem. 13, 2: 75-84.

Wess, J. (1998). "Molecular basis of receptor/G-protein-coupling selectivity." Pharmacol Ther 80(3): 231-64.

Applicants' Invention Based on GPCR Relationship to Amyloid Beta Peptides

As noted above, the present invention is based on the present inventors' discovery that the G-protein coupled receptor(s) ("GPCR(s)") are factors in the up-regulation and/or induction of amyloid beta precursor processing in mammalian, and principally, neuronal cells, and that the inhibition of the function of such polypeptides is effective in reducing levels of amyloid beta protein peptides.

The present inventors are unaware of any prior knowledge linking GPCRs, and more particularly GPR3, and amyloid beta peptide formation and secretion. As discussed in more detail in the Experimental section below, the present inventors demonstrate that the overexpression of GPR3 increases, and the knockdown of GPR3 reduces, amyloid beta 1-42 in the conditioned medium of transduced cells. The present invention is based on these findings and the recognition that the GPCRs are putative drug targets for Alzheimer's

disease, since the predominant expression of GPR3, GPR6 and GPR12 is in the tissue of the central nervous system.

One aspect of the present invention is a method based on the aforesaid discovery for identifying a compound that inhibits the processing of amyloid-beta precursor protein in a mammalian cell, and may therefore be useful in reducing amyloid beta peptide levels in a subject. The present method comprises contacting a drug candidate compound with a GPCR polypeptide, or a fragment of said polypeptide, and measuring a compound-polypeptide property related to the production of amyloid-beta protein. The "compound-polypeptide property" is a measurable phenomenon chosen by the person of ordinary skill in the art, and based on the recognition that GPCR activation and deactivation is a causative factor in the activation and deactivation, respectively, of amyloid beta protein precursor processing, and an increase and decrease, respectively, of amyloid beta peptide levels. The measurable property may range from the binding affinity for a peptide domain of the GPCR polypeptide, to the level of any one of a number of "second messenger" levels resulting from the activation or deactivation of the GPCR, to a reporter molecule property directly linked to the aforesaid second messenger, and finally to the level of amyloid beta peptide secreted by the mammalian cell contacted with the compound.

Depending on the choice of the skilled artisan, the present assay method may be designed to function as a series of measurements, each of which is designed to determine whether the drug candidate compound is indeed acting on the GPCR to amyloid beta peptide pathway. For example, an assay designed to determine the binding affinity of a compound to the GPCR, or fragment thereof, may be necessary, but not sufficient, to ascertain whether the test compound would be useful for reducing amyloid beta peptide levels when administered to a subject. Nonetheless, such binding information would be useful in identifying a set of test compounds for use in an assay that would measure a different property, further down the biochemical pathway. Such second assay may be designed to confirm that the test compound, having binding affinity for a GPCR peptide, actually down-regulates or inhibits, as an agonist or inverse agonist, GPCR function in a mammalian cell. This further assay may measure a second messenger that is a direct consequence of the activation or deactivation of the GPCR, or a synthetic reporter system responding to the messenger. Measuring a different second messenger, and/or confirming that the assay system itself is not being affected directly and not the GPCR pathway may further validate the assay. In this latter regard, suitable controls should always be in place to insure against false positive readings.

The order of taking these measurements is not believed to be critical to the practice of the present invention, which may be practiced in any order. For example, one may first perform a screening assay of a set of compounds for which no information is known respecting the compounds' binding affinity for GPCR. Alternatively, one may screen a set of compounds identified as having binding affinity for a GPCR peptide domain, or a class of compounds identified as being agonist or inverse agonists of a GPCR. It is not essential to know the binding affinity for GPCR due to the possible compound interaction in the intra-membrane domain of the GPCR polypeptide, which domain conformation may not be possible to reproduce in an affinity experiment. However, for the present assay to be meaningful to the ultimate use of the drug candidate compounds, a measurement of the second messenger(s), or the ultimate amyloid beta peptide levels, is necessary. Validation studies including controls, and measurements of binding affinity to GPCR are nonetheless useful in identifying a compound useful in any therapeutic or diagnostic application.

The present assay method may be practiced *in vitro*, using one or more of the GPCR proteins, or fragments thereof, or membrane preparations made from cells transduced with vectors over-expressing the GPCR polypeptides. The amino acid sequences of the GPCRs, and useful fragments thereof are found in SEQ ID NO: 4-6, 289-333. The binding affinity of the compound with the polypeptide can be measured by methods known in the art, such as using surface plasmon resonance biosensors (Biacore), by saturation binding analysis with a labeled compound (e.g. Scatchard and Lindmo analysis), by differential UV spectrophotometer, fluorescence polarization assay, Fluorometric Imaging Plate Reader (FLIPR[®]) system, Fluorescence resonance energy transfer, and Bioluminescence resonance energy transfer. The binding affinity of compounds can also be expressed in dissociation constant (K_d) or as IC_{50} or EC_{50} . The IC_{50} represents the concentration of a compound that is required for 50% inhibition of binding of another ligand to the polypeptide. The EC_{50} represents the concentration required for obtaining 50% of the maximum effect in any assay that measures receptor function. The dissociation constant, K_d , is a measure of how well a ligand binds to the polypeptide, it is equivalent to the ligand concentration required to saturate exactly half of the binding-sites on the polypeptide. Compounds with a high affinity binding have low K_d , IC_{50} and EC_{50} values, i.e. in the range of 100 nM to 1 pM; a moderate to low affinity binding relates to a high K_d , IC_{50} and EC_{50} values, i.e. in the micromolar range.

The present assay method may also be practiced in a cellular assay. A host cell expressing a GPCR polypeptide can be a cell with endogenous expression of the polypeptide or a cell over-expressing the polypeptide e.g. by transduction. When the endogenous expression of the polypeptide is not sufficient to determine a baseline that can easily be measured, one may use using host cells that over express GPCR. Overexpression has the advantage that the level of the second messenger is higher than the activity level by endogenous expression. Accordingly, measuring such levels using presently available techniques is easier. In such cellular assay, the biological activity of the GPCR may be measured using a second messenger, such as cyclic AMP or Ca^{2+} , cyclic GMP, inositol triphosphate (IP_3) and/or diacylglycerol (DAG). Cyclic AMP or Ca^{2+} are preferred second messengers to measure. Second messenger activation may be measured by several different techniques, either directly by ELISA or radioactive technologies or indirectly by reporter gene analysis, discussed below. Preferably the method further comprises contacting the host cell with an agonist for GPCR before determining the baseline level. The addition of an agonist further stimulates GPCR, thereby further increasing the activity level of the second messenger. Several such agonists (ligands) are known in the art; preferentially the agonist is spingosine-1-phosphate or dihydrosphingosine-1-phosphate. The GPCR polypeptides, when over expressed or activated the level of secreted amyloid beta peptides.

The present invention further relates to a method for identifying a compound that inhibits amyloid-beta precursor protein processing in a mammalian cell comprising:

- (a) contacting a compound with a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6,
- (b) determining the binding affinity of the compound to the polypeptide,
- (c) contacting a population of mammalian cells expressing said polypeptide with the compound that exhibits a binding affinity of at least 10 micromolar, and
- (d) identifying the compound that inhibits the amyloid-beta precursor protein processing in the cells.

A further embodiment of the present invention relates a method to identify a compound that inhibits the amyloid-beta precursor protein processing in a cell, wherein the activity level of the GPCR polypeptide is measured by determining the level of one or more second messengers, wherein the level of the one or second messenger is determined with a reporter controlled by a promoter, which is responsive to the second messenger. The reporter

is a reporter gene under the regulation of a promoter that responds to the cellular level of second messengers. Such preferred second messengers are Cyclic AMP or Ca^{2+} . The reporter gene should have a gene product that is easily detected, and that may be stably infected in the host cell. Such methods are well known by any person with ordinary skill in the art.

The reporter gene may be selected from alkaline phosphatase, green fluorescent protein (GFP), enhanced green fluorescent protein (eGFP), destabilized green fluorescent protein (dGFP), luciferase, beta-galactosidase among others. The reporter is preferably luciferase or beta-galactosidase, which are readily available and easy to measure over a large range. The promoter in the reporter construct is preferably a cyclic AMP-responsive promoter, an NF-KB responsive promoter, or a NF-AT responsive promoter. The cyclic-AMP responsive promoter is responsive to the cyclic-AMP levels in the cell. The NF-AT responsive promoter is sensitive to cytoplasmic Ca^{2+} -levels in the cell. The NF-KB responsive promoter is sensitive for activated NF- κ B levels in the cell.

A further embodiment of the present invention relates a method to identify a compound that inhibits the amyloid-beta precursor protein processing in a cell, wherein the activity level of the GPCR polypeptide is measured by determining the level of amyloid beta peptides. The levels of these peptides may be measured with specific ELISAs using antibodies specifically recognizing the different amyloid beta peptide species (see e.g. Example 1). Secretion of the various amyloid beta peptides may also be measured using antibodies that bind all peptides. Levels of amyloid beta peptides can also be measured by Mass spectrometry analysis.

For high-throughput purposes, libraries of compounds may be used such as antibody fragment libraries, peptide phage display libraries, peptide libraries (e.g. LOPAPTM, Sigma Aldrich), lipid libraries (BioMol), synthetic compound libraries (e.g. LOPACTM, Sigma Aldrich) or natural compound libraries (Specs, TimTec).

Preferred drug candidate compounds are low molecular weight compounds. Low molecular weight compounds, i.e. with a molecular weight of 500 Dalton or less, are likely to have good absorption and permeation in biological systems and are consequently more likely to be successful drug candidates than compounds with a molecular weight above 500 Dalton (Lipinski et al. (1997)). Peptides comprise another preferred class of drug candidate compounds, since peptides are known GPCRs antagonists. Peptides may be excellent drug

candidates and there are multiple examples of commercially valuable peptides such as fertility hormones and platelet aggregation inhibitors. Natural compounds are another preferred class of drug candidate compound. Such compounds are found in and extracted from natural sources, and which may thereafter be synthesized. The lipids are another preferred class of drug candidate compound. Lipids may be antagonists of the GPCRs listed in Table 2 (SEQ ID NO: 1-3, 4-6).

Another preferred class of drug candidate compounds is an antibody. The present invention also provides antibodies directed against the extracellular domains of the GPCR. These antibodies should specifically bind to one or more of the extra-cellular domains of the GPCRs, or as described further below, engineered to be endogenously produced to bind to the intra-cellular GPCR domain. These antibodies may be monoclonal antibodies or polyclonal antibodies. The present invention includes chimeric, single chain, and humanized antibodies, as well as FAb fragments and the products of a FAb expression library, and Fv fragments and the products of an Fv expression library.

In certain embodiments, polyclonal antibodies may be used in the practice of the invention. The skilled artisan knows methods of preparing polyclonal antibodies. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. Antibodies may also be generated against the intact GPCR protein or polypeptide, or against a fragment such as its extracellular domain peptides, derivatives including conjugates, or other epitope of the GPCR protein or polypeptide, such as the GPCR embedded in a cellular membrane, or a library of antibody variable regions, such as a phage display library.

It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants that may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). One skilled in the art without undue experimentation may select the immunization protocol.

In some embodiments, the antibodies may be monoclonal antibodies. Monoclonal antibodies may be prepared using methods known in the art. The monoclonal antibodies of the present invention may be "humanized" to prevent the host from mounting an immune response to the antibodies. A "humanized antibody" is one in which the complementarity
5 determining regions (CDRs) and/or other portions of the light and/or heavy variable domain framework are derived from a non-human immunoglobulin, but the remaining portions of the molecule are derived from one or more human immunoglobulins. Humanized antibodies also include antibodies characterized by a humanized heavy chain associated with a donor or acceptor unmodified light chain or a chimeric light chain, or vice versa. The humanization of
10 antibodies may be accomplished by methods known in the art (see, e.g. Mark and Padlan, (1994) "Chapter 4. Humanization of Monoclonal Antibodies", The Handbook of Experimental Pharmacology Vol. 113, Springer-Verlag, New York). Transgenic animals may be used to express humanized antibodies.

Human antibodies can also be produced using various techniques known in the art,
15 including phage display libraries (Hoogenboom and Winter, (1991) J. Mol. Biol. 227:381-8; Marks et al. (1991). J. Mol. Biol. 222:581-97). The techniques of Cole, et al. and Boerner, et al. are also available for the preparation of human monoclonal antibodies (Cole, et al. (1985) Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77; Boerner, et al (1991). J. Immunol., 147(1):86-95).

20 Techniques known in the art for the production of single chain antibodies can be adapted to produce single chain antibodies to the GPCR polypeptides and proteins of the present invention. The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The
25 heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain cross-linking. Alternatively; the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent cross-linking.

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens and preferably for a cell-surface
30 protein or receptor or receptor subunit. In the present case, one of the binding specificities is for one extracellular domain of the GPCR, the other one is for another extracellular domain of the same or different GPCR.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, (1983) Nature 305:537-9). Because of the random
5 assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. Affinity chromatography steps usually accomplish the purification of the correct molecule. Similar procedures are disclosed in Traunecker, et al. (1991) EMBO J. 10:3655-9.

10 According to another preferred embodiment, the assay method comprise using a drug candidate compound identified as having a binding affinity for GPCRs, and/or has already been identified as having down-regulating activity such as antagonist or inverse agonist activity vis-à-vis one or more GPCR. Examples of such compounds are the aryloxydithiourea compounds disclosed in US 6,420,563 (WO 01/62765), hereby
15 incorporated by reference with respect to the active inverse agonists disclosed therein.

Another aspect of the present invention relates to a method for reducing amyloid-beta precursor protein processing in a mammalian cell, comprising by contacting said cell with an expression-inhibiting agent that inhibits the translation in the cell of a polyribonucleotide encoding a GPCR polypeptide. A particular embodiment relates to a composition comprising
20 an polynucleotide including at least one antisense strand that functions to pair the agent with the target GPCR mRNA, and thereby down-regulate or block the expression of GPCR polypeptide. The inhibitory agent preferably comprises antisense polynucleotide, a ribozyme, and a small interfering RNA (siRNA), wherein said agent comprises a nucleic acid sequence complementary to, or engineered from, a naturally occurring polynucleotide
25 sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6.

A special embodiment of the present invention relates to a method wherein the expression-inhibiting agent is selected from the group consisting of antisense RNA, antisense oligodeoxynucleotide (ODN), a ribozyme that cleaves the polyribonucleotide coding for SEQ
30 ID NO: 4-6, a small interfering RNA (siRNA) that is sufficiently homologous to a portion of the polyribonucleotide corresponding to SEQ ID NO: 4-6 such that the siRNA interferes with the translation of the GPCR polyribonucleotide to the GPCR polypeptide.

Another embodiment of the present invention relates to a method wherein the expression-inhibiting agent is a nucleic acid expressing the antisense RNA, antisense oligodeoxynucleotide (ODN), a ribozyme that cleaves the polyribonucleotide coding for SEQ ID NO: 4-6, a small interfering RNA (siRNA) that is sufficiently homologous to a portion of the polyribonucleotide corresponding to SEQ ID NO: 4-6 such that the siRNA interferes with the translation of the GPCR polyribonucleotide to the GPCR polypeptide. Preferably the expression-inhibiting agent is an antisense RNA, ribozyme, antisense oligodeoxynucleotide, or siRNA comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 7-287.

The down regulation of gene expression using antisense nucleic acids can be achieved at the translational or transcriptional level. Antisense nucleic acids of the invention are preferably nucleic acid fragments capable of specifically hybridizing with all or part of a nucleic acid encoding a GPCR polypeptide or the corresponding messenger RNA. In addition, antisense nucleic acids may be designed which decrease expression of the nucleic acid sequence capable of encoding a GPCR polypeptide by inhibiting splicing of its primary transcript. Any length of antisense sequence is suitable for practice of the invention so long as it is capable of down-regulating or blocking expression of a nucleic acid coding for a GPCR. Preferably, the antisense sequence is at least about 17 nucleotides in length. The preparation and use of antisense nucleic acids, DNA encoding antisense RNAs and the use of oligo and genetic antisense is known in the art.

One embodiment of expression-inhibitory agent is a nucleic acid that is antisense to a nucleic acid comprising SEQ ID NO: 1-3. For example, an antisense nucleic acid (e.g. DNA) may be introduced into cells *in vitro*, or administered to a subject *in vivo*, as gene therapy to inhibit cellular expression of nucleic acids comprising SEQ ID NO: 1-3. Antisense oligonucleotides preferably comprise a sequence containing from about 17 to about 100 nucleotides and more preferably the antisense oligonucleotides comprise from about 18 to about 30 nucleotides. Antisense nucleic acids may be prepared from about 10 to about 30 contiguous nucleotides selected from the sequences of SEQ ID NO: 1-3, expressed in the opposite orientation.

The antisense nucleic acids are preferably oligonucleotides and may consist entirely of deoxyribo-nucleotides, modified deoxyribonucleotides, or some combination of both. The antisense nucleic acids can be synthetic oligonucleotides. The oligonucleotides may be chemically modified, if desired, to improve stability and/or selectivity. Since

oligonucleotides are susceptible to degradation by intracellular nucleases, the modifications can include, for example, the use of a sulfur group to replace the free oxygen of the phosphodiester bond. This modification is called a phosphorothioate linkage. Phosphorothioate antisense oligonucleotides are water soluble, polyanionic, and resistant to
5 endogenous nucleases. In addition, when a phosphorothioate antisense oligonucleotide hybridizes to its target site, the RNA-DNA duplex activates the endogenous enzyme ribonuclease (RNase) H, which cleaves the mRNA component of the hybrid molecule.

In addition, antisense oligonucleotides with phosphoramidite and polyamide (peptide) linkages can be synthesized. These molecules should be very resistant to nuclease
10 degradation. Furthermore, chemical groups can be added to the 2' carbon of the sugar moiety and the 5 carbon (C-5) of pyrimidines to enhance stability and facilitate the binding of the antisense oligonucleotide to its target site. Modifications may include 2'-deoxy, O-pentoxy, O-propoxy, O-methoxy, fluoro, methoxyethoxy phosphorothioates, modified bases, as well as other modifications known to those of skill in the art.

Another type of expression-inhibitory agent that reduces the levels of GPCRs are
15 ribozymes. Ribozymes are catalytic RNA molecules (RNA enzymes) that have separate catalytic and substrate binding domains. The substrate binding sequence combines by nucleotide complementarity and, possibly, non-hydrogen bond interactions with its target sequence. The catalytic portion cleaves the target RNA at a specific site. The substrate
20 domain of a ribozyme can be engineered to direct it to a specified mRNA sequence. The ribozyme recognizes and then binds a target mRNA through complementary base-pairing. Once it is bound to the correct target site, the ribozyme acts enzymatically to cut the target mRNA. Cleavage of the mRNA by a ribozyme destroys its ability to direct synthesis of the corresponding polypeptide. Once the ribozyme has cleaved its target sequence, it is released
25 and can repeatedly bind and cleave at other mRNAs.

Ribozyme forms include a hammerhead motif, a hairpin motif, a hepatitis delta virus, group I intron or RNaseP RNA (in association with an RNA guide sequence) motif or Neurospora VS RNA motif. Ribozymes possessing a hammerhead or hairpin structure are readily prepared since these catalytic RNA molecules can be expressed within cells from
30 eukaryotic promoters (Chen, et al. (1992) *Nucleic Acids Res.* 20:4581-9). A ribozyme of the present invention can be expressed in eukaryotic cells from the appropriate DNA vector. If desired, the activity of the ribozyme may be augmented by its release from the primary transcript by a second ribozyme (Ventura, et al. (1993) *Nucleic Acids Res.* 21:3249-55).

Ribozymes may be chemically synthesized by combining an oligodeoxyribonucleotide with a ribozyme catalytic domain (20 nucleotides) flanked by sequences that hybridize to the target mRNA after transcription. The oligodeoxyribonucleotide is amplified by using the substrate binding sequences as primers.

5 The amplification product is cloned into a eukaryotic expression vector.

Ribozymes are expressed from transcription units inserted into DNA, RNA, or viral vectors. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol (I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the
10 levels of a given pol II promoter in a given cell type will depend on nearby gene regulatory sequences. Prokaryotic RNA polymerase promoters are also used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Gao and Huang, (1993) Nucleic Acids Res. 21:2867-72). It has been demonstrated that ribozymes expressed from these promoters can function in mammalian cells (Kashani-Sabet, et al. (1992)
15 Antisense Res. Dev. 2:3-15).

A particularly preferred inhibitory agent is a small interfering RNA (siRNA). siRNAs mediate the post-transcriptional process of gene silencing by double stranded RNA (dsRNA) that is homologous in sequence to the silenced RNA. siRNA according to the present invention comprises a sense strand of 17-25 nucleotides complementary or
20 homologous to a contiguous 17-25 nucleotide sequence selected from the group of sequences described in SEQ ID NO: 1-3 and an antisense strand of 17-23 nucleotides complementary to the sense strand. The most preferred siRNA comprises sense and anti-sense strands that are 100 per cent complementary to each other and the target polynucleotide sequence. Preferably the siRNA further comprises a loop region linking the sense and the antisense strand.

25 A self-complementing single stranded siRNA molecule polynucleotide according to the present invention comprises a sense portion and an antisense portion connected by a loop region linker. Preferably, the loop region sequence is 4-30 nucleotides long, more preferably 5-15 nucleotides long and most preferably 8 nucleotides long. In a most preferred embodiment the linker sequence is UUGCUAUA (SEQ ID NO: 288). Self-complementary
30 single stranded siRNAs form hairpin loops and are more stable than ordinary dsRNA. In addition, they are more easily produced from vectors.

Analogous to antisense RNA, the siRNA can be modified to confer resistance to nucleolytic degradation, or to enhance activity, or to enhance cellular distribution, or to enhance cellular uptake, such modifications may consist of modified internucleoside linkages, modified nucleic acid bases, modified sugars and/or chemical linkage the siRNA to one or more moieties or conjugates. The nucleotide sequences are selected according to siRNA designing rules that give an improved reduction of the target sequences compared to nucleotide sequences that do not comply with these siRNA designing rules (For a discussion of these rules and examples of the preparation of siRNA, WO2004094636, published November 4, 2004, and UA20030198627, are hereby incorporated by reference.

The present invention also relates to compositions, and methods using said compositions, comprising a DNA expression vector capable of expressing a polynucleotide capable of inhibiting amyloid beta protein precursor processing and described hereinabove as an expression inhibition agent.

A special aspect of these compositions and methods relates to the down-regulation or blocking of the expression of a GPCR polypeptide by the induced expression of a polynucleotide encoding an intracellular binding protein that is capable of selectively interacting with the GPCR polypeptide. An intracellular binding protein includes any protein capable of selectively interacting, or binding, with the polypeptide in the cell in which it is expressed and neutralizing the function of the polypeptide. Preferably, the intracellular binding protein is a neutralizing antibody or a fragment of a neutralizing antibody having binding affinity to an intra-cellular domain of the GPCR polypeptide of SEQ ID NO: 4-6. More preferably, the intracellular binding protein is a single chain antibody.

A special embodiment of this composition comprises the expression-inhibiting agent selected from the group consisting of antisense RNA, antisense oligodeoxynucleotide (ODN), a ribozyme that cleaves the polyribonucleotide coding for SEQ ID NO: 4-6, and a small interfering RNA (siRNA) that is sufficiently homologous to a portion of the polyribonucleotide corresponding to SEQ ID NO: 4-6 such that the siRNA interferes with the translation of the GPCR polyribonucleotide to the GPCR polypeptide,

The polynucleotide expressing the expression-inhibiting agent or the encoding an intracellular binding protein is preferably included within a vector. The polynucleic acid is operably linked to signals enabling expression of the nucleic acid sequence and is introduced into a cell utilizing, preferably, recombinant vector constructs, which will express the

antisense nucleic acid once the vector is introduced into the cell. A variety of viral-based systems are available, including adenoviral, retroviral, adeno-associated viral, lentiviral, herpes simplex viral or a sendaviral vector systems, and all may be used to introduce and express polynucleotide sequence for the expression-inhibiting agents in target cells.

5 Preferably, the viral vectors used in the methods of the present invention are replication defective. Such replication defective vectors will usually lack at least one region that is necessary for the replication of the virus in the infected cell. These regions can either be eliminated (in whole or in part), or be rendered non-functional by any technique known to a person skilled in the art. These techniques include the total removal, substitution, partial
10 deletion or addition of one or more bases to an essential (for replication) region. Such techniques may be performed in vitro (on the isolated DNA) or in situ, using the techniques of genetic manipulation or by treatment with mutagenic agents. Preferably, the replication defective virus retains the sequences of its genome, which are necessary for encapsidating, the viral particles.

15 In a preferred embodiment, the viral element is derived from an adenovirus. Preferably, the vehicle includes an adenoviral vector packaged into an adenoviral capsid, or a functional part, derivative, and/or analogue thereof. Adenovirus biology is also comparatively well known on the molecular level. Many tools for adenoviral vectors have been and continue to be developed, thus making an adenoviral capsid a preferred vehicle for
20 incorporating in a library of the invention. An adenovirus is capable of infecting a wide variety of cells. However, different adenoviral serotypes have different preferences for cells. To combine and widen the target cell population that an adenoviral capsid of the invention can enter in a preferred embodiment, the vehicle includes adenoviral fiber proteins from at least two adenoviruses.

25 In a preferred embodiment, the nucleic acid derived from an adenovirus includes the nucleic acid encoding an adenoviral late protein or a functional part, derivative, and/or analogue thereof. An adenoviral late protein, for instance an adenoviral fiber protein, may be favorably used to target the vehicle to a certain cell or to induce enhanced delivery of the vehicle to the cell. Preferably, the nucleic acid derived from an adenovirus encodes for
30 essentially all adenoviral late proteins, enabling the formation of entire adenoviral capsids or functional parts, analogues, and/or derivatives thereof. Preferably, the nucleic acid derived from an adenovirus includes the nucleic acid encoding adenovirus E2A or a functional part, derivative, and/or analogue thereof. Preferably, the nucleic acid derived from an adenovirus

includes the nucleic acid encoding at least one E4-region protein or a functional part, derivative, and/or analogue thereof, which facilitates, at least in part, replication of an adenoviral derived nucleic acid in a cell. The adenoviral vectors used in the examples of this application are exemplary of the vectors useful in the present method of treatment invention.

5 Certain embodiments of the present invention use retroviral vector systems. Retroviruses are integrating viruses that infect dividing cells, and their construction is known in the art. Retroviral vectors can be constructed from different types of retrovirus, such as, MoMuLV ("murine Moloney leukemia virus" MSV ("murine Moloney sarcoma virus"), HaSV ("Harvey sarcoma virus"); SNV ("spleen necrosis virus"); RSV ("Rous sarcoma
10 virus") and Friend virus. Lentiviral vector systems may also be used in the practice of the present invention. Retroviral systems and herpes virus system may be preferred vehicles for transfection of neuronal cells.

In other embodiments of the present invention, adeno-associated viruses ("AAV") are utilized. The AAV viruses are DNA viruses of relatively small size that integrate, in a stable
15 and site-specific manner, into the genome of the infected cells. They are able to infect a wide spectrum of cells without inducing any effects on cellular growth, morphology or differentiation, and they do not appear to be involved in human pathologies.

In the vector construction, the polynucleotide agents of the present invention may be linked to one or more regulatory regions. Selection of the appropriate regulatory region or
20 regions is a routine matter, within the level of ordinary skill in the art. Regulatory regions include promoters, and may include enhancers, suppressors, etc.

Promoters that may be used in the expression vectors of the present invention include both constitutive promoters and regulated (inducible) promoters. The promoters may be prokaryotic or eukaryotic depending on the host. Among the prokaryotic (including
25 bacteriophage) promoters useful for practice of this invention are lac, lacZ, T3, T7, lambda P.sub.r, P.sub.1, and trp promoters. Among the eukaryotic (including viral) promoters useful for practice of this invention are ubiquitous promoters (e.g. HPRT, vimentin, actin, tubulin), intermediate filament promoters (e.g. desmin, neurofilaments, keratin, GFAP), therapeutic gene promoters (e.g. MDR type, CFTR, factor VIII), tissue-specific promoters (e.g. actin
30 promoter in smooth muscle cells, or Flt and Flk promoters active in endothelial cells), including animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in

pancreatic acinar cells (Swift, et al. (1984) Cell 38:639-46; Ornitz, et al. (1986) Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, (1987) Hepatology 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, (1985) Nature 315:115-22), immunoglobulin gene control region which is active in lymphoid cells
 5 (Grosschedl, et al. (1984) Cell 38:647-58; Adames, et al. (1985) Nature 318:533-8; Alexander, et al. (1987) Mol. Cell. Biol. 7:1436-44), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder, et al. (1986) Cell 45:485-95), albumin gene control region which is active in liver (Pinkert, et al. (1987) Genes and Devel. 1:268-76), alpha-fetoprotein gene control region which is active in liver
 10 (Krumlauf, et al. (1985) Mol. Cell. Biol., 5:1639-48; Hammer, et al. (1987) Science 235:53-8), alpha 1-antitrypsin gene control region which is active in the liver (Kelsey, et al. (1987) Genes and Devel., 1: 161-71), beta-globin gene control region which is active in myeloid cells (Mogam, et al. (1985) Nature 315:338-40; Kollias, et al. (1986) Cell 46:89-94), myelin basic protein gene control region which is active in oligodendrocyte cells in the
 15 brain (Readhead, et al. (1987) Cell 48:703-12), myosin light chain-2 gene control region which is active in skeletal muscle (Sani, (1985) Nature 314:283-6), and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason, et al. (1986) Science 234:1372-8).

Other promoters which may be used in the practice of the invention include promoters
 20 which are preferentially activated in dividing cells, promoters which respond to a stimulus (e.g. steroid hormone receptor, retinoic acid receptor), tetracycline-regulated transcriptional modulators, cytomegalovirus immediate-early, retroviral LTR, metallothionein, SV-40, E1a, and MLP promoters.

The vectors may also include other elements, such as enhancers, repressor systems,,
 25 and localization signals. A membrane localization signal is a preferred element when expressing a sequence encoding an intracellular binding protein, which functions by contacting the intracellular domain of the GPCR and is most effective when the vector product is directed to the inner surface of the cellular membrane, where its target resides. Membrane localization signals are well known to persons skilled in the art. For example, a
 30 membrane localization domain suitable for localizing a polypeptide to the plasma membrane is the C-terminal sequence CaaX for farnesylation (where "a" is an aliphatic amino acid residue, and "X" is any amino acid residue, generally leucine), for example, Cysteine-Alanine-Alanine-Leucine, or Cysteine-Isoleucine-Valine-Methionine. Other membrane

localization signals include the putative membrane localization sequence from the C-terminus of Bcl-2 or the C-terminus of other members of the Bcl-2 family of proteins.

Additional vector systems include the non-viral systems that facilitate introduction of polynucleotide agents into a patient. For example, a DNA vector encoding a desired
5 sequence can be introduced in vivo by lipofection. Synthetic cationic lipids designed to limit the difficulties encountered with liposome-mediated transfection can be used to prepare liposomes for in vivo transfection of a gene encoding a marker (Felgner, et. al. (1987) Proc. Natl. Acad. Sci. USA 84:7413-7); see Mackey, et al. (1988) Proc. Natl. Acad. Sci. USA 85:8027-31; Ulmer, et al. (1993) Science 259:1745-8). The use of cationic lipids may
10 promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Felgner and Ringold, (1989) Nature 337:387-8). Particularly useful lipid compounds and compositions for transfer of nucleic acids are described in International Patent Publications WO 95/18863 and WO 96/17823, and in U.S. Pat. No. 5,459,127. The use of lipofection to introduce exogenous genes into the specific
15 organs in vivo has certain practical advantages and directing transfection to particular cell types would be particularly advantageous in a tissue with cellular heterogeneity, for example, pancreas, liver, kidney, and the brain. Lipids may be chemically coupled to other molecules for the purpose of targeting. Targeted peptides, e.g., hormones or neurotransmitters, and proteins for example, antibodies, or non-peptide molecules could be coupled to liposomes
20 chemically. Other molecules are also useful for facilitating transfection of a nucleic acid in vivo, for example, a cationic oligopeptide (e.g., International Patent Publication WO 95/21931), peptides derived from DNA binding proteins (e.g., International Patent Publication WO 96/25508), or a cationic polymer (e.g., International Patent Publication WO 95/21931).

25 It is also possible to introduce a DNA vector in vivo as a naked DNA plasmid (see U.S. Pat. Nos. 5,693,622, 5,589,466 and 5,580,859). Naked DNA vectors for therapeutic purposes can be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun, or use of a DNA vector transporter (see,
30 e.g., Wilson, et al. (1992) J. Biol. Chem. 267:963-7; Wu and Wu, (1988) J. Biol. Chem. 263:14621-4; Hartmut, et al. Canadian Patent Application No. 2,012,311, filed Mar. 15, 1990; Williams, et al (1991). Proc. Natl. Acad. Sci. USA 88:2726-30). Receptor-mediated

DNA delivery approaches can also be used (Curiel, et al. (1992) Hum. Gene Ther. 3:147-54; Wu and Wu, (1987) J. Biol. Chem. 262:4429-32).

The present invention also provides biologically compatible compositions comprising the compounds identified as antagonists and/or inverse agonists of GPCR, and the expression-inhibiting agents as described hereinabove.

A biologically compatible composition is a composition, that may be solid, liquid, gel, or other form, in which the compound, polynucleotide, vector, and antibody of the invention is maintained in an active form, e.g., in a form able to effect a biological activity. For example, a compound of the invention would have inverse agonist or antagonist activity on the GPCR; a nucleic acid would be able to replicate, translate a message, or hybridize to a complementary mRNA of a GPCR; a vector would be able to transfect a target cell and expression the antisense, antibody, ribozyme or siRNA as described hereinabove; an antibody would bind a GPCR polypeptide domain.

A preferred biologically compatible composition is an aqueous solution that is buffered using, e.g., Tris, phosphate, or HEPES buffer, containing salt ions. Usually the concentration of salt ions will be similar to physiological levels. Biologically compatible solutions may include stabilizing agents and preservatives. In a more preferred embodiment, the biocompatible composition is a pharmaceutically acceptable composition. Such compositions can be formulated for administration by topical, oral, parenteral, intranasal, subcutaneous, and intraocular, routes. Parenteral administration is meant to include intravenous injection, intramuscular injection, intraarterial injection or infusion techniques. The composition may be administered parenterally in dosage unit formulations containing standard, well known non-toxic physiologically acceptable carriers, adjuvants and vehicles as desired.

A particularly preferred embodiment of the present composition invention is a cognitive-enhancing pharmaceutical composition comprising a therapeutically effective amount of an expression-inhibiting agent as described hereinabove, in admixture with a pharmaceutically acceptable carrier. Another preferred embodiment is a pharmaceutical composition for the treatment or prevention of a condition involving cognitive impairment or a susceptibility to the condition, comprising an effective amyloid beta peptide inhibiting amount of a GPCR antagonist or inverse agonist its pharmaceutically acceptable salts, hydrates, solvates, or prodrugs thereof in admixture with a pharmaceutically acceptable

carrier. A particularly preferred class of such compositions comprise an aryloxydithiourea compound.

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient. Pharmaceutical compositions for oral use can be prepared by combining active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethyl-cellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate. Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations that can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Preferred sterile injectable preparations can be a solution or suspension in a non-toxic parenterally acceptable solvent or diluent. Examples of pharmaceutically acceptable carriers are saline, buffered saline, isotonic saline (e.g. monosodium or disodium phosphate, sodium, potassium; calcium or magnesium chloride, or mixtures of such salts), Ringer's solution, dextrose, water, sterile water, glycerol, ethanol, and combinations thereof 1,3-butanediol and sterile fixed oils are conveniently employed as solvents or suspending media. Any bland

fixed oil can be employed including synthetic mono- or di-glycerides. Fatty acids such as oleic acid also find use in the preparation of injectables.

The composition medium can also be a hydrogel, which is prepared from any biocompatible or non-cytotoxic homo- or hetero-polymer, such as a hydrophilic polyacrylic acid polymer that can act as a drug absorbing sponge. Certain of them, such as, in particular, those obtained from ethylene and/or propylene oxide are commercially available. A hydrogel can be deposited directly onto the surface of the tissue to be treated, for example during surgical intervention.

Embodiments of pharmaceutical compositions of the present invention comprise a replication defective recombinant viral vector encoding the polynucleotide inhibitory agent of the present invention and a transfection enhancer, such as poloxamer. An example of a poloxamer is Poloxamer 407, which is commercially available (BASF, Parsippany, N.J.) and is a non-toxic, biocompatible polyol. A poloxamer impregnated with recombinant viruses may be deposited directly on the surface of the tissue to be treated, for example during a surgical intervention. Poloxamer possesses essentially the same advantages as hydrogel while having a lower viscosity.

The active expression-inhibiting agents may also be entrapped in microcapsules prepared, for example, by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences (1980) 16th edition, Osol, A. Ed.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM. (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for

over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37.degree. C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

The present invention also provides methods of inhibiting the processing of amyloid-beta precursor protein in a subject suffering or susceptible to the abnormal processing of said protein, which comprise the administration to said subject a therapeutically effective amount of an expression-inhibiting agent of the invention. Another aspect of the present method invention is the treatment or prevention of a condition involving cognitive impairment or a susceptibility to the condition. A special embodiment of this invention is a method wherein the condition is Alzheimer's disease.

As defined above, therapeutically effective dose means that amount of protein, polynucleotide, peptide, or its antibodies, agonists or antagonists, which ameliorate the symptoms or condition. Therapeutic efficacy and toxicity of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions that exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model is also used to achieve a desirable concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in

humans. The exact dosage is chosen by the individual physician in view of the patient to be treated. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Additional factors which may be taken into account include the severity of the disease state, age, weight and gender of the patient; diet, desired
5 duration of treatment, method of administration, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long acting pharmaceutical compositions might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

The pharmaceutical compositions according to this invention may be administered to
10 a subject by a variety of methods. They may be added directly to target tissues, complexed with cationic lipids, packaged within liposomes, or delivered to target cells by other methods known in the art. Localized administration to the desired tissues may be done by catheter, infusion pump or stent. The DNA, DNA/vehicle complexes, or the recombinant virus particles are locally administered to the site of treatment. Alternative routes of delivery
15 include, but are not limited to, intravenous injection, intramuscular injection, subcutaneous injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery. Examples of ribozyme delivery and administration are provided in Sullivan et al. WO 94/02595.

Antibodies according to the invention may be delivered as a bolus only, infused over
20 time or both administered as a bolus and infused over time. Those skilled in the art may employ different formulations for polynucleotides than for proteins. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

As discussed hereinabove, recombinant viruses may be used to introduce DNA encoding polynucleotide agents useful in the present invention. Recombinant viruses
25 according to the invention are generally formulated and administered in the form of doses of between about $10^{1.4}$ and about 10^{14} pfu. In the case of AAVs and adenoviruses, doses of from about $10^{6.6}$ to about $10^{11.1}$ pfu are preferably used. The term pfu ("plaque-forming unit") corresponds to the infective power of a suspension of virions and is determined by infecting an appropriate cell culture and measuring the number of plaques
30 formed. The techniques for determining the pfu titre of a viral solution are well documented in the prior art.

Still another aspect of the invention relates to a method for diagnosing a pathological condition involving cognitive impairment or a susceptibility to the condition in a subject, comprising determining the amount of polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6 in a biological sample, and comparing
5 the amount with the amount of the polypeptide in a healthy subject, wherein an increase of the amount of polypeptide compared to the healthy subject is indicative of the presence of the pathological condition.

Experimental Section

EXAMPLE 1: GPR3 Increases Amyloid Beta 1-42 Levels.

10 To identify novel drug targets that change the APP processing, a stable cell line over expressing APP is generated. This stable cell line is made by transfecting HEK293 cells with APP770wt cDNA cloned into pcDNA3.1, followed by selection with G418 for 3 weeks. At this time point colonies are picked and stable clones are expanded and tested for their secreted amyloid-beta peptide levels. One clone that secretes amyloid-beta at a high level,
15 HEK293 APPwt, is selected for experiments to identify drug targets. This is accomplished by transducing HEK293 APPwt with adenoviral cDNA libraries and measuring changes to the resulting amyloid beta 1-42 levels via ELISA.

Cells seeded in collagen-coated plates at a cell density of 15000 cells/well (384 well plate) in DMEM (10%FBS), are infected 24 h later with 1 μ l or 0.2 μ l of adenovirus
20 (corresponding to an average multiplicity of infection (MOI) of 120 and 24 respectively). The following day, the virus is washed away and DMEM (25 mM Hepes; 10%FBS) is added to the cells. Amyloid-beta peptides are allowed to accumulate during 24h. The ELISA plate is prepared by coating with a capture antibody (JRF/cAbeta42/26) (obtained from M Mercken, Johnson and Johnson Pharmaceutical Research and Development, B-2340 Beerse,
25 Belgium) overnight in buffer 42 (Table 1) at a concentration of 2.5 μ g/ml. The excess capture antibody is washed away the next morning with PBS and the ELISA plate is then blocked overnight with casein buffer (see Table 1) at 4°C. Upon removal of the blocking buffer, 30 μ l of the sample is transferred to the ELISA plate and incubated overnight at 4°C. After extensive washing with PBS-Tween20 and PBS, 30 μ l of the horseradish peroxidase
30 (HRP) labeled detection antibody (Peroxidase Labeling Kit, Roche), JRF/AbetaN/25-HRP (obtained from M Mercken, Johnson and Johnson Pharmaceutical Research and Development, B-2340 Beerse, Belgium) is diluted 1/5000 in buffer C (see Table 1) and added

to the wells for another 2h. Following the removal of excess detection antibody by a wash with PBS-Tween20 and PBS, HRP activity is detected via addition of luminol substrate (Roche), which is converted into a chemiluminescent signal by the HRP enzyme.

5 TABLE 1: buffers and solutions used for ELISA

Buffer 42	30mM NaHCO ₃ , 70mM Na ₂ CO ₃ , 0.05% NaN ₃ , pH9.6
Casein buffer	0.1% casein in PBS 1x
EC Buffer	20mM sodium phosphate, 2mM EDTA, 400mM NaCl, 0.2% BSA, 0.05% CHAPS, 0.4% casein, 0.05% NaN ₃ , pH7
Buffer C	20mM sodium phosphate, 2mM EDTA, 400mM NaCl, 1% BSA, pH7
PBS 10x	80g NaCl + 2g KCl + 11.5g Na ₂ HPO ₄ .7H ₂ O + 2g KH ₂ PO ₄ in 1 l milli Q, pH 7.4
PBST	PBS 1x with 0.05% Tween 20

In order to validate the assay, the effect of adenoviral over expression with random titer of two clinical PS1 mutants and BACE on amyloid beta 1-42 production is evaluated in the HEK293 APPwt cells. As is shown in Figure 2, all PS1 and BACE constructs induce
10 amyloid beta 1-42 levels as expected.

An adenoviral GPCR cDNA library was constructed as follows. DNA fragments covering the full coding region of the GPCRs, are amplified by PCR from a pooled placental and fetal liver cDNA library (InvitroGen). All fragments are cloned into an adenoviral vector as described in US 6,340,595, the contents of which are herein incorporated by reference, and
15 subsequently adenoviruses are made harboring the corresponding cDNAs. During the screening of the adenoviral GPCR library in the HEK293 APPwt cells, over expression of GPR3 lead to increased levels of amyloid beta 1-42 peptides in the conditioned medium of HEK293 APPwt cells. These results indicate that GPR3 was identified as a modulator of APP processing.

20 The stimulatory effect of GPR3 is confirmed upon re-screening of the viruses with a known titer (viral particles/ml), as determined by quantitative real time PCR. GPR3 virus is infected at MOIs ranging from 2 to 250 and the experiment is performed as described above. Amyloid beta 1-42 levels are 2 fold higher compared to the negative controls for Ad5/GPR3 (Figure 3A). In addition, the effect of GPR3 on amyloid beta 1-40, 11-42, 1-x and y-42
25 levels are checked under similar conditions as above (Figure 3B-3E). The respective ELISAs are performed as described above, except that the following antibodies were used: for the

amyloid beta 1-40 ELISA, the capture and detection antibody are respectively JRF/cAbeta40/10 and JRF/AbetaN/25-HRP (obtained from M Mercken, Johnson and Johnson Pharmaceutical Research and Development, B-2340 Beerse, Belgium), for the amyloid beta 11-42 ELISA, the capture and detection antibody are respectively JRF/cAbeta42/26 and JRF/hAb11/1 (obtained from M Mercken, Johnson and Johnson Pharmaceutical Research and Development, B-2340 Beerse, Belgium), for the amyloid beta y-42 ELISA (y ranges from 1-17), the capture and detection antibody are respectively JRF/cAbeta42/26 and 4G8-HRP (obtained respectively from M Mercken, Johnson and Johnson Pharmaceutical Research and Development, B-2340 Beerse, Belgium and from Signet, USA) while for the amyloid beta 1-x ELISA (x ranges from 24-42) the capture and detection antibodies are JRF/AbetaN/25 and 4G8-HRP, respectively (obtained respectively from M Mercken, Johnson and Johnson Pharmaceutical Research and Development, B-2340 Beerse, Belgium and from Signet, USA). The amyloid beta 1-x ELISA is used for the detection of amyloid peptides with a variable C-terminus (amyloid beta 1-37; 1-38; 1-39; 1-40; 1-42). The results of these experiments clearly show an increase of amyloid beta 1-40, 11-42, y-42 and 1-x species upon transduction of GPR3 (figure 3B-3E). The same procedure is used for the analysis of APP processing by GPR6 and GPR12.

EXAMPLE 2: Identification of homologues of GPR3.

The amino acid sequence of the human GPR3 receptor was used as query in a BLAST search against all the human GPCRs in order to find its closest homologues. Table 2 (SEQ ID NO: 5-6) shows the 2 closest homologues of the GPR3 receptor. Using ClustalW an alignment was constructed showing the degree of homology between the GPR3 and its closest homologues, the GPR6 and GPR12 (figure 5).

TABLE 2: GPCRs involved in APP processing (SEQ ID NO: 1-3; 4-6), Sequences for expression-inhibiting agent (SEQ ID NO: 7-287), the hairpin loop sequence of the RNAi (SEQ ID NO: 288), and the domains of GPR3, GPR6, and GPR 12 (SEQ ID NO: 289-333):

SEQID NO	Galapagos ID	Accession	Sequence		Type
1	1772	NM_005281		GPR3	DNA
2	1780	NM_005284		GPR6	DNA
3	1763	NM_005288		GPR12	DNA
4	1772	NP_005272		GPR3	Protein

5	1780	NP_005275		GPR6	Protein
6	1763	NP_005279		GPR12	Protein
7	NM_005281_idx127	NM_005281	TGGGATGTGGTGCTCTGCATC	GPR3	DNA
8	NM_005281_idx129	NM_005281	GGATGTGGTGCTCTGCATCTC	GPR3	DNA
9	NM_005281_idx172	NM_005281	AATGCGCTAGTGGTGGCCATC	GPR3	DNA
10	NM_005281_idx280	NM_005281	GTCCTGCACTTTGCTGCTGTC	GPR3	DNA
11	NM_005281_idx283	NM_005281	CTGCACTTTGCTGCTGTCTTC	GPR3	DNA
12	NM_005281_idx286	NM_005281	CACTTTGCTGCTGTCTTCTGC	GPR3	DNA
13	NM_005281_idx289	NM_005281	TTTGCTGCTGTCTTCTGCATC	GPR3	DNA
14	NM_005281_idx294	NM_005281	TGCTGTCTTCTGCATCGGCTC	GPR3	DNA
15	NM_005281_idx297	NM_005281	TGTCTTCTGCATCGGCTCAGC	GPR3	DNA
16	NM_005281_idx342	NM_005281	CGTGCTGGCAATGGCCTTTAC	GPR3	DNA
17	NM_005281_idx343	NM_005281	GTGCTGGCAATGGCCTTTACC	GPR3	DNA
18	NM_005281_idx352	NM_005281	ATGGCCTTTACCGCCAGCATC	GPR3	DNA
19	NM_005281_idx370	NM_005281	ATCGGCAGTCTACTGGCCATC	GPR3	DNA
20	NM_005281_idx376	NM_005281	AGTCTACTGGCCATCACTGTC	GPR3	DNA
21	NM_005281_idx379	NM_005281	CTACTGGCCATCACTGTGCGAC	GPR3	DNA
22	NM_005281_idx380	NM_005281	TACTGGCCATCACTGTGCGACC	GPR3	DNA
23	NM_005281_idx390	NM_005281	CACTGTGCGACCGCTACCTTTC	GPR3	DNA
24	NM_005281_idx392	NM_005281	CTGTGCGACCGCTACCTTCTC	GPR3	DNA
25	NM_005281_idx397	NM_005281	GACCGCTACCTTCTCTGTAC	GPR3	DNA
26	NM_005281_idx402	NM_005281	CTACCTTCTCTGTACAATGC	GPR3	DNA
27	NM_005281_idx403	NM_005281	TACCTTCTCTGTACAATGCC	GPR3	DNA
28	NM_005281_idx404	NM_005281	ACCTTCTCTGTACAATGCCC	GPR3	DNA
29	NM_005281_idx406	NM_005281	CTTCTCTGTACAATGCCCTC	GPR3	DNA
30	NM_005281_idx408	NM_005281	TTCTCTGTACAATGCCCTCAC	GPR3	DNA
31	NM_005281_idx409	NM_005281	TCTCTGTACAATGCCCTCACC	GPR3	DNA
32	NM_005281_idx412	NM_005281	CTGTACAATGCCCTCACCTAC	GPR3	DNA
33	NM_005281_idx417	NM_005281	CAATGCCCTCACCTACTATTC	GPR3	DNA
34	NM_005281_idx423	NM_005281	CCTCACCTACTATTCAGAGAC	GPR3	DNA
35	NM_005281_idx426	NM_005281	CACCTACTATTCAGAGACAAC	GPR3	DNA
36	NM_005281_idx432	NM_005281	CTATTCAGAGACAACAGTGAC	GPR3	DNA
37	NM_005281_idx434	NM_005281	ATTCAGAGACAACAGTGACAC	GPR3	DNA
38	NM_005281_idx438	NM_005281	AGAGACAACAGTGACACGGAC	GPR3	DNA
39	NM_005281_idx439	NM_005281	GAGACAACAGTGACACGGACC	GPR3	DNA
40	NM_005281_idx449	NM_005281	TGACACGGACCTATGTGATGC	GPR3	DNA
41	NM_005281_idx453	NM_005281	ACGGACCTATGTGATGCTGGC	GPR3	DNA
42	NM_005281_idx545	NM_005281	CCACATGTGGCGTGGTTTATC	GPR3	DNA
43	NM_005281_idx546	NM_005281	CACATGTGGCGTGGTTTATCC	GPR3	DNA
44	NM_005281_idx548	NM_005281	CATGTGGCGTGGTTTATCCAC	GPR3	DNA
45	NM_005281_idx550	NM_005281	TGTGGCGTGGTTTATCCACTC	GPR3	DNA
46	NM_005281_idx552	NM_005281	TGGCGTGGTTTATCCACTCTC	GPR3	DNA
47	NM_005281_idx553	NM_005281	GGCGTGGTTTATCCACTCTCC	GPR3	DNA
48	NM_005281_idx559	NM_005281	GTTTATCCACTCTCCAAGAAC	GPR3	DNA
49	NM_005281_idx560	NM_005281	TTTATCCACTCTCCAAGAACC	GPR3	DNA
50	NM_005281_idx563	NM_005281	ATCCACTCTCCAAGAACCATC	GPR3	DNA
51	NM_005281_idx572	NM_005281	CCAAGAACCATCTGGTAGTTC	GPR3	DNA
52	NM_005281_idx576	NM_005281	GAACCATCTGGTAGTTCTGGC	GPR3	DNA

53	NM_005281_idx577	NM_005281	AACCATCTGGTAGTTCTGGCC	GPR3	DNA
54	NM_005281_idx582	NM_005281	TCTGGTAGTTCTGGCCATTGC	GPR3	DNA
55	NM_005281_idx583	NM_005281	CTGGTAGTTCTGGCCATTGCC	GPR3	DNA
56	NM_005281_idx586	NM_005281	GTAGTTCTGGCCATTGCCTTC	GPR3	DNA
57	NM_005281_idx589	NM_005281	GTTCTGGCCATTGCCTTCTTC	GPR3	DNA
58	NM_008154_idx1099	NM_005281	GCCTTCTTCATGGTGTTTGGC	GPR3	DNA
59	NM_005281_idx604	NM_005281	TTCTTCATGGTGTTTGGCATC	GPR3	DNA
60	NM_005281_idx608	NM_005281	TCATGGTGTTTGGCATCATGC	GPR3	DNA
61	NM_005281_idx611	NM_005281	TGGTGTTTGGCATCATGCTGC	GPR3	DNA
62	NM_005281_idx614	NM_005281	TGTTTGGCATCATGCTGCAGC	GPR3	DNA
63	NM_005281_idx616	NM_005281	TTTGGCATCATGCTGCAGCTC	GPR3	DNA
64	NM_005281_idx619	NM_005281	GGCATCATGCTGCAGCTCTAC	GPR3	DNA
65	NM_005281_idx621	NM_005281	CATCATGCTGCAGCTCTACGC	GPR3	DNA
66	NM_005281_idx622	NM_005281	ATCATGCTGCAGCTCTACGCC	GPR3	DNA
67	NM_005281_idx628	NM_005281	CTGCAGCTCTACGCCCAAATC	GPR3	DNA
68	NM_005281_idx631	NM_005281	CAGCTCTACGCCCAAATCTGC	GPR3	DNA
69	NM_005281_idx632	NM_005281	AGCTCTACGCCCAAATCTGCC	GPR3	DNA
70	NM_005281_idx637	NM_005281	TACGCCCAAATCTGCCGCATC	GPR3	DNA
71	NM_005281_idx643	NM_005281	CAAATCTGCCGCATCGTCTGC	GPR3	DNA
72	NM_005281_idx644	NM_005281	AAATCTGCCGCATCGTCTGCC	GPR3	DNA
73	NM_005281_idx668	NM_005281	ATGCCCAGCAGATTGCCCTTC	GPR3	DNA
74	NM_005281_idx775	NM_005281	TGCTGGTTGCCCTTCACTGTC	GPR3	DNA
75	NM_005281_idx778	NM_005281	TGGTTGCCCTTCACTGTCTAC	GPR3	DNA
76	NM_005281_idx781	NM_005281	TTGCCCTTCACTGTCTACTGC	GPR3	DNA
77	NM_005281_idx782	NM_005281	TGCCCTTCACTGTCTACTGCC	GPR3	DNA
78	NM_005281_idx785	NM_005281	CCTTCACTGTCTACTGCCTGC	GPR3	DNA
79	NM_005281_idx816	NM_005281	CCACTCTCCACCTCTCTACAC	GPR3	DNA
80	NM_005281_idx817	NM_005281	CACTCTCCACCTCTCTACACC	GPR3	DNA
81	NM_005281_idx821	NM_005281	CTCCACCTCTCTACACCTATC	GPR3	DNA
82	NM_005281_idx825	NM_005281	ACCTCTCTACACCTATCTTAC	GPR3	DNA
83	NM_005281_idx826	NM_005281	CCTCTCTACACCTATCTTACC	GPR3	DNA
84	NM_005281_idx830	NM_005281	TCTACACCTATCTTACCTTGC	GPR3	DNA
85	NM_005281_idx832	NM_005281	TACACCTATCTTACCTTGCTC	GPR3	DNA
86	NM_005281_idx833	NM_005281	ACACCTATCTTACCTTGCTCC	GPR3	DNA
87	NM_005281_idx834	NM_005281	CACCTATCTTACCTTGCTCCC	GPR3	DNA
88	NM_005281_idx837	NM_005281	CTATCTTACCTTGCTCCCTGC	GPR3	DNA
89	NM_005281_idx838	NM_005281	TATCTTACCTTGCTCCCTGCC	GPR3	DNA
90	NM_005281_idx840	NM_005281	TCTTACCTTGCTCCCTGCCAC	GPR3	DNA
91	NM_005281_idx847	NM_005281	TTGCTCCCTGCCACCTACAAC	GPR3	DNA
92	NM_008154_idx1354	NM_005281	GCCACCTACAACCTCCATGATC	GPR3	DNA
93	NM_005281_idx859	NM_005281	ACCTACAACCTCCATGATCAAC	GPR3	DNA
94	NM_008154_idx1358	NM_005281	CCTACAACCTCCATGATCAACC	GPR3	DNA
95	NM_005281_idx861	NM_005281	CTACAACCTCCATGATCAACCC	GPR3	DNA
96	NM_005281_idx865	NM_005281	AACTCCATGATCAACCCTATC	GPR3	DNA
97	NM_005281_idx868	NM_005281	TCCATGATCAACCCTATCATC	GPR3	DNA
98	NM_005281_idx873	NM_005281	GATCAACCCTATCATCTACGC	GPR3	DNA
99	NM_005281_idx874	NM_005281	ATCAACCCTATCATCTACGCC	GPR3	DNA
100	NM_005281_idx877	NM_005281	AACCCTATCATCTACGCCTTC	GPR3	DNA

101	NM_005281_idx878	NM_005281	ACCCTATCATCTACGCCTTCC	GPR3	DNA
102	NM_005281_idx880	NM_005281	CCTATCATCTACGCCTTCCGC	GPR3	DNA
103	NM_005281_idx883	NM_005281	ATCATCTACGCCTTCCGCAAC	GPR3	DNA
104	NM_005281_idx884	NM_005281	TCATCTACGCCTTCCGCAACC	GPR3	DNA
105	NM_005281_idx902	NM_005281	ACCAGGATGTGCAGAAAGTGC	GPR3	DNA
106	NM_005281_idx909	NM_005281	TGTGCAGAAAGTGCTGTGGGC	GPR3	DNA
107	NM_005281_idx916	NM_005281	AAAGTGCTGTGGGCTGTCTGC	GPR3	DNA
108	NM_005281_idx941	NM_005281	GCTGTTCTCTTCCAAGATCC	GPR3	DNA
109	NM_005284_idx146	NM_005284	GAGCTAATGGGTCTCTGGAGC	GPR6	DNA
110	NM_005284_idx150	NM_005284	TAATGGGTCTCTGGAGCTGTC	GPR6	DNA
111	NM_005284_idx151	NM_005284	AATGGGTCTCTGGAGCTGTCC	GPR6	DNA
112	NM_005284_idx319	NM_005284	ATGTTCTGTGCTGGTAGGCAGC	GPR6	DNA
113	NM_005284_idx373	NM_005284	CTCATCTTGCACTTTGTGTTC	GPR6	DNA
114	NM_005284_idx374	NM_005284	TCATCTTGCACTTTGTGTTC	GPR6	DNA
115	NM_005284_idx379	NM_005284	TTGCACTTTGTGTTCCAGTAC	GPR6	DNA
116	NM_005284_idx386	NM_005284	TTGTGTTCCAGTACTTGGTGCC	GPR6	DNA
117	NM_005284_idx387	NM_005284	TGTGTTCCAGTACTTGGTGCC	GPR6	DNA
118	NM_005284_idx388	NM_005284	GTGTTCCAGTACTTGGTGCCC	GPR6	DNA
119	NM_005284_idx390	NM_005284	GTTCCAGTACTTGGTGCCCTC	GPR6	DNA
120	NM_005284_idx409	NM_005284	TCGGAGACTGTGAGTCTGCTC	GPR6	DNA
121	NM_005284_idx411	NM_005284	GGAGACTGTGAGTCTGCTCAC	GPR6	DNA
122	NM_005284_idx496	NM_005284	CGCTACCTGTCCCTGTATAAC	GPR6	DNA
123	NM_005284_idx498	NM_005284	CTACCTGTCCCTGTATAACGC	GPR6	DNA
124	NM_005284_idx500	NM_005284	ACCTGTCCCTGTATAACGCGC	GPR6	DNA
125	NM_005284_idx502	NM_005284	CTGTCCCTGTATAACGCGCTC	GPR6	DNA
126	NM_005284_idx504	NM_005284	GTCCCTGTATAACGCGCTCAC	GPR6	DNA
127	NM_005284_idx505	NM_005284	TCCCTGTATAACGCGCTCACC	GPR6	DNA
128	NM_005284_idx511	NM_005284	TATAACGCGCTCACCTATTAC	GPR6	DNA
129	NM_005284_idx513	NM_005284	TAACGCGCTCACCTATTACTC	GPR6	DNA
130	NM_005284_idx515	NM_005284	ACGCGCTCACCTATTACTCGC	GPR6	DNA
131	NM_005284_idx694	NM_005284	GCCGCTTCTTCATGGTCTTC	GPR6	DNA
132	NM_005284_idx697	NM_005284	GCCTTCTTCATGGTCTTCGGC	GPR6	DNA
133	NM_005284_idx700	NM_005284	TTCTTCATGGTCTTCGGCATC	GPR6	DNA
134	NM_005284_idx704	NM_005284	TCATGGTCTTCGGCATCATGC	GPR6	DNA
135	NM_005284_idx707	NM_005284	TGGTCTTCGGCATCATGCTGC	GPR6	DNA
136	NM_005284_idx709	NM_005284	GTCTTCGGCATCATGCTGCAC	GPR6	DNA
137	NM_005284_idx710	NM_005284	TCTTCGGCATCATGCTGCACC	GPR6	DNA
138	NM_005284_idx715	NM_005284	GGCATCATGCTGCACCTGTAC	GPR6	DNA
139	NM_005284_idx719	NM_005284	TCATGCTGCACCTGTACGTGC	GPR6	DNA
140	NM_005284_idx819	NM_005284	CACCAGAAAGGGTGTGGGTAC	GPR6	DNA
141	NM_005284_idx821	NM_005284	CCAGAAAGGGTGTGGGTACAC	GPR6	DNA
142	NM_005284_idx825	NM_005284	AAAGGGTGTGGGTACACTGGC	GPR6	DNA
143	NM_005284_idx877	NM_005284	CTGCCCTTCGCCATCTATTGC	GPR6	DNA
144	NM_005284_idx889	NM_005284	ATCTATTGCGTGGTGGGCAGC	GPR6	DNA
145	NM_005284_idx926	NM_005284	TCTACACTTACGCCACCCTGC	GPR6	DNA
146	NM_005284_idx956	NM_005284	CCTACAACCTCATGATCAATC	GPR6	DNA
147	NM_005284_idx957	NM_005284	CTACAACCTCATGATCAATCC	GPR6	DNA
148	NM_005284_idx958	NM_005284	TACAACCTCATGATCAATCCC	GPR6	DNA

149	NM_005284_idx961	NM_005284	AACTCCATGATCAATCCCATC	GPR6	DNA
150	NM_005284_idx964	NM_005284	TCCATGATCAATCCCATC	GPR6	DNA
151	NM_005284_idx969	NM_005284	GATCAATCCCATCATCTATGC	GPR6	DNA
152	NM_005284_idx970	NM_005284	ATCAATCCCATCATCTATGCC	GPR6	DNA
153	NM_000647_idx981	NM_005284	AATCCCATCATCTATGCCTTC	GPR6	DNA
154	NM_005284_idx974	NM_005284	ATCCCATCATCTATGCCTTCC	GPR6	DNA
155	NM_005284_idx976	NM_005284	CCCATCATCTATGCCTTCCGC	GPR6	DNA
156	NM_005284_idx979	NM_005284	ATCATCTATGCCTTCCGCAAC	GPR6	DNA
157	NM_005284_idx980	NM_005284	TCATCTATGCCTTCCGCAACC	GPR6	DNA
158	NM_005284_idx1024	NM_005284	CTCCTGCTCTGTGGCTGTTTC	GPR6	DNA
159	NM_005284_idx1025	NM_005284	TCCTGCTCTGTGGCTGTTTCC	GPR6	DNA
160	NM_005284_idx1029	NM_005284	GCTCTGTGGCTGTTTCCAGTC	GPR6	DNA
161	NM_005284_idx1030	NM_005284	CTCTGTGGCTGTTTCCAGTCC	GPR6	DNA
162	NM_005284_idx1037	NM_005284	GCTGTTTCCAGTCCAAAGTGC	GPR6	DNA
163	NM_005284_idx1038	NM_005284	CTGTTTCCAGTCCAAAGTGCC	GPR6	DNA
164	NM_005284_idx1039	NM_005284	TGTTTCCAGTCCAAAGTGCCC	GPR6	DNA
165	NM_005284_idx1043	NM_005284	TCCAGTCCAAAGTGCCCTTTC	GPR6	DNA
166	NM_005284_idx1047	NM_005284	GTCCAAAGTGCCCTTTCGTTTC	GPR6	DNA
167	NM_005284_idx1048	NM_005284	TCCAAAGTGCCCTTTCGTTCC	GPR6	DNA
168	NM_005284_idx1053	NM_005284	AGTGCCCTTTCGTTCCAGGTC	GPR6	DNA
169	NM_005284_idx1055	NM_005284	TGCCCTTTCGTTCCAGGTCTC	GPR6	DNA
170	NM_005284_idx1060	NM_005284	TTTCGTTCCAGGTCTCCCAGC	GPR6	DNA
171	NM_005288_idx115	NM_005288	GAGCCTGAGCTCGTAGTCAAC	GPR12	DNA
172	NM_005288_idx116	NM_005288	AGCCTGAGCTCGTAGTCAACC	GPR12	DNA
173	NM_005288_idx138	NM_005288	CTGGGACATTGTCTTGTGTAC	GPR12	DNA
174	NM_005288_idx139	NM_005288	TGGGACATTGTCTTGTGTACC	GPR12	DNA
175	NM_005288_idx141	NM_005288	GGACATTGTCTTGTGTACCTC	GPR12	DNA
176	NM_005288_idx147	NM_005288	TGTCTTGTGTACCTCGGGAAC	GPR12	DNA
177	NM_005288_idx148	NM_005288	GTCTTGTGTACCTCGGGAACC	GPR12	DNA
178	NM_005288_idx149	NM_005288	TCTTGTGTACCTCGGGAACCC	GPR12	DNA
179	NM_005288_idx151	NM_005288	TTGTGTACCTCGGGAACCCTC	GPR12	DNA
180	NM_005288_idx154	NM_005288	TGTACCTCGGGAACCCTCATC	GPR12	DNA
181	NM_005288_idx156	NM_005288	TACCTCGGGAACCCTCATCTC	GPR12	DNA
182	NM_005288_idx184	NM_005288	AATGCCATTGTGGTCCTTATC	GPR12	DNA
183	NM_005288_idx187	NM_005288	GCCATTGTGGTCCTTATCATC	GPR12	DNA
184	NM_005288_idx191	NM_005288	TTGTGGTCCTTATCATCTTCC	GPR12	DNA
185	NM_005288_idx193	NM_005288	GTGGTCCTTATCATCTTCCAC	GPR12	DNA
186	NM_005288_idx196	NM_005288	GTCCTTATCATCTTCCACAAC	GPR12	DNA
187	NM_005288_idx197	NM_005288	TCCTTATCATCTTCCACAACC	GPR12	DNA
188	NM_005288_idx198	NM_005288	CCTTATCATCTTCCACAACCC	GPR12	DNA
189	NM_005288_idx232	NM_005288	CCCATGTTCTGCTAATAGGC	GPR12	DNA
190	NM_005288_idx235	NM_005288	ATGTTCTGCTAATAGGCAGC	GPR12	DNA
191	NM_005288_idx236	NM_005288	TGTTCTGCTAATAGGCAGCC	GPR12	DNA
192	NM_005288_idx242	NM_005288	TGCTAATAGGCAGCCTGGCTC	GPR12	DNA
193	NM_005288_idx246	NM_005288	AATAGGCAGCCTGGCTCTTGC	GPR12	DNA
194	NM_005288_idx312	NM_005288	CTACCTGCTTCAGTCAGAAGC	GPR12	DNA
195	NM_005288_idx313	NM_005288	TACCTGCTTCAGTCAGAAGCC	GPR12	DNA
196	NM_005288_idx315	NM_005288	CCTGCTTCAGTCAGAAGCCAC	GPR12	DNA

197	NM_005288_idx316	NM_005288	CTGCTTCAGTCAGAAGCCACC	GPR12	DNA
198	NM_005288_idx320	NM_005288	TTCAGTCAGAAGCCACCAAGC	GPR12	DNA
199	NM_005288_idx325	NM_005288	TCAGAAGCCACCAAGCTGGTC	GPR12	DNA
200	NM_005288_idx327	NM_005288	AGAAGCCACCAAGCTGGTCAC	GPR12	DNA
201	NM_005288_idx343	NM_005288	GTCACGATCGGCCTCATTGTC	GPR12	DNA
202	NM_005288_idx352	NM_005288	GGCCTCATTGTCGCCTCTTTC	GPR12	DNA
203	NM_005288_idx354	NM_005288	CCTCATTGTCGCCTCTTCTC	GPR12	DNA
204	NM_005288_idx357	NM_005288	CATTGTCGCCTCTTCTCTGC	GPR12	DNA
205	NM_005288_idx358	NM_005288	ATTGTCGCCTCTTCTCTGCC	GPR12	DNA
206	NM_005288_idx360	NM_005288	TGTCGCCTCTTCTCTGCCTC	GPR12	DNA
207	NM_005288_idx364	NM_005288	GCCTCTTCTCTGCCTCTGTC	GPR12	DNA
208	NM_005288_idx367	NM_005288	TCTTCTCTGCCTCTGTCTGC	GPR12	DNA
209	NM_005288_idx370	NM_005288	TTCTCTGCCTCTGTCTGCAGC	GPR12	DNA
210	NM_005288_idx382	NM_005288	GTCTGCAGCTTGCTGGCTATC	GPR12	DNA
211	NM_005288_idx384	NM_005288	CTGCAGCTTGCTGGCTATCAC	GPR12	DNA
212	NM_005288_idx391	NM_005288	TTGCTGGCTATCACTGTTGAC	GPR12	DNA
213	NM_005288_idx392	NM_005288	TGCTGGCTATCACTGTTGACC	GPR12	DNA
214	NM_005288_idx394	NM_005288	CTGGCTATCACTGTTGACCGC	GPR12	DNA
215	NM_005288_idx397	NM_005288	GCTATCACTGTTGACCGCTAC	GPR12	DNA
216	NM_005288_idx398	NM_005288	CTATCACTGTTGACCGCTACC	GPR12	DNA
217	NM_005288_idx400	NM_005288	ATCACTGTTGACCGCTACCTC	GPR12	DNA
218	NM_005288_idx402	NM_005288	CACTGTTGACCGCTACCTCTC	GPR12	DNA
219	NM_005288_idx404	NM_005288	CTGTTGACCGCTACCTCTCAC	GPR12	DNA
220	NM_005288_idx409	NM_005288	GACCGCTACCTCTCACTGTAC	GPR12	DNA
221	NM_005288_idx412	NM_005288	CGCTACCTCTCACTGTACTAC	GPR12	DNA
222	NM_005288_idx414	NM_005288	CTACCTCTCACTGTACTACGC	GPR12	DNA
223	NM_005288_idx416	NM_005288	ACCTCTCACTGTACTACGCTC	GPR12	DNA
224	NM_005288_idx420	NM_005288	CTCACTGTACTACGCTCTGAC	GPR12	DNA
225	NM_005288_idx424	NM_005288	CTGTACTACGCTCTGACGTAC	GPR12	DNA
226	NM_005288_idx425	NM_005288	TGTACTACGCTCTGACGTACC	GPR12	DNA
227	NM_005288_idx429	NM_005288	CTACGCTCTGACGTACCATTC	GPR12	DNA
228	NM_005288_idx438	NM_005288	GACGTACCATTCGGAGAGGAC	GPR12	DNA
229	NM_005288_idx442	NM_005288	TACCATTCGGAGAGGACGGTC	GPR12	DNA
230	NM_005288_idx450	NM_005288	GGAGAGGACGGTCACGTTTAC	GPR12	DNA
231	NM_005288_idx451	NM_005288	GAGAGGACGGTCACGTTTACC	GPR12	DNA
232	NM_005288_idx457	NM_005288	ACGGTCACGTTTACCTATGTC	GPR12	DNA
233	NM_005288_idx461	NM_005288	TCACGTTTACCTATGTCATGC	GPR12	DNA
234	NM_005288_idx463	NM_005288	ACGTTTACCTATGTCATGCTC	GPR12	DNA
235	NM_005288_idx466	NM_005288	TTACCTATGTCATGCTCGTC	GPR12	DNA
236	NM_005288_idx470	NM_005288	CCTATGTCATGCTCGTCATGC	GPR12	DNA
237	NM_005288_idx472	NM_005288	TATGTCATGCTCGTCATGCTC	GPR12	DNA
238	NM_005288_idx571	NM_005288	GTCAGACCGCTACCAAGAAC	GPR12	DNA
239	NM_005288_idx574	NM_005288	AGACCGCTACCAAGAACAAC	GPR12	DNA
240	NM_005288_idx576	NM_005288	ACCGCTACCAAGAACAACGC	GPR12	DNA
241	NM_005288_idx583	NM_005288	ACCAAGAACAACGCGGCCATC	GPR12	DNA
242	NM_005288_idx586	NM_005288	AAGAACAACGCGGCCATCCTC	GPR12	DNA
243	NM_005288_idx601	NM_005288	ATCCTCTCGGTGTCCTTCCTC	GPR12	DNA
244	NM_005288_idx604	NM_005288	CTCTCGGTGTCCTTCCTCTTC	GPR12	DNA

245	NM_005288_idx612	NM_005288	GTCCTTCCTCTTCATGTTTGC	GPR12	DNA
246	NM_005288_idx614	NM_005288	CCTTCCTCTTCATGTTTGC	GPR12	DNA
247	NM_005288_idx616	NM_005288	TTCTCTTCATGTTTGC	GPR12	DNA
248	NM_005288_idx620	NM_005288	TCTTCATGTTTGC	GPR12	DNA
249	NM_005288_idx623	NM_005288	TCATGTTTGC	GPR12	DNA
250	NM_005288_idx626	NM_005288	TGTTTGC	GPR12	DNA
251	NM_005288_idx628	NM_005288	TTTGC	GPR12	DNA
252	NM_005288_idx631	NM_005288	GCGCTCATGCTTCAGCTCTAC	GPR12	DNA
253	NM_005288_idx634	NM_005288	CTCATGCTTCAGCTCTACATC	GPR12	DNA
254	NM_005288_idx635	NM_005288	TCATGCTTCAGCTCTACATCC	GPR12	DNA
255	NM_005288_idx640	NM_005288	CTTCAGCTCTACATCCAGATC	GPR12	DNA
256	NM_005288_idx659	NM_005288	TCTGTAAGATTGTGATGAGGC	GPR12	DNA
257	NM_005288_idx661	NM_005288	TGTAAGATTGTGATGAGGCAC	GPR12	DNA
258	NM_005288_idx663	NM_005288	TAAGATTGTGATGAGGCACGC	GPR12	DNA
259	NM_005288_idx664	NM_005288	AAGATTGTGATGAGGCACGCC	GPR12	DNA
260	NM_005288_idx665	NM_005288	AGATTGTGATGAGGCACGCCC	GPR12	DNA
261	NM_005288_idx668	NM_005288	TTGTGATGAGGCACGCCCATC	GPR12	DNA
262	NM_005288_idx685	NM_005288	CATCAGATAGCCCTGCAGCAC	GPR12	DNA
263	NM_005288_idx686	NM_005288	ATCAGATAGCCCTGCAGCACC	GPR12	DNA
264	NM_005288_idx691	NM_005288	ATAGCCCTGCAGCACCCTTC	GPR12	DNA
265	NM_005288_idx717	NM_005288	CACGTCGCACTATGTGACCAC	GPR12	DNA
266	NM_005288_idx718	NM_005288	ACGTCGCACTATGTGACCACC	GPR12	DNA
267	NM_005288_idx748	NM_005288	GTCTCCACCCTGGCTATCATC	GPR12	DNA
268	NM_005288_idx749	NM_005288	TCTCCACCCTGGCTATCATCC	GPR12	DNA
269	NM_005288_idx776	NM_005288	CGTTTGCTGCTTGCTGGATGC	GPR12	DNA
270	NM_005288_idx777	NM_005288	GTTTGCTGCTTGCTGGATGCC	GPR12	DNA
271	NM_005288_idx781	NM_005288	GCTGCTTGCTGGATGCCTTTC	GPR12	DNA
272	NM_005288_idx784	NM_005288	GCTTGCTGGATGCCTTTCACC	GPR12	DNA
273	NM_005288_idx811	NM_005288	TCCTTGATAGCGGATTACACC	GPR12	DNA
274	NM_005288_idx835	NM_005288	CCCTCCATCTATACCTACGCC	GPR12	DNA
275	NM_005288_idx838	NM_005288	TCCATCTATACCTACGCCACC	GPR12	DNA
276	NM_005288_idx839	NM_005288	CCATCTATACCTACGCCACCC	GPR12	DNA
277	NM_005288_idx842	NM_005288	TCTATACCTACGCCACCCTCC	GPR12	DNA
278	NM_005288_idx865	NM_005288	CCCGCCACCTACAATTCCATC	GPR12	DNA
279	NM_005288_idx868	NM_005288	GCCACCTACAATTCCATCATC	GPR12	DNA
280	NM_005288_idx872	NM_005288	CCTACAATTCCATCATCAACC	GPR12	DNA
281	NM_005288_idx877	NM_005288	AATTCCATCATCAACCCTGTC	GPR12	DNA
282	NM_005288_idx904	NM_005288	GCTTTCAGAAACCAAGAGATC	GPR12	DNA
283	NM_005288_idx912	NM_005288	AAACCAAGAGATCCAGAAAGC	GPR12	DNA
284	NM_005288_idx914	NM_005288	ACCAAGAGATCCAGAAAGCGC	GPR12	DNA
285	NM_005288_idx928	NM_005288	AAAGCGCTCTGTCTCATTTGC	GPR12	DNA
286	NM_005288_idx931	NM_005288	GCGCTCTGTCTCATTTGCTGC	GPR12	DNA
287	NM_005288_idx941	NM_005288	TCATTTGCTGCGGCTGCATCC	GPR12	DNA
288	Hairpin loop		TTGCTATA		DNA
289	N-term	MMWGAGSPLAWLSAGSGNVNVSSVGPAGPTGPAAPLPSPKA		GPR3	Protein
290	TM1	WDVVLCSGTLVSCENALVVAII		GPR3	Protein
291	IL1	VGTPAFRAPMFL		GPR3	Protein
292	TM2	LVGSLAVADLLAGLGLVLHFAAV		GPR3	Protein

293	EL1	FCIGSAEMS	GPR3	Protein
294	TM3	LVLVGVLMAMFTASIGSLLAITV	GPR3	Protein
295	IL2	DRYLSLYNALTYSETTVTR	GPR3	Protein
296	TM4	TYVMLALVWGGALGLLPLVLAW	GPR3	Protein
297	EL2	NCLDGLTTCGVVYPLSKNH	GPR3	Protein
298	TM5	LVVLAIAFFMVFVGIMLQLYAQIC	GPR3	Protein
299	IL3	RIVCRHAQQIALQRHLLPASHYVATRK	GPR3	Protein
300	TM6	GIATLAVVLGAFAACWLPFTVYC	GPR3	Protein
301	EL3	LLGDAHSP	GPR3	Protein
302	TM7	LYTYLTLLPATYNSMINPIIYAF	GPR3	Protein
303	C-term	RNQDVQKVLWAVCCCCSSSKIPIFRSRSPSDV	GPR3	Protein
304	N-term	MNASAAASLNDISQVVVAAEGAAAAATAAGGPDTEGWGPPAAAAALG AGGGANGSLELSSQLSAGPPGLLLPAVNP	GPR6	Protein
305	TM1	WDVLLCVSGTVIAGENALVVALI	GPR6	Protein
306	IL1	ASTPALRTPMFV	GPR6	Protein
307	TM2	LVGSLATADLLAGCGLILHVFVQ	GPR6	Protein
308	EL1	YLPSETVS	GPR6	Protein
309	TM3	LLTVGFLVASFAASVSSLLAITV	GPR6	Protein
310	IL2	DRYLSLYNALTYYSRRTLLG	GPR6	Protein
311	TM4	VHLLLAATWTVSLGLLPLVLGW	GPR6	Protein
312	EL2	NCLAERAACSVVRPLARSH	GPR6	Protein
313	TM5	VALLSAAFFMVFVGIMLHLYV	GPR6	Protein
314	IL3	RICQVVRHAHQIALQQHCLAPPHLAATRK	GPR6	Protein
315	TM6	GVGTLAVVLGTGASWLPFAIYC	GPR6	Protein
316	EL3	VVGSHEDPA	GPR6	Protein
317	TM7	VVGSHEDPAVYTYATLLPATYNSMINPIIYAF	GPR6	Protein
318	C-term	RNQEIQRALWLLLCGCFQSKVPFRSRSPSEV	GPR6	Protein
319	N-term	MNEDLKVNLISGLPRDYLDAAAENISAAVSSRVPVEPEPELVVNP	GPR12	Protein
320	TM1	WDIVLCTSGTLISCENAIIVLII	GPR12	Protein
321	IL1	FHNPSLRAPMFL	GPR12	Protein
322	TM2	LIGSLALADLLAGIGLITNFVFA	GPR12	Protein
323	EL1	YLLQSEATK	GPR12	Protein
324	TM3	LVTIGLIVASFASVCSLLAITV	GPR12	Protein
325	IL2	DRYLSLYYALTYHSERTVTF	GPR12	Protein
326	TM4	TYVMLVMLWGTSICLGLLPVMGW	GPR12	Protein
327	EL2	NCLRDESTCSVVRPLTKNN	GPR12	Protein
328	TM5	AAILSVSFLFMFALMLQLYIQIC	GPR12	Protein
329	IL3	KIVMRHAHQIALQHHFLATSHYVTTTK	GPR12	Protein
330	TM6	GVSTLAILLGTFAACWMPFTLYS	GPR12	Protein
331	EL3	LIADYTYP	GPR12	Protein
332	TM7	IYTYATLLPATYNSIINPVIYAF	GPR12	Protein
333	C-term	RNQEIQKALCLICCGCIPSSLAQRARSPSDV	GPR12	Protein

EXAMPLE 3: Amyloid Beta Peptide Reduction Via Knock Down Of GPCR Expression

The effect of an antagonist can be mimicked through the use of siRNA-based strategies, which result in decreased expression levels of the targeted protein. For example, transfection with GPR3 siRNA reduces amyloid beta 1-42.

HEK293 APPwt cells are transfected with a smart pool of siRNAs of GPR3 (Dharmacon, USA: Table 3), eGFP, Luciferase and BACE with Oligofectamine. 24 hours after transfection, the medium is refreshed and the cells are allowed to accumulate amyloid beta peptides in the conditioned medium for 24 hours prior to ELISA analysis as described above. The data clearly show that siRNA targeted against GPR3 RNA levels reduce amyloid beta 1-42 levels compared to the control conditions (Figure 4). In conclusion, these data show that GPR3 modulates the levels of secreted amyloid beta. The same procedure is used for the analysis of APP processing by GPR6 and GPR12.

TABLE 3: Specific siRNAs for GPR3 (Dharmacon, USA; SEQ ID NO: 334-337)

Gene symbol	NM number	Dharmacon Cat. number	Full sequence siRNA	SEQ ID NO:
GPR3	NM_005281	D-003951-01	GTTTATCCACTCTCCAAGA	334
GPR3	NM_005281	D-003951-02	TTTATCCACTCTCCAAGAA	335
GPR3	NM_005281	D-003951-03	CCACCTCTCTACACCTATC	336
GPR3	NM_005281	D-003951-04	ACCGCTACCTTTCTCTGTA	337

EXAMPLE 4: Expression Of GPR3 In The Human Brain.

Upon identification of a modulator of APP processing, it is important to evaluate whether the modulator is expressed in the tissue and the cells of interest. This can be achieved by measuring the RNA and/or protein levels in the tissue and cells. In recent years, RNA levels are being quantified through real time PCR technologies, whereby the RNA is first transcribed to cDNA and then the amplification of the cDNA of interest is monitored during a PCR reaction. The amplification plot and the resulting Ct value are indicators for the amount of RNA present in the sample. Determination of the levels of household keeping genes allows the normalization of RNA levels of the target gene between different RNA samples, represented as Δ Ct values.

To assess whether the GPCRs of the invention are expressed in the human brain, real time PCR with GAPDH specific primers and specific primers for each GPCR of the invention is performed on human total brain, human cerebral cortex, and human hippocampal total RNA (BD Biosciences)(see Table 4).

TABLE 4: Primers used in quantitative real time PCR-analysis of GPR3 expression levels (SEQ ID NO: 338-339)

Gene	Primer name	SEQ ID NO:	Primer sequence
GPR3	GPR3_Hs_For	338	GGCCTTTACCGCCAGCAT
	GPR3_Hs_Rev	339	TCTGAATAGTAGGTGAG GGCATTG

5 GAPDH is detected with a Taqman probe, while for the other GPCRs SybrGreen was used. In short, 40 ng of RNA is reverse-transcribed to DNA using the MultiScribe Reverse Transcriptase (50 U/ μ l) enzyme (Applied BioSystems). The resulting cDNA is amplified with AmpliTaq Gold DNA polymerase (Applied BioSystems) during 40 cycles using an ABI PRISM® 7000 Sequence Detection System.

10 Total brain, cerebral cortex and hippocampal total RNA are analyzed for the presence of the GPCR transcripts via quantitative real time PCR. For GPR3, the obtained Ct values indicate that it is detected in all RNA samples (Table 5).

To gain more insight into the specific cellular expression, immunohistochemistry (protein level) and/or *in situ* hybridization (RNA level) are carried out on sections from
15 human normal and Alzheimer's brain hippocampal, cortical and subcortical structures. These results indicate whether expression occurs in neurons, microglia cells, or astrocytes. The comparison of diseased tissue with healthy tissue indicates whether GPR3 is expressed in the diseased tissue and whether its expression level is changed compared to the non-pathological situation. The same procedure is used for expression profiling of GPR6 and GPR12.

20

TABLE 5: Quantitative real time PCR Ct values: Total human brain, cerebral cortex or hippocampus RNA tested for the presence of GPR3 RNA via quantitative real time PCR. GAPDH RNA is used to normalize all samples (Δ Ct).

Human Tissue	GAPDH Ct-values		GPR3 Ct-values		Δ Ct (+RT)
	+RT	-RT	+RT	-RT	
Total brain	21,29	NA	24,93	33,07	3,64
Hippocampus	21,65	NA	25,77	36,14	4,12
Cerebral cortex	20,97	NA	25,19	35,73	4,22

5 EXAMPLE 5: Amyloid Beta Production In Rat Primary Neuronal Cells.

In order to investigate whether GPR3 affects amyloid beta production in a real neuron, human or rat primary hippocampal or cortical neurons are transduced with adenovirus containing the GPR3 cDNA. Amyloid beta levels are determined by ELISA (see EXAMPLE 1). Since rodent APP genes carry a number of mutations in APP compared to the human sequence, they produce less amyloid beta 1-40 and 1-42. In order to achieve higher amyloid beta levels, co-transduction of GPR3 with human wild type APP or human Swedish mutant APP (which enhances amyloid beta production) cDNA is performed.

Rat primary neuron cultures are prepared from brain of E18-E19-day-old fetal Sprague Dawley rats according to Goslin and Banker (Culturing Nerve cells, second edition, 1998 ISBN 0-262-02438-1). Briefly, single cell suspensions obtained from the hippocampus or cortices are prepared. The number of viable cells is determined and plated on poly-L-lysine-coated plastic 96-well plates in minimal essential medium (MEM) supplemented with 10% horse serum. The cells are seeded at a density of 50,000 cells per well (i.e. about 166,000 cells/cm²). After 3–4 h, culture medium is replaced by 160 μ l serum-free neurobasal medium with B27 supplement (GIBCO BRL). Cytosine arabinoside (5 μ M) is added 24 h after plating to prevent nonneuronal (glial) cell proliferation.

Neurons are used at day 5 after plating. Before adenoviral transduction, 150 μ l conditioned medium of these cultures is transferred to the corresponding wells in an empty 96-well plate and 50 μ l of the conditioned medium is returned to the cells. The remaining 100 μ l/well is stored at 37°C and 5% CO₂. Hippocampal primary neuron cultures are infected with the crude lysate of Ad5C09Att00/A011200-GPR3_v3, Ad5C09Att00/A010801-

LacZ_v1, Ad5C09Att00/A010800-eGFP_v1 and Ad5C09Att00/A010800-luc_v17 viruses containing the cDNA of GPR3, LacZ, eGFP and luciferase respectively at different MOIs, ranging from 250 to 2000. In addition the cells are also infected with the purified adenovirus Ad5C01Att01/A010800 APP_v6 expressing human wild type APP695 at an MOI of 2000.

5 Sixteen to twenty-four hours after transduction, virus is removed and cultures are washed with 100 μ l pre-warmed fresh neurobasal medium. After removal of the wash solution, new medium, containing 50 μ l of the stored conditioned medium and 50 μ l of fresh neurobasal medium, is transferred to the corresponding cells. Medium was harvested after 48 and 72 hours. The cell number in the wells was determined by assessing the ATP levels. Amyloid
10 beta concentration was determined by amyloid beta 1-42 specific ELISA (see EXAMPLE 1). Amyloid beta 1-42 levels are normalized for cell number.

The data (Figure 6) clearly indicate that increased levels of over expression of GPR3 in the primary neurons result in a corresponding dose dependent increase of amyloid beta 1-42 levels compared to the negative control viruses.

15 EXAMPLE 6: Ligand Screens For GPCRs.

Reporter Gene Screen.

Mammalian cells such as HEK293 or CHO-K1 cells are either stably transfected with a plasmid harboring the luciferase gene under the control of a cAMP dependent promoter (CRE elements) or transduced with an adenovirus harboring a luciferase gene under the
20 control of a cAMP dependent promoter. In addition reporter constructs can be used with the luciferase gene under the control of a Ca^{2+} dependent promoter (NF-AT elements) or a promoter that is controlled by activated NF- κ B. These cells, expressing the reporter construct, are then transduced with an adenovirus harboring the cDNA of the GPCR of the present invention. Forty (40) hours after transduction the cells are treated with the following:

25 a) an agonist for the receptor (e.g. sphingosine 1 phosphate) and screened against a large collection of reference compounds comprising peptides (LOPAP, Sigma Aldrich), lipids (Biomol, TimTech), carbohydrates (Specs), natural compounds (Specs, TimTech), small chemical compounds (Tocris), commercially available screening libraries, and compounds that have been demonstrated to have binding affinity for a polypeptide
30 comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333, including compounds comprising aryloxydithiourea (see US 6,420,563), its salts, hydrates, or solvates, or

b) a large collection of reference compounds comprising peptides (LOPAP, Sigma Aldrich), lipids (Biomol, TimTech), carbohydrates (Specs), natural compounds (Specs, TimTech), small chemical compounds (Tocris), commercially available screening libraries, and compounds that have been demonstrated to have binding affinity for a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333, including compounds comprising aryloxydithiourea (see US 6,420,563), its salts, hydrates, or solvates, only, as GPR3 is considered to be a constitutively active GPCR.

Compounds, which decrease the agonist induced increase in luciferase activity or the constitutive activity, are considered to be antagonists or inverse agonists for the GPR3. These compounds are screened again for verification and screened against their effect on secreted amyloid beta peptide levels. The compounds are also screened to verify binding to the GPCR. The binding, amyloid-beta peptide and reporter activity assays can be performed in essentially any order to screen compounds.

In addition, cells expressing the NF-AT reporter gene can be transduced with an adenovirus harboring the cDNA encoding the α -subunit of G_{15} or chimerical $G\alpha$ subunits. G_{15} is a promiscuous G protein of the G_q class that couples to many different GPCRs and as such re-directs their signaling towards the release of intracellular Ca^{2+} stores. The chimerical G alpha subunits are members of the G_s and $G_{i/o}$ family by which the last 5 C-terminal residues are replaced by those of $G\alpha_q$, these chimerical G-proteins also redirect cAMP signaling to Ca^{2+} signaling.

FLIPR screen.

Mammalian cells such as HEK293 or CHO-K1 cells are stably transfected with an expression plasmid construct harboring the cDNA of a GPCR of the present invention. Cells are seeded, grown, and selected until sufficient stable cells can be obtained. Cells are loaded with a Ca^{2+} dependent fluorophore such as Fura3 or Fura4. After washing away the excess of fluorophore the cells are screened against a large collection of reference compounds comprising peptides (LOPAP, Sigma Aldrich), lipids (Biomol, TimTech), carbohydrates (Specs), natural compounds (Specs, TimTech), small chemical compounds (Tocris), commercially available screening libraries, and compounds that have been demonstrated to have binding affinity for a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333, including compounds comprising aryloxydithiourea (see US 6,420,563), its salts, hydrates, or solvates, by simultaneously

adding an agonist (alternatively no agonist need be added if the constitutive activity of the receptor is used) and a compound to the cells. Activation of the receptor is measured as an almost instantaneously increase in fluorescence due to the interaction of the fluorophore and the Ca^{2+} that is released. Compounds that reduce or inhibit the agonist induced increase in fluorescence (or constitutive fluorescence) are considered to be antagonists or inverse agonists for the receptor they are screened against. These compounds will be screened again to measure the amount of secreted amyloid beta peptide as well as binding to the GPCR.

AequoScreen.

CHO cells, stably expressing Apoequorin are stably transfected with a plasmid construct harboring the cDNA of a GPCR. Cells are seeded, grown, and selected until sufficient stable cells can be obtained. The cells are loaded with coelenterazine, a cofactor for apoequorin. Upon receptor activation intracellular Ca^{2+} stores will be emptied and the aequorin will react with the coelenterazine in a light emitting process. The emitted light is a measure for receptor activation. The CHO, stable expressing both the apoequorin and the receptor are screened against a large collection of reference compounds comprising peptides (LOPAP, Sigma Aldrich), lipids (Biomol, TimTech), carbohydrates (Specs), natural compounds (Specs, TimTech), small chemical compounds (Tocris), commercially available screening libraries, and compounds that have been demonstrated to have binding affinity for a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333, including compounds comprising aryloxydithiourea (see US 6,420,563), its salts, hydrates, or solvates, by simultaneously adding an agonist (alternatively no agonist need be added if the constitutive activity of the receptor is used) and a compound to the cells. Activation of the receptor is measured as an almost instantaneously light flash due to the interaction of the apoequorin, coelenterazine, and the Ca^{2+} that is released. Compounds that reduce or inhibit the agonist induced increase in light or the constitutive activity are considered to be antagonists or inverse agonists for the receptor they are screened against. These compounds will be screened again to measure the amount of secreted amyloid beta peptide as well as binding to the GPCR.

In addition, CHO cells stable expressing the apoequorin gene are stably transfected with a plasmid construct harboring the cDNA encoding the α -subunit of G_{15} or chimerical G_{α} subunits. G_{15} is a promiscuous G protein of the G_q class that couples to many different GPCRs and as such redirects their signaling towards the release of intracellular Ca^{2+} stores. The chimerical G alpha subunits are members of the G_s and $G_{i/o}$ family by which the last 5 C-

terminal residues are replaced by those of $G_{\alpha q}$, these chimerical G-proteins also redirect cAMP signaling to Ca^{2+} signaling.

Screening for compounds that bind to the GPCR polypeptides (displacement experiment)

Compounds are screened for binding to the GPCR polypeptides. The affinity of the compounds to the polypeptides is determined in a displacement experiment. In brief, the GPCR polypeptides are incubated with a labeled (radiolabeled, fluorescent labeled) ligand that is known to bind to the polypeptide (e.g., spingosine-1-phosphate or dihydrosphingosine-1-phosphate) and with an unlabeled compound. The displacement of the labeled ligand from the polypeptide is determined by measuring the amount of labeled ligand that is still associated with the polypeptide. The amount associated with the polypeptide is plotted against the concentration of the compound to calculate IC_{50} values. This value reflects the binding affinity of the compound to its target, i.e. the GPCR polypeptides. Strong binders have an IC_{50} in the nanomolar and even picomolar range. Compounds that have an IC_{50} of at least 10 micromol or better (nmol to pmol) are applied in beta amyloid secretion assay to check for their effect on the beta amyloid secretion and processing. The GPCR polypeptides can be prepared in a number of ways depending on whether the assay will be run on cells, cell fractions or biochemically, on purified proteins.

Screening for compounds that bind to the GPCR polypeptide (generic GPCR screening assay)

When a G protein receptor becomes constitutively active, it binds to a G protein (G_q , G_s , G_i , G_o) and stimulates the binding of GTP to the G protein. The G protein then acts as a GTPase and slowly hydrolyses the GTP to GDP, whereby the receptor, under normal conditions, becomes deactivated. However, constitutively activated receptors continue to exchange GDP to GTP. A non-hydrolyzable analog of GTP, $[^{35}S]GTP\gamma S$, can be used to monitor enhanced binding to membranes which express constitutively activated receptors. It is reported that $[^{35}S]GTP\gamma S$ can be used to monitor G protein coupling to membranes in the absence and presence of ligand. Moreover, a preferred approach is the use of a GPCR-G protein fusion protein. The strategy to generate a GPR3-G protein fusion protein is well known for those known in the art. Membranes expressing GPR3-G protein fusion protein are prepared for use in the direct identification of candidate compounds such as inverse agonist. Homogenized membranes with GPR3-G protein fusion protein are transferred in a 96-well plate. A pin-tool is used to transfer a candidate compound in each well plus $[^{35}S]GTP\gamma S$,

followed by incubation on a shaker for 60 minutes at room temperature. The assay is stopped by spinning of the plates at 4000 RPM for 15 minutes at 22°C. The plates are then aspirated and radioactivity is then read. The same procedure is used for analysis of GPR6 and GPR12.

Receptor Ligand Binding Study On Cell Surface

5 The receptor is expressed in mammalian cells (HEK293, CHO, COS7) by adenoviral transducing the cells (see US 6,340,595). The cells are incubated with both labeled ligand (iodinated, tritiated, or fluorescent) and the unlabeled compound at various concentrations, ranging from 10 pM to 10 μ M (3 hours at 4°C.: 25 mM HEPES, 140 mM NaCl, 1 mM CaCl₂, 5 mM MgCl₂ and 0.2% BSA, adjusted to pH 7.4). Reactions mixtures are aspirated onto PEI-
10 treated GF/B glass filters using a cell harvester (Packard). The filters are washed twice with ice cold wash buffer (25 mM HEPES, 500 mM NaCl, 1 mM CaCl₂, 5 mM MgCl₂, adjusted to pH 7.4). Scintillant (MicroScint-10; 35 μ l) is added to dried filters and the filters counted in a (Packard Topcount) scintillation counter. Data are analyzed and plotted using Prism software (GraphPad Software, San Diego, Calif.). Competition curves are analyzed and IC₅₀
15 values calculated. If one or more data points do not fall within the sigmoidal range of the competition curve or close to the sigmoidal range the assay is repeated and concentrations of labeled ligand and unlabeled compound adapted to have more data points close to or in the sigmoidal range of the curve.

Receptor Ligand Binding Studies On Membrane Preparations

20 Membranes preparations are isolated from mammalian cells (HEK293, CHO, COS7) cells over expressing the receptor is done as follows: Medium is aspirated from the transduced cells and cells are harvested in 1 x PBS by gentle scraping. Cells are pelleted (2500 rpm 5 min) and resuspended in 50 mM Tris pH 7.4 (10 x 10⁶ cells/ml). The cell pellet is homogenized by sonicating 3 x 5 sec (UP50H; sonotrode MS1; max amplitude: 140 μ m; max Sonic Power Density: 125W/cm²). Membrane fractions are prepared by centrifuging 20
25 min at maximal speed (13000 rpm ~15 000 to 20 000g or rcf). The resulting pellet is resuspended in 500 μ l 50 mM Tris pH 7.4 and sonicated again for 3 x 5 sec. The membrane fraction is isolated by centrifugation and finally resuspended in PBS. Binding competition and derivation of IC₅₀ values are determined as described above.

30 Internalization screen (1)

Activation of a GPCR-associated signal transduction pathway commonly leads to translocation of specific signal transduction molecules from the cytoplasm to the plasma

membrane or from the cytoplasm to the nucleus. Norak has developed their transfluor assay based on agonist-induced translocation of receptor- β -arrestin-GFP complex from the cytosol to the plasma membrane and subsequent internalization of this complex, which occurs during receptor desensitization. A similar assay uses GFP tagged receptor instead of β -arrestin.

5 HEK293 cells are transduced with a GPR3-eGFP vector that translates for a GPR3-eGFP fusion protein. 48 hours after transduction, the cells are set to fresh serum-free medium for 60 minutes and treated with a ligand (e.g. 100 nM sphingosine 1 phosphate) for 15, 30, 60 or 120 minutes at 37°C and 5% CO₂. After indicated exposure times, cells are washed with PBS and fixed with 5% paraformaldehyde for 20 minutes at RT. GFP fluorescence is
10 visualized with a Zeiss microscope with a digital camera. This method aims for the identification of compounds that inhibit a ligand-mediated (constitutive activity-mediated) translocation of the fusion protein to intracellular compartments. The same procedure is used for analysis of GPR6 and GPR12.

Internalization screen (2)

15 Various variations on translocation assays exists using β -arrestin and β -galactosidase enzyme complementation and BRET based assays with receptor as energy donor and β -arrestin as energy acceptor. Also the use of specific receptor antibodies labeled with pH sensitive dyes are used to detect agonist induced receptor translocation to acidic lysosomes. All of the translocation assays are used for screening for both agonistic and antagonistic
20 acting ligands.

Melanophore assay (Arena Pharmaceutical)

The melanophore assay is based on the ability of GPCRs to alter the distribution of melanin containing melanosomes in *Xenopus melanophores*. The distribution of the melanosomes depends on the exogenous receptor that is either Gi/o or Gs/q coupled. The distribution of
25 the melanosomes (dispersed or aggregated) is easily detected by measuring light absorption. This type of assay is used for both agonist as well as antagonist compound screens.

We claim:

1. A method for identifying a compound that inhibits the processing of amyloid-beta precursor protein in a mammalian cell, comprising
 - 5 (a) contacting a compound with a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333; and
 - (b) measuring a compound-polypeptide property related to the production of amyloid-beta protein.
2. The method according to claim 1, wherein said polypeptide comprises SEQ ID NO: 289-
10 333 in an *in vitro* cell-free preparation.
3. The method according to claim 1, wherein said polypeptide is membrane-bound.
4. The method according to claim 2, wherein said polypeptide is present as a transmembrane cell receptor in a mammalian cell.
5. The method of claims 1-4, wherein said property is a binding affinity of said compound to
15 said polypeptide.
6. The method of claims 1-4, wherein said property is activation of a biological pathway producing an indicator of the processing of amyloid-beta precursor protein.
7. The method of claim 6 wherein said indicator is a second messenger.
8. The method of claim 7 wherein said second messenger is cyclic AMP or Ca²⁺.
- 20 9. The method of claim 6 wherein said indicator is amyloid-beta peptide.
10. The method of claim 9 wherein said amyloid-beta protein is selected from the group consisting of one or more of amyloid-beta peptide 1-42, 1-40, 11-42 and 11-40.
11. The method of claim 10 wherein said amyloid-beta protein is amyloid-beta peptide 1-42.
12. The method according to claim 6 wherein said indicator induces the expression of a
25 reporter in said mammalian cell.
13. The method according to claim 12 wherein the reporter is selected from the group consisting of alkaline phosphatase, GFP, eGFP, dGFP, luciferase and β -galactosidase.
14. The method according to any of the claims 1-13, wherein said compound is selected from the group consisting of compounds of a commercially available screening library and

compounds that have been demonstrated to have binding affinity for a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6, 289-333.

15. The method according to claim 2, wherein said compound is a peptide in a phage display library or an antibody fragment library.

16. The method according to any of the claims 1-15, wherein said compound is an aryloxydithiourea, its salts, hydrates, or solvates.

17. An agent for the inhibition of amyloid-beta precursor processing selected from the group consisting of an antisense polynucleotide, a ribozyme, and a small interfering RNA (siRNA), wherein said agent comprises a nucleic acid sequence complementary to, or engineered from, a naturally occurring polynucleotide sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6.

18. The agent according to claim 17, wherein a vector in a mammalian cell expresses said agent.

19. The agent according to claim 18, wherein said vector is an adenoviral, retroviral, adeno-associated viral, lentiviral, a herpes simplex viral or a sendaiviral vector.

20. The agent according to claim 17, 18 or 19, wherein said antisense polynucleotide and said siRNA comprise an antisense strand of 17-25 nucleotides complementary to a sense strand, wherein said sense strand is selected from 17-25 continuous nucleotides of a naturally occurring nucleic acid sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4-6.

21. The agent according to any of the claims 17-20, wherein said siRNA further comprises said sense strand.

22. The agent according to claim 20, wherein said sense strand is selected from 17-25 continuous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1-3.

23. The agent according to any of the claims 17-22, wherein said siRNA further comprises a loop region connecting said sense and said antisense strand.

24. The agent according to claim 23 wherein said loop region comprises a nucleic acid sequence defined of SEQ ID NO: 288.

25. The agent according to any of the claims 17-24, wherein said agent is an antisense polynucleotide, ribozyme, or siRNA comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 7-287.

26. A cognitive enhancing pharmaceutical composition comprising a therapeutically effective amount of an agent of any of claims 17-25 in admixture with a pharmaceutically acceptable carrier.

27. The cognitive enhancing pharmaceutical composition according to claim 26 wherein said agent comprises a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 7-287, a polynucleotide complementary to said nucleic acid sequence, and a combination thereof.

28. A method of inhibiting the processing of amyloid-beta precursor protein in a subject suffering or susceptible to the abnormal processing of said protein, comprising administering to said subject a pharmaceutical composition according to claim 26 or 27.

29. A method according to claim 28 for treatment or prevention of a condition involving cognitive impairment or a susceptibility to the condition.

30. The method according to claim 29 wherein the condition is Alzheimer's disease.

31. A pharmaceutical composition for the treatment or prevention of a condition involving cognitive impairment or a susceptibility to the condition, comprising an effective amyloid-beta precursor processing-inhibiting amount of a GPCR antagonist or inverse agonist.

32. A composition according to claim 31, wherein said GPCR inverse agonist is an aryloxydithiourea, its pharmaceutically acceptable salts, hydrates, solvates, or prodrugs thereof in admixture with a pharmaceutically acceptable carrier.

33. Use of an agent as claimed in claims 17-25 for the preparation of a medicament for the inhibition of amyloid-beta precursor processing.

34. Use of an agent as claimed in claims 17-25 for the preparation of a medicament for the treatment or prevention of a condition involving cognitive impairment or a susceptibility to the condition.

35. Use of a GPCR antagonist or inverse agonist for the treatment or prevention of a condition involving cognitive impairment or a susceptibility to the condition.

36. Use as claimed in claim 34 or 35, wherein the condition is Alzheimer's disease.

37. Use as claimed in claim 35 or 36, wherein the GPCR inverse agonist is an aryloxydithiourea, its pharmaceutically acceptable salts, hydrates, solvates, or prodrugs thereof in admixture with a pharmaceutically acceptable carrier.

Figure 1A

1/7




Figure 1B




Figure 2

2/7





Figure 3:

3/6


A

Dose response screening of GPR3 in HEK293 APPwt cl 29 cells
Amyloid beta 1-42




B

Dose response screening of GPR3 in HEK293 APPwt cl29 cells
Amyloid beta 1-40




C

Dose response screening of GPR3 in HEK293 APPwt cl29 cells
Amyloid beta y-42



4/7

D



E





Figure 4

5/7



Essential multiple sequence parameters :

Algorithm: ClustalW

Substitution matrix : Blossum62

Gap open penalty: 10

Gap extension penalty : 0.05

	*	20	*	40	*	60	*	80	
GPR3 :	--	MWGC	S L A S	NVN	VSSV PAET	PT	PAA	PL SPKAWD	VVL C 48
GPR6 :	MN	SAASL	ND S VVAEECAAAATAA	GPDTGEWPPAAALAGC	NNSLELS	SQLSAGPPGLLLA	WDVLLC	: 80	
GPR12 :	--	MN D KVNLSGLRD	DAAA--	AENISVSSRVP	AVE	PEPEL-V	WDIVLC	: 52	
		6 G p a l Ag		g aa6g ga g		P	Lp vnpWD66LC	:	

	*	100	*	120	*	140	*	160						
GPR3 :	SGTLV	ENALVV	IIVG	PFR	PMFLLVGSLA	ADLLAG	GLV	FAAVFCIGSAE	LVL	VG	LAMA	F	ASIGSLL	: 128
GPR6 :	SGTVIAG	ENALVV	LIAS	PRT	PMFVLVGSLA	TADLLAG	CGLI	FQY	VPS	T	LL	VGFL	F	FASVSLL : 160
GPR12 :	TSGTLL	ENAI VVL	LI FHNPS	R	PMFLLIGSLA	ADLLAG	GLITNE	AY	LQS	ATKLV	IG	I	F	ASVC SLL : 132
	SGT66sc	ENA6Vva6I	tPalRa	PMF6L6GSLA	ADLLAG	GL6lhFvf	5l6 Se	sL6t6G	6vasF	AS6	SLL			

GPR3	:	A	I	T	V	D	R	Y	L	S	L	Y	A	L	T	Y	S	T	T	V	R	V	M	L	G	A	L	L	G	L	L	P	V	L	A	W	N	C	L	D	G	L	C	G	V	V	Y	P	L	K	V	V	L	A	F	V	F	I	:	208									
GPR6	:	A	I	T	V	D	R	Y	L	S	L	Y	A	L	T	Y	S	R	T	L	L	G	V	H	L	L	A	T	W	T	V	L	L	G	L	L	P	V	L	W	N	C	L	A	E	R	A	A	C	V	V	P	L	A	R	S	L	L	A	F	V	F	I	:	240				
GPR12	:	A	I	T	V	D	R	Y	L	S	L	Y	A	L	T	Y	H	S	T	V	F	V	M	L	V	W	T	I	C	L	G	L	L	P	V	M	W	N	C	L	R	D	E	C	V	V	P	L	K	N	A	I	L	S	F	L	F	M	F	A	L	:	212						
		A	I	T	V	D	R	Y	L	S	L	Y	A	L	T	Y	S	e	r	T	6	t	t	y	6	6	L	a	W	g	s	6	L	G	L	L	P	V	6	g	W	N	C	L	t	C	S	V	V	r	P	L	4	n	h	a	6	L	s	a	F	f	m	6	F	g	6		

GPR3 : ML ILYA IC IV CRHA QIALQ RHLL PA H TRKGIATLAVVLGAF A WLPF VY LL AHSPP LYTYL TLLPATY 320
 GPR6 : MLHL Y RICQ VVWRHA QIALQ QHCL PPHLA TRKGVCTLAVVLG F GAS WLPFAIY VV SHEDPAVYTY TLLPATY 320
 GPR12 : ML ILY IC IV MRHA QIALQ QHFL T H TTRKGVSTLAIILG F A WMPF LYSLIA YTYPS LYTY TLLPATY 320
 MLqLY qIC 6V RHahQIALQ H La shyvaTRKG6 TLA66LGtFaAcW6PFt6YC66gd P 6YTYaTLLPATY

GPR3	:	NSMINPIIYAFRNQD	VQKVL	AV	CCSSS	I	R	RSPS	V		360
GPR6	:	NSMINPIIYAFRNQ	IQR	L	LLC	C	FQS	V	R	RSPSEV	: 330
GPR12	:	NSLINPVIYAFRNQ	IQR	L	C	CIPSSL	AQRARSPS	V			: 362
		NS6INP6IYAFRNQe6Q4aLw16cCgC				Sk6pfRsrRSPsdv					: 334

Figure 6

7/7

