AUSTRALIAN PATENT OFFICE
AMENDED

LODGED AT SUB-OFFICE
10 FEB 1986
Melbourne

COMMONWEALTH of AUSTRALIA

PATENTS ACT 1952

APPLICATION FOR A STANDARD PATENT

594036

I
We PLANT GENETIC SYSTEMS N.V., of
Kunstlaan,
Avenue des Arts 46,
1040 Brussels,
Belgium.


hereby apply for the grant of a Standard Patent for an invention entitled:

"MODIFYING PLANTS BY GENETIC ENGINEERING TO COMBAT OR CONTROL INSECTS"

which is described in the accompanying ~~provisional~~ complete specification.

Details of basic application(s):—

Number	Convention Country	Date
692,759	United States of America	18th January 1985



The address for service is care of DAVIES & COLLISON, Patent Attorneys, of 1 Little Collins Street, Melbourne, in the State of Victoria, Commonwealth of Australia.

Dated this 10th day of FEBRUARY 19 86.

.....
(a member of the firm of DAVIES & COLLISON for and on behalf of the Applicant).

To: THE COMMISSIONER OF PATENTS

APPLICATION ACCEPTED AND AMENDMENTS ALLOWED 13.12.89

Davies & Collison, Melbourne and Canberra.

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

DECLARATION IN SUPPORT OF CONVENTION OR NON-CONVENTION APPLICATION FOR A PATENT

Insert title of invention

In support of the Application made for a patent for an invention entitled : "MODIFYING PLANTS BY GENETIC ENGINEERING TO COMBAT OR CONTROL INSECTS"

Insert full name(s) and address(es) of declarant(s) being the applicant(s) or person(s) authorized to sign on behalf of an applicant company.

I ~~We~~ ANNE-MARIE BOUCKAERT,
of PLANT GENETIC SYSTEMS, N.V.,
of Kunstlaan,
Avenue des Arts 46,
1040 Brussels,
Belgium

do solemnly and sincerely declare as follows :-

Cross out whichever of paragraphs 1(a) or 1(b) does not apply

1(a) relates to application made by individual(s)
1(b) relates to application made by company; insert name of applicant company.

1. ~~(a) I am / We are the applicant ..... for the patent~~

or (b) I am authorized by

PLANT GENETIC SYSTEMS, N.V.

the applicant..... for the patent to make this declaration on ~~their~~ its behalf.

Cross out whichever of paragraphs 2(a) or 2(b) does not apply

2(a) relates to application made by inventor(s)
2(b) relates to application made by company(s) or person(s) who are not inventor(s); insert full name(s) and address(es) of inventor(s)

2. ~~(a) I am / We are the actual inventor..... of the invention~~

or (b) Henri Marcel Jozef de Greve, of Lambeauxlaan 11, 1060 Brussels, Belgium; Maria Benita Leonor Fernandez Salgado of Almeda 44, Iguala, Guerrero, Mexico; Marc Charles Ernest van Montagu of De Strassartstraat 120, 1050 Brussels, Belgium; Mark Albert Vaeck of Aarschotsebaan 4, 2959 Zemst, Belgium; Marcus Florent Oscar Zabeau of Erpelsteeg, 70, 9000 Gent, Belgium Jan Jozef August Leemans of Ellebochten 38, 9210 Heusden, Belgium and Hermanus Fransiscus Paulus Hofte of Rozemarijnstraat 34, 9000 Gent, Belgium.

~~is~~ are the actual inventors..... of the invention and the facts upon which the applicant.....
~~is~~ are entitled to make the application are as follows :-

State manner in which applicant(s) derive title from inventor(s)

The applicant would if a patent were granted on an application made by the said actual inventors be entitled to have the patent assigned to it.

Cross out paragraphs 3 and 4 for non-convention applications. For convention applications, insert basic country(s) followed by date(s) and basic applicant(s).

3. The basic application..... as defined by Section 141 of the Act ~~were~~ was made
in United States of America on the 18th January, 1985
by Henri Marcel Jozef de Greve, Maria Benita Leonor Fernandez
in Salgado, Marc Charles Ernest van Montagu, Mark Albert Vaeck,
by and Marcus Florent Oscar Zabeau.
~~in~~ ..... ~~on the~~ .....
~~by~~ .....

4 The basic application..... referred to in paragraph 3 of this Declaration ~~were~~ was the first application, ... made in a Convention country in respect of the invention the subject of the application

Insert place and date of signature

Declared at Sent this August day of 21, 1989

Signature of declarant(s) (no attestation)

Note Initial all alterations.

DAVIES & COLLISON, MELBOURNE and CANBERRA.

(12) PATENT ABRIDGMENT (11) Document No. AU-B-52458/86
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 594036

(54) Title
TRANSFORMED PLANT PRODUCING B. THURINGIENSIS TOXIN

International Patent Classification(s)
(51)[4] C07H 021/04 A01H 001/00 A01H 005/10 A01N 063/02
A01N 065/00 C07G 017/00 C12N 001/20 C12N 001/21
C12N 005/00 C12N 005/10 C12N 015/00 C12N 015/32
C12P 021/02

(21) Application No. : 52458/86 (22) Application Date : 17.01.86

(30) Priority Data

(31) Number (32) Date (33) Country
692759 18.01.85 US UNITED STATES OF AMERICA

(43) Publication Date : 24.07.86

(44) Publication Date of Accepted Application : 01.03.90

(71) Applicant(s)
PLANT GENETIC SYSTEMS N.V.

(72) Inventor(s)
HENRI MARCEL JOZEF DE GREVE; MARIA BENITA LEONOR FERNANDEZ SALGADO; MARC CHARLES ERNEST VAN MONTAGU; MARK ALBERT VAECK; MARCUS FLORENT OSCAR ZABEAU; JAN JOZEF AUGUST LEEMANS; HERMANUS FRANSISCUS PAULUS HOFTE

(74) Attorney or Agent
DAVIES & COLLISON, MELBOURNE

(56) Prior Art Documents
AU 33473/84 874101 C12N 15/00
AU 39018/85 589115 C12N 15/00
AU 48044/85 C12N 15/00

(57) Claim

1. A chimeric gene capable of being expressed in cells of a plant as an insecticidal Bacillus thuringiensis polypeptide toxin and of thereby rendering the plant insect-resistant; the chimeric gene comprising:

(a) a first DNA fragment that encodes a Bacillus thuringiensis insecticidal crystal protein which has been truncated towards a trypsin cleavage site of the protein; and

(b) a promoter region and a 3' non-translated region containing a polyadenylation signal; the first DNA fragment being under the control of the promoter region; the promoter and 3' non-translated regions allowing the first DNA fragment to be expressed in the cells;

the chimeric gene being free of: (i) an ATG

triplet between the promoter region and an initiation codon of the first DNA fragment; and

(ii) a T-DNA oncogenic tumor-inducing gene.

13. A chimeric gene capable of being expressed in cells of a plant as an insecticidal Bacillus thuringiensis polypeptide toxin and of thereby rendering the plant insect-resistant; the cells being free of a T-DNA oncogenic tumor-inducing gene; the chimeric gene comprising:

(a) a first DNA fragment that encodes a Bacillus thuringiensis insecticidal crystal protein;

(b) a promoter region and a 3' non-translated region containing a polyadenylation signal; the first DNA fragment being under the control of the promoter region; the promoter and 3' non-translated regions allowing the first DNA fragment to be expressed in the cells; and

(c) a second DNA fragment which encodes an enzyme capable of being expressed in the cells and the expression of which can be identified in the cells; the second DNA fragment being fused to the first DNA fragment so that the first and second DNA fragments encode a fusion polypeptide;

whereby an identification of expression of the second DNA fragment in the cells provides an identification of expression of the first DNA fragment in the cells;

the chimeric gene being free of an ATG triplet between the promoter region and an initiation codon of the first DNA fragment.

25. A method for protecting a plant against an insect pest comprising the step of: providing genomes of cells of the plant with the chimeric gene of any one of claims 1-20, whereby the cells can express an insecticidal Bacillus thuringiensis polypeptide toxin.

27. A seed of an insect-resistant plant which is capable of being transformed by Agrobacterium, containing in its genome the chimeric gene of any one of claims 1-20.

28. A tissue of an insect-resistant plant which is capable of being transformed by Agrobacterium, having cells containing in their genome the chimeric gene of any one of claims 1-20, whereby the cells are capable of expressing an insecticidal Bacillus thuringiensis polypeptide toxin.

29. A plant cell containing in its genome the chimeric gene of any one of claims 1-20 whereby the cell is capable of expressing an insecticidal Bacillus thuringiensis polypeptide toxin.

594036

COMMONWEALTH OF AUSTRALIA

PATENT ACT 1952

COMPLETE SPECIFICATION

(Original)

FOR OFFICE USE

Class Int. Class

Application Number: 52458/86.
Lodged:

Complete Specification Lodged:
Accepted:
Published:

Priority:

Related Art:

This document contains the amendments made under Section 49 and is correct for printing.

Name of Applicant: PLANT GENETIC SYSTEMS N.V.



Address of Applicant: Kunstlaan,
Avenue des Arts 46,
1040 Brussels,
Belgium.

Actual Inventor(s):
Henri Marcel Jozef DE GREVE
Maria Benita Leonor Fernandez SALGADO
Marc Charles Ernest van MONTAGU
Mark Albert VAECK
Marcus Florent Oscar ZABEAU
Jan Jozef August LEEMANS
Hermanus Fransiscus Paulus HOFTE

Address for Service: DAVIES & COLLISON, Patent Attorneys,
1 Little Collins Street, Melbourne, 3000.

Complete Specification for the invention entitled:

"MODIFYING PLANTS BY GENETIC ENGINEERING TO COMBAT OR CONTROL INSECTS"

The following statement is a full description of this invention, including the best method of performing it known to us :-

MODIFYING PLANTS BY GENETIC ENGINEERING TO COMBAT OR CONTROL INSECTS

This invention is concerned with the use of genetic engeineering techniques in the modification of plants.

The invention provides chimeric genes, plasmid vectors, plants, plant seeds, plant tissue, cell cultures and plasmids arising in and from the use of such genetic engineering techniques in the modification of plants, and methods and compositions for controlling or combating insects in agriculture or horticulture.

The invention is particularly directed towards introduction and integration of a chimeric gene coding for a polypeptide toxin produced by Bacillus thuringiensis or having substantial sequence homology to a toxin gene described below in plant cells and obtaining an insect controlling or combating level of expression of said polypeptide toxin intracellularly by transformed plant cells and their progeny.

Recombinant DNA technology is currently used to genetically engineer certain microorganisms such as bacteria and yeast to synthesize specific proteins. Genetic engineering of higher organisms within the present state of technology requires that one or a few cells be genetically engineered from which the entire organisms can develop. Among higher organisms, the cells of certain plants exhibit excellent regeneration

capability and therefore are considered potentially good material for the genetic engineering of such plants. Furthermore, in higher plants, a known system is available to introduce foreign DNA into the plant genome. This system is provided by the tumor inducing plasmid from the gram negative soil bacterium Agrobacterium tumefaciens. Agrobacterium can genetically transform plant cells by stably integrating T-DNA, a well defined fragment of the Ti plasmid, into the plant cell genome. Recently, important progress has been made to facilitate the use of the Ti plasmid as a vector for plant genetic engineering. Small directly repeated sequences which flank the T-DNA (Border sequences) have been found to play a key role in the T-DNA integration. Nononcogenic Ti plasmid vectors have been constructed from which oncogenic tumor genes have been removed by an internal deletion in the T-DNA. These Ti plasmids still contain the border sequences and consequently transfer T-DNA without tumor induction. An example of such a Ti plasmid derived vector from plant genetic engineering is pGV3850 which contains a substitution of the internal T-DNA gene by the commonly used cloning vehicle pBR322. Several procedures have been developed to regenerate infected plants which contain the pGV3850. pGV3850 with the pBR322 sequences present in its T-DNA is an efficient acceptor plasmid for gene transfer experiments in plant cells. Indeed, genes cloned in pBR322 like plasmids are transferred to Agrobacterium and inserted via homologous recombination into the pGV3850 T-DNA in a single experimental step.

Another major advance in the development of plant engineering technique is the use of plant regulatory sequences to express chimeric genes in plants. In

general, these chimeric genes contain a promoter region derived from a gene which is naturally expressed in plant cells, the sequence to be expressed, and preferentially a 3' non-translated region containing a polyadenylation site of a gene which is naturally expressed in plant cells. For example, using the nopaline synthase promoter and bacterial antibiotic resistance genes, dominant selectable markers for plant cells have been constructed.

Although certain chimeric genes have now successfully been expressed in transformed plant cells, such expression is by no means straightforward. Various lines of evidence indicate that the level of expression of the foreign genes of non-plant origin not only varies greatly in different transformed tissues but are in general very low. Such low levels of gene expression could be due to several reasons: first, incomplete transcription of the gene resulting from inadvertent transcription termination signals; second, inefficient processing of the messenger RNA; third, impaired transport of the messenger RNA from the nucleus to the cytoplasm; fourth, instability of the cytoplasm messenger RNA; fifth, inefficient translation of the cytoplasm messenger RNA; and sixth, instability of the protein due to its susceptibility to plant specific proteins. Consequently, the successful transformation of plant cells using vectors such as those described above is not necessarily predictable prior to attempting a desired transformation.

Engineering of differentiated plant cells and their progeny to express the Bt2 polypeptide and/or a truncated version thereof and/or a polypeptide having substantial sequence homology thereto is far more difficult than other genes such as antibiotic

resistance genes or other plant genes such as thaumatin due to one or more of the following: (1) the large size of the Bt2 toxin, even in its truncated form; (2) the particular properties of the Bt2 polypeptide (such as, but not limited to, solubility of the polypeptide); (3) the potential toxicity of the Bt2 polypeptide toward the plant cells; or (4) the Bt2 polypeptide synthesized in plant cells and their progeny must retain substantially the same properties as the crystal protein synthesized in bacteria.

Bacillus thuringiensis (referred to at times herein as B.t.) bacteria includes approximately 19 known varieties that produce polypeptide toxins which form parasporal crystals during sporulation. The crystal protein made by B.t. is toxic to the larvae of certain insects. The toxins produced by a particular variety exhibit strong insecticidal activity, against certain Lepidoptera and/or Ceoleoptera and/or Diptera larva. See e.g., Tyrell D. J. et al., J. Bacteriology, (81) 145 (No. 2): p. 1052-1062. When ingested by insect larvae, the crystals are solubilized and processed in the insect midgut to yield at least one active polypeptide toxin which is believed to act on the midgut cell membrane. Studies have revealed that individual crystal polypeptides exhibit insecticidal activity. Yamamoto, T. et al., Current Microbiology, (83) 9: p. 279-284; Yamamoto, T. et al., Arch. Biochem. Biophysics, (83) 227: (No. 1): p. 233-241; Lilley, M. et al., J. Gen. Microbiol., (80) 118: p. 1-11; Bulla, L. A. et al., J. Biol. Chem., (81) 256 (No. 6): p. 3000-3004.

The toxic activity of the crystal polypeptide produced by Bacillus thuringiensis varieties is highly specific to particular insect species and is recognized as safe to higher vertebrates.

Preparations containing the crystals are used commercially as a biological insecticide. For example: Bactospeine, distributed by Biochem Products Ltd., Dipel Abbott Laboratories; and Thuricide, Sandoz AG. The efficacy of preparations obtained from bacterial hosts is, however, limited as adequate control of pests requires repeated and precisely timed applications. In addition, costs associated with the production of such preparations have made it difficult for them to compete effectively with other commercially available products, such as pyrethroid derivatives.

Molecular genetics studies have demonstrated that at least some polypeptide toxins produced by Bacillus thuringiensis are encoded by plasmids. Stahly, D. P. et al., (1978), Biochem. Biophys. Res. Commun., 84, p. 581-588; Debaboc, V. G. et al., (1977), Genetika, 13, p. 496-501. Genes encoding toxic crystal polypeptides from different B.t. strains have been cloned and expressed in other bacterial hosts. (Schnepf & Whiteley, PNAS (81) 78: 2993-2897. Klier, A. et al., EMBO J. (82) 1 (No. 7), p. 791-799; Adang et al., Gene, (36), p. 289, 1985; Schnepf et al., J. Biol. Chem., (20), p. 6264, 1985; Shibano et al., Gene, (34), 1985.

Considering the major importance of plants both for consumption and for production of valuable products, it would be highly desirable to genetically modify plants such that plant cells could synthesize

polypeptide toxins substantially similar to those toxins produced by Bacillus thuringiensis, without adverse effects to the plants. By stably integrating exogenous DNA fragments coding for polypeptide toxins produced by Bacillus thuringiensis into the plant cell genome and obtaining an insect controlling level of expression of said exogenous DNA fragments in plants, plant cells and their progeny so transformed would thereby become resistant to certain insect pests. Plant cells and their progeny genetically engineered in this way would provide an economically advantageous substitute to existing commercial varieties by substantially obviating the need for specific chemical or biological insecticides, and provide a more reliable means of controlling particular insect pests, while retaining normal morphological characteristics.

It is one object of this invention to provide novel chimeric genes coding for the polypeptide toxin produced by Bacillus thuringiensis, or coding for a polypeptide toxin having substantial sequence homology to a toxin gene described herein. The chimeric genes' plant regulatory sequences direct expression in transformed plant cells.

Another object of present invention is to provide novel hybrid plasmid vectors containing said chimeric genes that allow the introduction and integration and expression of said chimeric genes in a plant cell genome.

A further object of the present invention is to provide a process for preparing genetically transformed plant cells comprising the transformation of plant

cells with said hybrid plasmid vectors containing said chimeric genes.

Other objectives, features and advantages of the present invention will become apparent to those skilled in the art from the following description taken in conjunction with the accompanying drawings.

In accordance with the present invention, there are provided: chimeric genes capable of being expressed in differentiated plant cells comprising:

(a) a DNA fragment comprising a promotor region derived from a gene which is naturally expressed in a plant cell; and

(b) at least one DNA fragment coding for a polypeptide toxin produced by Bacillus thuringiensis or having substantial sequence homology thereto.

Said chimeric genes include those where DNA fragment (b) codes for a Bt2 protein, an insecticidally active truncated Bt2 protein, a DNA fragment having substantial sequence homology to Bt2 or the truncated Bt2, or where DNA fragment (b) is fused to a DNA fragment (c) coding for an enzyme capable of being expressed in differentiated plant cells and permitting identification of plant cells expressing DNA fragment (b) where said DNA fragments (b) and (c) encode a fusion polypeptide.

Also in accordance with the present invention there are provided: hybrid plasmid vectors comprising:

(a) a DNA fragment substantially homologous with that portion of a Ti plasmid essential for transfer of a T-region of a Ti plasmid to a plant cell genome (the virulence region of a Ti plasmid);

(b) at least one DNA fragment which delineates a DNA fragment to be integrated into a plant cell genome (the border sequences of the T-DNA portion of a Ti plasmid; where only one border sequence is present, preferably it is the right border sequence); and

(c) at least one chimeric gene comprising:
  (i) a DNA fragment comprising a promotor region derived from a gene which is naturally expressed in a plant cell; and
  (ii) at least one DNA fragment coding for a polypeptide toxin produced by Bacillus thuringiensis or at least one DNA fragment having substantial sequence homology thereto.

Said chimeric genes include those where DNA fragment (b) codes for a Bt2 protein, an insecticidally active truncated Bt2 protein, a DNA fragment having substantial sequence homology to Bt2 or the truncated Bt2, or where DNA fragment (b) is fused to a DNA fragment (c) coding for an enzyme capable of being

expressed in differentiated plant cells and permitting identification of plant cells expressing DNA fragment (b) where said DNA fragments (b) and (c) express a fusion polypeptide.

Further, in accordance with the present invention, there are provided: intermediate plasmid vectors containing at least one chimeric gene, said chimeric gene comprising:

(a) a DNA fragment comprising a promotor region derived from a gene which is naturally expressed in a plant cell; and

(b) at least one DNA fragment coding for a polypeptide toxin produced by Bacillus thuringiensis, or at least one DNA fragment having substantial sequence homology thereto.

Said chimeric genes include those where DNA fragment (b) codes for a Bt2 protein, an insecticidally active truncated Bt2 protein, a DNA fragment having substantial sequence homology to Bt2 or the truncated Bt2, or where DNA fragment (b) is fused to a DNA fragment (c) coding for an enzyme capable of being expressed in differentiated plant cells and permitting identification of plant cells expressing DNA fragment (b) where said DNA fragments (b) and (c) express a fusion polypeptide.

Further, in accordance with the present invention, there are provided insecticidal compositions and methods of using transformed plant cells and their progeny.

Still further in accordance with the present invention there are provided: plants which include in their cells genome and express the chimeric gene as described above; and plant seeds which are capable of germinating into a plant which expresses the chimeric gene as described above.

Transformed plant cells and their progeny intracellularly express a polypeptide toxin substantially similar to the polypeptide toxins produced by Bacillus thuringiensis and are substantially toxic to certain insects. Transformed plant cells and their progeny may be used in controlling said insects.

Brief Description of the Drawings

Figure 1 is a photograph showing a 7.5% SDS PAGE stained with Coomassie Blue.

Track 1: B.t. kurstaki crystal protein preparation;

Track 2: B.t. berliner crystal protein preparation;

Track 3: Molecular weight markers

a: phosphorylase B (92,500 dalton);

b: bovine serum albumin (66,200 dalton);

c: ovalbumin (45,000 dalton); and
d: carbonic anhydrase (31,000 dalton).

Figure 2 is a schematic diagram of plasmid pEcoR251. The EcoRI endonuclease gene (EndRI) is fused to the P_R promotor (P_R) and contains a unique BglII cloning site. Amp: beta-lactamase gene.

Figure 3 shows restriction enzyme maps of the inserts present in 4 immunopositive partial Sau3A digest clones of B.t. berliner 1715 plasmid DNA cloned in pEcoR251.

Figure 4 is a photograph showing a 7.5% SDS PAGE stained with Coomassie Blue.

Track 1: B.t. kurstaki crystal protein preparation (identicial with Figure 1, Track 1);

Track 2: B.t. berliner crystal protein preparation (identical with Figure 1, Track 2);

Track 3: Total lysate of E. coli K514 containing the Bt200 plasmid; and

Track 4: Control (total lysate of E. coli K514 without Bt200 plasmid).

Figure 5 is a photograph showing the results of immunoblotting experiment using a rabbit anti-B.t. kurstaki crystal serum.

Track 1: B.t. berliner crystal protein preparation;

Track 2: B.t. kurstaki crystal protein preparation;

Track 3: Total lysate of E. coli K514 harboring the pBt200 plasmid; and

Track 4: Control (total lysate of E. coli K514 without pBt200.

Figure 6 is a photograph showing the results of an immunoblotting experiment using a rabbit anti-kurstaki crystal serum (A) and a rabbit anti-berliner crystal serum (B). Part C shows a Coomassie staining of the 7.5% SDS PAGE after the blotting procedure (the same gel used for blotting shown in Part A).

Track 1: Bt2 protein (purified as described in Section 5-1);

Track 2: B.t. berliner crystal proteins; and

Track 3: B.t. kurstaki crystal proteins.

Figure 7 is a photograph showing a Coomassie staining of an SDS PAGE.

Track 1: Totally lysate of E. coli K514 harboring pBt200;

Track 2: Purified Bt2 protein prepared from the E. coli K514 harboring pBt200.

Figure 8 is a graph showing the results of an ELISA experiment. Binding curves of Bt2 protein and

solubilized B.t. crystal proteins using a goat anti-B.t. crystal serum as goat-antibody and a mouse anti-Bt2 serum as first antibody.

Figure 9 is a graph showing the results of an ELISA experiment showing reactivity of the different anti-berliner crystal monoclonal antibodies with:

(A) total berliner crystal proteins; and

(B) purified Bt2 protein.

Figure 10 shows a comparison of N-terminal amino acid sequences of the 130 Kd crystal proteins:

1) deduced from the DNA sequence published by Wong et al., J. Biol. Chem. 258, p. 1960-1967 (1983) (termed B.t. W);

2) determined for the Bt2 protein.

Figure 11 summarizes the immunological detection of polypeptides using Western blotting with a rabbit anti-B.t. berliner crystal serum. Polypeptides are encoded by pBt200 derivatives containing various deletions generated by restriction enzyme cleavage as indicated in the figure.

Figure 12 A - shows the restriction map of the HpaI-NdeI fragment containing the entire Bt2 gene indicated as a box. B - shows the sequenced regions of the Bt2 gene. Boxes represent the stretches which have been sequenced from each strand. C - shows sequencing strategy. Restriction fragments were end labeled with

polynucleotide kinase, strand isolated and sequenced using the Maxam and Gilbert method. The arrows indicate the length of the region sequenced in each experiment.

Figure 13 shows the DNA sequence of the complete Bt2 gene indicating the open reading frame (position 21 to 3605) and the corresponding deduced amino acid sequence (1155 amino acids). The amino acid sequence of the Bt2 protein which was experimentally determined is indicated by a line above the corresponding amino acids.

Figure 14 shows a comparison of the deduced amino acid sequences of the Bt2 gene (berliner) with the deduced sequences from three other B.t. crystal protein genes, cloned from other B.t. strains:

B.t. kurstaki HD73 (Adang et al., Gene 36, p. 289, 1985)

B.t. kurstaki HD1 (Schnepf et al., J.B.C. 20, p. 6264, 1985)

B.t. sotto (Shibano et al., Gene 34, p. 243, 1985)

In the latter 3 sequences, only those amino acids are represented that differ from those present in the Bt2 sequence at the homologous position. Eventually, gaps were introduced (marked by " - ") in order to align the sequences.

Figure 15 is a schematic outline of the construction of the Bt2 gene cassette pHD160.

Figure 16 is a schematic representation of the different Bt2 gene cassettes.

Figure 17 shows the experimental strategy used for the construction of plasmid pLB10. Also shown here is the structure of plasmids pLK54 and pLK57, described in J. Botterman et al. (in preparation).

Figure 18 shows the construction and structure of pLBKm25.

Figure 19 shows the strategy used to construct Bt:NPTII fusions and Bt2 deletions.

Figure 20 is a schematic representation of the different Bt2 3' end deletion mutants, used in the mapping of the 3' end of the minimal toxin encoding fragment. Arrows represent the positions of the 3' ends.

Figure 21 is a photograph showing the results of an immunoblotting experiment using a rabbit anti-berliner crystal serum. Samples analyzed are total extracts from Bt2 deletion clones specified in Figure 20 and in Section 7.

Figure 22 shows the 3' end points of deletion clones pLB834 and pLB879 on the Bt2 sequence, used to delineate the minimal gene fragment encoding an active toxin. Also shown is the deduced amino acid sequence and the position of a putative trypsin cleavage site.

Figure 23 is a schematic representation of the construction of the Bt2:NPTII fusion gene cassettes

pLBKm23, pLBKm33 and pLBKm14. Also represented are the 5' upstream sequences of the Bt2:NPTII fusions in the different constructs (sequences corresponding to a BamHI site are underlined).

Figure 24 is a schematic representation of different Bt:NPTII fusion gene cassettes.

Figure 25 is a photograph showing the results of a NPTII assay as described by Reiss et al. (Gene, 30, p. 217, 1984). The samples analyzed are the supernatants of cell extracts of bacterial clones producing NPTII or different Bt2-NPTII fusion proteins.

23 means K514 (lambda) (pLBKm23)
860 means K514 (lambda) (pLBKm860)
865 means K514 (lambda) (pLBKm865)
NPT means HB101 (lambda dv) (a gift from Julian Davis, formerly of Biogen)

Figure 26 shows the approximate positions of the 3' ends of the Bt sequences in different deletions and Bt:NPTII fusions (indicated by arrows).

Figure 27 shows the strategy used for the adaptation of the Bt2 and the Bt2:NPTII cassettes for expression in plant cells.

Figure 28 shows the DNA sequences at the junction between the promotor regions and the coding sequence of the Bt gene cassettes as they are present in the different engineered Ti plasmids. Sequences derived from the original promotor regions and from the coding

sequence of the Bt2 gene are underlined. Some relevant restriction enzyme sites which have been involved in the assembly of the chimeric genes are indicated. The ATG initiation codon is boxed.

Figure 29 is a schematic representation of the construction of pHD208 as described in Section 8 Example 2.

B: BamHI, Hp: HpaI, H: HindIII, E: EcoRI, Bg: BglII.

Figure 30 is a schematic representation of the construction of pGV831: pGV831 has been constructed by R. Deblaere, Lab of Genetical Virology, Free University Brussels, Belgium. It is a derivative of pGV700, as described in European Patent Application No. 83112985.3. Recombinant DNA techniques used followed Maniatis et al., Molecular Cloning (1982), Cold Spring Harbor Laboratory.

The HindIII fragment present in pGV700 was subcloned into pGV600 (Leemans et al., J. Mol. Appl. Genet., 1, 149-164, 1981). Recombinant plasmid pGV742 was isolated as a Cb^R Cm^S Tc^S recombinant. An internal deletion was created in pGV742 by digestion with BamHI and recirculization. This produced pGV744. An internal deletion was created in pGV744 by digestion with EcoRI and recirculization to yield pGV749. The HindIII-NruI fragment from pGV749 was cloned in pGV710. pGV710 had been digested with EcoRI, the 5' protruding end filled in using DNA polymerase and had been subsequently digested with HindIII. The resulting plasmid pGV815 was isolated as a Sm^R, Cb^R recombinant. Both the EcoRI site and the HindIII site of pGV815 were

removed by digestion with these enzymes and by filling in the protruding ends with DNA polymerase, followed by recirculization. Finally, a chimeric gene containing the nopaline synthase promotor and the neomycin phosfotransferase gene from Tn5 was isolated as a BclI-BamHI fragment from pKC7/:nos and was cloned in the BglII site from pGV825. The Sp^R, Km^R recombinant plasmid pGV831 was obtained.

Figure 31 is a schematic representation of the T region of Ti-plasmid pGV3850 and of the intermediate vector pHD205. The crossed lines indicate the regions which were involved in cointegration of pGV3850 with pHD205 to produce pHD1050. The T region of hybrid Ti plasmid pHD1050 is represented.

H: HindIII

Bt: chimeric Bt2 gene under control of the nopaline synthase promotor

nos: nopaline synthase gene

Ap, Km: genes encoding ampicillin and kanamycin resistance

Figure 32 is a schematic representation of the T region of Ti plasmid pGV2260 and of the intermediate vector pHD208. The crossed lines indicate the regions which were involved in cointegration of pGV2260 with pHD208 to produce pHD1076. The T-region of hybrid Ti-plasmid pHD1076 is represented.

1: vector fragment

2: T-DNA border region

3: Bt2 gene cassette

4: Pssu promotor fragment

5: Pnos promotor fragment
6: neomycin phosfotransferase gene cassette
7: T-DNA border region
8: vector fragment
Black triangles represent T-DNA border regions
Ap, Sp, Km: genes encoding respectively ampicillin, spectinomycin and kanamycin resistance
Pnos: nopaline synthase promotor
Pssu: small subunit of ribulose biphosphate carboxylase promotor
B.t.: Bt2 gene cassette

Figure 33 shows schematic representations of the different intermediate expression vectors.

Figure 34 is a graph showing the results of an ELISA assay of tobacco callus tissue transformed with C58Cl Rif^R pHD1076, as described in Section 11 Example 1. The coating antibody is goat anti-B.t. crystal serum. Rabbit anti-Bt2 is used as first antibody.

Numbers 1 to 14 are transformed calli.

The optical density (O.D.) corresponding to a level of 4 ng Bt2 protein per gram of tissue, determined in a reconstruction experiment, is indicated in the figure.

Figure 35 is a graph showing the results of an ELISA assay of tobacco callus tissue transformed with C58Cl Rif^R pHD1076, as described in Section 11 Example 1. The coating antibody is goat anti-B.t. crystal serum. Rabbit anti-Bt2 serum is used as first antibody.

Numbers 1 to 21 are transformed calli.

The O.D. value corresponding to a level of 4 ng Bt2 protein per gram of tissue, determined in a reconstruction experiment, is indicated in the figure.

Figure 36 is a graph showing the results of an ELISA assay of tobacco callus tissue transformed with C5 8ClRif^R pHD1076, as described in Section 11.2, Example 1. Coating antibody is goat anti-B.t. crystal serum. Different monoclonal antibodies were used as first antibody. Reactivity with untransformed SR1 callus tissue (used as a negative control) is also shown.

Figure 37 is a description of the experimental protocol used for the preparation of callus tissue extracts, used for the immunological detection of Bt2 expressed in this callus.

Figure 38 is a graph showing the growth rate of 1st instar M. sexta larvae feeding on leaves from transformed tobacco plants obtained as described in Section 10, Example 5. Open bars represent the number of larvae (on a total of 20 larvae tested) that went to the L2 stage after 3 days of feeding.

Figure 39 is a graph showing complete growth rate curves over a 4 day period, for M. sexta larvae feeding on leaves of transformed tobacco (data are from same experiments as those represented in Figure 38). The represented values are the numbers of larvae that were in the L2 stage at a certain point in time (per plant, 20 larvae were tested). C_1-C_4 are control plants

(transformed with the Pnos-NPTII gene only). The other numbers (N20-1, N20-46) refer to individual plants putatively transformed with pGS1110.

Figure 40 shows the DNA sequences of the P35S-1 and P35-2 promotor fragments derived from cauliflower mozaic virus Cm4-184 (Gardner et al., 1981, Nucl. Acid Res., 9, 2871-2888).

Figure 41 is a schematic representation of the construction of pGSH50.

Figure 42 is a schematic representation of the construction of pGV1500.

Figure 43 is a schematic representation of the construction of pGSH150 and pGSH151.

Figure 44 is a schematic representation of the construction of pAGS007 from Pssu301 whild type gene.

As used herein, the term "polypeptide" should be understood as meaning an intact protein or fragment(s) thereof.

"Plant" should be understood as referring to a multicellular differentiated organism capable of photosynthesis including angiosperms (monocots and dicots) and gymnosperms. "Plant cells" should be understood as referring to one or more cells derived from a plant. "Plant cell progeny" should be understood as referring to any cell or tissue derived

from plant cells including callus; plant parts such as stems, roots, fruits, leaves or flowers; plants; plant seed; pollen; and plant embryos. "Chimeric gene" should be understood as a hybrid DNA segment comprising a regulatory signal essential for transcription referred to as a promotor, fused to at least one structural gene sequence coding for a specific polypeptide. "Substantial sequence homology" should be understood as referring to either: a DNA fragment having a nucleotide sequence sufficiently similar to another DNA fragment to produce a protein having similar properties; or a polypeptide having an amino acid sequence sufficiently similar to another polypeptide to exhibit similar properties. "Identification" should be understood as referring to selection or scoring of cells harboring and expressing the desired gene. Selectable markers permit growth (selection) under otherwise lethal conditions such as kanamycin resistance (Km^R). Scorable markers add an identifiable trait (scoring) foreign to non-tranformed cells. "Naturally expressed gene" should be understood as meaning a DNA fragment whether originally part of a plant's genome or introduced by agents such as bacteria or viruses which produces RNA, protein or both in the plant in the absence of human intervention.

A chimeric gene may also include a nontranslated DNA fragment positioned on the 3' side (downstream) of the structural gene sequence, which in turn may include a regulatory signal referred to as a polyadenylation signal preferably derived from a gene which is naturally expressed in plants.

A naturally expressed gene includes a 3' non-

translated region which in turn includes a polyadenylation signal, both of which code for the corresponding messenger RNA (mRNA) regions. These corresponding mRNA regions are located on the 3' side of a stop codon in a monocistronic mRNA. The 3' non-translated region of mRNA is believed to be involved in the processing, stability and/or transport of the mRNA. This 3' non-translated region of mRNA is also believed to contain a sequence of bases, polyadenylation signal, which is recognized by an enzyme in the cell. This enzyme adds a substantial number of adenosine residues to the mRNA molecule to form a poly-A "tail" on the mRNA.

Generally, the process used to arrive at the present invention is described in European Patent Application Publication No. 0116718 entitled "Process for the Introduction of Expressible Genes into Plant Cell Genomes and Agrobacterium Strains Carrying Hybrid Ti Plasmid Vectors Useful for this Process." The introduction and integration of one or more chimeric genes coding for polypeptide toxins produced by Bacillus thuringiensis or having substantial sequence homology to Bt2 (see Figure 13) into a plant cell genome is achieved by:

(1) isolation of at least one DNA fragment from Bacillus thuringiensis coding for a polypeptide toxin by digestion of bacterial DNA and inserting the mixture of DNA fragments obtained into a cloning vehicle harbored in a bacterial host; and

(2) identification of bacterial clones harboring DNA fragments coding for said polypeptide toxin; and

(3) characterization of the structure of the DNA fragment coding for said polypeptide toxin; and

(4) removal of unwanted DNA sequences flanking the desired DNA fragment; or

(5) synthesis of a DNA fragment having substantial sequence homology and exhibiting a similar structure to a DNA fragment coding for Bt2; or

(6) construction of a DNA fragment containing the DNA fragment from (4) fused to a DNA fragment encoding an identification polypeptide to produce a fusion polypeptide; and

(7) insertion of said DNA fragment from (4) or (5) or (6) into plasmid vectors under the control of plant regulator sequences harbored in a bacterial host; and

(8) introduction of plasmids from (7) by conjugation (or mobilization) in a bacterial host harboring suitable helper plasmids; and

(9) conjugation of bacterial clones from (8) to Agrobacterium tumefaciens harboring an acceptor Ti plasmid vector; and

(10) identification of Agrobacterium tumefaciens which contain the desired chimeric gene; and

(11) contacting plant cells with Agrobacterium tumefaciens from (10); and

(12) identification of transformed plant cells from appropriate culture media; and

(13) immunological detection of Bt2 antigens present in extracts from transformed plant cells; and

(14) propagating transformed plant cells to regenerate a differentiated plant.

It is contemplated that cloning vectors and bacterial host strains other than those described below in the Examples can be used. Ti-based vectors like pGV3850 into which recombinant plasmids integrate before transfer to plant cells are known as cis-type vectors. There are also Ti-based vector systems in which the recombinant plasmids do not integrate into the resident Ti plasmid or in which large portions of the naturally occurring Ti plasmid are deleted. These binary-type systems, Hoekema et al., Nature, Vol. 303, 179 (1983), or mini-Ti plasmids, Framond et al., Biotechnology, Vol. 1, 262 (1983), have also been shown to introduce DNA into plant cells. These plasmids contain a border sequence (at least one, preferably two) flanking the gene to be introduced into plants. A marker which is selectable or scorable in plant cells is useful but not essential. Such plasmids are capable of autonomous replication in A. tumefaciens and need not integrate into a resident Ti plasmid. Virulence functions needed to effect transfer to DNA, such as the chimeric genes of the present invention, to plant cells

can be provided in trans. Hoekema et al., Nature, Vol. 303, 179 (1983). See also Fraley, R.T. et al., Biotechnology, Vol. 3, 629 (1985); and Klee et al., Biotechnology, Vol. 3, 637 (1985).

A. tumefaciens is not the only means of introducing genes into plants. DNA can be introduced by physical means such as electroporation or chemical means such as polyethylene glycol (PEG) fusion. It is believed any technique which introduces DNA, such as the chimeric genes of the present invention, can be used. Further, RNA viral vectors which introduce an RNA copy of an insecticidal chimeric gene may also be used.

Further, plasmid vectors containing plant regulatory sequences other than those described below in the Examples can be used. For example, enhancers can be included before, or after, or in such proximity to the chimeric gene to exert their function.

Plant cells transformed with the novel plasmid vectors of the present invention may then be cultured on suitable medium, preferably selectable growth medium, and plants which express the polypeptide toxin may be regenerated from the resulting callus. Subsequent generations of plant cells and their progeny should also exhibit expression of the polypeptide toxin.

Transformed plant cells and their progeny should express a polypeptide toxin substantially similar to polypeptide toxins being produced by Bacillus thuringiensis or a DNA fragment having substantial sequence homology to Bt2.

The present invention contemplates that the hybrid

plasmid transformation vectors may be used to develop plant cells and their progeny exhibiting insect resistant properties. It is contemplated that plants, particularly dicotyledonous plants, other than those described below in the examples can be transformed such as cotton, sugarbeet, soybean, rape and vegetables such as cabbage, lettuce and beans. Transformed plant cells and their progeny are protected against certain insect pests by expressing an insect controlling amount of polypeptide toxin. By controlling amount is meant a toxic (lethal) or combative (sublethal) amount of polypeptide toxin. In general in this description the words controlling and combating are used interchangeably and no distinction is intended except where, from the context, it is clear that a differentiation between a lethal and non-lethal dose is intended to be drawn. The transformed plants should be morphologically normal and may be cultivated in their usual manner for consumption and/or production of products. Further, said transformed plants should substantially reduce the need for chemical or biological insecticides directed toward combatting Lepidoptera and Coleoptera larvae. Since the genes coding for the polypeptide toxin are stably integrated in the plant cell genome and are thus heritable, seed obtained from said transformed plants should also produce plants expressing the polypeptide toxin at substantially the same level and thereby also be protected against certain insect pests.

In addition, it is contemplated that transformed plant cells and their progeny could be used to control certain insect pests by applying to the pests and/or the habitat of said pests (i.e., the locus to be protected, for example, to a growing crop or to an area where a crop is to be grown) an effective (controlling) amount of transformed plant matter alone or together with other components.

By way of example, but not limitation, transformed plant cells and their progeny could be used alone or as one component in a formulation or composition. For practical applications, plant cells and their progeny could be used as the active material or as a solid carrier in conventional pesticide compositions and formulations. Such compositions and formulations may also contain adjuvants such as surfactants and stabilizers. Examples of such composition and formulations include pastes, dusting powders, wettable powders, granules, baits and aerosol compositions.

Compositions and formulations are prepared in a known manner. The amount of transformed plant matter to be used depends on a variety of factors, for example, the kind of pest, the formulation or composition used, the state of the crop infected with the pest and the prevailing weather conditions. In general, transformed plant cells and their progeny may constitute from about 0.1 to about 100% by weight of the composition or formulation and preferably from about 1.0 to about 99% by weight.

Known insecticidal, fungicidal, biocidal, herbicidal and fertilizer compounds and compositions compatible with the polypeptide toxins may be included as components in the above described compositions and formulations to provide additional benefits and advantages.

In practice, certain Lepidoptera or Coleoptera larvae attempt to feed on transformed plants. A small amount of transformed plant matter is ingested. The ingested matter is processed in the insect midgut

yielding the active polypeptide toxin which acts on the midgut cell membrane to kill or inhibit growth of the pest.

Also in practice, when used alone or as one component in a formulation or composition, certain Lepidoptera and/or Coleoptera larva attempt to feed on plants treated with said formulations or compositions. A small amount of treated plant matter is ingested. The ingested matter containing the formulation or composition is processed in the insect midgut yielding the polypeptide toxin which acts on the midgut cell membrane to kill or inhibit growth of the pest.

Engineering of the present invention was generally accomplished as follows:

1. Isolation and preparation of antibodies specific for B.t. crystal polypeptides

   A. Isolation of Bacillus thuringiensis (B.t.) crystal polypeptides

   B. Preparation of antibodies (polyclonal and monoclonal) against B.t. crystal polypeptides

2. Preparations of B.t. Gene Bank

   A. Preparation of total DNA or plasmid DNA from B.t., preferably plasmid DNA

   B. Partial digestion of the purified DNA with a suitable restriction enzyme

C. Cloning DNA fragments into a suitable *E. coli* plasmid expression vector

3. Isolation of recombinant plasmids containing B.t. polypeptide genes

A. Screening of the transformed *E. coli* cells with anti-B.t. crystal protein serum

B. Identification and isolation of bacterial clones expressing the polypeptide

4. Characterization of Bt2 protein

A. Purification of the polypeptide encoded by the cloned B.t. gene

B. Testing to confirm that polypeptide expressed by clones is immunologically the same as B.t. crystal polypeptide

C. Testing to confirm that polypeptide expressed by clones is insecticidal

5. Mapping and subcloning of Bt2, including restriction enzyme analysis, subcloning and DNA sequence determination

6. Construction of toxin gene cassette including removal of undesired flanking ATG triplets preceding the initiator ATG and addition of suitable restriction enzyme cleavage sites using synthetic oligonucleotide linkers

7. Construction of Intermediate Vectors

8. Construction of Hybrid Ti Plasmids

9. Engineering of Plants

A. Identification of transformed plant tissues producing the toxin using the immunoassays and quantification of the toxin levels produced

B. Regeneration of plants from tissues

10. Detection of Bt2 toxin in engineered plants

11. Determine toxicity of engineered plants toward insects

Different types of chimeric genes (promotor-gene fusions), have been used to genetically transform plant cells, and basically 3 different types of plant specific promotors can be distinguished:

Promotors:

1. Ti plasmid derived promotors (Pnos, PTR at times referred to herein as PTR2)

2. Plant promotors (Pssu pea, Pssu301)

3. Plant virus promotors (P35S from cauliflower mozaic virus)

Types of chimeric genes:

1. Type I:
Straight promotor-gene fusions in which the entire Bt2 coding sequence is inserted behind the promotor fragment. Examples are: Pnos-Bt2 (pHD1050, pHD1060), Pssu pea-Bt2 (pHD1076), PTR2-Bt2 (pGS1161), Pssu301-Bt2 (pGS1181), P35S-1-Bt2 (pGS1261), P35S-2-Bt2 (pGS1271). Some of the constructs do not contain the intact 5' untranslated region of the original transcript (Pnos, Pssu pea), but others do (PTR, Pssu301).

2. Type II:
Chimeric Pssu-Tp-Bt2 gene fusion in which the Bt2 gene is fused to the transit peptide (Tp) sequence of the small subunit of RuBisco and expressed under the control of the Pssu promotor. In this case a fusion protein preferably is made from the natural translation initiation signal of the ssu gene. Van Den Broeck et al. (1985) demonstrated the transport of the bacterial NPTII protein into plant chloroplasts using a fusion between the transit peptide of the ssu of RuBisco and the NPTII coding region. In view of these results, we constructed the chimeric gene Pssu-Tp:Bt2. Both the Pssu promotor and the transit peptide (Tp) fragment were derived from the pea gene used by Van Den Broeck et al. (1985). The DNA sequence at the junction site is shown in Figure 28. It is worth mentioning that the original 5' untranslated region of the pea m-RNA is maintained in Pssu-Tp:Bt2, so that the chimeric gene is

translated from the genuine ssu translation initiation site (pHD1080).

3. Type III:
Straight promotor-gene fusions in which only part of the Bt2 coding sequence is used ("truncated Bt2"). Fragments of the Bt2 sequence still encoding an active toxin are inserted behind the plant specific promotors: The toxic polypeptides produced in the plant cells using these constructs should have biological and biophysical properties distinct from the intact Bt2 protein such as specific toxic activity or solubility.

Examples: pGS1162, pGS1163, pGS1262.

4. Type IV:
Straight promotor-gene fusions in which a Bt:NPTII fusion gene (also referred to at times at Bt2:NPTII) is inserted behind the promotor. Fusion genes were constructed, consisting of a fragment of the Bt2 coding sequence (still encoding an active toxin) fused to the coding sequence of the NPTII enzyme. The Bt:NPTII fusion genes used here, specify stable fusion proteins comprising amino terminal parts of the Bt2 protein fused to an intact Neomycin phosphotransferase (NTPII) enzyme. These fusion proteins have a specific toxicity comparable to the intact Bt2 protein and retain neomycin phosphotransferase enzyme activity. Thus, expression of the Bt:NPTII fusion proteins in plant cells allows

direct selection for the production of this protein by isolating Kanamycin resistant (Km^R) transformed cells. Furthermore, the level of Km^R should be directly correlated to the amount of protein synthesized. Thus, selection of plants resistant to a high level of Kanamycin should identify, among all possible transformations, those which produce high levels of the toxic fusion protein. Further, expression of the fusion protein by a Bt:NPTII fusion gene might have other desirable properties such as stability in plant cells; for example, mRNA may be more stable. Differences in results obtained with these Type IV fusion genes might be due to intrinsic differences in the properties of the fusion protein expressed as compared to the intact Bt2 protein.

Examples: pGS1110, pGS1151, pGS1152, pGS1171, pGS1251, pGS1253, pGS1281.

Alternative constructions of the desired transformation vectors described herein are also contemplated. For example, plant specific exogenous promotors other than those disclosed herein may be used. The use of a different exogenous promotor sequence may be useful for directing expression of the inserted exogenous DNA in a regulated fashion. Examples of other types of regulation which may be used include tissue-specific expression (leaves, roots, stems or flowers); and inducible expression (temperature, light or chemical factors). Additionally, given the DNA sequence data coding for

the polypeptide endotoxins produced by Bacillus thuringiensis, a transformation vector could be constructed containing an artificially created DNA fragment substantially similar to the Bt2 DNA fragment described herein. This artificially created DNA fragment could then be used to transform plants in substantially the same manner as described herein.

The following examples are offered by way of illustration and should not be construed as limiting the scope of the present invention.

EXPERIMENTAL

1. Isolation of Bacillus thuringiensis (B.t.) crystal proteins

Crystals were isolated and purified from spore preparations of strains B.t. berliner 1715 (received from Dr. A. Klier, EMBO J. 1, No. 7, p. 791-799, 1982) and B.t. var. kurstaki, (J. Bacteriol. 145, No. 2, p. 1052, 1981) as described by Mahillon and Delcour (J. Microbiol. Meth., Vol. 3, No. 2, p. 69-76, 1984). The crystal proteins were solubilized by incubating the purified crystals at 37°C for 2 h in 0.2 M thioglycolate, 0.1 M $NaHCO_3$ pH 9.5, whereafter the insoluble material was removed by low speed centrifugation. This procedure solubilizes more than 80% of the proteins present in the crystals. Solubilized crystal proteins were analyzed on 7.5% sodium dodecyl sulfate polyacrylamide gel (SDS PAGE). The crystal protein preparation from Bt berliner contained at least two major protein species in the high molecular weight region (apparent MW of 140 and 130 Kd) and a less abundant protein of about 120 Kd, as revealed by staining the gels with Coomassie brilliant

blue (Figure 1). The solubilized crystal proteins of strain kurstaki showed one major 130 Kd protein band and a weaker 60 Kd band (Figure 1).

These solubilized crystal proteins exhibited a strong toxic activity towards third instar larvae of the cabbage butterfly Pieris brassicae (L.D. 50 values of 0.5 ng/larva for kurstaki and 0.65 ng/larva for berliner) using the toxicity assay described in section 5.2 below.

2. Preparation of antibodies specific for B.t. crystal proteins

2.1 Polyclonal antisera

Antisera against B.t. crystal proteins (berliner 1715 and kurstaki) were prepared separately in rabbits and mice. Antiserum against B.t. crystal proteins (kurstaki) prepared in goat was received courtesy of Dr. L. Bulla, University of Idaho. To the best of applicant's knowledge and belief, the antiserum was prepared by known procedures substantially similar to those described for rabbit and mouse.

Rabbits were injected subcutaneously with 0.5 mg of a solubilized crystal protein preparation (.25 ml dialysed qagainst PBS pH 7.4) mixed with an equal volume of complete Freund's adjuvant (CFA). After three months, the rabbits received another injection of the same type of preparation, and three weeks later blood samples were taken. BALBc mice were injected intraperitoneally with 100 ug of crystal protein solution, mixed with CFA (1/1 vol.). Four to six weeks later they received a booster injection of 50 ug crystal protein PBS, and four days later blood samples

were taken. Antigen reactivity of the sera was confirmed by immunodiffusion tests (Ouchterlony assay). A strong crossreaction between berliner 1715 and kurstaki crystal protein preparations was observed, indicating that they contained antigenically related components.

Some of the mice were sacrificed and the spleens removed asceptically for cell fusion experiments (see 2.2).

2.2 Monoclonal antibodies

Although not essential for the identification of toxin expressing clones as described herein, hybridomas producing monoclonal antibodies against B.t. crystal proteins were generated following the procedure originally described by Koehler and Milstein (Nature 256: 495-497, 1975). Monoclonal antibodies were used as an additional and more specific means of determining toxin presence in bacterial clones and plant cells.

Spleen cells from immunized BALBc mice (see 2.1) were fused with the SP2/0 myeloma cell line (Shulman, M. et al., Nature 276, p. 269, 1978). Cells were plated at 3.10^5 per well in microtiter plates and 10-14 days later the supernatants were screened for the presence of anti-crystal protein antibodies using an enzyme immuno assay (Engvall and Pesce, Scand. J. Immunol., suppl. 7, 1978) with alkaline phosphatase labelled goat anti-mouse immunoglobulin as the second antibody (Sigma, A-5153). Approximately 4% of the wells were positive for the antigen (crystal protein). Positive clones were subcloned twice by limiting dilution. Positive subclones were selected, grown up and their

culture supernatants containing the monoclonal antibodies were collected. A total number of 17 hybridoma cell lines producing monoclonal antibodies reactive with B.t. berliner crystal proteins were generated.

3. Construction of a gene bank from plasmid DNA of B.t. strain berliner 1715

Kronstad et al., J. Bacteriol., 54, p. 419-428 (1983) reported that B.t. berliner 1715 contains two related toxin genes which are both located on plasmids. Intact endotoxin genes were isolated from a gene bank from total B.t. berliner 1715 plasmid DNA using partial Sau3A digests of plasmid DNA. B.t. berliner 1715 cells were grown in LB medium (Miller, Experiments in Molecular Genetics, (1972), Cold Spring Harbor Laboratory, New York) overnight at 37°C. Plasmid DNA was isolated from B.t. berliner 1715 using the denaturation-renaturation method described by Kronstad et al., J. Bacteriol., 54, p. 419-428 (1983). Analysis of the plasmid DNA on 0.5% agarose gels revealed that this plasmid DNA preparation contained several different plasmid species present in different molar concentrations. To construct the gene bank thirty ug of plasmid DNA was partially digested with Sau3A at 37°C in a total volume of 500 ul. 100 ul samples were taken after respectively 10, 20, 30, 45 and 60 minutes of incubation and phenol-chloroform extracted. The Sau3A digested DNA was size fractionated on a 10 to 40% sucrose gradient, and the size of the DNA fragments in the different fractions was estimated on a 0.8% agarose gel. The fractions containing DNA in the 6-10 Kb size range were pooled

and ligated to BglII digested pEcoR251 vector DNA. The pEcoR251 plasmid is a derivative of plasmid pBR322 in which the EcoRI-PuvII fragment has been replaced by a chimeric EcoRI endonuclease gene which is fused to a P_R promotor fragment derived from plasmid pLK5 (Zabeau and Stanley, EMBO Journal, 1, 1217-1224 (1982)) as depicted in Figure 2. The pEcoR251 contains a unique BglII site in the EcoRI endonuclease gene, where insertion will inactivate the gene. The pEcoR251 vector is a suicide vector similar to the positive-selection cloning vehicle pSCC31 described by Cheng and Modrich (J. Bacteriol. 154, 1005-1008, 1983). Sau3A DNA fragments were ligated into BglII digested pEcoR251. Recombinant plasmids were selected by transforming the ligation mix into competent E. coli K514 cells (Colson et al., Genetics 52, p. 1043-1050, 1965) as described by Dagert and Ehrlich, Gene 6 (1980), 23-28. Cells were plated on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York), supplemented with ampicillin (100 ug/ml).

Several gene banks were constructed each containing between 600 and 1500 recombinant clones. Analysis of the recombinant plasmids present in 12 randomly chosen clones confirmed that in each gene bank at least 10 out of the 12 clones contained inserted fragments with sizes ranging from 5 to 15 Kb.

4. Isolation of recombinant plasmids containing B.t. crystal protein genes

The colonies of the gene bank were screened for bacteria producing crystal proteins using a rabbit serum raised against purified B.t. berliner crystal

proteins (see Section 2.1 above). The procedures used are slightly modified from Helfman et al., (PNAS 80: 31-35 1983). Bacterial colonies, grown on 150 mm square Petri dishes, were replica plated on nitrocellulose sheets (Schleicher & Schuell, 0.45 um, 401196). Sheets were soaked in 0.1 M NaOH until colonies lysed. The sheets were then air dried, washed in phosphate buffered saline (PBS) pH 7.4 for 30 minutes and incubated overnight at 4°C or for 2 hours at room temperature with gentle agitation in PBS containing 1% crude ovalbumine (Sigma, A-5253). Nitrocellulose sheets were rinsed in PBS and incubated for 2 hours in rabbit anti-crystal serum diluted in PBS, 1% ovalbumin, 0.2% Triton X-100, at room temperature with gentle agitation. After additional washing the sheets were incubated with peroxidase-labeled goat anti-rabbit antibodies (Sigma, A-6154) (2 hours at room temperature). After extensive washing with PBS/0.2% Triton, the sheets were reacted with substrate solution (substrate was 4-chloro-1-naphtol, Sigma, C-8890). Positive colonies developed as dark blue dots. Using serial dilutions of purified crystal protein solution, the detection limit of this test was estimated to be 1-10 ng protein/ml. In total, 4 different immuno-positive clones were isolated from a gene bank of 1250 clones. Plasmid DNA was prepared from each clone following the procedure of Zabeau and Stanley, EMBO J., 1, 1217-1224, 1982. Primary restriction maps were constructed by performing single and simultaneous restriction enzyme digestions. Comparison of the restriction maps for the enzymes EcoRI, EcoRV, BamHI, SacI, MluI and Pst1 (See Figure 3) revealed that all 4 plasmids carried DNA fragments of different sizes which showed a clear region of overlap. These results show

that the Bt2 gene must be encoded by a 4.2 Kb region common in the 4 different recombinant plasmids. For further study we subcloned a 7.5 Kb BamHI-PstI fragment from clone B12 (see Figure 3) into the plasmid PUC8 (J. Viera and J. Messing, Gene, 19, p. 259-268, 1982) and this recombinant plasmid was termed pBt200.

5. Characterization of the Bt2 protein

5.1 Identification of a 130 Kd crystal protein encoded by pBt200

The E. coli strain K514 containing the pBt200 plasmid (see Section 4), showed a strong positive reaction in the colony assay. This was further confirmed using an enzyme linked immuno sorbent assay (ELISA) (Engvall & Pesce, 1978, Scand. J. Immunol., Suppl. 7). For the ELISA screening the following procedure was used: Flexible polyvinyl microtiter plates, coated with goat anti-Bt crystal protein antibodies, were incubated with lysate of bacterial colonies (lysates were obtained by freeze-thawing pelleted cells, followed by incubation in 0.1 M NaOH for 15 minutes, and subsequent neutralization with 0.1 M HCl). After washing, a diluted rabbit or mouse anti-B.t. crystal protein serum was added. After 1-2 hours incubation, plates were washed and incubated with rabbit or mouse anti-B.t. crystal serum (appropriately diluted). After 1-2 h incubation, plates were washed and incubated with goat anti-rabbit or anti-mouse IgG antibodies, alkaline phosphatase labeled (Sigma A-8025, A-5153). After incubation and washing the substrate (p-nitro phenyl phosphate, Sigma, 104-105) was added and the reaction monitored by measuring

optical density (O.D.) at 405 nm. Detection limit of the test for purified solubilized crystal protein was estimated to be in the range of 0.1-1 ng/ml.

Total cell protein extracts of E. coli strains harboring pBt200 were analyzed on SDS PAGE. An intense new protein band was visible in the high molecular weight range, corresponding to a M.W. of about 130 Kd. This band was not present in K514 cells containing the pUC8 vector plasmid without insert. This new protein also comigrated on SDS PAGE with one of the major crystal proteins of B.t. berliner and with the major crystal protein of Bt kurstaki (see Figure 4). The relationship of this protein, which was termed Bt2, with B.t. crystal proteins was confirmed by immunoblotting. Western blotting experiments were carried out using both rabbit anti-Bt kurstaki crystal serum and rabbit anti-Bt berliner crystal serum. Strong reaction of the Bt2 protein with both antisera was observed (see Figures 5 and 6).

These results demonstrate that the cloned Bt2 gene codes for one of the crystal proteins of B.t. berliner which is immunologically related to the 130Kd crystal protein of B.t. kurstaki.

The amount of positively reacting material in bacterial extracts was quantitated using an ELISA assay. Using purified crystal protein as a standard, the amount of crystal protein produced in E. coli harboring pBt200 was estimated to be in the range of 5-10% of the total cell protein content. The estimate agrees well with the observed intensity of the band the Bt2 protein band on SDS PAGE after staining with

Coomassie blue. To further characterize the 130Kd protein encoded by the pBt200 plasmid (termed Bt2 protein) we developed a rapid purification procedure, taking advantage of the relative insolubility of the protein. 5 g cells obtained from a 2 litre overnight culture of K514 (pBt200) were resuspended in 50 ml 50 mM TRIS pH 7.9, 50 mM EDTA, 15% sucrose, treated with lysozyme (100 ug/ml), sonicated (30 minutes at 400 watts in a Labsonic 1510), mixed with 200 ml of PBS, pH 7 containing 2% Triton X100 and incubated for 30 min. on ice. The lysate was centrifuged at 15000 g and the supernatant was discarded. The pellet containing the Bt2 protein was resuspended in the same buffer and the procedure was repeated. Whereafter the pellet was washed twice with 200 ml PBS. To solubilize the Bt2 protein the pellet was resuspended in 50 ml extraction buffer 0.2 N thioglycolate and 0.1 M $NaHCO_3$, pH 9.5 for 2 hr. at 37°C. An efficient (>90%) and selective solubilization of Bt2 protein was obtained in this way (Figure 7).

These semi-purified protein preparations were used for further studies. Antisera were raised against Bt2 protein in rabbits and mice using a similar immunization protocol as described in Section 2.1. These antisera reacted equally well with solubilized crystal proteins from B.t. berliner and kurstaki as with Bt2 itself, in the ELISA assay described above (Figure 8 shows results with the mouse serum).

A similar positive reaction was observed using antibodies purified, from anti-Bt crystal serum, by affinity chromatography on an immunoadsorbent of Bt2 (Bt2 protein coupled onto CNBr activated Sepharose 4B,

Pharmacia). These antibodies also reacted in Western blotting with a 130Kd protein present in both B.t. berliner and kurstaki crystals.

Finally, in the ELISA, 9 out of the 17 monoclonal antibodies raised against total B.t. berliner crystal proteins, were also reactive with the Bt2 protein. (Code numbers: 1F6, 167, 4D6, 4F3, 8G10, 10E3, 1.7, 4.8, C73) (Figure 9). The same 9 antibodies were also reactive with B.t. kurstaki crystal proteins.

In general both the Bt2 protein and the major 130 Kd crystal proteins from B.t. require alkaline pH and the presence of reducing reagents for complete solubilization. Also they both precipitate at pH 4-5.

Thus, the cloned gene product Bt2 exhibits biochemical properties similar to those of the major 130 Kd crystal protein from B.t. berliner and B.t. kurstaki and is immunologically related to these crystal proteins.

The Bt2 protein was purified further by DEAE-ion exchange chromatography and by Sephacryl gel filtration. The amino-terminal sequence of this purified protein was determined with the use of a gas-phase sequencer (Applied Biosystems), operated according to Hewick et al., J. Biol. Chem., 256, 7990-7997, 1981).

The sequence of the first 20 N-terminal amino acids was found to be substantially identical to the N-terminal sequence deduced from the DNA sequence of a

cloned B.t. kurstaki gene, Wong et al., J. Biol. Chem., 258 (3), 1960-1967 (1983) (Figure 10).

5.2 Insect Toxicity of the Bt2 protein

Crystals from B.t. are known to be particularly toxic against larvae of certain Lepidoptera species. In order to test whether Bt2 protein exhibited a similar toxic activity, toxicity tests were performed on larvae of the cabbage butterfly Pieris brassicae. Protein solutions of known concentration, expressed as ppm (1 ppm = 1 ug/ml) were serially diluted in water. Small discs (0.25 cm^2) were cut from fresh cabbage leaves and on each disc 5 ul of a test solution was applied. Discs were air dried and each disc was placed in a vial containing one larva. Third instar larvae were obtained from a synchronized culture of P. brassicae. During a 10 h period before moulting, these larvae were incubated in separate vials in the absence of food. Immediately after moulting they were given one leaf disc. When the first disc was consumed, the larva was offered a fresh disc without sample. For each sample dilution, 50 larvae were tested. Feeding and viability were monitored every 24 h up to 120 h. As can be seen from Table 1, Bt2 sample preparations exhibited similar degrees of toxicity for P. brassicae larvae as solubilized crystals from B.t. berliner 1715.

To test the effect of sublethal doses of Bt2 toxin on the growth of P. brassicae larvae, the following experimental design was used: cabbage leaves were dipped in a solution containing a known concentration of Bt2 protein (0.01-1 ppm) and dried. Groups of 100 third instar larvae (from synchronized cultures) were

fed on Bt2 coated leaves. The leaves were regularly replaced by new leaves treated in the same way. Growth of the larvae was followed over a period of seven days, which corresponds to the time period needed to develop from 3rd to 5th instar. As can be seen from the results presented in Table 2 the Bt2 protein induced a significant growth inhibition in P. brassicae larvae at doses that were sublethal. Growth inhibition was evident at a concentration of 0.01 ppm which corresponded to 2.67 ng protein/gram leaf. During the first 48 h the larvae feeding on leaves coated with 0.01 ppm ate 3.6 cm^2 of leaf (83 mg) and consequently ingested about 0.22 ng of Bt2 protein. At this time, 93% of the larvae were still in the L3 stage while only 33% of the control larvae were in this stage. Thus an inhibitory effect on growth can be observed with toxin doses that are significantly below the LD_{50} values (1.65 ng/larva, see Table 1).

These results indicate which levels of Bt2 protein synthesis must be reached in transformed plant cells in order to express insect resistance against P. brassicae. A level of 2.7 ng Bt2 protein/g tissue is sufficient to retard the growth of the larvae. This might already be adequate as such to halt a devastating spread of the larvae in the field. Toxicity assays with Bt2 protein were also performed on larvae of the Tobacco Hornworm, (Manduca sexta). As shown in Table 3, Bt2 protein is slightly more toxic than total berliner crystal proteins (100% mortality at 12.5 ng/cm^2). In addition, significant growth inhibition is observed at sublethal doses (2.5 ng/cm^2): 4.4 mg body weight after 7 days, as compared to 30.5 mg for control larvae. Due to the fact that Manduca is fed on an artificial diet, (ref:

Bell, R. A. & Joachim, F. G. (1976) Ann. Entomol. Soc. Am., 69: 365-373), results are expressed somewhat differently, namely as ng toxin applied per cm^2 of agar medium.

6. Characterization of the Bt2 gene

To locate the Bt2 toxin gene on the 7.2 Kb BamHI-PstI fragment of the pBt200 plasmid a series of deletions were made in the 7.2 Kb DNA fragment with respectively HpaI, KpnI and IbaI. The proteins encoded by these deletion plasmids were analyzed immunologically, using the ELISA technique and Western blotting (also referred herein to as immunoblotting) (Towbin et al., PNAS, USA, 76: 4350-4354, 1979 and Burnette, W. N. An. Biochemistry, 112, p. 195-203, 1981).

The results (diagrammed in Figure 11) can be summarized as follows: (1) Deletion of the HpaI fragment results in the synthesis of an intact Bt2 protein at a lower level. This finding indicates that the deletion only affects the regulatory region but not the structural part of the gene. (2) Deletion of the Kpn fragment results in a approximately 70 Kd protein fragment still detectable by immunoblotting. (3) The Xba deletions closer to the 5' end do not give rise to protein fragments detectable by Western blotting procedure. These results show that the intact gene encoding the 130 Kd protein is located on a 4.3 Kb HpaI-PstI fragment (see Figure 11). To determine the precise structure of the Bt2 gene, the complete nucleotide sequence of the 4,060 base pairs (bp) HpaI-NdeI fragment was determined by the Maxam and Gilbert sequencing method. The sequer ing strategy used is diagrammed in Figure 12.

The proposed nucleotide sequence was confirmed primarily by sequencing the complementary strand. Examination of the sequence revealed the presence of a single large open reading frame starting at position 141 and ending at position 3605, which could code for a protein of 1,155 amino acids with a molecular weight of 127 Kd. This is in agreement with the molecular weight of 130 Kd of the Bt2 protein as determined by SDS polyacrylamide gel electrophoresis. Furthermore, the amino-terminal amino acid sequence predicted from the nucleotide sequences agrees with the amino acid sequence determined on the purified Bt2 protein (see Figures 10 and 13).

The complete amino acid sequence of the Bt2 toxin shows extensive homology with the deduced amino acid sequences from 3 other B.t. crystal proteins from which the genes were cloned and sequenced: B.t. kurstaki HD1 (Dipel) (Schnepf et al., J. Biol. Chem., 20, p. 6264, 1985), B.t. kurstaki HD73 (Adang et al., Gene, 36, p. 289, 1985) and B.t. sotto (Shibano et al., Gene, 34, p. 243, 1985).

Comparison of these other B.t. sequences with our Bt2 at the amino acid level (Figure 14) reveals that they encode similar but distinct proteins, showing regions of striking homology but also stretches which diverge significantly.

7. Construction of the "Toxin Gene" cassettes

7.1 Construction of a cassette carrying the intact Bt2 gene

Inspection of the DNA sequence of the Bt2 gene revealed that the 160 bp region immediately upstream of the ATG translation initiation codon contains 5 ATG

triplets. Translation of eucaryotic genes usually starts at the first AUG in the message (In RNA U replaces T). These AUG triplets might act as initiator AUG's and could be recognized preferentially over the genuine Bt2 initiation codon and could thus reduce the level of expression in transformed plant cells. Moreover, these AUG's are in other reading frames and would give rise to nonsense polypeptides. To prevent initiation of translation at these AUG triplets, the sequences upstream of the Bt2 gene were removed by exonucleolytic treatment, prior to insertion of the pBt2 gene in the Ti expression vectors. To this end, deletion derivatives of the pBT200 plasmid in which upstream sequences were deleted up to the initiator ATG were constructed. Thirty-five ug of pBt200 DNA was digested with HpaI and treated with 6 units of Bal31 exonuclease (Biolabs, New England) for 1, 1.30, 2, 2.30 and 3 minutes in 300 ul of 12 mM $MgCl_2$, 12 mM $CaCl_2$, 0.6 M NaCl, 1 mM EDTA and 20 mM tris-HCl - pH 8.0, at 30°C. One ug of Bal31-treated molecules of each reaction were ligated at 4°C to 0.13 ug phosphorylated BamHI linkers (Biolabs, New England) with 2 units T4 DNA ligase in a total volume of 20 ul.

After the T4 ligase was inactivated at 68°C for 10 minutes, each ligation mix was digested with 20 units BamHI for 1 h at 37°C. Subsequently, 50 ng DNA was recircularized with 0.1 unit T4 DNA ligase in a total volume of 100 ul for 20 h at 4°C.

One-fifth of this ligation mixture was transformed into competent E. coli K514 cells (Colson et al., Genetics 52 (1965), 1043-1050) as described by Dagert and Ehrlich, Gene 6 (1980), 23-28. Cells were

plated on LB medium (Miller, Experiments in Molecular Genetics, (1972), Cold Spring Harbor Laboratory, New York), supplemented with carbenicillin (100 ug/ml).

The deletion end points in the plasmids were first analyzed by measuring the size of the newly generated EcoRI fragments of the recombinant plasmids on a 2% agarose gel. The nucleotide sequences of the exact deletion end points in plasmids with deletions ending just before the start of the Bt2 gene were determined. Clone pHD100 has a deletion ending 8 bp before the initiator ATG and removes all upstream non-initiator ATG's. Clone pBa3.3 contains the BamHI linker fused to the 4th bp of the coding sequence and clone pBa23-3 contains the Bam linker fused to bp -33.

In a second engineering step, the non-coding sequences at the 3' end of the toxin gene were deleted using Bal31 exonuclease (Biolabs, New England). Thirty ug of pHD100 plasmid DNA were digested with NdeI and treated with Bal31 exonuclease for 3, 4, 5, 6 and 8 minutes at 30°C in buffer. At each time interval, 60 ul aliquots (each containing 6 ug of Bal31 treated DNA molecules) were removed. After addition of phosphorylated BglII linkers (Biolabs, New England) to the Bal31 treated DNA molecules, the DNA molecules were recircularized with 0.1 U T4 ligase overnight at 4°C. The ligation mixture was transformed into competent E. coli K514 cells (Colson et al., Genetics 52 (1965), 1043-1050) as described by Dagert and Ehrlich, Gene 6 (1980), 23-28. Cells were plated on LB medium (Miller, Experiments in Molecular Genetics, (1973), Cold Spring Harbor Laboratory, New York) supplemented with carbenicillin (100 ug/ml). After determination of the

size of the deletion in several plasmids, using restriction enzyme digestion and agarose gel electrophoresis, pHD160, pHD162, pHD163 were retained for further experiments. In pHD160, the BglII site is positioned at approximately 300 bp behind the TAA stopcodon of the Bt2 gene; in pHD162 the BglII is at approximately 250 bp behind TAA; and in pHD163 the BglII is at position 3342 (bp) in the Bt2 coding sequence. Construction of pHD160 is schematically diagrammed in Figure 15. In this way, we constructed toxin gene cassettes carrying the Bt2 gene on a BamHI-BglII fragment which will be excised and inserted in the BamHI site of the Ti expression vectors. In order to construct pHD164, the BamHI-SacI fragment of pHD160 containing the 5' end of the coding sequence was replaced with the corresponding BamHI-SacI fragment of pBa3.3. To construct pHD159, the BamHI-SacI fragment of pHD163 was replaced by the BamHI-SacI fragment of pBa3.3 (Figure 16).

In order to create plasmid pDC3 (Figure 16), plasmid pHD164 was digested with DraI, ligated to BglII linkers, and the fragment containing the Bt2 gene was cloned in the BglII site of pLK57 (Figure 17). In this way, the BglII site of the BamHI-BglII cassette was placed in close proximity of the TAA stop codon of Bt2.

7.2 Construction of cassettes containing engineered Bt2 genes

7.2.1 Truncated Bt2 genes

7.2.1.1 Rational

Results from basic research on the functional properties of B.t. crystal proteins indicate that the

large approximately 130 Kd crystal proteins are relatively insoluble and, in addition, are protoxins which need processing in the insect midgut towards lower molecular weight active toxins, able to exert their toxic effects on the insects (Bulla, L. A., Jr., D. B. Bechtel, K. J. Kramer, Y. I. Shetna, A. I. Aronson and P. C. Fitz-James, 1980, Rev. Microbiol., 8:147-203; Bulla, L. A., Jr., K. J. Kramer, D. J. Cox, B. L. Jones, L. I. Davidson and G. L. Lookhart, 1981, Biol. Chem., 256:3000-3004; T. A. Angus, Can. J. Microbiol., 2:416 (1956); M. M. Lecadet, "Microbial Toxins", Vol. II, ed. by T. C. Montie and S. Kadis, Academic Press, Inc., New York and London, 1970, pp. 437-471). The specific activity of the Bt toxin when ingested by the insects as part of a composition of engineered plant material will be determined, not only by the total quantity of toxin present but also by the degree of accessibility of active toxin, released in the midgut. It has been shown that some insects species are more efficient than others in solubilizing and/or "processing" (enzymatically degrade) B.t. protoxins (Presentation by Dr. P. Luthy in "Second Workshop Bacterial Protein Toxins", Wepion, Belgium: June 30-July 4, 1985; to be published in congress proceedings). Therefore, it might be advantageous in the engineering of insect resistant plants to construct truncated toxins derived from Bt2 which have the properties of being: 1) already processed or partially processed toxin, exhibiting full toxic activity; and 2) more soluble than the original Bt2 protein. Plants expressing such truncated polypeptides might exhibit a higher specific toxicity against insects than plants expressing intact Bt2 at the same level.

7.2.1.2 Construction of the deletion mutants

1. Positioning of the toxin gene behind the P_L promotor

A gene coding for a 130 kD crystal protein toxin of B.t. berliner 1715 has been cloned into pUC8 (Viera and Messing, Gene 1, 259-268, 1982) giving rise to pBt200. Characteristics of this gene, called Bt2, and the resulting toxin (Bt2 protein) have been described in Sections 5 and 6.

In order to assure a regulatable, high-level expression in E. coli, the Bt2 gene was positioned behind the P_L promotor (Figure 17). To this end, the plasmid pBt200 carrying the Bt2 gene on a 7.7. Kb BamHI Pstl fragment was cut with Hpal, treated with Bal31, ligated to BamHI linkers, cut with BamHI and self-ligated (as described in Section 7.1). From the resulting clones, deletion derivatives with varying lengths of upstream sequences were selected, and inserted behind the P_L promotor of the expression plasmid pLK54 (see Figure 17 and Botterman et al., in press, Gene 1986) making use of the restriction enzymes BamHI and Pstl.

The resulting plasmids were assessed for the production of Bt2 protein and one of those producing the highest levels of Bt2, termed pLB10 was selected for further experiments. Plasmid pLB10 originated from pBa23-3 (Figure 17, Section 7.1).

2. Construction of deletions

From the internal deletions previously made in pBt200 with XbaI and KpnI, only the KpnI deletion gave

rise to immunologically detectable Bt2-derived protein (see Section 6). Deletions were made in pLB10 using restriction enzymes KpnI and HindIII. Western blotting analysis and ELISA showed that only the KpnI deletion mutant, containing the largest fragment extending from the start towards position 2167 of the Bt2 gene, produced a stable approximately 80 Kd polypeptide. The polypeptide encoded by the HindIII deletion derivative probably is highly sensitive to E. coli proteases.

Interestingly, the KpnI deletion mutant-encoded polypeptide exhibited an insecticidal activity that was equivalent to that of the intact Bt2 protein: in one experiment the LD_{50} value on 3rd instar P. brassicae larvae was determined to be 2.5 ng/larva for the Kpn deletion mutant as compared to 2 ng/larva for the intact Bt2. This result indicates that the truncated Bt2 gene product, arising from the KpnI deletion, comprises the entire active toxic unit.

The previous data suggests that the smallest gene fragment of Bt2, encoding an active toxin is contained within the KpnI deletion fragment but extends further than the HindIII site. To map the exact endpoint of the minimal fragment coding for the active toxin, deletion mutants were constructed which contained N-terminal fragments of decreasing size. To achieve this, we used a strategy which allowed us to construct simultaneously deletion-mutants and translational fusions to the NPTII-gene (see Section 7.2.2). The construction of the intermediate plasmid pLBKm25 is outlined in Figure 18. As shown in Figure 18, pLBKm25 is derived from pLB10 (see previous section) and pLKm91 which will be described in Section 7.2.2.2.

As shown, this plasmid is provided with a DNA sequence with stopcodons in the three reading frames behind an unique SalI site. This construct was cut with KpnI, digested with Bal31, cut with SalI, treated with Klenow polymerase and religated (Figure 19). In this way, the deleted coding region is fused to a stopcodon with a minimum of nonsense coding sequence. An overview of the deletion clones is given in Figure 20. Total cellular extracts were made of the clones (after induction) and analyzed in Western blotting and ELISA for the quantitative detection of Bt2-like polypeptides and in an insect toxicity assay to screen for active toxin. The results are presented in Figure 21 and indicate that detection of a stable polypeptide decreases gradually when the endpoint of the coding region is coming closer to the HindIII site.

From a certain position on (still downstream of HindIII), almost no Bt2-like protein was detectable anymore. Furthermore, toxicity of the extracted material from these clones, drops abruptly when the 3' endpoint is passing a particular position between HindIII and KpnI. The two clones characterizing the smallest toxic (pLB879) and the largest nontoxic (pLB834) polypeptide were verified by DNA sequence analysis. This analysis showed that the critical endpoint for a stable active toxin maps between positions 1797 and 1820 on the Bt2 gene (Figure 22). Therefore all N-terminal gene fragments of Bt2, ending downstream of position 1820 (bp) comprise a gene fragment encoding an active toxin. Interestingly, total cellular extract of one clone (pLB820) showed a much stronger reaction with a polyclonal antiserum and a monoclonal antibody in Western blotting. Moreover,

the protein produced by this clone was more soluble in E. coli than the KpnI deletion gene product and still exhibited full toxic activity.

7.2.2 Fusion genes to NPTII

7.2.2.1 Rationale

It is known that amino-terminal fusions at the NPTII gene can generate fusion proteins that still confer kanamycin resistance in bacteria (Reiss et al., EMBO J. 3, p. 3317, 1984).

Since NPTII is a most suitable selection marker in plant engineering, such gene fusions could have very promising applications. Indeed when using such NPTII fusion proteins to transform plants, a selection for high kanamycin resistance would allow direct selection for a high expression of the fusion product. Therefore, toxin gene fusions with NPTII might be used to transform plants and select for transformed plants expressing high levels of toxin, by selection for kanamycin resistance.

7.2.2.2 Construction of the fusion gene cassettes

Different fragments of the Bt2 gene were fused to the N-terminus of NPTII.

One of the fusion proteins termed Bt:NPT2 is described in more detail below.

1. Construction of the Bt:NPT2 fusion gene

The construction of the Bt:NPT2 gene is shown in

Figure 23. pLK54 is a pBR322 derivative containing the P_L promotor and 2 phage fd transcription terminators in tandem (Section 7.2.1.2). pKm109/90 contains the NPTII gene of Tn5 on pBR322 (Reiss et al., EMBO J., 1984) (Figure 24).

A 1141 bp gene fragment of pKm109/90 containing the NPTII gene was cloned in pLK54 giving rise to pLKm90. In order to create a BglII site behind the NPTII gene, BglII linkers were ligated at the XbaI and the SalI site after Klenow polymerase treatment. This gives rise to pLKm91.

pHD159 is a derivative of pBt200 (Section 7.1) whereby a BamHI linker has been fused to the 4th bp and a BglII linker to bp 3342 (after Bal31 treatment). The BamHI BglII fragment of this plasmid containing the deleted Bt2 gene was inserted in the BamHI site of pLKm91, in one orientation, giving rise to a Bt2:NPTII fusion gene on pLBKm10.

To construct pLBKm13 an Asp 728, Klenow treated BglII fragment was inserted between the BamHI site (after filling in) and the BglII site of pLKm91.

In order to produce the Bt:NPTII fusion proteins in E. coli, analogous constructs to pLBKm10 and 13 were made containing 5' leader sequences of the Bt2 gene with a ribosome binding site. Therefore, from another Bal31 deletion derivative of pBt200, pBa23-3 (Section 7.1), with the BamHI linker at position -33 we exchanged the BamHI-SacI fragment with pLBKm13, giving rise to pLBKm23.

For the expression of the fusion protein Bt:NPT2 behind the Pnos promotor and the 35S promotor, the BamHI-SacI fragment of pHD160 (described in Section 7.1) was cloned between the same sites in pLBKm13 giving pLBKm33.

Finally for the construction with the Petunia ssu-promotor (see Section 8) we used a modified Bt:NPTII cassette wherein the 3' non-coding region was removed up to the stopcodon of NPTII. To achieve this the NCo1-BglII fragment of pLBKm13 containing the 3' end of the NPTII gene was replaced by a NCo1 BglII- fragment generated from pLKm91 (Figure 23). This plasmid was cut with DdeI, treated with Klenow polymerase, and ligated to a BglII linker, whereafter the resulting DNA was cut by NCo1 and BglII. Figure 23 also shows the 5' Bt2 sequences in the different constructs.

In summary, the Bt:NPT2 gene contains (Figure 24):

1) The 5' end of the Bt2 gene starting 8 bp upstream of the initiation ATG codon or at pos +4 or at position -33 and extending towards nucleotide position 2173.

2) a 16 bp linker fragment.

3) the NPTII coding region starting at nucleotide position 13.

2. Characteristics of the fusion protein expressed in E. coli

The fusion gene Bt:NPT2, placed behind the P_L

promotor in plasmid construction pLBKm23 (Figure 23), was expressed in E. coli to study the properties of the fusion protein.

2.1 Identification of the fusion protein in E. coli

An E. coli clone transformed with pLBKm23 was analyzed in SDS-PAGE and in Western blotting. Coomassie staining of the complete bacterial extract in SDS PAGE showed the presence of a new protein band having an apparent molecular weight corresponding to the expected size of the fusion protein (approximately 110 Kd).

The protein could also be visualized in Western blotting using either anti-Bt2 serum or anti-NPTIII serum. Again the positively reacting band showed the expected size of the fusion protein. The protein is quite stable in E. coli since only limited amount of degradation products was detected (bands of lower M.W.).

2.2 Kanamycin resistance of the engineered E. coli clone and NPTII activity of the Bt:NPT2 protein

The E. coli clone containing pLBKm23 gene was indeed resistant to kanamycin. Bacteria were able to grow on 100 ug/ml Km (for comparison, the wild type NPTII gene confers resistance to more than 1000 ug/ml).

NPTII activity of the Bt-NPT2 fusion protein was evaluated using an NPTII assay as described elsewhere

(Reiss et al., Gene 30, p. 217, 1984). A cell extract of the E. coli clone expressing the Bt:NPT2 protein was run on gel in nondenaturing conditions, in parallel with an extract from an E. coli clone producing the wild type NPTII. Cell extracts were prepared as follows. The E. coli clones were grown during about 4 hrs. at 28°C in 20 ml cultures (containing LB medium), centrifuged and resuspensed in 1 ml TES buffer, sonicated for 2 times 2 minutes at 50 watts in Labsonic 1510 and centrifuged for 30 minutes at 15,000 rpm; the supernatant was used in our experiment. The position and NPTII-specific activity of the proteins was determined by in situ phosphorylation of kanamycin, using ^{32}P-ATP (Reiss et al., 1984, Gene, 30, 217-223). A gel containing the same samples was run in parallel and used in a Western blotting procedure (with anti-Bt2 and anti-NPTII antibodies). By comparison with dilution series of purified Bt2 and purified NPTII protein samples, which were included on this gel, the amount of protein present in the samples used for activity measurement could be determined; the results indicated that the specific activity of the Bt:NPT2 protein was roughly the same as that of NPTII (Figure 25).

2.3 Behavior of the Bt:NPT2 fusion protein in E. coli

The results in the previous section indicate that specific NPTII enzyme activities of intact NPTII enzyme and of the fusion protein were substantially equivalent. This seems somewhat incompatible with the "in vivo" behavior of the protein. While an E. coli producing wild type NPTII is resistant to more than 1000 ug/ml Km, the Bt:NPT2 engineered E. coli

expressing comparable levels of protein are only resistant to a level of 100 ug/ml.

It is known, however, that proteins produced in E. coli at high levels (such as is the case here, using the strong P_L promotor), tend to precipitate within the cell. One might expect that if a substantial fraction of the Bt:NPT2 protein is present in a precipitated form in E. coli (and most of the protein is enzymatically inactive) the result is a low resistant phenotype.

Analysis of a bacterial extract of the Bt:NPT2 clone using a Western blotting indicated that indeed nearly all the protein was present in an insoluble form. After lysis of the cells and centrifugation, most of detectable Bt:NPT2 was found in the pellet (this in contrast to the NPTII wild type protein which is mostly present in the supernatant fraction). Only a minor fraction of the Bt:NPT2 protein was present in the supernatant. To solubilize the fusion protein, present in the pellet fraction, 8 M urea + 2% ME (mercapto ethanol) was required. After dialysis, to renature the sclubilized protein an enriched Bt:NPT2 protein preparation was obtained which could be tested in insect toxicity assays (see 2.4).

2.4 Toxicity of the Bt:NPT2 protein

Previous toxicity assays have shown that the polypeptide encoded by the KpnI deletion fragment of Bt2 has still 100% of the toxic activity of the intact Bt2 protein (Section 7.2.1.2). In addition, our studies on the identification of minimal active toxic fragments have shown that this Kpn fragment comprises a

(approximately 60 Kd) active toxin which exhibits the complete toxic activity of the Bt2 molecule. In the following, we wanted to determine whether the Bt:NPT2 fusion protein had still the same degree of toxicity.

To this end, toxicity levels towards insect larvae, of enriched Bt:NPT2 and purified Bt2 protein were compared. The exact amount of fusion protein in the samples was determined using ELISA (with a monoclonal antibody specific for the N-terminal region of Bt2:4D6) and Western blotting (comparison of band intensity of dilutions of the fusion protein and purified Bt2). The results showed that the specific toxic activity on M. sexta and P. brassicae larvae was substantially the same for Bt:NPT2 fusion protein as for intact Bt2 (Tables 4 and 5).

Taken together the above data indicate that the Bt:NPT2 protein has NPTII activity both "in vivo" and "in vitro" and in addition that it has an equally potent insecticidal activity as the Bt2 toxin. Therefore, this truncated toxin clearly represents a valuable alternative in the engineering of plants expressing high level of insect toxicity.

2.5 Selection of additional Bt2:NPTII fusions expressing a higher kanamycin resistance phenotype

The results obtained with the previously constructed Bt:NPTII gene were very promising since this fusion protein conferred kanamycin resistance, showed normal levels of toxicity and was relatively stable. However, the kanamycin resistance conferred by

the Bt:NPT2 fusion was relatively low as compared to the wild type cells. Since the isolated purified Pnos-Bt:NPT2 protein had a specific NPTII activity comparable to the wild type NPTII protein, we concluded that the low kanamycin resistance was due to the low solubility of the fusion protein in E. coli. A relatively low resistance phenotype might interfere with an efficient selection system in plants. Therefore, we considered the possibility that other Bt-NPTII fusions could have different physicochemical properties leading to a higher Km resistance phenotype "in vivo". We designed an experiment to fuse the NPTII gene at random to the Bt2 sequence in the region between the KpnI site and the processing site. The experiment was as follows:

We used plasmid pLBKm25 (Figure 18) containing the following elements of interest: a P_L promotor and the Bt2 gene, fused at 1100 bp downstream of the KpnI site, to the NPTII gene (see Figure 19). A unique XhoI site separates the Bt from the NPTII gene. This plasmid was linearized by KpnI digestion and treated with Bal31 exonuclease. The Bal31 reaction was titrated such that the deletions did not proceed far beyond the HindIII site which is localized upstream of the C-terminal processing site. The Bal31 treated plasmid was ligated to XhoI linkers, digested with XhoI and self-ligated. As a result, the NPTII is fused to fragments of the Bt2 gene varying in size. These plasmids, transformed in E. coli, conferred kanamycin resistance on condition the NPTII gene was fused in frame to the Bt2 gene. Transformants were selected on plates containing low levels of kanamycin (20 ug/ml) and screened for the ability to grow on higher kanamycin concentrations.

145 kanamycin resistant transformants were screened for their ability to grow on higher kanamycin concentrations. 8 transformants proved more resistant and were able to grow on concentrations higher than 200 ug/ml of kanamycin. The fusion point in all 8 clones was determined by restriction enzyme mapping with an accuracy of 20 bp. Surprisingly 7 out of 8 clones had their fusion point around the HindIII site at position 1680 of the Bt gene. One clone (pLBKm860) mapped at position approximately 2050. Although the majority of the deletions were fused around position 1800, none of these conferred a higher kanamycin resistant phenotype. The 7 clones which have their fusion point positioned around the HindIII site are too short to encode an active toxin. However, one of the clones (pLBKm860) was:

More stable, since more protein per amount of total cellular extract was detected in Western blot analysis; and

More soluble since more truncated Bt protein was detected in the supernatant.

The positions of the 3' end points in the Bt2 coding sequences in clones 860 and 865 are represented in Figure 26.

Toxicity of the fusion proteins and truncated Bt2 gene products is illustrated in Table 6.

7.3 Adaption of cassettes containing truncated Bt genes or Bt gene fusions for expression in plant cells

Plasmids pLBKm860 and 865 were modified as described in Figure 27 to generated plasmids pLBKm1860 and pLBKm1865 respectively. pLBKm2860 was derived from (Figure 27) pLBKm860.

By replacing the BamHI-SacI fragment from pLB820 and 884 for the BamHI-SacI fragment of pLBKm14, the new plasmids called pLB1820 and 1884 respectively, were generated. pLB2820 was derived from pLB1820. As an example, the final constructs pLBKm1860, pLBKm1865 and pLBKm2860 are shown in Figure 24.

8. Construction of intermediate expression vectors containing the toxin gene

8.1 Overview

Table 7 gives an overview of the engineered plasmids which have been constructed and used in the plant tranformation experiments. Each engineered Ti plasmid is the result of a cointegration of a receptor Ti plasmid with an intermediate vector. Each intermediate vector contains a chimeric toxin gene comprising a plant promotor sequence derived from the indicated expression vector and a Bt gene cassette.

Assembly of chimeric genes

Table 7 gives the plasmids from which the Bt gene cassettes are isolated. The construction of these plasmids has been described in previous parts. The

detailed maps of the expression vectors mentioned in Table 7 are given in Figure 33 (A-J).

The detailed construction of the two chimeric genes present in pHD205 and pHD208 are described as an example. The experimental procedure uses standard recombinant DNA techniques and experimental protocols have been followed, according to Mariatis (Cold Spring Harbor, 1982).

The different plant promotors which have been used are (Table 7):

Pnos: a constitutive and relatively weak promotor derived from the T-DNA encoded nopaline synthase gene.

Pssu-pea: derived from a weakly expressed member of the family of small Subunit of Ribulose biphosphate-carboxylase genes of pea. Expression is light inducible in green tissues.

PTR2: A constitutive promotor derived from the T_R-DNA of the octopine Ti plasmid (Velten et al., 1984, EMBO J., 3, 2723). In the expression vectors pGSH150 and pGSH160 the dual promotor fragment is derived from pop443 (Velten et al., 1984, EMBO J., 3, 2723). The 5' untranslated region of the 2' promotor has been completed by adding a BamHI linker to produce the sequence

....ATCGATGGATCC
ClaI BamHI

Cloning into the BamHI site thus leaves the 5' untranslated region of the 2' gene intact and fuses the Bt gene cassette at the initiation ATG of the 2' gene.

The importance of an intact 5' untranslated region for obtaining high levels of expression of chimeric genes has been demonstrated (Jones et al., EMBO J, 4, 2411, 1985).

Pssu301: is the promotor of an rbs (ribulose bisphosphate carboxylase small subunit) gene of Petunia from which about 50% of the Petunia rbs mRNA is transcribed (Dean et al., EMBO J, 1984, ., No. 12). Via site directed mutagenesis the sequence around the initiation codon (TAACTATGGCTT) has been changed to TAA CCATGGCTT, creating an NCoI site on the initiation codon. Similarly, modification around the TAA stopcodon from GGCTTCTAAGTT to GGCTTCTAAGATCTT created a BglII site. BglII

As a result, the coding region of the ssu-301 gene could be precisely removed by NcoI-BglII digest and substituted by the Bt gene cassettes described under 7.2.

Isolation and characterization of the promotor fragment (Pssu PET) and 3' end (ssu-PET) is described in Dean et al., PNAS, USA, Vol. 82, 4964 (1985). Construction of pAGS007 (Figure 33H) is described in Figure 44.

P35S: is a strong constitutive promotor derived from cauliflower mosaic virus (CAMV). The P35S-1 promotor fragment is derived from CAMV isolate Cm4-184. It contains a fragment (nt6492-7454) of the CAMV genome (Gardner et al., 1981, Nucl. Acid Res., 9, 2871-2888) and a downstream BamHI site for cloning (Figure 40).

P35S-2 contains CAMV sequences from 6492 to 7466 followed by a ClaI site.

P35S-1 and P35S-2 thus only differ in the length of the 5' untranslated sequences of the 35S CAMV transcript.

Promotor-Bt gene junctions

Figure 28 gives the DNA sequences at the junction between the promotor regions and the coding sequence of the Bt gene cassettes as they are present in the different engineered Ti plasmids. Sequences derived from the original promotor regions and from the coding sequence of the Bt2 gene are underlined. Some relevant restriction enzyme sites which have been involved in the assembly of the chimeric genes are indicated. The ATG initiation codon is boxed.

Junctions between Bt gene cassettes and the 3' ends

All chimeric genes are provided at the 3' end with a sequence which contains the 3' untranslated region, including the polyadenylation site, of a gene which is naturally expressed in plant cells. The following sequences have been used:

3'ocs (in pHD1060, pHD1076, pHD1080)

A 706 bp PvuII fragment containing the 3' untranslated region of the octopine synthase gene (pos 11939-11233 according to Gielen et al., EMBO 4, p. 835, 1984).

3't7 (pGS1151, pGS1161, pGS1152, pGS1153, pGS1162, pGS1163, pGS1251, pGS1261, pGS1253, pGS1262, pGS1271, pGS1281

A 212 bp EcoRV-ClaI fragment containing the 3' untranslated region of T-DNA gene 7, cloned into the SmaI site of pUC8 and reisolated as a EcoRI-SalI fragment (pos 2317-2105 according to Gielen et al., EMBO 4, p. 835, 1984).

3'nos (in pGS1110)

A 182 bp Ta9I-ClaI fragment, containing the 3' untranslated region of the nopaline synthase gene (pos 1290-1472 according to Depicker et al., J.M.A.G. 1, p. 561, 1982)

3'SSu301 (in pGS1171, pGS1181)

A approximately 1.2 Kb BglII-BamHI fragment derived from the 3' end of the ssu301 gene was constructed by site-directed mutagenesis as follows:

	stop coding region (which is TAA)	
ssu301	...TTCTAAGTTATA	
coding sequence	...TTCTAAGATCTATA	
	BglII	Construction of a BglII site through site-directed mutagenesis

Example 8.1

This example describes the construction of pHD205, an intermediate vector containing a chimeric Bt2 toxin gene comprising: the nopaline synthase promotor, the Bt2 toxin gene cassette from pHD160 and a DNA fragment

containing the 3' untranslated region of the nopaline synthase gene including the polyadenylation site. In the chimeric gene the Bt2 gene cassette is oriented such that the expression of the Bt2 protein can be obtained from the nopaline synthase promotor. The Bt2 gene cassette was excised from pHD160 with BamHI and BglII and inserted in the BamHI site of pLGV2382 (Herrera-Estrella et al., EMBO J., 2, 987-995, 1983). Two ug of pHD160 DNA were totally digested with respectively 2 units of BglII and BamHI (Boehringer Mannheim) for 1 h at 37°C in a final volume of 20 ul, using the incubation buffer described by Maniatis et al. (Molecular Cloning (1982), Cold Spring Harbor Laboratory, 133-134). Five ug of pLGV2382 DNA was totally digested with BamHI under the same conditions. Subsequently the terminal 5' phosphates were removed from the DNA by treatment with calf intestinal alkaline phosphatase (CIP) (Boehringer Mannheim) using the conditions described by Maniatis et al., Molecular Cloning (1982), Cold Spring Harbor Laboratory, 133-134). One-fifth of BamHI digested and CIP treated pLGV2382 DNA was ligated to 0.1 ug of BamHI-BglII digested pHD160 DNA with 0.01 units of T4 DNA ligase (Boehringer Mannheim) in a final volume of 20 ul. Ligation buffer and incubation are as recommended by Boehringer Mannheim (Brochure "T4 ligase", Boehringer Mannheim, August 1980, 10.M.880.486). The ligation mixture was transformed into competent E. coli K514 cells (Colson et al., Genetics 52 (1965), 1043-1050) according to Dagert and Ehrlich, Gene, 6 (1980) 23-28. Cells are plated on LB medium (Miller, Experiments in Molecular Genetics (1972) Cold Spring Harbor Laboratory, New York) supplemented with carbenicillin (100 ug/ml). Transformants were screened for the presence of recombinant plasmids by microscale DNA preparations,

performed according to Birnboim and Doly (Nucl. Acids Res. 7 (1979), 1513-1523). The orientation of the BamHI-BglII fragment in the BamHI site of pLGV2382 was determined by BamHI-PstI double digestion. Double digestion pattern of recombinant plasmids shows 4 fragments after agarose gel electrophoresis. In the alpha-orientation there are fragments of approximately 5700 bp, 3000 bp, 2300 bp and 920 bp, whereas in the beta-orientation there are fragments of approximately 6200 bp, 3000 bp, 1800 bp and 920 bp. A recombinant plasmid with the alpha-orientation (the toxin gene under the control of the nopaline synthase promotor) is used in subsequent experiments and called pHD205.

Example 8.2

This example describes the construction of pHD208. The intermediate vector pHD208 contains a chimeric Bt2 toxin gene comprising: the promotor from a pea gene encoding a small subunit of ribulose biphosphate carboxylase (Pssu), the Bt2 toxin gene cassette from pHD160 and the 3' untranslated region of the octopine synthase gene including the polyadenylation site. The fragments of the chimeric gene were assembled in the cloning vector pGV831 as described in this example and as diagrammed in Figure 29. The construction of pGV831 is summarized in Figure 30.

Step 1: Insertion of a 706 bp PvuII fragment containing the 3' untranslated region of the octopine synthase gene into pGV831, to yield pGV858.

Five ug of pGV831 DNA was totally digested with 5u HpaI at 37°C for 1 h, using the incubation buffer

described by Maniatis et al., Molecular Cloning (1982). Subsequently, the terminal 5' phosphates are removed from the DNA by treatment with CIP using the conditions as described by Maniatis et al., Molecular Cloning, 1982. Twenty ug of pGV99 DNA (De Greve et al., J. Mol. Appl. Genet. 1, 499-512, 1982) was digested with 20 units of PvuII for 1 hour at 37°C.

The resulting DNA fragments were separated by electrophoresis on a horizontal agarose gel (0.8% agarose in TBE buffer). The agarose band containing the 706 bp PvuII fragment was cut out and the DNA was recovered by electroelution. After phenolisation and ether extraction, DNA was precipitated with ethanol by centrifugation in an Eppendorf centrifuge for 10 minutes, washed with 70% ethanol, dried and resuspended in 20 ul H_2O.

0.03 ug HpaI digested and CIP treated pGV831 was ligated to 0.1 ug of the purified 706 bp fragment with 1U of T4 ligase in a final volume of 10 ul. The ligation mixture was transformed into competent E. coli K514 cells. (Colson et al., Genetics 52 (1965), 1043-1050) as described by Dagert and Ehrlich, Gene 6 (1980), 23-28. Cells were plated on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York), supplemented with carbenicillin (100 ug/ml). The resulting recombinant plasmids were characterized by double digestion with PstI and ApaI digestion. One of the resulting plasmids, pGV858, yielded the desired digestion fragments of approximately 5300, 1700, 1500 and 900 bp and was used further.

Step 2: Construction of a modified Pssu fragment and insertion of this fragment in pGV858 to yield pHD503.

A BamHI site was positioned immediately downstream of the Pssu promotor by inserting the HindIII-BamHI polylinker from pUC8 (Vierra and Messing, Gene 19, p. 259-266, 1982) into pKC7::Pssu. pKC7::Pssu was provided by the Lab of Genetics, State University, Gent, Belgium. It contains the EcoRI-HindIII fragment which includes the promotor for one of the pea genes encoding the small subunit (ssu) of Ribulose biphosfate carboxylase (RUDP-case) (Herrera-Estrella, Nature 310; 115-120, 1984) cloned in vector pKC7 (Rao and Rogers, Gene 7, 1979). 1 ug of pKC7::Pssu was digested with 1 U HindIII and 1 U BamHI. 1 ug pUC8 DNA was digested with 1 U HindIII and 1 U BamHI at 37°C for 1 hr. 0.1 ug of each digested DNA were mixed and ligated with 0.01 unit of T4 DNA ligase in a final volume of 20 ul. The ligation mixture was transformed into competent E. coli K514 cells (Dagert and Erhlich, Gene 6 (1980) 23-18). Cells were plated on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with 100 ug/ml carbenicillin. In one of the resulting recombinant plasmids, pGV861, the HindIII-BamHI fragment containing the Km^R gene of pKC7 was substituted by the 20 bp HindIII-GamHI polylinker of pUC8.

Five ug of pGV858 were digested with 5 units of BamHI for 1 h at 37°C in a final volume of 20 ul, using the incubation buffer described by Maniatis et al., Molecular Cloning, Cold Spring Harbor Laboratory, 133-134, (1982). Subsequently, the terminal 5'

phosphates were removed from the DNA by treatment with CIP using the conditions described by Maniatis et al., Molecular Cloning, Cold Spring Harbor Laboratory, 133-134 (1982). Two ug of pGV861 were digested with 2 units of BglII, BamHI and PvuI for 1 h at 37°C in a final volume of 20 ul, using the incubation buffer described by Maniatis et al., Molecular Cloning, Cold Spring Harbor Laboratories 1982.

0.2 ug BamHI digested and CIP treated pGV858 was ligated to 0.05 ug BamHI-BglII-PvuI digested pGV861 with 0.01 units of T4 DNA ligase (Boehringer Mannheim) in a final volume of 20 ul. The ligation mixture was transformed into competent E. coli K514 cells (Colson et al., Genetics 52 (1965), 1043-1050) according to Dagert and Ehrlich, Gene, 6 (1980), 23-28. Cells are plated on LB medium (Miller, Experiments in Molecular Genetics (1972) Cold Spring Harbor Laboratory, New York) supplemented with carbenicillin (100 ug/ml). Carbenicillin resistant clones were screened for the presence of recombinant plasmids by restriction enzyme digestion of DNA prepared by the microscale technique described by Birnboim and Doly (Nucl. Acids. Res. 7 (1979), 1513-1523).

In one of the recombinant plasmids, pHD503, the BglII-BamHI fragment including the pea ssu promotor is inserted in the correct orientation in front of the 3' end of the octopine synthase gene. pHD503 contains a unique BamHI site, located between the Pssu promotor and the 3' end of the octopine synthase gene.

Step 3: Insertion of the BamHI-BglII Bt2 gene cassette into the BamHI site of pHD503 to yield the intermediate expression vector pHD208. Two ug of

pHD160 DNA were completely digested with 2 units of BglII and 2 units of BamHI for 1 hour at 37°C in a final volume of 20 ul. Five ug of pHD503 DNA were digested with 5 units of BamHI to completion under the same conditions, treated with CIP using the conditions described by Maniatis et al., Molecular Cloning (1982), Cold Spring Harbor Laboratory, 133-134) to remove the terminal 5' phosphates from the DNA. 0.1 ug of BamHI-BglII digested pHD160 DNA was ligated to 0.2 ug of BamHI digested and CIP treated pHD503 DNA with 0.01 U T4 DNA ligase in a final volume of 20 ul.

The ligation mixture was transformed into competent E. coli K514 cells (Dagert and Erhlich, Gene 6 (1980) 23-18). Cells were plated on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with streptomycin (20 ug/ml) and spectinomycin (50 mg/ml). Streptomycin-spectinomycin resistant clones were screened for the presence of recombinant plasmids by restriction enzyme digestion of DNA prepared from these clones by the microscale technique described by Birnboim and Doly (Nucl. Acids Res. 7, 1513-1523, 1979). pHD208, a recombinant plasmid containing the Bt2 gene cassette in the correct orientation with respect to the Pssu promotor was isolated and used in further experiments.

Example 8.3

This example describes the construction of pGSH151. The intermediate vector pGSH151 contains a chimeric Bt:NPTII fusion gene comprising: the promotor of transcript 2 of the TR-DNA of the octopine Ti plasmid (PTR2) (Velten et al., 1984, Embo J., 3, 2723),

the Bt:NPTII fusion gene cassette from pLBKm13 and the 3' untranslated region of the gene 7 of the T-DNA of the octopine Ti plasmid.

The fragments of the chimeric gene were assembled as described in this example. All the techniques were performed as described in Maniatis et al., Molecular Cloning (1982).

Step 1: Construction of pGSH50 (Figure 41)

This plasmid contains the TR promotor PTR2 with a completely intact 5' untranslated region, followed by an ATG-initiation codon, followed by a unique BamHI site, and the 3' untranslated end of the transcript 7 gene.

pOP443 (Velten et al., 1984) contains a ClaI-HdIII fragment comprising the PTR2 and the PTR1 of the octopine Ti plasmid. To eliminate the BamHI site, pOP443 was totally digested with BamHI and SalI, the sticky ends treated with the Klenow fragment of E. coli polymerase I and self-ligated with T4-ligase.

After transformation, ampicillin-resistant colonies were selected and their plasmids were screened for the absence of BamHI and SalI sites, yielding pOP4433SF.

In order to create a ClaI site in front of the 3' untranslated end of transcript 7 in pAP2034 (Velten et al., 1984), pAP2034 was totally digested with BamHI, treated with the Klenow fragment of E. coli polymerase I and ligated to kinated ClaI-linkers. The DNA was

subsequently totally digested with ClaI and self-ligated with T4-ligase; among the Amp^R transformants pAP2043C was selected.

From pOP443BSF, the ClaI-HindIII fragment containing the TR-promotors was cloned between the corresponding sites of pAP2034C giving rise to pGSH50.

Step 2: Construction of pGV1500 (Figure 42)

pGV825 is described in Deblaere et al., NAR, 13, 4777 (1985); to reduce its size, pGV825 was digested with PvuII and self-ligated. The resulting plasmid pGV956 contains a unique BamHI and a unique BglII-site within the T-DNA. pJB63 is described in Botterman et al. (in press, Gene, (1986)). The BamHI-BglII fragment containing several unique restriction sites was cloned between the corresponding sites in pGV956 giving rise to pGV1500.

Step 3: Construction of pGSH150 (Figure 43)

pGSH50 was digested with EcoRI, treated with the klenow fragment of E. coli polymerase I and digested with HindIII. The resulting fragment, containing the TR-promotors was cloned between the HpaI and the HindIII site of plasmid pGV1500.

Step 4: Construction of pGSH151 (Figure 3)

The BamHI-BglII fragment of pLBKm13 containing the Bt2 gene was cloned in the BamHI site of pGSH150 creating an in-frame fusion of the Bt2 gene starting at the 2nd codon to an ATG-initiation codon behind the PTR2.

9. Introduction of the intermediate expression vectors containing the toxin gene into Agrobacterium

The introduction of intermediate expression vectors into acceptor Ti plasmids of Agrobacterium is accomplished in two steps: first, the intermediate expression vector is transformed into E. coli strain GJ23 carrying two helper plasmids: R64 drd 11 containing tra functions and p GJ28 containing the mob functions (Finnegan et al., Mol. Gen. Genet. 185 (1982), 344-351). Secondly, the E. coli strain carrying all three plasmids is conjugated to an Agrobacterium strain containing an acceptor Ti plasmid carrying a region of homology with the intermediate expression vector essentially as described by Van Haute et al., (EMBO J. 2 411-418, 1983). The recombinant Ti plasmid, resulting from a single crossover event, is isolated by selecting for the antibiotic resistance marker carried by the intermediate expression vector.

As an example, the cointegration of pHD205 with pGV3850 and of pHD208 with pGV2260 is described. Intermediate vectors and receptor Ti plasmids used are listed in Table 7 and represented in Figures 31-33.

Example 9.1:

The intermediate expression vector pHD205 was inserted into the acceptor Ti plasmid pGV3850 to yield the hybrid Ti plasmid pHD1050. As diagrammed in Figure 31, pHD1050 contains the chimeric Bt2 gene under the control of the Pnos promotor, as well as the nopaline synthase gene positioned between T-DNA border fragments.

The plasmid pHD205 was introduced into competent E. coli GJ23 cells by transformation according to Dagert and Ehrlich (Gene 6 (1981, 23-28). To select for E. coli GJ23 cells transformed with pHD205, the cells were plated on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with carbenicillin (100 ug/ml).

Liquid LB medium was innoculated with one of the pHD205 transformed E. coli GJ23 colonies and cultured overnight (aobut 18 hours). 0.1 ml of this culture is conjugated with 0.1 ml of an overnight culture of the C58Cl Rif^R (also called GV3101, Van Larebeke et al., Nature 252, 169-170, 1974) containing (pGV3850) Zambryski et al (EMBO J. 2, 2143-2156, 1983) and cultured overnight at 28°C on solid LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York).

Agrobacterium strains containing hybrid Ti plasmids, resulting from a single cross-over event, were isolated by selecting for the kanamycin-neomycin marker carried by the pHD205 plasmid on minimal A medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with neomycin (400 ug/ml). After purification of transconjugants on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with rifampicin (100 ug/ml) and kanamycin (25 ug/ml). The physical structure of the T region of one of the transconjugants, pHD1050, was determined according to the method described by Dhaese et al., (Nucl. Acids Res. 7 (1979), 1837-1849) by hybridization of P^{32} labelled

pHD205 against HindIII digested to total DNA of C58C1 Ri f^R pHD1050. The structure of the T region of pHD1050 is diagrammed in Figure 31.

Example 9.2:

The intermediate expression vector pHD208 was inserted into the acceptor Ti plasmid pGV2260 to yield the hybrid Ti plasmid pHD1076. As diagrammed in Figure 32 pHD1076 contains the chimeric Bt2 gene under the control of the Pssu promotor as well as a chimeric gene containing the neomycin phosphotransferase gene under the control of the Pnos promotor, positioned between T-DNA border fragments. The Ti plasmid pGV2260 is described in European Patent Application Number 83112985.3 (Publication Number 0116718). The plasmid pHD208 was introduced into competent E. coli GJ23 cells by transformation according to Dagert and Ehrlich (Gene 6 (1980), 23-28). To select for E. coli GJ23 cells transformed with pHD208, the transformation mixture was plated on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold spring Harbor Laboratory, New York) supplemented with carbenicillin (100 ug/ml).

Liquid LB medium was inoculated by one of the transformed E. coli colonies and cultured overnight. 0.1 ml of the overnight culture of the E. coli strain carrying all 3 plasmids was conjugated overnight with an overnight culture of the C58C1 Rif^R (pGV2260) at 28°C on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York). Agrobacterium strains containing hybrid Ti plasmid, resulting from a single cross-over event between pGV2260 and pHD208 were isolated by selecting for the streptomycin-spectinomycin marker carried by

the pHD208 plasmid on minimal A medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with spectinomycin (300 ug/ml) and streptomycin (300 ug/ml) and streptomycin (1 ug/ml).

Transconjugants were purified on LB medium (Miller, Experiments in Molecular Genetics (1972), Cold Spring Harbor Laboratory, New York) supplemented with rifampicin (100 ug/ml), spectinomycin (100 ug/ml) and streptomycin (300 ug/ml). The physical structure of one of the transconjugants, pHD1076, was determined by hybridizing P^{32} labelled pHD208 against PstI digested total DNA of C58C1 Rif^R pHD1076 according to the method described by Dhaese et al., (Nucl. Acids Res. 7 (1979), 1837-1849). The physical structure of pHD1076 is shown in Figure 32.

Example 9.3

The intermediate expression vector pGSH151 was inserted intc the acceptor Ti plasmid pGV2260 to yield the hybrid Ti plasmid pGS1151.

The method used was a triparental cross according to Dittag et al. (1980), PNAS, 77, 7347-7351.

Liquid LB medium was inoculated with one of the pGSH151 transformed E. coli K514 colonies and cultured overnight at 37°C. 0.1 ml of this culture was plated together with 0.1 ml of overnight cultures of HB101 (pRK2013) Figurski & Helinski (1979), PNAS, 76, 1648-1652 and 0.1 ml of C58C1 Rif^R (Van Larebeke et al., Nature, 252, 169-170) on LB plates and grown overnight at 28°C.

The cells were collected from the LB plates and dilutions were plated on minimal A. medium (Miller, Experiments in Molecular Genetics, 1972, Cold Spring Harbor Laboratory, New York) supplemented with spectinomycin (300 ug/ml) and streptomycin (1 mg/ml). Transconjugants were purified on LB medium containing rifampicin (100 ug/ml), spectinomycin (100 ug/ml) and streptomycin (300 ug/ml). The physical structure of one of the transconjugants, pGS1151, was determined by hybridizing P^{32} labeled pGSH151 against PstI-BamHI digested total DNA of C58C1 Rif^R (pGS1151) according to Dhaese et al., N.A.R., 7 (1979) 1837-1849.

10. Isolation of plant cells and plants containing the chimeric toxin gene inserted in their genome

Procedures:

Two different protocols are described here for the transformation of tobacco plant cells with transformation vectors such as those described in Section 9 and for the generation of callus tissue and/or differentiated plants from these transformed cells.

Procedure 1: Cocultivation of protoplasts

This procedure describes the cocultivation of tobacco protoplasts with Agrobacterium C58C1 Rif^R and the isolation of transformed tobacco cell lines by screening for the presence of a scorable marker such as nopaline or for the expression of a selectable marker such as kanamycin resistance and the regeneration of whole plants from transformed callus lines.

Step 1: Preparation of Protoplasts

a) Grow 10-12 cm high Nocotiana tabacum cv. Petit Havana SR-1 aseptic plants for 4 weeks in vitro in medium containing half strength of the mineral components as well as half strength of the vitamins and sucrose of the Murashige and Skoog medium. (Murashige and Skoog, Physiol. Plant, 15, 473-497, (1962)).

b) Incubate leaf segments of 3 well developed young leaves with 20 ml of 1.4% cellulase Onozuka R-10 and 0.4% macerozyme Onozuka (both from Yakult Pharmaceutical Industry, Co., Ltd., Japan) in the following solution:

KCl 2.5 g/l
$MgSO_4.7H_2O$ 1 g/l
KH_2PO_4 0.136 g/l
Sorbitol 73 g/l
Polyvinyl pyrolidone - 10 0.3 g/l

c) Incubate overnight at 24°C in the dark;

d) Filter through a nylon filter with a mesh size of 50 micrometer;

e) Centrifuge in 15 ml tubes at 80 g for 10 minutes, remove the supernatant and resuspend the pellet in 20 ml of the same solution but without enzymes;

f) Centrifuge for 10 minutes at 80 g to remove excess of enzymes and remove the supernatant;

g) Resuspend pellet in 20 ml of 1/2 strength Murashige and Skoog medium supplemented with 0.22% $CaCl_2 \cdot 2H_2O$ and 0.4 M mannitol pH 5.6;

h) Centrifuge for 10 minutes at 80 g, remove supernatant;

i) Resuspend the pellets in 20 ml of medium 55 (see below);

j) Count protoplasts and dilute to a density of 10^5 pp/ml. Incubate in 5 cm petri dishes (2.5 ml per petri dish) in the dark about four days.

Step 2: Cocultivations with Agrobacterium strain C58Cl Rif^R containing the hybrid Ti plasmid (section 9).

a) A culture of Agrobacterium C58Cl Rif^R was grown until saturation in LB medium, centrifuged for 1 minute in an Eppendorf centrifuge, supernatant removed and the cells resuspended in an equal volume of 0.01 M $MgCl_2$. When about 30% of the protoplasts have started their first cell division, 50 ul of the bacterial suspension was added to 2.5 ml of the protoplast suspension (this represents about 100-500 bacteria per protoplast).

b) Incubate 48 hrs. in the dark.

c) Transfer the cell suspension to a centrifuge tube, wash the petri dish with the same volume of medium 55 supplemented with Claforan 500 mg/l, and add it to the centrifuge tube. Centrifuge for 10 minutes at 80 g, remove the supernatant and resuspend the pellet in the same volume of medium 55 supplemented with Claforan 500 mg/l.

d) Transfer to 5 cm petri dishes (2.5 ml/dish) at this moment the cell density is approximately 10^4 cells/ml. Incubate under 400 lux, 16 hours a day, at 23°C for 1-2 weeks until small aggregates of 4-8 cells are formed.

e) Add an equal volume of medium 56 (see below).

f) After 3-4 weeks colonies are plated on medium 56 solidified with 0.7% agarose, with reduced mannitol concentration (0.2 M instead of 0.44 M), and supplemented with Claforan 250 mg/l. At this stage the colonies must contain more than 50 cells/colony. In case Km^R is used as a selectable marker 50 ug/ml of Km is added to the medium as a selection agent.

g) Incubate 2-3 weeks at 800 lux, 16 hours a day, 23°C.

h) Transfer isolated calli to the same medium. Shoot induction occurs. At this stage, callus tissue is taken to screen for the presence of nopaline using the procedure as described by Aerts et al, Plant Sci. Lett. 17, 43-50 (1979), in case nopaline is used as scorable marker.

Step 3: Regeneration of transformed tobacco plants.

a) Grow nopaline positive or kanamycin resistant calli for 4 weeks.

b) Transfer the differentiating calli on hormone free Murashige and Skoog.

c) Grow for 3 weeks.

d) Separate shoots and transfer to the same medium, grow for 2-3 weeks till plants form roots.

e) At this stage small plants are transferred to grow in 250 ml containers containing 50 ml of half strength hormone free Murashige and Skoog medium.

f) Grow for 2-3 weeks. Remove a lower leaf for nopaline detection or screening of kanamycin resistance activity and for immunological detection of the toxin.

The leaf disc (also at times referred to herein as leaf segments) assay for testing Km resistance of a plant is performed as follows. Small discs are cut out from "in vitro" grown plants and transferred to petri dishes containing callus inducing medium (M&S macro and micronutrients and vitamins 3% sucrose, 500 mg/l Claforan, 1 mg/l NAA and 0.1 mg/l BAP) with various kanamycin sulphate concentrations (50-500 mg/l).

After three weeks incubation in a plant tissue culture room, callus growth on the leaf discs is monitored. The Km resistance level of the plant is determined as the highest concentration of Km on which the leaf discs still give rise to callus tissue.

Screening for the presence of nopaline (nopaline assay) is performed according to the procedures described in Aerts M., Jacobs M., Hernalsteens J-P., Van Montagu M. and Schell J. (1979) Plant Sci. Letters 17, 43-50.

Composition of medium 55:

- Half strength of the Macronutrients of the Murashige and Skoog salts

- 1 ml/l of 1000 x Micronutrients Heller modifed

- 1 ml/l of 1000 x vitamins Morel & Wetmore

- 100 ml/l Inositol

- 10 ml/l of a stock solution containing $FeSO_4$ 5.57 g/l and Na_2EDTA 7.45 g/l

- Benzylaminopurine 1 ml/l

Naphthalene acetic acid 3 mg/l

- Mannitol 80 g/l (0.44M)

Sucrose 20 g/l

1000 x Vitamins Morel and Wetmore for 100 ml	Micronutrients Heller modified (500 ml)
Ca pantotenate 100 mg;	500 mg $ZnSO_4 . 7H_2O$
Biotine 1 mg;	50 mg H_3BO_3;
Niacine 100 mg;	50 mg $MnSO_4 . 4H_2O$
Pyridoxine 100 mg;	50 mg $CuSO_4 . 5H_2O$
Thiamine 100 mg;	15 mg $AlCl_3$;
	15 mg $NiCl_2$

Composition of medium 56:

Medium 56 is the same as medium 55 except for the addition of naphthalene acetic acid at 0.2 mg/l and glutamine 1 mM.

Procedure 2: Infection of leaf segments with Agrobacterium strain C581 Rif^R containing a hybrid Ti plasmid

This procedure describes the infection of leaf segments with C58Cl Rif^R and the isolation of

transformed cell lines by selection on kanamycin containing medium.

Sterile Nicotiana tabacum cv. Petite Havana SR-1 plants were grown in vitro in plant nutrient agar containing half strength of the complete Murashige & Skoog (M&S) salt mixture complemented with half strength of the organic nutrients and sucrose of complete M&S medium. Twenty SR-1 leaf segments of approximately 1 cm^2 were floated on 5 ml liquid M&S medium (without hormones) in a 9 cm petri dish containing 0.1 ml of a washed bacterial suspension of C58C1 Rif^R. Incubation occurred on a shaker at 60 rmp in the dark for 48 h at 25°C. Subsequently, leaf segments were rinsed twice with M&S medium (without hormones) containing 500 mg/l Claforan, and then placed on a medium allowing both callus and shoot formation. This medium contains M&S macro- and micronutrients and vitamins, 3% sucrose, 500 mg/l Claforan, 500 mg/l kanamycinsulfate, 0.1 mg/l NAA and 1.0 mg/l BAP. The final pH of the medium is 5.8. Six leaf discs are placed per 9 cm petri dish containing about 30 ml medium and are incubated for 3 weeks at 23°C (approximately 1°C) under a 16 hours 2000 lux/day illumination cycle. After 3 weeks discs bearing callus and small shoots are transferred to the same medium for another 3 weeks. At that time shoots over 1 cm in length are transferred to M&S medium without hormones and without Km containing 500 mg/l Claforan. Afterwards, shoots are transferred about every three weeks on half strength M&S without hormones and the Claforan concentration is gradually decreased (1st transfer: 250 ug/ml, 2nd: 125 ug/ml, 3rd: 0 ug/ml Claforan). During the first transfer to 1/2 strength M&S, leaf material is removed to test kanamycin

resistance. Leaf discs are transferred to petri dishes containing callus inducing medium (M&S macro and micronutrients and vitamins, 3% sucrose, 500 mg/l Claforan, 1 mg/l NAA and 0.1 mg/l BAP) containing different kanamycin sulphate concentrations (50- 500 mg/l). Plants are retested for Km resistance on medium without Claforan when the material has been proved to be free of Agrobacteria.

Example 10.1: Callus and plants transformed with pHD1050.

T-DNA: Pnos-Bt2 (Bt2 gene fused to Pnos).

Marker: nopaline synthase as marker gene with additional border sequence between the Bt gene and the nos gene.

Transformation method: protoplast infection

Approximately 250 calli have been screened for nopaline and 19% were Nos^+, which represents a high efficiency of transformation.

In total 180 different callus lines, both nos^+ and nos^-, generated from these transformation experiments have been screened for the presence of Bt2 using the sensitive ELISA described above (Section 5.1). Most of the clones were tested early after transformation during the initial phase of propagation (when only 5 mm diameter) and some were retested after a period of subculturing (3 months later). On the basis of the immunoassay results, a number (25) of callus lines were selected for plant regeneration. From each callus several plants were regenerated, and each of them received a distinct number (total of 149 plants).

The 149 plants were propagated "in vitro" and subsequently 138 were transferred to the greenhouses.

All these plants appeared fully normal, flowered and set seeds. Some plants were tested for insect toxicity assays. From callus lines 161, 165 and 206, total DNA was prepared and the integration of the Bt2 gene was analyzed in Southern blotting. Integration of at least 1 copy of the Bt2 gene/genome was detected.

Example 10.2: Callus and plants transformed with pHD1060

T-DNA: Pnos-Bt2

Selectable marker: kanamycin resistance (Km)

Transformation method: protoplast infection (procedure 1) and leaf disc infection (procedure 2).

Following procedure 1, kanamycin resistant protoplast clones were obtained and grown as calli. Calli were selected at random and were put in generation medium for shoot formation. Shoots developed and isolated from these kanamycin resistant clones were propagated as plants "in vitro." Thereafter some of these plants were transferred to the greenhouse.

Following procedure 2, kanamycin resistant callus tissue and shoots were induced. Uncloned callus tissue was kept in continuous culture "in vitro." Kanamycin resistant shoots were isolated and were propagated "in vitro" as small plants (2-5 cm). These small plants were retested for kanamycin resistance using leaf disc assay (50 ug/ml Km). The shoots that were clearly resistant at this concentration of kanamycin were selected for further "in vitro" propagation. Plants were eventually transferred to the greenhouse. Using southern blotting analysis the presence of both the NPTII gene and the Bt2 gene was confirmed in the leaf tissue of these plants.

Example 10.3: Calli and plants transformed with pHD1076
T-DNA: Pssu-Bt2 (Bt2 gene fused to Pssu)
Selectable marker: kanamycin resistance
Transformation method: leaf disc infection.

Using conditions described in procedure 2 either callus transformation of shoot induction was performed on the infected leaf discs. Using the callus induction protocol, a number of calli were obtained by partial purification and maintained as separated semi clones. On the basis of positive immunoassay results 5 of these lines were selected for further propogation (1076-4, 10, 11, 12, 13). From the shoot induction protocol used in the initial stage of leaf disc infection a number (72) of kanamycin resistant plants were regenerated (selection on 50 ug/ml Km).

When retested by leaf disc assay 65% of these proved to be truly resistant to 50 ug/ml Km. From leaves of some "in vitro" propagated plants, callus tissue was generated and propagated "in vitro" for further testing.

Example 10.4: Calli and plants transformed with pHD1080
T-DNA: Pssu - Transit peptide (Tp) Bt2
Selectable marker: kanamycin resistance/(Nos)
Transformation method: leaf disc infection.

Kanamycin resistant calli and shoot were induced following procedure 2. Approximately 20 kanamycin resistant callus lines were analyzed for nopaline expression and all were found positive. 86 kanamycin resistant shoots were selected, propagated "in vitro" and retested for kanamycin resistance (using the leaf disc assay) and for nopaline expression.

52 plants (60%) were both kanamycin resistant and nopaline positive, and these were further propagated "in vitro." Approximately 10% of the plants expressed only one of the two markers.

Example 10.5: Plants transformed with pGS1110
T-DNA: Pnos-Bt:NPTII (fusion)
Selectable marker: kanamycin resistance/Nos
Tranformation method: leaf disc infection.

Leaf discs from "in vitro" maintained SR-1 plants were incubated during 48 hours with a suspension of Agrobacterium tumefaciens C58C1 Rif^R pGS1110 (procedure 2). Similar dilutions of different control strains containing chimeric genes encoding intact NPTII were included. After two weeks active shoot formation on M&S medium containing 50 mg/l kanamycin was observed both with the controls and pGS1110. However, after transfer to fresh selective M&S medium, a difference became apparent between the controls and pGS1110. Some shoots on discs inoculated with the latter strain turned yellow and were growing slowly. The best growing and green shoots were transferred to medium without kanamycin. Part of them could be rescued in this way and started growing normally after the second transfer on kanamycin free medium.

About 70 shoots were rescued from the pGS1110 transformation experiment. Screening among 35 of these shoots showed that 28 of these (85%) were real transformants since they produced nopaline. This important observation suggests that, although the shoots have not been maintained for a long period on Km containing medium, phenotypical selection for the expression of the fusion protein had occurred.

The obtained shoots were propagated "in vitro" as small plants on nonselective medium. A number of these plants were tested for Km^R resistance using the leaf disc assay. Most of them expressed a certain level of Km^R since they formed callus on Km containing medium. Variable resistance levels were recorded in the range of 50-500 mg Km/liter. However, most of the plants were only resistant to low levels of Km. Two out of a total of 61 plants showed resistance to 200 ug/ml Km and partial resistance to 500 ug/ml Km (very weak callus growth).

For a number of plants, copies were transferred into vermiculite pots. When reaching 10-15 cm height a first insect toxicity test was performed on leaves of these plants (see section 13).

Example 10.6: Plants transformed with pGS1161
T-DNA: PTR2-Bt2
Selectable marker: kanamycin resistance
Transformation method: leaf disc infection.

Leaf discs from "in vitro" maintained SR-1 plants were incubated during 48 h with a suspension of Agrobacterium tumefaciens C58C1 Rif^R pGS1161. As a control a A. tumefaciens C58C1 Rif^R pGS1160 containing NPTII under control of pTR was included. After two weeks shoot formation on medium containing 50 mg/l kanamycin sulphate was observed. After three weeks discs were transferred to fresh selective medium and after another three weeks the best growing shoots were transferred to kanamycin free medium. The level of Km^R is determined systematically using the leaf disc assay. Most plants showed high levels of resistance (callus formation on 500 ug/ml Km).

Example 10.7: Plants transformed with pGS1151
T-DNA: PTR2-Bt:NPT2 (fusion)
Selectable marker: kanamycin resistance
Transformation method: leaf disc infection.

Leaf discs from "in vitro" cultivated SR-1 plants were incubated during 48 hrs. with a suspension of Agrobacterium tumefaciens C58C1 Rif^R pGS1151. As a control A. tumefaciens C58C1 Rif^R pGS1160 containing NPTII under control of pTR was included.

Shoot formation and development of shoots on medium containing 50 mg/l kanamycin sulphate was slightly slower on discs treated with pGS1151 than in control discs (pGS1160). After three weeks discs were transferred to fresh selective medium and after another four weeks the best growing shoots were transferred to kanamycin free medium. The shoots were propagated "in vitro" as plants and the level of Km^R of these plants was determined systematically using the leaf disc assay. A number of plants were completely resistant to 500 ug/ml Km (normal callus growth). This data indicates that the PTR promotor directs higher levels of fusion protein expression in tobacco leaves than the Pnos promotor (pGS1110, Example 5 in this section).

Copies of the plants were transferred to pots and grown in the greenhouse. On a selected set of plants, those showing high Km resistance, detailed insect toxicity tests were performed (see Section 13). The level of Km^R is determined systematically using the leaf disc assay.

Example 10.8: Plants transformed with pGS1162 or pGS1163

T-DNA: PTR2-Bt2/820 - PTR2-Bt2/884

Selectable marker: kanamycin resistance

Transformation method: leaf disc infection.

Leaf discs obtained from "in vitro" grown SR-1 plants were infected with Agrobacterium tumefaciens C58Cl Rif^R pGS1162, pGS1163 or pGS1160 (as control). Discs were transferred to media containing different Km concentrations (50-100-200 mg/l). Shoots obtained on all three concentrations are transferred to Km free medium. Km resistance was checked by leaf disc test on callus inducing medium containing 50-500 ug/ml Km.

Example 10.9: Plants transformed with pGS1152

T-DNA: pTR2-Bt:NPT860

Selectable marker: kanamycin resistance

Transformation method: leaf disc infection.

Leaf discs obtained from "in vitro" grown SR-1 plants were infected with Agrobacterium tumefaciens C58Cl Rif^R pGS1152. Discs infected with Agrobacterium tumefaciens C58Cl Rif^R pGS460 were included as a control. Discs were transferred to media containing different Km concentrations (50-100-200 ml/l). Shoots were obtained on all three concentrations, although less abundant than in control discs infected with C58Cl Rif^R GS1160.

11. Immunological detection of Bt2 protein in engineered plant tissues

Expression of Bt2 in engineered plants (either callus tissue or differentiated plants) was monitored

using the ELISA described in Section 5 and adapted for assaying plant extracts.

Conditions for preparing and assaying plant extracts were established in reconstruction experiments in which purified Bt2 protein was mixed with plant extracts.

In reconstruction experiments we observed no significant loss in antigenic activity of Bt2 protein (less than 20%) due to the presence of plant extracts. In the ELISA assay, as little as 0.1 mg/ml purified Bt2 protein was still detectable. However, in reconstruction experiments a certain variability in background occurs, probably caused by plant proteins present in extracts. Therefore, reliable detection limit in these conditions was of the order of 1 ng/g tissue, which corresponds to a level of 2 ng Bt2 protein per g of plant tissue.

11.1 Screening of individual calli

For the immunological screening of individual calli, the following experimental procedure was established:

Two hundred mg of callus tissue was mixed with 150-200 ul of extraction buffer. Extraction buffer had the following composition: 50% of a solution of Na_2CO_3 500 mM and DIT 100 mM and 50% fetal calf serum. The tissue was homogenized by crunching with a spatula whereafter the cell debris were centrifuged. Fifty ul of supernatants was added to 50 ul of PBS pH 7.4 + 10% fetal calf serum in wells of a microtiter plate coated

with goat antibodies against B.t. crystal protein as described. During the entire procedure the samples were kept in ice and the microtiter plates were incubated at 4°C for 1.5 - 2 hours. Thereafter the ELISA procedure was continued as described in 5.1 for detection of Bt2 protein with either rabbit anti-Bt2 serum or with a mixture of monoclonal anti-Bt2 antibodies 4D6, 10E3, 1.7, and 4.8 (under the form of culture supernatants).

Example 11.1:
Analysis of calli transformed with C58C1 Rif^R pHD1050.

Transformed callus clones were obtained through the protoplast cocultivation method as described in Section 10 Example 10.1. Since 19% of the clones were found to express nopaline (Nos$^+$), at least 19% of them were transformed. However, due to an additional border sequence in the intermediate expression vector (pLGV2382) the nos gene and the Bt2 gene can be inserted independently as well as tandemly. Therefore both Nos$^+$ and Nos$^-$ clones were screened in the ELISA assay.

A total of 180 callus clones (130 nos$^-$, 50 nos$^+$) were tested. Some of the clones were retested once or twice at different time intervals after the initial propagation from protoplast culture. In none of the cases could a clear positive signal be recorded. When the substrate reaction times of the assay were prolonged (overnight incubation at 4°C) some of the clones (both nos$^+$ and nos$^-$) produced a very weak signal above the background (background being control callus without Bt2 gene). However, since the obtained values were clearly below the reliable detection limit

of the test system, no firm conclusions could be drawn concerning the expression of Bt2 protein in these calli.

Example 11.2:

Detection of Bt2 protein in tobacco callus tissue transformed with C58C1 Rif^R pHD1076.

Transformed callus tissue obtained from leaf segment infections using Agrobacterium strain C58C1 Rif^R (pHD1076) (see Section 10, Example 10.3), were screened immunologically for the presence of Bt2 protein.

After initial propagation calli were transferred for a second time after 20 days. When they reached optimal growth, 200 mg was used from each callus line for immunological screening in the ELISA. In a first experiment 9 out of 14 transformed calli showed a positive signal clearly above the background obtained with the 4 control calli (untransformed SR-1 callus), when reacted with a specific rabbit anti Bt2 serum. (see Figure 34). Three transformed calli generated a signal corresponding to approximately 5 ng Bt2 protein per gram tissue, as determined by comparison with a positive control (control SR-1 mixed with a known amount of Bt2 protein). All samples gave signals equal to background level signals (obtained with SR-1 control callus) when reacted with normal rabbit serum as a negative control. In a second experiment 13 out of 21 transformed calli yielded a signal significantly above background (Figure 35). One of the calli generated a signal corresponding to 4 ng of Bt2 per gram tissue. These results indicate that Bt2 protein is produced at

a detectable level in a fraction of the calli transformed with pHD1076.

About 5 weeks after the first ELISA experiments, 4 selected lines (1076-10, 11, 12 and 13) which in the initial screening gave high positive values, were retested in ELISA. From each line several "subclones" were tested (the original callus had been divided in pieces which were propagated independently in the next growth cycle; each new piece is referred herein to as a subclone). From 1076-10, one subclone was positive, one negative, from 1076-12, 2 subclones were positive, from 1076-13, 3 subclones were positive, 2 were negative. These results indicate that callus tissue originally scored as B.t. positive might, when further propagated, give rise to B.t. negative callus.

11.2 Detection of Bt2 in pooled callus extracts

In order to perform detailed immunoassay screenings with an increased sensitivity of detection, concentrated extracts from larger amounts of transformed callus tissues were prepared. The procedure developed here for obtaining an extract enriched in Bt2 protein, is based on the property of Bt2 to precipitate at pH 4-5.

Example 11.2.1: Calli transformed with pHD1076.

Transformed calli using pHD1076 were grown on medium containing 0.05 mg/ml kanamycin sulfate and 140 g of uncloned transformed calli were collected (a pool of callus lines 1076-4, 10, 11, 12, 13 and a number of unscreened lines). An extract was made by homogenizing the calli in the presence of 70 ml extraction buffer (Na_2CO_3 500 mM, pH 10, DIT 5 mM, PMFS 170 ug/ml).

The supernatant obtained after centrifugation at 15,000 rpm was diluted by adding 50 ml phosphate buffered saline pH 7.5. Subsequently the pH of the diluted extract was brought to pH 6 with 1 M HCl and it was incubated for 20 minutes at 0°C and the supernatant isolated by centrifugation and stored at 0°C (fraction pH 6). When pH was brought down to 4.5 a new precipitate was isolated (fraction pH 4.5) in the same way. The pellets were washed once with H_2O and subsequently incubated for 20 minutes at room temperature in the following buffer: Na_2CO_3 500 mM pH 10, DTT 50 mM, PMSF 170 ug/ml (pellet pH 6 in 1.5 ml and pellet pH 4.5 in 2 ml).

The material solubilizing in these conditions was isolated after centrifugation at 15,000 rpm and these samples were called 1076 pH 6 and 1076 pH 4.5 respectively.

A completely identical procedure was used to prepare extracts from normal SR-1 callus material (used here as a negative control) and resulted in two preparations called SR-1 pH 4.5. Total protein content in these samples was:

1076 pH 6	600 ug/ml
1076 pH 4.5	6560 ug/ml
SR-1 pH 6	380 ug/ml
SR-1 pH 4.5	3840 ug/ml

In order to evalaute the efficiency of the procedure a reconstruction experiment was done in which 1 ug of purified Bt2 was added to 20 g of SR-1 control callus tissue at the initiation of the sample homogenization. Presence of Bt2 protein in these extracts was determined using the ELISA (with goat anti-Bt crystal serum and rabbit anti Bt2, 6002). A strong reaction

was recorded in fraction 1076 pH 4.5 as compared to the negative control (SR-1). Fraction 1076 pH 6 gave a signal which was only slightly higher than SR-1 pH 6, indicating that this fraction only contained a minor part of the Bt2 protein content.

In the ELISA, fraction 1076 pH 4.5 also gave a significant reaction with five different monoclonal antibodies, specific for Bt2 protein, namely 1.7, 4D6, 4.8, 10E3 and 1F6 (see Figure 36). This strongly indicates that fraction 1076 pH 4.5 contains Bt2 protein which is in the same configuration as the bacterial Bt2.

In the following we attempted to remove Bt2 protein from the extract using a procedure of immunoprecipitation. A 5% volume of rabbit anti-Bt2 serum was added to the extract which was incubated at 4°C for 1 hour. Subsequently a 5% volume of goat anti-rabbit Ig serum was added, followed by 1.5 hours incubation at 4°C. The precipitate was removed by centrifugation and the supernatant was tested in the ELISA. This supernatant contained at least 10 times less Bt2 activity than the original 1076 pH 4.5 fraction, indicating that the material which generated the positive signals in ELISA could be specifically removed by anti-Bt2 antibodies, again confirming the Bt2 nature of the positively reacting substance in ELISA.

In a next experiment the above samples were dialysed against carbonate buffer pH 10. A quantitative determination of the Bt2 content of extract 1076 pH 4.5 was performed by testing dilutions of the extract and a solution of purified Bt2 protein

as a standard in the ELISA. The value determined was 122 ng Bt2/ml extract (total volume 2 ml). Reconstruction experiments (Bt2 added to SR-1 control callus at the beginning of the extraction), indicated that only 20% of Bt2 protein is lost during the extraction procedure and that 80% is contained in the pH 4.5 fraction. Based on these results, one could calculate that the total amount of Bt2 protein originally present in 140 g callus was 305 ng, which is 2.2 ng/g, a result that agrees well with the original estimates made for the screening of individual calli (see Figures 34 and 35). These data show that Bt2 protein present in extracts from transformed calli can be specifically concentrated using a precipitation procedure at pH 4.5 as described above, allowing us to quantify more accurately the amount of Bt2 protein produced in these plant tissues.

Example 11.2.2: Calli transformed with pHD1050.

A 500 g pool of selected callus clones (on the basis of previous ELISA tests on individual calli, approximately 25 callus lines, which gave values above background, were selected) and homogenized in the presence of 1000 ml extraction buffer using the same procedure as described in Example 1. Material which remained soluble at pH 6, but precipitated at pH 4.5 was isolated by centrifugation and subsequently redissolved in a small volume of carbonate buffer pH 10 (see Example 11.2.1). Analysis of the material in ELISA revealed positive signals corresponding to 60 ng/ml Bt2 or 1.2 ng/g callus tissue (Table 8).

Example 11.2.3: Calli transformed with pHD1060 and pHD1080.

In this example a slightly different and more extensive extraction protocol was used for the isolation of Bt2 protein from the engineered plant material. A protocol was developed to recover eventual residual Bt2 protein that would not be solubilized in a single extraction step as used in the procedures of Examples 11.2.1 and 11.2.2. Such could be the case, since Bt2 protein contains some highly hydrophobic regions which possibly interact with plant cell membrane structures and therefore would be difficult to solubilize in the absence of detergents. The step by step procedure used here would allow the recovery of additional proteins associated with insoluble plant cell structure. A schematic representative of the protocol is given in Figure 37.

A first protein fraction is obtained by extraction in carbonate buffer pH 10 + DTT and concentration through acid precipitation (pH 4.5) (fraction I). This fraction corresponds to the pH 4.5 extract obtained using the procedure in Examples 11.2.1 and 11.2.2.

Material not solubilized in this first extraction step and remaining in the pellet is then treated with the same extraction buffer containing 1% Triton X-100. Proteins, solubilizing in these conditions and precipitating at pH 4.5 are contained in fraction II. The last step involves solubilization in 2% SDS followed by acetone precipitation, yielding fraction III. Fractions I and II are analysed in ELISA and Western blotting; fraction III, which contains SDS, is only analysed in Western blotting.

ELISA results are given in Table 8: positive signals were detected in fractions I and II of both constructions 1060 and 1080, corresponding to Bt2 levels of respectively 1.9 and 1.4 ng/g original tissue (fr. I) and 0.27 and 0.29 ng/g (fr. II). Western blotting of the SDS solubilized material (fraction III) revealed the presence of a faint approximately 130 Kd band for both 1060 and 1080 callus material (using rabbit anti-Bt2 serum). Detection limit of the Western blotting was 10 ng/lane, therefore these fractions contained at least 0.39 ng/g for 1060 and 0.5 ng/g for 1080.

Western blotting of fractions I of 1050, 1060 and 1080 did not reveal the presence of a 130 Kd band probably because the concentration of Bt2 protein is too low in these fractions.

The present results indicate that low levels of Bt2 protein are indeed expressed in calli transformed with pHD1060 and pHD1080. Although small scale analysis of individual calli might not allow detection of immunopositive clones in these constructions, a more rigorous extraction and concentration procedure on a pool of selected calli clearly results in reliable and quantitative detection of Bt2 protein. A substantial fraction of the Bt2 was strongly bound to insoluble plant material and could only be released upon use of detergents such as Triton and SDS.

11.3 Detection of Bt2 protein in leaves of regenerated transformed plants

For the routine testing of leaf samples the following procedure was established:

Green leaf tissue (200-400 mg) was taken from "in vitro" grown plants (5-10 cm high as described in Section 10, Example 1) and homogenized in the presence of extraction buffer (200 ul), containing 50% of: Na_2CO_3 500 mM, 100mM DTT, 480 ug/ml leupeptine (Sigma, L-2884), 2 mM PMSF, 2mg/ml ascorbic acid (Sigma, A-7631), 2 mM EDTA and 50% of FCS. The tissue was homogenized by crunching with a spatula whereafter the cell debris were centrifuged. Thereafter the same ELISA procedure was followed as described for the screening of callus tissue (see Section 11.1).

Example 11.3.1: Detection of Bt2 protein in tobacco plants transformed with C58C1 Rif^R pHD 1050.

Leaves from the "in vitro" propagated plants (70 individual plants regenerated from 25 selected callus clones) were tested for Bt2 expression in the ELISA, using conditions described above.

From 70 plants tested, 5 gave a clear positive reaction above background obtained with leaves of untransformed SR-1 corresponding to Bt2 levels ranging from 6 to 25 ng per gram wet tissue (see Table 9).

One of the plants with the highest value (plant no 161-9: 25 ng) was studied in more detail. Leaf extract reacted positively with a rabbit anti-Bt2 serum and with a mixture of monoclonal antibodies specific for the Bt2 molecule (undiluted culture supernatants from clones 1.7, 4.8, 4D6, 10E3). Furthermore it did not react with a pool of monoclonals displaying an irrelevant specificity. The plant extract was retested at least twice after freezing at -20°C and thawing,

using the same reagents. Identical results were obtained each time, however, the level of Bt2 declines gradually with each cycle of freezing/thawing, probably as a result of Bt2 protein degradation.

Plant 161-9 was retested after it had been propagated in greenhouse conditions at a stage when it was about to flower. Again a clear positive signal was obtained, this time corresponding to a level of approximately 5 ng Bt2/gram tissue.

This result indicates that the levels of Bt2 protein detected in engineered plant leaves might vary considerably depending on the plants age, growth conditions, etc.

Example 11.3.2: Screening of tobacco plants transformed with C58C1 Rif^R pHD 1060.

Screening of 76 "in vitro" propagated kanamycin resistant plants did not yield clearly Bt2 positive plants using the ELISA method described in Example 11.3.1.

Example 11.3.3: Screening of tobacco plants transformed with C58C1 Rif^R pHD1076.

Leaf extracts from "in vitro" propagated plants obtained through the shoot induction method were screened in ELISA (method as in Example 11.3.1). Seventeen plants were tested and none of these gave a positive signal. Subsequently, an additional number (21) plants were screened when they were grown in pots and 15-20 cm high. Again no positive signal above background were recorded in ELISA upon screening of these plants.

Example 11.4: Screening of tobacco plants transformed with pHD1080.

About 30 "in vitro" propagated plants were screened in ELISA. One plant (plant nu 174) gave a positive signal corresponding to 20 ng/g Bt2 both with rabbit anti-Bt2 serum and a pool of monoclonal antibodies. The extract from this plant was consistently positive upon retesting in subsequent experiments.

11.4 Detection of Bt2 protein in callus tissue derived from leaves of transformed plants

The data outlined in Section 11.1, Example 11.1.2, 11.3, Example 11.3.3 suggests that the Pssu promotor constructions used here (pHD1076) are less active in differentiated plant leaves than in callus.

To investigate this further, new callus tissue was generated from leaves of the same plants used in the previous assays. Leaf discs were cultured on callus inducing medium and a few weeks later the callus material was collected and analyzed in a similar procedure (the only difference being that a Tris pH 7.5 buffer was used in the Step I extraction instead of a Na_2CO_3 pH 10/DTT buffer).

Callus, induced simultaneously from untransformed SR-1 leaves, was used as negative control.

ELISA analysis of the pHD1076 transformed calli revealed detectable amounts of Bt2 protein (Table 10).

These results show that a chimeric gene with the Pssu promotor from pea in the construction used herein is functional and induces expression of Bt2 protein in tobacco callus tissue derived from leaves, that did not express detectable amounts of the same protein. Thus a chimeric gene that directs expression in undifferentiated callus tissue may not necessarily be active to the same extent if at all in differentiated plant leaves.

12. Insecticidal activity of the Bt2 protein produced in engineered calli

Procedures:

Toxicity assays were performed on first instar larvae of Manduca sexta, fed on artificial diet. Three to four ml of liquid artificial diet (Bell, R.A. & Joachim, F.G. (1976) Ann. Entomol. Soc. Ann. 69: 365-373) were dispensed in each compartment (4 cm^2) of square Petri dishes. Formaldehyde was omitted from the diet. After the diet had become solid, 200 ul of a known dilution of sample was applied on the surface of the diet and dried in a cool air flow. Four newly hatched larvae were placed into each compartment. Growth and mortality were followed over a period of 3-5 days.

Example 12.1: Callus extract from calli transformed with pHD1076.

A concentrated extract from a pool of calli, transformed with pHD1076 was prepared as described in Section 11.2 Example 11.2.1. Extract dilutions were applied onto the surface of the diet and its toxicity

was evaluated. The extracted material 1076 pH 4.5 clearly had a toxic effect on the Manduca sexta larvae: at 12.5 ul/cm^2 all larvae showed growth inhibition and at 50 ul/cm^2 100% died (Table 11). Toxic activity of this material was significantly diminished after immunoprecipitation (100% normal growth at 12.5 and 25 ul/cm^2 and only 37% death at 50 ul/cm^2), indicating that the toxic activity can be depleted by anti-Bt2 antibodies. Extract from untransformed SR-1 callus tissue, the negative control, was completely nontoxic. Since the presence of Bt2 protein in the extracts was quantified immonologically, we could correlate the observed toxicity with the determined Bt2 conce ntration. The immunoassay values indicated that 1076 pH 4 contained 122 ng/ml Bt2. Thus, 50 ul extract per cm^2 corresponds to 6.1 ng Bt2 protein/cm^2. Previous toxicity assays with Bt2 on Manduca (Section 5.2, Table 3) indicated that Bt2 at 12 ng/cm^2 is 100% lethal while 2.5 ng/cm^2 induces growth inhibition.

Together these results indicate that the Bt2, expressed in engineered callus tissue, is a functional toxin and displays toxicity which is in the same range of potency as the bacterial Bt2 gene product.

13. Insecticidal activity exhibited by leaves of transformed tobacco plants

Procedures:

In order to evaluate the insecticidal activity expressed in leaves of transformed tobacco plants the growth rate and mortality of Manduca sexta larvae feeding on these leaves was recorded and compared with the growth rate of larvae feeding on untransformed SR-1 leaves. The following procedures were used:

Procedure 1

Subsequent experiments were carried out on leaf discs placed in Petri dishes. Four leaf discs of 4 cm diameter were punched out, placed on wet filter paper on a Petri dish together with 4 x 10 first instar larvae of M. sexta. Preferentially young leaves from the upper part of the plant were used. Twenty-four hours later a second disc was added. Between 48 hours and 100 hours after initiation of the experiment, the number of moulted insects were counted at regular time intervals. From this we could calculate the MT_{50} time at which 50% of the larvae had moulted. The whole experiment was conducted in a growth chamber at 26°C, 90% relative humidity and under a photoperiod of 16 hours light and 8 hours darkness.

In order to estimate the toxin levels required to have a notable effect on growth rate and viability of Manduca sexta larvae in the present experiment, a series of reconstruction experiments had to be included. To this end purified solubilized Bt2 protein (Section 5.1) was serially diluted in PBS containing 0.5% Triton X-100. Standard volumes of Bt2 solution were mechanically sprayed (to obtain a very homogenous coating) on tobacco leaf discs. Ten L1 (first instar) larvae were placed on each leaf disc, and 3 discs were used per Bt2 concentration. Growth rate and mortality of the larvae were followed over a 100 hour period.

Procedure 2

A procedure essentially similar to the previous one was also used. This experimental protocol was however somewhat more extensive in order to be more

effective in reliably detecting very small effects on larval growth rate. The set up was different from the previous one in the following aspects:

- care was taken that all plants were in exactly the same stage and condition so that the effects on larval growth caused by differences in the condition of the leaf tissue would be minimal.

- larval growth was followed up to the L_3 stage (unlike previous experiments where growth was only monitored up to L_2).

- not only the moulting time of the larvae was recorded but also larval weight in the final stage was measured.

The plants used in this set up were grown in the greenhouse until they reached a height of 60-80 cm, but were not flowering yet. Leaf discs were cut out, placed on wet filter paper in Petri dishes and 10 first instar larvae were placed on each disc. Per plant, five groups of 10 larvae were used (5 leaf discs).

Growth rate and mortality were followed over a 7 day period (at this time nearly 100% of the controls were in the L_3 stage).

Example 13.1

Plants transformed with pHD1050, 1060, 1076 and 1080 were screened in the insect assay following procedure 1. No significant effect on growth rate

and viability of the larvae could be recorded using this procedure. Results of a reconstruction experiment with purified bacterial Bt2 protein were as follows:

Growth inhibition but no mortalilty was observed at 25 ng/g and approximately 50% mortality at 50 ng/g.

Example 13.2

An extensive toxicity test using procedure 2 was done on a number of transformed plants that were previously scored as Bt+ in immunoassays. These plants were

161-9 (Pnos-Bt2, nos+) (pHD1050 Example 1 Section 10)

147 (Pnos-Bt2, nos+) (pHD1050 Example 1 Section 10)

174 (Pssu-Tp-Bt2, Km+, nos+) pHD1080 Example 4 Section 10)

As controls a Bt^- plant (161-6) and an untransformed SR-1 were used. Results are presented in Table 12.

A) The number of larvae that were still in the L_2 stage, or already went to L_3, or died, at 150 hours after initiation of the test. Clearly the L_2-L_3 transition is somewhat earlier in the groups of larvae feeding on SR-1 and 161-6 as compared to those feeding on the Bt^+ plants 161-9, 147 and 174. In none of the groups has significant mortality been recorded (10% or less is considered as background).

B) Mean larval weight at the end of the experiment is presented in the upper row (larvae in 5 groups of 10, deviation is calculated on the mean values of these groups).

Below are the weight values calculated for the 5 largest larvae from each group of 10. These results are compatible with the kinetics of the L3-L4 transition: control larvae are somewhat larger than larvae feeding on Bt^+ plants.

Example 13.3

Insect toxicity assays were done on leaves of plants obtained from the transformation procedure with pGS1110 (Section 10, Example 10.5). Plants were 10-20 cm and tests were performed following procedure 1. L1-L2 transition was monitored and 2 groups of 10 larvae were used per plant. A significant growth inhibition effect, exhibited by some of the plants on M. sexta larvae, was observed. Data on the L1-L2 ratio after about 3 days of feeding, are presented in Table 13. Control plants included in these experiments were transformed with vectors containing Pnos-NPTII only. In Exp. 1, from the 8 plants putatively transformed with pGS1110, 3 produced growth inhibition (N20-38, N20-22, N20-18) as compared to the 3 control plants (C1, C2, C3). In Exp. 2, one plant (N20-37) out of 6 produced growth inhibition when compared to the 4 control plants (C4, C5, C6, C7). The differences in growth rate are apparent when complete growth rate curves are compared (see Figures 38 and 39).

The same plants were also screened for the presence of nopaline and for resistance against

kanamycin, in order to determine whether they were real transformants. The results of the screening data on the plants used in the present insect tests are compiled in Table 14. All four plants that showed an effect on larval growth are among the positive transformants, since they are Kanamycin resistant (Km^R) and nopaline positive (nos+).

Example 13.4

Insect toxicity assays were performed on leaves of plants generated through transformation with pGS1151 (Section 10, Example 10.7). Plants were 15-30 cm high at the time of testing and had been grown in greenhouse conditions.

Two independent experiments are described below: some of the plants tested in the first experiment were retested in Experiment II, in order to confirm the observed toxicity effects.

Experiment I:

The test was performed as described in Procedure 2 (this section) except that only two groups of ten larvae were used per plant (newly hatched Manduca sexta larvae).

Growth rate and mortality of the larvae were followed over a 7 day period and the larval weight at the end of this period was determined. Detailed results from Experiment I are represented in Table 15 and indicate that larvae feeding on several plants transformed with pGS1151 show significant growth inhibition in the initial stage of the experiment, as

compared to larvae feeding on a control plant. For example, after 71 h, 60% of the larvae feeding on control plant N21-107 have gone to the L2 stage, while the number of L2 larvae is only 15% or less on plants N21-18, 43, 53, 50 and 11. When followed over a longer period, significant mortality was recorded in the larvae feeding on pGS1151 transformed plants. On one of the plants (N21-11), mortality reached 100% after less than 7 days. Mortality on the control plant only reached 15% on day 7 and 45% of the larvae had already gone to the L3 stage (this in contrast to the other plants having substantially no L3 larvae on day 7).

Experiment II:

Results from a second insect test (II) involving newly hatched M. sexta larvae was performed on some of the plants also used in Exp. I, following Procedure 1. Results are presented in Table 16. A high mortality rate was recorded in the plants transformed with pGS1151 (75-100% death) while nearly all the larvae feeding on the control plants N21-102, 104 and 107 were still viable after 4 days. A complete list of all the plants used in insect tests I and II is given in Table 17. Also indicated are the Km resistance levels determined for the plants transformed with pGS1151; the percentage mortality of the larvae feeding on these plants after several days; and the mean weight of the larvae that survived after 7 days in Experiment I.

Conclusion:

Tobacco plants transformed with pGS1151 and selected for high Km resistance clearly induce severe toxic effects on larvae feeding on these plants. The effects on insect larvae observed here, are the same as

those induced by the B.t. toxin of bacterial origin (see Section 5.2, Tables 2 and 3); that is, growth inhibition in the initial stage (retardation in the transition from one instar to the next) followed by death.

It is apparent from Table 17 that the plants exhibiting the highest levels of Km resistance (500 ug/ml Km) also induce the highest mortality rates. Thus, using the fusion protein construction, we were able to select for efficient expression of toxicity by selecting for Km resistance.

It should be noted that the use of a fusion protein, as described herein, may represent a particular advantage, not only because direct slection for transformants of interest can be done, but also because the fusion protein itself might have some intrinsic useful properties. For example, Bt2:NPTII fusion proteins might be more stable in plant cells than intact Bt2 protein and/or the messenger RNA derived from the fusion genes might be more stable than intact Bt2 RNA.

14. Stable inheritance of new phenotype, acquired through transformation

A substantial fraction of the plants transformed with the transformation vectors described herein will contain, stably inserted into their genome, a fragment of newly acquired DNA containing both a chimeric Bt toxin gene and a marker gene (nos, NPTII). This was confirmed by the results of southern blotting

experiments. The new phenotypic traits acquired through this transformation method (expression of Bt Toxin, antibiotic resistance, nopaline production) will be inherited according to classic Mendelian genetics. To verify stable inheritance of the new traits, F_1 descendants from transformed plants were analysed for the expression of Bt toxin and synthesis of nopaline.

Transformed tobacco plants were allowed to flower and give seed. Care was taken that no cross pollination occured. From 4 plants previously identified as Bt^+ (161-9, 10^{-1}, 147^{-8}, 174), seeds were germinated in agar medium and F_1 plants were analysed for the presence of nopaline (nopaline synthase being present as marker gene in the parental plants). Plants were tested 3 weeks after germination (approximately 1 cm in height) or later at 6-7 weeks (2-4 cm). The results are depicted in Table 18.

From plants 10-1 and 147-8 about 3/4 of the F_1 were nos^+, which is expected from Mendelian inheritance of a single locus (1:2:1). For F_1 plants from 161-9, the nopaline signal was very weak when plantlets were tested at approximately 3 weeks after germination. Due to this weak expression the nopaline signals were not clearly visible and therefore the number of positives might be underestimated at this stage. However at 7 weeks a clear positive signal was detected in a 3/4 of the plants. The reason for the low expression in the early age of the plants is not known.

In the F_1 from plant 174, of the 45 plants analysed, 43 were nos^+. This high percentage (95%) of no s^+ indicates that the nos gene is inserted in the

genome on more than one independent locus. F_1 plants were also analysed for the expression of Bt2 toxin using the ELISA. Data from ELISA assays on leaf tissue indicated that $Bt2^+$ phenotype was correlated with nos^+. Therefore the $Bt2^+$ trait is stably inherited.

Cultures of cells containing intermediate cloning vectors and hybrid plasmid vectors were deposited on 10th December 1984 with Deutsche Sammlung von Miko-organism (DSM) Gesellschaft fur Biotechnologische Forschung mbH, Grisbachstr 8D-3400, Gottingen, Federal Republic of Germany (having been despatched thereto on 5th December 1984) and have been assigned accession numbers as follows:

E. coli K514 (pHD208)	DSM 3127
E. coli K514 (pHD205)	DSM 3128
A. tumefaciens C58C1 Rif^R (pHD1076)	DSM 3129
A. tumefaciens C58C1 Rif^R (pHD1050)	DSM 3130

Cultures of B.t. berliner 1715 have also been deposited with the same depository and been assigned an accession number of DSM 3131. Nicotiana tabacum cv. Petit Havana SR-1 has been deposited with the United States Department of Agriculture, National Seed Storage Laboratory, Colorado State University, Ft. Collins, Colorado 80523 and assigned serial number 191197 and is freely available upon request.

Cultures of cells containing intermediate cloning vectors and hybrid plasmid vectors were deposited with American Type Culture Collection (ATCC) on 24th December 1985 and have been assigned accession numbers as follows:

E. coli K514 (pLBKm25)	ATCC 53390
E. coli K514 (pLBKm33)	(without lambda repressor) ATCC 53389

AUSTRALIAN PATENT OFFICE LS

E. coli K514 (pLBKm1820)	ATCC 53388
E. coli JM83 K12 (pSSU301)	ATCC 53391
E. coli K514 (pLBKm1860)	ATCC 53387
A. tumefaciens C58C1 Ery^R Cml^R (pHD1080)	ATCC 53385
A. tumefaciens C58C1 Rif^R (pGS1110)	ATCC 53386
A. tumefaciens C58C1 Rif^R (pGS1151)	ATCC 53392
A. tumefaciens C58C1 Rif^R (pGS1161)	ATCC 53393
A. tumefaciens C58C1 Rif^R (pGS1152)	ATCC 53394
A. tumefaciens C58C1 Rif^R (pGS1163)	ATCC 53395
A. tumefaciens C58C1 Rif^R (pGS1171)	ATCC 53396
A. tumefaciens C58C1 Rif^R (pGS1181)	ATCC 53397
A. tumefaciens C58C1 Rif^R (pGS1182)	ATCC 53398
A. tumefaciens C58C1 Rif^R (pGS1251)	ATCC 53399
A. tumefaciens C58C1 Rif^R (pGS1261)	ATCC 53400
A. tumefaciens C58C1 Rif^R (pGS1253)	ATCC 53401
A. tumefaciens C58C1 Rif^R (pGS1262)	ATCC 53402

Their viability was tested and confirmed on 27th December 1985.

Cultures of *E. coli* K514 are commercially available

TABLE 1

Toxicity (Toward P. brassicae Larvae)
of Bt2 and B.t. Crystal Proteins

Sample	Toxicity (mean value ± S.D.*) LD_{50} (ng/larva)
Solubilized B.t. berliner 1715 crystals	0.65 ± 0.35
Purified Bt2 protein	1.65 ± 1.3

* S.D. is Standard Deviation.

TABLE 2

Effect of Bt2 Protein on Growth Kinetics of P. brassicae Larvae
(Results Expressed in % of Larvae in a Certain Stage); 1 ppm = 267 ng/gram leaf

Bt2 Concentration

Time (hours)	0						0.01 ppm						0.1 ppm					
	Stage L3	WC	L4	WC	L5	% mort	L3	WC	L4	WC	L5	% mort	L3	WC	L4	WC	L5	% mort
24	100					0	100					0	100					0
48	33	67				0	93	7				0	100					-
52	16	50	34			0	70	30				0	100					-
57		30	70			0	55	45				0	100					-
71		3	97			0	44	15	41			0	100					60
77			100			0	15	18	67			0	100					-
95			100			0	3	5	92			0	100					85
102			89	11		0	3	5	92			0	100					85
119			63	30	7	0		3	97			0	100					95
127			36	40	24	0			97	3		0						100
143			7	58	35	0			45	51	4	0						
151			6	22	72	0			24	70	6	0						
167					100	0			15	27	58	0						

TABLE 3

Toxicity of Bt2 and Total B.t. berliner Crystal Proteins Towards Larvae of Manduca sexta, Expressed as Percentage Mortality

Time (days)	Control *E. coli* Extracts	Bt2					B.t. berliner Crystals				
	dose (ng protein/cm)										
	1250	2.5	12.5	25	125	250	2.5	12.5	25	125	250
1	0	0	0	0	0	0	0	0	0	0	0
2	0	0	4	8	28	36	0	0	0	20	20
3	0	0	64	92	100	100	0	32	64	92	100
4	0	4	80	100			0	72	92	100	
5	0	4	88				0	81	100		
6	0	8	100				0	88			
7	0	8					0	88			

TABLE 4

Toxicity of Bt:NPT2 Fusion Protein on 3rd Instar P. brassicae (% Mortality After 4 Days)

	Toxin dose (ug/ml)				
Bt protein	0.1	0.2	0.3	0.6	1
Bt2	70	NT(x)	90	NT	100
Bt:NPT2	NT	80	NT	100	NT

(x) NT = Not Tested

TABLE 5

Toxicity of Intact Bt2 Protein, 60 Kd "Processed" Bt2 Protein (Trypsin Digested) and Bt:NPT2 Fusion Protein on Larvae of Manduca sexta

% Mortality after 4 days

Toxin dose: (ng/cm^2)	0	0.67	2	6	18	54	162
130 Kd Bt2	0	0	0	0	3	8	100
60 Kd Processed Bt2	-	0	0	0	0	60	100
Bt:NPT2	-	0	0	0	0	83	100

Larval Weight after 4 days (mg/larva)

Toxic dose (ng/cm^2)	0	0.67	2	6	18
130 Kd Bt2	27.4	20.7	9.4	5.4	2.4
60 Kd Bt2	-	16.3	8.3	6.4	3.9
Bt:NPT2	-	26.5	15.8	7.7	4.5

Toxin dilutions were applied on artificial diet as described in Section 12. Thirty (30) 1st instar larvae were used per dilution.

TABLE 6

Toxicity of Bt:NPTII Fusion Proteins or Bt2 Deletions on 3rd Instar P. brassicae Larvae (% Mortality After 4 Days)

E. coli strain	Exp. 1	Dilution 1/100	Bacterial Extract 1/10	1/3
NF_1 (neg. control)		0	0	0
pLBKm860		100	98	100
pLBKm865		2	0	0
	Exp. 2	1/25	1/5	1/1
NF_1		14	2	2
pLB879		100	100	100
pLB834		2	2	0
	Exp. 3	1/100	1/10	1/1
NF_1		4	4	2
pLB879		8	50	98
pLB820		54	100	100
pLB884		74	100	100

TABLE 7

Summary of Engineered Ti Plasmids and Their Intermediate Vectors

Ti Plasmid	Ti Plasmid	Interme-diate	Expr. Vector	Bt Casette	Plant Prom.	Plant Marker	3' End
pHD1050	pVG3850	pHD205	pLGV2382	pHD160	Pnos	nos	-
pHD1060	pGV2260	pHD207	pGV857	pHD162	Pnos	Km	ocs
pHD1076	pGV2260	pHD208	pHD503	pHD160	Pssu pea	Km	ocs
pHD1080	pGV3850/ Km	pHD210	pAC6	pHD164	Pssu pea	Km	ocs
pGS1110	pGV3850	pGSH10	pGV874	pLBKm33	Pnos	KmF*	Nos
pGS1151	pGV2260	pGSH151	pGSH150	pLBKm33	PTR2	KmF	t7
pGS1161	pGV2260	pGSH161	pGSH160	pHD164	PTR2	Km	t7
pGS1152	pGV2260	pGSH152	pGSH150	pLBKm1860	PTR2	KmF	t7
pGS1162	pGV2260	pGSH162	pGSH160	pLB1820	PTR2	Km	t7
pGS1163	pGV2260	pGSH163	pGSH160	pLB1884	PTR2	Km	t7
pGS1171	pGV2260	pGSH171	pAGS007	pLBKm14	Pssu301	Hyg	ssu301
pGS1181	pGV2260	pGSH181	pAGS007	pDC3	Pssu301	Km	ssu301
pGS1182	pGV2260	pGSH182	pAGS007	pLB1820	Pssu301	Km	ssu301
pGS1251	pGV2260	pGSJ251	pGSJ250	pLBKm33	P35S*1	KmF	t7
pGS1261	pGV2260	pGSJ261	pGSJ260	pHD162	P35S*1	Km	t7
pGS1253	pGV2260	pGSJ253	pGSJ250	pLBKm2860	P35S*1	KmF	t7
pGS1262	pGV2260	pGSJ262	pGSJ260	pLB2820	P35S*1	Km	t7
pGS1271	pGV2260	pGSJ271	pGSJ270	pHD162	P35S*2	Km	t7
pGS1281	pGV2260	pGSJ281	pGSJ280	pLBKm33	P35S*2	KmF	t7

* KmF indicates Kanamycin fusions.

TABLE 8

Results Immunoassays on Pooled Callus Extracts

Construction	Extract Fraction	Protein Content ug/ml	Total Volume Extract (ml)	Bt2 in ELISA ng/ml	ng/g	Western Blotting Volume (ul)	130 Kd
pHD1050 (500 g)	I	9650	10	60	1.2	50	-
pHD1060 (392 g)	I	7800	8	95	1.9	50	-
	II	640	1	105	0.27	200	±
	III	N.D. (x)	0.3	N.D.	N.D.	20	+
pHD1080 (100 g)	I	4150	2	72	1.2	50	-
	II	326	1	29	0.29	N.D.	N.D.
	III	N.D.	0.5	N.D.	N.D.	100	+

(x) N.D. = Not Determined

TABLE 9

Levels of Bt2 Protein Detected in Leaves from 5 Immunopositive Plants Transformed by pHD1050

Plant Isolation Number	ng Bt2/g Plant Tissue
161-9	25.0
10-1	7.6
10-2	6.0
147-8	14.0
147-9	9.2

TABLE 10

Immunoassays on Extracts of Calli Derived from Leaves of Transformed Tobacco

Construction	Fraction	Protein Content (ug/ml)	Volume Extract (ml)	Bt2 Detected in ELISA (ng/g)
pHD1076 (59 g)	I	6200	7	1.6
	II	1520	1.5	0.4

TABLE 11

Toxicity of Callus Extract on Manduca Sexta Larvae

Extract	Volume Per cm^2 (ul)	Total Number Larvae	Results After L1	WC	L2	Dead
1076	12.5	4	3	1		
pH 4.5	50	4				4
	100	4				4
SR-1	50	8			8	
pH 4.5						
(Control No		44		1	43	
Plant Extract)						
After Immunoprec:						
1076	25	12			12	
pH 4.5	50	8		1	3	4
SR-1	50	8			8	
pH 4.5						

TABLE 12

Growth Rate and Mortality of Manduca Sexta Larvae Feeding on Transformed Tobacco Leaves

A. Larval Stage at 150 h: (Number of Larvae)

Plant	161-9	147	174	SR-1	161-6
L2	22	22	24	9	5
L3	25	27	23	36	41
Dead	3	1	3	5	4

B. Larval Weight at 164 h:

Mean Weight Per Larva (mg)	59.5 ±4.7	48.7 ±6.1	50.6 ±10.4	65.7 77.0	74.9 86.5
Mean Weight 5 Largest	67.6 ±6.5	61.9 ±6.4	60.0 ±1.3	77.0 ±2.5	86.5 ±7.2

TABLE 13

Growth Rate of Manduca sexta Larvae Feeding on Tobacco Leaves from Plants Transformed with pGS1110

Exp. 1: Number of Larvae in a Certain Stage After 87 h:

Plant No:	C1	C2	C3	N20-3	N20-46	N20-38	N20-22	N20-47	N20-18	N20-30	N20-31
Stage L1	6	0	3	7	5	15	12	4	14	6	0
L2	14	20	17	13	14	4	6	16	6	13	20
Dead	0	0	0	0	1	1	2	0	0	1	0

Exp. 2: Number of Larvae in a Certain Stage after 78 h:

Plant No:	C4	C5	C6	C7	N20-35	N20-37	N20-7(*)	N20-7(*)	N20-19	N20-13	N20-1
Stage L1	0	0	0	1	1	11	1	0	0	1	0
L2	20	20	20	19	19	9	19	20	20	19	20
Dead	0	0	0	0	0	0	0	0	0	0	0

(*) Two copies of this plant were tested in this experiment.

TABLE 14

Characteristics of Plants from Experiment No. 20

Plant Number	Nos.	Km^R	Insect Tox.
N20-4	+	+	-
N20-30	+	+	-
N20-18	N.T. (*)	+	+
N20-22	+	+	+
N20-3	-	+	-
N20-46	N.T.	N.T.	-
N20-38	+	+	+
N20-31	+	+	-
N20-37	+	+	+
N20-7	+	+	-
N20-35	+	+	-
N20-13	-	N.T.	-
N20-19	+	N.T.	-
N20-1	-	N.T.	-

(*) N.T. = Not Tested

TABLE 15

Growth Rate and Mortality of Manduca sexta Larvae Feeding on Leaves From Tobacco Plants Transformed with pGS1151 (Experiment I)

Represented are:

Numbers of larvae in a certain stage (L1, L2 or L3) or dead (D) from groups of 20 larvae after a period of feeding on the tobacco leaves.

Time	Plant N21-50				N21-35				N21-11				N21-56			
(Hours)	D	L1	L2	L3	D	L1	L2	L3	D	L1	L2	L3	D	L1	L2	L3
0		20				20				20				20		
55		20			5	15				20				20		
61	1	19			5	14	1		1	19				20		
66	1	19			5	11	4		1	19				19		
71	1	19			6	5	9		3	19			1	9	10	
76	1	18	1		7	4	9		5	15			1	8	11	
81	1	18	1		7	4	9		5	15			2	7	11	
87	1	18	1		7	4	9		5	15			2	7	11	
92	2	17	1		8	3	9		8	12			2	3	15	
119	11	7	2		12	1	7		18	2			3	1	16	
136	12	4	4		12		8		19	1			4		16	
144	12	4	4		15		5		19	1			4		16	
159	13	3	4		17		3		20				4		16	
168	15	1	4		17		2	1	20				4		15	1

TABLE 15 (cont'd)

Time	N21-107(*)				N21-18				N21-43				N21-53			
(Hours)	D	L1	L2	L3	D	L1	L2	L3	D	L1	L2	L3	D	L1	L2	L3
0		20				20				20				20		
55		20				20			1	19				20		
61		19	1			20			1	19				20		
66	1	10	9			20			1	19				20		
71	2	6	12		1	16	3		1	16	3			20		
76	2	6	12		1	14	5		2	15	3			16	4	
81	2	6	12		1	13	6		2	15	3		4	13	3	
87	2	2	16		1	12	7		3	14	3		5	11	4	
92	2		18		1	12	7		4	12	4		8	9	3	
119	2		18		9	3	8		6	7	7		17	1	2	
136	2		18		14		6		9	4	7		18	1	1	
144	2		18		16		4		10	4	6		18		2	
159	2		12	6	17		3		12	2	6		18		2	
168	3		8	9	17		3		15		5		18		2	

*Plant N21-107 is a control plant transformed with the same type of vector but comprising only a PTR:NPTII chimeric gene and no Bt2 sequences.

TABLE 16

Growth Rate and Mortality of Manduca sexta Larvae Feeding on Leaves from Tobacco Plants Transformed with pGS1151 (Experiment II) (See also Legend for Table 15)

Time	N21-50		N21-18		N21-43		N21-11		N21-56		N21-35		N21-53		N21-33		Controls N21-102			N21-104			N21-107		
(hours)	D	L1	D	L1	D	L1	D	L1	D	L1	D	L1	D	L1	D	L1	D	L1	L2	D	L1	L2	D	L1	L2
0		20		20		20		20		20		20		20		20		20			20			20	
29												20		20		20		20			20			20	
47											5	15	6	14	8	12		20			20			20	
51	9	11	8	12	2	18	18	2	2	18															
57											8	12	15	5	15	5		20			20			20	
69	16	4	16	4	10	10	20		3	17															
79	19	1	18	2	11	9	20		3	17	16	4	18	2	20			14	6		15	5		20	
96											17	3	20		20			4	16		12	8	1	18	1
100	19	1	18	2	14	6	20		10	10															
118	20		19	1	18	2	20		15	5															
120											18	2	20		20				20			20	1	13	6

*Plants N21-102, 104, 107 are control plants transformed with PTR:NPTII.

TABLE 17

Percentage mortality and mean weight of Manduca sexta larvae after a certain period of feeding on tobacco leaves from plants transformed with pGS1151. Complete results from the 2 independent Experiments I and II (Tables 15 and 16) are compiled. Kanamycin resistance levels of the plants expressing the Bt:NPT2 fusion protein are also given (ug/ml Km on which good callus growth still occurs).

Plant No.	Km^R (ug/ml Km)	% Mortality Exp. I (after 168 h)	Exp. II (after 118 h) (or 120 h*)	Mean Weight Surviving Larvae (mg/larva) Exp. I (after 168 h)
N21-3 34.0	200	15	N.T.	
5	200	30	N.T.	52.4
11	500	100	100	--
12	500	40	N.T.	16.6
16	200	45	N.T.	25.3
17	500	75	N.T.	13.4
18	500	85	95	9.0
23	500	90	100*	12.5
29	200	55	N.T.	21.9
32	200	50	N.T.	27.4
33	500	40	N.T.	27.7
35	500	85	90	18.7
40	200	20	N.T.	28.6
41	200	15	N.T.	29.1
42	200	55	N.T.	18.7
43	500	75	90	15.5

TABLE 17 (cont'd)

Plant No.	Km^R (ug/ml Km)	% Mortality Exp. I (after 168 h)	Exp. II (after 118 h) (or 120 h*)	Mean Weight Surviving Larvae (mg/larva) Exp. I (after 168 h)
45	200	30	N.T.	13.7
50	500	75	100	10.7
53	500	90	100*	12.5
56	200	20	75	22.4
Controls:				
N21-102	--	N.T.	0*	N.T.
104	--	N.T.	0*	N.T.
107	--	15	5*	44.1

N.T. = Not Tested

TABLE 18

Frequency of Nopaline Positive Plants in the F_1 Generation Derived from Transformed Tobacco Plants

Plant No. of Parental Plant	Age of the Seedlings Tested (wks)	Total Number of Plants Tested	Nopaline Positive	% Nopaline Positives
147-8	3	74	56	76%
	7	13	11	85%
10-1	3	25	20	80%
	7	9	7	78%
161-9	3	66	18 (x)	27%
	7	107	81	76%
174	6	45	43	95%

(x) Nopaline Signal Very Weak.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A chimeric gene capable of being expressed in cells of a plant as an insecticidal Bacillus thuringiensis polypeptide toxin and of thereby rendering the plant insect-resistant; the chimeric gene comprising:

(a) a first DNA fragment that encodes a Bacillus thuringiensis insecticidal crystal protein which has been truncated towards a trypsin cleavage site of the protein; and

(b) a promoter region and a 3' non-translated region containing a polyadenylation signal; the first DNA fragment being under the control of the promoter region; the promoter and 3' non-translated regions allowing the first DNA fragment to be expressed in the cells;

the chimeric gene being free of: (i) an ATG triplet between the promoter region and an initiation codon of the first DNA fragment; and (ii) a T-DNA oncogenic tumor-inducing gene.

2. The chimeric gene of claim 1, wherein the first DNA fragment encodes a polypeptide that has a molecular weight of approximately 60 to 80 kD.

3. The chimeric gene of claim 1 or claim 2, wherein the first DNA fragment encodes a polypeptide that has a molecular weight of approximately 60 kD.

4. The chimeric gene of any one of claims 1-3, wherein the first DNA fragment encodes a polypeptide active against Lepidoptera.

5. The chimeric gene of any one of claims 1-4, wherein the first DNA fragment encodes: a truncated 130 kD crystal protein of Bacillus thuringiensis berliner 1715; a truncated 130 kD crystal protein of Bacillus

STRALIAN
LS
ENT OFFICE

thuringiensis Kurstaki; or a truncated 130 kD crystal protein of Bacillus thuringiensis sotto.

6. The chimeric gene of any one of claims 1-5, wherein the first DNA fragment encodes a truncated Bt2 protein.

7. The chimeric gene of any one of claims 1-3, wherein the first DNA fragment encodes a polypeptide active against Coleoptera.

8. The chimeric gene of any one of claims 1-7, wherein the promoter region is Pnos, Pssu Pea, PTR2, Pssu301, P35S-1 or P35S-2.

9. The chimeric gene of any one of claims 1-8, also comprising a second DNA fragment which encodes an enzyme capable of being expressed in the cells and the expression of which can be identified in the cells; the second DNA fragment being fused to the first DNA fragment so that the first and second DNA fragments encode a fusion polypeptide; whereby an identification of expression of the second DNA fragment in the cells provides an identification of expression of the first DNA fragment in the cells.

10. The chimeric gene of claim 9, wherein the second DNA fragment encodes a selectable or scorable marker.

11. The chimeric gene of claim 9 or claim 10, wherein the second DNA fragment encodes, as a selectable marker, neomycin phosphotransferase.

12. The chimeric gene of any one of claims 9-11, wherein the second DNA fragment is fused to a side of the first DNA fragment containing a trypsin cleavage site.

STRALIAN LS TENT OFFICE

13. A chimeric gene capable of being expressed in cells of a plant as an insecticidal Bacillus thuringiensis polypeptide toxin and of thereby rendering the plant insect-resistant; the cells being free of a T-DNA oncogenic tumor-inducing gene; the chimeric gene comprising:

(a) a first DNA fragment that encodes a Bacillus thuringiensis insecticidal crystal protein;

(b) a promoter region and a 3' non-translated region containing a polyadenylation signal; the first DNA fragment being under the control of the promoter region; the promoter and 3' non-translated regions allowing the first DNA fragment to be expressed in the cells; and

(c) a second DNA fragment which encodes an enzyme capable of being expressed in the cells and the expression of which can be identified in the cells; the second DNA fragment being fused to the first DNA fragment so that the first and second DNA fragments encode a fusion polypeptide;

whereby an identification of expression of the second DNA fragment in the cells provides an identification of expression of the first DNA fragment in the cells;

the chimeric gene being free of an ATG triplet between the promoter region and an initiation codon of the first DNA fragment.

14. The chimeric gene of claim 13, wherein the second DNA fragment encodes a selectable or scorable marker.

15. The chimeric gene of claim 13 or claim 14, wherein the second DNA fragment encodes, as a selectable marker, neomycin phosphotransferase.

AUSTRALIAN PATENT OFFICE LS

16. The chimeric gene of any one of claims 13-15, wherein the first DNA fragment encodes a protein active against Lepidoptera.

17. The chimeric gene of any one of claims 13-16, wherein the first DNA fragment encodes: a 130 kd crystal protein of Bacillus thuringiensis berliner 1715; a 130 kD crystal protein of Bacillus thuringiensis Kurstaki; or a 130 kD crystal protein of Bacillus thuringiensis· sotto.

18. The chimeric gene of any one of claims 13-17, wherein the first DNA fragment encodes a Bt2 protein.

19. The chimeric gene of any one of claims 13-15, wherein the first DNA fragment encodes a protein active against Coleoptera.

20. The chimeric gene of any one of claims 13-19, wherein the promoter region is Pnos, Pssu Pea, PTR2, Pssu301, P35S-1 or P35S-2.

21. A vector comprising:

(a) the chimeric gene of any one of claims 1-20; and

(b) at least one border sequence of a T-DNA region of a Ti-plasmid allowing integration of the chimeric gene into a plant cell genome; the vector being free of an oncogenic tumor-inducing gene which can be expressed in a plant cell.

22. The vector of claim 21, also comprising:

(c) a virulence region of a Ti-plasmid allowing a transfer of the chimeric gene into a plant cell genome.

23. The vector of claim 21 or claim 22, comprising at least a right border sequence of the T-DNA region.

24. The vector of any one of claims 21-23, harbored in an Agrobacterium strain.

25. A method for protecting a plant against an insect pest comprising the step of: providing genomes of cells of the plant with the chimeric gene of any one of claims 1-20, whereby the cells can express an insecticidal Bacillus thuringiensis polypeptide toxin.

26. An insect-resistant plant which is capable of being transformed by Agrobacterium, having cells containing in their genome the chimeric gene of any one of claims 1-20, whereby the cells are capable of expressing an insecticidal Bacillus thuringiensis polypeptide toxin.

27. A seed of an insect-resistant plant which is capable of being transformed by Agrobacterium, containing in its genome the chimeric gene of any one of claims 1-20.

28. A tissue of an insect-resistant plant which is capable of being transformed by Agrobacterium, having cells containing in their genome the chimeric gene of any one of claims 1-20, whereby the cells are capable of expressing an insecticidal Bacillus thuringiensis polypeptide toxin.

29. A plant cell containing in its genome the chimeric gene of any one of claims 1-20 whereby the cell is capable of expressing an insecticidal Bacillus thuringiensis polypeptide toxin.

30. A culture of cells having the identifying characteristics of a culture of cells having an accession number of DSM 3127 or DSM 3128 or DSM 3129 or DSM 3130 or ATCC 53385 or ATCC 53386 or ATCC 53387 or ATCC 53388 or ATCC 53389 or ATCC 53390 or ATCC 53392 or ATCC 53393 or ATCC 53394 or ATCC 53395 or ATCC 53396 or ATCC 53397 or

LIAN U.S. PATENT OFFICE

ATCC 53398 or ATCC 53399 or ATCC 53400 or ATCC 53401 or ATCC 53402 (as hereinbefore described) from such a culture.

31. A plasmid having the identifying characteristics of plasmids harbored within a culture of cells having an accession number of DSM 3127 or DSM 3128 or DSM 3129 or DSM 3130 or ATCC 53385 or ATCC 53386 or ATCC 53387 or ATCC 53388 or ATCC 53389 or ATCC 53390 or ATCC 53392 or ATCC 53393 or ATCC 53394 or ATCC 53395 or ATCC 53396 or ATCC 53397 or ATCC 53398 or ATCC 53399 or ATCC 53400 or ATCC 53401 or ATCC 53402 (as hereinbefore described).

32. A chimeric gene of claim 1 or claim 13, a vector comprising a said chimeric gene, or a plant or plant tissue having cells containing in their genome a said chimeric gene, substantially as hereinbefore described with reference to the Examples.

Dated this 4th day of July, 1989.

PLANT GENETIC SYSTEMS N.V.,
By its Patent Attorneys
DAVIES & COLLISON

AUSTRALIAN PATENT OFFICE




FIG.1.

FIG.2.




FIG. 3.




FIG.4.




FIG.5.




FIG.6.




FIG.7.




FIG. 8.



O.D
2.0
1.5
1.0
0.5
0
Bt 2
Bt BERLINER CRY
Bt KURSTAKI CRY
EXTRACT E COLI -
(NEGATIVE CONTROL)
10⁴ 10³ 10² 10 1 01 001 0
Ng/ML PROTEIN

FIG. 9.



A
Bt BERLINER
O.D
1
0
B
Bt 2
1
0
1F6
1G7
2H5
4F6
4G6
4F3
4F11
4G2
4G3
4G11
4H9
8G10
9H11
10E3
1.7
4.8
C73
MONOCLONAL ANTIBODY CODE No:


FIG. 10.

COMPARISON OF N-TERMINAL AMINO ACID SEQUENCES OF 130 Kd CRYSTAL PROTEINS

1 Bt WHITELEY : Met-Asp-Asn-Asn-Pro-Asn-Ile-Asn-Glu-Cys-Ile-Pro-Tyr-Asn-Cys-Leu-Ser-Asn-Pro

2 Bt2 : ?-Asp-Asn-Asn-Pro-Asn-Ile-Asn-Glu-?-Ile-Pro-Tyr-Asn-?-Leu-?-Asn-Pro

FIG. 11.





C
B
A
Cla I
EcoRI
Xba I
Xba I
EcoRI
Acc I
Cla I
Sac I
EcoRI
EcoRI
Hiii
Bcl I
Kpn I
Xba I
Mlu I
Hiii
RSA I
Tag I
RUii
pVuII
Kpn I
Ndc I

FIG. 12.


-CAAAAATTGATATTTAGTAAAATTAGTTGCACTTTGTGCATTTTTTCATAAGATGAGTCATATGTTTTAAATTGTAGTAATGAAAAACAGTATTATATCATAATGAATTGGTATCTTAA
O
100
Met AspAsnAsnProAsnIleAsnGluCysIleProTyrAsnCysLeuSerAsnProGluValGluValLeuGlyGlyGluArgIleGluThrGlyTyr
TAAAAGAGATGGAGGTAACTTATGGATAACAATCCGAACATCAATGAATGCATTCCTTATAATTGTTTAAGTAACCCTGAAGTAGAAGTATTAGGTGGAGAAAGAATAGAAACTGGTTAC
200
ThrProIleAspIleSerLeuSerLeuThrGlnPheLeuLeuSerGluPheValProGlyAlaGlyPheValLeuGlyLeuValAspIleIleTrpGlyIlePheGlyProSerGlnTrp
ACCCCAATCGATATTTCCTTGTCGCTAACGCAATTTCTTTTGAGTGAATTTGTTCCCGGTGCTGGATTTGTGTTAGGACTAGTTGATATAATATGGGGAATTTTTGGTCCCTCTCAATGG
300
AspAlaPheLeuValGlnIleGluGlnLeuIleAsnGlnArgIleGluGluPheAlaArgAsnGlnAlaIleSerArgLeuGluGlyLeuSerAsnLeuTyrGlnIleTyrAlaGluSer
GACGCATTTCTTGTACAAATTGAACAGTTAATTAACCAAAGAATAGAAGAATTCGCTAGGAACCAAGCCATTTCTAGATTAGAAGGACTAAGCAATCTTTATCAAATTTACGCAGAATCT
400
PheArgGluTrpGluAlaAspProThrAsnProAlaLeuArgGluGluMetArgIleGlnPheAsnAspMetAsnSerAlaLeuThrThrAlaIleProLeuPheAlaValGlnAsnTyr
TTTAGAGAGTGGGAAGCAGATCCTACTAATCCAGCATTAAGAGAAGAGATGCGTATTCAATTCAATGACATGAACAGTGCCCTTACAACCGCTATTCCTCTTTTTGCAGTTCAAAATTAT
500
600
GlnValProLeuLeuSerValTyrValGlnAlaAlaAsnLeuHisLeuSerValLeuArgAspValSerValPheGlyGlnArgTrpGlyPheAspAlaAlaThrIleAsnSerArgTyr
CAAGTTCCTCTTTTATCAGTATATGTTCAAGCTGCAAATTTACATTTATCAGTTTTGAGAGATGTTTCAGTGTTTGGACAAAGGTGGGGATTTGATGCCGCGACTATCAATAGTCGTTAT
600
700
AsnAspLeuThrArgLeuIleGlyAsnTyrThrAspHisAlaValArgTrpTyrAsnThrGlyLeuGluArgValTrpGlyProAspSerArgAspTrpIleArgTyrAsnGlnPheArg
AATGATTTAACTAGGCTTATTGGCAACTATACAGATCATGCTGTACGCTGGTACAATACGGGATTAGAGCGTGTATGGGGACCGGATTCTAGAGATTGGATAAGATATAATCAATTTAGA
800
ArgGluLeuThrLeuThrValLeuAspIleValSerLeuPheProAsnTyrAspSerArgThrTyrProIleArgThrValSerGlnLeuThrArgGluIleTyrThrAsnProValLeu
AGAGAATTAACACTAACTGTATTAGATATCGTTTCTCTATTTCCGAACTATGATAGTAGAACGTATCCAATTCGAACAGTTTCCCAATTAACAAGAGAAATTTATACAAACCCAGTATTA
900
GluAsnPheAspGlySerPheArgGlySerAlaGlnGlyIleGluGlySerIleArgSerProHisLeuMetAspIleLeuAsnSerIleThrIleTyrThrAspAlaHisArgGlyGlu
GAAAATTTTGATGGTAGTTTTCGAGGCTCGGCTCAGGGCATAGAAGGAAGTATTAGGAGTCCACATTTGATGGATATACTTAACAGTATAACCATCTATACGGATGCTCATAGAGGAGAA
1000
TyrTyrTrpSerGlyHisGlnIleMetAlaSerProValGlyPheSerGlyProGluPheThrPheProLeuTyrGlyThrMetGlyAsnAlaAlaProGlnGlnArgIleValAlaGln
TATTATTGGTCAGGGCATCAAATAATGGCTTCTCCTGTAGGGTTTTCGGGGCCAGAATTCACTTTTCCGCTATATGGAACTATGGGAAATGCAGCTCCACAACAACGTATTGTTGCTCAA
1100
1200
LeuGlyGlnGlyValTyrArgThrLeuSerSerThrLeuTyrArgArgProPheAsnIleGlyIleAsnAsnGlnGlnLeuSerValLeuAspGlyThrGluPheAlaTyrGlyThrSer
CTAGGTCAGGGCGTGTATAGAACATTATCGTCCACTTTATATAGAAGACCTTTTAATATAGGGATAAATAATCAACAACTATCTGTTCTTGACGGACAGAATTTGCTTATGGAACCTCC
1200
1300
SerAsnLeuProSerAlaValTyrArgLysSerGlyThrValAspSerLeuAspGluIleProProGlnAsnAsnAsnValProProArgGlnGlyPheSerHisArgLeuSerHisVal
TCAAATTTGCCATCCGCTGTATACAGAAAAAGCGGAACGGTAGATTCGCTGGATGAAATACCGCCACAGAATAACAACGTGCCACCTAGGCAAGGATTTAGTCATCGATTAAGCCATGTT
1400
SerMetPheArgSerGlyPheSerAsnSerSerValSerIleIleArgAlaProMetPheSerTrpIleHisArgSerAlaGluPheAsnAsnIleIleProSerSerGlnIleThrGln
TCAATGTTTCGTTCAGGCTTTAGTAATAGTAGTGTAAGTATAATAAGAGCTCCTATGTTCTCTTGGATACATCGTAGTGCTGAATTTAATAATATAATTCCTTCATCACAAATTACACAA
1500
IleProLeuThrLysSerThrAsnLeuGlySerGlyThrSerValValLysGlyProGlyPheThrGlyGlyAspIleLeuArgArgThrSerProGlyGlnIleSerThrLeuArgVal
ATACCTTTAACAAAATCTACTAATCTTGGCTCTGGAACTTCTGTCGTTAAAGGACCAGGATTTACAGGAGGAGATATTCTTCGAAGAACTTCACCTGGCCAGATTTCAACCTTAAGAGTA
1600
AsnIleThrAlaProLeuSerGlnArgTyrArgValArgIleArgTyrAlaSerThrThrAsnLeuGlnPheHisThrSerIleAspGlyArgProIleAsnGlnGlyAsnPheSerAla
AATATTACTGCACCATTATCACAAAGATATCGGGTAAGAATTCGCTACGCTTCTACCACAAATTTACAATTCCATACATCAATTGACGGAAGACCTATTAATCAGGGGAATTTTTCAGCA
1700
1800
ThrMetSerSerGlySerAsnLeuGlnSerGlySerPheArgThrValGlyPheThrThrProPheAsnPheSerAsnGlySerSerValPheThrLeuSerAlaHisValPheAsnSer
ACTATGAGTAGTGGGAGTAATTTACAGTCCGGAAGCTTTAGGACTGTAGGTTTTACTACTCCGTTTAACTTTCAAATGGATCAAGTGTATTTACGTTAAGTGCTCATGTCTTCAATTCA
1800
1900
GlyAsnGluValTyrIleAspArgIleGluPheValProAlaGluValThrPheGluAlaGluTyrAspLeuGluArgAlaGlnLysAlaValAsnGluLeuPheThrSerSerAsnGln
GGCAATGAAGTTTATATAGATCGAATTGAATTTGTTCCGGCAGAAGTAACCTTTGAGGCAGAATATGATTTAGAAAGAGCACAAAAGGCGGTGAATGAGCTGTTTACTTCTTCCAATCAA
2000
IleGlyLeuLysThrAspValThrAspTyrHisIleAspGlnValSerAsnLeuValGluCysLeuSerAspGluPheCysLeuAspGluLysLysGluLeuSerGluLysValLysHis
ATCGGGTTAAAAACAGATGTGACGGATTATCATATTGATCAAGTATCCAATTTAGTTGAGTGTTTATCTGATGAATTTTGTCTGGATGAAAAAAAAGAATTGTCCGAGAAAGTCAAACAT
2100

FIG.13A.

98/85425




FIG.13B.

52458/86

FIG.14A.

SEQUENCE COMPARISON OF FOUR BACILLUS THURINGIENSIS TOXINS

```
          10                  20                  30                  40                  50
MDNNPNINEC  IPYNCLSNPE  VEVLGGERIE TGYTPIDISL  SLTQFLLSEF

          60                  70                  80                  90                 100
VPGAGFVLGL  VDIIWGIFGP  SQWDAFLVQI EQLINQRIEE  FARNQAISRL
                                    P

         110                 120                 130                 140                 150
EGLSNLYQIY  AESFREWEAD  PTNPALREEM RIQFNDMNSA  LTTAIPLFAV
                                                                                        L

         160                 170                 180                 190                 200
QNYQVPLLSV  YVQAANLMLS  VLRDVSVFGQ RWGFDAATIN  SRYNDLTRLI

         210                 220                 230                 240                 250
GNYTDHAVRW  YMTGLERVWG  PDSRDWIRYN QFRRELTLTV  LDIVSLFPNY
      Y                             V                             A
      Y                             V                             A   S
      Y                             V                             A   S
         260                 270                 280                 290                 300
DSRTYPIRTV  SQLTREIYTN  PVLENFDGSF RGSAQGIEGS  IRSPHLMDIL
   R                                                        R
   R                                                    M   R   QN      Q
   R                H                                   M   R   QN      Q
         310                 320                 330                 340                 350
NSITIYTDAH  RGEYYWSGHQ  IMASPVGFSG PEFTFPLYGT  MGNAAPQQRI
                    Y
         V          FN             T                   A       F N     A       PV-L
 R       V          FN             T                   A       VF N    A       PV-L
         360                 369                 379                 389                 398
VAQLGQGVYR  TLSSTLYRR-  PFNIGINNQQ LSVLDGTEFA  YGTSS-NLPS

 SLT L IF       P     I  ILGS P    E  F        S  FASLTT
 SLT L IF       P     I  ILGS P    E  F        S  FASLTT
         408                 418                 428                 438                 448
AVYRKSGTVD  SLDEIPPQNN  NVPPRQGFSH RLSHVSMFRS  GFSNSSVSII

TI   QR         V     D  S     A          T LSQ  AAGAVYTL--
TI   QR         V     D  S     A          T LSQ  AAGAVYTL--
         458                 468                 478                 488                 498
RAPMFSWIHR  SAEFNNIIPS  SQITQIPLTK STNLGSGTSV  VKGPGFTGGD
                    A   DS       AV  GNF FN -    IS
                                                                                        H
    T    Q
    T    Q
         508                 515                 523                 533                 543
ILRRTSPGQI  STLRVNI---  -TAPL-SQRY RVRIRYASTT  NLQFHTSIDG
LV LN S NN  IQN GY EVP  IHF ST T      V     V  PIHLMVNWGN

         553                 563                 573                 583                 593
RPINQGNFSA  TMSSGSNLQS  GSFRTVGFTT PFNFSNGSSV  FTLSAHVFNS
SS FSNTVP    AT LD      SD---F YFE SA AFTS LG  NIVGVRN SG
```

FIG. 14B.

```
          603                 613                 623                 633                 643
GNEVYIDRIE  FVPAEVTFEA EYDLERAQKA VNELFTSSNQ IGLKTDVTDY
TAG  I   F   I VTA  L      N              A    T   L     M

                                                    I
          653                 663                 673                 683                 693
HIDQVSNLVE  CLSDEFCLDE KKELSEKVKH AKRLSDERNL LQDPMFRGIN
         T  Y             R                          S   KD
                          Q
                          Q
          703                 713                 723                 733                 743
RQLDRGWRGS  TDITIQGGDD VFKENYVTLL GTFDECYLTY LYQKIDESKL
  PE    G     G                    S              P
                                                    P
                                                    P
          753                 763                 773                 783                 793
KAYTRYQLRG  YIEDSQDLEI  YLIRYNAKHE TVNVPGTGSL WRLSAPSPIG
  F                                                 P   Q
                                                    P   Q
                                                    P   Q
                                    797                 807                 817
----------  ----------  ------KCAH HSHHFSLDID VGCTDLNEDL
KCGEPNRCAP  HLEWNPDLDC  SCRDGE
KCGEPNRCAP  HLEWNPDLDC  SCRDGE                          H
KCGEPNRCAP  HLEWNPDLDC  SCRDGE   R
          827                 837                 847                 857                 867
GVWVIFKIKT  QDGHARLGNL  EFLEEKPLVG EALARVKRAE KKWRDKREKL

                                                    -
          877                 887                 897                 907                 917
EWETNIVYKE  AKESVDALFV NSQYDRLQAD TNIAMIHAAD  KRVHSIREAY
                                Q
                                Q
                        K                                   ***
          927                 937                 947                 957                 967
LPELSVIPGV  MAAIFEELEG RIFTAFSLYD  ARNVIKNGDF NNGLSCWNVK

          977                 987                 997                 1007                1017
GHVDVEEQNN  HRSVLVVPEW EAEVSQEVRV CPGRGYILRV TAYKEGYGEG
            Q
            Q       L

          1027                1037                1047                1057                1067
CVTIHEIENN  TDELKFSNCV EEEVYPNNTV TCNDYTATQE EYEGTYTSRN
                              I                 VN      G  A
                              I                 VN      G  A

          1077                1087                1097                1107                1117
RGYDGAYESN  SSVPADYASA  YEEKAYTDGR RDNPCESNRG YGDYTPLPAG
   NE PS--  --          V       S          E    F       R        V
   NE PS--  --          V       S          E    F       R        V

          1127                1137                1147        1155
YVTKELEYFP  ETDKVWIEIG ETEGTFIVDS VELLLMEE
```

FIG. 15.





BamHI
HpaI
Bt2
NdeI
PstI
pBt 200
HpaI digestion
Bal 31 treatment
add Bam linkers + T4 ligase
Bam digestion
T4 ligase
pHD 100
BamHI
Bt 2
NdeI
GGGATCCCGGTAACTTATG...
Bt2
Bam linker
Nde I digestion
Bal 3i treatment
BglII linkers + T4 ligase
BglII digestion
T4 ligase
BamHI
Bt2 cassette
BglII
pHD 160
± 300bp


FIG.16.


BamHI
SacI
BglII
pHD160
Bt2
GGATCCCG TAACTT'ATG' GAT'...
Bam HI
+1
Bt2
0
1000
2000
3000
3465
4000
kb
Bam
CGGGATCCCG
BglII
CAGATCTG
AGAGATGGAGGTAACTTATGGATAACAATCCG
Met Asp Asn Asn Pro
CACACCACTACCAGC
3660
3860
Bt cassette
1
2
3
4
5
pHD160
pHD162
pHD163
pHD164
pHD159
BamHI
Bt2
pHD164 : TAATATATGCTTTAAA
stop
DraI
pDC3 : TAATATAGCTTTCAGATCT
BglII

STRATEGY USED FOR THE POSITIONING OF THE TOXIN GENE BEHIND THE λ-P_L PROMOTOR.

FIG.17.





PLASMID	ADAPTOR
pLK54	AATTCCCGGGGATCCGTCGACCTGCAGCCAAGCTTGGTCTAGAGGTCGA EcoRI SmaI BamHI SalI PstI HindIII XbaI
pLK57	AATTCCCGGGAGAGCTCGATATCGCATGCGGTACCTCTAGAAG EcoRI SmaI SacI EcoRV SphI KpnI XbaI AAGCTTGGGATCCGTCGACCTGCAGATCTGCTAGAGGTCGA HindIII BamHI SalI PstI BglII

FIG.18.

Construction of pLBKm25




FIG. 19.

Construction of random Bt2:NPTII fusions and Bt2 deletions




FIG. 20.




BamHI
HpaI
HpaI
HpaI
Bt2
PstI
HpaI-160
-32
0
ClaI 110
XbaI 280
XbaI 660
EcoRV720
ClaI 1280
EcoRV1560
HindIII1680
KpnI 2170
XbaI2500
HindIII2740
TAA 3466
KpnI 3650
PstI 4340
pLB 10
pLB 12
pLB 16
HindIII
KpnI
1500
1600
1700
1800
1900
2000
2100
885
833
881
950.876
834
879.884
828
878
826
822
886
839
820
1 Kb

FIG.21.



1. Trypsin digest Bt2
2. pLB 950
3. pLB 834
4. pLB 879
5. pLB 884
6. pLB 828
7. pLB 886
8 NF1 control

FIG. 22.

Delineation of the toxic fragment using C-terminal deletions






FIG. 23.

Construction of Bt: NPT II Cassettes





Plasmid	5' ends of the Bt2 gene	Purpose
pLBKm 13	GGAT'CCC'GAT... +4 Bt2	Fusion at initiator ATG
pLBKm 23	GGATCCCGTGGTATCTTAATTAAAAGAGATG GAGGTAACTT'ATG'GAT ... +1 Bt2	Expression in E. coli
pLBKm 33	GGATCCCGTAACTT'ATG'GAT... BamHI +1 Bt2	Fusion to plant promotor

FIG.24.




pLBKm13
BamHI
Sac I
NCO I
DdeI
BglII
Bt2
NPT II
GGT ACG ATC CGG CCA AGC TTG GAT
2173
13
GGATCCC'GAT....
+4 Bt2
pLBKm14
BamHI
BglII
Bt2
NPT II
GGATCCCGTGGTATCTTAATTAAAAGAGATG
GAGGTAACTT'ATG'GAT.....
+1 Bt2
pLBKm860 (865)
Deletion
BamHI
SacI
NCOI
BglII
S.D.
Bt2
NPT II
pLBKm1860(1865)
BamHI
Sac I
NCO I
BglII
Bt2
NPT II
pLBKm2860
BamHI
SacI
NCO I
BglII
Bt2
NPTII
GGATCCCGTAACTT'ATG'GAT...
BamHI
+1 Bt2

FIG. 25.



23 = Bt : NPT2

860 = Bt : NPT860

865 = Bt : NPT865

NPT = NPT II intact

NF1 = control extract



HdIII
KpnI
HdIII
T
T
Bt2
0
1000
2000
3000
HdIII
T
KpnI
1500
1700
1900
2100
2300
pLB12
pLB884
pLB820
pLB16
pLBKm865
pLBKm22
pLBKm860
pLBKm23

FIG. 26.

FIG. 27.

Adaption of Bt2 and Bt:NPTII cassettes for expression in plant cells:




FIG. 28.

Promotor - Bt gene junction sequences

1) pHD1050, pHD1060, pGS1110: Pnos - Bt

CATAAATTCCCCTCGGTATCCAATTAGAGTTCTGATCGACGGATCCCGTAACTT'ATG'GAT

BamHI Bt2

2) pHD1076: Pssu pea - Bt

TAAAAACATTATATATAGCAAGTTTTAGCAGAAGCTTGGCTGCAGGTCGACGGATCCCGTAACTT'ATG'GAT

HindIII BamHI Bt2

3) pHD1080: Tp - Bt fusion

5'

TAAAAACATTATATATAGCAAGTTTTAGCAGAAGCTTTGCAATTCATACAGAAGTGAGAAAA

HindIII

'ATG'...'AGA'GTA'AAG 'TGC'ATG'GAT'CCC'GAT'AAC'AAT

TP BamHI +4 Bt2

4) pGS1151, pGS1152, pGS1153, pGS1161, pGS1162, pGS1163: PTR_2-Bt

ATACACCAAAATCGATGGAT'CCC'GAT

ClaI BamHI +4 Bt2

5) pGS1171, pGS1181: Pssu 301 - Bt

AAGCAAAATTCTTCTAACC'ATG'GAT'CCC'GAT'

NcoI +4 Bt2

6) pGS1251, pGS1252, pGS1261, pGS1262: P35S1 - Bt

CTGAAATCACCAGTCTCGGATCCCGTAACTT'ATG'GAT

BamHI Bt2

pos 22 from RNA start site

7) pGS1271, pGS1281: P35S2 - Bt

CAGTCTCTCTCTACAAATCGGATCCCGTAACTT'ATG'GAT

BamHI Bt2

pos 36 from RNA start site

FIG.29.



BamHI
HpaI
Neo
Ap
pGV831
Sp
HpaI
T4 Ligase
ocs
3'
pGV99
PvuII
3'
B
3'
Neo
Ap
pGV858
Sp
BamHI
BglII / BamHI
T4 -Ligase
E Bg H B
Pssu
Ap
pGV861
Km
T4 Ligase
Bg
E
H
Pssu
Ap
pKC7::Pssu
Km
B
H/B
pUC8 H/B
B
Pssu
3'
Neo
Ap
pHD503
Sp
BamHI
BamHI / BglII
T4 Ligase
Bt
B
Bg
pHD 160
B
Bt
3'
Pssu
Neo
pHD 208
Ap
Sp

FIG. 30.



T_L-DNA
T_R-DNA
H Bg Bg B B E B E H
5 7 2 1 4 6a 6b 3
pGV700
Cm
Cb
pGV600
HindIII
pGV742
Δ BamHI
pGV744
Δ EcoRI
pGV749
pGV710
HindIII/
NruI
EcoRI
DNA polym
HindIII
pGV815
Sm
Eco RI/DNA polym
HindIII/DNA polym
pKC7 nos
Neo
BclI/
BamHI
pGV825
BglII
pGV831
Tc
Su

FIG. 31.



H
H
H
H
pGV 3850
Ap
Bt
pHD 205
Km
Ap
H
H
H
H
H
H
H
pHD 1050
Ap
nos
Km
Ap
Bt
2kb

FIG. 32.




BamHI
Eco RI
pGV2260
Ap
Ap
B.t.
pHD 208
Sp
Pssu
Pnos
Km
Pst I
Pst I
Pst I
Pst I
Pst I
Pst I
Pst I
Sp
Ap
Ap
pHD 1076
1
2
3
4
5
6
7
8
9
T DNA border
Bt cassette
Pssu
Kan
Pnos
T DNA border
T-DNA
1kb

FIG.33A.


Expression Vectors



pLGV2382 : Herrera-Estrella et al, (1983) EMBOJ, 2, 987

FIG.33B.

Expression Vectors




derived from pGV831 (DeBlaere et al, 1985)

Pnos: bp BamHI-BclI fragment of pLGV23

3'ocs: 706 bp PVUII fragment of ocs in HpaI site of pGV831


◀: T-DNA Border

FIG.33C



derived from pGV 831 (DeBlaere et al, 1985)
Pssu: ~ 820 bp BglII – HindIII fragment of the pea ssu gene (Herrera-Estrella et al, 1984)
BamHI. HindIII: linker of puc 8
3'ocs: 706 bp pvuII fragment of ocs in HpaI site of pGV831
◀ : TDNA border


FIG.33D.



Pssu: ~ 970bp EcoRI-HindIII fragment of the pea ssu gene
Tp: sequence comprising transit peptide of the pea ssu gene (Van Den Brooke et al, 1985 Nature, 313, 358)
3'ocs: 706 bp PVUII fragment of ocs.
Km Tn 903: bacterial Km^R-gene
vector: pBR322


FIG. 33E.

Expression vector pGV874:



- "Pnos" fragment is the BclI-Bam HI fragment from pLGV2382 (Herrera-Estrella et al., EMBO J., 2, 987, 1983).
- "3' nos" fragment is the 687 bp TaQ1-Hind III fragments from the nos gene. (De Picker et al., J.M.A.G., 1, 561, 1982).
- "Sm-Sp" is the 2.3 Kb Hind III / Bam HI fragment from R702.
- The remaining of the vector is pBR322.

FIG.33F.



PLANT EXP. VECTOR

LEGEND


sequence from embl. databank
sequence from 1 to 6714
Sites from 1 to 6714
Maximal occurrence frequency: 2
All enzymes from ENZ. GOOD

1 L.B.

2 3' END T7

3 R.B.

FIG. 33G.




PLANT EXP. VECTOR

LEGEND

Sequence from embl. databank
Sequence from 1 to 8558
Sites from 1 to 8558
Maximal occurence frequency : 2
All enzymes from ENZ. GOOD


1. BETA-LACTAMASE
2. L. B.
3. 3' END T7
4. T-DNA
5. R. B.

FIG.33H.



Description of the isolation and characterization of the promoter fragment (Pssu PET) and 3'end SSU-PET See: C. Dean et al., EMBO J. Vol. 4, no. 12 (1985). Construction of pAGS 007 described in Figure 44.

FIG.331.



PLANT EXP. VECTOR

LEGEND


sequence from embl. databank
sequence from 1 to 7160
Sites from 1 to 7160
Maximal occurrence frequency: 2
All enzymes from ENZ. GOOD

1. L.B.
2. 3'END T7
3. T-DNA
4. R.B.

pGSJ 270 is identical to pGSJ 250 except that P35S-1 is replaced by P35S-2.

PLANT EXPR. VECTOR

FIG.33J



LEGEND

sequence from embl. databank
sequence from 1 to 9506
Sites from 1 to 9506
Maximal occurence frequency: 2
Some enzymes from ENZ. GOOD

1. BETA-LACTAMASE
2. L.B.
3. Pnos
4. 3' end T7
5. T-DNA
6. R.B.

pGSJ 280 is identical to pGSJ 260 except that P35S-1 is replaced by P35S-2.

FIG.34.



O.D.
0.2
0.1
Detection of Bt2 protein in engineered calli
4ng Bt2
1 2 3 4 5 6 7 8 9 10 11 12 13 14
callus nº
neg. control calli

FIG.35.




O.D.
4 ng B12
0.2
0.1
background level
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21
callus nº

FIG.36.


O.D.
1
0.5
0
1.7
4D6
4.8
10E3
AF6
6002
overnight
2h
Extract 1076
Extract SR1

PROTOCOL FOR PREPARATION OF PLANT EXTRACTS

-50 g callus material
-Homogenize at 0° C in 100 ml of the following buffer
  Na_2CO_3 pH 10 100mM
  PMFS 0.17 mg/ml
  EDTA 50mM
  DTT 10 mM

FIG. 37A.

-sonicate 2x3 min at 400 Watt on ice
-centrifuge 13,000 rpm ; 30 min

pellet I supernatant I

-Supernatant I
-Acid precipitation : bring pH down slowly to 4.5 by adding dropwise 1 M HCl
-Incubate for 30 min at 0°C
-Centrifuge 10,000 rpm, 30 min
-Wash on ice with cold distilled H_2O
-Resuspend pellet in small volume of buffer: Na_2CO_3 pH10 50mM DTT 5 mM PMFS 0.19 mg/ml
- Incubate for 1 h at 0°C, while regularly resuspending
- Centrifuge (in Eppendorf)

supernatant = fraction I

-Pellet I
-Resuspend in 100 ml of the following buffer (0°C):
  Na_2CO_3 pH 10 100mM
  DTT 10 mM
  PMFS 0.17 mg/ml
  EDTA 50 mM
  1% Triton x 100
-Sonicate 2x3 min at 400 Watt on ice
-Centrifuge 13,000 rpm 30 min

pellet II supernatant II

FIG. 37B.

Supernatant II:

- Acid precipitation: bring pH down slowly to 4.5 by adding dropwise 1 M HCl
- Incubate for 30 min at 0°C
- Centrifuge 10,000 rpm, 30 min
- Wash once with cold distilled H_2O
- Resuspend pellet in small volume of buffer: Na_2CO_3 pH 10 50 mM DTT 5 mM PMFS 0.17 mg/ml
- Incubate for 1 h at 0°C, while regularly resuspending
- Centrifuge (in Eppendorf)

→ supernatant = fraction II

Pellet II:

- Resuspend in 25 ml extraction buffer containing:
  2 % SDS
  Na_2CO_3 pH 10 100mM
  DTT 10 mM
  and agitate for 15 min

- Centrifuge 13,000 rpm, 30 min
- Supernatant ⟶ acetone precipitation:
  mix with 9 volumes of acetone 1/40 vol 1 MHCl
- Incubate overnight at −20°C
- Centrifuge 13,000 rpm, 20 min
- Resuspend pellet in small volume of buffer containing { 2% SDS; Na_2CO_3 pH10 100mM; DTT 10mM }
  and boil for 10 min
- Centrifuge ⟶ sup = fraction III

FIG. 38. Growth of M. sexta feeding on transformed tobacco leaves




FIG.39.



#L2 larvae
20
10
0
60
70
80
90
Plant nº
Dead larvae
C1 0
C2 0
C3 0
N20-3 0
N20-46 2
N20-38 1
N20-22 2
N20-47 0
N20-18 0
N20-30 1
XN20-31 0
20
10
0
60
70
80
90 t(h)
C1 0
C2 0
C3 0
C4 0
N20-35 0
N20-37 0
N20-7 0
N20-19 0
N20-7 0
N20-13 0
N20-1 0

FIG. 40.

35S PROMOTOR SEQUENCES AS PRESENT IN pGSJ250 AND pGSJ260

```
      6489      6499      6509      6519      6529      6539      6549      6559      6569      6579
          AGATCTCCTTTGCCCCGGAGATCACCATGGACGACTTTCTCTATCTCTACGATCTAGGAAGAAAGTTCGACGGAGAAGGTGACGATAC
          ^                   ^
          001-BglII           001-NcoI

      6589      6599      6609      6619      6629      6639      6649      6659      6669      6679
CATGTTCACGACCGATAATGAGAAGATTAGCCTCTTCAATTTCAGAAAGAATGCTGACCCACAGATGGTTAGAGAGGCCTACGCGGCAGGTCTCATCAAG
                                                                              ^ ^
                                                                              001-StuI
                                                                                001-BglI

      6689      6699      6709      6719      6729      6739      6749      6759      6769      6779
ACGATCTACCCGAGTAATAATCTCCAGGAGATCAAATACCTTCCCAAGAAGGTTAAAGATGCAGTCAAAAGATTCAGGACTAACTGCATGAAGAACACAG

      6789      6799      6809      6819      6829      6839      6849      6859      6869      6879
AGAAAGATATATTTCTCAAGATCAGAAGTACTATTCCAGTATGGACGATTCAAGGCTTGCTTCATAAACCAAGGCAAGTA  AGAGATTGGAGTCTCTAA
                          ^
                          001-ScaI

      6889      6899      6909      6919      6929      6939      6949      6959      6969      6979
GAAAGTAGTTCCTACTGAATCAAAGGCCATGGAGTCAAAAATTCAGATCGACGATCTAACAGAACTCGCCGTGAAGACTGGCGAACAGTTCATACAGAGT
                          ^
                          002-NcoI

      6989      6999      7009      7019      7029      7039      7049      7059      7069      7079
CTTTTACGACTCAATGACAAGAAGAAAATCTTCGTCAACATGGTGGAGCACGACACTCTCGTCTACTCCAAGAATATCAAAGATACAGTCTCAGAAGACC
                          ^
                          001-XmnI

      7089      7099      7109      7119      7129      7139      7149      7159      7169      7179
AAAGGGCTATTGAGACTTTTCAACAAAGGGTAATATCGGGAAACCTCCTCGGATTCCATTGCCCAGCTATCTGTCACTTCATCAAAAGGACAGTAGAAAA

      7189      7199      7209      7219      7229      7239      7249      7259      7269      7279
GGAAGGTGGCACCTACAAATGCCATCATTGCGATAAAGGAAAGGCTATCGTTCAAGATGCCTCTGCCGACAGTGGTCCCAAAGATGGACCCCCACCCACG

      7289      7299      7309      7319      7329      7339      7349      7359      7369      7379
AGGAGCATCGTGGAAAAAGAAGACGTTCCAACCACGTCTTCAAAGCAAGTGGATTGATGTGATATCTCCACTGACGTAAGGGATGACGCACAATCCCACT
                    ^                                       ^
                    002-XmnI                                001-EcoRV
                                                                                ClaI
                                                                                GAT ← P3   -2

      7389      7399      7409      7419      7429      7439      7449      7459      7469
ATCCTTCGCAAGACCCTTCCTCTATATAAGGAAGTTCATTTCATTTGGAGAGGACACGCTGAAATCACCAGTCTCTCTCTACAAATCTATC

NO MATCH FOR STRING                                                                  GGATCC ← P35S-1
                                                                                     BamHI
```

FIG. 41.



5'.. ATCG'ATG'GAC'... 3'
EcoRI. ClaI
BamHI. SalI. PstI. HindIII
PTR2'
PTR1'
ApR
pOP443
3200 bp
BamHI. SalI
PTR1
PTR2
NP
ApR
pAP2034
7700bp
3'ocs
St2/SpR
BamHI/SalI/Klenow pol./
Ligase/
pOP443 BSF
ClaI/HindIII
BamHI/Klenow pol./
ClaI/Linker/Ligase
ClaI/Ligase/
pAP 2034 C
ClaI/Hind III
Ligase
5'...ATCG'ATG'GAT'CCC...3'
ClaI
BamHI
BamHI ClaI
Eco RI
Pst I. Hind III
PTR2
PTR1
ApR
pGSH 50
St2/SpR

FIG.42. NcoI-BamHI-SacII-ApaI-XhoI-HindIII-BglII





FIG.43


pGSH 50
pGV1500
EcoRI / Klenow / HindIII
HpaI / HindIII
Ligase
pGSH150
(see figure 33F)
BamHI
pLBKm 13
BamHI / BglII
Ligase
SmaI
NruI
SalI
NcoI
ScaI
10,000
5
NPT II
NarI
1
6
7
8000
2000
NdeI
pGSH151
Sm-Sp-AD-TRANSF.
B12
4
2
3
6000
4000
HpaI
T-DNA
BamHI
BglII
LEGEND
sequence from embl. data-bank sequence from 1 to 10,035
Sites from 1 to 10,035
Maximal occurence frequency: 1
All enzymes from ENZ. GOOD
1 BETA-LACTAMASE
2 L.B.
3 PTR 1
4 PTR 2
5 3'- END T7
6 T-DNA
7 R.B.



SSU 301 type gene
R1 R1
RI
Nco I
R1
Cla1
Bgl II
Xho I
B
1Kb
5'
3'
subcloned the R1 fragmagment containing the 5' end of the gene into M13mp11
subcloned Bgl II-Bam fragment containing the 3' end of the gene into M13mp11
R1
R1
ATG
Bgl II
B
TAA
performed site-directed mutagenesis at the ATG to create an NcoI site
performed site-directed mutagenesis immediately 3' to the TAA codon to create a BglII site
wild type sequence
TCTAACTATGGCTTC
mutant sequence
TCTAACCATGGCTTC
wild type sequence
GGCTTCTAAGTTATATTAGGA
mutant sequence
GGCTTCTAAGATCTATTAGGA
reconstuct 5' flanking sequences (R1 partial) and mutagenize 5' Nco I site (NcoI partial and filling in)
R1
R1
Nco I
Bgl II
Xho I
B
joined mutagenized 5' and 3' flanking regions of SSU 301 together using a 600bp BglII+Nco I fragment of Tn5 as a linker
Hind Pst Sal Bam Sma
pUC 12 polylinker
RI
RI
Nco I Bgl II
Xho I
B
R I
To use: Cut with Nco I and Bgl II, clone in desired coding region making as perfect a fusion as possible. The complete fragment can be cloned into the AGS Agro bacterium binary vectors on a Bam HI, Sal/Xho or Sma/Xho fragment depending on sites in the coding region.

FIG.44.