



(86) Date de dépôt PCT/PCT Filing Date: 2005/02/18
(87) Date publication PCT/PCT Publication Date: 2005/09/09
(85) Entrée phase nationale/National Entry: 2006/08/02
(86) N° demande PCT/PCT Application No.: US 2005/005409
(87) N° publication PCT/PCT Publication No.: 2005/081905
(30) Priorité/Priority: 2004/02/20 (US60/546,339)

(51) Cl.Int./Int.Cl. *C12N 5/14* (2006.01),
A01H 5/06 (2006.01), *C12N 15/82* (2006.01),
C12N 15/87 (2006.01), *C12N 15/90* (2006.01)
(71) Demandeur/Applicant:
FRAUNHOFER USA INC., US
(72) Inventeurs/Inventors:
YUSIBOV, VIDADI, US;
SKARJINSKAIA, MARINA, US
(74) Agent: ROBIC

(54) Titre : SYSTEMES ET METHODES D'EXPRESSION CLONALE DANS DES PLANTES
(54) Title: SYSTEMS AND METHODS FOR CLONAL EXPRESSION IN PLANTS

(57) **Abrégé/Abstract:**

The present invention provides systems and methods for generating clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants and for expressing gene products in such cell lines and plants. According to certain of the inventive methods, a viral vector containing a polynucleotide of interest operably linked to a promoter is introduced into a plant or portion thereof. Plant material (e.g., leaf portions) containing the viral vector are used to generate clonal root lines, e.g., by contacting them with *A. rhizogenes* so as to form hairy roots. The hairy roots are derived from single ancestral cells and are therefore clonal. These clonal root lines contain the viral vector and express the polynucleotide of interest. Clonal root cell lines, clonal plant cell lines, and clonal plants are obtained from the clonal roots. According to certain other inventive methods, a viral vector containing a polynucleotide of interest operably linked to a promoter is introduced into cells of a plant cell line (e.g., pre-existing or newly derived) that is maintained in culture. Clonal plant cell lines that express the polynucleotide of interest are obtained, e.g., by successive rounds of enrichment until a population of cells exhibiting stable expression is obtained. Clonal plants whose cells contain the viral vector are obtained from the clonal plant cell lines using standard methods. The invention provides clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants generated using the inventive methods.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
9 September 2005 (09.09.2005)

(10) International Publication Number
WO 2005/081905 A3

(51) International Patent Classification:

C12N 5/14 (2006.01) *C12N 15/87* (2006.01)
A01H 5/06 (2006.01) *C12N 15/90* (2006.01)
C12N 15/82 (2006.01)

(21) International Application Number:

PCT/US2005/005409

(22) International Filing Date:

18 February 2005 (18.02.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/546,339 20 February 2004 (20.02.2004) US

(71) Applicant (for all designated States except US): **FRAUNHOFER USA INC.** [US/US]; 46025 Port Street, Plymouth, MI 48170 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **YUSIBOV, Vidadi** [RU/US]; 1808 Manor Road, Havertown, PA 19083 (US).
SKARJINSKAIA, Marina [UA/US]; 22 Fairway Road, Newark, DE 19711-5641 (US).

(74) Agent: **GERBER, Monica, R.**; Patent Group, Choate, Hall & Stewart LLP, Two International Place, Boston, MA 02110 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
22 June 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SYSTEMS AND METHODS FOR CLONAL EXPRESSION IN PLANTS

(57) Abstract: The present invention provides systems and methods for generating clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants and for expressing gene products in such cell lines and plants. According to certain of the inventive methods, a viral vector containing a polynucleotide of interest operably linked to a promoter is introduced into a plant or portion thereof. Plant material (e.g., leaf portions) containing the viral vector are used to generate clonal root lines, e.g., by contacting them with *A. rhizogenes* so as to form hairy roots. The hairy roots are derived from single ancestral cells and are therefore clonal. These clonal root lines contain the viral vector and express the polynucleotide of interest. Clonal root cell lines, clonal plant cell lines, and clonal plants are obtained from the clonal roots. According to certain other inventive methods, a viral vector containing a polynucleotide of interest operably linked to a promoter is introduced into cells of a plant cell line (e.g., pre-existing or newly derived) that is maintained in culture. Clonal plant cell lines that express the polynucleotide of interest are obtained, e.g., by successive rounds of enrichment until a population of cells exhibiting stable expression is obtained. Clonal plants whose cells contain the viral vector are obtained from the clonal plant cell lines using standard methods. The invention provides clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants generated using the inventive methods.

WO 2005/081905 A3

SYSTEMS AND METHODS FOR CLONAL EXPRESSION IN PLANTS

Cross-Reference to Related Application

[0001] This application claims priority to and the benefit of U.S. Provisional Patent Application No. 60/546,339 filed February 20, 2004, which is incorporated herein by reference.

Background of the Invention

[0002] Research to identify molecules with potential for preventative and therapeutic use (antibodies, enzymes, hormones and vaccine antigens) is of paramount importance to health and medicine. Historically, many of these molecules were recovered from human or animal sources. However, low quantities of target product in the source material coupled with immense costs, and more importantly, safety, have limited the availability of therapeutics and vaccines for prevention and treatment of many diseases around the world.

[0003] In the mid-1970s recombinant DNA technology revolutionized the process and made possible the production of target molecules predominantly in bacterial expression systems. Although prokaryotic expression systems continue to be a widely utilized method for recombinant protein production, this platform has its limitations because, for example, of the absence of eukaryotic posttranslational modifications and improper folding of many complex human proteins. During the last three decades many research laboratories have focused their interests on developing alternative systems for expressing recombinant proteins that could overcome the shortcomings of bacterial systems. Emerging out of these studies were animal and insect cell culture systems. Although a number of products such as monoclonal antibodies, vaccines and therapeutics have been produced using these systems, but the high cost of production combined with the requirement of highly sophisticated manufacturing facilities for each target protein motivated the search for different production systems.

[0004] In recent years, plants have been increasingly used as a host system for the expression of recombinant proteins. Such expression can be accomplished, for example, either by integrating the gene of interest into a plant genome, to create a

transgenic plant that stably expresses the desired protein, or by introducing the gene of interest into a plant vector that can be introduced into, and transiently maintained in, plant cells. Viral vector systems have proven to be particularly useful.

[0005] However, there remains a need for developing improved systems for expression of a molecule of interest in plants. For example, viruses may infect non-target plants, potentially posing significant environmental risks. Also, many available engineered plant viruses do not express inserted genes at desired levels, and/or in desired target plants or tissues. In addition, one disadvantage with various existing viral vector systems is that virus stability can be problematic. In general, there is a need in the art for plant expression systems that would allow for greater flexibility and control.

Summary of the Invention

[0006] The present invention encompasses the recognition that the availability of clonal expression systems for plants would offer a number of significant advantages. The invention provides methods and reagents for generating a variety of clonal entities derived from plants. These clonal entities include clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants. The invention further provides methods and reagents for expression of polynucleotide and polypeptide products in clonal cell lines derived from various plant tissues (e.g., roots, leaves), and in whole plants derived from single cells (clonal plants). The methods are based on the use of plant viral vectors of various types.

[0007] For example, in one aspect, the invention provides a method of obtaining a clonal root line that expresses a polynucleotide of interest comprising steps of: (i) introducing a viral vector that comprises a polynucleotide of interest into a plant or portion thereof; and (ii) generating one or more clonal root lines from the plant. The clonal root lines may be generated, for example, by infecting the plant or plant portion (e.g., a harvested piece of leaf) with an *Agrobacterium* (e.g., *A. rhizogenes*) that causes formation of hairy roots. Clonal root lines can be screened in various ways to identify lines that maintain the virus, lines that express the polynucleotide of interest at high levels, etc. The invention further provides clonal root lines, e.g., clonal root lines

produced according to the inventive methods and further encompasses methods of expressing polynucleotides and producing polypeptides of interest using the clonal root lines.

[0008] The invention further provides a method of generating a clonal root cell line that expresses a polynucleotide of interest comprising steps of: (i) generating a clonal root line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest; (ii) releasing individual cells from the clonal root line; and (iii) maintaining the cells under conditions suitable for root cell proliferation. The invention provides clonal root cell lines and methods of expressing polynucleotides and producing polypeptides using the clonal root cell lines.

[0009] In another aspect, the invention provides a method of generating a clonal plant cell line that expresses a polynucleotide of interest comprising steps of: (i) generating a clonal root line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest; (ii) releasing individual cells from the clonal root line; and (iii) maintaining the cells in culture under conditions appropriate for plant cell proliferation. The invention further provides a method of generating a clonal plant cell line that expresses a polynucleotide of interest comprising steps of: (i) introducing a viral vector that comprises a polynucleotide of interest into cells of a plant cell line maintained in culture; and (ii) enriching for cells that contain the viral vector. Enrichment may be performed, for example, by (i) removing a portion of the cells from the culture; (ii) diluting the removed cells so as to reduce the cell concentration; (iii) allowing the diluted cells to proliferate; and (iv) screening for cells that contain the viral vector. Clonal plant cell lines may be used for production of a polypeptide of interest.

[0010] The invention features a number of methods for generating clonal plants, cells of which contain a viral vector that comprises a polynucleotide of interest. For example, the invention provides a method of generating a clonal plant that expresses a polynucleotide of interest comprising steps of: (i) generating a clonal root line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest; (ii) releasing individual cells from the clonal root line; and (iii) maintaining the cells under conditions appropriate for formation of a plant. The invention further provides a method of generating a clonal plant that expresses a polynucleotide of interest

comprising steps of: (i) generating a clonal plant cell line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest; and (ii) maintaining the cells under conditions appropriate for formation of a plant. In general, the clonal plants can express any polynucleotide of interest. The clonal plants can be used for production of a polypeptide of interest.

[0011] This application refers to various patents, patent applications, and publications. The contents of all of these are incorporated herein by reference. In addition, the following publications are incorporated herein by reference: *Current Protocols in Molecular Biology*, *Current Protocols in Immunology*, *Current Protocols in Protein Science*, and *Current Protocols in Cell Biology*, all John Wiley & Sons, N.Y., edition as of July 2002; Sambrook, Russell, and Sambrook, *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 2001; Slater, A., Scott, N.W., and Fowler, M.R., *Plant Biotechnology*, Oxford University Press, 2003. In the event of a conflict between the instant specification and an incorporated reference the specification shall control.

Brief Description of the Drawings

[0012] *Figure 1* presents a schematic diagram of the engineering of a TMV based viral construct containing a polynucleotide of interest. The upper portion of the figure shows a diagram of the genomic organization of a TMV based virus construct, D4, and the lower portion shows the same construct following insertion of a polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as "target"). The 126/183 kDa protein is required for replication of the virus. The 30 kD protein is the movement protein (MP) that mediates cell-to-cell movement. Arrows indicate positions of the subgenomic promoters. Transcription of the inserted polynucleotide is under control of the TMV CP subgenomic promoter. The 3' portion of the construct includes TMV coat protein sequences and untranslated regions. These portions are optional.

[0013] *Figure 2* presents a schematic diagram of the engineering of a TMV based viral construct containing a polynucleotide of interest. The upper portion of the figure shows a schematic diagram of the genomic organization of a TMV based virus construct 30R. The lower portion shows the same construct following insertion of a

polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as “target”). The 126/183 kDa protein is required for replication of the virus. The 30 kD protein is the movement protein (MP) that mediates cell-to-cell movement. CP is the coat protein that mediates systemic spread. Arrows indicate positions of the subgenomic promoters. Transcription of the inserted polynucleotide is under control of an introduced promoter. CP expression is under control of the endogenous CP promoter. The 3' portion of the construct includes TMV coat protein sequences and untranslated regions. These portions are optional.

[0014] *Figure 3* presents a schematic diagram of the engineering of a TMV based viral construct containing a polynucleotide of interest and a gene encoding a marker for detection and/or selection. The upper portion of the figure shows the genomic organization of a TMV based virus construct, D4. The middle portion of the figure shows the same construct after insertion of a gene encoding a detectable marker (GFP) replacing the MP coding sequence. The lower portion of the figure shows the same construct following insertion of a polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as “target”). The 126/183 kDa protein is required for replication of the virus. Arrows indicate positions of the subgenomic promoters. Transcription of the detectable marker is under control of the MP subgenomic promoter. Transcription of the inserted polynucleotide of interest is under control of the TMV CP subgenomic promoter. The 3' portion of the construct includes TMV coat protein sequences and untranslated regions. These portions are optional.

[0015] *Figure 4* presents a schematic diagram of the engineering of a TMV based viral construct containing a polynucleotide of interest and a gene encoding a marker for detection and/or selection. The upper portion of the figure shows the genomic organization of a TMV based virus construct, D4. The middle portion of the figure shows the same construct after insertion of a gene encoding a selectable marker (gene encoding resistance to kanamycin) replacing the MP coding sequence. The lower portion of the figure shows the same construct following insertion of a polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as “target”). The 126/183 kDa protein is required for replication of the virus. Arrows indicate positions of the subgenomic promoters. Transcription of the selectable marker is under control of the TMV MP subgenomic promoter. Transcription of the inserted polynucleotide of

interest is under control of the TMV CP subgenomic promoter. The 3' portion of the construct includes TMV coat protein sequences and untranslated regions. These portions are optional.

[0016] *Figure 5* presents a schematic diagram of the engineering of AlMV based viral constructs containing a polynucleotide of interest either as an independent open reading frame or as a genetic fusion with AlMV CP coding sequences. The upper portion of the figure shows the genomic organization of RNA3 of AlMV, which includes genes encoding CP and MP as well as containing 5' and 3' UTRs and a subgenomic promoter. The left side of the figure shows a construct in which transcription of an mRNA containing separate open reading frames that encode a polypeptide of interest (indicated as "target") and the AlMV CP is under control of the AlMV subgenomic promoter. The right side of the figure shows a construct in which transcription of an mRNA containing a single open reading frame containing a polynucleotide of interest and CP coding sequences is under control of the AlMV CP subgenomic promoter. The open reading frame encodes a fusion protein in which a polypeptide of interest is fused to CP.

[0017] *Figures 6A-6E* illustrate steps in a method for generating clonal root lines for expression of a polynucleotide of interest (indicated as "target" in the figure). *Figure 6E* shows clonal root lines. *Figure 6F* shows clonal root lines at a higher level of magnification. *Figure 6G* shows GFP expression in a clonal root line, cells of which contain a viral vector that encodes GFP.

[0018] *Figures 7A – 7C* show Western blot analyses demonstrating GFP production in 3 clonal root lines derived from plant cells into which a viral vector whose genome contains a gene that encodes GFP under control of the TMV CP promoter was introduced. *Figure 7A* shows GFP expression in the clonal root lines after 30 days of propagation in culture (i.e., 30 days after separation of the root from the leaf from which it was derived). *Figure 7B* shows GFP expression in the clonal root lines after 60 days of propagation in culture (i.e., 60 days after separation of the root from the leaf from which it was derived). C- represents control lanes containing no protein. MWM represents molecular weight markers. GFP-R represents samples from clonal root lines. GFP-P represents GFP isolated from leaf tissue of a plant infected with the same

construct used for generation of the clonal root lines. *Figure 7C* is a control showing that the anti-GFP antibodies recognize commercially available GFP protein.

[0019] *Figures 8A and 8B* show photographs of clonal root lines producing hGH and GFP. *Figure 8A* shows a photograph of two clonal root lines taken under normal light conditions. The plate on the left shows a clonal root line derived from a plant cell into which a viral vector whose genome contains a gene that encodes human growth hormone (hGH) under control of the TMV CP promoter was introduced. The plate on the right shows a clonal root line derived from a plant cell into which a viral vector whose genome contains a gene that encodes green fluorescent protein (GFP) under control of the TMV CP promoter was introduced. *Figure 8B* shows a photograph of the same clonal root lines as shown in *Figure 8A* taken under UV light, demonstrating expression of GFP.

[0020] *Figure 9* shows a Western blot analysis to screen clonal root lines each derived from individual plant cells that were infected with a viral vector whose genome contains a gene that encodes human growth hormone (hGH) under control of the TMV CP promoter. Root lines were screened 30 days after separation of the root from the leaf from which it was derived. Root lines demonstrating high levels of expression are indicated with arrows. C- represents control lanes containing no protein. MWM represents molecular weight markers. hGH represents recombinant human growth hormone.

[0021] *Figure 10* shows a Western blot analysis demonstrating hGH production in selected clonal root lines derived from plant cells into which a viral vector whose genome contains a gene that encodes hGH under control of the TMV CP promoter was introduced. The analysis was performed following 10 subculturings after separation of the roots from the leaves from which they were derived. C- represents a control lane containing no protein. MWM represents molecular weight markers. hGH represents recombinant human growth hormone.

[0022] *Figures 11A and 11B* show Western blot analysis to screen clonal root lines each derived from individual plant cells that were infected with a viral vector whose genome contains a gene that encodes human growth hormone (GCSF) under control of the TMV CP promoter. Root lines were screened 30 days after separation of the root from the leaf from which it was derived. Root lines demonstrating high levels of

expression are indicated with arrows. C- represents control lanes containing no protein. MWM represents molecular weight markers. GCSF represents recombinant human granulocyte colony stimulating factor.

[0023] *Figure 12* shows a Western blot analysis demonstrating GCSF production in selected clonal root lines derived from plant cells into which a viral vector whose genome contains a gene that encodes GCSF under control of the TMV CP promoter was introduced. The analysis was performed following 10 subculturings after separation of the roots from the leaves from which they were derived. C- represents a control lane containing no protein. MWM represents molecular weight markers. GCSF represents recombinant human granulocyte colony stimulating factor.

[0024] *Figure 13* illustrates steps in a method for generating clonal plant cell lines for expression of a polynucleotide of interest and identifying cell lines that display expression. *Figure 13A* shows a viral vector in which a polynucleotide of interest is inserted under control of the TMV CP promoter. *Figure 13B* shows a protoplast suspension into which the vector was introduced (left plate) or a control protoplast suspension into which a vector lacking a GFP-encoding polynucleotide was introduced (right plate). The picture was taken under UV light and shows expression of GFP in the protoplasts containing the GFP-encoding expression vector. *Figure 13C* shows a protoplast suspension into which the GFP-encoding vector was introduced. The photo was taken under UV light. The inset (*Figure 13D*) shows a higher magnification of GFP-expressing cells, also taken under UV light. *Figure 13E* is a photograph showing enrichment for plant cell lines that express GFP. The photo was taken under normal light. of individual plant cell lines derived from the protoplast suspension shown in *Figure 13C*. *Figure 13F* is a photograph of the same plates as shown in *Figure 13C*, taken under UV light. Cultures enriched for cell lines expressing GFP are evident as green fluorescing spots.

[0025] *Figure 14* shows Western blot analyses demonstrating GCSF production in a clonal plant cell line derived from a plant cell into which a viral vector whose genome contains a gene that encodes GCSF under control of the TMV CP promoter was introduced. *Figure 14A* shows a Western blot performed 48 hours after introduction of the vector. *Figure 14B* shows a Western blot performed using the same cell lines as shown in *Figure 14A* performed 57 days after inoculation. GCSF-COM indicates a lane

in which recombinant GCSF protein was loaded as a positive control. MWM indicates molecular weight markers. C- indicates a lane in which plant extract made from plants not expressing GCSF was loaded.

[0026] *Figure 15* shows GFP production in plant cell lines derived from plant cells into which a viral vector whose genome contains a gene that encodes GFP under control of the TMV CP promoter was introduced. *Figure 15A* shows enrichment for plant cell lines that express GFP. *Figure 15B* shows a callus derived from a clonal plant cell line into which a similar viral vector, not encoding GFP, was introduced. The photographs were taken 3 months after the vector was introduced into the cells from which the clones in *Figure 15A* were derived. Both photographs were taken under UV light.

[0027] *Figure 16A* shows a clonal plant that was obtained from a clonal root line derived from a plant cell into which a viral vector encoding hGH was introduced. *Figure 16B* shows lesion formation in a sensitive host plant that was inoculated with a small leaf sample from the clonal plant, indicating that the clonal plant regenerated from the clonal root line maintains active viral replication. To test if the plant maintains virus replication a small leaf sample was used to inoculate a tobacco variety that is a host for formation of local lesions. Formation of lesions within 2 days of inoculation (see arrows) indicates that the clonal plant line regenerated from a clonal root line maintains active virus replication.

[0028] *Figure 17* presents a schematic representation of certain families of viruses that infect plants.

[0029] *Figure 18* shows representative examples of tobamovirus genomes.

Definitions

[0030] *Approximately:* "Approximately" in reference to a number includes numbers that fall within a range of 5% in either direction (greater than or less than) the number unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value). Where ranges are stated, the endpoints are included within the range unless otherwise stated or otherwise evident from the context.

[0031] *Clonal*: For the purpose of the present invention, the term clonal as applied, e.g., to a plant or plant tissue such as a root, leaf, stem, etc., means that the plant or plant tissue was derived from a single ancestral cell. In general, the cells or a clonal plant or plant tissue will be genetically identical with the exception of somatic mutations or other genetic alterations that may arise in descendant cells (e.g., through either natural or artificial introduction of a new gene into a descendant cell, telomere shortening, etc.). Typically the genome of the cells will be at least 95% identical, at least 98% identical, at least 99% identical, at least 99.5% identical, at least 99.9% identical.

[0032] *Gene*: For the purposes of the present invention, the term gene has its meaning as understood in the art. In general, a gene is taken to include gene regulatory sequences (e.g., promoters, enhancers, etc.) and/or intron sequences, in addition to coding sequences (open reading frames). It will further be appreciated that the definition of gene can include nucleic acids that do not encode proteins but rather provide templates for transcription of functional RNA molecules such as tRNAs, rRNAs, microRNAs (miRNAs), short hairpin RNAs (shRNAs), short interfering RNAs, (siRNAs), etc. For the purpose of clarity we note that, as used in the present application, the term “gene” generally refers to a nucleic acid that includes a portion that encodes a protein; the term may optionally encompass regulatory sequences such as promoters, enhancers, terminators, etc. This definition is not intended to exclude application of the term “gene” to non-protein coding expression units but rather to clarify that, in most cases, the term as used in this document refers to a protein coding nucleic acid.

[0033] *Gene product or expression product*: A gene product or expression product is, in general, an RNA transcribed from a gene or polynucleotide, or a polypeptide encoded by an RNA transcribed from the gene or polynucleotide. Expression of a gene or a polynucleotide refers to (i) transcription of RNA from the gene or polynucleotide; (ii) translation of RNA transcribed from the gene or polynucleotide, or both (i) and (ii). Other steps such as processing, translocation, etc., may also take place in the course of expression or thereafter.

[0034] *Isolated*: As used herein, the term “isolated” refers to a compound or entity that is 1) separated from at least some of the components with which it is normally

associated (e.g., purified); 2) synthesized *in vitro*; and/or 3) produced or prepared by a process that involves the hand of man.

[0035] *Naturally*: The term “naturally” or “naturally-occurring”, as used herein, refers to processes, events, or things that occur in their relevant form in nature. By contrast, “not-naturally-occurring”, “artificial”, or “synthetic” refers to processes, events, or things whose existence or form involves the hand of man.

[0036] *Operably linked*: As used herein, operably linked refers to a relationship between two nucleic acids or two polypeptides wherein the expression of one of the nucleic acids or polypeptides is controlled by, regulated by, modulated by, etc., the other nucleic acid or polypeptide. For example, the transcription of a nucleic acid sequence is directed by an operably linked promoter sequence; post-transcriptional processing of a nucleic acid is directed by an operably linked processing sequence; the translation of a nucleic acid sequence is directed by an operably linked translational regulatory sequence; the transport or localization of a nucleic acid or polypeptide is directed by an operably linked transport or localization sequence; and the post-translational processing of a polypeptide is directed by an operably linked processing sequence. Preferably a nucleic acid or polypeptide sequence that is operably linked to a second nucleic acid or polypeptide sequence is covalently linked, either directly or indirectly, to such a sequence, although any effective three-dimensional association is acceptable. It is noted that a single nucleic acid or polypeptide sequence can be operably linked to multiple other sequences. For example, a single promoter can direct transcription of multiple RNA species.

[0037] *Percent (%) identity*: In reference to polynucleotides, “percent (%) identity” is defined as the percentage of nucleotide residues in a polynucleotide sequence that are identical with the nucleotide residues in the specific nucleic acid sequence with which comparison is being made, after aligning the sequences and introducing gaps, as needed, to achieve the maximum percent sequence identity. In reference to polypeptides, “percent (%) identity” is defined as the percentage of amino acid residues in a polypeptide sequence that are identical with the amino acid residues in the specific polypeptide sequence with which comparison is being made, after aligning the sequences and introducing gaps, as needed, to achieve the maximum percent sequence identity.

[0038] Alignment can be performed in various ways known to those of skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. US Publication No. 20030211568 describes a number of suitable methods.

[0039] *Polynucleotide of interest*: As used herein, the term “polynucleotide of interest” refers to any target nucleic acid sequence to be expressed in plant cells, as described herein. In many embodiments, the polynucleotide of interest will be a protein-coding polynucleotide (in which case the encoded polypeptide may be referred to as a polypeptide or protein of interest) but may also be a sequence that provides a template for transcription of a structural RNA or an active RNA such as a ribozyme, interfering RNA strand, etc. Often, the polynucleotide will be a gene that is not expressed in nature in the relevant type of plant cell, or is not expressed at the level that the polynucleotide is expressed when expression is achieved by intervention of the hand of man, as described herein. In certain embodiments of the invention, the polynucleotide comprises gene sequences that are not naturally found in the relevant plant cell at all; often including gene sequences that are naturally found in other cell types or organisms. Alternatively or additionally, a polynucleotide of interest is one that is not naturally associated with the vector sequences with which it is associated according to the present invention. The word polynucleotide is used interchangeably with “nucleic acid” or “nucleic acid molecule” herein.

[0040] *Purified*: As used herein, “purified” means separated from one or more compounds or entities, e.g., one or more compounds or entities with which it is naturally found. A compound or entity may be *partially purified*, *substantially purified*, or *pure*, where it is *pure* when it is removed from substantially all other compounds or entities, i.e., is preferably at least about 90%, more preferably at least about 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or greater than 99% pure. In the context of a preparation of a nucleic acid molecule, a preparation may be considered substantially pure if the nucleic acid represents at least 50% of all nucleic acid molecules in the preparation, preferably at least 75%, yet more preferably at least 90%, or greater, as listed above, on a molecule per molecule basis, a w/w basis, or both.

In the context of a preparation of a polypeptide, a preparation may be considered substantially pure if the polypeptide represents at least 50% of all polypeptides in the preparation, preferably at least 75%, yet more preferably at least 90%, or greater, as listed above, on a molecule per molecule basis, a w/w basis, or both. A partially or substantially purified nucleic acid or polypeptide may be removed from at least 50%, at least 60%, at least 70%, or at least 80%, at least 90%, etc., of the material with which it is naturally found, e.g., cellular material such as other cellular proteins and/or nucleic acids.

[0041] *Recombinant*: A “recombinant” molecule refers to a molecule that has been altered by the hand of man or that is derived from (e.g., copied from) such a molecule. A recombinant polynucleotide typically contains sequences that are not found joined together in nature and/or that differ from a naturally occurring sequence. An amplified or assembled recombinant polynucleotide may be included in a suitable vector, and the vector can be used to transform a suitable cell, which may be referred to as a “recombinant cell”. The nucleotide may then be expressed in the recombinant cell to produce, e.g., a “recombinant polypeptide”. A recombinant polynucleotide may serve a non-coding function (e.g., promoter, origin of replication, ribosome-binding site, etc.) as well. A recombinant nucleic acid, e.g., a recombinant viral nucleic acid may be a viral nucleic acid in which one or more sequences present in the naturally occurring form has been deleted or replaced by a different sequence or into which a non-native sequence has been inserted. A “recombinant polypeptide” typically contains sequences that are not found joined together in nature and/or that differ from a naturally occurring sequence. One example of a recombinant polypeptide is a fusion protein, e.g., a protein containing two or more different proteins or peptides (which may be natural or synthetic and may be portions of a naturally occurring or synthetic polypeptide). A recombinant polynucleotide that encodes a fusion protein may be created by removing the stop codon from the polynucleotide that encodes the first protein or peptide and appending a polynucleotide that encodes the second protein or peptide in frame, so that the resulting recombinant polynucleotide encodes a single recombinant polypeptide comprising the two proteins or peptides.

[0042] The term “regulatory element” or “regulatory sequence” in reference to a nucleic acid is generally used herein to describe a portion of nucleic acid that directs or

increases one or more steps in the expression (particularly transcription, but in some cases other events such as splicing or other processing) of nucleic acid sequence(s) with which it is operatively linked. The term includes promoters and can also refer to enhancers and other transcriptional control elements. Promoters are regions of nucleic acid that include a site to which RNA polymerase binds before initiating transcription and that are typically necessary for even basal levels of transcription to occur.

Generally such elements comprise a TATA box. Enhancers are regions of nucleic acid that encompass binding sites for protein(s) that elevate transcriptional activity of a nearby or distantly located promoter, typically above some basal level of expression that would exist in the absence of the enhancer. In some embodiments of the invention, regulatory sequences may direct constitutive expression of a nucleotide sequence (e.g., expression in most or all cell types under typical physiological conditions in culture or in an organism); in other embodiments, regulatory sequences may direct cell or tissue-specific and/or inducible expression. For example, expression may be induced by the presence or addition of an inducing agent such as a hormone or other small molecule, by an increase in temperature, etc. Regulatory elements may also inhibit or decrease expression of an operatively linked nucleic acid.

[0043] In general, the level of expression may be determined using standard techniques for measuring mRNA or protein. Such methods include Northern blotting, *in situ* hybridization, RT-PCR, sequencing, immunological methods such as immunoblotting, immunodetection, or fluorescence detection following staining with fluorescently labeled antibodies, oligonucleotide or cDNA microarray or membrane array, protein array analysis, mass spectrometry, *etc.* A convenient way to determine expression level is to place a nucleic acid that encodes a readily detectable marker (e.g., a fluorescent or luminescent protein such as green fluorescent protein or luciferase, an enzyme such as alkaline phosphatase, etc.) in operable association with the regulatory element in an expression vector, introduce the vector into a cell type of interest or into an organism, maintain the cell or organism for a period of time, and then measure expression of the readily detectable marker, taking advantage of whatever property renders it readily detectable (e.g., fluorescence, luminescence, alteration of optical property of a substrate, etc.). Comparing expression in the absence and presence of the

regulatory element indicates the degree to which the regulatory element affects expression of an operatively linked sequence.

[0044] *Self-replicate*: As used herein, “self-replicate” refers to the ability of a vector to copy itself inside a host cell. A vector that can “self-replicate” carries sufficient information in its own genetic elements that it does not rely on other genetic elements (e.g., those utilized by the host cell to replicate its own genome) for its replication. In general, a vector that can self-replicate is one that includes at least one replicase gene such as an RNA polymerase and possibly additional replicase genes such as a helicase, methyltransferase, etc. In certain instances additional sequences, present in *cis* (i.e., as part of the vector sequence) are required or can facilitate self-replication. It will be understood that a self-replicating vector will typically utilize host cell components such as nucleotides, amino acids, etc., and may be dependent on certain functions and/or enzymes of the host cell that supply such components.

[0045] *Vector*: “Vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked and can include a plasmid, cosmid or viral vector. The vector may be capable of autonomous replication. Alternatively or additionally, a vector may provide one or more components necessary or sufficient for self-replication, or for replication or integration of another piece of nucleic acid. Vectors are typically nucleic acids, and may comprise DNA and/or RNA. Preferred vectors are maintained extrachromosomally.

[0046] *Viral nucleic acid*: The term “viral nucleic acid” refers to the genome of a virus, or a portion thereof (or, in the case of viruses whose genome comprises multiple segments, any of the segments or a portion of such a segment). The term encompasses both RNA and DNA forms of such nucleic acids and molecules having complementary sequences. DNA molecules identical to or complementary to viral RNA nucleic acids are considered viral nucleic acids, and RNA molecules identical to or complementary to viral DNA nucleic acids are considered viral nucleic acids, it being understood that DNA and RNA will contain T and U, respectively, at corresponding positions.

[0047] A viral nucleic acid may include one or more portions of non-viral origin (e.g., part or all of a naturally occurring gene, an entirely artificial sequence, or a combination of naturally occurring and artificial sequences) and may include portion(s) from multiple different virus types.

[0048] *Viral replicon*: The term “viral replicon” refers to a nucleic acid molecule comprising a portion or portions (*cis* sequences) sufficient for replication of the nucleic acid by viral replicase genes. Typically such sequences include a recognition site for a viral polymerase, e.g., a viral RNA polymerase in the case of viral replicons based on RNA viruses.

Detailed Description of Certain Embodiments of the Invention

[0049] *I. Clonal Plant and Plant Tissue Expression Systems*

[0050] As noted above, the present invention provides systems for expressing a polynucleotide or polynucleotides of interest in clonal root lines, clonal root cell lines, clonal plant cell lines (e.g., cell lines derived from leaf, stem, etc.), and in clonal plants. The polynucleotide of interest is introduced into an ancestral plant cell using a plant viral vector whose genome includes the polynucleotide of interest operably linked to (i.e., under control of) a promoter. A clonal root line or clonal plant cell line is established from the cell containing the virus according to any of several techniques further described below. The plant virus vector or portions thereof can be introduced into the plant cell by infection, by inoculation with a viral transcript or infectious cDNA clone, by electroporation, by T-DNA mediated gene transfer, etc.

[0051] The following sections describe plant viruses, plant viral vector, and methods for creating plant viral vectors for use in the present invention. The inventive methods for generating clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants that express a polynucleotide of interest are then described. A “root line” is distinguished from a “root cell line” in that a root line produces actual rootlike structures or roots while a root cell line consists of root cells that do not form rootlike structures. The use of the term “line” is intended to indicate that cells of the line can proliferate and pass genetic information on to progeny cells. Cells of a cell line typically proliferate in culture without being part of an organized structure such as those found in an intact plant. The use of the term “root line” is intended to indicate that cells in the root structure can proliferate without being part of a complete plant. It is noted that the term “plant cell” encompasses root cells. However, to distinguish the inventive methods for generating root lines and root cell lines from those used to directly generate plant cell lines from non-root tissue (as opposed to generating clonal

plant cell lines from clonal root lines or clonal plants derived from clonal root lines), the terms “plant cell” and “plant cell line” as used herein generally refer to cells and cell lines that consist of non-root plant tissue. The plant cells can be, for example, leaf, stem, shoot, flower part, etc. It is noted that seeds can be derived from the clonal plants generated as derived herein. Such seeds will also contain the viral vector as will plants obtained from such seeds. Methods for obtaining seed stocks are well known in the art. See, e.g., U.S.S.N. 10/294,314.

[0052] *A. Plant Viruses and Plant Viral Vectors*

[0053] A wide variety of viruses are known that infect various plant species, and can be employed for polynucleotide expression according to the present invention. Figure 17 presents a schematic representation of certain families of viruses that infect plants. Appendix A provides a representative list of plant virus families, based on the type of nucleic acid (e.g., dsDNA, ssDNA, ssRNA, dsRNA, or unassigned) that makes up the viral genome. Additional information can be found, for example, in *The Classification and Nomenclature of Viruses*”, Sixth Report of the International Committee on Taxonomy of Viruses” (Ed. Murphy et al.), Springer Verlag: New York, 1995, the entire contents of which are incorporated herein by reference (see also, Grierson et al., *Plant Molecular Biology*, Blackie, London, pp. 126-146, 1984; Gluzman et al., *Communications in Molecular Biology: Viral Vectors*, Cold Spring Harbor Laboratory, NY, pp. 172-189, 1988; Mathew, *Plant Viruses Online* (<http://image.fs.uidaho.edu/vide/>)).

[0054] In nature, in order to enter and infect a plant cell, plant viruses need to cross the cell wall, in addition to protective layers of waxes and pectins. Most or all plant viruses are thought to rely on mechanical breach of the cell wall, rather than on cell-wall-surface receptors, to enter a cell. Such a breach can be caused, for example, by physical damage to the cell, by an organism such as a bacterium, a fungus, a nematode, an insect, or a mite that can deliver the virus. In the laboratory, viruses are typically administered to plant cells simply by rubbing the virus on the plant.

[0055] Some plant viruses have segmented genomes, in which two or more physically separate pieces of nucleic acid together make up the plant genome. For example, many RNA plant virus genomes can be classified as mono-, bi-, or tri-partite, i.e., they may consist of 1, 2, or 3 nucleic acids respectively. In some cases, these

separate pieces are packaged together in the same viral capsid; in others (i.e., those with multipartite genomes), each genome segment is packaged into its own viral particle. Infection can typically be accomplished by delivery either of plant viral nucleic acid (e.g., RNA) or capsid containing such nucleic acid.

[0056] Once the virus has entered (infected) a cell, it typically replicates within the infected cell and then spreads locally (i.e., from cell to cell within leaves that were infected initially). Following local spread, the virus may move into uninfected leaves, e.g., upper leaves of the plant, which is referred to as systemic infection or systemic spread. In general, cell-to-cell spread of many plant viruses requires a functional movement protein (which allows movement of viral transcripts) while systemic spread requires a functional coat protein (and, generally, also a functional movement protein), which allows the formation of viral particles.

[0057] In addition to functional movement and coat protein encoding components, the viral genome may contain additional components that are required for local (e.g., cell-to-cell) or long distance (e.g., systemic) spread or facilitate such spread. These *cis*-acting components may be either coding or noncoding components. For example, they may correspond to portions of a 3' untranslated region (UTR, also referred to as NTR) of a viral transcript (i.e., they may provide a template for transcription of a 3' untranslated region of a viral transcript). Thus important viral components can be either coding or noncoding regions of a viral genome and include a variety of regulatory regions. Such regions may function in replication and/or processing or expression of mRNA. By "functional protein encoding component" is meant a polynucleotide comprising a coding portion that encodes a functionally active protein, operably linked to sufficient regulatory elements such as a promoter, so that expression is achieved.

[0058] In order to successfully establish either a local (intraleaf) or systemic infection a virus must be able to replicate. Many viruses contain genes encoding one or more proteins that participate in the replication process (referred to herein as replication proteins or replicase proteins). For example, many RNA plant viruses encode an RNA polymerase. Additional proteins may also be required, e.g., helicase or methyltransferase protein(s). The viral genome or segment may contain various sequence components, e.g., *cis*-acting sequences, in addition to functional genes

encoding replication proteins, which are also required for or facilitate replication. Viral genomes or segments may also contain *cis*-acting sequences that contribute to high levels of transcript and/or expression. It is noted that nucleic acids encoding various viral proteins, e.g., replicase proteins, movement protein, coat protein, may be present within different viral nucleic acid molecules, which may complement each other in *trans*. (See, e.g., WO 00/25574 and co-pending U.S. National Application Ser. No. 10/770,600, entitled "SYSTEM FOR EXPRESSION OF GENES IN PLANTS", filed February 3, 2004. Thus in certain embodiments of the invention rather than delivering a single viral vector to a plant cell, multiple different vectors are delivered which, together, allow for replication (and, optionally cell-to-cell and/or long distance movement) of the viral vector(s). Some or all of the proteins may be encoded by the genome of transgenic plants.

[0059] Viral vectors based on any virus that infects plants may be used to generate a clonal root line, clonal plant cell line or clonal plant that expresses a polynucleotide of interest in accordance with the present invention. Particularly preferred viruses are ssRNA viruses, most desirably with a (+)-stranded genome. Techniques and reagents for manipulating the genetic material present in such viruses are well known in the art. Typically, for example, a DNA copy of the viral genome is prepared and cloned into a microbial vector, particularly a bacterial vector. Certain ssDNA viruses, including particularly geminiviruses, may also be used. It will be appreciated that in general plant viral vectors and viral nucleic acids such as viral genomes may exist in RNA or DNA form. In addition, where reference is made to a feature such as a genome or portion thereof of an RNA virus, which is present within a DNA vector, it is to be understood that the feature is present as the DNA copy of the RNA form.

[0060] Preferred vectors are based on viruses such as members of the *Bromoviridae* (e.g., bromoviruses, alfamoviruses, ilarviruses) and *Tobamoviridae*. Certain preferred virus species include, for example, Alfalfa Mosaic Virus (AIMV), Apple Chlorotic Leaf Spot Virus, Apple Stem Grooving Virus, Barley Stripe Mosaic Virus, Barley Yellow Dwarf Virus, Beet Yellow Virus, Broad Bean Mottle Virus, Broad Bean Wilt Virus, Brome Mosaic Virus (BMV), Carnation Latent Virus, Carnation Mottle Virus, Carnation Ringspot Virus, Carrot Mottle Virus, Cassava Latent Virus (CLV), Cowpea Chlorotic Mottle Virus, Cowpea Mosaic Virus (CPMV), Cucumber Green Mottle

Mosaic Virus, Cucumber Mosaic Virus, Lettuce Infectious Yellow Virus, Maize Chlorotic Mottle Virus, Maize Rayado Fino Virus, Maize Streak Virus (MSV), Parsnip Yellow Fleck Virus, Pea Enation Mosaic Virus, Potato Virus X, Potato Virus Y, Raspberry Bushy Dwarf Virus, Rice Necrosis Virus (RNV), Rice Stripe Virus, Rice Tungro Spherical Virus, Ryegrass Mosaic Virus, Soil-borne Wheat Mosaic Virus, Southern Bean Mosaic Virus, Tobacco Etch Virus (TEV), Tobacco Mosaic Virus (TMV), Tobacco Necrosis Virus, Tobacco Rattle Virus, Tobacco Ring Spot Virus, Tomato Bushy Stunt Virus, Tomato Golden Mosaic Virus (TGMV), and Turnip Yellow Mosaic Virus (TYMV).

[0061] In certain embodiments of the invention a TMV-based viral vector (viral nucleic acid) is used. TMV is the type member of the tobamovirus group. Tobamoviruses have single-(+)-stranded RNA genomes, and produce rod-shaped virions consisting of the RNA genome and coat protein (CP) polypeptides. Tobamovirus genomes encode 4-5 polypeptides. Two of the polypeptides are translated from the same 5'-proximal initiation codon and function in viral replication. These polypeptides include an RNA-dependent RNA polymerase,. In addition, polypeptides having methyltransferase and RNA helicase activity are typically encoded. The other encoded proteins typically include a movement protein and the coat protein, each of which is translated from a separate subgenomic RNA. Representative examples of tobamovirus genomes are depicted in Figure 18.

Tobamoviruses other than TMV can be used in various embodiments of the invention.

[0062] The TMV genome is 6395 nucleotides long and is encapsidated with a 17.5 kD CP, which produces 300 nm-long rods. In addition to CP, TMV has three nonstructural proteins: 183 and 126 kD proteins are translated from genomic RNA and are required for viral replication. The 30 kD movement protein provides for the transfer of viral RNA from cell-to-cell. Plant species susceptible to infection with TMV include *Beta vulgaris*, *Capsicum frutescens*, *Chenopodium amaranticolor*, *Chenopodium hybridum*, *Chenopodium quinoa*, *Cucumis melo*, *Cucumis sativus*, *Cucurbita pepo*, *Datura stramonium*, *Lactuca sativa*, *Lucopersicon esculentum*, *Lycopersicon pimpinellifolium*, *Nicotiana benthamiana*, *Nicotiana bigelovii*, *Nicotiana clevelandii*, *Nicotiana debneyi*, *Nicotiana glutinosa*, *Nicotiana rustica*, *Nicotiana*

sylvestris, *Nicotiana tabacum*, *Papaver nudicaule*, *Phaseolus vulgaris*, *Physalis floridana*, *Physalis peruviana*, and *Solanum tuberosum*.

[0063] In various other embodiments of the invention an AlMV-based viral vector (viral nucleic acid) is used. AlMV is an *Alfamovirus*, closely related to the *Ilarvirus* group and is a member of the *Bromoviridae* family. The genome of AlMV consists of three positive-sense RNAs (RNAs 1-3). RNAs 1 and 2 encode replicase proteins P1 and P2, respectively; RNA3 encodes the cell-to-cell movement protein P3. A subgenomic RNA, RNA4, is synthesized from RNA3. This subgenomic RNA4 encodes the viral coat protein (CP). CP participates in viral genome activation to initiate infection, RNA replication, viral assembly, viral RNA stability, long-distance movement of viral RNA, and symptom formation. AlMV depends on a functional P3 protein for cell-to-cell movement, and requires the CP protein throughout infection. Depending on the size of the CP-encapsidated viral RNA, virions of AlMV can vary significantly in size (e.g., 30- to 60-nm in length and 18 nm in diameter) and form (e.g., spherical, ellipsoidal, or bacilliform).

[0064] The host range of AlMV is remarkably wide and includes the agriculturally valuable crops alfalfa (*Medicago sativa*), tomato (*Lycopersicon esculentum*), lettuce (*Lactuca sativa*), common bean (*Phaseolus vulgaris*), potato (*Solanum tuberosum*), white clover (*Trifolium repens*) and soybean (*Glycine max*). Particular susceptible host species include, for example, *Abelmoschus esculentus*, *Ageratum conyzoides*, *Amaranthus caudatus*, *Amaranthus retroflexus*, *Antirrhinum majus*, *Apium graveolens*, *Apium graveolens* var. *rapaceum*, *Arachis hypogaea*, *Astragalus glycyphyllos*, *Beta vulgaris*, *Brassica campestris* ssp. *rapa*, *Calendula officinalis*, *Capsicum annuum*, *Capsicum frutescens*, *Caryopteris incana*, *Catharanthus roseus*, *Celosia argentea*, *Cheiranthus cheiri*, *Chenopodium album*, *Chenopodium amaranticolor*, *Chenopodium murale*, *Chenopodium quinoa*, *Cicer arietinum*, *Cichorium endiva*, *Coriandrum sativum*, *Crotalaria spectabilis*, *Cucumis melo*, *Cucumis sativus*, *Cucurbita pepo*, *Cyamopsis tetragonoloba*, *Daucus carota* (var. *sativa*), *Dianthus barbatus*, *Dianthus caryophyllus*, *Emilia sagittata*, *Fagopyrum esculentum*, *Gomphrena globosa*, *Helianthus annuus*, *Lablab purpureus*, *Lathyrus odoratus*, *Lens culinaris*, *Linum usitatissimum*, *Lupinus albus*, *Macroptilium lathyroides*, *Malva parviflora*, *Matthiola incana*, *Medicago hispida*, *Melilotus albus*, *Nicotiana bigelovii*, *Nicotiana clevelandii*,

Nicotiana debneyi, *Nicotiana glutinosa*, *Nicotiana megalosiphon*, *Nicotiana rustica*, *Nicotiana sylvestris*, *Nicotiana tabacum*, *Ocimum basilicum*, *Petunia* × *hybrida*, *Phaseolus lunatus*, *Philadelphus*, *Physalis floridana*, *Physalis peruviana*, *Phytolacca americana*, *Pisum sativum*, *Solanum demissum*, *Solanum melongena*, *Solanum nigrum*, *Solanum nodiflorum*, *Solanum rostratum*, *Sonchus oleraceus*, *Spinacia oleracea*, *Stellaria media*, *Tetragonia tetragonoides*, *Trifolium dubium*, *Trifolium hybridum*, *Trifolium incarnatum*, *Trifolium pratense*, *Trifolium subterraneum*, *Tropaeolum majus*, *Viburnum opulus*, *Vicia faba*, *Vigna radiata*, *Vigna unguiculata*, *Vigna unguiculata* ssp. *sesquipedalis*, and

Zinnia elegans. While ALMV is a preferred viral vector, other alfamoviruses can also be used in various embodiments of the invention. Related viruses, such as ilarviruses can also be used.

[0065] *B. Creation of Plant Viral Expression Vectors*

[0066] Elements of these plant viruses are genetically engineered according to known techniques (see, for example, (see, for example, Sambrook et al., *Molecular Cloning*, 2nd Edition, Cold Spring Harbor Press, NY, 1989; Clover et al., *Molecular Cloning*, IRL Press, Oxford, 1985; Dason et al., *Virology*, 172:285-292, 1989; Takamatsu et al., *EMBO J.* 6:307-311, 1987; French et al., *Science* 231: 1294-1297, 1986; Takamatsu et al., *FEBS Lett.* 269:73-76, 1990; Yusibov and Loesch-Fries, *Virology*, 208(1): 405-7, 1995. Spitsin et al., *Proc Natl Acad Sci U S A*, 96(5): 2549-53, 1999, etc.) to generate viral vectors for use in accordance with the present invention. In general, a viral vector is a viral nucleic acid. Typically the viral vector is the genome, or a majority thereof (i.e., at least 50% of the genome), of a virus, or a nucleic acid molecule complementary in base sequence to such a nucleic acid molecule. In the case of segmented viruses, the viral vector may be a genome segment, or a majority thereof. The viral vector may be in RNA or DNA form.

[0067] Preferably the viral vector comprises a portion sufficient to support replication of the viral vector in the presence of the appropriate viral replicase proteins, i.e., constitutes a viral replicon. The ability of any particular portion of a viral genome to support replication of a nucleic acid that includes the portion, in the presence of viral replicase proteins, can readily be tested using methods known in the art, e.g., by making deletion mutants, by transferring the portion into a nucleic acid that does not

support replication and determining whether replication occurs, etc. The replicase proteins may be encoded by the vector, by another vector, or by a plant into which the vector is introduced. In certain preferred embodiments of the invention the vector is capable of self-replication, i.e., it encodes the necessary viral proteins for replication of the virus within an appropriate plant host. In certain embodiments of the invention the vector comprises a MP gene. In certain embodiments of the invention the vector comprises a CP gene. However, in certain embodiments of the invention neither an MP gene nor a CP gene is present in the vector. Since the clonal root lines, clonal plant lines, and clonal plants are derived from single ancestral cells into which the vector has been introduced, it is not necessary for the viral vector to have cell-to-cell or long distance movement capability. In particular, clonal plants can express the polynucleotide of interest throughout the plant even though the viral transcript does not move, since each cell is derived from a single ancestral cell that contains the viral vector.

[0068] In general, a polynucleotide of interest is inserted into a viral vector under control of (i.e., operably linked to), a promoter that directs transcription of the polynucleotide in a plant cell of interest. In certain preferred embodiments of the invention a plant viral promoter is used, e.g., a promoter for coat protein, movement protein, etc. The polynucleotide of interest may be inserted in place of the endogenous MP or CP coding sequence. For example, as described in more detail in the Examples, a TMV-based vector in which the TMV CP coding sequence has been replaced by a polynucleotide of interest, under control of the TMV CP promoter can be used. Alternately, the inserted polynucleotide may include its own promoter, which may be identical or similar to one of the naturally occurring viral promoters, may be from a different virus (e.g., the cauliflower mosaic virus), may be a non-viral promoter such as a promoter for a plant gene, or a synthetic promoter. In certain embodiments of the invention an inducible promoter is used. A variety of inducible promoters are known that function in plants. See, e.g., Zuo, J. and Chua, N-H., "Chemical-inducible systems for regulated expression of plant genes", *Curr. Op. in Biotechnol.*, 11:146-51, 2000. For example, promoters inducible by metals such as copper, or responsive to hormones such as estrogen, or systems responsive to other small molecules such as tetracycline

can be used. Other stimuli such as heat, light, etc., can be used. See U.S.S.N. 10/294,314.

[0069] In certain embodiments of the invention in any of its aspects, trans-activation is used to induce or increase expression of a polynucleotide of interest. For example, the expression cassette comprising the polynucleotide can be an inactive expression cassette that comprises an inactive or silenced foreign nucleic acid sequence, which is capable of directing expression of a polynucleotide of interest upon its activation. In certain embodiments of the invention trans-activation is accomplished by introducing a factor for activating or facilitating the expression of an inactive or silenced polynucleotide sequence into cells of the clonal entity. A promoter that can be activated in trans in such a manner is referred to as being “trans-activatable”. See U.S.S.N. 10/832,603, entitled “Expression of Foreign Sequences in Plants Using Trans-Activation System”, which is incorporated herein by reference, for further details of certain suitable methods. Such methods include techniques based on recombination (e.g., using a Lox/Cre or Flp/Frt recombinase system) and techniques based on proteins comprising a DNA binding domain such as GAL4 and a transcriptional activation domain such as VP16. A variety of other methods may be used for achieving trans-activation.

[0070] In certain embodiments of the invention the polynucleotide is inserted to create an independent open reading frame, while in other embodiments of the invention the polynucleotide is inserted to create an open reading frame in which a polynucleotide lacking a stop codon is inserted in frame with sequences encoding part or all of a viral protein such as CP, so that a fusion protein is produced upon translation. Multiple polynucleotides can be inserted. In certain preferred embodiments of the invention the TMV vector retains part or all of its 3' UTR and/or all or part of the CP coding sequence. In certain embodiments of the invention the polynucleotide of interest or a viral vector into which the polynucleotide of interest is inserted comprises a portion encoding a targeting sequence, e.g., a sequence that targets an encoded polypeptide to a particular intracellular organelle or compartment. For example, it may be desirable to target a polypeptide of interest to the endoplasmic reticulum, which may ultimately result in secretion of the polypeptide. The secreted polypeptide can then be harvested from culture medium or from interstitial fluid of a plant tissue.

[0071] Figures 1 – 5 show examples of engineering various plant virus vectors suitable for use in the present invention. Figure 1 shows a TMV based virus construct, D4, and the same construct following insertion of a polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as “target”) whose transcription is under control of the TMV CP subgenomic promoter. Details regarding the creation of such vectors are given in Example 1.

[0072] Figure 2 presents a schematic diagram of the engineering of a TMV based viral construct containing a polynucleotide of interest. The upper portion of the figure shows the genomic organization of a TMV based virus construct, 30B (Yusibov, V., Shivprasad, S., Turpen, T.H., Dawson, W., and Koprowski, H., “Plant viral vectors based on tobamoviruses”, in *Plant Biotechnology: New Products and Applications* (Eds. J. Hammond, P. McGarvey, and V. Yusibov), pp.81-94, Springer-Verlag, 1999). The lower portion shows the same construct following insertion of a polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as “target”). The 126/183 kDa protein is required for replication of the virus. The 30 kD protein is the movement protein (MP) that mediates cell-to-cell movement. CP is the coat protein that mediates systemic spread. Arrows indicate positions of the subgenomic promoters in certain embodiments of the invention. Transcription of the inserted polynucleotide is under control of an introduced promoter. CP expression is under control of the endogenous CP promoter in the construct shown in Figure 2.

[0073] Similar vectors in which polynucleotide of interest is in frame with the CP coding sequence so as to encode a fusion protein can also be used. In general, polynucleotides of interest (and their encoded proteins) can be expressed as independent open reading frames (see, e.g., Pogue, G.P., Lindbo, J.A., Dawson, W.O., and Turpen, T.H. “Tobamovirus transient expression vectors: tools for plant biology and high-level expression of foreign proteins in plants”, *Pl. Mol. Biol. Manual*. L4, 1-27., 1998) or as fusions with coat protein (Yusibov, V., Modelska, A., Steplewski, K., Agadjanyan, M., Weiner, D., Hooper, C. and Koprowski, H., “Antigens produced in plants by infection with chimeric plant viruses immunize against rabies virus and HIV-1”, *Proc. Natl. Acad. Sci. USA* 94, 5784-5788, 1997). In the vector described in the latter, target sequences are replicated from a second subgenomic promoter. In general, transcription of a polynucleotide of interest and/or an endogenous gene such as MP or

CP can be driven by endogenous promoters or inserted promoters (which may be identical to naturally occurring vectors from the same or a different virus or may be synthetic, or a combination of natural and synthetic sequences.

[0074] The 3' portion of the construct preferably includes the TMV 3' UTR, which may form stem-loop structure(s) as shown. The 3' portion of the construct may also include TMV coat protein sequences that contain a *cis* element that may be required for optimal replication. This sequence is optional.

[0075] Figure 3 presents a schematic diagram of the engineering of a TMV based viral construct containing a polynucleotide of interest and a gene encoding a marker, e.g., a marker that allows for detection and/or selection. The upper portion of the figure shows the genomic organization of a TMV based virus construct, D4. The middle portion of the figure shows the same construct after insertion of a gene encoding a detectable marker (GFP) replacing the MP coding sequence. The lower portion of the figure shows the same construct following insertion of a polynucleotide of interest (e.g., a gene encoding hGH, GCSF, GFP, etc., indicated as "target"). The 126/183 kDa protein is required for replication of the virus. Arrows indicate positions of the subgenomic promoters. Transcription of the detectable marker is under control of the MP subgenomic promoter. Transcription of the inserted polynucleotide of interest is under control of the TMV CP subgenomic promoter. However, other promoters could be used as described above. The 3' portion of the construct includes TMV coat protein sequences that contain a *cis* element that may be required for optimal replication and that may form stem-loop structure(s) as shown.

[0076] Figure 4 shows a vector similar to that shown in Figure 3 except that a selectable marker (a gene encoding a protein that confers resistance to kanamycin) is inserted instead of a gene encoding GFP. Including a gene that encodes a detectable or selectable marker in addition to a polynucleotide of interest is useful in the identification of clonal root lines and clonal plant cell lines that contain the vector and/or for identifying those lines that exhibit high and/or stable levels of expression.

[0077] In general, a wide variety of different markers can be used in accordance with the present invention. In general, a suitable marker for use in the invention is a detectable marker or a selectable marker. It is noted that in accordance with the practice in the art, the term "marker" can refer either to a nucleotide sequence, e.g., a

gene, that encodes a product (protein) that allows for detection or selection, or can be used to refer to the protein itself. The term “selectable marker” is used herein as it is generally understood in the art and refers to a marker whose presence within a cell or organism confers a significant growth or survival advantage or disadvantage on the cell or organism under certain defined culture conditions (selective conditions). For example, the conditions may be the presence or absence of a particular compound or a particular environmental condition such as increased temperature, increased radiation, presence of a compound that is toxic in the absence of the marker, etc. The presence or absence of such compound(s) or environmental condition(s) is referred to as a “selective condition” or “selective conditions”. By “growth advantage” is meant either enhanced viability (e.g., cells or organisms with the growth advantage have an increased life span, on average, relative to otherwise identical cells), increased rate of proliferation (also referred to herein as “growth rate”) relative to otherwise identical cells or organisms, or both. In general, a population of cells having a growth advantage will exhibit fewer dead or nonviable cells and/or a greater rate of cell proliferation than a population of otherwise identical cells lacking the growth advantage. Although typically a selectable marker will confer a growth advantage on a cell, certain selectable markers confer a growth disadvantage on a cell, e.g., they make the cell more susceptible to the deleterious effects of certain compounds or environmental conditions than otherwise identical cells not expressing the marker.

[0078] Antibiotic resistance markers are a non-limiting example of a class of selectable marker that can be used to select cells that express the marker. In the presence of an appropriate concentration of antibiotic (selective conditions), such a marker confers a growth advantage on a cell that expresses the marker. Thus cells that express the antibiotic resistance marker are able to survive and/or proliferate in the presence of the antibiotic while cells that do not express the antibiotic resistance marker are not able to survive and/or are unable to proliferate in the presence of the antibiotic. For example, a selectable marker of this type that is commonly used in plant cells is the NPTII protein, which encodes a protein that provides resistance against the antibiotic kanamycin. Additional selectable markers include proteins that confer resistance against carbenecillin (e.g., β -lactamases), proteins that confer resistance against gentamicin, hygromycin, etc.)

[0079] A second non-limiting class of selectable markers are nutritional markers. Such markers are generally enzymes that function in a biosynthetic pathway to produce a compound that is needed for cell growth or survival. In general, under nonselective conditions the required compound is present in the environment or is produced by an alternative pathway in the cell. Under selective conditions, functioning of the biosynthetic pathway in which the marker is involved is needed to produce the compound.

[0080] In general, a detectable marker is a marker whose presence within a cell can be detected through means other than subjecting the cell to a selective condition or directly measuring the level of the marker itself. Thus in general, the expression of a detectable marker within a cell results in the production of a signal that can be detected and/or measured. The process of detection or measurement may involve the use of additional reagents and may involve processing of the cell. For example, where the detectable marker is an enzyme, detection or measurement of the marker will typically involve providing a substrate for the enzyme. Preferably the signal is a readily detectable signal such as light, fluorescence, luminescence, bioluminescence, chemiluminescence, enzymatic reaction products, stainable products, or color. Thus preferred detectable markers for use in the present invention include fluorescent proteins such as green fluorescent protein (GFP) and variants thereof. Other suitable markers include luciferase, yellow fluorescent protein (YFP), lichenase, β -galactosidase, alkaline phosphatase, etc. Preferably the detectable marker is one that can be detected in intact, living root and/or plant cells.

[0081] Another example of a preferred viral vector for use in the present invention is an AlMV vector in which a polynucleotide of interest is inserted, as shown in Figure 5. For example, the polynucleotide of interest may replace the native AlMV CP encoding component in RNA3 of AlMV. Transcription of the polynucleotide of interest may be placed under control of the AlMV CP promoter. Alternately, the polynucleotide may replace the AlMV MP encoding component, and its transcription may be placed under control of the AlMV MP promoter. In other embodiments the inserted polynucleotide does not replace endogenous viral sequences. The polynucleotide of interest may be inserted in frame with CP coding sequences (complete or partial), so that a fusion protein is produced. In certain embodiments of

the invention the fusion protein comprises a cleavage site between the CP portion and the remainder, so that the fusion protein can be cleaved to yield a protein of interest free of CP sequences (or containing only a small number of such sequences). In certain embodiments of the invention the fusion protein assembles into particles, which can facilitate purification and/or antigen presentation (see, e.g., U.S. Patent Nos. 6,042,832 and 6,448,070).

[0082] Yet another example of a vector useful in the practice of the present invention is a cauliflower mosaic virus (CMV) viral vector in which a polynucleotide of interest is inserted under control of the CMV CP promoter, replacing the CMV CP encoding component found in the genome of naturally occurring CMV.

[0083] In certain embodiments of the invention it is desirable to insert a portion of coding or noncoding sequence from a viral vector of one virus type into a viral vector of another type. For example, certain sequences may enhance replication or expression, etc. Such sequences may comprise, for example, part or all of a viral transcript 5' or 3' UTR.

[0084] Generally, in order to preserve viral function and also simply for ease of genetic manipulation, viral vectors will be prepared by altering an existing plant virus genome, for example by removing particular genes and/or by disrupting or substituting particular sequences so as to inactivate or replace them. In such circumstances, the vectors will show very high sequence identity with natural viral genomes. Of course, completely novel vectors may also be prepared, for example, by separately isolating individual desired genetic elements and linking them together, optionally with the inclusion of additional elements. It is noted that when a plant virus vector is said to affirmatively express a particular protein or activity needed for viral replication, movement, or some other viral function, it is not necessary that the relevant gene be identical to the corresponding gene found in nature. So long as the protein is functional, it may be used in accordance with the present invention. Very high sequence identity with the natural protein, however, is generally preferred. For instance, large deletions (e.g., greater than about 25 amino acids) should generally be avoided according to certain embodiments of the invention. Typically, viral proteins expressed in accordance with the present invention will show at least 50%, preferably 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%

sequence identity with the corresponding natural viral protein. More particularly, the inventive viral protein should typically show 100% identity with critical functional portions (typically of at least several amino acids, often of at least 10, 20, 30, 40, 50 or more amino acids) of the relevant natural viral protein.

[0085] It is noted that in the case of many proteins a number of amino acid changes can be made without significantly affecting the functional activity and/or various other properties of the protein such as stability, etc. In particular, many proteins tolerate conservative amino acid changes, i.e., the substitution of an amino acid with a different amino acid having similar properties (conservative substitution) at many positions without significant reduction in activity. Conservative amino acid substitution is well known in the art and represents one approach to obtaining a polypeptide having similar or substantially similar properties to those of a given polypeptide while altering the amino acid sequence. In general, amino acids have been classified and divided into groups according to (1) charge (positive, negative, or uncharged); (2) volume and polarity; (3) Grantham's physico-chemical distance; and combinations of these. See, e.g., Zhang, J., *J. Mol. Evol.*, 50: 56-68, 2000; Grantham R., *Science*, 85: 862-864, 1974; Dagan, T., *et al.*, *Mol. Biol. Evol.*, 19(7), 1022-1025, 2002; *Biochemistry*, 4th Ed., Stryer, L., *et al.*, W. Freeman and Co., 1995; and U.S. Patent No. 6,015,692. For example, amino acids may be divided into the following 6 categories based on volume and polarity: special (C); neutral and small (A, G, P, S, T); polar and relatively small (N, D, Q, E), polar and relatively large (R, H, K), nonpolar and relatively small (I, L, M, V), and nonpolar and relatively large (F, W, Y). A conservative amino acid substitution may be defined as one that replaces one amino acid with an amino acid in the same group. Thus a variety of functionally equivalent proteins can be derived by making one or more amino acid substitutions, e.g., conservative amino acid substitutions, in a given viral protein.

[0086] *C. Clonal Root Lines*

[0087] The present invention provides methods for generating a clonal root line in which a plant viral vector is used to direct expression of a polynucleotide of interest. Figures 6A-6E show steps in the method according to certain embodiments of the invention. As shown in Figure 6, one or more viral expression vector(s) including a polynucleotide of interest operably linked to a promoter is introduced into a plant or a

portion thereof according to any of a variety of known methods. For example, as described in Example 2, plant leaves can be inoculated with viral transcripts. The vectors themselves may be directly applied to plants (e.g., via abrasive inoculations, mechanized spray inoculations, vacuum infiltration, particle bombardment, or electroporation). Alternatively, virions may be prepared (e.g., from already infected plants), and may be applied to other plants according to known techniques.

[0088] Where infection is to be accomplished by direct application of a viral genome to a plant, any available technique may be used to prepare the genome. For example, many viruses that are usefully employed in accordance with the present invention have ssRNA genomes. ssRNA may be prepared by transcription of a DNA copy of the genome, or by replication of an RNA copy, either *in vivo* or *in vitro*. Given the readily availability of easy-to-use *in vitro* transcription systems (e.g., SP6, T7, reticulocyte lysate, etc.), and also the convenience of maintaining a DNA copy of an RNA vector, it is expected that inventive ssRNA vectors will often be prepared by *in vitro* transcription, particularly with T7 or SP6 polymerase. Infectious cDNA clones can also be used. Agrobacterially mediated gene transfer can also be used to transfer viral nucleic acids such as viral vectors (either entire viral genomes or portions thereof) to plant cells using, e.g., agroinfiltration, according to methods known in the art.

[0089] Preferably the plant or plant portion is then maintained (e.g., cultured or grown) under conditions suitable for replication of the viral transcript. In certain embodiments of the invention the virus spreads beyond the initially inoculated cell, e.g., locally from cell to cell and/or systemically from an initially inoculated leaf into additional leaves. However, in other embodiments of the invention the virus does not spread. Thus the viral vector may contain genes encoding functional MP and/or CP, but may be lacking one or both of such genes. In general, the viral vector is introduced into (infects) multiple cells in the plant or portion thereof. Figure 6B shows a plant into which a viral vector (depicted schematically in Figure 6A) has been introduced.

[0090] Following introduction of the viral vector into the plant, leaves are harvested. Figure 6C shows leaf portions after harvesting from a virus-infected plant. In general, leaves may be harvested at any time following introduction of the viral vector. However, it may be preferable to maintain the plant for a period of time following introduction of the viral vector into the plant, e.g., a period of time sufficient

for viral replication and, optionally, spread of the virus from the cells into which it was initially introduced. A clonal root culture (or multiple cultures) is prepared, e.g., by known methods further described below and in Example 2.

[0091] In general, any available method may be used to prepare a clonal root culture from a plant or plant tissue into which a viral vector has been introduced. One such method employs genes that exist in certain bacterial plasmids. These plasmids are found in various species of *Agrobacterium* that infect and transfer DNA to a wide variety of organisms. As a genus, *Agrobacteria* can transfer DNA to a large and diverse set of plant types including numerous dicot and monocot angiosperm species and gymnosperms (See, Gelvin, S.B., “*Agrobacterium*-Mediated Plant Transformation: the Biology behind the “Gene-Jockeying” Tool”, *Microbiology and Molecular Biology Reviews*, 67(1): 16-37 (2003) and references therein, all of which are incorporated herein by reference). The molecular basis of genetic transformation of plant cells is transfer from the bacterium and integration into the plant nuclear genome of a region of a large tumor-inducing (Ti) or rhizogenic (Ri) plasmid that resides within various *Agrobacterial* species. This region is referred to as the T-region when present in the plasmid and as T-DNA when excised from the plasmid. Generally, a single-stranded T-DNA molecule is transferred to the plant cell in naturally occurring *Agrobacterial* infection and is ultimately incorporated (in double-stranded form) into the genome. Systems based on Ti plasmids are widely used for introduction of foreign genetic material into plants and for production of transgenic plants.

[0092] Infection of plants with various *Agrobacterial* species and transfer of the T-DNA has a number of effects. For example, *A. tumefaciens* causes crown gall disease while *A. rhizogenes* causes development of hairy roots at the site of infection, a condition known as “hairy root disease”. Each root arises from a single genetically transformed cell. Thus root cells in the roots are clonal, and each root represents a clonal population of cells. The roots produced by *A. rhizogenes* infection are characterized by a high growth rate and genetic stability. (Giri, A. and Narasu, M.L., *Biotechnology Advances*, 18: 1-22 (2000) and references therein, all of which are incorporated herein by reference). In addition, such roots are able to regenerate genetically stable plants (Giri 2000).

[0093] In general, the present invention encompasses the use of any strain of *Agrobacteria*, particularly *A. rhizogenes* strains, that is capable of inducing formation of roots from plant cells. As mentioned above, a portion of the Ri plasmid (Ri T-DNA) is responsible for causing hairy root disease. While transfer of this portion of the Ri plasmid to plant cells can conveniently be accomplished by infection with *Agrobacteria* harboring the Ri plasmid, the invention also encompasses the use of alternative methods of introducing the relevant region into a plant cell. Such methods include any available method of introducing genetic material into plant cells including, but not limited to, biolistics, electroporation, PEG-mediated DNA uptake, Ti-based vectors, etc. The relevant portions of the Ri T-DNA can also be introduced into plant cells by use of a viral vector. The Ri genes can be included in the same vector that contains the polynucleotide of interest or in a different viral vector, which can be the same or a different type to that of the vector that contains the polynucleotide of interest. It is noted that the entire Ri T-DNA may not be required for production of hairy roots, and the invention encompasses the use of portions of the Ri T-DNA, provided that such portions contain sufficient genetic material to induce root formation, as known in the art. Additional genetic material, e.g., genes present within the Ri plasmid but not within the T-DNA, may also be transferred to the plant cell in accordance with the invention, particularly genes whose expression products facilitate integration of the T-DNA into the plant cell DNA.

[0094] In order to prepare a clonal root line in accordance with certain embodiments of the invention, the harvested leaf portions are contacted with *A. rhizogenes* under conditions suitable for infection and transformation. Example 2 describes one method for generating root lines from leaves into which a viral vector has been introduced. The leaf portions are maintained in culture to allow development of hairy roots. Figure 6D shows hairy roots generated by individual cells in leaf portions infected with *A. rhizogenes*. Each root is clonal, i.e., cells in the root are derived from a single ancestral cell into which the Ri T-DNA was transferred. In accordance with the invention, a portion of such ancestral cells will also contain the viral vector. Thus cells in a root derived from such an ancestral cell will also contain the viral vector since it will be replicated and will be transmitted during cell division. Thus a high proportion, preferably at least 50%, more preferably at least 75%, at least 80%, at least 90%, at

least 95%, or all (100%) or substantially all (at least 98%) of the cells will contain the viral vector. It is noted that since the viral vector is inherited by daughter cells within the clonal root, movement of the viral vector within the root is not necessary to maintain the viral vector throughout the root.

[0095] Individual clonal hairy roots may be removed from the leaf portion and further cultured, as shown in Figures 6E and 6F. Such roots are also referred to herein as root lines. Figure 6E shows individual clonal roots placed in a line in a Petri dish. Figure 6F shows the same root lines at higher magnification. The roots continue to grow. These roots were derived from plants into which a viral vector containing a GFP gene had been introduced. Figure 6G shows a photograph of single root taken under UV light. Expression of GFP throughout the root is evident.

[0096] As described in Examples 2 – 4, a variety of different clonal root lines have been generated using the inventive methods. These root lines were generated using viral vectors containing polynucleotides of interest encoding GFP, hGH, and GCSF. The root lines were tested by Western blot. Root lines displayed a variety of different expression levels of the various polypeptides. Root lines displaying high expression were selected and further cultured. These root lines were subsequently tested again and shown to maintain high levels of expression over extended periods of time, indicating stability. The level of expression was comparable to or greater than expression in intact plants infected with the same viral vector used to generate the clonal root lines. In addition, the stability of expression of the root lines was superior to that obtained in plants infected with the same viral vector. Up to 80% of such virus-infected plants reverted to wild type after 2 – 3 passages. (Such passages involved inoculating plants with transcripts, allowing the infection (local or systemic) to become established, taking a leaf sample, and inoculating fresh plants that are subsequently tested for expression.)

[0097] The root lines may be cultured on a large scale for production of polypeptides of interest as discussed further below. It is noted that the clonal root lines (and cell lines derived from the clonal root lines) can generally be maintained in medium that does not include various compounds, e.g., plant growth hormones such as auxins, cytokinins, etc., that are typically employed in the culture of root and plant cells. This feature greatly reduces the expense associated with tissue culture, and the

inventors expect that it will contribute significantly to the economic feasibility of protein production using plants.

[0098] Any of a variety of methods may be used to select clonal roots that express a polynucleotide of interest. Western blots, ELISA assays, etc., can be used to detect an encoded polypeptide. In the case of detectable markers such as GFP, alternative methods such as visual screens can be performed. If a viral vector that contains a polynucleotide that encodes a selectable marker is used, an appropriate selection can be imposed (e.g., the leaf material and/or roots derived therefrom can be cultured in the presence of an appropriate antibiotic or nutritional condition and surviving roots identified and isolated). Certain viral vectors contain two or more polynucleotides of interest, e.g., two or more polynucleotides encoding different polypeptides. If one of these is a selectable or detectable marker, clonal roots that are selected or detected by selecting for or detecting expression of the marker will have a high probability of also expressing the second polynucleotide. Screening for root lines that contain particular polynucleotides can also be performed using PCR and other nucleic acid detection methods.

[0099] Alternatively, clonal root lines can also be screened for presence of the virus by inoculating host plants that will form local lesions as a result of virus infection (e.g., hypersensitive host plants). For example, 5 mg of root tissue can be homogenized in 50 μ l of phosphate buffer and used to inoculate a single leaf of a tobacco plant. If the virus is present in root cultures, within two to three days characteristic lesions will appear on the infected leaves. This means that the root line contains recombinant virus that carries the polynucleotide of interest (target gene). If no local lesions are formed, there is no virus, and the root line is rejected as negative. This method is highly time and cost efficient. After initially screening for the presence of virus, roots that contain the virus are subjected to secondary screening, e.g., by Western blot or ELISA to select high expressers. Additional screens, e.g., screens for rapid growth, growth in particular media or under particular environmental conditions, etc., can also be applied. These screening methods may, in general, be applied in the development of any of the clonal root lines, clonal root cell lines, clonal plant cell lines, and/or clonal plants described herein.

[00100] As will be evident to one of ordinary skill in the art, a variety of modifications may be made to the above description of the inventive methods for generating clonal root lines that contain a viral vector. Such modifications are within the scope of the invention. For example, while it is generally preferred to introduce the viral vector into an intact plant or portion thereof prior to introduction of the Ri T-DNA genes, in certain embodiments of the invention the Ri-DNA is introduced prior to introducing the viral vector. In addition, it is also possible to contact intact plants with *A. rhizogenes* rather than harvesting leaf portions and then exposing them to the bacterium.

[00101] Other methods of generating clonal root lines from single cells of the plant or portion thereof that harbor the viral vector can also be used (i.e., methods not using *A. rhizogenes* or genetic material from the Ri plasmid). For example, treatment with certain plant hormones or combinations of plant hormones is known to result in generation of roots from plant tissue.

[00102] In certain embodiments of the invention rather than introducing a single viral vector type into the plant, multiple different viral vectors are introduced. Such vectors may, for example, trans-complement each other with respect to functions such as replication, cell-to-cell movement, and/or long distance movement. The vectors may contain different polynucleotides of interest, e.g., polynucleotides that encode individual polypeptides that associate to form a single protein complex such as antibodies, etc., or polynucleotides that encode different enzymes in a biosynthetic pathway. Selection for roots that express multiple polypeptides of interest may be performed as described above for single polynucleotides or polypeptides.

[00103] *D. Clonal Cell Lines Derived from Clonal Root Lines*

[00104] As described above, the invention provides methods for generating clonal root lines, wherein cells in the root lines contain a viral vector. As is well known in the art, a variety of different cell lines can be generated from roots. For example, root cell lines can be generated from individual root cells obtained from the root using a variety of known methods. Such root cell lines may be obtained from various different root cell types within the root. In general, root material is harvested and dissociated (e.g., physically and/or enzymatically digested) to release individual root cells, which are then further cultured. Complete protoplast formation is generally not necessary. If

desired, root cells can be plated at very dilute cell concentrations, so as to obtain root cell lines from single root cells. Root cell lines derived in this manner are clonal root cell lines contain the viral vector. Such root cell lines therefore exhibit stable expression of the polynucleotide of interest. Clonal plant cell lines can also be obtained in a similar manner from the clonal roots, e.g., by culturing dissociated root cells in the presence of the appropriate plant hormones. Screens and successive rounds of enrichment can be used to identify cell lines that express the polynucleotide of interest at high levels. However, if the clonal root line from which the cell line is derived already expresses at high levels, such additional screens may be unnecessary.

[00105] As in the case of the clonal root lines, cells of a clonal root cell line are derived from a single ancestral cell that contains the viral vector and will, therefore, also contain the viral vector since it will be replicated and will be transmitted during cell division. Thus a high proportion, preferably at least 50%, more preferably at least 75%, at least 80%, at least 90%, at least 95%, or all (100%) or substantially all (at least 98%) of the cells will contain the viral vector. It is noted that since the viral vector is inherited by daughter cells within the clonal root cell line, movement of the viral vector among the cells is not necessary to maintain the viral vector. The clonal root cell lines can be used for production of a polynucleotide of interest as described below.

[00106] *E. Clonal Plant Cell Lines*

[00107] The present invention provides methods for generating a clonal plant cell line in which a plant viral vector is used to direct expression of a polynucleotide of interest. According to the inventive method, one or more viral expression vector(s) including a polynucleotide of interest operably linked to a promoter is introduced into cells of a plant cell line that is maintained in cell culture. A number of plant cell lines from various plant types are known in the art, any of which can be used. Newly derived cell lines can also be generated according to known methods for use in practicing the invention. A viral vector is introduced into cells of the plant cell line according to any of a number of methods. For example, as described in Example 5, protoplasts can be made and viral transcripts then electroporated into the cells. Other methods of introducing a plant viral vector into cells of a plant cell line can also be used.

[00108] Figure 13 shows steps in a method for generating clonal plant cell lines in accordance with the invention. Figure 13A shows a viral vector suitable for introduction into plant cells (e.g., protoplasts). Following introduction of the viral vector, the plant cell line may be maintained in tissue culture, e.g., as shown in Figures 13B and 13C. During this time the viral vector may replicate, and polynucleotides of interest may be expressed. Clonal plant cell lines are derived from the culture, e.g., by a process of successive enrichment. For example, as shown in Figure 13E, samples may be removed from the culture, optionally with dilution so that the concentration of cells is low, and plated in Petri dishes in individual droplets. The droplets are then maintained to allow cell division.

[00109] It will be appreciated that the droplets may contain a variable number of cells, depending on the initial density of the culture and the amount of dilution. The cells can be diluted such that most droplets contain either 0 or 1 cell if it is desired to obtain clonal cell lines expressing the polynucleotide of interest after only a single round of enrichment. However, it can be more efficient to select a concentration such that multiple cells are present in each droplet and then screen the droplets to identify those that contain expressing cells. In general, any appropriate screening procedure can be employed. For example, selection or detection of a detectable marker such as GFP can be used. Figure 13F is a photograph taken under UV light and showing individual droplets in which cell lines expressing GFP from a viral vector are present. Western blots or ELISA assays can also be used. Individual droplets (100 ul) contain more than enough cells for performance of these assays. Multiple rounds of enrichment are performed to isolate successively higher expressing cell lines. Single clonal plant cell lines (i.e., populations derived from a single ancestral cell) can be generated by further limiting dilution using standard methods for single cell cloning. However, it is not necessary to isolate individual clonal lines. A population containing multiple clonal cell lines can also be used for expression of a polynucleotide of interest.

[00110] In general, certain considerations described above for generation of clonal root lines also apply to the generation of clonal plant cell lines. For example, a diversity of viral vectors containing one or more polynucleotides of interest can be used as can combinations of multiple different vectors. Similar screening methods can also be used. As in the case of the clonal root lines and clonal root cell lines, cells of a

clonal plant cell line are derived from a single ancestral cell that contains the viral vector and will, therefore, also contain the viral vector since it will be replicated and will be transmitted during cell division. Thus a high proportion, preferably at least 50%, more preferably at least 75%, at least 80%, at least 90%, at least 95%, or all (100%) or substantially all (at least 98%) of the cells will contain the viral vector. It is noted that since the viral vector is inherited by daughter cells within the clonal plant cell line, movement of the viral vector among the cells is not necessary to maintain the viral vector. The clonal plant cell line can be used for production of a polypeptide of interest as described below.

[00111] *F. Clonal Plants*

[00112] Clonal plants can be generated from the clonal roots, clonal root cell lines, and/or clonal plant cell lines produced according to the various methods described above. Methods for the generation of plants from roots, root cell lines, and plant cell lines such as the clonal root lines, clonal root cell lines, and clonal plant cell lines described herein are well known in the art (See, e.g., Peres et al., *Plant Cell, Tissue, and Organ Culture* 65, 37-44, 2001 and standard reference works on plant molecular biology and biotechnology cited elsewhere herein. The invention therefore provides a method of generating a clonal plant comprising steps of (i) generating a clonal root line, clonal root cell line, or clonal plant cell line according to any of the inventive methods described above; and (ii) generating a whole plant from the clonal root line, clonal root cell line, or clonal plant. The clonal plants may be propagated and grown according to standard methods. Example 7 describes generation of a clonal plant from a clonal root line containing a viral vector that encodes human growth hormone.

[00113] As in the case of the clonal root lines, clonal root cell lines, and clonal plant cell lines, the cells of a clonal plant are derived from a single ancestral cell that contains the viral vector and will, therefore, also contain the viral vector since it will be replicated and will be transmitted during cell division. Thus a high proportion, preferably at least 50%, more preferably at least 75%, at least 80%, at least 90%, at least 95%, or all (100%) or substantially all (at least 98%) of the cells will contain the viral vector. It is noted that since the viral vector is inherited by daughter cells within the clonal plant, movement of the viral vector is not necessary to maintain the viral vector.

[00114] *II. Plant Species*

[00115] Any plant susceptible to viral infection may be utilized in accordance with the present invention. In general, it will often be desirable to utilize plants that are amenable to growth under defined conditions, for example in a greenhouse and/or in aqueous systems. It may also be desirable to select plants that are not typically consumed by human beings or domesticated animals and/or are not typically part of the human food chain, so that they may be grown outside without concern that the expressed polynucleotide may be undesirably ingested. In other embodiments, however, it will be desirable to employ edible plants.

[00116] Often, certain desirable plant characteristics will be determined by the particular polynucleotide to be expressed. To give but a few examples, when the polynucleotide encodes a protein to be produced in high yield (as will often be the case, for example, when therapeutic proteins are to be expressed), it will often be desirable to select plants with relatively high biomass (e.g., tobacco, which has the additional advantages that it is highly susceptible to viral infection, has a short growth period, and is not in the human food chain). Where the polynucleotide encodes a protein whose full activity requires (or is inhibited by) a particular post-translational modification, the ability (or inability) of certain plant species to accomplish the relevant modification (e.g., a particular glycosylation) may direct selection.

[00117] In certain preferred embodiments of the invention, crop plants, or crop-related plants are utilized. In some particularly preferred embodiments, edible plants are utilized.

[00118] Preferred plants for use in accordance with the present invention include Angiosperms, Bryophytes (e.g., Hepaticae, Musci, etc.), Pteridophytes (e.g., ferns, horsetails, lycopods), Gymnosperms (e.g., conifers, cycase, Ginko, Gnetales), and Algae (e.g., Chlorophyceae, Phaeophyceae, Rhodophyceae, Myxophyceae, Xanthophyceae, and Euglenophyceae). Particularly preferred are members of the family Leguminosae (Fabaceae; e.g., pea, alfalfa, soybean); Gramineae (Poaceae; e.g., corn, wheat, rice); Solanaceae, particularly of the genus *Lycopersicon* (e.g., tomato), *Solanum* (e.g., potato, eggplant), *Capsium* (e.e., pepper), or *Nicotiana* (e.g., tobacco); Umbelliferae, particularly of the genus *Daucus* (e.g., carrot), *Apium* (e.g., celery), or *Rutaceae* (e.g., oranges); Compositae, particularly of the genus *Lactuca* (e.g., lettuce);

Brassicaceae (Cruciferae), particularly of the genus *Brassica* or *Sinapis*. Particularly preferred Brassicaceae family members include *Brassica campestris*, *B. carinata*, *B. juncea*, *B. napus*, *B. nigra*, *B. oleraceae*, *B. tournifortii*, *Sinapis alba*, and *Raphanus sativus*.

[00119] *III. Polynucleotides and Polypeptides of Interest*

[00120] The teachings of the present invention may be employed to deliver to and/or express in plant cells any polynucleotide of interest. For example, protein-coding polynucleotides may express enzymes, antibodies, hormones, cytokines, regulatory factors, structural proteins, or any other protein or polypeptide of interest. Encoded proteins may be naturally-occurring proteins, or may be designed or engineered proteins, including for instance fusion proteins (e.g., fusion proteins incorporating part or all of a plant virus protein such as MP or CP). See, e.g., U.S. Pat. Nos. 6,448,070 and 6,660,500. Numerous types of fusion proteins may be encoded. A heterologous sequence may be fused to the 5' or 3' end of a plant virus protein or located internally. Numerous sequences of diverse origin may be included within a single fusion protein. The encoded protein may comprise a cleavage site, which may be encoded by the inserted polynucleotide or by the viral vector. See, e.g., U.S. Pat. No. 6,740,740. For example, the vector may comprise a portion that encodes a cleavage site upstream of a portion that encodes CP so that when a polynucleotide of interest is inserted between the CP promoter and the portion that encodes a cleavage site, the resulting open reading frame encodes a fusion protein containing a portion encoded by the polynucleotide of interest, a cleavage site, and part or all of the CP. Cleavage of the fusion protein at the cleavage site releases the encoded polypeptide of interest. The cleavage site may be a site for cleavage by chemical means (e.g., cyanogen bromide) or by enzymatic means (e.g., by a protease such as trypsin, chymotrypsin, thrombin, pepsin, *Staphylococcus aureus* V8 protease, and Factor Xa protease).

[00121] In certain embodiments of the invention the polynucleotide of interest comprises a portion encoding a tag, e.g., a 6X-His tag, HA tag, Myc tag, FLAG tag, etc. Such tags may simplify the detection, isolation and/or purification of the protein. In certain embodiments of the invention the tag is a cleavable tag, e.g., a tag cleavable by chemical means or by enzymatic means as described above. Including a cleavage site allows the tag to be readily be removed from the translated polypeptide, e.g., after

purification, resulting in a protein with wild type sequence. It is to be understood that the tag and/or cleavage site may be present within a viral vector into which a particular polynucleotide of interest is to be inserted and need not be present within the inserted polynucleotide itself. Once the polynucleotide is inserted, the entire portion comprising the region(s) that encode the tag, cleavage site, and newly inserted polynucleotide is considered a polynucleotide of interest.

[00122] In some instances, it may be desirable to utilize the inventive system to express more than one polypeptide chain in the same clonal root or plant cell line or clonal plant (e.g., using two different viral vectors each of which directs expression of a polynucleotide, inserting two different polynucleotides into one viral vector, utilizing a transgenic plant that expresses one or more polynucleotides to generate a clonal root or plant cell line or clonal plant), for example in order to produce a multimeric protein or to simultaneously produce two different proteins such as a protein of interest and a detectable or selectable marker).

[00123] In certain preferred embodiments of the invention, a polynucleotide that encodes a therapeutically active protein is employed. Exemplary proteins that have been approved for therapeutic uses include, for example, insulin, human growth hormone, interferons, albumin, tPA, erythropoietin, interleukins, factor VIII, DNase, factor IX, PDGF, FSH, TNF receptor (soluble form), calcitonin, and a variety of immunoglobulins. Of course, the invention is not limited to such approved proteins, but encompasses expression of any polynucleotide(s), whether protein-coding or not, and particularly encompasses expression of any polynucleotide encoding any therapeutically active protein, whether prokaryotic or eukaryotic in origin, etc.

[00124] Generally, the pharmaceutical proteins of interest include, but are not limited to, hormones (insulin, thyroid hormone, catecholamines, gonadotrophins, trophic hormones, prolactin, oxytocin, dopamine, bovine somatotropin, leptins and the like), growth hormones (e.g., human growth hormone), growth factors (e.g., epidermal growth factor, nerve growth factor, insulin-like growth factor and the like), growth factor receptors, cytokines and immune system proteins (e.g., interleukins, colony stimulating factor (CSF), granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), erythropoietin, tumor necrosis factor (TNF), interferons, integrins, addressins, selectins, homing receptors, T cell receptors,

immunoglobulins, soluble major histocompatibility complex antigens, immunologically active antigens such as bacterial, parasitic, or viral antigens or allergens), autoantigens, antibodies), enzymes (tissue plasminogen activator, streptokinase, cholesterol biosynthetic or degradative, steroidogenic enzymes, kinases, phosphodiesterases, methylases, de-methylases, dehydrogenases, cellulases, proteases, lipases, phospholipases, aromatases, cytochromes, adenylate or guanylate cyclases, neuramidases and the like), receptors (steroid hormone receptors, peptide receptors), binding proteins (steroid binding proteins, growth hormone or growth factor binding proteins and the like), transcription and translation factors, oncoproteins or proto-oncoproteins (e.g., cell cycle proteins), muscle proteins (myosin or tropomyosin and the like), myeloproteins, neuroactive proteins, tumor growth suppressing proteins (angiostatin or endostatin, both of which inhibit angiogenesis), anti-sepsis proteins (bactericidal permeability-increasing protein), structural proteins (such as collagen, fibronin, fibrinogen, elastin, tubulin, actin, and myosin), blood proteins (thrombin, serum albumin, Factor VII, Factor VIII, insulin, Factor IX, Factor X, tissue plasminogen activator, Protein C, von Willebrand factor, antithrombin III, glucocerebrosidase, erythropoietin granulocyte colony stimulating factor (GCSF) or modified Factor VIII, anticoagulants such as hirudin) and the like.

[00125] In one particular example, the present invention may be utilized to produce vaccine components. In general, it is desirable to include in vaccines proteins, or portions of proteins, to which a human or animal immune system is exposed when the human or animal is infected with a pathogen, or suffering some other undesirable event (e.g., development of a tumor). Thus, proteins or polypeptides that may be formulated in a vaccine include virtually any potentially antigenic protein or portion thereof, for example, viral coat proteins, viral fusion proteins, viral envelope proteins, viral glycoproteins, bacterial or fungal cell wall proteins, toxin proteins, parasite coat proteins, tumor-specific antigens, etc., or portions of any of the foregoing. See, e.g., WO9640229. Viruses of interest include HIV, respiratory syncytial virus (RSV), rabies virus, polio virus, pneumoviruses, metapneumoviruses, influenza viruses, poxviruses (including smallpox), rhinoviruses, coronaviruses, adenoviruses, herpesviruses, hantaviruses, Ebola virus, Yellow Fever virus, Dengue virus, hepatitis viruses (e.g., hepatitis A, B, C, D, E, F, or G virus) etc. Bacteria of interest include *Neisseria*,

Pneumococcus, *Streptococcus*, *H. influenzae*, *Staphylococcus*, anthrax, etc. Parasites of interest include *Plasmodium*, *Leishmania*, *Toxoplasma*, *Ascaris*, hookworm and other nematodes, ameba, flukes, etc.

[00126] In other embodiments, the inventive system may be used to express a polynucleotide encoding an enzyme that synthesizes or modifies a biologically active agent. For instance, certain enzymes (e.g., polyketide synthases, polypeptide synthetases, terpene synthases, etc.) synthesize small molecules with interesting biological activities, including therapeutic activities (e.g., antibiotic, anticancer, immunosuppressive activities, etc.). Also, a large number of enzymes that modify protein or small molecule substrates (e.g., kinases, hydrolases, transferases, etc.) is known. See U.S. Patent No. 6,500,644 for additional proteins that can be desirably expressed in plants using the inventive systems described herein.

[00127] In certain embodiments of the invention the polynucleotide encodes a component (e.g., an enzyme) in a biosynthetic pathway. Plants are a source of numerous natural products of use for medicinal and/or industrial purposes and others. It is of interest to increase the level or efficiency by which such products are produced. To this end, a polynucleotide of interest may encode a biosynthetic enzyme, e.g., an enzyme that catalyzes a rate-limiting step in a biosynthetic pathway, by which such natural product(s) are synthesized.

[00128] In other embodiments, the inventive system may be used to produce diagnostic or research reagents including, for example, antibodies.

[00129] In yet other embodiments of the invention the polynucleotide encodes a protein that enhances plant growth or survival in any of a variety of ways. For example, the protein may enhance the ability of the plant to extract nutrients from soil or culture medium, may confer resistance to an environmental condition such as temperature, salinity, etc., or may confer resistance to a pathogen such as a virus, bacterium, fungus, nematode, insect, etc. An example is the various plant peptides known as defensins (Thomma, B.P., *et al.*, *Planta*, 216(2):193-202, 2002) Such proteins include both endogenous plant proteins (i.e., proteins that are naturally expressed in the plant from which the clonal root line, clonal plant cell line, or clonal plant is derived) and non-endogenous proteins.

[00130] In yet other embodiments, the inventive system may be utilized to produce nutritionally relevant proteins or other products. Nutritionally relevant proteins include, for example, proteins that are found naturally in foods consumed by humans or domesticated animals (e.g., cats, dogs). Other examples include proteins having a balanced amino acid composition, e.g., proteins having an amino acid composition such as those used for total parenteral nutrition (TPN), etc.

[00131] In still other embodiments, the inventive system may be utilized to express polynucleotides that do not necessarily encode proteins, for example to produce active RNA species, e.g., ribozymes or interfering RNAs that silence gene expression (either long double-stranded RNAs or short interfering RNAs (siRNAs), microRNAs or microRNA precursors, short hairpin RNAs (shRNAs), etc. See, e.g., U.S. Pat. No. 6,531,647; 6,635,805 and U.S. Pub. No. 20040019930. In some embodiments, ribozymes or interfering RNAs may be produced that target plant genes, so that an altered plant is created, for example that does not express a particular receptor for a plant pathogen, or a particular allergenic protein.

[00132] *IV. Culturing or Growing Clonal Root Lines, Clonal Root Cell Lines, Clonal Plant Cell Lines, and Clonal Plants*

[00133] In general, standard methods known in the art may be used for culturing or growing the clonal root lines, clonal root cell lines, clonal plant cell lines, and clonal plants of the invention. A wide variety of culture media and bioreactors have been employed to culture hairy root cells, root cell lines, and plant cells. See, for example, Giri, A. and Narasu, M.L., *Biotechnol. Adv.* 18:1-22, 2000; Rao, S.R. and Ravishankar, G.A., *Biotechnol. Adv.* 20:101-153, 2002, and references in both of the foregoing, all of which are incorporated herein by reference. Clonal plants may be grown in any suitable manner.

[00134] *V. Isolation and/or Formulation of Polynucleotide Expression Products*

[00135] In many embodiments of the present invention, it will be desirable to isolate polynucleotide expression products from the plant tissue(s), e.g., roots, root cells, plants, plant cells, that express them. It may also be desirable to formulate such isolated products for their intended use (e.g., as a pharmaceutical or diagnostic agent, or as a reagent, etc.). In other embodiments, it will be desirable to formulate the products together with some or all of the plant tissues that express them.

[00136] Where it is desirable to isolate the expression product from some or all of the plant cells or tissues that express it, any available purification techniques may be employed. Those of ordinary skill in the art are familiar with a wide range of fractionation and separation procedures (see, for example, Scopes et al., *Protein Purification: Principles and Practice*, 3rd Ed., Janson et al., 1993; *Protein Purification: Principles, High Resolution Methods, and Applications*, Wiley-VCH, 1998; Springer-Verlag, NY, 1993; Roe, *Protein Purification Techniques*, Oxford University Press, 2001, each of which is incorporated herein by reference). Often, it will be desirable to render the product more than about 50%, preferably more than about 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% pure. See, e.g., U.S. Pat. Nos. 6,740,740 and 6,841,659 for discussion of certain methods useful for purifying substances from plant tissues or fluids.

[00137] Where it is desirable to formulate the product together with the plant material, it will often be desirable to have utilized a plant that is not toxic to the relevant recipient (e.g., a human or other animal). Relevant plant tissue (e.g., cells, roots, leaves) may simply be harvested and processed according to techniques known in the art, with due consideration to maintaining activity of the expressed product. In certain embodiments of the invention, it is desirable to have expressed the polynucleotide in an edible plant (and, specifically in edible portions of the plant) so that the material can subsequently be eaten. For instance, where the polynucleotide encodes a nutritionally relevant protein, or a therapeutic protein that is active after oral delivery (when properly formulated), it may be desirable to produce the protein in an edible plant portion, and to formulate the expressed polynucleotide for oral delivery together with the some or all of the plant material with which the polynucleotide was expressed.

[00138] Where the polynucleotide encodes or produces a therapeutic agent, it may be formulated according to known techniques. For example, an effective amount of a pharmaceutically active product can be formulated together with one or more organic or inorganic, liquid or solid, pharmaceutically suitable carrier materials. A pharmaceutically active product produced according to the present invention may be employed in dosage forms such as tablets, capsules, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, powder packets, liquid solutions,

solvents, diluents, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, and solid bindings, as long as the biological activity of the protein is not destroyed by such dosage form.

[00139] Materials that can serve as pharmaceutically acceptable carriers include, but are not limited to sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols such as propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening agents, flavoring agents, and perfuming agents, preservatives, and antioxidants can also be present in the composition, according to the judgment of the formulator (see also *Remington's Pharmaceutical Sciences*, Fifteenth Edition, E.W. Martin (Mack Publishing Co., Easton PA, 1975). For example, the polynucleotide expression product may be provided as a pharmaceutical composition by means of conventional mixing granulating dragee-making, dissolving, lyophilizing, or similar processes.

[00140] In certain preferred embodiments, it may be desirable to prolong the effect of a pharmaceutical preparation by slowing the absorption of the pharmaceutically active product (e.g., protein) that is subcutaneously or intramuscularly injected. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the product then depends upon its rate of dissolution, which in turn, may depend upon size and form. Alternatively, delayed absorption of a parenterally administered product is accomplished by dissolving or suspending the product in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the protein in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of product to polymer and the nature of the particular polymer employed, the rate of release can be

controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations may be prepared by entrapping the product in liposomes or microemulsions, which are compatible with body tissues.

[00141] Enterally administered preparations of pharmaceutically active products may be introduced in solid, semi-solid, suspension or emulsion form and may be compounded with any pharmaceutically acceptable carriers, such as water, suspending agents, and emulsifying agents. The expression products may also be administered by means of pumps or sustained-release forms, especially when administered as a preventive measure, so as to prevent the development of disease in a subject or to ameliorate or delay an already established disease.

[00142] Pharmaceutically active products, optionally together with plant tissue, are particularly well suited for oral administration as pharmaceutical compositions. Harvested plant material may be processed in any of a variety of ways (e.g., air drying, freeze drying, extraction etc.), depending on the properties of the desired therapeutic product and its desired form. In preferred embodiments, such compositions as described above are ingested orally alone or ingested together with food or feed or a beverage. Compositions for oral administration include plants; extractions of the plants, and proteins purified from infected plants provided as dry powders, foodstuffs, aqueous or non-aqueous solvents, suspensions, or emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil, fish oil, and injectable organic esters. Aqueous carriers include water, water-alcohol solutions, emulsions or suspensions, including saline and buffered medial parenteral vehicles including sodium chloride solution, Ringer's dextrose solution, dextrose plus sodium chloride solution, Ringer's solution containing lactose or fixed oils. Examples of dry powders include any plant biomass that has been dried, for example, freeze dried, air dried, or spray dried. For example, the plants may be air dried by placing them in a commercial air dryer at about 120 degrees Fahrenheit until the biomass contains less than 5% moisture by weight. The dried plants may be stored for further processing as bulk solids or further processed by grinding to a desired mesh sized powder. Alternatively, freeze-drying may be used for products that are sensitive to air-drying. Products may be freeze dried by placing them into a vacuum drier and dried frozen

under a vacuum until the biomass contains less than about 5% moisture by weight. The dried material can be further processed as described herein.

[00143] Plant-derived material may be administered as or together with one or more herbal preparations. Useful herbal preparations include liquid and solid herbal preparations. Some examples of herbal preparations include tinctures, extracts (e.g., aqueous extracts, alcohol extracts), decoctions, dried preparations (e.g., air-dried, spray dried, frozen, or freeze-dried), powders (e.g., lyophilized powder), and liquid. Herbal preparations can be provided in any standard delivery vehicle, such as a capsule, tablet, suppository, liquid dosage, etc. Those skilled in the art will appreciate the various formulations and modalities of delivery of herbal preparations that may be applied to the present invention.

[00144] Those skilled in the art will also appreciate that a particularly preferred method of obtaining the desired pharmaceutically active products is by extraction. Plant material (e.g., roots, leaves, etc.) may be extracted to remove the desired products from the residual biomass, thereby increasing the concentration and purity of the product. Plants may also be extracted in a buffered solution. For example, the plant material may be transferred into an amount of ice-cold water at a ratio of one to one by weight that has been buffered with, e.g., phosphate buffer. Protease inhibitors can also be added as required. The plant material can be disrupted by vigorous blending or grinding while suspended in the buffer solution and the extracted biomass removed by filtration or centrifugation. The product carried in solution can be further purified by additional steps or converted to a dry powder by freeze-drying or precipitation. Extraction can also be carried out by pressing. Plants or roots can also be extracted by pressing in a press or by being crushed as they are passed through closely spaced rollers. The fluids expressed from the crushed plants or roots are collected and processed according to methods well known in the art. Extraction by pressing allows the release of the products in a more concentrated form. However, the overall yield of the product may be lower than if the product were extracted in solution.

[00145] Inventive root lines, cell lines, plants, extractions, powders, dried preparations and purified protein or nucleic acid products, etc., can also be in encapsulated form with or without one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with

coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active product may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[00146] In other particularly preferred embodiments, a plant or portion thereof expressing a pharmaceutically active product according to the present invention, or biomass thereof, is administered orally as medicinal food. Such edible compositions are consumed by eating raw, if in a solid form, or by drinking, if in liquid form. In a preferred embodiment, the plant material is directly ingested without a prior processing step or after minimal culinary preparation. In an alternative embodiment, the plant biomass is processed and the material recovered after the processing step is ingested.

[00147] Processing methods preferably used in the present invention are methods commonly used in the food or feed industry. The final products of such methods still include a substantial amount of the expressed pharmaceutically active polynucleotide or polypeptide and are preferably conveniently eaten or drunk. The final product may also be mixed with other food or feed forms, such as salts, carriers, flavor enhancers, antibiotics, and the like, and consumed in solid, semi-solid, suspension, emulsion, or liquid form. In another preferred embodiment, such methods include a conservation step, such as, e.g., pasteurization, cooking, or addition of conservation and preservation agents. Any plant is used and processed in the present invention to produce edible or drinkable plant matter. The amount of pharmaceutically active polynucleotide or polypeptide expression product in a plant-derived preparation may be tested by methods standard in the art, e.g., gel electrophoresis, ELISA, or Western blot analysis, using a probe or antibody specific for the product. This determination may be used to standardize the amount of polynucleotide or protein ingested. For example, the amount

of therapeutically active product may be determined and regulated, for example, by mixing batches of product having different levels of product so that the quantity of material to be drunk or eaten to ingest a single dose can be standardized.

[00148] A pharmaceutically active polynucleotide or protein produced in a plant cell or tissue and eaten by a host is preferably absorbed by the digestive system. One advantage of the ingestion of plant tissue that has been only minimally processed is to provide encapsulation or sequestration of the polynucleotide or protein in cells of the plant. Thus, the product may receive at least some protection from digestion in the upper digestive tract before reaching the gut or intestine and a higher proportion of active product would be available for uptake.

[00149] The pharmaceutical compositions of the present invention can be administered therapeutically or prophylactically. In certain preferred embodiments, the compositions may be used to treat or prevent a disease. For example, any individual who suffers from a disease or who is at risk of developing a disease may be treated. It will be appreciated that an individual can be considered at risk for developing a disease without having been diagnosed with any symptoms of the disease. For example, if the individual has a particular genetic marker identified as being associated with increased risk for developing a particular disease, that individual will be considered at risk for developing the disease. Similarly, if members of an individual's family have been diagnosed with a particular disease, e.g., cancer, the individual may be considered to be at risk for developing that disease.

[00150] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butyleneglycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00151] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compositions of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active protein.

[00152] Dosage forms for topical or transdermal administration of a pharmaceutical composition of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active product, or preparation thereof, is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention.

Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a pharmaceutically active protein to the body. Such dosage forms can be made by suspending or dispensing the pharmaceutically active product in the proper medium. Absorption enhancers can also be used to increase the flux of the pharmaceutically active protein across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the pharmaceutically active protein in a polymer matrix or gel.

[00153] The compositions are administered in such amounts and for such time as is necessary to achieve the desired result. As described above, in certain embodiments of the present invention a "therapeutically effective amount" of a pharmaceutical composition is that amount effective for treating, attenuating, or preventing a disease in a host. Thus, the "amount effective to treat, attenuate, or prevent disease", as used herein, refers to a nontoxic but sufficient amount of the pharmaceutical composition to treat, attenuate, or prevent disease in any host. As but one example, the "therapeutically effective amount" can be an amount to treat, attenuate, or prevent diabetes, growth hormone deficiency, etc. As another example, the "therapeutically effective amount" can be an amount sufficient to cause an immune response in a subject, e.g., the production of antibodies that bind to a particular antigen. Preferably the antibodies protect against or reduce the severity of infection or protect against a disease or condition that may result from exposure to the antigen.

[00154] The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the stage of the disease, the particular pharmaceutical mixture, its mode of administration, and the like. The infected plants of the invention and/or protein preparations thereof are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form," as used herein, refers to a physically discrete unit of pharmaceutically active polynucleotide or polypeptide expression product appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compositions of the present invention is preferably decided by an attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient or organism may depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex of the patient, diet of the patient, pharmacokinetic condition of the patient, the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

[00155] It will also be appreciated that the pharmaceutical compositions of the present invention can be employed in combination therapies, that is, the pharmaceutical compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another anti-cancer agent), or they may achieve different effects.

Examples

Example 1: Construction of Recombinant Plant Virus Vectors

[00156] We employed vectors based on the Tobacco Mosaic Virus that are adapted

for insertion of a polynucleotide of interest to create a vector for use in generating clonal root lines, clonal root cell lines, clonal plant cell lines, and/or clonal plants that express a polynucleotide of interest according to the present invention. Figure 1 shows a schematic diagram of a TMV-based vector, D4, that was engineered to accept insertion of a polynucleotide of interest (Shivprasad *et al.*, *Virology*, 255(2):312-23, 1999), and illustrates insertion of various polynucleotides of interest into the vector. D4 contains a deletion of the TMV coat protein (CP) coding sequences but retains the TMV CP subgenomic promoter and the TMV 3' untranslated region (UTR), as indicated on the figure. The 126 and 183 kD proteins are required for TMV replication. The 30 kD protein is movement protein (MP), used for cell-to-cell movement. D4 contains *Pac* I and *Xho* I sites downstream of the CP subgenomic promoter, providing a site for convenient insertion of a polynucleotide of interest. Particular vectors created by inserting various polynucleotides of interest into D4 are described below.

[00157] D4C3GFP is a TMV-based expression vector that is deficient in CP production (Shivprasad *et al.*, 1999: TTT-GFP) as a result of deletion of the TMV CP coding region and its replacement with the C3GFP gene, which is placed under the control of the TMV CP subgenomic promoter. The C3GFP gene was recloned into D4 by overlapping PCR to eliminate the *Nco* I and *Xho* I sites in the C3GFP nucleotide sequence to facilitate further cloning steps. A polylinker *Pst* I-*Not* I-*Xho* I was introduced at the 3' end of C3GFP gene. The PCR product digested with *Pac* I-*Xho* I was cloned into D4 resulting in D4C3GFP.

[00158] The primers we used to modify the C3GFP gene and eliminate *Nco* I and *Xho* I sites are:

1) C3GFP.*Pac* I.For(N)

GGGAG.ATCTT.AATTA.ATGGC.TAGCA.AAGGA.GAAGA.A 36nt

2) C3GFP.*Xho* I.Rev(N)

CCCCT.CGAGC.GGCCG.CTGCA.GTTAT.TTGTA.GAGCT.CATCC.ATGCC 45nt

3) C3GFP.*Nco* I.For

GTTCC.CTGGC.CAACA.CTTGT.CAC 23nt

4) C3GFP.*Nco* I.Rev

TAGTG.ACAAG.TGTTG.GCCAG.GG 22nt

5) C3GFP.*Xho* I.For

GGACA.CAAAC.TGGAG.TACAA.CTATA 25nt

6) C3GFP.Xho1.Rev

AGTTA.TAGTT.GTACT.CCAGT.TTGTG 25nt

7) (BglII)-PacI

>AUG...HindIII...NcoI...NdeI...BsrGI...MluI...XhoI...BamHI...MfeI(MunI)...SalI
...SacI...TAA< PstI...NotI...XhoI

[00159] Three constructs that contained full-length or portions of the 3'-untranslated region (3' UTR) of AIMV RNA3 were also generated. In each of these constructs, sequences encoding C3GFP under control of the subgenomic TMV CP promoter were present upstream of AIMV RNA3 3'-UTR sequences (either full-length or a portion of the UTR), to allow us to precisely identify the sequences of the AIMV RNA3 3' UTR required for assembly and movement of TMV genomic RNA (either in *trans* or in *cis*). The RNA3 sequences were inserted between the *NotI* and *XhoI* sites of the new D4C3GFP vector as *NotI*-*SalI* fragments, resulting in the constructs SR25 (nts 1859-1941 of RNA3), SR26 (nts. 1859-1969 of RNA3) and SR27 (nts. 1859-2037 of RNA3, i.e., the entire 3' UTR) (Figure 11d). In addition to sequences from the AIMV RNA3 3' UTR, SR25, SR26, and SR27 also include sequences from the TMV 3' UTR (i.e., the UTR from the TMV genomic transcript) downstream of the inserted AIMV sequences. These sequences are TMV nucleotides 6192-6395, as in the D4 construct. The TMV-based viruses (SR25, SR26, and SR27) are defective in long-distance movement because the TMV coat protein is essential for effective phloem-mediated long distance transport and systemic infection of TMV.

[00160] The primers used to generate D4-based constructs with AIMV RNA3 3'-UTR sequences were:

1) SR -52 5' primer with *XhoI*-*PstI* sites at nt 1859 (plus sense)

5'-CCGCTCGAGCTGCAGTGTACCCCATTAATTTGG-3'

2) SR-53 3' primer at nt 1941 of AIMV RNA3 with *NotI*-*SalI* sites: minus sense

5'-CGGGTCGACGCGGCCGCGAATAGGACTTCATACCT-3'

3) SR-54 3' primer with *NotI*-*SalI* sites at nt 1969 of AIMV RNA3: minus sense

5'-CGGGTCGACGCGGCCGCAATATGAAGTCGATCCTA-3'

4) SR-55 3' primer with *NotI*-*SalI* sites at nt 2037 (minus sense)

5'-CGGGTCGACGCGGCCGCGCATCCCTTAGGGGCATT-3'.

[00161] Viral vectors in which polynucleotides of interest (e.g., GFP, hGH, GCSF) are inserted into SR25, SR26, and/or SR27 are in the process of being tested for generation of clonal root lines, clonal plant cell lines, and clonal plants as described herein.

[00162] To generate TMV-based constructs suitable for expression of human growth hormone (hGH) we inserted the gene for hGH into the D4 vector between the *Pac*I and *Xho*I sites. An AUG was introduced in the 5' primer used to amplify the gene from a plasmid, and the amino acids KDEL were introduced at the 3' end of the coding sequence in order to enhance translation due to retention in the ER. For the experiments described herein, hGH was cloned without its native leader sequence, resulting in D4-hGH, which was used in the experiments described herein.

[00163] Primer SR22 (5'-CCG TTAATTAATG TTC CCA ACT ATT CCA) was used to clone hGH without its leader, and introducing a *Pac*I site at the 5' end; primer SR23 (5'-CCG TTAATTAATG GCA ACT GGA TCA AGG) was used to clone hGH with its leader. Primer SR24 (5'-CGG CTC GAG TTA AAA ACC ACA TGA) was used to clone the hGH gene without KDEL and introducing a *Xho*I site at the 3' end; primer SR25 (5'-CGG CTC GAG TTC ATC TTT AAA ACC TGA TCC) was used to clone the gene with KDEL.

[00164] To generate TMV-based constructs suitable for expression of human granulocyte colony stimulating factor (GCSF), we first synthesized the entire open reading frame (ORF) encoding GCSF, without the signal peptide. The sequence of the synthesized gene was optimized for expression in plants. The ORF was synthesized with *Pac*I and *Xho*I sites at the 5 and 3' ends respectively. The gene was excised by *Pac*I / *Xho*I digestion and ligated into the D4 vector, which was linearized using *Pac*I and *Xho*I. The resulting vector (D4-GCSF) was used for the experiments described herein.

[00165] *Example 2: Generation and testing of clonal root lines expressing GFP*

[00166] Materials and Methods

[00167] *Synthesis of viral transcripts and viral infection.* *In vitro* transcripts of vector D4C3GFP, described above, which contains an open reading frame encoding GFP under control of the TMV CP subgenomic promoter, were synthesized using T7 polymerase. Approximately 10 µg of DNA was linearized with 30 units of *Kpn*I

overnight in a reaction volume of 100 μ l. Four μ l of the restriction digest was used to produce *in vitro* transcripts using the AmpliCap T7 High Yield message Maker Kit (Epicentre) according the manufacturers recommendations. Transcripts from one such reaction were used to infect six-week-old *Nicotiana benthamiana* plants by manually applying the transcripts dissolved in FES onto young, fully expanded leaves.

[00168] *Agrobacterium rhizogenes* stimulated root generation. *Agrobacterium rhizogenes* strain A4RSII was grown to OD₆₀₀ 0.8-1. Bacterial cells were pelleted and resuspended in MS-2 medium (MS salts, 2% sucrose, 10mM MES, pH 5.5) to a final OD₆₀₀ of 0.5. Acetosyringone was added to a final concentration 200 μ M 1 hour before transformation. Local or systemically infected leaves of *Nicotiana benthamiana* were harvested 5-14 days after inoculation with transcript. Leaves were surface sterilized for 6 min with 10% Clorox and washed several times with sterile distilled water.

[00169] Surface sterilized leaves of *N. benthamiana* were cut into pieces ~ 1 cm². They were dipped into bacterial suspension for 5 min, drained on filter paper and placed on the surface of solidified MS-2 medium. Plates were kept under dim light conditions at 24°C for 48 hours. After 48 hours the excess *Agrobacterial* suspension was removed, and leaf explants were placed on solid hormone free K₃ (Kao K.N. and Michayluk M.R., *Plants*, 115:355-367, 1974.) modified according to Nagy and Maliga, (Nagy J.J. and Maliga P., *Z.Pflanzenphysiol.* 78:453-455, 1976) and Menczel *et al.* (Menczel L., Nagy F., Kiss L.R. and Maliga P., *Theor. Appl. Genet.* 59:191-195, 1981) medium. Plates were maintained at 25°C with a 16 hr day/8 hr night light regime.

[00170] Three weeks after transformation, hairy roots were cut off and placed in a line on solid hormone free K₃ medium. Four to six days later, the most actively grown roots were isolated and transferred to liquid K₃ medium in individual Petri dishes. The roots were cultured on a rotary shaker at 24°C and subcultured $\sim$ weekly by dissecting and harvesting a portion of the root mass and transferring the harvested roots to a Petri dish containing fresh K₃ medium. Roots were screened for the presence of the protein of interest by Western blot analysis and/or by fluorescence under UV light, depending on the particular polynucleotide of interest.

[00171] *Western blot assays*: For Western blot assays 10 mg of fresh root material was placed into an Eppendorf tube and homogenized in 50 μ l of phosphate buffer, followed by the addition of 20 μ l of 5x loading buffer and 10 minutes of boiling. After

boiling, the homogenate was centrifuged for 5 to 10 minutes to clear the debris. Following centrifugation, 10 ul of sample was loaded on an SDS polyacrylamide gel, and proteins were separated by electrophoresis. Commercially available GFP protein (5 ng) (BD Biosciences Clontech) was loaded as a positive control. Leaf samples (10 mg) from *N. benthamiana* plants systemically infected with the same vector (D4C3GFP) were harvested at the time of peak expression, and an extract was prepared in an identical manner as described above for the root material and loaded on the gel for comparison with the root cell lines. Upon completion of electrophoresis proteins were electroblotted onto a nylon membrane, blocked using casein and reacted with GFP-specific antibodies (BD Biosciences Clontech). Proteins reacting with antibodies were visualized using a chemiluminescent substrate.

[00172] Results

[00173] Figures 6A – 6E show the overall method used for generating the clonal root lines (see Description). Figure 6G shows a photograph of a GFP-expressing clonal root line that was obtained by infecting *N. benthamiana* with viral vector D4C3GFP, harvesting leaf tissue from the infected region, infecting with *A. rhizogenes*, and culturing the pieces to allow development of hairy roots, which were then isolated and further cultured.

[00174] Figures 7A – 7C show Western blot analyses demonstrating GFP production in 3 clonal root lines derived from plant cells into which a viral vector whose genome contains a gene that encodes GFP under control of the TMV CP promoter (D4C3GFP) was introduced. Figure 7A shows GFP expression in the clonal root lines after 30 days of propagation in culture (i.e., 30 days after separation of the root from the leaf from which it was derived). Figure 7B shows GFP expression in the clonal root lines after 60 days of propagation in culture (i.e., 60 days after separation of the root from the leaf from which it was derived). C- represents control lanes containing no protein. MWM represents molecular weight markers. GFP-R represents samples from clonal root lines. GFP-P represents GFP isolated from leaf tissue of a plant infected with the same construct used for generation of the clonal root lines. Figure 7C is a control showing that the anti-GFP antibodies recognize commercially available GFP protein. These results demonstrate that the clonal root lines maintain high level expression of a protein

of interest (GFP) over an extended period of time, indicating the stability of the viral transcript in the clonal root lines.

[00175] Figures 8A and 8B show photographs of clonal root lines producing hGH (see Example 4) or GFP. Figure 8A shows a photograph of two clonal root lines taken under normal light conditions. The plate on the left shows a clonal root line derived from a plant cell into which a viral vector whose genome contains a gene that encodes human growth hormone (hGH) under control of the TMV CP promoter was introduced. The plate on the right shows a clonal root line derived from a plant cell into which a viral vector whose genome contains a gene that encodes green fluorescent protein (GFP) under control of the TMV CP promoter was introduced. Figure 8B shows a photograph of the same clonal root lines as shown in Figure 8A taken under UV light, demonstrating expression of GFP. These results demonstrate robust expression of GFP in the root mass and illustrate the convenience of fluorescence-based screening for expression of a polynucleotide of interest.

[00176] It is noted that Western analysis demonstrated expression of GFP throughout all portions of the root mass. However, when screened using a visual approach, expression generally appears stronger in the more mature portions of the root mass than in the growing tips, where cell division is proceeding rapidly. This appears to be due both to the time required for new cell to synthesize sufficient GFP for visibility and to the fact that when viewed from above, one is looking through multiple layers of cells in the thicker portion of the roots. It is also noted that the most mature portions of the roots may become somewhat “woody”, which can obscure visual detection of GFP.

[00177] *Example 3: Generation and testing of clonal root lines expressing hGH*

[00178] *N. benthamiana* plants were inoculated with a TMV-based vector, D4-hGH, containing an open reading frame encoding hGH under control of the TMV CP subgenomic promoter. Hairy roots were obtained and subcultured essentially as described in Example 2. Two weeks after separation from leaf discs, during the third round of subculture, the segments of roots were analyzed for hGH expression by Western blot assay (Figure 9) essentially as described in Example 2. Five ng hGH protein (Research Diagnostics) was used as a control in all Western blots in which expression of hGH was tested. Anti-hGH antibodies were from Research Diagnostics.

As can be seen from Figure 9, up to 80 % of the clonal root lines had detectable levels of hGH. We selected the highest producers and propagated them further. After 10 passages (subculturings), samples were taken and analyzed for hGH accumulation. Figure 10 shows a Western blot, demonstrating that the clonal root lines maintained stable expression of hGH after 10 passages in which hGH expression in selected lines was several fold higher (250 ug/gram fresh root tissue) than that in leaves infected with the same virus construct (70 ug/gram fresh leaf tissue) when compared by Western blot.

[00179] Figures 8A and 8B show photographs of clonal root lines producing hGH and GFP. *Figure 8A* shows a photograph of two clonal root lines taken under normal light conditions. The plate on the left shows a clonal root line derived from a plant cell into which a viral vector whose genome contains a gene that encodes human growth hormone (hGH) under control of the TMV CP promoter was introduced. The plate on the right shows a clonal root line derived from a plant cell into which a viral vector whose genome contains a gene that encodes green fluorescent protein (GFP) under control of the TMV CP promoter was introduced. *Figure 8B* shows a photograph of the same clonal root lines as shown in Figure 8A taken under UV light, demonstrating expression of GFP.

[00180] *Example 4: Generation and testing of clonal root lines expressing GCSF*

[00181] *N. benthamiana* plants were inoculated with a TMV-based vector, D4-GCSF, containing an open reading frame encoding GCSF under control of the TMV CP subgenomic promoter, and hairy roots were obtained essentially as described in Example 2. Two weeks after separation from leaf discs the segments of roots were analyzed for GCSF expression by Western blot assay (Figure 11). As can be seen from Figure 11, up to 80 % of the clonal root lines had detectable levels of GCSF. We selected the highest producers and propagated them further. After 10 passages (in which portions of the root mass were harvested and transferred to new Petri dishes containing fresh medium) samples were taken and analyzed for GCSF accumulation. Figure 12 shows a Western blot, demonstrating that the clonal root lines maintained stable expression of GCSF after 10 passages (subculturings). Five ng recombinant GCSF produced using an *E. coli* expression system was used as a control in all Westerns in which expression of GCSF was tested. Anti-GCSF antibodies were from Oncogene Science.

[00182] *Example 5: Generation and testing of clonal plant cell lines expressing GCSF*

[00183] *Cell culture and electroporation.* Cell lines derived from *Nicotiana tabacum* cv Bright yellow (BY-2) were maintained in MS medium (Murashige T. and Skoog F., *Physiol. Plant.* 15:473-497, 1962) supplemented with 0.2 mg/l 2,4-D and 0.1 mg/l Kinetin, 20mM MES, pH 5.6-5.8 on a shaker, 140 rpm at 25°C, and subcultured weekly. For electroporation, protoplasts were generated from cells that had been subcultured for 3 - 4 days. Cells were spun at 1000 rpm for 8 min, washed 2x with Mannitol 0.4M and MES 20mM, pH 5.5. Cells were then taken to 30-50 ml with filter sterilized protoplasting solution: 0.4M mannitol, MES 20mM, pH5.5, Cellulase Onozuka RS (Yakult Honsha Co.) 1%, Pectolyase Y23 (Seishin Pharmaceutical Co.) 0.1%. Cells were incubated in 250 ml flasks at 25°C for 20-25 min. The protoplast solution was filtered through a 100/ μ m sieve, spun at 700 rpm for 6 min, and washed 2x with ice-cold 0.4M Mannitol. Protoplasts were counted using a hemacytometer and resuspended in electroporation buffer: 10mM HEPES, 150 mM NaCl. 5 mM CaCl_2 , 0.4M mannitol, pH 7.2 to a final concentration 1×10^6 protoplasts/ml.

[00184] Transcript (25-30 μ l) was placed into an electroporation cuvette, 0.4 cm (Biorad) kept on ice, and after 10-15 min was mixed with 0.5 ml of protoplast suspension by Pasteur pipette and immediately used to electroporate cells.

Electroporation was performed using a Biorad Gene Pulser at 250 volts and 175 capacitance. Electroporated protoplasts were resuspended in 8 ml of PBS buffer containing .4 M mannitol and maintained for formation of the cell wall.

[00185] *Enrichment for stable producer cell lines.* Within 4 - 5 days following electroporation, dividing cells were diluted and sampled (10 μ l of infected cells into 100 μ l of medium) to enrich for cells that expressed the polynucleotide of interest (target molecule) at high levels. The diluted cells were spotted onto individual sections of a Petri dish, as shown in Figure 13E. Two to three weeks later each sample was tested by visual or other means (e.g., Western blot) for the presence of target molecule (e.g., GFP, GCSF, hGH, etc.). Stably infected cells producing target molecule were selected for further enrichment until producer cell line is obtained.

[00186] Results

[00187] Clonal plant cell lines were derived by introducing a TMV-based viral vector containing an open reading frame that encodes GCSF under control of the TMV CP subgenomic promoter into BY-2 cells. The overall process is shown in Figure 13. Enrichment for cells that express GCSF was performed using Western blot assays until populations of cells (either single clonal cell lines or populations containing several clonal cell lines) were obtained. *Figure 14* shows Western blot analyses demonstrating GCSF production in a plant cell population derived from plant cells into which a viral vector whose genome contains a gene that encodes GCSF under control of the TMV CP promoter was introduced. It is noted that the enriched plant cell population may contain either a single clonal cell line or multiple lines. Further enrichment, using more dilute samples, would result in clonal cell lines. *Figure 14A* shows a Western blot performed 48 hours after introduction of the vector. *Figure 14B* shows a Western blot performed using the same cell populations as shown in *Figure 14A* performed after further maintaining the cells in culture, i.e., 57 days after inoculation. GCSF-COM indicates a lane in which recombinant GCSF protein was loaded as a positive control. MWM indicates molecular weight markers. C- indicates a lane in which plant extract made from plants not expressing GCSF was loaded.

[00188] *Example 6: Generation and testing of clonal cell lines expressing GFP*

[00189] Results

[00190] Clonal plant cell lines were derived by introducing a TMV-based viral vector containing an open reading frame that encodes GFP under control of the TMV CP subgenomic promoter (D4C3GFP) into BY-2 cells essentially as described in Example 5. Enrichment for cells that express GFP was performed using a visual screen for fluorescence until populations of cells (either single clonal cell lines or populations containing several clonal cell lines) that stably express GFP were obtained. Figure 13C shows a protoplast suspension containing cells into which the viral vector was introduced. Figure 13E shows diluted samples from the suspension plated in individual droplets on Petri dishes. Figure 13F shows the same Petri dishes as in Figure 13E under UV light. Clonal plant cell lines expressing GFP are readily seen. It is noted that the droplets may contain either a single clonal plant cell line or multiple clonal plant cell lines. Single clonal plant cell lines (i.e, populations derived from a single ancestral

cell) can be generated by further limiting dilution using standard methods for single cell cloning.

[00191] *Figure 15* shows GFP production in plant cell lines derived from plant cells into which D4C3GFP. *Figure 15A* shows enrichment for plant cell lines that express GFP. *Figure 15B* shows a callus obtained from a clonal plant cell line that contains a similar viral vector that does not encode GFP. The photographs were taken 3 months after the vector was introduced into the cells from which the clones in *Figure 15A* were derived. Both photographs were taken under UV light.

[00192] *Example 7: Generation and testing of a clonal plant*

[00193] Clonal root lines expressing hGH were obtained as described in Example 3. Root cells were isolated by enzymatic digestion and cultured as described in Peres et al., *Plant Cell, Tissue, and Organ Culture* 65, 37-44, 2001, to generate clonal plants. *Figure 16A* shows a plant that was obtained from a clonal root line. To determine whether the plant contained the viral vector, a small leaf sample was used to inoculate a tobacco variety that is a sensitive host for formation of local lesions upon viral infection. Formation of lesions within 2 days of inoculation, as indicated by arrows in *Figure 16B*, indicated that the clonal plant regenerated from the clonal root line maintains active viral replication, strongly suggesting that the clonal plant also expresses hGH. Additional experiments showed that this was indeed the case (data not shown).

Equivalents

[00194] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the following claims.

[00195] We claim:

1. A clonal entity derived from a plant or portion thereof, wherein cells of the clonal entity comprise a viral vector that comprises a polynucleotide of interest.
2. The clonal entity of claim 1, wherein the clonal entity is a clonal root line.
3. The clonal root line of claim 2, wherein the viral vector is derived from TMV or AIMV.
4. The clonal root line of claim 2, wherein the polynucleotide of interest is operably linked to a CP promoter, an MP promoter, an inducible promoter, or a trans-activatable promoter.
5. The clonal root line of claim 2, wherein the viral vector is capable of self-replication.
6. The clonal entity of claim 1, wherein the clonal entity is a clonal root cell line.
7. The clonal root cell line of claim 6, wherein the viral vector is derived from TMV or AIMV.
8. The clonal root cell line of claim 6, wherein the polynucleotide of interest is operably linked to a CP promoter, an MP promoter, an inducible promoter, or a trans-activatable promoter.
9. The clonal root cell line of claim 6, wherein the viral vector is capable of self-replication.
10. The clonal entity of claim 1, wherein the clonal entity is a clonal plant cell line.
11. The clonal plant cell line of claim 10, wherein the viral vector is derived from TMV or AIMV.
12. The clonal plant cell line of claim 10, wherein the polynucleotide of interest is operably linked to a CP promoter, an MP promoter, an inducible promoter, or a trans-activatable promoter.

13. The clonal plant cell line of claim 10, wherein the viral vector is capable of self-replication.
14. The clonal entity of claim 1, wherein the clonal entity is a clonal plant.
15. The clonal plant of claim 14, wherein the viral vector is derived from TMV or AIMV.
16. The clonal plant of claim 14, wherein the polynucleotide of interest is operably linked to a CP promoter, an MP promoter, an inducible promoter, or a trans-activatable promoter.
17. The clonal plant of claim 14, wherein the viral vector is capable of self-replication.
18. A method of obtaining a clonal entity that expresses a polynucleotide of interest comprising steps of:
 - (i) introducing a viral vector that comprises a polynucleotide of interest into a plant or portion thereof; and
 - (ii) generating a clonal entity from the plant.
19. The method of claim 18, wherein the generating step comprises introducing a viral vector that comprises a polynucleotide of interest into a plant or portion thereof.
20. The method of claim 18, wherein the generating step comprises introducing an Ri T-DNA or portion thereof sufficient to generate hairy roots into the plant or portion thereof.
21. A method of producing a polynucleotide or polypeptide comprising steps of:
 - (i) generating a clonal entity derived from a plant, cells of which comprise a viral vector comprising a polynucleotide of interest;
 - (ii) maintaining the clonal entity in culture if the clonal entity is a clonal root line, clonal root cell line, or clonal plant cell line, or growing the clonal entity if the clonal entity is a clonal plant;
 - (iii) harvesting cells or culture medium if the if the clonal entity is a clonal root line, clonal root cell line, or clonal plant cell line or harvesting plant tissue if the clonal entity is a clonal plant; and

- (iv) isolating or purifying the polynucleotide or polypeptide from the harvested cells, culture medium, or plant tissue.
22. A method of obtaining a clonal root line that expresses a polynucleotide of interest comprising steps of:
- (i) introducing a viral vector that comprises a polynucleotide of interest into a plant or portion thereof; and
 - (ii) generating one or more clonal root lines from the plant.
23. The method of claim 22, wherein the step of generating comprises:
contacting the plant or a portion thereof with an *Agrobacterium*.
24. The method of claim 23, wherein the *Agrobacterium* is a causative agent of hairy root disease.
25. The method of claim 23, wherein the *Agrobacterium* is *Agrobacterium rhizogenes*.
26. The method of claim 1, wherein the step of generating comprises introducing an Ri T-DNA or portion thereof sufficient to generate hairy roots into the plant or portion thereof.
27. The method of claim 22, wherein the step of generating comprises exposing the plant or plant portion to hormonal stimulation.
28. The method of claim 22, wherein the viral vector is derived from TMV.
29. The method of claim 22, wherein the viral vector is derived from ALMV.
30. The method of claim 22, wherein the viral vector lacks sequences coding for a functional coat protein, functional movement protein, or both.
31. The method of claim 22, wherein the polynucleotide is operably linked to a CP promoter.
32. The method of claim 22, wherein the polynucleotide is operably linked to an MP promoter.

33. The method of claim 22, wherein the polynucleotide is operably linked to an inducible promoter.
34. The method of claim 22, wherein the step of introducing is performed prior to the step of generating.
35. The method of claim 22, wherein the viral vector or a portion thereof does not become integrated into host cell DNA.
36. The method of claim 22, wherein the viral vector is capable of self-replication.
37. The method of claim 22, wherein the viral vector comprises a polynucleotide that encodes a detectable or selectable marker.
38. The method of claim 1, further comprising the step of:
maintaining the plant for a period of time following introduction of the viral vector and prior to generating the clonal root line to allow replication of the viral vector.
39. The method of claim 22, further comprising the step of:
screening one or more clonal root lines derived from the plant for expression of the polynucleotide of interest.
40. A clonal root line obtained according to the method of claim 39.
41. A clonal root line, cells of which contain a viral vector comprising a polynucleotide of interest.
42. The clonal root line of claim 41, wherein at least 50% of the cells contain the viral vector.
43. The clonal root line of claim 41, wherein substantially all root cells contain the viral vector.
44. A method of generating a clonal root cell line that expresses a polynucleotide of interest comprising steps of:

- (i) generating a clonal root line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest;
 - (ii) releasing individual cells from the clonal root; and
 - (iii) maintaining the cells in culture under conditions appropriate for root cell proliferation.
45. A clonal root cell line obtained according to the method of claim 44.
46. A clonal root cell line, cells of which contain a viral vector comprising a polynucleotide of interest.
47. The clonal root cell line of claim 46, wherein at least 50% of the cells contain the viral vector.
48. The clonal root cell line of claim 46, wherein substantially all root cells contain the viral vector.
49. A method of generating a clonal plant cell line that expresses a polynucleotide of interest comprising steps of:
- (i) generating a clonal root line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest;
 - (ii) releasing individual cells from the clonal root; and
 - (iii) maintaining the cells in culture under conditions appropriate for plant cell proliferation.
50. A clonal plant cell line generated according to the method of claim 49.
51. A method of generating a clonal plant that expresses a polynucleotide of interest comprising steps of:
- (i) generating a clonal root line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest;
 - (ii) releasing individual cells from the clonal root; and
 - (iii) maintaining the cells under conditions appropriate for formation of a plant.

52. A clonal plant generated according to the method of claim 51.
53. A clonal plant, cells of which comprise a viral vector comprising a polynucleotide of interest.
54. The clonal plant of claim 53, wherein the viral vector is derived from TMV.
55. The clonal plant of claim 53, wherein the viral vector is derived from AIMV.
56. The clonal plant of claim 53, wherein the viral vector lacks sequences coding for a functional coat protein, functional movement protein, or both.
57. The clonal plant of claim 53, wherein the polynucleotide is operably linked to a CP promoter.
58. The clonal plant of claim 53, wherein the polynucleotide is operably linked to an MP promoter.
59. The clonal plant of claim 53, wherein the polynucleotide is operably linked to an inducible promoter.
60. A method of generating a clonal plant cell line that expresses a polynucleotide of interest comprising steps of:
- (i) introducing a viral vector that comprises a polynucleotide of interest into cells of a plant cell line maintained in culture; and
 - (ii) enriching for cells that contain the viral vector.
61. The method of claim 60, wherein the enriching step comprises:
- (i) removing a portion of the cells from the culture;
 - (ii) diluting the removed cells so as to reduce the cell concentration;
 - (iii) allowing the diluted cells to proliferate; and
 - (iv) screening for cells that contain the viral vector.
62. The method of claim 60, wherein the enriching step comprises:

determining whether cells removed from the culture express the polynucleotide of interest.

63. A clonal plant cell line generated according to the method of claim 60.
64. A clonal plant cell line, cells of which comprise a viral vector comprising a polynucleotide of interest.
65. The clonal plant cell line of claim 64, wherein the viral vector is derived from TMV.
66. The clonal plant cell line of claim 64, wherein the viral vector is derived from AIMV.
67. The clonal plant cell line of claim 64, wherein the viral vector lacks sequences coding for a functional coat protein, functional movement protein, or both.
68. The clonal plant cell line of claim 64, wherein the polynucleotide is operably linked to a CP promoter.
69. The clonal plant cell line of claim 64, wherein the polynucleotide is operably linked to an MP promoter.
70. The clonal plant cell line of claim 64, wherein the polynucleotide is operably linked to an inducible promoter.
71. A clonal plant generated from the clonal plant cell line of claim 64.
72. A method of generating a clonal plant that expresses a polynucleotide of interest comprising steps of:
 - (i) generating a clonal plant cell line, cells of which contain a viral vector whose genome comprises a polynucleotide of interest; and
 - (ii) maintaining the cells under conditions appropriate for formation of a plant.
73. A clonal plant generated according to the method of claim 72.
74. A method of producing a polypeptide of interest comprising steps of:

- (i) generating a clonal root line, cells of which comprise a viral vector comprising a polynucleotide that encodes a polypeptide of interest;
 - (ii) maintaining the clonal root line in culture;
 - (iii) harvesting root tissue or culture medium from the clonal root line; and
 - (iv) isolating or purifying the polypeptide of interest from the harvested root tissue or culture medium.
75. A method of producing a polypeptide of interest comprising steps of:
- (i) generating a clonal root cell line, cells of which comprise a viral vector comprising a polynucleotide that encodes a polypeptide of interest;
 - (ii) maintaining the clonal root cell line in culture;
 - (iii) harvesting cells or culture medium from the clonal root line; and
 - (iv) isolating or purifying the polypeptide of interest from the harvested root cells or culture medium.
76. A method of producing a polypeptide of interest comprising steps of:
- (i) generating a clonal plant cell line, cells of which comprise a viral vector comprising a polynucleotide that encodes a polypeptide of interest;
 - (ii) maintaining the clonal plant cell line in culture;
 - (iii) harvesting cells or culture medium from the clonal plant line; and
 - (iv) isolating or purifying the polypeptide of interest from the harvested plant cells or culture medium.
77. A method of producing a polypeptide of interest comprising steps of:
- (i) generating a clonal plant, cells of which comprise a viral vector comprising a polynucleotide that encodes a polypeptide of interest;
 - (ii) growing the clonal plant;
 - (iii) harvesting plant tissue from the clonal plant; and
 - (iv) isolating or purifying the polypeptide of interest from the harvested plant tissue.
78. A method of expressing a polynucleotide of interest in a root, root cell, plant cell line, or plant comprising steps of:

- (i) generating a clonal root, root cell, plant cell line, or plant, cells of which comprise a viral vector comprising a polynucleotide of interest;
- (ii) maintaining the clonal root, root cell, plant cell line, or plant under conditions suitable for expression of the polynucleotide of interest.

1/18

Figure 1





Figure 2



3/18




Figure 4




Figure 5




Figure 6
Clonal Root Line Technology (CRLT) for
Obtaining Target Producer Lines





Figure 7
Western blot analysis of GFP production in *clonal root line*




GFP-specific antibodies

8/18

Figure 8
Clonal Root Lines Producing hGH and GFP




Under normal light conditions



Same plates under UV light
10th subculture for GFP 53 (5 month
after separation from leaf discs)

Figure 9
Screening and Selection of hGH Producing *Clonal Root Lines*
by Western Blot Analysis



hGH-specific antibodies

→ Indicates selected lines
3d subculture

Figure 10
Western Blot Analysis of hGH Production in Selected *Clonal Root Lines*






Figure 11
Screening and Selection of GCSF Producing Clonal Root Lines
by Western Blot Analysis



GCSF-specific antibodies

→ Indicates selected lines
3d subculture

Figure 12
Western Blot Analysis of GCSF Production in Selected *Clonal Root Lines*



GCSF-specific antibodies

Figure 13
Clonal Cell Line Technology (CCLT) for Obtaining
Target Producer Lines






Figure 14
Western blot analysis of GCSF production in cell lines



GCSF-specific antibodies


15/18

Figure 15
Clonal cell lines producing GFP



Enriched clones under UV light
(3 month after infection)

Figure 16
Regeneration of clonal plant lines from clonal root lines



17/18




Figure 17

18/18

Figure 18

