




(86) Date de dépôt PCT/PCT Filing Date: 2007/07/19
(87) Date publication PCT/PCT Publication Date: 2008/01/24
(45) Date de délivrance/Issue Date: 2014/11/18
(85) Entrée phase nationale/National Entry: 2009/01/14
(86) N° demande PCT/PCT Application No.: US 2007/016341
(87) N° publication PCT/PCT Publication No.: 2008/011087
(30) Priorité/Priority: 2006/07/19 (US60/831,843)

(51) Cl.Int./Int.Cl. *A61K 47/24* (2006.01),
A61K 31/606 (2006.01), *A61K 47/44* (2006.01),
A61K 9/127 (2006.01), *A61K 9/16* (2006.01),
A61K 9/50 (2006.01)
(72) Inventeur/Inventor:
LICHTENBERGER, LENARD M., US
(73) Propriétaire/Owner:
THE BOARD OF REGENTS OF THE UNIVERSITY OF
TEXAS SYSTEM, US
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : PREPARATIONS DE PHOSPHOLIPIDES ET DE SUBSTANCES PHARMACEUTIQUES CONTENANT DE
L'ACIDE SALICYLIQUE 5-AMINE DESTINEES A TRAITER UNE MALADIE INTESTINALE INFLAMMATOIRE
(54) Title: PREPARATIONS OF PHOSPHOLIPIDS AND PHARMACEUTICALS CONTAINING 5-AMINO SALICYLIC ACID
FOR THE TREATMENT OF INFLAMMATORY BOWEL DISEASE



(57) **Abrégé/Abstract:**

A unique composition of a 5-amino salicylic acid (5 ASA) and a phospholipid is disclosed for treating Inflammatory Bowel Disease (IBD), where the composition can be a mixture, a molecular association complex or a covalent compound of 5ASA and a reactive phospholipid covalently bonded together via a diazo linkage and to methods for administering the compositions to treat symptoms of IBD.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
24 January 2008 (24.01.2008)

PCT

(10) International Publication Number
WO 2008/011087 A3

(51) International Patent Classification:

A61K 47/24 (2006.01) A61K 9/50 (2006.01)
A61K 47/44 (2006.01) A61K 31/606 (2006.01)
A61K 9/16 (2006.01) A61K 9/127 (2006.01)

(21) International Application Number:

PCT/US2007/016341

(22) International Filing Date: 19 July 2007 (19.07.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/831,843 19 July 2006 (19.07.2006) US

(71) Applicant (for all designated States except US): **THE BOARD OF REGENTS OF THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 W. 7th Street, Suite 820, Austin, TX 78701 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **LICHTENBERGER, Lenard, M.** [US/US]; 5427 Darnell, Houston, TX 77096 (US).(74) Agents: **CHWANG, T. Ling** et al.; Jackson Walker L.L.P., 901 Main Street, Suite 6000, Dallas, TX 75202 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

31 July 2008

(54) Title: PREPARATIONS OF PHOSPHOLIPIDS AND PHARMACEUTICALS CONTAINING 5-AMINO SALICYLIC ACID FOR THE TREATMENT OF INFLAMMATORY BOWEL DISEASE



(57) Abstract: A unique composition of a 5-amino salicylic acid (5 ASA) and a phospholipid is disclosed for treating Inflammatory Bowel Disease (IBD), where the composition can be a mixture, a molecular association complex or a covalent compound of 5ASA and a reactive phospholipid covalently bonded together via a diazo linkage and to methods for administering the compositions to treat symptoms of IBD.

WO 2008/011087 A3

- 1 -

**PREPARATIONS OF PHOSPHOLIPIDS AND PHARMACEUTICALS
CONTAINING 5-AMINO SALICYLIC ACID FOR THE TREATMENT OF
INFLAMMATORY BOWEL DISEASE**

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0002] The present invention relates to unique formulations for treating inflammatory bowel disease (IBD), such as ulcerative colitis, including 5-amino salicylic acid (5ASA) and a phospholipid.

[0003] The present invention relates to unique compositions of matter including 5ASA and phospholipid adapted to release 5ASA in a distal small intestine and/or colon, where lesions due to colitis or Inflammatory Bowel Disease (IBD) are present in order to enhance the anti-inflammatory efficacy of 5ASA. In certain embodiment, the phospholipid composition including from about 15wt.% to about 95wt.% of phosphatidylcholine (PC) in a bio-compatible carrier and/or resin. The present invention also relates to methods for making and administering the compositions of this invention orally, enterally and/or rectally.

2. Description of the Related Art

[0004] For a background of phospholipids and anti-inflammatory pharmaceuticals the reader is directed to United States Pat. Nos.: 4,918,063; 5,043,329; 4,950,656; 5,032,585; 5,763,422; and 5,955,451 and PCT/US01/51605.

[0005] Inflammatory Bowel Disease (IBD) represents a family of ulcerative diseases including Ulcerative Colitis and Crohn's Disease that affect the colon and distal small bowel. These diseases are manifested by episodes of GI bleeding, diarrhea, fever, infection and in the most advanced cases, GI fistulation are manifested by perforation and cancer. 5ASA has been used medically to manage patients with IBD, though its effectiveness to keep patients in remission has been limited.

[0006] Although 5ASA is an effective treatment, there is a need in the art for 5ASA preparations having improved efficacy to treat IBD and to keep patients in remission.

SUMMARY OF THE INVENTION

Certain exemplary embodiments provide the use of a 5-aminosalicylic acid (5ASA) containing component and a phospholipid-containing oil in the manufacture of a medicament for treating inflammatory bowel disease (IBD) characterized in that the phospholipid-containing oil is present in an amount to enhance the anti-inflammatory efficacy of the 5ASA.

Other exemplary embodiments provide the use of a 5-aminosalicylic acid (5ASA) containing component and a phospholipid or a phospholipid-containing oil in the manufacture of a non-liposomal medicament for treating inflammatory bowel disease (IBD) characterized in that the phospholipid or a phospholipid-containing oil is present in an amount to enhance the anti-inflammatory efficacy of the 5ASA.

Other exemplary embodiments provide a composition comprising 5-aminosalicylic acid (5ASA) and a phospholipid associated at a molecular level at a weight ratio of 5ASA to phospholipid of between 1:10 and 10:1 to form a complex.

Other exemplary embodiments provide a composition comprising a covalent compound including 5ASA and an active phospholipid, wherein the reactive phospholipid is covalently bonded to the 5ASA via a diazo linkage between the amino group of 5ASA and the reactive phospholipid, said linkage being cleavable by bacteria in the distal gut that catalyze the hydrolysis of the diazo linkage.

-2a-

[0007] The present invention provides compositions including a phospholipid (PL) and 5-amino salicylic acid (5ASA).

[0008] The present invention also provides compositions including a soy derived phospholipid component and a 5-amino salicylic acid (5ASA) containing component such as Mesalamine, Sulfasalazine, Olsalazine, Balsalazide, or mixtures thereof. The compositions can also include additional components such as a resin (*e.g.*, Eudragit®S) to facilitate release of the 5ASA-containing component in the distal gut.

[0009] The present invention also provides compositions including a soy derived phosphatidylcholine (PC) component and a 5-amino salicylic acid (5ASA) containing component such as Mesalamine, Sulfasalazine, Olsalazine, Balsalazide, or mixtures thereof, where the PC and 5ASA-containing components are surrounded or encapsulated or embedded in a resin (*e.g.*, Eudragit®S) to facilitate the release the components in the distal gut.

[0010] The present invention provides a method for making formulations including a 5ASA component and a phospholipid component.

[0011] The present invention also provides a method for making the formulations including a 5ASA and a PC enriched phospholipid component.

[0012] The present invention provides methods for administering 5ASA/PL formulations orally, enterally or rectally, where the administration can be a single administration, a periodic administration, an intermittent administration, or administration according to any administration protocol (administration according to a prescribed schedule, *e.g.*, one daily, twice daily, *etc.*).

[0013] The present invention provides methods of treating patients with IBD by administering a composition of this invention directly on a site of IBD injury to reduce inflammation, while: (1) reducing or preventing ulceration of the injury, (2) reducing or preventing further ulceration of the injury, (3) reducing or healing ulceration of the injury, (4) or to maintain an integrity of hydrophobic membranes and/or layers associated with the distal bowels.

-3-

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] The invention can be better understood with reference to the following detailed description together with the appended illustrative drawings in which like elements are numbered the same.

[0015] **Figure 1** depicts a graph demonstrating that placing mice on drinking water containing increasing concentrations (0% to 4%) of dextran sodium sulfate (DSS) for 4 days results in increased colonic inflammation as indicated by increased colon tissue weight per unit length.

[0016] **Figure 2** depicts that increasing concentrations of DSS in the drinking water over a 4 day period induces a dose-dependent increase in GI bleeding of mice as measured by fecal hemoglobin concentration.

[0017] **Figure 3** demonstrates the food consumption of mice eating diets containing 5ASA alone (100 mg/kg body weight) or 5ASA in combination on a 1:1 weight basis with three different soy PC preparations, P35, P53 or P75, containing 35%, 53% or 75% PC, respectively. All but one group labeled Sal/Sal had access to drinking water containing 3% DSS to induce colitis.

[0018] **Figure 4** demonstrates changes in colonic inflammation, as indicated by increased colon tissue weight per unit length in mice fed the experimental diets and placed on 3% DSS in the drinking water to induce colitis.

[0019] **Figure 5** demonstrates changes in fecal blood loss in mice fed the experimental diets and placed on 3% DSS in the drinking water to induce colitis.

[0020] **Figure 6** demonstrates changes in colonic myeloperoxidase (MPO) in mice fed the experimental diets and placed on 3% DSS in the drinking water to induce colitis.

[0021] **Figure 7** demonstrates changes in colonic mucosal histology, using a score from 0 (normal) to 4 (complete absence of mucosal glandular structure) in mice fed the experimental diets and placed on 3% DSS in the drinking water to induce colitis.

[0022] **Figure 8** depicts changes in hematocrit levels in Sprague Dawley male rats during a 6 day recovery period after DSS induced colitis for saline, 5ASA and two PC:5ASA formulations.

[0023] **Figure 9** depicts changes in fecal blood loss in Sprague Dawley male rats during the 6 day recovery period after DSS induced colitis for saline, 5ASA and two PC:5ASA formulations.

-4-

[0024] Figure 10 depicts changes in colonic tissue edema in Sprague Dawley male rats during the 6 day recovery period after DSS induced colitis for saline, 5ASA and two PC:5ASA formulations.

[0025] Figure 11 depicts changes in colonic MPO activity in Sprague Dawley male rats during the 6 day recovery period after DSS induced colitis for saline, 5ASA and two PC:5ASA formulations.

[0026] Figure 12 depicts changes in the number of inflammatory cells in lamina propria of colonic submucosa in Sprague Dawley male rats during the 6 day recovery period after DSS induced colitis for saline, 5ASA and two PC:5ASA formulations.

[0027] Figure 13 depicts changes in colonic mucosal injury in Sprague Dawley male rats as assessed histologically during the 6 day recovery period after DSS induced colitis for saline, 5ASA and two PC:5ASA formulations.

[0028] Figure 14 demonstrates changes in colon tissue weight per unit length in Sprague Dawley male rats during a recovery period after DSS induced colitis for saline with no DSS in the drinking water and saline, 5ASA and a PC:5ASA formulation with DSS in the drinking water.

[0029] Figure 15 depicts changes in colonic MPO activity in Sprague Dawley male rats during a recovery period after DSS induced colitis for saline with no DSS in the drinking water and saline, 5ASA and a PC:5ASA formulation with DSS in the drinking water.

[0030] Figure 16 depicts changes in colon contact angles in Sprague Dawley male rats during a recovery period after DSS induced colitis for saline with no DSS in the drinking water and saline, 5ASA and a PC:5ASA formulation with DSS in the drinking water.

[0031] Figure 17 depicts changes in hematocrit levels in Sprague Dawley male rats during a recovery period after DSS induced colitis for saline with no DSS in the drinking water and saline, 5ASA and a PC:5ASA formulation with DSS in the drinking water.

[0032] Figure 18 depicts changes in fecal hemoglobin in Sprague Dawley male rats during a recovery period after DSS induced colitis for saline with no DSS in the drinking water and saline, 5ASA and a PC:5ASA formulation with DSS in the drinking water.

-5-

DETAILED DESCRIPTION OF THE INVENTION

[0033] The inventor has found that compositions of 5-amino salicylic acid (5ASA) and 5ASA based pharmaceuticals and phospholipids have improved activity to reduce colitis in rodent using rodent colitis models. In certain embodiments, the phospholipids are phospholipids derived from soy bean and other sources. In other embodiments, the phospholipids are phospholipids enriched in phosphatidylcholine (PC) or a derivative of phosphatidylcholine. These 5ASA/PC formulations are capable of being administered orally, enterally, or rectally (enema) for treatment or amelioration of GI inflammation, ulceration, bleeding and other symptoms associated with inflammatory bowel disease (IBD), and it's associated sequelae of diarrhea, fever and pain. The compositions of the present invention including 5ASA and a phospholipid enhance anti-inflammatory benefits of the 5ASA and fortify hydrophobic barrier properties of affected mucosa, which are attenuated in IBD.

[0034] The invention relates to a method for preparing a composition including a phospholipid, in certain embodiments a phospholipid enriched in phosphatidylcholine (PC), and 5ASA or pharmaceutical formulations including 5ASA for the treatment of IBD. These formulations can be prepared by taking a powder containing 5ASA (*e.g.*, Mesalamine, Sulfasalazine, Olsalazine, Balsalazide, or mixtures thereof) or a larger molecule covalently bound to active 5ASA via a diazo bond and simply mixing it with a phospholipid. In certain embodiments, the 5ASA containing component is simply mixed with an oil-based phospholipid component such as a soy lecithin oil containing PC, *e.g.*, Phosphal 35 SB (P35). Generally, the weight ratio of 5ASA to phospholipid is between about 1:10 and about 10:1. In certain embodiments, the weight ratio is between about 1:5 and about 5:1. In other embodiments, the weight ratio is between about 2:1 and about 1:2. In yet other embodiments, the weight ratio is about 1:1.

[0035] In some embodiments, the compositions are prepared by heating the components together with mixing to facilitate formulation preparation. An alternative method is to heat the phospholipid and 5ASA to a temperature sufficient for the 5ASA to melt in the phospholipid, generally with good mixing to permit intimate mixing and the formation of associated complexes between 5ASA and phospholipid at the molecular level. The process can be performed in a high temperature solvent or a bio-compatible oil in the absence of air to facilitate the formation of 5ASA/phospholipid associated complexes without oxidative degradation. An alternative method of making the formulation is to make an aqueous

-6-


formulation including 5ASA and then add the aqueous solution to a vessel coated with a dried lipid film of PC, which will then be vigorously mixed, vortexed, sonicated or subjected to other means of agitation to make a lipidic suspension of the 5ASA/PC. Another method is to dissolve the 5ASA and phospholipid in a solvent that dissolves both materials. Mixing the solution with or without heating and then evaporating the solvent. The resulting material can then be ground and encapsulated with a resin for delayed release in the distal gut.

[0036] Yet another method of combining 5ASA and a phospholipid (PL) is to covalently bind the molecules together via a diazo bond between the amino group of 5ASA and a reactive phospholipid (PL), *e.g.*, phosphatidylethanolamine (PE), phosphatidylserine or other reactive phospholipids, using standard techniques to make a diazo bond. Similar to the other 5ASA precursors, such as sulfasalazine, osalazine, and basalazide, bacterial enzymes in the colon hydrolyze the diazo bond, liberating the 5ASA and the reactive PL such as PE. The released 5ASA can then cause its intended therapeutic effect, while the released phospholipid helps to maintain or restore the hydrophobic barrier of the affected mucosa of the distal gut, inhibit inflammation and promote healing.

[0037] Pre-clinical studies to date show that phospholipids such as PC enhance the efficacy of 5ASA to treat colitis.

Suitable Reagents

[0038] Suitable phospholipids for use in this invention include, without limitation, a phospholipid of general formula:



where R₁ and R₂ are saturated and/or unsaturated substitutions ranging from 8 to 32 carbon atoms, where one or more of the carbon atoms can be replaced by an oxygen atom, a sulfur atom, a nitrogen atom, or other main group element, such as B, C, Si, P, S, Ge, and Ga; R₃ is H or CH₃, and X is H or COOH; and R₄ is =O or H₂. Mixtures and combinations of the zwitterionic phospholipids of the general formula.

[0039] Examples of zwitterionic phospholipid of formula (II) include, without limitation, phosphatidylcholines such as phosphatidylcholine (PC), dipalmitoylphosphatidylcholine

-7-

(DPPC), other disaturated phosphatidylcholines or unsaturated phosphatidylcholines, phosphatidylethanolamines, phosphatidylinositol, phosphatidyl serines sphingomyelin or other ceramides, or various other zwitterionic phospholipids, phospholipid containing oils such as marine and/or fish oils enriched in Omega-3 fatty acids, lecithin oils derived from soy beans, dimyristoyl phosphatidylcholine, distearoylphosphatidylcholine, dilinoleoylphosphatidylcholine (DLL-PC), dipalmitoylphosphatidylcholine (DPPC), soy phosphatidylcholine (Soy-PC or PC_S) and egg phosphatidylcholine (Egg-PC or PC_E). In DPPC, a saturated phospholipid, the saturated aliphatic substitution R₁ and R₂ are CH₃-(CH₂)₁₄, R₃ is CH₃ and X is H. In DLL-PC, an unsaturated phospholipid, R₁ and R₂ are CH₃-(CH₂)₄-CH=CH-CH₂-CH=CH-(CH₂)₇, R₃ is CH₃ and X is H. In Egg PC, which is a mixture of unsaturated phospholipids, R₁ primarily contains a saturated aliphatic substitution (*e.g.*, palmitic or stearic acid), and R₂ is primarily an unsaturated aliphatic substitution (*e.g.*, oleic or arachidonic acid). In Soy-PC, which in addition to the saturated phospholipids (palmitic acid and stearic acid) is a mixture of unsaturated phospholipids, (oleic acid, linoleic acid and linolenic acid). In PC derived from marine animals (*e.g.* krill, salmon) the R₁ and R₂ groups may constitute the Omega -3 fatty acids: eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). The preferred zwitterionic phospholipid include, without limitation, dipalmitoyl phosphatidylcholine, phosphatidyl choline, or a mixture thereof.

EXAMPLES

Preparation of Compositions

[0040] Phosal™ 35 SB, a soy lecithin oil containing approximately 35 wt.% phosphatidylcholine (PC), was placed in glass beaker to which 5ASA (or a 5ASA containing pharmaceutical) was added under moderate heat, about 40°C, and stirring until a 1:1 weight ratio was attained. The resulting oil-based formulation was coated with a pH-sensitive polymer, *e.g.*, MultiSal™, or Eudragit®-S, to form polymer encapsulated micro-spheres of the 5ASA/PC composition. These encapsulated micro-spheres are formed so that the 5ASA and PC are released only when the drug reaches the distal gut, where the pH is 7 or above and the polymer coating degrades. This formulation was then added to the diet of mice or rats for evaluation. The mice or rats were placed on 3% dextran sodium sulfate (DSS) solution included in their drinking water to induce colitis.

[0041] An alternative method to the oil-based formulation described above is to suspend 5ASA or a 5ASA containing pharmaceutical in water or an appropriate aqueous solution.

-8-

The aqueous solution is then added to a second container coated with a dried lipid film of PC. The resulting mixture is then subjected to vigorous mixing, vortexing, sonication or other means of agitation to form an aqueous lipidic suspension for oral, enteral or rectal (enema) administration.

Examples Using C57BL/6 Mice in a Rodent Colitis Model

[0042] Groups of four C57BL/6 mice were exposed to differing amount of dextran sodium sulfate (DSS) in their drinking water for a four day period of time. The mice were then euthanized and colon tissue weight per length measurements for each group of mice were made. Referring now to **Figure 1**, DSS treatment resulted in colonic inflammation or edema as measured by increased colon tissue weight per length, where 3% DSS was shown to cause the greatest increase in the weight to length measurement.

[0043] Groups of C57BL/6 mice were exposed to differing amount of dextran sodium sulfate (DSS) in their drinking water for a four day period of time. The mice fecal hemoglobin measurements were made at the end of the four day period. Referring now to **Figure 2**, DSS treatment resulted in increased fecal hemoglobin with increasing concentration of DSS in their drinking, with 4% DSS showing the highest fecal hemoglobin.

[0044] Groups of C57BL/6 mice were exposed to different diets and their daily food consumption (grams/day) were measured over a four day period of time. Group 1 represented the control group receiving saline without DSS and regular food without any 5ASA containing formulation. Group 2 represented a DSS control receiving saline with 3% DSS and regular food without any 5ASA containing formulation. Group 3 represented a 5ASA treated group receiving saline containing 3% DSS and a diet including 5ASA MultiSal™ encapsulated micro-spheres. Groups 4 through 6 are designed to compare 5ASA to three oil-based 5ASA/PC formulations. Group 4 represented a 5ASA/P35 (a 1:1 weight ratio of 5ASA and Phosphal® 35SB, an oil-based soy lecithin product containing about 35 wt.% PC) treated group receiving saline containing 3% DSS and a diet including 5ASA/P35 MultiSal™ encapsulated micro-spheres. Group 5 represented a 5ASA/P35 (a 1:1 weight ratio of 5ASA and Phosal® 53 MCT, an oil-based soy lecithin product containing about 53 wt.% PC) treated group receiving saline containing 3% DSS and a diet including 5ASA/P53 MultiSal™ encapsulated micro-spheres. Group 6 represented a 5ASA/P35 (a 1:1 weight ratio of 5ASA and Phosal® 75 SA, an oil-based soy lecithin product containing about 75 wt.% PC) treated group receiving saline containing 3% DSS and a diet including 5ASA/P75

-9-

MultiSal™ encapsulated micro-spheres. Referring now to **Figure 3**, the food consumption (grams/day) of mice placed on the diets described above. The first three bars evidence feeding patterns of normal mice in the absence and presence of 5ASA in the feed, while the last three bars represent the feeding patterns of mice in the presence of the 5ASA/PC formulations. **Figure 3** indicates that the active materials did not significantly affect the eating patterns of the mice.

Mouse Data Comparing 5ASA and 5ASA:PL Formulations on Treating Induced Colitis

[0045] The previous examples show that drinking water comprising saline containing 3% dextran sodium sulfate (DSS) is capable of inducing a colitis type disorder in mice. Now, the 5ASA:PL formulations can be compared to 5ASA in treating the colitis type disorder in mice. Groups of mice were placed on drinking water comprising saline containing 3% DSS for four days to induce a colitis type disorder and then treated with the compositions of this invention. The efficacy of the treating compositions were determined by measuring colon tissue weight per length. Referring now to **Figure 4**, colonic inflammation as measured by an increased colon tissue weight per given length showed partial reversal when treated with saline, 5ASA, 5ASA:P35, 5ASA:P53 and 5ASA:P75 combined with DSS and administered to the mice in their diet with a group of non-DSS treated mice and normal diet. The data showed that the three 5ASA:PL formations are superior to 5ASA alone in reversing tissue inflammation as measured by an increased colon tissue weight to length.

[0046] Groups of mice were placed on drinking water comprising saline containing 3% DSS for four days to induce a colitis type disorder and then treated with the compositions of this invention. The appearance of blood as indicated by an increased hemoglobin content in the feces was measured to determine the efficacy of 5ASA versus the 5ASA:PL formulations. Referring now to **Figure 5**, hemoglobin content in mice feces was significantly reduced for the mice placed on a diet containing the 5ASA:P35 formulation and 5ASA:P75 compared to saline/DSS and 5ASA/DSS.

[0047] Groups of mice were placed on drinking water comprising saline containing 3% DSS for four days to induce a colitis type disorder and then treated with the compositions of this invention. Colonic inflammation as indicated by an increased tissue myeloperoxidase (MPO) activity was measured to determine the efficacy of saline, 5ASA and 5ASA:PC35. Referring now to **Figure 6**, colonic inflammation was reduced for the mice placed on diets containing the 5ASA:P35 formulation.

-10-

[0048] Groups of mice were placed on drinking water comprising saline containing 3% DSS for four days to induce a colitis type disorder and then treated with the compositions of this invention. Colonic mucosa were viewed through a light microscope and histologically scored using a scoring system, where 0 represents normal colonic mucosa and 4 represent a total loss of colonic glandular structure. Referring now to **Figure 7**, the 5ASA/P35 formulation significantly reduced microscopic injury to the colonic mucosa in mice placed on a diet containing 5ASA:P35.

[0049] The rodent data clearly shows a positive enhancement of the efficacy of 5ASA in the distal gut when the administered formulation includes a phospholipid.

Rat Data Comparing 5ASA and 5ASA:PL Formulations on Treating Induced Colitis

[0050] Rats were placed on drinking water containing 4% dextran sodium sulfate (DSS) for 5 days and 20 hours to induce colitis after which they were returned to normal drinking water. The animals were then intragastrically administered twice daily for 6 subsequent days with: (a) saline (DSS control); (b) 5ASA at a dose of 100 mg/kg; (c) a 1:1 wt. ratio of 5ASA and Phosal 35 SB (an oil-based formulation); or (d) a 1:2 wt. ratio of 5ASA and Phospholipon 90G (an aqueous lipidic suspension) at the same dose of 5ASA, 100 mg/kg, b.i.d. The formulations (c) and (d) are phosphatidylcholine (PC) associated 5ASA formulations, where the PC and 5ASA are associated at the molecular level. These treatments are designed to compare the efficacy of 5ASA and two different PC:5ASA formulations.

[0051] At 1, 3 and 6 days, rats were euthanized and colonic tissue, blood and fecal matter were collected for biochemical and histological analyses to assess colonic inflammation and GI bleeding. A separate group of rats were not placed on DSS and served as "normal controls."

[0052] Referring now to **Figure 8**, changes in hematocrit during the 6 day recovery period were measured from the euthanized rats. Changes in hematocrit levels represent an indirect estimate of anemia and internal (GI) bleeding among the groups over the 6 day recovery period. It can be appreciated that although a modest <10% decrease in the hematocrit levels were seen in all the groups post DSS exposure as compared to the no DSS or "normal control" hematocrit levels, there were no significant differences in this parameter among the test groups over the 6 day recovery period.

[0053] The changes in fecal hemoglobin are shown in **Figure 9**. It can be seen that fecal blood was detected during the first day of the recovery period in most of the test groups (vs

-11-

normal control values) with a suggestion that 5ASA $\pm$ 90G may have had a beneficial effect in reducing fecal blood loss.

[0054] Colonic inflammation and injury were assessed by measuring a number of biochemical and histological indices. First, tissue edema were assessed by measuring tissue wet weight /unit length. These results which are presented in **Figure 10**, demonstrate that intestinal edema was maximal 3 days post-DSS in saline-treated controls, and this inflammatory change was reduced by 5ASA (non significantly), and further decreased (significantly) in rats treated with the two PC-[5ASA] formulations. There were no significant differences among the groups in this parameter 6 days post-DSS, with some indication that the animals treated with 5ASA in combination with 90G had a rebound episode of inflammation.

[0055] Colonic myeloperoxidase (MPO) activity, a biochemical marker of neutrophil infiltration into the tissue, was measured as another marker of inflammation. The results presented in **Figure 11** demonstrate similar, but more dramatic changes to those described above, with 5ASA showing a reduction on day 3 and both PC-[5ASA] formulations further decreasing MPO activity to almost undetectable levels. MPO activity returned to normal control levels by 6 days in all groups, except for the group treated with 5ASA in combination with 90G.

[0056] Tissue histology was scored (under blinded conditions) for signs of injury and inflammation. To assess inflammation we scored the number of inflammatory cells in the lamina propria of the colonic submucosa using a scoring system from 1 (normal) to 4 (highly inflamed). These results, shown in **Figure 12**, demonstrate that 5ASA-treated rats had a reduced number of inflammatory cells with the 5ASA/P35 showing comparable or superior anti-inflammatory activity at 3 and 6 days and again 90G in combination with 5ASA causing a rebound of inflammation at 6 days.

[0057] To assess colonic mucosal injury, a histological scoring system was used, where: 0 = normal mucosa; 1 = surface injury; 2 = injury extending into the mid glandular region; 3 = injury extending into the deep glandular region; and 4 = injury extending into the serosal (muscle) layer. **Figure 13** shows that mucosal injury was increased in all groups day 1 post-DSS, and by day 3 there was a recovery in rats treated with 5 ASA $\pm$ PC with the 5ASA/P35 showing more consistent evidence of mucosal protection than 90G.

Conclusions

-12-

[0058] The protocol used in the above series of experiments, proves to be a very useful rodent model system to evaluate the therapeutic efficacy of 5ASA and PC-5ASA formulations to accelerate the recovery of experimentally (DSS)-induced colitis. Interestingly, it appears that inflammation progresses for the first 3 days after withdrawal from DSS, the colitis inducing agent, and subsides by day 6. Using a number of biochemical and histological markers, it appears that 5ASA at a dose of 100 mg/kg significantly reduces colonic inflammation and tissue injury 3 days post-DSS in comparison to saline-treated control values. The oil-based 5ASA/P35SB formulation showed a further improvement in the recovery from colitis (vs 5ASA alone) in 2 out of 4 markers of inflammation. In contrast, the 5ASA/90G formulation had a biphasic effect with a rebound of inflammation occurring 6 days post-DSS. It is our recommendation that this recovery experiment, using the above model system, be repeated using a lower less effective dose of 5ASA (50 mg/kg b.i.d.), and that we focus our efforts on the oil-based 5ASA/P35 formulation, to determine if a more clear and convincing enhancement in the anti-inflammatory efficacy of our PC-5ASA test formulation can be demonstrated in comparison to 5ASA alone.

Rat Data Comparing 5ASA and 5ASA:PL Formulations on Treating Induced Colitis

[0059] The following experiments were designed to compare the efficacy of 5ASA versus 5ASA:PC formulations in treating rats that had been subjected to colitis induced by adding 4% dextran sodium sulfate (DSS) to their drinking water for over 5 days (5 day, 20 hours) and then returned to normal drinking water. The rats were then treated with 5ASA, in this experiment at a lower dose (50 mg/kg, BID) than that previously studied, in the absence or presence to a PL component for 3 days post DSS induced colitis.

[0060] Thirty two healthy male rats used in the study. Rats comparable with respect to protocol, supplier, sex, and weight were divided serially into 4 groups. Group 1 were given drinking water without added DSS. Groups 2-4 received drinking water containing 4% DSS for 6 days. At day 6, Group 2 received 1 mL saline BID intra gastrically. Group 3 received 5ASA (50mg/Kg) BID intra gastrically. Group 4 received 5ASA(50mg/Kg): Phosal 35SB:MCT oil (1:1:0.3), referred to heretofore as 5ASA:PC BID intra gastrically, where MCT = Medium Chain Triglyceride.

[0061] At day 9, the rats were euthanized and the following were measured: (a) hematocrit, (b) fecal hemoglobin, (c) colon weight / length, (d) colon myeloperoxidase (MPO), (e) colonic surface hydrophobicity and (f) colon histology.

-13-

[0062] In this study, rats were placed on regular drinking water (Group 1) or water containing 4% DSS and then intragastrically administered: saline (Group 2); 5ASA (Group 3) at a dose of 50 mg/kg BID); or 5ASA:PC (P35SB at 1:1 wt ratio) (Group 4) for 3 days at which time the following changes were seen.

[0063] Referring now to **Figure 14**, the colon weight/length data is shown. This measure of mucosal edema demonstrated a significant increase when comparing the DSS saline-treated group (Group 2) with the absolute control (no DSS, Group 1). This DSS –induced mucosal edema was partially reversed by 5ASA at a dose of 50 mg/kg BID (Group 3), and completely reversed in the 5ASA:PC group (Group 4), with significant differences observed between the rats of Group 4 versus the rats of Groups 2 & 3, but not the rats of Group 1.

[0064] Referring now to **Figure 15**, the MPO Protein data is shown. This biochemical measure of inflammation showed the same pattern as outlined above, with DSS inducing a significant increase in MPO activity versus the control Group 1 (no DSS), and this inflammation was partially and completely reversed by 5ASA and 5ASA:PC, respectively. Significant differences ($p<0.05$) were observed between the rats of Group 4 versus the rats of Groups 2 & 3, but not the rats of Group 1.

[0065] Referring now to **Figure 16**, colon contact angle analysis data is shown. No differences in colonic mucosal surface hydrophobicity was observed between groups, possibly because this surface property had recovered fully by 3 days post-DSS induced colitis.

[0066] Referring now to **Figure 17**, hematocrit data is shown. This measure of anemia due to GI bleeding was not significantly different among the experimental groups treated with DSS versus control (no DSS), but did tend to be lower in the DSS-treated groups except for those administered 5ASA:PC (Group 4), where it was significantly higher.

[0067] Referring now to **Figure 18**, fecal hemoglobin data is shown. This direct measure of active GI bleeding demonstrated an increase in the saline-treated DSS group (Group 2) versus the control (no DSS) group (Group 1). This DSS-induced increase in GI fecal blood loss was partially reversed by 5ASA administration and completely reversed by 5ASA:PC administration. Significant differences ($p<0.05$) were observed between Group 4 versus Groups 2 & 3, but not Group 1.

[0068] DSS-induces colonic inflammation and bleeding in rats that is still evident 3 days post-DSS. 5ASA alone at a dose of 50 mg/kg administered twice daily intragastrically

-14-

promotes tissue recovery from DSS-induced colitis and reduces GI bleeding. PC-5ASA when administered intragastrically at the same dose of active 5ASA induced full recovery of the mucosa from DSS-induced colitis with little or no evidence of GI bleeding.

- 15 -

CLAIMS

1. Use of a 5-aminosalicylic acid (5ASA) containing component and a phospholipid-containing oil in the manufacture of a medicament for treating inflammatory bowel disease (IBD) characterized in that the phospholipid-containing oil is present in an amount to enhance the anti-inflammatory efficacy of the 5ASA.
2. The use of a 5-aminosalicylic acid (5ASA) containing component and a phospholipid or a phospholipid-containing oil in the manufacture of a non-liposomal medicament for treating inflammatory bowel disease (IBD) characterized in that the phospholipid or a phospholipid-containing oil is present in an amount to enhance the anti-inflammatory efficacy of the 5ASA.
3. The use of claim 1 or claim 2, wherein the weight ratio of 5ASA to phospholipid is between 1:10 and 10:1.
4. The use of claim 3, wherein the weight ratio of 5ASA to phospholipid is between 1:5 and 5:1.
5. The use of claim 4, wherein the weight ratio of 5ASA to phospholipid is between 2: 1 and 1:2.
6. The use of claim 5, wherein the weight ratio of 5ASA to phospholipid is about 1:1.
7. The use of claim 1 or claim 2, wherein the medicament comprises resin encapsulated microspheres of the 5ASA and phospholipid components capable of being released at pH values of 7 or higher.
8. The use of any one of claims 1 to 7, wherein the 5ASA and the phospholipid-containing oil components are in the form of a mixture.
9. The use of claim 1 or claim 2, wherein 5ASA and the phospholipid are associated at a molecular level to form a complex.

- 16 -


10. The use of claim 2, wherein 5ASA is covalently bonded to a reactive phospholipid via a diazo linkage capable of being released in the distal gut, due the presence of bacteria that catalyze the hydrolysis of the diazo linkage.
11. The use of any one of claims 1 to 10, wherein the medicament is for rectal administration.
12. The use of any one of claims 1 to 10, wherein the medicament is for oral administration.
13. The use of any one of claims 1 to 12, wherein the 5ASA component is selected from Mesalamine, Sulfasalazine, Olsalazine, Balsalazide, and mixtures thereof.
14. The use of claim 13, wherein the 5ASA component is Mesalamine.
15. The use of any one of claims 1 to 14, wherein the phospholipid component is a phosphatidylcholine-containing soy lecithin oil.
16. The use of claim 15, wherein the phospholipid component is a soy lecithin oil containing 35 wt. % to 75 wt. % phosphatidylcholine.
17. The use of claim 16, wherein the phospholipid component is a soy lecithin oil containing about 35 wt. % phosphatidylcholine.
18. The use of claim 16 or claim 17, wherein weight ratio of 5ASA to phosphatidylcholine-containing soy lecithin oil is 1:1.
19. A use of claim 2, wherein the medicament comprises a composition of the 5ASA and phospholipid components prepared by:
 - simply mixing a powder containing 5ASA or a larger molecule covalently bound to active 5ASA via a diazo bond with a phospholipid;
 - simply mixing the 5ASA containing component with an oil-based phospholipid component

- 17 -

heating the components together with mixing to facilitate formulation preparation;
heating the phospholipid and 5ASA to a temperature sufficient for the 5ASA to melt in the phospholipid; or
dissolving the 5ASA and phospholipid in solvent that dissolves both materials, mixing the solution with or without heating and then evaporating the solvent.

20. A use of claim 19, wherein the composition is as defined in any one of claims 3 to 8 and 11 to 14.
21. A composition comprising 5-aminosalicylic acid (5ASA) and a phospholipid associated at a molecular level at a weight ratio of 5ASA to phospholipid of between 1:10 and 10:1 to form a complex.
22. A covalent compound including 5ASA and an active phospholipid, wherein the reactive phospholipid is covalently bonded to the 5ASA via a diazo linkage between the amino group of 5ASA and the active phospholipid, said linkage being cleavable by bacteria in the distal gut that catalyze the hydrolysis of the diazo linkage.

Treatment of C57BL/6 Mice with DSS in Drinking Water for Four Days: Edema of Colon

**FIG. 1**

2/15

**DSS in Drinking Water of C57BL/6 Mice for 5
Days: Blood in Fecal Pellets**






FIG. 2



3/15

DSS Induced Colitis in C57BL/6 : Food Consumption

**FIG. 3**

**FIG. 4**

5/15

**FIG. 5****FIG. 6**

6/15

**DSS Induced Colitis in C57BL/6 Mice:
Mucosal Damage by H&E Stained Histology Slides**






FIG. 7

7/15

**FIG. 8**

8/15

**FIG. 9**

9/15




FIG. 10

10/15




FIG. 11

11/15




FIG. 12

12/15









FIG. 13

**FIG. 14****FIG. 15**

**FIG. 16****FIG. 17**

15/15

**FIG. 18**

