

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(10) International Publication Number  
**WO 2013/148330 A1**

(43) International Publication Date  
**3 October 2013 (03.10.2013)**

(51) International Patent Classification:  
**C07K 14/435 (2006.01) C12N 15/63 (2006.01)**  
**A23J 3/04 (2006.01) C12N 1/21 (2006.01)**  
**A23L 1/305 (2006.01)**

(21) International Application Number:  
**PCT/US2013/032218**

(22) International Filing Date:  
**15 March 2013 (15.03.2013)**

(25) Filing Language:  
**English**

(26) Publication Language:  
**English**

(30) Priority Data:  
**61/615,819 26 March 2012 (26.03.2012) US**

(71) Applicant: **PRONUTRIA, INC. [US/US]**; 840 Memorial Drive, 3rd Floor, Cambridge, MA 02139 (US).

(72) Inventors: **BERRY, David Arthur**; 20 Cameron Street, #402, Brookline, MA 02445 (US). **BOGHIGIAN, Brett Adam**; 344 W Broadway, Apt. #2, South Boston, MA 02127 (US). **SILVER, Nathaniel W.**; 1218 Cambridge Street, #2, Cambridge, MA 02139 (US). **VON MALTZAHN, Geoffrey**; 61 Salem St., Apt. 5, Boston, MA 02113 (US). **CHILLAKURU, Rajeev**; Pronutria, Inc., 840 Memorial Drive, 3rd Floor, Cambridge, MA 02139 (US). **HAMILL, Michael J.**; Pronutria, Inc., Drive, 3rd Floor, Cambridge, MA 02139 (US).

(74) Agents: **KABLER, Kevin, E. et al.**; Fenwick & West LLP, Silicon Valley Center, 801 California Street, Mountain View, California 94041 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with amended claims and statement (Art. 19(1))
- with sequence listing part of description (Rule 5.2(a))

**WO 2013/148330 A1**

(54) Title: **NUTRITIVE FRAGMENTS, PROTEINS AND METHODS**

(57) Abstract: Nutritive proteins are provided. In some embodiments the nutritive proteins comprise a first polypeptide sequence comprising a fragment of a naturally-occurring nutritive protein. In some embodiments the fragment comprises at least one of a) an enhanced ratio of branch chain amino acid residues to total amino acid residues present in the nutritive protein; b) an enhanced ratio of leucine residues to total amino acid residues present in the nutritive protein; and c) an enhanced ratio of essential amino acid residues to total amino acid residues present in the nutritive protein. In some embodiments, the fragment comprises at least one of a) a ratio of branch chain amino acid residues to total amino acid residues present in the nutritive protein; b) a ratio of leucine residues to total amino acid residues present in the nutritive protein; and c) a ratio of essential amino acid residues to total amino acid residues present in the nutritive protein, that is equal to or greater than the corresponding ratio present in a benchmark protein such as whey, egg or soy protein. Also provided are nucleic acids encoding the proteins, recombinant microorganisms that make the proteins, methods of making the proteins using recombinant microorganisms, compositions that comprise the proteins, and methods of using the proteins, among other things.

**NUTRITIVE FRAGMENTS, PROTEINS AND METHODS****REFERENCE TO RELATED APPLICATION:**

[0001] This application claims priority to U.S. Provisional Patent Application No. 61/615,819, filed March 26, 2012, which is hereby incorporated herein by reference in its 5 entirety.

**SEQUENCE LISTING:**

[0002] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on March 12, 2013, is named 1005.005-PCT\_SL.txt and is 3,234,600 10 bytes in size.

**INTRODUCTION:**

[0003] Dietary protein is an essential nutrient for human health and growth. The World Health Organization recommends that dietary protein should contribute approximately 10 to 15% of energy intake when in energy balance and weight stable. Average daily protein 15 intakes in various countries indicate that these recommendations are consistent with the amount of protein being consumed worldwide. Meals with an average of 20 to 30% of energy from protein are representative of high-protein diets when consumed in energy balance.

[0004] The body cannot synthesize certain amino acids that are necessary for health 20 and growth, and instead must obtain them from food. These amino acids, called “essential amino acids”, are Histidine (H), Isoleucine (I), Leucine (L), Lysine (K), Methionine (M), Phenylalanine (F), Threonine (T), Tryptophan (W), and Valine (V). Dietary proteins that provide all the essential amino acids are referred to as “high quality” proteins. Animal foods such as meat, fish, poultry, eggs, and dairy products are generally regarded as high quality 25 protein sources that provide a good balance of essential amino acids. Casein (a protein commonly found in mammalian milk, making up 80% of the proteins in cow milk) and whey (the protein in the liquid that remains after milk has been curdled and strained) are major sources of high quality dietary protein. Foods that do not provide a good balance of essential amino acids are referred to as “low quality” proteins. Most fruits and vegetables are poor 30 sources of protein. Some plants foods including beans, peas, lentils, nuts and grains (such as

wheat) are better sources of protein. Soy, a vegetable protein manufactured from soybeans, is considered by some to be a high quality protein.

[0005] Studies of the acute effects of consuming high amounts of protein in humans have shown that inclusion and in some cases increasing protein content in the diet can have 5 beneficial effects. For example, studies have shown that ingestion of protein can induce postprandial satiety (including by suppressing hunger), induce thermogenesis and reduce glycemic response in human subjects.

[0006] Studies of high protein diets for weight loss have shown that protein positively affects energy expenditure and lean body mass. Further studies have shown that overeating 10 produces significantly less weight gain in diets containing at least 5% of energy from protein, and that a high-protein diet decreases energy intake.

[0007] Clinical studies provide evidence that protein prevents muscle loss due to aging or bed rest. In particular, studies have shown that protein supplementation increases 15 muscle fractional synthetic rate (FSR) during prolonged bed rest, maintains leg mass and strength during prolonged bed rest increases lean body mass, improves functional measures of gait and balance, and may serve as a viable intervention for individuals at risk of sarcopenia due to immobility or prolonged bed rest.

[0008] Studies on increasing muscle protein anabolism in athletes have shown that protein provided following exercise promotes muscle hypertrophy to a greater extent than 20 that achieved by exercise alone. It has also been shown that protein provided following exercise supports protein synthesis without any increase in protein breakdown, resulting in a net positive protein balance and muscle mass accretion. While muscle protein synthesis appears to respond in a dose-response fashion to essential amino acid supplementation, not 25 all proteins are equal in building muscle. For example, milk proteins appear to be superior to soy in supporting muscle mass accretion with resistance training, while both are superior to carbohydrate alone. The amino acid leucine is an important factor in stimulating muscle protein synthesis.

[0009] Whole proteins commonly found in foods do not necessarily provide an amino acid composition that meets the amino acid requirements of a mammal, such as a human, in 30 an efficient manner. The result is that, in order to attain the minimal requirements of each essential amino acid, a larger amount of total protein must be consumed in the diet than

would be required if the quality of the dietary protein were higher. By increasing the quality of the protein in the diet it is possible to reduce the total amount of protein that must be consumed compared to diets that include lower quality proteins.

[0010] In general, proteins that have higher protein quality are considered more

5 beneficial in a mammalian diet than other proteins that do not. Such proteins are useful, for example, as components of a mammalian diet. Under certain circumstances such proteins promote maintenance of muscle mass, a healthy body mass index, and glycemic balance, among other things. Accordingly, there is a need for sources of proteins that have high protein quality.

10 [0011] Traditionally, desirable mixtures of amino acids, such as mixtures comprising essential amino acids, have been provided by hydrolyzing a protein with relatively high levels of essential amino acids, such as whey protein, and/or by combining free amino acids in a mixture that optionally also includes a hydrolyzed protein such as whey. Mixtures of this type may have a bitter taste and may be deemed unsuitable or undesirable for certain uses.

15 As a result, such mixtures sometimes include flavoring agents to mask the taste of the free amino acids and/or hydrolyzed protein. In some cases compositions in which a proportion of the amino acid content is provided by polypeptides or proteins are found to have a better taste than compositions with a high proportion of total amino acids provided as free amino acids and/or certain hydrolyzed proteins. The availability of such compositions has been limited,

20 however, because nutritional formulations have traditionally been made from protein isolated from natural food products, such as whey isolated from milk, or soy protein isolated from soy. The amino acid profiles of those proteins do not necessarily meet the amino acid requirements for a mammal. In addition, commodity proteins typically consist of mixtures of proteins and/or protein hydrolysates which can vary in their protein composition, thus leading

25 to unpredictability regarding their nutritional value. Moreover, the limited number of sources of such high quality proteins has meant that only certain combinations of amino acids are available on a large scale for ingestion in protein form.

[0012] The agricultural methods required to supply high quality animal protein

sources such as casein and whey, eggs, and meat, as well as plant proteins such as soy, also

30 require significant energy inputs and have potentially deleterious environmental impacts. Accordingly, it would be useful in certain situations to have alternative sources and methods of supplying proteins for mammalian consumption.

[0013] In theory, synthetic polypeptide sequences comprising a desired mixture of amino acids could be designed and produced in a laboratory setting. This approach may raise various concerns, however, and is therefore not always applicable. First, skilled artisans are aware that obtaining high levels of production of such synthetic sequences may be very 5 challenging. Second, even if such a synthetic protein were synthesized, its suitability for use in a nutritive product would be uncertain. For example, such a non-naturally occurring polypeptide could be an allergen or a toxin. Accordingly, in some embodiments this disclosure provides natural protein or polypeptide sequences, or variants thereof.

[0014] In this disclosure the inventors provide nutritive proteins comprising a first 10 polypeptide sequence that is homologous to a fragment of a naturally-occurring protein. The nutritive proteins and the fragments of which they are composed are made up of useful combinations of amino acids. The proteins can be produced by methods that do not rely solely on traditional agriculture for production. For example, the inventors have discovered and this disclosure provides nutritive proteins comprising a fragment of a naturally-occurring 15 protein and composed of combinations of amino acids that contain a useful level of at least one of a ratio of branch chain amino acids to total amino acids, a ratio of the amino acid leucine to total amino acids, and a ratio essential amino acids to total amino acids. In some embodiments the nutritive proteins comprise at least one of a level of a) a ratio of branch chain amino acid residues to total amino acid residues present in the nutritive protein equal to 20 or greater than the ratio of branch chain amino acid residues to total amino acid residues present in whey protein; b) a ratio of leucine residues to total amino acid residues present in the nutritive protein equal to or greater than the ratio of leucine residues to total amino acid residues present in whey protein; and c) a ratio of essential amino acid residues to total amino acid residues present in the nutritive protein equal to or greater than the ratio of essential 25 amino acid residues to total amino acid residues present in whey protein.

[0015] This disclosure also provides nucleic acids encoding the proteins, recombinant microorganisms that make the proteins, methods of making the proteins using recombinant microorganisms, compositions that comprise the proteins, and methods of using the proteins, among other things.

### 30 SUMMARY:

[0016] In a first aspect this disclosure provides isolated nutritive proteins comprising a first polypeptide sequence that is homologous to a fragment of a naturally-occurring

protein, wherein the first polypeptide sequence comprises at least one of: a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%; b. a ratio of Leu residues to total amino acid residues of at least 11%; and c. a ratio of essential amino acid residues to total amino acid residues of at least 49%. In some embodiments the first 5 polypeptide sequence further comprises at least one of each essential amino acid. In some embodiments the first polypeptide sequence comprises at least 70% homology to the fragment of a naturally-occurring protein. In some embodiments the first polypeptide sequence comprises at least 95% homology to the fragment of a naturally-occurring protein. In some embodiments the fragment of a naturally-occurring protein comprises at least 25 10 amino acid residues. In some embodiments the fragment of a naturally-occurring protein comprises at least 50 amino acid residues. In some embodiments the first polypeptide sequence comprises: a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%; b. a ratio of Leu residues to total amino acid residues of at least 11%; and c. a ratio of essential amino acid residues to total amino acid residues of at least 15 49%. In some embodiments the first polypeptide sequence comprises a fragment of a naturally-occurring nutritive protein. In some embodiments the first polypeptide sequence consists of a fragment of a naturally-occurring nutritive protein.

[0017] In some embodiments the first polypeptide sequence is not an allergen. In some embodiments the first polypeptide sequence has less than 50% global homology to a 20 known allergen.

[0018] In some embodiments the first polypeptide sequence is not a toxin. In some embodiments the first polypeptide sequence has less than 50% global homology to a known toxin.

[0019] In some embodiments the first polypeptide sequence has a simulated gastric 25 digestion half-life of less than 60 minutes. In some embodiments the first polypeptide sequence has a simulated gastric digestion half-life of less than 30 minutes. In some embodiments the first polypeptide sequence has a simulated gastric digestion half-life of less than 10 minutes. In some embodiments the first polypeptide sequence is completely digested in simulated gastric fluid. In some embodiments the first polypeptide sequence comprises at 30 least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site. In some embodiments the first polypeptide sequence comprises no cysteine residues. In some embodiments the first

polypeptide sequence comprises no disulfide bonds. In some embodiments the first polypeptide sequence does not comprise N-linked glycosylation. In some embodiments the first polypeptide sequence does not comprise O-linked glycosylation.

[0020] In some embodiments the first polypeptide sequence is resistant to

5 aggregation. In some embodiments the first polypeptide sequence is anionic at pH 7. In some embodiments the first polypeptide sequence has an aqueous solubility at pH 7 of at least 12.5 g/L. In some embodiments the first polypeptide sequence has a calculated solvation score of -20 or less. In some embodiments the first polypeptide sequence has a calculated aggregation score of 0.75 or less. In some embodiments the first polypeptide

10 sequence has a calculated aggregation score of 0.5 or less.

[0021] In some embodiments the first polypeptide sequence comprises an amino acid sequence selected from: i. an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and iii. a mutein of an amino acid sequence selected from **SEQ ID**

15 **NO: 1 to SEQ ID NO: 2609**. In some embodiments the first polypeptide sequence consists of an amino acid sequence selected from: i. an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and iii. a mutein of an amino acid sequence selected from v. In some embodiments the first polypeptide sequence is at least 40%, 45%,  
20 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

[0022] In some embodiments the first polypeptide sequence is a fragment of at least

25 25 amino acids of the naturally-occurring nutritive protein. In some embodiments the first polypeptide sequence is a fragment of at least 50 amino acids of the naturally-occurring nutritive protein. In some embodiments the isolated nutritive protein consists of the first polypeptide sequence. In some embodiments the isolated nutritive protein further comprises a polypeptide tag for affinity purification. In some embodiments the tag for affinity purification is a polyhistidine-tag.

30 [0023] In another aspect this disclosure provides isolated nutritive proteins

comprising a first polypeptide sequence that is homologous to a fragment of a naturally-

occurring protein, wherein the isolated nutritive protein comprises at least one of: a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%; b. a ratio of Leu residues to total amino acid residues of at least 11%; and c. a ratio of essential amino acid residues to total amino acid residues of at least 49%. In some embodiments the isolated nutritive protein further comprises at least one of each essential amino acid. In some embodiments the isolated nutritive protein comprises at least 70% homology to the fragment of a naturally-occurring protein. In some embodiments the isolated nutritive protein comprises at least 95% homology to the fragment of a naturally-occurring protein. In some embodiments the fragment of a naturally-occurring protein comprises at least 25 amino acid residues. In some embodiments the fragment of a naturally-occurring protein comprises at least 50 amino acid residues. In some embodiments the isolated nutritive protein comprises: a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%; b. a ratio of Leu residues to total amino acid residues of at least 11%; and c. a ratio of essential amino acid residues to total amino acid residues of at least 49%. In some embodiments the isolated nutritive protein comprises a fragment of a naturally occurring nutritive protein. In some embodiments the isolated nutritive protein consists of a fragment of a naturally occurring nutritive protein.

[0024] In some embodiments the isolated nutritive protein is not an allergen. In some embodiments the isolated nutritive protein has less than 50% global homology to a known allergen.

[0025] In some embodiments the isolated nutritive protein is not a toxin. In some embodiments the isolated nutritive protein has less than 50% global homology to a known toxin.

[0026] In some embodiments the isolated nutritive protein has a simulated gastric digestion half-life of less than 60 minutes. In some embodiments the isolated nutritive protein has a simulated gastric digestion half-life of less than 30 minutes. In some embodiments the isolated nutritive protein has a simulated gastric digestion half-life of less than 10 minutes. In some embodiments the isolated nutritive protein is completely digested in simulated gastric fluid. In some embodiments the isolated nutritive protein comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site. In some embodiments the isolated nutritive protein comprises no cysteine residues. In some embodiments the isolated nutritive

protein comprises no disulfide bonds. In some embodiments the isolated nutritive protein does not comprise N-linked glycosylation. In some embodiments the isolated nutritive protein does not comprise O-linked glycosylation.

[0027] In some embodiments the isolated nutritive protein is resistant to aggregation.

5 In some embodiments the isolated nutritive protein is anionic at pH 7. In some embodiments the isolated nutritive protein has an aqueous solubility at pH 7 of at least 12.5 g/L. In some embodiments the isolated nutritive protein has a calculated solvation score of -20 or less. In some embodiments the isolated nutritive protein has a calculated aggregation score of 0.75 or less. In some embodiments the isolated nutritive protein has a calculated aggregation score  
10 of 0.5 or less.

[0028] In some embodiments the isolated nutritive protein comprises an amino acid sequence selected from: i. an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and iii. a mutein of an amino acid sequence selected from **SEQ ID**

15 **NO: 1 to SEQ ID NO: 2609**. In some embodiments the isolated nutritive protein consists of an amino acid sequence selected from: i. an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and iii. a mutein of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**. In some embodiments the isolated nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%,  
20 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**. In some embodiments the isolated nutritive protein further comprises a polypeptide tag for affinity purification. In some embodiments the tag for affinity purification is a polyhistidine-tag.

25 [0029] In another aspect this disclosure provides isolated nucleic acids comprising a nucleic acid sequence that encodes a nutritive protein according to this disclosure. In some embodiments the isolated nucleic acid is selected from genomic DNA, cDNA, sense RNA and antisense RNA. In some embodiments the isolated nucleic acid is genomic DNA. In some embodiments the isolated nucleic acid is cDNA. In some embodiments the isolated nucleic acid further comprises an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein. In some embodiments the nucleic acid sequence that encodes a nutritive protein according to this disclosure is present in a  
30

vector. In some embodiments the vector further comprises an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein.

[0030] In another aspect this disclosure provides recombinant microorganisms comprising at least one of a nucleic acid and a vector according to this disclosure. In some 5 embodiments the recombinant microorganism is a prokaryote. In some embodiments the prokaryote is heterotrophic. In some embodiments the prokaryote is autotrophic. In some embodiments the prokaryote is a bacteria.

[0031] In another aspect this disclosure provides methods of making a nutritive protein according to this disclosure, the method comprising culturing a recombinant 10 microorganism according to this disclosure under conditions sufficient for production of the nutritive protein by the recombinant microorganism. In some embodiments the method further comprises isolating the nutritive protein from the culture.

[0032] In another aspect this disclosure nutritive compositions comprising a nutritive protein of of this disclosure and at least one second component. In some embodiments the at 15 least one second component is selected from a protein, a polypeptide, a peptide, a free amino acid, a carbohydrate, a fat, a mineral or mineral source, a vitamin, a supplement, an organism, a pharmaceutical, and an excipient. In some embodiments the at least one second component is a protein. In some embodiments the at least one second component is a nutritive protein. In some embodiments the at least one second component is a free amino acid selected from 20 essential amino acids, non-essential amino acids, branch chain amino acids, non-standard amino acids and modified amino acids. In some embodiments the at least one second component is a free amino acid selected from essential amino acids. In some embodiments the at least one second component is a free amino acid selected from branch chain amino acids. In some embodiments the at least one second component is Leu. In some 25 embodiments the at least one second component is a lipid. In some embodiments the lipid is selected from a fat, oil, triglyceride, cholesterol, phospholipid, and fatty acid. In some embodiments the at least one second component is selected from a mineral and a vitamin. In some embodiments the at least one second component is a supplement. In some embodiments the at least one second component is an organism. In some embodiments the at 30 least one second component is a pharmaceutical. In some embodiments the at least one second component is an excipient. In some embodiments the at least one excipient is selected from a buffering agent, a preservative, a stabilizer, a binder, a compaction agent, a

lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a coloring agent. In some embodiments the nutritive composition is formulated as a liquid solution, slurry, suspension, gel, paste, powder, or solid.

5 [0033] In another aspect this disclosure provides methods of making a nutritive composition of this disclosure, comprising providing a nutritive protein according to this disclosure and combining the nutritive protein with the at least one second component.

10 [0034] In another aspect this disclosure provides methods of maintaining or increasing at least one of muscle mass, muscle strength, and functional performance in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the 15 subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

20 [0035] In another aspect this disclosure provides methods of maintaining or achieving a desirable body mass index in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the nutritive protein of this disclosure, 25 nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

30 [0036] In another aspect this disclosure provides methods of providing protein to a subject with protein-energy malnutrition, the method comprising providing to the subject a

sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

5 [0037] In another aspect this disclosure provides methods of increasing thermogenesis in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

10 [0038] In another aspect this disclosure provides methods of inducing at least one of a satiation response and a satiety response in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

15 [0039] In another aspect this disclosure provides methods of treating at least one of cachexia, sarcopenia and frailty in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some

embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

#### **BRIEF DESCRIPTION OF THE DRAWINGS:**

5 [0040] **Figure 1** shows a two dimensional histogram indicating the relative likelihood (on a log scale) of a protein being expressed in an *E. coli* expression screen as a function of solvation score (y-axis) and aggregation score (x-axis).

[0041] **Figure 2** shows a two dimensional histogram indicating the relative likelihood (on a log scale) of a protein being solubly expressed in an *E. coli* expression screen as a 10 function of solvation score (y-axis) and aggregation score (x-axisNN).

#### **DETAILED DESCRIPTION:**

[0042] Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by 15 context, singular terms shall include the plural and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, biochemistry, enzymology, molecular and cellular biology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art. Certain references and other documents cited herein are expressly incorporated 20 herein by reference. Additionally, all UniProt/SwissProt records cited herein are hereby incorporated herein by reference. In case of conflict, the present specification, including definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0043] The methods and techniques of the present disclosure are generally performed 25 according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001); Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992, 30 and Supplements to 2002); Taylor and Drickamer, *Introduction to Glycobiology*, Oxford Univ. Press (2003); Worthington Enzyme Manual, Worthington Biochemical Corp.,

Freehold, N.J.; Handbook of Biochemistry: Section A Proteins, Vol I, CRC Press (1976); Handbook of Biochemistry: Section A Proteins, Vol II, CRC Press (1976); Essentials of Glycobiology, Cold Spring Harbor Laboratory Press (1999). Many molecular biology and genetic techniques applicable to cyanobacteria are described in Heidorn et al., "Synthetic 5 Biology in Cyanobacteria: Engineering and Analyzing Novel Functions," Methods in Enzymology, Vol. 497, Ch. 24 (2011), which is hereby incorporated herein by reference.

[0044] This disclosure refers to sequence database entries (e.g., UniProt/SwissProt) for certain protein and gene sequences that are published on the internet, as well as other information on the internet. The skilled artisan understands that information on the internet, 10 including sequence database entries, is updated from time to time and that, for example, the reference number used to refer to a particular sequence can change. Where reference is made to a public database of sequence information or other information on the internet, it is understood that such changes can occur and particular embodiments of information on the internet can come and go. Because the skilled artisan can find equivalent information by 15 searching on the internet, a reference to an internet web page address or a sequence database entry evidences the availability and public dissemination of the information in question.

[0045] Before the present proteins, compositions, methods, and other embodiments are disclosed and described, it is to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. It must 20 be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise.

[0046] The term "comprising" as used herein is synonymous with "including" or "containing", and is inclusive or open-ended and does not exclude additional, unrecited members, elements or method steps.

25 [0047] This disclosure makes reference to amino acids. The full name of the amino acids is used interchangeably with the standard three letter and one letter abbreviations for each. For the avoidance of doubt, those are: Alanine (Ala, A), Arginine (Arg, R), Asparagine (Asn, N), Aspartic acid (Asp, D), Cysteine (Cys, C), Glutamic Acid (Glu, E), Glutamine (Gln, Q), Glycine (Gly, G), Histidine (His, H), Isoleucine (Ile, I), Leucine (Leu, L), 30 Lysine (Lys, K), Methionine (Met, M), Phenylalanine (Phe, F), Proline (Pro, P), Serine (Ser, S), Threonine (Thr, T), Tryptophan (Trp, W), Tyrosine (Tyr, Y), Valine (Val, V).

[0048] As used herein, the term “in vitro” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, in a Petri dish, etc., rather than within an organism (e.g., animal, plant, or microbe).

5 [0049] As used herein, the term “in vivo” refers to events that occur within an organism (e.g., animal, plant, or microbe).

[0050] As used herein, the term “isolated” refers to a substance or entity that has been (1) separated from at least some of the components with which it was associated when initially produced (whether in nature or in an experimental setting), and/or (2) produced, prepared, and/or manufactured by the hand of man. Isolated substances and/or entities may 10 be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated agents are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a 15 substance is “pure” if it is substantially free of other components.

[0051] As used herein, a “branch chain amino acid” is an amino acid selected from Leucine, Isoleucine, and Valine.

[0052] As used herein, an “essential amino acid” is an amino acid selected from Histidine, Isoleucine, Leucine, Lysine, Methionine, Phenylalanine, Threonine, Tryptophan, 20 and Valine.

[0053] The term “peptide” as used herein refers to a short polypeptide, e.g., one that typically contains less than about 50 amino acids and more typically less than about 30 amino acids. The term as used herein encompasses analogs and mimetics that mimic structural and thus biological function.

25 [0054] The term “polypeptide” encompasses both naturally-occurring and non-naturally occurring proteins, and fragments, mutants, derivatives and analogs thereof. A polypeptide may be monomeric or polymeric. Further, a polypeptide may comprise a number of different domains each of which has one or more distinct activities. For the avoidance of doubt, a “polypeptide” may be any length greater than two amino acids.

[0055] The term “isolated protein” or “isolated polypeptide” is a protein or polypeptide that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) exists in a purity not found in nature, where purity can be adjudged with respect to the presence of other cellular material (e.g., is free of other proteins from the same species) (3) is expressed by a cell from a different species, or (4) does not occur in nature (e.g., it is a fragment of a polypeptide found in nature or it includes amino acid analogs or derivatives not found in nature or linkages other than standard peptide bonds). Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be “isolated” from its naturally associated components. A polypeptide or protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art. As thus defined, “isolated” does not necessarily require that the protein, polypeptide, peptide or oligopeptide so described has been physically removed from a cell in which it was synthesized.

[0056] The term “polypeptide fragment” as used herein refers to a polypeptide that has a deletion, e.g., an amino-terminal and/or carboxy-terminal deletion compared to a full-length polypeptide, such as a naturally occurring protein. In an embodiment, the polypeptide fragment is a contiguous sequence in which the amino acid sequence of the fragment is identical to the corresponding positions in the naturally-occurring sequence. Fragments typically are at least 5, 6, 7, 8, 9 or 10 amino acids long, or at least 12, 14, 16 or 18 amino acids long, or at least 20 amino acids long, or at least 25, 30, 35, 40 or 45, amino acids, or at least 50 or 60 amino acids long, or at least 70 amino acids long.

[0057] The term “fusion protein” refers to a polypeptide comprising a polypeptide or fragment coupled to heterologous amino acid sequences. Fusion proteins are useful because they can be constructed to contain two or more desired functional elements that can be from two or more different proteins. A fusion protein comprises at least 10 contiguous amino acids from a polypeptide of interest, or at least 20 or 30 amino acids, or at least 40, 50 or 60 amino acids, or at least 75, 100 or 125 amino acids. The heterologous polypeptide included within the fusion protein is usually at least 6 amino acids in length, or at least 8 amino acids in length, or at least 15, 20, or 25 amino acids in length. Fusions that include larger polypeptides, such as an IgG Fc region, and even entire proteins, such as the green fluorescent protein (“GFP”) chromophore-containing proteins, have particular utility. Fusion

proteins can be produced recombinantly by constructing a nucleic acid sequence which encodes the polypeptide or a fragment thereof in frame with a nucleic acid sequence encoding a different protein or peptide and then expressing the fusion protein. Alternatively, a fusion protein can be produced chemically by crosslinking the polypeptide or a fragment thereof to another protein.

[0058] As used herein, a protein has “homology” or is “homologous” to a second protein if the nucleic acid sequence that encodes the protein has a similar sequence to the nucleic acid sequence that encodes the second protein. Alternatively, a protein has homology to a second protein if the two proteins have similar amino acid sequences. (Thus, the term 10 “homologous proteins” is defined to mean that the two proteins have similar amino acid sequences.) As used herein, homology between two regions of amino acid sequence (especially with respect to predicted structural similarities) is interpreted as implying similarity in function.

[0059] When “homologous” is used in reference to proteins or peptides, it is 15 recognized that residue positions that are not identical often differ by conservative amino acid substitutions. A “conservative amino acid substitution” is one in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases 20 where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity or degree of homology may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well known to those of skill in the art. See, e.g., Pearson, 1994, Methods Mol. Biol. 24:307-31 and 25:365-89.

25 [0060] The following six groups each contain amino acids that are conservative substitutions for one another: 1) Serine, Threonine; 2) Aspartic Acid, Glutamic Acid; 3) Asparagine, Glutamine; 4) Arginine, Lysine; 5) Isoleucine, Leucine, Methionine, Alanine, Valine, and 6) Phenylalanine, Tyrosine, Tryptophan.

[0061] Sequence homology for polypeptides, which is also referred to as percent 30 sequence identity, is typically measured using sequence analysis software. See, e.g., the Sequence Analysis Software Package of the Genetics Computer Group (GCG), University of

Wisconsin Biotechnology Center, 910 University Avenue, Madison, Wis. 53705. Protein analysis software matches similar sequences using a measure of homology assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, GCG contains programs such as "Gap" and "Bestfit" which can 5 be used with default parameters to determine sequence homology or sequence identity between closely related polypeptides, such as homologous polypeptides from different species of organisms or between a wild-type protein and a mutein thereof. See, e.g., GCG Version 6.1.

[0062] An exemplary algorithm when comparing a particular polypeptide sequence to 10 a database containing a large number of sequences from different organisms is the computer program BLAST (Altschul et al., J. Mol. Biol. 215:403-410 (1990); Gish and States, Nature Genet. 3:266-272 (1993); Madden et al., Meth. Enzymol. 266:131-141 (1996); Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997); Zhang and Madden, Genome Res. 7:649-656 (1997)), especially blastp or tblastn (Altschul et al., Nucleic Acids Res. 25:3389-3402 15 (1997)).

[0063] Exemplary parameters for BLASTp are: Expectation value: 10 (default); Filter: seg (default); Cost to open a gap: 11 (default); Cost to extend a gap: 1 (default); Max. alignments: 100 (default); Word size: 11 (default); No. of descriptions: 100 (default); Penalty Matrix: BLOWSUM62. The length of polypeptide sequences compared for homology will 20 generally be at least about 16 amino acid residues, or at least about 20 residues, or at least about 24 residues, or at least about 28 residues, or more than about 35 residues. When searching a database containing sequences from a large number of different organisms, it may be useful to compare amino acid sequences. Database searching using amino acid sequences can be measured by algorithms other than blastp known in the art. For instance, polypeptide 25 sequences can be compared using FASTA, a program in GCG Version 6.1. FASTA provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. Pearson, Methods Enzymol. 183:63-98 (1990). For example, percent sequence identity between amino acid sequences can be determined using FASTA with its default parameters (a word size of 2 and the PAM250 scoring matrix), as provided in GCG 30 Version 6.1, herein incorporated by reference.

[0064] In some embodiments, polymeric molecules (e.g., a polypeptide sequence or nucleic acid sequence) are considered to be "homologous" to one another if their sequences

are at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, at least 5

30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% similar. The term “homologous” necessarily refers to a comparison between at least two sequences (nucleotides sequences or amino acid sequences). In some embodiments, two nucleotide sequences are considered to be homologous if the polypeptides they encode 10 are at least about 50% identical, at least about 60% identical, at least about 70% identical, at least about 80% identical, or at least about 90% identical for at least one stretch of at least about 20 amino acids. In some embodiments, homologous nucleotide sequences are characterized by the ability to encode a stretch of at least 4-5 uniquely specified amino acids. Both the identity and the approximate spacing of these amino acids relative to one another 15 must be considered for nucleotide sequences to be considered homologous. In some embodiments of nucleotide sequences less than 60 nucleotides in length, homology is determined by the ability to encode a stretch of at least 4-5 uniquely specified amino acids. In some embodiments, two protein sequences are considered to be homologous if the proteins are at least about 50% identical, at least about 60% identical, at least about 70% identical, at least about 80% identical, or at least about 90% identical for at least one stretch of at least 20 20 about 20 amino acids.

[0065] As used herein, a “modified derivative” refers to polypeptides or fragments thereof that are substantially homologous in primary structural sequence to a reference polypeptide sequence but which include, e.g., in vivo or in vitro chemical and biochemical 25 modifications or which incorporate amino acids that are not found in the reference polypeptide. Such modifications include, for example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labeling, e.g., with radionuclides, and various enzymatic modifications, as will be readily appreciated by those skilled in the art. A variety of methods for labeling polypeptides and of substituents or labels useful for such purposes 30 are well known in the art, and include radioactive isotopes such as  $^{125}\text{I}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , and  $^3\text{H}$ , ligands that bind to labeled antiligands (e.g., antibodies), fluorophores, chemiluminescent agents, enzymes, and antiligands that can serve as specific binding pair members for a labeled ligand. The choice of label depends on the sensitivity required, ease of conjugation with the

primer, stability requirements, and available instrumentation. Methods for labeling polypeptides are well known in the art. See, e.g., Ausubel et al., Current Protocols in Molecular Biology, Greene Publishing Associates (1992, and Supplements to 2002).

[0066] As used herein, “polypeptide mutant” or “mutein” refers to a polypeptide

5 whose sequence contains an insertion, duplication, deletion, rearrangement or substitution of one or more amino acids compared to the amino acid sequence of a reference protein or polypeptide, such as a native or wild-type protein. A mutein may have one or more amino acid point substitutions, in which a single amino acid at a position has been changed to another amino acid, one or more insertions and/or deletions, in which one or more amino acids are inserted or deleted, respectively, in the sequence of the reference protein, and/or truncations of the amino acid sequence at either or both the amino or carboxy termini. A mutein may have the same or a different biological activity compared to the reference protein.

10

[0067] In some embodiments, a mutein has, for example, at least 85% overall

15 sequence homology to its counterpart reference protein. In some embodiments, a mutein has at least 90% overall sequence homology to the wild-type protein. In other embodiments, a mutein exhibits at least 95% sequence identity, or 98%, or 99%, or 99.5% or 99.9% overall sequence identity.

[0068] As used herein, a “polypeptide tag for affinity purification” is any polypeptide

20 that has a binding partner that can be used to isolate or purify a second protein or polypeptide sequence of interest fused to the first “tag” polypeptide. Several examples are well known in the art and include a His-6 tag, a FLAG epitope, a c-myc epitope, a Strep-TAGII, a biotin tag, a glutathione 5-transferase (GST), a chitin binding protein (CBP), a maltose binding protein (MBP), or a metal affinity tag.

25 [0069] As used herein, “recombinant” refers to a biomolecule, e.g., a gene or protein,

that (1) has been removed from its naturally occurring environment, (2) is not associated with all or a portion of a polynucleotide in which the gene is found in nature, (3) is operatively linked to a polynucleotide which it is not linked to in nature, or (4) does not occur in nature. The term “recombinant” can be used in reference to cloned DNA isolates, chemically synthesized polynucleotide analogs, or polynucleotide analogs that are biologically synthesized by heterologous systems, as well as proteins and/or mRNAs encoded by such

nucleic acids. Thus, for example, a protein synthesized by a microorganism is recombinant, for example, if it is synthesized from an mRNA synthesized from a recombinant gene present in the cell.

[0070] The term “polynucleotide”, “nucleic acid molecule”, “nucleic acid”, or

5 “nucleic acid sequence” refers to a polymeric form of nucleotides of at least 10 bases in length. The term includes DNA molecules (e.g., cDNA or genomic or synthetic DNA) and RNA molecules (e.g., mRNA or synthetic RNA), as well as analogs of DNA or RNA containing non-natural nucleotide analogs, non-native internucleoside bonds, or both. The nucleic acid can be in any topological conformation. For instance, the nucleic acid can be  
10 single-stranded, double-stranded, triple-stranded, quadruplexed, partially double-stranded, branched, hairpinned, circular, or in a padlocked conformation.

[0071] A “synthetic” RNA, DNA or a mixed polymer is one created outside of a cell, for example one synthesized chemically.

[0072] The term “nucleic acid fragment” as used herein refers to a nucleic acid

15 sequence that has a deletion, e.g., a 5'-terminal or 3'-terminal deletion compared to a full-length reference nucleotide sequence. In an embodiment, the nucleic acid fragment is a contiguous sequence in which the nucleotide sequence of the fragment is identical to the corresponding positions in the naturally-occurring sequence. In some embodiments fragments are at least 10, 15, 20, or 25 nucleotides long, or at least 20, 30, 40, 50, 60, 70, 80,  
20 90, 100, 110, 120, 130, 140, or 150 nucleotides long. In some embodiments a fragment of a nucleic acid sequence is a fragment of an open reading frame sequence. In some embodiments such a fragment encodes a polypeptide fragment (as defined herein) of the protein encoded by the open reading frame nucleotide sequence.

[0073] As used herein, an endogenous nucleic acid sequence in the genome of an

25 organism (or the encoded protein product of that sequence) is deemed “recombinant” herein if a heterologous sequence is placed adjacent to the endogenous nucleic acid sequence, such that the expression of this endogenous nucleic acid sequence is altered. In this context, a heterologous sequence is a sequence that is not naturally adjacent to the endogenous nucleic acid sequence, whether or not the heterologous sequence is itself endogenous (originating  
30 from the same host cell or progeny thereof) or exogenous (originating from a different host cell or progeny thereof). By way of example, a promoter sequence can be substituted (e.g., by

homologous recombination) for the native promoter of a gene in the genome of a host cell, such that this gene has an altered expression pattern. This gene would now become “recombinant” because it is separated from at least some of the sequences that naturally flank it.

5 [0074] A nucleic acid is also considered “recombinant” if it contains any modifications that do not naturally occur to the corresponding nucleic acid in a genome. For instance, an endogenous coding sequence is considered “recombinant” if it contains an insertion, deletion or a point mutation introduced artificially, e.g., by human intervention. A “recombinant nucleic acid” also includes a nucleic acid integrated into a host cell

10 chromosome at a heterologous site and a nucleic acid construct present as an episome.

[0075] As used herein, the phrase “degenerate variant” of a reference nucleic acid sequence encompasses nucleic acid sequences that can be translated, according to the standard genetic code, to provide an amino acid sequence identical to that translated from the reference nucleic acid sequence. The term “degenerate oligonucleotide” or “degenerate primer” is used to signify an oligonucleotide capable of hybridizing with target nucleic acid sequences that are not necessarily identical in sequence but that are homologous to one another within one or more particular segments.

[0076] The term “percent sequence identity” or “identical” in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over a stretch of at least about nine nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32, and even more typically at least about 36 or more nucleotides. There are a number of different algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using FASTA, Gap or Bestfit, which are programs in Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wis. FASTA provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. Pearson, Methods Enzymol. 183:63-98 (1990). For instance, percent sequence identity between nucleic acid sequences can be determined using FASTA with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) or using Gap with its default parameters as provided in GCG Version 6.1, herein incorporated by reference. Alternatively, sequences can

be compared using the computer program, BLAST (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990); Gish and States, *Nature Genet.* 3:266-272 (1993); Madden et al., *Meth. Enzymol.* 266:131-141 (1996); Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997); Zhang and Madden, *Genome Res.* 7:649-656 (1997)), especially blastp or tblastn (Altschul et al.,

5 *Nucleic Acids Res.* 25:3389-3402 (1997)).

[0077] The term “substantial homology” or “substantial similarity,” when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 76%, 80%, 85%, or at least about 90%,  
10 or at least about 95%, 96%, 97%, 98% or 99% of the nucleotide bases, as measured by any well-known algorithm of sequence identity, such as FASTA, BLAST or Gap, as discussed above.

[0078] Alternatively, substantial homology or similarity exists when a nucleic acid or fragment thereof hybridizes to another nucleic acid, to a strand of another nucleic acid, or to  
15 the complementary strand thereof, under stringent hybridization conditions. “Stringent hybridization conditions” and “stringent wash conditions” in the context of nucleic acid hybridization experiments depend upon a number of different physical parameters. Nucleic acid hybridization will be affected by such conditions as salt concentration, temperature, solvents, the base composition of the hybridizing species, length of the complementary regions, and the number of nucleotide base mismatches between the hybridizing nucleic acids, as will be readily appreciated by those skilled in the art. One having ordinary skill in the art knows how to vary these parameters to achieve a particular stringency of hybridization.

[0079] In general, “stringent hybridization” is performed at about 25°C below the  
25 thermal melting point (Tm) for the specific DNA hybrid under a particular set of conditions. “Stringent washing” is performed at temperatures about 5°C lower than the Tm for the specific DNA hybrid under a particular set of conditions. The Tm is the temperature at which 50% of the target sequence hybridizes to a perfectly matched probe. See Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press,  
30 Cold Spring Harbor, N.Y. (1989), page 9.51, hereby incorporated by reference. For purposes herein, “stringent conditions” are defined for solution phase hybridization as aqueous hybridization (i.e., free of formamide) in 6xSSC (where 20xSSC contains 3.0 M NaCl and

0.3 M sodium citrate), 1% SDS at 65°C for 8-12 hours, followed by two washes in 0.2xSSC, 0.1% SDS at 65°C for 20 minutes. It will be appreciated by the skilled worker that hybridization at 65°C will occur at different rates depending on a number of factors including the length and percent identity of the sequences which are hybridizing.

5 [0080] As used herein, an “expression control sequence” refers to polynucleotide sequences which are necessary to affect the expression of coding sequences to which they are operatively linked. Expression control sequences are sequences which control the transcription, post-transcriptional events and translation of nucleic acid sequences. Expression control sequences include appropriate transcription initiation, termination, 10 promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (e.g., ribosome binding sites); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences 15 generally include promoter, ribosomal binding site, and transcription termination sequence. The term “control sequences” is intended to encompass, at a minimum, any component whose presence is essential for expression, and can also encompass an additional component whose presence is advantageous, for example, leader sequences and fusion partner sequences.

20 [0081] As used herein, “operatively linked” or “operably linked” expression control sequences refers to a linkage in which the expression control sequence is contiguous with the gene of interest to control the gene of interest, as well as expression control sequences that act in trans or at a distance to control the gene of interest.

25 [0082] As used herein, a “vector” is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid,” which generally refers to a circular double stranded DNA loop into which additional DNA segments may be ligated, but also includes linear double-stranded molecules such as those resulting from amplification by the polymerase chain reaction (PCR) or from treatment of a circular plasmid with a restriction enzyme. Other vectors include cosmids, bacterial artificial chromosomes (BAC) and yeast artificial chromosomes (YAC). Another 30 type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome (discussed in more detail below). Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., vectors having an origin of

replication which functions in the host cell). Other vectors can be integrated into the genome of a host cell upon introduction into the host cell, and are thereby replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply “expression vectors”).

[0083] The term “recombinant host cell” (or simply “recombinant cell” or “host cell”), as used herein, is intended to refer to a cell into which a recombinant nucleic acid such as a recombinant vector has been introduced. In some instances the word “cell” is replaced by a name specifying a type of cell. For example, a “recombinant microorganism” is a recombinant host cell that is a microorganism host cell and a “recombinant cyanobacteria” is a recombinant host cell that is a cyanobacteria host cell. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term “recombinant host cell,” “recombinant cell,” and “host cell”, as used herein. A recombinant host cell may be an isolated cell or cell line grown in culture or may be a cell which resides in a living tissue or organism.

[0084] As used herein, the term “heterotrophic” refers to an organism that cannot fix carbon and uses organic carbon for growth.

[0085] As used herein, the term “autotrophic” refers to an organism that produces complex organic compounds (such as carbohydrates, fats, and proteins) from simple inorganic molecules using energy from light (by photosynthesis) or inorganic chemical reactions (chemosynthesis).

[0086] As used herein, “muscle mass” refers to the weight of muscle in a subject’s body. Muscle mass includes the skeletal muscles, smooth muscles (such as cardiac and digestive muscles) and the water contained in these muscles. Muscle mass of specific muscles can be determined using dual energy x-ray absorptiometry (DEXA) (Padden-Jones et al., 2004). Total lean body mass (minus the fat), total body mass, and bone mineral content can be measured by DEXA as well. In some embodiments a change in the muscle mass of a specific muscle of a subject is determined, for example by DEXA, and the change is used as a

proxy for the total change in muscle mass of the subject. Thus, for example, if a subject consumes a nutritive protein as disclosed herein and experiences an increase over a period of time in muscle mass in a particular muscle or muscle group, it can be concluded that the subject has experienced an increase in muscle mass.

5 [0001] As used herein, a “muscle strength” refers to the amount of force a muscle can produce with a single maximal effort. There are two types of muscle strength, static strength and dynamic strength. Static strength refers to isometric contraction of a muscle, where a muscle generates force while the muscle length remains constant and/or when there is no movement in a joint. Examples include holding or carrying an object, or pushing against a  
10 wall. Dynamic strength refers to a muscle generating force that results in movement. Dynamic strength can be isotonic contraction, where the muscle shortens under a constant load or isokinetic contraction, where the muscle contracts and shortens at a constant speed. Dynamic strength can also include isoinertial strength.

15 [0002] Unless specified, “muscle strength” refers to maximum dynamic muscle strength. Maximum strength is referred to as “one repetition maximum” (1RM). This is a measurement of the greatest load (in kilograms) that can be fully moved (lifted, pushed or pulled) once without failure or injury. This value can be measured directly, but doing so requires that the weight is increased until the subject fails to carry out the activity to completion. Alternatively, 1RM is estimated by counting the maximum number of exercise  
20 repetitions a subject can make using a load that is less than the maximum amount the subject can move. Leg extension and leg flexion are often measured in clinical trials (Borsheim et al., “Effect of amino acid supplementation on muscle mass, strength and physical function in elderly,” Clin Nutr 2008;27:189-195; Paddon-Jones, et al., “Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bed  
25 rest,” J Clin Endocrinol Metab 2004;89:4351-4358).

30 [0003] As used herein, “functional performance” refers to a functional test that simulates daily activities. “Functional performance” is measured by any suitable accepted test, including timed-step test (step up and down from a 4 inch bench as fast as possible 5 times), timed floor transfer test (go from a standing position to a supine position on the floor and thereafter up to a standing position again as fast as possible for one repetition), and physical performance battery test (static balance test, chair test, and a walking test) (Borsheim et al., “Effect of amino acid supplementation on muscle mass, strength and physical function in elderly,” Clin Nutr 2008;27:189-195).

[0004] As used herein, a “body mass index” or “BMI” or “Quetelet index” is a subject’s weight in kilograms divided by the square of the subject’s height in meters ( $\text{kg}/\text{m}^2$ ).

- [0087] For adults, a frequent use of the BMI is to assess how much an individual's body weight departs from what is normal or desirable for a person of his or her height. The 5 weight excess or deficiency may, in part, be accounted for by body fat, although other factors such as muscularity also affect BMI significantly. The World Health Organization regards a BMI of less than 18.5 as underweight and may indicate malnutrition, an eating disorder, or other health problems, while a BMI greater than 25 is considered overweight and above 30 is considered obese. (World Health Organization. BMI classification. Accessed March 19, 10 2012 [http://apps.who.int/bmi/index.jsp?introPage=intro\\_3.html](http://apps.who.int/bmi/index.jsp?introPage=intro_3.html).) As used herein a “desirable body mass index” is a body mass index of from about 18.5 to about 25. Thus, if a subject has a BMI below about 18.5, then an increase in the subject’s BMI is an increase in the desirability of the subject’s BMI. If instead a subject has a BMI above about 25, then a decrease in the subject’s BMI is an increase in the desirability of the subject’s BMI.
- 15 [0088] As used herein, an “elderly” mammal is one who experiences age related changes in at least one of body mass index and muscle mass (e.g., age related sarcopenia). In some embodiments an “elderly” human is at least 50 years old, at least 60 years old, at least 65 years old, at least 70 years old, at least 75 years old, at least 80 years old, at least 85 years old, at least 90 years old, at least 95 years old, or at least 100 years old. In some 20 embodiments and an elderly animal, mammal, or human is a human who has experienced a loss of muscle mass from peak lifetime muscle mass of at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, or at least 60%. Because age related changes to at least one of body mass index and muscle mass are known to correlate with increasing age, in some embodiments an 25 elderly mammal is identified or defined simply on the basis of age. Thus, in some embodiments an “elderly” human is identified or defined simply by the fact that their age is at least 60 years old, at least 65 years old, at least 70 years old, at least 75 years old, at least 80 years old, at least 85 years old, at least 90 years old, at least 95 years old, or at least 100 years old, and without recourse to a measurement of at least one of body mass index and 30 muscle mass.

[0005] As used herein, a patient is “critically-medically ill” if the patient, because of medical illness, experiences changes in at least one of body mass index and muscle mass

(e.g., sarcopenia). In some embodiments the patient is confined to bed for at least 25%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% of their waking time. In some embodiments the patient is unconscious. In some embodiments the patient has been confined to bed as described in this paragraph for at least 1 day, 2 days, 3 days, 4 days, 5 days, 10 days, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 10 weeks or longer.

5 [0006] As used herein, “protein-energy malnutrition” refers to a form of malnutrition where there is inadequate protein intake. Types include Kwashiorkor (protein malnutrition predominant), Marasmus (deficiency in both calorie and protein nutrition), and Marasmic Kwashiorkor (marked protein deficiency and marked calorie insufficiency signs present, 10 sometimes referred to as the most severe form of malnutrition).

10 [0007] As used herein, “cachexia” refers to a multifaceted clinical syndrome that results in wasting and weight loss. It is a complex condition where protein catabolism exceeds protein anabolism, which makes muscle wasting a primary feature of the condition. In addition to the metabolic derangements in protein metabolism, it is also 15 characterized by anorexia and inflammation. These derangements plus impaired protein metabolism are responsive to nutrition therapy to varying degrees.

15 [0008] As used herein, “thermogenesis” is the process of heat production in a mammal. Thermogenesis is accompanied by an increase in energy expenditure. Thermogenesis is specifically the energy burned following the metabolism of a food 20 component (such as protein). This may also be referred to as the thermic effect of food. Total energy expenditure by an individual equals the sum of resting energy expenditure (energy consumed at rest in a fasting state to support basal metabolism), the thermic effect of food, and energy expenditure related to physical activity. Resting energy expenditure accounts for about 65-75% of total energy expenditure in humans. The amount and activity 25 of muscle mass is one influencer of resting energy expenditure. Adequate protein consumption to support muscle also influences resting energy expenditure. The ingestion of protein tends to increase energy expenditure following a meal; this is the thermic effect of food. The thermic effect of food accounts for about 10% of total energy expenditure in humans. While this is a small proportion of total energy expenditure, small increases in this 30 value can impact body weight. Protein has a higher thermic effect than fat or carbohydrate; this effect along with other metabolic influences of protein make it a useful substrate for weight control, diabetes management and other conditions.

[0009] As used herein, “satiation” is the act of becoming full while eating or a reduced desire to eat. This halts or diminishes eating.

[0010] As used herein, “satiety” is the act of remaining full after a meal which manifests as the period of no eating follow the meal.

5 [0011] As used herein, “exercise” is, most broadly, any bodily activity that enhances or maintains physical fitness and overall health and wellness. Exercise is performed for various reasons including strengthening muscles and the cardiovascular system, honing athletic skills, weight loss or maintenance, as well as for the purpose of enjoyment.

10 [0012] As used herein, a “sufficient amount” is an amount of a protein or polypeptide disclosed herein that is sufficient to cause a desired effect. For example, if an increase in muscle mass is desired, a sufficient amount is an amount that causes an increase in muscle mass in a subject over a period of time. A sufficient amount of a protein or polypeptide fragment can be provided directly, i.e., by administering the protein or polypeptide fragment to a subject, or it can be provided as part of a composition comprising the protein or 15 polypeptide fragment. Modes of administration are discussed elsewhere herein.

20 [0089] As used herein, the term “mammal” refers to any member of the taxonomic class mammalia, including placental mammals and marsupial mammals. Thus, “mammal” includes humans, primates, livestock, and laboratory mammals. Exemplary mammals include a rodent, a mouse, a rat, a rabbit, a dog, a cat, a sheep, a horse, a goat, a llama, cattle, a primate, a pig, and any other mammal. In some embodiments, the mammal is at least one of a transgenic mammal, a genetically-engineered mammal, and a cloned mammal.

#### A. Nutritive Proteins

25 [0090] For the purposes of this disclosure, a “nutritive protein” is a protein that contains a desirable amount of essential amino acids. In some embodiments, the nutritive protein comprises at least 30% essential amino acids by weight. In some embodiments, the nutritive protein comprises at least 40% essential amino acids by weight. In some embodiments, the nutritive protein comprises at least 50% essential amino acids by weight. In some embodiments the nutritive protein comprises or consists of a protein or fragment of a protein that naturally occurs in an edible species. In its broadest sense, an “edible species” 30 encompasses any species known to be eaten without deleterious effect by at least one type of mammal. A deleterious effect includes a poisonous effect and a toxic effect. In some

embodiments an edible species is a species known to be eaten by humans without deleterious effect. Some edible species are an infrequent but known component of the diet of only a small group of a type of mammal in a limited geographic location while others are a dietary staple throughout much of the world. In other embodiments an edible species is one not known to be previously eaten by any mammal, but that is demonstrated to be edible upon testing. Edible species include but are not limited to *Gossypium turneri*, *Pleurotus cornucopiae*, *Glycine max*, *Oryza sativa*, *Thunnus obesus*, *Abies bracteata*, *Acomys ignitus*, *Lathyrus aphaca*, *Bos gaurus*, *Raphicerus melanotis*, *Phoca groenlandica*, *Acipenser sinensis*, *Viverra tangalunga*, *Pleurotus sajor-caju*, *Fagopyrum tataricum*, *Pinus strobus*, *Ipomoea nil*, *Taxus cuspidata*, *Ipomoea wrightii*, *Mya arenaria*, *Actinidia deliciosa*, *Gazella granti*, *Populus tremula*, *Prunus domestica*, *Larus argentatus*, *Vicia villosa*, *Sargocentron punctatissimum*, *Silene latifolia*, *Lagenodelphis hosei*, *Spisula solidissima*, *Crossarchus obscurus*, *Phaseolus angularis*, *Lathyrus vestitus*, *Oncorhynchus gorbuscha*, *Alligator mississippiensis*, *Pinus halepensis*, *Larus canus*, *Brassica napus*, *Silene cucubalus*, *Phoca fasciata*, *Gazella bennettii*, *Pinus taeda*, *Taxus canadensis*, *Zamia furfuracea*, *Pinus yunnanensis*, *Pinus wallichiana*, *Asparagus officinalis*, *Capsicum baccatum*, *Pinus longaeva*, *Taxus baccata*, *Pinus sibirica*, *Citrus sinensis*, *Sargocentron xantherythrum*, *Bison bison*, *Gazella thomsonii*, *Vicia sativa*, *Branta canadensis*, *Apium graveolens*, *Acer campestre*, *Coriandrum sativum*, *Silene conica*, *Lactuca sativa*, *Capsicum chinense*, *Abies veitchii*, *Capra hircus*, *Gazella spekei*, *Oncorhynchus keta*, *Ipomoea obscura*, *Cucumis melo* var. *conomon*, *Phoca hispida*, *Vulpes vulpes*, *Ipomoea quamoclit*, *Solanum habrochaites*, *Populus* sp., *Pinus rigida*, *Quercus lyrata*, *Phaseolus coccineus*, *Larus ridibundus*, *Sargocentron spiniferum*, *Thunnus thynnus*, *Vulpes lagopus*, *Bos gaurus frontalis*, *Acer opalus*, *Acer palmatum*, *Quercus ilex*, *Pinus mugo*, *Grus antigone*, *Pinus uncinata*, *Prunus mume*, *Oncorhynchus tschawytscha*, *Gazella subgutturosa*, *Vulpes zerda*, *Pinus coulteri*, *Gossypium barbadense*, *Acer pseudoplatanus*, *Oncorhynchus nerka*, *Sus barbatus*, *Fagopyrum esculentum* subsp. *Ancestrale*, *Cynara cardunculus*, *Phaseolus aureus*, *Populus nigra*, *Gossypium schwendimanii*, *Solanum chacoense*, *Quercus rubra*, *Cucumis sativus*, *Equus burchelli*, *Oncorhynchus kisutch*, *Pinus radiata*, *Phoca vitulina richardsi*, *Grus nigricollis*, *Abies grandis*, *Oncorhynchus masou*, *Spinacia oleracea*, *Solanum chilense*, *Addax nasomaculatus*, *Ipomoea batatas*, *Equus grevyi*, *Abies sachalinensis*, *Pinus pinea*, *Hipposideros commersoni*, *Crocus nudiflorus*, *Citrus maxima*, *Acipenser transmontanus*, *Gossypium gossypoides*, *Viverra zibetha*, *Quercus cerris*, *Anser indicus*, *Pinus balfouriana*, *Silene otites*, *Oncorhynchus* sp., *Viverra megaspila*, *Bos mutus grunniens*, *Pinus elliottii*,

*Equus hemionus kulan, Capra ibex ibex, Allium sativum, Raphanus sativus, Pinus echinata, Prunus serotina, Sargocentron diadema, Silene gallica, Brassica oleracea, Daucus carota, Oncorhynchus mykiss, Brassica oleracea var. alboglabra, Gossypium hirsutum, Abies alba, Citrus reticulata, Cichorium intybus, Bos sauveli, Lama glama, Zea mays, Acorus gramineus,*

**5** *Vulpes macrotis, Ovis ammon darwini, Raphicerus sharpei, Pinus contorta, Bos indicus, Capra sibirica, Pinus ponderosa, Prunus dulcis, Solanum sotgarandinum, Ipomoea aquatica, Lagenorhynchus albirostris, Ovis canadensis, Prunus avium, Gazella dama, Thunnus alalunga, Silene pratensis, Pinus cembra, Crocus sativus, Citrullus lanatus, Gazella rufifrons, Brassica tournefortii, Capra falconeri, Bubalus mindorensis, Pinus palustris,*

**10** *Prunus laurocerasus, Grus vipio, Ipomoea purpurea, Pinus leiophylla, Lagenorhynchus obscurus, Raphicerus campestris, Brassica rapa subsp. Pekinensis, Acmella radicans, Ipomoea triloba, Pinus patula, Cucumis melo, Pinus virginiana, Solanum lycopersicum, Pinus densiflora, Pinus engelmannii, Quercus robur, Ipomoea setosa, Pleurotus djamor, Hipposideros diadema, Ovis aries, Sargocentron microstoma, Brassica oleracea var. italica,*

**15** *Capra cylindricornis, Populus kitakamiensis, Allium textile, Vicia faba, Fagopyrum esculentum, Bison priscus, Quercus suber, Lagophylla ramosissima, Acrantophis madagascariensis, Acipenser baerii, Capsicum annuum, Triticum aestivum, Xenopus laevis, Phoca sibirica, Acipenser naccarii, Actinidia chinensis, Ovis dalli, Solanum tuberosum, Bubalus carabanensis, Citrus jambhiri, Bison bonasus, Equus asinus, Bubalus*

**20** *depressicornis, Pleurotus eryngii, Solanum demissum, Ovis vignei, Zea mays subsp. Parviflumis, Lathyrus tingitanus, Welwitschia mirabilis, Grus rubicunda, Ipomoea coccinea, Allium cepa, Gazella soemmerringii, Brassica rapa, Lama vicugna, Solanum peruvianum, Xenopus borealis, Capra caucasica, Thunnus albacares, Equus zebra, Gallus gallus, Solanum bulbocastanum, Hipposideros terasensis, Lagenorhynchus acutus, Hippopotamus*

**25** *amphibius, Pinus koraiensis, Acer monspessulanum, Populus deltoides, Populus trichocarpa, Acipenser guldenstadtii, Pinus thunbergii, Brassica oleracea var. capitata, Abyssocottus korotneffi, Gazella cuvieri, Abies homolepis, Abies holophylla, Gazella gazella, Pinus parviflora, Brassica oleracea var. acephala, Cucurbita pepo, Pinus armandii, Abies mariesii, Thunnus thynnus orientalis, Citrus unshiu, Solanum cheesmanii, Lagenorhynchus*

**30** *obliquidens, Acer platanoides, Citrus limon, Acrantophis dumerili, Solanum commersonii, Gossypium arboreum, Prunus persica, Pleurotus ostreatus, Abies firma, Gazella leptoceros, Salmo salar, Homarus americanus, Abies magnifica, Bos javanicus, Phoca largha, Sus cebifrons, Solanum melongena, Phoca vitulina, Pinus sylvestris, Zamia floridana, Vulpes corsac, Allium porrum, Phoca caspica, Vulpes chama, Taxus chinensis, Brassica oleracea*

*var. botrytis, Anser anser anser, Phaseolus lunatus, Brassica campestris, Acer saccharum, Pinus pumila, Solanum pennellii, Pinus edulis, Ipomoea cordatotriloba, Populus alba, Oncorhynchus clarki, Quercus petraea, Sus verrucosus, Equus caballus przewalskii, Populus euphratica, Xenopus tropicalis, Taxus brevifolia, Lama guanicoe, Pinus banksiana, Solanum 5 nigrum, Sus celebensis, Brassica juncea, Lagenorhynchus cruciger, Populus tremuloides, Pinus pungens, Bubalus quarlesi, Quercus gamelliflora, Ovis orientalis musimon, Bubalus bubalis, Pinus luchuensis, Sus philippensis, Phaseolus vulgaris, Salmo trutta, Acipenser persicus, Solanum brevidens, Pinus resinosa, Hippotragus niger, Capra nubiana, Asparagus scaber, Ipomoea platensis, Sus scrofa, Capra aegagrus, Lathyrus sativus, Sargocentron tiere, 10 Hippoglossus hippoglossus, Acorus americanus, Equus caballus, Bos taurus, Barbarea vulgaris, Lama guanicoe pacos, Pinus pinaster, Octopus vulgaris, Solanum crispum, Hippotragus equinus, Equus burchellii antiquorum, Crossarchus alexandri, Ipomoea alba, Triticum monococcum, Populus jackii, Lagenorhynchus australis, Gazella dorcas, Quercus coccifera, Anser caerulescens, Acorus calamus, Pinus roxburghii, Pinus tabuliformis, Zamia 15 fischeri, Grus carunculatus, Acomys cahirinus, Cucumis melo var. reticulatus, Gallus lafayettei, Pisum sativum, Pinus attenuata, Pinus clausa, Gazella saudiya, Capra ibex, Ipomoea trifida, Zea luxurians, Pinus krempfii, Acomys wilsoni, Petroselinum crispum, Quercus palustris, Triticum timopheevi, Meleagris gallopavo, Brassica oleracea, Brassica oleracea, Beta vulgaris, Solanum lycopersicum, Phaseolus vulgaris, Xiphias gladius, Morone 20 saxatilis, Micropterus salmoides, Placopecten magellanicus, Sprattus sprattus, Clupea harengus, Engraulis encrasicolus, Cucurbita maxima, Agaricus bisporus, Musa acuminata x balbisiana, Malus domestica, Cicer arietinum, Anas platyrhynchos, Vaccinium macrocarpum, Rubus idaeus x strigosus, Vaccinium angustifolium, Fragaria ananassa, Rubus fruticosus, Cucumis melo, Ananas comosus, Cucurbita pepo, Cucurbita moschata, Sus 25 scrofa domesticus, Ocimum basilicum, Rosmarinus officinalis, Foeniculum vulgare, Rheum rhabarbarum, Carica papaya, Mangifera indica, Actinidia deliciosa, Prunus armeniaca, Prunus avium, Cocos nucifera, Olea europaea, Pyrus communis, Ficus carica, Passiflora edulis, Oryza sativa subsp. Japonica, Oryza sativa subsp. Indica, Coturnix coturnix, Saccharomyces cerevisiae.*

30 [0013] In some embodiments the nutritive protein comprises or consists of a derivative or mutein of a protein or fragment of a protein that naturally occurs in an edible species. Such a nutritive protein may be referred to as an “engineered nutritive protein.” In such embodiments the natural protein or fragment thereof is a “reference” protein or polypeptide and the engineered nutritive protein or a first polypeptide sequence thereof

comprises at least one sequence modification relative to the amino acid sequence of the reference protein or polypeptide. For example, in some embodiments the engineered nutritive protein or first polypeptide sequence thereof is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one reference nutritive protein amino acid sequence. Typically the ratio of at least one of branch chain amino acid residues to total amino acid residues, essential amino acid residues to total amino acid residues, and leucine residues to total amino acid residues, present in the engineered nutritive protein or a first polypeptide sequence thereof is greater than the corresponding ratio of at least one of branch chain amino acid residues to total amino acid residues, essential amino acid residues to total amino acid residues, and leucine residues to total amino acid residues present in the reference nutritive protein or polypeptide sequence.

[0091] In some embodiments the nutritive protein is an abundant protein in food or a derivative or mutein thereof, or is a fragment of an abundant protein in food or a derivative or mutein thereof. An abundant protein is a protein that is present in a higher concentration in a food relative to other proteins present in the food. The food can be a known component of the diet of only a small group of a type of mammal in a limited geographic location, or a dietary staple throughout much of the world. In some embodiments the abundant protein in food is selected from chicken egg proteins such as ovalbumin, ovotransferrin, and ovomucoid; meat proteins such as myosin, actin, tropomyosin, collagen, and troponin; cereal proteins such as casein, alpha1 casein, alpha2 casein, beta casein, kappa casein, beta-lactoglobulin, alpha-lactalbumin, glycinin, beta-conglycinin, glutelin, prolamine, gliadin, glutenin, albumin, globulin; chicken muscle proteins such as albumin, enolase, creatine kinase, phosphoglycerate mutase, triosephosphate isomerase, apolipoprotein, ovotransferrin, phosphoglucomutase, phosphoglycerate kinase, glycerol-3-phosphate dehydrogenase, glyceraldehyde 3-phosphate dehydrogenase, hemoglobin, cofilin, glycogen phosphorylase, fructose-1,6-bisphosphatase, actin, myosin, tropomyosin a-chain, casein kinase, glycogen phosphorylase, fructose-1,6-bisphosphatase, aldolase, tubulin, vimentin, endoplasmic, lactate dehydrogenase, destrin, transthyretin, fructose bisphosphate aldolase, carbonic anhydrase, aldehyde dehydrogenase, annexin, adenosyl homocysteinase; pork muscle proteins such as actin, myosin, enolase, titin, cofilin, phosphoglycerate kinase, enolase, pyruvate dehydrogenase, glycogen phosphorylase, triosephosphate isomerase, myokinase; and fish

proteins such as parvalbumin, pyruvate dehydrogenase, desmin, and triosephosphate isomerase.

[0092] Three natural sources of protein generally regarded as good sources of high quality amino acids are whey protein, egg protein, and soy protein. Each source comprises 5 multiple proteins. Table 1 presents the weight proportional representation of each amino acid in the protein source (g AA / g protein) expressed as a percentage.

**Table 1**

Amino Acid	Whey	Egg	Soy
Isoleucine	6.5%	5.5%	5.0%
Leucine	11.0%	8.6%	8.0%
Lysine	9.1%	7.2%	6.3%
Methionine	2.1%	3.1%	1.3%
Phenylalanine	3.4%	5.3%	1.2%
Threonine	7.0%	4.8%	3.7%
Tryptophan	1.7%	1.2%	1.3%
Valine	6.2%	6.1%	4.9%
Histidine	2.0%	2.4%	2.7%
<i>Other</i>	51.7%	49.5%	60.4%

[0093] Table 2 presents the weight proportion of each protein source that is essential 10 amino acids, branched chain amino acids (L, I, and V), and leucine (L) (alone).

**Table 2**

<b>Protein Source</b>	<b>Essential Amino Acids</b>	<b>Branch Chain Amino Acids</b>	<b>Leucine</b>
Whey	49.0%	23.7%	11.0%
Egg	50.5%	20.1%	8.6%
Soy	39.6%	17.9%	8.0%

[0094] The sources relied on to determine the amino acid content of Whey are: Belitz HD., Grosch W., and Schieberle P. Food Chemistry (4th Ed). Springer-Verlag, Berlin Heidelberg 2009; <http://www.gnc.com/product/index.jsp?productId=2986027>;

5        [http://www.nutrabio.com/Products/whey\\_protein\\_concentrate.htm](http://www.nutrabio.com/Products/whey_protein_concentrate.htm); and [http://nutrabio.com/Products/whey\\_protein\\_isolate.htm](http://nutrabio.com/Products/whey_protein_isolate.htm). The amino acid content values from those sources were averaged to give the numbers presented in Tables 1 and 2. The source for soy protein is Egg, National Nutrient Database for Standard Reference, Release 24 (<http://ndb.nal.usda.gov/ndb/foods/list>). The source for soy protein is Self Nutrition Data

10      (<http://nutritiondata.self.com/facts/legumes-and-legume-products/4389/2>).

[0014] As used herein, “whey protein” or “whey” means a protein mixture comprising an amino acid composition according to Tables 1 and 2. As used herein, whey protein comprises 49% essential amino acids, 24% branch chain amino acids, and 11% leucine, by weight.

15      [0015] As used herein, “egg protein” or “egg” means a protein mixture comprising an amino acid composition according to Tables 1 and 2. As used herein, egg protein comprises 51% essential amino acids, 20% branch chain amino acids, and 9% leucine, by weight.

[0095] As used herein, “soy protein” or “soy” means a protein mixture comprising an amino acid composition according to Tables 1 and 2. As used herein, soy protein comprises 20 40% essential amino acids, 18% branch chain amino acids, and 8% leucine, by weight.

[0096] In some instances herein the portion of amino acid(s) of a particular type within a polypeptide, protein or a composition is quantified based on the weight ratio of the type of amino acid(s) to the total weight of amino acids present in the polypeptide, protein or composition in question. This value is calculated by dividing the weight of the particular

amino acid(s) in the polypeptide, protein or a composition by the weight of all amino acids present in the polypeptide, protein or a composition.

[0097] In other instances the ratio of a particular type of amino acid(s) residues present in a polypeptide or protein to the total number of amino acids present in the 5 polypeptide or protein in question is used. This value is calculated by dividing the number of the amino acid(s) in question that is present in each molecule of the polypeptide or protein by the total number of amino acid residues present in each molecule of the polypeptide or protein. A skilled artisan appreciates that these two methods are interchangeable and that the weight proportion of a type of amino acid(s) present in a polypeptide or protein can be 10 converted to a ratio of the particular type of amino acid residue(s), and vice versa.

[0098] In certain embodiments herein the weight proportion of branched chain amino acids, leucine, and/or essential amino acids in whey, egg, or soy is used as a benchmark to measure the amino acid composition of a polypeptide, a protein, or a composition comprising at least one of a polypeptide and a protein. In those embodiments it is understood that the 15 two measures are not completely equivalent, but it is also understood that the measures result in measurements that are similar enough to use for this purpose. For example, when a protein of interest is characterized as comprising a ratio of branch chain amino acid residues to total amino acid residues that is equal to or greater than 24% (the weight proportion of branch chain amino acid residues present in whey), that is a precise description of the branch chain 20 amino acid content of the protein. At the same time, the weight proportion of branch chain amino acid residues present in that protein is not necessarily exactly equal to 24%. Even so, the skilled artisan understands that this is a useful comparison. If provided with the total number of amino acid residues present in the protein of interest the skilled artisan can also determine the weight proportion of branch chain amino acid residues in the protein of 25 interest.

[0099] In some embodiments herein a fragment of a naturally-occurring nutritive protein is selected and optionally isolated. In some embodiments the fragment comprises at least 25 amino acids. In some embodiments the fragment comprises at least 50 amino acids. In some embodiments the fragment consists of at least 25 amino acids. In some 30 embodiments the fragment consists of at least 50 amino acids. In some embodiments an isolated recombinant nutritive protein is provided. In some embodiments the nutritive protein comprises a first polypeptide sequence, and the first polypeptide sequence comprises a fragment

of at least 25 or at least 50 amino acids of a naturally-occurring nutritive protein. In some embodiments the nutritive proteins is isolated. In some embodiments the nutritive proteins are recombinant. In some embodiments the nutritive proteins comprise a first polypeptide sequence comprising a fragment of at least 50 amino acids of a naturally-occurring nutritive protein. In some embodiments the nutritive proteins are isolated recombinant nutritive proteins. In some embodiments the isolated recombinant nutritive proteins disclosed herein are provided in a non-isolated and/or non-recombinant form

5 [00100] In some embodiments a nutritive protein according to this disclosure comprises a first polypeptide sequence comprising a fragment of a naturally-occurring nutritive protein. In some embodiments of the nutritive protein, the nutritive protein consists of the first polypeptide sequence. In some embodiments of the nutritive protein, the nutritive protein consists of the fragment of a naturally-occurring nutritive protein.

10 [00101] In some embodiments a nutritive protein according to this disclosure comprises a first polypeptide sequence that comprises ratio of branch chain amino acid residues to total amino acid residues that is equal to or greater than the ratio of branch chain amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. Thus, in such embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues that is equal to or greater than a ratio selected from 24%, 20%, and 18%.

15 [00102] In some embodiments a nutritive protein according to this disclosure comprises a first polypeptide sequence that comprises a ratio of L residues to total amino acid residues that is equal to or greater than the ratio of L residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. Thus, in such embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of L residues to total amino acid residues that is equal to or greater than a ratio selected from 11%, 9%, and 8%.

20 [00103] In some embodiments a nutritive protein according to this disclosure comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. Thus, in such embodiments the nutritive protein comprises a first polypeptide

sequence that comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than a ratio selected from 49%, 51%, and 40%.

[00104] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues that is equal to or greater than the ratio of branch chain amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein; and comprises a first polypeptide sequence that comprises a ratio of L residues to total amino acid residues that is equal to or greater than the ratio of L residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. In some such 5 embodiments the nutritive protein further comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. In some such 10 embodiments the nutritive protein further comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein.

[00105] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24% and a ratio of L residues to total amino acid residues that is equal to or greater than 11%. In some such embodiments the nutritive protein further comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%. 15

[00106] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 20% and a ratio of L residues to total amino acid residues that is equal to or greater than 9%. In some such embodiments the nutritive protein further comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues 20 to total amino acid residues equal to or greater than 51%.

[00107] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 18% and a ratio of L residues to total amino acid residues that is equal to or greater than 8%. In some such embodiments the nutritive protein further 30 comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues equal to or greater than 40%.

[00108] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues that is equal to or greater than the ratio of branch chain amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein; and

5 comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid

10 residues equal to or greater than 24% and a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than 49%. In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 20% and a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than 51%. In some

15 embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 18% and a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than 40%.

[00109] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of L residues to total amino acid residues that is equal to or greater than the ratio of L residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein; and comprises a first polypeptide sequence that comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present

25 in at least one of whey protein, egg protein, and soy protein. In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of L residues to total amino acid residues equal to or greater than 11% and a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than 49%. In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of L amino acid residues to total amino acid residues equal to or greater than 9% and a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than 51%. In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises a ratio of L amino acid residues to total amino acid residues equal to or greater

than 8% and a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than 40%.

[00110] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24%, a ratio of L residues to total amino acid residues that is equal to or greater than 11%, a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%, and comprises at least one of every essential amino acid, and the first polypeptide sequence is selected from **SEQ ID NO: 1 to SEQ ID NO: 1000**. In some embodiments the first polypeptide sequence is selected from a modified derivative of **SEQ ID NO: 1 to SEQ ID NO: 1000**. In some embodiments the first polypeptide sequence is selected from a mutein of **SEQ ID NO: 1 to SEQ ID NO: 1000**.

[00111] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24%, a ratio of L residues to total amino acid residues that is equal to or greater than 11%, and a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%, and the first polypeptide sequence is selected from **SEQ ID NO: 1001 to SEQ ID NO: 1414**. In some embodiments the first polypeptide sequence is selected from a modified derivative of **SEQ ID NO: 1001 to SEQ ID NO: 1414**. In some embodiments the first polypeptide sequence is selected from a mutein of **SEQ ID NO: 1001 to SEQ ID NO: 1414**.

[00112] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24%, a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%, and comprises at least one of every essential amino acid, and the first polypeptide sequence is selected from **SEQ ID NO: 1415 to SEQ ID NO: 1899**. In some embodiments the first polypeptide sequence is selected from a modified derivative of **SEQ ID NO: 1415 to SEQ ID NO: 1899**. In some embodiments the first polypeptide sequence is selected from a mutein of **SEQ ID NO: 1415 to SEQ ID NO: 1899**.

[00113] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49% and a ratio of L residues to total amino acid residues that is equal to or greater than 11%, and comprises at least one of every

5 essential amino acid, and the first polypeptide sequence is selected from **SEQ ID NO: 1900 to SEQ ID NO: 2102**. In some embodiments the first polypeptide sequence is selected from a modified derivative of **SEQ ID NO: 1900 to SEQ ID NO: 2102**. In some embodiments the first polypeptide sequence is selected from a mutein of **SEQ ID NO: 1900 to SEQ ID NO: 2102**.

10 [00114] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24% and a ratio of L residues to total amino acid residues that is equal to or greater than 11%, and comprises at least one of every essential amino acid, and the first polypeptide sequence is selected from **SEQ ID NO: 2103 to SEQ ID NO: 2518**.

15 In some embodiments the first polypeptide sequence is selected from a modified derivative of **SEQ ID NO: 2103 to SEQ ID NO: 2518**. In some embodiments the first polypeptide sequence is selected from a mutein of **SEQ ID NO: 2103 to SEQ ID NO: 2518**.

20 [00115] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24%, a ratio of L residues to total amino acid residues that is equal to or greater than 11%, and comprises at least one of every essential amino acid.

25 [00116] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24% and a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%.

30 [00117] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of L residues to total amino acid residues that is equal to or greater than 11% and a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%.

[00118] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49% and comprises at least one of every essential amino acid.

5 [00119] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24% and comprises at least one of every essential amino acid.

10 [00120] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of L residues to total amino acid residues that is equal to or greater than 11% and comprises at least one of every essential amino acid.

15 [00121] In some embodiments of the nutritive protein, the first polypeptide sequence comprises a first polypeptide sequence comprising a ratio of L residues to total amino acid residues that is equal to or greater than 11%.

20 [00122] In some embodiments of the nutritive protein, the nutritive protein further comprises a second polypeptide sequence. In some embodiments the second polypeptide sequence consists of from 3 to 10, 5 to 20, 10 to 30, 20 to 50, 25 to 75, 50 to 100 or 100 to 200 amino acids. In some embodiments the second polypeptide sequence is not derived from a naturally-occurring nutritive protein. In some embodiments the second polypeptide sequence is selected from a tag for affinity purification, a protein domain linker, and a protease recognition site. In some embodiments the tag for affinity purification is a polyhistidine-tag. In some embodiments the protein domain linker comprises at least one copy of the sequence GGSG. In some embodiments the protease is selected from pepsin, trypsin, and chymotrypsin.

25

[00123] In some embodiments the recombinant nutritive protein consists of the first polypeptide sequence and the second polypeptide sequence.

30 [00124] In some embodiments the nutritive protein further comprises a third polypeptide sequence comprising a fragment of at least 25 or at least 50 amino acids of a naturally-occurring nutritive protein. In some embodiments the first and third polypeptide

sequences are the same. In some embodiments the first and third polypeptide sequences are different. In some embodiments the first and third polypeptide sequences are derived from the same naturally-occurring nutritive protein. In some embodiments the order of the first and third polypeptide sequences in the isolated recombinant nutritive protein is the same as  
5 the order of the first and third polypeptide sequences in the naturally-occurring nutritive protein. In some embodiments the order of the first and third polypeptide sequences in the nutritive protein is different than the order of the first and third polypeptide sequences in the naturally-occurring nutritive protein. In some embodiments the first and third polypeptide sequences are derived from different naturally-occurring nutritive proteins. In some  
10 embodiments the second polypeptide sequence is flanked by the first and third polypeptide sequences.

[00125] In some embodiments of the nutritive protein, the first and/or third polypeptide sequence comprises a fragment of a naturally-occurring nutritive protein and the ratio of branch chain amino acid residues to total amino acid residues present in the first  
15 and/or third polypeptide sequence is equal to or greater than the ratio of branch chain amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein.

[00126] In some embodiments of the nutritive protein, the first and/or third polypeptide sequence comprises a first and/or third polypeptide sequence comprising a  
20 fragment of a naturally-occurring nutritive protein and the ratio of L residues to total amino acid residues present in the first and/or third polypeptide sequence is equal to or greater than the ratio of L residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein.

[00127] In some embodiments of the nutritive protein, the first and/or third  
25 polypeptide sequence comprises a first and/or third polypeptide sequence comprising a fragment of a naturally-occurring nutritive protein and the ratio of essential amino acid residues to total amino acid residues present in the first and/or third polypeptide sequence is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein.

30 [00128] In some embodiments of the nutritive protein, the first and/or third polypeptide sequence comprises a first and/or third polypeptide sequence comprising a

fragment of a naturally-occurring nutritive protein and the ratio of branch chain amino acid residues to total amino acid residues present in the first and/or third polypeptide sequence is equal to or greater than the ratio of branch chain amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein; and the first and/or third polypeptide sequence comprises a ratio of L residues to total amino acid residues that is equal to or greater than the ratio of L residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein. In some such embodiments the first and/or third polypeptide sequence further comprises a ratio of essential amino acid residues to total amino acid residues that is equal to or greater than the ratio of essential amino acid residues to total amino acid residues present in at least one of whey protein, egg protein, and soy protein.

[00129] In some embodiments of the nutritive protein, the first and/or third polypeptide sequence comprises a first and/or third polypeptide sequence comprising a ratio of branch chain amino acid residues to total amino acid residues equal to or greater than 24% and a ratio of L residues to total amino acid residues that is equal to or greater than 11%. In some such embodiments the first and/or third polypeptide sequence further comprises a ratio of essential amino acid residues to total amino acid residues equal to or greater than 49%.

[00130] In some embodiments the naturally-occurring nutritive protein from which a fragment is derived is a nutritive protein other than at least one nutritive protein selected from egg proteins such as ovalbumin, ovotransferrin, and ovomucoid; meat proteins such as myosin, actin, tropomyosin, collagen, and troponin; milk proteins such as whey and casein; cereal proteins such as casein, alpha1 casein, alpha2 casein, beta casein, kappa casein, beta-lactoglobulin, alpha-lactalbumin, glycinin, beta-conglycinin, glutelin, prolamine, gliadin, glutenin, albumin, globulin; chicken muscle proteins such as albumin, enolase, creatine kinase, phosphoglycerate mutase, triosephosphate isomerase, apolipoprotein, ovotransferrin, phosphoglucomutase, phosphoglycerate kinase, glycerol-3-phosphate dehydrogenase, glyceraldehyde 3-phosphate dehydrogenase, hemoglobin, cofilin, glycogen phosphorylase, fructose-1,6-bisphosphatase, actin, myosin, tropomyosin a-chain, casein kinase, glycogen phosphorylase, fructose-1,6-bisphosphatase, aldolase, tubulin, vimentin, endoplasmic, lactate dehydrogenase, destrin, transthyretin, fructose bisphosphate aldolase, carbonic anhydrase, aldehyde dehydrogenase, annexin, adenosyl homocysteinase; pork muscle proteins such as actin, myosin, enolase, titin, cofilin, phosphoglycerate kinase, enolase, pyruvate

dehydrogenase, glycogen phosphorylase, triosephosphate isomerase, myokinase; and fish proteins such as parvalbumin, pyruvate dehydrogenase, desmin, and triosephosphate isomerase.

[00131] Phenylketonuria (PKU) is an autosomal recessive metabolic genetic disorder characterized by a mutation in the gene for the hepatic enzyme phenylalanine hydroxylase (PAH), rendering it nonfunctional. This enzyme is necessary to metabolize phenylalanine to tyrosine. When PAH activity is reduced, phenylalanine accumulates and is converted into phenylpyruvate (also known as phenylketone), which is detected in the urine. Untreated children are normal at birth, but fail to attain early developmental milestones, develop microcephaly, and demonstrate progressive impairment of cerebral function. Hyperactivity, EEG abnormalities and seizures, and severe learning disabilities are major clinical problems later in life. A characteristic odor of skin, hair, sweat and urine (due to phenylacetate accumulation); and a tendency to hypopigmentation and eczema are also observed. All PKU patients must adhere to a special diet low in Phe. Accordingly, nutritive proteins comprising a low number or no Phe residues are desirable for PKU patients. Such proteins can be obtained by selecting nutritive proteins provided herein that have few or no Phe residues. Accordingly, in some embodiments the nutritive protein comprises a ratio of Phe residues to total amino acid residues equal to or lower than 5%, 4%, 3%, 2%, or 1%. In some embodiments the nutritive protein comprises 10 or fewer Phe residues, 9 or fewer Phe residues, 8 or fewer Phe residues, 7 or fewer Phe residues, 6 or fewer Phe residues, 5 or fewer Phe residues, 4 or fewer Phe residues, 3 or fewer Phe residues, 2 or fewer Phe residues, 1 Phe residue, or no Phe residues. In some embodiments, the nutritive protein comprises no Phe residues.

[00132] Arginine is a conditionally nonessential amino acid, meaning most of the time it can be manufactured by the human body, and does not need to be obtained directly through the diet. Individuals who have poor nutrition, the elderly, or people with certain physical conditions (e.g., sepsis) may not produce sufficient amounts of arginine and therefore need to increase their intake of foods containing arginine. Arginine is believed to have beneficial health properties, including reducing healing time of injuries (particularly bone), and decreasing blood pressure, particularly high blood pressure during high risk pregnancies (pre-eclampsia). In addition, studies have shown that dietary supplementation with L-arginine is beneficial for enhancing the reproductive performance of pigs with naturally occurring in-

trauterine growth retardation, enhancing protein deposition and postnatal growth of milk-fed piglets, normalizing plasma glucose levels in streptozotocin-induced diabetic rats, reducing fat mass in obese Zucker diabetic fatty (ZDF) rats, and improving vascular function in diabetic rats. In order to combine these benefits with at least one utility of the nutritive 5 proteins disclosed herein, in some embodiments of the nutritive proteins disclosed herein the nutritive protein comprises a ration of Arginine residues to total amino acid residues in the nutritive protein of equal to or greater than 3%, equal to or greater than 4%, equal to or greater than 5%, equal to or greater than 6%, equal to or greater than 7%, equal to or greater than 8%, equal to or greater than 9%, equal to or greater than 10%, equal to or greater than 10 11%, or equal to or greater than 12%.

[00133] Digestibility is a parameter relevant to the nutritive benefits and utility of nutritive proteins. Information relating to the relative completeness of digestion can serve as a predictor of peptide bioavailability (Daniel, H., 2003. Molecular and Integrative Physiology of Intestinal Peptide Transport. Annual Review of Physiology, Volume 66, pp. 361-384). In 15 some embodiments nutritive proteins disclosed herein are screened to assess their digestibility. Digestibility of nutritive proteins can be assessed by any suitable method known in the art. In some embodiments digestability is assessed by a physiologically relevant in vitro digestion reaction that includes one or both phases of protein digestion, simulated gastric digestion and simulated intestinal digestion (see, e.g., Moreno, et al., 2005. 20 Stability of the major allergen Brazil nut 2S albumin (Ber e 1) to physiologically relevant in vitro gastrointestinal digestion. FEBS Journal, pp. 341-352; Martos, G., Contreras, P., Molina, E. & Lopez-Fandino, R., 2010. Egg White Ovalbumin Digestion Mimicking 25 Physiological Conditions. Journal of Agricultural and food chemistry, pp. 5640-5648; Moreno, F. J., Mackie, A. R. & Clare Mills, E. N., 2005. Phospholipid interactions protect the milk allergen a-Lactalbumin from proteolysis during in vitro digestion. Journal of agricultural and food chemistry, pp. 9810-9816). Briefly, test proteins are sequentially exposed to a simulated gastric fluid (SGF) for 120 minutes (the length of time it takes 90% of a liquid meal to pass from the stomach to the small intestine; see Kong, F. & Singh, R. P., 2008. Disintegration of Solid Foods in Human Stomach. Journal of Food Science, pp. 67-80) and 30 then transferred to a simulated duodenal fluid (SDF) to digest for an additional 120 minutes. Samples at different stages of the digestion (e.g., 2, 5, 15, 30, 60 and 120 min) are analyzed by electrophoresis (e.g., chip electrophoresis or SDS-PAGE) to monitor the size and amount of intact protein as well as any large digestion fragments (e.g., larger than 4 kDa). The

disappearance of protein over time indicates the rate at which the protein is digested in the assay. By monitoring the amount of intact protein observed over time, the half-life ( $\tau_{1/2}$ ) of digestion is calculated for SGF and, if intact protein is detected after treatment with SGF, the  $\tau_{1/2}$  of digestion is calculated for SIF. This assay can be used to assess comparative

5 digestibility (i.e., against a benchmark protein such as whey) or to assess absolute digestibility. In some embodiments the digestibility of the nutritive protein is higher (i.e., the SGF  $\tau_{1/2}$  and/or SIF  $\tau_{1/2}$  is shorter) than whey protein. In some embodiments the nutritive protein has a SGF  $\tau_{1/2}$  of 30 minutes or less, 20 minutes or less, 15 minutes or less, 10 minutes or less, 5 minutes or less, 4 minutes or less, 3 minutes or less, 2 minutes or less or 1 minute or less. In some embodiments the nutritive protein has a SIF  $\tau_{1/2}$  of 30 minutes or less, 20 minutes or less, 15 minutes or less, 10 minutes or less, 5 minutes or less, 4 minutes or less, 3 minutes or less, 2 minutes or less or 1 minute or less. In some embodiments the nutritive protein is not detectable in one or both of the SGF and SIF assays by 2 minutes, 5 minutes, 15 minutes, 30 minutes, 60 minutes, or 120 minutes. In some embodiments the 10 nutritive protein is digested at a constant rate and/or at a controlled rate in one or both of SGF and SIF. In such embodiments the rate of digestion of the nutritive protein may not be optimized for the highest possible rate of digestion. In such embodiments the rate of absorption of the protein following ingestion by a mammal may be slower and the total time period over which absorption occurs following ingestion may be longer than for nutritive 15 proteins of similar amino acid composition that are digested at a faster initial rate in one or both of SGF and SIF. In some embodiments the nutritive protein is completely or substantially completely digested in SGF. In some embodiments the nutritive protein is substantially not digested or not digested by SGF; in most such embodiments the protein is digested in SIF.

20

25 [00134] Assessing protein digestibility can also provide insight into a protein's potential allergenicity, as proteins or large fragments of proteins that are resistant to digestive proteases can have a higher risk of causing an allergenic reaction (Goodman, R. E. et al., 2008. Allergenicity assessment of genetically modified crops - what makes sense? *Nature Biotechnology*, pp. 73-81). To detect and identify peptides too small for chip electrophoresis 30 analysis, liquid chromatography and mass spectrometry can be used. In SGF samples, peptides can be directly detected and identified by LC/MS. SIF protein digestions may require purification to remove bile acids before detection and identification by LC/MS.

[00135] In some embodiments digestibility of a nutritive protein is assessed by identification and quantification of digestive protease recognition sites in the protein amino acid sequence. In some embodiments the nutritive protein comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

[00136] As used herein, a “pepsin recognition site” is any site in a polypeptide sequence that is experimentally shown to be cleaved by pepsin. In some embodiments it is a peptide bond after (i.e., downstream of) an amino acid residue selected from Phe, Trp, Tyr, Leu, Ala, Glu, and Gln, provided that the following residue is not an amino acid residue selected from Ala, Gly, and Val.

[00137] As used herein, a “trypsin recognition site” is any site in a polypeptide sequence that is experimentally shown to be cleaved by trypsin. In some embodiments it is a peptide bond after an amino acid residue selected from Lys or Arg, provided that the following residue is not a proline.

[00138] As used herein, a “chymotrypsin recognition site” is any site in a polypeptide sequence that is experimentally shown to be cleaved by chymotrypsin. In some embodiments it is a peptide bond after an amino acid residue selected from Phe, Trp, Tyr, and Leu.

[00139] Disulfide bonded cysteine residues in a protein tend to reduce the rate of digestion of the protein compared to what it would be in the absence of the disulfide bond. For example, it has been shown that the rate of digestion of the protein b-lactoglobulin is increased when its disulfide bridges are cleaved (I. M. Reddy, N. K. D. Kella, and J. E. Kinsella. “Structural and Conformational Basis of the Resistance of B-Lactoglobulin to Peptic and Chymotryptic Digestion”. *J. Agric. Food Chem.* 1988, 36, 737-741). Accordingly, digestibility of a nutritive protein with fewer disulfide bonds tends to be higher than for a comparable nutritive protein with a greater number of disulfide bonds. In some embodiments the nutritive proteins disclosed herein are screened to identify the number of cysteine residues present in each and in particular to allow selection of a nutritive protein comprising a relatively low number of cysteine residues. For example, naturally occurring nutritive proteins or fragments may be identified that comprise a no Cys residues or that comprise a relatively low number of Cys residues, such as 10 or fewer Cys residues, 9 or fewer Cys residues, 8 or fewer Cys residues, 7 or fewer Cys residues, 6 or fewer Cys

residues, 5 or fewer Cys residues, 4 or fewer Cys residues, 3 or fewer Cys residues, 2 or fewer Cys residues, 1 Cys residue, or no Cys residues. In some embodiments one or more Cys residues in a naturally occurring nutritive protein or fragment thereof is removed by deletion and/or by substitution with another amino acid. In some embodiments 1 Cys residue  
5 is deleted or replaced, 1 or more Cys residues are deleted or replaced, 2 or more Cys residues are deleted or replaced, 3 or more Cys residues are deleted or replaced, 4 or more Cys residues are deleted or replaced, 5 or more Cys residues are deleted or replaced, 6 or more Cys residues are deleted or replaced, 7 or more Cys residues are deleted or replaced, 8 or more Cys residues are deleted or replaced, 9 or more Cys residues are deleted or replaced, or  
10 10 or more Cys residues are deleted or replaced. In some embodiments the nutritive protein of this disclosure comprises a ratio of Cys residues to total amino acid residues equal to or lower than 5%, 4%, 3%, 2%, or 1%. In some embodiments the nutritive protein comprises 10 or fewer Cys residues, 9 or fewer Cys residues, 8 or fewer Cys residues, 7 or fewer Cys residues, 6 or fewer Cys residues, 5 or fewer Cys residues, 4 or fewer Cys residues, 3 or  
15 fewer Cys residues, 2 or fewer Cys residues, 1 Cys residue, or no Cys residues. In some embodiments, the nutritive protein comprises 1 or fewer Cys residues. In some embodiments, the nutritive protein comprises no Cys residues.

[00140] Alternatively or in addition, disulfide bonds that are or may be present in a nutritive protein may be removed. Disulfides can be removed using chemical methods by  
20 reducing the disulfide to two thiol groups with reducing agents such as beta-mercaptoethanol, dithiothreitol (DTT), or tris(2-carboxyethyl)phosphine (TCEP). The thiols can then be covalently modified or “capped” with reagents such as iodoacetamide, N-ethylmaleimide, or sodium sulfite (see, e.g., Crankshaw, M. W. and Grant, G. A. 2001. Modification of Cysteine. Current Protocols in Protein Science. 15.1.1–15.1.18).

25 [00141] Eukaryotic proteins are often glycosylated, and the carbohydrate chains that are attached to proteins serve various functions. N-linked and O-linked glycosylation are the two most common forms of glycosylation occurring in proteins. N-linked glycosylation is the attachment of a sugar molecule to a nitrogen atom in an amino acid residue in a protein. N-linked glycosylation occurs at Asparagine and Arginine residues. O-linked glycosylation is  
30 the attachment of a sugar molecule to an oxygen atom in an amino acid residue in a protein. O-linked glycosylation occurs at Threonine and Serine residues.

[00142] Glycosylated proteins are often more soluble than their un-glycosylated forms.

In terms of protein drugs, proper glycosylation usually confers high activity, proper antigen binding, better stability in the blood, etc. However, glycosylation necessarily means that a protein “carries with it” sugar moieties. Such sugar moieties may reduce the usefulness of

5 the nutritive proteins of this disclosure including recombinant nutritive proteins. For example, as demonstrated in the examples, a comparison of digestion of glycosylated and non-glycosylated forms of the same proteins shows that the non-glycosylated forms are digested more quickly than the glycosylated forms. For these reasons, in some embodiments the nutritive proteins according to the disclosure comprise low or no glycosylation. For  
10 example, in some embodiments the nutritive proteins comprise a ratio of non-glycosylated to total amino acid residues of at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%. In some embodiments the nutritive proteins to not comprise any glycosylation.

[00143] In some embodiments, the nutritive protein according to the disclosure is de-

15 glycosylated after it is produced or after it is isolated. Nutritive proteins of low or no glycosylation may be made by any method known in the art. For example, enzymatic and/or chemical methods may be used (Biochem. J. (2003) 376, p339-350.). Enzymes are produced commercially at research scales for the removal of N-linked and O-linked oligosaccharides. Chemical methods include use of trifluoromethanesulfonic acid to selectively break N-linked  
20 and O-linked peptide-saccharide bonds. This method often results in a more complete deglycosylation than does the use of enzymatic methods.

[00144] In other embodiments, the nutritive protein according to the disclosure is produced with low or no glycosylation by a host organism. Most bacteria and other prokaryotes have very limited capabilities to glycosylate proteins, especially heterologous

25 proteins. Accordingly, in some embodiments of this disclosure a nutritive protein is made recombinantly in a microorganism such that the level of glycosylation of the recombinant protein is low or no glycosylation. In some embodiments the level of glycosylation of the recombinant nutritive protein is lower than the level of glycosylation of the protein as it occurs in the organism from which it is derived.

30 [00145] In some embodiments a nutritive protein or polypeptide according to the disclosure comprises a ratio of amino acids selected from Asn, Arg, Ser, and Thr to total amino acids of 20% or less, 19% or less, 18 % or less, 17% or less, 16% or less, 15% or less,

14% or less, 13% or less, 12% or less, 11% or less, 10% or less, 9% or less, 8% or less, 7% or less, 6% or less, 5% or less, 4% or less, 3% or less, 2% or less, 1% or less. In some embodiments, a nutritive protein or polypeptide according to the disclosure comprises no amino acids selected from Asn, Arg, Ser, and Thr. In some embodiments a nutritive protein

5 or polypeptide according to the disclosure comprises fewer than 20, fewer than 19, fewer than 18, fewer than 17, fewer than 16, fewer than 15, fewer than 14, fewer than 13, fewer than 12, fewer than 11, fewer than 10, fewer than 9, fewer than 8, fewer than 7, fewer than 6, fewer than 5, fewer than 4, fewer than 3, fewer than 2, fewer than 1, or no amino acids selected from Asn, Arg, Ser, and Thr.

10 [00146] In some embodiments of the nutritive proteins disclosed herein the nutritive protein is soluble. Solubility can be measured by any method known in the art. In some embodiments solubility is examined by centrifuge concentration followed by protein concentration assays. Samples of nutritive proteins in 20 mM HEPES pH 7.5 are tested for protein concentration according to protocols using two methods, Coomassie Plus (Bradford) 15 Protein Assay (Thermo Scientific) and Bicinchoninic Acid (BCA) Protein Assay (Sigma-Aldrich). Based on these measurements 10 mg of protein is added to an Amicon Ultra 3 kDa centrifugal filter (Millipore). Samples are concentrated by centrifugation at 10,000 Xg for 30 minutes. The final, now concentrated, samples are examined for precipitated protein and then tested for protein concentration as above using two methods, Bradford and BCA.

20 [00147] In some embodiments the nutritive proteins have a final solubility limit of at least 5 g/L, 10 g/L, 20 g/L, 30 g/L, 40 g/L, 50 g/L, or 100 g/L at physiological pH. In some embodiments the nutritive proteins are greater than 50%, greater than 60%, greater than 70%, greater than 80%, greater than 90%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, greater than 99%, or greater than 99.5% soluble with no precipitated 25 protein observed at a concentration of greater than 5 g/L, or 10 g/L, or 20 g/L, or 30 g/L, or 40 g/L, or 50 g/L, or 100 g/L at physiological pH. In some embodiments, the solubility of the nutritive protein is higher than those typically reported in studies examining the solubility limits of whey (12.5 g/L; Pelegrine et al., Lebensm.-Wiss. U.-Technol. 38 (2005) 77-80) and soy (10 g/L; Lee et al., JAOCs 80(1) (2003) 85-90).

30 [00148] As used herein, a “stable” protein is one that resists changes (e.g., unfolding, oxidation, aggregation, hydrolysis, etc.) that alter the biophysical (e.g., solubility), biological

(e.g., digestibility), or compositional (e.g. proportion of Leucine amino acids) traits of the protein of interest.

[00149] Protein stability can be measured using various assays known in the art and nutritive proteins disclosed herein and having stability above a threshold can be selected. In 5 some embodiments a protein is selected that displays thermal stability that is comparable to or better than that of whey protein. Thermal stability is a property that can help predict the shelf life of a nutritive protein. In some embodiments of the assay stability of nutritive protein samples is determined by monitoring aggregation formation using size exclusion chromatography (SEC) after exposure to extreme temperatures. Aqueous samples of the 10 protein to be tested are placed in a heating block at 90°C and samples are taken after 0, 1, 5, 10, 30 and 60 min for SEC analysis. Protein is detected by monitoring absorbance at 214nm, and aggregates are characterized as peaks eluting faster than the protein of interest. No overall change in peak area indicates no precipitation of protein during the heat treatment. Whey protein has been shown to rapidly form ~ 80% aggregates when exposed to 90°C in 15 such an assay.

[00150] In some embodiments the thermal stability of a nutritive protein is determined by heating a sample slowly from 25°C to 95°C in presence of a hydrophobic dye (e.g., ProteoStat® Thermal shift stability assay kit, Enzo Life Sciences) that binds to aggregated proteins that are formed as the protein denatures with increasing temperature (Niesen, F. H., 20 Berglund, H. & Vadadi, M., 2007. The use of differential scanning fluorimetry to detect ligand interactions that promote protein stability. *Nature Protocols*, Volume 2, pp. 2212-2221). Upon binding, the dye's fluorescence increases significantly, which is recorded by an rtPCR instrument and represented as the protein's melting curve (Lavinder, J. J., Hari, S. B., 25 Suillivan, B. J. & Magilery, T. J., 2009. High-Throughput Thermal Scanning: A General, Rapid Dye-Binding Thermal Shift Screen for Protein Engineering. *Journal of the American Chemical Society*, pp. 3794-3795). After the thermal shift is complete, samples are examined for insoluble precipitates and further analyzed by analytical size exclusion chromatography (SEC).

[00151] In some embodiments a nutritive protein of this disclosure shows resistance to 30 aggregation, exhibiting, for example, less than 80% aggregation, 10% aggregation, or no detectable aggregation at elevated temperatures (e.g., 50°C, 60°C, 70°C, 80°C, 85°C, 90°C, or 95°C).

[00152] One benefit of stable nutritive proteins as disclosed herein is that they may be able to be stored for an extended period of time before use, in some instances without the need for refrigeration or cooling. In some embodiments, nutritive proteins are processed into a dry form (e.g., by lyophilization). In some embodiments, nutritive proteins are stable upon 5 lyophilization. In some embodiments, such lyophilized nutritive proteins maintain their stability upon reconstitution (e.g., liquid formulation).

[00153] For most embodiments it is preferred that the nutritive protein not exhibit inappropriately high allergenicity. Accordingly, in some embodiments the potential allergenicity of the nutritive protein is assessed. This can be done by any suitable method 10 known in the art. In some embodiments an allergenicity score is calculated. The allergenicity score is a primary sequence based metric based on WHO recommendations (<http://www.fao.org/ag/agn/food/pdf/allergygm.pdf>) for assessing how similar a protein is to any known allergen, the primary hypothesis being that high percent identity between a target and a known allergen is likely indicative of cross reactivity. For a given protein, the 15 likelihood of eliciting an allergic response can be assessed via one or both of a complimentary pair of sequence homology based tests. The first test determines the protein's percent identity across the entire sequence via a global-global sequence alignment to a database of known allergens using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. It has been suggested that 20 proteins with less than 50% global homology are unlikely to be allergenic (Goodman R. E. et al. Allergenicity assessment of genetically modified crops—what makes sense? *Nat. Biotech.* 26, 73-81 (2008); Aalberse R. C. Structural biology of allergens. *J. Allergy Clin. Immunol.* 106, 228-238 (2000)).

[00154] In some embodiments of a nutritive protein, the nutritive protein has less than 25 50% global homology to any known allergen in the database used for the analysis. In some embodiments a cutoff of less than 40% homology is used. In some embodiments a cutoff of less than 30% homology is used. In some embodiments a cutoff of less than 20% homology is used. In some embodiments a cutoff of less than 10% homology is used. In some 30 embodiments a cutoff of from 40% to 50% is used. In some embodiments a cutoff of from 30% to 50% is used. In some embodiments a cutoff of from 20% to 50% is used. In some embodiments a cutoff of from 10% to 50% is used. In some embodiments a cutoff of from 5% to 50% is used. In some embodiments a cutoff of from 0% to 50% is used. In some

embodiments a cutoff of greater than 50% global homology to any known allergen in the database used for the analysis is used. In some embodiments a cutoff of from 50% to 60% is used. In some embodiments a cutoff of from 50% to 70% is used. In some embodiments a cutoff of from 50% to 80% is used. In some embodiments a cutoff of from 50% to 90% is used. In some embodiments a cutoff of from 55% to 60% is used. In some embodiments a cutoff of from 65% to 70% is used. In some embodiments a cutoff of from 70% to 75% is used. In some embodiments a cutoff of from 75% to 80% is used.

[00155] The second test assesses the local allergenicity along the protein sequence by determining the local allergenicity of all possible contiguous 80 amino acid fragments via a global-local sequence alignment of each fragment to a database of known allergens using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. The highest percent identity of any 80 amino acid window with any allergen is taken as the final score for the protein of interest. The WHO guidelines suggest using a 35% identity cutoff with this fragment test. In some embodiments of a nutritive protein, all possible fragments of the nutritive protein have less than 35% local homology to any known allergen in the database used for the analysis using this test. In some embodiments a cutoff of less than 30% homology is used. In some embodiments a cutoff of from 30% to 35% homology is used. In some embodiments a cutoff of from 25% to 30% homology is used. In some embodiments a cutoff of from 20% to 25% homology is used. In some embodiments a cutoff of from 15% to 20% homology is used. In some embodiments a cutoff of from 10% to 15% homology is used. In some embodiments a cutoff of from 5% to 10% homology is used. In some embodiments a cutoff of from 0% to 5% homology is used. In some embodiments a cutoff of greater than 35% homology is used. In some embodiments a cutoff of from 35% to 40% homology is used. In some embodiments a cutoff of from 40% to 45% homology is used. In some embodiments a cutoff of from 45% to 50% homology is used. In some embodiments a cutoff of from 50% to 55% homology is used. In some embodiments a cutoff of from 55% to 60% homology is used. In some embodiments a cutoff of from 65% to 70% homology is used. In some embodiments a cutoff of from 70% to 75% homology is used. In some embodiments a cutoff of from 75% to 80% homology is used.

[00156] Skilled artisans are able to identify and use a suitable database of known allergens for this purpose. In some embodiments the database is custom made by selecting proteins from more than one database source. In some embodiments the custom database

comprises pooled allergen lists collected by the Food Allergy Research and Resource Program (<http://www.allergenonline.org/>), UNIPROT annotations (<http://www.uniprot.org/docs/allergen>), and the Structural Database of Allergenic Proteins (SDAP, [http://fermi.utmb.edu/SDAP/sdap\\_lnk.html](http://fermi.utmb.edu/SDAP/sdap_lnk.html)). This database includes all currently

5 recognized allergens by the International Union of Immunological Societies (IUIS, <http://www.allergen.org/>) as well as a large number of additional allergens not yet officially named. In some embodiments the database comprises a subset of known allergen proteins available in known databases; that is, the database is a custom selected subset of known allergen proteins. In some embodiments the database of known allergens comprises at least

10 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100, proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 1,100 proteins, at least 1,200 proteins, at least 1,300 proteins, at least 1,400 proteins, at least 1,500 proteins, at least 1,600 proteins, at least 1,700

15 proteins, at least 1,800 proteins, at least 1,900 proteins, or at least 2,000 proteins. In some embodiments the database of known allergens comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, or from 1,000 to 2,000 proteins.

[00157] In some embodiments all (or a selected subset) of contiguous amino acid windows of different lengths (e.g., 70, 60, 50, 40, 30, 20, 10, 8 or 6 amino acid windows) of a nutritive protein are tested against the allergen database and peptide sequences that have 100% identity, 95% or higher identity, 90% or higher identity, 85% or higher identity, 80% or higher identity, 75% or higher identity, 70% or higher identity, 65% or higher identity, 60% or higher identity, 55% or higher identity, or 50% or higher identity matches are

25 identified for further examination of potential allergenicity.

[00158] Another method of predicting the allergenicity of a protein is to assess the homology of the protein to a protein of human origin. The human immune system is exposed to a multitude of possible allergenic proteins on a regular basis and has the intrinsic ability to differentiate between the host body's proteins and exogenous proteins. The exact nature of

30 this ability is not always clear, and there are many diseases that arise as a result of the failure of the body to differentiate self from non-self (e.g. arthritis). Nonetheless, the fundamental hypothesis is that proteins that share a degree of sequence homology to human proteins are

less likely to elicit an immune response. In particular, it has been shown that for some protein families with known allergenic members (tropomyosins, parvalbumins, caseins), those proteins that bear more sequence homology to their human counterparts relative to known allergenic proteins, are not thought to be allergenic (Jenkins J. A. et al. Evolutionary distance from human homologs reflects allergenicity of animal food proteins. *J. Allergy Clin Immunol.* 120 (2007): 1399-1405). For a given protein, a human homology score is measured by determining the maximum percent identity of the protein to a database of human proteins (e.g., the UNIPROT database) from a global-local alignment using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. According to Jenkins et al. (Jenkins J. A. et al. Evolutionary distance from human homologs reflects allergenicity of animal food proteins. *J. Allergy Clin Immunol.* 120 (2007): 1399-1405) proteins with a sequence identity to a human protein above about 62% are less likely to be allergenic. Skilled artisans are able to identify and use a suitable database of known human proteins for this purpose, for example, by searching the UNIPROT database (<http://www.uniprot.org>). In some embodiments the database is custom made by selecting proteins from more than one database source. Of course the database may but need not be comprehensive. In some embodiments the database comprises a subset of human proteins; that is, the database is a custom selected subset of human proteins. In some embodiments the database of human proteins comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100 proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 2,000 proteins, at least 3,000 proteins, at least 4,000 proteins, at least 5,000 proteins, at least 6,000 proteins, at least 7,000 proteins, at least 8,000 proteins, at least 9,000 proteins, or at least 10,000 proteins. In some embodiments the database comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, from 1,000 to 2,000 proteins, from 1,000 to 5,000 proteins, or from 5,000 to 10,000 proteins. In some embodiments the database comprises at least 90%, at least 95%, or at least 99% of all known human proteins.

30 [00159] In some embodiments of a nutritive protein, the nutritive protein is at least 20% homologous to a human protein. In some embodiments a cutoff of at least 30% homology is used. In some embodiments a cutoff of at least 40% homology is used. In some embodiments a cutoff of at least 50% homology is used. In some embodiments a cutoff of at

least 60% homology is used. In some embodiments a cutoff of at least 70% homology is used. In some embodiments a cutoff of at least 80% homology is used. In some embodiments a cutoff of at least 62% homology is used. In some embodiments a cutoff of from at least 20% homology to at least 30% homology is used. In some embodiments a cutoff of from at least 30% homology to at least 40% homology is used. In some embodiments a cutoff of from at least 50% homology to at least 60% homology is used. In some embodiments a cutoff of from at least 60% homology to at least 70% homology is used. In some embodiments a cutoff of from at least 70% homology to at least 80% homology is used.

- 5 [00160] For most embodiments it is preferred that the nutritive protein not exhibit inappropriately high toxicity. Accordingly, in some embodiments the potential toxicity of the nutritive protein is assessed. This can be done by any suitable method known in the art. In some embodiments a toxicity score is calculated by determining the protein's percent identity to databases of known toxic proteins (e.g., toxic proteins identified from the UNIPROT 10 database). A global-global alignment of the protein of interest against the database of known 15 toxins is performed using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. In some embodiments of a nutritive protein, the nutritive protein is less than 35% homologous to a known toxin. In some embodiments a cutoff of less than 35% homology is used. In some embodiments a cutoff of from 30% to 35% homology is used. In some embodiments a cutoff of from 25% to 20 35% homology is used. In some embodiments a cutoff of from 20% to 35% homology is used. In some embodiments a cutoff of from 15% to 35% homology is used. In some embodiments a cutoff of from 10% to 35% homology is used. In some embodiments a cutoff of from 5% to 35% homology is used. In some embodiments a cutoff of from 0% to 35% 25 homology is used. In some embodiments a cutoff of greater than 35% homology is used. In some embodiments a cutoff of from 35% to 40% homology is used. In some embodiments a cutoff of from 35% to 50% homology is used. In some embodiments a cutoff of from 35% to 55% homology is used. In some embodiments a cutoff of from 35% to 60% homology is used. In some 30 embodiments a cutoff of from 35% to 70% homology is used. In some embodiments a cutoff of from 35% to 75% homology is used. In some embodiments a cutoff of from 35% to 80% homology is used. Skilled artisans are able to identify and use a suitable database of known toxins for this purpose, for example, by searching the UNIPROT database

(<http://www.uniprot.org>). In some embodiments the database is custom made by selecting proteins identified as toxins from more than one database source. In some embodiments the database comprises a subset of known toxic proteins; that is, the database is a custom selected subset of known toxic proteins. In some embodiments the database of toxic proteins

- 5 comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100, proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 2,000 proteins, at least 3,000 proteins, at least 4,000 proteins, at least 5,000 proteins, at least 6,000 proteins, at least 7,000 proteins, at least 8,000 proteins, at least 9,000 proteins, or at least 10,000 proteins. In some embodiments the database comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, from 1,000 to 2,000 proteins, from 1,000 to 5,000 proteins, or from 5,000 to 10,000 proteins.
- 10

- [00161] For some embodiments it is preferred that the nutritive protein not exhibit 15 anti-nutritional activity (“anti-nutricity”), i.e., proteins that have the potential to prevent the absorption of nutrients from food. Examples of anti-nutritive factors include protease inhibitors, which inhibit the actions of trypsin, pepsin and other proteases in the gut, preventing the digestion and subsequent absorption of protein. Accordingly, in some embodiments the potential anti-nutricity of the nutritive protein is assessed. This can be done 20 by any suitable method known in the art. In some embodiments an anti-nutricity score is calculated by determining the protein’s percent identity to databases of known protease inhibitors (e.g., protease inhibitors identified from the UNIPROT database). A global-global alignment of the protein of interest against the database of known protease inhibitors is performed using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open 25 penalty of 10, and a gap extension penalty of 2, to identify whether the nutritive protein is homologous to a known anti-nutritive protein. In some embodiments of a nutritive protein, the nutritive protein has less than 35% global homology to any known anti-nutritive protein (e.g., any known protease inhibitor) in the database used for the analysis. In some embodiments a cutoff of less than 35% identify is used. In some embodiments a cutoff of 30 30 from 30% to 35% is used. In some embodiments a cutoff of from 25% to 35% is used. In some embodiments a cutoff of from 20% to 35% is used. In some embodiments a cutoff of from 15% to 35% is used. In some embodiments a cutoff of from 10% to 35% is used. In some embodiments a cutoff of from 5% to 35% is used. In some embodiments a cutoff of

from 0% to 35% is used. In some embodiments a cutoff of greater than 35% identify is used. In some embodiments a cutoff of from 35% to 40% is used. In some embodiments a cutoff of from 35% to 45% is used. In some embodiments a cutoff of from 35% to 50% is used. In some embodiments a cutoff of from 35% to 55% is used. In some embodiments a cutoff of from 35% to 60% is used. In some embodiments a cutoff of from 35% to 70% is used. In some embodiments a cutoff of from 35% to 75% is used. In some embodiments a cutoff of from 35% to 80% is used. Skilled artisans are able to identify and use a suitable database of known protease inhibitors for this purpose, for example, by searching the UNIPROT database (<http://www.uniprot.org>). In some embodiments the database is custom made by selecting 5 proteins identified protease-inhibitors as from more than one database source. In some embodiments the database comprises a subset of known protease inhibitors available in databases; that is, the database is a custom selected subset of known protease inhibitor proteins. In some embodiments the database of known protease inhibitor proteins comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 10 proteins, at least 100, proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 1,100 proteins, at least 1,200 proteins, at least 1,300 proteins, at least 1,400 proteins, at least 1,500 proteins, at least 1,600 proteins, at least 1,700 proteins, at least 1,800 proteins, at least 1,900 proteins, or at least 2,000 proteins. 15 20 In some embodiments the database of known protease inhibitor proteins comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, or from 1,000 to 2,000 proteins, or from 2,000 to 3,000 proteins.

[00162] In other embodiments a nutritive protein that does exhibit some degree of protease inhibitor activity is used. For example, in some embodiments such a protein may be 25 useful because it delays protease digestion when the nutritive protein is consumed such that the nutritive protein traverses a greater distance within the GI tract before it is digested, thus delaying absorption. For example, in some embodiments the nutritive protein inhibits gastric digestion but not intestinal digestion.

[00163] Delaney B. et al. (Evaluation of protein safety in the context of agricultural 30 biotechnology. *Food. Chem. Toxicol.* 46 (2008: S71-S97)) suggests that one should avoid both known toxic and anti-nutritive proteins when assessing the safety of a possible food protein. In some embodiments of a nutritive protein, the nutritive protein has a favorably low

level of global homology to a database of known toxic proteins and/or a favorably low level of global homology to a database of known anti-nutritive proteins (e.g., protease inhibitors), as defined herein.

[00164] One feature that can enhance the utility of a nutritive protein is its charge (or 5 per amino acid charge). Nutritive proteins with higher charge can in some embodiments exhibit desirable characteristics such as increased solubility, increased stability, resistance to aggregation, and desirable taste profiles. For example, a charged nutritive protein that exhibits enhanced solubility can be formulated into a beverage or liquid formulation that includes a high concentration of nutritive protein in a relatively low volume of solution, thus 10 delivering a large dose of protein nutrition per unit volume. A charged nutritive protein that exhibits enhanced solubility can be useful, for example, in sports drinks or recovery drinks wherein a user (e.g., an athlete) wants to ingest nutritive protein before, during or after physical activity. A charged nutritive protein that exhibits enhanced solubility can also be particularly useful in a clinical setting wherein a subject (e.g., a patient or an elderly person) 15 is in need of protein nutrition but is unable to ingest solid foods or large volumes of liquids.

[00165] For example, the net charge (ChargeP) of a polypeptide at pH 7 can be calculated using the following formula:

[00166] 
$$\text{ChargeP} = -0.002 - (C)(0.045) - (D)(0.999) - (E)(0.998) + (H)(0.091) + (K)(1.0) + (R)(1.0) - (Y)(-0.001)$$

[00167] where C is the number of cysteine residues, D is the number of aspartic acid residues, E is the number of glutamic acid residues, H is the number of histidine residues, K is the number of lysine residues, R is the number of arginine residues and Y is the number of tyrosine residues in the polypeptide. The per amino acid charge (ChargeA) of the 20 polypeptide can be calculated by dividing the net charge (ChargeP) by the number of amino acid residues (N), i.e.,  $\text{ChargeA} = \text{ChargeP}/N$ . (See Bassi S (2007), "A Primer on Python for 25 Life Science Researchers." PLoS Comput Biol 3(11): e199. doi:10.1371/journal.pcbi.0030199).

[00168] One metric for assessing the hydrophilicity and potential solubility of a given 30 protein is the solvation score. Solvation score is defined as the total free energy of solvation (i.e. the free energy change associated with transfer from gas phase to a dilute solution) for all amino acid side chains if each residue were solvated independently, normalized by the total

number of residues in the sequence. The side chain solvation free energies are found computationally by calculating the electrostatic energy difference between a vacuum dielectric of 1 and a water dielectric of 80 (by solving the Poisson-Boltzmann equation) as well as the non-polar, Van der Waals energy using a linear solvent accessible surface area model (D. Sitkoff, K. A. Sharp, B. Honig. "Accurate Calculation of Hydration Free Energies Using Macroscopic Solvent Models". *J. Phys. Chem.* 98, 1994). For amino acids with ionizable sidechains (Arg, Asp, Cys, Glu, His, Lys and Tyr), an average solvation free energy is used based on the relative probabilities for each ionization state at the specified pH. Solvation scores start at 0 and continue into negative values, and the more negative the solvation score, the more hydrophilic and potentially soluble the protein is predicted to be. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -30 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at pH 7.

[00169] The solvation score is a function of pH by virtue of the pH dependence of the molar ratio of undissociated weak acid ([HA]) to conjugate base ([A-]) as defined by the Henderson-Hasselbalch equation:

$$[00170] \quad pH = pK_a + \log \left( \frac{[A^-]}{[H_A]} \right)$$

[00171] All weak acids have different solvation free energies compared to their conjugate bases, and the solvation free energy used for a given residue when calculating the solvation score at a given pH is the weighted average of those two values.

[00172] Accordingly, in some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a

solvation score of -25 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -30 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a

5 solvation score of -40 or less at acidic pH.

[00173] Accordingly, in some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at

10 a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -30 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at a basic pH. In some

15 embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at basic pH.

[00174] Accordingly, in some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8,

20 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein

25 has a solvation score of -30 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8,

30 8-9, 9-10, 10-11, and 11-12.

[00175] The aggregation score is a primary sequence based metric for assessing the hydrophobicity and likelihood of aggregation of a given protein. Using the Kyte and

Doolittle hydrophobicity scale (Kyte J, Doolittle RF (May 1982) "A simple method for displaying the hydropathic character of a protein". *J. Mol. Biol.* 157 (1): 105-32), which gives hydrophobic residues positive values and hydrophilic residues negative values, the average hydrophobicity of a protein sequence is calculated using a moving average of five residues.

- 5 The aggregation score is drawn from the resulting plot by determining the area under the curve for values greater than zero and normalizing by the total length of the protein. The underlying hypothesis is that aggregation is the result of two or more hydrophobic patches coming together to exclude water and reduce surface exposure, and the likelihood that a protein will aggregate is a function of how densely packed its hydrophobic (i.e., aggregation prone) residues are. Aggregation scores start at 0 and continue into positive values, and the smaller the aggregation score, the less hydrophobic and potentially less prone to aggregation the protein is predicted to be. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 2 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 1.5 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 1 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.9 or less. In some 10 embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.8 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.7 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.6 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.5 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.4 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.3 or less. In some 15 embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.2 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.1 or less.
- 20 [00176] In some cases, soluble expression is desirable because it can increase the amount and/or yield of the nutritive protein and facilitate one or more of the isolation and purification of the nutritive protein. In some embodiments, the nutritive proteins of this disclosure are solubly expressed in the host organism. Solvation score and aggregation score can be used to predict soluble expression of recombinant nutritive proteins in a host 25 organism. As shown in Example 8, this disclosure provides evidence suggesting that nutritive proteins with solvation scores of  $\leq -20$  and aggregation scores of  $\leq 0.75$  are more

likely to be recombinantly expressed in a particular *E. coli* expression system. Moreover, the data also suggests that nutritive proteins with solvation scores of  $\leq -20$  and aggregation scores of  $\leq 0.5$  are more likely to be solubly expressed in this system. Therefore, in some embodiments the nutritive protein of this disclosure has a solvation score of -20 or less. In 5 some embodiments the nutritive protein has an aggregation score of 0.75 or less. In some embodiments the nutritive protein has an aggregation score of 0.5 or less. In some embodiments the nutritive protein has a solvation score of -20 or less and an aggregation score of 0.75 or less. In some embodiments the nutritive protein has a solvation score of -20 or less and an aggregation score of 0.5 or less.

10 [00177] Certain free amino acids and mixtures of free amino acids are known to have a bitter or otherwise unpleasant taste. In addition, hydrolysates of common proteins (e.g., whey and soy) often have a bitter or unpleasant taste. In some embodiments, nutritive proteins disclosed and described herein do not have a bitter or otherwise unpleasant taste. In some embodiments, nutritive proteins disclosed and described herein have a more acceptable 15 taste as compared to at least one of free amino acids, mixtures of free amino acids, and/or protein hydrolysates. In some embodiments, nutritive proteins disclosed and described herein have a taste that is equal to or exceeds at least one of whey protein.

[00178] Proteins are known to have tastes covering the five established taste modalities: sweet, sour, bitter, salty and umami. The taste of a particular protein (or its lack 20 thereof) can be attributed to several factors, including the primary structure, the presence of charged side chains, and the electronic and conformational features of the protein. In some embodiments, nutritive proteins disclosed and described herein are designed to have a desired taste (e.g., sweet, salty, umami) and/or not to have an undesired taste (e.g., bitter, sour). In this context “design” includes, for example, selecting naturally occurring proteins embodying 25 features that achieve the desired taste property, as well as creating muteins of naturally-occurring proteins that have desired taste properties. For example, nutritive proteins can be designed to interact with specific taste receptors, such as sweet receptors (T1R2-T1R3 heterodimer) or umami receptors (T1R1-T1R3 heterodimer, mGluR4, and/or mGluR1). Further, nutritive proteins may be designed not to interact, or to have diminished interaction, 30 with other taste receptors, such as bitter receptors (T2R receptors).

[00179] Nutritive proteins disclosed and described herein can also elicit different physical sensations in the mouth when ingested, sometimes referred to as “mouth feel”. The

mouth feel of the nutritive proteins may be due to one or more factors including primary structure, the presence of charged side chains, and the electronic and conformational features of the protein. In some embodiments, nutritive proteins elicit a buttery or fat-like mouth feel when ingested.

amino acids, at least 150 amino acids, at least 155 amino acids, at least 160 amino acids, at least 165 amino acids, at least 170 amino acids, at least 175 amino acids, at least 180 amino acids, at least 185 amino acids, at least 190 amino acids, at least 195 amino acids, at least 200 amino acids, at least 205 amino acids, at least 210 amino acids, at least 215 amino acids, at 5 least 220 amino acids, at least 225 amino acids, at least 230 amino acids, at least 235 amino acids, at least 240 amino acids, at least 245 amino acids, or at least 250 amino acids.

## B. Nucleic Acids

[00181] Also provided herein are nucleic acids encoding nutritive polypeptides or proteins. In some embodiments the nucleic acid is isolated. In some embodiments the 10 nucleic acid is purified.

[00182] In some embodiments of the nucleic acid, the nucleic acid comprises a nucleic acid sequence that encodes a first polypeptide sequence disclosed in Section A above. In some embodiments of the nucleic acid, the nucleic acid consists of a nucleic acid sequence that encodes a first polypeptide sequence disclosed in Section A above. In some 15 embodiments of the nucleic acid, the nucleic acid comprises a nucleic acid sequence that encodes a nutritive protein disclosed in Section A above. In some embodiments of the nucleic acid, the nucleic acid consists of a nucleic acid sequence that encodes a nutritive protein disclosed in Section A above. In some embodiments of the nucleic acid the nucleic acid sequence that encodes the first polypeptide sequence is operatively linked to at least one 20 expression control sequence. For example, in some embodiments of the nucleic acid the nucleic acid sequence that encodes the first polypeptide sequence is operatively linked to a promoter such as a promoter described in Section D below.

[00183] Accordingly, in some embodiments the nucleic acid molecule of this disclosure encodes a polypeptide or protein that itself is a nutritive polypeptide or protein. 25 Such a nucleic acid molecule may be referred to as a “nutritive nucleic acid”. In some embodiments the nutritive nucleic acid encodes a polypeptide or protein that itself comprises at least one of: a) a ratio of branch chain amino acid residues to total amino acid residues of at least 24%; b) a ratio of Leu residues to total amino acid residues of at least 11%, and c) a ratio of essential amino acid residues to total amino acid residues of at least 49%. In some 30 embodiments the nutritive nucleic acid comprises at least 10 nucleotides, at least 20 nucleotides, at least 30 nucleotides, at least 40 nucleotides, at least 50 nucleotides, at least 60 nucleotides, at least 70 nucleotides, at least 80 nucleotides, at least 90 nucleotides, at least

100 nucleotides, at least 200 nucleotides, at least 300 nucleotides, at least 400 nucleotides, at least 500 nucleotides, at least 600 nucleotides, at least 700 nucleotides, at least 800 nucleotides, at least 900 nucleotides, at least 1,000 nucleotides. In some embodiments the nutritive nucleic acid comprises from 10 to 100 nucleotides, from 20 to 100 nucleotides, 5 from 10 to 50 nucleotides, or from 20 to 40 nucleotides. In some embodiments the nutritive nucleic acid comprises all or part of an open reading frame that encodes a naturally occurring nutritive polypeptide or protein. In some embodiments the nutritive nucleic acid consists of an open reading frame that encodes a fragment of a naturally occurring nutritive protein, wherein the open reading frame does not encode the complete naturally occurring nutritive 10 protein.

[00184] In some embodiments the nutritive nucleic acid is a cDNA.

[00185] In some embodiments nucleic acid molecules are provided that comprise a sequence that is at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 99.9% identical to a naturally occurring nutritive nucleic acid. In some embodiments nucleic 15 acids are provided that hybridize under stringent hybridization conditions with at least one reference nutritive nucleic acid.

[00186] The nutritive nucleic acids and fragments thereof provided in this disclosure display utility in a variety of systems and methods. For example, the fragments may be used as probes in various hybridization techniques. Depending on the method, the target nucleic 20 acid sequences may be either DNA or RNA. The target nucleic acid sequences may be fractionated (e.g., by gel electrophoresis) prior to the hybridization, or the hybridization may be performed on samples *in situ*. One of skill in the art will appreciate that nucleic acid probes of known sequence find utility in determining chromosomal structure (e.g., by Southern blotting) and in measuring gene expression (e.g., by Northern blotting). In such 25 experiments, the sequence fragments are preferably detectably labeled, so that their specific hybridization to target sequences can be detected and optionally quantified. One of skill in the art will appreciate that the nucleic acid fragments of this disclosure may be used in a wide variety of blotting techniques not specifically described herein.

[00187] It should also be appreciated that the nucleic acid sequence fragments disclosed herein also find utility as probes when immobilized on microarrays. Methods for creating microarrays by deposition and fixation of nucleic acids onto support substrates are 30

well known in the art. Reviewed in DNA Microarrays: A Practical Approach (Practical Approach Series), Schena (ed.), Oxford University Press (1999) (ISBN: 0199637768); Nature Genet. 21(1)(suppl):1-60 (1999); Microarray Biochip: Tools and Technology, Schena (ed.), Eaton Publishing Company/BioTechniques Books Division (2000) (ISBN: 1881299376), the disclosures of which are incorporated herein by reference in their entireties. Analysis of, for example, gene expression using microarrays comprising nucleic acid sequence fragments, such as the nucleic acid sequence fragments disclosed herein, is a well-established utility for sequence fragments in the field of cell and molecular biology. Other uses for sequence fragments immobilized on microarrays are described in Gerhold et al., Trends Biochem. Sci. 24:168-173 (1999) and Zweiger, Trends Biotechnol. 17:429-436 (1999); DNA Microarrays: A Practical Approach (Practical Approach Series), Schena (ed.), Oxford University Press (1999) (ISBN: 0199637768); Nature Genet. 21(1)(suppl):1-60 (1999); Microarray Biochip: Tools and Technology, Schena (ed.), Eaton Publishing Company/BioTechniques Books Division (2000) (ISBN: 1881299376).

**15 C. Vectors**

[00188] Also provided are vectors, including expression vectors, which comprise at least one of the nucleic acid molecules disclosed herein, as described further herein. In some embodiments, the vectors comprise at least one isolated nucleic acid molecule encoding a nutritive protein as disclosed herein. In alternative embodiments, the vectors comprise such a nucleic acid molecule operably linked to one or more expression control sequence. The vectors can thus be used to express at least one recombinant protein in a recombinant microbial host cell.

[0016] Suitable vectors for expression of nucleic acids in microorganisms are well known to those of skill in the art. Suitable vectors for use in cyanobacteria are described, for example, in Heidorn et al., "Synthetic Biology in Cyanobacteria: Engineering and Analyzing Novel Functions," Methods in Enzymology, Vol. 497, Ch. 24 (2011). Exemplary replicative vectors that can be used for engineering cyanobacteria as disclosed herein include pPMQAK1, pSL1211, pFC1, pSB2A, pSCR119/202, pSUN119/202, pRL2697, pRL25C, pRL1050, pSG111M, and pPBH201.

[0017] Other vectors such as pJB161 which are capable of receiving nucleic acid sequences disclosed herein may also be used. Vectors such as pJB161 comprise sequences which are homologous with sequences present in plasmids endogenous to certain

photosynthetic microorganisms (e.g., plasmids pAQ1, pAQ3, and pAQ4 of certain *Synechococcus* species). Examples of such vectors and how to use them is known in the art and provided, for example, in Xu et al., “Expression of Genes in Cyanobacteria: Adaptation of Endogenous Plasmids as Platforms for High-Level Gene Expression in *Synechococcus* sp.

- 5 PCC 7002,” Chapter 21 in Robert Carpentier (ed.), “Photosynthesis Research Protocols,” Methods in Molecular Biology, Vol. 684, 2011, which is hereby incorporated herein by reference. Recombination between pJB161 and the endogenous plasmids *in vivo* yield engineered microbes expressing the genes of interest from their endogenous plasmids. Alternatively, vectors can be engineered to recombine with the host cell chromosome, or the
- 10 10 vector can be engineered to replicate and express genes of interest independent of the host cell chromosome or any of the host cell's endogenous plasmids.

[0018] A further example of a vector suitable for recombinant protein production is the pET system (Novagen®). This system has been extensively characterized for use in *E. coli* and other microorganisms. In this system, target genes are cloned in pET plasmids under 15 control of strong bacteriophage T7 transcription and (optionally) translation signals; expression is induced by providing a source of T7 RNA polymerase in the host cell. T7 RNA polymerase is so selective and active that, when fully induced, almost all of the microorganism's resources are converted to target gene expression; the desired product can comprise more than 50% of the total cell protein a few hours after induction. It is also 20 possible to attenuate the expression level simply by lowering the concentration of inducer. Decreasing the expression level may enhance the soluble yield of some target proteins. In some embodiments this system also allows for maintenance of target genes in a transcriptionally silent un-induced state.

[0019] In some embodiments of using this system, target genes are cloned using hosts 25 that do not contain the T7 RNA polymerase gene, thus alleviating potential problems related to plasmid instability due to the production of proteins potentially toxic to the host cell. Once established in a non-expression host, target protein expression may be initiated either by infecting the host with λCE6, a phage that carries the T7 RNA polymerase gene under the control of the λ pL and pI promoters, or by transferring the plasmid into an expression host 30 containing a chromosomal copy of the T7 RNA polymerase gene under lacUV5 control. In the second case, expression is induced by the addition of IPTG or lactose to the bacterial culture or using an autoinduction medium. Other plasmids systems that are controlled by the *lac* operator, but do not require the T7 RNA polymerase gene and rely upon *E. coli*'s native

RNA polymerase include the pTrc plasmid suite (Invitrogen) or pQE plamid suite (QIAGEN).

[0020] In other embodiments it is possible to clone directly into expression hosts. Two types of T7 promoters and several hosts that differ in their stringency of suppressing 5 basal expression levels are available, providing great flexibility and the ability to optimize the expression of a wide variety of target genes.

[00189] Suitable vectors for expression of nucleic acids in mammalian cells typically comprise control functions provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma virus, Adenovirus 2, cytomegalovirus, or Simian 10 Virus 40.

#### D. Promoters

[00190] Promoters useful for expressing the recombinant genes described herein include both constitutive and inducible/repressible promoters. Examples of inducible/repressible promoters include nickel-inducible promoters (e.g., PnrsA, PnrsB ; see, 15 e.g., Lopez-Mauy et al., Cell (2002) v.43: 247-256) and urea repressible promoters such as PnirA (described in, e.g., Qi et al., Applied and Environmental Microbiology (2005) v.71: 5678-5684). Additional examples of inducible/repressible promoters include PnirA (promoter that drives expression of the nirA gene, induced by nitrate and repressed by urea) and Psuf (promoter that drives expression of the sufB gene, induced by iron stress). 20 Examples of constitutive promoters include Pcp (promoter that drives expression of the cpc operon), Prbc (promoter that drives expression of rubisco), PpsbAII (promoter that drives expression of PpsbAII), Pcro (lambda phage promoter that drives expression of cro). In other embodiments, a PaphII and/or a lacIq-Ptrc promoter can be used to control expression. Where multiple recombinant genes are expressed in an engineered microorganism, the different genes 25 can be controlled by different promoters or by identical promoters in separate operons, or the expression of two or more genes may be controlled by a single promoter as part of an operon.

[0021] Further non-limiting examples of inducible promoters may include, but are not limited to, those induced by expression of an exogenous protein (e.g., T7 RNA polymerase, SP6 RNA polymerase), by the presence of a small molecule (e.g., IPTG, galactose, 30 tetracycline, steroid hormone, abscisic acid), by absence of small molecules (e.g., CO<sub>2</sub>, iron, nitrogen), by metals or metal ions (e.g., copper, zinc, cadmium, nickel), and by environmental factors (e.g., heat, cold, stress, light, darkness), and by growth phase. In some

embodiments, the inducible promoter is tightly regulated such that in the absence of induction, substantially no transcription is initiated through the promoter. In some embodiments, induction of the promoter does not substantially alter transcription through other promoters. Also, generally speaking, the compound or condition that induces an 5 inducible promoter is not naturally present in the organism or environment where expression is sought.

[0022] In some embodiments, the inducible promoter is induced by limitation of CO<sub>2</sub> supply to a cyanobacteria culture. By way of non-limiting example, the inducible promoter may be the promoter sequence of *Synechocystis* PCC 6803 that are up-regulated 10 under the CO<sub>2</sub>-limitation conditions, such as the *cmp* genes, *ntp* genes, *ndh* genes, *sbt* genes, *chp* genes, and *rbc* genes, or a variant or fragment thereof.

[0023] In some embodiments, the inducible promoter is induced by iron starvation or by entering the stationary growth phase. In some embodiments, the inducible promoter may be variant sequences of the promoter sequence of cyanobacterial genes that are up-regulated 15 under Fe-starvation conditions such as *isiA*, or when the culture enters the stationary growth phase, such as *isiA*, *phrA*, *sigC*, *sigB*, and *sigH* genes, or a variant or fragment thereof.

[0024] In some embodiments, the inducible promoter is induced by a metal or metal ion. By way of non-limiting example, the inducible promoter may be induced by copper, zinc, cadmium, mercury, nickel, gold, silver, cobalt, and bismuth or ions thereof. In some 20 embodiments, the inducible promoter is induced by nickel or a nickel ion. In some embodiments, the inducible promoter is induced by a nickel ion, such as Ni<sup>2+</sup>. In another exemplary embodiment, the inducible promoter is the nickel inducible promoter from *Synechocystis* PCC 6803. In another embodiment, the inducible promoter may be induced by copper or a copper ion. In yet another embodiment, the inducible promoter may be induced 25 by zinc or a zinc ion. In still another embodiment, the inducible promoter may be induced by cadmium or a cadmium ion. In yet still another embodiment, the inducible promoter may be induced by mercury or a mercury ion. In an alternative embodiment, the inducible promoter may be induced by gold or a gold ion. In another alternative embodiment, the inducible promoter may be induced by silver or a silver ion. In yet another alternative embodiment, the inducible 30 promoter may be induced by cobalt or a cobalt ion. In still another alternative embodiment, the inducible promoter may be induced by bismuth or a bismuth ion.

- [0025] In some embodiments, the promoter is induced by exposing a cell comprising the inducible promoter to a metal or metal ion. The cell may be exposed to the metal or metal ion by adding the metal to the microbial growth media. In certain embodiments, the metal or metal ion added to the microbial growth media may be efficiently recovered from the media.
- 5 In other embodiments, the metal or metal ion remaining in the media after recovery does not substantially impede downstream processing of the media or of the bacterial gene products.

[00191] Further non-limiting examples of constitutive promoters include constitutive promoters from Gram-negative bacteria or a bacteriophage propagating in a Gram-negative bacterium. For instance, promoters for genes encoding highly expressed Gram-negative gene products may be used, such as the promoter for Lpp, OmpA, rRNA, and ribosomal proteins.

10 Alternatively, regulatable promoters may be used in a strain that lacks the regulatory protein for that promoter. For instance  $P_{lac}$ ,  $P_{tac}$ , and  $P_{trc}$ , may be used as constitutive promoters in strains that lack LacI. Similarly, P22  $P_R$  and  $P_L$  may be used in strains that lack the lambda C2 repressor protein, and lambda  $P_R$  and  $P_L$  may be used in strains that lack the lambda C1 repressor protein. In one embodiment, the constitutive promoter is from a bacteriophage. In another embodiment, the constitutive promoter is from a *Salmonella* bacteriophage. In yet another embodiment, the constitutive promoter is from a cyanophage. In some embodiments, the constitutive promoter is a *Synechocystis* promoter. For instance, the constitutive promoter may be the *PpsbA* promoter or its variant sequences, the *Prbc* promoter or its variant sequences, the  $P_{cpc}$  promoter or its variant sequences, and the *PrnpB* promoter or its variant sequences.

15

20

#### E. Host Cells

[00192] Also provided are host cells transformed with the nucleic acid molecules or vectors disclosed herein, and descendants thereof. In some embodiments the host cells are microbial cells. In some embodiments, the host cells carry the nucleic acid sequences on vectors, which may but need not be freely replicating vectors. In other embodiments, the nucleic acids have been integrated into the genome of the host cells and/or into an endogenous plasmid of the host cells. The transformed host cells find use, e.g., in the production of recombinant nutritive proteins disclosed herein.

25

30 [0026] “Microorganisms” includes prokaryotic and eukaryotic microbial species from the Domains *Archaea*, *Bacteria* and *Eucarya*, the latter including yeast and filamentous

fungi, protozoa, algae, or higher Protista. The terms “microbial cells” and “microbes” are used interchangeably with the term microorganism.

[0027] A variety of host microorganisms may be transformed with a nucleic acid sequence disclosed herein and may in some embodiments be used to produce a recombinant nutritive protein disclosed herein. Suitable host microorganisms include both autotrophic and heterotrophic microbes. In some applications the autotrophic microorganisms allows for a reduction in the fossil fuel and/or electricity inputs required to make a nutritive protein encoded by a recombinant nucleic acid sequence introduced into the host microorganism. This, in turn, in some applications reduces the cost and/or the environmental impact of producing the nutritive protein and/or reduces the cost and/or the environmental impact in comparison to the cost and/or environmental impact of manufacturing alternative nutritive proteins, such as whey, egg, and soy. For example, the cost and/or environmental impact of making a nutritive protein disclosed herein using a host microorganism as disclosed herein is in some embodiments lower than the cost and/or environmental impact of making whey protein in a form suitable for human consumption by processing of cows milk.

[0028] Non-limiting examples of heterotrophs include *Escherichia coli*, *Salmonella typhimurium*, *Bacillus subtilis*, *Bacillus megaterium*, *Corynebacterium glutamicum*, *Streptomyces coelicolor*, *Streptomyces lividans*, *Streptomyces venezuelae*, *Streptomyces roseosporus*, *Streptomyces fradiae*, *Streptomyces griseus*, *Streptomyces calvuligerus*, *Streptomyces hygroscopicus*, *Streptomyces platensis*, *Saccharopolyspora erythraea*, *Corynebacterium glutamicum*, *Aspergillus niger*, *Aspergillus nidulans*, *Aspergillus oryzae*, *Aspergillus terreus*, *Aspergillus sojae*, *Penicillium chrysogenum*, *Trichoderma reesei*, *Clostridium acetobutylicum*, *Clostridium beijerinckii*, *Clostridium thermocellum*, *Fusibacter paucivorans*, *Saccharomyces cerevisiae*, *Saccharomyces boulardii*, *Pichia pastoris*, and *Pichia stipitis*.

[0029] Photoautotrophic microorganisms include eukaryotic algae, as well as prokaryotic cyanobacteria, green-sulfur bacteria, green non-sulfur bacteria, purple sulfur bacteria, and purple non-sulfur bacteria.

[0030] Extremophiles are also contemplated as suitable organisms. Such organisms withstand various environmental parameters such as temperature, radiation, pressure, gravity, vacuum, desiccation, salinity, pH, oxygen tension, and chemicals. They include hyperthermophiles, which grow at or above 80°C such as *Pyrolobus fumarii*; thermophiles,

which grow between 60-80°C such as *Synechococcus lividis*; mesophiles, which grow between 15-60°C; and psychrophiles, which grow at or below 15°C such as *Psychrobacter* and some insects. Radiation tolerant organisms include *Deinococcus radiodurans*. Pressure-tolerant organisms include piezophiles, which tolerate pressure of 130 MPa. Weight-tolerant 5 organisms include barophiles. Hypergravity (e.g., >1g) hypogravity (e.g., <1g) tolerant organisms are also contemplated. Vacuum tolerant organisms include tardigrades, insects, microbes and seeds. Dessicant tolerant and anhydrobiotic organisms include xerophiles such as *Artemia salina*; nematodes, microbes, fungi and lichens. Salt-tolerant organisms include halophiles (e.g., 2-5 M NaCl) *Halobacteriaceae* and *Dunaliella salina*. pH-tolerant organisms 10 include alkaliphiles such as *Natronobacterium*, *Bacillus firmus* OF4, *Spirulina* spp. (e.g., pH > 9) and acidophiles such as *Cyanidium caldarium*, *Ferroplasma* sp. (e.g., low pH). Anaerobes, which cannot tolerate O<sub>2</sub> such as *Methanococcus jannaschii*; microaerophils, which tolerate some O<sub>2</sub> such as *Clostridium* and aerobes, which require O<sub>2</sub> are also contemplated. Gas-tolerant organisms, which tolerate pure CO<sub>2</sub> include *Cyanidium caldarium* 15 and metal tolerant organisms include metalotolerants such as *Ferroplasma acidarmanus* (e.g., Cu, As, Cd, Zn), *Ralstonia* sp. CH34 (e.g., Zn, Co, Cd, Hg, Pb). Gross, Michael. *Life on the Edge: Amazing Creatures Thriving in Extreme Environments*. New York: Plenum (1998) and Seckbach, J. "Search for Life in the Universe with Terrestrial Microbes Which Thrive Under Extreme Conditions." In Cristiano Batalli Cosmovici, Stuart Bowyer, and Dan Wertheimer, 20 eds., *Astronomical and Biochemical Origins and the Search for Life in the Universe*, p. 511. Milan: Editrice Compositori (1997).

[0031] Mixotrophic organisms are also suitable organisms. Mixotrophic organisms can utilize a mix of different sources of energy and carbon, for example, photo- and chemotrophy, litho- and organotrophy, auto- and heterotrophy, and combinations 25 thereof or a combination of it. Mixotrophs can be either eukaryotic or prokaryotic. Additionally, mixotrophs can be obligate or facultative. Suitable mixotrophic organisms include mixotrophic algae and mixotrophic bacteria.

[0032] Algae and cyanobacteria include but are not limited to the following genera: Acanthoceras, Acanthococcus, Acaryochloris, Achnanthes, Achnanthidium, Actinastrum, 30 Actinochloris, Actinocyclus, Actinotaenium, Amphichrysis, Amphidinium, Amphikrikos, Amphipleura, Amphiprora, Amphithrix, Amphora, Anabaena, Anabaenopsis, Aneumastus, Ankistrodesmus, Ankyra, Anomoeoneis, Apatococcus, Aphanizomenon, Aphanocapsa, Aphanochaete, Aphanothece, Apiocystis, Apistonema, Arthrodesmus, Artherospira,

Ascochloris, Asterionella, Asterococcus, Audouinella, Aulacoscira, Bacillaria, Balbiania, Bambusina, Bangia, Basichlamys, Batrachospermum, Binuclearia, Bitrichia, Blidingia, Botriopsis, Botrydium, Botryococcus, Botryosphaerella, Brachiomonas, Brachysira, Brachytrichia, Brebissonia, Bulbochaete, Bumilleria, Bumilleriopsis, Caloneis, Calothrix,  
5 Campylodiscus, Capsosiphon, Carteria, Catena, Cavinula, Centritractus, Centronella, Ceratium, Chaetoceros, Chaetochloris, Chaetomorpha, Chaetonella, Chaetonema, Chaetopeltis, Chaetophora, Chaetosphaeridium, Chamaesiphon, Chara, Characiocloris, Characiopsis, Characium, Charales, Chilomonas, Chlainomonas, Chlamydoblepharis, Chlamydocapsa, Chlamydomonas, Chlamydomonopsis, Chlamydomyxa, Chlamydonephris,  
10 Chlorangiella, Chlorangiopsis, Chlorella, Chlorobotrys, Chlorobrachis, Chlorochytrium, Chlorococcum, Chlorogloea, Chlorogloeopsis, Chlorogonium, Chlorolobion, Chloromonas, Chlorophysema, Chlorophyta, Chlorosaccus, Chlorosarcina, Choricystis, Chromophyton, Chromulina, Chroococcidiopsis, Chroococcus, Chroodactylon, Chroomonas, Chroothece, Chrysamoeba, Chrysopsis, Chrysidiastrum, Chrysocapsa, Chrysocapsella, Chrysochaete,  
15 Chrysochromulina, Chrysococcus, Chrysocrinus, Chrysolepidomonas, Chrysolykos, Chrysonebula, Chrysophyta, Chrysopyxis, Chrysosaccus, Chrysphaerella, Chrysostephanosphaera, Clodophora, Clastidium, Closteriopsis, Closterium, Coccomyxa, Coccineis, Coelastrella, Coelastrum, Coelosphaerium, Coenochloris, Coenococcus, Coenocystis, Colacium, Coleochaete, Collodictyon, Compsogonopsis, Compsopogon,  
20 Conjugatophyta, Conochaete, Coronastrum, Cosmarium, Cosmioneis, Cosmocladium, Crateriportula, Craticula, Crinalium, Crucigenia, Crucigeniella, Cryptoaulax, Cryptomonas, Cryptophyta, Ctenophora, Cyanodictyon, Cyanonephron, Cyanophora, Cyanophyta, Cyanothece, Cyanothomonas, Cyclonexis, Cyclostephanos, Cyclotella, Cylindrocapsa, Cylindrocystis, Cylindrospermum, Cylindrotheca, Cymatopleura, Cymbella,  
25 Cymbellonitzschia, Cystodinium Dactylococcopsis, Debarya, Denticula, Dermatochrysis, Dermocarpa, Dermocarpella, Desm tractum, Desmidium, Desmococcus, Desmonema, Desmosiphon, Diacanthos, Diacronema, Diadesmis, Diatoma, Diatomella, Dicellula, Diclobothrix, Dichotomococcus, Dicranochaete, Dictyochloris, Dictyococcus, Dictyosphaerium, Didymocystis, Didymogenes, Didymosphenia, Dilabifilum,  
30 Dimorphococcus, Dinobryon, Dinococcus, Diplochloris, Diploneis, Diplostauron, Distri onella, Docidium, Draparnaldia, Dunaliella, Dysmorphococcus, Ecballocystis, Elakatothrix, Ellerbeckia, Encyonema, Enteromorpha, Entocladia, Entomoneis, Entophysalis, Epichrysis, Epipyxis, Epithemia, Eremosphaera, Euastropsis, Euastrum, Eucapsis, Eucocconeis, Eudorina, Euglena, Euglenophyta, Eunotia, Eustigmatophyta, Eutreptia,

Fallacia, Fischerella, Fragilaria, Fragilariforma, Franceia, Frustulia, Curcilla, Geminella,  
Genicularia, Glaucocystis, Glauco phyta, Glenodiniopsis, Glenodinium, Gloeocapsa,  
Gloeochaete, Gloeochrysis, Gloeococcus, Gloeocystis, Gloeodendron, Gloeomonas,  
Gloeoplax, Gloeothece, Gloeotila, Gloeotrichia, Gloiodictyon, Golenkinia, Golenkiniopsis,  
5 Gomontia, Gomphocymbella, Gomphonema, Gomphosphaeria, Gonatozygon, Gongrosia,  
Gongrosira, Goniochloris, Gonium, Gonyostomum, Granulochloris, Granulocystopsis,  
Groenbladia, Gymnodinium, Gymnozyga, Gyrosigma, Haematococcus, Hafniomonas,  
Hallassia, Hammatoidea, Hannaea, Hantzschia, Hapalosiphon, Haplotaenium, Haptophyta,  
Haslea, Hemidinium, Hemitoma, Heribaudiella, Heteromastix, Heterothrix, Hibberdia,  
10 Hildenbrandia, Hillea, Holopedium, Homoeothrix, Hormanthonema, Hormotila,  
Hyalobrachion, Hyalocardium, Hyalodiscus, Hyalogonium, Hyalotheca, Hydrianum,  
Hydrococcus, Hydrocoleum, Hydrocoryne, Hydrodictyon, Hydrosera, Hydrurus, Hyella,  
Hymenomonas, Isthmochloron, Johannesbaptistia, Juranyiella, Karayevia, Kathablepharis,  
Katodinium, Kephryion, Keratococcus, Kirchneriella, Klebsormidium, Kolbesia, Koliella,  
15 Komarekia, Korshikoviella, Kraskella, Lagerheimia, Lagynion, Lamprothamnium, Lemanea,  
Lepocinclis, Leptosira, Lobococcus, Lobocystis, Lobomonas, Luticola, Lyngbya,  
Malleochloris, Mallomonas, Mantiella, Marssonella, Martyana, Mastigocoleus,  
Gastogloia, Melosira, Merismopedia, Mesostigma, Mesotaenium, Micractinium,  
Micrasterias, Microchaete, Microcoleus, Microcystis, Microglena, Micromonas, Microspora,  
20 Microthamnion, Mischooccus, Monochrysis, Monodus, Monomastix, Monoraphidium,  
Monostroma, Mougeotia, Mougeotiopsis, Myochloris, Myromecia, Myxosarcina,  
Naegeliella, Nannochloris, Nautococcus, Navicula, Neglectella, Neidium, Nephroclamys,  
Nephrocytium, Nephrodiella, Nephroelmis, Netrium, Nitella, Nitellopsis, Nitzschia,  
Nodularia, Nostoc, Ochromonas, Oedogonium, Oligochaetophora, Onychonema, Oocardium,  
25 Oocystis, Opephora, Ophiocytium, Orthoseira, Oscillatoria, Oxyneis, Pachycladella,  
Palmella, Palmodictyon, Pnadorina, Pannus, Paralia, Pascherina, Paulschulzia, Pediastrum,  
Pedinella, Pedinomonas, Pedinopera, Pelagodictyon, Penium, Peranema, Peridiniopsis,  
Peridinium, Peronia, Petroneis, Phacotus, Phacus, Phaeaster, Phaeodermatium, Phaeophyta,  
Phaeosphaera, Phaeothamnion, Phormidium, Phycopeltis, Phyllariochloris, Phyllocardium,  
30 Phyllomitas, Pinnularia, Pitophora, Placoneis, Planctonema, Planktosphaeria, Planothidium,  
Plectonema, Pleodorina, Pleurastrum, Pleurocapsa, Pleurocladia, Pleurodiscus, Pleurosigma,  
Pleurosira, Pleurotaenium, Pocillomonas, Podohedra, Polyblepharides, Polychaetophora,  
Polyedriella, Polyedriopsis, Polygoniochloris, Polyepidomonas, Polytaenia, Polytoma,  
Polytomella, Porphyridium, Posteriochromonas, Prasinochloris, Prasinocladus, Prasinophyta,

Prasiola, Prochlorphyta, Prochlorothrix, Protoderma, Protosiphon, Provasoliella, Prymnesium, Psammodictyon, Psammothidium, Pseudanabaena, Pseudenoclonium, Psuedocarteria, Pseudochate, Pseudocharacium, Pseudococcoomyxa, Pseudodictyosphaerium, Pseudokephyrion, Pseudoncobyrsa, Pseudoquadrigula, Pseudosphaerocystis,

5 Pseudostaurastrum, Pseudostaurosira, Pseudotetrastrum, Pteromonas, Punctastruata, Pyramichlamys, Pyramimonas, Pyrrophyta, Quadrichloris, Quadricoccus, Quadrigula, Radiococcus, Radiofilm, Raphidiopsis, Raphidocelis, Raphidonema, Raphidophyta, Peimeria, Rhabdoderma, Rhabdomonas, Rhizoclonium, Rhodomonas, Rhodophyta, Rhoicosphenia, Rhopalodia, Rivularia, Rosenvingiella, Rossithidium, Roya, Scenedesmus,

10 Scherffelia, Schizochlamydella, Schizochlamys, Schizomeris, Schizothrix, Schroederia, Scolioneis, Scottiella, Scotiellopsis, Scourfieldia, Scytonema, Selenastrum, Selenochloris, Sellaphora, Semiorbis, Siderocelis, Diderocystopsis, Dimonsenia, Siphononema, Sirocladium, Sirogonium, Skeletonema, Sorastrum, Spennatozopsis, Sphaerellocystis, Sphaerellopsis, Sphaerodinium, Sphaeroplea, Sphaerozosma, Spiniferomonas, Spirogyra,

15 Spirotaenia, Spirulina, Spondylomorum, Spondylosium, Sporotetras, Spumella, Staurastrum, Staurodesmus, Stauroneis, Staurosira, Staurosirella, Stenopterobia, Stephanocostis, Stephanodiscus, Stephanoporus, Stephanosphaera, Stichococcus, Stichogloea, Stigeoclonium, Stigonema, Stipitococcus, Stokesiella, Strombomonas, Stylochrysalis, Stylocladium, Styloysis, Stylosphaeridium, Surirella, Sykidion, Symploca, Synechococcus, Synechocystis,

20 Synedra, Synochromonas, Synura, Tabellaria, Tabularia, Teilingia, Temnogametum, Tetmemorus, Tetrachlorella, Tetracyclus, Tetraedromus, Tetraedriella, Tetraedron, Tetraselmis, Tetraspora, Tetrastrum, Thalassiosira, Thamniochaete, Thorakochloris, Thorea, Tolypella, Tolypothrix, Trachelomonas, Trachydiscus, Trebouxia, Trentepholia, Treubaria, Tribonema, Trichodesmium, Trichodiscus, Trochiscia, Tryblionella, Ulothrix, Uroglena,

25 Uronema, Urosolenia, Urospora, Uva, Vacuolaria, Vaucheria, Volvox, Volvulina, Westella, Woloszynskia, Xanthidium, Xanthophyta, Xenococcus, Zygema, Zyg nemopsis, and Zygionium.

[0033] Additional cyanobacteria include members of the genus Chamaesiphon, Chroococcus, Cyanobacterium, Cyanobium, Cyanothece, Dactylococcopsis, Gloeobacter, 30 Gloeocapsa, Gloeothece, Microcystis, Prochlorococcus, Prochloron, Synechococcus, Synechocystis, Cyanocystis, Dermocarrella, Stanieria, Xenococcus, Chroococcidiopsis, Myxosarcina, Arthrosira, Borzia, Crinalium, Geitlerinemia, Leptolyngbya, Limnothrix, Lyngbya, Microcoleus, Oscillatoria, Planktothrix, Prochlorothrix, Pseudanabaena, Spirulina,

Starria, Symploca, Trichodesmium, Tychonema, Anabaena, Anabaenopsis, Aphanizomenon, Cyanospira, Cylindrospermopsis, Cylindrospermum, Nodularia, Nostoc, Scytonema, Calothrix, Rivularia, Tolypothrix, Chlorogloeopsis, Fischerella, Geitiera, Iyengariella, Nostochopsis, Stigonema and Thermosynechococcus.

5 [0034] Green non-sulfur bacteria include but are not limited to the following genera: Chloroflexus, Chloronema, Oscillochloris, Heliothrix, Herpetosiphon, Roseiflexus, and Thermomicrombium.

[0035] Green sulfur bacteria include but are not limited to the following genera: *Chlorobium*, *Clathrochloris*, and *Prosthecochloris*.

10 [0036] Purple sulfur bacteria include but are not limited to the following genera: Allochromatium, Chromatium, Halochromatium, Isochromatium, Marichromatium, Rhodovulum, Thermochromatium, Thiocapsa, Thiorhodococcus, and Thiocystis.

[0037] Purple non-sulfur bacteria include but are not limited to the following genera: Phaeospirillum, Rhodobaca, Rhodobacter, Rhodomicrobium, Rhodopila,

15 Rhodopseudomonas, Rhodothalassium, Rhodospirillum, Rodovibrio, and Roseospira.

[0038] Aerobic chemolithotrophic bacteria include but are not limited to nitrifying bacteria such as *Nitrobacteraceae* sp., *Nitrobacter* sp., *Nitrospina* sp., *Nitrococcus* sp., *Nitrospira* sp., *Nitrosomonas* sp., *Nitrosococcus* sp., *Nitrosospira* sp., *Nitrosolobus* sp., *Nitrosovibrio* sp.; colorless sulfur bacteria such as, *Thiovulum* sp., *Thiobacillus* sp.,

20 *Thiomicrospira* sp., *Thiosphaera* sp., *Thermothrix* sp.; obligately chemolithotrophic hydrogen bacteria such as *Hydrogenobacter* sp., iron and manganese-oxidizing and/or depositing bacteria such as *Siderococcus* sp., and magnetotactic bacteria such as *Aquaspirillum* sp.

[0039] Archaeobacteria include but are not limited to methanogenic archaeobacteria such as *Methanobacterium* sp., *Methanobrevibacter* sp., *Methanothermus* sp., *Methanococcus*

25 sp., *Methanomicrobium* sp., *Methanospirillum* sp., *Methanogenium* sp., *Methanosarcina* sp., *Methanolobus* sp., *Methanothrix* sp., *Methanococcoides* sp., *Methanoplanus* sp.; extremely thermophilic S-Metabolizers such as *Thermoproteus* sp., *Pyrodictium* sp., *Sulfolobus* sp., *Acidianus* sp. and other microorganisms such as, *Bacillus subtilis*, *Saccharomyces cerevisiae*, *Streptomyces* sp., *Ralstonia* sp., *Rhodococcus* sp., *Corynebacteria* sp., *Brevibacteria* sp.,

30 *Mycobacteria* sp., and oleaginous yeast.

[0040] Yet other suitable organisms include synthetic cells or cells produced by synthetic genomes as described in Venter et al. US Pat. Pub. No. 2007/0264688, and cell-like systems or synthetic cells as described in Glass et al. US Pat. Pub. No. 2007/0269862.

[0041] Still other suitable organisms include *Escherichia coli*, *Acetobacter aceti*,  
5 *Bacillus subtilis*, yeast and fungi such as *Clostridium ljungdahlii*, *Clostridium thermocellum*,  
*Penicillium chrysogenum*, *Pichia pastoris*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pseudomonas fluorescens*, or *Zymomonas mobilis*. In some embodiments those organisms are engineered to fix carbon dioxide while in other embodiments they are not.

[00193] In some embodiments eukaryotic cells, such as insect cells or mammalian  
10 cells, such as human cells are used as host cells. Vectors and expression control sequences including promoters and enhancers are well known for such cells. Examples of useful mammalian host cell lines for this purpose are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster  
15 kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells  
20 (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

#### **F. Production of Nutritive Proteins**

[00194] Skilled artisans are aware of many suitable methods available for culturing recombinant cells to produce (and optionally secrete) a recombinant nutritive protein as disclosed herein, as well as for purification and/or isolation of expressed recombinant proteins. The methods chosen for protein purification depend on many variables, including the properties of the protein of interest, its location and form within the cell, the vector, host  
25 strain background, and the intended application for the expressed protein. Culture conditions can also have an effect on solubility and localization of a given target protein. Many

approaches can be used to purify target proteins expressed in recombinant microbial cells as disclosed herein, including without limitation ion exchange and gel filtration.

[00195] In some embodiments a peptide fusion tag is added to the recombinant protein making possible a variety of affinity purification methods that take advantage of the peptide fusion tag. In some embodiments, the use of an affinity method enables the purification of the target protein to near homogeneity in one step. Purification may include cleavage of part or all of the fusion tag with enterokinase, factor Xa, thrombin, or HRV 3C proteases, for example. In some embodiments, before purification or activity measurements of an expressed target protein, preliminary analysis of expression levels, cellular localization, and solubility of the target protein is performed. The target protein may be found in any or all of the following fractions: soluble or insoluble cytoplasmic fractions, periplasm, or medium. Depending on the intended application, preferential localization to inclusion bodies, medium, or the periplasmic space can be advantageous, in some embodiments, for rapid purification by relatively simple procedures.

[00196] While *Escherichia coli* is widely regarded as a robust host for heterologous protein expression, it is also widely known that over-expression of many proteins in this host is prone to aggregation in the form of insoluble inclusion bodies. One of the most commonly used methods for either rescuing inclusion body formation, or to improve the titer of the protein itself, is to include an amino-terminal maltose-binding protein (MBP) [Austin BP, Nallamsetty S, Waugh DS. Hexahistidine-tagged maltose-binding protein as a fusion partner for the production of soluble recombinant proteins in *Escherichia coli*. *Methods Mol Biol.* 2009;498:157-72], or small ubiquitin-related modifier (SUMO) [Saitoh H, Uwada J, Azusa K. Strategies for the expression of SUMO-modified target proteins in *Escherichia coli*. *Methods Mol Biol.* 2009;497:211-21; Malakhov MP, Mattern MR, Malakhova OA, Drinker M, Weeks SD, Butt TR. SUMO fusions and SUMO-specific protease for efficient expression and purification of proteins. *J Struct Funct Genomics.* 2004;5(1-2):75-86; Panavas T, Sanders C, Butt TR. SUMO fusion technology for enhanced protein production in prokaryotic and eukaryotic expression systems. *Methods Mol Biol.* 2009;497:303-17] fusion to the protein of interest. These two proteins are expressed extremely well, and in the soluble form, in *Escherichia coli* such that the protein of interest is also effectively produced in the soluble form. The protein of interest can be cleaved by designing a site specific protease recognition

sequence (such as the tobacco etch virus (TEV) protease) in-between the protein of interest and the fusion protein [1].

[00197] In some embodiments the recombinant protein is initially not folded correctly or is insoluble. A variety of methods are well known for refolding of insoluble proteins.

5 Most protocols comprise the isolation of insoluble inclusion bodies by centrifugation followed by solubilization under denaturing conditions. The protein is then dialyzed or diluted into a non-denaturing buffer where refolding occurs. Because every protein possesses unique folding properties, the optimal refolding protocol for any given protein can be empirically determined by a skilled artisan. Optimal refolding conditions can, for example, 10 be rapidly determined on a small scale by a matrix approach, in which variables such as protein concentration, reducing agent, redox treatment, divalent cations, etc., are tested. Once the optimal concentrations are found, they can be applied to a larger scale solubilization and refolding of the target protein.

[00198] In some embodiments the nutritive protein does not comprise a tertiary structure. In some embodiments less than half of the amino acids in the nutritive protein participate in a tertiary structure. In some embodiments the nutritive protein does not comprise a secondary structure. In some embodiments less than half of the amino acids in the nutritive protein participate in a secondary structure. Recombinant nutritive proteins may be isolated from a culture of cells expressing them in a state that comprises one or more of these 20 structural features. In some embodiments the tertiary structure of a recombinant nutritive protein is reduced or eliminated after the protein is isolated from a culture producing it. In some embodiments the secondary structure of a recombinant nutritive protein is reduced or eliminated after the protein is isolated from a culture producing it.

[00199] In some embodiments a CAPS buffer at alkaline pH in combination with N-lauroylsarcosine is used to achieve solubility of the inclusion bodies, followed by dialysis in the presence of DTT to promote refolding. Depending on the target protein, expression conditions, and intended application, proteins solubilized from washed inclusion bodies may be > 90% homogeneous and may not require further purification. Purification under fully denaturing conditions (before refolding) is possible using His•Tag® fusion proteins and 30 His•Bind® immobilized metal affinity chromatography (Novogen®). In addition, S•Tag™, T7•Tag®, and Strep•Tag® II fusion proteins solubilized from inclusion bodies using 6 M urea can be purified under partially denaturing conditions by dilution to 2 M urea (S•Tag and

T7•Tag) or 1 M urea (Strep•Tag II) prior to chromatography on the appropriate resin. Refolded fusion proteins can be affinity purified under native conditions using His•Tag, S•Tag, Strep•Tag II, and other appropriate affinity tags (e.g., GST•Tag<sup>TM</sup>, and T7•Tag) (Novogen®).

5 [00200] In some embodiments the recombinant nutritive protein is an endogenous protein of the host cell used to express it. That is, the cellular genome of the host cell comprises an open reading frame that encodes the recombinant nutritive protein. In some embodiments regulatory sequences sufficient to increase expression of the nutritive protein are inserted into the host cell genome and operatively linked to the endogenous open reading  
10 frame such that the regulatory sequences drive overexpression of the recombinant nutritive protein from a recombinant nucleic acid. In some embodiments heterologous nucleic acid sequences are fused to the endogenous open reading frame of the nutritive protein and cause the nutritive protein to be synthesized comprising a heterologous amino acid sequence that changes the cellular trafficking of the recombinant nutritive protein, such as directing it to an  
15 organelle or to a secretion pathway. In some embodiments an open reading frame that encodes the endogenous host cell protein is introduced into the host cell on a plasmid that further comprises regulatory sequences operatively linked to the open reading frame. In some embodiments the recombinant host cell expresses at least 2 times, at least 3 times, at least 4 times, at least 5 times, at least 10 times, or at least 20 times, at least 30 times, at least  
20 40 times, at least 50 times, or at least 100 times more of the recombinant nutritive protein than the amount of the nutritive protein produced by a similar host cell grown under similar conditions.

25 [00201] In some embodiments nutritive proteins of this disclosure are synthesized chemically without the use of a recombinant production system. Protein synthesis can be carried out in a liquid-phase system or in a solid-phase system using techniques known in the art (see, e.g., Atherton, E., Sheppard, R.C. (1989). Solid Phase peptide synthesis: a practical approach. Oxford, England: IRL Press; Stewart, J.M., Young, J.D. (1984). Solid phase peptide synthesis (2nd ed.). Rockford: Pierce Chemical Company.

#### **G. Production of Recombinant Nutritive Proteins in Plants**

30 [00202] The nucleic acid molecules comprising a nucleic acid sequence encoding a nutritive protein of this disclosure enable production of transgenic plants comprising the nucleic acid sequence. Accordingly, this disclosure also provides plant comprising a

recombinant nucleic acid molecule comprising a nucleic acid sequence encoding a nutritive protein of this disclosure. The plant can be any plant that is subject to transformation and regeneration and include, but are not limited to, Acacia, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, 5 broccoli, brussels sprouts, cabbage, canola, cantaloupe, carrot, cassaya, cauliflower, celery, Chinese cabbage, cherry, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, forest trees, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom, nut, oat, okra, onion, orange, an ornamental plant, papaya, parsley, pea, 10 peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, rapeseed, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet corn, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, turf, a vine, watermelon, wheat, yams, and zucchini. In preferred embodiments, the plant is a bean, 15 broccoli, cabbage, canola, carrot, cauliflower, celery, Chinese cabbage, corn, cotton cucumber, eggplant, leek, lettuce, melon, pea, pepper, pumpkin, radish, spinach, soybean, squash, sugarcane, sweet corn, tomato, watermelon, and wheat plant. In some embodiments, the plant is a corn plant. In some embodiments, the plant is a soybean plant. In some embodiments, the plant is a cotton plant. In some embodiments, the plant is a canola plant. In 20 some embodiments the plant is a member of a genus selected from *Arabidopsis*, *Beta*, *Glycine*, *Jatropha*, *Miscanthus*, *Panicum*, *Phalaris*, *Populus*, *Saccharum*, *Salix*, *Simmondsia* and *Zea*.

[00203] Numerous promoters that are active in plant cells have been described in the literature. These include promoters present in plant genomes as well as promoters from other 25 sources, including nopaline synthase (NOS) promoter and octopine synthase (OCS) promoters carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*, caulimovirus promoters such as the cauliflower mosaic virus. For instance, see U.S. Pat. Nos. 5,858,742 and 5,322,938, which disclose versions of the constitutive promoter derived from cauliflower mosaic virus (CaMV35S), U.S. Pat. No. 5,641,876, which discloses a rice actin promoter, 30 U.S. Patent Application Publication 2002/0192813A1, which discloses 5', 3' and intron elements useful in the design of effective plant expression vectors, U.S. patent application Ser. No. 09/757,089, which discloses a maize chloroplast aldolase promoter, U.S. patent application Ser. No. 08/706,946, which discloses a rice glutelin promoter, U.S. patent

application Ser. No. 09/757,089, which discloses a maize aldolase (FDA) promoter, and U.S. patent application Ser. No. 60/310,370, which discloses a maize nicotianamine synthase promoter. These and numerous other promoters that function in plant cells are known to those skilled in the art and available for use in recombinant nucleic acids to provide for expression 5 of nutritive proteins in transgenic plants.

[00204] For some applications preferential expression in plant green tissues is desired. Promoters of interest for such uses include those from genes such as *Arabidopsis thaliana* ribulose-1,5-bisphosphate carboxylase (Rubisco) small subunit (Fischhoff et al. (1992) *Plant Mol. Biol.* 20:81-93), aldolase and pyruvate orthophosphate dikinase (PPDK) (Taniguchi et 10 al. (2000) *Plant Cell Physiol.* 41(1):42-48).

[0036] Furthermore, the promoters may be altered to contain at least one enhancer sequence to assist in elevating gene expression. Such enhancers are known in the art. By including an enhancer sequence with such constructs, the expression of the nutritive protein may be enhanced. These enhancers often are found 5' to the start of transcription in a promoter that 15 functions in eukaryotic cells, but can often be inserted upstream (5') or downstream (3') to the coding sequence. In some instances, these 5' enhancing elements are introns. Particularly useful as enhancers are the 5' introns of the rice actin 1 (see U.S. Pat. No. 5,641,876) and rice actin 2 genes, the maize alcohol dehydrogenase gene intron, the maize heat shock protein 70 gene intron (U.S. Pat. No. 5,593,874) and the maize shrunken 1 gene.

20 [00205] For some applications expression in plant seed tissues is desired to effect modify seed composition. Exemplary promoters for use for seed composition modification include promoters from seed genes such as napin (U.S. Pat. No. 5,420,034), maize L3 oleosin (U.S. Pat. No. 6,433,252), zein Z27 (Russell et al. (1997) *Transgenic Res.* 6(2):157-166), globulin 1 (Belanger et al (1991) *Genetics* 129:863-872), glutelin 1 (Russell (1997) *supra*), 25 and peroxiredoxin antioxidant (Per1) (Stacy et al. (1996) *Plant Mol. Biol.* 31(6):1205-1216)

[00206] Recombinant nucleic acid constructs prepared in accordance with the disclosure will also generally include a 3' element that typically contains a polyadenylation signal and site. Well-known 3' elements include those from *Agrobacterium tumefaciens* genes such as nos 3', tml 3', tmr 3', tms 3', ocs 3', tr7 3', for example disclosed in U.S. Pat. 30 No. 6,090,627; 3' elements from plant genes such as wheat (*Triticum aestivum*) heat shock protein 17 (Hsp17 3'), a wheat ubiquitin gene, a wheat fructose-1,6-biphosphatase gene, a rice glutelin gene a rice lactate dehydrogenase gene and a rice beta-tubulin gene, all of which

are disclosed in U.S. published patent application 2002/0192813 A1; and the pea (*Pisum sativum*) ribulose biphosphate carboxylase gene (rbs 3'), and 3' elements from the genes within the host plant.

5 [00207] Constructs and vectors may also include a transit peptide for targeting of a gene target to a plant organelle, particularly to a chloroplast, leucoplast or other plastid organelle. For descriptions of the use of chloroplast transit peptides see U.S. Pat. No. 5,188,642 and U.S. Pat. No. 5,728,925. For description of the transit peptide region of an *Arabidopsis* EPSPS gene, see Klee, H. J. et al (MGG (1987) 210:437-442).

10 [00208] Numerous methods for transforming plant cells with recombinant DNA are known in the art and may be used in the present disclosure. Two commonly used methods for plant transformation are Agrobacterium-mediated transformation and microprojectile bombardment. Microprojectile bombardment methods are illustrated in U.S. Pat. Nos. 5,015,580 (soybean); 5,550,318 (corn); 5,538,880 (corn); 5,914,451 (soybean); 6,160,208 (corn); 6,399,861 (corn) and 6,153,812 (wheat) and Agrobacterium-mediated transformation 15 is described in U.S. Pat. Nos. 5,159,135 (cotton); 5,824,877 (soybean); 5,591,616 (corn); and 6,384,301 (soybean). For *Agrobacterium tumefaciens* based plant transformation system, additional elements present on transformation constructs will include T-DNA left and right border sequences to facilitate incorporation of the recombinant polynucleotide into the plant genome.

20 [00209] In general it is useful to introduce recombinant DNA randomly, i.e. at a non-specific location, in the genome of a target plant line. In special cases it may be useful to target recombinant DNA insertion in order to achieve site-specific integration, for example to replace an existing gene in the genome, to use an existing promoter in the plant genome, or to insert a recombinant polynucleotide at a predetermined site known to be active for gene 25 expression. Several site specific recombination systems exist which are known to function in plants, including cre-lox as disclosed in U.S. Pat. No. 4,959,317 and FLP-FRT as disclosed in U.S. Pat. No. 5,527,695.

30 [00210] Transformation methods are generally practiced in tissue culture on media and in a controlled environment. "Media" refers to the numerous nutrient mixtures that are used to grow cells in vitro, that is, outside of the intact living organism. Recipient cell targets include, but are not limited to, meristem cells, callus, immature embryos and gametic cells

such as microspores, pollen, sperm and egg cells. It is contemplated that any cell from which a fertile plant may be regenerated is useful as a recipient cell. Callus may be initiated from tissue sources including, but not limited to, immature embryos, seedling apical meristems, microspores and the like. Cells capable of proliferating as callus are also recipient cells for 5 genetic transformation. Practical transformation methods and materials for making transgenic plants of this disclosure, for example various media and recipient target cells, transformation of immature embryo cells and subsequent regeneration of fertile transgenic plants are disclosed in U.S. Pat. Nos. 6,194,636 and 6,232,526.

[00211] The seeds of transgenic plants can be harvested from fertile transgenic plants 10 and used to grow progeny generations of transformed plants that produce the recombinant nutritive protein of this disclosure. In addition to direct transformation of a plant with a recombinant DNA, transgenic plants can be prepared by crossing a first plant having a recombinant DNA with a second plant lacking the DNA. For example, recombinant DNA can be introduced into first plant line that is amenable to transformation to produce a transgenic 15 plant which can be crossed with a second plant line to introgress the recombinant DNA into the second plant line. A transgenic plant with recombinant DNA encoding a nutritive protein of this disclosure, can be crossed with transgenic plant line having other recombinant DNA that confers another trait, for example herbicide resistance or pest resistance, or production of a second nutritive product such as an oil, to produce progeny plants having recombinant 20 DNA that confers both traits. Typically, in such breeding for combining traits the transgenic plant donating the additional trait is a male line and the transgenic plant carrying the base traits is the female line. The progeny of this cross will segregate such that some of the plants will carry the DNA for both parental traits and some will carry DNA for one parental trait; such plants can be identified by markers associated with parental recombinant DNA, e.g. 25 marker identification by analysis for recombinant DNA or, in the case where a selectable marker is linked to the recombinant, by application of the selecting agent such as a herbicide for use with a herbicide tolerance marker, or by selection for the enhanced trait. Progeny plants carrying DNA for both parental traits can be crossed back into the female parent line multiple times, for example usually 6 to 8 generations, to produce a progeny plant with 30 substantially the same genotype as one original transgenic parental line but for the recombinant DNA of the other transgenic parental line.

- [00212] In the practice of transformation DNA is typically introduced into only a small percentage of target plant cells in any one transformation experiment. Marker genes are used to provide an efficient system for identification of those cells that are stably transformed by receiving and integrating a transgenic DNA construct into their genomes. Preferred marker genes provide selective markers which confer resistance to a selective agent, such as an antibiotic or herbicide. Any of the herbicides to which the transformed plants may be resistant are useful agents for selective markers. Potentially transformed cells are exposed to the selective agent. In the population of surviving cells will be those cells where, generally, the resistance-conferring gene is integrated and expressed at sufficient levels to permit cell survival. Cells may be tested further to confirm stable integration of the exogenous DNA. Commonly used selective marker genes include those conferring resistance to antibiotics such as kanamycin and paromomycin (nptII), hygromycin B (aph IV) and gentamycin (aac3 and aacC4) or resistance to herbicides such as glufosinate (bar or pat) and glyphosate (aroA or EPSPS). Examples of such selectable are illustrated in U.S. Pat. Nos. 5,550,318; 5,633,435; 5,780,708 and 6,118,047. Selectable markers which provide an ability to visually identify transformants can also be employed, for example, a gene expressing a colored or fluorescent protein such as a luciferase or green fluorescent protein (GFP) or a gene expressing a beta-glucuronidase or uidA gene (GUS) for which various chromogenic substrates are known.
- [00213] Plant cells that survive exposure to the selective agent, or plant cells that have been scored positive in a screening assay, may be cultured in regeneration media and allowed to mature into plants. Developing plantlets regenerated from transformed plant cells can be transferred to plant growth mix, and hardened off, for example, in an environmentally controlled chamber. Plants are regenerated from about 6 weeks to 10 months after a transformant is identified, depending on the initial tissue. Plants may be pollinated using conventional plant breeding methods known to those of skill in the art and seed produced, for example self-pollination is commonly used with transgenic corn. The regenerated transformed plant or its progeny seed or plants can be tested for expression of the recombinant DNA and selected for the presence of a heterologous nutritive protein.
- [00214] Transgenic plants derived from the plant cells of this disclosure are grown to generate transgenic plants comprising the heterologous nucleic acid that encodes a nutritive protein of this disclosure and produce transgenic seed and haploid pollen comprising the

heterologous nucleic acid sequence. Such plants with enhanced traits are identified by selection of transformed plants or progeny seed for the enhanced trait. Transgenic plants grown from transgenic seed provided herein demonstrate improved agronomic traits that contribute to increased yield or other traits that provide increased plant value, including, for 5 example, improved protein quality such as increasing the content of at least one of essential amino acids, branch chain amino acids, or Leu.

[00215] The transgenic plants are useful as sources of nutritive proteins. For example, in some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of total protein compared to a control non- 10 transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of essential amino acids compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of branch chain amino acids compared to a control non-transgenic plant. In some 15 embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of Leu compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises at least one of: a) an increased ratio of branch chain amino acid residues to total amino acid residues compared to a control non-transgenic plant; b) an 20 increased ratio of Leu residues to total amino acid residues compared to a control non-transgenic plant; and c) an increased ratio of essential amino acid residues to total amino acid residues compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises: a) an increased ratio of branch chain amino acid residues to total amino acid residues compared to a control 25 non-transgenic plant; b) an increased ratio of Leu residues to total amino acid residues compared to a control non-transgenic plant; and c) an increased ratio of essential amino acid residues to total amino acid residues compared to a control non-transgenic plant.

[00216] Accordingly, the transgenic plants are useful as sources of high quality protein. The plants may be harvested and used in mammalian diets with or without further 30 processing. For example, flour made from transgenic wheat, cornmeal made from transgenic corn, or rice or rice flour derived from transgenic rice is enriched in at least one of protein, essential amino acids, branch chain amino acids, and Leu compared to similar products made

from plants that do not comprise the recombinant nutritive protein. In some embodiments the recombinant nutritive protein is a plant protein or comprises a polypeptide sequence of a plant protein or a derivative or mutein thereof, such as but not necessarily a protein or polypeptide sequence of the same type of plant. In other embodiments the recombinant nutritive protein is not a plant protein or a derivative or mutein thereof.

[00217] In some embodiments the recombinant nutritive protein is recovered or partially recovered from the transgenic plant before it is consumed by a mammal.

## H. Compositions

[00218] At least one nutritive protein disclosed herein may be combined with at least one second component to form a nutritive composition. In some embodiments the only source of amino acid in the composition is the at least one nutritive protein disclosed herein. In such embodiments the amino acid composition of the composition will be the same as the amino acid composition of the at least one nutritive protein disclosed herein. In some embodiments the composition comprises at least one nutritive protein disclosed herein and at least one second protein. In some embodiments the at least one second protein is a second nutritive protein disclosed herein, while in other embodiments the at least one second protein is not a nutritive protein disclosed herein. In some embodiments the composition comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nutritive proteins disclosed herein. In some embodiments the composition comprises 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more proteins that are not nutritive proteins disclosed herein. In some embodiments the composition comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nutritive proteins and the composition comprises 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more proteins that are not nutritive proteins disclosed herein.

[00219] In some embodiments the nutritive composition as described in the preceding paragraph, further comprises at least one of at least one polypeptide, at least one peptide, and at least one free amino acid. In some embodiments the nutritive composition comprises at least one polypeptide and at least one peptide. In some embodiments the nutritive composition comprises at least one polypeptide and at least one free amino acid. In some embodiments the nutritive composition comprises at least one peptide and at least one free amino acid. In some embodiments the at least one polypeptide, at least one peptide, and/or at least one free amino acid comprises amino acids selected from 1) branch chain amino acids,

2) leucine, and 3) essential amino acids. In some embodiments the at least one polypeptide, at least one peptide, and/or at least one free amino acid consists of amino acids selected from 1) branch chain amino acids, 2) leucine, and 3) essential amino acids. In some embodiments, the nutritive composition comprises at least one modified amino acid or a non-standard 5 amino acid. Modified amino acids include amino acids that have modifications to one or more of the carboxy terminus, amino terminus, and/or side chain. Non-standard amino acids may be selected from those that are formed by post-translational modification of proteins, for example, carboxylated glutamate, hydroxyproline, or hypusine. Other non-standard amino acids are not found in proteins. Examples include lanthionine, 2-aminoisobutyric 10 acid, dehydroalanine, gamma-aminobutyric acid, ornithine and citrulline. In some embodiments, the nutritive composition comprises one or more D-amino acids. In some embodiments, the nutritive composition comprises one or more L-amino acids. In some embodiments, the nutritive composition comprises a mixture of one or more D-amino acids and one or more L-amino acids.

15 [00220] By adding at least one of a polypeptide, a peptide, and a free amino acid to a nutritive composition the proportion of at least one of branch chain amino acids, leucine, and essential amino acids, to total amino acid, present in the composition can be increased.

[00221] In some embodiments the composition comprises at least one carbohydrate. A “carbohydrate” refers to a sugar or polymer of sugars. The terms “saccharide,” 20 “polysaccharide,” “carbohydrate,” and “oligosaccharide” may be used interchangeably. Most carbohydrates are aldehydes or ketones with many hydroxyl groups, usually one on each carbon atom of the molecule. Carbohydrates generally have the molecular formula  $C_nH_{2n}O_n$ . A carbohydrate may be a monosaccharide, a disaccharide, trisaccharide, oligosaccharide, or polysaccharide. The most basic carbohydrate is a monosaccharide, such 25 as glucose, sucrose, galactose, mannose, ribose, arabinose, xylose, and fructose. Disaccharides are two joined monosaccharides. Exemplary disaccharides include sucrose, maltose, cellobiose, and lactose. Typically, an oligosaccharide includes between three and six monosaccharide units (e.g., raffinose, stachyose), and polysaccharides include six or more monosaccharide units. Exemplary polysaccharides include starch, glycogen, and cellulose. 30 Carbohydrates may contain modified saccharide units such as 2'-deoxyribose wherein a hydroxyl group is removed, 2'-fluororibose wherein a hydroxyl group is replaced with a fluorine, or N-acetylglucosamine, a nitrogen-containing form of glucose (e.g., 2'-

fluororibose, deoxyribose, and hexose). Carbohydrates may exist in many different forms, for example, conformers, cyclic forms, acyclic forms, stereoisomers, tautomers, anomers, and isomers.

[00222] In some embodiments the composition comprises at least one lipid. As used herein a “lipid” includes fats, oils, triglycerides, cholesterol, phospholipids, fatty acids in any form including free fatty acids. Fats, oils and fatty acids may be saturated, unsaturated (cis or trans) or partially unsaturated (cis or trans). In some embodiments the lipid comprises at least one fatty acid selected from lauric acid (12:0), myristic acid (14:0), palmitic acid (16:0), palmitoleic acid (16:1), margaric acid (17:0), heptadecenoic acid (17:1), stearic acid (18:0), 10 oleic acid (18:1), linoleic acid (18:2), linolenic acid (18:3), octadecatetraenoic acid (18:4), arachidic acid (20:0), eicosenoic acid (20:1), eicosadienoic acid (20:2), eicosatetraenoic acid (20:4), eicosapentaenoic acid (20:5) (EPA), docosanoic acid (22:0), docosenoic acid (22:1), docosapentaenoic acid (22:5), docosahexaenoic acid (22:6) (DHA), and tetracosanoic acid (24:0). In some embodiments the composition comprises at least one modified lipid, for 15 example a lipid that has been modified by cooking.

[00223] In some embodiments the composition comprises at least one supplemental mineral or mineral source. Examples of minerals include, without limitation: chloride, sodium, calcium, iron, chromium, copper, iodine, zinc, magnesium, manganese, molybdenum, phosphorus, potassium, and selenium. Suitable forms of any of the foregoing 20 minerals include soluble mineral salts, slightly soluble mineral salts, insoluble mineral salts, chelated minerals, mineral complexes, non-reactive minerals such as carbonyl minerals, and reduced minerals, and combinations thereof.

[00224] In some embodiments the composition comprises at least one supplemental vitamin. The at least one vitamin can be fat-soluble or water soluble vitamins. Suitable 25 vitamins include but are not limited to vitamin C, vitamin A, vitamin E, vitamin B12, vitamin K, riboflavin, niacin, vitamin D, vitamin B6, folic acid, pyridoxine, thiamine, pantothenic acid, and biotin. Suitable forms of any of the foregoing are salts of the vitamin, derivatives of the vitamin, compounds having the same or similar activity of the vitamin, and metabolites of the vitamin.

[00225] In some embodiments the composition comprises at least one organism.

Suitable examples are well known in the art and include probiotics (e.g., species of *Lactobacillus* or *Bifidobacterium*), spirulina, chlorella, and porphyra.

[00226] In some embodiments the composition comprises at least one dietary supplement. Suitable examples are well known in the art and include herbs, botanicals, and certain hormones. Non limiting examples include ginko, gensing, and melatonin.

[00227] In some embodiments the composition comprises an excipient. Non-limiting examples of suitable excipients include a buffering agent, a preservative, a stabilizer, a binder, a compaction agent, a lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a coloring agent.

[00228] In some embodiments the excipient is a buffering agent. Non-limiting examples of suitable buffering agents include sodium citrate, magnesium carbonate, magnesium bicarbonate, calcium carbonate, and calcium bicarbonate.

[00229] In some embodiments the excipient comprises a preservative. Non-limiting examples of suitable preservatives include antioxidants, such as alpha-tocopherol and ascorbate, and antimicrobials, such as parabens, chlorobutanol, and phenol.

[00230] In some embodiments the composition comprises a binder as an excipient. Non-limiting examples of suitable binders include starches, pregelatinized starches, gelatin, polyvinylpyrrolidone, cellulose, methylcellulose, sodium carboxymethylcellulose, ethylcellulose, polyacrylamides, polyvinyloxoazolidone, polyvinylalcohols, C12-C18 fatty acid alcohol, polyethylene glycol, polyols, saccharides, oligosaccharides, and combinations thereof.

[00231] In some embodiments the composition comprises a lubricant as an excipient. Non-limiting examples of suitable lubricants include magnesium stearate, calcium stearate, zinc stearate, hydrogenated vegetable oils, sterotex, polyoxyethylene monostearate, talc, polyethyleneglycol, sodium benzoate, sodium lauryl sulfate, magnesium lauryl sulfate, and light mineral oil.

[00232] In some embodiments the composition comprises a dispersion enhancer as an excipient. Non-limiting examples of suitable dispersants include starch, alginic acid,

polyvinylpyrrolidones, guar gum, kaolin, bentonite, purified wood cellulose, sodium starch glycolate, isoamorphous silicate, and microcrystalline cellulose as high HLB emulsifier surfactants.

[00233] In some embodiments the composition comprises a disintegrant as an excipient. In some embodiments the disintegrant is a non-effervescent disintegrant. Non-limiting examples of suitable non-effervescent disintegrants include starches such as corn starch, potato starch, pregelatinized and modified starches thereof, sweeteners, clays, such as bentonite, micro-crystalline cellulose, alginates, sodium starch glycolate, gums such as agar, guar, locust bean, karaya, pectin, and tragacanth. In some embodiments the disintegrant is an effervescent disintegrant. Non-limiting examples of suitable effervescent disintegrants include sodium bicarbonate in combination with citric acid, and sodium bicarbonate in combination with tartaric acid.

[00234] In some embodiments the excipient comprises a flavoring agent. Flavoring agents incorporated into the outer layer can be chosen from synthetic flavor oils and flavoring aromatics; natural oils; extracts from plants, leaves, flowers, and fruits; and combinations thereof. In some embodiments the flavoring agent is selected from cinnamon oils; oil of wintergreen; peppermint oils; clover oil; hay oil; anise oil; eucalyptus; vanilla; citrus oil such as lemon oil, orange oil, grape and grapefruit oil; and fruit essences including apple, peach, pear, strawberry, raspberry, cherry, plum, pineapple, and apricot.

[00235] In some embodiments the excipient comprises a sweetener. Non-limiting examples of suitable sweeteners include glucose (corn syrup), dextrose, invert sugar, fructose, and mixtures thereof (when not used as a carrier); saccharin and its various salts such as the sodium salt; dipeptide sweeteners such as aspartame; dihydrochalcone compounds, glycyrrhizin; Stevia Rebaudiana (Stevioside); chloro derivatives of sucrose such as sucralose; and sugar alcohols such as sorbitol, mannitol, sylitol, and the like. Also contemplated are hydrogenated starch hydrolysates and the synthetic sweetener 3,6-dihydro-6-methyl-1,2,3-oxathiazin-4-one-2,2-dioxide, particularly the potassium salt (acesulfame-K), and sodium and calcium salts thereof.

[00236] In some embodiments the composition comprises a coloring agent. Non-limiting examples of suitable color agents include food, drug and cosmetic colors (FD&C),

drug and cosmetic colors (D&C), and external drug and cosmetic colors (Ext. D&C). The coloring agents can be used as dyes or their corresponding lakes.

[00237] The weight fraction of the excipient or combination of excipients in the formulation is usually about 50% or less, about 45% or less, about 40% or less, about 35% or less, about 30% or less, about 25% or less, about 20% or less, about 15% or less, about 10% or less, about 5% or less, about 2% or less, or about 1% or less of the total weight of the amino acids in the composition.

[00238] The nutritive proteins and nutritive compositions disclosed herein can be formulated into a variety of forms and administered by a number of different means. The compositions can be administered orally, rectally, or parenterally, in formulations containing conventionally acceptable carriers, adjuvants, and vehicles as desired. The term “parenteral” as used herein includes subcutaneous, intravenous, intramuscular, or intrasternal injection and infusion techniques. In an exemplary embodiment, the nutritive protein or composition is administered orally.

[00239] Solid dosage forms for oral administration include capsules, tablets, caplets, pills, troches, lozenges, powders, and granules. A capsule typically comprises a core material comprising a nutritive protein or composition and a shell wall that encapsulates the core material. In some embodiments the core material comprises at least one of a solid, a liquid, and an emulsion. In some embodiments the shell wall material comprises at least one of a soft gelatin, a hard gelatin, and a polymer. Suitable polymers include, but are not limited to: cellulosic polymers such as hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose (HPMC), methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropylmethyl cellulose phthalate, hydroxypropylmethyl cellulose succinate and carboxymethylcellulose sodium; acrylic acid polymers and copolymers, such as those formed from acrylic acid, methacrylic acid, methyl acrylate, ammonio methylacrylate, ethyl acrylate, methyl methacrylate and/or ethyl methacrylate (e.g., those copolymers sold under the trade name “Eudragit”); vinyl polymers and copolymers such as polyvinyl pyrrolidone, polyvinyl acetate, polyvinylacetate phthalate, vinylacetate crotonic acid copolymer, and ethylene-vinyl acetate copolymers; and shellac (purified lac). In some embodiments at least one polymer functions as taste-masking agents.

[00240] Tablets, pills, and the like can be compressed, multiply compressed, multiply layered, and/or coated. The coating can be single or multiple. In one embodiment, the coating material comprises at least one of a saccharide, a polysaccharide, and glycoproteins extracted from at least one of a plant, a fungus, and a microbe. Non-limiting examples include corn 5 starch, wheat starch, potato starch, tapioca starch, cellulose, hemicellulose, dextrans, maltodextrin, cyclodextrins, inulins, pectin, mannans, gum arabic, locust bean gum, mesquite gum, guar gum, gum karaya, gum ghatti, tragacanth gum, funori, carrageenans, agar, alginates, chitosans, or gellan gum. In some embodiments the coating material comprises a protein. In some embodiments the coating material comprises at least one of a fat and an oil.

10 In some embodiments the at least one of a fat and an oil is high temperature melting. In some embodiments the at least one of a fat and an oil is hydrogenated or partially hydrogenated. In some embodiments the at least one of a fat and an oil is derived from a plant. In some embodiments the at least one of a fat and an oil comprises at least one of glycerides, free fatty acids, and fatty acid esters. In some embodiments the coating material comprises at least one 15 edible wax. The edible wax can be derived from animals, insects, or plants. Non-limiting examples include beeswax, lanolin, bayberry wax, carnauba wax, and rice bran wax. Tablets and pills can additionally be prepared with enteric coatings.

[00241] Alternatively, powders or granules embodying the nutritive proteins and nutritive compositions disclosed herein can be incorporated into a food product. In some 20 embodiments the food product is be a drink for oral administration. Non-limiting examples of a suitable drink include fruit juice, a fruit drink, an artificially flavored drink, an artificially sweetened drink, a carbonated beverage, a sports drink, a liquid dairy product, a shake, an alcoholic beverage, a caffeinated beverage, infant formula and so forth. Other suitable means for oral administration include aqueous and nonaqueous solutions, creams, pastes, emulsions, 25 suspensions and slurries, , each of which may optionally also containin at least one of suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, coloring agents, and flavoring agents.

[00242] In some embodiments the food product is a solid foodstuff. Suitable examples of a solid foodstuff include without limitation a food bar, a snack bar, a cookie, a brownie, a 30 muffin, a cracker, a biscuit, a cream or paste, an ice cream bar, a frozen yogurt bar, and the like.

[00243] In some embodiments, the nutritive proteins and nutritive compositions disclosed herein are incorporated into a therapeutic food. In some embodiments, the therapeutic food is a ready-to-use food that optionally contains some or all essential macronutrients and micronutrients. In some embodiments, the nutritive proteins and nutritive compositions disclosed herein are incorporated into a supplementary food that is designed to be blended into an existing meal. In some embodiments, the supplemental food contains some or all essential macronutrients and micronutrients. In some embodiments, the nutritive proteins and nutritive compositions disclosed herein are blended with or added to an existing food to fortify the food's protein nutrition. Examples include food staples (grain, salt, sugar, cooking oil, margarine), beverages (coffee, tea, soda, beer, liquor, sports drinks), snacks, sweets and other foods.

[00244] The compositions disclosed herein can be utilized in methods to increase at least one of muscle mass, strength and physical function, thermogenesis, metabolic expenditure, satiety, mitochondrial biogenesis, weight or fat loss, and lean body composition for example.

## I. Methods of Use

[00245] In some embodiments the nutritive proteins and nutritive compositions disclosed herein are administered to a patient or a user (sometimes collectively referred to as a "subject"). As used herein "administer" and "administration" encompasses embodiments in which one person directs another to consume a nutritive protein or nutritive composition in a certain manner and/or for a certain purpose, and also situations in which a user uses a nutritive protein or nutritive composition in a certain manner and/or for a certain purpose independently of or in variance to any instructions received from a second person. Non-limiting examples of embodiments in which one person directs another to consume a nutritive protein or nutritive composition in a certain manner and/or for a certain purpose include when a physician prescribes a course of conduct and/or treatment to a patient, when a trainer advises a user (such as an athlete) to follow a particular course of conduct and/or treatment, and when a manufacturer, distributor, or marketer recommends conditions of use to an end user, for example through advertisements or labeling on packaging or on other materials provided in association with the sale or marketing of a product.

[00246] In some embodiments the nutritive proteins or nutritive compositions are provided in a dosage form. In some embodiments the dosage form is designed for

administration of at least one nutritive protein disclosed herein, wherein the total amount of nutritive protein administered is selected from 0.1g to 1g, 1g to 5g, from 2g to 10g, from 5g to 15g, from 10g to 20g, from 15g to 25g, from 20g to 40g, from 25-50g, and from 30-60g. In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of nutritive protein administered is selected from about 0.1g, 0.1g-1g, 1g, 2g, 3g, 4g, 5g, 6g, 7g, 8g, 9g, 10g, 15g, 20g, 25g, 30g, 35g, 40g, 45g, 50g, 55g, 60g, 65g, 70g, 75g, 80g, 85g, 90g, 95g, and 100g.

[00247] In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of essential amino acids administered is selected from 0.1g to 1g, from 1g to 5g, from 2g to 10g, from 5g to 15g, from 10g to 20g, and from 1-30 g. In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of nutritive protein administered is selected from about 0.1g, 0.1-1g, 1g, 2g, 3g, 4g, 5g, 6g, 7g, 8g, 9g, 10g, 15g, 20g, 25g, 30g, 35g, 40g, 45g, 50g, 55g, 60g, 65g, 70g, 75g, 80g, 85g, 90g, 95g, and 100g.

[00248] In some embodiments the nutritive protein or nutritive composition is consumed at a rate of from 0.1g to 1g a day, 1g to 5 g a day, from 2g to 10g a day, from 5g to 15g a day, from 10g to 20g a day, from 15g to 30g a day, from 20g to 40g a day, from 25g to 50g a day, from 40g to 80g a day, from 50g to 100g a day, or more.

[00249] In some embodiments, of the total protein intake by the subject, at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or about 100% of the total protein intake by the subject over a dietary period is made up of at least one nutritive protein according to this disclosure. In some embodiments, of the total protein intake by the subject, from 5% to 100% of the total protein intake by the subject, from 5% to 90% of the total protein intake by the subject, from 5% to 80% of the total protein intake by the subject, from 5% to 70% of the total protein intake by the subject, from 5% to 60% of the total protein intake by the subject, from 5% to 50% of the total protein intake by the subject, from 5% to 40% of the total protein intake by the subject, from 5% to 30% of the total protein intake by the subject, from 5% to 20% of the total protein intake by the subject, from 5% to 10% of the total protein intake by the subject, from 10% to 100% of the total protein intake by the subject, from 10%

to 100% of the total protein intake by the subject, from 20% to 100% of the total protein intake by the subject, from 30% to 100% of the total protein intake by the subject, from 40% to 100% of the total protein intake by the subject, from 50% to 100% of the total protein intake by the subject, from 60% to 100% of the total protein intake by the subject, from 70% to 100% of the total protein intake by the subject, from 80% to 100% of the total protein intake by the subject, or from 90% to 100% of the total protein intake by the subject, over a dietary period, is made up of at least one nutritive protein according to this disclosure. In some embodiments the at least one nutritive protein of this disclosure accounts for at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, or at least 50% of the subject's calorie intake over a dietary period.

[00250] In some embodiments the at least one nutritive protein according to this disclosure comprises at least 2 nutritive proteins of this disclosure, at least 3 nutritive proteins of this disclosure, at least 4 nutritive proteins of this disclosure, at least 5 nutritive proteins of this disclosure, at least 6 nutritive proteins of this disclosure, at least 7 nutritive proteins of this disclosure, at least 8 nutritive proteins of this disclosure, at least 9 nutritive proteins of this disclosure, at least 10 nutritive proteins of this disclosure, or more.

[00251] In some embodiments the dietary period is 1 meal, 2 meals, 3 meals, at least 1 day, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 2 weeks, at least 3 weeks, at least 4 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, or at least 1 year. In some embodiments the dietary period is from 1 day to 1 week, from 1 week to 4 weeks, from 1 month, to 3 months, from 3 months to 6 months, or from 6 months to 1 year.

[00252] Clinical studies provide evidence that protein prevents muscle loss due to aging or bed rest. In particular, studies have shown that protein supplementation increases muscle fractional synthetic rate (FSR) during prolonged bed rest, maintains leg mass and strength during prolonged bed rest, increases lean body mass, improves functional measures of gait and balance, and may serve as a viable intervention for individuals at risk of sarcopenia due to immobility or prolonged bed rest. (See, e.g., Paddon-Jones D, et al. J Clin Endocrinol Metab 2004, 89:4351–4358; Ferrando, A et al. Clinical Nutrition 2009 1-6; Katsanos C et al. Am J Physiol Endocrinol Metab. 2006, 291: 381-387).

[00253] Studies on increasing muscle protein anabolism in athletes have shown that protein provided following exercise promotes muscle hypertrophy to a greater extent than that achieved by exercise alone. It has also been shown that protein provided following exercise supports protein synthesis without any increase in protein breakdown, resulting in a 5 net positive protein balance and muscle mass accretion. While muscle protein synthesis appears to respond in a dose-response fashion to essential amino acid supplementation, not all proteins are equal in building muscle. For example, the amino acid leucine is an important factor in stimulating muscle protein synthesis. (See, e.g., Borsheim E et al. Am J Physiol Endocrinol Metab 2002, 283: E648-E657; Borsheim E et al. Clin Nutr. 2008, 27: 10 189-95; Esmarck B et al J Physiol 2001, 535: 301-311; Moore D et al. Am J Clin Nutr 2009, 89: 161-8).

[00254] In another aspect this disclosure provides methods of maintaining or increasing at least one of muscle mass, muscle strength, and functional performance in a subject. In some embodiments the methods comprise providing to the subject a sufficient 15 amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a 20 method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00255] In another aspect this disclosure provides methods of maintaining or achieving 25 a desirable body mass index in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the sufficient 30 amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this

disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00256] In another aspect this disclosure provides methods of providing protein to a subject with protein-energy malnutrition. In some embodiments the methods comprise 5 providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

10 [00257] The need for essential amino acid supplementation has been suggested in cancer patients and other patients suffering from cachexia. Dietary studies in mice have shown survival and functional benefits to cachectic cancer-bearing mice through dietary intervention with essential amino acids. Beyond cancer, essential amino acid supplementation has also shown benefits, such as improved muscle function and muscle gain, 15 in patients suffering from other diseases who have difficulty exercising and therefore suffer from muscular deterioration, such as chronic obstructive pulmonary disease, chronic heart failure, HIV, and other disease states.

20 [00258] Studies have shown that specific amino acids have advantages in managing cachexia. A relatively high content of BCAAs and Leu in diets are thought to have a positive effect in cachexia by promoting total protein synthesis by signaling an increase in translation, enhancing insulin release, and inhibiting protein degradation. Thus, consuming increased dietary BCAAs in general and/or Leu in particular will contribute positively to reduce or reverse the effects of cachexia. Because nitrogen balance is important in countering the underlying cause of cachexia it is thought that consuming increased dietary glutamine and/or 25 arginine will contribute positively to reduce or reverse the effects of cachexia. (See, e.g., Op den Kamp C, Langen R, Haegens A, Schols A. "Muscle atrophy in cachexia: can dietary protein tip the balance?" Current Opinion in Clinical Nutrition and Metabolic Care 2009, 12:611-616; Poon RT-P, Yu W-C, Fan S-T, et al. "Long-term oral branched chain amino acids in patients undergoing chemoembolization for hepatocellular carcinoma:a randomized 30 trial." Aliment Pharmacol Ther 2004; 19:779-788; Tayek JA, Bistrian BR, Hehir DJ, Martin R, Moldawer LL, Blackburn GL. "Improved protein kinetics and albumin synthesis by branched chain amino acid-enriched total parenteral nutrition in cancer cachexia." Cancer.

1986;58:147-57; Xi P, Jiang Z, Zheng C, Lin Y, Wu G "Regulation of protein metabolism by glutamine: implications for nutrition and health." *Front Biosci.* 2011 Jan 1;16:578-97).

[00259] Accordingly, also provided herein are methods of treating cachexia in a subject. In some embodiments a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure for a subject with cachexia is an amount such that the amount of protein of this disclosure ingested by the person meets or exceeds the metabolic needs (which are often elevated). A protein intake of 1.5 g/kg of body weight per day or 15-20% of total caloric intake appears to be an appropriate target for persons with cachexia. In some embodiments all of the protein consumed by the subject is a nutritive protein according to this disclosure. In some embodiments nutritive protein according to this disclosure is combined with other sources of protein and/or free amino acids to provide the total protein intake of the subject. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the subject suffers from a disease that makes exercise difficult and therefore causes muscular deterioration, such as chronic obstructive pulmonary disease, chronic heart failure, HIV, cancer, and other disease states. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to this disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00260] Sarcopenia is the degenerative loss of skeletal muscle mass (typically 0.5-1% loss per year after the age of 25), quality, and strength associated with aging. Sarcopenia is a component of the frailty syndrome. The European Working Group on Sarcopenia in Older People (EWGSOP) has developed a practical clinical definition and consensus diagnostic criteria for age-related sarcopenia. For the diagnosis of sarcopenia, the working group has proposed using the presence of both low muscle mass and low muscle function (strength or performance). Sarcopenia is characterized first by a muscle atrophy (a decrease in the size of the muscle), along with a reduction in muscle tissue "quality," caused by such factors as replacement of muscle fibres with fat, an increase in fibrosis, changes in muscle metabolism, oxidative stress, and degeneration of the neuromuscular junction. Combined, these changes

lead to progressive loss of muscle function and eventually to frailty. Frailty is a common geriatric syndrome that embodies an elevated risk of catastrophic declines in health and function among older adults. Contributors to frailty can include sarcopenia, osteoporosis, and muscle weakness. Muscle weakness, also known as muscle fatigue, (or “lack of strength”) 5 refers to the inability to exert force with one’s skeletal muscles. Weakness often follows muscle atrophy and a decrease in activity, such as after a long bout of bedrest as a result of an illness. There is also a gradual onset of muscle weakness as a result of sarcopenia.

[00261] The nutritive proteins of this disclosure are useful for treating sarcopenia or frailty once it develops in a subject or for preventing the onset of sarcopenia or frailty in a 10 subject who is a member of an at risk groups. In some embodiments all of the protein consumed by the subject is a nutritive protein according to this disclosure. In some embodiments nutritive protein according to this disclosure is combined with other sources of protein and/or free amino acids to provide the total protein intake of the subject. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from 15 protein-energy malnutrition. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to this disclosure, the nutritive composition according to disclosure, or the nutritive composition 20 made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00262] Obesity is a multifactorial disorder associated with a host of comorbidities including hypertension, type 2 diabetes, dyslipidemia, coronary heart disease, stroke, cancer (eg, endometrial, breast, and colon), osteoarthritis, sleep apnea, and respiratory problems. 25 The incidence of obesity, defined as a body mass index  $>30$  kg/m<sup>2</sup>, has increased dramatically in the United States, from 15% (1976–1980) to 33% (2003–2004), and it continues to grow. Although the mechanisms contributing to obesity are complex and involve the interplay of behavioral components with hormonal, genetic, and metabolic processes, obesity is largely viewed as a lifestyle-dependent condition with 2 primary causes: excessive 30 energy intake and insufficient physical activity. With respect to energy intake, there is evidence that modestly increasing the proportion of protein in the diet, while controlling total energy intake, may improve body composition, facilitate fat loss, and improve body weight

maintenance after weight loss. Positive outcomes associated with increased dietary protein are thought to be due primarily to lower energy intake associated with increased satiety, reduced energy efficiency and/or increased thermogenesis, positive effects on body composition (specifically lean muscle mass), and enhanced glycemic control.

- 5 [00263] Dietary proteins are more effective in increasing post-prandial energy expenditure than isocaloric intakes of carbohydrates or fat (see, e.g., Dauncey M, Bingham S. "Dependence of 24 h energy expenditure in man on composition of the nutrient intake." Br J Nutr 1983, 50: 1-13; Karst H et al. "Diet-induced thermogenesis in man: thermic effects of single proteins, carbohydrates and fats depending on their energy amount." Ann Nutr 10 Metab. 1984, 28: 245-52; Tappy L et al "Thermic effect of infused amino acids in healthy humans and in subjects with insulin resistance." Am J Clin Nutr 1993, 57 (6): 912-6). This property along with other properties (satiety induction; preservation of lean body mass) make protein an attractive component of diets directed at weight management. The increase in energy expenditure caused by such diets may in part be due to the fact that the energy cost of 15 digesting and metabolizing protein is higher than for other calorie sources. Protein turnover, including protein synthesis, is an energy consuming process. In addition, high protein diets may also up-regulate uncoupling protein in liver and brown adipose, which is positively correlated with increases in energy expenditure. It has been theorized that different proteins may have unique effects on energy expenditure.
- 20 [00264] Studies suggest that ingestion of protein, particularly proteins with high EAA and/or BCAA content, leads to distinct effects on thermogenesis and energy expenditure (see, e.g., Mikkelsen P. et al. "Effect of fat-reduced diets on 24 h energy expenditure: comparisons between animal protein, vegetable protein and carbohydrate." Am J Clin Nutr 2000, 72:1135-41; Acheson K. et al. "Protein choices targeting thermogenesis and metabolism." Am J Clin 25 Nutr 2011, 93:525-34; Alfenas R. et al. "Effects of protein quality on appetite and energy metabolism in normal weight subjects" Arg Bras Endocrinol Metabol 2010, 54 (1): 45-51; Lorenzen J. et al. "The effect of milk proteins on appetite regulation and diet-induced thermogenesis." J Clin Nutr 2012 66 (5): 622-7). Additionally, L-tyrosine has been identified as an amino acid that plays a role in thermogenesis (see, e.g., Belza A. et al. "The beta- 30 adrenergic antagonist propranolol partly abolishes thermogenic response to bioactive food ingredients." Metabolism 2009, 58 (8):1137-44). Further studies suggest that Leucine and Arginine supplementation appear to alter energy metabolism by directing substrate to lean

body mass rather than adipose tissue (Dulloo A. "The search for compounds that stimulate thermogenesis in obesity management: from pharmaceuticals to functional food ingredients." *Obes Rev* 2011 12: 866-83).

5 [00265] Collectively the literature suggests that different protein types leads to distinct effects on thermogenesis. Because proteins or peptides rich in EAAs, BCAA, and/or at least one of Tyr, Arg, and Leu are believed to have a stimulatory effect on thermogenesis, and because stimulation of thermogenesis is believed to lead to positive effects on weight management, this disclosure also provides products and methods useful to stimulation thermogenesis and/or to bring about positive effects on weight management in general.

10 [00266] More particularly, this disclosure provides methods of increasing thermogenesis in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

15

20 [00267] At the basic level, the reason for the development of an overweight condition is due to an imbalance between energy intake and energy expenditure. Attempts to reduce food at any particular occasion (satiation) and across eating occasions (satiety) have been a major focus of recent research. Reduced caloric intake as a consequence of feeling satisfied during a meal and feeling full after a meal results from a complex interaction of internal and external signals. Various nutritional studies have demonstrated that variation in food properties such as energy density, content, texture and taste influence both satiation and satiety.

25 [00268] There are three macronutrients that deliver energy: fat, carbohydrates and proteins. A gram of protein or carbohydrate provides 4 calories while a gram of fat 9 calories. Protein generally increases satiety to a greater extent than carbohydrates or fat and

therefore may facilitate a reduction in calorie intake. However, there is considerable evidence that indicates the type of protein matters in inducing satiety (see, e.g., W.L. Hall, et al. "Casein and whey exert different effects on plasma amino acid profiles, gastrointestinal hormone secretion and appetite." *Br J Nutr.* 2003 Feb, 89(2):239-48; R. Abou-Samra, et al.

- 5 "Effect of different protein sources on satiation and short-term satiety when consumed as a starter." *Nutr J.* 2011 Dec 23, 10:139; T. Akhavan, et al. "Effect of premeal consumption of whey protein and its hydrolysate on food intake and postmeal glycemia and insulin responses in young adults." *Am J Clin Nutr.* 2010 Apr, 91(4):966-75, Epub 2010 Feb 17; MA Veldhorst "Dose-dependent satiating effect of whey relative to casein or soy" *Physiol Behav.*
- 10 2009 Mar 23, 96(4-5):675-82). Evidence indicates that protein rich in Leucine is particularly effective at inducing satiety (see, e.g., Fromentin G et al "Peripheral and central mechanisms involved in the control of food intake by dietary amino acids and proteins." *Nutr Res Rev* 2012 25: 29-39).

[00269] In some embodiments a nutritive protein of this disclosure is consumed by a 15 subject concurrently with at least one pharmaceutical or biologic drug product. In some embodiments the beneficial effects of the nutritive protein and the at least one pharmaceutical or biologic drug product have an additive effect while in some embodiments the beneficial effects of the nutritive protein and the at least one pharmaceutical or biologic drug product have a synergistic effect. Examples of pharmaceutical or biologic drug products that can be 20 administered with the nutritive proteins of this disclosure are well known in the art. For example, when a nutritive protein of this disclosure is used to maintain or increase at least one of muscle mass, muscle strength, and functional performance in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to maintain or increase at least 25 one of muscle mass, muscle strength, and functional performance in a subject, such as an anabolic steroid. When a nutritive protein of this disclosure is used to maintain or achieve a desirable body mass index in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to maintain or achieve a desirable body mass index in a subject, such as 30 orlistat, lorcaserin, sibutramine, rimonabant, metformin, exenatide, or pramlintide. When a nutritive protein of this disclosure is used to induce at least one of a satiation response and a satiety response in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product

indicated to induce at least one of a satiation response and a satiety response in a subject, such as rimonabant, exenatide, or pramlintide. When a nutritive protein of this disclosure is used to treat at least one of cachexia, sarcopenia and frailty in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to treat at least one of cachexia, sarcopenia and frailty, such as omega-3 fatty acids or anabolic steroids. Because of the role of dietary protein in inducing satiation and satiety, the nutritive proteins and nutritive compositions disclosed herein can be used to induce at least one of a satiation response and a satiety response in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00270] In some embodiments incorporating a least one nutritive protein or nutritive composition of this disclosure into the diet of a subject has at least one effect selected from inducing postprandial satiety (including by suppressing hunger), inducing thermogenesis, reducing glycemic response, positively affecting energy expenditure positively affecting lean body mass, reducing the weight gain caused by overeating, and decreasing energy intake. In some embodiments incorporating a least one nutritive protein or nutritive composition of this disclosure into the diet of a subject has at least one effect selected from increasing loss of body fat, reducing lean tissue loss, improving lipid profile, and improving glucose tolerance and insulin sensitivity in the subject.

## EXAMPLES

[00271] The following examples serve to more fully describe the manner of using the invention. These examples are presented for illustrative purposes and should not serve to limit the true scope of the invention.

**Example 1: Identification of Protein Fragments Containing Ratios of Essential Amino Acids, Branch Chain Amino Acids, and Leucine Greater Than or Equal to Whey and Containing All Essential Amino Acids**

[00272] The UniProtKB/Swiss-Prot (a collaboration between the European Bioinformatics Institute and the Swiss Institute of Bioinformatics) is a manually curated and reviewed protein database, and was used as the starting point for protein identification. Proteins from the edible species *Solanum lycopersicum*, *Zea mays*, *Oryza sativa subsp. Japonica*, *Glycine max*, *Ovis aries*, *Pisum sativum*, *Spinacia oleracea*, *Oryza sativa subsp. Indica*, *Triticum aestivum*, *Sus scrofa*, *Prunus persica*, *Capsicum annuum*, *Malus domestica*, *Thunnus albacares*, *Capra hircus*, *Cicer arietinum*, *Salmo salar*, *Meleagris gallopavo*, *Solanum tuberosum*, and *Agaricus bisporus* having greater than or equal to fifty (50) amino acids were considered as viable targets. Roughly 8,415 such proteins were selected for evaluation and the proteins were then fragmented into roughly fifty (50), roughly one hundred (100), or roughly one hundred and fifty (150) amino acids portions of the original protein, designed to contain digestive enzyme cleavage sites on both ends. In addition, the proteins were screened against a database of known allergens to determine whether any had greater than 50% global homology to a known allergen. A total of 1,000 fragments were identified that contain greater than or equal to 51% EAA, greater than or equal to 25% BCAA, and greater than or equal to 13% Leu, and contain all essential amino acids, and have less than 50% global homology to known allergens (**SEQ ID NOS: 1 to 1000**). (Note that for Examples 1-5, 51% EAA/25% BCAA/13% Leu represent values that are each 1-2% greater than those of whey (defined herein to be 49% EAA/24% BCAA/11% Leu). These values were used to identify nutritive protein fragments of interest for the purposes of this Example only, in order to ensure that the identified proteins have a higher content of EAA, BCAA, and Leu than whey). For the set of proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated, and the total number of Cys residues (“C”) were determined. The characteristics of 75 representative fragments thus identified are presented in **Tables 3A and 3B**.

**Table 3A**

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C
1	Q29550	101:301	0.53	0.27	0.13	3
2	Q29078	1:176	0.52	0.27	0.14	5
3	Q29550	201:426	0.56	0.25	0.13	2
4	Q29078	1:76	0.60	0.34	0.20	3
5	Q9N2D1	1:76	0.57	0.27	0.18	3
6	Q29079	1:76	0.56	0.29	0.18	1
7	Q29550	101:176	0.53	0.31	0.15	1
8	Q70BM6	201:326	0.58	0.26	0.15	0
9	Q70BM6	201:426	0.59	0.30	0.17	3
10	Q70BM6	101:326	0.56	0.28	0.14	2
11	O81122	101:301	0.53	0.29	0.16	1
12	Q9M7M1	101:201	0.55	0.28	0.19	1
13	O81122	101:201	0.55	0.28	0.19	1
14	Q9M7M1	101:301	0.53	0.28	0.16	1
15	Q8WNV7	151:226	0.51	0.29	0.14	1
16	P50118	176:276	0.52	0.26	0.15	3
17	P12263	451:526	0.55	0.26	0.13	0
18	Q29197	26:126	0.52	0.29	0.18	0
19	O02799	151:301	0.54	0.26	0.19	2
20	P80229	201:276	0.55	0.27	0.18	1
21	Q29624	226:326	0.59	0.25	0.13	2
22	A5GFQ5	26:151	0.52	0.25	0.15	3
23	Q5GN48	1601:1676	0.57	0.26	0.15	0
24	Q5GN48	1026:1101	0.55	0.26	0.16	2
25	A6M931	176:276	0.54	0.26	0.13	1
26	P49039	1:101	0.51	0.27	0.14	0
27	P79430	226:326	0.62	0.25	0.14	3
28	P61291	1:151	0.53	0.27	0.14	2
29	B5X2B8	126:326	0.52	0.25	0.13	3
30	Q1A730	226:301	0.52	0.28	0.20	2
31	D2SW95	276:476	0.52	0.31	0.13	3
32	O02799	251:326	0.57	0.29	0.22	1

33	B5RI54	126:251	0.54	0.26	0.13	2
34	P21999	76:176	0.51	0.28	0.13	5
35	Q07511	151:276	0.52	0.27	0.13	2
36	A5A779	351:476	0.53	0.31	0.24	4
37	Q6ITT4	476:601	0.54	0.30	0.16	1
38	O46374	876:1076	0.55	0.26	0.13	2
39	Q4U116	901:1076	0.55	0.25	0.14	2
40	P04045	376:476	0.53	0.27	0.13	1
41	Q29121	151:226	0.52	0.30	0.13	2
42	Q2VEC3	1601:1726	0.54	0.27	0.16	1
43	B5X737	26:176	0.57	0.26	0.13	2
44	Q9XSW2	201:401	0.53	0.25	0.15	8
45	P19756	276:351	0.60	0.32	0.14	0
46	P67931	101:201	0.51	0.26	0.19	1
47	Q9AWA5	401:476	0.58	0.27	0.13	0
48	O46374	976:1051	0.56	0.26	0.19	2
49	Q9XSW2	301:401	0.59	0.29	0.16	3
50	Q8MIQ9	551:651	0.56	0.28	0.15	1
51	O02668	526:626	0.53	0.26	0.14	0
52	Q6UAQ8	51:226	0.51	0.27	0.13	3
53	P21343	326:401	0.54	0.33	0.17	1
54	P79430	126:326	0.58	0.26	0.13	5
55	P10691	26:176	0.52	0.25	0.14	0
56	O02697	626:701	0.52	0.27	0.19	1
57	Q5VI41	676:751	0.53	0.30	0.13	1
58	O97716	201:376	0.52	0.25	0.13	4
59	Q66UW5	51:151	0.52	0.25	0.16	2
60	D2SW95	1:201	0.54	0.27	0.14	5
61	O46491	201:351	0.53	0.25	0.15	1
62	Q52QT8	1:151	0.53	0.26	0.16	5
63	Q9P4T7	1:76	0.53	0.25	0.17	0
64	Q2VEE8	201:276	0.60	0.25	0.14	0
65	P37831	251:376	0.52	0.26	0.16	0
66	B9EPI1	1:126	0.56	0.30	0.16	4
67	B5X2B8	126:201	0.61	0.30	0.15	0

68	O02799	526:651	0.57	0.26	0.16	1
69	Q99028	76:151	0.53	0.30	0.16	3
70	Q07511	51:276	0.51	0.26	0.13	4
71	P08454	1:76	0.57	0.28	0.13	1
72	Q5GN48	2251:2501	0.51	0.26	0.14	0
73	O64390	276:426	0.52	0.25	0.15	1
74	P09954	1:101	0.53	0.31	0.14	0
75	P29196	801:876	0.57	0.28	0.18	0

Table 3B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
1	-16.14	0.70	0.58	0.40	0.23	0.83
2	-15.72	0.57	0.54	0.00	0.25	0.50
3	-18.83	0.61	0.45	0.30	0.00	0.76
4	-18.32	0.80	0.43	0.28	0.29	0.46
5	-10.78	0.78	0.42	0.42	0.29	0.78
6	-18.03	0.62	0.41	0.28	0.28	0.48
7	-14.86	0.78	0.40	0.43	0.26	0.79
8	-17.96	0.60	0.36	0.00	0.25	0.81
9	-13.70	0.90	0.36	0.00	0.21	0.82
10	-15.03	0.80	0.36	0.00	0.22	0.81
11	-18.93	0.53	0.36	0.00	0.21	0.00
12	-22.36	0.47	0.35	0.21	0.00	0.00
13	-22.36	0.47	0.35	0.21	0.00	0.00
14	-18.90	0.52	0.35	0.00	0.24	0.00
15	-12.53	0.69	0.35	0.00	0.00	0.82
16	-20.41	0.43	0.35	0.20	0.25	0.86
17	-18.65	0.43	0.25	0.43	0.24	0.83
18	-28.11	0.37	0.31	0.00	0.00	0.98
19	-26.99	0.37	0.33	0.24	0.22	0.81
20	-26.56	0.35	0.34	0.29	1.00	0.79
21	-26.33	0.75	0.30	0.29	0.22	0.84
22	-25.32	0.42	0.29	0.23	0.29	0.95
23	-25.11	0.34	0.29	0.00	0.00	0.93

24	-24.91	0.25	0.28	0.00	0.26	0.88
25	-24.78	0.47	0.31	0.00	0.20	1.00
26	-24.60	0.44	0.29	0.25	0.31	0.25
27	-24.55	0.79	0.28	0.24	0.27	0.77
28	-24.41	0.38	0.33	0.00	0.25	1.00
29	-24.09	0.41	0.31	0.00	0.25	0.50
30	-23.74	0.38	0.27	0.26	0.26	0.99
31	-23.68	0.62	0.31	0.00	0.23	1.00
32	-23.68	0.38	0.27	0.00	0.24	0.80
33	-23.57	0.39	0.32	0.00	0.00	0.89
34	-23.51	0.64	0.33	0.17	0.27	0.99
35	-23.41	0.45	0.34	0.21	0.00	0.29
36	-23.38	0.50	0.32	0.23	0.26	0.98
37	-23.11	0.63	0.30	0.22	0.24	1.00
38	-23.11	0.41	0.33	0.22	0.00	0.99
39	-23.02	0.72	0.29	0.00	0.23	0.99
40	-22.98	0.36	0.30	0.00	0.28	0.57
41	-22.88	0.59	0.30	0.27	0.29	1.00
42	-22.87	0.49	0.32	0.00	0.23	0.00
43	-22.79	0.42	0.31	0.21	0.23	0.59
44	-22.74	0.56	0.31	0.00	0.21	0.88
45	-22.73	0.80	0.26	0.00	0.30	0.91
46	-22.72	0.50	0.29	0.00	0.30	0.73
47	-22.62	0.38	0.23	0.23	0.00	0.24
48	-22.55	0.42	0.24	0.27	0.24	1.00
49	-22.53	0.71	0.31	0.00	0.26	0.96
50	-22.43	0.59	0.30	0.00	0.00	0.90
51	-22.38	0.48	0.30	0.00	1.00	0.76
52	-22.32	0.47	0.33	0.00	0.00	0.96
53	-22.21	0.49	0.26	0.00	0.26	0.30
54	-22.18	0.64	0.31	0.20	0.21	0.78
55	-22.18	0.45	0.32	0.20	0.24	0.00
56	-22.16	0.52	0.24	0.00	0.23	1.00
57	-22.06	0.82	0.27	0.24	0.27	0.79
58	-22.05	0.59	0.30	0.00	0.22	1.00

59	-22.01	0.36	0.29	0.00	0.00	0.96
60	-21.91	0.62	0.33	0.00	0.21	1.00
61	-21.88	0.46	0.33	0.21	0.18	0.73
62	-21.83	0.49	0.30	0.24	0.22	0.99
63	-21.80	0.39	0.29	0.00	0.30	0.51
64	-21.75	0.44	0.28	0.00	0.24	0.22
65	-21.74	0.58	0.31	0.00	0.26	0.00
66	-21.65	0.52	0.33	0.25	0.26	0.82
67	-21.59	0.58	0.24	0.00	0.27	0.71
68	-21.58	0.40	0.30	0.26	0.25	0.85
69	-21.57	0.64	0.25	0.22	0.28	0.96
70	-21.54	0.46	0.31	0.22	0.00	0.28
71	-21.50	0.73	0.25	0.28	1.00	0.28
72	-21.47	0.34	0.32	0.22	0.00	0.86
73	-21.43	0.46	0.29	0.22	0.25	0.38
74	-21.41	0.62	0.31	0.00	0.27	0.85
75	-21.41	0.58	0.28	0.00	0.00	0.29

**Example 2: Identification of Protein Fragments Containing Ratios of Essential Amino Acids, Branch Chain Amino Acids, and Leucine Greater Than or Equal to Whey**

[00273] Fragments from the set of 8,415 proteins described in Example 1 were evaluated for amino acid content, percentage of essential amino acids (“EAA”), the percentage of branched chain amino acids (“BCAA”), the percentage of leucine (“L”), and whether the protein contained all of the essential amino acids were calculated for each protein. In addition, the protein fragments were screened against a database of known allergens to determine whether any had greater than 50% global homology to a known allergen. A total of 414 protein fragments were identified that contain greater than or equal to 51% EAA, greater than or equal to 25% BCAA, and greater than or equal to 13% Leu, and have less than 50% global homology to known allergens (**SEQ ID NOS: 1001 to 1414**). For the set of proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were

calculated, and the total number of Cys residues (“C”) were determined. The characteristics of 75 representative proteins thus identified are presented in **Tables 4A and 4B**.

**Table 4A**

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C
1001	P21641	1:126	0.52	0.30	0.19	1
1002	Q41782	151:276	0.52	0.25	0.14	4
1003	O24174	176:251	0.53	0.29	0.15	3
1004	Q41784	201:276	0.55	0.26	0.17	3
1005	P09315	276:351	0.51	0.35	0.14	1
1006	Q43695	201:276	0.53	0.26	0.17	3
1007	P18025	201:276	0.54	0.26	0.17	3
1008	P18026	201:276	0.54	0.26	0.17	3
1009	Q43697	201:276	0.54	0.26	0.17	3
1010	Q41783	201:276	0.54	0.26	0.17	3
1011	Q41785	201:276	0.54	0.26	0.17	3
1012	O24562	126:276	0.53	0.28	0.14	1
1013	Q6ZHS4	126:301	0.52	0.27	0.13	1
1014	Q6ERW7	226:301	0.55	0.29	0.13	1
1015	Q01595	51:126	0.52	0.36	0.22	7
1016	Q6ERW9	226:301	0.54	0.29	0.13	1
1017	Q0J6T3	151:276	0.55	0.28	0.13	1
1018	P93203	201:276	0.55	0.25	0.18	0
1019	Q7XWU3	151:276	0.52	0.27	0.14	1
1020	Q41874	126:251	0.51	0.29	0.16	2
1021	Q8H859	151:276	0.54	0.25	0.14	1
1022	P0CD58	301:476	0.53	0.32	0.18	3
1023	P0CD59	301:476	0.53	0.32	0.18	3
1024	O04161	251:376	0.58	0.26	0.13	5
1025	Q2R114	151:276	0.54	0.26	0.13	1
1026	Q5Z987	1051:1276	0.54	0.30	0.14	3
1027	Q41808	176:276	0.51	0.28	0.17	2
1028	Q41807	176:276	0.51	0.29	0.17	2
1029	P54773	101:276	0.55	0.29	0.15	0

1030	Q5Z987	1151:1251	0.60	0.34	0.14	2
1031	Q5Z987	1151:1351	0.54	0.33	0.16	4
1032	P31542	326:401	0.53	0.30	0.14	0
1033	P31541	126:201	0.51	0.32	0.15	0
1034	P04706	1:76	0.51	0.31	0.17	2
1035	O64411	1:76	0.53	0.32	0.18	0
1036	Q0JI49	351:426	0.60	0.26	0.17	0
1037	Q9XGC9	476:551	0.57	0.26	0.13	0
1038	P93648	326:401	0.51	0.25	0.17	1
1039	Q6H7U5	376:451	0.56	0.26	0.14	0
1040	Q43143	76:176	0.52	0.26	0.14	2
1041	Q9LGV5	51:126	0.52	0.27	0.14	0
1042	P0C8M8	126:201	0.51	0.26	0.16	4
1043	Q05761	51:126	0.56	0.32	0.16	0
1044	Q84NP7	326:426	0.54	0.26	0.16	1
1045	Q75HC2	276:351	0.51	0.27	0.14	2
1046	Q6QNU4	126:201	0.52	0.27	0.14	2
1047	Q5JMF2	51:126	0.55	0.26	0.14	4
1048	Q7XU38	276:351	0.54	0.25	0.16	1
1049	Q8GUQ5	1076:1151	0.53	0.27	0.16	2
1050	P51059	76:176	0.52	0.27	0.14	0
1051	P0C8M8	26:176	0.54	0.27	0.14	6
1052	P49037	26:101	0.51	0.29	0.15	0
1053	O49187	101:176	0.56	0.30	0.18	0
1054	Q0J4I1	26:176	0.52	0.25	0.14	2
1055	Q6ZAG3	51:176	0.53	0.25	0.13	1
1056	P29620	51:126	0.54	0.27	0.14	0
1057	P33544	201:276	0.54	0.26	0.15	1
1058	O24573	1:101	0.55	0.28	0.14	2
1059	Q3HRN8	126:201	0.55	0.26	0.15	0
1060	Q3HRP5	1:101	0.53	0.27	0.15	3
1061	P93647	801:876	0.53	0.32	0.16	0
1062	Q9ZS62	951:1026	0.53	0.35	0.16	0
1063	Q6Z4U2	276:351	0.53	0.29	0.13	1
1064	P29185	201:301	0.57	0.31	0.13	1

1065	Q75LU8	1:76	0.51	0.31	0.19	4
1066	Q43298	201:301	0.55	0.31	0.13	1
1067	P49971	26:101	0.52	0.27	0.13	2
1068	P46641	1:76	0.64	0.26	0.16	1
1069	Q2QMX9	701:776	0.52	0.25	0.14	0
1070	Q43270	226:301	0.53	0.25	0.13	0
1071	P93841	76:176	0.52	0.26	0.15	1
1072	P10581	151:226	0.52	0.29	0.14	1
1073	P93647	151:251	0.51	0.26	0.14	0
1074	Q2QQS5	126:201	0.52	0.30	0.13	0
1075	Q0JNK5	301:426	0.52	0.34	0.15	1

**Table 4B**

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
1001	-7.76	1.07	0.63	0.00	0.00	0.28
1002	-16.11	0.62	0.49	0.24	0.26	0.87
1003	-11.65	0.84	0.48	0.00	0.29	0.54
1004	-13.93	0.66	0.46	0.00	0.27	0.87
1005	-15.37	0.78	0.45	0.32	0.00	0.47
1006	-14.07	0.63	0.45	0.00	0.27	0.86
1007	-14.01	0.63	0.45	0.00	0.27	0.87
1008	-14.01	0.63	0.45	0.00	0.27	0.87
1009	-14.01	0.63	0.45	0.00	0.27	0.87
1010	-14.01	0.63	0.45	0.00	0.27	0.87
1011	-14.01	0.63	0.45	0.00	0.27	0.87
1012	-14.02	0.70	0.44	0.24	0.27	0.29
1013	-13.33	0.73	0.42	0.00	0.26	0.27
1014	-10.77	0.79	0.42	0.00	0.30	0.34
1015	-14.67	0.92	0.40	0.31	0.29	0.28
1016	-13.59	0.79	0.39	0.00	0.29	0.35
1017	-13.51	0.76	0.39	0.00	0.00	0.28
1018	-35.15	0.19	0.38	0.00	0.00	0.35
1019	-16.25	0.65	0.38	0.22	0.00	0.27
1020	-13.96	0.54	0.38	0.00	0.00	0.83

1021	-15.75	0.75	0.37	0.22	0.22	0.27
1022	-9.84	1.10	0.36	0.00	0.25	0.26
1023	-9.84	1.10	0.36	0.00	0.25	0.26
1024	-9.79	0.97	0.36	0.27	0.24	0.29
1025	-14.86	0.77	0.36	0.21	0.00	0.27
1026	-20.01	0.60	0.36	0.00	0.00	0.20
1027	-11.76	0.58	0.36	0.00	0.00	0.85
1028	-11.75	0.58	0.36	0.00	0.00	0.85
1029	-14.00	0.90	0.36	0.00	0.25	0.26
1030	-19.70	0.84	0.35	0.00	0.21	0.00
1031	-19.59	0.71	0.35	0.00	0.17	0.24
1032	-24.74	0.63	0.28	0.67	0.29	0.00
1033	-20.05	0.57	0.24	0.40	0.28	0.30
1034	-4.01	0.67	0.31	0.40	0.38	0.43
1035	-10.91	1.01	0.29	0.39	0.34	0.42
1036	-30.64	0.41	0.27	0.00	0.30	0.00
1037	-27.88	0.25	0.23	0.00	0.00	0.31
1038	-27.52	0.30	0.27	0.00	0.27	0.64
1039	-27.42	0.47	0.29	0.00	0.00	0.00
1040	-27.42	0.29	0.32	0.00	0.27	0.00
1041	-26.41	0.31	0.26	0.00	0.29	0.53
1042	-26.11	0.38	0.28	0.22	0.28	0.66
1043	-25.86	0.56	0.25	0.00	0.26	0.79
1044	-25.75	0.31	0.32	0.00	0.28	0.58
1045	-25.74	0.51	0.27	0.29	0.30	0.31
1046	-25.62	0.48	0.26	0.30	0.21	0.66
1047	-25.19	0.41	0.27	0.32	0.29	0.26
1048	-25.11	0.72	0.28	0.00	0.28	0.36
1049	-24.99	0.51	0.21	0.27	0.27	0.34
1050	-24.92	0.42	0.28	0.00	0.00	0.00
1051	-24.77	0.34	0.30	0.26	0.24	0.50
1052	-24.70	0.55	0.26	0.00	0.22	0.32
1053	-24.58	0.52	0.25	0.00	0.00	0.00
1054	-24.51	0.39	0.33	0.00	0.00	0.52
1055	-24.44	0.27	0.31	0.26	0.00	0.46

1056	-24.32	0.51	0.27	0.00	0.00	0.56
1057	-24.32	0.57	0.25	0.00	0.31	0.00
1058	-24.14	0.55	0.31	0.00	0.28	0.48
1059	-24.14	0.48	0.29	0.00	0.30	0.41
1060	-23.96	0.56	0.34	0.00	0.26	0.30
1061	-23.86	0.55	0.26	0.00	0.26	0.58
1062	-23.81	0.67	0.23	0.00	0.24	0.00
1063	-23.35	0.63	0.23	0.26	0.26	0.28
1064	-23.30	0.56	0.28	0.00	0.00	0.56
1065	-23.28	0.63	0.33	0.26	0.30	0.27
1066	-23.25	0.56	0.29	0.00	0.27	0.55
1067	-23.24	0.32	0.28	0.29	0.28	0.50
1068	-23.20	0.71	0.29	0.33	0.25	0.27
1069	-22.91	0.44	0.27	0.00	0.30	0.53
1070	-22.86	0.47	0.28	0.31	0.25	0.34
1071	-22.79	0.51	0.29	0.25	0.25	0.25
1072	-22.76	0.62	0.25	0.00	0.00	0.00
1073	-22.76	0.47	0.30	0.00	0.00	0.31
1074	-22.72	0.56	0.27	0.00	0.21	0.31
1075	-22.70	0.75	0.30	0.00	0.00	0.51

**Example 3: Identification of Protein Fragments Containing Ratios of Essential Amino Acids and Branch Chain Amino Acids Greater Than or Equal to Whey and Containing All Essential Amino Acids**

[00274] Fragments from the set of 8,415 proteins described in Example 1 were evaluated for amino acid content, percentage of essential amino acids (“EAA”), the percentage of branched chain amino acids (“BCAA”), the percentage of leucine (“L”), and whether the protein contained all of the essential amino acids were calculated for each protein. In addition, the protein fragments were screened against a database of known allergens to determine whether any had greater than 50% global homology to a known allergen. A total of 485 proteins were identified that contain greater than or equal to 51% EAA and greater than or equal to 25% BCAA, and that contain all essential amino acids, and have less than 50% global homology to known allergens (**SEQ ID NOS: 1415 to 1899**). For the set of protein fragments the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as

described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated, and the total number of Cys residues (“C”) were determined. The 5 characteristics of 75 representative proteins thus identified are presented in **Table 5A and 5B**.

**Table 5A**

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C
1415	Q96576	151:326	0.52	0.27	0.10	2
1416	P15722	176:276	0.53	0.28	0.04	1
1417	Q8GUQ5	876:1001	0.51	0.25	0.12	1
1418	P17801	476:751	0.52	0.26	0.12	3
1419	Q6ERW7	201:351	0.53	0.26	0.10	1
1420	Q7XWU3	51:276	0.51	0.25	0.09	6
1421	Q6ER94	76:226	0.52	0.26	0.10	1
1422	Q43272	176:376	0.52	0.26	0.10	6
1423	Q43272	76:326	0.52	0.28	0.11	7
1424	Q6ERW9	226:351	0.55	0.26	0.11	1
1425	Q10717	251:326	0.51	0.30	0.10	1
1426	Q6ERW9	126:276	0.51	0.25	0.06	5
1427	Q6ERW5	26:101	0.52	0.25	0.03	4
1428	P16243	226:476	0.51	0.27	0.12	4
1429	Q43272	76:226	0.51	0.28	0.12	4
1430	O04161	151:376	0.55	0.25	0.11	5
1431	Q9ATN1	101:251	0.56	0.29	0.11	2
1432	Q9ATN1	1:251	0.51	0.25	0.10	5
1433	P29185	26:301	0.52	0.25	0.08	3
1434	Q08451	126:276	0.53	0.27	0.08	2
1435	Q96575	1:276	0.52	0.26	0.12	4
1436	P09607	1:276	0.51	0.26	0.12	5
1437	Q6ETN3	401:501	0.52	0.25	0.06	0
1438	Q5BQN5	151:226	0.60	0.26	0.11	1
1439	Q2MI42	976:1176	0.64	0.25	0.10	0
1440	P28523	26:176	0.51	0.26	0.09	3

1441	Q8H6B1	126:226	0.57	0.26	0.10	1
1442	P54767	176:251	0.54	0.26	0.11	1
1443	Q67W82	401:526	0.51	0.26	0.06	0
1444	P0C8M8	1:226	0.53	0.25	0.11	7
1445	Q9FYT6	551:651	0.53	0.25	0.09	0
1446	P0C8M8	151:226	0.52	0.26	0.12	2
1447	Q10S72	426:551	0.52	0.26	0.10	1
1448	Q42982	426:551	0.53	0.26	0.06	0
1449	Q2MI46	26:101	0.51	0.25	0.09	4
1450	B4G0F3	276:401	0.52	0.26	0.07	1
1451	P93648	76:251	0.52	0.26	0.12	1
1452	C1K5M3	551:751	0.55	0.25	0.12	5
1453	Q41741	151:276	0.53	0.27	0.12	1
1454	O22424	151:251	0.60	0.25	0.05	0
1455	Q9XFH3	51:201	0.54	0.26	0.10	3
1456	Q7XTS4	201:451	0.51	0.25	0.11	3
1457	Q9FUZ0	176:276	0.52	0.25	0.12	2
1458	P05118	1:76	0.59	0.26	0.10	1
1459	Q8S7E1	51:126	0.52	0.27	0.12	1
1460	C1K5M2	101:201	0.55	0.25	0.10	3
1461	Q6Z4U2	151:351	0.52	0.26	0.12	4
1462	Q8H6B1	26:176	0.56	0.25	0.11	0
1463	Q0J7J7	176:351	0.51	0.27	0.11	5
1464	P10581	651:751	0.51	0.27	0.12	1
1465	Q2RAS0	26:251	0.51	0.25	0.09	2
1466	O24594	351:426	0.53	0.29	0.08	2
1467	Q0J7J7	276:351	0.54	0.31	0.12	3
1468	Q6XZ79	176:276	0.51	0.25	0.10	1
1469	P19862	651:801	0.53	0.26	0.11	3
1470	P28554	276:426	0.53	0.26	0.11	2
1471	C1K5M3	451:651	0.52	0.25	0.12	4
1472	P31927	451:551	0.51	0.26	0.12	1
1473	Q65X71	451:526	0.61	0.31	0.12	0
1474	Q00497	76:226	0.51	0.25	0.11	4
1475	Q7XKC0	76:176	0.51	0.26	0.06	3

1476	P10582	276:451	0.55	0.27	0.12	3
1477	P29535	176:401	0.51	0.26	0.11	8
1478	Q43298	26:301	0.51	0.25	0.08	3
1479	Q7XRA1	176:251	0.61	0.29	0.12	0
1480	Q852N2	826:1001	0.52	0.27	0.12	6
1481	Q65X71	451:726	0.52	0.26	0.09	5
1482	Q852N2	826:901	0.51	0.25	0.10	2
1483	Q65X71	51:326	0.52	0.27	0.10	3
1484	Q8H6B1	801:901	0.57	0.29	0.12	1
1485	P54767	176:451	0.52	0.26	0.10	2
1486	P61242	1626:1876	0.51	0.27	0.12	4
1487	Q65X71	451:626	0.54	0.26	0.09	3
1488	P04712	326:426	0.52	0.27	0.11	1
1489	Q75LR2	351:426	0.55	0.28	0.12	0

Table 5B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
1415	-16.93	0.58	0.54	0.53	0.24	0.00
1416	-19.58	0.41	0.53	0.00	0.21	0.00
1417	-21.91	0.46	0.47	0.27	0.27	0.37
1418	-19.73	0.54	0.46	0.00	0.00	0.34
1419	-11.15	0.74	0.43	0.00	0.25	0.29
1420	-16.57	0.61	0.42	0.22	0.00	0.27
1421	-20.09	0.54	0.41	0.23	0.00	0.63
1422	-16.36	0.68	0.41	0.00	0.24	0.34
1423	-16.55	0.67	0.40	0.00	0.21	0.32
1424	-11.31	0.79	0.40	0.24	0.00	0.27
1425	-17.05	0.64	0.39	0.28	0.35	0.53
1426	-11.49	0.73	0.38	0.00	0.26	0.35
1427	-17.25	0.63	0.38	0.25	0.28	0.41
1428	-19.07	0.59	0.38	0.21	0.00	0.60
1429	-16.46	0.67	0.36	0.25	0.23	0.33
1430	-11.00	0.89	0.36	0.00	0.00	0.26
1431	-6.06	1.08	0.36	0.27	0.27	0.36

1432	-8.76	0.87	0.36	0.23	0.23	0.28
1433	-21.28	0.49	0.36	0.00	0.00	0.60
1434	-10.55	0.95	0.35	0.25	0.24	0.44
1435	-17.94	0.61	0.34	0.37	0.00	0.00
1436	-18.54	0.63	0.33	0.36	0.19	0.00
1437	-28.54	0.46	0.30	0.00	0.23	0.38
1438	-25.54	0.40	0.27	0.00	0.00	0.26
1439	-25.12	0.37	0.33	0.24	0.00	0.19
1440	-25.09	0.34	0.31	0.00	0.21	0.80
1441	-24.93	0.38	0.32	0.00	0.25	0.27
1442	-24.55	0.45	0.22	0.00	0.00	0.32
1443	-24.41	0.49	0.29	0.24	0.00	0.33
1444	-24.26	0.36	0.31	0.00	0.00	0.49
1445	-23.95	0.53	0.28	0.00	0.00	0.00
1446	-23.76	0.45	0.30	0.00	0.26	0.70
1447	-23.69	0.45	0.30	0.24	0.27	0.39
1448	-23.58	0.48	0.31	0.00	0.24	0.36
1449	-23.51	0.56	0.27	0.25	0.27	0.36
1450	-23.40	0.48	0.31	0.25	0.00	0.43
1451	-23.21	0.41	0.30	0.00	0.22	0.27
1452	-23.11	0.50	0.30	0.00	0.23	0.00
1453	-22.98	0.53	0.30	0.29	0.26	0.71
1454	-22.62	0.40	0.27	0.00	0.00	0.70
1455	-22.54	0.48	0.31	0.24	0.24	0.42
1456	-22.52	0.37	0.33	0.24	0.00	0.30
1457	-22.50	0.44	0.30	0.00	0.00	0.44
1458	-22.49	0.61	0.27	0.24	1.00	0.26
1459	-22.48	0.47	0.23	0.00	0.27	0.24
1460	-22.30	0.61	0.30	0.23	0.19	0.36
1461	-22.29	0.45	0.30	0.21	0.24	0.00
1462	-22.25	0.48	0.31	0.00	0.21	0.29
1463	-22.20	0.44	0.30	0.00	0.22	0.00
1464	-22.05	0.53	0.28	0.00	0.00	0.00
1465	-22.05	0.69	0.30	0.00	0.00	0.29
1466	-22.02	0.45	0.29	0.00	0.32	0.54

1467	-22.00	0.48	0.27	0.00	0.30	0.32
1468	-21.97	0.50	0.29	0.00	0.00	0.31
1469	-21.83	0.54	0.30	0.23	0.00	0.00
1470	-21.80	0.58	0.31	0.25	0.00	0.23
1471	-21.78	0.56	0.30	0.21	0.23	0.00
1472	-21.60	0.54	0.30	0.00	0.00	0.00
1473	-21.53	0.69	0.24	0.00	0.26	0.33
1474	-21.44	0.48	0.33	0.21	0.00	0.27
1475	-21.39	0.65	0.28	0.26	0.00	0.59
1476	-21.27	0.57	0.30	0.00	0.24	0.00
1477	-21.26	0.50	0.33	0.00	0.23	0.30
1478	-21.26	0.49	0.32	0.00	0.00	0.60
1479	-21.13	0.55	0.28	0.00	0.00	0.28
1480	-21.08	0.61	0.33	0.17	0.24	0.27
1481	-21.00	0.50	0.31	0.19	0.22	0.39
1482	-20.97	0.43	0.24	0.17	0.26	0.00
1483	-20.94	0.68	0.33	0.00	0.23	0.34
1484	-20.89	0.61	0.29	0.00	0.26	0.51
1485	-20.84	0.48	0.31	0.00	0.00	0.23
1486	-20.74	0.51	0.30	0.00	0.22	0.22
1487	-20.65	0.51	0.31	0.28	0.27	0.34
1488	-20.58	0.44	0.32	0.00	0.00	0.00
1489	-20.54	0.44	0.28	0.00	0.00	0.21

**Example 4: Identification of Protein Fragments Containing Ratios of Essential Amino Acids and Leucine Greater Than or Equal to Whey and Containing All Essential Amino Acids.**

5 [00275] Fragments from the set of 8,415 proteins described in Example 1 were evaluated for amino acid content, percentage of essential amino acids (“EAA”), the percentage of branched chain amino acids (“BCAA”), the percentage of leucine (“L”), and whether the protein contained all of the essential amino acids were calculated for each protein. In addition, the protein fragments were screened against a database of known allergens to determine whether any had greater than 50% global homology to a known 10 allergen. A total of 203 protein fragments were identified that contain greater than or equal

to 51% EAA and greater than or equal to 13% Leu, and that contain all essential amino acids, and have less than 50% global homology to known allergens (**SEQ ID NOS: 1900 to 2102**). For the set of protein fragments the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as

described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated, and the total number of Cys residues (“C”) were determined. The characteristics of 75 representative protein fragments thus identified are presented in **Table 6A and 6B**.

**10 6A and 6B.**

**Table 6A**

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C
1900	Q41784	201:301	0.51	0.22	0.15	4
1901	P18025	201:301	0.51	0.22	0.15	4
1902	Q43697	201:301	0.51	0.22	0.15	4
1903	Q41783	201:301	0.51	0.22	0.15	4
1904	Q41785	201:301	0.51	0.22	0.15	4
1905	P18026	201:301	0.51	0.22	0.15	4
1906	P10708	76:176	0.51	0.24	0.13	1
1907	P27525	51:151	0.53	0.22	0.14	0
1908	Q3HRN9	51:201	0.52	0.23	0.14	1
1909	Q9ZSV1	726:801	0.51	0.23	0.14	1
1910	Q5BQN5	51:176	0.52	0.20	0.13	1
1911	P56669	1:76	0.53	0.18	0.14	1
1912	Q75LU8	1:201	0.52	0.24	0.13	4
1913	Q8S0F0	601:851	0.51	0.23	0.14	3
1914	Q9ZSV1	626:801	0.51	0.23	0.13	1
1915	Q0JI49	301:451	0.53	0.23	0.14	0
1916	P93703	201:401	0.51	0.24	0.14	1
1917	Q2MI42	426:601	0.52	0.23	0.13	1
1918	Q8S0F0	501:751	0.52	0.23	0.13	4
1919	Q2MI42	1601:1701	0.51	0.23	0.14	1
1920	Q7XTS3	401:476	0.57	0.21	0.13	2

1921	P49037	1:201	0.51	0.24	0.14	1
1922	Q9FQ11	1:76	0.53	0.24	0.16	1
1923	Q7X996	201:376	0.55	0.22	0.13	1
1924	P27061	126:201	0.53	0.24	0.14	0
1925	P53682	326:401	0.51	0.24	0.13	1
1926	Q84ZL0	1201:1476	0.51	0.24	0.16	4
1927	Q8S0F0	701:851	0.51	0.24	0.14	1
1928	Q6ZAG3	51:201	0.54	0.24	0.13	1
1929	Q69Q02	276:451	0.52	0.22	0.13	1
1930	P10581	276:526	0.51	0.24	0.13	2
1931	P49037	201:276	0.52	0.24	0.15	2
1932	P61242	776:951	0.51	0.23	0.15	2
1933	Q5Z987	1826:1926	0.51	0.19	0.14	1
1934	Q5JMM1	251:326	0.52	0.22	0.14	0
1935	Q84N48	426:526	0.51	0.24	0.14	0
1936	Q7G6K7	826:1101	0.51	0.23	0.14	4
1937	Q0J4I1	1:201	0.53	0.24	0.14	2
1938	Q7XWS7	1226:1476	0.51	0.22	0.13	3
1939	Q0JF02	76:276	0.51	0.24	0.14	4
1940	Q2QN26	201:351	0.51	0.24	0.14	5
1941	P49037	101:276	0.53	0.22	0.14	2
1942	Q6K5F8	351:626	0.51	0.24	0.14	3
1943	Q10QA2	101:226	0.51	0.22	0.13	6
1944	Q9XHR2	201:301	0.52	0.24	0.14	0
1945	Q0DCT8	176:351	0.52	0.24	0.15	5
1946	Q6AU53	476:576	0.52	0.24	0.16	0
1947	A7LFZ6	276:451	0.52	0.23	0.15	6
1948	Q6ZKB2	451:726	0.52	0.22	0.13	2
1949	Q8S983	176:251	0.51	0.22	0.14	0
1950	Q3LXA7	526:626	0.53	0.24	0.14	2
1951	P93703	201:276	0.55	0.24	0.14	1
1952	A3B9A0	76:226	0.51	0.21	0.13	2
1953	Q9XGD6	51:126	0.52	0.22	0.13	0
1954	P18485	301:376	0.54	0.20	0.14	2
1955	Q32SG6	876:951	0.52	0.23	0.16	0

1956	Q5VS72	501:576	0.54	0.24	0.18	2
1957	Q43503	401:476	0.54	0.21	0.15	1
1958	Q42896	101:326	0.52	0.24	0.13	4
1959	Q9MAX6	151:251	0.52	0.24	0.14	3
1960	Q9LWX9	1:201	0.52	0.22	0.14	5
1961	P09562	176:301	0.52	0.23	0.15	3
1962	Q7XWS7	1226:1376	0.51	0.22	0.13	1
1963	Q7XU38	151:351	0.51	0.24	0.14	4
1964	Q69LX2	926:1151	0.51	0.24	0.13	5
1965	Q7XD96	601:726	0.51	0.22	0.14	5
1966	Q7XI73	26:101	0.52	0.17	0.13	5
1967	Q42881	301:376	0.53	0.20	0.14	3
1968	C1K5M3	176:301	0.53	0.23	0.14	4
1969	Q6L4F8	226:301	0.53	0.22	0.15	2
1970	P48022	1:101	0.51	0.23	0.14	0
1971	Q8LHA8	51:201	0.51	0.24	0.14	3
1972	Q6ZAG3	151:301	0.52	0.24	0.14	3
1973	P04711	751:901	0.53	0.24	0.13	0
1974	Q5N870	576:801	0.53	0.23	0.13	8

Table 6B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
1900	-14.31	0.56	0.48	0.00	0.25	0.88
1901	-14.31	0.54	0.46	0.00	0.25	0.88
1902	-14.31	0.54	0.46	0.00	0.25	0.88
1903	-14.31	0.54	0.46	0.00	0.25	0.88
1904	-14.31	0.54	0.46	0.00	0.25	0.88
1905	-14.42	0.53	0.46	0.00	0.24	0.87
1906	-16.21	0.63	0.41	0.00	0.00	0.28
1907	-14.10	0.61	0.39	0.00	0.25	0.00
1908	-24.00	0.41	0.36	0.22	0.26	0.31
1909	-27.98	0.27	0.25	0.29	0.33	0.49
1910	-27.07	0.17	0.33	0.23	0.24	0.00
1911	-27.03	0.25	0.22	0.00	0.27	0.41

1912	-25.61	0.41	0.35	0.23	0.25	0.32
1913	-24.50	0.36	0.33	0.00	0.00	0.30
1914	-24.43	0.35	0.35	0.00	0.23	0.39
1915	-24.41	0.37	0.32	0.24	0.23	0.00
1916	-24.40	0.42	0.32	0.00	0.24	0.30
1917	-24.34	0.23	0.33	0.00	0.23	0.00
1918	-23.77	0.39	0.32	0.00	0.00	0.34
1919	-23.65	0.44	0.31	0.00	0.22	0.00
1920	-23.49	0.37	0.28	0.00	0.00	0.33
1921	-23.32	0.40	0.33	0.22	0.25	0.00
1922	-23.31	0.48	0.25	0.26	0.24	0.00
1923	-23.23	0.39	0.30	0.00	0.21	0.27
1924	-23.18	0.42	0.24	0.00	0.00	0.00
1925	-23.13	0.29	0.28	0.25	0.26	0.27
1926	-23.12	0.37	0.34	0.00	0.00	0.32
1927	-23.00	0.35	0.34	0.00	0.25	0.29
1928	-22.83	0.31	0.31	0.00	0.22	0.49
1929	-22.74	0.36	0.29	0.00	0.23	0.00
1930	-22.67	0.39	0.31	0.00	0.00	0.00
1931	-22.62	0.50	0.26	0.25	0.27	0.26
1932	-22.44	0.44	0.33	0.24	0.22	0.00
1933	-22.39	0.41	0.31	0.22	0.25	0.34
1934	-22.25	0.36	0.22	0.00	0.33	0.33
1935	-22.24	0.40	0.30	0.00	0.25	0.00
1936	-22.14	0.35	0.32	0.00	0.00	0.31
1937	-22.12	0.41	0.34	0.00	0.00	0.48
1938	-21.99	0.35	0.31	0.00	0.00	0.29
1939	-21.98	0.49	0.33	0.00	0.23	0.00
1940	-21.79	0.55	0.32	0.26	0.24	0.36
1941	-21.63	0.40	0.29	0.00	0.21	0.00
1942	-21.60	0.42	0.33	0.00	0.24	0.65
1943	-21.54	0.62	0.30	0.25	0.24	0.33
1944	-21.23	0.37	0.28	0.00	0.00	0.42
1945	-21.20	0.48	0.32	0.00	0.25	0.40
1946	-21.19	0.55	0.30	0.00	0.25	0.00

1947	-21.18	0.43	0.30	0.00	0.25	0.00
1948	-21.10	0.36	0.32	0.00	0.00	0.31
1949	-20.98	0.41	0.24	0.00	0.26	0.24
1950	-20.85	0.56	0.30	0.00	0.24	0.37
1951	-20.78	0.47	0.26	0.00	0.23	0.31
1952	-20.67	0.40	0.30	0.23	0.22	0.00
1953	-20.65	0.55	0.26	0.00	0.00	0.37
1954	-20.62	0.51	0.28	0.28	0.26	0.30
1955	-20.55	0.33	0.26	0.00	0.29	0.00
1956	-20.51	0.45	0.26	0.28	0.00	0.46
1957	-20.51	0.66	0.26	0.00	0.25	0.26
1958	-20.48	0.48	0.32	0.00	0.20	0.25
1959	-20.40	0.55	0.32	0.00	0.26	0.56
1960	-20.37	0.48	0.29	0.24	0.22	0.00
1961	-20.36	0.45	0.29	0.25	0.23	0.00
1962	-20.34	0.33	0.30	0.23	0.24	0.30
1963	-20.33	0.59	0.30	0.00	0.22	0.25
1964	-20.31	0.50	0.30	0.21	0.20	0.25
1965	-20.18	0.36	0.29	0.30	0.21	0.32
1966	-20.15	0.38	0.28	0.31	0.29	0.27
1967	-20.14	0.50	0.31	0.00	0.27	0.36
1968	-20.11	0.36	0.27	0.00	0.00	0.00
1969	-19.97	0.40	0.23	0.23	0.28	0.63
1970	-19.89	0.73	0.30	0.00	0.00	0.29
1971	-19.81	0.55	0.32	0.22	0.22	0.31
1972	-19.49	0.46	0.33	0.00	0.25	0.51
1973	-19.43	0.48	0.34	0.26	0.20	0.00
1974	-19.42	0.46	0.32	0.00	0.00	0.26

**Example 5: Identification of Proteins Containing Ratios of Branch Chain Amino Acids and Leucine Greater Than or Equal to Whey and Containing All Essential Amino Acids.**

[00276] Fragments from the set of 8,415 proteins described in Example 1 were evaluated for amino acid content, percentage of essential amino acids (“EAA”), the percentage of branched chain amino acids (“BCAA”), the percentage of leucine (“L”), and whether the protein contained all of the essential amino acids were calculated for each

protein. In addition, the proteins were screened against a database of known allergens to determine whether any had greater than 50% global homology to a known allergen. A total of 415 protein fragments were identified that contain greater than or equal to 25% BCAA and greater than or equal to 13% Leu, and that contain all essential amino acids, and have less than 50% global homology to known allergens (**SEQ ID NOS: 2103 to 2518**). For the set of protein fragments the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated, and the total number of Cys residues (“C”) were determined. The characteristics of 75 representative protein fragments thus identified are presented in **Table 7A and 7B**.

**Table 7A**

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C
2103	O24174	76:251	0.49	0.26	0.14	4
2104	P24396	1:126	0.48	0.27	0.13	2
2105	P17801	501:776	0.49	0.26	0.13	3
2106	O24585	501:751	0.47	0.26	0.14	3
2107	O24585	626:751	0.46	0.29	0.14	1
2108	P42211	1:101	0.50	0.27	0.20	2
2109	Q75L42	51:226	0.48	0.28	0.13	3
2110	Q9LGV5	1:251	0.48	0.26	0.13	4
2111	Q9LGV5	101:226	0.45	0.26	0.14	3
2112	Q10PS6	101:276	0.47	0.26	0.13	5
2113	Q5JLQ9	76:226	0.47	0.26	0.15	3
2114	Q2QMI0	26:226	0.43	0.26	0.15	3
2115	Q0D4B2	201:301	0.47	0.27	0.14	3
2116	Q8GUQ5	351:626	0.49	0.30	0.20	6
2117	Q8GUQ5	451:626	0.47	0.31	0.22	3
2118	Q8GUQ5	451:726	0.46	0.28	0.18	4
2119	Q2QY53	126:226	0.46	0.25	0.17	3
2120	Q2RAX3	126:226	0.46	0.25	0.17	3
2121	Q75GK4	1:226	0.46	0.25	0.16	3

2122	P16243	176:376	0.49	0.28	0.13	4
2123	Q41807	151:401	0.49	0.26	0.13	3
2124	Q69Q47	51:226	0.49	0.27	0.14	3
2125	P46420	1:201	0.48	0.25	0.13	3
2126	Q6Z9F4	51:226	0.47	0.27	0.13	3
2127	P16025	1026:1226	0.50	0.26	0.14	3
2128	P16025	926:1151	0.45	0.26	0.13	2
2129	P49174	1:76	0.47	0.29	0.14	0
2130	P48186	1:176	0.49	0.25	0.13	1
2131	P93648	276:401	0.50	0.26	0.16	2
2132	P49037	551:626	0.49	0.26	0.15	1
2133	Q2MIA9	201:276	0.48	0.28	0.15	1
2134	P16024	201:276	0.48	0.28	0.15	1
2135	P16024	201:376	0.44	0.26	0.13	2
2136	P49037	1:101	0.49	0.27	0.14	1
2137	Q84N48	376:451	0.49	0.27	0.13	2
2138	Q94IW5	251:351	0.45	0.27	0.14	1
2139	Q75LD5	576:726	0.44	0.25	0.13	1
2140	Q84ZX8	1:76	0.47	0.25	0.17	0
2141	Q336R3	26:101	0.43	0.26	0.20	3
2142	P93231	576:851	0.48	0.26	0.14	6
2143	P93648	276:551	0.48	0.25	0.13	4
2144	P16024	201:476	0.47	0.28	0.13	3
2145	Q2NM15	26:251	0.47	0.26	0.16	3
2146	P93231	576:751	0.50	0.25	0.14	3
2147	C7E5V8	26:251	0.47	0.26	0.16	3
2148	C7E5V7	26:251	0.48	0.27	0.17	3
2149	P30792	126:226	0.49	0.28	0.15	0
2150	Q5VS72	151:276	0.46	0.25	0.14	1
2151	Q9XGC9	426:501	0.45	0.27	0.16	0
2152	P61242	1626:1776	0.47	0.28	0.14	3
2153	Q9XGD6	126:251	0.46	0.27	0.14	2
2154	B6SJQ0	1:176	0.40	0.26	0.16	4
2155	P93207	1:126	0.42	0.25	0.14	2
2156	Q653U3	801:876	0.44	0.26	0.14	1

2157	P51059	351:576	0.44	0.25	0.15	4
2158	P07920	76:226	0.48	0.25	0.15	2
2159	P16024	301:451	0.48	0.28	0.13	2
2160	P29390	176:251	0.50	0.25	0.14	1
2161	Q5VQ78	576:676	0.47	0.25	0.14	0
2162	P16025	926:1026	0.46	0.27	0.13	1
2163	P51059	476:576	0.46	0.25	0.15	1
2164	P17847	326:451	0.38	0.25	0.16	1
2165	P05116	76:226	0.47	0.25	0.15	2
2166	P24157	76:226	0.47	0.25	0.15	2
2167	Q9XGC9	301:401	0.48	0.25	0.15	1
2168	P04711	351:551	0.45	0.25	0.13	3
2169	Q84ZW8	376:551	0.50	0.26	0.14	5
2170	Q6H6V4	801:876	0.45	0.26	0.13	1
2171	Q9XGD6	26:251	0.48	0.25	0.14	2
2172	Q41342	301:401	0.49	0.27	0.15	0
2173	Q42883	451:626	0.45	0.25	0.14	6
2174	Q2MI98	26:201	0.45	0.25	0.13	1
2175	Q7XTS4	251:426	0.49	0.26	0.13	3
2176	Q5Z987	1276:1451	0.47	0.25	0.14	8
2177	Q6AVM3	276:351	0.47	0.26	0.16	2

**Table 7B**

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
2103	-17.26	0.66	0.55	0.00	0.27	0.46
2104	-14.60	0.68	0.49	0.25	0.26	0.23
2105	-21.22	0.41	0.46	0.00	0.00	0.36
2106	-19.98	0.48	0.44	0.20	0.23	0.35
2107	-18.81	0.59	0.44	0.00	0.00	0.40
2108	-15.83	0.61	0.43	0.29	0.27	0.40
2109	-20.27	0.45	0.42	0.00	0.25	0.55
2110	-21.01	0.39	0.42	0.00	0.00	0.51

2111	-19.90	0.44	0.42	0.25	0.00	0.59
2112	-13.77	0.63	0.41	0.00	0.19	0.28
2113	-18.33	0.59	0.40	0.27	0.23	0.53
2114	-19.91	0.56	0.40	0.25	0.25	0.44
2115	-17.67	0.47	0.39	0.00	0.24	0.59
2116	-12.97	0.46	0.39	0.00	0.24	0.32
2117	-12.09	0.48	0.39	0.00	0.24	0.35
2118	-13.41	0.44	0.39	0.00	0.22	0.31
2119	-17.69	0.50	0.38	0.00	0.00	0.55
2120	-17.69	0.50	0.38	0.00	0.00	0.55
2121	-18.70	0.55	0.38	0.21	0.00	0.42
2122	-18.11	0.64	0.38	0.00	0.24	0.61
2123	-16.10	0.45	0.38	0.00	0.00	0.78
2124	-19.89	0.47	0.37	0.24	0.23	0.48
2125	-18.94	0.60	0.37	0.24	0.25	0.28
2126	-21.69	0.54	0.37	0.19	0.25	0.52
2127	-17.96	0.48	0.36	0.24	0.00	0.00
2128	-19.80	0.48	0.36	0.20	0.00	0.00
2129	-10.98	0.80	0.36	0.31	0.28	0.34
2130	-20.67	0.47	0.35	0.00	0.23	0.00
2131	-29.02	0.26	0.33	0.26	0.24	0.60
2132	-28.96	0.53	0.26	0.00	0.27	0.26
2133	-28.54	0.45	0.28	0.23	0.26	0.30
2134	-28.35	0.49	0.25	0.27	0.00	0.23
2135	-27.16	0.36	0.31	0.00	0.24	0.28
2136	-25.78	0.45	0.33	0.00	0.25	0.30
2137	-25.22	0.53	0.27	0.19	0.00	0.33
2138	-24.85	0.53	0.30	0.00	0.26	0.29
2139	-24.69	0.52	0.33	0.20	0.00	0.00
2140	-24.65	0.46	0.24	0.00	0.30	0.27
2141	-24.35	0.46	0.24	0.29	0.27	0.27
2142	-24.29	0.47	0.30	0.00	0.00	0.40
2143	-24.22	0.34	0.33	0.22	0.00	0.67
2144	-23.92	0.43	0.31	0.21	0.25	0.27
2145	-23.69	0.40	0.32	0.00	0.00	0.00

2146	-23.69	0.50	0.30	0.00	0.20	0.43
2147	-23.61	0.41	0.32	0.00	0.00	0.23
2148	-23.59	0.42	0.33	0.00	0.00	0.00
2149	-23.57	0.39	0.31	0.00	0.27	0.00
2150	-23.48	0.33	0.30	0.00	0.24	0.34
2151	-23.44	0.29	0.30	0.00	0.25	0.26
2152	-23.42	0.51	0.30	0.00	0.23	0.27
2153	-23.41	0.52	0.29	0.27	0.00	0.40
2154	-23.34	0.47	0.32	0.00	0.26	0.00
2155	-23.10	0.47	0.34	0.20	0.22	0.54
2156	-23.09	0.45	0.28	0.25	0.28	0.00
2157	-23.07	0.47	0.32	0.00	0.21	0.00
2158	-23.01	0.33	0.31	0.00	0.25	0.24
2159	-22.95	0.47	0.31	0.24	0.21	0.32
2160	-22.81	0.26	0.26	0.00	0.00	0.42
2161	-22.75	0.46	0.33	0.00	0.21	0.53
2162	-22.74	0.44	0.27	0.27	0.28	0.00
2163	-22.72	0.54	0.30	0.00	0.28	0.24
2164	-22.69	0.36	0.30	0.24	0.00	0.32
2165	-22.64	0.34	0.31	0.25	0.00	0.00
2166	-22.64	0.34	0.31	0.25	0.00	0.00
2167	-22.52	0.32	0.29	0.00	0.00	0.47
2168	-22.29	0.48	0.31	0.00	0.25	0.00
2169	-22.23	0.51	0.32	0.19	0.22	0.00
2170	-22.16	0.48	0.29	0.25	0.27	0.22
2171	-22.13	0.49	0.32	0.00	0.00	0.38
2172	-22.05	0.63	0.34	0.00	0.00	0.00
2173	-21.94	0.45	0.34	0.00	0.26	0.37
2174	-21.89	0.34	0.29	0.00	0.21	0.00
2175	-21.84	0.41	0.33	0.00	0.25	0.28
2176	-21.74	0.50	0.32	0.00	0.25	0.30
2177	-21.74	0.50	0.32	0.30	0.28	0.38

**Example 6: Protein Expression**

[00277] Genes encoding nutritive proteins disclosed herein were codon optimized for expression in *Escherichia coli* and synthesized by either LifeTechnologies/GeneArt or DNA 2.0. Genes were designed to contain one of two amino-terminal tags to facilitate purification:

5 [00278] MGSHHHHHHH (SEQ ID NO: 2615)

[00279] MGSSHHHHHSSGLVPRGSH (SEQ ID NO: 2614)

[00280] These gene constructs were inserted into the pET15b plasmid vector (Novagen) using *NcoI*-*BamHI* restriction sites (in case of the first tag) or using the *NdeI*-*BamHI* restriction sites (in the case of the second tag). All restriction enzymes were 10 purchased from New England Biolabs. Plasmids were transformed into *Escherichia coli* T7 Express (New England Biolabs) and selected on lysogeny broth (LB) plates containing 100 mg/l carbenicillin. A single colony was picked, grown to  $OD_{600nm} \approx 0.6$  in LB with 100 mg/l carbencillin, and stored as a glycerol stock (in LB with 10% glycerol (v/v)) at  $-80^{\circ}C$ , to serve as a master cell stock.

15 [0042] 2ml LB with 100 mg/l carbenicillin (in a 14mm $\times$ 100mm culture tube) was inoculated with a stab from the glycerol stock and grown overnight at  $37^{\circ}C$  and 250rpm. The next day, 2ml LB with 100 mg/l carbenicillin (in a 14mm $\times$ 100mm culture tube) was inoculated with the overnight culture to  $OD_{600nm} = 0.05$  and grown at  $30^{\circ}C$  or  $37^{\circ}C$  and 250rpm. At  $OD_{600nm} \approx 0.8$ , heterologous gene-expression was initiated with 1mM isopropyl

20  $\beta$ -D-1-thiogalactopyranoside (IPTG) and grown for another 2 hr (when grown at  $37^{\circ}C$ ) or 4 hr (when grown at  $30^{\circ}C$ ) until harvest. Upon harvesting,  $OD_{600nm}$  was measured, a 1ml aliquot was centrifuged, and the supernatant was decanted. Cells were re-suspended to  $OD_{600nm} = 1.50$  for SDS-PAGE analysis to evaluate expression level. 10 $\mu$ l of resuspended culture was loaded onto either: 1) a Novex<sup>®</sup> NuPAGE<sup>®</sup> 12% Bis-Tris gel (Life

25 Technologies), or 2) a Novex<sup>®</sup> 16% Tricine gel (Life Technologies), and run using standard manufacturer's protocols. Gels were stained using SimplyBlue<sup>™</sup> SafeStain (Life Technologies) using the standard manufacturer's protocol and imaged using the Molecular Imager<sup>®</sup> Gel Doc<sup>™</sup> XR+ System (Bio-Rad). Over-expressed heterologous protein was identified by comparison against a molecular weight marker and control cultures.

30 [00281] Using this method, recombinant expression of the proteins listed in Tables 8A and 8B was observed.

**Table 8A**

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C
2519	P15989	388:487	0.52	0.36	0.15	0
2520	P15989	341:442	0.48	0.36	0.16	0
2521	-	-	0.50	0.41	0.16	1
2522	P10587	1287:1386	0.49	0.26	0.17	0
2523	Q27991	1353:1452	0.46	0.23	0.17	0
2524	-	-	0.64	0.33	0.21	0
2525	P15989	398:447	0.57	0.41	0.18	0
2526	P02662	2:51	0.59	0.31	0.18	1
2527	P02662	3:52	0.59	0.33	0.18	1
2528	P02662	4:53	0.57	0.31	0.16	1
2529	P02662	5:54	0.55	0.29	0.14	1
2530	P02662	6:55	0.55	0.29	0.16	1
2531	P02662	79:128	0.41	0.28	0.12	0
2532	P02662	86:135	0.41	0.27	0.13	0
2533	Q9FRT9	1:90	0.32	0.11	0.07	10
2534	P09643	1:322	0.44	0.20	0.08	10
2535	Q9HD67	472:521	0.49	0.29	0.20	1
2536	Q02440	461:511	0.50	0.23	0.13	1
2537	Q13402	452:509	0.52	0.24	0.11	1
2538	Q02440	413:512	0.52	0.22	0.09	2
2539	Q28970	427:527	0.51	0.21	0.10	1
2540	Q90584	436:486	0.60	0.41	0.32	0
2541	P22281	82:134	0.58	0.46	0.24	1
2542	Q17R14	270:321	0.64	0.39	0.20	0
2543	Q27991	1396:1446	0.49	0.31	0.23	0
2544	P12106	170:220	0.70	0.32	0.14	1
2545	P32191	112:163	0.58	0.31	0.15	1
2546	P19524	1398:1449	0.56	0.25	0.18	2
2547	Q03262	225:277	0.65	0.30	0.13	1
2548	P12863	82:132	0.51	0.33	0.17	1
2549	P12106	159:209	0.67	0.29	0.14	1
2550	A6QR56	589:691	0.47	0.34	0.21	2

2551	P32492	1218:1319	0.67	0.25	0.14	1
2552	P79114	688:788	0.48	0.26	0.15	2
2553	Q5MIB5	492:592	0.53	0.27	0.14	2
2554	P04119	28:129	0.51	0.26	0.17	2
2555	P06642	14:115	0.55	0.27	0.14	2
2556	Q2XQV4	108:213	0.49	0.26	0.15	2
2557	P32492	1086:1287	0.58	0.29	0.15	1
2558	Q76FS2	59:260	0.46	0.23	0.12	7
2559	Q36967	1:105	0.65	0.46	0.26	0
2560	Q31721	1:58	0.63	0.38	0.19	2
2561	Q5ZMN0	44:93	0.53	0.36	0.26	0
2562	Q5ZMN0	46:95	0.54	0.36	0.26	0
2563	Q5ZMN0	47:96	0.52	0.36	0.26	0
2564	Q5ZMN0	1:146	0.53	0.30	0.21	2
2565	Q5ZMN0	1:147	0.52	0.30	0.21	2
2566	Q5ZMN0	1:148	0.53	0.30	0.21	2
2567	Q5ZMN0	1:149	0.52	0.29	0.21	2
2568	Q5ZMN0	1:150	0.52	0.29	0.21	2
2569	Q5ZMN0	1:151	0.52	0.29	0.20	2
2570	Q5ZMN0	1:152	0.51	0.29	0.20	2
2571	Q5ZMN0	1:153	0.51	0.29	0.20	2
2572	Q5ZMN0	1:154	0.51	0.28	0.20	2
2573	Q5ZMN0	1:161	0.49	0.27	0.19	2
2574	Q9JLT0	127:231	0.48	0.21	0.15	0
2575	Q27991	141:190	0.51	0.28	0.23	0
2576	Q27991	136:185	0.51	0.28	0.21	0
2577	Q27991	116:185	0.51	0.27	0.21	0
2578	Q27991	146:200	0.51	0.29	0.21	0
2579	Q27991	146:210	0.52	0.27	0.20	0
2580	Q27991	136:190	0.50	0.27	0.21	0
2581	Q27991	146:195	0.50	0.30	0.21	0
2582	Q27991	126:190	0.52	0.26	0.19	0
2583	Q27991	141:200	0.51	0.28	0.21	0
2584	Q27991	161:210	0.50	0.24	0.19	0
2585	Q27991	126:185	0.53	0.27	0.19	0

2586	Q27991	111:210	0.50	0.26	0.19	0
2587	Q27991	116:215	0.50	0.26	0.18	0
2588	Q27991	126:225	0.46	0.22	0.15	0
2589	Q27991	126:237	0.45	0.21	0.15	0
2590	Q9JLT0	194:244	0.45	0.20	0.14	0
2591	Q61879	154:252	0.41	0.19	0.14	0
2592	P15989	99:208	0.52	0.37	0.17	0
2593	P15989	99:203	0.51	0.37	0.16	0
2594	P15989	49:166	0.44	0.28	0.15	0
2595	Q90339	201:265	0.45	0.21	0.18	0
2596	Q9BE41	216:265	0.48	0.21	0.18	0
2597	Q9TV62	241:290	0.48	0.17	0.15	0
2598	Q5SX39	236:285	0.45	0.19	0.17	0
2599	Q9TV61	151:265	0.52	0.21	0.16	0
2600	Q27991	20:289	0.38	0.20	0.15	0
2601	Q0WVK7	53:105	0.57	0.27	0.12	2
2602	Q94A52	149:261	0.54	0.26	0.12	2
2603	P38111	1093:1165	0.59	0.30	0.13	3
2604	P38111	1093:1182	0.57	0.27	0.13	3
2605	P38111	1093:1162	0.57	0.27	0.13	3
2606	P38111	1092:1166	0.59	0.29	0.13	3
2607	P38111	1093:1168	0.58	0.29	0.13	3
2608	P38111	1091:1164	0.59	0.30	0.13	3
2609	P38111	1089:1164	0.58	0.29	0.13	3

Table 8B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutritrity	Human Homology
2519	-15.79	0.80	0.31	0.00	1.00	0.67
2520	-16.50	0.84	0.29	0.00	1.00	0.62
2521	-21.97	1.00	0.30	0.00	0.25	0.39
2522	-28.44	0.16	0.36	0.00	0.00	0.84
2523	-36.33	0.16	0.33	0.00	0.00	0.87

2524	-27.36	0.49	0.34	0.00	0.30	0.41
2525	-17.59	0.99	0.20	0.00	1.00	0.64
2526	-17.98	0.90	0.63	0.00	0.00	0.44
2527	-16.59	0.95	0.63	0.00	0.32	0.44
2528	-16.79	0.87	0.63	0.00	0.33	0.44
2529	-18.37	0.77	0.63	0.00	0.30	0.42
2530	-18.37	0.71	0.63	0.00	0.32	0.40
2531	-24.84	0.25	0.63	0.26	0.00	0.35
2532	-26.10	0.22	0.63	0.26	0.00	0.33
2533	-15.45	0.39	0.32	1.00	0.34	0.34
2534	-19.21	0.44	0.99	0.00	0.00	0.96
2535	-25.30	0.46	0.22	0.33	0.34	1.00
2536	-22.49	0.43	0.21	0.31	0.35	0.94
2537	-22.06	0.51	0.21	0.28	0.27	1.00
2538	-21.01	0.41	0.31	0.00	0.28	0.94
2539	-23.16	0.43	0.28	0.00	0.00	0.98
2540	-15.02	1.17	0.22	0.34	0.35	0.75
2541	-10.60	1.38	0.22	0.00	0.34	0.28
2542	-18.91	0.89	0.20	0.00	0.32	0.94
2543	-31.57	0.29	0.21	0.00	0.31	0.88
2544	-17.13	0.77	0.20	0.31	0.28	0.61
2545	-16.98	0.76	0.19	0.25	0.17	0.52
2546	-20.09	0.24	0.21	0.29	0.30	0.43
2547	-19.93	0.52	0.22	0.32	0.32	0.33
2548	-16.57	0.79	0.58	0.33	0.24	0.59
2549	-16.05	0.65	0.20	0.28	0.31	0.65
2550	-17.15	0.83	0.30	0.30	0.00	0.84
2551	-21.64	0.55	0.26	0.26	0.26	0.00
2552	-24.45	0.54	0.32	0.24	0.25	0.91
2553	-23.71	0.37	0.30	0.27	0.29	0.93
2554	-22.23	0.52	0.74	0.00	0.19	0.44
2555	-18.41	0.58	0.29	0.29	0.25	0.78
2556	-17.59	0.58	0.60	0.00	0.23	0.98
2557	-19.98	0.57	0.31	0.00	0.24	0.00
2558	-17.02	0.48	0.53	0.00	0.22	0.88

2559	-5.92	1.63	0.30	0.00	0.27	0.61
2560	-6.10	1.53	0.21	0.00	0.26	0.40
2561	-18.69	0.54	0.18	0.24	0.35	0.84
2562	-17.31	0.54	0.20	0.24	0.33	0.84
2563	-17.40	0.47	0.20	0.22	0.34	0.84
2564	-22.65	0.39	0.32	0.25	0.25	0.76
2565	-22.50	0.39	0.32	0.25	0.25	0.76
2566	-22.35	0.39	0.32	0.25	0.26	0.76
2567	-22.74	0.39	0.32	0.00	0.25	0.76
2568	-22.59	0.39	0.32	0.00	0.25	0.75
2569	-22.97	0.38	0.32	0.24	0.25	0.76
2570	-23.34	0.38	0.32	0.24	0.25	0.75
2571	-23.25	0.38	0.32	0.24	0.25	0.75
2572	-23.61	0.37	0.32	0.00	0.25	0.75
2573	-24.11	0.36	0.32	0.23	0.26	0.75
2574	-33.59	0.15	0.34	0.25	0.00	1.00
2575	-36.26	0.21	0.21	0.29	0.33	0.92
2576	-34.27	0.27	0.22	0.25	0.00	0.90
2577	-32.73	0.22	0.25	0.31	0.00	0.87
2578	-34.50	0.23	0.21	0.00	0.00	0.89
2579	-34.08	0.25	0.26	0.00	0.00	0.91
2580	-34.67	0.22	0.23	0.25	0.00	0.91
2581	-33.39	0.29	0.20	0.00	0.31	0.88
2582	-37.57	0.19	0.25	0.31	0.00	0.92
2583	-35.42	0.21	0.24	0.21	0.00	0.90
2584	-33.68	0.26	0.23	0.00	0.00	0.92
2585	-37.47	0.23	0.25	0.00	0.00	0.92
2586	-33.38	0.20	0.31	0.00	0.00	0.87
2587	-32.78	0.18	0.33	0.27	0.29	0.88
2588	-36.89	0.16	0.35	0.00	0.00	0.92
2589	-37.66	0.15	0.39	0.00	0.00	0.93
2590	-29.21	0.16	0.25	0.00	0.27	1.00
2591	-30.24	0.12	0.33	0.00	0.00	0.99
2592	-16.02	0.94	0.32	0.00	1.00	0.63
2593	-16.69	0.94	0.30	0.00	1.00	0.63

2594	-17.69	0.65	0.31	0.00	1.00	0.62
2595	-34.37	0.07	0.29	0.00	0.00	0.98
2596	-38.56	0.08	0.24	0.00	0.00	0.98
2597	-41.59	0.04	0.26	0.00	0.00	0.98
2598	-40.29	0.04	0.23	0.28	0.00	0.98
2599	-35.70	0.11	0.36	0.26	0.22	0.99
2600	-34.97	0.13	0.40	0.23	0.22	0.93
2601	-26.91	0.45	0.18	0.30	0.22	0.31
2602	-25.52	0.46	0.29	0.00	0.25	0.81
2603	-23.00	0.80	0.27	0.00	0.24	0.26
2604	-20.82	0.80	0.27	0.00	0.25	0.27
2605	-23.98	0.70	0.25	0.27	0.25	0.27
2606	-22.45	0.79	0.27	0.27	0.25	0.25
2607	-22.10	0.80	0.27	0.28	0.24	0.25
2608	-22.75	0.77	0.28	0.31	0.25	0.25
2609	-23.28	0.71	0.28	0.28	0.25	0.24

**Example 7: Scaled Up Production of Recombinant Nutritive Proteins**

[00282] A representative protocol for producing quantities of nutritive proteins as described in this disclosure is as follows

[00283] 5ml LB with 100 mg/l carbenicillin (in a 50ml baffled Pyrex shake flask) is inoculated with a stab from the glycerol stock of a recombinant *E. coli* strain comprising a recombinant gene encoding a nutritive protein and grown until late exponential phase (OD<sub>600nm</sub> ≈ 2) at 37°C and 250rpm. A 2.5l Ultra Yield Flask (Thomson Instrument Company) is inoculated with 500ml sterile water and enough EnBase EnPresso™ tablets (BioSilta) to formulate 500ml growth medium. This medium is supplemented with 100 mg/l carbenicillin, 0.001% Industrol 204 antifoam, and 0.6 U/l EnzI'm (BioSilta). The shake flask is inoculated to OD<sub>600nm</sub> = 0.05 and grown 16hr at 30°C and 250rpm. At this point, OD<sub>600nm</sub> ≈ 10 had been reached, the growth medium is further supplemented with EnPresso™ Booster tablets (BioSilta), 1.2 U/l EnzI'm, and 1mM IPTG to induce heterologous protein production. After another 8-24hr of shaking at 30°C and 250rpm, the flask is harvested by centrifugation at 10,000rpm, the supernatant is decanted, and the wet cell weight is measured. At this point, approximately 20gWCW (grams wet cell weight)/l medium is recovered.

[00284] The harvested cells from each shake-flask fermentation are suspended in 25 mL of IMAC Equilibration Solution (30 mM Imidazole, 50 mM Phosphate, 0.5 M NaCl, pH 7.5). The suspended cells are then lysed by sonication at 90 Watts for 90 seconds, on ice. The lysed cells are centrifuged at 15,500 RCF for 60 minutes, and decanted. The cell debris 5 is discarded, and the supernatants are 0.2  $\mu$ m filtered. Filters are then flushed with an additional 10 mL of IMAC Equilibration Solution. These filtered protein solutions are then purified by immobilized metal affinity chromatography (IMAC).

[00285] IMAC resin (GE Healthcare, IMAC Sepharose 6 Fast Flow) is charged with nickel and equilibrated. 30 mL of each protein solution is loaded onto a 5 mL IMAC column, 10 and washed with additional equilibration solution to remove unbound impurities. The protein of interest is then eluted with 15 mL of 0.5 M NaCl, 0.2 M Imidazole, pH 7.5. The purified proteins are shown to be 90% pure, by SDS-PAGE. Approximately 20 to 60 mg of each protein is recovered in the IMAC elution fractions. Each IMAC elution fraction is buffer exchanged by dialysis into a formulation solution (20 mM HEPES, pH 7.5). After buffer 15 exchange, these protein solutions are recovered for all downstream processing.

#### **Example 8: Prediction of Soluble Expression of Nutritive Proteins**

[00286] Open reading frames encoding a set of 292 nutritive proteins were cloned and introduced into *E. coli* to assess recombinant protein expression using the method of Example 6. In the system used, 163 proteins were identified as expressed while 129 were not. Of the 20 163 proteins that expressed, 125 were tested for soluble expression. It was found that 75 were solubly expressed while 50 were not.

[00287] Figure 1 shows a two dimensional histogram of protein expression in the *E. coli* expression screen. Figure 1 shows the relative likelihood (on a log scale) of a protein being expressed as a function of solvation score (y-axis) and aggregation score (x-axis). A 25 darker mark on the histogram indicates a higher number of proteins expressed, while a lighter mark indicates a fewer number of proteins expressed. Figure 1 shows that those proteins that were successfully expressed tend to cluster in the top left region of the plot, where the solvation score is more negative ( $\leq -20$ ) and the aggregation score is smaller ( $\leq 0.75$ ). There were few examples of proteins that were successfully expressed with less negative solvation 30 scores ( $\geq -15$ ) and large aggregation scores ( $\geq 1$ ). This result suggests that nutritive proteins with solvation scores of -20 or less and aggregation scores of 0.75 or less are more likely to be expressed in this system.

[00288] Figure 2 shows a two dimensional histogram of the number of soluble protein expression in the *E. coli* expression screen. Figure 2 shows the relative likelihood (on a log scale) of a protein being solubly expressed as a function of solvation score (y-axis) and aggregation score (x-axis). Again, a darker mark on the histogram indicates a higher number of proteins expressed, while a lighter mark indicates a fewer number of proteins expressed.

5 Figure 1 shows that those proteins that were expressed solubly tended to cluster in the top left region of the plot, where the solvation score is more negative ( $\leq -20$ ) and the aggregation score is smaller ( $\leq 0.5$ ). There were few examples of proteins that were expressed solubly with less negative solvation scores ( $\geq -15$ ) and large aggregation scores ( $\geq 0.75$ ). This result

10 suggests that nutritive proteins with solvation scores of -20 or less and aggregation scores of 0.5 or less are more likely to be solubly expressed in this system.

#### **Example 9: Solubility Screening**

[00289] The solubility of nine nutritive proteins produced as described Examples 6 and 7 was examined by centrifuge concentration followed by protein concentration assays.

15 Samples in 20 mM HEPES pH 7.5 were tested for protein concentration according to the protocol for Coomassie Plus (Bradford) Protein Assay (Thermo Scientific) and absorbance at 280 nm (if applicable). Based on these measurements 10 mg of protein was added to an Amicon Ultra 3 kDa centrifugal filter (Millipore). Samples were concentrated by centrifugation at 10,000 Xg for 30 minutes. The final concentrated samples were examined

20 for precipitated protein and color, and then tested for protein concentration as described above. The results are shown in Table 9.

**Table 9**

Seq ID No	Appearance	Concentration (g/L)
2580	Clear Faint Yellow	44
2582	Clear Faint Yellow	166
2587	Clear Colorless	107
2595	Clear Colorless	60
2596	Clear Colorless	29
2598	Clear Colorless	151
2599	Clear Colorless	207
2600	Clear Yellow	95

2603	Clear Yellow	191
------	--------------	-----

[00290] The solubilities of these nutritive proteins were found to be significantly higher than concentrations typically found for whey (12.5 g/L) and soy (10 g/L) (Pelegrine, D. H. G. & Gasparetto, C. A., 2005. Whey proteins solubility as function of temperature and pH. LWT - Food Science and Technology, p. 77–80; Lee, K. H., Ryu, H. S. & and Rhee, K. C., 2003. Protein solubility characteristics of commercial soy protein products. Journal of the American Oil Chemists' Society, pp. 85-90). This demonstrates the usefulness of the nutritive proteins disclosed herein. For example, the solubility of nutritive proteins may improve compliant delivery of high quality protein in as small of a volume as possible while avoiding the “chalkyness” that often characterizes proteins delivered in this manner. This may, for example, be useful to deliver proteins to the elderly or other subjects.

#### **Example 10: Stability Screening**

[00291] Thermal stability of nutritive proteins provides insight regarding whether the protein is likely to have a useful shelf life. Samples of proteins produced as described in Examples 6 and 7 were screened in parallel using a rapid thermal stability screening method. In this method proteins were heated slowly from 25°C to 95°C in two representative formulations in the presence of a hydrophobic dye (Enzo Life Sciences, ProteoStat® Thermal shift stability assay kit) that binds to aggregated proteins that form as the protein denatures with increasing temperature (Niesen, F. H., Berglund, H. & Vadadi, M., 2007. The use of differential scanning fluorimetry to detect ligand interactions that promote protein stability. *Nature Protocols*, Volume 2, pp. 2212-2221.). Upon binding, the dye's fluorescence increases significantly, which is then recorded by the rtPCR instrument and represented as the protein's melting curve (Lavinder, J. J., Hari, S. B., Suillivan, B. J. & Magilery, T. J., 2009. High-Throughput Thermal Scanning: A General, Rapid Dye-Binding Thermal Shift Screen for Protein Engineering. *Journal of the American Chemical Society*, pp. 3794-3795.). After the thermal shift is complete samples were examined for insoluble precipitates and further analyzed by analytical size exclusion chromatography (SEC).

[0043] Protein solutions (12.5 mg/ml) were prepared in both PBS and 20 mM HEPES pH 7.7 buffers, each containing 1X ProteoStat TS Detection Reagent. Samples of each solution were heated slowly from 25°C – 95°C, 0.5°C/30 seconds using a real-time PCR (rtPCR) thermocycler while monitoring the fluorescence of the dye. From this thermal scan

the temperature of aggregation was determined ( $T_{agg}$ ) from the temperature with the strongest slope if an increase in fluorescence was observed. To supplement the assay, samples were taken before and after the thermal shift and analyzed by SEC (GE Healthcare - Superdex 75 5/150) which can detect large soluble aggregates. The results for seven nutritive proteins of 5 this disclosure and a whey standard are presented in Table 10. The presence of soluble aggregates detected by SEC is noted by a “yes” if observed or “no” if not observed, and the “n/a” entries for the whey standard indicate the production of insoluble precipitates such that no SEC analysis was performed.

**Table 10**

10

Seq ID No	HEPES - $T_{agg}$	PBS - $T_{agg}$	HEPES - SEC Agg?	PBS - SEC Agg?
2582	95	95	No	No
2587	95	95	No	No
2596	45	95	No	No
2598	95	95	No	No
2599	95	95	No	No
2600	95	95	Yes	Yes
2603	95	95	Yes	Yes
whey	79	81.5	n/a	n/a

[00292] As shown in Table 10, SEQ ID NOS: 2582, 2587, 2596, 2598, and 2599 had much higher HEPES  $T_{agg}$  than whey (in fact, the proteins did not form any aggregates at 95°C, which was the upper limit of the assay), and thus are expected to be more stable than 15 whey. The SEQ ID NOS: 2600 and 2603 proteins had higher HEPES  $T_{agg}$  and PBS  $T_{agg}$  values than whey, but both formed soluble aggregates in solution while whey formed insoluble aggregates that precipitated from solution. Thus, even in an aggregated state, these two nutritive proteins would be expected to remain in solution for a greater period of time than whey.

**20 Example 11: Digestibility Screening – Determination of Digestion Half-Life**

[00293] The goal of screening for protein digestibility is to eliminate potentially unsafe allergenic proteins and to determine the relative completeness of digestion as a predictor of peptide bioavailability. This screening method utilizes a physiologically relevant *in vitro* digestion reaction that includes both phases of protein digestion, simulated gastric digestion 25 and simulated intestinal digestion (Moreno, J. F. et al., 2005. Stability of the major allergen

Brazil nut 2S albumin (Ber e 1) to physiologically relevant in vitro gastrointestinal digestion. FEBS Journal, pp. 341-352.). Samples can be taken throughout the reaction and analyzed for intact protein and peptide fragments using chip electrophoresis and LC-QTOF-MS. Proteins with allergenic properties can be assessed by identifying proteins or large fragments of

5 proteins that are resistant to digestive proteases and thus have a higher risk of causing an allergenic reaction (Goodman, R. E. et al., 2008. Allergenicity assessment of genetically modified crops - what makes sense?. Nature Biotechnology, pp. 73-81.). Digestibility is measured by determining how efficiently the protein is broken down into peptides (Daniel, H., 2003. Molecular and Integrative Physiology of Intestinal Peptide Transport. Annual

10 Review of Physiology, Volume 66, pp. 361-384.).

[0044] The method used an automated assay for in vitro digestions of proteins wherein assay conditions and protease concentrations are physiologically relevant (Moreno, F. J., Mackie, A. R. & Clare Mills, E. N., 2005. Phospholipid interactions protect the milk allergen  $\alpha$ -Lactalbumin from proteolysis during in vitro digestion. Journal of agricultural and food chemistry, pp. 9810-9816; Martos, G., Contreras, P., Molina, E. & Lopez-Fandino, R., 2010. Egg White Ovalbumin Digestion Mimicking Physiological Conditions. Journal of Agricultural and food chemistry, pp. 5640-5648; Moreno, J. F. et al., 2005. Stability of the major allergen Brazil nut 2S albumin (Ber e 1) to physiologically relevant in vitro gastrointestinal digestion. FEBS Journal, pp. 341-352.). The first phase of digestion is in

15 simulated gastric fluid (SGF) and formulated at pH 1.5 and with a pepsin:substrate ratio of (1:10 w/w). The second phase of digestion is in simulated intestinal fluid (SIF) is formulated with bile salts at pH 6.5 and with an trypsin:chymotrypsin:substrate ratio of (1:4:400 w/w). The protein is treated for 120 mins in the simulated gastric fluid, which is how long it takes for 90% of a liquid meal to pass from the stomach to the small intestine (Kong, F. & Singh,

20 R. P., 2008. Disintegration of Solid Foods in Human Stomach. *Journal of Food Science*, pp. 67-80), and then treated with simulated intestinal fluid for 120 mins. Sample time points are taken throughout both reactions and quenched for analysis. Bovine serum albumin, which is readily digested by pepsin, is the positive control for the SGF solution, and beta-lactoglobulin, which is naturally resistant to pepsin but digested in SIF, is the positive control

25 for SIF solution. Intact protein and large fragments were detected using electrophoresis. For chip electrophoresis, a Caliper Labchip GXII equipped with a HT Low MW Protein Assay Kit was used to monitor the size and amount of intact protein as well as any digestion fragments larger than 4 kDa. By monitoring the amount of intact protein observed over time,

the half-life ( $\tau_{1/2}$ ) of digestion was calculated for SGF and, if intact protein is detected after SGF digestion, in SIF.

[0045] This method was used to analyze the digestion half-lives of thirteen nutritive proteins of this disclosure (SEQ ID NOS: 2547, 2575, 2578, 2580, 2582, 2587, 2595, 2596, 2598, 2599, 2600, 2602, and 2603) produced as described in Examples 6 and 7, as well as native and recombinant ovalbumin (OVA and rOVA, respectively; SEQ ID NO: 2610) and beta-lactoglobulin (BLG and rBLG, respectively; SEQ ID NO: 2611) proteins and a whey standard. The results of these experiments are summarized in Table 11. An “n/a” entry in the Simulated Intestinal Fluid field indicates that no intact protein was detected after SGF digestion.

**Table 11**

Seq ID No.	Digestion $\tau_{1/2}$ (min.)	
	Simulated Gastric Fluid	Simulated Intestinal Fluid
2547	0.3	n/a
2575	0.3	n/a
2578	2	n/a
2580	6	n/a
2582	0.5	n/a
2587	0.7	n/a
2595	10	n/a
2596	0.6	n/a
2598	0.7	n/a
2599	0.3	n/a
2600	29	1
2602	1	n/a
2603	6	n/a
BLG (2611)	77	4
rBLG (2611)	50	0.7
OVA (2610)	18	1
rOVA (2610)	5	n/a
whey	99	4

[0046] The results shown in Table 11 indicate that twelve of the thirteen nutritive proteins of this disclosure were all completely digested by SGF and have SGF half lives of ten minutes or less. By comparison whey is not completely digested by SGF and has an SGF half life of 99 minutes and a SIF half life of 4 minutes. This study suggests that the nutritive 5 proteins of this disclosure are likely to be readily digested and not likely to elicit an allergic response when ingested.

[00294] The results in Table 11 also show that the recombinant beta-lactoglobulin and ovalbumin produced according to this disclosure were both more readily digested than their naturally-occurring counterparts. The speed in which a protein is broken down can be 10 controlled by selecting for properties that improve or limit accessibility of the gastrointestinal proteases. This capability can be demonstrated for two typical protein properties, glycosylation and disulfide cross-linking. Like many naturally occurring proteins, naturally occurring OVA and BLG are glycosylated by their host organisms. In contrast, the recombinant proteins produced according to the present disclosure are not glycosylated 15 because the host organism (*E. coli* in this case) does not glycosylate. The lack of glycosylation in recombinant nutritive proteins according to this disclosure may result in proteins that are more readily digested. Furthermore, BLG has four disulfide bonds that are known to slow down or interfere with digestion. When these disulfide bonds are disrupted, the rate of digestion increases (Reddy, I. M., Kella, N. K. D. & Kinsella, J. E., 1988. 20 Structural and conformational Basis of the Resistance of b-Lactoglobulin to Peptic and Chymotryptic Digestion. *J. Agric. Food Chem.*, Volume 36, pp. 737-741). A lack or disruption of disulfide bond formation in recombinant nutritive proteins according to this disclosure may result in proteins that are more readily digested.

#### **Example 12: Digestibility Screening – Analysis of Digestion Products**

25 [00295] Two nutritive proteins produced as described in Examples 6 and 7 (SEQ ID NOS: 2612 and 2613) were subjected to SGF and SIF digestion as described in Example 11. Both proteins were completely digested in SGF, and the SGF half lives are shown in Table 12.

**Table 12**

Seq ID No	Digestion $\tau_{1/2}$ (min.)	
	Simulated Gastric Fluid	Simulated Intestinal Fluid
2612	0.7	n/a
2613	6	n/a

[00296] To detect and identify peptides that were present after SGF and SIF digestion, samples of the SGF and SIF digests were analyzed by LC/Q-TOF MS/MS. Samples from the SGF digests were directly analyzed by LC/Q-TOF MS/MS, while SIF protein digestions required purification by SCX to remove bile acids before detection and identification by LC/Q-TOF MS/MS. Peptides were extracted from the chromatograms and identified using Bioconfirm Software (Aglient). The sequence assignment of peptides were based on accurate mass match ( $\pm 10$  ppm) and further confirmed by MS/MS fragmentation. The results are shown in Tables 13 and 14 below.

10

**Table 13**  
**(SEQ ID NO: 2612)**

SGF Peptides 120 min	SEQ ID NO	SIF Peptides 120 min	SEQ ID NO
LL		SE	
LAL		PSE	
HVL		HVL	
LEL		FKV	
LALA	2616	HQI	
LLLD	2617	PSEA	2618
IAEF	2619	REV	
IQQF	2620	FDK	
YDKL	2621	AEFK	2622
SNLTE	2623	LKHV	2624
ELLEA	2625	SSSEL	2626
EELAL	2627	FKVF	2628
DDLLL	2629	AELKH	2630
LAYDK	2631	LKHVL	2632
TKTRL	2633	FKEAF	2634
DLDHQ	2635	NGSISSS	2636

GTLENL	2637	SLGLSPS	2638
EKLTDA	2639	GEKLTD	2640
AEVDDM	2641	ELATVM	2642
LDDLLL	2643	GSGEINI	2644
LDLDHQ	2645	RSLGLSP	2646
DLKKKL	2647	ELKHVL	2648
KTKTRL	2649	KLTDAEV	2650
SQRLEE	2651	FKVFDK	2652
AEVDDML	2653	AEVDDML	2654
QQELDDL	2655	DAEVDDM	2656
LEKTKTR	2657	AELKHVL	2658
LEKTKTR	2659	HVLTSIGE	2660
GTLENLEE	2661	SIGEKLT	2662
QQELDDLL	2663	RSLGLSPSE	2664
KLEKTKTRLQ	2665	VLTSIGEKL	2666
SRQLKSNDSEQ	2667	TSIGEKLT	2668
EKTKTRLQQEL	2669	HVLTSIGEK	2670
YDKLEKTKTRL	2671	AELKHVLTS	2672
LAYDKLEKTKTRL	2673	RSLGLSPSEA	2674
EELKKKLLKDDEL	2675	SSNLTEEQIA	2676
EELKKKLLKDLELL	2677	RSLGLSPSEAE	2678
AELKHVLTSIGEKLT	2679	KVFDKNGDGLISA	2680
MGSHHHHHHHHSSNL	2681	FDKDNNGSISSSEL	2682
LREVSDGSGEINIQQF	2683	REVSDGSGEINIQQ	2684
AAELKHVLTSIGEKLT	2685	DVDGNHQIEFSEF	2686
AELKHVLTSIGEKLTDAE	2687	LREVSDGSGEINIQQ	2688
AYDKLEKTKTRLQQEL	2689	REVSDGSGEINIQQF	2690
AAELKHVLTSIGEKLTDAE	2691	LREVSDGSGEINIQQFAALLS	2692
LENLEELKKKLLKDDEL	2693		
KLEKTKTRLQQELDDL	2694		
DKLEKTKTRLQQELDDL	2695		
LAYDKLEKTKTRLQQELDDL	2696		
LALAYDKLEKTKTRLQQELDDL	2697		

**Table 14**  
**(SEQ ID NO: 2613)**

SGF Peptides 120 min	SEQ ID NO	SIF Peptides 120 min	SEQ ID NO
GVL		TKH	
ALL		INDI	2698
LVL		HLVL	2699
IGVL	2700	TIKF	2701
TIKF	2702	IGVLD	2703
TIKF	2704	RNLD	2705
EVYDL	2706	IGVLDV	2707
LNDSVQ	2708	QTIKF	2709
IWVIND	2710	VQTIKF	2711
DLNDSVQ	2712	SVQTIKF	2713
SVQTIKF	2714	KCAKCISMIGVL	2715
HHLVLGALLD	2716	EKCAKCISMIGV	2717
HHHHHHHLVL	2718	HEFKRTTYSE	2719
HHHHHHHHHLVL	2720	SHKFRNLDKDL	2721
DVTKHEFKRTTY	2722	ISMIGVLDVTKHE	2723
SHHHHHHHHLVL	2724		
TKHEFKRTTYSEN	2725		
GSHHHHHHHHLVLG	2726		
MGSHHHHHHHHHLVL	2727		
KRTTYSENEVYDLN	2728		

5 [00297] As can been seen in Tables 13 and 14, each protein was digested into multiples smaller peptide fragments ranging in size from 2 to 22 amino acids (SEQ ID NO: 2612) or 2 to 13 amino acids (SEQ ID NO: 2613). None of these peptide fragments was found to be homologous to any known allergen.

10 [00298] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular

situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

**CLAIMS:**

1. An isolated nutritive protein comprising a first polypeptide sequence that is homologous to a fragment of a naturally-occurring protein, wherein the first polypeptide sequence comprises at least one of:
  - a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;
  - b. a ratio of Leu residues to total amino acid residues of at least 11%; and
  - c. a ratio of essential amino acid residues to total amino acid residues of at least 10 49%.
2. The isolated nutritive protein of claim 1, wherein the first polypeptide sequence further comprises at least one of each essential amino acid.
- 15 3. The isolated nutritive protein of claim 1 or 2, wherein the first polypeptide sequence comprises at least 70% homology to the fragment of a naturally-occurring protein.
4. The isolated nutritive protein of claim 1 or 2, wherein the first polypeptide sequence comprises at least 95% homology to the fragment of a naturally-occurring protein.
- 20 5. The isolated nutritive protein of any one of claims 1 to 4, wherein the fragment of a naturally-occurring protein comprises at least 25 amino acid residues.
- 25 6. The isolated nutritive protein of any one of claims 1 to 4, wherein the fragment of a naturally-occurring protein comprises at least 50 amino acid residues.
7. The isolated nutritive protein of any one of claims 1 to 6, wherein the first polypeptide sequence comprises:
  - a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;
  - b. a ratio of Leu residues to total amino acid residues of at least 11%; and
  - c. a ratio of essential amino acid residues to total amino acid residues of at least 30 49%.

8. The isolated nutritive protein of claim any one of claims 1 to 7, wherein the first polypeptide sequence comprises the fragment of a naturally-occurring nutritive protein.

5 9. The isolated nutritive protein of any one of claims 1 to 7, wherein the first polypeptide sequence consists of the fragment of a naturally-occurring nutritive protein.

10. The isolated nutritive protein of any one of claims 1 to 9, wherein the first polypeptide sequence is not an allergen.

10 11. The isolated nutritive protein of claim 10, wherein the first polypeptide sequence has less than 50% global homology to a known allergen.

15 12. The isolated nutritive protein of any one of claims 1 to 11, wherein the first polypeptide sequence is not a toxin.

13. The isolated nutritive protein of claim 12, wherein the first polypeptide sequence has less than 50% global homology to a known toxin.

20 14. The isolated nutritive protein of any one of claims 1 to 13, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 60 minutes.

15. The isolated nutritive protein of claim 14, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 30 minutes.

25 16. The isolated nutritive protein of claim 15, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 10 minutes.

17. The isolated nutritive protein of any one of claims 1 to 16, wherein the first 30 polypeptide sequence is completely digested in simulated gastric fluid.

18. The isolated nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

19. The isolated nutritive protein of any one of claims 1 to 18, wherein the first polypeptide sequence comprises no cysteine residues.

5 20. The isolated nutritive protein of any one of claims 1 to 19, wherein the first polypeptide sequence comprises no disulfide bonds.

21. The isolated nutritive protein of any one of claims 1 to 20, wherein the first polypeptide sequence does not comprise N-linked glycosylation.

10 22. The isolated nutritive protein of any one of claims 1 to 21, wherein the first polypeptide sequence does not comprise O-linked glycosylation.

15 23. The isolated nutritive protein of any one of claims 1 to 22, wherein the first polypeptide sequence is resistant to aggregation.

24. The isolated nutritive protein of any one of claims 1 to 23, wherein the first polypeptide sequence is anionic at pH 7.

20 25. The isolated nutritive protein of any one of claims 1 to 24, wherein the first polypeptide sequence has an aqueous solubility at pH 7 of at least 12.5 g/L.

26. The isolated nutritive protein of any one of claims 1 to 25, wherein the first polypeptide sequence has a calculated solvation score of -20 or less.

25 27. The isolated nutritive protein of any one of claims 1 to 26, wherein the first polypeptide sequence has a calculated aggregation score of 0.75 or less.

28. The isolated nutritive protein of any one of claims 1 to 27, wherein the first 30 polypeptide sequence has a calculated aggregation score of 0.5 or less.

29. The isolated nutritive protein of any one of claims 1 to 28, wherein the first polypeptide sequence comprises an amino acid sequence selected from:

i. an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609;

- ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and
- iii. a mutein of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

5

30. The isolated nutritive protein of any one of claims 1 to 29, wherein the first polypeptide sequence consists of an amino acid sequence selected from:

- i. an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**;
- ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and
- iii. a mutein of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

10

31. The isolated nutritive protein of any one of claims 1 to 30, wherein the first polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

20

32. The isolated nutritive protein of any one of claims 1 to 31, wherein the first polypeptide sequence is a fragment of at least 25 amino acids of the naturally-occurring nutritive protein.

25

33. The isolated nutritive protein of any one of claims 1 to 32, wherein the first polypeptide sequence is a fragment of at least 50 amino acids of the naturally-occurring nutritive protein.

34. The isolated nutritive protein of any one of claims 1 to 33, wherein the isolated nutritive protein consists of the first polypeptide sequence.

30

35. The isolated nutritive protein of any of claims 1 to 34, wherein the isolated nutritive protein further comprises a polypeptide tag for affinity purification.

36. The isolated nutritive protein of claim 35, wherein the tag for affinity purification is a polyhistidine-tag.

37. An isolated nutritive protein comprising a first polypeptide sequence that is 5 homologous to a fragment of a naturally-occurring protein, wherein the isolated nutritive protein comprises at least one of:

- a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;
- b. a ratio of Leu residues to total amino acid residues of at least 11%; and
- c. a ratio of essential amino acid residues to total amino acid residues of at least 10 49%.

38. The isolated nutritive protein of claim 37, wherein the isolated nutritive protein further comprises at least one of each essential amino acid.

15

39. The isolated nutritive protein of claim 37 or 38, wherein the isolated nutritive protein comprises at least 70% homology to the fragment of a naturally-occurring protein.

20

40. The isolated nutritive protein of claim 37 or 38, wherein the isolated nutritive protein comprises at least 95% homology to the fragment of a naturally-occurring protein.

41. The isolated nutritive protein of any one of claims 37 to 40, wherein the fragment of a naturally-occurring protein comprises at least 25 amino acid residues.

25

42. The isolated nutritive protein of any one of claims 37 to 40, wherein the fragment of a naturally-occurring protein comprises at least 50 amino acid residues.

43. The isolated nutritive protein of any one of claims 37 to 42, wherein the isolated nutritive protein comprises:

30

- a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;
- b. a ratio of Leu residues to total amino acid residues of at least 11%; and
- c. a ratio of essential amino acid residues to total amino acid residues of at least 49%.

44. The isolated nutritive protein of any one of claims 37 to 43, wherein the isolated nutritive protein comprises a fragment of a naturally occurring nutritive protein.

5 45. The isolated nutritive protein of any one of claims 37 to 43, wherein the isolated nutritive protein consists of a fragment of a naturally occurring nutritive protein.

46. The isolated nutritive protein of any one of claims 37 to 45, wherein the isolated nutritive protein is not an allergen.

10

47. The isolated nutritive protein of claim 46, wherein the isolated nutritive protein has less than 50% global homology to a known allergen.

15

48. The isolated nutritive protein of any one of claims 37 to 47, wherein the isolated nutritive protein is not a toxin.

49. The isolated nutritive protein of claim 48, wherein the isolated nutritive protein has less than 50% global homology to a known toxin.

20

50. The isolated nutritive protein of any one of claims 37 to 49, wherein the isolated nutritive protein has a simulated gastric digestion half-life of less than 60 minutes.

51. The isolated nutritive protein of claim 50, wherein the isolated nutritive protein has a simulated gastric digestion half-life of less than 30 minutes.

25

52. The isolated nutritive protein of claim 51, wherein the isolated nutritive protein has a simulated gastric digestion half-life of less than 10 minutes.

30

53. The isolated nutritive protein of any one of claims 37 to 52, wherein the isolated nutritive protein is completely digested in simulated gastric fluid.

54. The isolated nutritive protein of any one of claims 37 to 53, wherein the isolated nutritive protein comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

55. The isolated nutritive protein of any one of claims 37 to 54, wherein the isolated nutritive protein comprises no cysteine residues.

5 56. The isolated nutritive protein of any one of claims 37 to 55, wherein the isolated nutritive protein comprises no disulfide bonds.

57. The isolated nutritive protein of any one of claims 37 to 56, wherein the isolated nutritive protein does not comprise N-linked glycosylation.

10

58. The isolated nutritive protein of any one of claims 37 to 57, wherein the isolated nutritive protein does not comprise O-linked glycosylation.

15

59. The isolated nutritive protein of any one of claims 37 to 58, wherein the isolated nutritive protein is resistant to aggregation.

60. The isolated nutritive protein of any one of claims 37 to 59, wherein the isolated nutritive protein is anionic at pH 7.

20

61. The isolated nutritive protein of any one of claims 37 to 60, wherein the isolated nutritive protein has an aqueous solubility at pH 7 of at least 12.5 g/L.

62. The isolated nutritive protein of any one of claims 37 to 61, wherein the isolated nutritive protein has a calculated solvation score of -20 or less.

25

63. The isolated nutritive protein of any one of claims 37 to 62, wherein the isolated nutritive protein has a calculated aggregation score of 0.75 or less.

30

64. The isolated nutritive protein of any one of claims 37 to 63, wherein the isolated nutritive protein has a calculated aggregation score of 0.5 or less.

65. The isolated nutritive protein of any one of claims 37 to 64, wherein the isolated nutritive protein comprises an amino acid sequence selected from:

- i. an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609;

- ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and
- iii. a mutein of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

5

66. The isolated nutritive protein of any one of claims 37 to 65, wherein the isolated nutritive protein consists of an amino acid sequence selected from:
- i. an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**;
  - ii. a modified derivative of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**; and
  - iii. a mutein of an amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

10

67. The isolated nutritive protein of any one of claims 37 to 66, wherein the isolated nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from **SEQ ID NO: 1 to SEQ ID NO: 2609**.

20

68. The isolated nutritive protein of any one of claims 37 to 67, wherein the isolated nutritive protein further comprises a polypeptide tag for affinity purification.

69. The isolated nutritive protein of claim 68, wherein the tag for affinity purification is a polyhistidine-tag.

25

70. An isolated nucleic acid comprising a nucleic acid sequence that encodes a nutritive protein according to any one of claims 1 to 69.

30

71. The isolated nucleic acid of claim 70, wherein the isolated nucleic acid is selected from genomic DNA, cDNA, sense RNA and antisense RNA.

72. The isolated nucleic acid of claim 71, wherein the isolated nucleic acid is genomic DNA.

73. The isolated nucleic acid of claim 71, wherein the isolated nucleic acid is cDNA.

74. The isolated nucleic acid of any one of claims 70 to 73, further comprising an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein.

75. A vector comprising a nucleic acid sequence that encodes a nutritive protein of any one of claims 1 to 69.

10

76. The vector of claim 75, further comprising an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein

15

77. A recombinant microorganism comprising at least one of a nucleic acid of any one of claims 70 to 74 and a vector of any one of claims 75 and 76.

78. The recombinant microorganism according to claim 77, wherein the recombinant microorganism is a prokaryote.

20

79. The recombinant prokaryote according to claim 78, wherein the prokaryote is heterotrophic.

80. The recombinant prokaryote according to claim 78, wherein the prokaryote is autotrophic.

25

81. The recombinant prokaryote according to claim 78, wherein the prokaryote is a bacteria.

30

82. A method of making a nutritive protein according to any one of claims 1 to 69, the method comprising culturing a recombinant microorganism according to any one of claims 77 to 81 under conditions sufficient for production of the nutritive protein by the recombinant microorganism.

83. The method of claim 82, further comprising isolating the nutritive protein from the culture.

84. A nutritive composition comprising a nutritive protein of any one of claims 1  
5 to 69 and at least one second component.

85. The nutritive composition of claim 84, wherein the at least one second component is selected from a protein, a polypeptide, a peptide, a free amino acid, a carbohydrate, a fat, a mineral or mineral source, a vitamin, a supplement, an organism, a  
10 pharmaceutical, and an excipient.

86. The nutritive composition of claim 85, wherein the at least one second component is a protein.

15 87. The method of claim 86, wherein the at least one second component is a nutritive protein.

88. The nutritive composition of claim 85 wherein the at least one second component is a free amino acid selected from essential amino acids, non-essential amino  
20 acids, branch chain amino acids, non-standard amino acids and modified amino acids.

89. The nutritive composition of claim 88, wherein the at least one second component is a free amino acid selected from essential amino acids.

25 90. The nutritive composition of claim 88, wherein the at least one second component is a free amino acid selected from branch chain amino acids.

91. The nutritive composition of claim 88, wherein the at least one second component is Leu.

30 92. The nutritive composition of claim 85, wherein the at least one second component is a lipid.

93. The nutritive composition of claim 92, wherein the lipid is selected from a fat, oil, triglyceride, cholesterol, phospholipid, and fatty acid.

94. The nutritive composition of claim 85, wherein the at least one second  
5 component is selected from a mineral and a vitamin.

95. The nutritive composition of claim 85, wherein the at least one second component is a supplement.

10 96. The nutritive composition of claim 85, wherein the at least one second component is an organism.

97. The nutritive composition of claim 85, wherein the at least one second component is a pharmaceutical.

15 98. The nutritive composition of claim 85, wherein the at least one second component is an excipient.

99. The nutritive composition of claim 98, wherein the at least one excipient is  
20 selected from a buffering agent, a preservative, a stabilizer, a binder, a compaction agent, a lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a coloring agent.

100. A nutritive composition of any one of claims 84 to 99, wherein the nutritive  
25 composition is formulated as a liquid solution, slurry, suspension, gel, paste, powder, or solid.

101. A method of making a nutritive composition of any one of claims 84 to 100, comprising providing a nutritive protein according to any one of claims 1 to 69 and  
30 combining the nutritive protein with the at least one second component.

102. A method of maintaining or increasing at least one of muscle mass, muscle strength, and functional performance in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive

composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

103. A method of maintaining or achieving a desirable body mass index in a  
5 subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

104. The method of claim 102 or 103, wherein the subject is at least one of elderly,  
10 critically-medically ill, and suffering from protein-energy malnutrition.

105. A method of providing protein to a subject with protein-energy malnutrition, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a  
15 nutritive composition made by the method of claim 101.

106. A method of increasing thermogenesis in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made  
20 by the method of claim 101.

107. A method of inducing at least one of a satiation response and a satiety response in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.  
25

108. The method of claim 106 or 107, wherein the subject is obese.

109. The method of any one of claims 102 to 108, wherein the nutritive protein of  
30 any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101 is consumed by the subject in coordination with performance of exercise.

110. A method of treating at least one of cachexia, sarcopenia and frailty in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

5

111. The method of any one of claims 102 to 110, wherein the nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101 is consumed by the subject by an oral, enteral, or parenteral route.

10

received by the International Bureau on 04 September 2013 (04.09.2013)

1. An isolated nutritive protein comprising a first polypeptide comprising a first polypeptide sequence that is homologous to a fragment of a naturally-occurring protein, wherein the fragment is at least 25 amino acids in length, wherein the first polypeptide sequence comprises at least one of:

5 a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;

b. a ratio of Leu residues to total amino acid residues of at least 11%; and

c. a ratio of essential amino acid residues to total amino acid residues of at least

10 49%,

and wherein the first polypeptide sequence has less than 90% global homology to a known allergen.

2. The isolated nutritive protein of claim 1, wherein the first polypeptide

15 sequence further comprises at least one of each essential amino acid.

3. The isolated nutritive protein of claim 1 or 2, wherein the first polypeptide sequence comprises at least 70% homology to the fragment of a naturally-occurring protein.

20 4. The isolated nutritive protein of claim 1 or 2, wherein the first polypeptide sequence comprises at least 95% homology to the fragment of a naturally-occurring protein.

5. The isolated nutritive protein of any one of claims 1 to 4, wherein the fragment of a naturally-occurring protein comprises at least 25 amino acid residues.

25

6. The isolated nutritive protein of any one of claims 1 to 4, wherein the fragment of a naturally-occurring protein comprises at least 50 amino acid residues.

30 7. The isolated nutritive protein of any one of claims 1 to 6, wherein the first polypeptide sequence comprises:

a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;

b. a ratio of Leu residues to total amino acid residues of at least 11%; and

c. a ratio of essential amino acid residues to total amino acid residues of at least 49%.

5 8. The isolated nutritive protein of claim any one of claims 1 to 7, wherein the first polypeptide sequence comprises the fragment of a naturally-occurring nutritive protein.

9. The isolated nutritive protein of any one of claims 1 to 7, wherein the first polypeptide sequence consists of the fragment of a naturally-occurring nutritive protein.

10 10. The isolated nutritive protein of any one of claims 1 to 9, wherein the first polypeptide sequence is not an allergen.

11. The isolated nutritive protein of any one of claims 1 to 10, wherein the first polypeptide sequence has less than 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 75, 80, 85, 15 or 90% global homology to a known allergen.

12. The isolated nutritive protein of any one of claims 1 to 11, wherein the first polypeptide sequence is not a toxin.

20 13. The isolated nutritive protein of claim 12, wherein the first polypeptide sequence has less than 50% global homology to a known toxin.

14. The isolated nutritive protein of any one of claims 1 to 13, wherein the first polypeptide sequence has a simulated gastric digestion half-life shorter than whey.

25 15. The isolated nutritive protein of claim 14, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 60 minutes or less than 30 minutes.

30 16. The isolated nutritive protein of claim 15, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 10 minutes.

17. The isolated nutritive protein of any one of claims 1 to 16, wherein the first polypeptide sequence is completely digested in simulated gastric fluid.

18. The isolated nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

5

19. The isolated nutritive protein of any one of claims 1 to 18, wherein the first polypeptide sequence comprises 0 or 1 cysteine residues.

10 20. The isolated nutritive protein of any one of claims 1 to 19, wherein the first polypeptide sequence comprises no disulfide bonds.

21. The isolated nutritive protein of any one of claims 1 to 20, wherein the first polypeptide sequence does not comprise N-linked glycosylation.

15 22. The isolated nutritive protein of any one of claims 1 to 21, wherein the first polypeptide sequence does not comprise O-linked glycosylation.

20 23. The isolated nutritive protein of any one of claims 1 to 22, wherein the first polypeptide sequence is resistant to aggregation.

24. The isolated nutritive protein of any one of claims 1 to 23, wherein the first polypeptide sequence is anionic at pH 7.

25 25. The isolated nutritive protein of any one of claims 1 to 24, wherein the first polypeptide sequence has an aqueous solubility at pH 7 of at least 12.5 g/L.

26. The isolated nutritive protein of any one of claims 1 to 25, wherein the first polypeptide sequence has a calculated solvation score of -20 or less.

30 27. The isolated nutritive protein of any one of claims 1 to 26, wherein the first polypeptide sequence has a calculated aggregation score of 0.75 or less.

28. The isolated nutritive protein of any one of claims 1 to 27, wherein the first polypeptide sequence has a calculated aggregation score of 0.5 or less.

29. The isolated nutritive protein of any one of claims 1 to 28, wherein the first polypeptide sequence comprises an amino acid sequence selected from:

- 5 i. an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609;
- ii. a modified derivative of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609; and
- iii. a mutein of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609.

10 30. The isolated nutritive protein of any one of claims 1 to 29, wherein the first polypeptide sequence consists of an amino acid sequence selected from:

- i. an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609;
- ii. a modified derivative of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609; and
- 15 iii. a mutein of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609.

20 31. The isolated nutritive protein of any one of claims 1 to 30, wherein the first polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609.

25 32. The isolated nutritive protein of any one of claims 1 to 31, wherein the first polypeptide sequence is a fragment of at least 25 amino acids of the naturally-occurring nutritive protein.

30 33. The isolated nutritive protein of any one of claims 1 to 32, wherein the first polypeptide sequence is a fragment of at least 50 amino acids of the naturally-occurring nutritive protein.

34. The isolated nutritive protein of any one of claims 1 to 33, wherein the isolated nutritive protein consists of the first polypeptide sequence and/or wherein the nutritive protein is a nutritive protein other than ovalbumin, ovotransferrin, ovomucoid,

myosin, actin, tropomyosin, collagen, troponin, whey, casein, alpha1 casein, alpha2 casein, beta casein, kappa casein, beta-lactoglobulin, alpha-lactalbumin, glycinin, beta-conglycinin, glutelin, prolamine, gliadin, glutenin, albumin, globulin, albumin, enolase, creatine kinase, phosphoglycerate mutase, triosephosphate isomerase, apolipoprotein, phosphoglucomutase, 5 phosphoglycerate kinase, glycerol-3-phosphate dehydrogenase, glyceraldehyde 3-phosphate dehydrogenase, hemoglobin, cofilin, glycogen phosphorylase, fructose-1,6-bisphosphatase, tropomyosin a-chain, casein kinase, aldolase, tubulin, vimentin, endoplasmic, lactate dehydrogenase, destrin, transthyretin, fructose bisphosphate aldolase, carbonic anhydrase, aldehyde dehydrogenase, annexin, adenosyl homocysteinase, titin, cofilin, phosphoglycerate 10 kinase, pyruvate dehydrogenase, glycogen phosphorylase, myokinase, parvalbumin, and/or desmin.

35. The isolated nutritive protein of any of claims 1 to 34, wherein the isolated nutritive protein further comprises a polypeptide tag for affinity purification.

15

36. The isolated nutritive protein of claim 35, wherein the tag for affinity purification is a polyhistidine-tag.

37. An isolated nutritive protein comprising a first polypeptide comprising a first 20 polypeptide sequence that is homologous to a fragment of a naturally-occurring protein, wherein the fragment is at least 25 amino acids in length, wherein the isolated nutritive protein comprises at least one of:

- a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;
- b. a ratio of Leu residues to total amino acid residues of at least 11%; and
- c. a ratio of essential amino acid residues to total amino acid residues of at least 49%,

and wherein the first polypeptide sequence has less than 90% global homology to a known allergen.

38. The isolated nutritive protein of claim 37, wherein the isolated nutritive protein further comprises at least one of each essential amino acid.

39. The isolated nutritive protein of claim 37 or 38, wherein the isolated nutritive protein comprises at least 70% homology to the fragment of a naturally-occurring protein.

40. The isolated nutritive protein of claim 37 or 38, wherein the isolated nutritive protein comprises at least 95% homology to the fragment of a naturally-occurring protein.

5 41. The isolated nutritive protein of any one of claims 37 to 40, wherein the fragment of a naturally-occurring protein comprises at least 25 amino acid residues.

42. The isolated nutritive protein of any one of claims 37 to 40, wherein the fragment of a naturally-occurring protein comprises at least 50 amino acid residues.

10 43. The isolated nutritive protein of any one of claims 37 to 42, wherein the isolated nutritive protein comprises:

- a. a ratio of branch chain amino acid residues to total amino acid residues of at least 24%;
- 15 b. a ratio of Lcu residues to total amino acid residues of at least 11%; and
- c. a ratio of essential amino acid residues to total amino acid residues of at least 49%.

44. The isolated nutritive protein of any one of claims 37 to 43, wherein the isolated nutritive protein comprises a fragment of a naturally occurring nutritive protein.

45. The isolated nutritive protein of any one of claims 37 to 43, wherein the isolated nutritive protein consists of a fragment of a naturally occurring nutritive protein.

25 46. The isolated nutritive protein of any one of claims 37 to 45, wherein the isolated nutritive protein is not an allergen.

47. The isolated nutritive protein of any one of claims 37 to 46, wherein the isolated nutritive protein has less than 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 75, 80, 30 85, or 90% global homology to a known allergen.

48. The isolated nutritive protein of any one of claims 37 to 47, wherein the isolated nutritive protein is not a toxin.

49. The isolated nutritive protein of claim 48, wherein the isolated nutritive protein has less than 50% global homology to a known toxin.

50. The isolated nutritive protein of any one of claims 37 to 49, wherein the isolated nutritive protein has a simulated gastric digestion half-life shorter than whey.

51. The isolated nutritive protein of claim 50, wherein the isolated nutritive protein has a simulated gastric digestion half-life of less than 60 minutes or less than 30 minutes.

10

52. The isolated nutritive protein of claim 51, wherein the isolated nutritive protein has a simulated gastric digestion half-life of less than 10 minutes.

15

53. The isolated nutritive protein of any one of claims 37 to 52, wherein the isolated nutritive protein is completely digested in simulated gastric fluid.

54. The isolated nutritive protein of any one of claims 37 to 53, wherein the isolated nutritive protein comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

20

55. The isolated nutritive protein of any one of claims 37 to 54, wherein the isolated nutritive protein comprises 0 or 1 cysteine residues.

25

56. The isolated nutritive protein of any one of claims 37 to 55, wherein the isolated nutritive protein comprises no disulfide bonds.

57. The isolated nutritive protein of any one of claims 37 to 56, wherein the isolated nutritive protein does not comprise N-linked glycosylation.

30

58. The isolated nutritive protein of any one of claims 37 to 57, wherein the isolated nutritive protein does not comprise O-linked glycosylation.

59. The isolated nutritive protein of any one of claims 37 to 58, wherein the isolated nutritive protein is resistant to aggregation.

60. The isolated nutritive protein of any one of claims 37 to 59, wherein the isolated nutritive protein is anionic at pH 7.

5 61. The isolated nutritive protein of any one of claims 37 to 60, wherein the isolated nutritive protein has an aqueous solubility at pH 7 of at least 12.5 g/L.

62. The isolated nutritive protein of any one of claims 37 to 61, wherein the isolated nutritive protein has a calculated solvation score of -20 or less.

10 63. The isolated nutritive protein of any one of claims 37 to 62, wherein the isolated nutritive protein has a calculated aggregation score of 0.75 or less.

15 64. The isolated nutritive protein of any one of claims 37 to 63, wherein the isolated nutritive protein has a calculated aggregation score of 0.5 or less.

65. The isolated nutritive protein of any one of claims 37 to 64, wherein the isolated nutritive protein comprises an amino acid sequence selected from:

- 20 i. an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609;  
ii. a modified derivative of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609; and  
iii. a mutein of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609.

25 66. The isolated nutritive protein of any one of claims 37 to 65, wherein the isolated nutritive protein consists of an amino acid sequence selected from:  
i. an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609;  
ii. a modified derivative of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609; and  
iii. a mutein of an amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609.

30 67. The isolated nutritive protein of any one of claims 37 to 66, wherein the isolated nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%,

86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from SEQ ID NO: 1 to SEQ ID NO: 2609.

5        68.      The isolated nutritive protein of any one of claims 37 to 67, wherein the isolated nutritive protein further comprises a polypeptide tag for affinity purification.

10       69.      The isolated nutritive protein of claim 68, wherein the tag for affinity purification is a polyhistidine-tag.

15       70.      An isolated nucleic acid comprising a nucleic acid sequence that encodes a nutritive protein according to any one of claims 1 to 69.

20       71.      The isolated nucleic acid of claim 70, wherein the isolated nucleic acid is selected from genomic DNA, cDNA, sense RNA and antisense RNA.

25       72.      The isolated nucleic acid of claim 71, wherein the isolated nucleic acid is genomic DNA.

30       73.      The isolated nucleic acid of claim 71, wherein the isolated nucleic acid is cDNA.

35       74.      The isolated nucleic acid of any one of claims 70 to 73, further comprising an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein.

40       75.      A vector comprising a nucleic acid sequence that encodes a nutritive protein of any one of claims 1 to 69.

45       76.      The vector of claim 75, further comprising an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein

50       77.      A recombinant microorganism comprising at least one of a nucleic acid of any one of claims 70 to 74 and a vector of any one of claims 75 and 76.

78. The recombinant microorganism according to claim 77, wherein the recombinant microorganism is a prokaryote.

5 79. The recombinant prokaryote according to claim 78, wherein the prokaryote is heterotrophic.

80. The recombinant prokaryote according to claim 78, wherein the prokaryote is autotrophic.

10

81. The recombinant prokaryote according to claim 78, wherein the prokaryote is a bacteria.

15

82. A method of making a nutritive protein according to any one of claims 1 to 69, the method comprising culturing a recombinant microorganism according to any one of claims 77 to 81 under conditions sufficient for production of the nutritive protein by the recombinant microorganism.

20

83. The method of claim 82, further comprising isolating the nutritive protein from the culture.

84. A nutritive composition comprising a nutritive protein of any one of claims 1 to 69 and at least one second component.

25

85. The nutritive composition of claim 84, wherein the at least one second component is selected from a protein, a polypeptide, a peptide, a free amino acid, a carbohydrate, a fat, a mineral or mineral source, a vitamin, a supplement, an organism, a pharmaceutical, and an excipient.

30

86. The nutritive composition of claim 85, wherein the at least one second component is a protein.

87. The method of claim 86, wherein the at least one second component is a nutritive protein.

88. The nutritive composition of claim 85 wherein the at least one second component is a free amino acid selected from essential amino acids, non-essential amino acids, branch chain amino acids, non-standard amino acids and modified amino acids.

5

89. The nutritive composition of claim 88, wherein the at least one second component is a free amino acid selected from essential amino acids.

90. The nutritive composition of claim 88, wherein the at least one second component is a free amino acid selected from branch chain amino acids.

10 91. The nutritive composition of claim 88, wherein the at least one second component is Leu.

15 92. The nutritive composition of claim 85, wherein the at least one second component is a lipid.

93. The nutritive composition of claim 92, wherein the lipid is selected from a fat, oil, triglyceride, cholesterol, phospholipid, and fatty acid.

20

94. The nutritive composition of claim 85, wherein the at least one second component is selected from a mineral and a vitamin.

95. The nutritive composition of claim 85, wherein the at least one second component is a supplement.

96. The nutritive composition of claim 85, wherein the at least one second component is an organism.

30 97. The nutritive composition of claim 85, wherein the at least one second component is a pharmaceutical.

98. The nutritive composition of claim 85, wherein the at least one second component is an excipient.

99. The nutritive composition of claim 98, wherein the at least one excipient is selected from a buffering agent, a preservative, a stabilizer, a binder, a compaction agent, a lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a  
5 coloring agent.

100. A nutritive composition of any one of claims 84 to 99, wherein the nutritive composition is formulated as a liquid solution, slurry, suspension, gel, paste, powder, or solid.

10

101. A method of making a nutritive composition of any one of claims 84 to 100, comprising providing a nutritive protein according to any one of claims 1 to 69 and combining the nutritive protein with the at least one second component.

15

102. A method of maintaining or increasing at least one of muscle mass, muscle strength, and functional performance in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

20

103. A method of maintaining or achieving a desirable body mass index in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

25

104. The method of claim 102 or 103, wherein the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition.

30

105. A method of providing protein to a subject with protein-energy malnutrition, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

106. A method of increasing thermogenesis in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

5

107. A method of inducing at least one of a satiation response and a satiety response in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

10

108. The method of claim 106 or 107, wherein the subject is obese.

109. The method of any one of claims 102 to 108, wherein the nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101 is consumed by the subject in coordination with performance of exercise.

110. A method of treating at least one of cachexia, sarcopenia and frailty in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101.

111. The method of any one of claims 102 to 110, wherein the nutritive protein of any one of claims 1 to 69, a nutritive composition of any one of claims 84 to 100, or a nutritive composition made by the method of claim 101 is consumed by the subject by an oral, enteral, or parenteral route.

112. A method of making a nutritive protein of any one of claims 1 to 69, the method comprising chemically synthesizing the protein.

30

113. A method of making a nutritive protein of any one of claims 1 to 69, the method comprising isolating the protein.

114. The isolated nutritive protein of any one of claims 1 to 69, wherein the first polypeptide sequence is at least 25 amino acids in length.

**STATEMENT UNDER ARTICLE 19 (1)**

The above amendments to the Claims are being submitted in accordance with the Patent Cooperation Treaty Article 19.

The differences between the claims as filed and the claims as amended are provided herein.

Claims 2-10, 12-13, 16-18, 20-33, 35-36, 38-46, 48-49, 52-54, 56-111 are unchanged.

Claims 1, 11, 14-15, 19, 34, 37, 47, 50-51 and 55 have been replaced by amended claims 1, 11, 14-15, 19, 34, 37, 47, 50-51 and 55.


Claims 112-114 are added.

The above-described amendments do not go beyond the disclosure of the international application as filed, and entry of these amendments is respectfully requested.


Replacement sheets effecting the above-described amendments are being transmitted herewith.

1/1

WHOLE PROTEINS/FRAGMENTS - RELATIVE LOG LIKELIHOOD OF EXPRESSION

**FIG. 1**

WHOLE PROTEINS/FRAGMENTS - RELATIVE LOG LIKELIHOOD OF SOLUBLE EXPRESSION

**FIG. 2**

## INTERNATIONAL SEARCH REPORT

International application No.  
**PCT/US2013/032218****A. CLASSIFICATION OF SUBJECT MATTER****C07K 14/435(2006.01)i, A23J 3/04(2006.01)i, A23L 1/305(2006.01)i, C12N 15/63(2006.01)i, C12N 1/21(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

C07K 14/435; A23L 1/29; C07K 4/12; A23L 1/305; A23J 3/34; A61K 38/14; A23J 3/16; A23J 1/04; A23K 1/16; A23J 3/04; C12N 15/63; C12N 1/21

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Korean utility models and applications for utility models  
Japanese utility models and applications for utility modelsElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
**eKOMPASS(KIPO internal) & Keywords:** nutritive protein, homologous, naturally occurring protein, branch chain amino acid, leucine, essential amino acid, muscle, body mass index, thermogenesis, etc.**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2010-126353 A1 (N. V. NUTRICIA) 04 November 2010 See claims 1, 9 and 10.	1-4, 37-40
A	KR 10-2010-0057630 A (AMINOTECH AS) 31 May 2010 See claims 1, 9, 10, and 13-15.	1-4, 37-40
A	US 2010-0317597 A1 (NEY, DENISE M. et al.) 16 December 2010 See claims 1, 2, and 6.	1-4, 37-40
A	EP 1552753 B1 (CENTRO DE INVESTIGACION EN ALIMENTACION Y DESARROLLO A.C.) 13 July 2005 See claims 1 and 3.	1-4, 37-40
A	EP 1455603 B1 (NUTRICA N.V.) 04 October 2006 See claims 1-3, 11, and 12.	1-4, 37-40

 Further documents are listed in the continuation of Box C. See patent family annex.

- \* Special categories of cited documents:  
 "A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier application or patent but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
 "&" document member of the same patent family

Date of the actual completion of the international search  19 July 2013 (19.07.2013)	Date of mailing of the international search report  <b>22 July 2013 (22.07.2013)</b>
Name and mailing address of the ISA/KR  Korean Intellectual Property Office 189 Cheongsa-ro, Seo-gu, Daejeon Metropolitan City, 302-701, Republic of Korea  Facsimile No. +82-42-472-7140	Authorized officer  HEO Joo Hyung  Telephone No. +82-42-481-8150

**INTERNATIONAL SEARCH REPORT**

International application No.

**PCT/US2013/032218****Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: 102-111  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims 102-111 pertain to methods for treatment of the human body by therapy, and thus relate to a subject matter which this International Searching Authority is not required, under Article 17(2)(a)(i) of the PCT and Rule 39.1(iv) of the Regulations under the PCT, to search.
2.  Claims Nos.: 11,13,15,16,36,47,49,51,52,69,71-73,76,78-81,83,85-99  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
Claims 11, 13, 15, 16, 36, 47, 49, 51, 52, 69, 71-73, 76, 78-81, 83, and 85-99 are unclear since they refer to claims which are not searchable due to not being drafted in accordance with the second and third sentence of Rule 6.4(a).
3.  Claims Nos.: 5-10,12,14,17-35,41-46,48,50,53-68,70,74,75,77,82,84,100-111  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No.

**PCT/US2013/032218**

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010-126353 A1	04/11/2010	CN 102413716 A CN 102458159 A EP 2424384 A1 EP 2424386 A1 MX 2011011320 A MX 2011011455 A US 2012-0094901 A1 US 2012-0283180 A1 WO 2010-126362 A1 WO 2010-131952 A1	11/04/2012 16/05/2012 07/03/2012 07/03/2012 20/01/2012 20/01/2012 19/04/2012 08/11/2012 04/11/2010 18/11/2010
KR 10-2010-0057630 A	31/05/2010	CA 2695444 A1 CN 101815441 A EP 2185004 A1 JP 2011-511620 A US 2011-0183040 A1 WO 2009-020394 A1	12/02/2009 25/08/2010 19/05/2010 14/04/2011 28/07/2011 12/02/2009
US 2010-0317597 A1	16/12/2010	AU 2011-259975 A1 CA 2764571 A1 CN 102481338 A EP 2440232 A1 MX 2011013126 A WO 2010-144821 A1	22/12/2011 16/12/2010 30/05/2012 18/04/2012 13/02/2012 16/12/2010
EP 1552753 B1	29/11/2006	AU 2004-314601 A1 DE 60216523 D1 DE 60216523 T2 EP 1552753 A1 WO 04-000033 A1	06/01/2004 11/01/2007 04/10/2007 13/07/2005 31/12/2003
EP 1455603 B1	04/10/2006	AT 341226 T AU 2004-290447 A1 AU 290447 B2 CA 2510760 A1 CA 2510760 C CN 100353866 C CN 1753625 A DE 60308810 D1 DE 60308810 T2 DK 1455603 T3 EP 1455603 A1 ES 2274299 T3 HK 1065447 A1 JP 2006-510711 A US 2004-0122097 A1 US 2008-0161398 A1 US 7288570 B2 WO 2004-056208 A1	15/10/2006 14/07/2004 04/10/2007 08/07/2004 20/09/2011 12/12/2007 29/03/2006 16/11/2006 02/08/2007 05/02/2007 15/09/2004 16/05/2007 09/02/2007 30/03/2006 24/06/2004 03/07/2008 30/10/2007 08/07/2004

## 摘要

本發明提供營養性蛋白質。在一些實施方案中，所述營養性蛋白質包含含有天然存在的營養性蛋白質的片段的第一多肽序列。在一些實施方案中，所述片段包含以下至少一者：a)所述營養性蛋白質中存在的支鏈氨基酸殘基與總氨基酸殘基的比率增大；b)所述營養性蛋白質中存在的亮氨酸殘基與總氨基酸殘基的比率增大；和 c)所述營養性蛋白質中存在的必需氨基酸殘基與總氨基酸殘基的比率增大。在一些實施方案中，所述片段包含以下至少一者：a)所述營養性蛋白質中存在的支鏈氨基酸殘基與總氨基酸殘基的比率；b)所述營養性蛋白質中存在的亮氨酸殘基與總氨基酸殘基的比率；和 c)所述營養性蛋白質中存在的必需氨基酸殘基與總氨基酸殘基的比率等於或大於如乳清、蛋或大豆蛋白質的基準蛋白質中存在的相應比率。也尤其提供編碼所述蛋白質的核酸、製備所述蛋白質的重組微生物、使用重組微生物製備所述蛋白質的方法、包含所述蛋白質的組合物和使用所述蛋白質的方法。