



(86) Date de dépôt PCT/PCT Filing Date: 1997/09/29
(87) Date publication PCT/PCT Publication Date: 1998/04/09
(45) Date de délivrance/Issue Date: 2003/03/11
(85) Entrée phase nationale/National Entry: 1999/03/29
(86) N° demande PCT/PCT Application No.: JP 1997/003467
(87) N° publication PCT/PCT Publication No.: 1998/014437
(30) Priorité/Priority: 1996/09/30 (08-278948) JP

(51) Cl.Int.⁶/Int.Cl.⁶ C07D 285/06, A01N 43/82, C07D 417/04


(72) Inventeurs/Inventors:
TSUBATA, KENJI, JP;
SHIMAOKA, TAKASHI, JP;
SAMPEI, OSAMU, JP;
TANIYAMA, YOSHITAKA, JP;
TAKAGI, KAZUHIRO, JP;
NISHIGUCHI, TSUTOMU, JP;
TAJIMA, SOHKICHI, JP

(73) Propriétaire/Owner:
NIHON NOHYAKU CO., LTD., JP

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : DERIVES DE 1,2,3-THIADIAZOLE ET SELS DE CES DERIVES, AGENTS EN USAGE DANS L'AGRICULTURE ET L'HORTICULTURE POUR LUTTER CONTRE LES MALADIES VEGETALES, ET PROCEDE D'UTILISATION CORRESPONDANT

(54) Title: 1,2,3-THIADIAZOLE DERIVATIVES OR SALTS THEREOF AND AGROHORTICULTURAL DISEASE CONTROLLER AND METHOD FOR USING THE SAME



(I)

(57) Abrégé/Abstract:

1,2,3-Thiadiazole derivatives of general formula (I) and salts thereof, disease controlling agents containing the derivatives or salts as the active ingredients, and a method for the use of the same, wherein R¹ is H, C₁-C₄ (halo)alkyl, a 5- or 6-membered heterocyclic group containing 1 to 3 heteroatoms which may be the same or different from each other and are selected from among O, N and S, or the like; A is (a),(b),(c),(d) (wherein R² and R³ are each H, halogeno, cyano, formyl, alkylthio, alkoxy carbonyl, substituted or unsubstituted phenyl or the like, or alternatively R² and R³ may be united to form a C₂-C₆ alkylene which is optionally interrupted by -O-, -S(O)_n- (wherein n is 0 to 2), -CO- or -NR⁸- (wherein R⁸ is H, halogeno or the like), thus forming a 3- to 7-membered ring); B is cyano, an optionally substituted 5- or 6-membered heterocyclic group containing 1 to 3 heteroatoms selected from among O, N and S, or the like; and m is 0 to 4.



PCT

特許協力条約に基づいて公開された国際出願

(51) 国際特許分類6 C07D 285/06, 417/04, A01N 43/82	A1	(11) 国際公開番号 WO98/14437 (43) 国際公開日 1998年4月9日(09.04.98)
(21) 国際出願番号 PCT/JP97/03467 (22) 国際出願日 1997年9月29日(29.09.97) (30) 優先権データ 特願平8/278948 1996年9月30日(30.09.96) JP (71) 出願人 (米国を除くすべての指定国について) 日本農薬株式会社(NIHON NOHYAKU CO., LTD.)(JP/JP) 〒103 東京都中央区日本橋1丁目2番5号 Tokyo, (JP) (72) 発明者; および (75) 発明者/出願人 (米国についてのみ) 津幡健治(TSUBATA, Kenji)(JP/JP) 〒586 大阪府河内長野市大矢船北町8-9 Osaka, (JP) 嶋岡孝史(SHIMAOKA, Takashi)(JP/JP) 〒591 大阪府堺市北花田町4丁目90-18 Osaka, (JP) 三瓶 修(SANPEI, Osamu)(JP/JP) 〒586 大阪府河内長野市市町1395-1-205 Osaka, (JP) 谷山吉隆(TANIYAMA, Yoshitaka)(JP/JP) 〒274 千葉県船橋市薬円台6丁目8-13-101 Chiba, (JP) 高木和裕(TAKAGI, Kazuhiro)(JP/JP) 〒550 大阪府大阪市西区北堀江4丁目12-10-1124 Osaka, (JP)		西口 勉(NISHIGUCHI, Tsutomu)(JP/JP) 〒586 大阪府河内長野市市町1395-1-203 Osaka, (JP) 田島崇吉(TAJIMA, Soukichi)(JP/JP) 〒558 大阪府大阪市住吉区长居2丁目10-9-301 Osaka, (JP) (74) 代理人 弁理士 浅村 皓, 外(ASAMURA, Kiyoshi et al.) 〒100 東京都千代田区大手町2丁目2番1号 新大手町ビル331 Tokyo, (JP) (81) 指定国 AU, CA, CN, KR, RU, US, 欧州特許 (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). 添付公開書類 国際調査報告書
(54)Title: 1,2,3-THIADIAZOLE DERIVATIVES AND SALTS THEREOF, DISEASE CONTROLLING AGENTS FOR AGRICULTURAL AND HORTICULTURAL USE, AND METHOD FOR THE USE THEREOF (54)発明の名称 1, 2, 3-チアジアゾール誘導体又はその塩類及び農園芸用病害防除剤ならびにその使用方法 (I) (a) (b) (c) (d) (57) Abstract 1,2,3-Thiadiazole derivatives of general formula (I) and salts thereof, disease controlling agents containing the derivatives or salts as the active ingredients, and a method for the use of the same, wherein R¹ is H, C₁-C₄ (halo)alkyl, a 5- or 6-membered heterocyclic group containing 1 to 3 heteroatoms which may be the same or different from each other and are selected from among O, N and S, or the like; A is (a), (b), (c), (d) (wherein R² and R³ are each H, halogeno, cyano, formyl, alkylthio, alkoxycarbonyl, substituted or unsubstituted phenyl or the like, or alternatively R² and R³ may be united to form a C₂-C₆ alkylene which is optionally interrupted by -O-, -S(O)ₙ- (wherein n is 0 to 2), -CO- or -NR⁸- (wherein R⁸ is H, halogeno or the like), thus forming a 3- to 7-membered ring); B is cyano, an optionally substituted 5- or 6-membered heterocyclic group containing 1 to 3 heteroatoms selected from among O, N and S, or the like; and m is 0 to 4.		

DESCRIPTION

1,2,3-THIADIAZOLE DERIVATIVES OR SALTS THEREOF
AND AGROHORTICULTURAL DISEASE CONTROLLER
AND METHOD FOR USING THE SAME

TECHNICAL FIELD

The present invention relates to 1,2,3-thiadiazole derivatives and salts thereof, and agrohorticultural disease controller containing said
5 compound as an active ingredient, and method for using the disease controller.

BACKGROUND ART


1,2,3-Thiadiazoles are disclosed in JP-A-2-149579/1990 as an agent for treating central nervous
10 system diseases, in JP-A-54-9272/1979, JP-A-3-181463/1991, JA-A-4-234881/1992, Canadian Patent 947297, etc. as a herbicide or a plant growth regulator, and in WO 9501340 and JP-A-7-252242/1995 as a fungicide.

DISCLOSURE OF THE INVENTION

15 With the aim of creating a novel agrohorticultural disease controller, the present inventors have conducted extensive studies to find that the 1,2,3-thiadiazole derivatives of the present invention represented by general formula (I) or salts
20 thereof which are novel compounds not found in

literature are useful as an agrohorticultural disease controller. Based on this finding, the present invention has been accomplished.

The present invention relates to 1,2,3-
 5 thiadiazole derivatives represented by the following general formula (I) or salts thereof, an agrohorticultural disease controller containing said compounds as an active ingredient, and a method for using said disease controller:




(I)

10 wherein R¹ represents hydrogen atom, C₁-C₈ alkyl group, halo C₁-C₄ alkyl group, phenyl group, substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, nitro group, C₁-C₄ alkyl
 15 group, halo C₁-C₄ alkyl group, C₁-C₄ alkoxy group, halo C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from oxygen atom, nitrogen atom and sulfur atom, or substituted 5- or 6-membered heterocycle
 20 containing 1 to 3 same or different heteroatoms selected from oxygen atom, nitrogen atom and sulfur atom and having 1 to 4 same or different substituents selected from the group consisting of halogen atom, cyano group,

hydroxyl group, mercapto group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, formyl group, carboxyl group, carboxy C₁-C₄ alkyl group, C₁-C₄ alkoxycarbonyl group, C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, amino group, substituted amino group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl C₁-C₄ alkyl group, and substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups;

A represents a group of the following formula:



(wherein R² and R³, same or different, represent hydrogen atom, halogen atom, cyano group, formyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, C₁-C₄ alkylsulfinyl group, C₁-C₄ alkylsulfonyl group, C₁-C₄ alkylcarbonyl group, C₁-C₄ alkoxycarbonyl group, phenyl group, substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group,

nitro group, C₁-C₄ alkyl group, halo C₁-C₄ alkyl group, C₁-C₄ alkoxy group, halo C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, phenylcarbonyl group, or substituted phenylcarbonyl group having, on the ring
5 thereof, 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, C₁-C₄ alkyl group and C₁-C₄ alkoxycarbonyl group; further, R² and R³ may be taken conjointly to form a 3- to 7-membered ring including a
10 C₂-C₆ alkylene group which may be intercepted by

-O-,

-S(O)_n- in which n is an integer of 0 to 2,

-CO- or


-NR⁸- in which R⁸ represents hydrogen atom,

15 formyl group, C₁-C₈ alkyl group, halo C₁-C₄ alkyl group, C₂-C₄ alkenyl group, halo C₂-C₄ alkenyl group, C₂-C₄ alkynyl group, halo C₂-C₄ alkynyl group, C₁-C₄ alkylthio group, C₁-C₄ alkylsulfinyl group, C₁-C₄ alkylsulfonyl group, C₁-C₄ alkylcarbonyl group, C₁-C₄ alkoxycarbonyl
20 group, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl group and phenyl group, phenyl group, substituted phenyl group having 1 to 5 same or different substituents selected from the
25 group consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, phenyl C₁-C₄ alkyl group, substituted phenyl C₁-C₄ alkyl group having, on the ring


25711-791

5


thereof, 1 to 5 same or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, phenylcarbonyl group, or substituted phenylcarbonyl group having, on the ring thereof, 1 to 5 same or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group; and said 3- to 7-membered ring may be substituted by one or more same or different substituents selected from the group consisting of halogen atom and C₁-C₄ alkyl group; and m represents an integer of 0 to 4), or



in which R² and R³ are as defined above, or




15 in which R² and R³ are as defined above, or



in which R² and R³ are as defined above; and

when A is -(CR²R³)_m- and m is 0, then B represents cyano group, 5- or 6-membered

heterocycle having 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom, substituted 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom and having 1 to 4 same or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, mercapto group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, formyl group, carboxyl group, carboxy C₁-C₄ alkyl group, C₁-C₄ alkoxycarbonyl group, C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, amino group, substituted amino group having 1 or 2 same or different C₁-C₄ alkyl or phenyl groups, carbamoyl group, substituted carbamoyl group having 1 or 2 C₁-C₄ alkyl or phenyl groups, carbamoyl C₁-C₄ alkyl group and substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different C₁-C₄ alkyl or phenyl groups, or



(in which R⁴ represents hydrogen atom, C₁-C₈ alkyl group, halo C₁-C₄ alkyl group, C₂-C₄ alkenyl group, halo C₂-C₄ alkenyl group, C₂-C₄ alkynyl group, halo C₂-C₄ alkynyl group, phenyl group, substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, phenyl C₁-C₄ alkyl group, substituted phenyl C₁-C₄ alkyl group having 1 to 5 same

or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom, or substituted 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom and having 1 to 4 same or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, mercapto group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, formyl group, carboxyl group, carboxy C₁-C₄ alkyl group, C₁-C₄ alkoxycarbonyl group, C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, amino group, substituted amino group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl group and phenyl group, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl group and phenyl group, carbamoyl C₁-C₄ alkyl group, and substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl group and phenyl group, and

X represents O, S,

N-R⁶ in which R⁶ is as defined later,

NO-R^6 in which R^6 is as defined later,

$\text{N}(\rightarrow\text{O})-\text{R}^6$ in which R^6 is as defined later,

$\text{NN}(\text{R}^6)\text{R}^7$ in which R^6 and R^7 are as defined later, or

- 5 $\text{NN}=\text{C}(\text{R}^6)\text{R}^7$ (in which R^6 and R^7 , same or different, represent hydrogen atom, halogen atom, formyl group, $\text{C}_1\text{-C}_4$ alkyl group, $\text{C}_2\text{-C}_4$ alkenyl group, $\text{C}_2\text{-C}_4$ alkynyl group, $\text{C}_1\text{-C}_4$ alkylsulfonyl group, halo $\text{C}_1\text{-C}_4$ alkylsulfonyl group, $\text{C}_1\text{-C}_{20}$ alkylcarbonyl group, $\text{C}_1\text{-C}_4$ alkoxy carbonyl group, phenyl group, substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, nitro group, $\text{C}_1\text{-C}_4$ alkyl group, halo $\text{C}_1\text{-C}_4$ alkyl group, $\text{C}_1\text{-C}_4$ alkoxy group, halo $\text{C}_1\text{-C}_4$ alkoxy group, amino group and $\text{C}_1\text{-C}_4$ alkoxy carbonyl group, phenyl $\text{C}_1\text{-C}_4$ alkyl group, substituted phenyl $\text{C}_1\text{-C}_4$ alkyl group having, on the ring thereof, 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, nitro group, $\text{C}_1\text{-C}_4$ alkyl group, halo $\text{C}_1\text{-C}_4$ alkyl group, $\text{C}_1\text{-C}_4$ alkoxy group, halo $\text{C}_1\text{-C}_4$ alkoxy group, amino group and $\text{C}_1\text{-C}_4$ alkoxy carbonyl group, phenyl carbonyl group, substituted phenyl carbonyl group having 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, nitro group, $\text{C}_1\text{-C}_4$ alkyl group, halo $\text{C}_1\text{-C}_4$ alkyl group, $\text{C}_1\text{-C}_4$ alkoxy group, halo $\text{C}_1\text{-C}_4$ alkoxy group and $\text{C}_1\text{-C}_4$ alkoxy carbonyl group, carbamoyl group, substituted

25711-791

9

carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl group, phenyl group and substituted phenyl group having, on the ring thereof, 1 to 5 same or different substituents selected from the group consisting of halogen atom, C₁-C₄ alkyl group, halo C₁-C₄ alkyl group, C₁-C₄ alkoxy group and halo C₁-C₄ alkoxy group, phenylsulfonyl group, substituted phenylsulfonyl group having 1 to 5 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, nitro group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, thio-carbamoyl group, substituted thiocarbamoyl group substituted by same or different C₁-C₄ alkyl or phenyl group, C₁-C₄ alkoxy carbonimidoyl group, substituted C₁-C₄ alkoxy carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, C₁-C₄ alkylthio carbonimidoyl group, substituted C₁-C₄ alkylthio carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, C₁-C₄ alkylsulfinyl carbonimidoyl group, substituted C₁-C₄ alkylsulfinyl carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, C₁-C₄ alkylsulfonyl carbonimidoyl group, substituted C₁-C₄ alkylsulfonyl

25711-791

10

carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, amidino group, substituted amidino group having same or
5 different substituents selected from the group consisting of C₁-C₆ alkyl and phenyl groups, 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom, or
10 substituted 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom and having 1 to 4 same or different substituents selected from the group consisting of halogen atom,
15 cyano group, hydroxyl group, mercapto group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, formyl group, carboxyl group, carboxy C₁-C₄ alkyl group, C₁-C₄ alkoxycarbonyl group, C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, amino group, substituted amino group having 1 or
20 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl
25 C₁-C₄ alkyl group, and substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; and R⁶ and R⁷ may be taken conjointly to

25711-791

11

form a 3- to 7-membered ring including a C₂-C₆ alkylene group which may be intercepted by


-O-,

-S(O)_n- in which n is as defined above,

5 -CO-, or


-NR⁸- in which R⁸ is as defined above, and said 3- to 7-membered ring may have same or different substituents selected from the group consisting of halogen atom and C₁-C₄ alkyl group)); and

10 when A is -(CR²R³)_m- and m is 1 or greater or A is



then B represents hydrogen atom, halogen atom, cyano group, 5- or 6-membered heterocycle containing 1 to 3 same or
 15 different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom, substituted 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom and having 1 to 3
 20 same or different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, mercapto group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, formyl group, carboxyl group, carboxy C₁-C₄ alkyl group, C₁-C₄ alkoxycarbonyl group, C₁-C₄ alkoxycarbonyl
 25 C₁-C₄ alkyl group, amino group, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl C₁-C₄ alkyl group, and substituted carbamoyl C₁-C₄ alkyl group having 1 or 2

same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups,



(in which X is as defined above; and R⁵ represents

5 hydrogen atom, C₁-C₈ alkyl group, halo C₁-C₄ alkyl group, C₂-C₄ alkenyl group, halo C₂-C₄ alkenyl group, C₂-C₄ alkynyl group, halo C₂-C₄ alkynyl group, phenyl group, substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of

10 halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, phenyl C₁-C₄ alkyl group, substituted phenyl C₁-C₄ alkyl group having, on the ring thereof, 1 to 5 same or different substituents selected from the group

15 consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and C₁-C₄ alkoxycarbonyl group, 5- to 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom,

20 nitrogen atom and sulfur atom, substituted 5- or 6-membered heterocycle containing 1 to 3 same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom and having 1 to 4 same or different substituents selected from the


25 group consisting of halogen atom, cyano group, hydroxyl group, mercapto group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group, C₁-C₄ alkylthio group, formyl group, carboxyl group, carboxy C₁-C₄ alkyl group, C₁-C₄ alkoxycarbonyl

group, C₁-C₄ alkoxy carbonyl C₁-C₄ alkyl group, amino group, substituted amino group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, carbamoyl C₁-C₄ alkyl group, and substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups,

O-R⁶ in which R⁶ is as defined above,

S(O)_n-R⁶ in which n and R⁶ are as defined above, or

N(R⁶)R⁷ in which R⁶ and R⁷ are as defined above),



(wherein R⁵ is as defined above; and R⁹ and R¹⁰, same or different, represent formyl group, C₁-C₄ alkyl carbonyl group, C₁-C₄ alkoxy carbonyl group, phenyl carbonyl group, or substituted phenyl carbonyl group having, on the ring thereof, 1 to 5 same or different substituents selected from the group consisting of halogen atom, cyano group, hydroxyl group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and

25711-791

14

C₁-C₄ alkoxy carbonyl group; further, R⁹ and R¹⁰ may be taken conjointly to form a 3- to 7-membered ring including a C₂ to C₆ alkylene group which may be intercepted by

-O-,

5 -S(O)_n- in which n is as defined above,

-CO-, or

-N(R⁸) - in which R⁸ is as defined above, and said 3- to 7-membered ring may have one or more same or different substituents selected from the group consisting of halogen
10 atom and C₁-C₄ alkyl group),

O-R⁶ in which R⁶ is as defined above,

S(O)_n-R⁶ in which R⁶ and n are as defined above,


N(R⁶)R⁷ in which R⁶ and R⁷ are as defined above,

N=C(R⁶)R⁷ in which R⁶ and R⁷ are as defined above, or

15 ON=C(R⁶)R⁷ in which R⁶ and R⁷ are as defined above,

provided that (1) when R¹ is a hydrogen atom or C₁-C₄ alkyl group, A is

20




m is 1, both of R² and R³ are hydrogen atoms or either one of R² and R³ is a hydrogen atom and other is a C₁-C₄ alkyl group, then B is not a chlorine or bromine atom;

25711-791


14a

(2) when R^1 is a C_1 - C_4 alkyl group or halo C_1 - C_4 alkyl group, A is



m is 1, either one of R^2 and R^3 is a hydrogen atom and other is a C_1 - C_4 alkyl group, then B is not a hydrogen atom; and

(3) when R^1 is a hydrogen atom, furan-2-yl, thiophen-2-yl, phenyl, or phenyl having 1 to 3 same or different substituents selected from a hydroxyl group and C_1 - C_4 alkoxy group, A is



and m is 0, then B is not a furan-2-yl, thiophen-2-yl, phenyl, or phenyl having 1 to 3 same or different substituents selected from a hydroxyl group and C_1 - C_4 alkoxy group.

BEST MODE FOR CARRYING OUT THE INVENTION

In the definitions of the substituents in the 1,2,3-thiadiazole derivatives of the present invention represented by general formula (I), the term "halogen atom" means chlorine atom, bromine atom, iodine atom or fluorine

25711-791

14b

atom; the term "C₁-C₈ alkyl group" means a straight or branched chain alkyl group having 1 to 8 carbon atoms such as methyl, ethyl, n-propyl, i-propyl,

n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl and the like; the term "halo C₁-C₄ alkyl group" means a straight or branched chain alkyl group having 1 to 4 carbon atoms and substituted by one or more same or different halogen atoms, the term "C₂-C₄ alkenyl group" means a straight or branched chain alkenyl group having 2 to 4 carbon atoms and a double bond, and the term "halo C₂-C₄ alkenyl group" means a straight or branched chain alkenyl group having 2 to 4 carbon atoms and substituted by at least one same or different halogen atom.

The term "C₂-C₄ alkynyl group" means a straight or branched chain alkynyl group having 2 to 4 carbon atoms and a triple bond, the term "halo C₂-C₄ alkynyl group" means a straight or branched chain alkynyl group having 2 to 4 carbon atoms and substituted by one or more same or different halogen atoms, and the term "5- or 6-membered heterocycle containing at least one same or different hetero atoms selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom" means a 5- or 6-membered ring such as furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridine, pyridazine, pyrimidine, pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithiolane, dithiane, piperazine, dioxolane, imidazolidine and the like.


As examples of the salt of 1,2,3-thiadiazole derivative represented by general formula (I), there can be referred to salts of alkali metals such as sodium, potassium and the like, salts of alkaline earth metals such as calcium, magnesium and the like, ammonium salts, substituted ammonium salts substituted by one or more same or different substituents selected from the group consisting of C₁-C₁₂ alkyl group, phenyl group, substituted phenyl group, benzyl group or substituted benzyl group, guanidium salt and the like.

As preferred substituents of the present invention, the following are referred to. Thus, as R¹, methyl, ethyl, n-propyl, cyclopropyl and the like are preferred. As A, a single bond (B is directly linked to thiadiazole ring) or methylene are preferred. In a case where A is a single bond, B is preferably cyano, formyl, hydroxyiminomethyl, benzyloxyiminomethyl, hydrazonomethyl, phenylhydrazonomethyl, imidazolin-2-yl, 4-methylthiazol-2-yl and the like. In a case where A is methylene, B is preferably chlorine atom, hydroxyl group, methoxy group, benzyloxy group, acetoxy group, benzoyloxy group, ethoxycarbonyloxy group, amino group and hydrochloride thereof, diethylamino group, dipropenylamino group, benzoylamino group, p-toluenesulfonylamino group, 3-phenylureido group, 4-methyl-1,2,3-thiadiazol-5-ylmethoxy group and the like.

The 1,2,3-thiadiazole derivatives of the present invention represented by general formula (I) and

salts thereof can be produced, for example, by the Production processes exemplified below.


Production process 1



wherein R^1 is as defined above and R^{11} represents C_1-C_8 alkyl group.

A 1,2,3-thiadiazole derivative represented by general formula (I-1) can be produced by reducing a compound of general formula (II) in the presence of a reductant to form a 1,2,3-thiadiazole derivative represented by general formula (I-1) and, after isolating or without isolating the thus formed (I-1), oxidizing (I-1) in the presence of an oxidant.

This reaction can be performed according to the description of Shin Jikken Kagaku Koza, Vol. 15 (II), p. 191 (Maruzen K. K.); Jikken Kagaku Koza (4th edition), Vol. 23(V), p. 43 (Maruzen K. K.), etc.

Production process 2


wherein R^1 and R^{11} are as defined above, and $R^{4'}$
 represents phenyl group, substituted phenyl group having
 1 to 5 same or different substituents selected from the
 5 group consisting of halogen atom, cyano group, hydroxyl
 group, C_1 - C_4 alkyl group, C_1 - C_4 alkoxy group and C_1 - C_4
 alkoxycarbonyl group, 5- or 6-membered heterocycle
 containing 1 to 3 same or different hetero atoms
 selected from the group consisting of oxygen atom,
 10 nitrogen atom and sulfur atom, or substituted 5- or 6-
 membered heterocycle containing 1 to 3 same or different
 hetero atoms selected from the group consisting of
 oxygen atom, nitrogen atom and sulfur atom and having 1
 to 4 same or different substituents selected from the
 15 group consisting of halogen atom, cyano group, hydroxyl
 group, mercapto group, C_1 - C_4 alkyl group, C_1 - C_4 alkoxy
 group, C_1 - C_4 alkylthio group, formyl group, carboxyl
 group, carboxy C_1 - C_4 alkyl group, C_1 - C_4 alkoxycarbonyl
 group, C_1 - C_4 alkoxycarbonyl C_1 - C_4 alkyl group, amino
 20 group, substituted amino group having 1 or 2 same or

different substituents selected from the group consisting of C_1 - C_4 alkyl and phenyl groups, carbamoyl group, substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C_1 - C_4 alkyl and phenyl groups, carbamoyl C_1 - C_4 alkyl group, and substituted carbamoyl C_1 - C_4 alkyl group having 1 or 2 same or different substituents selected from the group consisting of C_1 - C_4 alkyl and phenyl groups.

A 1,2,3-thiadiazole derivative represented by general formula (I-3) can be produced by hydrolyzing a compound of general formula (II) and then reacting it with thionyl chloride to form a compound represented by general formula (III), and after isolating compound (III) or without isolating (III), reacting (III) with a compound of general formula (IV) in the presence of a catalyst.

This reaction can be performed according to the description of "Shin Jikken Kagaku Koza", Vol. 14(II), p. 794.

Production process 3




wherein R^1 , R^4 and R^6 are as defined above.

A 1,2,3-thiadiazole derivative represented by general formula (I-5) can be produced by reacting a 1,2,3-thiadiazole derivative of general formula (I-4) with a compound represented by general formula (V).

This reaction can be performed according to the description of "Jikken Kagaku Koza" (4th edition), Vol. 20 (V), p. 353.


Production process 4



wherein R^1 , R^4 , R^6 and R^7 are as defined above.

A 1,2,3-thiadiazole derivative represented by general formula (I-6) can be produced by reacting a 1,2,3-thiadiazole derivative of general formula (I-4) with a compound represented by general formula (VI).

This reaction can be performed according to the description of Org. Synth. Coll. VI, p.12 (1988).

Production process 5


wherein R^1 and R^6 are as defined above and Hal represents halogen atom.

A 1,2,3-thiadiazole derivative represented by
 5 general formula (I-8) or (I-9) can be produced by

treating a 1,2,3-thiadiazole derivative represented by
 general formula (I-1) with a halogenating agent such as
 thionyl chloride or phosphorus tribromide to form a
 compound represented by general formula (III'), and
 5 after isolating the compound (III') or without isolating
 (III'), reacting the compound (III') with sodium
 formimide and then with an acid such as hydrochloric
 acid to form a salt of a 1,2,3-thiadiazole derivative
 represented by general formula (I-7), and then reacting
 10 the salt (I-7) with a compound of formula (VII).
 Further, if a salt of the 1,2,3-thiadiazole derivative
 of formula (I-7) is reacted with an isocyanate of
 formula (VIII), a 1,2,3-thiadiazole derivative of
 general formula (I-8') can be obtained.

15 This reaction can be performed according to
 the description of "Jikken Kagaku Koza" (4th edition),
 Vol. 19 (I), p. 438; Synthesis, 615 (1990); "Jikken
 Kagaku Koza" (4th edition), Vol. 22 (II), p. 1142; "Shin
 Jikken Kagaku Koza", Vol. 14 (III), p. 1631; etc.

20 Production process 6




wherein R^1 , R^6 and Hal are as defined above.

A 1,2,3-thiadiazole derivative represented by general formula (I-10) can be produced by reacting a 1,2,3-thiadiazole derivative of formula (I-1) with a compound represented by formula (VII) in the presence of a base.

This reaction can be performed according to the description of "Jikken Kagaku Koza" (4th edition), Vol. 22 (V), p. 50.

10 Production process 7




wherein R^1 , R^6 and Hal are as defined above and n' represents an integer of 1-2.

A 1,2,3-thiadiazole derivative represented by general formula (I-12) can be produced by reacting a 1,2,3-thiadiazole derivative of general formula (I-1) with a compound represented by general formula (IX) in the presence of a base to form a 1,2,3-thiadiazole derivative represented by general formula (I-11), and after isolating the compound (I-11) or without isolation

it, oxidizing the derivative (I-11) in the presence of an oxidant.

This reaction can be performed according to the description of "Shin Jikken Kagaku Koza", Vol. 14 (III), p. 1716; "Jikken Kagaku Koza" (4th edition), Vol. 24 (VI), p. 365; etc.

Production process 8




wherein R^1 and Hal are as defined above; and R^{12} and R^{13} , same or different, represent C_1 - C_4 alkyl group, carboxyl group, carboxy C_1 - C_4 alkyl group, C_1 - C_4 alkoxycarbonyl group, C_1 - C_4 alkoxycarbonyl C_1 - C_4 alkyl group, carbamoyl group, substituted carbamoyl having 1 or 2 same or different substituents selected from the group consisting of C_1 - C_4 alkyl and phenyl groups, carbamoyl C_1 - C_4 alkyl group, and substituted carbamoyl C_1 - C_4 alkyl group having 1 or 2 same or different substituents selected from the group consisting of C_1 - C_4 alkyl and phenyl groups.

A 1,2,3-thiadiazole derivative represented by general formula (I-13) can be produced by reacting a compound represented by general formula (X) with phosphorus pentasulfide and then with a compound represented by general formula (XI).

This reaction can be performed according to the description of "Shin Jikken Kagaku Koza", Vol. 14 (IV), p. 2198; Org. Synth., Coll. Vol. III, p. 332 (1955); etc.

10 Production process 9



wherein R^1 is as defined above, R^{14} represents hydrogen atom or C_1 - C_4 alkyl, and W represents C_3 - C_6 alkylene group which may be substituted by C_1 - C_4 alkyl group and may be intercepted by

15

-O-,

- NR^6 - in which R^6 is as defined above,

- $S(O)_n$ - in which n is as defined above, or


-CO-.

A 1,2,3-thiadiazole derivative represented by

general formula (I-15) can be produced by subjecting a compound of general formula (X) to a dehydrating reaction to form a 1,2,3-thiadiazole derivative represented by general formula (I-14), and after
 5 isolating the derivative (I-14) or without isolating it, reacting the derivative (I-14) with a compound represented by general formula (XII).

This reaction can be performed according to the description of "Shin Jikken Kagaku Koza", Vol. 14
 10 (II), p. 1190.

Production process 10




wherein R¹, R², R³ and B are as defined above.

A 1,2,3-thiadiazole derivative represented by general formula (I-17) can be produced by reacting a
 15 1,2,3-thiadiazole derivative represented by general

formula (I-16) with a compound represented by formula (XIII) or formula (XIV).

This reaction can be performed according to the description of "Shin Jikken Kagaku Koza", Vol. 14 (I), p. 224; JP-A-7-165757/1995; etc.

The compound of general formula (II) used as a starting compound for production of the 1,2,3-thiadiazine derivative represented by general formula (I) can be produced, for example, by the following method.



wherein R^1 is as defined above, R^{11} represents C_1 - C_8 alkyl group, and R^{12} represents C_1 - C_4 alkyl group, C_1 - C_4 alkoxy group or amino group which may have a substituent.

A compound represented by general formula (II) can be produced by reacting a compound represented by general formula (XV) with a compound represented by general formula (XVI) to form a compound represented by general formula (XVII), and after isolating the compound

(XVII) or without isolating it, subjecting the compound (XVII) to ring closure using thionyl chloride.













This reaction can be preformed according to the description of J. Am. Chem. Soc., 77, 5359 (1955),
 5 etc.

















Next, typical examples of the 1,2,3-thiadiazole derivative of the present invention represented by general formula (I) are mentioned below. The invention is by no means limited by these examples.

10 In Table 1, meaning of the abbreviations are as follows.


Ph: phenyl group

cyclo: alicyclic hydrocarbon





General formula (I)



(I)

Table 1

No	R ¹	A	B	Property
1	H	—	COPh	m. p. 60-61 °C
2	H	CH ₂	Cl	
3	H	CH ₂	OH	
4	H	CH ₂	OCH ₃	
5	H	CH ₂	OCH ₂ Ph	

Table 1 (Continued)

No	R ¹	A	B	Property
6	H	CH ₂	OCH ₂ -Q ₁	
7	CH ₃	—	CHO	nD 1.5555(26°C)
8	CH ₃	—	COCH ₃	
9	CH ₃	—	COPh	nD 1.6145(22°C)
10	CH ₃	—	CO(2-Cl-Ph)	
11	CH ₃	—	CO(4-Cl-Ph)	
12	CH ₃	—	CO(2-CH ₃ -Ph)	
13	CH ₃	—	CO(4-CH ₃ -Ph)	
14	CH ₃	—	CO(4-i-C ₃ H ₇ -Ph)	
15	CH ₃	—	CO(4-CH ₃ O-Ph)	
16	CH ₃	—	CO(4-t-C ₄ H ₉ -Ph)	
17	CH ₃	—	CO-(1-CH ₃ -Q ₂)	m. p. 85-90 °C
18	CH ₃	—	CN	m. p. 50.4°C
19	CH ₃	—	CH=NOH	m. p. 270 °C(dec.)
20	CH ₃	—	CH=NOCH ₃	m. p. 97°C
21	CH ₃	—	CH=NOCH ₂ Ph	nD 1.5998(25°C)
22	CH ₃	—	CH=NOCH ₂ (2-Cl-Ph)	
23	CH ₃	—	CH=NOCH ₂ (3-Cl-Ph)	
24	CH ₃	—	CH=NOCH ₂ (4-Cl-Ph)	m. p. 70-80 °C
25	CH ₃	—	CH=NOCH ₂ (2-CH ₃ -Ph)	
26	CH ₃	—	CH=NOCH ₂ (3-CH ₃ -Ph)	
27	CH ₃	—	CH=NOCH ₂ (4-CF ₃ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
28	CH ₃	—	CH=NOCH ₂ (4-OH-Ph)	
29	CH ₃	—	CH=NOCH ₂ (4-CH ₃ O-Ph)	
30	CH ₃	—	CH=NOCH ₂ (4-CF ₃ O-Ph)	
31	CH ₃	—	CH=NOCH ₂ (4-NH ₂ -Ph)	
32	CH ₃	—	CH=NOCH ₂ (4-NO ₂ -Ph)	
33	CH ₃	—	CH=NOCH ₂ (4-CN-Ph)	
34	CH ₃	—	CH=NOCH ₂ (4-COOCH ₃ -Ph)	
35	CH ₃	—	CH=NNH ₂	m. p. 81 °C
36	CH ₃	—	CH=NNHCH ₃	
37	CH ₃	—	CH=NNHPh	m. p. 218 °C
38	CH ₃	—	CH=NNH(2-Cl-Ph)	
39	CH ₃	—	CH=NNH(2-F-4-Cl-Ph)	
40	CH ₃	—	CH=NNH(3,5-Cl ₂ -Ph)	
41	CH ₃	—	CH=NNH(3-Cl-Ph)	
42	CH ₃	—	CH=NNH(3-i-C ₃ H ₇ O-Ph)	
43	CH ₃	—	CH=NNH(4-CF ₃ -Ph)	
44	CH ₃	—	CH=NNH(4-Cl-Ph)	
45	CH ₃	—	CH=NNHCOOCH ₃	
46	CH ₃	—	CH=NNHCOOC ₄ H ₉ -t	
47	CH ₃	—	CH=NNHCONHCH ₃	
48	CH ₃	—	CH=NNHCONHC ₂ H ₅	
49	CH ₃	—	CH=NNHCON(CH ₃)Ph	

Table 1 (Continued)

No	R ¹	A	B	Property
50	CH ₃	—	CH=NNHCONH(4-Cl-Ph)	
51	CH ₃	—	CH=NNHCONH(4-t-C ₄ H ₉ -Ph)	
52	CH ₃	—	CH=NNHCONH(4-CF ₃ -Ph)	
53	CH ₃	—	CH=NNHCONH(4-i-C ₃ H ₇ O-Ph)	
54	CH ₃	—	CH=NNHCONH(4-CF ₃ O-Ph)	
55	CH ₃	—	CH=NNHCSNHCH ₃	
56	CH ₃	—	CH=NNHC(N(CH ₃) ₂)=NCH ₃	
57	CH ₃	—	CH=NNHC(OCH ₃)=NCH ₃	
58	CH ₃	—	CH=NNHC(SCH ₃)=NCH ₃	
59	CH ₃	—	CH=NNHC(SOCH ₃)=NCH ₃	
60	CH ₃	—	CH=NNH-(4, 6-(CH ₃) ₂ -Q ₃)	m. p. 182-183 °C
61	CH ₃	—	C(Ph)=NOH	m. p. 172 °C
62	CH ₃	—	C(Ph)=NOCH ₃	nD 1.5905(22.8°C)
63	CH ₃	—	C(Ph)=NOC ₈ H ₁₇	
64	CH ₃	—	C(Ph)=NOCH ₂ CH=CH ₂	
65	CH ₃	—	C(Ph)=NOCH ₂ CCl=CH ₂	
66	CH ₃	—	C(Ph)=NOCH ₂ C≡CH	nD 1.5945(22.8°C)
67	CH ₃	—	C(Ph)=NOCH ₂ C≡CBr	
68	CH ₃	—	C(Ph)=NOCH ₂ C≡CCl	
69	CH ₃	—	C(Ph)=NOCH ₂ Ph	nD 1.6183(23.9°C)
70	CH ₃	—	C(Ph)=NOCH ₂ (4-Cl-Ph)	
71	CH ₃	—	C(Ph)=NOCH ₂ (3, 4-Cl ₂ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
72	CH ₃	—	C(Ph)=NOCH ₂ (2, 4, 6-Cl ₃ -Ph)	
73	CH ₃	—	C(Ph)=NOCH ₂ (2-CH ₃ -Ph)	
74	CH ₃	—	C(Ph)=NOCH ₂ (3-CH ₃ -Ph)	
75	CH ₃	—	C(Ph)=NOCH ₂ (4-CH ₃ -Ph)	
76	CH ₃	—	C(Ph)=NOCH ₂ (3-CH ₃ O-Ph)	
77	CH ₃	—	$\begin{array}{c} \text{C}(=\text{N}^+ \text{CH}_3)\text{-Ph} \\ \downarrow \\ \text{O}^- \end{array}$	m. p. 180.1 °C
78	CH ₃	—	$\begin{array}{c} \text{C}(=\text{N}^+ \text{CH}_2\text{Ph})\text{Ph} \\ \downarrow \\ \text{O}^- \end{array}$	m. p. 116.3 °C
79	CH ₃	—	C(Ph)=NNH ₂	m. p. 72.3 °C
80	CH ₃	—	C(Ph)=NNHCOOC ₂ H ₅	
81	CH ₃	—	C(Ph)=NNHCSNHPh	
82	CH ₃	—	3-CN-Q ₂	m. p. 190-200 °C
83	CH ₃	—	Q ₄	
84	CH ₃	—	5-CH ₃ -5-i-C ₃ H ₇ -4-(=O)-Q ₄	m. p. 59 °C
85	CH ₃	—	5-(CH ₃) ₂ -Q ₅	m. p. 114 °C
86	CH ₃	—	4-(CH ₃) ₂ -5-(=O)-Q ₆	m. p. 107 °C
87	CH ₃	—	4, 6-(CH ₃) ₂ -Q ₃	m. p. >180 °C
88	CH ₃	—	4-CH ₃ -Q ₇	m. p. 89 °C
89	CH ₃	—	4-t-C ₄ H ₉ -Q ₇	m. p. 61 °C
90	CH ₃	—	4-(2-Cl-Ph)-Q ₇	m. p. 136 °C
91	CH ₃	—	4-(4-Cl-Ph)-Q ₇	m. p. 125 °C

Table 1 (Continued)

No	R'	A	B	Property
92	CH ₃	—	4-(2,4-Cl ₂ -Ph)-Q ₇	m. p. 170 °C
93	CH ₃	—	4-(4-CH ₃ -Ph)-Q ₇	m. p. 118 °C
94	CH ₃	—	5-CH ₃ -4-Ph-Q ₇	m. p. 151 °C
95	CH ₃	—	4-CONHCH ₃ -Q ₇	m. p. 144 °C
96	CH ₃	—	4-CH ₂ COOH-Q ₇	m. p. 196 °C
97	CH ₃	—	4-CH ₂ CONH ₂ -Q ₇	m. p. 182 °C
98	CH ₃	—	4-CH ₂ COOC ₂ H ₅ -Q ₇	m. p. 98°C
99	CH ₃	—	4-CH ₂ CONHCH ₃ -Q ₇	m. p. 155 °C
100	CH ₃	—	4-CH ₂ CON(C ₂ H ₅) ₂ -Q ₇	m. p. 132 °C
101	CH ₃	—	5-Ph-Q ₈	m. p. 154 °C
102	CH ₃	—	5-SH-Q ₈	m. p. 182 °C
103	CH ₃	—	5-SCH ₃ -Q ₈	m. p. 65°C
104	CH ₃	CH ₂	Cl	nD 1.5590(26°C)
105	CH ₃	CH ₂	OH	nD 1.553(21 °C)
106	CH ₃	CH ₂	OCH ₃	nD 1.5180(26°C)
107	CH ₃	CH ₂	OC ₄ H ₉ -n	
108	CH ₃	CH ₂	OC ₈ H ₁₇ -n	
109	CH ₃	CH ₂	OCH ₂ Ph	nD 1.5730(25°C)
110	CH ₃	CH ₂	OCH ₂ (2-Cl-Ph)	
111	CH ₃	CH ₂	OCH ₂ (3-Cl-Ph)	
112	CH ₃	CH ₂	OCH ₂ (4-Cl-Ph)	
113	CH ₃	CH ₂	OCH ₂ (2-CH ₃ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
114	CH ₃	CH ₂	OCH ₂ (3-CH ₃ -Ph)	
115	CH ₃	CH ₂	OCH ₂ (4-t-C ₄ H ₉ -Ph)	
116	CH ₃	CH ₂	OCH ₂ (4-CF ₃ -Ph)	
117	CH ₃	CH ₂	OCH ₂ (2,4-(CH ₃) ₂ -Ph)	
118	CH ₃	CH ₂	OCH ₂ (3,4-(CH ₃) ₂ -Ph)	
119	CH ₃	CH ₂	OCH ₂ (4-OH-Ph)	
120	CH ₃	CH ₂	OCH ₂ (4-CH ₃ O-Ph)	
121	CH ₃	CH ₂	OCH ₂ (4-CF ₃ O-Ph)	
122	CH ₃	CH ₂	OCH ₂ (2,4-(CH ₃ O) ₂ -Ph)	
123	CH ₃	CH ₂	OCH ₂ (3,4-(CH ₃ O) ₂ -Ph)	
124	CH ₃	CH ₂	OCH ₂ (4-NH ₂ -Ph)	
125	CH ₃	CH ₂	OCH ₂ (4-NO ₂ -Ph)	
126	CH ₃	CH ₂	OCH ₂ (4-COOCH ₃ -Ph)	
127	CH ₃	CH ₂	OCH ₂ (4-COOC ₄ H ₉ -t-Ph)	
128	CH ₃	CH ₂	OCH ₂ (4-CN-Ph)	
129	CH ₃	CH ₂	OCH ₂ (4-CH ₃ -Q ₁)	nD 1.547(26.3 °C)
130	CH ₃	CH ₂	OPh	
131	CH ₃	CH ₂	O(2-Cl-Ph)	
132	CH ₃	CH ₂	O(3-Cl-Ph)	
133	CH ₃	CH ₂	O(4-Cl-Ph)	
134	CH ₃	CH ₂	O(2-CH ₃ -Ph)	
135	CH ₃	CH ₂	O(3-CH ₃ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
136	CH ₃	CH ₂	O(4-t-C ₄ H ₉ -Ph)	
137	CH ₃	CH ₂	O(4-CF ₃ -Ph)	
138	CH ₃	CH ₂	O(2, 4-(CH ₃) ₂ -Ph)	
139	CH ₃	CH ₂	O(3, 4-(CH ₃) ₂ -Ph)	
140	CH ₃	CH ₂	O(4-OH-Ph)	
141	CH ₃	CH ₂	O(4-CH ₃ O-Ph)	
142	CH ₃	CH ₂	O(4-CF ₃ O-Ph)	
143	CH ₃	CH ₂	O(2, 4-(CH ₃ O) ₂ -Ph)	
144	CH ₃	CH ₂	O(3, 4-(CH ₃ O) ₂ -Ph)	
145	CH ₃	CH ₂	O(4-NH ₂ -Ph)	
146	CH ₃	CH ₂	O(4-NO ₂ -Ph)	
147	CH ₃	CH ₂	O(4-COOCH ₃ -Ph)	
148	CH ₃	CH ₂	O(4-COOC ₄ H ₉ -t-Ph)	
149	CH ₃	CH ₂	O(4-CN-Ph)	
150	CH ₃	CH ₂	O-COCH ₃	nD 1.3580(25°C)
151	CH ₃	CH ₂	O-COC ₂ H ₅	
152	CH ₃	CH ₂	O-COC ₃ H ₇ -n	
153	CH ₃	CH ₂	O-COC ₃ H ₇ -i	
154	CH ₃	CH ₂	O-COC ₄ H ₉ -t	
155	CH ₃	CH ₂	O-COC ₄ H ₉ -s	
156	CH ₃	CH ₂	O-COC ₁₇ H ₃₅ -n	
157	CH ₃	CH ₂	O-COC ₆ H ₁₁ -cyclo	

Table 1 (Continued)

No	R ¹	A	B	Property
158	CH ₃	CH ₂	O-COCH=CH ₂	
159	CH ₃	CH ₂	O-COCH=CHPh	
160	CH ₃	CH ₂	O-COCH ₂ Cl	
161	CH ₃	CH ₂	O-COCF ₃	
162	CH ₃	CH ₂	O-COPh	m. p. 50-53 °C
163	CH ₃	CH ₂	O-CO(2-Cl-Ph)	
164	CH ₃	CH ₂	O-CO(3-Cl-Ph)	
165	CH ₃	CH ₂	O-CO(4-Cl-Ph)	
166	CH ₃	CH ₂	O-CO(2-CH ₃ -Ph)	
167	CH ₃	CH ₂	O-CO(3-CH ₃ -Ph)	
168	CH ₃	CH ₂	O-CO(4-t-C ₄ H ₉ -Ph)	
169	CH ₃	CH ₂	O-CO(4-CF ₃ -Ph)	
170	CH ₃	CH ₂	O-CO(2, 4-(CH ₃) ₂ -Ph)	
171	CH ₃	CH ₂	O-CO(3, 4-(CH ₃) ₂ -Ph)	
172	CH ₃	CH ₂	O-CO(4-CH ₃ O-Ph)	
173	CH ₃	CH ₂	O-CO(4-CF ₃ O-Ph)	
174	CH ₃	CH ₂	O-CO(2, 4-(CH ₃ O) ₂ -Ph)	
175	CH ₃	CH ₂	O-CO(3, 4-(CH ₃ O) ₂ -Ph)	
176	CH ₃	CH ₂	O-CO(4-NO ₂ -Ph)	
177	CH ₃	CH ₂	O-CO(4-COOCH ₃ -Ph)	
178	CH ₃	CH ₂	O-CO(4-COOC ₄ H ₉ -t-Ph)	
179	CH ₃	CH ₂	O-CO(4-CN-Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
180	CH ₃	CH ₂	O-COOC ₂ H ₅	nD 1.5041(26°C)
181	CH ₃	CH ₂	O-COOC ₄ H ₉ -i	
182	CH ₃	CH ₂	ON=C(CH ₃) ₂	
183	CH ₃	CH ₂	ON=C(CH ₃)C ₂ H ₅	
184	CH ₃	CH ₂	ON=C ₆ H ₁₀ -cyclo	
185	CH ₃	CH ₂	ON=CHPh	
186	CH ₃	CH ₂	ON=CH(2-Cl-Ph)	
187	CH ₃	CH ₂	ON=CH(3-Cl-Ph)	
188	CH ₃	CH ₂	ON=CH(4-Cl-Ph)	
189	CH ₃	CH ₂	ON=CH(2-CH ₃ -Ph)	
190	CH ₃	CH ₂	ON=CH(3-CH ₃ -Ph)	
191	CH ₃	CH ₂	ON=CH(4-C ₄ H ₉ -t-Ph)	
192	CH ₃	CH ₂	ON=CH(4-CF ₃ -Ph)	
193	CH ₃	CH ₂	ON=CH(2, 4-(CH ₃) ₂ -Ph)	
194	CH ₃	CH ₂	ON=CH(3, 4-(CH ₃) ₂ -Ph)	
195	CH ₃	CH ₂	ON=CH(4-OH-Ph)	
196	CH ₃	CH ₂	ON=CH(4-CH ₃ O-Ph)	
197	CH ₃	CH ₂	ON=CH(4-CF ₃ O-Ph)	
198	CH ₃	CH ₂	ON=CH(2, 4-(CH ₃ O) ₂ -Ph)	
199	CH ₃	CH ₂	ON=CH(3, 4-(CH ₃ O) ₂ -Ph)	
200	CH ₃	CH ₂	ON=CH(4-NH ₂ -Ph)	
201	CH ₃	CH ₂	ON=CH(4-NO ₂ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
202	CH ₃	CH ₂	ON=CH(4-COOCH ₃ -Ph)	
203	CH ₃	CH ₂	ON=C(CH ₃)Ph	
204	CH ₃	CH ₂	OSO ₂ CH ₃	
205	CH ₃	CH ₂	OSO ₂ CF ₃	
206	CH ₃	CH ₂	OSO ₂ Ph	
207	CH ₃	CH ₂	OSO ₂ (4-Br-Ph)	
208	CH ₃	CH ₂	OSO ₂ (4-CH ₃ -Ph)	Paste
209	CH ₃	CH ₂	SPh	nD 1.636(24.6 °C)
210	CH ₃	CH ₂	SOPh	
211	CH ₃	CH ₂	SO ₂ Ph	m. p. 118 °C
212	CH ₃	CH ₂	SO ₂ (4-CH ₃ -Ph)	
213	CH ₃	CH ₂	SO ₂ (4-NO ₂ -Ph)	
214	CH ₃	CH ₂	SCH ₂ Ph	nD 1.612(24.6 °C)
215	CH ₃	CH ₂	SO ₂ CH ₂ Ph	m. p. 116 °C
216	CH ₃	CH ₂	SCON(CH ₂ CH=CH ₂) ₂	
217	CH ₃	CH ₂	SCO-Q ₉	
218	CH ₃	CH ₂	SCOOC ₂ H ₅	
219	CH ₃	CH ₂	SCSN(C ₂ H ₅) ₂	
220	CH ₃	CH ₂	NH ₂ · HCl	m. p. >250°C
221	CH ₃	CH ₂	N(C ₂ H ₅) ₂	Paste
222	CH ₃	CH ₂	N(CH ₂ CH=CH ₂) ₂	Paste
223	CH ₃	CH ₂	NHPh	

Table 1 (Continued)

No	R ¹	A	B	Property
224	CH ₃	CH ₂	NH(4-CH ₃ -Ph)	
225	CH ₃	CH ₂	NHCOCH ₃	
226	CH ₃	CH ₂	NHCOC ₂ H ₅	
227	CH ₃	CH ₂	NHCOC ₃ H ₇ -n	
228	CH ₃	CH ₂	NHCOC ₃ H ₇ -i	
229	CH ₃	CH ₂	NHCOC ₄ H ₉ -t	m. p. 93-94 °C
230	CH ₃	CH ₂	NHCOC ₄ H ₉ -s	
231	CH ₃	CH ₂	NHCOC ₁₇ H ₃₅ -n	
232	CH ₃	CH ₂	NHCOC ₃ H ₅ -cyclo	m. p. 75-78 °C
233	CH ₃	CH ₂	NHCO(2,2-Cl ₂ -C ₃ H ₅ -cyclo)	
234	CH ₃	CH ₂	NHCOC ₆ H ₁₁ -cyclo	
235	CH ₃	CH ₂	NHCOCH=CH ₂	
236	CH ₃	CH ₂	NHCOCH=CHPh	
237	CH ₃	CH ₂	NHCOCH ₂ Cl	
238	CH ₃	CH ₂	NHCOCF ₃	
239	CH ₃	CH ₂	NHCOPh	m. p. 96°C
240	CH ₃	CH ₂	NHCO(2-Cl-Ph)	
241	CH ₃	CH ₂	NHCO(3-Cl-Ph)	
242	CH ₃	CH ₂	NHCO(4-Cl-Ph)	
243	CH ₃	CH ₂	NHCO(2-CH ₃ -Ph)	
244	CH ₃	CH ₂	NHCO(3-CH ₃ -Ph)	
245	CH ₃	CH ₂	NHCO(4-t-C ₄ H ₉ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
246	CH ₃	CH ₂	NHCO(3-CF ₃ -Ph)	
247	CH ₃	CH ₂	NHCO(4-CF ₃ -Ph)	
248	CH ₃	CH ₂	NHCO(2, 4-(CH ₃) ₂ -Ph)	
249	CH ₃	CH ₂	NHCO(3, 4-(CH ₃) ₂ -Ph)	
250	CH ₃	CH ₂	NHCO(4-OH-Ph)	
251	CH ₃	CH ₂	NHCO(4-CH ₃ O-Ph)	
252	CH ₃	CH ₂	NHCO(4-CF ₃ O-Ph)	
253	CH ₃	CH ₂	NHCO(2, 4-(CH ₃ O) ₂ -Ph)	
254	CH ₃	CH ₂	NHCO(3, 4-(CH ₃ O) ₂ -Ph)	
255	CH ₃	CH ₂	NHCO(4-NH ₂ -Ph)	
256	CH ₃	CH ₂	NHCO(4-NO ₂ -Ph)	
257	CH ₃	CH ₂	NHCO(4-COOCH ₃ -Ph)	
258	CH ₃	CH ₂	NHCO(2, 4-Cl ₂ -Ph)	
259	CH ₃	CH ₂	NHCO(3, 4-Cl ₂ -Ph)	
260	CH ₃	CH ₂	N(CHO) ₂	m. p. 70-75 °C
261	CH ₃	CH ₂	Q ₁₀	
262	CH ₃	CH ₂	3, 3, 4, 4, 5, 5-F ₆ -Q ₁₁	
263	CH ₃	CH ₂	Q ₁₂	m. p. 124-125 °C
264	CH ₃	CH ₂	NHCOOCH ₃	
265	CH ₃	CH ₂	NHCOOC ₂ H ₅	Paste
266	CH ₃	CH ₂	NHCOOC ₄ H ₉ -i	
267	CH ₃	CH ₂	NHCOOC ₄ H ₉ -t	

Table 1 (Continued)

No	R ¹	A	B	Property
268	CH ₃	CH ₂	NHCOO(4-NO ₂ -Ph)	
269	CH ₃	CH ₂	NHCOOPh	
270	CH ₃	CH ₂	NHCONHCH ₃	
271	CH ₃	CH ₂	NHCONHC ₂ H ₅	
272	CH ₃	CH ₂	NHCONHC ₃ H ₇ -i	
273	CH ₃	CH ₂	NHCONHPh	m. p. 145 °C
274	CH ₃	CH ₂	NHCONH(2-Cl-Ph)	
275	CH ₃	CH ₂	NHCONH(3-Cl-Ph)	
276	CH ₃	CH ₂	NHCONH(4-Cl-Ph)	
277	CH ₃	CH ₂	NHCONH(2-CH ₃ -Ph)	
278	CH ₃	CH ₂	NHCONH(3-CH ₃ -Ph)	
279	CH ₃	CH ₂	NHCONH(4-t-C ₄ H ₉ -Ph)	
280	CH ₃	CH ₂	NHCONH(3-CF ₃ -Ph)	
281	CH ₃	CH ₂	NHCONH(4-CF ₃ -Ph)	
282	CH ₃	CH ₂	NHCONH(2, 4-(CH ₃) ₂ -Ph)	
283	CH ₃	CH ₂	NHCONH(3, 4-(CH ₃) ₂ -Ph)	
284	CH ₃	CH ₂	NHCONH(4-OH-Ph)	
285	CH ₃	CH ₂	NHCONH(3-CH ₃ O-Ph)	
286	CH ₃	CH ₂	NHCONH(4-i-C ₃ H ₇ O-Ph)	
287	CH ₃	CH ₂	NHCONH(4-CF ₃ O-Ph)	
288	CH ₃	CH ₂	NHCONH(2, 4-(CH ₃ O) ₂ -Ph)	
289	CH ₃	CH ₂	NHCONH(3, 4-(CH ₃ O) ₂ -Ph)	

Table 1 (Continued)

No	R ¹	A	B	Property
290	CH ₃	CH ₂	NHCONH(4-NH ₂ -Ph)	
291	CH ₃	CH ₂	NHCONH(4-NO ₂ -Ph)	
292	CH ₃	CH ₂	NHCONH(4-COOCH ₃ -Ph)	
293	CH ₃	CH ₂	NHCONH(2, 4-Cl ₂ -Ph)	
294	CH ₃	CH ₂	NHCSNHCH ₃	
295	CH ₃	CH ₂	NHCSNHPh	m. p. 171-172 °C
296	CH ₃	CH ₂	NHC(=NCH ₃)OC ₂ H ₅	
297	CH ₃	CH ₂	NHC(=NPh)N(C ₂ H ₅) ₂	
298	CH ₃	CH ₂	NHC(=NPh)SCH ₃	
299	CH ₃	CH ₂	NHSO ₂ (4-CH ₃ -Ph)	Paste
300	CH ₃	CH ₂	NHNH ₂	
301	CH ₃	CH ₂	NHN=C(CH ₃) ₂	
302	CH ₃	CH ₂	NHN=CHPh	
303	CH ₃	CH ₂	NHN=C(CH ₃)Ph	
304	CH ₃	CH ₂	Q ₁₃	
305	CH ₃	CH ₂	Q ₁₄	
306	CH ₃	CH ₂	Q ₁₅	
307	CH ₃	CH ₂	CHO	
308	CH ₃	CH ₂	CN	
309	CH ₃	CH ₂	COOCH ₃	
310	CH ₃	CH ₂	COOC ₂ H ₅	
311	CH ₃	CH ₂	COOC ₃ H ₇ -i	

Table 1 (Continued)

No	R ¹	A	B	Property
312	CH ₃	CH ₂	COOC ₈ H _{17-n}	
313	CH ₃	CH ₂	COOCH ₂ CH=CH ₂	
314	CH ₃	CH ₂	CONH ₂	
315	CH ₃	CH ₂	CONHCH ₃	
316	CH ₃	CH ₂	CONHC ₃ H _{7-i}	
317	CH ₃	CH ₂	CON(CH ₃) ₂	
318	CH ₃	CH ₂	CO-Q ₁₆	
319	CH ₃	CH ₂	CO-Q ₁₇	
320	CH ₃	CH ₂	CO-Q ₁₈	
321	CH ₃	CH ₂	CO-Q ₁₉	
322	CH ₃	CH ₂	CH(COCH ₃) ₂	
323	CH ₃	CH ₂	CH(COOC ₂ H ₅) ₂	
324	CH ₃	CH ₂	C(NHCOCH ₃)(COOC ₂ H ₅) ₂	m. p. 93-97 °C
325	CH ₃	CH ₂	CH(SCH ₃)(SOCH ₃)	
326	CH ₃	CH ₂	Q ₂₀	
327	CH ₃	CHCl	COOCH ₃	nD 1.5456(23°C)
328	CH ₃	CHCH ₃	CN	
329	CH ₃	CHCH ₃	COOCH ₃	
330	CH ₃	C(CH ₃) ₂	CN	
331	CH ₃	C(CH ₃) ₂	COOCH ₃	
332	CH ₃	CHPh	Cl	nD 1.5979(21.8°C)
333	CH ₃	CHPh	OH	m. p. 93.2°C

Table 1 (Continued)

No	R ¹	A	B	Property
334	CH ₃	CHPh	OCH ₃	
335	CH ₃	CHPh	CN	
336	CH ₃	CHPh	COOH	
337	CH ₃	CH ₂ CH ₂	COOCH ₃	
338	CH ₃	CH=CH (E)	COOCH ₃	m. p. 70 °C
339	CH ₃	CH=CH (E)	CON(CH ₃) ₂	
340	CH ₃	CH=CH (E)	CONHPh	
341	CH ₃	CH=CH (Z)	COOCH ₃	m. p. 80-85°C
342	CH ₃	CH=CH (E, Z mixture)	CHO	m. p. 77-78°C
343	CH ₃	C=C(SCH ₃) ₂	CN	
344	CH ₃	Q ₂₁	CN	
345	CH ₃	C=C(COOCH ₃) ₂	H	
346	CH ₃	C=C(COOCH ₃)CN	H	Paste
347	CH ₃	Q ₂₂	CN	
348	C ₂ H ₅	CH ₂	Cl	
349	C ₂ H ₅	CH ₂	OH	
350	C ₂ H ₅	CH ₂	OCH ₃	
351	C ₂ H ₅	CH ₂	OCH ₂ Ph	
352	C ₂ H ₅	CH ₂	OCH ₂ (4-C ₂ H ₅ -Q ₁)	
353	C ₃ H ₇ -i	CH ₂	F	
354	C ₃ H ₇ -i	CH ₂	Cl	
355	C ₃ H ₇ -i	CH ₂	Br	

Table 1 (Continued)

No	R ¹	A	B	Property
356	C ₃ H ₇ -i	CH ₂	I	
357	C ₃ H ₇ -i	CH ₂	OH	
358	C ₃ H ₇ -i	CH ₂	OCH ₃	
359	C ₃ H ₇ -i	CH ₂	OCH ₂ Ph	
360	C ₃ H ₇ -i	CH ₂	OCH ₂ (4-i-C ₃ H ₇ -Q ₁)	
361	C ₃ H ₇ -i	CH ₂	NH ₂ · HCl	
362	C ₃ H ₇ -i	CH ₂	N(CH ₃) ₂	
363	C ₃ H ₇ -i	CH ₂	N(CH ₂ CH=CH ₂) ₂	
364	C ₃ H ₇ -i	CH ₂	NHCOCH ₃	
365	CH ₂ Cl	CH ₂	OH	Paste
366	CF ₃	CH ₂	OH	
367	Ph	—	Q ₄	m. p. 150 °C
368	Ph	—	Q ₅	m. p. 158 °C
369	Ph	—	5-(CH ₃) ₂ -Q ₅	m. p. 147 °C
370	Ph	—	4-CH ₃ -Q ₇	m. p. 161 °C
371	Ph	—	4-t-C ₄ H ₉ -Q ₇	m. p. 110 °C
372	Ph	—	4-COOCH ₃ -Q ₇	m. p. 124 °C
373	Ph	—	4-CONHCH ₃ -Q ₇	m. p. 187 °C
374	Ph	—	4-CH ₂ COOH-Q ₇	m. p. 137 °C
375	Ph	—	4-CH ₂ CONH ₂ -Q ₇	m. p. 137 °C
376	Ph	—	4-CH ₂ CONHCH ₃ -Q ₇	m. p. 154 °C
377	Ph	—	5-CH ₃ -Q _{2,3}	m. p. 85°C

Table 1 (Continued)

No	R ¹	A	B	Property
378	Ph	CH ₂	Cl	
379	Ph	CH ₂	OH	
380	Ph	CH ₂	OCH ₃	
381	Ph	CH ₂	OCH ₂ Ph	
382	Ph	CH ₂	OCH ₂ (4-Ph-Q ₁)	
383	Ph	CH ₂	COOCH ₃	
384	Ph	CH ₂	CON(CH ₃)Ph	
385	Ph	CH ₂	CONH(4-CH ₃ O-Ph)	m. p. 129 °C
386	2-Cl-Ph	CH ₂	OH	
387	3-Cl-Ph	CH ₂	OH	
388	4-Cl-Ph	CH ₂	OH	
389	4-CH ₃ -Ph	CH ₂	OH	
390	3-CH ₃ -Ph	CH ₂	OH	
391	4-CH ₃ -Ph	CH ₂	OH	
392	4-t-C ₄ H ₉ -Ph	CH ₂	OH	
393	2-CF ₃ -Ph	CH ₂	OH	
394	4-CF ₃ -Ph	CH ₂	OH	
395	2,4-(CH ₃) ₂ -Ph	CH ₂	OH	
396	3,4-(CH ₃) ₂ -Ph	CH ₂	OH	
397	4-OH-Ph	CH ₂	OH	
398	4-CH ₃ O-Ph	CH ₂	OH	
399	4-CF ₃ O-Ph	CH ₂	OH	

Table 1 (Continued)

No	R ¹	A	B	Property
400	4-PhO-Ph	CH ₂	OH	
401	2,4-(CH ₃ O) ₂ -Ph	CH ₂	OH	
402	3,4-(CH ₃ O) ₂ -Ph	CH ₂	OH	
403	4-COOCH ₃ -Ph	CH ₂	OH	
404	2,4-Cl ₂ -Ph	CH ₂	OH	
405	3,4-Cl ₂ -Ph	CH ₂	OH	
406	Q _{2 4}	CH ₂	OH	
407	Q _{2 5}	CH ₂	OH	
408	Q _{2 6}	CH ₂	OH	
409	Q _{2 7}	CH ₂	OH	m. p. 158 °C
410	Q _{2 8}	CH ₂	OH	
411	6-Cl-Q _{2 7}	CH ₂	OH	
412	Q _{2 7}	—	CN	m. p. 129 °C
413	4-F-Ph	CH ₂	O-CO-(4-Cl-Ph)	m. p. 79 °C
414	4-F-Ph	CH ₂	O-(4-CH ₃ O-Ph)	m. p. 86 °C
415	4-F-Ph	CH ₂	O-CO-CH ₂ -(4-CH ₃ -Ph)	m. p. 113 °C
416	4-F-Ph	CH ₂	O-CO-NH-Ph	m. p. 99 °C
417	4-F-Ph	CH ₂	ON=C(CH ₃)(4-Cl-Ph)	m. p. 91 °C
418	4-F-Ph	CH ₂	OH	m. p. 100 °C
419	4-F-Ph	CH ₂	Br	m. p. 88 °C
420	4-F-Ph	CH ₂	O-CO-CH ₃	m. p. 59 °C
421	Q _{2 6}	—	CN	m. p. 190 °C

Table 1 (Continued)

No	R ¹	A	B	Property
422	Ph	CH ₂	CONH-(4-CH ₃ O-Ph)	m. p. 129 °C
423	Ph	—	C=NOH(NH ₂)	m. p. 130 °C
424	Ph	—	CN	
425	CH ₃	—	Q ₈	Paste
426	CH ₃	—	5-CH ₃ -Q ₈	nD 1.5661(26.4°C)
427	CH ₃	—	SCH ₂ CH=CH ₂	Paste
428	CH ₃	—	SCH ₂ C≡CH	Paste
429	CH ₃	—	SCH ₂ -Ph	m. p. 89°C
430	CH ₃	—	SCH ₂ -(4-t-C ₄ H ₉ -Ph)	m. p. 79°C
431	CH ₃	—	S-C ₃ H ₇ -i	Paste
432	CH ₃	—	S-CH ₂ -Q ₂₇	m. p. 78°C
433	CH ₃	—	S-CH ₂ -Q ₂₈	m. p. 97°C

In some of the compounds shown in Table 1, property is expressed by a term of "Paste". Table 2 summarizes the data concerning these compounds.

Table 2

No	H-NMR[CDCl ₃ /TMS, δ value (ppm)]
208	2.46(s, 3H), 2.63(s, 3H), 5.35(s, 2H), 7.35(d, 2H), 7.77(d, 2H).
221	1.05(t, 6H), 2.54(q, 4H), 2.63(s, 3H), 3.78(s, 2H).
222	2.61(s, 3H), 3.10-3.12(m, 4H), 3.80(s, 2H), 5.16-5.24(m, 4H), 5.78-5.89(m, 2H).
265	1.25(t, 3H), 2.63(s, 3H), 4.15(q, 2H), 4.59(d, 2H), 5.2-5.35(br, 1H).
299	2.44(s, 3H), 2.58(s, 3H), 4.42(d, 2H), 5.37(br, t, 1H), 7.31(d, 2H), 7.70(d, 2H).
346	1.41(t, 3H), 2.90(s, 3H), 4.20(q, 2H), 8.34(s, 1H).
365	2.58(t, 1H), 5.03(s, 2H), 5.16(d, 2H).
426	3.08(s, 3H), 3.96(d, 2H), 5.27(d, 1H), 5.43(d, 1H), 6.00(m, 1H).
427	3.08(s, 3H), 5.23(d, 2H), 6.28(s, 1H).
430	1.55(d, 6H), 3.08(s, 3H), 4.02(m, 1H).

The 1,2,3-thiadiazole derivatives of the general formula (I) or salts thereof according to the present invention are useful for agricultural and horticultural disease control. For example, these
5 compounds have a very high controlling effect against various diseases, for instance, rice blast (Pyricularia oryzae), rice sheath blight (Rhizoctonia solani), rice helminthosporium leaf spot (Cochiobolus miyabeanus), powdery mildew of various host plants such as powdery
10 mildew of barley and wheat (Erysiphe graminis), oats crown rust (Puccinia coronata) and rust of other plants, tomato late blight (Phytophthora infestans) and late blight or Phytophthora rots of other plants, downy mildew of various plants such as cucumber downy mildew
15 (Pseudoperonospora cubensis) and grape downy mildew (Plasmopara viticola), apple scab (Venturia inaequalis), apple alternaria leaf spot (Alternaria mali), pear black spot (Alternaria kikuchiana), citrus melanose (Diaporthe citri), bacterial diseases including Pseudomonas
20 diseases such as cucumber bacterial blight (Pseudomonas syringae pv. lachrymans) and tomato bacterial wilt (Pseudomonas solanacearum), Xanthomonas diseases such as cabbage black rot (Xanthomonas campestris), rice bacterial leaf blight (Xanthomonas oryzae) and citrus
25 canker (Xanthomonas citri) and Erwinia diseases such as cabbage bacterial soft rot (Erwinia carotovora), and viral diseases such as tobacco mosaic (Tobacco mosaic virus).

The agrohorticultural disease controller containing the 1,2,3-thiadiazole derivative of general formula (I) or salts thereof as an active ingredient according to the present invention exhibits a marked
5 controlling effect against the above-exemplified diseases which damage paddy field crops, upland field crops, fruit trees, vegetables, other crop plants, flowers, ornamental plants, and the like. Accordingly, the desired effects of the agrohorticultural disease
10 controller of the present invention can be exhibited by applying the disease controller to the paddy field water, stalks and leaves or soil of the paddy field, upland field, fruit trees, vegetables, other crops, flowers and ornamental plants at a season at which the
15 diseases are expected to occur, either before their occurrence or at the time when their occurrence has been confirmed.

The agrohorticultural disease controller of the present invention is generally put to use after
20 being prepared into a conveniently usable form according to an ordinary manner for preparation of pesticides.

That is, the 1,2,3-thiadiazole derivative represented by the general formula (I) or a salt thereof according to the present invention and, optionally, an
25 adjuvant are blended with an appropriate inert carrier in a proper proportion and dissolved, separated, suspended, mixed, impregnated, adsorbed or stuck, prepared into a suitable preparation form such as

suspension, emulsifiable concentrate, soluble concentrate, wettable powder, granule, dust, tablet, etc. and put to use.

The inert carrier used in the present invention may be either solid or liquid. As the material usable as solid carrier, there can be exemplified soybean flour, cereal flour, wood flour, bark flour, saw dust, powdered tobacco stalks, powdered walnut shells, bran, powdered cellulose, extraction residue of vegetables, powdered synthetic polymers of synthetic resins and the like, clays such as kaolin, bentonite, acid clay and the like, talcs such as talc, pyrophyllite and the like, silica materials such as diatomaceous earth, siliceous sand, mica, white carbon (namely, synthetic, high-dispersion silicic acid, also called finely divided hydrated silica or hydrated silicic acid, some of the commercially available products thereof contain calcium silicate as the major component), activated carbon, powdered sulfur, pumice, calcined diatomaceous earth, crushed brick, fly ash, sand, inorganic mineral powders such as calcium carbonate, calcium phosphate and the like, chemical fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, ammonium chloride and the like, and compost. These carriers may be used alone or as a mixture thereof.

The material constituting the liquid carrier is selected from materials having a dissolving ability

25711-791

55

in themselves and materials which have no dissolving ability in themselves but can disperse the active ingredient compound by the aid of an adjuvant. The following are typical examples of the liquid carrier material, which can be used either alone or in the form of a mixture: water; alcohols such as methanol, ethanol, isopropanol, butanol, ethylene glycol and the like; ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone, diisobutyl ketone, cyclohexanone and the like; ethers such as ethyl ether, dioxane, Cellosolve*, dipropyl ether, tetrahydrofuran and the like; aliphatic hydrocarbons such as kerosene, mineral oils and the like; aromatic hydrocarbons such as benzene, toluene, xylene, solvent naphtha, alkyl-naphthalene and the like; halogenated hydrocarbons such as dichloroethene, chloroform, carbon tetrachloride, chlorobenzene and the like; esters such as ethyl acetate, diisopropyl phthalate, dibutyl phthalate, dioctyl phthalate and the like; amides such as dimethylformamide, diethylformamide, dimethylacetamide and the like; nitriles such as acetonitrile and the like; dimethyl sulfoxide; etc.

As the adjuvant, the following can be referred to as typical ones. These adjuvants are used depending upon purposes, either alone or in combination of two or more in some cases. It is also possible to use no adjuvant at all, in some cases.

A surfactant is used for the purpose of

*Trade-mark

emulsifying, dispersing, solubilizing and/or wetting an active ingredient compound. As the surfactant, there can be exemplified polyoxyethylene alkyl ethers, polyoxyethylene alkylaryl ethers, polyoxyethylene higher fatty acid esters, polyoxyethylene resin acid esters, polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate, alkylarylsulfonates, naphthalene-sulfonic acid condensates, ligninsulfonic acid salts, higher alcohol sulfuric esters, etc.

The following adjuvants can also be used for the purpose of stabilizing, tackifying and/or binding the dispersion of active ingredient compound: casein, gelatin, starch, methyl cellulose, carboxymethyl cellulose, gum arabic, polyvinyl alcohol, turpentine, bran oil, bentonite, ligninsulfonic acid salts and the like.

Further, wax, stearic acid salts, alkyl phosphates and the like may be used as an adjuvant for the purpose of improving flowability of a solid product.

Further, naphthalenesulfonic acid condensates, polycondensed phosphoric acid salts and the like may also be used as a peptizer for dispersible products.

Adjuvants such as silicone oils may also be used as a defoaming agent.

The content of the active ingredient compound may be increased or decreased according to the need. For example, in dusts and granules, the suitable content thereof is from 0.01 to 50% by weight. In emulsifiable

concentrates and wettable powders, too, the suitable content thereof is from 0.01 to 50% by weight.

For controlling various diseases, the agrohorticultural disease controller of the present invention itself or its appropriate dilution or suspension in water or the like is applied to a crop on which the diseases are expected to occur or a site where occurrence of the diseases is undesirable, in an amount effective for disease control. For example, for controlling the diseases of paddy rice, said disease controller can be used by a method such as direct application to regular paddy field, application to a rice nursery bed, dressing of seeds for direct sowing to flooded paddy field, seed disinfection, etc. For controlling the diseases of barley, wheat, oat or the like, the disease controller of the present invention is applied to stalks and leaves or used for soil treatment where the disease controller is absorbed from the roots.


The application amount of the agrohorticultural disease controller of the present invention may vary depending on various factors including purpose of application, objective disease, state of plant growth, tendency of prevalence of the disease, weather, environmental conditions, preparation form, method of application, site of application, time of application, etc. The application amount, however, should be properly chosen in the range of from 0.1 g to 10 kg per 10 ares as expressed in terms of amount of

active ingredient compound, depending on purposes.

In order to expand the spectrum of controllable diseases and the time period of effective application or to reduce the dosage, it is also possible to use the disease controller of the present invention in the form of a mixture with other agrohorticultural disease controllers.

Next, typical examples, formulation examples and test examples of the 1,2,3-thiadiazole derivatives of general formula (I) or salts thereof according to the present invention are presented below. The present invention is by no means limited by these examples.

Example 1: Production of 4-methyl-1,2,3-thiadiazole-5-carbaldehyde (Compound No. 7)




In 25 ml of methylene chloride were suspended 9.9 g (46 mmol) of pyridinium chlorochromate and 10 g of silica gel. After adding 3.0 g (23 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol, the mixture was stirred at room temperature for 4 hours to make a reaction progress. After the reaction was completed, the reaction mixture was diluted with ether, the resulting florisil was filtered off, the filtrate was concentrated

under reduced pressure, and the residue was purified by silica gel column chromatography using 3:1 mixture of n-hexane and ethyl acetate. Thus, 2.0 g of the objective compound was obtained.

5 Property: nD 1.5555 (26°C)

Yield: 62%

Example 2: Production of 5-benzoyl-4-methyl-1,2,3-thiadiazole (Compound No. 9)



To 2.0 g (15 mmol) of thionyl chloride was
10 added 0.50 g (3.5 mmol) of 4-methyl-1,2,3-thiadiazole-5-carboxylic acid. After heating the mixture with stirring for 2 hours, the thionyl chloride was distilled off under reduced pressure, and the residue was added to a solution of 1.8 g (15 mmol) of aluminum chloride in 20
15 ml benzene and made to react with heating for 3 hours. After the reaction was completed, the reaction mixture was poured into ice water, the objective product was extracted with ethyl acetate, the organic layer was washed with a saturated aqueous solution of sodium
20 chloride and dried, the solvent was distilled off under reduced pressure, and the residue was purified by silica

25711-791

60


gel column chromatography using 4:1 mixture of n-hexane and ethyl acetate. Thus, 0.69 g of the objective compound was obtained.

Property: n_D 1.6145 (22°C)

5

Yield: 97%

Example 3: Production of O-benzyl-4-methyl-1,2,3-thiadiazole-5-carbaldehyde oxime (Compound No. 21)




To 10 ml of ethanol were added 0.40 g (3.1 mmol) of 4-methyl-1,2,3-thiadiazole-5-carbaldehyde
10 obtained in Example 1, 0.60 g (3.8 mmol) of benzyloxyamine hydrochloride and 0.50 g (3.8 mmol) of sodium acetate. The mixture was stirred for 4 hours with heating under reflux. After the reaction was completed, the reaction mixture was diluted with ethyl
15 acetate and filtered to remove the precipitate, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography using 4:1 mixture of n-hexane and ethyl acetate. Thus, 0.66 g of the objective compound was
20 obtained.

Property: n_D 1.5998 (25°C)

61

Yield: 91%

Example 4: Production of 5-(5-mercapto-1,3,4-oxadiazol-2-yl)-4-methyl-1,2,3-thiadiazole (Compound No. 102)




5 A solution of 0.3 g of potassium hydroxide in 1 ml water and 0.6 ml of carbon disulfide were successively added to a solution of 1.1 g (7.2 mmol) of 4-methyl-1,2,3-thiadiazole-5-carbohydrazide in 15 ml ethanol. The mixture was stirred for 6.5 hours with
 10 heating under reflux. After the reaction was completed, the solvent was distilled off under reduced pressure, water was added to the residue, acetic acid was added for the sake of acidification, and the deposited
 15 0.82 g of the objective compound was obtained.

Property: m.p. 182°C

Yield: 59%

Example 5: Production of 4-methyl-5-(5-methylthio-1,3,4-oxadiazol-2-yl)-1,2,3-thiadiazole (Compound No.
 20 103)




To 10 ml of acetone were added 0.40 g (2.0 mmol) of 5-(5-mercapto-1,3,4-thiadiazol-2-yl)-4-methyl-1,2,3-thiadiazole, 0.43 g (3.0 mmol) of methyl iodide and 0.83 g (6.0 mmol) of potassium carbonate. The mixture was stirred for 5 hours with heating under reflux. After the reaction was completed, the reaction mixture was filtered to remove the insoluble matter, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography using 2:1 mixture of n-hexane and ethyl acetate. Thus, 0.41 g of the objective compound was obtained.

Property: m.p. 65°C

Yield 96%

Example 6: Production of 5-chloromethyl-4-methyl-1,2,3-thiadiazole (Compound No. 104)




While cooling the reaction system with ice, 2 ml of thionyl chloride was added to 0.50 g (3.8 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol. Then, at room temperature, the mixture was stirred for 2 hours. Then, 5 water was added to stop the reaction, the objective product was extracted with ethyl acetate, the organic layer was washed with water and dried, the solvent was distilled off under reduced pressure, and the residue thus obtained was purified by silica gel column 10 chromatography using 3:1 mixture of n-hexane and ethyl acetate. Thus, 0.46 g of the objective compound was obtained.

Property: n_D 1.5590 (26°C)

Yield: 81%

15 Example 7: Production of 4-methyl-1,2,3-thiadiazol-5-ylmethanol (Compound No. 105)




To a solution of 0.10 g (2.5 mmol) of sodium hydroxide in 10 ml ethanol was added 1.0 g (5.8 mmol) of ethyl 4-methyl-1,2,3-thiadiazole-5-carboxylate. Then, 20 0.33 g (8.7 mmol) of sodium borohydride was added and stirred at room temperature. After the reaction was

completed, water was added to the reaction mixture to decompose the excessive sodium borohydride, after which the objective product was extracted with ethyl acetate. The organic layer was washed with water and dried, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 2:1 mixture of n-hexane and ethyl acetate. Thus, 0.36 g of the objective compound was obtained.

Property: n_D 1.553 (21°C)

Yield: 48%

Example 8: Production of 4-methyl-5-benzyloxymethyl-1,2,3-thiadiazole (Compound No. 109)



In 10 ml of acetone was dissolved 0.40 g (3.1 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol. After adding 1.6 g (12 mmol) of potassium carbonate and 0.80 g (4.5 mmol) of benzyl bromide, the resulting mixture was stirred for 10 hours with heating under reflux. After the reaction was completed, the insoluble matter was filtered off, the solvent was distilled off from the filtrate under reduced pressure, and the residue was purified by silica gel column chromatography using 3:1


65

mixture of n-hexane and ethyl acetate. Thus, 0.65 g of the objective compound was obtained.

Property: n_D 1.5730 (25°C)

Yield 96%

- 5 Example 9: Production of 4,4'-dimethyl-5,5'-oxybis(methyl-1,2,3-thiadiazole) (Compound No. 129)



In 20 ml of tetrahydrofuran were dissolved 2.0 g (15 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol and 3.9 g (39 mmol) of triethylamine. While cooling the system with ice, 3.5 g (18 mmol) of tosyl chloride was added and stirred at room temperature for 24 hours. Then, a saturated aqueous solution of sodium chloride was added to the reaction mixture to stop the reaction, the objective product was extracted with ethyl acetate, the extract solution was washed successively with dilute hydrochloric acid, a saturated aqueous solution of sodium hydrogen carbonate and a saturated aqueous solution of sodium chloride, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 4:1 mixture of n-hexane and ethyl acetate. Thus, 1.1 g of the


66

objective compound was obtained.

Property: n_D 1.547 (26°C)

Yield: 57%

Example 10: Production of 4-methyl-1,2,3-thiadiazol-5-ylmethyl acetate (Compound No.150)




In 10 ml of tetrahydrofuran were dissolved 0.50 g (3.8 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol, 0.50 g (5.7 mmol) of acetyl chloride and 0.80 g (7.6 mmol) of triethylamine. The solution thus obtained was stirred at room temperature for 18 hours. Then, a saturated aqueous solution of sodium chloride was added to the reaction mixture to stop the reaction, the objective product was extracted with ethyl acetate, the organic layer was washed with water and dried, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 3:1 mixture of n-hexane and ethyl acetate. Thus, 0.70 g of the objective compound was obtained.

Property: n_D 1.3580 (25°C)

Yield: 100%

Example 11: Production of ethyl 4-methyl-1,2,3-thiadiazol-5-ylmethyl carbonate (Compound No. 180)




In 5 ml of tetrahydrofuran was dissolved 0.50 g (3.8 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol. After adding 0.80 g (7.6 mmol) of triethylamine and 0.60 g (5.7 mmol) of ethyl chlorocarbonate, the resulting mixture was stirred at room temperature for 22 hours. Then, water was added to the reaction mixture to stop the reaction, the objective product was extracted with ethyl acetate, the organic layer was washed with water and dried, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 5:1 mixture of n-hexane and ethyl acetate. Thus, 0.50 g of the objective compound was obtained.

Property: n_D 1.5041 (26°C)

Yield: 68%

Example 12: Production of 4-methyl-5-phenylthiomethyl-1,2,3-thiadiazole (Compound No. 209)




At an ice-cooled temperature, 3 ml of thionyl chloride was added to 0.50 g (3.8 mmol) of 4-methyl-1,2,3-thiadiazol-5-ylmethanol and stirred for 4 hours. Then, the reaction mixture was concentrated under reduced pressure, and 1.30 g (11.5 mmol) of potassium carbonate, 0.38 g (3.4 mmol) of thiophenol and 10 ml of acetone were added to the concentrate and reacted for 7 hours with heating under reflux. After the reaction was completed, the reaction mixture was cooled to room temperature, the precipitated matter was filtered off, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 5:1 mixture of n-hexane and ethyl acetate. Thus, 0.64 g of the objective compound was obtained.

Property: n_D 1.636 (25°C)

Yield: 85%

Example 13: Production of 4-methyl-5-phenylsulfonfylmethyl-1,2,3-thiadiazole (Compound No. 211)




In 5.0 ml of methylene chloride was dissolved 0.20 g (0.99 mmol) of 4-methyl-5-phenylthiomethyl-1,2,3-thiadiazole. Then, 0.40 g (2.3 mmol) of m-chloroperbenzoic acid was added to the solution obtained
 5 above, and was stirred for 20 hours at room temperature. Then, an aqueous solution of sodium hydrogen sulfite was added to the reaction mixture to stop the reaction, the objective product was extracted with ethyl acetate, the organic layer was washed with a saturated aqueous
 10 solution of sodium hydrogen carbonate, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 2:1 mixture of n-hexane and ethyl acetate. Thus, 0.23 g of the objective compound was obtained.

15 Property: m.p. 118°C

Yield: 100%

Example 14: Production of 5-aminomethyl-4-methyl-1,2,3-thiadiazole hydrochloride (Compound No. 220)




To 50 ml of acetonitrile were added 7.0 g (47 mmol) of 5-chloromethyl-4-methyl-1,2,3-thiadiazole and 9.0 g (94 mmol) of sodium formimide. After stirring the resulting mixture for 16 hours with heating under
 5 reflux, the reaction mixture was cooled to the room temperature, a saturated aqueous solution of sodium chloride was added, the objective product was extracted with ethyl acetate, the organic layer was washed with water and dried, and the solvent was distilled off under
 10 reduced pressure. Then, while stirring the resulting crystalline product together with 40 ml of water, 30 ml of concentrated hydrochloric acid was added. After ten minutes, water and hydrochloric acid were distilled off, and the residue was washed with ethyl acetate. Thus,
 15 6.6 g of the objective compound was obtained.

Property: m.p. > 250°C

Yield: 77%

Example 15: Production of 5-(N,N-diallylamino)methyl-4-methyl-1,2,3-thiadiazole (Compound No. 222)




To 15 ml of acetone were added 0.50 g (3.0 mmol) of 5-aminomethyl-4-methyl-1,2,3-thiadiazole hydrochloride, 1.10 g (9.0 mmol) of allyl bromide and 2.1 g (15 mmol) of potassium carbonate. The resulting mixture was stirred for 20 hours with heating under reflux. After the reaction was completed, the reaction mixture was filtered to remove the precipitated matter, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 3:1 mixture of n-hexane and ethyl acetate. Thus, 0.42 g of the objective compound was obtained.

Property: Pasty product

Yield: 67%

H-NMR [CDCl₃/TMS, δ value (ppm)] 2.61 (s, 3H), 3.10-3.12 (m, 4H), 3.80 (s, 2H), 5.16-5.24 (m, 4H), 5.78-5.89 (m, 2H)

Example 16: Production of methyl (E)-3-(4-methyl-1,2,3-thiadiazol-5-yl)-2-propenate (Compound No. 338) and methyl (Z)-3-(4-methyl-1,2,3-thiadiazol-5-yl)-2-propenate (Compound No. 341)



To 0.40 g (3.1 mmol) of 4-methyl-1,2,3-thiadiazol-5-carbaldehyde dissolved in tetrahydrofuran was added 1.1 g (3.4 mmol) of methyl triphenylphosphol-anilideneacetate. The solution thus obtained was

5 stirred for 16 hours at room temperature. After the reaction was completed, the reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium chloride was added, and the objective product was extracted with ethyl acetate. The organic layer was


10 washed with water and dried, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 5:1 mixture of n-hexane and ethyl acetate. Thus, the E-compound and the Z-compound, both the objective compounds, were

15 obtained in yields of 0.49 g and 0.05 g, respectively.

Property: E-compound m.p. 70°C, Yield 90%

Z-compound m.p. 80-85°C, Yield 10%

Example 17: Production of ethyl 2-cyano-3-(4-methyl-1,2,3-thiadiazol-5-yl)-2-propenate (Compound No. 346)




To 50 ml of toluene were added 1.5 g (12 mmol) of 4-methyl-1,2,3-thiadiazole-5-carbaldehyde, 1.5 g (13 mmol) of ethyl cyanoacetate, 0.40 g (6.0 mmol) of pyrrolidine and 0.20 g (1.2 mmol) of tosylic acid. In a reactor equipped with a Dean-Stark dehydrating apparatus, the mixture obtained above was stirred for 2 hours with heating under reflux. After the reaction was completed, a saturated aqueous solution of sodium chloride was added to the reaction mixture, the objective product was extracted with ethyl acetate, the organic layer was washed successively with dilute hydrochloric acid, saturated aqueous solution of sodium hydrogen carbonate and saturated aqueous solution of sodium chloride, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography using 3:1 mixture of n-hexane and ethyl acetate. Thus, 0.73 g of the objective compound was obtained.

Property: Pasty product,

Yield: 30%

H-NMR [CDCl_3/TMS , δ value (ppm)] 1.41 (t, 3H), 2.90 (s, 3H), 4.20 (q, 2H), 8.34 (s, 1H)

Example 18: Production of 4-phenyl-3,4,5,6-tetrahydropyrimidin-2-yl-1,2,3-thiadiazole (Compound No. 368)




In 10 ml of ethanol were dissolved 0.95 g (5.0 mmol) of 4-phenyl-1,2,3-thiadiazole-5-carbonitrile and 3.8 g (50 mmol) of 1,3-propanediamine. After adding 0.05 g of sodium ethoxide, the solution thus obtained was stirred for 8 hours with heating under reflux. After the reaction was completed, the reaction mixture was cooled to room temperature, the solvent was distilled off under reduced pressure, and the objective product was extracted with ethyl acetate. The organic layer was concentrated under reduced pressure, and the oily product thus obtained was recrystallized from a small quantity of ethyl acetate to obtain 0.68 g of the objective compound.

Property: m.p. 158°C

Yield: 55%

Example 19: Production of methyl 2-(4-phenyl-1,2,3-thiadiazol-5-yl)thiazole-4-carboxylate (Compound No. 372)



In 20 ml of chlorobenzene were dissolved 1.1 g (5.0 mmol) of 4-phenyl-1,2,3-thiadiazole-5-carbothioamide and 0.92 g (5.0 mmol) of methyl bromopyruvate. The solution thus obtained was stirred for 3 hours with heating under reflux. After the reaction was completed, the reaction mixture was cooled to room temperature, a saturated aqueous solution of sodium hydrogen carbonate was added thereto, and the objective product was extracted with ethyl acetate. The solvent was distilled off from the extract solution under a reduced pressure, and the residue was purified by silica gel column chromatography using 3:1 mixture of n-hexane and ethyl acetate. Thus, 0.74 g of the objective compound was obtained.

Property: m.p. 124°C

Yield: 41%

Next, typical formulation examples and test examples of the present invention are mentioned below.

In the formulation examples, the term "parts" means "parts by weight".

Formulation Example 1

	The compound shown in Table 1	50 parts
	Xylene	40 parts
5	Mixture of polyoxyethylene nonyl-phenyl ether and calcium alkyl-benzenesulfonate	10 parts

The ingredients mentioned above are uniformly mixed and dissolved together to prepare an emulsifiable concentrate.

10 Formulation Example 2

	The compound shown in Table 1	3 parts
	Clay powder	82 parts
	Diatomaceous earth powder	15 parts

15 The ingredients mentioned above are uniformly mixed together and pulverized to prepare a dust.

Formulation Example 3

	The compound shown in Table 1	5 parts
	Powdery mixture of bentonite and clay	90 parts
20	Calcium ligninsulfonate	5 parts

The ingredient mentioned above are kneaded together with a proper quantity of water, and then granulated and dried to prepare a granular composition.

Formulation Example 4

25	The compound shown in Table 1	20 parts
----	-------------------------------	----------

Kaolinite and synthetic high-
dispersion silicic acid 75 parts
Mixture of polyoxyethylene
nonylphenyl ether and calcium
5 alkylbenzenesulfonate 5 parts

The ingredients mentioned above are uniformly
mixed together and pulverized to prepare a wettable
powder.

Test Example 1: Rice blast-controlling test by submerged
10 application

A chemical agent containing the compound shown
in Table 1 as an active ingredient was applied to paddy
rice plants of the 5- to 6-leaved stage, cultivated in
1/10000 are pots, by the method of submerged application
15 at a dosage of 200 g/10 ares as expressed in terms of
active ingredient. After standing in a greenhouse for a
week, the plants were inoculated with a spore suspension
of rice blast fungus (Pyricularia oryzae) by the method
of spraying.

20 After the inoculation, the plants were allowed
to stand in a moist chamber for one day and then in a
greenhouse for 6 days to cause the disease sufficiently.
Then, lesions on each leaf were counted and compared
with those in the untreated plot, from which the
25 controlling degree was calculated, whereby the effect
was judged according to the following criterion:

78

Effect	Controlling degree (%)
A	100-95
B	94-85
C	84-60
D	59- 0


The results of the above test demonstrate that the compounds listed in Table 1 were found to have a marked blast-controlling activity. Of these compounds, the following were rated C or higher: Compound Nos. 7, 17-19, 21, 24, 35, 36, 60, 62, 66-69, 77, 83, 84, 87-89, 91, 93-100, 102, 104-106, 109, 129, 150, 162, 180, 209, 220-222, 229, 239, 260, 263, 273, 295, 299, 327, 338, 341, 342, 346, 365, 367, 369, 371 and 376. Especially, the following exhibited so excellent a blast-controlling activity as rated A: Compound Nos. 7, 37, 88, 89, 91, 104, 105, 109, 129, 150, 220, 221, 222, 239, 260, 263, 273, 299, 327, 342, 367 and 371.

25711-791

79

CLAIMS:

1. A 1,2,3-thiadiazole derivative represented by the following general formula (I) or a salt thereof:



wherein:

R^1 represents (1) a hydrogen atom; (2) a C_1-C_8 alkyl group; (3) a halo C_1-C_4 alkyl group; (4) a phenyl group; (5) a substituted phenyl group having 1 to 5 same or
 10 different substituents selected from the group consisting of halogen atom, hydroxyl group, cyano group, nitro group, C_1-C_4 alkyl group, halo C_1-C_4 alkyl group, C_1-C_4 alkoxy group, halo C_1-C_4 alkoxy group and C_1-C_4 alkoxycarbonyl group; (6) a 5- or 6-membered heterocycle containing 1 to 3 same or
 15 different heteroatoms selected from an oxygen atom, a nitrogen atom and a sulfur atom; or (7) a substituted 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from an oxygen atom, a nitrogen atom and a sulfur atom and having 1 to 4 same or different
 20 substituents selected from the group consisting of a halogen atom, a cyano group, hydroxyl group, a mercapto group, a C_1-C_4 alkyl group, a C_1-C_4 alkoxy group, a C_1-C_4 alkylthio group, a formyl group, a carboxyl group, a carboxy C_1-C_4 alkyl group, a C_1-C_4 alkoxycarbonyl group, a C_1-C_4
 25 alkoxycarbonyl C_1-C_4 alkyl group, an amino group, a substituted amino group having 1 or 2 same or different substituents selected from the group consisting of C_1-C_4 alkyl and phenyl groups, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents
 30 selected from the group consisting of C_1-C_4 alkyl and phenyl groups, a carbamoyl C_1-C_4 alkyl group, and a substituted

25711-791

80

carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups;

A represents a group of the following formula:



(wherein:

R² and R³, which are the same or different,
 10 represent (1) a hydrogen atom; (2) a halogen atom; (3) a cyano group; (4) a formyl group; (5) a C₁-C₄ alkyl group; (6) a C₁-C₄ alkoxy group; (7) a C₁-C₄ alkylthio group; (8) a C₁-C₄ alkylsulfinyl group; (9) a C₁-C₄ alkylsulfonyl group; (10) a C₁-C₄ alkylcarbonyl group; (11) a C₁-C₄ alkoxycarbonyl group;
 15 (12) a phenyl group; (13) a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group
 20 and a C₁-C₄ alkoxycarbonyl group; (14) a phenylcarbonyl group; or (15) a substituted phenylcarbonyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄ alkyl group
 25 and a C₁-C₄ alkoxycarbonyl group; or

R² and R³ may be taken conjointly to form a 3- to 7-membered ring including a C₂-C₆ alkylene group which may be intercepted by -O-, -S(O)_n- in which n is an integer of 0 to 2, -CO- or -NR⁸- in which R⁸ represents (1) a hydrogen atom;
 30 (2) a formyl group; (3) a C₁-C₈ alkyl group; (4) a halo C₁-C₄

25711-791

81


alkyl group; (5) a C₂-C₄ alkenyl group; (6) a halo C₂-C₄ alkenyl group; (7) a C₂-C₄ alkynyl group; (8) a halo C₂-C₄ alkynyl group; (9) a C₁-C₄ alkylthio group; (10) a C₁-C₄ alkylsulfinyl group; (11) a C₁-C₄ alkylsulfonyl group; (12) a C₁-C₄ alkylcarbonyl group; (13) a C₁-C₄ alkoxycarbonyl group; (14) a carbamoyl group; (15) a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of a C₁-C₄ alkyl group and a phenyl group; (16) a phenyl group; (17) a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; (18) a phenyl C₁-C₄ alkyl group; (19) a substituted phenyl C₁-C₄ alkyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; (20) a phenylcarbonyl group; or (21) a substituted phenylcarbonyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group, wherein the 3- to 7-membered ring may be substituted by one or more same or different substituents selected from the group consisting of a halogen atom and a C₁-C₄ alkyl group; and

m represents an integer of 0 to 4), or

A represents a group of the formula:

25711-791

82



in which R^2 and R^3 are as defined above; and

5 when A is $-(\text{CR}^2\text{R}^3)_m-$ and m is 0, then B represents:

(1) a cyano group;

(2) a 5- or 6-membered heterocycle selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole,

10 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine, pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithialane, dithiane, piperazine, dioxolane and imidazolidine;


15 (3) a substituted 5- or 6-membered heterocycle selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine, pyrazine, pyrrolidine, piperidine, morpholine, 20 thiomorpholine, dithialane, dithiane, piperazine, dioxolane and imidazolidine and having 1 to 4 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a mercapto group, a C_1 - C_4 alkyl group, a C_1 - C_4 alkoxy group, a C_1 - C_4 25 alkylthio group, a carboxyl group, a formyl group, a carboxy C_1 - C_4 alkyl group, a C_1 - C_4 alkoxycarbonyl group, a C_1 - C_4 alkoxycarbonyl C_1 - C_4 alkyl group, an amino group, a substituted amino group having 1 or 2 same or different C_1 - C_4 30 alkyl or phenyl groups, a carbamoyl group, a substituted

25711-791

83

carbamoyl group having 1 or 2 C₁-C₄ alkyl or phenyl groups, a carbamoyl C₁-C₄ alkyl group and a substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different C₁-C₄ alkyl or phenyl groups; or

5 (4) a group of the formula:



(in which:

R⁴ represents a hydrogen atom; a C₁-C₈ alkyl group; a halo C₁-C₄ alkyl group; a C₂-C₄ alkenyl group; a halo C₂-C₄ alkenyl group; a C₂-C₄ alkynyl group; a halo C₂-C₄ alkynyl group; a phenyl group; a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxy carbonyl group; a phenyl C₁-C₄ alkyl group; a substituted phenyl C₁-C₄ alkyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxy carbonyl group; a 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom; or a substituted 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom and having 1 to 4 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a mercapto group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a C₁-C₄ alkylthio group, a formyl group, a carboxyl group, a carboxy C₁-C₄ alkyl group, a C₁-C₄ alkoxy carbonyl group, a C₁-C₄ alkoxy carbonyl C₁-C₄ alkyl group, an amino group, a substituted amino group

25711-791

84

having 1 or 2 same or different substituents selected from the group consisting of a C₁-C₄ alkyl group and a phenyl group, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of a C₁-C₄ alkyl group and a phenyl group, a carbamoyl C₁-C₄ alkyl group, and a substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of a C₁-C₄ alkyl group and a phenyl group; and

10 X represents O; S; N-R⁶; NO-R⁶; N(→O)-R⁶; NN(R⁶)R⁷; or NN=C(R⁶)R⁷ (in which R⁶ and R⁷, same or different, represent a hydrogen atom; a halogen atom; a formyl group; a C₁-C₄ alkyl group; a C₂-C₄ alkenyl group; a C₂-C₄ alkynyl group; a C₁-C₄ alkylsulfonyl group; a halo C₁-C₄ alkylsulfonyl group; a C₁-C₂₀ alkylcarbonyl group; a C₁-C₄ alkoxycarbonyl group; a phenyl group; a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group, an amino group and a C₁-C₄ alkoxycarbonyl group; a phenyl C₁-C₄ alkyl group; a substituted phenyl C₁-C₄ alkyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group, an amino group and a C₁-C₄ alkoxycarbonyl group; a phenylcarbonyl group; a substituted phenylcarbonyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a carbamoyl group; a

25711-791

85

substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of a C₁-C₄ alkyl group, a phenyl group and a substituted phenyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a halo C₁-C₄ alkoxy group; a phenylsulfonyl group; a substituted phenylsulfonyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a thiocarbamoyl group; a substituted thiocarbamoyl group substituted by same or different C₁-C₄ alkyl or phenyl group; a C₁-C₄ alkoxy carbonimidoyl group; a substituted C₁-C₄ alkoxy carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; a C₁-C₄ alkylthio carbonimidoyl group; a substituted C₁-C₄ alkylthio carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; a C₁-C₄ alkylsulfinyl carbonimidoyl group; a substituted C₁-C₄ alkylsulfinyl carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; a C₁-C₄ alkylsulfonyl carbonimidoyl group; a substituted C₁-C₄ alkylsulfonyl carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; an amidino group; a substituted amidino group having same or different substituents selected from the group consisting of C₁-C₆ alkyl and phenyl groups; a 5- or 6-membered heterocycle selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole,

25711-791

86

imidazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine, pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithialane, dithiane, piperazine, dioxolane

5 and imidazolidine; or a 5- or 6-membered heterocycle selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine,

10 pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithialane, dithiane, piperazine, dioxolane and imidazolidine and having 1 to 4 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a mercapto group, a

15 C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a C₁-C₄ alkylthio group, a formyl group, a carboxyl group, a carboxy C₁-C₄ alkyl group, a C₁-C₄ alkoxycarbonyl group, a C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, an amino group, a substituted amino group having 1 or 2 same or different

20 substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl C₁-C₄ alkyl group, and a substituted


25 carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; and R⁶ and R⁷ may be taken conjointly to form a 3- to 7-membered ring including a C₂-C₆ alkylene group which may be intercepted by -O-, -S(O)_n- in

30 which n is as defined above, -CO-, or -NR⁸- in which R⁸ is as defined above, and the 3- to 7-membered ring may have same or different substituents selected from the group consisting of halogen atom and C₁-C₄ alkyl group); and

25711-791


87

when A is $-(CR^2R^3)_m-$ and m is an integer of 1 to 4,
 or A is



5 $\begin{array}{c} \text{---C}=\text{C---R}^3 \\ | \quad | \\ \text{R}^2 \end{array}$, then B represents (1) a hydrogen atom;

(2) a halogen atom; (3) a cyano group; (4) a 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom; (5) a substituted 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom and having 1 to 3 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a mercapto group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a C₁-C₄ alkylthio group, a formyl group, a carboxyl group, a carboxy C₁-C₄ alkyl group, a C₁-C₄ alkoxycarbonyl group, a C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, an amino group, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl C₁-C₄ alkyl group, and a substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; (6) a group of the formula:



(in which X is as defined above; and R⁵ represents a hydrogen atom; a C₁-C₈ alkyl group; a halo C₁-C₄ alkyl group; a C₂-C₄ alkenyl group; a halo C₂-C₄ alkenyl group; a C₂-C₄ alkynyl

25711-791

87a

group; a halo C₂-C₄ alkynyl group; a phenyl group; a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a phenyl C₁-C₄ alkyl group; a substituted phenyl C₁-C₄ alkyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen

25711-791

88

atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a 5- to 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom; a substituted 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom and having 1 to 4 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a mercapto group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a C₁-C₄ alkylthio group, a formyl group, a carboxyl group, a carboxy C₁-C₄ alkyl group, a C₁-C₄ alkoxycarbonyl group, a C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, an amino group, a substituted amino group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl C₁-C₄ alkyl group, and a substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; O-R⁶ in which R⁶ is as defined above; S(O)_n-R⁶ in which n and R⁶ are as defined above; or N(R⁶)R⁷ in which R⁶ and R⁷ are as defined above); (7) a group of the formula:




(wherein R⁵ is as defined above; and R⁹ and R¹⁰, same or different, represent a formyl group; a C₁-C₄ alkylcarbonyl group; a C₁-C₄ alkoxycarbonyl group; a phenylcarbonyl group; or a substituted phenylcarbonyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group,

25711-791

89

a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; further, R⁹ and R¹⁰ may be taken conjointly to form a 3- to 7-membered ring including a C₂ to C₆ alkylene group which may be intercepted by -O-,
 5 -S(O)_n- in which n is as defined above, -CO-, or -N(R⁸)- in which R⁸ is as defined above, and the 3- to 7-membered ring may have one or more same or different substituents selected from the group consisting of halogen atom and C₁-C₄ alkyl group); (8) O-R⁶ in which R⁶ is as defined above; (9) S(O)_n-R⁶
 10 in which R⁶ and n are as defined above; (10) N(R⁶)R⁷ in which R⁶ and R⁷ are as defined above; (11) N=C(R⁶)R⁷ in which R⁶ and R⁷ are as defined above; or (12) ON=C(R⁶)R⁷ in which R⁶ and R⁷ are as defined above,


provided that (1) when R¹ is a hydrogen atom or C₁-
 15 C₄ alkyl group, A is



20 m is 1, both of R² and R³ are hydrogen atoms or either one of R² and R³ is a hydrogen atom and other is a C₁-C₄ alkyl group, then B is not a chlorine or bromine atom;

(2) when R¹ is a C₁-C₄ alkyl group or halo C₁-C₄ alkyl group, A is

25




25711-791

90

m is 1, either one of R^2 and R^3 is a hydrogen atom and other is a C_1 - C_4 alkyl group, then B is not a hydrogen atom; and

(3) when R^1 is a hydrogen atom, furan-2-yl, thiophen-2-yl, phenyl, or phenyl having 1 to 3 same or different substituents selected from a hydroxyl group and C_1 - C_4 alkoxy group, A is




10

and m is 0, then B is not a furan-2-yl, thiophen-2-yl, phenyl, or phenyl having 1 to 3 same or different substituents selected from a hydroxyl group and C_1 - C_4 alkoxy group.

15 2. The 1,2,3-thiadiazole derivative or salt thereof according to claim 1, wherein:

R^1 represents a hydrogen atom; a C_1 - C_8 alkyl group or a halo C_1 - C_4 alkyl group;

A represents a group of the following formula:



20

(wherein R^2 and R^3 , which are the same or different, represent: (1) a hydrogen atom; (2) a halogen atom; (3) a cyano group; (4) a formyl group; (5) a C_1 - C_4 alkyl group; (6)

25

25711-791

91

a C₁-C₄ alkoxy group; (7) a C₁-C₄ alkylthio group; (8) a C₁-C₄ alkylsulfinyl group; (9) a C₁-C₄ alkylsulfonyl group; (10) a C₁-C₄ alkylcarbonyl group; (11) a C₁-C₄ alkoxycarbonyl group; (12) a phenyl group; (13) a substituted phenyl group having
 5 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; (14) a phenylcarbonyl
 10 group; or (15) a substituted phenylcarbonyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄ alkyl group and a C₁-C₄ alkoxycarbonyl group; and


15 m represents an integer of 0 to 4); and

when m is 0, B represents (1) a cyano group; (2) a 5- or 6-membered heterocycle selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole, 1,2,3-thiadiazole, 1,2,4-
 20 thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine, pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithialane, dithiane, piperazine, dioxolane and imidazolidine; (3) a substituted 5- or 6-membered heterocycle selected from the group
 25 consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine, pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithialane, dithiane,
 30 piperazine, dioxolane and imidazolidine and having 1 to 4 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a mercapto group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy

25711-791

92

group, a C₁-C₄ alkylthio group, a formyl group, a carboxyl group, a carboxy C₁-C₄ alkyl group, a C₁-C₄ alkoxycarbonyl group, a C₁-C₄ alkoxycarbonyl C₁-C₄ alkyl group, an amino group, a substituted amino group having 1 or 2 same or
 5 different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl C₁-C₄ alkyl group, and a
 10 substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; or (4) a group of the formula:



(in which R⁴ represents a hydrogen atom; a C₁-C₈ alkyl group;
 15 a halo C₁-C₄ alkyl group; a phenyl group; a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; or a 5- or 6-
 20 membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom; and

X represents O; S; NO-R⁶; or NN(R⁶)(R⁷) (in which R⁶ and R⁷, same or different represent a hydrogen atom; a
 25 halogen atom; a formyl group; a C₁-C₄ alkyl group; a C₂-C₄ alkenyl group; a C₂-C₄ alkynyl group; a C₁-C₄ alkylsulfonyl group; a halo C₁-C₄ alkylsulfonyl group; a C₁-C₂₀ alkylcarbonyl group; a C₁-C₄ alkoxycarbonyl group; a phenyl group; a substituted phenyl group having 1 to 5 same or
 30 different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄

25711-791

93

alkoxy group, a halo C₁-C₄ alkoxy group, an amino group and a C₁-C₄ alkoxycarbonyl group; a phenyl C₁-C₄ alkyl group; a substituted phenyl C₁-C₄ alkyl group having, on the phenyl ring thereof, 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group, an amino group and a C₁-C₄ alkoxycarbonyl group; a phenylcarbonyl group; a substituted phenylcarbonyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a halo C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a carbamoyl group; a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl group, a phenyl group and a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a C₁-C₄ alkyl group, a halo C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and halo C₁-C₄ alkoxy group; a phenylsulfonyl group; a substituted phenylsulfonyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a nitro group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a thiocarbamoyl group; a substituted thiocarbamoyl group substituted by same or different C₁-C₄ alkyl or phenyl group; a C₁-C₄ alkoxy carbonimidoyl group; a substituted C₁-C₄ alkoxy carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; a C₁-C₄ alkylthio carbonimidoyl group; a substituted C₁-C₄ alkylthio carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group

25711-791

94


consisting of C₁-C₄ alkyl and phenyl groups; a C₁-C₄ alkylsulfinyl carbonimidoyl group; a substituted C₁-C₄ alkylsulfinyl carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; a C₁-C₄ alkylsulfonyl carbonimidoyl group; a substituted C₁-C₄ alkylsulfonyl carbonimidoyl group having, on the nitrogen atom thereof, same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; an amidino group; a substituted amidino group having same or different substituents selected from the group consisting of C₁-C₆ alkyl and phenyl groups; or a 5- or 6-membered heterocycle selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, isothiazole, pyrazole, imidazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,4-triazole, pyridazine, pyrazine, pyrrolidine, piperidine, morpholine, thiomorpholine, dithialane, dithiane, piperazine, dioxolane and imidazolidine)); and

when m is an integer of 1 to 4, B represents (1) a hydrogen atom; (2) a halogen atom; (3) a cyano group; (4) a 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom; (5) a substituted 5- or 6-membered heterocycle containing 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom and having 1 to 3 same or different substituents selected from the group consisting of a halogen atom, a hydroxyl group, a cyano group, a mercapto group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group, a C₁-C₄ alkylthio group, a carbonyl group, a carboxyl group, a carboxy C₁-C₄ alkyl group, a C₁-C₄ alkoxycarbonyl group, a C₁-C₄ alkoxycarbonyl

25711-791

95

C₁-C₄ alkyl group, an amino group, a carbamoyl group, a substituted carbamoyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups, a carbamoyl C₁-C₄ alkyl group, and a substituted carbamoyl C₁-C₄ alkyl group having 1 or 2 same or different substituents selected from the group consisting of C₁-C₄ alkyl and phenyl groups; (6) a group of the formula:



(in which X is as defined above; and R⁵ represents a hydrogen atom; a C₁-C₈ alkyl group; a halo C₁-C₄ alkyl group; a phenyl group; a substituted phenyl group having 1 to 5 same or different substituents selected from the group consisting of a halogen atom, a cyano group, a hydroxyl group, a C₁-C₄ alkyl group, a C₁-C₄ alkoxy group and a C₁-C₄ alkoxycarbonyl group; a 5- or 6-membered heterocycle having 1 to 3 same or different heteroatoms selected from the group consisting of an oxygen atom, a nitrogen atom and a sulfur atom; O-R⁶ in which R⁶ is as defined above; S(O)_n-R⁶ in which n and R⁶ are as defined above; or N(R⁶)R⁷ in which R⁶ and R⁷ are as defined above); (7) O-R⁶ in which R⁶ is as defined above; (8) S(O)_n-R⁶ in which n and R⁶ are as defined above; (9) N(R⁶)R⁷ in which R⁶ and R⁷ are as defined above; (10) N=C(R⁶)R⁷ in which R⁶ and R⁷ are as defined above; or (11) ON=C(R⁶)R⁷ in which R⁶ and R⁷ are as defined above.

3. An agrohorticultural disease controller comprising:

(a) an effective amount of the 1,2,3-thiadiazole derivative or a salt thereof according to claim 1, and

(b) an agrohorticulturally appropriate inert carrier.

25711-791

96

4. An agrohorticultural disease controller comprising:

(a) an effective amount of the 1,2,3-thiadiazole derivative or a salt thereof according to claim 2, and

5 (b) an agrohorticulturally appropriate inert carrier.

5. A method for controlling a plant disease which comprises treating a plant on which occurrence of plant disease is undesirable with an effective quantity of the
10 agrohorticultural disease controller according to claim 3.

6. A method for controlling a plant disease which comprises treating a plant on which occurrence of plant disease is undesirable with an effective quantity of the agrohorticultural disease controller according to claim 4.

15 7. The 1,2,3-thiadiazole derivative or salt according to claim 1, wherein R^1 is C_1 - C_8 alkyl group.

8. The 1,2,3-thiadiazole derivative or salt according to claim 7, wherein R^1 is methyl, ethyl, n-propyl or cyclopropyl.

20 9. The 1,2,3-thiadiazole derivative or salt according to claim 1, 7 or 8, wherein A is a single bond.

10. The 1,2,3-thiadiazole derivative or salt according to claim 9, wherein B is cyano, formyl, hydroxyiminomethyl, benzyloxyiminomethyl, hydrazonomethyl, phenylhydrazonomethyl,
25 imidazolin-2-yl or 4-methylthiazol-2-yl.

11. The 1,2,3-thiadiazole derivative or salt according to claim 1, 7 or 8, wherein A is methylene.

25711-791

97

12. The 1,2,3-thiadiazole derivative or salt according to claim 11, wherein B is chlorine, hydroxyl, methoxy, benzyloxy, acetoxy, benzoyloxy, ethoxycarbonyloxy, amino, diethylamino, dipropenylamino, benzoylamino,
5 p-toluenesulfonylamino, 3-phenylureido, or 4-methyl-1,2,3-thiadiazol-5-ylmethoxy.

13. An agrohorticultural disease controller comprising:


(a) an effective amount of the 1,2,3-thiadiazole
10 derivative or a salt thereof according to any one of claims 7 to 12, and

(b) an agrohorticulturally appropriate inert carrier.

14. A method for controlling a plant disease which
15 comprises treating a plant on which occurrence of plant disease is undesirable with an effective quantity of the agrohorticultural disease controller according to any one of claims 7 to 12.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS



(I)