



(12)

Oversættelse af europæisk patent

Patent- og
Varemærkestyrelsen

(51) Int.Cl.: **C 07 J 43/00 (2006.01)** **A 61 K 31/58 (2006.01)** **A 61 P 35/00 (2006.01)**

(45) Oversættelsen bekendtgjort den: **2015-01-26**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2014-10-22**

(86) Europæisk ansøgning nr.: **10150763.0**

(86) Europæisk indleveringsdag: **2006-03-02**

(87) Den europæiske ansøgnings publiceringsdag: **2010-07-14**

(30) Prioritet: **2005-03-02 US 657390 P**

(62) Stamansøgningsnr: **06736460.4**

(84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR**

(73) Patenthaver: **University of Maryland, Baltimore, 620 W. Lexington Street, 4th floor, Baltimore, MD 21201, USA**

(72) Opfinder: **BRODIE, Angela, 8873 Reservoir Road, Fulton, Maryland 20759-9622, USA**
NJAR, Vincent, 7908 Springridge Road, Glen Burnie, Maryland 21060-8344, USA

(74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Højbro Plads 10, 1200 København K, Danmark**

(54) Benævnelse: **FARMACEUTISK SAMMENSÆTNING OMFATTENDE 3-BETA-HYDROXY-17-(1-H-BENZIMIDAZOL-1-YL)ANDROSTA-5, 16-DIEN**

(56) Fremdragne publikationer:
US-A- 5 994 335
CLEMENT, OMOSHILE O. ET AL: "Three Dimensional Pharmacophore Modeling of Human CYP17 Inhibitors. Potential Agents for Prostate Cancer Therapy" JOURNAL OF MEDICINAL CHEMISTRY, vol. 46, no. 12, 10 May 2003 (2003-05-10), pages 2345-2351, XP002535273
HANDRATTA, VENKATESH D. ET AL: "Novel C-17-Heteroaryl Steroidal CYP17 Inhibitors/Antiandrogens: Synthesis, in Vitro Biological Activity, Pharmacokinetics, and Antitumor Activity in the LAPC4 Human Prostate Cancer Xenograft Model" JOURNAL OF MEDICINAL CHEMISTRY, vol. 48, no. 8, 25 March 2005 (2005-03-25), pages 2972-2984, XP002535274
POTTER G A ET AL: "A CONVENIENT, LARGE-SCALE SYNTHESIS OF ABIRATERONE ACETATE 3.BETA.-ACETOXY-17-(3-PYRIDYL)ANDROSTA-5,1 6-DIENE, A POTENTIAL NEW DRUG FOR THE TREATMENT OF PROSTATE CANCER" ORGANIC PREPARATIONS AND PROCEDURES INTERNATIONAL, ORGANIC PREPARATION AND PROCEDURES CO., NEWTON HIGHLANDS, MA, US, vol. 290, no. 1, 1 January 1997 (1997-01-01), pages 123-128, XP008056111 ISSN: 0030-4948
VINCENT C. O. NJAR, KATSUYA KATO, IVO P. NNANE, DMITRY N. GRIGORYEV, BRIAN J. LONG, AND ANGELA M. H. BRODIE: "Novel 17-Azolyl Steroids, Potent Inhibitors of Human Cytochrome 17alpha-Hydroxylase-C17,20-lyase (P45017alpha): Potential Agents for the Treatment of Prostate Cancer" JOURNAL OF MEDICINAL CHEMISTRY, vol. 41, no. 6, 24 February 1998 (1998-02-24), pages 902-912, XP002535275
NJAR V C O ET AL: "Nucleophilic vinylic addition-elimination substitution reaction of 3beta-acetoxy-17-chloro-16-formylandrosta- 5,16-diene: a novel and general route to 17-substituted steroids. part 1 - synthesis of novel

Fortsættes ...

17-azolyl-DELTA<16> steroids; inhibitors of 17alpha-hydroxylase/17,20 lyase (17alpha-lyase)", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 6, no. 22, 19 November 1996 (1996-11-19), pages 2777-2782, XP004135894, ISSN: 0960-894X, DOI: 10.1016/S0960-894X(96)00512-4
LING Y-Z ET AL: "17-Imidazolyl, pyrazolyl, and isoxazolyl androstene derivatives. Novel steroidal inhibitors of human cytochrome C17,20-lyase (P450(17 alpha)", JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 40, no. 20, 26 September 1997 (1997-09-26), pages 3297-3304, XP002270332, ISSN: 0022-2623, DOI: 10.1021/JM970337K
HARTMANN R W ET AL: "Synthesis and evaluation of novel steroidal oxime inhibitors of P450 17 (17alpha-hydroxylase/C17-20-Lyase) and 5alpha-reductase types 1 and 2", JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 43, no. 22, 2 November 2000 (2000-11-02), pages 4266-4277, XP002329767, ISSN: 0022-2623, DOI: 10.1021/JM001008M
MAURIZIO RECANATINI ET AL: 'A New Class of Nonsteroidal Aromatase Inhibitors: Design and Synthesis of Chromone and Xanthone Derivatives and Inhibition of the P450 Enzymes Aromatase and 17[alpha]-Hydroxylase/C17,20-Lyase' JOURNAL OF MEDICINAL CHEMISTRY vol. 44, no. 5, 01 March 2001, pages 672 - 680, XP055022294 DOI: 10.1021/jm000955s ISSN: 0022-2623

Description

[0001] This application claims the benefit of the filing date of U.S. Provisional Application Serial No. 60/657390 filed March 2, 2005.

5 [0002] The U.S. Government has a paid-up license in this invention and the right in limited circumstances to require the patent owner to license others on reasonable terms as provided for by the terms of Grant No. CA27440 awarded by the National Institutes of Health (NIH).

10 [0003] The present document describes new chemical entities, particularly steroidal C-17 benzoazoles, pyrimidinoazoles (azabenzooazoles) and diazines. Described are also methods for the synthesis of the benzoazoles, pyrimidinoazoles and diazines. In one instance, the methods for synthesizing benzoazoles or pyrimidinoazoles comprise a step of nucleophilic vinylic "addition-elimination" substitution reaction of 3 β -acetoxy-17-chloro-16- formylandrosta-5,16-diene or analogs thereof and benzoazole or pyrimidinoazole nucleophiles. In another instance, the methods for synthesizing diazines comprise a 15 palladium catalyzed cross-coupling reaction of 17-iodoandrosta-5,16-dien-3 β -ol or analogs thereof with tributylstannyl diazines.

20 [0004] The compounds described herein are potent inhibitors of human CYP17 enzyme as well as potent antagonists of both wild type and mutant androgen receptors (AR). The most potent CYP17 inhibitors were: 3 β -hydroxy-17-(1H-benzimidazole-1-yl)androsta-5,16-diene (**5**, code named **VN/124-1**), 3 β -hydroxy-17-(5¹-pyrimidyl)androsta-5,16-diene (**15**) and 17-(1H-benzimidazole-1-yl)androsta-4,16-diene-3-one (**6**), with IC₅₀ values of 300, 500 and 915 nM, respectively. Compounds **5**, **6**, **14** and **15** were effective at preventing binding of ³H- R1881 (methyltrienolone, a stable synthetic androgen) to both the mutant and LNCaP AR and the wild-type AR, but with a 2.2 to 5-fold higher 25 binding efficiency to the latter. Compounds **5** and **6** were also shown to be potent pure AR antagonists. The cell growth studies showed that **5** and **6** inhibit the growth of DHT-stimulated LNCaP and LAPC4 prostate cancer cells with IC₅₀ values in the low micromolar range (i.e., < 10 μ M). Their inhibitory 30 potencies were comparable to that of casodex but remarkably superior to that of flutamide. The pharmacokinetics of compounds **5** and **6** in mice was investigated. Following s.c. administration of 50 mg/kg of **5** and **6**, peak plasma levels of 16.82 and 5.15 ng/mL, respectively occurred after 30 to 60 min, both compounds were cleared rapidly from plasma (terminal half-lives of 44.17 and 39.93 min, respectively) and neither was detectable at 8 h. Remarkably, compound **5** was rapidly converted into a metabolite tentatively identified as 17-(1H-benzimidazol-1-yl)androsta-3-one. When tested *in vivo*, **5** proved to be very effective at inhibiting the growth of androgen-dependent LAPC4 human 35 prostate tumor xenograft, while **6** was ineffective. Compound **5** (50 mg/kg/twice daily) resulted in a 93.8% reduction (P = 0.00065) in the mean final tumor volume compared with controls, and it was also

significantly more effective than castration. To our knowledge, this is the first example of an anti-hormonal agent (an inhibitor of androgen synthesis (CYP 17 inhibitor)/antiandrogen) that is significantly more effective than castration in suppression of androgen-dependent prostate tumor growth. In view of these impressive anti-cancer properties, compound 5 and others can be used for the treatment of human prostate cancer.

[0005] Prostate cancer (PCA) is the most common malignancy and age-related cause of cancer death worldwide. Apart from lung cancer, PCA is the most common form of cancer in men and the second leading cause of death in American men. In the United States this year (2004), an estimated 230,000 new case of prostate cancer will be diagnosed and about 23,000 men will die of this disease (Jemal et al., Cancer Statistics, 2004. CA Cancer J. Clin., 2004, 54, 8-29). During the period of 1992 to 1999, the average annual incidence of PCA among African American men was 59% higher than among Caucasian men, and the average annual death rate was more than twice that of Caucasian men (American Cancer Society - Cancer Facts and Figures 2003). Androgens play an important role in the development, growth, and progression of PCA (McConnell, J. D., "Physiological basis of endocrine therapy for prostatic cancer", Urol. Clin. North Am., 1991, 18: 1-13). The two most important androgens in this regard are testosterone (T) and dihydrotestosterone (DHT). The testes synthesize about 90% of T and the rest (10 %) is synthesized by the adrenal glands. T is further converted to the more potent androgen DHT by the enzyme steroid 5 α -reductase that is localized primarily in the prostate (Bruchovsky et al., "The conversion of testosterone to 5 α -androstan-17 β -ol-3-one by rat prostate in vivo and in vitro", J. Biol. Chem., 1968, 243, 2012-2021). Huggins *et al.* introduced androgen deprivation as therapy for advanced and metastatic PCA in 1941 (Huggins *et al.* "Studies on prostatic cancer: 2. The effects of castration on advanced carcinoma of the prostate gland.", Arch. Surg., 1941, 43, 209-212). Thereafter, androgen ablation therapy has been shown to produce the most beneficial responses in multiple settings in PCA patients (Denmeade *et al.* "A history of prostate cancer treatment." Nature Rev. Cancer, 2002, 2: 389-396). Orchidectomy (either surgical or medical with a GnRH agonist) remains the standard treatment option for most prostate cancer patients. Medical and surgical orchidectomy reduces or eliminates androgen production by the testes but does not affect androgen synthesis in the adrenal glands. Several studies have reported that a combination therapy of orchidectomy with antiandrogens, to inhibit the action of adrenal androgens, significantly prolongs the survival of PCA patients (Crawford, et al., "A controlled trial of leuprolide with and without flutamide in prostatic carcinoma", N. Engl. J. Med., 1989, 321, 419-424; Crawford, et al., "Treatment of newly diagnosed stage D2 prostate cancer with leuprolide and flutamide or leuprolide alone, Phase III: intergroup study 0036", J. Urol., 1992, 147: 417A; and Denis, L., "Role of maximal androgen blockade in advanced prostate cancer", Prostate, 1994, 5 (Suppl.), 17s-22s). In a recent featured article by Mohler and colleagues (Mohler *et al.*, "The androgen axis in recurrent prostate cancer", Clin. Cancer Res., 2004, 10, 440-448) it was clearly demonstrated that T and DHT occur in recurrent PCA tissues at levels

sufficient to activate androgen receptor. In addition, using microarray-based profiling of isogenic PCA xenograft models, Sawyer and colleagues (Chen et al., "Molecular determinants of resistance to antiandrogen therapy." *Nat. Med.*, 2004, 10, 33-39) found that a modest increase in androgen receptor mRNA was the only change consistently associated with the development of resistance to antiandrogen therapy. Potent and specific compounds that inhibit androgen synthesis in the testes, adrenals, and other tissue may be more effective for the treatment of PCA (Njar, V.C.O.; Brodie, A. M. H., "Inhibitors of 17 α -hydroxylase-C17,20-lyase (CYP17): Potential agents for the treatment of prostate cancer", *Current Pharm. Design*, 1999, 5: 163-180).


[0006] In the testes and adrenal glands, the last step in the biosynthesis of T involves two key reactions, which act sequentially and they are both catalyzed by a single enzyme, the cytochrome P450 monooxygenase 17 α -hydroxylase/17,20-lyase (CYP17) (Hall, P. F., "Cytochrome P-450 C21scc: one enzyme with two actions: Hydroxylase and lyase", *J. Steroid Biochem. Molec. Biol.*, 1991, 40, 527-532). Ketoconazole, as an antifungal agent and by virtue of inhibiting P450 enzymes, is also a modest CYP17 inhibitor and has been used clinically for the treatment of PCA (Trachtenberg et al., "Ketoconazole: A novel and rapid treatment for advanced prostatic cancer", *J. Urol.* 1983, 130, 152-153). It is reported that careful scheduling of treatment can produce prolonged responses in otherwise hormone-refractory prostate cancer patients (Muscati et al., "Optimal dosing of ketocanazole and hydrocortisone leads to long responses in hormone refractory prostate cancer", *Proc. Am. Assoc. Cancer Res.*, 1994, 13: 22 (Abstract)). Furthermore, ketoconazole was found to retain activity in advanced PCA patients with progression despite flutamide withdrawal (Small et al., "Ketoconazole retains activity in advanced prostate cancer patients with progression despite flutamide withdrawal", *J. Urol.*, 1997, 157, 1204-1207). Although, ketoconazole has now been withdrawn from use because of liver toxicity and other side effects this suggests that more potent and selective inhibitors of CYP 17 could provide useful agents for treating this disease, even in advanced stages and in some patients who may appear to be hormone refractory.

[0007] A variety of potent steroid and non-steroidal inhibitors of CYP17 have been reported and some have been shown to be potent inhibitors of testosterone production in rodent models (Njar and Brodie, above). Recently, Jarman and colleagues have described the hormonal impact of their most potent CYP17 inhibitor, abiraterone in patients with prostate cancer (O'Donnell et al., "Hormonal impact of the 17 α -hydroxylase/C17,20-lyase inhibitors abiraterone acetate (CB7630) in patients with prostate cancer", *Br. J. Cancer*, 2004, 90: 2317- 2325). Some of our potent CYP17 inhibitors have been shown to also inhibit 5 α -reductase and/or are potent antiandrogens with potent antitumor activity (Njar and Brodie, above, and Long et al., "Antiandrogenic effects of novel androgen synthesis inhibitors on hormone- dependent prostate cancer." *Cancer Res.*, 2000, 60, 6630-6640). Further illustrative of the background of the invention are U.S. Patent Nos. 5,994,335; 6,200,965; and, 6,444,683. Attempts to

model the three dimensional pharmacophore of human CYP17 inhibitors have been reported (Clement et al, J. Med. Chem 2003, 46, 2345-2351). This document discloses a range of compounds as inhibitors of the CYP17 enzyme in vitro.

5 [0008] We have discovered a series of potent CYP17 inhibitors/antiandrogens, the 17- benzoazoles, 17-pyrimidinoazoles and 17-diazines (see, e.g., Schemes 1 and 2, for examples of preparation of compounds which can be analogously applied to other structures, as described below). The stimulus for preparing these C- 17 heteroaryl steroids was based on our desire to incorporate benzimidazole, benzotriazole, pyrimidinoazole and diazine moieties, so-called "privileged substructures" (Nicolaou et 10 al., "Natural product-like combinatorial libraries based on privileged structures. 1. General principles and solid-phase synthesis of benzopyrans", J. Am. Chem. Soc, 2000, 122, 9939-9953. "Privileged structures", a term originally introduced by Evans et al. (J Med. Chem., 1988, 31, 2235-2246) to describe structural motifs capable of interacting with a variety of unrelated molecular targets) in the new molecules. These scaffolds, especially the benzimidazole scaffold, continue to receive extensive 15 attention in medicinal chemistry because of their diverse portfolio of biological activities and also as entities of a variety of useful drugs (Nicolaou et al., above).

[0009] Described are C-17 heteroaryl steroid compounds of the following general formula I:



20 wherein:

- the ABC ring structure is the A, B and C ring portions of a steroid or analog thereof, which are optionally substituted;
- the — bond at the 16,17 position is a double bond or, when the compound is 17-(1H-benzimidazol-1-yl)androst-3-one, a single bond; and
- X is an optionally substituted benzimidazole, benzotriazole, pyrimidinoimidazole (purine), pyrimidinotriazole or diazine; the benzimidazole, benzotriazole, and pyrimidinoimidazole groups being bonded to the steroid residue through a nitrogen atom on the 5-membered ring; and, the diazine groups being bonded to the steroid residue through a carbon atom on the diazine ring.


[0010] Pharmaceutically acceptable salts of these compounds are also described.

[0011] The optional substitution for the ABC ring structure includes one or more of: alkyl and


5 halogenated alkyl (preferably C₁₋₆); alkenyl and halogenated alkenyl (preferably C₁₋₆) including where the double bond is directly attached to the ring structure; halogen; amino; aminoalkylene; hydroxyimino; and hydroxy. Further optionally, hydrogen substituents on adjacent carbon atoms of the ABC ring structure may be removed and replaced by an additional bond between the adjacent carbon atoms to result in a double bond between these carbons in the ring structure. Preferred optional
10 substitutions on the ABC ring structure are methyl groups at the 10 and/or 13 positions of the ring structure.

[0012] The optional substitution for the benzimidazole, benzotriazole, pyrimidinoimidazole, pyrimidinotriazole or diazine structures include halogen, amino, aminoalkylene, hydroxy, -SH, -S-alkyl, 15 alkyl and halogenated alkyl (preferably C₁₋₆). These optional substituents will be on ring carbon atoms of the benzimidazole, benzotriazole, pyrimidinoimidazole, pyrimidinotriazole or diazine structures.


[0013] The benzimidazole, benzotriazole, pyrimidinoimidazole, pyrimidinotriazole or diazine structures are of the following formulae, respectively:




20 benzimidazole




benzotriazole




pyrimidinoimidazole



pyrimidinotriazole



pyrimidine




























pyrazine

wherein the * indicates the point of attachment to the steroid residue.

25 [0014] In one instance, the ABC ring structure has a C ring which has no substitution except for preferably alkyl, particularly methyl, substitution at the carbon shared with the D ring which is adjacent the attachment to the C-17 heteroaryl substitution, i.e., the 13 -position.

[0015] In another instance, the A, B and C rings of the ABC ring structure have a conventional structure based on 3β -hydroxy-androsta-5,16-diene or 3-oxo-androsta-5,16-diene. But in another instance the A and B rings have one of the following structures 1- 25:



[0016] The following lists the chemical names of compounds having the AB rings as in 1-25, and the C and D rings conventional, wherein X is benzimidazole. Analogous compounds wherein X is benzotriazole, pyrimidinoimidazole, pyrimidinotriazole, pyrazine or pyrimidine are also described.

15 **Compound 1:** 3β -Hydroxy- 3α -methyl- 17-(1*H*-benzimidazol-1-yl)-androsta-5,16- diene

Compound 2: 3β -Fluoro-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

Compound 3: 3β -Chloro-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

Compound 4: 3β -Bromo-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

Compound 5: 3β -Iodo-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

Compound 6: 3β -Amino-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

Compound 7: 17-(1*H*-benzimidazol-1-yl)-androsta-3,5,16-triene

Compound 8: 17-(1*H*-benzimidazol-1-yl)-androsta-2,4, 16-triene

Compound 9: 17-(1*H*-benzimidazol-1-yl)-3-methyleneandrosta-5,16-triene

Compound 10: 17-(1*H*-benzimidazol-1-yl)-3-methyleneandrosta-4,16-triene

Compound 11: 3,3-Difluoro-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

Compound 12: 3 ,3-Difluoro- 17-(1*H*-benzimidazol-1-yl)-androsta-4,16-diene

Compound 13: 17-(1*H*-benzimidazol-1-yl)-3-methyleneandrosta-2,4,16-triene

Compound 14: 17-(1*H*-benzimidazol-1-yl)-3-methyleneandrosta-2,4,6,16-tetraene

Compound 15: 3,3-Difluoro-17-(1*H*-benzimidazol-1-yl)-androsta-2,4,16-triene

Compound 16: 3,3-Difluoro-17-(1*H*-benzimidazol-1-yl)-androsta-2,4,6,16-tetraene

Compound 17: 3-Hydroxyimino-17-(1*H*-benzimidazol-1-yl)-androsta-5,16-diene

5 **Compound 18:** 3-Hydroxyimino-17-(1*H*-benzimidazol-1-yl)-androsta-4,16-diene

Compound 19: 3-Hydroxyimino-17-(1*H*-benzimidazol-1-yl)-androsta-2,4,16-triene

Compound 20: 3-Hydroxyimino-17-(1*H*-benzimidazol-1-yl)-androsta-2,4,6,16-diene

Compound 21: 3-Hydroxy-17-(1*H*-benzimidazol-1-yl)-estra-1,3,5(10),16-tetraene




Compound 22: 3-Fluoro-17-(1*H*-benzimidazol-1-yl)-estra-1,3,5(10),16-tetraene

10 **Compound 23:** 3-Chloro-17-(1*H*-benzimidazol-1-yl)-estra-1,3,5(10),16-tetraene

Compound 24: 3-Bromo-17-(1*H*-benzimidazol-1-yl)-estra-1,3,5(10),16-tetraene


Compound 25: 3-Iodo 17-(1*H*-benzimidazol-1-yl)-estra-1,3,5(10),16-tetraene

15 [0017] Examples of optional substituents for the heteroaryl ring, X, are shown by the following structures **26 - 40** wherein X is benzimidazole. Analogous compounds wherein X is substituted benzotriazole, pyrimidinoimidazole, pyrimidinotriazole, pyrazine or pyrimidine are also described.




20 [0018] Other examples of optional substituents for the heteroaryl ring, X, are shown by the following structures **41 - 46** wherein X is substituted C-17-azabenzimidazole (i.e., pyrimidinoimidazole or purine). Analogous compounds wherein X is substituted benzimidazole, benzotriazole,

pyrimidinotriazole, pyrazine or pyrimidine are also described.



[0019] Described are compounds of the following structures M5, M6, M9 and M10.



5

[0020] The inhibitory activities of these compounds compared to CYP17 and steroid 5 α -reductases, the binding to and transactivation of androgen receptors, and their antiproliferative effects against two human prostate cancer cell lines, LNCaP and LAPC-4 were studied. The pharmacokinetics of compounds 5 and 6 were evaluated in mice and the in vivo antitumor activities against human LAPC-4 prostate carcinoma were also evaluated in mice. To our knowledge, all the compounds described here, with the exception of compound 15 represent novel entities (Haidar et al., "Novel steroid pyrimidyl inhibitors of P450 17 (17 α -hydroxylase/C17-20-lyase)", Arch. Pharm. Med Chem., 2001, 334, 373-374; and Haidar et al., "Effects of novel 17 α -hydroxylase/C17,20-lyase (P45017, CYP17) inhibitors on androgen biosynthesis in vitro and in vivo", J Steroid Biochem. Molec. Biol, 2003, 84, 555-562).

[0021] The preparation of the new 17-benzoazoles and 17-diazines is outlined in Schemes 1 and 2, respectively. These methods can be applied analogously to other analogs described herein.

[0022] The key intermediate in our synthesis of the 17-benzoazoles, 3 β -acetoxy-17-chloro-16-formylandrosta-5,16-diene (2) was obtained from (1) by our routine procedure as previously described (Njar et al., "Nucleophilic vinylic "addition-elimination" substitution reaction of 3 β -acetoxy-17-chloro-16-formylandrosta-5,16-diene: A novel and general route to 17-substituted- Δ 16-steroids. Part 1. Synthesis of novel 17-azolyl- Δ 16 steroids; inhibitors of 17 α -hydroxylase/17,20-lyase (P45017 α)", Bioorg. Med. Chem. Lett., 1996, 6, 2777 - 2782; and "Novel 17-azolyl steroids; potent inhibitors of cytochrome P450 17 α -hydroxylase/17,20-lyase (P45017 α): Potential agents for the treatment of prostate cancer", J. Med. Chem., 1998, 41, 902 - 912). Treatment of 2 with benzimidazole in the presence of K₂CO₃ in DMF at approximately 80°C gave the desired 3 β -acetoxy-17-1H-bezimidazole 3 in near

quantitative yield. Compound **3** was smoothly deformylated with 10% palladium on activated charcoal in refluxing benzonitrile to give compound **4** in 93% yield, from which hydrolysis gave the required 3 β -hydroxy 17-benzimidazole **5**. Modified Oppenauer oxidation of **5** afforded the corresponding Δ^4 -3-oxo analog, **6**.

5

[0023] The reaction of **2** with benzotriazole in the presence of K_2CO_3 in DMF at approximately 80°C gave the desired 3 β -acetoxy-17-benzo-1*H*-1,2,3-triazole **7b** in excellent yield, together with the 2*H*-1,2,3-triazole regioisomer **7a** in approx 5% yield. These two regioisomers were readily separated by flash column chromatography (FCC) on silica gel and were also easily identified by their respective 10 proton NMR spectra. Thus, the four aromatic protons of the symmetrical 2*H*-1,2,3-triazole **7a** appeared as two pairs of doublets at δ 7.43, 7.45, 7.88 and 7.90 while the four aromatic protons of the unsymmetrical 1*H*-1,2,3-triazole **7b** appeared as multiplet at δ 7.46 (2H) and doublets at δ 7.57 (1H) and 8.15 (1H), respectively. In addition, the 16-CHO proton in **7a** was significantly shifted downfield to δ 10.66 compared to that in **7b** at δ 9.59. Deformylation of **7b** with in situ generation of Rh(1,3-bis(diphenylphosphino)propane)₂⁺Cl⁻ catalyst [Rh(dppp)₂⁺Cl] in refluxing xylenes gave compound **8**, and following hydrolysis of the 3 β -acetoxy group, we obtained the target 3 β -hydroxy-17-(benzo-1*H*-1,2,3-triazol-1-yl)androsta-5,16-diene (**9**) in 90% yield. Oxidation of **9** afforded **10** in good yield.

[0024] Synthesis of the 17-diazines, (17-diazine **14** and 17-pyrimidine **15**) commenced from the readily

20 available dehydroepiandrosterone (**11**, Scheme 2), which was converted to the corresponding 17 hydrazone **12** by treatment with hydrazine hydrate and hydrazine sulfate as previously described by Potter et al., A convenient, large-scale synthesis of abiraterone acetate [3 β -acetoxy-17(3-pyridyl)androsta-5,16-diene], a potential new drug for the treatment of prostate cancer. Org. Prep. Proc. Int., 1997, 29, 123-128. Treatment of **12** with iodine in the presence of 1,1,3,3-tetramethylguanidine 25 gave the vinyl 17-iodide **13** in excellent yield. The palladium catalyzed cross-coupling reactions (Choshi et al., "Total synthesis of Grossularines-1 and -2." J Org. Chem., 1995, 60, 5899-5904) of **13** with (2-tributylstannyl)pyrazine or (5-tributylstannyl)pyrimidine proceeded to give 3 β -hydroxy-17-(2-pyrazyl)-androsta-5,16-diene (14, 15%), and 3 β -hydroxy-17-(5-pyrimidyl)-androsta-5,16-diene (15, 10%), respectively. The low yields of these two cross-coupling reactions may be due to instability 30 of the stannyldiazine reagents under the reaction conditions employed. The structures of the target compounds, **14** and **15** were readily identified by their proton NMR spectra: The three nonequivalent protons of the 17-pyrazine moiety in **14** appeared as three singlet at δ 8.35, 8.48 and 8.70, while for the three protons of the 17-pyrimidine moiety in **15**, two equivalent protons appear as a singlet at δ 8.73 and one proton appeared at δ 9.07. Furthermore, the 17-diazine groups of **14** and **15** exhibit different 35 influences on the chemical shifts of their respective 16-olefinic protons with respect to the 16-proton of the precursor Δ^{16} -17-iodide **13**: the 16-H in **14** appeared as a singlet at δ 6.77, being significantly deshielded compared to the 16-H in **13** (δ 6.14); the 16-H in **15** appeared at δ 6.11, similar to **13**. As

indicated above, compound 15 was previously reported Haidar et al., however, it was synthesized by a procedure that is different from the one described herein.

5 [0025] A representative sample of the novel compounds were then subjected to extensive in vitro and in vivo studies as described in detail in the following sections.

10 [0026] A method of treating prostate cancer or prostate hyperplasia comprising administering to a subject in need thereof an effective amount of a compound. The term "treating" is used as described herein is also disclosed conventionally, e.g., the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving, etc., one or more of the symptoms associated with the prostate disease. Examples of prostate diseases that can be treated include, e.g., prostatic hyperplasia (BPH), and prostate cancer (e.g., prostatic adenocarcinoma).

15 The specific dose level and frequency of dosage may vary, depending upon a variety of factors, including the activity of the specific active compound, its metabolic stability and length of action, rate of excretion, mode and time of administration, the age, body weight, health condition, gender, diet, etc., of the subject, and the severity of the prostate cancer or hyperplasia. Any effective amount of the compound can be administered, e.g., from about 1 mg to about 500 mg per day, more specifically about 50 mg to about 150 mg per day. The compounds can be administered in any form by any effective route, including, e.g., oral, parenteral, enteral, intraperitoneal, topical, transdermal (e.g., using any 20 standard patch), ophthalmic, nasally, local, non-oral, such as aerosol, spray, inhalation, subcutaneous, intravenous, intramuscular, buccal, sublingual, rectal, vaginal, intra-arterial, and intrathecal, etc. A compound of the present invention can be administered alone, or in combination with any other ingredient(s), active or inactive, for example, with physiologically acceptable vehicles to make suitable pharmaceutical compositions.

25

30 [0027] The invention provides a pharmaceutical composition which comprises 3 β -hydroxy-17-(1H-benzimidazole-1-yl)androsta-5, 16-diene or a pharmaceutically acceptable salt thereof for use in the treatment of an androgen-dependent tumor by inhibition of both wild-type and mutant androgen receptors. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

35 [0028] In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius and, all parts and percentages are by weight, unless otherwise indicated.

EXAMPLES

Biological Studies

[0029] CYP17 Inhibition Studies: A CYP17 inhibition assay is performed according to our previously reported procedure, in which intact cytochrome P450c17-expressing *E. coli* is used as the enzyme source (Grigoryev et al., "Cytochrome P450c17-expressing Escherichia coli as a fist-step screening system for 17 α -hydroxylase-C17,20-lyase inhibitors", Anal. Biochem.; 1999, 267, 319-330; and "Effects of new 17 α -hydroxylase/C17,20-lyase inhibitors on LNCaP prostate cancer cell growth in vitro and in vivo", Br. J. Cancer, 1999, 81, 622-630. IC₅₀ values of the compounds are determined from dose-response curves and are listed in Table 1. The IC₅₀ values for ketoconazole, abiraterone (a CYP17 inhibitor in clinical trials (O'Donnell, above), Chart 1) and 3 β -hydroxy-17-(1H-imidazole-1-yl)androsta-5,16-diene (VN/85-1, compound **16**, Chart 1, believed to be the most potent CYP17 inhibitor (Njar et al., Current Pharm. Design, 1999, 5: 163-180; and J. Med. Chem., 1998, 41, 902 - 912, above) are also determined in the same assay system for comparison. Some of the new 17-heterocycles exhibit potent inhibition of CYP17 with IC₅₀ values of 300 - 915 nM. The benzimidazoles, **5** and **6** are 4- to 6-fold more potent than the benzotriazoles **9** and **10**. This result suggests that the electronic nature of the 17-heterocycle influence inhibitory activity. Furthermore, compounds with the Δ^5 -3 β -ol functionality, **5** and **9** are at least 3-fold more potent than the corresponding analogs with Δ^4 -3-one functionality, **6** and **10**, respectively. These results are in contrast to our previous results for the simple 17-azole CYP17 inhibitors. In that series of inhibitors, there is no marked difference in the inhibitory potencies between the Δ^5 -3 β -ol azoles and the corresponding Δ^4 -3-one analogs (Njar et al., J. Med. Chem., 1998, 41, 902 - 912, above). A possible explanation is that the bulkier benzoazoles bind differently at the active site of the enzyme such that the interaction(s) of the moiety at the 3-position is important for binding.

[0030] The binding of the substrate or inhibitory ligands to the heme component of some P450 cytochromes is investigated using UV-vis difference spectroscopy (Jefcoat C. R., "Measurement of substrate and inhibitor binding to microsomal cytochrome P450 by optical difference spectroscopy", Methods Enzymol., 1978, 52, 258-279). This approach is extended following standard procedure previously reported by us (Njar et al., Bioorg. Med. Chem. Lett., 1996, 6, 2777 - 2782; and J. Med. Chem., 1998, 41, 902 - 912). Compounds **5** and **9** each induce a type II difference spectrum, indicating coordination of steroidal nitrogen (N-3 of benzimidazole or benzotriazole ring) to the heme iron of CYP117, with formation of low-spin iron. The peak positions for the Soret maximum for the enzyme complex with **5** and **9** (426 nM) is in agreement with available data for the binding of nitrogen ligands to CYP systems, and is also in agreement with our results with other 17-azolyl CYP 17 inhibitors (Njar et al., Bioorg. Med. Chem. Lett., 1996, 6, 2777 - 2782; and J. Med. Chem., 1998, 41, 902 - 912). The interaction of the benzoazole nitrogen with the heme iron of CYP 17 suggests bulk tolerance at the 17-

position, because the binding affinities of **5** and **9** are identical to that of the less sterically demanding **16**, with a 17-imidazole group.

[0031] Of the two 17-diazines tested, the 17-pyrimidine **15** with an IC_{50} value of 500 nM is about 8-fold more potent than the 17-pyrazine **14** (IC_{50} = 3810 nM). As with the benzoazoles, this result suggests that the electronic nature of the 17-heterocycle influence inhibitory activity. Finally, IC_{50} values in the same assay system for ketoconazole, and abiraterone are evaluated (Table 1). The most potent compound in this series, 17-benzimidazole **5**, exhibits about 4 and about 3 -fold improvements in CYP17 inhibition over these compounds, respectively, although it is less potent than **16**.

10

[0032] **Inhibition of human 5 α -reductase isozymes type 1 and 2 in vitro:** On the basis of previous findings that some CYP17 inhibitors are able to inhibit human 5 α -reductase enzymes, we briefly evaluated this new series of CYP17 inhibitors. The inhibitory activities of compounds **5**, **6**, **9**, **10** and finasteride as a reference are determined using the DU-145 cell line (human type 1 enzyme) and human homogenates of BPH tissue (human type 2 enzyme) as described by Hartmann et al., "Synthesis and evaluation of 2'-substituted 4-(4'-carboxy- or 4'-carboxymethylbenzylidene)-N-acylpiperidines: Highly potent and in vivo active steroid 5 α -reductase type 2 inhibitors", J. Med. Chem., 2002, 45, 3406-3417. The IC_{50} values or the percent inhibition values at a concentration of 10 μ M for some compound are presented in Table 1. Only compound **6** exhibits potent inhibition of both type **1** and **2** enzymes (IC_{50} = 770 and 480 nM, respectively), although it is several fold less potent than finasteride (IC_{50} = 60 and 2 nM, respectively).

15

20

[0033] **LNCaP and PC-3AR androgen receptor binding assays:** Because we had previously demonstrated that some of our CYP 17 inhibitors are potent antiandrogens for both the mutant and wild-type AR (Long et al., Gregoriyev et al. and Njar et al., J. Med. Chem., 1998, 41, 902 - 912, above) it was of interest to assess the ability of this series of CYP 17 inhibitors to bind to these receptors. AR competition is determined using labeled R1881 ($[^3H]$ -R1881) in the androgen-sensitive LNCaP cells, that express mutant AR, and the androgen-independent PC-3 cells stably transfected with the wild-type AR (designated PC-3AR). Compounds **5**, **6**, **14** and **15**, in the nanomolar concentration range, compete effectively with labeled R1881 for binding to both types of ARs in a dose-dependent manner (Figure not shown). Compounds **5**, **6**, **14** and **15**, with IC_{50} values of 384, 242, 336 and 374 nM, respectively (Table 1), *versus* the wild type AR are 29 to 45-fold more potent than with the clinically used antiandrogen, flutamide (IC_{50} = 10,985 nM). As shown in Table 1, the binding affinities for the mutant AR of **5** and **6** are comparable to that of casodex, a currently used antiandrogen, but again superior to that of flutamide. However, the biological activity of flutamide is derived mainly from a metabolite, hydroxyflutamide, which is a much more potent AR antagonist.

25

30

35

[0034] **Effects of agents on LNCaP mutant AR-mediated transcription:** Next, we asked whether compounds **5** and **6** are acting as AR agonists or antagonists. A study on androgen-regulated transcriptional activation is performed in LNCaP cells transiently transected with a probasin luciferase reporter construct AARZ-Luc (luciferase activity assay) (Kim et al., "Synergism of cytoplasmic kinases in IL6-induced ligand-independent activation of androgen receptor in prostate cancer cells", *Oncogene*, 2004, 23:1838-1844; and Zhang et al., "A Small composite probasin promoter confers high levels of prostate-specific gene expression through regulation by androgens and glucocorticoids in Vitro and in Vivo", *Endocrinology*, 2000, 141: 4698-4710). Compounds **5**, **6** or casodex each at 0.1 and 10.0 μ M have no effect on luciferase activity, whereas luciferase expression is increased approximately 99.6-fold after treatment with 1.0 nM DHT for 18 h (Figure 1). Furthermore, luciferase expression induced by exposure to 1.0 nM DHT is decreased in a concentration-dependent manner by **5**, **6**, and casodex and in a similar fashion (Figure 1). Together, these results suggest that compounds **5** and **6** like casodex do not possess AR agonistic or partial agonistic activity and may be considered as strong, pure androgen antagonists. Although we did not test the compounds with PC-3AR/LU cells, which express wild-type AR, it is likely that they may also behave in a similar fashion. We have previously shown that some of our CYP17 inhibitors were more comparable to casodex than to flutamide (Long et al., above), and this appears to be the case with these new compounds. In general, our novel compounds interact strongly with both AR types, an indication that the compounds may be useful for the treatment of patients with tumors expressing either wild-type or mutated AR, or for patients with amplified AR expression.

[0035] **Effects of benzoazoles on the growth of LNCaP and LAPC-4 prostate cancer cells in vitro:** The abilities of compounds **5** and **6** to inhibit proliferation in mutant LNCaP cells stimulated by 1 nM DHT is examined. This concentration of DHT stimulated LNCaP cell proliferation by about 2-fold compared to vehicle-treated cells (Figure 2A). As shown in Figure 2A, compounds **5** and **6** each inhibit the DHT-induced LNCaP cell proliferation in a dose-dependent fashion, with IC₅₀ values of 6.0 and 1.8 μ M, respectively. Casodex is used as a positive control, and it exhibits similar inhibition of DHT-induced LNCaP cell proliferation (Figure 2A, IC₅₀ = 8.6 μ M). Treating the androgen-sensitive LAPC4 prostate cell line with 10 nM DHT, surprisingly, does not significantly induce cell proliferation (Figure 2B). Other investigators have also reported that the response of LAPC4 cells to androgens is not as pronounced as that observed in LNCaP cells (Thompson et al., "Androgen antagonist activity by the antioxidant moiety of vitamin E, 2,2,5,7,8-pentamethyl-6-chromanol in human prostate carcinoma cells", *Molec. Cancer Ther.*, 2003, 2, 797-803). However, compounds **5**, **6** and casodex each exhibit a dose-dependent inhibition of this cell line (Figure 2B) as with the LNCaP cells. The order of inhibitory potency of LAPC4 cell proliferation is **6** > **5** > casodex, with IC₅₀ values of 1.0, 3.2 and 10 μ M, respectively. Together, these results suggest that **5** and **6** may be acting to block the action of DHT in stimulating cell proliferation, in correlation with their androgen receptor binding and activation.

properties described above. Compounds **5** and **6** are amongst the most potent antiandrogens described to date.

[0036] Pharmacokinetics of **5 and **6** and metabolism of **5**:** The pharmacokinetic properties in male

5 SCID mouse for the two lead compounds, **5** and **6** are studied following our recently described procedure for other CYP17 inhibitors (Nnane et al., "Pharmacokinetic profile of 3 β -hydroxy-17-(1H-123-triazol-1-yl)androsta-5,16-diene (VN/87-1), a potent androgen synthesis inhibitor in mice", J. Steroid Biochem. Molec. Biol., 2001, 71, 145-152; and Handratta et al., "Potent CYP17 inhibitors: improved syntheses, pharmacokinetics and antitumor activity in the LNCaP human prostate cancer 10 model Steroid Biechein. Molec. Biol., 2004, 92, 155-165. The results are summarized in Table 2 and Figures 3-5.

[0037] On reverse phase HPLC, **5** [retention time (rt) = 21.6 min] is well resolved from the internal standard (**16**, rt = 11.5 min), a metabolite (rt = 17.3 min) and other endogenous compounds in mouse

15 plasma (**Figure 3**). The calibration curves derived for **5** are linear and reproducible (data not shown), the inter- and intra-assay variability is less than 10% and its limit of detection is 100 ng/ml. The HPLC assay is validated and used to monitor **5** in mice plasma.

[0038] Following subcutaneous administration, the plasma concentration of **5** declines exponentially

20 with a mean half-life of about 44.17 min and elimination rate constant of 56.5 min⁻¹. Compound **5** is cleared at a rate of 1986.14 ml/h/kg from the systemic circulation and was not detected 6 h after administration. The calculated non-compartmental pharmacokinetic parameters based on the plasma concentration profile following subcutaneous administration of **5** are shown in Table 2. The plasma concentration-time curves after s.c. administration of **5** (50 and 100 mg/kg) to male SCID mice are also 25 shown in Figure 4. After s.c. administration of **5**, the observed plasma concentration in mice reach peak levels 30.0 min post dose. Compound **5** is well absorbed from the subcutaneous site and the area under the curve for the plasma concentration versus time profiles after s.c. administration increases proportionately to dose as the administration dose is changed from 50 to 100 mg/kg. Furthermore, the elimination half-life, and mean residence time are relatively constant as the dose of **5** increases from 50 30 to 100 mg/kg (Table 1). These results indicate that the pharmacokinetic profile of **5** is dose independent.

[0039] Figure 5 shows that a significant amount of a polar metabolite [retention time, 17.3 min, see

Figure 3)] is formed from **5** and present in the plasma during the *in vivo* pharmacokinetic studies. The

maximum amount of the metabolite is 67.72 %, attained about 2 h post dose. This metabolite shows

35 identical retention time as compound **6**. This metabolite is tentatively identified by LC-MS; its molecular mass (m/z 391 = M + H⁺) is consistent with the structure of 3-oxo- $\Delta^{5,16}$ -tetrahydro compound **5**(i.e., 17-(1H-benzimidazol-1-yl)androst-3-one). The metabolite may have been formed

from **5** via oxidation of the 3β -OH \rightarrow 3-oxo, followed by reduction (reductases) of both Δ^5 and Δ^{16} double bonds. A similar metabolite was previously identified (formed as a result of oxidation of a 3β -OH \rightarrow 3-oxo, followed by isomeraization of Δ^5 double bond) in male mice of a closely related steroid 17-imidazole (Handratta et al., above).

5

[0040] A major metabolite of **5**, i.e., 17-(1H-benzimidazol-1-yl)androst-3-one, may be synthesized from trans-androsterone; see Scheme 3. It is also expected to have analogous activity.

10 **[0041]** The *in vivo* pharmacokinetics of **6** in mice is unlike that of compound **5** due to the relatively low C_{max} and significantly higher elimination rate (Figure 4 and Table 2). In addition, we did not detect any metabolism(s) of compound **6** in the plasma, in contrast to our observation with compound **5**.

15 **[0042] Effects of 5 and 6 on LAPC4 Xenografts grown in SCID mice:** On the basis of impressive multiple *in vitro* biological activities, i.e., potent inhibition of CYP17, strong antiproliferative prostate cancer cell activity and antiandrogenic activities, **5** and **6** are selected for *in vivo* antitumor efficacy studies in androgen-depended LAPC4 human prostate cancer xenograft model.

20 **[0043]** In the first experiment, the effect of compounds **5** and **6** on the growth of well-established LAPC4 prostate cancer tumors in SCID mice is determined, and castration is used as the reference treatment. Tumor-bearing mice are assigned ($n = 5$ /group) to receive one of two doses of **5** or **6** (0.15 mmol/kg once-daily or 0.15 mmol/kg twice-daily). Tumor volumes are measured weekly and compared with controls receiving vehicle or castrated mice.

25 **[0044]** Castration leads to a 55% reduction of final tumor volume, as compared to the control (Figure 6). Administration of 0.15 mmol/kg once daily and 0.15 mmol/kg twice daily of **5** results in reduction of average final tumor volumes of 41% and 86.5%, respectively, compared to tumors in vehicle-treated control animals (Figure 6). In contrast to the excellent tumor growth inhibition for **5** treated mice, mice treated with compound **6** are either ineffective at the low dose or even show stimulation of tumor growth compared to control (Figure 6). The inability of **6** to inhibit LAPC4 tumor growth *in vivo* is especially disappointing because the compound is very effective at inhibiting PCA cell growth *in vitro*, and is a highly potent pure antiandrogen (see Figure 1). The highly significant disparity in the *in vivo* antitumor efficacy of **5** and **6** cannot easily be attributable to differences in the pharmacokinetic properties of the two compounds. The underlying reason(s) for the dramatic differences in *in vivo* antitumor efficacy of these two closely related compounds is unknown at this time. However, it may be attributable to **6** being converted in the animals to metabolite(s) that may be a strong agonist of androgen receptor thus causing tumor growth stimulation. During the study, all mice were weighed once per week. The body weights of all treated groups increased slightly and were similar to the increase observed with the control group.

All mice appeared healthy and no adverse effects were observed suggesting that the compounds were without significant toxicity.

[0045] The second *in vivo* experiment tests the ability of **5** to inhibit the growth of LAPC4 prostate

5 cancer cells growing in SCID mice, and **16** (Chart 1), a previously identified potent CYP17
inhibitor/antiandrogen (Gregoriyev et al. and Njar et al., *J. Med. Chem.*, 1998, 41, 902 - 912, above)
and castration are used as reference treatments. In this experiment, treatment begins on the day that
mice were inoculated subcutaneously with hormone-dependent LAPC4 cells and are castrated or
injected sc twice daily with **5** or **16**. Figure 7 shows the effects of the various treatments on the
10 emergence and on the size of tumors during the 21 weeks of therapy.

[0046] All other groups develop palpable and measurable tumor at week 10 of therapy except for the
group treated with **16** (0.15 mmol/kg twice-daily) that develop palpable and measurable tumors at week
11. Total tumor volume in the control mice increases by 8-fold over 14 weeks of treatment when mice

15 are sacrificed because of the large tumors. Thus, the tumor volumes for the other groups are compared
to those of the control group at week 14 of treatment. Tumor volume in the castrated mice increases by
only 4.1 fold (about 50% reduction compared to control), and is similar to the 3.7 fold increase (53.8%
reduction compared to control) observed in mice treated with **16**. In the mice treated with **5** (0.15
mmol/kg twice-daily), tumor volume increases by only 0.5 fold, which represents a 93.8%

20 reduction *versus* control mice ($P= 0.00065$). At week 16, the mean tumor volume in the compound **5**-
treated animals is found to be lower (almost negligible and dormant) than their mean tumor volume at
week 10 when measurable tumors emerge. Furthermore, **5** causes a significant inhibitory effect on
tumors, compared to **16** or castration, $P = 0.005$ and 0.05 , respectively. In general, tumors in the
control, castration and compound **16** treated mice grow rapidly, while the tumor of the **5** treated mice
25 grows very slowly and in a biphasic manner (Figure 7). Compound **5** is the most effective agent, and
significantly is much more effective than castration at inhibition of tumor growth. It is interesting to
note that although **16** is 6 times more potent than **5** in CYP 17 inhibition, the latter exhibits a superior *in*
vivo antitumor activity. The reason(s) responsible for this phenomenon is unknown at this time, but may
be in part due to better pharmacokinetic and or pharmacodynamic properties of **5**.

30

[0047] To determined whether the "dormant" compound **5**-treated prostate tumors (see Figure 7, week
16) are able to grow on a lower dose of **5**, its dose was reduced to 0.15 mmol/kg thrice a week (a 78.6%
reduction in dosage) from weeks 16 - 9, and the tumor volumes measured weekly. During this period of
35 treatment with reduced dose of the compound, tumors resume growth (Figure 7). After this 3-week
interval, drug treatment with the usual dose is resumed, and the tumor growth slows and reaches a
plateau. These data suggest a cytostatic nature of this treatment and infer the need for continuous
administration to achieve the antitumor effect.

[0048] At the end of the experiment, the levels of **5** in the tumors and organs of the **5**-treated mice are determined. The **5** levels by HPLC in the tumors, testis and liver 1 h after administration of the final dose (insert of Figure 7) are measured. Interestingly, a small (~ 15% relative to **5**) amount of metabolite is detected only in the liver tissues. This metabolite has the same retention time as the metabolite

5 observed in the plasma (*vide supra*). The highest concentration of $39.0 \pm 8.4 \mu\text{g}/\text{mg}$ tissue of **5** is measured in the s.c. tumors. The concentrations in the liver and testis are lower but detectable. The level of **5** in tumors is significantly higher than the levels measured in the plasma, which may be a result of accumulation of the compound through the period of the experiment. Thus, inhibition of tumor growth by **5** can be explained in part higher concentrations in tumor xenografts, which may exert direct
10 cytotoxic/cytostatic effect on the prostate cancer cells. It should be stated that there is evidence to suggest a possible direct cytotoxic effect of ketoconazole (a modest CYP 17 inhibitor) on prostate cancer cells.³³ In addition; the accumulation of **5** in the testes would enable inhibition of testosterone synthesis in the animals.

15 [0049] Although it is well established that LAPC4 are androgen-dependent, these cells can become androgen-independent, and as such represent a suitable model that mimics prostate cancer development in patients (Chen et al., above, and Kline et al., "Progression of metastatic human prostate cancer to androgen independence in immunodeficient SCID mice," *Nat. Med.*, 1997, 3, 402-408). As shown in Figure 7, we are able to replicate this phenomenon. Furthermore, our results show that treatment
20 with **16** or castration effectively suppresses tumor growth for a certain period (androgen-dependent phase), but was ineffective thereafter (possibly as a result of an androgen-independent phase) since the tumors grow rapidly just as in intact control mice. Tumor growth in the mice treated with **5** is strongly suppressed throughout the treatment period. This suggests that **5** may have effects on androgen-independent prostate cancer. However, it is also plausible that treatment with this compound enables
25 LAPC4 tumors to remain androgen dependent for a longer period and therefore responsive to antiandrogen therapy.

[0050] Recent studies that clearly demonstrate the up-regulation and involvement of AR in advanced and recurrent PCA (Mohler et al., and Chen et al., above) have renewed interest in the androgen
30 receptor as a target for development of drugs to treat PCA (Tindall et al., "Symposium on androgen action in prostate cancer", *Cancer Res.*, 2004, 64, 7178-7180). Because of its potent properties, **5** may be an excellent candidate.

Conclusions:

[0051] The data reinforce our earlier concept of modification of the C17 substituent of Δ^{16} steroids to
35 produce potent inhibitors of CYP17 as well as potent AR antagonists. The 17-benzimidazoles **5** and **6** are shown to coordinate the heme iron of CYP17, a property that may in part be

responsible for their enzyme inhibitory activity. Compounds **5** and **6** exhibit almost equipotent *in vitro* activities for CYP17 inhibition, AR antagonism, and inhibition of prostate cancer cell growth. Surprisingly, the compounds are very different in their antitumor activities, as **5** causes marked suppression of LAPC4 tumor xenograft growth, and in contrast, **6**(0.15 mmol/kg twice daily) enhances tumor growth. The present study provides compelling evidence that **5** is a potent inhibitor of human prostate tumor growth and is remarkably more effective than castration. This is the first example of a CYP17 inhibitor/antiandrogen demonstrating *in vivo* antitumor activity against a prostate cancer tumor to an extent that is superbly more effective than castration. These impressive biological activities, makes **5** a strong candidate for further development as a potential drug for the treatment of prostate cancer in humans. The excellent antitumor activity of compound **5**, containing a benzimidazole group makes the benzimidazoles a preferred group. However, analogs of **5** as discussed above are expected to have related activity.

Experimental Section

[0052] Chemistry: General procedures and techniques were identical with those previously reported (Njar et al., J. Med. Chem., 1998, 41, 902 - 912). Infra red spectra are recorded on a Perkin Elmer 1600 FTIR spectrometer using solutions in CHCl₃. High-resolution mass spectra (HRMS) are determined on a 3-Tesla Finnigan FTMS-2000 FT mass spectrometer, ESI mode (Ohio State University, Department of Chemistry). As a criterion of purity for key target compounds, we provided high resolution mass spectral data with HPLC chromatographic data indicating compound homogeneity. Low-resolution mass spectra (LRMS) are determined on a Finnegan LCR-MS. Melting points (mp) are determined with a Fischer Johns melting point apparatus and are uncorrected. Dehydroepiandrosterone and dehydroepiandrosterone acetate were purchased from Aldrich, Milwaukee, WI. 5-Tributylstannylypyrimidine and 2-tributylstannylypyrazine were purchased from Frontier Scientific, Inc., Logan, UT.

25

[0053] 3 β -Acetoxy-17-chloro-16-formylandrosta-5,16-diene (2): This compound prepared from 3 β -acetoxyandrost-5-en-17-one (1) as previously described, provided spectral and analytical data as described (Njar et al., J. Med. Chem., 1998, 41, 902 - 912).

[0054] 3 β -Acetoxy-17-(1H-benzimidazol-1-yl)-16-formylandrosta-5,16-diene (3): A mixture of 3 β -Acetoxy-17-chloro-16-formylandrosta-5,16-diene (**2**, 2.5 g, 6.65 mmol), benzimidazole (2.35 g, 19.9 mmol), and K₂CO₃ (2.76 g, 23.9 mmol) in dry DMF (20 mL) is stirred at ca. 80 °C under Ar for 1.5 h. After cooling to room temperature, the reaction mixture is poured onto ice-cold water (250 mL) and the resulting precipitate is filtered, washed with water, and dried to give a crude dirty white solid (ca. 2.9 g). Purification by FCC [petroleum ether/EtOAc/Et₃N (6:4:0.3)] gives 2.7 g (88.7%) of pure compound **3**: mp 227-230 °C; IR (CHCl₃) 3691, 3024, 2951, 2359, 1725, 1670, 1604, 1491, 1452,

1375, 1253, 1032, 897, 852, 818, 700, 657, 618, 576, 565, 550, 529, 511, 476 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.07 (s, 6H, 18- and 19- CH_3), 2.04 (s, 3H, 3 β - OCH_3), 4.60 (m, 1H, 3 α -H), 5.43 (br s, 1H, 6-H), 7.35 (br. s, 2H, aromatic-Hs), 7.85 (s, 1H, aromatic-H), 7.98 (s, 1H, aromatic-H), 7.98 (s, 1H, 2 1 -H) and 9.59 (s, 1H, 16-CHO). HRMS calcd 481.2462 ($\text{C}_{29}\text{H}_{34}\text{O}_3\text{N}_2\text{Na}^+$), found 481.2454.

5

[0055] **3 β -Acetoxy-17-(1*H*-benzimidazol-1-yl)androsta-5,16-diene (4):** A solution of 3 β -Acetoxy-17-(1*H*-benzimidazol-1-yl)-16-formylandrosta-5,16-diene (**3**, 2.04 g, 4.45 mmol) in dry benzonitrile (10 mL) was refluxed in the presence of 10% palladium on activated charcoal (1.02 g, i.e., 50% weight of **3**) for 5 h. After cooling to room temperature, the catalyst was removed by filtration through a Celite pad. The filtrate was evaporated, and the residue was purified by FCC [petroleum ether/EtOAc/Et₃N (7.5:3:0.5)] gave 1.41 g (73.8%) of pure compound **4**: mp 159-160 °C; IR (CHCl_3) 3687, 2947, 2854, 2358, 2340, 1725, 1633, 1609, 1557, 1489, 1454, 1373, 1291, 1253, 1195, 1136, 1031, 985, 910, 839, 735, 665, 590, 544, 533, 513, 502, 488 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.02 (s, 3H, 18- CH_3), 1.07 (s, 3H, 19- CH_3), 2.04 (s, 3H, 3 β - OCH_3), 4.62 (m, 1H, 3 α -H), 5.43 (br s, 1H, 6-H), 5.98 (s, 1H, 16-H), 7.30 (m, 2H, aromatic-Hs), 7.49 (s, 1H, aromatic-H), 7.81 (s, 1H, aromatic-H), and 7.95 (s, 1H, 2 1 -H). HRMS calcd 453.2512 ($\text{C}_{28}\text{H}_{34}\text{O}_2\text{N}_2\text{Na}^+$), found 453.2511.

[0056] **3 β -Hydroxy-17-(1*H*-benzimidazol-1-yl)androsta-5,16-dtene (5):** The acetate **4** (1.3 g 3.02 mmol) was dissolved in methanol (20 mL) under an inert Ar atmosphere, and the resulting solution treated with 10% methanolic KOH (8 mL). The mixture was stirred at room temperature for 1.5 h, and then concentrated under reduced pressure at approx. 40 °C to a volume of 10 mL. This solution was poured into ice water (300 mL), and the resulting white precipitate was filtered, washed with water and dried. Crystallization from EtOAc/MeOH gave **5** (1.10 g, 94%), mp 189-190 °C; IR (CHCl_3) 2934, 2339, 1609, 1490, 1453, 1291, 1040, 837, 808, 705, 663, 608, 578, 550, 517 cm^{-1} ; ^1NMR (300 MHz, CDCl_3) δ 1.02 (s, 3H, 18- CH_3), 1.07 (s, 3H, 19- CH_3), 3.55 (m, 1H, 3 α -H), 5.41 (br s, 1H, 6-H), 5.99 (s, 1H, 16-H), 7.30 (m, 2H, aromatic-Hs), 7.54 (s, 1H, aromatic-H), 7.80 (s, 1H, aromatic-H), and 7.96 (s, 1H, 2 1 -H). HRMS calcd 411.2407 ($\text{C}_{26}\text{H}_{32}\text{ON}_2\text{Na}^+$) found 411.2396.

[0057] **17-(1*H*-benzimidazol-1-yl)androsta-4,16-diene-3-one (6):** From a mixture of compound **5** (660 mg, 1.70 mmol), 1-methyl-4-piperidone (2.5 mL), and toluene (40 mL) was distilled off ca. 10 mL. Aluminum isopropoxide (521 mg, 2.55 mmol) was then added, and the mixture was refluxed under Ar for 4 h. After cooling, the mixture was diluted with EtOAc (50 mL), washed successively with 5% aqueous NaHCO_3 (x3) and brine (x2), and then dried (Na_2SO_4). The solvent was evaporated, and the crude product was purified by FCC [CH_2Cl_2 /FtOH (25:1)] to give the title compound **6** (544 mg, 82%): mp 201-204 °C; IR (CHCl_3) 2946, 2858, 1622, 1611, 1490, 1453, 1376, 1291, 1270, 1228, 1189, 893, 850, 837, 722, 662, 615, 568, 553, 537, 519 cm^{-1} ; ^1NMR (300 MHz, CDCl_3) δ 1.04 (s, 3H, 18- CH_3), 1.24 (s, 3H, 19- CH_3), 5.78 (s, 1H, 4-H), 5.99 (s, 1H, 16-H), 7.31 (m,

2H, aromatic-Hs), 7.48 (m, 1H, aromatic-H), 7.81 (s, 1H, aromatic-H), and 7.95 (s, 1H, 2¹-H). HRMS calcd 409.2250 (C₂₆H₃₀ON₂.Na⁺), found 409.2250.

[0058] **Reaction of 3 β -acetoxy-17-chloro-16-formylandrosta-5,16-diene (2) with benzo-1H-1,2,3-triazole and K₂CO₃: 3 β -Acetoxy-17-(benzo-2H-1,2,3-triazol-2-yl)-16-formylandrosta-5,16-diene (7a) and 3 β -Acetoxy-17-(benzo-1N-1,2,3-triazol-yl)-16-formylandrosta-5,16-diene (7b):**

A mixture of compound 2 (2.5 g, 6.65 mmol), benzotriazole (2.35g, 19.9 mmol), and K₂CO₃ (2.76 g, 23.9 mmol) in dry DMF (20 mL) was stirred at ca. 80 °C under Ar for 45 min. After cooling to room temperature, the reaction mixture was poured onto ice-cold water (250 mL) and the resulting precipitate was filtered, washed with water, and dried to give a crude dirty white solid. Purification by FCC [pet. Ether/EtOAc, (4:1)] first gave 3 β -acetoxy-17-(benzo-2H-1,2,3-triazol-2-yl)-16-formylandrosta-5,16-diene (**7a**, 0.3 g, 9.8 %) as minor product; mp 248-250 °C; IR (CHCl₃) 3023, 2945, 2358, 1725, 1657, 1600, 1375, 1257, 1032, 728, 656, 584, 564, 540, 526, 506, 498 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.11 (s, 3H, 18-CH₃), 1.37 (s, 3H, 19-CH₃), 2.04 (s, 3H, 3 β -OCH₃), 4.62 (m, 1H, 3 α -H), 5.43 (br s, 1H, 6-H), 7.43 (d, 1H, *J* = 2.4 Hz, aromatic-Hs), 7.45 (d, 1H, *J* = 2.7 Hz, aromatic-H), 7.88 (d, 1H, *J* = 2.7 Hz, aromatic-H), 7.90 (d, 1H, *J* = 2.4 Hz, aromatic-H) and 10.66 (s, 1H, 16-CHO). HRMS calcd 482.2414 (C₂₈H₃₃O₃N₃.Na⁺), found 482.2413. Further elution with the same solvent system afforded the major product, 3 β -acetoxy-17-(benzo-1H-1,2,3-triazol-yl)-16-formylandrosta-5,16-diene (**7b**, 2.3 g, 75.4 %); mp: 186-188 °C; IR(CHCl₃) 3023, 2948, 1725, 1670, 1604, 1488, 1450, 1374, 1253, 1196, 1032, 846, 824, 720, 658, 619, 548, 527, 504, 497 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.07 (s, 6H, 18- and 19-CH₃), 2.04 (s, 3H, 3 β -OCH₃), 4.60 (m, 1H, 3 α -H), 5.43 (br s, 1H, 6-H), 7.46 (m, 2H, aromatic-Hs), 7.57 (d, 1H, *J* = 6.9 Hz, aromatic-H), 8.15 (d, 1H, *J* = 8.4 Hz, aromatic-H), and 9.59 (s, 1H, 16-CHO). HRMS calcd 482.2414 (C₂₈H₃₃O₃N₃.Na⁺), found 482.2416.

[0059] **3 β -Acetoxy-17-(beuzo-1H-1,2,3-triazol-1-yl)androsta-5,16-diene (8):** A mixture of bis(triphenylphosphine)rhodium(I) carbonyl chloride (303 mg, 0.438 mmol) and 1,3-bis(diphenylphosphino)propane (394 mg, 0.954 mmol) in dry xylene (40 mL) was stirred at 80 °C under Ar for 15 min when fine yellow precipitate formed. Compound 7b (1.71 g, 3.72 mmol) was added, and the mixture was refluxed under Ar for 18 h, and then concentrated under reduced pressure. The crude product was purified by FCC [pet ether/BtOAc/Et₃N, (8.9:1:0.1)] to give 1.2 g (74.7 %) of pure compound **8**; mp 184-186 °C. IR (CHCl₃) 3063, 2918, 2389, 2358, 1725, 1458, 1373, 1254, 1069, 1031, 843, 809, 786, 692, 646, 560, 535, 528, 512, 494 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.10 (s, 3H, 18-CH₃), 1.25 (s, 3H, 19-CH₃), 2.04 (s, 3H, 3 β -OCH₃), 4.64 (m, 1H, 3 α -H), 5.43 (br s, 1H, 6-H), 6.01 (s, 1H, 16-H), 7.40 (t, 1H, *J* = 7.8 Hz, aromatic-H), 7.51 (t, 1H, *J* = 7.8 Hz, aromatic-H), 7.67 (d, 1H, *J* = 8.1 Hz, aromatic-H), and 8.10 (d, 1H, *J* = 8.1 Hz, aromatic-H). HRMS calcd 454.2465 (C₂₇H₃₃O₂N₃.Na⁺), found 454.2469.

[0060] **3 β -Hydroxy-17-(benzo-1*H*-1,2,3-triazol-1-yl)androsta-5,16-diene (9):** The method followed that described for compound 5 but using 3 β -acetoxy-17-(benzo-1*H*-1,2,3-triazol-1-yl)androsta-5,16-diene (8; 700 mg, 1.62 mmol). Recrystallization from EtOAc/MeOH give the title compound 9 (600

5 mg, 95%); mp 241-244 °C; IR (CHCl₃) 3603, 2937, 2859, 1609, 1488, 1451, 1373, 1287, 1243, 1069, 1040, 1007, 953, 845, 805, 715, 665, 618, 570, 553, 517 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.09 (s, 3H, 18-CH₃), 1.24 (s, 3H, 19-CH₃), 3.55 (m, 1H, 3 α -H), 5.41 (br s, 1H, 6-H), 6.06 (s, 1H, 16-H), 7.40 (t, 1H, *J* = 7.8 Hz, aromatic-H), 7.52 (t, 1H, *J* = 7.8 Hz, aromatic-H), 7.67 (d, 1H, *J* = 8.1 Hz, aromatic-H), and 8.10 (d, 1H, *J* = 8.1 Hz, aromatic-H). HRMS calcd 412.2359 (C₂₅H₃₁ON₃.Na⁺), found 10 412.2365.

[0061] **17-(benzo-1*H*-1,2,3-triazol-1-yl)androsta-4,16-diene-3-one (10):** The method followed that described for compound 6 but using β -hydroxy-17-(benzo-1*H*-1,2,3-triazol-1-yl)androsta-5,16-diene (9; 500 mg, 1.28 mmol). Purification of the crude product by FCC [CH₂Cl₂/EtOH, (50:1)] afforded the 15 titled compound 10 (420 mg, 84.4%); mp: 280-283 °C; IR (CHCl₃) 2944, 1658, 1450, 1070, 8444, 825, 721, 624, 589, 564, 554, 541, 521 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.26 (s, 3H, 18-CH₃), 1.27 (s, 3H, 19-CH₃), 5.77 (s, 1H, 4-H), 6.01 (s, 1H, 16-H), 7.40 (t, 1H, *J* = 7.8 Hz, aromatic-H), 7.52 (t, 1H, *J* = 7.8 Hz, aromatic-H), 7.67 (d, 1H, *J* = 7.8 Hz, aromatic-H), and 8.10 (d, 1H, *J* = 8.1 Hz, aromatic-H). HRMS calcd 410.2203(C₂₅H₂₉ON₃.Na⁺), found 20 410.2185.

[0062] **Dehydroepiandrosterone-17 hydrozone (12):** Dehydroepiandrosterone (11, 3.5 g, 12.2 mmol) was dissolved in ethanol (60 mL); and the resulting solution was treated with hydrazine hydrate (2.37 mL, 0.049 mol) followed by a solution of hydrazine sulfate (7.9 mg, 0.061 mmol) in 0.25 mL of water. The mixture was stirred at room temperature for 12 h and then poured into ice water. The resulting 25 precipitate was filtered, washed with water, and dried to give white crystals of the titled compound 12; mp: 242-244 °C (lit. 204-206 °C);²² ¹H NMR (300 MHz, CDCl₃): δ 0.76 (s, 3H, 18-CH₃), 1.05 (s, 3H, 19-CH₃), 3.74 (br s, 1H, 3-H) and 5.35 (s, 1H, 6-H).

[0063] **17-Iodoandrosta-5,16-diene-3 β -ol (13):** A stirred solution of iodine (12.16 g, 0.0203 mol) in 30 dry of THF (144 mL) and dry of Et₂O (72 mL) was cooled in an ice bath to 0 °C and the solution was treated with 1,1,3,3, tetramethylguanidine (6.72 mL, 6.24 g, 0.054 mole). A solution of compound 12 (3.0 g, 9.9 mmol) in THF (81 mL) was added dropwise to the iodine solution over 2 h maintaining the reaction temperature at 0 °C. The reaction mixture was then concentrated under vacuum, cooled in an ice-bath, and then dried to under vacuum at room temperature to afford a yellow solid (13, 3.65 g, 35 92.4%). mp: 169-171 °C. (lit. 175-176 °C);²² IR (CHCl₃) 2935, 1371, 1039, 862, 843, 799, 715, 665, 582, and 566 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 0.76 (s, 3H, 18-CH₃), 1.05 (s, 3H, 19-CH₃), 3.50 (br s, 1H, 3 α -H), 5.35 (s, 1H, 6-H) and 6.14 (s, 1H, 16-H).

[0064] **3 β -Hydroxy-17-(2-pyrazyl)-androsta-5,16-diene (14):** A mixture of 17-idoandrosta-5,16-diene-3 β -ol (**13**; 0.5 g, 1.257 mmol) in solution with dry dimethylformamide (DMF, 10 mL) along with tetrakis(triphenylphosphine) palladium ($\text{Pd}(\text{PPh}_3)_4$) (71.6 mg, 0.062 mmol) and (2-tributylstannylyl) pyrazine (774.6 mg, 2.099 mmol) was heated at 120 °C for 20 h. After cooling, the mixture was diluted with cold water (50 mL), and extracted with EtOAc (30 mL x 3). The combined EtOAc extract was washed with brine and water, dried over Na_2SO_4 and then concentrated to give a brownish solid. This crude product was purified by flash column chromatography [FCC, pet.ether/EtOAc/Et₃N (3:2:0.15)] to give **14** (66 mg, 15%); mp: 199-201 °C. ¹HNMR (300 MHz, CDCl_3): δ 0.94 (s, 3H, 18-CH₃), 1.08 (s, 3H, 19-CH₃), 3.52 (br s, 1H, 3 α -H), 5.40 (s, 1H, 6-H), 6.77 (s, 1H, 16-H), 8.35 (s, 1H, pyrazine-H), 8.48 (s, 1H, pyrazine-H), 8.70 (s, 1H, pyrazine-H). HRMS calcd 350.2358 ($\text{C}_{23}\text{H}_{30}\text{ON}_2$), found 350.2354.

[0065] **3 β -Hydroxy-17-(5-pyrimidyl)-androsta-5,16-diene (15):** Reaction of **13** (0.645 g, 1.623 mmol) as described above for **14**, but using (5-tributylstannylyl) pyrimidine (1.0 g, 2.710 mmol) dissolved in 10 mL of dry DMF along with CPd($\text{PPh}_3)_4$ (92.88 mg, 0.0804 mmol) and (5-tributylstannylyl) pyrimidine (1.0 g, 2.710 mmol) and following purification by [FCC, pet.ether/EtOAc/Et₃N (3:2:0.15)] gave 3 β -hydroxy-17-(5-pyrimidyl)-androsta-5,16-diene **15** (44 mg, 10%); mp: 231-233 °C (lit. 240-242 °C); ¹HNMR (300 MHz, CDCl_3): δ 1.05 (s, 3H, 18-CH₃), 1.08 (s, 3H, 19-CH₃), 3.83 (br s, 1H, 3 α -H), 5.39 (s, 1H, 6-H), 7.26 (s, 1H, 16-H), 8.73 (s, 2H, 4¹-H and 6¹-H) and 9.07 (s, 1H, 2¹-H). HRMS calcd 350.2358 ($\text{C}_{23}\text{H}_{30}\text{ON}_2$), found 350.2348,

[0066] **In Vitro Assay of CYP17:** The *in vitro* CYP17 inhibitory activities of the compounds are evaluated using our rapid acetic acid releasing assay (AARA), utilizing intact P450c17-expressing *E. coli* as the enzyme source (Grigoryev, above). It involves the use of [21-³H]-17 α -hydroxypregnolone as the substrate and CYP 17 activity is measured by the amount of tritiated acetic acid formed during the cleavage of the C-21 side chain of the substrate. This establishes that the method is comparable in terms of accuracy and reliability to the HPLC analysis procedure used by researchers in the field (Grigoryev, above). IC₅₀ values are obtained directly from plots relating percentage inhibition versus inhibitor concentration over appropriate ranges. Each compound is tested at a minimum of five different concentrations. The assays are performed in triplicate, and the IC₅₀ values reported are the mean of triplicate experiments. The standard deviations were \pm 5% of the mean values.

[0067] **Human 5 α -Reductase Type 1 and 2 Assay:** The inhibitory activities of compounds and finasteride as reference are determined using the DU145 cell line (for human type 1 enzyme) and human prostate homogenate (BPH tissue for type 2 enzyme) according to the procedure described by Hartmann and colleagues (Picard et al., "Synthesis and evaluation of 2'-substituted 4-(4'-carboxy- or 4'-

carboxymethylbenzylidene)-N-acylpiperidines: Highly potent and in vivo active steroid 5 α -reductase type 2 inhibitors", J. Med. Chem., 2002, 45, 3406-3417). The percent inhibition values at a concentration of 10 μ M or, in case of more potent compounds, the IC₅₀ values are determined.

5 [0068] Competitive androgen receptor (AR) binding and luciferase assays: AR

Binding/Competition Assay: Wells in 24-well multiwell dishes are coated with poly-L-lysine (0.05mg/ml) for 5 minutes, dried, rinsed with sterilized, distilled, water, and dried for 2 hours. To determine the kinetics of R1881 binding to the LNCaP AR and the wild-type AR, LNCaP and PC3AR cells are plated (2-3 x 10⁵) in 24 well multiwell dishes in steroid-free medium and allowed to attach.

10 The following day the medium is replaced with serum-free, steroid free RPMI supplemented with 0.1%BSA and containing [³H]R1881 (0.01-10nM) in the presence or absence of a 200 fold excess of cold DHT, to determine nonspecific binding, and 1 μ M triamcinolone acetonide to saturate progesterone and glucocorticoid receptors. Following a 2-hour incubation period at 37°C, cells are washed twice with ice-cold DPBS and solubilized in DPBS containing 0.5% SDS and 20% glycerol. Extracts are removed
15 and cell associated radioactivity counted in a scintillation counter. The data is analyzed, including Kd and Bmax determination, by nonlinear regression using Graphpad Prism software. When the concentration required to almost saturate AR in both cell lines is established, the ability of the test compounds (0.1nM-10 μ M) to displace [³H]R1881 (5.0nM) from the receptors is determined as described above. The IC₅₀ of each compound is determined by nonlinear regression with Graphpad
20 Prism software (GraphPad Software, Inc, San Diego, CA). *Luciferase Transactivation Assay:* Transcriptional activation assay is carried out as described previously by Kim *et al.*, above, with minor modifications. The Probasin luciferase reporter construct ARR2-Luc is generated by insertion of the minimal probasin promoter ARR2, kindly provided by Dr R Matusik of Vanderbilt University Medical Center (Endocrinology, 2000, 141: 4698-4710) into the polyclonal linker region of PGL3-
25 enhaticer vector (Promega). The pRL-null (Promega) is used as the internal control. Briefly, LNCaP cells grown in 24-well plates coated with poly-L-lysine were transfected with ARR2-Luc in the phenol-red free RPMI 1640 medium containing 5% charcoal-stripped FBS (Hyclone). 24 h post-transfection, the cells are incubated with fresh phenol-red free serum-free RPMI 1640 medium with or without DHT and inhibitors for 18 h. Luciferase activities are measured in triplicates by using dual luciferase assay
30 system according to the manufacturer's instruction (Promega). The results are presented as the fold induction, that is, the relative luciferase activity of the treated cells divided by that of the control.

[0069] Cell Culture and Viability Assay: LNCaP cells are grown in RPMI 1640 medium

supplemented with 10% FBS and 1% penicillin/streptomycin solution. To determine the effect of novel
35 compounds on cell proliferation, cells are transferred into steroid-free medium three days prior to the start of the experiments. Steroid-free medium consisted of phenol red free RPMI supplemented with 5% dextran-coated, charcoal treated serum, and 1 % penicillin/streptomycin solution. Growth studies are

then performed by plating cells (3×10^4) in 24-well multi-well dishes (Corning, Inc. Corning, NY).

After a 24 hours attachment period, the medium is aspirated and replaced with steroid-free medium containing vehicle or the indicated concentration of DHT (1nM) and compounds (0.1 μ M-10 μ M).

Control wells are treated with vehicle (ethanol). This medium is changed every three days and the

5 number of viable cells is compared by WST-1 [4-[3-(4-iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazolio]-1,3-benzene disulfonate] assay on the seventh day. Following incubation of cells for the above-mentioned time, 10% WST-1 solution is added to each well and incubated at 37°C for three hours.

Following incubation, plates are slightly shaken and immediately read at 450 nm with a scanning multi-well spectrophotometer. All results represent the average of a minimum of three wells. Additional

10 control consists of medium alone with no cells.

[0070] Pharmacokinetic Studies: All animal studies are performed according to the guidelines and approval of the Animal Care Committee of the University of Maryland School of Medicine, Baltimore.

Male SCID mice weighing 20-22 gm (8-10 weeks old) obtained from NCI, Frederick, MD, USA are

15 maintained in a controlled environment of about 25°C, 50% relative humidity and 12 h of light and 12 h of dark cycles and allowed free access to food and water. Compounds 5 and 6 are formulated in 40% β -cyclodextrin in water and a single subcutaneous dose is given to mice. The animals are sacrificed at various times up to 6 h after drug administration and blood was obtained by cardiac puncture under light halothane (Ayerst, New York, NY, USA) anesthesia.

20

[0071] HPLC Analysis: Chromatographic separations and quantification of the steroids and the appropriate internal standards are achieved by a reverse phase HPLC method on a

Waters® Novapak® C18 column (3.9 x 150 mm) protected by Waters® guard cartridge packed with pellicle C18 as previously described. Briefly, the HPLC system used in this study consisted of

25 Waters® solvent delivery system, Waters® controller (Milford, MA), coupled to a Waters® 717^{plus} autosampler and a Waters® 996 photodiode array detector operated at 242.7 nm. The mobile phase composition is Water/McOH/CH₃CN (35:35:30, v/v/v + 200 μ L of Et₃N and 0.77 g of NH₄OAc per 1000 mL of mobile phase) at a flow rate of 1.0 mL/min. The HPLC analysis is performed at ambient temperature and data acquisition and management is achieved with a Waters® millennium

30 chromatography manager. **Sample Preparation:** Test tubes containing mouse plasma (200 μ L), **5** or **6** and VN/85-1 (internal standard, 10 μ L of 100 μ g/mL), are extracted with diethyl ether (2 x 2 mL) using a vortex mixer for 3 minutes and centrifuged at 3000 g for 5 min. The organic layers are evaporated to dryness under a gentle stream of air. The residue is reconstituted in an aliquot of the mobile phase (100 μ L) and filtered using 0.2 μ m Teflon filters before HPLC analysis. **Calibration**

35 **Curve and HPLC Assay Validation:** The calibration curves for **5** in plasma and tissue and for **6** in plasma are constructed by spiking varying amounts of the compounds into extraction tubes (duplicate) containing plasma (200 μ L) and tissue preparations (200 μ L) from untreated animals to give final

concentrations of 0.1 - 100.0 μ g/mL. Appropriate blank extraction tubes are also prepared and an aliquot of the internal standard is added into each extraction tube to give a final concentration of 5 μ g/ml. The calibration samples are taken through the sample preparation procedure as described above. An aliquot of the reconstituted extract (50 μ l) is injected into the HPLC system and the ratio of

5 the peak areas for each analyte to that of the internal standard are plotted against concentrations of **5** or **6**. The precision and accuracy of the assays are determined from a range of known concentrations of the inhibitors in blank plasma and taken through the HPLC procedure. The study is repeated on three separate occasions.

10 **[0072] Data Analysis:** Pharmacokinetic calculations are performed as previously described. The non-compartmental pharmacokinetic calculations are performed using WinNONlin (Scientific Consulting Inc.). One-way analysis of variance (ANOVA) on SigmaStat for Windows version 1.0 is used to compare different treatment groups at the 95%'confidence level. The Bonferroni post-hoc test is used for determination of significance. A P-value of less than 0.05 is considered as statistically significant. **In Vivo Antitumor Studies (LAPC-4 Prostate Cancer Xenografts):** All animal studies are performed according to the guidelines and approval of the Animal Care Committee of the University of Maryland School of Medicine, Baltimore. Male severe combined immunodeficient (SCID) mice 4-6 weeks of age purchased from the National Cancer Institute-Frederick Cancer Research and Development Center (Fredrick, MD) are housed in a pathogen-free environment under controlled 15 conditions of light and humidity and allowed free access to food and water. Tumors are developed from LAPC4 cells inoculated subcutaneously (s.c.) in the mice essentially as previously described (21). LAPC4 cells are grown in IMEM with 15% FBS plus 1% PS and 10nm DHT until 80% confluent. Cells are scraped into DPBS, collected by centrifugation, and resuspended in Matrigel (10 mg/ml) at 3 x 20 10⁷cells/ml. Mice are injected s.c. with 100 μ l of the cell suspension at one site on each flank. Tumors are measured weekly with calipers, and tumor volumes are calculated by the formula: $4/3\pi \times r_1^2 \times r_2$ ($r_1 < r_2$). 25

30 **[0073]** In the first experiment, LAPC4 tumors are allowed to grow for 8-10 weeks following inoculation. Groups of 5 mice with comparable total tumor volumes are either castrated or treated with **5** and **6** (0.15 mmol/kg once-daily and 0.15 mmol/kg twice-daily, 9 a.m. and 5 p.m.). Mice are castrated under methoxyfluorane anesthesia. Compounds **5** and **6** were prepared at 17.2 mg/ml in a 0.3% solution of hydroxypropyl cellulose in saline, and mice receiv s.c. injections daily. Control and castrated mice are treated with vehicle only. Tumors are measured weekly for the 4 weeks of treatment and tumor volumes are calculated. At the end of the treatment period, the animals are sacrificed under 35 halothane anesthesia; tumors are excised, weighed and stored at -80°C. Animals are also weighed weekly and monitored for general health status and signs of possible toxicity due to treatment.

[0074] In the second experiment, mice are inoculated with LAPC4 cells and are divided into four groups of 5 mice each. The control and castrated group receive vehicle, while the other two groups receive either VN/85-1 (0.15 mmol/kg twice-daily, 9 a.m. and 5 p.m.) or **5** (0.15 mmol/kg twice-daily, 9 a.m. and 5 p.m.). These treatments are initiated one day after LAPC4 cell inoculation; continued for 14 weeks for control group, 19 weeks (for VN/85-1 and castration groups) and for 21 weeks for 5 treated group and tumors are measured and processed as described above.

[0075] **Measurement of 5 (VN/124-1) levels in tumor, liver and testes:** The animals in the VN/124-1-treated group are sacrificed 1 h after the last VN/124-1 administration, and tumor, liver and testis are harvested and snap frozen in liquid nitrogen. Tissue samples are homogenized in phosphate buffer (pH = 7.4, 0.5 ml/mg of tissue). Homogenized tissue (200 μ l) is spiked with the internal standard, VN/85-1 (10 μ L from 100 μ g/mL stock solution), and then extracted with Et₂O (2 x 2 mL) by vortexing for 3 min followed by centrifugation at 3000g for 5 min. The Et₂O extract is separated and evaporated to dryness under a gentle stream of air. The residue is reconstituted in 100 μ L of the HPLC mobile phase, filtered through 0.2 μ m Teflon filters and then analyzed by HPLC as described above.

[0076] The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

Table 1: CYP17 and 5 α -reductase activities and androgen receptor binding of novel 17-beteroaryl compounds.

Compound ^a	CYP17 IC ₅₀ (nM) ^b	5 α -Reductase % inhibition at 10 μ M [IC ₅₀ (nM) ^b]		AR Binding IC ₅₀ (nM) ^c	
		type 1 ^d	type 2 ^e	LNCaP	PC3-AR
5	300.0	4	53	845	384
6	915.0	[770]	[480]	1200	242
9	1250.0	n ^f	17	-	-
10	5817.4	21	56	-	-
14	3810.0	-	-	-	366
15	500.0	-	-	-	374
<i>For comparison</i>					

Compound ^a	CYP17 IC ₅₀ (nM) ^b	5 α -Reductase % inhibition at 10 μ M [IC ₅₀ (nM) ^b]		AR Binding IC ₅₀ (nM) ^c	
		type 1 ^d	type 2 ^e	LNCaP	PC3-AR
VN/85-1	50.0	-	-	-	-
Abiraterone	800.0	-	-	-	-
Ketoconazole	1100.0	-	-	-	-
Finasteride	-	[60.0]	[2.0]	-	-
Casodex	-	-	-	940	-
Flutamide	-	-	-	11600	10985

^a We have previously reported the synthesis of VN/85-1 (Njar et al., above)

Abiraterone was synthesized as described by Potter et al (A convenient, large-scale synthesis of abiraterone acetate [3 β -acetoxy-17(3-pyridyl)androsta-5,16-diene], a potential new drug for the treatment of prostate cancer. Org. Prep. Proc. Int., 1997, 29, 123-128).

^b IC₅₀ is the concentration of inhibitor required to inhibit the enzyme activity by 50%, each in duplicate for CYP7, triplicate for 5 α -reductase and AR binding.

^c IC₅₀ is the concentration of compound required for a 50% displacement of [³H]R1881 from the androgen receptor.

^d Prostatic tumor cell line (DU-145) expressing type 1 enzyme; substrate: 5 nM [1 β -³H]androstenedione.

^e Enzyme from BPH tissue (type 2 enzyme), 125 μ g of protein, substrate: 210 nM [1 β ,2 β -³H]testosterone.

^f ni = no inhibition up to 10 μ M. - = not determined.

Table 2: Pharmacokinetic parameters for **5** (50 and 100 mg/kg) and **6** (50 mg/kg) after s.c. administration.

Parameter ^a	5		6
	50 mg/kg	100 mg/kg	50 mg/kg
- t _{1/2} (min)	44.7 \pm 1.15	36.6 \pm 1.6	37.93 \pm 1.15
K _{el} (min ⁻¹)	56.5 \pm 0.94	68.49 \pm 1.26	0.0183 \pm 0.004
AUC (mm. μ g/mL)	1440.00 \pm 60.23	1813.94 \pm 10.94	647.10 20.23
T _{max} (min)	30.00 \pm 0.0	30.00 \pm 0.0	60.00 \pm 0.00

Parameter ^a	5		6
	50 mg/kg	100 mg/kg	50 mg/kg
C _{max} (μg/mL)	16.82 ± 0.37	32.23 ± 0.34	5.15 ± 0.09
MRT (min)	65.40 ± 0.60	60.46 ± 1.54	79.95 ± 0.01
V _d (mL/kg)	2098.99 ± 4.11	3276.39 ± 26.71	4207.24 ± 6.25

^a Values are expressed as mean ± S.E., n = 5.

Brief Description of Schemes and Figures

[0077]

Chart 1:

5 Structures of abirateron and **VN/85-1 (16)**

Scheme 1:

Synthesis of 17-benzoazole compounds (**5**, **6**, **9** and **10**).

Scheme 2:

Synthesis of 17-diazine compounds (**14** and **15**).

10 **Scheme 3:**

Synthesis of metabolites of trans-andro sterone, including VN LG/81.

Figure 1:

The effects of **5**, **6** and casodex on transcriptional activity of luciferase mediated through LNCaP-AR in LNCaP-ARR2-lu prostate cancer cells. Cells in steroid-free medium were treated with vehicle, or increasing concentrations of either **5** or casodex with and without 1 nM DHT for 18h. Cells were then assayed for luciferase activity as described in "Materials and Methods". The bars represent the mean light units [counts per second (cps)/unit protein, i.e., relative luciferase activity] in triplicate wells from three separate experiments.

Figure 2:

20 The effects of **5**, **6** and casodex on (a) LNCaP and (b) LAPC4 prostate cancer cell growth. Cells were grown in steroid-free medium before plating. Triplicate wells were then co-treated with increasing concentrations of **5**, **6** or casodex and DHT as described in "Materials and Methods." The percentage (compared to control) of growth inhibition after 7 days of treatment was determined using WST-1 assay, The results represent the average and standard deviation of 25 three experiments performed in triplicate.

Figure 3:

Typical HPLC chromatogram of **5**, **16** (internal standard) and metabolite extracted from mouse plasma. The retention times for **16**, metabolite, and **5** were 11.5, 17.3 and 21.6 min, respectively.

Figure 4:

Pharmacokinetic profiles of **5** and **6** following administration of a single subcutaneous bolus dose to male SCID mice. Each data point represents the mean plasma concentrations obtained from three mice. The standard deviations (not shown) were \pm 5-8% of the mean values.

5 Figure 5:

Pharmacokinetic profiles of **5** and metabolite following a single subcutaneous bolus dose (100 mg/kg.bw) of **5** to male mice.

Figure 6:

In vivo antitumor activity of **5**, **6** and orchidectomy on the growth of LAPC4 prostate tumors in male SCID mice. Groups of five mice with LAPC-4 tumors were treated with **5** (0.15 mmol/kg/day or 0.30 mmol/kg/day. Tumors volumes were measured weekly, and the percentage of change in tumor volume was determined after 28 days of treatment. The standard deviations of tumor volumes (not shown) were \pm 10-12 % of the mean values.

Figure 7:

The effects of **5**, **16**, and orchidectomy on the formation and growth of LAPC4 prostate tumors in male SCID mice. 3×10^7 LAPC-4 cells were injected s.c. into the dorsal flank of SCID mice. One group of mice was castrated. The other groups of mice received either vehicle or **5** (0.15 mmol/kg twice-daily) or **16** (0.15 mmol/kg twice-daily). Daily treatment with **5** or **16** was initiated 1 day after cell inoculation. Tumors volumes were measured weekly, and the percentage of change in tumor volume was determined after 16 weeks of treatment. * Indicates significant difference of **5** versus control, castration and **16** at week 14 ($P = 0.00065$, 0.05 and 0.0097, respectively). ** Indicates significant difference of **5** versus castration and **16** at week 16 ($P = 0.047$ and 0.0047, respectively). \downarrow - $\downarrow\downarrow$: Period of reduced administered dose of **5**.






Chart 1. Structures of Abiraterone and VN/85-1




Scheme 1. (i) POCl_3 -DMF, CHCl_3 , Ar, reflux; (ii) benzimidazole, K_2CO_3 , DMF, Ar, 80°C ;
 (iii) 10% Pd on activated charcoal, PhCN, reflux; (iv) 10% Methanolic KOH, Ar, rt.;
 (v) $\text{Al}(i\text{-PrO})_3$, 1-methyl-4-piperidone, toluene, reflux; (vi) benzo- $1H$ -1,2,3-triazole,
 K_2CO_3 , DMF, Ar, 80°C ; (vii) $(\text{PPh}_3)_2\text{RhCOCl}\text{-Ph}_2(\text{CH}_2)_3\text{PPh}_2$, xylene, Ar, reflux.




Scheme 2: i) $\text{N}_2\text{H}_4\text{H}_2\text{O}$, $\text{N}_2\text{H}_4\text{H}_2\text{SO}_4$, EtOH; ii) I_2 /THF, TG; iii) (2-tributylstannylypyrazine/Pd(PPh_3)₄; iv) (5-tributylstannylypyrimidine/Pd(PPh_3)₄

Scheme 3: Synthesis of Metabolites of trans-androsterone, including VNLG/81



Patentkrav

1. Farmaceutisk sammensætning, der omfatter 3β -hydroxy-17-(1*H*-benzimidazol-1-yl)androsta-5, 16-dien eller et farmaceutisk acceptabelt salt deraf til anvendelse i behandling af en androgen-
5 afhængig tumor ved inhibering af både vildtype og muterede androgenreceptorer.
2. Farmaceutisk sammensætning ifølge krav 1 til anvendelse ifølge krav 1 ved oral administration.
3. Farmaceutisk sammensætning ifølge krav 1 til anvendelse ifølge krav 1, hvor
10 sammensætningen ligeledes omfatter et andet fysiologisk aktivt middel.

**FIG. 1**




FIG. 2a




FIG. 2b






FIG. 3

**FIG. 4****FIG. 5**




FIG. 6




FIG. 7