

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
6 March 2003 (06.03.2003)

PCT

(10) International Publication Number
WO 03/018549 A2(51) International Patent Classification⁷: **C07D**

(21) International Application Number: PCT/US02/28007

(22) International Filing Date: 29 August 2002 (29.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/316,386 30 August 2001 (30.08.2001) US(71) Applicant (for all designated States except US): **CHEMO-CENTRYX, INC.** [US/US]; 1539 Industrial Road, San Carlos, CA 94070 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MCMASTER, Brian, E.** [US/US]; 120 Walker Drive, Mountain View, CA 94043 (US). **SCHALL, Thomas, J.** [US/US]; 530 Webster Place, #5, Palo Alto, CA 94301 (US). **PENFOLD, Mark** [AU/US]; 822 Calderon Avenue, Mountain View, CA 94041 (US). **WRIGHT, J., J.** [AU/US]; 740 Bair Island Road #301, Redwood City, CA 94063 (US). **DAIRAGHI, Daniel, J.** [US/US]; 178 El Dorado Avenue, Palo Alto, CA 94306 (US).(74) Agents: **KEZER, William, B.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, Eighth Floor, San Francisco, CA 94111 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).**Declaration under Rule 4.17:**


— of inventorship (Rule 4.17(iv)) for US only

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BICYCLIC COMPOUNDS AS INHIBITORS OF CHEMOKINE BINDING TO US28



(57) **Abstract:** Methods for treating CMV or a CMV-related disease are provided that use compounds having the formula (I) wherein Ar is a substituted or unsubstituted 5-14 membered heteroaryl group having from 1 to 5 heteroatoms as ring members; R¹ is selected from the group consisting of substituted or unsubstituted aryl(C₁-C₄)alkyl, heteroaryl(C₁-C₄)alkyl, -C(O)R¹¹, and -C(O)NR¹¹R¹², wherein each R¹¹ and R¹² independently is substituted or unsubstituted aryl, substituted or unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl; R² is H or (C₁-C₈)alkyl; and Z^N is a substituted or unsubstituted hetero(C₆-C₁₀)bicycloalkyl group.



WO 03/018549 A2

BICYCLIC COMPOUNDS AS INHIBITORS OF CHEMOKINE BINDING TO US28

CROSS-REFERENCES TO RELATED APPLICATIONS

- 5 **[0001]** This application claims the benefit of U.S. Provisional Application Ser. No. 60/316,386, filed August 30, 2001, the contents of which are incorporated herein by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

- 10 **[0002]** The United States government may have certain rights to inventions described herein pursuant to DARPA grant No. N66001-01-C-8009.

REFERENCE TO A "SEQUENCE LISTING," A TABLE, OR A COMPUTER PROGRAM LISTING APPENDIX SUBMITTED ON A COMPACT DISK.

- 15 **[0003]** NOT APPLICABLE

BACKGROUND OF THE INVENTION

- 20 **[0004]** Cytomegalovirus (CMV) is an important human pathogen and a major opportunist which emerges to cause disease in the immuno-compromised such as AIDS patients, neonates, and individuals who have been given immunosuppressive drugs as part of a transplantation regimen. In these individuals, the consequences of CMV in acute or re-emerging infections can be dire, including retinitis, encephalitis, and pneumocystis,
- 25 among other pathologies. Furthermore, in immuno-competent hosts, CMV establishes a persistent lifelong infection through which it has been linked to a variety of inflammatory conditions including coronary artery occlusion following heart transplant and atherectomy and restenosis following angioplasty. CMV interacts with leukocytes during acute infection of the host as well as during lifelong latency. As such, leukocytes are
- 30 important players in CMV-induced disease and have been implicated in the acute phase of infection as vehicles for dissemination of virus and as sites of residence during lifelong latency.

[0005] CMV harbors in its genome an open reading frame (ORF), designated US28, which encodes a protein that acts as a functional receptor for certain human and viral chemokines. Upon infection of a cell by CMV, US28 is expressed on the surface of the infected cell and becomes capable of responding to chemokines in the environment.

5 Because the virus on its own is inherently non-motile, and because chemokines and their receptors encoded by human cells are known to regulate the migration of leukocytes and other cells through the body, CMV US28 is thought to be encoded by the virus to facilitate the dissemination of CMV through the body during and after infection. Therefore, agents which block the binding of chemokines to US28 should prove useful in
10 inhibiting viral dissemination during acute or re-emerging CMV infection.

[0006] CMV US28 has been shown to bind a variety of human, murine, and virus-encoded CC chemokines in a variety of assay formats. In addition, the CX3C chemokine, Fractalkine, binds with a very high affinity ($K_I \sim 50$ pM) to US28.

Fractalkine is expressed on certain endothelial cell surfaces and on populations of
15 dendritic cells (DC), and may thus define a portal through which CMV infected cells go from the circulation to the tissue space, as well as find residence in the DC.

[0007] Since the US28 receptor is expressed on cytomegalovirus infected cells, and also in view of its ability to bind multiple chemokines, a small molecule inhibitor for this receptor would have significant use as an anti-CMV agent.

20

SUMMARY OF THE INVENTION

[0008] In one aspect, the present invention provides methods for treating or preventing viral dissemination from CMV infection. The methods typically involve
25 administering to a patient an effective formulation of one or more of the compounds of formula I:



wherein Ar is a substituted or unsubstituted 5-14 membered heteroaryl group having from 1 to 5 heteroatoms as ring members; R^1 is selected from the group consisting of
30 substituted or unsubstituted aryl(C_1 - C_4)alkyl, heteroaryl(C_1 - C_4)alkyl, $-C(O)R^{11}$, and $-C(O)NR^{11}R^{12}$, wherein each R^{11} and R^{12} independently is substituted or unsubstituted

aryl, substituted or unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl; R² is H or (C₁-C₈)alkyl; and Z^N is a substituted or
5 unsubstituted hetero(C₆-C₁₀)bicycloalkyl group.

[0009] Additionally, the invention provides compositions of the compounds above in combination with a pharmaceutically acceptable carrier or excipient.

BRIEF DESCRIPTION OF THE DRAWINGS

10

[0010] None

DETAILED DESCRIPTION OF THE INVENTION

15 Abbreviations and Definitions

[0011] The term "alkyl," by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or combination thereof, and can include di- and multivalent radicals, having the number of
20 carbon atoms designated (*i.e.* C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, cyclohexyl, (cyclohexyl)methyl, cyclopropylmethyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An "unsaturated alkyl" group (*e.g.*, an alkenyl or alkynyl group) is one having one or more
25 double bonds or triple bonds. Examples of unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers.

[0012] The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by -CH₂CH₂CH₂CH₂-. Typically,
30 an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower alkylene" is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms.

[0013] The terms “alkoxy,” “alkylamino” and “alkylthio” (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively.

[0014] The term “heteroalkyl,” by itself or in combination with another term,
5 means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and from one to three heteroatoms selected from the group consisting of O, N, Si and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N and S may be placed
10 at any interior position of the heteroalkyl group. The heteroatom Si may be placed at any position of the heteroalkyl group, including the position at which the alkyl group is attached to the remainder of the molecule. Examples include -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-
15 CH₃. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. Similarly, the term “heteroalkylene” by itself or as part of another substituent means a divalent radical derived from heteroalkyl, as exemplified by -CH₂-CH₂-S-CH₂CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (*e.g.*, alkyleneoxy,
20 alkylenedioxy, alkyleneamino, alkylenediamino, and the like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking group is implied.

[0015] The terms “cycloalkyl” and “heterocycloalkyl”, by themselves or in combination with other terms, represent, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl”, respectively. Additionally, for heterocycloalkyl, a heteroatom
25 can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like.
30

[0016] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl,” are meant to include monohaloalkyl and

polyhaloalkyl. For example, the term "halo(C₁-C₄)alkyl" is mean to include trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0017] The term "aryl" means, unless otherwise stated, a polyunsaturated, typically aromatic, hydrocarbon substituent which can be a single ring or multiple rings (up to three rings) which are fused together or linked covalently. The term "heteroaryl" refers to aryl groups (or rings) that contain from one to five heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom. Generally, the aryl groups will have from six to fourteen carbon atoms as ring members, while the heteroaryl groups will have from six to fourteen ring members selected from carbon, nitrogen, sulfur and oxygen. Non-limiting examples of aryl and heteroaryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, and 6-quinolyl. When the aryl or heteroaryl groups are provided as being "substituted or unsubstituted" the substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below.

[0018] For brevity, the term "aryl" when used in combination with other terms (*e.g.*, aryloxy, arylthioxy, arylalkyl) includes both aryl and heteroaryl rings as defined above. Thus, the term "arylalkyl" is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, pyridylmethyl and the like) including those alkyl groups in which a carbon atom (*e.g.*, a methylene group) has been replaced by, for example, an oxygen atom (*e.g.*, phenoxymethyl, 2-pyridyloxymethyl, 3-(1-naphthyloxy)propyl, and the like).

[0019] Each of the above terms (*e.g.*, "alkyl," "heteroalkyl," "aryl" and "heteroaryl") are often noted as "substituted or unsubstituted" and will include substituents for each type of radical as provided below.

[0020] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be a variety of groups

selected from: -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''',
 -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-
 C(O)NR''R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -
 S(O)R', -S(O)₂R', -S(O)₂NR'R'', -CN and -NO₂ in a number ranging from zero to
 5 (2m'+1), where m' is the total number of carbon atoms in such radical. R', R'' and R'''
 each independently refer to hydrogen, unsubstituted (C₁-C₈)alkyl and heteroalkyl,
 unsubstituted aryl, aryl substituted with 1-3 halogens, unsubstituted alkyl, alkoxy or
 thioalkoxy groups, or aryl-(C₁-C₄)alkyl groups. When R' and R'' are attached to the same
 nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-
 10 membered ring. For example, -NR'R'' is meant to include 1-pyrrolidinyl and 4-
 morpholinyl. From the above discussion of substituents, one of skill in the art will
 understand that the term "substituted alkyl" is meant to include groups such as haloalkyl
 (e.g., -CF₃ and -CH₂CF₃) and acyl (e.g., -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the
 like).

15 **[0021]** Similarly, substituents for the aryl and heteroaryl groups are varied and are
 selected from: -halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R',
 -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR''R''',
 -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R',
 -S(O)₂NR'R'', -N₃, -CH(Ph)₂, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a
 20 number ranging from zero to the total number of open valences on the aromatic ring
 system; and where R', R'' and R''' are independently selected from hydrogen,
 (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-(C₁-
 C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl.

[0022] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring
 25 may optionally be replaced with a substituent of the formula -T-C(O)-(CH₂)_q-U-, wherein
 T and U are independently -NH-, -O-, -CH₂- or a single bond, and q is an integer of from
 0 to 2. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl
 ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein
 A and B are independently -CH₂-, -O-, -NH-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single
 30 bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so
 formed may optionally be replaced with a double bond. Alternatively, two of the
 substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced
 with a substituent of the formula -(CH₂)_s-X-(CH₂)_t-, where s and t are independently
 integers of from 0 to 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The

substituent R' in -NR'- and -S(O)₂NR'- is selected from hydrogen or unsubstituted (C₁-C₆)alkyl.

[0023] As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

5 [0024] The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient
10 amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of
15 the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively
20 nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge, S.M., et al, "Pharmaceutical Salts", *Journal of*
25 *Pharmaceutical Science*, 1977, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0025] The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner.

30 The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

[0026] In addition to salt forms, the present invention provides compounds which are in a prodrug form. Prodrugs of the compounds described herein are those compounds

that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.

5 [0027] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple
10 crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0028] Certain compounds of the present invention possess asymmetric carbon
15 atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are all intended to be encompassed within the scope of the present invention.

[0029] The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such
20 compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

25 General

[0030] In related co-owned applications Ser. Nos. 09/944,163, 09/944,051 and 09/944,049, all filed August 30, 2001 and incorporated herein by reference, assays for identifying compounds useful for blocking CMV dissemination in a host were provided. Typically, these assays determine whether the compound inhibits the binding of a
30 chemokine to US28 or a US28 fragment, and are run as a competitive binding assay using a labeled chemokine. A variety of chemokines are known to bind to US28 and are useful in such assays. Preferably, the chemokine is fractalkine and the assay is a radioligand binding assay.

[0031] Following methods outlined in the above-noted applications, compounds have now been identified that are effective in blocking CMV dissemination in a host.

Description of the Embodiments

5

A. Methods of treating CMV infection

[0032] In one aspect, the present invention provides novel methods for treating or preventing CMV infection or diseases associated with, or resultant from, CMV infection.

10 Without intending to be bound by theory, it is believed that the compounds and compositions provided below, exert their effect by inhibiting US28-mediated viral dissemination. The methods typically involve administering to a patient an effective formulation of one or more of the subject compounds or compositions described in more detail below (e.g., compounds of formula I).

15 [0033] The invention provides methods of using the compounds and compositions described below to treat disease or provide medicinal prophylaxis to individuals who possess a compromised immune system or are expected to suffer immunosuppressed conditions, such as patients prior to undergoing immunosuppressive therapy in connection with organ transplantation or anticancer chemotherapy. Additional methods
20 are provided for the treatment of CMV-associated diseases (e.g., atherosclerosis or cardiovascular diseases) in non-immunocompromised individuals. These methods generally involve administering to the host an effective amount of the subject compounds or pharmaceutically acceptable compositions.

[0034] The compositions and compounds described herein and the
25 pharmaceutically acceptable salts thereof can be administered in any effective way such as via oral, parenteral or topical routes. Generally, the compounds are administered in dosages ranging from about 2 mg up to about 2,000 mg per day, although variations will necessarily occur depending on the disease target, the patient, and the route of administration. Preferred dosages are administered orally in the range of about 0.05
30 mg/kg to about 20 mg/kg, more preferably in the range of about 0.05 mg/kg to about 2 mg/kg, most preferably in the range of about 0.05 mg/kg to about 0.2 mg per kg of body weight per day.

[0035] Therapeutic and prophylactic methods of this invention comprise the step of treating patients in a pharmaceutically acceptable manner with those compounds or

compositions. Such compositions may be in the form of tablets, capsules, caplets, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid preparations, such as oral or sterile parenteral solutions or suspensions. Compounds of the invention may also be administered via an intraocular implant for treating retinitis as a result of CMV infection. In particular, compounds may be embedded in a polymer based implant which will be release into the eye over an extended period of time.

[0036] Physicians will determine the dosage of the present therapeutic agents which will be most suitable. Dosages may vary with the mode of administration and the particular compound chosen. In addition, the dosage may vary with the particular patient under treatment. The dosage of the compound used in the treatment will vary, depending on viral load, the weight of the patient, the relative efficacy of the compound and the judgment of the treating physician. Such therapy may extend for several weeks or months, in an intermittent or uninterrupted manner.

B. *Compounds which block CMV dissemination*

[0037] Compounds that are useful in blocking CMV dissemination have been identified using the assays described below, and have been found to possess a structure of general formula I:



wherein Ar is a substituted or unsubstituted 5-14 membered heteroaryl group having from 1 to 5 heteroatoms as ring members; R¹ is selected from the group consisting of substituted or unsubstituted aryl(C₁-C₄)alkyl, heteroaryl(C₁-C₄)alkyl, -C(O)R¹¹, and -C(O)NR¹¹R¹², wherein each R¹¹ and R¹² independently is substituted or unsubstituted aryl, substituted or unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl; R² is H or (C₁-C₈)alkyl; and Z^N is a substituted or unsubstituted hetero(C₆-C₁₀)bicycloalkyl group. For each of the terms hetero(C₄-C₈)cycloalkyl and hetero(C₆-C₁₀)bicycloalkyl, the parenthetical referring (e.g., C₄-C₈ or C₆-C₁₀) refers to the number of ring atoms in the cyclo or bicyclo group, regardless of

whether the atoms are carbon atoms or a heteroatom. For example, a C₅ heterocycloalkyl refers to, among others, a pyrrolidinyl group (four carbon atoms and one nitrogen atom).

[0038] In a first group of preferred embodiments, the compounds are represented by formula (I) wherein Ar is a monocyclic or fused bicyclic nitrogen heteroaryl group.


- 5 More preferably, Ar is a substituted or unsubstituted ring selected from pyridine, pyrimidine, quinoline, quinazoline, isoquinoline, benzimidazole and indole. In the most preferred embodiments, Ar is a substituted or unsubstituted quinoline, preferably attached to the remainder of the molecule at the 4-position of the quinoline ring.

- [0039]** The R¹ group of formula (I), as noted above, represents substituted or
10 unsubstituted aryl(C₁-C₄)alkyl, heteroaryl(C₁-C₄)alkyl, -C(O)R¹¹ or -C(O)NR¹¹R¹², wherein each R¹¹ and R¹² independently represents a substituted or unsubstituted aryl, substituted or unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl. In one group of preferred embodiments, R¹ is substituted or
15 unsubstituted aryl(C₁-C₄)alkyl or heteroaryl(C₁-C₄)alkyl. More preferably, substituted or unsubstituted benzyl, phenethyl or pyridylmethyl. Preferred substituents for each of these groups are selected from halogen, NO₂, CN, R, OR, NR₂, CO₂R, C(O)R, OC(O)R, NRC(O)R and NRC(O)NR₂, wherein each R is independently selected from H and (C₁-C₈)alkyl (and including haloalkyl groups such as fluoromethyl, difluoromethyl, trifluoromethyl, chlorodifluoromethyl and the higher homologs thereof). In another group of preferred embodiments, R¹ is -C(O)R¹¹ or -C(O)NR¹¹R¹², wherein each R¹¹ and R¹² independently represents a substituted or unsubstituted aryl, substituted or
20 unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl. Within this group of preferred embodiments, R¹ is a substituted or unsubstituted benzoyl, phenylacetyl, 2-picolinyl, 3-picolinyl, 4-picolinyl, 2-pyridylacetyl, 3-pyridylacetyl and 4-pyridylacetyl. As above, preferred substituents in this group of
25 embodiments are selected from halogen, NO₂, CN, R, OR, NR₂, CO₂R, C(O)R, OC(O)R, NRC(O)R and NRC(O)NR₂, wherein each R is independently selected from H and (C₁-C₈)alkyl and (C₁-C₈)haloalkyl.

- [0040]** Z^N is a substituted or unsubstituted hetero(C₆-C₁₀)bicycloalkyl group. Preferably the hetero(C₆-C₁₀)bicycloalkyl group is selected from 1-


azabicyclo[2.2.2]octane (quinuclidine), 2-azabicyclo[2.2.2]octane, 1-azabicyclo[3.2.2]nonane, 2-azabicyclo[3.2.2]nonane, 1-azabicyclo[2.2.1]heptane and 2-azabicyclo[2.2.1]heptane. When present, substituents are preferably, substituted or unsubstituted (C₁-C₈)alkyl, (C₂-C₈)alkenyl and (C₂-C₈)alkynyl. Particularly preferred substituents are (C₂-C₈)alkenyl groups such as vinyl, allyl and 3-buten-1-yl. In the most preferred embodiments, Z^N is a quinuclidine ring having a vinyl substituent.

[0041] In view of the preferred groups noted above, certain particular compounds stand out as being preferred. Those compounds are represented by the formula:



wherein the subscript n is an integer of from 0 to 3; each R¹³ is independently selected from the group consisting of halogen, NO₂, CN, R, OR, NR₂, CO₂R, C(O)R, OC(O)R, NRC(O)R and NRC(O)NR₂, wherein each R is independently selected from H and (C₁-C₈)alkyl; and R¹⁴ is selected from the group consisting of H and substituted or unsubstituted (C₁-C₈)alkyl. More preferably, R¹⁴ is unsaturated (C₂-C₈)alkyl (e.g., alkenyl). Most preferably, R¹⁴ is vinyl.

[0042] In the most preferred embodiments of this group, the compound is selected from the following:



[0043] Within the groups of compounds above, evaluation of efficacy can be accomplished by assays to determine whether the compound inhibits the binding of a chemokine to US28 or a suitable US28 fragment.

5


Preparation of Compounds of Formula I

[0044] Compounds useful in the present methods and compositions can be prepared using generally accepted synthetic techniques starting with commercially available materials (e.g., from Aldrich Chemical Co., Milwaukee, WI, USA or Fluka Chemical Co.).

10

[0045] Scheme I illustrates the preparation of compounds of formula I:

Scheme I



[0046] According to Scheme I, a bromo-substituted hetero(C₆-C₁₀)bicycloalkane (a) can be converted to the corresponding Grignard derivative (b) according to established methods. The Grignard reagent can then be reacted with an aryl aldehyde or ketone to provide alcohols of formula (c) wherein R² represents a hydrogen atom or a lower alkyl group (e.g., one to eight carbon atoms that does not sterically interfere with the reaction at the carbonyl center). Alkylation or acylation of (c) provides compounds of formula I. Alternatively, certain compounds of formula (c) are commercially available and can be used in the alkylation or acylation step. For example, cincholine and cinchonidine are both commercially available and can be used as described in the examples below.

C. Assays for identifying compounds which block viral dissemination

[0047] Assays are provided herein for identifying a compound capable of blocking CMV dissemination in a host, by determining whether the compound inhibits the binding of a chemokine to US28 or a US28 fragment.

[0048] These assays are typically cell-based assays in which a cell which stably expresses US28 is treated with a candidate compound, or more preferably a compound having formula I above and a chemokine in a competitive binding format. Selection of the most potent compounds can be made according to established practices for receptor-ligand competitive binding studies. A variety of other assay formats are also useful in the

present invention. For example, substrate-bound or support-bound chemokines (or ligands) can be contacted with a labeled cell or liposome having an associated US28 or US28 fragment

[0049] A variety of cell lines can be used in this aspect of the invention. In one group of embodiments, the cell line is a mouse cell line. In other embodiments, the cell line is a human or primate cell line (e.g., human foreskin fibroblasts (HFF), human diploid lung fibroblasts (MRC-5 and WI-38), or HUVECs), 293 and COS-7 cells. The cell lines described are transfected with US28 cDNA, typically under the control of a CMV promoter, using conventional methods. The cells are cultured in a suitable buffer (e.g., IMDM-5% FBS, DMEM 10% FCS, HUVEC complete medium, and the like) then centrifuged and resuspended in assay buffer (e.g., HEPES with NaCl, CaCl₂, MgCl₂, and BSA) to a concentration of from about 5×10^5 to about 5×10^7 , preferably from about 2×10^6 to about 8×10^6 . Aliquots of the cells are then contacted with the candidate compounds and labeled chemokine.

[0050] A variety of chemokines can be used in this aspect of the invention, including, for example, fractalkine, RANTES, MCP-3, MIP-1 α and MCP-1. Preferably, the labeled chemokine is labeled fractalkine. Additionally, a variety of labels can also be used with the chemokines described above. Typically, the label will be a fluorescence label, a phosphorescence label, a radiolabel, a colorimetric label, or the like. In preferred embodiments the labeled chemokine is a radiolabeled fractalkine, more preferably, ¹²⁵I-fractalkine.

[0051] After contacting the cells with one or more candidate compounds in the presence of labeled chemokine, the assay mixture is typically incubated for a period of time of from about 1 to about 6 hours at a temperature of from about 1 to about 10°C. Preferably the mixture is incubated for a period of from about 2 to about 4 hours at a temperature of about 4°C. One of skill in the art will understand that a variety of assay conditions can be employed, depending on the cell line used, the concentrations of the compounds and chemokine and the concentration of the cells themselves.

[0052] Following incubation the assay wells (for those embodiments carried out on 96-, 384-, 1536-well or larger plates) are typically harvested under vacuum using filter plates, pre-soaked with PEI solution. Scintillation fluid (for radiolabel assays) is added, the plates are sealed and the wells are counted.

D. Compositions useful in the treatment of CMV infection

[0053] The present invention also provides compositions useful for preventing CMV dissemination in a host, which comprises a pharmaceutically acceptable carrier or adjuvant and an effective amount of a compound identified using the assays described herein. Preferably, the compound is a compound of formula I.

[0054] Typically, the compositions contain from about 0.1% to about 99% by weight of active compound, and preferably from about 10% to about 60% by weight depending on which method of administration is employed.

[0055] A CMV dissemination-inhibiting amount is that amount of active compound required to slow the progression of viral dissemination or reduce the amount of viral dissemination from that which would otherwise occur without administration of the compound. Or, it is an amount of active compound required to slow the progression or reduce the intensity of symptoms resulting from CMV infection or reactivation or elimination thereof.

[0056] CMV dissemination-inhibiting activity of compounds of the invention can be determined according to the assays described herein. The assays provide an indication of chemokine binding to US28, more typically fractalkine binding to US28. The compounds provided herein inhibit the binding of fractalkine to US28 with activity expressed as IC₅₀ (that amount of compound that reduces fractalkine binding by 50%). The compounds provided herein will typically exhibit an IC₅₀ of approximately 50 μ M or less, preferably 25 μ M or less, more preferably 10 μ M or less, and most preferably less than 1 μ M.

[0057] For the compositions of the invention, the proportion of each carrier, diluent or adjuvant is determined by the solubility and chemical nature of the compound and the route of administration according to standard pharmaceutical practice. In order to obtain consistency of administration, however, it is preferred that a composition of the invention is in the form of a unit dose. For example, the unit dose presentation forms for oral administration may be tablets and capsules and may contain conventional excipients such as binding agents (e.g., acacia, gelatin, sorbitol, or polyvinylpyrrolidone), fillers (e.g., lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine), tableting lubricants (e.g., magnesium stearate), disintegrants (e.g., starch, polyvinylpyrrolidone, sodium starch glycoallate or microcrystalline cellulose), or pharmaceutically acceptable wetting agents (e.g., sodium lauryl sulfate).

[0058] The compounds may be injected parenterally; this being intramuscularly, intravenously, or subcutaneously. For parenteral administration, the compound may be used in the form of sterile solutions containing other solutes, for example, sufficient saline or glucose to make the solution isotonic. The amount of active ingredient administered parenterally will be approximately 0.01 to 250 mg/kg/day, preferably about 1 to 10 mg/kg/day, more preferably about 0.5 to 30 mg/kg/day, and most preferably about 1-20 mg/kg/day.

[0059] The compounds may be administered orally in the form of tablets, capsules, or granules containing suitable excipients such as starch, lactose, white sugar and the like. The compounds may be administered orally in the form of solutions which may contain coloring and/or flavoring agents. The compounds may also be administered sublingually in the form of tracheas or lozenges in which each active ingredient is mixed with sugar or corn syrups, flavoring agents and dyes, and then dehydrated sufficiently to make the mixture suitable for pressing into solid form. The amount of active ingredient administered orally will depend on bioavailability of the specific compound.

[0060] The solid oral compositions may be prepared by conventional methods of blending, filling, tableting, or the like. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art. The tablets may be coated according to methods well known in normal pharmaceutical practice, in particular with an enteric coating.

[0061] Oral liquid preparations may be in the form of emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may or may not contain conventional additives. For example suspending agents, such as sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminum stearate gel, or hydrogenated edible fats; emulsifying agents, such as sorbitan monooleate or acaci; non-aqueous vehicles (which may include edible oils), such as almond oil, fractionated coconut oil, oily esters selected from the group consisting of glycerine, propylene glycol, ethylene glycol, and ethyl alcohol; preservatives, for instance methyl parahydroxybenzoate, ethyl para-hydroxybenzoate, n-propyl parahydroxybenzoate, or n-butyl parahydroxybenzoate of sorbic acid; and, if desired, conventional flavoring or coloring agents.

[0062] The compounds of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

[0063] For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of the present invention are employed. As used herein, topical application is also meant to include the use of mouth washes and gargles.

[0064] In another embodiment, the invention provides the subject compounds in the form of a pro-drug, which can be metabolically or chemically converted to the subject compound by the recipient host. A wide variety of pro-drug derivatives are known in the art such as those that rely on hydrolytic cleavage or oxidative activation of the prodrug.

[0065] The compositions may be advantageously combined and/or used in combination with other antiviral agents which are either therapeutic or prophylactic agents, and different from the subject compounds. The compositions may also be advantageously combined and/or used in combination with agents that treat or induce conditions often associated with the viral infections that are sensitive to the present compounds, such as anti-HIV agents or immunosuppressive agents. In many instances, administration in conjunction with the subject compositions enhances the efficacy of such agents. Exemplary antiviral agents include ganciclovir, foscarnet and cidofovir. Exemplary anti-HIV agents include indinavir, ritonavir, AZT, lamivudine and saquinavir. Exemplary immunosuppressive agents include cyclosporin and FK-506. The compositions may also be advantageously used as antiviral prophylactic treatment in combination with immunosuppressive protocols such as bone-marrow destruction (either by radiation or chemotherapy).

[0066] To further assist in understanding the present invention, the following non-limiting examples are provided.

EXAMPLES

Example 1

[0067] Radioligand binding assays were carried out in the following manner. The target cells typically used in these assays were either Rhesus dermal fibroblasts which had


been infected with Rhesus CMV for 2-4 days, or hUS28 transfected murine cells. Lots of cells verified for fractalkine binding were frozen until use. For the assay, cells were thawed, washed, and resuspended in assay buffer (20 mM HEPES pH 7.1, 140 mM NaCl, 1 mM CaCl₂, 5 mM MgCl₂, and with 0.2% bovine serum albumin) to a concentration of
5 from 4×10^5 to 5×10^6 cells/ml. Compounds were prepared as 10X solutions in 20% DMSO, and 0.02 ml was placed in assay plates. Next 0.09 ml of cells was added to the assay plates containing the compounds. Lastly 0.09 ml of ¹²⁵I-fractalkine diluted in assay buffer (final concentration ~50 pM, with 20,000-50,000 cpm per well) was added, the plates sealed and incubated for 2-4 hours at 4°C on a shaker platform. The assay plates
10 were harvested using Packard GF/B filter plates, pre-soaked in 0.3% polyethyleneimine solution, on a Packard vacuum cell harvester. Scintillation fluid was added to all wells, the plates were sealed and counted in a Top Count scintillation counter. Control wells containing either diluent only (for total counts) or excess unlabeled fractalkine (1 µg/ml, for non-specific binding) were used to calculate the percent of total inhibition. IC₅₀
15 values are those concentrations required to reduce the binding of labeled fractalkine to the receptor by 50%, and were determined by nonlinear regression curve-fitting of the dose-response assays.

Example 2

20

[0068] This example illustrates one method for the preparation of compounds of formula I, from commercially available starting materials.


3-Nitrobenzoic acid cinchonine ester (I-i)



[0069] 3-Nitrobenzoyl chloride (186 mg, 1.0 mmol) was added in one portion to a mixture of cinchonine (294 mg, 1.0 mmol, Fluka) in toluene (3.5 mL) at room temperature then heated to reflux for 24 hr. After cooling, the crude product which had
30 precipitated was collected by filtration and washed three times with cold toluene and air-


dried under ambient conditions to provide the desired product as a HCl salt (427 mg, about 89% as a white solid).

[0070] 4-Methylbenzoic acid cinchonidine ester (I-ii)



[0071] Cinchonidine (1.00 g, 3.40 mmol) was suspended in dry toluene (30.0 ml) containing triethylamine (0.68 g, 6.72 mmol). 4-Methylbenzoyl chloride (0.58 g, 3.75 mmol) was added dropwise to the above suspension with efficient stirring for about 5 min (warmed spontaneously to about 30-40°C) and the stirring was continued for 15 hrs at room temperature. The mixture was diluted with dichloromethane (~100 ml), washed with 5% aq. Na₂CO₃ solution and distilled water, respectively. The organic layer collected was dried over MgSO₄ and evaporated to dryness under reduced pressure. The obtained crude product was purified by column chromatography. Yield: 1.20 g (86%) solid (R_f = 0.59 (CHCl₃-methanol 10:1). The structural identity and the purity of the product was determined by ¹H-NMR (approved, 98%) and HPLC-MS spectroscopy (100% with correct MS signal)).


[0072] 2-Chlorobenzoic acid cinchonidine ester (I-iii)



[0073] Cinchonidine (1.00 g, 3.40 mmol) was suspended in dry toluene (30.0 ml) containing triethylamine (0.68 g, 6.72 mmol). 2-Chlorobenzoyl chloride (0.65 g, 3.71 mmol) was added dropwise to the above suspension with efficient stirring for about 5 min (the mixture spontaneously warmed up to about 30-40°C) and the stirring was continued


for 15 hrs at room temperature. The mixture was diluted with dichloromethane (~100 ml), washed with 5% aq. Na_2CO_3 solution and pure water, respectively. The organic layer collected was dried over MgSO_4 and evaporated to dryness under reduced pressure. The obtained crude product was purified by column chromatography. Yield: 0.93 g (63%) solid ($R_f = 0.56$ (CHCl_3 -methanol 10:1)). The structural identity and the purity of the product was determined by $^1\text{H-NMR}$ (approved, 98%) and HPLC-MS spectroscopy (97% with correct MS signal).

[0074] 2-Chloro-4-nitrobenzoic acid cinchonidine ester (**I-iv**)




[0075] Cinchonidine (1.00 g, 3.40 mmol) was suspended in dry toluene (30.0 ml) containing triethylamine (0.68 g, 6.72 mmol). 2-Chloro-4-nitrobenzoyl chloride (0.82 g, 3.73 mmol) was added dropwise to the above suspension with efficient stirring for about 5 min (the mixture spontaneously warmed up to about 30-40°C) and the stirring was continued for 15 hrs at room temperature. The mixture was diluted with dichloromethane (~100 ml), washed with 5% aq. Na_2CO_3 solution and distilled water, respectively. The organic layer collected was dried over MgSO_4 and evaporated to dryness under reduced pressure. The obtained crude product was purified by column chromatography. Yield: 1.25 g (77%) solid ($R_f = 0.62$ (CHCl_3 -methanol 10:1)). The structural identity and the purity of the product was determined by $^1\text{H-NMR}$ (approved, 97%) and HPLC-MS spectroscopy (98% with correct MS signal).

[0076] Phenylacetic acid cinchonidine ester (**I-v**)




[0077] Cinchonidine (1.00 g, 3.40 mmol) was suspended in dry toluene (30.0 ml) containing triethylamine (0.68 g, 6.72 mmol). Phenylacetyl chloride (1.16 g, 7.50 mmol) was added dropwise to the above suspension with efficient stirring for about 5 min (the mixture spontaneously warmed up to about 30-40°C) and the stirring was continued for 5 15 hrs at room temperature. The solvent was removed in vacuum, the residue was dissolved in 3% aq. HCl solution (50 ml) and washed twice with diethylether (2x50 ml). The aqueous phase collected was made basic by the addition of 5 N aq. NaOH solution to pH ~14 and extracted 3 times with dichloromethane (3x50 ml). The combined organic layers were dried over MgSO₄ and evaporated to dryness under reduced pressure. The 10 obtained crude product was purified by column chromatography. Yield: 0.53 g (38%) oil (R_f = 0.57 (CHCl₃-methanol 10:1)). The structural identity and the purity of the product was determined by ¹H-NMR (approved, pure) and HPLC-MS spectroscopy (86% at 220 nm with correct MS signal).

15 [0078] Phenylacetic acid cinchonine ester (I-vi)




[0079] Cinchonine (1.00 g, 3.40 mmol) was suspended in dry toluene (30.0 ml) containing triethylamine (0.68 g, 6.72 mmol). Phenylacetyl chloride (0.58 g, 3.75 mmol) was added dropwise to the above suspension with efficient stirring for about 5 min (the mixture spontaneously warmed up to about 30-40°C) and the stirring was continued for 15 hrs at room temperature. The solvent was removed in vacuum, the residue was diluted with 3% aq. HCl solution (50 ml) and washed twice with diethylether (2x50 ml). The aqueous phase collected was made basic by the addition of 5 N aq. NaOH solution to pH 20 ~14 and extracted 3 times with dichloromethane (3x50 ml). The combined organic layers were dried over MgSO₄ and evaporated to dryness under reduced pressure. The crude product was purified by column chromatography. Yield: 0.35 g (25%) oil (R_f = 0.56 (CHCl₃-methanol 10:1)). The structural identity and the purity of the product was 25 determined by ¹H-NMR (approved, 98%) and HPLC-MS spectroscopy (100% with 30 correct MS signal).

[0080] 3-Cyclopentylpropionic acid cinchonidine ester (**I-vii**)



- 5 **[0081]** Cinchonidine (1.00 g, 3.40 mmol) was dissolved in dry pyridine (30.0 ml). 3-Cyclopentylpropionyl chloride (0.60 g, 3.73 mmol) was added dropwise to it and the mixture was stirred at room temperature for 15 hrs. The solvent was removed in vacuum. The residue was diluted with 3% aq. HCl solution (50 ml) and washed twice with diethylether (2x50 ml). The aqueous phase collected was made basic by the addition of 5
- 10 N aq. NaOH solution to pH ~14 and extracted 3 times with dichloromethane (3x50 ml). The combined organic layers were dried over MgSO₄ and evaporated to dryness under reduced pressure. The obtained crude product was purified by column chromatography (Merck Kieselgel 60, 0.063-0.2 mm, eluent mixture: CHCl₃-methanol 20:1). Yield: 0.52 g (37%) oil (R_f = 0.65 (CHCl₃-methanol 10:1)). The structural identity and the purity of
- 15 the product was determined by ¹H-NMR (approved, 97%) and HPLC-MS spectroscopy (100% with correct MS signal).

[0082] 2-Thiopheneacetic acid cinchonine ester (**I-viii**)



20

- [0083]** Cinchonine (1.00 g, 3.40 mmol) was dissolved in dry pyridine (30.0 ml). 2-Thiopheneacetyl chloride (0.60 g, 3.74 mmol) was added dropwise to it and the mixture was stirred at room temperature for 15 hrs. The solvent was removed in vacuum. The residue was diluted with 3% aq. HCl solution (50 ml) and washed twice with diethylether
- 25 (2x50 ml). The aqueous phase collected was made basic by the addition of 5 N aq. NaOH solution to pH ~14 and extracted 3 times with dichloromethane (3x50 ml). The combined





organic layers were dried over MgSO_4 and evaporated to dryness under reduced pressure. The obtained crude product was purified by column chromatography. Yield: 0.13 g (9.1%) oil ($R_f = 0.58$ (CHCl_3 -methanol 10:1)). The structural identity and the purity of the product was determined by $^1\text{H-NMR}$ (approved, 91%) and HPLC-MS spectroscopy (93% with correct MS signal).













[0084] The above method can be readily carried out using other substituted benzoyl chlorides (e.g., 4-toluoyl chloride, 2-chlorobenzoyl chloride, 2-chloro-4-nitrobenzoyl chloride, 3,4-dichlorobenzoyl chloride and the like), substituted or unsubstituted phenylacetyl chlorides (e.g., phenylacetyl chloride, 4-fluorophenylacetyl chloride, 4-methoxyphenylacetyl chloride, and the like), substituted or unsubstituted nicotinoyl chlorides (e.g., isonicotinoyl chloride, nicotinoyl chloride, 6-chloronicotinoyl chloride, and the like), and substituted or unsubstituted cycloalkyl(C_1 - C_6)alkanoyl chlorides (e.g., 3-cyclopentylpropionyl chloride).













[0085] Additionally, other cinchonine or quinine derivatives are well known in the art and can be used for starting materials. See for example, U.S. Patent Nos. 4,818,441 and 3,953,453.













[0086] Following the general methods outlined above and substituting commercially-available or literature reagents, the following compounds were prepared.











Table 1

Number	Structure	Number	Structure
101		136	
102		138	







103		139	
104		140	
105		141	
106		142	
107		171	
108		145	

109		146	
110		147	
111		148	
112		149	
113		150	
114		151	

115		152	
116		153	
117		154	
118		155	
119		156	
120		157	

121		158	
122		159	
123		160	
124		161	
125		162	

126		163	
127		164	
128		165	
129		166	
130		167	
131		168	

132		169	
133		170	
134		135	

Example 3

- 5 [0087] The table below illustrates the activity associated with various compounds of formula I.

Compound	Rhesus CMV IC ₅₀ (μM)	hUS28 IC ₅₀ (μM)
I-i	+	+++
I-ii	++	+++
I-iii	+++	++
I-iv	+++	++
I-v	+++	++
I-vi	++	+++
I-vii	++	++
I-viii	++	+++
101	+	+++

102	++	+++
103	+	+++
104	++	+++
105	++	+++
106	+	+++
107	++	+++
108	++	+++
109	++	+++
110	+++	+++
111	++	+++
112	+	+++
113	++	+++
114	++	+++
115	+	++
116	++	++
117	++	++
118	+	++
119	+++	++
120	++	++
121	++	++
122	++	++
123	+	++
124	+	++
125	++	++
126	++	++
127	+++	++
128	+++	++
129	++	++
130	++	++
131	+	++
132	++	++
133	+++	++

134	+	++
135	++	++
136	++	++
137	+++	++
138	+++	++
139	++	++
140	++	++
141	+++	++
142	++	++
145	++	++
146	++	++
147	++	++
148	+	++
149	+	++
150	++	++
151	+	++
152	++	++
153	++	++
154	++	++
155	+	++
156	++	++
157	++	++
158	+	++
159	+	++
160	+	++
161	+++	++
162	+	++
163	+++	++
164	++	++
165	+	++
166	+++	++
167	+	++

168	+	+
169	+	+
170	++	+
171	+	+

Rh CMV: +, 10-40 micromolar; ++, 1-10 micromolar; +++, less than 1 micromolar
hUS28: +, 10-40 micromolar; ++, 1-10 micromolar; +++, less than 1 micromolar

[0088] It is understood that the examples and embodiments described herein are
5 for illustrative purposes only and that various modifications or changes in light thereof
will be suggested to persons skilled in the art and are to be included within the spirit and
purview of this application and scope of the appended claims. All publications, patents,
and patent applications cited herein are hereby incorporated by reference for all purposes.

WHAT IS CLAIMED IS:

- 1 1. A method for treating CMV or CMV-related diseases in a host,
2 comprising administering to said host an effective amount of a compound of formula I:



3
4 wherein

- 5 Ar is a substituted or unsubstituted 5-14 membered heteroaryl group having from
6 1 to 5 heteroatoms as ring members;
7 R¹ is selected from the group consisting of substituted or unsubstituted aryl(C₁-
8 C₄)alkyl, heteroaryl(C₁-C₄)alkyl, -C(O)R¹¹, and -C(O)NR¹¹R¹², wherein
9 each R¹¹ and R¹² independently is substituted or unsubstituted aryl,
10 substituted or unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted
11 (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl,
12 substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or
13 unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl;
14 R² is H or (C₁-C₈)alkyl; and
15 Z^N is a substituted or unsubstituted hetero(C₆-C₁₀)bicycloalkyl group.


- 1 2. A method in accordance with claim 1, wherein Ar is selected from
2 the group consisting of quinoline, isoquinoline, pyridine and pyrimidine.

- 1 3. A method in accordance with claim 1, wherein Z^N is a substituted
2 or unsubstituted hetero(C₆-C₁₀)bicycloalkyl group selected from the group consisting of
3 1-azabicyclo[2.2.2]octane, 2-azabicyclo[2.2.2]octane, 1-azabicyclo[3.2.2]nonane, 2-
4 azabicyclo[3.2.2]nonane, 1-azabicyclo[2.2.1]heptane and 2-azabicyclo[2.2.1]heptane.

- 1 4. A method in accordance with claim 1, wherein R¹ is selected from
2 the group consisting of -C(O)R¹¹ and -C(O)NR¹¹R¹².

- 1 5. A method in accordance with claim 1, wherein R² is hydrogen and
2 R¹ is selected from the group consisting of -C(O)R¹¹ and -C(O)NR¹¹R¹².

- 1 6. A method in accordance with claim 1, wherein said compound has
2 the formula:




- 3
4 wherein the subscript n is an integer of from 0 to 3; each R¹³ is independently selected
5 from the group consisting of halogen, NO₂, CN, R, OR, NR₂, CO₂R, C(O)R, OC(O)R,
6 NRC(O)R and NRC(O)NR₂, wherein each R is independently selected from H and (C₁-
7 C₈)alkyl; and R¹⁴ is selected from the group consisting of H and substituted or
8 unsubstituted (C₁-C₈)alkyl.

- 1 7. A method in accordance with claim 6, wherein R¹⁴ is unsaturated
2 (C₂-C₈)alkyl.

- 1 8. A method in accordance with claim 7, wherein R¹⁴ is vinyl.

- 1 9. A method in accordance with claim 1, wherein said compound is
2 selected from the group consisting of:




3

1 **10.** A method in accordance with claim **1**, wherein said compound
2 inhibits chemokine binding to US28 on the surface of an infected cell or virion.

1 **11.** A method in accordance with claim **10**, wherein said chemokine is
2 fractalkine.

1 **12.** A pharmaceutical composition comprising a pharmaceutically
2 acceptable carrier and a compound of formula I:



3

(I)

4 wherein

5 Ar is a substituted or unsubstituted 5-14 membered heteroaryl group having from
6 1 to 5 heteroatoms as ring members;

7 R¹ is selected from the group consisting of substituted or unsubstituted aryl(C₁-
8 C₄)alkyl, heteroaryl(C₁-C₄)alkyl, -C(O)R¹¹, and -C(O)NR¹¹R¹², wherein
9 each R¹¹ and R¹² independently is substituted or unsubstituted aryl,
10 substituted or unsubstituted aryl(C₁-C₄)alkyl, substituted or unsubstituted

- 11 (C₄-C₈)cycloalkyl(C₁-C₄)alkyl, substituted or unsubstituted heteroaryl,
12 substituted or unsubstituted heteroaryl(C₁-C₄)alkyl and substituted or
13 unsubstituted hetero(C₄-C₈)cycloalkyl(C₁-C₄)alkyl;
14 R² is H or (C₁-C₈)alkyl;
15 Z^N is a substituted or unsubstituted hetero(C₆-C₁₀)bicycloalkyl group.