




(86) Date de dépôt PCT/PCT Filing Date: 2002/11/29
(87) Date publication PCT/PCT Publication Date: 2003/06/05
(45) Date de délivrance/Issue Date: 2012/05/01
(85) Entrée phase nationale/National Entry: 2004/05/27
(86) N° demande PCT/PCT Application No.: CA 2002/001829
(87) N° publication PCT/PCT Publication No.: 2003/046560
(30) Priorité/Priority: 2001/11/30 (US60/334,098)

(51) Cl.Int./Int.Cl. *C07K 19/00* (2006.01),
A61K 39/395 (2006.01), *C07K 14/245* (2006.01),
C07K 14/705 (2006.01), *C07K 16/00* (2006.01),
C07K 16/18 (2006.01), *C07K 16/26* (2006.01),
C07K 16/46 (2006.01), *G01N 33/53* (2006.01),
A61K 38/00 (2006.01)
(72) Inventeurs/Inventors:
MACKENZIE, C. ROGER, CA;
ZHANG, JIANBING, CA
(73) Propriétaire/Owner:
NATIONAL RESEARCH COUNCIL OF CANADA, CA
(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : COMPLEXES DE LIAISONS MULTIMERES D'AUTO-ASSEMBLAGE DERIVES DE MEMBRES DE LA
FAMILLE DES TOXINES AB5
(54) Title: SELF-ASSEMBLING MULTIMERIC BINDING COMPLEXES DERIVED FROM AB5 TOXIN FAMILY MEMBERS



<i>ompA</i>	signal sequence	MKKTAIAIAVALAGFATVAQA	seq. ID 2
1	linker 1	DVQLQ	seq. ID 3
VTB	verotoxin B subunit	EMBL M16625: nt 1349 - 1555	
2	linker 2	GGGGS	seq. ID 6
PTH50	single domain antibody PTH50	GenBank AF447918	
3	linker 3	GS	seq. ID 7
<i>myc</i>	c-myc detection tag	EQKLISEEDLN	seq. ID 8
H	His ₅ purification tag	HHHHH	seq. ID 9

(57) Abrégé/Abstract:

There is provided a method of forming a multimeric complex having affinity for a target. The method comprises: obtaining a plurality of self-assembly molecules, said self-assembly molecules including complementary self-assembly units such as verotoxin subunit B, each of which is operatively connected to an interaction domain such as a single domain antibody specific for the target; and combining said self-assembly molecules such that at least three said self-assembly units simultaneously bind to one another so as to permit the single domain antibodies to bind the target.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 03/046560 A3

(51) International Patent Classification⁷: **G01N 33/53**,
C07K 19/00

(21) International Application Number: PCT/CA02/01829

(22) International Filing Date:
29 November 2002 (29.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/334,098 30 November 2001 (30.11.2001) US

(71) Applicant (*for all designated States except US*): **NATIONAL RESEARCH COUNCIL OF CANADA** [CA/CA]; Montreal Road, Bldg. M58, Room EG12, Intellectual Property Services, Ottawa, Ontario K1A 0R6 (CA).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MACKENZIE, Roger, C.** [CA/CA]; 2195 Hamelin Crescent, Ottawa, Ontario K1J 6K9 (CA). **ZHANG, Jianbing** [CN/CA]; 5963 Pineglade Crescent, Ottawa, Ontario K1W 1G7 (CA).

(74) Agent: **MCKAY, Margaret**; National Research Council of Canada, Building M-58. Rm EG-06B, 1200 Montreal Road, Ottawa, Ontario K1A 0R6 (CA).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- with amended claims

(88) Date of publication of the international search report:
4 September 2003

[Continued on next page]

(54) Title: SELF-ASSEMBLY MOLECULES



<i>ompA</i>	signal sequence
1	linker 1
VTB	verotoxin B subunit
2	linker 2
PTH50	single domain antibody PTH50
3	linker 3
<i>myc</i>	c-myc detection tag
H	His ₅ purification tag

MKKTAIAIAVALAGFATVAQA	seq. ID 2
DVQLQ	seq. ID 3
EMBL M16625: nt 1349 - 1555	
GGGGS	seq. ID 6
GenBank AF447918	
GS	seq. ID 7
EQKLISEEDLN	seq. ID 8
HHHHH	seq. ID 9

(57) Abstract: There is provided a method of forming a multimeric complex having affinity for a target. The method comprises: obtaining a plurality of self-assembly molecules, said self-assembly molecules including complementary self-assembly units such as verotoxin subunit B, each of which is operatively connected to an interaction domain such as a single domain antibody specific for the target; and combining said self-assembly molecules such that at least three said self-assembly units simultaneously bind to one another so as to permit the single domain antibodies to bind the target.



WO 03/046560 A3

WO 03/046560 A3



Date of publication of the amended claims: 16 October 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE OF THE INVENTION

SELF-ASSEMBLING MULTIMERIC BINDING COMPLEXES
DERIVED FROM AB5 TOXIN FAMILY MEMBERS

5

FIELD OF THE INVENTION

The invention relates to self-assembly molecules, and particularly self-assembly molecules useful to form complexes having an affinity to targets of interest.

10

BACKGROUND OF THE INVENTION

A variety of molecules having affinity for particular binding sites in cells are known. For example, antibodies and cell-surface receptors can bind to
15 specific targets. However, the strength of the interaction between a given molecule and its target will in some instances be low.

Efforts to increase the affinity of antibody fragments to their target have resulted in the production of dimerized antibody fragments, wherein the dimer has two binding sites, each of which is specific for the antibody target.
20 Dimerization has been conducted by a variety of means, generally involving modification of a "tail" region attached to the antibody fragment. Thus, self-associating secondary structures such as helix bundles have been employed in an effort to produce dimerizable units which retain their specificity and ability to bind to a target of interest. However, the use of known self-associating domains
25 can present several problems, including unwanted and non-functional aggregation of the molecules, as well as difficulties in obtaining optimal spacing between molecules, resulting from limits on control over the geometry of the resultant structure. It is often desirable that the binding region of each molecule in a dimer is located on the same face of the dimer, with sufficient spacing
30 between the molecules to allow engagement of their target molecules.

Successful efforts at forming oligomers of more than two self-associating subunits in association with specific-binding regions have been limited. In one instance, a tetramer of subunits comprising a modified helix of the transcription factor GCN4 together with a "miniantibody" was produced. Similarly, the coiled-
35 coil

- 2 -

assembly domain of the cartilage oligomeric matrix protein fused to a small peptide has been used to form pentamers. However, the structure of the cartilage oligomeric assembly is believed to be thin and rod-shaped which may render it unsuitable for use with larger peptides or proteins for which greater inter-unit spacing may be needed.

5 A discussion of efforts to produce self-assembling molecules can be found in Pluckthun *et al.*, 1997, ref. 1, as well as Terskikh *et al.*, 1997 ref.2.

SUMMARY OF THE INVENTION

Avidity, the dramatic contribution of multivalency to the strength of biomolecular interactions, is a key feature of antigen-antibody interactions. There is provided herein
10 a method for greatly improving the binding properties of interaction domains such as single domain antibodies (dAbs) through the introduction of avidity. The interaction domains are fused to a suitable self-assembly unit such as the B-subunit of *Escherichia coli* verotoxin ("VTB"). VTB self-assembles to form a homopentamer. When VTB is fused to an interaction domain, the resultant molecules tend to
15 pentamerize. Such pentamerized molecules are referred to as pentabodies.

Introduction of avidity is a very good way of improving the antigen binding properties of antibody fragments. Multimerization is a particularly appealing strategy for the improvement of dAb antigen binding properties.

In an embodiment of the invention there is provided a method of forming a self-
20 assembling multimeric complex, said method comprising obtaining suitable self-assembly molecules, at least three of the self-assembly molecules comprising a complementary self-assembly unit operatively connected to an interaction domain; and combining the self-assembly molecules such that at least three self-assembly units simultaneously bind to one another to form the complex.

25 In another embodiment of the invention there is provided self-assembly molecules, each comprising a complementary self-assembly unit operatively connected to an interaction domain. More particularly, the present invention provides a self-assembly molecule comprising a plurality of proteinaceous or peptide portions

- 3 -

operatively connected together, a first said portion comprising a self-assembly region capable of binding to the self-assembly regions of at least two other self-assembly molecules of the same or different composition, to form a complex of at least three self-assembly molecules, and a second said portion comprising an
5 interaction domain adapted to specifically interact with a target region on another, different molecule.

BRIEF DESCRIPTION OF THE DRAWINGS

10 These and other advantages of the invention will become apparent upon reading the following detailed description and upon referring to the drawings in which: FIGURE 1 is a depiction of the believed primary structure of an embodiment of the VTB-dAb fusion protein.

FIGURE 2 is a ribbon-type depiction of the believed structure of an
15 embodiment of the VTB oligomerization domains in complex with oligosaccharide and with fused dAbs. The saccharide binding face is shown in Figure 2A, the side view is Figure 2B, and the believed modeled structure of the embodiment of the PTH50 pentabody is Figure 2C.

FIGURE 3 is a graphical description of the Superdex TM200
20 chromatography and SDS-PAGE results for an embodiment of PTH50-VTB (Figure 3A), and PTH50 (Figure 3B). PTH50-VTB fusion product is hereinafter referred to as 1 V5, for simplicity.

FIGURE 4 is a graphical depiction of the BIACORETM analysis of the
oligosaccharide binding properties of an embodiment of 2nM VTB and an
25 embodiment of 2nM 1 V5.

FIGURE 5 is a graphical depiction of the BIACORE analysis of the antigen
binding properties of PTH50 and an embodiment of 1V5. Figure 5A relates to
binding of 0.25-2 μ M PTH50 to immobilized peptide antigen; Figure 5B relates
to the binding of 2-10 μ M peptide to immobilized 1V5 ; and Figure 5C relates to
30 the binding of 2.5-15nM 1V5 to immobilized peptide.

FIGURE 6 is a depiction of embodiments of certain sequence listings of
interest.

- 4 -

FIGURE 7 is the heat induced denaturation curve for pentabody 1V5 – see Example 1, thermal stability studies.

FIGURE 8 shows protease susceptibility of pentameric sdAB – see Example 1, protease stability studies.

5 FIGURE 9 is a diagrammatic presentation of the primary structure of pentabody 1V12, according to Example 2.

FIGURE 10 is the result of size exclusion chromatography measurements on iV12.

10 FIGURE 11 is the result of surface plasmon resonance analysis of 1V12 binding.

FIGURE 12 is a diagrammatic presentation of the products of Example 3 herein.

FIGURE 13 shows the construction and size exclusion chromatography of decavalent single domain antibody 1V13, Example 3 herein.

15 FIGURE 14 is a comparison by size exclusion chromatography of 1V13 and its variations, Example 4 herein.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

20 As used herein the terms “self-assembly units” and “self-assembling units” refer to peptides or proteins (which may be in association with carbohydrate or other modifications), which are adapted to interact with other such regions on separate molecules to form one or more structures having a substantially defined geometry and including three or more units.

25 As used herein the term “interaction domains” refer to peptides, or proteins (which may be glycosylated or otherwise modified), which are adapted to specifically interact with target regions (or targets) on other molecules differing from themselves. Preferably the other molecules are not self-assembly units.

- 5 -

As used herein the term "homopentamer" refers to a structure containing five units specifically interacting with one another to form a structure having substantially defined geometry, each unit being substantially identical to the others.

As used herein the term "multimerization" refers to the process by which more than two self-assembling units combine together to form a larger complex of defined geometry.

As used herein "sdAb, and "dAb" refer to single domain antibody fragments.

As used herein the term "complementary self-assembly units" means at least three self-assembly units which are adapted to associate with one another to form a complex having defined geometry.

As used herein the phrase "a structure having substantially defined geometry" means a structure the approximate size and shape of which is consistent when it is formed from the same components under the same conditions.

As used herein the term "derived from genetic material encoding" refers to something which includes a peptide or protein which could have been substantially produced by transcription of DNA and/or translation of RNA encoding that peptide or protein, or a larger protein of which it forms a part, followed if necessary by cleavage (natural or unnatural) and/or post-translational modification. It will be apparent that a peptide or protein will be derived from genetic material even if the actual genetic material encoding it differs through degeneracy in the genetic code or conservative substitution or the like.

Self-Assembly Molecules

The invention provides novel self-assembly molecules (also called "self-assembling molecules" or "subunits") which in one embodiment may comprise a self-assembly unit operatively connected to an interaction domain. Each self-assembly molecule has a portion which under the correct conditions can specifically bind a target and a portion which under the correct conditions can bind to another self-assembly molecule to form a complex of at least three self-assembly molecules. The self-

- 6 -

assembly unit may be connected to the interaction domain by a linker region" (also referred to as a "linking region", "linkers", "linking domain" or "linker domain").

It will be readily understood that the self-assembly molecules of the present invention may in some instances be fusion proteins. Such fusion proteins can be synthesized by a variety of means, including chemical synthesis followed by chemical
5 modification as needed, synthesis in bacterial or other culture, or synthesis in a mammalian host, for example through the administration of nucleotides encoding the fusion protein of interest to a mammalian host by gene therapy according to standard techniques. The self-assembly molecules of the present invention can also be
10 produced by any suitable means, including chemical linking of the interaction domain to a region of the self-assembly unit so as to preserve target binding and specificity and self-assembly.

Self-Assembly Unit

The self-assembling unit is defined above. In some instances it will be a protein
15 adapted to form an oligomer of three or more units with similar proteins, such that the resulting structure has a defined geometry.

In some instances it will be desirable to select self-assembling units which assemble so as to substantially orient either their N-terminus or their C-terminus on a single face of the structure. In some instances it will be desirable to select self-
20 assembly units which assemble to substantially align their N-termini on a first face and their C-termini on a second face. In some instances it will be desirable to use self-assembly units which assemble to substantially align a mixture of N-terminal regions and C-terminal regions on a single face of the molecule.

In some instances it will be desirable to use self-assembly units which are not
25 identical to one another. For example, each subunit of the final assembly need not be identical, so long as the binding region of each self-assembling unit is complementary to others, such that three or more self-assembling units can form a complex having a defined geometry.

- 7 -

In some instances it will be desirable to use self-assembling units selected from the self-assembly regions of members of the AB5 toxin family, including but not limited to verotoxin, shiga toxin, heat labile enterotoxin, cholera toxin, and pertussis toxin. In some instances it will be desirable to employ self-assembly units which are, or are derived from, proteins naturally encoded by the same family of organisms as is being used to express the subunits.

In some instances it will be desirable to select self-assembly units which will provide self-assembly molecules which form multimeric structures of 4 to 100 self-assembly molecules. In some instances self-assembly units providing multimeric structures of 5 to 50 self-assembly molecules will be desired, and in some instances, selection to provide multimeric structures of 5 to 15 self-assembly molecules will be desired.

In general, multimeric complexes formed from larger numbers of self-assembly molecules will tend to bind target more strongly than complexes of fewer molecules. However, multimeric complexes which are very large may be less able to pass through matrices or tissues and may be less likely to be taken up by cells.

In some instances it will be desirable to use self-assembling units having a constant of dissociation (" K_D " or KD ") in the sub-nanomolar range. In some instances it will be desirable to use self-assembling units having a constant of dissociation in the picomolar range.

In some instances it will be desirable to select self-assembling units to provide self-assembly molecules wherein K_D of the interaction of the oligomerization domains is between 10 μ M and 1fM. In some instances a K_D of between 100nM and 100fM will be desirable. In some instances a K_D of between 100nM and 1pM will be desirable.

In light of the disclosure herein, it is within the capacity of one skilled in the art to determine a suitable K_D and select a suitable self-assembly unit based on the intended application and concentration to be used.

Some self-assembly units, such as verotoxin B-subunit will have a native tendency to bind to a particular cell-surface marker or other target. Where binding to this native target is not desired, it may be disrupted by conventional means, such as

- 8 -

mutation of the target binding site while preserving affinity for other self-assembly units.

In one embodiment, the invention provides self-assembly units suitable for use with rather bulky interaction domains, and particularly interaction domains comprising 15 or more amino acids. In another embodiment there is provided self-assembly units suitable for use with interaction domains having 25-500 amino acids. In another embodiment there is provided self-assembly units suitable for use with interaction domains having 50-300 amino acids. In another embodiment there is provided self-assembly units suitable for use with interaction domains having 75-200 amino acids. In another embodiment of the invention there is provided self-assembly units suitable for use with a bulky interaction domain comprising a peptide or protein region which is glycosylated, or has some other modification thereto.

In one embodiment there is provided self-assembly units suitable for use with an interaction domain and a marker. In one embodiment there is provided self-assembly units suitable for use with an interaction domain and a destructive material.

In many instances it will be desired to use self-assembly units which are the smallest which can be used without over-crowding the interaction domains. Over-crowding may result if excessively small self-assembly units draw the interaction domains too close together. In general, self-assembly units which form short squat complexes will generally be preferable to those which form tall slim complexes, as the interaction domains will generally be fused to the same region of each self-assembly molecule, and will therefore frequently appear on the same face. Where long slim self-assembly units are used, the limited space available to the interaction domains on each face may cause interference between them which may reduce the availability of these interaction domains for binding with the target, or may result in the interaction domains being too close together to optimally bind target on the surface.

Without limiting the invention to any particular mechanism or binding region, it is believed that the region of VTB responsible for oligomerization is the beta sheet (AA11-15) and the anti-parallel beta sheet (65-68). These two regions appear to associate on adjacent monomers. Additionally, the circle of five alpha-helices (one

- 9 -

from each monomer, AA36-46) appear to play a role in the selective association of these individual self-assembly units with one another. Similar analysis of other self-assembly units, followed by routine testing to confirm the binding site location are possible in light of the present invention, thereby making the identification of significant binding regions possible for a skilled technician.

While this is believed to be the first description of antibody fragment pentamerization, the homopentameric cartilage oligomeric matrix protein has been employed to produce oligomeric peptides termed peptabodies (Terskikh et al., 1997, ref 2). However, with a diameter of approximately 20Å (Efimov et.al., 1994, ref 3), pentamerization of an antibody fragment with this protein would require a long linker which could result in further complications. A 24 amino acid linker was used for peptide oligomerization. In contrast, a molecular mass of 38.5 kDa the VT B-pentamer is similar in size to the 31.3 kDa matrix protein but has a much different geometry with a diameter of approximately 56 Å. This diameter and the peripheral positions of the N- and C-termini are generally preferable as they allow for presentation of five dAbs without a complicated requirement for long linker sequences.

Thus, self-assembly units may be specifically crafted to provide the geometry, diameter, and N-terminal or C-terminal orientation desired, provided that the regions responsible for association between the self-assembly units, and the geometry, hydrophobicity and charge characteristics necessary for association are preserved.

In light of the disclosure herein, it is within the capacity of a person skilled in the art to identify suitable self-assembly units. While a variety of methods are possible, one reasonable approach is to look at the three-dimensional structure formed when three or more self-assembling units associate, to determine if the C- or N-termini are aligned in a desirable manner, and to identify the spacing between such termini. This information can then be compared to the orientation and space requirements of the desired interaction domain (and linker region, marker and/or destructive material if used) to select suitable self-assembly units. In some instances it will be desirable to select self-assembling units which provide good levels of soluble product when produced from *E.coli*.

- 10 -

Interaction Domains

A variety of interaction domains are contemplated, and will be apparent to one skilled in the art in light of the disclosure herein. For example, interaction domains such as antibodies, antibody fragments, single domain antibody fragments ("dAbs" or "sdAbs"), single chain polypeptides encoding the V_H and the V_L region of an antibody ("scFv"), peptides or proteins derived from the binding region of an antibody, portions of cell surface receptors, the binding region of cell surface receptors, molecules or the binding portion thereof, which specifically bind to cell surface receptors, and other molecules having an affinity for a specific target.

In some instances, such single domain antibody fragments may be produced by isolating antibody fragments from a library encoding antibody variable domains. The size of sdAbs may vary, and can be determined by sequencing the DNA which encodes them.

Single domain antibodies will sometimes be preferable to scFvs for oligomerization purposes. Since they are generally about half the size of scFvs, oligomeric forms of sdAbs are generally much smaller than their scFv counterparts. Also, the yields of soluble product in *E. coli* tend to be much higher for sdAbs than for scFvs. More importantly, however, sdAbs generally exist entirely as monomers whereas scFvs often form dimers, trimers etc. in which the V_L of one scFv associates with the V_H of a second scFv and *vice versa*. This property can be exploited by carefully choosing the linker length between the V_L and V_H so as to create quite pure dimeric and trimeric scFvs, termed diabodies and triabodies (Hudson et.al., 1999, ref.4). However, introduction of another layer of oligomerization can also lead to undesired complexity.

It will be apparent in light of the disclosure herein to one skilled in the art that self-assembly molecules recognizing more than one target may also be formed substantially according to the method disclosed herein by linking one interaction domain (such as an sdAb, or the like) to the C-terminal region of a self-assembly unit, and a second interaction domain to the N-terminal region.

- 11 -

When selecting interaction domains, it will frequently be desirable to avoid interaction domains which tend to associate with each other, to the detriment of target binding. Direct binding between interaction domains can not only reduce target binding, but may change the shape of the overall oligomer, and reduce the ability of other interaction domains within the oligomer to bind their target.

The desired constant of dissociation between the interaction domain and its target will depend on several factors, including target abundance, the extent of binding desired, the average and duration of binding desired.

In some instances, an interaction domain will be selected to provide a K_D (for a single, non-oligomerized) self-assembly molecule of between about 1fM and 100 μ M. In some instances a K_D of between about 1nM and 10 μ M will be desired.

In some instances it will be desirable to select an interaction domain and a self-assembly unit to form self-assembly molecules which, in monomer form bind only very weakly to target (e.g. K_D of between about 10 μ M to 10mM, or between about 100 μ M to 1mM), but bind target strongly (e.g. K_D of between about 1fM to 100nM or between about 1pM to 10nM) in oligomeric form. In this way, a linked toxin or radioisotope could be targeted to cells expressing the target at high levels, while largely sparing those expressing it at only low levels. This is useful in situations such as some cancers where the abundance of a cell surface marker and not its mere presence, characterizes the undesired cells. A discussion of the use of a different system to provide selective recognition of cells over-expressing a target can be found in Kaminski *et al.*, 1999, ref.5.

In some instances it may be desirable to select an interaction region or self-assembly unit including a "spoiler". A "spoiler" is anything present on or adjacent or operatively connected to the binding region whereby the binding region is initially unable to bind strongly and is only released from this state upon entry into a cell or tissue type of interest. For example, a change in the accessibility of the binding site could be effected through the action of a tissue-specific phosphatase, such that the dephosphorylated binding site was able to bind effectively whereas in its phosphorylated state it could not.

- 12 -

Similarly, a peptide region which ordinarily blocked a binding site could be selectively cleaved by a tissue-specific protease to enable binding. For example, self-assembly molecules having interaction domains specific for a target found both on prostate cancer cells and on healthy cells in remote tissues could be designed to include a peptide which would ordinarily block the interaction domain binding site for the target on the prostate cancer cell. This peptide could include a sequence sensitive to cleavage by PSA (a prostate-specific protease), such that PSA causes release of the peptide. Thus, the self assembly molecules would not significantly bind target in non-prostate tissue, but would be free to bind cancer cells in the prostate. Where self-assembly molecules include destructive material such as toxins or radio-isotopes, such tissue-specific binding may help to reduce the side effects of cancer therapy.

Linker Region

The connections between a self-assembly unit and an interaction domain may include the use of a "linker region" joining the self-assembly unit and the interaction domain. Linker regions may be selected from any number of peptide sequences or other suitable materials. The length of a linker region will depend on several factors, including the geometry of the self-assembling units, the size of the interaction domain and the size and positioning of a marker or destructive material, if used. It is generally desirable to provide a linker region sufficient to allow the interaction domain operatively connected to each self-assembly unit to orient towards its target, permitting the binding of several interaction domains to their targets when their self-assembly units are engaged to one another.

In some instances it will be desirable to use linkers between four and three hundred amino acids in length, in some instances it will be desirable to use linkers between four amino acids and two hundred amino acids in length, and in some instances it will be desirable to use linkers between five amino acids and twenty amino acids in length. When selecting a linker, it will sometimes be desirable to select a sequence which is resistant to proteases.

- 13 -

In some instances it will be desirable to use linkers of the general formula (GGGGS)_n, where "n" is preferably between about 1 and 50. It will sometimes be preferable to select "n" to be between 1 and 25, sometimes between 1 and 10 and sometimes between 1 and 4. In some instances a C-terminal linker sequence
5 GGGGS and an N-terminal linker sequence GGGGSGGGGS will be desirable.

Linker regions will preferably be selected to allow maximum target accessibility to the binding sites of the interaction domains. Linkers will generally be selected for resistance to proteases where *in vivo* applications are contemplated. However, there may be instances where linkers with sensitivity to a tissue-specific protease may be
10 employed, for example where a dissociation of the interaction domain from the rest of the complex is desired in a particular tissue.

Linkers may also be used to join a marker (such as biotin, or a radioactive or fluorescently labeled moiety or compound) or a destructive material (such as a toxin or radioactive material of sufficient activity) to a self-assembly unit. In one embodiment,
15 the linker may be secured (for example as a fusion protein) to the opposite terminus of the self-assembly unit from the interaction domain.

Size

The total size of individual self-assembly molecules containing self-assembly units and interaction domains, and linkers (and markers and/or destructive materials)
20 where applicable, will sometimes be of concern. In particular, where the passage of the self-assembly molecule through a matrix or through tissues is desired, smaller self-assembly molecules may be preferable.

In some instances it will be desirable to select subunits to provide a multimeric complex having a diameter between 10 and 200Å. In some instances a diameter of
25 between 20 and 180Å will be desirable, in some instances a diameter of between 50 and 150Å will be desirable. In some instances a diameter of between 60 and 100Å will be desirable.

- 14 -

It will be readily apparent that the specific embodiments described in detail herein may be conveniently modified to suit other circumstances. For example, the five amino acid sequence present at the N-terminus of the Fig. 1 construct, present only for cloning convenience, can be removed where desired. The *ompA* sequence is removed during secretion of the mature protein to the *E. coli* periplasm. The His₅ tails will not always be required as an affinity purification method based on the fusion protein's retention of the P^k trisaccharide binding of VTB (Fig. 4) could be applied instead. Given the availability of anti-VTB monoclonal antibodies (see Soltyk et.al., 2002, ref.6), the c-myc detection tags will not always be necessary. (Together these modifications would reduce the size of the pentavalent molecule by 13.5 kDa.)

Variations, modifications and alternative embodiments are specifically contemplated and will be apparent to one skilled in the art in light of the disclosure herein.

Internalization

In some instances it will be desirable to select self-assembly units, interaction domains (and where needed linkers, markers and/or destructive material) to provide self-assembly molecules which are capable of being internalized into cells in their monomer form. In some instances it will be desirable to design self-assembly molecules such that multimeric complexes are readily internalized into cells.

Specific triggers for internalization of bound molecules are known. For example, mechanisms for the internalization of materials bound to certain cell surface receptors are known. Thus, in light of the disclosure herein, it is within the capacity of a technician skilled in the art to produce a subunit suitable for internalization in a particular cell type in monomer or oligomer form. Internalization may be desirable, for example, where the subunits include types of toxins or radioisotopes which preferentially act from within the cell.

- 15 -

Diagnostic Use

The self-assembly molecules and multimeric complexes of the present invention are useful for the diagnosis of conditions, as well as the identification of proteins and other materials of interest in biological material. For example, an
5 interaction domain specific for a tumour antigen may be fused to a suitable self-assembly unit and to a radioactive or chemically recognizable marker subunit to provide self-assembly molecules having diagnostic applications. For example, self-assembly molecules of this type will tend to form multimeric complexes in regions where the antigen is found at high levels on a surface, such as a tumour cell. The
10 presence of the multimeric complex can be determined through identification of the marker and can be used to diagnose the presence of tumour cells either in culture, or within an individual. Similarly, subunits containing interaction domains specific for known pathogens in food, water, or similar materials can be used to assay the safety of samples of these products.

15 The formation of the multimeric complex may improve the avidity of binding to target, thereby improving the sensitivity of these assays, as well as providing for a stronger signal strength, making identification of contaminated products easier.

Therapeutic Use

The self-assembly molecules and multimeric complexes of the present
20 invention may be used in the therapy of a number of conditions in mammalian subjects. Self-assembly molecules having an interaction domain specific for a marker of a cell type of particular interest (or specific for a marker which is more highly expressed on a cell type of particular interest, such that binding of the subunit to the cell type of interest will occur preferentially to binding to other cell types) may be
25 employed for therapeutic use.

Where it is desired to destroy a particular cell type, a toxic or otherwise destructive "payload" may be added to the subunit. For example, where the self-assembly unit is the verotoxin B-subunit, or a variation thereof, it may be desirable to

- 16 -

include the verotoxin A-subunit as well. Verotoxin B-subunit will ordinarily be modified to eliminate or substantially reduce its inherent binding to the P_K antigen. The verotoxin A-subunit is toxic, and when added to a subunit specific for an undesired cell type, can be used to eliminate cells of that type within a patient, or within a culture of patient cells destined for re-administration to the patient.

For example, the VEGF receptor is overexpressed in the vasculature of many tumour types. Thus, the destruction of cells over expressing this receptor could be used to compromise blood flow to tumours, thereby potentially reducing tumour load and improving therapeutic outcome. Similarly, HER-2 is overexpressed in many breast cancers and would be a suitable target in order to reduce the load of breast cancer tumour cells in a subject. Interaction domains specific for particular cell surface targets may be readily identified, once the target itself is known. In many instances an sdAb specific for cell surface targets will be the preferred interaction domain.

In some instances it will be desirable to use multimeric complexes to bring two different cell types into close proximity. For example, it may be desirable to bring a killer cell to an undesired cell type, such as a cancer cell. This may be readily accomplished in light of the present disclosure by forming a multivalent molecule which displays an interaction domain specific for the undesired cell type on one face and an interaction domain specific for the killer cell on the other face. For example, a decavalent molecule displaying an sdAb recognizing the cancer cell on the C-terminal face of VTB and having a second sdAb which recognized the killer cell on the N-terminal face of VTB would allow the killer cell and tumour cell to be brought together, facilitating destruction of the cancer cell.

In light of the disclosure herein and the position of the N-terminus and C-terminus (Fig. 2A and 2B), it is within the capacity of a skilled technician to generate such decavalent structures.

When used with respect to therapeutic treatments and compositions, the term "effective amount" refers to an amount which, when administered to the patient over a two-week period causes a significant reduction in the number or viability of the undesired or over-expressed cell or substance.

- 17 -

In some instances, multivalent-bispecific antibodies will have advantages over bivalent-bispecific antibodies, particularly where there is multivalent antigen presentation, such as on cell surfaces.

The oligomerization strategy described here, particularly when used in conjunction with phage display techniques, provides a relatively rapid means of isolating antibody reagents for use in proteomics. Furthermore, immunotoxins in which interaction domains are fused to a destructive material or toxin such as the highly toxic VTA subunit or incorporate or are bound to a radioisotope can be produced in light of the disclosure herein, and such immunotoxins may provide for wide-ranging therapeutic and diagnostic uses.

In some instances it will be desirable to use non-identical self-assembly molecules. The individual self-assembly units will still preferably specifically recognize one another, enabling the formation of a multimer of predictable geometry and size. However, one or both of the interaction domain or the self-assembly unit will differ between the molecules.

For example, it may be desirable to express several different interaction domains and have them come together in multimeric complexes. This could be accomplished by producing various fusion proteins having the same self-assembly unit and a different interaction domain. Alternatively, and particularly where it is desired to maintain a certain stoichiometry between the different interaction domains, it may be desirable to use different self-assembly units which nonetheless assemble together to form a multimeric complex of known geometry and size. For example, pertussis toxin is a heteropentameric product, containing four different B-subunits, one of which is present in duplicate. Thus, if it was desired to form a pentamer having four different interaction domains, one of which was present in two copies in the pentamer whereas the other were only present in a single copy, this could be accomplished by forming four different fusion proteins corresponding to the four different B-subunits of pertussis toxin, with each subunit being fused (through a suitable linker domain if needed) to a different interaction domain. Such an assembly could be particularly useful in cases where several different target molecules are expressed in very close proximity to one

- 18 -

another, and it is their co-expression, or their close relation to one another which is particularly indicative of the cell type of interest.

Example 1 – Summary

In one embodiment of the invention there is provided a novel procedure to
5 improve the avidity of interaction domains by fusing the antibody fragment to the C-terminus of the verotoxin B-subunit. This generates pentameric sdAbs, termed pentabodies. The pentabody described here bound immobilized antigen 7,000 fold more strongly than the monomeric sdAb. This technology can be easily applied to
10 other sdAbs, single chain variable fragments, as well as other suitable interaction domains. Antigen binding affinities can also be improved by *in vitro* affinity maturation (refs. 7 and 8) although this is a time consuming process that involves the construction of and panning of sub-libraries. In some instances *in vitro* affinity improvement may modify fine-specificity or introduce unwanted cross-reactivity (ref. 9).

In the instance of an sdAb specific for a peptide antigen, pentamerization
15 resulted in a 7,000-fold increase in functional affinity for immobilized antigen. The pentavalent sdAb was expressed in high yield in *Escherichia coli* and displayed excellent physical properties. This technology in conjunction with phage display techniques provides a rapid means of generating novel antigen binding molecules with subnanomolar affinities for immobilized antigen. While phage display frequently offers
20 a more efficient means of isolating monoclonal antibodies than hybridoma technologies (ref. 10), the dissociation constants of antibody fragments isolated by phage techniques are typically in the 10^{-5} to 10^{-7} M⁻¹ range (refs 11 and 12) and may be too low for many applications.

Single domain antibodies are typically based on the variable domains of heavy
25 chain antibodies whose variable domains have excellent physical properties that relate to their natural existence in absence of a light chain partner. One useful source of antibodies is a single domain antibody library, derived from the llama heavy chain antibody repertoire and displayed on phage [Tanha (2001), ref. 13]. This library was

- 19 -

the source of the antibody selected for evaluation of the oligomerization strategy described here.

The verotoxin B-subunit monomer ("VTB") was chosen as a non-limiting example of a self-assembly domain because of its relatively small size and the structure of the homopentamer that it forms by self-assembly. The verotoxin B-pentamer has a doughnut-shaped structure with the N- and C-termini exposed on opposite sides of and at the periphery of the molecule.

Verotoxin, or shiga-like toxin, is an AB₅ toxin in which the A subunit is the toxic entity and the pentameric B-subunit mediates binding to the glycolipid globotriaosylceramide, abbreviated Gb₃, (Gal α 1 $\rightarrow$ 4Gal β 1 $\rightarrow$ 4Glc β 1 $\rightarrow$ ceramide). Native *E.coli* verotoxin subunit B binds to eukaryotic cell membranes via the glycolipid Gb₃ receptors. Verotoxin has several varieties. Specific work reported herein was conducted with VT-1, although the process is believed equally applicable to other varieties, and such are within the scope of the invention.

It is possible to overcome Gb₃ binding by mutation of VTB. For example, the W $\rightarrow$ A mutation of amino acid 34, combined with either a D17E mutation or a A56Y mutation abolish detectable binding to glycolipid (Soltyk, *et al.*, Ref. 6). Thus, it is within the capacity of one skilled in the art in light of the disclosure herein to prepare VTB mutants which do not significantly bind Gb₃.

For purposes of illustrating the full potential of a pentavalent sdAb in terms of binding to an immobilized or cell surface antigen, an sdAb recognizing a relatively small antigen that could be immobilized at quite high density was chosen. Accordingly, an sdAb specific for a modified 31 amino acid peptide sequence from human parathyroid hormone ("PTH50") was isolated from a llama sdAb library (ref. 13) by phage display. The peptide has the sequence ¹SVSEIQLMHNLGKHLNSMER**VE**WLR**K**LLQDV³¹ (SEQ. ID. NO. 1) with a β -lactam bond linking the side chains of ²²E and ²⁶K (shown in bold). The anti-PTH sdAb (PTH50) expressed extremely well in *E. coli* (approximately 200 mg/L) and did not aggregate.

- 20 -

The anti-PTH sdAb was fused to VT B-subunit monomer as diagrammed in Fig. 1. The dAb was fused to the VT B C-terminus so as to position the sdAbs away from the oligosaccharide binding sites of the B-pentamer since retention of carbohydrate binding activity provides a convenient means of fusion protein purification by affinity chromatography (Fig. 2A). For cloning convenience arising from the presence of a PstI restriction site near the 5'-end of the sdAb gene, the VT B gene was inserted after sdAb residue 5, which placed five amino acid N-terminal extensions on the B-pentamer. The five displaced sdAb amino acids were replaced in the sdAb, which was fused to the VT B by a five amino acid spacer and followed by detection and purification tags.

A modeled structure of the pentabody (Fig. 2C) shows the five sdAbs radiating out from the C-terminal side and periphery of the VT B core. The highly symmetric representation of the pentabody is considered to be a snapshot of a highly dynamic structure. The sdAbs are thought to be highly flexible since modeling the fusion PTH50 to VT B via linker 2 without molecular overlaps in space was relatively easy.

Size exclusion chromatographic analyses showed that both PTH50 and 1V5 were very homogeneous with respect to oligomerization state. This is shown Fig. 3, Superdex 200 chromatography and SDS PAGE (12%) of an embodiment of 1V5 (A) and PTH50 (B). Based on the molecular mass markers separated under the same conditions, the mass of 1V5 was estimated to be 128 kDa, which is very close to the predicted size of 114.5 kDa for the pentameric fusion protein.

Retention of the full saccharide binding activity of VT B by 1V5 confirmed that the fusion protein was pentameric. The crystal structure of VT B in complex with a Globotriaosylceramide, Gal- α (1-4) Gal- β (1-4) Glc- β -ceramide ("Gb₃") analogue (ref. 14) shows the presence of fifteen Gb₃ trisaccharides, also known as the P^k trisaccharide, per VT B pentamer (Fig. 2A). SDS-PAGE showed that while 5 μ M PTH50 did not bind to the Synsorb™ P^k resin, the vast majority of 1V5 did bind at this concentration. At this concentration the pentameric structure is required for effective oligosaccharide binding (ref. 6). Somewhat surprisingly, surface plasmon resonance ("SPR") data showed that the K_D of the 1 V5 trisaccharide interaction was

- 21 -

approximately 1.7 nM compared to 6.6 nM for the VTB saccharide interaction. Possibly the five amino acid N-terminal extension on 1V5 (Fig. 1) interacts with the spacer on the neoglycoconjugate, resulting improved binding of the fusion protein to trisaccharide, relative to VTB. In any case, the data are consistent with the formation of a fully functional pentameric molecule. Fig. 4 presents the BIACORE analysis of an embodiment of the oligosaccharide binding properties of 2 nM VTB and 2 nM 1V5. Fitting of the data to a 1:1 interaction model gave k_a s of $2.6 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ and $5.1 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ for VTB and 1V5, respectively, and k_d s of 1×10^{-2} and 8.8×10^{-4} for VTB and 1V5, respectively.

The antigen binding profiles of PTH50 and 1V5 confirmed that the VTB fusion protein bound immobilized peptide much more effectively than the monomeric sdAb (Fig. 5 and Table 1). PTH50 binding to immobilized peptide (Fig. 5A) displayed rapid but analyzable kinetics. The K_D of the interaction was determined to be 2.5 μM (Table 1). Although rate constants could not be derived for the interaction of peptide with immobilized pentabody (Fig. 5B), the K_D was determined to be 3.6 μM (Table 1) indicating that pentamerization did not significantly alter dAb binding site accessibility. The binding of 1V5 to immobilized peptide was analyzed at low concentrations in order to maximize the binding valency and assess avidity gain conferred by dAb pentavalency (Fig 5C). Under these conditions of antigen surplus, the pentavalency conferred an avidity gain of approximately 7,000 (Table 1). Fig. 5 presents BIACORE analysis of an embodiment of the antigen binding properties of an embodiment of PTH50 and 1V5. (A) – binding of 0.25 – 2 μM PTH50 to immobilized peptide antigen; (B) – binding of 2 – 10 μM peptide to immobilized 1V5 and (c) – binding of 2.5 – 15 nM 1V5 to immobilized peptide.

BIACORE analysis was carried out according to standard procedures using CM5 sensor chips. Results as shown in Figures 4 and 5 show response units (RU) on the y axis. 1RU is a result of pg/mm^2 of protein binding to the immobilized ligand.

Detail of Example 1

Isolation of PTH50. The PTH50 sdAb was isolated from a non-immunized llama dAb library constructed as described elsewhere (ref. 13). Panning was performed with 100 µg of streptavidin-coated paramagnetic particles (SA-PMPs) (Promega, Madison, WI) that had been washed according to the manufacturer's instructions. Tubes were shaken frequently to maintain the SA-PMPs in suspension during all steps involving SA-PMPs. To reduce background binding, the library phage were pre-incubated in the absence of antigen with SA-PMPs that had been blocked with 100 µl 2% milk in phosphate buffered saline (MPBS) for 1 hr at room temperature. Also, the SA-PMPs used in panning were blocked in 400 µl MPBS at 37°C for 2 hr. The panning mixtures contained pre-adsorbed phage (10^{12} tu in the first round and 10^{11} tu in subsequent rounds), 20 mg/ml BSA, 0.05% TWEEN™ 20 and 1 µg/ml biotinylated antigen in a total volume of 150 µl MPBS. The phage-biotinylated antigen complexes were captured by transferring the mixtures to tubes containing blocked SA-PMPs followed by incubation at room temperature for 30 min. The SA-PMPs were washed five times with PBS containing 0.05% Tween-20 and then five times with PBS. Bound phage were eluted and propagated on agarose top plates as described (ref. 13). After overnight incubation at 37°C the phage particles were eluted from the plates, purified and tittered (ref. 13).

Screening of phage clones for antigen binding activity was performed by ELISA as described by Tanha *et al.* (ref. 13). Biotinylated peptide, 1 µg/ml, was captured at room temperature for 30 min in wells that had been blocked for 2 h with MPBS at 37°C following overnight coating with 5 µg/ml streptavidin at 4°C. In control experiments biotinylated antigen was replaced with appropriate buffer. Several peptide specific antibodies were identified⁴, one of which, PTH50, was selected for this study.

Construction of VTB-PTH50. Standard cloning techniques (ref. 15) were used to generate the VTB-PTH50 or 1V5 gene (Figure 1). The VTB gene (Accession EMBL M16625) was amplified by PCR with primers that introduced *Pst*I sites at both ends and added a sequence encoding DVQLQ (SEQ. ID. NO. 3) at the C-terminus of VTB.

- 23 -

The PCR product was digested by *Pst*I and ligated into the PTH50 gene (GenBank AF447918) linearized with the same enzyme. The *Pst*I site encodes residues 5 and 6 of PTH50 (Fig. 1) and hence the DVQLQ C-terminal extension (SEQ. ID. NO. 3) on the VTB PCR product. A clone in which VTB was inserted in the correct orientation was identified and designated clone pJR1V5.

Molecular modeling. The structure of 1V5 (SEQ. ID. NO. 2) was modeled using the BIOPOLYMER™ and DISCOVERY 3™ modules of INSIGHT II™ on a R10,000 SGI workstation. The PTH50 model is SEQ. ID. NO. 4 (protein) and SEQ. ID. NO. 5 (DNA). The spacers were constructed using the Insight II amino acid database. Potentials were fixed using the AMBER force field and energy minimizations were performed at all steps of the model building.

Expression and purification. *Escherichia coli* TG1 and ER2537 (New England Biolabs) harboring plasmids encoding PTH50, and PTH50-VTB were cultured and induced with 1mM isopropyl- β -D-thiogalactopyranoside. Periplasmic proteins were extracted by osmotic shock substantially according to the method of Skerra *et al.* 1991 – ref. 16 and the recombinant proteins were purified by immobilized metal affinity chromatography (IMAC) (HI-TRAP™, Amersham Pharmacia). Purifications were monitored by Western blotting with an anti-c-myc monoclonal antibody and an anti-mouse IgG/alkaline phosphatase conjugate. The purified proteins were concentrated by CENTRICON 10™ ultrafiltration (AMICON™), dialysed against 10 mM HEPES, pH 7.4, containing 150 mM NaCl and 3.4 mM EDTA and analyzed by SDS-PAGE.

Size exclusion chromatography. The oligomerization states of PTH50 and 1V5 were assessed by SUPERDEX 200™ (Amersham Pharmacia) size exclusion chromatography. In both instances separations were carried out in 10 mM HEPES, pH 7.4, containing 150 mM NaCl, 3.4 mM EDTA and 0.05% TWEEN 20™.

Oligosaccharide and antigen binding activities. To determine if 1V5 retained the oligosaccharide binding activity of VTB and if this activity could be exploited for dAb-VTB fusion protein purification 5 μ M PTH50-VTB, as well as 5 μ M PTH50, were mixed with SYNSORB P^k™, a resin comprised of the P^k trisaccharide immobilized on CHROMOSORB P™. The binding of the two proteins to the resin was monitored by

- 24 -

SDS-PAGE. The oligosaccharide binding activities of VTB and PTH50-VTB were compared by surface plasmon resonance (SPR). Analyses were performed with a BIACORE 3000™ biosensor system (Johnson) (Biacore, Inc., Piscataway, NJ) on proteins that were purified by size exclusion chromatography as described above. A
 5 neoglycoconjugate, BSA-(spacer-O-Gal α 1-4Gal β 1-4Glc β)_n (Glycorex AB), was immobilized, at a surface density of 13,000 RUs, on research grade CM5™ sensor chips (Biacore, Inc.) at a concentration of 100 μ g/ml in 10 mM acetate, pH 4.5. by amine coupling according to the manufacturer's instructions. Analyses were carried out at 25°C in 10 mM HEPES, pH 7.4, containing 150 mM NaCl, 3 mM EDTA and
 10 0.005% P-20 at a flow rate of 10 μ l per minute. The surface was regenerated by contact with 100 mM HCl for 3s.

The antigen binding profiles of PTH50 and 1V5 were analyzed by SPR. Peptide antigen and 1V5 were immobilized at surface densities of 1,300 RU and 6,000 RU, respectively, on research grade CM5 sensor chips by amine coupling and
 15 according to the manufacturer's instructions. Immobilizations were carried out at protein or peptide concentrations of 50 μ g/ml in 10 mM acetate, pH 4.5. Analyses and data analyses were performed as described above. Regeneration was not required with the 1V5 surface. The peptide surface was regenerated by contact with 10 mM borate, pH8.5, containing 1M NaCl and 0.1% P-20 for 30s. Data were evaluated with
 20 the BIAEVALUATION 3.0™ software from Biacore, Inc. Kinetics and affinity results are shown in the following Table 1.

Table 1. Kinetics and affinities of PTH50 and 1V5 binding to peptide antigen.

25	Ligand	Analyte	k_a (M ⁻¹ s ⁻¹) ^a	k_d (s ⁻¹) ^a	K_D (M)	Relative K_D
	Peptide	PTH50	-	-	7×10^{-6b} 1	
	VT1B-PTH50	peptide	-	-	4×10^{-6b} 0.6	
	peptide	1V5	3×10^5	3×10^{-4}	1×10^{-9}	0.00014

- 25 -

^aRate constants derived by data fitting to a 1:1 interaction model

^bDetermined by Scatchard analysis because the kinetics were too rapid for determination of rate constants

Thermal Stability of VTB-PTH 50 and Its Building Blocks

5 For the determination of thermal stability of the pentabody 1V5, circular dichroism (CD) spectra of 1V5 as well as its building blocks, PTH50 sdAb and the VT1B pentamer, were measured at various temperatures. Circular dichroism (CD) spectra were recorded with a Jasco J-600 spectropolarimeter connected to Neslab RTE-110 water bath. Experiments were performed in 10 mM sodium phosphate buffer
10 pH 7.0 using circular cuvettes with pathlengths of 5 cm (for PTH50 and VT1B at concentrations of 1.8 and 3 μ g/ml, respectively) and 1 cm (for 1V5 at a concentration of 9 μ g/ml). Spectra were recorded from 215 – 260 nm at 0.2 nm intervals, a scan speed of 20 nm per min, a bandwidth of 2 nm and an integration time of 1 s. Protein spectra were subtracted from a blank spectrum and subsequently smoothed by Jasco
15 software. To determine T_m s, sample temperatures were gradually increased from 30 °C to 82 °C. Spectra were recorded at various temperatures following a 10 min temperature equilibration time. An average of five ellipticity values at 235, 234.8, 234.6, 234.4 and 234.2 nm was used to plot the sigmoidal graph of ellipticity versus temperature and T_m s determined as the temperature corresponding to 50% unfolding -
20 see Fig. 7.

The data collected at 234.2, 234.4, 234.6, 234.8 and 235 nm were used to obtain denaturing curves for the proteins. No obvious changes in the CD spectra of VT1B were observed, even at temperatures as high as 70°C. Above 72°C, a sharp increase in signal was observed because of protein precipitation (data not shown).
25 This result indicates that the VT1B pentamer is a very stable structure although a T_m could not be determined. The abrupt loss of solubility at high temperature without any indication of denaturation suggests that maintenance of the structure is dependent on pentamer formation. The melting temperature of PTH50 spans a relatively wide

- 26 -

temperature range, which is typical of non-cooperative conformational change and frequently observed with small peptides. The melting temperature was calculated to be 59.7 C, indicating that the protein has very good thermostability. The fusion protein made of the two building blocks has a typical heat-denaturation curve with a T_m of 52 C (Figure 7, the heat-induced denaturation curve for pentabody 1V5). Although this number is lower than the T_m of PTH50 the fusion protein is less thermostable than VT1B, 1V5 is nonetheless a very heat stable molecule.

The thermal stability of the pentabody described here is a good indicator for use of these molecules in various applications. While high thermostability is always a useful property it is very important for *in vivo* medical applications. Despite high antigen binding affinity, a tumor-specific scFv failed to localize in xenographs because of its insufficient thermal stability (Willuda et al., 1999, ref 17). Grafting the antigen binding loops of the tumor-specific scFv onto the framework of a highly stable scFv produced a molecule with good serum stability and tumor localization.

Protease Stability of Pentabodies

Tryptic and chymotryptic digestion experiments were carried out at enzyme:pentameric sdAb ratio of 1:200 for 1 h at 37°C using sequencing grade trypsin and chymotrypsin purchased from Boehringer Mannheim. Digestion mixtures contained approximately 2 μ g/ml of 1V5 in 100 mM Tris-HCl buffer, pH 7.8. The chymotrypsin digestion mixture was supplemented with 50 mM CaCl_2 . The reactions were terminated by adding 10 μ l of 0.1 μ g/ml trypsin-chymotrypsin inhibitor (Sigma). For molecular weight determinations by mass spectrometry, DTT was added to a final concentration of 200 mM and the samples were processed as described previously by Tanha, J. et al, ref 18).

Fig. 8 shows the protease susceptibility of pentameric sdAb. 1V5 was analyzed by SDS-PAGE following digestion with trypsin for 0 and 2 hours (A and B respectively) and chymotrypsin for 0 and 2 h (C and D respectively).

- 27 -

1V5 exhibited very good resistance to digestion by trypsin and chymotrypsin. It was observed that trypsin rapidly converted 1V5 to a product with a slightly lower molecular size with no evidence of further digestion under the conditions employed (Fig. 8A and 8B). Mass spectrometric analyses indicated that the C-terminal tags had been removed. Trypsin treatment decreased the size of the 1V5 monomer by 1602 Da which corresponds to removal of the C-terminal LISEEDLNHHHHH sequence (Fig. 8C) from 1V5. No cleavage products were observed following 1 h of incubation with chymotrypsin (Fig. 8C and 8D). Mass spectrometric analyses confirmed that the bands shown in Fig. 8C and 8D corresponded to the 1V5 monomer.

The pentameric sdAb described here displayed surprising resistance to trypsin and chymotrypsin. Although there are multiple sites for both enzymes in the pentamer, there was no any evidence of degradation by either enzyme under the conditions employed here. This observation highlights one of the advantages of single domain antibodies, the smallest antigen binding fragments from conventional antibodies, namely scFvs that contain protease-sensitive linkers. Resistance to proteases is highly desirable for *in vivo* applications, such as tumor imaging. In terms of conferring protease resistance, VT1B is a logical choice as an oligomerization domain because of its natural existence in digestive environments.

Example 2 - Pentabodies in Which the sdAbs are Fused to the N-terminus of VT1B

To determine if the strategy described here is a generally suitable strategy for sdAb pentamerization, one can examine whether pentameric antibodies can be formed when 1) antibodies were fused to the N-terminus of VTB, 2) antibody molecules other than PTH50 are fused to VTB and 3) if mutant versions of VTB can be used as pentamerization domains.

For this purpose:

- (a) Protein 1V12, which is the fusion of another sdAb molecule (PTH61), to the N-terminus of a mutant version of VTB, was constructed (Fig. 9

- 28 -

illustrates the primary structure. The *ompA* sequence is removed during secretion of the mature protein to the *E. Coli* periplasm). PTH61 is a parathyroid hormone-binding sdAb isolated from the same panning experiment as PTH50. Protein 1V12 was isolated from *E. coli* TG1 cells harboring the gene encoding for 1V12 as described for the isolation of 1V5. Size exclusion chromatography was run on a Superdex 200 column, and indicated that, as for 1V5, 1V12 forms a very homogenous pentamer (Fig. 10). Protein markers (in kDa) were run on the same column and under the same conditions. Data obtained from SPR analysis show that the protein 1V12, the pentameric form of PTH61, binds much more strongly than PTH61 to its antigen (Fig. 11, surface plasmon resonance analysis of the binding of monovalent and pentavalent (1V12) to immobilized antigen).

(b) A total of eight pentabodies (1V5, 1V11, 1V12, 1V14, PES1, PVTGL10, PJS5, PSJ6) were made using both wild type and mutant versions of VTB and with different sdAb molecules fused to either N- or to the C- terminus of VT1B. All formed pentabodies that expressed well in *E. coli* TG1 cells, have little or no sign of aggregation and which bind to their antigens with a much higher functional affinity than observed for their monomeric counterparts.

Example 3 - Construction of Decabodies.

The fact that many, if not all, sdAbs can be easily pentamerized by fusing them to either the N- or to the C-terminus of different versions of VTB allows constructing decavalent antibodies by fusing one sdAb to the N-terminus and another sdAb to the C-terminus terminus of VT1B. This concept is diagrammatically illustrated in Fig. 12, a diagrammatic representation of N-terminal pentabodies , C-terminal pentabodies and a bispecific decabody. To exploit this, 1V13, a decavalent antibody, termed a decabody, was constructed. This is illustrated in Fig.13; showing the primary structure and sequence of 1V13 and the size exclusion chromatography (Superdex

- 29 -

200 column, Amersham Pharmacia) of 1V12. Protein markers (in kDa) were run on the same column and under the same conditions. 1V13 has sdAb PTH22 fused to the C-terminus and sdAb PTH61 fused to the N-terminus of a D17E/W34A mutant of VT1B (Fig 13A). PTH22 is another parathyroid hormone binding sdAb isolated from
5 the same panning experiment as PTH50 and PTH61. 1V13 was isolated from *E. coli* TG1 cells harboring the gene encoding 1V13. Size exclusion chromatography showed that 1V13 forms a pentamer, which is however not very homogenous (Fig. 13B),

Example 4 - Improved Homogeneity of Decabodies

10 Because sdAbs and VT1B, the two building units for decabody construction, are very homogeneous and are expressed well in *E. coli* cells, changes are mainly made to linkers that connect the building blocks. Four new decabodies, PJR9, PJR10, PJR13 and PJR14, were constructed using different linkers, the sequences of which are shown in Fig. 14. Fig. 14 presents a comparison by size exclusion chromatography
15 of 1V13 decavalent and its variations. The sequences of linkers connecting the sdAbs and VTB are indicated above each chromatogram. As shown in Fig. 14, the four proteins form very homogenous pentamers with no or little sign of aggregation. Protein 1V9, with the linker GPGGGGS to connect PTH61 and VTB and a linker AKRVAPELLGGPSG to connect VTB and PTH22, has the best size exclusion profile.

20 The concept of preparation and use of decavalent antibodies (decabodies) can be extended, according to another embodiment of the present invention, by use of known oligovalent, water soluble carbohydrate ligands analogous to the carbohydrate receptor of VT1B and which form complexes with VT1B, e.g. STARFISH – see Kitov, Pavel I. et.al.; 2000, ref. 19. Since STARFISH cross-links two pentameric VT1Bs, it
25 provides another means of making decabodies.

STARFISH will bind to the pentabodies of the present invention, but not to the individual units. Accordingly, a labeled version of STARFISH can be used for detecting the pentabodies described herein. Pentabodies bound to a target such as a type of

- 30 -

cancer cell, a bacterial pathogen, an anthrax spore etc., will form a complex with added, labeled STARFISH to allow detection of the STARFISH-pentabody-target complex. After addition and binding of STARFISH, residual unbound STARFISH is washed out of the system, and the bound STARFISH detected by means of its label (fluorescence, radioactivity, etc.), to measure bound pentabodies. Moreover, STARFISH-type ligands can also be used in conjunction with pentabodies of the present invention in therapeutic applications. For example, one could have two separate and distinct types of self-assembly molecules according to the invention, one to recognize and to bind to the target and another to recognize and to bind to a killer cell or compound for the target. Both types form pentabodies, for strong binding. Addition of STARFISH to cause binding together of the two types of pentabodies will bring the target and killer cells into close proximity, for interaction between them.

Thus, it is apparent that there has been provided in accordance with the invention a Novel Self-Assembly Molecules that fully satisfies the objects, aims and advantages set forth above. While the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art in light of the foregoing description. Accordingly, it is intended to embrace all such alternatives, modifications and variations as fall within the spirit and broad scope of the invention.

References

1. Pluckthun, A and Pack, P., New protein engineering approaches to multivalent and bispecific antibody fragments..., *Immunotechnology* 3, (1997) 83-105.
2. Terskikh *et.al.*, "Peptabody: a new type of high avidity binding protein, *Proc. Natl. Acad. Sci. (USA)* Vol. 94,(1997) 1663-1668.
3. Efimov, V.P., Lustig, A. & Engel, J. The thrombospondin-like chains of cartilage oligomeric matrix protein are assembled by a five-stranded alpha-helical bundle between residues 20 and 83. *FEBS Lett.* **341**, 54-58 (1994).

- 31 -

4. Hudson, P.J. & Kortt, A.A. High avidity scFv multimers; diabodies and triabodies. *J. Immunol. Methods* **231**, 177-189 (1999).
5. Kaminski, M. J. *et al.*, The role of homophilic binding in anti-tumor antibody R24 recognition of molecular surfaces, *J. Biol. Chem.*, Vol **274** No. 9 (1999) 5597-5604.
6. Soltyk, A.M. *et al.* A mutational analysis of the globotriaosylceramide binding sites of verotoxin VT1. *J. Biol. Chem.* **277**, 5351-5359 (2002).
7. Yang, W.P. *et al.* CDR walking mutagenesis for the affinity maturation of a potent human anti-HIV-1 antibody into the picomolar range. *J. Mol. Biol.* **254**, 392-403 (1995).
8. Schier, R. *et al.* Isolation of picomolar affinity anti-c-erbB-2 single-chain Fv by molecular evolution of the complementarity determining regions in the center of the antibody binding site. *J. Mol. Biol.* **263**, 551-567 (1996).
9. Ohlin, M., Owman, H., Mach, M. & Borrebaeck, C.A. Light chain shuffling of a high affinity antibody results in a drift in epitope recognition. *Mol. Immunol.* **33**, 47-56 (1996).
10. McCafferty, J., Griffiths, A.D., Winter, G. & Chiswell, D.J. Phage antibodies: filamentous phage displaying antibody variable domains. *Nature* **348**, 552-554 (1990).
11. Griffiths, A.D. *et al.* Isolation of high affinity human antibodies directly from large synthetic repertoires. *EMBO J.* **13**, 3245-3260 (1994).
12. Hoogenboom, H.R. & Chames, P. Natural and designer binding sites made by phage display technology. *Immunol. Today* **21**, 371-378 (2000).

- 32 -

13. Tanha, J., Dubuc, G., Hiram, T. Narang, S.A. & MacKenzie, C.R. Selection by phage display of llama conventional V_H fragments with heavy chain antibody V_HH properties. *J. Immunol. Methods* **263**, 97-109 (2002).
14. Ling, H. *et al.* Structure of the shiga-like toxin I B-pentamer complexed with an
5 analogue of its receptor Gb3. *Biochemistry* **37**, 1777-1788 (1998).
15. Sambrook, J. & Russel, D.W. Molecular Cloning, a laboratory manual.
16. Skerra, A., Pfitzinger, I. & Pluckthun, A. The functional expression of antibody Fv fragments in *Escherichia coli*: improved vectors and a generally applicable purification technique. *Biotechnology (N. Y.)* **9**, 273-278 (1991).
- 10 17. Willuda *et. al.*, High Thermal Stability is Essential for Tumor Targeting of Antibody fragments: engineering of a humanized anti-epithelial glycoprotein-2 (epithelial cell adhesion molecule) single-chain Fv Fragment, 1999 *Cancer Research* **59**, 5758-5767.
- 15 18. Tanha *et. al.*, Optimal Design Features of Camelized Human Single-Domain antibody Libraries, *J. Biol. Chem.* **276**, 24774-24780.
19. Kitov, Pavel I. *et.al.*, *Nature*, **403**, February 2000, 669 – 672.

- 33 -

SEQUENCE LISTING

Applicant: National Research Council of Canada

Inventors:

<110> Mackenzie, C. Roger

Zhang, Jianbing

Title:

<120> Self-Assembly Molecules

Correspondence Address:

Ridout & Maybee LLP
150 Metcalfe Street, 19th Floor,
Ottawa, Ontario K2P 1P1
Canada
Telephone: 613.236.1995
Fax: 613.236.2485

Our File Reference:

<130> 36361-0033

International Application No.:

<140> PCT/CA02/01829

Filing Date:

<141> 2002-11-29

Priority Information:

<150> US 60/334,098

<151> 2001-11-30

- 34 -

Number of Sequences:

<160> 8

Software:

<170> PatentIn version 3.1

<210> 1

<211> 31

<212> PRT

<213> mutated human

<400> 1

Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn
1 5 10 15

Ser Met Glu Arg Val Glu Trp Leu Arg Lys Leu Leu Gln Asp Val
20 25 30

<210> 2

<211> 21

<212> PRT

<213> Escherichia coli

<220>

<223> none

- 35 -

<400> 2

Met Lys Lys Thr Ala Ile Ala Ile Ala Val Ala Leu Ala Gly Phe Ala
1 5 10 15

Thr Val Ala Gln Ala
20

<210> 3

<211> 5

<212> PRT

<213> llama

<220>

<223> none

<400> 3

Asp Val Glu Leu Glu
1 5

<210> 4

<211> 116

<212> PRT

<213> llama

<220>

<223> none

<400> 4

Asp Val Gln Leu Glu Ala Ser Gly Gly Gly Ser Val Gln Ala Gly Asp
1 5 10 15

- 36 -

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Pro Phe Ser Ser Phe
20 25 30

Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45

Ala Ala Ile Ser Ala Ser Gly Gly Glu Thr Tyr Tyr Thr Gly Ser Leu
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Val Tyr
65 70 75 80

Leu Gln Met Asp Ser Leu Lys Pro Glu Asp Thr Gly Val Tyr Tyr Cys
85 90 95

Ala Ala Thr Ile Asn Gly Ala Ala Arg Arg Gly Gln Gly Thr Glu Val
100 105 110

Thr Val Ser Ser
115

<210> 5

<211> 348

<212> DNA

<213> llama

<220>

<223> none

<400> 5

gatgtgcagc tgcaggcgctc tgggggagga tcggtgcagg ctgggggactc tctgagactc 60

tctgtgcag cctctggacg ccccttcagt agctttgccca tgggctgggt cgcgccaggct 120

ccaggaaagg agcgtgagtt ttagctgct attagcgca gtggtggcga gacatactat 180

- 37 -

accggctccc tgaagggccg attcaccatc tccagagaca acgccaagaa cacggtatat 240

ctgcaaattgg acagcctgaa acctgaggat acaggcgtct attactgtgc agccaccatt 300

aacgggggcgg cccgacgagg ccaggggacc caggtcaccg tctcctca 348

<210> 6

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> none

<400> 6

Gly Gly Gly Gly Ser

1

5

<210> 7

<211> 2

<212> PRT

<213> Artificial Sequence

<220>

<223> none

<400> 7

Gly Ser

1

<210> 8

<211> 11

- 38 -

<212> PRT

<213> human

<400> 8

Glu	Gln	Lys	Leu	Ile	Ser	Glu	Glu	Asp	Leu	Asn
1		5				10				

CLAIMS:

1. A method of forming a multimeric complex having affinity for a target, said method comprising:
 - obtaining a plurality of self-assembly molecules derived from a member of the AB5 toxin family, said self-assembly molecules including complementary self-assembly units, each of which is operatively connected to an interaction domain specific for the target;
 - combining said self-assembly molecules such that at least three said self-assembly units substantially simultaneously bind to one another and permit the interaction domain to bind the target.
2. The method of claim 1, wherein the interaction domain is derived from an antibody region or a cell surface receptor.
3. The method of claim 1 or 2, wherein the interaction domains in said units recognize at least two different targets.
4. The method of any one of claims 1 to 3, wherein in each self-assembly molecule, the self-assembly unit is operatively connected to a single domain antibody at either the C-terminus or the N-terminus of the self-assembly unit.
5. The method of any one of claims 1 to 4, wherein the multimeric complex is a pentamer.
6. The method of any one of claims 1 to 5, wherein the multimeric complex is a homopentamer.
7. The method of any one of claims 1 to 6, wherein the interaction domain is a cell-surface binding region.
8. The method of any one of claims 1 to 6, wherein the interaction domain is a single domain antibody specific for the target.
9. The method of any one of claims 1 to 6, wherein the interaction domain is a single chain polypeptide encoding the V_H and the V_L regions of an antibody.

10. A self-assembly molecule having an affinity for one or more known targets and suitable for use in the formation of a multimeric complex, comprising:

a self-assembly unit derived from a member of the AB5 toxin family; and,
an interaction domain operatively connected to said self-assembly unit so as to permit binding of said interaction domain to the target and to permit binding of said self-assembly unit to another self-assembly unit.

11. A self-assembly molecule comprising a plurality of proteinaceous or peptide portions operatively connected together, a first said portion comprising a self-assembly region and a second said portion comprising an interaction domain, wherein the self-assembly region is derived from a member of the AB5 toxin family and is capable of binding to the self-assembly region derived from a member of the AB5 toxin family of at least two other self-assembly molecules comprising the same or different interaction domain, to form a complex of at least three self-assembly molecules; and wherein the interaction domain is adapted to specifically interact with a target region on another, different molecule.

12. A self-assembly molecule having an affinity for one or more known targets and suitable for use in formation of a multimeric complex, comprising:

a self-assembly unit derived from a member of the AB5 toxin family comprising a monomer of the B subunit of an AB5 toxin able to bind to another self-assembly unit derived from a member of the AB5 toxin family comprising a monomer of the B subunit of the AB5 toxin; and
an interaction domain fused directly or via a linker to said monomer of the B subunit of the AB5 toxin so as to enhance binding of the interaction domain to the target,

wherein said interaction domain comprises an antibody, antibody fragment, single domain antibody fragment, a single chain polypeptide of a VH or VL region of an antibody, a peptide or protein derived from a binding region of an antibody, or a molecule or binding portion thereof that specifically binds to cell surface receptors.

13. The self-assembly molecule of any one of claims 11 to 12, wherein the interaction domain is operatively connected at one of the N-terminus or the C-terminus of the self-assembly molecule.

14. The self-assembly molecule of any one of claims 11 to 13, wherein the self-assembly unit is derived from verotoxin.
15. The self-assembly molecule of any one of claims 11 to 13, wherein the self-assembly unit is derived from pertussis toxin.
16. The self-assembly molecule of any one of claims 11 to 15, wherein the interaction domain is a single domain antibody.
17. The self-assembly molecule of any one of claims 11 to 15, wherein the interaction domain is a cell-surface binding peptide.
18. The self-assembly molecule of any one of claims 11 to 17, further including a linker joining said interaction domain to said self-assembly unit.
19. The self-assembly molecule of any one of claims 11 to 18 further including a second interaction domain operatively connected to said self-assembly unit so as to permit binding of each interaction domain to a target.
20. The self-assembly molecule of claim 19, wherein said second interaction domain comprises an antibody, antibody fragment, single domain antibody fragment, a single chain polypeptide of a VH or VL region of an antibody, a peptide or protein derived from a binding region of an antibody, or a molecule or binding portion thereof that specifically binds to cell surface receptors.
21. The self-assembly molecule of claim 19 or 20, wherein one of said interaction domains is fused to the N-terminus and the other interaction domain is fused to the C-terminus of the self-assembly unit.
22. The self-assembly molecule according to any one of claims 11 to 21, in pentameric form.
23. The self-assembly molecule according to claim 22, wherein the pentameric form is dimerized to form a decabody.

24. The self-assembly molecule according to claim 23, wherein the dimerization is via an intermediary which binds to said molecule in pentameric form but does not bind to the individual self-assembly units in their monomeric form.
25. A method of detecting the presence of a target molecule on a surface, said method comprising:
- obtaining a plurality of self-assembly molecules derived from a member of the AB5 toxin family, each self-assembly molecule including a complementary self-assembly unit, an interaction domain specific for the target molecule and connected to the self-assembly unit, and a marker;
 - exposing said surface to the self-assembly molecules;
 - permitting said self-assembly molecules to self-assemble into a multimeric complex; and,
 - assaying for the marker.
26. The method of claim 25, wherein the marker is a radio-isotope.
27. The method of claim 25, wherein the marker is biotin.
28. A self-assembly molecule comprising a plurality of proteinaceous or peptide portions operatively connected together, a first said portion comprising a self-assembly region and a second said portion comprising an interaction domain, wherein the self-assembly region is derived from a member of the AB5 toxin family, and wherein the interaction domain is adapted to specifically interact with a target region on another, different molecule.
29. A kit comprising:
- at least one self-assembly molecule according to any one of claims 10 to 24 or claim 28; and,
 - instructions for the use of said self-assembly molecules in the specific recognition of a target.





FIG. 1

2/11

**FIG. 2A**

3/11




FIG. 2B

4/11




FIG. 2C(i)







FIG. 2C(ii)

5/11

**FIG. 3****FIG. 4**

6/11

**FIG. 5**

7/11

SEQ.ID. NO. 1:

¹SVSEIQLMHNLGKHLNSMERVEWLRKLLQDV³¹

SEQ. ID. NO. 2:

MKKTAIAIAVALAGFATVAQA

SEQ. ID. NO. 3:

DVQLQ

SEQ ID NO.4:

DVQLQASGGGSVQAGDSLRLSCAASGRPFSSFAMGWFRQAPGKEREFVAAISA
SGGETYYTGSLKGRFTISRDNANKNTVYLQMDSLKPEDTGVYYCAATINGAARRG
QGTQVTVSS

SEQ ID NO. 5:

GATGTGCAGCTGCAGGCGTCTGGGGGAGGATCGGTGCAGGCTGGGGACTCT
CTGAGACTCTCCTGTGCAGCCTCTGGACGCCCCCTTCAGTAGCTTTGCCATGG
GCTGGTTCCGCCAGGCTCCAGGAAAGGAGCGTGAGTTTGTAGCTGCTATTAG
CGCGAGTGGTGGCGAGACATACTATAACCGGCTCCCTGAAGGGCCGATTCAC
CATCTCCAGAGACAACGCCAAGAACACGGTATATCTGCAAATGGACAGCCTG
AAACCTGAGGATACAGGCGTCTATTACTGTGCAGCCACCATTAACGGGGCGG
CCCGACGAGGCCAGGGGACCCAGGTCACCGTCTCCTCA

SEQ. ID. NO. 6:

GGGGS

SEQ. ID. NO. 7:



GS

SEQ. ID. NO. 8:

EQKLISEEDLN

FIG. 6

8/11

**FIG. 7****FIG. 8**

9/11




FIG. 9




FIG. 10




FIG. 11

10/11




FIG. 12

A



PTH61	dsAb PTH61	TREMBL:AF447920
L1	spacer 1	GPGGGSGGGGS
NTB*	D17E/W34A mutant of NTB	NTB: Swissnew:P08027, AA21 -89
L2	spacer 2	GGGGSGLAGS
PTH22	sdAb PTH22	TREMBL:AF447916
m-H	C-myc and histidine tag	EQKLJSEEDLNHHHHH




FIG. 13

11/11

PTH61	L1	VTB (D17E,W34A)	L2	PTH22	m-H
-------	----	-----------------	----	-------	-----





FIG. 14



<i>ompA</i>	signal sequence	MKKTAIAIAVALAGFATVAQA	seq. ID 2
1	linker 1	DVQLQ	seq. ID 3
VTB	verotoxin B subunit	EMBL M16625: nt 1349 - 1555	
2	linker 2	GGGGS	seq. ID 6
PTH50	single domain antibody PTH50	GenBank AF447918	
3	linker 3	GS	seq. ID 7
<i>myc</i>	c-myc detection tag	EQKLISEEDLN	seq. ID 8
H	His ₅ purification tag	HHHHH	seq. ID 9