




(86) Date de dépôt PCT/PCT Filing Date: 2013/12/18
(87) Date publication PCT/PCT Publication Date: 2014/06/26
(45) Date de délivrance/Issue Date: 2021/08/10
(85) Entrée phase nationale/National Entry: 2015/06/08
(86) N° demande PCT/PCT Application No.: US 2013/076068
(87) N° publication PCT/PCT Publication No.: 2014/100147
(30) Priorité/Priority: 2012/12/21 (US61/745,013)

(51) Cl.Int./Int.Cl. *A01N 43/40* (2006.01),
A01N 43/08 (2006.01), *A01N 43/824* (2006.01),
A01P 13/00 (2006.01)
(72) Inventeurs/Inventors:
BANGEL, BRYSTON L., US;
SATCHIVI, NORBERT M., US
(73) Propriétaire/Owner:
DOW AGROSCIENCES LLC, US
(74) Agent: SMART & BIGGAR LLP

(54) Titre : COMPOSITIONS HERBICIDES COMPRENANT DE L'ACIDE 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL)-5-FLUOROPYRIDINE-2-CARBOXYLIQUE OU UN DERIVE DE CELUI-CI ET DE LA FLURTAMONE, DU DIFLUFENICAN OU DU PICOLINAFEN
(54) Title: HERBICIDAL COMPOSITIONS COMPRISING 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL)-5-FLUOROPYRIDINE-2-CARBOXYLIC ACID OR A DERIVATIVE THEREOF AND FLURTAMONE, DIFLUFENICAN, OR PICOLINAFEN




(57) Abrégé/Abstract:

Provided herein is a synergistic herbicidal composition comprising a herbicidally effective amount of (a) a benzyl ester of the compound of formula (I) and (b) picolinafen or diflufenican, or flurtamone and diflufenican. A method of controlling undesirable vegetation which comprises applying to an area where control is desired a herbicidally effective amount of said composition is also provided.

(see formula I)

ABSTRACT

Provided herein is a synergistic herbicidal composition comprising a herbicidally effective amount of (a) a benzyl ester of the compound of formula (I) and (b) picolinafen or diflufenican, or flurtamone and diflufenican. A method of controlling undesirable vegetation which comprises applying to an area where control is desired a herbicidally effective amount of said composition is also provided.




HERBICIDAL COMPOSITIONS COMPRISING 4-AMINO-3-CHLORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL)-5-FLUOROPYRIDINE-2-CARBOXYLIC ACID OR A DERIVATIVE THEREOF AND FLURTAMONE, DIFLUFENICAN, OR PICOLINAFEN

Background

[0001] The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem, researchers in the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use. However, there remains a need for compositions and methods that are effective in controlling undesirable vegetation.

Summary

[0002] Provided herein are herbicidal compositions comprising a herbicidally effective amount of a combination of (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester thereof and (b) one or more compounds selected from the group consisting of flurtamone, diflufenican, and picolinafen. In some embodiments the compositions contain (a) a compound of the formula (I) or an agriculturally acceptable salt or ester thereof and two or more compounds selected from the group consisting of flurtamone, diflufenican, picolinafen, and flufenacet. The compositions may also contain an agriculturally acceptable adjuvant or carrier.


[0003] Also provided herein is a method of controlling undesirable vegetation which comprises applying to the area where control is desired a herbicidally effective amount of a combination comprising (a) a compound of formula (I) or an agriculturally acceptable ester or salt thereof and (b) one or more compounds selected from the group consisting of flurtamone, diflufenican, and picolinafen. In some embodiments the combination is applied to the vegetation or the locus thereof. In some embodiments the combination is applied to

soil or water to prevent the emergence or growth of the vegetation. In some embodiments two or more compounds selected from the group consisting of flurtamone, diflufenican, picolinafen, and flufenacet are applied with the compound of formula (I) or an agriculturally acceptable ester or salt thereof.

Detailed Description

DEFINITIONS

[0004] As used herein, the compound of formula (I) has the following structure:



The compound of formula (I) can be identified by the name 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid and has been described in U.S. Patent 7,314,849 (B2). An exemplary form of the compound of formula (I) is the benzyl ester. Exemplary uses of the compound of the formula (I) include controlling undesirable vegetation, including grass, broadleaf and sedge weeds, in multiple non-crop and cropping situations.

[0005] As used herein, flurtamone is (±)-5-(methylamino)-2-phenyl-4-[3-(trifluoromethyl)phenyl]-3(2*H*)-furanone. Its herbicidal activity is exemplified in Tomlin, C. D. S., Ed. *The Pesticide Manual: A World Compendium*, 15th ed.; BCPC: Alton, 2009 (hereafter “*The Pesticide Manual*, Fifteenth Edition, 2009”). Exemplary uses of flurtamone include its use for pre-plant incorporated, pre-emergence or post emergence control of broad-leaved and some grass weeds in small grains, peanuts, cotton, peas and sunflowers.

[0006] As used herein, diflufenican is *N*-(2,4-difluorophenyl)-2-[3-(trifluoromethyl)phenoxy]-3-pyridinecarboxamide. As described in *The Pesticide Manual*, Fifteenth Edition, 2009, page 362, diflufenican is a selective contact and residual herbicide that is used pre- and early post-emergence in autumn-sown wheat and barley to control grass and broad-leaved weeds. It is typically used in combination with other cereal herbicides, *e.g.* flufenacet.

[0007] As used herein, picolinafen is *N*-(4-fluorophenyl)-6-[3-(trifluoromethyl)phenoxy]-2-pyridinecarboxamide. As described in *The Pesticide Manual*,

Fifteenth Edition, 2009, page 910, picolinafen is a post-emergence herbicide used either alone or in mixtures for broad-spectrum weed control in cereals.

[0008] As used herein, flufenacet is *N*-(4-fluorophenyl)-*N*-(1-methylethyl)-2-[[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]oxy]acetamide. As described in *The Pesticide Manual*, Fifteenth Edition, 2009, page 522, flufenacet is a systemic herbicide used, for example, post-emergence in maize, wheat, and rice.

[0009] As used herein, BACARA® (Bayer CropScience) is a suspension concentrate formulation containing 250 grams per liter (g/L) (22.4% weight per weight (w/w)) flurtamone and 100 g/L (8.9% w/w) diflufenican.

[0010] As used herein, BACARA® FORTE (Bayer CropScience) is a suspension concentrate containing 120 g/L flurtamone, 120 g/L diflufenican, and 120 g/L flufenacet.

[0011] As used herein, herbicide means a compound, *e.g.*, active ingredient that kills, controls or otherwise adversely modifies the growth of plants.


[0012] As used herein, a herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect to the vegetation *e.g.*, causing deviations from natural development, killing, effecting regulation, causing desiccation, causing retardation, and the like.

[0013] As used herein, controlling undesirable vegetation means preventing, reducing, killing, or otherwise adversely modifying the development of plants and vegetation. Described herein are methods of controlling undesirable vegetation through the application of certain herbicide combinations or compositions. Methods of application include, but are not limited to applications to the vegetation or locus thereof, *e.g.*, application to the area adjacent to the vegetation, as well as pre-emergence, post-emergence, foliar, and in-water applications.

[0014] As used herein, plants and vegetation include, but are not limited to, germinant seeds, emerging seedlings, plants emerging from vegetative propagules, immature vegetation, and established vegetation.

[0015] As used herein, agriculturally acceptable salts and esters refer to salts and esters that exhibit herbicidal activity, or that are or can be converted in plants, water, or soil to the referenced herbicide. Exemplary agriculturally acceptable esters are those that are or can be hydrolyzed, oxidized, metabolized, or otherwise converted, *e.g.*, in plants, water, or soil, to the corresponding carboxylic acid which, depending upon the pH, may be in the dissociated or undissociated form.

[0016] Exemplary salts include those derived from alkali or alkaline earth metals and those derived from ammonia and amines. Exemplary cations include sodium, potassium, magnesium, and aminium cations of the formula:




wherein R^1 , R^2 , R^3 and R^4 each, independently represents hydrogen or C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl or C_3 - C_{12} alkynyl, each of which is optionally substituted by one or more hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio or phenyl groups, provided that R^1 , R^2 , R^3 and R^4 are sterically compatible. Additionally, any two of R^1 , R^2 , R^3 and R^4 together may represent an aliphatic difunctional moiety containing one to twelve carbon atoms and up to two oxygen or sulfur atoms. Salts can be prepared by treatment with a metal hydroxide, such as sodium hydroxide, with an amine, such as ammonia, trimethylamine, diethanolamine, 2-methylthiopropylamine, bisallylamine, 2-butoxyethylamine, morpholine, cyclododecylamine, or benzylamine or with a tetraalkylammonium hydroxide, such as tetramethylammonium hydroxide or choline hydroxide.

[0017] Exemplary esters include those derived from C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl, C_3 - C_{12} alkynyl or C_7 - C_{10} aryl-substituted alkyl alcohols, such as methyl alcohol, isopropyl alcohol, 1-butanol, 2-ethylhexanol, butoxyethanol, methoxypropanol, allyl alcohol, propargyl alcohol, cyclohexanol or unsubstituted or substituted benzyl alcohols. Benzyl alcohols may be substituted with from 1-3 substituents independently selected from halogen, C_1 - C_4 alkyl or C_1 - C_4 alkoxy. Esters can be prepared by coupling of the acids with the alcohol using any number of suitable activating agents such as those used for peptide couplings such as dicyclohexylcarbodiimide (DCC) or carbonyl diimidazole (CDI); by reacting the acids with alkylating agents such as alkylhalides or alkylsulfonates in the presence of a base such as triethylamine or lithium carbonate; by reacting the corresponding acid chloride of an acid with an appropriate alcohol; by reacting the corresponding acid with an appropriate alcohol in the presence of an acid catalyst or by transesterification.

[0018] As used herein when expressing weight ratios of the compound of formula (I) or a salt or ester thereof to other compounds, in cases where a salt or ester of the compound of formula (I) is involved, the weight is the acid equivalent weight.

COMPOSITIONS AND METHODS

[0019] Provided herein are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester thereof, and (b) a compound selected from the group consisting of flurtamone, diflufenican, and picolinafen. In certain embodiments, the composition comprises (a) the compound of formula (I) or an agriculturally acceptable salt or ester thereof and (b) diflufenican. In certain embodiments, the composition comprises (a) the compound of formula (I) or an agriculturally acceptable salt or ester thereof and (b) picolinafen. In certain embodiments, the composition comprises (a) the compound of formula (I) or an agriculturally acceptable salt or ester thereof and (b) flurtamone.

[0020] Also provided are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I) or an agriculturally acceptable salt or ester thereof and two or more compounds selected from the group consisting of flurtamone, diflufenican, picolinafen, and flufenacet. In certain embodiments, the composition comprises (a) the compound of formula (I) or an agriculturally acceptable salt or ester thereof, (b) flurtamone, and (c) diflufenican. In certain embodiments, the composition comprises (a) the compound of formula (I) or an agriculturally acceptable salt or ester thereof, (b) flurtamone, (c) diflufenican, and (d) flufenacet.

[0021] Also provided are methods of controlling undesirable vegetation comprising contacting the vegetation or the locus thereof, *i.e.*, area adjacent to the vegetation, with or applying to the soil or water to prevent the emergence or growth of vegetation a herbicidally effective amount of a combination comprising the compound of formula (I) or an agriculturally acceptable salt or ester thereof and (b) one or more compounds selected from the group consisting of flurtamone, diflufenican, picolinafen, and flufenacet. In certain embodiments, the combination comprises (a) the benzyl ester of the compound of formula (I) and (b) diflufenican. In certain embodiments, the combination comprises (a) the benzyl ester of the compound of formula (I) and (b) picolinafen. In certain embodiments, the combination comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, and (c) diflufenican. In certain embodiments, the combination comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, (c) diflufenican, and (d) flufenacet.

[0022] Furthermore, in some embodiments, the two component and multiple component combinations described above exhibit synergism, *i.e.*, the herbicidal active ingredients are more effective in combination than when applied individually. Synergism has been defined as “an interaction of two or more factors such that the effect when combined is greater than the predicted effect based on the response of each factor applied separately.” Senseman, S., Ed. *Herbicide Handbook*. 9th ed. Lawrence: Weed Science Society of America, 2007. In certain embodiments, the compositions exhibit synergy as determined by the Colby equation (Colby, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. *Weeds* **1967**, *15*, 20-22).

[0023] More specifically, the following equation is used to calculate the expected activity of mixtures containing two herbicidal active ingredients:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of a first active ingredient at the same concentration as used in the mixture;

B = observed efficacy of the second active ingredient (or product, which may be a combination of active ingredients) at the same concentration as used in the mixture.

[0024] In certain embodiments of the compositions and methods described herein, the compound of formula (I), *i.e.*, the carboxylic acid, is employed. In certain embodiments, a carboxylate salt, *e.g.* the potassium salt, of the compound of formula (I) is employed. In certain embodiments, an arylalkyl or alkyl ester is employed. In certain embodiments, a benzyl, substituted benzyl, or C₁₋₄ alkyl, *e.g.*, *n*-butyl ester is employed. In certain embodiments, the benzyl ester is employed.

[0025] In some embodiments, the herbicidal active ingredients are formulated in one composition, tank-mixed, applied simultaneously, or applied sequentially.

[0026] Herbicidal activity is exhibited by the herbicidal compositions when they are applied directly to the plant or to the locus of the plant at any stage of growth. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted to promote non-selective or selective herbicidal action. In some embodiments, the compositions described herein are

applied as a post-emergence application, pre-emergence application, or in-water application to flooded paddy rice or water bodies (*e.g.*, ponds, lakes and streams), to relatively immature undesirable vegetation to achieve the maximum control of weeds.

[0027] In some embodiments, the compositions and methods provided herein are utilized to control weeds in crops, *e.g.* cereals, including but not limited to rice, wheat, triticale, barley, oats, rye, sorghum, and corn/maize, and in pastures, grasslands, rangelands, fallowland, industrial vegetation management (IVM) and rights-of-way.

[0028] The compositions and methods described herein be used to control undesirable vegetation on glyphosate-tolerant-, glufosinate-tolerant-, dicamba-tolerant-, phenoxy auxin-tolerant-, pyridyloxy auxin-tolerant-, aryloxyphenoxypropionate-tolerant-, acetyl CoA carboxylase (ACCase) inhibitor-tolerant-, imidazolinone-tolerant-, acetolactate synthase (ALS) inhibitor-tolerant-, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitor -tolerant-, protoporphyrinogen oxidase (PPO) inhibitor-tolerant-, triazine-tolerant-, and bromoxynil-tolerant- crops, for example, in conjunction with glyphosate, dicamba, phenoxy auxins, pyridyloxy auxins, aryloxyphenoxypropionates, ACCase inhibitors, imidazolinones, ALS inhibitors, HPPD inhibitors, PPO inhibitors, triazines, and bromoxynil. The compositions and methods may be used in controlling undesirable vegetation in crops possessing multiple or stacked traits conferring tolerance to multiple chemistries and/or inhibitors of multiple modes-of-action. In some embodiments, the compound of formula (I) or salt or ester thereof and complementary herbicide or salt or ester thereof are used in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. In some embodiments, the compositions described herein and other complementary herbicides are applied at the same time, either as a combination formulation or as a tank mix.

[0029] The compositions and methods provided herein are utilized to control undesirable vegetation. Undesirable vegetation includes, but is not limited to, undesirable vegetation that occurs in rice, cereals, range and pasture, and non-crop settings, (*e.g.*, rights-of-way, IVM).

[0030] In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation including, for example, *Brassica*, *Sinapis*, *Kochia*, *Lamium*, *Salsola*, *Veronica*, *Chenopodium*, *Viola*, *Cirsium*, *Matricaria*, *Galium*, *Papaver*, and/or *Stellaria*.

[0031] In some embodiments, the methods provided herein are utilized to control undesirable vegetation in cereals. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Apera spica-venti* (L.) Beauv. (windgrass, APESV), *Avena fatua* L. (wild oat, AVEFA), *Bromus tectorum* L. (downy brome, BROTE), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Lolium rigidum* (rigid ryegrass), *Lolium multiflorum subsp. Gaudini* (annual ryegrass, LOLMG), *Phalaris minor* Retz. (littleseed canarygrass, PHAMI), *Poa annua* L. (annual bluegrass, POAAN), *Setaria pumila* (Poir.) Roemer & J.A. Schultes (yellow foxtail, SETLU), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Amaranthus retroflexus* (redroot pigweed, AMARE), *Chenopodium album* (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Galium aparine* L. (catchweed bedstraw, GALAP), *Kochia scoparia* (L.) Schrad. (kochia, KCHSC), *Lamium purpureum* L. (purple deadnettle, LAMPU), *Matricaria recutita* L. (wild chamomile, MATCH), *Matricaria matricarioides* (Less.) Porter (pineappleweed, MATMT), *Papaver rhoeas* L. (common poppy, PAPRH), *Polygonum convolvulus* L. (wild buckwheat, POLCO), *Salsola tragus* L. (Russian thistle, SASKR), *Sinapis arvensis* (wild mustard, SINAR), *Stellaria media* (L.) Vill. (common chickweed, STEME), *Veronica hederifolia* (Ivy-leaved speedwell, VERHE), *Veronica persica* Poir. (Persian speedwell, VERPE), *Viola arvensis* Murr. (field violet, VIOAR), or *Viola tricolor* L. (wild violet, VIOTR).

[0032] In some embodiments, the methods provided herein are utilized to control undesirable vegetation in rice. In certain embodiments, the undesirable vegetation is *Brachiaria platyphylla* (Groseb.) Nash (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa colonum* (L.) LINK (junglerice, ECHCO), *Echinochloa oryzoides* (Ard.) Fritsch (early watergrass, ECHOR), *Echinochloa oryzicola* (Vasinger) Vasinger (late watergrass, ECHPH), *Ischaemum rugosum* Salisb. (saramollagrass, ISCRU), *Leptochloa chinensis* (L.) Nees (Chinese sprangletop, LEFCH), *Leptochloa fascicularis* (Lam.) Gray (bearded sprangletop, LEFFA), *Leptochloa panicoides* (Presl.) Hitchc. (Amazon sprangletop, LEFPA), *Panicum dichotomiflorum* (L.) Michx. (fall panicum, PANDI), *Paspalum dilatatum* Poir. (dallisgrass, PASDI), *Cyperus difformis* L. (smallflower flatsedge, CYPDI), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus iria* L. (rice flatsedge, CYPRI), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Eleocharis* species (ELOSS), *Fimbristylis miliacea* (L.) Vahl (globe fringerush, FIMMI), *Schoenoplectus*

juncoides Roxb. (Japanese bulrush, SPCJU), *Schoenoplectus maritimus* L. (sea clubrush, SCPMA), *Schoenoplectus mucronatus* L. (ricefield bulrush, SCPMU), *Aeschynomene* species, (jointvetch, AESSS), *Alternanthera philoxeroides* (Mart.) Griseb. (alligatorweed, ALRPH), *Alisma plantago-aquatica* L. (common waterplantain, ALSPA), *Amaranthus* species, (pigweeds and amaranths, AMASS), *Ammannia coccinea* Rottb. (redstem, AMMCO), *Eclipta alba* (L.) Hassk. (American false daisy, ECLAL), *Heteranthera limosa* (SW.) Willd./Vahl (ducksalad, HETLI), *Heteranthera reniformis* R. & P. (roundleaf mudplantain, HETRE), *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory, IPOHE), *Lindernia dubia* (L.) Pennell (low false pimpernel, LIDDU), *Monochoria korsakowii* Regel & Maack (monochoria, MOOKA), *Monochoria vaginalis* (Burm. F.) C. Presl ex Kuhth, (monochoria, MOOVA), *Murdannia nudiflora* (L.) Brenan (doveweed, MUDNU), *Polygonum pensylvanicum* L., (Pennsylvania smartweed, POLPY), *Polygonum persicaria* L. (ladysthumb, POLPE), *Polygonum hydropiperoides* Michx. (POLHP, mild smartweed), *Rotala indica* (Willd.) Koehne (Indian toothcup, ROTIN), *Sagittaria* species, (arrowhead, SAGSS), *Sesbania exaltata* (Raf.) Cory/Rydb. Ex Hill (hemp sesbania, SEBEX), or *Sphenoclea zeylanica* Gaertn. (gooseweed, SPDZE).

[0033] In some embodiments, the methods provided herein are utilized to control undesirable vegetation in range and pasture. In certain embodiments, the undesirable vegetation is *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Cassia obtusifolia* (sickle pod, CASOB), *Centaurea maculosa* auct. non Lam. (spotted knapweed, CENMA), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Convolvulus arvensis* L. (field bindweed, CONAR), *Euphorbia esula* L. (leafy spurge, EPHES), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Plantago lanceolata* L. (buckhorn plantain, PLALA), *Rumex obtusifolius* L. (broadleaf dock, RUMOB), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Sonchus arvensis* L. (perennial sowthistle, SONAR), *Solidago* species (goldenrod, SOOSS), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Urtica dioica* L. (common nettle, URTDI).

[0034] In some embodiments, the methods provided herein are utilized to control undesirable vegetation found in row crops. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Avena fatua* L. (wild oat, AVEFA), *Brachiaria platyphylla* (Groseb.) Nash (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv.

(barnyardgrass, ECHCG), *Echinochloa colonum* (L.) Link (junglerice, ECHCO), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Panicum dichotomiflorum* Michx. (fall panicum, PANDI), *Panicum miliaceum* L. (wild-proso millet, PANMI), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Sorghum halepense* (L.) Pers. (Johnsongrass, SORHA), *Sorghum bicolor* (L.) Moench ssp. *Arundinaceum* (shattercane, SORVU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Abutilon theophrasti* Medik. (velvetleaf, ABUTH), *Amaranthus* species (pigweeds and amaranths, AMASS), *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Ambrosia psilostachya* DC. (western ragweed, AMBPS), *Ambrosia trifida* L. (giant ragweed, AMBTR), *Asclepias syriaca* L. (common milkweed, ASCSY), *Chenopodium album* L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Commelina benghalensis* L. (tropical spiderwort, COMBE), *Datura stramonium* L. (jimsonweed, DATST), *Daucus carota* L. (wild carrot, DAUCA), *Euphorbia heterophylla* L. (wild poinsettia, EPHHL), *Erigeron bonariensis* L. (hairy fleabane, ERIBO), *Erigeron canadensis* L. (Canadian fleabane, ERICA), *Helianthus annuus* L. (common sunflower, HELAN), *Jacquemontia tamnifolia* (L.) Griseb. (smallflower morningglory, IAQTA), *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory, IPOHE), *Ipomoea lacunosa* L. (white morningglory, IPOLA), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Portulaca oleracea* L. (common purslane, POROL), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Solanum ptychanthum* Dunal (eastern black nightshade, SOLPT), or *Xanthium strumarium* L. (common cocklebur, XANST).

[0035] In some embodiments, the compositions and methods provided herein are used to control *Brachiaria platyphylla* (Griseb.) Nash (broadleaf signalgrass, BRAPP), *Chamomilla chamomilla* (L.) Rydb. (scented mayweed, MATCH), *Cyperus difformis* L. (smallflower umbrella sedge, CYPDI), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Cyperus iria* L. (rice flatsedge, CYPPIR), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) Beauv. (barnyardgrass, ECHCG), *Ipomoea hederacea* Jacq. (ivyleaf morningglory, IPOHE), *Lamium purpureum* (L.) (purple deadnettle, LAMPU), *Leptochloa chinensis* (L.) Nees (Chinese sprangletop, LEFCH), *Leptochloa panicoides* (J. Presl) A.S. Hitchc. (Amazon sprangletop, LEFPA), *Salsola iberica* (L.) (Russian thistle, SASKR), *Schoenoplectus*

juncoides (Roxb.) Palla (Japanese bulrush, SCPJU), *Veronica persica* Poir. (bird's-eye speedwell, VERPE) and/or *Viola tricolor* (L.) (wild pansy, VIOTR).

[0036] In certain embodiments, the methods and compositions provided herein are used to control *Ipomoea hederacea* (ivyleaf morningglory, IPOHE), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Abutilon theophrasti* Medik. (velvetleaf, ABUTH), *Euphorbia heterophylla* L. (wild poinsettia, EPHHL), *Amaranthus retroflexus* L. (redroot pigweed, AMARE), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Chenopodium album* L. (common lambsquarters, CHEAL), *Viola tricolor* L. (wild violet, VIOTR), *Stellaria media* (L.) Vill. (common chickweed, STEME), and/ or *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR).

[0037] The compounds of formula (I) or agriculturally acceptable salt or ester thereof may be used to control herbicide resistant or tolerant weeds. The methods employing the combination of a compound of formula (I) or agriculturally acceptable salt or ester thereof and the compositions described herein may also be employed to control herbicide resistant or tolerant weeds. Exemplary resistant or tolerant weeds include, but are not limited to, biotypes resistant or tolerant to acetolactate synthase (ALS) inhibitors, photosystem II inhibitors, acetyl CoA carboxylase (ACCase) inhibitors, synthetic auxins, photosystem I inhibitors, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, microtubule assembly inhibitors, lipid synthesis inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, carotenoid biosynthesis inhibitors, very long chain fatty acid (VLCFA) inhibitors, phytoene desaturase (PDS) inhibitors, glutamine synthetase inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors, mitosis inhibitors, cellulose biosynthesis inhibitors, herbicides with multiple modes-of-action such as quinclorac, and unclassified herbicides such as arylaminopropionic acids, difenzoquat, endothall, and organoarsenicals. Exemplary resistant or tolerant weeds include, but are not limited to, biotypes with resistance or tolerance to multiple herbicides, multiple chemical classes, and multiple herbicide modes-of-action.

[0038] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a two component combination with flurtamone. In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone is from about 1-40 of (a) to about 30-250 of (b). In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone is from about 1.25-10 of (a) to about 30-250 of (b). In certain

embodiments, the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone is from about 2.5-7.5 of (a) to about 60-125 of (b). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I) and (b) flurtamone, wherein the weight ratio of the two components is from about 2.5-7.5 of (a) to about 60 -125 of (b).

[0039] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a three component combination with flurtamone and flufenacet. In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) flufenacet is from about 1-40 of (a) to about 30-250 of (b) to about 15-240 of (c). In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) flufenacet is from about 1.25-10 of (a) to about 30-250 of (b) to about 15-240 of (c). In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) flufenacet is from about 2.5-7.5 of (a) to about 60-125 of (b) to about 30-60 of (c). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, and (c) flufenacet, wherein the weight ratio of the three components is from about 2.5-7.5 of (a) to about 60-125 of (b) to about 30-90 of flufenacet.

[0040] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a three component combination with flurtamone and diflufenican. In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) diflufenican is from about 1-40 of (a) to about 30-250 of (b) to about 12.5-100 of (c). In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) diflufenican is from about 1.25-10 of (a) to about 30-250 of (b) to about 12.5-100 of (c). In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) diflufenican is from about 2.5-7.5 of (a) to about 60-125 of (b) to about 25-75 of diflufenican. In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, and (c) diflufenican, wherein the weight

ratio of the three components is from about 2.5-7.5 of (a) to about 60-125 of (b) to about 25-75 of diflufenican. In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, and (c) diflufenican, wherein the weight ratio of the three components is from about 1.0-40 of (a) to about 30-250 of (b) to about 1.25-100 of (c).

[0041] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a four component combination with flurtamone, diflufenican, and flufenacet. In some embodiments, the four components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) diflufenican to (d) flufenacet is from about 1-40 of (a) to about 30-250 of (b) to about 12.5-100 of (c) to about 15-240 of (d). In some embodiments, the four components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) diflufenican to (d) flufenacet is from about 1.25-10 of (a) to about 30-250 of (b) to about 12.5-100 of (c) to about 15-240 of (d). In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to (b) flurtamone to (c) diflufenican to (d) flufenacet is from about 2.5-7.5 of (a) to about 60-125 of (b) to about 25-75 of (c) to about 30-90 of (d). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, (c) diflufenican, and (d) flufenacet, wherein the weight ratio of the three components is from about 2.5-7.5 of (a) to about 60-125 of (b) to about 25-75 of (c) to about 30-90 of (d). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) flurtamone, (c) diflufenican, and (d) flufenacet, wherein the weight ratio of the three components is from about 1.0-40 of (a) to about 30-250 of (b) to about 1.25-100 of (c) to about 15-240 of (d).

[0042] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a two component combination with picolinafen. In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) picolinafen is from about 1-40 of (a) to about 1.25-100 of (b). In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) picolinafen is from about 1.25-10 of (a) to about 1.25-100 of (b). In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) picolinafen is from about 1.25-10 of (a)

to about 12.5-100 of (b). In certain embodiments, the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) picolinafen is from about 2.5-7.5 of (a) to about 25-75 of (b). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I) and (b) picolinafen, wherein the weight ratio of the two components is from about 2.5-7.5 of (a) to about 25-75 of (b). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I) and (b) picolinafen, wherein the weight ratio of the two components is from about 1.0-40 of (a) to about 12.5-100 of (b).

[0043] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a two component combination with diflufenican. In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican is from about 1-40 of (a) to about 1.25-100 of (b). In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican is from about 1-40 of (a) to about 12.5-100 of (b). In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican is from about 1.25-10 of (a) to about 1.25-100 of (b). In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican is from about 1-40 of (a) to about 12.5-100 of (b). In certain embodiments, the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican is from about 2.5-7.5 of (a) to about 25-75 of (b). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I) and (b) diflufenican, wherein the weight ratio of the two components is from about 2.5-7.5 of (a) to about 25-75 of (b). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I) and (b) diflufenican, wherein the weight ratio of the two components is from about 1.0-40 of (a) to about 1.25-100 of (b).

[0044] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a two component combination with flufenacet. In some embodiments, the two components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flufenacet is from about 1-40 of (a) to about 15-240 of (b). In some embodiments, the two components are used

in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flufenacet is from about 1.25-10 of (a) to about 15-240 of (b). In certain embodiments, the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) flufenacet is from about 2.5-7.5 of (a) to about 30-90 of (b). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I) and (b) flufenacet, wherein the weight ratio of the two components is from about 2.5-7.5 of (a) to about 30-90 of (b).

[0045] In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in a three component combination with diflufenican and flufenacet. In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican to (c) flufenacet is from about 1.0-40 of (a) to about 12.5-100 of (b) to about 15-240 of (c). In some embodiments, the three components are used in amounts such that the weight ratio of (a) the compound of formula (I) or salt or ester thereof to (b) diflufenican to (c) flufenacet is from about 1.25-10 of (a) to about 12.5-100 of (b) to about 15-240 of (c). In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to (b) diflufenican to (c) flufenacet is from about 2.5-7.5 of (a) to about 25-75 of (b) to about 30-90 of (c). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) diflufenican, and (c) flufenacet, wherein the weight ratio of the three components is from about 2.5-7.5 of (a) to about 25-75 of (b) to about 30-90 of (c). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) diflufenican, and (c) flufenacet, wherein the weight ratio of the three components is from about 1.0-40 of (a) to about 12.5-100 of (b) to about 15-240 of (c). In one embodiment, the composition comprises (a) the benzyl ester of the compound of formula (I), (b) diflufenican, and (c) flufenacet, wherein the weight ratio of the three components is from about 1.0-40 of (a) to about 12.5-100 of (b) to about 15-240 of (c).

[0046] With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof with the herbicidally active components or applying the herbicidally active components of a composition described herein to the soil or water to prevent the emergence or growth of vegetation. In some embodiments, the composition is applied at an application rate from about 30 grams active ingredient per

hectare (g ai/ha) to about 500 g ai/ha based on the total amount of herbicidal active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 60 g ai/ha to about 200 g ai/ha based on the total amount of active ingredients in the composition.

[0047] In some two component embodiments, picolinafen is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1 grams acid equivalent per hectare (g ae/ha) to about 40 g ae/ha. In some two component embodiments, picolinafen is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 grams acid equivalent per hectare (g ae/ha) to about 10 g ae/ha. In some embodiments, picolinafen is applied at a rate from about 25 g ai/ha to about 75 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester and picolinafen. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and picolinafen, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, and picolinafen is applied at a rate from about 25 g ai/ha to about 75 g ai/ha.

[0048] In some two component embodiments, diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.0 g ae/ha to about 40 g ae/ha. In some two component embodiments, diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from

about 1.25 g ae/ha to about 10 g ae/ha. In some embodiments, diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester and diflufenican. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and diflufenican, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, and diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha.

[0049] In some three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.0 g ae/ha to about 40 g ae/ha, diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha, and flufenacet is applied at a rate from about 15 g ai/ha to about 240 g ai/ha. In some three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 10 g ae/ha, diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha, and flufenacet is applied at a rate from about 15 g ai/ha to about 240 g ai/ha. In some embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, the diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha, and flufenacet is applied at a rate from about 30 g ai/ha to about 90 g ai/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester, diflufenican, and flufenacet. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I), diflufenican, and flufenacet wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha, and flufenacet is applied at a rate from about 30 g ai/ha to about 90 g ai/ha.

[0050] In some two component embodiments, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.0 g ae/ha to about 40 g ae/ha. In some two component embodiments, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 10 g ae/ha. In some embodiments, flurtamone is applied at a rate from about 60 g ai/ha to about 125 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester and flurtamone. In one

embodiment, the methods utilize the benzyl ester of the compound of formula (I) and flurtamone, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, and flurtamone is applied at a rate from about 60 g ai/ha to about 125 g ai/ha.

[0051] In some three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.0 g ae/ha to about 40 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, and diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha. In some three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 10 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, and diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha. In other three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 10 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, and diflufenican is applied at a rate from about 1.25 g ai/ha to about 100 g ai/ha. In some embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, the flurtamone is applied at a rate from about 60g ai/ha to about 125 g ai/ha, and diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester, flurtamone, and diflufenican. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I), flurtamone, and diflufenican wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, flurtamone is applied at a rate from about 60 g ai/ha to about 125 g ai/ha, and diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha.

[0052] In some three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.0 g ae/ha to about 40 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, and flufenacet is applied at a rate from about 15 g ai/ha to about 240 g ai/ha. In some three component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 10 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, and flufenacet is applied at a rate from about 15 g ai/ha to about 240 g ai/ha. In some embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, the flurtamone is applied at a rate from about 60g ai/ha

to about 125 g ai/ha, and flufenacet is applied at a rate from about 30 g ai/ha to about 90 g ai/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester, flurtamone, and flufenacet. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I), flurtamone, and flufenacet wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, flurtamone is applied at a rate from about 60 g ai/ha to about 125 g ai/ha, and flufenacet is applied at a rate from about 30 g ai/ha to about 90 g ai/ha.

[0053] In some four component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.0 g ae/ha to about 40 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha, and flufenacet is applied at a rate from about 15 g ai/ha to about 240 g ai/ha. In some four component embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 1.25 g ae/ha to about 10 g ae/ha, flurtamone is applied at a rate from about 30 g ai/ha to about 250 g ai/ha, diflufenican is applied at a rate from about 12.5 g ai/ha to about 100 g ai/ha, and flufenacet is applied at a rate from about 15 g ai/ha to about 240 g ai/ha. In some embodiments, the compound of formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, the flurtamone is applied at a rate from about 60g ai/ha to about 125 g ai/ha, diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha, and flufenacet is applied at a rate from about 30 g ai/ha to about 90 g ai/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl ester, flurtamone, and diflufenican. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I), flurtamone, and diflufenican, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 2.5 g ae/ha to about 7.5 g ae/ha, flurtamone is applied at a rate from about 60 g ai/ha to about 125 g ai/ha, diflufenican is applied at a rate from about 25 g ai/ha to about 75 g ai/ha, and flufenacet is applied at a rate from about 30 g ai/ha to about 90 g ai/ha.

[0054] The components of the mixtures described herein can be applied either separately or as part of a multipart herbicidal system.

[0055] The mixtures described herein can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank-mixed with the other herbicide or herbicides or applied sequentially with

the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the compositions and methods described herein include, but are not limited to: 4-CPA, 4-CPB, 4-CPP, 2,4-D, 2,4-D choline salt, 2,4-D esters and amines, 2,4-DB, 3,4-DA, 3,4-DB, 2,4-DEB, 2,4-DEP, 3,4-DP, 2,3,6-TBA, 2,4,5-T, 2,4,5-TB, acetochlor, acifluorfen, acetonifene, acrolein, alachlor, allidochlor, alloxydim, allyl alcohol, alorac, ametrifone, ametryn, amibuzin, amicarbazone, amidosulfuron, aminocyclopyrachlor, aminopyralid, amiprofos-methyl, amitrole, ammonium sulfamate, anilofos, anisuron, asulam, atraton, atrazine, azafenidin, azimsulfuron, aziprotryne, barban, BCPC, beflubutamid, benazolin, bencarbazone, benfluralin, benfuresate, bensulfuron-methyl, bensulide, benthioncarb, bentazon-sodium, benzadox, benzfendazole, benzipram, benzobicyclon, benzofenap, benzofluor, benzoylprop, benzthiazuron, bicyclopiron, bifenox, bilanafos, bispyribac-sodium, borax, bromacil, bromobonil, bromobutide, bromofenoxim, bromoxynil, brompyrazon, butachlor, butafenacil, butamifos, butenachlor, buthidazole, buthiuron, butralin, butoxydim, buturon, butylate, cacodylic acid, cafenstrole, calcium chlorate, calcium cyanamide, cambendichlor, carbasulam, carbetamide, carboxazole, chlorprocarb, carfentrazone-ethyl, CDEA, CEPC, chlormethoxyfen, chloramben, chloranocryl, chlorazifop, chlorazine, chlorbromuron, chlorbufam, chloreturon, chlorfenac, chlorfenprop, chlorflurazole, chlorflurenol, chloridazon, chlorimuron, chlornitrofen, chloropon, chlorotoluron, chloroxuron, chloroxynil, chlorpropham, chlorsulfuron, chlorthal, chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, cisanilide, clethodim, cliodinate, clodinafop-propargyl, clofop, clomazone, clomeprop, cloprop, cloproxydim, clopyralid, cloransulam-methyl, CMA, copper sulfate, CPMF, CPPC, credazine, cresol, cumyluron, cyanatryl, cyanazine, cycloate, cyclosulfamuron, cycloxydim, cycluron, cyhalofop-butyl, cyperquat, cyprazine, cyprazole, cypromid, daimuron, dalapon, dazomet, delachlor, desmedipham, desmetryn, di-allate, dicamba, dichlobenil, dichloralurea, dichlormate, dichlorprop, dichlorprop-P, diclofop-methyl, diethamquat, diethatyl, difenopenten, difenoxuron, difenzoquat, diflufenzopyr, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimexano, dimidazon, dinitramine, dinofenat, dinoprop, dinosam, dinoseb, dinoterb, diphenamid, dipropetryn, diquat, disul, dithiopyr, diuron, DMPA, DNOC, DSMA, EBEP, eglinazine, endothal, epronaz, EPTC, erbon, esprocarb, ethalfluralin, ethbenzamide, ethametsulfuron, ethidimuron, ethiolate, ethobenzamid, etobenzamid, ethofumesate, ethoxyfen, ethoxysulfuron, etinofen, etniproamid, etobenzanid, EXD, fenasulam, fenoprop, fenoxaprop, fenoxaprop-P-ethyl, fenoxaprop-P-

ethyl + isoxadifen-ethyl, fenoxasulfone, fenquinotrione, fenteracol, fenthiaprop, fentrazamide, fenuron, ferrous sulfate, flamprop, flamprop-M, flazasulfuron, fluazifop, fluazifop-P-butyl, fluazolate, flucarbazone, flucetosulfuron, fluchloralin, flufenican, flufenpyr-ethyl, flumezin, flumiclorac-pentyl, flumioxazin, flumipropyn, fluometuron, fluorodifen, fluoroglycofen, fluoromidine, fluoronitrofen, fluothiuron, flupoxam, flupropacil, flupropanate, flupyrsulfuron, fluridone, flurochloridone, fluroxypyr, fluthiacet, fomesafen, foramsulfuron, fosamine, fumiclorac, furyloxyfen, glufosinate, glufosinate-ammonium, glufosinate-P-ammonium, glyphosate, halosafen, halosulfuron-methyl, haloxydine, haloxyfop-methyl, haloxyfop-P-methyl, hexachloroacetone, hexaflurate, hexazinone, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazosulfuron, indanofan, indaziflam, iodobonil, iodomethane, iodosulfuron, iodosulfuron-ethyl-sodium, iofensulfuron, ioxynil, ipazine, ipfencarbazone, iprymidam, isocarbamid, isocil, isomethiozin, isonoruron, isopollinate, isopropalin, isoproturon, isouron, isoxaben, isoxachlortole, isoxaflutole, isoxapyrifop, karbutilate, ketospiradox, lactofen, lenacil, linuron, MAA, MAMA, MCPA esters and amines, MCPA-thioethyl, MCPB, mecoprop, mecoprop-P, medinoterb, mefenacet, mefluidide, mesoprazine, mesosulfuron, mesotrione, metam, metamifop, metamitron, metazachlor, metazosulfuron, metflurazon, methabenzthiazuron, methalpropalin, methazole, methiobencarb, methiozolin, methiuron, methometon, methoprottryne, methyl bromide, methyl isothiocyanate, methylidymron, metobenzuron, metobromuron, metolachlor, metoxuron, metribuzin, metsulfuron, metsulfuron-methyl, molinate, monalide, monisouron, monochloroacetic acid, monolinuron, monuron, morfamquat, MSMA, naproanilide, napropamide, napropamide-M, naptalam, neburon, nicosulfuron, nipyraclufen, nitralin, nitrofen, nitrofluorfen, norflurazon, noruron, OCH, orbencarb, *ortho*-dichlorobenzene, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxapyrazon, oxasulfuron, oxaziclomefone, oxyfluorfen, paraflufen-ethyl, parafluron, paraquat, pebulate, pelargonic acid, pendimethalin, pentachlorophenol, pentanochlor, pentoxazone, perfluidone, pethoxamid, phenisopham, phenmedipham, phenmedipham-ethyl, phenobenzuron, phenylmercury acetate, picloram, picolinafen, pinoxaden, piperophos, potassium arsenite, potassium azide, potassium cyanate, pretilachlor, primisulfuron-methyl, procyzazine, prodiamine, profluazol, profluralin, profoxydim, proglinazine, prohexadione-calcium, prometon, prometryn, pronamide, propachlor, propanil, propaquizafop, propazine, propham, propisochlor, propoxycarbazone, propyrisulfuron, propyzamide, prosulfalin, prosulfocarb, prosulfuron, proxan, prynachlor, pydanon, pyracilonil, pyraflufen-ethyl, pyrasulfotole, pyrazogyl, pyrazolynate,

pyrazosulfuron-ethyl, pyrazoxyfen, pyribenzoxim, pyributicarb, pyriclor, pyridafof, pyridate, pyriftalid, pyriminobac, pyrimisulfan, pyriothiobac-sodium, pyroxasulfone, quinclorac, quinmerac, quinclamine, quinonamid, quizalofop, quizalofop-P-ethyl, rhodethanil, rimsulfuron, saflufenacil, S-metolachlor, sebuthylazine, sebumeton, sethoxydim, siduron, simazine, simeton, simetryn, SMA, sodium arsenite, sodium azide, sodium chlorate, sulcotrione, sulfallate, sulfentrazone, sulfometuron, sulfosate, sulfosulfuron, sulfuric acid, sulglycapin, swep, TCA, tebutam, tebuthiuron, tefuryltrione, tembotrione, tepraloxym, terbacil, terbucarb, terbutachlor, terbuturon, terbuthylazine, terbutryn, tetrafluron, thenylchlor, thiazafluron, thiazopyr, thidiazimin, thidiazuron, thiencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thiobencarb, tiocarbamil, tioclorim, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron, triaziflam, tribenuron, tribenuron-methyl, tricamba, triclopyr choline salt, triclopyr esters and salts, tridiphane, trietazine, trifloxysulfuron, trifluralin, triflusulfuron, trifop, trifopside, trihydroxytriazine, trimeturon, tripropindan, tritosulfuron, vernolate, xylachlor and salts, esters, optically active isomers and mixtures thereof.

[0056] In some embodiments, the compositions described herein are employed in combination with one or more herbicide safeners, such as AD-67 (MON 4660), benoxacor, benthicarb, brassinolide, cloquintocet (mexyl), cyometrinil, daimuron, dichlorim, dicyclonon, dimepiperate, disulfoton, fenclorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, jiecaowan, jiecaoxi, mefenpyr-diethyl, mephentate, naphthalic anhydride (NA), oxabetrinil, R29148, 1-[4-(*N*-(2-methoxybenzoyl)sulfamoyl)phenyl]-3-methylurea, *N*-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide and *N*-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity. In some embodiments, the safeners are employed in rice, cereal, corn, or maize settings. In some embodiments, the safener is cloquintocet or an ester or salt thereof. In certain embodiments, cloquintocet is utilized to antagonize harmful effects of the compositions on rice and cereals. In some embodiments, the safener is cloquintocet (mexyl).

[0057] In some embodiments, compositions provided herein further comprise at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such mixtures can

be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water-dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions. They can also be provided as a pre-mix or tank mixed.

[0058] Suitable agricultural adjuvants and carriers include, but are not limited to, crop oil concentrate; nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C₉-C₁₁ alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C₁₂-C₁₆) ethoxylate; di-*sec*-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

[0059] Liquid carriers that can be employed include water and organic solvents. The organic solvents include, but are not limited to, petroleum fractions or hydrocarbons such as mineral oil, aromatic solvents, paraffinic oils, and the like; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; esters of the above vegetable oils; esters of monoalcohols or dihydric, trihydric, or other lower polyalcohols (4-6 hydroxy containing), such as 2-ethyl hexyl stearate, *n*-butyl oleate, isopropyl myristate, propylene glycol dioleate, di-octyl succinate, di-butyl adipate, di-octyl phthalate and the like; esters of mono, di and polycarboxylic acids and the like. Specific organic solvents include, but are not limited to toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, *N*-methyl-2-pyrrolidinone, *N,N*-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. In certain embodiments, water is the carrier for the dilution of concentrates.

[0060] Suitable solid carriers include but are not limited to talc, pyrophyllite clay, silica, attapulugus clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, cellulose, and the like.

[0061] In some embodiments, the compositions described herein further comprise one or more surface-active agents. In some embodiments, such surface-active agents are employed in both solid and liquid compositions, and in certain embodiments those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Surfactants which may also be used in the present formulations are described, *inter alia*, in “McCutcheon’s Detergents and Emulsifiers Annual,” MC Publishing Corp., Ridgewood, New Jersey, 1998 and in “Encyclopedia of Surfactants,” Vol. I-III, Chemical Publishing Co., New York, 1980-81. Surface-active agents include, but are not limited to salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C₁₈ ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C₁₆ ethoxylate; soaps, such as sodium stearate; alkyl naphthalene-sulfonate salts, such as sodium dibutyl naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono and dialkyl phosphate esters; vegetable or seed oils such as soybean oil, rapeseed/canola oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils, and in certain embodiments, methyl esters.

[0062] In some embodiments, these materials, such as vegetable or seed oils and their esters, can be used interchangeably as an agricultural adjuvant, as a liquid carrier or as a surface active agent.

[0063] Other exemplary additives for use in the compositions provided herein include but are not limited to compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

[0064] In some embodiments, the concentration of the active ingredients in the compositions described herein is from about 0.0005 to 98 percent by weight. In some embodiments, the concentration is from about 0.0006 to 90 percent by weight. In compositions designed to be employed as concentrates, the active ingredients, in certain embodiments, are present in a concentration from about 0.1 to 98 weight percent, and in certain embodiments about 0.5 to 90 weight percent. Such compositions are, in certain embodiments, diluted with an inert carrier, such as water, before application. The diluted compositions usually applied to weeds or the locus of weeds contain, in certain embodiments, about 0.0003 to 1.5 weight percent active ingredient and in certain embodiments contain about 0.0008 to 1.0 weight percent.

[0065] The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation or paddy water, and by other conventional means known to those skilled in the art.

[0066] The described embodiments and following examples are for illustrative purposes and are not intended to limit the scope of the claims. Other modifications, uses, or combinations with respect to the compositions described herein will be apparent to a person of ordinary skill in the art without departing from the spirit and scope of the claimed subject matter.

EXAMPLES

[0067] Results in Tables 1-8 are greenhouse trial results for foliar applied compositions. The compositions tested, application rates employed, plant species tested, and results are given in Tables 1-8.

[0068] The following abbreviations are used in Tables 1-8:

CHEAL	<i>Chenopodium album</i> L.	common lambsquarters
CIRAR	<i>Cirsium arvense</i> (L.) Scop.	Canada thistle
GALAP	<i>Galium aparine</i> L.	catchweed bedstraw
HORVS	<i>Hordeum vulgare</i> L.	spring barley
KCHSC	<i>Kochia scoparia</i> (L.) Schrad.	kochia
LAMPU	<i>Lamium purpureum</i> L.	purple deadnettle
MATCH	<i>Matricaria recutita</i> L.	wild chamomile
PAPRH	<i>Papaver rhoeas</i> L.	common poppy
SASKR	<i>Salsola tragus</i> L.	Russian thistle

SINAR	<i>Sinapis arvensis</i> L.	wild mustard
TRZAS	<i>Triticum aestivum</i> (spring)	spring wheat
VERPE	<i>Veronica persica</i> Poir.	Persian speedwell
VIOTR	<i>Viola tricolor</i> L.	wild violet

g /ha = grams acid equivalent per hectare (g ae/ha) for Cmpd 1 and grams active ingredient per hectare (g ai/ha) for diflufenican, picolinafen, BACARA® and BACARA® FORTE

Obs = observed value

Exp = expected value as calculated by the equations set forth above in paragraph [0024].

Cmpd 1 = benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropyridine-2-carboxylate

Table 1: Synergistic activity of foliar-applied combination of Cmpd 1 and diflufenican

Application Rate (g/ha)		TRZAS		HORVS		SINAR		LAMPUI		GALAP		CHEAL	
Cmpd 1	Diflufenican	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	0	-	0	-	60	-	80	-	60	-	50	-
5	0	5	-	0	-	70	-	80	-	60	-	60	-
0	50	5	-	5	-	30	-	0	-	0	-	10	-
2.5	50	15	5	10	5	90	72	87	80	70	60	80	55
5	50	30	10	15	5	97	79	90	80	70	60	90	64

Table 2: Synergistic activity of foliar-applied combination of Cmpd 1 and diflufenican

Application Rate (g/ha)		KCHSC		SASKR		VERPE		VIOTR		MATCH		PAPRH		CIRAR	
Cmpd 1	Diflufenican	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	20	-	10	-	0	-	10	-	0	-	20	-	10	-
5	0	40	-	60	-	5	-	10	-	10	-	70	-	20	-
0	50	40	-	70	-	10	-	15	-	0	-	0	-	0	-
2.5	50	95	52	85	73	30	10	25	24	0	0	30	20	50	10
5	50	90	64	85	88	65	15	50	24	20	10	85	70	70	20

Table 3: Synergistic activity of foliar-applied combination of Cmpd 1 and picolinafen

Application Rate (g/ha)		TRZAS		HORVS		SINAR		LAMPUP		CHEAL	
Cmpd 1	Picolinafen	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	0	-	0	-	60	-	80	-	50	-
5	0	5	-	0	-	70	-	80	-	60	-
0	50	10	-	10	-	70	-	0	-	70	-
2.5	50	15	10	10	10	93	88	90	80	95	85
5	50	15	15	0	10	95	91	95	80	97	88

Table 4: Synergistic activity of foliar-applied combination of Cmpd 1 and picolinafen

Application Rate (g/ha)		KCHSC		SASKR		VERPE		VIOTR		MATCH		CIRAR	
Cmpd 1	Picolinafen	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	20	-	10	-	0	-	10	-	0	-	10	-
5	0	40	-	60	-	5	-	10	-	10	-	20	-
0	50	20	-	30	-	10	-	10	-	0	-	15	-
2.5	50	60	36	75	37	30	10	20	19	20	0	60	24
5	50	70	52	85	72	40	15	40	19	30	10	70	32

Table 5: Synergistic activity of foliar-applied combination of Cmpd 1 and BACARA® (flurtamone+diflufenican)

Application Rate (g/ha)		TRZAS		HORVS		SINAR		LAMPU		CHEAL		KCHSC	
Cmpd 1	BACARA®	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	0	-	0	-	60	-	80	-	50	-	20	-
5	0	5	-	0	-	70	-	80	-	60	-	40	-
0	175	15	-	0	-	20	-	0	-	10	-	60	-
2.5	175	10	15	0	0	95	68	95	80	95	55	93	68
5	175	15	19	0	0	97	76	97	80	97	64	97	76

Table 6: Synergistic activity of foliar-applied combination of Cmpd 1 and BACARA® (flurtamone+diflufenican)

Application Rate (g/ha)		SASKR		VERPE		VIOTR		PAPRH		STEME		CIRAR	
Cmpd 1	BACARA®	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	10	-	0	-	10	-	20	-	20	-	10	-
5	0	60	-	5	-	10	-	70	-	30	-	20	-
0	175	65	-	10	-	10	-	0	-	0	-	15	-
2.5	175	80	69	50	10	20	19	50	20	30	20	50	24
5	175	95	86	30	15	30	19	85	70	40	30	90	32

A rate of 175 g/ha of BACARA® = 50 g ai/ha of diflufenican + 125 g ai/ha of flurtamone.

Table 7: Synergistic activity of foliar-applied combination of Cmpd 1 and BACARA® FORTE (diflufenican + flufenacet+ flurtamone)

Application Rate (g/ha)		TRZAS		HORVS		SINAR		LAMPU		CHEAL		KCHSC	
Cmpd 1	BACARA FORTE	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	0	-	0	-	60	-	80	-	50	-	20	-
5	0	5	-	0	-	70	-	80	-	60	-	40	-
0	180	10	-	0	-	50	-	10	-	65	-	80	-
2.5	180	10	10	10	0	100	80	90	82	90	83	90	84
5	180	0	15	0	0	100	85	95	82	93	86	95	88


Table 8: Synergistic activity of foliar-applied combination of Cmpd 1 and BACARA® FORTE (diflufenican + flufenacet+ flurtamone)

Application Rate (g/ha)		VERPE		VIOTR		PAPRH		STEME		CIRAR	
Cmpd 1	BACARA® FORTE	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	0	-	10	-	20	-	20	-	10	-
5	0	5	-	10	-	70	-	30	-	20	-
0	180	35	-	40	-	0	-	10	-	10	-
2.5	180	60	35	60	46	87	20	70	28	85	19
5	180	70	38	60	46	93	70	80	37	90	28

A rate of 180 g/ha of BACARA® FORTE = 60 g ai/ha diflufenican + 60 g ai/ha flufenacet + 60 g ai/ha flurtamone

CLAIMS:

1. A synergistic herbicidal composition comprising a herbicidally effective amount of (a) a benzyl ester of the compound of formula (I)




and (b) picolinafen or diflufenican, or flurtamone and diflufenican.

2. The composition of claim 1, further comprising (c) flufenacet.
3. The composition of claim 1 or 2, wherein (b) is flurtamone and diflufenican.
4. The composition of claim 1 or 2, wherein (b) is picolinafen.
5. The composition of claim 1 or 2, wherein (b) is diflufenican.
6. The composition of claim 1 or 2, further comprising a herbicide safener.
7. The composition of claim 1, wherein (b) is picolinafen, and the weight ratio of (a) to (b) is from about 1.0 to about 40 of (a) to about 1.25 to about 100 of (b).
8. The composition of claim 1, wherein (b) is picolinafen, and the weight ratio of (a) to (b) is from about 1.0 to about 40 of (a) to about 12.5 to about 100 of (b).
9. The composition of claim 1, wherein (b) is diflufenican, and the weight ratio of (a) to (b) is from about 1.0 to about 40 of (a) to about 1.25 to about 100 of (b).
10. The composition of claim 1, wherein (b) is diflufenican, and the weight ratio of (a) to (b) is from about 1.0 to about 40 of (a) to about 12.5 to about 100 of (b).

11. The composition of claim 2, wherein (b) is diflufenican, and the weight ratio of (a) to (b) to (c) is from about 1.0 to about 40 of (a) to about 1.25 to about 100 of diflufenican to about 15 to about 240 of (c) flufenacet.
12. The composition of claim 2, wherein (b) is diflufenican, and the weight ratio of (a) to (b) to (c) is from about 1.0 to about 40 of (a) to about 12.5 to about 100 of diflufenican to about 15 to about 240 of (c) flufenacet.
13. The composition of claim 1, wherein (b) is flurtamone and diflufenican, and the weight ratio of (a) to flurtamone to diflufenican is from about 1.0 to about 40 of (a) to about 30 to about 250 of flurtamone to about 1.25 to about 100 of diflufenican.
14. The composition of claim 1, wherein (b) is flurtamone and diflufenican, and the weight ratio of (a) to flurtamone to diflufenican is from about 1.0 to about 40 of (a) to about 30 to about 250 of flurtamone to about 12.5 to about 100 of diflufenican.
15. The composition of claim 2, wherein (b) is flurtamone and diflufenican, and the weight ratio of (a) to flurtamone to diflufenican to (c) is from about 1.0 to about 40 of (a) to about 30 to about 250 of flurtamone to about 1.25 to about 100 of diflufenican to about 15 to about 240 of (c) flufenacet.
16. The composition of claim 2, wherein (b) is flurtamone and diflufenican, and the weight ratio of (a) to flurtamone to diflufenican to (c) is from about 1.0 to about 40 of (a) to about 30 to about 250 of flurtamone to about 12.5 to about 100 of diflufenican to about 15 to about 240 of (c) flufenacet.
17. The composition of any one of claims 1–16, which is synergistic as determined by the Colby equation.
18. A method of controlling undesirable vegetation which comprises applying to an area where control is desired a herbicidally effective amount of the composition of any one of claims 1–17.

19. A method of controlling undesirable vegetation which comprises applying to an area where control is desired a herbicidally effective amount of a combination comprising (a) a benzyl ester of a compound of formula (I)



and (b) picolinafen or diflufenican, or flurtamone and diflufenican.

20. The method of claim 19, further comprising (c) flufenacet.

21. The method of claim 19 or 20, wherein the undesirable vegetation is controlled in rice, wheat, triticale, barley, oats, rye, corn, maize, cereals, pastures, grasslands, rangelands, fallowland, and, industrial vegetation management or rights-of-way.

22. The method of any one of claims 19–21, wherein the undesirable vegetation is immature.

23. The method of any one of claims 19–21, wherein the combination is applied pre-emergently.

24. The method of any one of claims 19–21, wherein the combination is applied post-emergently.

25. The method of any one of claims 19–21, wherein the undesirable vegetation is controlled in a glyphosate-, glufosinate-, dicamba-, phenoxy auxin-, pyridyloxy auxin-, aryloxyphenoxypropionate-, acetyl CoA carboxylase (ACCase) inhibitor-, imidazolinone-, acetolactate synthase (ALS) inhibitor-, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitor-, protoporphyrinogen oxidase (PPO) inhibitor-, triazine-, or bromoxynil- tolerant crop.

