



US 20050153330A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0153330 A1**

Bass et al.

(43) **Pub. Date: Jul. 14, 2005**

(54) **NOVEL DKR POLYPEPTIDES**

Related U.S. Application Data

(75) Inventors: **Michael Brian Bass**, Thousand Oaks, CA (US); **John Kevin Sullivan**, Newbury Park, CA (US); **Lars Eyde Theill**, Thousand Oaks, CA (US); **Daguang Wang**, Thousand Oaks, CA (US)

(60) Continuation of application No. 09/976,736, filed on Oct. 9, 2001, now abandoned, which is a division of application No. 09/161,241, filed on Sep. 25, 1998, now Pat. No. 6,344,541.

Publication Classification

Correspondence Address:
AMGEN INC.
LAW DEPARTMENT
1201 AMGEN COURT WEST
SEATTLE, WA 98119 (US)

(51) **Int. Cl.⁷** **C12Q 1/68**; C07H 21/04;
C07K 14/705
(52) **U.S. Cl.** **435/6**; 435/69.1; 435/320.1;
435/325; 530/350; 536/23.5

(73) Assignee: **Amgen Inc.**

(57) **ABSTRACT**

(21) Appl. No.: **10/998,271**

Disclosed are nucleic acid molecules encoding novel DKR polypeptides. Also disclosed are methods of preparing the nucleic acid molecules and polypeptides, and methods of using these molecules.

(22) Filed: **Nov. 24, 2004**

FIG. 1

1 ATGCAGCGGC TCGGGGGTAT TTTGCTGTGT AACTGCTGG CGGCGCGGT
51 CCCCACTGCT CCTGCTCCTT CCCCACGGT CACTTGGACT CCGCGGAGC
101 CGGGCCCAGC TCTCAACTAC CCTCAGGAGG AAGCTACGCT CAATGAGATG
151 TTTCGAGAGG TGGAGGAGCT GATGGAAGAC ACTCAGCACA AACTGCGCAG
201 TGCCGTGGAG GAGATGGAGG CGGAAGAAGC AGCTGCTAAA ACGTCCCTCTG
251 AGGTGAACCT GGCAAGCTTA CCTCCCACT ATCACAATGA GACCAGCACG
301 GAGACCCAGG TGGGAAATAA CACAGTCCAT GTGCACCAGG AAGTTCACAA
351 GATAACCAAC AACCAGAGTG GACAGGTGGT CTTTCTGAG ACAGTCATTA
401 CATCTGTAGG GGATGAAGAA GGCAAGAGGA GCCATGAATG TATCATTTGAT
451 GAAGACTGTG GGCCACCAG GTACTGCCAG TTCTCCAGCT TCAAGTACAC
501 CTGCCAGCCA TGCCGGGACC AGCAGATGCT ATGCACCCGA GACAGTGAGT
551 GCTGTGGAGA CCAGCTGTGT GCCTGGGGTC ACTGCACCCA AAAGGCCACC
601 AAAGGTGGCA ATGGGACCAT CTGTGACAA CAGAGGGATT GCCAGCCTGG
651 CCTGTGTTGT GCCTTCCAAA GAGGCCCTGCT GTTCCCCGTG TGCACACCCC
701 TGCCCGTGGA GGGAGAGCTC TGCCATGACC CCACCAGCCA GCTGCTGGAT
751 CTCATCACCT GGGAACTGGA GCCTGAAGGA GCTTTGGACC GATGCCCCCTG
801 CGCCAGTGGC CTCCTATGCC AGCCACACAG CCACAGTCTG GTGTACATGT
851 GCAAGCCAGC CTTCTGTGGC AGCCATGACC ACAGTGAGGA GAGCCAGCTG
901 CCCAGGGAGG CCCC GGATGA GTACGAAGAT GTTGGCTTCA TAGGGGAAGT
951 GCGCCAGGAG CTGGAAGACC TGGAGCGGAG CCTAGCCCCAG GAGATGGCAT
1001 TTGAGGGGCC TGCCCTGTG GAGTCACTAG GCGGAGAGGA GGAGATTAG

FIG. 2

1 ATGCAGCGGC TTGGGGCCAC CCTGCTGTGC CTGCTGTCTG CGGGGGCGGT
51 CCCACGGCC CCCGCGCCG CTCCGACGGC GACCTCGGCT CCAGTCAAGC
101 CCGGCCCGGC TCTCAGCTAC CCGCAGGAGG AGGCCACCCCT CAATGAGATG
151 TTCCGCGAGG TTGAGGAACT GATGGAGGAC ACGCAGCACA AATTGCGCAG
201 CGCGGTGGAA GAGATGGAGG CAGAAGAAGC TGCTGCTAAA GCATCATCAG
251 AAGTGAACCT GGCAAACTTA CCTCCAGCT ATCACAATGA GACCAACACA
301 GACACGAAGG TTGGAAATAA TACCATCCAT GTGCACCGAG AAATTCACAA
351 GATAACCAAC AACCAGACTG GACAAATGGT CTTTTCAGAG ACAGTTATCA
401 CATCTGTGG AGACGAAGAA GGCAGAAGGA GCCACGAGTG CATCATCGAC
451 GAGGACTGTG GGCCAGCAT GTACTGCCAG TTTGCCAGCT TCCAGTACAC
501 CTGCCAGCCA TGCCGGGGCC AGAGGATGCT CTGCACCCGG GACAGTGAGT
551 GCTGTGGAGA CCAGCTGTGT GTCTGGGGTC ACTGCACCAA AATGGCCACC
601 AGGGCAGCA ATGGACCAT CTGTGACAA CAGAGGACT GCCAGCCGGG
651 GCTGTGCTGT GCCTTCCAGA GAGGCCCTGCT GTTCCCTGTG TGCACACCCC
701 TGCCCCGTGA GGGCGAGCTT TGCCATGACC CCGCCAGCCG GCTTCTGGAC
751 CTCATCACCT GGGAGCTAGA GCCTGATGGA GCCTTGACC GATGCCCTTG
801 TGCCAGTGGC CTCCTCTGCC AGCCCCACAG CCACAGCCTG GTGTATGTGT
851 GCAAGCCGAC CTTCTGTGGG AGCCGTGACC AAGATGGGA GATCCTGCTG
901 CCCAGAGAGG TCCCCGATGA GTATGAAGTT GGCAGCTTCA TGGAGGAGGT
951 GCGCCAGGAG CTGGAGGACC TGGAGAGGAG CCTGACTGAA GAGATGGCGC
1001 TGGGGGAGCC TGGGGCTGCC GCCGCTGCAC TGCTGGGAGG GGAAGAGATT
1051 TAG

FIG. 3

1 ATGATGGCTC TGGGCGCAGC GGGAGCTACC CGGGTCTTTG TCGCGATGGT
51 AGCGCGGCT CTCGGCGGC ACCCTCTGCT GGGAGTGAGC GCCACCTTGA
101 ACTCGGTTCT CAATTCCAAC GCTATCAAGA ACCTGCCCCC ACCGCTGGGC
151 GCGCTGCGG GGCACCCAGG CTCTGCAGTC AGCGCCGCGC CGGGAATCCT
201 GTACCCGGGC GGAATAAGT ACCAGACCAT TGACAACTAC CAGCCGTACC
251 CGTGCGCAGA GGACGAGGAG TCGGCACTG ATGAGTACTG CGTAGTCCC
301 ACCCGCGGAG GGGACGCGG CGTGCAATC TGTCTCGCCT GCAGGAAGCG
351 CCGAAAACGC TGCAATGCGTC ACGCTATGTG CTGCCCCGGG AATTACTGCA
401 AAAATGGAAT ATGTGTGTCT TCTGATCAA ATCATTTCCG AGGAGAAATT
451 GAGGAAACCA TCACTGAAAG CTTTGGTAAT GATCATAGCA CCTTGGATGG
501 GTATTCCAGA AGAACCACCT TGTCTTCAA AATGTATCAC ACCAAAGGAC
551 AAGAAGGTTT TGTTTGTCTC CGGTCATCAG ACTGTGCCCTC AGGATTGTGT
601 TGTGCTAGAC ACTTCTGGTC CAAGATCTGT AAACCTGTCC TGAAGAAGG
651 TCAAGTGTGT ACCAAGCATA GGAGAAAAGG CTCTCATGGA CTAGAAATAT
701 TCCAGCGTTG TTAATGTGGA GAAGTCTGT CTTGCCGGAT ACAGAAAGAT
751 CACCATCAAG CCAGTAATTC TTCTAGGCTT CACACTTGTC AGAGACACTA
801 A

FIG. 4

```
1  ATGGCCGCGC TGATGCGGGT CAAGGATTCA TCCC GCTGCC TTCTCCTACT
51  GGCCGCGGTG CTGATGGTGG AGAGCTCACA GCTAGGCAGC TCGCGGGCCA
101 AACTCAACTC CATCAAGTCC TCCTAGGAG GGGAGACTCC TGCTCAGTCA
151 GCCAACCGAT CTGCAGGCAT GAACCAAGGA CTGGCTTTCG GCGGCAGTAA
201 GAAGGGCAA AGCCTGGGC AGCCTACCC TTGCAGCAGT GATAAGGAAT
251 GTGAAGTTGG AAGATACTGC CACAGTCCCC ACCAAGGATC ATCAGCCTGC
301 ATGCTCTGTA GGAGGAAAA GAAACGATGC CACAGAGATG GGATGTGTG
351 CCCTGGTACC CGCTGCAATA ATGGAATCTG CATCCCAGTC ACTGAGAGCA
401 TCCTCACCCC ACATATCCCA GCTCTGGATG GCACCCGGCA TAGAGATCGC
451 AACCATGGTC ACTATTCCAA CCATGACCTG GGATGGCAGA ATCTAGGAAG
501 GCCACACTCC AAGATGCCCTC ATATAAAGG ACATGAAGGA GACCCATGCC
551 TACGGTCATC AACTGCATT GATGGGTTT GTTGTGCTCG CCACTTCTGG
601 ACCAAAATCT GCAAACCAGT GCTCCATCAG GGGGAAGTCT GTACCAACA
651 ACGCAAGAAG GGTTCGCACG GGCTGGAGAT TTTC CAGAGG TGTGACTGTG
701 CAAAGGGCCT GTCCTGCAA GTGTGGAAG ATGCCACCTA CTC TTCCAAA
751 GCCAGACTCC ATGTATGCCA GAAGATCTGA
```

FIG. 5

1 ATGGCCGCGT TGATGCCGAG CAAGGATTCTG TCCTGCTGCC TGCTCCTACT
51 GGCCGCGGTG CTGATGGTGG AGAGCTCACA GATCGGCAGT TCGCGGGCCA
101 AACTCAACTC CATCAAGTCC TCTCTGGGCG GGGAGACGCC TGGTCAGGCC
151 GCCAATCGAT CTGCGGGCAT GTACCAAGGA CTGGCATTCTG GCGGCAGTAA
201 GAAGGGCAAA AACCTGGGCG AGGCTACCC TTGTAGCAGT GATAAGGAGT
251 GTGAAGTTGG GAGGTATTGC CACAGTCCCC ACCAAGGATC ATCGGCCTGC
301 ATGGTGTGTC GGAGAAAAAA GAAGCGCTGC CACCGAGATG GCATGTGCTG
351 CCCCAGTACC CGCTGCAATA ATGGCATCTG TATCCCAGTT ACTGAAAGCA
401 TCTTAACCC TCACATCCCG GCTCTGGATG GTACTCGGCA CAGAGATCGA
451 AACCACGGTC ATTACTCAA CCATGACTTG GGATGGCAGA ATCTAGGAAG
501 ACCACACACT AAGATGTCAC ATATAAAAGG GCATGAAGGA GACCCCTGCC
551 TACGATCATC AGACTGCATT GAAGGGTTT GCTGTGCTCG TCATTCTGG
601 ACCAAAATCT GCAAACCAGT GCTCCATCAG GGGGAAGTCT GTACCAAACA
651 ACGCAAGAAG GGTCTCATG GGCTGGAAT TTTCCAGCGT TCGGACTGTG
701 CGAAGGGCCT GTCTTGCAA GTATGGAAG ATGCCACCTA CTCCTCCAAA
751 GCCAGACTCC ATGTGTGTCA GAAAATTGA

FIG. 6

1 ATGGCCGCGT TGATGCGGAG CAAGGATTCG TCCTGCTGCC TGCTCCTACT
51 GGCCGCGGTG CTGATGGTGG AGAGCTCACA GATCGGCAGT TCGCGGGCCA
101 AACTCAACTC CATCAAGTCC TCTCTGGGCG GGGAGACGCC TGGTCAGGCC
151 GCCAATCGAT CTGCGGGCAT GTACCAAGGA CTGGCATTCG GCGGCAGTAA
201 GAAGGGCAA AACCTGGGCG AGGCCCTACCC TTGTAGCAGT GATAAGGAGT
251 GTGAAGTTGG GAGGTATTGC CACAGTCCCC ACCAAGGATC ATCGGCCTGC
301 ATGGTGTGTC GGAGAAAAA GAAGCGCTGC CACCGAGATG GCATGTGCTG
351 CCCAGTACC CGCTGCAATA ATGGGCATGA AGGAGACCCC TGCCCTACGAT
401 CATCAGACTG CATTGAAGGG TTTTGCTGTG CTCGTCAATT CTGGACCAA
451 ATCTGCAAAC CAGTGCTCCA TCAGGGGGA GTCTGTACCA AACAAACGCA
501 GAAGGGTTCT CATGGGCTGG AAATTTTCCA GCGTTGCGAC TGTGCCAAGG
551 GCGTGTCTTG CAAAGTATGG AAAGATGCCA CCTACTCCTC CAAAGCCAGA
601 CTCCATGTGT GTCAGAAAAAT TTGA

FIG. 7

```
1  ATGGTGGCGG CCGTCCTGCT GGGGCTGAGC TGGCTCTGCT CTCCCCTGGG
51  AGCTCTGGTC CTGGACTTCA ACAACATCAG GAGCTCTGCT GACCTGCATG
101  GGGCCCGGAA GGGCTCACAG TGCCTGTCTG ACACGGACTG CAATACCAGA
151  AAGTCTGCC TCCAGCCCCG CGATGAGAAG CCGTCTGTG CTACATGTCG
201  TGGGTTGCCG AGGAGGTGCC AGCGAGATGC CATGTGCTGC CCTGGGACAC
251  TCTGTGTGAA CGATGTTTGT ACTACGATGG AAGATGCAAC CCCAATATTA
301  GAAAGGCAGC TTGATGAGCA AGATGGCACA CATGCAGAAG GAACAACCTGG
351  GCACCCAGTC CAGGAAAAACC AACCCAAAAG GAAGCCAAGT ATTAAGAAAT
401  CACAAGGCAG GAAGGGACAA GAGGGAGAAA GTTGTCTGAG AACTTTTGAC
451  TGTGGCCCTG GACTTTGCTG TGCTCGTCAT TTTTGGACGA AAATTGTAA
501  GCCAGTCCTT TTGGAGGGAC AGGCTGCTC CAGAAGAGGG CATAAAGACA
551  CTGCTCAAGC TCCAGAAATC TTCCAGCGTT GCGACTGTGG CCTGGACTA
601  CTGTGTCGAA GCCAATTGAC CAGCAATCGG CAGCATGCTC GATTAAGAGT
651  ATGCCAAAAA ATAGAAAAGC TATAA
```


FIG. 8

1	MQRLGGILLC	TLAAAVPTA	PAPSTVTWT	PAEPGPAINY	PQEEATLNEM
51	FREVEEELMED	TQHKLRSAVE	EMEAEEAAAK	TSSEVNILASL	PPNYHNETST
101	ETRVGNNTVH	VHQEVHKITN	NQSGQVVFSE	TVITSVGDEE	GKRSHECIID
151	EDCGPTRYCQ	FSSFKYTCQP	CRDQQMLCTR	DSECCGDQLC	AWGHCTQKAT
201	KGGNGTICDN	QRDCQPGLCC	AFQRGLLFPV	CTPLPVEGEL	CHDPTSQLLD
251	LITWELEPEG	ALDRCPCASG	LLCQPHSHSL	VYMCKPAFVG	SHDHSEESQL
301	PREAPDEYED	VGFIGEVRQE	LEDLERSLAQ	EMAFEGPAPV	ESLGEEEEI*

FIG. 9

1	MQRLGATLLC	LLLAADVPTA	PAPAPTATSA	PVKPGPALS	PQEEATLNEM
51	FREVEELMED	TQHKLRSAVE	EMEAEEAAAK	ASSEVNLANL	PPSYHNETNT
101	DTKVGNNNTIH	VHREIHKITN	NQTGQMVFESE	TVITSVGDEE	GRRSHECIID
151	EDCGPSMYCQ	FASFQYTCQP	CRGQRLCTR	DSECCGDQLC	VWGHCTKMAT
201	RGSNGTICDN	QRDCQPGGCC	AFQRGLLFPV	CTPLPVEGEL	CHDPASRLLD
251	LITWELEPDG	ALDRCPCASG	LLCQPHSHSL	VYVCKPTFVG	SRDQDGEILL
301	PREVPDEYEV	GSFMEEVRQE	LEDLERSLTE	EMALGEPAAA	AAALLGGEET
351	*				

FIG. 10

1	MMALGAAGAT	RVFVAMVAAA	LGGHPLLGV	ATLNSVLNSN	AIKNLPPPLG
51	GAAGHPGSAV	SAAPGILYPG	GNKYQTIDNY	QYPYCAEDEE	CGTDEYCASP
101	TRGGDAGVQI	CLACRKRKR	CMRHAMCCPG	NYCKNGICVS	SDQNHFRGEI
151	EETITESFGN	DHSTLDGYSR	RTTLSSKMYH	TKGQEGSVCL	RSSDCASGLC
201	CARHFWSKIC	KPVLKEGQVC	TKHRRKGSYG	LEIFQRCYCG	EGLSCRIQKD
251	HHQASNSSRL	HTCQRH*			

FIG. 11

1	MAALMRVKDS	SRCLLLLA	AV	LMV	ESSQLGS	SRAKLN	SIKS	SLGGETPAQS
51	ANRSAGMNQG	LAFGGSKKGK	SLGQAYPCSS	DKECEVGRYC	HSPHQGSSAC			
101	MLCRRKKKRC	HRDGMCCPGT	RCNNGICIPV	TESILTPHIP	ALDGT	RRDR		
151	NHGHYSNHD	L	GWQNLGRPHS	KMPHIKGHEG	DPCLRSSDCI	DGFCCARHFW		
201	TKICKPVLHQ	GEVCTKQ	RKK	GSHGLEIFQR	CDCAKGLSCK	VWKDATYSSK		
251	ARLHVCQKI*							

FIG. 12

1	MAALMRSKDS	SCCLLLAAV	LMVSSQIGS	SRAKLNSIKS	SLGGETPGQA
51	ANRSAGMYQG	LAFGGSKKGK	NLGQAYPCSS	DKECEVGRYC	HSPHQGSSAC
101	MVCCRKKKRC	HRDGMCCPST	RCNNGICIPV	TESILTPHIP	ALDGTTRHRDR
151	NHGHYSNHDL	GWQNLGRPHT	KMSHIKGHEG	DPCLRSSDCI	EGFCCARHEW
201	TKICKPVLHQ	GEVCTKQRRK	GSHGLEIFQR	CDCAKGLSCK	VWKDATYSSK
251	ARLHVCQKI*				

FIG. 13

1	MAALMRKDS	SCCLLLAAV	LMVSSQIGS	SRAKLNSIKS	SLGGETPGQA
51	ANRSAGMYQG	LAFGGSKKGK	NLGQAYPCSS	DKECEVGRYC	HSPHQGSSAC
101	MVCRRKKKRC	HRDGMCCPST	RCNNGHEGDP	CLRSSDCIEG	FCCARHFWTK
151	ICKPVLHQGE	VCTKQRRKGS	HGLEIFQRC	CAKGLSCKVW	KDATYSSKAR
201	LHVCQKI*				

FIG. 14

1 MVAAVLLGLS WLCSP LGALV LDFENNIRSSA DLHGARKGSQ CLSDTDCNTR
51 KFCLQPRDEK PFCATCRGLR RRCQRDAMCC PGTL CVNDVC TTMEDATPIL
101 ERQLDEQDGT HAEGTTGHPV QENQPKRKPS IKKSQGRKGQ EGESCLRTFD
151 CGPGLCCARH FWTKICKPVL LEGQVCSRRG HKDTAQAPEI FQRCD CGPGL
201 LCRSQLTSNR QHARLRVCQK IEKL*

FIG. 15A




FIG. 15B




FIG. 15C




FIG. 15D




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20

1 2 3 4 5 6 7 8 9 10




FIG. 21




FIG. 22

1 ATGATGGCTC TGGGTGCTGC TGGTGCTACC CGTGTTTTCG TTGCTATGGT
51 TGCTGCTGCT CTGGGTGGTC ACCCGCTGCT GGGTGTTTCC GCTACCCCTGA
101 ACTCCGTTCT GAACTCCAAC GCTATCAAAA ACCTGCCGCC GCCGCTGGGT
151 GGTGCTGCTG GTCACCCGGG TTCCGCTGTT TCCGCTGCTC CGGGTATCCT
201 GTACCCGGGT GGTAACAAAT ACCAGACCAT CGACAATAC CAGCCGTACC
251 CGTGCGCTGA AGACGAAGAA TGGGTACCG ACGAATACTG CGCTTCCCCG
301 ACCCGTGGTG GTGACGCTGG TGTTCAGATC TGCCCTGGCTT GCCGTAAACG
351 TCGTAAACGT TGCATGCGTC ACGCTATGTG CTGCCCGGGT AACTACTGCA
401 AAAACGGTAT CTGCGTTTCC TCCGACCAGA ACCACTTCCG TGGTGAATC
451 GAAGAAACCA TCACCCGAATC CTTCCGGTAA GACCACCTCCA CCTGGACGG
501 TTACTCCCGT CGTACCACCC TGTCCCTCAA AATGTACCAC ACCAAAGGTC
551 AGGAAGGTTT CGTTTGCCCTG CGTTCCTCCG ACTGCGCTTC CGGTCGTGC
601 TGGGCTCGTC ACTTCTGGTC CAAAATCTGC AAACCGGTTT TGAAGAAGG
651 TCAGGTTTGC ACCAAACACC GTCGTAAAGG TTCCCACGGT CTGGAATCT
701 TCCAGCGTTG CTA CTGCGGT GAGGTCTGT CCTGCCGTAT CCAGAAAGAC
751 CACCACCAGG CTTCCAATC CTCCCCTCTG CACACCTGCC AGCGTCAC

FIG. 23

1 ATGGCTGCTC TGATGCGTTC CAAAGACTCC TCCTGCTGCC TGCTGCTGCT
51 GGCTGCTGTT CTGATGGTTG AATCCTCCCA GATCGGTTCC TCCCGTGCTA
101 AACTGAACCTC CATCAAAATCC TCCCTGGGTG GTGAAACCCC GGGTCAGGCT
151 GCTAACCGTT CCGCTGGTAT GTACCAGGGT CTGGCTTTTCG GTGGTTCCAA
201 AAAAGGTAAA AACCTGGGTC AGGCTTACCC GTGCTCCTCC GACAAAGAAAT
251 GCGAAGTTGG TCGTTACTGC CACTCCCCGC ACCAGGGTTC CTCCGCTTGC
301 ATGGTTTGCC GTCGTAAAAA AAAACGTTGC CACCGTGACG GTATGTGCTG
351 CCCGTCCACC CGTTGCAACA ACGGTATCTG CATCCC GGTT ACCGAAATCCA
401 TCCTGACCCC GCACATCCCG GCTCTGGACG GTACCCCGTCA CCGTGACCGT
451 AACCAACGGTC ACTACTCCAA CCACGACCTG GGTTGGCAGA ACCTGGGTCG
501 TCCGCACACC AAAATGTCCC ACATCAAAGG TCACGAAGGT GACCCGTGCC
551 TCGGTTCCCTC CGACTGCATC GAAGGTTTCT GCTGCGCTCG TCACTTCTGG
601 ACCAAAATCT GCAAACCGGT TCTGCACCCAG GGTAAGTTT GCACCAACA
651 GCGTAAAAAA GGTTCCACG GTCTGGAAT CTTCACGCGT TGCGACTGCG
701 CTAAAGGTCT GTCCTGCAAA GTTTGGAAG ACGCTACCTA CTCCTCCAAA
751 GCTCGTCTGC ACGTTTGCCA GAAATC

FIG. 24

1 ATGCAGCGTC TGGGTGCTAC CCTGCTGTGC CTGCTGCTGG CTGCTGCTGT
51 TCCGACCGCT CCGGCTCCGG CTCCGACCGC TACCCTCCGT CCGTTAAAC
101 CGGTCCGGC TCTGTCCCTAC CCGCAGGAAG AAGCTACCCCT GAACGAAATG
151 TTCCGTGAAG TTGAAGAAGT GATGGAAGAC ACCAGCACAA AACTGCGTTC
201 CGCTGTTGAA GAAATGGAAG CTGAAGAAGC TGCTGCTAAA GCTTCCTCCG
251 AAGTTAACCT GGCTAACCTG CCGCGTCCT ACCACAACGA AACCAACACC
301 GACACCAAAG TTGGTAACAA CACCATCCAC GTTCACCCGTG AAATCCACAA
351 AATCACC AAC ACCAGACCG GTCAGATGGT TTTCCTCCGAA ACCGTTATCA
401 CCTCCGTTGG TGACGAAGAA GGTGTCGTT CCCACGAATG CATCATCGAC
451 GAAGACTGCG GTCCGTCCAT GTA CTGCCAG TTGCTTCCCT TCCAGTACAC
501 CTGCCAGCCG TGCCGTGGTC AGCGTATGCT GTGCACCCGT GACTCCGAAT
551 GCTGCGGTGA CCAGCTGTGC GTTTGGGGTC ACTGCACCAA AATGCTACC
601 CGTGGTTCCA ACGGTACCAT CTGCGACAAAC CAGCGTGAAT GCCAGCCGGG
651 TCTGTGCTGC GCTTTCCAGC GTGGTCTGCT GTTCCCGGTT TGCACCCCGC
701 TGCCGGTTGA AGGTGAATG TGCCACGACC CCGCTTCCCG TCTGCTGAC
751 CTGATCACCT GGAACCTGGA ACCGACGGT GCTCTGGACC GTTGCCCGTG
801 CGCTTCCGGT CTGCTGTGCC AGCCGCACTC CCACCTCCCTG GTTTACGTTT
851 GCAACCCGAC CTTCGTTGGT TCCCGTGACC AGGACGGTGA AATCCCTGCTG
901 CCGCGTGAAG TTCCGGACGA ATACGAAGTT GGTTCCTTCA TGAAGAAGT
951 TCGTCAGGAA CTGGAAGACC TGAACGTTT CCTGACCCGAA GAAATGGCTC
1001 TGGGTGAACC GGCTGCTGCT GCTGCTGCTC TGCTGGGTGG TGAAGAATC

FIG. 25

1 ATGGTTGCTG CTGTTCTGCT GGGTCTGTCC TGGCTGTGCT CCCCCTGGG
51 TGCTCTGGTT CTGGACTTCA ACAACATCCG TTCTTCCGCT GACCTGCACG
101 GTGCTCGTAA AGGTTCCAG TGCCTGTCCG ACACCGACTG CAACACCCGT
151 AAATTCTGCC TGCAGCCCGG TGACGAAAAA CCGTCTGCG CTACCTGCCG
201 TGGTCTGCGT CGTCGTTGCC AGCGTGACGC TATGTGCTGC CCGGTACCC
251 TGTGCCGTTAA CGACGTTTGC ACCACCATGG AAGACGCTAC CCCGATCCTG
301 GAACGTCAGC TGGACGAACA GGACGGTACC CACGCTGAAG GTACCACCCG
351 TCACCCCGTT CAGGAAAACC AGCCGAAACG TAAACCGTCC ATCAAAAAAT
401 CCCAGGGTCG TAAAGGTCAG GAAGGTGAAT CCTGCCCTGCG TACCTTCGAC
451 TCGGGTCCGG GTCTGTGCTG CGCTCGTCAC TTCTGGACCA AAATCTGCAA
501 ACCGGTTCTG CTGGAAGGTC AGGTTTGCTC CCGTCGTGGT CACAAGACA
551 CCGCTCAGGC TCCGGAAATC TTCCAGCGTT GCGACTGCGG TCCGGGTCTG
601 CTGTGCCGTT CCCAGCTGAC CTCCAACCGT CAGCACGCTC GTCTGCGTGT
651 TTGCCAGAAA ATCGAAAAAC TG

NOVEL DKR POLYPEPTIDES

[0001] This application is a continuation of U.S. Ser. No. 09/976,736, filed Oct. 9, 2001, which is a divisional of U.S. Ser. No. 09/161,241, filed Sep. 25, 1998, which are hereby incorporated by reference.

FIELD OF THE INVENTION

[0002] This invention relates generally to novel genes encoding proteins that have use as anti-cancer therapeutics.

BACKGROUND

[0003] Related Art

[0004] One of the hallmarks of cells that have become cancerous is the change in the gene expression pattern in those cells as compared to normal, non-cancerous cells. An intricate series of cell signaling events leads to this so called "differential gene expression", resulting in conversion of a normal cell to a cancer cell (also known as "oncogenesis" or "cell transformation"). A number of cell signaling pathways have been implicated in the process of cell transformation, such as, for example, the cadherin pathway, the delta/jagged pathway, the hedgehog/sonic hedgehog pathway, and the wnt/wingless pathway (Hunter, *Cell*, 88: 333-346 [1997]; Currie, *J. Mol. Med.*, 76: 421-433 [1998]; Peifer, *Science*, 275: 1752-1753 [1997]. Interestingly, these same pathways are involved in cell morphogenesis, or cell differentiation, during embryo development (Hunter, supra; Cadigan et al., *Genes and Develop.*, 11: 3286-3305 [1997]).

[0005] The wnt genes encode glycoproteins that are secreted from the cell. These glycoproteins are found in both vertebrate and invertebrate organisms. Currently, there are at least 20 wnt family members, and these members are believed to function variously in control of growth and in tissue differentiation. Recently, discovery of a novel gene was identified in *Xenopus* and mouse and has been termed dickkopf-1 ("dkk-1"). This gene is purportedly a potent antagonist of wnt-8 signaling (Glinka et al., *Nature*, 391: 357-362 [1998]). Interestingly, this gene is also purportedly involved in morphogenesis in the developing embryo (Glinka et al., supra). This gene thus represents a novel growth factor which may be useful in tissue regeneration, and also represents a means for potentially inhibiting cell transformation via wnt signaling.

[0006] The Frzb proteins and the protein Cerberus are examples of secreted proteins that purportedly inhibit wnt signaling (Brown, *Curr. Opinion Cell Biol.*, 10: 182-187 [1998]).

[0007] PCT WO 98/35043, published 13 Aug. 1998 describes human SDF-5 proteins which are purportedly useful in regulating the binding of wnt polypeptides to their receptors.

[0008] PCT WO 98/23730, published 4 Jun. 1998, describes transfecting tumors cells with wnt-5a to purportedly decrease tumorigenicity. Wnt-5a purportedly is an antagonist of other wnts.

[0009] In view of the devastating effects of cancer, there is a need in the art to identify additional genes that may serve as antagonists of proteins involved in cell transformation.

[0010] Accordingly, it is an object of this invention to provide nucleic acid molecules and polypeptides that may be useful as anti-cancer compounds.

[0011] It is a further object to provide methods of altering the level of expression and/or activity of such polypeptides in the human body.

[0012] Other related objects will readily be apparent from a reading of this disclosure.

SUMMARY OF THE INVENTION

[0013] In one embodiment, the present invention provides an isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group consisting of:

[0014] (a) the nucleic acid molecule comprising SEQ ID NO:1;

[0015] (b) the nucleic acid molecule comprising SEQ ID NO:2;

[0016] (c) the nucleic acid molecule comprising SEQ ID NO:3;

[0017] (d) the nucleic acid molecule comprising SEQ ID NO:4;

[0018] (e) the nucleic acid molecule comprising SEQ ID NO:5;

[0019] (f) the nucleic acid molecule comprising SEQ ID NO:6;

[0020] (g) the nucleic acid molecule comprising SEQ ID NO:7;

[0021] (h) the nucleic acid molecule comprising SEQ ID NO:75;

[0022] (i) the nucleic acid molecule comprising SEQ ID NO:76;

[0023] (j) the nucleic acid molecule comprising SEQ ID NO:77;

[0024] (k) the nucleic acid molecule comprising SEQ ID NO:78;

[0025] (l) the nucleic acid molecule encoding the polypeptide of SEQ ID NO:8;

[0026] (m) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:9;

[0027] (n) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:10, or a biologically active fragment thereof;

[0028] (o) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:11, or a biologically active fragment thereof;

[0029] (p) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:12, or a biologically active fragment thereof;

[0030] (q) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:13, or a biologically active fragment thereof;

[0031] (r) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:14, or a biologically active fragment thereof;

[0032] (s) a nucleic acid molecule that encodes a polypeptide that is at least 85 percent identical to the polypeptide of SEQ ID NOs: 10, 11, 12, 13, or 14;

[0033] (t) a nucleic acid molecule that encodes a biologically active DKR polypeptide that has 1-100 amino acid substitutions and/or deletions as compared with the polypeptide of any of SEQ ID NOs: 8, 9, 10, 11, 12, 13, or 14; and

[0034] (u) a nucleic acid molecule that hybridizes under conditions of high stringency to any of (c), (d), (e), (f), (g), (h), (i), (k), (l), (m), (n), (o), (p), (q), (r), (s), and (t) above.

[0035] In another embodiment, the invention provides an isolated nucleic acid molecule that is the complement of any of the nucleic acid molecules above.

[0036] In yet another embodiment, the invention provides an isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group of: amino acids 16-350, 21-350, 22-350, 23-350, 33-350, or 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 150-290, 300-350, or 310-350 of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-359 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

[0037] In other embodiments, the invention provides vectors comprising the nucleic acid molecules, and host cells comprising the vectors.

[0038] In still another embodiment, the invention provides a process for producing a biologically active DKR polypeptide comprising the steps of:

[0039] (a) expressing a polypeptide encoded by any of the nucleic acid molecules herein in a suitable host; and

[0040] (b) isolating the polypeptide.

[0041] In still one other embodiment, the invention provides a biologically active DKR polypeptide selected from the group consisting of:

[0042] (a) the polypeptide of SEQ ID NO:8;

[0043] (b) the polypeptide of SEQ ID NO:9;

[0044] (c) the polypeptide of SEQ ID NO:10;

[0045] (d) the polypeptide of SEQ ID NO:11;

[0046] (e) the polypeptide of SEQ ID NO:12;

[0047] (f) the polypeptide of SEQ ID NO:13;

[0048] (g) the polypeptide of SEQ ID NO:14;

[0049] (h) a polypeptide that has 1-100 amino acid substitutions or deletions as compared with the polypeptide of any of (a)-(g) above; and

[0050] (i) a polypeptide that is at least 85 percent identical to any of the polypeptides of (c)-(h) above.

[0051] In still one other embodiment, the invention provides the following polypeptides: a polypeptide that is amino acids 16-350, 21-350, 22-350, 23-350, 33-350, or 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, or 310-350 of FIG. 9, a polypeptide that is amino acids 15, 266, 24-266, or 32-266

of FIG. 10, a polypeptide that is amino acids 17-259, 26-259, or 34-259 of FIG. 12, and a polypeptide that is amino acids 19-224, 20-224, 21-224, or 22-224 of FIG. 14.

BRIEF DESCRIPTION OF THE DRAWINGS

[0052] FIG. 1 (SEQ ID NO:1) depicts the cDNA sequence of mouse DKR-3.

[0053] FIG. 2 (SEQ ID NO:2) depicts the cDNA sequence of human DKR-3.

[0054] FIG. 3 (SEQ ID NO:3) depicts the DNA sequence of human DKR-1.

[0055] FIG. 4 (SEQ ID NO:4) depicts the cDNA sequence of mouse DKR-2.

[0056] FIG. 5 (SEQ ID NO:5) depicts the cDNA sequence of human DKR-2.

[0057] FIG. 6 (SEQ ID NO:6) depicts the cDNA sequence of human DKR-2a, a splice variant of the DKR-2 gene.

[0058] FIG. 7 (SEQ ID NO:7) depicts the cDNA sequence of human DKR-4.

[0059] FIG. 8 (SEQ ID NO:8) depicts the amino acid sequence of mouse DKR-3 as translated from the corresponding cDNA.

[0060] FIG. 9 (SEQ ID NO:9) depicts the amino acid sequence of human DKR-3 as translated from the corresponding cDNA.

[0061] FIG. 10 (SEQ ID NO:10) depicts the amino acid sequence of human DKR-1 as translated from the corresponding cDNA.

[0062] FIG. 11 (SEQ ID NO:11) depicts the amino acid sequence of mouse DKR-2 as translated from the corresponding cDNA.

[0063] FIG. 12 (SEQ ID NO:12) depicts the amino acid sequence of human DKR-2 as translated from the corresponding cDNA.

[0064] FIG. 13 (SEQ ID NO:13) depicts the amino acid sequence of human DKR-2a as translated from the corresponding cDNA.

[0065] FIG. 14 (SEQ ID NO:14) depicts the amino acid sequence of human DKR-4 as translated from the corresponding cDNA.

[0066] FIGS. 15A-15D are photographs of Northern blots which were probed with human DKR-3. FIG. 15A shows the transcript level of DKR-3 in various human normal (Lanes 1-2) and immortal (Lanes 3-4) cell lines, and in human estrogen receptor plus ("ER+"; Lanes 5-9) and estrogen receptor minus ("ER-"; Lanes 10-16) breast cancer cell lines. FIG. 15B shows the transcript level of human DKR-3 in human normal lung cells (Lane 1), and in various human non-small cell lung cancer ("NSCLC"; Lanes 2-9) and small cell lung cancer ("SCLC"; Lanes 10-15) cell lines. FIG. 15C shows the amount of transcript of human DKR-3 in five glioblastoma cell lines; three of these lines (SNB-19, U-87MG, and U-373MG) are capable of forming tumors in nude mice, while the other two lines (Hs 683 and A 172) are not. FIG. 15D shows the transcript level of human DKR-3 in

human immortal (non-cancerous) and normal cervical cells, and in human cervical cancer cell lines (indicated as “tumor cells”).

[0067] FIG. 16 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

[0068] FIG. 17 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

[0069] FIG. 18 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

[0070] FIG. 19 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

[0071] FIG. 20 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

[0072] FIG. 21 is a photograph of a Western blot. Contents of the Lanes are indicated in the Examples herein.

[0073] FIG. 22 (SEQ ID NO:75) is a nucleic acid sequence of human DKR-1 with codons optimized for expression in *E. coli*.

[0074] FIG. 23 (SEQ ID NO:76) is a nucleic acid sequence of human DKR-2 with codons optimized for expression in *E. coli*.

[0075] FIG. 24 (SEQ ID NO:77) is a nucleic acid sequence of human DKR-3 with codons optimized for expression in *E. coli*.

[0076] FIG. 25 (SEQ ID NO:78) is a nucleic acid sequence of human DKR-4 with codons optimized for expression in *E. coli*.

DETAILED DESCRIPTION OF THE INVENTION

[0077] Included in the scope of this invention are DKR polypeptides such as the polypeptides of SEQ ID NOs:8-14, and related biologically active polypeptide fragments, variants, and derivatives thereof.

[0078] Also included within the scope of the present invention are nucleic acid molecules that encode DKR polypeptides such as the nucleic acid molecules of SEQ ID Nos:1-7.

[0079] Additionally included within the scope of the present invention are non-human mammals such as mice, rats, rabbits, goats, or sheep in which the gene (or genes) encoding a native DKR polypeptide has (have) been disrupted (“knocked out”) such that the level of expression of this gene or genes is (are) significantly decreased or completely abolished. Such mammals may be prepared using techniques and methods such as those described in U.S. Pat. No. 5,557,032. The present invention further includes non-human mammals such as mice, rats, rabbits, goats, or sheep in which the gene (or genes) encoding DKR polypeptides in which either the native form of the gene(s) for that mammal or a heterologous DKR polypeptide gene(s) is (are) over expressed by the mammal, thereby creating a “transgenic” mammal. Such transgenic mammals may be prepared using

well known methods such as those described in U.S. Pat. No. 5,489,743 and PCT patent application no. WO94/28122, published 8 Dec. 1994. The present invention further includes non-human mammals in which the promoter for one or more of the DKR polypeptides of the present invention is either activated or inactivated (using homologous recombination methods as described below) to alter the level of expression of one or more of the native DKR polypeptides.

[0080] The DKR polypeptides of the present invention are expected to have utility as anti-cancer therapeutics for those cancers such as mammary tumors, stem cell tumors, or other cancers in which the wnt and/or sonic hedgehog (shh) signal transduction pathways are activated. Specific wnt members can transform mammary tissue (Hunter, supra) and are abnormally expressed in many human tumors (Huguet, *Cancer Res.*, 54: 2615-2621 [1994]; Dale, *Cancer Res.*, 56: 4320-4323 [1996]; see also PCT WO 97/39357). Such activity is expected in view of data presented herein in which the level of DKR-3 transcript is decreased or not detectable at all in many cancer cell lines as compared to similar normal cell lines. Further, such activity is expected in view of the relationship of the genes and polypeptides of the present invention to the gene dickkopf-1 (which, as mentioned above, is purportedly a potent antagonist of wnt-8). DKR-1, a novel gene of the present invention, is a human ortholog of dkk-1. DKR-2, DKR-3, and DKR-4, all novel genes of the present invention, are each related to DKR-1 by their cysteine pattern. In particular, these DKR polypeptides may be of use for treatment of stem cell tumors, mammary tumors, and other cancers in which wnt genes are expressed, and in cancers where wnt and/or shh signaling is activated.

[0081] The DKR polypeptides of the present invention may also be administered as agents that can induce and/or enhance tissue differentiation, such as bone formation, cartilage formation, muscle tissue formation, nerve tissue formation, and hematopoietic cell formation. Such activities are expected in view of the fact that a) *Xenopus* dkk-1 purportedly promotes head induction, heart formation, and differentiation or the developing CNS (Glinka, supra); and b) certain wnt polypeptides appear to function in embryo development (Cadigan, *Genes and Devel.*, 11: 3286-3305 [1997]), specifically development of the pituitary (Treier, *Genes and Devel.*, 12: 1691-1704 [1998]), myogenesis (Munsterberg et al., *Genes and Devel.*, 9: 2911-2922 [1995]), osteogenesis (PCT WO 95/17416; PCT WO98/16641), kidney development (Stark et al., *Nature* 372: 679-683 [1994]), development of the CNS (Dickinson et al., *Development*, 120: 1453-1471 [1994]), and hematopoiesis (PCT WO 98/06747). Thus, addition of certain DKR polypeptides in such cell cultures or tissues may serve to modify the activity of various wnt polypeptides in cellular differentiation processes.

[0082] The DKR polypeptides herein may be used in either an in vivo manner or an ex vivo manner for such applications. For example, one or more of the DKR polypeptides of the present invention may be added to a culture of cartilage tissue or nerve tissue, or hematopoietic stem cells, either alone, or in combination with other growth factors and/or other tissue differentiation factors, so as to induce or enhance the regeneration of such tissues. Alternatively, such DKR polypeptides of the present invention may, for example, be injected directly into a joint in need of cartilage,

into the spinal cord where the cord has been damaged, into damaged brain tissue, or into bone marrow to enhance hematopoiesis.

[0083] The term “DKR polypeptides” as used herein refers to any protein or polypeptide having the properties described herein for DKR polypeptides. The DKR polypeptides may or may not have amino terminal methionines, depending on the manner in which they are prepared. By way of illustration, DKR polypeptides refers to (1) a biologically active polypeptide encoded by any of the DKR polypeptides nucleic acid molecules as defined in any of items (a)-(f) below; (2) naturally occurring allelic variants and synthetic variants of any of DKR polypeptide in which one or more amino acid substitutions, deletions, and/or insertions are present as compared to the DKR polypeptides of SEQ ID NOS:8-14, and/or (3) biologically active polypeptides, or fragments or variants thereof, that have been chemically modified.

[0084] As used herein, the term “DKR polypeptide fragment” refers to a peptide or polypeptide that is less than the full length amino acid sequence of a naturally occurring DKR polypeptide but has the biological activity of any of the DKR polypeptides provided herein. Such a fragment may be truncated at the amino terminus, the carboxy terminus, and/or internally (such as by natural splicing), and may be a variant or a derivative of any of the DKR polypeptides. Such DKR polypeptides fragments may be prepared with or without an amino terminal methionine. In addition, DKR polypeptides fragments can be naturally occurring fragments such as DKR polypeptide splice variants (SEQ ID NO:13), other splice variants, and fragments resulting from naturally occurring in vivo protease activity. Preferred DKR polypeptide fragments include amino acids 16-350, 21-350, 22-350, 23-350, 33-350, 42-350, 21-145, 40-145, 40-150, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, and 310-350, of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-359 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

[0085] As used herein, the term “DKR polypeptide variants” refers to DKR polypeptides whose amino acid sequences contain one or more amino acid sequence substitutions, deletions, and/or insertions as compared to the DKR polypeptides amino acid sequences set forth in SEQ ID NOS:8-14. Such DKR polypeptides variants can be prepared from the corresponding DKR polypeptides nucleic acid molecule variants, which have a DNA sequence that varies accordingly from the DNA sequences for wild type DKR polypeptides as set forth in SEQ ID NOS:7-14. Preferred variants of the human DKR polypeptides include alanine substitutions at one or more of amino acid positions. Other preferred substitutions include conservative substitutions at the amino acid positions indicated in the Examples herein, as well as those encoded by DKR nucleic acid molecules as described below.

[0086] As used herein, the term “DKR polypeptide derivatives” refers to DKR polypeptides, variants, or fragments thereof, that have been chemically modified, as for example, by addition of one or more polyethylene glycol molecules, sugars, phosphates, and/or other such molecules, where the molecule or molecules are not naturally attached to wild-type DKR polypeptides.

[0087] As used herein, the terms “biologically active DKR polypeptides”, “biologically active DKR polypeptide fragments”, “biologically active DKR polypeptide variants”, and “biologically active DKR polypeptide derivatives” refer to DKR polypeptides that have the ability to decrease cancer cell proliferation in the Anchorage Independent Growth Assay of Example 12 herein, or in the In Vivo Tumor Assay of Example 13 herein, or in both assays.

[0088] As used herein, the term “DKR polypeptide nucleic acids” when used to describe a nucleic acid molecule refers to a nucleic acid molecule or fragment thereof that (a) has the nucleotide sequence as set forth in any of SEQ ID NOS:1-7; (b) has a nucleic acid sequence encoding a polypeptide that is at least 85 percent identical, but may be greater than 85 percent, i.e., 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent identical to the polypeptide encoded by any of SEQ ID NOS:10-14; (c) is a naturally occurring allelic variant or alternate splice variant of (a) or (b); (d) is a nucleic acid variant of (a)-(c) produced as provided for herein; (e) has a sequence that is complementary to (a)-(d); (f) hybridizes to any of (a)-(e) under conditions of high stringency and/or (g) has a nucleic acid sequence encoding 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or up to 100 amino acid substitutions and/or deletions of any mature DKR polypeptide (i.e., a DKR polypeptide with its endogenous signal peptide removed).

[0089] Percent sequence identity can be determined by standard methods that are commonly used to compare the similarity in position of the amino acids of two polypeptides. By way of example, using a computer algorithm such as GAP (Genetic Computer Group, University of Wisconsin, Madison, Wis.), the two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the “matched span”, as determined by the algorithm). A gap opening penalty (which is calculated as 3× the average diagonal; the “average diagonal” is the average of the diagonal of the comparison matrix being used; the “diagonal” is the score or number assigned to each perfect amino acid by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix (see Dayhoff et al., in: *Atlas of Protein Sequence and Structure*, vol. 5, supp.3 [1978] for the PAM250 comparison matrix; see Henikoff et al., *Proc. Natl. Acad. Sci USA*, 89: 10915-10919 [1992] for the BLOSUM 62 comparison matrix) is also used by the algorithm. The percent identity is then calculated by the algorithm by determining the percent identity as follows:

$$\frac{\text{Total number of identical matches in the matched span}}{[\text{length of the longer sequence within the matched span}] + [\text{number of gaps introduced into the longer sequence in order to align the two sequences}]} \times 100$$

[0090] Polypeptides that are at least 85 percent identical will typically have one or more amino acid substitutions, deletions, and/or insertions as compared with any of the wild type DKR polypeptides. Usually, the substitutions of the native residue will be either alanine, or a conservative amino acid so as to have little or no effect on the overall net charge, polarity, or hydrophobicity of the protein. Conservative substitutions are set forth in Table I below.

TABLE I

Conservative Amino Acid Substitutions	
Basic:	arginine lysine histidine
Acidic:	glutamic acid aspartic acid
Uncharged Polar:	glutamine asparagine serine threonine tyrosine
Non-Polar:	phenylalanine tryptophan cysteine glycine alanine valine proline methionine leucine isoleucine

[0091] The term “conditions of high stringency” refers to hybridization and washing under conditions that permit binding of a nucleic acid molecule used for screening, such as an oligonucleotide probe or cDNA molecule probe, to highly homologous sequences. An exemplary high stringency wash solution is 0.2×SSC and 0.1 percent SDS used at a temperature of between 50° C.-65° C.

[0092] Where oligonucleotide probes are used to screen cDNA or genomic libraries, one of the following two high stringency solution may be used. The first of these is 6×SSC with 0.05 percent sodium pyrophosphate at a temperature of 35° C.-62° C., depending on the length of the oligonucleotide probe. For example, 14 base pair probes are washed at 35-40° C., 17 base pair probes are washed at 45-50° C., 20 base pair probes are washed at 52-57° C., and 23 base pair probes are washed at 57-63° C. The temperature can be increased 2-3° C. where the background non-specific binding appears high. A second high stringency solution utilizes tetramethylammonium chloride (TMAC) for washing oligonucleotide probes. One stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and 0.2 percent SDS. The washing temperature using this solution is a function of the length of the probe. For example, a 17 base pair probe is washed at about 45-50° C.

[0093] As used herein, the terms “effective amount” and “therapeutically effective amount” refer to the amount of a DKR polypeptide necessary to support one or more biological activities of the DKR polypeptides as set forth above.

[0094] A full-length DKR polypeptide or fragment thereof can be prepared using well known recombinant DNA technology methods such as those set forth in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring

Harbor Laboratory Press, Cold Spring Harbor, N.Y. [1989]) and/or Ausubel et al., eds., (*Current Protocols in Molecular Biology*, Green Publishers Inc. and Wiley and Sons, NY [1994]). A gene or cDNA encoding a DKR polypeptide or fragment thereof may be obtained for example by screening a genomic or cDNA library, or by PCR amplification. Probes or primers useful for screening the library can be generated based on sequence information for other known genes or gene fragments from the same or a related family of genes, such as, for example, conserved motifs found in other DKR polypeptides such as the cysteine pattern. In addition, where a gene encoding DKR polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify homologous genes from other species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the DKR gene. Typically, conditions of high stringency will be employed for screening to minimize the number of false positives obtained from the screen.

[0095] Another means to prepare a gene encoding a DKR polypeptide or fragment thereof is to employ chemical synthesis using methods well known to the skilled artisan such as those described by Engels et al. (*Angew. Chem. Intl. Ed.*, 28: 716-734 [1989]). These methods include, inter alia, the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the DKR polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full length DKR polypeptide. Usually, the DNA fragment encoding the amino terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of the DKR polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell.

[0096] In some cases, it may be desirable to prepare nucleic acid and/or amino acid variants of the naturally occurring DKR polypeptides. Nucleic acid variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point mutations (see Sambrook et al., supra, and Ausubel et al., supra, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels et al., supra, may also be used to prepare such variants. Other methods known to the skilled artisan may be used as well. Preferred nucleic acid variants are those containing nucleotide substitutions accounting for codon preference in the host cell that is to be used to produce the DKR polypeptide(s). Such “codon optimization” can be determined via computer algorithms which incorporate codon frequency tables such as “Ecohigh. Cod” for codon preference of highly expressed bacterial genes as provided by the University of Wisconsin Package Version 9.0, Genetics Computer Group, Madison, Wis. Other useful codon frequency tables include “Celegans_high.cod”, “Celegans_low.cod”, “Drosophila_high.cod”, “Human_high.cod”, “Maize_high.cod”, and “Yeast_high.cod”. Other preferred variants are those encoding conservative amino acid changes as described above (e.g., wherein the charge or polarity of the naturally occurring amino acid side chain is not altered

substantially by substitution with a different amino acid) as compared to wild type, and/or those designed to either generate a novel glycosylation and/or phosphorylation site(s), or those designed to delete an existing glycosylation and/or phosphorylation site(s).

[0097] The gene, cDNA, or fragment thereof encoding the DKR polypeptide can be inserted into an appropriate expression or amplification vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). The gene, cDNA or fragment thereof encoding the DKR polypeptide may be amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether the DKR polypeptide or fragment thereof is to be glycosylated and/or phosphorylated. If so, yeast, insect, or mammalian host cells are preferable.

[0098] Typically, the vectors used in any of the host cells will contain 5' flanking sequence (also referred to as a "promoter") and other regulatory elements as well such as an enhancer(s), an origin of replication element, a transcriptional termination element, a complete intron sequence containing a donor and acceptor splice site, a signal peptide sequence, a ribosome binding site element, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these elements is discussed below. Optionally, the vector may contain a "tag" sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the DKR polypeptide coding sequence; the oligonucleotide molecule encodes polyHis (such as hexaHis), or other "tag" such as FLAG, HA (hemagglutinin Influenza virus) or myc for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as means for affinity purification of the DKR polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified DKR polypeptide by various means such as using certain peptidases.

[0099] The human immunoglobulin hinge and Fc region could be fused at either the N-terminus or C-terminus of the DKR polypeptides by one skilled in the art. The subsequent Fc-fusion protein could be purified by use of a Protein A affinity column. Fc is known to exhibit a long pharmacokinetic half-life in vivo and proteins fused to Fc have been found to exhibit a substantially greater half-life in vivo than the unfused counterpart. Also, fusion to the Fc region allows for dimerization/multimerization of the molecule that may be useful for the bioactivity of some molecules.

[0100] The 5' flanking sequence may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of 5' flanking sequences from more than one source), synthetic, or it may be the native DKR polypeptides gene 5' flanking sequence. As such, the source of the 5' flanking sequence may be any unicellular prokaryotic or eukaryotic organism, any verte-

brate or invertebrate organism, or any plant, provided that the 5' flanking sequence is functional in, and can be activated by, the host cell machinery.

[0101] The 5' flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, 5' flanking sequences useful herein other than the DKR gene flanking sequence will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of the 5' flanking sequence may be known. Here, the 5' flanking sequence may be synthesized using the methods described above for nucleic acid synthesis or cloning.

[0102] Where all or only a portion of the 5' flanking sequence is known, it may be obtained using PCR and/or by screening a genomic library with suitable oligonucleotide and/or 5' flanking sequence fragments from the same or another species.

[0103] Where the 5' flanking sequence is not known, a fragment of DNA containing a 5' flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion using one or more carefully selected enzymes to isolate the proper DNA fragment. After digestion, the desired fragment may be isolated by agarose gel purification, Qiagen® column or other methods known to the skilled artisan. Selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

[0104] The origin of replication element is typically a part of prokaryotic expression vectors purchased commercially, and aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for optimal expression of the DKR polypeptide. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector.

[0105] The transcription termination element is typically located 3' of the end of the DKR polypeptide coding sequence and serves to terminate transcription of the DKR polypeptide. Usually, the transcription termination element in prokaryotic cells is a G-C rich fragment followed by a poly T sequence. While the element is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described above.

[0106] A selectable marker gene element encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells, (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene.

[0107] The ribosome binding element, commonly called the Shine-Dalgarno sequence (prokaryotes) or the Kozak

sequence (eukaryotes), is usually necessary for translation initiation of mRNA. The element is typically located 3' to the promoter and 5' to the coding sequence of the DKR polypeptide to be synthesized. The Shine-Dalgarno sequence is varied but is typically a polypurine (i.e., having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth above and used in a prokaryotic vector.

[0108] In those cases where it is desirable for DKR polypeptide to be secreted from the host cell, a signal sequence may be used to direct the DKR polypeptide out of the host cell where it is synthesized, and the carboxy-terminal part of the protein may be deleted in order to prevent membrane anchoring. Typically, the signal sequence is positioned in the coding region of the DKR gene or cDNA, or directly at the 5' end of the DKR gene coding region. Many signal sequences have been identified, and any of them that are functional in the selected host cell may be used in conjunction with the DKR gene or cDNA. Therefore, the signal sequence may be homologous or heterologous to the DKR gene or cDNA, and may be homologous or heterologous to the DKR polypeptides gene or cDNA. Additionally, the signal sequence may be chemically synthesized using methods set forth above.

[0109] In most cases, secretion of the polypeptide from the host cell via the presence of a signal peptide will result in the removal of the amino terminal methionine from the polypeptide.

[0110] In many cases, transcription of the DKR gene or cDNA is increased by the presence of one or more introns in the vector; this is particularly true where the DKR polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the DKR gene, especially where the gene used is a full length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron(s) may be obtained from another source. The position of the intron with respect to the 5' flanking sequence and the DKR gene is generally important, as the intron must be transcribed to be effective. As such, where the DKR gene is inserted into the expression vector is a cDNA molecule, the preferred position for the intron is 3' to the transcription start site, and 5' to the polyA transcription termination sequence. Preferably for DKR cDNA, the intron or introns will be located on one side or the other (i.e., 5' or 3') of the cDNA such that it does not interrupt the this coding sequence. Any intron from any source, including any viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell(s) into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

[0111] Where one or more of the elements set forth above are not already present in the vector to be used, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the elements are well known to the skilled artisan and are comparable to the methods set forth above (i.e., synthesis of the DNA, library screening, and the like).

[0112] The final vectors used to practice this invention are typically constructed from a starting vectors such as a

commercially available vector. Such vectors may or may not contain some of the elements to be included in the completed vector. If none of the desired elements are present in the starting vector, each element may be individually ligated into the vector by cutting the vector with the appropriate restriction endonuclease(s) such that the ends of the element to be ligated in and the ends of the vector are compatible for ligation. In some cases, it may be necessary to "blunt" the ends to be ligated together in order to obtain a satisfactory ligation. Blunting is accomplished by first filling in "sticky ends" using Klenow DNA polymerase or T4 DNA polymerase in the presence of all four nucleotides. This procedure is well known in the art and is described for example in Sambrook et al., *supra*.

[0113] Alternatively, two or more of the elements to be inserted into the vector may first be ligated together (if they are to be positioned adjacent to each other) and then ligated into the vector.

[0114] One other method for constructing the vector to conduct all ligations of the various elements simultaneously in one reaction mixture. Here, many nonsense or nonfunctional vectors will be generated due to improper ligation or insertion of the elements, however the functional vector may be identified and selected by restriction endonuclease digestion.

[0115] Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, pCR3, and pcDNA3.1 (Invitrogen Company, San Diego, Calif.), pBSII (Stratagene Company, La Jolla, Calif.), pET15b (Novagen, Madison, Wis.), pGEX (Pharmacia Biotech, Piscataway, N.J.), pEGFP-N2 (Clontech, Palo Alto, Calif.), pETL (BlueBacII; Invitrogen), and pFastBacDual (Gibco/BRL, Grand Island, N.Y.).

[0116] After the vector has been constructed and a nucleic acid molecule encoding full length or truncated DKR polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression.

[0117] Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast cell, an insect cell, or a vertebrate cell). The host cell, when cultured under appropriate conditions, can synthesize DKR polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). After collection, the DKR polypeptide can be purified using methods such as molecular sieve chromatography, affinity chromatography, and the like.

[0118] Selection of the host cell for DKR polypeptide production will depend in part on whether the DKR polypeptide is to be glycosylated or phosphorylated (in which case eukaryotic host cells are preferred), and the manner in which the host cell is able to "fold" the protein into its native tertiary structure (e.g., proper orientation of disulfide bridges, etc.) such that biologically active protein is prepared by the DKR polypeptide that has biological activity, the DKR polypeptide may be "folded" after synthesis using appropriate chemical conditions as discussed below.

[0119] Suitable cells or cell lines may be mammalian cells, such as Chinese hamster ovary cells (CHO), human embry-

onic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from in vitro culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines.

[0120] Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 α , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas* spp., other *Bacillus* spp., *Streptomyces* spp., and the like may also be employed in this method.

[0121] Many strains of yeast cells known to those skilled in the art are also available as host cells for expression of the polypeptides of the present invention.

[0122] Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described for example in Kitts et al. (*Biotechniques*, 14: 810-817 [1993]), Lucklow (*Curr. Opin. Biotechnol.*, 4: 564-572 [1993]) and Lucklow et al. (*J. Virol.*, 67: 4566-4579 [1993]). Preferred insect cells are Sf-9 and Hi5 (Invitrogen, Carlsbad, Calif.).

[0123] Insertion (also referred to as "transformation" or "transfection") of the vector into the selected host cell may be accomplished using such methods as calcium chloride, electroporation, microinjection, lipofection or the DEAE-dextran method. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., supra.

[0124] The host cells containing the vector (i.e., transformed or transfected) may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells are for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells are RPMI 1640, MEM, DMEM, all of which may be supplemented with serum and/or growth factors as required by the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

[0125] Typically, an antibiotic or other compound useful for selective growth of the transformed cells only is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin.

[0126] The amount of DKR polypeptide produced in the host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, HPLC separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

[0127] If the DKR polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. Polypeptides prepared in this way will typically not possess an amino terminal methionine, as it is removed during secretion from the cell. If however, the DKR polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for gram negative bacteria host cells) and may have an amino terminal methionine.

[0128] For DKR polypeptide situated in the host cell cytoplasm and/or nucleus, the host cells are typically first disrupted mechanically or with detergent to release the intra-cellular contents into a buffered solution. DKR polypeptide can then be isolated from this solution.

[0129] Purification of DKR polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (DKR polypeptide/hexaHis) or other small peptide such as FLAG (Eastman Kodak Co., New Haven, Conn.) or myc (Invitrogen, Carlsbad, Calif.) at either its carboxyl or amino terminus, it may essentially be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag or for the polypeptide directly (i.e., a monoclonal antibody specifically recognizing DKR polypeptide). For example, polyhistidine binds with great affinity and specificity to nickel, thus an affinity column of nickel (such as the Qiagen® nickel columns) can be used for purification of DKR polypeptide/polyHis. (See for example, Ausubel et al., eds., *Current Protocols in Molecular Biology*, Section 10.11.8, John Wiley & Sons, New York [1993]).

[0130] Where the DKR polypeptide is prepared without a tag attached, and no antibodies are available, other well known procedures for purification can be used. Such procedures include, without limitation, ion exchange chromatography, molecular sieve chromatography, HPLC, native gel electrophoresis in combination with gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific). In some cases, two or more of these techniques may be combined to achieve increased purity.

[0131] If it is anticipated that the DKR polypeptide will be found primarily intracellularly, the intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

[0132] If the DKR polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation.

The pellet material can then be treated at pH extremes or with chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The DKR polypeptide in its now soluble form can then be analyzed using gel electrophoresis, immunoprecipitation or the like. If it is desired to isolate the DKR polypeptide, isolation may be accomplished using standard methods such as those set forth below and in Marston et al. (*Meth. Enz.*, 182: 264-275 [1990]). In some cases, the DKR polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages, can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization but usually at a lower concentration and is not necessarily the same chaotrope as used for the solubilization. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent plus its' oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridge(s). Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol (DTT)/dithiane DTT, 2-mercaptoethanol(bME)/dithio-b(ME). In many instances a cosolvent is necessary to increase the efficiency of the refolding and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, and arginine.

[0133] If DKR polypeptide inclusion bodies are not formed to a significant degree in the host cell, the DKR polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate, and the DKR polypeptide can be isolated from the supernatant using methods such as those set forth below.

[0134] In those situations where it is preferable to partially or completely isolate the DKR polypeptide, purification can be accomplished using standard methods well known to the skilled artisan. Such methods include, without limitation, separation by electrophoresis followed by electroelution, various types of chromatography (immunoaffinity, molecular sieve, and/or ion exchange), and/or high pressure liquid chromatography. In some cases, it may be preferable to use more than one of these methods for complete purification.

[0135] In addition to preparing and purifying DKR polypeptide using recombinant DNA techniques, the DKR polypeptides, fragments, and/or derivatives thereof may be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield et al., (*J. Am. Chem. Soc.*, 85: 2149 [1963]), Houghten et al. (*Proc Natl Acad. Sci. USA*, 82: 5132 [1985]), and Stewart and Young (*Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, Ill. [1984]). Such polypeptides may be synthesized with or without a methionine on the amino terminus. Chemically synthesized DKR polypeptides or fragments may be oxidized using methods set forth in these references to form disulfide bridges. The DKR polypeptides or fragments are expected to have bio-

logical activity comparable to DKR polypeptides produced recombinantly or purified from natural sources, and thus may be used interchangeably with recombinant or natural DKR polypeptide.

[0136] Chemically modified DKR polypeptide compositions in which DKR polypeptide is linked to a polymer are included within the scope of the present invention. The polymer selected is typically water soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. The polymer selected is usually modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the degree of polymerization may be controlled as provided for in the present methods. The polymer may be of any molecular weight, and may be branched or unbranched. Included within the scope of DKR polypeptide polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

[0137] The water soluble polymer or mixture thereof may be selected from the group consisting of, for example, polyethylene glycol (PEG), monomethoxy-polyethylene glycol, dextran, cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (e.g., glycerol) and polyvinyl alcohol.

[0138] For the acylation reactions, the polymer(s) selected should have a single reactive ester group. For reductive alkylation, the polymer(s) selected should have a single reactive aldehyde group. A preferred reactive aldehyde is polyethylene glycol propionaldehyde, which is water stable, or mono C1-C10 alkoxy or aryloxy derivatives thereof (see U.S. Pat. No. 5,252,714).

[0139] Pegylation of DKR polypeptides may be carried out by any of the pegylation reactions known in the art, as described for example in the following references: *Focus on Growth Factors* 3: 4-10 (1992); EP 0 154 316; and EP 0 401 384. Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described below.

[0140] A particularly preferred water-soluble polymer for use herein is polyethylene glycol, abbreviated PEG. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol.

[0141] In general, chemical derivatization may be performed under any suitable conditions used to react a biologically active substance with an activated polymer molecule. Methods for preparing pegylated DKR polypeptides will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby DKR polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of PEG: protein, the greater the percentage of poly-pegylated product.

[0142] Generally, conditions which may be alleviated or modulated by administration of the present polymer/polypeptides include those described herein for DKR polypeptides molecules. However, the polymer/ DKR polypeptides molecules disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

[0143] The DKR polypeptides, fragments thereof, variants, and derivatives, may be employed alone, together, or in combination with other pharmaceutical compositions. The DKR polypeptides, fragments, variants, and derivatives may be used in combination with cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the indication being treated.

[0144] DKR nucleic acid molecules, fragments, and/or derivatives that do not themselves encode polypeptides that are active in activity assays may be useful as hybridization probes in diagnostic assays to test, either qualitatively or quantitatively, for the presence of DKR DNA or corresponding RNA in mammalian tissue or bodily fluid samples.

[0145] DKR polypeptide fragments, variants, and/or derivatives that are not themselves active in activity assays may be useful for preparing antibodies that recognize DKR polypeptides.

[0146] The DKR polypeptides, fragments, variants, and/or derivatives may be used to prepare antibodies using standard methods. Thus, antibodies that react with the DKR polypeptides, as well as reactive fragments of such antibodies, are also contemplated as within the scope of the present invention. The antibodies may be polyclonal, monoclonal, recombinant, chimeric, single-chain and/or bispecific. Typically, the antibody or fragment thereof will either be of human origin, or will be "humanized", i.e., prepared so as to prevent or minimize an immune reaction to the antibody when administered to a patient. The antibody fragment may be any fragment that is reactive with DKR polypeptides of the present invention, such as, F_{ab} , F_{ab}' , etc. Also provided by this invention are the hybridomas generated by presenting any DKR polypeptide or fragments thereof as an antigen to a selected mammal, followed by fusing cells (e.g., spleen cells) of the mammal with certain cancer cells to create immortalized cell lines by known techniques. The methods employed to generate such cell lines and antibodies directed against all or portions of a human DKR polypeptide of the present invention are also encompassed by this invention.

[0147] The antibodies may be used therapeutically, such as to inhibit binding of the DKR polypeptide to its binding partner. The antibodies may further be used for in vivo and in vitro diagnostic purposes, such as in labeled form to detect the presence of DKR polypeptide in a body fluid or cell sample.

[0148] Preferred antibodies are human antibodies, either polyclonal or monoclonal.

[0149] Therapeutic Compositions and Administration

[0150] Therapeutic compositions of DKR polypeptides are within the scope of the present invention. Such compositions may comprise a therapeutically effective amount of the polypeptide or fragments, variants, or derivatives in admixture with a pharmaceutically acceptable carrier. The

carrier material may be water for injection, preferably supplemented with other materials common in solutions for administration to mammals. Typically, a DKR polypeptide therapeutic compound will be administered in the form of a composition comprising purified polypeptide, fragment, variant, or derivative in conjunction with one or more physiologically acceptable carriers, excipients, or diluents. Neutral buffered saline or saline mixed with serum albumin are exemplary appropriate carriers. Preferably, the product is formulated as a lyophilizate using appropriate excipients (e.g., sucrose). Other standard carriers, diluents, and excipients may be included as desired. Other exemplary compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor.

[0151] The DKR polypeptide compositions can be administered parenterally. Alternatively, the compositions may be administered intravenously or subcutaneously. When systemically administered, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such pharmaceutically acceptable protein solutions, with due regard to pH, isotonicity, stability and the like, is within the skill of the art.

[0152] Therapeutic formulations of DKR polypeptide compositions useful for practicing the present invention may be prepared for storage by mixing the selected composition having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*Remington's Pharmaceutical Sciences*, 18th Edition, A. R. Gennaro, ed., Mack Publishing Company [1990]) in the form of a lyophilized cake or an aqueous solution. Acceptable carriers, excipients or stabilizers are nontoxic to recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

[0153] An effective amount of the DKR polypeptide composition(s) to be employed therapeutically will depend, for example, upon the therapeutic objectives such as the indication for which the DKR polypeptide is being used, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage may range from about 0.1 $\mu\text{g/kg}$ to up to 100 mg/kg or more, depending on the factors mentioned above. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of DKR polypeptide) over time, or as a continuous infusion via implantation device or catheter.

[0154] As further studies are conducted, information will emerge regarding appropriate dosage levels for treatment of

various conditions in various patients, and the ordinary skilled worker, considering the therapeutic context, the type of disorder under treatment, the age and general health of the recipient, will be able to ascertain proper dosing.

[0155] The DKR polypeptide composition to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using these methods may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration ordinarily will be stored in lyophilized form or in solution.

[0156] Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0157] The route of administration of the composition is in accord with known methods, e.g. oral, injection or infusion by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, or intralesional routes, or by sustained release systems or implantation device which may optionally involve the use of a catheter. Where desired, the compositions may be administered continuously by infusion, bolus injection or by implantation device.

[0158] Alternatively or additionally, the composition may be administered locally via implantation into the affected area of a membrane, sponge, or other appropriate material on to which DKR polypeptide has been absorbed.

[0159] Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of DKR polypeptide may be directly through the device via bolus, or via continuous administration, or via catheter using continuous infusion.

[0160] DKR polypeptide may be administered in a sustained release formulation or preparation. Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al, *Biopolymers*, 22: 547-556 (1983)), poly (2hydroxyethyl-methacrylate) (Langer et al., *J. Biomed. Mater. Res.*, 15: 167-277 [1981] and Langer, *Chem. Tech.*, 12: 98-105 [1982]), ethylene vinyl acetate (Langer et al., supra) or poly-D(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also may include liposomes, which can be prepared by any of several methods known in the art (e.g., Eppstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688-3692 [1985]; EP 36,676; EP 88,046; EP 143,949).

[0161] In some cases, it may be desirable to use DKR polypeptide compositions in an ex vivo manner. Here, cells, tissues, or organs that have been removed from the patient are exposed to DKR polypeptide compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

[0162] In other cases, DKR polypeptide may be delivered through implanting into patients certain cells that have been genetically engineered, using methods such as those

described herein, to express and secrete the polypeptides, fragments, variants, or derivatives. Such cells may be animal or human cells, and may be derived from the patient's own tissue or from another source, either human or non-human. Optionally, the cells may be immortalized. However, in order to decrease the chance of an immunological response, it is preferred that the cells be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow release of the protein product(s) but prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

[0163] Methods used for membrane encapsulation of cells are familiar to the skilled artisan, and preparation of encapsulated cells and their implantation in patients may be accomplished without undue experimentation. See, e.g., U.S. Pat. Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT WO 91/10425 (Aebischer et al.). Techniques for formulating a variety of other sustained or controlled delivery means, such as liposome carriers, bio-erodible particles or beads, are also known to those in the art, and are described, for example, in U.S. Pat. No. 5,653,975 (Baetge et al., Cyto-Therapeutics, Inc.). The cells, with or without encapsulation, may be implanted into suitable body tissues or organs of the patient.

[0164] As discussed above, it may be desirable to treat isolated cell populations such as stem cells, lymphocytes, red blood cells, chondrocytes, neurons, and the like with one or more DKR polypeptides, variants, derivatives and/or fragments. This can be accomplished by exposing the isolated cells to the polypeptide, variant, derivative, or fragment directly, where it is in a form that is permeable to the cell membrane. Alternatively, gene therapy can be employed as described below.

[0165] One manner in which gene therapy can be applied is to use the DKR gene (either genomic DNA, cDNA, and/or synthetic DNA encoding a DKR polypeptide, or a fragment, variant, or derivative thereof) which may be operably linked to a constitutive or inducible promoter to form a "gene therapy DNA construct". The promoter may be homologous or heterologous to the endogenous DKR gene, provided that it is active in the cell or tissue type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include, as required, DNA molecules designed for site-specific integration (e.g., endogenous flanking sequences useful for homologous recombination), tissue-specific promoter, enhancer(s) or silencer(s), DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell targeting) cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as factors to enable vector manufacture.

[0166] This gene therapy DNA construct can then be introduced into the patient's cells (either ex vivo or in vivo). One means for introducing the gene therapy DNA construct is via viral vectors. Suitable viral vectors typically used in gene therapy for delivery of gene therapy DNA constructs include, without limitation, adenovirus, adeno-associated

virus, herpes simplex virus, *lentivirus*, papilloma virus, and retrovirus vectors. Some of these vectors, such as retroviral vectors, will deliver the gene therapy DNA construct to the chromosomal DNA of the patient's cells, and the gene therapy DNA construct can integrate into the chromosomal DNA; other vectors will function as episomes and the gene therapy DNA construct will remain in the cytoplasm. The use of gene therapy vectors is described, for example, in U.S. Pat. No. 5,672,344 (30 Sep. 1997; Kelly et al., University of Michigan), U.S. Pat. No. 5,399,346 (21 Mar. 1995; Anderson et al., U.S. Dept. Health and Human Services), U.S. Pat. No. 5,631,236 (20 May 1997; Woo et al., Baylor College of Medicine), and U.S. Pat. No. 5,635,399 (3 Jun. 1997; Kriegler et al., Chiron Corp.).

[0167] Alternative means to deliver gene therapy DNA constructs to a patient's cells without the use of viral vectors include, without limitation, liposome-mediated transfer, direct injection of naked DNA, receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation, and microparticle bombardment (e.g., "gene gun"). See U.S. Pat. No. 4,970,154 (13 Nov. 1990; Chang, Baylor College of Medicine), WO 96/40958 (19 Dec. 1996; Smith et al., Baylor College of Medicine) U.S. Pat. No. 5,679,559 (21 Oct. 1997; Kim et al., University of Utah) U.S. Pat. No. 5,676,954 (14 Oct. 1997; Brigham, Vanderbilt University), and U.S. Pat. No. 5,593,875 (14 Jan. 1997; Wurm et al., Genentech).

[0168] Another means to increase endogenous DKR polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the DKR polypeptide promoter, where the enhancer element(s) can serve to increase transcriptional activity of the DKR polypeptides gene. The enhancer element(s) used will be selected based on the tissue in which one desires to activate the gene(s); enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a DKR polypeptide is to be "turned on" in T-cells, the *lck* promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing the DKR polypeptide promoter (and optionally, vector, 5' and/or 3' flanking sequence, etc.) using standard cloning techniques. This construct, known as a "homologous recombination construct" can then be introduced into the desired cells either *ex vivo* or *in vivo*.

[0169] Gene therapy can be used to decrease DKR polypeptide expression by modifying the nucleotide sequence of the endogenous promoter(s). Such modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the DKR gene(s) selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. Here, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing transcription of the corresponding DKR gene. Deletion of the TATA box or transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the DKR polypeptide promoter(s) (from the same or a related species as the DKR gene(s) to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated

via substitution, deletion and/or insertion of one or more nucleotides such that the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' flanking regions of the promoter segment that has been modified, may be introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described above. Typically, integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' flanking DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to the endogenous chromosomal DNA.

[0170] Other gene therapy methods may also be employed where it is desirable to inhibit one or more DKR polypeptides. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of the selected DKR polypeptide gene(s) can be introduced into the cell. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected DKR gene. When the antisense molecule then hybridizes to the corresponding DKR polypeptides mRNA, translation of this mRNA is prevented.

[0171] Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more of the DKR polypeptides. In this situation, the DNA encoding a mutant full length or truncated polypeptide of each selected DKR polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described above. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

[0172] Samples of the *E. coli* cell lines GM121 and GM94 have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, Va., USA on DATE as accession numbers X and Y, respectively.

[0173] The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

EXAMPLES

Example 1

Cloning of the Mouse DKR-3 Gene

[0174] About 120 adult mice with an average body weight of about 18 grams were each injected intraperitoneally with a kainate solution (prepared as a stock solution of about 1 mg/ml kainate in sterile PBS) at a dose of about 25 mg kainate per kilogram body weight. About six hours after injection, the mice were sacrificed, and the hippocampus was dissected from each mouse. Total RNA was extracted from hippocampal tissue using the Trizol method (Gibco BRL, Grand Island, N.Y.). The poly(A⁺) mRNA fraction was isolated from total RNA using Message Maker (Gibco BRL, Grand Island, N.Y.) according to the manufacturer's recommended procedure. Hippocampal tissue was also obtained from control mice (which received an injection of PBS only), and poly(a⁺) mRNA was obtained from this tissue as well using the same procedures.

[0175] Two random primed cDNA libraries were prepared; one from the kainate-treated and one from the control

poly (A+) mRNA using the Superscript® plasmid system (Gibco BRL, Gaithersburg, Md.). A random cDNA primer containing an internal NotI restriction site was used to initiate first strand synthesis and had the following sequence:

GGAAGGAAAAAGCGCGCGCAACANNNNNNNN (SEQ ID NO: 15)

[0176] where N is A, G, C, or T.

[0177] Both first strand cDNA synthesis and second strand cDNA synthesis were performed according to the manufacturer's recommended protocol. After second strand synthesis, the reaction products were extracted with phenol:chloroform:isoamyl alcohol (in a volume ratio of 25: 24: 1), followed by ethanol precipitation. The double strand cDNA products were ligated using standard ligation procedures to the following double stranded oligonucleotide adapter (obtained from Gibco BRL, Grand Island, N.Y.):

TCGACCCACGCGTCCG (SEQ ID NO:16)

GGGTGCGCAGGC (SEQ ID NO:17)

[0178] After ligation, the cDNA was digested to completion with NotI, and size fractionated on a 1 percent agarose gel. The cDNA products between about 250 and 800 base pairs were selected and purified from the gel using the Qiagen® gel extraction kit (Qiagen, Chatsworth, Calif.). The purified cDNA products were directionally ligated into the vector pYY41L (American Type Culture Collection, "ATCC"; 10801 University Blvd., Manassas, Va., USA; accession number 209636) which had been previously digested with NotI and SalI. The ligated cDNA was then introduced into electrocompetent ElectroMax® DH108 *E. coli* cells (Gibco-BRL, Grand Island, N.Y.) via standard electroporation techniques. The library was then titered by a serial dilution of the transformation cell mixture.

[0179] About one million primary clones were divided into 20 pools (50,000 clones each pool) and each pool was plated on 245 mm×245 mm square plate containing MR2001 medium (MacConnel Research, San Diego, Calif.) and about 60 ug/ml carbonocillin. After incubation overnight at 37 C, the colonies were scraped off the plate in about 20 ml SOC(SOC contains about 2 percent Bactotryptone, 0.5 percent yeast extract, 10 mM sodium chloride, 2.5 mM potassium chloride, and 10 mM magnesium sulfate) and were pelleted by centrifugation at about 6000 rpm for about 10 minutes. The plasmids were then recovered from the cells using Qiagen® maxi prep columns (Qiagen, Chatsworth, Calif.) according to the protocol suggested by the manufacturer.

[0180] About two hundred and fifty thousand clones (50 ug total plasmids/10 ug from each pool) were used to transform yeast strain YPH499 (ATCC accession number 90834) and an amylase-based signal trap assay was conducted as follows (see co-pending U.S. Ser. No. 09/026,959 filed 20 Feb. 1998 for a detailed description of this technique). Around 1000 transformants were plated on a single starch-containing selection plate (15 cm diameter with a medium containing about 0.6 percent yeast nitrogen base, 2 percent glucose, 0.1 percent CAA, 1.0xtrp dropout solution, 0.7 percent potato starch azure, and 1.5 percent agarose).

The plates were incubated at about 30 C for 4-5 days until full development of halos was observed. The colonies in the center of the halo were picked and restreaked on a fresh plate to form single colonies. The single colonies with halos were then picked and arrayed into 96 well microtiter plates containing about 100 ul of water per well, thereby generating the "yeast colony solutions".

[0181] About ten microliters of each well of each yeast colony solution was used as template to recover the cDNA fragment from that colony through PCR. Therefore, ninety-six PCR reactions were independently performed using PCR-Ready Beads® (96 well format, Amersham-Pharmacia Biotech, Piscataway, N.J.) and the following oligonucleotides according to the manufacturer's protocol:

ACTAGCTCCAGTGATCTC (SEQ ID NO:18)

CGTCATTGTTCTCGTTCC (SEQ ID NO:19)

[0182] PCR was conducted using a Perkin-Elmer 9600 thermocycler with the following cycle conditions: 94 C for 10 minutes followed by 35 cycles of 94 C for 30 seconds, 55 C for 30 seconds and 72 C for 1 minute, after which a final extension cycle of 72 C for 10 minutes was conducted. Most PCR reactions contained a single PCR product. The amplified cDNA products were purified using the Qiagen® PCR purification kit (Qiagen, Chatsworth, Calif.). These products were sequenced on an Applied Biosystems 373A automated DNA sequencer using the following oligonucleotide primer:

GCTATACCAAGCATACAATC (SEQ ID NO:35)

[0183] Taq dye-terminator sequencing reactions (Applied Biosystems, Foster City, Calif.) were conducted following the manufacturer's recommended procedures.

[0184] Each PCR fragment was translated in all six possible ways to identify those fragments which (1) had a potential signal peptide in the same direction as reporter gene; (2) had a stop codon(s) upstream of the putative methionine translation start site; and (3) appeared to lack a transmembrane domain.

[0185] One clone that met these criteria, termed "ymrs2-00009-c4", was selected for further analysis. This clone contained 5' sequence, including a putative signal sequence, but was lacking 3' sequence.

[0186] To obtain the 3' sequence of this clone, a 3' RACE reaction was performed using as a template pool number 4 from the YmHK2 cDNA library. This YmHK2 library was prepared as follows: First strand cDNA synthesis was performed using about 2 micrograms of the RNA obtained from the hippocampus of the kainate treated mice and about 1 ug of Not I primer-adaptor having the following sequence:

GACTAGTCTAGATCGCGAGCGCGCCCTTTTTT (SEQ ID NO:42)

TTTTTTTT

[0187] Both the first strand and second strand cDNA synthesis reactions were performed using the Superscript® plasmid system (Gibco BRL, Grand Island, N.Y.). After

second strand synthesis, the double stranded cDNA products were ligated into the double stranded adapters of SEQ ID NOs:16 and 17.

[0188] After ligation, the cDNA was digested to completion with Not I, and size fractionated on a 0.8 percent agarose gel. The cDNA products larger than about 800 base pairs were selected and purified from the gel using the Qiagen® gel extraction kit (Qiagen, Chatsworth, Calif.). The purified cDNA products were directly ligated into Sal I and Not I digested pSport® vector (Gibco BRL, Grand Island, N.Y.).

[0189] The ligated cDNA products were then introduced into electrocompetent *E. coli* cells called ElectroMax® DH10B (Gibco BRL, Grand Island, N.Y.). The library was then titered.

[0190] About twelve million primary clones were obtained, and expanded into about 250 ml of LB containing about 100 ug/ml ampicillin. After overnight incubation at 37 C, the plasmids were recovered using the Qiagen® maxi-prep kit (Qiagen, Chatsworth, Calif.).

[0191] About 20 ng of the plasmid library were used to transform the ElectroMax® DH10B electrocompetent *E. coli* cells using standard electroporation techniques. About two million transformants were divided into 40 pools (containing approximately 50,000 plasmids/pool). Each pool was then expanded into about 3 ml of LB medium containing about 100 ug/ml ampicillin. After overnight incubation at 37 C, the plasmids were recovered using the Qiagen® mini-prep kit. The DNA from each pool were then stored at about minus 20 C for future use.

[0192] The 3' RACE reaction was performed using about 1.5 ng of pool #4 of the YmHK2 library as a template, and using the Advantage® cDNA PCR kit (Clontech, Palo Alto, Calif.) with the following oligonucleotides:

CCAGCTGCTCTGTGGCAGCCAG (SEQ ID NO:20)

CCCAGTCACGACGTTGTAAACGACGGCC (SEQ ID NO:21)

[0193] The reaction was conducted in a standard thermocycler (Perkin-Elmer 9600) for thirty five cycles under the following conditions: 94 C for 1 minute; 94 C for 5 seconds, and 72 C for 5 minutes. This was followed by a final extension at 72 C for 10 minutes. About one microliter of the reaction products was diluted to 50 ul using TE buffer (10 mM TRIS pH 8.0 and 1 mM EDTA).

[0194] To enrich the RACE reaction for the gene of interest, a nested PCR reaction was conducted using about five microliters of the TE solution (containing the RACE reaction products as described in the preceding paragraph) together with the following oligonucleotides:

AACATGCAGCGGCTCGGGG (SEQ ID NO:22)

GGTGACACTATAGAAGAGCTATGACGTCGC (SEQ ID NO:23)

[0195] The nested PCR reaction was incubated in a thermocycler (Perkin-Elmer 9600) using the following protocol: 94 C for one minute; five cycles of 94 C for 5 seconds followed by 72 C for 5 minutes; five cycles of 94 C for five

seconds, followed by 70 C for 5 minutes; and 20-25 cycles of 94 C for 5 seconds followed by 68 C for 5 minutes. After this PCR, the 3' RACE products and the nested PCR products were analyzed using standard agarose gel electrophoresis.

[0196] A PCR product of about 3.3 kb was identified from the nested PCR reaction. This fragment was purified using Qiagen® Gel Extraction Kit (Qiagen, Chatsworth, Calif.) and ligated into the vector pCRII-TOPO (Invitrogen, Carlsbad, Calif.) according to the procedures recommended by the manufacturer. After ligation, the products were transformed into One Shot® *E. coli* cells (Invitrogen, Carlsbad, Calif.) and plated on a LB (Luria broth) plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal. After overnight incubation at 37 C, 12 white colonies and one blue colony were selected, and screened using PCR-Ready Beads® (Amersham-Pharmacia Biotech, Piscataway, N.J.) according to the manufacturer's recommended protocol using oligonucleotide SEQ ID NO:20 together with the following primer:

GTGCTGAGTGCTCTCCATCAGC (SEQ ID NO:24)

[0197] Two colonies were picked that had yielded PCR products of the expected size of about 192 base pairs. These colonies were inoculated into about 3 ml of LB medium containing about 100 ug/ml ampicillin, and were incubated at 37 C. The cultures were placed on a shaker for about 16 hours, and the plasmids were recovered using Qiagen® mini prep columns (Qiagen, Chatsworth, Calif.) according to the manufacturer's protocol. Plasmid DNA was then sequenced as described above.

[0198] A contiguous stretch of DNA of about 3366 nucleotides was assembled by combining the sequence of clone ymrs2-00009-c4 (containing 5' sequence) together with the nested PCR fragment of 3.3 kb containing 3' sequence. Within this contiguous sequence is an open reading frame of 349 amino acids. The nucleotide sequence of this novel mouse gene, referred to as DKR-3, is set forth in FIG. 1. The putative amino acid sequence, as translated from the DNA sequence, is set forth in FIG. 8.

[0199] A BLAST search of the Genbank database using the amino acid sequence of DKR-3 revealed that this open reading frame has homology to a gene known as human rig-like 7-1 mRNA (Genbank accession number AF034208; see also Ligon et al., *J NeuroVirology*, 4: 217-226 [1998]). DKR-3 also has homology to the gene for chicken lens fiber protein clfst4 (Genbank accession number D26311); the overall identity to this protein is about 50 percent with the highest homology in the middle of the protein.

Example 2

Cloning of the Human DKR-3 Gene

[0200] Mouse DKR-3 DNA can be used to search a public EST database for human homologs, resulting in the identification of the following Genbank accession numbers:

[0201] AA628979

[0202] AA349552

[0203] AA633061

[0204] AA351624

[0205] W61032

[0206] T30923

[0207] AA683017

[0208] AA324686

[0209] T08793

[0210] T31076

[0211] R14945

[0212] AA226979

[0213] W45085

[0214] AA424460

[0215] R58671

[0216] R57834

[0217] AF034208

[0218] These EST sequences were analyzed and assembled to create a putative sequence for human DKR-3. Based on this putative sequence, two oligonucleotides were designed for use in PCR in an attempt to clone the human DKR-3 gene. The sequence of these oligonucleotides is:

GAGATGCAGCGGCTTGGGGCCACCC (SEQ ID NO:25)

GCCTGGTCAGCCACGCCTAAAG (SEQ ID NO:26)

[0219] PCR was performed using the Advantage® cDNA PCR kit (Clontech, Palo Alto, Calif.) together with human fetal brain Quick-Clone® cDNA (Clontech). PCR was conducted in a thermocycler (Perkin-Elmer 9600) under the following cycle conditions: 94 C for 2 minute; 94 C for 30 seconds, and 72 C for 2 minutes. Thirty-five cycles were conducted after which samples were treated at 72 C for 10 minutes. A single fragment of about 1150 base pairs was visible when the PCR products were visualized on a 1 percent agarose gel. This fragment was purified using the Qiagen® Gel Extraction Kit (Qiagen, Chatsworth, Calif.) and ligated into the vector pCRII-TOPO (Invitrogen, Carlsbad, Calif.). After ligation, the products were transformed into One Shot *E. coli*® (Invitrogen, Carlsbad, Calif.) and plated on a LB plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal. After overnight incubation at 37 C, 2 white colonies were picked and inoculated into about 3 ml of LB medium containing about 100 ug/ml ampicillin. The cultures were kept on a shaker at about 37 C for about 16 hours. The plasmids were isolated using Qiagen® mini-prep columns (Qiagen, Chatsworth, Calif.) according to the manufacturer's recommended protocol, and the inserts were then sequenced using methods described above.

[0220] The cloned fragment is 1141 bp in length and contains an open reading frame of 350 amino acids. The nucleotide sequence is set forth in FIG. 2, and the putative amino acid sequence, as translated from the DNA sequence, is set forth in FIG. 9. This amino acid sequence is about 80 percent identical to the mouse DKR-3 gene. In addition, human DKR-3 is identical to the human rig-like protein fragment described by Lignon et al., supra between amino acids 157 and 308 of DKR-3. Significantly, the rig-like

protein has an amino terminal start corresponding to amino acid 156 of DKR-3. Rig-like does not appear to be a secreted protein, and the carboxy terminal region of rig-like protein has no homology to human DKR-3. Just as for mouse DKR-3, human DKR-3 is about 54 percent identical to the chicken lens fiber protein clfest. Human DKR-3 appears to be secreted, with a signal peptide cleavage site after either amino acid 20 or 21. Other potential cleavage sites (due to signal peptides or to other endogenous processing sites are after amino acid 16, 22, 32, and/or 41). There appear to be N-linked glycosylation sites at amino acids 96, 106, 121, and 204, which would render them preferable sites for generating substitution mutants. Human DKR-3 and mouse DKR-3 amino acid sequences differ at amino acid positions 6, 7, 11, 24, 27, 29, 30, 32, 33, 39, 81, 89, 93, 99, 101, 103, 109, 113, 115, 123, 126, 142, 156, 157, 162, 165, 173, 175, 191, 197, 198, 201, 203, 245, 247, 259, 283, 287, 292, 294, 295, 296, 298, 299, 304, 310, 311, 312, 314, 315, 329, 330, 334, 335, 336, 339, 340, 341, 342, 343, 345, and 347 (all with respect to the human DKR-3 sequence), which renders these positions preferable for generating human DKR-3 substitution or deletion variants. Based on computer analysis of the amino acid sequence of DKR-3, significant regions of the molecule include the span from about amino acids 21-145 (a potential alpha-helical region and region of potential N-linked glycosylation) such as for example amino acids 21-145, 40-145, 40-150, 45-145, and 45-150, the span from about amino acids 145-350, such as, for example 145-290, 145-300, and 145-350, and the span from about amino acids 300-350 (a second potential alpha-helical region), such as for example amino acids 310-350. Such regions would be suitable fragments of full length DKR-3.

[0221] Northern blot analysis was conducted to assess the tissue specific expression of human DKR-3. A probe for use in Northern blot analysis was prepared by PCR of human fetal brain Quick-Clone® cDNA (Clontech, Palo Alto, Calif.) using the following oligonucleotides:

CCTGCTGCTGGCGGCGGCGGTCCCCACGGC (SEQ ID NO:27)

GCCTGGTCAGCCACGCCTAAAG (SEQ ID NO:28)

[0222] The PCR reaction was conducted in a thermocycler (Perkin-Elmer 9600). PCR conditions were: 94 C for 2 minute; 94 C for 30 seconds, and 72 C for 2 and ½ minutes. Thirty-five cycles were conducted followed by a final extension treatment at 72 C for 10 minutes. PCR products were run on a one percent agarose gel, and a band of about 1100 bp was gel purified using the Qiagen gel extraction kit (Qiagen®, Chatsworth, Calif.), cloned into the vector CR11-TOPO (Invitrogen, Carlsbad, Calif.) and sequenced to confirm that the band contained the human DKR-3 open reading frame minus the amino terminal 10 amino acids.

[0223] About twenty-five nanograms of this probe was denatured by heating to about 100 C for about 5 minutes, followed by placing on ice, and then radioactively labeled with alpha-32P-dCTP using the Rediprime® labeling kit (Amersham, Arlington Heights, Ill.) and following the manufacturer's instructions. A human multiple tissue Northern blot was purchased (Clontech, Palo Alto, Calif.) and was first prehybridized in about 5 ml of Clontech Express® hybridization buffer at about 68 C for 30-60 minutes. After prehybridization, the labeled probe was added to the solu-

tion and allowed to hybridize for about 60 minutes. After hybridization, the blot was first washed with 2×SSC plus 0.05 percent SDS at room temperature for about 30 minutes, then washed with 0.1×SSC plus 0.1 percent SDS at about 65 C for about 30 minutes. The blot was dried briefly and then exposed to a Phosphorimager screen (Molecular Dynamics, Sunnyvale, Calif.). After overnight exposure, the image of the blot was analyzed on a Storm 820 machine (Molecular Dynamics, Sunnyvale, Calif.) with Imagequant software (Molecular Dynamics, Sunnyvale, Calif.).

[0224] The size of the human DKR-3 RNA transcript is about 2.6 kb. The results of the Northern blot analysis indicate that human DKR-3 is highly expressed in adult heart and brain, although weak expression in placenta, adult lung, skeletal muscle, kidney, and pancreas is also apparent. A second smaller transcript is apparent in adult pancreas, and could result from degradation of the full length transcript.

[0225] To evaluate the role of this gene in cancer, a variety of human cancer cell lines were analyzed for the presence or absence of DKR-3 RNA transcript.

[0226] The glioblastoma cell lines Hs 683; A 172; SNB-19; U-87MG; and U-373MG are all from ATCC, and cultured in the media recommended by ATCC.

[0227] Normal human mammary epithelial cells (NMECs) derived from reduction mammoplasties were purchased from Clonetics Corp. (San Diego, Calif.) and the Corriel Institute (Camden, N.J.). The immortalized breast epithelial cell line MCF-10 and the ER+ cell line MCF-7 can be obtained from the American Type Culture Collection. The ER+ BT20T cells were provided by Dr. K. Keyomarsi (N.Y. State Dept. of Health). Immortalized 184A1 and other breast cancer cells including T47-D, ZR75-1, and BT474, MDA-MB-157, MDA-MB-231, MDA-MB-361, MDA-MB-453, MD-MBA-468, HS578T and SKBr3 were all obtained from the American Type Culture Collection (10801 University Blvd., Manassas, Va.).

[0228] NMECs, 184A1 and MCF10 cells were cultured in a modified DME/F12 medium (Gibco/BRL, Grand Island, N.Y.) supplemented with 10 mM Hepes, 2 mM glutamine, 0.1 mM nonessential amino acids, 0.5 mM ethanolamine, 5 mg/ml transferrin, 1 mg/ml Bovine serum albumin, 5.0 ng/ml sodium selenite, 20 ng/ml triiodothyronine, 10 ng/ml EGF, 5 µg/ml insulin and 0.5 µg/ml hydrocortisone (DMEM/F12C) (Ethier et al, *Cancer Letters*, 74: 189-195 [1993]). The ER+ and ER+ breast cancer cells were cultured in Alpha or Richter improved minimal essential medium (MEM) (Gibco/BRL) supplemented with 10 mM Hepes, 2 mM glutamine, 0.1 mM nonessential amino acids, 10 percent fetal bovine serum and 1 µg/ml insulin.

[0229] Normal human bronchial and cervical epithelial cells were purchased from Clonetics Corp. (San Diego, Calif.). Normal cervical epithelial cells were culture in KBM2 (Clonetics Corp. San Diego, Calif.) supplemented with 13 mg/ml bovine pituitary extract, 0.5 µg/ml hydrocortisone, 2 ng/ml EGF, 0.5 mg/ml epinephrine, 0.1 ng/ml retinoic acid, 5 µg/ml transferrin, 6.5 ng/ml triiodothyronine and 5 µg/ml insulin. Normal bronchial epithelial cells were cultured in BEBM (Clonetics Corp., San Diego, Calif.) supplemented with 0.5 mg/ml hydrocortisone, 0.5 ng/ml EGF, 0.5 µg/ml epinephrine, 10 µg/ml transferrin, 5 µg/ml insulin, 0.1 ng/ml retinoic acid and 5.5 ng/ml triiodothyronine.

[0230] The lung cancer cell lines H1299, H23, H358, H441, H460, H520, H522, H727, H146, H209, H446, H510A, H526, and H889 and the cervical cancer cells Caski, C-4-I, MS751, SiHa and C-33-A were all obtained from the American Type Culture Collection. The lung cancer cells were cultured in RPMI (MEM) (Gibco/BRL) supplemented with 10 mM Hepes, 2 mM glutamine, 0.1 mM nonessential amino acids and 10 percent fetal bovine serum (FBS). The cervical cancer cells were cultured in Earles MEM supplemented with 0.1 mM nonessential amino acids, 1 mM sodium pyruvate and 10 percent FBS. All cells were routinely screened for mycoplasma contamination and maintained at about 37° C. in an atmosphere of about 6.5 percent CO₂.

[0231] Total RNA was prepared by lysing cell monolayers in guanidinium isothiocyanate and centrifuging over a 5.7 M CsCl cushion as described previously (Gudas, *Proc. Natl. Acad. Sci USA*, 85: 4705-4709 [1988]). RNA (about 20 µg) was electrophoresed on denaturing formaldehyde gels, transferred to MagnaNT membranes (Micron Separations Inc., Westboro, Mass.) and cross-linked with UV irradiation.

[0232] The blots were prehybridized, probed, and washed under the same conditions as those set forth above for the tissue blot. The blots were dried briefly and then exposed to a Phosphorimager screen (Molecular Dynamics, Sunnyvale, Calif.). After overnight exposure, the image of the blot was analyzed on a Storm 820 machine with Imagequant software (both from Molecular Dynamics).

[0233] The results are shown in FIGS. 15A-15D. As can be seen in FIG. 15A, expression of DKR-3 is decreased in most of the breast cancer cell lines as compared to the normal cell lines. FIG. 15B indicates that DKR-3 expression is decreased in the non-small cell lung cancer cell lines, and in most of the small cell lung cancer cell lines as well. FIG. 15C indicates that expression of DKR-3 is decreased in three glioblastoma cell lines (SNB-19, U-87MG, and U-373MG) that are capable of forming tumors in nude mice (the other two cell lines, Hs 683 and A 172 do not form tumors in nude mice). FIG. 15D indicates that expression of DKR-3 is reduced in cervical cancer cell lines as compared to normal and immortalized cells.

Example 3

Cloning of the Human DKR-1 Gene

[0234] Human and mouse DKR-3 cDNA and amino acid sequences were used to search Genbank using the BLAST program in an attempt to identify DKR-3 related genes.

[0235] A number of EST (expressed sequence tags) were found and were analyzed to determine whether the sequences overlapped. Using the following human EST accessions, a novel gene, termed DKR-1, was predicted.

[0236] AA336797

[0237] R27865

[0238] W39690

[0239] AA043027

[0240] HUM517HO4B

[0241] AA143670

- [0242] W51876
- [0243] N94525
- [0244] AA641247
- [0245] AA137219
- [0246] AA115249
- [0247] AA031969
- [0248] AA136192
- [0249] AA032060
- [0250] AA035583
- [0251] AA207078
- [0252] AA371363
- [0253] AA037322
- [0254] AA088618
- [0255] W46873
- [0256] AA115337
- [0257] AA693679
- [0258] W30750
- [0259] H83554

[0260] PCR was conducted in an attempt to clone the full length gene, and the following two oligonucleotides were used for PCR:

CCCGGACCCTGACTCTGCAGCCG (SEQ ID NO:29)
 GAGGAAAAATAGGCAGTGCAGCACC (SEQ ID NO:30)

[0261] PCR was performed using the Advantage® cDNA PCR kit (Clontech, Palo Alto, Calif.) containing the oligonucleotides listed above and human placenta Quick-Clone® cDNA (Clontech, Palo Alto, Calif.). The reaction was conducted according to the manufacturer's recommendations. Thirty-five cycles of PCR were conducted in a thermocycler (Perkin-Elmer 9600) under the following conditions: 94 C for 2 minutes; 94 C for 30 seconds, and 72 C for 1-½ minutes, followed by a final extension at 72 C for 10 minutes.

[0262] After cycling, the PCR products were analyzed on a one percent agarose gel. A single band of about 1200 base pairs in length was detected after agarose gel electrophoresis. This fragment was purified using the Qiagen® gel extraction kit (Qiagen, Chatsworth, Calif.) and ligated into the vector pCR11-TOPO (Invitrogen, Carlsbad, Calif.) using standard ligation procedures. After ligation, the products were transformed into One Shoot® competent *E. coli* cells according to the procedures recommended by manufacturer (Invitrogen, Carlsbad, Calif.). The transformed *E. coli* cells were plated on a LB plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal.

[0263] After overnight incubation at about 37 C, two white colonies were picked and inoculated into about 3 ml of TB containing 100 ug/ml ampicillin. The culture was incubated at about 37 C for about 16 hours, plasmids were

then recovered using Qiagen® mini-prep columns (Qiagen, Chatsworth, Calif.) and sequenced. Both colonies contained the same insert.

[0264] The insert is 1193 base pairs, and is referred to as human DKR-1. The sequence of this gene is set forth in FIG. 3. This gene contains an open reading frame of 266 amino acids. The amino acid sequence is set forth in FIG. 10. A stop codon is present upstream of the first methionine, indicating the first methionine is likely to be the amino terminus of the protein. Human DKR-1 has a predicted signal peptide with a predicted signal peptide cleavage site between amino acids 19 and 20.

[0265] The gene has about 80 percent homology to the mouse gene dkk-1 (Glinka et al., supra), however the mouse dkk-1 gene is 272 amino acids in length while human DKR-1 is 266 amino acids in length. Human DKR-1 differs from mouse dkk-1 at amino acid positions 3, 4, 5, 7, 8, 10, 12, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 29, 53, 55, 62, 66, 69, 77, 93, 98, 101, 105, 106, 123, 139, 140, 143, 144, 153, 155, 157, 158, 163, 164, 165, 169, 175, 178, 197, 224, and 244. In addition, the alignment of human DKR-1 and mouse dkk-1 shows one gap in human DKR-1 between amino acids 37 and 38, and two gaps between 103 and 104, 146 and 147, and 165 and 166. Glinka et al. state on page 362 of their article that "Coordinates of *Xenopus* dkk family members have been deposited in Genbank with the following accession numbers . . . hdkk-1 AA207078." However, forward three frame translations of AA207078 by the inventors herein showed no homology to the published mouse and *Xenopus* dkk-1 sequences, or to the human DKR-1 sequence, except in the 3' end of this accession, which exhibits a 95 percent identity to human DKR-1 from amino acids 81-179, indicating that AA207078 does not encode full length human dkk-1. Significantly, AA207078 is missing amino acids 1-90 and 180-350 of human DKR-1 which includes the signal peptide and the second cysteine right domain respectively.

Example 4

Cloning of the Mouse DKR-2 Gene

[0266] Genbank accession number AA265561 (a mouse sequence) has homology to both human DKR-1 and human DKR-3 at the amino acid level based primarily on its cysteine pattern.

[0267] To extend this EST sequence in both the 5' and 3' directions, the following oligonucleotides were designed:

GCCACAGTCCCCACCAAGGATCATC (SEQ ID NO:31)
 GATGATCCTTGGTGGGACTGTGGC (SEQ ID NO:32)
 CTGCAAACCAAGTCTCCATCAGGG (SEQ ID NO:33)
 CCCTGATGGAGCACTGGTTTCAG (SEQ ID NO:34)

[0268] Separately, 5' RACE and 3' RACE reactions were performed according to the manufacturer's protocol using mouse heart Marathon-Ready® cDNA and the Advantage® cDNA PCR kit (both from Clontech, Palo Alto, Calif.) and using oligonucleotide SEQ ID NOs: 31 and 34. The RACE reactions were incubated in a thermocycler (Perkin-Elmer 9600) using the following cycling conditions: 94 C for one

minute; five cycles of 94 C for 5 seconds followed by 72 C for 5 minutes; five cycles of 94 C for five seconds, followed by 70 C for 5 minutes; and 20-25 cycles of 94 C for 5 seconds followed by 68 C for 5 minutes.

[0269] To enrich each RACE reaction for the desired product, about one microliter of each of the RACE PCR products was added together, and the mixture was diluted to about 50 ul using TE buffer. About five microliters of this solution were used to conduct nested PCR reactions. The Advantage® cDNA PCR kit (Clontech, Palo Alto, Calif.) and oligonucleotide SEQ ID NOs: 32 and 33 were used for the 5' and 3' nesting reactions, respectively. The nested PCR reactions were incubated in a thermocycler (Perkin-Elmer 9600) using the following program for thirty five cycles: 94 C for 1 minute; 94 C for 5 seconds; and 72 C for 2 minutes. A final extension was then conducted at 72 C for 10 minutes. The PCR products were analyzed using a one percent agarose gel.

[0270] Several fragments ranging from about 500 bp to about 1500 base pairs were obtained from the 5' nested PCR reaction, and two fragments of about 1900 bp and 450 bp were obtained from the 3' nested PCR reaction. These PCR products were purified using the Qiagen® PCR purification kit (Qiagen, Chatsworth, Calif.) and were then ligated into the vector pCR11-TOPO (Invitrogen). The ligation products were transformed into OneShot® *E. coli* cells (Invitrogen, Carlsbad, Calif.), and the cells were then plated on to two X-gal containing plates (one for each reaction) as described above.

[0271] Eight white colonies from each plate were picked and PCR selected via RACE reactions using the Clontech primer AP2 and the oligonucleotide SEQ ID NO:32 (for the 5' RACE) or the oligonucleotide SEQ ID NO:33 (for the 3' RACE). Three colonies from each plate that contained the correct size fragments were cultured, and the plasmids were isolated and sequenced using procedures described above.

[0272] Three clones, 9813302, 9813304 and 9813305 contained sequence which extended the EST sequence in the 5' direction. One clone, 9813308, contained sequence which extended the EST sequence in the 3' direction. A continuous sequence of 2678 base pairs was thus assembled using the sequence of clones 9813308, 9813304, and the EST AA265561. This full length DNA has been termed DKR-2, and the sequence is set forth in FIG. 4. The corresponding amino acid sequence is set forth in FIG. 11.

[0273] Within the amino acid sequence is an open reading frame of 259 amino acids. This protein has approximately 38 percent identity with mouse dkk-1 at the amino acid level. Mouse DKR-2 has a predicted signal peptide with a signal peptide cleavage site between amino acids 33 and 34.

Example 5

Cloning of the Human DKR-2 Gene

[0274] The Genbank EST database was searched using the BLAST program with both DNA and amino acid sequences from human DKR-1 and human DKR-3, and one human EST, W55979, was identified that showed homology to both human DKR-1 and human DKR-3 at the amino acid level based on its cysteine pattern. W55979 is about 88 percent identical to mouse DKR-2 at the DNA level, and about 93 percent identical to mouse DKR-2 at the amino acid level.

[0275] A BLAST search of Genbank W55979 indicated that W55979 has homology to BAC clone number B284B3 (Genbank accession number AC003099). BAC clone B284B3 is 95129 base pairs in length. Three portions of W55979 are homologous to three different regions of BAC clone B284B3, indicating that human DKR-2 has at least three exons. A 3' sequence of 556 bp in length was assembled based on the sequences of both BAC clone B284B3 and W55979, and it was determined that this sequence is the 3' portion of the human ortholog of mouse DKR-2. Within this 3' sequence of human DKR-2 is an open reading frame of 174 amino acids, and a stop codon is present after amino acid 174. This 3' sequence of human DKR-2 is about 97 percent identical to mouse DKR-2.

[0276] To obtain the 5' end sequence of human DKR-2, a 5' RACE reaction was performed using Clontech human heart Marathon-Ready® cDNA and the Advantage® cDNA PCR kit, together with oligonucleotide SEQ ID NO:34. The RACE reaction was performed according to the manufacturer's protocol. The 5' RACE reaction products were then subjected to nesting PCR to enrich for the 5' sequence using the Advantage® cDNA PCR kit and oligonucleotide SEQ ID NO:32. The PCR conditions for both the 5' RACE reaction and the nested PCR reaction were the same as those described in Example 4.

[0277] The nested PCR products were purified using the Qiagen® (Qiagen, Chatsworth, Calif.) PCR purification kit, and were ligated into the vector Zero-Blunt® (Invitrogen, San Diego, Calif.) according to the procedures recommended by the manufacturer. The ligation products were transformed into OneShot® *E. coli* cells which were then plated on X-gal containing plates as described above.

[0278] After overnight culturing, three white colonies were picked and were inoculated into about 3 ml of TB containing about 100 ug/ml ampicillin. The cultures were allowed to grow for about 16 hours, after which the plasmids were isolated using Qiagen® mini-prep columns (Qiagen, Chatsworth, Calif.) according to the manufacturer's protocol. The sequence of each insert was then obtained.

[0279] One of the 5'-RACE clones, termed 9812826, extended the human DKR-2 sequence 5'-terminally. A contiguous sequence of 1531 bp in length was assembled using this clone 9812826 together with the human DKR-2 3' sequence. Within this contiguous sequence is an open reading frame of 259 amino acids. The human DKR-2 gene has a predicted signal peptide of about 33 amino acids, with a predicted cut site between amino acids 33 and 34, and is about 95 percent identical to mouse DKR-2 at the amino acid level. The amino acid positions that differ between human and mouse DKR-2 include (with respect to the numbering of the human sequence) 7, 12, 28, 48, 50, 58, 71, 102, 119, 170, 173, and 191, rendering these positions preferable for generating amino acid substitution or deletion variants.

[0280] An alternative spliced isoform of human DKR-2 was discovered when PCR was conducted using human heart Marathon-Ready® cDNA (Clontech, Palo Alto, Calif.) and the Advantage® cDNA PCR kit (Clontech, Palo Alto, Calif.) together with the following oligonucleotides:

GGGTTGAGGGAACACAATCTGCAAG (SEQ ID NO:36)

GTCTGCAATTGATGATGTTCTCCTCAATGG (SEQ ID NO:37)

[0281] PCR was conducted using parameters set forth in the manufacturer's protocol. PCR products were analyzed by agarose gel electrophoresis, and two PCR products were obtained. The bands corresponding to these products were gel purified as described above, amplified and purified as described above, and then sequenced. One product corresponded to full length DKR-2, however, the other band corresponded to an isoform of DKR-2. This isoform has an open reading frame of 207 amino acids, and appears to be missing an exon. This isoform is referred to as human DKR-2a. The DNA sequence of human DKR-2a is set forth in FIG. 6, and the amino acid sequence as translated from the DNA is set forth in FIG. 13.

Example 6

Cloning of the Human DKR-4 Gene

[0282] A human EST that showed significant homology to human DKR-1 and human DKR-3 on protein level was identified in Genbank. This sequence, Genbank accession number AA565546, has a cysteine pattern that is similar to that of human DKR-1 and human DKR-3.

[0283] A BLAST search of Genbank showed no human ESTs overlapping with AA565546. Therefore, to extend the EST sequence in the 5' direction, a 5' RACE reaction was performed using human heart Marathon-Ready® cDNA (Clontech, Palo Alto, Calif.) together with the Advantage® cDNA PCR kit (Clontech, Palo Alto, Calif.) and the following oligonucleotide:

CCAGGGCCACAGTCGCAACGCTGG (SEQ ID NO:38)

[0284] The RACE reaction was performed according to the protocol provided with the Advantage® kit. After 5' RACE, the products were nested to enrich for the desired 5' sequence using the Advantage® cDNA PCR kit according to the manufacturer's recommendations, together with the following oligonucleotide:

CTCCCTCTTGTCCTTCCTGCCTTG (SEQ ID NO:39)

[0285] After the nested PCR reaction, the products were purified using the Qiagen® PCR purification kit (Qiagen, Chatsworth, Calif.), ligated into the vector pCRII-TOPO (Invitrogen, Carlsbad, Calif.), and transformed into OneShot® *E. coli* cells as described above. After transformation, the cells were plated on a LB plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal.

[0286] After overnight incubation at 37 C, four white colonies were picked from the plate and were inoculated in about 3 ml TB containing about 100 ug/ml ampicillin. The cultures were incubated at about 37 C for about 16 hours. The plasmids were then recovered using Qiagen® mini-prep columns (Qiagen, Chatsworth, Calif.) and sequenced.

[0287] Two clones, termed 9813563 and 9853564, were found to contain the 5' sequence of human DKR-4.

[0288] To obtain the 3' sequence of human DKR-4, a 3' RACE reaction was performed using human uterus Marathon-Ready® cDNA (Clontech, Palo Alto, Calif.) together with the Advantage® cDNA PCR kit (Clontech) and the following oligonucleotide:

CAAGGCAGGAAGGGACAAGAGGGAG (SEQ ID NO:40)

[0289] The 3' RACE reaction was performed according to the manufacturer's recommendations. After the RACE reaction, the products were nested using the Advantage® cDNA PCR kit and the following oligonucleotide:

CCAGCGTTGCGACTGTGGCCCTGG (SEQ ID NO:41)

[0290] The parameters for PCR were 94 C for 1 minute followed by thirty five cycles of 94 C for 5 seconds and then 72 C for 2 minutes, after which a final extension of 70 C for 10 minutes was conducted. After the nesting reaction, the products were analyzed on a 1 percent agarose gel. A single band of about 1200 bp in length was observed. This band was purified from the gel using methods described above, and was then cloned into the vector pCR2.1-TOPO (Invitrogen, Carlsbad, Calif.) and sequenced. Sequence of this band indicated that it contained the 3' sequence of human DKR-4, and this sequence was assembled together with the 5' sequence (from clones 9813563 and 9853564) to generate the full length sequence of human DKR-4. This sequence is set forth in FIGS. 7 (DNA sequence) and 14 (translated amino acid sequence). The polypeptide is 224 amino acids in length and is about 34 percent identical to human DKR-1 at the amino acid sequence level.

Example 7

Expression of Human DKR-1 in Bacteria

[0291] PCR amplification employing the primer pairs and template described below were used to generate a recombinant form of human DKR-1. One primer of each pair introduces a TAA stop codon and a unique BamHI site following the carboxy terminus of the gene. The other primer of each pair introduces a unique NdeI site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique NdeI and BamHI sites of vector pAMG21 (ATCC accession no. 98113) and transformed into the prototrophic *E. coli* host GM121 (deposited with the American Type Culture Collection on XX as accession number XX). Other commonly used *E. coli* expression vectors and host cells are also suitable for expression by one skilled in the art. After transformation, positive clones were selected and examined for expression of the recombinant gene product.

[0292] The construct pAMG21-human DKR-1-24-266 was engineered to be 244 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-His-Pro-Leu-Leu-Gly (SEQ ID NO:43)

Thr-Cys-Gln-Arg-His (SEQ ID NO:44)

[0293] The template used for PCR was human DKR-1 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

GTTCTCCTCATATGCATCCATTATTAGGCGTAAGTG (SEQ ID NO:45)

CCACCTTGAACCTCGGTTCTCAAT

TACGCACTGGATCCTTAGTGTCTCTGACAAGTGTGA (SEQ ID NO:46)

AG

[0294] Transformed *E. coli* strain GM121 containing pAMG21-human DKR-1-24-266 were grown in 2xYT media containing 20 micrograms/ml kanamycin at 30 C until the culture reached an optical density of about 600 nm of about 0.5. Induction of DKR-1 protein expression was achieved by addition of *Vibrio fischeri* synthetic autoinducer to 100 ng/ml final and incubation of the culture at either 30° C. or 37° C. for about 9 hours further with shaking. In addition, as a uninduced control, for each culture no auto-inducer was added to an aliquot of the culture, but the culture was also incubated for about 9 hours further at about 30 C with shaking along with the induced cultures. After about 9 hours, the optical density of cultures were measured at 600 nm, an aliquot of cultures were examined by oil emersion microscopy at 1600x magnification, and aliquots of cultures were pelleted by centrifugation. Bacterial pellets of cultures were processed for SDS-polyacrylamide gel electrophoresis on a 14 percent gel to examine levels of protein produced in crude lysates and for N-terminal sequencing confirmation of the recombinant gene product. The gel was stained with Coomassie blue.

[0295] The results are shown in the photo of FIG. 16. Lane 1 contains molecular weight markers; Lanes 2 and 5

Example 8

Expression of DKR-2 in Bacteria

[0296] PCR amplification employing the primer pairs and templates described below were used to generate various forms of DKR-2. One primer of each pair introduces a TAA stop codon and a unique BamHI site following the carboxy terminus of the gene. The other primer of each pair introduces a unique NdeI site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique NdeI and BamHI sites of vector pAMG21 (ATCC accession no. 98113) and transformed into either prototrophic *E. coli* host GM121 or GM94 (GM 94 was deposited with the ATCC on XX as accession number XX). Other commonly used *E. coli* expression vectors and host cells are also suitable for expression. After transformation, positive clones were selected and examined for expression of the recombinant gene product.

[0297] The construct pAMG21-human DKR-2-26-259 was engineered to be 235 amino acids in length and have the following N-terminal and the following C-terminal amino acids, respectively:

Met-Ser-Gln-Ile-Gly-Ser (SEQ ID NO:47)

Val-Cys-Gln-Lys-Ile. (SEQ ID NO:48)

[0298] The template used for PCR was human DKR-2 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct.

GTTCTCCTCATATGTCTCAAATTGGTAGTTCTCGTGCCAAACTCAACTCCATCAA (SEQ ID NO:49)

G

TACGCACTGGATCCTTAAATTTTCTGACACACATGGAGT (SEQ ID NO:50)

contain crude lysates of uninduced control cells incubated at 30 C; Lanes 3 and 6 are crude lysates of induced cells cultured at 30 C; Lanes 4 and 7 are crude lysates of induced cells cultured at 37 C. The arrow on the left of Lane 1 indicates the expected location of human DKR-1-24-266. As can be seen, large amounts of recombinant protein were observed in crude lysates of induced cultures at both 30° C. and 37° C. (Lanes 3 and 6, and 4 and 7). Microscopic analysis of bacterial cells revealed most cells contained at least one inclusion body, suggesting that at least some of the protein may be produced in the insoluble fraction of *E. coli*.

[0299] The construct pAMG21 mouse DKR-2-26-259 was engineered to be 235 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Ser-Gln-Leu-Gly-Ser (SEQ ID NO:51)

Val-Cys-Gln-Lys-Ile (SEQ ID NO:52)

[0300] The template used for PCR was mouse DKR-2 cDNA, and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct.

GTTCTCCTCATATGTCTCAATTAGGTAGCTCTCGTGCTAAACTCAACTCCATCAA (SEQ ID NO:53)

GTCC

TACGCACTGGATCCTTAGATCTTCTGGCATAATGGAGT (SEQ ID NO:54)

[0301] Transformed *E. coli* GM121 or GM94 containing either pAMG21-human DKR-2-26-259 or pAMG21-mouse DKR-2-26-259 plasmid were grown in 2×YT media containing 20 µg/ml kanamycin at 30° C. until the culture reached an optical density at 600 nm of about 0.5. Induction of DKR-2 protein expression was achieved by addition of *Vibrio fischeri* synthetic autoinducer to 100 ng/ml final and incubation of the culture at either 30 C or 37 C for about 5 or 9 hours further with shaking. In addition, as a uninduced control, for each culture no autoinducer was added to an aliquot of the culture, but the culture was also incubated for about 5 or 9 hours further at 30 C with shaking along with the induced cultures. After either 5 or 9 hours incubation, the optical density of cultures were measured at about 600 nm, an aliquot of cultures were examined by oil emersion microscopy at 1600× magnification, and aliquots of cultures were pelleted by centrifugation. Bacterial pellets of cultures were processed for SDS-polyacrylamide gel electrophoresis on a 14 percent gel to examine levels of protein produced in crude lysates and for N-terminal sequencing confirmation of the recombinant gene product. The gel was stained with Coomassie blue.

[0302] The results are shown in **FIG. 16**, Lanes 8-10 (human DKR-2 polypeptide) and in **FIG. 17** (mouse DKR-2 polypeptide). In **FIG. 16**, Lane 8 contains crude lysate of uninduced control cells; Lane 9 contains crude lysate of induced cells cultured at 30 C, and Lane 10 contains crude lysate of induced cells cultured at 37 C. The arrow to the left of Lane 10 indicates the expected location of human DKR-

2-26-259. As can be seen, significant amounts of polypeptide were generated in the induced cultures whether grown at 30 C or 37 C, while the uninduced cells did not produce a large amount of polypeptide. **FIG. 17** shows the results of polypeptide production of mouse DKR-2-26-259. Lane 1 is molecular weight markers. Lanes 2-4 are one clone of *E. coli* cells transfected with the DKR-2 plasmid, while Lanes 5-7 are a second clone transfected with the same plasmid. Lanes 2 and 5 are crude lysates of uninduced control cells; Lanes 3 and 6 are crude lysates of induced cells cultured at 30 C; and Lanes 4 and 7 are crude lysates of cells cultured at 37

C. The arrows to the left of Lanes 4 and 7 indicate the expected location of the DKR-2 polypeptide. As can be seen, large amounts of recombinant protein were observed in crude lysates of induced cultures at 37 C but not at 30 C. Microscopic analysis of bacterial cells revealed most cells contained at least one inclusion body, suggesting that at least some of the protein may be produced in the insoluble fraction of *E. coli*.

Example 9

Expression of DKR-3 in Bacteria

[0303] PCR amplification employing the primer pairs and templates described below were used to generate various forms of DKR-3. One primer of each pair introduces a TAA stop codon and a unique SacII site following the carboxy terminus of the gene. The other primer of each pair introduces a unique NdeI site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique NdeI and SacII sites of vector pAMG21 (ATCC accession no. 98113) and transformed into the prototrophic *E. coli* host GM121. Other commonly used *E. coli* expression vectors and host cells are also suitable for expression by one skilled in the art. After transformation, positive clones were selected, plasmid DNA was isolated and the sequence of the DKR-3 gene insert was confirmed.

[0304] The construct pAMG21-human DKR-3-23-350 was engineered to be 329 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Pro-Ala-Pro-Thr-Ala (SEQ ID NO:55)

Gly-Gly-Glu-Glu-Ile. (SEQ ID NO:56)

[0305] The template used for PCR was human DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct.

GTTCTCCTCATATGCCTGCTCCAACCTTCGGCTCCAGTCAAGCCCGGCC (SEQ ID NO:57)

TACGCACTCCGCGTTAAATCTCTTCCCTCCAGCA (SEQ ID NO:58)

[0306] The construct pAMG21-human DKR-3-33-350 was engineered to be 319 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Lys-Pro-Gly-Pro-Ala (SEQ ID NO:59)

Gly-Gly-Glu-Glu-Ile (SEQ ID NO:60)

[0307] The template used for PCR was human DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

GTTCTCCTCATATGAAACCAGGTCCAGCCTTAAGCTACCCGAGGAGAGGCCA (SEQ ID NO:61)

TACGCACTCCGCGTTAAATCTCTTCCCTCCAGCA (SEQ ID NO:62)

[0308] The construct pAMG21-human DKR-3-42-350 was engineered to be 310 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Gln-Glu-Glu-Ala-Thr (SEQ ID NO:63)

Gly-Gly-Glu-Glu-Ile (SEQ ID NO:64)

[0309] The template used for PCR was human DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

GTTCCTCTCATATGCAAGAAGAAGCTACTCTGAATGAGATGTTCCGCGAGGTT (SEQ ID NO:65)

TACGCACTCCGCGGTTAAATCTCTTCCCTCCAGCA (SEQ ID NO:66)

[0310] The construct pAMG21-mouse DKR-3-33-349 was engineered to be 318 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Glu-Pro-Gly-Pro-Ala (SEQ ID NO:67)

Gly-Glu-Glu-Glu-Ile (SEQ ID NO:68)

[0311] The template used for PCR was mouse DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

GTTCCTCTCATATGGAACCAGGTCCAGCTTTAAACTACCTCAGGAGGAAGCTA (SEQ ID NO:69)

TACGCACTCCGCGGTTAAATCTCTCTCTCCGCCTA (SEQ ID NO:70)

[0312] Transformed *E. coli* GM121 containing the various pAMG21 DKR-3 plasmids described above were grown in 2×YT media containing 20 micrograms/ml kanamycin at 30° C. until the culture reached an optical density at 600 nm of about 0.5. Induction of DKR-3 polypeptide expression was achieved by addition of *Vibrio fischeri* synthetic auto-inducer to 100 ng/ml final concentration and incubation of the culture at either 30 or 37 C for about 6 hours further with shaking. In addition, as a uninduced control, for each culture no autoinducer was added to an aliquot of the culture, but the culture was also incubated for about 6 hours further at 30 C with shaking along with the induced cultures. After about 6 hours, the optical density of cultures were measured at about 600 nm, an aliquot of cultures were examined by oil emersion microscopy at 1600× magnification, and aliquots of cultures were pelleted by centrifugation. Bacterial pellets of cultures were processed for SDS-polyacrylamide gel electrophoresis to examine levels of protein produced in crude lysates, or bacterial pellets were processed to determine whether the recombinant protein was in the soluble or insoluble fraction of *E. coli* and for N-terminal sequencing confirmation of the recombinant gene product. The results are shown as photos of the SDS gels in **FIGS. 18 and 19**. In **FIG. 18**, Lane 10 is molecular weight markers, and Lanes 1-9 are crude lysates of bacterial cells. Lane 1 is crude lysate of uninduced control cells; Lanes 2, 4, 6, and 8 are crude lysates of induced cells cultured at 30 C; Lanes 3, 5, 7, and 9 are induced cells cultured at 37 C. Lanes 1-5 contain lysates of cells transfected with the pAMG21-human DKR-3-23-350 construct; and Lanes 6-9 contain lysates of cells transfected with the pAMG21-human DKR-3-33-350 construct. The arrows to the left of Lane 2 and the right of Lane 9 indicate the expected location of the DKR-3 polypeptides. **FIG. 19** contains molecular weight markers in Lane 10; Lanes 1-5 are crude lysates of cultured cells transfected with the pAMG21-human DKR-3-42-350 construct; Lanes 6-9

are crude lysates of cells transfected with the pAMG21-mouse DKR-3-33-349 construct. Lanes 1 and 6 are uninduced controls; Lanes 2, 4, 7, and 8 are crude lysates of

induced cells cultured at 30 C (two different clones of each construct); Lanes 3, 5, and 9 are crude lysates of induced cells cultured at 37 C (two separate clones of the human DKR-3-42-350 construct in Lanes 3 and 5). The arrow to the right of Lane 9 indicates the expected location of the mouse DKR-3 polypeptides; the arrow to the left of Lane 4 indicates the expected location of human DKR-3 polypeptide. As can be seen, all DKR-3 constructs produced large amounts of recombinant protein in *E. coli*. No inclusion bodies could be detected by oil emersion microscopy, and

the recombinant polypeptides were mostly found in the soluble fraction of the cells.

Example 10

Expression of DKR-4 in Bacteria

[0313] PCR amplification employing the primer pairs and template described below were used to generate a recombinant form of human DKR-4. One primer of each pair introduces a TAA stop codon and a unique BamHI site following the carboxy terminus of the gene. The other primer of each pair introduces a unique NdeI site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique NdeI and BamHI sites of the vector pAMG21 (ATCC accession no. 98113) and transformed into the prototrophic *E. coli* host GM94. Other commonly used *E. coli* expression vectors and host cells are also suitable for expression. After transformation, positive clones were selected and will be examined for expression of the recombinant gene product.

[0314] The construct pAMG21-human DKR-4-19-224 was engineered to be 207 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Leu-Val-Leu-Asp-Phe (SEQ ID NO:71)

Lys-Ile-Glu-Lys-Leu (SEQ ID NO:72)

[0315] The template used for PCR was human DKR-4 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

(SEQ ID NO:73)
 GTTCTCCTCATATGTTAGTTTGGATTTCACACATCAGGAGCTCT

(SEQ ID NO:74)
 TACGCACTGGATCCTTACAGTTTTTCTATTTTTGGCATACTCTTAATC

[0316] It is anticipated that DKR-4 polypeptide could be prepared using the PCR product as described above for the other DKR polypeptides.

Example 11

Production and Purification of DKR-3 Polypeptide in Mammalian Cells

[0317] Human DKR-3 cDNA was cloned onto the mammalian expression vector pcDNA3.1(-)/mycHis (Invitrogen, Carlsbad, Calif.) and the vector construct was amplified using the Qiagen maxi-prep kit (Qiagen, Chatsworth, Calif.) standard ligation techniques.

[0318] Human embryonic kidney 293T cells (American Type Culture Collection) were cultured in 10 cm dishes, and grown to about 80 percent confluence. The cells were then transfected with the vector construct using the DMRIE-C® liposome formulation (Gibco BRL, Grand Island, N.Y.) as follows. About 240 microliters of DMRIE-C® were added to 8 ml of Optimem medium. About 40 ul (equivalent to about 56 micrograms) of purified vector construct was then added to another 8 ml of Optimem. After mixing and incubation at room temperature for about 15 minutes, 2 ml of this solution was added to each of 8 plates. After about 5 hours, the medium was aspirated and 10 ml of DME medium containing about 10 percent fetal calf serum was added. The cells were incubated 16-18 hours after which the medium was removed and about 10 ml of SF Optimem medium per well without phenol red were added. After about 24 hours, this medium, the "conditioned medium" was harvested, passed over a 0.22 micron filter and stored at 4° C. The cells were then incubated in another 10 ml of SF Optimem per plate. After 24 hours, this medium was collected, filtered and also stored at 4° C.

[0319] The conditioned media was added to a buffer containing 50 mM NaPO₄, pH8, and 250 mM sodium chloride, and passed over a column of nickel-Sephadex (Qiagen, Chatsworth, Calif.). Non-specifically bound proteins were eluted using the same buffer containing 10 mM imidazole, followed by the same buffer containing 20 mM imidazole. DKR-3 was then eluted using 125 mM-250 mM imidazole. Fractions from the column were subjected to 12 percent SDS gel electrophores and silver stained. The results are shown in FIG. 20. Lane 2 contains material that was loaded on to the gel. Lane 3 contains the flow through fraction after loading the column with conditioned medium, Lanes 4, 5, 6, and 7 contain column fractions after treatment with 10, 20, 125, and 250 mM imidazole. Molecular weight standards are shown in Lane 8. As can be seen a single band of protein of the correct molecular weight is seen in Lanes 5 and 6, indicating that this procedure resulted in generation of purified DKR-3 protein (attached to myc and His tags). Smearing of the protein band may be due to glycosylation. Separately, a Western blot was run to confirm that the purified protein did indeed have a His tag (indicating that the fusion protein DKR-3 mycHis had been produced). The

Western blot was prepared using standard procedures and was probed with a polyclonal anti-His-HRP antibody (Invitrogen, Carlsbad, Calif.). A photo of the Western blot is shown in FIG. 21; the Lanes correspond to that for the gel (described immediately above). As can be seen, there is antibody binding in Lanes 2, 5, and 6, indicating that DKR-3 mycHis was loaded on to the column and was eluted in the 20 and 125 mM imidazole washes.

Example 12

Anchorage Independent Growth Assay

[0320] A distinguishing feature of many cancer cell lines is their ability to grow in an anchorage independent manner. Whereas normal cells will only grow and divide until they come in contact with their neighbors, cancer cells continue to grow and divide after contact, thereby forming tumors. Thus, one assay for cancer cell growth inhibitor compounds measures the ability of cancer cells to grow and divide in the presence of the compound. There are many ways known to the skilled artisan in which this assay can be conducted, however two preferred methods are set forth below.

[0321] A. Stably Transfected Cell Assay

[0322] In this procedure, any human cancer cell line that does not express the DKR gene to be tested (either human DKR-1, 2, 3, 4, or a fragment or variant thereof) is transfected with the DKR gene under evaluation, where the DKR gene is inserted into a vector such as pcDNA3.1 (Invitrogen, Carlsbad, Calif.) or other suitable mammalian expression vector. Transfection can be conducted as described herein. The transfected cancer cells are cultured to generate a stably transfected cell line. Once a stably transfected cell line has been established, the cells are added to Noble or equivalent agar (about 0.35 percent) prepared in standard mammalian cell culture medium such as RPMI. The cell/agar solution is poured on to petri plates containing solidified agar ban (about 0.5 percent agar). Colony formation is evaluated daily to determine the rate of growth of the cells, and culture medium is added to each plate as needed. Separately, the same cells are transfected with vector only (containing no DKR gene). These "control" cells are then treated in an identical manner to the DKR gene containing cells and can be used as a standard of comparison for the DKR gene containing cells.

[0323] Examples of suitable cancer cell lines for conducting this assay include, without limitation, human breast cancer cell line MCF7 and the glioblastoma cell line U-87MG.

[0324] B. Protein Assay

[0325] An alternate or additional assay to measure the growth of cancer cell lines treated with a DKR polypeptide is as follows. Any human cancer cell line not expressing the DKR polypeptide under evaluation can be cultured and prepared with an agar solution as described above. The cells can then be plated as described, and a solution of DKR polypeptide (either full length, or a fragment or variant thereof) in culture medium can be added to the agar either daily, every other day, or once per week for three weeks. Typically, a concentration of about 10 nM will be added, although a series of dilutions ranging from 1 nM to 1 mM

can be used. Control plates will receive a solution of culture medium only. The plates can be monitored daily for up to about three weeks to evaluate cell colony formation. After three weeks, control and experimental plates can be compared for the number and size of cell colonies. It is anticipated that those plates receiving DKR polypeptide that is biologically active will have fewer cell colonies, and the colonies will be smaller, as compared to control plates.

Example 13

In Vivo Tumor Assay

[0326] The ability of each DKR polypeptide to inhibit tumor growth in vivo can be evaluated as follows. Tumor cells not expressing the DKR gene under evaluation can be transfected using procedures described herein with a DKR nucleic acid construct encoding a full length DKR gene, or a fragment or variant thereof. The transfected cells can be maintained in culture (as described herein) until ready for use.

[0327] Male or female athymic nude mice (Charles River Labs, Boston, Mass.) are kept in a sterile environment. The mice are then injected with about 2×10^6 cells (either DKR transfected cells or control "vector only" transfected cells) in a total volume of about 0.1 ml can be injected sub-cutaneously. The mice can then be examined daily for appearance of (a) tumor(s) and for the size of the tumor. Preferably, the mice will be examined for up to about six months so as to provide time for tumor growth (and regression where DKR polypeptides are effective at decreasing tumor growth). The tumor(s), where present, can then be removed, weighed and examined for (1) the presence of DKR polypeptide, and (2) morphology. Tumors from mice containing DKR construct transfected cells can be compared to tumors from mice containing cells transfected with vector only. It is anticipated that DKR polypeptides, due to their similarity with dkk-1, a potent wnt8 antagonist, will be able to decrease the size of the tumor as compared with controls.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 78

<210> SEQ ID NO 1

<211> LENGTH: 1050

<212> TYPE: DNA

<213> ORGANISM: Mouse

<400> SEQUENCE: 1

```

atgcagcggc tcgggggtat ttgctgtgt aactgctgg cggcggcggc cccactgct      60
cctgtcctt cccgacggc cacttgact ccggcggagc cgggccagc tctcaactac      120
cctcaggagg aagctacgt caatgagatg ttctgagagg tggaggagct gatggaagac      180
actcagcaca aactgcgcag tgccgtggag gagatggagg cggaagaagc agctgctaaa      240
acgtcctctg aggtgaacct ggcaagctta cctcccaact atcacaatga gaccagcacg      300
gagaccaggg tgggaataa cacagtccat gtgcaccagg aagttcaca gataaccaac      360
aaccagagtg gacaggtggt cttttctgag acagtcatta catctgtagg ggatgaagaa      420
ggcaagagga gccatgaatg tatcattgat gaagactgtg ggcccaccag gtactgccag      480
ttctccagct tcaagtacac ctgccagcca tgccgggacc agcagatgct atgcacccga      540
gacagtgagt gctgtggaga ccagctgtgt gcctggggtc actgcacca aaaggccacc      600
aaagtggtgca atgggacat ctgtgacaac cagagggatt gccagcctgg cctgtgttgt      660
gccttccaaa gaggcctgct gttcccctgt tgcacacccc tgcccgtgga gggagagctc      720
tgccatgacc ccaccagcca gctgctggat ctcatcacct gggaactgga gcctgaagga      780
gctttggacc gatgccctg cgccagtggc ctctatgcc agccacacag ccacagtctg      840
gtgtacatgt gcaagccagc cttcgtgggc agccatgacc acagtgagga gagccagctg      900
cccagggagg ccccgatga gtacgaagat gttggcttca taggggaagt gcgccaggag      960
ctggaagacc tggagcggag cctagcccag gagatggcat ttgaggggccc tgcccctgtg     1020
gagtcactag gcggagagga ggagatttag                                     1050

```

<210> SEQ ID NO 2

-continued

<211> LENGTH: 1053
 <212> TYPE: DNA
 <213> ORGANISM: Human

<400> SEQUENCE: 2

```

atgcagcggc ttggggccac cctgctgtgc ctgctgctgg cggcggcggg cccacgccc      60
cccgcgcccg ctccgacggc gacctcggct ccagtcaagc ccggcccggc tctcagctac      120
ccgcaggagg aggccaccct caatgagatg ttccgcgagg ttgaggaact gatggaggac      180
acgcagcaca aattgcgcag cgcggtggaa gagatggagg cagaagaagc tgctgctaaa      240
gcatcatcag aagtgaacct ggcaactta cctcccagct atcacaatga gaccaacaca      300
gacacgaagg ttggaaataa taccatccat gtgcaccgag aaattcaca gataaccaac      360
aaccagactg gacaaatggt cttttcagag acagttatca catctgtggg agacgaagaa      420
ggcagaagga gccacgagtg catcatcgac gaggactgtg ggcccagcat gtactgccag      480
tttgccagct tccagtacac ctgccagcca tgcgggggcc agaggatgct ctgcacccgg      540
gacagtgagt gctgtggaga ccagctgtgt gtctggggtc actgcaccaa aatggccacc      600
aggggcagca atgggaccat ctgtgacaac cagagggact gccagccggg gctgtgctgt      660
gccttcaga gaggcctgct gttccctgtg tgcacacccc tgcccgtgga gggcgagctt      720
tgccatgacc ccgccagccg gcttctggac ctcatcacct gggagctaga gcctgatgga      780
gccttggaac gatgcccttg tgccagtggc ctccctgtgc agcccacag ccacagcctg      840
gtgtatgtgt gcaagccgac ctctcgtggg agccgtgacc aagatgggga gatcctgctg      900
cccagagagg tccccgatga gtatgaagtt ggcagcttca tggaggaggt gcgccaggag      960
ctggaggacc tggagaggag cctgactgaa gagatggcgc tgggggagcc tgcggctgcc      1020
gccgctgcac tgctgggagg ggaagagatt tag                                     1053

```

<210> SEQ ID NO 3
 <211> LENGTH: 801
 <212> TYPE: DNA
 <213> ORGANISM: Human

<400> SEQUENCE: 3

```

atgatggctc tgggcgcagc gggagctacc cgggtccttg tcgcgatggt agcggcggct      60
ctcggcggcc accctctgct gggagttagc gccaccttga actcggttct caattccaac      120
gctatcaaga acctgcccc accgctgggc ggcgtgcggg ggcacccagg ctctgcagtc      180
agcgccgcgc cggaatcct gtaccgggc ggaataagt accagaccat tgacaactac      240
cagccgtacc cgtgcgcaga ggacgaggag tgcggcactg atgagtactg cgctagtccc      300
acccgaggag gggacgcggg cgtgcaaacc tgtctcgctt gcaggaagcg ccgaaaacgc      360
tgcatgcgtc acgctatgtg ctgccccggg aattactgca aaaatggaat atgtgtgtct      420
tctgatcaaa atcatttcgg aggagaaatt gaggaacca tcaactgaaag ctttgtaat      480
gatcatagca ccttgtagtg gtattccaga agaaccacct tgtcttcaaa aatgtatcac      540
accaaaggac aagaaggttc tgtttgtctc cggtcacgag actgtgcctc aggattgtgt      600
tgtgctagac acttctgtgc caagatctgt aaacctgtcc tgaagaagg tcaagtgtgt      660
accaagcata ggagaaaagg ctctcatgga ctagaaatat tccagcgttg ttactgtgga      720
gaaggctctg cttgccggat acagaaagat caccatcaag ccagtaattc ttctaggctt      780

```

-continued

cacacttgtc agagacacta a 801

<210> SEQ ID NO 4
 <211> LENGTH: 780
 <212> TYPE: DNA
 <213> ORGANISM: Mouse

<400> SEQUENCE: 4

atggccgcgc tgatgcgggt caaggattca tcccgtgcc ttctcctact ggccgcgggtg 60
 ctgatggtgg agagctcaca gctaggcagc tcgcggggcca aactcaactc catcaagtcc 120
 tctctaggag gggagactcc tgctcagtca gccaacccgat ctgcaggcat gaaccaagga 180
 ctggccttcg gcggcagtaa gaaggcaca agcctggggc aggcctaccc ttgcagcagt 240
 gataaggaat gtgaagttag aagatactgc cacagtcccc accaaggatc atcagcctgc 300
 atgctctgta ggaggaaaaa gaaacgatgc cacagagatg ggatgtgttg ccttggtacc 360
 cgctgcaata atggaatctg catcccgatc actgagagca tcctcaccac acatatccca 420
 gctctggatg gcacccggca tagagatcgc aaccatggtc actattccaa ccatgacctg 480
 ggatggcaga atctaggaag gccacactcc aagatgcctc atataaaagg acatgaagga 540
 gaccatgcc tacggatcgc agactgcatt gatgggtttt gttgtgctcg ccacttcttg 600
 accaaaaatc gcaaaccagt gctccatcag ggggaagtct gtaccaaaca acgcaagaag 660
 ggttcgcacg ggctggagat tttccagagg tgtgactgtg caaaggccct gtcttgcaaa 720
 gtgtgaaaag atgccaccta ctcttccaaa gccagactcc atgtatgcca gaagatctga 780

<210> SEQ ID NO 5
 <211> LENGTH: 780
 <212> TYPE: DNA
 <213> ORGANISM: Human

<400> SEQUENCE: 5

atggccgcgt tgatgcggag caaggattcg tcctgctgcc tgctcctact ggccgcgggtg 60
 ctgatggtgg agagctcaca gatcggcagt tcgcggggcca aactcaactc catcaagtcc 120
 tctctgggag gggagacgcc tggtcaggcc gccaatcgat ctgcgggcat gtaccaagga 180
 ctggcattcg gcggcagtaa gaaggcaca aacctggggc aggcctaccc ttgtagcagt 240
 gataaggagt gtgaagttag gaggtattgc cacagtcccc accaaggatc atcggcctgc 300
 atggtgtgtc ggagaaaaa gaagcgtgc caccgagatg gcatgtgtcg cccagtagc 360
 cgctgcaata atggcatctg tatcccgatt actgaaagca tcttaacccc tcacatcccg 420
 gctctggatg gtactcggca cagagatcga aaccacggtc attactcaaa ccatgacttg 480
 ggatggcaga atctaggaag accacacact aagatgtcac atataaaagg gcatgaagga 540
 gaccctgcc tacgatcgc agactgcatt gaagggtttt gctgtgctcg tcatttcttg 600
 accaaaaatc gcaaaccagt gctccatcag ggggaagtct gtaccaaaca acgcaagaag 660
 ggttctcatg ggctggaat tttccagcgt tgcgactgtg cgaaggccct gtcttgcaaa 720
 gtatgaaaag atgccaccta ctcttccaaa gccagactcc atgtgtgtca gaaaatttga 780

<210> SEQ ID NO 6
 <211> LENGTH: 624
 <212> TYPE: DNA
 <213> ORGANISM: Human

-continued

<400> SEQUENCE: 6

```

atggccgcgt tgatcggag caaggattcg tcctgctgcc tgctcctact ggccgcggtg      60
ctgatggtgg agagctcaca gatcggcagt tcgcggggcca aactcaactc catcaagtcc      120
tctctgggcg gggagacgcc tggtcaggcc gccaatcgat ctgcgggcat gtaccaagga      180
ctggcattcg gggcgagtaa gaagggcaaa aacctggggc aggcctaccc ttgtagcagt      240
gataaggagt gtgaagttag gaggtattgc cacagtcccc accaaggatc atcggcctgc      300
atggtgtgtc ggagaaaaaa gaagcgctgc caccgagatg gcatgtgctg cccagttacc      360
cgctgaata atgggcatga aggagacccc tgcctacgat catcagactg cattgaaggg      420
ttttgtgtg ctcgtcatth ctggacaaa atctgcaaac cagtgtctca tcagggggaa      480
gtctgtacca aacaacgcaa gaagggttct catgggctgg aaattttcca gcgttgcgac      540
tgtgcgaagg gcctgtcttg caaagtatgg aaagatgcca cctactctc caaagccaga      600
ctccatgtgt gtcagaaaat ttga                                           624

```

<210> SEQ ID NO 7

<211> LENGTH: 675

<212> TYPE: DNA

<213> ORGANISM: Human

<400> SEQUENCE: 7

```

atggtggcgg ccgtcctgct ggggctgagc tggctctgct ctcccctggg agctctggtc      60
ctggacttca acaacatcag gagctctgct gacctgcatg gggcccgaa gggctcacag      120
tgctgtctg acacggactg caataccaga aagttctgcc tccagccccg cgatgagaag      180
ccgttctgtg ctacatgtcg tgggttgagg aggaggtgcc agcgagatgc catgtgctgc      240
cctgggacac tctgtgtgaa cgatgtttgt actacgatgg aagatgcaac cccaatatta      300
gaaaggcagc ttgatgagca agatggcaca catgcagaag gaacaactgg gcacccagtc      360
caggaaaacc aacccaaaag gaagccaagt attaagaaat cacaaggcag gaagggacaa      420
gaggagaaaa gttgtctgag aacttttgac tgtggccctg gactttgctg tgctcgtcat      480
ttttggacga aaattttaa gccagtcctt ttggagggac aggtctgctc cagaagaggg      540
cataaagaca ctgctcaagc tccagaaatc ttccagcgtt gcgactgtgg ccctggacta      600
ctgtgtcgaa gccaattgac cagcaatcgg cagcatgctc gattaagagt atgccaaaaa      660
atagaaaagc tataa                                           675

```

<210> SEQ ID NO 8

<211> LENGTH: 349

<212> TYPE: PRT

<213> ORGANISM: Mouse

<400> SEQUENCE: 8

```

Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala
 1             5             10             15
Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala
                20             25             30
Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn
 35             40             45
Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
 50             55             60

```

-continued

```

Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys
 65          70          75          80

Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn
          85          90          95

Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His
 100          105          110

Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe
 115          120          125

Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser
 130          135          140

His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln
 145          150          155          160

Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met
 165          170          175

Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp
 180          185          190

Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys
 195          200          205

Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
 210          215          220

Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
 225          230          235          240

Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu
 245          250          255

Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
 260          265          270

Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe
 275          280          285

Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala
 290          295          300

Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu
 305          310          315          320

Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly
 325          330          335

Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
 340          345

```

```

<210> SEQ ID NO 9
<211> LENGTH: 350
<212> TYPE: PRT
<213> ORGANISM: Human

```

```

<400> SEQUENCE: 9

```

```

Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala
 1          5          10          15

Val Pro Thr Ala Pro Ala Pro Ala Pro Thr Ala Thr Ser Ala Pro Val
 20          25          30

Lys Pro Gly Pro Ala Leu Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn
 35          40          45

Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
 50          55          60

Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys
 65          70          75          80

```

-continued

Ala Ser Ser Glu Val Asn Leu Ala Asn Leu Pro Pro Ser Tyr His Asn
 85 90 95
 Glu Thr Asn Thr Asp Thr Lys Val Gly Asn Asn Thr Ile His Val His
 100 105 110
 Arg Glu Ile His Lys Ile Thr Asn Asn Gln Thr Gly Gln Met Val Phe
 115 120 125
 Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Arg Arg Ser
 130 135 140
 His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln
 145 150 155 160
 Phe Ala Ser Phe Gln Tyr Thr Cys Gln Pro Cys Arg Gly Gln Arg Met
 165 170 175
 Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp
 180 185 190
 Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys
 195 200 205
 Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
 210 215 220
 Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
 225 230 235 240
 Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu
 245 250 255
 Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
 260 265 270
 Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe
 275 280 285
 Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val
 290 295 300
 Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu
 305 310 315 320
 Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Gly Glu
 325 330 335
 Pro Ala Ala Ala Ala Ala Ala Leu Leu Gly Gly Glu Glu Ile
 340 345 350

<210> SEQ ID NO 10

<211> LENGTH: 266

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 10

Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met
 1 5 10 15
 Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr
 20 25 30
 Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro
 35 40 45
 Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro
 50 55 60
 Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr
 65 70 75 80
 Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr

-continued

85					90					95					
Cys	Ala	Ser	Pro	Thr	Arg	Gly	Gly	Asp	Ala	Gly	Val	Gln	Ile	Cys	Leu
			100					105					110		
Ala	Cys	Arg	Lys	Arg	Arg	Lys	Arg	Cys	Met	Arg	His	Ala	Met	Cys	Cys
		115					120					125			
Pro	Gly	Asn	Tyr	Cys	Lys	Asn	Gly	Ile	Cys	Val	Ser	Ser	Asp	Gln	Asn
	130					135					140				
His	Phe	Arg	Gly	Glu	Ile	Glu	Glu	Thr	Ile	Thr	Glu	Ser	Phe	Gly	Asn
	145					150					155				160
Asp	His	Ser	Thr	Leu	Asp	Gly	Tyr	Ser	Arg	Arg	Thr	Thr	Leu	Ser	Ser
				165					170						175
Lys	Met	Tyr	His	Thr	Lys	Gly	Gln	Glu	Gly	Ser	Val	Cys	Leu	Arg	Ser
			180					185					190		
Ser	Asp	Cys	Ala	Ser	Gly	Leu	Cys	Cys	Ala	Arg	His	Phe	Trp	Ser	Lys
		195					200					205			
Ile	Cys	Lys	Pro	Val	Leu	Lys	Glu	Gly	Gln	Val	Cys	Thr	Lys	His	Arg
	210					215					220				
Arg	Lys	Gly	Ser	His	Gly	Leu	Glu	Ile	Phe	Gln	Arg	Cys	Tyr	Cys	Gly
	225					230					235				240
Glu	Gly	Leu	Ser	Cys	Arg	Ile	Gln	Lys	Asp	His	His	Gln	Ala	Ser	Asn
				245					250						255
Ser	Ser	Arg	Leu	His	Thr	Cys	Gln	Arg	His						
			260					265							

<210> SEQ ID NO 11
 <211> LENGTH: 259
 <212> TYPE: PRT
 <213> ORGANISM: Mouse

<400> SEQUENCE: 11

Met	Ala	Ala	Leu	Met	Arg	Val	Lys	Asp	Ser	Ser	Arg	Cys	Leu	Leu	Leu
	1			5					10					15	
Leu	Ala	Ala	Val	Leu	Met	Val	Glu	Ser	Ser	Gln	Leu	Gly	Ser	Ser	Arg
			20					25					30		
Ala	Lys	Leu	Asn	Ser	Ile	Lys	Ser	Ser	Leu	Gly	Gly	Glu	Thr	Pro	Ala
		35					40					45			
Gln	Ser	Ala	Asn	Arg	Ser	Ala	Gly	Met	Asn	Gln	Gly	Leu	Ala	Phe	Gly
	50					55					60				
Gly	Ser	Lys	Lys	Gly	Lys	Ser	Leu	Gly	Gln	Ala	Tyr	Pro	Cys	Ser	Ser
	65					70			75					80	
Asp	Lys	Glu	Cys	Glu	Val	Gly	Arg	Tyr	Cys	His	Ser	Pro	His	Gln	Gly
				85					90					95	
Ser	Ser	Ala	Cys	Met	Leu	Cys	Arg	Arg	Lys	Lys	Lys	Arg	Cys	His	Arg
			100				105						110		
Asp	Gly	Met	Cys	Cys	Pro	Gly	Thr	Arg	Cys	Asn	Asn	Gly	Ile	Cys	Ile
		115					120					125			
Pro	Val	Thr	Glu	Ser	Ile	Leu	Thr	Pro	His	Ile	Pro	Ala	Leu	Asp	Gly
	130					135					140				
Thr	Arg	His	Arg	Asp	Arg	Asn	His	Gly	His	Tyr	Ser	Asn	His	Asp	Leu
	145					150			155						160
Gly	Trp	Gln	Asn	Leu	Gly	Arg	Pro	His	Ser	Lys	Met	Pro	His	Ile	Lys
			165						170					175	

-continued

Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Asp Gly
 180 185 190

Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
 195 200 205

His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
 210 215 220

Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
 225 230 235 240

Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
 245 250 255

Gln Lys Ile

<210> SEQ ID NO 12
 <211> LENGTH: 259
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 12

Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys Cys Leu Leu Leu
 1 5 10 15

Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile Gly Ser Ser Arg
 20 25 30

Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly Glu Thr Pro Gly
 35 40 45

Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly Leu Ala Phe Gly
 50 55 60

Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr Pro Cys Ser Ser
 65 70 75 80

Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
 85 90 95

Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
 100 105 110

Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile
 115 120 125

Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly
 130 135 140

Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu
 145 150 155 160

Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys
 165 170 175

Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly
 180 185 190

Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
 195 200 205

His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
 210 215 220

Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
 225 230 235 240

Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
 245 250 255

Gln Lys Ile

-continued

<210> SEQ ID NO 13
 <211> LENGTH: 207
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 13

```

Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys Cys Leu Leu Leu
 1           5           10          15

Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile Gly Ser Ser Arg
 20          25          30

Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly Glu Thr Pro Gly
 35          40          45

Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly Leu Ala Phe Gly
 50          55          60

Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr Pro Cys Ser Ser
 65          70          75          80

Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
 85          90          95

Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
100          105          110

Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly His Glu Gly
115          120          125

Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly Phe Cys Cys Ala
130          135          140

Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu His Gln Gly Glu
145          150          155          160

Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly Leu Glu Ile Phe
165          170          175

Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys Val Trp Lys Asp
180          185          190

Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys Gln Lys Ile
195          200          205

```

<210> SEQ ID NO 14
 <211> LENGTH: 224
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 14

```

Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu
 1           5           10          15

Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu
 20          25          30

His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn
 35          40          45

Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala
 50          55          60

Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys
 65          70          75          80

Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala
 85          90          95

Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala
100          105          110

Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys

```

-continued

115	120	125
Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser		
130	135	140
Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His		
145	150	155
Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys		
165	170	175
Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln		
180	185	190
Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser		
195	200	205
Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu		
210	215	220

<210> SEQ ID NO 15
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide primer

<400> SEQUENCE: 15

ggaaggaaaa aagcggccgc aacannnnnn nnn

33

<210> SEQ ID NO 16
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide adapter

<400> SEQUENCE: 16

tcgacccacg cgtccg

16

<210> SEQ ID NO 17
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide adapter

<400> SEQUENCE: 17

gggtgcgcag gc

12

<210> SEQ ID NO 18
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide primer

<400> SEQUENCE: 18

actagctcca gtgatctc

18

<210> SEQ ID NO 19
 <211> LENGTH: 18
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 19
cgtcattggt ctcgttcc 18

<210> SEQ ID NO 20
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 20
ccagctgctc tgtggcagcc cag 23

<210> SEQ ID NO 21
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 21
cccagtcacg acgttgtaaa acgacggcc 29

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 22
aacatgcagc ggctcggggg 20

<210> SEQ ID NO 23
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 23
ggtgacacta tagaagagct atgacgtcgc 30

<210> SEQ ID NO 24
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 24
gtgctgagtg tcttccatca gc 22

<210> SEQ ID NO 25

-continued

<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probes

<400> SEQUENCE: 25
gagatgcagc ggcttggggc cacc
25

<210> SEQ ID NO 26
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probes

<400> SEQUENCE: 26
gcctggtcag cccacgccta aag
23

<210> SEQ ID NO 27
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probes

<400> SEQUENCE: 27
cctgctgctg gcggcggcgg tccccacggc
30

<210> SEQ ID NO 28
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probes

<400> SEQUENCE: 28
gcctggtcag cccacgccta aag
23

<210> SEQ ID NO 29
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probes

<400> SEQUENCE: 29
cccgaccct gactctgcag ccg
23

<210> SEQ ID NO 30
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probes

<400> SEQUENCE: 30
gaggaaaaat aggcagtgcg gcacc
25

-continued

<210> SEQ ID NO 31
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primers

<400> SEQUENCE: 31

gccacagtcc ccaccaagga tcatc 25

<210> SEQ ID NO 32
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primers

<400> SEQUENCE: 32

gatgatcctt ggtggggact gtggc 25

<210> SEQ ID NO 33
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primers

<400> SEQUENCE: 33

ctgcaaacca gtgctccatc aggg 24

<210> SEQ ID NO 34
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primers

<400> SEQUENCE: 34

ccctgatgga gcactggttt gcag 24

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer

<400> SEQUENCE: 35

gctataccaa gcatacaatc 20

<210> SEQ ID NO 36
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 36

-continued

gggttgaggg aacacaatct gcaag 25

<210> SEQ ID NO 37
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 37

gtctgcaatt gatgatgttc ctcaatgg 28

<210> SEQ ID NO 38
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 38

ccagggccac agtcgcaacg ctgg 24

<210> SEQ ID NO 39
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 39

ctccctcttg tcccttctg ccttg 25

<210> SEQ ID NO 40
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 40

caaggcagga agggacaaga gggag 25

<210> SEQ ID NO 41
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 41

ccagcgttgc gactgtggcc ctgg 24

<210> SEQ ID NO 42
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide primer/adaptor

-continued

<400> SEQUENCE: 42

gactagtcttct agatcgcgag cggccgccct tttttttttt tttt 44

<210> SEQ ID NO 43

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 43

Met His Pro Leu Leu Gly
1 5

<210> SEQ ID NO 44

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 44

Thr Cys Gln Arg His
1 5

<210> SEQ ID NO 45

<211> LENGTH: 59

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 45

gttctcctca tatgcatcca ttattaggcg taagtgccac cttgaactcg gttctcaat 59

<210> SEQ ID NO 46

<211> LENGTH: 38

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 46

tacgcactgg atccttagtg tctctgacaa gtgtgaag 38

<210> SEQ ID NO 47

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 47

Met Ser Gln Ile Gly Ser
1 5

<210> SEQ ID NO 48

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 48

Val Cys Gln Lys Ile
1 5

<210> SEQ ID NO 49

-continued

<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 49
gttctcctca tatgtctcaa attggtagtt ctcgtgccaa actcaactcc atcaag 56

<210> SEQ ID NO 50
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 50
tacgcactgg atccttaaatt tttctgacac acatggagt 39

<210> SEQ ID NO 51
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mouse

<400> SEQUENCE: 51
Met Ser Gln Leu Gly Ser
1 5

<210> SEQ ID NO 52
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mouse

<400> SEQUENCE: 52
Val Cys Gln Lys Ile
1 5

<210> SEQ ID NO 53
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 53
gttctcctca tatgtctcaa ttaggtagct ctcgtgctaa actcaactcc atcaagtcc 59

<210> SEQ ID NO 54
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 54
tacgcactgg atccttagat cttctggcat acatggagt 39

<210> SEQ ID NO 55
<211> LENGTH: 6
<212> TYPE: PRT

-continued

<213> ORGANISM: Human

<400> SEQUENCE: 55

Met Pro Ala Pro Thr Ala
1 5

<210> SEQ ID NO 56

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 56

Gly Gly Glu Glu Ile
1 5

<210> SEQ ID NO 57

<211> LENGTH: 54

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 57

gttctcctca tatgcctgct ccaactgcaa cttcggctcc agtcaagccc ggcc 54

<210> SEQ ID NO 58

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 58

tacgcactcc gcggttaaatt ctcttcccct cccagca 37

<210> SEQ ID NO 59

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 59

Met Lys Pro Gly Pro Ala
1 5

<210> SEQ ID NO 60

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Human

<400> SEQUENCE: 60

Gly Gly Glu Glu Ile
1 5

<210> SEQ ID NO 61

<211> LENGTH: 54

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 61

-continued

gtttctcctca tatgaaacca ggtccagcct taagctaccc gcaggaggag gccca 54

<210> SEQ ID NO 62
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 62

tacgcactcc gcggttaaat ctcttcccct cccagca 37

<210> SEQ ID NO 63
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 63

Met Gln Glu Glu Ala Thr
1 5

<210> SEQ ID NO 64
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 64

Gly Gly Glu Glu Ile
1 5

<210> SEQ ID NO 65
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 65

gtttctcctca tatgcaagaa gaagctactc tgaatgagat gttccgcgag gtt 53

<210> SEQ ID NO 66
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 66

tacgcactcc gcggttaaat ctcttcccct cccagca 37

<210> SEQ ID NO 67
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mouse

<400> SEQUENCE: 67

Met Glu Pro Gly Pro Ala
1 5

-continued

<210> SEQ ID NO 68
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mouse

<400> SEQUENCE: 68

Gly Glu Glu Glu Ile
1 5

<210> SEQ ID NO 69
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 69

gttctcctca tatggaacca ggtccagctt taaactaccc tcaggaggaa gcta 54

<210> SEQ ID NO 70
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 70

tacgcactcc gcggttaaatt ctctctctct cgccta 37

<210> SEQ ID NO 71
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 71

Met Leu Val Leu Asp Phe
1 5

<210> SEQ ID NO 72
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 72

Lys Ile Glu Lys Leu
1 5

<210> SEQ ID NO 73
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 73

gttctcctca tatgttagtt ttggatttca acaacatcag gagctct 47

<210> SEQ ID NO 74
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide probe

<400> SEQUENCE: 74

taccgactgg atccttacag tttttctatt ttttggcata ctcttaatc 49

<210> SEQ ID NO 75

<211> LENGTH: 798

<212> TYPE: DNA

<213> ORGANISM: Human

<400> SEQUENCE: 75

atgatggctc tgggtgctgc tgggtgctacc cgtgttttcg ttgctatggg tgctgctgct 60
ctgggtgggc acccgctgct ggggtgttcc gctaccctga actccgttct gaactccaac 120
gctatcaaaa acctgccgcc gccgctgggt ggtgctgctg gtcacccggg ttccgctggt 180
tccgctgctc cgggtatcct gtaccgggt ggtaacaaat accagaccat cgacaactac 240
cagccgtacc cgtgcgtga agacgaagaa tgcggtagcg acgaatactg cgcttccccg 300
acccgtgggtg gtgacgctgg tgttcagatc tgcctggctt gccgtaaacg tcgtaaacgt 360
tgcatgctc acgctatgtg ctgccgggt aactactgca aaaacgggtat ctgctgttcc 420
tccgaccaga accacttccg tgggtgaaatc gaagaaacca tcaccgaatc ctccgtaac 480
gaccactcca ccctggacgg ttactcccgt cgtaccaccc tgcctccaa aatgtaccac 540
accaaaggtc aggaagggtc cgtttgcctg cgttccctcg actgcgttc cggctctgtg 600
tgcgctcgct acttctggtc caaaatctgc aaaccgggtc tgaagaagg tcaggtttgc 660
accaaacacc gtcgtaaaagg ttcccacggt ctggaaatct tccagcgttg ctactgcggt 720
gaaggtctgt cctgcggtat ccagaaagac caccaccagg cttccaactc ctcccgtctg 780
cacacctgcc agcgtcac 798

<210> SEQ ID NO 76

<211> LENGTH: 777

<212> TYPE: DNA

<213> ORGANISM: Human

<400> SEQUENCE: 76

atggctgctc tgatgcgttc caaagactcc tcctgctgcc tgctgctgct ggctgctggt 60
ctgatgggtt aatcctccca gatcgggttc tcccgtgcta aactgaactc catcaaatcc 120
tccctgggtg gtgaaacccc gggtcaggct gctaaccgtt ccgctgggtat gtaccagggt 180
ctggctttcg gtggttccaa aaaaggtaaa aacctgggtc aggttaccct gtgctcctcc 240
gacaaagaat gcgaagtggc tcgttactgc cactccccgc accagggttc ctccgcttgc 300
atggtttgcc gtcgtaaaaa aaaacgttgc caccgtgacg gtatgtgctg cccgtccacc 360
cgttgcaaca acggtatctg catccgggt accgaatcca tcctgacccc gcacatcccg 420
gctctggacg gtacccgta ccgtgaccgt aaccacggtc actactccaa ccacgacctg 480
ggttggcaga acctgggtcg tccgcacacc aaaatgtccc acatcaaagg tcacgaagg 540
gacctgccc tgcgttccct cgactgcac gaaggtttct gctgcgctcg tcacttctgg 600
acaaaaatct gaaacgggt tctgcaccag ggtgaagttt gcacaaaaca gcgtaaaaa 660
ggttcccacg gtctggaat cttccagcgt tgcgactgcg ctaaagggtc gtccctgcaa 720

-continued

gtttgaaag acgctaccta ctctccaaa gctcgtctgc acgtttgcc aaaaatc 777

<210> SEQ ID NO 77
<211> LENGTH: 1050
<212> TYPE: DNA
<213> ORGANISM: Human

<400> SEQUENCE: 77

atgcagcgtc tgggtgctac cctgctgtgc ctgctgctgg ctgctgctgt tccgaccgct 60
cgggtccgg ctccgaccgc tacctccgct cgggttaaac cgggtccggc tctgtcctac 120
ccgcaggaag aagctaccct gaacgaaatg ttccgtgaag ttgaagaact gatggaagac 180
accagcaca aactgcgttc cgctgttgaa gaaatggaag ctgaagaagc tgctgctaaa 240
gttctctccg aagttaacct ggctaacctg ccgccgtcct accacaacga aaccaacacc 300
gacaccaaag ttggtataca caccatccac gttcaccgtg aaatccacaa aatcaccaac 360
aaccagaccg gtcagatggt tttctccgaa accgttatca cctccgttgg tgacgaagaa 420
ggctgctggt ccacgaatg catcatcgac gaagactgcg gtccgtccat gtactgccag 480
ttcgcttctt tccagtacac ctgccagccg tgccgtggtc agcgtatgct gtgcaccctg 540
gactccgaat gctgcgggtg ccagctgtgc gtttggggtc actgcaccaa aatgggtacc 600
cgtgggtcca acggtaccat ctgcgacaac cagcgtgact gccagccggg tctgtgctgc 660
gttttccagc gtggtctgct gttcccggtt tgcaccccgc tgccgggtga aggtgaactg 720
tgccacgacc cggcttcccg tctgtgggac ctgatcacct gggaaactga accggacggg 780
gctctggacc gttgcccggt cgcttccggt ctgctgtgcc agccgcactc ccaactccctg 840
gtttacgttt gaaaccgac ctctggtggt tcccgtagac aggacgggtg aatcctgctg 900
ccgcgtgaag ttccggacga atacgaagtt ggttccttca tggagaagat tcgtcaggaa 960
ctggaagacc tggaacgttc cctgaccgaa gaaatggctc tgggtgaacc ggctgctgct 1020
gctgctgctc tgctgggtgg tgaagaaatc 1050

<210> SEQ ID NO 78
<211> LENGTH: 672
<212> TYPE: DNA
<213> ORGANISM: Human

<400> SEQUENCE: 78

atggttgctg ctgttctgct gggctgtgcc tggctgtgct ccccgctggg tgctctggtt 60
ctggacttca acaacatccg ttcctccgct gacctgcacg gtgctcgtaa aggttcccag 120
tgctgtccg acaccgactg caacacccgt aaattctgcc tgcagccggc tgacgaaaaa 180
ccgttctgcg ctacctgccg tggctctgct cgctgttgc agcgtgacgc tatgtgctgc 240
ccgggtaccc tgtgcgttaa cgacgtttgc accaccatgg aagacgctac cccgatcctg 300
gaacgtcagc tggacgaaca ggacggtacc cacgtgaag gtaccaccgg tcacccggtt 360
caggaaaacc agccgaaacg taaaccgtcc atcaaaaaat cccagggctc taaaggtcag 420
gaagtgtaat cctgcctgcg taccttcgac tgcggtcagg gtctgtgctg cgctcgtcac 480
ttctggacca aaatctgcaa accggttctg ctggaaggtc aggtttgctc ccgtcgtggt 540
cacaaagaca ccgtcagcg tccggaaatc ttccagcgtt gcgactgcgg tccgggtctg 600

-continued

```

ctgtgccggt cccagctgac ctccaaccgt cagcacgctc gtctgctgt ttgccagaaa    660
atcgaaaaac tg                                                         672

```

We claim:

1. An isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group consisting of:

- (a) the nucleic acid molecule comprising SEQ ID NO:1;
- (b) the nucleic acid molecule comprising SEQ ID NO:2;
- (c) the nucleic acid molecule comprising SEQ ID NO:3;
- (d) the nucleic acid molecule comprising SEQ ID NO:4;
- (e) the nucleic acid molecule comprising SEQ ID NO:5;
- (f) the nucleic acid molecule comprising SEQ ID NO:6;
- (g) the nucleic acid molecule comprising SEQ ID NO:7;
- (h) the nucleic acid molecule comprising SEQ ID NO:75;
- (i) the nucleic acid molecule comprising SEQ ID NO:76;
- (j) the nucleic acid molecule comprising SEQ ID NO:77;
- (k) the nucleic acid molecule comprising SEQ ID NO:78;
- (l) the nucleic acid molecule encoding the polypeptide of SEQ ID NO:8;
- (m) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:9;
- (n) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:10, or a biologically active fragment thereof;
- (o) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:11, or a biologically active fragment thereof;
- (p) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:12, or a biologically active fragment thereof;
- (q) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:13, or a biologically active fragment thereof;
- (r) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:14, or a biologically active fragment thereof;
- (s) a nucleic acid molecule that encodes a polypeptide that is at least 85 percent identical to the polypeptide of SEQ ID NOs: 10, 11, 12, 13, or 14;
- (t) a nucleic acid molecule that encodes a biologically active DKR polypeptide that has 1-100 amino acid substitutions and/or deletions as compared with, the polypeptide of any of SEQ ID NOs:8, 9, 10, 11, 12, 13, or 14; and
- (u) a nucleic acid molecule that hybridizes under conditions of high stringency to any of (c), (d), (e), (f), (g), (h), (i), (k), (l), (m), (n), (o), (p), (q), (r), (s), and (t) above.

2. An isolated nucleic acid molecule that is the complement of the nucleic acid molecule of claim 1.

3. An isolated nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or SEQ ID NO:7.

4. An isolated nucleic acid molecule encoding the polypeptide of SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, or SEQ ID NO:14.

5. An isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group consisting of: amino acids 16-350, 21-350, 22-350, 23-350, 33-350, or 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, or 310-350 of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-359 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

6. A vector comprising the nucleic acid molecule of claim 1.

7. A vector comprising the nucleic acid molecule of claim 2.

8. A vector comprising the nucleic acid molecule of claim 3.

9. A vector comprising the nucleic acid molecule of claim 4.

10. A vector comprising the nucleic acid molecule of claim 5.

11. A host cell comprising the vector of claim 6.

12. A host cell comprising the vector of claim 7.

13. A host cell comprising the vector of claim 8.

14. A host cell comprising the vector of claim 9.

15. A host cell comprising the vector of claim 10.

16. A process for producing a biologically active DKR polypeptide comprising the steps of:

(a) expressing a polypeptide encoded by the nucleic acid of claim 1 in a suitable host; and

(b) isolating the polypeptide.

17. The process of claim 16 wherein the polypeptide is SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13 or SEQ ID NO:14.

18. A biologically active DKR polypeptide selected from the group consisting of:

- (a) the polypeptide of SEQ ID NO:8;
- (b) the polypeptide of SEQ ID NO:9;
- (c) the polypeptide of SEQ ID NO:10;
- (d) the polypeptide of SEQ ID NO:11;
- (e) the polypeptide of SEQ ID NO:12;
- (f) the polypeptide of SEQ ID NO:13;

(g) the polypeptide of SEQ ID NO:14;

(h) a polypeptide that has 1-100 amino acid substitutions or deletions as compared with the polypeptide of any of (a)-(h) above; and

(i) a polypeptide that is at least 85 percent identical to any of the polypeptides of (c)-(h) above.

19. The polypeptide of claim 18 that does not possess an endogenous signal peptide.

20. A polypeptide selected from the group consisting of amino acids 16-350, 21-350, 22-350, 23-350, 33-350, 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, or 310-350 of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-259 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

* * * * *