

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2005303610 B2

(54) Title
Guanidine derivatives as inhibitors of DDAH

(51) International Patent Classification(s)
C07C 279/14 (2006.01) A61P 9/10 (2006.01)
A61K 31/16 (2006.01) A61P 11/06 (2006.01)
A61K 31/198 (2006.01) A61P 19/02 (2006.01)
A61K 31/223 (2006.01) C07C 279/16 (2006.01)
A61K 31/40 (2006.01) C07C 279/18 (2006.01)
A61K 31/445 (2006.01) C07C 323/44 (2006.01)
A61K 31/4453 (2006.01) C07C 323/52 (2006.01)
A61K 31/5375 (2006.01) C07D 295/215 (2006.01)

(21) Application No: **2005303610** (22) Date of Filing: **2005.11.11**

(87) WIPO No: **WO06/051314**

(30) Priority Data

(31) Number **0510348.6** (32) Date **2005.05.20** (33) Country **GB**
0425039.5 **2004.11.12** **GB**

(43) Publication Date: **2006.05.18**
(44) Accepted Journal Date: **2011.07.21**

(71) Applicant(s)
UCL Business PLC

(72) Inventor(s)
Vallance, Patrick John Thompson;Selwood, David Lawrence;Hartzoulakis, Basil;Leiper, James Mitchell;Rossiter, Sharon

(74) Agent / Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(56) Related Art
US 5352796
US 4686283
US 4659693

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

**(19) World Intellectual Property Organization
International Bureau**



A standard 1D barcode is located at the top of the page, spanning most of the width. It is used for document tracking and identification.

**(43) International Publication Date
18 May 2006 (18.05.2006)**

PCT

(10) International Publication Number
WO 2006/051314 A3

(51) International Patent Classification:

C07C 279/14 (2006.01) **A61P 19/02** (2006.01)
C07C 279/16 (2006.01) **A61K 31/40** (2006.01)
C07C 279/18 (2006.01) **A61K 31/4453** (2006.01)
C07C 323/44 (2006.01) **A61K 31/5375** (2006.01)
C07D 295/215 (2006.01) **A61K 31/198** (2006.01)
A61P 9/10 (2006.01) **A61K 31/223** (2006.01)
A61P 11/06 (2006.01) **A61K 31/16** (2006.01)

Hatfield, Herts AL10 9AB (GB). **HARTZOULAKIS, Basil** [GR/GB]; c/o NCE Discovery Ltd, 418 Science Park, Milton Road, Cambridge CB4 0PZ (GB).

(21) International Application Number:

PCT/GB2005/004361

(22) International Filing Date:

11 November 2005 (11.11.2005)

(74) **Agents:** WOODS, Geoffrey, Corlett et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5JJ (GB).

(74) **Agents:** WOODS, Geoffrey, Corlett et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5JJ (GB).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

(88) Date of publication of the international search report:
22 June 2006


(48) Date of publication of this corrected version:
28 September 2006

(15) Information about Correction:

see PCT Gazette No. 39/2006 of 28 September 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GUANIDINE DERIVATIVES AS INHIBITORS OF DDAH



(57) Abstract: Compounds of formula (I) have been found to be useful as inhibitors of DDAH. The present invention thus provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment of a disease whose pathology is affected by DDAH.

CHEMICAL COMPOUNDS

Field of the Invention


The present invention relates to compounds that are inhibitors of DDAH and
 5 which are therefore useful in treating diseases whose pathology is affected by
 DDAH.

Background of the Invention

Endogenous methylarginines $N^G N^G$ dimethyl-L-arginine (ADMA) and N^G
 10 monomethyl-L-arginine (L-NMMA) which are released by proteolysis of methylated
 protein arginine residues, are inhibitors of all nitric oxide synthase (NOS) isoforms.
 The enzyme dimethylarginine dimethylaminohydrolase (DDAH) metabolises the
 two NOS-inhibiting methylarginines, but not the related $N^G N^G$ dimethylarginine
 (SDMA) which is physiologically inactive.
 15 Thus DDAH, which controls levels of asymmetrically methylated arginine
 derivatives, may have therapeutic potential through its ability to indirectly influence
 the activity of NOS.

Summary of the Invention

20 It has now surprisingly been found that compounds of the general formula (I)
 set out below act as inhibitors of DDAH. Accordingly, the present invention
 provides the use of a compound of formula (I), or a pharmaceutically acceptable salt
 thereof, in the manufacture of a medicament for use as an inhibitor of DDAH



25

wherein:

- either (a) R^1 is C_3-C_{12} alkyl, C_3-C_{12} alkenyl, C_3-C_{12} alkynyl, C_6-C_{10} aryl,
 C_3-C_8 carbocyclyl, 5- to 10-membered heterocyclyl, 5- to 10-membered
 30 heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A or -L-Y-A, wherein:
 L is C_1-C_6 alkyl, C_2-C_6 alkenyl or C_2-C_6 alkynyl;

L' is C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
A is C₆-C₁₀ aryl, C₃-C₈ carbocyclyl, 5- to 10-membered heterocyclyl
or 5- to 10-membered heteroaryl;
Het is -O-, -S- or -NR'-, wherein R' is hydrogen, C₁-C₆ alkyl, C₂-C₆
5 alkenyl, C₂-C₆ alkynyl; and
Y is -CO-, -SO-, -SO₂-, -CO-O-, -CO-NR'-, -O-CO- or -NR'-CO-,
wherein R' is as defined above;

10 - R² is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl; and
- R³ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl,
- or (b) R¹ and R² form, together with the nitrogen to which they are attached, a
5- to 10-membered heterocyclic or heteroaryl ring, and R³ is as defined
above,
- or (c) R¹ and R³ form, together with the -N-C=N- moiety to which they are
attached, a 5- to 10-membered heterocyclic or heteroaryl ring, and R² is as
15 defined above;

15 - R⁴ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
- B is C₁-C₆ alkyl, C₂-C₆ alkenyl, -O-L-, -S-L- or -L-Het-L'-, wherein L, L' and
Het are as defined above;
- X is -O-, -S- or -NR'-, wherein R' is as defined above;

20 - R⁵ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, aryl, C₃-C₈
carbocyclyl, 5- to 10-membered heterocyclyl, 5- to 10-membered heteroaryl,
-L-A, -L-Het-L', -L-Y-L', -L-Het-A, -L-Y-A, -L-A-Het-A, -L-A-L'-A or
-L-A-Het-L'-A, wherein L, L', Het and Y are as defined above and wherein
each A is the same or different and is as defined above;

25 - R⁶ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
- R⁷ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl, or R⁷ is a bond, a
C₁-C₄ alkyl or C₂-C₄ alkenyl moiety which is joined to one of the carbon
atoms of B to form, together with the -N-C- moiety to which R⁷ and B are
attached, a 5- to 10-membered heterocyclic ring; and


30 - R⁸ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl;
wherein:
- the alkyl, alkenyl and alkynyl groups and moieties in the substituents R¹ to
R⁷, X and B are unsubstituted or substituted by one, two or three substituents

which are the same or different and are selected from halogen, hydroxy, amino and thio substituents; and

- the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R¹ and R⁵, the heterocyclic or heteroaryl moieties formed by R¹ together with R² and R¹ together with R³ and heterocyclic moieties formed by R⁷ together with B are unsubstituted or substituted by one, two or three substituents selected from halogen, hydroxy, amino, thio, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, nitro, cyano, C₁-C₆ alkylamino, di-(C₁-C₆ alkyl)amino, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy and C₁-C₆ haloalkylthio substituents.

10 Examples of compounds of formula (I) are compounds as defined above wherein R⁵ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, aryl, C₃-C₈ carbocyclyl, 5- to 10-membered heterocyclyl, 5- to 10-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A or -L-Y-A, wherein L, L', A, Het and Y are as defined above.

15 According to an aspect of the present invention there is provided use of a compound of formula (I), or a pharmaceutically acceptable salt thereof in the manufacture of a medicament, for use in the treatment of a disease whose pathology is affected by DDAH



wherein:

- either (a) R¹ is C₃-C₈ alkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A or -L-Y-A, wherein:

L is C₁-C₆ alkyl;

L' is C₁-C₆ alkyl;

A is phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl or 5- to 6-membered heteroaryl;

Het is -O-, -S- or -NR'-, wherein R' is hydrogen or C₁-C₆ alkyl; and

Y is -CO-, -SO-, -SO₂-, -CO-O-, -CO-NR'-, -O-CO- or -NR'-CO-, wherein

R' is as defined above;

30 - R² is hydrogen or C₁-C₆ alkyl;

- R³ is hydrogen or C₁-C₆ alkyl;

- or (b) R¹ and R² form, together with the nitrogen to which they are

attached, a 5- to 6-membered heterocyclic or heteroaryl ring, and R^3 is as defined above;

- or (c) R^1 and R^3 form, together with the $-N-C=N-$ moiety to which they are attached, a 5- to 6-membered heterocyclic or heteroaryl ring, and R^2 is as defined above;

- R^4 is hydrogen or C_1 - C_6 alkyl;

- B is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, $-O-L-$, $-S-L-$ or $L-Het-L'$, wherein L , L' and Het are as defined above;

- X is $-O-$, $-S-$ or $-NR'$, wherein R' is as defined above;

- R^5 is hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, phenyl, C_3 - C_6 carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, $-L-A$, $-L-Het-L'$, $-L-Y-L'$, $-L-Het-A$, $-L-Y-A$ or $-L-A-Het-A$, wherein L , L' , Het and Y are as defined above and wherein each A is the same or different and is as defined above;

- R^6 is hydrogen or C_1 - C_6 alkyl;

- R^7 is hydrogen or C_1 - C_6 alkyl or R^7 is a bond, a C_1 - C_4 alkyl or C_2 - C_4 alkenyl moiety which is joined to one of the carbon atoms of B to form, together with the $-N-C-$ moiety to which R^7 and B are attached, a 5- to 6-membered heterocyclic ring; and

R^8 is hydrogen or C_1 - C_6 alkyl,

wherein:

- the alkyl, alkenyl and alkynyl groups and moieties in the substituents R^1 to R^7 , X and B are unsubstituted or substituted by one, two or three substituents which are the same or different and are selected from fluorine, chlorine, bromine, hydroxy, amino and thio substituents; and

- the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R^1 and R^5 , the heterocyclic or heteroaryl moieties formed by R^1 together with R^2 and R^1 together with R^3 and the heterocyclic moieties formed by R^7 together with B are typically unsubstituted or substituted by one, two or three substituents selected from fluorine, chlorine, bromine, hydroxy, amino, thio, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylamino, di-(C_1 - C_4 alkyl)amino, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy and C_1 - C_4 haloalkylthio substituents.

The present invention also relates to a compound of formula (I) for use in the manufacture of a medicament for use in the treatment of a disease whose pathology is affected by DDAH.

The present invention also relates to a compound of formula (I) for use in the manufacture of a medicament for use in the treatment of ischaemia-reperfusion injury of the brain or heart, cancer, lethal hypotension in severe inflammatory conditions, local and systemic inflammatory disorders, neurodegeneration, asthma, pain or sepsis.

5

Description of the Figures

Figure 1 is a graph showing the nitrate and nitrite (NOx) production from isolated rat aortic rings in static culture in the presence of either media alone (untreated); media plus 10 μ g/ml LPS (LPS) and media plus 10 μ g/ml LPS and 500 μ M SR257 or SR291. n \geq 6;
*p<0.0001 One Way ANOVA plus Bonferroni post hoc correction.

10 Figures 2a and 2b are traces illustrating the tension in isolated aortic ring demonstrating contraction in response to phenylephrine EC80 and subsequent iNOS mediated dilatation in response to LPS treatment over 4 hour timecourse (Figure 2a)

and response of vessel to D-SR257, L-SR257 and L-arginine (Figure 2b - expanded trace from 2a).

Figure 3 is a trace illustrating blood pressure in anaesthetized rat demonstrating initial effect of 40mg/kg endotoxin (ETX) administration on blood pressure followed by a recovery and gradual fall in blood pressure over time. Effects 5 of intervention after 2-3 hours (when blood pressure had fallen to 80% of baseline pre LPS blood pressure) with 30mg/kg/hr infusion of SR291.

Detailed Description of the Invention

10 The present invention relates to the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use as an inhibitor of DDAH.

For the avoidance of doubt, the orientation of the group Y in the moiety -L-Y-A is such that the left hand side of the depicted group is attached to L. Thus, 15 for example, when Y is -CO-NR'-, the group -L-Y-A is -L-CO-NR'-A. Similarly, for the avoidance of doubt, the orientation of the group Y in the moiety -L-Y-L' is such that the left hand side of the depicted group is to L. Thus, for example, when Y is -CO-NR'-, the group -L-Y-L' is -L-CO-NR'-L'.

For the avoidance of doubt, when the group B is -O-L- or -S-L- the left hand 20 side of the depicted group is attached to the guanadine nitrogen. Thus, for example, when B is -O-L-, the moiety -R⁴N-B-CR⁸(NR⁶R⁷)- is -R⁴N-O-L-CR⁸(NR⁶R⁷)-.

As used herein, a C₁-C₁₂ alkyl group or moiety is a linear or branched alkyl group or moiety containing from 1 to 12 carbon atoms, such as a C₃-C₁₂, a C₁-C₆ or C₁-C₄ alkyl group or moiety. Examples of such alkyl groups or moieties are methyl, 25 ethyl, n-propyl, i-propyl, n-butyl, n-octyl and -CH₂CMe₃. Examples of C₃-C₁₂ alkyl groups or moieties are n-propyl, i-propyl, n-butyl, n-octyl and -CH₂CMe₃. In one embodiment, a C₃-C₁₂ alkyl group is preferably a C₅-C₁₂ alkyl group and more preferably a C₆-C₁₂ alkyl group. A divalent alkyl group or moiety (or alkylene group or moiety) can be attached via the same carbon atom, via adjacent carbon atoms or 30 via non-adjacent carbon atoms. Examples of divalent alkyl moieties are methylene, 1,2-ethyl and 1,3-propyl moieties. Preferred divalent alkyl moieties are 1,2-ethyl and 1,3-propyl moieties.

As used herein, a C₂-C₁₂ alkenyl group or moiety is a linear or branched alkenyl group or moiety containing from 2 to 12 carbon atoms, such as a C₃-C₁₂, C₂-C₆ or C₂-C₄ alkenyl group or moiety, for example ethenyl, *n*-propenyl and *n*-butenyl. A preferred C₂-C₆ alkenyl group is ethenyl. A preferred C₃-C₁₂ alkenyl group is allyl. In one embodiment, a C₃-C₁₂ alkenyl group is preferably a C₄-C₁₂ alkenyl group and more preferably a C₅-C₁₂ alkenyl group. Typically an alkenyl group has only one double bond. This double bond is typically located at the α -position of the alkenyl group. A divalent alkenyl group (or alkenylene group) can be attached via the same carbon atom, via adjacent carbon atoms or via non-adjacent carbon atoms.

As used herein, a C₂-C₁₂ alkynyl group or moiety is a linear or branched alkenyl group or moiety containing from 2 to 12 carbon atoms, such as a C₃-C₁₂, C₂-C₆ or C₂-C₄ alkynyl group or moiety, for example ethynyl. In one embodiment, a C₃-C₁₂ alkynyl group is preferably a C₄-C₁₂ alkynyl group and more preferably a C₅-C₁₂ alkynyl group. Typically an alkynyl group has only one triple bond. This triple bond is typically located at the α -position of the alkynyl group. A divalent alkynyl group (or alkynylene group) can be attached via the same carbon atom, via adjacent carbon atoms or via non-adjacent carbon atoms.

As used herein, a halogen is typically chlorine, fluorine, bromine or iodine and is preferably chlorine or fluorine. As used herein, a said C₁-C₆ alkoxy group is typically a said C₁-C₆ alkyl group attached to an oxygen atom. A said C₁-C₆ alkylthio group is typically a said C₁-C₆ alkyl group attached to a thio group.

As used herein, a C₁-C₆ haloalkyl group is typically a said C₁-C₆ alkyl group, for example a C₁-C₄ alkyl group, substituted by one or more said halogen atoms. Typically, it is substituted by 1, 2 or 3 said halogen atoms. Examples of haloalkyl groups include perhaloalkyl groups such as -CX₃ wherein X is a said halogen atom, for example -CF₃. Preferred haloalkyl groups include monohaloalkyl groups such as -CH₂-CH₂F and perhaloalkyl groups such as -CF₃. Examples of preferred haloalkyl groups include monohaloalkyl groups such as -CH₂-CH₂F.

As used herein, a C₁-C₆ haloalkoxy group is typically a said C₁-C₆ alkoxy group, for example a C₁-C₄ alkoxy group, substituted by one or more said halogen atoms. Typically, it is substituted by 1, 2 or 3 said halogen atoms. Preferred

haloalkoxy groups include perhaloalkoxy groups such as -OCX₃ wherein X is a said halogen atom. Particularly preferred haloalkoxy groups are -OCF₃ and -OCCl₃.

As used herein, a C₁-C₆ haloalkylthio group is typically a said C₁-C₆ alkylthio group, for example a C₁-C₄ alkylthio group, substituted by one or more said halogen atoms. Typically, it is substituted by 1, 2 or 3 said halogen atoms. Preferred haloalkylthio groups include perhaloalkylthio groups such as -SCX₃ wherein X is a said halogen atom. Particularly preferred haloalkylthio groups are -SCF₃ and -SCCl₃.

As used herein, a C₃-C₆ carbocyclyl group or moiety is a non-aromatic saturated or unsaturated hydrocarbon ring, having from 3 to 6 carbon atoms.

Preferably it is a saturated group, i.e. a C₃-C₆ cycloalkyl group. Examples include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Preferred carbocyclyl groups are cyclopropyl and cyclohexyl. In one embodiment, a C₃-C₆ carbocyclyl group is preferably a C₄-C₆ carbocyclyl group and more preferably a C₅-C₆ carbocyclyl group.

As used herein, a 5- to 10-membered heterocyclyl group or moiety is a non-aromatic, saturated or unsaturated C₅-C₁₀ carbocyclic ring, for example a 5- or 6-membered ring, in which one or more, for example 1, 2 or 3, of the carbon atoms are replaced by a heteroatom selected from N, O and S. Saturated heterocyclyl groups are preferred. Examples of suitable heterocyclyl groups include piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, tetrahydrofuranyl, imidazolidinyl, thiazolidinyl, 1,4 dioxanyl, 1,3 dioxolanyl and dihydroimidazolyl. Preferred heterocyclyl groups are piperidinyl, morpholinyl, pyrrolidinyl, tetrahydrofuran and dihydroimidazolyl. Examples of preferred heterocyclyl groups are piperidinyl, morpholinyl, pyrrolidinyl and dihydroimidazolyl. Particularly preferred heterocyclyl groups are piperidinyl, morpholinyl, tetrahydrofuran and pyrrolidinyl. Examples of particularly preferred heterocyclyl groups are piperidinyl, morpholinyl and pyrrolidinyl.

As used herein, a C₆-C₁₀ aryl group or moiety is typically a phenyl or naphthyl group or moiety. Preferably, it is a phenyl moiety.

As used herein, a 5- to 10-membered heteroaryl group is a 5- to 10-membered aromatic ring, such as a 5- or 6-membered ring, containing at least one heteroatom, for example 1, 2 or 3 heteroatoms, selected from O, S and N. Examples

include pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, furanyl, thienyl, imidazolyl, pyrazolidinyl, pyrrolyl, oxadiazolyl, oxazolyl, isoxazolyl, thiadiazolyl, thiazolyl and pyrazolyl groups. Thienyl groups are preferred.

Typically, the alkyl, alkenyl and alkynyl groups and moieties in the substituents R¹ to R⁷, X and B are unsubstituted or substituted by one, two or three substituents which are the same or different and are selected from fluorine, chlorine, bromine, hydroxy, amino and thio substituents. Preferably, the alkyl, alkenyl and alkynyl groups and moieties in the substituents R¹ to R⁷, X and B are unsubstituted or substituted by a single hydroxy substituent or by one, two or three substituents which are the same or different and are selected from fluorine and chlorine substituents. More preferably, the alkyl, alkenyl and alkynyl groups and moieties in the substituents R¹ to R⁷, X and B are unsubstituted or substituted by a single fluoro substituent.

Typically, when a said aryl, heteroaryl, heterocyclyl or carbocyclyl group or moiety carries a nitro or cyano substituent, only one of the substituents on the aryl, heteroaryl, heterocyclyl, or carbocyclyl group is a nitro or cyano group. Further, the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R¹ and R⁵, the heterocyclic or heteroaryl moieties formed by R¹ together with R² and R¹ together with R³ and the heterocyclic moieties formed by R⁷ together with B are typically unsubstituted or substituted by one, two or three substituents selected from fluorine, chlorine, bromine, hydroxy, amino, thio, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylamino, di-(C₁-C₄ alkyl)amino, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy and C₁-C₄ haloalkylthio substituents.

Preferably, the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R¹ and R⁵, the heterocyclic or heteroaryl moieties formed by R¹ together with R² and R¹ together with R³ and the heterocyclic moieties formed by R⁷ together with B are typically unsubstituted or substituted by one, two or three substituents selected from fluorine, chlorine, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy and C₁-C₄ haloalkyl substituents.

More preferably, the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R¹, the heterocyclic or heteroaryl moieties formed by R¹ together with R² and R¹ together with R³ and the heterocyclic moieties formed by R⁷ together with B are unsubstituted and the aryl, carbocyclyl, heterocyclyl and

heteroaryl groups and moieties in the substituent R⁵ are unsubstituted or substituted by one or two substituents selected from fluorine, C₁-C₂ alkyl, C₁-C₂ alkoxy and C₁-C₂ haloalkyl substituents.

It is further preferred that the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R¹ and R⁵, the heterocyclic or heteroaryl moieties formed by R¹ together with R² and R¹ together with R³ and the heterocyclic moieties formed by R⁷ together with B are unsubstituted.

Typically, the substituents on the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties are themselves unsubstituted.

Typically, R' in the or each Het or Y moiety is hydrogen or C₁-C₆ alkyl. Preferably, R' in the or each Het or Y moiety is hydrogen or C₁-C₄ alkyl. More preferably, R' in the or each Het or Y moiety is hydrogen or methyl.

Typically, L in the or each R¹, R⁵ or B moiety is C₁-C₆ alkyl. Preferably, L in the or each R¹, R⁵ or B moiety is C₁-C₄ alkyl. More preferably, L in the or each R¹, R⁵ or B moiety is C₁-C₂ alkyl.

Typically, L' in the or each R¹, R⁵ or B moiety is C₁-C₆ alkyl. Preferably, L' in the or each R¹, R⁵ or B moiety is C₁-C₄ alkyl. More preferably, L' in the or each R¹, R⁵ or B moiety is C₁-C₂ alkyl.

Typically, A in the or each R¹ or R⁵ moiety is phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl or 5- to 6-membered heteroaryl. Preferably, A in the or each R¹ or R⁵ moiety is phenyl, 5- to 6-membered heterocyclyl or 5- to 6-membered heteroaryl. Examples of preferred A groups in the or each R¹ or R⁵ moiety are phenyl and 5- to 6-membered heteroaryl. More preferably, A in the or each R¹ or R⁵ moiety is phenyl, tetrahydrofuranyl or thienyl. Examples of more preferred A groups in the or each R¹ or R⁵ moiety are phenyl and thienyl.

Typically, Het in the or each R¹, R⁵ or B moiety is -O-, -S- or -NR'-, wherein R' is as defined above. Preferably, Het in the or each R¹, R⁵ or B moiety is -O-, -S- or -NMe-.

Typically, Y in the or each R¹ or R⁵ moiety is -CO-, -SO-, -SO₂-, -CO-O-, -CO-NR'-, -O-CO- or -NR'-CO-, wherein R' is as defined above. Preferably, Y in the or each R¹ or R⁵ moiety is -CO-, -CO-O- or -CO-NR'-, wherein R' is as defined above. More preferably, Y in the or each R¹ or R⁵ moiety is -CO-, -CO-O-, -CO-NH- or -CO-NMe-.

Typically, -L-Y-L' in the or each R¹ or R⁵ moiety is -(C₁-C₆ alkyl)-Y-(C₁-C₆ alkyl), wherein Y is as defined above. Preferably, -L-Y-L' in the or each R¹ or R⁵ moiety is -(C₁-C₄ alkyl)-Y-(C₁-C₄ alkyl), wherein Y is as defined above. More preferably, -L-Y-L' in the or each R¹ or R⁵ moiety is -(C₁-C₂ alkyl)-CO-(C₁-C₂ alkyl), -(C₁-C₂ alkyl)-CO-O-(C₁-C₂ alkyl), -(C₁-C₂ alkyl)-CO-NH-(C₁-C₂ alkyl) or -(C₁-C₂ alkyl)-CO-NMe-(C₁-C₂ alkyl).

Typically, -L-Het-L' in the or each R¹ or R⁵ moiety is -(C₁-C₆ alkyl)-Het-(C₁-C₆ alkyl), wherein Het is as defined above. Preferably, -L-Het-L' in the or each R¹ or R⁵ moiety is -(C₁-C₄ alkyl)-Het-(C₁-C₄ alkyl), wherein Het is as defined above. More preferably, -L-Het-L' in the or each R¹ or R⁵ moiety is -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl), -(C₁-C₂ alkyl)-S-(C₁-C₂ alkyl) or -(C₁-C₂ alkyl)-NMe-(C₁-C₂ alkyl).

Typically, -L-Het-A in the or each R¹ or R⁵ moiety is -(C₁-C₆ alkyl)-Het-A, wherein Het and A are as defined above. Preferably, -L-Het-A in the or each R¹ or R⁵ moiety is -(C₁-C₄ alkyl)-Het-A, wherein Het is as defined above and A is a phenyl or 5- to 6-membered heteroaryl group. More preferably, -L-Het-A in the or each R¹ or R⁵ moiety is -(C₁-C₂ alkyl)-O-phenyl.

Typically, -L-Y-A in the or each R¹ or R⁵ moiety is -(C₁-C₆ alkyl)-Y-A, wherein Y and A are as defined above. Preferably, -L-Y-A in the or each R¹ or R⁵ moiety is -(C₁-C₄ alkyl)-Y-A, wherein Y is as defined above and A is a phenyl or 5- to 6-membered heteroaryl group. More preferably, -L-Het-A in the or each R¹ or R⁵ moiety is -(C₁-C₂ alkyl)-CO-phenyl, -(C₁-C₂ alkyl)-CO-O-phenyl, -(C₁-C₂ alkyl)-CO-NH-phenyl or -(C₁-C₂ alkyl)-CO-NMe-phenyl.

Typically, -L-A in the or each R¹ or R⁵ moiety is -(C₁-C₆ alkyl)-A, wherein A is as defined above. Preferably, -L-A in the or each R¹ or R⁵ moiety is -(C₁-C₄ alkyl)-A, wherein A is a phenyl, 5- to 6-membered heterocycl or 5- to 6-membered heteroaryl group. An example of a preferred -L-A in the or each R¹ or R⁵ moiety is -(C₁-C₄ alkyl)-A, wherein A is a phenyl or 5- to 6-membered heteroaryl group. More preferably, -L-A in the R¹ moiety is -(C₁-C₂ alkyl)-phenyl, -(C₁-C₂ alkyl)-tetrahydrofuran or -(C₁-C₂ alkyl)-thienyl. Examples of more preferred -L-A in the R¹ moiety is -(C₁-C₂ alkyl)-phenyl or -(C₁-C₂ alkyl)-thienyl. Further, more preferably, -L-A in the R⁵ moiety is -(C₁-C₂ alkyl)-phenyl.

Typically, -L-A-Het-A in the R⁵ moiety is -(C₁-C₆ alkyl)-A-Het-A wherein Het is as defined above and each A is the same or different and is as defined above.

Preferably, -L-A-Het-A in the R⁵ moiety is -(C₁-C₄ alkyl)-A-Het-A, wherein Het is as defined above and each A is the same or different and is a phenyl or 5- to 6-membered heteroaryl group. More preferably, -L-A-Het-A in the R⁵ moiety is -(C₁-C₂ alkyl)-phenyl-O-phenyl.

5 Typically, R¹ is C₃-C₈ alkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A or -L-Y-A, wherein L, L', A, Het and Y are as defined above.

10 Preferably, R¹ is C₃-C₈ alkyl, C₃-C₆ alkenyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L' or -L-Het-A, wherein L, L', A and Het are as defined above.

15 In one embodiment, R¹ is preferably phenyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L' or -L-Het-A, wherein L, L', A and Het are as defined above. It is particularly preferred that R¹ is -L-Het-L' or -L-Het-A, wherein L, L', A and Het are as defined above. It is more preferred that R¹ is -L-Het-L' wherein L, L' and Het are as defined above. It is further preferred that R¹ is -(C₁-C₄ alkyl)-Het-(C₁-C₄ alkyl), wherein Het is -O-, -S- or -NMe-.

20 More preferably, R¹ is C₃-C₈ alkyl, allyl, phenyl, cyclopropyl, cyclohexyl, -(C₁-C₂ alkyl)-phenyl, -(C₁-C₂ alkyl)-thienyl, -(C₁-C₂ alkyl)-tetrahydrofuranyl, -(C₁-C₂ alkyl)-O-phenyl, -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl), -(C₁-C₂ alkyl)-S-(C₁-C₂ alkyl) or -(C₁-C₂ alkyl)-NMe-(C₁-C₂ alkyl). Examples of more preferred R¹ groups are C₃-C₈ alkyl, allyl, phenyl, cyclopropyl, cyclohexyl, -(C₁-C₂ alkyl)-phenyl, -(C₁-C₂ alkyl)-thienyl, -(C₁-C₂ alkyl)-O-phenyl, -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl), -(C₁-C₂ alkyl)-S-(C₁-C₂ alkyl) and -(C₁-C₂ alkyl)-NMe-(C₁-C₂ alkyl).

25 Typically, R² is hydrogen or C₁-C₆ alkyl. Preferably, R² is hydrogen or C₁-C₄ alkyl. More preferably, R² is hydrogen, methyl or ethyl.

Typically, R³ is hydrogen or C₁-C₆ alkyl. Preferably, R³ is hydrogen or C₁-C₂ alkyl. More preferably, R³ is hydrogen.

When R¹ and R² form, together with the nitrogen to which they are attached, 30 a 5- to 10-membered heterocyclic or heteroaryl ring, they typically form a 5- to 6-membered heterocyclic or heteroaryl ring. Preferably, they form a 5- to 6-membered heterocyclic ring. Examples of such rings are piperidinyl, morpholinyl and

pyrrolidinyl rings. Typically, said heterocyclic and heteroaryl rings are unsubstituted.

When R¹ and R³ form, together with the -N-C=N- moiety to which they are attached, a 5- to 10-membered heterocyclic or heteroaryl ring, they typically form a 5- to 6-membered heterocyclic or heteroaryl ring. Preferably, they form a 5- to 6-membered heterocyclic ring. Examples of such rings are dihydroimidazole and tetrahydropyrimidine rings with dihydroimidazole rings being preferred. Typically, said heterocyclic and heteroaryl rings are unsubstituted.

Typically, R⁴ is hydrogen or C₁-C₆ alkyl. Preferably, R⁴ is hydrogen or C₁-C₂ alkyl. More preferably, R⁴ is hydrogen.

Typically, -L-Het-L'- in the moiety B is -(C₁-C₄ alkyl)-Het-(C₁-C₄ alkyl)-, wherein Het is as defined above. Preferably, -L-Het-L'- in the moiety B is -(C₁-C₂ alkyl)-Het-(C₁-C₂ alkyl)-, wherein Het is as defined above.

Typically, -O-L- in the moiety B is -O-(C₁-C₄ alkyl)-. Preferably, -O-L- in the moiety B is -O-(C₁-C₂ alkyl)-.

Typically, -S-L- in the moiety B is -S-(C₁-C₄ alkyl)-. Preferably, -S-L- in the moiety B is -S-(C₁-C₂ alkyl)-.

Typically, B is C₂-C₆ alkyl. Preferably, B is C₂-C₄ alkyl. More preferably, B is 1,2-ethyl or 1,3-propyl. In a preferred embodiment, B is unsubstituted.

Typically, X is -O-, -S- or -NR'-, wherein R' is as defined above. Preferably, X is -O- or -NH-. Most preferably, X is -O-.

Typically, R⁵ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A, -L-Y-A or -L-A-Het-A, wherein L, L', Het and Y are as defined above and wherein each A is the same or different and is as defined above. Examples of typical R⁵ groups are hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Het-A, -L-Y-L', -L-Het-A and L-Y-A, wherein L, L', A, Het and Y are as defined above. Preferably, R⁵ is hydrogen, C₁-C₆ alkyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Het-A or -L-A-Het-A, wherein L, L', and Het are as defined above and wherein each A is the same or different and is as defined above. Examples of preferred R⁵ groups are hydrogen, C₁-C₆ alkyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered

heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L' and -L-Het-A, wherein L, L', A and Het are as defined above. More preferably, R⁵ is hydrogen, C₁-C₄ alkyl or -(C₁-C₂ alkyl)-phenyl, -(C₁-C₂ alkyl)-O-(C₁-C₂ alkyl) or -(C₁-C₂ alkyl)-phenyl-O-phenyl. Examples of more preferred R⁵ groups are hydrogen, C₁-C₄ alkyl and 5 -(C₁-C₂)-phenyl.

Typically, R⁶ is hydrogen or C₁-C₆ alkyl. Preferably, R⁶ is hydrogen or C₁-C₂ alkyl. More preferably, R⁶ is hydrogen.


Typically, R⁷ is hydrogen or C₁-C₆ alkyl. Preferably, R⁷ is hydrogen or C₁-C₂ alkyl. More preferably, R⁷ is hydrogen.

10 When R⁷ is a bond, a C₁-C₄ alkyl or C₂-C₄ alkenyl moiety which is joined to one of the carbon atoms of B to form, together with the -N-C- moiety to which R⁷ and B are attached, a 5- to 10-membered heterocyclic ring, it typically forms a 5- to 6-membered heterocyclic ring such as a pyrrolidine or piperidine ring. Pyrrolidine rings are preferred. Typically, said heterocyclic rings are unsubstituted.

15 When R⁷ is a bond, a C₁-C₄ alkyl or C₂-C₄ alkenyl moiety which is joined to one of the carbon atoms of B to form, together with the -N-C- moiety to which R⁷ and B are attached, a 5- to 10-membered heterocyclic ring, R⁷ is typically a bond, or a C₁-C₄ alkyl moiety. Preferably, R⁷ is a C₁-C₂ alkyl moiety.


Typically, R⁸ is hydrogen or C₁-C₆ alkyl. Preferably, R⁸ is hydrogen or C₁-C₂ alkyl. More preferably, R⁸ is hydrogen.

In one embodiment of the present invention, the moiety




in the compounds of formula (I) typically represents

25




preferably represents



5

and more preferably represents



10 wherein R⁶, R⁷ and R⁸ are as defined above.

In one embodiment, the compound of formula (I) is not N^G -allyl-L-arginine or N^G -cyclopropyl-L-arginine. In a second embodiment, the compound of formula (I) is not N^G -allyl-L-arginine, N^G -propyl-L-arginine or N^G -propargyl-L-arginine. In a further embodiment, the compound of formula (I) is not N^G -propyl-L-arginine or N^G -butyl-L-arginine. In a preferred embodiment, the compound of formula (I) is not N^G -propyl-L-arginine, N^G -allyl-L-arginine, N^G -propargyl-L-arginine, N^G -butyl-L-arginine or N^G -cyclopropyl-L-arginine. More specifically, if the compound of formula (I) is used in the manufacture of a medicament for use in the treatment of pruritus, it is preferred that the compound of formula (I) is not N^G -allyl-L-arginine or N^G -cyclopropyl-L-arginine. If the compound of formula (I) is used in the manufacture of a medicament for use in the treatment of stroke, Alzheimer's disease and other neurodegenerative diseases, septic shock, inflammatory arthritis or colitis, it is preferred that the compound of formula (I) is not N^G -allyl-L-arginine, N^G -propyl-L-arginine or N^G -propargyl-L-arginine. If the compound of formula (I) is used in the manufacture of a medicament for use in the treatment of solid tumors, it is preferred that the compound of formula (I) is not N^G -propyl-L-arginine or N^G -butyl-L-arginine.

Preferred compounds of formula (I) are those wherein:

- either (a) R^1 is C_3 - C_8 alkyl, C_3 - C_8 alkenyl, C_3 - C_8 alkynyl, phenyl, C_3 - C_6 carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, - L - A , - L -Het- L' , - L - Y - L' , - L -Het- A or - L - Y - A , wherein:
 - 5 L is C_1 - C_6 alkyl;
 - L' is C_1 - C_6 alkyl;
 - A is phenyl, C_3 - C_6 carbocyclyl, 5- to 6-membered heterocyclyl or 5- to 6-membered heteroaryl;
 - Het is - O -, - S - or - NR' -, wherein R' is hydrogen or C_1 - C_6 alkyl; and
 - Y is - CO -, - SO -, - SO_2 -, - $CO-O$ -, - $CO-NR'$ -, - $O-CO$ - or - $NR'-CO$ -, wherein R' is as defined above;
- R^2 is hydrogen or C_1 - C_6 alkyl.
- R^3 is hydrogen or C_1 - C_6 alkyl.
- or (b) R^1 and R^2 form, together with the nitrogen to which they are attached, a 5- to 6-membered heterocyclic or heteroaryl ring, and R^3 is as defined above;
- 15 - or (c) R^1 and R^3 form, together with the - $N-C=N-$ moiety to which they are attached, a 5- to 6-membered heterocyclic or heteroaryl ring, and R^2 is as defined above;
- R^4 is hydrogen or C_1 - C_6 alkyl;
- B is C_2 - C_6 alkyl;
- 20 - X is - O -, - S - or - NR' -, wherein R' is as defined above;
- R^5 is hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, phenyl, C_3 - C_6 carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, - L - A , - L -Het- L' , - L - Y - L' , - L -Het- A , - L - Y - A or - L - A -Het- A , wherein L , L' , Het and Y are as defined above and wherein each A is the same or different and is as defined above;
- 25 - R^6 is hydrogen or C_1 - C_6 alkyl;
- R^7 is hydrogen or C_1 - C_6 alkyl or R^7 is a bond, a C_1 - C_4 alkyl or C_2 - C_4 alkenyl moiety which is joined to one of the carbon atoms of B to form, together with the - $N-C-$ moiety to which R^7 and B are attached, a 5- to 6-membered heterocyclic ring; and
- 30 - R^8 is hydrogen or C_1 - C_6 alkyl,

wherein:

- the alkyl, alkenyl and alkynyl groups and moieties in the substituents R^1 to R^7 , X and B are unsubstituted or substituted by one, two or three substituents which are the same or different and are selected from fluorine, chlorine, bromine, hydroxy, amino and thio substituents; and
- 5 - the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R^1 and R^5 , the heterocyclic or heteroaryl moieties formed by R^1 together with R^2 and R^1 together with R^3 and the heterocyclic moieties formed by R^7 together with B are typically unsubstituted or substituted by one, two or three substituents selected from fluorine, chlorine, bromine, hydroxy, amino, thio, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylamino, di-(C_1 - C_4 alkyl)amino, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy and C_1 - C_4 haloalkylthio substituents.

Examples of such compounds are the preferred compounds of formula (I) as defined above wherein R⁵ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A or L-Y-A, wherein L, L', A, Het and Y are as defined above.

More preferred compounds of formula (I) are those wherein:

20 - either (a) R¹ is C₃-C₈ alkyl, C₃-C₆ alkenyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L' or -L-Het-A, wherein:

L is C₁-C₄ alkyl;

L' is C₁-C₄ alkyl;

A is phenyl, 5- to 6-membered heterocyclyl or 5- to 6-membered heteroaryl;

Het is -O-, -S- or -NMe-;

30 - R^2 is hydrogen or C_1 - C_4 alkyl;
- R^3 is hydrogen or C_1 - C_2 alkyl;
- or (b) R^1 and R^2 form, together with the nitrogen to which they are attached, a 5- to 6-membered heterocyclic ring and R^3 is as defined above;
. - or (c) R^1 and R^3 form, together with the $-N-C=N-$ moiety to which they are attached, a 5- to 6-membered heterocyclic ring and R^2 is as defined above;
- R^4 is hydrogen or C_1 - C_2 alkyl;


- B is C₂-C₄ alkyl;;
- X is -O- or -NH-;
- R⁵ is hydrogen, C₁-C₆ alkyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Het-A or -L-A-Het-A, wherein L, L' and Het are as defined above and wherein each A is the same or different and is as defined above;
- 5 - R⁶ is hydrogen or C₁-C₂ alkyl;
- R⁷ is hydrogen or C₁-C₂ alkyl or R⁷ is a C₁-C₂ alkyl moiety which is joined to one of the carbon atoms of B to form, together with the -N-C- moiety to which R⁷ and B are attached, a pyrrolidine or piperidine ring; and
- 10 - R⁸ is hydrogen or C₁-C₂ alkyl;
- wherein:
 - the alkyl, alkenyl and alkynyl groups and moieties in the substituents R¹ to R⁷ and B are unsubstituted or substituted by a single hydroxy substituent or by one, two or three substituents which are the same or different and are selected from fluorine and chlorine substituents; and
 - the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R¹ and R⁵, the heterocyclic or heteroaryl moieties formed by R¹ together with R² and R¹ together with R³ and the heterocyclic moieties 15 formed by R⁷ together with B are typically unsubstituted or substituted by one, two or three substituents selected from fluorine, chlorine, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy and C₁-C₄ haloalkyl substituents.

Examples of such compounds are the more preferred compounds of formula (I) as defined above wherein:

25 A is phenyl or 5- to 6-membered heteroaryl; and

R⁵ is hydrogen, C₁-C₆ alkyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L' or -L-Het-A, wherein L, L', A and Het are as defined above.

Particularly preferred compounds of formula (I) are compounds of formula 30 (II)



wherein:

- either (a) R^{11} is C_3 - C_8 alkyl, allyl, phenyl, cyclopropyl, cyclohexyl, $-(C_1$ - C_2 alkyl)-phenyl, $-(C_1$ - C_2 alkyl)-thienyl, $-(C_1$ - C_2 alkyl)-tetrahydrofuran, $-(C_1$ - C_2 alkyl)-O-phenyl, $-(C_1$ - C_2 alkyl)-O- $(C_1$ - C_4 alkyl), $-(C_1$ - C_2 alkyl)-S- $(C_1$ - C_2 alkyl) or $-(C_1$ - C_2 alkyl)-NMe- $(C_1$ - C_2 alkyl);
- R^{21} is hydrogen, methyl or ethyl;
- or (b) R^{11} and R^{21} form, together with the nitrogen to which they are attached, a piperidinyl, morpholinyl or pyrrolidinyl ring;
- B^1 is a 1,2-ethyl or 1,3-propyl moiety; and
- R^{51} is hydrogen, C_1 - C_4 alkyl, $-(C_1$ - C_2 alkyl)-phenyl, $-(C_1$ - C_2 alkyl)-O- $(C_1$ - C_2 alkyl) or $-(C_1$ - C_2 alkyl)-phenyl-O-phenyl;

wherein:

- 15 - the alkyl groups and moieties in the substituents R^{11} , R^{21} , B and R^{51} are unsubstituted or substituted by a single fluoro substituent; and
- the phenyl and carbocyclyl groups in R^{11} and the heterocyclic moieties formed by R^{11} together with R^{21} are unsubstituted and the phenyl groups in R^{51} are unsubstituted or substituted by one or two substituents selected from fluorine, C_1 - C_2 alkyl, C_1 - C_2 alkoxy and C_1 - C_2 haloalkyl substituents.

Examples of such compounds are the particularly preferred compounds of formula (II) as defined above wherein:

- R^{11} is C_3 - C_8 alkyl, allyl, phenyl, cyclopropyl, cyclohexyl, $-(C_1$ - C_2 alkyl)-phenyl, $-(C_1$ - C_2 alkyl)-thienyl, $-(C_1$ - C_2 alkyl)-O-phenyl, $-(C_1$ - C_2 alkyl)-O- $(C_1$ - C_4 alkyl), $-(C_1$ - C_2 alkyl)-S- $(C_1$ - C_2 alkyl) or $-(C_1$ - C_2 alkyl)-NMe- $(C_1$ - C_2 alkyl); and
- R^{51} is hydrogen, C_1 - C_4 alkyl or $-(C_1$ - C_2 alkyl)-phenyl, wherein:
- the phenyl and carbocyclyl groups in R^{11} , the phenyl groups in R^{51} and the heterocyclic moieties formed by R^{11} together with R^{21} are unsubstituted.

In one embodiment, R¹¹ is preferably phenyl, -(C₁-C₂ alkyl)-phenyl, -(C₁-C₂ alkyl)-thienyl, -(C₁-C₂ alkyl)-tetrahydrofuranyl, -(C₁-C₂ alkyl)-O-phenyl, -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl), -(C₁-C₂ alkyl)-S-(C₁-C₂ alkyl) or -(C₁-C₂ alkyl)-NMe-(C₁-C₂ alkyl). In a preferred embodiment of formula (II), R¹¹ is -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl). In a particularly preferred embodiment of formula (II), R¹¹ is -(CH₂)₂-O-CH₃.

In a further preferred embodiment of formula (II), R⁵¹ is C₁-C₄ alkyl or -(C₁-C₂)-phenyl. In a particularly preferred embodiment of formula (II), R⁵¹ is -CH₂-phenyl.

In a preferred embodiment of formula (II), B¹ is a 1,3-propyl moiety.

10 In a further preferred embodiment of formula (II), R¹¹ is -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl) and R⁵¹ is C₁-C₄ alkyl or -(C₁-C₂)-phenyl. In a particularly preferred embodiment of formula (II), R¹¹ is -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl) and R⁵¹ is C₁-C₄ alkyl.

Examples of these particularly preferred compounds of the invention are:

15 (S)-2-Amino-4-(N'-isopropylguanidino)butanoic acid;
 (S)-2-Amino-4-(N'-propylguanidino)butanoic acid;
 (S)-2-Amino-4-(N'-benzylguanidino)butanoic acid;
 (S)-2-Amino-4-(N'-cyclohexylguanidino)butanoic acid;
 (S)-2-Amino-4-(N'-phenylguanidino)butanoic acid;
 20 (S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid;
 (S)-2-Amino-4-(N'-(2-thiophenemethyl)guanidino)butanoic acid;
 (S)-2-Amino-4-(N'-octylguanidino)butanoic acid;
 (S)-2-Amino-4-(N'-cyclopropylguanidino)butanoic acid
 (S)-2-Amino-4-(N'-(2'-dimethylaminoethyl)guanidino)butanoic acid;
 25 (S)-2-Amino-4-(N'-2,2-dimethylpropyl)guanidino)butanoic acid;
 (S)-2-Amino-4-(N'-(2-phenoxyethyl)guanidino)butanoic acid;
 (S)-2-Amino-4-(N'-(2-methylthioethyl)guanidino)butanoic acid;
 (S)-2-Amino-4-(N'-(2-isopropoxyethyl)guanidino)butanoic acid;
 (S)-2-Amino-4-(N'-allylguanidino)butanoic acid;
 30 (S)-2-Amino-4-[(piperidine-1-carboximidoyl)-amino]butanoic acid;
 (S)-2-Amino-4-[(pyrrolidine-1-carboximidoyl)-amino]butanoic acid;
 (S)-2-Amino-4-[(morpholine-4-carboximidoyl)-amino]butanoic acid;
 (S)-4-(N'-(2-methoxyethyl)guanidino)pyrrolidine-2-carboxylic acid;

(*S*)-4-*N*^G-(2-methoxyethyl)guanidinobutanoic acid;
N^G-(2-methoxyethyl)-L-arginine;
(*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid methyl ester;
(*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid ethyl ester;
5 (*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid propyl ester;
(*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid butyl ester;
(*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid benzyl ester;
(*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid isopropyl ester;
N^G-(2-Methoxyethyl)-L-arginine methyl ester;
10 *N*^G-(2-Methoxyethyl)-L-arginine benzyl ester;
(*S*)-2-Amino-5-(*N*^G-(2-methoxyethyl)guanidino)pentanoic acid methylamide;
(*S*)-2-Amino-4-(*N*^G-(2-methoxyethyl)guanidino)butanoic acid benzylamide;
(*S*)-*N*^G-((Tetrahydrofuran-2-yl)methyl)-arginine;
(2*S*)-2-Amino-4-*N*^G-((tetrahydrofuran-2-yl)methyl)guanidinobutanoic acid;
15 2-Methoxybenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate;
4-Methoxybenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate;
2-Fluorobenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate;
3-Trifluoromethyl-4-fluorobenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))
guanidino]butanoate;
20 3-Phenoxybenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate;
3-Methylbenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate;
3-Trifluoromethylbenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate;
3-Fluorobenzyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate; and
25 2-Methoxyethyl (*S*)-2-amino-4-[(*N*^G-(2-methoxyethyl))guanidino]butanoate,
and pharmaceutically acceptable salts thereof.

As used herein, a pharmaceutically acceptable salt is a salt with a pharmaceutically acceptable acid or base. Pharmaceutically acceptable acids include both inorganic acids such as hydrochloric, sulphuric, phosphoric, diphosphoric, hydrobromic or nitric acid and organic acids such as citric, fumaric, maleic, malic, ascorbic, succinic, tartaric, benzoic, acetic, methanesulphonic, ethanesulphonic, 30 benzenesulphonic or p-toluenesulphonic acid. Pharmaceutically acceptable bases include alkali metal (e.g. sodium or potassium) and alkali earth metal (e.g. calcium or


magnesium) hydroxides and organic bases such as alkyl amines, aralkyl amines or heterocyclic amines.

The compounds of the present invention have the chirality shown in formula (I). However, the compounds can also have chiral centres at other points in the molecule. For the avoidance of doubt, the chemical structures depicted herein are intended to embrace all stereoisomers of the compounds shown, provided they have the chirality at the carbon α to the carbonyl group which is shown in formula (I).

The compounds of formula (I) may be prepared by conventional routes, for example those set out in schemes 1 and 2 below.

10


Reaction Scheme 1



Compounds of formula (I) may be prepared from a substituted guanidine in a two step reaction as shown in Reaction Scheme 1, wherein R^1 to R^8 , B and X are as defined above. In the first step, a substituent is introduced onto the carboxamide nitrogen by reaction with the appropriate alcohol under Mitsunobu conditions. In the second step a further substituent is introduced by displacement of the pyrazole moiety with a primary or secondary amine. Where necessary, when R^2 , R^3 , R^5 or R^7 are hydrogen, they can be replaced by a suitable protecting group which can then be removed after the second step. A suitable protecting group is Boc which can be removed under acid conditions, such as treatment with HCl in 1,4-dioxane. The skilled person will readily understand when such protection is necessary.

Compounds of formulae (II), (III) and (V) are known compounds or can be prepared by the skilled person using known methods. For example, compounds of formula (I) can be prepared from bis-*tert*-butoxycarbonylpyrazole-1-carboxamidine.

5 Reaction Scheme 2



Reaction Scheme 2 is a variant of Reaction Scheme 1 and can be used to
10 prepare compounds of formula (I) wherein R¹ is a primary or secondary alkyl group and R² to R⁸, B and X are as defined above. Compounds of formulae (VI), (VII) and (IX) are known compounds or can be prepared by the skilled person using known methods. For example, compounds of formula (VI) can be prepared from bis-*tert*-butoxycarbonylpyrazole-1-carboxamidine.

15 The thus obtained compounds of formula (I) may be salfied by treatment with an appropriate acid or base. Racemic mixtures obtained by any of the above processes can be resolved by standard techniques, for example elution on a chiral chromatography column.

An inhibitor of a DDAH is one which, when present, produces a measurable
20 reduction in DDAH activity.

Preferred inhibitors are those which reduce DDAH activity by at least 10%, at least 20%, at least 30%, at least 40% at least 50%, at least 60%, at least 70%,

at least 80%, at least 90%, at least 95% or at least 99% at a concentration of the inhibitor of 1 $\mu\text{g ml}^{-1}$, 10 $\mu\text{g ml}^{-1}$, 100 $\mu\text{g ml}^{-1}$, 500 $\mu\text{g ml}^{-1}$, 1mg ml^{-1} , 10mg ml^{-1} or 100mg ml^{-1} .

5 The percentage inhibition represents the percentage decrease in activity in a comparison of assays in the presence and absence of the test substance. Any combination of the above mentioned degrees of percentage inhibition or activation and concentration of inhibitor or activator may be used to define an inhibitor or activator of the invention, with greater inhibition at lower concentrations being preferred.

10 Inhibition may occur if, for example, the inhibitor resembles the substrate and binds at the active site of the DDAH. The substrate is thus prevented from binding to the same active site and the rate of catalysis is reduced by reducing the proportion of enzyme molecules bound to substrate (competitive inhibition). An inhibitor may also exert its effects by non-competitive inhibition where the inhibitor and substrate 15 can bind simultaneously to DDAH and thus bind at different non-overlapping sites. Inhibition occurs as the turnover number of the DDAH decreases.

Typically, the compounds of the present invention cause substantially no inhibition of NOS expression or activity.

20 A suitable assay for inhibition of NOS activity is the [^{14}C]Arginine - [^{14}C]citrulline method which is described in D.S. Bredt and S.H. Snyder, Nitric oxide mediates glutamate-linked enhancement of cGMP levels in the cerebellum, Proc Natl Acad Sci U S A, 1989 Nov, 86(22), 9030-3.

25 Typically, the compounds of the present invention achieve less than 50% inhibition, preferably less than 20 % inhibition, more preferably less than 10% inhibition of NOS and particularly preferably less than 5% inhibition. In a preferred embodiment, the compounds of the present invention cause no detectable inhibition of NOS expression or activity. Generally, such results are achieved at a concentration of compound of 0.01 to 10 μm , for example 0.1 to 5, or 1 to 2 μm .

30 In one embodiment it is preferred that an inhibitor for use in the invention shows at least 100x, 1000x or at least 10^6 x greater percentage inhibition of DDAH methylarginase activity and/or expression at a given concentration of inhibitor than the percentage inhibition of NOS at that same concentration.

In one embodiment, a compound of the invention has substantially no effect, activatory or inhibitory, upon NOS activity and/or substantially no effect upon NOS expression.

The compounds of the invention are found to be inhibitors of DDAH. The 5 compounds of the present invention can therefore be used to treat a condition treatable by a DDAH inhibitor. The compounds of the invention are therefore therapeutically useful. Accordingly, the present invention provides a compound of the formula (I), as defined above, or a pharmaceutically acceptable salt thereof, for use in the treatment of the human or animal body. In particular, in the treatment of a 10 condition in which the abnormal metabolism of NO is implicated. The present invention relates to the use of a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof for use in the manufacture of a medicament for use in the treatment of a disease treatable by a DDAH inhibitor. In one embodiment, the present invention provides a compound of the formula (I), as 15 defined above, or a pharmaceutically acceptable salt thereof, for use in the treatment of the human or animal body, provided that the compound of formula (I) N^G -propyl-L-arginine, N^G -allyl-L-arginine, N^G -propargyl-L-arginine, N^G -butyl-L-arginine or N^G -cyclopropyl-L-arginine. The compounds of the invention are believed to be novel and the present invention thus provides for compounds of formula (I), as 20 defined above, or a pharmaceutically acceptable salt thereof. In one embodiment, the present invention provides a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof provided that the compound of formula (I) is not N^G -propyl-L-arginine, N^G -allyl-L-arginine, N^G -propargyl-L-arginine, N^G -butyl-L-arginine or N^G -cyclopropyl-L-arginine. Also provided is a pharmaceutical 25 composition comprising a compound of the formula (I), as defined above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent. In one embodiment, the present invention provides a pharmaceutical composition comprising a compound of the formula (I), as defined above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent, provided that the compound of formula (I) is not N^G -propyl-L-arginine, 30 N^G -allyl-L-arginine, N^G -propargyl-L-arginine, N^G -butyl-L-arginine or N^G -cyclopropyl-L-arginine. Said pharmaceutical composition typically contains up to 85 wt% of a compound of the invention. More typically, it contains up to 50 wt% of

a compound of the invention. Preferred pharmaceutical compositions are sterile and pyrogen free. Further, the pharmaceutical compositions provided by the invention typically contain a compound of the invention which is a substantially pure optical isomer.

5 As used herein, the term "disease whose pathology is affected by DDAH" refers to any disease wherein the level and/or activity of DDAH is not at the required amount to cause reversal of the pathology of the disease. In one embodiment, such diseases are associated with an increase in the expression of DDAH. Such diseases can be identified by measuring the expression of DDAH in a patient with a particular 10 disease and comparing the results with a control. In another embodiment, a disease whose pathology is affected by DDAH is a disease wherein alteration in the level and/or activity of DDAH causes a change in the level and/or activity of another species which is associated with the disease. An example of such a species is NO. Such diseases can be identified by measuring the amount of this species in a patient 15 before and after the administration of a compound of the present invention and determining whether there has been a change in the level and/or activity of said species.

The compounds of the invention may be administered in a variety of dosage forms. Thus, they can be administered orally, for example as tablets, troches, 20 lozenges, aqueous or oily suspensions, dispersible powders or granules. Preferred pharmaceutical compositions of the invention are compositions suitable for oral administration, for example tablets and capsules.

Compositions suitable for oral administration may, if required, contain a colouring or flavoring agent. Typically, a said capsule or tablet comprises from 5 to 25 500 mg, preferably 10 to 500 mg, more preferably 15 to 100 mg, of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

The compounds of the invention may also be administered parenterally, whether subcutaneously, intravenously, intramuscularly, intrasternally, transdermally or by infusion techniques. The compounds may also be administered as 30 suppositories.

One preferred route of administration is inhalation. The major advantages of inhaled medications are their direct delivery to the area of rich blood supply in comparison to many medications taken by oral route. Thus, the absorption is very

rapid as the alveoli have an enormous surface area and rich blood supply and first pass metabolism is bypassed.

Preferred pharmaceutical compositions of the invention therefore include those suitable for inhalation. The present invention also provides an inhalation 5 device containing such a pharmaceutical composition. Typically said device is a metered dose inhaler (MDI), which contains a pharmaceutically acceptable chemical propellant to push the medication out of the inhaler. Typically, said propellant is a fluorocarbon.

Further preferred inhalation devices include nebulizers. Nebulizers are 10 devices capable of delivering fine liquid mists of medication through a "mask" that fits over the nose and mouth, using air or oxygen under pressure. They are frequently used to treat those with asthma who cannot use an inhaler, including infants, young children and acutely ill patients of all ages.

Said inhalation device can also be, for example, a rotary inhaler or a dry 15 powder inhaler, capable of delivering a compound of the invention without a propellant.

Typically, said inhalation device contains a spacer. A spacer is a device which enables individuals to inhale a greater amount of medication directly into the lower airways, where it is intended to go, rather than into the throat. Many spacers 20 fit on the end of an inhaler; for some, the canister of medication fits into the device. Spacers with withholding chambers and one-way valves prevent medication from escaping into the air. Many people, especially young children and the elderly, may have difficulties coordinating their inhalation with the action necessary to trigger a puff from a metered dose inhaler. For these patients, use of a spacer is particularly 25 recommended.

Another preferred route of administration is intranasal administration. The nasal cavity's highly permeable tissue is very receptive to medication and absorbs it quickly and efficiently, more so than drugs in tablet form. Nasal drug delivery is less painful and invasive than injections, generating less anxiety among patients. Drugs 30 can be delivered nasally in smaller doses than medication delivered in tablet form. By this method absorption is very rapid and first pass metabolism is bypassed, thus reducing inter-patient variability. Nasal delivery devices further allow medication to be administered in precise, metered doses. Thus, the pharmaceutical compositions of

the invention are typically suitable for intranasal administration. Further, the present invention also provides an intranasal device containing such a pharmaceutical composition.

A further preferred route of administration is transdermal administration. The 5 present invention therefore also provides a transdermal patch containing a compound of the invention, or a pharmaceutically acceptable salt thereof. Also preferred is sublingual administration. The present invention therefore also provides a sub-lingual tablet comprising a compound of the invention or a pharmaceutically acceptable salt thereof.

10 A compound of the invention is typically formulated for administration with a pharmaceutically acceptable carrier or diluent. For example, solid oral forms may contain, together with the active compound, diluents, e.g. lactose, dextrose, saccharose, cellulose, corn starch or potato starch; lubricants, e.g. silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; binding agents; 15 e.g. starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinyl pyrrolidone; disaggregating agents, e.g. starch, alginic acid, alginates or sodium starch glycolate; effervescent mixtures; dyestuffs; sweeteners; wetting agents, such as lecithin, polysorbates, laurylsulphates; and, in general, non toxic and pharmacologically inactive substances used in pharmaceutical formulations. Such 20 pharmaceutical preparations may be manufactured in known manner, for example, by means of mixing, granulating, tableting, sugar coating, or film coating processes.

Liquid dispersions for oral administration may be syrups, emulsions and suspensions. The syrups may contain as carriers, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

25 Suspensions and emulsions may contain as carrier, for example a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspension or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol, and if desired, 30 a suitable amount of lidocaine hydrochloride.

A further preferred route of administration is intravenous administration. Solutions for injection or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions.

The compounds of the invention are inhibitors of DDAH and may be used in the treatment of conditions in which increased NO production is implicated. In particular, conditions such as ischaemia-reperfusion injury of the brain or heart, cancer, lethal hypotension in severe inflammatory conditions such as septic shock or 5 multi-organ failure, or local and systemic inflammatory disorders including arthritis, skin disorders, inflammatory cardiac disease or migraine may be treated. Further, conditions such as neurodegeneration and asthma, wherein inhibition of NO has been suggested as a therapeutic option, can also be treated. The compounds of the invention can also be used in the treatment of ischaemia-reperfusion injury of the 10 brain or heart, cancer, lethal hypotension in severe inflammatory conditions, local and systemic inflammatory disorders, neurodegeneration, asthma, pain or sepsis. The compounds of the present invention can further be used in the treatment of peritoneal inflammation. Preferably, the compounds of the present invention are used in the treatment of ischaemia-reperfusion injury of the brain or heart, cancer, 15 lethal hypotension in severe inflammatory conditions, local and systemic inflammatory disorders, pain or sepsis.

Increased DDAH expression has also been implicated in an increase in tumour growth and angiogenesis (V. Kostourou, S.P. Robinson, J.E. Cartwright and G. St.J. Whitley, British Journal of Cancer 2002, 87, 673-680) and has been further implicated in 20 enhanced tumour hypoxia (S.P. Robinson, V. Kostourou, H. Troy, J.F. Murray, G. Whitley and J.R. Griffiths, Clinical Cancer Research, 2003, 9, 6209s). DDAH over-expression has also been shown to enhance VEGF expression and the tube-forming capacity of cells. The compounds of the present invention can therefore be used in the treatment of conditions associated with tumour growth, angiogenesis and hypoxia and also 25 enhanced VEGF expression. The compounds of the present invention can therefore be used for anti-angiogenic and anti-cancer therapy. In particular, the types of cancer that can be treated by the compounds of the present invention are those which are angiogenic in nature. The compounds of the present invention can further be used in the treatment of a tumour which overexpresses DDAH, for example DDAH1 overexpressing glioma (brain 30 tumour) or pancreatic cancer.

The compounds of the present invention are thought to be particularly useful in the treatment of solid tumours. Examples of such solid tumours include ovarian cancer, colorectal cancer, breast cancer, brain cancer, liver cancer, kidney cancer, stomach cancer,

prostate cancer, lung cancer, thyroid cancer, pancreatic cancer, Kaposi's sarcoma and skin cancer. The compounds of the present invention are also thought to be useful in the treatment of head, neck and oesophageal solid tumours. Skin cancer is a preferred example a solid tumour for treatment by the compounds of the present invention and a 5 preferred type of skin cancer is melanoma. A further preferred solid tumour for treatment by the compounds of the present invention is pancreatic cancer.

Inhibition of DDAH will result in an increase in the concentration of L-NMMA. L-NMMA has been shown to be effective in treating conditions such as inflammatory disease states, pain and sepsis. The compounds of the present invention can therefore be 10 used in the treatment of such conditions.

Alternatively, the compounds of the present invention could be used as a joint therapy together with an inhibitor of NOS activity (for example, a methylarginine). For example, a specific inhibitor of a DDAH isoform could be used with the methylarginine L-NMMA. This approach may radically alter the activity profile of L-NMMA and may 15 result in L-NMMA having an increased inhibitory effect for a specific NOS isoform. Thus, the invention provides products containing an inhibitor of a DDAH activity and/or expression and a methylarginine as a combined preparation for simultaneous, separate or sequential use in the treatment of ischaemia-reperfusion injury of the brain or heart, cancer, lethal hypotension in severe inflammatory conditions such as septic shock or 20 multi-organ failure, or local and systemic inflammatory disorders including arthritis, skin disorders, inflammatory cardiac disease or migraine.

The compounds of the present invention are thought to be particularly suitable for acute clinical use.

The compounds of the present invention may also be used as antimicrobial agents, 25 for example antibacterial agents. Therefore, the invention also provides a chemical entity for use in the treatment of a bacterial infection.

An inhibition in DDAH activity and/or expression has also been implicated in the treatment of inflammatory pain. A compound of the present invention is therefore useful for combating pain in a human or animal. An agent may act to alleviate existing pain in 30 an individual or may be administered in anticipation of a painful condition. Thus treatment may be therapeutic or prophylactic. Preferably the compound is suitable for use in a mammal. In particular, it is preferred that the subject for treatment is human. However, the invention will also be of veterinary use for treating livestock and domestic animals.

For example, the invention may be of use for treating cattle, pigs, sheep, horses, dogs, cats or rabbits.

Generally the pain to be treated is one in which pain signals are processed at least in part by the thalamus. For example, the pain may be "fast pain" for example, 5 sharp pain, pricking pain, electric pain, or "slow pain", for example, burning pain, aching pain, throbbing pain, nauseous pain.

Attempts have been made to classify different types of pain. Among those classes which have been broadly identified, but which nevertheless overlap to an extent, are acute and chronic pain.

10 A definition of acute pain (Halpern (1984) *Advances in Pain Research and Therapy* Vol.7, Ed. C. Bendetti et al, p147) which is not intended to be limiting, is as a constellation of unpleasant sensory, perceptual and emotional experiences of certain 15 associate autonomic (reflex) responses, and of psychological and behavioural reactions provoked by injury or disease. Tissue injury provokes a series of noxious stimuli which are transduced by nociceptors to impulses transmitted to the spinal cord and then to the upper part of the nervous system. Examples of acute pain are dental pain, post-operative pain, obstetric pain, headaches, neuralgia and myalgia.

20 A definition of chronic pain (Halpern (1984) *ibid.*) also not limiting, is pain that persists beyond the usual course of an acute disease or beyond a reasonable time for an 25 injury to heal. Chronic pain is typically a result of persistent dysfunction of the nociceptive pain system. Examples of chronic pain include trigeminal neuralgia, post-herpetic neuralgia (a form of chronic pain accompanied by skin changes in a dermatomal distribution following damage by acute Herpes Zoster disease), diabetic neuropathy, causalgia, "phantom limb" pain and pain associated with osteoarthritis, rheumatoid arthritis and cancer. Some of these, for example, trigeminal neuralgia, diabetic 30 neuropathic pain, causalgia, phantom limb pain and central post-stroke pain, have also been classified as neurogenic pain. One non-limiting definition of neurogenic pain is pain caused by dysfunction of the peripheral or central nervous system in the absence of nociceptor stimulation by trauma or disease.

Physiological and pathological pain have been defined in terms of their origin in the plasticity of the nervous system. The latter is defined in turn as the alteration in structure or function of the nervous system caused by development, experience or injury and can be either adaptive or maladaptive (Cervero, F.(1991) *Eur. J. Neurosci Suppl 4,*

162). Adaptive plasticity underlies the ability of the nervous system to compensate for damage or to produce changes in function which are appropriate to environmental change. Physiological pain, considered to be a sensation which reflects specific peripheral stimuli, is based on adaptive plasticity.

5 Maladaptive plasticity comprises those changes in the nervous system which lead to a disruption of function and therefore effectively constitute a disease state. Pathological pain is considered to be a sensation resulting from changes within the nervous system which bring about an alteration in the way in which information from the periphery, some of which is quite normal, is handled. Pathological pain is therefore based on maladaptive 10 plasticity (Woolf, C.J. (1989) *Br. J. Anaesth.* 63, 139-146).

Maladaptive plasticity of the nociceptive system has also been shown, in experimental models, to be present in states of chronic pain. For instance, multiple injections of hyperalgesic substances such as PGE₂ into the paw of a rat have been shown to induce sustained hyperalgesia to mild pressure (e.g. Nakamura-Craig and Smith (1989) 15 *Pain* 38, 91-98; Ferreira et al (1990) *Pain* 42, 365-371; Nakamura-Craig and Gill (1991) *Neurosci. Lett.* 124, 49-51).

A number of animal models have been developed of neuropathic pain in particular peripheral neuropathic pain, which suggest that this pain is often associated with partial denervation (Decosterd I and Woolf C J (2000) *Pain* 00 : 1-10). Such models mimic in 20 particular, pain associated with, for example diabetic neuropathy, postherpetic neuralgia, toxic neuropathies, compression neuropathies and trauma, characterised by spontaneous lancinating, burning pain and shock like pain, as well as pain hypersensitivity including tactile allodynia, pinprick hyperalgesia and hyperpathia.

The compounds of the present invention can be used in a method of treating, 25 including preventing, pain in a human or animal, the method comprising administering thereto a therapeutically or prophylactically effective amount of the agent. The compounds can be used in the therapeutic or prophylactic treatment of pathological conditions in which pain occurs. The condition of a human or animal to which the compound is administered can thereby be improved.

30 The compounds of the present invention are useful for treating or preventing different types of pain including chronic pain and acute pain. Examples of chronic pain which can be targeted using the present inhibitors include trigeminal neuralgia, post-herpetic neuralgia, painful diabetic neuropathy, causalgia, central post-stroke pain,

“phantom limb” pain, atypical facial pain, back pain, headaches, neuralgia and pain associated with osteoarthritis, rheumatoid arthritis and cancer.

Examples of acute pain which can be treated using the compounds of the present invention include dental pain, post-operative pain, obstetric pain, headaches, neuralgia and myalgia. In particular, the compounds can be administered pre-operatively to counteract the acute pain associated with surgical operations, including dental surgery and labour pain. In a preferred embodiment the present invention comprises a method of administering a therapeutically effective amount of a compound to a patient prior to a dental operation, for example a tooth extraction. Virtually no analgesic agents currently in use are effective in controlling pain when administered pre-operatively in this way.

The invention also provides a method for treating pain according to which a compound of the present invention is administered jointly together with one or more other agents. For example, a compound may be administered with a known anti-pain agent. In one embodiment a compound of the present invention is administered with an inhibitor of NOS activity such as a methylarginine. For example, a compound of the present invention may be used in conjunction with L-NMMA and/or ADMA. This approach may radically alter the activity profile of L-NMMA and/or ADMA and may result in L-NMMA and/or ADMA having an increased inhibitory effect for NOS.

A compound of the present invention may be administered in parallel with another agent (such as methylarginine), or the agents may be administered sequentially, one following on from the other. Alternatively, a compound of the present invention and another agent (for example methylarginine), while acting together for the same purpose, may be administered separately. Thus, the invention provides products containing a compound of the present invention and a modulator of DDAH methylarginase activity and/or expression and a further agent (such as methylarginine) as a combined preparation for simultaneous, separate or sequential use in the treatment of pain in a human or animal.

As described, a compound of the present invention may be used to improve the condition of a patient experiencing pain or to prevent or reduce future pain. The formulation of a compound of the present invention for administration in preventing or alleviating pain will depend largely upon the nature of the compound. Formulation is also influenced, for example, by whether a pharmaceutical or

veterinary use is intended, and by the requirement for passage across the blood-brain barrier. A compound may also be formulated for simultaneous, separate or sequential use with another substance such as a methylarginine. In general, it is preferred that the compound is directed or administered to neuronal cells, especially 5 those of the CNS.

In one embodiment, the activity of the compounds of the invention is restricted only to particular cell types. For example, activity may be limited to neuronal cells especially those of the CNS, in particular thalamic cells. A compound may be formulated for uptake specifically by such cells. Alternatively, a compound of the present invention 10 may be administered directly to such cells. For example, an agent may be injected into the thalamus.

The term treatment in this context is deemed to cover any effect from a cure of said condition to alleviation of any or all of the symptoms. The compounds of the invention may, where appropriate, be used prophylactically to reduce the incidence 15 or severity of said conditions.

A therapeutically effective amount of a compound of the invention is administered to a patient. A typical dose is from about 0.001 to 50 mg per kg of body weight, for example 0.01 to 10 mg, according to the activity of the specific compound, the age, weight and conditions of the subject to be treated, the type and severity of the disease and the 20 frequency and route of administration. Preferably, daily dosage levels are from 5 mg to 2 g. A daily dose may be given in one or more than one, for example, 2, 3 or 4 administrations.

The following Examples illustrate the invention. They do not, however, limit the invention in any way. In this regard, it is important to understand that the particular assays 25 used in the Examples section are designed only to provide an indication of activity in inhibiting DDAH. A negative result in any one particular assay is not determinative.

EXAMPLES

General. All starting materials were either commercially available or reported previously in the literature unless noted. Solvents and reagents were used without further purification. Reactions were monitored by TLC on precoated silica gel plates (Kieselgel 60 F₂₅₄, Merck). Purification was performed by flash chromatography using silica gel (particle size 40-63 µM, Merck). ¹H, and ¹³C NMR spectra were recorded on a Bruker AMX-300 or Bruker AMX-400 spectrometer. Chemical shifts are reported as ppm (δ) relative to TMS as an internal standard. Mass spectra were recorded on either a VG ZAB SE spectrometer (electron impact and FAB) or a Micromass Quattro electrospray LC-mass spectrometer. Melting points were determined on a Gallenkamp melting point apparatus and are uncorrected. Microanalysis was carried out by the Analytical Services Section, University College London. All yields reported in the experimental section are non-optimized, isolated yields.

Method A: General method for synthesis of *N*^G-monosubstituted guanidine amino acids

Diethyl azodicarboxylate (3 mmol) was added dropwise to a solution of *N,N'*-bis-*tert*-butoxycarbonylpyrazole-1-carboxamidine (2 mmol), triphenylphosphine (3 mmol) and the appropriate alcohol (2 mmol) at 0°C, with stirring. The mixture was then stirred at room temperature for 3-16h, concentrated on the rotary evaporator and then subjected to flash column chromatography (10% ethyl acetate/cyclohexane) to give the product as a colourless oil.

The *N*-alkyl substituted pyrazolecarboxamidine (1.5 mmol), Boc-diaminoalkanoic acid *tert*-butyl ester (1.5 mmol) and diisopropylethylamine (2.0 mmol) were stirred in acetonitrile (10 mL) for 24 hours. The mixture was concentrated on the rotary evaporator, and then subjected to flash column chromatography (20-30% ethyl acetate/cyclohexane) to give the protected guanidino-amino acid as a colourless gum.

The protected amino acid was stirred in excess 4M hydrogen chloride/dioxane for 24-72 hours. Removal of solvent and byproducts *in vacuo* gave the amino acid as a

white hygroscopic solid.

Method B: General method for synthesis of N^G -disubstituted 2-amino-4-guanidinobutanoic acids and N^G -aryl 2-amino-4-guanidinobutanoic acids

5 Diethyl azodicarboxylate (3 mmol) was added dropwise to a solution of N,N' -bis-*tert*-butoxycarbonylpyrazole-1-carboxamidine (2 mmol), triphenylphosphine (3 mmol) and *N*-Boc-homoserine-*tert*-butyl ester (2 mmol) (prepared from the acid by the method described in Mathias, L. J. *Synthesis* 1979, 561-576) at 0°C, with stirring. The mixture was stirred at room temperature for 3-16 hours, concentrated on the 10 rotary evaporator and then subjected to flash column chromatography (20% ethyl acetate/cyclohexane) to give the product as a colourless oil.

The *N*-substituted pyrazolecarboxamidine (1.5 mmol), was stirred with the appropriate secondary or aryl amine (1.5 mmol) in acetonitrile (10 mL) for 24 hours. The mixture was concentrated on the rotary evaporator, and then subjected to flash 15 column chromatography (20-30% ethyl acetate/cyclohexane) to give the protected guanidino-amino acid as a colourless gum.

The protected amino acid was stirred in excess 4M hydrogen chloride/dioxane for 24-72 hours. Removal of solvent *in vacuo* gave the amino acid as a white hygroscopic solid.

20

Example 1 (S)-2-Amino-4-(*N'*-isopropylguanidino)butanoic acid

Yield 31% (Method A). ^1H NMR (CD₃OD, 300 MHz): δ 4.00 (1H, t, *J* 6.5 Hz, CH), 3.67 (1H, septet, *J* 6.4 Hz, NCH(Me)₂), 3.43 (2H, t, *J* 6.9 Hz, CH₂), 2.21-2.03 (2H, m, CH₂), 1.16 (6H, d, *J* 6.4 Hz, (CH₃)₂); ^{13}C NMR (75 MHz, CD₃OD, 27°C) δ 171.6 (C), 157.1 (C), 51.9 (CH), 46.7 (CH), 39.4 (CH₂), 31.5 (CH₂), 23.1 (2 x CH₃); MS (FAB+) found *m/z* 203.15101 (M+H), C₈H₁₈N₄O₂ M+H calc. 203.15709.

Example 2 (S)-2-Amino-4-(*N'*-propylguanidino)butanoic acid

Yield 35% (Method A). ^1H NMR (CD₃OD, 300 MHz): δ 4.07 (1H, t, *J* 6.8 Hz, CH), 3.50 (2H, t, *J* 7.0 Hz, CH₂), 3.18 (2H, t, *J* 7.1 Hz, CH₂), 2.26-2.13 (2H, m, CH₂), 1.62 (2H, td, *J* 7.3, 7.1 Hz, CH₂), 0.98 (3H, t, *J* 7.3 Hz, CH₃); ^{13}C NMR (75 MHz, CD₃OD, 27°C) δ 171.6 (C), 158.0 (C), 51.8 (CH), 44.8 (CH₂), 39.4 (CH₂), 31.4

(CH₂), 23.6 (CH₂), 11.8 (CH₃); MS (FAB+) found *m/z* 202.14300 (M+H), C₈H₁₈N₄O₂ M+H calc. 202.14297.

Example 3 (S)-2-Amino-4-(*N'*-benzylguanidino)butanoic acid

5 Yield 55% (Method A). ¹H NMR (CD₃OD, 300 MHz): δ 7.53-7.22 (5H, m, ArH), 4.59 (2H, s, PhCH₂), 4.15 (1H, t, *J* 6.7 Hz, CH), 3.66 (2H, t, *J* 7.2 Hz, CH₂), 2.41-2.24 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.5 (C), 158.0 (C), 137.9 (C), 130.3 (CH), 129.4 (CH), 128.8 (CH), 51.7 (CH), 46.5 (CH₂), 39.5 (CH₂), 31.4 (CH₂); MS (FAB+) found *m/z* 251.15045 (MH⁺), C₁₂H₁₈N₄O₂ M+H calc. 251.15079.

10

Example 4 (S)-2-Amino-4-(*N'*-cyclohexylguanidino)butanoic acid

Yield 12% (Method A). ¹H NMR (CD₃OD, 300 MHz): δ 3.99 (1H, t, *J* 6.2 Hz, CH), 3.43 (2H, t, *J* 6.8 Hz, CH₂), 2.21-2.06 (2H, m, CH₂), 1.87 (2H, br d, 2 x cyclohexyl H), 1.72 (2H, br d, 2 x cyclohexyl H), 1.55 (1H, m, NCH cyclohexyl), 1.41-1.14 (6H, m, 6 x cyclohexyl H); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 52.5 (CH), 52.0 (CH), 39.4 (CH₂), 34.2 (CH₂), 31.5 (CH₂), 26.6 (CH₂), 26.1 (CH₂); MS (FAB+) found *m/z* 243.18175 (M+H), C₁₁H₂₂N₄O₂ M+H calc. 243.18209.

Example 5 (S)-2-Amino-4-(*N'*-phenylguanidino)butanoic acid

20 Yield 17% (Method B). ¹H NMR (CD₃OD, 300 MHz): δ 7.47 (2H, t, *J* 7.3 Hz Ar C2H, C5H), 7.37 (1H, t, *J* 7.3 Hz, Ar C4H), 7.31 (2H, m, Ar C3H, C5H), 4.10 (1H, t, *J* 6.3 Hz, CH), 3.59 (2H, t, *J* 7.0 Hz, CH₂), 2.33-2.14 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.5 (C), 131.5 (CH), 129.2 (CH), 127.1 (CH), 51.9 (CH), 39.8 (CH₂), 31.4 (CH₂); MS (FAB+) found *m/z* 237.13543 (M+H), C₁₁H₁₆N₄O₂ M+H calc. 237.13514.

Example 6 (S)-2-Amino-4-(*N'*-(2-methoxyethyl)guanidino)butanoic acid

Yield 41% (Method A). ¹H NMR (CD₃OD, 300 MHz): δ 4.29 (1H, t, *J* 6.7 Hz, CH), 3.78-3.68 (4H, m, 2 x CH₂), 3.63 (2H, m, OCH₂), 3.61 (3H, s, OCH₃), 2.51-2.33 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.6 (C), 158.0 (C), 72.6 (CH₂), 59.6 (CH₃), 51.8 (CH), 43.5 (CH₂), 39.5 (CH₂), 31.4 (CH₂); MS (FAB+) found *m/z* 219.14585 (M+H), C₈H₁₈N₄O₃ M+H calc. 219.14585.

Example 7 (S)-2-Amino-4-(N'-octylguanidino)butanoic acid

Yield 18% (Method A). ^1H NMR (CD₃OD, 300 MHz): δ 4.13 (1H, t, *J* 6.5 Hz, CH), 3.56 (2H, t, *J* 6.7 Hz, NCH₂), 3.28 (2H, t, *J* 7.1 Hz, NCH₂), 2.37-2.17 (2H, m, CH₂), 1.42-1.38 (10H, m, 5 x CH₂), 0.97 (3H, m, CH₃); ^{13}C NMR (75 MHz, CD₃OD, 27°C) δ ; MS (FAB+), found *m/z* 273.22849 (M+H), C₁₃H₂₈N₄O₂ M+H calc. 273.22903.

Example 8 (S)-2-Amino-4-(N'-cyclopropylguanidino)butanoic acid

Yield 41% (Method B). ^1H NMR (CD₃OD, 300 MHz): δ 4.07 (1H, t, *J* 6.5 Hz, CH), 3.50 (2H, t, *J* 6.9 Hz, CH₂), 2.52 (1H, m, NCH cyclopropyl), 2.27-2.11 (2H, br m, CH₂), 0.89 (2H, m, 2 x CH cyclopropyl), 0.66 (2H, m, 2 x CH cyclopropyl); ^{13}C NMR (75 MHz, CD₃OD, 27°C) δ 171.6 (C), 159.3 (C), 51.9 (CH), 39.4 (CH₂), 31.4 (CH₂), 23.9 (CH), 8.2 (CH₂); MS (FAB+) found *m/z* 201.13519 (M+H), C₈H₁₆N₄O₂ M+H calc. 201.13514.

Example 9 (S)-2-Amino-4-(N'-(2'-dimethylaminoethyl)guanidino)butanoic acid
Yield .29% (Method B). ^1H NMR (CD₃OD, 300 MHz): δ 4.15 (1H, t, *J* 6.7 Hz, CH), 3.73 (2H, t, *J* 6.2 Hz, NCH₂), 3.56 (2H, t, *J* 7.1 Hz, NCH₂), 3.43 (2H, t, *J* 6.2 Hz, NCH₂), 2.96 (6H, s, N(CH₃)₂), 2.32-2.19 (2H, m, CH₂); ^{13}C NMR (75 MHz, CD₃OD, 27°C) δ 171.2 (C), 158.0 (C), 56.9 (CH₂), 51.4 (CH), 43.9 (CH₃), 39.2 (CH₂), 37.9 (CH₂), 30.8 (CH₂); MS (FAB+), found *m/z* 232.17283 (M+H), C₉H₂₁N₅O₂ M+H calc 232.17734.

Example 10 (S)-2-Amino-4-(N'-2,2-dimethylpropyl)guanidino)butanoic acid

Yield 18% (Method A). ^1H NMR (CD₃OD, 300 MHz): δ 4.18 (1H, t, *J* 6.7 Hz, CH), 3.62 (2H, t, *J* 7.3 Hz, CH₂), 3.14 (2H, s, NCH₂(Me)₃), 2.40-2.21 (2H, m, CH₂), 1.09 (9H, s, (CH₃)₃); ^{13}C NMR (75 MHz, CD₃OD, 27°C) δ 171.5 (C), 158.0 (C), 54.2 (CH₂), 51.8 (CH₂), 39.5 (CH₂), 33.5 (C), 31.5 (CH₂), 27.7 (3 x CH₃); MS (FAB+) found *m/z* 231.18222 (M+H), C₁₀H₂₂N₄O₂ M+H calc. 231.18209.

Example 11 (S)-2-Amino-4-(N'-(2-phenoxyethyl)guanidino)butanoic acid

Yield 56% (Method A). ^1H NMR (CD₃OD, 300 MHz): δ 7.31-7.24 (2H, m, 2 x ArH), 6.97-6.92 (3H, m, 3 x ArH), 4.14 (2H, t, *J* 5.1 Hz, CH₂), 4.09 (1H, t, *J* 6.9 Hz, CH), 3.66 (2H, t, *J* 5.1 Hz, CH₂), 3.52 (2H, t, *J* 7.1 Hz, CH₂), 2.30-2.12 (2H, m,

CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.2 (C), 159.8 (C), 158.0 (C), 130.6 (CH), 122.4 (CH), 115.6 (CH), 67.4 (CH₂), 51.4 (CH), 42.5 (CH₂), 39.1 (CH₂), 30.9 (CH₂); MS (FAB+) found *m/z* 281.16180 (M+H), C₁₃H₂₀N₄O₃ M+H calc. 281.16136.

5 Example 12 (*S*)-2-Amino-4-(*N'*-(2-methylthioethyl)guanidino)butanoic acid
Yield 38% (Method A). ¹H NMR (CD₃OD, 300 MHz): δ 4.09 (1H, t, *J* 6.6 Hz, CH),
3.51 (2H, t, *J* 7.1 Hz, CH₂), 3.46 (2H, t, *J* 6.7 Hz, CH₂), 2.72 (2H, t, *J* 6.7 Hz, SCH₂),
2.30-2.12 (2H, m, CH₂), 2.13 (3H, s, SCH₃); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ
171.2 (C), 157.6 (C), 51.4 (CH), 41.8 (CH₂), 39.0 (CH₂), 33.9 (CH₂), 30.9 (CH₂),
10 15.2 (CH₃); MS (FAB+) found *m/z* 235.12235 (M+H), C₈H₁₈N₄O₂S M+H calc.
235.12287.

20 Example 13 (*S*)-2-Amino-4-(*N'*-(2-isopropoxyethyl)guanidino)butanoic acid
Yield 41% (Method A). ¹H NMR (CD₃OD, 300 MHz): δ 4.08 (1H, t, *J* 6.5 Hz, CH),
15 3.65 (1H, septet, *J* 6.1 Hz, OCH), 3.59 (2H, t, *J* 5.1 Hz, CH₂), 3.50 (2H, t, *J* 7.1 Hz,
CH₂), 3.39 (2H, t, *J* 5.1 Hz, CH₂), 2.30-2.11 (2H, m, CH₂), 1.17 (5H, d, *J* 6.1 Hz, 2 x
CH₃); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.2 (C), 158.4 (C), 73.7 (CH), 51.4
(CH), 43.6 (CH₂), 39.1 (CH₂), 31.0 (CH₂), 22.3 (CH₃); MS (FAB+) found *m/z*
247.17699 (M+H), C₁₀H₂₂N₄O₃ M+H calc. 247.17701.

20 Example 14 (*S*)-2-Amino-4-(*N'*-allylguanidino)butanoic acid
Yield 50% (Method A). ¹H NMR (CD₃OD, 300 MHz): δ 5.99-5.87 (1H, m, vinyl
CH), 5.34 (1H, d, *J* 17.1 Hz, cis vinyl CH), 5.27 (1H, d, *J* 10.4 Hz, trans vinyl CH),
4.13 (1H, t, *J* 6.6 Hz, NCH), 3.93 (2H, d, *J* 4.9 Hz. NCH₂), 3.57 (2H, t, *J* 6.0 Hz,
25 CH₂), 2.37-2.15 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.5 (C),
158.0 (C), 134.1 (CH), 117.9 (CH₂), 51.8 (CH), 45.0 (CH₂), 39.4 (CH₂), 35.2 (CH₂),
31.4 (CH₂); MS (FAB+) found *m/z* 201.13558 (M+H), C₈H₁₆N₄O₂ M+H calc.
201.13514.

30 Example 15 (*S*)-2-Amino-4-[(piperidine-1-carboximidoyl)-amino]butanoic acid
Yield 32% (Method B). ¹H NMR (CD₃OD, 300 MHz): δ 4.08 (1H, t, *J* 6.5 Hz,
NCH), 3.54 (2H, t, *J* 7.1 Hz, CH₂), 3.48-3.44 (4H, m, 2 x CH₂), 2.31-2.13 (2H, m,
CH₂), 1.68-1.59 (6H, m, 3 x CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.7 (C),

157.5 (C), 51.9 (CH), 40.3 (CH₂), 31.3 (CH₂), 26.9 (CH₂), 25.2 (CH₂); MS (FAB+) found *m/z* 229.16582 (M+H), C₁₀H₂₀N₄O₂ M+H calc. 229.16644.

Example 16 (S)-2-Amino-4-[(pyrrolidine-1-carboximidoyl)-amino]butanoic acid

5 Yield 23% (Method B). ¹H NMR (CD₃OD, 300 MHz): δ 4.11 (1H, t, *J* 6.2 Hz, CH), 3.54 (2H, t, *J* 6.8 Hz, CH₂), 3.44 (4H, t, *J* 6.4 Hz, 2 x NCH₂ pyrr), 2.32-2.05 (2H, m, CH₂), 2.03 (4H, t, *J* 6.4 Hz, 2 x CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.4 (C), 155.0 (C), 51.5 (CH), 48.5 (CH₂), 39.4 (CH₂), 30.9 (CH₂), 26.2 (CH₂); MS (FAB+) found *m/z* 215.15070 (M+H), C₉H₁₈N₄O₂ M+H calc. 215.15079.

10

Example 17 (S)-2-Amino-4-[(morpholine-4-carboximidoyl)-amino]butanoic acid

Yield 30% (Method B). ¹H NMR (CD₃OD, 300 MHz): δ 4.09 (1H, t, *J* 6.2 Hz, CH), 3.74 (4H, t, *J* 4.9 Hz, 2 x CH₂ morph), 3.55 (2H, t, *J* 6.8 Hz, CH₂), 3.48 (4H, t, *J* 4.9 Hz, 2 x CH₂ morph), 2.30-2.19 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 66.9 (CH₂), 51.5 (CH), 47.6 (CH₂), 39.9 (CH₂), 30.8 (CH₂); MS (FAB+) found *m/z* 231.14593 (M+H), C₉H₁₈N₄O₃ M+H calc. 231.14571.

Example 18 (S)-4-(*N*⁷-methylguanidino)pyrrolidine-2-carboxylic acid

Yield 38% (Method B, from *N*-Boc-Hydroxyproline-OBu^t). ¹H NMR (CD₃OD, 300 MHz): δ 4.55-4.43 (2H, m, CH₂), 3.77-3.68 (1H, m, CH), 3.44-3.31 (1H, m, CH), 2.94-2.84 (1H, m, CH) 2.87 (3H, s, CH₃), 2.27-2.17 (1H, m, CH); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ; 170.4 (C), 158 (C), 59.6 (CH₂), 51.6 (CH₃) 50.6 (CH₂), 35.2 (CH₂), 28.6 (CH₃); MS (FAB+) found *m/z* 187.10587 (M+H), C₇H₁₄N₄O₂ M+H calc. 187.11950.

25

Example 19 (S)-4-(*N*⁷-(2-methoxyethyl)guanidino)pyrrolidine-2-carboxylic acid

Yield 26% (Method B, from *N*-Boc-Hydroxyproline-OBu^t). ¹H NMR (CD₃OD, 300 MHz): δ 4.55-4.45 (2H, m, CH₂), 3.75-3.71 (1H, m, CH), 3.54 (2H, t, *J* 4.8 Hz, CH₂), 3.44-3.40 (3H, m, CH₂ + CH), 3.38 (3H, s, OCH₃), 2.89-2.85 (1H, m, CH), 2.28-2.21 (1H, m, CH); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 170.4 (C), 159.0 (C), 72.1 (CH₂), 59.6 (CH), 59.2 (CH₃), 51.7 (CH), 50.7 (CH₂), 43.2 (CH₂), 35.2 (CH₂); MS (FAB+) found *m/z* 231.14592 (M+H), C₉H₁₈N₄O₃ M+H calc. 231.14571.

Example 20 (S)-4-N⁷-(2-methoxyethyl)guanidinobutanoic acid

Yield 59% (Method A, from 4-amino-*tert*-butylbutanoate). ¹H NMR (CD₃OD, 300 MHz): δ 3.53 (2H, t, *J* 4.7 Hz, CH₂), 3.39-3.34 (2H, m, CH₂), 3.36 (3H, s, OCH₃), 3.24 (2H, t, *J* 7.1 Hz, CH₂), 2.39 (2H, t, *J* 7.1 Hz, CH₂), 1.86 (2H, quintet, *J* 7.1 Hz, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 176.7 (C), 158.0 (C), 72.7 (CH₂), 59.2 (CH₃), 42.9 (CH₂), 41.9 (CH₂), 31.5 (CH₂), 25.2 (CH₂); MS (FAB+) found *m/z* 204.13477 (M+H), C₈H₁₇N₃O₃ M+H calc. 204.13481.

Example 21 N^G-(2-methoxyethyl)-L-arginine

10 Yield 44% (Method A, from Boc-Orn-OBu^t). ¹H NMR (CD₃OD, 300 MHz): δ 4.03 (1H, t, *J* 6.0 Hz, CH), 3.53 (2H, t, *J* 5.0 Hz, CH₂), 3.39 (2H, t, *J* 5.0 Hz, CH₂), 3.37 (3H, s, OCH₃), 3.30 (2H, m, CH₂), 2.03-1.94 (2H, m, CH₂), 1.83-1.72 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 171.5 (C), 158.2 (C), 72.2 (CH₂), 59.2 (CH₃), 53.5 (CH), 42.9 (CH₂), 41.9 (CH₂), 28.7 (CH₂), 25.8 (CH₂); MS (FAB+) found *m/z* 15 233.16097 (M+H), C₉H₂₀N₄O₃ M+H calc. 233.16136.

General method for preparation of esters

To a solution of the acid (typically 0.5 mmol) in the appropriate alcohol (2 mL) at 20 0°C was added thionyl chloride (1.1 equivalents), with stirring. The solution was stirred for 30 minutes at 0°C and then heated under reflux for 1 hour and stirred at room temperature overnight. Solvent was then removed *in vacuo* to give the ester as a white or yellow solid.

Example 22 (S)-2-Amino-4-(N⁷-(2-methoxyethyl)guanidino)butanoic acid methyl 25 ester
Yield 83%. ¹H NMR (CD₃OD, 300 MHz): δ 4.15 (1H, t, *J* 6.5 Hz, CH), 3.86 (3H, s, OCH₃), 3.54 (2H, t, *J* 4.9 Hz, CH₂), 3.47 (2H, t, *J* 7.1 Hz, CH₂), 3.42 (2H, t, *J* 6.0 Hz, CH₂), 3.38 (3H, s, OCH₃), 2.29-2.11 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 170.3 (C), 157.5 (C), 72.1 (CH₂), 59.2 (CH₃), 54.0 (CH₃), 51.5 (CH), 43.0 30 (CH₂), 38.8 (CH₂), 30.9 (CH₂); MS (FAB+) found *m/z* 233.16097 (M+H), C₉H₂₀N₄O₃ M+H calc. 233.16136.

Example 23 (S)-2-Amino-4-(N⁷-(2-methoxyethyl)guanidino)butanoic acid ethyl ester

Yield 84%. ¹H NMR (CD₃OD, 300 MHz): δ 4.33 (2H, q, *J* 7.1 Hz, CH₂), 4.13 (1H, t, *J* 6.2 Hz, CH), 3.55-3.39 (6H, m, 3 x CH₂) 3.38 (3H, s, OCH₃), 2.28-2.20 (2H, m, 5 CH₂), 1.34 (3H, t, *J* 7.1 Hz, CH₃); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 169.9 (C), 158.3 (C), 72.1 (CH₂), 64.1 (CH₂), 59.2 (CH₃), 51.5 (CH), 43.1 (CH₂), 38.9 (CH₂), 30.9 (CH₂), 14.4 (CH₃); MS (FAB+) found *m/z* 247.17723 (M+H), C₁₀H₂₂N₄O₃ M+H calc. 247.17701.

10 Example 24 (S)-2-Amino-4-(N⁷-(2-methoxyethyl)guanidino)butanoic acid propyl ester

Yield 80%. ¹H NMR (CD₃OD, 300 MHz): δ 4.25-4.15 (3H, m, CH, CH₂), 3.56-3.39 (6H, m, 3 x CH₂), 3.38 (3H, s, OCH₃), 2.27-2.15 (2H, m, CH₂), 1.73 (2H, m, CH₂), 0.98 (3H, t, *J* 7.1 Hz, CH₃); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 170.0 (C), 158.1 (C), 72.1 (CH₂), 69.5 (CH₂), 59.3 (CH₃), 51.6 (CH), 43.1 (CH₂), 38.9 (CH₂), 30.9 (CH₂), 22.9 (CH₂), 10.7 (CH₃); MS (FAB+) found *m/z* 261.19258 (M+H), C₁₁H₂₄N₄O₃ M+H calc. 261.19266.

20 Example 25 (S)-2-Amino-4-(N⁷-(2-methoxyethyl)guanidino)butanoic acid butyl ester

Yield 60%. ¹H NMR (CD₃OD, 300 MHz): δ ; ¹³C NMR (75 MHz, CD₃OD, 27°C) δ; MS (FAB+) found *m/z* 275.20818 (M+H), C₁₂H₂₆N₄O₃ M+H calc. 275.20831.

25 Example 26 (S)-2-Amino-4-(N⁷-(2-methoxyethyl)guanidino)butanoic acid benzyl ester

Yield 79%. ¹H NMR (CD₃OD, 300 MHz): δ 7.45-7.27 (5H, m, ArH), 5.23 (2H, s, PhCH₂O), 4.20 (1H, t, *J* 6.2 Hz, CH), 3.59-3.42 (6H, m, 3 x CH₂), 3.30 (3H, s, OCH₃), 2.27-2.11 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 169.9 (C), 158.0 (C), 136.3 (C), 129.9 (CH), 129.4 (CH), 129.0 (CH), 72.1 (CH₂), 69.5 (CH₂), 30 59.2 (CH₃), 51.6 (CH), 43.0 (CH₂), 38.9 (CH₂), 30.9 (CH₂); MS (FAB+) found *m/z* 309.19466 (M+H), C₁₅H₂₄N₄O₃ M+H calc. 309.19266.

Example 27 (S)-2-Amino-4-(N⁷-(2-methoxyethyl)guanidino)butanoic acid isopropyl ester

Yield 72%. ¹H NMR (CD₃OD, 300 MHz): δ 5.11 (1H, septet, *J* 6.2 Hz, (Me)₂CH), 4.12 (1H, t, *J* 6.7 Hz, CH), 3.56-3.42 (6H; m, 3 x CH₂), 3.38 (3H, s, OCH₃), 2.29-5 2.04 (2H, m, CH₂), 1.33 (6H, d, *J* 6.2 Hz, (CH₃)₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 169.5 (C), 158.1 (C), 72.5 (CH₂), 72.1 (CH) 59.3 (CH₃), 51.6 (CH), 43.1 (CH₂), 39.0 (CH₂), 31.0 (CH₂), 21.9 (CH₃); MS (FAB+) found *m/z* 261.19378 (M+H), C₁₁H₂₄N₄O₃ M+H calc. 261.19266.

10 Example 28 N^G-(2-Methoxyethyl)-L-arginine methyl ester

Yield 80%. ¹H NMR (CD₃OD, 300 MHz): δ 4.12 (1H, t, *J* 6.0 Hz, CH), 3.85 (3H, s, OCH₃), 3.53 (2H, t, *J* 4.7 Hz, CH₂), 3.40 (2H, t, *J* 4.7 Hz, CH₂), 3.38 (3H, s, OCH₃), 3.31-3.27 (2H, m, CH₂), 2.10-1.92 (2H, m, CH₂), 1.84-1.69 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 170.7 (C), 158.1 (C), 72.2 (CH₂), 59.3 (CH₃), 53.9 15 (CH₃), 53.6 (CH), 43.0 (CH₂), 41.9 (CH₂), 28.7 (CH₂), 25.8 (CH₂); MS (FAB+) found *m/z* 247.17725 (M+H), C₁₀H₂₂N₄O₃ M+H calc. 247.17701.

General method for preparation of amides

Amine derivatives of the guanidine amino acids can be synthesised using a modified 20 version of Method A. For example, 3-N⁷-(2-methoxyethyl)guanidinopropylamine can be synthesised from Method A using Boc-1,3-diaminopropane.

Preparative Example 1 3-N⁷-(2-Methoxyethyl)guanidinopropylamine

Yield 63% (Method A, from Boc-1,3-diaminopropane). ¹H NMR (CD₃OD, 300 25 MHz): δ 3.53 (2H, t, *J* 5.0 Hz, CH₂), 3.40 (2H, t, *J* 5.0 Hz, CH₂), 3.38 (3H, s, OCH₃), 3.34 (2H, t, *J* 7.0 Hz, CH₂), 3.02 (2H, t, *J* 7.6 Hz, CH₂), 1.96 (2H, tt, *J* 7.0, 7.6 Hz, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 158.2 (C), 72.2 (CH₂), 59.2 (CH₃), 43.0 (CH₂), 39.7 (CH₂), 38.2 (CH₂), 28.0 (CH₂); MS (FAB+) found *m/z* 175.15591 (M+H), C₇H₁₈N₄O M+H calc. 175.15588.

30

Isobutyl chloroformate (1 mmol) and N-methylmorpholine (1 mmol) were added to L-Boc-ornithine(Fmoc) (1 mmol) in chloroform (5 mL) at -10°C, with stirring. The mixture was stirred at -10°C for 15 minutes, and then the required amine (1 mmol)

was added and the mixture stirred for a further 3 hours with gradual warming to room temperature. The solvent was removed *in vacuo* and the residue subjected to flash column chromatography (1:1 cyclohexane: ethyl acetate) to give the amide as a white powder.

5 The amide (0.5 mmol) was stirred with piperidine (5 equivalents) in chloroform (3 mL) with monitoring by tlc until the starting material was consumed. Solvent and excess piperidine were removed *in vacuo* and the entire crude product was used in the next step.

The *N*-alkyl substituted pyrazolecarboxamidine (0.5 mmol), and the crude Boc-
10 protected amide (~0.5 mmol) were stirred in acetonitrile (5 mL) for 48 hours. After removal of solvent, the residue was subjected to flash column chromatography to give the Boc-protected guanidino- amino amide.

The Boc-protected amide was stirred in excess HCl/dioxane for 72 hours, then the solvent removed *in vacuo* to give the product as a white solid.

15

Example 29 (S)-2-Amino-5-(*N*⁷-(2-methoxyethyl)guanidino)pentanoic acid methylamide

Yield 36%. ¹H NMR (CD₃OD, 300 MHz): δ 3.92 (1H, t, *J* 6.5 Hz, CH), 3.53 (2H, t, *J* 4.7 Hz, CH₂), 3.40 (2H, t, *J* 5.0 Hz, CH₂), 3.37 (3H, s, NCH₃), 3.32-3.26 (2H, m, CH₂), 2.79 (3H, s, CH₃), 1.96-1.89 (2H, m, CH₂), 1.74-1.66 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 170.3 (C), 158.1 (C), 72.2 (CH₂), 59.3 (CH₃), 54.1 (CH), 43.0 (CH₂), 42.0 (CH₂), 29.7 (CH₂), 26.4 (CH₃), 25.6 (CH₂); MS (FAB+) found *m/z* 246.19305 (M+H), C₁₀H₂₃N₅O₂ M+H calc. 246.19299.

25 Example 30 (S)-2-Amino-4-(*N*⁷-(2-methoxyethyl)guanidino)butanoic acid benzylamide

Yield 33%. ¹H NMR (CD₃OD, 300 MHz): 7.33-7.22 (5H, m, ArH), 4.43 (2H, d, *J* 5.5 Hz, CH₂), 4.09 (1H, t, *J* 6.7 Hz, CH), 3.53-3.40 (4H, m, 2 x CH₂), 3.36 (3H, s, OMe), 2.24-2.09 (2H, m, CH₂); ¹³C NMR (75 MHz, CD₃OD, 27°C) δ 169.9 (C), 158.1 (C), 139.3 (C), 129.6 (Ar CH), 128.9 (Ar CH), 128.5 (Ar CH), 72.1 (CH₂), 59.3 (CH₃), 52.3 (CH), 44.4 (CH₂), 43.0 (CH₂), 38.8 (CH₂), 32.1 (CH₂); MS (FAB+) found *m/z* 308.21039 (M+H), C₁₅H₂₅N₅O₂ M+H calc. 308.20864.

Example 31 (S)-N^G-((Tetrahydrofuran-2-yl)methyl)-arginine

tert-Butyl (2S)-2-Boc-amino-5-(N,N'-bis-Boc-N^G-((tetrahydrofuran-2-yl)methyl)guanidino) pentanoate (0.30 g, 0.5 mmol, 1 eq.), was dissolved in dioxane (3 mL). 4M HCl/dioxane (5 mL, 20 mmol, 40 eq) was added and the solution stirred

5 for 48 hours. All excess reagents and solvents were removed *in vacuo*. The compound was purified by freeze-drying to yield a sticky, hygroscopic solid (74.2 mg, 67.3%). ¹H NMR (d₃-MeOD): δ 1.64 (m, 2H); 1.79 (m, 2H); 1.93 (m, 2H); 1.98 (m, 2H); 3.44 (m, 2H); 3.58 (m, 1H); 3.88 (m, 2H); 3.93 (m, 1H); 4.00 (m, 2H). ¹³C NMR (d₃-MeOD): δ 25.8 (CH₂); 26.7 (CH₂); 28.8 (CH₂); 29.4 (CH₂); 41.9 (CH₂); 10 48.2 (CH₂); 54.1 (CH); 69.4 (CH). N.B. Values for C=N and C=O are too weak to be determined.

Example 32 (2S)-2-Amino-4-N^G-((tetrahydrofuran-2-yl)methyl)guanidinobutanoic acid

15 Synthesised using an adaptation of the procedure of Example 31. Yield 56.3%. ¹H NMR (d₃-MeOD): δ 1.63 (m, 2H); 1.91 (m, 2H); 2.21 (m, 2H); 3.51 (t, *J* = 6 Hz, 2H); 3.60 (t, *J* = 6.3 Hz, 1H); 3.75 (m, 2H); 3.88 (m, 1H); 4.80 (t, *J* = 7.5 Hz, 2H). ¹³C NMR (d₃-MeOD): δ 26.7 (CH₂); 29.9 (CH₂); 31.0 (CH₂); 39.0 (CH₂); 46.8 (CH₂); 51.4 (CH); 69.4 (CH₂); 70.7 (CH); 158.3 (C); 171.2 (C).

20

Example 33 2-Methoxybenzyl (S)-2-amino-4-[(N^G-(2-methoxyethyl))guanidino] butanoate

(2S)-2-Amino-4-N^G-(2-methoxyethyl)-guanidinobutanoic acid (100 mg, 0.34 mmol, 1 eq.) was dissolved in excess 2-methoxybenzyl alcohol (2 mL). The solution was cooled to 0°C and thionyl chloride (0.2 mL, 2.75 mmol, 8 eq.) was added slowly. Once calm, the solution was heated to 70°C for 48 hours. The product was purified by trituration (EtOAc) and freeze-dried to yield a sticky, hygroscopic solid (56.1 mg, 48%). ¹H NMR (d₃-MeOD): δ 2.20 (m, 2H); 3.30 (m, 2H); 3.41 (m, 3H); 3.54 (m, 5H); 3.81 (m, 2H); 4.05 (t, *J* = 6.3 Hz, 2H); 4.58 (m, 1H); 6.67 (m, 2H); 7.32 (m, 1H); 7.36 (m, 1H). ¹³C NMR (d₃-MeOD): δ 31.0 (CH₂); 39.1 (CH₂); 43.0 (CH₂); 51.8 (CH); 55.7 (CH₃); 59.2 (CH₃); 60.4 (CH₂); 72.1 (CH₂); 111.2; 112.3; 121.4; 129.0; 30 129.6; 132.1 (Aromatic Peaks); 158.2 (C); 171.4 (C). LCMS: 339.4 (M + H).

Example 34 4-Methoxybenzyl (S)-2-amino-4-[(*N*⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 80%. ¹H NMR (d₃-MeOD): δ 2.22 (m, 2H); 3.30 (m, 2H); 3.41 (m, 2H); 3.52 (m, 5H); 3.77 (m, 5H); 4.43 (m, 1H); 6.96 (m, 2H); 7.25 (m, 2H). ¹³C NMR (d₃-MeOD): δ 31.0 (CH₂); 39.0 (CH₂); 43.0 (CH₂); 43.8 (CH₂); 51.4 (CH); 55.7 (CH₃); 59.3 (CH₃); 72.2 (CH₂); 114.8; 115.3; 115.6; 129.9; 130.6; 133.0 (Aromatic Peaks); 158.3 (C); 171.2 (C). LCMS: 339.3 (M + H).

10 Example 35 2-Fluorobenzyl (S)-2-amino-4-[(*N*⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 42.7%. ¹H NMR (d₃-MeOD): δ 2.20 (m, 2H); 3.30 (m, 2H); 3.39 (m, 5H); 3.52 (m, 2H); 4.06 (t, J = 7.4 Hz, 1H); 4.20 (t, J = 7.0 Hz, 1H); 4.66 (s, 1H); 7.19 (m, 2H); 7.47 (m, 2H). ¹³C NMR (d₃-MeOD): δ 31.0 (CH₂); 39.0 (CH₂); 43.0 (CH₂); 51.5 (CH); 58.7; 59.2 (CH₃); 63.2 (CH₂); 72.1 (CH₂); 116.7; 117.1; 125.7; 130.5; 131.0; 133.2 (Aromatic Peaks); 158.3 (C); 171.9 (C). LCMS: 327.3 (M + H).

20 Example 36 3-Trifluoromethyl-4-fluorobenzyl (S)-2-amino-4-[(*N*⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 28.3%. ¹H NMR (d₃-MeOD): δ 2.05 (m, 2H); 3.31 (m, 2H); 3.40 (m, 5H); 3.53 (m, 2H); 3.61 (m, 2H); 3.63 (m, 1H); 7.32 (m, 1H); 7.67 (m, 2H). ¹³C NMR (d₃-MeOD): δ 22.8; 32.2 (CH₂); 39.8 (CH₂); 42.3 (CH₂); 53.1 (CH); 59.2 (CH₃); 72.1 (CH₂); 158.1 (C). 25 N.B. The integration values for the NMR revealed the presence of 50% starting material, indicating the reaction had not gone to completion. LCMS: 395.4 (M + H).

Example 37 3-Phenoxybenzyl (S)-2-amino-4-[(*N*⁷-(2-methoxyethyl))guanidino] butanoate

30 Synthesised using an adaptation of the procedure of Example 33. Yield 52.8%. ¹H NMR (d₃-MeOD): δ 2.20 (m, 2H); 3.30 (m, 1H); 3.35 (m, 1H); 3.42 (m, 2H); 3.52 (m, 5H); 4.03 (t, J = 7 Hz, 1H); 4.21 (t, J = 7 Hz, 1H); 4.57 (s, 1H); 6.99 (m, 3H); 7.13 (m, 3H); 7.37 (m, 3H). ¹³C NMR (d₃-MeOD): δ 31.1 (CH₂); 39.1 (CH₂); 43.0

(CH₂); 51.6 (CH); 59.2 (CH₃); 69.0 (CH₂); 72.1 (CH₂); 118.1; 118.5; 119.8; 120.1; 122.7; 124.5; 124.8; 130.8; 131.0; 131.3; 138.3; 145.1 (Aromatic Peaks); 158.7 (C); 169.8 (C). LCMS: 401.3 (M + H).

5 Example 38 3-Methylbenzyl (S)-2-amino-4-[(N⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 64.1%. ¹H NMR (d₃-MeOD): δ 2.22 (m, 2H); 2.32 (d, *J* = 6.4 Hz, 3H); 3.30 (b, 2H); 3.35 (s, 2H); 3.38 (b, 2H); 3.45 (b, 2H); 3.51 (b, 3H); 4.55 (s, 1H); 7.13 (m, 4H). ¹³C NMR (d₃-MeOD): δ 21.5 (CH₃); 31.0 (CH₂); 38.9 (CH₂); 43.0 (CH₂); 51.6 (CH); 59.2 (CH₃); 69.6 (CH₂); 72.1 (CH₂); 125.1; 126.9; 128.9; 130.5; 136.2; 139.0 (Aromatic Peaks); 142.6 (C); 170.0 (C). LCMS: 323.3 (M + H).

15 Example 39 3-Trifluoromethylbenzyl (S)-2-amino-4-[(N⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 11.0%. ¹H NMR (d₃-MeOD): δ 2.20 (m, 2H); 3.38 (m, 7H); 3.53 (t, *J* = 4.7, 2H); 3.60 (t, *J* = 7.1 Hz, 2H); 4.67 (s, 1H); 7.70 (m, 4H). The NMR integrations indicate the presence of 50% starting material and hence, it can be concluded that the reaction did not go to completion. LCMS: 377.4 (M + H).

20 Example 40 3-Fluorobenzyl (S)-2-amino-4-[(N⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 60.7%. ¹H NMR (d₃-MeOD): δ 2.20 (m, 2H); 3.30 (m, 2H); 3.42 (m, 2H); 3.54 (m, 5H); 4.06 (t, *J* = 6.5 Hz, 1H); 4.24 (t, *J* = 6.9 Hz, 1H); 6.96 (m, 1H); 7.12 (m, 1H); 7.31 (m, 2H). ¹³C NMR (d₃-MeOD): δ 30.9 (CH₂); 38.9 (CH₂); 43.0 (CH₂); 51.6 (CH); 59.2 (CH); 64.4 (CH₂); 72.1 (CH₂); 114.5; 116.5; 123.4; 125.5; 131.0; 138.2 (Aromatic peaks); 169.8 (C); 171.3 (C). LCMS: 327.3 (M + H).

30

Example 41 2-Methoxyethyl (S)-2-amino-4-[(N⁷-(2-methoxyethyl))guanidino] butanoate

Synthesised using an adaptation of the procedure of Example 33. Yield 66.7%. ¹H NMR (d₃-MeOD): δ 2.20 (m, 2H); 3.38 (s, 6H); 3.41 (t, J = 5.1 Hz, 2H); 3.47 (t, J = 5.1 Hz, 2H); 3.52 (t, J = 5.1 Hz, 2H); 3.67 (t, J = 5.1 Hz, 2H); 4.18 (t, J = 7.5 Hz, 1H); 4.41 (m, 2H). ¹³C NMR (d₃-MeOD): δ 31.0 (CH₂); 43.0 (CH₂); 51.5 (CH); 59.1 (CH₃); 59.2 (CH₃); 64.3 (CH₂); 66.4 (CH₂); 71.0 (CH₂); 72.1 (CH₂); 158.1 (C); 169.9 (C). LCMS: 277.4 (M + H).

Results

Biological Screening

10

DDAH activity assay

Rat kidney was homogenized in PBS containing PMSF 1mM, leupeptin 5μg/ml, pepstatin 5μg/ml, chymostatin 5μg/ml, and the lysate was centrifuged (4000 rpm, 20 min then supernatant further centrifuged for 40000rpm, 30min, 4°C). The 15 supernatants were analyzed for DDAH activity; [¹⁴C]L-NMMA (1 μmol/L) was added to the cell lysates (final volume 100 μl) and incubated for 1 h at 37°C. Dowex resin (0.5 ml) at neutral pH was immediately added to samples to terminate the reaction and bind unmetabolized L-NMMA. [¹⁴C]Citrulline formation was determined by scintillation counting (Hewlett Packard). The results are shown in 20 Tables 1 and 1A.

Cell assays

RAW cells were grown to 70% confluence in T₁₅₀ flasks then stimulated with LPS (5 μg/ml), TNFα (10 ng/ml) and IFNγ (100 U/ml) for 24h, cells were scraped in cold 25 100 mM Tris (pH 7.6), 1 mM DTT, PMSF 1mM, leupeptin 5μg/ml, pepstatin 5μg/ml, chymostatin 5μg/ml and CaCl₂ 200 μM. Cells were lysed by three cycles of freeze / thaw and cleared by microcentrifugation. Protein concentration in the supernatants was determined by Biorad protein assay prior to use in the NOS assay.

30 **Cell viability assay**

RAW cells were treated with DDAH inhibitor compound (0.5 mM) in complete media for 24h; cells were subsequently assessed for viability by MTT assay. Cell

culture media was removed from cells and replaced with 200 μ l complete media containing 3-[4,5-Dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT 0.2 mg /ml) for 30 min at 37°C. MTT containing media was removed and 100ul DMSO was added to solubilize cells; after shaking, plates were read at OD $_{550\text{nm}}$.

5 There was no significant difference between untreated cells and those exposed to DDAH inhibitors. The results of the assay are shown in Table 2.

iNOS activity Greiss assay

RAW cells were stimulated for 24 h with LPS / IFN and TNF, in the presence of the 10 DDAH inhibitor compounds (0.5 mM). Levels of NO_x secreted into the media by the cells were determined by Griess assay: Griess reagent [1:1 sulfanilamide (10%) in ortho-phosphoric acid (50%) and N-(1-napthyl)ethylenediamine dihydrochloride (1%)] was added as 100 μ l volume to 100 μ l cell culture media and the plate was read at 550 nm and analysed in relation to sodium nitrite and nitrate standards.

15 There was no significant difference between untreated cells and those exposed to DDAH inhibitors (n=10). The results of the assay are shown in Table 3.

eNOS assay

Compounds to be tested were added to NOS assay buffer (HEPES 50 mM pH 7.2, 20 FAD 5 μ M, FMN 5 μ M, BH4 10 μ M, NADPH 1mM, DTT 0.5 mM, CaCl₂ 1mM, MgCl₂ 1mM, calmodulin 50 nM, [¹⁴C] arginine 100 μ M) in a final volume of 100 μ l and the reaction was initiated by addition of eNOS (eNOS 3.3. U / reaction, Alexis Biochemicals). Reactions were incubated for 1 h at 37°C and the reaction was terminated by the addition of Dowex pH 5.5 (0.85 ml) to each tube. Tubes were 25 vortexed and microfuged <12000g, 5 min. 100 μ l of the Dowex supernatant was added to 5 ml scintillation cocktail and was counted in a scintillation counter (Hewlett Packard). The compounds were tested at 1mM and 0.1mM concentrations. Activity is expressed as % of control where control contained no inhibitor, n=4. The DDAH inhibitors of the present invention tested had no significant effect on eNOS 30 activity. The results are shown in Table 4.

In vivo studies

Male Wistar rats (250-300g; Charles River, Margate, Kent, UK) were anaesthetised using Isoflurane. The jugular vein was cannulated. Animals were injected intravenously with a bolus dose of inhibitor (60mg/kg) or saline. Anaesthesia was 5 maintained for 4 hours at the end of which blood was obtained via cardiac puncture for the determination of serum dimethylarginine levels. [All animals received care in compliance with The British Home Office Regulations and Principles of Laboratory Animal Care. (Project licenses PPL/5344, PPL/4824, PPL 70/5580)]

10 The results of the study are shown in Table 5.

As can be seen from Table 5, ADMA levels were higher in the treated animals, which is consistent with the inhibition of DDAH. SDMA is not a substrate for DDAH and the levels of SDMA would be expected to be the same of the treated and 15 control animals, and this was the case. Since there is a great deal of variability of the "normal" ADMA and SDMA level, it is useful to measure the ratio of ADMA and SDMA in each animal to provide a reliable indicator of the relative changes of these two substances.

20 **Inhibition of DDAH reduces peritoneal inflammation**

The effects of a DDAH inhibitor on zymosan-induced peritonitis were tested in two different strains of mice. C57/Blk6 or TO mice, male 8-10 weeks were placed in a heat box for 10 mins at 38°C, and treated with 60mg/kg of SR291 [N^G -(2-methoxyethyl)-L-arginine methyl ester] i.v. 30 mins later they were given 1mg of 25 zymosan i.p. A second dose of SR 291 (60mg/kg) was administered 2h later and after 4 h mice were anaesthetised with isofluorane and blood withdrawn by cardiac puncture. Mice were sacrificed and had 2mls of ice cold PBS injected into their peritoneal cavity, the cavity was then gently massaged and cut open, the peritoneal washout was harvested and put on ice. Blood was spun down at 400g, 10 mins at 30 4°C. The cells were spun at 4000rpm for 5 mins at 4°C. Plasma was removed from the blood and frozen at -80°C. The exudates from the washout were removed and frozen at -80°C, the cell pellet was resuspended in 1ml ice cold PBS and were counted on a haemocytometer. The cells were spun as before, the supernatant

discarded and the cells frozen at -80°C. Polymorphonuclear (PMN) cell infiltrate into the peritoneal cavity was used to measure inflammation. In three separate experiments each involving 8-12 mice (half treated with saline half with active drug) SR291 significantly reduced PMN infiltrate. The results of the study are shown in 5 Tables 5 to 10. Table 6 shows the results obtained for zymosan peritonitis (C57's male 8-10 weeks) 60mg/kg SR 291 iv -30min taken off at 4 hrs. Table 7 shows the cumulative results of the dataset that appears in Table 6. Table 8 shows a combination of two datasets for the C57 mice. Table 9 shows the results obtained for zymosan peritonitis (TO mice male 8-10 weeks) 60mg/kg SR 291 iv -30min, +2hrs. 10 Taken off at 4 hrs. Table 10 shows the cumulative results of the dataset that appears in Table 9. Table 11 shows the results of a second study obtained for zymosan peritonitis (TO mice male 8-10 weeks) 60mg/kg SR 291 iv -30min, +2hrs. Taken off at 4 hrs. Table 12 shows the cumulative results of the dataset that appears in Table 11. Table 13 shows the cumulative results of the two datasets that appear in Tables 9 15 and 11.

DDAH Inhibition restores vascular reactivity and maintains BP in endotoxaemia

Effects of two compounds SR 291 [N^G -(2-methoxyethyl)-L-arginine methyl ester] 20 and SR 257 [N^G -(2-methoxyethyl)-L-arginine]) have been tested in a series of models of endotoxin-induced vascular dysfunction. Treatment of rat aortic vascular rings in static culture with endotoxin (10 μ g/ml) led to an increase in generation of nitric oxide (as measured by nitrite and nitrate). This increase in NO was prevented by treatment with 500 μ M SR257 or SR291. The results of this study are shown in 25 Figure 1.

Incubation of rat aortic rings with endotoxin (0.3 μ g/ml) led to the development of hyporeactivity to vasoconstrictors. This effect was reversed dose dependently by inhibition of DDAH using 100 μ M and 200 μ M SR291 or SR257 whilst the inactive 30 D-isomers of each compound showed no effect. This is shown in Figures 2a and 2b.

Injection of endotoxin (40mg/kg) into rats caused a progressive fall in systemic BP. The decline was halted by infusion of 30mg/kg/hr SR291. This is shown in Figure 3.

These data show that inhibition of DDAH may be used to prevent vascular collapse in sepsis.

Table 1

Compound Name	Inhibition at 1mM, %	IC ₅₀ , μ M
(S)-2-Amino-4-(N'-isopropylguanidino)butanoic acid	38	
(S)-2-Amino-4-(N'-propylguanidino)butanoic acid	41	
(S)-2-Amino-4-(N'-2,2-dimethylpropyl)guanidino)butanoic acid	19	
(S)-2-Amino-4-(N'-octylguanidino)butanoic acid	27	
(S)-2-Amino-4-(N'-cyclohexylguanidino)butanoic acid	51	1470
(S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid	90	189
(S)-2-Amino-4-(N'-(2-phenoxyethyl)guanidino)butanoic acid	38	
(S)-2-Amino-4-(N'-(2-isopropoxyethyl)guanidino)butanoic acid	97	301
(S)-2-Amino-4-(N'-(2-methylthioethyl)guanidino)butanoic acid	19	
(S)-2-Amino-4-(N'-phenylguanidino)butanoic acid	65	1000
(S)-2-Amino-4-(N'-benzylguanidino)butanoic acid	22	
(S)-2-Amino-4-(N'-(2-thiophenemethyl)guanidino)butanoic acid	34	
(S)-2-Amino-4-(N'-allylguanidino)butanoic acid	70	998
(S)-2-Amino-4-(N'-(2'-dimethylaminoethyl)guanidino)butanoic acid	25	
(S)-2-Amino-4-(N'-cyclopropylguanidino)butanoic acid	65	1020
(S)-2-Amino-4-[(piperidine-1-carboximidoyl)-amino]butanoic acid	74	264
(S)-2-Amino-4-[(morpholine-4-carboximidoyl)-amino]butanoic acid	60	686
(S)-2-Amino-4-[(pyrrolidine-1-carboximidoyl)-amino]butanoic acid	79	500
N ^G -(2-methoxyethyl)-L-arginine	100	22
N ^G -(2-methoxyethyl)-L-arginine methyl ester	83	20
(S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid methyl ester	75	96
(S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid ethyl ester	44	159
(S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid propyl ester	64	111
(S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid butyl ester	40	113
(S)-2-Amino-4-(N'-(2-methoxyethyl)guanidino)butanoic acid benzyl ester	96	27

Table 1A

Compound Name	Inhibition at 100 μ M (%)
(S)- N^G -((Tetrahydrofuran-2-yl)methyl)-arginine	30
(2S)-2-Amino-4- N^G -((tetrahydrofuran-2-yl)methyl)guanidinobutanoic acid	10
2-Methoxybenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	47.4
4-Methoxybenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	22.1
2-Fluorobenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino]butanoate	39.8
3-Trifluoromethyl-4-fluorobenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	50.6
3-Phenoxybenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	42.7
3-Methylbenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	48.7
3-Trifluoromethylbenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	45.3
3-Fluorobenzyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	33.2
2-Methoxyethyl (S)-2-amino-4-[($N^$ -(2-methoxyethyl))guanidino] butanoate	59

Table 2

	OD units drug \pm sd	Number of experiments
Control	2.54 \pm 0.26	3
(S)-2-Amino-4-($N^$ -(2 methoxyethyl)guanidino) butanoic acid methyl ester	2.33 \pm 0.19	3
(S)-2-Amino-4-($N^$ -(2-methoxyethyl)guanidino) butanoic acid propyl ester	2.40 \pm 0.21	3
(S)-2-Amino-4-($N^$ -(2-methoxyethyl)guanidino) butanoic acid benzyl ester	2.40 \pm 0.17	3
N^G -(2-methoxyethyl)-L-arginine	2.15 \pm 0.13	3
N^G -(2-methoxyethyl)-L-arginine methyl ester	2.41 \pm 0.10	3
L- N -5-(1-iminoethyl)ornithine [NIO]	1.89 \pm 0.41	3
ADMA	2.22 \pm 0.10	3
L-NMMA	2.22 \pm 0.08	3

Table 3

	OD units drug ± sd	Number of experiments
Control	0.367 ± 0.021	5
(S)-2-Amino-4-(N ^G -(2 methoxyethyl)guanidino) butanoic acid methyl ester	0.329 ± 0.013	5
(S)-2-Amino-4-(N ^G -(2-methoxyethyl)guanidino) butanoic acid propyl ester	0.351 ± 0.013	5
(S)-2-Amino-4-(N ^G -(2-methoxyethyl)guanidino) butanoic acid benzyl ester	0.327 ± 0.026	5
N ^G -(2-methoxyethyl)-L-arginine	0.326 ± 0.014	5
N ^G -(2-methoxyethyl)-L-arginine methyl ester	0.325 ± 0.009	5
NIO	0.103 ± 0.002	5
ADMA	0.195 ± 0.014	5
L-NMMA	0.121 ± 0.003	5

Table 4

	% of control ± sd [number of experiments]	
	1mM	0.1 mM
Control	100.0 ± 14.95 [6]	100.0 ± 14.95 [6]
(S)-2-Amino-4-(N ^G -(2 methoxyethyl)guanidino)butanoic acid methyl ester	84.05 ± 8.09 [4]	87.23 ± 18.05 [2]
(S)-2-Amino-4-(N ^G -(2-methoxyethyl)guanidino)butanoic acid propyl ester	93.97 ± 12.93 [4]	113.5 ± 7.93 [2]
(S)-2-Amino-4-(N ^G -(2-methoxyethyl)guanidino)butanoic acid benzyl ester	82.88 ± 10.08 [4]	101.2 ± 38.07 [2]
N ^G -(2-methoxyethyl)-L-arginine	84.81 ± 22.74 [4]	96.84 ± 30.49 [2]
N ^G -(2-methoxyethyl)-L-arginine methyl ester	112.5 ± 27.47 [4]	120.8 ± 17.80 [2]
NIO	6.25 ± 7.52 [5]	4.14 ± 8.74 [3]
L-NMMA	16.17 ± 15.78 [4]	

5 Table 5

	ADMA:SDMA ± sd	Number of experiments
Control	1.81 ± 0.23	9
N ^G -(2-methoxyethyl)-L-arginine	3.50 ± 0.48	8
N ^G -(2-methoxyethyl)-L-arginine methyl ester	2.73 ± 0.44	4

Table 6

drug	cells million /ml				
	1	2	3	4	5
Saline	14	12	7	6.5	6
SR291	3	4	4.5	6.5	9

Table 7

	cells million/ml	
	mean	SEM
Saline	9.1	1.631
SR291	5.4	1.0654

Table 8

	cells 10^6 /ml	
	mean	sem
saline	6.7455	0.9746
SR291	4.2	0.6097

5

Table 9

animal	cells 10^6 /ml					
	1	2	3	4	5	6
Saline	4.2	5	4.9	4.8	4.75	5.05
SR291	3.05	3.35	3.2	1.4	3.9	4.3

Table 10

	cells 10^6 /ml	
	mean	sem
Saline	4.783333	0.12561
SR291	3.2	0.407636

10

Table 11

	cells 10^6 /ml					
	1	2	3	4	5	6
Saline	6	6.15	6.5	6.3	5.9	5.9
SR291	4.3	4.6	4.85	5.2	4.3	3.9

Table 12

	cells 10^6 /ml	
	mean	sem
Saline	6.125	0.098107
SR291	4.525	0.187861


15

Table 13

	cells 10^6 /ml	
	mean	sem
saline	5.454167	0.216065
SR291	3.8625	0.292723

The claims defining the invention are as follows:

1. Use of a compound of formula (I), or a pharmaceutically acceptable salt thereof in the manufacture of a medicament, for use in the treatment of a disease whose pathology is affected by DDAH



wherein:

10 - either (a) R¹ is C₃-C₈ alkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl, phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A or -L-Y-A, wherein:

- L is C₁-C₆ alkyl;
- L' is C₁-C₆ alkyl;

15 - A is phenyl, C₃-C₆ carbocyclyl, 5- to 6-membered heterocyclyl or 5- to 6-membered heteroaryl;

- Het is -O-, -S- or -NR'-, wherein R' is hydrogen or C₁-C₆ alkyl; and

- Y is -CO-, -SO-, -SO₂-, -CO-O-, -CO-NR'-, -O-CO- or -NR'-CO-, wherein R' is as defined above;

20 - R² is hydrogen or C₁-C₆ alkyl;

- R³ is hydrogen or C₁-C₆ alkyl;

- or (b) R¹ and R² form, together with the nitrogen to which they are attached, a 5- to 6-membered heterocyclic or heteroaryl ring, and R³ is as defined above;

- or (c) R¹ and R³ form, together with the -N-C=N- moiety to which they are attached, a 5- to 6-membered heterocyclic or heteroaryl ring, and R² is as defined above;

25 - R⁴ is hydrogen or C₁-C₆ alkyl;

- B is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, -O-L-, -S-L- or L-Het-L', wherein L, L' and Het are as defined above;
- X is -O-, -S- or -NR'-, wherein R' is as defined above;
- R^5 is hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, phenyl, C_3 - C_6 carbocyclyl, 5- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Y-L', -L-Het-A, -L-Y-A or -L-A-Het-A, wherein L, L', Het and Y are as defined above and wherein each A is the same or different and is as defined above;
- R^6 is hydrogen or C_1 - C_6 alkyl;
- 10 - R^7 is hydrogen or C_1 - C_6 alkyl or R^7 is a bond, a C_1 - C_4 alkyl or C_2 - C_4 alkenyl moiety which is joined to one of the carbon atoms of B to form, together with the -N-C- moiety to which R^7 and B are attached, a 5- to 6-membered heterocyclic ring; and
- R^8 is hydrogen or C_1 - C_6 alkyl,

15 wherein:

- the alkyl, alkenyl and alkynyl groups and moieties in the substituents R^1 to R^7 , X and B are unsubstituted or substituted by one, two or three substituents which are the same or different and are selected from fluorine, chlorine, bromine, hydroxy, amino and thio substituents; and
- 20 - the aryl, carbocyclyl, heterocyclyl and heteroaryl groups and moieties in the substituents R^1 and R^5 , the heterocyclic or heteroaryl moieties formed by R^1 together with R^2 and R^1 together with R^3 and the heterocyclic moieties formed by R^7 together with B are typically unsubstituted or substituted by one, two or three substituents selected from fluorine, chlorine, bromine, hydroxy, amino, thio, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylamino, di-(C_1 - C_4 alkyl)amino, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy and C_1 - C_4 haloalkylthio substituents.


2. Use according to claim 1 wherein the compound of formula (I) is not an inhibitor of NOS.

3. Use according to either claim 1 or claim 2 wherein the disease is ischaemia-reperfusion injury of the brain or heart, cancer, lethal hypotension in severe inflammatory conditions, local and systemic inflammatory disorders, neurodegeneration, asthma, pain or sepsis.

5

4. Use according to any of the preceding claims wherein the disease is peritonitis or sepsis.

5. Use according to any one of the preceding claims wherein the compound of
10 formula (I) is a compound of formula (II)



15 wherein:

- either (a) R^{11} is C_3 - C_8 alkyl, allyl, phenyl, cyclopropyl, cyclohexyl, $-(C_1$ - C_2 alkyl)-phenyl, $-(C_1$ - C_2 alkyl)-thienyl, $-(C_1$ - C_2 alkyl)-tetrahydrofuranyl, $-(C_1$ - C_2 alkyl)-O-phenyl, $-(C_1$ - C_2 alkyl)-O- $(C_1$ - C_4 alkyl), $-(C_1$ - C_2 alkyl)-S- $(C_1$ - C_2 alkyl) or $-(C_1$ - C_2 alkyl)-NMe- $(C_1$ - C_2 alkyl);

20 - R^{21} is hydrogen, methyl or ethyl;

- or (b) R^{11} and R^{21} form, together with the nitrogen to which they are attached, a piperidinyl, morpholinyl or pyrrolidinyl ring;

- B^1 is a 1,2-ethyl or 1,3-propyl moiety; and

- R^{51} is hydrogen, C_1 - C_4 alkyl, $-(C_1$ - C_2 alkyl)-phenyl, $-(C_1$ - C_2 alkyl)-O- $(C_1$ - C_2 alkyl) or $-(C_1$ - C_2 alkyl)-phenyl-O-phenyl;

25 wherein:

- the alkyl groups and moieties in the substituents R^{11} , R^{21} , B and R^{51} are unsubstituted or substituted by a single fluoro substituent; and

- the phenyl and carbocycll groups in R¹¹ and the heterocyclic moieties formed by R¹¹ together with R²¹ are unsubstituted and the phenyl groups in R⁵¹ are unsubstituted or substituted by one or two substituents selected from fluorine, C₁-C₂ alkyl, C₁-C₂ alkoxy and C₁-C₂ haloalkyl substituents.

5

6. Use according to claim 5 wherein R¹¹ is -(C₁-C₂ alkyl)-O-(C₁-C₄ alkyl) and/or R⁵¹ is C₁-C₄ alkyl or -(C₁-C₂)-phenyl.

7. Use according to claim 1 wherein B is C₂-C₆ alkyl.

10

8. Use according to either claim 1 or claim 7 wherein R¹ is C₃-C₈ alkyl, C₃-C₆ alkenyl, phenyl, C₃-C₆ carbocycll, 5- to 6-membered heterocycll, 5- to 6-membered heteroaryl, -L-A, -L-Het-L' or -L-Het-A.

15

9. Use according to claim 1, claim 7, or claim 8 wherein R⁵ is hydrogen, C₁-C₆ alkyl, phenyl, C₃-C₆ carbocycll, 5- to 6-membered heterocycll, 5- to 6-membered heteroaryl, -L-A, -L-Het-L', -L-Het-A or -L-A-Het-A.

20

10. Use according to any one of claims 1, 8 and 9 wherein R² is hydrogen or C₁-C₄ alkyl and/or R³ is hydrogen or C₁-C₂ alkyl and/or R¹ and R² form, together with the nitrogen to which they are attached, a 5- to 6-membered heterocyclic ring and/or R⁴ is hydrogen or C₁-C₂ alkyl and/or B is C₂-C₄ alkyl and/or R⁶ is hydrogen or C₁-C₂ alkyl and/or R⁷ is hydrogen or C₁-C₂ alkyl or R⁷ is a C₁-C₂ alkyl moiety which is joined to one of the carbon atoms of B to form, together with the -N-C- moiety to which R⁷ and B are attached, a pyrrolidine or piperidine ring and/or R⁸ is hydrogen or C₁-C₂ alkyl and/or R' is hydrogen or C₁-C₄ alkyl and/or Het is -O-, -S- or -NMe- and/or A is phenyl, 5- to 6-membered heterocycll or 5- to 6- membered heteroaryl and/or L and L' are the same or different and are C₁-C₄ alkyl and/or Y is -CO-, -CO-O- or CO-NR'-.


25

11. A method of treating a patient suffering from or susceptible to a disease as defined in any one of claims 1, 3 and 4, which method comprises administering to said

patient an effective amount of a compound as defined in any one of the preceding claims, or a pharmaceutically acceptable salt thereof.

12. A compound of formula (I), or a pharmaceutically acceptable salt thereof.

5



wherein R¹ is -L-Het-L', -L-Y-L', -L-Het-A or -L-Y-A and R², R³, R⁴, B, X, R⁵, R⁶, R⁷, R⁸, L, Het, L', Y and A are as defined in any one of claims 1 to 10.

10

13. A compound according to claim 12 which is *N*^C-(2-methoxyethyl)-L-arginine or *N*^C-(2-Methoxyethyl)-L-arginine methyl ester, or a pharmaceutically acceptable salt thereof.

15 14. A pharmaceutical composition comprising a compound of formula (I), as defined in claim 12 or claim 13, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

Dated 22 June, 2011
UCL Business PLC

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON




Figure 1

2/3

Fig.2a.





Fig.2b.



3/3

Fig.3.

