



(86) Date de dépôt PCT/PCT Filing Date: 2011/04/01
(87) Date publication PCT/PCT Publication Date: 2011/10/06
(45) Date de délivrance/Issue Date: 2018/05/15
(85) Entrée phase nationale/National Entry: 2012/09/28
(86) N° demande PCT/PCT Application No.: CA 2011/000357
(87) N° publication PCT/PCT Publication No.: 2011/120153
(30) Priorité/Priority: 2010/04/01 (US61/320,223)


(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
A61K 31/506 (2006.01), *A61K 31/519* (2006.01),
A61P 31/18 (2006.01), *C07D 239/46* (2006.01),
C07D 413/04 (2006.01)

(72) Inventeurs/Inventors:
DANTER, WAYNE, CA;
THRELFALL, CLINTON, CA;
GUIZZETTI, SYLVAIN, FR;
MARIN, JULIEN, FR


(73) Propriétaire/Owner:
CRITICAL OUTCOME TECHNOLOGIES INC., CA

(74) Agent: AIRD & MCBURNEY LP

(54) Titre : COMPOSES ET METHODES POUR LE TRAITEMENT DU VIH
(54) Title: COMPOUNDS AND METHOD FOR TREATMENT OF HIV



Formula I



Formula II

(57) **Abrégé/Abstract:**

The invention relates to a compound of Formulae I and/or II: Formula I Formula II and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof; X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S, and the remaining substituents are described herein; and a composition thereof. The invention also relates to a method of administration thereof; and use thereof to treat HIV.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
6 October 2011 (06.10.2011)

PCT

(10) International Publication Number
WO 2011/120153 A1

(51) International Patent Classification:

C07D 471/04 (2006.01) *A61P 31/18* (2006.01)
A61K 31/506 (2006.01) *C07D 239/46* (2006.01)
A61K 31/519 (2006.01) *C07D 413/04* (2006.01)

(21) International Application Number:

PCT/CA2011/000357

(22) International Filing Date:

1 April 2011 (01.04.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/320,223 1 April 2010 (01.04.2010) US

(71) **Applicant** (for all designated States except US): **CRITICAL OUTCOME TECHNOLOGIES INC.** [CA/CA]; Suite 213, 700 Collip Circle, London, Ontario N6G 4X8 (CA).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **DANTER, Wayne** [CA/CA]; 147 Chesham Avenue, London, Ontario N6G 3V2 (CA). **THRELFALL, Clinton** [US/CA]; 370 Victoria Street, London, Ontario N6A 2C9 (CA). **GUIZZETTI, Sylvain** [FR/FR]; c/o NovAliX-BioParc, Boulevard Sebastien Brant, BP 30170, F-67405 Illkirch Cedex (FR). **MARIN, Julien** [FR/FR]; c/o NovAliX-BioParc, Boulevard Sebastien Brant, BP 30170, F-67405 Illkirch Cedex (FR).

(74) **Agents:** **MCMANUS, Kimberly, A.** et al.; Sim & McBurney, 330 University Avenue 6th Floor, Toronto, Ontario M5G 1R7 (CA).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))


Published:

— with international search report (Art. 21(3))

(54) Title: COMPOUNDS AND METHOD FOR TREATMENT OF HIV



Formula I



Formula II

(57) **Abstract:** The invention relates to a compound of Formulae I and/or II: Formula I Formula II and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof; X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S, and the remaining substituents are described herein; and a composition thereof. The invention also relates to a method of administration thereof; and use thereof to treat HIV.

COMPOUNDS AND METHOD FOR TREATMENT OF HIV

FIELD OF THE INVENTION

The present invention relates generally to therapeutic compounds and compositions, as well as methods for treatment of HIV.

BACKGROUND OF THE INVENTION


5 Human immunodeficiency viruses (HIV) are lentiviruses from the family of retroviridae. Transmission is initiated by the passage of HIV across the mucosal barrier of sexual organs or placenta when exposed to infectious body fluids such as semen, vaginal secretions, or blood.

HIV, particularly the strains known as HIV type-1 (HIV-1) and type-2
10 (HIV-2) viruses, is the etiological agent of the complex disease that includes progressive destruction of the immune system AIDS (acquired immune deficiency syndrome) and degeneration of the central and peripheral nervous system. A common feature of retrovirus replication is the action of integrase, a viral enzyme that inserts the viral genome into the DNA of the mammal cell.
15 Since integration is a vital step in retroviral replication, blocking it can halt further spread of the virus.

Toxicity or undesirable side effects of the common drugs for treating HIV infection, e.g., AZT or HIV protease inhibitors, are incompatible with their antiviral activity when used at an effective pharmaceutical concentration.
20 Thus, there is still a need in the art for alternative compounds for treating HIV.

SUMMARY OF THE INVENTION

In an aspect, there is provided a compound of Formula I:




Formula I

5 and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

wherein:

when X is Se, N-OH, NH, or N-CN,  is a double bond, and

10 R¹, R², R³, R⁴ and R⁵ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or
15 unsubstituted heteroaromatic group;

when X is selected from NO₂, CN or N=O,  is a single bond, and


R¹, R², R³, R⁴ and R⁵ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or
20 unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

when X is O or S,  is a double bond,

R¹, R², R³, and R⁴ are each independently selected from H, halo,
25 hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or

unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

R^5 is $-(C=O)NR^6R^7$ or



5


wherein R^6 and R^7 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

wherein Z and A are each independently selected from CR^9 or N, and R^8 and R^9 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

In another aspect, R^1 , R^2 , R^3 , and R^4 are each independently selected from H, halo, hydroxyl, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted haloalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkoxy, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, a substituted or unsubstituted heteroaromatic group, carboxyl, alkylcarbonyl, arylcarbonyl, cycloalkylcarbonyl, heterocyclcarbonyl, amino, aminoalkyl, alkylaminoalkyl, heterocyclalkyl, aralkyl, arylalkenyl, arylalkynyl, alkylthio, alkylamino, arylamino, heteroaryl amino, aralkylamino, alkylaminoalkylamino, arylthio, aralkylthio, aryloxy, aralkoxy, heterocyclalkoxy, heterocyclloxyalkyl, cycloalkyl, and cycloalkenyl.

In a further aspect, R^1 , R^2 , R^3 , and R^4 are each independently selected from H or substituted or unsubstituted alkyl. In another aspect, X is selected from Se, N-OH, NH, NO_2 , CN, N-CN, or N=O; and R^5 is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In yet another aspect, R^5 is $-(C=O)Y$ and Y is selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In still another aspect, Y is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In another aspect, the $-(C=O)Y$ is $-(C=O)NR^6R^7$ and R^6 and R^7 are each independently selected from H or the substituted or unsubstituted hydrocarbon group. In another aspect, the substituted or unsubstituted hydrocarbon group is selected from a substituted or unsubstituted alkyl group. In another aspect, the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. In another aspect, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group.


In a further aspect, R^5 is $-(C=O)OR^6$ and R^6 is selected from H or a substituted or unsubstituted hydrocarbon group. In another aspect, R^6 is selected from a substituted or unsubstituted alkyl group. In yet another aspect, R^6 is H. In another aspect, R^1 , R^2 , R^3 , and R^4 are each independently selected from a substituted or unsubstituted alkylaryl. In another aspect, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group. In another aspect, the compound is selected from:




and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.

- In another aspect, X is selected from Se, N-OH, NH, or N=O. In another aspect, X is O or S; and R^6 and R^7 are each independently selected from H or the substituted or unsubstituted hydrocarbon group. In another aspect, the substituted or unsubstituted hydrocarbon group is selected from a substituted or unsubstituted alkyl group. In yet another aspect, the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. In another aspect, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group. In another aspect, the halo group is selected from bromo, chloro, fluoro or iodo.


In another aspect, the compound is selected from:




15



,




,




,


5



,




,




,


5



,




5




and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical
 10 isomer, E-isomer, Z-isomer, or combination thereof.

In another aspect, X is selected from Se, N-OH, NH, or N-CN; and R⁵
 is selected from H, a substituted or unsubstituted hydrocarbon group, a
 substituted or unsubstituted heterogeneous group, a substituted or
 unsubstituted aromatic group, or a substituted or unsubstituted
 15 heteroaromatic group. In another aspect, R⁵ is selected from a substituted or
 unsubstituted aromatic group, or a substituted or unsubstituted
 heteroaromatic group. In another aspect, R⁵ is a substituted or unsubstituted
 heteroaromatic group. In another aspect, R⁵ is




20 wherein Z and A are each independently selected from CR⁹ or N, and R⁸ and
 R⁹ are each independently selected from H, halo, hydroxyl, amino, a

substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In
 5 another aspect, X is O or S and R⁵ is the




wherein Z and A are each independently selected from CR⁹ or N, and R⁸ and R⁹ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In yet
 10 another aspect, at least one of Z and A is N. In another aspect, both Z and A are N. In another aspect, R⁸ is a substituted or unsubstituted hydrocarbon group. In another aspect, R⁸ is selected from a substituted or unsubstituted alkyl group. In another aspect, the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. In another aspect, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is
 15 substituted with a halo group. In another aspect, the halo group is selected from bromo, chloro, fluoro or iodo.
 20

In yet another aspect, the compound is selected from:



and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical
 5 isomer, E-isomer, Z-isomer, or combination thereof.

In still another aspect, there is provided a compound of Formula II:



10

Formula II

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical
 isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

15 wherein:

when X is Se, N-OH, NH, N-CN, or S,  is a double bond, and


R¹, R², and R³ are each independently selected from H, halo, hydroxyl,
 amino, a substituted or unsubstituted hydrocarbon group, a substituted or
 unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic
 20 group, a substituted or unsubstituted heterocyclic group, a substituted or

unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group;

when X is selected from NO₂, CN or N=O,  is a single bond, and

R¹, R², and R³ are each independently selected from H, halo, hydroxyl,

5 amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and


10 when X is O,  is a double bond,

R¹ and R² are each independently selected from H, halo, hydroxyl,

amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or

15 unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

R³ is




wherein Z and A are each independently selected from CR¹⁰ or N, and

20 R⁴ and R¹⁰ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

25 In another aspect, R¹ and R² are each independently selected from H, halo, hydroxyl, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted haloalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkoxy, a substituted or unsubstituted heterocyclic group, a

substituted or unsubstituted aromatic group, a substituted or unsubstituted heteroaromatic group, carboxyl, alkylcarbonyl, arylcarbonyl, cycloalkylcarbonyl, heterocyclylcarbonyl, amino, aminoalkyl, alkylaminoalkyl, heterocyclylalkyl, aralkyl, arylalkenyl, arylalkynyl, alkylthio, alkylamino, arylamino, heteroaryl amino, aralkyl amino, alkylaminoalkyl amino, arylthio, aralkylthio, aryloxy, aralkoxy, heterocyclylalkoxy, heterocyclyloxyalkyl, cycloalkyl, and cycloalkenyl. In yet another aspect, R^1 and R^2 are each independently selected from H or substituted or unsubstituted alkyl. In another aspect, X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, or S; and R^3 is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In another aspect, R^3 is selected from a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In another aspect, R^3 is a substituted or unsubstituted heteroaromatic group. In still another aspect, R^3 is



wherein Z and A are each independently selected from CR¹⁰ or N, and R^4 and R^{10} are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In another aspect, X is O. In another aspect, at least one of Z and A is N. In another aspect, both Z and A are N. In another aspect, R^4 is a substituted or unsubstituted hydrocarbon group. In another aspect, R^4 is selected from a substituted or unsubstituted alkyl group. In another aspect, the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. In another aspect, the substituted

or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group. In another aspect, the halo group is selected from bromo, chloro, fluoro or iodo. In another aspect, X is selected from Se, N-OH, NH, N-CN, or S; and R³ is selected from



5 H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In another aspect, R³ is $-(C=O)Y$ and Y is selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted

10 heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In still another aspect, Y is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a


15 substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group. In another aspect, the $-(C=O)Y$ is $-(C=O)NR^8R^9$ and R⁸ and R⁹ are each independently selected from H or the substituted or unsubstituted hydrocarbon group. In another aspect, the substituted or unsubstituted hydrocarbon group is selected from a substituted or

20 unsubstituted alkyl group. In another aspect, the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. In another aspect, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group. In another aspect, the


25 compound is selected from:




5




10



,




,




,

5


10




,



,



,



,

5

10

15

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.

In another aspect, wherein the compound above is an HIV integrase
5 inhibitor.

In another aspect, there is provided a pharmaceutical composition comprising the compound above, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent. In
10 another aspect, there is provided the pharmaceutical composition in combination with at least one other anti-HIV agent.

In another aspect, there is provided a method for the treatment of HIV in a mammal comprising administering to said mammal an anti-HIV effective treatment amount of the compound above, and/or a pharmaceutically-
15 acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent.

In another aspect, there is provided a method for salvage therapy in the treatment of HIV in a mammal comprising administering to said mammal
20 an anti-HIV effective treatment amount of a compound above, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent. In another aspect, the mammal is human. In another aspect, the method further comprises co-administering at least one
25 other anti-HIV agent in combination and/or in alternation with the compound.

In yet another aspect, there is provided a use of an anti-HIV effective treatment amount of the compound above, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable
30 carrier or diluent, in the manufacture of a medicament for the treatment of HIV in a mammal.


In another aspect, there is provided a use of an anti-HIV effective treatment amount of the compound above, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, in the manufacture of a medicament for salvage therapy in the treatment of HIV in a mammal.

In another aspect, there is provided a use of an anti-HIV effective treatment amount of the compound above, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, for the treatment of HIV in a mammal.

In another aspect, there is provided a use of an anti-HIV effective treatment amount of the compound above, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, for salvage therapy in the treatment of HIV in a mammal.

In another aspect, the mammal is human. In another aspect, the use further comprises use of at least one other anti-HIV agent in combination and/or in alternation with the compound.

In yet another aspect, there is provided a method for making the compound of Formula III:



and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S, wherein:

5 when X is selected from Se, N-OH, NH, N-CN, O or S, $\text{---}\text{X}\text{---}$ is a double bond,

when X is selected from NO₂, CN or N=O, $\text{---}\text{X}\text{---}$ is a single bond,


R¹, R², R³, and R⁴ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

Z and A are each independently selected from CR⁹ or N, and

15 R⁸ and R⁹ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group,

20 wherein the method comprises:


a) reacting a compound of Formula VII with an amine of NH₂AHC(O)R⁸ to form an intermediate of Formula VIII:



b) reacting the intermediate of Formula VIII with base to yield Formula III, whereby X is oxygen;

c) reacting Formula III, whereby X is oxygen, under conditions to replace the oxygen of the carbonyl group with any X group defined herein to form the compounds of Formula III. In yet another aspect, wherein c) comprises reacting Formula III, whereby X is oxygen, with NH_2OH to form the compound of Formula III, wherein X is N-OH.

In another aspect, there is provided a method for making the compound of Formula IV:



Formula IV

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO_2 , CN, N-CN, N=O, O or S, wherein:

when X is Se, N-OH, NH, or N-CN, ---X--- is a double bond,

when X is selected from NO_2 , CN or N=O, ---X--- is a single bond,


when X is O or S, ---X--- is a double bond,

R^1 , R^2 , R^3 , and R^4 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

wherein R^6 and R^7 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group,


wherein the method comprises:

a) reacting a compound of Formula VII with an amine of NHR^6R^7 to form an intermediate of Formula IX:



b) reacting the intermediate of Formula IX under conditions to replace the oxygen of the carbonyl group in the ring with any X group defined herein to form the compound of Formula IV. In another aspect, wherein b) comprises reacting Formula IX with NH_2OH to form the compound of Formula IV, wherein X is N-OH.

In yet another aspect, there is provided a method for making the compound of Formula V:



Formula V

5 and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

wherein:

when X is Se, N-OH, NH, N-CN, or S,  is a double bond, and


10 when X is selected from NO₂, CN or N=O,  is a single bond,

R¹ and R² are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

wherein R⁶ and R⁷ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group,

wherein the method comprises:

a) reacting a compound of Formula XII with an amine of NHR⁶R⁷ to
25 form an intermediate of Formula XIII:




Formula XII

Formula XIII

- b) reacting the intermediate of Formula XIII under conditions to replace the oxygen of the carbonyl group in the ring with any X group defined herein to form the compound of Formula V. In yet another aspect, wherein b) comprises reacting Formula XIII with NH₂OH to form the compound of Formula V, wherein X is N-OH.

In another aspect, there is provided a method for making the compound of Formula VI:



Formula VI

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;


- 15 X is N-OH and is a double bond,

R¹ and R² are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or

unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

wherein Z and A are each independently selected from CR¹⁰ or N, and R⁴ and R¹⁰ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, wherein the method comprises:

- 10 a) reacting a compound of Formula X, whereby R is a benzyl group, with an amine of NH₂AHC(O)R⁴ to form an intermediate of Formula XI:



Formula X

Formula XI

- b) reacting the intermediate of Formula XI with base to yield Formula VI, whereby X is oxygen;
- 15 c) reacting Formula VI, whereby X is oxygen, with Lawesson's reagent, to yield Formula VI, whereby X is sulfur;
- d) reacting Formula VI, whereby X is sulfur and R is a benzyl group, with NH₂OCH₂Ar, to yield Formula VI, whereby X is N-O-CH₂Ar; and
- e) reducing Formula VI, whereby X is N-O-CH₂Ar, to yield Formula VI, 20 whereby X is N-OH and R is a H.

Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating embodiments of the invention are given by way of illustration only, 25 since various changes and modifications within the spirit and scope of the

invention will become apparent to those skilled in the art from the detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Certain embodiments of the present invention will now be described more fully with reference to the accompanying drawings:

Figure 1 shows a graph of the effect of COTI-HIV1Doxo in an HIV-1 integrase assay; and

10 Figure 2 shows plots of the log-transformation of drug concentrations and best-fit curve for COTI-HIV-1 (A), COTI-HIV28233 (B), COTI-HIV28236 (C), and two sodium azide controls (D & E) to allow calculation of IC₅₀ using non-linear regression.

DETAILED DESCRIPTION

15 The present invention is directed to new compounds, compositions comprising such compounds, a method of administration thereof, and use thereof to treat HIV. The compounds of this invention may function as inhibitors of HIV integrase.

Definitions

20 When describing the compounds, compositions, methods and uses of this invention, the following terms have the following meanings unless otherwise indicated.

25 The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

30 The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described, for example, in: E. L. Eliel and S. H. Wilen, Stereo-chemistry of Carbon Compounds, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic

mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, E isomers, and Z isomers, being included in the present invention. In addition, the compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be
5 encompassed by the scope of the invention, even though only one tautomeric structure may be depicted.

Generally, reference to a certain element such as hydrogen or H is meant to, if appropriate, include all isotopes of that element.

Where the term "alkyl group" is used, either alone or within other terms
10 such as "haloalkyl group" and "alkylamino group", it encompasses linear or branched carbon radicals having, for example, one to about twenty carbon atoms or, in specific embodiments, one to about twelve carbon atoms. In other embodiments, alkyl groups are "lower alkyl" groups having one to about six carbon atoms. Examples of such groups include, but are not limited
15 thereto, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl and the like. In more specific embodiments, lower alkyl groups have one to four carbon atoms.

The term "alkenyl group" encompasses linear or branched carbon radicals having at least one carbon-carbon double bond. The term "alkenyl
20 group" can encompass conjugated and non-conjugated carbon-carbon double bonds or combinations thereof. An alkenyl group, for example and without being limited thereto, can encompass two to about twenty carbon atoms or, in a particular embodiment, two to about twelve carbon atoms. In embodiments, alkenyl groups are "lower alkenyl" groups having two to about four carbon
25 atoms. Examples of alkenyl groups include, but are not limited thereto, ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The terms "alkenyl group" and "lower alkenyl group", encompass groups having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations.

The term "alkynyl group" denotes linear or branched carbon radicals
30 having at least one carbon-carbon triple bond. The term "alkynyl group" can encompass conjugated and non-conjugated carbon-carbon triple bonds or

combinations thereof. Alkynyl group, for example and without being limited thereto, can encompass two to about twenty carbon atoms or, in a particular embodiment, two to about twelve carbon atoms. In embodiments, alkynyl groups are "lower alkynyl" groups having two to about ten carbon atoms.

- 5 Some examples are lower alkynyl groups having two to about four carbon atoms. Examples of such groups include propargyl, butynyl, and the like.

The term "halo" means halogens such as fluorine, chlorine, bromine or iodine atoms.

- The term "haloalkyl group" encompasses groups wherein any one or
10 more of the alkyl carbon atoms is substituted with halo as defined above. Specifically encompassed are monohaloalkyl, dihaloalkyl and polyhaloalkyl groups including perhaloalkyl. A monohaloalkyl group, for one example, may have either an iodo, bromo, chloro or fluoro atom within the group. Dihalo and polyhaloalkyl groups may have two or more of the same halo atoms or a
15 combination of different halo groups. "Lower haloalkyl group" encompasses groups having 1- 6 carbon atoms. In some embodiments, lower haloalkyl groups have one to three carbon atoms. Examples of haloalkyl groups include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl,
20 difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl.

- The term "hydroxyalkyl group" encompasses linear or branched alkyl groups having, for example and without being limited thereto, one to about ten carbon atoms, any one of which may be substituted with one or more hydroxyl
25 groups. In embodiments, hydroxyalkyl groups are "lower hydroxyalkyl" groups having one to six carbon atoms and one or more hydroxyl groups. Examples of such groups include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl.

- The term "alkoxy group" encompasses linear or branched oxy-
30 containing groups each having alkyl portions of, for example and without being limited thereto, one to about ten carbon atoms. In embodiments, alkoxy

groups are "lower alkoxy" groups having one to six carbon atoms. Examples of such groups include methoxy, ethoxy, propoxy, butoxy and tert-butoxy. In certain embodiments, lower alkoxy groups have one to three carbon atoms. The "alkoxy" groups may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" groups. In other embodiments, lower haloalkoxy groups have one to three carbon atoms. Examples of such groups include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy, and fluoropropoxy.

The term "aromatic group" or "aryl group" means an aromatic group having one or more rings wherein such rings may be attached together in a pendent manner or may be fused. In particular embodiments, an aromatic group is one, two or three rings. Monocyclic aromatic groups may contain 4 to 10 carbon atoms, typically 4 to 7 carbon atoms, and more typically 4 to 6 carbon atoms in the ring. Typical polycyclic aromatic groups have two or three rings. Polycyclic aromatic groups having two rings typically have 8 to 12 carbon atoms, preferably 8 to 10 carbon atoms in the rings. Examples of aromatic groups include, but are not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, biphenyl, phenanthryl, anthryl or acenaphthyl.

The term "heteroatom" means an atom other than carbon. Typically, heteroatoms are selected from the group consisting of sulfur, phosphorous, nitrogen and oxygen atoms. Groups containing more than one heteroatom may contain different heteroatoms.

The term "heteroaromatic group" or "heteroaryl group" means an aromatic group having one or more rings wherein such rings may be attached together in a pendent manner or may be fused, wherein the aromatic group has at least one heteroatom. Monocyclic heteroaromatic groups may contain 4 to 10 member atoms, typically 4 to 7 member atoms, and more typically 4 to 6 member atoms in the ring. Typical polycyclic heteroaromatic groups have two or three rings. Polycyclic aromatic groups having two rings typically have 8 to 12 member atoms, more typically 8 to 10 member atoms in the rings. Examples of heteroaromatic groups include, but are not limited thereto,

pyrrole, imidazole, thiazole, oxazole, furan, thiophene, triazole, pyrazole, isoxazole, isothiazole, pyridine, pyrazine, pyridazine, pyrimidine, triazine, indole, benzofuran, benzothiophene, benzimidazole, benzthiazole, quinoline, isoquinoline, quinazoline, quinoxaline and the like.

5 The term "carbocyclic group" means a saturated or unsaturated carbocyclic hydrocarbon ring. Carbocyclic groups are not aromatic. Carbocyclic groups are monocyclic or polycyclic. Polycyclic carbocyclic groups can be fused, spiro, or bridged ring systems. Monocyclic carbocyclic groups may contain 4 to 10 carbon atoms, typically 4 to 7 carbon atoms, and
10 more typically 5 to 6 carbon atoms in the ring. Bicyclic carbocyclic groups may contain 8 to 12 carbon atoms, typically 9 to 10 carbon atoms in the rings.

 The term "heterocyclic group" means a saturated or unsaturated ring structure containing carbon atoms and 1 or more heteroatoms in the ring. Heterocyclic groups are not aromatic. Heterocyclic groups are monocyclic or
15 polycyclic. Polycyclic heterocyclic groups can be fused, spiro, or bridged ring systems. Monocyclic heterocyclic groups may contain 4 to 10 member atoms (i.e., including both carbon atoms and at least 1 heteroatom), typically 4 to 7, and more typically 5 to 6 in the ring. Bicyclic heterocyclic groups may contain 8 to 18 member atoms, typically 9 or 10 member atoms in the rings.
20 Representative heterocyclic groups include, by way of example, pyrrolidine, imidazolidine, pyrazolidine, piperidine, 1,4-dioxane, morpholine, thiomorpholine, piperazine, 3-pyrroline and the like.

 The term "heterogeneous group" means a saturated or unsaturated chain of non-hydrogen member atoms comprising carbon atoms and at least
25 one heteroatom. Heterogeneous groups typically have 1 to 25 member atoms. More typically, the chain contains 1 to 12 member atoms, 1 to 10, and most typically 1 to 6. The chain may be linear or branched. Typical branched heterogeneous groups have one or two branches, more typically one branch. Typically, heterogeneous groups are saturated. Unsaturated heterogeneous
30 groups may have one or more double bonds, one or more triple bonds, or both. Typical unsaturated heterogeneous groups have one or two double

bonds or one triple bond. More typically, the unsaturated heterogeneous group has one double bond.

The term "hydrocarbon group" or "hydrocarbyl group" means a chain of 1 to 25 carbon atoms, typically 1 to 12 carbon atoms, more typically 1 to 10 carbon atoms, and most typically 1 to 8 carbon atoms. Hydrocarbon groups may have a linear or branched chain structure. Typical hydrocarbon groups have one or two branches, typically one branch. Typically, hydrocarbon groups are saturated. Unsaturated hydrocarbon groups may have one or more double bonds, one or more triple bonds, or combinations thereof.

Typical unsaturated hydrocarbon groups have one or two double bonds or one triple bond; more typically unsaturated hydrocarbon groups have one double bond.

When the term "unsaturated" is used in conjunction with any group, the group may be fully unsaturated or partially unsaturated. However, when the term "unsaturated" is used in conjunction with a specific group defined herein, the term maintains the limitations of that specific group. For example, an unsaturated "carbocyclic group", based on the limitations of the "carbocyclic group" as defined herein, does not encompass an aromatic group.

The terms "carboxy group" or "carboxyl group", whether used alone or with other terms, such as "carboxyalkyl group", denotes $-(C=O)-O-$.

The term "carbonyl group", whether used alone or with other terms, such as "aminocarbonyl group", denotes $-(C=O)-$.

The terms "alkylcarbonyl group" denotes carbonyl groups which have been substituted with an alkyl group. In certain embodiments, "lower alkylcarbonyl group" has lower alkyl group as described above attached to a carbonyl group.

The term "aminoalkyl group" encompasses linear or branched alkyl groups having one to about ten carbon atoms any one of which may be substituted with one or more amino groups. In some embodiments, the aminoalkyl groups are "lower aminoalkyl" groups having one to six carbon

atoms and one or more amino groups. Examples of such groups include aminomethyl, aminoethyl, aminopropyl, aminobutyl and aminohexyl.

The term "alkylaminoalkyl group" encompasses aminoalkyl groups having the nitrogen atom independently substituted with an alkyl group. In
5 certain embodiments, the alkylaminoalkyl groups are "loweralkylaminoalkyl" groups having alkyl groups of one to six carbon atoms. In other embodiments, the lower alkylaminoalkyl groups have alkyl groups of one to three carbon atoms. Suitable alkylaminoalkyl groups may be mono or dialkyl substituted, such as N-methylaminomethyl, N, N-dimethyl-aminoethyl, N, N-
10 diethylaminomethyl and the like.

The term "aralkyl group" encompasses aryl-substituted alkyl groups. In embodiments, the aralkyl groups are "lower aralkyl" groups having aryl groups attached to alkyl groups having one to six carbon atoms. In other
15 embodiments, the lower aralkyl groups phenyl is attached to alkyl portions having one to three carbon atoms. Examples of such groups include benzyl, diphenylmethyl and phenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy.

The term "arylalkenyl group" encompasses aryl-substituted alkenyl groups. In embodiments, the arylalkenyl groups are "lower arylalkenyl" groups
20 having aryl groups attached to alkenyl groups having two to six carbon atoms. Examples of such groups include phenylethenyl. The aryl in said arylalkenyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy.

The term "arylalkynyl group" encompasses aryl-substituted alkynyl
25 groups. In embodiments, arylalkynyl groups are "lower arylalkynyl" groups having aryl groups attached to alkynyl groups having two to six carbon atoms. Examples of such groups include phenylethynyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The terms benzyl and phenylmethyl are interchangeable.

30 The term "alkylthio group" encompasses groups containing a linear or branched alkyl group, of one to ten carbon atoms, attached to a divalent sulfur

atom. In certain embodiments, the lower alkylthio groups have one to three carbon atoms. An example of "alkylthio" is methylthio, (CH₃S-).

The term "alkylamino group" denotes amino groups which have been substituted with one alkyl group and with two alkyl groups, including terms "N-alkylamino" and "N,N-dialkylamino". In embodiments, alkylamino groups are
5 "lower alkylamino" groups having one or two alkyl groups of one to six carbon atoms, attached to a nitrogen atom. In other embodiments, lower alkylamino groups have one to three carbon atoms. Suitable "alkylamino" groups may be mono or dialkylamino such as N-methylamino, N-ethylamino, N,N-
10 dimethylamino, N,N-diethylamino and the like.

The term "arylamino group" denotes amino groups which have been substituted with one or two aryl groups, such as N-phenylamino. The "arylamino" groups may be further substituted on the aryl ring portion of the group.

15 The term "heteroarylamino" denotes amino groups which have been substituted with one or two heteroaryl groups, such as N-thienylamino. The "heteroarylamino" groups may be further substituted on the heteroaryl ring portion of the group.

The term "aralkylamino group" denotes amino groups which have been
20 substituted with one or two aralkyl groups. In other embodiments, there are phenyl-C₁-C₃-alkylamino groups, such as N-benzylamino. The "aralkylamino" groups may be further substituted on the aryl ring portion of the group.

The term "alkylaminoalkylamino group" denotes alkylamino groups which have been substituted with one or two alkylamino groups. In
25 embodiments, there are C₁-C₃-alkylamino- C₁-C₃-alkylamino groups.

The term "arylthio group" encompasses aryl groups of six to ten carbon atoms, attached to a divalent sulfur atom. An example of "arylthio" is phenylthio. The term "aralkylthio group" encompasses aralkyl groups as described above, attached to a divalent sulfur atom. In certain embodiments
30 there are phenyl- C₁-C₃-alkylthio groups. An example of "aralkylthio" is benzylthio.

The term "aryloxy group" encompasses optionally substituted aryl groups, as defined above, attached to an oxygen atom. Examples of such groups include phenoxy.

5 The term "aralkoxy group" encompasses oxy-containing aralkyl groups attached through an oxygen atom to other groups. In certain embodiments, aralkoxy groups are "lower aralkoxy" groups having optionally substituted phenyl groups attached to lower alkoxy group as described above.

10 The term "cycloalkyl group" includes saturated carbocyclic groups. In certain embodiments, cycloalkyl groups include C₃-C₆ rings. In embodiments, there are compounds that include, cyclopentyl, cyclopropyl, and cyclohexyl.

The term "cycloalkenyl group" includes carbocyclic groups that have one or more carbon-carbon double bonds; conjugated or non-conjugated, or a combination thereof. "Cycloalkenyl" and "cycloalkyldienyl" compounds are included in the term "cycloalkenyl". In certain embodiments, cycloalkenyl
15 groups include C₃-C₆ rings. Examples include cyclopentenyl, cyclopentadienyl, cyclohexenyl and cycloheptadienyl. The "cycloalkenyl" group may have 1 to 3 substituents such as lower alkyl, hydroxyl, halo, haloalkyl, nitro, cyano, alkoxy, lower alkylamino, and the like.

20 The term "suitable substituent", "substituent" or "substituted" used in conjunction with the groups described herein refers to a chemically and pharmaceutically acceptable group, i.e., a moiety that does not negate the therapeutic activity of the inventive compounds. It is understood that substituents and substitution patterns on the compounds of the invention may be selected by one of ordinary skill in the art to provide compounds that are
25 chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below. If a substituent is itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon/member atom or on different carbons/member atoms, as long as a stable structure results. Illustrative
30 examples of some suitable substituents include, cycloalkyl, heterocyclyl, hydroxyalkyl, benzyl, carbonyl, halo, haloalkyl, perfluoroalkyl, perfluoroalkoxy,

alkyl, alkenyl, alkynyl, hydroxy, oxo, mercapto, alkylthio, alkoxy, aryl or heteroaryl, aryloxy or heteroaryloxy, aralkyl or heteroaralkyl, aralkoxy or heteroaralkoxy, HO--(C=O)--, amido, amino, alkyl- and dialkylamino, cyano, nitro, carbamoyl, alkylcarbonyl, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylcarbonyl, aryloxycarbonyl, alkylsulfonyl, and arylsulfonyl. Typical substituents include aromatic groups, substituted aromatic groups, hydrocarbon groups including alkyl groups such as methyl groups, substituted hydrocarbon groups such as benzyl, and heterogeneous groups including alkoxy groups such as methoxy groups.

10 The term "fused" means in which two or more carbons/member atoms are common to two adjoining rings, e.g., the rings are "fused rings".


 The pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

 The pharmaceutically acceptable salts of the compounds of this invention can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with the appropriate inorganic or organic base.

The present invention includes pharmaceutically acceptable salts, solvates and prodrugs of the compounds of the invention and mixtures thereof.

The terms "comprising", "having" and "including", and various endings thereof, are meant to be open ended, including the indicated component but not excluding other elements.

A compound of the invention is represented by a compound of Formula I:



10


Formula I

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

15 X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S, wherein:

when X is Se, N-OH, NH, or N-CN,  is a double bond, and

R¹, R², R³, R⁴ and R⁵ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group;


25 when X is selected from NO₂, CN or N=O,  is a single bond, and R¹, R², R³, R⁴ and R⁵ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a

substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

5 when X is O or S,  is a double bond,

R^1 , R^2 , R^3 , and R^4 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and


R^5 is $-(C=O)NR^6R^7$ or



wherein R^6 and R^7 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

20 wherein Z and A are each independently selected from CR^9 or N, and R^8 and R^9 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

A compound of the invention is also represented by a compound of Formula II:




Formula II

5 and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

wherein:


when X is Se, N-OH, NH, N-CN, or S,  is a double bond, and

10 R¹, R², and R³ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted

15 heteroaromatic group;

when X is selected from NO₂, CN or N=O,  is a single bond, and


R¹, R², and R³ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

when X is O,  is a double bond,

25 R¹ and R² are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or


unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

R^3 is



- 5 wherein Z and A are each independently selected from CR^{10} or N, and R^4 and R^{10} are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.
- 10

In one embodiment, there is provided a compound represented by a compound of Formula III:




15

Formula III

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

- 20 X is selected from Se, N-OH, NH, NO_2 , CN, N-CN, N=O, O or S, wherein:


when X is selected from Se, N-OH, NH, N-CN, O or S,  is a double bond,

when X is selected from NO₂, CN or N=O,  is a single bond,
 R¹, R², R³, and R⁴ are each independently selected from H, halo,
 hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a
 substituted or unsubstituted heterogeneous group, a substituted or
 5 unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic
 group, a substituted or unsubstituted aromatic group, or a substituted or
 unsubstituted heteroaromatic group, and

Z and A are each independently selected from CR⁹ or N, and
 R⁸ and R⁹ are each independently selected from H, halo, hydroxyl, amino, a
 10 substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted
 heterogeneous group, a substituted or unsubstituted carbocyclic group, a
 substituted or unsubstituted heterocyclic group, a substituted or unsubstituted
 aromatic group, or a substituted or unsubstituted heteroaromatic group.

In another embodiment, both Z and A are N and R⁸ is a substituted or
 15 unsubstituted hydrocarbon group. In a more specific embodiment, R⁸ is
 selected from a substituted or unsubstituted alkyl group, such as a substituted
 or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl.
 More specifically, the substituted or unsubstituted alkylaryl is a substituted or
 unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo
 20 group such as a 4-fluorobenzyl group. R¹, R², R³, and R⁴ may be specifically
 H or substituted or unsubstituted alkyl groups.

In a further embodiment, there is provided a compound represented by
 a compound of Formula IV:



25

Formula IV

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

5 wherein:

when X is Se, N-OH, NH, or N-CN,  is a double bond,

when X is selected from NO₂, CN or N=O,  is a single bond,


when X is O or S,  is a double bond,

R¹, R², R³, and R⁴ are each independently selected from H, halo,
10 hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and


15 wherein R⁶ and R⁷ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or
20 unsubstituted heteroaromatic group.

In another embodiment, R⁶ and R⁷ are each independently selected from a substituted or unsubstituted alkyl group. The substituted or unsubstituted alkyl group may be a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. More specifically, the substituted
25 or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group such as a 4-fluorobenzyl group. R¹, R², R³, and R⁴ may be specifically H or substituted or unsubstituted alkyl groups.

Specific embodiments include, without being limited thereto,
30 compounds such as:




COTI-HIV28236-SeA




5

COTI-HIV28236-SA




COTI-HIV28236

10




COTI-HIV28233 (COTI-HIV1oxime)

5



COTI-HIV1oxo


,



COTI-HIV1thio


,

10




COTI-HIV1Se ,

5




COTI-HIV1imine ,

10



COTI-HIV2oxo

5




COTI-HIV2imine

10

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.




In a further embodiment, there is provided a compound represented by
15 a compound of Formula IVA:



Formula IV

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S, wherein:


- 5 when X is Se, N-OH, NH, or N-CN,  is a double bond,
 when X is selected from NO₂, CN or N=O,  is a single bond,
 when X is O or S,  is a double bond,

- R¹, R², R³, and R⁴ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a
10 substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

- wherein R⁶ is selected from H, halo, a substituted or unsubstituted
15 hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.


- In another embodiment, R⁶ is selected from a substituted or
20 unsubstituted alkyl group. In yet another embodiment, R⁶ is H. In another embodiment, R¹, R², R³, and R⁴ are each independently selected from a substituted or unsubstituted alkylaryl. In another embodiment, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group.

- 25 In a further embodiment, the compound is selected from:



and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.

In yet another embodiment, there is provided a compound represented
5 by a compound of Formula V:



Formula V

10

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,
wherein:

15

when X is Se, N-OH, NH, N-CN, or S, ----- is a double bond, and

when X is selected from NO₂, CN or N=O, ----- is a single bond,


R¹ and R² are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or

unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

- 5 wherein R^6 and R^7 are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or
10 unsubstituted heteroaromatic group.

 In another embodiment, R^6 and R^7 are each independently selected from a substituted or unsubstituted alkyl group. The substituted or unsubstituted alkyl group may be a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl. More specifically, the substituted
15 or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group such as a 4-fluorobenzyl group. R^1 and R^2 may be specifically H or substituted or unsubstituted alkyl groups.

 In another embodiment, there is provided a compound of Formula VI:



20

Formula VI

and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical
25 isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO_2 , CN, N-CN, N=O, O or S,

wherein:

when X is Se, N-OH, NH, N-CN, O or S,  is a double bond,

when X is selected from NO₂, CN or N=O,  is a single bond,

R¹ and R² are each independently selected from H, halo, hydroxyl,


5 amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

10 wherein Z and A are each independently selected from CR¹⁰ or N, and R⁴ and R¹⁰ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

In another embodiment, both Z and A are N and R⁴ is a substituted or unsubstituted hydrocarbon group. In a more specific embodiment, R⁴ is selected from a substituted or unsubstituted alkyl group, such as a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl.


20 More specifically, the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group is substituted with a halo group such as a 4-fluorobenzyl group. R¹ and R² may be specifically H or substituted or unsubstituted alkyl groups.

Specific embodiments include, without being limited thereto,
25 compounds such as:



COTI-HIV1DOxime


,



5

COTI-HIV1Doxo


,



10


COTI-HIV1Dthio

,




COTI-HIV1Dcyano ,

5




COTI-HIV1Dimino ,

10



COTI-HIV2DOxime ,


15



COTI-HIV2Doxo

,


5



COTI-HIV2Dthio

,


10



COTI-HIV2Dcyano


,

15



5 The compounds of this invention may be prepared by employing reactions and standard manipulations that are known in the literature or exemplified herein.


10 a) reacting a compound of Formula VII with an amine of $\text{NH}_2\text{AHC}(\text{O})\text{R}^8$
to form an intermediate of Formula VIII:



c) reacting Formula III, whereby X is oxygen, with NH_2OH (or under conditions to replace the oxygen of the carbonyl group in the ring with any X group mentioned herein (e.g. Se, S, etc.)) to form the compounds of Formula

The compounds of Formula IV described herein can be prepared as follows:

a) reacting a compound of Formula VII with an amine of NHR^6R^7 to form an intermediate of Formula IX:




5

b) reacting the intermediate of Formula IX with NH_2OH (or under conditions to replace the oxygen of the carbonyl group in the ring with any X group mentioned herein (e.g. Se, S, etc.)) to form the compound of Formula IV.

10 The compounds of Formula V described herein can be prepared as follows:

a) reacting a compound of Formula XII with an amine of NHR^6R^7 to form an intermediate of Formula XIII:




15

b) reacting the intermediate of Formula XIII with NH_2OH (or under conditions to replace the oxygen of the carbonyl group in the ring with any X group mentioned herein (e.g. Se, S, etc.)) to form the compound of Formula V.

The compounds of Formula VI described herein can be prepared as follows:

a) reacting a compound of Formula X (R is a benzyl group) with an amine of $\text{NH}_2\text{AHC}(\text{O})\text{R}^4$ to form an intermediate of Formula XI:



Formula X

Formula XI

b) reacting the intermediate of Formula XI with base to yield Formula VI, whereby X is oxygen and R is a benzyl group;

c) reacting Formula VI, whereby X is oxygen and R is a benzyl group, with Lawesson's reagent, to yield Formula VI, whereby X is sulfur and R is a benzyl group;

d) reacting Formula VI, whereby X is sulfur and R is a benzyl group, with $\text{NH}_2\text{OCH}_2\text{Ar}$, to yield Formula VI, whereby X is $\text{N-O-CH}_2\text{Ar}$ and R is a benzyl group; and

e) reacting Formula VI, whereby X is $\text{N-O-CH}_2\text{Ar}$ and R is a benzyl group, with H_2/Pd , to yield Formula VI, whereby X is N-O-H and R is a H.

The compounds of the present invention are useful in the treatment of HIV. Certain compounds of the present invention may exhibit reduced toxicity as compared with conventionally administered agents.

The methods and uses described herein can be specifically directed to inhibiting HIV integrase in a mammal in need thereof. Such methods and uses may prevent, treat or delay the onset of AIDS in a mammal in need thereof. The present invention also includes a compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof for use in, for use as a medicament for, and/or for use in the

preparation of a medicament for: inhibiting HIV integrase, preventing or treating infection by HIV, or preventing, treating or delaying the onset of AIDS.

A composition can comprise the compound(s) of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, 5 tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally in combination with a pharmaceutically acceptable carrier or diluent. The compounds of this invention may be administered, for example, to mammals, typically humans, either alone or, in combination with pharmaceutically acceptable carriers or diluents, optionally with known 10 adjuvants, such as alum, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, and subcutaneous routes of administration.

The term "administration" (e.g., "administering" a compound) in 15 reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the mammal in need of treatment. For the purpose of inhibiting HIV integrase, preventing or treating HIV infection or preventing, treating or delaying the onset of AIDS, the compounds of the present invention can be administered by any means that 20 produces contact of the active agent with the agent's site of action. They can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but typically are administered with a pharmaceutical carrier selected on the basis 25 of the chosen route of administration and standard pharmaceutical practice. The compounds of the invention can, for example, be administered orally, parenterally (including subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques), by inhalation spray, or rectally, in the form of a unit dosage of a pharmaceutical composition containing an 30 effective amount of the compound and conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. Liquid

preparations suitable for oral administration (e.g., suspensions, syrups, elixirs and the like) can be prepared according to techniques known in the art and can employ any of the usual media such as water, glycols, oils, alcohols and the like. Solid preparations suitable for oral administration (e.g., powders, pills, capsules and tablets) can be prepared according to techniques known in the art and can employ such solid excipients as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like. Parenteral compositions can be prepared according to techniques known in the art and typically employ sterile water as a carrier and optionally other ingredients, such as a solubility aid. Injectable solutions can be prepared according to methods known in the art wherein the carrier comprises a saline solution, a glucose solution or a solution containing a mixture of saline and glucose. Further description of methods suitable for use in preparing pharmaceutical compositions of the present invention and of ingredients suitable for use in said compositions is provided in Remington's Pharmaceutical Sciences, 18th edition, edited by A. R. Gennaro, Mack Publishing Co., 1990.

The compounds of this invention can be administered orally in a dosage range of 0.001 to 1000 mg/kg of mammal (e.g., human) body weight per day in a single dose or in divided doses. One preferred dosage range is 0.01 to 500 mg/kg body weight per day orally in a single dose or in divided doses. Another preferred dosage range is 0.1 to 100 mg/kg body weight per day orally in single or divided doses. For oral administration, the compositions can be provided in the form of tablets or capsules containing 1.0 to 500 milligrams of the active ingredient, particularly 1, 5, 10, 15, 20, 25, 50, 75, 100, 150, 200, 250, 300, 400, and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and

time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

The compounds of the present invention may be used with one or more agents useful in the treatment of HIV infection or AIDS. As the compounds of the present invention can be HIV integrase inhibitors, such compounds are also useful in salvage therapy for patients whose virus has mutated and acquired resistance to other drugs. Such inhibitors target a distinct step in the retroviral life cycle and therefore, may be taken in combination with other types of HIV drugs to minimize adaptation by the virus.

For example, the compounds of this invention may be effectively administered, whether at periods of pre-exposure and/or post-exposure, in combination with effective amounts of one or more HIV/AIDS antivirals, immunomodulators, anti-infectives, or vaccines useful for treating HIV infection or AIDS, such as those disclosed in Table 1 of International Patent Application No. WO 01/38332 or in the Table in International Patent Application WO 02/30930. Suitable HIV/AIDS antivirals for use in combination with the compounds of the present invention include, for example, HIV protease inhibitors (e.g., indinavir, atazanavir, lopinavir optionally with ritonavir, saquinavir, or nelfinavir), nucleoside HIV reverse transcriptase inhibitors (e.g., abacavir, lamivudine (3TC), zidovudine (AZT), or tenofovir), and non-nucleoside HIV reverse transcriptase inhibitors (e.g., efavirenz or nevirapine). It will be understood that the scope of combinations of the compounds of this invention with HIV/AIDS antivirals, immunomodulators, anti-infectives or vaccines is not limited to the foregoing substances or to the list in the above-referenced Tables, but includes in principle any combination with any pharmaceutical composition useful for the treatment of AIDS. The HIV/AIDS antivirals and other agents will typically be employed in these combinations in their conventional dosage ranges and regimens as reported in the art, including, for example, the dosages described in the Physicians' Desk Reference, 57th edition, Thomson PDR, 2003. The

dosage ranges for a compound of the invention in these combinations are the same as those set forth above.

Other embodiments of the present invention include a pharmaceutical composition comprising an effective amount of a compound of the present invention described herein and a pharmaceutically acceptable carrier and/or diluent. The composition may further comprise an effective amount of an HIV infection/AIDS treatment agent selected from the group consisting of HIV/AIDS antiviral agents, immunomodulators, and anti-infective agents. The HIV infection/AIDS treatment agent is an antiviral selected from the group consisting of HIV protease inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, and nucleoside HIV reverse transcriptase inhibitors. The HIV infection/AIDS treatment agent and the compound of the present invention can each be employed in an amount that renders the combination effective for inhibiting HIV integrase, for treating or preventing infection by HIV, or for preventing, treating or delaying the onset of AIDS.

Another embodiment includes a method for the treatment of HIV in a mammal comprising administering to the mammal an anti-HIV effective treatment amount of the compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent. There is also a method for salvage therapy in the treatment of HIV in a mammal comprising administering to said mammal an anti-HIV effective treatment amount of a compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent. The mammal is typically a human. If the compound of the present invention is co-administered with at least one other anti-HIV agent, this can be done in combination and/or in alternation with the compound.

In other embodiments, there is provided the use of an anti-HIV effective treatment amount of the compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a
5 pharmaceutically acceptable carrier or diluent, in the manufacture of a medicament for the treatment of HIV in a mammal. There is also the use of an anti-HIV effective treatment amount of the compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination
10 thereof, optionally with a pharmaceutically acceptable carrier or diluent, in the manufacture of a medicament for salvage therapy in the treatment of HIV in a mammal.

In other embodiments, there is provided the use of an anti-HIV effective treatment amount of the compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical
15 isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, for the treatment of HIV in a mammal. There is also the use of an anti-HIV effective treatment amount of the compound of the present invention described herein, and/or a
20 pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, for salvage therapy in the treatment of HIV in a mammal.

With respect to the uses of the compound of the present invention
25 described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, the mammal is typically a human. If the compound of the present invention described herein, and/or a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof is co-
30 administered with at least one other anti-HIV agent, this can be done in combination and/or in alternation with the compound.


When introducing elements disclosed herein, the articles “a”, “an”, “the”, and “said” are intended to mean that there are one or more of the elements.

The above disclosure generally describes the present invention. A more
5 complete understanding can be obtained by reference to the following specific Examples. These Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such
10 terms are intended in a descriptive sense and not for purposes of limitation.

EXAMPLES

Example 1

15 Synthesis of the compound of the following formula:




COTI-HIV1DOxime

20 is as follows:

Synthesis of Starting Materials

Oxalic acid tert-butyl ester methyl ester



5

To a solution of methyl chlorooxoacetate in TBME (1mmol/ml) was added at 0 - 5 °C tert-butanol (1.5 equiv.), followed by dropwise addition of pyridine (1.5 equiv.). The resulting mixture was stirred at room temperature for 20 h. Water (15 vol.) was added, the mixture was extracted with TBME twice. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum to give a colorless oil.


10

Synthesis of Oxalic acid tert-butyl ester methyl ester

Starting material	Conditions	Isolated Product
1 g chlorooxoacetate	1.5 eq. t-BuOH 1.5 eq. Pyridine 8 ml TBME 0° to rt 20 h	¹ HNMR conforms
20 g chlorooxoacetate	1.5 eq. t-BuOH 1.5 eq. Pyridine 160 ml TBME 0° to rt 20 h	22.7 g (87% yield) ¹ HNMR conforms
40 g chlorooxoacetate	1.5 eq. t-BuOH 1.5 eq. Pyridine 8 ml TBME 0° to rt 20 h	41.5 g (79.3% yield) ¹ HNMR conforms

Benzyloxy-acetic acid methyl ester


- 5 Benzyloxyacetyl chloride was added slowly to 10 vol of MeOH at 0 - 5 °C. The mixture was stirred at room temperature for 20 h. MeOH was removed under vacuum to give quantitatively the desired product.



Synthesis of Benzyloxy-acetic acid methyl ester

Starting material	Conditions	Isolated Product
1 g Benzyloxyacetyl chloride	10 ml MeOH	0.98 g (100%) ¹ HNMR conforms
20 g Benzyloxyacetyl chloride	200 ml MeOH	19.5 g (100% yield) ¹ HNMR conforms
25 g benzyloxyacetyl chlorid	10 ml MeOH	24.5 g (100%) ¹ HNMR conforms

Coupling of Oxalic acid tert-butyl ester methyl ester and Benzyloxyacetic acid methyl ester



- 10 Following the procedure described in U.S. Patent Publication No. 2005/0090668, a mixture of benzyloxy-acetic acid methyl ester (1 equiv.) and oxalic acid tert-butyl ester methyl ester (1.5 equiv.) in THF (12 vol.) was treated with 1.5 equiv. of LDA (prepared freshly from diisopropyl amine and n-BuLi in THF) at -78 °C for 1 h. The mixture was then warmed to room
- 15 temperature for 1 h and quenched with cold HCl (1N, 11 vol.). The mixture was extracted with EtOAc (3 x 12 vol.), the organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated to an oil. Chromatography of the crude product on silica gel (eluted with 15-60%


EtOAc/Hexane) gave 78% of the coupled product as a colorless oil. As described in the patent, the ^1H NMR is complex due to multiple enol tautomers and enol olefin configurations. TLC of the purified product showed a single spot. The results are summarized in the following table.

5

Coupling of Oxalic acid tert-butyl ester methyl ester and Benzyloxyacetic acid methyl ester

Starting material	Conditions	Isolated Product
0.9 g Benzyloxyacetic acid methyl ester 1.2 g Oxalic acid tert-butyl ester methyl ester	1.5 eq. LDA 12 vol THF -78°C 1h to rt 1 h	Crude yield 1.35 g (87%) TLC 1 major spot Used in the next step without purification
1.8 g Benzyloxyacetic acid methyl ester 2.4 g Oxalic acid tert-butyl ester methyl ester	1.5 eq. LDA 12 vol THF -78°C 1h to rt 1 h	Chromatographed on silica gel 2.4 g (78% yield) TLC 1 single spot
14.2 g Benzyloxyacetic acid methyl ester 18.9 g Oxalic acid tert-butyl ester methyl ester	1.5 eq. LDA 12 vol THF -78°C 1h to rt 1 h	Crude yield 23.8 g (98%) TLC 1 major spot 3.3 g used in the next step without purification. 20.5 g Chromatographed on silica gel to give 16 g product
14.2 g Benzyloxyacetic acid methyl ester 18.9 g Oxalic acid tert-butyl ester methyl ester	1.5 eq. LDA 12 vol THF -78°C 1h to rt 1 h	10 g of 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester isolated after chromatography
12 g Benzyloxyacetic acid methyl ester 16 g Oxalic acid tert-butyl ester methyl ester	1.5 eq. LDA 12 vol THF -78°C 1h to rt 1 h	Chromatographed on silica gel 14 g (68.3% yield) TLC 1 single spot
12 g Benzyloxyacetic acid methyl ester 16 g Oxalic acid tert-butyl ester methyl ester	1.5 eq. LDA 12 vol THF -78°C 1h to rt 1 h	Chromatographed on silica gel 14.5 g (70.7% yield) TLC 1 single spot

**Synthesis of 5-Benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid
tert-butyl ester**




- 5 Initial exploratory work on the synthesis of 5-Benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester was performed on the crude coupled product with potassium t-butoxide as base using the procedure described in U.S. Patent Publication No. 2005/0090668. A mixture of the coupled product (1 equiv.) and formamidine HCl (1.5 equiv.) in MeOH (10
- 10 vol.) was treated with 4.5 equiv. of t-BuOK (1M in t-BuOH) at 0 °C. The resulting mixture was stirred at room temperature for 20 h, saturated NH₄Cl was added, the mixture was extracted with EtOAc. The combined organic extracts were washed with brine, dried and concentrated to give a beige solid. Chromatography of the crude product gave 30% yield of a white solid. ¹HNMR
- 15 conforms except for small amount of unknown contaminant. Mass spec: m/z = 247, 303, 605 corresponds to (M-tBu)+1, M+1 and 2M+1. The reaction was repeated in larger scale with NaOMe as base to give 30% yield of the desired product and 4% of the corresponding methyl ester. Subsequently, the yield was improved to 65.4% using NaOMe as base. The results are summarized
- 20 in the following table.

Synthesis of 5-Benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester

Starting material	Conditions	Isolated Product
885 mg 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester 342 mg formamidine HCl	12.75 ml t-BuOK (1M) 8 ml MeOH 0°C 0.25h to rt 20 h	Chromatographed 250 mg (30%) ¹ HNMR conforms Mass spec: m/z = 247 (M-tBu)+1, 303 (M+1) and 605 (2M+1)
1.78 g 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester 511.8 mg formamidine HCl	3.97 ml NaOMe (25% w/w) 6 ml MeOH 0°C 0.25h to rt 20 h	Chromatographed 517 mg (30%) ¹ HNMR conforms 58 mg (4%) of the methyl ester also isolated
3.3 g crude 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester 962 mg formamidine HCl	7.4 ml NaOMe (25% w/w) 12 ml MeOH 0°C 0.25h to rt 60 h	Quench with 1 N HCl, product precipitate and filtered. Filtrate extracted with EtOAc and chromatographed. Combined product swish with 20% EtOAc and filtered to give 527 mg methyl ester
16 g purified 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester 5.02 g formamidine HCl	35 ml NaOMe (25% w/w) 60 ml MeOH 0°C 0.25h to rt 17 h	Quench with 1 N HCl, extracted with EtOAc. Concentrate to a solid, swished with 40% EtOAc and filtered: 6.45 g. Filtrate concentrated and chromatographed to give 2.57 g t-Bu ester and 0.73 g mixture of t-Bu and Me ester.
8.4 g 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester 2.92 g formamidine HCl	18.7 ml NaOMe (25% w/w) 32 ml MeOH 0°C 0.25h to rt 17 h	5 g of 5-Benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester isolated plus 3.3 g ML
14 g purified 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl	32 ml NaOMe (25% w/w) 60 ml MeOH	Quench with 1 N HCl, extracted with EtOAc. Concentrate to a solid, swished with 40% EtOAc and

ester 4.4 g formamidine HCl	0°C 0.25h to rt 17 h	filtered: 7.75 g. 56.3% yield + M.L.
14.3 g purified 2-Benzyloxy-3-hydroxy-but-2-enedioic acid 4-tert-butyl ester 1-methyl ester 4.4 g formamidine HCl	32 ml NaOMe (25% w/w) 60 ml MeOH 0°C 0.25h to rt 17 h	Quench with 1 N HCl, extracted with EtOAc. Concentrate to a solid, swished with 40% EtOAc and filtered: 9.0 g. 65.4% yield + M.L.

Synthesis of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester



Using a procedure described in U.S. Patent Publication No. 2005/0090668, a solution of 5-benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester was first treated with NaH (1.3 equiv.) in DMF followed by the addition of dimethylsulfate (1.3 equiv.) to give mainly the desired product after 2 h at room temperature. The reaction mixture was quenched with cold HCl (1N, 30 vol.) and extracted with EtOAc (3x30 vol.). The combined organic extracts were washed with dilute HCl and brine, dried and concentrated to an oil. Chromatography of the crude product (eluted with 40-60% EtOAc/hexane) gave 88% yield of the N-methylated compound (Synthon A).

Synthesis of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester

Starting material	Conditions	Isolated Product
151 mg 5-benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester 82 mg Me ₂ SO ₄	26 mg NaH (60% in oil) 2 ml DMF 0°C 0.25h to rt 1.5 h	TLC showed one major product. Crude ¹ HNMR conforms. Crude yield 173 mg
273 mg 5-benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester 148 mg Me ₂ SO ₄	47 mg NaH (60% in oil) 4 ml DMF 0°C 0.25h to rt 2 h	Crude product combined with product above and chromatographed to give 392 mg (88%) yield of desired product. ¹ HNMR conforms
3.03 g 5-benzyloxy-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester 1.64 g Me ₂ SO ₄	520 mg NaH (60% in oil) 30 ml DMF 0°C 0.25h to rt 1.5 h	Chromatography gave 2.15 g Synthon A (64.7%)


Synthesis of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid N'-[2-(4-fluoro-phenyl)-acetyl]-hydrazide

Synthon A was reacted with 1 equiv. of 1N sodium hydroxide to yield the 5-benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid.

Synthesis of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid

Starting material	Conditions	Isolated Product
50 mg of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester	1 equiv. NaOH (1N), 10 v EtOH 25 °C for 20 h	5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid isolated
1.7 g of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester	1 equiv. NaOH (1N), 10 v EtOH 25 °C for 20 h	1.34 g of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid isolated
4.8 g of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester	1 equiv. NaOH (1N), 10 v EtOH 25 °C for 20 h	3.38 g of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid isolated <u>85.6% yield.</u>
8.3 g of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid tert-butyl ester	1 equiv. NaOH (1N), 10 v EtOH 25 °C for 20 h	6.7 g of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid isolated <u>98.2% yield.</u>

Synthesis of (4-Fluoro-phenyl)-acetic acid hydrazide




- 5 (4-Fluoro-phenyl)-acetyl chloride was readily prepared from the corresponding acid by refluxing the acid in thionyl chloride for 1 h. However, the acid chloride was too reactive. Addition of hydrazine hydrate to a solution of the acid chloride gave only the dimer. Reverse addition of acid chloride to hydrazine hydrate at 0 °C gave the desired hydrazide with some dimer. The
- 10 acid chloride was then transformed to the corresponding ethyl ester. Reaction of the ethyl ester with 2 equiv. of hydrazine hydrate in refluxing EtOH gave cleanly the (4-Fluoro-phenyl)-acetic acid hydrazide in 76% yield. The results are summarized below.

Synthesis of (4-Fluoro-phenyl)-acetic acid hydrazide

Starting material	Conditions	Isolated Product
2.5 g (4-Fluoro-phenyl)-acetic acid	6 ml SOCl ₂ 80 °C for 1h, evaporate to dryness	2.78 g of (4-Fluoro-phenyl)-acetyl chloride isolated
172 mg of (4-Fluoro-phenyl)-acetyl chloride	Acid chloride added to H ₂ NNH ₂ .H ₂ O in 5 ml CH ₂ Cl ₂ at 0 °C. Stir at 0 °C for 1 h, 20 °C for 0.5 h	114 mg (4-Fluoro-phenyl)-acetic acid hydrazide isolated plus 37 mg dimer
2.4 g of (4-Fluoro-phenyl)-acetyl chloride	10 ml EtOH 20 °C for 1 h	2.55 g (4-Fluoro-phenyl)-acetic acid ethyl ester
2.55 g (4-Fluoro-phenyl)-acetic acid ethyl ester	2 equiv. of H ₂ NNH ₂ .H ₂ O, 4 v EtOH, stir at 85 °C for 5 h	1.8 g (4-Fluoro-phenyl)-acetic acid hydrazide isolated
10 g (4-Fluoro-phenyl)-acetic acid	18 ml SOCl ₂ 80 °C for 1h, evaporate to dryness	11.88 g of (4-Fluoro-phenyl)-acetyl chloride isolated
11.88 g of (4-Fluoro-phenyl)-acetyl chloride	50 ml EtOH 20 °C for 1 h	11.3 g (4-Fluoro-phenyl)-acetic acid ethyl ester
CKL-3-44 11 g (4-Fluoro-phenyl)-acetic acid ethyl ester	2 equiv. of H ₂ NNH ₂ .H ₂ O, 4 v EtOH, stir at 85 °C for 5 h	9.2 g (4-Fluoro-phenyl)-acetic acid hydrazide isolated

The carboxylic acid was transformed to the corresponding acid chloride by refluxing the latter in thionyl chloride. Reaction of the acid chloride with (4-fluoro-phenyl)-acetic acid hydrazide gave almost quantitatively the desired bishydrazide.

5




Synthesis of 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid N'-[2-(4-fluoro-phenyl)-acetyl]-hydrazide

10

Starting material	Conditions	Isolated Product
1. 100 mg 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid, 2 ml SOCl ₂	Reflux 1 h, concentrated under vacuum, used as is.	160 mg isolated, NMR MS conformed.
2. 63.8 mg (4-Fluorophenyl)-acetic acid hydrazide (1.0 equiv.), 72 mg Et ₃ N (2.0 equiv.)	Add acid chloride solution in CH ₂ Cl ₂ (1 ml) to a suspension of hydrazide in CH ₂ Cl ₂ (1 ml), at 0 °. Stir at rt for 1 h.	160 mg isolated, NMR MS conformed
1. 1.15 g 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid, 10 ml SOCl ₂	Reflux 1 h, concentrated under vacuum, used as is.	
2. 742 mg (4-Fluorophenyl)-acetic acid	Add acid chloride solution in CH ₂ Cl ₂ (10 ml) to a suspension of	1.8 g isolated (99%), NMR conformed.

hydrazide (1.0 equiv.), 892 mg Et ₃ N (2.0 equiv.)	hydrazide in CH ₂ Cl ₂ (10 ml), at 0 °. Stir at rt for 1 h.	
1. 3.3 g 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid, 30 ml SOCl ₂ 2. 2.13 g (4-Fluorophenyl)-acetic acid hydrazide (1.0 equiv.), 3.5 mL Et ₃ N (2.0 equiv.)	Reflux 1 h, concentrated under vacuum, used as is. Add acid chloride solution in CH ₂ Cl ₂ (10 ml) to a suspension of hydrazide in CH ₂ Cl ₂ (10 ml), at 0 °. Stir at rt for over night.	4.97 g isolated <u>yield 95.6%</u> NMR conformed.
1. 6.7 g 5-Benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid , 30 ml SOCl ₂ 2. 4.33 g (4-Fluorophenyl)-acetic acid hydrazide (1.0 equiv.), 3.5 mL Et ₃ N (2.0 equiv.)	Reflux 1 h, concentrated under vacuum, used as is. Add acid chloride solution in CH ₂ Cl ₂ (10 ml) to a suspension of hydrazide in CH ₂ Cl ₂ (10 ml), at 0 °. Stir at rt for over night.	9.44 g isolated <u>yield 89.3%</u> NMR conformed.

Synthesis of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one



5

Following the procedure described in European Patent No. 1698628, 5-benzyloxy-1-methyl-6-oxo-1,6-dihydro-pyrimidine-4-carboxylic acid N'-[2-(4-fluoro-phenyl)-acetyl]-hydrazide was treated with CCl_4 (7 equiv.), Ph_3P (3 equiv.) and Et_3N (3 equiv.) in CH_3CN (10 vol) at room temperature for 1.5 h.

10 TLC showed only one product formed. The reaction mixture was diluted with EtOAc , washed with aqueous NaHCO_3 and brine, dried and concentrated to give a crude solid. For the 1.8 g scale reaction, chromatography (4 columns) plus recrystallization twice was needed to isolate the desired product without $\text{Ph}_3\text{P}=\text{O}$ contamination. NMR and MS confirmed the structure.

15

Starting material	Conditions	Isolated Product
160 mg bis hydrazide 0.262 ml CCl ₄ (7 eq.) 0.164 mL Et ₃ N (3 eq.) 306 mg Ph ₃ P (3 eq.)	Ph ₃ P was added to a suspension of bis hydrazide, CCl ₄ and Et ₃ N at rt. Stir for 1 h at rt.	40 mg of cyclised oxadiazole contaminated by Ph ₃ P isolated after chromatography (eluted with 100% EtOAc). NMR, MS conformed.
1.8 g bis hydrazide 3 ml CCl ₄ (7 eq.) 1.8 mL Et ₃ N (3 eq.) 3.45 g Ph ₃ P (3 eq.)	Ph ₃ P was added to a suspension of bis hydrazide, CCl ₄ and Et ₃ N at rt. Stir for 1.5 h at rt.	First column chromatography gave 1.91 g of desired product contaminated by small amount of Ph ₃ P=O and 3.5 g of Ph ₃ P=O containing small amount of product. The Ph ₃ P=O fraction was repurified by chromatography to give more desired product. All the impure fractions were combined and chromatographed, the product isolated was first crystallized from EtOAc and then recrystallization in toluene gave a combined yield of 1.3 g (75%) of product.
4.95 g bis hydrazide 8.13 ml CCl ₄ (7 eq.) 5.04 mL Et ₃ N (3 eq.) 9.49 g Ph ₃ P (3 eq.)	Ph ₃ P was added to a suspension of bis hydrazide, CCl ₄ and Et ₃ N at rt. Stir over night at rt.	First column chromatography gave 1.6 g of desired product contaminated by small amount of Ph ₃ P=O and 5.2 g of desired product with Ph ₃ P=O. Recrystallisation of 1.6 g with toluene gives 1.3 g of pure product.


9.4 g bis hydrazide 8.13 ml CCl ₄ (7 eq.) 5.04 mL Et ₃ N (3 eq.) 9.49 g Ph ₃ P (3 eq.)	Ph ₃ P was added to a suspension of bis hydrazide, CCl ₄ and Et ₃ N at rt. Stir over night at rt.	First column chromatography gave 5.0 g of desired product contaminated by small amount of Ph ₃ P=O and 7.5 g of desired product with Ph ₃ P=O. <u>Recrystallisation of 5.0g with toluene gives 4.1 g of pure product.</u>
---	--	--

Synthesis of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione

5


5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one was treated with 0.62 equiv of Lawesson's reagent at 80-85 °C in toluene for 5-7 h. TLC showed one major product formation with some substrate still present. Chromatography (eluted with 70%-90% EtOAc/hexane)

10 gave ~50% yield of the title compound.



Starting material	Conditions	Isolated Product
100 mg 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 63 mg Lawesson's reagent (0.62 eq.)	3 ml toluene, 80-85 °C for 5 h	50 mg of a yellow powder. NMR, MS conformed.
392 mg 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 63 mg Lawesson's reagent (0.62 eq.)	3 ml toluene, 80-85 °C for 5 h	50 mg of a yellow powder. NMR, MS conformed.
350 mg 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 233 mg Lawesson's reagent (0.65 eq.)	10 ml toluene, 80-85 °C for 7 h	269 mg (74%) of a yellow powder
190 mg 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 126 mg Lawesson's reagent (0.65 eq.)	6 ml toluene, 80-85 °C for 14 h	150 mg (76%) of a yellow powder. NMR, MS conformed.
2.0g 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 1.28 g Lawesson's reagent (0.62 eq.)	55 ml toluene, 80-85 °C for 8 h, r.t. for 16 hrs.	1.2 g (57.7%) of a yellow powder. NMR, MS conformed.
3.75g 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 2.51 g Lawesson's reagent (0.62 eq.)	94 ml toluene, 80-85 °C for 8 h, r.t. for 16 hrs.	2.75 g (70.5%) of a yellow powder. NMR, MS conformed.

Reaction of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione with N, O-bis(trimethylsilyl)hydroxylamine in the presence of Hg(OAc)₂.



5

Reaction of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione with N, O-bis(trimethylsilyl)hydroxylamine in the presence of $\text{Hg}(\text{OAc})_2$, the desired amidoxime derivative was isolated as a mixture of syn- and anti-isomers. Mass spec: $m/z=408$ ($M+1$) conforms to structure expected.

10

Starting material	Conditions	Isolated Product
50 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione (75% pure) 83 mg N,O-bis(trimethylsilyl)hydroxylamine (5 equiv.) 40 mg Hg(Oac) ₂ 1.4 equiv.	2.5 ml dioxane, 4h rt	Small amount of expected product as mixture of isomers. MS: m/z = 408 (M+1)
400 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione (pure) 1.05 mL N,O-bis(trimethylsilyl)hydroxylamine (5 equiv.) 400 mg Hg(Oac) ₂ 1.4 equiv.	20 ml dioxane, 3h rt	Small amount of expected product. MS: m/z = 408 (M+1) ~20 mg of pure product
400 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione (pure) 2.0 mL N,O-bis(trimethylsilyl)hydroxylamine (10 equiv.) 400 mg Hg(Oac) ₂ 1.4 equiv.	20 ml dioxane, 3h rt	Small amount of expected product. MS: m/z = 408 (M+1) ~10 mg of pure product
200 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione (pure) 0.5 mL N,O-bis(trimethylsilyl)hydroxylamine (5 equiv.) 200 mg Hg(Oac) ₂ 1.4 equiv.	20 ml toluene, 2h, 90°C. same TLC	n/a
200 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione (pure) 0.5 mL N,O-bis(trimethylsilyl)hydroxylamine (5 equiv.) 200 mg Hg(Oac) ₂ 1.4 equiv.	20 ml THF, 2h, r.t. same TLC	Purification of above and this one give ~50 mg of good product.
2.48 g 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidine-4-thione (pure)	125 ml Dioxane, 1h, r.t. same TLC	Purification by silica gives 190 mg of desire product. (7.66%)


6.48 mL N,O-bis(trimethylsilyl)hydroxylamine (5 equiv.) 2.66 g Hg(OAc) ₂ 1.4 equiv.		yield).
---	--	---------

6-[5-(4-Fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H-pyrimidin-4-one oxime (COTI-HIV-1DOxime)

5

5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one oxime was hydrogenated at 1 atmosphere of hydrogen in the presence of equal weight of 10% Pd/C to give 68% yield of 6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H-pyrimidin-4-one oxime (COTI-HIV-1DOxime or, also referred to as COTI-HIV-1). Results are summarized below.

10




Synthesis of 6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H-pyrimidin-4-one oxime (COTI-HIV-1DOxime)

Starting material	Conditions	Isolated Product
~ 50 to 100 mg of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one oxime (+/- pure) 100% w/w Pd/C 10% and 50% water.	25 ml THF, rt 1 atm of H ₂ .	28 mg COTI-HIV-1 + by-product 56 mg of +/- pure COTI-HIV-1DOxime and by product
190 mg of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one oxime (pure) 100% w/w Pd/C 10% and 50% water.	25 ml THF, rt 1 atm of H ₂ .	The crude is purifying by silica with the above products to give 100 mg (68%) of pure desire product. (M/Z = 318) NMR ¹ H conform

Example 1A

- 5 Synthesis of the compound of the following formula:




COTI-HIV1Doxo

is as follows:

5 **6-[5-(4-Fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H**
pyrimidin-4-one

a) Preparation of 6-[5-(4-Fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-
methyl-3H-pyrimidin-4-one by acid catalyzed debenzylation

10




Reaction of 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-
3H-pyrimidin-4-one and a 1:1 mixture of TFA/CH₂Cl₂ gave the debenzylated
15 product in reasonable yield. Results are summarized below.

Preparation of 6-[5-(4-Fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H-pyrimidin-4-one

Starting material	Conditions	Isolated Product
24 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one	0.3 ml CH ₂ Cl ₂ , 0.3 ml TFA, 20 h, rt. Concentrate under vacuum, swished with 20% EtOAc/hexane	Beige solid, ¹ HNMR and MS conformed to expected product
109 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one	1 ml CH ₂ Cl ₂ , 1 ml TFA, 48 h, rt. Concentrate under vacuum, co-distill with EtOAc/heptane, swished with 30% EtOAc/hexane	65 mg beige solid (77%), ¹ HNMR conformed.
100 mg 5-benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one	1 ml CH ₂ Cl ₂ , 1 ml TFA, 48 h, rt. Concentrate under vacuum, co-distill with EtOAc/heptane, swished with 30% EtOAc/hexane for 20 h	69 mg beige solid , ¹ HNMR showed an unknown impurity. Re-purify by swishing in 4 ml pure EtOAc for 20 h to give 56 mg of pure product. ¹ HNMR conformed.

5 b) Preparation of 6-[5-(4-Fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H-pyrimidin-4-one by palladium catalyzed debenzylation



Hydrogenation of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one in THF with 25% w/w of 10% Pd/C gave 67.5% of the desired product. Results are shown below.

Preparation of 6-[5-(4-Fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-5-hydroxy-3-methyl-3H-pyrimidin-4-one by palladium catalyzed debenzylation


Starting material	Conditions	Isolated Product
1.0 g of 5-Benzyloxy-6-[5-(4-fluoro-benzyl)-[1,3,4]oxadiazol-2-yl]-3-methyl-3H-pyrimidin-4-one 250 mg mg 10% Pd/C	200 ml THF, r.t. 1 atm 30 mins	0.52 g of pure product (67.5%) 0.35g of +/- pure product.

10

Example 2

Synthesis of COTI-HIV28236

Synthesis of the compound of the following formula:




15

COTI-HIV28236

was conducted according to the following synthetic methodology:

20

Synthesis of the common intermediate 4c:**Synthesis of Compound 63:**

- 5 To a suspension of lithium 3-fluoropyridine-2-carboxylate (10.1 g ; 69 mmol ; 1 eq.) in methanol (170 mL) were successively added 1,2-dimethoxypropane (17.0 mL ; 138 mmol ; 2 eq.) and concentrated sulphuric acid (18.0 mL ; 345 mmol ; 5 eq.). The reaction mixture became homogeneous and turned yellow after addition of sulphuric acid. The mixture
- 10 was then heated at reflux for 20 hours. After cooling down to room temperature, the solvent was removed under vacuum. The oily residue was slowly poured into a mixture of ethyl acetate (200 mL) and a saturated solution of NaHCO_3 (400 mL). At the end of the addition, the aqueous layer was again saturated by addition of solid NaHCO_3 . The layers were then
- 15 separated and the aqueous phase was extracted with ethyl acetate (2×200 mL). The combined organic phases were dried over Na_2SO_4 , filtered and evaporated to dryness to afford compound **63** as a light yellow solid (9.5 g ; yield = 89%).

Synthesis of Compound 66:

- 20 To a solution of LDA (1.8M in THF /heptane ; 72.0 mL ; 129 mmol ; 2.1 eq.) in THF (60 mL) at -78°C was added *tert*-butylacetate (20.0 mL ; 148 mmol ; 2.4 eq.). The deep red solution was stirred at -78°C for 45 minutes and maintained at that temperature during the slow addition of a solution of

compound **63** (9.5 g ; 62 mmol ; 1 eq.) in THF (60 mL). The reaction mixture was then stirred for an additional 2 hours and quenched at -78°C by addition of MTBE (150 mL) and water/acetic acid (1:1 ; 150 mL). After warming up to room temperature, the layers were separated and the aqueous phase was
5 extracted with MTBE (2 × 100 mL). The combined organic phases were washed with water (300 mL) and brine (300 mL), dried over Na₂SO₄, filtered and evaporated to dryness to afford compound **66** as an yellow oil (16.2 g ; contaminated with 8.5% ^w/_w of acetic acid ; yield = 100%).

Synthesis of Compound 67:

10 To a solution of crude compound **66** (16.2 g ; 62 mmol ; 1 eq.) in acetonitrile (190 mL) at 0°C was successively added triethylamine (14.6 mL ; 112 mmol ; 1.8 eq.) and ABSA (16.3 g ; 68 mmol ; 1.1 eq.). At the end of the addition, the reaction mixture was allowed to warm up to room temperature. A white solid precipitated progressively from the orange solution. After 18
15 hours, the reaction mixture was filtered on a Büchner funnel. The white cake was washed with ethyl acetate (450 mL). The mother liquor was decanted and the layers were separated. The organic phase was washed with a saturated solution of NH₄Cl (400 mL) and brine (400 mL), dried over Na₂SO₄, filtered and evaporated to dryness. The orange oily residue was triturated in diethyl
20 ether. After filtration of the brown solid, the evaporation of the mother liquor afforded compound **67** as an orange oil (18.9 g contaminated with 9% w/w of *p*-AcHN-Ph-SO₂NH₂, yield = 100%).

Synthesis of Compound 68:

To a solution of crude compound **67** (18.9 g ; 62 mmol ; 1 eq.) in THF
25 (125 mL) at room temperature was slowly added trimethylphosphine (1M/THF ; 68.0 mL ; 68 mmol ; 1.1 eq.). At the end of the addition, the reaction mixture was stirred for 15 minutes and quenched by addition of water (11.0 mL ; 616 mmol ; 10 eq.). The mixture was stirred for an additional 15 minutes prior to the addition of silica gel (45 g). The solvent was then evaporated under
30 vacuum and the residue, which was adsorbed on silica gel, was purified by


flash chromatography on silica gel (eluent: dichloromethane/methanol, 100/0 to 96/4) to afford compound **68** as an orange oil (15.7 g ; yield = 96%).

Synthesis of Compound 4d:

To a solution of compound **68** (15.7 g ; 59 mmol ; 1 eq.) in acetonitrile
5 (300 mL) at room temperature were successively added DBU (18.5 mL ; 124 mmol ; 2.1 eq.) and methyl iodide (3.8 mL ; 62 mmol ; 1.05 eq.). The reaction mixture became deep red. After stirring for 18 hours at room temperature, the reaction was quenched by addition of a saturated solution of NH₄Cl (500 mL). The resulting mixture was extracted with dichloromethane (3 × 500 mL) and
10 the combined organic phases were washed with brine (1 L), dried over Na₂SO₄, filtered and evaporated to dryness. The purification of the residue by chromatography on silica gel (eluent: dichloromethane/methanol, 100/0 to 90/10) afforded two fractions containing compound **4d**. After evaporation to dryness, the first one was triturated in diethyl ether to afford a first batch of
15 pure compound **4d** as a pale orange solid (2.6 g ; yield = 17%). The second one was dissolved in dichloromethane (500 mL) and the resulting solution was washed with a saturated solution of NH₄Cl (2 × 500 mL), dried over Na₂SO₄, filtered and evaporated to dryness to afford a second batch of pure compound **4d** as a pale orange solid (3.1 g ; yield = 20%). The two batches
20 were combined (yield = 37%).

Synthesis of Compound 4c:

To a solution of compound **4d** (3.1 g ; 12 mmol ; 1 eq.) in dichloromethane (110 mL) at room temperature was added trifluoroacetic acid
25 (12.0 mL ; 157 mmol ; 13.3 eq.). The reaction mixture turned brown. After stirring at room temperature for 18 hours, the reaction mixture was diluted by addition of diethyl ether (150 mL). The solvent were then removed under vacuum. The beige residue was triturated in diethyl ether to afford compound **4c** as a beige solid (2.4 g ; yield = 100%). ¹H NMR (DMSO, ppm): 4.29 (s,
30 3H); 8.02 (m, 1H); 8.52 (m, 1H); 9.01 (m, 1H); and 14.32 (m, COOH).

Synthesis of COTI-HIV28236:**5 Synthesis of Compound 76:**

To a solution of compound **4c** (907 mg ; 4.4 mmol ; 1 eq.) in DMSO (6 mL) at room temperature was successively added triethylamine (2.4 mL ; 17.6 mmol ; 4 eq.), *para*-fluorobenzylamine (610 μ L ; 5.3 mmol ; 1.2 eq.) and TBTU (1.70 g ; 5.3 mmol ; 1.2 eq.). After stirring the mixture at room temperature for 16 hours, the reaction was quenched by addition of a saturated solution of NH₄Cl (50 mL). The aqueous phase was then extracted with dichloromethane (5 $\times$ 100 mL). The combined organic phases were combined, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was dissolved in ethyl acetate (100 mL) and the solution was washed with water (100 mL), brine (100 mL), dried over Na₂SO₄, filtered and evaporated to dryness. The residue was then triturated in diethyl ether to afford compound **76** as an yellow solid (480 mg ; yield = 35%). All aqueous layers were then combined and extracted with chloroform (3 $\times$ 250 mL). The combined chloroform phases were dried over Na₂SO₄, filtered and evaporated to dryness in the presence of silica gel (3 g). The purification by chromatography on silica gel (eluent: dichloromethane/methanol, 98/2 to 90/10) afforded compound **76** as an yellow solid (454 mg ; yield = 33%). The two batches were combined (yield = 68%).

Synthesis of Compound COTI-HIV28236:


To a suspension of compound **76** (228 mg ; 73 μ mol; 1 eq.) in toluene (3.7 mL) at room temperature was added hydroxylamine hydrochloride (76 mg ; 1.10 mmol ; 1.5 eq.) and titanium (IV) isopropoxide (330 μ L; 1.10 mmol ;

1.5 eq.). The reaction mixture was refluxed for 72 hours and evaporated to dryness. The red residue is dissolved in a mixture of a saturated solution of NaHCO₃/ethyl acetate/methanol (5:5:1 ; 55 mL). After separation of the layers, the aqueous phase was extracted with ethyl acetate (3 × 50 mL). The
 5 combined organic phases were washed with brine (100 mL), dried over Na₂SO₄, filtered and evaporated to dryness. The purification of the residue on silica gel (eluent: dichloromethane/methanol, 100/0 to 98/2) followed by a trituration in methanol afforded compound **COTI-HIV28236** as a yellow solid (42 mg ; yield = 17%). ¹H NMR (CDCl₃, ppm): 3.82 (s, 3H); 4.61 (s, 2H); 6.99
 10 (m, 2H); 7.34 (m, 1H); 7.44 (m, 1H), 7.55 (m, 1H), 8.24 (m, 1H), 9.82 (m, 1H) and 15.71 (m, NOH). ¹³C NMR (CDCl₃, ppm): 42.3, 43.3, 115.3, 115.6, 121.4, 127.5, 129.7, 129.8, 134.3, 135.8, 138.1, 138.7, 139.7, 160.9, 161.6, and 163.4. MS (ES + IC): m/z (M+H)⁺ 328.

15 Example 3

Synthesis of COTI-HIV28233


Synthesis of the compound of the following formula:



20

COTI-HIV28233 (COTI-HIV1oxime)

Synthesis of the compound of COTI-HIV28233 was conducted according to the following synthetic methodology. Synthesis of the common intermediate **4c** is as described above with respect to Example 2.

Synthesis of COTI-HIV28233:5 Synthesis of compound 78:

To a solution of compound **77** (5.0 g ; 33 mmol ; 1 eq.) in dichloromethane (100 mL) at room temperature was successively added triethylamine (13.0 mL ; 100 mmol ; 3 eq.), *tert*-butylcarbazate (4.8 g ; 36 mmol ; 1.1 eq.), EDCI (6.9 g ; 36 mmol ; 1.1 eq.) and HOBt (445 mg ; 3.3 mmol ; 0.1 eq.). After stirring the mixture for 18 hours at room temperature, the reaction was quenched by addition of a saturated solution of NH_4Cl (200 mL). The layers were separated and the aqueous phase was extracted with dichloromethane (3 $\times$ 200mL). The combined organic phases were dried over Na_2SO_4 , filtered and evaporated to dryness to afford compound **78** as a colourless oil (9.1 g ; yield = 100%).

Synthesis of Compound 79:

To a solution of compound **78** (9.1 g ; 33 mmol ; 1 eq.) in dichloromethane (300 mL) at room temperature was slowly added trifluoroacetic acid (33.0 mL ; 434 mmol ; 13.2 eq.). The reaction mixture turned light brown. After stirring for 3 hours, the reaction mixture was
5 evaporated to dryness. The colourless residue was dissolved in ethyl acetate (400 mL) and the solution was washed with a saturated solution of NaHCO₃ (3 × 500 mL), dried over Na₂SO₄, filtered and evaporated to dryness. The resulting white foam was triturated in diethyl ether to afford compound **79** as a white solid (3.2 g ; yield = 57%).

10 Synthesis of Compound 80:

To a solution of compound **4c** (1.5 g ; 7.3 mmol ; 1 eq.) in DMSO (11 mL) at room temperature was successively added triethylamine (3.8 mL ; 29 mmol ; 4 eq.), compound **79** (1.5 g ; 8.8 mmol ; 1.2 eq.) and TBTU (2.8 g ; 8.8 mmol ; 1.2 eq.). The reaction mixture became homogeneous for a few
15 minutes prior to the precipitation of a solid. After stirring for 16 hours at room temperature, the precipitate was filtered on a Büchner funnel. The cake was washed with ethyl acetate and dried to afford compound **80** as a yellow solid (1.7 g ; yield = 64%).

Synthesis of compound 81:

20 To a suspension of compound **80** (1.7 g ; 4.7 mmol ; 1 eq.) in acetonitrile (33 mL) at room temperature was successively added N,N-diisopropylethylamine (4.7 mL ; 28.2 mmol ; 6 eq.), triphenylphosphine (2.2 g ; 8.5 mmol ; 1.8 eq.) and hexachloroethane (1.7 g ; 7.0 mmol ; 1.5 eq.). The reaction mixture became homogeneous for few minutes prior to the
25 precipitation of a solid. After stirring for 16 hours at room temperature, the precipitate was filtered on a Büchner funnel. The cake was washed with diethyl ether to afford compound **81** as a beige solid (1.1 g ; yield = 71%).


Synthesis of compound COTI-HIV28233:

To a suspension of compound **81** (236 mg ; 700 μ mol ; 1 eq.) in toluene (3.5 mL) at room temperature was added hydroxylamine hydrochloride (73 mg ; 1.05 mmol ; 1.5 eq.) and titanium (IV) isopropoxide (310 μ L ; 1.10 mmol ; 1.5 eq.). The reaction mixture was refluxed for 18 hours and evaporated to dryness. The red residue was dissolved in a mixture of a saturated solution of NaHCO₃/ethyl acetate/methanol (5:5:1 ; 55 mL). After separation of the layers, the aqueous phase was extracted with ethyl acetate (3 $\times$ 50 mL). The combined organic phases were washed with brine (100 mL), dried over Na₂SO₄, filtered and evaporated to dryness. The purification of the residue on silica gel (eluent: dichloromethane/methanol, 100/0 to 98/2) followed by a trituration in methanol afforded compound **COTI-HIV28233** as an yellow solid (39 mg ; yield = 16%). ¹H NMR (CDCl₃, ppm): 3.71 (s, 3H); 4.27 (s, 2H); 7.00 (m, 2H); 7.35 (m, 3H); 7.52 (m, 1H), 8.22 (m, 1H), 15.23 (m, NOH). MS (ES + IC): m/z (M+H)⁺353.

Example 4

Synthesis of COTI-HIV28236-SA


Synthesis of the compound of COTI-HIV28236-SA can be conducted according to the following synthetic methodology:




Example 5

Synthesis of COTI-HIV28236-SeA


Synthesis of the compound of COTI-HIV28236-SeA can be conducted according to the following synthetic methodology:




Strategy A:



Strategy B:



Strategy C:



5

COTI-HIV28236-SeA

Example 6: Computational Assessment of Properties

A computational assessment of the properties of compounds according to the present invention was performed using the CHEMSAS[®] computational platform. CHEMSAS[®] is a robust proprietary computational platform for accelerated drug discovery, optimization and lead selection based upon a unique combination of traditional and modern pharmacology principles, statistical modeling and machine learning technologies. At the centre of the CHEMSAS[®] platform is a hybrid machine learning technology that may be used to: find, profile and optimize new targeted lead compounds; find novel uses for known compounds; and, solve problems with existing or potential

drugs. In using the CHEMSAS[®] platform, first a therapeutic target was selected, in this case HIV and more particularly the critical enzyme HIV integrase. The second step involved the design of a candidate molecule library containing thousands of potential compounds through the assembly of privileged molecular fragments. Thirdly, the candidate library was profiled and optimized using a combination of validated computational models and traditional expert medicinal chemistry. In this step, the CHEMSAS[®] platform developed about 315 molecular descriptors for each candidate therapeutic compound. For example, molecular properties relating to a candidate compound's therapeutic efficacy, expected human toxicity, oral absorption, cumulative cellular resistance and/or kinetics were assessed. In some instances, comparative properties relating to commercially relevant benchmark compounds were also assessed. Potential lead compounds were then selected from the candidate library using a proprietary decision making tool designed to identify candidates with the optimal physical chemical properties, efficacy, ADME/Toxicity profile, etc. according to a pre-determined set of design criteria. The lead compounds selected from the candidate library were then synthesized for further pre-clinical development.

The properties of certain compounds according to the present invention, specifically, those indicated in Tables 1-8 that were assessed using the CHEMSAS[®] computational platform are shown below. Some of the predicted properties are validated by the experimental data provided herein, while other properties have been validated elsewhere during the development of other clinical candidates. The CHEMSAS[®] platform therefore provides a means of determining, predicting and/or testing the properties of a compound, particularly when used to determine the properties of compounds according to the present invention. The CHEMSAS[®] platform is also particularly useful in comparing the properties of compounds according to the invention with prior art compounds on a relative basis *in silico*.

Tables 1A and 1B: Physical Chemical Properties

Tables 1A and 1B shows that the compounds are all “drug-like” with good drug like physical properties.

Table 1A

MolID	Formula	MolWt	ALogP	HBndAcc
COTI-HIV1oxo	C17H12FN5O2	337.31	2.45	7
COTI-HIV1thio	C17H12FN5OS	353.38	3.35	6
COTI-HIV1Se	C17H12FN5OSe	400.27	2.48	6
COTI-HIV1imine	C17H13FN6O	336.33	2.51	7
COTI-HIV1Oxime	C17H13FN6O2	352.33	2.54	8
COTI-HIV2oxo	C16H13FN4O2	312.30	1.98	6
COTI-HIV28236-SA	C16H13FN4OS	328.37	2.88	5
COTI-HIV28236-SeA	C16H13FN4OSe	375.26	2.00	5
COTI-HIV2imine	C16H14FN5O	311.32	2.03	6
COTI-HIV28236	C16H14FN5O2	327.32	2.06	7
COTI-HIV1DOxime	C14H12FN5O3	317.28	1.15	8
COTI-HIV1Doxo	C14H11FN4O3	302.26	0.54	7
COTI-HIV1Dthio	C14H11FN4O2S	318.33	1.96	6
COTI-HIV1Dcyano	C15H11FN6O2	326.29	1.35	8
COTI-HIV1Dimino	C14H12FN5O2	301.28	1.12	7
COTI-HIV2DOxime	C13H13FN4O3	292.27	0.68	7
COTI-HIV2Doxo	C13H12FN3O3	277.25	0.07	6
COTI-HIV2Dthio	C13H12FN3O2S	293.32	1.49	5
COTI-HIV2Dcyano	C14H12FN5O2	301.28	0.89	7
COTI-HIV2Dimino	C13H13FN4O2	276.27	0.65	6

Table 1B

MolID	Formula	HBndDon	RotBnds	TPSA
COTI-HIV1oxo	C17H12FN5O2	0	3	78.31
COTI-HIV1thio	C17H12FN5OS	0	3	61.01
COTI-HIV1Se	C17H12FN5OSe	0	3	61.01
COTI-HIV1imine	C17H13FN6O	1	3	84.10
COTI-HIV1Oxime	C17H13FN6O2	1	4	93.15
COTI-HIV2oxo	C16H13FN4O2	1	4	73.35
COTI-HIV28236-SA	C16H13FN4OS	1	4	56.05
COTI-HIV28236-SeA	C16H13FN4OSe	1	4	56.05
COTI-HIV2imine	C16H14FN5O	2	4	79.14
COTI-HIV28236	C16H14FN5O2	2	5	88.19
COTI-HIV1DOxime	C14H12FN5O3	2	5	102.71
COTI-HIV1Doxo	C14H11FN4O3	1	4	87.87
COTI-HIV1Dthio	C14H11FN4O2S	1	4	70.57
COTI-HIV1Dcyano	C15H11FN6O2	1	5	104.83
COTI-HIV1Dimino	C14H12FN5O2	2	4	93.66
COTI-HIV2DOxime	C13H13FN4O3	3	6	97.74
COTI-HIV2Doxo	C13H12FN3O3	2	5	82.90
COTI-HIV2Dthio	C13H12FN3O2S	2	5	65.60
COTI-HIV2Dcyano	C14H12FN5O2	2	6	99.86
COTI-HIV2Dimino	C13H13FN4O2	3	5	88.69

Legend for Table 1:

- 5 MolWt stands for Molecular Weight measured in Daltons and is a size descriptor;
- ALogP is calculated lipophilicity/solubility estimates;
- HBndDon stands for Hydrogen Bond Donor and refers to the number of atoms able to donate electrons to potentially form Hydrogen bonds;

HBndAcc stands for Hydrogen Bond Acceptor and refers to the number of atoms able to accept electrons to potentially form Hydrogen bonds;

TPSA stands for Topological Polar Surface Area and is a measure of Molecular Surface Charge/Polarity; and

- 5 RotBnds stands for Rotatable Bonds which is a count of freely rotatable single bonds in the molecule.

Table 2: Solubility Properties

10 Table 2 shows that all of the compounds are expected to have acceptable solubility values for drug-like compounds.

Table 2A

MolID	Formula	LogD(pH 7.4)	LogSw
COTI-HIV1oxo	C17H12FN5O2	2.52	-3.94
COTI-HIV1thio	C17H12FN5OS	2.60	-4.35
COTI-HIV1Se	C17H12FN5OSe	2.15	-4.22
COTI-HIV1imine	C17H13FN6O	1.79	-3.89
COTI-HIV1Oxime	C17H13FN6O2	2.03	-3.94
COTI-HIV2oxo	C16H13FN4O2	2.66	-3.46
COTI-HIV28236-SA	C16H13FN4OS	2.87	-3.65
COTI-HIV28236-SeA	C16H13FN4OSe	1.97	-3.55
COTI-HIV2imine	C16H14FN5O	2.00	-3.52
COTI-HIV28236	C16H14FN5O2	1.93	-3.42
COTI-HIV1DOxime	C14H12FN5O3	1.47	-3.44
COTI-HIV1Doxo	C14H11FN4O3	1.57	-3.19
COTI-HIV1Dthio	C14H11FN4O2S	1.84	-3.75
COTI-HIV1Dcyano	C15H11FN6O2	1.58	-3.71
COTI-HIV1Dimino	C14H12FN5O2	0.15	-3.36
COTI-HIV2DOxime	C13H13FN4O3	1.76	-2.63
COTI-HIV2Doxo	C13H12FN3O3	0.54	-2.45
COTI-HIV2Dthio	C13H12FN3O2S	1.17	-3.04

COTI-HIV2Dcyano	C14H12FN5O2	1.88	-2.73
COTI-HIV2Dimino	C13H13FN4O2	-1.05	-2.59


Legend for Table 2:

LogD(7.4) is a measure of relative solubility in octanol vs water at a specific pH, in this case pH= 7.4; and

- 5 LogSw is the logarithm of the calculated solubility in pure water usually measured at 25°C.


Table 3: Efficacy (LogIC50)

- Referring to Table 3, compounds with a high value for Prob_LogIC50<-6 (meaning a value that is greater than 0.70, preferably greater than 0.80, more preferably greater than 0.90) are predicted to have a high likelihood of sub-micromolar activity against HIV Integrase. Similarly, Prob_LogIC50<-7 represents the probability that the IC50 is less than or equal to 100 nM. *In-silico* comparative data is also shown for compounds described in PCT
- 15 Publication No. WO2003/062204: Compound A; PCT Publication No. WO 2004/101512: Compound B; and Laquinimod™. These compounds are as follows:




Compound A

5



Compound B

10



15

LaquinimodTM**Table 3**

MolID	Formula	Prob_LogIC50<-6	Prob_LogIC50<-7
COTI-HIV1oxo	C17H12FN5O2	0.911	0.000
COTI-HIV1thio	C17H12FN5OS	0.202	0.000
COTI-HIV1Se	C17H12FN5OSe	0.000	0.000
COTI-HIV1imine	C17H13FN6O	0.944	0.934
COTI-HIV1Oxime	C17H13FN6O2	0.935	0.912
COTI-HIV2oxo	C16H13FN4O2	0.986	0.000
COTI-HIV28236-SA	C16H13FN4OS	0.018	0.000
COTI-HIV28236-SeA	C16H13FN4OSe	0.000	0.000
COTI-HIV2imine	C16H14FN5O	0.938	0.000
COTI-HIV28236	C16H14FN5O2	0.965	0.973
COTI-HIV1DOxime	C14H12FN5O3	0.989	0.947
COTI-HIV1Doxo	C14H11FN4O3	0.996	0.957
COTI-HIV1Dthio	C14H11FN4O2S	0.776	0.000

COTI-HIV1Dcyano	C15H11FN6O2	0.829	0.011
COTI-HIV1Dimino	C14H12FN5O2	0.979	0.895
COTI-HIV2DOxime	C13H13FN4O3	0.996	0.581
COTI-HIV2Doxo	C13H12FN3O3	0.995	0.994
COTI-HIV2Dthio	C13H12FN3O2S	0.203	0.000
COTI-HIV2Dcyano	C14H12FN5O2	0.980	0.001
COTI-HIV2Dimino	C13H13FN4O2	0.985	0.955
Compound A	C17H14N3O3	0.999	0.996
Compound B	C18H13FN4O2	0.946	0.076
Laquinimod TM	C18H17N3O3	0.897	0.000

Legend for Table 3:

- Prob_LogIC₅₀<-6 is the probability that the IC₅₀ value of the compound (i.e. the amount required to reduce the initial concentration by 50%) will be less than 10⁻⁶ (i.e. sub-micromolar activity against the chosen target).
- Prob_LogIC₅₀<-7 is the probability that the IC₅₀ value of the compound (i.e. the amount required to reduce the initial concentration by 50%) will be less than 10⁻⁷ (i.e. less than or equal to 100 nM activity against the chosen target).

10

Tables 4A and 4B: Oral Absorption and BBB penetration

- Referring to Tables 4A and 4B, compounds with a high P_{OrlAvail}>30% are likely to have acceptable oral absorption. Compounds with a high P_{Clr}<30ml/min/kg are expected to have an acceptable clearance rate and are therefore expected to remain in circulation longer than those with a low P_{Clr}<30ml/min/kg. All compounds were predicted to be acceptable in both attributes. EST_PPb% is the estimated percentage plasma protein binding of the compound. Estimates of greater than 95% plasma protein binding would be considered to be highly protein bound. P_{BBPenetration} is an estimate of the probability that the drug will penetrate the blood brain barrier and enter the central nervous system (CNS). Predicted values of greater than 0.7 P_{BBPenetration}

20

indicate that the drug is likely to pass the blood brain barrier and enter the central nervous system (CNS).

Table 4A

MolID	Formula	P_OrlAvail>30%	P_Clr<30ml/min/kg
COTI-HIV1oxo	C17H12FN5O2	0.970	0.961
COTI-HIV1thio	C17H12FN5OS	0.954	0.925
COTI-HIV1Se	C17H12FN5OSe	0.980	0.906
COTI-HIV1imine	C17H13FN6O	0.970	0.934
COTI-HIV1Oxime	C17H13FN6O2	0.933	0.917
COTI-HIV2oxo	C16H13FN4O2	0.971	0.981
COTI-HIV28236-SA	C16H13FN4OS	0.968	0.980
COTI-HIV28236-SeA	C16H13FN4OSe	0.975	0.980
COTI-HIV2imine	C16H14FN5O	0.973	0.967
COTI-HIV28236	C16H14FN5O2	0.966	0.957
COTI-HIV1DOxime	C14H12FN5O3	0.992	0.966
COTI-HIV1Doxo	C14H11FN4O3	0.992	0.984
COTI-HIV1Dthio	C14H11FN4O2S	0.993	0.984
COTI-HIV1Dcyano	C15H11FN6O2	0.992	0.928
COTI-HIV1Dimino	C14H12FN5O2	0.992	0.966
COTI-HIV2DOxime	C13H13FN4O3	0.950	0.976
COTI-HIV2Doxo	C13H12FN3O3	0.940	0.994
COTI-HIV2Dthio	C13H12FN3O2S	0.959	0.994
COTI-HIV2Dcyano	C14H12FN5O2	0.960	0.952
COTI-HIV2Dimino	C13H13FN4O2	0.955	0.986

Table 4B

MolID	Formula	EST_PPB%	P_BBBPenetration
COTI-HIV1oxo	C17H12FN5O2	94.10	0.605
COTI-HIV1thio	C17H12FN5OS	95.83	0.817
COTI-HIV1Se	C17H12FN5OSe	95.42	0.863
COTI-HIV1imine	C17H13FN6O	86.83	0.777
COTI-HIV1Oxime	C17H13FN6O2	93.92	0.640
COTI-HIV2oxo	C16H13FN4O2	48.00	0.694
COTI-HIV28236-SA	C16H13FN4OS	89.71	0.887
COTI-HIV28236-SeA	C16H13FN4OSe	86.05	0.984
COTI-HIV2imine	C16H14FN5O	50.21	0.713
COTI-HIV28236	C16H14FN5O2	51.05	0.275
COTI-HIV1DOxime	C14H12FN5O3	94.53	0.927
COTI-HIV1Doxo	C14H11FN4O3	65.97	0.966
COTI-HIV1Dthio	C14H11FN4O2S	94.37	0.842
COTI-HIV1Dcyano	C15H11FN6O2	93.61	0.946
COTI-HIV1Dimino	C14H12FN5O2	91.38	0.971
COTI-HIV2DOxime	C13H13FN4O3	53.38	0.928
COTI-HIV2Doxo	C13H12FN3O3	50.40	0.920
COTI-HIV2Dthio	C13H12FN3O2S	71.37	0.134
COTI-HIV2Dcyano	C14H12FN5O2	58.95	0.915
COTI-HIV2Dimino	C13H13FN4O2	50.41	0.895

Legend for Table 4:

5 P_OrlAvail>30% is the probability that the oral availability of the compound will be greater than 30%.

P_Clr<30ml/min/kg is the probability that the compound will be cleared from circulation at a rate of less than 30ml/min/kg.

EST_PPB% is the estimated percentage plasma protein binding of the compound.

P_BBBPenetration is an estimate of the probability that the drug will penetrate the blood brain barrier and enter the central nervous system (CNS).

Table 5: Metabolic Stability (Percent remaining at 60 minutes and calculated half life in hours)

Table 5 shows that *in vitro* metabolic stability is expected to be adequate for all compounds. The compounds with a high Est_T1/2(Hrs) are expected to take longer to be reduced to one half of their initial concentration than those with a low Est_T1/2(Hrs). All predicted half lives were considered to be acceptable.

MolID	Formula	Est_T1/2(Hrs)
COTI-HIV1oxo	C17H12FN5O2	12.5
COTI-HIV1thio	C17H12FN5OS	12.1
COTI-HIV1Se	C17H12FN5OSe	14.5
COTI-HIV1imine	C17H13FN6O	16.0
COTI-HIV1Oxime	C17H13FN6O2	13.3
COTI-HIV2oxo	C16H13FN4O2	5.6
COTI-HIV28236-SA	C16H13FN4OS	8.2
COTI-HIV28236-SeA	C16H13FN4OSe	7.2
COTI-HIV2imine	C16H14FN5O	6.4
COTI-HIV28236	C16H14FN5O2	6.1
COTI-HIV1DOxime	C14H12FN5O3	8.5
COTI-HIV1Doxo	C14H11FN4O3	7.0
COTI-HIV1Dthio	C14H11FN4O2S	12.8
COTI-HIV1Dcyano	C15H11FN6O2	16.5
COTI-HIV1Dimino	C14H12FN5O2	8.2
COTI-HIV2DOxime	C13H13FN4O3	7.0
COTI-HIV2Doxo	C13H12FN3O3	6.0
COTI-HIV2Dthio	C13H12FN3O2S	6.7

COTI-HIV2Dcyano	C14H12FN5O2	8.4
COTI-HIV2Dimino	C13H13FN4O2	6.7

Legend for Table 5:

Est_T1/2(hrs) is a calculated estimate of the half life of the drug measured in
5 hours.

Table 6: Probability of Toxicity

The low values provided in Table 6 for HepToxProb show that none of the compounds are expected to cause any Hepatic Toxicity. The compounds
10 with a low AcuteLD50(oral) are potentially more toxic than those with a high AcuteLD50(oral). It takes more of a compound with a high AcuteLD50(oral) to cause death in 50% of the rats to which it is administered, therefore such compounds are less toxic than those with a low AcuteLD50(oral).

Compounds with an LD50 of greater than 500 mg/kg are predicted to have
15 acceptable toxicity for further development.

MolID	Formula	AcuteLD50(oral)	HepToxProb
COTI-HIV1oxo	C17H12FN5O2	698.32	0.131
COTI-HIV1thio	C17H12FN5OS	1063.08	0.109
COTI-HIV1Se	C17H12FN5OSe	760.55	0.155
COTI-HIV1imine	C17H13FN6O	682.51	0.106
COTI-HIV1Oxime	C17H13FN6O2	790.04	0.077
COTI-HIV2oxo	C16H13FN4O2	1276.26	0.022
COTI-HIV28236-SA	C16H13FN4OS	1157.14	0.028
COTI-HIV28236-SeA	C16H13FN4OSe	839.67	0.023
COTI-HIV2imine	C16H14FN5O	1101.07	0.033
COTI-HIV28236	C16H14FN5O2	2362.44	0.070
COTI-HIV1DOxime	C14H12FN5O3	1226.94	0.147
COTI-HIV1Doxo	C14H11FN4O3	1805.56	0.059

COTI-HIV1Dthio	C14H11FN4O2S	1256.64	0.091
COTI-HIV1Dcyano	C15H11FN6O2	1148.98	0.311
COTI-HIV1Dimino	C14H12FN5O2	1356.28	0.048
COTI-HIV2DOxime	C13H13FN4O3	1985.67	0.017
COTI-HIV2Doxo	C13H12FN3O3	658.87	0.322
COTI-HIV2Dthio	C13H12FN3O2S	953.29	0.040
COTI-HIV2Dcyano	C14H12FN5O2	1417.21	0.115
COTI-HIV2Dimino	C13H13FN4O2	1202.51	0.011

Legend for Table 6:

AcuteLD50 (oral) is the calculated point estimate of the dose of the drug in mg/kg that would cause death in 50% of healthy test lab rats when the drug is given orally.

HepToxProb is the average calculated probability from an ensemble of models that the drug in question will cause any liver toxicity.

Table 7: Animal and Human Toxicity Predictions

The expected maximum recommended therapeutic daily dose in humans is provided in the column labeled hMRTD and is measured in mg/kg/d. hERG(IC50>10umol) is the probability that the IC50 value (i.e. the amount required to produce a 50% decrease in initial concentration) for hERG will be greater than 10 umol. A high value indicates that it is likely that the compound will have little hERG channel activity. All of the compounds met the acceptance criterion of the probability being greater than 0.70. A high predicted hERG IC50 value is associated with a low probability of ventricular tachycardia (VentTach).

MolID	Formula	hMRTD (mg/kg/ day)	VentTach	hERG (IC50>10umol)
COTI-HIV1oxo	C17H12FN5O2	1.29	0.018	0.934
COTI-HIV1thio	C17H12FN5OS	0.87	0.469	0.907
COTI-HIV1Se	C17H12FN5OSe	1.23	0.526	0.771
COTI-HIV1imine	C17H13FN6O	1.18	0.008	0.835
COTI-HIV1Oxime	C17H13FN6O2	1.64	0.012	0.814
COTI-HIV2oxo	C16H13FN4O2	1.85	0.013	0.988
COTI-HIV28236- SA	C16H13FN4OS	2.46	0.086	0.932
COTI-HIV28236- SeA	C16H13FN4OSe	2.16	0.040	0.896
COTI-HIV2imine	C16H14FN5O	1.56	0.031	0.959
COTI-HIV28236	C16H14FN5O2	2.15	0.009	0.969
COTI- HIV1DOxime	C14H12FN5O3	7.62	0.002	0.978
COTI-HIV1Doxo	C14H11FN4O3	7.81	0.003	0.993
COTI-HIV1Dthio	C14H11FN4O2S	1.01	0.007	0.981
COTI-HIV1Dcyano	C15H11FN6O2	4.01	0.009	0.928
COTI-HIV1Dimino	C14H12FN5O2	5.04	0.006	0.987
COTI- HIV2DOxime	C13H13FN4O3	14.49	0.003	0.998
COTI-HIV2Doxo	C13H12FN3O3	12.39	0.002	1.000
COTI-HIV2Dthio	C13H12FN3O2S	7.62	0.007	0.995
COTI-HIV2Dcyano	C14H12FN5O2	6.34	0.009	0.994
COTI-HIV2Dimino	C13H13FN4O2	7.15	0.003	0.998

Legend for Table 7:

hMRTD is the calculated maximum recommended therapeutic daily dose of the drug in milligrams per kg per day for the average 60Kg human adult;

VentTach is the probability that the drug would produce ventricular

5 trachycardia, which is a potentially lethal cardiac rhythm disturbance.

hERG(IC₅₀>10 μ mol) is the probability that the IC₅₀ value (i.e. the amount required to produce a 50% decrease in initial concentration) for hERG will be greater than 10 μ mol. A high value indicates that it is likely that the compound will have little activity against hERG.

10

Table 8: Predicted Bio-pharmaceutical Classification (BCS)

Table 8 shows the predicted BCS for all of the structures. Referring to the Legend below, the probability of belonging to a given BCS class is

provided. The majority of the compounds are predicted to be in Class 2. This

15 implies that the compounds are predicted to have good permeability and low solubility and provides guidance for formulation and further development.

MolID	Formula	P_Class 1	P_Class 2	P_Class 3	P_Class 4
COTI-HIV1oxo	C17H12FN5O2	0.000	0.994	0.000	0.005
COTI-HIV1thio	C17H12FN5OS	0.010	0.983	0.000	0.006
COTI-HIV1Se	C17H12FN5OSe	0.000	0.993	0.000	0.006
COTI-HIV1imine	C17H13FN6O	0.005	0.986	0.000	0.009
COTI-HIV1Oxime	C17H13FN6O2	0.008	0.983	0.000	0.009
COTI-HIV2oxo	C16H13FN4O2	0.000	0.079	0.000	0.921
COTI-HIV28236- SA	C16H13FN4OS	0.000	0.741	0.000	0.259
COTI-HIV28236- SeA	C16H13FN4OSe	0.010	0.537	0.008	0.445
COTI-HIV2imine	C16H14FN5O	0.001	0.316	0.003	0.680
COTI-HIV28236	C16H14FN5O2	0.000	0.831	0.000	0.169
COTI- HIV1DOxime	C14H12FN5O3	0.000	0.197	0.000	0.803
COTI-HIV1Doxo	C14H11FN4O3	0.000	0.523	0.000	0.476
COTI-HIV1Dthio	C14H11FN4O2S	0.000	0.995	0.000	0.004
COTI- HIV1Dcyano	C15H11FN6O2	0.003	0.991	0.000	0.007
COTI-HIV1Dimino	C14H12FN5O2	0.000	0.656	0.000	0.343
COTI- HIV2DOxime	C13H13FN4O3	0.004	0.838	0.001	0.157
COTI-HIV2Doxo	C13H12FN3O3	0.001	0.710	0.000	0.288
COTI-HIV2Dthio	C13H12FN3O2S	0.000	0.255	0.001	0.744
COTI- HIV2Dcyano	C14H12FN5O2	0.003	0.774	0.001	0.222
COTI-HIV2Dimino	C13H13FN4O2	0.002	0.817	0.001	0.180

Legend for Table 8:

P_Class 1 is the probability that the compound would be in BCS Class 1 and have a high solubility and a high permeability.

5 P_Class 2 is the probability that the compound would be in BCS Class 2 and have a low solubility and a high permeability.


P_Class 3 is the probability that the compound would be in BCS Class 3 and have a high solubility and a low permeability.

P_Class 4 is the probability that the compound would be in BCS Class 4 and have a low solubility and a low permeability.

10

Example 7: HIV Integrase Assay *in vitro*

The purpose of this study was to analyze the capacity of the COTI-HIV1Doxo (a COTI-HIV1 Ketone Derivative) to inhibit HIV-1 integrase in an *in vitro* assay. It should be noted that this compound is a precursor to all of the
15 COTI-HIV1D- structures. The structure of COTI-HIV1Doxo is shown below:



COTI-HIV1Doxo

Methods:

The capacity of the compound to inhibit HIV integrase in a cell-free
20 system was assessed. An HIV Integrase Assay System (XpressBio Life Science Products; Thurmont, Maryland, www.expressbio.com) was used. In

brief, double-stranded HIV-1 donor substrate (the U5 long terminal repeat) was immobilized in 96-well plates. Recombinant HIV-1 integrase protein was added, along with an oligonucleotide target substrate. The integrase cleaves the donor substrate and integrates it into the target substrate, which is now in solution phase – the integrated product (the amount is indicative of integrase activity) was detected using an antibody specific for the recombinant target/donor sequence. Inhibitors added along with the donor substrate reduced the amount of product. The degree of inhibition by the compound at each concentration tested (from 1 nM to 100 μ M) was determined and, if sufficient inhibition was achieved, an IC₅₀ of the inhibitor was determined. The capacity of a compound known to inhibit HIV integrase activity (NaN₃) was assessed as a positive control. All assessed inhibitions were compared to controls treated with vehicle only (compounds dissolved in 100% DMSO, then diluted in buffer to a maximum DMSO concentration of 0.1% DMSO). Concentrations of DMSO up to 5% have no significant effect on HIV integrase activity.

An advantage of the assay is that the capacity of a test compound to directly inhibit HIV integrase can be determined, without complications arising from the effects of the test compound on cell viability that are a complicating factor when testing effects on integration in living cells. However, it should be kept in mind that indirect HIV integrase inhibition (which might possibly occur in living cells) will not be detected.

Results and discussion:


COTI-HIV1Doxo was assayed for HIV-1 integrase activity, as described above. All measurements were made in quadruplicate (n=4) to allow statistical analysis of significance. NaN₃ (1.0% or 3.3%), a known HIV-1 integrase inhibitor, effectively decreased HIV-1 integrase activity *in vitro*, indicating the reliability of the assay (Fig. 1). COTI-HIV1Doxo inhibited HIV-1 integrase at a concentration of 100 nM (p<0.001, Student's t-test) but not at 10 nM (Rank Sum Test, p=1.000, performed after failure of an equal variance

test on the raw data). All higher tested compound concentrations (up to 100 μ M) significantly decreased HIV-1 integrase activity ($p < 0.001$ in all cases). Interestingly, a low concentration of COTI-HIV-1 ketone derivative (1 nM) enhanced HIV-1 integrase activity, for unknown reasons ($p < 0.001$).


5 Overall, COTI-HIV1Doxo appears to be an effective inhibitor of HIV-1 integrase *in vitro* with capacity to significantly reduce HIV-1 integrase activity at a nanomolar (100 nM) concentration. Complete inhibition of integrase activity was not observed, indicating that increasing COTI-HIV1Doxo concentrations did not reduce integrase function in a linear fashion. In fact,
 10 50% reduction in HIV-1 integrase activity was not achieved until the compound concentrations were in the 10 μ M range. Finally, very low concentrations of COTI-HIV-1 ketone derivative (1 nM) slightly but significantly enhanced HIV-1 integrase activity. It did so in a reproducible fashion, in 3 independent experiments. This is consistent with the
 15 computational predictions provided for Prob_LogIC50 < -6 in Table 3, above.

Example 8: HIV-1 Integrase Assay of COTI-HIV-1, COTI-HIV28233, and COTI-HIV28236


The purpose of this study was to analyze the capacity of COTI-HIV-1 (COTI-
 20 HIV1DOxime), COTI-HIV28233 (COTI-HIV1oxime), and COTI-HIV28236, to inhibit HIV-1 integrase in a cell-free/in-vitro assay system.



COTI-HIV-1
 Formula: $C_{14}H_{12}FN_5O_3$
 Mol. weight: 317.09 g/mol



COTI-HIV28233
 Formula: $C_{17}H_{13}FN_6O_2$
 Mol. weight: 352.32 g/mol



COTI-HIV28236
 Formula: $C_{16}H_{14}FN_5O_2$
 Mol. weight: 327.31 g/mol

Methods:

Compound Preparation: COTI-HIV-1 (8mg; MW:317.27; Lot # JFM-16-177-FIN) COTI-HIV28233 (8mg; Lot # ND10/CO04164D) and COTI-HIV28236 (20mg; MW:327.31; Lot # ND10/CO04139I) were received at room
5 temperature in lyophilized form. Both compounds were kept in this form at room temperature prior to testing (< 1 week).

Test Samples: Integrase only negative control; No integrase negative control; Sodium azide positive control: a known HIV-1 integrase inhibitor; COTI-HIV-1
10 (100.000 µM, 10.000 µM, 1.000 µM, 0.100 µM, 0.010 µM, 0.001 µM); COTI-HIV28233 (100.000 µM, 10.000 µM, 1.000 µM, 0.100 µM, 0.010 µM, 0.001 µM); COTI-HIV28236 (100.000 µM, 10.000 µM, 1.000 µM, 0.100 µM, 0.010 µM, 0.001 µM)

15 HIV-1 integrase assay kit procedure: The effect of each compound on HIV integrase activity was assayed using the "HIV-1 Integrase Assay Kit" version 2.0, catalogue number EZ-1700 obtained from XpressBio Life Science Products. Immediately prior to the start of the assay, both compounds were dissolved in dimethyl sulfoxide (DMSO) and used immediately in the HIV
20 integration assay. The assay was performed according to the instructions listed by the manufacturer (XpressBio Life Science Products, Maryland, USA). In brief, the required streptavidin-coated strip wells were coated with DS oligo and placed overnight at 4°C. The plates were then incubated with integrase and followed by the addition of test articles at the desired concentration. The
25 plates were developed using the supplied HRP antibody and TMB peroxidase substrate as per the manufacturer's instructions. The reactions were stopped with TMB stop solution and immediately read (<10 min) using a VICTOR XXX. The absorbance of each well was read at 450 nm for 0.5 sec. All measurements were performed in triplicate. Sodium azide, a known HIV-1
30 integrase inhibitor, was used as a positive control for inhibiting HIV integrase

activity and was found to effectively decrease HIV-1 integrase activity *in vitro*, indicating the reliability of the assay.

Data analysis: The background absorbance was subtracted from the raw data of each sample to yield "background corrected" absorbance values. The mean absorbance of the test samples divided by the mean integrase control activity multiplied by the associated CV yielded the percent adjusted SD. The CV = %SD for the 100% integrase alone control.

Results and Discussion:

To determine the IC₅₀ of each test compound, non-linear regression analysis was used. The concentration data (x-axis) was log-transformed (Figure 2) and analyzed using the equation $\log(\text{inhibitor})$ vs. normalized response ($Y=100/(1+10^{((X-\text{LogIC}_{50}))})$) to determine the IC₅₀ (Table 1). Since the data was compared to the integrase alone activity, it was presented as a percent of the control activity. Therefore, the activity observed with the integrase alone control was defined at 100% and the activity observed in the absence of integrase as 0%. Since we have a well-defined 0% and 100% and good control data, the response of the test samples were normalized to run between 0% and 100%.

Figure 2D and 2E show the results of the sodium azide control, which was run in parallel with the test samples. As observed, in the absence of sodium azide, the percent control activity was 100% and with increasing concentration of sodium azide, the percent control activity decreased, reaching 0% activity with 10% sodium azide. The calculated IC₅₀ for both sodium azide controls were similar, 0.12% and 0.14%. The data obtained from all test compounds did not form a full sigmoidal curve reaching 0% control activity (Figure 2). Because the top and the bottom plateaus can be defined by solid control data (eg. sodium azide), the data was fitted to a normalized model in order to calculate the IC₅₀. This model assumes that the dose response curve has a standard slope, equal to a Hill slope of -1.0. Since 6 compound concentrations were tested in triplicate, the data was analyzed

using the standard slope model, in contrast with the variable slope model. The IC_{50} values for COTI-HIV-1, COTI-HIV28233 and COTI-HIV28236 were 58.7 μ M, 23.9 μ M and 14.0 μ M (Table 9). COTI-HIV-1 and COTI-HIV28236 both significantly inhibited HIV-1 integrase at all concentrations tested

- 5 ($P < 0.03$ and $P < 0.02$ t-test, respectively). COTI-HIV28233 significantly inhibited HIV-1 integrase at 100 μ M ($P = 0.039$). Of the three compounds, COTI-HIV28233 achieved the lowest percent of the total control activity (34.82%; $P = 0.039$) at a concentration of 100 μ M.

10 **Table 9: Summary of the IC_{50} values for COTI-HIV-1, COTI-HIV28233 and COTI-HIV28236**

Test Sample	IC_{50} (μ M)
COTI-HIV-1	58.7
COTI-HIV28233	23.9
COTI-HIV28236	14.0


- Overall, COTI-HIV-1, COTI-HIV28233 and COTI-HIV28236 appear to
 15 be effective inhibitors of HIV-1 integrase *in vitro* with capacity to significantly reduce HIV-1 integrase activity. COTI-HIV-1 and COTI-HIV28236 were found to significantly inhibit HIV-1 integrase activity *in vitro* at all concentrations tested, with the lowest concentration being 1 nM. COTI-HIV28233 was found to modestly inhibit HIV-1 integrase activity at low concentrations, but to
 20 significantly reduce activity in the range of 100 μ M. The lack of attaining statistical significance at lower concentrations of COTI-HIV28233 can likely be attributed to higher standard deviations in the values obtained.

At least 50% reduction in HIV-1 integrase activity was achieved at 100 μM for both COTI-HIV28233 and COTI-HIV28236. Non-linear regression analysis of the data from COTI-HIV-1, COTI-HIV28233 and COTI-HIV28236 allowed for the calculation of their IC_{50} values, which were 58.7 μM , 23.9 μM and 14.0 μM respectively. Overall, it appears as though COTI-HIV28236 was the most potent drug of the three compounds tested.

The above-described embodiments of the present invention are intended to be examples only. Alterations, modifications and variations may be effected to the particular embodiments by those of skill in the art without departing from the scope of the invention, which is defined solely by the claims appended hereto.

We Claim:

1. A compound of Formula I:



Formula I

a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

wherein:

when X is Se, N-OH, NH, or N-CN,  is a double bond, and

R¹, R², R³, R⁴ and R⁵ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group;


when X is selected from NO₂, CN or N=O,  is a single bond, and

R¹, R², R³, R⁴ and R⁵ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group; and

when X is O or S,  is a double bond,

R¹, R², R³, and R⁴ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

R⁵ is $-(C=O)NR^6R^7$ or



wherein R⁶ and R⁷ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

wherein Z and A are each independently selected from CR⁹ or N, and R⁸ and R⁹ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

2. The compound of claim 1, wherein R¹, R², R³, and R⁴ are each independently selected from H, halo, hydroxyl, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted haloalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkoxy, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, a substituted or unsubstituted heteroaromatic group, carboxyl, alkylcarbonyl, arylcarbonyl, cycloalkylcarbonyl, heterocyclylcarbonyl, amino, aminoalkyl, alkylaminoalkyl, heterocyclylalkyl, aralkyl, arylalkenyl, arylalkynyl,

alkylthio, alkylamino, arylamino, heteroaryl amino, aralkylamino, alkylaminoalkylamino, arylthio, aralkylthio, aryloxy, aralkoxy, heterocyclalkoxy, heterocycloxyalkyl, cycloalkyl, and cycloalkenyl.

3. The compound of claim 1 or 2, wherein R^1 , R^2 , R^3 , and R^4 are each independently selected from H or substituted or unsubstituted alkyl.

4. The compound according to any one of claims 1 to 3, wherein X is selected from Se, N-OH, NH, NO₂, CN, N-CN, or N=O; and R^5 is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.


5. The compound according to claim 4, wherein R^5 is $-(C=O)Y$ and Y is selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

6. The compound according to claim 5, wherein Y is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

7. The compound according to claim 5, wherein the $-(C=O)Y$ is $-(C=O)NR^6R^7$ and R^6 and R^7 are each independently selected from H or the substituted or unsubstituted hydrocarbon group.

8. The compound according to claim 7, wherein the substituted or unsubstituted hydrocarbon group is selected from a substituted or unsubstituted alkyl group.

9. The compound according to claim 8, wherein the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl.
10. The compound according to claim 9, wherein the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group of the substituted alkylphenyl is substituted with a halo group.
11. The compound according to claim 4, wherein R^5 is $-(C=O)OR^6$ and R^6 is selected from H or a substituted or unsubstituted hydrocarbon group.
12. The compound according to claim 11, wherein R^6 is selected from a substituted or unsubstituted alkyl group.
13. The compound according to claim 11, wherein R^6 is H.
14. The compound according to any one of claims 11 to 13, wherein R^1 , R^2 , R^3 , and R^4 are each independently selected from a substituted or unsubstituted alkylaryl.
15. The compound according to claim 14, wherein the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group of the substituted alkylphenyl is substituted with a halo group.
16. The compound according to claim 11, wherein the compound is selected from:



a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.

17. The compound according to any one of claims 1 to 15, wherein X is selected from Se, N-OH, NH, or N=O.

18. The compound according to any one of claims 1 to 3, wherein X is O or S; and R⁶ and R⁷ are each independently selected from H or the substituted or unsubstituted hydrocarbon group.

19. The compound according to claim 18, wherein the substituted or unsubstituted hydrocarbon group is selected from a substituted or unsubstituted alkyl group.




20. The compound according to claim 19, wherein the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl.




21. The compound according to claim 20, wherein the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein



the phenyl group of the substituted alkylphenyl is substituted with a halo group.



22. The compound according to claim 10, 15 or 21, wherein the halo group is selected from bromo, chloro, fluoro or iodo.

23. The compound according to claim 1, wherein the compound is selected from:










a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.

24. The compound according to any one of claims 1 to 3, wherein X is selected from Se, N-OH, NH, or N-CN; and R⁵ is selected from H, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

25. The compound according to claim 24, wherein R⁵ is selected from a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.


26. The compound according to claim 25, wherein R⁵ is a substituted or unsubstituted heteroaromatic group.

27. The compound according to claim 26, wherein R⁵ is



wherein Z and A are each independently selected from CR⁹ or N, and R⁸ and R⁹ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

28. The compound according to any one of claims 1 to 3, wherein X is O or S and R⁵ is the




wherein Z and A are each independently selected from CR⁹ or N, and R⁸ and R⁹ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group.

29. The compound according to claim 27 or 28, wherein at least one of Z and A is N.

30. The compound according to claim 29, wherein both Z and A are N.

31. The compound according to any one of claims 27 to 30, wherein R⁸ is a substituted or unsubstituted hydrocarbon group.
32. The compound according to claim 31, wherein R⁸ is selected from a substituted or unsubstituted alkyl group.
33. The compound according to claim 32, wherein the substituted or unsubstituted alkyl group is a substituted or unsubstituted alkylaryl or a substituted or unsubstituted alkylheteroaryl.
34. The compound according to claim 33, wherein the substituted or unsubstituted alkylaryl is a substituted or unsubstituted alkylphenyl, wherein the phenyl group of the substituted alkylphenyl is substituted with a halo group.
35. The compound according to claim 34, wherein the halo group is selected from bromo, chloro, fluoro or iodo.
36. The compound according to claim 1, wherein the compound is selected from:



a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof.

37. The compound according to any one of claims 1 to 36, wherein the compound is an HIV integrase inhibitor.

38. A pharmaceutical composition comprising the compound of any one of claims 1 to 37, a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent.

39. The pharmaceutical composition according to claim 38, in combination with at least one other anti-HIV agent.

40. Use of an anti-HIV effective treatment amount of the compound of any one of claims 1 to 37, a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, in the manufacture of a medicament for the treatment of HIV in a mammal.

41. Use of an anti-HIV effective treatment amount of the compound of any one of claims 1 to 37, a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, in the manufacture of a medicament for salvage therapy in the treatment of HIV in a mammal.


42. Use of an anti-HIV effective treatment amount of the compound of any one of claims 1 to 37, a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, for the treatment of HIV in a mammal.

43. Use of an anti-HIV effective treatment amount of the compound of any one of claims 1 to 37, a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof, optionally with a pharmaceutically acceptable carrier or diluent, for salvage therapy in the treatment of HIV in a mammal.

44. The use according to any one of claims 40 to 43, wherein the mammal is human.

45. The use according to any one of claims 40 to 44, further comprising use of at least one other anti-HIV agent in combination and/or in alternation with the compound.

46. A method for making the compound of Formula III:



Formula III


a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S, wherein:

when X is selected from Se, N-OH, NH, N-CN, O or S,  is a double bond,

when X is selected from NO₂, CN or N=O,  is a single bond,

48. A method for making the compound of Formula IV:




Formula IV

a pharmaceutically-acceptable salt, hydrate, solvate, tautomer, optical isomer, E-isomer, Z-isomer, or combination thereof;

X is selected from Se, N-OH, NH, NO₂, CN, N-CN, N=O, O or S,

wherein:

when X is Se, N-OH, NH, or N-CN,  is a double bond,

when X is selected from NO₂, CN or N=O,  is a single bond,


when X is O or S,  is a double bond,

R¹, R², R³, and R⁴ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group, and

wherein R⁶ and R⁷ are each independently selected from H, halo, hydroxyl, amino, a substituted or unsubstituted hydrocarbon group, a substituted or unsubstituted heterogeneous group, a substituted or unsubstituted carbocyclic group, a substituted or unsubstituted heterocyclic group, a substituted or unsubstituted aromatic group, or a substituted or unsubstituted heteroaromatic group,

wherein the method comprises:


a) reacting a compound of Formula VII with an amine of NHR^6R^7 to form an intermediate of Formula IX:



b) reacting the intermediate of Formula IX under conditions to replace the oxygen of the carbonyl group in the ring with any X group defined herein to form the compound of Formula IV.


49. The method of claim 48, wherein b) comprises reacting Formula IX with NH_2OH to form the compound of Formula IV, wherein X is N-OH.


1/2

Figure 1


2/2

Figure 2





Formula I



Formula II