



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2304303 A1 2001/09/30

(21) **2 304 303**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(22) Date de dépôt/Filing Date: 2000/03/31

(41) Mise à la disp. pub./Open to Public Insp.: 2001/09/30

(51) Cl.Int.⁷/Int.Cl.⁷ A23K 1/14, A21D 2/38, C09F 5/02

(71) Demandeurs/Applicants:
HK RUOKATALO OYJ, FI;
ELIXI OIL OY, FI

(72) Inventeur/Inventor:
MYLLYMAKI, OLAVI, FI

(74) Agent: CASSAN MACLEAN

(54) Titre : TRAITEMENT DE GRAINES DE LIN

(54) Title: PROCESSING OF FLAXSEED

(57) **Abrégé/Abstract:**

The invention relates to a method for removing the flaxseed husk from the seed core by means of abrasion. In this method, the outer husk portion firstly removed by abrasion is separated as a mucilage fraction and the subsequently removed inner portion is separated as a fibre fraction. The fibre fraction is particularly rich in lignans.



(57) Abstract

The invention relates to a method for removing the flaxseed husk from the seed core by means of abrasion. In this method, the outer husk portion firstly removed by abrasion is separated as a mucilage fraction and the subsequently removed inner portion is separated as a fibre fraction. The fibre fraction is particularly rich in lignans.

PROCESSING OF FLAXSEED

Technological background

5

The invention relates to the processing of flaxseed (*Linum sp.*) in order to produce a food supplement.

10

The flaxseed husk consists of three separate anatomic parts. The outer part consists of a water-soluble mucilage, the epidermic layer. Under the mucilage, there is the actual husk substance, the spermodermis, consisting of four layers. The inner husk surface is the endosperm, and the seed core consists of the cotyledon, which accounts for about half of the weight of the seed. The major portion of the oil and protein of the seed is located in the cotyledon. Seeds have been used as such or in a ground form in animal feed and human nutrition. The mucilage separated from the

15

Flaxseed has also been husked in order to separate the husk portion from the inner portion (Wanasundra & Shahidi, Food Chem, 59 (1997) 47-55). Nevertheless, husking is a laborious and awkward process due particularly to the seed shape. Abrasive methods have been implemented in the husking.

20

Oil can be separated from flaxseed by pressing or extracting. The solid residue has been mainly used as an animal feed. It has been generally considered inapt for use as a foodstuff (Oomah & Mazza, Food. Chem., 48 (1993), 109-114), but has been used all the same, especially in bakery products. Its use is restricted especially by the flavour and colour changes in the foodstuff, which are felt to be unpleasant.

25

Flaxseed also contains lignans in an amount of 80-370 mg/100 g (Mazur et al., Anal. Biochem., 233 (1996) 169-180). In nutrition, lignans have proved to provide health benefits, i.a. anti-carcinogenic effects.

30

General description of the invention

35

A method has now been found in which the husk layer is removed from whole flaxseed and the husk is subsequently divided into a firstly removed mucilage fraction and a secondly removed fibre fraction. The mucilage fraction comprises the outermost husk layers and is rich in water-dispersible carbohydrates (mucilage). The

fibre fraction comprises the inner husk layers and is particularly rich in fibres and lignans. The abrasion can be performed in a mill specifically equipped with millstones.

- 5 The husk removing method of the invention serves to detach separate fractions suitable for different purposes of use in one single process.

10 A method has now also been found, in which flaxseed solid substance obtained by separating oil from the seed is bleached by treating it with a mixture of water and alcohol. The bleaching mixture may contain hydrogen peroxide in addition. Husks or husk fractions separated from whole seeds can also be bleached with the bleaching method of the invention.

15 The bleaching method of the invention yields a product, which, used as a food supplement, causes less changes in colour and flavour than a non-processed solid substance does.

Detailed description of the invention

20 The outermost flaxseed layer comprises principally water-soluble carbohydrates and forms the flaxseed mucilage. Under this, there is a layer comprising mainly fibres. When a whole unbroken flaxseed is abraded, the outer layer is first removed. This layer is separated as a mucilage fraction. Then the inner layer comes off, being separated as a fibre fraction. This fraction is also rich in lignans. The remainder
25 comprises the inner seed portion, which is rich in oil and proteins. The inner portion accounts for approximately half of the weight of the seed.

30 The mucilage fraction is usable in foodstuffs especially as a viscosity-increasing agent in breads, for instance. The mucilage fraction is also usable in cosmetic and medicinal products. The rheological behaviour of the mucilage fraction is similar to that of arabic gum.

35 The mucilage fraction accounts for about 0.5-10% of the weight of the seed, such as 5-8%. The mucilage fraction draws along some of the inner portion of the husk.

The fibre fraction is usable as a food supplement particularly when an increase in fibres and lignans is desired. Lignans have proved to provide health benefits. They seem to act as anti-carcinogenes, among other things. The fibre fraction naturally

comprises a certain amount of oil. The oil can be removed by means of hexane for instance. A fraction particularly rich in fibres can also be separated from the fibre fraction. A lignan concentrate can be further produced from the fibre fraction e.g. by screening or grading, especially if the oil is first removed. Lignan can be concentrated also by means of extraction.

The fibre fraction accounts for about 20-70%, such as 30-50%, especially 35-45% of the weight of the seed.

Oil can be separated from the inner part of the seed, the residue being rich in proteins. A protein part with a particularly high, up to 70% protein content, and a mucilage fraction can be further separated from the residue.

The abrasion is performed by seeking to maintain the inner part of the seed as complete as possible to allow it to be separated from the abraded powder. The abrasion can be performed using commonly used abrading devices. The devices can be modified rice polishing devices, for instance. The abrasion is performed by seeking to detach the husk portion as completely as possible, while removing only a minimum of the inner portion. The abrasion can be performed as a continuous process.

The seeds can be abraded for instance in a mill equipped with millstones. The rough surfaces of the millstones abrade the seeds at a given rate, preferably for about 1-3 minutes. The abraded powder has been found to have a particularly high lignan content after an abrasion period of about 1 minute.

The fibre fraction is a brownish powder. It can be bleached and simultaneously deflavoured, degreased and upgraded.

In the bleaching, the powder is moisturised with water, the ratio of water to powder varying e.g. in the range from 1:1 to 3:1. Enzymes such as lipase or protease can be added to the water. The water is allowed to act for e.g. 0.5-12 h, and then the paste is homogenised in alcohol, such as C1-C5 alcohol, especially ethanol or isopropanol, particularly in isopropanol. The alcohol amount is e.g. 2-5 times the powder amount. The homogenised alcohol-containing suspension is separated from the solution e.g. by filtering or centrifugating. The separated powder is washed e.g. with a 2-6-fold alcohol amount, and after this the suspension is preferably filtered

and centrifugated. The alcoholic mass is dried e.g. at 20-90 °C. The dried paste is ground or granulated to the desired grain size if necessary.

5 The alcohol breaks down the viscosity generated by the alcohol-insoluble but water-soluble carbohydrate constituent. This makes it easy to homogenise the paste to a non-viscous suspension and to perform the mechanical separating treatments of the suspension, such as filtering and centrifugation. The water treatment results in the solid structure opening up so that the major portion of the remaining oil in the paste is removed during the water-alcohol treatment. The more finely ground the powder,
10 the stronger the effect. The alcohol added after the water treatment serves to precipitate the water-soluble carbohydrate, detaching it from the surrounding matrix. This allows for instance the dry product to be concentrated into a water-soluble component by mechanical means, such as grading.

15 The bleaching also removes cyanogens from the product.

A better result than with mere water-alcohol treatment is achieved by adding e.g. 2-7% of hydrogen peroxide to the aqueous solution at the elutriation stage. Hydrogen peroxide added before alcohol detaches mucilages from the surface efficiently.

20 Before the bleaching, the powder can be thermally treated in order to inactivate any enzyme activity affecting the flavour. The temperature range may be e.g. 40-80 °C and the treatment period e.g. 0.5-2 h. The treatment can be performed in vacuum. Besides inactivating enzymes, the temperature and possible vacuum have the joint
25 effect of removing low-molecular compounds affecting the taste.

A solid residue remaining after oil has been separated from whole flaxseed (flaxseed groats), especially a residue obtained by pressing, can also be bleached with the method of the invention. The groats are preferably ground before the
30 bleaching e.g. by means of an emery mill with a pair of millstones.

Example 1

A device designed for oat abrasion was used in the flaxseed abrasion. The seeds
35 were abraded between the millstones and the metal mesh in a batch device (batches of 80 g) equipped with two millstones. At the end of the abrasion, the seeds were removed from the device through a hatch at the mesh bottom. During the process, the abraded powder passed through the mesh into a receiver. Table 1 shows the

abraded fraction yield and the protein, fat and lignan concentrations as a function of abrasion time.

5 Table 1. Abrased laxseed (*Linum usitatissimum*, Helmi ssp.) powder yield and protein, fat and lignan concentration calculated on the dry substance.

Abrasion time (min)	Protein (% d.s.)	Fat (% d.s.)	Lignan mg/100 g	Yield (%)
1	12	18	1480	4
2	15	27	1360	9
3	17	29	1230	12
4	18	31	1070	16
5	19	34	1070	18
6	19	34	960	21
7	19	36	1030	22
8	20	36	800	24

10 The powder obtained is usable as a reactant in further lignin concentration. By extracting the oil from the abraded powder e.g. by hexane extraction, the lignin concentration of the powder will increase by an amount equal to the oil content, i.e. by 18% after 1 minute of abrasion. The powder can be further bleached for instance as described in example 2.

Example 2

15

Flaxseed groats obtained in a cold-drawing process were bleached. The groats were subjected to thermal treatment in order to inactivate any enzyme activity affecting the flavour of the groats. The best sensory result was obtained by thermally treating the groats for 1 h in vacuum at 60 °C. The dark brown groats obtained were
 20 elutriated in water at room temperature, forming a paste-like powder. After a settling of about 0.5-2 hours the paste-like powder was elutriated in isopropanol at a ratio of 1.2-3 (w/w). The crushed flaxseed suspension obtained was filtrated in vacuum. The deposit obtained, i.e. the filterpress cake, was washed during homogenisation in isopropanol, was refiltered in vacuum or centrifugated in a
 25 decanter centrifuge. The filtrated powder was dried at room temperature or using a closed drying technique, allowing isopropanol to be recovered.

A better result than the one obtained with mere water-alcohol treatment was obtained by adding hydrogen peroxide to the aqueous solution at the elutriation stage. A very light powder was obtained by elutriating the groats in an aqueous solution with a 4-5% hydrogen peroxide content. After an elutriation time of 0.5-2 h
5 the paste-like powder was homogenised in a double amount of isopropanol, filtered in vacuum, washed in isopropanol and refiltered. The filtered powder was dried.

A product particularly rich in proteins was obtained when isopropanol accounted for 50% of the solution and peroxide for about 5% of the solution.

Claims

1. A method for removing the flaxseed husk from the seed core by abrasion, characterised in that the firstly removed outer portion of the husk is separated as a mucilage fraction and then the secondly removed inner portion is separated as a fibre fraction.
2. A method as defined in claim 1, in which the separated mucilage fraction accounts for 3-10%, such as 5-8% of the weight of the seed.
3. A method as defined in claim 1 or 2, in which the separated fibre fraction accounts for 20-70%, such as 30-50%, especially 35-45% of the weight of the seed.
4. A method as defined in any of claims 1-4, in which a high-fibre fraction rich in fibres is further separated from the fibre fraction.
5. A method as defined in any of claims 1-4, in which a lignan fraction rich in lignan is further separated from the fibre fraction.
6. A method as defined in any of claims 1-5, in which oil is separated from the fibre fraction, especially by means of extraction.
7. A method as defined in any of claims 1-6, in which oil is separated from the inner portion remaining after the abrasion.
8. A method as defined in claim 7, in which a protein fraction and carbohydrate fraction of the core are separated from the residue remaining after the oil separation.
9. A method as defined in any of claims 1-8, in which the abrasion is performed in a mill equipped with millstones.
10. A method as defined in any of claims 1-9, in which the separated fibre fraction is bleached.
11. A method as defined in claim 10, in which the bleaching is performed with a bleaching solution containing water and alcohol.

12. A method as defined in claim 11, in which the alcohol is C1-C5 alcohol, such as ethanol or isopropanol, especially isopropanol.
13. A method as defined in claims 11 or 12, in which the alcohol amount is 2-5
5 times the amount of fibre fraction.
14. A method as defined in any of claims 11-13, in which the bleaching solution also contains hydrogen peroxide, e.g. 2-7%, such as 4-6%, especially about 5%.
- 10 15. A method as defined in any of claims 11-14, in which the fibre fraction is treated with an aqueous solution for e.g. 0.5-12 h before the treatment with the bleaching solution.
- 15 16. A method as defined in claim 15, in which the aqueous solution contains enzymes such as lipases or proteases.
17. A method as defined in claims 15 or 16, in which the slurry obtained in the water treatment is mixed with said alcohol in order to perform bleaching.
- 20 18. A method as defined in claims 14 and 17, in which the hydrogen peroxide is added to said aqueous solution.
- 25 19. A method as defined in any of claims 10-18, in which the fibre fraction is thermally treated before the bleaching, e.g. for 0.5-2 h, at a temperature of e.g. 40-80 °C, especially in vacuum.