



US010822394B2

(12) **United States Patent**
Grillberger et al.

(10) **Patent No.:** **US 10,822,394 B2**
(45) **Date of Patent:** ***Nov. 3, 2020**

(54) **METHOD OF PRODUCING RECOMBINANT HIGH MOLECULAR WEIGHT VWF IN CELL CULTURE**

(71) Applicants: **Baxalta Incorporated**, Bannockburn, IL (US); **Baxalta GmbH**, Glattpark (Opfikon) (CH)

(72) Inventors: **Leopold Grillberger**, Vienna (AT); **Manfred Reiter**, Vienna (AT); **Wolfgang Mundt**, Vienna (AT)

(73) Assignees: **Baxalta GmbH**, Glattpark (Opfikon) (CH); **Baxalta Incorporated**, Bannockburn, IL (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/125,141**

(22) Filed: **Sep. 7, 2018**

(65) **Prior Publication Data**

US 2019/0085052 A1 Mar. 21, 2019

Related U.S. Application Data

(60) Continuation of application No. 15/591,022, filed on May 9, 2017, now Pat. No. 10,100,099, which is a continuation of application No. 15/201,125, filed on Jul. 1, 2016, now Pat. No. 9,834,591, which is a division of application No. 14/339,319, filed on Jul. 23, 2014, now Pat. No. 9,409,971, which is a division of application No. 13/179,386, filed on Jul. 8, 2011, now Pat. No. 8,852,888.

(60) Provisional application No. 61/362,635, filed on Jul. 8, 2010.

(51) **Int. Cl.**

A61K 48/00 (2006.01)
C07K 14/755 (2006.01)
C12N 9/64 (2006.01)
A61K 9/00 (2006.01)
A61K 38/36 (2006.01)
A61K 38/37 (2006.01)
C12P 21/00 (2006.01)
A61K 38/00 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 14/755** (2013.01); **A61K 9/0019** (2013.01); **A61K 38/36** (2013.01); **A61K 38/37** (2013.01); **C12N 9/6489** (2013.01); **C12P 21/00** (2013.01); **A61K 38/00** (2013.01); **C12N 2500/24** (2013.01); **C12N 2511/00** (2013.01); **C12Y 304/24087** (2013.01)

(58) **Field of Classification Search**

CPC ... **C07K 14/755**; **C12N 9/6489**; **A61K 9/0019**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,767,704 A 8/1988 Cleveland et al.
5,804,420 A 9/1998 Chan et al.

6,100,061 A 8/2000 Reiter et al.
6,156,570 A 12/2000 Hu et al.
6,171,825 B1 1/2001 Chan et al.
6,528,286 B1 3/2003 Ryll
6,531,577 B1 3/2003 Kaersgaard et al.
6,936,441 B2 8/2005 Reiter et al.
7,335,634 B2 2/2008 Walter et al.
7,709,229 B2 5/2010 Casatorres Hernandez et al.
8,039,231 B2 10/2011 Luan et al.
8,313,926 B2 11/2012 Grillberger et al.
8,486,699 B2 7/2013 Talbot et al.
8,580,554 B2 11/2013 Grillberger et al.
8,759,026 B2 6/2014 Grillberger et al.
8,852,888 B2 10/2014 Grillberger et al.
2003/0077806 A1 4/2003 Selden et al.
2005/0084928 A1 4/2005 Birch et al.
2005/0266528 A1 12/2005 Laemmie et al.
2007/0190057 A1 8/2007 Wu et al.
2007/0212770 A1 9/2007 Grillberger et al.
2008/0009040 A1 1/2008 Grillberger et al.
2008/0064644 A1 3/2008 Castorres Hernandez et al.
2009/0068705 A1 3/2009 Drapeau et al.
2009/0176269 A1 7/2009 Giovagnoli et al.
2009/0291064 A1 11/2009 Talbot et al.
2009/0317867 A1 12/2009 Luan et al.
2010/0227819 A1 9/2010 Hernandez et al.
2011/0086413 A1 4/2011 Grillberger et al.
2011/0086813 A1 4/2011 Glaser et al.
2011/0151512 A1 6/2011 Grillberger et al.
2012/0035110 A1 2/2012 Grillberger et al.

FOREIGN PATENT DOCUMENTS

EP 0 481 791 A2 4/1992
EP 0 197 592 B1 2/1993
EP 1 096 017 A2 5/2001
KR 10-2007-0073930 A 7/2007
KR 10-2007-0085220 A 8/2007
WO WO 86/06096 A1 10/1986
WO WO 96/15231 A2 5/1996
WO WO 96/26266 A1 8/1996
WO WO 98/08934 A1 3/1998
WO WO 98/15614 A1 4/1998

(Continued)

OTHER PUBLICATIONS

Printout from Wikipedia. En.wikipedia.org/wiki/Von_Wellebrand_factor. pp. 1-9. Printed Jan. 23, 2020. (Year: 2020).*

(Continued)

Primary Examiner — Marcia S Noble
(74) *Attorney, Agent, or Firm* — Morgan, Lewis & Bockius LLP

(57) **ABSTRACT**

Among other aspects, the present invention relates to cell culture conditions for producing high molecular weight vWF, in particular, highly multimericWF with a high specific activity and ADAMTS13 with a high specific activity. The cell culture conditions of the present invention can include, for example, a cell culture medium with an increased copper concentration and/or cell culture supernatant with a low ammonium (NH₄⁺) concentration. The present invention also provides methods for cultivating cells in the cell culture conditions to express high molecular weight vWF and rA13 having high specific activities.

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 00/03000	A2	1/2000
WO	WO 01/23527	A1	4/2001
WO	WO 2002/042441	A2	5/2002
WO	WO 2007/077217	A2	7/2007
WO	WO 2008/082669	A2	7/2008
WO	WO 2008/109410	A1	9/2008
WO	WO 2009/086309	A2	7/2009
WO	WO 2009/086309	A3	7/2009
WO	WO 2009/088713	A1	7/2009
WO	WO 2011/012725	A1	2/2011
WO	WO 2011/014838	A1	2/2011
WO	WO 2012/006591	A1	1/2012
WO	WO 2012/006594	A1	1/2012

OTHER PUBLICATIONS

Sadler. Annu. Rev. Biochem. 1998. 67:395-424 (Year: 1998).*

Altamirano, C. et al., "Improvement of CHO Cell Culture Medium Formulation: Simultaneous Substitution of Glucose and Glutamine," *Biotechnol. Prog.*, 2000, vol. 16, pp. 69-75.

Bevitt, D.J. et al., "Expression of ADAMTS metalloproteinases in retinal pigment epithelium derived cell line ARPE-19: transcriptional regulation by TNF α ," *Biochimica et Biophysica Acta*, 2003, vol. 1626, pp. 8391.

Franěk, F. et al., "Plant Protein Hydrolysates: Preparation of Defined Peptide Fractions Promoting Growth and Production in Animal Cells Cultures," *Biotechnol. Prog.*, 2000, vol. 16, pp. 688-692.

International Search Report dated Sep. 27, 2011, for International Patent Application No. PCT/US2011/043455 filed Jul. 8, 2011, 5 pages.

International Search Report dated Sep. 30, 2011, for International Patent Application No. PCT/US2011/043459 filed Jul. 8, 2011, 4 pages.

Jeong, Y-H. et al., "In Situ Removal of Ammonium Ions From Hybridoma Cell Culture Media: Selection of Adsorbent," *Biotechnology Techniques*, Jul./Aug. 1992, vol. 6, No. 4, pp. 341-346.

Kaufman, R.J. et al., "Effect of von Willebrand Factor Coexpression on the Synthesis and Secretion of Factor VIII in Chinese Hamster Ovary Cells," *Mol. Cell Biol.*, Mar. 1989 9(3):1233-1242.

Kokame, K. et al., FRET-S-VWF73, a first fluorogenic substrate for ADAMTS13 assay, "*British Journal of Haematology*," 2005, vol. 129, pp. 93-100.

Lao, M-S. et al., "Effects of Ammonium and Lactate on Growth and Metabolism of a Recombinant Chinese Hamster Ovary Cell Culture," *Biotechnol. Prog.*, 1997, vol. 13, pp. 688-691.

Martinelle, K. et al., "Mechanisms of ammonia and ammonium ion toxicity in animal cells: transport across cell membranes," *Journal of Biotechnology*, 1993, 30:339-350.

Mayadas, T.N. et al., "In Vitro Multimerization of von Willebrand Factor Is Triggered by Low pH, Importance of the Propolypeptide and Free Sulfhydryls," *The Journal of Biological Chemistry*, Aug. 15, 1989, vol. 264, No. 23, pp. 13497-13503.

Mignot, G. et al., "Production of recombinant Von Willebrand factor by CHO cells cultured in macroporous microcarriers," *Cytotechnology*, 1990, vol. 4, pp. 163-171.

Neifar, M. et al., "Effect of culturing processes and copper addition on laccase production by the white-rot fungus *Fomes fomentarius* MUCL 35117," *The Society for Applied Microbiology, Letters in Applied Microbiology*, 2009, vol. 49, pp. 73-78.

Plaimauer, B. et al., "Cloning, expression, and functional characterization of the von Willebrand factor-cleaving protease (ADAMTS13)," *Blood*, Nov. 15, 2002, vol. 100, No. 10, pp. 3626-3632.

Plaimauer, B. et al., "Expression and Characterization of Recombinant Human ADAMTS-13," *Seminars in Hematology*, Jan. 2004, vol. 41, No. 1, pp. 24-33.

Schneider, M. et al., "The importance of ammonia in mammalian cell culture," *Journal of Biotechnology*, 1996, vol. 46, pp. 161-185.

Turecek, P.L. et al., "Development of a plasma- and albumin-free recombinant von Willebrand factor," *Hämostaseologie*, 2009, vol. 29 (Suppl 1): S32-38.

Wagner, D.D. et al., "Initial Glycosylation and Acidic pH in the Golgi Apparatus Are Required for Multimerization of von Willebrand Factor," *The Journal of Cell Biology*, Apr. 1986, vol. 102, pp. 1320-1324.

Altamirano, C. et al. "Advances in improving mammalian cells metabolism for recombinant protein production" *Electronic Journal of Biotechnology* [Online], 16 May 14, 2013.

Davami, F. et al. "Effects of Peptone Supplementation in Different Culture Media on Growth, Metabolic Pathway and Productivity of CHO DG44 Cells; a New Insight into Amino Acid Profiles." *Iranian biomedical journal* vol. 19,4 (2015): 194-205.

Sadler, J.E. "von Willebrand factor assembly and secretion." *Journal of thrombosis and haemostasis* : JTH vol. 7 Suppl 1 (2009): 24-7.

Sakai, Y., et al. "Growth Inhibition and Heavy Metal Accumulation in CHO Cells." In: Beuvery E.C., Griffiths J.B., Zeijlemaker W.P. (eds) *Animal Cell Technology: Developments Towards the 21st Century*. Springer, Dordrecht. (1995).

Thermo Fisher Catalog, 11320—DMEM/F-12.

Voorberg, et al. "Domains involved in multimer assembly of von Willebrand factor (VWF)—multimerization is independent of dimerization." *The EMBO journal*. 9. 797-803. (1990).

Fischer, B. E. "Effect of multimerization of human and recombinant von Willebrand factor on platelet aggregation, binding to collagen and binding of coagulation factor VIII," 1996, pp. 55-66, vol. 84, No. 1, *Thrombosis Research*.

* cited by examiner




FIG. 1A

Lane Number	Band Number	Trace Int x mm	Relative Qty
1	1	809715.2	74.5
1	2	277519.7	25.5
2	1	681110.1	83.7
2	2	132376.5	16.3
3	1	568177	90.2
3	2	61619.3	9.8
4	1	500725.4	94.0
4	2	31815.3	6.0
5	1	433194.4	95.9
5	4	18307.3	4.1
6	1	412190.2	95.9
6	2	17441	4.1
7	1	842215.1	72.0
7	2	327960.1	28.0
8	1	723836.4	70.5
8	2	303095.4	29.5
9	1	739912.7	68.7
9	2	337575.1	31.3
10	1	929848.4	68.6
10	2	425784.2	31.4
11	1	684679.9	78.6
11	2	185964.5	21.4
12	3	512298.6	66.7
12	4	256326.7	33.3

FIG. 1B




FIG. 2A




FIG. 2B

4 μ g/L 1 μ g/L 0.66 μ g/L




FIG. 3A

4 μ g/L 1 μ g/L 0.66 μ g/L




FIG. 3B




FIG. 4




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 5F




FIG. 5G




FIG. 5H




FIG. 5I




FIG. 5J




FIG. 5K




FIG. 6A

Lane Number	Band Number	Trace Int x mm	Relative Qty
1	1	155657.0	97.2
1	2	4453.3	2.8
2	1	419998.6	87.6
2	2	59416.6	12.4
3	1	225344.4	97.4
3	2	5898.2	2.6
4	1	557340.0	73.2
4	2	203807.5	26.8
5	1	248653.8	97.8
5	2	5599.8	2.2
6	1	434040.2	95.4
6	2	20823.4	4.6
7	1	238580.2	97.9
7	2	5159.7	2.1
8	1	322940.6	77.3
8	2	94677.8	22.7
9	1	250356.6	98.0
9	2	5183.5	2.0
10	1	275819.1	97.7
10	2	6469.7	2.3
11	1	207447.3	97.3
11	2	5687.5	2.7
12	1	604779.1	76.2
12	2	189356.4	23.8

FIG. 6B

METHOD OF PRODUCING RECOMBINANT HIGH MOLECULAR WEIGHT VWF IN CELL CULTURE

CROSS REFERENCES TO APPLICATIONS

This application is a Continuation of U.S. patent application Ser. No. 15/591,022, filed May 9, 2017 (now issued as U.S. Pat. No. 10,100,099), which is a Continuation of U.S. patent application Ser. No. 15/201,125, filed Jul. 1, 2016, (now issued as U.S. Pat. No. 9,834,591) which is a Divisional of U.S. patent application Ser. No. 14/339,319, filed Jul. 23, 2014, (now issued as U.S. Pat. No. 9,409,971), which is a Divisional of U.S. patent application Ser. No. 13/179,386, filed Jul. 8, 2011 (now issued as U.S. Pat. No. 8,852,888), which claims priority to U.S. Provisional Patent Application Ser. No. 61/362,635, filed Jul. 8, 2010, the disclosures of which are hereby incorporated herein by reference in their entireties for all purposes.

BACKGROUND OF THE INVENTION

The recombinant expression of therapeutic proteins in cell culture (particularly large-scale cell cultures), including eukaryotic cell culture, and more specifically mammalian cell culture, requires the use of special culture media providing nutrient substances for efficient growth of cells. Cell culture media formulations are often supplemented with a range of additives, including fetal calf serum (FCS), animal derived proteins and/or protein hydrolysates of bovine origin as well as protein hydrolysates derived from plants or yeast. One challenge with such cultures is that the amount of protein and the total and specific activity of the protein produced are often variable across different cell cultures, even when the formulation for the cell culture media is not changed. This variability is especially apparent in the case of large-scale manufacturing processes utilizing cell culture volumes of 10 liters to over 20,000 liters. Cell culture media containing hydrolysates are particularly prone to variability from one cell culture to the next, leading to decreased production of total protein as well as decreased total and specific activity.

One potential reason for the variability seen across different cell cultures is that contaminants in additives such as hydrolysates vary from one batch to the next. In general, serum or serum-derived substances, such as, e.g., albumin, transferrin or insulin, may comprise unwanted agents that can contaminate the cell cultures and the biological products obtained thereof. Furthermore, human serum derived additives have to be tested for all known viruses, including hepatitis viruses and HIV which can be transmitted via serum. Moreover, bovine serum and products derived thereof bear the risk of BSE contamination. In addition, all serum-derived products can be contaminated by unknown substances. When using serum or protein additives derived from human or animal sources in cell culture, there are numerous problems (e.g., the varying quality in composition of different batches and the risk of contamination with *mycoplasma*, viruses or BSE), particularly if the cells are used in the manufacture of drugs or vaccines for human administration. Therefore, many attempts have been made to provide efficient host systems and cultivation conditions, which do not require serum or other animal protein compounds.

Such serum-free media have been developed on the basis of protein extracts derived from plants or yeast. For example, soy hydrolysates are known to be useful for

fermentation processes and can enhance the growth of many fastidious organisms, yeasts and fungi. WO 96/26266 describes that papain digests of soy meal are a source of carbohydrate and nitrogen and many of the components can be used in tissue culture. Franek et al. (Biotechnology Progress (2000) 16, 688-692) describe growth and productivity promoting effects of defined soy and wheat hydrolysate peptide fractions.

WO 96/15231 discloses a serum-free medium composed of a synthetic minimal essential medium and a yeast extract for the propagation of vertebrate cells and a virus production process. A medium formulation composed of a basal cell culture medium comprising a rice peptide and an extract of yeast and an enzymatic digest thereof, and/or a plant lipid for growth of animal cells is disclosed in WO 98/15614. A medium comprising purified soy hydrolysate for the cultivation of recombinant cells is disclosed in WO 01/23527. WO 00/03000 discloses a medium that comprises a soy hydrolysate and a yeast extract, but also requires the presence of recombinant forms of animal proteins, such as growth factors.

EP-A-0 481 791 describes a biochemically defined culture medium for culturing engineered CHO cells, which is free from protein, lipid and carbohydrate isolated from an animal source, further comprising a recombinant insulin or insulin analogue, 1% to 0.025% w/v papain digested soy peptone and putrescine. WO 98/08934 describes a serum-free eukaryotic cell culture comprising hydrolyzed soy peptides (1-1000 mg/L), 0.01 to 1 mg/L putrescine and a variety of animal-derived components, including albumin, fetuin, various hormones and other proteins. In this context, it should be noted that putrescine is also known to be comprised in standard media like DMEM/Ham's F12 in a concentration of 0.08 mg/L.

The plant and/or yeast hydrolysates, however, are undefined mixtures of oligopeptides and other unknown components and contaminants. Moreover, the quality of commercially available lots of hydrolysates varies extremely. As a result, there are large variations in the production of recombinant proteins or viral products (a variation of up to a factor of three) as a function of the lots of hydrolysates used ("lot-to-lot variation"). This drawback affects the proliferation of the cells as well as the protein expression of each cell. US 2007/0212770 describes various animal protein-free and oligopeptide-free, chemically defined culture mediums that are useful for the large-scale production of recombinant protein biopharmaceuticals.

Hemostasis involves the interaction of various hemostatic reaction routes finally leading to thrombus formation. Thrombi are deposits of blood components on the surface of the vascular wall that mainly consist of aggregated blood platelets and insoluble cross-linked fibrin. Fibrin formation is the result of the restricted proteolysis of fibrinogen by thrombin, a coagulation enzyme. Thrombin is the end product of the coagulation cascade, a succession of zymogen activations occurring on the surfaces of activated blood platelets and leucocytes, and a variety of vascular cells (for a survey, cf. K. G. Mann et al., Blood, 1990, Vol. 76, pp. 1-16).

An important function in the coagulation cascade resides in the activation of Factor X by the complex of activated Factor IX (Factor IXa) and activated Factor VIII (Factor VIIIa). A deficiency or a dysfunction of the components of this complex is associated with the blood disease known as hemophilia (J. E. Sadler & E. W. Davie: Hemophilia A, Hemophilia B, and von Willebrand's Disease, in G. Stamatoyannopoulos et al., (Eds.): The molecular basis of blood

diseases. W.B. Saunders Co., Philadelphia, 1987, pp. 576-602). Hemophilia A is related to a deficiency of Factor VIII activity, whereas Hemophilia B is related to a Factor IX deficiency. Current treatment consists of a replacement therapy using pharmaceutical preparations comprised of the normal coagulation factor. Of these thrombopathies, Hemophilia A occurs more frequently, affecting approximately one out of 10,000 men. Replacement therapy in Hemophilia A patients involves the repeated administration of preparations containing normal Factor VIII by intravenous infusion. The interval between the infusions is a function of the degradation of the Factor VIII activity in blood circulation. The half-life of the Factor VIII activity after an infusion differs from one individual to another, ranging from 10 to 30 hours. Thus, a prophylactic therapy requires an infusion every two to three days. This constitutes a heavy load on the life of hemophilic patients, in particular, if the venous access has become difficult due to local citratization following frequent needle punctures for intravenous infusions.

It would be particularly advantageous if the frequency of infusions could be lowered by using Factor VIII having extended half-lives. It is well known in the art that the half-life of the non-activated Factor VIII heterodimer strongly depends on the presence of von Willebrand Factor, which exhibits a strong affinity to Factor VIII (yet not to Factor VIIIa) and serves as a carrier protein (J. E. Sadler and E. W. Davie: Hemophilia A, Hemophilia B and von Willebrand's disease, in G. Stamatoyannopoulos et al. (Eds.): The molecular basis of blood diseases. W.B. Saunders Co., Philadelphia, 1987, pp. 576-602). It is known that patients suffering from von Willebrand's disease type 3, who do not have a detectable von Willebrand Factor in their blood circulation, also suffer from a secondary Factor VIII deficiency. In addition, the half-life of intravenously administered Factor VIII in those patients is 2 to 4 hours, which is considerably shorter than the 10 to 30 hours observed in Hemophilia A patients. From these findings results that Factor VIII tends to a rapid clearance from the blood circulation and that this process is to some extent inhibited by complexation with its natural carrier, von Willebrand Factor.

Von Willebrand factor (vWF) is a glycoprotein circulating in plasma as a series of multimers typically ranging in size from about 500 to 20,000 kD (or 2 to 40 dimers of vWF). Dimers and multimeric forms of vWF are composed of 250 kD polypeptide subunits linked together by disulfide bonds. vWF mediates the initial platelet adhesion to the sub-endothelium of the damaged vessel wall; only the larger multimers also exhibiting hemostatic activity. Multimerized vWF binds to the platelet surface glycoprotein Gp1b α , through an interaction in the A1 domain of vWF, in order to facilitate platelet adhesion. It is assumed that endothelial cells secrete large polymeric forms of vWF and that those forms of vWF which have a low molecular weight (low molecular weight vWF) have arisen from proteolytic cleavage. The multimers having large molecular masses are stored in the Weibel-Palade bodies of the endothelial cells and liberated upon stimulation.

Reduction of FVIII binding activity, due to either reduced vWF protein levels or lowered FVIII binding affinity, results in one of three types of von Willebrand's Disease. In addition to, or alternatively, certain types of von Willebrand's disease are characterized by an increase or decrease in the level of Gp1b α -mediated platelet association, namely in Types 2A, 2B, and 2M (summarized in Castaman et al., Disorders of Hemostasis 88(1):94-108 (2003)). As such, the modulation of vWF interactions with both FVIII and Gp1b α

is a viable strategy for the treatment of both Haemophilia and von Willebrand's Disease.

Given the biological importance of vWF, there is a constant need in the art to improve ways for producing vWF for therapeutic applications. It is well known that vWF can be isolated from endogenous sources, such as human blood plasma. The isolated vWF is advantageous in that it has a high specific activity for carrying out its biological function and can, therefore, be used effectively as a therapeutic protein for treating related diseases, such as von Willebrand's disease. Typically, plasma vWF has a specific Ristocetin activity of about 100 mU/ μ g, but isolation from human blood plasma has disadvantages because, for example, the plasma can contain a variety of viruses, such as HIV and/or hepatitis viruses, which can be transferred to the patient. Furthermore, plasma is a limited resource and, thus, shortages of plasma can be problematic in providing enough vWF for treatment. As such, recombinant methods for producing vWF are advantageous in addressing some of the problems associated with relying on plasma as a source for vWF. For recombinant production, the full length of cDNA of vWF was cloned; the polypeptide corresponds to amino acid residues 23 to 764 of the full length prepro-vWF (Eikenboom et al (1995) Haemophilia 1, 77-90).

Unfortunately, vWF is a molecule with complex post-translational modifications. Also, the multimerization of the vWF dimers to large and ultralarge multimers in the Golgi apparatus is a challenge for expression in mammalian cells. For example, high molecular weight vWF expressed in cell culture of, e.g., human (primary) endothelial cells depends on the specific storage of ultralarge vWF molecules in Weibel-Palade bodies. Such cell cultures are not suitable for the production of therapeutic proteins. Other cell culture methods have been reported, and it is known that cell culture conditions can affect the production of vWF in a variety of ways. For instance, high concentrations of ammonium (NH_4^+) have been shown to disturb posttranslational modifications. Mayadas et al. (J. Biol. Chem., 264(23):13497-13503, 1989) demonstrated that levels of 25 mM ammonium resulted in reduced vWF multimerization in endothelial cells, which also negatively affects the specific Ristocetin activity of recombinant vWF. Reduction of multimerization is generally associated in reduction of activity, particularly specific Ristocetin activity, of recombinant vWF.

It still remains difficult to predict which parameters can positively or negatively affect production of a particular protein, especially complex glycoproteins like Factor VIII and vWF. For example, certain components of a cell culture medium have been shown to affect production of Factor VIII. As disclosed in U.S. Pat. No. 5,804,420, the addition of polyol, copper, and other trace metals can positively affect production yield of Factor VIII. As also described in WO 2009/086309, cell culture processes using copper in have been shown to improve production of Factor VIII. Expression of vWF in recombinant CHO cells has also been reported by Mignot et al. (1989). However, none of these examples provide information regarding the specific activity of vWF or its multimeric distribution.

The ADAMTS (a disintegrin and metalloproteinase with thrombospondin type I motifs) proteins are a family of metalloproteinases containing a number of conserved domains, including a zinc-dependant catalytic domain, a cysteine-rich domain, a disintegrin-like domain, and at least one, and in most cases multiple, thrombospondin type I repeats (for review, see Nicholson et al., BMC Evol Biol. 2005 Feb. 4; 5(1):11). These proteins, which are evolutionarily related to the ADAM and MMP families of metallo-

proteinases (Jones G C, *Curr Pharm Biotechnol.* 2006 February; 7(1):25-31), are secreted enzymes that have been linked to a number of diseases and conditions including thrombotic thrombocytopenic purpura (TTP) (Moake J L, *Semin Hematol.* 2004 January; 41(1):4-14), connective tissue disorders, cancers, inflammation (Nicholson et al.), and severe *Plasmodium falciparum* malaria (Larkin et al., *PLoS Pathog.* 2009 March; 5(3):e1000349). Because of these associations, the ADAMTS enzymes have been recognized as potential therapeutic targets for a number of pathologies (Jones G C, *Curr Pharm Biotechnol.* 2006 February; 7(1): 25-31). Accordingly, methods of producing large yields of ADAMTS proteins having high specific activities, which are free of contaminants such as viruses, BSE, and pathogens like *Mycoplasma* bacteria, are needed.

One ADAMTS family member, ADAMTS13, cleaves von Willebrand factor (vWF) between residues Tyr 1605 and Met 1606, a function responsible for the degradation of large vWF multimers in vivo. Loss of ADAMTS13 activity has been linked to a number of conditions, such as TTP (Moake J L, *Semin Hematol.* 2004 January; 41(1):4-14), acute and chronic inflammation (Chauhan et al., *J Exp Med.* 2008 Sep. 1; 205(9):2065-74), and most recently, severe *Plasmodium falciparum* malaria (Larkin et al., *PLoS Pathog.* 2009 March; 5(3):e1000349).

The ADAMTS13 protease is a 190 kDa glycosylated protein produced predominantly by the liver (Levy et al., *Nature.* 2001; 413:488-494; Fujikawa et al., *Blood.* 2001; 98:1662-1666; Zheng et al., *J Biol Chem.* 2001; 276:41059-41063; Soejima et al., *J Biochem (Tokyo).* 2001; 130:475-480; and Gerritsen et al., *Blood.* 2001; 98:1654-1661). Much like the higher order rVWF multimers, recombinant expression of the large ADAMTS13 in mammalian cell culture presents many challenges.

Therefore, there is a need to provide cell culture conditions, particularly large-scale manufacturing culture conditions, that provide consistent total protein yield and/or consistent total and specific activity of the proteins produced between different cell cultures. Consistency among cultures in large-scale manufacturing processes is of importance in the manufacture of therapeutic proteins. There is also a need for cell culture conditions for large-scale production of rVWF with a multimeric distribution and specific Ristocetin activity comparable or higher than VWF as it is present in normal human plasma. Similarly, as ADAMTS proteins have been implicated in a number of diseases and conditions, there is a need in the art for methods of large scale production of recombinant ADAMTS proteins having high specific activities, which are suitable for pharmaceutical formulation and administration. The present invention satisfies these and other needs in the art for the production of recombinant Von Willebrand Factor and recombinant ADAMTS13.

BRIEF SUMMARY OF INVENTION

In certain aspects, the present invention is based on the surprising finding that supplementation of cell culture media used to express recombinant Von Willebrand Factor (rVWF) and recombinant ADAMTS13 (rA13) results in significantly improved protein expression and enzymatic activity.

In a first aspect, the present invention provides a method for producing a recombinant Von Willebrand Factor (rVWF) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper to provide a final copper concentration of at least 2.4 µg/L; (c) providing one or more

cells comprising a nucleic acid encoding a rVWF protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rVWF is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 30 mU/µg rVWF.

In one embodiment of the methods provided above, the method further comprises a step of supplementing the basal cell culture media with a hydrolysate prior to culturing the one or more cells.

In one embodiment of the methods provided above, the hydrolysate is a plant hydrolysate. In a specific embodiment, the hydrolysate is a soy hydrolysate.

In one embodiment of the methods provided above, the basal cell culture media is an animal protein free culture media.

In one embodiment of the methods provided above, the basal cell culture media is a protein free culture media.

In one embodiment of the methods provided above, the basal cell culture media is a chemically defined culture media.

In one embodiment of the methods provided above, the final copper concentration of the copper supplemented basal cell culture media is at least 4 µg/L copper.

In one embodiment of the methods provided above, the final copper concentration of the copper supplemented basal cell culture media is between 2.4 µg/L and 20 µg/L copper.

In one embodiment of the methods provided above, the copper supplementing the basal cell culture media is provided as a copper salt, a copper chelate, or a combination thereof.

In one embodiment of the methods provided above, the copper salt is selected from the group consisting of copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide.

In one embodiment of the methods provided above, the one or more cells are mammalian cells. In a specific embodiment, the mammalian cells are CHO cells.

In one embodiment of the methods provided above, culturing the one or more cells comprises batch cultivation of the cells.

In one embodiment of the methods provided above, culturing the one or more cells comprises continuous cultivation of the cells. In a specific embodiment, the continuous cultivation of cells is performed in chemostatic mode. In another specific embodiment, the continuous cultivation of cells is performed in perfusion mode.

In one embodiment of the methods provided above, the one or more cells is cultured in at least 100 L of the supplemented basal cell culture media.

In one embodiment of the methods provided above, the cell density is maintained at less than 2.5×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 2.0×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 1.5×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the step of recovering at least a portion of the culture supernatant comprises filtration or centrifugation to remove cells from the portion of culture supernatant.

In one embodiment of the methods provided above, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 40 mU/µg rVWF. In a specific

embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μ g rVWF. In a more specific embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 60 mU/ μ g rVWF. In a more specific embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μ g rVWF. In a yet more specific embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 80 mU/ μ g rVWF.

In one embodiment of the methods provided above, at least 10% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers. In a specific embodiment, at least 15% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 20% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 25% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 30% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers.

In one embodiment of the methods provided above, the supernatant contains high molecular weight VWF multimers of 14 to 22 dimers.

In one embodiment of the methods provided above, the NH_4^+ content of the culture supernatant is maintained at a concentration below 10 mM.

In one embodiment of the methods provided above, the NH_4^+ content of the culture supernatant is maintained at a concentration below 4 mM.

In one embodiment of the methods provided above, rVWF is co-expressed with recombinant Factor VIII (rFVIII). In a specific embodiment, the method further comprises a step of purifying rVWF away from at least 50% of the rFVIII present in the recovered supernatant. In one embodiment, the ratio of rVWF to rFVIII after the purification step is at least 10:1.

In one embodiment of the methods provided above, the method further comprises a rVWF enrichment step.

In a second aspect, the present invention provides a recombinant Von Willebrand Factor (rVWF) composition prepared by a method provided herein.

In one embodiment of the compositions provided above, the composition further comprises recombinant Factor VIII (rFVIII). In a specific embodiment, the ratio of rVWF to rFVIII is at least 10:1.

In one embodiment of the compositions provided above, the composition is formulated for pharmaceutical administration. In a specific embodiment, the composition is formulated for intravenous administration.

In a third aspect, the present invention provides a cell culture supernatant comprising recombinant Von Willebrand Factor (rVWF), wherein the supernatant is prepared by a method provided herein.

In a fourth aspect, the present invention provides a cell culture supernatant comprising recombinant Von Willebrand Factor (rVWF), wherein at least 10% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers. In a specific embodiment, at least 15% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 20% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 25% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific

embodiment, at least 30% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers. In yet another specific embodiment of the supernatants provided above, the supernatant is prepared according to a method provided herein.

In a fifth aspect, the present invention provides a cell culture supernatant comprising recombinant Von Willebrand Factor (rVWF), wherein the supernatant contains at least 0.4 IU ristocetin cofactor activity per mL. In a specific embodiment, the supernatant contains at least 0.5 IU ristocetin cofactor activity per mL. In another specific embodiment, the supernatant contains at least 0.6 IU ristocetin cofactor activity per mL. In another specific embodiment, the supernatant contains at least 0.7 IU ristocetin cofactor activity per mL. In yet another specific embodiment of the supernatants provided above, the supernatant is prepared according to a method provided herein.

In a sixth aspect, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper to provide a final copper concentration of at least 1.0 μ g/L; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment of the methods provided above, the basal cell culture media is an animal protein free culture media.

In one embodiment of the methods provided above, the basal cell culture media is a protein free culture media.

In one embodiment of the methods provided above, the basal cell culture media is a chemically defined culture media.

In one embodiment of the methods provided above, the final copper concentration of the supplemented basal cell culture media is at least 1 μ g/L copper.

In one embodiment of the methods provided above, the final copper concentration of the supplemented basal cell culture media is at least 2 μ g/L copper.

In one embodiment of the methods provided above, the final copper concentration of the supplemented basal cell culture media is at least 4 μ g/L copper.

In one embodiment of the methods provided above, the final copper concentration of the supplemented basal cell culture media is between 1 μ g/L and 6 μ g/L copper.

In one embodiment of the methods provided above, the final copper concentration of the supplemented basal cell culture media is between 2 μ g/L and 4 μ g/L copper.

In one embodiment of the methods provided above, copper supplementing the basal cell culture media is provided as a copper salt, a copper chelate, or a combination thereof. In a specific embodiment, the copper salt is selected from the group consisting of copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide.

In one embodiment of the methods provided above, the one or more cells are mammalian cells. In a specific embodiment, the mammalian cells are CHO cells.

In one embodiment of the methods provided above, culturing the one or more cells comprises batch cultivation of the cells.

In one embodiment of the methods provided above, culturing the one or more cells comprises continuous cultivation of the cells. In a specific embodiment, the continuous cultivation of cells is performed in chemostatic mode. In another specific embodiment, the continuous cultivation of cells is performed in perfusion mode.

In one embodiment of the methods provided above, the one or more cells is cultured in at least 100 L of the supplemented basal cell culture media.

In one embodiment of the methods provided above, the cell density is maintained at less than 4.0×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 3.5×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 3.0×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 2.5×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 2.0×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the cell density is maintained at less than 1.5×10^6 cells per mL during the step of culturing the one or more cells.

In one embodiment of the methods provided above, the step of recovering at least a portion of the culture supernatant comprises filtration or centrifugation to remove cells from the portion of culture supernatant.

In one embodiment of the methods provided above, at least 2000 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment of the methods provided above, at least 2500 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment of the methods provided above, the recovered supernatant has a rA13 specific FRETS-VWF73 activity of at least 800 mU/ μ g.

In a preferred embodiment of the methods provided above, the recovered supernatant has a rA13 specific FRETS-VWF73 activity of at least 1200 mU/ μ g.

In a more preferred embodiment of the methods provided above, the recovered supernatant has a rA13 specific FRETS-VWF73 activity of at least 1600 mU/ μ g.

In one embodiment of the methods provided above, the NH_4^+ content of the culture supernatant is maintained at a concentration below 10 mM.

In one embodiment of the methods provided above, the NH_4^+ content of the culture supernatant is maintained at a concentration below 5 mM.

In one embodiment of the methods provided above, the NH_4^+ content of the culture supernatant is maintained at a concentration below 4 mM.

In one embodiment of the methods provided above, the method further comprises a rA13 enrichment step.

In a seventh aspect, the present invention provides a cell culture supernatant comprising recombinant ADAMTS13 (rA13), wherein the supernatant is prepared by a method provided herein.

In an eighth aspect, the present invention provides a cell culture supernatant comprising recombinant ADAMTS13 (rA13), wherein the supernatant contains at least 5 U FRETS-VWF73 activity per mL. In a specific embodiment,

the supernatant contains at least 6 U FRETS-VWF73 activity per mL. In another specific embodiment, the supernatant contains at least 7 U FRETS-VWF73 activity per mL. In another specific embodiment, the supernatant contains at least 8 U FRETS-VWF73 activity per mL. In another specific embodiment, the supernatant contains at least 9 U FRETS-VWF73 activity per mL. In another specific embodiment, the supernatant contains at least 10 U FRETS-VWF73 activity per mL. In yet another specific embodiment of the supernatants provided above, the supernatant is prepared according to a method provided herein.

In a ninth aspect, the present invention provides a cell culture supernatant comprising recombinant ADAMTS13 (rA13), wherein the supernatant contains at least 2 μ g rA13 per mL. In a specific embodiment, the supernatant contains at least 3 μ g rA13 per mL. In another specific embodiment, the supernatant contains at least 4 μ g rA13 per mL. In another specific embodiment, the supernatant contains at least 5 μ g rA13 per mL. In another specific embodiment, the supernatant contains at least 6 μ g rA13 per mL. In yet another specific embodiment of the supernatants provided above, the supernatant is prepared according to a method provided herein.

In a tenth aspect, the present invention provides a recombinant ADAMTS13 (rA13) composition prepared by a method according to any one of the methods described above.

In one embodiment of the compositions provided above, the composition is formulated for pharmaceutical administration. In a specific embodiment, the composition is formulated for intravenous administration.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1A-1B. (1A) Low resolution (1%) agarose gel electrophoresis of rVWF expressed in mammalian cell culture in the presence of low (1.0 μ g/L) and high (4.3 μ g/L) copper concentration, as described in Example 2. Note that culture Day 3 is equivalent to batch day 1 of Table 7 and Table 8. (1B) The relative amounts of VWF multimers having 1 to 10 dimers (band number 1) and more than 10 dimers (band number 2), as indicated by the bands defined in FIG. 1A, was quantitated by densitometric analysis.

FIG. 2A-2B. (2A) Interval plot of the average rVWF specific activity present in rVWF cell culture supernatants grown at high and low cell densities in the presence of high or low levels of copper. (2B) Interval plot of the average NH_4^+ concentration found in rVWF cell culture supernatants grown at high and low cell densities in the presence of high or low levels of copper.

FIG. 3A-3B. Supernatants from cell cultures expressing recombinant ADAMTS13 in the presence of increasing levels of copper were investigated by SDS-PAGE analysis. rA13 was visualized after SDS-PAGE by (3A) silver staining and (3B) anti-A13 western blotting.

FIG. 4. Plot of volumetric productivity (P Frets) data versus copper concentration showing an extrapolation (solid line) of optimal copper concentration effect on rA13 productivity.

FIG. 5A-5K. Bar graphs from continuous suspension (chemostat) cell cultures expressing rA13 over a culture time of 8 weeks comparing the effects of basal levels of copper (0.66 μ g/L) to that of cultures supplemented to a final concentration of 2 μ g/L copper. Each bar represents the mean data of one week of chemostatic culturing. The legend refers to the particular week represented in the data.

FIG. 6A-6B. (6A) Low resolution (1%) agarose gel electrophoresis of rVWF expressed in mammalian cell culture in the presence of low (1.0 $\mu\text{g/L}$) and high (4.3 $\mu\text{g/L}$) copper concentration under high and low cell densities as described in Example 3. Note that culture Day 8 and 17 (“CST8” and “CST17”) is equivalent to Day 8 and Day 17 of Table 10 to Table 13. (6B) The relative amounts of VWF multimers having 1 to 10 dimers and more than 10 dimers, as indicated by the bands defined in FIG. 6A, was quantitated by densitometric analysis.

DETAILED DESCRIPTION OF INVENTION

I. Introduction

Recombinant vWF (rVWF) and recombinant ADAMTS13 (rA13) can be produced by expression in large-scale mammalian cell cultures. However, the activity of these proteins when produced using standard cell culture conditions often varies across cell cultures, even when the general formulations of the media are not changed, and the specific activities of the recombinant proteins are often not equal to that of vWF and rA13 derived from blood plasma. In addition, rVWF expressed in mammalian cell cultures tends to produce protein compositions with low (under 10%) percentages of higher order multimers (higher order multimers include molecules containing more than 10 VWF dimers). These drawbacks in standard production methods of rVWF and rA13 are particularly problematic when developing cultures for large scale production (i.e., from 10 to over 20,000 liter cultures).

One potential source of the variability often seen in different cell culture batches is the presence of contaminants in components of the cell culture media. These contaminants may be present in different amounts in different batches, leading to variable results in the production of rVWF and rA13. After investigating the different contaminants found in various cell culture media additives, the present inventors found that the presence of hydrolysates leads to variation in copper concentrations in the cell media. Further investigation provided the surprising result that supplementing copper concentrations in cell media to produce a total copper concentration of at least about 1 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$ consistently increased the total and specific activity of rVWF and rA13 and/or could also lead to increased total protein yield. Thus, the present invention provides methods and compositions for high yield production of rVWF and rA13 proteins with high specific activity.

In one aspect, the present invention provides cell culture methods and compositions for producing large quantities of rVWF and rA13 with activity that is comparable to or higher than that seen with plasma derived vWF (pdVWF) or plasma derived ADAMTS13 (pdA13). In further aspects, the rVWF and rA13 proteins produced in accordance with the present invention show consistently higher activity than proteins produced using standard cell culture methods in media that has not been supplemented with copper or other supplements described in further detail herein. Advantageously, in certain embodiments of the methods and compositions provided herein, the rVWF and rA13 proteins produced in accordance with the present invention show consistently higher specific activity (i.e., U/mg protein) than proteins produced using standard cell culture methods in media that has not been supplemented with copper or other supplements described in further detail herein. Likewise, the methods provided herein for the production of rVWF and rA13 provide higher yields of activity per volume culture

(i.e., U/L/D), as compared to standard cell culture methods utilizing media that has not been supplemented with copper or other supplements described in further detail herein.

In a further aspect, the present invention provides cell culture methods in which a basal cell culture medium is supplemented with copper to result in a total concentration of at least about 1 $\mu\text{g/L}$. In other embodiments, the basal cell culture medium is supplemented with copper to result in a total concentration of at least about 2 $\mu\text{g/L}$. In yet other embodiments, the basal cell culture medium is supplemented with copper to result in a total concentration of at least about 1 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$. In some embodiments, the total concentration of copper is from about 1.5-4.5 $\mu\text{g/L}$. In certain embodiments, the cell culture medium is supplemented to result in about 1, 1.2, 1.4, 1.6, 1.8, 2.0, 2.2, 2.4, 2.6, 2.8, 3, 3.2, 3.4, 3.6, 3.8, 4, 4.2, 4.4, 4.6, 4.8, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 $\mu\text{g/L}$ copper, or more. Basal cell culture media generally have a trace copper concentration of under 1 $\mu\text{g/L}$.

In some embodiments, the present invention provides cell culture methods in which a basal cell culture medium is supplemented with from about 1.0 to about 20 $\mu\text{g/L}$ copper for production of rVWF. In further embodiments, the basal cell culture medium is supplemented with from about 1.5-15, 2.0-10, 2.5-8, 3.0-6, 4.0-5.0 $\mu\text{g/L}$ copper for production of rVWF. In still further embodiments, the basal cell culture medium may, in addition to the supplemented copper, also include one or more hydrolysates.

In other embodiments, the present invention provides cell culture methods in which a basal cell culture medium is supplemented with from about 1.5 to about 4 $\mu\text{g/L}$ copper for production of rA13. In further embodiments, the basal cell culture medium is supplemented with from about 1.6-3.8, 1.7-3.6, 1.8-3.4, 1.9-3.2, 2.0-3.0, 2.1-2.8, 2.2-2.6, 2.3-2.4 $\mu\text{g/L}$ copper for production of rA13. In still further embodiments, the basal cell culture medium may, in addition to the supplemented copper, also include one or more hydrolysates. In yet further embodiments, the basal cell culture medium includes, in addition to copper and/or one or more hydrolysates, about 1.0 to about 30 μM zinc. In still further embodiments, the basal cell culture medium further includes, in addition to copper and/or one or more hydrolysates and/or zinc, between about 0.5 to about 5.0 mM calcium.

In a still further aspect and in accordance with any of the above, the present invention provides cell culture methods in which the ammonium levels of the cell culture solution are low (under 10 mM). In certain embodiments, cell culture methods of the present invention utilize cell culture media having over 1, 2, 3, 4, or 5 $\mu\text{g/L}$ copper in combination with low levels of ammonium.

One of the advantages of the methods and compositions of the present invention is that they are amenable to large scale cell culture production. These large scale cell cultures are at least 10 L, 50 L, 100 L, 150 L, 200 L, 250 L, 500 L, 750 L, 1,000 L, 1,500 L, 2,000 L, 5,000 L, 10,000 L, or 20,000 liter cultures.

In certain aspects, the methods of the invention do not necessarily result in a higher amount of recombinant protein overall, but the recombinant protein (either rVWF or rA13) that is produced shows higher total and specific activity than is seen in proteins produced using standard cell cultures, particularly as compared to proteins produced in cell cultures in which the cell culture medium has not been supplemented with additional copper. In further aspects, the rVWF and rA13 proteins produced in cells cultured in copper supplemented media show consistently increased activity

per liter of cell culture as compared to cells cultured in basal cell culture media that has not been supplemented with copper. In still further aspects, the copper supplemented media of the present invention result in increased protein yield, increased number of cells in the culture, and/or increased total activity per liter of culture as compared to media that has not been supplemented with copper.

Still further advantages of methods and compositions of the invention is that they result a population of proteins containing a high percentage (over 10%) of highly multimerized rVWF.

Although much of the discussion herein regarding ADAMTS proteins is in terms of ADAMTS13 (A13), it will be appreciated that because all ADAMTS proteins share a common core domain architecture and common structure-function relationships, the methods and compositions described herein are applicable for production of any ADAMTS proteins, not only rA13.

II. Definitions

As used herein, “recombinant vWF” includes vWF obtained via recombinant DNA technology. In certain embodiments, vWF proteins of the invention can comprise a construct, for example, prepared as in WO 1986/06096 published on Oct. 23, 1986 and U.S. patent application Ser. No. 07/559,509, filed on Jul. 23, 1990, in the name of Ginsburg et al., which is incorporated herein by reference with respect to the methods of producing recombinant vWF. The vWF in the present invention can include all potential forms, including the monomeric and multimeric forms. It should also be understood that the present invention encompasses different forms of vWF to be used in combination. For example, the vWF of the present invention may include different multimers, different derivatives and both biologically active derivatives and derivatives not biologically active.

The term “recombinant” when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

In the context of the present invention, the recombinant vWF embraces any member of the vWF family from, for example, a mammal such as a primate, human, monkey, rabbit, pig, rodent, mouse, rat, hamster, gerbil, canine, feline, and biologically active derivatives thereof. In a preferred embodiment, the recombinant vWF is human vWF. Mutant and variant vWF proteins having activity are also embraced, as are functional fragments and fusion proteins of the vWF proteins. Furthermore, the vWF of the invention may further comprise tags that facilitate purification, detection, or both. The vWF described herein may further be modified with a therapeutic moiety or a moiety suitable imaging in vitro or in vivo.

The terms “highly multimeric vWF,” “high molecular weight vWF,” and “HMW vWF” may be used interchangeably and refer to covalently attached vWF multimers containing more than 10 vWF dimers. In certain embodiments, HMW vWF contains at least 11 vWF dimers, or at least 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or more vWF dimers.

As used herein, an “ADAMTS protein” refers to a polypeptide of the disintegrin and metalloproteinase with thrombospondin type I motifs family of metalloproteinases. Members of this family include the human proteins ADAMTS1 (NM_006988), ADAMTS2 (NM_014244; NM_021599), ADAMTS3 (NM_014243), ADAMTS4 (NM_005099), ADAMTS5 (NM_007038), ADAMTS6 (NM_014273), ADAMTS7 (NM_0142727), ADAMTS8 (NM_007037), ADAMTS9 (NM_182920; NM_182921; NM_020249), ADAMTS10 (NM_030957), ADAMTS12 (NM_030955), ADAMTS13 (NM_139025; NM_139026; NM_139027; NM_139028), ADAMTS14 (NM_139155; NM_080722), ADAMTS15 (NM_139055), ADAMTS16 (NM_139056), ADAMTS17 (NM_139057), ADAMTS18 (NM_199355; NM_139054), ADAMTS19 (NM_133638), and ADAMTS20 (NM_025003, NM_175851). ADAMTS proteins include both full-length proteins and partial polypeptides that display at least partial biological activity, for example, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more of the activity demonstrated by the full-length protein, in particular the protease activity demonstrated by the full length protein. In certain instances, an ADAMTS protein will be post-translationally modified either in vivo or in vitro, for example, by enzymatic or chemical means. It is understood that the ADAMTS proteins of the present invention include alternatively spliced isoforms, conservatively modified proteins, substantially identical proteins, homologues, and the like.

In the context of the present invention, an ADAMTS protein embraces any member of the ADAMTS family from, for example, a mammal such as a primate, human, monkey, rabbit, pig, rodent, mouse, rat, hamster, gerbil, canine, feline, and biologically active derivatives thereof. Mutant and variant ADAMTS proteins having activity are also embraced, as are functional fragments and fusion proteins of the ADAMTS proteins. Furthermore, the ADAMTS proteins of the invention may further comprise tags that facilitate purification, detection, or both. The ADAMTS proteins described herein may further be modified with a therapeutic moiety or a moiety suitable imaging in vitro or in vivo.

As used herein, an “ADAMTS13 protein” refers to any protein or polypeptide with ADAMTS13 activity, particularly the ability to cleave the peptide bond between residues Tyr-842 and Met-843 of vWF. In an exemplary embodiment, an ADAMTS13 protein refers to a polypeptide comprising an amino acid sequence that is highly similar to that of NP 620594 (ADAMTS13 isoform 1, preproprotein) or amino acids 75 to 1427 of NP 620594 (ADAMTS13 isoform 1, mature polypeptide). In another embodiment, an ADAMTS13 protein refers to a polypeptide comprising an amino acid sequence that is highly similar to that of NP 620596 (ADAMTS13 isoform 2, preproprotein) or amino acids 75 to 1371 of NP 620594 (ADAMTS13 isoform 2, mature polypeptide). In yet another embodiment, ADAMTS13 proteins include polypeptides comprising an amino acid sequence highly similar to that of NP 620595 (ADAMTS13 isoform 3, preproprotein) or amino acids 75 to 1340 of NP 620595 (ADAMTS13 isoform 1, mature polypeptide). As used herein, an ADAMTS13 protein includes natural variants with vWF cleaving activity and artificial constructs with vWF cleaving activity. As used in the present invention, ADAMTS13 encompasses any natural variants, alternative sequences, isoforms or mutant proteins that retain some basal activity. Examples of ADAMTS13 mutations found in the human population include, without limitation, R7W, V88M, H96D, R102C, R193W, T196I, H234Q, A250V, R268P, W390C, R398H, Q448E, Q456H, P457L,

C508Y, R528G, P618A, R625H, I673F, R692C, A732V, S903L, C908Y, C951G, G982R, C1024G, A1033T, R1095W, R1123C, C1213Y, T1226I, G1239V, R1336W, many of which have been found associated with thrombotic thrombocytopenic purpura (TTP). ADAMTS13 proteins also includes polypeptides containing post-translational modifications. For example, ADAMTS13 has been shown to be modified by N-acetylglucosamine (GlcNAc) at residues 614, 667, and 1354, and it has been predicted that residues 142, 146, 552, 579, 707, 828, and 1235 may also be modified in this fashion.

Proteolytically active recombinant ADAMTS13 may be prepared by expression in mammalian cell cultures, as described in Plaimauer et al., (2002, *Blood*. 15; 100(10): 3626-32) and US 2005/0266528, the disclosures of which are herein incorporated by reference in their entireties for all purposes. Methods for the expression of recombinant ADAMTS13 in cell culture are disclosed in Plaimauer B, Scheiflinger F. (*Semin Hematol*. 2004 January; 41(1):24-33) and US 2011/0086413, the disclosures of which are herein incorporated by reference in their entireties for all purposes.

As used herein, the term "biologically active derivative", when used in the context of an ADAMTS protein, also embraces polypeptides obtained via recombinant DNA technology. This may include any method known in the art for (i) the production of recombinant DNA by genetic engineering, e.g., via reverse transcription of RNA and/or amplification of DNA, (ii) introducing recombinant DNA into prokaryotic or eukaryotic cells by transfection, i.e., via electroporation or microinjection, (iii) cultivating said transformed cells, e.g., in a continuous or batch-wise manner, (iv) expressing an ADAMTS protein, e.g., constitutively or upon induction, and (v) isolating said ADAMTS protein, e.g., from the culture medium or by harvesting the transformed cells, in order to (vi) obtain substantially purified recombinant ADAMTS protein, e.g., via ion exchange chromatography, size exclusion chromatography, affinity chromatography, hydrophobic interaction chromatography, and the like. The term "biologically active derivative" includes also chimeric molecules such as e.g. an ADAMTS protein, or functional fragment thereof, in combination with a second polypeptide, e.g., an immunoglobulin Fc domain or an albumin domain, in order to improve the biological/pharmacological properties such as e.g., half life of the ADAMTS protein in the circulation system of a mammal, particularly a human.

The terms "isolated," "purified," or "biologically pure" refer to material that is substantially or essentially free from components that normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. In one embodiment, rVWF is the predominant species present in a preparation is substantially purified. In another embodiment, rA13 is the predominant species present in a preparation is substantially purified. The term "purified" in some embodiments denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. In other embodiments, it means that the nucleic acid or protein is at least 50% pure, more preferably at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or more pure. "Purify" or "purification" in other embodiments means removing at least one contaminant from the composition to be purified. In this sense, purification does not require that the purified compound be homogeneous, e.g., 100% pure.

The biological activity of vWF can be measured by known in vitro assays. For example, the Ristocetin Cofactor assay is based on the agglutination of fresh or formalin-fixed platelets induced by the antibiotic ristocetin in the presence of vWF. The degree of platelet agglutination depends on the vWF concentration and can be measured by the turbidimetric method, e.g. by use of an aggregometer (Weiss et al., *J. Clin. Invest.* 52: 2708-2716, 1973; Macfarlane et al., *Thromb. Diath. Haemorrh.* 34: 306-308, 1975). As provided herein, the specific Ristocetin Cofactor activity of the vWF of the present invention is described in terms of mU/ μ g of vWF, as measured using in vitro assays.

As used herein, "one unit of ADAMTS activity" is defined as the amount of activity in 1 mL of pooled normal human plasma, regardless of the assay being used. For example, when the ADAMTS protein is ADAMTS13, one unit of ADAMTS13 FRET-VWF73 activity is the amount of activity needed to cleave the same amount of FRET-VWF73 substrate (Kokame et al., *Br J Haematol*. 2005 April; 129(1):93-100) as is cleaved by one mL of pooled normal human plasma. Conveniently, ADAMTS13 activity may be determined by functional assays, such as functional assays employing modified von Willebrand factor peptides as substrate for ADAMTS13 (Triodi et al. *J Thromb Haemost.* 2008 September; 6(9): 1534-41). A preferred method of determining recombinant human ADAMTS13 activity is disclosed in Gerritsen et al. (Assay of von Willebrand factor (vWF)-cleaving protease based on decreased collagen binding affinity of degraded vWF: a tool for the diagnosis of thrombotic thrombocytopenic purpura (TTP). *Thromb Haemost* 1999; 82: 1386-1389). In one embodiment, to be considered as an ADAMTS13 protein as defined above, a polypeptide or protein must have at least 1% of the vWF cleaving activity of native ADAMTS13. In other embodiments, an ADAMTS13 protein will contain at least 10% of the activity of native ADAMTS13. In yet other embodiments, an ADAMTS13 protein will contain at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of the activity of native ADAMTS13. The quantity of an ADAMTS13 protein may also be determined by measurement of an ADAMTS13 antigen, for example using the ELISA method disclosed in Rieger et al., (2006, *Thromb Haemost.* 2006 95(2):212-20). A well established convention in the art is that 1 mL of pooled normal human plasma contains 1 μ g of ADAMTS13. Thus, the convention in the art is that 1 μ g of plasma-derived ADAMTS13 has one unit of ADAMTS13 activity.

The terms "cell culture solution," "cell culture medium or media," and "cell culture supernatant" refer to aspects of cell culture processes generally well known in the art. In the context of the present invention, a cell culture solution can include cell culture media and cell culture supernatant. The cell culture media are externally added to the cell culture solution, optionally together with supplements, to provide nutrients and other components for culturing cells expressing rVWF or rA13. The cell culture supernatant refers to a cell culture solution comprising the nutrients and other components from the cell culture medium as well as products released, metabolized, and/or excreted from the cells during culture, but not the cells themselves. Thus in one context, a cell culture supernatant can refer to the liquid phase of a cell culture solution (i.e., the cell culture solution excluding the cells). For example, the ammonium concentration of a culture supernatant generally refers to the ammonium concentration present in the cell culture solution. In other contexts, a cell culture supernatant refers to a cell

culture solution from which the cells have been removed (i.e., a recovered cell culture supernatant).

As used herein, the terms “vitamin B3,” “nicotinamide,” “niacinamide,” “niacin,” and “nicotinic acid” may be used interchangeably to refer to any member of the B3 family of vitamins. Accordingly, any member of this family may be used to supplement medium used in the methods of the present invention.

As used herein, the term “chemically defined medium” or “chemically defined media” refers to a synthetic growth medium in which the identity and concentration of all the components are known. Chemically defined mediums do not contain bacterial, yeast, animal, or plant extracts, although they may or may not include individual plant or animal-derived components (e.g., proteins, polypeptides, etc.). Non-limiting examples of commercially available chemically defined mediums include, various Dulbecco’s Modified Eagle’s (DME) mediums (Sigma-Aldrich Co; SAFC Biosciences, Inc), Ham’s Nutrient Mixture (Sigma-Aldrich Co; SAFC Biosciences, Inc), combinations thereof, and the like. Methods of preparing chemically defined culture mediums are known in the art, for example in U.S. Pat. Nos. 6,171,825 and 6,936,441, WO 2007/077217, and U.S. Patent Application Publication Numbers 2008/0009040 and 2007/0212770, the disclosures of which are incorporated herein by reference in their entireties for all purposes.

As used herein, the term “oligopeptide-free culture medium” or “oligopeptide-free culture media” refers to a protein-free medium that does not comprise oligopeptides, such as, e.g., oligopeptides derived from a protein hydrolysate. In one embodiment, the medium does not comprise oligopeptides having twenty or more amino acids. In one embodiment of the present invention, the medium does not comprise oligopeptides having fifteen or more amino acids. In another embodiment of the invention, the medium does not comprise oligopeptides having ten or more amino acids. In one embodiment the medium does not comprise oligopeptides having seven or more amino acids. In another embodiment the medium does not comprise oligopeptides having five or more amino acids. In still another embodiment the medium does not comprise oligopeptides having three or more amino acids. According to a further embodiment of the present invention, the medium does not comprise oligopeptides having two or more amino acids. Methods of preparing oligopeptide-free culture medium are known in the art, for example in U.S. Pat. Nos. 6,171,825 and 6,936,441, WO 2007/077217, and U.S. Patent Application Publication Numbers 2008/0009040 and 2007/0212770, the disclosures of which are incorporated herein by reference in their entireties for all purposes.

As used herein, the term “serum-free culture medium” or “serum-free culture media” refers to a culture medium that is not supplemented with an animal serum. Although oftentimes serum-free mediums are chemically defined mediums, serum-free mediums may be supplemented with discrete animal or plant proteins or protein fractions. Methods of preparing serum-free culture medium are known in the art, for example in U.S. Pat. Nos. 6,171,825 and 6,936,441, WO 2007/077217, and U.S. Patent Application Publication Numbers 2008/0009040 and 2007/0212770, the disclosures of which are incorporated herein by reference in their entireties for all purposes.

As used herein, the term “animal protein-free culture medium” or “animal protein-free culture media” refers to a culture medium that is not supplemented with an animal serum, protein, or protein fraction. Although oftentimes animal protein-free culture mediums are chemically defined

mediums, animal protein-free culture mediums may contain plant or yeast hydrolysates. Methods of preparing animal protein-free culture medium are known in the art, for example in U.S. Pat. Nos. 6,171,825 and 6,936,441, WO 2007/077217, and U.S. Patent Application Publication Numbers 2008/0009040 and 2007/0212770, the disclosures of which are incorporated herein by reference in their entireties for all purposes.

As used herein, the term “basal cell culture medium” or “basal cell culture media” refers to a chemically defined culture medium, an oligopeptide-free culture medium, a serum-free culture medium, or an animal protein-free culture medium that has not been supplemented with a hydrolysate, e.g., a plant or yeast hydrolysate. basal mediums are well known in the art, e.g., DMEM, Ham’s F12, DMEM/Ham’s F12, Medium 199, McCoy, or RPMI. The basal medium can include a number of ingredients, including amino acids, vitamins, organic and inorganic salts, and sources of carbohydrate. Each ingredient can be present in an amount that supports the cultivation of a cell, such amounts being generally known to a person skilled in the art. The medium can include auxiliary substances, such as buffer substances, e.g., sodium bicarbonate, antioxidants, stabilizers to counteract mechanical stress, or protease inhibitors. If necessary, a non-ionic surfactant such as copolymers and/or mixtures of polyethylene glycols and polypropylene glycols can be added.

III. Cell Culture Media and Cell Culture Supernatant

One aspect of the present invention relates to cell culture media for producing rVWF and/or rA13 that have increased activity as compared to rVWF and rA13 produced with basal cell culture media. In one aspect, the present invention relates to cell culture media for producing rVWF and/or rA13 in which basal cell culture media is supplemented with one or more additional substances. In specific embodiments and as is discussed in further detail below, cell culture conditions of the present invention include basal cell culture media supplemented to have at least 1.0 µg/L copper. In further embodiments, cell culture media of use and supernatants derived from the processes in the present invention also comprise low levels (under 10 mM) of ammonium. In a particular embodiment, the cell culture conditions used for expressing rVWF and/or rA13 are controlled such that the cell culture supernatant maintains a low level of ammonium, i.e., less than 10 mM and preferably less than 5 mM.

The culture media of the present invention can be based on a suitable basal medium well known in the art, e.g., DMEM, Ham’s F12, DMEM/Ham’s F12, Medium 199, McCoy, or RPMI. The basal medium can include a number of ingredients, including amino acids, vitamins, organic and inorganic salts, and sources of carbohydrate. Each ingredient can be present in an amount that supports the cultivation of a cell, such amounts being generally known to a person skilled in the art. The medium can include auxiliary substances, such as buffer substances, e.g., sodium bicarbonate, antioxidants, stabilizers to counteract mechanical stress, or protease inhibitors. If necessary, a non-ionic surfactant such as copolymers and/or mixtures of polyethylene glycols and polypropylene glycols can be added.

Generally, basal media contain less than 1 µg/L of copper—for example, DMEM/Ham’s F12 has a copper concentration of about 0.3 µg/L. Such concentrations of copper do

not provide enough copper ions to support production of rVWF and rA13 proteins of the present invention, which show high specific activity.

Copper can be provided to cell culture media of the present invention through a variety of ways, such as through addition of a medium supplement. In some embodiments, the medium supplement can contain hydrolysate, which can be provided to increase the copper concentration in the media. Hydrolysates can include any hydrolysate well known in the art, such as plant hydrolysates, soy hydrolysates, and wheat gluten hydrolysate. In certain embodiments, addition of hydrolysate can contribute an increased copper concentration of about 0.2 to about 10 µg/L of Cu²⁺. In some embodiments, the amount of copper provided by the hydrolysate can depend on the amount of copper in the hydrolysate as well as the amount of hydrolysate added. The copper content of a hydrolysate can be determined by elemental analysis, e.g., atom adsorption spectroscopy (GFAA: graphite furnace atomic adsorption), or mass spectroscopy methods (e.g., ICP-MS).

In certain embodiments, copper can be provided to the culture media alone or in addition to hydrolysate by providing a medium supplement including a suitable copper salt or copper chelate. Suitable copper salts can include but are not limited to copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide. Suitable copper chelators can include but are not limited to albumin, ethylenediaminetetraacetic acid (EDTA), polyamine chelating agents, ethylenediamine, diethylenetriamine, triethylenetetramine, triethylenediamine, tetraethylenepentamine, aminoethyl ethanolamine, aminoethylpiperazine, pentaethylenhexamine, triethylenetetramine-hydrochloride, tetraethylenepentamine-hydrochloride, pentaethylenhexamine-hydrochloride, tetraethylpentamine, captopril, penicilamine, N,N'-bis(3-aminopropyl)-1,3-propanediamine, N,N-Bis (2 animoethyl) 1,3 propane diamine, 1,7-dioxa-4,10-diazacyclododecane, 1,4,8,11-tetraaza cyclotetradecane-5,7-dione, 1,4,7-triazacyclononane trihydrochloride, 1-oxa-4,7,10-triazacyclododecane, 1,4,8,12-tetraaza cyclopentadecane, and 1,4,7,10-tetraaza cyclododecane.

In certain embodiments, basal cell culture media is supplemented with copper to result in a total copper concentration of about 1.0 to about 20 µg/L. In a specific embodiment, the basal cell media is supplemented with copper to a final concentration of between about 1.0 to about 10 µg/L. In further embodiments, the basal cell culture media is supplemented to result in a final copper concentration of about 1.0-5.0, 1.2-4.0, 1.3-3.0, 1.4-2.9, 1.5-2.8, 1.6-2.7, 1.7-2.6, 1.8-2.5, 1.9-2.4, 2.0-2.3, 2.1-2.2 µg/L copper. In further embodiments, the basal cell culture media used in methods of the present invention are supplemented

to result in about 1.2-9.5, 1.4-9, 1.6-8.5, 1.8-8, 2.0-7.5, 2.2-7, 2.4-6.5, 2.6-6.0, 2.8-5.5, 3.0-5.0, 3.2-4.5, 3.4-4, and 2-4 µg/L copper. In yet other embodiments, the basal cell culture media used in methods of the present invention are supplemented to result in about 1-6, 2-5, 3-4 µg/L copper. In one embodiment, the basal cell culture media is supplemented with copper to result in a total copper concentration of at least 1 µg/L. In another embodiment, the basal cell culture media is supplemented with copper to result in a total copper concentration of at least 2 µg/L. In yet another embodiment, the basal cell culture media is supplemented with copper to result in a total copper concentration of at least 4 µg/L. In certain embodiments, the basal cell culture media is supplemented with copper to result in a total copper concentration of at least 1 µg/L, or at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 µg/L copper, or more. In certain embodiments and as discussed in further detail herein, cultures for producing rA13 may contain from about 2-4 µg/L of copper, whereas cultures for producing rVWF may contain at least 2 µg/L of copper.

The concentrations indicated above are the respective concentrations of pure copper, in cupric form (Cu²⁺). If a copper derivative, e.g., a hydrated salt, or a compound comprising copper, e.g., a copper chelator, is used, the amount of derivative or chelator is added such that the final concentration of the copper is in the ranges described herein. For example, 2 µg/L of CuSO₄·5H₂O is equivalent to a copper concentration of about 0.51 µg/L (without sulfate and 5H₂O).

Advantageously, it has been found that the use of cell culture medium in cell culture processes resulting in low ammonium (NH₄⁺) concentrations in the cell culture solution (i.e., in the culture supernatant) results in the expression of recombinant VWF and/or rA13 with higher specific activities. Accordingly, in certain embodiments, the NH₄⁺ concentration of the supernatant is no higher than 10 mM. In a preferred embodiment, the NH₄⁺ concentration of the supernatant is no higher than 5 mM. In a preferred embodiment, the NH₄⁺ concentration of the supernatant is no higher than 4 mM. In yet other embodiments, the NH₄⁺ concentration of the supernatant is no higher than 10 mM, 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less.

Accordingly, in certain embodiments, the methods and compositions provided herein rely on the use of basal cell culture media supplemented with copper (e.g., to a final concentration of at least 2 µg/L) used in a process that results in an NH₄⁺ concentration of no higher than 10 mM in the supernatant. In yet other embodiments, the basal cell culture medium is supplemented to provide a final copper and ammonium concentration according to any one of variations 1 to 440, as provided in Table 1.

TABLE 1

Exemplary embodiments of copper and ammonium concentrations present in culture media and supernatant useful for the expression of a recombinant proteins as provided herein.			Ammonium Concentration									
		N.D.	NMT 10 mM	NMT 9 mM	NMT 8 mM	NMT 7 mM	NMT 6 mM	NMT 5 mM	NMT 4 mM	NMT 3 mM	NMT 2 mM	NMT 1 mM
Copper Concentration	AL	Var. 1	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1 µg/L		41	81	121	161	201	241	281	321	361	401
	AL	Var. 2	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2 µg/L		42	82	122	162	202	242	282	322	362	402
AL	Var. 3	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	3 µg/L		43	83	123	163	203	243	283	323	363	403

TABLE 1-continued

Exemplary embodiments of copper and ammonium concentrations present in culture media and supernatant useful for the expression of a recombinant proteins as provided herein.											
		Ammonium Concentration									
	N.D.	NMT 10 mM	NMT 9 mM	NMT 8 mM	NMT 7 mM	NMT 6 mM	NMT 5 mM	NMT 4 mM	NMT 3 mM	NMT 2 mM	NMT 1 mM
AL	Var. 4	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
4 µg/L		44	84	124	164	204	244	284	324	364	404
AL	Var. 5	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
5 µg/L		45	85	125	165	205	245	285	325	365	405
AL	Var. 6	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
6 µg/L		46	86	126	166	206	246	286	326	366	406
AL	Var. 7	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
7 µg/L		47	87	127	167	207	247	287	327	367	407
AL	Var. 8	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
8 µg/L		48	88	128	168	208	248	288	328	368	408
AL	Var. 9	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
9 µg/L		49	89	129	169	209	249	289	329	369	409
AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
10 µg/L	10	50	90	130	170	210	250	290	330	370	410
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
1 µg/L	11	51	91	131	171	211	251	291	331	371	411
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
1.5 µg/L	12	52	92	132	172	212	252	292	332	372	412
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
2 µg/L	13	53	93	133	173	213	253	293	333	373	413
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
2.5 µg/L	14	54	94	134	174	214	254	294	334	374	414
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
3 µg/L	15	55	95	135	175	215	255	295	335	375	415
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
3.5 µg/L	16	56	96	136	176	216	256	296	336	376	416
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
4 µg/L	17	57	97	137	177	217	257	297	337	377	417
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
4.5 µg/L	18	58	98	138	178	218	258	298	338	378	418
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
5 µg/L	19	59	99	139	179	219	259	299	339	379	419
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
5.5 µg/L	20	60	100	140	180	220	260	300	340	380	420
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
6 µg/L	21	61	101	141	181	221	261	301	341	381	421
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
7 µg/L	22	62	102	142	182	222	262	302	342	382	422
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
8 µg/L	23	63	103	143	183	223	263	303	343	383	423
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
9 µg/L	24	64	104	144	184	224	264	304	344	384	424
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
10 µg/L	25	65	105	145	185	225	265	305	345	385	425
1-20 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
26		66	106	146	186	226	266	306	346	386	426
2-20 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
27		67	107	147	187	227	267	307	347	387	427
1-10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
28		68	108	148	188	228	268	308	348	388	428
2-10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
29		69	109	149	189	229	269	309	349	389	429
1-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
30		70	110	150	190	230	270	310	350	390	430
2-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
31		71	111	151	191	231	271	311	351	391	431
3-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
32		72	112	152	192	232	272	312	352	392	432
4-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
33		73	113	153	193	233	273	313	353	393	433
1-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
34		74	114	154	194	234	274	314	354	394	434
2-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
35		75	115	155	195	235	275	315	355	395	435
3-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
36		76	116	156	196	236	276	316	356	396	436
4-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
37		77	117	157	197	237	277	317	357	397	437
1-4 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
38		78	118	158	198	238	278	318	358	398	438
2-4 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
39		79	119	159	199	239	279	319	359	399	439

TABLE 1-continued

Exemplary embodiments of copper and ammonium concentrations present in culture media and supernatant useful for the expression of a recombinant proteins as provided herein.											
		Ammonium Concentration									
N.D.		NMT 10 mM	NMT 9 mM	NMT 8 mM	NMT 7 mM	NMT 6 mM	NMT 5 mM	NMT 4 mM	NMT 3 mM	NMT 2 mM	NMT 1 mM
3-4 µg/L	Var. 40	Var. 80	Var. 120	Var. 160	Var. 200	Var. 240	Var. 280	Var. 320	Var. 360	Var. 400	Var. 440

*ND = not defined

*NMT = no more than

*AL = at least

In some embodiments, the copper supplemented media of the present invention are produced by supplementing basal media that is animal protein-free and/or chemically defined. Methods of preparing animal protein-free and chemically defined culture mediums are known in the art, for example in U.S. Pat. Nos. 6,171,825 and 6,936,441, WO 2007/077217, and U.S. Patent Application Publication Numbers 2008/0009040 and 2007/0212770, the disclosures of which are incorporated herein by reference in their entireties for all purposes. In one embodiment, the basal culture medium used in the methods described herein is animal protein-free or oligopeptide-free medium. In certain embodiments, the culture medium may be chemically defined. In certain embodiments, the culture media may contain at least one polyamine at a concentration of about 0.5 mg/L to about 10 mg/L.

In further embodiments and in addition to any of the description provided above, culture media of the invention are provided in which a basal medium is supplemented with copper and at least one of calcium, zinc, and/or vitamin B3. In certain embodiments, the medium may be an animal protein-free, oligopeptide-free, or chemically defined medium. In certain embodiments, the animal protein-free or oligopeptide free medium is prepared as taught in U.S. Pat. Nos. 6,171,825 and 6,936,441, WO 2007/077217, and U.S. Patent Application Publication Numbers 2008/0009040 and 2007/0212770, the disclosures of which are incorporated herein by reference in their entireties for all purposes, both of which are incorporated herein by reference in their entireties for all purposes, and supplemented with additional copper and optionally one or more of calcium, zinc, and vitamin B3. In a specific embodiment, the chemically defined culture medium may be similar to a Dulbecco's Modified Eagle's Media/Ham's F12 1:1 mixture (DMEM/Ham's F12), which has been supplemented with additional copper and optionally calcium, zinc, and/or vitamin B3, in order to increase the specific activity of rVWF or rA13 expressed in a cell cultured in the medium. In yet other embodiments, the culture medium is animal component free. In another embodiment, the culture medium contains protein, e.g., animal protein from serum such as fetal calf serum. In another embodiment, the culture has recombinant proteins exogenously added. In another embodiment, the proteins are from a certified pathogen free animal.

In certain embodiments, the culture media contains at least one polyamine at a concentration of at or about between 0.5 mg/L and 30 mg/L. In another embodiment, the culture medium contains at least one polyamine at or about between 0.5 mg/L and 10 mg/L. In one embodiment, the culture medium contains at least one polyamine at or about between 2 mg/L and 8 mg/L. In certain embodiments the polyamine is from the group of ornithine, putrescine, sper-

mine or spermidine, or the like. In a preferred embodiment, the polyamine is putrescine. In a specific embodiment, the culture medium contains at or about between 2 mg/L and 8 mg/L putrescine.

In one embodiment, the culture media contains at least one polyamine at a concentration of at or about between 0.5 mg/L and 30 mg/L and a copper and ammonium combination according to any one of variations 1 to 440, as set forth in Table 1. In another embodiment, the culture medium contains at least one polyamine at or about between 0.5 mg/L and 10 mg/L and a copper and ammonium combination according to any one of variations 1 to 440, as set forth in Table 1. In one embodiment, the culture medium contains at least one polyamine at or about between 2 mg/L and 8 mg/L and a copper and ammonium combination according to any one of variations 1 to 440, as set forth in Table 1. In certain embodiments the polyamine is from the group of ornithine, putrescine, spermine or spermidine, or the like. In a preferred embodiment, the polyamine is putrescine. In a specific embodiment, the culture medium contains at or about between 2 mg/L and 8 mg/L putrescine and a copper and ammonium combination according to any one of variations 1 to 440, as set forth in Table 1.

In further aspects, in addition to copper, cell culture media of use in the present invention may further include one or more of: additional calcium, zinc, one or more vitamins, and any combination thereof.

Generally, any calcium salt may be used to supplement the media of the invention, non-limiting examples of acceptable salts include CaCl_2 , CaCl_2 , $\text{CaFPO}_3 \cdot 2\text{H}_2\text{O}$, CaI_2 , CaBr_2 , $(\text{C}_2\text{H}_3\text{O}_2)_2\text{Ca}$, $(\text{CHO}_2)_2\text{Ca}$, $(\text{C}_6\text{H}_7\text{O}_6)_2\text{Ca}$, $(\text{C}_6\text{H}_5\text{O}_7)_2\text{Ca} \cdot 2\text{H}_2\text{O}$, and the like. In certain embodiments, a pharmaceutically acceptable salt of calcium is used to supplement the culture mediums of the invention.

Generally, any zinc salt may be used to supplement the media of the invention, non-limiting examples of acceptable salts include, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, $\text{ZnSO}_3 \cdot 2\text{H}_2\text{O}$, $(\text{C}_6\text{H}_5\text{O}_7)_2\text{Zn} \cdot 2\text{H}_2\text{O}$, ZnBr_2 , $\text{ZnBr}_2 \cdot 2\text{H}_2\text{O}$, ZnCl_2 , $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$, $\text{Zn}(\text{H}_2\text{PO}_4)_2 \cdot \text{H}_2\text{O}$, $(\text{C}_2\text{H}_3\text{O}_2)_2\text{Zn} \cdot \text{H}_2\text{O}$, and the like. In certain embodiments, a pharmaceutically acceptable salt of zinc is used to supplement the culture mediums of the invention. In other embodiments, a zinc containing peptide or protein preparation, for example insulin, may be used to supplement the culture provided herein.

In still further aspects, basal cell media supplemented with copper and one or more of the additional materials discussed above may further be used in cultures with low ammonium levels in the supernatant. In certain embodiments, supplemented cell culture media of use in the present invention result in ammonium levels in the cell culture solution of under 10 mM. In further embodiments, the supplemented cell culture media of the present invention is

used with cell culture ammonium levels from about 0.5-9.5, 1.0-9.0, 1.5-8.5, 2.0-8.0, 2.5-7.5, 3.0-7.0, 3.5-6.5, 4.0-6.0, 4.5-5.5 mM.

In one embodiment, the copper and ammonium concentrations of a cell culture media and a cell culture supernatant are maintained for an extended period of time during the manufacturing process. In a specific embodiment, the copper and ammonium concentrations of a cell culture are maintained for the duration of a manufacturing process, i.e., during the time in which rVWF or rA13 is being expressed and recovered from a large-scale cell culture. In certain embodiments, the copper and ammonium concentrations are maintained in the culture solution at a level according to any one of variations 1 to 440, as set forth in Table 1. In a preferred embodiment the copper and ammonium concentrations are maintained for the entire time of such a production process.

In some embodiments, the culture medium provided by the invention may be provided in a liquid or a dry or powder form. The medium may be pre-aliquoted in an amount suitable for single use or provided in a larger quantity that may be used for more than one cell-culture. Generally, the medium of the invention will be provided in a sterile fashion.

Specific details of cell culture media of use for producing rVWF or rA13 are discussed below. Although the following are discussed with respect to either rVWF or rA13, it will be appreciated that any of the discussion provided below with respect to rVWF is applicable for rA13, and vice versa.

A. Recombinant VWF Cell Culture Media

One aspect of the present invention relates to a cell culture solution for producing recombinant vWF, more specifically, high molecular weight vWF having a high specific activity, which are further described herein. In one embodiment, the present invention provides a cell culture solution for producing high molecular weight, recombinant vWF, comprising a cell culture medium comprising a copper concentration of at least about 2.4 $\mu\text{g/L}$ and a plurality of cells expressing highly multimeric vWF comprising about 14 to about 22 dimers and a specific Ristocetin co-factor activity of at least about 30 mU/ μg .

In one embodiment, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In a preferred embodiment, the cell culture comprises an ammonium concentration of no more than 5 mM. In yet other embodiments, the cell culture comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less. In yet other embodiments, the cell culture has a copper and ammonium concentration according to any one of variations 1 to 440, as set forth in Table 1. In certain embodiments, the ammonium concentration of the cell culture is maintained for an extended period at a concentration as provided above. For example, in one embodiment, the ammonium concentration is maintained at a low concentration for at least 3 days, or at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In a specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 5 mM for at least 14 days. In another specific embodiment, the ammonium concentration of the cell cul-

ture is maintained at an ammonium concentration of no more than 4 mM for at least 14 days. In yet another embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rVWF).

In one embodiment of the invention, the cell culture media can comprise a copper concentration of at least about 2.4 $\mu\text{g/L}$, in another embodiment at least about 3 $\mu\text{g/L}$, in yet another embodiment at least about 4 $\mu\text{g/L}$, in yet another embodiment at least about 8 $\mu\text{g/L}$, in yet another embodiment at least about 10 $\mu\text{g/L}$, in yet another embodiment at least about 15 $\mu\text{g/L}$, and in a further embodiment at least about 20 $\mu\text{g/L}$.

In other embodiments, the copper concentration in the cell culture media of the present invention can range from about 2.4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$, in another embodiment from about 2.4 $\mu\text{g/L}$ to about 15 $\mu\text{g/L}$, in yet another embodiment from about 2.4 $\mu\text{g/L}$ to about 10 $\mu\text{g/L}$, in yet another embodiment from about 2.4 $\mu\text{g/L}$ to about 8 $\mu\text{g/L}$, in yet another embodiment from about 2.4 $\mu\text{g/L}$ to about 6 $\mu\text{g/L}$, in yet another embodiment from about 2.4 $\mu\text{g/L}$ to about 4 $\mu\text{g/L}$, in yet another embodiment from about 4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$, in yet another embodiment from about 4 $\mu\text{g/L}$ to about 15 $\mu\text{g/L}$, in yet another embodiment from about 4 $\mu\text{g/L}$ to about 10 $\mu\text{g/L}$, in yet another embodiment from about 4 $\mu\text{g/L}$ to about 8 $\mu\text{g/L}$, and in a further embodiment from about 4 $\mu\text{g/L}$ to about 6 $\mu\text{g/L}$.

The present invention also provides kits for the expression or production of rVWF, the kits comprising a culture medium suitable for the expression of rVWF having high specific activity.

B. ADAMTS13 (A13) Cell Culture Media

In one aspect, the present invention provides culture media that are useful for the expression of ADAMTS proteins having high specific activities. Advantageously, it has been found that by supplementing a culture medium with copper, that the activities of recombinant ADAMTS (e.g., rADAMTS13) enzymes expressed in cells cultured in the supplemented medium are greatly enhanced, while the enzymes are expressed at levels as high, if not higher, than cells cultured in non-supplemented mediums.

In one aspect, the present invention provides cell culture media supplemented with copper for the expression of recombinant ADAMTS13 protein with high specific activity. In one embodiment, the media are supplemented to result in a total copper concentration of from about 2 to about 4 $\mu\text{g/L}$. In further embodiments, the media are supplemented to result in a total copper concentration of from about 1-3, 2-3, 3-4 $\mu\text{g/L}$. In one embodiment, the media contain a copper concentration of at least 1 $\mu\text{g/L}$. In another embodiment, the media contains at least 2 $\mu\text{g/L}$ copper. In another embodiment, the media contains at least 4 $\mu\text{g/L}$ copper. In other embodiments, the media contains between 2 $\mu\text{g/L}$ and 20 $\mu\text{g/L}$ copper. In another embodiment, the media contains between 1 $\mu\text{g/L}$ and 6 $\mu\text{g/L}$ copper. In another embodiment, the media contains between 2 $\mu\text{g/L}$ and 5 $\mu\text{g/L}$ copper. In another embodiment, the media contains between 3 $\mu\text{g/L}$ and 4 $\mu\text{g/L}$ copper. In yet other embodiments, the media contains at least 1 $\mu\text{g/L}$ copper, or at least 2 $\mu\text{g/L}$, 3 $\mu\text{g/L}$, 4 $\mu\text{g/L}$, 5 $\mu\text{g/L}$, 6 $\mu\text{g/L}$, 7 $\mu\text{g/L}$, 8 $\mu\text{g/L}$, 9 $\mu\text{g/L}$, 10 $\mu\text{g/L}$, 11 $\mu\text{g/L}$, 12 $\mu\text{g/L}$, 13 $\mu\text{g/L}$, 14 $\mu\text{g/L}$, 15 $\mu\text{g/L}$, 16 $\mu\text{g/L}$, 17 $\mu\text{g/L}$, 18 $\mu\text{g/L}$, 19 $\mu\text{g/L}$, 20 $\mu\text{g/L}$, or higher concentrations of copper.

In one embodiment, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM.

In yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less. In yet other embodiments, the cell culture solution has a copper and ammonium concentration according to any one of variations 1 to 440, as set forth in Table 1. In certain embodiments, the ammonium concentration of the cell culture is maintained for an extended period at a concentration as provided above. For example, in one embodiment, the ammonium concentration is maintained at a low concentration for at least 3 days, or at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In a specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 5 mM for at least 14 days. In another specific embodiment,

the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 14 days. In yet another embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rA13).

In one embodiment, a culture medium is provided for the expression of a recombinant ADAMTS protein (e.g., rADAMTS13) containing at least 1 µg/L copper and at least 2 µM zinc. In other embodiments, the media contains at least 2 µg/L copper or at least 4 µg/L copper. In another embodiment wherein the media is supplemented with copper, the culture medium also contains at least at or about 5 µM zinc. In one embodiment, the culture medium also contains at or about between 2 µM and 12 µM zinc. In another embodiment, the culture medium also contains at or about between 5 µM and 12 µM zinc. In yet other embodiments, the culture medium also may contain at least at or about 2 µM, or at least at or about 3 µM, 4 µM, 5 µM, 6 µM, 7 µM, 8 µM, 9 µM, 10 µM, 11 µM, 12 µM, 13 µM, 14 µM, 15 µM, 20 µM, 25 µM, 30 µM, or more zinc. In one embodiment, the culture medium contains a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2.

TABLE 2

		Zinc Concentration										
		AL 2 µM	AL 3 µM	AL 4 µM	AL 5 µM	AL 6 µM	AL 7 µM	AL 8 µM	AL 9 µM	AL 10 µM	2-12 µM	5-12 µM
Copper Concentration	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1 µg/L	441	481	521	561	601	641	681	721	761	801	841
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2 µg/L	442	482	522	562	602	642	682	722	762	802	842
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	3 µg/L	443	483	523	563	603	643	683	723	763	803	843
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	4 µg/L	444	484	524	564	604	644	684	724	764	804	844
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	5 µg/L	445	485	525	565	605	645	685	725	765	805	845
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	6 µg/L	446	486	526	566	606	646	686	726	766	806	846
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	7 µg/L	447	487	527	567	607	647	687	727	767	807	847
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	8 µg/L	448	488	528	568	608	648	688	728	768	808	848
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	9 µg/L	449	489	529	569	609	649	689	729	769	809	849
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	10 µg/L	450	490	530	570	610	650	690	730	770	810	850
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1 µg/L	451	491	531	571	611	651	691	731	771	811	851
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1.5 µg/L	452	492	532	572	612	652	692	732	772	812	852
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2 µg/L	453	493	533	573	613	653	693	733	773	813	853
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2.5 µg/L	454	494	534	574	614	654	694	734	774	814	854
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	3 µg/L	455	495	535	575	615	655	695	735	775	815	855
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	3.5 µg/L	456	496	536	576	616	656	696	736	776	816	856
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	4 µg/L	457	497	537	577	617	657	697	737	777	817	857
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	4.5 µg/L	458	498	538	578	618	658	698	738	778	818	858
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	5 µg/L	459	499	539	579	619	659	699	739	779	819	859
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	5.5 µg/L	460	500	540	580	620	660	700	740	780	820	860
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	6 µg/L	461	501	541	581	621	661	701	741	781	821	861

TABLE 2-continued

Exemplary embodiments of copper and zinc concentrations present in culture media useful for the expression of a recombinant ADAMTS13 protein.											
	Zinc Concentration										
	AL 2 μM	AL 3 μM	AL 4 μM	AL 5 μM	AL 6 μM	AL 7 μM	AL 8 μM	AL 9 μM	AL 10 μM	2-12 μM	5-12 μM
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
7 μg/L	462	502	542	582	622	662	702	742	782	822	862
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
8 μg/L	463	503	543	583	623	663	703	743	783	823	863
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
9 μg/L	464	504	544	584	624	664	704	744	784	824	864
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
10 μg/L	465	505	545	585	625	665	705	745	785	825	865
1-20 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
466	506	546	586	626	666	706	746	786	826	866	
2-20 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
467	507	547	587	627	667	707	747	787	827	867	
1-10 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
468	508	548	588	628	668	708	748	788	828	868	
2-10 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
469	509	549	589	629	669	709	749	789	829	869	
1-6 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
470	510	550	590	630	670	710	750	790	830	870	
2-6 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
471	511	551	591	631	671	711	751	791	831	871	
3-6 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
472	512	552	592	632	672	712	752	792	832	872	
4-6 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
473	513	553	593	633	673	713	753	793	833	873	
1-5 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
474	514	554	594	634	674	714	754	794	834	874	
2-5 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
475	515	555	595	635	675	715	755	795	835	875	
3-5 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
476	516	556	596	636	676	716	756	796	836	876	
4-5 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
477	517	557	597	637	677	717	757	797	837	877	
1-4 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
478	518	558	598	638	678	718	758	798	838	878	
2-4 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
479	519	559	599	639	679	719	759	799	839	879	
3-4 μg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
480	520	560	600	640	680	720	760	800	840	880	

*AL = at least

In one embodiment, the cell culture solution further comprises a low ammonium concentration. In one embodiment, the cell culture solution comprises an ammonium concentration of less than 10 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a specific embodiment, the ammonium is maintained at a concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 6 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a specific embodiment, the ammonium is maintained at a concentration of no more than 6 mM for at least 7 days. In another preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a specific embodiment, the ammonium is maintained at a concentration of no more than 5 mM for at least 7 days. In another preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a specific embodiment, the ammonium is maintained at a concentration of no more than 4 mM for at least 7 days. In

yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In yet another specific embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rA13).

In one embodiment, a culture medium is provided for the expression of a recombinant ADAMTS protein (e.g., rAD-AMTS13) containing at least 1 μg/L copper and at least at or about 0.5 mM calcium. In other embodiments, the media contains at least 2 μg/L copper or at least 4 μg/L copper. In another embodiment wherein the media is supplemented with copper, the culture medium also contains at least 1.5 mM calcium. In one embodiment, the culture medium contains at or about between 0.5 mM and 1.5 mM calcium. In yet other embodiments, the culture medium may contain at least at or about 0.5 mM, or at least at or about 0.6 mM, 0.7 mM, 0.8 mM, 0.9 mM, 1.0 mM, 1.1 mM, 1.2 mM, 1.3 mM, 1.4 mM, 1.5 mM, 1.6 mM, 1.7 mM, 1.8 mM, 1.9 mM, 2.0 mM, 2.25 mM, 2.5 mM, 2.75 mM, 3.0 mM, 3.5 mM, 4.0

mM, 4.5 mM, 5.0 mM, or more calcium. In one embodiment, the culture medium contains a copper and calcium

concentration according to any one of variations 881 to 1320, as set forth in Table 3.

TABLE 3

		Exemplary embodiments of copper and calcium concentrations present in culture media useful for the expression of a recombinant ADAMTS13 protein.										
		Calcium Concentration										
		AL 0.5 mM	AL 0.75 mM	AL 1.0 mM	AL 1.25 mM	AL 1.5 mM	AL 2.0 mM	AL 2.5 mM	AL 3.0 mM	AL 4 mM	AL 5 mM	AL 0.5-1.5 mM
Copper Concentration	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1 µg/L	881	921	961	1001	1041	1081	1121	1161	1201	1241	1281
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2 µg/L	882	922	962	1002	1042	1082	1122	1162	1202	1242	1282
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	3 µg/L	883	923	963	1003	1043	1083	1123	1163	1203	1243	1283
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	4 µg/L	884	924	964	1004	1044	1084	1124	1164	1204	1244	1284
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	5 µg/L	885	925	965	1005	1045	1085	1125	1165	1205	1245	1285
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	6 µg/L	886	926	966	1006	1046	1086	1126	1166	1206	1246	1286
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	7 µg/L	887	927	967	1007	1047	1087	1127	1167	1207	1247	1287
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	8 µg/L	888	928	968	1008	1048	1088	1128	1168	1208	1248	1288
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	9 µg/L	889	929	969	1009	1049	1089	1129	1169	1209	1249	1289
	AL	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	10 µg/L	890	930	970	1010	1050	1090	1130	1170	1210	1250	1290
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1 µg/L	891	931	971	1011	1051	1091	1131	1171	1211	1251	1291
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1.5 µg/L	892	932	972	1012	1052	1092	1132	1172	1212	1252	1292
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2 µg/L	893	933	973	1013	1053	1093	1133	1173	1213	1253	1293
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	2.5 µg/L	894	934	974	1014	1054	1094	1134	1174	1214	1254	1294
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	3 µg/L	895	935	975	1015	1055	1095	1135	1175	1215	1255	1295
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	3.5 µg/L	896	936	976	1016	1056	1096	1136	1176	1216	1256	1296
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	4 µg/L	897	937	977	1017	1057	1097	1137	1177	1217	1257	1297
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	4.5 µg/L	898	938	978	1018	1058	1098	1138	1178	1218	1258	1298
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	5 µg/L	899	939	979	1019	1059	1099	1139	1179	1219	1259	1299
	About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	5.5 µg/L	900	940	980	1020	1060	1100	1140	1180	1220	1260	1300
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
6 µg/L	901	941	981	1021	1061	1101	1141	1181	1221	1261	1301	
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
7 µg/L	902	942	982	1022	1062	1102	1142	1182	1222	1262	1302	
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
8 µg/L	903	943	983	1023	1063	1103	1143	1183	1223	1263	1303	
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
9 µg/L	904	944	984	1024	1064	1104	1144	1184	1224	1264	1304	
About	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
10 µg/L	905	945	985	1025	1065	1105	1145	1185	1225	1265	1305	
1-20 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	906	946	986	1026	1066	1106	1146	1186	1226	1266	1306	
2-20 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	907	947	987	1027	1067	1107	1147	1187	1227	1267	1307	
1-10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	908	948	988	1028	1068	1108	1148	1188	1228	1268	1308	
2-10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	909	949	989	1029	1069	1109	1149	1189	1229	1269	1309	
1-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	910	950	990	1030	1070	1110	1150	1190	1230	1270	1310	
2-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	911	951	991	1031	1071	1111	1151	1191	1231	1271	1311	
3-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	912	952	992	1032	1072	1112	1152	1192	1232	1272	1312	
4-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	913	953	993	1033	1073	1113	1153	1193	1233	1273	1313	
1-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	
	914	954	994	1034	1074	1114	1154	1194	1234	1274	1314	

TABLE 3-continued

Exemplary embodiments of copper and calcium concentrations present in culture media useful for the expression of a recombinant ADAMTS13 protein.											
Calcium Concentration											
	AL 0.5 mM	AL 0.75 mM	AL 1.0 mM	AL 1.25 mM	AL 1.5 mM	AL 2.0 mM	AL 2.5 mM	AL 3.0 mM	AL 4 mM	AL 5 mM	AL 0.5-1.5 mM
2-5 µg/L	Var. 915	Var. 955	Var. 995	Var. 1035	Var. 1075	Var. 1115	Var. 1155	Var. 1195	Var. 1235	Var. 1275	Var. 1315
3-5 µg/L	Var. 916	Var. 956	Var. 996	Var. 1036	Var. 1076	Var. 1116	Var. 1156	Var. 1196	Var. 1236	Var. 1276	Var. 1316
4-5 µg/L	Var. 917	Var. 957	Var. 997	Var. 1037	Var. 1077	Var. 1117	Var. 1157	Var. 1197	Var. 1237	Var. 1277	Var. 1317
1-4 µg/L	Var. 918	Var. 958	Var. 998	Var. 1038	Var. 1078	Var. 1118	Var. 1158	Var. 1198	Var. 1238	Var. 1278	Var. 1318
2-4 µg/L	Var. 919	Var. 959	Var. 999	Var. 1039	Var. 1079	Var. 1119	Var. 1159	Var. 1199	Var. 1239	Var. 1279	Var. 1319
3-4 µg/L	Var. 920	Var. 960	Var. 1000	Var. 1040	Var. 1080	Var. 1120	Var. 1160	Var. 1200	Var. 1240	Var. 1280	Var. 1320

*AL = at least

In one embodiment, the cell culture solution further comprises a low ammonium concentration. In one embodiment, the cell culture solution comprises an ammonium concentration of less than 10 mM and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In a specific embodiment, the ammonium is maintained at a concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture medium comprises an ammonium concentration of no more than 6 mM and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In a specific embodiment, the ammonium is maintained at a concentration of no more than 6 mM for at least 7 days. In another preferred embodiment, the cell culture medium comprises an ammonium concentration of no more than 5 mM and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In a specific embodiment, the ammonium is maintained at a concentration of no more than 5 mM for at least 7 days. In another preferred embodiment, the cell culture medium comprises an ammonium concentration of no more than 4 mM and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In a specific embodiment, the ammonium is maintained at a concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture medium comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In yet another specific embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rA13).

In one embodiment, the cell culture medium is supplemented with copper, zinc and calcium. In a specific embodiment, the culture medium has a calcium concentration of at least 0.5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a calcium concentration of at least 1.5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a calcium concentration between 0.5 mM and 1.5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In yet other embodiments, the culture medium has a calcium concentration of at least 0.6 mM, 0.7 mM, 0.8 mM, 0.9 mM, 1.0 mM, 1.1 mM, 1.2 mM, 1.3 mM, 1.4 mM, 1.5 mM, 1.6 mM, 1.7 mM, 1.8 mM, 1.9 mM, 2.0 mM, 2.25 mM, 2.5 mM, 2.75 mM, 3.0 mM, 3.5 mM, 4.0 mM, 4.5 mM, 5.0 mM, or more, and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2.

In one embodiment, a culture medium is provided for the expression of a recombinant ADAMTS protein (e.g., rADAMTS13) containing at least 1 µg/L copper and at least 2 mg/L nicotinamide (vitamin B3). In other embodiments, the media contains at least 2 µg/L copper or at least 4 µg/L copper. In another embodiment wherein the media is supplemented with copper, the culture medium also contains at least 7 mg/L nicotinamide (vitamin B3). In one embodiment, the culture medium contains at or about between 2 mg/L and 10 mg/L nicotinamide (vitamin B3). In yet other embodiments, the culture medium may contain at least at or about 2 mg/L, 3 mg/L, 4 mg/L, 5 mg/L, 6 mg/L, 7 mg/L, 8 mg/L, 9 mg/L, 10 mg/L, 15 mg/L, 20 mg/L, or higher concentrations of nicotinamide (vitamin B3). In one embodiment, the culture medium contains a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4.

TABLE 4

Exemplary embodiments of copper and nicotinamide concentrations present in culture media useful for the expression of a recombinant ADAMTS13 protein.												
Calcium Concentration												
	AL 2 mg/mL	AL 3 mg/mL	AL 4 mg/mL	AL 5 mg/mL	AL 6 mg/mL	AL 7 mg/mL	AL 8 mg/mL	AL 9 mg/mL	AL 10 mg/mL	AL 15 mg/mL	AL 2-10 mg/mL	
Copper Concentration	AL 1 µg/L	Var. 1321	Var. 1361	Var. 1401	Var. 1441	Var. 1481	Var. 1521	Var. 1561	Var. 1601	Var. 1641	Var. 1681	Var. 1721

TABLE 4-continued

	Calcium Concentration										
	AL 2 mg/mL	AL 3 mg/mL	AL 4 mg/mL	AL 5 mg/mL	AL 6 mg/mL	AL 7 mg/mL	AL 8 mg/mL	AL 9 mg/mL	AL 10 mg/mL	AL 15 mg/mL	2-10 mg/mL
AL 2 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1322	1362	1402	1442	1482	1522	1562	1602	1642	1682	1722
AL 3 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1323	1363	1403	1443	1483	1523	1563	1603	1643	1683	1723
AL 4 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1324	1364	1404	1444	1484	1524	1564	1604	1644	1684	1724
AL 5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1325	1365	1405	1445	1485	1525	1565	1605	1645	1685	1725
AL 6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1326	1366	1406	1446	1486	1526	1566	1606	1646	1686	1726
AL 7 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1327	1367	1407	1447	1487	1527	1567	1607	1647	1687	1727
AL 8 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1328	1368	1408	1448	1488	1528	1568	1608	1648	1688	1728
AL 9 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1329	1369	1409	1449	1489	1529	1569	1609	1649	1689	1729
AL 10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1330	1370	1410	1450	1490	1530	1570	1610	1650	1690	1730
About 1 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1331	1371	1411	1451	1491	1531	1571	1611	1651	1691	1731
About 1.5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1332	1372	1412	1452	1492	1532	1572	1612	1652	1692	1732
About 2 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1333	1373	1413	1453	1493	1533	1573	1613	1653	1693	1733
About 2.5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1334	1374	1414	1454	1494	1534	1574	1614	1654	1694	1734
About 3 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1335	1375	1415	1455	1495	1535	1575	1615	1655	1695	1735
About 3.5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1336	1376	1416	1456	1496	1536	1576	1616	1656	1696	1736
About 4 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1337	1377	1417	1457	1497	1537	1577	1617	1657	1697	1737
About 4.5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1338	1378	1418	1458	1498	1538	1578	1618	1658	1698	1738
About 5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1339	1379	1419	1459	1499	1539	1579	1619	1659	1699	1739
About 5.5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1340	1380	1420	1460	1500	1540	1580	1620	1660	1700	1740
About 6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1341	1381	1421	1461	1501	1541	1581	1621	1661	1701	1741
About 7 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1342	1382	1422	1462	1502	1542	1582	1622	1662	1702	1742
About 8 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1343	1383	1423	1463	1503	1543	1583	1623	1663	1703	1743
About 9 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1344	1384	1424	1464	1504	1544	1584	1624	1664	1704	1744
About 10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1345	1385	1425	1465	1505	1545	1585	1625	1665	1705	1745
1-20 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1346	1386	1426	1466	1506	1546	1586	1626	1666	1706	1746
2-20 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1347	1387	1427	1467	1507	1547	1587	1627	1667	1707	1747
1-10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1348	1388	1428	1468	1508	1548	1588	1628	1668	1708	1748
2-10 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1349	1389	1429	1469	1509	1549	1589	1629	1669	1709	1749
1-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1350	1390	1430	1470	1510	1550	1590	1630	1670	1710	1750
2-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1351	1391	1431	1471	1511	1551	1591	1631	1671	1711	1751
3-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1352	1392	1432	1472	1512	1552	1592	1632	1672	1712	1752
4-6 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1353	1393	1433	1473	1513	1553	1593	1633	1673	1713	1753
1-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1354	1394	1434	1474	1514	1554	1594	1634	1674	1714	1754
2-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1355	1395	1435	1475	1515	1555	1595	1635	1675	1715	1755
3-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1356	1396	1436	1476	1516	1556	1596	1636	1676	1716	1756
4-5 µg/L	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.	Var.
	1357	1397	1437	1477	1517	1557	1597	1637	1677	1717	1757

TABLE 4-continued

Exemplary embodiments of copper and nicotinamide concentrations present in culture media useful for the expression of a recombinant ADAMTS13 protein.											
Calcium Concentration											
	AL 2 mg/mL	AL 3 mg/mL	AL 4 mg/mL	AL 5 mg/mL	AL 6 mg/mL	AL 7 mg/mL	AL 8 mg/mL	AL 9 mg/mL	AL 10 mg/mL	AL 15 mg/mL	2-10 mg/mL
1-4 µg/L	Var. 1358	Var. 1398	Var. 1438	Var. 1478	Var. 1518	Var. 1558	Var. 1598	Var. 1638	Var. 1678	Var. 1718	Var. 1758
2-4 µg/L	Var. 1359	Var. 1399	Var. 1439	Var. 1479	Var. 1519	Var. 1559	Var. 1599	Var. 1639	Var. 1679	Var. 1719	Var. 1759
3-4 µg/L	Var. 1360	Var. 1400	Var. 1440	Var. 1480	Var. 1520	Var. 1560	Var. 1600	Var. 1640	Var. 1680	Var. 1720	Var. 1760

*AL = at least

In one embodiment, the cell culture solution further comprises a low ammonium concentration. In one embodiment, the culture medium comprises an ammonium concentration of less than 10 mM and a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4. In a specific embodiment, the ammonium is maintained at a concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 6 mM and a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4. In a specific embodiment, the ammonium is maintained at a concentration of no more than 6 mM for at least 7 days. In another preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM and a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4. In a specific embodiment, the ammonium is maintained at a concentration of no more than 5 mM for at least 7 days. In another preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM and a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4. In a specific embodiment, the ammonium is maintained at a concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less and a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4. In yet another specific embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rA13).

In one embodiment, the cell culture medium is supplemented with copper, zinc and nicotinamide. In a specific embodiment, the culture medium has a nicotinamide concentration of at least 2 mg/mL and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a nicotinamide concentration of at least 7 mg/mL mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a nicotinamide concentration between 2 mg/mL and 10 mg/mL and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In yet other embodiments, the culture medium has a nicotinamide concentration of at least 2 mg/mL, 3 mg/mL, 4 mg/mL,

5 mg/mL, 6 mg/mL, 7 mg/mL, 8 mg/mL, 9 mg/mL, 10 mg/mL, 11 mg/mL, 12 mg/mL, 13 mg/mL, 14 mg/mL, 15 mg/mL, or more, and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2.

In one embodiment, the cell culture medium is supplemented with copper, calcium and nicotinamide. In a specific embodiment, the culture medium has a nicotinamide concentration of at least 2 mg/mL and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In another specific embodiment, the culture medium has a nicotinamide concentration of at least 7 mg/mL mM and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In another specific embodiment, the culture medium has a nicotinamide concentration between 2 mg/mL and 10 mg/mL and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In yet other embodiments, the culture medium has a nicotinamide concentration of at least 2 mg/mL, 3 mg/mL, 4 mg/mL, 5 mg/mL, 6 mg/mL, 7 mg/mL, 8 mg/mL, 9 mg/mL, 10 mg/mL, 11 mg/mL, 12 mg/mL, 13 mg/mL, 14 mg/mL, 15 mg/mL, or more, and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3.

IV. Methods for the Production of Blood Factors Having High Specific Activity

A. Cell Cultivation Methods

The present invention provides methods for large-scale production of recombinant proteins (such as rVWF and rA13). In certain embodiments, such large-scale production methods utilize stirred/agitated tank reactors for manufacture of these therapeutic recombinant proteins.

In certain embodiments, the methods of the present invention can comprise the use of a cell culture system operated under a batch or continuous mode of operation. For example, when batch cell cultures are utilized, they may be operated under single batch, fed-batch, or repeated-batch mode. Likewise, continuous cell cultures may be operated under, for example, perfusion, turbidostat or chemostat mode. Batch and continuous cell cultivation may be performed under either suspension or adherence conditions. When operated under suspension conditions, the cells will be freely suspended and mixed within the culture medium. Alternatively, under adherence conditions, the cells will be bound to a solid phase, for example, a microcarrier, a porous microcarrier, disk carrier, ceramic cartridge, hollow fiber, flat sheet, gel matrix, and the like.

A batch culture is typically a large scale cell culture in which a cell inoculum is cultured to a maximum density in a tank or fermenter, and harvested and processed as a single batch. A fed-batch culture is typically a batch culture which is supplied with either fresh nutrients (e.g., growth-limiting substrates) or additives (e.g., precursors to products). The feed solution is usually highly concentrated to avoid dilution of the bioreactor. In a repeated-batch culture, the cells are placed in a culture medium and grown to a desired cell density. To avoid the onset of a decline phase and cell death, the culture is then diluted with complete growth medium before the cells reach their maximum concentration. The amount and frequency of dilution varies widely and depends on the growth characteristics of the cell line and convenience of the culture process. The process can be repeated as many times as required and, unless cells and medium are discarded at subculture, the volume of culture will increase stepwise as each dilution is made. The increasing volume may be handled by having a reactor of sufficient size to allow dilutions within the vessel or by dividing the diluted culture into several vessels. The rationale of this type of culture is to maintain the cells in an exponentially growing state. Serial subculture is characterized in that the volume of culture is always increasing stepwise, there can be multiple harvests, the cells continue to grow and the process can continue for as long as desired. In certain embodiments, a recombinant ADAMTS protein (e.g., rADAMTS13) may be recovered after harvesting the supernatant of a batch culture. In other embodiments, a recombinant VWF may be recovered after harvesting the supernatant of a batch culture.

A continuous culture can be a suspension culture that is continuously supplied with nutrients by the inflow of fresh medium, wherein the culture volume is usually kept constant by the concomitant removal of spent medium. In chemostat and turbidostat methods, the extracted medium contains cells. Thus, the cells remaining in the cell culture vessel must grow to maintain a steady state. In the chemostat method, the growth rate is typically controlled by controlling the dilution rate, i.e., the rate at which fresh medium is added. The growth rate of the cells in the culture may be controlled, for example, at a sub-maximal growth rate, by alteration of the dilution rate. In contrast, in the turbidostat method, the dilution rate is set to permit the maximum growth rate that the cells can achieve at the given operating conditions, such as pH and temperature. In certain embodiments, the rVWF or rA13 is recovered after harvesting the supernatant of a continuous culture. An exemplary method for continuous cell cultivation is described in WO/2011/012725 (Grillberger et al.), the content of which is hereby incorporated by reference in its entirety for all purposes.

In a perfusion culture, the extracted medium is depleted of cells, which are retained in the culture vessel, for example, by filtration or by centrifugal methods that lead to the reintroduction of the cells into the culture. However, typically membranes used for filtration do not retain 100% of cells, and so a proportion is removed when the medium is extracted. It may not be crucial to operate perfusion cultures at very high growth rates, as the majority of the cells are retained in the culture vessel. In certain embodiments, the rVWF or rA13 is recovered after harvesting the supernatant of a perfusion culture.

Stirred-tank reactor system can be used for batch and continuous cell cultures operated under suspension or adherent modes. Generally, the stirred-tank reactor system can be operated as any conventional stirred-tank reactor with any type of agitator such as a Rushton, hydrofoil, pitched blade, or marine.

In certain embodiments, the cell-culture methods of the invention may comprise the use of a microcarrier. In some embodiments, the cell-cultures of the embodiments can be performed in large bioreactors under conditions suitable for providing high volume-specific culture surface areas to achieve high cell densities and protein expression. One means for providing such growth conditions is to use microcarriers for cell-culture in stirred tank bioreactors. The concept of cell-growth on microcarriers was first described by van Wezel (van Wezel, A. L., *Nature* 216:64-5 (1967)) and allows for cell attachment on the surface of small solid particles suspended in the growth medium. These methods provide for high surface-to-volume ratios and thus allow for efficient nutrient utilization. Furthermore, for expression of secreted proteins in eukaryotic cell lines, the increased surface-to-volume ratio allows for higher levels of secretion and thus higher protein yields in the supernatant of the culture. Finally, these methods allow for the easy scale-up of eukaryotic expression cultures.

The cells expressing vWF and/or rA13 can be bound to a spherical or a porous microcarrier during cell culture growth. The microcarrier can be a microcarrier selected from the group of microcarriers based on dextran, collagen, plastic, gelatine and cellulose and others as described in Butler (1988. In: Spier & Griffiths, *Animal Cell Biotechnology* 3:283-303). It is also possible to grow the cells to a biomass on spherical microcarriers and subculture the cells when they have reached final fermenter biomass and prior to production of the expressed protein on a porous microcarrier or vice versa. Suitable spherical microcarriers can include smooth surface microcarriers, such as Cytodex™ 1, Cytodex™ 2, and Cytodex™ 3 (GE Healthcare) and macroporous microcarriers such as Cytopore™ 1, Cytopore™ 2, Cytoline™ 1, and Cytoline™ 2 (GE Healthcare).

As described above, the present invention includes cell culture media having an increased copper concentration. It is understood that all of the embodiments and concentrations described in the "Cell Culture Media" section above can be applied to the methods of the present invention described herein.

In certain embodiments, the culture can be maintained for at least about 7 days, or at least about 14 days, 21 days, 28 days, or at least about 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, or at least about 2 months, or 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 months or longer. The cell density at which a cell-culture is maintained at for production of a recombinant vWF or recombinant rA13 protein will depend upon the culture-conditions and medium used for protein expression. One of skill in the art will readily be able to determine the optimal cell density for a cell-culture producing rVWF or rA13. In one embodiment, the culture is maintained at a cell density of between about 0.5×10^6 and 4×10^7 cells/mL for an extended period of time. In other embodiments, the cell density is maintained at a concentration of between about 1.0×10^6 and about 1.0×10^7 cells/mL for an extended period of time. In other embodiments, the cell density is maintained at a concentration of between about 1.0×10^6 and about 4.0×10^6 cells/mL for an extended period of time. In other embodiments, the cell density is maintained at a concentration of between about 1.0×10^6 and about 4.0×10^6 cells/mL for an extended period of time. In yet other embodiments, the cell density may be maintained at a concentration between about 2.0×10^6 and about 4.0×10^6 cells/mL, or between about 1.0×10^6 and about 2.5×10^6 cells/mL, or between about 1.5×10^6 and about 3.5×10^6 cells/mL, or any other similar range, for an extended period of time.

of a continuous cell culture provided herein is maintained at a concentration of no more than 2.0×10^6 cells/mL for at least 8 weeks. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 2.0×10^6 cells/mL for at least 9 weeks.

In one embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 7 days. In a specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 14 days. In a more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 21 days. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 28 days. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 5 weeks. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 6 weeks. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 7 weeks. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 8 weeks. In a yet more specific embodiment, the cell density of a continuous cell culture provided herein is maintained at a concentration of no more than 1.5×10^6 cells/mL for at least 9 weeks.

The following provides specific details on methods for producing rVWF and rA13. As will be appreciated, although the conditions are presented specifically for rVWF or rA13, the conditions for rVWF can be used for producing rA13 and vice versa.

B. Methods of Producing High Molecular Weight Recombinant vWF

In another aspect, the present invention further relates to methods for producing vWF under cell culture conditions comprising a cell culture medium having an increased copper concentration. In certain embodiments, the culture also comprises a low ammonium concentration. As used herein, the term "cell culture" and "cell culture solution" are used interchangeably.

In one embodiment, the present invention provides a method of producing a high molecular weight, recombinant vWF, comprising: a) providing a culture of cells; b) introducing a nucleic acid sequence coding for vWF; c) selecting the cells carrying the nucleic acid sequence; and, d) expressing vWF in the cells under cell culture conditions comprising a cell culture medium comprising a copper concentration of at least about $2.4 \mu\text{g/L}$ and a cell culture supernatant comprising an ammonium concentration less than about 10 mM, wherein the vWF is highly multimeric vWF comprising about 14 to about 22 dimers and a specific Ristocetin activity of at least about $30 \text{ mU}/\mu\text{g}$. In further embodiments, the multimeric rVWF produced using methods of the present invention comprise about 10-30, 12-28, 14-26, 16-24, 18-22, 20-21 dimers. In still further embodiments, the rVWF produced in accordance with the present invention has a specific activity of at least about 20, 22.5, 25, 27.5, 30, 32.5, 35, 37.5, 40, 42.5, 45, 47.5, 50, 52.5, 55, 57.5, 60, 62.5, 65, 67.5, 70, 72.5, 75, 77.5, 80, or more $\text{mU}/\mu\text{g}$. In a specific embodi-

ment, the cell density of the continuous cell culture for production of rVWF is maintained at a concentration of no more than 2.5×10^6 cells/mL for an extended period. In other specific embodiments, the cell density is maintained at no more than 2.0×10^6 cells/mL, 1.5×10^6 cells/mL, 1.0×10^6 cells/mL, 0.5×10^6 cells/mL, or less. In one embodiment, the cell density is maintained at between 1.5×10^6 cells/mL and 2.5×10^6 cells/mL.

In one embodiment, the present invention provides a method for producing a recombinant Von Willebrand Factor (rVWF) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper to provide a final copper concentration of at least $2.0 \mu\text{g/L}$; (c) providing one or more cells comprising a nucleic acid encoding a rVWF protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rVWF is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least $30 \text{ mU}/\mu\text{g}$ rVWF. In one embodiment, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In another specific embodiment, the ammonium concentration of the cell culture solution is maintained at an ammonium concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM. In another specific embodiment, the ammonium concentration of the cell culture solution is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less. In yet another embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rVWF). In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least $50 \text{ mU}/\mu\text{g}$ rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least $70 \text{ mU}/\mu\text{g}$ rVWF.

In one embodiment of the method described above, the basal cell culture media is supplemented with copper to provide a final copper concentration of at least $2.4 \mu\text{g/L}$. In one embodiment, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In another specific embodiment, the ammonium concentration of the cell culture solution is maintained at an ammonium concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM. In another specific embodiment, the ammonium concentration of the cell culture solution is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM. In another specific embodiment, the ammonium concentration of the cell culture solution is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture solution comprises an ammo-

10 $\mu\text{g/L}$. In one embodiment, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM. In another specific embodiment, the ammonium concentration of the cell culture solution is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less. In yet another embodiment, the ammonium concentration of the cell culture solution is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rVWF). In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the basal cell culture media is supplemented with copper to provide a final copper concentration of between 4 $\mu\text{g/L}$ and 7.5 $\mu\text{g/L}$. In one embodiment, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less. In yet another embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rVWF). In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment, the present invention provides a method for producing a recombinant Von Willebrand Factor (rVWF) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper; (c) providing one or more cells comprising a nucleic acid encoding a rVWF protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rVWF is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture

supernatant, wherein the NH_4^+ concentration of the supernatant is maintained at a low level for at least 7 days, and further wherein the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 30 mU/ μg rVWF. In a specific embodiment, the copper concentration and NH_4^+ concentration of the cell culture solution is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 14 days. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 21 days. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 28 days. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 5 weeks. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 6 weeks. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 7 weeks. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 8 weeks. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, the copper concentration and NH_4^+ concentration of the cell culture is maintained at a concentration according to any one of variations 1 to 440, as set forth in Table 1 for at least 9 weeks. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment, the present invention provides a method for producing a recombinant Von Willebrand Factor (rVWF) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper to provide a final copper concentration of at least 2.0 $\mu\text{g}/\text{L}$; (c) providing one or more cells comprising a nucleic acid encoding a rVWF protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rVWF is expressed and excreted from the cells into a culture supernatant; (e) monitoring the ammonium concentration of the culture supernatant; and (f) recovering at least a portion of the culture supernatant, wherein culture supernatant comprising an ammonium concentration of more than 10 mM is not used for producing the rVWF composition, and further wherein the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 30 mU/ μg rVWF. In certain embodiments, the final copper concentration of the supplemented basal culture media is at least 2.4 $\mu\text{g}/\text{L}$, 3 $\mu\text{g}/\text{L}$, 4 $\mu\text{g}/\text{L}$, 5 $\mu\text{g}/\text{L}$, 6 $\mu\text{g}/\text{L}$, 7 $\mu\text{g}/\text{L}$, 8 $\mu\text{g}/\text{L}$, 9 $\mu\text{g}/\text{L}$, 10 $\mu\text{g}/\text{L}$, 15 $\mu\text{g}/\text{L}$, 20 $\mu\text{g}/\text{L}$, or higher. In other embodiments, the final copper concentration of the supplemented basal culture media is between 2-20 $\mu\text{g}/\text{L}$, 2-10 $\mu\text{g}/\text{L}$, 3-8 $\mu\text{g}/\text{L}$, or 4-6 $\mu\text{g}/\text{L}$. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, culture supernatant comprising an ammonium concentration of more than 6 mM is not used for producing the rVWF composition. In certain embodiments, the final copper concentration of the supplemented basal culture media is at least 2.4 $\mu\text{g}/\text{L}$, 3 $\mu\text{g}/\text{L}$, 4 $\mu\text{g}/\text{L}$, 5 $\mu\text{g}/\text{L}$, 6 $\mu\text{g}/\text{L}$, 7 $\mu\text{g}/\text{L}$, 8 $\mu\text{g}/\text{L}$, 9 $\mu\text{g}/\text{L}$, 10 $\mu\text{g}/\text{L}$, 15 $\mu\text{g}/\text{L}$, 20 $\mu\text{g}/\text{L}$, or higher. In other embodiments, the final copper concentration of the supplemented basal culture media is between 2-20 $\mu\text{g}/\text{L}$, 2-10 $\mu\text{g}/\text{L}$, 3-8 $\mu\text{g}/\text{L}$, or 4-6 $\mu\text{g}/\text{L}$. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, culture supernatant comprising an ammonium concentration of more than 5 mM is not used for producing the rVWF composition. In certain embodiments, the final copper concentration of the supplemented basal culture media is at least 2.4 $\mu\text{g}/\text{L}$, 3 $\mu\text{g}/\text{L}$, 4 $\mu\text{g}/\text{L}$, 5 $\mu\text{g}/\text{L}$, 6 $\mu\text{g}/\text{L}$, 7 $\mu\text{g}/\text{L}$, 8 $\mu\text{g}/\text{L}$, 9 $\mu\text{g}/\text{L}$, 10 $\mu\text{g}/\text{L}$, 15 $\mu\text{g}/\text{L}$, 20 $\mu\text{g}/\text{L}$, or higher. In other

embodiments, the final copper concentration of the supplemented basal culture media is between 2-20 $\mu\text{g}/\text{L}$, 2-10 $\mu\text{g}/\text{L}$, 3-8 $\mu\text{g}/\text{L}$, or 4-6 $\mu\text{g}/\text{L}$. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

In one embodiment of the method described above, culture supernatant comprising an ammonium concentration of more than 4 mM is not used for producing the rVWF composition. In certain embodiments, the final copper concentration of the supplemented basal culture media is at least 2.4 $\mu\text{g}/\text{L}$, 3 $\mu\text{g}/\text{L}$, 4 $\mu\text{g}/\text{L}$, 5 $\mu\text{g}/\text{L}$, 6 $\mu\text{g}/\text{L}$, 7 $\mu\text{g}/\text{L}$, 8 $\mu\text{g}/\text{L}$, 9 $\mu\text{g}/\text{L}$, 10 $\mu\text{g}/\text{L}$, 15 $\mu\text{g}/\text{L}$, 20 $\mu\text{g}/\text{L}$, or higher. In other embodiments, the final copper concentration of the supplemented basal culture media is between 2-20 $\mu\text{g}/\text{L}$, 2-10 $\mu\text{g}/\text{L}$, 3-8 $\mu\text{g}/\text{L}$, or 4-6 $\mu\text{g}/\text{L}$. In a preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 50 mU/ μg rVWF. In a more preferred embodiment, the recovered supernatant has a rVWF specific ristocetin cofactor activity of at least 70 mU/ μg rVWF.

Recombinant vWF can be produced by expression in a suitable eukaryotic host system. Examples of eukaryotic cells include, without limitation, mammalian cells, such as CHO, COS, HEK 293, BHK, SK-Hep, and HepG2; insect cells, e.g., SF9 cells, SF21 cells, S2 cells, and High Five cells; and yeast cells, e.g., *Saccharomyces* or *Schizosaccharomyces* cells. In one embodiment, the vWF can be expressed in yeast cells, insect cells, avian cells, mammalian cells, and the like. For example, in a human cell line, a hamster cell line, or a murine cell line. In one particular embodiment, the cell line is a CHO, BHK, or HEK cell line. Typically, mammalian cells, e.g., CHO cells from a continuous cell line, can be used to express the vWF of the present invention.

In certain embodiments, the nucleic acid sequence comprising a sequence coding for vWF can be a vector. The vector can be delivered by a virus or can be a plasmid. The nucleic acid sequence coding for the protein can be a specific gene or a biologically functional part thereof. In one embodiment, the protein is at least a biologically active part of vWF.

A wide variety of vectors can be used for the expression of the vWF and can be selected from eukaryotic expression vectors. Examples of vectors for eukaryotic expression include: (i) for expression in yeast, vectors such as pAO, pPIC, pYES, pMET, using promoters such as AOX1, GAP, GAL1, AUG1, etc; (ii) for expression in insect cells, vectors such as pMT, pAc5, pIB, pMIB, pBAC, etc., using promoters such as PH, p10, MT, Ac5, OpIE2, gp64, polh, etc., and (iii) for expression in mammalian cells, vectors such as pSVL, pCMV, pRc/RSV, pcDNA3, pBPV, etc., and vectors derived from viral systems such as vaccinia virus, adeno-associated viruses, herpes viruses, retroviruses, etc., using promoters such as CMV, SV40, EF-1, UbC, RSV, ADV, BPV, and β -actin. An exemplary vector for expressing rVWF is described by Kaufman et al. (Mol Cell Biol. 1989 March; 9(3):1233-42), the content of which is hereby incorporated by reference in its entirety for all purposes.

In some embodiments of the present invention, the nucleic acid sequence further comprises other sequences suitable for a controlled expression of a protein such as promoter sequences, enhancers, TATA boxes, transcription initiation sites, polylinkers, restriction sites, poly-A-sequences, protein processing sequences, selection markers, and the like which are generally known to a person of ordinary skill in the art.

In addition to cell culture media comprising an increased copper concentration, the cell culture conditions of the present invention can include an ammonium concentration of less than about 25 mM throughout an entire upstream process in culture systems. In one embodiment, the cell culture conditions include an ammonium concentration of less than about 25 mM, in another embodiment less than about 20 mM, in yet another embodiment less than about 15 mM, in yet another embodiment less than about 10 mM, and in a further embodiment less than about 5 mM.

In some embodiments, the ammonium concentrations of the present invention are kept constant throughout the entire upstream process of the cell culture system. The cells used according to the present invention can be cultivated, e.g., by methods that are modified as compared to conventional batch-cultivation and feed-batch-cultivation, each of which are generally known in the field. However, such conventional techniques can produce high concentrations of ammonium at the end of the culture. The methods of the present invention overcome this problem by employing production systems that can provide a continuous supply of culture media through techniques such as, e.g., perfusion or chemostat cultures. Following culture of the host cells, the vWF can be recovered from the spent medium using standard methodologies, such as ultrafiltration or centrifugation. If desired, the vWF can be purified by, e.g., ion exchange and/or size exclusion chromatography and the like.

A continuous culture (e.g., a perfusion or chemostat culture) can be a suspension culture that is continuously supplied with nutrients by the inflow of fresh medium, wherein the culture volume is usually constant. Similarly, continuous fermentation can refer to a process in which cells or micro-organisms are maintained in culture in the exponential growth phase by the continuous addition of fresh medium that is exactly balanced by the removal of cell suspension from the bioreactor. Furthermore, a stirred-tank reactor system can be used for suspension, perfusion, chemostatic, and/or microcarrier cultures. Generally, the stirred-tank reactor system can be operated as any conventional stirred-tank reactor with any type of agitator such as a Rushton, hydrofoil, pitched blade, or marine.

C. Methods of Producing Recombinant ADAMTS13 (A13)

In another aspect, the present invention further relates to methods for producing rA13 under cell culture conditions comprising a cell culture medium having an increased copper concentration. In certain embodiments, the culture also comprises a low ammonium concentration.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper to provide a final copper concentration of at least 1.0 $\mu\text{g/L}$; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day (i.e., 1500 Units FRETS-VWF73 activity per day per liter cell culture; P FRETS) is present in the recovered culture supernatant. In certain embodiments, the final copper concentration of the supplemented basal culture media is at least 2 $\mu\text{g/L}$, 3 $\mu\text{g/L}$, 4 $\mu\text{g/L}$, 5 $\mu\text{g/L}$, 6 $\mu\text{g/L}$, or higher. In other embodiments, the final copper concentration of the supplemented basal culture

media is between 1-6 $\mu\text{g/L}$, 2-5 $\mu\text{g/L}$, 2-4 $\mu\text{g/L}$, or 3-4 $\mu\text{g/L}$. In a preferred embodiment, at least 2000 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000 Units FRETS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In certain embodiments, the methods provide sustainably improved volumetric FRETS-VWF73 productivity (P FRETS). For example, in certain embodiments, at least 1500 Units FRETS-VWF73 activity per liter supplemented basal cell culture media is recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a preferred embodiment, at least 2000 Units FRETS-VWF73 activity per liter supplemented basal cell culture media is recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a more preferred embodiment, at least 2500 Units FRETS-VWF73 activity per liter supplemented basal cell culture media is recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a most preferred embodiment, at least 3000 Units FRETS-VWF73 activity per liter supplemented basal cell culture media is recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a specific embodiment, the cell density of the continuous cell culture for production of rA13 is maintained at a concentration of no more than 4.0×10^6 cells/mL for an extended period. In other specific embodiments, the cell density is maintained at no more than 3.5×10^6 cells/mL, 3.0×10^6 cells/mL, 2.5×10^6 cells/mL, 2.0×10^6 cells/mL, 1.5×10^6 cells/mL, 1.0×10^6 cells/mL, or less. In one embodiment, the cell density is maintained at between 3.0×10^6 cells/mL and 4.0×10^6 cells/mL.

In one embodiment of the methods described above, the recovered supernatant has at least 4 Units FRETS-VWF73 activity per mL supernatant (FRETS). In a preferred embodiment, the recovered supernatant has at least 6 Units FRETS-VWF73 activity per mL supernatant. In a more preferred embodiment, the recovered supernatant has at least 8 Units FRETS-VWF73 activity per mL supernatant. In a most preferred embodiment, the recovered supernatant has at least 10 Units FRETS-VWF73 activity per mL supernatant. In certain embodiments, the methods provide sustainably improved FRETS production. For example, in certain embodiments, supernatant having at least 4 Units FRETS-VWF73 activity per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a preferred embodiment, supernatant having at least 6 Units FRETS-VWF73 activity per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a more preferred embodiment, supernatant having at least 8 Units FRETS-VWF73 activity per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a most preferred embodiment, supernatant having at least 10 Units FRETS-VWF73 activity per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days.

In one embodiment of the methods described above, the cell culture results in the production of at least 800 mU of FRETS-VWF73 activity per 10^6 cells present in the culture per day (i.e., q FRETS). In a preferred embodiment, the cell culture results in the production of at least 1 U of FRETS-VWF73 activity per 10^6 cells present in the culture per day.

In a more preferred embodiment, the cell culture results in the production of at least 1.2 U of FRET5-VWF73 activity per 10^6 cells present in the culture per day. In a most preferred embodiment, the cell culture results in the production of at least 1.4 U of FRET5-VWF73 activity per 10^6 cells present in the culture per day. In certain embodiments, the methods provide sustainably improved cell-specific FRET5-VWF73 productivity (q FRET5). For example, in certain embodiments, the cell culture results in the production of at least 800 mU of FRET5-VWF73 activity per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a preferred embodiment, the cell culture results in the production of at least 1 U of FRET5-VWF73 activity per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a more preferred embodiment, the cell culture results in the production of at least 1.2 U of FRET5-VWF73 activity per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a most preferred embodiment, the cell culture results in the production of at least 1.4 U of FRET5-VWF73 activity per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days.

In one embodiment of the methods described above, the cell culture results in the production of at least 1 mg rA13, as measured by ELISA, per liter culture per day (P ELISA). In a preferred embodiment, the cell culture results in the production of at least 1.5 mg rA13, as measured by ELISA, per liter culture per day. In a more preferred embodiment, the cell culture results in the production of at least 2 mg rA13, as measured by ELISA, per liter culture per day. In certain embodiments, the methods provide sustainably improved rA13 production. For example, in certain embodiments, the cell culture results in the production of at least 1 mg rA13, as measured by ELISA, per liter culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a preferred embodiment, the cell culture results in the production of at least 1.5 mg rA13, as measured by ELISA, per liter culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a more preferred embodiment, the cell culture results in the production of at least 2 mg rA13, as measured by ELISA, per liter culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days.

In one embodiment of the methods described above, the cell culture results in the production of at least 0.5 μ g rA13, as measured by ELISA, per 10^6 cells present in the culture per day (i.e., q ELISA). In a preferred embodiment, the cell culture results in the production of at least 0.7 μ g rA13, as measured by ELISA, per 10^6 cells present in the culture per day. In a more preferred embodiment, the cell culture results in the production of at least 0.9 μ g rA13, as measured by ELISA, per 10^6 cells present in the culture per day. In certain embodiments, the methods provide sustainably improved q ELISA production. For example, in certain embodiments, the cell culture results in the production of at least 0.5 μ g rA13, as measured by ELISA, per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a preferred embodiment, the cell culture results in the production of at least 0.7 μ g rA13, as measured by ELISA, per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a more preferred embodiment, the cell culture results in the production of at least 0.9 μ g rA13,

as measured by ELISA, per 10^6 cells present in the culture daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days.

In one embodiment of the methods described above, the recovered supernatant has at least 3 μ g rA13, as measured by ELISA, per mL supernatant. In a preferred embodiment, the recovered supernatant has at least 4 μ g rA13, as measured by ELISA, per mL supernatant. In a more preferred embodiment, the recovered supernatant has at least 5 μ g rA13, as measured by ELISA, per mL supernatant. In a most preferred embodiment, the recovered supernatant has at least 6 μ g rA13, as measured by ELISA, per mL supernatant. In certain embodiments, the methods provide sustainably improved rA13 production. For example, in certain embodiments, supernatant having at least 3 μ g rA13 per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a preferred embodiment, supernatant having at least 4 μ g rA13 per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a more preferred embodiment, supernatant having at least 5 μ g rA13 per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days. In a most preferred embodiment, supernatant having at least 6 μ g rA13 per mL are recovered daily for at least 7 days, or at least 14, 21, 28, 35, 42, 49, 56, 63, 70, or more days.

In one embodiment of the methods described above, the cell culture solution further comprises an ammonium concentration of less than 10 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 10 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 5 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 5 mM for at least 7 days. In a preferred embodiment, the cell culture solution comprises an ammonium concentration of no more than 4 mM. In another specific embodiment, the ammonium concentration of the cell culture is maintained at an ammonium concentration of no more than 4 mM for at least 7 days. In yet other embodiments, the cell culture solution comprises an ammonium concentration of no more than 10 mM, or no more than 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, 1 mM, or less. In yet another embodiment, the ammonium concentration of the cell culture is maintained at a low level for the duration of the process (i.e., for the entire time the culture is being used to produce rA13). In a particular embodiment, the culture solution has a copper and ammonium concentration according to any one of variations 1 to 440, as provided in Table 1.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper and zinc; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In one embodiment, the culture medium contains at least 1 μ g/L copper and at least 2 μ M zinc. In other embodiments, the media contains at least 2 μ g/L copper or at least 4 μ g/L copper. In one embodiment

wherein the media is supplemented with copper, the culture medium also contains at least at or about 5 μM zinc. In one embodiment, the culture medium also contains at or about between 2 μM and 12 μM zinc. In another embodiment, the culture medium also contains at or about between 5 μM and 12 μM zinc. In yet other embodiments, the culture medium also may contain at least at or about 2 μM , or at least at or about 3 μM , 4 μM , 5 μM , 6 μM , 7 μM , 8 μM , 9 μM , 10 μM , 11 μM , 12 μM , 13 μM , 14 μM , 15 μM , 20 μM , 25 μM , 30 μM , or more zinc. In one embodiment, the culture medium contains a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a preferred embodiment, at least 2000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper and calcium; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In one embodiment, the culture medium contains at least 1 $\mu\text{g/L}$ copper and at least 0.5 mM calcium. In other embodiments, the media contains at least 2 $\mu\text{g/L}$ copper or at least 4 $\mu\text{g/L}$ copper. In another embodiment wherein the media is supplemented with copper, the culture medium also contains at least 1.5 mM calcium. In one embodiment, the culture medium contains at or about between 0.5 mM and 1.5 mM calcium. In yet other embodiments, the culture medium may contain at least at or about 0.5 mM, or at least at or about 0.6 mM, 0.7 mM, 0.8 mM, 0.9 mM, 1.0 mM, 1.1 mM, 1.2 mM, 1.3 mM, 1.4 mM, 1.5 mM, 1.6 mM, 1.7 mM, 1.8 mM, 1.9 mM, 2.0 mM, 2.25 mM, 2.5 mM, 2.75 mM, 3.0 mM, 3.5 mM, 4.0 mM, 4.5 mM, 5.0 mM, or more calcium. In one embodiment, the culture medium contains a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In a preferred embodiment, at least 2000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper, zinc, and calcium; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that

rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In one embodiment, the culture medium has a calcium concentration of at least 0.5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a calcium concentration of at least 1.5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a calcium concentration between 0.5 mM and 1.5 mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In yet other embodiments, the culture medium has a calcium concentration of at least 0.6 mM, 0.7 mM, 0.8 mM, 0.9 mM, 1.0 mM, 1.1 mM, 1.2 mM, 1.3 mM, 1.4 mM, 1.5 mM, 1.6 mM, 1.7 mM, 1.8 mM, 1.9 mM, 2.0 mM, 2.25 mM, 2.5 mM, 2.75 mM, 3.0 mM, 3.5 mM, 4.0 mM, 4.5 mM, 5.0 mM, or more, and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a preferred embodiment, at least 2000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper and nicotinamide; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In one embodiment, the culture medium contains at least 1 $\mu\text{g/L}$ copper and at least 2 mg/L nicotinamide (vitamin B3). In other embodiments, the media contains at least 2 $\mu\text{g/L}$ copper or at least 4 $\mu\text{g/L}$ copper. In another embodiment wherein the media is supplemented with copper, the culture medium also contains at least 7 mg/L nicotinamide (vitamin B3). In one embodiment, the culture medium contains at or about between 2 mg/L and 10 mg/L nicotinamide (vitamin B3). In yet other embodiments, the culture medium may contain at least at or about 2 mg/L, 3 mg/L, 4 mg/L, 5 mg/L, 6 mg/L, 7 mg/L, 8 mg/L, 9 mg/L, 10 mg/L, 15 mg/L, 20 mg/L, or higher concentrations of nicotinamide (vitamin B3). In one embodiment, the culture medium contains a copper and nicotinamide concentration according to any one of variations 1321 to 1760, as set forth in Table 4. In a preferred embodiment, at least 2000 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRETTS-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000

Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper, zinc, and nicotinamide; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In one embodiment, the cell culture medium has a nicotinamide concentration of at least 2 mg/mL and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a nicotinamide concentration of at least 7 mg/mL mM and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In another specific embodiment, the culture medium has a nicotinamide concentration between 2 mg/mL and 10 mg/mL and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In yet other embodiments, the culture medium has a nicotinamide concentration of at least 2 mg/mL, 3 mg/mL, 4 mg/mL, 5 mg/mL, 6 mg/mL, 7 mg/mL, 8 mg/mL, 9 mg/mL, 10 mg/mL, 11 mg/mL, 12 mg/mL, 13 mg/mL, 14 mg/mL, 15 mg/mL, or more, and a copper and zinc concentration according to any one of variations 441 to 880, as set forth in Table 2. In a preferred embodiment, at least 2000 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

In one embodiment, the present invention provides a method for producing a recombinant ADAMTS13 (rA13) composition, the method comprising the steps of: (a) providing a basal cell culture media; (b) supplementing the basal cell culture media with copper, calcium, and nicotinamide; (c) providing one or more cells comprising a nucleic acid encoding a rA13 protein; (d) culturing the one or more cells in the copper supplemented cell culture media such that rA13 is expressed and excreted from the cells into a culture supernatant; and (e) recovering at least a portion of the culture supernatant, wherein at least 1500 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In one embodiment, the cell culture medium has a nicotinamide concentration of at least 2 mg/mL and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In another specific embodiment, the culture medium has a nicotinamide concentration of at least 7 mg/mL mM and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In another specific embodiment, the culture medium has a nicotinamide concentration between 2 mg/mL and 10 mg/mL and a copper and calcium concentration according to any one of variations 881 to

1320, as set forth in Table 3. In yet other embodiments, the culture medium has a nicotinamide concentration of at least 2 mg/mL, 3 mg/mL, 4 mg/mL, 5 mg/mL, 6 mg/mL, 7 mg/mL, 8 mg/mL, 9 mg/mL, 10 mg/mL, 11 mg/mL, 12 mg/mL, 13 mg/mL, 14 mg/mL, 15 mg/mL, or more, and a copper and calcium concentration according to any one of variations 881 to 1320, as set forth in Table 3. In a preferred embodiment, at least 2000 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a more preferred embodiment, at least 2500 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant. In a most preferred embodiment, at least 3000 Units FRET5-VWF73 activity per liter supplemented basal cell culture media per day is present in the recovered culture supernatant.

Recombinant ADAMTS proteins can be produced by expression in any suitable prokaryotic or eukaryotic host system. Examples of eukaryotic cells include, without limitation, mammalian cells, such as CHO, COS, HEK 293, BHK, SK-Hep, and HepG2; insect cells, for example SF9 cells, SF21 cells, S2 cells, and High Five cells; and yeast cells, for example *Saccharomyces* or *Schizosaccharomyces* cells. In one embodiment, the ADAMTS proteins can be expressed in bacterial cells, yeast cells, insect cells, avian cells, mammalian cells, and the like. For example, in a human cell line, a hamster cell line, or a murine cell line. In one particular embodiment, the cell line is a CHO, BHK, or HEK cell line. In a preferred embodiment, the cell line is a CHO cell line. In a specific embodiment, CHO clones capable of stably expressing rA13 are prepared by cotransfecting a CHO cell with coding sequences for rA13 and a dihydrofolate reductase (e.g., a murine dhfr gene) and selecting for growth in the presence of increasing levels of methotrexate.

In one embodiment, the cells may be any mammalian cell that can be cultured, preferably in a manufacturing process (i.e., at least 10 liter, preferably at least 100 liters), to produce a desired ADAMTS protein such as ADAMTS13. Examples include the monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR, such as the DUKX-B11 subclone (CHO, Uriaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse Sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HeLa, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., *Annals N. Y. Acad. Sci.*, 383:44-68 (1982)); MRC 5 cells; FS4 cells; and the human hepatoma line (Hep G2). Preferably, the cell line is a rodent cell line, especially a hamster cell line such as CHO or BHK.

A wide variety of vectors can be used for the expression of an ADAMTS protein (e.g., ADAMTS13) and can be selected from eukaryotic and prokaryotic expression vectors. In certain embodiments, a plasmid vector is contemplated for use in expressing an ADAMTS protein (e.g., ADAMTS13). In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with

these hosts. The vector can carry a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. The plasmid will comprise a nucleotide sequence encoding an ADAMTS protein (e.g., ADAMTS13) operable linked to one or more control sequences, for example, a promoter.

A preferred method of preparing stable CHO cell clones expressing a recombinant ADAMTS protein is as follows. A DHFR deficient CHO cell line DUKX-B11 is transfected with a DHFR expression vector to allow for expression of the relevant recombinant protein. An exemplary method is described by Plaimauer et al. (Blood. 2002 Nov. 15; 100(10):3626-32. Epub 2002 Jul. 12), the content of which is hereby incorporated by reference in its entirety for all purposes. Selection is carried out by growth in Hypoxanthine/Thymidine (HT) free media and amplification of the relevant region coding for expression of the recombinant ADAMTS protein and DHFR gene is achieved by propagation of the cells in increasing concentrations of methotrexate. Where appropriate, CHO cell lines may be adapted for growth in serum and/or protein free medium, essentially as described in U.S. Pat. No. 6,100,061 (Reiter et al., Immuno Aktiengesellschaft).

In another preferred embodiment, stable HEK293 cells are prepared by transfecting with a construct containing a hygromycin selectable marker and selecting transformants by antibiotic resistance.

The ability of certain viruses to infect cells or enter cells via receptor-mediated endocytosis, and to integrate into host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign nucleic acids into cells (e.g., mammalian cells). Accordingly, in certain embodiments, a viral vector is used to introduce a nucleotide sequence encoding an ADAMTS protein (e.g., ADAMTS13) into a host cell for expression. The viral vector will comprise a nucleotide sequence encoding an ADAMTS protein (e.g., ADAMTS13) operable linked to one or more control sequences, for example, a promoter. Alternatively, the viral vector may not contain a control sequence and will instead rely on a control sequence within the host cell to drive expression of the ADAMTS protein. Non-limiting examples of virus vectors that may be used to deliver a nucleic acid include Adenoviral vectors, AAV vectors, and Retroviral vectors.

In one embodiment, an Adenovirus expression vector include those constructs containing adenovirus sequences sufficient to support packaging of the construct and to ultimately express an ADAMTS construct that has been cloned therein. Adenoviral vectors allow for the introduction of foreign sequences up to 7 kb (Grunhaus et al., Seminar in Virology, 200(2):535-546, 1992).

In another embodiment, an adeno-associated virus (AAV) can be used to introduce a nucleotide sequence encoding an ADAMTS protein (e.g., ADAMTS13) into a host cell for expression. AAV systems have been described previously and are generally well known in the art (Kelleher and Vos, Biotechniques, 17(6):1110-7, 1994; Cotten et al., *Proc Natl Acad Sci USA*, 89(13):6094-6098, 1992; Curiel, *Nat Immunol*, 13(2-3):141-64, 1994; Muzyczka, *Curr Top Microbiol Immunol*, 158:97-129, 1992). Details concerning the generation and use of rAAV vectors are described, for example, in U.S. Pat. Nos. 5,139,941 and 4,797,368, each incorporated herein by reference in their entireties for all purposes.

In one embodiment, a retroviral expression vector can be used to introduce a nucleotide sequence encoding an ADAMTS protein (e.g., ADAMTS13) into a host cell for expression. These systems have been described previously

and are generally well known in the art (Mann et al., *Cell*, 33:153-159, 1983; Nicolas and Rubinstein, In: *Vectors: A survey of molecular cloning vectors and their uses*, Rodriguez and Denhardt, eds., Stoneham: Butterworth, pp. 494-513, 1988; Temin, In: *Gene Transfer*, Kucherlapati (ed.), New York: Plenum Press, pp. 149-188, 1986). In a specific embodiment, the retroviral vector is a lentiviral vector (see, for example, Naldini et al., *Science*, 272(5259):263-267, 1996; Zufferey et al., *Nat Biotechnol*, 15(9):871-875, 1997; Blomer et al., *J Virol*, 71(9):6641-6649, 1997; U.S. Pat. Nos. 6,013,516 and 5,994,136).

Non-limiting examples of vectors for prokaryotic expression include plasmids such as pRSET, pET, pBAD, etc., wherein the promoters used in prokaryotic expression vectors include lac, trc, trp, recA, araBAD, etc. Examples of vectors for eukaryotic expression include: (i) for expression in yeast, vectors such as pAO, pPIC, pYES, pMET, using promoters such as AOX1, GAP, GAL1, AUG1, etc; (ii) for expression in insect cells, vectors such as pMT, pAc5, pIB, pMIB, pBAC, etc., using promoters such as PH, p10, MT, Ac5, OpIE2, gp64, polh, etc., and (iii) for expression in mammalian cells, vectors such as pSVL, pCMV, pRc/RSV, pcDNA3, pBPV, etc., and vectors derived from viral systems such as vaccinia virus, adeno-associated viruses, herpes viruses, retroviruses, etc., using promoters such as CMV, SV40, EF-1, UbC, RSV, ADV, BPV, and β -actin. An exemplary vector for expressing rA13 is described by Plaimauer et al. (Blood. 2002 Nov. 15; 100(10):3626-32. Epub 2002 Jul. 12), the content of which is hereby incorporated by reference in its entirety for all purposes.

In certain embodiments, the cell-culture methods of the invention may comprise the use of a microcarrier. The present invention provides, among other aspect, methods of large-scale ADAMTS protein expression. In some embodiments, the cell-cultures of the embodiments can be performed in large bioreactors under conditions suitable for providing high volume-specific culture surface areas to achieve high cell densities and protein expression. One means for providing such growth conditions is to use microcarriers for cell-culture in stirred tank bioreactors. In another embodiment, these growth requirements are met via the use of a suspension cell culture.

V. Specific Embodiments

A. Recombinant Von Willebrand Factor (rVWF)

Recombinant vWF can be expressed in mammalian cells, but the specific activity of the vWF can vary widely depending on the cell culture conditions and has not been shown to be comparable or equal to that of vWF isolated from blood plasma. The present invention is based in-part on the surprising result that cell culture media having at least 2.4 $\mu\text{g/L}$ of copper provides an advantageous effect of promoting expression of high molecular weight vWF having a high specific activity. In particular, the high molecular weight, recombinant vWF of the present invention can include a highly multimeric form comprising about 14 to about 22 dimers and a specific Ristocetin activity of at least about 30 mU/ μg . The cell culture processes according to the present invention also allow for maintaining low NH_4^+ levels (e.g., less than 10 mM) during the upstream process in cell culture systems, thereby reducing deleterious effects to post-translational modifications. It is believed that the present invention provides for the first time cell culture conditions comprising a medium having a suitable copper concentration in

combination with appropriate levels of ammonium in the supernatant to express a highly multimeric vWF with a high specific activity.

In one aspect, the present invention relates to cell culture conditions for producing recombinant, high molecular weight vWF with a high specific activity. The cell culture conditions of the present invention can include, for example, a cell culture medium with an increased copper concentration and/or cell culture supernatant with a low ammonium (NH_4^+) concentration. The present invention also provides methods for cultivating cells in cell culture conditions to express a high molecular weight vWF with a high specific activity.

In one aspect, the present invention provides a cell culture solution for producing high molecular weight, recombinant vWF protein, the cell culture solution comprising: a cell culture medium comprising a copper concentration of at least about 2.4 $\mu\text{g/L}$; a cell culture supernatant comprising an ammonium concentration of less than 10 mM; and a plurality of cells expressing highly multimeric vWF protein, wherein the vWF protein comprises a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the cell cultures described above, the cell culture solution comprises a medium supplement comprising copper.

In one specific embodiment of the cell cultures described above, the medium supplement comprises a hydrolysate, optionally a soy hydrolysate.

In one specific embodiment of the cell cultures described above, the medium supplement comprises a copper salt, a copper chelate, or a combination thereof.

In one specific embodiment of the cell cultures described above, the copper salt is selected from the group consisting of copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide.

In one specific embodiment of the cell cultures described above, the copper concentration is at least about 4 $\mu\text{g/L}$.

In one specific embodiment of the cell cultures described above, the copper concentration is from about 2.4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$.

In one specific embodiment of the cell cultures described above, the vWF protein comprises about 14 to about 22 dimers.

In one aspect, the present invention provides a method of producing a high molecular weight, recombinant vWF protein, the method comprising the steps of: a) providing a culture of cells comprising a nucleic acid encoding recombinant vWF protein; b) expressing vWF protein in the cells under cell culture conditions comprising a cell culture medium comprising a copper concentration of at least about 2.4 $\mu\text{g/L}$ and a cell culture supernatant comprising an ammonium concentration less than about 10 mM, wherein the vWF protein is highly multimeric vWF protein and comprises a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the methods described above, the cells are mammalian cells.

In one specific embodiment of the methods described above, the cells are from a continuous cell line.

In one specific embodiment of the methods described above, the cells are CHO cells.

In one specific embodiment of the methods described above, the copper concentration is at least about 4 $\mu\text{g/L}$.

In one specific embodiment of the methods described above, the copper concentration is from about 2.4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$.

In one specific embodiment of the methods described above, the recombinant vWF protein has a specific Ristocetin Cofactor activity of at least about 50 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the methods described above, the recombinant vWF protein has a specific Ristocetin Cofactor activity from about 30 $\text{mU}/\mu\text{g}$ to about 100 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the methods described above, the recombinant vWF protein comprises about 14 to about 22 dimers.

In one aspect, the present invention provides a high molecular weight, recombinant vWF protein produced by a process, the process comprising the steps of: a) providing a culture of cells comprising a nucleic acid encoding recombinant vWF protein; and b) expressing vWF protein in the cells under cell culture conditions comprising a cell culture medium comprising a copper concentration of at least 2.4 $\mu\text{g/L}$ and a cell culture supernatant comprising an ammonium concentration less than 10 mM, wherein the vWF protein is highly multimeric vWF protein and comprises a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the rVWF compositions described above, the recombinant vWF protein has a specific Ristocetin Cofactor activity of at least about 50 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the rVWF compositions described above, the recombinant vWF protein has a specific Ristocetin Cofactor activity from about 30 $\text{mU}/\mu\text{g}$ to about 100 $\text{mU}/\mu\text{g}$.

In one specific embodiment of the rVWF compositions described above, the recombinant vWF protein comprises about 14 to about 22 dimers.

In one aspect, the present invention provides a cell culture solution for producing high molecular weight, recombinant vWF protein, the cell culture solution comprising: a cell culture medium comprising a copper concentration of at least about 2.4 $\mu\text{g/L}$; a cell culture supernatant comprising an ammonium concentration of less than 10 mM; and a plurality of cells expressing highly multimeric vWF protein, wherein the vWF protein comprises about 14 to about 22 dimers and a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$.

In one aspect, the present invention relates to cell culture conditions for producing recombinant high molecular weight vWF that is in highly multimeric form with a high specific activity. The cell culture conditions of the present invention can include, for example, a cell culture medium with an increased copper concentration and cell culture supernatant with a low ammonium (NH_4^+) concentration. The present invention also provides methods for cultivating cells in cell culture conditions to express a high molecular weight vWF with a high specific activity.

In one aspect, the present invention includes a cell culture solution for producing high molecular weight, recombinant vWF, comprising a cell culture medium comprising a copper concentration of at least about 2.4 $\mu\text{g/L}$; a cell culture supernatant comprising an ammonium concentration of less than 10 mM; and a plurality of cells expressing highly multimeric vWF comprising about 14 to about 22 dimers and a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$.

In one embodiment of the cell culture solutions described above, at least 10% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In a specific embodiment, at least 15% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 20% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at

least 25% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 30% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In certain embodiments, the copper concentration can be at least about 4 $\mu\text{g/L}$ or the copper concentration can range from about 2.4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$. In some embodiments, the cell culture media comprise a medium supplement comprising copper. In certain embodiments, the medium supplement can comprise a hydrolysate or a copper salt, copper chelate, or a combination thereof. In some embodiments, the copper salt can include copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, or copper oxide. In certain embodiments, the cells that can be from a continuous cell line and can include mammalian cells, such as CHO cells. In some embodiments, the recombinant vWF has a specific Ristocetin Cofactor activity of at least about 50 $\text{mU}/\mu\text{g}$, or the specific Ristocetin Cofactor activity can range from about 30 $\text{mU}/\mu\text{g}$ to about 100 $\text{mU}/\mu\text{g}$.

In another aspect, the present invention includes a method of producing a high molecular weight, recombinant vWF, comprising a) providing a culture of cells; b) introducing a nucleic acid sequence coding for vWF; c) selecting the cells carrying the nucleic acid sequence; and, d) expressing vWF in the cells under cell culture conditions comprising a cell culture medium comprising a copper concentration of at least about 2.4 $\mu\text{g/L}$ and a cell culture supernatant comprising an ammonium concentration less than about 10 mM, wherein the vWF is highly multimeric vWF comprising about 14 to about 22 dimers and a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$. In one embodiment of the cell culture supernatant described above, at least 10% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In a specific embodiment, at least 15% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 20% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 25% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 30% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In some embodiments, the cells that can be from a continuous cell line and can include mammalian cells, such as CHO cells. In certain embodiments, the copper concentration can be at least about 4 $\mu\text{g/L}$ or the copper concentration can range from about 2.4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$. In some embodiments, the cell culture media comprise a medium supplement comprising copper. In certain embodiments, the medium supplement can comprise a hydrolysate or a copper salt, copper chelate, or a combination thereof. In some embodiments, the copper salt can include copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, or copper oxide. In some embodiments, the recombinant vWF has a specific Ristocetin Cofactor activity of at least about 50 $\text{mU}/\mu\text{g}$, or the specific Ristocetin Cofactor activity can range from about 30 $\text{mU}/\mu\text{g}$ to about 100 $\text{mU}/\mu\text{g}$.

In yet another aspect, the present invention includes a high molecular weight, recombinant vWF produced by a process, comprising the steps of: a) providing a culture of cells; b) introducing a nucleic acid sequence coding for vWF; c) selecting the cells carrying the nucleic acid sequence; and, d) expressing vWF in the cells under cell culture conditions comprising a cell culture medium com-

prising a copper concentration of at least 2.4 $\mu\text{g/L}$ and a cell culture supernatant comprising an ammonium concentration less than 10 mM, wherein the vWF is highly multimeric vWF comprising about 14 to about 22 dimers and a specific Ristocetin activity of at least about 30 $\text{mU}/\mu\text{g}$. In one embodiment of the cell culture supernatant described above, at least 10% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In a specific embodiment, at least 15% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 20% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 25% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 30% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In some embodiments, the cells that can be from a continuous cell line and can include mammalian cells, such as CHO cells. In certain embodiments, the copper concentration can be at least about 4 $\mu\text{g/L}$ or the copper concentration can range from about 2.4 $\mu\text{g/L}$ to about 20 $\mu\text{g/L}$. In some embodiments, the cell culture media comprise a medium supplement comprising copper. In certain embodiments, the medium supplement can comprise a hydrolysate or a copper salt, copper chelate, or a combination thereof. In some embodiments, the copper salt can include copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, or copper oxide. In some embodiments, the recombinant vWF has a specific Ristocetin Cofactor activity of at least about 50 $\text{mU}/\mu\text{g}$, or the specific Ristocetin Cofactor activity can range from about 30 $\text{mU}/\mu\text{g}$ to about 100 $\text{mU}/\mu\text{g}$.

In one aspect, the present invention provides a composition comprising recombinant Von Willebrand Factor (rVWF) having a specific ristocetin cofactor activity of at least 30 $\text{mU}/\mu\text{g}$. In a preferred embodiment, the composition has a specific ristocetin cofactor activity of at least 40 $\text{mU}/\mu\text{g}$. In a more preferred embodiment, the composition has a specific ristocetin cofactor activity of at least 50 $\text{mU}/\mu\text{g}$. In a more preferred embodiment, the composition has a specific ristocetin cofactor activity of at least 60 $\text{mU}/\mu\text{g}$. In a more preferred embodiment, the composition has a specific ristocetin cofactor activity of at least 70 $\text{mU}/\mu\text{g}$. In yet a more preferred embodiment, the composition has a specific ristocetin cofactor activity of at least 80 $\text{mU}/\mu\text{g}$.

In one embodiment of the compositions described above, at least 10% of the rVWF in the composition is present in a high molecular weight VWF multimer of more than 10 dimers. In a specific embodiment, at least 15% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 20% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 25% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers. In another specific embodiment, at least 30% of the rVWF is present in a high molecular weight VWF multimer of more than 10 dimers.

In one embodiment of the compositions described above, the composition comprises a culture supernatant. In one specific embodiment the culture supernatant is a mammalian cell culture supernatant. In a more specific embodiment, the mammalian cell culture supernatant is a CHO cell supernatant.

In one embodiment of the compositions described above, the rVWF is expressed in a cell culture comprising at least 2.4 $\mu\text{g/L}$ copper. In a specific embodiment, the cell culture comprises at least 4 $\mu\text{g/L}$ copper. In a more specific embodiment, the culture comprises between 2.4 $\mu\text{g/L}$ and 20 $\mu\text{g/L}$ copper. In one embodiment, the copper is provided as a copper salt, a copper chelate, or a combination thereof. In a specific embodiment, the copper salt is selected from the group consisting of copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide.

In one embodiment of the compositions described above, the cell culture is a batch culture.

In one embodiment of the compositions described above, the cell culture is a continuous culture. In a specific embodiment, the continuous culture is performed in chemostatic mode. In another specific embodiment, the continuous culture is performed in perfusion mode.

In one embodiment of the compositions described above, the level of NH_4^+ in the culture is maintained at a concentration below 4 mM.

In one embodiment of the compositions described above, the cell density of the culture is maintained at less than 2.5×10^6 cells per mL.

In one embodiment of the compositions described above, the cell density of the culture is maintained at less than 2.0×10^6 cells per mL.

In one embodiment of the compositions described above, the culture is maintained at less than 1.5×10^6 cells per mL.

In one embodiment of the compositions described above, the rVWF is co-expressed with recombinant Factor VIII (rFVIII). In a specific embodiment, a majority of the co-expressed rFVIII has been removed. In a more specific embodiment, the ratio of rVWF to rFVIII in the composition is at least 10:1.

In one embodiment of the compositions described above, the composition is formulated for pharmaceutical administration. In a specific embodiment, the composition is formulated for intravenous, subcutaneous, or intramuscular administration.

In one embodiment of the compositions described above, the composition is lyophilized.

In yet another aspect, the present invention includes a high molecular weight, recombinant vWF produced by a process, comprising the steps of: a) providing a culture of cells; b) introducing a nucleic acid sequence coding for vWF; c) selecting the cells carrying the nucleic acid sequence; and, d) expressing vWF in the cells under cell culture conditions comprising a cell culture medium comprising a copper concentration of at least 2.4 $\mu\text{g/L}$ and a cell culture supernatant comprising an ammonium concentration less than 10 mM, wherein the vWF is highly multimeric vWF comprising about 14 to about 22 dimers and a specific Ristocetin activity of at least about 30 mU/ μg . It is understood that all of the embodiments and concentrations described in the "Cell Culture Media" and "Methods of Producing Recombinant vWF" sections above can apply here.

The recombinant vWF of the present invention can include a high molecular weight recombinant vWF protein having a high specific activity. In one embodiment, the vWF of the present invention is a highly multimeric form of vWF. In some embodiments, the highly multimeric form of vWF includes at least up to about 14 dimers and in other embodiments at least up to about 22 dimers. In yet other embodiments, the highly multimeric form of vWF can range from about 10 to about 20 dimers, or from about 15 to about 25

dimers, or from about 20 to about 40 dimers. In certain embodiments, the recombinant vWF is comparable to plas-matic vWF.

As described herein, the present invention provides the surprising result that increased copper concentration in cell culture media can produce high molecular weight vWF with high specific activity. Cell culture media comprising a copper concentration, e.g., greater than about 2.4 $\mu\text{g/L}$ can increase the yield of recombinant multimeric vWF, as compared to media without copper. In certain embodiments, the percentage of multimeric vWF (i.e., rVWF comprising at least 2 dimers) can be greater than about 50%, or greater than about 75%, or greater than about 90%. The multimeric distribution of the vWF can be analyzed using standard techniques such as, e.g., in Agarose electrophoresis under non-reducing conditions.

As provided herein, the recombinant vWF produced by the methods of the present invention can have a high specific activity, e.g., a high specific Ristocetin Cofactor activity. In one embodiment, the recombinant vWF produced by the methods of the present invention can include a specific Ristocetin Cofactor activity of at least 30 mU/ μg and in another embodiment at least 50 mU/ μg . In other embodiments, the specific Ristocetin Cofactor activity can range from about 30 mU/ μg to about 100 mU/ μg or from about 50 mU/ μg to about 100 mU/ μg .

B. Recombinant ADAMTS13 (rA13)

The ADAMTS proteins (i.e., ADAMTS-1 to ADAMTS-20) are a family of secreted zinc metalloproteinases that share a common modular domain organization (for review, see, Flannery C. R., *Front Biosci.* 2006 Jan. 1; 11:544-69). All of the ADAMTS protein share a common core domain architecture, consisting of a signal peptide, followed by a prodomain, a zinc-dependent metalloproteinase catalytic domain, a disintegrin-like domain, a thrombospondin type I repeat, a cysteine-rich domain, and a spacer domain (Apte S. S., *J Biol Chem.* 2009 Nov. 13; 284(46):31493-7). Additionally, all but ADAMTS-4 contain at least one more thrombospondin type I repeat domain, and many of the ADAMTS protein contain one or more additional ancillary domains. Notably, it has been reported that all ADAMTS protein appear to contain at least one calcium binding site and at least one zinc binding site located within the metalloproteinase catalytic domain (Andreini et al., *J. Proteome Res.*, 2005, 4 (3), pp 881-888).

Biological roles for ADAMTS proteins have been reported for various diseases and conditions, including, Antiangiogenesis, Renal interstitial fibrosis, Bone remodeling, Ovarian folliculogenesis, Atherosclerosis, Urogenital development, and Tumor growth/remodeling (ADAMTS-1); Ehler-Danlos syndrome type 7C and Bovine dermatopraxia (ADAMTS-2); Arthritis, Atherosclerosis, and Tendinopathy (ADAMTS-4); Arthritis and Glioblastoma (ADAMTS-5); Arthritis (ADAMTS-7); Antiangiogenesis, Brain malignancy, Arthritis, and Atherosclerosis (ADAMTS-8); Arthritis (ADAMTS-9, -12); Thrombotic thrombocytopenic purpura (ADAMTS-13); and Antithrombosis/stroke (ADAMTS18) (for review, see, Lin and Liu, *Open Access Rheumatology Research and Reviews* 2009:1 121-131).

Recombinant ADAMTS13 (A13) has been expressed before in mammalian cells, however the specific activity varies widely dependent on the cell culture conditions. It has been found that many commercially available culture mediums are not sufficient for expression of rA13 with high specific activities, expressed as the ratio of activity, measured by FRET-VWF73 assay, to antigen content, as determined by ELISA. In one aspect, the methods provided

herein are based on several advantageous findings that allow for cell-culture expression of rA13 having increased levels of total and specific activity.

Accordingly, due to the shared structure-function relationship between the ADAMTS family of secreted metalloproteinases, the methods provided by the present invention allow for the expression of all ADAMTS proteins in cell culture and recovery from the cell medium.

In one aspect, the present invention provides a composition comprising recombinant ADAMTS13 (rA13) having a specific FRETTS-VWF activity of at least 1600 mU/ μ g.

In one embodiment of the compositions described above, the rA13 has a specific FRETTS-VWF activity of at least 800 mU/ μ g.

In one embodiment of the compositions described above, the composition comprises a culture supernatant. In a specific embodiment, the culture supernatant is a mammalian cell culture supernatant. In a more specific embodiment, the mammalian cell culture supernatant is a CHO cell supernatant.

In one embodiment of the compositions described above, the rA13 is expressed in a cell culture comprising at least 1 μ g/L copper. In a specific embodiment, the cell culture comprises at least 2 μ g/L copper. In a more specific embodiment, the culture comprises between 2 μ g/L and 20 μ g/L copper.

In one embodiment of the compositions described above, the copper is provided as a copper salt, a copper chelate, or a combination thereof. In a specific embodiment, the copper salt is selected from the group consisting of copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide.

In one embodiment of the compositions described above, the cell culture is a batch culture.

In one embodiment of the compositions described above, the cell culture is a continuous culture. In a specific embodiment, the continuous culture is performed in chemostatic mode. In another specific embodiment, the continuous culture is performed in perfusion mode.

In one embodiment of the compositions described above, the level of NH_4^+ in the culture is maintained at a concentration below 5 mM.

In one embodiment of the compositions described above, the level of NH_4^+ in the culture is maintained at a concentration below 4 mM.

In one embodiment of the compositions described above, the cell density of the culture is maintained at less than 4.0×10^6 cells per mL. In a specific embodiment, the cell density of the culture is maintained at less than 3.0×10^6 cells per mL. In a specific embodiment, the cell density of the culture is maintained at less than 2.0×10^6 cells per mL. In a more specific embodiment, the cell density of the culture is maintained at less than 1.5×10^6 cells per mL.

In one embodiment of the compositions described above, the composition is formulated for pharmaceutical administration. In a specific embodiment, the composition is formulated for intravenous, subcutaneous, or intramuscular administration.

In one embodiment of the compositions described above, the composition is lyophilized.

VI. Formulations

In one aspect, the formulations comprising the recombinant therapeutic proteins rVWF or rA13 of the present invention are lyophilized prior to administration. Lyophilization is carried out using techniques common in

the art and should be optimized for the composition being developed [Tang et al., *Pharm Res.* 21:191-200, (2004) and Chang et al., *Pharm Res.* 13:243-9 (1996)].

Methods of preparing pharmaceutical formulations can include one or more of the following steps: adding a stabilizing agent as described herein to said mixture prior to lyophilizing, adding at least one agent selected from a bulking agent, an osmolarity regulating agent, and a surfactant, each of which as described herein, to said mixture prior to lyophilization. A lyophilized formulation is, in one aspect, at least comprised of one or more of a buffer, a bulking agent, and a stabilizer. In this aspect, the utility of a surfactant is evaluated and selected in cases where aggregation during the lyophilization step or during reconstitution becomes an issue. An appropriate buffering agent is included to maintain the formulation within stable zones of pH during lyophilization.

The standard reconstitution practice for lyophilized material is to add back a volume of pure water or sterile water for injection (WFI) (typically equivalent to the volume removed during lyophilization), although dilute solutions of antibacterial agents are sometimes used in the production of pharmaceuticals for parenteral administration [Chen, *Drug Development and Industrial Pharmacy*, 18:1311-1354 (1992)]. Accordingly, methods are provided for preparation of reconstituted recombinant VWF compositions comprising the step of adding a diluent to a lyophilized recombinant VWF composition of the invention.

The lyophilized material may be reconstituted as an aqueous solution. A variety of aqueous carriers, e.g., sterile water for injection, water with preservatives for multi dose use, or water with appropriate amounts of surfactants (for example, an aqueous suspension that contains the active compound in admixture with excipients suitable for the manufacture of aqueous suspensions). In various aspects, such excipients are suspending agents, for example and without limitation, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents are a naturally-occurring phosphatide, for example and without limitation, lecithin, or condensation products of an alkylene oxide with fatty acids, for example and without limitation, polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example and without limitation, heptadecaethyl-eneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example and without limitation, polyethylene sorbitan monooleate. In various aspects, the aqueous suspensions also contain one or more preservatives, for example and without limitation, ethyl, or n-propyl, p-hydroxybenzoate.

To administer compositions to human or test animals, in one aspect, the compositions comprises one or more pharmaceutically acceptable carriers. The phrases "pharmaceutically" or "pharmacologically" acceptable refer to molecular entities and compositions that are stable, inhibit protein degradation such as aggregation and cleavage products, and in addition do not produce allergic, or other adverse reactions when administered using routes well-known in the art, as described below. "Pharmaceutically acceptable carriers" include any and all clinically useful solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like, including those agents disclosed above.

The pharmaceutical formulations are administered intravenously, orally, topically, transdermally, parenterally, by inhalation spray, vaginally, rectally, or by intracranial injection. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intracisternal injection, or infusion techniques. Administration by intravenous, intradermal, intramuscular, intramammary, intraperitoneal, intrathecal, retrobulbar, intrapulmonary injection and or surgical implantation at a particular site is contemplated as well. Generally, compositions are essentially free of pyrogens, as well as other impurities that could be harmful to the recipient.

Single or multiple administrations of the compositions are carried out with the dose levels and pattern being selected by the treating physician. For the prevention or treatment of disease, the appropriate dosage depends on the type of disease to be treated, the severity and course of the disease, whether drug is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the drug, and the discretion of the attending physician.

In one aspect, formulations of the invention are administered by an initial bolus followed by a continuous infusion to maintain therapeutic circulating levels of drug product. As another example, the inventive compound is administered as a one-time dose. Those of ordinary skill in the art will readily optimize effective dosages and administration regimens as determined by good medical practice and the clinical condition of the individual patient. The frequency of dosing depends on the pharmacokinetic parameters of the agents and the route of administration. The optimal pharmaceutical formulation is determined by one skilled in the art depending upon the route of administration and desired dosage. See for example, Remington's Pharmaceutical Sciences, 18th Ed. (1990, Mack Publishing Co., Easton, Pa. 18042) pages 1435-1712, the disclosure of which is hereby incorporated by reference. Such formulations influence the physical state, stability, rate of in vivo release, and rate of in vivo clearance of the administered agents. Depending on the route of administration, a suitable dose is calculated according to body weight, body surface area or organ size. Appropriate dosages may be ascertained through use of established assays for determining blood level dosages in conjunction with appropriate dose-response data. The final dosage regimen is determined by the attending physician, considering various factors which modify the action of drugs, e.g. the drug's specific activity, the severity of the damage and the responsiveness of the patient, the age, condition, body weight, sex and diet of the patient, the severity of any infection, time of administration and other clinical factors. By way of example, a typical dose of a recombinant vWF of the present invention is approximately 50 U/kg, equal to 500 µg/kg. As studies are conducted, further information will emerge regarding the appropriate dosage levels and duration of treatment for various diseases and conditions.

VII. Methods of Treatment

The present invention further contemplates methods of treating a patient in need of rVWF or rA13 produced according to the methods described herein. Such methods of

treatment can include administration of pharmaceutical formulations comprising the recombinant rA13 or the high molecular weight, recombinant vWF of the present invention.

In another aspect, the present invention provides methods for therapeutic or prophylactic treatments comprising the administration of an rVWF or rA13 composition provided herein. Generally, for therapeutic applications, formulations are administered to a subject with a disease or condition associated with ADAMTS13 or VWF dysfunction or otherwise in need thereof, in a "therapeutically effective dose." Formulations and amounts effective for these uses will depend upon the severity of the disease or condition and the general state of the patient's health. Single or multiple administrations of the formulations may be administered depending on the dosage and frequency as required and tolerated by the patient.

In one embodiment, the present invention provides methods of treating or preventing a disease or condition associated with an ADAMTS13 or VWF dysfunction. In a further embodiment, the pharmaceutical formulations comprising recombinant vWF can be administered to treat diseases related to vWF, such as von Willebrand's disease or hemophilia. In another embodiment, the invention provides methods of treating or preventing a disease or condition associated with the formation and/or presence of one or more thrombus, comprising the administration of a rA13 composition provided herein. In another embodiment, the invention provides methods of disintegrating one or more thrombus in a subject in need thereof. In yet other embodiments, the invention provides methods of treating or preventing an infarction in subject in need thereof. Generally, the methods provided by the invention comprise administering an rADAMTS13 composition as provided herein to a subject in need thereof.

Non-limiting examples of disorders associated with the formation and/or the presence of one or more thrombus are hereditary thrombotic thrombocytopenic purpura (TTP), acquired TTP, arterial thrombosis, acute myocardial infarction (AMI), stroke, sepsis, and disseminated intravascular coagulation (DIC).

Non-limiting examples of disorders associated with an infarction, include without limitation, myocardial infarction (heart attack), pulmonary embolism, cerebrovascular events such as stroke, peripheral artery occlusive disease (such as gangrene), antiphospholipid syndrome, sepsis, giant-cell arteritis (GCA), hernia, and *volvulus*.

VIII. Examples

The present invention will now be further illustrated in the following examples, without being limited thereto.

Example 1

Continuous cell culture experiments were performed using cultures of a recombinant CHO cell line expressing vWF. The basal medium was DMEM/F12, which contained about 0.3 µg/L of Cu²⁺. The medium was supplemented with

soy hydrolysate and copper sulfate (CuSO₄·5H₂O) to bring the final copper concentration in the medium to greater than at least 2.4 µg/L.

Recombinant CHO cells expressing vWF were cultivated by continuous cell cultures so that the ammonium levels (NH₄⁺) were kept at a concentration less than about 10 mM. It was found that production systems providing a continuous supply of medium (e.g., perfusion or chemostat cultures) were preferable because conventional batch or fed batch techniques produced high NH₄⁺ concentrations at the end of the culture. At the end of the culture, the highly multimeric vWF was isolated and the specific Ristocetin Cofactor activity of the vWF was measured.

Example 2

Recombinant Factor VIII (rFVIII) and Von Willebrand Factor (rVWF) were co-expressed in batch cultures of GD8/6 cells to determine the effect of the composition of the culture medium on VWF expression and activity. Briefly, GD8/6 cells were batch cultivated in BAV-SP medium composed of a modified DMEM/F12 basal powder (Table 5) and additional supplements also containing 4 g/L of soy hydrolysate, see Table 6, with and without additional copper supplementation. To test the effect of low copper concentrations on rVWF expression and activity, basal BAV-SP media was used. The basal media contained 0.3 µg/L copper and was supplemented with soy hydrolysate, which contributed an additional 0.7 µg/L copper, as determined experimentally, providing a final copper concentration of 1.0 µg/L. As a comparison, the BAV-SP media used for the batch cultures was further supplemented with an additional 3.3 µg/L of Cu²⁺, providing a final copper concentration of 4.3 µg/L, to determine the effect high copper concentrations on VWF expression and activity.

TABLE 5

Composition of BAV-SP Culture Media	
BAV-SP-Medium Components	mg/L
<u>Amino Acids</u>	
L-Alanine	4.45
L-Arginine HCl	147.50
L-Asparagine-H ₂ O	30.21
L-Aspartic Acid	6.65
L-Cystèine HCl—H ₂ O	32.55
L-Cystine 2HCl	57.35
L-Glutamic Acid	7.35
Glycine	18.75
L-Histidine-H ₂ O HCl	31.48
Hydroxy-L-Proline	
L-Isoleucine	54.47
L-Leucine	59.05
L-Lysine HCl	91.25
L-Methionine	17.24
L-Phenylalanine	35.48
L-Proline	52.24
L-Serine	26.25
L-Threonine	53.45
L-Tryptophan	29.01
L-Tyrosine 2Na 2H ₂ O	55.79
L-Valine	52.85

TABLE 5-continued

Composition of BAV-SP Culture Media	
BAV-SP-Medium Components	mg/L
<u>Vitamins</u>	
Ascorbic Acid	3.499
Biotin	0.0035
Choline Chloride	8.980
D-Ca-Pantothenate	2.240
Folic Acid	2.650
I-Inositol	12.600
Nicotinamide	2.020
Pyridoxine HCl	2.031
Riboflavin	0.219
Thiamine HCl	2.170
Vitamin B12	0.680
<u>Anorganic Salts</u>	
Calcium Chloride (CaCl ₂)	116.600
Copper Sulfate (CuSO ₄ —5H ₂ O)	0.0013
Ferric Nitrate (Fe(NO ₃) ₃ —9H ₂ O)	0.050
Ferrous Sulfate (FeSO ₄ —7H ₂ O)	0.417
Magnesium Chloride (MgCl ₂)	28.640
Magnesium Sulfate (MgSO ₄)	48.840
Potassium Chloride (KCl)	311.800
Sodium Chloride (NaCl)	6995.500
Sodium phosphate (Na ₂ HPO ₄)	71.020
Sodium phosphate (NaH ₂ PO ₄)	62.500
Zinc Sulfate Heptahydrate (ZnSO ₄ —7H ₂ O)	0.432
Sodium selenite	0.0131
<u>Others</u>	
D-Glucose	3151
Linoleic Acid	0.042
Lipoic Acid	0.105
Putrescine 2HCl	0.081
Thymidine	0.365
Hypoxantine Na	2.390
Sodium Pyruvate	55.000

TABLE 6

Composition of BAV-SP Supplement	
Componets for final formulation	
L-Glutamine	600.00
Ethanolamine	1.530
Synperonic F68	250.000
Sodium Bicarbonate (NaHCO ₃)	2000.000
Soy peptone	4000.00
Total formulation	18596.8

GD8/6 cells expressing rVWF were grown in either low copper media (Table 7) or high copper media (Table 8) for 7 days. After 2 days the cultures were subcultured to perform a batch culture for a period of 5 days. Samples of the culture supernatant were tested daily for rVWF content (vWF ELISA), total (Ristocetin) and specific (Specific Activity) via a ristocetin cofactor assay. Various culture parameters, including cell count, cell viability, and ammonium concentration were also monitored daily (except for batch days 3 and 4).

Unexpectedly, cell cultures grown under high copper concentrations produced supernatants containing significantly higher total and specific rVWF activity (compare results in Table 7 and Table 8). For example, at batch day 4, the cell culture grown under high copper concentration contained 1.52 IU rVWF activity/mL, as compared to 0.2 IU rVWF activity/mL for the low copper culture. This is despite

the fact that the low copper cell culture produced nearly twice the amount of rVWF as the high copper cell culture. Furthermore, the specific activity of the supernatant obtained from the high copper culture was more than 13 times greater than that of the low copper culture supernatant (831 mU/10 μ g rVWF v. 62 mU/10 μ g rVWF).

As seen in Table 7 and Table 8, the total and specific ristocetin cofactor activity of the high copper culture was twice that of the low copper culture at batch day 1. Furthermore, in contrast to the subsequent increases in activity observed in the high copper culture, no increase in the amount of ristocetin cofactor activity was seen after batch day 1 for the low copper cell culture. Consistent with this result, agarose gel electrophoresis analysis of the rVWF multimer state revealed that a low concentration of high molecular weight rVWF, relative to the concentration of low molecular weight rVWF species, was present in the supernatant of the low copper cell culture at batch day 1, and that the relative concentration further decreased over time (FIGS. 1A and 1B). In contrast, supplementation of the culture medium with Cu^{2+} resulted in a consistent formation of ristocetin cofactor (RiCoF) active antigen through the fourth day. Consistently, no loss in the amount of stable high molecular weight rVWF multimers occurred through day 4 of the batch culture (FIGS. 1A and 1B). The densitometric results of the agarose electrophoresis gel shown in FIG. 1B revealed that under low copper conditions the culture was only able to produce a rVWF population in which more than 10% (i.e., 16.3%) of the rVWF was present in molecules

having more than 10 dimers for one day of the batch culture (i.e. "day3" sample in Lane 2 of the agarose gel of FIG. 1A), and notably, this population fell to only 4% at day 5 of the batch culture ("day 7" sample in Lane 6). In contrast under high copper conditions the relative amount of vWF multimers with more than 10 dimers is consistently around 30% through day 4 of the batch culture ("day 3" to "day 6" in Lanes 7-10; 28% to 31.4%).

Notably, beginning at batch day 5 of the high copper batch culture, when NH_4^+ levels exceeded 100 mg/L (greater than about 5.0 mM), the expression of additional antigen (18.3 to 35.4 μ g/L; compare batch days 4 and 5 of Table 8) did not result in a concomitant increase in the RiCoF activity present in the supernatant. Consistent with this result, the level of high molecular weight rVWF multimers at batch day 5 (day 7 of the culture) decreased relative to the concentration of low molecular weight rVWF multimers. Also only on batch day 5 ("day 7" sample of the culture in Lane 11) the relative amount of the vWF with more than 10 dimers is reduced to 21.4%.

Taken together, the data provided above demonstrate that the supplementation of copper concentrations in cell cultures expressing rVWF dramatically increases the total and specific rVWF ristocetin cofactor activity, as well the stable production of high molecular weight rVWF multimers. Furthermore, the data show a correlation between the presence of high NH_4^+ concentrations in the cell culture and the loss of rVWF ristocetin cofactor activity and high molecular weight rVWF multimer production.

TABLE 7

Expression of rVWF in mammalian cell culture performed in batch mode using BAV-SP media with a low copper concentration (1.0 μ g/L).						
batch day	Cell Count [10E6 cells/mL]	NH_4^+ [mg/l]	Viability [%]	vWF ELISA [μ g/ml]	Ristocetin [IU/mL]	Specific Activity mU/10 μ g
0	0.42	21	98.8	n.d.	n.d.	n.d.
1	0.78	43	n.d.	6.3	0.23	365
2	1.24	64	99.1	12.8	0.23	180
3	1.86	n.d.	n.d.	22.7	0.21	93
4	2.49	n.d.	n.d.	32.2	0.20	62
5	3.11	106	98.3	44.4	0.20	45

TABLE 8

Expression of rVWF in mammalian cell culture performed in batch mode using BAV-SP media with a high copper concentration (4.3 μ g/L).						
batch day	Cell Count [10E6 cells/mL]	NH_4^+ [mg/l]	Viability [%]	vWF ELISA [μ g/ml]	Ristocetin [IU/mL]	Specific Activity mU/10 μ g
0	0.40	21	99.3	n.d.	n.d.	n.d.
1	0.71	42	n.d.	4.8	0.40	833
2	1.23	63	99.5	8.7	0.62	713
3	1.85	n.d.	n.d.	15.0	1.07	713
4	2.54	n.d.	n.d.	18.3	1.52	831
5	3.32	109	98.9	35.4	1.55	438

Example 3

Recombinant Factor VIII (rFVIII) and Von Willebrand Factor (rVWF) were co-expressed in continuous cultures of GD8/6 cells operated under chemostatic conditions to determine the effect of the composition of the culture medium on VWF expression and activity. Briefly, GD8/6 cells were cultivated in BAV-SP medium containing 4 g/L of soy hydrolysate with and without copper supplementation as described in Example 2. To test the effect of low copper concentrations on rVWF expression and activity, basal BAV-SP media was used. The basal media contained 0.3 $\mu\text{g/L}$ copper and was supplemented with soy hydrolysate, which contributed an additional 0.7 $\mu\text{g/L}$ copper, providing a final copper concentration of 1.0 $\mu\text{g/L}$. As a comparison, the

BAV-SP media used for the batch cultures was further supplemented with an additional 3.3 $\mu\text{g/L}$ of Cu^{2+} , providing a final copper concentration of 4.3 $\mu\text{g/L}$, to determine the effect high copper concentrations on VWF expression and activity. Cultures grown in the presence of high and low copper concentrations were cultivated under both high (2.8×10^6 cells/mL) and low (appr. 1.4×10^6 cells/mL) cell densities.

As before, samples of the culture supernatant were tested for rVWF content (vWF ELISA), total (Ristocetin) and specific (Specific Activity) activity via a ristocetin cofactor assay. Various culture parameters, including cell count, cell viability, and ammonium concentration were also monitored. Data were generated from the steady state phase of weeks 2 and 3 of the chemostat cultures (Table 9 to Table 13).

TABLE 9

Mean data for rVWF expression in chemostatic cell culture during weeks 2 and 3.							
Cell Count	Copper Concentration	Cell Count [10E6 cells/mL]	NH4+ [mM]	Viability [%]	vWF		specific activity [mU/10 μg]
					ELISA [$\mu\text{g/ml}$]	Ristocetin [IU/mL]	
high	low	2.88	3.88	97.74	44.56	0.10	21.62
high	high	2.79	4.04	98.25	38.38	0.19	53.11
low	low	1.55	3.33	98.63	18.96	0.10	50.16
low	high	1.43	3.17	98.59	11.76	0.70	598.76

TABLE 10

rVWF expression in chemostatic cell culture under high cell count, low copper conditions during weeks 2 and 3.							
Day	Cell Count [10E6 cells/mL]	NH4+ [mM]	Viability [%]	vWF ELISA [$\mu\text{g/ml}$]	Ristocetin [IU/mL]	specific activity [mU/10 μg]	
8	2.54	3.7	98.20	39.8	0.095	23.86934673	
9	3.02	4.2	97.40				
10	2.97	3.9	97.90	41.3	0.095	23.00242131	
11	2.78						
13	2.91	3.8	97.60	41.7	0.095	22.78177458	
14	2.90	3.8	97.40				
15	3.05	3.9	97.70	44.7	0.095	21.25279642	
16	2.99	3.8	98.30				
17	2.76	3.9	97.40	55.3	0.095	17.17902351	
	2.88	3.88	97.74	44.56	0.10	21.62	

TABLE 11

rVWF expression in chemostatic cell culture under high cell count, high copper conditions during weeks 2 and 3.							
Day	Cell Count [10E6 cells/mL]	NH4+ [mM]	Viability [%]	vWF ELISA [$\mu\text{g/ml}$]	Ristocetin [IU/mL]	specific activity [mU/10 μg]	
8	2.52	3.9	98.30	31.8	0.35	110.0628931	
9	2.92	4.3	98.50				
10	2.80	4.2	98.60	37.4	0.32	85.56149733	
11	2.57						
13	2.81	3.9	98.50	37.6	0.095	25.26595745	
14	2.92	3.9	97.80				
15	2.77	3.9	97.80	43.0	0.095	22.09302326	
16	2.72	4.0	98.30				
17	3.04	4.1	98.20	42.1	0.095	22.56532067	
	2.79	4.04	98.25	38.38	0.19	53.11	

TABLE 12

rVWF expression in chemostatic cell culture under low cell count, low copper conditions during weeks 2 and 3.

Day	Cell Count [10E6 cells/mL]	NH4+ [mM]	Viability [%]	vWF ELISA [µg/ml]	Ristocetin [IU/mL]	specific activity [mU/10 µg]
8	1.61	3.3	99.10	19.3	0.095	49.22279793
9	1.65	3.3	98.00			
10	1.49	3.3	98.90	18.6	0.095	51.07526882
11	1.58					
13	1.58	3.4	98.10	18.1	0.095	52.48618785
14	1.56	3.3	99.30			
15	1.52	3.3	98.50	18.9	0.095	50.26455026
16	1.50	3.3	98.40			
17	1.50	3.4	98.70	19.9	0.095	47.73869347
	1.55	3.33	98.63	18.96	0.10	50.16

TABLE 13

rVWF expression in chemostatic cell culture under low cell count, high copper conditions during weeks 2 and 3.

Day	Cell Count [10E6 cells/mL]	NH4+ [mM]	Viability [%]	vWF ELISA [µg/ml]	Ristocetin [IU/mL]	specific activity [mU/10 µg]
8	1.46	3.2	98.80	11.6	0.73	629.3103448
9	1.45	3.2	98.20			
10	1.37	3.1	98.90	11.0	0.7	636.3636364
11	1.43					
13	1.33	3.2	98.10	11.1	0.68	612.6126126
14	1.39	3.2	97.20			
15	1.51	3.2	99.70	12.4	0.69	556.4516129
16	1.49	3.2	98.60			
17	1.43	3.2	99.20	12.7	0.71	559.0551181
	1.43	3.17	98.59	11.76	0.70	598.76

As shown in FIG. 2A, supernatant harvested from continuous rVWF cell culture grown at low cell densities and high copper concentrations contained high specific activity (average of 600 mU/10 µg), while supernatants harvested from rVWF cell cultures grown at high cell densities in the presence of high or low copper concentrations and rVWF cell cultures grown at low cell density in the presence of low copper concentration contained low specific activities (less than 100 mU/10 µg). Consistent with the results observed for batch cultures, FIG. 2B shows that continuous mammalian cell cultures expressing rVWF with high specific activities have lower NH₄⁺ concentrations than do cultures producing rVWF with low specific activities. This data further strengthen the correlation between NH₄⁺ concentration in the cell culture and the specific activity of rVWF produced by the culture. Notably, the combination of high copper concentration and low ammonium concentration in the cell culture allowed for the production significantly improved rVWF activity.

Consistent with this result, agarose gel electrophoresis analysis of the rVWF multimer state of chemostat cultures (FIG. 6A-6B) revealed that only supernatants from cultures operated at high copper and low cell density and thus low ammonium resulted in a consistent expression high multimeric vWF (about 23% to 27% on CST day 8, 17 and 24 in Lanes 4, 8, and 12, respectively). All other conditions do not reach consistently high amounts of more than 10% of vWF with more than 10 dimers over an extended cultivation period.

Example 4

To determine the effect of culture media copper concentration on the expression and specific activity of rA13, mammalian cell cultures expression rA13 were grown for 4 weeks under chemostatic continuous culture conditions in ADAMTS13 medium comprising a modified DMEM/F12 basal media BESP845 and additional supplements (Table 14) containing copper concentrations ranging from 0.66 µg/L (without additional copper supplementation) and with additional copper supplementation to 4 µg/L. As shown in Table 15, increasing copper concentration in the cell culture media resulted in a significant increase in volumetric (P) and specific (q) productivity expressed as the total rA13 activity produced per liter culture and day and the total rA13 activity produced per cell and day, respectively.

TABLE 14

Composition of ADAMTS13 medium.

DMEM/F12 BESP845	mg/L
<u>Amino Acids</u>	
L-Alanine	13.3500
L-Arginine HCl	147.5000
L-Asparagine-H ₂ O	45.1600
L-Aspartic Acid	19.9500
L-Cystéine HCl—H ₂ O	32.5500
L-Cystine 2HCl	102.3500
L-Glutamic Acid	22.0500
Glycine	26.2500

TABLE 14-continued

Composition of ADAMTS13 medium.	
DMEM/F12 BESP845	mg/L
L-Histidine-H ₂ O HCl	51.4800
L-Isoleucine	74.4700
L-Leucine	119.0500
L-Lysine HCl	146.2500
L-Methionine	100.0000
L-Phenylalanine	60.4800
L-Proline	63.7400
L-Serine	36.7500
L-Threonine	53.4500
L-Tryptophan	29.0100
L-Tyrosine 2Na 2H ₂ O	75.7900
L-Valine	82.8500
salts	
Calcium Chloride (CaCl ₂)	116.6000
Copper Sulfate (CuSO ₄ —5H ₂ O)	0.0026
Ferric Nitrate (Fe(NO ₃) ₃ —9H ₂ O)	0.0500
Ferrous Sulfate (FeSO ₄ —7H ₂ O)	1.0170
Magnesium Chloride (MgCl ₂)	28.6400
Magnesium Sulfate (MgSO ₄)	48.8400
Potassium Chloride (KCl)	311.8000
Sodium Chloride (NaCl)	5495.5000
Na ₂ HPO ₄ Anhydrous	213.0200
NaH ₂ PO ₄ Anhydrous	54.3500
Zinc Sulfate Heptahydrate (ZnSO ₄ —7H ₂ O)	0.4320
Sodium selenite*anhydrous	0.0087
Vitamins	
Ascorbic Acid	3.4990
Biotin	0.2035
Choline Chloride	26.9800
D-Ca-Pantothenate	6.2400
Folic Acid	6.6500
I-Inositol	36.6000
Nicotinamide	7.0200
Pyridoxine HCl	6.0310
Riboflavin	0.6590
Thiamine HCl	6.5100
Vitamin B12	2.6800
miscellaneous	
D-Glucose	5000.0000
Linoleic Acid	0.0420
Lipoic Acid	1.0050
Putrescine 2HCl	3.6810
Thymidine	0.3650
Hypoxanthine Na	2.3900
Sodium Pyruvate	55.0000
DMEM/F12 BESP845: Total	12738.3
ADAMTS-13 Medium Supplements	
L-Glutamin	1300.0000
Pluronic F68	1000.0000
Etanolamin	1.5300
ZnSo ₄ *7 H ₂ O	1.0000
Na-Hydrogencarbonat	1500.0000
Supplements Total	3802.5
Total amount of Ingredients	16540.8

To determine if increased copper concentrations affected the integrity of the expressed rA13, supernatant harvested from rA13 cell cultures grown in media containing 0.66 µg/L, 1 µg/L, and 4 µg/L copper was examined by SDS-PAGE analysis. As seen in FIG. 3A (silver stain) and FIG. 3B (anti-A13 western blot), no obvious change in product quality by gel electrophoresis was observed. Specifically, no increase in the level of truncated 170 kD or other low MW rA13 variants and no other additional or increased HCP bands resulted from rA13 expression in the presence of increased copper concentrations.

To estimate an optimal concentration of copper on the activity of rA13, an extrapolation of data from Table 15 of P Frets versus copper concentration (FIG. 4) shows that optimum effect is likely reached at about 2 µg/L, with a conservative estimation of a negative effect above about 4 µg/L.

TABLE 15

Expression of rA13 in mammalian cell culture performed in chemostatic continuous culture mode using media containing variable copper concentrations.					
	10 L Scale R&D Experiment Mean of 4 CST weeks	ZZ-NC [10E6 cells/mL]	P Frets [U/(L*d)]	q Frets [U/(10E9 cells*d)]	Spec. Activity U/mg
0.66 µg/L Cu ²⁺		1.35	1322	1028	1759
1 µg/L Cu ²⁺		1.64	1962	1247	1690
4 µg/L Cu ²⁺		2.26	2960	1338	1768

Based on the results obtained above, an additional experiment was performed, which compared the expression of rA13 in cell culture media containing 0.66 µg/L copper with 2.0 µg/L copper. As shown in Table 16, supplementation of the basal cell media with copper, to a final concentration of 2.0 µg/L copper, resulted in a significant increase in volumetric (P) and specific (q) productivity expressed as the total rA13 activity produced per liter culture and day and the total rA13 activity produced per cell and day, respectively, over a period of 8 weeks. The specific data for each week of rA13 production in the two cultures is shown in FIG. 5A-5K. From these data, it is clear that copper supplementation has a measurable beneficial effect on cell metabolism, specific growth rate and rA13 productivity.

TABLE 16

Expression of rA13 in mammalian cell culture performed in chemostatic continuous culture mode using media containing variable copper concentrations.					
	10 L Scale Mean of 8 CST weeks	ZZ-NC [10E6 cells/mL]	P Frets [U/(L*d)]	q Frets [U/(10E9 cells*d)]	Spec. Activity U/mg
0.66 µg/L Cu ²⁺		1.29	1049	821	1862
2 µg/L Cu ²⁺		2.17	2470	1146	1749

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

What is claimed is:

1. A recombinant Von Willebrand Factor (rVWF) composition comprising isolated soluble rVWF, wherein the composition has a NH₄⁺ concentration of below 4 mM or below 10 mM, wherein the composition contains at least 0.5 IU ristocetin cofactor activity per mL, and wherein the composition is prepared by a method comprising the steps of:

- (a) providing a basal cell culture media;
- (b) supplementing the basal cell culture media with copper to provide a final copper concentration of at least 2.4 µg/L;

81

- (c) providing one or more cells comprising a nucleic acid encoding a rVWF protein;
- (d) culturing the one or more cells in the copper supplemented cell culture media such that rVWF is expressed and excreted from the cells into a culture supernatant and the NH₄⁺ content of the cell culture supernatant remains at a concentration below 4 mM or below 10 mM; and
- (e) recovering at least a portion of the culture supernatant, wherein at least 20% of the isolated soluble rVWF in the composition is present in a high molecular weight VWF multimer of more than 10 dimers.
2. The rVWF composition of claim 1, further comprising the step of supplementing the basal cell culture media with a hydrolysate prior to culturing the one or more cells.
3. The rVWF composition of claim 2, wherein the hydrolysate is a plant hydrolysate.
4. The rVWF composition of claim 3, wherein the hydrolysate is a soy hydrolysate.
5. The rVWF composition of claim 1, wherein the basal cell culture media is an animal protein free culture media.
6. The rVWF composition of claim 1, wherein the basal cell culture media is a protein free culture media.
7. The rVWF composition of claim 1, wherein the basal cell culture media is a chemically defined culture media.
8. The rVWF composition of claim 1, wherein the final copper concentration of the copper supplemented basal cell culture media is at least 4 μg/L copper.
9. The rVWF composition of claim 1, wherein the final copper concentration of the copper supplemented basal cell culture media is between 2.4 μg/L and 20 μg/L copper.
10. The rVWF composition of claim 1, wherein the copper supplementing the basal cell culture media is provided as a copper salt, a copper chelate, or a combination thereof.
11. The rVWF composition of claim 10, wherein the copper salt is selected from the group consisting of copper sulfate, copper acetate, copper carbonate, copper chloride, copper hydroxide, copper nitrate, and copper oxide.
12. The rVWF composition of claim 1, wherein the one or more cells are mammalian cells.
13. The rVWF composition of claim 12, wherein the mammalian cells are CHO cells.
14. The rVWF composition of claim 1, wherein culturing the one or more cells comprises batch cultivation of the cells.
15. The rVWF composition of claim 1, wherein culturing the one or more cells comprises continuous cultivation of the cells.
16. The rVWF composition of claim 15, wherein the continuous cultivation of cells is performed in chemostatic mode.
17. The rVWF composition of claim 15, wherein the continuous cultivation of cells is performed in perfusion mode.
18. The rVWF composition of claim 1, wherein the one or more cells is cultured in at least 100 L of the supplemented basal cell culture media.
19. The rVWF composition of claim 1, wherein the cell density is maintained at less than 2.5×10^6 cells per mL during the step of culturing the one or more cells.

82

20. The rVWF composition of claim 19, wherein the cell density is maintained at less than 2.0×10^6 cells per mL during the step of culturing the one or more cells.
21. The rVWF composition of claim 19, wherein the cell density is maintained at less than 1.5×10^6 cells per mL during the step of culturing the one or more cells.
22. The rVWF composition of claim 1, wherein the step of recovering at least a portion of the culture supernatant comprises filtration or centrifugation to remove cells from the portion of culture supernatant.
23. The rVWF composition of claim 1, wherein at least 30% of the rVWF in the supernatant is present in a high molecular weight VWF multimer of more than 10 dimers.
24. The rVWF composition of claim 1, wherein the supernatant contains high molecular weight VWF multimers of 14 to 22 dimers.
25. The rVWF composition of claim 1, wherein rVWF is co-expressed with recombinant Factor VIII (rFVIII).
26. The rVWF composition of claim 25, further comprising a step of purifying rVWF away from at least 50% of the rFVIII present in the recovered supernatant.
27. The rVWF composition of claim 26, wherein the ratio of rVWF to rFVIII after the purification step is at least 10:1.
28. The rVWF composition of claim 1, wherein the method further comprises a rVWF enrichment step.
29. The rVWF composition of claim 1, wherein the composition is formulated for pharmaceutical administration.
30. The rVWF composition of claim 1, wherein the composition is formulated for intravenous administration.
31. A rVWF composition comprising isolated soluble rVWF having a specific ristocetin cofactor activity of at least 40 mU/μg, wherein the composition has a NH₄⁺ concentration of below 4 mM or below 10 mM, and wherein the composition is prepared by a method comprising the steps of:
- providing a basal cell culture media;
 - supplementing the basal cell culture media with copper to provide a final copper concentration of at least 2.4 μg/L;
 - providing one or more cells comprising a nucleic acid encoding a rVWF protein;
 - culturing the one or more cells in the copper supplemented cell culture media such that rVWF is expressed and excreted from the cells into a culture supernatant and the NH₄⁺ content of the cell culture supernatant remains at a concentration below 4 mM or below 10 mM; and
 - recovering at least a portion of the culture supernatant, wherein at least 20% of the isolated soluble rVWF in the composition is present in a high molecular weight VWF multimer of more than 10 dimers.
32. The rVWF composition according to claim 1, wherein the isolated soluble rVWF is not plasma derived VWF (pdVWF) or ultra-large VWF in Weibel-Palade bodies or Golgi apparatus.
33. The rVWF composition according to claim 31, wherein the isolated soluble rVWF is not plasma derived VWF (pdVWF) or ultra-large VWF in Weibel-Palade bodies or Golgi apparatus.

* * * * *