

(19) **DANMARK**

(10) **DK/EP 2598633 T3**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 12 N 5/074 (2010.01)** **A 61 K 35/407 (2015.01)**
- (45) Oversættelsen bekendtgjort den: **2019-07-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-05-08**
- (86) Europæisk ansøgning nr.: **11764293.4**
- (86) Europæisk indleveringsdag: **2011-07-29**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-06-05**
- (86) International ansøgning nr.: **IB2011002167**
- (87) Internationalt publikationsnr.: **WO2012014076**
- (30) Prioritet: **2010-07-29 US 368736 P** **2010-07-29 EP 10171265**
2011-06-10 US 201161520569 P
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Koninklijke Nederlandse Akademie van Wetenschappen, Hubrecht Institute , Uppsalalaan 8, 3584 CT Utrecht, Holland**
- (72) Opfinder: **HUCH ORTEGA, Meritxell, Willem Barentszstraat 25-11, 3572 PB Utrecht, Holland**
CLEVERS, Johannes, Carolus, Ruysdaellaan 7, 3712 AP Huis ter Heide, Holland
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **LEVERORGANOID, ANVENDELSER DERAFF OG DYRKNINGSFREMGANGSMÅDE TIL OPNÅELSE DERAFF**
- (56) Fremdragne publikationer:
WO-A2-2004/087896
WO-A2-2007/127454
WO-A2-2009/022907
SNYKERS SARAH ET AL: "In Vitro Differentiation of Embryonic and Adult Stem Cells into Hepatocytes: State of the Art", STEM CELLS (MIAMISBURG), vol. 27, no. 3, 2009, pages 577-605, XP002603406, ISSN: 1066-5099
MITAKA TOSHIHIRO: "Reconstruction of hepatic organoid by hepatic stem cells", JOURNAL OF HEPATO-BILIARY-PANCREATIC SURGERY, SPRINGER VERLAG, BERLIN, DE, vol. 9, no. 6, 1 January 2002 (2002-01-01) , pages 697-703, XP002331148, ISSN: 0944-1166, DOI: DOI:10.1007/S005340200096
SNYKERS S ET AL: "Differentiation of neonatal rat epithelial cells from biliary origin into immature hepatic cells by sequential exposure to hepatogenic cytokines and growth factors reflecting liver development", TOXICOLOGY IN VITRO, ELSEVIER SCIENCE, GB, vol. 30, 4 April 2007 (2007-04-04), pages 1-7, XP002453247, ISSN: 0887-2333
LEMAIGRE ET AL: "Mechanisms of Liver Development: Concepts for Understanding Liver Disorders and Design of Novel Therapies", GASTROENTEROLOGY, ELSEVIER, PHILADELPHIA, PA, vol. 137, no. 1, 1 July 2009 (2009-07-01), pages 62-79, XP026418789, ISSN: 0016-5085 [retrieved on 2009-03-27]

Fortsættes ...

WILLIAMS JENNIFER M ET AL: "The Role of the Wnt Family of Secreted Proteins in Rat Oval "Stem" Cell-Based Liver Regeneration Wnt1 Drives Differentiation", AMERICAN JOURNAL OF PATHOLOGY, vol. 176, no. 6, June 2010 (2010-06), pages 2732-2742, XP002678410,
BARKER N ET AL: "Identification of stem cells in small intestine and colon by marker gene Lgr5", NATURE: INTERNATIONAL WEEKLY JOURNAL OF SCIENCE, NATURE PUBLISHING GROUP, UNITED KINGDOM, vol. 449, 25 October 2007 (2007-10-25), pages 1003-1008, XP002457164, ISSN: 0028-0836, DOI: 10.1038/NATURE06196

DESCRIPTION

FIELD OF THE INVENTION

[0001] The invention relates to a liver organoid, uses thereof and method for obtaining them.

BACKGROUND OF THE INVENTION

[0002] The basic architectural unit of the liver is the liver lobule. Each lobule consists of plates of hepatocytes lined by sinusoidal capillaries that radiate toward a central efferent vein. Liver lobules are roughly hexagonal with each of six corners demarcated by the presence of a portal triad (portal vein, bile duct, and hepatic artery). Although hepatocytes are the major parenchymal cell type of the liver they function in concert with cholangiocytes (biliary epithelial cells), endothelial cells, sinusoidal endothelial cells, Kupffer cells, natural killer cells and hepatic stellate cells. This complex architecture is crucial for hepatic function.

[0003] The existence of liver stem cells remains controversial. On one hand, tissue maintenance in the liver and liver regeneration upon certain types of injury, are not driven by stem cells but rather by division of the mature cells (hepatocytes or cholangiocytes). However, liver injury models in which hepatocyte proliferation has been inhibited also demonstrated the ability of the organ to regenerate in response to damage. This suggests that the liver can be considered as an organ with facultative stem cells.

[0004] So far, liver cultures derived from hepatocytes, or by differentiation of Embryonic Stem cells (ES) or induced Pluripotent Stem Cells, do not expand and self renew for longer periods.

[0005] Recently, in the small intestine, the gene *Lgr5* was identified which is specifically expressed in cycling Crypt Base Columnar (CBC) cells, which are small cells that are interspersed between the Paneth cells (Barker et al., 2007. Nature 449: 1003-1007). Using a mouse in which a GFP/tamoxifen-inducible *Cre* recombinase cassette was integrated into the *Lgr5* locus, it was shown by lineage tracing that the *Lgr5*⁺ CBC cells constitute multipotent stem cells which generate all cell types of the epithelium even when assessed 14 months after *Cre* induction.

[0006] It was recently discovered that also *Lgr6*, besides *Lgr5*, but not *Lgr4*, is a unique marker for adult stem cells. While *Lgr5* is expressed in stem cells of brain, kidney, liver, retina, stomach, intestine, pancreas, breast, hair follicle, ovary, adrenal medulla, and skin, *Lgr6* is expressed in stem cells of brain, lung, breast, hair follicle, and skin.

[0007] Here we have developed a method to culture adult liver progenitors and to obtain a liver organoid that shows longer-lived maintenance, are able to differentiate to both, hepatocyte and

cholangiocyte lineages and preserve the basic physiology of isolated liver fragments.

SUMMARY OF THE INVENTION

[0008] The invention provides a method for obtaining a liver organoid, wherein said method comprises:

culturing one or more liver epithelial stem cell, or a liver fragment or a liver biliary duct, in contact with an extracellular matrix in the presence of an expansion medium, the medium comprising a basal medium for animal or human cells to which is added: epidermal growth factor (EGF), a fibroblast growth factor (FGF) able to bind to FGFR2 or FGFR4, Nicotinamide, and a Wnt agonist selected from Rspondin 1, Rspondin 2, Rspondin 3 or Rspondin 4.

[0009] The invention also provides a cell culture medium comprising a basal medium for animal or human cells to which is added: EGF, an FGF able to bind to FGFR2 or FGFR4, Nicotinamide, and a Wnt agonist selected from R-spondin 1, R-spondin 2, R-spondin 3 and R-spondin 4.

[0010] The invention also provides a liver organoid comprising cells expressing Lgr5, obtainable by culturing one or more liver epithelial stem cell, or a liver fragment or a liver biliary duct, in contact with an extracellular matrix in the presence of an expansion medium, the medium comprising a basal medium for animal or human cells to which is added: epidermal growth factor (EGF), a fibroblast growth factor (FGF) able to bind to FGFR2 or FGFR4, Nicotinamide, a Wnt agonist selected from Rspondin 1, Rspondin 2, Rspondin 3 or Rspondin 4, and HGF, wherein the liver organoid can be cultured for at least 3 months.

[0011] The invention also provides the use of a liver organoid of the invention in a drug discovery screen.

[0012] The invention also provides the use of a liver organoid of the invention in a toxicity assay.

[0013] The invention also provides a liver organoid of the invention for use in treating a liver disorder, condition or disease or for use in regenerative medicine.

SUMMARY OF THE DISCLOSURE

[0014] The invention relates to a method for obtaining a liver organoid, wherein the method comprises culturing cells in a first "expansion" culture medium (also referred to herein as EM), preferably followed by culturing the cells in a second "differentiation" culture medium (also referred to herein as DM).

[0015] The liver organoid may be obtained by culturing a single Lgr5+ stem cell, a population of cells comprising at least one Lgr5+ stem cell, and/or a liver fragment. Herein, where "cells in/of the culture medium" are referred to, the meaning includes a single Lgr5+ stem cell, a population of cells comprising at least one Lgr5+ stem cell, and/or a liver fragment.

[0016] In one embodiment, the expansion medium comprises EGF, a Wnt agonist, FGF, and Nicotinamide. Preferably, the Wnt agonist is R-spondin 1 and so the expansion medium is referred to as "ERFNic". A particularly preferred expansion medium additionally comprises HGF and is referred to as "ERFHNic".

[0017] In one embodiment, the differentiation medium comprises EGF, a TGF-beta inhibitor, FGF and a Notch inhibitor. In one embodiment, the TGF-beta inhibitor is A83-01 and/or the Notch inhibitor is DAPT. This differentiation medium is referred to herein as "EAFD" and is a preferred differentiation medium of the invention. FGF may optionally be replaced by HGF or alternatively both FGF and HGF may be present or absent in the differentiation medium. Dexamethasone may also be added.

[0018] In a preferred embodiment, the cells may initially be cultured in an expansion medium that additionally contains Wnt and Noggin, for example an "ENRW" medium containing EGF, Noggin, R-spondin and Wnt, and optionally FGF, HGF and Nicotinamide. The inventors have found that this medium is optimum for stimulating initial expansion of cells for the first few days. Therefore, this first expansion medium is sometimes referred to herein as EM1. In some embodiments, the Wnt and Noggin are removed after approximately 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days or more, for example 2 week, 1 month, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 months or more, 14 months or more. The cells may then be expanded in an expansion medium of the invention, as described above that does not contain Wnt or Noggin. This second expansion medium is sometimes referred to herein as EM2. In some embodiments, the cells are cultured in EM2 for approximately 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days or a longer time period, such as 3, 4, 5, 10, 20 or more weeks. The culture medium may then be changed to an optimised differentiation medium, as described above, that contains a TGF-beta inhibitor and a Notch inhibitor. Typically, the differentiation medium does not contain a Wnt agonist, R-spondin or Nicotinamide. This encourages the differentiation of the cells towards mature hepatocytes and cholangiocytes. These cells are suitable for transplantation into humans or animals.

[0019] The invention also relates to a liver organoid. The present application describes the first time that liver organoids have been grown *ex vivo*.

DESCRIPTION OF THE INVENTION

[0020] In a first aspect, the invention provides a method for obtaining a liver organoid, wherein said method comprises:
culturing one or more liver epithelial stem cell, or a liver fragment or a liver biliary duct in

contact with an extracellular matrix in the presence of a medium, the medium comprising a basal medium for animal or human cells to which is added: Epidermal Growth Factor, an FGF able to bind to FGFR2 or FGFR4, preferably FGF10 as a mitogenic growth factor, Nicotinamide, and a Wnt agonist, selected from R-spondin 1, R-spondin 2, R-spondin 3 or R-spondin 4 and optionally Wnt-3a. Preferably, HGF is also added.

[0021] For example, in one embodiment, the invention provides a method for obtaining a liver organoid, wherein said method comprises:

culturing one or more liver epithelial stem cell, or a liver fragment or a liver biliary ductin contact with an extracellular matrix in the presence of a medium, the medium comprising a basal medium for animal or human cells to which is added a BMP inhibitor, a Wnt agonist selected from R-spondin 1, R-spondin 2, R-spondin 3 or R-spondin 4, Epidermal Growth Factor, a FGF able to bind to FGFR2 or FGFR4, preferably FGF10 and HGF as mitogenic growth factors, gastrin, Nicotinamide, B27, N2 and N-Acetylcysteine.

[0022] It has surprisingly been found by the present inventors that a method of the invention allows culturing of epithelial stem cells, and/or isolated fragments from the liver comprising said stem cells, while preserving the presence of stem cells that retain an undifferentiated phenotype and self-maintenance capabilities. Even more surprising was the observation that a method of the invention allows the outgrowth of a single, isolated epithelial stem cell into a liver organoid in the absence of a stem cell niche.

[0023] Stem cells are found in many organs of adult humans and mice. Although there may be great variation in the exact characteristics of adult stem cells in individual tissues, adult stem cells share at least the following characteristics: they retain an undifferentiated phenotype; their offspring can differentiate towards all lineages present in the pertinent tissue; they retain self-maintenance capabilities throughout life; and they are able to regenerate the pertinent tissue after injury. Stem cells reside in a specialized location, the stem cell niche, which supplies the appropriate cell-cell contacts and signals for maintenance of said stem cell population.

[0024] Epithelial stem cells are able to form the distinct cell types of which the epithelium is composed. Some epithelia, such as skin or intestine, show rapid cell turnover, indicating that the residing stem cells must be continuously proliferating. Other epithelia, such as the liver or pancreas, show a very slow turnover under normal conditions.

[0025] The isolation of liver epithelial stem cells may be carried out by a number of different protocols. Although the inventors do not wish to be bound by any particular theory, it is hypothesised herein that a population of stem cells exists within the liver that upon tissue injury, is responsible for liver regeneration. It is thought that the cell population responsible for this injury-responsive regeneration expresses the marker Lgr5 when activated.

[0026] By injecting Lgr5-LacZ knock-in mice with the hepatotoxic compound CC14 to simulate liver damage, the inventors have demonstrated for the first time that liver damage induces Lgr5 expression in the liver at sites of active regeneration (see example 4). The inventors have

also shown that injection of the Wnt agonist R-spondin causes upregulation of Wnt signalling expression in the liver ducts. Although the inventors do not wish to be bound by any particular theory, it is hypothesised herein that regenerative stem cells/progenitors may be activated by Wnt signalling after liver injury. The Lgr5 positive cells may then be isolated from the liver/liver fragment, if required, as described herein. Thus the disclosure describes a method for obtaining/isolating Lgr5+ cells from the liver comprising inducing liver injury or by stimulating Wnt signalling, for example with R-spondin. This is useful because in some embodiments Lgr5+ cells are the starting point for obtaining liver organoids *in vitro* (although the use of a culture medium of the invention allows generation of a Lgr5+ cell by adding a Wnt agonist such as R-spondin and so it is not essential to obtain an Lgr5+ cell from the liver in order to practice the present invention). Lgr5+ cells obtained by this method are also provided. In some embodiments, such cells preferably do not express markers of stellate cells, for example SMA. Instead, such cells preferably express one or more liver progenitor markers or stem cell markers such as Sox9. Preferably, the expression of one or more liver progenitor markers or stem cell markers such as Sox9 is upregulated in said cells compared to adult liver.

[0027] The disclosure also describes a method for regenerating the liver comprising stimulating Wnt signalling. Such a method may be useful in the treatment of liver disorders. The induction of Lgr5 expression in liver cells by injury or by stimulating Wnt signalling may be carried out *in vivo*, *ex vivo* in an isolated liver, or *in vitro* in a liver fragment or population of liver cells. In embodiments in which the induction of Lgr5 expression is carried out *ex vivo* or *in vitro*, the Lgr5 positive cells may be administered to a patient in need thereof by any suitable means, for example, by injection or by implantation. In some embodiments, implanted cells are contained in biocompatible matrix. In some embodiments, the cells are expanded *ex vivo* prior to being used in therapy.

[0028] In a preferred method of the invention, said epithelial stem cells are isolated from a liver fragment or a liver biliary duct, more preferably from biliary duct tissue.

[0029] Methods for the isolation of bile duct tissue are known to those of skill in the art. For example, biliary duct may be isolated from a liver as described in the examples enclosed herein. Briefly, an adult liver tissue may be washed in a cold (4-10°C) culture medium, preferably Advanced-DMEM/F12 (Invitrogen) and then, the tissue can be chopped into pieces of around 5mm and further washed with cold dissociation buffer (collagenase, dispase, FBS in DMEM media). The tissue fragments are preferably incubated with the dissociation buffer for about 2 h at about 37°C. Then, the tissue fragments can be vigorously suspended in 10 ml of cold (4-10°C) isolation buffer with a 10 ml pipette. The first supernatant containing dead cells is preferably discarded and the sediment preferably suspended with dissociation buffer (e.g. 10-15 ml). After further vigorous suspension of the tissue fragments the supernatant is enriched in biliary ducts. A suspension containing biliary ducts can in this way be obtained and biliary ducts are collected under the microscope and retained in cold media (DMEM + 5-10% FBS). This procedure may be repeated until at least 10-20 biliary ducts/well are collected. Then, the isolated biliary ducts may be precipitated. Isolated bile ducts are preferably seeded in 50ul of matrigel at an approximate ratio of 20 biliary ducts/ well.

[0030] In contrast to mature hepatocytes, which grow to confluence for a short period of time, before dying, liver epithelial stem cells isolated according to the disclosure are self-renewing and grow indefinitely. It has been found that the self-renewing population of cells are those which are capable of expressing Lgr5 on their surface. Lgr5 negative cells do not self-renew. The term "self-renewing" should be understood to represent the capacity of a cell to reproduce itself whilst maintaining the original proliferation and differentiation properties of cells of the invention. Such cells proliferate by dividing to form clones, which further divide into clones and therefore expand the size of the cell population without the need for external intervention, without evolving into cells with a more restricted differentiation potential.

[0031] A preferred method is based on the fact that liver stem cells according to the invention express Lgr5 and/or Lgr6 on their surface; these proteins belong to the large G protein-coupled receptor (GPCR) superfamily (see, for example, WO2009/022907, the contents of which are incorporated herein in their entirety). The Lgr subfamily is unique in carrying a large leucine-rich ectodomain important for ligand binding. A preferred method therefore comprises preparing a cell suspension from said epithelial tissue as described above, contacting said cell suspension with an Lgr5 and/or 6 binding compound (such as an antibody), isolating the Lgr5 and/or 6 binding compound, and isolating the stem cells from said binding compound.

[0032] It is preferred that a single cell suspension comprising the epithelial stem cells is mechanically generated from the isolated biliary duct. Small organoid fragments generated in this way by mechanical disruption are preferably split at a ratio of approximately 1:6. If necessary, such fragments can be incubated for a short time (only 2 or 3 minutes) in trypsin at a dilution of approximately 1:2. It has been found that at this stage epithelial stem cells treated with trypsin yielded rather low survival rates: if the cells are split into individual cells, then only those expressing Lgr5 survive. This fraction is rather small (approximately 12% of the total cell population).

[0033] Preferred Lgr5 and/or 6 binding compounds comprise antibodies, such as monoclonal antibodies that specifically recognize and bind to the extracellular domain of either Lgr5 or Lgr6, such as monoclonal antibodies including mouse and rat monoclonal antibodies (see, for example, WO2010/016766, the contents of which are incorporated herein in their entirety). Using such an antibody, Lgr5 and/or Lgr6-expressing stem cells can be isolated, for example with the aid of magnetic beads or through fluorescence-activated cell sorting, as is clear to a skilled person. Using a method of the disclosure, it is possible to isolate one single Lgr5 and/or Lgr6 expressing cell and to apply a method of the invention to it. A liver organoid may therefore be derived from one single cell.

Stem Cell Niche

[0034] Isolated stem cells are preferably cultured in a microenvironment that mimics at least in part a cellular niche in which said stem cells naturally reside. This cellular niche may be

mimicked by culturing said stem cells in the presence of biomaterials, such as matrices, scaffolds, and culture substrates that represent key regulatory signals controlling stem cell fate. Such biomaterials comprise natural, semi-synthetic and synthetic biomaterials, and/or mixtures thereof. A scaffold provides a two-dimensional or three dimensional network. Suitable synthetic materials for such a scaffold comprise polymers selected from porous solids, nanofibers, and hydrogels such as, for example, peptides including self-assembling peptides, hydrogels composed of polyethylene glycol phosphate, polyethylene glycol fumarate, polyacrylamide, polyhydroxyethyl methacrylate, polycellulose acetate, and/or copolymers thereof (see, for example, Saha et al., 2007. *Curr Opin Chem Biol.* 11(4): 381-387; Saha et al., 2008. *Biophysical Journal* 95: 4426-4438; Little et al., 2008. *Chem. Rev* 108, 1787-1796). As is known to a skilled person, the mechanical properties such as, for example, the elasticity of the scaffold influences proliferation, differentiation and migration of stem cells. A preferred scaffold comprises biodegradable (co)polymers that are replaced by natural occurring components after transplantation in a subject, for example to promote tissue regeneration and/or wound healing. It is furthermore preferred that said scaffold does not substantially induce an immunogenic response after transplantation in a subject. Said scaffold is supplemented with natural, semi-synthetic or synthetic ligands, which provide the signals that are required for proliferation and/or differentiation, and/or migration of stem cells. In a preferred embodiment, said ligands comprise defined amino acid fragments. Examples of said synthetic polymers comprise Pluronic® F127 block copolymer surfactant (BASF), and Ethisorb® (Johnson and Johnson).

[0035] A cellular niche is in part determined by the stem cells and surrounding cells, and the extracellular matrix (ECM) that is produced by the cells in said niche. In a preferred method of the invention, isolated liver fragments or isolated biliary duct or epithelial stem cells are attached to an ECM. ECM is composed of a variety of polysaccharides, water, elastin, and glycoproteins, wherein the glycoproteins comprise collagen, entactin (nidogen), fibronectin, and laminin. ECM is secreted by connective tissue cells. Different types of ECM are known, comprising different compositions including different types of glycoproteins and/or different combination of glycoproteins. Said ECM can be provided by culturing ECM-producing cells, such as for example fibroblast cells, in a receptacle, prior to the removal of these cells and the addition of isolated liver fragments or isolated biliary duct or isolated epithelial stem cells. Examples of extracellular matrix-producing cells are chondrocytes, producing mainly collagen and proteoglycans, fibroblast cells, producing mainly type IV collagen, laminin, interstitial procollagens, and fibronectin, and colonic myofibroblasts producing mainly collagens (type I, III, and V), chondroitin sulfate proteoglycan, hyaluronic acid, fibronectin, and tenascin-C. Alternatively, said ECM is commercially provided. Examples of commercially available extracellular matrices are extracellular matrix proteins (Invitrogen) and basement membrane preparations from Engelbreth-Holm-Swarm (EHS) mouse sarcoma cells (e.g. Matrigel™ (BD Biosciences)). A synthetic extracellular matrix material, such as ProNectin (Sigma Z378666) may be used. Mixtures of extracellular matrix materials may be used, if desired. The use of an ECM for culturing stem cells enhanced long-term survival of the stem cells and the continued presence of undifferentiated stem cells. In the absence of an ECM, stem cell cultures could not be cultured for longer periods and no continued presence of undifferentiated stem cells was

observed. In addition, the presence of an ECM allowed culturing of three-dimensional tissue organoids, which could not be cultured in the absence of an ECM. The extracellular matrix material will normally be coated onto a cell culture vessel, but may (in addition or alternatively) be supplied in solution. A fibronectin solution of about 1mg/ml may be used to coat a cell culture vessel, or between about 1 $\mu\text{g}/\text{cm}^2$ to about 250 $\mu\text{g}/\text{cm}^2$, or at about 1 $\mu\text{g}/\text{cm}^2$ to about 150 $\mu\text{g}/\text{cm}^2$. In some embodiments, a cell culture vessel is coated with fibronectin at between 8 $\mu\text{g}/\text{cm}^2$ and 125 $\mu\text{g}/\text{cm}^2$.

[0036] A preferred ECM for use in a method of the invention comprises at least two distinct glycoproteins, such as two different types of collagen or a collagen and laminin. The ECM can be a synthetic hydrogel extracellular matrix or a naturally occurring ECM. A further preferred ECM is provided by Matrigel™ (BD Biosciences), which comprises laminin, entactin, and collagen IV.

[0037] The compositions of the invention may comprise serum, or may be serum-free and/or serum-replacement free, as described elsewhere herein. Culture media and cell preparations are preferably GMP processes in line with standards required by the FDA for biologics products and to ensure product consistency.

Culture Media

[0038] A cell culture medium that is used in a method of the invention comprises any suitable cell culture medium, subject to the limitations provided herein. Cell culture media typically contain a large number of ingredients, which are necessary to support maintenance of the cultured cells. Suitable combinations of ingredients can readily be formulated by the skilled person, taking into account the following disclosure. A culture medium according to the invention will generally be a nutrient solution comprising standard cell culture ingredients, such as amino acids, vitamins, inorganic salts, a carbon energy source, and a buffer, as described in more detail in the literature and below.

[0039] A culture medium of the invention will normally be formulated in deionized, distilled water. A culture medium of the invention will typically be sterilized prior to use to prevent contamination, *e.g.* by ultraviolet light, heating, irradiation or filtration. The culture medium may be frozen (*e.g.* at -20°C or -80°C) for storage or transport. The medium may contain one or more antibiotics to prevent contamination. The medium may have an endotoxin content of less than 0.1 endotoxin units per ml, or may have an endotoxin content less than 0.05 endotoxin units per ml. Methods for determining the endotoxin content of culture media are known in the art.

[0040] A preferred cell culture medium is a defined synthetic medium that is buffered at a pH of 7.4 (preferably with a pH 7.2 - 7.6 or at least 7.2 and not higher than 7.6) with a carbonate-based buffer, while the cells are cultured in an atmosphere comprising between 5 % and 10%

CO₂, or at least 5% and not more than 10% CO₂, preferably 5 % CO₂.

[0041] The skilled person will understand from common general knowledge the types of culture media that might be used for as the basal medium in the cell culture mediums of the invention. Potentially suitable cell culture media are available commercially, and include, but are not limited to, Dulbecco's Modified Eagle Media (DMEM), Minimal Essential Medium (MEM), Knockout-DMEM (KO-DMEM), Glasgow Minimal Essential Medium (G-MEM), Basal Medium Eagle (BME), DMEM/Ham's F12, Advanced DMEM/Ham's F12, Iscove's Modified Dulbecco's Media and Minimal Essential Media (MEM), Ham's F-10, Ham's F-12, Medium 199, and RPMI 1640 Media.

[0042] A preferred cell culture medium is selected from DMEM/F12 and RPMI 1640 supplemented with glutamine, insulin, Penicillin/streptomycin and transferrin. In a further preferred embodiment, Advanced DMEM/F12 or Advanced RPMI is used, which is optimized for serum free culture and already includes insulin. In this case, said Advanced DMEM/F12 or Advanced RPMI medium is preferably supplemented with glutamine and Penicillin/streptomycin.

[0043] In some embodiments, the basal medium comprises Advanced DMEM F12, hepes, penicillin/streptomycin, Glutamin, NAcetyl Cystein, B27, N2 and Gastrin. In some embodiments, culture is initiated with a basal medium comprising N2 and Gastrin and penicillin/streptomycin but these are later withdrawn. For example, in some embodiments, N2 and Gastrin and penicillin/streptomycin are present in an EM1 medium of the invention but not in an EM2 or DM. For example, in some embodiments, N2 and Gastrin and penicillin/streptomycin are present in an EM1 and EM2 medium of the invention but not in a DM. In particularly preferred embodiments, the basal medium is Advanced DMEM/F12 or a DMEM variant supplemented with penicillin/streptomycin, N2, B27, glutamine and gastrin.

[0044] In preferred embodiments, the basal medium comprises Gastrin. In some embodiments, the basal medium also comprises NAc and/or B27.

[0045] It is furthermore preferred that said cell culture medium is supplemented with a purified, natural, semi-synthetic and/or synthetic growth factor and does not comprise an undefined component such as fetal bovine serum or fetal calf serum. Various different serum replacement formulations are commercially available and are known to the skilled person. Where a serum replacement is used, it may be used at between about 1% and about 30% by volume of the medium, according to conventional techniques.

Expansion Medium (EM2):

[0046] In one aspect of the present invention there is provided a cell culture medium which comprises or consists of a basal medium for animal or human cells to which is added: Epidermal Growth Factor, an FGF able to bind to FGFR2 or FGFR4, preferably FGF10 as a

mitogenic growth factor, Nicotinamide, and a Wnt agonist, selected from R-spondin 1, R-spondin 2, R-spondin 3 and R-spondin 4, preferably R-spondin 1. This medium is referred to as EM2.

[0047] Preferably, the Wnt agonist is R-spondin 1. A medium comprising EGF, R-spondin 1, FGF and Nicotinamide is referred to herein as ERFNic.

[0048] In some embodiments, the EM2 medium does not comprise noggin, and more preferably does not comprise a BMP-inhibitor. In some embodiments, the EM2 medium does not comprise Wnt, for example Wnt-3a.

[0049] In some embodiments, HGF is present in addition to FGF. A preferred medium comprising HGF in addition to FGF is ERFHNic (EGF + R-spondin (preferably R-spondin1) + FGF (preferably FGF10) + HGF + Nicotinamide). The inventors have found that the ERFHNic medium is the optimal medium for long-term expansion of cells. In the absence of HGF, cells did not remain viable in culture for longer than three months. Further, in the absence of HGF, after 10 passages, cells showed a growth disadvantage compared to cells cultured in the presence of HGF as evidenced by a lower proliferation ratio. In particular, after 15 passages, the cells were not growing organoids at the same speed ratio in the absence of HGF as in the presence of HGF. Thus, HGF was found to be essential for maintaining a good proliferation rate during long-term culture. Thus the invention provides the use of an ERFHNic medium of the invention for culturing cells for at least 2 weeks, at least 1 month, at least 2 months, more preferably at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 20, at least 24, at least 25, at least 30 or more months, for example 3 or more years. In practice, some embodiments of the invention comprise the use of E2 for around 20-30 passages of the cells. For example, the cells may be split 20-30 times, generally once a week. Preferably the cells will expand at a rate of about 4 fold per week or two population doublings a week. The invention further provides the use of an ERFHNic medium of the invention for culturing cells for at least 10 passages, for example, at least 11, at least 12, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60 passages or for between 15-35 passages, for example approximately 20-30 passages. In some embodiments, a TGF-beta inhibitor such as A83-01 is additionally present in the EM2 medium. This is particularly useful when human cells or organoids are being cultured. In some embodiments, the A83-01 is present at a concentration of between 400-600 nM, for example 450-550 nM, 470-530 nM or approximately 500 nM. In embodiments in which a TGF-beta inhibitor is present in EM2, a Notch inhibitor is preferably not present.

Expansion Medium (EM1):

[0050] In one aspect, the disclosure describes a cell culture medium comprising or consisting of a basal medium for animal or human cells to which is added EGF, a BMP inhibitor, R-spondin and Wnt. Preferably, the BMP inhibitor is Noggin and the EM1 medium is termed "ENRW" (EGF, Noggin, R-spondin and Wnt). This medium is referred to as EM1. The inventors

have found that a medium containing Wnt and Noggin is ideal for stimulating initial expansion of cells. Thus, in some embodiments, the EM1 medium is used for just 1 passage or 1 week but it is also envisaged that EM1 medium can be used for around a year because it is not harmful for the cells. Thus, in some embodiments, an EM1 medium is used for culturing cells from day 0 to day 10, for example from days 0-7, days 0-6, days 0-5, days 0-4, days 0-3, days 0-2, days 0-1, wherein day 0 is the day that the cells are isolated from their tissue of origin and day 1 is the subsequent day or is used for 1 or more weeks for example 2, 3, 4 or more weeks or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or more months. In some embodiments, the EM1 medium is used only for the first day or first two days of culture. In some embodiments, EM1 medium is used for 1 or more passages, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30 or more passages, for example, 20-30 passages. In some embodiments, the EM1 medium is used subsequent to a freezing step or any other transportation step involving a medium or temperature change that does not combine with optimal growth.

[0051] The EM1 medium is supplemented with one or more of the compounds selected from the group consisting of FGF, HGF, Nicotinamide, gastrin, B27, N-acetylcystein and N2. In the case of starting the cultures from a frozen stock or from a single cell, the EM1 media is preferably supplemented with a ROCK inhibitor. Y27632 is the preferred ROCK inhibitor for use in the invention.

[0052] Thus, in one embodiment there is provided a cell culture medium which comprises or consists of a basal medium for animal or human cells to which is added: Epidermal Growth Factor, an FGF able to bind to FGFR2 or FGFR4, preferably FGF10 and HGF as mitogenic growth factors,

gastrin, Nicotinamide, B27, N2 and N-Acetylcysteine, and preferably;

a BMP inhibitor, preferably Noggin; and

a Wnt agonist selected from R-spondin 1, R-spondin 2, R-spondin 3 and R-spondin 4, preferably R-spondin 1, and optionally Wnt-3a.

[0053] B27 (Invitrogen), N-Acetylcysteine (Sigma) and N2 (Invitrogen), Gastrin (Sigma) and Nicotinamide (Sigma) are also added to the medium defined above and are believed to stimulate proliferation of the cells. In the context of the invention, Nicotinamide is also referred to herein as "Nic".

[0054] 'N2 Supplement' is available from Invitrogen, Carlsbad, CA; www.invitrogen.com; catalog no. 17502-048; and from PAA Laboratories GmbH, Pasching, Austria; www.paa.com; catalog no. F005-004; Bottenstein & Sato, PNAS, 76(1):514-517, 1979. N2 Supplement is supplied by PAA Laboratories GmbH as a 100x liquid concentrate, containing 500µg/ml human transferrin, 500µg/ml bovine insulin, 0.63µg/ml progesterone, 1611µg/ml putrescine, and 0.52µg/ml sodium selenite. N2 Supplement may be added to a culture medium as a concentrate or diluted before addition to a culture medium. It may be used at a 1x final concentration or at other final concentrations. Use of N2 Supplement is a convenient way to incorporate transferrin, insulin, progesterone, putrescine and sodium selenite into a culture medium of the invention.

[0055] 'B27 Supplement' (available from Invitrogen, Carlsbad, CA; www.invitrogen.com; currently catalog no. 17504-044; and from PAA Laboratories GmbH, Pasching, Austria; www.paa.com; catalog no. F01-002; Brewer et al., J Neurosci Res., 35(5):567-76, 1993) may be used to formulate a culture medium that comprises biotin, cholesterol, linoleic acid, linolenic acid, progesterone, putrescine, retinol, retinyl acetate, sodium selenite, tri-iodothyronine (T3), DL-alpha tocopherol (vitamin E), albumin, insulin and transferrin. B27 Supplement is supplied by PAA Laboratories GmbH as a liquid 50x concentrate, containing amongst other ingredients biotin, cholesterol, linoleic acid, linolenic acid, progesterone, putrescine, retinol, retinyl acetate, sodium selenite, tri-iodothyronine (T3), DL-alpha tocopherol (vitamin E), albumin, insulin and transferrin. Of these ingredients at least linolenic acid, retinol, retinyl acetate and tri-iodothyronine (T3) are nuclear hormone receptor agonists. B27 Supplement may be added to a culture medium as a concentrate or diluted before addition to a culture medium. It may be used at a 1x final concentration or at other final concentrations. Use of B27 Supplement is a convenient way to incorporate biotin, cholesterol, linoleic acid, linolenic acid, progesterone, putrescine, retinol, retinyl acetate, sodium selenite, tri-iodothyronine (T3), DL-alpha tocopherol (vitamin E), albumin, insulin and transferrin into a culture medium of the invention.

BMP Inhibitors

[0056] A component that is preferably added to the basal culture medium is a BMP inhibitor. BMPs bind as a dimeric ligand to a receptor complex consisting of two different receptor serine/threonine kinases, type I and type II receptors. The type II receptor phosphorylates the type I receptor, resulting in the activation of this receptor kinase. The type I receptor subsequently phosphorylates specific receptor substrates (SMAD), resulting in a signal transduction pathway leading to transcriptional activity.

[0057] A BMP inhibitor is defined as an agent that binds to a BMP molecule to form a complex wherein the BMP activity is neutralized, for example by preventing or inhibiting the binding of the BMP molecule to a BMP receptor. Alternatively, said inhibitor is an agent that acts as an antagonist or reverse agonist. This type of inhibitor binds with a BMP receptor and prevents binding of a BMP to said receptor. An example of a latter agent is an antibody that binds a BMP receptor and prevents binding of BMP to the antibody-bound receptor.

[0058] A BMP inhibitor may be added to the media in an amount effective to inhibit a BMP-dependent activity in a cell to at most 90%, more preferred at most 80%, more preferred at most 70%, more preferred at most 50%, more preferred at most 30%, more preferred at most 10%, more preferred 0%, relative to a level of a BMP activity in the absence of said inhibitor, as assessed in the same cell type. As is known to a skilled person, a BMP activity can be determined by measuring the transcriptional activity of BMP, for example as exemplified in Zilberberg et al., 2007. BMC Cell Biol. 8:41.

[0059] Several classes of natural BMP-binding proteins are known, including Noggin

(Peprotech), Chordin and chordin-like proteins (R&D systems) comprising chordin domains, Follistatin and follistatin-related proteins (R&D systems) comprising a follistatin domain, DAN and DAN-like proteins (R&D systems) comprising a DAN cysteine-knot domain, sclerostin /SOST (R&D systems), decorin (R&D systems), and alpha-2 macroglobulin (R&D systems).

[0060] A preferred BMP inhibitor for use in a method of the invention is selected from Noggin, DAN, and DAN-like proteins including Cerberus and Gremlin (R&D systems). These diffusible proteins are able to bind a BMP ligand with varying degrees of affinity and inhibit their access to signalling receptors. The addition of any of these BMP inhibitors to the basal culture medium prevents the loss of stem cells.

[0061] A preferred BMP inhibitor is Noggin. In the context of a culture medium of the invention, Noggin is also referred to herein as "N". Noggin is preferably added to the basal culture medium at a concentration of at least 10 ng/ml, more preferred at least 20 ng/ml, more preferred at least 50 ng/ml, more preferred at least 100 ng/ml. A still more preferred concentration is approximately 100 ng/ml or exactly 100 ng/ml. During culturing of stem cells, said BMP inhibitor may be added to the culture medium when required, for example, daily or every other day. The BMP inhibitor is preferably added to the culture medium every second day. The culture medium may be refreshed when required, for example, daily or every other day, although is preferably refreshed every fourth day.

Wnt Agonists

[0062] A further component that is added to the basal culture medium is a Wnt agonist. In the context of a culture medium of the invention, the Wnt agonist is also referred to herein as "W". The Wnt signalling pathway is defined by a series of events that occur when a Wnt protein binds to a cell-surface receptor of a Frizzled receptor family member. This results in the activation of Dishevelled family proteins which inhibit a complex of proteins that includes axin, GSK-3, and the protein APC to degrade intracellular β -catenin. The resulting enriched nuclear β -catenin enhances transcription by TCF/LEF family transcription factors.

[0063] A Wnt agonist is defined as an agent that activates TCF/LEF-mediated transcription in a cell. Wnt agonists are therefore selected from true Wnt agonists that bind and activate a Frizzled receptor family member including any and all of the Wnt family proteins, an inhibitor of intracellular β -catenin degradation, and activators of TCF/LEF. Said Wnt agonist is added to the media in an amount effective to stimulate a Wnt activity in a cell by at least 10%, more preferred at least 20%, more preferred at least 30%, more preferred at least 50%, more preferred at least 70%, more preferred at least 90%, more preferred at least 100%, relative to a level of said Wnt activity in the absence of said molecule, as assessed in the same cell type. As is known to a skilled person, a Wnt activity can be determined by measuring the transcriptional activity of Wnt, for example by pTOPFLASH and pFOPFLASH Tcf luciferase reporter constructs (Korinek et al., 1997. Science 275:1784-1787).

[0064] A Wnt agonist may comprise a secreted glycoprotein including Wnt-1/Int-1; Wnt-2/Irp (Int-1-related Protein); Wnt-2b/13; Wnt-3/Int-4; Wnt-3a (R&D systems); Wnt-4; Wnt-5a; Wnt-5b; Wnt-6 (Kirikoshi H et al. 2001. *Biochem Biophys Res Com* 283: 798-805); Wnt-7a (R&D systems); Wnt-7b; Wnt-8a/8d; Wnt-8b; Wnt-9a/14; Wnt-9b/14b/15; Wnt-10a; Wnt-10b/12; Wnt-11; and Wnt-16. An overview of human Wnt proteins is provided in "THE WNT FAMILY OF SECRETED PROTEINS", R&D Systems Catalog, 2004.

[0065] Further Wnt agonists include the R-spondin family of secreted proteins, which is implicated in the activation and regulation of Wnt signaling pathway and which is comprised of 4 members (R-spondin 1 (NU206, Nuvelo, San Carlos, CA), R-spondin 2 ((R&D systems), R-spondin 3, and R-spondin-4); and Norrin (also called Norrie Disease Protein or NDP) (R&D systems), which is a secreted regulatory protein that functions like a Wnt protein in that it binds with high affinity to the Frizzled-4 receptor and induces activation of the Wnt signaling pathway (Kestutis Planutis et al. (2007) *BMC Cell Biol.* 8: 12).

[0066] Compounds that mimic the activity of R-spondin may be used as Wnt agonists of the invention. It has recently been found that R-spondin interacts with Lgr5. Thus, Lgr5 agonists such as agonistic anti-Lgr5 antibodies are examples of Wnt agonists that may be used in the invention.

[0067] In the context of a culture medium of the invention, R-spondin is also referred to herein as "R".

[0068] A small-molecule agonist of the Wnt signaling pathway, an aminopyrimidine derivative, was recently identified and is also expressly included as a Wnt agonist (Liu et al. (2005) *Angew Chem Int Ed Engl.* 44, 1987-90).

[0069] Known GSK-inhibitors comprise small-interfering RNAs (siRNA; Cell Signaling), lithium (Sigma), kenpaullone (Biomol International; Leost, M. et al. (2000) *Eur. J. Biochem.* 267, 5983-5994), 6-Bromoindirubin-30-acetoxime (Meijer, L. et al. (2003) *Chem. Biol.* 10, 1255-1266), SB 216763 and SB 415286 (Sigma-Aldrich), and FRAT-family members and FRAT-derived peptides that prevent interaction of GSK-3 with axin. An overview is provided by Meijer et al., (2004) *Trends in Pharmacological Sciences* 25, 471-480. Methods and assays for determining a level of GSK-3 inhibition are known to a skilled person and comprise, for example, the methods and assay as described in Liao et al 2004, *Endocrinology*, 145(6): 2941-9).

[0070] In a preferred embodiment, said Wnt agonist is selected from one or more of a R-spondin 1-4 (for example, R-spondin 1 or R-spondin 4), and optionally further comprises a Wnt family member, Norrin, and/or a GSK-inhibitor. It was found by the inventors that the addition of at least one Wnt agonists to the basal culture medium is important for proliferation of the liver epithelial stem cells or isolated biliary duct or isolated liver fragments.

[0071] In a further preferred embodiment, said Wnt agonist comprises or consists of R-spondin 1. R-spondin 1 is preferably added to the basal culture medium at a concentration of at least

200 ng/ml, more preferred at least 300 ng/ml, more preferred at least 500 ng/ml. A still more preferred concentration of R-spondin 1 is approximately 500 ng/ml or exactly 500 ng/ml. During culturing of stem cells, said Wnt family member may be added to the culture medium when required, for example, daily or every other day. The Wnt family member is preferably added to the culture medium every second day. The culture medium may be refreshed when required, for example, daily or every other day, although is preferably refreshed every fourth day.

[0072] In a preferred embodiment, a Wnt agonist is selected from the group consisting of: R-spondin, Wnt-3a and Wnt-6. More preferably, R-spondin and Wnt-3a are both used as a Wnt agonist. This combination is particularly preferred since this combination surprisingly has a synergistic effect on organoid formation. Preferred concentrations are 1-500 ng/ml, for example, 1-10 ng/ml, 10-100 ng/ml, 1-50 ng/ml, 10-200 ng/ml, 200 to 500 ng/ml, 30 ng/ml for R-spondin and 100 ng/ml to 1000 ng/ml, for example, 100 ng/ml or 1000 ng/ml for Wnt3a.

[0073] A Wnt agonist is preferably a Wnt ligand, such as for example Wnt3a, and may be freshly added to a culture medium. Alternatively, a Wnt ligand is expressed in a cell line by transfecting or infected a cell line with a suitable expression construct expressing said Wnt ligand. Said cell line is cultured and the culture medium comprising the secreted Wnt ligand is harvested at suitable time intervals. For example, cells will produce Wnt3a as soon as they reach confluency and stop growing. Culture medium from cells that were not transfected or infected with said expression construct is used as a control. The conditioned medium is harvested and tested, for example in an assay wherein luciferase expression is controlled by TCF responsive elements to test for the presence of a Wnt agonist such as Wnt3a (Korinek et al., 1997. Science 275:1784-1787). The medium is diluted when used in the cultures to regenerate tissue. As is known to the skilled person, the addition of an excess of Wnt ligand sometimes is as detrimental for the culture as is the addition of too little Wnt ligand. Therefore, the actual dilution of the conditioned medium will depend on the amount of Wnt ligand that is determined in the test.

Mitogenic Growth Factors

[0074] Yet a further component that is added to the basal culture medium is a combination of mitogenic growth factors, including epidermal growth factor (EGF; (preferably from Peprotech), a fibroblast growth factor (FGF) able to bind to FGFR2 or FGFR4, and optionally hepatocyte growth factor (HGF) (also preferably from Peprotech). An FGF able to bind to FGFR2 (FGF receptor) or FGFR4 is preferably FGF4, FGF7 or FGF10 (preferably from Peprotech), most preferably FGF10. Preferably all 3 of EGF, an FGF and HGF are used. In the context of a culture medium of the invention, EGF is also referred to herein as "E", FGF is also referred to herein as "F" and HGF is also referred to herein as "H". EGF is a potent mitogenic factor for a variety of cultured ectodermal and mesodermal cells and has a profound effect on the differentiation of specific cells *in vivo* and *in vitro* and of some fibroblasts in cell culture. The EGF precursor exists as a membrane-bound molecule which is proteolytically cleaved to

generate the 53-amino acid peptide hormone that stimulates cells. EGF exerts its effects in the target cells by binding to the plasma membrane located EGF receptor. The EGF receptor is a transmembrane protein tyrosine kinase. Binding of EGF to the receptor causes activation of the kinase and subsequently receptor autophosphorylation. The autophosphorylation is essential for the interaction of the receptor with its substrates. The signal transduction pathways activated by EGF include the phosphatidylinositol pathway, leading to activation of protein kinase C and to increase in the intracellular Ca^{2+} concentration, and to the ras pathway leading to MAP kinase activation. These pathways are involved in regulating cellular proliferation, differentiation, and survival (Herbst, 2004, Int Journal of radiation oncology, 59, 2, S21-S26).

[0075] EGF is preferably added to the basal culture medium at a concentration of between 5 and 500 ng/ml or of at least 5 and not higher than 500 ng/ml. A preferred concentration is at least 10, 20, 25, 30, 40, 45, or 50 ng/ml and not higher than 500, 450, 400, 350, 300, 250, 200, 150, or 100 ng/ml. A more preferred concentration is at least 50 and not higher than 100 ng/ml. An even more preferred concentration is about 50 ng/ml or 50 ng/ml.

[0076] FGF10 is a protein that belongs to the fibroblast growth factor (FGF) family of proteins. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGFs stimulate cells by interacting with cell surface tyrosine kinase receptors (FGFR). Four closely related receptors (FGFR1-FGFR4) have been identified. FGFR1-FGFR3 genes have been shown to encode multiple isoforms, and these isoforms can be critical in determining ligand specificity. Most FGFs bind more than one receptor (Ornitz J Biol Chem. 1998 Feb 27;273 (9):5349-57). However, FGF10 and FGF7 are unique among FGFs in that they interact only with a specific isoform of FGFR2, designated FGFR2b which is expressed exclusively by epithelial cells (Igarashi, J Biol Chem. 1998 273(21):13230-5). FGF10 is a preferred FGF able to bind to FGFR2 or FGFR4.

[0077] Hepatocyte growth factor/scatter factor (HGF/SF) is a morphogenic factor that regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor.

[0078] The same concentrations may be used for FGF10 and HGF. Preferred concentrations for FGF10 are 20, 50, 100, 500 ng/ml, not higher than 500ng/ml. Preferred concentrations for HGF are 1, 10, 20, 50 ng/ml, not higher than 50ng/ml. During culturing of stem cells, said combination of mitogenic growth factors (i.e. EGF, FGF10 and HGF) is preferably added to the culture medium when required, for example, daily or every other day. It is preferable that it is added every second day. The culture medium may be refreshed when required, for example, daily or every other day, although is preferably refreshed every fourth day.

[0079] In some embodiments, a TGF-beta inhibitor such as A83-01 is additionally present in the EM1 medium. This is particularly useful when human cells or organoids are being cultured. In some embodiments, the A83-01 is present at a concentration of between 400-600 nM, for

example 450-550 nM, 470-530 nM or approximately 500 nM. In embodiments in which a TGF-beta inhibitor is present in EM1, a Notch inhibitor is preferably not present.

Preferred Expansion Media of the Invention

[0080] Preferred culture media and methods of using these are described in the Examples.

[0081] For example, an cell culture medium may comprise or consist of a basal medium to which is added: EGF and R-spondin 1 supplemented with FGF10, HGF and Nicotinamide; for example, EGF (50 ng/ml) and R-spondin 1 (1ug/ml) supplemented with FGF10 (100ng/ml), HGF (25-50ng/ml) and Nicotinamide (1-10mM). The inventors have found that this medium is preferable for long-term expansion of cells. Thus, this cell culture medium is preferred for use as an EM2 of the invention. The basal medium is preferably supplemented with B27, N2 and 200ng/ml N-Acetylcysteine. In some embodiments, the basal medium is Advanced-DMEM/F12. However, any other suitable basal medium may be used.

[0082] Another preferred cell culture medium, and method of using this medium, is described in the examples, and comprises or consists of Advanced-DMEM/F12 preferably supplemented with B27, N2, 200ng/ml N-Acetylcysteine, 50ng/ml EGF, 1 µg/ml R-spondin1, 10 nM gastrin, 100 ng/ml FGF10, 10mM Nicotinamide and 50ng/ml HGF and 50% Wnt conditioned media and, preferably 10-100ng/ml Noggin. Wnt conditioned media comprises Advanced DMEM, P/S, B27, N2 and also FCS. 293T cells transfected with Wnt3A expression plasmid produce Wnt. The whole medium is taken after a few days (i.e. with secreted Wnt) and used as the Wnt source.

[0083] The invention therefore provides a cell culture medium, comprising or consisting of a basal medium for animal or human cells to which is added:

Epidermal Growth Factor, an FGF able to bind to FGFR2 or FGFR4, preferably FGF10 and HGF as mitogenic growth factors,

gastrin, Nicotinamide, B27, N2 and N-Acetylcystein, and preferably

a BMP inhibitor more preferably Noggin and

a Wnt agonist selected from R-spondin 1, R-spondin 2, R-spondin 3 and R-spondin 4, more preferably R-spondin 1, and optionally Wnt-3a.

[0084] The invention thus encompasses a first preferred culture medium comprising or consisting of a basal medium for animal or human cells to which is added:

Epidermal Growth Factor, FGF10 and HGF as mitogenic growth factors,

gastrin, Nicotinamide, B27, N2 and N-Acetylcysteine,

a BMP inhibitor more preferably Noggin and

a Wnt agonist selected R-spondin 1, R-spondin 2, R-spondin 3 and R-spondin 4, more preferably comprising R-spondin 1 and Wnt-3a.

[0085] This medium may be used as an EM1 cell culture medium of the invention to stimulate initial expansion of cells.

[0086] In some embodiments, the medium used as an EM1 cell culture medium comprises all the components of an EM2 culture medium of the invention and additionally comprises Wnt-3a and Noggin.

[0087] In embodiments in which the basal medium is supplemented with N-Acetylcysteine, B27 and N2, the following are preferably added to the culture media: EGF, R-spondin1, gastrin, FGF10, Nicotinamide and HGF and Wnt-conditioned media. Preferably, the basal medium is supplemented with N-Acetylcysteine, EGF, R-spondin1, gastrin, FGF10, Nicotinamide and HGF and Wnt-conditioned media in accordance with the quantities described hereinabove.

[0088] For example, in some embodiments the basal medium may be supplemented with 150ng/ml to 250 ng/ml N-Acetylcysteine; preferably, the basal medium is supplemented with, about or exactly 200ng/ml N-Acetylcysteine. For example, in some embodiments the basal medium may be supplemented with 40ng/ml to 60ng/ml EGF; preferably, the basal medium is supplemented with about or exactly 50ng/ml EGF. For example, in some embodiments the basal medium may be supplemented with 0.5 µg/ml to 1.5 µg/ml R-spondin1; preferably, the basal medium is supplemented with about or exactly 1 µg/ml R-spondin1. For example, in some embodiments the basal medium may be supplemented with 5nM to 15nM gastrin; preferably, the basal medium is supplemented with about or exactly 10nM gastrin. For example, in some embodiments the basal medium may be supplemented with 25-200ng/ml FGF10, for example 70 ng/ml to 130 ng/ml FGF10; preferably, the basal medium is supplemented with about or exactly 100 ng/ml FGF10. For example, in some embodiments the basal medium may be supplemented with 5mM to 15mM Nicotinamide; preferably, the basal medium is supplemented with about or exactly 10mM Nicotinamide. For example, in some embodiments the basal medium may be supplemented with 25ng/ml to 100 ng/ml HGF, for example 35ng/ml to 65ng/ml HGF; preferably, the basal medium is supplemented with about or exactly 50ng/ml HGF. For example, in some embodiments the basal medium may be supplemented with 35% to 65% Wnt-conditioned media; preferably, the basal medium is supplemented with about or exactly 50% Wnt-conditioned media.

[0089] In some embodiments one or both of gastrin and N2 are not present in the cell culture medium.

[0090] Preferably, the basal medium is advanced-DMEM/F12.

[0091] This first culture medium (for example, EM1, EM2 or both EM1 and EM2) is preferably used during the first two weeks of the culture method of the invention. However, it may be used for a shorter time period, such as for 1, 2, 3, 5, 7, or 10 days, or a longer time period, such as 3, 4, 5, 10, 20 or more weeks, 5 months or more, for example, 6, 7, 8, 9, 10, 11, 12 months or more.

Differentiation Medium (DM):

[0092] In some embodiments, the method of the invention involves a second cell culture medium which comprises or consists of a basal medium for animal or human cells to which is added: EGF, a TGF-beta inhibitor, and a Notch inhibitor. The inventors have found that this medium is useful for differentiating cells. The medium used for differentiating the cells may be referred to herein as DM.

[0093] Preferably, the second cell culture medium also comprises FGF and/or HGF.

[0094] In one embodiment, the second culture medium comprises or consists of a basal medium for animal or human cells to which is added:

Epidermal Growth Factor, FGF10 and HGF as mitogenic growth factors;

a Notch inhibitor; and

a TGF-beta inhibitor.

[0095] In one embodiment, the TGF-beta inhibitor is A83-01 and/or the Notch inhibitor is DAPT. In another embodiment, the DM cell culture medium additionally comprises Dexamethasone.

[0096] A preferred second cell culture medium, and method of using this medium, is described in the examples, and comprises or consists of a basal medium to which is added: 50ng/ml EGF, 100 ng/ml FGF10, 50 nM A8301 and 10 uM DAPT. Advanced-DMEM/F12 may be used as the basal medium as may any other suitable basal medium.

[0097] In some embodiments, the second cell culture medium does not comprise R-spondin or Wnt. In some embodiments, the second cell culture medium does not comprise a Wnt agonist. In some embodiments, the second cell culture medium does not comprise Nicotinamide. In some embodiments, the second cell culture medium does not comprise a BMP inhibitor.

[0098] The inventors have discovered that R-spondin1 and Nicotinamide both inhibit the

expression of the mature hepatocyte marker CYP3A11 and yet promote the expression of the hepatoblast marker albumin. Therefore, to increase differentiation of the cells to more mature liver fates, the inventors removed R-spondin and Nicotinamide from the cell culture. The inventors have also discovered that the expression of specific biliary transcription factors is highly upregulated in expansion cultures containing R-spondin1, indicating that the culture gene expression was unbalanced towards a more biliary cell fate. Notch and TGF-beta signaling pathways have been implicated in biliary cell fate *in vivo*. In fact, deletion of Rbpj (essential to achieve active Notch signalling) results in abnormal tubulogenesis (Zong Y. Development 2009) and the addition of TGF-beta to liver explants facilitates the biliary differentiation *in vitro* (Clotman F. Genes and Development 2005). Since both Notch and TGF-beta signalling pathways were highly upregulated in the liver cultures (Figure 9) the inventors reasoned that inhibition of biliary duct cell-fate might trigger the differentiation of the cells towards a more hepatocytic phenotype. It was found that addition of a TGF-beta inhibitor (such as A8301) and a Notch inhibitor (such as DAPT) to a differentiation medium that preferably does not contain R-spondin or Wnt, enhances the expression of mature hepatocyte markers and increases the number of hepatocyte-like cells (for example, see example 2).

[0099] Notch is a transmembrane surface receptor that can be activated through multiple proteolytic cleavages, one of them being cleavage by a complex of proteins with protease activity, termed gamma-secretase. Gamma-secretase is a protease that performs its cleavage activity within the membrane. Gamma-secretase is a multicomponent enzyme and is composed of at least four different proteins, namely, presenilins (presenilin 1 or 2), nicastrin, PEN-2 and APH-I. Presenilin is the catalytic centre of gamma-secretase. On ligand binding the Notch receptor undergoes a conformational change that allows ectodomain shedding through the action of an ADAM protease which is a metalloprotease. This is followed immediately by the action of the gamma-secretase complex which results in the release of the Notch intracellular domain (NICD). NICD translocates to the nucleus where it interacts with CSL (C-promoter-binding factor/recombinant signal-sequence binding protein Jk/Suppressor-of-Hairless/Lag1). The binding of NICD converts CSL from a transcriptional repressor to an activator which results in the expression of Notch target genes. Examples of preferred Notch inhibitors that can be used in the context of this invention are: gamma-secretase inhibitors, such as DAPT or dibenzazepine (DBZ) or benzodiazepine (BZ) or LY-411575, an inhibitor capable of diminishing ligand mediated activation of Notch (for example via a dominant negative ligand of Notch or via a dominant negative Notch or via an antibody capable of at least in part blocking the interacting between a Notch ligand and Notch), or an inhibitor of ADAM proteases.

[0100] TGF-beta signalling is involved in many cellular functions, including cell growth, cell fate and apoptosis. Signalling typically begins with binding of a TGF-beta superfamily ligand to a type II receptor which recruits and phosphorylates a type I receptor. The type 1 receptor then phosphorylates SMADs, which act as transcription factors in the nucleus and regulate target gene expression. Alternatively, TGF-beta signalling can activate MAP kinase signalling pathways, for example, via p38 MAP kinase. The TGF-beta superfamily ligands comprise bone morphogenic proteins (BMPs), growth and differentiation factors (GDFs), anti-müllerian

hormone (AMH), activin, nodal and TGF-betas.

[0101] A TGF-beta inhibitor is an agent that reduces the activity of the TGF-beta signalling pathway. There are many ways of disrupting the TGF-beta signaling pathway that are known in the art and that can be used in conjunction with this invention. For example, the TGF-beta signaling may be disrupted by: inhibition of TGF-beta expression by a small-interfering RNA strategy; inhibition of furin (a TGF-beta activating protease); inhibition of the pathway by physiological inhibitors, such as inhibition of BMP by Noggin, DAN or DAN-like proteins; neutralisation of TGF-beta with a monoclonal antibody; inhibition with small-molecule inhibitors of TGF-beta receptor kinase 1 (also known as activin receptor-like kinase, ALK5), ALK4, ALK6, ALK7 or other TGF-beta-related receptor kinases; inhibition of Smad 2 and Smad 3 signaling by overexpression of their physiological inhibitor, Smad 7, or by using thioredoxin as an Smad anchor disabling Smad from activation (Fuchs, O. Inhibition of TGF-Signaling for the Treatment of Tumor Metastasis and Fibrotic Diseases. Current Signal Transduction Therapy, Volume 6, Number 1, January 2011, pp. 29-43(15)).

[0102] Various methods for determining if a substance is a TGF-beta inhibitor are known and might be used in conjunction with the invention. For example, a cellular assay may be used in which cells are stably transfected with a reporter construct comprising the human PAI-1 promoter or Smad binding sites, driving a luciferase reporter gene. Inhibition of luciferase activity relative to control groups can be used as a measure of compound activity (De Gouville et al., Br J Pharmacol. 2005 May; 145(2): 166-177).

[0103] A TGF-beta inhibitor according to the present invention may be a protein, peptide, small-molecules, small-interfering RNA, antisense oligonucleotide, aptamer or antibody. The inhibitor may be naturally occurring or synthetic. Examples of preferred small-molecule TGF-beta inhibitors that can be used in the context of this invention include the small molecule inhibitors listed in table 1:

Table 1: Small-molecule TGF-beta inhibitors targeting receptor kinases

Inhibitor	Targets	IC50 (nM)	Mol Wt	Name	Formula
A83-01	ALK5 (TGF- β R1)	12	421.52	3-(6-Methyl-2-pyridinyl)-N-phenyl-4-(4-quinolinyl)-1H-pyrazole-1-carbothioamide	C25H19N5S
	ALK4	45			
	ALK7	7.5			
SB-431542	ALK5	94	384.39	4-[4-(1,3-benzodioxol-5-yl)-5-(2-pyridinyl)-1H-imidazol-2-yl]benzamide	C22H16N4O3
	ALK4				
	ALK7				
SB-505124	ALK5	47	335.4	2-(5-benzo[1,3]dioxol-5-yl-2-tert-butyl-3Himidazol-4-yl)-6-methylpyridine hydrochloride hydrate	C20H21N3O2
	ALK4	129			

Inhibitor	Targets	IC50 (nM)	Mol Wt	Name	Formula
SB-525334	ALK5	14.3	343.42	6-[2-(1,1-Dimethylethyl)-5-(6-methyl-2-pyridinyl)-1H-imidazol-4-yl]quinoxaline	C21H21N5
SD-208	ALK5	49	352.75	2-(5-Chloro-2-fluorophenyl)-4-[(4-pyridyl) amino]pteridine	C17H10C1FN6
LY-36494	TGR- β RI	59	272.31	4-[3-(2-Pyridinyl)-1H-pyrazol-4-yl]-quinoline	C17H12N4
	TGF- β RII	400			
	MLK-7K	1400			
SJN-2511	ALK5	23	287.32	2-(3-(6-Methylpyridine-2-yl)-1H-pyrazol-4-yl)-1,5-naphthyridine	C17H13N5

[0104] A method of the invention may comprise the use of one or more of any of the inhibitors listed in table 1. A method of the invention may comprise the use of any combination of one inhibitor with another inhibitor listed. For example, a method of the invention may comprise the use of SB-525334 or SD-208 or A83-01; or a method of the invention may comprise the use of SD-208 and A83-01; or a method of the invention may comprise the use of SD-208 and A83-01. The skilled person will appreciate that a number of other small-molecule inhibitors exist that are primarily designed to target other kinases, but at high concentrations may also inhibit TGF-beta receptor kinases. For example, SB-203580 is a p38 MAP kinase inhibitor that, at high concentrations (for example, approximate 10 μ M or more) is thought to inhibit ALK5. Any such inhibitor that inhibits the TGF-beta signalling pathway can also be used in the context of this invention.

[0105] A83-01 may be added to the culture medium at a concentration of between 10 nM and 10 μ M, or between 20 nM and 5 μ M, or between 50 nM and 1 μ M. For example, A83-01 may be added to the culture medium at approximately 500 nM. When used in an EM, A83-01 may be added to the culture medium at a concentration of between 350-650 nM, for example, 450-550 nM, more preferably approximately 500 nM. When used in the DM, A83-01 may be added to the culture medium at a concentration of between 25-75 nM, for example, 40-60 nM or approximately 50nM.

[0106] SB-431542 may be added to the culture medium at a concentration of between 80 nM and 80 μ M, or between 100 nM and 40 μ M, or between 500 nM and 10 μ M. For example, SB-431542 may be added to the culture medium at approximately 1 μ M.

[0107] SB-505124 may be added to the culture medium at a concentration of between 40 nM and 40 μ M, or between 80 nM and 20 μ M, or between 200 nM and 1 μ M. For example, SB-505124 may be added to the culture medium at approximately 500 nM.

[0108] SB-525334 may be added to the culture medium at a concentration of between 10 nM and 10 μ M, or between 20 nM and 5 μ M, or between 50 nM and 1 μ M. For example, SB-525334 may be added to the culture medium at approximately 100 nM.

[0109] LY 364947 may be added to the culture medium at a concentration of between 40 nM and 40 μ M, or between 80 nM and 20 μ M, or between 200 nM and 1 μ M. For example, LY 364947 may be added to the culture medium at approximately 500 nM.

[0110] SD-208 may be added to the culture medium at a concentration of between 40 nM and 40 μ M, or between 80 nM and 20 μ M, or between 200 nM and 1 μ M. For example, SD-208 may be added to the culture medium at approximately 500 nM.

[0111] SJN 2511 may be added to the culture medium at a concentration of between 20 nM and 20 μ M, or between 40 nM and 10 μ M, or between 100 nM and 1 μ M. For example, A83-01 may be added to the culture medium at approximately 200 nM.

[0112] In a further aspect, the invention involves an alternative second culture medium comprising or consisting of a basal medium for animal or human cells to which is added:

Epidermal Growth Factor, FGF10 and HGF as mitogenic growth factors;

gastrin, Nicotinamide, B27, N2 and N-Acetylcysteine.

[0113] This alternative second culture medium is useful as a differentiation medium (DM). This alternative second culture medium is preferably used after the two first weeks of culture. At this stage, it seems that a BMP inhibitor and a Wnt ligand are no longer needed. Thus, in some embodiments, the alternative second culture medium does not comprise a BMP inhibitor or a Wnt ligand. In some embodiments, the alternative second culture medium does not comprise a BMP inhibitor or a Wnt agonist.

[0114] As with the first culture medium (EM, i.e. EM1 and EM2 or EM1 or EM2), the basal medium may in some embodiments be supplemented with N-Acetylcysteine, EGF, gastrin, FGF10, Nicotinamide and HGF in accordance with the quantities described hereinabove. For example, preferably, the basal medium is supplemented with about or exactly 200ng/ml N-Acetylcysteine. Preferably, the basal medium is supplemented with about or exactly 50ng/ml EGF. Preferably, the basal medium is supplemented with about or exactly 10nM gastrin. Preferably, the basal medium is supplemented with about or exactly 100 ng/ml FGF10. Preferably, the basal medium is supplemented with about or exactly 10mM Nicotinamide. Preferably, the basal medium is supplemented with about or exactly 50ng/ml HGF.

[0115] A first and a second cell culture medium used according to the invention allows the survival and/or proliferation and/or differentiation of epithelial stem cells or liver epithelial stem

cells or isolated biliary duct or isolated liver fragments on an extracellular matrix.

[0116] A medium allowing for survival and/or proliferation is a medium which preferably induces or promotes the survival and/or proliferation of cells during at least 5, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200 days of culture. Proliferation can be assessed using techniques known in the art such as BrdU staining, Edu staining, Ki67 staining and the use of growth curves assay can be done. For example, we have shown that from 10 biliary ducts, it is possible after 6 days to dilute to 6 wells with 10 organoids per well (60 new organoids). In each passage of 6 days we can generate around 360 organoids. By performing 1 passage/week over 32 weeks we are able to generate over 11,000 (11520) new organoid structures in only 7 months. This is important for the industry, since the availability of cells and organoids for transplantation poses a significant problem. For a mouse transplant, for example, a minimum of 10^5 cells are required. Possibly 10^6 , or 10×10^6 might be required for a human transplant, in order for a graft to be successful.

[0117] Put another way, media used according to the invention is capable of expanding a population of stem cells to form liver organoids for at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100, passages under appropriate conditions.

[0118] The second medium as identified above preferably induces or promotes a specific differentiation of cells during at least five days of culture. Differentiation may be measured by detecting the presence of a specific marker associated with the liver lineage as defined herein. Differentiation may be measured by detecting the presence of a specific marker associated with the liver lineage as defined herein. Depending on the identity of the marker, the expression of said marker may be assessed by RTPCR or immuno-histochemistry after at least 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 days of culture in a first medium or in a first subsequently in a second medium as defined herein.

[0119] A differentiation medium is a medium which preferably induces or promotes a specific differentiation of cells during at least five days of culture. Differentiation may be measured by detecting the presence of a specific marker associated with the liver lineage as defined herein.

[0120] Within the context of the invention, the term cell culture medium is synonymous with medium, culture medium or cell medium or basal medium or basal cell medium or basal cell culture medium or expansion medium or differentiation medium.

[0121] According to a still further aspect of the invention, there is provided a hermetically-sealed vessel containing a culture medium of the invention. In some embodiments, the culture medium is an expansion medium. In some embodiments, the culture medium is a differentiation medium. Hermetically-sealed vessels may be preferred for transport or storage of the culture media, to prevent contamination. The vessel may be any suitable vessel, such as a flask, a plate, a bottle, a jar, a vial or a bag.

Methods for Obtaining and/or Culturing Stem Cells

[0122] The method for obtaining and/or culturing a liver fragment or a liver organoid, comprises culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of one or more cell culture media according to the invention.

[0123] In some embodiments, a method for obtaining and/or culturing a liver fragment or a liver organoid comprises culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of EM1 medium and then EM2 medium and then DM medium.

[0124] In some embodiments, a method for obtaining and/or culturing a liver fragment or a liver organoid comprises culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of EM2 medium and then DM medium without the use of EM1 medium.

[0125] In some embodiments, a method for obtaining and/or culturing a liver fragment or a liver organoid comprises culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of EM1 medium and then DM medium without the use of EM2 medium.

[0126] The method for obtaining and/or culturing a liver fragment or a liver organoid, may comprise:

1. i) culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of a medium, the medium comprising a basal medium for animal or human cells to which is added: Epidermal Growth Factor, an FGF able to bind to FGFR2 or FGFR4, preferably FGF10 and HGF as mitogenic growth factors, Nicotinamide, and preferably, a Wnt agonist, preferably any one of R-spondin 1-4; and subsequently
2. ii) culturing the stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of a second cell culture medium as defined here (a DM medium).

[0127] In some embodiments, prior to the step i), the method comprises culturing the epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in the presence of a medium comprising or consisting of a basal medium for animal or human cells to which is added EGF, a BMP inhibitor, R-spondin and Wnt. Preferably, the BMP inhibitor is Noggin and the EM1 medium is termed "ENRW" (EGF, Noggin, R-spondin and Wnt).

[0128] In one embodiment there is provided a method for obtaining and/or culturing a liver fragment or a liver organoid, wherein said method comprises:

culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in an EM2 culture medium as described herein, for example, a medium comprising a Wnt agonist, such as R-spondin, Epidermal Growth Factor, an FGF able to bind to FGFR2 or FGFR4, preferably FGF10, and nicotinamide. Preferably, the medium also comprises HGF.

[0129] In a further embodiment there is provided a method for obtaining and/or culturing a liver fragment or a liver organoid, wherein said method comprises:

culturing epithelial stem cells, and/or isolated tissue fragments comprising said epithelial stem cells in contact with an extracellular matrix in a medium comprising a BMP inhibitor, a Wnt agonist, Epidermal Growth Factor, a FGF able to bind to FGFR2 or FGFR4, preferably FGF10 and HGF as mitogenic growth factors, gastrin, Nicotinamide, B27, N2 and N-Acetylcysteine.

[0130] In another preferred method, after at least two weeks of culture in an EM1 medium as defined above, the culture is continued in a medium that does not comprise a BMP inhibitor and a Wnt agonist. In some embodiments, this culture is continued in an EM2 of the invention. In alternative embodiments, the culture is continued in a second medium of the invention (DM), which additionally comprises a TGF-beta inhibitor, such as A83-01, and a Notch inhibitor, such as DAPT. In some embodiments, this culture is continued in an EM2 and then a DM medium of the invention.

[0131] Therefore in a preferred method for obtaining and/or culturing a liver organoid, epithelial stem cells, and/or isolated liver tissue fragments comprising said epithelial stem cells are cultured in a first step in the first medium (EM), then subsequently in a discrete second step in the second medium (DM). The first culture step may have a duration of at least two weeks and may be longer, for example, 3 weeks or longer, 4 weeks or longer or 1, 2, 3, 4, 5, 6, 7, 8 months or longer. The second step preferably has a duration of 6 weeks or less, more preferably 1 month or less, for example 3 weeks or less, 2 weeks or less, 1 week or less. It has been found that many cells start dying after 2 weeks in DM and so preferably, the second step is 2 weeks or less or at most 1 month. Thus, it is only in less preferred embodiments, that the second step may have a duration of 8, 9, 10, 11, 12, 13, 14, 15, 16 days or 3 weeks or 1, 2, 3, 4, 5, 6 months or longer. Each step is preferably carried out using an extracellular matrix as defined herein, and using the culture media described in detail herein.

[0132] Alternatively in another embodiment, said method is carried out in the first medium, without switch to the second medium. This method may have a duration of 8, 9, 10, 11, 12, 13, 14, 15, 16 days or 3 weeks or 1, 2, 3, 4, 5, 6, 7, 8 months or longer.

[0133] In some embodiments, the step of culturing in the first medium of the invention comprises both culturing in EM1 medium and then culturing in EM2 medium. Thus, the periods described above may refer to the total period in which the cells or fragments are cultured in the

EM1 and subsequently in the EM2 media. In some embodiments, the step of culturing in the EM1 culture medium may have a duration of less than 10 days, for example, less than 9, less than 8, less than 7, less than 6, less than 5, less than 4, less than 3, less than 2 or less than 1 day. In some embodiments, the step of culturing in the EM1 culture medium may have a duration of between 1 and 10 days, for example, 1-7 days, 2-6 days, 3-5 days, for example 4 or 5 days. In some embodiments, the step of culturing in the EM2 medium may have a duration of 10 days or longer, for example, 11, 12, 13, 14, 15, 16 days or longer, or 1-3 weeks or 2-4 weeks or 3-6 weeks or 1, 2, 3, 4, 5, 6 months or longer.

[0134] Preferred concentrations of each compound present in each medium have already been defined herein in the description or in the examples.

[0135] The methods described herein for obtaining and/or culturing a liver fragment or a liver organoid may also be used to obtain and/or culture a population of cells expressing Lgr5 and such methods are encompassed by the invention. A population of cells expressing Lgr5 obtained by a method of the invention is thus also disclosed.

[0136] As will be apparent to the skilled reader, the preferred culture methods of the invention are advantageous because feeder cells are not required. Feeder cell layers are often used to support the culture of stem cells, and to inhibit their differentiation. A feeder cell layer is generally a monolayer of cells that is co-cultured with, and which provides a surface suitable for growth of, the cells of interest. The feeder cell layer provides an environment in which the cells of interest can grow. Feeder cells are often mitotically inactivated (e.g. by irradiation or treatment with mitomycin C) to prevent their proliferation. The use of feeder cells is undesirable, because it complicates passaging of the cells (the cells must be separated from the feeder cells at each passage, and new feeder cells are required at each passage). The use of feeder cells can also lead to contamination of the desired cells with the feeder cells. This is clearly problematic for any medical applications, and even in a research context, complicates analysis of the results of any experiments performed on the cells. As noted elsewhere herein, the culture media of the invention are particularly advantageous because they can be used to culture cells without feeder cell contact, i.e. the methods of the invention do not require a layer of feeder cells to support the cells whose growth is being sponsored.

[0137] Accordingly, the compositions of the invention may be feeder cell-free compositions. A composition is conventionally considered to be feeder cell-free if the liver cells in the composition have been cultured for at least one passage in the absence of a feeder cell layer. A feeder cell-free composition of the invention will normally contain less than about 5%, less than about 4%, less than about 3%, less than about 2%, less than about 1% feeder cells (expressed as a % of the total number of cells in the composition) or preferably no feeder cells at all.

[0138] In one preferred embodiment, for example, if a liver organoid is to be used for regenerative medicine, the method may start from epithelial cells or from an isolated liver fragment. The cells or liver fragment may be autologous or allogeneic. "Autologous" cells are

cells which originated from the same organism into which they are being re-introduced for cellular therapy, for example in order to permit tissue regeneration. An autologous cell does not, in principle, require matching to the patient in order to overcome the problems of immune rejection, and/or reduces the need for immune suppression interventions upon transplant. "Allogeneic" cells are cells which originated from an individual which is different from the individual into which the cells are being introduced for cellular therapy, for example in order to permit tissue regeneration, although of the same species. Some degree of patient matching may still be required to prevent the problems of rejection. Techniques for minimising tissue rejection will be known to those of skill in the art.

[0139] A preferred method of the invention encompasses a method for culturing an isolated liver fragment comprising said epithelial stem cells.

[0140] Another preferred method encompasses a method for obtaining a liver organoid from epithelial stem cells and/or isolated liver fragments comprising said epithelial stem cells.

[0141] Another preferred method encompasses a method for obtaining and culturing a liver organoid from epithelial stem cells and/or isolated liver fragments comprising said epithelial stem cells.

[0142] In a preferred method as earlier identified herein, the BMP inhibitor is selected from Noggin, DAN, and DAN-like proteins including Cerberus and Gremlin, preferably comprises or is Noggin, and/or said Wnt agonist is selected from one or more of Wnt, preferably Wnt-3a, R-spondin 1-4, Norrin, and a GSK-inhibitor, more preferably comprises or is Wnt-3a and/or R-spondin.

[0143] In another preferred method, a liver organoid originates from one single cell, preferably a cell expressing Lgr5, more preferably wherein the single cell comprises a nucleic acid construct comprising a nucleic acid molecule of interest. The cell may also express liver-specific markers such as Hnf1 α , and Hnf4.

[0144] The isolation of certain cell types expressing Lgr5 has already been described previously (see, for example, WO2009/022907 and WO2010/016766). However, liver specific stem cells expressing Lgr5 of the type disclosed herein have not previously been described. Accordingly, the invention provides an organoid comprising a population of adult stem cells characterised by natural expression of at least Lgr5 and one or more of the following markers Hnf1 α , Hnf4a, Sox9, KRT7 and KRT19 at a significant level. This cell population also expresses markers of progenitor populations common to the small intestine and stomach, such as Cd44 and Sox9 (Barker & Huch et al Cell stem cell 2010). These are highly expressed in the stem cells according to the invention, but are not expressed in adult liver, reinforcing the self-renewal capacity of the liver cultures described herein. Cells according to this aspect of the invention may also up-regulate Wnt target genes, including for example, MMP7, Sp5 and Tnfrs19. This provides strong evidence of the requirement for an active and robust canonical Wnt signalling activity to maintain the self renewing capacity of these cultures.

[0145] By "natural expression" is meant that the cells have not been manipulated recombinantly in any way, i.e., the cells have not been artificially induced to express these markers or to modulate these markers' expression by introduction of exogenous genetic material, such as introduction of heterologous (non-natural) or stronger promoters or other regulatory sequences operably linked to either the endogenous genes or exogenously-introduced forms of the genes. Natural expression is from genomic DNA within the cells, including introns between the exon coding sequences where these exist. Natural expression is not from cDNA. Natural expression can if necessary be proven by any one of various methods, such as sequencing out from within the reading frame of the gene to check that no extraneous heterogenous sequence is present. "Adult" means post-embryonic. With respect to the stem cells of the present invention, the term "adult stem cell" means that the stem cell is isolated from a tissue or organ of an animal at a stage of growth later than the embryonic stage.

[0146] This stem cell population can also be characterised by a lack of natural expression of certain markers at any significant level, many of which are associated with cellular differentiation. Specifically, the cells of the isolated adult stem cell population do not naturally express one or more of Cd11b, CD13, CD14, AFP, Pdx1, any CYP member (e.g. CYP3A11, CYP 11A1) at a significant level. As defined herein, these markers are said to be negative markers.

[0147] The term "expressed" is used to describe the presence of a marker within a cell. In order to be considered as being expressed, a marker must be present at a detectable level. By "detectable level" is meant that the marker can be detected using one of the standard laboratory methodologies such as PCR, blotting or FACS analysis. A gene is considered to be expressed by a cell of the population of the invention if expression can be reasonably detected after 30 PCR cycles, which corresponds to an expression level in the cell of at least about 100 copies per cell. The terms "express" and "expression" have corresponding meanings. At an expression level below this threshold, a marker is considered not to be expressed. The comparison between the expression level of a marker in a cell of the invention, and the expression level of the same marker in another cell, such as for example an embryonic stem cell, may preferably be conducted by comparing the two cell types that have been isolated from the same species. Preferably this species is a mammal, and more preferably this species is human. Such comparison may conveniently be conducted using a reverse transcriptase polymerase chain reaction (RT-PCR) experiment.

[0148] Any one of a number of physical methods of separation known in the art may be used to select the cells of this aspect of the invention and distinguish these from other cell types. Such physical methods may involve FACS and various immuno-affinity methods based upon markers specifically expressed by the cells. As described above, Lgr5, Hnf1 α and Hnf4 are 3 of the cell markers expressed at high levels in the cells. Therefore, by way of illustration only, the cells may be isolated by a number of physical methods of separation, which rely on the presence of these markers.

[0149] In one embodiment, the cells may be isolated by FACS utilizing an antibody, for example, against one of these markers. As will be apparent to one skilled in the art, this may be achieved through a fluorescent labeled antibody, or through a fluorescent labeled secondary antibody with binding specificity for the primary antibody. Examples of suitable fluorescent labels includes, but is not limited to, FITC, Alexa Fluor® 488, GFP, CFSE, CFDA-SE, DyLight 488, PE, PerCP, PE-Alexa Fluor® 700, PE-Cy5 (TRI-COLOR®), PE-Cy5.5, PI, PE-Alexa Fluor® 750, and PE-Cy7 . This list is provided by way of example only, and is not intended to be limiting.

[0150] It will be apparent to a person skilled in the art that FACS analysis using an anti-Lgr5 antibody will provide a purified cell population. However, in some embodiments, it may be preferable to purify the cell population further by performing a further round of FACS analysis using one or more of the other identifiable markers, preferably Hnf1 α and Hnf4, but others may also be used.

[0151] In another embodiment, the cells may be isolated by immuno-affinity purification, which is a separation method well known in the art. By way of illustration only, the cells may be isolated by immuno-affinity purification directed towards c-kit. As will be apparent to one skilled in the art, this method relies upon the immobilisation of antibodies on a purification column. The cell sample is then loaded onto the column, allowing the appropriate cells to be bound by the antibodies, and therefore bound to the column. Following a washing step, the cells are eluted from the column using a competitor which binds preferentially to the immobilised anti-c-kit antibody, and permits the cells to be released from the column.

[0152] It will be apparent to a person skilled in the art that immuno-affinity purification using an immobilised antibody will provide a purified cell population. However, in some embodiments, it may be preferable to purify the cell population further by performing a further round of immuno-affinity purification using one or more of the other identifiable markers, for example Hnf4, and use an aliquot of the isolated clones to ascertain the expression of other relevant intracellular markers.

[0153] It will be apparent to a person skilled in the art that the sequential purification steps are not necessarily required to involve the same physical method of separation. Therefore, it will be clear that, for example, the cells may be purified through a FACS step using an anti-Lgr5 antibody, followed by an immuno-affinity purification step using a SSEA-1 affinity column. In certain embodiments, the cells may be cultured after isolation for at least about 15, at least about 20 days, at least about 25 days, or at least about 30 days. In certain aspects, the cells are expanded in culture longer to improve the homogeneity of the cell phenotype in the cell population.

[0154] Other features of this method are defined in the part of the description dedicated to definitions. Single-cell suspensions or small clusters of cells (2-50 cells/cluster) will normally be seeded, rather than large clusters of cells, as is known in the art. As they divide, such cells will be seeded onto a support at a density that promotes cell proliferation. Typically, when single

cells are isolated the plating density of at least 1-500 cells/well is used, the surface of the well being 0.32 cm². When clusters are seeded the plating density is preferably 250-2500 cells/cm². For replating, a density of between about 2500 cells/ cm² and about 5,000 cells/ cm² may be used. During replating, single-cell suspensions or small cluster of cells will normally be seeded, rather than large clusters of cells, as is known in the art.

Organoids of the Invention

[0155] The cells described above grow into bodies which are herein termed "organoids". Accordingly, a liver organoid obtainable by a method of the invention is a further aspect of the invention. To the best of our knowledge, this is the first time that a liver organoid has been obtained that is functional and alive after such an extended period of time (i.e. at least 7 months of culture; see examples included herein). Functionality is preferably characterized by the presence of a liver marker as defined herein and/or by the structure of said organoid as defined herein. Since the final amount of liver organoids obtained correlates with the duration of culture, the skilled person will understand that the invention is a pioneer invention and potentially opens new possibilities in for example regenerative medicine.

[0156] For example, an organoid according to the present invention may comprise a population of cells of at least 1x10³ cells, at least 1 x 10⁴ cells, at least 1x10⁵ cells, at least 1x10⁶ cells, at least 1x10⁷ cells or more. In some embodiments, each organoid comprises between approximately 1x10³ cells and 5x10³ cells; generally, 10-20 organoids may be grown together in one well of a 24 well plate.

[0157] Cells and organoids according to the present invention may be non-human animal or human. The inventors have shown, for the first time, that it is possible to grow and maintain both animal and human liver organoids *in vitro*, using the culture media and methods of the invention.

[0158] Illustrative examples of organoids generated according to the invention are given in the accompanying figures. It can be seen that organoids according to the invention may possess a cystic structure, with on the outside, a layer of cells with at least one bud and a central lumen. The organoids in the outside of the matrigel tend to be larger than the organoids in the center of the matrigel, perhaps because they have better access to the necessary growth factors. Structurally, organoids according to the invention are often elongated in shape. They may include one or more budding structure - a single cell epithelial layer which has a structure not unlike a bile duct. Under confocal microscopy, the structures may stain positive for keratin. They may include cells with polarised nuclei and small cytoplasm. The organoids may have a section which is formed of multiple layers; such cells often tend to have their nuclei more central to the cells, i.e. not polarized. The cells in the multilayer section may organise themselves to include a gap, or lumen between the cells.

[0159] A liver organoid preferably comprises a hepatocyte and a cholangiocyte cell, more preferably wherein at least one of the following markers could be detected: at least one hepatocyte marker such as albumin, transthyretin, B-1 integrin and Glutamine synthetase and/or at least one of CYP3A11, FAH, tbx3, TAT and Gck and/or at least one cholangiocyte maker such as Keratin 7 and 19. The skilled person knows how to detect each of these markers (i.e. RT-PCR and/or immunohistochemistry). Preferably the expression of each of these markers is assessed as carried out in the experimental part. Each of these markers is usually expressed after at least two weeks, three weeks or one month of culture using a method of the invention. Microarray analysis of the organoids in both culture conditions showed that liver organoids resemble adult liver tissue.

[0160] In some embodiments, approximately 35% of the cells in a liver organoid express a hepatocyte surface marker, for example, 25-45%, 30-40%, 33-37%, 35% or less, or 15-35% of cells.

[0161] Preferably, cells and organoids generated according to the invention also possess hepatocyte functions, such as expressing or staining positive for the mature hepatic markers albumin, B-1 integrin, CK-8, CK-18, transthyretin (TTR), glucose 6P, Met, Glutamine synthase (Glul), transferrin, Fahd1, Fahd2a, K7, K19 and cytochrome P450 isoforms 3A13 (CYP3A13), 51 (CYP51) 2D10 (CYP2D10), 2j6 (CYP2j6), 39A1 (CYP39A1), 4A10 (CYP4A10), 4F13 (CYP4F13) 4F16 (CYP4F16), CYP4B1 and 20A1 (CYP20A1). Also, embryonic liver gene AFP is in some embodiments not detected in neither of both culture conditions, as in adult liver. In some embodiments, the expression of alpha fetal protein is just above the background gene expression.

[0162] Also, the well known liver transcription factors as HNF1a, HNF1b and HNF4a are highly expressed in both conditions.

[0163] Since liver and pancreas are closely related organs, we investigated whether our liver cultures also expressed pancreas-specific genes. The pancreas is functionally divided into endocrine and exocrine pancreas. The endocrine pancreas is mainly characterized for expressing insulin, glucagon and somatostatin. The expression of these hormones is tightly regulated by a set of endocrine pancreas-specific transcription factors, the most important being Pdx1 and NeuroD. The exocrine pancreas is formed by acinar and ductal compartments responsible of producing the digestive enzymes amylase, pancreatic lipase and chymotrypsin, among others. The expression of these genes is also regulated by specific exocrine pancreatic genes as Ptf1.

[0164] The pancreas specific genes Ptf1a, pancreatic amylase (Amy2a4), pancreatic lipase (Pnlip), insulin (ins1 and ins2), glucagon (Gcg), chymotrypsin (cela1), Pdx1 and NeuroD were absent in the liver cultures here described.

[0165] In some embodiments, one or more or all of the following genes are expressed in the liver organoids at a similar level to the corresponding gene in adult liver hepatocytes: Aqp1,

Bmp2, Apo3, Apol7a, Sord, C3, Ppara, Pparg, tbx3, Igf1, Il17rb, Il1b, Tgfb, ApoA1, ApoA4, Apob, Cyp26b1, Cyp27a1, Cyp2b13, Cyp2b9, Cyp2c37, Cyp2f2, Cyp2g1, Cyp2j13, Cyp3a11, Cyp4a10 and Cypf14. For example, see Figure 16A.

[0166] In some embodiments, one or more of the following genes is expressed in the liver organoids at a similarly shut down level compared to the corresponding gene in adult liver hepatocytes: Ccl2, Osmr, Icam1 and Cxcl2.

[0167] In some embodiments, one or both of the following genes is differentially expressed in both a liver organoid and newborn liver: mKi67 and cdkn3.

[0168] In some embodiments, one, two or all of the following genes are expressed at a similar level in a liver organoid and a newborn liver: cyp2j6, olfm4 and Lefty 1. For example, see Figure 16B.

[0169] In some embodiments, a liver organoid of the invention has a ductal phenotype when cultured in expansion medium of the invention (e.g. EM1 or EM2).

[0170] In some embodiments, a liver organoid of the invention expresses adult liver markers when cultured in a differentiation medium of the invention.

[0171] In one embodiment, a liver organoid of the invention has a gene expression profile as shown in Figure 16C.

[0172] In a particularly preferred embodiment, a mouse liver cell population or organoid of the invention has the gene expression profile as shown in Figure 18. For example, in one preferred embodiment, a mouse liver cell population or organoid of the invention:

1. a) expresses at least one (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11), preferably all of the following stem cell markers: lgr5, lgr4, epcam, Cd44, Tnfrsf19, Sox9, Sp5, Cd24a, Prom1, Cdca7 and Elf3; and/or
2. b) does not express the following stem cell marker: lgr6; and/or
3. c) expresses at least one (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19), preferably all of the following hepatocyte or cholangiocyte markers when grown in expansion medium of the invention: Hnf1a, Hnf1b, Hnf4a, Hhex, Onecut1, Onecut2, Prox1, Cdh1, Foxa2, Gata6, Foxm1, Cebpa, Cebpb, Cebp, Cebp, Glul, Krt7, Krt19 and Met; and/or
4. d) does not express at least one (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17) of the following genes when grown in expansion medium of the invention: afp, Ins1, Ins2, Gcg, Ptfla, Cela1, Cela2a, Cela3b, Neurod1, Neurod2, Neurog1, Neurog2, Neurog3, Amy2a4, Igflr, Igf2 and Cd34; and/or
5. e) expresses at least one (e.g. 1, 2 or 3) of the following reprogramming genes: Klf4, Myc and Pou5f1 and/or
6. f) does not express the following reprogramming gene: Sox2.

wherein the expression of the genes is preferably detected by measuring expression at the mRNA level, for example, using a microarray.

[0173] More preferably a mouse liver cell population or organoid of the invention has all of features a) to f) above.

[0174] In some embodiments, the gene expression profile described above for a mouse cell population or organoid of the invention is for a mouse cell population or organoid cultured in expansion medium of the invention.

[0175] In some embodiments, there is provided a human liver cell population or organoid of the invention that has the gene expression signature shown in Figure 19. For example, a human liver cell population or organoid cultured in EM1 of the invention preferably expresses the genes indicated in Figure 19 as being expressed in EM1 cell culture medium. For example, a human liver cell population or organoid cultured in EM2 of the invention preferably expresses the genes indicated in Figure 19 as being expressed in EM2 cell culture medium. For example, a human liver cell population or organoid cultured in DM preferably expresses the genes indicated in Figure 19 as being expressed in DM cell culture medium.

[0176] For example, in one preferred embodiment, a human liver cell population or organoid of the invention:

1. a) expresses at least one (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9), preferably all of the following stem cell signature genes: LGR4, TACSTD1/Epcam, CD44, SOX9, SP5, CD24, PROM1, CDCA7 and ELF3; and/or
2. b) expresses at least one (e.g. 1, 2, 3, 4), preferably all of the following reprogramming genes: KLF4, MYC, POU5F1 and SOX2; and/or
3. c) expresses at least one (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19), preferably all of the following hepatocyte/cholangiocyte specific genes: HNF1A, HNF1B, HNF4A, HHEX, ONECUT1, ONECUT2, PROX1, CDH1, FOXA2, GATA6, FOXM1, CEBPA, CEBPB, CEBPD, CEBPG, GLUL, KRT7, KRT19 and MET; and/or
4. d) does not express at least one (e.g. 1, 2, 3, 4, 5, 6), preferably all of the following hepatocyte/cholangiocyte specific genes: NEUROG2, IGF1R and CD34, AFP, GCG and PTF1A, for example, it does not express NEUROG2, IGF1R and CD34; and/or
5. e) expresses at least one (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18), preferably all of the following hepatocyte specific genes: TTR, ALB, FAH, TAT, CYP3A7, APOA1, HMGCS1, PPARG, CYP2B6, CYP2C18, CYP2C9, CYP2J2, CYP3A4, CYP3A5, CYP3A7, CYP4F8, CYP4V2 and SCARB1;

wherein the expression of the genes is preferably detected by measuring expression at the mRNA level, for example, using a microarray.

[0177] More preferably a human liver cell population or organoid of the invention has all of features a) to e) above.

[0178] In some embodiments, the genes in a human liver cell population or organoid of the invention are upregulated or downregulated relative to expression of a reference RNA as shown in Figure 19. Preferably, the reference RNA is Universal Human Reference RNA (Stratagene, Catalog #740000). In some embodiments, a gene is upregulated or downregulated relative to the reference RNA if it is also shown in Figure 19 as being upregulated or downregulated relative to the reference RNA but the extent of upregulation or downregulation need not be the same. In other embodiments, the extent of upregulation or downregulation is $\pm 35\%$, $\pm 30\%$, $\pm 25\%$, $\pm 20\%$, $\pm 20\%$, $\pm 15\%$, $\pm 10\%$, $\pm 5\%$, ± 3 or approximately the same as shown in Figure 19. In other embodiments, the absolute level of expression of the genes in a human organoid of the invention is $\pm 35\%$, $\pm 30\%$, $\pm 25\%$, $\pm 20\%$, $\pm 15\%$, $\pm 10\%$, $\pm 5\%$, ± 3 or approximately the same as shown in Figure 19.

[0179] The human liver cell population or organoids of the invention also preferably express Lgr5 and/or Tnfrsf19, preferably both. In some embodiments, the human liver cell population or organoids, when cultured in expansion medium of the invention express Lgr5 and/or Tnfrsf19, preferably both. Preferably, expression of Lgr5 and/or Tnfrsf19 is detected by RT PCR. In some embodiments, Lgr5 and/or Tnfrsf19 are present at much lower levels of expression in organoids or cells when cultured in the differentiation medium compared to their level of expression organoids or cells when cultured in the expansion medium (for example at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, at least 15-fold lower).

[0180] Cells and organoids according to the present invention may preferably be capable of secreting albumin, for example, at a rate of between approximately $1\mu\text{g}$ per hour per 10^6 cells and $10\mu\text{g}$ per hour per 10^6 cells, preferably between $2\mu\text{g}$ and $6\mu\text{g}$ per hour per 10^6 cells.

[0181] Furthermore, such cells and organoids may secrete urea. For example, in a 35mm dish of cells, the activity of urea synthesis may be between $1\mu\text{g}$ and $50\mu\text{g}$ in 48 hours, preferably between $5\mu\text{g}$ and $30\mu\text{g}$.

[0182] Cells and organoids according to the invention may show visible glycogen stores, for example, when stained. The capacity for cells and organoids according to the invention to synthesize glycogen actively can be tested by switching the culture media from low-glucose differentiation media to high-glucose DMEM supplemented with 10% FBS and $0.2\mu\text{M}$ dexamethasone for two days.

[0183] Cells and organoids according to the invention may possess inducible cytochrome P450 activity (e.g. CYP1A). Such activity may be tested, for example, using an ethoxyresorufin-O-deethylase (EROD) assay (Cancer Res, 2001, 61: 8164-8170). For example, cells or organoids may be exposed to a P450 substrate such as 3-methylcholanthrene and the levels of EROD activity compared to control cells.

[0184] Morphologically, the cells appear hepatocyte-like.

[0185] A preferred liver organoid comprises or consists of a cystic structure with on the outside a layer of cells with buds and a central lumen as depicted in Figure 2. This liver organoid may have one or more (e.g. 2, 3, or all 4) of the following characteristics: (a) having a cell density of $>5 \times 10^5$ cells/cm³, preferably $>10 \times 10^5$ cells/cm³; (b) having a thickness equivalent to 2-30 layers of cells, preferably a thickness equivalent to 2-15 layers of cells; (c) the cells mutually contact in three dimensions, (d) demonstrate a function inherent to healthy liver tissue, (e) have an elongated shape, with 2 defined domains, i.e. a single layered epithelial domain where highly polarized cells are detected and keratin markers are expressed (this domain resembles the bile duct domain) and the other domain constitutes the main body of the organoid and is formed by a multilayered epithelia with non-polarized cells wherein albumin expression may be detected. It is clear to the skilled person that such a liver organoid is preferably not a liver fragment and/or does not comprise a blood vessel, and/or does not comprise a liver lobule or a bile duct.

[0186] Within the context of the invention, a liver fragment is a part of an adult liver, preferably a human adult liver. Preferably a liver organoid as identified herein is therefore not a liver fragment. A liver organoid is preferably obtained using a cell from an adult liver, preferably an epithelial stem cell from an adult liver, more preferably an epithelial stem cell from an adult liver expressing Lgr5.

[0187] In some embodiments, a liver organoid comprises cells that express Lgr5. For example, in some embodiments, at least 2%, more preferably at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the cells in the liver organoid express Lgr5. Similarly, the invention provides a cell or a population of cells which express Lgr5, wherein said cells are obtained from a liver organoid of the invention. The progeny of such cells is also encompassed by the invention.

[0188] In an embodiment, a liver organoid is a liver organoid which is still being cultured using a method of the invention and is therefore in contact with an extracellular matrix. Preferably, a liver organoid is embedded in a non-mesenchymal extracellular matrix. Within the context of the invention, "in contact" means a physical or mechanical or chemical contact, which means that for separating said liver organoid from said extracellular matrix a force needs to be used.

[0189] In a preferred embodiment, a liver organoid could be cultured during at least 2, 3, 4, 5, 6, 7, 8, 9, 10 weeks or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 months or longer.

[0190] In another preferred embodiment, a liver organoid originates from a single cell, preferably expressing Lgr5, more preferably wherein the single cell comprises a nucleic acid construct comprising a nucleic acid molecule of interest.

[0191] The invention further provides the use of a culture medium according to the invention

for culturing epithelial stem cells or isolated organoid structures that comprise these stem cells on an extracellular matrix, whereby said stem cells preferably do not comprise human embryonic stem cells. Preferred are human adult stem cells. Furthermore, single sorted epithelial stem cells from the liver are also able to initiate these 3 dimensional organoids in a culture medium according to the invention. The invention further provides the use of a culture medium according to the invention for culturing liver fragments comprising stem cells.

[0192] It is preferred that said stem cells are liver stem cells, or epithelial stem cells. A culture medium according to the invention allowed the establishment of long-term culture conditions under which a liver organoid is formed in which all differentiated cell types are present. Using a culture method according to the invention allowed culture periods of at least seven months, at least eight months, at least nine months, at least ten months.

[0193] The invention further provides a liver organoid, preferably comprising at least 50% viable cells, more preferred at least 60% viable cells, more preferred at least 70% viable cells, more preferred at least 80% viable cells, more preferred at least 90% viable cells. Viability of cells may be assessed using Hoechst staining or Propidium Iodide staining in FACS.

[0194] The viable cells preferably possess hepatic functions, or characteristics of hepatocytes, as described above.

Uses of Cells and Organoids of the Invention

[0195] In a further aspect, the invention provides the use of a liver cell or organoid according to the invention as described above in a drug discovery screen, toxicity assay or in regenerative medicine. The invention furthermore provides the use of the progeny of liver organoids of the invention, in toxicity assays. Such toxicity assays may be *in vitro* assays using a cell derived from a liver organoid or a liver organoid or part thereof. Such progeny and liver organoids are easy to culture and more closely resemble primary epithelial cells than, for example, epithelial cell lines such as Caco-2 (ATCC HTB-37), I-407 (ATCC CCL6), and XBF (ATCC CRL 8808) which are currently used in toxicity assays. It is anticipated that toxicity results obtained with liver organoids more closely resemble results obtained in patients. A cell-based toxicity test is used for determining organ specific cytotoxicity. Compounds that are tested in said test comprise cancer chemopreventive agents, environmental chemicals, food supplements, and potential toxicants. The cells are exposed to multiple concentrations of a test agent for certain period of time. The concentration ranges for test agents in the assay are determined in a preliminary assay using an exposure of five days and log dilutions from the highest soluble concentration. At the end of the exposure period, the cultures are evaluated for inhibition of growth. Data are analyzed to determine the concentration that inhibited end point by 50 percent (TC50).

[0196] For example, induction of cytochrome P450 enzymes in liver hepatocytes is a key factor that determines the efficacy and toxicity of drugs. In particular, induction of P450s is an

important mechanism of troublesome drug-drug interactions, and it is also an important factor that limits drug efficacy and governs drug toxicity. Cytochrome P450 induction assays have been difficult to develop, because they require intact normal human hepatocytes. These cells have proven intractable to production in numbers sufficient to sustain mass production of high throughput assays.

[0197] For example, according to this aspect of the invention, a candidate compound may be contacted with cell or organoid as described herein, and any change to the cells or in to activity of the cells may be monitored. Examples of other non-therapeutic uses of the cells or organoids of the present invention include research of liver embryology, liver cell lineages, and differentiation pathways; gene expression studies including recombinant gene expression; mechanisms involved in liver injury and repair; research of inflammatory and infectious diseases of the liver; studies of pathogenetic mechanisms; and studies of mechanisms of liver cell transformation and aetiology of liver cancer.

[0198] For high-throughput purposes, said liver organoids are cultured in multiwell plates such as, for example, 96 well plates or 384 well plates. Libraries of molecules are used to identify a molecule that affects said organoids. Preferred libraries comprise antibody fragment libraries, peptide phage display libraries, peptide libraries (e.g. LOPAP™, Sigma Aldrich), lipid libraries (BioMol), synthetic compound libraries (e.g. LOP AC™, Sigma Aldrich) or natural compound libraries (Specs, TimTec). Furthermore, genetic libraries can be used that induce or repress the expression of one of more genes in the progeny of the adenoma cells. These genetic libraries comprise cDNA libraries, antisense libraries, and siRNA or other non-coding RNA libraries. The cells are preferably exposed to multiple concentrations of a test agent for certain period of time. At the end of the exposure period, the cultures are evaluated. The term "affecting" is used to cover any change in a cell, including, but not limited to, a reduction in, or loss of, proliferation, a morphological change, and cell death. Said liver organoids can also be used to identify drugs that specifically target epithelial carcinoma cells, but not said liver organoids.

[0199] Liver organoids according to the invention can further replace the use of cell lines such as Caco-2 cells in toxicity assays of potential novel drugs or of known or novel food supplements.

[0200] Furthermore, such liver organoids can be used for culturing of a pathogen.

[0201] Cultures comprising liver organoids are useful in regenerative medicine, for example in post-radiation and/or post-surgery repair of the liver epithelium, in the repair of the epithelium in patients suffering from chronic or acute liver failure or disease. Liver diseases include, but are not limited to Hepatocellular Carcinoma, Alagille Syndrome, Alpha- 1- Antitrypsin Deficiency, Autoimmune Hepatitis, Biliary Atresia, Chronic Hepatitis, Cancer of the Liver , Cirrhosis Liver Cysts Fatty Liver, Galactosemia Gilbert's Syndrome, Primary Biliary Cirrhosis, Hepatitis A, Hepatitis B, Hepatitis C, Primary Sclerosing Cholangitis, Reye's Syndrome, Sarcoidosis, Tyrosinemia, Type I Glycogen Storage Disease, Wilson's Disease, Neonatal

Hepatitis, Non-alcoholic SteatoHepatitis, Porphyrin, and Hemochromatosis.

[0202] Genetic conditions that lead to liver failure could benefit from cell-based therapy in the form of partial or full cell replacement using cells cultured according to the media and/or methods of the invention. A non-limiting list of genetic conditions that lead to liver failure includes: Progressive familial intrahepatic cholestasis, Glycogen storage disease type III, Tyrosinemia, Deoxyguanosine kinase deficiency, Pyruvate carboxylase deficiency, Congenital dyserythropoietic anemia, Polycystic Liver Disease Polycystic Kidney Disease, Alpha-1 antitrypsin deficiency, Urea cycle defects, Organic acidemia, lysosomal storage diseases, and Fatty Acid Oxidation Disorders. Other conditions that may also benefit from cell-based therapy include Wilson's Disease and Hereditary Amyloidosis (FAP).

[0203] Other non-hepatocyte related causes of liver failure that would require a full liver transplant to reach full therapeutic effect, may still benefit from some temporary restoration of function using cell-based therapy using cells cultured according to the media and/or methods of the invention. A non-limiting list of examples of such conditions includes: Primary Biliary Cirrhosis, Primary Sclerosing Cholangitis, Agalactemia syndrome, Homozygous Familial hypercholesterolemia, Hepatitis B with cirrhosis, Hepatitis C with cirrhosis, Budd-Chiari syndrome, Primary hyperoxaluria, Autoimmune Hepatitis, and Alcoholic liver disease.

[0204] The liver organoids of the invention may be used in a method of treating a hereditary disease that involves malfunctioning hepatocytes. Such diseases may be early onset or late onset. Early onset disease include metabolite related organ failure (e.g. alpha-1-antitrypsin deficiency), glycogen storage diseases (e.g. GSD II, Pompe's disease), tyrosinemia, mild DGUOK, CDA type I, Urea cycle defects (e.g. OTC deficiency), organic acidemia and fatty acid oxidation disorders. Late onset diseases include primary hyperoxaluria, familial hypercholesterolemia, Wilson's disease, Hereditary Amyloidosis and Polycystic liver disease. Partial or full replacement with healthy hepatocytes arising from liver organoids of the invention may be used to restore liver function or to postpone liver failure.

[0205] The liver organoids of the invention may be used in a method of treating chronic liver failure arising due to hereditary metabolic disease or as a result of hepatocyte infection. Treatment of a hereditary metabolic disease may involve administration of genetically modified autologous liver organoids of the invention. Treatment of hepatocyte infections may involve administration of allogeneic liver organoids of the invention. In some embodiments, the liver organoids are administered over a period of 2-3 months.

[0206] The liver organoids of the invention may be used to treat acute liver failure, for example, as a result of liver intoxication which may result from use of paracetamol, medication or alcohol. In some embodiments, the therapy to restore liver function will comprise injecting hepatocyte suspension from frozen, ready to use allogeneic hepatocytes obtained from organoids of the invention. The ability to freeze suitable organoids means that the organoids can be available for immediate delivery and so it is not necessary to wait for a blood transfusion.

[0207] In the case of replacement or correction of deficient liver function, it may be possible to construct a cell-matrix structure from one or more liver organoids generated according to the present invention. It is thought that only about 10% of hepatic cell mass is necessary for adequate function. This makes implantation of organoid unit compositions into children especially preferable to whole organ transplantation, due to the relatively limited availability of donors and smaller size of juvenile organs. For example, an 8-month-old child has a normal liver that weighs approximately 250g. That child would therefore need about 25g of tissue. An adult liver weighs approximately 1500g; therefore, the required implant would only be about 1.5% of the adult liver. When organoid units according to the invention are implanted, optionally attached to a polymer scaffold, proliferation in the new host will occur, and the resulting hepatic cell mass replaces the deficient host function. The inventors have shown, for the first time, that it is possible to generate mature hepatocytes from adult liver stem cells or liver tissue fragments comprising stem cells that are suitable for transplantation into non-human animals or humans. Using the first culture medium according to the invention, the inventors have demonstrated that it is possible to maintain and expand a population of liver stem cells. Using the second culture medium according to the invention, the inventors have shown that hepatoblasts can be differentiated *in vivo* to mature hepatocytes suitable for transplantation purposes. Hence, the inventors provide a new source of hepatocytes for liver regeneration, replacement or correction of deficient liver function.

[0208] The inventors have also demonstrated successful transplantation of the organoids, grown by methods of the present invention, into immunodeficient mice (see example 7), with transplanted organoid-derived cells generating both cholangiocytes and hepatocytes *in vivo*. Therefore, in one embodiment the invention provides organoids or organoid-derived cells of the invention for transplanting into human or animals.

[0209] The use of human liver organoids for transplantation purposes is advantageous over the use of fetal or adult hepatocytes for a number of reasons. Firstly, the culture methods of the invention provide unlimited expansion of cells and hence, an unlimited supply. In particular, the inventors have shown that under the correct culture conditions (e.g. using the expansion culture medium of the invention), that Lgr5⁺ cells can undergo more than 1000 divisions *in vitro*. Therefore, Lgr5⁺ cells can be extracted from the liver organoids and repassaged providing a continual self-renewing source of transplantable hepatocyte and cholangiocyte-generating cells. By contrast, fetal or adult hepatocytes are derived from donor livers which only provide a single round of transplantation. Furthermore, donor cells can only be kept alive for a few days but lose their hepatocyte properties. This means the transplants must be made as soon as the donor becomes available. Organoid-derived cells, on the other hand, retain their phenotype over multiple divisions and over prolonged periods of time meaning that they are ready and available for transplantation at any stage. This could also allow the organoid-derived cells to be used as a temporary liver treatment to extend the lifespan of patients for patients on the waiting list for liver transplants. A further advantage of the liver organoids of the invention is that they can be frozen and later be defrosted without loss of function. This enables cell banking, easy storage and rapid availability for acute use. This could be useful for

example, in the preparation of an "off-the-shelf" product that might be used for the treatment of acute liver toxicity. Organoids can also be grown from cells or tissue fragments taken as small biopsies from live donors minimising any ethical objections to the treatment. The donor may even be from the patient that is to be treated, which could reduce any negative side-effects associated with transplantation of foreign cells and organs and reduce the need for immunosuppressive drugs.

[0210] Accordingly, included within the scope of the invention are organoids for use in methods of treatment of a human or animal patient through cellular therapy. The term "animal" here denotes all mammalian animals, preferably human patients. It also includes an individual animal in all stages of development, including embryonic and foetal stages. For example, the patient may be an adult, or the therapy may be for pediatric use (e.g. newborn, child or adolescent). Such cellular therapy encompasses the administration of cells or organoids generated according to the invention to a patient through any appropriate means. Specifically, such methods of treatment involve the regeneration of damaged tissue. The term "administration" as used herein refers to well recognized forms of administration, such as intravenous or injection, as well as to administration by transplantation, for example transplantation by surgery, grafting or transplantation of tissue engineered liver derived from cells or organoids according to the present invention. In the case of cells, systemic administration to an individual may be possible, for example, by infusion into the superior mesenteric artery, the celiac artery, the subclavian vein via the thoracic duct, infusion into the heart via the superior vena cava, or infusion into the peritoneal cavity with subsequent migration of cells via subdiaphragmatic lymphatics, or directly into liver sites via infusion into the hepatic arterial blood supply or into the portal vein.

[0211] Between 10^4 and 10^{13} cells per 100 kg person may be administered per infusion. Preferably, between about $1\text{-}5 \times 10^4$ and $1\text{-}5 \times 10^7$ cells may be infused intravenously per 100 kg person. More preferably, between about 1×10^4 and 10×10^6 cells may be infused intravenously per 100 kg person. In some embodiments, a single administration of cells or organoids is provided. In other embodiments, multiple administrations are used. Multiple administrations can be provided over an initial treatment regime, for example, of 3-7 consecutive days, and then repeated at other times.

[0212] In some embodiments it is desirable to repopulate/replace 10-20% of a patient's liver with healthy hepatocytes arising from a liver organoid of the invention.

[0213] It is also possible to reconstitute a liver organoid from one single cell expressing Lgr5 as defined herein. This single cell may have been modified by introduction of a nucleic acid construct as defined herein, for example, to correct a genetic deficiency or mutation. It would also be possible to specifically ablate expression, as desired, for example, using siRNA. Potential polypeptides to be expressed could be any of those that are deficient in metabolic liver diseases, including, for example, AAT (alpha antitrypsin). For elucidating liver physiology, we might also express or inactivate genes implicated in the Wnt, EGF, FGF, BMP or notch

pathway. Also, for screening of drug toxicity, the expression or inactivation of genes responsible for liver drug metabolism (for example, genes in the CYP family) would be of high interest

[0214] In one embodiment, the expanded epithelial stem cells may be reprogrammed into related tissue fates such as, for example, liver cells including a hepatocyte and a cholangiocyte cell. Thus far, it has not been possible to regenerate liver cells from adult stem cells. The culturing methods of the present invention will enable to analyse for factors that trans-differentiate the closely related epithelial stem cell to a liver cell, including a hepatocyte and a cholangiocyte cell.

[0215] It will be clear to a skilled person that gene therapy can additionally be used in a method directed at repairing damaged or diseased tissue. Use can, for example, be made of an adenoviral or retroviral gene delivery vehicle to deliver genetic information, like DNA and/or RNA to stem cells. A skilled person can replace or repair particular genes targeted in gene therapy. For example, a normal gene may be inserted into a nonspecific location within the genome to replace a non functional gene. In another example, an abnormal gene sequence can be replaced for a normal gene sequence through homologous recombination. Alternatively, selective reverse mutation can return a gene to its normal function. A further example is altering the regulation (the degree to which a gene is turned on or off) of a particular gene. Preferably, the stem cells are ex vivo treated by a gene therapy approach and are subsequently transferred to the mammal, preferably a human being in need of treatment. For example, organoid-derived cells may be genetically modified in culture before transplantation into patients.

[0216] The inventors have found that Lgr5 is not detectable in healthy liver, although residual Lgr5 may be detected. Thus, the disclosure further describes a method of diagnosing liver injury comprising detecting whether Lgr5 is expressed, wherein the expression of Lgr5 protein indicates liver injury. The disclosure also describes a method of monitoring the repair or regeneration of the liver by monitoring the expression of Lgr5 in the liver. Lgr5 expression may be detected by any suitable method, for example, flow cytometry, immunohistochemistry or by use of PCR methods.

[0217] The invention also provides a composition or cell culture vessel comprising cells and/or organoids according to any one of the aspects of the invention described above, and a culture medium according to any one of the aspects of the invention described above. For example, such a composition or cell culture vessel may comprise any number of cells or organoids cultured according to a method of the invention, in a culture medium as described above. For example, a preferred culture medium comprises or consists of Advanced-DMEM/F12 supplemented with B27, N2, 200ng/ml N-Acetylcysteine, 50ng/ml EGF, 1 µg/ml R-spondin1, 10 nM gastrin, 100 ng/ml FGF10, 10mM Nicotinamide and 50ng/ml HGF and 50% Wnt conditioned media.

Definitions

[0218] A nucleic acid construct comprises a nucleic acid molecule of interest and will ensure expression of the given nucleic acid molecule in the cells wherein it had been introduced. A particularly preferred nucleic acid construct is an expression vector wherein a nucleic acid molecule encoding a polypeptide is operably linked to a promoter capable of directing expression of said nucleic acid molecule (i.e a coding sequence) in said cells. The phrase "expression vector" generally refers to a nucleic acid molecule that is capable of effecting expression of a gene/nucleic acid molecule it contains in a cell compatible with such sequences. These expression vectors typically include at least suitable promoter sequences and optionally, transcription termination signals. A nucleic acid or DNA or nucleotide sequence encoding a polypeptide is incorporated into a DNA/nucleic acid construct capable of introduction into and expression in an *in vitro* cell culture as identified in a method of the invention. A DNA construct prepared for introduction into a particular cell typically include a replication system recognized by said cell, an intended DNA segment encoding a desired polypeptide, and transcriptional and translational initiation and termination regulatory sequences operably linked to the polypeptide-encoding segment. A DNA segment is "operably linked" when it is placed into a functional relationship with another DNA segment. For example, a promoter or enhancer is operably linked to a coding sequence if it stimulates the transcription of the sequence. DNA for a signal sequence is operably linked to DNA encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of a polypeptide. Generally, a DNA sequence that is operably linked are contiguous, and, in the case of a signal sequence, both contiguous and in reading phase. However, enhancers need not be contiguous with a coding sequence whose transcription they control. Linking is accomplished by ligation at convenient restriction sites or at adapters or linkers inserted in lieu thereof.

[0219] The selection of an appropriate promoter sequence generally depends upon the host cell selected for the expression of a DNA segment. Examples of suitable promoter sequences include eukaryotic promoters well known in the art (see, e.g. Sambrook and Russell, 2001, *supra*). A transcriptional regulatory sequence typically includes a heterologous enhancer or promoter that is recognised by the cell. Suitable promoters include the CMV promoter. An expression vector includes the replication system and transcriptional and translational regulatory sequences together with the insertion site for the polypeptide encoding segment can be employed. Examples of workable combinations of cell lines and expression vectors are described in Sambrook and Russell (2001, *supra*) and in Metzger et al. (1988) Nature 334: 31-36.

[0220] The invention is not limited to a specific polypeptide or nucleic acid molecule of interest to be expressed in a cell expressing Lgr5 as identified herein. Depending on the aim of the method, one may envisage to express a polypeptide in a liver cell and/or to inactivate the expression of a polypeptide in a liver cell. Potential polypeptides to be expressed could be all of those that are deficient in metabolic liver diseases, as eg AAT (alpha antitrypsin). For elucidating the liver physiology, we might also express or inactivate genes implicated in the Wnt, EGF, FGF, BMP or notch pathway. Also, for screening of drug toxicity, the expression or

inactivation of genes responsible for liver drug metabolism (e.g. genes of the CYP family) would be of high interest.

[0221] Some aspects of the invention concern the use of a nucleic acid construct or expression vector comprising a nucleotide sequence as defined above, wherein the vector is a vector that is suitable for gene therapy. Vectors that are suitable for gene therapy are described in Anderson 1998, *Nature* 392: 25-30; Walther and Stein, 2000, *Drugs* 60: 249-71; Kay et al., 2001, *Nat. Med.* 7: 33-40; Russell, 2000, *J. Gen. Virol.* 81: 2573-604; Amado and Chen, 1999, *Science* 285: 674-6; Federico, 1999, *Curr. Opin. Biotechnol.* 10: 448-53; Vigna and Naldini, 2000, *J. Gene Med.* 2: 308-16; Marin et al., 1997, *Mol. Med. Today* 3: 396-403; Peng and Russell, 1999, *Curr. Opin. Biotechnol.* 10: 454-7; Sommerfelt, 1999, *J. Gen. Virol.* 80: 3049-64; Reiser, 2000, *Gene Ther.* 7: 910-3; and references cited therein. Examples include integrative and non-integrative vectors such as those based on retroviruses, adenoviruses (AdV), adeno-associated viruses (AAV), lentiviruses, pox viruses, alphaviruses, and herpes viruses.

[0222] A particularly suitable gene therapy vector includes an Adenoviral (Ad) and Adeno-associated virus (AAV) vector. These vectors infect a wide number of dividing and non-dividing cell types including liver cells. In addition adenoviral vectors are capable of high levels of transgene expression. However, because of the episomal nature of the adenoviral and AAV vectors after cell entry, these viral vectors are most suited for therapeutic applications requiring only transient expression of the transgene (Russell, 2000, *J. Gen. Virol.* 81: 2573-2604; Goncalves, 2005, *Virol J.* 2(1):43) as indicated above. Preferred adenoviral vectors are modified to reduce the host response as reviewed by Russell (2000, *supra*). Safety and efficacy of AAV gene transfer has been extensively studied in humans with encouraging results in the liver, muscle, CNS, and retina (Manno et al *Nat medicine* 2006, Stroes et al *ATVB* 2008, Kaplitt, Feigin, *Lancet* 2009; Maguire, Simonelli et al *NEJM* 2008; Bainbridge et al *NEJM* 2008). AAV2 is the best characterized serotype for gene transfer studies both in humans and experimental models. AAV2 presents natural tropism towards skeletal muscles, neurons, vascular smooth muscle cells and hepatocytes. AAV2 is therefore a good choice of vector to target liver tissues. Other examples of adeno-associated virus-based non integrative vectors include AAV1, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11 and pseudotyped AAV. The use of non-human serotypes, like AAV8 and AAV9, might be useful to overcome these immunological responses in subjects, and clinical trials have just commenced (ClinicalTrials.gov Identifier: NCT00979238). For gene transfer into a liver cell, an adenovirus serotype 5 or an AAV serotype 2, 7 or 8 have been shown to be effective vectors and therefore a preferred Ad or AAV serotype (Gao, *Molecular Therapy* (2006) 13, 77-87).

[0223] A preferred retroviral vector for application in the present invention is a lentiviral based expression construct. Lentiviral vectors have the unique ability to infect non-dividing cells (Amado and Chen, 1999 *Science* 285: 674-6). Methods for the construction and use of lentiviral based expression constructs are described in U.S. Patent No.'s 6,165,782, 6,207,455, 6,218,181, 6,277,633 and 6,323,031 and in Federico (1999, *Curr Opin Biotechnol* 10: 448-53) and Vigna et al. (2000, *J Gene Med* 2000; 2: 308-16).

[0224] Generally, gene therapy vectors will be as the expression vectors described above in the sense that they comprise a nucleotide sequence encoding a polypeptide of the invention to be expressed, whereby a nucleotide sequence is operably linked to the appropriate regulatory sequences as indicated above. Such regulatory sequence will at least comprise a promoter sequence. Suitable promoters for expression of a nucleotide sequence encoding a polypeptide from gene therapy vectors include e.g. cytomegalovirus (CMV) intermediate early promoter, viral long terminal repeat promoters (LTRs), such as those from murine moloney leukaemia virus (MMLV) rous sarcoma virus, or HTLV-1, the simian virus 40 (SV 40) early promoter and the herpes simplex virus thymidine kinase promoter. Suitable promoters are described below.

[0225] Several inducible promoter systems have been described that may be induced by the administration of small organic or inorganic compounds. Such inducible promoters include those controlled by heavy metals, such as the metallothionein promoter (Brinster et al. 1982 *Nature* 296: 39-42; Mayo et al. 1982 *Cell* 29: 99-108), RU-486 (a progesterone antagonist) (Wang et al. 1994 *Proc. Natl. Acad. Sci. USA* 91: 8180-8184), steroids (Mader and White, 1993 *Proc. Natl. Acad. Sci. USA* 90: 5603-5607), tetracycline (Gossen and Bujard 1992 *Proc. Natl. Acad. Sci. USA* 89: 5547-5551; U.S. Pat. No. 5,464,758; Furth et al. 1994 *Proc. Natl. Acad. Sci. USA* 91: 9302-9306; Howe et al. 1995 *J. Biol. Chem.* 270: 14168-14174; Resnitzky et al. 1994 *Mol. Cell. Biol.* 14: 1669-1679; Shockett et al. 1995 *Proc. Natl. Acad. Sci. USA* 92: 6522-6526) and the tTAER system that is based on the multi-chimeric transactivator composed of a tetR polypeptide, as activation domain of VP16, and a ligand binding domain of an estrogen receptor (Yee et al., 2002, US 6,432,705).

[0226] Suitable promoters for nucleotide sequences encoding small RNAs for knock down of specific genes by RNA interference (see below) include, in addition to the above mentioned polymerase II promoters, polymerase III promoters. The RNA polymerase III (pol III) is responsible for the synthesis of a large variety of small nuclear and cytoplasmic non-coding RNAs including 5S, U6, adenovirus VA1, Vault, telomerase RNA, and tRNAs. The promoter structures of a large number of genes encoding these RNAs have been determined and it has been found that RNA pol III promoters fall into three types of structures (for a review see Geiduschek and Tocchini-Valentini, 1988 *Annu. Rev. Biochem.* 57: 873-914; Willis, 1993 *Eur. J. Biochem.* 212: 1-11; Hernandez, 2001, *J. Biol. Chem.* 276: 26733-36). Particularly suitable for expression of siRNAs are the type 3 of the RNA pol III promoters, whereby transcription is driven by cis-acting elements found only in the 5'-flanking region, i.e. upstream of the transcription start site. Upstream sequence elements include a traditional TATA box (Mattaj et al., 1988 *Cell* 55, 435-442), proximal sequence element and a distal sequence element (DSE; Gupta and Reddy, 1991 *Nucleic Acids Res.* 19, 2073-2075). Examples of genes under the control of the type 3 pol III promoter are U6 small nuclear RNA (U6 snRNA), 7SK, Y, MRP, HI and telomerase RNA genes (see e.g. Myslinski et al., 2001, *Nucl. Acids Res.* 21: 2502-09).

[0227] A gene therapy vector may optionally comprise a second or one or more further nucleotide sequence coding for a second or further polypeptide. A second or further polypeptide may be a (selectable) marker polypeptide that allows for the identification, selection and/or screening for cells containing the expression construct. Suitable marker

proteins for this purpose are e.g. the fluorescent protein GFP, and the selectable marker genes HSV thymidine kinase (for selection on HAT medium), bacterial hygromycin B phosphotransferase (for selection on hygromycin B), Tn5 aminoglycoside phosphotransferase (for selection on G418), and dihydrofolate reductase (DHFR) (for selection on methotrexate), CD20, the low affinity nerve growth factor gene. Sources for obtaining these marker genes and methods for their use are provided in Sambrook and Russel (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, New York.

[0228] Alternatively, a second or further nucleotide sequence may encode a polypeptide that provides for fail-safe mechanism that allows a subject from the transgenic cells to be cured, if deemed necessary. Such a nucleotide sequence, often referred to as a suicide gene, encodes a polypeptide that is capable of converting a prodrug into a toxic substance that is capable of killing the transgenic cells in which the polypeptide is expressed. Suitable examples of such suicide genes include e.g. the *E.coli* cytosine deaminase gene or one of the thymidine kinase genes from Herpes Simplex Virus, Cytomegalovirus and Varicella-Zoster virus, in which case ganciclovir may be used as prodrug to kill the IL-10 transgenic cells in the subject (see e.g. Clair et al., 1987, Antimicrob. Agents Chemother. 31: 844-849).

[0229] For knock down of expression of a specific polypeptide, a gene therapy vector or other expression construct is used for the expression of a desired nucleotide sequence that preferably encodes an RNAi agent, i.e. an RNA molecule that is capable of RNA interference or that is part of an RNA molecule that is capable of RNA interference. Such RNA molecules are referred to as siRNA (short interfering RNA, including e.g. a short hairpin RNA). Alternatively, a siRNA molecule may directly, e.g. in a pharmaceutical composition that is administered within or in the neighborhood of a cell expressing Lgr5, or of a liver cell (i.e. hepatocyte or cholangiocyte) or of a liver organoid.

[0230] A desired nucleotide sequence comprises an antisense code DNA coding for the antisense RNA directed against a region of the target gene mRNA, and/or a sense code DNA coding for the sense RNA directed against the same region of the target gene mRNA. In a DNA construct of the invention, an antisense and sense code DNAs are operably linked to one or more promoters as herein defined above that are capable of expressing an antisense and sense RNAs, respectively. "siRNA" preferably means a small interfering RNA that is a short-length double-stranded RNA that are not toxic in mammalian cells (Elbashir et al., 2001, Nature 411: 494-98; Caplen et al., 2001, Proc. Natl. Acad. Sci. USA 98: 9742-47). The length is not necessarily limited to 21 to 23 nucleotides. There is no particular limitation in the length of siRNA as long as it does not show toxicity. "siRNAs" can be, e.g. at least 15, 18 or 21 nucleotides and up to 25, 30, 35 or 49 nucleotides long. Alternatively, the double-stranded RNA portion of a final transcription product of siRNA to be expressed can be, e.g. at least 15, 18 or 21 nucleotides and up to 25, 30, 35 or 49 nucleotides long.

[0231] "Antisense RNA" is preferably an RNA strand having a sequence complementary to a target gene mRNA, and thought to induce RNAi by binding to the target gene mRNA. "Sense

RNA" has a sequence complementary to the antisense RNA, and annealed to its complementary antisense RNA to form siRNA. The term "target gene" in this context preferably refers to a gene whose expression is to be silenced due to siRNA to be expressed by the present system, and can be arbitrarily selected. As this target gene, for example, genes whose sequences are known but whose functions remain to be elucidated, and genes whose expressions are thought to be causative of diseases are preferably selected. A target gene may be one whose genome sequence has not been fully elucidated, as long as a partial sequence of mRNA of the gene having at least 15 nucleotides or more, which is a length capable of binding to one of the strands (antisense RNA strand) of siRNA, has been determined. Therefore, genes, expressed sequence tags (ESTs) and portions of mRNA, of which some sequence (preferably at least 15 nucleotides) has been elucidated, may be selected as the "target gene" even if their full length sequences have not been determined.

[0232] The double-stranded RNA portions of siRNAs in which two RNA strands pair up are not limited to the completely paired ones, and may contain nonpairing portions due to mismatch (the corresponding nucleotides are not complementary), bulge (lacking in the corresponding complementary nucleotide on one strand), and the like. A non-pairing portions can be contained to the extent that they do not interfere with siRNA formation. The "bulge" used herein preferably comprise 1 to 2 non-pairing nucleotides, and the double-stranded RNA region of siRNAs in which two RNA strands pair up contains preferably 1 to 7, more preferably 1 to 5 bulges. In addition, the "mismatch" used herein is preferably contained in the double-stranded RNA region of siRNAs in which two RNA strands pair up, preferably 1 to 7, more preferably 1 to 5, in number. In a preferable mismatch, one of the nucleotides is guanine, and the other is uracil. Such a mismatch is due to a mutation from C to T, G to A, or mixtures thereof in DNA coding for sense RNA, but not particularly limited to them. Furthermore, a double-stranded RNA region of siRNAs in which two RNA strands pair up may contain both bulge and mismatched, which sum up to, preferably 1 to 7, more preferably 1 to 5 in number. Such non-pairing portions (mismatches or bulges, etc.) can suppress the below-described recombination between antisense and sense code DNAs and make the siRNA expression system as described below stable. Furthermore, although it is difficult to sequence stem loop DNA containing no non-pairing portion in the double-stranded RNA region of siRNAs in which two RNA strands pair up, the sequencing is enabled by introducing mismatches or bulges as described above. Moreover, siRNAs containing mismatches or bulges in the pairing double-stranded RNA region have the advantage of being stable in *E. coli* or animal cells.

[0233] The terminal structure of siRNA may be either blunt or cohesive (overhanging) as long as siRNA enables to silence the target gene expression due to its RNAi effect. The cohesive (overhanging) end structure is not limited only to the 3' overhang, and the 5' overhanging structure may be included as long as it is capable of inducing the RNAi effect. In addition, the number of overhanging nucleotide is not limited to the already reported 2 or 3, but can be any numbers as long as the overhang is capable of inducing the RNAi effect. For example, the overhang consists of 1 to 8, preferably 2 to 4 nucleotides. Herein, the total length of siRNA having cohesive end structure is expressed as the sum of the length of the paired double-stranded portion and that of a pair comprising overhanging single-strands at both ends. For

example, in the case of 19 bp double-stranded RNA portion with 4 nucleotide overhangs at both ends, the total length is expressed as 23 bp. Furthermore, since this overhanging sequence has low specificity to a target gene, it is not necessarily complementary (antisense) or identical (sense) to the target gene sequence. Furthermore, as long as siRNA is able to maintain its gene silencing effect on the target gene, siRNA may contain a low molecular weight RNA (which may be a natural RNA molecule such as tRNA, rRNA or viral RNA, or an artificial RNA molecule), for example, in the overhanging portion at its one end.

[0234] In addition, the terminal structure of the "siRNA" is necessarily the cut off structure at both ends as described above, and may have a stem-loop structure in which ends of one side of double-stranded RNA are connected by a linker RNA (a "shRNA"). The length of the double-stranded RNA region (stem-loop portion) can be, e.g. at least 15, 18 or 21 nucleotides and up to 25, 30, 35 or 49 nucleotides long. Alternatively, the length of the double-stranded RNA region that is a final transcription product of siRNAs to be expressed is, e.g. at least 15, 18 or 21 nucleotides and up to 25, 30, 35 or 49 nucleotides long. Furthermore, there is no particular limitation in the length of the linker as long as it has a length so as not to hinder the pairing of the stem portion. For example, for stable pairing of the stem portion and suppression of the recombination between DNAs coding for the portion, the linker portion may have a clover-leaf tRNA structure. Even though the linker has a length that hinders pairing of the stem portion, it is possible, for example, to construct the linker portion to include introns so that the introns are excised during processing of precursor RNA into mature RNA, thereby allowing pairing of the stem portion. In the case of a stem-loop siRNA, either end (head or tail) of RNA with no loop structure may have a low molecular weight RNA. As described above, this low molecular weight RNA may be a natural RNA molecule such as tRNA, rRNA, snRNA or viral RNA, or an artificial RNA molecule.

[0235] To express antisense and sense RNAs from the antisense and sense code DNAs respectively, a DNA construct comprise a promoter as defined above. The number and the location of the promoter in the construct can in principle be arbitrarily selected as long as it is capable of expressing antisense and sense code DNAs. As a simple example of a DNA construct, a tandem expression system can be formed, in which a promoter is located upstream of both antisense and sense code DNAs. This tandem expression system is capable of producing siRNAs having the aforementioned cut off structure on both ends. In the stem-loop siRNA expression system (stem expression system), antisense and sense code DNAs are arranged in the opposite direction, and these DNAs are connected via a linker DNA to construct a unit. A promoter is linked to one side of this unit to construct a stem-loop siRNA expression system. Herein, there is no particular limitation in the length and sequence of the linker DNA, which may have any length and sequence as long as its sequence is not the termination sequence, and its length and sequence do not hinder the stem portion pairing during the mature RNA production as described above. As an example, DNA coding for the above-mentioned tRNA and such can be used as a linker DNA.

[0236] In both cases of tandem and stem-loop expression systems, the 5' end may be have a sequence capable of promoting the transcription from the promoter. More specifically, in the

case of tandem siRNA, the efficiency of siRNA production may be improved by adding a sequence capable of promoting the transcription from the promoters at the 5' ends of antisense and sense code DNAs. In the case of stem-loop siRNA, such a sequence can be added at the 5' end of the above-described unit. A transcript from such a sequence may be used in a state of being attached to siRNA as long as the target gene silencing by siRNA is not hindered. If this state hinders the gene silencing, it is preferable to perform trimming of the transcript using a trimming means (for example, ribozyme as are known in the art). It will be clear to the skilled person that an antisense and sense RNAs may be expressed in the same vector or in different vectors. To avoid the addition of excess sequences downstream of the sense and antisense RNAs, it is preferred to place a terminator of transcription at the 3' ends of the respective strands (strands coding for antisense and sense RNAs). The terminator may be a sequence of four or more consecutive adenine (A) nucleotides.

[0237] In this document and in its claims, the verb "to comprise" and its conjugations is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded. In addition the verb "to consist" may be replaced by "to consist essentially of" meaning that a product as defined herein may comprise additional component(s) than the ones specifically identified, said additional component(s) not altering the unique characteristic of the invention. In addition a method as defined herein may comprise additional step(s) than the ones specifically identified, said additional step(s) not altering the unique characteristic of the invention. In addition, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there be one and only one of the elements. The indefinite article "a" or "an" thus usually means "at least one". The word "about" or "approximately" when used in association with a numerical value (about 10) preferably means that the value may be the given value of 10 more or less 1% of the value.

[0238] The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

DESCRIPTION OF FIGURES

[0239]

Fig. 1 Liver organoids growth factor requirement. **(A)** DIC images of liver organoids maintained with EGF (E), R-spondin 1 (R), Noggin (N), Wnt3A conditioned media (W) or the combination of them, supplemented with FGF10, HGF and Nicotinamide. **(B)** The number of organoids was counted weekly and passaged when required. Results are shown as mean $\pm$ SEM of 3 independent experiments. **(C)** gene expression analysis by RTPCR of Lgr5, Keratin 7 (K7) and Albumin (Alb) genes. **(D)** Isolated biliary ducts growing into organoids. Differential interference contrast images from the corresponding days after seeding. Magnification 10x (days 0, 1, 3 and 5). Days 15 on magnification 4x. Cultures were passage every 4-7 days by mechanical dissociation. Cultures have been grown at least for 8 months.

Fig. 2 Morphology of liver organoids. (A) Upper panels: paraffin section of a mouse liver showing the different domains (PT= portal triad, CV= central vein). Lower panels: Paraffin section of a liver organoid showing different domains b (single layered epithelia) and h (stratified epithelia) (B) Right pannel: Ecadherin staining in the liver organoids. Two different domains can be identified. Domain b, formed by a single layered epithelia that resembles the bile duct structures in the liver. This bile duct domain is formed by highly polarized cells that shows positive staining for pancytokeratin (PCK) (lower panel). Left panels show the presence of a second domain within the liver organoids. This h domain is formed by a stratified epithelia with non-polarized cells. The cells are organized around a central lumen and express the hepatocyte marker Alb. Magnification 10 x.

Fig. 3 Wnt signalling in the liver cultures. Lac Z expression was detected in cultures derived from Lgr5-LacZ or Axin2 LacZ mice. No positive staining was detected in liver cultures derived from a B16 mice. Magnification 4x, inset 20x.

Fig. 4 Expression of liver differentiating markers. (A) immunohistochemical and immunofluorescence anaylisis of the expression of the cholangiocyte marker keratin 7 (K7) and the hepatocyte markers keratin 8 (K8) and albumin (Alb). (B) analysis of the gene expression of hepatocyte markers: Albumin (Alb), transthyretrin (Ttr), Glutamine synthetase (Glul), glucose 6 phosphatase (G6P) and Cytocrome p450 isoform 3A11 (CYP3A11); and cholangiocyte markers keratin 7 (K7) and Keratin 19 (K19).

Fig. 5 Liver single cell cultures. (A) flow cytometry plot indicating the area of the sorted cells. (B) single cell growing into organoids at the time points indicated. Magnification 40 x (day 1-3), 10x (day 16), 4 x (day 21-on). (C & D) representative image of the colony formation efficiency of a Lgr5GFP single sorted cells. 100 cells were seeded in triplicate and colonies were counted 10 days later.

Fig. 6 Microarray analysis of the liver cultures. Analysis of the gene expression profile of adult liver tissue and liver organoid cultures maintained for 1 month in the ER or ER media supplemented with Noggin (ENR) or with noggin and Wnt (ENRW). The genetic profile was compared between the different samples and the genetic profile of Brow adipose tissue (BAT) white adipose tissue (WAT), muscle and new born liver. (A) hit map analysis showing that the cultures present a similar profile to the adult liver but a different profile to non-liver related tissues as muscle and BAT and WAT, (B) List of hepatocyte markers and cholangyocyte markers in the different conditions.

Figure 7: Mouse liver organoid culture shows stable karyotyping after long-term culture.

1. A - DIC images of liver organoids maintained in EGF (E) and R-spondin 1 (R), supplemented with FGF10, HGF and Nicotinamide (left figure, ER) or maintained in the same combination supplemented with Noggin (N) and Wnt3A conditioned media (W) (right figure, ENRW) for a period of 24 months.
2. B - Karyotype analysis of mouse liver organoids after 8 months in culture. Normal chromosomal counts (n=40, left panel figure) and polyploidy, a typical hepatocyte

feature, were found (n=80, right panel figure)

Figure 8: Supplemental factors FGF10, HGF and Nicotinamide; effect on growth and differentiation.

1. A - Diagram depicting the genes differentially expressed during the 3 stages of liver development, from hepatoblast to mature hepatocyte.
2. B - Scheme showing the protocol used. Cultures were seeded in expansion medium EM2 (ERFHNic: EGF (E) and R-spondin 1 (R), supplemented with FGF10, HGF and Nicotinamide; ERFHNic is indicated as 'ER' in Figure 8B) 2 days prior the experiment. Two days later, culture media was changed to either EGF (E) alone or EGF supplemented with R-spondin 1 (ER) with or without additional supplements chosen from FGF10 (F) or HGF (H) or Nicotinamide (Nic) or a combination of these at the concentrations stated in the text. Five days later cultures were split and replated at 1:4 ratio for each condition. Under these conditions, cultures have been split and replated every 7 days for a total period of 10 weeks
3. C - First day after first split in each of the culture conditions tested. Results shows that EGF and R-spondin 1 combined with FGF10 or HGF or Nicotinamide or a combination of these are essential to achieve at least 1 passage.
4. D - After long-term culture, the combination of ER supplemented with FNic or ER supplemented with FHNic, both result in high passage numbers. After passage 10, the growth rate is better for the culture condition including the 3 supplemental factors; ERFHNic (figure 15A, B).
5. E - RT-PCR analysis showing the expression of different hepatocyte markers (CYP3A11, Alb, FAH) and cholangiocyte marker (K19) and stem cell marker IGR5 5 days after the withdrawal of certain factors (starting point was ERFHNic). Note that only the condition EF showed expression of all hepatocyte markers tested. HPRT was used as a housekeeping gene to normalize for gene expression.

Figure 9: Table showing the quantification of different hepatocyte and cholangiocyte specific transcription factors in cells from three different liver culture conditions and in adult liver tissue. Also shown is the expression of the key components of the Notch and TGF-beta signalling pathways. E=EFHNic, ER=ERFHNic, ENRW=ENRWFHNic.

Figure 10: Differentiation protocol

1. A - Scheme showing the protocol used. Cultures were seeded in expansion medium (ERFHNic: EGF (E) and R-spondin 1 (R), supplemented with FGF10, HGF and Nicotinamide; ERFHNic is indicated as 'ER' in Figure 10A) 2 days prior to the experiment. Two days later, culture media was changed to EGF (E) supplemented with either A8301 (A), or DAPT (D), or FGF10 (F) or HGF (H) or Nicotinamide (Nic) or R-spondin 1 (R) or Wnt3A or Noggin (N) or a combination of these at the concentrations shown. RNA was isolated at several time points. Mouse liver tissue was taken as positive control (+) whereas water was taken as negative control (-).
2. B - RT-PCR analysis showing the expression of the hepatocyte markers CYP3A11, Alb, FAH, tbx3, TAT and Gck 7 days after differentiation conditions. Note that only the

condition EADF showed an expression of all hepatocyte markers tested. HPRT was used as a housekeeping gene to normalize for gene expression.

3. C - Time course expression analysis after differentiation conditions. At days 2, 5 and 8 days after differentiation, the expression of the hepatocyte markers CYP3A11, Alb, FAH, and the cholangyocyte marker K19, was analysed by RTPCR. Note that the expression of the liver markers CYP3A11 and FAH starts at day 5 and peaks at day 8 after. HPRT was used as a housekeeping gene to normalize for gene expression. A; A8301, D; DAPT, F; FGF10, H; HGF, De; Dexamethasone
4. D - Titration experiment showing the expression of the hepatocyte markers CYP3A11, Alb, FAH, tbx3, TAT, G6P and Gck 7 days after different concentrations of the differentiation compounds A and D. HPRT was used as a housekeeping gene to normalize for gene expression.
5. E - Immunofluorescent staining for the liver markers K19, Albumin and hepatocyte surface marker. Hoeschst was used to stain nuclei.
6. F - Xgal staining on Albcreert2LacZ mice liver-derived organoids. Albumin positive cells (arrows) were detected after EADF differentiation in tamoxifen induced Albcreert2LacZ derived cultures.

Figure 11: Human-derived liver cultures under ERFHNic culture conditions

Figure 12: Liver response to Wnt signaling stimulation under physiological conditions or during regeneration after injury

1. A: Injection of Lgr5 ligand R-spondin1 in Axin2 LacZ mice shows that liver cells are responsive to Wnt stimulation (arrows pointing X-gal positive cells). There was no Lgr5 expression so the inventors hypothesise that Lgr4 was used to initiate the response.
2. B: CCL4 injection in Axin2 LacZ mice shows that during the regeneration response Wnt signalling is activated

Figure 13: Lgr5 upregulation following liver injury-regeneration model.

Adult Lgr5-LacZ KI mice were injected with 0.8 ml/kg of the hepatotoxic compound CCL4. The pictures show that in non injected (undamaged) livers the Wnt pathway is active only in cells lining the ducts. After damage by CCL4 cells also cells not lining duct have an activated Wnt pathway.

1. A - Time course experiment showing upregulation of Lgr5 in CCL4 damaged livers (arrows showing x-gal positive cells). Control CCL4 injected WT mice and placebo-injected Lgr5LacZ Ki mice did not show any staining (right-hand panel).
2. B - Lgr5 co-staining with liver markers.

Figure 14: Isolated duct staining for K19

Lgr5LacZ duct isolation. K19 staining confirms that the isolated and seeded structures are indeed intrahepatic ducts.

Figure 15: Growth factor requirement

The 3 supplemental factors (FGF10, HGF and Nicotinamide) are essential for long term self-

maintenance of liver cultures. After long-term culture, the combination of ER including FNic (\$) or ERFHNic (\$\$), both result in high passage numbers. After passage 10, the growth rate is better for the culture condition including the 3 supplemental factors; ERFHNic (see figure 16B).

Figure 16: Gene expression profile of mouse liver organoids under differentiation conditions resemble the adult and newborn liver profile

1. A - Gene clusters showing the genes similarly expressed (a) or similarly shut down (b) between the differentiation condition EADF and adult or newborn liver.
2. B - Gene clusters showing the genes differentially expressed between the liver organoids and adult or newborn liver (a) and the genes similarly expressed between EADF and newborn liver (b).
3. C - Raw signal data from a microarray analysis, comparing the expression levels of selected ductal markers, transcription factors necessary for Ngn3 expression and endocrine markers in adult liver, adult pancreas, pancreas organoids and liver organoids in expansion media.

Figure 17: Transplantation of the cells into mouse model of liver disease

Organoids were transplanted into the mouse model: adult FGR mice (FAH^{-/-}RAG^{-/-} IL2R^{-/-}). Hepatocytes were transplanted into the mice as a control.

A - K19 positive cells (left top panel) and Fah positive cells (middle panel) derived from the liver organoids transplanted into FAH knock out mice. Hepatocyte transplanted control (top right panel). Lower flow Cytometry plots show that the % of hepatocyte positive cells was higher in the group that resulted in positive FAH engrafted hepatocytes.

C & D - Flow Cytometry analyses of cells transplanted. (C) Clone 1, obtained from Lgr5-GFP mouse, and (D) clone 2, obtained from Lgr5-lacZ mouse. The hepatocyte surface marker shows a positive subpopulation that comprises large cells and highly granular cells, i.e. cells that represent the phenotype of mature hepatocytes.

E -Transplantation schedule.

Figure 18: Mouse liver signature genes

Table showing a) markers expressed in mouse liver stem cells; b) markers not expressed in mouse liver stem cells; c) hepatocyte and cholangiocyte markers expressed in mouse liver stem cell signature for mouse liver organoids in expansion media; d) hepatocyte and cholangiocyte markers not expressed in mouse liver stem cell signature for mouse liver organoids in expansion media; e) reprogramming genes expressed in mouse liver organoids; f) reprogramming genes not expressed in mouse liver organoids. The results were obtained using a liver microarray using the Universal Mouse Reference RNA (Stratagene, Catalog #740100) as a reference RNA. If the absolute figures detected were less than 100, the gene was considered as undetected.

Figure 19: Human liver signature genes

Table showing results of liver microarray of human organoids. From left to right, the results are shown for a) expansion medium EM1, b) expansion medium EM2, c) differentiation

medium, d) adult liver.

The numbers (log2) in the left four columns are the result of a comparison between the sample and a reference (commercial) RNA sample which is used for all arrays. The relative expression of mRNA in each sample compared to the RNA present in the reference sample is shown.. The reference RNA used was Universal Human Reference RNA (Stratagene, Catalog #740000). Thus, negative numbers in these columns do not relate to real expression levels it just means there is less of that RNA then in the Reference sample. The 4 columns on the right are absolute figures. If they are below 100, they are considered as undetected.

EXAMPLES

Example 1 - An expansion medium for liver organoid growth and expansion

[0240] After isolation, biliary ducts (see Fig. 1) were suspended in Matrigel and cultured in different growth factor conditions. The combination of EGF (50ng/ml) and R-spondin 1 (lug/ml) supplemented with FGF10 (100ng/ml), HGF (25-50ng/ml) and Nicotinamide (1-10mM), (ERFHNic) were essential for the long term maintenance of the cultures, indicating that Wnt signalling and EGF signalling are strictly required to maintain adult liver progenitor proliferation *in vitro*. The addition of Noggin (100ng/ml) and Wnt conditioned media (50%) also showed long term maintenance of the cultures (see Figs. 1A and 1B). Under these conditions that supported long-term maintenance, Lgr5 expression as well as hepatocyte markers (Albumin) and cholangiocyte markers (K7) were detected by RT-PCR (see Fig. 1C). Under these conditions liver organoids have been weekly passaged by mechanical or enzymatic dissociation, at 1:8 dilution, and have been grown for many months (Fig. 1D).

[0241] We analysed the expression of the Wnt target genes Axin2 and Lgr5 in the cultures. Cultures of both Axin2LacZ and Lgr5-LacZ livers revealed the presence of Axin2- and Lgr5-positive cells in the liver organoids 1 month after seeding, thus confirming that the Wnt signalling is active and required for culture growth (Fig. 3). The liver cultures also express hepatocyte markers (e.g. albumin, transthyretin, Glutamine synthetase) and cholangiocyte makers (Keratin 7 and 19) (see Fig 4).

[0242] When single Lgr5 cells from a Lgr5LacZ or Lgr5GFP mouse were sorted, single colonies grew into organoids. These cultures also express markers of cholangiocyte and hepatocyte lineages and have been maintained and regularly split into 1:6-1:8 for more than 4 months (see Figs. 5A & B). Interestingly, only the cultures derived from Lgr5 positive cells grew into organoids (Figs 5C & D). These data indicate that Lgr5 cells are progenitor cells of these cultures and able to propagate progeny of the 2 different liver lineages.

[0243] Having established that the liver organoids are derived from Lgr5+ve cells we set out to

determine their individual gene signature as compared to the adult liver signature. RNA was isolated from adult liver and from liver organoids grown in ER or ENRW media supplemented with FGF10, Nicotinamide and Hepatocyte Growth Factor. The genetic signature of the adult liver and the 2 liver culture conditions was subsequently derived via comparative gene expression profiling in respect to the expression of a Universal RNA reference. The use of the same reference RNA for the hybridization to all the samples allowed us to compare the 3 independent samples among them (adult liver, ER and ENRW). The heat map analysis revealed that the expression profile of both culture conditions highly resemble the adult liver tissue expression profile, whereas they do not share the same profile when compared to muscle or adipose tissue profile (see Fig 6). Among the similar gene expression profile between the adult liver and the liver cultures, liver specific genes as HNF1a, HNF1b, HNF4, Alb, Glul, Met, G6P, Fahd1, Fahd2a, CYP4B1, K7 and K19 are detected. The heat map analysis reveals that both culture conditions present similar expression pattern among each other and when compared to the adult liver sample. However, when analyzing the data in detail, we can observe that the condition without Wnt and without noggin shows a more differentiated pattern than the condition including both growth factors. This is in agreement with the data shown in Fig 1C where hepatocyte differentiation (by means of albumin expression) is almost absent in the presence of Wnt. This result would indicate that Wnt is favouring the self-renewal of the culture in detriment of the differentiation.

[0244] Also, in both culture conditions as well as in the adult liver, non-specific adult liver genes as AFP, and non-liver transcription factors as Pdx1 or NeuroD can be detected.

[0245] It is remarkable that, in both culture conditions but not in the adult liver, the stem cell marker Lgr5 was one of the most highly enriched genes in the liver culture signature. Also, cell markers of progenitor populations in small intestine and stomach as Cd44 and Sox9 (Barker & Huch et al Cell stem cell 2010) were highly expressed in both culture conditions but not in adult liver, indicating again the self-renewal capacity of the liver cultures as well as the quiescent status of the normal adult liver.

[0246] Additionally, apart from Lgr5, multiple Wnt target genes were also highly upregulated in the liver cultures compared to the adult liver including MMP7, Sp5 and Tnfrs19, among others, providing strong evidence of the requirement of an active and robust canonical Wnt signaling activity to maintain the self renewing capacity of the cultures.

Example 2 - An improved differentiation medium

[0247] Under ER or ENRW conditions the liver cultures self-renew, and can be maintained and expanded in a weekly basis, for up to 1 year (figure 7A). The karyotypic analysis after 1 year shows no evidence of chromosomal aberrations. More than 66% of the cells analysed presented normal chromosomal counts and 13% of them also showed polyploidy, a characteristic trait of hepatocytes (Figure 7B).

[0248] The combination of EGF (50ng/ml) and R-spondin 1 (lug/ml) supplemented with FGF10 (100ng/ml), HGF (25-50ng/ml) and Nicotinamide (1-10mM), were preferable for the long term maintenance of the cultures. Under these conditions, we obtained long-lived cell cultures that express biliary duct and some hepatoblast or immature-hepatocyte markers (Glul, Albumine). However, the number of cells positive for these hepatocyte markers was very low. Under these culture conditions, no mature hepatocyte markers (e.g. p450 Cytochromes) were detected. These results suggest that the culture conditions described here facilitate the expansion of liver progenitors able to generate hepatocyte-like cells, albeit at lower numbers, but not fully mature hepatocytes (Fig.8A).

[0249] To enhance the hepatocytic nature of the cultures and obtain mature hepatocytes *in vitro*, we first determined whether the three supplemental factors (FGF10, HGF and Nicotinamide) added to EGF and R-spondin1 were exerting either a positive or negative effect on the hepatocyte expression, as well as on the self-renewal of the culture. We generated liver organoid cultures and cultured them either with EGF or EGF and R-spondin1 plus FGF10 or HGF or Nicotinamide or the combination of these, and we split the cultures once a week for a total period of 10 weeks. At each time-point we also analysed the expression of several mature hepatocyte markers (FAH, CYP3A11) and hepatoblast markers (albumin) (Figure 8B).

[0250] In agreement with the data in figure 1 (see example 1), we observed that R-spondin1 and Nicotinamide combined with FGF10 are essential for the growth and self-renewal of the liver cultures (Figure 8C&D). R-spondin1 and Nicotinamide both inhibit the expression of the mature marker CYP3A11 and yet promote the expression of the hepatoblast marker albumin. The addition of either FGF10 or HGF to media containing only EGF (without R-spondin1 and without nicotinamide), facilitated the expression of the mature marker CYP3A11, albeit at very low levels (figure 8E). To identify additional compounds that might facilitate hepatocyte differentiation, we used two different approaches, both based upon base conditions of: EGF + HGF and/or FGF10.

[0251] The first approach involved testing a series of compounds in addition to the EGF + FGF10 or HGF condition. A complete list of the compounds analysed is shown in table 2.

Table 2

Compounds	Signal		Concentration	Result	
				<i>Alb</i>	<i>CYP3A11</i>
Exendin4	Glucagon like peptide 2 analog	Sigma E7144	0.1-1uM		
Retinoic Acid	RAR-RXR receptor ligand	Sigma	25nM		
Retinoic Acid + Exendin 4					
Sonic Hedgehog		Invitrogen C25II	500-100ng/ml		

Compounds	Signal		Concentration	Result	
				<i>Alb</i>	<i>CYP3AII</i>
BMP4	BMP signaling	Peprotech 120-05	20ng/ml		
DAPT	Gamma-secretase inhibitor	Sigma D5942	10 nM		
A8301	Alk5/4/7 inhibitor	Tocris Bioscience 2939	50 nM		
DAPT + A8301				+++	+++
FGF4	FGFR1,2 ligand	Peprotech	50ng/ml		
FGF1	FGFR1,2,3,4 ligand	Peprotech 450-33A	100ng/ml		
Dexamethasone		Sigma D4902 25MG	10 µM-1mM		
Oncostatin M (OSM)		R&D systems 495-MO- 025	10-1000 ng/ml		
FGF4+OSM+Dexa					
IGF		peprotech	100ng/ml		
Valproic acid	histone deacetylase inhibitor and regulator of ERK, PKC Wnt/β- catenin pathways	Stemgent 04-0007	250 µM		
Sodium Butyrate	histone deacetylase inhibitor	Stemgent 04-0005	250 µM		
BIX01294	G9a HMTase inhibitor	Stemgent 04-0002	1 µM		
RG 108	DNA methyltransferase inhibitor	Stemgent 04-0001	1 µM		
TSA			100 nM	+	-
Hydrocortisone	glucocorticoid	Sigma H6909	5nM		
Oncostatin M (OSM)		R&D systems 495-MO-	10-1000 ng/ml		

Compounds	Signal		Concentration	Result	
				Alb	CYP3AII
		025			
ARA		Sigma A 0937	500 nM		
R 59022	Diacylglycerol kinase inhibitor	Sigma D 5919	500nM-50nM	+	+
Arterenol bitartre:----	andrenoreceptor agonist	sigma A 0937	500nM-50nM-5nM		
LIF			10 ³		
PD 035901	MEK1 inhibitor	Axon Medchem cat n 1386	500nM		
CHIR99021	GSK3 inhibitor	Axon Medchem cat n 1408	3uM		
DMSO			1%		
L-Ascorbic acid		Sigma 077K13021	1mM		
VEGF		Peprotech			
Matrigel 50%					
Matrigel 20%					
VEGF+DEXA					

[0252] The second approach took into account knowledge from published developmental studies regarding the expression of the transcription factors essential to achieve biliary and hepatocyte differentiation *in vivo*. A comparative analysis of the expression of transcription factors in the organoids under E or ER or ENRW conditions supplemented with FGF10, HGF and Nicotinamide is shown in figure 8. All the transcription factors required for Hepatocyte specification were present, besides *tbx3* and *prox1*. However, we also noticed that the expression of specific biliary transcription factors was highly upregulated in the cultures containing R-spondin1 (R), indicating that the culture gene expression was unbalanced towards a more biliary cell fate.

[0253] Notch and TGF-beta signaling pathways have been implicated in biliary cell fate *in vivo*. In fact, deletion of *Rbpj* (essential to achieve active Notch signalling) results in abnormal tubulogenesis (Zong Y. Development 2009) and the addition of TGF-beta to liver explants facilitates the biliary differentiation *in vitro* (Clotman F. Genes and Development 2005). Since both Notch and TGF-beta signalling pathways were highly upregulated in the liver cultures

(Figure 9) we reasoned that inhibition of biliary duct cell-fate might trigger the differentiation of the cells towards a more hepatocytic phenotype. A8301 was selected as an inhibitor of TGF-beta receptor ALK5, 4, and 7 and DAPT as inhibitor of the gamma-secretase, the active protease essential to activate the Notch pathway. We first cultured the cells for 2 days in the expansion conditions (ER media) and at day 2 (figure 10A) we started the differentiation conditions by adding the combination of the different compounds. Media was changed every other day, and the expression of differentiated markers was analysed 8-9 days later. The ER and ENRW conditions were used as negative control.

[0254] The combination of EGF + FGF10 with DAPT and A8301 resulted in surprisingly large enhancement of expression of the hepatocyte markers analysed (CYP3A11, TAT, Albumin) (figure 10B). The effect was already detectable by day 5 and peaked at days 8-9 (figure 10C). The maximal concentration efficiency was achieved at 10uM (DAPT) and 50 nM (A8301) (figure 10D) respectively. The addition of dexamethasone (a known hepatocyte differentiation molecule) did not result in any improvement in gene expression. The combination of EGF, FGF10, A8301 and DAPT not only enhances the expression but also increases the number of hepatocyte-like cells, as assessed by immunofluorescent against the hepatocyte markers albumin and 2F8, and Xgal staining on AlbCreLacZ derived organoids (figure 10E & F). Therefore, we can conclude that the aforementioned differentiation protocol facilitates the generation of hepatocyte-like cells *in vitro* from liver stem cell cultures.

Example 3 - Human liver organoids

[0255] Using these expansion conditions (ERFHNic and ENRWFHNic) we have also been able to expand human biliary-derived cultures (figure 11) with the addition of 500 uM TGF beta inhibitor (A83-01) to the expansion medium.

MATERIAL AND METHODS (For Examples 1-3)

Liver culture-biliary duct isolation

[0256] Isolated adult liver tissue was washed in cold Advanced-DMEM/F12 (Invitrogen) and then, the tissue was chopped into pieces of around 5mm animals and further washed with cold dissociation buffer (collagenase, dispase, FBS in DMEM media). The tissue fragments were incubated with the dissociation buffer for 2 h at 37°C. Then, the tissue fragments were vigorously suspended in 10 ml of cold isolation buffer with a 10 ml pipette. The first supernatant containing death cells was discarded and the sediment was suspended with 10-15ml of dissociation buffer. After further vigorous suspension of the tissue fragments the supernatant is enriched in biliary ducts. This procedure is repeated until enough biliary ducts are obtained.

[0257] Isolated biliary ducts are pelleted and mixed with 50µl of Matrigel (BD Bioscience), seeded on 24-well tissue culture plates and incubated for 5-10 min at 37°C until complete polymerization of the Matrigel. After polymerization, 500 µl of tissue culture media are overloaded.

[0258] Media composition: Advanced-DMEM/F12 supplemented with B27, N2, 200ng/ml N-Acetylcysteine, 50ng/ml EGF, 1 µg/ml R-spondin1, gastrin: 10 nM, FGF10 100 ng/ml, Nicotinamide 10mM and HGF: 50ng/ml and 50% Wnt conditioned media.

[0259] The entire medium was changed every 2 days. After 1 week, Wnt conditioned media is withdrawal and the formed organoids removed from the Matrigel using a 1000 µl pipette and were dissociated mechanically into small fragments and transferred to fresh Matrigel. Passage was performed in 1:4 split ratio once or twice per week. Under these conditions cultures have been maintained for at least 6 month.

Reagents

[0260] Human Hepatocyte Growth Factor (HGF) was purchased from Peprotech, EGF invitrogen, R-Spondin Nuvelo, Noggin peprotech , FGF10 Peprotech, gastrin Sigma Aldrich, nicotinamide Sigma.

Microarray

[0261] For the expression analysis of Lgr5-derived liver cultures, RNA was isolated using a Qiagen RNAase kit, from adult liver or from liver cultures cultured in media without Wntcm and Noggin (ER) or with Wntcm and Noggin (ENRW). 150ng of total RNA was labelled with low RNA Input Linear Amp kit (Agilent Technologies, Palo Alto, CA). Universal mouse Reference RNA (Agilent) was differentially labelled and hybridized to either adult liver tissue or ER or ENRW treated cultures. A 4X44K Agilent Whole Mouse Genome dual colour Microarrays (G4122F) was used. Labelling, hybridization, and washing were performed according to Agilent guidelines.

Example 4 - Lgr5 expression is upregulated following liver injury

[0262] In the liver, Wnt signalling is active in central vein areas. We have recently observed that Wnt signaling plays a key role in liver metabolism (Boj et al. personal communication). In the liver duct cells, Wnt signalling is activated following liver injury (Hu et al 2007, Gastroenterology, 133(5): 1579-91). Similarly, using the *Axin2-LacZ* allele, which represents a faithful, general reporter for Wnt signalling, we also have observed upregulation of Wnt

signaling in the whole liver parenchyma after injection of the Wnt agonist Rspo1 (see figure 12A) or following liver injury by the hepatotoxic compound carbon tetrachloride (CCl₄) (see figure 12B).

[0263] The Wnt target gene *Lgr5* marks stem cells in several actively self-renewing tissues, but has not previously been reported to be expressed upon injury. Our previously described *Lgr5-LacZ* knockin mice (Barker et al, 2007, Nature 449 (7165): 1003-7) show that *Lgr5* is essentially undetectable in healthy liver although a residual mRNA expression is detected by qPCR. Following injection of CCl₄ on *Lgr5-LacZ* knockin mice (see Barker et al, 2007, supra for LacZ mice and Furuyama K et al., Nat Genetics, 43, 34-41, 2001 for description of CCl₄ method), we observed a clear expression of the reporter in newly formed bud structures in the liver (see figure 13A), peaking at day 6.5 after injury and being maintained up to day 9 to show a clear decay once the liver is completely regenerated at day 13 after injury (see figure 13A, top right panel). No expression of the reporter was detected in wild-type littermates undergoing similar injury protocol (see figure 13A, bottom right panel).

[0264] The appearance of *Lgr5* expression at sites of active regeneration, suggested that *Lgr5* might herald *de novo* activation by Wnt of regenerative stem cells/progenitors upon injury. Indeed, we found that the *de novo* appearing *Lgr5* cells do not express markers of mature liver cells (K19 or FAH) or stellate cells (SMA) but instead, they are positive for the recently described liver progenitor marker Sox9 (figure 13B). This means that *Lgr5*⁺ cells, which are the starting point for obtaining *in vitro* organoids, can be obtained from liver fragments by inducing liver injury or by stimulating Wnt signalling with R-spondin. The induction of *Lgr5* expression in liver cells by injury or by R-spondin may be carried out *in vivo* before the cells are obtained, *ex vivo* in an isolated liver, or *in vitro* in a liver fragment or population of liver cells.

Example 5 - Long-term expansion of liver organoid cultures

[0265] In example 1, it was found that the combination of EGF (50 ng/ml) and R-spondin 1 (lug/ml) supplemented with FGF10 (100ng/ml), HGF (25-50ng/ml) and Nicotinamide (1-10mM), were preferable for the long term maintenance of the cultures. We now also have evidence that the three supplemental factors (FGF10, HGF and Nicotinamide) added to EGF and R-spondin1 are all necessary for the expansion of the cultures for longer than 3 months. To assess that, we isolated biliary ducts from the liver parenchyma, as shown in figure 14 (K19 staining was used to confirm the identity of the isolated structures), and generated liver organoid cultures by culturing them with: i) EGF; or ii) EGF and R-spondin1 plus FGF10 or HGF or Nicotinamide; or iii) EGF and R-spondin1 plus FGF10 and HGF and Nicotinamide (ERFHNic). We have split the cultures once a week for a total period of 14 weeks. Results confirmed, as reported in examples 1 and 2, that EGF, R-spondin1 and Nicotinamide combined with FGF10 are essential for the growth and self-renewal of the liver cultures. After 10 passages, the cultures lacking HGF showed a growth disadvantage compared to the cultures supplemented with HGF. Although still viable, the proliferation ratio decreased to 1:2-1:4 compared to the 1:6-1:8 of the cultures supplemented with the complete combination (FGF10,

HGF, and Nicotinamide). After 15 passages, the cultures with ERFNic not supplemented with HGF were no longer viable. Therefore, these results suggest that HGF is essential for maintaining a good proliferating rate after long-term maintenance (figure 15).

Example 6 - Markers expressed in liver organoids under differentiation conditions

[0266] Using the differentiation protocol described in example 2, we were able to detect a hepatoblast marker (albumin) and a hepatocyte surface marker in the liver organoids. To quantify the number of these hepatocyte-like cells, we performed flow cytometry analysis of the cultures using a hepatocyte surface marker. We observed that, whereas in the expansion culture condition almost no hepatocyte surface marker-positive cells were detected, after differentiation, up to 35% of the cells were positive for this hepatocyte surface marker (see Figure 17B&C).

[0267] We then analysed the gene expression profile of the mouse liver organoids under these differentiation conditions (figure 16 and figure 1, we see strong upregulation of eg Alb, FAH, and TAT and the Cyp3 genes). We found that the gene expression of the mouse liver organoids after differentiation resemble that of mature mouse hepatocytes and/or mouse liver.

Example 7 - Transplantation of liver organoids into mice

[0268] Cells were taken from the organoids that had been grown using ERFHNic expansion conditions and EAFD differentiation conditions and were transplanted into immunodeficient strain of mice deficient in the tyrosine catabolic enzyme fumarylacetoacetate hydrolase (FAH), a mouse model for Tyrosinemia type I human disease (Azuma et al. 2007, Nature Biotech, 25(8), 903-910). The transplantation schedule is shown in figure 17D. Preliminary results show that scattered FAH positive cells can be found in the liver parenchyma of the FAH deficient mice, indicating that liver cells derived from the organoid cultures have engrafted into the recipient livers (see figure 17A, right-hand side). Furthermore, significantly increased numbers of K19 positive cells were also detected in the livers of the recipient mice. This suggests that the organoid-derived transplanted cells are able to generate both lineages *in vivo*: hepatocytes (as demonstrated by the FAH marker) and cholangiocytes (as demonstrated by the K19 marker) (see figure 17A, left top panel). This was further supported by flow cytometry analysis of transplanted cells that had come from two separate clones from two separate cultures (figures 17B and 17C respectively). The Lgr5+ cells were transduced with a virus containing GFP and flow cytometry analysis was carried out after differentiation. Cells that were positive for the hepatocyte surface marker show a larger scatter indicating larger cells, which represent granularity and maturity i.e. mature hepatocyte cells. The cells that were negative for the hepatocyte surface marker resulted in less scattering indicating smaller cells i.e. less mature progenitors. Therefore, all cell types are present (mature and immature cells) in a differentiating culture. The rest of the differentiated cells, so the cells not used for FACS

analysis were used for the transplantation experiments.

Example 8

[0269] Organoids from mouse liver cultured in accordance with a method of the invention were analysed using microarray analysis to determine which genes are expressed and which genes are not expressed.

Example 9

[0270] Organoids from human liver cultured using the EM1, EM2 and DM media of the invention and human liver were analysed using oligonucleotide microarray analysis to determine which genes are expressed and which genes are not expressed. A significantly different gene expression profile was noticeable between the genes expressed in expansion media, the genes expressed in differentiation medium and the genes expressed in adult liver. The trend for hepatocyte gene expression is roughly the same as for in the mouse but the differentiation of the organoids was less than in the mouse liver organoids. This may be due to use of the human cell used.

[0271] As often happens in an analysis using an oligonucleotide microarray, Lgr5 and Tnfrsf19 were not detected. However, they were found to be present in organoids cultured in the expansion medium.

MATERIALS & METHODS (for Examples 4 to 7)

Animal treatment

[0272] Two-Eight month old Lgr5LacZ or Axin2-LacZ or WT littermates BL6/Balbc Flmice received an intraperitoneal injection of 0.8ml/kg of CCL4 dissolved in corn oil (n=) or corn oil alone (n=). Mice were sacrificed 2 or 5 or 9 or 13 days later and the liver was isolated and further processed for RNA or bgalactosidase staining.

β-galactosidase (lacZ) staining

[0273] Liver tissues were isolated and immediately incubated for 2 hours in a 20-fold volume of ice-cold fixative (1 % Formaldehyde; 0.2 % Gluteraldehyde; 0.02 % NP40 in PBS0) at 4 °C on a rolling platform. The fixative was removed and the tissues washed twice in washing buffer

(PES0; 2 mM MgCl₂; 0.02 % NP40; 0.1 % NaDeoxycholate) for 20 minutes at room temperature on a rolling platform. The β -galactosidase substrate (5 mM K₃Fe(CN)₆; 5 mM K₄Fe(CN)₆·3H₂O; 2 mM MgCl₂; 0.02 % NP40; 0.1 % NaDeoxycholate; 1 mg/ml X-gal in PBS0) was then added and the tissues incubated in the dark at 37°C for 2h and overnight at room temperature. The substrate was removed and the tissues washed twice PBS0 for 20 minutes at room temperature on a rolling platform. The tissues were then fixed overnight in a 20-fold volume of 4 % Paraformaldehyde (PFA) in PBS0 at 4°C in the dark on a rolling platform. The PFA was removed and the tissues washed twice in PBS0 for 20 minutes at room temperature on a rolling platform.

[0274] The stained tissues were transferred to tissue cassettes and paraffin blocks prepared using standard methods. Tissue sections (4 μ M) were prepared and counterstained with neutral red.

R-spondin1 treatment

[0275] Axin2-lacZ mice aged 6-8 weeks were injected IP with 100 μ g of purified human R-spondin1 and sacrificed 48 hours later for LacZ expression analysis in the liver.

RT-PCR

[0276] RNA was extracted from gastric cell cultures or freshly isolated tissue using the RNeasy Mini RNA Extraction Kit (Qiagen) and reverse-transcribed using Moloney Murine Leukemia Virus reverse transcriptase (Promega). cDNA was amplified in a thermal cycler (GeneAmp PCR System 9700; Applied Biosystems, London, UK) as previously described (Huch et al., 2009). Primers used are shown in Table 3 below.

Table 3: Primers for RT-PCR

Gene name	Gene Symbol	Sequence	PCR product (bp)
cytochrome P450, family 3, subfamily a, polypeptide 11	CYP3A11	fw TGGTCAAACGCCCTCTCCTTGCT G rv ACTGGGCCAAAATCCCGCCG	100
Glucose-6-phosphatase	G6P	fw GAATTACCAAGACTCCAGG rv TGAGACAATACTCCGGAGG	581
Keratin 19	Krt19	fw GTCCTACAGATTGACAATGC rv CACGCTCTGGATCTGTGACA	549

Albumin	Alb	fw	GCGCAGATGACAGGGCGGAA	358
		rv	GTGCCGTAGCATGCGGGAGG	
t-box 3	Tbx3	fw	AGCGATCACGCAACGTGGCA	441
		rv	GGCTTCGCTGGGACACAGATCT	
		TT		
Prospero-related-homeobox protein 1	Prox1	fw	TTCAACAGATGCATTACC	270
		rv	TCTTTGCCCCGCGATGATG	
Fumarylacetoacetate-hydrolase	Fah	fw	ACGACTGGAGCGCACGAGAC	183
		rv	AGGGCTGGCTGTGGCAGAGA	
Tyrosine aminotransferase	Tat	fw	TTTGGCAGTGGCTGAAAGGCA	258
		rv	GGGCCCAGGATCCGCTGACT	
Tryptophan 2,3-dioxygenase	Tdo2	fw	ACTCCCCGTAGAAGGCAGCGA	583
		rv	TCTTTCCAGCCATGCCTCCACT	
Leucine-rich repeat-containing G-protein coupled receptor 5	Lgr5	fw	GGAAATGCTTTGACACACATTC	413
		rv	GGAAGTCATCAAGGTTATTATA	
		A		
Transthyretin	TTR	fw	ATGGTCAAAGTCCTGGATGC	220
		rv	AATTCATGGAACGGGGAAAT	
Glucokinase	Gck	fw	AAGATCATTGGCGGAAAG	193
		rv	GAGTGCTCAGGATGTTAAG	
hypoxanthine phosphoribosyltransferase	Hprt	fw	AAGCTTGCTGGTGAAAAGGA	186
		rv	TTGCGCTCATCTTAGGCTTT	

Immunohistochemistry

[0277] Immunostaining procedure used here was previously described in Huch et al. 2009. Briefly, five-micrometer sections were deparaffinized, rehydrated, and tissue sections were permeabilized using PBS-T (PBS; Tween20 0.1 %). When required, sections were treated with 10 mM citrate buffer (pH 6.0) for antigen retrieval, blocked using Universal blocking buffer (BioGenex)) and incubated with the primary antibody. Then, sections were washed twice with

PBS and incubated with peroxidase conjugated secondary antibodies. DAB+ (DAKO) was used as a chromogen substrate. Sections were counterstained with Mayer's hematoxylin and visualized on a Leica DMR microscope. The primary antibodies used were rabbit anti-Sox9 (1:600; 1h at RT, Millipore), mouse anti-SMA (1:1000, overnight at 4°C, Sigma), rabbit anti-FAH (1:5000; overnight 37°C, gift from M. Grompe), rabbit anti-K19 (1:500; overnight 4°C, gift from M. Grompe). The peroxidase conjugated secondary antibodies used were Mouse or Rabbit Brightvision (Immunologic).

Immunofluorescence

[0278] For whole mount staining, organoids or isolated biliary ducts were fixed with acetone (organoids) or PFA4% (biliary ducts) for 30min, washed once with PBS, permeabilized with PBS 0.3%Triton-X100 for 5min, blocked using Universal blocking solution (Power block HK085-5KE BioGenex) and incubated overnight with the primary antibodies diluted in PBS1%FBS. Following several washes in PBS, samples were incubated with the secondary antibody. Nuclei were stained with Hoescht33342. Images were acquired using confocal microscopy (Leica, SP5). Three-dimensional reconstruction was performed using Volocity Software (Improvision). The primary antibodies used were rabbit anti-K19 (1:500; gift from M. Grompe), rat anti-hepatocyte surface marker (1:50, gift M. Grompe), goat anti-albumin (1:50, santa Cruz). The secondary antibodies used were all raised in donkey and conjugated to different Alexa fluorophores (donkey anti-goat 568, donkey anti rat-488, donkey anti rabbit-647, Molecular probes).

Flow cytometry

[0279] Dissociated cells were resuspended at 1×10^4 cells per milliliter in 1ml of DMEM + 2% FBS prior to the addition of MIC1-1C3 hybridoma supernatant at a 1:20 dilution or OC2-2F8 hybridoma supernatant at a 1:50 dilution, and incubated for 30 min at 4°C. After a wash with cold Dulbecco's Phosphate Buffered Saline (DPBS), cells were resuspended in DMEM + 2% FBS containing a 1:200 dilution of APC-conjugated goat anti-rat secondary antibody adsorbed against mouse serum proteins (Jackson ImmunoResearch). Propidium iodide staining was used to label dead cells for exclusion. Cells were analyzed and sorted with a Cytopeia inFluxV-GS (Becton-Dickenson).

Transplantation assay

[0280] The injection of sorted cell populations to the spleen and the withdrawal of NTBC to induce hepatocyte selection were performed as described previously (Overturf et al. 1996). Drug withdrawal was done in periods of 3 wk, followed by readministration until normal weight was restored in the recipient animals.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2009022907A](#) [\[0031\]](#) [\[0144\]](#)
- [WO2010016766A](#) [\[0033\]](#) [\[0144\]](#)
- [US6165782A](#) [\[0223\]](#)
- [US6207455B](#) [\[0223\]](#)
- [US6218181B](#) [\[0223\]](#)
- [US6277633B](#) [\[0223\]](#)
- [US6323031B](#) [\[0223\]](#)
- [US5464758A](#) [\[0225\]](#)
- [US6432705B](#) [\[0225\]](#)

Non-patent literature cited in the description

- **BARKER et al.** Nature, 2007, vol. 449, 1003-1007 [\[0005\]](#)
- **SAHA et al.** Curr Opin Chem Biol., 2007, vol. 11, 4381-387 [\[0034\]](#)
- **SAHA et al.** Biophysical Journal, 2008, vol. 95, 4426-4438 [\[0034\]](#)
- **LITTLE et al.** Chem. Rev, 2008, vol. 108, 1787-1796 [\[0034\]](#)
- **BOTTENSTEINS** ATOPNAS, 1979, vol. 76, 1514-517 [\[0054\]](#)
- **BREWER et al.** J Neurosci Res., 1993, vol. 35, 5567-76 [\[0055\]](#)
- **ZILBERBERG et al.** BMC Cell Biol., 2007, vol. 8, 41- [\[0058\]](#)
- **KORINEK et al.** Science, 1997, vol. 275, 1784-1787 [\[0063\]](#) [\[0073\]](#)
- **KIRIKOSHI H et al.** Biochem Biophys Res Com, 2001, vol. 283, 798-805 [\[0064\]](#)
- **KESTUTIS PLANUTIS et al.** BMC Cell Biol., 2007, vol. 8, 12- [\[0065\]](#)
- **LIU et al.** Angew Chem Int Ed Engl., 2005, vol. 44, 1987-90 [\[0068\]](#)
- **LEOST, M. et al.** Eur. J. Biochem., 2000, vol. 267, 5983-5994 [\[0069\]](#)
- **MEIJER, L. et al.** Chem. Biol., 2003, vol. 10, 1255-1266 [\[0069\]](#)
- **MEIJER et al.** Trends in Pharmacological Sciences, 2004, vol. 25, 471-480 [\[0069\]](#)
- **LIAO et al.** Endocrinology, 2004, vol. 145, 62941-9 [\[0069\]](#)

- **HERBST**Int Journal of radiation oncology, 2004, vol. 59, 2S21-S26 [0074]
- **ORNITZ**J Biol Chem., 1998, vol. 273, 95349-57 [0076]
- **IGARASHI**J Biol Chem., 1998, vol. 273, 2113230-5 [0076]
- **ZONG Y.**Development, 2009, [0098] [0253]
- **CLOTMAN F.**Genes and Development, 2005, [0098] [0253]
- **FUCHS, O.**Inhibition of TGF-Signaling for the Treatment of Tumor Metastasis and Fibrotic DiseasesCurrent Signal Transduction Therapy, 2011, vol. 6, 129-43 [0101]
- **DE GOUVILLE et al.**Br J Pharmacol., 2005, vol. 145, 2166-177 [0102]
- **BARKERHUCH et al.**Cell stem cell, 2010, [0144] [0245]
- **Cancer Res**, 2001, vol. 61, 8164-8170 [0183]
- **METZGER et al.**Nature, 1988, vol. 334, 31-36 [0219]
- **ANDERSON**Nature, 1998, vol. 392, 25-30 [0221]
- **WALTHERSTEIN**Drugs, 2000, vol. 60, 249-71 [0221]
- **KAY et al.**Nat. Med., 2001, vol. 7, 33-40 [0221]
- **RUSSELL**J. Gen. Virol., 2000, vol. 81, 2573-604 [0221]
- **AMADOCHEN**Science, 1999, vol. 285, 674-6 [0221] [0223]
- **FEDERICO**Curr. Opin. Biotechnol., 1999, vol. 10, 448-53 [0221]
- **VIGNANALDINI**J. Gene Med., 2000, vol. 2, 308-16 [0221]
- **MARIN et al.**Mol. Med. Today, 1997, vol. 3, 396-403 [0221]
- **PENGRUSSELL**Curr. Opin. Biotechnol., 1999, vol. 10, 454-7 [0221]
- **SOMMERFELT**J. Gen. Virol., 1999, vol. 80, 3049-64 [0221]
- **REISER**Gene Ther., 2000, vol. 7, 910-3 [0221]
- **RUSSELL**J. Gen. Virol., 2000, vol. 81, 2573-2604 [0222]
- **GONCALVES**Virol J., 2005, vol. 2, 143- [0222]
- **MANNO et al.**Nat medicine, 2006, [0222]
- **STROES et al.**ATVB, 2008, [0222]
- **KAPLITT, FEIGIN**Lancet, 2009, [0222]
- **MAGUIRE, SIMONELLI et al.**NEJM, 2008, [0222]
- **BAINBRIDGE et al.**NEJM, 2008, [0222]
- **GAO**Molecular Therapy, 2006, vol. 13, 77-87 [0222]
- **FEDERICO**Curr Opin Biotechnol, 1999, vol. 10, 448-53 [0223]
- **VIGNA et al.**J Gene Med, 2000, vol. 2, 308-16 [0223]
- **BRINSTER et al.**Nature, 1982, vol. 296, 39-42 [0225]
- **MAYO et al.**Cell, 1982, vol. 29, 99-108 [0225]
- **WANG et al.**Proc. Natl. Acad. Sci. USA, 1994, vol. 91, 8180-8184 [0225]
- **MADERWHITE**Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 5603-5607 [0225]
- **GOSSENBUJARD**Proc. Natl. Acad. Sci. USA, 1992, vol. 89, 5547-5551 [0225]
- **FURTH et al.**Proc. Natl. Acad. Sci. USA, 1994, vol. 91, 9302-9306 [0225]
- **HOWE et al.**J. Biol. Chem., 1995, vol. 270, 14168-14174 [0225]
- **RESNITZKY et al.**Mol. Cell. Biol., 1994, vol. 14, 1669-1679 [0225]
- **SHOCKETT et al.**Proc. Natl. Acad. Sci. USA, 1995, vol. 92, 6522-6526 [0225]
- **GEIDUSCHEK TOCCHINI-VALENTINI**Annu. Rev. Biochem., 1988, vol. 57, 873-914 [0226]
- **WILLIS**Eur. J. Biochem., 1993, vol. 212, 1-11 [0226]

- **HERNANDEZ**J. Biol. Chem., 2001, vol. 276, 26733-36 [0226]
- **MATTAJ et al.**Cell, 1988, vol. 55, 435-442 [0226]
- **GUPTAREDDY**Nucleic Acids Res., 1991, vol. 19, 2073-2075 [0226]
- **MYSLINSKI et al.**Nucl. Acids Res., 2001, vol. 21, 2502-09 [0226]
- **SAMBROOKRUSSEL**Molecular Cloning: A Laboratory ManualCold Spring Harbor Laboratory Press20010000 [0227]
- **CLAIR et al.**Antimicrob. Agents Chemother, 1987, vol. 31, 844-849 [0228]
- **ELBASHIR et al.**Nature, 2001, vol. 411, 494-98 [0230]
- **CAPLEN et al.**Proc. Natl. Acad. Sci. USA, 2001, vol. 98, 9742-47 [0230]
- **HU et al.**Gastroenterology, 2007, vol. 133, 51579-91 [0262]
- **BARKER et al.**Nature, 2007, vol. 449, 71651003-7 [0263]
- **FURUYAMA K et al.**Nat Genetics, 2001, vol. 43, 34-41 [0263]
- **AZUMA et al.**Nature Biotech, 2007, vol. 25, 8903-910 [0268]

Patentkrav

- 1.** Fremgangsmåde til opnåelse af et leverorganoid, hvilken fremgangsmåde omfatter:
- dyrkning af én eller flere leverepitelstamceller, eller et leverfragment eller en levergaldegang, i kontakt med en ekstracellulær matrix i tilstedeværelse af et ekspansionsmedium, mediet omfatter et basalmedium til dyre- eller humane celler til hvilke tilsættes:
- epidermal vækstfaktor (EGF), en fibroblast vækstfaktor (FGF) der er i stand til at binde til FGFR2 eller FGFR4, nicotinamid, og en Wnt-agonist valgt fra Rspondin 1, Rspondin 2, Rspondin 3 eller Rspondin 4.
- 2.** Fremgangsmåden ifølge krav 1, hvor ekspansionsmediet yderligere omfatter HGF.
- 3.** Fremgangsmåden ifølge krav 1 eller krav 2, hvor en TGF-beta inhibitor yderligere er til stede i mediet og, eventuelt, hvor TGF-beta-inhibitoren er en lille molekyleinhibitor valgt fra gruppen bestående af: A83-01, SB-431542, SB-505124, SB-525334, LY 364947, SD-208, SJN 2511.
- 4.** Fremgangsmåden ifølge et hvilket som helst af kravene 1 til 4, hvor fremgangsmåden omfatter dyrkning af den ene eller flere epitelstamceller i et celledyrkningsmedium omfattende eller bestående af et basalmedium til dyre- eller humane celler til hvilke tilsættes EGF, en BMP inhibitor, R-spondin og Wnt; og efterfølgende i et ekspansionsmedium der omfatter eller består af et basalmedium til dyre- eller humane celler til hvilke tilsættes: EGF, en FGF, og HGF, nicotinamid, og et R-spondin; og efterfølgende i et differentieringsmedium der omfatter eller består af et basalmedium til dyre- eller humane celler til hvilke tilsættes: EGF, FGF og/eller HGF, en TGF-beta-inhibitor, og en Notch-inhibitor.
- 5.** Fremgangsmåden ifølge krav 4, hvor differentieringsmediet yderligere omfatter dexamethason.

6. Celledyrkningsmedium omfattende et basalmedium til dyre- eller humane celler til hvilke tilsættes: EGF, en FGF der er i stand til at binde til FGFR2 eller FGFR4, nicotinamid, og en Wnt-agonist valgt fra R-spondin 1, R-spondin 2, R-spondin 3 og R-spondin 4.

5

7. Celledyrkningsmedium ifølge til krav 6, hvor HGF tilsættes til mediet.

8. Celledyrkningsmedium ifølge krav 6 eller 7, hvor én, to, tre, fire eller flere fra gruppen gastrin, B27, N2, og N-acetylcystein tilsættes til basalmediet.

10

9. Leverorganoid omfattende celler der udtrykker Lgr5, opnåelige ved dyrkning af én eller flere leverepitelstamceller, eller et leverfragment eller en levergaldegang, i kontakt med en ekstracellulær matrix i tilstedeværelse af et

ekspansionsmedium, mediet omfatter et basal medium til dyre- eller humane

15 celler til hvilke tilsættes: en epidermal vækstfaktor (EGF), en fibroblast vækstfaktor (FGF) der er i stand til at binde til FGFR2 eller FGFR4, nicotinamid, en Wnt-agonist valgt fra Rspondin 1, Rspondin 2, Rspondin 3 eller Rspondin 4, and HGF, hvor leverorganoidet kan dyrkes i mindst 3 måneder.

20 **10.** Leverorganoidet ifølge til krav 9, der omfatter en cystisk struktur og på ydersiden, et lag af celler med mindst en knop og et centralt lumen.

11. Leverorganoidet ifølge til krav 9 eller krav 10, der har én eller flere (fx 2, 3, 4 eller alle 5) af følgende karakteristika: (a) med en celledensitet på $>5 \times 10^5$

25 celler/cm^3 , fortrinsvis $>10 \times 10^5 \text{ celler/cm}^3$; (b) med en tykkelse ækvivalent til 2-30 lag celler, fortrinsvis en tykkelse ækvivalent med 2-15 lag celler; (c) cellerne i gensidig kontakt i tre dimensioner, (d) demonstrerer en funktion inhærent med rask levervæv, (e) med en aflang form, med to definerede domæner.

30 **12.** Leverorganoidet ifølge et hvilket som helst af kravene 9 til 11, hvor:

(a) organoidet omfatter et enkeltlaget epiteldomæne hvor polariserede celler kan påvises og keratinmarkører udtrykkes og et andet domæne, der udgør hovedlegemet af organoidet, dannet af flerlaget epitel med ikke-polariserede celler, eventuelt udtrykkende albumin, og/eller

(b) organoidet opnås fra en epitelstamcelle fra en voksen lever udtrykkende Lgr5, og som omfatter en population på mellem mindst 1×10^3 celler og 5×10^4 .

5 **13.** Anvendelse af et leverorganoid ifølge et hvilket som helst af kravene 9 til 12 i en screening til opdagelse af nye lægemidler.

14. Anvendelse af et leverorganoid ifølge et hvilket som helst af kravene 9 til 12 i et toxicitetsassay.

10

15. Leverorganoidet ifølge et hvilket som helst af kravene 9 til 12 til anvendelse i behandling af en leverlidelse, tilstand eller sygdom eller til anvendelse i regenerativ medicin.

15

DRAWINGS




Fig 1a

Fig 1b




Fig 1c






Fig 1d*Fig 2a*



Fig 2b*Fig 3*

Fig 4a




Fig 4b






Fig 5a*Fig 5b*



Fig 5c*Fig 5d*

Fig 6a




Fig 6b

	ER	ENR	ENRW	Liver (adult)
G6pc2	119.5	122	125	117.5
G6pc3	4679.5	4340.5	4139.5	685.5
Glul	923.5	839.5	1387	18345
Met	5337	5743.5	4122	1828.5
Hnf1a	10980	9865	6176	8334
Hnf1b	2357	2632	2770.5	361
Hnf4a	1336	1283	1086	4033
Cyp39a1	1190.5	1167	1391	1051
Cyp2j6	938.5	918	998.5	1164
Cyp3a13	344	426.5	244	10787
Cyp4b1	1567	1567	370.5	2247
Lap3	4704	4659.5	5069.5	20262
Krt7	163790	161041	195354	386.5
Krt19	88264	79467.5	54949.5	347

Figure 7

Mouse liver culture- 24months old

A



B

n chromosomes	counts
40	10/15
80	2/15
42	1/15
36	1/15
85	1/15






Figure 8

A



B




Figure 8

C After first split






Figure 8

D




E



5 days after differentiation

Figure 9



HC specific TF

Gata6	4981,5	5381	5124	5184	740
Cebpa	1431	1188	856	946	10640
Hnf1a	11640	10980	9865	6176	8334
Hnf4a	2573	1336	1283	1086	4033

Prox1	495	219,5	379,5	319	2139
Tbx3	196,5	214	208	180	1840,5
Nr5a2	132	139	131,5	141	1234



Notch signaling

Jag1	6832,5	3638,5	4111	3394	343
Dll1	4035	3758	2791,5	2714,5	415,5
Notch3	557	718	476	788	234

TGFB signaling

Tgfb1	3439	2971	4281	3664	733
Tgfb1	245	234	243	210	116
Tgfb2	8442	9942	13874	11392	1364


Figure 10



E=EGF 50ng/ml
A=A8301 50nM
D=DAPT 10nM
F=FGF10 100ng/ml
H=HGF 50ng/ml

Nic=Nicotinamide 10nM
R=Rspondin1 500ng/ml
Wnt= Wnt conditioned media 50%

Figure 10



E=EGF 50ng/ml

A=A8301 50nM

D=DAPT 10nM

F=FGF10 100ng/ml

H=HGF 50ng/ml

Nic=Nicotinamide 10nM

R=Rspondin1 500ng/ml

Wnt= Wnt conditioned media 50%

De= dexamethasone 10mM

Figure 10E

Staining for liver markers after 8 days of differentiation condition EADF




Figure 10F

F Albcreert2LacZ derived organoids, (tamoxifen inducible system)







Figure 11 - Human-derived liver cultures ERFHNic culture conditions



5 days after seeding 4x



P1 (day 10) 4x



P2 (day 20) 4x




Figure 12

Figure 13A





Figure 13B






Figure 14




Figure 16A






Figure 16C

	Adult liver	Adult pancreas	Liver pancreas organoid
Ductal markers			
ref Mus musculus leucine aminopeptidase 3 (Lap3), mRNA [NM_024434]	11825	922	3286
ref Mus musculus keratin 19 (Krt19), mRNA [NM_008471]	302	566	76191
ref Mus musculus keratin 7 (Krt7), mRNA [NM_033073]	237	369	147227
ref Mus musculus SRY-box containing gene 9 (Sox9), mRNA [NM_011448]	235	185	2365
ref Mus musculus HNF1 homeobox B (Hnf1b), mRNA [NM_009330]	234	115	608
ref Mus musculus one cut domain, family member 1 (Onecut1), mRNA [NM_008262]	1239	120	1297
Transcription Factors necessary for Ngn3 expression			
ref Mus musculus SRY-box containing gene 9 (Sox9), mRNA [NM_011448]	235	185	2365
ref Mus musculus forkhead box A2 (Foxa2), mRNA [NM_010446]	8695	1415	12948
ref Mus musculus HNF1 homeobox B (Hnf1b), mRNA [NM_009330]	234	115	608
ref Mus musculus one cut domain, family member 1 (Onecut1), mRNA [NM_008262]	1239	120	1297
Endocrine markers			
ref Mus musculus neurogenin 3 (Neurog3), mRNA [NM_009719]	226.5	108	110
gb Mouse hepatocyte nuclear factor 1 (HNF-1) mRNA, complete cds. [M57966]	5709.5	693	4929
ref Mus musculus hepatic nuclear factor 4, alpha (Hnf4a), mRNA [NM_008261]	2947	90.5	385
ref Mus musculus pancreatic and duodenal homeobox 1 (Pdx1), mRNA [NM_008814]	137	102	201.5
ref Mus musculus neurogenic differentiation 1 (Neurod1), mRNA [NM_010894]	87	76	88
ref Mus musculus ISL1 transcription factor, LIM/homeodomain (Isl1), mRNA [NM_021459]	82	88.5	93
ref Mus musculus NK2 transcription factor related, locus 2 (Drosophila) (Nkx2-2), mRNA [NM_010919]	100	105	158
ref Mus musculus NK2 transcription factor related, locus 6 (Drosophila) (Nkx2-6), mRNA [NM_010920]	159	93	94
ref Mus musculus paired box gene 6 (Pax6), mRNA [NM_013627]	157	88.5	100.5
ref Mus musculus insulin II (Ins2), mRNA [NM_008387]	195.5	113183	107.5
ref Mus musculus solute carrier family 2 (facilitated glucose transporter), member 2 (Slc2a2), mRNA [NM_031197]	1212.5	100	188
ref Mus musculus pyruvate kinase liver and red blood cell (Pklr), transcript variant 1, mRNA [NM_013631]	237	124.5	288
ref Mus musculus glucokinase (Gck), mRNA [NM_010292]	1771	184	261

Figure 17A




Figure 17B

Flow Cytometry analysis for Hepatocyte (HC) surface marker on transplanted cells derived from Clone 1




Figure 17C

Flow Cytometry analysis for Hepatocyte (HC) surface marker on transplanted cells derived from Clone 2





Figure 17D

Mouse model: adult FGR MICE (FAH-/-RAG-/-IL2R-/-). Transplant of organoids or hepatocytes as Control

Transplantation schedule---



EM1= expansion media 1=ENRWFHnic

EM2= expansion media 2= ERFHnic

DM= differentiation media= EADF

Figure 18

<u>stem cell signature mouse liver</u>		<u>liver/HC CG specific signature</u>		<u>reprogramming genes</u>	
<u>positive for</u>	<u>negative for</u>	<u>positive for in expansion</u>	<u>negative for</u>	<u>positive</u>	<u>negative</u>
Igf5	Igf6	Hnf1a	afp	Klf4	Pou5f1
Igf4		Hnf1b	Ins1	Myc	Sox2
epcam		Hnf4a	Ins2		
Cd44		Hhex	Gcg		
Tnfrsf19		Onecut1	Ptf1a		
Sox9		Onecut2	Cela1		
Sp5		Prox1	Cela2a		
Cd24a		Cdh1	Cela3b		
Prom1		Foxa2	Neurod1		
Cdca7		Gata6	Neurod2		
Elf3		Foxm1	Neurog1		
		Cebpa	Neurog2		
		Cebpb	Neurog3		
		Cebpd	Amy2a4		
		Cebpg	Igf1r		
		Glul	Igf2		
		Krt7	Cd34		
		Krt19			
		Met			

Figure 19

*background cutoff >3.5fold (gMedian); dyeswap to Cy3/Cy5 (Cy3-samples; Cy5-reference RNA)														
Feature Number	Systematic Name	Description	Gene Name	ENRW_pat1051	ER_pat1051	EADF_pat1051	liver tissue pat1051	ENRW_pat1051	gMedian Signal	ER_pat1051	gMedian Signal	EADF_pat1051	gMedian Signal	liver tissue pat1051
		refHomo sapiens leucine-rich repeat-containing G protein-coupled receptor 5 (LGR5), mRNA [NM_003667]	LGR5						85	82.5	83			
40197	NM_003667	refHomo sapiens leucine-rich repeat-containing G protein-coupled receptor 4 (LGR4), mRNA [NM_018490]	LGR4	1.3710	1.5826	1.8852	1.1145		1602	1684	2689			80
40602	NM_018490	refHomo sapiens tumor-associated calcium signal transducer 1 (TACSTD1), mRNA [NM_002354]	TACSTD1/IE pcam	3.0978	3.5264	4.2412	-3.2251		16341	23939	42144			118
4087	NM_002354	refHomo sapiens CD44 molecule (Indian blood group) (CD44), transcript variant 1, mRNA [NM_000610]	CD44	0.8742	0.8495	-1.5025	-3.8416		10117	11624.5	2586.5			146.5
18697	NM_000610	refHomo sapiens tumor necrosis factor receptor superfamily, member 19 (TNFRSF19), transcript variant 1, mRNA [NM_018647]	TNFRSF19						83	85.5	81			81
31576	NM_018647	refHomo sapiens SRY (sex determining region Y)-box 9 (campomelic dysplasia, autosomal sex-reversal) (SOX9), mRNA [NM_000346]	SOX9	2.2553	2.3538	1.5394	-2.0445		484	448.5	355			88
2504	NM_000346	refHomo sapiens SP5 transcription factor (SP5), mRNA [NM_007003845]	SP5	-0.0509	0.4409	0.0402	3.3653		162	217.5	199			239.5
6370	NM_001003	gbHomo sapiens CD24 signal transducer mRNA, complete cds and 3' region, [L33930]	CD24	3.2316	3.5288	1.9579	-2.4236		15141.5	17882	7049			125
40879	A_23_P8525_0													
1705	NM_005017	refHomo sapiens prominin 1 (PPOM1), mRNA [NM_005017]	PPOM1	3.2512	2.6437	2.4209	-1.5556		445	340	346			84

43572	NM_031942	reflHomo sapiens cell division cycle associated 7 (CDCA7), transcript variant 1, mRNA [NM_031942]	CDCA7	-2.4618	-3.6148	-7.7338	-5.0979	476	218	87	83
1226	NM_004433	reflHomo sapiens E74-like factor 3 (ets domain transcription factor, epithelial-specific) (ELF3), transcript variant 1, mRNA [NM_004433]	ELF3	3.3588	3.9163	4.2213	0.4499	6376.5	8700.5	14527	161
Reprogramming genes											
19017	NM_004235	reflHomo sapiens Kruppel-like factor 4 (gub) (KLF4), mRNA [NM_004235]	KLF4	0.3754	2.3397	1.9646	-0.9306	291	1176.5	919	105
6084	NM_002467	reflHomo sapiens v-myc myelocytomatosis viral oncogene homolog (avian) (MYC), mRNA [NM_002467]	MYC	-1.1416	-0.8254	-3.3457	-2.1032	3006	3896	871	323
11740	NM_002701	reflHomo sapiens POU class 5 homeobox 1 (POU5F1), transcript variant 1, mRNA [NM_002701]	POU5F1	-3.9024	-3.5281	-3.8577	-1.8681	1643	2251.5	2261	1173
34630	NM_003106	reflHomo sapiens SRY (sex determining region Y)-box 2 (SOX2), mRNA [NM_003106]	SOX2	-4.0411	-4.0509	-4.3534	-2.6317	131	131.5	129	98
Liver/HC CG specific signature: positive for expansion											
22258	NM_000545	reflHomo sapiens HNF1 homeobox A (HNF1A), mRNA [NM_000545]	HNF1A	0.1593	0.4237	0.3115	2.1337	287.5	366.5	367.5	240
19971	NM_000458	reflHomo sapiens HNF1 homeobox B (HNF1B), mRNA [NM_000458]	HNF1B	3.3524	3.3315	3.1792	0.0000	200.5	200	141	86
7418	NM_000457	reflHomo sapiens hepatocyte nuclear factor 4, alpha (HNF4A), transcript variant 2, mRNA [NM_000457]	HNF4A	-0.9981	1.7168	1.0493	2.0374	84.5	119.5	108	103
25738	NM_002729	reflHomo sapiens hematopoietically expressed homeobox (HHEX), mRNA [NM_002729]	HHEX	0.8347	0.9290	1.4544	0.2789	1025	1263	1931	178
28260	NM_004498	reflHomo sapiens one cut homeobox 1 (ONECUT1), mRNA [NM_004498]	ONECUT1	0.5189	2.1647	2.9438	5.8053	120	211	327	387

30379	NM_004852	ref Homo sapiens one cut homeobox 2 (ONECUT2), mRNA [NM_004852]	ONECUT2	3.4671	3.7373	3.5642	2.8300	4693	6955	6460	498
20882	NM_002763	ref Homo sapiens prospero homeobox 1 (PROX1), mRNA [NM_002763]	PROX1	2.4930	1.0227	1.1415	2.4817	98	98	97.5	101
12860	NM_004360	ref Homo sapiens cadherin 1, type 1, E-cadherin (epithelial) (CDH1), mRNA [NM_004360]	CDH1	3.3253	3.6614	4.1391	2.4935	10803	16648	25328	1076.5
33903	NM_021784	ref Homo sapiens forkhead box A2 (FOXA2), transcript variant 1, mRNA [NM_021784]	FOXA2	2.8637	3.2434	3.0108	2.6774	289.5	306	398	129.5
39806	NM_005257	ref Homo sapiens GATA binding protein 6 (GATA6), mRNA [NM_005257]	GATA6	3.0052	3.5536	3.8552	1.2789	1620	2038	3585	136
16668	NM_202002	ref Homo sapiens forkhead box M1 (FOXM1), transcript variant 1, mRNA [NM_202002]	FOXM1	-1.9203	-3.6288	-5.0418	-4.5112	329	156.5	117.5	94
9214	NM_004364	ref Homo sapiens CCAAT/enhancer binding protein (C/EBP), alpha (CEBPA), mRNA [NM_004364]	CEBPA	-0.9322	1.2903	-0.0086	4.2303	304	1257	647	1586.5
14125	NM_005194	ref Homo sapiens CCAAT/enhancer binding protein (C/EBP), beta (CEBPB), mRNA [NM_005194]	CEBPB	-0.1111	-0.2252	0.4127	2.5221	5257	5653.5	10124	5163.5
37458	NM_005195	ref Homo sapiens CCAAT/enhancer binding protein (C/EBP), delta (CEBPD), mRNA [NM_005195]	CEBPD	-0.1615	0.9115	0.4583	4.1610	1525	3232	2733	2651
145	NM_001806	ref Homo sapiens CCAAT/enhancer binding protein (C/EBP), gamma (CEBPG), mRNA [NM_001806]	CEBPG	0.9187	2.0261	1.8780	0.7494	2185	4317.5	4980	364
41435	NM_002065	ref Homo sapiens glutamate-ammonia ligase (glutamine synthetase) (GLUL), transcript variant 1, mRNA [NM_002065]	GLUL	-0.1834	0.0758	-0.8704	0.4736	1365.5	1338	876	311
30469	NM_005556	ref Homo sapiens keratin 7 (KRT7), mRNA [NM_005556]	KRT7	5.1396	4.8373	4.3723	-0.0401	7538.5	4712	2627	96
13920	NM_002276	ref Homo sapiens keratin 19 (KRT19), mRNA [NM_002276]	KRT19	4.1830	3.7196	3.6914	-2.4731	365708	424094	308350	1571
5427	NM_000245	ref Homo sapiens met proto-oncogene (hepatocyte growth factor receptor) (MET), mRNA [NM_000245]	MET	1.3614	1.0945	1.2510	-0.3692	7623.5	6639	8627	406

[illegible]

1710	NM_000039	ref Homo sapiens apolipoprotein A-I (APOA1), mRNA [NM_000039]	APOA1	-3.5478	-2.8913	-2.1400	5.2314	370	536	1014.5	24580
8511	NM_002130	ref Homo sapiens 3-hydroxy-3-methylglutaryl-Coenzyme A synthase 1 (soluble) (HMGCS1), transcript variant 2, mRNA [NM_002130]	HMGCS1	1.9834	1.8579	-1.2536	1.9013	1518	1248	264	271
2331	NM_138711	ref Homo sapiens peroxisome proliferator-activated receptor gamma (PPARG), transcript variant 3, mRNA [NM_138711]	PPARG	1.9376	3.5421	1.0259	-0.9723	794	2235	489	102
42261	NM_000767	ref Homo sapiens cytochrome P450, family 2, subfamily B, polypeptide 5 (CYP2B6), mRNA [NM_000767]	CYP2B6	3.1696	3.8429	3.7438	7.0424	691	835	1376	23535
40086	NM_000772	ref Homo sapiens cytochrome P450, family 2, subfamily C, polypeptide 18 (CYP2C18), mRNA [NM_000772]	CYP2C18	7.6099	8.9559	7.5195	8.1267	441	938	458.5	231
18649	NM_000771	ref Homo sapiens cytochrome P450, family 2, subfamily C, polypeptide 9 (CYP2C9), mRNA [NM_000771]	CYP2C9	7.4072	9.3744	7.5571	10.9138	371.5	1144	765	3195
1249	NM_000775	ref Homo sapiens cytochrome P450, family 2, subfamily J, polypeptide 2 (CYP2J2), mRNA [NM_000775]	CYP2J2	2.5839	2.5692	2.5672	5.4100	839	595.5	802	682.5
37457	NM_017460	ref Homo sapiens cytochrome P450, family 3, subfamily A, polypeptide 4 (CYP3A4), mRNA [NM_017460]	CYP3A4	6.0960	6.9956	7.3093	9.3567	400.5	1260	1142	496.5
33810	ENST00000439761	gb Homo sapiens CYP3A5 mRNA, allele CYP3A5*3, exon 4B and partial cds, alternatively spliced. [AF355801]	CYP3A5	-0.3739	0.1067	0.0937	1.4438	736	962.5	1172	403.5
2423	NM_000765	ref Homo sapiens cytochrome P450, family 3, subfamily A, polypeptide 7 (CYP3A7), mRNA [NM_000765]	CYP3A7	5.5511	7.1260	6.6750	8.9567	3849	10927	9853	5989
14586	NM_007253	ref Homo sapiens cytochrome P450, family 4, subfamily F, polypeptide 8 (CYP4F8), mRNA [NM_007253]	CYP4F8	2.7132	4.1535	2.5950	5.9628	696	2541.5	825.5	1176
14415	NM_207352	ref Homo sapiens cytochrome P450, family 4, subfamily V, polypeptide 2 (CYP4V2), mRNA [NM_207352]	CYP4V2	2.9194	2.9333	2.7663	5.2822	853	995	923.5	822
29103	NM_005505	ref Homo sapiens scavenger receptor class B, member 1 (SCARB1), transcript variant 1, mRNA [NM_005505]	SCARB1	-0.8468	-0.4215	-1.0325	2.0324	14042	23763	13137	22012