



(86) Date de dépôt PCT/PCT Filing Date: 2005/01/05
(87) Date publication PCT/PCT Publication Date: 2005/07/28
(85) Entrée phase nationale/National Entry: 2006/06/08
(86) N° demande PCT/PCT Application No.: US 2005/000147
(87) N° publication PCT/PCT Publication No.: 2005/067741
(30) Priorités/Priorities: 2004/01/09 (US10/754,396);
2005/01/04 (US11/030,271)

(51) Cl.Int./Int.Cl. *A23L 3/3508* (2006.01),
A23B 4/24 (2006.01), *A23B 4/20* (2006.01),
A23B 4/30 (2006.01)

(71) Demandeur/Applicant:
ECOLAB INC., US


(72) Inventeurs/Inventors:
MAN, VICTOR F., US;
MAGNUSON, JOSHUA P., US;
HILGREN, JOHN D., US

(74) Agent: CASSAN MACLEAN

(54) Titre : PROCEDES DE LAVAGE DE VOLAILLE DURANT LE TRAITEMENT AVEC DES COMPOSITIONS D'ACIDE
PEROXYCARBOXYLIQUE A CHAINE MOYENNE
(54) Title: METHODS FOR WASHING POULTRY DURING PROCESSING WITH MEDIUM CHAIN
PEROXYCARBOXYLIC ACID COMPOSITIONS

(57) **Abrégé/Abstract:**

The present invention relates to methods for reducing microbial contamination on poultry, or on surfaces used in processing poultry, employing compositions including medium chain peroxycarboxylic acid, and to the compositions. The methods include applying a medium chain peroxycarboxylic acid composition to poultry or to the surfaces.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
28 July 2005 (28.07.2005)

PCT

(10) International Publication Number
WO 2005/067741 A1

(51) International Patent Classification⁷: **A23L 3/3508**,
A23B 4/20, 4/24, 4/30

(21) International Application Number:
PCT/US2005/000147

(22) International Filing Date: 5 January 2005 (05.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/754,396 9 January 2004 (09.01.2004) US
11/030,271 4 January 2005 (04.01.2005) US

(71) Applicant (for all designated States except US): **ECO-LAB INC.** [US/US]; Ecolab Center, 370 N. Wabasha Street, Saint Paul, MN 55102 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MAN, Victor, F.** [US/US]; 1410 Carling Drive, Apt. 207, St. Paul, MN 55108 (US). **MAGNUSON, Joshua, P.** [US/US]; 258 Syndicate Street S., St. Paul, MN 55105 (US). **HILGREN, John, D.** [US/US]; 925 Tanglewood Drive, Shoreview, MN 55116 (US).

(74) Agent: **SORENSEN, Andrew, D.**; ECOLAB INC., 840 Sibley Memorial Highway, Mendota Heights, MN 55118 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR WASHING POULTRY DURING PROCESSING WITH MEDIUM CHAIN PEROXYCARBOXYLIC ACID COMPOSITIONS

(57) Abstract: The present invention relates to methods for reducing microbial contamination on poultry, or on surfaces used in processing poultry, employing compositions including medium chain peroxycarboxylic acid, and to the compositions. The methods include applying a medium chain peroxycarboxylic acid composition to poultry or to the surfaces.

WO 2005/067741 A1

METHODS FOR WASHING POULTRY DURING PROCESSING WITH MEDIUM CHAIN PEROXYCARBOXYLIC ACID COMPOSITIONS

Field of the Invention

The present invention relates to methods for reducing microbial contamination
5 on poultry, or on surfaces used in processing poultry, employing compositions
including medium chain peroxycarboxylic acid, and to the compositions. The methods
include applying a medium chain peroxycarboxylic acid composition to poultry or to
the surfaces.

Background of the Invention

All poultry carcasses entering the processing environment are contaminated
with microorganisms, some with pathogenic bacteria such as *Salmonella*. Fecal matter
and dirt are the main sources of this contamination. As a result of such contamination,
poultry is typically washed at any of several steps during the process of converting a
15 live bird to an edible food product. Such washing aims to remove dirt, offal, blood,
viscera, other debris, and microbes from the poultry. Removing or reducing microbes
aids the safe storage and consumption of poultry, yet many existing washing
procedures fail to significantly reduce the microbe burden on poultry. The potential for
poultry skin to become cross-contaminated is worsened by the ability of all types of
20 microorganisms to adhere within only 15 seconds of contact. Once in the processing
environment, a significant number of carcasses can become cross-contaminated with
pathogens during handling, scalding, mechanical processing, and chilling. Current
methods for many of these procedures also fail to significantly reduce the microbe
burden on poultry.

25 Water used for washing or these other procedures is often used repeatedly over
time, which provides yet another opportunity for spreading, rather than reducing,
microbial burden on poultry. For example, the water becomes contaminated with
organic matter and microbes from the poultry, and the organic matter provides nutrients
for microbial growth in the water over time or through additional use. These microbes
30 can grow on and contaminate additional poultry and processing equipment. In
particular, water left untreated in a submersion bath tends to decontaminate poultry
early in a shift but contaminates poultry later in the shift. In fact, such water has been

identified as a potential source of coliform, *E. coli* and *Salmonella* contamination or cross contamination during poultry processing. *Salmonella* and other microorganisms are generally undesirable to the poultry, the water, and can cause buildup on all water contact surfaces of slime or biofilm, which requires frequent cleaning to remove.

5 Microbial contamination or cross contamination of poultry via water continues to be a major concern for poultry processors and end users. Although washing, cooling, or heating poultry carcasses with water can reduce potential contamination, the processing water can also serve as a source of contamination or cross contamination. If pathogenic microorganisms in water are not removed, inactivated or otherwise
10 controlled, they can spread to other poultry, potentially contaminating them. Further, handling or processing steps that pool many individual poultry parts tend to increase the risk that a single contaminated item may contaminate the entire lot. Immersing or spray-washing poultry in fresh water can help reduce surface populations of microorganisms. However sterilization by repeated washing, even with sterile water,
15 cannot be achieved because microorganisms within tissues of poultry remain in place.

 The addition of antimicrobial agents to wash or process water can inactivate vegetative bacteria cells in water, helping avoid contamination. Ideally, an antimicrobial agent or compound used in such a system will have several important properties in addition to its antimicrobial efficacy. The compound or agent should have
20 no technical effect on the final food product. Residual activity implies the presence of a film of antimicrobial material which will continue to have antimicrobial effect which may require further rinsing of the food product. The antimicrobial agent can also be odor free to prevent transfer of undesirable odors onto food stuffs. If direct food contact occurs, the antimicrobial agent should also be composed of food additive
25 materials which will not affect food wholesomeness, nor affect humans should incidental ingestion result. In addition, the antimicrobial agent can be composed of naturally occurring or innocuous ingredients, which are chemically compatible with the environment and cause no concerns for toxic residues within the water.

 In the past, poultry wash or process waters have generally been treated with
30 chlorinated compounds, organic acids, acidified sodium chlorite, trisodium phosphate, or ozone. Generally, these materials are effective in reducing microbial contamination on poultry. However, the use rate of these antimicrobials is very high because they are

not effective at low concentrations or they tend to be rapidly consumed by the high organic load included with the poultry. Excessive chlorination of food processing water with hypochlorite has prompted concern over production of toxic or carcinogenic organochlorine compounds and other by-products.

5 Further, the efficacy of conventional antimicrobial agents on the surface of poultry is often limited. For example, it has been reported that, generally, concentrations of more than 4 wt-% of organic acids or of 5 to 10 wt-% of trisodium phosphate are required to effectively reduce contamination of poultry skin by *S. typhimurium*. Antimicrobial agents such as peroxides or lactic acid can result in
10 discoloring, bleaching, or bloating of poultry tissue.

The EPA approved a peroxyacetic acid-based composition in 1996 for controlling microbial growth and reducing biofilm formation in fruit and vegetable transport or process waters. From a historical perspective, peroxyacetic acid has been used for food contact surface sanitizing, aseptic packaging and medical device cold-
15 sterilization. In addition to its biocidal properties, the environmentally-friendly decomposition byproducts and good stability in the presence of organic matter helped gain acceptance of this technology among fruit and vegetable packers, handlers, and processors. Such conventional peroxycarboxylic acid compositions typically include short chain peroxycarboxylic acids or mixtures of short chain peroxycarboxylic acids
20 and medium chain peroxycarboxylic acids (see, e.g., U.S. Patent Nos. 5,200,189, 5,314,687, 5,409,713, 5,437,868, 5,489,434, 6,674,538, 6,010,729, 6,111,963, and 6,514,556).

Nevertheless, there remains a need for improved antimicrobial compositions for addition to waters used for washing or processing poultry.

25

Summary of the Invention

The present invention relates to methods for reducing microbial contamination on poultry, or on surfaces used in processing poultry, employing compositions including medium chain peroxycarboxylic acid, and to the compositions. The methods
30 include applying a medium chain peroxycarboxylic acid composition to poultry or to the surface. The compositions and methods of the invention provide an antimicrobial agent useful in water for washing or processing poultry, that has a high degree of

antimicrobial efficacy, and that is safely ingestible by humans while imposing no unacceptable environmental incompatibility.

The compositions of the invention and other medium chain peroxycarboxylic acid antimicrobial compositions can be employed in methods for reducing microbial contamination on poultry and in water used for washing or processing poultry. These methods include applying to the poultry during processing a medium chain peroxycarboxylic acid antimicrobial composition, for example in an amount and time sufficient to reduce the microbial population. The composition can be applied by methods including submersing, rinsing, spraying, or air chilling the poultry, or a combination of these routes. During processing, the composition can be applied to whole, dismembered, portioned, or boned poultry.

In an embodiment, the method includes recovering a medium chain peroxycarboxylic acid antimicrobial composition previously applied to poultry. The recovered composition can be treated by adding a sufficient amount of a medium chain peroxycarboxylic acid to yield a recycled medium chain peroxycarboxylic acid antimicrobial composition. The recycled medium chain composition includes a reduced level of microbes, such as human pathogens, and can be disposed of more safely. Alternatively, the recycled medium chain composition can be applied to poultry during processing. In an embodiment, the medium chain peroxycarboxylic acid added to form the recycled composition is formed by adding a concentrate composition of medium chain peroxycarboxylic acid to form a composition with suitable use antimicrobial levels of the medium chain peroxycarboxylic acid.

In an embodiment, water (e.g., medium chain peroxycarboxylic acid antimicrobial composition) that has been used to wash poultry can be recovered and recycled to wash poultry or the apparatus or plant used to transport or process poultry. In an embodiment, the method of recycling poultry wash water includes employing wash water recovered from one or more wash procedures at an earlier wash procedure in poultry processing. In an embodiment, the method of recycling poultry wash water includes employing wash water recovered from one or more wash procedures for washing apparatus in the poultry processing plant or parts of the plant itself.

In an embodiment, the antimicrobial composition of the present invention includes medium chain peroxycarboxylic acid, solubilizer, oxidizing agent, and

acidulant. Such a composition can include about 0.0005 to about 5 wt-% medium chain peroxydicarboxylic acid; about 0.001 to about 10 wt-% medium chain dicarboxylic acid; about 0 to about 99.99 wt-% water; and about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium chain peroxydicarboxylic acid and the medium chain dicarboxylic acid. The composition can include a microemulsion and/or about 2 or more parts by weight of medium chain peroxydicarboxylic acid for each 7 parts by weight of medium chain dicarboxylic acid. In use form, the medium chain peroxydicarboxylic acid composition can include about 2 to about 500 ppm medium chain peroxydicarboxylic acid, about 5 to about 2000 ppm medium chain dicarboxylic acid, about 95 to about 99.99 wt-% water; and about 2 to about 16,000 ppm solubilizer.

Detailed Description of the Invention

Definitions

As used herein, the phrase “medium chain dicarboxylic acid” refers to a dicarboxylic acid that: 1) has reduced or is lacking odor compared to the bad, pungent, or acrid odor associated with an equal concentration of small chain dicarboxylic acid, and 2) has a critical micellar concentration greater than 1 mM in aqueous buffers at neutral pH. Medium chain dicarboxylic acids exclude dicarboxylic acids that are infinitely soluble in or miscible with water at 20 °C. Medium chain dicarboxylic acids include dicarboxylic acids with boiling points (at 760 mm Hg pressure) of 180 to 300 °C. In an embodiment, medium chain dicarboxylic acids include dicarboxylic acids with boiling points (at 760 mm Hg pressure) of 200 to 300 °C. In an embodiment, medium chain dicarboxylic acids include those with solubility in water of less than 1 g/L at 25 °C. Examples of medium chain dicarboxylic acids include pentanedioic acid, hexanedioic acid, heptanedioic acid, octanedioic acid, nonanedioic acid, decanedioic acid, undecanedioic acid, and dodecanedioic acid.

As used herein, the phrase “medium chain peroxydicarboxylic acid” refers to the peroxydicarboxylic acid form of a medium chain dicarboxylic acid.

As used herein, the phrase “short chain dicarboxylic acid” refers to a dicarboxylic acid that: 1) has characteristic bad, pungent, or acrid odor, and 2) is infinitely soluble in or miscible with water at 20 °C. Examples of short chain dicarboxylic acids include formic acid, acetic acid, propionic acid, and butyric acid.

As used herein, the phrase “short chain peroxycarboxylic acid” refers to the peroxycarboxylic acid form of a short chain carboxylic acid.

As used herein, the term “solubilizer” refers to a component of the present compositions to that makes soluble or increases the solubility in a carrier (e.g., water) of the medium chain carboxylic acid, medium chain peroxycarboxylic acid, or mixture thereof. For example, in an embodiment, the solubilizer can keep a composition including medium chain carboxylic acid, medium chain peroxycarboxylic acid, or mixture thereof in solution or can keep the composition finely and evenly dispersed under ordinary storage conditions without forming a separate layer. The solubilizer can, for example, solubilize a medium chain carboxylic acid to an extent sufficient to allow it to react with an oxidizing agent, such as hydrogen peroxide. A solubilizer can be identified by a test that measures phase separation under ordinary storage conditions, such as room temperature, 100 °F, or 60 °C. As used herein, the term “solubilizer” does not include short chain carboxylic acids; they are not solubilizers.

As used herein, the term “microemulsion” refers to a thermodynamically stable dispersion of one liquid phase into another stabilized by an interfacial film of surfactant. The dispersion can be oil-in-water or water-in-oil. Microemulsions are typically clear solutions when the droplet diameter is approximately 100 nanometers or less. In an embodiment, the present microemulsion composition is a shear thinning viscoelastic gel that has a blue tyndall appearance.

As used herein, the phrases “blue tyndall appearance” or “blue tyndall” refer to a bluish hue due to scattering of blue light or the blue region of the light spectrum.

As used herein, the phrases “viscoelastic gel” and “viscoelastic liquid” refer to a liquid composition that exhibits both viscous and elastic characteristics or responses, which is indicative of long range order or structure.

As used herein, a composition or combination “consisting essentially” of certain ingredients refers to a composition including those ingredients and lacking any ingredient that materially affects the basic and novel characteristics of the composition or method. The phrase “consisting essentially of” excludes from the claimed compositions and methods short chain carboxylic acids, short chain peroxycarboxylic acids, or mixtures thereof; unless such an ingredient is specifically listed after the phrase.

As used herein, a composition or combination “substantially free of” one or more ingredients refers to a composition that includes none of that ingredient or that includes only trace or incidental amounts of that ingredient. Trace or incidental amounts can include the amount of the ingredient found in another ingredient as an impurity or that is generated in a minor side reaction during formation or degradation of the medium chain peroxy-carboxylic acid.

As used herein, the phrase “a level insufficient to solubilize” refers to a concentration of an ingredient at which the ingredient is not sufficient to solubilize an insoluble material and to keep the composition substantially in one phase.

As used herein, the phrases “objectionable odor”, “offensive odor”, or “malodor” refer to a sharp, pungent, or acrid odor or atmospheric environment from which a typical person withdraws if they are able to. Hedonic tone provides a measure of the degree to which an odor is pleasant or unpleasant. An “objectionable odor”, “offensive odor”, or “malodor” has an hedonic tone rating it as unpleasant as or more unpleasant than a solution of 5 wt-% acetic acid, propionic acid, butyric acid, or mixtures thereof.

As used herein, the term “microorganism” refers to any noncellular or unicellular (including colonial) organism. Microorganisms include all prokaryotes. Microorganisms include bacteria (including cyanobacteria), lichens, fungi, protozoa, virinos, viroids, viruses, phages, and some algae. As used herein, the term “microbe” is synonymous with microorganism.

As used herein the term “poultry” refers to all forms of any bird kept, harvested, or domesticated for meat or eggs, and including chicken, turkey, ostrich, game hen, squab, guinea fowl, pheasant, quail, duck, goose, emu, or the like and the eggs of these birds. Poultry includes whole, sectioned, processed, cooked or raw poultry, and encompasses all forms of poultry flesh, by-products, and side products. The flesh of poultry includes muscle, fat, organs, skin, bones and body fluids and like components that form the animal. Forms of animal flesh include, for example, the whole or part of animal flesh, alone or in combination with other ingredients. Typical forms include, for example, processed poultry meat, such as cured poultry meat, sectioned and formed products, minced products, finely chopped products and whole products.

As used herein, the phrase “poultry processing surface” refers to a surface of a tool, a machine, equipment, a structure, a building, or the like that is employed as part of a poultry processing, preparation, or storage activity. Examples of poultry processing surfaces include surfaces of poultry processing or preparation equipment, of poultry processing wares or utensils, and of floors, walls, or fixtures of structures in which poultry processing occurs. Poultry processing utensils or tools include stunning apparatus, knife, sharpening steel, sharpening stone, scabbard and belt for holding knives, meat saw, cleaver, conveyor, bucket, working platform, scalding barrel or tank, pot, barrel or system for boiling water, scrapers, scraping table or platform, thermometer, hook, bleeding hooks, blood-catching trough, wash trough, hand wash-basin, rack, table, tank, cutting table, tote, bin, wrapping table, paper or plastic foil/bags for poultry wrapping, tool holder, metal mask, safety glove, boning apron, safety apron, and the like.

As used herein, the phrase “poultry processing machinery” refers to equipment used from first processing or slaughter through chilling and includes conveyor, shackle, killing machine, blood tunnel, scalding machine, picking machine, post-picking rinse cabinet, feet “hock” cutter, head cutter, neck breaker/oil (preen) gland remover, vent cutter, opening machine, evisceration machine, pack puller, crop remover, lung remover, carcass washer, inside-out washer, antimicrobial rinse cabinet, chiller; and also refers to equipment used for second processing or processing parts (e.g., deboning, and portion control) and including carcass splitter, deboner, cutters, and packaging.

As used herein, the phrase “poultry debris” refers to any debris, residue, material, dirt, offal, poultry part, poultry waste, poultry viscera, poultry organ, fragments or combinations of such materials, and the like removed from a poultry carcass or portion during processing and that enters a waste stream.

As used herein, the terms “mixed” or “mixture” when used relating to “peroxycarboxylic acid composition” or “peroxycarboxylic acids” refer to a composition or mixture including more than one peroxycarboxylic acid, such as a composition or mixture including peroxyacetic acid and peroxyoctanoic acid.

As used herein, the phrase “densified fluid” refers to a fluid in a critical, subcritical, near critical, or supercritical state. The fluid is generally a gas at standard conditions of one atmosphere pressure and 0 °C. As used herein, the phrase

“supercritical fluid” refers to a dense gas that is maintained above its critical temperature, the temperature above which it cannot be liquefied by pressure. Supercritical fluids are typically less viscous and diffuse more readily than liquids. In an embodiment, a densified fluid is at, above, or slightly below its critical point. As
5 used herein, the phrase “critical point” is the transition point at which the liquid and gaseous states of a substance merge into each other and represents the combination of the critical temperature and critical pressure for a substance. The critical pressure is a pressure just sufficient to cause the appearance of two phases at the critical
10 temperature. Critical temperatures and pressures have been reported for numerous organic and inorganic compounds and several elements.

As used herein, the terms "near critical" fluid or "subcritical" fluid refer to a fluid material that is typically below the critical temperature of a supercritical fluid, but remains in a fluid state and denser than a typical gas due to the effects of pressure on the fluid. In an embodiment, a subcritical or near critical fluid is at a temperature
15 and/or pressure just below its critical point. For example, a subcritical or near critical fluid can be below its critical temperature but above its critical pressure, below its critical pressure but above its critical temperature, or below both its critical temperature and pressure. The terms near critical and subcritical do not refer to materials in their ordinary gaseous or liquid state.

As used herein, the term "about" modifying the quantity of an ingredient in the compositions of the invention or employed in the methods of the invention refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use
20 solutions in the real world; through inadvertent error in these procedures; through differences in the manufacture, source, or purity of the ingredients employed to make the compositions or carry out the methods; and the like. The term about also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial mixture. Whether or not modified by the term "about", the claims include equivalents to the quantities.

30 Differentiation of antimicrobial "-cidal" or "-static" activity, the definitions which describe the degree of efficacy, and the official laboratory protocols for measuring this efficacy are considerations for understanding the relevance of

antimicrobial agents and compositions. Antimicrobial compositions can effect two kinds of microbial cell damage. The first is a lethal, irreversible action resulting in complete microbial cell destruction or incapacitation. The second type of cell damage is reversible, such that if the organism is rendered free of the agent, it can again multiply. The former is termed microbiocidal and the later, microbistatic. A sanitizer and a disinfectant are, by definition, agents which provide antimicrobial or microbiocidal activity. In contrast, a preservative is generally described as an inhibitor or microbistatic composition.

For the purpose of this patent application, successful microbial reduction is achieved when the microbial populations are reduced by at least about 50%, or by significantly more than is achieved by a wash with water. Larger reductions in microbial population provide greater levels of protection for processed poultry product.

As used herein, a composition or combination "consisting essentially" of certain ingredients refers to a composition including those ingredients and lacking any ingredient that materially affects the basic and novel characteristics of the composition or method. The phrase "consisting essentially of" excludes from the claimed compositions and methods: a coupling agent; an ingredient that cannot be employed in food products or in food wash, handling, or processing according to U.S. government rules or regulations; and/or a peroxycarboxylic acid or carboxylic acid with 10 or more carbon atoms; unless such an ingredient is specifically listed after the phrase.

Poultry Processing Employing Medium Chain Peroxycarboxylic Acid

Compositions

The concentrate and use compositions of the present invention can be employed for a variety of antimicrobial purposes, for example as or for forming water-based systems for processing and/or washing poultry. The present compositions and methods can be employed for processing poultry and/or poultry meat at any step from gathering the live birds through packaging the final product. For example, the present compositions and methods can employed for washing, rinsing, chilling, or scalding poultry carcasses, poultry carcass parts, or poultry organs for reducing contamination of these items with spoilage/decay-causing microorganisms, and pathogenic microorganisms.

Before processing, live poultry are generally transported to and gathered at the beginning of a processing line. Poultry can be washed before entering the processing line. Processing typically begins with sacrificing the bird, for example, by electrical stunning, followed by neck cutting and bleeding. A first washing step, known as
5 scalding (e.g. submersion or immersion scalding) can follow bleeding and loosens attachment of feathers to poultry skin. Submersion scalding can be accomplished according to the methods and employing compositions of the present invention. Submersion scalding can include immersing a stunned and bled bird into a scalding hot bath of water or a liquid antimicrobial composition, for example, at a temperature of
10 about 50 to about 80 °C or about 50 to about 60 °C. The liquid antimicrobial composition in the bath can be agitated, sonicated, or pumped to increase contact of the composition with the carcass. Scalding can be conducted in a scald tank or trough, which contains the scalding liquid with sufficient liquid depth to completely submerge the poultry carcass. The carcass can be transported through the tank or trough by
15 conveyor at a speed that provides a few minutes in the scalding liquid.

According to the present invention, the scalding bath can include a medium chain peroxycarboxylic acid antimicrobial composition, for example, a composition of the present invention. In an embodiment, the scalding hot bath contains a medium chain peroxycarboxylic acid antimicrobial composition with about 2 to about 50 ppm,
20 about 10 to about 200 ppm, or about 10 to about 20 ppm of peroxycarboxylic, and amounts and additional ingredients as described herein. The scalding bath can also include one or more of the additional ingredients permitted in scalding baths.

After submersion scalding, the poultry can be picked, rinsed, and, optionally, singed before the next washing process. This second washing process is generally
25 known as "dress" rinsing, "New York dress" rinsing, or post-pick rinsing, which rinses residual feathers and follicle residues from the carcass. Dress rinsing can include spraying a picked carcass with water, for example, at a temperature of about 5 to about 30 °C. To increase contact with the carcass, the antimicrobial compositions in the spray water can be applied at higher pressures, flow rates, temperatures, or with agitation or
30 ultrasonic energy. Dress rinsing can be accomplished with a washing apparatus such as a wash or spray cabinet with stationary or moving spray nozzles. Alternatively, a

“flood”-rinsing or liquid submersion washing apparatus may be used immediately after picking.

According to the present invention, picker rinsing or dress rinsing can be accomplished employing a medium chain peroxycarboxylic acid antimicrobial composition, for example a composition of the present invention. For example, the dress rinsing can employ a medium chain peroxycarboxylic acid antimicrobial composition with about 2 to about 500, about 10 to about 200, about 50 to about 300 ppm, or about 100 to about 200 ppm of peroxycarboxylic acid, and additional ingredients as described herein.

Dress rinsing can be a final washing step before dismembering the poultry. Dismembering can include removing the head, the feet, eviscerating, and removing the neck, in any order commonly employed in poultry processing. Washing can also occur during dismembering. The dismembered and eviscerated poultry can then be subjected to a washing step known as inside-outside bird washing (IOBW). Inside-outside bird washing washes the interior (body cavity) and exterior of the bird. Inside-outside bird washing can include rinsing the interior and exterior surfaces of the carcass with streams or floods of water, for example, at a temperature of about 5 to about 30 °C. To increase contact with the carcass, the antimicrobial compositions in the spray water can be applied at higher pressures, flow rates, temperatures, or with agitation or ultrasonic energy. Inside-outside bird washing can be accomplished by an apparatus that floods the bird carcass with streams of water in the inner cavity and over the exterior of the carcass. Such an apparatus can include a series of fixed spray nozzles to apply antimicrobial composition to the exterior of the bird and a rinse probe or bayonet that enters and applies antimicrobial composition to the body cavity.

According to the present invention, inside-outside bird washing can be accomplished employing a medium chain peroxycarboxylic acid antimicrobial composition, for example, a composition of the present invention. For example, inside-outside bird washing can employ a medium chain peroxycarboxylic acid antimicrobial composition with about 2 to about 500, about 10 to about 200, about 20 to about 200 ppm, or about 50 to about 100 ppm of peroxycarboxylic acid, and additional ingredients as described herein.

After inside-outside bird washing, both the interior and the exterior of the bird can be subjected to further decontamination. This further decontamination can be accomplished in part by a step commonly known as antimicrobial spray rinsing, sanitizing rinsing, or finishing rinsing. Such rinsing can include spraying the interior and exterior surfaces of the carcass with water, for example, at a temperature of about 5 to about 30 °C. To increase contact with the carcass, the antimicrobial compositions in the spray water can be applied using fixed or articulating nozzles, at higher pressures, flow rates, temperatures, with agitation or ultrasonic energy, or with rotary brushes. Spray rinsing can be accomplished by an apparatus such as a spray cabinet with stationary or moving spray nozzles. The nozzles create a mist, vapor, or spray that contacts the carcass surfaces.

According to the present invention, antimicrobial spray rinsing, sanitizing rinsing, or finishing rinsing can be accomplished employing a medium chain peroxycarboxylic acid antimicrobial composition, for example, a composition of the present invention. For example, spray rinsing can employ a medium chain peroxycarboxylic acid antimicrobial composition with about 2 to about 500, about 10 to about 200, about 50 to about 300 ppm, or about 100 to about 200 ppm of peroxycarboxylic acid, and additional ingredients as described herein.

After spray rinsing, the bird can be made ready for packaging or for further processing by chilling, specifically submersion chilling or air chilling. Submersion chilling both washes and cools the bird to retain quality of the meat. Submersion chilling can include submersing the carcass completely in water or slush, for example, at a temperature of less than about 5 °C, until the temperature of the carcass approaches that of the water or slush. Chilling of the carcass can be accomplished by submersion in a single bath, or in two or more stages, each of a lower temperature. Water can be applied with agitation or ultrasonic energy to increase contact with the carcass. Submersion chilling can be accomplished by an apparatus such as a tank containing the chilling liquid with sufficient liquid depth to completely submerge the poultry carcass. The carcass can be conveyed through the chiller by various mechanisms, such as an auger feed or a drag bottom conveyor. Submersion chilling can also be accomplished by tumbling the carcass in a chilled water cascade.

According to the present invention, submersion chilling can be accomplished employing a medium chain peroxycarboxylic acid antimicrobial composition, for example, a composition of the present invention. For example, submersion chilling can employ a medium chain peroxycarboxylic acid antimicrobial composition with about 2 to about 500, about 10 to about 200, about 2 to about 100 ppm, or about 2 to about 30 ppm of peroxycarboxylic acid, and additional ingredients as described herein.

Like submersion chilling, air chilling or cryogenic chilling cools the bird to retain quality of the meat. Air cooling can be less effective for decontaminating the bird, as the air typically would not dissolve, suspend, or wash away contaminants. Air chilling with a gas including an antimicrobial agent can, however, reduce the burden of microbial, and other, contaminants on the bird. Air chilling can include enclosing the carcass in a chamber having a temperature below about 5 °C until the carcass is chilled. Air chilling can be accomplished by applying a cryogenic fluid or gas as a blanket or spray.

According to the present invention, air chilling can be accomplished employing a medium chain peroxycarboxylic acid antimicrobial composition, e.g., a composition of the present invention. For example, air chilling compositions can include a gaseous or densified fluid medium chain peroxycarboxylic acid antimicrobial composition.

After chilling, the bird can be subjected to additional processing steps including post-chill submersion, post-chill spray, weighing, quality grading, allocation, portioning, deboning, and the like. This further processing can also include methods or compositions according to the present invention for washing with medium chain peroxycarboxylic acid compositions. For example, it can be advantageous to wash poultry portions, such as legs, breast quarters, wings, and the like, formed by portioning the bird. Such portioning forms or reveals new meat, skin, or bone surfaces which may be subject to contamination and benefit from treatment with medium chain peroxycarboxylic acid antimicrobial composition. Similarly, deboning a poultry carcass or a portion of a poultry carcass can expose additional areas of the meat or bone to microbial contamination. Washing the deboned poultry carcass or portion with a medium chain peroxycarboxylic acid composition can advantageously reduce any such contamination. In addition, during any further processing, the deboned meat can also come into contact with microbes, for example, on contaminated surfaces. Washing the

deboned meat with a medium chain peroxycarboxylic acid composition can reduce such contamination. Washing can be accomplished by spraying, immersing, tumbling, or a combination thereof, or by applying a gaseous or densified fluid antimicrobial composition.

5 Usable side products of poultry include heart, liver, and gizzard (e.g. giblets), neck, feet, and the like. These are typically harvested later in processing, and are sold as food products. Of course, microbial contamination of such food products is undesirable. Thus, these side products can also be washed with a medium chain peroxycarboxylic acid composition in methods of the present invention. Typically,
10 these side products will be washed after harvesting from the poultry carcass and before packaging. They can be washed by submersion or spraying, or transported in a flume including the antimicrobial composition. They can be contacted with an antimicrobial composition according to the invention in a giblet chiller or ice chiller.

 The poultry, poultry product, poultry portion, poultry side product, or the like
15 can be packaged before sending it to more processing, to another processor, into commerce, or to the consumer. Any such poultry can be washed with a water based medium chain peroxycarboxylic acid antimicrobial composition, which can then be removed (e.g., drained, blown, or blotted) from the poultry. In certain circumstances wetting the poultry before packaging is disadvantageous. In such circumstances, a
20 gaseous or densified fluid form of the peroxycarboxylic acid antimicrobial composition can be employed for reducing the microbial burden on the poultry. Such a gaseous composition can be employed in a variety of processes known for exposing poultry to a gas before or during packaging, such as modified atmosphere packaging.

 The advantageous stability of medium chain peroxycarboxylic acid
25 compositions in such methods, which include the presence of poultry debris or residue, makes these compositions competitive with cheaper, less stable, and potentially toxic chlorinated compounds. Embodiments of methods of the present invention include agitation or sonication of the use composition, particularly as a concentrate is added to water to make the use composition. In an embodiment, the present methods include
30 water systems that have some agitation, spraying, or other mixing of the solution. The poultry product can be contacted with the compositions of the invention effective to result in a reduction significantly greater than is achieved by washing with water, or at

least a 50% reduction, at least a 90% reduction, or at least a 99% reduction in the resident microbial preparation.

The present methods can employ a certain minimal contact time of the composition with poultry for occurrence of significant antimicrobial effect. The contact time can vary with concentration of the use composition, method of applying the use composition, temperature of the use composition, amount of soil on the poultry, number of microorganisms on the poultry, or the like. In an embodiment, the exposure time is at least about 5 to about 15 seconds.

10 Spraying Poultry

An embodiment of the present method for washing poultry employs a pressure spray of the medium chain peroxycarboxylic acid composition. During application of the spray solution on the poultry product, the surface of the poultry product can be moved with mechanical action, e.g., agitated, rubbed, brushed, etc. Agitation can be by physical scrubbing of the poultry product, through the action of the spray solution under pressure, through sonication, or by other methods. Agitation increases the efficacy of the spray solution in killing micro-organisms, perhaps due to better exposure of the solution into the crevasses or small colonies containing the micro-organisms. The spray solution, before application, can also be heated to a temperature of about 15 to 20 °C or about 20 to 60 °C to increase efficacy.

Application of the material by spray can be accomplished using a manual spray wand application, an automatic spray of poultry product moving along a production line using multiple spray heads to ensure complete contact, or other spray apparatus. One automatic spray application involves the use of a spray booth. The spray booth substantially confines the sprayed composition to within the parameter of the booth. The production line moves the poultry product through the entryway into the spray booth in which the poultry product is sprayed on all its exterior surfaces with sprays within the booth. After a complete coverage of the material and drainage of the material from the poultry product within the booth, the poultry product can then exit the booth in a fully treated form. The spray booth can include steam jets that can be used to apply the antimicrobial compositions of the invention. These steam jets can be used in combination with cooling water to ensure that the treatment reaching the poultry

product surface is less than 65°C, or less than 60°C. The temperature of the spray on the poultry product can ensure that the poultry product is not substantially altered (cooked) by the temperature of the spray. The spray pattern can be virtually any useful spray pattern.

5

Immersing Poultry

During processing of the poultry product, the poultry product can be immersed into a tank containing a quantity of washing solution. The washing solution can be agitated to increase the efficacy of the solution and the speed in which the solution reduces micro-organisms accompanying to the poultry product. Agitation can be obtained by conventional methods, including ultrasonics, aeration by bubbling air through the solution, by mechanical methods, such as strainers, paddles, brushes, pump driven liquid jets, or by combinations of these methods. The washing solution can be heated to increase the efficacy of the solution in killing micro-organisms. In an embodiment, the poultry product can be immersed in the washing solution after the poultry product has been eviscerated and before any cooling process such as a chiller tank or a chill water spray. In an embodiment, the poultry product can be immersed in the washing composition post-chill.

Foam Treating Poultry

In another alternative embodiment of the present invention, the poultry product can be treated with a foaming version of the composition. The foam can be prepared by mixing foaming surfactants with the washing solution at time of use. The foaming surfactants can be nonionic, anionic or cationic in nature. Examples of useful surfactant types include, but are not limited to the following: alcohol ethoxylates, alcohol ethoxylate carboxylate, amine oxides, alkyl sulfates, alkyl ether sulfate, sulfonates, quaternary ammonium compounds, alkyl sarcosines, betaines and alkyl amides. The foaming surfactant can be mixed at time of use with the washing solution. Use solution levels of the foaming agents is from about 50 ppm to about 2.0 wt-%. At time of use, compressed air can be injected into the mixture, then applied to the poultry product surface through a foam application device such as a tank foamer or an aspirated wall mounted foamer.

Gel Treating Poultry

In another alternative embodiment of the present invention, the poultry product can be treated with a thickened or gelled version of the composition. In the thickened or gelled state the washing solution remains in contact with the poultry product surface for longer periods of time, thus increasing the antimicrobial efficacy. The thickened or gelled solution will also adhere to vertical surfaces. The composition or the washing solution can be thickened or gelled using existing technologies such as: xanthan gum, polymeric thickeners, cellulose thickeners or the like. Rod micelle forming systems such as amine oxides and anionic counter ions could also be used. The thickeners or gel forming agents can be used either in the concentrated product or mixing with the washing solution, at time of use. Typical use levels of thickeners or gel agents range from about 100 ppm to about 10 wt-%.

Light Treating Poultry

In another alternative embodiment of the present invention, the poultry product can be exposed to an activating light (or other electromagnetic radiation) source following application of the washing solution. The activating light (or other electromagnetic radiation) can improve the antimicrobial efficacy of the washing solution. The light can be ultraviolet light, infrared light, visible light, or a combination thereof. Other forms of electromagnetic radiation include radar and microwave.

Processing Poultry Wash Water

Washing poultry can employ a large volume of water, or another carrier. Poultry wash water can be used more than once (recycled), provided the water can be treated so that it does not transfer undesirable microbes to the poultry being washed with the recycled wash water. One way to prevent the transfer of such undesirable microbes, is to reduce the microbial burden of the recycled wash water by adding a mixture of peroxycarboxylic acids. For example, if the fluid to be recycled is water-based and lacking any peroxycarboxylic acid, a medium chain peroxycarboxylic acid concentrate composition can be added to result in an effective antimicrobial concentration of peroxycarboxylic acid in the fluid to be recycled. Alternatively, if the

fluid to be recycled already includes or has included a peroxycarboxylic acid, a medium chain peroxycarboxylic acid concentrate composition can be added to increase any concentration of peroxycarboxylic acid to an effective antimicrobial level. It may be that the peroxycarboxylic acid in the solution to be recycled has been totally depleted,
5 in which case more of the medium chain peroxycarboxylic acid composition is added.

In some circumstances, the water to be recycled includes a substantial burden of organic matter or microbes. If this is the case, the water may be unsuitable for recycling. However, if the water is to be recycled, the operator adds a sufficient quantity of the medium chain peroxycarboxylic acid composition to provide an
10 effective antimicrobial amount of the peroxycarboxylic acid after a certain amount is consumed by the organic burden or microbes already present. Then, the recycled fluid can be used with antimicrobial effect. Routine testing can be employed for determining levels of peroxycarboxylic acid, or of organic burden.

In each case, the method of recycling the poultry wash water includes
15 recovering the poultry wash water, adding a composition including medium chain peroxycarboxylic acids, and reusing the poultry wash water for washing poultry, for example, as described above. The poultry wash water can be recovered from steps in poultry processing including submersion scalding, dress rinsing, inside-outside bird washing, spray rinsing, and submersion chilling. Methods of recovering wash water
20 from these steps are well-known to those skilled in the poultry washing and/or processing arts. The wash water can also be strained, filtered, diluted, or otherwise cleaned and processed during recycling.

In an embodiment, water (e.g., medium chain peroxycarboxylic acid composition) that has been used to wash poultry can be recovered and recycled to wash
25 poultry or the apparatus or plant used to transport or process poultry. In this embodiment, the water can be treated so that it does not transfer undesirable microbes to the poultry, apparatus, or plant being washed with the recycled wash water. Not transferring undesirable microbes be accomplished by employing a recycled composition that is free of pathogenic microorganisms, that is free of fecal coliform
30 organisms, or free of both. Not transferring undesirable microbes can include reducing contamination (e.g., physical, chemical, or microbiological contamination) of the recycled composition to prevent adulteration of product. Not transferring undesirable

microbes can include reducing contamination (e.g., physical, chemical, or microbiological contamination) of the recycled composition to prevent contamination or adulteration of product. The water can be treated by adding medium chain peroxycarboxylic acid concentrate composition, by filtering or straining the water, and/or by light treating the water. The method of recycling the poultry wash water then includes reusing the poultry wash water for washing poultry, for example, as described above.

In an embodiment, the method of recycling poultry wash water (e.g., medium chain peroxycarboxylic acid composition) includes employing wash water recovered from one or more wash procedures at an earlier wash procedure in poultry processing. For example, the present method can include recovering poultry wash water from submersion chilling; treating the water; and employing the treated wash water in at least one of submersion scalding, dress rinsing, inside-outside bird washing, and spray rinsing. For example, the present method can include recovering poultry wash water from spray rinsing; treating the water; and employing the treated wash water in at least one of submersion scalding, dress rinsing, and inside-outside bird washing. For example, the present method can include recovering poultry wash water from inside-outside bird washing; treating the water; and employing the treated wash water in at least one of submersion scalding and dress rinsing. For example, the present method can include recovering poultry wash water from dress rinsing; treating the water; and employing the treated wash water in submersion scalding.

In an embodiment, the method of recycling poultry wash water (e.g., medium chain peroxycarboxylic acid composition) includes employing wash water recovered from one or more wash procedures for washing apparatus in the poultry processing plant or parts of the plant itself (e.g., floors, walls, exterior pavement, or the like). For example, the present method can include recovering poultry wash water from at least one of submersion scalding, dress rinsing, inside-outside bird washing, spray rinsing, and submersion chilling; treating the water; and employing the treated water for washing processing apparatus. For example, the present method can include recovering poultry wash water from at least one of submersion scalding, dress rinsing, inside-outside bird washing, spray rinsing, and submersion chilling; treating the water; and employing the treated water for washing a portion of the processing plant, such as a

floor, a wall, or exterior pavement. For example, the present method can include recovering poultry wash water from at least one of submersion scalding, dress rinsing, inside-outside bird washing, spray rinsing, and submersion chilling; treating the water; and employing the treated water for washing trucks or cages, for example, those in which poultry was transported to the plant. For example, the present method can include recovering poultry wash water from at least one of submersion scalding, dress rinsing, inside-outside bird washing, spray rinsing, and submersion chilling; treating the water; and employing the treated water for washing poultry entering the plant.

10 Test Methods for Reducing Microbial Populations on Poultry

Spray application of an antimicrobial composition of the invention can be tested and shown to significantly reduce bacterial pathogen contamination on poultry carcass samples. Carcass samples can be contaminated with, for example, *Salmonella typhimurium* ATCC 13311, *Escherichia coli* serotype O157:H7 ATCC 43895, or *Listeria monocytogenes* (Petite Scott A) ATCC 49594. Identities of these bacteria can be confirmed based on gram stain reactions, microscopic morphology and growth characteristics using the appropriate selective medium. These strains can be grown in culture, by conventional techniques, and adjusted to yield, for example, $\geq 10^7$ colony forming units per milliliter (CFU/mL).

Carcass samples can be prepared by inoculating the exterior of a thawed piece of chicken skin with a suitable volume of a pathogen culture. The square of skin can be depressed to form a bowl, and the culture can be allowed to sit in this bowl for, for example, 5 min to allow attachment of the bacteria. After attachment, culture can be removed from the skin sample and each sample can be placed on a metal stand, epidermal side/inoculated side up.

The skin sample can be then sprayed with an antimicrobial composition. For use, the composition can be diluted to, for example, 200 ± 10 ppm total medium chain peroxycarboxylic acid. The poultry carcass sample can be sprayed with the composition for, for example, 15 seconds at 60 psi and room temperature.

After spraying, the skin samples can be aseptically removed and placed into solution including an agent that inactivates the peroxycarboxylic acids without killing bacteria. The remaining bacteria can be suspended by vortexing and serial dilutions of

this solution can be plated for growth of the bacteria. Dilutions can be prepared using phosphate buffered dilution water (PBDW). *S. typhimurium* and *L. monocytogenes* plates can be incubated at 37 °C for 48 hours. *E. coli* O157:H7 plates can be incubated at 37 °C for 24 hours. For each sample, the number of colony forming units per skin square can be calculated.

Analysis of the numbers of bacteria on skin samples can demonstrate that spray application of the antimicrobial composition of the invention significantly reduces levels of, for example, *Salmonella typhimurium*, *Escherichia coli*, and *Listeria monocytogenes*.

Submersion application of an antimicrobial composition of the invention can be tested and shows significant reduction of bacterial pathogen contamination on poultry carcass samples.

Bacteria can be selected and cultured generally as described above, except that the bacteria can be diluted to, for example, $\geq 10^6$ colony forming units per milliliter (CFU/mL) for inoculation of carcass samples.

Carcass samples can be prepared by thawing frozen chicken wings and livers. The thawed samples can be inoculated by submersing in the suspension of bacteria. Other surfaces can be not inoculated. The inoculated surface can be marked for identification and allowed to drain and sit for 5 min at room temperature.

The antimicrobial composition can be as described above. The use solution can be diluted to, for example, 30 ppm total medium chain peroxycarboxylic acid. Antimicrobial agent can be applied by submersing the inoculated and uninoculated surfaces, for example, for 60 min in the same 2 liters of a use solution of the antimicrobial agent at 4 °C. The same experiment can be performed using water without the antimicrobial agent.

After submersing, the inoculated and uninoculated surfaces can be removed from the antimicrobial agent use solution or water and gently agitated solution including an agent that inactivates the peroxycarboxylic acids without killing bacteria. Removing and analyzing the uninoculated surfaces can be necessary to measure cross contamination of bacteria from the inoculated surfaces. Serial dilutions of this solution can be plated for growth of the bacteria as described above in Example 2. Cross contamination log reduction can be calculated by subtracting the Log number of

bacteria on uninoculated surfaces submersed in the antimicrobial use solution from the Log number of surviving bacteria on uninoculated surfaces submersed in water.

Analysis of the numbers of bacteria on carcass samples can demonstrate that submersion application of the antimicrobial composition of the invention significantly reduces levels of, for example, *Salmonella typhimurium*, *Escherichia coli*, and *Listeria monocytogenes*.

Medium chain peroxycarboxylic acid antimicrobial composition in water can be used for spraying or submersing eviscerated chicken carcasses and can provide a reduction of, for example, total aerobic bacteria, coliform bacteria, and *Escherichia coli*.

Freshly collected chicken carcasses can be subjected to spraying with or submersion in an antimicrobial composition described above. Use solutions, spray time and pressure, and submersion temperature and duration can be as described above. Some carcasses can be both sprayed and submersed. Control carcasses can be untreated.

Each carcass can be then placed in a collection bag using freshly gloved hands. In the bag, the carcass can be rinsed with Butterfield's Phosphate Diluent (BPD) and the BPD solution collected for microbiology testing. Known, standard procedures for quantifying, for example, total aerobic bacteria, coliform bacteria, and *Escherichia coli* can be employed.

Analysis of the numbers of bacteria on carcass samples can demonstrate that spray, submersion, and combination application of the antimicrobial composition of the invention significantly reduces levels of, for example, total aerobic bacteria, coliform bacteria, and *Escherichia coli*.

Medium Chain Peroxycarboxylic Acid Antimicrobial Compositions

The present invention includes medium chain peroxycarboxylic acid compositions. The present medium chain peroxycarboxylic acid compositions can include increased levels of medium chain peroxycarboxylic acid compared to conventional peroxycarboxylic acid compositions. The inventive compositions can include medium chain peroxycarboxylic acid and a solubilizer. The solubilizer can increase or maintain the solubility of the medium chain peroxycarboxylic acid. The

present medium chain peroxycarboxylic acid compositions can include a microemulsion or a surfactant that can form a microemulsion. The present medium chain peroxycarboxylic acid compositions need not include substantial amounts of short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof. It is
5 believed that, in conventional mixed peroxycarboxylic acid compositions, the short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof can solubilize medium chain peroxycarboxylic acid.

In an embodiment, the present compositions include medium chain peroxycarboxylic acid. These compositions can also include medium chain carboxylic
10 acid. Such compositions can include advantageously high levels of medium chain peroxycarboxylic acid. In an embodiment, the present compositions include about 2 or more parts by weight of medium chain peroxycarboxylic acid for each 7 parts by weight of medium chain carboxylic acid. In an embodiment, the present compositions include about 2 or more parts by weight of medium chain peroxycarboxylic acid for
15 each 6 parts by weight of medium chain carboxylic acid. In an embodiment, the present compositions include about 2 or more parts by weight of medium chain peroxycarboxylic acid for each 5 parts by weight of medium chain carboxylic acid. In an embodiment, the present compositions include about 2 or more parts by weight of medium chain peroxycarboxylic acid for each 4 parts by weight of medium chain
20 carboxylic acid. In an embodiment, the present compositions include about 2 parts by weight of medium chain peroxycarboxylic acid for each 3 parts by weight of medium chain carboxylic acid.

In an embodiment, the present compositions include medium chain peroxycarboxylic acid and solubilizer. The solubilizer can include a solvent, a
25 surfactant, or a mixture thereof. Suitable solvents include any of a variety of solvents that solubilize and do not significantly degrade the medium chain peroxycarboxylic acid. In certain embodiments, suitable solvents include polyalkylene oxide, capped polyalkylene oxide, mixtures thereof, or the like. Suitable solvents include nonionic surfactant, such as alkoxyated surfactant. Suitable alkoxyated surfactants include, for
30 example, EO/PO copolymer, capped EO/PO copolymer, alcohol alkoxyate, capped alcohol alkoxyate, mixtures thereof, or the like. When employed as a solvent a

surfactant, such as a nonionic surfactant, can be at concentrations higher than those conventionally employed.

The solubilizer can include surfactant (e.g., microemulsion forming surfactant). Suitable surfactants include anionic surfactant, nonionic surfactant, cationic surfactant, amphoteric surfactant, zwitterionic surfactant, mixtures thereof, or the like. The solubilizer can include a microemulsion forming surfactant. Suitable microemulsion forming surfactants include anionic surfactant, cationic surfactant, amphoteric surfactant, zwitterionic surfactant, mixtures thereof, or the like. Suitable microemulsion forming surfactants include anionic surfactants, such as sulfate surfactant, sulfonate surfactant, phosphate surfactant (phosphate ester surfactant), and carboxylate surfactant, mixtures thereof, or the like.

In an embodiment, the present composition need not include substantial amounts of short chain peroxydicarboxylic acid. For example, the present compositions can be free of added short chain peroxydicarboxylic acid. As used herein, free of added material refers to a composition that includes the material only as a incidental or trace quantity found, for example, as an ingredient of or impurity in another named ingredient or incidentally generated from a minor side reaction.

In an embodiment, the present composition includes only relatively small amounts of short chain peroxydicarboxylic acid. For example, the present composition can include about 1 or more parts of medium chain peroxydicarboxylic acid for each 8 parts of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. For example, the present composition can include short chain peroxydicarboxylic acid at a level insufficient to cause odor offensive to a typical person.

In certain embodiments, the present composition does not include substantial amounts of peroxyacetic acid, is free of added peroxyacetic acid, includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 8 parts of peroxyacetic acid, or includes peroxyacetic acid at a level insufficient to cause odor offensive to a typical person.

In an embodiment, the present composition need not include substantial amounts of short chain dicarboxylic acid. For example, the present compositions can be free of added short chain dicarboxylic acid. In an embodiment, the present composition includes only relatively small amounts of short chain dicarboxylic acid. By way of

further example, the present composition can include about 1 or more parts of medium chain peroxydicarboxylic acid for each 8 parts of short chain dicarboxylic acid. For example, the present composition can include short chain dicarboxylic acid at a level insufficient to cause odor offensive to a typical person.

5 In certain embodiments, the present composition does not include substantial amounts of acetic acid, is free of added acetic acid, includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 8 parts of acetic acid, or includes acetic acid at a level insufficient to cause odor offensive to a typical person. In certain
10 embodiments, the present compositions include, for example, less than 10 wt-%, less than less than 5 wt-%, less than 2 wt-%, or less than 1 wt-% acetic acid. In certain
embodiments, the present use compositions include, for example, less than 40 ppm, less than 20 ppm, less than 10 ppm, or less than 5 ppm acetic acid.

 In an embodiment, the present composition need not include substantial amounts of short chain peroxydicarboxylic acid, short chain dicarboxylic acid, or mixture
15 thereof. For example, the present compositions can be free of added short chain peroxydicarboxylic acid, short chain dicarboxylic acid, or mixture thereof. For example, the present composition can include short chain dicarboxylic acid, short chain
peroxydicarboxylic acid, or mixture thereof at a level insufficient to cause odor offensive to a typical person. In certain embodiments, the present composition does not include
20 substantial amounts of acetic acid, peroxyacetic acid, or mixtures thereof; is free of added acetic acid, peroxyacetic acid, or mixtures thereof; includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 8 parts of acetic acid,
peroxyacetic acid, or mixtures thereof; or includes acetic acid, peroxyacetic acid, or mixtures thereof at a level insufficient to cause odor offensive to a typical person.

25 In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 8 parts of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for
each 7 parts of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or
30 mixture thereof. In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 6 parts of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. In an embodiment, the

present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 5 parts of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 4 parts of short chain
5 dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 3 parts of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 2 parts
10 of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof. In an embodiment, the present composition includes about 1 or more parts of medium chain peroxydicarboxylic acid for each 1 part of short chain dicarboxylic acid, short chain peroxydicarboxylic acid, or mixture thereof.

In an embodiment, the present composition has an odor less unpleasant than
15 (e.g., as measured by an hedonic tone rating) than 5, 4, 3, 2, or 1 wt-% acetic acid in water. In an embodiment, the present composition has an odor less unpleasant than (e.g., as measured by an hedonic tone rating) than 5 wt-% acetic acid in water. In an embodiment, the present composition has an odor less unpleasant than (e.g., as measured by an hedonic tone rating) than 4 wt-% acetic acid in water. In an
20 embodiment, the present composition has an odor less unpleasant than (e.g., as measured by an hedonic tone rating) than 3 wt-% acetic acid in water. In an embodiment, the present composition has an odor less unpleasant than (e.g., as measured by an hedonic tone rating) than 2 wt-% acetic acid in water. In an embodiment, the present composition has an odor with an odor less unpleasant than
25 (e.g., as measured by an hedonic tone rating) than 1 wt-% acetic acid in water.

In certain embodiments, the present composition includes one or more (e.g., at least one) of oxidizing agent, acidulant, stabilizing agent, mixtures thereof, or the like. The present composition can include any of a variety of oxidizing agents, for example, hydrogen peroxide. The oxidizing agent can be effective to convert a medium chain
30 dicarboxylic acid to a medium chain peroxydicarboxylic acid. The oxidizing agent can also have antimicrobial activity, although it may not be present at a concentration sufficient to exhibit such activity. The present composition can include any of a variety of

acidulants, for example, an inorganic acid. The acidulant can be effective to bring the pH of the present concentrate composition to less than 1, or to bring the pH of the present use composition to about 5 or below, about 4 or below, or about 3 or below. The acidulant can augment the antimicrobial activity of the present composition. The present composition can include any of a variety of stabilizing agents, for example, sequestrant, for example, phosphonate sequestrant. The sequestrant can be effective to stabilize the peroxycarboxylic acid.

In an embodiment, the present composition exhibits advantageous stability of the peroxycarboxylic acid. It is believed that in approximately one year at ambient conditions or room temperature (or 1 week at 60 °C) the amount of peroxycarboxylic acid in the compositions can be about 80% or more, about 85 % or more, about 90% or more, or about 95% or more of the initial values or use composition levels. Such aged compositions are included in the scope of the present invention.

In an embodiment, the present composition exhibits advantageous efficacy compared to other antimicrobial compositions at the same level of active. In certain embodiments, the present composition has reduced or no volatile organic compounds compared to conventional peroxycarboxylic acid compositions. In an embodiment, the present composition has a higher flash point compared to conventional peroxycarboxylic acid compositions. In an embodiment, the present composition exhibits improved operator or user safety compared to conventional peroxycarboxylic acid compositions. In an embodiment, the present composition exhibits improved storage or transportation safety compared to conventional peroxycarboxylic acid compositions.

In certain embodiments, the present composition includes about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid, about 0.3 to about 7 wt-% medium chain peroxycarboxylic acid, about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid, about 0.5 to about 4 wt-% medium chain peroxycarboxylic acid, about 0.8 to about 3 wt-% medium chain peroxycarboxylic acid, about 1 to about 3 wt-% medium chain peroxycarboxylic acid, or about 1 to about 2 wt-% medium chain peroxycarboxylic acid. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 8 wt-% medium chain carboxylic acid, about 1 to about 10 wt-% medium chain carboxylic acid, about 1 to about 8 wt-% medium chain carboxylic acid, about 1.5 to about 6 wt-% medium chain carboxylic acid, about 2 to about 8 wt-% medium chain carboxylic acid, about 2 to about 6 wt-% medium chain carboxylic acid, about 2 to about 4 wt-% medium chain carboxylic acid, about 2.5 to about 5 wt-% medium chain carboxylic acid, about 3 to about 6 wt-% medium chain carboxylic acid, or about 3 to about 5 wt-% medium chain carboxylic acid. The composition can include any of these ranges or amounts not modified by about.

10 In certain embodiments, the present composition includes about 0 to about 98 wt-% carrier, about 0.001 to about 99.99 wt-% carrier, about 0.2 to about 60 wt-% carrier, about 1 to about 98 wt-% carrier, about 5 to about 99.99 wt-% carrier, about 5 to about 97 wt-% carrier, about 5 to about 90 wt-% carrier, about 5 to about 70 wt-% carrier, about 5 to about 20 wt-% carrier, about 10 to about 90 wt-% carrier, about 10 to about 80 wt-% carrier, about 10 to about 50 wt-% carrier, about 10 to about 20 wt-% carrier, about 15 to about 70 wt-% carrier, about 15 to about 80 wt-% carrier, about 20 to about 70 wt-% carrier, about 20 to about 50 wt-% carrier, about 20 to about 40 wt-% carrier, about 20 to about 30 wt-% carrier, about 30 to about 75 wt-% carrier, about 30 to about 70 wt-% carrier, about 40 to about 99.99 wt-% carrier, about 40 to about 90 wt-% carrier, or about 60 to about 70 wt-% carrier. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 80 wt-% solubilizer, about 0.001 to about 60 wt-% solubilizer, about 1 to about 80 wt-% solubilizer, about 1 to about 25 wt-% solubilizer, about 1 to about 20 wt-% solubilizer, about 2 to about 70 wt-% solubilizer, about 2 to about 60 wt-% solubilizer, about 2 to about 20 wt-% solubilizer, about 3 to about 65 wt-% solubilizer, about 3 to about 15 wt-% solubilizer, about 4 to about 10 wt-% solubilizer, about 4 to about 20 wt-% solubilizer, about 5 to about 70 wt-% solubilizer, about 5 to about 60 wt-% solubilizer, about 5 to about 20 wt-% solubilizer, about 10 to about 70 wt-% solubilizer, about 10 to about 65 wt-% solubilizer, about 10 to about 20 wt-% solubilizer, about 20 to about 60 wt-% solubilizer, or about 40 to about 60 wt-% solubilizer. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 30 wt-% oxidizing agent, about 0.001 to about 10 wt-% oxidizing agent, 0.002 to about 10 wt-% oxidizing agent, about 2 to about 30 wt-% oxidizing agent, about 2 to about 25 wt-% oxidizing agent, about 2 to about 20 wt-% oxidizing agent, about 4 to about 20 wt-% oxidizing agent, about 5 to about 10 wt-% oxidizing agent, or about 6 to about 10 wt-% oxidizing agent. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 50 wt-% acidulant, about 0.001 to about 30 wt-% acidulant, about 1 to about 50 wt-% acidulant, about 1 to about 30 wt-% acidulant, about 2 to about 40 wt-% acidulant, about 2 to about 10 wt-% acidulant, about 3 to about 40 wt-% acidulant, about 5 to about 40 wt-% acidulant, about 5 to about 25 wt-% acidulant, about 10 to about 40 wt-% acidulant, about 10 to about 30 wt-% acidulant, about 15 to about 35 wt-% acidulant, about 15 to about 30 wt-% acidulant, or about 40 to about 60 wt-% acidulant. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 50 wt-% stabilizing agent, about 0.001 to about 5 wt-% stabilizing agent, about 0.5 to about 50 wt-% stabilizing agent, about 1 to about 50 wt-% stabilizing agent, about 1 to about 30 wt-% stabilizing agent, about 1 to about 10 wt-% stabilizing agent, about 1 to about 5 wt-% stabilizing agent, about 1 to about 3 wt-% stabilizing agent, about 2 to about 10 wt-% stabilizing agent, about 2 to about 5 wt-% stabilizing agent, or about 5 to about 15 wt-% stabilizing agent. The composition can include any of these ranges or amounts not modified by about.

25 **Compositions of Medium Chain Carboxylic Acids and/or Peroxycarboxylic Acids**

Peroxycarboxylic (or percarboxylic) acids generally have the formula $R(CO_3H)_n$, where, for example, R is an alkyl, arylalkyl, cycloalkyl, aromatic, or heterocyclic group, and n is one, two, or three, and named by prefixing the parent acid with peroxy. The R group can be saturated or unsaturated as well as substituted or unsubstituted. The composition and methods of the invention can employ medium chain peroxycarboxylic acids containing, for example, 6 to 12 carbon atoms. For example, medium chain peroxycarboxylic (or percarboxylic) acids can have the

formula $R(\text{CO}_3\text{H})_n$, where R is a $\text{C}_5\text{-C}_{11}$ alkyl group, a $\text{C}_5\text{-C}_{11}$ cycloalkyl, a $\text{C}_5\text{-C}_{11}$ arylalkyl group, $\text{C}_5\text{-C}_{11}$ aryl group, or a $\text{C}_5\text{-C}_{11}$ heterocyclic group; and n is one, two, or three.

Peroxy-carboxylic acids can be made by the direct action of an oxidizing agent on a carboxylic acid, by autoxidation of aldehydes, or from acid chlorides, and hydrides, or carboxylic anhydrides with hydrogen or sodium peroxide. In an embodiment, the medium chain percarboxylic acids can be made by the direct, acid catalyzed equilibrium action of hydrogen peroxide on the medium chain carboxylic acid. Scheme 1 illustrates an equilibrium between carboxylic acid and oxidizing agent (Ox) on one side and peroxy-carboxylic acid and reduced oxidizing agent (Ox_{red}) on the other:



Scheme 2 illustrates an embodiment of the equilibrium of scheme 1 in which the oxidizing agent is hydrogen peroxide on one side and peroxy-carboxylic acid and water on the other:



In conventional mixed peroxy-carboxylic acid compositions it is believed that the equilibrium constant for the reaction illustrated in scheme 2 is about 2.5, which may reflect the equilibrium for acetic acid. Although not limiting to the present invention, it is believed that the present compositions have an equilibrium constant of about 4.

Peroxy-carboxylic acids useful in the compositions and methods of the present invention include peroxy-pentanoic, peroxy-hexanoic, peroxy-heptanoic, peroxy-octanoic, peroxy-nonanoic, peroxy-decanoic, peroxy-undecanoic, peroxy-dodecanoic, peroxy-ascorbic, peroxy-adipic, peroxy-citric, peroxy-pimelic, or peroxy-suberic acid, mixtures thereof, or the like. The alkyl backbones of these medium chain peroxy-carboxylic acids can be straight chain, branched, or a mixture thereof. Peroxy forms of carboxylic acids with more than one carboxylate moiety can have one or more (e.g., at least one) of the carboxyl moieties present as peroxy-carboxyl moieties.

Peroxy-octanoic (or perooctanoic) acid is a peroxy-carboxylic acid having the formula, for example, of n-peroxy-octanoic acid: $\text{CH}_3(\text{CH}_2)_6\text{COOOH}$. Peroxy-octanoic acid can be an acid with a straight chain alkyl moiety, an acid with a branched alkyl

moiety, or a mixture thereof. Peroxyoctanoic acid is surface active and can assist in wetting hydrophobic surfaces, such as those of microbes.

The composition of the present invention can include a carboxylic acid. Generally, carboxylic acids have the formula $R-COOH$ wherein the R can represent any number of different groups including aliphatic groups, alicyclic groups, aromatic groups, heterocyclic groups, all of which can be saturated or unsaturated as well as substituted or unsubstituted. Carboxylic acids can have one, two, three, or more carboxyl groups. The composition and methods of the invention typically employ medium chain carboxylic acids containing, for example, 6 to 12 carbon atoms. For example, medium chain carboxylic acids can have the formula $R-COOH$ in which R can be a C_5-C_{11} alkyl group, a C_5-C_{11} cycloalkyl group, a C_5-C_{11} arylalkyl group, C_5-C_{11} aryl group, or a C_5-C_{11} heterocyclic group.

Suitable medium chain carboxylic acids include pentanoic, hexanoic, heptanoic, octanoic, nonanoic, decanoic, undecanoic, dodecanoic, ascorbic, citric, adipic, pimelic, and suberic acid. The alkyl backbones of these medium chain carboxylic acids can be straight chain, branched, or a mixture thereof. Carboxylic acids which are generally useful are those having one or two carboxyl groups where the R group is a primary alkyl chain having a length of C_4 to C_{11} . The primary alkyl chain is that carbon chain of the molecule having the greatest length of carbon atoms and directly appending carboxyl functional groups.

The present compositions and methods include a medium chain peroxycarboxylic acid. The medium chain peroxycarboxylic acid can include or be a C_6 to C_{12} peroxycarboxylic acid. The C_6 to C_{12} peroxycarboxylic acid can include or be peroxyhexanoic acid, peroxyheptanoic acid, peroxyoctanoic acid, peroxyononanoic acid, peroxydecanoic acid, peroxyundecanoic acid, peroxydodecanoic acid, or mixture thereof. The medium chain peroxycarboxylic acid can include or be a C_7 to C_{12} peroxycarboxylic acid. The C_7 to C_{12} peroxycarboxylic acid can include or be peroxyheptanoic acid, peroxyoctanoic acid, peroxyononanoic acid, peroxydecanoic acid, peroxyundecanoic acid, peroxydodecanoic acid, or mixture thereof. The medium chain peroxycarboxylic acid can include or be a C_6 to C_{10} peroxycarboxylic acid. The C_6 to C_{10} peroxycarboxylic acid can include or be peroxyhexanoic acid, peroxyheptanoic acid, peroxyoctanoic acid, peroxyononanoic acid, peroxydecanoic acid, or mixture

thereof. The medium chain peroxycarboxylic acid can include or be a C8 to C10 peroxycarboxylic acid. The C8 to C10 peroxycarboxylic acid can include or be peroxyoctanoic acid, peroxydecanoic acid, or mixture thereof. In certain embodiments, the medium chain peroxycarboxylic acid includes or is
5 peroxyoctanoic acid, peroxydecanoic acid, or mixture thereof. In an embodiment, the medium chain peroxycarboxylic acid includes or is peroxyoctanoic acid.

In certain embodiments, the present composition includes about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid, about 0.3 to about 7 wt-% medium chain peroxycarboxylic acid, about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid,
10 about 0.5 to about 4 wt-% medium chain peroxycarboxylic acid, about 0.8 to about 3 wt-% medium chain peroxycarboxylic acid, about 1 to about 3 wt-% medium chain peroxycarboxylic acid, or about 1 to about 2 wt-% medium chain peroxycarboxylic acid. The composition can include any of these ranges or amounts not modified by about.

15 In an embodiment, the present compositions and methods include a medium chain carboxylic acid. The medium chain carboxylic acid can include or be a C6 to C12 carboxylic acid. The C6 to C12 carboxylic acid can include or be hexanoic acid, heptanoic acid, octanoic acid, nonanoic acid, decanoic acid, undecanoic acid, dodecanoic acid, or mixture thereof. The medium chain carboxylic acid can include or
20 be a C7 to C12 carboxylic acid. The C7 to C12 carboxylic acid can include or be heptanoic acid, octanoic acid, nonanoic acid, decanoic acid, undecanoic acid, dodecanoic acid, or mixture thereof. The medium chain peroxycarboxylic acid can include or be a C6 to C10 carboxylic acid. The C6 to C10 carboxylic acid can include or be hexanoic acid, heptanoic acid, octanoic acid, nonanoic acid, decanoic acid, or
25 mixture thereof. The medium chain carboxylic acid can include or be a C8 to C10 carboxylic acid. The C8 to C10 carboxylic acid can include or be octanoic acid, nonanoic acid, decanoic acid, or mixture thereof. In certain embodiments, the medium chain carboxylic acid includes or is octanoic acid, decanoic acid, or mixture thereof. In an embodiment, the medium chain carboxylic acid includes or is octanoic acid.

30 In certain embodiments, the present composition includes about 0.001 to about 8 wt-% medium chain carboxylic acid, about 1 to about 10 wt-% medium chain carboxylic acid, about 1 to about 8 wt-% medium chain carboxylic acid, about 1.5 to

about 6 wt-% medium chain carboxylic acid, about 2 to about 8 wt-% medium chain carboxylic acid, about 2 to about 6 wt-% medium chain carboxylic acid, about 2 to about 4 wt-% medium chain carboxylic acid, about 2.5 to about 5 wt-% medium chain carboxylic acid, about 3 to about 6 wt-% medium chain carboxylic acid, or about 3 to about 5 wt-% medium chain carboxylic acid. The composition can include any of these ranges or amounts not modified by about.

In an embodiment, the compositions and methods include a medium chain peroxy-carboxylic acid and the corresponding medium chain carboxylic acid.

In an embodiment, the present composition includes an amount of medium chain peroxy-carboxylic acid effective for killing one or more (e.g., at least one) of the food-borne pathogenic bacteria associated with a food product, such as *Salmonella typhimurium*, *Salmonella javiana*, *Campylobacter jejuni*, *Listeria monocytogenes*, and *Escherichia coli* O157:H7, yeast, mold, and the like. In an embodiment, the present composition includes an amount of medium chain peroxy-carboxylic acid effective for killing one or more (e.g., at least one) of the pathogenic bacteria associated with a health care surfaces and environments, such as *Salmonella typhimurium*, *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Escherichia coli*, mycobacteria, yeast, mold, and the like. The compositions and methods of the present invention have activity against a wide variety of microorganisms such as Gram positive (for example, *Listeria monocytogenes* or *Staphylococcus aureus*) and Gram negative (for example, *Escherichia coli* or *Pseudomonas aeruginosa*) bacteria, yeast, molds, bacterial spores, viruses, etc. The compositions and methods of the present invention, as described above, have activity against a wide variety of human pathogens. The present compositions and methods can kill a wide variety of microorganisms on a food processing surface, on the surface of a food product, in water used for washing or processing of food product, on a health care surface, or in a health care environment.

Embodiments of the present invention include medium chain carboxylic acid and medium chain peroxy-carboxylic acid, and certain embodiments specifically exclude short chain peroxy-carboxylic acid, short chain carboxylic acid, or mixture thereof. Nonetheless embodiments of the present compositions can include short chain peroxy-carboxylic acid, short chain carboxylic acid, or mixture thereof. It is not

intended that addition of short chain peroxycarboxylic acid, short chain carboxylic acid, or mixture thereof to a composition should necessarily take a composition outside the spirit and scope of the present invention.

5 Solubilizers

The present compositions can include a solubilizer. The present invention relates to solubilizers for medium chain carboxylic acids and medium chain peroxycarboxylic acids. In an embodiment, the solubilizer can increase or maintain the solubility in the composition of the medium chain peroxycarboxylic acid or the medium
10 chain carboxylic acid. The present compositions and methods can include any of a variety of suitable solubilizers. For example, the solubilizer can include a solvent, a surfactant, or a mixture thereof. In an embodiment, the surfactant can be employed as a solvent. In an embodiment, the surfactant can form a microemulsion. In an embodiment, the composition including the present solubilizer takes the form of a
15 viscoelastic gel or liquid. In an embodiment, the solubilizer is effective to dissolve octanoic acid at a concentration of 5 wt-% in water. In an embodiment, the solubilizer is effective to dissolve octanoic acid at a concentration of 4 wt-% in water. In an embodiment, the solubilizer is effective to dissolve octanoic acid at a concentration of 3 wt-% in water. In an embodiment, the solubilizer is effective to dissolve octanoic acid
20 at a concentration of 2 wt-% in water.

In certain embodiments, the present composition includes about 0.001 to about 80 wt-% solubilizer, about 0.001 to about 60 wt-% solubilizer, about 1 to about 80 wt-% solubilizer, about 1 to about 25 wt-% solubilizer, about 1 to about 20 wt-% solubilizer, about 2 to about 70 wt-% solubilizer, about 2 to about 60 wt-% solubilizer,
25 about 2 to about 20 wt-% solubilizer, about 3 to about 65 wt-% solubilizer, about 3 to about 15 wt-% solubilizer, about 4 to about 10 wt-% solubilizer, about 4 to about 20 wt-% solubilizer, about 5 to about 70 wt-% solubilizer, about 5 to about 60 wt-% solubilizer, about 5 to about 20 wt-% solubilizer, about 10 to about 70 wt-% solubilizer, about 10 to about 65 wt-% solubilizer, about 10 to about 20 wt-% solubilizer, about 20
30 to about 60 wt-% solubilizer, or about 40 to about 60 wt-% solubilizer. The composition can include any of these ranges or amounts not modified by about.

Solvent Solubilizers and Compositions Including Them

In an embodiment, the present compositions and methods can include as solubilizer one or more (e.g., at least one) solvents. Suitable solvents include any of a variety of solvents that solubilize but do not significantly degrade the medium chain
5 peroxycarboxylic acid. Suitable solvents include polyalkylene oxide, capped polyalkylene oxide, glycol ether, nonionic surfactant, mixtures thereof, or the like.

In an embodiment, the present composition includes medium chain peroxycarboxylic acid; medium chain carboxylic acid; carrier; and polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof. For example, the
10 present composition can include about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid; about 1 to about 10 wt-% medium chain carboxylic acid; about 1 to about 98 wt-% carrier; and about 1 to about 80 wt-% polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof. For example, the present composition can include about 0.5 to about 5 wt-% medium chain peroxycarboxylic
15 acid; about 1 to about 10 wt-% medium chain carboxylic acid; about 5 to about 35 wt-% carrier; and about 20 to about 65 wt-% polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof. For example, the present composition can include about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid; about 1 to about 10 wt-% medium chain carboxylic acid; about 10 to about 35 wt-% carrier; and
20 about 40 to about 60 wt-% polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof. In an embodiment, the present composition includes solvent solubilizer and less than or equal to 35 wt-% carrier (e.g., water). The composition can include any of these ranges or amounts not modified by about.

In an embodiment, the present composition includes C8 peroxycarboxylic acid; C8 carboxylic acid; water; and polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof. For example, the present composition can include about 0.5 to about 5 wt-% C8 peroxycarboxylic acid; about 1 to about 10 wt-% C8 carboxylic acid; about 1 to about 98 wt-% water; and about 1 to about 80 wt-% polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof.
25 For example, the present composition can include about 0.5 to about 5 wt-% C8 peroxycarboxylic acid; about 1 to about 10 wt-% C8 carboxylic acid; about 5 to about 35 wt-% water; and about 20 to about 65 wt-% polyalkylene oxide, capped
30

polyalkylene oxide, nonionic surfactant, or mixture thereof. For example, the present composition can include about 0.5 to about 5 wt-% C8 peroxycarboxylic acid; about 1 to about 10 wt-% C8 carboxylic acid; about 10 to about 35 wt-% water; and about 40 to about 60 wt-% polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 80 wt-% solvent as solubilizer, about 0.001 to about 60 wt-% solvent as solubilizer, about 1 to about 80 wt-% solvent as solubilizer, about 5 to about 70 wt-% solvent as solubilizer, about 10 to about 65 wt-% solvent as solubilizer, or about 20 to about 60 wt-% solvent as solubilizer. The composition can include any of these ranges or amounts not modified by about.

In an embodiment, when the present compositions and methods include a solvent as solubilizer, they need not include a significant amount, or even any, of a short chain peroxycarboxylic acid, a short chain carboxylic acid, or a mixture thereof. Examples of short chain carboxylic acids include formic acid, acetic acid, propionic acid, and butanoic acid. Short chain carboxylic acids and peroxycarboxylic acids include those with 4 or fewer carbon atoms. In an embodiment, the present compositions and methods including a solvent solubilizer need not include substantial amounts of short chain peroxycarboxylic acid. In an embodiment, the present compositions and methods including a solvent solubilizer can be free of added short chain peroxycarboxylic acid.

In an embodiment, the present compositions and methods including a solvent solubilizer can include medium chain peroxycarboxylic acid in greater proportion compared to the short chain peroxycarboxylic acid than found in conventional compositions. For example, the present compositions and methods can include solvent solubilizer and about 1 or more parts of medium chain peroxycarboxylic acid for each 8 parts of short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof. For example, the present compositions and methods can include solvent solubilizer and short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof at a level insufficient to cause odor offensive to a typical person.

Polyalkylene Oxide Solubilizers

Suitable polyalkylene oxides include polyethylene glycol, polypropylene glycol, polybutylene glycol, mixtures thereof, or the like. Suitable capped polyalkylene oxides include mono-alkyl and di-alkyl ethers of the respective polyalkylene oxides, such as

5 mono- and di-methyl ethers of polyalkylene glycol, mono- and di-ethyl ethers of polyalkylene glycol, mono- and di-propyl ethers of polyalkylene glycol, mono- and di-butyl ethers of polyalkylene glycol, mixtures thereof, or the like. Suitable capped polyalkylene oxides include methyl polyethylene glycol (e.g., the monomethyl ether of polyethylene glycol), dimethyl polyethylene glycol (e.g., the dimethyl ether of

10 polyethylene glycol), mixtures thereof, or the like.

Glycol Ether Solubilizers

Suitable solvent solubilizers include glycol ethers. Suitable glycol ethers include diethylene glycol n-butyl ether, diethylene glycol n-propyl ether, diethylene

15 glycol ethyl ether, diethylene glycol methyl ether, diethylene glycol t-butyl ether, dipropylene glycol n-butyl ether, dipropylene glycol methyl ether, dipropylene glycol ethyl ether, dipropylene glycol propyl ether, dipropylene glycol tert-butyl ether, ethylene glycol butyl ether, ethylene glycol propyl ether, ethylene glycol ethyl ether, ethylene glycol methyl ether, ethylene glycol methyl ether acetate, propylene glycol n-

20 butyl ether, propylene glycol ethyl ether, propylene glycol methyl ether, propylene glycol n-propyl ether, tripropylene glycol methyl ether and tripropylene glycol n-butyl ether, ethylene glycol phenyl ether (commercially available as DOWANOL EPH™ from Dow Chemical Co.), propylene glycol phenyl ether (commercially available as DOWANOL PPH™ from Dow Chemical Co.), and the like, or mixtures thereof.

25 Additional suitable commercially available glycol ethers (all of which are available from Union Carbide Corp.) include Butoxyethyl PROPASOL™, Butyl CARBITOL™ acetate, Butyl CARBITOL™, Butyl CELLOSOLVE™ acetate, Butyl CELLOSOLVE™, Butyl DIPROPASOL™, Butyl PROPASOL™, CARBITOL™ PM-600, CARBITOL™ Low Gravity, CELLOSOLVE™ acetate, CELLOSOLVE™, Ester EEP™, FILMER

30 IBT™, Hexyl CARBITOL™, Hexyl CELLOSOLVE™, Methyl CARBITOL™, Methyl CELLOSOLVE™ acetate, Methyl CELLOSOLVE™, Methyl DIPROPASOL™, Methyl

PROPASOL[™] acetate, Methyl PROPASOL[™], Propyl CARBITOL[™], Propyl CELLOSOLVE[™], Propyl DIPROPASOL[™] and Propyl PROPASOL[™].

Nonionic Surfactants


5 Suitable nonionic surfactants for use as solvents include alkoxyated surfactants. Suitable alkoxyated surfactants include EO/PO copolymers, capped EO/PO copolymers, alcohol alkoxyates, capped alcohol alkoxyates, mixtures thereof, or the like. Suitable alkoxyated surfactants for use as solvents include EO/PO block copolymers, such as the Pluronic and reverse Pluronic surfactants; alcohol alkoxyates, 10 such as Dehypon LS-54 (R-(EO)₅(PO)₄) and Dehypon LS-36 (R-(EO)₃(PO)₆); and capped alcohol alkoxyates, such as Plurafac LF221 and Tegoten EC11; mixtures thereof, or the like. When employed as a solvent a surfactant, such as a nonionic surfactant, can be at concentrations higher than those conventionally employed as surfactant.

15

Semi-Polar Nonionic Surfactants

 The semi-polar type of nonionic surface active agents are another class of nonionic surfactant useful in compositions of the present invention. Semi-polar nonionic surfactants include the amine oxides, phosphine oxides, sulfoxides and their 20 alkoxyated derivatives.

 Amine oxides are tertiary amine oxides corresponding to the general formula:



wherein the arrow is a conventional representation of a semi-polar bond; and, R¹, R², and R³ may be aliphatic, aromatic, heterocyclic, alicyclic, or combinations thereof.

25 Generally, for amine oxides of detergent interest, R¹ is an alkyl radical of from about 8 to about 24 carbon atoms; R² and R³ are alkyl or hydroxyalkyl of 1-3 carbon atoms or a mixture thereof; R² and R³ can be attached to each other, e.g. through an oxygen or nitrogen atom, to form a ring structure; R⁴ is an alkylene or a hydroxyalkylene group containing 2 to 3 carbon atoms; and n ranges from 0 to about 20.

Useful water soluble amine oxide surfactants are selected from the octyl, decyl, dodecyl, isododecyl, coconut, or tallow alkyl di-(lower alkyl) amine oxides, specific examples of which are octyldimethylamine oxide, nonyldimethylamine oxide, decyldimethylamine oxide, undecyldimethylamine oxide, dodecyldimethylamine oxide, 5 iso-dodecyldimethyl amine oxide, tridecyldimethylamine oxide, tetradecyldimethylamine oxide, pentadecyldimethylamine oxide, hexadecyldimethylamine oxide, heptadecyldimethylamine oxide, octadecyldimethylamine oxide, dodecyldipropylamine oxide, tetradecyldipropylamine oxide, hexadecyldipropylamine oxide, tetradecyldibutylamine oxide, 10 octadecyldibutylamine oxide, bis(2-hydroxyethyl)dodecylamine oxide, bis(2-hydroxyethyl)-3-dodecoxy-1-hydroxypropylamine oxide, dimethyl-(2-hydroxydodecyl)amine oxide, 3,6,9-trioctadecyldimethylamine oxide and 3-dodecoxy-2-hydroxypropyldi-(2-hydroxyethyl)amine oxide.

15 Surfactant Solubilizers and Compositions Including Them

In an embodiment, the present compositions and methods can include as solubilizer one or more (e.g., at least one) surfactants, e.g., a microemulsion forming surfactant. Suitable surfactants include anionic surfactant, cationic surfactant, amphoteric surfactant, zwitterionic surfactant, mixtures thereof, or the like. Suitable 20 microemulsion forming surfactants include anionic surfactant, cationic surfactant, amphoteric surfactant, zwitterionic surfactant, mixtures thereof, or the like. Suitable microemulsion forming surfactants include anionic surfactant. A microemulsion forming surfactant can form a microemulsion in a composition including a medium chain peroxycarboxylic acid, a medium chain carboxylic acid, or a mixture thereof. In 25 an embodiment, the present composition includes a microemulsion.

In an embodiment, the present composition can be determined to be a microemulsion by testing the composition for being a shear thinning viscoelastic gel or liquid that has a blue tyndall appearance. Although not limiting to the present invention, blue tyndall appearance is believed to indicate a heterogeneous system of a 30 small, suspended dispersion (e.g., a microemulsion), which is effective in scattering blue light.

In an embodiment, the present composition can be determined to be a microemulsion by testing the ability to form a physically stable composition at different concentrations of surfactant solubilizer. A microemulsion can yield a curve with a maximum of physical stability at a concentration with unstable compositions at higher and lower concentrations. Typically, mixtures of solvents and surfactants (e.g., acetic acid and surfactant) do not form microemulsions.

In an embodiment, the composition including surfactant solubilizer takes the form of a viscoelastic gel or liquid. Increasing the concentration of the medium chain carboxylic acid, medium chain peroxycarboxylic acid, or mixture thereof can increase the degree to which the composition is a viscoelastic gel or liquid. Increasing the concentration of the surfactant solubilizer can increase the degree to which the composition is a viscoelastic gel or liquid. In an embodiment, the gel can be sufficiently viscoelastic to hold its molded shape. Alkyl benzene sulfonate surfactant (e.g., LAS) can be employed to form a viscoelastic gel or liquid that can hold its molded shape. In an embodiment, the alkyl benzene sulfonate surfactant containing viscoelastic gel can hold its shape even at 60 °C.

Although not limiting to the present invention, the present compositions may include medium chain peroxycarboxylic acid sequestered in the surfactant of the microemulsion. This can stabilize the peroxycarboxylic acid by keeping it away from impurities or reducing agents in the bulk water. This can increase the production of peroxycarboxylic acid by pulling it out of solution. Although not limiting to the present invention, it is believed that one explanation for the viscoelastic properties of gels of the present compositions is that they are due to repulsive forces between the dispersions/droplets that are stabilized by the microemulsion-forming surfactant. Surfactants that are charged may increase the electrostatic repulsion. Suitable charged surfactants include anionic surfactants.

In an embodiment, the present composition includes anionic surfactant and another surfactant or surfactants. For example, the present compositions can include anionic surfactant and nonionic surfactant or semi-polar nonionic surfactant.

In an embodiment, the present composition includes medium chain peroxycarboxylic acid; medium chain carboxylic acid; carrier; and one or more (e.g., at least one) surfactants, e.g., microemulsion forming surfactants. For example, the

present composition can include about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid; about 1 to about 10 wt-% medium chain carboxylic acid; about 5 to about 97 wt-% carrier; and about 1 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant. For example, the present composition can include about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid; about 1 to about 10 wt-% medium chain carboxylic acid; about 15 to about 80 wt-% carrier; and about 1 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant. For example, the present composition can include about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid; about 1 to about 10 wt-% medium chain carboxylic acid; about 30 to about 70 wt-% carrier; and about 2 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant. In an embodiment, the present composition includes surfactant or microemulsion former solubilizer and greater than or equal to 35 wt-% carrier (e.g., water). The composition can include any of these ranges or amounts not modified by about.

15 In an embodiment, the present composition includes C8 peroxycarboxylic acid; C8 carboxylic acid; water; and one or more (e.g., at least one) surfactants, e.g., microemulsion forming surfactants. For example, the present composition can include about 0.5 to about 5 wt-% C8 peroxycarboxylic acid; about 1 to about 10 wt-% C8 carboxylic acid; about 5 to about 97 wt-% water; and about 1 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant. For example, the present composition can include about 0.5 to about 5 wt-% C8 peroxycarboxylic acid; about 1 to about 10 wt-% C8 carboxylic acid; about 15 to about 80 wt-% water; and about 1 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant. For example, the present composition can include about 0.5 to about 5 wt-% C8 peroxycarboxylic acid; about 1 to about 10 wt-% C8 carboxylic acid; about 30 to about 70 wt-% water; and about 2 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant. The composition can include any of these ranges or amounts not modified by about.

30 In certain embodiments, the present composition includes about 0.001 to about 60 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, about 1 to about 25 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, about 1 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, about 2 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant, as

solubilizer, about 3 to about 15 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, about 4 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, about 4 to about 10 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, about 5 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer, or about 10 to about 20 wt-% surfactant, e.g., microemulsion forming surfactant, as solubilizer. The composition can include any of these ranges or amounts not modified by about.

Anionic Surfactants

10 The present composition can include an anionic surfactant as solubilizer. Suitable anionic surfactants include organic sulfonate surfactant, organic sulfate surfactant, phosphate ester surfactant, carboxylate surfactant, mixtures thereof, or the like. In an embodiment, the anionic surfactant includes alkyl sulfonate, alkylaryl sulfonate, alkylated diphenyl oxide disulfonate, alkylated naphthalene sulfonate, 15 alcohol alkoxylate carboxylate, sarcosinate, taurate, acyl amino acid, alkanolic ester, phosphate ester, sulfuric acid ester, salt or acid form thereof, or mixture thereof. The particular salts will be suitably selected depending upon the particular formulation and the needs therein.

 Suitable anionic surfactants include sulfonic acids (and salts), such as 20 isethionates (e.g. acyl isethionates), alkylaryl sulfonic acids and salts thereof, alkyl sulfonates, secondary alkane sulfonates, and the like.

 Examples of suitable synthetic, water soluble anionic detergent compounds include the ammonium and substituted ammonium (such as mono-, di- and triethanolamine) and alkali metal (such as sodium, lithium and potassium) salts of the 25 alkyl mononuclear aromatic sulfonates such as the alkyl benzene sulfonates containing from about 5 to about 18 carbon atoms in the alkyl group in a straight or branched chain, e.g., the salts of alkyl benzene sulfonates or of alkyl toluene, xylene, cumene and phenol sulfonates; alkyl naphthalene sulfonate, diamyl naphthalene sulfonate, and dinonyl naphthalene sulfonate and alkoxylated derivatives or their free acids. Suitable 30 sulfonates include olefin sulfonates, such as long chain alkene sulfonates, long chain hydroxyalkane sulfonates or mixtures of alkenesulfonates and hydroxyalkane-sulfonates. Suitable sulfonates include secondary alkane sulfonates.

In certain embodiments, the present compositions including an anionic surfactant, such as a normal C8 sulfonate, can be non-foam or low foam compositions. Such compositions can be advantageous for applications such as clean in place, machine warewashing, destaining, and sanitizing, laundry washing, destaining, and
5 sanitizing, etc.


For applications in which foaming is desirable, a foaming agent can be added as part of the present composition or separately. In a two-step offering, a foaming agent can be combined with a dilution of the non-foam or low foam composition to form a foaming use solution. In a one-step offering, the foaming agent can be incorporated
10 into the concentrated composition. One suitable foaming agent is LAS acid. LAS acid can form a microemulsion in the present compositions. LAS acid can form a viscoelastic gel or liquid in the present compositions. Additional suitable foaming agents include secondary alkane sulfonate, alkylated diphenyl oxide disulfonate (e.g., C12 alkyl diphenyl oxide disulfonate), alkyl ether sulfate (e.g., with n=1-3) (e.g.,
15 sodium laureth sulfate (with n=1, 2, or 3)), sodium lauryl sulfate, or the like.


In an embodiment, such foaming agents provide a foaming composition with one or more desirable foaming characteristics. Desirable foaming characteristics include, for example, foam being visible for about 5 min after forming the foam; foam with continuous and good drainage (e.g., when applied to a vertical surface); foam that
20 dries to a clear appearance, e.g., that leaves no visible residue on a stainless steel surface; and/or foam that can be applied with a moderate or low odor compared to a conventional foam containing peroxyacetic acid.


Anionic sulfate surfactants suitable for use in the present compositions include alkyl ether sulfates, alkyl sulfates, the linear and branched primary and secondary alkyl
25 sulfates, alkyl ethoxysulfates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, the C₅-C₁₇ acyl-N-(C₁-C₄ alkyl) and -N-(C₁-C₂ hydroxyalkyl) glucamine sulfates, and sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside, and the like. Also included are the alkyl sulfates, alkyl
poly(ethyleneoxy) ether sulfates and aromatic poly(ethyleneoxy) sulfates such as the
30 sulfates or condensation products of ethylene oxide and nonyl phenol (usually having 1 to 6 oxyethylene groups per molecule).

Anionic carboxylate surfactants suitable for use in the present compositions include carboxylic acids (and salts), such as alkanoic acids (and alkanoates), ester carboxylic acids (e.g. alkyl succinates), ether carboxylic acids, and the like. Such carboxylates include alkyl ethoxy carboxylates, alkyl aryl ethoxy carboxylates, alkyl polyethoxy polycarboxylate surfactants and soaps (e.g. alkyl carboxyls). Secondary carboxylates useful in the present compositions include those which contain a carboxyl unit connected to a secondary carbon. The secondary carbon can be in a ring structure, e.g. as in p-octyl benzoic acid, or as in alkyl-substituted cyclohexyl carboxylates. The secondary carboxylate surfactants typically contain no ether linkages, no ester linkages and no hydroxyl groups. Further, they typically lack nitrogen atoms in the head-group (amphiphilic portion). Suitable secondary soap surfactants typically contain 11-13 total carbon atoms, although more carbons atoms (e.g., up to 16) can be present. Suitable carboxylates also include acylamino acids (and salts), such as acylgluamates, acyl peptides, sarcosinates (e.g. N-acyl sarcosinates), taurates (e.g. N-acyl taurates and fatty acid amides of methyl tauride), and the like.

Suitable anionic surfactants include alkyl or alkylaryl ethoxy carboxylates of Formula 3:



in which R is a C₈ to C₂₂ alkyl group or , in which R¹ is a C₄-C₁₆ alkyl group; n is an integer of 1-20; m is an integer of 1-3; and X is a counter ion, such as hydrogen, sodium, potassium, lithium, ammonium, or an amine salt such as monoethanolamine, diethanolamine or triethanolamine. In an embodiment, in Formula 3, n is an integer of 4 to 10 and m is 1. In an embodiment, in Formula 3, R is a C₈-C₁₆ alkyl group. In an embodiment, in Formula 3, R is a C₁₂-C₁₄ alkyl group, n is 4, and m is 1.

In an embodiment, in Formula 3, R is  and R¹ is a C₆-C₁₂ alkyl group. In an embodiment, in Formula 3, R¹ is a C₉ alkyl group, n is 10 and m is 1.

Such alkyl and alkylaryl ethoxy carboxylates are commercially available. These ethoxy carboxylates are typically available as the acid forms, which can be readily converted to the anionic or salt form. Commercially available carboxylates include, Neodox 23-4, a C₁₂₋₁₃ alkyl polyethoxy (4) carboxylic acid (Shell Chemical), and Emcol CNP-110, a C₉ alkylaryl polyethoxy (10) carboxylic acid (Witco Chemical). Carboxylates are also available from Clariant, e.g. the product Sandopan[®] DTC, a C₁₃ alkyl polyethoxy (7) carboxylic acid.

Amphoteric Surfactants

Amphoteric, or ampholytic, surfactants contain both a basic and an acidic hydrophilic group and an organic hydrophobic group. These ionic entities may be any of anionic or cationic groups described herein for other types of surfactants. A basic nitrogen and an acidic carboxylate group are the typical functional groups employed as the basic and acidic hydrophilic groups. In a few surfactants, sulfonate, sulfate, phosphonate or phosphate provide the negative charge.

Amphoteric surfactants can be broadly described as derivatives of aliphatic secondary and tertiary amines, in which the aliphatic radical may be straight chain or branched and wherein one of the aliphatic substituents contains from about 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfo, sulfato, phosphato, or phosphono. Amphoteric surfactants are subdivided into two major classes known to those of skill in the art and described in "Surfactant Encyclopedia" Cosmetics & Toiletries, Vol. 104 (2) 69-71 (1989). The first class includes acyl/dialkyl ethylenediamine derivatives (e.g. 2-alkyl hydroxyethyl imidazoline derivatives) and their salts. The second class includes N-alkylamino acids and their salts. Some amphoteric surfactants can be envisioned as fitting into both classes.

Amphoteric surfactants can be synthesized by methods known to those of skill in the art. For example, 2-alkyl hydroxyethyl imidazoline is synthesized by condensation and ring closure of a long chain carboxylic acid (or a derivative) with dialkyl ethylenediamine. Commercial amphoteric surfactants are derivatized by subsequent hydrolysis and ring-opening of the imidazoline ring by alkylation -- for example with chloroacetic acid or ethyl acetate. During alkylation, one or two

Long chain imidazoline derivatives having application in the present invention generally have the general formula:



Commercially prominent imidazoline-derived amphoteric surfactants that can be employed in the present compositions include for example: Cocoamphopropionate, Cocoamphocarboxy-propionate, Cocoamphoglycinate, Cocoamphocarboxy-glycinate, Cocoamphopropyl-sulfonate, and Cocoamphocarboxy-propionic acid.

15

Long chain N-alkylamino acids are readily prepared by reaction RNH_2 , in which $\text{R}=\text{C}_8\text{-C}_{18}$ straight or branched chain alkyl, fatty amines with halogenated carboxylic acids. Alkylation of the primary amino groups of an amino acid leads to secondary and tertiary amines. Alkyl substituents may have additional amino groups that provide more than one reactive nitrogen center. Most commercial N-alkylamine acids are alkyl derivatives of beta-alanine or beta-N(2-carboxyethyl) alanine.

25

containing from about 8 to about 18 carbon atoms, and M is a cation to neutralize the charge of the anion.

Suitable amphoteric surfactants include those derived from coconut products such as coconut oil or coconut fatty acid. Additional suitable coconut derived surfactants include as part of their structure an ethylenediamine moiety, an alkanolamide moiety, an amino acid moiety, e.g., glycine, or a combination thereof; and an aliphatic substituent of from about 8 to 18 (e.g., 12) carbon atoms. Such a surfactant can also be considered an alkyl amphodicarboxylic acid. These amphoteric surfactants can include chemical structures represented as: $C_{12}\text{-alkyl-C(O)-NH-CH}_2\text{-CH}_2\text{-N}^+(\text{CH}_2\text{-CH}_2\text{-CO}_2\text{Na})_2\text{-CH}_2\text{-CH}_2\text{-OH}$ or $C_{12}\text{-alkyl-C(O)-N(H)-CH}_2\text{-CH}_2\text{-N}^+(\text{CH}_2\text{-CO}_2\text{Na})_2\text{-CH}_2\text{-CH}_2\text{-OH}$. Disodium cocoampho dipropionate is one suitable amphoteric surfactant and is commercially available under the tradename Miranol™ FBS from Rhodia Inc., Cranbury, N.J. Another suitable coconut derived amphoteric surfactant with the chemical name disodium cocoampho diacetate is sold under the tradename Mirataine™ JCHA, also from Rhodia Inc., Cranbury, N.J.


A typical listing of amphoteric classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

Zwitterionic Surfactants

Zwitterionic surfactants can be thought of as a subset of the amphoteric surfactants and can include an anionic charge. Zwitterionic surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. Typically, a zwitterionic surfactant includes a positive charged quaternary ammonium or, in some cases, a sulfonium or phosphonium ion; a negative charged carboxyl group; and an alkyl group. Zwitterionics generally contain cationic and anionic groups which ionize to a nearly equal degree in the isoelectric region of the molecule and which can develop strong "inner-salt" attraction between positive-negative charge centers. Examples of such zwitterionic synthetic surfactants include derivatives of aliphatic quaternary

ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight chain or branched, and wherein one of the aliphatic substituents contains from 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Betaine and sultaine surfactants
 5 are exemplary zwitterionic surfactants for use herein.


A general formula for these compounds is:



wherein R^1 contains an alkyl, alkenyl, or hydroxyalkyl radical of from 8 to 18 carbon atoms having from 0 to 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety; Y
 10 is selected from the group consisting of nitrogen, phosphorus, and sulfur atoms; R^2 is an alkyl or monohydroxy alkyl group containing 1 to 3 carbon atoms; x is 1 when Y is a sulfur atom and 2 when Y is a nitrogen or phosphorus atom, R^3 is an alkylene or hydroxy alkylene or hydroxy alkylene of from 1 to 4 carbon atoms and Z is a radical selected from the group consisting of carboxylate, sulfonate, sulfate, phosphonate, and
 15 phosphate groups.

Examples of zwitterionic surfactants having the structures listed above include: 4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate; 5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxypentane-1-sulfate; 3-[P,P-diethyl-P-3,6,9-trioxatetracosanephosphonio]-2-hydroxypropane-1-phosphate; 3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropyl-ammonio]-propane-1-phosphonate; 3-(N,N-dimethyl-N-hexadecylammonio)-propane-1-sulfonate; 3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxy-propane-1-sulfonate; 4-[N,N-di(2(2-hydroxyethyl)-N(2-hydroxydodecyl)ammonio)-butane-1-carboxylate; 3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-propane-1-phosphate; 3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-phosphonate; and S[N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-hydroxy-pentane-1-sulfate. The alkyl groups contained in said detergent surfactants can be straight or branched and saturated or unsaturated.

The zwitterionic surfactant suitable for use in the present compositions includes a betaine of the general structure:



These surfactant betaines typically do not exhibit strong cationic or anionic characters at pH extremes nor do they show reduced water solubility in their isoelectric range.

Unlike "external" quaternary ammonium salts, betaines are compatible with anionics.

- 5 Examples of suitable betaines include coconut acylamidopropyldimethyl betaine; hexadecyl dimethyl betaine; C₁₂₋₁₄ acylamidopropylbetaine; C₈₋₁₄ acylamidohexyldiethyl betaine; 4-C₁₄₋₁₆ acylmethyldiethylammonio-1-carboxybutane; C₁₆₋₁₈ acylamidodimethylbetaine; C₁₂₋₁₆ acylamidopentanedimethylbetaine; and C₁₂₋₁₆ acylmethyldiethylbetaine.

- 10 Sultaines useful in the present invention include those compounds having the formula (R(R¹)₂ N⁺ R²SO³⁻), in which R is a C₆-C₁₈ hydrocarbyl group, each R¹ is typically independently C₁-C₃ alkyl, e.g. methyl, and R² is a C₁-C₆ hydrocarbyl group, e.g. a C₁-C₃ alkylene or hydroxyalkylene group.

- 15 A typical listing of zwitterionic classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

In an embodiment, the composition of the present invention includes a betaine. For example, the composition can include cocoamidopropyl betaine.

20

Embodiments of Compositions

- Some examples of representative constituent concentrations for embodiments of the present compositions can be found in Tables A-C, in which the values are given in wt-% of the ingredients in reference to the total composition weight. In certain
- 25 embodiments, the proportions and amounts in Tables A-C can be modified by "about".

Table A

Ingredient	wt-%	wt-%	wt-%	wt-%
medium chain peroxycarboxylic acid	0.3-7	0.5-5	0.5-4	1-3
medium chain carboxylic acid	1-10	2-8	2-6	2.5-5
solubilizer	1-80	2-70	3-65	5-60
carrier	0-98	5-90	10-80	20-70

Table B

Ingredient	wt-%	wt-%	wt-%	wt-%
medium chain peroxycarboxylic acid	0.3-7	0.5-5	0.5-4	1-3
medium chain carboxylic acid	1-10	2-8	3-6	3-5
solubilizer	1-80	5-70	10-65	20-60
carrier	0-98	0.2-60	5-20	20-40

5

Table C

Ingredient	wt-%	wt-%	wt-%	wt-%
medium chain peroxycarboxylic acid	0.3-7	0.5-5	0.5-4	1-2
medium chain carboxylic acid	1-10	1-8	1.5-6	2-4
solubilizer	1-25	2-20	3-15	4-10
carrier	5-97	10-90	15-70	30-75

Some examples of representative constituent concentrations for additional embodiments of the present compositions can be found in Tables D-F, in which the values are given in wt-% of the ingredients in reference to the total composition weight.

10 In certain embodiments, the proportions and amounts in Tables D-F can be modified by “about”.

Table D

Ingredient	wt-%	wt-%	wt-%	wt-%
medium chain peroxycarboxylic acid	0.3-7	0.5-5	0.5-4	1-3
medium chain carboxylic acid	1-10	2-8	2-6	2.5-5
solubilizer	1-80	2-70	3-65	5-60
carrier	0-98	5-90	10-80	20-70
oxidizing agent	2-30	2-25	4-20	6-10
acidulant	1-50	2-40	3-40	5-40
stabilizing agent	1-50	1-10	1-5	1-3

Table E

Ingredient	wt-%	wt-%	wt-%	wt-%
medium chain peroxycarboxylic acid	0.3-7	0.5-5	0.5-4	1-3
medium chain carboxylic acid	1-10	2-8	3-6	3-5
solubilizer	1-80	5-70	10-65	20-60
carrier	0-98	0.2-60	5-20	20-40
oxidizing agent	2-30	2-25	4-20	6-10
acidulant	1-50	2-40	3-40	5-40
stabilizing agent	1-50	1-10	1-5	1-3

Table F

Ingredient	wt-%	wt-%	wt-%	wt-%
medium chain peroxycarboxylic acid	0.3-7	0.5-5	0.5-4	1-2
medium chain carboxylic acid	1-10	1-8	1.5-6	2-4
solubilizer	1-25	2-20	3-15	4-10
carrier	5-97	10-90	15-70	30-75
oxidizing agent	2-30	2-25	4-20	6-10
acidulant	1-50	2-40	3-35	5-30
stabilizing agent	1-50	1-15	1-5	1-3

In an embodiment, the compositions of the present invention include only ingredients that can be employed in food products or in food wash, handling, or processing, for example, according to government (e.g. FDA or USDA) rules and regulations, 21 CFR §170-178. In an embodiment, the compositions of the present invention can include only ingredients at the concentrations approved for incidental food contact by the USEPA, 40 CFR §180.940.

The present compositions can take the form of a liquid, solid, gel, paste, unit dose, gel pack, or the like. The present compositions can be supplied in any of a variety of containers or media, such as in a 2 compartment dispenser or as a pre-moistened wipe, towelette, or sponge.

Carrier

The composition of the invention can also include a carrier. The carrier provides a medium which dissolves, suspends, or carries the other components of the composition. For example, the carrier can provide a medium for solubilization, suspension, or production of peroxycarboxylic acid and for forming an equilibrium mixture. The carrier can also function to deliver and wet the antimicrobial composition of the invention on an object. To this end, the carrier can contain any component or components that can facilitate these functions.

In certain embodiments, the carrier includes primarily water which can promote solubility and work as a medium for reaction and equilibrium. The carrier can include or be primarily an organic solvent, such as simple alkyl alcohols, e.g., ethanol, isopropanol, n-propanol, and the like. Polyols are also useful carriers, including
 5 glycerol, sorbitol, and the like.

Suitable carriers include glycol ethers. Suitable glycol ethers include diethylene glycol n-butyl ether, diethylene glycol n-propyl ether, diethylene glycol ethyl ether, diethylene glycol methyl ether, diethylene glycol t-butyl ether, dipropylene glycol n-butyl ether, dipropylene glycol methyl ether, dipropylene glycol ethyl ether,
 10 dipropylene glycol propyl ether, dipropylene glycol tert-butyl ether, ethylene glycol butyl ether, ethylene glycol propyl ether, ethylene glycol ethyl ether, ethylene glycol methyl ether, ethylene glycol methyl ether acetate, propylene glycol n-butyl ether, propylene glycol ethyl ether, propylene glycol methyl ether, propylene glycol n-propyl ether, tripropylene glycol methyl ether and tripropylene glycol n-butyl ether, ethylene
 15 glycol phenyl ether (commercially available as DOWANOL EPH™ from Dow Chemical Co.), propylene glycol phenyl ether (commercially available as DOWANOL PPH™ from Dow Chemical Co.), and the like, or mixtures thereof. Additional suitable commercially available glycol ethers (all of which are available from Union Carbide Corp.) include Butoxyethyl PROPASOL™, Butyl CARBITOL™ acetate, Butyl
 20 CARBITOL™, Butyl CELLOSOLVE™ acetate, Butyl CELLOSOLVE™, Butyl DIPROPASOL™, Butyl PROPASOL™, CARBITOL™ PM-600, CARBITOL™ Low Gravity, CELLOSOLVE™ acetate, CELLOSOLVE™, Ester EEP™, FILMER IBT™, Hexyl CARBITOL™, Hexyl CELLOSOLVE™, Methyl CARBITOL™, Methyl CELLOSOLVE™ acetate, Methyl CELLOSOLVE™, Methyl DIPROPASOL™, Methyl
 25 PROPASOL™ acetate, Methyl PROPASOL™, Propyl CARBITOL™, Propyl CELLOSOLVE™, Propyl DIPROPASOL™ and Propyl PROPASOL™.

In certain embodiments, the carrier makes up a large portion of the composition of the invention and may be the balance of the composition apart from the active antimicrobial components, solubilizer, oxidizing agent, adjuvants, and the like. Here
 30 again, the carrier concentration and type will depend upon the nature of the composition as a whole, the environmental storage, and method of application including concentration of the medium chain peroxycarboxylic acid, among other

factors. Notably the carrier should be chosen and used at a concentration which does not inhibit the antimicrobial efficacy of the medium chain peroxycarboxylic acid in the composition of the invention.

In certain embodiments, the present composition includes about 0 to about 98 wt-% carrier, about 0.001 to about 99.99 wt-% carrier, about 0.2 to about 60 wt-% carrier, about 1 to about 98 wt-% carrier, about 5 to about 99.99 wt-% carrier, about 5 to about 97 wt-% carrier, about 5 to about 90 wt-% carrier, about 5 to about 70 wt-% carrier, about 5 to about 20 wt-% carrier, about 10 to about 90 wt-% carrier, about 10 to about 80 wt-% carrier, about 10 to about 50 wt-% carrier, about 10 to about 20 wt-% carrier, about 15 to about 70 wt-% carrier, about 15 to about 80 wt-% carrier, about 20 to about 70 wt-% carrier, about 20 to about 50 wt-% carrier, about 20 to about 40 wt-% carrier, about 20 to about 30 wt-% carrier, about 30 to about 75 wt-% carrier, about 30 to about 70 wt-% carrier, about 40 to about 99.99 wt-% carrier, about 40 to about 90 wt-% carrier, or about 60 to about 70 wt-% carrier. The composition can include any of these ranges or amounts not modified by about.

Oxidizing Agent

The present compositions and methods can include any of a variety of oxidizing agents. The oxidizing agent can be used for maintaining or generating peroxycarboxylic acids.

Examples of inorganic oxidizing agents include the following types of compounds or sources of these compounds, or alkali metal salts including these types of compounds, or forming an adduct therewith:

hydrogen peroxide;

group 1 (IA) oxidizing agents, for example lithium peroxide, sodium peroxide, and the like;

group 2 (IIA) oxidizing agents, for example magnesium peroxide, calcium peroxide, strontium peroxide, barium peroxide, and the like;

group 12 (IIB) oxidizing agents, for example zinc peroxide, and the like;

group 13 (IIIA) oxidizing agents, for example boron compounds, such as perborates, for example sodium perborate hexahydrate of the formula

$\text{Na}_2[\text{B}_2(\text{O}_2)_2(\text{OH})_4] \cdot 6\text{H}_2\text{O}$ (also called sodium perborate tetrahydrate and formerly

written as $\text{NaBO}_3 \cdot 4\text{H}_2\text{O}$); sodium peroxyborate tetrahydrate of the formula $\text{Na}_2\text{B}_2(\text{O}_2)_2[(\text{OH})_4] \cdot 4\text{H}_2\text{O}$ (also called sodium perborate trihydrate, and formerly written as $\text{NaBO}_3 \cdot 3\text{H}_2\text{O}$); sodium peroxyborate of the formula $\text{Na}_2[\text{B}_2(\text{O}_2)_2(\text{OH})_4]$ (also called sodium perborate monohydrate and formerly written as $\text{NaBO}_3 \cdot \text{H}_2\text{O}$); and the
5 like; in an embodiment, perborate;

group 14 (IVA) oxidizing agents, for example persilicates and peroxycarbonates, which are also called percarbonates, such as persilicates or peroxycarbonates of alkali metals; and the like; in an embodiment, percarbonate; in an embodiment, persilicate;

10 group 15 (VA) oxidizing agents, for example peroxynitrous acid and its salts; peroxyphosphoric acids and their salts, for example, perphosphates; and the like; in an embodiment, perphosphate;

group 16 (VIA) oxidizing agents, for example peroxysulfuric acids and their salts, such as peroxymonosulfuric and peroxydisulfuric acids, and their salts, such as
15 persulfates, for example, sodium persulfate; and the like; in an embodiment, persulfate;

group VIIa oxidizing agents such as sodium periodate, potassium perchlorate and the like.

Other active inorganic oxygen compounds can include transition metal peroxides; and other such peroxygen compounds, and mixtures thereof.

20 In an embodiment, the compositions and methods of the present invention employ one or more (e.g., at least one) of the inorganic oxidizing agents listed above. Suitable inorganic oxidizing agents include ozone, hydrogen peroxide, hydrogen peroxide adduct, group IIIA oxidizing agent, group VIA oxidizing agent, group VA oxidizing agent, group VIIA oxidizing agent, or mixtures thereof. Suitable examples of
25 such inorganic oxidizing agents include percarbonate, perborate, persulfate, perphosphate, persilicate, or mixtures thereof.

Hydrogen peroxide presents one suitable example of an inorganic oxidizing agent. Hydrogen peroxide can be provided as a mixture of hydrogen peroxide and water, e.g., as liquid hydrogen peroxide in an aqueous solution. Hydrogen peroxide is
30 commercially available at concentrations of 35%, 70%, and 90% in water. For safety, the 35% is commonly used. The present compositions can include, for example, about 2 to about 30 wt-% or about 5 to about 20 wt-% hydrogen peroxide.

In an embodiment, the inorganic oxidizing agent includes hydrogen peroxide adduct. For example, the inorganic oxidizing agent can include hydrogen peroxide, hydrogen peroxide adduct, or mixtures thereof. Any of a variety of hydrogen peroxide adducts are suitable for use in the present compositions and methods. For example, 5 suitable hydrogen peroxide adducts include percarbonate salt, urea peroxide, peracetyl borate, an adduct of H_2O_2 and polyvinyl pyrrolidone, sodium percarbonate, potassium percarbonate, mixtures thereof, or the like. Suitable hydrogen peroxide adducts include percarbonate salt, urea peroxide, peracetyl borate, an adduct of H_2O_2 and polyvinyl pyrrolidone, or mixtures thereof. Suitable hydrogen peroxide adducts include sodium 10 percarbonate, potassium percarbonate, or mixtures thereof, for example sodium percarbonate.

In an embodiment, the present compositions and methods can include hydrogen peroxide as oxidizing agent. Hydrogen peroxide in combination with the percarboxylic acid can provide certain antimicrobial action against microorganisms. Additionally, 15 hydrogen peroxide can provide an effervescent action which can irrigate any surface to which it is applied. Hydrogen peroxide can work with a mechanical flushing action once applied which further cleans the surface of an object. An additional advantage of hydrogen peroxide is the food compatibility of this composition upon use and decomposition.

20 In certain embodiments, the present composition includes about 0.001 to about 30 wt-% oxidizing agent, about 0.001 to about 10 wt-% oxidizing agent, 0.002 to about 10 wt-% oxidizing agent, about 2 to about 30 wt-% oxidizing agent, about 2 to about 25 wt-% oxidizing agent, about 2 to about 20 wt-% oxidizing agent, about 4 to about 20 wt-% oxidizing agent, about 5 to about 10 wt-% oxidizing agent, or about 6 to about 10 25 wt-% oxidizing agent. The composition can include any of these ranges or amounts not modified by about.

Acidulant

In an embodiment, the present composition can include an acidulant. The 30 acidulant can act as a catalyst for conversion of carboxylic acid to peroxycarboxylic acid. The acidulant can be effective to form a concentrate composition with pH of about 1 or less. The acidulant can be effective to form a use composition with pH of

about 5, about 5 or less, about 4, about 4 or less, about 3, about 3 or less, about 2, about 2 or less, or the like. In an embodiment, the acidulant includes an inorganic acid.

Suitable inorganic acids include sulfuric acid, phosphoric acid, nitric acid, hydrochloric acid, methane sulfonic acid, ethane sulfonic acid, propane sulfonic acid, butane sulfonic acid, xylene sulfonic acid, benzene sulfonic acid, mixtures thereof, or the like.

In an embodiment, the acidulant includes a carboxylic acid with pK_a less than 4. Suitable carboxylic acids with pK_a less than 4 include hydroxyacetic acid, hydroxypropionic acid, other hydroxycarboxylic acids, mixtures thereof, or the like. Such an acidulant is present at a concentration where it does not act as a solubilizer.

In certain embodiments, the present composition includes about 0.001 to about 50 wt-% acidulant, about 0.001 to about 30 wt-% acidulant, about 1 to about 50 wt-% acidulant, about 1 to about 30 wt-% acidulant, about 2 to about 40 wt-% acidulant, about 2 to about 10 wt-% acidulant, about 3 to about 40 wt-% acidulant, about 5 to about 40 wt-% acidulant, about 5 to about 25 wt-% acidulant, about 10 to about 40 wt-% acidulant, about 10 to about 30 wt-% acidulant, about 15 to about 35 wt-% acidulant, about 15 to about 30 wt-% acidulant, or about 40 to about 60 wt-% acidulant. The composition can include any of these ranges or amounts not modified by about.

Stabilizing Agent

One or more stabilizing agents can be added to the composition of the invention, for example, to stabilize the peracid and hydrogen peroxide and prevent the premature oxidation of this constituent within the composition of the invention.

Suitable stabilizing agents include chelating agents or sequestrants. Suitable sequestrants include organic chelating compounds that sequester metal ions in solution, particularly transition metal ions. Such sequestrants include organic amino- or hydroxy-polyphosphonic acid complexing agents (either in acid or soluble salt forms), carboxylic acids (e.g., polymeric polycarboxylate), hydroxycarboxylic acids, or aminocarboxylic acids.

The sequestrant can be or include phosphonic acid or phosphonate salt.

Suitable phosphonic acids and phosphonate salts include 1-hydroxy ethylidene-1,1-diphosphonic acid ($\text{CH}_3\text{C}(\text{PO}_3\text{H}_2)_2\text{OH}$) (HEDP); ethylenediamine tetrakis methylenephosphonic acid (EDTMP); diethylenetriamine pentakis

methylenephosphonic acid (DTPMP); cyclohexane-1,2-tetramethylene phosphonic acid; amino[tri(methylene phosphonic acid)]; (ethylene diamine[tetra methylene-phosphonic acid]); 2-phosphene butane-1,2,4-tricarboxylic acid; or salts thereof, such as the alkali metal salts, ammonium salts, or alkylol amine salts, such as mono, di, or
5 tetra-ethanolamine salts; or mixtures thereof.

Suitable organic phosphonates include HEDP.

Commercially available food additive chelating agents include phosphonates sold under the trade name DEQUEST® including, for example, 1-hydroxyethylidene-1,1-diphosphonic acid, available from Monsanto Industrial Chemicals Co., St. Louis,
10 MO, as DEQUEST® 2010; amino(tri(methylenephosphonic acid)), (N[CH₂PO₃H₂]₃), available from Monsanto as DEQUEST® 2000; ethylenediamine[tetra(methylenephosphonic acid)] available from Monsanto as DEQUEST® 2041; and 2-phosphonobutane-1,2,4-tricarboxylic acid available from Mobay Chemical Corporation, Inorganic Chemicals Division, Pittsburgh, PA, as
15 Bayhibit AM.

The sequestrant can be or include aminocarboxylic acid type sequestrant. Suitable aminocarboxylic acid type sequestrants include the acids or alkali metal salts thereof, e.g., amino acetates and salts thereof. Suitable aminocarboxylates include N-hydroxyethylaminodiacetic acid; hydroxyethylenediaminetetraacetic acid,
20 nitrilotriacetic acid (NTA); ethylenediaminetetraacetic acid (EDTA); N-hydroxyethyl-ethylenediaminetriacetic acid (HEDTA); diethylenetriaminepentaacetic acid (DTPA); and alanine-N,N-diacetic acid; and the like; and mixtures thereof.

The sequestrant can be or include a polycarboxylate. Suitable polycarboxylates
25 include, for example, polyacrylic acid, maleic/olefin copolymer, acrylic/maleic copolymer, polymethacrylic acid, acrylic acid-methacrylic acid copolymers, hydrolyzed polyacrylamide, hydrolyzed polymethacrylamide, hydrolyzed polyamide-methacrylamide copolymers, hydrolyzed polyacrylonitrile, hydrolyzed polymethacrylonitrile, hydrolyzed acrylonitrile-methacrylonitrile copolymers,
30 polymaleic acid, polyfumaric acid, copolymers of acrylic and itaconic acid, phosphino polycarboxylate, acid or salt forms thereof, mixtures thereof, and the like.

In certain embodiments, the present composition includes about 0.5 to about 50 wt-% sequestrant, about 1 to about 50 wt-% sequestrant, about 1 to about 30 wt-% sequestrant, about 1 to about 15 wt-% sequestrant, about 1 to about 5 wt-% sequestrant, about 1 to about 4 wt-% sequestrant, about 2 to about 10 wt-% sequestrant, about 2 to about 5 wt-% sequestrant, or about 5 to about 15 wt-% sequestrant. The composition can include any of these ranges or amounts not modified by about.

In certain embodiments, the present composition includes about 0.001 to about 50 wt-% stabilizing agent, about 0.001 to about 5 wt-% stabilizing agent, about 0.5 to about 50 wt-% stabilizing agent, about 1 to about 50 wt-% stabilizing agent, about 1 to about 30 wt-% stabilizing agent, about 1 to about 10 wt-% stabilizing agent, about 1 to about 5 wt-% stabilizing agent, about 1 to about 3 wt-% stabilizing agent, about 2 to about 10 wt-% stabilizing agent, about 2 to about 5 wt-% stabilizing agent, or about 5 to about 15 wt-% stabilizing agent. The composition can include any of these ranges or amounts not modified by about.

15

Adjuvants

The antimicrobial composition of the invention can also include any number of adjuvants. Specifically, the composition of the invention can include antimicrobial solvent, antimicrobial agent, wetting agent, defoaming agent, thickener, a surfactant, foaming agent, solidification agent, aesthetic enhancing agent (i.e., colorant (e.g., pigment), odorant, or perfume), among any number of constituents which can be added to the composition. Such adjuvants can be preformulated with the antimicrobial composition of the invention or added to the system simultaneously, or even after, the addition of the antimicrobial composition. The composition of the invention can also contain any number of other constituents as necessitated by the application, which are known and which can facilitate the activity of the present invention.

25

Antimicrobial Solvent

Any of a variety of solvents can be useful as antimicrobial solvents in the present compositions. Antimicrobial solvent can be added to use compositions before use. Suitable antimicrobial solvents include acetamidophenol; acetanilide; acetophenone; 2-acetyl-1-methylpyrrole; benzyl acetate; benzyl alcohol; benzyl

30

benzoate; benzyloxyethanol; essential oils (e.g., benzaldehyde, pinenes, terpeneols, terpinenes, carvone, cinnamaldehyde, borneol and its esters, citrals, ionenes, jasmine oil, limonene, dipentene, linalool and its esters); diester dicarboxylates (e.g., dibasic esters) such as dimethyl adipate, dimethyl succinate, dimethyl glutarate (including
5 products available under the trade designations DBE, DBE-3, DBE-4, DBE-5, DBE-6, DBE-9, DBE-IB, and DBE-ME from DuPont Nylon), dimethyl malonate, diethyl adipate, diethyl succinate, diethyl glutarate, dibutyl succinate, and dibutyl glutarate; dimethyl sebacate, dimethyl pimelate, dimethyl suberate; dialkyl carbonates such as dimethyl carbonate, diethyl carbonate, dipropyl carbonate, diisopropyl carbonate, and
10 dibutyl carbonate; organo-nitriles such as acetonitrile and benzonitrile; and phthalate esters such as dibutyl phthalate, diethylhexyl phthalate, and diethyl phthalate. Mixtures of antimicrobial solvents can be used if desired.

The antimicrobial solvent can be selected based upon the characteristics of the surface and microbes to which the antimicrobial composition will be applied and upon
15 the nature of any coating, soil or other material that will be contacted by the antimicrobial composition and optionally removed from the surface. Polar solvents, and solvents that are capable of hydrogen bonding typically will perform well on a variety of surfaces and microbes and thus, for such applications, can be selected. In certain applications, the antimicrobial solvent can be selected for a high flashpoint (e.g.,
20 greater than about 30°C, greater than about 50°C, or greater than about 100°C), low odor, and low human and animal toxicity.

In an embodiment, the antimicrobial solvent is compatible as an indirect or direct food additive or substance; especially those described in the Code of Federal Regulations (CFR), Title 21--Food and Drugs, parts 170 to 186. The compositions of
25 the invention should contain sufficient antimicrobial solvent to provide the desired rate and type of microbial reduction.

The present composition can include an effective amount of antimicrobial solvent, such as about 0.01 wt-% to about 60 wt-% antimicrobial solvent, about 0.05 wt-% to about 15 wt-% antimicrobial solvent, or about 0.08 wt-% to about 5 wt-%
30 antimicrobial solvent.

Additional Antimicrobial Agent

The antimicrobial compositions of the invention can contain an additional antimicrobial agent. Additional antimicrobial agent can be added to use compositions before use. Suitable antimicrobial agents include carboxylic esters (e.g., p-hydroxy
5 alkyl benzoates and alkyl cinnamates), sulfonic acids (e.g., dodecylbenzene sulfonic acid), iodo-compounds or active halogen compounds (e.g., elemental halogens, halogen oxides (e.g., NaOCl, HOCl, HOBr, ClO₂), iodine, interhalides (e.g., iodine monochloride, iodine dichloride, iodine trichloride, iodine tetrachloride, bromine chloride, iodine monobromide, or iodine dibromide), polyhalides, hypochlorite salts,
10 hypochlorous acid, hypobromite salts, hypobromous acid, chloro- and bromo-hydantoins, chlorine dioxide, and sodium chlorite), organic peroxides including benzoyl peroxide, alkyl benzoyl peroxides, ozone, singlet oxygen generators, and mixtures thereof, phenolic derivatives (e.g., o-phenyl phenol, o-benzyl-p-chlorophenol, tert-amyl phenol and C₁-C₆ alkyl hydroxy benzoates), quaternary ammonium
15 compounds (e.g., alkyldimethylbenzyl ammonium chloride, dialkyldimethyl ammonium chloride and mixtures thereof), and mixtures of such antimicrobial agents, in an amount sufficient to provide the desired degree of microbial protection.

The present composition can include an effective amount of antimicrobial agent, such as about 0.001 wt-% to about 60 wt-% antimicrobial agent, about 0.01 wt-
20 % to about 15 wt-% antimicrobial agent, or about 0.08 wt-% to about 2.5 wt-% antimicrobial agent.

Wetting or Defoaming Agents

Also useful in the composition of the invention are wetting and defoaming
25 agents. Wetting agents function to increase the surface contact or penetration activity of the antimicrobial composition of the invention. Wetting agents which can be used in the composition of the invention include any of those constituents known within the art to raise the surface activity of the composition of the invention.

Suitable defoamers which can be used in accordance with the invention include
30 silica and silicones; aliphatic acids or esters; alcohols; sulfates or sulfonates; amines or amides; halogenated compounds such as fluorochlorohydrocarbons; vegetable oils, waxes, mineral oils as well as their sulfated derivatives; fatty acid soaps such as alkali,

alkaline earth metal soaps; and phosphates and phosphate esters such as alkyl and alkaline diphosphates, and tributyl phosphates among others; and mixtures thereof.

In an embodiment, the present compositions can include antifoaming agents or defoamers which are of food grade quality given the application of the method of the invention. To this end, one of the more effective antifoaming agents includes silicones. Silicones such as dimethyl silicone, glycol polysiloxane, methylphenol polysiloxane, trialkyl or tetraalkyl silanes, hydrophobic silica defoamers and mixtures thereof can all be used in defoaming applications. Commercial defoamers commonly available include silicones such as Ardefoam® from Armour Industrial Chemical Company which is a silicone bound in an organic emulsion; Foam Kill® or Kresseo® available from Krusable Chemical Company which are silicone and non-silicone type defoamers as well as silicone esters; and Anti-Foam A® and DC-200 from Dow Corning Corporation which are both food grade type silicones among others. These defoamers can be present at a concentration range from about 0.01 wt-% to 5 wt-%, from about 0.01 wt-% to 2 wt-%, or from about 0.01 wt-% to about 1 wt-%.

Thickening or Gelling Agents

The present compositions can include any of a variety of known thickeners. Suitable thickeners include natural gums such as xanthan gum, guar gum, or other gums from plant mucilage; polysaccharide based thickeners, such as alginates, starches, and cellulosic polymers (e.g., carboxymethyl cellulose); polyacrylates thickeners; and hydrocolloid thickeners, such as pectin. In an embodiment, the thickener does not leave contaminating residue on the surface of an object. For example, the thickeners or gelling agents can be compatible with food or other sensitive products in contact areas. Generally, the concentration of thickener employed in the present compositions or methods will be dictated by the desired viscosity within the final composition. However, as a general guideline, the viscosity of thickener within the present composition ranges from about 0.1 wt-% to about 1.5 wt-%, from about 0.1 wt-% to about 1.0 wt-%, or from about 0.1 wt-% to about 0.5 wt-%.

30

Solidification Agent

The present compositions can include a solidification agent, which can participate in maintaining the compositions in a solid form. Suitable solidification agents include a solid polyethylene glycol (PEG), a solid EO/PO block copolymer, and the like; an amide, such as stearic monoethanolamide, lauric diethanolamide, an alkylamide, or the like; starches that have been made water-soluble through an acid or alkaline treatment process; celluloses that have been made water-soluble; an inorganic agent, or the like; poly(maleic anhydride/methyl vinyl ether); polymethacrylic acid; other generally functional or inert materials with high melting points; and the like.

In certain embodiments, the solidification agent includes solid PEG, for example PEG 1500 up to PEG 20,000. In certain embodiments, the PEG includes PEG 1450, PEG 3350, PEG 4500, PEG 8000, PEG 20,000, and the like. Additional suitable solidification agents include EO/PO block copolymers such as those sold under the tradenames Pluronic 108, Pluronic F68; amides such as lauric diethanolamide or cocodiethylene amide; and the like. In certain embodiments, the solidification agent includes a combination of solidification agents, such as combination of PEG and an EO/PO block copolymer (such as a Pluronic) and combination of PEG and an amide (such as lauric diethanolamide amide or stearic monoethanol amide).

Fragrance

In an embodiment, the present composition includes a fragrance. The fragrance can be selected to avoid undesirable effects on the stability or efficacy of the composition. Suitable fragrances include amyl acetate, iso-bornyl acetate, and alkyl salicylates, such as methyl salicylate. In an embodiment, the fragrance can include an alkylsalicylate.

Additional Embodiments of the Medium Chain Peroxycarboxylic Acid Compositions

The present invention relates to compositions including medium chain peroxycarboxylic acid, methods for making these compositions, and methods for reducing the population of a microorganism. In certain embodiments, the compositions

can include advantageously high levels of the medium chain peroxycarboxylic acid, can be readily made, and/or can exhibit reduced odor.

In an embodiment, the present compositions can include medium chain peroxycarboxylic acid, medium chain carboxylic acid, carrier, and solubilizer. In
5 certain embodiments, the present compositions include about 2 or more parts of medium chain peroxycarboxylic acid for each 7 parts of medium chain carboxylic acid; about 2 or more parts of medium chain peroxycarboxylic acid for each 5 parts of medium chain carboxylic acid; about 2 or more parts of medium chain
10 peroxycarboxylic acid for each 4 parts of medium chain carboxylic acid; or about 2 parts of medium chain peroxycarboxylic acid for each 3 parts of medium chain carboxylic acid.

In an embodiment, the solubilizer includes solvent, surfactant, or mixture thereof. In an embodiment, the surfactant solubilizer includes a microemulsion forming surfactant, e.g., an anionic surfactant. In an embodiment, the composition
15 includes a microemulsion. In an embodiment, the solubilizer includes polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, anionic surfactant, or mixture thereof. In an embodiment, the solvent solubilizer includes polyalkylene oxide, capped polyalkylene oxide, nonionic surfactant, or mixture thereof.

In an embodiment, the present compositions include no, only insignificant, or
20 relatively small amounts of short chain peroxycarboxylic acid, short chain carboxylic acid, or mixture thereof. For example, in an embodiment, the composition can be substantially free of added short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof. For example, in an embodiment, the composition can include short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof at a
25 level insufficient to solubilize medium chain peroxycarboxylic acid. For example, in an embodiment, the composition can include short chain carboxylic acid, short chain peroxycarboxylic acid, or mixture thereof at a level insufficient to cause objectionable odor. For example, in an embodiment, the composition can include about 1 or more parts of medium chain peroxycarboxylic acid for each 8 parts of short chain carboxylic
30 acid, short chain peroxycarboxylic acid, or mixture thereof.

In an embodiment, the composition also includes oxidizing agent, inorganic acid, stabilizing agent, another adjuvant or additive, or mixture thereof.

In an embodiment, the present invention includes a method of making a medium chain peroxy-carboxylic acid composition. The method can include reacting medium chain carboxylic acid and oxidizing agent in the presence of carrier, solubilizer, acidulant, stabilizing agent, or mixture thereof. The method can form advantageously high levels of medium chain peroxy-carboxylic acids in advantageously short times. For example, in an embodiment, the present method includes converting 20% or more of the medium chain carboxylic acid to medium chain peroxy-carboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method includes converting about 25% or more of the medium chain carboxylic acid to medium chain peroxy-carboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method includes converting about 30% or more of the medium chain carboxylic acid to medium chain peroxy-carboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method includes converting about 35% or more of the medium chain carboxylic acid to medium chain peroxy-carboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method includes converting about 40% of the medium chain carboxylic acid to medium chain peroxy-carboxylic acid in about 24 or fewer hours.

In an embodiment, the present invention includes a method of using a medium chain peroxy-carboxylic acid composition. The method can include contacting an object with the present composition (e.g., a use composition) and can result in reducing the population of one or more microorganisms on the object.

Use Compositions

The present compositions include concentrate compositions and use compositions. For example, a concentrate composition can be diluted, for example with water, to form a use composition. In an embodiment, a concentrate composition can be diluted to a use solution before to application to an object. For reasons of economics, the concentrate can be marketed and an end user can dilute the concentrate with water or an aqueous diluent to a use solution.

The level of active components in the concentrate composition is dependent on the intended dilution factor and the desired activity of the medium chain peroxy-carboxylic acid compound. Generally, a dilution of about 1 fluid ounce to about

20 gallons of water to about 5 fluid ounces to about 1 gallon of water is used for aqueous antimicrobial compositions. Higher use dilutions can be employed if elevated use temperature (greater than 25 °C) or extended exposure time (greater than 30 seconds) can be employed. In the typical use locus, the concentrate is diluted with a major proportion of water using commonly available tap or service water mixing the materials at a dilution ratio of about 3 to about 20 ounces of concentrate per 100 gallons of water.

For example, a use composition can include about 0.01 to about 4 wt-% of a concentrate composition and about 96 to about 99.99 wt-% diluent; about 0.5 to about 4 wt-% of a concentrate composition and about 96 to about 99.5 wt-% diluent; about 0.5, about 1, about 1.5, about 2, about 2.5, about 3, about 3.5, or about 4 wt-% of a concentrate composition; about 0.01 to about 0.1 wt-% of a concentrate composition; or about 0.01, about 0.02, about 0.03, about 0.04, about 0.05, about 0.06, about 0.07, about 0.08, about 0.09, or about 0.1 wt-% of a concentrate composition. Amounts of an ingredient in a use composition can be calculated from the amounts listed above for concentrate compositions and these dilution factors.

The present methods can employ medium chain peroxycarboxylic acid at a concentration effective for reducing the population of one or more microorganisms. Such effective concentrations include about 2 to about 500 ppm medium chain peroxycarboxylic acid, about 2 to about 300 ppm medium chain peroxycarboxylic acid, about 5 to about 100 ppm medium chain peroxycarboxylic acid, about 5 to about 60 ppm medium chain peroxycarboxylic acid, about 5 to about 45 ppm medium chain peroxycarboxylic acid, about 5 to about 35 ppm medium chain peroxycarboxylic acid, about 5 to about 25 ppm medium chain peroxycarboxylic acid, about 8 to about 50 ppm medium chain peroxycarboxylic acid, about 10 to about 500 ppm medium chain peroxycarboxylic acid, about 10 to about 50 ppm medium chain peroxycarboxylic acid, about 40 to about 140 ppm medium chain peroxycarboxylic acid, about 100 to about 250 ppm medium chain peroxycarboxylic acid, or about 200 to about 300 ppm medium chain peroxycarboxylic acid. In an embodiment, the use composition can include about 2 to about 500 ppm medium chain peroxycarboxylic acid, about 5 to about 2000 ppm medium chain carboxylic acid, about 95 to about 99.99 wt-% carrier and/or diluent

(e.g., water); and about 2 to about 23,000 ppm polyalkylene oxide, capped polyalkylene oxide, alkoxyated surfactant, anionic surfactant, or mixture thereof.

The level of reactive species, such as peroxycarboxylic acids and/or hydrogen peroxide, in a use composition can be affected, typically diminished, by organic matter that is found in or added to the use composition. For example, when the use composition is a bath or spray used for washing an object, soil on the object can consume peroxy acid and peroxide. Thus, the present amounts of ingredients in the use compositions refer to the composition before or early in use, with the understanding that the amounts will diminish as organic matter is added to the use composition.

In an embodiment, the present use composition can be made more acidic by passing the concentrate through an acidifying column, or by adding additional acidulant to the use composition.

Other Fluid Compositions

The present and compositions can include a critical, near critical, or supercritical (densified) fluid and an antimicrobial agent or a gaseous composition of an antimicrobial agent. The densified fluid can be a near critical, critical, supercritical fluid, or another type of fluid with properties of a supercritical fluid. Fluids suitable for densification include carbon dioxide, nitrous oxide, ammonia, xenon, krypton, methane, ethane, ethylene, propane, certain fluoroalkanes (e.g., chlorotrifluoromethane and monofluoromethane), and the like, or mixtures thereof. Suitable fluids include carbon dioxide.

In an embodiment, the present compositions or methods include densified carbon dioxide, medium chain peroxycarboxylic acid, and medium chain carboxylic acid. Such a composition can be referred to as a densified fluid medium chain peroxycarboxylic acid composition. In another embodiment, the antimicrobial composition includes the fluid, an antimicrobial agent, and any of the optional or added ingredients, but is in the form of a gas.

Densified fluid antimicrobial compositions can be applied by any of several methods known to those of skill in the art. Such methods include venting at an object a vessel containing densified fluid and antimicrobial agent. The aqueous phase, which includes hydrogen peroxide, is advantageously retained in the device. The vented gas

includes an effective amount of antimicrobial agent making the densified fluid peroxycarboxylic acid compositions effective antimicrobial agents.

Because of the high pressure nature of the densified fluid compositions of the invention, these compositions are typically applied by venting a vessel containing the composition through a pressure relief device that is designed to promote rapid efficient coverage of an object. Devices including such a pressure relief device include sprayers, foggers, foamers, foam pad applicators, brush applicators or any other device that can permit the expansion of the fluid materials from high pressure to ambient pressure while applying the material to an object. The densified fluid peroxycarboxylic acid composition can also be applied to an object by any of a variety of methods known for applying gaseous agents to an object.

Densified fluid antimicrobial compositions can be made by reacting an oxidizable substrate with an oxidizing agent in a medium comprising a densified fluid to form an antimicrobial composition. This reaction is typically carried out in a vessel suitable for containing a densified fluid. Reacting can include adding to the vessel the oxidizable substrate and the oxidizing agent, and adding fluid to the vessel to form the densified fluid. In an embodiment, the reaction is between a medium chain carboxylic acid and hydrogen peroxide to form the corresponding peroxycarboxylic acid. The hydrogen peroxide is commonly supplied in the form of an aqueous solution of hydrogen peroxide.

Supercritical, subcritical, near supercritical, and other dense fluids and solvents that can be employed with such fluids are disclosed in U.S. Patent No. 5,306,350, issued April 26, 1994 to Hoy et al., which is incorporated by reference herein for such disclosure. Supercritical and other dense forms of carbon dioxide, and cosolvents, co-surfactants, and other additives that can be employed with these forms of carbon dioxide are disclosed in U.S. Patent No. 5,866,005, issued February 2, 1999 to DeSimone et al., which is incorporated by reference herein for such disclosure.

Making Medium Chain Peroxycarboxylic Acid Compositions

The compositions of or used in the methods of the invention can be made by combining or reacting the medium chain carboxylic acid and the oxidizing agent, such as hydrogen peroxide. Combining or reacting medium chain carboxylic acid and oxidizing

agent results in production of medium chain peroxycarboxylic acid. In an embodiment, combining includes mixing. The formulation combined for making the present compositions can also include the solubilizer, the acidulant, the carrier, stabilizing agent, mixtures thereof, or the like. In an embodiment, the formulation includes solubilizer.

- 5 Alternatively, one or more (e.g., at least one) of the solubilizer, the acidulant, the carrier, or mixtures thereof, can be added after production of some or all of the peroxycarboxylic acid.

In an embodiment, the present invention includes a method of making a medium chain peroxycarboxylic acid. The method can include combining or reacting
10 medium chain carboxylic acid, carrier (e.g., water), oxidizing agent (e.g., hydrogen peroxide), solubilizer, acidulant, and stabilizing agent. The method can include mixing the ingredients at concentrations of about 1 to about 10 wt-% medium chain carboxylic acid, about 0 to about 98 wt-% carrier, about 2 to about 30 wt-% oxidizing agent, about 1 to about 80 wt-% solubilizer, about 1 to about 50 wt-% acidulant, and about 0.5 to
15 about 50 wt-% stabilizing agent. The method can include mixing the ingredients at concentrations about 1 to about 10 wt-% medium chain carboxylic acid, about 5 to about 97 wt-% carrier, about 2 to about 30 wt-% oxidizing agent, about 1 to about 20 wt-% solubilizer (e.g., microemulsion forming surfactant), about 1 to about 50 wt-% acidulant, and about 0.5 to about 50 wt-% stabilizing agent. The present compositions
20 also include compositions in which these combinations of ingredients have come to equilibrium forming medium chain peroxycarboxylic acid.

In an embodiment, the present method produces advantageously high levels of medium chain peroxycarboxylic acid in advantageously short times. Advantageously short times include, for example, about 24 or fewer hours, about 6 or fewer hours,
25 about 3 or fewer hours, or about 0.5 hr. In an embodiment, high levels of medium chain peroxycarboxylic acid can be achieved nearly instantaneously. High levels of medium chain peroxycarboxylic acid be achieved by converting 20% or more, 25% or more, 30% or more, 35% or more, or 40% of the medium chain carboxylic acid to medium chain peroxycarboxylic acid. Such conversions can be achieved at room
30 temperature or in a reaction started at room temperature and warmed by an exotherm. Lower temperatures can require a longer time to reach the same amount of conversion.

The amount of time is typically measured from the time that the carboxylic acid, oxidizing agent, solubilizer, and acidulant are combined or reacted.

For example, in an embodiment, the present method can convert 20% or more of the medium chain carboxylic acid to medium chain peroxycarboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method can convert about 25% or more of the medium chain carboxylic acid to medium chain peroxycarboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method can convert about 30% or more of the medium chain carboxylic acid to medium chain peroxycarboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method can convert about 35% or more of the medium chain carboxylic acid to medium chain peroxycarboxylic acid in about 24 or fewer hours. For example, in an embodiment, the present method can convert about 40% of the medium chain carboxylic acid to medium chain peroxycarboxylic acid in about 24 or fewer hours.

In an embodiment, making the present compositions includes forming a microemulsion. A microemulsion can be formed by mixing the desired ingredients including a microemulsion forming surfactant. The method can include combining or mixing the ingredients at concentration of about 1 to about 10 wt-% medium chain carboxylic acid, about 5 to about 97 wt-% carrier (e.g., water), about 2 to about 30 wt-% oxidizing agent, about 1 to about 20 wt-% microemulsion forming surfactant, and about 1 to about 50 wt-% stabilizer. The present compositions also include compositions in which these combinations of ingredients have come to equilibrium forming medium chain peroxycarboxylic acid. The components can be added in any of a variety of orders. In an embodiment, formation of the medium chain peroxy carboxylic acid can proceed rapidly after the addition of the microemulsion forming surfactant. Although not limiting to the present invention, it is believed that the formation of the microemulsion can significantly increase the effective surface area of the medium chain carboxylic acid (as micro-droplets) for reaction.

The present compositions can be made in a plant as a concentrate and shipped to an end user who need only dilute the concentrate to form a use composition. The present medium chain peroxycarboxylic acid compositions can also be made at the site of use. For example, the product can be shipped as a two or more part composition or

as a kit. The user can then combine the two or more compositions or components of the kit to produce the present medium chain peroxycarboxylic acid compositions. Alternatively, a system of formulating equipment and containers of raw materials can be provided at the site of use, and programmed or operated to mix and disperse the
5 present medium chain peroxycarboxylic acid compositions.

In an embodiment, the product can be supplied as a two or more part composition. One composition can include carboxylic acid and one or more (e.g., at least one) of solubilizer, acidulant, carrier, stabilizing agent, mixtures thereof, or the like. The second composition can include oxidizing agent and one or more (e.g., at
10 least one) of solubilizer, acidulant, carrier, stabilizing agent mixtures thereof, or the like. Alternatively, the solubilizer, acidulant, carrier, stabilizing agent mixtures thereof, or the like can be supplied as additional composition(s).

In an embodiment, the pH of a concentrate composition can be less than about 1 or about 2. In an embodiment, the pH of a 1% or 1.5% solution of the mixture in water
15 is about 1 or 2 to about 7, depending on the other components of the 1% solution. In an embodiment, the pH of a use composition can be from about 2 to about 7 depending on the other components.

Some examples of representative concentrations of ingredients useful in the present methods of making medium chain peroxycarboxylic acid compositions can be
20 found in Tables G and H, in which the values are given in wt-% of the ingredients in reference to the total composition weight. In certain embodiments, the proportions and amounts in Tables G-H can be modified by "about". The present compositions also include compositions in which these combinations of ingredients have come to equilibrium forming medium chain peroxycarboxylic acid.

25

Table G

Ingredient	wt-%	wt-%	wt-%	wt-%	wt-%	wt-%	wt-%	wt-%
medium chain carboxylic acid	1-10	3-8	4-6	2-8	3-6	1-10	3-8	3-6
solubilizer	1-80	2-70	3-65	5-70	10-65	1-25	3-15	4-10
carrier	0-98	5-90	10-80	0.2-60	5-20	5-97	15-70	30-75

Table H

Ingredient	wt-%	wt-%	wt-%	wt-%	wt-%	wt-%	wt-%	wt-%
medium chain carboxylic acid	1-10	3-8	4-6	2-8	3-6	1-10	3-8	3-6
solubilizer	1-80	2-70	3-65	5-70	10-65	1-25	3-15	4-10
carrier	0-98	5-90	10-80	0.2-60	5-20	5-97	15-70	30-75
oxidizing agent	2-30	2-25	4-20	2-25	4-20	2-30	4-20	6-10
acidulant	1-50	2-40	3-40	2-40	3-40	1-50	3-35	5-30
stabilizing agent	1-50	1-10	1-5	1-10	1-5	1-50	1-5	1-3

The present invention may be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

EXAMPLES

Example 1 - - Compositions Including Medium Chain Peroxycarboxylic Acid and Solubilizer

Tables 1-5 present illustrative examples of the present compositions including medium chain peroxycarboxylic acid and solubilizer. Quantities in the tables are in wt-%.

Table 1 - Examples of Compositions Including Solvent Solubilizer

Ingredient	A	B	C	D	E
Medium Chain Peroxycarboxylic Acid	1.8	1.6	1.4	1.6	2.9
Medium Chain Carboxylic Acid	3.4	3.6	3.7	3.6	2.4
Solubilizer	60	40	60	60	40
Carrier	25	22	25	22	22
Oxidizing Agent	7.0	6.6	7.0	6.9	6.9
Acidulant	2	25	2	5	25
Stabilizing Agent	1.2	1.2	1.2	1.2	1.2

In each of compositions A-Q: the medium chain peroxycarboxylic acid was peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier was water; the oxidizing agent was hydrogen peroxide (supplied from a 35% solution); and the stabilizing agent was HEDP (supplied as Dequest 2010 which includes 60 wt-%
5 HEDP).

In each of compositions A-L, O, P, and Q: the acidulant was concentrated sulfuric acid. In compositions M and N, the acidulant was phosphoric acid (supplied as 85% and 75% phosphoric acid, respectively).

The solubilizer was varied among these compositions. In compositions A and B, the
10 solubilizer was polyethyleneglycol 300. In compositions C, D, and E, the solubilizer was monomethyl ether of polyethyleneglycol (MPEG 550). In composition F, the solubilizer was nonionic surfactant, specifically Pluronic 17R4 an $(PO)_x(EO)_y(PO)_x$ reverse triblock copolymer with 40% EO and 60% PO. In composition G, the solubilizer was polyethyleneglycol 300 plus LAS acid (98% linear dodecylbenzene
15 sulfonic acid). In composition H, the solubilizer was polyethyleneglycol 300 plus 1-octane sulfonate (supplied under the tradename NAS-FAL as 38% active). In composition I, the solubilizer was polyethyleneglycol 300 plus Dowfax Hydrotrope acid (C_6 alkylated diphenyl oxide disulfonic acid). In composition J, the solubilizer was dimethyl ether of polyethyleneglycol (PolyDME250) and LAS acid. In
20 composition K, the solubilizer was dimethyl ether of polyethyleneglycol (PolyDME250) and NAS-FAL. In composition L, the solubilizer was dimethyl ether of polyethyleneglycol (PolyDME250) and Dowfax Hydrotrope acid. In compositions M, N, O and P, the solubilizer was dimethyl ether of polyethyleneglycol (PolyDME250) and NAS-FAL. In composition Q, the solubilizer was dimethyl ether of
25 polyethyleneglycol (PolyDME250) and NAS acid (supplied as 93% 1-octane sulfonic acid).

These compositions were made from a composition including 5 wt-% medium chain carboxylic acid.

In each of compositions R-Z: the medium chain peroxycarboxylic acid was
30 peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier was water; the oxidizing agent was hydrogen peroxide (supplied from a 35% solution);

and the stabilizing agent was HEDP (supplied as Dequest 2010 which includes 60 wt-% HEDP).

In compositions R and S, the acidulant was phosphoric acid (supplied as 75% phosphoric acid). In each of compositions T, U, and V, the acidulant was reagent
5 grade,

Table 2 - Examples of Compositions Including Solvent Solubilizer and Surfactant Solubilizer

Ingredient	F	G	H	I	J	K	L	M	N	O	P	Q
Medium Chain Peroxycarboxylic Acid	0.8	0.7	1.1	1.1	0.9	2.1	1.6	0.7	0.9	5.0	not measured	5.0
Medium Chain Carboxylic Acid	4.3	4.4	4.0	4.0	4.2	4.2	3.1	4.4	4.2	0.2	<5	0.2
Solvent Solubilizer	0	40	40	40	42	44	42	34	29	28	28	28
Surfactant Solubilizer	45	5	2	5	8	6	7	6	4	6	6	10
Carrier	37	30	33	30	29	21	24	26	28	28	26	24
Oxidizing Agent	7.0	6.9	6.8	6.9	6.1	6.4	6.5	6.7	6.5	6.9	8.7	6.9
Acidulant	5	7	7	7	8	15	15	21	26	25	25	25
Stabilizing Agent	1.2	6	6	6	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2

Table 3 - Examples of Compositions Including Surfactant Solubilizer

Ingredient	R	S	T	U	V	W	X	Y	Z
Medium Chain Peroxycarboxylic Acid	0.5	0.4	1.0	1.0	0.7	3.8	3.7	3.8	3.5
Medium Chain Carboxylic Acid	4.6	4.6	3.1	3.1	3.4	2.6	2.7	2.6	2.9
Surfactant Solubilizer	17	20	20	20	20	20	20	20	20
Carrier	32	29	27	27	27	24	24	24	24
Oxidizing Agent	8.0	8.3	9.2	9.2	9.3	8.6	8.7	8.6	8.7
Acidulant	36	36	38	38	38	39	39	39	39
Stabilizing Agent	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4

Table 4 - Examples of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer

Ingredient	AA	AA-O	BB	CC	DD	EE	FF	GG	HH	II	JJ	KK
Medium Chain Peroxycarboxylic Acid	1.5	2.0	1.2	1.4	1.3	1.4	1.1	1.0	1.2	1.0	1.1	1.3
Medium Chain Carboxylic Acid	3.6	2.7	2.9	2.5	2.6	2.5	2.8	2.9	2.9	3.1	3.0	2.6
Solubilizer	8	5	5	9	4	4	6	4	5	5	5	4
Carrier	41	45	69	52	59	60	62	56	67	67	67	55
Oxidizing Agent	7.7	7.4	6.3	7.8	8.0	7.6	7.9	8.0	7.8	7.3	7.8	8.1
Acidulant	36	36	14	25	23	23	18	26	14	15	14	27
Stabilizing Agent	2.4	2.4	1.8	2.3	2.3	2.3	2.3	2.3	1.8	1.8	1.8	2.0

Table 4 - continued

Ingredient	LL	MM	NN	OO	PP	QQ	RR	SS	TT	UU	VV
Medium Chain Peroxycarboxylic Acid	1.4	1.1	1.5	not determined	0.9	0.5	0.54	3.4	0.2	1.0	0.4
Medium Chain Carboxylic Acid	2.5	2.7	2.3	<3.8	3.1	3.3	3.3	0.5	3.6	2.8	3.4
Solubilizer	4	4	4	5	1	2	4	10	6	10	22
Carrier	56	57	57	40-50	60	59	58	53	54	51	39
Oxidizing Agent	7.8	6.9	6.5	<8	7.1	7.5	7.5	5.6	7.8	8.0	7.7
Acidulant	26	26	26	26	26	26	26	26	1.8	1.8	1.8
Stabilizing Agent	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	27	27	27

98%, concentrated sulfuric acid (15 wt-%) and phosphoric acid (23 wt-%) (supplied as 75% phosphoric acid). In compositions W, X, Y, and Z, the acidulant was concentrated sulfuric acid (25 wt-%) and phosphoric acid (14 wt-%) (supplied as 75% phosphoric acid).

5 The solubilizer was varied among these compositions. In composition R, the solubilizer was 1-octane sulfonate (1.9 wt-%) and Tegotens EC-11 (a butoxy capped alcohol ethoxylate, a fast wetting surfactant) (15 wt-%). In compositions S, T, and W the solubilizer was Tegotens EC-11. In compositions U and Y, the solubilizer was Dehypon LS-54 ($(R(EO)_5(PO)_4$, a fast wetting surfactant). In compositions V and Z, the
10 solubilizer was Dehypon LT-104 (a butyl capped alcohol ethoxylate). In composition X, the solubilizer was LF-221 (a butoxy capped alcohol ethoxylate).

 In each of compositions AA-VV: the medium chain peroxy-carboxylic acid was peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier was water; the oxidizing agent was hydrogen peroxide (supplied as 35% hydrogen
15 peroxide in water); and the stabilizing agent was HEDP (supplied as Dequest 2010, which includes 60 wt-% HEDP).

 In each of compositions AA, AA-O, DD, EE, GG, KK, LL, MM, NN, OO, PP, QQ, RR, SS, TT, UU, and VV the acidulant was phosphoric acid (supplied as 75% phosphoric acid). In composition BB, HH the acidulant was concentrated sulfuric acid
20 (reagent grade, 98%). In composition CC, the acidulant was methane sulfonic acid (99.5 % + Aldrich). In composition FF, the acidulant was nitric acid (supplied as 70% nitric acid). In composition II, the acidulant was concentrated sulfuric acid (technical grade, 93%). In composition JJ, the acidulant was sulfuric acid (supplied as 50% sulfuric acid).

25 The solubilizer was varied among these compositions. In compositions AA, AA-O, BB, CC, DD, FF, LL, HH, II, and JJ, the solubilizer was 1-octane sulfonate. In compositions EE and GG, the solubilizer was 1-octane sulfonate (3.8 wt-%) and Dehypon LS-54 (0.2 wt-%). In composition KK, the solubilizer was 1-octane sulfonate (NAS-FAL). In composition MM, the solubilizer was 1-octane sulfonate (3.8 wt-%)
30 and Barlox 12 (dodecyldimethyl amine oxide, 30% active) (0.25 wt-%). In composition NN, the solubilizer was 1-octane sulfonate (3.8 wt-%) and Barlox 12 (0.5 wt-%). In composition OO, the solubilizer was 1-octane sulfonate (3.8 wt-%) and

Barlox 12 (1 wt-%). In compositions PP, QQ, RR, and SS, the solubilizer was LAS-acid. In composition TT, the solubilizer was disodium cocoampho dipropionate (supplied under the tradename Miranol[®] FBS, which includes 39% solids). In composition UU, the solubilizer was an aminopropionate betaine (supplied under the tradename Mirataine[®] JC-HA, which includes 42% solids). In composition VV, the solubilizer C12-13 alcohol 4 mole EO carboxylic acid (supplied under the tradename Neodox 23-4, which includes 90% active).

The quantities of medium chain peroxy-carboxylic acid were determined in compositions PP, QQ, RR, and SS after 7.5 days at 60 °C.

Table 5 - Examples of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer plus Strong Organic Acidulant

Ingredient	WW	XX	YY	ZZ	BA
Medium Chain Peroxy-carboxylic Acid	1.5	1.3	0.5	0.5	0.8
Medium Chain Carboxylic Acid	2.5	2.7	3.5	3.5	3.2
Solubilizer	4	4	4	4	4
Carrier	58	58	56	57	71
Oxidizing Agent	7.7	7.6	7.7	8.1	8.2
Acidulant	24	24	26	25	11
Stabilizing Agent	1.8	1.8	1.8	1.8	1.8

In each of compositions WW, XX, YY, ZZ, and BA: the medium chain peroxy-carboxylic acid was peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier was water; the oxidizing agent was hydrogen peroxide (supplied as 35% hydrogen peroxide in water); the stabilizing agent was HEDP (supplied as Dequest 2010, which includes 60 wt-% HEDP); and the solubilizer was NAS-FAL.

The acidulant was varied among these compositions. In composition WW, the acidulant was hydroxyacetic acid (supplied as 75% hydroxyacetic acid) (19 wt-%) and sulfuric acid (reagent grade, 98%) (5 wt-%). In composition XX, the acidulant was hydroxyacetic acid (supplied as 75% hydroxyacetic acid) (19 wt-%) and methane

sulfonic acid (99.5 % + Aldrich) (5 wt-%). In composition YY, the acidulant was hydroxyacetic acid (supplied as 75% hydroxyacetic acid). In composition ZZ, the acidulant was purified hydroxyacetic acid. In composition BA, the acidulant was hydroxypropionic acid (supplied as 22% 3-hydroxypropionic acid).

5 In these compositions the hydroxycarboxylic acids contributed virtually no solubilization of the medium chain carboxylic acid. The compositions required solubilizer.

Making the Exemplified Compositions

10 Table 6 shows the rapid generation of peroxyoctanoic acid achieved in making composition KK.

Table 6 - Generation of Peroxyoctanoic Acid with Time at Room Temperature and at 120 °F (Composition KK)

Minutes at RT	[POOA] wt-%	Minutes at 120 °F	[POOA] wt-%
11	0.61	30	1.46
53	1.09	45	1.38
97	1.11	60	1.23
130	1.1	90	1.47
235	1.24	120	1.31
293	1.27		
330	1.46		
366	1.39		
395	1.5		

15

When a high level of sulfuric acid was used as the acidulant (Examples include B, E, O, and Q), a strong exotherm was obtained, and the medium chain peroxy carboxylic acid was generated rapidly, for example, virtually instantaneously. For some of these compositions, the sulfuric acid needed to be added slowly and with
 20 cooling to keep the temperature below 170 °F or below 120 °F. Such formulas that can generate medium chain peroxy carboxylic acids, rapidly or almost instantaneously can be employed for on site generation at the use location.

The concentrations of peroxyoctanoic acid reported in the present examples were determined by a well established and standardized titration protocol. First,
 25 hydrogen peroxide content was determined by an oxidation-reduction titration with

ceric sulfate. After the endpoint of this titration was reached, an excess of potassium iodide was added to the solution. The potassium iodide reacts with peroxycarboxylic acids to liberate iodine. The liberated iodine was titrated with a standard solution of sodium thiosulfate to yield the concentration of peroxycarboxylic acid. The remaining
5 level of carboxylic acid can be calculated.

The octanoic acid employed in the present examples was obtained from sources including Procter & Gamble Chemicals and includes a minimum of 95% octanoic acid with minor amounts of hexanoic acid (ca. 2%), decanoic acid (ca. 2%), and dodecanoic acid (<0.5%).

10

Example 2 - - Stability of Compositions Including Medium Chain

Peroxycarboxylic Acid and Solubilizer

Compositions according to the present invention were evaluated and demonstrated physical stability and advantageous stability of the medium chain
15 peroxycarboxylic acid.

Materials and Methods

Several of the present medium chain peroxycarboxylic acid compositions were evaluated for stability of the medium chain peroxycarboxylic acid. A sealed container
20 including the composition was placed in an oven at an elevated temperature or was left at room temperature for a period of time. The temperatures and times are reported in the tables below. One week at 60 °C can be considered equivalent to a year at room temperature (RT). The quantity of peroxycarboxylic acid was determined by titration.

Several of the present medium chain peroxycarboxylic acid compositions were
25 also evaluated for physical stability. The sample were visually inspected at intervals at which peroxycarboxylic acid level was also determined.

Results

The results obtained for determinations of stability of the medium chain
30 peroxycarboxylic acid and of physical stability are reported below in Tables 7 and 8.

The results presented in Table 7 for compositions M and N indicate that stability of the medium chain peroxycarboxylic acid decreases when phosphoric acid

increases from 25 % to 35%. This suggests that the compositions including solvent solubilizer are susceptible to degradation caused by impurities present in the technical grade phosphoric acid.

5 The results presented in Table 8, specifically the blue tyndall appearance, indicates that each of these compositions was in the form of a microemulsion.

A study of accelerated aging of a mixed peroxydicarboxylic acid composition demonstrated that peroxyoctanoic acid in a mixed peracid composition underwent significant degradation at 60 °C in 7 days. After 7 days, three samples underwent 20, 23, and 54 % degradation.

10 The microemulsion compositions were less susceptible to degradation by impurities. For example, compositions KK and LL included technical grade phosphoric acid and exhibited good stability. In contrast, if phosphoric acid is to be used in conventional formulations of peroxydicarboxylic acids, high purity grade is required to avoid unacceptable degradation.

15 Compositions A, B, C, D, and E were two phase compositions.

Table 7 - Advantageous Stability of Medium Chain Peroxycarboxylic Acid in the Present Compositions Including Solvent Solubilizer

Composition	Starting [POOA] (wt-%)	Days at 100 °F	Wt-% Remaining, 100 °F	Days at RT	Wt-% Remaining, RT
A	1.8 (after 1 day at 100 °F)	22	1	46	2.3
B	1.6	37	0.8	37	2.1
C	1.4	36	0.9	36	1.3
D	1.6	36	0.7	36	1.4
E	2.9	36	0.4	36	1.8
F	0.8	31	1.1	31	0.9
J	0.9 (after 3 days at RT)	33	1.2	13	1.2
K	2.1 (after 3 days at RT)	33	1.1	17	2.0
L	1.6 (after 3 days at RT)	22	1.2	13	1.5
M	0.7	28	1	8	1.1
N	0.9	28	0.7	7	1.4

Table 8 - Stability of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer

Composition	Starting [POOA] (wt-%)	Days at 60 °C	Wt-% Remaining	Appearance
LL	1.4 (after 1 day at 60 °C)	7	1.4	1 phase, hazy blue tyndall
HH	1.2 (after 3 days at 60 °C)	7	1.2	Blue tyndall gel with no bubbles in solution. Slightly hazy.
KK	1.3	7	1.3	1 phase, hazy blue tyndall

5 **Example 3 - - Shear Thinning Viscosity of Compositions Including
Medium Chain Peroxycarboxylic Acid and Solubilizer**

Compositions according to the present invention were evaluated and demonstrated to have advantageous shear thinning viscosity, which is characteristic of microemulsions.

10

Materials and Methods

Several of the present medium chain peroxycarboxylic acid compositions were evaluated for viscosity as a function of rate of spindle rotation using an LVT viscometer and an N2 spindle. The temperature of the compositions was room temperature (about 75 °F).

15

Results

The results obtained for determinations of viscosity of the present compositions are reported below in Table 7. Decreasing viscosity with increasing spindle rotation rate indicates shear thinning, which is characteristic of a microemulsion. Each of the compositions tested showed shear thinning viscosity.

20

Table 9 - Shear Thinning Viscosity of Composition LL

rpm	Viscosity (cp)		rpm	Viscosity (cp)
0.6	3875		2	2260
1.5	2600		2.5	1952
3	1700		4	1380
6	1300		5	1208
12	863		10	736
30	483		20	468
60	308		50	280
			100	204

Table 10 - Shear Thinning Viscosity of Composition HH

rpm	Viscosity (cp)		rpm	Viscosity (cp)
0.6	7000		2	3500
1.5	3500		2.5	2848
3	2200		4	1950
6	1500		5	1648
12	950		10	976
30	515		20	600
60	315		50	324
			100	212

Table 11 - Shear Thinning Viscosity of Composition KK

rpm	Viscosity (cp)
0.5	4080
1	3120
2	2240
2.5	2016
4	1570
5	1344
10	820
20	520
50	320
100	218

Conclusions

The shear thinning viscosity of the present compositions is characteristic of a structured composition, such as a microemulsion.

Example 4 - - Antimicrobial Efficacy of the Present Compositions Including Medium Chain Peroxycarboxylic Acid and Solubilizer

Compositions according to the present invention were evaluated and demonstrated advantageous antimicrobial activity against microbes such as gram negative bacteria, gram positive bacteria, fungi, spores, viruses, and mycobacteria.

Materials and Methods

Antimicrobial activity was determined according to two well established methods. The first method was the procedure set out in *Germicidal and Detergent Sanitizing Action of Disinfectants*, Official Methods of Analysis of the Association of Official Analytical Chemists, paragraph 960.09 and applicable sections, 15th Edition, 1990 (EPA Guideline 91-2). The second method was the procedure described in *A.O.A.C. Use Dilution Methods*, Official Methods of Analysis of the Association of Official Analytical Chemists, paragraph 955.14 and applicable sections, 15th Edition, 1990 (EPA Guideline 91-2). Briefly, antimicrobial activity of the present compositions

was determined by exposing a one mL aliquot containing the target microorganism to 99 mL of the desired concentration of the test substance at the desired temperature. After the specified contact time, one mL of the test solution containing the microorganism was neutralized and enumerated for survivors.

5 The hospital disinfectant efficacy of the present compositions was determined by drying the target microorganism on a stainless steel carrier and exposing the carrier to 10 mL of the desired concentration of test composition at the desired temperature for the specified contact time. Then, the carrier was aseptically transferred to a neutralizer/subculture medium.

10 Antiviral activity against Herpes Simplex Virus Type 1 was determined by known procedures. Briefly: Herpes Simplex Virus Type 1 was dried on a glass surface. The film of virus was exposed to test substance for 10 min at room temperature. Then, the mixture of film and test substance were subjected to gel filtration to separate small molecules from virus particles. The recovered virus was
15 assayed for infectivity by an accepted assay method.

 Antiviral activity against Poliovirus Type 1 was determined by known procedures. Briefly: Poliovirus Type 1 was dried on a glass surface. The film of virus was exposed to test substance for 5 min at room temperature. Then, the mixture of film and test substance were subjected to gel filtration to separate small molecules from
20 virus particles. The recovered virus was assayed for infectivity by an accepted assay method.

Results

 Tables 12-21 include data showing that the present medium chain
25 peroxycarboxylic acid compositions had antimicrobial activity when tested against bacteria, fungi, and spores in several different types of tests.

 The data presented in Table 12 demonstrate that the present compositions exhibited significant antimicrobial activity when diluted with a diluent to pH less than 4. Efficacy was not as great if the composition was diluted and then the pH was
30 brought to less than or equal to 4. These results illustrate that present compositions with significant levels of acidulant exhibited, under certain circumstances, advantageous activity.

The data presented in Table 13 demonstrate that the present compositions exhibited significant antimicrobial activity at pH of 2.6 to 3.5. These results indicate that at a pH of 6.1, 11 ppm of peroxyoctanoic acid (POOA) is still effective at reducing *S. aureus* by >7.04 log. The data presented in Table 14 demonstrate that efficacy of this composition was not as great against *E. coli* if it was diluted and then the pH was brought to less than 4.

The data presented in Table 15 demonstrate that the present compositions exhibited significant antimicrobial activity. All formulas tested achieved >5 log reductions of *Escherichia coli* in 30 seconds at 0.069% when diluted in 500 ppm synthetic hard water. Also, these compositions achieved complete kill (>7 log reduction) of *Pseudomonas aeruginosa* in 30 seconds at 0.082% when diluted in 500 ppm synthetic hard water. The combination of higher pH and lower ppm in one composition may have contributed to the lower log reduction.

The data presented in Table 16 demonstrate that the present compositions exhibited significant antimicrobial activity against several fungi and bacteria. The present compositions exhibited broad spectrum antimicrobial activity against bacteria and fungi at low levels of medium chain peroxydicarboxylic acid. These results indicate that composition 106 is more effective than composition DD. Composition BB achieved higher reductions of *A. niger* and *P. aeruginosa* at similar levels of peroxydicarboxylic acid.

The data presented in Table 17 demonstrate that the present compositions exhibited significant antimicrobial activity against several fungi and several bacteria.

The data presented in Table 18 demonstrate that one of the present compositions (KK) exhibited significant antimicrobial activity against *E. coli* O157:H7, *S. typhimurium*, and *L. monocytogenes*. This composition achieved more than 99.999% reduction within a 30 second exposure time.

The data presented in Table 19 demonstrate that the present compositions exhibited significant antimicrobial activity against several bacteria in a hospital disinfectant test. The hospital disinfectant test measures whether the composition killed all of the microbes on a stainless steel carrier. A composition listed as 10/10 killed all of the bacteria on each of 10 carriers. Likewise a result of 60/60 indicates that a composition kills all of the bacteria on each of 60 carriers. These results present a

greater challenge for an antimicrobial agent because it requires activity in the presence of 5% fetal bovine serum. Therefore, it indicates that the present compositions were effective as a hospital disinfectant in the presence of blood soil.

5 The data presented in Table 20 demonstrate that one of the present compositions exhibited superior antimicrobial activity against several bacteria in a hospital disinfectant test compared to a conventional, commercially available antimicrobial agent. The hospital disinfectant test measures whether the composition killed all of the microbes on a particular carrier. The composition according to the present invention, AA-O, passed the hospital disinfectant test, with complete kill on 59 of 60 carriers.
10 The conventional antimicrobial agent (containing hydrogen peroxide as active) did not pass the test. It yielded complete kill on only 58 of 60 carriers. These results indicate that in the presence of fetal bovine serum and when diluted in synthetic hard water the current composition was more effective than the commercially available hospital disinfectant.

15 The data presented in Table 21 demonstrate that the present compositions exhibited significant antimicrobial activity against bacterial spores. Bacterial spores are difficult to kill. These results indicate that at elevated temperatures the effectiveness of the present compositions increased, which provided for effective kill at reduced contact times.

20 The data presented in Table 22 demonstrate that the present compositions exhibited superior antimicrobial activity against bacterial spores compared to conventional peroxide and peroxycarboxylic acid antimicrobials. The present composition resulted in greater kill at equal or lower concentrations of antimicrobial active. These results indicate that the present compositions exhibited superior
25 antimicrobial activity compared to conventional antimicrobials.

The data presented in Table 23 demonstrate that the present compositions exhibited effective antimicrobial activity against *Mycobacterium bovis*. The present composition (B) provided complete kill of M. bovis BCG at dilutions of 1 oz per 4 gal and 1 oz per 6 gal with exposure times as short as 6 min. These results indicate that the
30 compositions of the present invention can be employed as a tuberculocidal agent.

Tests against Herpes Simplex Virus Type 1 resulted in complete kill of this virus. The virus was dried onto a hard surface. The virus on the hard surface was

contacted for 10 min with composition B diluted at 1 oz per 6 gallons or 1 oz per 8 gallons. Both dilutions resulted in complete kill, a greater than 5.3 log reduction in virus. Virus and cells survived in appropriate controls. These results indicate that the present compositions are effective virucides.

5 Tests against Poliovirus Type 1 resulted in nearly complete kill of this virus. The virus was dried onto a hard surface. The virus on the hard surface was contacted for 10 min with composition LL diluted at 1 oz per 1 gallon or 1 oz per 0.5 gallons. The dilution of 1 oz to 1 gallon completely killed the poliovirus at 5 different titers, killed no virus at the highest titer, and resulted in incomplete kill at the second and third
10 highest titers. This dilution exhibited 1.5 log reduction in virus titer. The dilution of 1 oz to 0.5 gallons completely killed poliovirus at all titers tested. This dilution resulted in >4 log reduction in virus titer. Virus and cells survived in appropriate controls. These results indicate that the present compositions are effective general virucides.

The data presented in Table 24 demonstrate that the present compositions
15 exhibited antimicrobial activity superior to that of compositions including synthetic medium chain peroxycarboxylic acid that had been added to a composition. Better efficacy was found in the solutions with the lower pH, which were made up with Milli-Q water. The 60 ppm sample almost achieved a 5 log reduction in 30 seconds. However, this data indicates that the pH of the test solution can be more important than
20 the ppm of active POOA.

The data presented in Table 25 demonstrate that the present compositions exhibited antimicrobial activity superior to that of compositions including synthetic medium chain peroxycarboxylic acid that had been added to a composition. These data further suggest that POOA exhibited greater activity against *Escherichia coli* a pH of
25 ~4.0 and a concentration >5 ppm no matter what diluent is used. Against *Staphylococcus aureus* POOA achieved 5 log reductions at a concentration of 5 ppm and at a pH of ~5. There was no difference between the reductions seen in Milli-Q water and soft water for either organism.

Table 12 - Antimicrobial Activity of Compositions Including Solvent Solubilizer
Against *E. coli* and *S. aureus* with 30 Second Exposure at Room Temperature

Composition	[POOA] (ppm)	Diluent	pH	Log Reduction of <i>E. coli</i>	Log Reduction of <i>S. aureus</i>
F stored at RT for 31 days 0.92% POOA	5	HW - pH 5.0	3.19	2.45	6.10
		HW - pH 7.8	7.74	0.10	3.52
		HW - adjusted to pH 4.0 after dosing	3.98	0.10	5.62
	8	HW - pH 5.0	3.03	7.15	>6.70
		HW - pH 7.8	6.16	0.07	5.62
		HW - adjusted to pH 4.0 after dosing	4.00	0.65	>6.40
	12	HW - pH 5.0	2.86	>7.15	>6.70
		HW - pH 7.8	4.41	0.59	6.70
		HW - adjusted to pH 4.0 after dosing	3.96	2.84	6.40
F stored at 100 °F for 31 days 1.13% POOA	7	HW - pH 5.0	3.19	1.39	5.80
		HW - pH 7.8	6.80	0.15	2.09
		HW - adjusted to pH 4.0 after dosing	3.89	0.15	5.24
	10	HW - pH 5.0	3.01	>6.84	6.70
		HW - pH 7.8	6.14	0.10	5.24
		HW - adjusted to pH 4.0 after dosing	3.89	0.39	5.49
	14	HW - pH 5.0	2.85	>7.15	>6.70
		HW - pH 7.8	4.28	0.28	>6.40
		HW - adjusted to pH 4.0 after dosing	4.07	1.40	6.22

HW = 500 ppm synthetic hard water

Table 13 - Antimicrobial Activity of Compositions Including Solvent Solubilizer
Against *E. coli* and *S. aureus* with a 30 Second Exposures at Room Temperature - Tests
Conducted Using pH Adjusted Synthetic Hard Water

Composition	pH of Diluent	pH of Test Substance	Log Reduction of <i>E. coli</i>	Log Reduction of <i>S. aureus</i>
K (0.086 wt-%) 16 ppm POOA	3.9 - 4.0	2.64	>7.11	>7.04
	4.9 - 5.1	2.74	>7.11	>7.04
	5.9 - 6.1	2.75	>7.11	>7.04
	7.7 - 7.9	3.50	>7.11	>7.04
K (0.057 wt-%) 11 ppm POOA	3.9 - 4.0	2.80	>7.11	>7.04
	4.9 - 5.1	2.83	>7.11	>7.04
	5.9 - 6.1	2.97	>7.11	>7.04
	7.7 - 7.9	6.12	0.21	>7.04

- 5 Table 14 - Antimicrobial Activity of Compositions Including Solvent Solubilizer
Against *E. coli* and *S. aureus* with 30 Second Exposure at Room Temperature - Tests
Conducted With pH is Adjustment After Dosing

Composition	Natural pH	Adjusted pH	Log Reduction of <i>E. coli</i>	Log Reduction of <i>S. aureus</i>
K (0.050 wt-%)	5.09	3.91*	2.84	>6.84
K (0.057 wt-%)	4.92	3.85**	4.61	>6.84

* 2 drops of 1.0 N HCl

** 5 drops of 1.0 N HCl

Table 15 - Antimicrobial Activity of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer Against *Pseudomonas aeruginosa* and *Escherichia coli* with 30 Second Exposure at Room Temperature to a Composition Made with 500 ppm Synthetic Hard Water at pH 7.60

Composition	Use-Solution [POOA] ppm	pH	Log Reduction of <i>E. coli</i>	Log Reduction of <i>P. aeruginosa</i>
T	13	2.9	5.16*	Not Tested
U	13	3.1	>7.28	
V	12	3.0	>7.28	
T	16	2.8	Not tested	>7.15
U	16	2.8		>7.15
V	15	2.9		4.75

5

* = Duplicate plate counts were not consistent

Table 16 - Antimicrobial Activity of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer Against Several Fungi and *Pseudomonas aeruginosa* with a 30 Second Exposure at Room Temperature

Composition	[POOA] (ppm)	Log Kill of <i>S. cerevisiae</i> , (30 sec, RT)	Log Kill of <i>C. albicans</i> , (30 sec, RT)	Log Kill of <i>A. niger</i> , (5 min, RT)	Log Kill of <i>P. aeruginosa</i> (30 sec, RT)
BB	22	>5.6	>6.1	1.6	
	20	5.1	>6.1	1.4	
	18	4.7	>6.1	1.2	>7.2
	17				>7.2
	16				>7.2
	15	4.1	4.2	1.0	>7.2
	14				>7.2
	13				4.7
DD	16			0	5.6
	15			0	3.5
	14			0	1.8
	13			0	0.73

Table 17 - Antimicrobial Activity of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer Against Several Fungi and Several Bacteria with a 30 Second Exposure at Room Temperature

Composition	[POOA] (ppm)	Log Kill of <i>S. cerevisiae</i> , (30 sec, RT)	Log Kill of <i>C. albicans</i> , (30 sec, RT)	Log Kill of <i>A. niger</i> , (5 min, RT)	Log Kill of <i>P.</i> <i>aeruginosa</i> (30 sec, RT)	Log Kill of <i>E. coli</i> O157:H7 (30 sec, RT)	Log Kill of <i>L.</i> <i>monocytogenes</i> (30 sec, RT)	Log Kill of <i>S. aureus</i> (30 sec, RT)
LL	34	>5.6	>6.1	3.0				
	30	>5.6	>6.1	2.3				
	27	>5.6	>6.1	1.7				
	23	4.6	>6.1	1.4	5	>7	>7	
	21					>7	>7	
HH	26	>5.4	>5.8	3.4				
	21	4.2	>5.8	2.2				
	17	4.1	>5.8	1.4	>7.0	>7*	>7.0	6.4
	16				>7.0	>7*	>7.0	4.5

* also killed a less virulent strain of *E. coli*;

Table 18 - Antimicrobial Activity of Composition Including Anionic Surfactant and/or Microemulsion Solubilizer Against Several Bacteria 30 and 60 Second Exposure at Room Temperature

Composition	[POOA] (ppm)	Log Kill of <i>E. coli</i> O157:H7, (30 and 60 sec, RT)	Log Kill of <i>S. typhimurium</i> , (30 and 60 sec, RT)	Log Kill of <i>L. monocytogenes</i> , (30 and 60 sec, RT)
KK	17	>6.9	>7.2	>6.6

5

Table 19 - Antimicrobial Activity of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer Against Several Bacteria in a Hospital Disinfectant Test

Composition	[POOA] (ppm)	<i>P. aeruginosa</i> (kill tubes/total tubes)	<i>S. aureus</i> (methicillin resistant) (kill tubes/total tubes)	<i>E. faecalis</i> (vancomycin resistant) (kill tubes/total tubes)
BB	130	60/60		
	89	59/60		
	59	60/60	10/10	10/10
	44	58/60	10/10	10/10
DD	140	60/60		
	93	60/60		
	62	60/60		
	47	58/60		
LL	91		10/10	10/10
	68		10/10	10/10

Table 20 - Antimicrobial Activity of Composition Including Anionic Surfactant and/or Microemulsion Solubilizer and of Conventional Antimicrobial Composition Against Several Bacteria in a Hospital Disinfectant Test

Composition	[POOA] (ppm)	<i>P. aeruginosa</i> (kill tubes/total tubes)	<i>S. aureus</i> (kill tubes/total tubes)
AA-O (0.98 wt-%)	196	60/60	59/60
Virox 5 (1:16 dilution)	0	58/60	58/60

5 Table 21 - Antimicrobial Activity of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer Against Bacterial Spores

Composition	[POOA] (ppm)	Log Kill of <i>Bacillus cereus</i> spores (30 sec at 40 °C)	Log Kill of <i>Bacillus cereus</i> spores (10 sec at 60 °C)
BB	200	2.1	4.7
	150	0.21	2.0
HH	240	4.2	5.6
	180	0.94	2.6
DD	200	4.5	6.0
	150	0.53	4.1
LL	290	4.7	5.7
	220	0.88	4.3

Table 22 - Antimicrobial Activity of Compositions Including Anionic Surfactant and/or Microemulsion Solubilizer and of Conventional Compositions Against Bacterial Spores

Composition	Concentration of Antimicrobial	pH	Exposure Temperature (°C)	Exposure Time (sec)	Log Reduction
H ₂ O ₂	35%	3.32	40	30	1.19
				60	2.94
				120	>6.30
			60	10	1.59
				20	4.85
				30	4.89
			80	10	>6.30
				20	>6.30
				30	>6.30
KK (2.0 wt-%)	250 ppm POOA 1400 ppm H ₂ O ₂ 520 ppm OA	1.85	40	30	2.33
				60	6.30
				120	>6.30
			60	10	5.30
				20	>6.30
				30	>6.30
			80	10	>6.30
				20	>6.30
				30	>6.30
Conventional Mixed Peroxycarboxylic Acid (1.5 wt-%)	750 ppm peracid 1000 ppm H ₂ O ₂ 555 ppm OA	3.06	40	30	1.02
				60	2.80
				120	4.22
			60	10	3.96
				20	5.22
				30	>6.30
			80	10	>6.30
				20	>6.30
				30	>6.30

Composition	Concentration of Antimicrobial	pH	Exposure Temperature (°C)	Exposure Time (sec)	Log Reduction
Conventional Peroxyacetic Acid (4.5 wt-%)	2610 ppm POAA 1.26% H ₂ O ₂	2.61	40	30	0.30
				60	0.30
				120	0.75
			60	10	0.58
				20	1.85
				30	2.64
			80	10	4.70
				20	>6.30
				30	>6.30

Table 23 - Antimicrobial Activity of Compositions Including Solvent Solubilizer Against Mycobacteria

Composition	[POOA] (ppm)	Exposure Time at Room Temperature (min)	Log Kill of <i>M. bovis</i>
B	39	5	>6.5
		10	>6.5
		15	>6.5
		20	>6.5
B	26	5	6.2
		10	6.2
		15	>6.5
		20	>6.5

Table 24 - Antimicrobial Activity of Compositions Including POOA from Pure Crystals at 60, 40 and 20 ppm in Milli-Q and Synthetic Hard Water

Test Substance	Concentration	Diluent	pH	Log Reduction of E. coli
Pure POOA Crystals	60 ppm	500 ppm Synthetic Hard Water, pH 7.77	7.54	1.12
	40 ppm		7.61	0.93
	20 ppm		7.68	0.62
	60 ppm	Milli-Q water	5.08	4.68
	40 ppm		5.28	2.61
	20 ppm		5.58	0.55

Table 25 - Antimicrobial Activity of Compositions Including of POOA from Pure Crystals in Milli-Q and Soft Water at Differing pH Values Against Two Bacteria with a 30 Second Exposure at Room Temperature

Test Substance	Concentration	Diluent	Post Test pH	Log Reduction of <i>E. coli</i>	Log Reduction of <i>S. aureus</i>
Pure POOA Crystals	5 ppm	Milli-Q water pH 6.60	6.24	0.09	6.04
		Milli-Q water pH 5.98	5.89	0.11	4.44
		Milli-Q water pH 5.00	5.03	0.07	5.01
		Milli-Q water pH 4.04	4.09	1.34	6.28
		Soft water pH 9.29	9.12	0.07	0.1
		Soft water pH 5.91*	6.68	0.08	4.19
		Soft water pH 5.08*	5.79	0.09	5.16
		Soft water pH 3.91	4.01	1.26	5.82
	10 ppm	Milli-Q water pH 6.60	5.80	0.06	>6.82
		Milli-Q water pH 5.98	5.90	0.1	6.52
		Milli-Q water pH 5.00	4.98	0.07	>6.82
		Milli-Q water pH 4.04	4.08	6.04	>6.82
		Soft water pH 9.29	9.09	0.07	0.26
		Soft water pH 5.91	6.68	0.24	>6.82
		Soft water pH 5.08	5.67	0.55	6.12
		Soft water pH 3.91	4.01	6.34	6.28

*Indicates a pH drift of ~0.7 pH units during the 5 hours the test was performed.

Example 5 - - Compositions Including Medium Chain
Peroxycarboxylic Acid and Solubilizer

Table 26 presents additional illustrative examples of the present compositions including medium chain peroxycarboxylic acid and solubilizer. Quantities in the tables
5 are in wt-%.

In each of compositions AB-AQ: the medium chain peroxycarboxylic acid was peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier was water; the oxidizing agent was hydrogen peroxide (supplied from a 35% solution); the stabilizing agent was HEDP (supplied as Dequest 2010 which includes 60 wt-%
10 HEDP); and the acidulant was phosphoric acid (supplied as 75% phosphoric acid). Composition AC included fragrance (1 wt-%), specifically a mint apple fragrance.

The solubilizer was varied among these compositions. In each of compositions AB-AD, AH, AI, AN, the solubilizer was LAS acid. In compositions AE and AJ, the solubilizer was LAS acid plus C8 amine oxide. In composition AF, the solubilizer was
15 LAS acid plus n-octyl amine. In composition AG, the solubilizer was LAS plus C8-dimethyl amine. In composition AK, the solubilizer was LAS acid plus alkylated diphenyl oxide disulfonate (acid form). In composition AL, the solubilizer was alkylated diphenyl oxide disulfonate (acid form). In composition AM, the solubilizer was LAS acid plus alkylated diphenyl oxide disulfonate (acid form) and C8 amine
20 oxide. In composition AO, the solubilizer was sodium laureth sulfate; suitable sodium laureth sulfates tested include those with n=1 and 3. In composition AP, the solubilizer was alkylated diphenyl oxide disulfonate (salt form). In composition AQ, the solubilizer was alkylated diphenyl oxide disulfonate (salt form) plus NAS-FAL.

In each of compositions AR-AW: the carrier was water; the oxidizing agent
25 was hydrogen peroxide (supplied from a 35% solution); the stabilizing agent was HEDP (supplied as Dequest 2010 which includes 60 wt-% HEDP); the acidulant was phosphoric acid (supplied as 75% phosphoric acid), and the solubilizer was LAS acid.

The medium chain peroxycarboxylic acid and medium chain carboxylic acid were varied among these compositions. In composition AR, the medium chain
30 peroxycarboxylic acid was peroxy-nonanoic acid and the medium chain carboxylic acid was nonanoic acid (straight chain nonanoic acid). In compositions AS-AW, the medium chain

Table 26 - Examples of Compositions Including Surfactant Solubilizer (quantities in wt-%)

Ingredient	AB	AC	AD	AE	AF	AG	AH	AI	AJ	AK	AL	AM	AN
Medium Chain Peroxycarboxylic Acid	1.0	1.1	3.1	1.2	1.5	0.9	1.2	1.1	nd	0.9	0.9	nd	0.9
Medium Chain Carboxylic Acid	2.8	2.7	2.0	2.6	2.3	2.9	2.6	2.7	<3.8	2.9	2.9	<3.8	2.6
Solubilizer	7.8	9.7	11	8.2	7.9	7.9	7	6.5	8-12	5.7	6.3	8.6	7.8
Carrier	52	51	34	52	52	52	53	53	48-52	54	54	52	52
Oxidizing Agent	8.0	8.1	11	8.1	8.2	8.1	8.0	8.1	8	8.1	8.1	8	7.9
Acidulant	27	27	36	27	27	27	27	27	27	27	27	27	27
Stabilizing Agent	2.0	2.0	2.7	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0

5 Table 26, continued - Examples of Compositions Including Surfactant Solubilizer

Ingredient	AO	AP	AQ	AR	AS	AT	AU	AV	AW	AX	AY	AZ	BC
Medium Chain Peroxycarboxylic Acid	1.0	0.9	0.9	1.0	nd	nd	1.0	1.0	nd	nd	nd	0.7	0.7
Medium Chain Carboxylic Acid	2.8	2.9	2.9	2.8	<4.3	<4.8	2.9	3.0	<3.8	<3.8	<3.8	3.1	3.1
Solubilizer	8-9	4.5	4.3	7.8	7.8	7.8	7.8	7.8	8	8.3	8.6	7.4	7.8
Carrier	52	56	56	52	52	52	52	52	52	52	52	53	52
Oxidizing Agent	8.1	8.2	8.2	8.0	8	8	8.2	8.2	8	8	8	8.2	8.2
Acidulant	27	27	27	27	27	27	27	27	27	27	27	27	27
Stabilizing Agent	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0

Table 26, continued - Examples of Compositions Including Surfactant Solubilizer

Ingredient	BD	BE	BF	BG	BH	BI	BJ	BK
Medium Chain Peroxycarboxylic Acid	1.0	1.0	1.0	0.9	0.9	1.0	1.0	1.1
Medium Chain Carboxylic Acid	2.8	2.8	2.9	2.9	2.9	2.8	2.8	2.7
Solubilizer	12	10	9	10	13	15	14	16
Carrier	48	50	51	50	47	45	46	44
Oxidizing Agent	7.8	8.2	7.6	8.3	8.3	8.3	8.2	8.1
Acidulant	27	27	27	14	14	14	14	14
Stabilizing Agent	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0

peroxycarboxylic acid was peroxyoctanoic acid and peroxyonanoic acid and the medium chain carboxylic acid was octanoic acid and nonanoic acid; nonanoic acid (as
5 isononanoic acid (which is believed to be a 6 carbon main chain with three pendant methyl groups)) was present at 0.5, 1, 0.1, 0.2, and 0.3 wt-% for AS-AW, respectively.

In each of compositions AX-AZ and BC-BF: the medium chain peroxycarboxylic acid was peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier was water; the oxidizing agent was hydrogen peroxide
10 (supplied from a 35% solution); the stabilizing agent was HEDP (supplied as Dequest 2010 which includes 60 wt-% HEDP); and the acidulant was phosphoric acid (supplied as 75% phosphoric acid).

The solubilizer was varied among these compositions. In composition AX, the solubilizer was LAS acid plus sodium lauryl sulfate. In composition AY, the
15 solubilizer was LAS acid plus sodium lauryl sulfate and C8 dimethyl amine. In compositions AZ and BC-BF, the solubilizer was secondary alkane sulfonate (a mixture of sulfonated paraffins sold under the tradename Hostapur SAS).

In each of compositions BG-BK: the medium chain peroxycarboxylic acid was peroxyoctanoic acid; the medium chain carboxylic acid was octanoic acid; the carrier
20 was water; the oxidizing agent was hydrogen peroxide (supplied from a 35% solution); the stabilizing agent was HEDP (supplied as Dequest 2010 which includes 60 wt-% HEDP); the solubilizer was secondary alkane sulfonate (a mixture of sulfonated

paraffins sold under the tradename Hostapur SAS) plus NAS-FAL; and the acidulant was sulfuric acid.

The compositions that included LAS, secondary alkane sulfonate, alkylated diphenyl oxide disulfonate, or sodium lauryl sulfate as solubilizer were foaming
5 compositions. Specifically, compositions AB and AC are foaming compositions

Most of the compositions were phase stable. In particular: Compositions AX and AY were determined to be phase stable at 60 °C. For example, several of the compositions for which the wt-% of medium chain peroxycarboxylic acid was not determined (nd) were not phase stable. That is, they separated into more than one
10 phase after a predetermined time at one or more (e.g., at least one) of 40 °F, room temperature, 100 °F, or 140 °F (60 °C).

The concentrations of peroxyoctanoic acid reported in the present examples were determined by a well established and standardized titration protocol. First, hydrogen peroxide content was determined by an oxidation-reduction titration with
15 potassium permanganate. After the endpoint of this titration was reached, an excess of potassium iodide was added to the solution. The potassium iodide reacts with peroxycarboxylic acids to liberate iodine. The liberated iodine was titrated with a standard solution of sodium thiosulfate to yield the concentration of peroxycarboxylic acid. The remaining level of carboxylic acid can be (and was) calculated.

20 The peroxycarboxylic acid was titrated at a time after formulation that was practical in the laboratory. For example, the peroxycarboxylic acid was titrated for compositions AB, AD, AE, AF, AG, AH, AK, AL, AO, AP, AQ, AU, AV, AZ, BC, and BD after the sample had sat at room temperature for 0, 2 (BD), or 3 (AP, AU, and AV) days. For example, the peroxycarboxylic acid was titrated for compositions AC
25 and BG-BK after the sample had sat at 100 °F for 4 days (AC) or 7 days (BG-BK). For example, the peroxycarboxylic acid was titrated for compositions AI, AN, AR, BE and BF after the sample had sat at 140 °F (60 °C) for 1 day (AI, AR, and BE) or 4 days (AN and BF).

For composition AB, no decomposition of peroxycarboxylic acid was observed
30 upon aging the composition for 7 days at 140 °F (60 °C). For composition AC, no decomposition of peroxycarboxylic acid was observed upon aging the composition for

34 days at 100 °F. Other compositions were also observed to include stable peroxydicarboxylic acid.

The octanoic acid employed in the present examples was obtained from sources including Procter & Gamble Chemicals and includes a minimum of 95% octanoic acid with minor amounts of hexanoic acid (ca. 2%), decanoic acid (ca. 2%), and dodecanoic acid (<0.5%).

Fragrance

Certain of the compositions were evaluated for phase stability and for smell after addition of a fragrance. In particular, compositions AB and AG were evaluated. 10
Fragrances evaluated included Green Meadow (Klabi); Vinegar Mask I (J&E Sozio); Vinegar Mask II (J&E Sozio); amyl acetate; iso-bornyl acetate; and methyl salicylate.

Composition AC included fragrance (1 wt-%), specifically a mint apple fragrance which is believed to be or include an alkyl salicylate. Composition AC 15
altered to include 10 wt-% LAS remained single phase at 40 °F, room temperature, and 70 °F.

Foaming

The results in Table 27 show that the present medium chain peroxydicarboxylic acid composition produced foam with desirable qualities. This study employed a 20
“FOAM IT” brand tank foamer set to produce slightly wet foam, 2 turns from the mid point. The foam was dispensed from use composition at 95-98 °F. The foam was sprayed on a vertical stainless steel surface (approximately 15 ft by 15 ft) from a distance of about 10 ft. The results of Table 27 demonstrate that the present 25
compositions provided foam with desirable hang time and density. Each of the compositions tested at 1 oz/ 6 gal. provided foam with desirable characteristics, such as the breaking foam was visible for about 5 min, the foam drained well from the vertical surface, exhibited good sheeting down vertical surface, and dried evenly to no visible residue.

30

**Example 6 - - Antimicrobial Efficacy of the Present Compositions Including
Medium Chain Peroxycarboxylic Acid and Solubilizer**

Additional compositions according to the present invention were evaluated and demonstrated advantageous antimicrobial activity against microbes such as gram
5 negative bacteria, gram positive bacteria, fungi, spores, viruses, and mycobacteria.

Table 27 - Foaming by the Present Medium Chain Peroxycarboxylic Acid
Compositions.

Composition	Amount in Use Solution (oz/gal)	Break Time	Foam Dry Time (min)	Odor	Initial Appearance of Foam	Comments
AB	0.17	slow, about 2 min	>10	moderate	Covers well, wet, about 1/16 inch thick	foam breaks to spotty foam, dries to no visible residue
AG	0.17	slow, about 2 min	>10	moderate	Covers well, wet, about 1/16 inch thick	foam breaks to spotty foam, dries to no visible residue
AH	0.17	faster, < 2 min	95% dry at 10 min	moderate	Covers well, wetter than above	foam breaks to spotty foam, dries to no visible residue
AK	0.17	fast, about 1 min	95% dry at 10 min	moderate	Wetter than above	no visible residue
AY	0.17	fast, about 10 sec	95% dry at 10 min	strong	Very wet, lays flat	no visible residue
AB	0.13	fast, < 1 min	about 10 min	low	Covers, wet	spotty foam
AG	0.13	fast, < 1 min	about 10 min	low	Covers, wet	streaky foam
AH	0.13	very fast, < 1 min	about 10 min	low	Extremely wet	very spotty foam
AK	0.13	very fast, < 1 min	about 10 min	low	Extremely wet	very spotty foam
AY	0.13	fast, about 10 sec	95% dry at 10 min	strong	Very wet, lays flat	no visible residue

Materials and Methods

Antimicrobial activity was determined as described above in Example 4.

Results

5 Tables 28-29 include data showing that the present medium chain
peroxycarboxylic acid compositions had antimicrobial activity when tested against
bacteria, fungi, and spores in several different types of tests.

 The data presented in Table 28 demonstrate that the present compositions
exhibited significant antimicrobial activity. Test 1 included 5 min exposure of the
10 microbe to composition AB at room temperature. The microbes in test 1 included *E.*
aerogenes ATCC 13048 and *S. aureus* ATCC 6538. Test 2 included 30 sec exposure
of the microbe to composition AB at room temperature. The microbes in test 2
included *S. aureus* ATCC 6538, *E. coli* ATCC 11229, and *P. aeruginosa* ATCC 13442.

 The data presented in Table 29 demonstrate sporicidal activity of a composition
15 according to the present invention.

 Tests against Poliovirus Type 1 resulted in complete kill of this virus. The virus
was dried onto a hard surface. The virus on the hard surface was contacted for 10 min
with composition AG diluted at 1 oz per 1 gallon or 1 oz per 0.5 gallons. Composition
AG demonstrated complete inactivation of Poliovirus type 1 following either 3 min or 5
20 min exposure at 20 °C. The composition produced >6 and >5.3 log reduction in 3 and 5
min, respectively. Virus and cells survived in appropriate controls. These results
indicate that the present compositions are effective general virucides.

 The compositions that included fragrance showed no negative effect on
antimicrobial efficacy from the fragrance. Several additional compositions were tested
25 for antimicrobial activity and exhibited results similar to those reported in this
Example.

Table 28 - Activity of Composition AB against Several Microorganisms

Test	Dilution	Log Reduction of <i>E. aerogenes</i>	Log Reduction of <i>S. aureus</i>	Log Reduction of <i>E. coli</i>	Log Reduction of <i>P. aeruginosa</i>
1	500 ppm in synthetic hard water	4.5	5.4		
2	1 oz/9 gal water		>6.7	>7.3	5.8
2	1 oz/9.5 gal water		>6.7	>7.3	5.7
2	1 oz/10 gal water		>6.7	>7.3	5.2
2	1 oz/10.5 gal water		>6.7	>7.3	1.7

Table 29 - Activity of Composition KK against Spores of *B. subtilis* ATCC 49760

Composition	Dilution	Exposure Time (min)	Log Reduction of <i>B. subtilis</i> spores
KK plus 8 wt-% NAS FAL	1 oz/6 gal	30	0.5
		60	0.6
		120	0.6
KK plus 10 wt-% LAS	1 oz/6 gal	30	0.8
		60	1.5
		120	3.0

It should be noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to a composition containing "a compound" includes a mixture of two or more compounds. It should also be noted that
5 the term "or" is generally employed in its sense including "and/or" unless the content clearly dictates otherwise.

All publications and patent applications in this specification are indicative of the level of ordinary skill in the art to which this invention pertains.

The invention has been described with reference to various specific and
10 preferred embodiments and techniques. However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention.

WE CLAIM:

1. A method of reducing a microbial population on poultry during processing comprising:

5 applying to the poultry during processing a medium chain peroxycarboxylic acid antimicrobial composition in an amount and time sufficient to reduce the microbial population;

the medium chain peroxycarboxylic acid antimicrobial composition comprising:

about 2 to about 500 ppm peroxyoctanoic acid;
10 about 5 to about 2000 ppm octanoic acid;
about 95 to about 99.99 wt-% water; and
about 2 to about 16,000 ppm at least one of polyalkylene oxide, monoalkyl ether of polyalkylene oxide, dialkyl ether of polyalkylene oxide, nonionic surfactant, and anionic surfactant;

15 the composition comprising at least about 2 parts by weight of peroxyoctanoic acid for each 7 parts by weight of octanoic acid.

2. A method of recycling water previously applied to poultry, the method comprising:

20 recovering a medium chain peroxycarboxylic acid antimicrobial composition previously applied to poultry; and

adding to the recovered composition a sufficient amount of a medium chain peroxycarboxylic acid composition to yield a recycled medium chain peroxycarboxylic acid antimicrobial composition;

25 the added medium chain peroxycarboxylic acid composition comprising:

about 0.0005 to about 5 wt-% peroxyoctanoic acid;
about 0.001 to about 10 wt-% octanoic acid;
about 5 to about 99.99 wt-% water;
about 0.001 to about 60 wt-% at least one of polyalkylene oxide,
30 monoalkyl ether of polyalkylene oxide, dialkyl ether of polyalkylene oxide, nonionic surfactant, and anionic surfactant;
about 0.002 to about 10 wt-% oxidizing agent;

about 0.001 to about 30 wt-% inorganic acid; and
about 0.001 to about 5 wt-% sequestrant;
the composition comprising at least about 2 parts by weight of
peroxyoctanoic acid for each 7 parts by weight of octanoic acid.

5

3. A method of recycling water previously applied to poultry, the method comprising:

recovering a medium chain peroxycarboxylic acid antimicrobial composition previously applied to poultry; and

10 adding to the recovered composition a sufficient amount of a medium chain peroxycarboxylic acid composition to yield a recycled medium chain peroxycarboxylic acid antimicrobial composition;

the added medium chain peroxycarboxylic acid composition comprising:

about 0.5 to about 5 wt-% peroxyoctanoic acid;

15 about 1 to about 10 wt-% octanoic acid;

about 5 to about 97 wt-% water;

about 1 to about 20 wt-% anionic surfactant;

about 5 to about 10 wt-% oxidizing agent;

about 15 to about 35 wt-% inorganic acid; and

20 about 1 to about 5 wt-% sequestrant;

the composition comprising a microemulsion.

4. A method of recycling water previously applied to poultry, the method comprising:

25 recovering a medium chain peroxycarboxylic acid antimicrobial composition previously applied to poultry; and

adding to the recovered composition a sufficient amount of a medium chain peroxycarboxylic acid composition to yield a recycled medium chain peroxycarboxylic acid antimicrobial composition;

30 the added medium chain peroxycarboxylic acid composition comprising:

about 0.0005 to about 5 wt-% peroxyoctanoic acid;

about 0.001 to about 10 wt-% octanoic acid;

about 40 to about 99.99 wt-% water;

about 0.001 to about 60 wt-% at least one of polyalkylene oxide,
monoalkyl ether of polyalkylene oxide, dialkyl ether of polyalkylene oxide,
anionic surfactant, and nonionic surfactant;

5 about 0.002 to about 10 wt-% oxidizing agent;
 about 0.001 to about 30 wt-% inorganic acid; and
 about 0.001 to about 5 wt-% sequestrant.

5. An antimicrobial concentrate composition comprising:

10 a medium chain peroxycarboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;

the composition comprising:

about 0.0005 to about 5 wt-% peroxyoctanoic acid;

about 0.001 to about 10 wt-% octanoic acid;

15 about 5 to about 99.99 wt-% water;

about 0.001 to about 60 wt-% at least one of polyalkylene oxide, monoalkyl
ether of polyalkylene oxide, dialkyl ether of polyalkylene oxide, nonionic surfactant,
and anionic surfactant;

about 0.002 to about 10 wt-% oxidizing agent;

20 about 0.001 to about 30 wt-% inorganic acid; and

about 0.001 to about 5 wt-% sequestrant;

the composition comprising at least about 2 parts by weight of peroxyoctanoic
acid for each 7 parts by weight of octanoic acid.

25 6. An antimicrobial concentrate composition comprising:

a medium chain peroxycarboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;

the composition comprising:

about 0.5 to about 5 wt-% peroxyoctanoic acid;

30 about 1 to about 10 wt-% octanoic acid;

about 5 to about 97 wt-% water;

about 1 to about 20 wt-% anionic surfactant;

about 5 to about 10 wt-% oxidizing agent;
about 15 to about 35 wt-% inorganic acid; and
about 1 to about 5 wt-% sequestrant;
the composition comprising a microemulsion.

5

7. An antimicrobial concentrate composition comprising:
a medium chain peroxycarboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;

the composition comprising:

10

about 0.0005 to about 5 wt-% peroxyoctanoic acid;

about 0.001 to about 10 wt-% octanoic acid;

about 40 to about 99.99 wt-% water;

about 0.001 to about 60 wt-% at least one of polyalkylene oxide, monoalkyl
ether of polyalkylene oxide, dialkyl ether of polyalkylene oxide, anionic surfactant, and
nonionic surfactant;

15

about 0.002 to about 10 wt-% oxidizing agent;

about 0.001 to about 30 wt-% inorganic acid; and

about 0.001 to about 5 wt-% sequestrant.

20

8. A method of reducing a microbial population on poultry during
processing comprising:

applying to the poultry during processing a medium chain peroxycarboxylic
acid antimicrobial composition in an amount and time sufficient to reduce the microbial
population.

25

9. The method of claim 8, wherein the poultry being processed comprises
chicken, turkey, ostrich, game hen, squab, guinea fowl, pheasant, duck, goose, emu, or
a combination thereof.

30

10. The method of claim 8, comprising applying the medium chain
peroxycarboxylic acid composition by submersing the poultry.

11. The method of claim 10, comprising applying the medium chain peroxycarboxylic acid composition by submersion scalding, by submersion chilling, by hydro-cooling or chilling, tumble immersion, or by a combination thereof.

5 12. The method of claim 10, comprising applying the medium chain peroxycarboxylic acid composition for a duration and at a concentration selected to yield visually imperceptible darkening of subcutaneous bruises, pooled blood, or a combination thereof.

10 13. The method of claim 8, comprising applying the medium chain peroxycarboxylic acid composition by rinsing or spraying the poultry.

14. The method of claim 13, comprising applying the medium chain peroxycarboxylic acid composition with a de-feathering picker, by inside-outside bird
15 washing, by dress rinsing, by spray rinsing, or a combination thereof.

15. The method of claim 8, comprising applying the medium chain peroxycarboxylic acid composition to a whole poultry carcass.

20 16. The method of claim 15, comprising applying the medium chain peroxycarboxylic acid composition to a poultry carcass that has been subjected to stunning, bleeding, scalding, picking, singeing, or a combination thereof.

17. The method of claim 8, comprising applying the medium chain
25 peroxycarboxylic acid composition to one or more dismembered parts of a poultry carcass.

18. The method of claim 17, comprising applying the medium chain peroxycarboxylic acid composition to a poultry carcass that has been subjected to
30 beheading, removing feet, eviscerating, neck-cropping, portioning, or a combination thereof.

19. The method of claim 18, comprising applying the medium chain peroxycarboxylic acid composition to a poultry leg, thigh, breast quarter, wing, or combination thereof of a poultry that has been subjected to portioning.

5 20. The method of claim 17, comprising applying the medium chain peroxycarboxylic acid composition to a poultry that has also been subjected to boning.

21. The method of claim 20, comprising applying the medium chain peroxycarboxylic acid composition to a boned poultry leg, thigh, breast, wing, or
10 combination thereof.

22. The method of claim 8, comprising applying the medium chain peroxycarboxylic acid composition by air chilling.

15 23. The method of claim 22, wherein the medium chain peroxycarboxylic acid composition comprises peroxyoctanoic acid.

24. The method of claim 22, wherein air chilling comprises applying a gaseous or densified fluid antimicrobial composition.

20 25 The method of claim 8, further comprising exposing the poultry to activated light.

26. The method of claim 25, wherein the activated light comprises
25 ultraviolet light, infrared light, visible light, or a combination thereof.

27. The method of claim 8, wherein the medium chain peroxycarboxylic acid antimicrobial composition comprises:
about 2 to about 500 ppm medium chain peroxycarboxylic acid;
30 about 5 to about 2000 ppm medium chain carboxylic acid;
about 95 to about 99.99 wt-% water; and
about 2 to about 16,000 ppm solubilizer.

28. The method of claim 27, wherein the medium chain peroxycarboxylic acid antimicrobial composition further comprises stabilizing agent, wetting agent, thickener, foaming agent, acidulant, pigment, dye, or a combination thereof.

5

29. The method of claim 8, wherein the microbial population is the result of contamination by fecal matter or digestive tract content.

30. The method of claim 29, wherein the microbial population is reduced in
10 a continuous online process.

31. The method of claim 8, further comprising, after applying:
recovering the applied medium chain peroxycarboxylic acid antimicrobial
composition; and

15 adding to the recovered composition a sufficient amount of a medium chain
peroxycarboxylic acid to yield a recycled medium chain peroxycarboxylic acid
antimicrobial composition.

32. The method of claim 31, further comprising applying the recycled
20 composition to poultry during processing.

33. The method of claim 31, wherein the medium chain peroxycarboxylic
acid comprises:

25 about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid;
about 1 to about 10 wt-% medium chain carboxylic acid;
about 5 to about 97 wt-% water; and
about 1 to about 20 wt-% microemulsion former;
the composition comprising a microemulsion.

30 34. The method of claim 31, wherein the medium chain peroxycarboxylic
acid comprises:

about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid;

about 0.001 to about 10 wt-% medium chain carboxylic acid;
about 0 to about 99.99 wt-% water; and
about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium
chain peroxydicarboxylic acid and the medium chain carboxylic acid;
5 the composition comprising about 2 or more parts by weight of medium chain
peroxydicarboxylic acid for each 7 parts by weight of medium chain carboxylic acid.

35. The method of claim 31, wherein the medium chain peroxydicarboxylic
acid comprises:
10 about 0.0005 to about 5 wt-% medium chain peroxydicarboxylic acid;
about 0.001 to about 10 wt-% medium chain carboxylic acid;
about 40 to about 99.99 wt-% water; and
about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium
chain peroxydicarboxylic acid and the medium chain carboxylic acid.

15 36. The method of claim 31, wherein the recycled medium chain
peroxydicarboxylic acid antimicrobial composition comprises:
about 2 to about 500 ppm medium chain peroxydicarboxylic acid;
about 5 to about 2000 ppm medium chain carboxylic acid;
20 about 95 to about 99.99 wt-% water; and
about 2 to about 16,000 ppm solubilizer.

37. A method of recycling water previously applied to poultry, the method
comprising:
25 recovering a medium chain peroxydicarboxylic acid antimicrobial composition
previously applied to poultry; and
adding to the recovered composition a sufficient amount of a medium chain
peroxydicarboxylic acid to yield a recycled medium chain peroxydicarboxylic acid
antimicrobial composition.

30 38. The method of claim 37, further comprising applying the recycled
composition to poultry during processing.

39. The method of claim 37, wherein the medium chain peroxycarboxylic acid composition comprises:

- about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid;
5 about 0.001 to about 10 wt-% medium chain carboxylic acid;
about 0 to about 99.99 wt-% water; and
about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium chain peroxycarboxylic acid and the medium chain carboxylic acid;
the composition comprising about 2 or more parts by weight of medium chain
10 peroxycarboxylic acid for each 7 parts by weight of medium chain carboxylic acid.

40. The method of claim 37, wherein the medium chain peroxycarboxylic acid composition comprises:

- about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid;
15 about 1 to about 10 wt-% medium chain carboxylic acid;
about 5 to about 97 wt-% water; and
about 1 to about 20 wt-% microemulsion former;
the composition comprising a microemulsion.

20 41. The method of claim 37, wherein the medium chain peroxycarboxylic acid composition comprises:

- about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid;
about 0.001 to about 10 wt-% medium chain carboxylic acid;
about 40 to about 99.99 wt-% water; and
25 about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium chain peroxycarboxylic acid and the medium chain carboxylic acid.

42. The method of claim 37, wherein the composition was previously applied by a carcass wash or rinse.

30

43. The method of claim 37, wherein the composition was previously applied by an inside-outside bird wash.

44. An antimicrobial concentrate composition comprising:
a medium chain peroxycarboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;
5 the composition comprising:
about 0.5 to about 5 wt-% medium chain peroxycarboxylic acid;
about 1 to about 10 wt-% medium chain carboxylic acid;
about 5 to about 97 wt-% water; and
about 1 to about 20 wt-% microemulsion former;
10 the composition comprising a microemulsion.
45. An antimicrobial concentrate composition comprising:
a medium chain peroxycarboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;
15 the composition comprising:
about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid;
about 0.001 to about 10 wt-% medium chain carboxylic acid;
about 0 to about 99.99 wt-% water; and
about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium
20 chain peroxycarboxylic acid and the medium chain carboxylic acid;
the composition comprising about 2 or more parts by weight of medium chain
peroxycarboxylic acid for each 7 parts by weight of medium chain carboxylic acid.
46. An antimicrobial concentrate composition comprising:
25 a medium chain peroxycarboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;
the composition comprising:
about 0.0005 to about 5 wt-% medium chain peroxycarboxylic acid;
about 0.001 to about 10 wt-% medium chain carboxylic acid;
30 about 40 to about 99.99 wt-% water; and
about 0.001 to about 80 wt-% solubilizer effective for solubilizing the medium
chain peroxycarboxylic acid and the medium chain carboxylic acid.

47. An antimicrobial use composition comprising:
a medium chain peroxy-carboxylic acid composition effective for reducing the
microbial burden on a surface of poultry;

5 the composition comprising:

about 2 to about 500 ppm medium chain peroxy-carboxylic acid;

about 5 to about 2000 ppm medium chain carboxylic acid;

about 95 to about 99.99 wt-% water; and

about 2 to about 16,000 ppm solubilizer.

10

48. A method of reducing a microbial population on poultry processing
surface, the method comprising:

applying to the poultry processing surface medium chain peroxy-carboxylic acid
antimicrobial composition in amount and for time sufficient to reduce the microbial

15 population.