



US 20130121983A1

(19) **United States**(12) **Patent Application Publication**
JONES et al.(10) **Pub. No.: US 2013/0121983 A1**(43) **Pub. Date: May 16, 2013**(54) **MODIFIED FORMS OF PSEUDOMONAS
EXOTOXIN A**(75) Inventors: **Timothy David JONES**, Babraham
(GB); **Francis Joseph Carr**, Balmedie
(GB)(73) Assignee: **Intrexon Corporation**, Blacksburg, VA
(US)(21) Appl. No.: **13/604,173**(22) Filed: **Sep. 5, 2012****Related U.S. Application Data**(60) Provisional application No. 61/531,576, filed on Sep.
6, 2011.**Publication Classification**(51) **Int. Cl.**
C12N 9/10 (2006.01)(52) **U.S. Cl.**CPC **C12N 9/1077** (2013.01)USPC **424/94.5**; 435/193; 435/188; 536/23.2;
435/320.1; 435/252.33; 530/350; 435/15(57) **ABSTRACT**

Pseudomonas exotoxin A or "PE" is a 66 kD, highly potent, cytotoxic protein secreted by the bacterium *Pseudomonas aeruginosa*. Various forms of PE have been coupled to other proteins, such as antibodies, to generate therapeutically useful cytotoxin conjugates that selectively target cells of a desired phenotype (such as tumor cells). In the present invention, peptides spanning the sequence of an approximately 38 kD form of *Pseudomonas* exotoxin A protein were analyzed for the presence of immunogenic CD4+ T cell epitopes. Six immunogenic T cell epitopes were identified. Residues were identified within each epitope for introduction of targeted amino acid substitutions to reduce or prevent immunogenic T-cell responses in PE molecules which may be administered to a heterologous host.

1 AEEAFDLWNE CAKACVLDLK DQVRSSRMSV DPAIADTNGQ GVLHYSMVLE GCNDALKLAI
61 DNALSITS DG LTIIRLEGGVE PNKPVRYST ROARGSWSLN WLVPICHEKP SNIKVFIHEL
121 NAGNQLSHMS PIYTIEMGDE LLAKLARDAT FFVRAHESNE MQPTLAISHA GVSVMMAQTO
181 PRREKRWSEW ASGKVLCLLD PLDGVVNYLA QQRNLDDTW EGKIYRVLAG NPAKHDLDIK
241 PTVISHRLHF PE GGSALAALT AHQACHLPLE TFTRHRQPRG WEQLEQCGYP VQRLVALYLA
301 ARLSWNQVDQ VIRNALASPG SGGDLGEAIR EQPEQARLAL TLAAAESERF VRQGTGNDIA
361 GAAN advvsl tcpvaageca GPADSGDALL ERNYPTGAE F LGDGGDVSEF TRGTYNWIVE
421 RLICAHRQLE ERGYVEVGYH GTFLEAAQSI VEGGVARSQ DLDATWGEFY IAGDPALAYG
481 YACDQEPDAR GRIRNGALLR VYVPRSSLPG FYRTSLTLAA PEAAGEVERL IGHPLPLRID
541 AITGPEEEGG RLETILGWPL AERTVVIPSA IPTDPRNVGG DLDPSSIPDK EQAISALEDY
601 ASQPGKPPRE DLK - 613 (SEQ ID NO:133)

.....
Alternative carboxy-terminal tails:

609 REDLK - 613 (SEQ ID NO: 135)
609 REDL - 612 (SEQ ID NO: 136)
609 KDEL - 612 (SEQ ID NO: 137)

.....
Amino Acids (AA):

1~252 = Domain IA (cell binding domain; underlined)
253-364 = Domain II (cytosolic translocation; *italics*)
365-399 = Domain IB (dashed underling; SEQ ID NO:139)
365-380 = optional deletion of 365- ADVVSLTCPVAAGECA -380
(SEQ ID NO:138) (lowercase letters; dashed underling)
400-613 = Domain III (cytotoxic portion; **bold, double-underline**)
609-613 or 609-612 = Alternative carboxy-terminal tails

FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6





FIG. 7






FIG. 9




FIG. 10




FIG. 11

MODIFIED FORMS OF PSEUDOMONAS EXOTOXIN A

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Appl. No. 61/531,576, filed Sep. 6, 2011 which is herein incorporated by reference in its entirety.

REFERENCE TO RELATED APPLICATION

[0002] This application claims benefit of and priority based on U.S. Provisional Patent Application Ser. No. 61/531,576, filed Sep. 6, 2011, the contents of which are herein incorporated by reference in their entirety.

NAMES OF THE PARTIES IN A JOINT RESEARCH AGREEMENT

[0003] The claimed invention was made pursuant to a joint research agreement, as defined in 35 U.S.C. §103 (c)(3), that was in effect on or before the date the claimed invention was made, and as a result of activities undertaken within the scope of the joint research agreement, by or on behalf of the Intraxon Corp. (Foster City, Calif., U.S.A.) and Antitope Ltd. (Cambridge, UK).

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

[0004] A Sequence Listing is submitted electronically via EFS-Web as an ASCII formatted sequence listing in a file named "OT050-PCT_SEQLIST.txt", created on Sep. 4, 2012, and having a file size of 295,678 bytes which is filed concurrently with the present specification, claims, abstract and figures provided herewith. The sequence listing contained in this ASCII formatted document is part of the specification and is herein incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

Immune System and T Cell Epitopes

[0005] Immune responses to biological therapeutic agents are wide ranging, and can be

[0006] directed against agents that are both non-human and human in origin. These responses include those that elicit a weak clinical effect and those that limit efficacy which can occasionally result in morbidity or even mortality in patients. In particular, serious complications can arise with the production of neutralizing antibodies, especially when they target recombinant self proteins and therefore have the potential to cross react with the patient's own endogenous protein (Lim, 2005). Problems associated with immunogenicity to biologics (i.e., therapeutic medical products; such as, antibodies and recombinant proteins/polypeptides) have been reduced largely due to advances in molecular biology. There are, however, many recombinant protein biologics that are identical to endogenously expressed human sequences that still elicit potent neutralizing immune responses in patients (Hochuli, 1997; Schellekens et al, 1997; Namaka et al, 2006). The mechanism by which immunogenicity is triggered remains unclear although the tolerance to self proteins may be broken by a number of factors linked to both the product and the patient, (reviewed in Chester et al, 2006; Baker and Jones, 2007). For the product, these include dose, frequency of

administration, route, immunomodulatory capacity of the protein therapeutic, and the formulation (Jaber and Baker, 2007). For the patient, factors such as immune competence (i.e. whether the patient is receiving immunosuppressive treatment), patient's MHC haplotype and intrinsic tolerance to the protein therapeutic will influence immunogenicity. Regardless of how immunogenicity is triggered, one of the single most important factors in the development of an ensuing immune response is the presence of epitopes that are able to effectively stimulate a potent CD4+ T cell response (reviewed Baker and Jones, 2007).

[0007] T cells or T lymphocytes are a subset of white blood cells known as lymphocytes. (The abbreviation "T" in T cell is for "thymus" since this is the primary organ responsible for T cell maturation.) T cells play a central role in cell-mediated immunity. They can be distinguished from other types of lymphocytes (such as B cells and natural killer cells (NK cells)), by the presence of cell-surface proteins called T cell receptors (TCRs). Different types of T cells have also been identified; these can be distinguished based on the differing functions they serve (e.g., CD4+ T cells (a.k.a., T_H or T helper cells), CD8+ cytotoxic T cells (CTLs), memory T cells, regulatory T cells (T_{reg} cells), natural killer cells (NK cells), and gamma delta T cells (γδ T cells)).

[0008] T helper (T_H) cells are so named because they aid other white blood cells in immunologic processes including, *inter alia*, assisting the maturation of B cells into plasma and B memory cells, and activation of cytotoxic T cells and macrophages. T_H cells are also known as CD4+ T cells because they express CD4 protein on the cell-surface. CD4+ T cells are activated when peptide antigens are presented by MHC class II molecules expressed on the surface of Antigen Presenting Cells (APCs). Once activated, CD4+ T cells divide rapidly and secrete chemokines that further assist in activating or regulating immune responses.

[0009] T cell epitope analysis is becoming increasingly important particularly in the pre-clinical analysis of biologics and may, in time, become a requirement for regulatory approval for clinical trials. To this end, a pre-clinical *ex vivo* T cell assay (EPISCREEN™) has been used to provide an effective technology for predicting T cell immunogenicity by identifying linear T cell epitopes present in protein sequences. Synthetic overlapping peptides typically of about 15 amino acids in length are tested against a cohort of community blood donors carefully selected based on MHC class II haplotypes to provide a quantitative analysis of T cell epitopes present in protein sequences. This technology has been used successfully to compare protein variants for the potential to induce an immune response *in vivo*. By providing a high degree of sensitivity along with high reproducibility, the EPISCREEN™ assay allows an accurate pre-clinical assessment of the potential for immunogenicity of biologics. See, Baker & Carr, "Preclinical Considerations in the Assessment of Immunogenicity for Protein Therapeutics," *Current Drug Safety*; 5(4): 1-6 (2010); Bryson et al., "Prediction of Immunogenicity of Therapeutic Proteins: Validity of Computational Tools," *Biodrugs* 24(1)1-8 (2010); Holgate & Baker, "Circumventing Immunogenicity in the Development of Therapeutic Antibodies," *IDrugs* 12(4):233-237 (2009); Perry et al., "New Approaches to Prediction of Immune Responses to Therapeutic Proteins during Preclinical Development," *Drugs R D* 9(6):385-396 (2008); and, Baker & Jones, "Identification, and removal of immunogenicity in

therapeutic proteins,” *Current Opinion in Drug Discovery & Development* 10(2):219-227 (2007).

Pseudomonas Exotoxin A

[0010] *Pseudomonas* exotoxin A (PE-A) is a highly potent, 66 kD, cytotoxic protein secreted by the bacterium *Pseudomonas aeruginosa*. PE-A causes cell death by inhibiting protein synthesis in eukaryotic cells via inactivation of translation elongation factor 2 (EF-2), which is mediated by PE-A catalyzing ADP-ribosylation of EF-2 (i.e., transfer of an ADP ribosyl moiety onto EF-2). PE-A typically produces death by causing liver failure.

[0011] PE-A has at least three different structural domains responsible for various biological activities (FIG. 1). See e.g., Siegall et al, *Biochemistry*, vol. 30, pp. 7154-7159 (1991); Theuer et al. *Jour. Biol. Chem.*, vol 267, no. 24, pp. 16872-16877 (1992); and, U.S. Pat. No. 5,821,238. PE-A domain IA (amino acids 1-252 (see e.g., SEQ ID NO:133)) is responsible for cell binding. Domain U (amino acids 253-364 (see e.g., SEQ ID NO:133)) is responsible for translocation of PE-A into the cell cytosol. Domain III, the cytotoxic domain (amino acids 400-613 (see e.g., SEQ ID NO:133)), is responsible for ADP ribosylation of Elongation Factor 2 (EF2); which thereby inactivates EF2, subsequently causing cell death. Additionally, a function for domain IB (amino acids 365-399 (SEQ ID NO:139)) has not been established. Indeed, it has been reported that amino acids 365-380 (SEQ ID NO:138) within domain IB can be deleted without producing an identifiable loss of function. See, Siegall et al, *Biochemistry*, vol. 30, pp. 7154-7159(1991).

[0012] It has also been reported that PE-A may comprise any one of at least three different carboxy-terminal tails (FIG. 1); these appear to be essential for maintaining or recycling proteins into the endoplasmic reticulum. See, Theuer et al, *J. Biol. Chem.*, vol 267, no. 24, pp. 16872-16877 (1992); Chaudhary et al., *Proc. Natl. Acad. Sci. USA*, vol. 87, pp. 308-312 (1990); and, Seetharam et al. *Jour. Biol. Chem.*, vol 266, 17376-17381 (1991). In particular, in correspondence with the exemplary sequence shown in FIG. 1 (SEQ ID NO:133) these alternative carboxy-terminal tails comprise amino acid sequences:

```
609 - REDLK - 613;      (SEQ ID NO: 135)
609 - REDL - 612;      (SEQ ID NO: 136)
and
609 - KDEL - 612.      (SEQ ID NO: 137)
```

[0013] Variants of PE-A, modified to lack the cell binding domain but coupled to heterologous cell-specific targeting molecules (e.g., antibodies), have been, shown to have reduced levels of non-specific toxicity. See e.g., U.S. Pat. No. 4,892,827.

[0014] Various forms of PE-A (e.g., truncated/deletion forms with molecular weights of ~37 kD, 38 kD, 40 kD, et cetera) have been combined with a number of growth factors, antibodies, and oilier proteins to generate cytotoxins which selectively target cells of a desired phenotype. See, for example:

[0015] Kreilman et al., “Recombinant immunotoxins and other therapies for relapsed/refractory hairy cell leukemia,” *Leuk. Lymphoma*, Suppl. 2:82-86 (June-2011);

[0016] Itoi et al., “Targeting of locus ceruleus noradrenergic neurons expressing human interleukin-2 receptor α -subunit in transgenic mice by a recombinant immunotoxin anti-Tac(Fv)-PE38,” *J. Neurosci.*, 31 (16):632-6139 (April-2011);

[0017] Shapira et al., “An immunoconjugate of anti-CD24 and *Pseudomonas* exotoxin selectively kills human colorectal tumors in mice,” *Gastroenterology*, 140(3):935-946 (March-2011);

[0018] Kuan et al., “Affinity-matured anti-glycoprotein NMB recombinant immunotoxins targeting malignant gliomas and melanomas,” *Int. J. Cancer*, 129(1): 111-21 (July-2011);

[0019] Hu et al., “Investigation of a plasmid containing a novel immunotoxin VEGF165-PE38 gene for antiangiogenic therapy in a malignant glioma model,” *Int. J. Cancer*, 127(9): 2222-2229 (November-2010);

[0020] Mareeva et al., “A novel composite immunotoxin that suppresses rabies virus production by the infected cells,” *J. Immunol. Methods*, 353(1-2):78-86 (February-2010);

[0021] Zielinski et al., “Affitoxin—a novel recombinant, HER2-specific, anticancer agent for targeted therapy of HER2-positive tumors,” *J. Immunother.* 32(8):817-825 (October-2009);

[0022] Theuer et al, *J. Biol. Chem.*, 267(24): 16872-16877 (1992);

[0023] Pastan et al., “Recombinant toxins for cancer treatment,” *Science*, 254; 1173-1177 (1991);

[0024] U.S. Pat. No. 5,821,238 (“Recombinant *Pseudomonas* Exotoxins with Increased Activity”); and

[0025] U.S. Pat. No. 4,892,827 (“Recombinant *Pseudomonas* Exotoxins: Construction of an Active Immunotoxin with Low Side Effects”).

[0026] A significant disadvantage in using PE-A for treatment of disease, however, is that

[0027] it is a foreign (non-self) protein being introduced into a heterologous host (e.g., a human). Introduction of non-self proteins into heterologous hosts commonly elicits host immune reactions, such as the generation of antibodies (“neutralizing antibodies”) or immune cell reactions (e.g., cytotoxic T cell responses) which are directed at eliminating the non-self protein (i.e., PE-A). Accordingly, it would be advantageous if elements of PE-A (PE-A epitopes) which are recognized and targeted as “non-self” could be removed prior to use of this molecule as a therapeutic agent.

Deimmunization of PE

[0028] Some investigators have previously attempted to identify and remove immunogenic determinants from PE-A (i.e., to “deimmunize” PE-A). See, for example:

[0029] Pastan et al., “Immunotoxins with decreased immunogenicity and improved activity,” *Leukemia and Lymphoma*, 52(S2):87-90 (June-2011);

[0030] Onda et al., “Recombinant immunotoxin against B-cell malignancies with no immunogenicity in mice by removal of B-cell epitopes,” *Proc. Natl. Acad. Sci. USA*, 108(14): 5742-5747 (April-2011);

[0031] Hansen al., “A recombinant immunotoxin targeting CD22 with low immunogenicity, low nonspecific toxicity, and high antitumor activity in mice,” *J. Immunother.* 33(3): 297-304 (April-2011);

[0032] Stish et al., “Design and modification of EGF4KDEL 7Mut, a novel bispecific ligand-directed

toxin, with decreased immunogenicity and potent anti-mesothelioma activity," *Br. J. Cancer*, 101(7): 1114-1123 (October-2009);

[0033] Nagata et al., "Removal of B cell, epitopes as a practical approach for reducing the immunogenicity of foreign protein-based therapeutics," *Adv. Drug Deliv. Rev.*, 61(11):977-985 (September-2009);

[0034] Onda et al., "An immunotoxin with greatly reduced immunogenicity by identification and removal of B cell epitopes," *Proc. Natl. Acad. Sci. USA*, 105(32): 11311-11316 (August-2008); and

[0035] Pastan et al., "Mutated *Pseudomonas* Exotoxins with Reduced Antigenicity," U.S. Patent Application No. 2009/0142341.

[0036] Despite progress in the area of deimmunization of PE-A, there remains a need for the development of optimized, less immunogenic or non-immunogenic, biologically active forms of this useful cytotoxin. The invention described herein addresses this need.

BRIEF SUMMARY OF THE INVENTION

[0037] Peptides spanning the sequence of an approximately 38 kD (predicted molecular weight) form of *Pseudomonas* exotoxin A protein (SEQ ID NO:1) were analyzed for the presence of immunogenic CD4+ T cell epitopes. A total of 120 overlapping 15 mer peptides spanning this sequence (SEQ ID NO:1), but also including an amino terminal (Gly_{x5}-Ser)_{x2} linker sequence (SEQ ID NO:3) to produce a 359 amino acid Gly-Set-PE38 polypeptide sequence (SEQ ID NO:2), were tested against a cohort of healthy human donors. CD4+ T cell responses against individual peptides were measured via proliferation assays. Assay data was used to compile a T cell epitope map of the PE38 sequence. Six immunogenic T cell epitopes were identified. Residues were then identified, within each of these epitopes for use in targeted amino acid substitutions to reduce or prevent PE38-induced immunogenicity. Reduction or prevention of PE immunogenicity should allow for multiple therapeutic administrations of cytotoxic PE for use, for example, in the targeted destruction of cancer cells in vivo (such as when administered as an immunoconjugate or cell-surface targeted fusion protein).

BRIEF DESCRIPTION OF THE DRAWINGS

[0038] FIG. 1. Example of a *Pseudomonas* exotoxin A protein and domains which may be contained therein.

[0039] FIG. 2. Comparison of the frequency of donor allo-types expressed in the IEX01 study cohort (n=52) and the world population.

[0040] FIGS. 3A & 3B. CD4+ T cell epitope map of IEX01 PE38 sequence using overlapping 15 mer peptides tested against 52 healthy donors. The non-adjusted (FIG. 3A) and adjusted (FIG. 3B) proliferation assay data for the PE38 sequence is shown. Peptides inducing positive (SI $\geq$ 2.00, p<0.05, including borderline responses) T cell proliferation responses at a frequency above the background response rate (mean positive T cell responses plus SD) contain T cell epitopes (dotted line indicates the background response threshold). KLH induced positive responses in (SI $\geq$ 2.00, p<0.05) 75% of (non-adjusted) donors.

[0041] FIG. 4. Alignment of peptides 50, 52 and 53 showing the predicted HLA-DR core 9 mer binding register. Predicted core 9 mer sequences are bracketed by p1 and p9

anchor residues. Peptides that stimulated positive T cell responses in the adjusted data set are shown. Amino acid numbering (residues 153 and 161) correspond to SEQ ID NO:2 (PE38 of SEQ ID NO:1 plus amino-terminal linker GGGGGSGGGGS (SEQ ID NO:3)).

[0042] FIG. 5. Alignment of peptides 65, 67 and 68 showing one predicted HLA-DR core 9 mer binding register. Predicted core 9 mer sequences are bracketed by p1 and p9 anchor residues. Peptides that stimulated positive T cell responses in the adjusted data set are shown. Amino acid numbering (residues 196 and 204) correspond to SEQ ID NO:2 (PE38 of SEQ ID NO:1 plus amino-terminal linker GGGGGSGGGGS (SEQ ID NO:3)).

[0043] FIG. 6. Alignment of peptides 81 and 82 showing the potential HLA-DR core 9 mer binding register. Predicted core 9 mer sequences are bracketed by p1 and p9 anchor residues. Peptides that stimulated positive T cell responses in the adjusted data set are shown. Amino acid numbering (residues 244 and 252) correspond to SEQ ID NO:2 (PE38 of SEQ ID NO:1 plus amino-terminal linker GGGGGSGGGGS (SEQ ID NO:3)).

[0044] FIG. 7. Alignment of peptides 110 and 111 showing a predicted HLA-DR core 9 mer binding register. Predicted core 9 mer sequences are bracketed by p1 and p9 anchor residues. Peptides that stimulated positive T cell responses in the adjusted data set are shown. Amino acid numbering (residues 333 and 341) correspond to SEQ ID NO:2 (PE38 of SEQ ID NO:1 plus amino-terminal linker GGGGGSGGGGS (SEQ ID NO:3)).

[0045] FIG. 8. Position of CD4+ T cell epitopes within the PE38 sequence. T cell epitopes identified by EPISCREEN™ T cell epitope mapping are shown as shaded bars above the sequence. The frequency of donors responding (SI $\geq$ 2.00, p<0.05) to each epitope are indicated by the shading of the bars; light grey<10%, mid grey 10-14%; dark grey $\geq$ 14%. Numbers assigned to each individual donor (that responded to a corresponding epitope) are included within each shaded bar.

[0046] FIG. 9. In vivo Transcription/Translation (IVTT) shows that circular plasmid expression vector encoding PE38-IL2 fusion protein was slightly better at inhibiting Luciferase protein synthesis compared to linearized plasmid encoding the same PE38-IL2 fusion protein.

[0047] FIG. 10. Luciferase activity measure in counts per second (CPS) in In vitro Transcription/Translation (IVTT) assays of genes encoding either Wild-Type (WT) PE or encoding amino acid substituted PE.

[0048] FIG. 11. Analysis of production of cytokines IL-2 and IL-6 stimulated in response to expression of genes encoding either Wild-Type (WT) PE or encoding amino acid substituted PE.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions and Descriptions

[0049] Unless specifically indicated otherwise, as used herein the term "PE" or "PE-A" is intended to indicate a polypeptide comprising a cytotoxic polypeptide sequence derived from a wild-type or naturally occurring form of *Pseudomonas aeruginosa* exotoxin A protein. In addition to cytotoxic polypeptide sequences, PE polypeptides may comprise additional naturally occurring or heterologous polypeptide sequences. Additional naturally occurring polypeptide sequences may include sequences such as are found in full-length *Pseudomonas* exotoxin A protein, for example, amino

acid sequences responsible for cytosolic translocation and cell-specific targeting (as discussed further herein)). Additional heterologous polypeptide sequences may include sequences with which at least a PE cytotoxic polypeptide is fused to impart additional functions or properties. (For example, a PE cytotoxic polypeptide may be fused to antigen binding polypeptide sequences such as an scFv antibody.) Examples of sequences comprising a cytotoxic portion of PE can be found in SEQ ID NO:1 and SEQ ID NO:4 spanning amino acid residues Phe-134 to Lys-347. Examples of sequences comprising a cytotoxic portion of PE can also be found in SEQ ID NO:133 and SEQ ID NO:134 spanning amino acid residues Phe-400 to Lys-613.

[0050] As used herein in reference to PE, unless indicated otherwise, a “cytotoxic polypeptide” or “cytotoxic polypeptide sequence” is intended to indicate a polypeptide (or portion thereof) which is capable of inactivating translation elongation factor 2 (EF-2), mediating ADP-ribosylation of EF-2, inhibiting protein synthesis, or inducing cell, death. For example, it has been demonstrated that PE domain III, comprised of amino acid residues 400-613 of SEQ ID NO:133, is sufficient to mediate ADP-ribosylation of EF-2 and thereby cause cell death. See, Theuer et al., *J. Biol. Chem.*, vol. 267, no. 24, pp. 16872-16877 (1992) and Hwang et al., *Cell* vol. 48, pp. 129-136 (1987).

[0051] Cytotoxic polypeptide sequences in the present invention may also comprise alternative carboxy-terminal sequences. See, Theuer et al., Chaudhary et al. and, Seetharam et al. In particular embodiments, examples of carboxy-terminal tails of PE38 in the present invention may comprise sequences as shown in FIG. 1 (SEQ ID NO:133). Hence, exemplary alternative carboxy-terminal tails may comprise amino acid sequences:

609-REDLK-613 (SEQ ID NO:135; numbers 609-613 correspond to SEQ ID NO:133)

609-REDL-612 (SEQ ID NO:136; numbers 609-612 correspond to SEQ ID NO:133); and

609-KDEL-612 (SEQ ID NO:137; numbers 609-612 correspond to SEQ ID NO:133)

[0052] Unless specifically indicated otherwise, as used herein the term “PE38” is intended to indicate a *Pseudomonas aeruginosa* exotoxin A (PE (or PE-A)) molecule comprising an amino acid sequence as shown in SEQ ID NO:1. The amino acid sequence used to generate peptide sequences referenced in the Examples is shown in SEQ ID NO:2. SEQ ID NO:2 comprises an amino terminal GGGGSGGGGS linker sequence (SEQ ID NO:3) fused to the PE38 amino acid sequence of SEQ ID NO:1. A variant form of PE38 is shown in SEQ ID NO:4. SEQ ID NO:4 differs from SEQ ID NO:1 by comprising a Ser-to-Asn change at position 114, a Ile-to-Val change at position 141, and a Gly-to-Ser change at position 249.

[0053] As used herein, unless specifically stated otherwise, “biological activity” in reference to *Pseudomonas* exotoxin A (PE-A), PE or PE38 is intended to indicate at least one of the biological activities exhibited by naturally occurring forms of the *Pseudomonas aeruginosa* exotoxin A molecule. These activities include, for example, cell killing or cell cytotoxic activity (a.k.a., cell cytotoxicity), inactivation of translation elongation factor EF-2, ADP-ribosylation of EF-2, and inhibition of protein synthesis. The biological activity of PE and PE38 polypeptides (and modified forms thereof; e.g., PE and PE38 amino acid substituted variants and fusion proteins) can be measured using assays and experiments which are well-

known and routinely used by those skilled in the art. Examples of some of these assays and experiments are further described and referenced herein, without limitation, in the Examples sections included herein.

[0054] As used herein, the term “having *Pseudomonas* exotoxin A (PE-A) biological activity” (or “PE biological activity”) is intended to indicate molecules exhibiting about 5% or more of at least one biological activity compared to a corresponding wild-type, naturally occurring, or non-amino acid substituted form of PE or PE-A. In some embodiments, molecules “having *Pseudomonas* exotoxin A biological activity” (or “PE biological activity”) exhibit 5% or more, about 10% or more, 10% or more, about 15% or more, 15% or more, about 20% or more, 20% or more, about 25% or more, 25% or more, about 30% or more, 30% or more, about 35% or more, 35% or more, about 40% or more, 40% or more, about 45% or more, 45% or more, about 50% or more, 50% or more, about 60% or more, 60% or more, about 70% or more, 70% or more, about 75% or more, 75% or more, about 80% or more, 80% or more, about 85% or more, 85% or more, about 90% or more, 90% or more, about 95% or more, 95% or more, about 100%, or 1.00% of at least one biological activity compared to a corresponding wild-type, naturally occurring, or non-amino acid substituted forms of PE or PE-A.

[0055] As used herein, the term “wild-type” *Pseudomonas* exotoxin A (PE-A) (or “wild-type” PE) biological activity is intended to indicate at least one or more biological activities exhibited by naturally occurring forms of the *Pseudomonas* exotoxin A (PE-A) or PE polypeptides. These include, for example, without limitation, activities such as cell killing or cell cytotoxic activity (a.k.a., cell cytotoxicity), inactivation of translation elongation factor EF-2, ADP-ribosylation of EF-2, and inhibition of protein synthesis. Two examples, without limitation, of polypeptide sequences representing “wild-type” or non-amino acid substituted forms of PE-A are shown in SEQ ID NO:133 and SEQ ID NO:134. Two examples, without limitation, of polypeptide sequences representing “wild-type” or non-amino acid substituted forms of PE are shown in SEQ ID NO:1 (PE38) and SEQ ID NO:4 (variant of PE38).

[0056] As used or claimed herein the term “a” or “an” in reference to the subsequent recited entity refers to one or more of that entity; for example, “a PE38 antibody” or “a polynucleotide encoding PE38” is understood to indicate one or more PE38 antibody molecules and one or more polynucleotides encoding PE38, not a single PE38 antibody molecule nor a single polynucleotide molecule encoding PE38, respectively. As such, the terms “a” (or “an”), “one or more,” and “at least one” can be used interchangeably herein.

[0057] Likewise, as used herein, the term “polypeptide” is intended to encompass a singular “polypeptide” as well as plural “polypeptides,” and refers to a molecule composed of monomers (amino acids) linked by amide bonds (also known as peptide bonds). The term “polypeptide” refers to any chain or chains of two or more amino acids, and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides, “protein,” “amino acid chain,” or any other term used to refer to a chain or chains of two or more amino acids, are included within the definition of “polypeptide,” and the term “polypeptide” may be used instead of, or interchangeably with any of these terms. The term “polypeptide” is also intended to refer to the products of post-expression modifications of the polypeptide, such as, but without limitation glycosylation, acetylation, phosphorylation, ami-

dation, et cetera. A “polypeptide” unless specifically described otherwise herein, may be derived from a natural biological source or produced by recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It may be generated in any manner, including by chemical synthesis.

[0058] Polypeptides may have a defined three-dimensional structure, although they do not necessarily have such structure. Polypeptides with a defined three-dimensional structure may be referred to as “folded” or having a “tertiary” structure. Polypeptides not configured into a three-dimensional structure, are referred to as unfolded. As used herein, the term glycoprotein refers to a protein coupled to at least one carbohydrate moiety attached to the protein via a covalent bond.

[0059] The term “isolated” is intended to indicate a biological component no longer in its naturally occurring milieu. For example, an “isolated polypeptide” or “isolated polynucleotide” is intended to indicate a polypeptide or polynucleotide, respectively, which has been removed from its naturally occurring milieu and which may have been inserted within a non-naturally occurring milieu. By way of example, this would include, without limitation, a polynucleotide which has been removed from a naturally occurring location within a host genome, and subsequently inserted, for example, into an expression vector or inserted into a new host genome location or into the genome of a heterologous host organism. The “isolation” of a polypeptide or polynucleotide, as used herein, requires no particular level of purification. For example, recombinantly produced polypeptides expressed in host cells are considered isolated for purposes of the invention, as are native or recombinant polypeptides which have been separated, fractionated, or partially or substantially purified by any suitable technique.

[0060] Polypeptide embodiments also include fragments, derivatives, analogs, variants and fusion proteins; preferably but not necessarily wherein such embodiments retain one or more biological activities associated with a corresponding full-length or naturally occurring polypeptide. Fragments include proteolytic fragments, deletion fragments, and fragments encoded by synthetically or recombinantly produced polynucleotides. Variants may occur naturally or be non-naturally occurring. Non-naturally occurring variants may be produced using art-known mutagenesis techniques. Variant polypeptides may comprise conservative or non-conservative amino acid substitutions, deletions, or additions. Derivatives include, but are not limited to, polypeptides which contain one or more non-naturally occurring amino acids, non-standard amino acids, and, amino acid analogs. Polypeptide embodiments may comprise amino acid sequences which are at least 60% identical, at least 70% identical, at least 80% identical, at least 85% identical, at least 90% identical, at least 95% identical, at least 97% identical, at least 98% identical, or at least 99% identical to SEQ ID NO:1

[0061] Unless specifically defined otherwise, the term “polynucleotide” is intended to indicate nucleic acid molecules or constructs as routinely used and understood, by those of skill in the art. For example, nucleic acids include, but are not limited to, molecules such as messenger RNA (mRNA), plasmid DNA (pDNA), complementary DNA (cDNA), and genomic DNA (gDNA). A polynucleotide may comprise a conventional phosphodiester bond or a non-conventional bond (e.g., an amide bond, such as found in peptide nucleic acids (PNA)). The terms “polynucleotide” and “nucleic acid” are intended to include embodiments wherein

any one or more sequences of poly-nucleotide or nucleic acid segments are contained, or comprised within, a larger polynucleotide or nucleic acid sequence. For example, but without limitation, and unless stated otherwise to the contrary herein, reference to a nucleic acid such as “a polynucleotide encoding PE38” is intended to include nucleic acids comprising “a polynucleotide encoding PE38” wherein such polynucleotide may also be part of a larger nucleic acid or polynucleotide, such as an expression vector or a polynucleotide/nucleic acid encoding an PE fusion protein.

[0062] An “isolated” nucleic acid or polynucleotide is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, a recombinant polynucleotide encoding an antibody contained in a vector is considered isolated for the purposes of the present invention. Further examples of an isolated polynucleotide include recombinant polynucleotides maintained in heterologous host cells or purified (partially or substantially) polynucleotides in solution. Isolated RNA molecules include in vivo or in vitro synthesized RNA molecules; including synthetically produced molecules.

[0063] As used herein, a “coding region” is a portion of nucleic acid containing codons which may be translated into amino acids, although “stop codons” (TAG, TGA, or TAA) are not translated into an amino acids, but may also be considered to be part of a coding region. Unless stated otherwise herein, promoters, ribosome binding sites, transcriptional terminators, introns, and the like, are not considered part of a coding region. Two or more coding regions of the present invention can be present in a single polynucleotide construct, e.g., on a single vector, or in separate polynucleotide constructs, e.g., on separate (different) vectors. Furthermore, any vector may contain a single coding region, or may comprise two or more coding regions, e.g., a single vector may separately encode an immunoglobulin heavy chain variable region and an immunoglobulin light chain variable region. In addition, a vector, polynucleotide, or nucleic acid embodiments may encode heterologous coding regions, either fused or unfused to a nucleic acid encoding a different heterologous polypeptide. Heterologous coding regions include without limitation specialized elements or motifs, such as a secretory signal peptide or a heterologous functional domains.

[0064] In certain embodiments, the polynucleotide or nucleic acid is DNA. In the case of DNA, a polynucleotide comprising a nucleic acid which encodes a polypeptide normally may include a promoter and/or other transcription or translation control elements operably associated with one or more coding regions. An operable association is when a coding region for a gene product, e.g., a polypeptide, is associated with one or more regulatory sequences in such a way as to place expression of the gene product under the influence or control of the regulatory sequence(s). Two DNA fragments (such as a polypeptide coding region and a promoter associated therewith) are “operably associated” if induction of promoter function results in the transcription of mRNA encoding the desired gene product. Thus, a promoter region would be operably associated with a nucleic acid encoding a polypeptide if the promoter was capable of effecting transcription of that nucleic acid. Other transcription control elements, besides a promoter, include for example, but without limitation, enhancers, operators, repressors, and transcription termination signals, can be operably associated with the poly-

nucleotide to direct cell-specific transcription. Suitable promoters and other transcription control regions are disclosed herein.

[0065] The terms “antibody” and “immunoglobulin” may be used interchangeably herein. An antibody or immunoglobulin comprises at least the antigen-binding elements (e.g., complementarity determining regions or CDRs) of the variable domain of a heavy chain and/or of the variable domain of a light chain. Basic immunoglobulin structures in vertebrate systems are well understood by those of skill in the art. See, e.g., Harlow & Lane, *Using Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 1999 (ISBN 0879695447)); see also, Harlow & Lane, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988). The term “immunoglobulin” or “antibody” comprises various broad classes of antibody molecules, such as, but without limitation, IgG, IgM, IgA, IgG, and IgE classes of antibodies; as well as antibody subclasses (isotypes), such as, TgG1, IgG2, IgG3, IgG4, IgA1, et cetera.

[0066] Antibodies or antigen-binding fragments, variants, or derivatives thereof of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized, primatized, or chimeric antibodies, single chain antibodies, epitope-binding fragments, e.g., Fab, Fab' and F(ab'), Fd, Fvs, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv), fragments comprising either a VL or VH domain, fragments produced by a Fab expression library, and anti-idiotypic (anti-Id) antibodies.

[0067] As used herein, an “epitope” or “antigenic determinant” is the part of a polypeptide, antigen, or molecule that is recognized by the immune system, specifically by antibodies, B cells, or T cells. Epitopes of polypeptide antigens may function as conformational epitopes or linear epitopes. A conformational epitope is comprised of non-linear sections of a target molecule (such as that formed via the tertiary structure of a folded polypeptide). In contrast, amino acids that make up a linear epitope may be comprised of a continuous sequence of amino acids or may be comprised only of particular amino acid residues critical to antibody/B cell/T cell binding.

[0068] By “specifically binds,” it is generally meant that an antibody binds to an epitope via its antigen binding domain, and that the binding entails some complementarity between the antigen binding domain and the epitope. According to this definition, an antibody is said to “specifically bind” to an epitope when it binds to that epitope, via its antigen binding domain more readily than it would bind to a random, unrelated epitope. The term “specificity” may be used herein to qualify the relative affinity by which a certain antibody binds to a certain epitope. For example, antibody “A” may be deemed to have a higher specificity for a given epitope than antibody “B,” or antibody “A” may be said to bind to epitope “C” with a higher specificity than it has for related epitope “D.”

[0069] By “preferentially binds,” it is meant that the antibody specifically binds to an epitope more readily than it would bind to a related, similar, homologous, or analogous epitope. Thus, an antibody which “preferentially binds” to a given epitope would more likely bind to that epitope than to a related epitope, even though such an antibody may cross-react with the related epitope.

[0070] An antibody is said to competitively inhibit binding of a reference antibody to a given epitope if it preferentially binds to that epitope to the extent that it blocks, to some

degree, binding of the reference antibody to the epitope. Competitive inhibition may be determined by any method known in the art, for example, competition ELISA assays. An antibody may be said to competitively inhibit binding of the reference antibody to a given epitope by at least 90%, at least 80%, at least 70%, at least 60%, or at least 50%.

[0071] As used herein, the term “affinity” refers to a measure of the strength of the binding of an individual epitope with the CDR of an immunoglobulin molecule. See, e.g., Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed, 1988) at pages 27-28. As used herein, the term “avidity” refers to the overall stability of the complex between a population of immunoglobulins and an antigen, that is, the functional combining strength of an immunoglobulin mixture with the antigen. See, e.g., Harlow at pages 29-34. Avidity is related to both the affinity of individual immunoglobulin molecules in the population with specific epitopes, and also the valencies of the immunoglobulins and the antigen. For example, the interaction between a bivalent monoclonal antibody and an antigen with a highly repeating epitope structure, such as a polymer, would be one of high avidity.

[0072] The term “cross-reactivity” refers to the ability of an antibody, specific for one antigen, to react with a second antigen: a measure of relatedness between two different antigenic substances. Thus, an antibody is cross reactive if it binds to an epitope other than the one that induced its formation. The cross reactive epitope generally contains many of the same complementary structural features as the inducing epitope, and in some cases, may actually fit better than the original.

[0073] As used herein, the terms “linked,” “fused” or “fusion” may be used interchangeably. These terms refer to the joining together of two more elements or components, by whatever means including chemical conjugation or recombinant means. An “in-frame fusion” refers to the joining of two or more polynucleotide open reading frames (ORFs) to form a continuous longer ORF, in a manner that maintains the correct translational reading frame of the original ORFs. Thus, a recombinant fusion protein is a single protein containing two or more segments that correspond to polypeptides encoded by the original ORFs (which segments are not normally so joined in nature.) Although the reading frame is thus made continuous throughout the fused segments, the segments may be physically or spatially separated by, for example, in-frame linker sequence. For example, polynucleotides encoding the CDRs of an immunoglobulin variable region may be fused, in-frame, but be separated by a polynucleotide encoding at least one immunoglobulin framework region or additional CDR regions, as long as the “fused” CDRs are co-translated as part of a continuous polypeptide.

[0074] In the context of polypeptides, a “linear sequence” or a “sequence” is an order of amino acids in a polypeptide in an amino to carboxyl terminal direction in which residues that neighbor each other in the sequence are contiguous in the primary structure of the polypeptide.

[0075] A “variant” of a polypeptide or protein refers to any analogue, fragment, derivative, or mutant which is derived from a polypeptide or protein and which retains at least one biological property of the polypeptide or protein. Different variants of the polypeptide or protein may exist in nature or may be generated artificially (e.g., via synthetic or genetic engineering). Variants may be allelic variations characterized by differences in the nucleotide sequences of the structural

gene coding for the protein, or may involve differential splicing or post-translational modification. The skilled artisan can produce variants having single or multiple amino acid substitutions, deletions, additions, or replacements. Variants may include, inter alia: (a) variants in which one or more amino acid residues are substituted with, for example, conservative amino acids, non-conservative amino acids, or amino acid analogs (b) variants in which one or more amino acids are added to the polypeptide or protein, (c) variants in which one or more of the amino acids includes a substituent group, and (d) variants in which the polypeptide or protein is fused with another polypeptide such as serum albumin. The techniques for obtaining these variants, including genetic (suppressions, deletions, mutations, etc.), chemical, and enzymatic techniques, are known to those of skill in the art.

[0076] The term “expression” as used herein refers to a process by which a gene produces a biochemical, for example, an RNA or polypeptide. It includes without limitation transcription of the gene into RNA molecules such as, for example, messenger RNA (mRNA), transfer RNA (tRNA) or any other RNA product, and the translation of mRNA into polypeptide(s).

[0077] Expression of a gene produces a “gene product.” As used herein, a gene product can be either a nucleic acid, e.g., a messenger RNA produced by transcription of a gene, or a polypeptide which is translated from a transcript. Gene products described herein further include nucleic acids with post transcriptional modifications, e.g., polyadenylation, or polypeptides with post translational modifications, e.g., methylation, glycosylation, the addition of lipids, association with other protein subunits, proteolytic cleavage, et cetera.

[0078] As used herein, the term “gene” refers to a polynucleotide comprising nucleotides that encode a functional molecule, including functional molecules produced by transcription only (e.g., a bioactive RNA species) or by transcription and translation (e.g. a polypeptide). The term “gene” encompasses cDNA and genomic DNA nucleic acids. “Gene” also refers to a nucleic acid fragment that expresses a specific RNA, protein or polypeptide, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers to any gene that is not a native gene, comprising regulatory and/or coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. A chimeric gene may comprise coding sequences derived from different sources and/or regulatory sequences derived from different sources. “Endogenous gene” refers to a native gene in its natural location in the genome of an organism. A “foreign” gene or “heterologous” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

[0079] “RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary

transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. “Messenger RNA (mRNA)” refers to the RNA that is without introns and that can be translated into protein by the cell. “cDNA” refers to a double-stranded DNA that is complementary to and derived from mRNA. “Sense” RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. “Antisense RNA” refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, or the coding sequence.

[0080] A “vector” refers to any vehicle for the cloning of and/or transfer of a nucleic acid into a host cell. A vector may be a replicon to which another DNA segment may be attached so as to bring about the replication of the attached segment. A “replicon” refers to any genetic element (e.g. plasmid, phage, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication in vivo, i.e., capable of replication under its own control. The term “vector” includes both viral and nonviral vehicles for introducing the nucleic acid into a cell in vitro, ex vivo or in vivo. A large number of vectors known in the art may be used to manipulate nucleic acids, incorporate coding sequences into genes, et cetera. Possible vectors include, for example, plasmids or modified viruses including, for example bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives, or the Bluescript vector. Another example of vectors that are useful in the present invention is the Ultra Vector™ Production System (Intrexon Corp., Blacksburg, Va.) as described in WO 2007/038276, incorporated by reference herein. For example, the insertion of the DNA fragments corresponding to response elements and promoters into a suitable vector for in vitro and/or in vivo expression of modified forms of PE (and fragments thereof) as described herein (including fusion proteins, conjugates, and otherwise linked forms) can be accomplished by ligating the appropriate DNA fragments into a chosen vector that has complementary cohesive termini. Alternatively, the ends of the DNA molecules may be enzymatically modified or any site may be produced by ligating nucleotide sequences (linkers) into the DNA termini. Such vectors may be engineered to contain selectable marker genes that provide for the selection of cells that have incorporated the marker into the cellular genome. Such markers allow identification and/or selection of host cells that incorporate and express the proteins encoded by the marker.

[0081] Viral vectors, and particularly retroviral vectors, have been used in a wide variety of gene delivery applications in cells, as well as living animal subjects. Viral vectors that can be used to express embodiments of the invention described herein include, but are not limited to, retrovirus, adeno-associated virus, pox, baculovirus, vaccinia, herpes simplex, Epstein-Barr, adenovirus, geminivirus, and caulimovirus vectors. Non-viral vectors include plasmids, liposomes, electrically charged lipids (cytofectins) DNA-protein complexes, and biopolymers. In addition to a nucleic acid, a vector may also comprise one or more regulatory regions, and/or selectable markers useful in selecting, measuring, and monitoring nucleic acid transfer results (e.g., monitoring transfer to target or non-target tissues, duration of expression, et cetera).

[0082] The term “plasmid” refers to an extra-chromosomal element often carrying a gene that is not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome integrating sequences, phage or nucleotide sequences, linear, circular, or supercoiled, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell.

[0083] A “cloning vector” refers to a “replicon,” which is a unit length of a nucleic acid, preferably DNA, that replicates sequentially and which comprises an origin of replication, such as a plasmid, phage or cosmid, to which another nucleic acid segment may be attached so as to bring about the replication of the attached segment. Cloning vectors may be capable of replication in one cell type and expression in another (“shuttle vector”). Cloning vectors may comprise one or more sequences that can be used for selection of cells comprising the vector and/or one or more multiple cloning sites for insertion of sequences of interest. The term “expression vector” refers to a vector, plasmid or vehicle designed to enable the expression of an inserted nucleic acid sequence following transformation into the host. The cloned gene, i.e., the inserted nucleic acid sequence, is usually placed under the control of control elements such as a promoter, a minimal promoter, an enhancer, or the like. Initiation control regions or promoters, which are useful to drive expression of a nucleic acid in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving expression of these genes can be used in an expression vector, including but not limited to, viral promoters, bacterial promoters, animal promoters, mammalian promoter's, synthetic promoters, constitutive promoters, tissue specific promoters, pathogenesis or disease related promoters, developmental specific promoters, inducible promoters, light regulated promoters. CYC1, HIS3, GAL1, GAL4, GAL10, ADH1, PGK, PH05, GAPDH, ADC1, TRP1, URA3, LEU2, ENO, TP1, alkaline phosphatase promoters (useful for expression in *Saccharomyces*); AOX1 promoter (useful for expression in *Pichia*); β -lactamase, lac, ara, tet, trp, IP_L , IP_R , T7, tac, and trc promoters (useful for expression in *Escherichia coli*); light regulated-, seed specific-, pollen specific-, ovary specific-, cauliflower mosaic virus 35S, CMV 35S minimal, cassava vein mosaic virus (CsVMV), chlorophyll a/b binding protein, ribulose 1,5-bisphosphate carboxylase, shoot-specific, rot specific, chitinase, stress inducible, rice tungro bacilliform virus, plant super-promoter, potato leucine aminopeptidase, nitrate reductase, mannopine synthase, nopaline synthase, ubiquitin, zein protein, and anthocyanin promoters (useful for expression in plant cells); animal and mammalian promoters known in the art including, but are not limited to, the SV40 early (SV40e) promoter region, the promoter contained in the 3' long terminal repeat (LTR) of Rous sarcoma virus (RSV), the promoters of the E1A or major late promoter (MLP) genes of adenoviruses (Ad), the cytomegalovirus (CMV) early promoter, the herpes simplex virus (HSV) thymidine kinase (TK) promoter, a baculovirus IE1 promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, an albumin promoter, the regulatory sequences of the mouse metallothionein-L promoter and

transcriptional control regions, the ubiquitous promoters (HPRT, vimentin, α -actin, tubulin and the like), the promoters of the intermediate filaments (desmin, neurofilaments, keratin, GFAP, and the like), the promoters of therapeutic genes (of the MDR, CFTR or factor VII type, and the like), pathogenesis or disease related-promoters, and promoters that exhibit tissue specificity and have been utilized in transgenic animals, such as the elastase I gene control region which is active in pancreatic acinar cells; insulin gene control region active in pancreatic beta cells, immunoglobulin gene control region active in lymphoid cells, mouse mammary tumor virus control region active in testicular, breast, lymphoid and mast cells; albumin gene, Apo AI and Apo AII control regions active in liver, alpha-fetoprotein gene control region active in liver, alpha 1-antitrypsin gene control region active in the liver, beta-globin gene control region active in myeloid cells, myelin basic protein gene control region active in oligodendrocyte cells in the brain, myosin light chain-2 gene control region active in skeletal muscle, and gonadotropin releasing hormone gene control region active in the hypothalamus, pyruvate kinase promoter, villin promoter, promoter of the fatty acid binding intestinal protein, promoter of the smooth muscle cell α -actin, and the like. In addition, these expression sequences may be modified by addition of enhancer or regulatory sequences and the like.

[0084] Vectors comprising polynucleotides of the invention may be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., *J. Biol. Chem.* 267:963 (1992); Wu et al., *J. Biol. Chem.* 263: 14621 (1988); and Hartmut et al., Canadian Patent No. 2,012, 311).

[0085] Vectors and polynucleotides of the invention may be introduced in vivo by lipofection. For example, via use of liposomes for encapsulation and transfection of nucleic acids in vitro. Synthetic cationic lipids designed to limit the difficulties encountered with liposome-mediated transfection can be used to prepare liposomes for in vivo transfection of a gene encoding a marker (Feigner et al., *Proc. Natl. Acad. Sci. USA.* 84:7413 (1987); Mackey et al., *Proc. Natl. Acad. Sci. USA.* 85:8027 (1988); and Ulmer et al., *Science* 259:1745 (1993)). Use of cationic lipids may promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Feigner et al., *Science* 337:387 (1989)). Particularly useful lipid compounds and compositions for transfer of nucleic acids are described in WO95/18863, WO96/17823 and U.S. Pat. No. 5,459,127.

[0086] Other molecules are also useful for facilitating transfection of a nucleic acid in vivo, such as a cationic oligopeptide (e.g., WO95/21931), peptides derived from DNA binding proteins (e.g., WO96/25508), or a cationic polymer (e.g., WO95/21931).

[0087] It is also possible to introduce a vector in vivo as a naked DNA plasmid (see U.S. Pat. Nos. 5,693,622, 5,589,466 and 5,580,859). Receptor-mediated DNA delivery approaches can also be used (Curiel et al., *Hum. Gene Ther.* 3:147 (1992); and Wu et al., *J. Biol. Chem.* 262:4129 (1987)).

[0088] The term “transfection” refers to the uptake of exogenous or heterologous RNA or DNA by a cell. A cell has been “transfected” by exogenous or heterologous RNA or DNA when such RNA or DNA has been introduced inside the cell.

[0089] “Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” or “recombinant” or “transformed” organisms.

[0090] In addition, recombinant vector comprising polynucleotides of the invention may include one or more origins for replication in the cellular hosts in which their amplification or their expression is sought, markers or selectable markers.

[0091] The term “selectable marker” refers to an identifying factor, usually an antibiotic or chemical resistance gene, that is able to be selected for based upon the marker gene’s effect, i.e., resistance to an antibiotic, resistance to a herbicide, colorimetric markers, enzymes, fluorescent markers, and the like, wherein the effect is used to track the inheritance of a nucleic acid of interest and/or to identify a cell or organism that has inherited the nucleic acid of interest. Examples of selectable marker genes known and used in the art include: genes providing resistance to ampicillin, streptomycin, gentamycin, kanamycin, hygromycin, bialaphos herbicide, sulfonamide, and the like; and genes that are used as phenotypic markers, i.e., anthocyanin regulatory genes, isopentenyl transferase gene, and the like.

[0092] The term “reporter gene” refers to a nucleic acid encoding an identifying factor that is able to be identified based upon the reporter gene’s effect, wherein the effect is used to track the inheritance of a nucleic acid of interest, to identify a cell or organism that has inherited the nucleic acid of interest, and/or to measure gene expression induction or transcription. Examples of reporter genes known and used in the art include: luciferase (Luc), green fluorescent protein (GFP), chloramphenicol acetyltransferase (CAT), β -galactosidase (LacZ), β -glucuronidase (Gus), and the like. Selectable marker genes may also be considered reporter genes.

[0093] “Promoter and “promoter sequence” are used interchangeably and refer to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental or physiological conditions. Promoters that cause a gene to be expressed in most cell types at most times are commonly referred to as “constitutive promoters.” Promoters that cause a gene to be expressed in a specific cell type are commonly referred to as “cell-specific promoters” or “tissue-specific promoters.” Promoters that cause a gene to be expressed at a specific stage of development or cell differentiation are commonly referred to as “developmentally-specific promoters” or “cell differentiation-specific promoters.” Promoters that are induced and cause a gene to be expressed following exposure or treatment of the cell with an agent, biological molecule, chemical, ligand, light, or the like that induces the promoter are commonly referred to as “inducible promoters” or “regulatable promoters.” It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

[0094] The promoter sequence is typically bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease SI), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

[0095] A coding sequence is “under the control” of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced (if the coding sequence contains nitrons) and translated into the protein encoded by the coding sequence.

[0096] “Transcriptional and translational control sequences” refer to DNA regulatory sequences, such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

[0097] The term “response element” (“RE”) refers to one or more cis-acting DNA elements which confer responsiveness on a promoter mediated through interaction with the DNA-binding domains of a transcription factor. This DNA element may be either palindromic (perfect or imperfect) in its sequence or composed of sequence motifs or half sites separated by a variable number of nucleotides. The half sites can be similar or identical and arranged as either direct or inverted repeats or as a single half site or multimers of adjacent half sites in tandem. The response element may comprise a minimal promoter isolated from different organisms depending upon the nature of the cell or organism into which the response element will be incorporated. The DNA binding domain of the transcription factor binds, in the presence or absence of a ligand, to the DNA sequence of a response element to initiate or suppress transcription of downstream gene(s) under the regulation of this response element. Examples of DNA sequences for response elements of the natural ecdysone receptor include: RRGG/TTCANTGAC/ACY (SEQ ID NO:40) (see Cherbas et. al., *Genes Dev.* 5:120-131 (1991)); AGGTCAN(n)AGGTCA (SEQ ID NO:141), where N(n) can be one or more spacer nucleotides (see D'Avino et al., *Mol. Cell. Endocrinol.* 113:1 (1995)); and CGGTTGAATGAATT (SEQ ID NO:142) (see Antoniewski et al., *Mol. Cell Biol.* 14:4465 (1994)).

[0098] The terms “operably linked,” “operably associated,” “through operable association,” and the like refer to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

[0099] The terms “cassette,” “expression cassette” and “gene expression cassette” refer to a segment of DNA that can be inserted into a nucleic acid or polynucleotide at specific restriction sites or by homologous recombination. The segment of DNA comprises a polynucleotide that encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure insertion of the cassette in the proper reading frame for transcription and translation. “Transformation cassette” refers to a specific vector comprising a poly-

nucleotide that encodes a polypeptide of interest and having elements in addition to the polynucleotide that facilitate transformation of a particular host cell, Cassettes, expression cassettes, gene expression cassettes and transformation cassettes of the invention may also comprise elements that allow for enhanced expression of a polynucleotide encoding a polypeptide of interest in a host cell. These elements may include, but are not limited to: a promoter, a minimal promoter, an enhancer, a response element, a terminator sequence, a polyadenylation sequence, and the like.

[0100] For purposes of expressing polynucleotides and polypeptides under control of a gene switch mechanism, the term “gene switch” refers to the combination of a response element associated with a promoter, and a ligand-dependent transcription factor-based system which, in the presence of one or more ligands, modulates the expression of a gene into which the response element and promoter are incorporated. Stated otherwise, a “gene switch” refers to a peptide, protein or polypeptide complex that functions to (a) bind an activating ligand, and (b) regulate the transcription of a gene of interest in a ligand-dependent fashion.

[0101] As used herein with respect to gene switch regulation systems, the term “dimerizes with the ligand binding domain that binds an activating ligand” refers to a selective protein-protein interaction that is induced by the presence of activating ligand.

[0102] As used herein, the term “ligand binding domain that binds an activating ligand” refers to an amino acid sequence that selectively binds an activating ligand. In the methods disclosed herein, an activating ligand binds to a ligand binding domain, e.g., an ecdysone receptor ligand binding domain, that is part of a ligand-dependent transcriptional activation complex that regulates the expression of a polynucleotide sequence that encodes a gene of interest. Hence, the expression of the gene of interest is regulated in a ligand-dependent fashion.

[0103] The term “ecdysone receptor-based,” with respect to a gene switch, refers to a gene switch comprising at least a functional part of a naturally occurring or synthetic ecdysone receptor ligand binding domain and which regulates gene expression in response to a ligand that binds to the ecdysone receptor ligand binding domain.

[0104] The terms “modulate” and “modulates” mean to induce, reduce or inhibit nucleic acid or gene expression, resulting in the respective induction, reduction or inhibition of protein or polypeptide production.

[0105] Polynucleotides or vectors comprising sequences encoding polypeptides of the present invention may further comprise at least one promoter suitable for driving expression of a gene in a modified cell.

[0106] Enhancers that may be used in embodiments of the invention include but are not limited to: an SV40 enhancer, a cytomegalovirus (CMV) enhancer, an elongation factor 1 (EF1) enhancer, yeast enhancers, viral gene enhancers, et cetera.

[0107] “Regulatory region” refers to a nucleic acid sequence that regulates the expression of a second nucleic acid sequence. A regulatory region may include sequences which are naturally responsible for expressing a particular nucleic acid (a homologous region) or may include sequences of a different origin that are responsible for expressing different proteins or even synthetic proteins (a heterologous region). In particular, the sequences can be sequences of prokaryotic, eukaryotic, or viral genes or derived sequences

that stimulate or repress transcription of a gene in a specific or non-specific manner and in an inducible or non-inducible manner. Regulatory regions include origins of replication, RNA splice sites, promoters, enhancers, transcriptional termination sequences, and signal sequences which direct the polypeptide into the secretory pathways of the target cell.

[0108] The term “exogenous gene” or “heterologous gene” means a gene foreign to the subject or organism, that is, a gene which is introduced into the subject through a transformation process, an unmutated version of an endogenous mutated gene or a mutated version of an endogenous unmutated gene. The method of transformation is not critical to this invention and may be any method suitable for the subject known to those in the art. Exogenous genes can be either natural or synthetic genes which are introduced into the subject in the form of DNA or RNA which may function through a DNA intermediate such as by reverse transcriptase. Such genes can be introduced into target cells, directly introduced into the subject, or indirectly introduced by the transfer of transformed cells into the subject.

[0109] Polynucleotides and polypeptides of the invention may be expressed in vivo under control of a “gene switch” control mechanism, such as those described in, for example, but not limited to:

- [0110]** WO 2009/045370 (PCT/US2008/011270);
- [0111]** WO 2009/025866 (PCT/US2008/010040);
- [0112]** WO 2008/073154 (PCT/US2007/016747);
- [0113]** WO 2005/108617 (PCT/US2005/015089);
- [0114]** WO 2003/027289 (PCT/US2002/005026);
- [0115]** WO 2002/066615 (PCT/US2002/005708);
- [0116]** WO 2003/027266 (PCT/US2002/05234);
- [0117]** WO 2002/066612 (PCT/US2002/005090);
- [0118]** WO 2002/066614 (PCT/US/2002/005706);
- [0119]** WO 2002/066613 (PCT/US2002/005090);
- [0120]** WO 2002/029075 (PCT/US2001/030608); and
- [0121]** WO 2001/070816 (PCT/US2001/090500),

[0122] each of which are incorporated by reference herein.

[0123] The term “ligand-dependent transcription factor complex” or “LDTFC” refers to a transcription factor comprising one or more protein subunits, which complex can regulate gene expression driven by a “factor-regulated promoter” as defined herein. A model LDTFC is an “ecdysone receptor complex” generally refers to a heterodimeric protein complex having at least two members of the nuclear receptor family, ecdysone receptor (“EcR”) and ultraspiracle (“USP”) proteins (see Yao et al., *Nature* 366:476 (1993)); Yao et al., *Cell* 77:63 (1992)). A functional LDTFC such as an EcR complex may also include additional protein(s) such as immunophilins. Additional members of the nuclear receptor family of proteins, known as transcriptional factors (such as DHR38, betaFTZ-1 or other insect homologs), may also be ligand dependent or independent partners for EcR and/or USP. A LDTFC such as an EcR complex can also be a heterodimer of EcR protein and the vertebrate homolog of ultraspiracle protein, retinoic acid-X-receptor (“RXR”) protein or a chimera of USP and RXR. The terms “LDTFC” and “EcR complex” also encompass homodimer complexes of the EcR protein or USP, as well as single polypeptides or trimers, tetramer, and other multimers serving the same function.

[0124] A LDTFC such as an EcR complex can be activated by an active ecdysteroid or non-steroidal ligand bound to one of the proteins of the complex, inclusive of EcR, but not excluding other proteins of the complex. As used herein, the term “ligand,” as applied to LDTFC-based gene switches e.g.,

EcD complex based gene switches, describes small and soluble molecules having the capability of activating a gene switch to stimulate expression of a polypeptide encoded therein. Examples of ligands include, without limitation, an ecdysteroid, such as ecdysone, 20-hydroxyecdysone, ponasterone A, muristerone A, and the like, 9-cis-retinoic acid, synthetic analogs of retinoic acid, N,N'-diacylhydrazines such as those disclosed in U.S. Pat. Nos. 6,013,836; 5,117,057; 5,530,028; 5,378,726; and 7,304,161 and U.S. Pat. No. 7,456,315, oxadiazolines as described in U.S. Pat. No. 7,304,162; dibenzoylalkyl cyanohydrazines such as those disclosed in European Patent No. 461,809B1; N-alkyl-N,N'-diaroylhydrazines such as those disclosed in U.S. Pat. No. 5,225,443; N-acyl-N-alkylcarbonylhydrazines such as those disclosed in European Patent No. 234,994B1; N-aroyl-N-alkyl-N'-aroylhydrazines such as those described in U.S. Pat. No. 4,985,461; amidoketones such as those described in U.S. Pat. No. 7,375,093; each of which is incorporated herein by reference and other similar materials including 3,5-di-tert-butyl-4-hydroxy-N-isobutyl-benzamide, 8-O-acetylharpagide, oxysterols, 22(R) hydroxycholesterol, 24(S) hydroxycholesterol, 25-epoxycholesterol, T0901317, 5- α -6- α -epoxycholesterol-3-sulfate (ECHS), 7-ketocholesterol-3-sulfate, famesol, bile acids, 1,1-biphosphonate esters, juvenile hormone III, and the like. Examples of diacylhydrazine ligands useful in the present invention include RG-115819 (3,5-Dimethyl-benzoic acid N-(1-ethyl-2,2-dimethyl-propyl)-N'-(2-methyl-3-methoxy-benzoyl)-hydrazide), RG-L15932 (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-ethyl-3-methoxy-benzoyl)-hydrazide) and RG-115830 (3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-ethyl-3-methoxy-benzoyl) hydrazide). See, e.g., U.S. Pat. No. 8,076,517 (Publication No. 2009/0163592), and PCT Appl. No. PCT/US2008/006757 (WO 2008/153801), both of which are incorporated herein by reference in their entireties.

[0125] A LDFTC such as an EcR complex includes proteins which are members of the nuclear receptor superfamily wherein all members are characterized by the presence of one or more polypeptide subunits comprising an amino-terminal transactivation domain ("AD," "TD," or "TA," used interchangeably herein), a DNA binding domain ("DBD"), and a ligand binding domain ("LBD"). The AD may be present as a fusion with a "heterodimerization partner" or "HP." A fusion protein comprising an AD and HP of the invention is referred to herein as a "coactivation protein" or "CAP." The DBD and LBD may be expressed as a fusion protein, referred to herein as a "ligand-inducible transcription factor ("LTF"). The fusion partners may be separated by a linker, e.g., a hinge region. Some members of the LTF family may also have another transactivation domain on the carboxy-terminal side of the LBD. The DBD is characterized by the presence of two cysteine zinc fingers between which are two amino acid motifs, the P-box and the D-box, which confer specificity for ecdysone response elements. These domains may be either native, modified, or chimeras of different domains of heterologous receptor proteins.

[0126] EcR ligands, when used with a LDFTC, e.g., an EcR complex, which in turn is bound to the response element linked to an exogenous gene (e.g., a reporter gene), provide the means for external temporal regulation of expression of the exogenous gene. The order in which the various components bind to each other, that is, ligand to receptor complex and receptor complex to response element, is not critical. Typically, modulation of expression of the exogenous gene is

in response to the binding of a LDFTC, e.g., an EcR complex, to a specific control, or regulatory, DNA element. The EcR protein, like other members of the nuclear receptor family, possesses at least three domains, an AD, a DBD, and a LBD. This receptor, like a subset of the nuclear receptor family, also possesses less well-defined regions responsible for heterodimerization properties (referred to herein as a "heterodimerization partner" or "HP"). Binding of the ligand to the ligand binding domain of a LTF, e.g., an EcR protein, after heterodimerization with a CAP including, e.g., an AD and/or an HP, e.g., a USP or RXR protein, enables the DNA binding domains of the heterodimeric proteins to bind to the response element in an activated form, thus resulting in expression or suppression of the exogenous gene. This mechanism does not exclude the potential for ligand binding to individual subunits, e.g., LTF or CAP, e.g., an EcR or USP, and the resulting formation of active homodimer complexes (e.g. EcR+EcR or USP+USP). In one embodiment, one or more of the receptor domains can be varied producing a chimeric gene switch. Typically, one or more of the three domains may be chosen from a source different than the source of the other domains so that the chimeric receptor is optimized in the chosen host cell or organism for transactivating activity, complementary binding of the ligand, and recognition of a specific response element. In addition, the response element itself can be modified or substituted with response elements for other DNA binding protein domains such as the GAL-4 protein from yeast (see Sadowski et al., *Nature* 335:563 (1988)) or LexA protein from *E. coli* (see Brent et al., *Cell* 43:729-736 (1985)) to accommodate chimeric LDFTCs, e.g., EcR complexes. Another advantage of chimeric systems is that they allow choice of a promoter used to drive the exogenous gene according to a desired end result. Such double control can be particularly important in areas of gene therapy, especially when cytotoxic proteins are produced, because both the timing of expression as well as the cells wherein expression occurs can be controlled. When exogenous genes, operatively linked to a suitable promoter, are introduced into the cells of the subject, expression of the exogenous genes is controlled by the presence of the ligand of this invention. Promoters may be constitutively or inducibly regulated or may be tissue-specific (that is, expressed only in a particular type of cell) or specific to certain developmental stages of the organism.

[0127] For in vivo use, the ligands described herein may be taken up in pharmaceutically acceptable carriers, such as, for example, solutions, suspensions, tablets, capsules, ointments, elixirs, and injectable compositions. Pharmaceutical compositions may contain from 0.01% to 99% by weight of the ligand. Compositions may be either in single or multiple dose forms. The amount of ligand in any particular pharmaceutical composition will depend upon the effective dose, that is, the dose required to elicit the desired gene expression or suppression.

[0128] Suitable routes of administering the pharmaceutical preparations include oral, rectal, topical (including dermal, buccal and sublingual), vaginal, parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural) and by naso-gastric tube. It will be understood by those skilled in the art that the preferred route of administration will depend upon the condition being treated and may vary with factors such as the condition of the recipient.

[0129] As used herein, the terms "treat" or "treatment" refer to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow

down (lessen) an undesired physiological change or disorder, such as the development, progression or spread (i.e., metastasis) of cancer. Beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total). "Treatment" can also mean prolonging survival as compared to expected survival if not receiving treatment. Those in need of treatment include those already with the condition or disorder as well as those prone to have the condition or disorder or those in which the condition or disorder is to be prevented.

[0130] The terms "subject," "individual," "animal," "patient," or "mammal," is meant any subject, particularly a mammalian subject, for whom diagnosis, prognosis, or therapy is desired. Mammalian subjects include, without limitation, humans, domestic animals, farm animals, and zoo, sports, or pet animals such as dogs, cats, guinea pigs, rabbits, rats, mice, horses, cattle, cows, et cetera.

[0131] The terms "hyperproliferative disease or disorder" is intended to encompass all neoplastic cell growth and proliferation, whether malignant or benign, including all transformed cells and tissues and all cancerous cells and tissues. Hyperproliferative diseases or disorders include, but are not limited to, precancerous lesions, abnormal cell growths, tumors (whether benign or malignant), "cancer" and other hyperplasias.

[0132] The term "cancer" includes, but is not limited to, primary, malignant cells or tumors (e.g., those whose cells have not migrated to sites in the subject's body other than the site of the original malignancy or tumor) and secondary malignant cells or tumors (e.g., those arising from metastasis, the migration of malignant cells or tumor cells to secondary sites that are different from the site of the original tumor).

[0133] A tumor or tumor tissue may also comprise "tumor-associated non-tumor cells", e.g., vascular cells which form blood vessels to supply the tumor or tumor tissue. Non-tumor cells may be induced to replicate and develop by tumor cells, for example, the induction of angiogenesis in a #tumor or tumor tissue.

[0134] Some examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia or lymphoid malignancies. More particular examples of such cancers are noted below and include: squamous cell cancer (e.g. epithelial squamous cell cancer), lung cancer including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, rectal cancer, colorectal cancer, endometrial cancer or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, as well as head (e.g., brain) and neck cancer.

[0135] Other examples of cancers or malignancies include, but are not limited to: Acute Childhood Lymphoblastic Leukemia, Acute Lymphoblastic Leukemia, Acute Lymphocytic Leukemia, Acute Myeloid Leukemia, Adrenocortical Carcinoma, Adult (Primary) Hepatocellular Cancer, Adult (Primary) Liver Cancer, Adult Acute Lymphocytic Leukemia, Adult Acute Myeloid Leukemia, Adult Hodgkin's Disease, Adult Hodgkin's Lymphoma, Adult Lymphocytic Leukemia,

Adult Non-Hodgkin's Lymphoma, Adult Primary Liver Cancer, Adult Soft Tissue Sarcoma, AIDS-Related Lymphoma, AIDS-Related Malignancies, Anal Cancer, Astrocytoma, Bile Duct Cancer, Bladder Cancer, Bone Cancer, Brain Stem Glioma, Brain Tumors, Breast Cancer, Cancer of the Renal Pelvis and Ureter, Central Nervous System (Primary) Lymphoma, Central Nervous System Lymphoma, Cerebellar Astrocytoma, Cerebral Astrocytoma, Cervical Cancer, Childhood (Primary) Hepatocellular Cancer, Childhood (Primary) Liver Cancer, Childhood Acute Lymphoblastic Leukemia, Childhood Acute Myeloid Leukemia, Childhood Brain Stem Glioma, Childhood Cerebellar Astrocytoma, Childhood Cerebral Astrocytoma, Childhood Extracranial Germ Cell Tumors, Childhood Hodgkin's Disease, Childhood Hodgkin's Lymphoma, Childhood Hypothalamic and Visual Pathway Glioma, Childhood Lymphoblastic Leukemia, Childhood Medulloblastoma, Childhood Non-Hodgkin's Lymphoma, Childhood Pineal and Supratentorial Primitive Neuroectodermal Tumors, Childhood Primary Liver Cancer, Childhood Rhabdomyosarcoma, Childhood Soft Tissue Sarcoma, Childhood Visual Pathway and Hypothalamic Glioma, Chronic Lymphocytic Leukemia, Chronic Myelogenous Leukemia, Colon Cancer, Cutaneous T cell Lymphoma, Endocrine Pancreas Islet Cell Carcinoma, Endometrial Cancer, Ependymoma, Epithelial Cancer, Esophageal Cancer, Ewing's Sarcoma and Related Tumors, Exocrine Pancreatic Cancer, Extracranial Germ Cell Tumor, Extragonadal Germ Cell Tumor, Extrahepatic Bile Duct Cancer, Eye Cancer, Female Breast Cancer, Gaucher's Disease, Gallbladder Cancer, Gastric Cancer, Gastrointestinal Carcinoid Tumor, Gastrointestinal Tumors, Germ Cell Tumors, Gestational Trophoblastic Tumor, Hairy Cell Leukemia, Head and Neck Cancer, Hepatocellular Cancer, Hodgkin's Disease, Hodgkin's Lymphoma, Hypergammaglobulinemia, Hypopharyngeal Cancer, Intestinal Cancers, Intraocular Melanoma, Islet Cell Carcinoma, Islet Cell Pancreatic Cancer, Kaposi's Sarcoma, Kidney Cancer, Laryngeal Cancer, Lip and Oral Cavity Cancer, Liver Cancer, Lung Cancer, Lymphoproliferative Disorders, Macroglobulinemia, Male Breast Cancer, Malignant Mesothelioma, Malignant Thymoma, Medulloblastoma, Melanoma, Mesothelioma, Metastatic Occult Primary Squamous Neck Cancer, Metastatic Primary Squamous Neck Cancer, Metastatic Squamous Neck Cancer, Multiple Myeloma, Multiple Myeloma/Plasma Cell Neoplasm, Myelodysplastic Syndrome, Myelogenous Leukemia, Myeloid Leukemia, Myeloproliferative Disorders, Nasal Cavity and Paranasal Sinus Cancer, Nasopharyngeal Cancer, Neuroblastoma, Non-Hodgkin's Lymphoma During Pregnancy, Nonmelanoma Skin Cancer, Non-Small Cell Lung Cancer, Occult Primary Metastatic Squamous Neck Cancer, Oropharyngeal Cancer, Osteo-/Malignant Fibrous Sarcoma, Osteosarcoma/Malignant Fibrous Histiocytoma, Osteosarcoma/Malignant Fibrous Histiocytoma of Bone, Ovarian Epithelial Cancer, Ovarian Germ Cell Tumor, Ovarian Low Malignant Potential Tumor, Pancreatic Cancer, Paraproteinemias, Purpura, Parathyroid Cancer, Pheochromocytoma, Pituitary Tumor, Plasma Cell Neoplasm/Multiple Myeloma, Primary Central Nervous System Lymphoma, Primary Liver Cancer, Prostate Cancer, Rectal Cancer, Renal Cell Cancer, Renal Pelvis and Ureter Cancer, Retinoblastoma, Rhabdomyosarcoma, Salivary Gland Cancer, Sarcoidosis Sarcomas, Sezary Syndrome, Skin Cancer, Small Cell Lung Cancer, Small Intestine Cancer, Soft Tissue Sarcoma, Squamous Neck Cancer, Stomach Cancer, Supratentorial Primitive Neuroectodermal and

Pineal Tumors, T cell Lymphoma, Testicular Cancer, Thyoma, Thyroid Cancer, Transitional Cell Cancer of the Renal Pelvis and Ureter, Transitional Renal Pelvis and Ureter Cancer, Trophoblastic Tumors, Ureter and Renal Pelvis Cell Cancer, Urethral Cancer, Uterine Cancer, Uterine Sarcoma, Vaginal Cancer, Visual Pathway and Hypothalamic Glioma, Vulvar Cancer, Waldenstrom's Macroglobulinemia, Wilms' Tumor, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

[0136] Naturally Occurring Amino Acid Substitutions

[0137] List of naturally occurring amino acids and some of their biochemical properties.

Amino Acid	3-Letter Code	1-Letter Code	Side-chain polarity*	Side-chain charge (pH 7.4)*	Hydropathy index**
Alanine	Ala	A	nonpolar	neutral	1.8
Arginine	Arg	R	polar	positive	-4.5
Asparagine	Asn	N	polar	neutral	-3.5
Aspartic acid	Asp	D	polar	negative	-3.5
Cysteine	Cys	C	polar	neutral	2.5
Glutamic acid	Glu	E	polar	negative	-3.5
Glutamine	Gln	Q	polar	neutral	-3.5
Glycine	Gly	G	nonpolar	neutral	-0.4
Histidine	His	H	polar	positive(10%) neutral(90%)	-3.2
Isoleucine	Ile	I	nonpolar	neutral	4.5
Leucine	Leu	L	nonpolar	neutral	3.8
Lysine	Lys	K	polar	positive	-3.9
Methionine	Met	M	nonpolar	neutral	1.9
Phenylalanine	Phe	F	nonpolar	neutral	2.8
Proline	Pro	P	nonpolar	neutral	-1.6
Serine	Ser	S	polar	neutral	-0.8
Threonine	Thr	T	polar	neutral	-0.7
Tryptophan	Trp	W	nonpolar	neutral	-0.9
Tyrosine	Tyr	Y	polar	neutral	-1.3
Valine	Val	V	nonpolar	neutral	4.2

*Hausman & Cooper, (2004), *The Cell: A Molecular Approach*, Washington, D.C: ASM Press, p. 51 (2004)(ISBN 0-87893-214-3).

**Kyte & Doolittle, "A simple method for displaying the hydropathic character of a protein," *Journal of Molecular Biology*, 157(1): 105-132 (May 1982).

[0138] Conservative Amino Acid Substitutions

[0139] Polypeptides may be made to differ by introduction of conservative or non-conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar amino acid side chains. "Conservative amino acid substitutions" refer to substitutions of one or more amino acids in a native amino acid sequence (e.g., wild-type or naturally occurring form of PE) with other amino acid(s) having similar side chains (e.g., side chains similar in terms of size, charge, element composition, and/or hydrophobicity/hydrophilicity).

[0140] Conserved substitutes for an amino acid within a native amino acid sequence can be selected from other members of the group to which the naturally occurring amino acid belongs. For example, conservative amino acid residue substitution groups include:

[0141] (1) Alanine (A)-Glycine (G)-Serine (S)-Threonine;

[0142] (2) Aspartic acid (D)-Glutamic acid (E);

[0143] (3) Asparagine (N)-Glutamine (Q);

[0144] (4) Arginine (R)-Lysine (K)-Histidine (H);

[0145] (5) Isoleucine (I)-Leucine (L)-Methionine (M)-Valine (V); and

[0146] (6) Phenylalanine (F)-Tyrosine (Y)-Tryptophan (W).

[0147] Other substitution groups of amino acids can be envisioned. For example, amino acids can be grouped by similar function or chemical structure or composition (e.g., acidic, basic, aliphatic, aromatic, sulfur-containing). For example, an Aliphatic grouping may comprise: Glycine (G), Alanine (A), Valine (V), Leucine (L), Isoleucine (I). Other groups containing amino acids that are considered conservative substitutions for one another include:

[0148] Aromatic: Phenylalanine (F)-Tyrosine (Y)-Tryptophan (W);

[0149] Sulfur-containing: Methionine (M)-Cysteine (C);

[0150] Basic: Arginine (R)-Lysine (K)-Histidine (H);

[0151] Acidic: Aspartic acid (D)-Glutamic acid (E);

[0152] Non-polar uncharged residues: Cysteine (C)-Methionine (M)-Proline (P); and

[0153] Hydrophilic Uncharged Residues: Serine (S)-Threonine (T)-Asparagine (N)-Glutamine (Q).

[0154] Exemplary embodiments of conservative amino acid substitutions include the interchangeability of: valine-leucine, valine-isoleucine-leucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

[0155] Examples of Amino Acid Analogs and Non-Standard Amino Acid Residues

[0156] Examples of a few of the many possible amino acid analogs routinely known to those of skill in the art include, for example, but without limitation, analogs such as: 4-hydroxyproline which may be substituted for proline; 5-hydroxylysine which may be substituted for lysine; 3-methylhistidine which may be substituted for histidine; homoserine which may be substituted for serine; and ornithine which may be substituted for lysine.

[0157] Examples of a few of the many possible non-standard amino acids routinely known to those of skill in the art include, for example, but without limitation, molecules such as: ornithine, citrulline, lanthionine, 2-aminoisobutyric acid, dehydroalanine, γ -aminobutyric acid, β -alanine (3-amino-propanoic acid), selenocysteine and pyrrolysine.

[0158] Substitution mutations may be made by any technique for mutagenesis known in the art including, for example, but not limited to, in vitro site-directed mutagenesis (Hutchinson et al, *J. Biol. Chem.* 255:6551 (1978); Zoller et al, *DNA* 3:479 (1984); Oliphant et al, *Gene* 44:177 (1986); Hutchinson et al, *Proc. Natl. Aca. Sci. USA* 83:710 (1986)), use of TAB® linkers (Pharmacia), restriction endonuclease digestion/fragment deletion and substitution, PCR-mediated/oligonucleotide-directed mutagenesis, et cetera. PCR-based techniques are preferred for site-directed mutagenesis (see Higuchi, 1989, "Using PCR to Engineer DNA", in PCR. Technology: Principles and Applications for DNA Amplification, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70).

Embodiments of the Invention

[0159] Embodiments of the invention include isolated polypeptides (proteins) comprising or consisting of a modified form of *Pseudomonas* exotoxin A, or a fragment thereof wherein said modified form, or fragment thereof, comprises an epitope selected from the group consisting of:

ISFSTRGTQ; (epitope 1; SEQ ID NO: 5)

GTQNWTVR; (epitope 2; SEQ ID NO: 6)

-continued

IVFGGVRAR;	(epitope 3; SEQ ID NO: 7)
ARSQDLDAI;	(epitope 4; SEQ ID NO: 8)
LRVYVPRSS; and	(epitope 5; SEQ ID NO: 9)
IPDKEQAIS	(epitope 6; SEQ ID NO: 10)

[0160] wherein one or more amino acid residues in any one or more of these epitopes are substituted with a different amino acid residue.

[0161] Embodiments of the invention include isolated polypeptides (proteins) comprising or consisting of a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form, or fragment thereof, comprises an epitope selected from the group consisting of:

GDGGDISFSTRGTQN	(peptide 50 (epitope 1);;	SEQ ID NO: 60)
SFSTRGTQNWTVRL	(peptide 52 (epitope 2);;	SEQ ID NO: 62)
TRGTQNWTVRLLQA	(peptide 53 (epitope 2);;	SEQ ID NO: 63)
AQSIVFGGVRARSQD	(peptide 65 (epitope 3);;	SEQ ID NO: 75)
GGVRARSQDLDAIWR	(peptide 67 (epitope 4);;	SEQ ID NO: 77)
RARSQDLDAIWRGFY	(peptide 68 (epitope 4);;	SEQ ID NO: 78)
NGALLRVYVPRSSLP	(peptide 81 (epitope 5);;	SEQ ID NO: 91)
LLRVYVPRSSLPGFY	(peptide 82 (epitope 5);;	SEQ ID NO: 92)
LDPSSIPDKEQAISA	(peptide 110 (epitope 6);;	SEQ ID NO: 120)
SSIPDKEQAISALPD	(peptide 111 (epitope 6);;	SEQ ID NO: 121)
ISFSTRGTQNWTVR	(overlapping epitopes 1 and 2;;	SEQ ID NO: 131)
IVFGGVRARSQDLDAI	(overlapping epitopes 3 and 4;;	SEQ ID NO: 132)

[0162] wherein one or more amino acid residues in any one or more of these epitopes are substituted with a different amino acid residue.

[0163] Embodiments of the invention include isolated polypeptides (proteins) comprising or consisting of a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form, or fragment thereof comprises an epitope selected from the group consisting of

[0164] a) ISESTRGTQ (SEQ ID NO:5), wherein amino acid residues at one or more of positions 1, 6 and 9 are substituted with a different amino acid residue;

[0165] b) GTQNWTVR (SEQ ID NO:6), wherein amino acid residues at one or more of positions 3, 4 and 6 are substituted with a different amino acid residue;

[0166] c) IVFGGVRAR (SEQ ID NO:7), wherein amino acid residues at one or more of positions 1 and 6 are substituted with a different amino acid residue;

[0167] d) ARSQDLDAI (SEQ ID NO:8), wherein amino acid residues at one or more of positions 4 and 7 are substituted with a different amino acid residue;

[0168] e) LRVYVPRSS (SEQ ID NO:9), wherein amino acid residues at one or more of positions 1, 2 and 9 are substituted with a different amino acid residue;

[0169] f) IPDKEQAIS (SEQ ID NO:10), wherein amino acid residues at one or more of positions 1, 4, 6 and 7 are substituted with a different amino acid residue;

[0170] g) ISFSTRGTQNWTVR (SEQ ID NO:131), wherein amino acid residues at one or more of positions 1, 6, 9, 10 and 12 are substituted with a different amino acid residue; and

[0171] h) IVFGGVRARSQDLDAI (SEQ ID NO:132), wherein amino acid residues at one or more of positions 1, 6, 11, and 14 are substituted with a different amino acid residue.

[0172] Embodiments of the invention include isolated polypeptides (proteins) comprising or consisting of a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form, or fragment thereof, comprises an epitope selected from the group consisting of

[0173] a) ISFSTRGTQ (SEQ ID NO:5), wherein amino acid residues at one or more of positions 1, 6 and 9 are substituted with a conservative amino acid substitution;

[0174] b) GTQNWTVR (SEQ ID NO:6), wherein amino acid residues at one or more of positions 3, 4 and 6 are substituted with a conservative amino acid substitution;

[0175] c) IVFGGVRAR (SEQ ID NO:7), wherein amino acid residues at one or more of positions 1 and 6 are substituted with a conservative amino acid substitution;

[0176] d) ARSQDLDAI (SEQ ID NO:8), wherein amino acid residues at one or more of positions 4 and 7 are substituted with a conservative amino acid substitution;

[0177] e) LRVYVPRSS (SEQ ID NO:9), wherein amino acid residues at one or more of positions 1, 2 and 9 are substituted with a conservative amino acid substitution;

[0178] f) IPDKEQAIS (SEQ ID NO:10), wherein amino acid residues at one or more of positions 1, 4, 6 and 7 are substituted with a conservative amino acid substitution;

[0179] g) ISFSTRGTQNWTVR (SEQ ID NO:131), wherein amino acid residues at one or more of positions 1, 6, 9, 10 and 12 are substituted with a conservative amino acid substitution; and

[0180] h) IVFGGVRARSQDLDAI (SEQ ID NO:132), wherein amino acid residues at one or more of positions 1, 6, 11, and 14 are substituted with a conservative amino acid substitution.

[0181] Embodiments of the invention include isolated polypeptides (proteins) comprising or consisting of a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form, or fragment thereof, comprises an epitope selected from the group consisting of:

[0182] a) ISFSTRGTQ (SEQ ID NO:5), wherein amino acid residues at one or more of positions 1, 6 and 9 are substituted with a conservative amino acid substitution;

[0183] b) CTQNWTVR (SEQ ID NO:6), wherein amino acid residues at one or more of positions 3, 4 and 6 are substituted with a conservative amino acid substitution;

- [0184] c) IVEGGAVRAR (SEQ ID NO:7), wherein amino acid residues at one or more of positions 1 and 6 are substituted with a conservative amino acid substitution;
- [0185] d) ARSQDLDAI (SEQ ID NO:8), wherein amino acid residues at one or more of positions 4 and 7 are substituted with a conservative amino acid substitution;
- [0186] e) LRVYVPRSS (SEQ ID NO:9), wherein amino acid residues at one or more of positions 1, 2 and 9 are substituted with a conservative amino acid substitution;
- [0187] f) IPDKEQAIS (SEQ ID NO:10), wherein amino acid residues at one or more of positions 1, 4, 6 and 7 are substituted with a conservative amino acid substitution;
- [0188] g) ISFSTRGTQNWTVR (SEQ ID NO:131), wherein amino acid residues at one or more of positions 1, 6, 9, 10 and 12 are substituted with a conservative amino acid substitution; and
- [0189] h) ITVFGGVRARSQDLDAI (SEQ ID NO:132) wherein amino acid residues at one or more of positions 1, 6, 1, and 14 are substituted with a conservative amino acid substitution.
- [0190] wherein the conservative amino acid substitution at one or more of said positions in a) through f) is selected from the group consisting of:
- [0191] 1) A is substituted with any one of G, I, L, S, T or V;
 - [0192] 2) D is substituted with E;
 - [0193] 3) I is substituted with any one of L, M or V;
 - [0194] 4) K is substituted with any one of H or R;
 - [0195] 5) L is substituted with any one of A, G, I, M or V;
 - [0196] 6) N is substituted with any one of S, T or Q;
 - [0197] 7) Q is substituted with any one of S, T or N;
 - [0198] 8) R is substituted with any one of K or H;
 - [0199] 9) S is substituted with any one of A, G, N, T or Q;
 - [0200] 10) T is substituted with any one of A, G, N, Q or S; and
 - [0201] 11) V is substituted with any one of A, G, I, L or M;
- [0202] Embodiments of the invention also comprise or consist of isolated poly-peptides and peptides comprising or consisting of the above-referenced amino acids sequences, except wherein one or more amino acids have been substituted with conservative amino acids substitutions. Embodiments of the invention further comprise or consist of isolated polypeptides (proteins) and peptides comprising or consisting of the above-referenced amino acids sequences, except wherein one or more amino acids have been substituted with amino acids which are naturally occurring non-occurring, non-natural/occurring, non-standard amino acids, or amino acid analogs.
- [0203] Embodiments of the invention include isolated polypeptides (proteins) comprising or consisting of a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form, or fragment thereof, comprises an epitope selected from the group consisting of:
- [0204] a) ISFSTRGTQ (SEQ ID NO:5), wherein amino the acid residue at position 1 (I) is substituted with A, N, T, Q or H, or wherein the amino acid residue at position 6 (R) is substituted with Q, or wherein the amino acid residue at position 9 (Q) is substituted with N or T, or wherein the amino acid sequence ISSFTRGTQ (SEQ ID NO:5) comprises two or more of said substitutions in any combination;
 - [0205] b) GTQNWTVR (SEQ ID NO:6), wherein the amino acid residue at position 3 (Q) is substituted with N or T, wherein amino the acid residue at position 4 (N) is substituted with K or R, or wherein the amino acid residue at position 6 (T) is substituted with K or R, or wherein the amino acid sequence GTQNWTVR (SEQ ID NO:6) comprises two or more of said substitutions in any combination;
 - [0206] c) IVFGGVRAR (SEQ ID NO:7), wherein amino the acid residue at position 1 (I) is substituted with A or N, or wherein the amino acid residue at position 6 (V) is substituted with D, M, or N, or wherein the amino acid sequence IVFGGVRAR (SEQ ID NO:7) comprises substitutions at both positions in any combination of amino acid residues A or N at position 1 (I) and D, M, or N at position 6 (V);
 - [0207] d) ARSQDLDAI (SEQ ID NO:8), wherein amino the acid residue at position 4 (Q) is substituted with K or R, or wherein the amino acid residue at position 7 (D) is substituted with K or R, or wherein the amino acid sequence ARSQDLDAI (SEQ ID NO:8) comprises substitutions with K or R in any combination at both positions 4 (Q) and 7 (D);
 - [0208] e) LRVYVPRSS (SEQ ID NO:9), wherein amino the acid residue at position 1 (L) is substituted with A, or wherein the amino acid residue at position 2 (R) is substituted with D, S or A, or wherein the amino acid residue at position 9 (S) is substituted with D, E, N, K, P or T, or wherein the amino acid sequence LRVYVPRSS (SEQ ID NO:9) comprises two or more of said substitutions in any combination;
 - [0209] f) IPDKEQAIS (SEQ ID NO:10), wherein amino acid residues at one or more of positions 1, 4, 6 and 7 are substituted with a different amino acid residue, wherein amino the acid residue at position 1 (I) is substituted with A, N, T, Q or H, or wherein the amino acid residue at position 4 (K) is substituted with T, or wherein, the amino acid residue at position 6 (Q) is substituted with D or wherein the amino acid residue at position 7 (A) is substituted with D, or wherein the amino acid sequence IPDKEQAIS (SEQ ID NO:10) comprises two or more of said substitutions in any combination;
 - [0210] g) ISFSTRGTQNWTVR (SEQ ID NO:131), wherein amino acid residues at one or more of positions 1, 6, 9, 10 and 12 are substituted with a different amino acid residues wherein amino the acid residue at position 1 (I) is substituted with A, N, T, Q or H, or wherein the amino acid residue at position 6 (R) is substituted with Q, or wherein the amino acid residue at position 9 (Q) is substituted with N or T, or wherein amino the acid residue at position 10 (N) is substituted with K or R, or wherein the amino acid residue at position 12 (T) is substituted with K or R, or wherein the amino acid sequence ISFSTRGTQNWTVR (SEQ ID NO:131) comprises two or more of said substitutions in any combination; and
 - [0211] h) IVFGGVRARSQDLDAI (SEQ ID NO:132), wherein amino the acid residue at position 1 (I) is substituted with A or N, or wherein the amino acid residue at position 6 (V) is substituted with D, M, or N, wherein amino the acid residue at position 11 (Q) is substituted with K or R, or wherein the amino acid residue at position 14 (D) is substituted with K, or R, or wherein the amino acid sequence IVFGGVRARSQDLDAI (SEQ ID NO:132) comprises two or more of said substitutions in any combination.

[0212] Embodiments of the invention include an isolated polypeptide comprising a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form or fragment thereof, comprises one or more amino acid substitutions selected from the group consisting of:

[0213] a) I at position 141 changed to any amino acid residue; (epitope 1)

[0214] b) R at position 146 changed to any amino acid residue; (epitope 1)

[0215] c) Q at position 149 changed to any amino acid residue; (epitope 1)

[0216] d) N at position 150 changed to any amino acid residue; (epitope 2)

[0217] e) T at position 152 changed to any amino acid residue; (epitope 2)

[0218] f) I at position 184 changed to any amino acid residue; (epitope 3)

[0219] g) V at position 189 changed to any amino acid residue; (epitope 3)

[0220] h) Q at position 194 changed to any amino acid residue; (epitope 4)

[0221] i) D at position 197 changed to any amino acid residue; (epitope 4)

[0222] j) L at position 233 changed to any amino acid residue; (epitope 5)

[0223] k) R at position 234 changed to any amino acid residue; (epitope 5)

[0224] l) S at position 241 changed to any amino acid residue; (epitope 5)

[0225] m) I at position 321 changed to any amino acid residue; (epitope 6)

[0226] n) K at position 324 changed to any amino acid residue; (epitope 6)

[0227] o) Q at position 326 changed to any amino acid residue; (epitope 6)

[0228] p) A at position 327 changed to any amino acid residue; (epitope 6)

[0229] q) any combination of one or more of a) through ao),

[0230] wherein the amino acid numbering corresponds to SEQ ID NO:1.

[0231] Embodiments of the invention include an isolated polypeptide comprising a modified form of *Pseudomonas* exotoxin A, or a fragment thereof, wherein said modified form, or fragment thereof, comprises one or more amino acid substitutions selected from the group consisting of:

[0232] a) I at position 141 changed to A; (epitope 1)

[0233] b) I at position 141 changed to N; (epitope 1)

[0234] c) I at position 141 changed to T; (epitope 1)

[0235] d) I at position 141 changed to Q; (epitope 1)

[0236] e) I at position 141 changed to H; (epitope 1)

[0237] f) R at position 146 changed to Q; (epitope 1)

[0238] g) Q at position 149 changed to N; (epitope 1)

[0239] h) Q at position 149 changed to T; (epitope 1)

[0240] i) N at position 150 changed to R; (epitope 2)

[0241] j) N at position 150 changed to K; (epitope 2)

[0242] k) T at position 152 changed to R; (epitope 2)

[0243] l) T at position 152 changed to K; (epitope 2)

[0244] m) I at position 184 changed to A; (epitope 3)

[0245] n) I at position 184 changed to N; (epitope 3)

[0246] o) V at position 189 changed to D; (epitope 3)

[0247] p) V at position 189 changed to M; (epitope 3)

[0248] q) V at position 189 changed to N; (epitope 3)

[0249] r) Q at position 194 changed to R; (epitope 4)

[0250] s) Q at position 194 changed to K; (epitope 4)

[0251] t) D at position 197 changed to R; (epitope 4)

[0252] u) D at position 197 changed to K; (epitope 4)

[0253] v) L at position 233 changed to A; (epitope 5)

[0254] w) R at position 234 changed to D; (epitope 5)

[0255] x) R at position 234 changed to S; (epitope 5)

[0256] y) R at position 234 changed to A; (epitope 5)

[0257] z) S at position 241 changed to D; (epitope 5)

[0258] ab) S at position 241 changed to E; (epitope 5)

[0259] ac) S at position 241 changed to N; (epitope 5)

[0260] ad) S at position 241 changed to K; (epitope 5)

[0261] ae) S at position 241 changed to P; (epitope 5)

[0262] af) S at position 241 changed to T; (epitope 5)

[0263] ag) I at position 321 changed to A; (epitope 6)

[0264] ah) I at position 321 changed to N; (epitope 6)

[0265] ai) I at position 321 changed to T; (epitope 6)

[0266] ak) I at position 321 changed to Q; (epitope 6)

[0267] al) I at position 321 changed to H; (epitope 6)

[0268] am) K at position 324 changed to T; (epitope 6)

[0269] an) Q at position 326 changed to D; (epitope 6)

[0270] ao) A at position 327 changed to D; (epitope 6)

[0271] ap) any combination of one or more of a) through ao),

[0272] wherein the amino acid numbering corresponds to SEQ ID NO:1.

[0273] Embodiments of the invention comprise isolated polypeptides as described above, including polypeptides comprising amino acid substitutions introduced at each of amino acid positions 141, 146, 149, 150, 152, 184, 189, 194, 197, 233, 234, 241, 321, 324, 326 and 327 (in comparison to the amino acid sequence of SEQ ID NO:1).

[0274] Embodiments of the invention include isolated polypeptides (proteins) and peptides comprising, or consisting of, the following amino acid sequences:

GGGGGSGGGGSPGEG;	(peptide 1; SEQ ID NO: 11)
GGSGGGGSGGGSLS;	(peptide 2; SEQ ID NO: 12)
GGGGGSGGGSLSAAL;	(peptide 3; SEQ ID NO: 13)
GGSPGGSLSAALTAH;	(peptide 4; SEQ ID NO: 14)
PEGGSLSAALTAHQAC;	(peptide 5; SEQ ID NO: 15)
GSLAALTAHQACHLP;	(peptide 6; SEQ ID NO: 16)
AALTAHQACHLPLET;	(peptide 7; SEQ ID NO: 17)
TAHQACHLPLETFTTR;	(peptide 8; SEQ ID NO: 18)
QACHLPLETFTTRHRQ;	(peptide 9; SEQ ID NO: 19)
HLPLETFTTRHRQPRG;	(peptide 10; SEQ ID NO: 20)
LETFTTRHRQPRGWEQ;	(peptide 11; SEQ ID NO: 21)
FTRHRQPRGWEQLEQ;	(peptide 12; SEQ ID NO: 22)
HRQPRGWEQLEQCGY;	(peptide 13; SEQ ID NO: 23)
PRGWEQLEQCGYPVQ;	(peptide 14; SEQ ID NO: 24)
WEQLEQCGYPVQRLV;	(peptide 15; SEQ ID NO: 25)
LEQCGYPVQRLVALY;	(peptide 16; SEQ ID NO: 26)
CGYPVQRLVALYLAA;	(peptide 17; SEQ ID NO: 27)
PVQRLVALYLAAARLS;	(peptide 18; SEQ ID NO: 28)

-continued

RLVALYLAARLSWNQ; (peptide 19; SEQ ID NO: 29)
 ALYLAARLSWNQVDQ; (peptide 20; SEQ ID NO: 30)
 LAARLSWNQVDQVIR; (peptide 21; SEQ ID NO: 31)
 RLSWNQVDQVIRNAL; (peptide 22; SEQ ID NO: 32)
 WNQVDQVIRNALASP; (peptide 23; SEQ ID NO: 33)
 VDQVIRNALASPGSG; (peptide 24; SEQ ID NO: 34)
 VIRNALASPGSGGDL; (peptide 25; SEQ ID NO: 35)
 NALASPGSGGDLGEA; (peptide 26; SEQ ID NO: 36)
 ASPGSGGDLGEAIRE; (peptide 27; SEQ ID NO: 37)
 GSGGDLGEAIREQPE; (peptide 28; SEQ ID NO: 38)
 GDLGEAIREQPEQAR; (peptide 29; SEQ ID NO: 39)
 GEAIREQPEQARLAL; (peptide 30; SEQ ID NO: 40)
 IREQPEQARLALTLA; (peptide 31; SEQ ID NO: 41)
 QPEQARLALTLAAAE; (peptide 32; SEQ ID NO: 42)
 QARLALTLAAAESER; (peptide 33; SEQ ID NO: 43)
 LALTLAAAESERFVR; (peptide 34; SEQ ID NO: 44)
 TLAAAESERFVRQGT; (peptide 35; SEQ ID NO: 45)
 AAESERFVRQGTGND; (peptide 36; SEQ ID NO: 46)
 SERFVRQGTGNDDEAG; (peptide 37; SEQ ID NO: 47)
 FVRQGTGNDDEAGAAS; (peptide 38; SEQ ID NO: 48)
 QGTGNDDEAGAASGPA; (peptide 39; SEQ ID NO: 49)
 GNDDEAGAASGPADSG; (peptide 40; SEQ ID NO: 50)
 EAGAASGPADSGDAL; (peptide 41; SEQ ID NO: 51)
 AASGPADSGDALLER; (peptide 42; SEQ ID NO: 52)
 GPADSGDALLERNYP; (peptide 43; SEQ ID NO: 53)
 DSGDALLERNYPTGA; (peptide 44; SEQ ID NO: 54)
 DALLERNYPTGAEFL; (peptide 45; SEQ ID NO: 55)
 LERNYPTGAEFLGDG; (peptide 46; SEQ ID NO: 56)
 NYPTGAEFLGDGGDI; (peptide 47; SEQ ID NO: 57)
 TGAEFLGDGGDISFS; (peptide 48; SEQ ID NO: 58)
 EFLGDGGDISFSTRG; (peptide 49; SEQ ID NO: 59)
 GDISFSTRGTQNWTV; (peptide 51; SEQ ID NO: 61)
 TQNWTVRLLQAHQR; (peptide 54; SEQ ID NO: 64)
 WTVRLLQAHQRLEE; (peptide 55; SEQ ID NO: 65)
 ERLQAHQRLEERGY; (peptide 56; SEQ ID NO: 66)
 LQAHQRLEERGYVVF; (peptide 57; SEQ ID NO: 67)
 HRQLEERGYVFGYH; (peptide 58; SEQ ID NO: 68)
 LEERGYVFGYHGTF; (peptide 59; SEQ ID NO: 69)
 RGYVFGYHGTFLEA; (peptide 60; SEQ ID NO: 70)

-continued

VFGYHGTFLEAAQS; (peptide 61; SEQ ID NO: 71)
 GYHGTFLEAAQSIVF; (peptide 62; SEQ ID NO: 72)
 GTFLEAAQSIVFGGV; (peptide 63; SEQ ID NO: 73)
 LEAAQSIVFGGVRAR; (peptide 64; SEQ ID NO: 74)
 IVFGGVRARSQDLDA; (peptide 66; SEQ ID NO: 76)
 SQDLDAIWRGFYIAG; (peptide 69; SEQ ID NO: 79)
 LDAIWRGFYIAGDPA; (peptide 70; SEQ ID NO: 80)
 IWRGFYIAGDPALAY; (peptide 71; SEQ ID NO: 81)
 GFYIAGDPALAYGYA; (peptide 72; SEQ ID NO: 82)
 IAGDPALAYGYAQDQ; (peptide 73; SEQ ID NO: 83)
 DPALAYGYAQDQEPD; (peptide 74; SEQ ID NO: 84)
 LAYGYAQDQEPDARG; (peptide 75; SEQ ID NO: 85)
 GYAQDQEPDARGRIR; (peptide 76; SEQ ID NO: 86)
 QDQEPDARGRIRNGA; (peptide 77; SEQ ID NO: 87)
 EPDARGRIRNGALLR; (peptide 78; SEQ ID NO: 88)
 ARGRIRNGALLRVYV; (peptide 79; SEQ ID NO: 89)
 RIRNGALLRVYVPRS; (peptide 80; SEQ ID NO: 90)
 VYVPRSSLPGFYRTG; (peptide 83; SEQ ID NO: 93)
 PRSSLPGFYRTGLTL; (peptide 84; SEQ ID NO: 94)
 SLPGFYRTGLTLAAP; (peptide 85; SEQ ID NO: 95)
 GFYRTGLTLAAPEAA; (peptide 86; SEQ ID NO: 96)
 RTGLTLAAPEAAGEV; (peptide 87; SEQ ID NO: 97)
 LTLAAPEAAGEVERL; (peptide 88; SEQ ID NO: 98)
 AAPEAAGEVERLIGH; (peptide 89; SEQ ID NO: 99)
 EAAGEVERLIGHPLP; (peptide 90; SEQ ID NO: 100)
 GEVERLIGHPLPLRL; (peptide 91; SEQ ID NO: 101)
 ERLIGHPLPLRLDAI; (peptide 92; SEQ ID NO: 102)
 IGHPLPLRLDAITGP; (peptide 93; SEQ ID NO: 103)
 PLPLRLDAITGPPEEE; (peptide 94; SEQ ID NO: 104)
 LRLDAITGPPEEGGR; (peptide 95; SEQ ID NO: 105)
 DAITGPPEEGGRLET; (peptide 96; SEQ ID NO: 106)
 TGPEEGGRLETILG; (peptide 97; SEQ ID NO: 107)
 EEEGRLETILGWPL; (peptide 98; SEQ ID NO: 108)
 GGRLETILGWPLAER; (peptide 99; SEQ ID NO: 109)
 LETILGWPLAERTVV; (peptide 100; SEQ ID NO: 110)
 ILGWPLAERTVVIPS; (peptide 101; SEQ ID NO: 111)
 WPLAERTVVIPS AIP; (peptide 102; SEQ ID NO: 112)
 AERTVVIPS AIPTDP; (peptide 103; SEQ ID NO: 113)
 TVVIPS AIPTDPRNV; (peptide 104; SEQ ID NO: 114)

-continued

IPSAIPTDPRNVGGD; (peptide 105; SEQ ID NO: 115)
 AIPTDPRNVGGDLDP; (peptide 106; SEQ ID NO: 116)
 TDPNRVGGDLDPSSI; (peptide 107; SEQ ID NO: 117)
 RNVGGDLDPSSIPDK; (peptide 108; SEQ ID NO: 118)
 GGDLDPSSIPDKEQA; (peptide 109; SEQ ID NO: 119)
 PDKEQAISALPDYAS; (peptide 112; SEQ ID NO: 122)
 EQAISALPDYASQPG; (peptide 113; SEQ ID NO: 123)
 ISALPDYASQPGKPP; (peptide 114; SEQ ID NO: 124)
 LPDYASQPGKPPRED; (peptide 115; SEQ ID NO: 125)
 YASQPGKPPREDLK; (peptide 116; SEQ ID NO: 126)
 ITGPEEEGGRLDTIL; (peptide 117; SEQ ID NO: 127)
 PEEEGGRLDTILGWP; (peptide 118; SEQ ID NO: 128)
 EGGRLDTILGWPLAE; (peptide 119; SEQ ID NO: 129)
 and
 RLDTILGWPLAERTV. (peptide 120; SEQ ID NO: 130)

[0275] Embodiments of the invention also comprise or consist of isolated polypeptides (proteins) and peptides comprising or consisting of the above-referenced amino acids sequences, except wherein one or more amino acids have been substituted with conservative amino acids substitutions. Embodiments of the invention also comprise or consist of isolated polypeptides (proteins) and peptides comprising or consisting of the above-referenced amino acids sequences, except wherein one or more amino acids have been substituted with amino acids which are naturally occurring, non-naturally occurring, non-standard amino acids, or amino acid analogs.

[0276] Embodiments of the invention include polypeptides comprising a PE-A Domain III (i.e., a cytotoxic domain; see e.g., FIG. 1). Examples of sequences comprising a cytotoxic portion of PE can be found in SEQ ID NO:1 and SEQ ID NO:4 spanning amino acid residues Phe-134 to Lys-347. Examples of sequences comprising a cytotoxic portion of PE can also be found in SEQ ID NO:133 and SEQ ID NO:134 spanning amino acid residues Phe-400 to Lys-613.

[0277] Embodiments of the invention include polypeptides comprising a PE-A Domain III (i.e., a cytotoxic domain) and one or more PE-A domains selected from the group consisting of:

[0278] (a) Domain II (i.e., a cytosolic translocation domain; e.g., amino acids corresponding to Gly-3 to Ser-14 in SEQ ID NO:1 or amino acids corresponding to Gly-3 to Asn-114 in SEQ ID NO:4, see e.g., FIG. 1);

[0279] (b) Carboxy-terminal portion of Domain IB (e.g., amino acids corresponding to Gly-115 to Glu-133 in SEQ ID NO:1 or SEQ ID NO:4; and amino acids corresponding to Gly-381 to Glu-399 in SEQ ID NO:133 or SEQ ID NO:134; see e.g., FIG. 1);

[0280] (c) Amino-terminal portion of Domain IB (e.g., amino acids corresponding to Ala-365 to Ala-380 in SEQ ID NO:133 or SEQ ID NO:134; see e.g., FIG. 1);

[0281] (d) Domain IB (i.e., amino acid sequences intervening between Domains II and III; e.g., amino acids

corresponding to Ala-365 to Glu-399 in SEQ ID NO:133 or SEQ ID NO:134; see e.g., FIG. 1);

[0282] (e) a carboxy-terminal tail selected from the group consisting of:

(i) Arg-Glu-Asp-Leu-Lys; (SEQ ID NO: 135)

(ii) Arg-Glu-Asp-Leu; (SEQ ID NO: 136)
 and

(iii) Lys-Asp-Glu-Leu, (SEQ ID NO: 137)

[0283] wherein one or more of said domains has been modified with amino acid substitutions, as described herein, to reduce or eliminate immunogenicity.

[0284] Embodiments of the invention further comprise PE variants. For example, such variants include, without limitation, PE polypeptide examples as shown in SEQ ID Nos: 143 to 163 and SEQ ID No:175.

[0285] Embodiments of the invention comprise any one or more of the PE-A domains indicated in the preceding paragraphs, wherein said one or more domains are chemically linked, covalently coupled, or fused (i.e., as in-frame fusion proteins) with a heterologous polypeptide (for example, such as ligand or antigen-binding polypeptide).

[0286] Embodiments of the invention include polypeptides comprising PE wherein one or more amino acids are substituted with any combination of one or more conservative amino acid substitutions, non-conservative amino acid substitutions, non-naturally occurring amino acid substitutions, non-standard amino acids, and/or substitutions with amino acid analogs and further wherein said polypeptides are non-immunogenic or exhibit reduced immunogenicity as determined and assayed by comparison to immunogenicity of corresponding non-amino acid substituted forms of PE; as measured using in vitro or in vivo assays. In particular embodiments, amino acid substituted forms of PE are at least 25%, at least about 25%, at least 50%, at least about 50%, at least 75%, or at least about 75% less immunogenic compared to corresponding non-amino acid substituted forms of PE. In particular embodiments, amino acid substituted forms of PE are at least 2-fold, at least about 2-fold, at least 3-fold, at least about 3-fold, at least 4-fold, at least about 4-fold, at least 5-fold, at least about 5-fold, at least 10-fold, at least about 10-fold, at least 50-fold, at least about 50-fold, at least 100-fold, at least about 100-fold, at least 500-fold, at least about 500-fold, at least 1,000-fold, or at least about 1000-fold less immunogenic compared to corresponding non-amino acid substituted forms of PE. In one embodiment, amino acid substituted forms of PE are non-immunogenic or exhibit undetectable immunogenicity compared to corresponding non-amino acid substituted forms of PE.

[0287] The immunogenicity of substituted peptides may be measured, via assays routinely known and used by those of skill in the art. For example, immunogenicity may be assayed by methods including, but not limited to, the proliferation assays described in Example 1 herein.

[0288] Additionally, methods for predicting, and assays for assessing, immunogenicity include those methods and assays such as described or referenced in:

[0289] Baker M P and Jones T D. Identification and removal of immunogenicity in therapeutic proteins. *Curr. Opin. Drug. Disc. Dev.* 2007 10(2): 219-227.

- [0290] Bryson C J, Jones T D, Baker M P. Prediction of immunogenicity of therapeutic proteins: validity of computational tools. *BioDrugs*. 2010; 24(1): 1-8.
- [0291] Chester, K, Baker, M P and Mayer A. Overcoming the immunologic response to foreign enzymes in cancer therapy. *Expert Rev. Clin. Immunol.* 2006 1(4): 549-559.
- [0292] Hochuli E. Interferon immunogenicity: technical evaluation of interferon-alpha 2a. *J Interferon Cytokine Res.* 1997 17 Suppl 1: 15-21
- [0293] Jaber A and Baker M P. Assessment of the immunogenicity of different interferon beta-1a formulations using ex vivo T cell assays. *J Pharm Biomed Anal* 2007 43(4): 1256-61.
- [0294] Perry L C, Jones T D and Baker M P. New approaches to prediction of immune responses to therapeutic proteins during preclinical development. *Drugs R D.* 2008 9(6):385-96.
- [0295] Schellekens, H., Ryff, J C., and Van Der Meide, P. H. Assays for antibodies to human interferon-alpha: the need for standardization *J. Interferon Cytokine Res.* 1997 17(Suppl. 1), S5-S8.
- [0296] Embodiments of the invention include polypeptides comprising PE wherein one or more amino acids are substituted with any combination of one or more conservative amino acid substitutions, non-conservative amino acid substitutions, non-naturally occurring amino acid substitutions, non-standard amino acids, and/or substitutions with amino acid analogs and further wherein said polypeptides retain biological activity as determined and assayed by comparison to biological activities of corresponding non-amino acid substituted forms of PE; such as, but not limited to, cell killing activity, cell cytotoxicity, inactivation of the translation elongation factor EF-2, ADP-ribosylation of EF-2, and inhibition of protein synthesis as measured using in vitro or in vivo assays. In particular embodiments, amino acid substituted forms of PE exhibit 100% or about 100% of biological activity compared to corresponding non-amino acid substituted forms of PE. In particular embodiments, amino acid substituted forms of PE exhibit at least 95%, or at least about 95% of biological activity compared to corresponding non-amino

acid substituted forms of PE. In particular embodiments, amino acid substituted forms of PE exhibit at least 90%, at least about 90%, at least 85% at least about 85%, at least 80%, at least about 80%, at least 75%, at least about 75%, at least 70%, at least about 70%, at least 60%, at least about 60%, at least 50%, or at least about 50% of biological activity compared to corresponding non-amino acid substituted forms of PE.

[0297] Embodiments of the invention further comprise fusion proteins, conjugates, covalently-linked, and non-covalently linked amino acid substituted forms of PE, or fragments thereof, as described herein. Amino acid substituted forms of PE may be fused, conjugated or otherwise linked with any artificial, recombinant, or naturally occurring molecule or polypeptide to modify PE activity and/or PE localization/targeting, such as by conferring to PE, via said fusion or conjugation, the tissue targeting, cell targeting, or sub-cellular localization properties of the molecule to which PE is fused, conjugated or otherwise linked. For example, but without limitation, amino acid substituted forms, or fragments thereof, of PE may be fused, conjugated, or otherwise linked with any type of antibody or antigen-binding fragments thereof, cell-surface receptor, secreted or cell-surface ligand, or fragments thereof.

[0298] In one embodiment, amino acid substituted forms of PE, including amino acid substituted forms of PE fused, conjugated or otherwise linked to another molecule or polypeptide are useful in the treatment of cancer; including, but not limited to, types of cancer described herein. In one embodiment, amino acid substituted forms of PE as described herein are useful for the preparation of a medicament for the treatment of cancer; including, but not limited to, types of cancer described herein.

[0299] In one embodiment, amino acid substituted forms of PE, or fragments thereof, may be fused, conjugated, or otherwise linked, without limitation, antigen-binding moieties such as antibodies, or fragments thereof, which specifically or preferentially bind to disease associated antigens. Such molecules include, for example, but without limitation, antibodies indicated in Table 1.

TABLE 1

Examples of Antibodies and Therapeutic Uses			
NAME	TRADE NAME	Putative Antigen Targets	Example(s) of Therapeutic Use
3F8		GD2	neuroblastoma
ABAGOVOMAB		CA-125 (imitation)	ovarian cancer
ABCIXIMAB	REOPRO	CD41 (integrin alpha-IIb)	platelet aggregation inhibitor
ADALIMUMAB	HUMIRA	TNF- α	rheumatoid arthritis etc.
ADECATUMUMAB		EpCAM	prostate and breast cancer
AFELIMOMAB		TNF- α	sepsis
AFUTUZUMAB		CD20	lymphoma
ALACIZUMAB		VEGFR2	cancer
PEGOL			
ALD518		IL-6	rheumatoid arthritis
ALEMTUZUMAB	CAMPATH, MABCAMPATH	CD52	CLL, CTCL
ALTUMOMAB	HYBRI-CEAKER	CEA	colorectal cancer (diagnosis)
PENTETATE			
ANATUMOMAB		TAG-72	non-small cell lung carcinoma
MAFENATOX			
ANRUKINZUMAB		IL-13	antigen-induced pulmonary inflammation, asthma

TABLE 1-continued

Examples of Antibodies and Therapeutic Uses			
NAME	TRADE NAME	Putative Antigen Targets	Example(s) of Therapeutic Use
APOLIZUMAB ARCITUMOMAB	CEA-SCAN	HLA-DR CEA	hematological cancers gastrointestinal cancers (diagnosis)
ASELIZUMAB		L-selectin (CD62L)	severely injured patients
ATLIZUMAB	ACTEMRA, ROACTEMRA	IL-6 receptor	rheumatoid arthritis
ATOROLIMUMAB		Rhesus factor	hemolytic disease of the newborn
BAPINEUZUMAB BASILIXIMAB	SIMULECT	beta amyloid CD25 (α chain of IL-2 receptor)	Alzheimer's disease prevention of organ transplant rejections
BAVITUXIMAB BECTUMOMAB		phosphatidylserine CD22	cancer, viral infections non-Hodgkin's lymphoma (detection)
BELIMUMAB	BENLYSTA, LYMPHOSTAT-B	BAFF	non-Hodgkin lymphoma etc.
BENRALIZUMAB BERTILIMUMAB BESILESOMAB		CD125 CCL11 (eotaxin-1)	asthma severe allergic disorders
BEVACIZUMAB BICIROMAB	SCINTIMUN	CEA-related antigen	inflammatory lesions and metastases (detection)
BIVATUZUMAB MERTANSINE BLINATUMOMAB BRENTUXIMAB VEDOTIN BRIAKINUMAB		VEGF-A fibrin II, beta chain	metastatic cancer thromboembolism (diagnosis)
	AVASTIN FIBRISCINT	CD44 v6	squamous cell carcinoma
		CD19 CD30 (TNFRSF8)	cancer hematologic cancers
	ILARIS	IL-12, IL-23	psoriasis, rheumatoid arthritis, inflammatory bowel diseases, multiple sclerosis
CANAKINUMAB CANTUZUMAB MERTANSINE CAPROMAB PENDETIDE CATUMAXOMAB		IL-1 mucin CanAg	rheumatoid arthritis colorectal cancer etc.
	PROTASCINT	prostatic carcinoma cells	prostate cancer (detection)
CC49 CEDELIZUMAB	REMOVAB	EpCAM, CD3	ovarian cancer, malignant ascites, gastric cancer tumor detection
	CIMZIA	TAG-72 CD4	prevention of organ transplant rejections, treatment of autoimmune diseases
CERTOLIZUMAB PEGOL CETUXIMAB		TNF- α	Crohn's disease
	ERBITUX	EGFR	metastatic colorectal cancer and head and neck cancer
CITATUZUMAB BOGATOX CIXUTUMUMAB CLENOLIXIMAB CLIVATUZUMAB TETRAXETAN CONATUMUMAB CR6261		EpCAM	ovarian cancer and other solid tumors
	ZENAPAX	IGF-1 receptor CD4 MUC1	solid tumors rheumatoid arthritis pancreatic cancer
		TRAIL-R2 Influenza A hemagglutinin	cancer infectious disease/influenza A
DACETUZUMAB DACLIZUMAB	PROLIA	CD40 CD25 (α chain of IL-2 receptor)	hematologic cancers prevention of organ transplant rejections
DARATUMUMAB		CD38 (cyclic ADP ribose hydrolase)	myeloma, CD38-positive multiple myeloma
DENOSUMAB	SOLIRIS	RANKL	osteoporosis, bone metastases etc.
DETUMOMAB DORLIMOMAB ARTOX DORLIXIZUMAB		B-lymphoma cell auto immune associated antigen	lymphoma auto immune disorders
	SOLIRIS	CD3	type 1 diabetes, autoimmune diseases
ECROMEXIMAB ECULIZUMAB		GD3 ganglioside C5	malignant melanoma paroxysmal nocturnal hemoglobinuria

TABLE 1-continued

Examples of Antibodies and Therapeutic Uses			
NAME	TRADE NAME	Putative Antigen Targets	Example(s) of Therapeutic Use
EDOBACOMAB		Endotoxin	sepsis caused by Gram-negative bacteria
EDRECOLOMAB	PANOREX	EpCAM	colorectal carcinoma
EFALIZUMAB	RAPTIVA	LFA-1 (CD11a)	psoriasis (blocks T-cell migration)
EFUNGUMAB	MYCOGRAB	Hsp90	invasive <i>Candida</i> infection
ELOTUZUMAB		SLAMF7	multiple myeloma
ELSILIMOMAB		IL-6	Lymphoma, myeloma
ENLIMOMAB/ ENLIMOMAB PEGOL		ICAM-1 (CD54)	stroke
EPITUMOMAB		Episialin	cancer
CITUXETAN			
EPRATUZUMAB		CD22	cancer, SLE
ERLIZUMAB		ITGB2 (CD18)	heart attack, stroke, traumatic shock
ERTUMAXOMAB	REXOMUN	HER2/neu, CD3	breast cancer etc.
ETARACIZUMAB	ABEGRIN	integrin $\alpha v \beta 3$	melanoma, prostate cancer, ovarian cancer etc.
EXBIVIRUMAB		hepatitis B surface antigen	hepatitis B
FANOLESOMAB	NEUTROSPEC	CD15	appendicitis (diagnosis)
FARALIMOMAB		interferon receptor	autoimmune disorders
FARLETUZUMAB		folate receptor I	ovarian cancer
FELVIZUMAB		respiratory syncytial virus	respiratory syncytial virus infection
FEZAKINUMAB		IL-22	rheumatoid arthritis, psoriasis
FIGITUMUMAB		IGF-1 receptor	adrenocortical carcinoma, non-small cell lung carcinoma etc.
FONTOLIZUMAB	HUZAF	IFN- γ	Crohn's disease etc.
FORAVIRUMAB		rabies virus glycoprotein	rabies (prophylaxis)
FRESOLIMUMAB		TGF- β	idiopathic pulmonary fibrosis, local segmental glomerulosclerosis, cancer
GALIXIMAB		CD80	B-cell lymphoma
GANTENERUMAB		beta amyloid	Alzheimer's disease
GAVILIMOMAB		CD147 (basigin)	graft versus host disease
GEMTUZUMAB	MYLOTARG	CD33	acute myelogenous leukemia
OZOGAMICIN			
GIRENTUXIMAB	RENCAREX	carbonic anhydrase 9 (CA-IX)	clear cell renal cell carcinoma
GLEMBATUMUMAB		GPNMB	melanoma, breast cancer
VEDOTIN			
GOLIMUMAB	SIMPONI	TNF- α	rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis
GOMILIXIMAB		CD23 (IgE receptor)	allergic asthma
IBALIZUMAB		CD4	HIV infection
IBRITUMOMAB	ZEVALIN	CD20	non-Hodgkin's lymphoma
TIUXETAN			
IGOVOMAB	INDIMACIS-125	CA-125	ovarian cancer (diagnosis)
IMCIROMAB	MYOSCINT	cardiac myosin	cardiac imaging
INFLIXIMAB	REMICADE	TNF- α	rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, psoriasis, Crohn's disease, ulcerative colitis
INTETUMUMAB		CD51	solid tumors (prostate cancer, melanoma)
INOLIMOMAB		CD25 (α chain of IL-2 receptor)	graft versus host disease
INOTUZUMAB		CD22	cancer
OZOGAMICIN			
IPILIMUMAB	YERVOY	CD152	melanoma
IRATUMUMAB		CD30 (TNFRSF8)	Hodgkin's lymphoma
KELIXIMAB		CD4	chronic asthma

TABLE 1-continued

Examples of Antibodies and Therapeutic Uses			
NAME	TRADE NAME	Putative Antigen Targets	Example(s) of Therapeutic Use
LABETUZUMAB LEBRIKIZUMAB LEMALESOMAB	CEA-CIDE	CEA IL-13 NCA-90 (granulocyte antigen)	colorectal cancer asthma diagnostic agent
LERDELIMUMAB		TGF beta 2	reduction of scarring after glaucoma surgery
LEXATUMUMAB LIBIVIRUMAB		TRAIL-R2 hepatitis B surface antigen	cancer hepatitis B
LINTUZUMAB LORVOTUZUMAB MERTANSINE LUCATUMUMAB		CD33 CD56	cancer cancer
		CD40	multiple myeloma, non-Hodgkin's lymphoma, Hodgkin's lymphoma
LUMILIXIMAB		CD23 (IgE receptor)	chronic lymphocytic leukemia
MAPATUMUMAB MASLIMOMAB MATUZUMAB		TRAIL-R1 T-cell receptor EGFR	cancer autoimmune disorders colorectal, lung and stomach cancer
MEPOLIZUMAB		IL-5	asthma and white blood cell diseases
METELIMUMAB MILATUZUMAB		TGF beta 1 CD74	systemic scleroderma multiple myeloma and other hematological malignancies
MINRETUMOMAB MITUMOMAB MOROLIMUMAB MOTAVIZUMAB		TAG-72 GD3 ganglioside Rhesus factor	cancer small cell lung carcinoma disease antigen
MUROMONAB-CD3 NACOLOMAB TAFENATOX NAPTUMOMAB ESTAFENATOX	NUMAX ORTHOCLONE OKT3	respiratory syncytial virus CD3	respiratory syncytial virus (prevention) prevention of organ transplant rejections
		C242 antigen	colorectal cancer
NATALIZUMAB	TYSABRI	5T4	non-small cell lung carcinoma, renal cell carcinoma
NEBACUMAB NECITUMUMAB		integrin $\alpha 4$	multiple sclerosis, Crohn's disease
NERELIMOMAB NIMOTUZUMAB		Endotoxin EGFR	sepsis non-small cell lung carcinoma
	THERACIM, THERALOC	TNF- α EGFR	auto immune disorders squamous cell carcinoma, head and neck cancer, nasopharyngeal cancer, glioma
NOFETUMOMAB MERPENTAN OCRELIZUMAB		cancer-associated antigen CD20	cancer (diagnosis) rheumatoid arthritis, lupus erythematosus etc.
ODULIMOMAB		LFA-1 (CD11a)	prevention of organ transplant rejections, immunological diseases
OFATUMUMAB	ARZERRA	CD20	chronic lymphocytic leukemia
OLARATUMAB OMALIZUMAB OPORTUZUMAB MONATOX	XOLAIR	PDGF-R α IgE Fc region EpCAM	cancer allergic asthma cancer
OREGOVOMAB OTELIXIZUMAB PAGIBAXIMAB PALIVIZUMAB		CA-125 CD3 lipoteichoic acid respiratory syncytial virus	ovarian cancer diabetes mellitus type 1 sepsis (<i>Staphylococcus</i>) respiratory syncytial virus (prevention)
PANTUMUMAB PANOACUMAB	SYNAGIS, ABBOSYNAGIS VECTIBIX	EGFR <i>Pseudomonas aeruginosa</i>	colorectal cancer <i>Pseudomonas aeruginosa</i> infection
PASCOLIZUMAB PEMTUMOMAB		IL-4 MUC1	asthma cancer

TABLE 1-continued

Examples of Antibodies and Therapeutic Uses			
NAME	TRADE NAME	Putative Antigen Targets	Example(s) of Therapeutic Use
PERTUZUMAB PEXELIZUMAB	OMNITARG	HER2/neu C5	cancer reduction of side effects of cardiac surgery
PINTUMOMAB		adenocarcinoma antigen	adenocarcinoma
PRILIXIMAB		CD4	Crohn's disease, multiple sclerosis
PRITUMUMAB PRO 140		vimentin	brain cancer
RAFIVIRUMAB	LUCENTIS	CCR5	HIV infection
RAMUCIRUMAB		rabies virus glycoprotein	rabies (prophylaxis)
RANIBIZUMAB		VEGFR2	solid tumors
RAXIBACUMAB		VEGF-A	macular degeneration (wet form)
REGAVIRUMAB		anthrax toxin, protective antigen	anthrax (prophylaxis and treatment)
RESLIZUMAB		cytomegalovirus glycoprotein B	cytomegalovirus infection
RILOTUMUMAB	MABTHERA, RITUXAN	IL-5	inflammations of the airways, skin and gastrointestinal tract
RITUXIMAB		HGF	solid tumors
ROBATUMUMAB		CD20	lymphomas, leukemias, some autoimmune disorders
RONTALIZUMAB		IGF-1 receptor	cancer
ROVELIZUMAB	LEUKARREST ANTOVA	IFN- α	systemic lupus erythematosus
RUPLIZUMAB		CD11, CD18	haemorrhagic shock
SATUMOMAB		CD154 (CD40L)	rheumatic diseases
PENDETIDE		TAG-72	cancer
SEVIRUMAB		cytomegalovirus	cytomegalovirus infection
SIBROTUZUMAB		FAP	cancer
SIFALIMUMAB		IFN- α	SLE, dermatomyositis, polymyositis
SILTUXIMAB		IL-6	cancer
SIPLIZUMAB	LEUKOSCAN	CD2	psoriasis, graft-versus-host disease (prevention)
SOLANEZUMAB		beta amyloid	Alzheimer's disease
SONEPCIZUMAB		sphingosine-1-phosphate	choroidal and retinal neovascularization
SONTUZUMAB		episialin	disease antigen
STAMULUMAB		myostatin	muscular dystrophy
SULESOMA		NCA-90 (granulocyte antigen)	osteomyelitis (imaging)
TACATUZUMAB		alpha-fetoprotein	cancer
TETRAXETAN	AFP-CIDE	integrin α IIb β 3	percutaneous coronary intervention
TADOCIZUMAB		IgE	allergic reaction
TALIZUMAB		NGF	pain
TANEZUMAB		CD19	cancer
TAPLITUMOMAB			
PAPTOX			
TEFIBAZUMAB	AUREXIS	clumping factor A	<i>Staphylococcus aureus</i> infection
TELIMOMAB		autoimmune antigen	autoimmune disorders
ARITOX		tenascin C	cancer
TENATUMOMAB		CD40	autoimmune disorders
TENELIXIMAB		CD3	diabetes mellitus type 1
TEPLIZUMAB		CD2	chronic lymphocytic leukemia, rheumatoid arthritis
TGN1412			
TICILIMUMAB	ACTEMRA, ROACTEMRA	CTLA-4	cancer
TIGATUZUMAB		TRAIL-R2	cancer
TNX-650		IL-13	Hodgkin's lymphoma
TOCILIZUMAB		IL-6 receptor	rheumatoid arthritis
TORALIZUMAB		CD154 (CD40L)	rheumatoid arthritis, lupus nephritis

TABLE 1-continued

Examples of Antibodies and Therapeutic Uses			
NAME	TRADE NAME	Putative Antigen Targets	Example(s) of Therapeutic Use
TOSITUMOMAB	BEXXAR	CD20	follicular lymphoma
TRASTUZUMAB	HERCEPTIN	HER2/neu	breast cancer
TREMELIMUMAB		CTLA-4	cancer
TUCOTUZUMAB		EpCAM	cancer
CELMOLEUKIN			
TUVIRUMAB		hepatitis B virus	chronic hepatitis B
URTOXAZUMAB		<i>Escherichia coli</i>	diarrhoea caused by <i>E. coli</i>
USTEKINUMAB	STELARA	IL-12, IL-23	multiple sclerosis, psoriasis, psoriatic arthritis
VAPALIXIMAB		AOC3 (VAP-1)	autoimmune disorders
VEDOLIZUMAB		integrin $\alpha 4\beta 7$	Crohn's disease, ulcerative colitis
VELTUZUMAB		CD20	non-Hodgkin's lymphoma
VEPALIMOMAB		AOC3 (VAP-1)	inflammation
VISILIZUMAB	NUVION	CD3	Crohn's disease, ulcerative colitis
VOLOCIXIMAB		integrin $\alpha 5\beta 1$	solid tumors
VOTUMUMAB	HUMASPECT	tumor antigen CTAA16.88	colorectal tumors
ZALUTUMUMAB	HUMAX-EGFR	EGFR	squamous cell carcinoma of the head and neck
ZANOLIMUMAB	HUMAX-CD4	CD4	rheumatoid, arthritis, psoriasis, T-cell lymphoma
ZIRALIMUMAB		CD147 (basigin)	autoimmune disorders
ZOLIMOMAB		CD5	systemic lupus
ARITOX			erythematosis, graft-versus-host disease

[0300] In certain embodiments, amino acid substituted forms of PE, or fragments thereof, may be fused, conjugated, or otherwise linked, without limitation, to naturally occurring normal or disease related molecules such as secreted, extracellular, intracellular, transmembrane, or cell-surface-bound molecules or fragments thereof (or non-naturally occurring variants and fragments thereof), such as without limitation: ligands, receptors, receptor extracellular domains, cytokines, growth factors, cell signaling proteins, extracellular and intracellular enzymes, structural proteins, cell adhesion proteins and molecules, cluster of differentiation (CD) molecules, mitogens, cell division regulating molecules, cancer/tumor markers and antigens, et cetera. In certain embodiments, molecules which are normally transmembrane and cell-surface bound polypeptides may be fused or conjugated to amino acid substituted forms of PE as polypeptide fragments lacking at least their transmembrane domains or polypeptide regions responsible for cell-surface binding.

[0301] In certain embodiments, molecules may be fused or conjugated to amino acid substituted forms of PE wherein such molecules possess or retain the ability (even as fusion proteins or protein conjugates) to form multimeric complexes (such as hetero- and homopolymers including, but not limited to, dimers, trimers, tetramers, pentamers, hexamers, et cetera.)

[0302] In certain embodiments, amino acid substituted forms of PE, or fragments thereof, may be generated as in-frame polypeptide fusion proteins with molecules (such as, but not limited to, those referenced above) wherein the PE moiety is either an amino-terminal portion or a carboxyl-terminal portion of the fusion protein. Determination of which of these two configurations provides the desired results and/or biological activities may be determined by routine experimentation practiced by those skilled in the art.

[0303] In certain embodiments, amino acid substituted forms of PE, or fragments thereof, may be generated as fusion proteins wherein heterologous amino acid sequences (such as cell targeting sequences) are inserted within the amino acid substituted form of PE (i.e., heterologous amino acids are flanked at the amino terminus and at the carboxy terminus by PE amino acid sequences). An example of a non-amino acid substituted form of PE in such a configuration is demonstrated in U.S. Pat. No. 8,854,044 wherein a TGF- α polypeptide is incorporated at amino acid residues 607 to 604 within a "PE37" polypeptide sequence. See e.g., U.S. Pat. No. 8,854,044, FIG. 1.

[0304] Some examples of molecules which may be fused, conjugated, or otherwise linked to amino acid substituted forms of PE, include for example, but without limitation, those such as indicated in Table 2.

TABLE 2

Examples of Potential Compounds and Indications to which Amino Acid Substituted Forms of PE May Be Fused or Conjugated for Therapeutic Use		
Note: The potential indications and nucleic acid and amino acid sequences shown in Table 2 (as well as accession numbers listed) are presented for purposes of providing a few illustrative examples only. Thus, embodiments of the invention may or may not comprise these indications and sequences. Accordingly, it is envisioned that embodiments of the invention comprise other indication uses as well as other molecules and sequence variants (e.g., naturally occurring variants (such as allelic or polymorphic variants) and non-naturally occurring variants (such as genetically engineered or mutated variants)) of these sequences wherein one or multiple amino acids are changed and/or wherein only a fragment or fragments of such sequences are fused or conjugated to amino acid substituted forms of PE. Hence, the examples shown in Table 2 should in no manner be considered limiting with respect to potential therapeutic indications or protein fusions and conjugates of amino acid modified forms of PE.		
Example Molecule (Nucleotide Accession*) [Protein Accession**]	Potential Indications	Example Amino Acid Sequence
Mesothelin (NM_013404) [NP_037536]	Pancreatic cancer Ovarian cancer	MALPTARPLLGSCGTPALGSLFLFLFSLGWVQPS RTLGETGQEAAPLDGVLANPPNIISSLSRQLLG FPCAESVGLSTERVRELAVALAQKNVKLSTEQRL CLAHRLSEPPEDLDALPLDLLFLNPDASFSGPQA CTRFFSRITKANVDLLPRGAPERQRLPAALACW GVRGSLLEADVRLGGLACDLPGRFVAESAELV LPRLVSCPGPLDQDQQAARAALQGGGPPYPGPS TWSVSTMDALRGLLPVLGQPIIRSIPQGIVAAWR QRSSRDPSWRQPRTILRPRFRREVEKTACPSGK KAREIDESLIFYKKWELEACVDAALLATQMDRVN AIPFTYEQLDVLKHKLDLEYPQGYPESVIQHLGY LFLKMSPEDIRKWNVTSLTLKALLEVNESHMS PQAPRRPLPQVATLIDRFVKGRGQLDKDTLDTLT AFYPGYLCSLSPEELSSVPPSSIWAVRPQDLDT DPRQLDVLYPKARLAFQNMNGSEYFVKIQSFLGG APTEDLKALSQQNVSMDLATFMKLRTDAVLPLTV AEVQKLLGPHVEGLKAEERHRPVRDWILRQRQDD LDTLGLGLQGGIPNGYLVLDLSMQEALSGTPCLL GPSPVLTVLALLASTLA (SEQ ID NO: 167)
CD24 (NM_013230) [AAH64619]	Liver cancer Colorectal cancer Pancreatic cancer	MGRAMVARLGLGLLLALLPTQIYSSETTTGTS SNSSQSTNSGLAPNPTNATTKAAGGALQSTASL FVVSLSLLHLYS (SEQ ID NO: 168)
CD22 (AB013007) [BAA36576]	Hairy Cell Leukemia Chronic Lymphocytic Leukemia Non-Hodgkin's Lymphoma	VRRAPLSEGPSLGCYNPMMEDGISYTTLRFPPEM NIPRTG (SEQ ID NO: 169)
CD25 aka., Interleukin 2 receptor, alpha chain (NM_000417) [NP_000408]	Hodgkin's Lymphoma Hairy Cell Leukemia Chronic Lymphocytic Leukemia Cutaneous T-cell Lymphoma Adult T-cell leukemia	MDSYLLMWGLLTFIMVPGCAELCDDDPPEIPHA TFKAMAYKEGTMNLNCECKRGFRRIKSGSLYMLCT GNSSHSSWDNQCQTSSATRNNTKQVTPQPEEQK ERKTTTEMQSPMQPVDQASLPGHCREPPWENEAT ERIYHFVVQGMVYVYQCVQGYRALHRGPAESVCKM THGKTRWTQPQLICTGEMETSQFPGEKQPASFE GRPESETSCLVTTTDFQIQTEMAATMETSIFTE YQVAVAGCVFLLISVLLLSGLTQRRQRKSRRTI (SEQ ID NO: 170)
CD174 a.k.a., Lewis Y, galactoside 3(4)-L-fucosyltransferase (NM_000149) [NP_000140]	Bladder cancer Breast cancer Colorectal cancer Esophageal cancer Gastric cancer Lung cancer Pancreatic cancer	MDPLGAAPQWPWRRCLAALLFQLLVAVCFFSYL RVSRDDATGSPRAPSGSSRQDTPTTRPTLLILLW TWPFHIPVALSRCSEMVPGTADCHITADRKVPYQ ADTVIVHHWDIMSNPKSRLPPSPRPQGQPWIWFN LEPPPNQCQHLEALDRYFNLTMSYRSDSDIFTYPG WLEPWSGQPAHPPLNLNSAKTELVAWAVSNWKPDS ARVRYYQSLQAHLKVDVYGRSHKPLPKGTMMETL SRYKPYLAFENSLHPDIYTEKLWRNALEAWAVPV

TABLE 2-continued

Examples of Potential Compounds and Indications to which Amino Acid Substituted Forms of PE May Be Fused or Conjugated for Therapeutic Use		
Note: The potential indications and nucleic acid and amino acid sequences shown in Table 2 (as well as accession numbers listed) are presented for purposes of providing a few illustrative examples only. Thus, embodiments of the invention may or may not comprise these indications and sequences. Accordingly, it is envisioned that embodiments of the invention comprise other indication uses as well as other molecules and sequence variants (e.g., naturally occurring variants (such as allelic or polymorphic variants) and non-naturally occurring variants (such as genetically engineered or mutated variants)) of these sequences wherein one or multiple amino acids are changed and/or wherein only a fragment or fragments of such sequences are fused or conjugated to amino acid substituted forms of PE. Hence, the examples shown in Table 2 should in no manner be considered limiting with respect to potential therapeutic indications or protein fusions and conjugates of amino acid modified forms of PE.		
Example Molecule (Nucleotide Accession*) [Protein Accession**]	Potential Indications	Example Amino Acid Sequence
		VLGPRSRSNYERFLPPDAFIHVDDFQSPKDLARYL QELDKDHARYLSYFRWRETLRPRSFSWALDFCKA CWKLQQESRYQTVRSIAAWFT (SEQ ID NO: 171)
TPBG a.k.a., oncofetal antigen 5T4, 5T4 oncofetal trophoblast glycoprotein (NM_006670) [CAA09930]	Non small cell lung cancer Renal carcinoma Pancreatic cancer	MPGGCSRGPAGDGRRLRLARLALVLLGWVSSSP TSSASSFSSAPFLASAVSAQPPLPDQCPALCEC SEAARTVKCVNRNLTEVPTDLPAYVRNLFITGNQ LAVLPAGAFARRPPLAELALNLSGSRLDEVRA AFEHLPSLRQLDLSHNPLADLSPFAFSGSNASVS APSPLVELILNHI VPPEDERQNRSEFQMVAALL AGRALQGLRRL ELASNHFYLPDVL AQLP SLRH LDLSNNSLVSLTYVSRNLTHLES LHL EDNALKV LHNGTLAELQGLPHIRVFLDNNPWCDCMHMADMV TWLKETE VVGKDRLTCAYPEKMRNRVLELNSA DLDCDPI LPPSLQTSYVFLGIVLALIGAIFLLVL YLNRRGIIKKWMHNIRDACRDMHEGYHTRYEINAD PRLTNLSSNSDV (SEQ ID NO: 172)
CD56 a.k.a., NCAM1, neural cell adhesion molecule 1 isoform 1 precursor (NM_000615) [NP_000606]	Small cell lung cancer Merkel cell carcinoma Ovarian cancer Neuroendocrine tumors Multiple Myeloma	MLQTKDLIWTLFFLGTA VSLQVDIVPSQGEISVG ESKFFLCVAGDAKD KDISWFS PNEGKLT PNPQQR ISVVWDDSSSTLT IYNANID DAGIYKCVVTGED GSESEATVNVKIFQKLMFNAPTQEFREGEDAV IVCDVVS LPPPTI IWKHKGRDVILKKDVRFIVLS NNYLQIRGIKKTDEGTYRCEGRILARGEINFKDI QVIVNVPTIQRQNI V NATANLGQS VTLVCDAE GFPEPTMSWTKDGEQIEQEEDDEK YIFSDSSQL TIKKVDKNDEAEYICIAENKAGEQDATIHLKVFA KPKITYVENQTAMELEEQVTLTCEASGDPIPSIT WRTSTRNISSEKTL DGHMVVRSHARVSSLTKS IQYTDAGEYICTASNTIGQDSQSMYLEVQYAPKL QGPVAVYTWEQNQVNI TCEVFAYPSATI SWFRDG QLLPSSNYSNIKIYNTPSASYLEVTPDSENDFGN YNCTAVNRIGQSELEFILVQADTPSSPSIDQVEP YSTAQVQFDEPEATGGVPILKYKA EWRAVGEEV WSKWDYAKEASMEGI VTI VGLK PETTYAVRLAA LNGKSLGEISAASEFKTQPVQGEPSAPKLEGQMG EDGNSIKVNLIKQDDGGSPIRHYLVRYRALSSEW KPEIRLP SGSDHVMLKSLDWNAEYEVYVAENQQ SKSKAAHFVFR TSAQPTAIPANGSP TSGLSTGAI VGILIVIFVLLLVVDITCYFLNKCGLFMCIAVN LCGKAGPGAKGKMEEGKAAFSKDESKEPIVEVR TEEERTPNHDGKGKHTPEPNETTPTLEPEKGPVEAK PECQETETKPAPAEVKTVPN DATQT KENESKA (SEQ ID NO: 173)
C-type lectin- like molecule-1 a.k.a., CLL-1 (AY547296) [AAT11783]	Acute myeloid leukemia	MWIDFFTYSSMSEEVTYADLQFQNSSEMEKIPEI GKFGEKAPPAPSHVWRPAALFLLCLLLIGLG VLASMFHVTLKIEMKMKNLQNI SEELQRNISLQ LMSNMNINSKIRNLSTTLQT IATKLCRELYSKEQ EHKCKPCPRRWIWHKDCYFLSDDVQ TWQESKMA

TABLE 2-continued

Examples of Potential Compounds and Indications to which Amino Acid Substituted Forms of PE May Be Fused or Conjugated for Therapeutic Use		
Note: The potential indications and nucleic acid and amino acid sequences shown in Table 2 (as well as accession numbers listed) are presented for purposes of providing a few illustrative examples only. Thus, embodiments of the invention may or may not comprise these indications and sequences. Accordingly, it is envisioned that embodiments of the invention comprise other indication uses as well as other molecules and sequence variants (e.g., naturally occurring variants (such as allelic or polymorphic variants) and non-naturally occurring variants (such as genetically engineered or mutated variants)) of these sequences wherein one or multiple amino acids are changed and/or wherein only a fragment or fragments of such sequences are fused or conjugated to amino acid substituted forms of PE. Hence, the examples shown in Table 2 should in no manner be considered limiting with respect to potential therapeutic indications or protein fusions and conjugates of amino acid modified forms of PE.		
Example Molecule (Nucleotide Accession*) [Protein Accession**]	Potential Indications	Example Amino Acid Sequence
		CAAQNASLLKINNKNALFEIKSQSRSYDYWLGLS PEEDSTRGMRVDNIINSSAWVIRNAPDLNNMYCG YINRLYVQYYHCTYKQRMICEKMANPVQLGSTYF REA (SEQ ID NO: 174)

*"Nucleotide Accession" refers to the NCBI Reference Sequence accession number associated with the corresponding nucleic acid sequence as found in the "Nucleotide" database provided for public access and searching (via the Internet) through the National Center for Biotechnology Information., U.S. National Library of Medicine (8600 Rockville Pike, Bethesda MD, 20894 USA (www.ncbi.nlm.nih.gov)).

**"Protein Accession" refers to the NCBI Reference Sequence accession number associated with the corresponding amino acid sequence as found in the "Protein" database provided for public access and searching (via the Internet) through the National Center for Biotechnology Information, U.S. National Library of Medicine (8600 Rockville Pike, Bethesda MD 20894 USA (www.ncbi.nlm.nih.gov)).

[0305] In one embodiment, the present invention includes isolated nucleic acids and methods of expressing nucleic acids encoding any of the herein-referenced modified forms of PE, including fusions, conjugates, and otherwise linked molecules; whether such forms are expressed from a single or one more separate polynucleotide sequences; whether such polynucleotide sequences are expressed from a single or one or more separate expression vectors.

[0306] Expression Vectors

[0307] In one embodiment, the present invention includes methods of making and using recombinant expression vectors to express nucleic acids encoding polypeptides comprising any of the herein-referenced modified forms of PE, including fusions, conjugates, and otherwise linked molecules. Use of a wide variety of expression vectors are well-known and routinely used by those skilled in the art. A few examples of the types of expression vectors which may be used include, but are not limited to: derivatives of human or animal viruses (such as retrovirus, adeno-associated virus, pox, baculovirus, vaccinia, herpes simplex, Epstein-Barr, adenovirus, geminivirus, and caulimovirus vectors) and insect viruses (such as baculovirus); yeast vectors; bacteriophage vectors (e.g., bacteriophage lambda); plasmids; cosmids; artificial chromosomes; liposomes; electrically charged lipids (cytofectins); DNA-protein, complexes, and biopolymers.

[0308] Gene Delivery and Expression Systems

[0309] A wide variety of methods (i.e., gene delivery systems) are available and well-known to those of skill in the art; any of such methods may be used for introducing nucleic acids encoding modified forms of PE into a cell, tissue, or

organism for in vitro, in vivo, in situ, or ex vivo expression. The methods referenced below represent examples of ways in which nucleic acid(s) encoding modified forms of PE may be introduced into a cell. These examples are in no way intended to limit the scope of that may be used for gene delivery and expression of modified forms of PE in cells, tissues, or organisms; these examples are presented to illustrate the many available methods.

[0310] —Viral-Based Delivery of Target Nucleic Acids—

[0311] Gene therapy based methods can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other poly nucleotides, as needed, to allow expression of the same). As one example, polynucleotides operably encoding the target nucleic acid can be delivered to a tissue or organism either as "naked nucleic acid" or as part of an expression vector. The term vector includes for example, but is not limited to, vectors such as plasmid vectors, cosmid vectors, artificial chromosome vectors, and viral vectors. Some examples of viral vectors include adenovirus, herpes simplex virus (HSV), alphavirus, simian virus 40, picomavirus, vaccinia virus, retrovirus, lentivirus, and adeno-associated virus. Vectors encoding modified forms of PE may be capable of replication in a cell in which it is introduced, or it may be preferred that the vector is not capable of replication. Vectors encoding modified forms of PE may be capable of integration into the genomic DNA of a cell (and subsequent expression therefrom), or it may be preferred that the vector is not capable of integrating into the host genome. An example of a vector that can integrate into the genomic DNA of a cell is a retroviral vector, in which an integrase enzyme mediates integration of the retroviral vector

sequences. A vector may also contain transposon sequences that facilitate integration of the coding region into the genomic DNA of a host cell. Liposomes represent another manner in which target DNA may be delivered to a subject.

[0312] Selection of a vector depends upon a variety of desired characteristics in the resulting construct, such as a selection marker, vector replication rate, type of target host cell, species of host organism, desired duration of protein expression. An expression vector optionally includes expression control sequences operably linked to the coding sequence such that the coding region is expressed in the cell. The invention is not limited by the use of any particular promoter, and a wide variety is known. Promoters act as regulatory signals that bind RNA polymerase in a cell to initiate transcription of a downstream (3' direction) operably linked coding sequence. The promoter used in the invention may be a constitutive or an inducible promoter. It can be, but need not be, heterologous with respect to the cell to which it is introduced.

[0313] In certain embodiments, adenovirus expression vectors can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). The terms "adenovirus expression vector" is meant to include those constructs containing nucleic acid sequences sufficient to (a) support packaging of the construct and (b) to ultimately express a recombinant gene construct that has been inserted therein. In contrast to retroviruses, use of adenovirus vectors does not result in chromosomal integration because adenovirus DNA replicates in an episomal manner. Moreover, adenoviruses are considered to be structurally stable with no genome rearrangement occurring even after extensive virus reproduction and amplification. Methods of constructing and using adenovirus vectors as gene delivery systems are well-known to those of skill in the art.

[0314] In certain embodiments, adeno-associated virus (AAV) expression vectors can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). AAV may be desirable for a number of reasons; for example, because AAV vectors exhibit a high frequency of integration, can infect nondividing cells, and have a broad host range. AAV is a dependent parvovirus in that it requires coinfection with another virus (either adenovirus or a member of the herpes virus family) to undergo a productive infection in cultured cells. In the absence of coinfection with helper virus, the wild-type AAV genome integrates through its ends into a human chromosome where it resides as a latent provirus. When a cell containing latent AAV provirus is superinfected with a helper virus, the AAV genome is "rescued" from the chromosome and a normal productive infection is established. Methods of constructing and using AAV vectors as gene delivery systems are well-known to those of skill in the art.

[0315] In certain embodiments, retrovirus expression vectors can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). Retroviruses are a group of single-stranded RNA viruses characterized by the ability to convert their genomic RNA to double-stranded DNA in infected cells through a reverse-transcription process. The resulting DNA stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. Retroviral integration results in the retention

of viral gene sequences in the recipient cell and in its descendants. Retroviral vectors are able to infect a broad variety of cell types. Methods of constructing and using retroviruses as gene delivery systems are well-known to those of skill in the art.

[0316] Many other expression vectors can also be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). For example, vectors derived from viruses such as vaccinia viruses, herpes viruses, equine encephalitis viruses, hepatitis viruses and lentiviruses can be used. Methods of constructing and using viral expression vectors as gene delivery systems are well-known to those of skill in the art. The examples of such vectors referenced herein are not intended to be limiting with respect to the means by which modified forms of PE may be delivered and expressed in various host cells, tissues, or organisms.

[0317] —Non-Viral Delivery of Modified Target Nucleic Acids—

[0318] In addition to viral delivery of modified target nucleic acid, the following are additional methods of recombinant gene delivery can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). Methods of constructing and using non-viral gene delivery systems are well-known to those of skill in the art. See, for example, Al-Dosari et al., "Nonviral gene delivery: principle, limitations, and recent progress," *AAPS Journal*, 11(4):671-681 (2009); and references cited therein.

[0319] In certain embodiments, electroporation can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). Methods of using electroporation are well-known to those of skill in the art. See, for example, Bodles-Brakhop et al., "Electroporation for the delivery of DNA-based vaccines and immunotherapeutics: current clinical developments," *Mol. Ther.*, 17(4): 585-592 (2009); and references cited therein. See also, Golzio et al., "Observations of the mechanisms of electromediated DNA uptake from vesicles to tissues," *Curr Gene Ther.*, 10(4):256-266 (2010); and references cited therein. See also, Andre et al., "Nucleic acids electrotransfer in vivo: mechanisms and practical aspects," *Curr Gene Ther.*, 10(4): 267-280 (2010) and references cited therein. See also, Wells, "Electroporation and ultrasound enhanced non-viral gene delivery in vitro and in vivo," *Cell Biol Toxicol.*, 26(1):21-28 (2010); and references cited therein.

[0320] In certain embodiments, particle bombardment can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). This method depends on the ability to accelerate nucleic acid-coated microprojectiles to a sufficient velocity to allow them to pierce cell membranes, thereby delivering nucleic acid "payloads," without killing them. Some typical microprojectiles consist of biologically inert substances such as tungsten, platinum, and gold beads. Methods of using particle bombardment are well-known to those of skill in the art. See, for example, Klein et al., "Particle bombardment: a universal approach for gene transfer to cells and tissues," *Curr. Opin. Biotechnol.*, 4(5):583-590 (1993); and references cited therein.

[0321] In certain embodiments, a variety of methods incorporating calcium phosphate co-precipitation can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). Methods of using calcium phosphate co-precipitation are well-known to those of skill in the art. See for example, Uskoković et al., "Nanosized hydroxyapatite and other calcium phosphates: chemistry of formation and application as drug and gene delivery agents," *J. Biomed. Mater. Res. B Appl. Biomater.* 96(1):152-191 (2011); and references cited therein. See also, Colosimo et al., "Transfer and expression of foreign genes in mammalian cells," *Biotechniques*, 29(2):314-8, 320-322 (2000); and references cited therein.

[0322] In certain embodiments, microinjection and sonication methods can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). Methods of using microinjection and sonication are well-known to those of skill in the art. See, for example, Rochlitz et al., "Gene therapy of cancer," *Swiss Med. Wkly.*, 131(1-2):4-9 (2001); and references cited therein. See also, Donnelly et al., "Microneedle-based drug delivery systems: microfabrication, drug delivery, and safety," *Drug.Deliv.*, 17(4):187-207 (2010); and references cited therein. See also, Miller et al., "Sonoporation: mechanical DNA delivery by ultrasonic cavitation," *Somat. Cell Mol. Genet.*, 27(1-6):115-34 (2002); and references cited therein.

[0323] In certain embodiments, liposomes and lipid formulations can be used to deliver (into a host cell, tissue or organism) target nucleic acids encoding modified forms of PE (and other polynucleotides, as needed, to allow expression of the same). Liposomes are vesicular structures characterized by a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. An example of a commonly used, commercially available lipid formulation is Lipofectamine (Gibco BRL). Methods of using liposomes and lipid formulations to deliver nucleic acids to cells, tissues and organisms are well-known to those of skill in the art. See, for example, Xiong et al., "Cationic liposomes as gene delivery system: transfection efficiency and new application," *Pharmazie*, 66(3):158-64 (2011); and references cited therein. See also, Pichon et al., "Chemical vectors for gene delivery: uptake and intracellular trafficking," *Curr Opin Biotechnol.*, 21(5):640-645 (2010); and references cited therein. See also, Pathak et al., "Recent trends in non-viral vector-mediated gene delivery," *Biotechnol J.*, 4(11):1559-1572 (2009).

[0324] Expression of Modified Forms of PE via Gene Switch Modulation Systems

[0325] Expression of modified forms of PE, including fusions, conjugates, and otherwise linked molecules, may be expressed in host cells, tissues, and organisms using gene switch expression systems. Some examples, without limitation, of such gene expression systems, and genetically engineered cells comprising gene switch expression systems, which can be used to express polynucleotides and polypeptides of the present invention, are described in the following publications; each of which are hereby incorporated by reference herein: WO 2009/045370 (PCT/US2008/011270); WO 2009/025866 (PCT/US2009/010040); WO 2002/066614 (PCT/US2002/005706); WO 2008/073154 (PCT/US2007/

016747); 2002/066613 (PCT/US2002/005090); WO 2005/108617 (PCT/US2005/015089); WO 2002/029075 (PCT/US2001/030608); WO 2003/027289 (PCT/US2002/005026); WO 2001/070816 (PCT/US2001/090500); WO 2002/066615 (PCT/US2002/005708); WO 2009/048560 (PCT/US2008/011563); WO 2003/027266 (PCT/US2002/05234); WO 2010/042189 (PCT/US2009/005510); and WO 2002/066612 (PCT/US2002/005090); WO 2011/119773 (PCT/US2011/029682).

[0326] For purposes of expressing polynucleotides and polypeptides under control of a gene switch mechanism, the term "gene switch" refers to the combination of a response element associated with a promoter, and a ligand-dependent transcription factor-based system which, in the presence of one or more ligands, modulates the expression of a gene into which the response element and promoter are incorporated. Stated otherwise, a "gene switch" refers to a peptide, protein or polypeptide complex that functions to (a) bind an activating ligand, and (b) regulate the transcription of a gene of interest in a ligand-dependent fashion.

[0327] In one embodiment, the polynucleotide encoding a gene switch is a recombinant polynucleotide, i.e., a polynucleotide, that has been engineered, by molecular biological manipulation, to encode the gene switch. In another embodiment, the recombinant polynucleotide is a synthetic polynucleotide.

[0328] As used herein with respect to gene switch regulation systems, the term "dimerizes with the ligand binding domain that binds an activating ligand" refers to a selective protein-protein interaction that is induced by the presence of activating ligand.

[0329] As used herein, the term "ligand binding domain that binds an activating ligand" refers to an amino acid sequence that selectively binds an activating ligand. In the methods disclosed herein, an activating ligand binds to a ligand binding domain, e.g., an ecdysone receptor ligand binding domain, that is part of a ligand-dependent transcriptional activation complex that regulates the expression of a polynucleotide sequence that encodes a gene of interest. Hence, the expression of the gene of interest is regulated in a ligand-dependent fashion.

[0330] The term "ecdysone receptor-based," with respect to a gene switch, refers to a gene switch comprising at least a functional part of a naturally occurring or synthetic ecdysone receptor ligand binding domain and which regulates gene expression in response to a ligand that binds to the ecdysone receptor ligand binding domain.

[0331] As used herein, "selective binding" of an activating ligand to a ligand binding domain in a gene switch means that the ligand has an EC50 of about 700 nanomolar (nM), 650 nM, 600 nM, 550 nM, 500 nM, 450 nM, 400 nM, 350 nM, 300 nM, 250 nM, 225 nM, 200 nM, 175 nM, 150 nM, 125 nM, 100 nM, 95 nM, 90 nM, 85 nM, 80 nM, 75 nM, 70 nM, 65 nM, 60 nM, 55 nM, 50 nM, 45 nM, 40 nM, 35 nM, 30 nM, 25 nM, 20 nM, 15 nM, 10 nM, 9 nM, 8 nM, 7 nM, 6 nM, 5 nM, 4 nM, 3 nM, 2 nM or 1 nM, or less, in a gene switch assay.

[0332] As used herein, "EC50" is the "half maximal effective concentration" which refers to the concentration of an activating ligand that induces a gene switch-regulated change in expression of a polynucleotide encoding a gene of interest (e.g. modified forms of PE, including fusions, conjugates, et cetera), that is halfway between the baseline level of expression and the maximum level of expression after a specified exposure time. Examples of cellular assays for measuring

gene switch-regulated gene expression are well known to those of skill in the art. See, for example, Katzenowski et al., *BioTechniques* 39: 191-200 (2005).

[0333] In one embodiment, the ligand binding domain that binds an activating ligand, e.g., an ecdysone receptor ligand binding domain, dimerizes with another ligand binding domain, e.g., a retinoid X receptor ligand binding domain, to form a protein-protein complex.

[0334] In one embodiment, the expression of the gene of interest is regulated by an activating ligand in an on/off fashion that is independent of the concentration or dosage of an activating ligand. In another embodiment, the expression of the gene of interest is regulated by an activating ligand in a concentration (or dosage)-dependent fashion, i.e., there is a dose-response relationship between the concentration (or dosage) of an activating ligand and the level of gene expression of the gene of interest. See, e.g., US Patent Publication No. 2009/0123441 (see a/to, WO 2009/048560 (PCT/USUS2008/011563)).

[0335] The term “operably linked” refers to the association of polynucleotide sequences on a single polynucleotide so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

[0336] In one embodiment, an activating ligand, or a composition thereof, is administered to a subject orally. In another embodiment, an activating ligand, or a composition thereof, is administered to a subject parenterally. In another embodiment, an activating ligand, or a composition thereof, is administered subcutaneously, intramuscularly, intravenously, intraperitoneally, transdermally, or intratumorally.

[0337] In one embodiment, the ligand binding domain in the gene switch is a Group H nuclear receptor ligand binding domain, or a mutant thereof that binds an activating ligand. In another embodiment, the Group H nuclear receptor ligand binding domain is selected from the group consisting of an ecdysone receptor ligand binding domain, a ubiquitous receptor ligand binding domain, an orphan receptor-1 ligand binding domain, an NER-1 ligand binding domain, a receptor-interacting protein-15 ligand binding domain, a liver X receptor-3 ligand binding domain, a steroid hormone receptor-like protein ligand binding domain, a liver X receptor ligand binding domain, a liver X receptor ligand binding domain, a farnesoid X receptor ligand binding domain, a receptor-interacting protein-14 ligand binding domain, and a farnesol receptor ligand binding domain ligand binding domain, or a mutant thereof that binds an activating ligand.

[0338] In another embodiment, the Group H nuclear receptor ligand binding domain is an ecdysone receptor ligand binding domain, or a mutant thereof that binds an activating ligand. In another embodiment, the ecdysone receptor ligand binding domain is selected from the group consisting of an Arthropod ecdysone receptor ligand binding domain a Lepidopteran ecdysone receptor ligand binding domain, a Dipteran ecdysone receptor ligand binding domain, an Orthopteran ecdysone receptor ligand binding domain, a Homopteran ecdysone receptor ligand binding domain and a Hemipteran ecdysone receptor ligand binding domain, a spruce budworm *Choristoneura fumiferana* ecdysone receptor ligand binding domain, a beetle *Tenebrio molitor* ecdys-

one receptor ligand binding domain, a *Manduca sexta* ecdysone receptor ligand binding domain, a *Heliothis virescens* ecdysone receptor ligand binding domain, a midge *Chironomus tentans* ecdysone receptor ligand binding domain, a silk moth *Bombyx mori* ecdysone receptor ligand binding domain, a squinting bush brown *Bicyclus anynana* ecdysone receptor ligand binding domain, a buckeye *Junonia coenia* ecdysone receptor ligand binding domain, a fruit fly *Drosophila melanogaster* ecdysone receptor ligand binding domain, a mosquito *Aedes aegypti* ecdysone receptor ligand binding domain, a blowfly *Lucilia capitata* ecdysone receptor ligand binding domain, a blowfly *Lucilia cuprina* ecdysone receptor ligand binding domain, a blowfly *Calliphora vicina* ecdysone receptor ligand binding domain, a Mediterranean fruit fly *Ceratitis capitata* ecdysone receptor ligand binding domain, a locust *Locusta migratoria* ecdysone receptor ligand binding domain, an aphid *Myzus persicae* ecdysone receptor ligand binding domain, a fiddler crab *Celula pugilator* ecdysone receptor ligand binding domain, an ixodid tick *Amblyomma americanum* ecdysone receptor ligand binding domain, a whitely *Bamecia argentifolia* ecdysone receptor ligand binding domain, a leafhopper *Nephotetix cincticeps* ecdysone receptor ligand binding domain, or a mutant thereof that binds An activating ligand.

[0339] In another embodiment, the ecdysone receptor ligand binding domain is a spruce budworm *Choristoneura fumiferana* ecdysone receptor ligand binding domain, for which the amino acid sequence is: Leu Thr Ala Asn Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Gln Gln Pro Ser Asp Gln Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln Ala Asp Asp Gln Asn Gln Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Gln Phe Ala Lys Gly Len Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Let Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala Ala Ser Asp Ser Val (position 107) Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met ala Tyr (position 127) Val Ile Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met ala Leu Asp Asn Ile His Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Pro Phe Leu Glu Ile Trp Asp Val (SEQ ID NO:1), which is also set forth as SEQ NO: 1 in U.S. Patent Publication No. 2006/0100416 A1 (see also, WO 2002/066612 (PCT/US2002/005090)).

[0340] Exemplary ecdysone receptor ligand binding domains include those disclosed, for example, in U.S. Pat. No. 7,935,510 (see also, WO 2003/0/27289 (PCT/US2002/005026)); U.S. Pat. No. 7,919,269 (see also, WO 2003/027266 (PCT/US2002/05234); U.S. Pat. No. 7,563,879 (see also, WO 2003/0/27289 (PCT/US2002/05026)); and in U.S. Patent Publication No. 2006/0100416 A1 (see also, WO 2002/066612 (PCT/US2002/005090)), each of which is hereby incorporated by reference in its entirety.

[0341] In one embodiment, the ecdysone receptor ligand binding domain is a mutant of an ecdysone receptor ligand binding domain that binds the activating compound. In another embodiment, the ecdysone receptor ligand binding domain is a mutant of the spruce budworm *Choristoneura fumiferana* ecdysone receptor ligand binding domain that binds the activating compound.

[0342] In one embodiment, the gene switch comprises a *Choristoneura fumiferana* ecdysone receptor ligand binding domain that is engineered to contain the mutations V107I and Y127E of the *Choristoneura fumiferana* ecdysone receptor sequence as set forth in SEQ ID NO:1 of U.S. Patent Publication No. 2006/0100416 (see also, WO 2002/066612 (PCT/US2002/005090)). The term "V107I" means that the valine amino acid residue at position 107 (as set forth in SEQ ID NO:1 of U.S. Patent Publication No. 2006/0100416) is changed to isoleucine. The term "Y127E" means that the tyrosine amino acid residue at position 127 (as set forth in SEQ ID NO:1 of U.S. Patent Publication No. 2006/0100416) is changed to glutamate.

[0343] Exemplary mutant ecdysone receptor ligand binding domains are disclosed, for example, in US 2006/0100416 A1 (see also, WO 2002/066612 (PCT/US2002/005090)) and U.S. Pat. No. 7,935,510 (Pub. No. 2005/0266457) (see also, WO 2005/108617 (PCT/US2005/015089)) each of which is incorporated by reference in its entirety.

[0344] In one embodiment, the gene switch comprises a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand. In one embodiment, the ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand is a Group B nuclear receptor ligand binding domain. In another embodiment, the Group B nuclear receptor ligand binding domain is selected from the group consisting of a retinoid, X receptor ligand binding domain, an H-2 region II binding protein ligand binding domain, a nuclear receptor co-regulator-1 ligand binding domain, an ultraspiracle protein ligand binding domain, a 2C1 nuclear receptor ligand binding domain, and a chorion factor 1 ligand binding domain. In another embodiment, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand is not an ecdysone receptor ligand binding domain.

[0345] In one embodiment, the ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand is a retinoic X receptor ligand binding domain. In another embodiment, the retinoic X receptor ligand binding domain is a vertebrate retinoic X receptor ligand binding domain. In another embodiment, the retinoic X receptor ligand binding domain is a *Homo sapiens* retinoic X receptor ligand binding domain. In another embodiment, the retinoic X receptor ligand binding domain is a retinoic X receptor α isoform. In another embodiment, the retinoic X receptor ligand binding domain is a retinoic X receptor β isoform. In another embodiment, the retinoic X receptor ligand binding domain is a retinoic X receptor γ isoform.

[0346] In another embodiment, the retinoic X receptor ligand binding domain is an invertebrate retinoic X receptor ligand binding domain. In another embodiment, the invertebrate retinoic X receptor ligand binding domain is a *Locusta migratoria* retinoic X receptor ligand binding domain.

[0347] In another embodiment, the invertebrate retinoic X receptor ligand binding domain is a non-Lepidopteran, non-Dipteran retinoic X receptor ligand binding domain.

[0348] In one embodiment, the retinoid receptor ligand binding domain is a vertebrate retinoid X receptor ligand binding domain, an invertebrate retinoid X receptor ligand binding domain, an ultraspiracle protein ligand binding domain, or a chimeric retinoid X receptor ligand binding domain.

[0349] In one embodiment, the chimeric retinoid X receptor ligand binding domain comprises two polypeptide frag-

ments, wherein the first polypeptide fragment is from a vertebrate retinoid X receptor ligand binding domain, an invertebrate retinoid X receptor ligand binding domain, or an ultraspiracle protein ligand binding domain, and the second polypeptide fragment is from a different vertebrate retinoid X receptor ligand binding domain, a different invertebrate retinoid X receptor ligand binding domain, or a different ultraspiracle protein ligand binding domain.

[0350] In another embodiment, the chimeric retinoid X receptor ligand binding domain is one that is disclosed in U.S. Pat. No. 7,531,326, which is hereby incorporated by reference in its entirety.

[0351] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-6, helices 1-7, helices 1-8, helices 1-9, helices 1-10, helices 1-11, or helices 1-12 of a first species of retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 7-12, helices 8-12, helices 9-12, helices 10-12, helices 11-12, helix 12, or F domain of a second species of retinoid X receptor, respectively.

[0352] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-6 of a first species RXR according to the disclosure, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 7-12 of a second species of retinoid X receptor.

[0353] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-7 of a first species retinoid X receptor according to the disclosure, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 8-12 of a second species retinoid X receptor.

[0354] In another embodiment, the first polypeptide fragment of the chimeric retinoid, X receptor ligand binding domain comprises helices 1-8 of a first species of retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 9-12 of a second species of retinoid X receptor.

[0355] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-9 of a first species of retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 10-12 of a second species of retinoid X receptor.

[0356] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-10 of a first species of retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 11-12 of a second species of retinoid X receptor.

[0357] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-11 of a first species of retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helix 12 of a second species of retinoid X receptor.

[0358] In another preferred embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-12 of a first species of retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises an F domain of a second species of retinoid X receptor.

[0359] In one embodiment, the first polypeptide fragment in the chimeric retinoid X receptor ligand binding domain is human retinoid X receptor sequence, and the second polypeptide fragment in the chimeric retinoid X receptor ligand binding domain is invertebrate retinoid X receptor sequence. In another embodiment, the invertebrate retinoid X receptor sequence is *Locusta migratoria* retinoid X receptor sequence.

[0360] In another embodiment, the first polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 1-8 of a human retinoid X receptor, and the second polypeptide fragment of the chimeric retinoid X receptor ligand binding domain comprises helices 9-12 of *Locusta migratoria* retinoid X receptor.

[0361] In one embodiment, the gene switch further comprises a DNA binding domain ("DBD"). In another embodiment, the DBD is selected from the group consisting of a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, and a yeast DBD.

[0362] In one embodiment, the gene switch further comprises a transactivation domain ("TD"). In another embodiment, the transactivation domain is selected from the group consisting of a VP16 TD, a GAL4 TD, an NF- κ B TD, a BP64 TD, and a B42 acidic TD.

[0363] In one embodiment, a DNA binding domain, the ligand binding domain that binds an activating ligand, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand, and a transactivation domain are encoded by polynucleotide sequences that are contained in the same polynucleotide.

[0364] In another embodiment, a DNA binding domain, a ligand binding domain that binds an activating ligand, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand, and a transactivation domain are encoded by polynucleotide sequences that are contained in two or more separate polynucleotide sequences.

[0365] In another embodiment, a DNA binding domain, a ligand binding domain that binds an activating ligand, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand, and a transactivation domain are encoded by polynucleotide sequences that are contained in two separate polynucleotide sequences.

[0366] In another embodiment, a DNA binding domain and a ligand binding domain that binds an activating ligand are encoded by polynucleotide sequences that are contained in a first polynucleotide sequence, and a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand and a transactivation domain are encoded by polynucleotide sequences that are contained in a second polynucleotide sequence.

[0367] In another embodiment, a DNA binding domain and a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand are encoded by polynucleotide sequences that are contained in a first polynucleotide sequence, and a ligand binding domain that binds an activating ligand and a transactivation domain are encoded by polynucleotide sequences that are contained in a second polynucleotide sequence.

[0368] In embodiments in which one or more of the DNA binding domain, a ligand binding domain that binds an activating ligand, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand, and a transactivation domain are encoded by polynucleotide sequences that are contained in one or more separate poly-

nucleotide sequences, then the one or more separate polynucleotide sequences are operably linked to one or more separate promoters. In another embodiment, the one or more separate polynucleotide sequences are operably linked to one or more separate enhancer elements. In another embodiment, the promoter(s) and/or the enhancer(s) are constitutively active. In another embodiment, the promoter(s) and/or the enhancer(s) are tissue specific promoters and/or enhancers.

[0369] In one embodiment, the gene switch comprises a DNA binding domain, an ecdysone receptor ligand binding domain, a ligand binding domain that dimerizes with the ecdysone receptor ligand binding domain, and a transactivation domain.

[0370] In another embodiment, the gene switch comprises a DNA binding domain, an ecdysone receptor ligand binding domain, a retinoid X receptor ligand binding domain, and a transactivation domain.

[0371] In another embodiment, the gene switch comprises a DNA binding domain, an ecdysone receptor ligand binding domain, a chimeric vertebrate/invertebrate retinoid X receptor ligand binding domain, and a transactivation domain.

[0372] In another embodiment, the gene switch comprises a GAL4 DNA binding domain, a *Choristoneura fumiferana* ecdysone receptor ligand binding domain that is engineered to contain the mutations V107I and V127E of the *Choristoneura fumiferana* ecdysone receptor sequence set forth in SEQ ID NO:1, a chimeric *Homo sapiens/Locusta migratoria* retinoid X receptor ligand binding, and a VP16 transactivation domain.

[0373] In another embodiment, the host cell further comprises a polynucleotide encoding a peptide, protein or polypeptide whose expression is regulated by the gene switch. A promoter that binds the gene switch complex is operably linked to the polynucleotide encoding a peptide, protein or polypeptide whose expression is regulated by the gene switch.

[0374] In another embodiment, the polynucleotide encoding a peptide, protein or polypeptide whose expression is regulated by the gene switch is contained in the same polynucleotide as a polynucleotide that encodes one or more of a DNA binding domain, the ligand binding domain that binds an activating ligand, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand, and a transactivation domain. Such constructs are disclosed, for example, in U.S. Patent Publication No. 2009/0123441 (see also, WO 2009-048560 (PCT/USUS2008/011563)).

[0375] In another embodiment, the polynucleotide encoding a peptide, protein or polypeptide whose expression is regulated by the gene switch is contained in a different nucleic acid molecule than a nucleic acid molecule that encodes one or more of a DNA binding domain, the ligand binding domain that binds an activating ligand, a ligand binding domain that dimerizes with the ligand binding domain that binds an activating ligand, and a transactivation domain.

[0376] In one embodiment, the gene switch is more sensitive to an activating ligand than to a steroid hormone. In another embodiment, the gene switch is more sensitive to an activating ligand than to another diacylhydrazine compound.

[0377] The sensitivity of a gene switch to an activating ligand, relative to another ligand, can readily be determined in an in vitro assay, for example, an in vitro assay that employs a reporter gene, such as firefly luciferase. Examples of such in

vitro assays are well known to those of ordinary skill in the art. See, for example, Karzenowski et al., *BioTechniques* 39: 3191-200 (2005).

[0378] In one embodiment, the polynucleotide encoding the gene switch is contained in a vector. In one embodiment, the vector selected from the group consisting of a plasmid, an expression vector, a replicon, a phage vector, a cosmid, a viral vector, a liposome, an electrically charged lipid (e.g., a cytofectin), a DNA-protein complex, and a biopolymer.

[0379] In another embodiment, the vector is a retroviral vector. In another embodiment, the vector is selected from the group consisting of an adeno-associated viral vector, a pox viral vector, a baculoviral vector, a vaccinia viral vector, a herpes simplex viral vector, an Epstein-Barr viral vector, an adenoviral vector, a gemini viral vector, and a caulimo viral vector.

[0380] In one embodiment, a composition of the invention comprises one or more polynucleotides that encode two or more orthogonal gene switches. Two or more individually operable gene regulation systems are said to be "orthogonal" when (a) modulation of each of the given gene switches by its respective ligand results in a measurable change in the magnitude of expression of the gene that is regulated by that gene switch, and (b) the change is statistically significantly different than the change in expression of all other gene switches that are in the host cell. In one embodiment, regulation of each individually operable gene switch system effects a change in gene expression at least 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 20-fold, 50-fold, 70-fold, 100-fold, 200-fold, 300 fold, 400-fold or 500-fold greater than all of the other operable gene switches in the host cell. Non-limiting examples of orthogonal gene switch systems are set forth in U.S. Pat. No. 8,105,825 (Publication No. US 2002/0110861 A1).

[0381] As used herein, an "activating ligand" is a compound that binds selectively to the ligand binding domain of a gene switch.

[0382] In one embodiment, the activating ligand is administered to the subject within an hour of the time at which the priming dosage is administered to the subject. In another embodiment, the activating ligand is administered to the subject within about 24, 48, 96, 120, 144 or 168 hours of the time at which the priming dosage is administered to the subject. In another embodiment, the activating ligand is administered to the subject within about 1, 2, 3, 4 or 5 weeks of the time at which the priming dosage is administered to the subject.

[0383] In one embodiment, the activating ligand is administered to the subject within an hour of the time at which the first of the at least one boosting dosage is administered to the subject. In another embodiment, the activating ligand is administered to the subject within about 24, 48, 96, 120, 144 or 168 hours of the time at which the first of the at least one boosting dosage is administered to the subject. In another embodiment, the activating ligand is administered to the subject within about 1, 2, 3, 4 or 5 weeks of the time at which the first of the at least one boosting dosage is administered to the subject.

[0384] In another embodiment, a composition of the invention is contained within a container. In one embodiment, the container is a vial. In another embodiment the container is a multiple-use vial. In another embodiment, the container displays an expiration date for the composition. In another embodiment, the container contains instructions for using the composition.

[0385] In one embodiment, a composition of the invention is a unit dosage composition. In one embodiment, a unit dosage composition is a composition that is manufactured to supply a single dosage of the composition of the invention. In another embodiment, the unit dosage composition is manufactured to provide more than one measured dosages of the composition of the invention.

[0386] The present application also provides an article of manufacture comprising more than one of the unit dosage compositions of the invention. In one embodiment, the article of manufacture is a container. In another embodiment, the article of manufacture is a box. In another embodiment, the article of manufacture displays an expiration date for the unit dosage composition.


[0387] The present invention also provides a kit comprising more than one of the composition or unit dosage of the present invention. In one embodiment, the kit displays an expiration date for the composition or unit dosage. In another embodiment, the kit displays and/or contains instructions for using the composition or unit dosage. In another embodiment, the kit also comprises an activating ligand that binds to the ligand binding domain of the gene switch encoded by the polynucleotide in the composition or unit dosage.

[0388] The present invention also provides a drug label for the composition or unit dosage of the present invention. In one embodiment, the drug label displays an expiration date for the composition or unit dosage. In another embodiment, the drug label displays instructions for using the composition or unit dosage. In another embodiment, the drug label displays the approved indication(s) for the composition or unit dosage. In another embodiment, the said label is in paper form. In another embodiment, the drug label is in digital or computer-readable form.

[0389] The term "activating ligand" as used herein refers to a compound that shows activity as an ecdysone receptor agonist, i.e., a compound that is able to mimic 20-hydroxyecdysone biological activity, and binds to a gene switch ligand binding domain. Activating ligands for use in the present invention include both ecdysteroids and non-steroidal compounds, e.g., tebufenozide and methoxyfenozide.

[0390] In one embodiment, the activating ligand is an ecdysone receptor agonist disclosed in U.S. Pat. No. 8,076,517 (Publication No. 2009/0163592). No. 2009/0298175. No. 2005/0228016 and in U.S. Pat. No. 6,258,603, U.S. Pat. No. 7,375,093, U.S. Pat. No. 7,456,315, U.S. Pat. No. 7,304,161, and U.S. Pat. No. 7,304,162; each of which are hereby incorporated by reference herein.

[0391] In certain embodiments, the activating ligand is a compound having Formula I:



wherein:

[0392] A is alkoxy, arylalkoxy, aryloxy, arylalkyl, optionally substituted aryl or optionally substituted heteroaryl;

[0393] B is optionally substituted aryl or optionally substituted heteroaryl;


[0394] E is CR¹R²R³;

[0395] R¹ is optionally substituted alkyl, arylalkyl, hydroxyalkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted heterocycle, optionally substituted aryl or optionally substituted heteroaryl; and

[0396] R² and R³ are independently hydrogen, optionally substituted alkyl, arylalkyl, hydroxyalkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted heterocycle, optionally substituted aryl or optionally substituted heteroaryl; or

[0397] R¹ and R² taken together form an optionally substituted alkenyl group.

[0398] In one embodiment, the activating ligand is a compound having Formula I:



wherein:

[0399] A is selected from the group consisting of 2,3,6-tri-F-phenyl-; 2,3-di-CH₃-phenyl-; 2,6-di-F-phenyl-; 2-Br, 3,4-ethylenedioxy-phenyl-; 2-CH=CH₂, 3-OCH₃-phenyl-; 2-CH₂CH₃, 3,4-ethylenedioxy-phenyl-; 2-CH₂CH₃, 3-OCH₃-phenyl-; 2-CH₂Cl, 3-OCH₃-phenyl-; 2-CH₂F, 3-OCH₃-phenyl-; 2-CH₂NHCH₃, 3-OCH₃-phenyl-; 2-CH₂NMe₂, 3-OCH₃-phenyl-; 2-CH₂OAc, 3-OCH₃-phenyl-; 2-CH₂OCH₂CH=CH₂, 3-OCH₃-phenyl-; 2-CH₂OH, 3-OCH₃-phenyl-; 2-CH₂OMe, 3-OCH₃-phenyl-; 2-CH₂OMe, 3-OMe-phenyl-; 2-CH₂S(O)₂CH₃, 3-OCH₃-phenyl-; 2-CH₂S(O)CH₃, 3-OCH₃-phenyl-; 2-CH₂SCH₃, 3-OCH₃-phenyl-; 2-CH₃, 3,4-ethylenedioxy-phenyl-; 2-CH₃, 3,4-OCH₂O-phenyl-; 2-CH₃, 3-Ac-phenyl-; 2-CH₃, 3-CH₂CH₂CH₂O-4-phenyl-; 2-CH₃, 3-CH₃-phenyl-; 2-CH₃, 3-Cl-phenyl-; 2-CH₃, 3-Et-phenyl-; 2-CH₃, 3-I-phenyl-; 2-CH₃, 3-NMe₂-phenyl-; 2-CH₃, 3-NO₂-phenyl-; 2-CH₃, 3-OAc-phenyl-; 2-CH₃, 3-OCF₃-phenyl-; 2-CH₃, 3-OCH₂OCH₂-4-phenyl-; 2-CH₃, 3-OCH₃-phenyl-; 2-CH₃, 3-OH-phenyl-; 2-CH₃, 3-Oi-Pr-phenyl-; 2-CH₃, 3-OMe-phenyl-; 2-CH₃, 4,5-methylenedioxy-phenyl-; 2-CH₃, 3-OCH₃-phenyl-; 2-Cl, 3-CH₂OCH₂O-4-phenyl-; 2-CH₃OCH₂O-4-phenyl-; 2-Cl, 3-OMe-phenyl-; 2-Et, 3,4-ethylenedioxy-phenyl-; 2-Et, 3,4-OCH(CH₃)O-phenyl-; 2-Et, 3,4-OCH₂O-phenyl-; 2-Et, 3-OCH₃-phenyl-; 2-F, 3,4-CH₂OCH₂O-phenyl-; 2-F, 4-CH₂CH₃-phenyl-; 2-F, 4-Et-phenyl-; 2-I, 3-OMe-phenyl-; 2-NH₂, 3-OMe-phenyl-; 2-NO₂, 3-OMe-phenyl-; 2-Vinyl, 3-OMe-phenyl-; 3,4-(CH₂)₄-phenyl-; 3,4-di-Et-phenyl-; 3,4-ethylenedioxy-phenyl-; 3,4-OCF₂O-phenyl-; 3,4-OCH(CH₃)O-phenyl-; 3,4-OCH₂O-phenyl-; 3-Cl, 4-Et-phenyl-; 3,4-NH-C≡C-4-phenyl-3-OCH(CH₃)CH₂O-4-phenyl-; 3-OCH₃, 4-CH₃-phenyl-; 3,4-S-C≡N-phenyl-; 4-Br-phenyl-; 4-C(O)CH₃-phenyl-; 4-CH(OH)CH₃-phenyl-; 4-CH₂CH₃-phenyl-;

4-CH₂CN-phenyl-; 4-CH₃-phenyl-; 4-Cl-phenyl-; 4-Et-phenyl-; 4-OCH-phenyl-; phenyl-; and benzo[1,2,5]oxadiazole-5-yl;

[0400] B is selected from the group consisting of 1-trityl-5-benzimidazolyl-; 3-trityl-5-benzimidazolyl-1H-indazole-3-yl-; 1-methyl-1H-indole-2-yl-; 1-methyl-2-oxo-6-trifluoromethyl-3-pyridyl-; 1-trityl-1H-indazole-3-yl-; 2,3,4,5-phenyl-; 2,3,4,5-tetra-F-phenyl-; 2,3,4-F-phenyl-; 2,3-F-phenyl-; 2,3-OCH₂O-phenyl-; 2,4,5-F-phenyl-; 2,4-di-Cl-5-F-phenyl-; 2,5-di-OCH₃-phenyl-; 2,5-F-phenyl-; 2,6-di-Cl-4-pyridyl-; 2,6-dimethoxy-4-pyrimidinyl-2,6-di-OCH₃-3-pyridyl-2,6-F-phenyl-; 2-Cl, 5-NO₂-phenyl-; 2-Cl-3-pyridyl-2-Cl-4-F-phenyl-; 2-Cl-5-CH₃-phenyl-; 2-Cl-6-CH₃-4-pyridyl-; 2-Et-phenyl-; 2-F, 4-Cl-phenyl-; 2-F, 5-CH₃-phenyl-; 2-methoxy-6-trifluoromethyl-3-pyridyl-; 2-NO₂-3,5-di-OCH₃, 4-CH₃-phenyl-; 2-NO₂-4-Cl-phenyl-; 2-NO₂-5-CH₃-phenyl-; 2-NO₂-5-Cl-phenyl-; 2-NO₂-5-F-phenyl-; 2-NO₂-phenyl-; 2-OCH₂CF₃, 5-OCH₃-phenyl-; 2-OCH₃-3-pyridyl-2-OCH-4-CH₃-phenyl-; 2-OCH₃-4-Cl-phenyl-; 2-OCH₃-4-F-phenyl-; 2-OCH₃-5-CH₃-phenyl-; 2-OCH₃-5-Cl-phenyl-; 2-OCH₃-phenyl-; 2-S(O)CH₃-phenyl-; 2-SO₃H-phenyl-; 3,4,5-F-phenyl-; 3,4,5-tri-OCH₃-phenyl-; 3,4-di-CH₃-5-Cl-phenyl-; 3,4-F-phenyl-; 3,4-methylenedioxy-phenyl-; 3,5-di(CH₂OH)-phenyl-; 3,5-di-CH₃-4-Cl-phenyl-; 3,5-di-CH₃-phenyl-; 3,5-di-Cl-F-phenyl-; 3,5-di-Cl-phenyl-; 3,5-di-CO₂H-phenyl-; 3,5-di-F-phenyl-; 3,5-di-OCH₃, 4-CH₃-phenyl-; 3,5-OCH₃-4-OAc-phenyl-; 3,5-di-OCH₃-phenyl-; 3,6-dichloro-4-pyridazinyl-; 3,6-dimethoxy-4-pyridazinyl-; 3-Br-phenyl-; 3-CF₃, 5-F-phenyl-; 3-CF₃-4-F-phenyl-; 3-CF₃-4-F-phenyl-3-CF₃-phenyl-; 3-CH=NNHCOCONH₂, 5-CH₃-phenyl-; 3-CH=NNHCONH₂, 5-CH₃-phenyl-; 3-CH=NOH, 5-CH₃-phenyl-; 3-CH₂OAc, 5-CH₃-phenyl-; 3-CH₃, 5-Br-phenyl-; 3-CH₃, 5-CH₃-phenyl-; 3-CH₃, 5-Cl-phenyl-; 3-CH₃-4-Br-phenyl-; 3-CH₃-phenyl-; 3-chloro-6-methylsulfanyl-pyrazine-2-yl-; 3-Cl, 5-Br-phenyl-; 3-Cl, 5-Cl-phenyl-; 3-Cl-5-OCH₃-4-pyridyl-; 3-Cl-phenyl-; 3-CN-phenyl-; 3-F, 5-F-phenyl-; 3-F-phenyl-; 3-NO₂-phenyl-; 3-OCH₃-4-CH₃-phenyl-; 3-OCH₃-4-pyridyl-; 3-OCH₃-phenyl-; 3-OMe, 5-CH₃-phenyl-; 3-OMe, 5-OMe-phenyl-; 3-oxo-6-methoxy-4-pyridazinyl-; 4,6-dimethyl-pyridyl-; 4-CH₂-phenyl-; 4-F-phenyl-; 4-pyridazinyl-; 5-benzimidazolyl-; 5-methoxycarbonyl-2-pyridyl-; 5-methyl-1-phenyl-1H-pyrazole-3-yl-; 5-methyl-pyrazine-2-yl-; 6-CH₃-2-pyridyl-phenyl-; and pyrazine-2-yl; and

[0401] E is selected from the group consisting of C(CH₃)₂C(O)OEt; C(CH₃)₂CH=NCH₂CH₂OH; C(CH₃)₂CH=NNHC(O)C(O)NH₂; C(CH₃)₂CH=NNHC(O)NH₂; C(CH₃)₂CH=NOH; C(CH₃)₂CH₂OC(O)CH₃C(CH₃)₂CH₂OCH₃; C(CH₃)₂CH₂OH; C(CH₃)CH₂OSi(CH₃)₂tBu; C(CH₃)₂CHO; C(CH₃)₂CN; C(CH₃)₂COOH; CH(CH₃)C(CH₃)₃; CH(Et)(n-Bu); CH(Et)(t-Bu); CH(n-Bu)(t-Bu); CH(n-Pr)(t-Bu); CH(Ph)(t-Bu); and t-Bu.

[0402] In another embodiment, the activating ligand is a compound having Formula I wherein A, B, and E are defined according to Table 3.

TABLE 3

Ligand Components		
A	B	E
4-Cl—Ph	Ph	t-Bu
4-Et—Ph	2-NO ₂ —Ph	t-Bu
4-CH ₃ —Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
4-Et—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2,6-di-F—Ph	3-Cl, 5-Cl—Ph	t-Bu
2-CH ₃ , 3-Cl—Ph	3-Cl—Ph	t-Bu
2-Cl, 3-OMe—Ph	2-Cl-5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-Cl—Ph	3-CH ₃ -4-Br—Ph	t-Bu
4-Et—Ph	3,5-di-CH ₃ -4-Cl—Ph	t-Bu
4-Et—Ph	3,4-di-CH ₃ -5-Cl—Ph	t-Bu
4-OCH ₃ —Ph	2-Cl-4-F—Ph	t-Bu
4-Et—Ph	3-CH ₃ , 5-Cl—Ph	t-Bu
4-Et—Ph	2-Et—Ph	t-Bu
4-OCH ₃ —Ph	3-Cl, 5-Cl—Ph	t-Bu
4-Et—Ph	2-NO ₂ -5-CH ₃ —Ph	t-Bu
4-CH ₂ CN—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	3-CH ₃ —Ph	t-Bu
4-Br—Ph	3-Cl, 5-Cl—Ph	t-Bu
2-CH ₃ , 3-NO ₂ —Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₃ —Ph	2,5-di-OCH ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₃ —Ph	2-OCH ₃ -5-Cl—Ph	t-Bu
2-NO ₂ , 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₃ —Ph	3-OMe, 5-OMe—Ph	t-Bu
3-Cl, 4-Et—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
4-CH(OH)CH ₃ —Ph	3-F, 5-F—Ph	t-Bu
2-CH ₃ , 3-NMe ₂ —Ph	3-Cl, 5-Cl—Ph	t-Bu
2-CH ₃ , 3-Ac—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OAc—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-I—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	3-Cl, 5-Br—Ph	t-Bu
2-CH ₃ , 3-Oi-Pr—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OCH ₃ —Ph	2-Cl-3-pyridyl	t-Bu
2-CH ₃ , 3-OMe—Ph	2-OCH ₃ -5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	2,5-F—Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	2-Et—Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	3-CH ₃ , 5-Br—Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	3-OMe, 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	2-OCH ₃ -4-Cl—Ph	t-Bu
2-CH ₃ , 3-OCF ₃ —Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	3-OCH ₃ -4-CH ₃ —Ph	t-Bu
3-OCH ₃ , 4-CH ₃ —Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	2-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OCH ₃ —Ph	2,6-di-Cl-4-pyridyl	t-Bu
2-CH ₃ , 3-OMe—Ph	2-NO ₂ -5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	2-F-4-Cl—Ph	t-Bu
3,4-OCH ₂ O—Ph	2-Cl-4-F—Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-Et—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
3-CH ₂ CH ₂ O-4-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	3,5-di-Cl-4-F—Ph	t-Bu
2-CH ₃ , 3,4-OCH ₂ O—Ph	4-F—Ph	t-Bu
2-Et, 3,4-OCH ₂ O—Ph	2-OCH ₃ —Ph	t-Bu
3,4-di-Et—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	4-F—Ph	t-Bu
2-Et, 3-OMe—Ph	2-OCH ₃ —Ph	t-Bu
2-CH ₃ , 3-OMe—Ph	2-OCH ₃ -4-F—Ph	t-Bu
2-Et, 3-OCH ₃ —Ph	2-Cl-6-CH ₃ -4-pyridyl	t-Bu
2-Et, 3-OMe—Ph	3-OMe, 5-OMe—Ph	t-Bu
2-I, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
3,4-ethylenedioxy-Ph	2-OCH ₃ —Ph	t-Bu
3,4-(CH ₂) ₄ —Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	2,3-OCH ₂ O—Ph	t-Bu
2-F, 4-Et—Ph	4-F—Ph	t-Bu
2-Et, 3-OMe—Ph	3,4-methylenedioxy-Ph	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	4-F—Ph	t-Bu
3,4-OCH(CH ₃)O—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Et, 3,4-OCH(CH ₃)O—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	3-OCH ₃ —Ph	t-Bu
3-OCH(CH ₃)CH ₂ O-4-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Br, 3,4-ethylenedioxy-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	3-CH ₃ , 5-Cl—Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	3-CH ₃ —Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2-OCH ₃ —Ph	t-Bu

TABLE 3-continued

Ligand Components		
A	B	E
2-Et, 3,4-ethylenedioxy-Ph	3-OCH ₃ —Ph	t-Bu
3-S—C≡N-4-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	2-OCH ₃ -4-Cl—Ph	t-Bu
2-Et, 3-OMe—Ph	2,5-di-OCH ₃ —Ph	t-Bu
2-CH ₃ , 4,5-methylenedioxy-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
3-CH ₂ OCH ₂ O-4-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-OCH ₂ OCH ₂ -4-Ph	2-OCH ₃ —Ph	t-Bu
2-Et, 3-OCH ₂ OCH ₂ -4-Ph	4-F—Ph	t-Bu
2-Cl 4,5-methylenedioxy-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2,3,6-tri-F—Ph	2-Cl-4-F—Ph	t-Bu
2-Et, 3-OMe—Ph	2,6-F—Ph	t-Bu
2-Et, 3-OMe—Ph	3-F—Ph	t-Bu
2-Et, 3-OMe—Ph	3-Br—Ph	t-Bu
2-Et, 3-OMe—Ph	2-NO ₂ —Ph	t-Bu
2-Et, 3-OMe—Ph	2,3-F—Ph	t-Bu
2-Et, 3-OMe—Ph	3,4,5-tri-OCH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-CF ₃ , 5-F—Ph	t-Bu
2-Et, 3-OMe—Ph	3-CN—Ph	t-Bu
2-Vinyl, 3-OMe—Ph	2,4-di-Cl-5-F—Ph	t-Bu
2-Et, 3-OCH ₂ OCH ₂ -4-Ph	Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ C(O)OEt
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ CH ₂ OH
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ CHO
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ CH ₂ OCH ₃
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ CH=NOH
2-NH ₂ , 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH ₂ OAc, 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ CH ₂ OC(O)CH ₃
2-CH ₃ , 3-OH—Ph	2,3,4-F—Ph	t-Bu
2-CH ₃ , 3-OH—Ph	3-Cl-5-OCH ₃ -4-pyridyl	t-Bu
2-CH ₃ , 3-OH—Ph	2,6-di-Cl-4-pyridyl	t-Bu
2-CH ₃ , 3-OH—Ph	3-OCH ₃ -4-pyridyl	t-Bu
2-CH ₃ , 3-OH—Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2-OCH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	2,4-di-Cl-5-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2,4-di-Cl-5-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2-F, 5-CH ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2,5-F—Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2,3,4-F—Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2,3,4,5--Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	3-CF ₃ -4-F—Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2,6-di-Cl-4-pyridyl	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2-OCH ₃ —Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2,4-di-Cl-5-F—Ph	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	2-F, 4-Cl—Ph	t-Bu
2-CH ₃ , 3-OAc—Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	2-OCH ₃ -5-Cl—Ph	t-Bu
2-Et, 3,4-OCH ₂ O—Ph	2-OCH ₃ -4-Cl—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2-OCH ₃ -5-Cl—Ph	t-Bu
2-Et, 3-OMe—Ph	2-NO ₂ -5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	2-NO ₂ -4-Cl—Ph	t-Bu
2-Et, 3-OMe—Ph	2-NO ₂ -5-Cl—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2-NO ₂ -5-CH ₃ —Ph	t-Bu
Benzo[1,2,5]oxadiazole-5-yl	2-OCH ₃ -4-Cl—Ph	t-Bu
2-Vinyl, 3-OMe—Ph	2-Cl, 5-NO ₂ —Ph	t-Bu
2-Vinyl, 3-OMe—Ph	2-OCH ₃ -4-Cl—Ph	t-Bu
2-Et, 3-OCH ₃ —Ph	1-methyl-1H-indole-2-yl	t-Bu
2-Et, 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-Cl, 3-CH ₂ OCH ₂ O-4-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
2-F, 4-Et—Ph	3-NO ₂ —Ph	t-Bu
2-F, 4-Et—Ph	3-OCH ₃ —Ph	t-Bu
2-Cl, 3-CH ₂ OCH ₂ O-4-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-F, 4-Et—Ph	2,6-di-Cl-4-pyridyl	t-Bu
2-F, 4-Et—Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-F, 4-Et—Ph	3,4,5-F—Ph	t-Bu
2-F, 4-Et—Ph	3-CH ₃ —Ph	t-Bu
2-F, 4-Et—Ph	2-OCH ₃ —Ph	t-Bu
2-F, 4-Et—Ph	2-NO ₂ -5-F—Ph	t-Bu
2-F, 4-Et—Ph	2-OCH ₂ CF ₃ , 5-OCH ₃ —Ph	t-Bu
2-F, 4-Et—Ph	2-Cl-6-CH ₃ -4-pyridyl	t-Bu
2-F, 4-Et—Ph	2,6-di-OCH ₃ -3-pyridyl	t-Bu
3-NH—C≡C-4-Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu

TABLE 3-continued

Ligand Components		
A	B	E
2-Et, 3-OMe—Ph	2-S(O)CH ₃ —Ph	t-Bu
3,4-OCF ₂ O—Ph	2-NO ₂ —Ph	t-Bu
3,4-OCF ₂ O—Ph	3-CH ₃ , 5-CH ₃ —Ph	t-Bu
3,4-OCF ₂ O—Ph	3-OCH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-Br—Ph	—C(CH ₃) ₂ CN
2-CH ₃ OMe, 3-OMe—Ph	3,5-di-Cl—Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH=NOH, 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH=NNHCONH ₂ , 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH=NNHCOCONH ₂ , 5-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3-CH ₃ , 5-CH ₃ —Ph	—C(CH ₃) ₂ CN
2-Et, 3-OMe—Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—C(CH ₃) ₂ CN
2-Et, 3-OCH ₂ OCH ₂ -4-Ph	2-OCH ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2,4,5-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	3,4,5-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	3-F—Ph	t-Bu
2-Et, 3,4-OCH ₂ O—Ph	3-CF ₃ —Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	4-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	3,4-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	3,5-di-F—Ph	t-Bu
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	2,3,4,5-tetra-F—Ph	t-Bu
2-Et, 3-OCH ₂ OCH ₂ -4-Ph	4-CH ₃ —Ph	t-Bu
2-Et, 3-OMe—Ph	3,5-di-OCH ₃ -4-OAc—Ph	t-Bu
2-Et, 3-OMe—Ph	3,5-di-OCH ₃ -OH—Ph	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	2,6-di-OCH ₃ -3-pyridyl	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	2,6-di-Cl-4-pyridyl	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	3-F—Ph	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	3-CF ₃ , 5-F—Ph	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	2-NO ₂ -5-CH ₃ —Ph	t-Bu
2-ethyl, 3-methoxy	4,6-dimethyl-pyridyl	t-Bu
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	2-methoxy-6-trifluoromethyl-3-pyridyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	1-methyl-2-oxo-6-trifluoromethyl-3-pyridyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	2,6-dimethoxy-4-pyrimidinyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,6-dimethoxy-4-pyridazinyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,6-dichloro-4-pyridazinyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	4-pyridazinyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3-oxo-6-methoxy-4-pyridazinyl	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,5-di-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,5-di-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ —Ph	3,5-di-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ —Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ —Ph	3,5-di-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ —Ph	3,5-di-OCH ₃ -4-CH ₃ —Ph	—CH(n-Pr)C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃ —Ph	3,5-di-CH ₃ —Ph	—CH(Et)C(CH ₃) ₃
2-CH ₃ , 3-OH—Ph	3-OCH ₃ -4-pyridyl	—C(CH ₃) ₃
4-CH(OH)CH ₃ —Ph	3,5-di(CH ₂ OH)—Ph	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ —Ph	2-S(O)CH ₃ —Ph	—C(CH ₃) ₃
4-C(O)CH ₃ —Ph	3,5-di-CO ₂ H—Ph	—C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	2,6-di-OCH ₃ -3-pyridyl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3,4-ethylenedioxy-Ph	3-CF ₃ -4-F-phenyl	—C(CH ₃) ₃
2-F, 4-CH ₂ CH ₃ —Ph	3,5-di-CH ₃ —Ph	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ —Ph	2-SO ₃ H—Ph	—C(CH ₃) ₃
2-CH ₃ , 3-CH ₂ CH ₂ CH ₂ O-4-Ph	3,5-di-CH ₃ —Ph	—C(CH ₃) ₃
4-CH ₂ CH ₃ —Ph	3,5-di-CH ₃ —Ph	—CH(CH ₃)C(CH ₃) ₃
2-CH ₂ CH ₃ , 3,4-ethylenedioxy-Ph	3-CH ₃ —Ph	—C(CH ₃) ₃
2,3-di-CH ₃ —Ph	Ph	—CH(Et)(n-Bu)
2,3-di-CH ₃ —Ph	3-CH ₃ —Ph	—CH(Et)(t-Bu)
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ , 4-OH—Ph	—C(CH ₃) ₃
2-F, 3-CH ₂ OCH ₂ O-4-Ph	3,5-di-CH ₃ —Ph	—C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	2-S(O)CH ₃ —Ph	—C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-OCH ₃ , 4-CH ₃ —Ph	—C(CH ₃) ₂ CN

TABLE 3-continued

Ligand Components		
A	B	E
2-CH ₂ CH ₃ -3-OCH ₃ -Ph	6-CH ₃ -2-pyridyl-	—C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	2-NO ₂ -3,5-di-OCH ₃ , 4-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₃ , 3,4-ethylenedioxy-Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
4-CH ₂ CH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₃ -3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
4-CH ₂ CH ₃ -Ph	3,5-di-CH ₃ -Ph	—CH(Et)(t-Bu)
4-CH ₂ CH ₃ -Ph	2-OCH ₃ -3-pyridyl	—CH(Et)(t-Bu)
4-CH ₂ CH ₃ -Ph	3,5-di-CH ₃ -Ph	—CH(n-Bu)(t-Bu)
4-CH ₂ CH ₃ -Ph	3,5-di-OCH ₃ , 4-CH ₃ -Ph	—CH(n-Bu)(t-Bu)
4-CH ₂ CH ₃ -Ph	2-OCH ₃ -3-pyridyl	—CH(n-Bu)(t-Bu)
4-CH ₂ CH ₃ -Ph	3,5-di-CH ₃ -Ph	—CH(Ph)(t-Bu)
4-CH ₂ CH ₃ -Ph	3,5-di-OCH ₃ , 4-CH ₃ -Ph	—CH(Ph)(t-Bu)
4-CH ₂ CH ₃ -Ph	2-OCH ₃ -3-pyridyl	—CH(Ph)(t-Bu)
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	5-benzimidazolyl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	1-(or 3-)-trityl-5-benzimidazolyl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	5-methyl-1-phenyl-1H-pyrazole-3-yl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	3-chloro-6-methylsulfanyl-pyrazine-2-yl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	1H-indazole-3-yl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	1-trityl-1H-indazole-3-yl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	5-methoxycarbonyl-2-pyridyl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	pyrazine-2-yl	—C(CH ₃) ₃
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₂ CH ₂ OSi(CH ₃) ₂ tBu
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₂ CH=NNHC(CH ₃) ₂ OH
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₂ CH=NNHC(O)NH ₂
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₂ CH=NNHC(O)C(O)NH ₂
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₂ COOH
2-CH ₂ S(O)CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₃ S(O) ₂ CH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ NMe ₂ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ NHCH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH=CH ₂ , 3-OCH ₃ -Ph—	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ OMe, 3-OCH ₃ -Ph—	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ SCH ₃ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ OCH ₂ CH=CH ₂ , 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ Cl, 3-OCH ₃ -Ph—	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ OH, 3-OCH ₃ -Ph—	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ OAc, 3-OCH ₃ -Ph	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₂ F, 3-OCH ₃ -Ph—	3,5-di-CH ₃ -Ph	—C(CH ₃) ₃
2-CH ₃ , 3-OCH ₃	3,5-di-CH ₃	—CH(n-Bu)(t-Bu)
2-CH ₃ , 3-OCH ₃	3,5-di-OCH ₃ , 4-CH ₃	—CH(n-Bu)(t-Bu)
2-CH ₂ CH ₃ , 3-OCH ₃ -Ph	5-Methyl-pyrazine-2-yl-	—C(CH ₃) ₃

[0403] In another embodiment, the activating ligand is a compound having Formula I selected from the group consisting of:

[0404] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(3-hydroxymethyl-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0405] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-[3-(tert-butyl-dimethyl-silanyloxymethyl)-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl]-hydrazide;

[0406] 7-[N'-tert-Butyl—N-(3,5-dimethyl-benzoyl)-hydrazinocarbonyl]-8-methyl-2,3-dihydro-benzo[1,4]dioxine-2-carboxylic acid;

[0407] 7-[N'-tert-Butyl-N'-(3,5-dimethyl-benzoyl)-hydrazinocarbonyl]-8-methyl-2,3-dihydro-benzo[1,4]dioxine-2-carboxylic acid methyl ester;

[0408] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(3-semicarbazidomethyl-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0409] Phenyl-carbamic acid 7-[N'-tert-butyl-N'-(3,5-dimethylbenzoyl)-hydrazinocarbonyl]-8-methyl-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl ester;

[0410] 3,5-Dimethyl-benzoic acid N'-[3-(2-amino-ethyl)-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl]-N-tert-butyl-hydrazide;

[0411] 7-[N'-tert-Butyl-N'-(3,5-dimethyl-benzoyl)-hydrazinocarbonyl]-8-methyl-2,3-dihydro-benzo[1,4]dioxine-2-carboxylic acid pentafluorophenyl ester;

[0412] 7-[N'-tert-butyl-N'-(3,5-dimethyl-benzoyl)-hydrazinocarbonyl]-8-methyl-2,3-dihydro-benzo[1,4]dioxine-2-carboxylic acid methylamide;

[0413] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(3-formyl-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0414] Toluene-4-sulfonic acid 7-[N'-tert-butyl-N'-(3-dimethylbenzoyl)-hydrazinocarbonyl]-8-methyl-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl ester;

[0415] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(3-hydroxylamino-methyl)-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0416] 3,5-Dimethyl-benzoic acid N'-tert-butyl-N'-(3-cyanomethyl-5-methyl-2,3-dihydro benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0417] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(5-methyl-3-methylsulfanylmethyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0418] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(3-methanesulfonylmethyl-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0419] 3,5-Dimethyl-benzoic acid N-tert-butyl-N'-(3-fluoromethyl-5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;

[0420] 3,5-Dimethyl-benzoic acid N-(1-tert-butyl-heptyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide;

[0421] 3,5-Dimethyl-benzoic acid N-(1-tert-butyl-heptyl)-N'-(4-ethyl-benzoyl)-hydrazide;

[0422] 3,5-Dimethoxy-4-methyl-benzoic acid-N-(1-tert-butyl-heptyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide;

[0423] 3,5-Dimethoxy-4-methyl-benzoic acid-N-(1-tert-butyl-heptyl)-N'-(4-ethyl-benzoyl)-hydrazide;

[0424] 2-Methoxy-nicotinic acid N-(1-tert-butyl-heptyl)-N'-(4-ethyl-benzoyl)-hydrazide;


[0425] 3,5-Dimethyl-benzoic acid N-(1-tert-butyl-3,4,4-trimethyl-pent-2-enyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide;

[0426] 3,5-Dimethyl-benzoic acid N-(1-tert-butyl-2-cyano-vinyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide;

[0427] 3,5-Dimethyl-benzoic acid N-(1-butyl-2,2-dimethyl-pentyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide; and

[0428] 3,5-Dimethyl-benzoic acid N-(1-butyl-2,2-dimethyl-pent-4-enyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide.

[0429] In another embodiment, the activating ligand is an enantiomerically enriched compound having Formula II: U, 2



II

wherein:

[0430] A is alkoxy, arylalkyloxy, aryloxy, arylalkyl, optionally substituted aryl or optionally substituted heteroaryl;


[0431] B is optionally substituted aryl or optionally substituted heteroaryl; and

[0432] R¹ and R² are independently optionally substituted alkyl, arylalkyl, hydroxyalkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted heterocycle, optionally substituted aryl or optionally substituted heteroaryl;

[0433] with the proviso that R¹ does not equal R²;

[0434] wherein the absolute configuration at the asymmetric carbon atom bearing R¹ and R² is predominantly S.

[0435] In another embodiment, the activating ligand is an enantiomerically enriched compound having Formula III:



III

wherein:

[0436] A is alkoxy, arylalkyloxy, aryloxy, arylalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

[0437] B is optionally substituted aryl or optionally substituted heteroaryl; and


[0438] R¹ and R² are independently optionally substituted alkyl, arylalkyl, hydroxyalkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted heterocycle, optionally substituted aryl or optionally substituted heteroaryl;

[0439] with the proviso that R¹ does not equal R²;


[0440] wherein the absolute configuration at the asymmetric carbon atom bearing R¹ and R² is predominantly R.

[0441] In another embodiment, the activating ligand is an enantiomerically enriched compound having Formula III wherein:

[0442] A is:



[0443] B is:



[0444] R^{3a}, R^{3b}, R^{3c}, R^{3d}, R^{3e}, R^{3f}, R^{3g}, R^{3h}, R³ⁱ and R^{3j} are independently selected from hydrogen, halo, (C₁-C₄)alkyl, or (C₁-C₄)alkoxy;

[0445] R¹ is (C₁-C₆)alkyl, hydroxy(C₁-C₄)alkyl, or (C₂-C₄)alkenyl; and


[0446] R² is optionally substituted (C₁-C₆)alkyl.

[0447] In another embodiment, the activating ligand is a compound having Formula III selected from the group consisting of:

[0448] (R)-N'-(1-ter-Butyl-butyl)-N'-(3,5-dimethyl-benzoyl)-hydrazinecarboxylic acid benzyl ester;

- [0449] (R)—N'-(1-tert-Butyl-butyl)-N'-(3,5-dimethyl-benzoyl)-hydrazinecarboxylic acid tert-butyl ester;
- [0450] (R)—N'-(1-tert-Butyl-4-hydroxy-butyl)-N'-(3,5-dimethyl-benzoyl)-hydrazine carboxylic acid benzyl ester;
- [0451] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-ethyl-3-methoxy-benzoyl)-hydrazide;
- [0452] (R)—3,5-Dimethyl-benzoic acid N'-benzoyl-N-(1-tert-butyl-butyl)-hydrazide;
- [0453] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-methyl-benzoyl)-hydrazide;
- [0454] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-methoxy-benzoyl)-hydrazide;
- [0455] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-fluoro-benzoyl)-hydrazide;
- [0456] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-chloro-benzoyl)-hydrazide;
- [0457] (R)-3,5-Dimethyl-benzoic acid N'-(2-bromo-benzoyl)-N-(1-tert-butyl-butyl)-hydrazide;
- [0458] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methyl-benzoyl)-hydrazide;
- [0459] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methoxy-benzoyl)-hydrazide;
- [0460] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-chloro-benzoyl)-hydrazide;
- [0461] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-methyl-benzoyl)-hydrazide;
- [0462] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-ethyl-benzoyl)-hydrazide;
- [0463] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-methoxy-benzoyl)-hydrazide;
- [0464] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-chloro-benzoyl)-hydrazide;
- [0465] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2,6-difluoro-benzoyl)-hydrazide;
- [0466] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2,6-dichloro-benzoyl)-hydrazide;
- [0467] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3,4-dimethoxy-benzoyl)-hydrazide;
- [0468] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3,5-difluoro-benzoyl)-hydrazide;
- [0469] (R)-3,5-Dimethyl benzoic acid N-(1-tert-butyl-butyl)-N'-(3,5-dimethoxy-4-methyl-benzoyl)-hydrazide;
- [0470] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-methyl-benzo[1,3]dioxole-5-carbonyl)-hydrazide;
- [0471] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(5-methyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;
- [0472] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(5-ethyl-2,3-dihydro-benzo[1,4]dioxine-6-carbonyl)-hydrazide;
- [0473] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(naphthalene-1-carbonyl)-hydrazide;
- [0474] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(naphthalene-2-carbonyl)-hydrazide;
- [0475] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(thiophene-2-carbonyl)-hydrazide;
- [0476] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2,5-dimethyl-furan-3-carbonyl)-hydrazide;
- [0477] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-chloro-pyridine-3-carbonyl)-hydrazide;
- [0478] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(6-chloro-pyridine-3-carbonyl)-hydrazide;

- [0479] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide;
- [0480] (R)-3,5-Dimethoxy-4-methyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide; and
- [0481] (R)-3,5-Dimethyl-benzoic acid N'-(4-ethyl-benzoyl)-N-(1-phenethyl-but-3-enyl)-hydrazide.
- [0482] In another embodiment, the activating ligand is a compound having Formula IV:



IV

wherein:

- [0483] Q is O or S;
- [0484] R¹ is selected from the group consisting of hydrogen, (C₁-C₁₂)alkyl, (C₃-C₁₂)cycloalkyl, (C₃-C₁₂)cycloalkyl(C₁-C₃)alkyl, (C₁-C₁₂)haloalkyl, (C₂-C₁₂)alkenyl, (C₃-C₁₂)cycloalkenyl, (C₂-C₁₂)haloalkenyl, (C₂-C₁₂)alkynyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₆)alkylthio(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, succinimidylmethyl, benzosuccinimidylmethyl, optionally substituted phenyl, optionally substituted 1-naphthyl, optionally substituted 2-naphthyl, optionally substituted phenyl(C₁-C₃)alkyl, optionally substituted phenyl(C₂-C₃)alkenyl, optionally substituted naphthyl(C₁-C₃)alkyl, optionally substituted phenoxy (C₁-C₃)alkyl, optionally substituted phenylamino, and optionally substituted heterocycle;
- [0485] R² and R³ are each independently selected from hydrogen, (C₁-C₆)alkyl, and (C₁-C₆)haloalkyl;
- [0486] R⁴ is hydrogen, (C₁-C₆)alkyl, or (C₁-C₆)haloalkyl;
- [0487] R⁵ and R⁶ are each independently selected from the group consisting of hydrogen, (C₁-C₁₂)alkyl, (C₃-C₁₂)cycloalkyl, (C₁-C₁₂)haloalkyl, (C₂-C₁₂)alkenyl, (C₃-C₁₂)cycloalkenyl, (C₂-C₁₂)haloalkenyl, (C₂-C₁₂)alkynyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₆)alkylthio (C₁-C₆)alkyl, aminocarbonyl, aminothiocarbonyl, formyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylcarbonyl, cyclo(C₃-C₆)alkylcarbonyl, halo (C₁-C₆)alkylcarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkoxycarbonylcarbonyl, or phenyl(C₂-C₃)alkenylcarbonyl, optionally substituted phenyl, optionally substituted 1-naphthyl, optionally substituted 2-naphthyl, optionally substituted phenyl(C₁-C₃)alkyl, optionally substituted phenyl(C₂-C₃)alkenyl, optionally substituted phenylcarbonyl, or optionally substituted heterocycle;
- [0488] R⁷, R⁸, R⁹, and R¹⁰ are each independently selected from the group consisting of hydrogen, cyano, nitro, halogen, (C₁-C₁₂)alkyl, (C₃-C₁₂)cycloalkyl, (C₁-C₁₂)haloalkyl, (C₂-C₁₂)alkenyl, (C₃-C₁₂)cycloalkenyl, (C₂-C₁₂)haloalkenyl, (C₂-C₁₂)alkynyl, halo(C₂-C₆)

alkyl, hydroxy, (C₁-C₆)alkoxy, halo(C₁-C₆)alkoxy, (C₂-C₆)alkenyl, halo(C₂-C₆)alkenyl, (C₂-C₆)alkynyl, halo(C₂-C₆)alkynyl, aryloxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₆)alkylthio, halo(C₁-C₆)alkylthio, (C₂-C₆)alkenylthio, halo(C₂-C₆)alkenylthio, (C₂-C₆)alkynylthio, halo(C₂-C₆)alkynylthio, (C₁-C₆)alkylsulfinyl, halo(C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, halo(C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₃)alkoxy(C₁-C₃)alkyl, (C₁-C₆)alkylthio(C₁-C₆)alkyl, (C₁-C₆)alkylsulfinyl(C₁-C₃)alkyl, (C₁-C₃)alkylsulfinyl(C₁-C₃)alkyl, (C₁-C₃)alkylamino(C₁-C₃)alkyl, di(C₁-C₂)alkylamino(C₁-C₃)alkyl, (C₁-C₆)alkylcarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, or (C₁-C₆)alkoxycarbonyl, optionally substituted phenyl, optionally substituted 1-naphthyl, optionally substituted 2-naphthyl, optionally substituted phenyl(C₁-C₃)alkyl, optionally substituted phenyl(C₂-C₃)alkenyl, or optionally substituted heterocycle.

[0489] In another embodiment, the activating ligand is a compound having Formula IV wherein:

[0490] Q is O;

[0491] R¹ is selected from the group consisting of 4-fluorophenyl, 3-fluorophenyl, 4-fluoro-3-methylphenyl, 4-fluoro-3-(trifluoromethyl)phenyl, 4-fluoro-3-iodophenyl, 3-fluoro-4-iodophenyl, 3,4-di-fluorophenyl, 4-ethylphenyl, 3-fluoro-4-methylphenyl, 3-fluoro-4-ethylphenyl, 3-chloro-4-fluorophenyl, 3-fluoro-4-chlorophenyl, 2-methyl-3-methoxyphenyl, 2-ethyl-3-methoxyphenyl, 2-ethyl-3,4-ethylenedioxyphenyl, 3-nitrophenyl, 4-iodophenyl, 3-fluoro-4-trifluoromethylphenyl, 3-methylphenyl, 4-methylphenyl, 4-chlo-

rophenyl, 3-trifluoromethylphenyl, 3-methoxyphenyl, 3-chloro-6-pyridyl, 2-chloro-4-pyridyl, phenylamino, 3-chlorophenylamino, 3-methylphenylamino, 4-chlorophenylamino, and 4-methylphenylamino;

[0492] R² is hydrogen, methyl or CF₃;

[0493] R³ is hydrogen, methyl or CF₃;

[0494] R⁴ is hydrogen;

[0495] R⁵ is optionally substituted phenyl, wherein the substituents are selected from the group consisting of cyano, nitro, halogen, (C₁-C₃)alkyl, halo(C₁-C₃)alkyl, (C₁-C₃)alkoxy, halo(C₁-C₃)alkoxy, (C₃)alkenyl, (C₃)alkynyl, (C₁-C₃)alkylthio, halo(C₁-C₃)alkylthio, (C₁-C₃)alkylsulfinyl, halo(C₁-C₃)alkylsulfinyl, (C₁-C₃)alkylsulfonyl, halo(C₁-C₃)alkylsulfonyl, (C₁-C₃)alkoxy(C₁-C₃)alkyl, (C₁-C₂)alkylthio(C₁-C₂)alkyl, and (C₁-C₃)alkoxycarbonyl;

[0496] R⁶ is selected from the group consisting of hydrogen, formyl, (C₁-C₃)alkylcarbonyl, and cyclo(C₃-C₆)alkylcarbonyl; and

[0497] R⁷, R⁸, R⁹, R¹⁰ are independently selected from the group consisting of hydrogen, cyano, nitro, chlorine, fluorine, methyl, trifluoromethyl, difluoromethyl, methoxy, trifluoromethoxy, difluoromethoxy, methylthio, trifluoromethylthio, difluoromethylthio, methylsulfinyl, trifluoromethylsulfinyl, difluoromethylsulfinyl, methylsulfonyl, trifluoromethylsulfonyl, difluoromethylsulfonyl, methoxymethyl, and methoxycarbonyl, or R⁷/R⁸, R⁸/R⁹, or R⁹/R¹⁰ form a 5- or 6-membered heterocyclic ring.

[0498] In another embodiment, the activating ligand is a compound having Formula IV wherein Q is O, R² is methyl and R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, and R¹⁰ are defined according to Table 4.

TABLE 4

Ligand Components					
R ¹	R ³ , R ⁴ , R ⁷ , R ⁹ , and R ¹⁰	R ⁵	R ⁶	R ⁸	Stereo. ¹
n-Hexyl	H	Ph	H	H	cis
n-Heptyl	H	Ph	H	H	cis
n-Bu	H	Ph	H	H	cis
3-CF ₃ -4-F—Ph	H	4-F—Ph	H	F	trans
3-CF ₃ -4-F—Ph	H	4-F—Ph	H	F	cis
3-Cl-4-F—Ph	H	4-F—Ph	H	F	trans
3-Cl-4-F—Ph	H	4-F—Ph	H	F	cis
4-F—Ph	H	4-F—Ph	H	F	trans
4-F—Ph	H	4-F—Ph	H	F	cis
Ph	H	4-F—Ph	H	F	trans
Ph	H	4-F—Ph	H	F	cis
3-F-4-Me—Ph	H	4-F—Ph	H	F	cis
3-Me-4-F—Ph	H	4-F—Ph	H	F	cis
3-F-4-Me—Ph	H	4-F—Ph	H	F	trans
3,4-di-F—Ph	H	Ph	H	H	cis
3-F-4-Me—Ph	H	Ph	H	H	cis
3-F-4-CF ₃ —Ph	H	Ph	H	H	cis
3,4-di-F—Ph	H	Ph	H	H	trans
3-F-4-Me—Ph	H	Ph	H	H	trans
3-F-4-CF ₃ —Ph	H	Ph	H	H	trans
3,4-di-F—Ph	H	4-Me—Ph	H	Me	cis
3-F-4-Me—Ph	H	4-Me—Ph	H	Me	cis
3-F-4-CF ₃ —Ph	H	4-Me—Ph	H	Me	cis
3,4-di-F—Ph	H	4-F—Ph	H	F	trans
3-F-4-CF ₃ —Ph	H	4-F—Ph	H	F	trans
3,4-di-F—Ph	H	4-F—Ph	H	F	cis
3-F-4-CF ₃ —Ph	H	4-F—Ph	H	F	cis
3-F-4-Me—Ph	H	4-Me—Ph	H	Me	trans
4-Cl—Ph	H	Ph	H	H	trans
4-CH ₃ OC(O)—Ph	H	Ph	H	H	trans

TABLE 4-continued

Ligand Components					
R ¹	R ³ , R ⁴ , R ⁷ , R ⁹ , and R ¹⁰	R ⁵	R ⁶	R ⁸	Stereo. ¹
3,4-OCH ₂ O—Ph	H	Ph	H	H	trans
4-Cl—Ph	H	4-Me—Ph	H	Me	trans
4-CH ₃ OC(O)—Ph	H	4-Me—Ph	H	Me	trans
3,4-OCH ₂ O—Ph	H	4-Me—Ph	H	Me	trans
4-Cl—Ph	H	4-F—Ph	H	F	trans
4-Et—Ph	H	4-F—Ph	H	F	trans
4-CH ₃ OC(O)—Ph	H	4-F—Ph	H	F	trans
3,4-OCH ₂ O—Ph	H	4-F—Ph	H	F	trans
4-Me—Ph	H	Ph	H	H	80:20 cis:trans
4-Me—Ph	H	4-F—Ph	H	H	75:25 cis:trans
4-Me—Ph	H	2-Cl—Ph	H	H	80:20 cis:trans
4-Me—Ph	H	3-Cl—Ph	H	H	50:50 cis: trans
4-Me—Ph	H	4-Cl—Ph	H	H	80:20 cis:trans
4-Me—Ph	H	3-Me—Ph	H	H	60:40 cis:trans
4-Me—Ph	H	4-Me—Ph	H	H	70:30 cis:trans
4-Me—Ph	H	3-MeO—Ph	H	H	60:40 cis: trans
4-Me—Ph	H	4-MeO—Ph	H	H	80:20 cis:trans
3-F-4-Me—Ph	H, (R ⁹ = Cl)	4-F—Ph	H	H	60:40 cis:trans
3-F-4-CF ₃ —Ph	H	4-Me—Ph	H	Me	trans
2-Me-3-MeO—Ph	H	Ph	H	H	cis
2-F—Ph	H	4-Me—Ph	H	Me	cis
2-Me—Ph	H	4-Me—Ph	H	Me	cis
2-MeO—Ph	H	4-Me—Ph	H	Me	cis
2-Me-3-MeO—Ph	H	4-Me—Ph	H	Me	cis
2-F—Ph	H	4-F—Ph	H	F	cis
2-Me—Ph	H	4-F—Ph	H	F	cis
2-MeO—Ph	H	4-F—Ph	H	F	cis
2-Me-3-MeO—Ph	H	4-F—Ph	H	F	cis
4-Et—Ph	H	Ph	H	H	trans
4-Et—Ph	H	4-Me—Ph	H	Me	trans
4-Cl—Ph	H	Ph	H	H	cis
4-Et—Ph	H	Ph	H	H	cis
4-Cl—Ph	H	4-Me—Ph	H	Me	cis
4-Et—Ph	H	4-Me—Ph	H	Me	cis
4-Cl—Ph	H	4-F—Ph	H	F	cis
4-Et—Ph	H	4-F—Ph	H	F	cis
Ph	H	Ph	H	H	cis
3-F—Ph	H	Ph	H	H	cis
2-CF ₃ —Ph	H	Ph	H	H	cis
3-CF ₃ —Ph	H	Ph	H	H	cis
4-CF ₃ —Ph	H	Ph	H	H	cis
Ph	H	4-Me—Ph	H	Me	cis
3-F—Ph	H	4-Me—Ph	H	Me	cis
2-CF ₃ —Ph	H	4-Me—Ph	H	Me	cis
3-CF ₃ —Ph	H	4-Me—Ph	H	Me	cis
4-CF ₃ —Ph	H	4-Me—Ph	H	Me	cis
3-F—Ph	H	4-F—Ph	H	F	cis
2-CF ₃ —Ph	H	4-F—Ph	H	F	cis
3-CF ₃ —Ph	H	4-F—Ph	H	F	cis
4-CF ₃ —Ph	H	4-F—Ph	H	F	cis
3-MeO—Ph	H	Ph	H	H	cis
4-Me—Ph	H	Ph	H	H	cis
4-MeO—Ph	H	Ph	H	H	cis
4-CH ₃ OC(O)—Ph	H	Ph	H	H	cis
3-Me—Ph	H	4-Me—Ph	H	Me	cis
3-MeO—Ph	H	4-Me—Ph	H	Me	cis
4-Me—Ph	H	4-Me—Ph	H	Me	cis
4-MeO—Ph	H	4-Me—Ph	H	Me	cis
4-CH ₃ OC(O)—Ph	H	4-Me—Ph	H	Me	cis
3-Me—Ph	H	4-F—Ph	H	F	cis
3-MeO—Ph	H	4-F—Ph	H	F	cis
4-Me—Ph	H	4-F—Ph	H	F	cis

TABLE 4-continued

Ligand Components					
R ¹	R ³ , R ⁴ , R ⁷ , R ⁹ , and R ¹⁰	R ⁵	R ⁶	R ⁸	Stereo. ¹
4-MeO—Ph	H	4-F—Ph	H	F	cis
4-CH ₃ OC(O)—Ph	H	4-F—Ph	H	F	cis
4-MeO—Ph	H	Ph	H	H	trans
4-Me—Ph	H	Ph	H	H	trans
Ph	H	Ph	H	H	trans
4-MeO—Ph	H	4-Me—Ph	H	Me	trans
4-Me—Ph	H	4-Me—Ph	H	Me	trans
Ph	H	4-Me—Ph	H	Me	trans
4-MeO—Ph	H	4-F—Ph	H	F	trans
4-Me—Ph	H	4-F—Ph	H	F	trans
6-Cl-3-pyridyl	H	Ph	H	H	cis
5-isoxazolyl	H	Ph	H	H	cis
3-F-4-Cl—Ph	H	Ph	H	H	cis
2-Cl-4-pyridyl	H	Ph	H	H	cis
2-Et-3-MeO—Ph	H	Ph	H	H	cis
3-Cl-6-pyridyl	H	4-Me—Ph	H	Me	cis
5-isoxazolyl	H	4-Me—Ph	H	Me	cis
3-F-4-Cl—Ph	H	4-Me—Ph	H	Me	cis
2-Cl-4-pyridyl	H	4-Me—Ph	H	Me	cis
2-Et-3-MeO—Ph	H	4-Me—Ph	H	Me	cis
3-Cl-6-pyridyl	H	4-F—Ph	H	F	cis
5-isoxazolyl	H	4-F—Ph	H	F	cis
3-F-4-Cl—Ph	H	4-F—Ph	H	F	cis
2-Cl-4-pyridyl	H	4-F—Ph	H	F	cis
2-Et-3-MeO—Ph	H	4-F—Ph	H	F	cis
2-Thienyl	H	Ph	Ac	H	
Styryl	H	Ph	Ac	H	
4-Cl—Ph	H	Ph	4-MeO—Ph—C(O)	H	
furan-2-ylvinyl	H	Ph	H	H	
2-Thienyl	H	Ph	H	H	
4-t-butyl—Ph	H	Ph	Ac	H	
4-F—Ph	H	4-Me—Ph	H	Me	
Benzosuccinimidyl-methyl	H	4-Me—Ph	H	Me	
n-Pr	H	4-F—Ph	benzoyl	H	
n-Octyl	H	Ph	H	H	cis
Me	H	Ph	4-F—Ph—C(O)	H	
2-Cl—PhOCH ₂	H	Ph	B	H	
Benzyl	H	Ph	H	H	
4-MeO—Ph	H	Ph	2-thiophenyl-C(O)	H	
Me	H	Ph	4-Me—Ph—C(O)	H	
3-MeO—Ph	H	Ph	n-hexanoyl	H	
4-t-butyl—Ph	H	Ph	H	H	cis
4-MeO—Ph	H, (R ¹⁰ = Me)	2-Me—Ph	H	H	
3-F—Ph	H	Ph	3-F—Ph(CO)	H	
Ph	H	3-MeO—Ph	H	H	
4-n-pentyl—Ph	H	Ph	H	H	
2-furanyl	H	Ph	H	H	
Ph	H	3-MeO—Ph	Ac	H	
4-Me—Ph	H	Ph	3-MeO—PhC(O)	H	
Me	H	Ph	3-MeO—PhC(O)	H	
4-Me—Ph	H	Ph	4-F—Ph—C(O)	H	
4-Cl—Ph	H	4-Me—Ph	H	Me	
CO ₂ Et	H	Ph	EtOC(O)C(O)	H	
3,4-di-MeO-styryl	H	Ph	H	H	cis
Styryl	H	Ph	styryl-C(O)	H	
3-Br—Ph	H	Ph	H	H	
Ph	H	4-Me—Ph	Ac	H	
4-MeO-styryl	H	Ph	Ac	H	
Benxosuccinimidyl-methyl	H	Ph	H	H	
4-MeO—Ph	H	4-Me—Ph	H	Me	trans
4-MeO—Ph	H	Ph	4-MeO—Ph—C(O)	H	
3-NO ₂ —Ph	H	4-Me—Ph	H	Me	
cyclopropyl	H	Ph	cyclopropyl-C(O)	H	
Me	H	3-MeO—Ph	benzoyl	H	
4--n-propyl	H, (R ¹⁰ = Me)	2-Me—Ph	H	H	
3-NO ₂ —Ph	H	Ph	H	H	cis
4-F—PhOCH ₂	H	Ph	H	H	
n-Pr	H	Ph	3-MeO—PhC(O)	H	

TABLE 4-continued

Ligand Components					
R ¹	R ³ , R ⁴ , R ⁷ , R ⁹ , and R ¹⁰	R ⁵	R ⁶	R ⁸	Stereo. ¹
4-Cl—Ph	H	Ph	4-Me—Ph—C(O)	H	
4-Et—Ph	H, (R ¹⁰ = Me)	2-Me—Ph	styryl-C(O)	H	
Styryl	H	Ph	H	H	cis
3-Me—Ph	H	Ph	3-Me—Ph—C(O)	H	
3,4-di-Cl—Ph	H	Ph	H	H	cis
3-OH—Ph	H	Ph	3-Br—Ph(CO)	H	
succinimidyImethyl	H	Ph	H	H	
4-I—Ph	H	Ph	H	H	cis
1-naphthylmethyl	H	Ph	H	H	
cyclohexylethyl	H	Ph	H	H	
CO ₂ Et	H	Ph	H	H	
4-F—Ph	H	Ph	4-F—Ph—C(O)	H	
4-n-propyl—Ph	H	Ph	H	H	
3-F—Ph	H	4-Me—Ph	H	Me	trans
4-CH ₃ S(O ₂)NH—Ph	H	Ph	H	H	
NHPh	H	Ph	H	H	
4-MeO—styryl	H	Ph	H	H	cis
i-Pr	H	4-NO ₂ —Ph	benzoyl	H	
3-Cl-benzofuran-2-yl	H	Ph	3-Cl- benzothiophen-2-yl	H	
4-Cl—PhOCH ₂	H	Ph	H	H	
4-MeO—Ph	H	Ph	4-MeO—styryl	H	
CF ₃	H	Ph	CF ₃ C(O)	H	
Et	H	Ph	4-NO ₂ —Ph—C(O)	H	
Ph	H, (R ¹⁰ = Me)	2-Me—Ph	H	H	cis
Me	H	Ph	2-F—Ph—C(O)	H	
n-pentyl	H	Ph	2-F—Ph—C(O)	H	
4-Me—Ph	H, (R ¹⁰ = Me)	2-Me—Ph	H	H	cis
3-F-4-Me—Ph	H	4-Me—Ph	3-F-4-Me—Ph(CO)	Me	trans
3-F-4-CF ₃ —Ph	H	4-Me—Ph	3-F-4-CF ₃ —Ph—C(O)	Me	trans
4-Cl—Ph	H	Ph	4-Cl—Ph—C(O)	H	trans
4-Et—Ph	H	Ph	4-Et—Ph—C(O)	H	trans
4-Cl—Ph	H	4-Me—Ph	4-Cl—Ph—C(O)	Me	trans
4-Et—Ph	H	4-Me—Ph	4-Et—Ph—C(O)	Me	trans
3,4-OCH ₂ O—Ph	H	4-Me—Ph	3,4-OCH ₂ O—Ph—C(O)	Me	trans
3-F-4-Me—Ph	H	4-F—Ph	Ac	F	trans
3-F-4-Me—Ph	H	4-F—Ph	3-F-4-Me—Ph(CO)	F	trans
3-F-4-Me—Ph	H	4-F—Ph	3-F-4-Me—Ph(CO)	F	cis
3-Me—Ph	H	Ph	H	H	trans
3-F—Ph	H	Ph	H	H	trans
3-MeO—Ph	H	Ph	H	H	trans
3-CF ₃ —Ph	H	Ph	H	H	trans
3-Me—Ph	H	4-Me—Ph	H	Me	trans
3-F—Ph	H	4-Me—Ph	H	Me	trans
3-MeO—Ph	H	4-Me—Ph	H	Me	trans
3-CF ₃ —Ph	H	4-Me—Ph	H	Me	trans
3-Me—Ph	H	4-F—Ph	H	F	trans
3-F—Ph	H	4-F—Ph	H	F	trans
3-MeO—Ph	H	4-F—Ph	H	F	trans
3-CF ₃ —Ph	H	4-F—Ph	H	F	trans
NHEt	H	Ph	H	H	cis
NHPh	H	Ph	H	H	cis
4-Cl—Ph—NH	H	Ph	H	H	cis
3-Cl—Ph—NH	H	Ph	H	H	cis
4-Me—Ph—NH	H	Ph	H	H	cis
3-Me—Ph—NH	H	Ph	H	H	cis
NHPh	H	4-Me—Ph	H	Me	cis
4-Cl—Ph—NH	H	4-Me—Ph	H	Me	cis
3-Cl—Ph—NH	H	4-Me—Ph	H	Me	cis
4-Me—Ph—NH	H	4-Me—Ph	H	Me	cis
3-Me—Ph—NH	H	4-Me—Ph	H	Me	cis
NHPh	H	4-F—Ph	H	F	cis
4-Cl—Ph—NH	H	4-F—Ph	H	F	cis
3-Cl—Ph—NH	H	4-F—Ph	H	F	cis
4-Me—Ph—NH	H	4-F—Ph	H	F	cis
3-Me—Ph—NH	H	4-F—Ph	H	F	cis

¹Relative stereochemistry at 2- and 4-positions

halocycloalkylamino, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₆)althylthio(C₁-C₆)alkyl, (C₁-C₆)alkylsulfinyl(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl(C₁-C₆)alkyl, (C₁-C₆)alkylamino(C₃-C₆)alkyl, di(C₁-C₆)alkylamino(C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl(C₁-C₆)alkyl, cyano(C₁-C₆)alkyl, optionally substituted phenyl, optionally substituted phenyl(C₁-C₆)alkyl, optionally substituted heterocycle, optionally substituted phenoxy, optionally substituted heterocycloxy, optionally substituted phenylthio, optionally substituted heterocyclithio, optionally substituted naphthyl, optionally substituted phenylamino, optionally substituted heterocyclylamino, optionally substituted N-phenyl-N-(C₁-C₆)alkylamino, and optionally substituted N-heterocyclyl-N-(C₁-C₆)alkylamino.

[0508] In another embodiment, the activating ligand is a compound having Formula V, wherein:

[0509] Q¹ is O;

[0510] R¹ is substituted phenyl wherein the substituents are independently selected from the group consisting of (C₁-C₂)alkyl and (C₁-C₂)alkoxy; or two adjacent positions are joined together with the atoms to which they are attached to form an unsubstituted or substituted, unsaturated, partially unsaturated, or saturated 5-, 6- or 7-membered carbocyclic or heterocyclic ring, wherein the heterocyclic ring contains from one to two oxygen atoms and one to four substituents are independently selected from the group consisting of: cyano, (C₁-C₂)alkyl, (C₁-C₂)alkylamino, di(C₁-C₂)alkylamino, (C₁-

C₂)alkoxycarbonyl, (C₁-C₂)alkylaminocarbonyl, di(C₁-C₂)alkylaminocarbonyl, oxo, and methoxyimino;

[0511] R² and R³ are independently selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₃)alkoxy(C₁-C₃)alkyl, (C₁-C₃)althylthio(C₁-C₃)alkyl, (C₁-C₃)alkylsulfinyl(C₁-C₃)alkyl, (C₁-C₃)alkylsulfonyl(C₁-C₃)alkyl, (C₁-C₃)alkylamino(C₁-C₃)alkyl, di(C₁-C₃)alkylamino(C₁-C₃)alkyl, (C₁-C₆)alkylcarbonyl, (C₁-C₃)alkylcarbonyl(C₁-C₃)alkyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₃)alkylaminocarbonyl(C₁-C₃)alkyl, di(C₁-C₃)alkylaminocarbonyl(C₁-C₃)alkyl, (C₁-C₃)alkylcarbonylamino(C₁-C₃)alkyl, (C₁-C₆)alkoxycarbonyl, and C₁-C₃)alkoxycarbonyl(C₁-C₃)alkyl; or

[0512] R² and R³ may be joined together with the carbon to which they are attached to form an unsubstituted or substituted, partially unsaturated or saturated 5-, 6- or 7-membered carbocyclic or heterocyclic ring, wherein the heterocyclic ring contains one heteroatom selected from O or S; and one to four substituents are independently selected from the group consisting of (C₁-C₃)alkyl, (C₁-C₃)alkylamino, di(C₁-C₃)alkylamino, (C₁-C₄)alkoxycarbonyl, (C₁-C₃)alkylaminocarbonyl, and di(C₁-C₃)alkylaminocarbonyl, and

[0513] R⁴ is selected from optionally substituted phenyl or pyridyl wherein the substituents are independently selected from the group consisting of (C₁-C₃)alkyl and (C₁-C₃)alkoxy;

[0514] In another embodiment, the activating ligand is a compound having Formula V, wherein Q is oxygen, and R¹, R², R³, and R⁴ are defined according to Table 5

TABLE 5

Ligand Components			
R ¹	R ²	R ³	R ⁴
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3-Me—Ph
4-Et—Ph	—(CH ₂) ₄ —		Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₃ —		3-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3-Me—Ph
2-Me-3-MeO—Ph	Bn	Me	3-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₂ —		3-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	Bn	Me	3,5-diMe—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₂ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₃ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		4-Me—Ph
2-Me-3-MeO—Ph	Bn	Me	4-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3-Me-4-F—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3-Me-4-F—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₂ —		3-Me-4-F—Ph
2-Me-3-MeO—Ph	i-Pr	Me	3,5-diMe—Ph
2-Et-3-MeO—Ph	—(CH ₂) ₄ —		Ph
2-Et-3,6-OCH ₂ CH ₂ O—Ph	—(CH ₂) ₄ —		Ph
2-Me-3,4-OCH ₂ O—Ph	—(CH ₂) ₄ —		Ph
2-Me-3-MeO—Ph	—(CH ₂) ₂ —		4-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ OCH ₂ CH ₂ —		3-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ SCH ₂ CH ₂ —		3-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ OCH ₂ CH ₂ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ SCH ₂ CH ₂ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ C(OCH ₂ CH ₂ O)CH ₂ CH ₂ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	i-Pr	Me	2-MeO—Ph
2-Me-3-MeO—Ph	i-Pr	Me	3-Me—Ph
2-Me-3-MeO—Ph	i-Pr	Me	3-MeO—Ph
2-Me-3-MeO—Ph	i-Pr	Me	4-Me—Ph
2-Me-3-MeO—Ph	i-Pr	Me	Ph

TABLE 5-continued

Ligand Components			
R ¹	R ²	R ³	R ⁴
2-Me-3-MeO—Ph	—CH ₂ CH ₂ C(OCH ₂ CH ₂ O)CH ₂ CH ₂ —		3-Me—Ph
2-Me-3-MeO—Ph	Et	Et	2-Me—Ph
2-Me-3-MeO—Ph	Et	Et	2-MeO—Ph
2-Me-3-MeO—Ph	Et	Et	4-F—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		2-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		2-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		4-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		4-F—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3,4-OCH ₂ O—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		2-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		2-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		4-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3,4-OCH ₂ O—Ph
2-Me-3-MeO—Ph	Et	Et	3-Me—Ph
2-Me-3-MeO—Ph	Et	Et	3-MeO—Ph
2-Me-3-MeO—Ph	Et	Et	3-Me-4-F—Ph
2-Me-3-MeO—Ph	Et	Et	3,5-diMe—Ph
2-Me-3-MeO—Ph	i-Bu	Me	3-Me—Ph
2-Me-3-MeO—Ph	i-Bu	Me	3-MeO—Ph
2-Me-3-MeO—Ph	i-Bu	Me	3-Me-4-F—Ph
2-Me-3-MeO—Ph	i-Bu	Me	3,5-diMe—Ph
2-Me-3-MeO—Ph	i-Pr	Me	3-Me-4-F—Ph
2-Me-3-MeO—Ph	Ph	i-Pr	3-Me—Ph
2-Me-3-MeO—Ph	Et	Et	4-MeO—Ph
2-Me-3-MeO—Ph	Et	Et	3,4-OCH ₂ O—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		4-F—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ C(=O)CH ₂ CH ₂ —		3-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ S(=O) ₂ CH ₂ CH ₂ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	i-Pr	Me	2-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		2,6-diMeO-3-pyridyl
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3,5-diMeO-4-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3,5-diMeO-4-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		3-MeO-4,5-diF—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		3-MeO-4,5-diF—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		Ph
2-Me-3-MeO—Ph	—(CH ₂) ₆ —		2-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₆ —		3,5-diMe—Ph
2-Me-3-MeO—Ph	4-F—Ph	Me	2-MeO—Ph
2-Me-3-MeO—Ph	4-F—Ph	Me	3,5-diMe—Ph
2-Me-3-MeO—Ph	Me	Me	2-MeO—Ph
2-Me-3-MeO—Ph	Me	Me	3,5-diMe—Ph
2-Me-3-MeO—Ph	Me	Me	Ph
2-Me-3-MeO—Ph	Et	Et	4-Me—Ph
2-Me-3-MeO—Ph	Et	Et	Ph
2-Me-3-MeO—Ph	—(CH ₂) ₄ —		4-Me—Ph
2-Et-3,4-OCH ₂ CH ₂ O—Ph	—(CH ₂) ₅ —		3,5-di-Me—Ph
2-Me-3,4-OCH ₂ O—Ph	—(CH ₂) ₅ —		3,5-di-Me—Ph
3,4-OCH ₂ CH ₂ O—Ph	—(CH ₂) ₅ —		3,5-di-Me—Ph
3,4-CH ₂ OCH ₂ O—Ph	—(CH ₂) ₅ —		3,5-di-Me—Ph
2-Et-3,4-OCH ₂ CH ₂ O—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
2-Me-3,4-OCH ₂ O—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
3,4-OCH ₂ CH ₂ O—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
3,4-CH ₂ OCH ₂ O—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
3,4-OCH ₂ O—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
2-Me—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
Ph	t-Bu	H	4-Cl—Ph
4-Cl—Ph	—(CH ₂) ₄ —		Ph
Me	Ph	H	4-Me—Ph
Me	4-Me—Ph	H	Ph
Me	Ph	H	Ph
4-Cl—Ph	Me	Me	Ph
4-Me—Ph	t-Bu	H	Ph
2,3-di-Me—Ph	t-Bu	H	Ph
4-NO ₂ —Ph	t-Bu	H	Ph
2-Me-3-MeO—Ph		—(CH ₂) ₂ —	3-MeO—Ph
2-Me-3-MeO—Ph	Benzyl	Me	3-MeO—Ph
2-Me-3-MeO—Ph		—(CH ₂) ₂ —	2-Me—Ph
3-Me-benzofuran-2-yl		—(CH ₂) ₄ —	Ph
Ph	Me	Me	Ph
2-Me—Ph	Me	Me	Ph
3,4-OCH ₂ O—Ph	Me	Me	Ph
3-MeO—Ph	Me	Me	Ph
4-Et—Ph	Me	Me	Ph

TABLE 5-continued


Ligand Components			
R ¹	R ²	R ³	R ⁴
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N(C(O)OtBu)CH ₂ CH ₂ —		3-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N(C(O)OtBu)CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	i-Pr	Me	3,4-OCH ₂ O—Ph
2-Me-3-MeO—Ph	i-Pr	Me	Me
2-Me-3-MeO—Ph	t-Bu	H	3-Me—Ph
2-Me-3-MeO—Ph	t-Bu	H	3-MeO—Ph
2-Me-3-MeO—Ph	t-Bu	H	3,5-di-Me—Ph
2-MeO—Ph	Me	Me	3-Me—Ph
2-MeO—Ph	Me	Me	3-MeO—Ph
2-Me-3-MeO—Ph	i-Bu	Me	4-MeO—Ph
2-MeO—Ph	Me	Me	3,5-di-Me—Ph
2-Me-3-MeO—Ph	(CH ₂) ₅		n-Bu
Ph	Me	Me	Et
3-MeO—Ph	Me	Me	Et
3,4-OCH ₂ O—Ph	Me	Me	Et
2-Me—Ph	Me	Me	Et
4-Et—Ph	Me	Me	Et
Ph	Me	Me	3,5-di-Me—Ph
2-Me—Ph	Me	Me	3,5-di-Me—Ph
3-MeO—Ph	Me	Me	3,5-di-Me—Ph
4-Et—Ph	Me	Me	3,5-di-Me—Ph
3,4-OCH ₂ O—Ph	Me	Me	3,5-di-Me—Ph
Ph	—(CH ₂) ₄ —		Et
2-Me—Ph	—(CH ₂) ₄ —		Et
3-MeO—Ph	—(CH ₂) ₄ —		Et
4-Et—Ph	—(CH ₂) ₄ —		Et
3,4-OCH ₂ O—Ph	—(CH ₂) ₄ —		Et
Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
3-MeO—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
4-Et—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
Ph	—(CH ₂) ₄ —		Ph
2-Me—Ph	—(CH ₂) ₄ —		Ph
3-MeO—Ph	—(CH ₂) ₄ —		Ph
3,4-OCH ₂ O—Ph	—(CH ₂) ₄ —		Ph
2-Et-3-MeO—Ph	—(CH ₂) ₅ —		3,5-di-Me—Ph
2-Et-3-MeO—Ph	—(CH ₂) ₄ —		3,5-di-Me—Ph
CF ₃	—(CH ₂) ₄ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ N[(C=O)Ot-Bu]CH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ NHCH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ NHCH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[(C=O)CH ₃]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[(C=O)(C=O)OEt]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[S(O) ₂ CH ₃]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[CH ₂ (C=O)OEt]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ N[(C=O)CH ₃]CH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ N[(C=O)(C=O)OEt]CH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ N[S(O) ₂ CH ₃]CH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ N[CH ₂ (C=O)OCH ₃]CH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[(C=O)NH ₂]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[(C=O)OiPr]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ N[CH ₂ CN]CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ N[(C=O)NH ₂]CH ₂ CH ₂ CH ₂ —		3,5-di-Me—Ph
2-Me-3-MeO—Ph	—CH ₂ CH ₂ CH ₂ N(CH ₃)CH ₂ —		3,5-di-Me—Ph
2-NH ₂ —Ph	Et	H	Ph
4-Et—Ph	—(CH ₂) ₅ —		3,5-di-Cl—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		2-MeO-5-F—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		2-MeO-5-Me—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		2,5-di-MeO—Ph
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		4-Me-2-pyridyl
2-Me-3-MeO—Ph	—(CH ₂) ₅ —		6-Me-2-pyridyl
4-Et—Ph	—(CH ₂) ₅ —		2-MeO-5-F—Ph
4-Et—Ph	—(CH ₂) ₅ —		2-MeO-5-Me—Ph
4-Et—Ph	—(CH ₂) ₅ —		2,5-di-MeO—Ph
4-Et—Ph	—(CH ₂) ₅ —		4-Me-2-pyridyl
4-Et—Ph	—(CH ₂) ₅ —		6-Me-2-pyridyl
4-Et—Ph	—(CH ₂) ₅ —		2-MeO—Ph
4-Et—Ph	—(CH ₂) ₅ —		3,5-di-Me—Ph
4-Et—Ph	—(CH ₂) ₅ —		3-Me—Ph
2-Me-3-MeO—Ph	i-Pr	Et	2-MeO—Ph
2-Me-3-MeO—Ph	i-Pr	Et	3,5-di-Me—Ph

[0515] In another embodiment, the activating ligand is a compound having Formula VI, wherein a is 2, and R^2 , R^3 , R^4 , and R^5 are defined according to Table 6.

TABLE 6

Ligand Components		
R^2/R^3	R^4	R^5
$-(CH_2)_5-$	3,5-di-Me-Ph	4H-benzo[1,3]dioxine-6-yl
$-(CH_2)_4-$	3,5-di-Me-Ph	4-Me-Ph
$-(CH_2)_5-$	3,5-di-Cl-Ph	4-Me-Ph
$-(CH_2)_5-$	3,5-di-Cl-Ph	3-MeO-Ph

[0516] In another embodiment, the activating ligand is a compound having Formula VIII:



VIII

wherein:

[0517] X and X' are independently O or S;

[0518] R^1 is selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_1-C_6) haloalkyl, (C_1-C_6) cyanoalkyl, (C_1-C_6) alkoxycarbonyl (C_1-C_6) alkyl, (C_1-C_6) alkoxy, benzyloxy, optionally substituted phenyl, optionally substituted naphthyl wherein the substituents are independently 1 to 3 halo, nitro, (C_1-C_6) alkoxy, (C_1-C_6) alkyl, or amino, optionally substituted benzothiophene-2-yl, benzothiophene-3-yl, benzofuran-2-yl, or benzofuran-3-yl wherein the substituents are independently 1 to 3 halo, nitro, hydroxy, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, carboxy, or (C_1-C_6) alkoxycarbonyl $(-CO_2R^a)$, optionally substituted 2-, 3-, or 4-pyridyl wherein the substituents are independently 1 to 3 halo, nitro, cyano, nitro, hydroxy, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, or (C_1-C_6) haloalkoxy, optionally substituted 5-membered heterocycle selected from furyl, thiophenyl, triazolyl, pyrrolyl, isopyrrolyl, pyrazolyl, isoimidazolyl, thiazolyl, isothiazolyl, oxazolyl, or isooxazolyl wherein the substituents are independently 1 to 3 halo, nitro, hydroxy, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, carboxy, (C_1-C_6) alkoxycarbonyl $(-CO_2R^a)$, or unsubstituted or substituted phenyl wherein the substituents are independently 1 to 3 halo, nitro, (C_1-C_6) alkyl, (C_1-C_6) haloalkyl, (C_1-C_6) alkoxy, (C_1-C_6) haloalkoxy, carboxy, (C_1-C_4) alkoxycarbonyl $(-CO_2R^a)$, or amino $(-NR^aR^b)$, aromatic substituted or unsubstituted phenyl (C_1-C_6) alkyl, phenyl (C_1-C_6) alkoxy (C_1-C_6) alkyl, or phenoxy (C_1-C_6) alkyl wherein the aromatic substituents are independently 1 to 3 halo, nitro, (C_1-C_6) alkoxy, (C_1-C_6) alkyl, or amino, and aromatic substituted or unsubstituted phenylamino, phenyl (C_1-C_6) alkylamino, or phenylcarbonylamino wherein the aromatic substituents are independently 1 to 3 halo, nitro, (C_1-C_6) alkoxy, (C_1-C_6) alkyl, or amino;

[0519] wherein R^a , R^b , and R^c are independently H, (C_1-C_6) alkyl, or phenyl;

[0520] R^2 and R^3 are independently H, (C_1-C_6) alkyl, (C_1-C_6) haloalkyl, (C_3-C_6) cyanoalkyl, (C_1-C_6) hydroxyalkyl, (C_1-C_6) alkoxy (C_1-C_6) alkyl, phenyl, or together as an alkane linkage $(-CH_2)_x-$, an alkoxyalkyl linkage $(-CH_2)_yO(CH_2)_z-$, an alkylaminoalkyl linkage $(-CH_2)_yNR^a(CH_2)_z-$, or an alkylbenzoalkyl linkage $(-CH_2)_y-1-benzo-2-(CH_2)_z-$ form a ring with the carbon atom to which they are attached, wherein $x=3$ to 7, $y=1$ to 3, $z=1$ to 3, and R^a is H, (C_1-C_6) alkyl, or phenyl; and

[0521] R^4 is optionally substituted phenyl, wherein the substituents are independently 1 to H; halo; nitro; cyano; hydroxy; amino $(-NR^aR^b)$; (C_1-C_6) alkyl; (C_1-C_6) haloalkyl; (C_1-C_6) cyanoalkyl; (C_1-C_6) hydroxyalkyl; (C_1-C_6) alkoxy; phenoxy; (C_1-C_6) haloalkoxy; (C_1-C_6) alkoxy (C_1-C_6) alkyl; (C_1-C_6) alkoxy (C_1-C_6) alkoxy; (C_1-C_6) alkanoyloxy (C_1-C_6) alkyl; (C_2-C_6) alkenyl optionally substituted with halo, cyano, (C_1-C_4) alkyl, or (C_1-C_4) alkoxy; (C_2-C_6) alkynyl optionally substituted with halo or (C_1-C_4) alkyl; formyl; carboxy; (C_1-C_6) alkylcarbonyl; (C_1-C_6) haloalkylcarbonyl; benzoyl; (C_1-C_6) alkoxycarbonyl; (C_1-C_6) haloalkoxycarbonyl; (C_1-C_6) alkanoyloxy $(-OCOR^a)$; carboxamido $(-CONR^aR^b)$; amido $(-NR^aCOR^b)$; alkoxycarbonylamino $(-NR^aCO_2R^b)$; alkylaminocarbonylamino $(-NR^aCONR^bR^c)$ mercapto; (C_1-C_6) alkylthio; (C_1-C_6) alkylsulfonyl; (C_1-C_6) alkylsulfoxido $(-S(O)R^a)$; sulfamido $(-SO_2NR^aR^b)$; or optionally substituted phenyl wherein the substituents are independently 1 to 3 halo, nitro, (C_1-C_6) alkoxy, (C_1-C_6) alkyl, or amino; or when two adjacent positions on the phenyl ring are substituted with alkoxy groups, these groups, together with the carbon atoms to which they are attached, may be joined to form a 5- or 6-membered dioxolano $(-OCH_2O-)$ or dioxano $(-OCH_2CH_2O-)$ heterocyclic ring; wherein R^a , R^b , and R^c are independently H, (C_1-C_6) alkyl, or phenyl.

[0522] In another embodiment, the activating ligand is a compound having Formula VIII, wherein:

[0523] X and X' are O;

[0524] R^1 is phenyl, 4-chlorophenyl-, 4-ethylphenyl-, 2-ethyl-3,4-ethylenedioxyphenyl-, 3-fluorophenyl-, 2-fluoro-4-ethylphenyl-, 2-methyl-3-methoxyphenyl-, 2-ethyl-3-methoxyphenyl-, 3-methylphenyl-2-methoxyphenyl-, 2-nitrophenyl-, 3-nitrophenyl-, 2-furanyl-, benzyl-, benzothiophene-2-yl-, phenylamino-, benzyloxymethyl-, phenoxyethyl-, 3-toluylamino-, benzylamino-, benzoylamino-, ethoxycarbonyl-, or 3-chloro-2,2,3,3-tetrafluoroethyl;

[0525] R^2 and R^3 are independently methyl, ethyl, or together as a tetramethylene $(-CH_2)_4-$, 4-pyrano $(-CH_2CH_2OCH_2CH_2-)$, or methylenebenzoethylene $(-CH_2-1-benzo-2-CH_2CH_2-)$ linkage form a ring with the carbon atom to which they are attached; and

[0526] R^4 is phenyl, 4-biphenyl, 4-chlorophenyl, 2,4-dimethoxyphenyl, 3,5-dimethylphenyl, 2-methoxyphenyl, 3,4-methylenedioxyphenyl, 3-trifluoromethylphenyl, or 4-trifluoromethoxyphenyl;


[0527] In another embodiment, the activating ligand is a compound having Formula VIII selected from the group consisting of:

- [0528] 1-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0529] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-fluoro-benzamide;
- [0530] Furan-2-carboxylic acid [3-(3,5-dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-amide;
- [0531] 3-Chloro-N-[3-(3,5-dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-2,2,3,3-tetrafluoro-propionamide;
- [0532] N-[3-(3,5-Dimethyl-phenyl)-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-4-ethyl-benzamide;
- [0533] 2-Benzyloxy-N-[5,5-dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-acetamide;
- [0534] N-[3-(3,5-Dimethyl-phenyl)-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0535] 2-Benzyloxy-N-[3-(3,5-dimethyl-phenyl)-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-acetamide;
- [0536] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-benzamide;
- [0537] Furan-2-carboxylic acid [3-(2-methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0538] 2-Phenoxy-N-(3-phenyl-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-acetamide;
- [0539] N-(3-Phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-succinamic acid ethyl ester;
- [0540] N-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0541] 2-Ethyl-3-methoxy-N-[3-(2-methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0542] 1-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-3-phenyl-urea;
- [0543] 2-Benzyloxy-N-[3-(2-methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-acetamide;
- [0544] N-[3-(3,5-Dimethyl-phenyl)-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-benzamide;
- [0545] N-(3-Biphenyl-4-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-2-ethyl-3-methoxy-benzamide;
- [0546] N-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-2-phenyl-acetamide;
- [0547] N-[5,5-Dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0548] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-2-ethyl-3-methoxy-benzamide;
- [0549] 4-Chloro-N-[3-(3,5-dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0550] 1-[3-(2-Methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0551] 4-Ethyl-N-[3-(2-methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0552] 1-Phenyl-3-(3-phenyl-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-urea;
- [0553] N-[5,5-Dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-2-phenoxy-acetamide;
- [0554] 2-Phenyl-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-acetamide;
- [0555] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-succinamic acid ethyl ester;
- [0556] N-[5,5-Dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0557] 2-Benzyloxy-N-(3-phenyl-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-acetamide;
- [0558] N-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-4-ethyl-benzamide;
- [0559] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,5]-7,8-benzo-dec-2-en-4-yl]-3-methoxy-2-methyl-benzamide;
- [0560] N-[3-(2,4-Dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-succinamic acid ethyl ester;
- [0561] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0562] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-4-ethyl-benzamide;
- [0563] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-2-phenoxy-acetamide;
- [0564] N-(5,5-Dimethyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-3-methoxy-2-methyl-benzamide;
- [0565] N-(3-Phenyl-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-benzamide;
- [0566] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-methoxy-2-methyl-benzamide;
- [0567] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-2-phenyl-acetamide;
- [0568] Benzo[b]thiophene-2-carboxylic acid [3-(2-methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0569] N-[3-(3,5-Dimethyl-phenyl)-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-2-phenoxy-acetamide;
- [0570] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0571] 2-Benzyloxy-N-[3-(3,5-dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-acetamide;
- [0572] 1-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0573] 2-Benzyloxy-N-[3-(3,5-dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-acetamide;
- [0574] 1-[3-(3,5-Dimethyl-phenyl)-1,8-dioxa-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-3-phenyl-urea;
- [0575] N-[5,5-Dimethyl-3-(4-trifluoromethoxy-phenyl)-3,1,2,4]oxadiazol-4-yl]-4-ethyl-benzamide;
- [0576] 1-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-m-tolyl-urea;
- [0577] N-[3-(2-Methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-phenoxy-acetamide;
- [0578] N-[3-(2,4-Dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0579] 3-Chloro-N-[5,5-dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-2,2,3,3-tetrafluoro-propionamide;
- [0580] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-4-ethyl-benzamide;
- [0581] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-4-ethyl-benzamide;
- [0582] 3-Chloro-2,2,3,3-tetrafluoro-N-[3-(2-methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-propionamide;
- [0583] 3-Chloro-2,2,3,3-tetrafluoro-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-propionamide;
- [0584] 2-Benzyloxy-N-[5,5-dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-acetamide;
- [0585] 1-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0586] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0587] Furan-2-carboxylic acid [5,5-dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-amide;
- [0588] Furan-2-carboxylic acid (3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-amide;

- [0589] 1-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0590] 3-Chloro-N-[3-(4-chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2,2,3,3-tetrafluoro-propionamide;
- [0591] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-methoxy-benzamide;
- [0592] 2-Ethyl-N-(5-ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-3-methoxy-benzamide;
- [0593] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-methyl-benzamide;
- [0594] N-[3-(2,4-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-phenyl-acetamide;
- [0595] N-[3-(2,4-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-phenoxy-acetamide;
- [0596] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0597] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-2-phenyl-acetamide;
- [0598] Furan-2-carboxylic acid [3-(4-chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0599] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-succinamic acid ethyl ester;
- [0600] N-[3-(3,5-Dimethyl-phenyl)-1,8-dioxo-2,4-diazaspiro[4,5]dec-2-en-4-yl]-2-phenyl-acetamide;
- [0601] N-[3-(3,5-Dimethyl-phenyl)-1,8-dioxo-2,4-diazaspiro[4,5]dec-2-en-4-yl]-3-methoxy-2-methyl-benzamide;
- [0602] Benzo[b]thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0603] 1-Benzyl-3-[3-(3,5-dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-urea;
- [0604] N-(3-Phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-benzamide;
- [0605] 3-Chloro-N-[3-(3,5-dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-2,2,3,3-tetrafluoro-propionamide;
- [0606] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-nitro-benzamide;
- [0607] 2-Ethyl-3-methoxy-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-benzamide;
- [0608] N-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0609] Furan-2-carboxylic acid [5,5-dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-amide;
- [0610] 1-(5-Ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-3-phenyl-urea;
- [0611] N-[3-(2,4-Dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0612] N-[3-(3,5-Dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-nitro-benzamide;
- [0613] N-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-ethyl-3-methoxy-benzamide;
- [0614] Furan-2-carboxylic acid (5-ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-amide;
- [0615] Furan-2-carboxylic acid [3-(2,4-dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0616] N-(5-Ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-2-phenoxy-acetamide;
- [0617] Furan-2-carboxylic acid [3-(3,5-dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0618] Benzo[b]thiophene-2-carboxylic acid [5,5-dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-amide;
- [0619] Benzo[b]thiophene-2-carboxylic acid [5,5-dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-amide;
- [0620] 2-Benzoyl-N-[3-(2,4-dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-acetamide;
- [0621] 1-Benzoyl-3-[3-(3,5-dimethyl-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-urea;
- [0622] 1-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-3-phenyl-urea;
- [0623] 1-[3-(2,4-Dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0624] N-(5,5-Dimethyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-4-ethyl-benzamide;
- [0625] 2-Benzoyloxy-N-[3-(4-chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-acetamide;
- [0626] N-(5-Ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-benzamide;
- [0627] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-2-phenyl-acetamide;
- [0628] N-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-phenyl-acetamide;
- [0629] 1-[5,5-Dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-3-phenyl-urea;
- [0630] 4-Ethyl-N-(3-phenyl-1,8-dioxo-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-benzamide;
- [0631] 4-Ethyl-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-benzamide;
- [0632] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-succinamic acid ethyl ester;
- [0633] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-2-phenoxy-acetamide;
- [0634] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-4-ethyl-benzamide;
- [0635] Benzo[b]thiophene-2-carboxylic acid [3-(2,4-dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0636] 2-Phenyl-N-(3-phenyl-1,8-dioxo-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-acetamide;
- [0637] 1-Phenyl-3-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-urea;
- [0638] Benzo[b]thiophene-2-carboxylic acid (5-ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-amide;
- [0639] N-[3-(2,4-Dimethoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-4-ethyl-benzamide;
- [0640] 4-Ethyl-N-(5-ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-benzamide;
- [0641] Furan-2-carboxylic acid [3-(3,5-dimethyl-phenyl)-1,8-dioxo-2,4-diaza-spiro[4,5]dec-2-en-4-yl]-amide;
- [0642] Benzo[b]thiophene-2-carboxylic acid (3-benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-amide;
- [0643] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-2-phenoxy-acetamide;
- [0644] N-(3-Biphenyl-4-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-4-ethyl-benzamide;
- [0645] N-[3-(2-Methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-succinamic acid ethyl ester;
- [0646] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-2-benzoyloxy-acetamide;
- [0647] N-(5-Ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-2-phenyl-acetamide;

- [0648] N-[3-(2-Methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0649] N-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-4-ethyl-benzamide;
- [0650] Furan-2-carboxylic acid (3-benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-amide;
- [0651] Benzo[b]thiophene-2-carboxylic acid (3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-amide;
- [0652] N-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-benzamide;
- [0653] Benzo[b]thiophene-2-carboxylic acid [3-(3,5-dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-amide;
- [0654] N-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-succinamic acid ethyl ester;
- [0655] 2-Benzyloxy-N-(5-ethyl-5-methyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-acetamide;
- [0656] 2-Benzyloxy-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-acetamide;
- [0657] N-(3-Benzo[1,3]dioxol-5-yl-5,5-dimethyl-[1,2,4]oxadiazol-4-yl)-benzamide;
- [0658] N-[3-(2-Methoxy-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-phenyl-acetamide;
- [0659] 2-Phenoxy-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-acetamide;
- [0660] 2-Ethyl-3-methoxy-N-(3-phenyl)-1,8-dioxo-2,4-diaza-spiro[4,5]dec-2-en-4-yl)-benzamide;
- [0661] N-[5,5-Dimethyl-3-(4-trifluoromethoxy-phenyl)-[1,2,4]oxadiazol-4-yl]-2-phenyl-acetamide;
- [0662] Benzo[b]thiophene-2-carboxylic acid [3-(3,5-dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-amide;
- [0663] N-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-2-phenoxy-acetamide;
- [0664] N-[5,5-Dimethyl-3-(3-trifluoromethyl-phenyl)-[1,2,4]oxadiazol-4-yl]-2-phenoxy-acetamide;
- [0665] N-[3-(4-Chloro-phenyl)-5,5-dimethyl-[1,2,4]oxadiazol-4-yl]-succinamic acid ethyl ester;
- [0666] N-[3-(3,5-Dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-4-ethyl-2-fluoro-benzamide;
- [0667] 4-Ethyl-2-fluoro-N-(3-phenyl-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl)-benzamide;
- [0668] N-[3-(3,5-Dimethyl-phenyl)-1-oxa-2,4-diaza-spiro[4,4]non-2-en-4-yl]-4-ethyl-2-fluoro-benzamide;
- [0669] N-(5,5-Dimethyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-4-ethyl-2-fluoro-benzamide;
- [0670] 5-Ethyl-2,3-dihydro-benzo[1,4]dioxine-6-carboxylic acid (5,5-diethyl-3-phenyl-[1,2,4]oxadiazol-4-yl)-amide; and
- [0671] 5-Ethyl-2,3-dihydro-benzo[1,4]dioxine-6-carboxylic acid [3-(3,5-dimethyl-phenyl)-5-ethyl-5-methyl-[1,2,4]oxadiazol-4-yl]-amide.

[0672] In another embodiment, the activating ligand is a compound having Formula IX or X:



[0673] wherein R^1 , R^2 , R^3 , and R^4 are each independently:

[0674] a) H, (C_1-C_6) alkyl; (C_1-C_6) haloalkyl; (C_1-C_6) cycanoalkyl; (C_1-C_6) hydroxyalkyl; (C_1-C_4) alkoxy (C_1-C_6) alkyl; (C_2-C_6) alkenyl optionally substituted with halo, cyano, hydroxyl, or (C_1-C_4) alkyl; (C_2-C_6) alkynyl optionally substituted with halo, cyano, hydroxyl, or (C_1-C_4) alkyl; (C_3-C_5) cycloalkyl optionally substituted with halo, cyano, hydroxyl, or (C_1-C_4) alkyl; oxiranyl optionally substituted with halo, cyano, or (C_1-C_4) alkyl; or

[0675] b) unsubstituted or substituted benzyl wherein the substituents are independently 1 to 5H, halo, nitro, cyano, hydroxyl, (C_1-C_6) alkyl, or (C_1-C_6) alkoxy; and

[0676] R^5 is H; OH; F; Cl; or (C_1-C_6) alkoxy.

[0677] In another embodiment, the activating ligand is a compound selected from the group consisting of 20-hydroxyecdysone-2-methyl ether; 20-hydroxyecdysone-3-methyl ether; 20-hydroxyecdysone-14-methyl ether; 20-hydroxyecdysone-2,22-dimethyl ether; 20-hydroxyecdysone-3,22-dimethyl ether; 20-hydroxyecdysone-1,4,22-dimethyl ether; 20-hydroxyecdysone-22,25-dimethyl ether; 20-hydroxyecdysone-2,3,14,22-tetramethyl ether; 20-hydroxyecdysone-22-n-propyl ether; 20-hydroxyecdysone-22-n-butyl ether; 20-hydroxyecdysone-22-allyl ether; 20-hydroxyecdysone-22-benzyl ether; 20-hydroxyecdysone-22-benzyl ether; 20-hydroxyecdysone-22-(28R,S)-2'-ethyloxiranyl ether; ponasterone A-2-methyl ether; ponasterone A-14-methyl ether; ponasterone A-22-methyl ether; ponasterone A-2,22-dimethyl ether; ponasterone A-3,22-dimethyl ether; ponasterone A-14,22-dimethyl ether; dactyriansterone-22-me-

parameters normally encountered in medical treatment and gene expression systems and which are obvious to those skilled in the art are within the spirit and scope of the invention.

[0680] Pharmaceutical Compositions

[0681] In certain embodiments, polynucleotides and polypeptides of the invention can be administered as part of a medicament or pharmaceutical composition. Medicaments and pharmaceutical compositions of the invention comprise one or more pharmaceutically acceptable carriers, diluents, excipients or additives.

[0682] The term “excipient” as used herein is typically an inert substance added to a composition to facilitate processing, handling, administration, et cetera of a pharmaceutically acceptable composition. Useful excipients include, but are not limited to, adjuvants, anti-adherents, binders, carriers, disintegrants, fillers, flavors, colors, diluents, lubricants, glidants, preservatives, sorbents, solvents, surfactants, and sweeteners.

[0683] A few examples of pharmaceutically acceptable carriers, diluents, excipients and additives include, without limitation, water, saline, Ringer's solution, dextrose solution, buffers (such as phosphates (e.g., calcium phosphates such as tricalcium phosphate or calcium hydrogen phosphate)), citrate, succinate, acetic acid, and other organic acids or their salts), antioxidants, proteins and other high molecular weight molecules (such as serum albumin, gelatin, or immunoglobulins), hydrophilic polymers (such as polyvinylpyrrolidone), amino acids (such as glycine, glutamic acid, aspartic acid, and arginine), saccharides (for example monosaccharides, disaccharides, lactose, sucrose, mannitol, sorbitol, other carbohydrates and sugar-alcohols, cellulose or its derivatives, glucose, mannose, and dextrans), chelating agents (such as EDTA), sugar alcohols (such as mannitol or sorbitol), counterions (such as sodium), surfactants (such as polysorbates, poloxamers, or polyethylene glycol (PEG)), and binders (such as starch paste (e.g., maize starch, wheat starch, rice starch, potato starch)), gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone).

[0684] Pharmaceutically acceptable carriers, diluents, excipients and additives may include: disintegrating agents such as the above-mentioned starches as well as compounds such as carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate; and, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. In one embodiment, dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

[0685] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as

glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules or nanoparticles which may optionally be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In one embodiment, the is dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin, optionally with stabilizers.

[0686] Fatty oils may comprise mono-, di- or triglycerides. Mono-, di- and triglycerides include those that are derived from C6, C8, C10, C12, C14, C16, C18, C20 and C22 acids. Exemplary diglycerides include, in particular, diolein, dipalmitolein, and mixed caprylin-caprin diglycerides. Preferred triglycerides include vegetable oils, fish oils, animal fats, hydrogenated vegetable oils, partially hydrogenated vegetable oils, synthetic triglycerides, modified triglycerides, fractionated triglycerides, medium and long-chain triglycerides, structured triglycerides, and mixtures thereof. Exemplary triglycerides include: almond oil; babassu oil; borage oil; blackcurrant seed oil; canola oil; castor oil; coconut oil; corn oil; cottonseed oil; evening primrose oil; grapeseed oil; groundnut oil; mustard seed oil; olive oil; palm oil; palm kernel oil; peanut oil; rapeseed oil; safflower oil; sesame oil; shark liver oil; soybean oil; sunflower oil; hydrogenated castor oil; hydrogenated coconut oil; hydrogenated palm oil; hydrogenated soybean oil; hydrogenated vegetable oil; hydrogenated cottonseed and castor oil; partially hydrogenated soybean oil; partially soy and cottonseed oil; glyceryl tricaproate; glyceryl tricaprylate; glyceryl tricaprinate; glyceryl triundecanoate; glyceryl trilaurate; glyceryl trioleate; glyceryl trilinoleate; glyceryl trilinolenate; glyceryl tricaprylate/caprinate; glyceryl tricaprylate/caprinate/laurate; glyceryl tricaprylate/caprinate/linoleate; and glyceryl tricaprylate/caprinate/stearate.

[0687] In one embodiment, the triglyceride is the medium chain triglyceride available under the trade name LABRAFAC CC. Other triglycerides include neutral oils, e.g., neutral plant oils, in particular fractionated coconut oils such as known and commercially available under the trade name MIGLYOL, including the products: MIGLYOL 810; MIGLYOL 812; MIGLYOL 818; and CAPTEX 355. Other triglycerides are caprylic-capric acid triglycerides such as known and commercially available under the trade name MYRITOL, including the product MYRITOL 813. Further triglycerides of this class are CAPMUL MCT, CAPTEX 200, CAPTEX 300, CAPTEX 800, NEOBEE MS and MAZOL 1400.

[0688] Pharmaceutical compositions comprising triglycerides may further comprise lipophilic and/or hydrophilic surfactants which may form clear solutions upon dissolution with an aqueous solvent. One such surfactant is tocopheryl polyethylene glycol 1000 succinate (vitamin E TPGS). Examples of such compositions are described in U.S. Pat. No. 6,267,985.

[0689] In another embodiment, the pharmaceutically acceptable carrier comprises LABRASOL (Gattefosse SA), which is PEG-8 caprylic/capric glycerides. In another embodiment, the pharmaceutically acceptable carrier comprises PL90G, vitamin E TPGS, and Miglyol 812N.

[0690] Pharmaceutical compositions can be administered in any suitable manner as determined by those skilled in the art, such as, but without limitation, by oral, rectal, vaginal, topical (including dermal, buccal and sublingual), parenteral, intravenous, intraperitoneal, intramuscular, intratumoral,

intraarticular, subcutaneous, intranasal, inhalation, intradermal, intrathecal, epidural, and by naso-gastric routes.

[0691] Methods and compositions for preparation, formulation, and delivery of pharmaceutically acceptable compositions and medicaments are well-known and routinely practiced by those skilled in the art. A few examples of textbooks and manuals providing information and instruction on such methods and compositions include: Rowe et al. (Editor), "Handbook of Pharmaceutical Excipients," Pharmaceutical Press, 6th Ed. (August 2009) University of the Sciences in Philadelphia (Editor), "Remington The Science and Practice of Pharmacy," Lippincott Williams & Wilkins, 21st Ed. (2005), "Physicians' Desk Reference 2011," PDR Network (2010) "Physicians' Desk Reference 2012," PDR Network (2011); O'Neil, "The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals," 14th Ed. (2006); Allen et al (Editor) "Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems," Lippincott Williams & Wilkins 9th Ed. (2011); and, Ash et al. (Editor), "Handbook of Pharmaceutical Additives, Third Edition," Synapse Information Resources, Inc.; 3rd Ed. (2007).

[0692] Protocols for general molecular biology methods can be found in: *Methods in Molecular Biology*, series editor JM Walker, Humana Press, New York.

[0693] Embodiments of the invention comprise any amino acid substituted form of PE as indicated by, or represented in, Table 13. Embodiments of the invention further comprise any amino acid substituted form of PE which may comprise any combination of amino acid substitutions indicated by, or represented in, Table 13.

[0694] Embodiments of the invention also comprise variants, derivatives, or biologically active fragments of any amino acid substituted form of PE as indicated by, or represented in, Table 13, wherein said variant, derivative, or biologically active fragment of PE is at least 80% identical, at least 85% identical, at least 90% identical, at least 95% identical, at least 97% identical, at least 98% identical, at least 99% identical, or is at least 100% identical to an amino acid substituted form of PE, or a fragment thereof, as indicated by, or represented in, Table 13. For example, embodiments of the invention comprise variants, derivatives, or biologically active fragments of any amino acid substituted form of PE as indicated by, or represented in, Table 13, wherein said variant, derivative, or biologically active fragment of PE is at least 80% identical, at least 85% identical, at least 90% identical, at least 95% identical, at least 97% identical, at least 98% identical, at least 99% identical, or is at least 100% identical to PE constructs, or fragments thereof, as represented by pIEX02-003 through pIEX02-248 in Table 13 (such as, for example, as shown in SEQ ID NO:177 (pIEX02-228), SEQ ID NO:178 (pIEX02-244), and SEQ ID NO:179 (pIEX02-246)).

[0695] Embodiments of the invention include methods of making, methods of using, methods of treatment using, medicaments comprising, pharmaceutically acceptable compositions comprising, therapeutically useful compositions comprising, and kits comprising any of the amino acid substituted forms of PE referenced, or otherwise described or provided for, herein.

[0696] Embodiments of the invention also include (where "E" indicates "embodiment"):

[0697] E1. An isolated polypeptide having *Pseudomonas* exotoxin A biological activity, wherein said polypeptide comprises an epitope selected from the group consisting of:

a)	ISFSTRGTQ;	(SEQ ID NO: 5)
b)	GTQNWTVER;	(SEQ ID NO: 6)
c)	IVFGGVRAR;	(SEQ ID NO: 7)
d)	ARSQDLDAI;	(SEQ ID NO: 8)
e)	LRVYVPRSS;	(SEQ ID NO: 9)
f)	IPDKEQAIS;	(SEQ ID NO: 10)
g) and	ISFSTRSTQNWTVER;	(SEQ ID NO: 131)
h)	IVFGGVRARSQDLDAI	(SEQ ID NO: 132)

[0698] wherein one or more amino acid residues in any one or more of said epitopes in a) through h) are substituted with a different amino acid residue.

[0699] E2. An isolated polypeptide having *Pseudomonas* exotoxin A biological activity, wherein said polypeptide comprises an epitope selected from the group consisting of:

- [0700] a) ISFSTRGTQ (SEQ ID NO:5), wherein amino acid residues at one or more of positions 1, 6 and 9 are substituted with a different amino acid residue;
- [0701] b) GTQNWTVER (SEQ ID NO:6), wherein amino acid residues at one or more of positions 3, 4 and 6 are substituted with a different amino acid residue;
- [0702] c) IVFGCVRAR (SEQ ID NO:7), wherein amino acid residues at one or more of positions 1 and 6 are substituted with a different amino acid residue;
- [0703] d) ARSQDLDAI (SEQ ID NO:8), wherein amino acid residues at one or more of positions 4 and 7 are substituted with a different amino acid residue;
- [0704] e) LRVYVPRSS (SEQ ID NO:9), wherein amino acid residues at one or more of positions 1, 2 and 9 are substituted with a different amino acid residue;
- [0705] j) IPDKEQAIS (SEQ ID NO:10), wherein amino acid residues at one or more of positions 1, 4, 6 and 7 are substituted with a different amino acid residue;
- [0706] g) ISFSTRGTNWTVER (SEQ ID NO:131), wherein amino acid residues at one or more of positions 1, 6, 9, 10 and 12 are substituted, with a different amino acid residue; and
- [0707] h) IVFGGVRARSQDLDAI (SEQ ID NO:132), wherein amino acid residues at one or more of positions 1, 6, 11, and 14 are substituted with a different amino acid residue.

[0708] E3. The isolated polypeptide of embodiment E1 or E2, wherein said different amino acid residue is a conservative amino acid substitution.

[0709] E4. The isolated polypeptide of embodiment E3, wherein said conservative amino acid substitution is one or more substitutions selected from the group consisting of:

- [0710] a) A is substituted with any one of G, I, L, S, T or V;
- [0711] b) D is substituted with E;
- [0712] c) I is substituted with any one of L, M or V;
- [0713] d) K is substituted with any one of H or R;

- [0714] e) L is substituted with any one of A, G, I, M or V;
- [0715] f) N is substituted with any one of S, T or Q;
- [0716] g) Q is substituted with any one of S, T or N;
- [0717] h) R is substituted with any one of K or H;
- [0718] i) S is substituted with any one of A, G, N, T or Q;
- [0719] j) T is substituted with any one of A, G, N, Q or S; and
- [0720] k) V is substituted with any one of A, G, L or M.
- [0721] E5. An isolated polypeptide having *Pseudomonas* exotoxin A biological activity, wherein said polypeptide comprises an epitope selected from the group consisting of:
- [0722] a) ISFSTRGTQ (SEQ ID NO:5), wherein amino the acid residue at position 1 (I) is substituted with A, N, T, Q or H, or wherein the amino acid residue at position 6 (R) is substituted with Q, or wherein the amino acid residue at position 9 (Q) is substituted with N or T, or wherein the amino acid sequence ISFSTRGTQ (SEQ ID NO:5) comprises two or more of said substitutions in any combination;
- [0723] b) GTQNWTVR (SEQ ID NO:6), wherein the amino acid residue at position 3 (Q) is substituted with N or T, wherein amino the acid residue at position 4 (N) is substituted with K or R, or wherein the amino acid residue at position 6 (T) is substituted with K or R, or wherein the amino acid sequence GTQNWTVR (SEQ ID NO:6) comprises two or more of said substitutions in any combination;
- [0724] c) IVFGGVRAR (SEQ ID NO:7), wherein amino the acid residue at position 1 (I) is substituted with A or N, or wherein the amino acid residue at position 6 (V) is substituted with D, M, or N, or wherein the amino acid sequence IVFGGVRAR (SEQ ID NO:7) comprises substitutions at both positions in any combination of amino acid residues A or N at position 1 (I) and D, M, or N at position 6 (V);
- [0725] d) ARSQDLDAI (SEQ ID NO:8), wherein amino the acid residue at position 4 (Q) is substituted with K or R, or wherein the amino acid residue at position 7 (D) is substituted with K or R, or wherein the amino acid sequence ARSQDLDAI (SEQ ID NO:8) comprises substitutions with K or R in any combination at both positions 4 (Q) and 7 (D);
- [0726] e) LRVYVPRSS (SEQ ID NO:9), wherein amino the acid residue at position 1 (L) is substituted with A, or wherein the amino acid residue at position 2 (R) is substituted with D, S or A, or wherein the amino acid residue at position 9 (S) is substituted with D, E, N, K, P or T, or wherein the amino acid sequence LRVYVPRSS (SEQ ID NO:9) comprises two or more of said substitutions in any combination;
- [0727] f) IPDKEQAIS (SEQ ID NO:10), wherein amino acid residues at one or more of positions 1, 4, 6 and 7 are substituted with a different amino acid residue, wherein amino the acid residue at position 1 (I) is substituted with A, N, T, Q or H, or wherein the amino acid residue at position 4 (K) is substituted with T, or wherein the amino acid residue at position 6 (Q) is substituted with D, or wherein the amino acid residue at position 7 (A) is substituted with D, or wherein the amino acid sequence IPDKEQAIS (SEQ ID NO:10) comprises two or more of said substitutions in any combination;
- [0728] g) ISFSTRGTQNWTVER (SEQ ID NO:131), wherein amino acid residues at one or more of positions 1, 6, 9, 10 and 12 are substituted with a different amino acid residues wherein amino the acid residue at position 1 (I) is substituted with A, N, T, Q or H, or wherein the amino acid residue at position 6 (R) is substituted with Q, or wherein the amino acid residue at position 9 (Q) is substituted with N or T, or wherein amino the acid residue at position 10 (N) is substituted with K or R, or wherein the amino acid residue at position 12 (T) is substituted with K or R, or wherein the amino acid sequence ISFSTRGTQNWTVER (SEQ ID NO:131) comprises two or more of said substitutions in any combination; and
- [0729] h) IVFGGVRARSQDLDAI (SEQ ID NO:132), wherein amino the acid residue at position 1 (I) is substituted with A or N, or wherein the amino acid residue at position 6 (V) is substituted with D, M, M, or N, wherein amino the acid residue at position 11 (Q) is substituted with K or R, or wherein the amino acid residue at position 14 (D) is substituted with K or R, or wherein the amino acid sequence IVFGGVRARSQDLDAI (SEQ ID NO:132) comprises two or more of said substitutions in any combination.
- [0730] E6. An isolated polypeptide having *Pseudomonas* exotoxin A biological activity, wherein said polypeptide comprises an epitope selected from the group consisting of:
- [0731] a) I at position 141 is a different amino acid;
- [0732] b) R at position 146 is a different amino acid;
- [0733] c) Q at position 149 is a different amino acid;
- [0734] d) N at position 150 is a different amino acid;
- [0735] e) T at position 152 is a different amino acid;
- [0736] f) I at position 184 is a different amino acid;
- [0737] g) V at position 189 is a different amino acid;
- [0738] h) Q at position 194 is a different amino acid;
- [0739] i) L at position 197 is a different amino acid;
- [0740] j) L at position 233 is a different amino acid;
- [0741] k) R at position 234 is a different amino acid;
- [0742] l) S at position 241 is a different amino acid;
- [0743] m) I at position 321 is a different amino acid;
- [0744] n) K at position 324 is a different amino acid;
- [0745] o) Q at position 326 is a different amino acid;
- [0746] p) A at position 327 is a different amino acid;
- [0747] q) any combination of one or more of a) through p),
- [0748] wherein the amino acid numbering corresponds to SEQ ID NO:1.
- [0749] E7. An isolated polypeptide comprising an amino acid sequence identical to SEQ ID NO:1, except for one or more amino acid substitutions selected from the group consisting of:
- [0750] a) I at position 141 is substituted with a conservative amino acid substitution;
- [0751] b) R at position 146 is substituted with a conservative amino acid substitution;
- [0752] c) Q at position 149 is substituted with a conservative amino acid substitution;
- [0753] d) N at position 150 is substituted with a conservative amino acid substitution;

- [0754] e) T at position 152 is substituted with a conservative amino acid substitution;
- [0755] f) I at position 184 is substituted with a conservative amino acid substitution;
- [0756] g) V at position 189 is substituted with a conservative amino acid substitution;
- [0757] h) Q at position 194 is substituted with a conservative amino acid substitution
- [0758] i) at d position 197 is substituted with a conservative amino acid substitution;
- [0759] j) L at position 233 is substituted with a conservative amino acid substitution
- [0760] k) R at position 234 is substituted with a conservative amino acid substitution;
- [0761] l) S at position 241 is substituted with a conservative amino acid substitution;
- [0762] m) I at position 321 is substituted with a conservative amino acid substitution;
- [0763] n) K at position 324 is substituted with a conservative amino acid substitution;
- [0764] o) Q at position 326 is substituted with a conservative amino acid substitution;
- [0765] p) A at position 327 is substituted with a conservative amino acid substitution;
- [0766] q) any combination of one or more of a) through p),
- [0767] wherein the amino acid numbering corresponds to SEQ ID NO:1.
- [0768] E8. The isolated polypeptide of embodiment E7, wherein said conservative amino acid substitution is one or more substitutions selected from the group consisting of:
- [0769] a) A is substituted with any one of G, I, L, S, T or V;
- [0770] b) D is substituted with E;
- [0771] c) I is substituted with any one of L, M or V;
- [0772] d) K is substituted with any one of H or R;
- [0773] e) I is substituted with any one of A, G, I, M or V;
- [0774] f) N is substituted with any one of S, T or Q;
- [0775] g) Q is substituted with any one of S, T or N;
- [0776] h) R is substituted with any one of K or H;
- [0777] i) S is substituted with any one of A, O, N, T or Q;
- [0778] j) T is substituted with any one of A, O, N, Q or S; and
- [0779] k) V is substituted with any one of A, G, I, L or M.
- [0780] E9. An isolated polypeptide comprising an amino acid sequence identical to SEQ ID NO:1, except for one or more amino acid substitutions selected, from the group consisting of:

- | | |
|----|-------------------------|
| a) | I at position 141 is A; |
| b) | I at position 141 is N; |
| c) | I at position 141 is T; |
| d) | I at position 141 is Q; |
| e) | I at position 141 is H; |
| f) | R at position 146 is Q; |
| g) | Q at position 149 is N; |
| h) | Q at position 149 is T; |
| i) | N at position 150 is R; |
| j) | N at position 150 is K; |
| k) | T at position 152 is R; |

-continued

- | | |
|-----|--|
| l) | T at position 152 is K; |
| m) | I at position 184 is A; |
| n) | I at position 184 is N; |
| o) | V at position 189 is D; |
| p) | V at position 189 is M; |
| q) | V at position 189 is N; |
| r) | Q at position 194 is R; |
| s) | Q at position 194 is K; |
| t) | D at position 197 is R; |
| u) | D at position 197 is K; |
| v) | L at position 233 is A; |
| w) | R at position 234 is D; |
| x) | R at position 234 is S; |
| y) | R at position 234 is A; |
| z) | S at position 241 is D; |
| ab) | S at position 241 is E; |
| ac) | S at position 241 is N; |
| ad) | S at position 241 is K; |
| ae) | S at position 241 is P; |
| af) | S at position 241 is T; |
| ag) | I at position 321 is A; |
| ah) | I at position 321 is N; |
| ai) | I at position 321 is T; |
| ak) | I at position 321 is Q; |
| al) | I at position 321 is H; |
| am) | K at position 324 is T; |
| an) | Q at position 326 is D; |
| ao) | A at position 327 is D; |
| ap) | any combination, of one or more of a) through ao). |

[0781] wherein the amino acid numbering corresponds to SEQ ID NO:1.

[0782] E10. The polypeptide in any one of embodiments E1 to E9, comprising a number of amino acid substitutions selected from the group consisting of:

- | | |
|----|----------------------------------|
| a) | 1 amino acid substitution; |
| b) | 2 amino acid substitutions; |
| c) | 3 amino acid substitutions; |
| d) | 4 amino acid substitutions; |
| e) | 5 amino acid substitutions; |
| f) | 6 amino acid substitutions; |
| g) | 7 amino acid substitutions; |
| h) | 8 amino acid substitutions; |
| i) | 9 amino acid substitutions; |
| j) | 10 amino acid substitutions; |
| k) | 11 amino acid substitutions; |
| l) | 12 amino acid substitutions; |
| m) | 13 amino acid substitutions; |
| n) | 14 amino acid substitutions; |
| o) | 15 amino acid substitutions; and |
| p) | 16 amino acid substitutions. |

[0783] E1. The polypeptide of embodiment E9, comprising amino acid substitutions present at each of amino acid positions 141, 146, 149, 150, 152, 184, 189, 194, 197, 233, 234, 241, 321, 324, 326 and 327.

[0784] E12. The polypeptide of any one of embodiments E1 to E11, wherein said polypeptide comprises the amino acid sequence in SEQ ID NO:1, except for amino acid substitutions indicated in embodiments E1 to E11.

[0785] E13. The polypeptide in any one of embodiments E1 to E11, wherein said polypeptide is a variant or fragment of a *Pseudomonas* exotoxin A polypeptide.

[0786] E14. The polypeptide of embodiment E13, wherein said variant or fragment comprises a number of epitopes selected from the group consisting of:

[0787] a) at least one epitope;

[0788] b) at least two epitopes;

- [0789] c) at least three epitopes;
 [0790] d) at least four epitopes;
 [0791] e) at least five epitopes; and
 [0792] f) at least six epitopes.
- [0793] E15. The polypeptide of embodiment E14, wherein said polypeptide comprises a contiguous number of amino acids selected from the group consisting of:
 [0794] a) at least 20 contiguous amino acids;
 [0795] b) at least 30 contiguous amino acids;
 [0796] c) at least 40 contiguous amino acids;
 [0797] d) at least 50 contiguous amino acids;
 [0798] e) at least 60 contiguous amino acids;
 [0799] f) at least 70 contiguous amino acids;
 [0800] g) at least 80 contiguous amino acids;
 [0801] h) at least 90 contiguous amino acids;
 [0802] i) at least 100 contiguous amino acids;
 [0803] j) at least 125 contiguous amino acids;
 [0804] k) at least 150 contiguous amino acids;
 [0805] l) at least 175 contiguous amino acids;
 [0806] m) at least 200 contiguous amino acids;
 [0807] n) at least 225 contiguous amino acids;
 [0808] o) at least 250 contiguous amino acids;
 [0809] p) at least 275 contiguous amino acids;
 [0810] q) at least 300 contiguous amino acids;
 [0811] r) at least 325 contiguous amino acids; and
 [0812] s) at least 350 contiguous amino acids.
- [0813] E16. An isolated polypeptide comprising a *Pseudomonas* exotoxin A (PE-A) cytotoxic domain (Domain III), wherein the cytotoxic domain comprises one or more amino acid substitutions which prevent or reduce host immunogenic responses compared to the same polypeptide without said one or more amino acid substitutions.
- [0814] E17. The polypeptide of embodiment E16, wherein said one or more amino acid substitutions are introduced into a cytotoxic domain sequence selected from the group consisting of:
 [0815] (a) amino acid residues Phe-134 to Lys-347 of SEQ ID NO:1;
 [0816] (b) amino acid residues Phe-134 to Lys-347 of SEQ ID NO:4;
 [0817] (c) amino acid residues Phe-400 to Lys-613 of SEQ ID NO:133; and
 [0818] (d) amino acid residues Phe-400 to Lys-613 of SEQ ID NO:134.
- [0819] E18. The polypeptide of embodiment E17, wherein the last five amino acids in said cytotoxic domain are replaced with an amino acid sequence selected from the group consisting of:
 (i) Arg-Glu-Asp-Leu; (SEQ ID NO: 136)
 and
 (ii) Lys-Asp-Glu-Leu. (SEQ ID NO: 137)
- [0820] E19. The polypeptide in any one of embodiments E16 to E18, wherein said polypeptide further comprises one or more PE-A domains selected from the group consisting of:
 [0821] (a) a cytosolic translocation domain (Domain II);
 [0822] (b) a carboxy-terminal portion of Domain IB;
 [0823] (c) an amino-terminal portion of Domain IB; and
 [0824] (d) a complete Domain IB.
- [0825] wherein one or more of said domains has been modified with amino acid substitutions, as described herein, to reduce or eliminate immunogenicity.
- [0826] E20. The polypeptide of embodiment E19, wherein said cytosolic translocation domain (Domain II) comprises an amino acid sequence selected from the group consisting of:
 [0827] (a) amino acids Gly-3 to Ser-114 of SEQ ID NO:1; and
 [0828] (b) amino acids Gly-3 to Asn-114 of SEQ ID NO:4.
- [0829] E21. The polypeptide of embodiment E19, wherein said carboxy-terminal portion of Domain IB comprises the amino acid sequence of Gly-115 to Glu-133 of SEQ ID NO:1 or wherein said amino-terminal portion of Domain IB comprises the amino acid sequence of Ala-365 to Ala-380 of SEQ ID NO:1.33.
- [0830] E22. The polypeptide of embodiment E19, wherein said complete Domain IB comprises the amino acid sequence of Ala-365 to Glu-399 of SEQ ID NO:1.33.
- [0831] E23. The polypeptide in any one of embodiments E16 to E22, wherein said polypeptide is a variant or fragment of a *Pseudomonas* exotoxin A polypeptide.
- [0832] E24. The polypeptide of any one of embodiments E1 to E23, wherein said polypeptide has one or more biological activities selected from the group consisting of:
 [0833] a) eukaryotic cell killing activity (cell cytotoxicity);
 [0834] b) inhibits translation elongation factor EF-2 biological activity;
 [0835] c) induces or catalyzes ADP-ribosylation of EF-2; and
 [0836] d) inhibits protein synthesis.
- [0837] E25. The polypeptide of any one of embodiments E1 to E24, wherein said one or more amino acid substitutions prevent or reduce host immunogenic responses compared to the same polypeptide without the corresponding said one or more amino acid substitutions.
- [0838] E26. The polypeptide of any one of embodiments E1 to E24, wherein said one or more amino acid substitutions prevent or reduce host immunogenic responses compared to a polypeptide comprising an amino acid sequence selected from the group consisting of:
 [0839] (a) SEQ ID NO:1;
 [0840] (b) SEQ ID NO:4;
 [0841] (c) SEQ ID NO:133; and
 [0842] (d) SEQ ID NO:134.
- [0843] E27. The polypeptide of any one of embodiments E1 to E26, wherein said polypeptide is a fusion protein.
- [0844] E28. The fusion protein of embodiment E27, wherein the amino-terminal end of said polypeptide in any one of embodiments E1 to E26 is fused to the carboxyl-terminal end of a different polypeptide.
- [0845] E29. The fusion protein of embodiment E27, wherein the carboxyl-terminal end of said polypeptide in any one of embodiments E1 to E26 is fused to the amino-terminal end of a different polypeptide.

- [0846] E30. The fusion protein in embodiment E28 or E29, wherein said different polypeptide comprises an antigen binding moiety.
- [0847] E31. The fusion protein of embodiment E30, wherein said antigen binding moiety is an antibody or fragment thereof.
- [0848] E32. The fusion protein of embodiment E31, wherein said antibody, or fragment thereof, is an antibody selected from the list in Table 1, or is a fragment thereof.
- [0849] E33. The fusion protein of embodiment E31, wherein said antibody, or fragment thereof, specifically binds to a cancer-specific or tumor-specific antigen.
- [0850] E34. The fusion protein of embodiment E33, wherein said cancer-specific or tumor-specific antigen is a breast cancer antigen.
- [0851] E35. The fusion protein of embodiment E34, wherein said breast cancer antigen is HER2.
- [0852] E36. The fusion protein of embodiment E31, wherein said antibody, or fragment thereof is selected from the group consisting of:
- [0853] a) ERTUMAXOMAB (Rexotmun);
- [0854] b) PERTUZUMAB (Omnitarg); and
- [0855] c) TRASTUZUMAB (Herceptin).
- [0856] E37. The fusion protein of any one of embodiments E27 to E29, wherein said different polypeptide comprises a polypeptide selected from the group consisting of:
- [0857] a) Mesothelin;
- [0858] b) CD24;
- [0859] c) CD22;
- [0860] d) CD25;
- [0861] e) CD174;
- [0862] f) TPBG
- [0863] g) CD56; and
- [0864] h) C-type lectin-like molecule-1.
- [0865] E38. An isolated polynucleotide encoding the polypeptide or fusion protein in any one of embodiments E1 to E37.
- [0866] E39. An expression vector comprising the polynucleotide of embodiment E38.
- [0867] E40. A host cell comprising the expression vector of embodiment E39.
- [0868] E41. A method of producing the polypeptide or fusion protein in any one of embodiments E1 to E37, wherein said method comprises:
- [0869] a) obtaining a host cell comprising a polynucleotide encoding said polypeptide or fusion protein;
- [0870] b) exposing said host cell to conditions wherein said polypeptide or fusion protein is produced.
- [0871] E42. A method of producing the polypeptide or fusion protein in any one of embodiments E1 to E37, wherein said method comprises use of an expression system comprising:
- [0872] (A) a first polynucleotide encoding a first hybrid polypeptide comprising:
- [0873] (i) a first ligand binding domain; and
- [0874] (ii) a DNA-binding domain;
- [0875] (B) a second polynucleotide encoding a second hybrid polypeptide comprising:
- [0876] (i) a second ligand binding domain; and
- [0877] (ii) a transactivation domain;
- [0878] (C) a third polynucleotide encoding the polypeptide or fusion protein in any one of embodiments E1 to E37, wherein said third polynucleotide is operably associated with a response element capable of being bound by the DNA-binding domain of said first hybrid polypeptide;
- [0879] wherein the first ligand binding domain and the second ligand binding domain are capable of ligand-induced dimerization,
- [0880] wherein expression of the polypeptide or fusion protein in any one of embodiments E1 to E37 is modulated by a ligand which induces dimerization of said first and said second ligand binding domains,
- [0881] wherein the polypeptide or fusion protein in any one of embodiments E1 to E37 is produced by allowing said ligand to contact said first and said second ligand binding domains.
- [0882] E43. A single expression vector or two or more expression vectors comprising the first, second, and third polynucleotides of embodiment E42.
- [0883] E44. The expression vector or expression vectors of embodiment E43, wherein one or more of the vectors is a viral expression vector.
- [0884] E45. A host cell comprising the expression vector or expression vectors of embodiments E43 or E44.
- [0885] E46. A method of treating a disease or disorder comprising administering to a subject in need thereof the polypeptide or fusion protein in any one of embodiments E1 to E37, the polynucleotide of embodiment E38, the vector of embodiment E39, the host cell of embodiment E40, or a polypeptide or fusion protein produced by the method of embodiment E41,
- [0886] E47. A method of treating a disease or disorder comprising delivering to a subject in need thereof a polypeptide or fusion protein produced by the method of embodiment E42, wherein said method comprises administration of the ligand to said subject.
- [0887] E48. The method of embodiment E47, wherein the polypeptide or fusion protein is delivered to the subject by first administering the first, second, and third polynucleotides.
- [0888] E49. The method of embodiment E47, wherein the polypeptide or fusion protein is delivered to the subject by first administering the expression vector or expression vectors of embodiments E43 or E44.
- [0889] E50. The method of embodiment E47, wherein said polypeptide or fusion protein is delivered to the subject by first administering the host cell of embodiment E45.
- [0890] E51. A pharmaceutical composition comprising the polypeptide or fusion protein in any one of embodiments E1 to E37, comprising the polynucleotide of embodiment E38, comprising the expression vector or expression vectors in any one of embodiments E39, E43 or E44, or comprising the host cell of embodiments E40 or E45, and a pharmaceutically acceptable carrier, diluent or excipient.
- [0891] E52. A medicament comprising the polypeptide or fusion protein in any one of embodiments E1 to E37, comprising the polynucleotide of embodiment E38, comprising the expression vector or expression vectors in any one of embodiments E39, E43 or E44, or comprising the host cell of embodiments E40 or E45.

- [0892] E53. Use of the medicament of embodiment E52, wherein said use is for the treatment of a disease or disorder.
- [0893] E54. Use of the medicament according to embodiment E53, wherein the disease or disorder is cancer.
- [0894] E55. A polypeptide having at least one *Pseudomonas* exotoxin A (PE-A) biological activity, wherein said polypeptide comprises one or more amino acid substitutions compared to a wild-type PE-A polypeptide, wherein said one or more amino acid substitutions is a substitution of a different amino acid at one or more positions corresponding to amino acid residues in the polypeptide of SEQ ID NO:1, wherein said substitution positions are selected from the group consisting of:
- [0895] a) isoleucine (I) at position 141 ;
 - [0896] b) arginine (R) at position 146;
 - [0897] c) glycine (G) at position 147;
 - [0898] d) glutamine (Q) at position 149;
 - [0899] e) asparagine (N) at position 150;
 - [0900] f) threonine (T) at position 152;
 - [0901] g) valine (V) at position 189;
 - [0902] h) arginine (R) at position 192;
 - [0903] i) glutamine (Q) at position 194;
 - [0904] j) aspartic acid (D) at position 197;
 - [0905] k) serine (S) at position 241;
 - [0906] l) isoleucine (I) at position 321; and
 - [0907] m) glutamine (Q) at position 326.
- [0908] E56. The polypeptide of embodiment E55, wherein said one or more amino acid substitutions is a conservative amino acid substitution.
- [0909] E57. The polypeptide of embodiment E55, wherein said one or more amino acid substitutions is selected from the group consisting of:
- [0910] a) isoleucine (I) at position 141 is substituted with alanine (A), threonine (T), or histidine (H);
 - [0911] b) arginine (R) at position 146 is substituted with glutamine (Q) or alanine (A);
 - [0912] c) glycine (G) at position 147 is substituted with serine (S);
 - [0913] d) glutamine (Q) at position 149 is substituted with threonine (T);
 - [0914] e) asparagine (N) at position 150 is substituted with alanine (A);
 - [0915] f) threonine (T) at position 152 is substituted with alanine (A) or arginine (R);
 - [0916] g) valine (V) at 189 is substituted with alanine (A);
 - [0917] h) arginine (R) at position 192 is substituted with alanine (A) or glutamine (Q);
 - [0918] i) glutamine (Q) at position 194 is substituted with arginine (R);
 - [0919] j) aspartic acid (D) at position 197 is substituted with lysine (K);
 - [0920] k) serine (S) at position 241 is substituted with threonine (T), asparagine (N), lysine (K), or proline (P);
 - [0921] l) isoleucine (I) at position 321 is substituted with alanine (A), asparagine (N), histidine (H), threonine (T), or glutamine (Q); and
 - [0922] m) glutamine (Q) at position 326 is substituted with glutamic acid (E).
- [0923] E58. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution for isoleucine (I) at position 141, a substitution for threonine (T) at position 152, a substitution for arginine (R) at position 192, a substitution for aspartic acid (D) at position 197, a substitution for serine (S) at position 241, and a substitution for glutamine (Q) at position 326.
- [0924] E59. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of threonine (T) or alanine (A) for isoleucine (I) at position 141, a substitution alanine (A) or arginine (R) for threonine (T) at position 152, a substitution of alanine (A) for arginine (R) at position 192, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.
- [0925] E60. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of threonine (T) for isoleucine (I) at position 141, a substitution alanine (A) for threonine (T) at position 152, a substitution of alanine (A) for arginine (R) at position 192, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.
- [0926] E61. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of alanine (A) for isoleucine (I) at position 141, a substitution alanine (A) for threonine (T) at position 152, a substitution of alanine (A) for arginine (R) at position 192, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.
- [0927] E62. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution for isoleucine (I) at position 141, a substitution for threonine (T) at position 152, a substitution for aspartic acid (D) at position 197, a substitution for serine (S) at position 241, and a substitution for glutamine (Q) at position 326.
- [0928] E63. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution for isoleucine (I) at position 141, a substitution for threonine (T) at position 152, a substitution for arginine (R) at position 192, a substitution for aspartic acid (D) at position 197, and a substitution for serine (S) at position 241.
- [0929] E64. The isolated polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of alanine (A) or threonine (T) for isoleucine (I) at position 141, a substitution of arginine (R) or alanine (A) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.
- [0930] E65. The isolated polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of alanine (A) for isoleucine (I) at position 141, a substitution of arginine (R) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.

- [0931] E66. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of alanine (A) for isoleucine (I) at position 141, a substitution of alanine (A) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.
- [0932] E67. The polypeptide of embodiment E55, wherein said polypeptide comprises a substitution of threonine (T) for isoleucine (I) at position 141, a substitution of alanine (A) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.
- [0933] E68. The polypeptide in any one of embodiments E55 to E67, wherein the at least one *Pseudomonas* exotoxin A (PE-A) biological activity comprises the ability to inhibit in vitro transcription/translation compared to a corresponding wild-type or non-substituted PE-A polypeptide, wherein said ability to inhibit in vitro transcription/translation is in an amount selected from the group consisting of:
- [0934] (a) at least 5% inhibition;
 - [0935] (b) at least 10% inhibition;
 - [0936] (c) at least 15% inhibition;
 - [0937] (d) at least 20% inhibition;
 - [0938] (e) at least 25% inhibition;
 - [0939] (f) at least 30% inhibition;
 - [0940] (g) at least 40% inhibition;
 - [0941] (h) at least 50% inhibition;
 - [0942] (i) at least 60% inhibition;
 - [0943] (j) at least 70% inhibition;
 - [0944] (k) at least 380% inhibition
 - [0945] (l) at least 90% inhibition;
 - [0946] (m) at least 100% inhibition;
 - [0947] (n) about 100% inhibition; and
 - [0948] (o) 100% inhibition.
- [0949] E69. The polypeptide in any one of embodiments E55 to E68, comprising a number of amino acid substitutions selected from the group consisting of:
- [0950] a) 1 amino acid substitution;
 - [0951] b) 2 amino acid substitutions;
 - [0952] c) 3 amino acid substitutions;
 - [0953] d) 4 amino acid substitutions;
 - [0954] e) 5 amino acid substitutions; and
 - [0955] f) 6 amino acid substitutions.
- [0956] E70. The polypeptide in any one of embodiments E55 to E69, wherein said polypeptide comprises one or more amino acid substitutions which prevent or reduce host immunogenic responses compared to the same polypeptide without said one or more amino acid substitutions.
- [0957] E71. The polypeptide of embodiment E70 wherein host immunogenic responses are prevented or reduced in a human host.
- [0958] E72. The polypeptide in any one of embodiments E55 to E71 wherein the last five or six amino acids in said polypeptide comprise one or more amino acid sequences selected from the group consisting of:
- [0959] (i) Arg-Glu-Asp-Leu-Lys;
 - [0960] (ii) Arg-Glu-Asp-Leu;
 - [0961] (iii) Lys-Asp-Glu-Leu;
 - [0962] (iv) Glu-Asp-Leu-Lys; and
 - [0963] (v) a dimer, trimer, pentamer, hexamer, septamer, or octamer of (i), (ii), or (iii), or any combination thereof
- [0964] E73. The polypeptide of any one of embodiments E55 to E72, wherein said polypeptide has one or more biological activities selected from the group consisting of:
- [0965] a) eukaryotic cell killing activity (cell cytotoxicity);
 - [0966] b) inhibits translation elongation factor EF-2 biological activity;
 - [0967] c) induces or catalyzes ADP-ribosylation of EF-2; and
 - [0968] d) inhibits protein synthesis.
- [0969] E74. The polypeptide of any one of embodiments E55 to E72, wherein said one or more amino acid substitutions prevent or reduce host immunogenic responses compared to the same polypeptide without the corresponding said one or more amino acid substitutions.
- [0970] E75. A polypeptide comprising a biologically active fragment of the polypeptide in any one of embodiments E55 to E74.
- [0971] E76. A polypeptide comprising a variant or derivative of the polypeptide in any one of embodiments E55 to E75, wherein said variant or derivative shares amino acid sequence identity with the polypeptide in any one of embodiments E55 to E75, wherein said shared amino acid sequence identity is selected from the group consisting of:
- [0972] a) at least 80% identity;
 - [0973] b) at least 85% identity;
 - [0974] c) at least 90% identity;
 - [0975] d) at least 95% identity;
 - [0976] e) at least 97% identity
 - [0977] f) at least 98% identity; and
 - [0978] g) at least 99% identity.
- [0979] E77. The polypeptide of any one of embodiments E55 to E76, wherein said one or more amino acid substitutions prevent or reduce host immunogenic responses compared to a polypeptide comprising an amino acid sequence selected from the group consisting of:
- [0980] (a) SEQ ID NO:1;
 - [0981] (b) SEQ ID NO:4;
 - [0982] (c) SEQ ID NO:133; and
 - [0983] (d) SEQ ID NO:134.
- [0984] E78. The polypeptide of any one of embodiments E55 to E77, wherein said polypeptide is a fusion protein.
- [0985] E79. The fusion protein of embodiment E78, wherein the amino-terminal end of said polypeptide in any one of embodiments E55 to E78 is fused to the carboxyl-terminal end of a different polypeptide.
- [0986] E80. The fusion protein of embodiment E78, wherein the carboxyl-terminal end of said polypeptide in any one of embodiments E55 to E78 is fused to the amino-terminal end of a different polypeptide.
- [0987] E81. The fusion protein in embodiment E79 or E80, wherein said different polypeptide comprises an antigen binding moiety.
- [0988] E82. The fusion protein of embodiment E81, wherein said antigen binding moiety is an antibody or fragment thereof.

- [0989] E83. The fusion protein of any one of embodiments E78 to E82, wherein said antibody, or fragment thereof, is an antibody selected from the list in Table 1, or is a fragment thereof.
- [0990] E84. The fusion protein of embodiment E82, wherein said antibody, or fragment thereof, specifically binds to a cancer-specific or tumor-specific antigen.
- [0991] E85. The fusion protein of embodiment E84, wherein said cancer-specific or tumor-specific antigen is a breast cancer antigen.
- [0992] E86. The fusion protein of embodiment E85, wherein said breast cancer antigen is HER2.
- [0993] E87. The fusion protein of embodiment E82, wherein said antibody, or fragment thereof is selected from the group consisting of:
- [0994] a) ERTUMAXOMAB (Rexomun);
 - [0995] b) PERTUZUMAB (Omnitarg); and
 - [0996] c) TRASTUZUMAB (Herceptin).
- [0997] E88. The fusion protein of any one of embodiments E78 to E80, wherein said different polypeptide comprises a polypeptide selected from the group consisting of
- [0998] a) Mesothelin;
 - [0999] b) CD24;
 - [1000] c) CD22;
 - [1001] d) CD25;
 - [1002] e) CD174;
 - [1003] j) TPBG;
 - [1004] g) CD56; and
 - [1005] h) C-type lectin-like molecule-1.
- [1006] E89. A polynucleotide encoding the polypeptide or fusion protein in any one of embodiments E55 to E88.
- [1007] E90. An expression vector comprising the polynucleotide of embodiment E89,
- [1008] E91. A host cell comprising the expression vector of embodiment E90.
- [1009] E92. A method of producing the polypeptide or fusion protein in any one of embodiments E55 to E88, wherein said method comprises:
- [1010] a) obtaining a host cell comprising a polynucleotide encoding said polypeptide or fusion protein;
 - [1011] b) exposing said host cell to conditions wherein said polypeptide or fusion protein is produced.
- [1012] E93. A method of producing the polypeptide or fusion protein in any one of embodiments E55 to E88, wherein said method comprises use of an expression system comprising:
- [1013] (A) a first polynucleotide encoding a first hybrid polypeptide comprising:
 - [1014] (i) a first ligand binding domain; and
 - [1015] (ii) a DNA-binding domain;
 - [1016] (B) a second polynucleotide encoding a second hybrid polypeptide comprising:
 - [1017] (i) a second ligand binding domain; and
 - [1018] (ii) a transactivation domain;
 - [1019] (C) a third polynucleotide encoding the polypeptide or fusion protein in any one of embodiments E55 to E88, wherein said third polynucleotide is operably associated with a response element capable of being bound by the DNA-binding domain of said first hybrid polypeptide;
- [1020] wherein the first ligand binding domain and the second ligand binding domain are capable of ligand-induced dimerization,
- [1021] wherein expression of the polypeptide or fusion protein in any one of embodiments E55 to E88 is modulated by a ligand which induces dimerization of said first and said second ligand binding domains,
- [1022] wherein the polypeptide or fusion protein in any one of embodiments E55 to E88 is produced by allowing said ligand to contact said first and said second ligand binding domains.
- [1023] E94. A single expression vector or two or more expression vectors comprising the first, second, and third polynucleotides of embodiment E93.
- [1024] E95. The expression vector or expression vectors of embodiment E94, wherein one or more of the vectors is a viral expression vector.
- [1025] E96. A host cell comprising the expression vector or expression vectors of embodiments E94 or E95.
- [1026] E97. A method of treating a disease or disorder comprising administering to a subject in need thereof the polypeptide or fusion protein in any one of embodiments E55 to E88, the polynucleotide of embodiment E89, the expression vector or expression vectors in any one of embodiments E90, E94 or E95, the host cell of embodiments E91 or E96, or a polypeptide or fusion protein produced by the method of embodiment E92,
- [1027] E98. A method of treating a disease or disorder comprising delivering to a subject in need thereof a polypeptide or fusion protein produced by the method of embodiment E93, wherein said method comprises administration of the ligand to said subject.
- [1028] E99. The method of embodiment E44, wherein the polypeptide or fusion protein is delivered to the subject by first administering the first, second, and third polynucleotides.
- [1029] E100. The method of embodiment E98, wherein the polypeptide or fusion protein is delivered to the subject by first administering the expression vector or expression vectors of embodiments E94 or E95.
- [1030] E101. The method of embodiment E98, wherein said polypeptide or fusion protein is delivered to the subject by first administering the host cell of embodiments E91 or E96.
- [1031] E102. A pharmaceutical composition comprising the polypeptide or fusion protein in any one of embodiments E55 to E88, comprising the polynucleotide of embodiment E89, comprising the expression vector or expression vectors in any one of embodiments E90, E94 or E95, or comprising the host cell of embodiments E91 or E96, and a pharmaceutically acceptable carrier, diluent or excipient.
- [1032] E103. A medicament comprising the polypeptide or fusion protein in any one of embodiments E55 to E88, comprising the polynucleotide of embodiment E89, comprising the expression vector or expression vectors in any one of embodiments E90, E94 or E95, or comprising the host cell of embodiments E91 or E96.
- [1033] E104. Use of the pharmaceutical composition of embodiment E102 or the medicament of embodiment E103, wherein said use is for the treatment of a disease or disorder.

[1034] E105. Use of the pharmaceutical composition or the medicament according to embodiment E104, wherein the disease or disorder is cancer.

[1035] E106. An *Pseudomonas* exotoxin A (PE-A) polypeptide, wherein said polypeptide comprises a mutation at a position corresponding to amino acid position E184 in SEQ ID NO:1 (or position E196 in SEQ ID NO:2) wherein an isoleucine at position E184 (or position 196 in SEQ ID NO:2) is substituted with a different amino acid.

[1036] E107. The polypeptide of embodiment E106, wherein said polypeptide does not have PE-A biological activity.

[1037] E108. A method for assaying the immunogenicity of a mutated form of *Pseudomonas* exotoxin A (PE-A), wherein said method comprises:

[1038] (a) contacting immune cells with a mutated form of PE-A; and

[1039] (b) assaying immune cell stimulation,

[1040] wherein said mutated form of PE-A comprises a mutation at a position corresponding to amino acid position E184 in SEQ ID NO:1 (or position 196 in SEQ ID NO:2) wherein an isoleucine at position E184 (or position 196 in SEQ ID NO:2) is substituted with a different amino acid, and wherein said mutated form of PE-A also comprises one or more additional amino acid substitutions compared to a wild-type form of PE-A.

[1041] E109. The method of embodiment E108, wherein said immune cells are human immune cells.

[1042] E110. The method of embodiment E109, wherein said immune cells are human T-cells, cells of a human T-cell lineage, human B-cells, or cells of a human B-cell lineage.

[1043] E111. The method of embodiment E47, wherein the ligand is a compound having Formula I, or a pharmaceutically acceptable salt thereof.


[1044] E112. The method of embodiment E47, wherein the ligand is a compound having Formula II, or a pharmaceutically acceptable salt thereof.

[1045] E113. The method of embodiment E47, wherein the ligand is a compound having Formula III, or a pharmaceutically acceptable salt thereof.


[1046] E114. The method of embodiment E47, wherein the ligand is a compound of Table 3, or a pharmaceutically acceptable salt thereof.

[1047] E115. The method of embodiment E47, wherein the ligand is a compound having Formula II, wherein:

[1048] A is:



[1049] B is:



[1050] R^{3a} , R^{3b} , R^{3c} , R^{3d} , R^{3e} , R^{3f} , R^{3g} , R^{3h} , R^{3i} and R^{3j} are independently selected from hydrogen, halo, (C_1-C_4) alkyl, or (C_1-C_4) alkoxy;

[1051] R^1 is (C_1-C_6) alkyl, hydroxy (C_1-C_4) alkyl, or (C_2-C_4) alkenyl; and

[1052] R^2 is optionally substituted (C_1-C_6) alkyl, or a pharmaceutically acceptable salt thereof.

[1053] E116. The method of embodiment E47, wherein the ligand is a compound selected from the group consisting of:

[1054] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-2-ethyl-3-methoxy-benzoyl-hydrazide;

[1055] (R)-3,5-Dimethyl-benzoic acid N'-benzoyl-N-(1-tert-butyl-butyl)-hydrazide;

[1056] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-methyl-benzoyl)-hydrazide;

[1057] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-methoxy-benzoyl)-hydrazide;

[1058] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-fluoro-benzoyl)-hydrazide;

[1059] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-chloro-benzoyl)-hydrazide;

[1060] (R)-3,5-Dimethyl-benzoic acid N'-(2-bromo-benzoyl)-N-(1-tert-butyl-butyl)-hydrazide;

[1061] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methyl-benzoyl)-hydrazide;

[1062] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-chloro-benzoyl)-hydrazide;

[1063] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-methyl-benzoyl)-hydrazide;

[1064] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-ethyl-benzoyl)-hydrazide;

[1065] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-methoxy-benzoyl)-hydrazide;

[1066] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-chloro-benzoyl)-hydrazide;

[1067] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2,6-difluoro-benzoyl)-hydrazide;

[1068] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2,6-dichloro-benzoyl)-hydrazide;

- [1069] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3,4-dimethoxy-benzoyl)-hydrazide;
- [1070] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3,5-difluoro-benzoyl)-hydrazide;
- [1071] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3,5-dimethoxy-4-methyl-benzoyl)-hydrazide;
- [1072] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(4-methyl-benzo[1,3]dioxole-5-carbonyl)-hydrazide;
- [1073] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(5-methyl-2,3-dihydro-benzo[1,4]dioxole-6-carbonyl)-hydrazide;
- [1074] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(5-ethyl-2,3-dihydro-benzo[1,4]dioxole-6-carbonyl)-hydrazide;
- [1075] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(naphthalene-1-carbonyl)-hydrazide;
- [1076] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(naphthalene-2-carbonyl)-hydrazide;
- [1077] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(thiophene-2-carbonyl)-hydrazide;
- [1078] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2,5-dimethyl-furan-3-carbonyl)-hydrazide;
- [1079] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(2-chloro-pyridine-3-carbonyl)-hydrazide;
- [1080] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(6-chloro-pyridine-3-carbonyl)-hydrazide;
- [1081] (R)-3,5-Dimethyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide;
- [1082] (R)-3,5-Dimethoxy-4-methyl-benzoic acid N-(1-tert-butyl-butyl)-N'-(3-methoxy-2-methyl-benzoyl)-hydrazide; and
- [1083] (R)-3,5-Dimethyl-benzoic acid N'-(4-ethyl-benzoyl)-N-(1-phenethyl-but-3-enyl)-hydrazide,
- [1084] or a pharmaceutically acceptable salt thereof.

EXAMPLES

Example 1

T Cell Epitope Mapping of PE

[1085] Peptides spanning the sequence of an approximately 38 kD form of *Pseudomonas* exotoxin A protein ("PE38") were analyzed for the presence of immunogenic CD4+ T cell epitopes using EPISCREEN™ T cell epitope mapping analysis (Antitope Ltd, Cambridge, UK).

[1086] EPISCREEN™ is a proprietary technology commercially available through Antitope Ltd, Cambridge, UK, to map T cell epitopes within a protein sequence to determine potential for immunogenicity (based on the number and potency of T cell epitopes within a sequence). EPISCREEN™ T cell epitope mapping typically uses CD8+ T cell depleted PBMCs from a minimum of 50 HLA-typed donors (selected to represent the human population of interest). Typically, 15 mer peptides with 12 amino acid overlaps spanning a protein sequence are analyzed in a large number of replicate cultures for in vitro CD4+ T cell stimulation by 3H TdR incorporation. CD4+ T cell stimulation is often detected in two or three adjacent and overlapping peptides since the core 9 mer that binds the MHC class II binding groove will be present in more than one peptide sequence. Following identification of peptides that stimulate CD4+ T cells in vitro in silico technology can be used to design epitope-depleted (de-

immunized) variants by determining the precise location of core 9 mer sequences and the location of key MHC class II anchor residues.

[1087] A total of 120 overlapping 15 mer peptides spanning the entire PE38 sequence (SEQ ID NO:2), including 4 peptides covering a null mutation and 4 peptides spanning an N-terminal linker sequence (SEQ ID NO:3) were tested against a cohort of 52 healthy donors. CD4+ T cell responses against individual peptides were measured using proliferation assays (3H-thymidine incorporation). The proliferation assay data was used to compile a T cell epitope map of the PE38 sequence and six T cell epitopes were identified.

[1088] EPISCREEN™ Donor Selection

[1089] Peripheral blood mononuclear cells (PBMC) were isolated from healthy community donor buff coats (from blood drawn within 24 hours) obtained from the UK. National Blood Transition Service (Addenbrooke's Hospital, Cambridge, UK) and according to approval granted by Addenbrooke's Hospital Local Research Ethics Committee, PBMC were isolated from buffy coats by LYMPHOPREP™ (Axis-Shield UK, Dundee, Scotland) density centrifugation. (LYMPHOPREP™ is a ready-made, sterile and endotoxin tested solution for the isolation of human mononuclear cells from blood. See, Axis-Shield, package insert for LYMPHOPREP™ density gradient media No. 619. March 3. Div.-1114740.) CD8+ T cells were depleted using CD8+ ROSETTESEPT™ (STEMCELL™, Technologies Inc, Manchester, UK) to remove CD8+ cells from the isolated mononuclear cells. See e.g., StemCell Technologies Inc., ROSETTESEPT™ procedure for Human CD8+ T Cell Enrichment Cocktail (Catalog #15023/15063: Procedure version 13.0, "#28572 (May 2011)).

[1090] HLA allotypes of donors were characterized using the Biotest HLA SSP-PCR tissue-typing kit (Biotest, Solihull, UK, catalogue number 826215). T cell responses to a reproducibility control neo-antigen were also determined using Imject mariculture keyhole limpet haemocyanin (KLH) (Pierce (Perbio Science UK, Ltd)), Cramlington, UK, catalogue number 77600) with the KLH diluted to a final concentration of 100 µg/ml. PBMC were then frozen and stored in liquid nitrogen until required.

[1091] A cohort of 52 donors was selected to best represent the number and frequency of HLA-DR allotypes expressed in the world population. Analysis of the allotypes expressed in the cohort against those expressed in the world population revealed that coverage of >80% was achieved and that all major HLA-DR alleles (individual allotypes with a frequency >5% expressed in the world population) were well represented. Details of individual donor haplotypes and a comparison of the frequency of MHC class II haplotypes expressed in the world population and the sample population are shown in Table 7 and FIG. 2, respectively.

[1092] Table 7. Donor details and haplotypes. Donor responses (SI) to KLH are shown for two independent proliferation assays. Test 1 was performed using KLH on freshly isolated PBMC and IEX01 is the KLH re-test performed in the current study on PBMC recovered from liquid nitrogen storage as indicated above. Responses that did not produce the same result (i.e. positive including borderline SI >1.90 p < 0.05 or negative) in both tests are highlighted in grey (i.e., donors 3, 7, 9, 33 and 44),

TABLE 7

Donor Haplotypes and Responses			
Donor	Haplotype	KLH	
		Test 1	IEX01
1	DRB1*15, DRB1*16; DRB5*	18.10	1.97
2	DRB1*03, DRB1*07; DRB3*; DRB4*	2.49	5.74
3	DRB1*11, DRB1*13; DRB3*; DRB4*	0.81	4.00
4	DRB1*03; DRB3*	1.73	1.78
5	DRB1*01, DRB1*13; DRB3*; DRB4*	0.99	1.05
6	DRB1*03, DRB1*14; DRB3*	2.91	2.01
7	DRB1*13, DRB1*14; DRB3*; DRB4*	3.13	1.20
8	DRB1*01, DRB1*07; DRB4*	2.49	5.74
9	DRB1*03, DRB1*07; DRB3*; DRB4*	0.81	4.00
10	DRB1*03, DRB1*15; DRB5	6.16	6.16
11	DRB1*01, DRB1*13; DRB3*	7.15	17.34
12	DRB1*13; DRB4*; DRB5*	7.98	2.76
13	DRB1*13, DRB1*14; DRB4*	1.00	1.81
14	DRB1*03, DRB1*13; DRB3*	2.28	2.70
15	DRB1*04, DRB1*11; DRB3*; DRB4*	8.96	1.91
16	DRB1*04, DRB1*14; DRB3*; DRB4*	5.85	4.01
17	DRB1*13, DRB1*15; DRB3*; DRB5*	19.16	2.69
18	DRB1*11, DRB1*13; DRB3*	10.32	6.48
19	DRB1*04, DRB1*15; DRB4*	2.70	3.39
20	DRB1*04, DRB1*07; DRB4*	0.51	1.27
21	DRB1*01, DRB1*04; DRB4*	1.71	1.05
22	DRB1*03; DRB3*	1.05	1.59
23	DRB1*04, DRB1*15; DRB4*; DRB5*	2.83	2.34
24	DRB1*01	1.63	1.09
25	DRB1*04, DRB1*15; DRB4*; DRB5*	1.12	1.44
26	DRB1*03, DRB1*07; DRB3*; DRB4*	1.18	0.84
27	DRB1*11, DRB1*13; DRB3*	8.80	14.30
28	DRB1*01, DRB1*07; DRB4*	3.68	4.53
29	DRB1*12, DRB1*13; DRB3*	3.68	2.40
30	DRB1*11; DRB3*	7.68	2.71
31	DRB1*03, DRB1*11; DRB3*	3.04	4.16
32	DRB1*13; DRB3*	1.96	2.22
33	DRB1*15; DRB4*	1.31	3.13

TABLE 7-continued

Donor Haplotypes and Responses			
34	DRB1*03, DRB1*04; DRB3*; DRB4*	0.97	1.35
35	DRB1*12; DRB3*; DRB4*	3.51	2.55
36	DRB1*07; DRB3*; DRB4*	6.63	8.90
37	DRB1*04, DRB1*04; DRB4*	44.94	6.28
38	DRB1*01, DRB1*15; DRB3*; DRB4*	1.36	1.30
39	DRB1*07, DRB1*13; DRB3*; DRB4*	12.62	2.29
40	DRB1*03, DRB1*04; DRB3*; DRB4*	1.39	1.37
41	DRB1*07, DRB1*08; DRB4*	3.40	3.47
42	DRB1*07, DRB1*13; DRB3*; DRB4*	40.32	7.36
43	DRB1*13, DRB1*15; DRB3*; DRB5*	3.56	3.21
44	DRB1*11, DRB1*14; DRB3*	1.15	2.86
45	DRB1*03, DRB1*13; DRB3*	8.78	8.72
46	DRB1*03, DRB1*13; DRB3*	11.47	3.11
47	DRB1*03, DRB1*04; DRB3*; DRB4*	6.27	2.03
48	DRB1*04, DRB1*15; DRB5*	10.29	3.77
49	DRB1*07, DRB1*15; DRB4*; DRB5*	2.59	2.32
50	DRB1*04, DRB1*15; DRB4*; DRB5*	2.49	2.42
51	DRB1*11; DRB3*; DRB4*	8.30	2.09
52	DRB1*03, DRB1*13; DRB3*	3.99	6.22

[1093] EPISCREEN™ Analysis Proliferation Assay

[1094] PBMC from each donor were thawed, counted and viability was assessed. Cells were revived in room temperature AIM V® culture medium (Invitrogen, Paisley, UK) before adjusting the cell density to $2-3 \times 10^5$ PBMC/ml (proliferation cell stock). Peptides were synthesized on a 1-3 mg scale with free N-terminal amine and C-terminal carboxylic acid. Peptides were dissolved in DMSO to a concentration of 10 mM and peptide culture stocks prepared by diluting into AIM V® culture medium to a final concentration of 5 μ M in the well. For each peptide and each donor, sextuplicate cultures were established in a flat bottomed 96 well plate. Both positive and negative control cultures were also tested in sextuplicate. For each donor, three control antigen/peptides (KLH protein and peptides derived from influenza A and Epstein Barr viruses) were also included.

[1095] Cultures were incubated for a total, of 6 days before adding 0.75 μ Ci 3[H]-thymidine (PERKIN ELMER®, Beaconsfield, UK) to each well. Cultures were incubated for a further 18 hours before harvesting onto filter mats using a TOMTEC MACH® III cell harvester (TOMTEC®, Hamden, Conn., USA). Counts per minute (cpm) for each well were determined by Meltilex™ (PERKIN ELMER®) scintillation counting on a Microplate Beta Counter (PERKIN ELMER®) in paralux, low background counting mode.

[1096] EPISCREEN™ Data Analysis

[1097] For proliferation assays, an empirical threshold of a stimulation index (SI) equal to or greater than 2 ($SI \geq 2.00$) has been previously established whereby samples inducing proliferative responses above this threshold are deemed positive (where included, borderline $SI \geq 1.90$ are highlighted). Extensive assay development and previous studies have

shown that this is the minimum signal to noise threshold allowing maximum sensitivity without detecting large numbers of false positive responses. Positive responses are defined by the following statistical and empirical thresholds:

[1098] 1. Significance ($p < 0.05$) of the response by comparing cpm of test wells against medium control wells using unpaired two sample Student's t-test.

[1099] 2. Stimulation index greater than 2.00 ($SI \geq 2.00$), where $SI = \text{mean cpm of test wells} / \text{mean cpm medium control wells}$. Data presented in this way is indicated as $SI \geq 2.00$, $p < 0.05$.

[1100] In addition, intra-assay variation was assessed by calculating the coefficient of variance and standard deviation (SD) of the raw data from replicate cultures.

[1101] Proliferation assays were set up in sextuplicate cultures ("non-adjusted data"). To ensure that intra-assay variability was low, the data was also analyzed after removing the maximum and minimum cpm values ("adjusted data") and the SI of donor responses was compared using both data sets.

[1102] T cell epitopes were identified by calculating the average frequency of positive responses (defined above) to all peptides in the study plus standard deviation (SD) to give a background response threshold. Any peptide that induced a frequency of positive proliferation responses above this threshold in both the adjusted and non-adjusted data was considered to contain an immunogenic T cell epitope (and, thus, potentially represents an immunogenicity inducing epitope which could give rise to immunogenic responses in vivo).

[1103] In Silico Analysis of Peptides

[1104] The sequences of peptides that were positive in the proliferation assay were analyzed using Antitope's predictive iTOPE™ software (Perry et al. 2008). This software predicts favorable interactions between amino acid side chains of the peptide and specific binding pockets within the MHC class II binding groove. Analysis of the peptide sequences using iTOPE™ was performed with overlapping 9 mers spanning the peptides which were tested against each of the 34 MHC class II alleles. Each 9 mer was scored based on the potential "fit" and interactions with the MHC class II molecules. 9 mers that produced a high mean binding score were identified and, from the T cell proliferation data, 9 mers which were considered as critical to T cell responses ("core 9 mers") were highlighted. iTOPE™ analysis was then repeated with a range of amino acid changes in the core 9 mers in order to determine preferred amino acid substitutions for use in deimmunization.

[1105] Results and Discussion

[1106] A total of 120 peptides were synthesized spanning the entire PE38 sequence. The peptides were designed as 15 mers to span the sequence in overlapping increments of 12 amino acids. These peptides were then tested for the presence of CD4+ T cell epitopes by EPISCREEN™ T cell epitope

mapping analysis. Positive T cell responses were defined by donors that produced a significant ($p < 0.05$) response with a $SI \geq 2.00$ to any given peptide ($SI \geq 2.00$, $p < 0.05$). T cell epitopes were identified by calculating the average frequency of the positive responses to all peptides in the study plus SD (termed 'background response threshold'). This was calculated to be 10.8% in the raw 'non-adjusted' data and 10.7% in the adjusted data (where maximum and minimum values were removed and the mean cpm calculated on the remaining four wells). Thus, peptides containing a T cell epitope induced positive T cell proliferation responses ($SI \geq 2.00$, $p < 0.05$) in ≥ 6 donors in the non-adjusted and adjusted data sets. Inter-assay variability was assessed using KLH as a reproducibility control where the frequency of positive T cell responses against KLH were compared in two separate EPISCREEN™ assays (Table 7). The results show that inter-assay variability is within the acceptable range and consistent with previous studies ($\leq 10\%$). The frequency of T cell responses against the two control peptides C3 (EBNA derived epitope) and C32 (Influenza derived epitope) ranged between 23-31% (non-adjusted) and between 21-29% (adjusted) for the two peptides, respectively (FIG. 3). This is within the typical range observed for these two peptides in T cell epitope mapping studies.

[1107] The output from non-adjusted and adjusted data analysis was examined to ensure that intra-assay variability was low and that positive responses were not the result of spurious proliferation in individual wells. The results from each analysis showed, in most cases, only small differences between the methods and donor responses for both non-adjusted and adjusted analysis. Table 10 provides a summary of individual donor responses to each of the peptides. The proliferation assay data showing the frequency of positive donor responses to each peptide is shown in FIG. 3. For all peptides that induced a high frequency of positive ($SI \geq 2.00$, $p < 0.05$, including borderline responses) T cell proliferation responses above the background response threshold, additional in silico analysis was performed to aid in the identification of the precise location of MHC class II core 9 mer binding registers (using iTOPE™), and to identify peptides that are homologous to sequences containing T cell epitopes that have been tested in previous EPISCREEN™ cell epitope mapping assays (using TCED™).

[1108] Table 8. Summary of individual donor responses to PE38 peptides. Positive responses ($SI \geq 2.00$, $p < 0.05$, including borderline responses) are indicated by the donor number and individual SI are shown in parentheses next to the corresponding donor. The background response rate was 10.8% in the non-adjusted data and 1.0.7% in the adjusted data peptides inducing positive T cell proliferation above this frequency (positive response in ≥ 6 donors) contained T cell epitopes (indicated with bold text: i.e., peptides 50, 52, 53, 65, 67, 68, 81, 82 and 110 (also as indicated in FIGS. 3-6)).

TABLE 8

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
1	11 (2.25), 19 (2.42) 25 (3.24), 35 (2.77) 36 (2.21)	11 (2.34), 19 (2.72), 25 (3.33), 35 (2.76), 36 (2.10)	GGGGSGGGGSPEG (SEQ ID NO: 11)

TABLE 8-continued

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
2	19(2.44), 36(1.94)	19(2.67), 48(2.06)	GGSGGGGSPEGGSL (SEQ ID NO: 12)
3	11(1.98), 16(1.92), 19(2.54)	11(1.99), 19(2.78), 38(5.37)	GGGGGSPEGGSLAAL (SEQ ID NO: 13)
4	16(2.33), 17(2.02), 19(2.29), 35(2.33)	16(2.13), 17(2.02), 19(2.46), 35(2.35) 38(3.71)	GSPEGGSLAALTAH (SEQ ID NO: 14)
5	7(1.97), 10(2.24), 17(2.24), 24(1.90)	7(2.05), 10(2.46), 17(2.20)	PEGGSLAALTAHQAC (SEQ ID NO: 15)
6		3(2.00)	GSLAALTAHQACHLP (SEQ ID NO: 16)
7	3(2.11), 17(2.39), 24(2.20), 45(2.11)	3(2.30), 17(2.22), 24(2.11), 45(1.90)	AALTAHQACHLPLET (SEQ ID NO: 17)
8	17(2.23), 45(2.12)	17(2.04)	TAHQACHLPLETFTR (SEQ ID NO: 18)
9	—	—	QACHLPLETFTRHRQ (SEQ ID NO: 19)
10	—	—	HLPLETFTRHRQPRG (SEQ ID NO: 20)
11	10(2.48), 24(2.13)	7(1.98), 10(2.90), 24(2.12)	LETFTTRHRQPRGWEQ (SEQ ID NO: 21)
12	—	7(1.95)	FTRHRQPRGWEQLEQ (SEQ ID NO: 22)
13	10(3.20), 24(2.12)	10(7.11), 24(2.12)	HRQPRGWEQLEQCGY (SEQ ID NO: 23)
14	10(2.14), 17(2.47), 24(2.06)	1(2.22), 10(2.04), 17(2.25), 24(2.17), 51(1.91)	PRGWEQLEQCGYPVQ (SEQ ID NO: 24)
15	17(2.04), 24(2.24)	17(2.05), 24(2.21)	WEQLEQCGYPVQRLV (SEQ ID NO: 25)
16	—	—	LEQCGYPVQRLVALY (SEQ ID NO: 26)
17	—	—	CGYPVQRLVALYLAA (SEQ ID NO: 27)
18	—	—	PVQRLVALYLAARLS (SEQ ID NO: 28)
19	—	—	RLVALYLAARLSWNQ (SEQ ID NO: 29)
20	3(2.35), 40(1.96)	3(9.55), 8(2.06), 9(1.90), 19(1.94), 40(2.07)	ALYLAARLSWNQVDQ (SEQ ID NO: 30)
21	11(1.90)	3(4.22), 8(2.22), 11(2.17), 40(1.96)	LAARLSWNQVDQVIR (SEQ ID NO: 31)
22	3(2.20), 40(2.10)	3(4.32), 6(2.17), 19(2.00), 40(2.15)	RLSWNQVDQVIRNAL (SEQ ID NO: 32)
23	6(2.24), 19(1.97)	34(4.67), 6(2.40), 19(2.22),	WNQVDQVIRNALASP (SEQ ID NO: 33)
24	3(3.08)	3(3.99)	VDQVIRNALASPGSG (SEQ ID NO: 34)

TABLE 8-continued

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
25	3(1.90), 6(1.92)	6(2.36)	VIRNALASPGSGGDL (SEQ ID NO: 35)
26	—	—	NALASPGSGGDLGEA (SEQ ID NO: 36)
27	—	—	ASPGSGGDLGEAIRE (SEQ ID NO: 37)
28	24(2.62)	9(1.99), 11(1.92), 24(3.47),	GSGGDLGEAIREQPE (SEQ ID NO: 38)
29	4(2.05), 17(2.07), 45(2.18)	4(1.91), 17(1.96), 24(2.29), 45(2.18)	GDLGEAIREQPEQAR (SEQ ID NO: 39)
30	17(1.94)	8(2.54), 17(2.18), 24(2.44)	GEAIREQPEQARLAL (SEQ ID NO: 40)
31	—	—	IREQPEQARLALTLA (SEQ ID NO: 41)
32	—	31(1.96)	QPEQARLALTAAAE (SEQ ID NO: 42)
33	—	—	QARLALTAAAESER (SEQ ID NO: 43)
34	—	—	LALTAAAESERFVR (SEQ ID NO: 44)
35	4(1.91), 35(1.91), 37(2.18), 42(2.05)	29(2.42), 35(1.98), 36(2.05), 37(2.27), 42(2.07)	TLAAAESERFVRQGT (SEQ ID NO: 45)
36	3(2.14), 42(1.91)	3(2.12), 29(3.71), 36(1.91)	AAESERFVRQGTGND (SEQ ID NO: 46)
37	37(2.34), 42(1.96)	13(1.91), 29(3.91), 35(2.06), 37(2.20)	SERFVRQGTGNDEAG (SEQ ID NO: 47)
38	37(2.20), 42(2.12)	29(2.28), 36(1.95), 37(2.20), 42(2.06)	FVPQGTGNDEAGAAS (SEQ ID NO: 48)
39	42(1.90)	—	QGTGNDEAGAASGPA (SEQ ID NO: 49)
40	1(2.21)	1(2.08)	GNDEAGAASGPADSG (SEQ ID NO: 50)
41	1(2.28)	1(2.16)	EAGAASGPADSGDAL (SEQ ID NO: 51)
42	—	—	AASGPADSGDALLER (SEQ ID NO: 52)
43	—	—	GPADSGDALLERNYP (SEQ ID NO: 53)
44	17(2.08), 22(1.95), 42(2.02)	17(2.13), 22(2.00), 37(1.98)	DSGDALLERNYPTGA (SEQ ID NO: 54)
45	—	—	DALLERNYPTGAEFL (SEQ ID NO: 55)
46	31(2.23)	31(1.98)	LERNYPTGAEFLGDG (SEQ ID NO: 56)
47	—	—	NYPTGAEFLGDGGDI (SEQ ID NO: 57)
48	2(2.63)	—	TGAEFLGDGGDISFS (SEQ ID NO: 58)

TABLE 8-continued

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
49	—	—	EFLGDGGDISFSTRG (SEQ ID NO: 59)
50	10(2.54), 11(1.93), 19(2.29), 36(2.36), 37(1.92), 39(2.17), 42(2.66), 45(1.96)	10(2.56), 11(2.33), 19(2.37), 36(2.37), 39(2.13), 42(2.63), 45(1.95), 46(1.93)	GDGGDISFSTRGTQN (SEQ ID NO: 60)
51	19(2.03), 42(2.25), 45(1.93)	2(2.57), 11(2.06), 19(1.97), 42(2.20), 45(1.90)	GDISFSTRGTQNWTV (SEQ ID NO: 61)
52	3(7.10), 11(2.76), 16(2.41), 19(2.36), 42(1.97), 44(1.92)	2(1.95), 3(6.19), 11(3.01), 16(2.58), 19(2.45), 42(2.02), 44(2.05)	SFSTRGTQNWTVRL (SEQ ID NO: 62)
53	2(2.13), 3(5.19), 11(1.98), 16(2.12), 19(2.19), 27(2.09), 45(1.92)	2(2.27), 3(4.50), 11(2.01), 16(1.94), 19(2.10), 27(2.46)	TRGTQNWTVRLLQA (SEQ ID NO: 63)
54	—	3(1.90), 11(1.95), 16(1.92)	TQNWTVRLLQAHRQ (SEQ ID NO: 64)
55	3(1.98)	—	WTVRLLQAHRQLEE (SEQ ID NO: 65)
56	—	—	ERLLQAHRQLEERGY (SEQ ID NO: 66)
57	—	—	LQAHRQLEERGYVFPV (SEQ ID NO: 67)
58	10(2.67), 11(2.90)	9(1.99), 10(2.55), 11(3.46), 4(1.90)	HRQLEERGYVFGYH (SEQ ID NO: 68)
59	9(2.27), 37(2.56), 42(2.70)	9(2.38), 11(2.15), 37(2.71), 42(3.01)	LEERGYVFGYHGTGTF (SEQ ID NO: 69)
60	—	16(2.09)	RGYVFGYHGTGFLEA (SEQ ID NO: 70)
61	—	11(2.07)	VFGYHGTGFLEAAQS (SEQ ID NO: 71)
62	—	—	GYHGTGFLEAAQSIVF (SEQ ID NO: 72)
63	3(2.88)	11(1.97), 16(2.02)	GTFLEAAQSIVFGGV (SEQ ID NO: 73)
64	—	—	LEAAQSIVFGGVRAR (SEQ ID NO: 74)
65	2(2.17), 4(1.94), 14(3.63), 17(2.19), 18(2.46), 36(2.06), 39(1.97), 51(7.91)	2(2.30), 4(1.93), 11(2.10), 14(3.65), 18(2.31), 36(2.09), 39(2.04), 51(6.71)	AQSIVFGGVRARSQD (SEQ ID NO: 75)
66	18(1.95), 19(1.90), 36(1.94), 51(10.38)	19(2.02), 36(1.98), 47(1.91), 51(9.41)	IVFGGVRARSQDLDA (SEQ ID NO: 76)
67	6(2.07), 14(2.62), 16(2.21), 17(2.11), 18(2.60), 42(1.95), 47(1.93), 51(6.83)	14(2.68), 16(2.55), 18(2.42), 19(2.06), 38(1.95), 47(1.95), 51(5.22)	GGVRARSQDLDAIWR (SEQ ID NO: 77)
68	2(2.07), 14(2.24), 18(2.70), 38(1.94), 39(2.05), 42(2.10), 51(3.69)	2(2.12), 11(2.11), 14(2.04), 16(2.00), 19(2.06), 38(2.15), 39(2.17), 51(3.62)	RARSQDLDAIWRGFY (SEQ ID NO: 78)

TABLE 8-continued

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
69	31(1.95), 42(1.99), 51(2.47)	31(1.93), 51(2.19)	SQDLDAIWRGFYIAG (SEQ ID NO: 79)
70		24(2.22)	LDAIWRGFYIAGDPA (SEQ ID NO: 80)
71		—	IWRGFYIAGDPALAY (SEQ ID NO: 81)
72	—	—	GFYIAGDPALAYGYA (SEQ ID NO: 82)
73	6(1.91), 14(2.70), 17(2.13), 39(1.98)	11(2.02), 14(2.77), 17(1.94), 39(2.01)	IAGDPALAYGYAQDQ (SEQ ID NO: 83)
74	6(1.99), 14(2.77), 38(1.99), 39(2.25), 42(1.90)	14(2.89), 16(2.01), 39(2.27)	DPALAYGYAQDQEEPD (SEQ ID NO: 84)
75	6(2.22), 14(2.27), 17(1.93), 39(2.05)	14(2.24), 16(2.26), 39(2.07)	LAYGYAQDQEPDARG (SEQ ID NO: 85)
76	14(2.20), 17(1.98)	14(2.40), 39(1.94)	GYAQDQEPDARGRIR (SEQ ID NO: 86)
77	—	38(1.90)	QDQEPDARGRIRNGA (SEQ ID NO: 87)
78	—	—	EPDARGRIRNGALLR (SEQ ID NO: 88)
79	—	24(1.91)	ARGRIRNGALLRVYV (SEQ ID NO: 89)
80	9(2.89), 11(2.85), 19(2.18), 36(2.63), 42(2.23), 45(2.23)	9(2.84), 19(2.03), 36(2.59), 42(2.29), 45(2.20)	RIRNGALLRVYVPRS (SEQ ID NO: 90)
81	1(2.08), 8(1.96), 9(2.05), 11(3.22), 13(2.09), 19(2.09), 36(2.21), 42(2.31), 45(2.13), 49(2.07)	1(2.13), 9(2.05), 13(2.10), 19(2.10), 36(2.15), 42(2.31), 45(1.94), 49(2.00), 51(2.25)	NGALLRVYVPRSSLP (SEQ ID NO: 91)
82	9(1.93), 10(2.01), 11(2.41), 13(2.09), 16(2.11), 19(2.01), 36(2.25), 45(1.97), 49(2.44)	9(1.90), 10(2.00), 13(2.21), 16(2.23), 19(1.98), 36(2.28), 45(1.98), 49(2.37)	LLRVYVPRSSLPGFY (SEQ ID NO: 92)
83	33(2.02), 42(2.14), 46(1.90), 49(1.92)	33(1.97), 42(2.14), 49(1.95)	VKYVPRSSLPGFYRTG (SEQ ID NO: 93)
84	11(1.93)	—	PRSSLPGFYRTGLTL (SEQ ID NO: 94)
85	—	—	SLPGFYRTGLTLAAP (SEQ ID NO: 95)
86	—	—	GFYRTGLTLAAPEAA (SEQ ID NO: 96)
87	—	—	RTGLTLAAPEAAGEV (SEQ ID NO: 97)
88	9(2.59), 11(3.03), 42(2.03), 51(2.30)	9(2.47), 42(2.01)	LTAAPEAAGEVERL (SEQ ID NO: 98)
89	9(1.91), 11(4.31), 42(2.34), 49(2.09), 51(5.22)	11(2.05), 13(2.28), 42(2.16), 51(6.48)	AAPEAAGEVERLIGH (SEQ ID NO: 99)
90	11(2.59), 14(2.07), 49(2.12), 51(7.78)	14(2.11), 49(2.11), 51(6.45)	EAAGEVERLIGHPLP (SEQ ID NO: 100)

TABLE 8-continued

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
91	11(4.15), 42(1.99), 51(4.84),	42(2.06), 51(4.07)	GEVERLIGHPLPLRL (SEQ ID NO: 101)
92	11(2.19), 49(1.99)	—	ERLIGHPLPLRLDAI (SEQ ID NO: 102)
93	—	—	IGHPLPLRLDAITGP (SEQ ID NO: 103)
94	—	—	PLPLRLDAITGPTEE (SEQ ID NO: 104)
95	3(2.10), 7(1.91), 18(1.90), 19(2.07), 35(2.17)	3(2.00), 7(1.95), 19(2.04), 45(2.03)	LRLDAITGPTEEGGR (SEQ ID NO: 105)
96	3(2.62), 13(2.19), 16(2.18), 39(1.96)	3(2.34), 13(2.46), 16(2.24), 31(1.93), 39(2.10)	DAITGPTEEGGRLET (SEQ ID NO: 106)
97	13(2.29), 16(2.32), 19(2.20), 35(2.43), 45(2.13)	13(2.48), 16(2.44), 19(2.31), 45(2.16)	TGPTEEGGRLETILG (SEQ ID NO: 107)
98	11(1.92), 13(1.97), 16(2.26), 35(1.91), 50(1.98)	11(2.26), 13(2.04), 16(2.33)	EEEGGRLETILGWPL (SEQ ID NO: 108)
99	35(2.33)	—	GGRLETILGWPLAER (SEQ ID NO: 109)
100	35(2.20)	—	LETILGWPLAERTVV (SEQ ID NO: 110)
101	—	—	ILGWPLAERTVVIPS (SEQ ID NO: 111)
102	27(1.93)	—	WPLAERTVVIPSAIP (SEQ ID NO: 112)
103			AERTVVIPSAIPTDP (SEQ ID NO: 113)
104	3(2.40), 16(2.20), 22(1.98), 49(1.91)	3(2.17), 13(2.05), 16(2.15)	TVVIPSAIPTDPRNV (SEQ ID NO: 114)
105	16(2.43), 22(1.96), 45(1.97)	16(2.30), 45(1.95), 49(1.96)	IPSAIPTDPRNVGGD (SEQ ID NO: 115)
106	16(2.02), 19(2.02)	16(1.95), 19(1.90)	AIPTDPRNVGGDLDP (SEQ ID NO: 116)
107	19(2.00), 27(2.06)	19(1.93), 27(1.99)	TDPRNVGGDLDPSSI (SEQ ID NO: 117)
108	—	—	RNVGGDLDPSSIPDK (SEQ ID NO: 118)
109	—	—	GGDLDPSSIPDKEQA (SEQ ID NO: 119)
110	8(2.07), 9(2.35), 11(2.27), 13(2.13), 16(1.91), 19(3.00), 35(1.90)	9(2.46), 10(2.04), 13(2.11), 19(1.99), 35(1.94), 38(1.95), 50(2.01)	LDPSSIPDKEQAISA (SEQ ID NO: 120)
111	3(2.29), 8(2.20), 9(1.93), 11(2.08), 16(2.19), 19(2.60)	3(2.33), 8(2.74), 9(2.02), 13(1.98)	SSIPDKEQAISALPD (SEQ ID NO: 121)
112	11(2.47), 16(3.07), 19(2.61)	9(1.90), 16(2.11), 50(1.97)	PDKEQAISALPDYAS (SEQ ID NO: 122)
113	3(2.07), 11(2.61), 16(2.44), 19(2.62)	3(2.04), 11(1.93), 45(1.90)	EQAISALPDYASQPG (SEQ ID NO: 123)

TABLE 8-continued

Donor Responses to PE38 Peptides			
Peptide #	Proliferation Non-Adjusted	Proliferation Adjusted	Peptide Sequence
114	19 (2.04)	—	ISALPDYASQPGKPP (SEQ ID NO: 124)
115	16 (1.99)	—	LPDYASQPGKPPRED (SEQ ID NO: 125)
116	—	—	YASQPGKPPREDLK (SEQ ID NO: 126)
117	—	—	ITGPEEEGGRDITIL (SEQ ID NO: 127)
118	9 (2.04), 11 (2.26), 16 (2.12), 39 (1.93), 5	9 (2.27)	PEEEGGRDITILGWP (SEQ ID NO: 128)
119	16 (2.11), 39 (2.13)	14 (1.90), 38 (1.96), 39 (2.10)	EGGRDITILGWPLAE (SEQ ID NO: 129)
120	11 (2.13), 39 (2.05)	39 (2.07)	RLDITILGWPLAERTV (SEQ ID NO: 130)

[1109] T Cell Epitope Map**[1110]** Epitopes 1 and 2—

[1111] Peptides 50, 52 and 53 induced a high number of positive T cell proliferation responses in the study cohort (Table 8 and FIG. 3). Peptide 50 showed the highest number of positive responses with 15.38% donors responding in the non-adjusted dataset, and 15.38% in the adjusted data set, ($SI \geq 2.00$, $p < 0.05$). From in silico analysis, the proposed core 9 mer in this region is ISFSTRGTQ (SEQ ID NO:5). Peptides 52 and 53 induced lower frequencies of response with 11.54% and 13.46% positive donor responses in the non-adjusted dataset, and 13.46% and 11.54% in the adjusted datasets, respectively. A core 9 mer was identified in peptide 50 but was only partially present in peptides 52 and 53 suggesting that these peptides must contain a different T cell epitope. In silico analysis of peptides 52 and 53 did not identify any core HLA-DR restricted 9 mers so it is likely that the positive T cell responses seen are due to a HLA-DQ restricted T cell epitope.

[1112] The magnitude of T cell proliferation responses can provide an indication as to the T cell precursor frequency. In general, peptides that induce high frequency (of positive responses in the study cohort) and high magnitude T cell proliferation responses are a characteristic of ‘recall-like’ T cell responses in which the T cell pre-cursor frequency is high. In contrast, naive T cell responses are generally characterized by low magnitude T cell proliferation responses (with low T cell precursor frequencies). Peptides 52 and 53 induced moderately high magnitude T cell proliferation responses where the mean SI for positive ($SI \geq 2.00$, $p < 0.05$) T cell responses in the non-adjusted and adjusted data sets were 3.09-2.89 (peptide 52) and 2.51-2.55 (peptide 53) (Table 9). Thus these peptides may induce T cell responses in clones that are present in high frequencies in healthy individuals and may be indicative of a memory T cell response. Peptide 50 induced lower magnitude T cell proliferation responses where the mean SI were 2.23 and 2.28 in the non-adjusted and adjusted datasets respectively suggesting that this peptide

may induce a naïve T cell response (Table 9).

[1113] Epitopes 3 and 4

[1114] A cluster of T cell responses were observed around peptides 65-68 and the subsequent analysis revealed the presence of two T cell epitopes in this region. Peptide 65 stimulated positive T cell proliferation responses in 15.38% of the study cohort for both non-adjusted and adjusted datasets (Table 8 and FIG. 3) ($SI \geq 2.00$, $p < 0.05$). The positive responses were high magnitude (mean SI of positive responses ranged from 3.04-2.89 in the non-adjusted and adjusted data sets) suggesting that the T cell precursor frequency in healthy donors against this epitope is high (Table 9). In silico analysis revealed a potential core 9 mer comprising IVFGGVRAR (FIG. 5: SEQ ID NO:7). Peptides 67 and 68 induced frequencies of response with 15.38% and 13.46% positive donor responses in the non-adjusted dataset, and 13.46% and 15.38% in the adjusted datasets, respectively, in silico analysis of these peptides did not identify any core HLA-DR 9 mers so it is likely that the positive T cell responses seen are due to a HLA-DQ restricted T cell epitope.

[1115] Epitope 5

[1116] Peptides 81 and 82 stimulated a number of T cell responses in the study cohort (Table 8 and FIG. 3). Peptide 81 had the highest frequency of response of all the peptides tested with a frequency of positive responses of 19.23% in the non-adjusted and 17.31% in the adjusted data set. For peptide 82, the frequency of positive response was 17.31% and 15.38% in the non-adjusted and adjusted data sets respectively. The positive responses were relatively low in magnitude (mean SI of positive responses ranged from 2.12 to 2.22 in the non-adjusted and adjusted data sets) suggesting that the T cell precursor frequency in healthy donors against this epitope is relatively low (Table 9). Adjacent peptide 80 induced a sub-threshold response. In silico analysis of peptides 81 and 82 suggested a core 9 mer of LRVYVPRSS (FIG. 6; SEQ ID NO:9).

[1117] Epitope 6

[1118] Peptide 110 induced positive T cell responses in 13.46% of the study cohort in non-adjusted and 13.46% in adjusted datasets (Table 8 and FIG. 3). The magnitude of positive proliferation responses was low with a mean SI of 2.23 for the non-adjusted dataset and 2.07 for the adjusted dataset (Table 9). There was also a sub-threshold response to peptide 111. In silico analysis of the peptides sequence revealed a core 9 mer, IPDKEQAIS (FIG. 7; SEQ ID NO:10) which, in addition to peptide 110, was also present in peptide 111.

[1119] Table 9. Summary of magnitude (mean SI and standard deviation) and frequency (% donor response) of positive T cell proliferation responses against peptides containing T cell epitopes for PE38. The position of p1 in potential core 9 mers are shown as underlined/bolded text (as predicted by iTOPE™) in peptides 50, 65, 81, 82 and 110.

rate in the adjusted data in the study cohort and was also restricted to allotypes expressed at higher frequencies (>5%) in the study population.

[1122] Analysis of responding donor allotypes (Table 10 and FIG. 8) revealed that there was a possible association between T cell responses to peptides 81, and 82 and MHC class II allotype HLA DRB1*07 which was expressed at twice the percentage of positively responding donors compared to the study population. Peptide 53 also had a possible association with DRB1*11, and peptides 82 and 110 showed possible associations with DRB1*15. It should be noted that further studies (such as MHC class II binding analysis) would be required to show conclusively that responses to the T cell epitope are associated with these allotypes as the present analysis was performed on a small group of responding donors.

TABLE 9

Magnitude and Frequency of Donor Responses						
		Response Frequency		Mean ($\pm$ SD)	Mean ($\pm$ SD)	
Peptide	Peptide Sequence	Non-Adjusted	Adjusted	Non-Adjusted Data	Adjusted Data	
50	GDGGD <u>IS</u> FSFSTRGTQN (SEQ ID NO: 60)	15.38%	15.38%	2.23 $\pm$ 0.28	2.28 $\pm$ 0.26	
52	SFSTRGTQNWTVERL (SEQ ID NO: 62)	11.54%	13.46%	3.09 $\pm$ 1.99	2.89 $\pm$ 1.50	
53	TRGTQNWTVERLLQA (SEQ ID NO: 63)	13.46%	11.54%	2.51 $\pm$ 1.18	2.55 $\pm$ 0.98	
65	AQSI <u>IV</u> FGGVARSQD (SEQ ID NO: 75)	15.38%	15.38%	3.04 $\pm$ 2.04	2.89 $\pm$ 1.64	
67	GGVARSQDLDAIWR (SEQ ID NO: 77)	15.38%	13.46%	2.79 $\pm$ 1.65	2.69 $\pm$ 1.15	
68	RARSQDLDAIWRGFY (SEQ ID NO: 78)	13.46%	15.38%	2.40 $\pm$ 0.62	2.28 $\pm$ 0.54	
81	NGALL <u>LR</u> VYVPRSSLP (SEQ ID NO: 91)	19.23%	17.31%	2.22 $\pm$ 0.36	2.12 $\pm$ 0.12	
82	<u>LL</u> RVYVPRSSLPGFY (SEQ ID NO: 92)	17.31%	15.38%	2.14 $\pm$ 0.19	2.12 $\pm$ 0.17	
110	LDPSS <u>IP</u> DKEQAISA (SEQ ID NO: 120)	13.46%	13.46%	2.23 $\pm$ 0.38	2.07 $\pm$ 0.18	

[1120] HLA Analysis

[1121] Analysis of the responding donor haplotypes was performed whereby an association between MHC class II allotype and a response to a particular peptide was considered possible if the frequency of the allotype within the responding population was double the frequency observed in the study cohort. This analysis was only carried out for peptides that induced positive responses above the background response

[1123] Table 10, Frequency (expressed as a percentage) of responding donor allotypes compared to the frequency of allotypes expressed in the IEX01 study cohort. An association between MHC class II allotype and a response to a particular epitope was considered if the frequency of the allotype within the responding population was double the frequency observed in the study population in the adjusted data set. Possible associations are indicated in heavily bordered boxes. The analysis has been restricted to allotypes expressed at higher frequencies (>5%) in the study population.

TABLE 10

Frequency of responding donor allotypes versus frequency of allotypes in the IEX01 study cohort.								
Frequency (%) of HLA alleles expressed within:	DRB1*03	DRB1*04	DRB1*07	DRB1*11	DRB1*15	DRB3	DRB4	DRB5
Study population	8	8	7	5	7	20	18	6
Peptide 50	8	8	12	0	8	23	15	4
Peptide 52	4	8	8	8	4	24	20	0
Peptide 53	5	10	5	10	5	24	19	0
Peptide 65	12	0	12	8	0	32	16	0
Peptide 67	8	13	0	8	8	25	21	0
Peptide 68	7	7	7	4	7	25	21	0
Peptide 81	7	0	14	7	7	21	21	7
Peptide 82	8	8	15	0	15	15	23	8
Peptide 110	4	13	4	0	17	13	25	8

[1124] Results

[1125] The results show that six T cell epitopes were present in the PE38 sequence. Table 6 Table 11 and FIG. 8 summarize the location of the putative core 9 mers in each sequence along with the frequency and magnitude of T cell responses against each epitope. The T cell epitopes identified in PE38 were prioritized according to their potency based on the frequency and magnitude (mean SI) of positive donor responses to each peptide. However since the responding donor magnitudes were similar (Table 4 Table 9) for most epitopes, the ranking was mainly based on frequency of positive donor responses (from highest to lowest):

[1126] Epitope 5>Epitope 4>Epitope 3>Epitope 1>Epitope 2>Epitope 6

[1127] Deimmunization Strategy

[1128] The six epitope core 9 mer sequences were analyzed by proprietary software (iTOPE™) in order to identify mutations that remove the T cell epitopes by eliminating or significantly reducing binding to MHC class II (Table 11). As part of the strategy as to which residues to mutate, location within the structure was considered, especially whether the residue is buried, on the surface, or near active sites.

[1129] Table 11. Projected mutations to remove MHC class II binding (based upon iTOPE™ and crystal structure data).

TABLE 11-continued

Location of Core 9-mers and Projected Mutations				
Epi-tope	Amino Acids in Sequence	Anchor Residues	Projected Mutations	Notes:
2	G	1		HLA-DQ epitopes have a strong negative preference for positively charged residues in key anchor positions. All four mutations are equally preferred.
	T			
	Q			
	N	4	K, R	
	W	6	K, R	
	V	7		
3	E			P1 is buried, therefore A is preferred. P6 V is partially exposed. All mutations should be tolerated. Preference is D > M > N.
	R	9		
	I	1	A, N	
	V			
	F			
	G	4		
4	G			HLA-DQ epitopes have a strong negative preference for positively charged residues in key anchor positions. All four mutations are equally preferred.
	V	6	D, M, N	
	R	7		
	A			
	R	9		
	A	1		
5	R			P1 is buried and close in the structure to epitope 3 P1, therefore changes are limited. For this epitope, changes at P2 affect binding (D > S > A). P9 is mostly surface exposed. Preferred changes are D, E, N, then K > P > T.
	S			
	Q	4	K, R	
	D			
	L	6		
	D	7	K, R	
6	A			P1 I is partially surface exposed, therefore all alternatives should be possible. P4, P6 and P9 changes are less preferred than P1
	I	9		
	L	1	A	
	R		D, S, A	
	V			
	Y	4		
	V			P1 I is partially surface exposed, therefore all alternatives should be possible.
	P	6		
	R	7		
	S			
	S	9	D, E, N, K, P, T	
	I	1	A, N, T, Q, H	
	P			P4, P6 and P9 changes are less preferred than P1
	D			
	K	4	T	
	E			

TABLE 11

Location of Core 9-mers and Projected Mutations				
Epi-tope	Amino Acids in Sequence	Anchor Residues	Projected Mutations	Notes:
1	I	1	A, N, T, Q, H	P1 (Ile) is partially surface exposed, therefore all alternatives should be possible. P6 and P9 changes perform equally well, but are less preferred than P1 changes.
	S			
	F			
	S	4		
	T			
	R	6	Q	
	G	7		
	T			
	Q	9	N, T	

TABLE 11-continued

Location of Core 9-mers and Projected Mutations				
Epi- tope	Amino Acids in Sequence	Anchor Residues	Projected Mutations	Notes:
	Q	6	D	changes. P6 D $\geq$ P7 D > P4 T.
	A	7	D	
	I			
	S	9		

[1130] Conclusions

[1131] EPISCREEN™ T cell epitope mapping of 120 overlapping 15 mer peptides including 112 spanning the entire PE38 sequence suggested six novel T cell epitopes. In silico analysis was used to identify potential core 9 mers for MHC binding and, together with structural analysis, was used as a basis for design of changes for re-engineering and deimmunizing PE38 in particular, and PE molecules in general.

Example 2

T Cell Epitope Mapping of Deimmunized/Amino Acid Substituted Forms of PE

[1132] The immunogenicity of amino acid substituted forms of PE can be assessed using the same procedures as described in Example 1. Accordingly, EPISCREEN™ T cell epitope mapping analysis (Antitope Ltd, Cambridge, UK) analysis permits identification of amino acid substituted epitopes in PE polypeptides, wherein the introduced amino acid changes result in reduced or undetectable immunogenicity (i.e., for generating deimmunized forms of PE) as compared to epitopes in corresponding forms of non-amino acid substituted PE polypeptides.

[1133] EPISCREEN™ is a proprietary technology commercially available through Antitope Ltd, Cambridge, UK, to map T cell epitopes within a protein sequence to determine potential for immunogenicity (based on the number and potency of T cell epitopes within a sequence). EPISCREEN™ T cell epitope mapping typically uses CD8+ T cell depleted PBMCs from a minimum of 50 HLA-typed donors (selected to represent the human population of interest). Typically, 15 mer peptides with 12 amino acid overlaps spanning a protein sequence are analyzed in a large number of replicate cultures for in vitro CD4+ T cell stimulation by 3H TdR incorporation. CD4+ T cell stimulation is often detected in two or three adjacent and overlapping peptides since the core 9 mer that binds the MHC class II binding groove will be present in more than one peptide sequence. Following identification of peptides that stimulate CD4+ T cells in vitro, in silico technology can be used to design epitope-depleted (deimmunized) variants by determining the precise location of core 9 mer sequences and the location of key MHC class II anchor residues.

[1134] In this case, amino acid substituted PE peptides are analyzed for the presence of immunogenic CD4+ T cell epitopes using EPISCREEN™ T cell epitope mapping analysis. For example, amino acid substituted 15 mer peptides (compared to non-substituted 15 mer peptides corresponding to a non-amino acid substituted form of PE) are tested against a cohort of healthy donors. CD4+ T cell responses against individual peptides are measured using proliferation assays (3H-thymidine incorporation). Proliferation assay data is used to compile a T cell epitope map of varying responses to

amino acid substituted forms of PE to determine those amino acid changes producing reduced or abrogated immunogenic responses.

[1135] EPISCREEN™ Donor Assessments

[1136] Peripheral blood mononuclear cells (PBMC) are isolated from healthy donor buffy coats (e.g., from blood drawn within 24 hours). For example, PBMC are isolated from buffy coats using density gradient centrifugation using LYMPHOPREP™ (Axis-Shield UK, Dundee, Scotland) or a similar density gradient centrifugation media for the isolation of human mononuclear cells from blood (such methods, media and products are well known and routinely used by those skilled in the art). See e.g., Axis-Shield, package insert for LYMPHOPREP™ density gradient media No. 619. March 3, Div.-1114740.) To remove CD8+ cells from the isolated mononuclear cells, CD8+ T cells are depleted using CD8+ ROSETTESEP™ kit (STEMCELL™ Technologies Inc, Manchester, UK.) or similar CD8+ selection methods and techniques (such methods, media and products are well known and routinely used by those skilled in the art). See e.g., StemCell Technologies Inc., ROSETTESEP™ procedure for Human CD8+ T Cell Enrichment Cocktail (Catalog #15023/15063 Procedure version 1.3.0, “#28572 (May 2011)).

[1137] Donors HLA-DR haplotypes are determined using methods or kits well-known and routinely used by those skilled in the art. For example, Donors HLA-DR haplotypes are determined using a Biotest HLA SSP-PCR tissue-typing kit (Biotest, Solihull, UK, catalogue number 826215). T cell responses to a reproducibility control antigen are measured using, for example neo-antigen, using Inject mariculture keyhole limpet haemocyanin (KLH) (Pierce (Perbio Science UK, Ltd), Cramlington, UK, catalogue number 77600), or other similar control antigen (such antigens and methods are well known and routinely used by those skilled in the art). PBMC are frozen and stored in liquid nitrogen until ready for use in to measuring immunogenicity of amino acid substituted forms of PE.

[1138] A cohort of donors are selected to best represent the number and frequency of HLA-DR allotypes expressed in the world population. It is desirable that allotypes expressed in the cohort represent a coverage of >80% of all major HLA-DR alleles in the world population (i.e., individual allotypes with a frequency >5% expressed in the world population are well represented). Records of individual donor haplotypes and comparison of the frequency of MHC class II haplotypes expressed in the world population and the sample population are recorded and assessed.

[1139] Donor responses (SI) to a control antigen (such as KLH) are assessed by comparing two independent proliferation assays. Test-1 is performed using the control antigen (such as KLH) on freshly isolated PBMC and Test-2 is the control antigen re-test performed on PBMC recovered from liquid nitrogen storage, the latter of which are used in assessing immunogenicity of amino acid substituted epitopes in PE. Responses that do not produce the same result in these two tests (i.e. positive including borderline SI >1.90 p < 0.05 or negative) in both tests are disregarded.

[1140] EPISCREEN™ Analysis: Proliferation Assay

[1141] PBMC from each donor are thawed, counted and viability is assessed. Cells are revived in room temperature AIM V® Culture Medium (INVITROGEN™, Paisley, UK) before adjusting cell density to 2-3×10⁶ PBMC/ml (proliferation cell stock). Peptides are synthesized on a 1-3 mg scale with free N-terminal amine and C-terminal carboxylic acid.

Peptides are dissolved in DMSO to a concentration of 10 mM and peptide culture stocks are prepared by diluting into AIM V® Culture Medium to a final concentration of 5 µM per well. For each peptide and each donor, sextuplicate cultures are established in a flat bottomed 96 well plate. Both positive and negative control cultures are tested in sextuplicate. For each donor, three control antigen/peptides (KLH protein and peptides derived from Influenza A and Epstein Barr viruses) are also included.

[1142] Cultures are incubated for 6 days before adding 0.75 µCi 3[H]-thymidine (PERKINELMER®, Beaconsfield, UK) to each well. Cultures are incubated a further 18 hours before harvesting onto filter mats using a TOMTEC MACH® III cell harvester (TOMTEC®, Hamden, CT, USA). Counts per minute (cpm) for each well are determined by MELTILEX™ (PERKIN ELMER®) scintillation counting on a Microplate Beta Counter (PERKIN ELMER®) in paralux, low background counting mode.

[1143] EPISCREEN™ Data Analysis

[1144] In proliferation assays, an empirical threshold of stimulation index (SI) equal to or greater than 2 ($SI \geq 2.00$) is considered to represent an induced proliferative response; samples registering values above this threshold are deemed positive (values of $SI < 2.00$ but ≥ 1.90 are considered borderline). Extensive assay development and previous studies have shown that this is the minimum signal to noise threshold allowing maximum sensitivity without detecting large numbers of false positive responses. Positive responses are defined by the following statistical and empirical thresholds:

[1145] 1. Significance ($p < 0.05$) of the response by comparing cpm of test wells against medium control wells using unpaired two sample Student's t-test.

[1146] 2. Stimulation index greater than 2.00 ($SI \geq 2.00$), where $SI = \text{mean cpm of test wells} / \text{mean cpm medium control wells}$. Thus, data presented is indicated as $SI \geq 2.00$, $p < 0.05$.

[1147] In addition, intra-assay variation is assessed by calculating the coefficient of variance and standard deviation (SD) of raw data from replicate cultures.

[1148] Proliferation assays are set up in sextuplicate cultures from which "non-adjusted data" is gathered. To ensure intra-assay variability is low, data is also analyzed after removing maximum and minimum cpm values (to produce "adjusted data") and the SI of donor responses is compared using both data sets.

[1149] Reactive T cell epitopes are identified by calculating the average frequency of positive responses (defined above) to all peptides in the study plus standard deviation (SD) to give a background response threshold. Any peptide inducing a frequency of positive proliferation responses above the threshold in both adjusted and non-adjusted data is considered to contain an immunogenic T cell epitope (and, thus, potentially represents an immunogenicity inducing epitope which could give rise to immunogenic responses in vivo). Output from non-adjusted and adjusted data is examined to ensure that intra-assay variability is low and that positive responses are not the result of spurious proliferation in individual wells. An example of this type of analysis is provided in Example 1.

[1150] A comparison of corresponding forms of non-amino acid substituted PE immunogenic epitope responses versus responses obtained with amino acid substituted PE peptides is used to assess and predict the effects of various

amino acid substitutions in reducing or eliminating the immunogenicity of PE polypeptides (i.e., for making deimmunized forms of PE).

[1151] Assays for measuring and testing the immunogenicity of amino acid substituted forms of PE may also be done as described and exemplified in Example 1 (i.e., via proliferation assays quantitating CD4+ T cell responses) wherein amino acid substituted forms of PE (i.e., "deimmunized PE" or "DI-PE"), and/or DI-PE conjugates and fusion proteins (e.g., fusions of DI-PE to antibodies or antigen-binding fragments thereof) are tested and measured for the presence and potency of immunogenic responses compared to responses induced by corresponding forms of non-amino acid substituted PE peptides, polypeptides, and fusion or conjugation constructs.

[1152] Assays for measuring immunogenicity of amino acid substituted forms of PE specifically (as indicated above), or PE molecules, generally, may also be done according to methods routinely used and well-known to those of skill in the art. For example, immunogenicity of amino acid substituted forms of PE, in particular, or PE molecules, in general, (as indicated above) may be measured in vivo in non-human primates and/or in transgenic mouse model systems.

Example 3

Measuring Biological Activity of Amino Acid Substituted Forms of PE

[1153] Assays for measuring the biological activity of amino acid substituted/deimmunized forms of PE, may be done according to methods routinely used and well-known to those of skill in the art. Measured biological activities of deimmunized ("DI") forms of PE ("DI-PE"), in particular, or PE molecules, in general, may include, for example, assays to measure:

[1154] a) general or specific inhibition of protein synthesis (i.e., measuring inhibition of synthesis of a specific protein (or specific proteins) or inhibition of overall (mass) protein synthesis;

[1155] b) inhibition of translation elongation factor EF-2 biological activity;

[1156] c) induction or catalysis of ADP-ribosylation of EF-2; and

[1157] d) eukaryotic cell killing activity (cell cytotoxicity).

[1158] Assays for Biological Activity: Inhibition of Protein Synthesis

[1159] In one example, measurement of inhibition of protein synthesis may be done via use of in vitro transcription/translation assays (which are routinely used and well-known to those of skill in the art). For example, a cell-free assay may be used to measure DI-PE induced inhibition of in vitro transcription/translation of a target plasmid (such as, but not limited to, T7-luc). In the case of using a T7-luc transcription/translation assay, the biological activity readout would be chemiluminescent measurement of luciferase activity wherein amino acid substituted forms of PE are compared to corresponding non-amino acid substituted forms of PE for ability/inability to inhibit translation of the luciferase enzyme in vitro. In such assays, the PE polypeptides being assayed can be introduced via expression from template DNA (e.g., a PCR product) encoding the toxin-conjugate gene, or by directly introducing quantified amounts of PE proteins. Such assays may be used to assess IC50 values* of the various

forms of PE tested (*IC50=concentration at which 50% of protein synthesis is inhibited versus standardized control samples lacking PE).

[1160] Some examples of kits and reagents available for in vitro transcription/translation assays include, but are not limited to:

[1161] TNT® SP6 Coupled Reticulocyte Lysate System (e.g., PROMEGA® catalog #L4610 (PROMEGA® Corp., Madison, Wis., USA)) allows for eukaryotic cell-free protein expression in a single-tube, as a coupled transcription/translation process. More traditional rabbit reticulocyte lysate translations commonly use RNA synthesized in vitro from SP6, T3 or T7 RNA polymerase promoters and require three separate reactions with several steps between each reaction. The TNT® System bypasses many of these steps by incorporating transcription directly into the translation mix. See e.g., PROMEGA® Technical Bulletin #TB126 (Revised December 2010) which is incorporated by reference herein. See also, Pelham et al., *Eur. J. Biochem.* 67, 247-56 (1976); Krieg et al., (1984) *Nuc. Acids Res.* 12, 7057-7070 (1984). See also, U.S. Pat. Nos. 5,324,637; 5,492,817; 5,641,641; and, 5,650,289.

[1162] TNT® T7 Quick Coupled Transcription/Translation System (e.g., PROMEGA® catalog #1170 (PROMEGA® Corp., Madison, Wis., USA)) further simplifies in vitro transcription-translation reactions by combining RNA polymerase, nucleotides, salts and Recombinant RNasin® Ribonuclease Inhibitor with the reticulocyte lysate to form a single TnT® Quick Master Mix. The TnT® Quick Coupled Transcription/Translation System may be used with plasmids for transcription and translation of genes cloned downstream from either the T7 or SP6 RNA polymerase promoters. The TnT® Quick System includes a luciferase-encoding control plasmid and Luciferase Assay Reagent, which can be used in a non-radioactive assay for rapid (<30 seconds) detection of functionally active luciferase protein. Starting with either circular plasmid DNA or PCR-generated DNA, in vitro transcription/translation results may be obtained in 5-6 hours. See e.g., PROMEGA® Technical Bulletin #TM045 (Revised May 2011) which is incorporated by reference herein.

[1163] STEADY-GLO® Luciferase Assay System (e.g., PROMEGA® catalog #E2510) (PROMEGA® Corp., Madison, Wis., USA)) allows for high-throughput quantitation of firefly (*Photinus pyralis*) luciferase expression in mammalian cells via batch processing of 96- and 384-well plates. The STEADY-GLO®, Luciferase Assay System provides signal half-lives of over 5 hours in commonly used cell culture media without prior sample processing. Throughput rates of several thousand samples per hour may be achieved with high reproducibility under standard laboratory conditions. See e.g., PROMEGA® Technical Bulletin #TM051. (Revised March 2009 & Revised September 2011) which is incorporated by reference herein. See also, U.S. Pat. Nos. 5,641,641; 5,650,289; 5,583,024; 5,674,713; and 5,700,673.

[1164] Full protocols for use of such kits are provided by the manufacturer with each kit. A brief example of a typical experimental procedure may include:

[1165] Assembling kit reagents (except target T7-luc plasmid), plus PE test samples (using an experimentally

determined, titration of PE test samples; e.g., in a range of 0-500 ng DNA per reaction for PCR templates or using a PE protein titre in a range to be determined experimentally), in a total volume of 12.5 ul RNase-free water in PCR tubes or cell wells on plates.

[1166] For plasmid DNA: Pre-incubate for required time (e.g., 30-60 min, time to be determined experimentally) at 30° C. to allow pre-reaction transcription/translation to occur.

[1167] For purified protein: No pre-incubation step required.

[1168] Add target plasmid T7-luc (e.g. 250 ng/reaction, determined experimentally) and incubate further (e.g. 30-60 min, time to be determined experimentally) at 30° C.

[1169] Stop reaction by placing on ice. Increase sample volume to 50 ul with RNase-free water.

[1170] Add luciferase reagent (e.g. SteadyGlo, 50 ul per well) to each well, incubate according to manufacturer's instructions, transfer to 96 well black/white plate and read chemiluminescent signal via chemiluminescence platereader.

[1171] Compare to 'zero toxin.' control samples (i.e., no PE present) to determine the % inhibition of transcription/translation (i.e., as a function of inhibition of luciferase activity).

[1172] Compare inhibition of transcription translation values of amino acid substituted/deimmunized forms of PE compared to corresponding forms of non-amino acid substituted PE.

[1173] Comparative protein synthesis inhibition values may show that various forms of DI-PE exhibit 100% or about 100% of biological activity (inhibition of protein synthesis) compared to corresponding forms of non-amino acid substituted PE. Comparative protein synthesis inhibition values may also show that various forms of DI-PE exhibit at least 95%, or at least about 95%, at least 90%, at least about 90%, at least 85% at least about 85%, at least 80%, at least about 80%, at least 75%, at least about 75%, at least 70%, at least about 70%, at least 60%, at least about 60%, at least 50%, or at least about 50% of biological activity compared to corresponding forms of non-amino acid substituted PE.

[1174] Assays for Biological Activity: Cell Cytotoxic Activity

[1175] In one example, measurement of cell cytotoxic activity may be done via use of in vitro cell based assays wherein deimmunized PE (DI-PE)-antibody conjugates are assayed in comparison to non-amino acid substituted PE-antibody conjugates. The antibody portion of such conjugates would be antibodies, or antigen-binding fragments thereof, which specifically bind antigens expressed on the cell-surface of cell types used in such in vitro assays. Cell cytotoxicity may be quantitated, for example, by measuring cell lysis wherein the biological readout is represented by measurement of, for example, based on chemiluminescent (LUMI), fluorometric (FL), and colorimetric (COL) outputs; such as can be practiced using commercially available kits routinely used and well-known to those of skill in the art.

[1176] Some examples of kits available for measurement and comparison of DI-PE versus non-amino acid substituted PE cell cytotoxicity include, without limitation:

[1177] TOXLIGHT® BioAssay Kit (e.g., Catalog #LT07-117 (Lonza Rockland, Inc., Rockland, Me., USA)) is a non-destructive bioluminescent cytotoxicity

assay that quantitatively measures release of Adenylate Kinase (AK) from damaged mammalian cells and cell lines in vitro. The assay is based on the bioluminescent measurement of AK which is present in all cells. A loss of cell integrity, through damage to the plasma membrane, results in the leakage of a number of factors from cells cultured in vitro into the surrounding medium. The measurement of the release of AK from the cells allows the accurate and sensitive determination of cytotoxicity and cytolysis. The reaction involves two steps. The first involves the addition of ADP as a substrate for AK. In the presence of the enzyme, AK, the ADP is converted to ATP for assay by bioluminescence. The bioluminescent part of the assay utilizes the enzyme Luciferase, which catalyses the formation of light from ATP and luciferin. By combining these two reactions, the emitted light intensity is linearly related to the AK concentration and can be measured using a luminometer or beta counter. See, "TOXILIGHT® BioAssay Kit: Instructions for Use," ©2007 Lonza Rockland, Inc., which is incorporated by reference herein. See also, Crouch, et al., *J. Immunol. Methods*, 160(1):81-88 (1993); Olsson, T. et al., *J. Appl. Biochem* 5, 347-445 (1983); and, Squirrel et al., *A Practical Guide to industrial Uses of ATP Luminescence in Rapid Microbiology*, p. 107-113 (1997).

[1178] CYTOTOX-GLO® (e.g., PROMEGA® catalog #G9290 (PROMEGA® Corp., Madison, Wis., USA)) is a luminescent cytotoxicity assay that measures the relative number of dead cells in cell populations. The assay measures extracellular activity of a distinct intracellular protease activity (dead-cell protease) when the protease is released from membrane-compromised cells. A luminogenic cell-impermeant peptide substrate (AAF-aminoluciferin) is used to measure dead-cell protease activity. The liberated aminoluciferin product is measured as "glow type" luminescence generated by ULTRA-GLO® Recombinant Luciferase provided in the assay reagent. The AAF-aminoluciferin substrate cannot cross the intact membrane of viable cells and does not generate appreciable signal from the live-cell population. The amount of luminescence directly correlates with the percentage of cells undergoing cytotoxic stress. With the addition of a lysis reagent (provided with the kit), the CYTOTOX-GLO™ Assay provides a luminescent signal associated with the total number of cells in each assay well. Viability can be calculated by subtracting the luminescent dead-cell signal from the total luminescent value, thus allowing normalization of assay data to cell number and mitigation of assay interferences. The cytotoxicity protease biomarker is constitutive and conserved across cell lines. See e.g., PROMEGA® Technical Bulletin Nos. TB359 (Revised May 2009 & Revised October 2011) which is incorporated by reference herein. See also, Niles, A. et al. (2007) *Anal. Biochem.*, 366, 197-206 (2007) and U.S. Pat. Nos. 6,602,677 and 7,241,584.

[1179] CYTOTOX-ONE™ kit (e.g., PROMEGA® catalog #G7891 (PROMEGA® Corp., Madison, Wis., USA)) allows performance of homogeneous membrane integrity assays wherein a fluorometric method may be used to estimate the number of nonviable cells present in multiwell plates. This assay measures the release of lactate dehydrogenase (LDH) from cells with damaged membranes, LDH released into the culture medium is

measured with a coupled enzymatic assay that results in the conversion of resazurin into a fluorescent resorufin product. The amount of fluorescence produced is proportional to the number of lysed cells (which may be monitored using a 96- or 384-well plate formats). The CYTOTOX-ONE® Reagent does not damage normal healthy cells. Therefore, reactions to measure released quantities of LDH can be performed directly in a homogeneous format in assay wells containing a mixed population of viable and damaged cells. See e.g., PROMEGA® Technical Bulletin #TB306 (Revised May 2009) which is incorporated by reference herein. See also, U.S. Pat. Nos. 6,982,152 and 7,282,348.

[1180] CELLTITER GLO® Luminescent Cell Viability Assay (e.g., PROMEGA® catalog #G7571 (PROMEGA® Corp., Madison, Wis., USA)) provides a homogeneous method for determining the number of viable cells in a culture based on quantitation of the amount of ATP present (an indicator of metabolically active cells). The CELLTITER GLO® Assay is particularly useful for automated high-throughput screening (HTS), cell proliferation and cytotoxicity assays. The homogeneous assay procedure involves adding the single reagent (CELLTITER GLO® Reagent) directly to cells cultured in serum-supplemented medium. The assay allows for detection of as few as 15 cells/well in a 384-well format in 10 minutes after adding reagent and mixing. The homogeneous "add-mix-measure" format results in cell lysis and generation of a luminescent signal proportional to the amount of ATP present (which is directly proportional to the number of cells present in culture). The CellTiter-Glo® Assay generates a "glow-type" luminescent signal, which has a half-life generally greater than five hours, depending on cell type and medium used. See e.g., PROMEGA® Technical Bulletin Nos. TB288 (Revised June 2009 & Revised August 2011) which are incorporated by reference herein. See also: U.S. Pat. Nos. 6,602,677; 7,241,584; 7,700,310; 7,083,911; 7,452,663; 7,732,128; 7,741,067; 5,583,024, 5,674,713; and 5,700,673.

[1181] VIALIGHT® Plus Kit (e.g., Catalog #LT07-221 (Lonza Rockland, Inc., Rockland, Me., USA)) may be used for rapid detection of cytotoxicity in mammalian cells and cell lines in culture via determination of ATP levels. Any form of cell injury results in a rapid decrease in cytoplasmic ATP levels. Therefore, the VIALIGHT® Plus Kit may be used to measure a wide range of biological activities effecting cell viability. The kit is formulated for use with a microtitre plate reading luminometer for assay automation. The assay is based on bioluminescent measurement of ATP is present in all metabolically active cells. The bioluminescent method utilizes an enzyme, luciferase, which catalyses the formation of light from ATP and luciferin according to the following reaction:



The emitted light intensity is linearly related to the ATP concentration and can be measured using a luminometer or beta counter. The assay is conducted at ambient temperature (18° C.-22° C.), the optimal temperature for luciferase enzymes. See, "VIALIGHT® Plus Kit: Instructions for Use," ©2007 Lonza Rockland, Inc., which is incorporated by reference herein.

[1182] Full protocols for use of such kits are provided by the manufacturer with each kit. A brief example of a typical experimental procedure may include:

[1183] Plate cells to test plate (e.g., 96 well plates) in growth medium.

[1184] Incubate cells with titrations of amino acid substituted forms of PE-toxin conjugates (including zero toxin and non-amino acid substituted PE controls (up to a maximum toxicity point, e.g. 100% cell lysis) for required time (determined experimentally, e.g. 48-72 hr).

[1185] Add kit reagents for cytotoxicity measurements as per manufacturer's instructions.

[1186] Transfer test samples to 96 well black/white walled plate (as appropriate) and read reaction signal output.

[1187] Compare cell cytotoxicity values obtained for substituted/deimmunized forms of PE versus corresponding non-amino acid substituted forms of PE.

[1188] Comparative cell cytotoxicity values may show that various forms of DI-PE exhibit 100% or about 100% of biological activity (induction of cell cytotoxicity) compared to corresponding forms of non-amino acid substituted PE. Comparative cell cytotoxicity values may also show that various forms of DI-PE exhibit at least 95%, or at least about 95%, at least 90%, at least about 90%, at least 85%, at least about 85%, at least 80%, at least about 80%, at least 75%, at least about 75%, at least 70%, at least about 70%, at least 60%, at least about 60%, at least 50%, or at least about 50% of biological activity compared to corresponding forms of non-amino acid substituted PE.

Example 4

Measuring Ability of Deimmunized PE Variants to Inhibit Protein Synthesis

[1189] Quantitative in vitro transcription/translation (IVTT) assays to assess the biological activity of deimmunized variants of PE in inhibiting protein synthesis (i.e., possess wild-type PE biological activity) may be performed using the TNT® Quick Coupled Transcription/Translation Systems assay from PROMEGA® Corp. (Madison, Wis., USA), See. PROMEGA® Technical Bulletin #TB126 (Revised December 2010) which is incorporated by reference herein.

Example 5

Measuring Ability of a PE-IL2 Fusion Protein to Inhibit Protein Synthesis in an In Vitro Transcription/Translation (IVTT) Assay

[1190] A preliminary experiment was performed to compare the ability of a PE-IL2 fusion protein to inhibit protein synthesis in an in vitro transcription/translation assay when a commercially available PE-IL fusion protein is translated in vitro following transcription from either a circular plasmid expression vector or a linearized plasmid expression vector. The PE-IL2 expression vector in this experiment is referred to as "VVN-52431." A few examples of IL2-PE fusion construct are shown in SEQ ID NO:164, 165 and 166. The aim of this experiment was to determine if circular or linearized plasmids produced significantly different quantities of PE-IL protein in the PROMEGA® Corp. TNT® Quick Coupled Transcription Translation Systems assay. A commercially available T7 Pro-

moter/Luciferase expression vector (PROMEGA® Corp.; hereinafter "T7-Luc DNA") was used to measure the ability of PE-IL2 to inhibit protein synthesis in vitro.

[1191] Based on a pilot IVTT experiment, it was determined that 0.2 µg T7-Luc DNA provided optimal RLU (Relative Light Units) in a 90 minute IVTT reaction. In this experiment, VVN-52431 was linearized using the restriction enzyme Fsp-I. Linearized and circular VVN-52431 DNA were used as templates in the IVTT reactions. Reactions were done in triplicate, using 0.5, 1 and 2 µg of DNA. The T7 control reaction was performed using 1 µg DNA. Reactions were analyzed via SDS-PAGE and by Luciferase assay.

[1192] Materials:

Item	Vendor	Lot #
Nuclease-free water (1000 ml)	Ambion	1105062
TNT T7 Quick Coupled T/T system	PROMEGA ®	328577
T7 luciferase plasmid DNA (From same kit)	PROMEGA ®	
Fsp I	NEB	0571101
Dual Glo ® Luciferase Assay System	PROMEGA ®	322310
Ultrapure Water	GIBCO	896656
Tris-Glycine SDS Sample Buffer, 2x	Invitrogen	743995
10x Reducing Agent	Invitrogen	897034
Criterion Tris HCl 4-15%, 1 mm, 12 + 2 well	Bio-Rad	400059499
Precision Plus Protein Standards, Kaleidoscope	Bio-Rad	310009928
10x Tris/Glycine/SDS Buffer	Bio-Rad	210007884
Gelcode Blue Safe Protein Stain	ThermoFisher	LL152043

[1193] Equipment:

Hera	Vendor	ID #
P20, P200, P1,000	Rainin	N/A
Water bath		
Luminometer		
Power Pac HC	Bio-Rad	N/A
Heat block	VWR	N/A
Microcentrifuge, refrigerated	Eppendorf	N/A
Platform Adjustable Tilt Rocker	Labnet	N/A
Thermal cycler	MJ Research	N/A

[1194] Procedure:

[1195] Per manufacturer's instructions: Except for the actual transcription/translation incubation, all handling of the TNT® Quick Master Mix was performed at 4° C. Unused Master Mix was refrozen as soon as possible after thawing to minimize loss of translational activity.

[1196] Restriction Digest:

[1197] In PCR tubes, the following were combined:

[1198] VVN-52431 was linearized by combining the following:

Rxn	NF H ₂ O (µL)	VVN52431 (µL)	Fsp I dig.	final	Reaction Product
1	5	5 (5 µg)	0	0.5 µg/µl	Circular Vector - No Restriction

-continued

Rxn	NF H ₂ O (μL)	VVN52431 (μL)	Fsp I dig.	final	Reaction Product
2	4	5 (5 μg)	1	0.5 μg/μl	enzyme added Linearized Vector

[1199] Reactions were incubated at 37° C. for 60 min.

[1200] Reactions were heat inactivated at 65° C. for 20 min.

[1201] IVTT Reactions:

[1202] 1. In nuclease-free 1.5 ml eppendorf tubes, the following were combined according to the chart below:

[1203] Diluted T7-Luc DNA in NF (nuclease free) water at 1:5. Final DNA concentration=0.1 μg/μl.

[1204] Serial diluted unlinearized and linearized VVN-52431 DNA (0.5 μg/μl) in NF water at 1:5. Final concentration=0.5 μg/μl, 0.1 μg/μl, 0.02 μg/μl.

Rxn	NF H ₂ O (μL)	T7 Luc DNA (μL)	VVN52431 (μL)	Fsp I dig.	Methi- onine (μL)	T7 TNT rex (μL)
1	7	2 (0.2 μg)			1	40
2	7		2 (1 μg)	-	1	40
3	7		2 (1 μg)	+	1	40
4	5	2 (0.2 μg)	2 (1 μg)	-	1	40
5	5	2 (0.2 μg)	2 (0.2 μg)	-	1	40
6	5	2 (0.2 μg)	2 (0.04 μg)	-	1	40
7	5	2 (0.2 μg)	2 (1 μg)	+	1	40
8	5	2 (0.2 μg)	2 (0.2 μg)	+	1	40
9	5	2 (0.2 μg)	2 (0.04 μg)	+	1	40
10	9				1	40

[1205] 2. Reactions were incubated at 30° C. for 90 minutes in a water bath.

[1206] 3. Reactions were analyzed for the synthesis of functional Luciferase using a standard Luciferase assay.

[1207] Luciferase Assay:

[1208] 1. Luciferase assay substrate was prepared according to manufacturer's instructions:

[1209] Reagent Kit was thawed at room temperature.

[1210] Dual-Glo® Luciferase Buffer was transferred into the Dual-Glo® Luciferase Substrate bottle and shaken slightly to ensure the substrate dissolved.

[1211] Dual-Glo® Stop & Glo® substrate was transferred into the Dual-Glob Stop & Glo® buffer and mixed well.

[1212] Rehydrated reagent was aliquoted into 15 ml centrifuge tube (10 ml/tube) and wrapped with Aluminum foil.

[1213] Rehydrated reagent was stored at -80° C. until ready for use (the reagent is good for 6 months).

[1214] 2. 5 μl of reaction end products/well were transferred into a 96-well white plate

[1215] 3. 100 μL of the Luciferase Assay Reagent was dispensed per well and mixed by pipetting 2-3x.

[1216] 4. RLU's for each well on plate were read within 10 minutes.

[1217] Results:

[1218] Results are shown in FIG. 9. Circular plasmid expression vector encoding PE-IL2 fusion protein was slightly better at inhibiting Luciferase protein synthesis compared to linearized plasmid encoding the same (at all

Luciferase vector: VVN-52431 vector ratios). These results also demonstrate the ability of to test and compare the biological activity of PE-fusion proteins in inhibiting protein synthesis.

[1219] In addition to measuring inhibition of protein synthesis as a measure of light production catalyzed by Luciferase, quantitative analysis of inhibition of protein synthesis was also performed by separating polypeptide reaction products on SDS-PAGE gels, staining, and assessing amounts of polypeptide produced (data not shown).

[1220] Assays such as these may be used to compare the ability of amino acid substituted (e.g., deimmunized) forms of PE (alone or as fusion proteins) to retain biological activity (such as inhibition of protein synthesis) compared to corresponding non-amino acid substituted forms of PE (alone or as fusion proteins).

Example 6

In Vitro Transcription/Translation (IVTT) Assay to Measure and Compare Ribosylation Activity of Amino Acid Substituted Variants of PE

[1221] Purpose:

[1222] This protocol provides an example of they type of methods which may be used to measure and compare the ribosylation activity (i.e., inhibition of protein synthesis) of amino acid substituted forms of PE compared to corresponding non-amino acid substituted forms of PE.

[1223] Background:

[1224] The IVTT assay measures PE mediated inhibition of in vitro transcription/translation of a target plasmid, T7-Luc. The level of inhibition (or lack thereof) is determined by chemiluminescent measurement of luciferase activity (i.e., the transcribed and translated protein). In this assay, a lowered level of transcription and translation (and thereby, lowered levels of chemiluminescent light output) corresponds to increased inhibition of protein synthesis. IVTT can be performed using template DNA encoding PE, or by directly using quantified protein. This assay may be used to rank different PE variants against each other and to compare their biological activities to corresponding non-amino acid substituted forms.

[1225] Materials:

[1226] Test sample: Vectors comprising either circular plasmids with an SP6 promoter or linearized plasmids with a T7 promoter.

Reagents and Materials	Vendor
Nuclease-free water (1000 ml)	Ambion
TNT SP6 Quick Coupled T/T system	PROMEGA ®
SP6 luciferase plasmid DNA (From same kit)	PROMEGA ®
RNase-free 1.5 ml microfuge tubes	Ambion
Dual Glo ® Luciferase Assay System	PROMEGA ®

[1227] Equipment:

Item	Vendor
P20, P200, P1,000 96 well white plate Water bath (circulation) Luminometer Microcentrifuge, refrigerated	Rainin Costar Eppendorf

[1228] Reagent Preparation:**[1229] Luciferase Preparation:**

[1230] 1. Thaw reagent Kit on ice or at 4 degrees C.

[1231] 2. Transfer entire Dual-Glo® Luciferase Buffer into Dual-Glo® Luciferase Substrate bottle and shake bottle slightly to ensure substrate completely dissolved.

[1232] 3. Transfer entire Dual-Glo® Stop & Glo® substrate into Dual-Glo® Stop & Glo® buffer and mix well.

[1233] 4. Aliquot rehydrated reagent into 15 ml centrifuge tube (10 ml/tube) and wrap tube with Aluminum foil.

[1234] 5. Store rehydrated reagent in -80° C. Freezer (reagent is good for 6 months)

[1235] Procedure:

[1236] Per manufacturer's recommendations: Except for the actual transcription/translation incubation, all handling of TNTR Quick Master Mix should be done at 4° C. Any unused Master Mix should be refrozen as soon as possible after thawing to minimize loss of translational activity. Do not freeze-thaw the Master Mix more than two times.

[1237] Plasmid DNA Dilution:

[1238] Dilute plasmid DNA and Luc plasmid DNA in nuclease free water to final concentration of 0.1 µg/µl.

[1239] IVTT Reactions

[1240] 1. In 1.5 ml NF (nuclease free) eppendorf tubes, the following are combined for each test sample:

[1241] a. 5 µL of NF H₂O;

[1242] b. 2 µl, 0.1 µg/µl test plasmid (for increased accuracy of results test a dilution series of samples);

[1243] c. 1µl mM Methionine;

[1244] d. 40 µL TNT quick master mix;

[1245] e. Negative control (reaction mix only. NF water 9 µl, methionine 1 µl, reaction mixture 40 µl;

[1246] 2. Incubate reaction mixes at 30° C. for 15 minutes in water bath;

[1247] 3. Add 2 µl, 0.1 µg/µl Luc plasmid to each reaction mix

[1248] 4. Incubate reactions at 30° C. for 90 minutes in water bath; and

[1249] 5. Transfer all samples onto ice to stop reaction.

[1250] Luciferase Assay

[1251] 6. Transfer 5 µl of end product/well into 96-well white plates in triplicate;

[1252] 7. Dispense 100 µL of Luciferase Assay Reagent per well. Mix by pipetting 2-3x;

[1253] 8. Read entire plate within 10 minutes.

[1254] Calculations

[1255] 1. Calculate percent inhibition based on relative luminescence units (RFU) of the test sample divided by the RFU of the LUC plasmid with no test sample, then subtract the result from 100.

[1256] 2. Calculate the percent of activity of non-amino acid substituted PE by percent inhibition of the test

sample divided by the percent inhibition of non-amino acid substituted, then multiply the result by 100,

[1257] 3. If a dilution series of samples is tested, calculate the IC₅₀ (half maximal inhibition concentration) for each sample using the RFU of the test sample divided by the RFU of the LUC plasmid alone, then subtract the result from 100. Determine the concentration which results in 50% inhibition.

[1258] 4. Calculate the percent of non-amino acid substituted PE inhibition by dividing the IC₅₀ of the non-amino acid substituted PB by the IC₅₀ of the test sample and multiplying the result by 100.

Example 7

Ex Vivo Assays to Assess Immunogenicity of Amino Acid Substituted Forms of PE (i.e. Deimmunized PE) Versus Corresponding Non-Amino Acid Substituted Forms

[1259] The immunogenicity of amino acid substituted forms of PE (alone or as PE-fusion proteins) are assessed using methods well-known and routinely used by those skilled in the art. For example, ELISA assays are used wherein serum is assayed ex vivo (following extraction from organisms in which amino acid substituted forms of PE, or non-amino acid substituted PE, (alone or as fusion proteins) are administered) to determine whether or not antibodies that specifically bind the administered protein are produced. It is noted that in the case of PE-fusion proteins it is necessary to use, as the ELISA assay target antigen, not only intact PE-fusion proteins (i.e., amino acid substituted or non-amino acid substituted PB), but to also test the PE component and the polypeptide fusion component separately to determine whether or not antibodies produced specifically bind the PB portion or the fusion polypeptide portion (e.g., IL2 as used in a previous example of a PE-IL2 fusion protein). Accordingly, it is most desirable to identify amino acid substituted forms of PE which do not result in host production of antibodies that specifically bind modified forms of PE (i.e., deimmunized forms of PE).

[1260] Organisms in which amino acid substituted forms of PE may be administered (alone or as fusion proteins) include, for example, without limitation: mice (including transgenic mice expressing human immunoglobulin genes), rats, rabbits, dogs, goats, sheep, horses, cows (and other bovine species), non-human primates, and humans.

Example 8

In Vitro and In Vivo Assays to Assess Cytotoxicity of Amino Acid Substituted Forms of PE

[1261] The cytotoxicity of amino acid substituted forms of PE (alone or as PE-fusion proteins) are assessed using methods well-known and routinely used by those skilled in the art. For example, the cytotoxic effects of amino acid substituted forms of PE (alone or as PE-fusion proteins) administered to cells in vitro or organisms in vivo, may be assessed with reference to cytotoxic (cell killing) effects on target cells, organs, tissues, or tumors against which PE or PE fusion proteins are expected to produce a cytotoxic effect. For example, the therapeutically beneficial cytotoxic effects of amino acid substituted PE-Mesothelin fusions may be assessed by monitoring and measuring reduction or elimination of tumor or cancer cells or tissues (in vitro or in vivo) in

response to administration of amino acid substituted forms of PE-Mesothelin versus wild-type PE-Mesothelin fusion.

[1262] Organisms in which amino acid substituted forms of PE may be administered (alone or as fusion proteins) include, for example, without limitation: mice (including transgenic mince expressing human immunoglobulin genes), rats, rabbits, dogs, goats, sheep, horses, cows (and other bovine species), non-human primates, and humans.

Examples 9-13

[1263] Oligonucleotides referenced in the following examples are listed in Table 12.

Example 9

Generation of cDNAs Encoding Amino Acid Substituted Forms of PE

[1264] The Kozak sequence in vector pET14b (EMD Millipore catalog #69660, Darmstadt, Germany) was modified by introducing a linker made up of annealed oligonucleotides 5'-CATGGT

GGCTCTCCTTCTTAAAGTTAAACAAAATTATTT-3' (SEQ ID NO:239) (OL2216 in Table 12) and

[1265] 5'-CTAGAAATAATTTGTTTAACTTTAA-GAAGGAGAGCCAC-3' (SEQ ID NO:240) (OL2217 in Table 12) (underlined letters indicate nucleotides changed in the Kozak sequence*) via NcoI and XbaI restriction sites into vector pET14b resulting in a modified Kozak sequence (SEQ ID NO:176) by mutation of three nucleotides at positions 587 to 589. The resulting vector was named pET14b-K.

[1266] *Kozak sequence=(gcc)gccRccAUGG (SEQ ID NO:286), where R is a purine (i.e., adenine or guanine) three

bases upstream of the start codon (AUG), which is followed by another 'G'. See, Kozak, *Nucleic Acids Res.* 15 (20); 8125-8148 (1987).

[1267] Oligonucleotides for generation of genes encoding amino acid substituted forms of PE are listed as OL2164 to OL2194 and OL2281 to OL2366 in Table 12. A wild-type (WT) PE gene (SEQ ID NO:1) was made by gene synthesis and amplified using oligonucleotides: 5'-ATTGTCCATATGCCAGAAGGCGGTAGCCTGGC-3' (SEQ ID NO:215) (OL2154 in Table 12) to introduce a NdeI site, and 5'-ATCCTCGAGTTACTTCAGGTCCTCACGCGGCG-3' (SEQ ID NO:222) (OL2167 in Table 12) to introduce a XhoI site. The resulting DNA fragment was subcloned into pGEMT®-T (PROMEGA® catalogue #A1360, PROMEGA®, Southampton, UK). Colonies were screened by PCR using M13 primers OL0001 and OL0002 (Table 12). Subsequently the wild-type (WT) PE gene was subcloned into pET14b-K using NdeI and XhoI restriction enzymes (Fermentas catalog #FD0583 and FD0695, respectively) and resulting in a gene encoding an N-terminal His6 tag fused to the WT PE sequence. The resulting vector was named pET1.4b-K-WT PE.

[1268] Oligonucleotides for generation of genes encoding amino acid substituted forms of PE were generated using overlapping PCR with the WT PE gene in pET14b-K as template. Pairs of primers from the oligonucleotides of Table 12 (as noted in the "application" column of Table 12) were annealed to WT PE DNA and the amino acid substituted genes were PCR amplified using terminal oligonucleotides: 5'-ATCTCCCTCTAGAAATAATTTTGTTTAACTTTAAGAAG-3' (SEQ ID NO:241) (OL2268 in Table 12) and 5'-ATCCTCAGTTACTTCAGACCAGGTCCTCACCGGCG-3' (SEQ ID NO:216) (OL2161 in Table 12). PCR fragments were and cloned into pET14b-K using XbaI and XhoI restriction sites.

TABLE 12

Oligonucleotides				
Name	sequence	length	application	
OL	001	CGCCAGGGTTTTCCAGTCAC GAC (SEQ ID NO: 205)	24	M13 FOR
OL	002	AGCGGATAACAATTTCACACA GGA (SEQ ID NO: 206)	24	M13 REV
OL	2043	GAAGTGCAGCTGGTGGAG (SEQ ID NO: 207)	18	RFB4 VH5' PCR primer sequence
OL	2044	CAGAGCCACCTCCGCCTGAAC CGCCTCCACCTGAGGAGACA GTGACCAG (SEQ ID NO: 208)	49	RFB4 VH3' PCR primer sequence
OL	2045	CAGGCGGAGGTGGCTCTGGC GGTGGCGGATCGGATATCCA GATGACCCAG (SEQ ID NO: 209)	50	RFB4 VK 5' PCR Primer Sequence
OL	2046	TTTGATCTCCAGCTTGGTG (SEQ ID NO: 210)	19	RFB4 VK 3' PCR Primer sequence
OL	2047	CCCAGCCGGCCATGGCGGAA GTGCAGCTGGTGGAG (SEQ ID NO: 211)	35	RFB4 Pull through Primer (FOR))
OL	2048	GGTGCTCGAGTGGCGCCGCC GTTTGATCTCCAGCTTGGTG (SEQ ID NO: 212)	41	RFB Pull through Primer (REV)

TABLE 12-continued

Oligonucleotides			
Name	sequence length	application	
OL 2097	AACCGCCCGGCCGTTCTTCTC 39 CGTGTGCCCCGAAAGCC (SEQ ID NO: 213)	IEX02 GroEL/ES REV	
OL 2098	GGGCCAAAGCTTGTTCTTGTT 36 TGAGTCCACTCATGG (SEQ ID NO: 214)	IEX02 GroEL/ES FOR	
OL 2154	ATTGTCCATATGCCAGAAGGC 32 GGTAGCCTGGC (SEQ ID NO: 215)	IEX02 PE38 FOR, introducing NdeI	
OL 2161	ATCCTCGAGTTACTTCAGGTC 32 CTCACGCGGCG (SEQ ID NO: 216)	IEX02 PE38 REV, introducing XhoI	
OL 2162	GGGTGGTCGCCCTGGACACTAT 30 CCTGGGTTG (SEQ ID NO: 217)	TEX02 PE38 NM E229D FOR	
OL 2163	CAACCCAGGATAGTGTCCAG 30 GCGACCAACC (SEQ ID NO: 218)	IEX02 PE38 NM E229D REV	
OL 2164	CAGTACGATAGAAACCCGGC 40 AGATTGCTGCGCGGTACGTA (SEQ ID NO: 219)	IEX02 PE38 S253N	
OL 2165	CAGTACGATAGAAACCCGGC 40 AGCTTGCTGCGCGGTACGTA (SEQ ID NO: 220)	IEX02 PE38 S253K	
OL 2166	CAGTACGATAGAAACCCGGC 40 AGAGGGCTGCGCGGTACGTA (SEQ ID NO: 221)	IEX02 PE38 S253P	
OL 2167	CAGTACGATAGAAACCCGGC 40 AGGGTGCTGCGCGGTACGTA (SEQ ID NO: 222)	IEX02 PE38 S253T	
OL 2168	GTACGTCGTCGTAGCAGAGAC 33 CTGGATGCCATC (SEQ ID NO: 223)	IEX02 PE38 Q206R	
OL 2169	GATGGCATCCAGGTCTCTGCT 33 ACGAGCACGTAC (SEQ ID NO: 224)	IEX02 PE38 Q206R	
OL 2170	CGTAGCCAGGACCTGAAGGC 33 CATCTGGCGTGGC (SEQ ID NO: 225)	IEX02 PE38 D209K	
OL 2171	GCCACGCCAGATGGCCTTCAG 33 GTCCTGGCTACG (SEQ ID NO: 226)	IEX02 PE38 D209K	
OL 2183	GAAGCTGCTCAGTCTGCCGTG 32 TTCGGTGGCGT (SEQ ID NO: 227)	IEX02 PE38 I196A FOR, to pair with OL2161	
OL 2184	ACGCCACCGAACACGGCAGA 32 CTGAGCAGCTTC (SEQ ID NO: 228)	IEX02 PE38 I196A REV, to pair with OL2268	
OL 2185	GAAGCTGCTCAGTCTAACGTG 32 TTCGGTGGCGT (SEQ ID NO: 229)	IEX02 PE38 I196N FOR, to pair with OL2161	
OL 2186	ACGCCACCGAACACGTTAGA 32 CTGAGCAGCTTC (SEQ ID NO: 230)	IEX02 PE38 I196N REV, to pair with OL2268	

TABLE 12-continued

Oligonucleotides				
Name	sequence	length	application	
OL 2187	GGTGATGGCGCGATGCCTCT TTTTCTACCCGC (SEQ ID NO: 231)	33	IEX02 to introduce I153A FOR	
OL 2188	GCGGGTAGAAAAAGAGGCAT CGCCGCCATCACC (SEQ ID NO: 232)	33	IEX02 to introduce I153A REV	
OL 2189	GGTGATGGCGCGATACCTCT TTTTCTACCCGC (SEQ ID NO: 233)	33	IEX02 to introduce I153T FOR	
OL 2190	GCGGGTAGAAAAAGAGGTAT CGCCGCCATCACC (SEQ ID NO: 234)	33	IEX02 to introduce I153T REV	
OL 2191	GGTGATGGCGCGCATCACTCT TTTTCTACCCGC (SEQ ID NO: 235)	33	IEX02 to introduce I153H FOR	
OL 2192	GCGGGTAGAAAAAGAGTGAT CGCCGCCATCACC (SEQ ID NO: 236)	33	IEX02 to introduce I153H REV	
OL 2193	GCACCCAGAACTGGAGAGTT GAACGTCTGCTG (SEQ ID NO: 237)	32	IEX02 to introduce T164R FOR	
OL 2194	CAGCAGACGTTCAACTCTCCA GTTCTGGGTGC (SEQ ID NO: 238)	32	IEX02 to introduce T164R REV	
OL 2216	CATGTGGCTCTCCTTCTTAA AGTTAAACAAAATTATTT (SEQ ID NO: 239)	39	IEX02 Linker to optimize Kozak in pET14b, to anneal with OL2217	
OL 2217	CTAGAAATAATTTTGTTTAAC TTTAAGAAGGAGAGCCAC (SEQ ID NO: 240)	39	IEX02 Linker to optimize Kozak in pET14b, to anneal with OL2216	
OL 2268	ATCTCCCTCTAGAAATAATTT TGTTTAACTTTAAGAAG (SEQ ID NO: 241)	38	IEX02 outside FOR spans over XbaI site (pET14b) - to be paired with OL2161	
OL 2279	GAAGCTGCTCAGTCTATCGTG TTCGGTGGCGT (SEQ ID NO: 242)	32	IEX02 FOR oligo to remove TM to be paired with OL2161	
OL 2280	ACGCCACCGAACACGATAGA CTGAGCAGCTTC (SEQ ID NO: 243)	32	IEX02 REV oligo to remove TM to be paired with OL2268	
OL 2281	CTCTGCTACGAGCACGGGCGC CACCGAACACG (SEQ ID NO: 244)	32	IEX02 A201 REV, ONLY for templates having Q206	
OL 2282	CGTGTTCGGTGGCGCCCGTGC TCGTAGCAGAG (SEQ ID NO: 245)	32	IEX02 A201 FOR, ONLY for templates having Q206	
OL 2283	CATCCAGGTCTCTGCTGGCAG CACGTACGCCAC (SEQ ID NO: 246)	33	IEX02 A204 REV, ONLY for templates having Q206	
OL 2284	GTGGCGTACGTGCTGCCAGCA GAGACCTGGATG (SEQ ID NO: 247)	33	IEX02 A204 FOR, ONLY for templates having Q206	
OL 2285	CATCCAGGTCTCTGCTCTGAG CACGTACGCCAC (SEQ ID NO: 248)	33	IEX02 Q204 REV, ONLY for templates having Q206	

TABLE 12-continued

Oligonucleotides			
Name	sequence length		application
OL 2286	GTGGCGTACGTGCTCAGAGCA 33 GAGACCTGGATG (SEQ ID NO: 249)	IEX02 Q204 FOR, ONLY for templates having Q206	
OL 2287	CCAGTTCTGGGTGCCGCGGT 31 AGAAAAAGAG (SEQ ID NO: 250)	IEX02 A158 REV	
OL 2288	CTCTTTTCTACCGCCGCAC 31 CCAGAACTGG (SEQ ID NO: 251)	IEX02 A158 FOR	
OL 2289	CCAGTTCTGGGTGCCCTGGGT 36 AGAAAAAGAGATATC (SEQ ID NO: 252)	IEX02 Q158 REV	
OL 2290	GATATCTCTTTTCTACCCAG 36 GGCACCAGAACTGG (SEQ ID NO: 253)	IEX02 Q158 FOR	
OL 2291	GTCCAGTTCTGGGTGGAGCGG 38 GTAGAAAAAGAGATATC (SEQ ID NO: 254)	IEX02 S159 REV	
OL 2292	GATATCTCTTTTCTAC- 38 CCGCT CCACCAGAACTGGAC (SEQ ID NO: 255)	IEX02 S159 FOR	
OL 2293	ACCACCCAGAACTGGACCGTT 25 GAAC (SEQ ID NO: 256)	IEX02 T159 REV	
OL 2294	CCAGTTCTGGGTGGTGCGGGT 31 AGAAAAAGAG (SEQ ID NO: 257)	IEX02 T159 FOR	
OL 2295	GAGCTTGGGTCCAGATCGCCA 23 CC (SEQ ID NO: 258)	IEX02 generic REV oligo for mutations at 333	
OL 2296	CTGGACCCAAGCTCTGCCCCG 31 GATAAAGAAC (SEQ ID NO: 259)	IEX02 A333 FOR	
OL 2297	CTGGACCCAAGCTCTAACCCG 28 GATAAAG (SEQ ID NO: 260)	IEX02 N333 FOR	
OL 2298	CTGGACCCAAGCTCTACCCG 28 GATAAAG (SEQ ID NO: 261)	IEX02 T333 FOR	
OL 2299	CTGGACCCAAGCTCTCAGCCG 31 GATAAAGAAC (SEQ ID NO: 262)	IEX02 Q333 FOR	
OL 2300	CTGGACCCAAGCTCTCACCCG 28 GATAAAG (SEQ ID NO: 263)	IEX02 H333 FOR	
OL 2301	CTGGACCCAAGCTCTATCCCG 48 GATAAAGAAAACGCTATTCT GCCCTG (SEQ ID NO: 264)	IEX02 N338 FOR	
OL 2302	CTGGACCCAAGCTCTATCCCG 46 GATAAAGAGAGGCTATTCT GCCC (SEQ ID NO: 265)	IEX02 E338 FOR	
OL 2303	CTGGCTACGAGCACGGGCGC 28 CACCGAAC (SEQ ID NO: 266)	IEX02 V201A REV	

TABLE 12-continued

Oligonucleotides			
Name	sequence length	application	
OL 2304	GTTCGGTGGCGCCCGTGCTCG 28 TAGCCAG (SEQ ID NO: 267)	IEX02	V201A FOR
OL 2305	CTTCAGGTCCTGGCTGGCAGC 30 ACGTACGCC (SEQ ID NO: 268)	IEX02	R204A REV, ONLY for templates having D209K
OL 2306	GGCGTACGTGCTGCCAGCCAG 30 GACCTGAAG (SEQ ID NO: 269)	IEX02	R204A FOR, ONLY for templates having D209K
OL 2307	CTTCAGGTCCTGGCTCTGAGC 29 ACGTACGC (SEQ ID NO: 270)	IEX02	R204Q REV, ONLY for templates having D209K
OL 2308	GCGTACGTGCTCAGAGCCAG 29 GACCTGAAG (SEQ ID NO: 271)	IEX02	R204Q FOR, ONLY for templates having D209K
OL 2309	GTTCAACGGTCCAGTTGTTGG 33 TGCCGCGGGTAG (SEQ ID NO: 272)	IEX02	Q161N REV
OL 2310	CTACCCGCGGCACCAACAAC 33 GGACCGTTGAAC (SEQ ID NO: 273)	IEX02	Q161N FOR
OL 2311	GTTCAACGGTCCAGTTGTTGG 33 TGCCGCGGGTAG (SEQ ID NO: 274)	IEX02	Q161T REV
OL 2312	CTACCCGCGGCACCAACAAC 33 GGACCGTTGAAC (SEQ ID NO: 275)	IEX02	Q161T FOR
OL 2313	CAGCAGACGTTCAACGGCCC 31 AGTTCTGGGTG (SEQ ID NO: 276)	IEX02	T164A REV
OL 2314	CACCCAGAACTGGGCCGTTGA 31 ACGTCTGCTG (SEQ ID NO: 277)	IEX02	T164A FOR
OL 2315	GACGTTCAACGGTCCAGGCCT 33 GGGTGCCGCGGG (SEQ ID NO: 278)	IEX02	N162A REV
OL 2316	CCCGCGGCACCCAGGCCTGG 33 ACCGTTGAACGTC (SEQ ID NO: 279)	IEX02	N162A FOR
OL 2318	ATTGCCACCATGGCGGAAGTG 22 C (SEQ ID NO: 280)	IEX02	FOR RFB4 (NcoI)
OL 2320	CACCAGGCCGCTGCTTTTGAT 31 CTCCAGCTTG (SEQ ID NO: 281)	IEX02	REV to create RFB4 for RFB4-PE38-8xHis to pair with OL2318
OL 2321	CAAGCTGGAGATCAAAAGCA 31 GCGGCCTGGTG (SEQ ID NO: 282)	IEX02	FOR to create RFB4-PE38-8xHis to pair with OL2322
OL 2322	CGATTCTCGAGTTACTTCAGG 66 TCCTCGTGGTGGTGGTGATGA TGATGATGACGCGGCGGTTTA CCC (SEQ ID NO: 283)	IEX02	REV introducing 8xHis C-terminus of PE, introducing XhoI

TABLE 12-continued

Oligonucleotides				
Name	sequence	length	application	
OL	2323	CAAGCTGGAGATCAAAGCTC ATGGGGGCGAGCCATCATCATC ATC (SEQ ID NO: 284)	44	IEX02 FOR to create REB4-6xHis PE38 fusions (pIEX02-302 and pIEX02- 304) to pair with OL2161
OL	2324	GATGATGATGATGGCTGCCCC CATGAGCTTTGATCTCCAGCT TG (SEQ ID NO: 285)	44	IEX02 REV to create RFB4-6xHis PE38 fusions (pIEX02-302 and pIEX02- 304) to pair with OL2318

Example 10

Analysis of Genes Encoding Amino Acid Substituted
Forms of PE by an In Vitro Transcription/Translation
(IVTT) Assay

[1269] The cell-free in vitro transcription/translation (IVTT) assay was performed with a TnT® T7 Coupled Reticulocyte Lysate System (PROMEGA® catalog #L4610) following the procedure described in the User's Manual provided in the kit. See, PROMEGA®, Technical Bulletin #TB126, Revised December 2010, pp. 1-28 (2010) which is incorporated by reference herein.

[1270] WT PE in plasmid pET14b-K was used as standard on every plate and tested at concentrations ranging from 0.08 ng to 10 ng in a 12.5 microliter final volume reaction. All test samples were run in triplicate. DNAs encoding WT or amino acid substituted PE in plasmid pET14b-K were added to the IVTT reaction mix supplemented with NAD⁺ (final concentration 0.15 mM; Fisher Scientific catalog #BPE9746-212) and incubated at 30° C. for 1.5 min. Following a subsequent cooling step at 4° C. for 5 min, 250 ng of T7 Luciferase plasmid (Luciferase T7 control DNA supplied in the TnT® T7 Coupled Reticulocyte Lysate System) was added to each reaction and incubated at 30° C. for 90 min. The reaction was stopped by placing the samples on ice. Samples were analyzed using the STEADY-GLO® Luciferase Assay (PROMEGA® catalog #E2510) according to the protocol provided by the manufacturer. See, PROMEGA®, Technical Bulletin #TM051, revised 9/11, pp. 1-23 (2011) which is incorporated by reference herein. Luminescence was measured in a FLUOstar OPTIMA plate reader (BMG Labtech Ltd., Aylesbury, UK)

[1271] A representative result is shown in FIG. 10 which shows the results expressed in CPS (counts per second as read from the FLUOstar OPTIMA fluorescence plate reader) for luciferase activity from IVTT assays of genes encoding either WT PE (pIEX02-001 (SEQ ID NO:1)) or encoding amino acid substituted PE (pIEX02-228 (SEQ ID NO: 177), pIEX02-244 (SEQ ID NO:178), pIEX02-246 (SEQ ID NO:179)); which were expressed as fusion proteins comprising a histidine polymer and a linker sequence preceding a sequence of WT PE or amino acid substituted PE; see, pIEX02-001 PE WT (SEQ ID NO:189 (DNA) and SEQ ID NO:190 (AA)), pIEX02-228 amino acid substituted PE (SEQ ID NO:193 (DNA) and SEQ ID NO:194 (AA)), pIEX02-244 amino acid substituted PE (SEQ ID NO:197 (DNA) and SEQ ID NO:198 (AA)), pIEX02-246 amino acid substituted PE (SEQ ID NO:201 (DNA) and SEQ ID NO:202 (AA)). See c/s, Table 13.

[1272] For the analysis of various amino acid substituted PEs, the potency of each mutated PE in inhibiting IVTT was expressed as relative inhibition exhibited via expression from 2.5 nanograms of DNA encoding amino acid substituted PE compared to expression from 2.5 nanograms WT PE DNA as shown in Table 13 (data expressed as % inhibition of IVTT for the DNA encoding amino acid substituted PE compared to wild-type PE). In order to identify "inhibitory" amino acid substituted PE polypeptides (i.e., genes encoding amino acid substituted forms of PE which inhibit IVTT), selected mutations in each T cell epitope as identified in Table 11 were initially tested using various combinations of epitope 5 mutations (e.g., corresponding to S241N, S241K, S241P and S241T in SEQ ID NO: along with mutations in either: epitope 4; epitopes 4 and 3; epitopes 4 and 1; epitopes 4 and 2; or, epitopes 4 and 6 (see, Table 13). For all combinations except those including amino acid substitutions in epitope 3 at 1184 (SEQ ID NO:1; or, 1196 in SEQ ID NO:2) (which produced 0% inhibition), one or more inhibitory PE polypeptides were identified (Table 13). (Note: "Inhibitory PE polypeptides" indicates amino acid substituted forms of PE which exhibit PE biological activity in the inhibition of IVTT). From structural analysis of PE, it was noted that residue 1184 (SEQ ID NO:1; or, 1196 in SEQ ID NO:2) (anchor residue 1, Table 11) was located within the active enzymatic site of PE. In view of this result, alternative mutations distal to the active site were sought at anchor residues 6 and 9 (V189 and R192 in SEQ ID NO:1; or, V201 and R204 in SEQ ID NO:2). Alternative epitope 3 mutations at V189 and R192 in SEQ ID NO:1 were tested in combination with other epitope mutations. These mutations confirmed that inhibitory PE polypeptides with epitope 3 mutations could also be generated (Table 13, pIEX02-173 to -248). A range of combinations of DNAs encoding multiple amino acid substituted forms of PE were tested progressively leading to a final analysis of mutations in four of six, five of six, and six of six identified immunogenic epitopes. See, Table 13, "Quadruplicates", "Quintuplicates" and "Sextuplicates". In this regard, quadruplicate epitope mutations were identified which exhibited IVTT inhibitory activity ranging from 0% to about 70%. Quintuplicate mutations were identified that exhibited IVTT inhibitory activity ranging from about 5% to about 35%. Sextuplicate mutations were identified that exhibited IVTT inhibitory activity ranging from about 5% to about 20%. It is also noted that multiple single, double, and triple epitope mutations also resulted in amino acid substituted forms of PE exhibiting PE biological activity in the inhibition of IVTT such that the percent (%) inhibitor activity ranged from 0% to 100% (or about 100%) see Table 13.

[1273] Three different “candidates” (i.e., amino acid substituted forms of PE or DNA constructs encoding the same) were selected for use as examples in performing subsequent experiments described further herein. In particular, additional experiments were performed using the sextuplicate AA substituted candidate pIEX02-244 (SEQ ID NO:178; see also, Table 13); which retained approximately 20% of the WT PE inhibitor activity. Likewise, additional experiments were also performed using the sextuplicate AA substituted candidate pIEX02-246 (SEQ ID NO:179; see also, Table 13) which retained approximately 8% of the WT PE inhibitory activity; and using the quintuplicate AA substituted candidate pIEX02-228 (SEQ ID NO:177; see also, Table 13) which retained approximately 36% of the WT PE inhibitory activity. These were expressed as fusion proteins comprising a histidine polymer and a linker sequence preceding a sequence of WT PE or amino acid substituted PE; see, pIEX02-001 PE WT (SEQ ID NO:189 (DNA) and SEQ ID NO:190 (AA)), pIEX02-228 amino acid substituted PE (SEQ ID NO:193 (DNA) and SEQ ID NO:194 (AA)), pIEX02-244 amino acid substituted PE (SEQ ID NO:197 (DNA) and SEQ ID NO:198

(AA)), pIEX02-246 amino acid substituted PE (SEQ ID NO:201 (DNA) and SEQ ID NO:202 (AA)). These AA substituted PE polypeptides, and DNA constructs encoding, them may be referenced herein as “228”, “244” or “246” using simply these three numbers, or using these numbers and a prefix or suffix included therewith.

[1274] Moreover, it is noted that in view of the highly cytotoxic nature of wild-type PE, IVTT inhibition activity (i.e., cytotoxicity) as low as about 5% (or higher) of WT (e.g., 8%, 20%, and 36%) in amino acid substituted forms of PE may provide a therapeutically effective polypeptide. See, for example, Thomas et al., “Abrogation of Head and Neck Squamous Cell Carcinoma Growth by Epidermal Growth Factor Receptor Ligand Fused to *Pseudomonas* Exotoxin Transforming Growth Factor α -PE38,” *Clin. Cancer Res.* 10:7079-7087 (2004); Siegall et al., “Cell-specific toxicity of a chimeric protein composed of interleukin-6 and *Pseudomonas* exotoxin (IL6-PE40) on tumor cells”, *Mol. Cell. Biol.* 10(6): 2443-2447 (1990); and, Weldon & Pastan, “A Guide to Tam- ing a Toxin—Recombinant Immunotoxins Constructed From *Pseudomonas* Exotoxin A for the Treatment of Cancer”, *FEBS Journal* 278(23):4683-4700 (2011).

TABLE 13

Examples of Amino Acid Substituted Forms of PE and Associated Cell Cytotoxic Activity.								
Epitopes changed	pIEX02 - ###	Epitope 5	Epitope 4	Epitope 3	Epitope 1	Epitope 2	Epitope 6	% Inhibition of IVTT
Wild-Type (WT)								
	001 (WT)							100.00
Single Substitutions								
5	003	S241N [S253N]						100.00
5	004	S241K [S253K]						100.00
5	005	S241P [S253P]						99.71
5	006	S241T [S253T]						99.99
Double Substitutions								
4, 5	007	S241N [S253N]	Q194R [Q206R]					41.63
4, 5	008	S241K [S253K]	Q194R [Q206R]					69.51
4, 5	009	S241P [S253P]	Q194R [Q206R]					20.58
4, 5	010	S241T [S253T]	Q194R [Q206R]					21.44
4, 5	011	S241N [S253N]	D197K [D209K]					99.88
4, 5	012	S241K [S253K]	D197K [D209K]					42.49
4, 5	013	S241P [S253P]	D197K [D209K]					74.87
4, 5	014	S241T [S253T]	D197K [D209K]					98.84
Triple Substitutions								
3, 4, 5	015	S241N [S253N]	Q194R [Q206R]	I184A [I196A]				0.00
3, 4, 5	016	S241K [S253K]	Q194R [Q206R]	I184A [I196A]				0.00
3, 4, 5	017	S241P [S253P]	Q194R [Q206R]	I184A [I196A]				0.00
3, 4, 5	018	S241T [S253T]	Q194R [Q206R]	I184A [I196A]				0.00
3, 4, 5	019	S241N [S253N]	Q194R [Q206R]	I184N [I196N]				0.00

TABLE 13-continued

Examples of Amino Acid Substituted Forms of PE and Associated Cell Cytotoxic Activity.								
Epitopes changed	pIEX02 - ###	Epitope 5	Epitope 4	Epitope 3	Epitope 1	Epitope 2	Epitope 6	% Inhibition of IVTT
3, 4, 5	020	S241K [S253K]	Q194R [Q206R]	I184N [I196N]				0.00
3, 4, 5	021	S241P [S253P]	Q194R [Q206R]	I184N [I196N]				0.00
3, 4, 5	022	S241T [S253T]	Q194R [Q206R]	I184N [I196N]				0.00
3, 4, 5	024	S241K [S253K]	D197K [D209K]	I184A [I196A]				0.00
3, 4, 5	027	S241N [S253N]	D197K [D209K]	I184N [I196N]				0.00
3, 4, 5	028	S241K [S253K]	D197K [D209K]	I184N [I196N]				0.00
3, 4, 5	029	S241P [S253P]	D197K [D209K]	I184N [I196N]				0.00
3, 4, 5	030	S241T [S253T]	D197K [D209K]	I184N [I196N]				0.00
1, 4, 5	127	S241T [S253T]	Q194R [Q206R]		I141A [I153A]			15.58
1, 4, 5	128	S241T [S253T]	Q194R [Q206R]		I141T [I153T]			14.16
1, 4, 5	129	S241T [S253T]	Q194R [Q206R]		I141H [I153H]			56.73
1, 4, 5	130	S241T [S253T]	D197K [D209K]		I141H [I153H]			20.46
1, 4, 5	131	S241T [S253T]	D197K [D209K]		I141T [I153T]			86.84
1, 4, 5	132	S241T [S253T]	D197K [D209K]		I141A [I153A]			88.15
1, 4, 5	139	S241T [S253T]	D197K [D209K]		R146Q [R158Q]			21.81
1, 4, 5	140	S241T [S253T]	D197K [D209K]		G147S [G159S]			42.81
1, 4, 5	143	S241T [S253T]	D197K [D209K]		Q149T [Q161T]			49.33
1, 4, 5	170	S241T [S253T]	Q194R [Q206R]		R146A [R158A]			39.74
1, 4, 5	171	S241T [S253T]	Q194R [Q206R]		R146Q [R158Q]			6.01
1, 4, 5	172	S241T [S253T]	Q194R [Q206R]		G147S [G159S]			1.49
2, 4, 5	144	S241T [S253T]	D197K [D209K]			T152A [T164A]		100.67
2, 4, 5	145	S241T [S253T]	D197K [D209K]			N150A [N162A]		70.69
2, 4, 5	133	S241T [S253T]	Q194R [Q206R]			T152R [T164R]		17.46
2, 4, 5	134	S241T [S253T]	D107K [D209K]			T152R [T164R]		23.78
6, 4, 5	146	S241T [S253T]	D197K [D209K]				I321A [I333A]	104.47
6, 4, 5	147	S241T [S253T]	D197K [D209K]				I321N [I333N]	99.97
6, 4, 5	148	S241T [S253T]	D197K [D209K]				I321T [I333T]	53.19
6, 4, 5	149	S241T [S253T]	D197K [D209K]				I321Q [I333Q]	99.91
6, 4, 5	150	S241T [S253T]	D197K [D209K]				I321H [I333H]	89.43
6, 4, 5	152	S241T [S253T]	D197K [D209K]				Q326E [Q338E]	99.97
3, 4, 5	173	S241T [S253T]	Q194R [Q206R]	R192A [R204A]				23.15
3, 4, 5	174	S241T [S253T]	Q194R [Q206R]	R192Q [R204Q]				16.37
3, 4, 5	175	S241T [S253T]	Q194R [Q206R]	V189A [V201A]				6.50
Quadruplicate Substitutions								
1, 2, 4, 5	156	S241N [S253N]	Q194R [Q206R]		I141T [I153T]	T152R [T164R]		6.16
1, 2, 4, 5	166	S241N [S253N]	D197K [D209K]		I141A [I153A]	T152R [T164R]		7.90

TABLE 13-continued

Examples of Amino Acid Substituted Forms of PE and Associated Cell Cytotoxic Activity.								
Epitopes changed	pIEX02 - ###	Epitope 5	Epitope 4	Epitope 3	Epitope 1	Epitope 2	Epitope 6	% Inhibition of IVTT
1, 3, 4, 5	105	S241N [S253N]	D197K [D209K]	I184A [I196A]	I141A [I153A]			0.00
1, 3, 4, 5	106	S241K [S253K]	D197K [D209K]	I184A [I196A]	I141A [I153A]			0.00
1, 3, 4, 5	110	S241K [S253K]	D197K [D209K]	I184A [I196A]	I141T [I153T]			0.00
1, 3, 4, 5	111	S241P [S253P]	D197K [D209K]	I184A [I196A]	I141T [I153T]			0.00
1, 3, 4, 5	112	S241T [S253T]	D197K [D209K]	I184A [I196A]	I141T [I153T]			0.00
1, 3, 4, 5	114	S241K [S253K]	D197K [D209K]	I184A [I196A]	I141H [I153H]			0.00
1, 3, 4, 5	115	S241P [S253P]	D197K [D209K]	I184A [I196A]	I141H [I153H]			0.00
1, 3, 4, 5	117	S241N [S253N]	D197K [D209K]	I184N [I196N]	I141A [I153A]			0.00
1, 3, 4, 5	118	S241K [S253K]	D197K [D209K]	I184N [I196N]	I141A [I153A]			0.00
1, 3, 4, 5	120	S241T [S253T]	D197K [D209K]	I184N [I196N]	I141A [I153A]			0.00
1, 3, 4, 5	121	S241N [S253N]	D197K [D209K]	I184N [I196N]	I141T [I153T]			0.00
1, 3, 4, 5	122	S241K [S253K]	D197K [D209K]	I184N [I196N]	I141T [I153T]			0.00
1, 3, 4, 5	123	S241P [S253P]	D197K [D209K]	I184N [I196N]	I141T [I153T]			0.00
1, 3, 4, 5	124	S241T [S253T]	D197K [D209K]	I184N [I196N]	I141T [I153T]			0.00
1, 3, 4, 5	125	S241N [S253N]	D197K [D209K]	I184N [I196N]	I141H [I153H]			0.00
2, 3, 4, 5	179	S241T [S253T]	Q194R [Q206R]	V189A [V201A]		T152R [T164R]		13.37
2, 3, 4, 5	180	S241T [S253T]	Q194R [Q206R]	R192A [R204A]		T152R [T164R]		58.93
2, 3, 4, 5	181	S241T [S253T]	Q194R [Q206R]	R192Q [R204Q]		T152R [T164R]		13.70
1, 3, 4, 5	183	S241T [S253T]	D197K [D209K]	R192A [R204A]	I141A [I153A]			36.87
1, 3, 4, 5	188	S241T [S253T]	D197K [D209K]	R192A [R204A]	I141T [I153T]			20.75
1, 2, 4, 5	195	S241T [S253T]	D197K [D209K]		I141T [I153T]	T152A [T164A]		42.90
1, 4, 5, 6	200	S241T [S253T]	D197K [D209K]		I141T [I153T]		I321A [I333A]	22.04
1, 4, 5, 6	201	S241T [S253T]	D197K [D209K]		I141T [I153T]		I321N [I333N]	58.30
1, 4, 5, 6	204	S241T [S253T]	D197K [D209K]		I141T [I153T]		I321H [I333H]	12.76
1, 2, 4, 5	208	S241T [S253T]	D197K [D209K]		I141A [I153A]	T152A [T164A]		49.49
1, 4, 5, 6	213	S241T [S253T]	D197K [D209K]		I141A [I153A]		I321A [I333A]	18.03
1, 4, 5, 6	214	S241T [S253T]	D197K [D209K]		I141A [I153A]		I321N [I333N]	0.18
1, 4, 5, 6	215	S241T [S253T]	D197K [D209K]		I141A [I153A]		I321T [I333T]	5.87
1, 4, 5, 6	216	S241T [S253T]	D197K [D209K]		I141A [I153A]		I321Q [I333Q]	20.21
1, 4, 5, 6	217	S241T [S253T]	D197K [D209K]		I141A [I153A]		I321H [I333H]	11.22
1, 4, 5, 6	219	S241T [S253T]	D197K [D209K]		I141A [I153A]		Q326E [Q338E]	70.65
Quintuplicate Substitutions								
	222	S241T [S253T]	D197K [D209K]		G147S [G159S]	T152A [T164A]	Q326E [Q338E]	4.87
	224	S241T [S253T]	D197K [D209K]		Q149T [Q161T]	T152A [T164A]	Q326E [Q338E]	3.69
	226	S241T [S253T]	D197K [D209K]		I141A [I153A]	N150A [N162A]	Q326E [Q338E]	11.23
1, 2, 4, 5, 6	228	S241T [S253T]	D197K [D209K]		I141A [I153A]	T152R [T164R]	Q326E [Q338E]	36.27

TABLE 13-continued

Examples of Amino Acid Substituted Forms of PE and Associated Cell Cytotoxic Activity.								
Epitopes changed	pIEX02 - ###	Epitope 5	Epitope 4	Epitope 3	Epitope 1	Epitope 2	Epitope 6	% Inhibition of IVTT
1, 2, 4, 5, 6	229	S241T [S253T]	D197K [D209K]		I141A [I153A]	T152A [T164A]	Q326E [Q338E]	18.11
1, 2, 3, 4, 5	221	S241T [S253T]	Q194R [Q206R]	R192A [R204A]	I141A [I153A]	T152R [T164R]		4.79
1, 2, 4, 5, 6	242	S241T [S253T]	D197K [D209K]		I141T [I153T]	T152A [T164A]	Q326E [Q338E]	21.64
Sextuplicate Substitutions								
1-6	244	S241T [S253T]	D197K [D209K]	R192A [R204A]	I141T [I153T]	T152A [T164A]	Q326E [Q338E]	20.53
1-6	246	S241T [S253T]	D197K [D209K]	R192A [R204A]	I141A [I153A]	T152A [T164A]	Q326E [Q338E]	7.95
1-6	248	S241T [S253T]	D197K [D209K]	R192A [R204A]	I141A [I153A]	T152R [T164R]	Q326E [Q338E]	4.45

Note:

Non-bracketed amino acid substitution positions correspond to the PE polypeptide sequence in SEQ ID NO: 1. Bracketed [amino acid substitution positions] correspond to the PE polypeptide sequence in SEQ ID NO: 2 (comprising an N-terminal 12-amino acid linker).

Example 11

Ex Vivo Human T Cell Assay to Assess Immunogenicity of Wild-Type (WT) and Amino Acid Substituted PE

[1275] Ex vivo human T cell assays (EPISCREEN™; see e.g., preceding Examples) were performed to assess the immunogenicity of whole proteins corresponding to pIEX02-244 (SEQ ID NO:178) pIEX02-246 (SEQ ID NO:179) and pIEX02-228 (SEQ ID NO:177) (Example 10). In order to avoid direct cytotoxicity to cells used in the assay, “null mutants” were generated for the three candidates and WT PE by overlapping PCR as in Example 9 using primers OL2162 and 2163 (Table 12) to introduce an amino acid substitution of E287D in the three candidates and WT PE (to give SEQ ID NOs: 180 to 183). (Note: “Null mutants” is intended to indicate mutated forms of PE which lack cell cytotoxic biological activity; the amino acid substitution used to generate null mutants corresponds to a change of E287D in SEQ ID NO:1 or E299 in SEQ ID NO:2.) The E287D (SEQ ID NOs: 180 to 183) encoding genes were cloned into pET14b-K as in Example 9. WT PE sequence is shown in SEQ ID NO:180; pIEX02-228 sequence is shown in SEQ ID NO:181; pIEX02-244 sequence is shown in SEQ ID NO:182; and, pIEX02-246 sequence is shown in SEQ ID NO:183. These were expressed as fusion proteins comprising a histidine polymer and a linker sequence preceding a “null mutant” sequence of WT PE or amino acid substituted PE; see, pIEX02-001 PE WT null mutant (SEQ ID NO:191 (DNA) and SEQ ID NO:192 (AA)), pIEX02-228 null mutant (SEQ ID NO:195 (DNA) and SEQ ID NO:196 (AA)), pIEX02-244 null mutant (SEQ ID NO:199 (DNA) and SEQ ID NO:200 (AA)), pIEX02-246 null mutant (SEQ ID NO:203 (DNA) and SEQ ID NO:204 (AA)).

[1276] The host cell for expression of the PE E287D genes was an *Escherichia coli* BL21 derivative strain called SHuffle™ T7 Express (NEB catalog #C3029H, New England Biolabs UK Ltd., Knowl Piece, UK) which was altered to overexpress the chaperonins GroEL/ES by amplification of the GroEL/ES operon, including its promoter/regulatory sites, from *E. coli* DH5alpha™ (Invitrogen catalog #18265-017, Life Technologies Ltd., Paisley, UK.) using OL2097 (introducing EagI site) and OL2098, introducing HindIII site

(Table 12). The resulting PCR fragment was subcloned into pACYC184 (NEB catalog #E4152S) which was then transformed into SHuffle™ T7 with selection for chloramphenicol resistance. The PE E287D (SEQ ID NOs: 180 to 183) genes in pET14b-K were transformed into the SHuffle™ T7/GroEL/ES strain with selection for ampicillin resistance. Single colonies were grown in 2×YT medium (Sigma-Aldrich catalog #Y2627-1KG) and protein expression was induced at OD_{600 nm} 1.0 by adding isopropyl-β-D-thio-galactoside (IPTG) to a final concentration of 0.4 mM. Cultures were then grown at 16 degrees C. for 17 h before harvesting by centrifugation. Cell pellets were resuspended in 35 ml of binding buffer (50 mM Tris pH 8.0, 500 mM NaCl and 10 mM imidazole) supplemented with protease inhibitors (cOmplete protease inhibitor tablets, Roche catalog #11873580001, Roche Diagnostics Ltd., Burgess Hill, UK (mixture of several protease inhibitors for inhibition of serine and cysteine proteases)). Cells were lysed by sonication (SONICATOR®, Misonix catalog #XL2020, Misonix Inc., Farmingdale, N.Y.), and cell debris and insoluble material removed by centrifugation. Proteins were purified from the soluble fraction by nickel chelate affinity chromatography using HISTRAP® FF Crude columns (GE Healthcare catalog #11-004-58, GE Healthcare Life Sciences, Little Chalfont, UK). After loading, the columns were washed with 50 mM Tris (pH 8.0) containing 500 mM NaCl and 20 mM imidazole and bound protein was eluted with 50 mM Tris (pH 8.0) containing 500 mM NaCl and 500 mM imidazole. Following buffer exchange to 20 mM Tris (pH 8.0) using Zeba Spin desalting columns (7K. MWCO, Pierce catalog #89893), a negative purification step was employed using anion exchange chromatography on Q-Sepharose (1 ml, HISTRAP® Q FF column (GE Healthcare catalog #17-5053-01) with 20 mM Tris pH 8.0 and an NaCl gradient from 0 M to 1.5 M. For each protein, the column flow through was concentrated using an AMICON® Ultra centrifugal filter (EMD Millipore catalog #UFC 800 396, EMD Millipore, Feltham, UK) and further purified by size-exclusion chromatography (120 ml, HiLoad 16/60 SUPERDEX® 75 pg (GE Healthcare catalog #28-9893-33)) using 1×PBS pH 7.4 (PAA catalog #H15-002, PAA Laboratories Ltd, Yeovil, UK). For each protein, the protein peak was collected and concentrated.

[1277] Endotoxin levels were determined using an ENDOSAFE®-PTS™ Portable Test System reader (Charles River Laboratories Inc., Wilmington, Mass.) with ENDOSAFE® Licensed PTS Endotoxin cartridges (Charles River catalog #PTS20F). Endotoxins were reduced to a value below 5 endotoxin units (EU)/mg by repeatedly performing a phase separation using Triton X-114, (Aida Y. and Pabst M. J., *Journal of Immunological Methods*, 132 (1990) 191-195). Triton X-114 was removed using PIERCE® Detergent Removal Spin Columns according to the manufacturer's provided protocol (PIERCE® catalog #87779; Thermo Fisher Scientific/PIERCE® Biotechnology, Rockford, Ill.; see, Thermo Scientific instructions manual #2164.3). Protein concentration was quantified by absorbance at 280 nm using a BIOMATE™ 3 UV-Visible spectrophotometer (Thermo Fisher Scientific) and a conversion factor of $OD_{280} 1.0 = 1.5$ mg/ml derived from the calculated molar extinction coefficient of 6xHis PE (Pace C. N. et al. *Protein Science* 1995 4:2411-2423).

[1278] Ex vivo human T cell assays (EPISCREEN®) were performed using PBMC isolated from healthy community donor buffy coats as in Example 2. A cohort of 20 donors was selected to best represent the number and frequency of HLA-DR allotypes expressed in the world population. The haplotypes of the 20 donors in the assay is shown in Table 14. PBMCs from each donor were thawed, counted and viability assessed. Cells were revived in room temperature AIM-V® culture medium (INVITROGEN®, Paisley, UK.), washed and resuspended in AIM-V® to $4-6 \times 10^6$ PBMC/ml. For each donor, 1 ml of cells were dispensed into multiple wells of a 24 well plate. 0.5 ml of proteins were added at 50 micrograms/ml per sample together with 0.5 ml of AIM-V® culture medium. For each donor, a reproducibility control (cells incubated with 100 micrograms/ml keyhole limpet hemocyanin (KLH), an "intermediate" positive control (expected to give 20-30% T cell responses) of humanized A33 antibody (Welt e al. *Clinical Cancer Research*, 9 (2003) p 1338-1343) (cells were incubated with 50 micrograms/ml humanized A33), and a culture medium only control well were also included. Cultures were incubated for a total of 8 days at 37° C. with 5% CO₂.

TABLE 14

Donor Haplotypes			
Donor		KLH	
No	Haplotype	Test 1	IEX02
1	DRB1*01, DRB1*11; DRB3*	1.95	5.41
2	DRB1*11, DRB1*15; DRB3*; DRB5*	N/D	8.39
3	DRB1*04, DRB1*11; DRB3*; DRB4*	6.04	4.58
4	DRB1*08, DRB1*14; DRB3*	1.78	1.35
5	DRB1*07, DRB1*13; DRB3*; DRB4*	5.57	6.77
6	DRB1*04; DRB4*	12.36	11.25
7	DRB1*03, DRB1*04; DRB3*; DRB4*	1.48	1.12
8	DRB1*03, DRB1*13; DRB3*	2.73	1.63
9	DRB1*03, DRB1*07; DRB3*; DRB4*	3.59	3.07
10	DRB1*04, DRB1*12; DRB3*; DRB4*	3.35	3.26
11	DRB1*01, DRB1*07	13.67	15.34
12	DRB1*01, DRB1*14; DRB3*	6.05	50.13
13	DRB1*07, DRB1*09; DRB4*	9.17	19.32
14	DRB1*15; DRB5*	2.83	3.97
15	DRB1*03, DRB1*15; DRB3*; DRB5*	3.36	3.09
16	DRB1*07, DRB1*13; DRB3*; DRB4*	2.18	6.76
17	DRB1*15, DRB1*13, DRB3*; DRB5*	1.93	7.04
18	DRB1*01, DRB1*04; DRB4*	2.49	28.59

TABLE 14-continued

Donor Haplotypes			
Donor		KLH	
No	Haplotype	Test 1	IEX02
19	DRB1*01, DRB1*11; DRB3*	0.83	4.50
20	DRB1*01	2.03	3.18

[1279] For the T cell proliferation assay, on days 5, 6, 7 and 8, the cells in each well were gently resuspended and triplicate 100 microliter aliquots were transferred to each well of a round bottomed 96 well plate. The cultures were pulsed with 0.75 microCi [3H]-Thymidine (PERKIN ELMER®, Beaconsfield, UK) in 100 microliters AIM-V® culture medium and incubated for a further 18 hours before harvesting onto filter mats (Perkin Elmer®) using a TOMTEC® HARVESTER 96™ Mach III cell harvester (TOMTEC® Inc., Hamden, Conn., USA). Counts per minute (cpm) for each well were determined using MELTILEX® solid scintillator (PERKIN ELMER® Life and Analytical Sciences, Shelton, CT, USA) via scintillation counting on a Wallac 1450 Microbeta Trilux Microplate Scintillation and Luminescence Counter (Perkin Elmer®) in paralux, low background counting.

[1280] For proliferation assays, an empirical Stimulation Index (S) threshold of equal to, or greater than, 2 ($SI \geq 2.0$) was used whereby samples inducing proliferative responses above this threshold at any day after addition of protein were deemed positive. (The Stimulation Index is a ratio of stimulated proliferative response compared to a background index; an SI of 1=background or "noise".) For the triplicate proliferation data for each time point with each protein, the significance ($p < 0.05$) of positive responses was defined by statistical and empirical thresholds by comparing CPM of test protein wells against medium-only control wells using unpaired two sample Student's T-Test.

[1281] The results of the proliferation assay are shown in Table 15. The results demonstrate a significantly reduced level of T cell responses from the amino acid substituted PE molecules: pIEX02-228 (SEQ ID NO:181) 5% donor responses; pIEX02-244 (SEQ ID NO:182) 10% donor responses; and, pIEX02-246 (SEQ ID NO:183) 20% donor responses, compared to WT PE (SEQ ID NO:180)) which induced T cell responses in 70% of donors.

TABLE 15

Relative T-cell Stimulated Proliferative Responses to Amino Acid Substituted variants of PE compared to Wild-Type (WT) PE.					
	WT PE	pIEX02-228	pIEX02-244	pIEX02-246	Hu A33
Donor 1	P				
Donor 2	P*				P
Donor 3	P*		P	P	P
Donor 4					
Donor 5					
Donor 6	P				
Donor 7					
Donor 8	P				
Donor 9	P				
Donor 10	P				
Donor 11	P			P	P
Donor 12	P				P

TABLE 15-continued

Relative T-cell Stimulated Proliferative Responses to Amino Acid Substituted variants of PE compared to Wild-Type (WT) PE.					
	WT PE	pIEX02-228	pIEX02-244	pIEX02-246	Hu A33
Donor 13	P				
Donor 14	P	P		P	
Donor 15	P				
Donor 16	P				P
Donor 17	P				
Donor 18					
Donor 19			P	P	P
Donor 20					
% Donor Proliferation	70	5	10	20	30

*Positive T cell responses for proliferation (SI ≥ 2.00 , significant $p < 0.05$) during the entire time course days 5-8 ("P") are shown.

**Borderline responses (significant $p < 0.05$ with SI ≥ 1.90) are shown (*).

[1282] In addition to the proliferation assay, additional analysis of the cytokines IL-2 and IL-6 was performed using aliquots of culture supernatant taken on day 6. The analysis was performed using the BD Cytometric Bead Array (CBA) Enhanced Sensitivity Flex Set Systems for IL-2 and IL-6 (BD Bioscience, Oxford, UK) according to the manufacturer's instructions. The enhanced sensitivity standards from the CBA kit were reconstituted and serially diluted before adding 50 microliters of supernatant or standard to 20 microliters of mixed capture beads in 96 well filter plates (Millipore, Watford, UK) and incubating for 2 hours. Mixed human detection reagent (20 microliters) was then added to each well and incubated for a further 2 hours. Plates were washed twice and enhanced detection (100 microliters) reagent added to each well for a final 1 hour incubation. Plates were washed before reading on an Accuri C6 instrument (BD Biosciences).

[1283] Data was analysed using FCAP v3.0 software (BD Biosciences). For each individual donor, data was expressed as pg/ml of cytokine for each donor and plotted on a log scale with a median of cytokine levels depicted as a line. The results are shown in FIG. 11 which shows a significantly reduced level of the cytokines IL-2 and IL-6 from the amino acid substituted PE molecules pIEX02-228 (SEQ ID NO:181), pIEX02-244 (SEQ ID NO:182) and pIEX02-246 (SEQ ID NO:183) compared to WT PE (SEQ ID NO:180).

[1284] The proliferation and cytokine results both independently demonstrate that the amino acid substitutions in PE result in greatly reduced level of T cell responses when using amino acid substituted forms of PE. These results considered and expected to correlate with low or reduced PE immunogenicity in human subjects.

Example 12

Cytotoxicity Analysis of Amino Acid Substituted PE in Dendritic Cells

[1285] Amino acid substituted forms of PE and WT PE may be generated as in Example 11. For a dendritic cell cytotoxicity assay, PBMC are isolated from healthy community donor buffy coats (preferably from blood drawn within 24 hours), for example, by Lymphoprep (Axis-shield, Dundee, UK) density centrifugation. To prepare monocyte-derived dendritic cells (DC), CD14+ cells (monocytes) may be isolated from donor PBMC preparations using Miltenyi Monocyte Isolation Kit II (human) and LS columns (Miltenyi Biotech GmbH, Bergisch Gladbach, Germany; catalog #130-

091-153). Monocytes are resuspended in an appropriate culture medium, such as AIM-V® cell culture medium supplemented with 1000 IU/ml IL-4 and 1000 IU/ml GM-CSF ("DC culture medium") to $4\text{--}6 \times 10^6$ cells/ml and then distributed in 24 well plates (e.g., 2 ml final culture volume). Cells are fed on day 2 by half volume DC culture medium change. On day 3, amino acid substituted PE and WT PE proteins are added to semi-mature DC to selected final concentrations, such as 1 micrograms/ml or 10 micrograms/ml. Semi-mature DC are incubated for a period of time, such as 24-72 hours, after which cells are assessed for cytotoxicity by viability, such as via use of Trypan Blue (Sigma, Dorset, UK) dye exclusion and by propidium iodide (P) and Annexin V staining (Invitrogen, Paisley UK) followed by FACS analysis.

Example 13

Cytotoxicity Analysis of Anti-CD22 scFv Fused to Amino Acid Substituted PE in RAJI Cells

[1286] WT PE and amino acid substituted PE encoded by pIEX02-244 (SEQ ID NO:1.78) are fused to an anti-CD22 single-chain Fv (scFv). The VH and VL (V_{kappa}) regions of RFB4 (Campana D. et al., *J. Immunol.*, 134:1524-1530 (1985); Mansfield, E., et al., *Blood*, 90:2020-2026 (1997) are synthesized. RFB4 VH is amplified using oligonucleotides; OL2043 and OL2044 (Table 12). RFB4 and VL (Vk) is amplified using oligonucleotides OL2045 and OL2046 (Table 12). The RFB4 scFv is obtained using a pull-through PCR reaction using oligonucleotides OL2047 introducing a NcoI site and OL2048 introducing a XhoI site. The resultant PCR product is subcloned into pET14b-K via NcoI and XhoI restriction enzymes (Fermentas catalog #FD0573 and FD0695, respectively).

[1287] The gene encoding RFB4 scFv is fused to genes encoding either WT PE or amino acid substituted PE encoded by pIEX02-244 (SEQ ID NO:178) having a C-terminal 8xHis-tag followed by the sorting signal EDLK to give fusion sequences SEQ ID NO:184 and SEQ ID NO:186, respectively, which are cloned into the expression vector pET14b-K by fusion PCR. To create these RFB4-PE fusions a fusion PCR is carried out. The RFB4 scFv gene is amplified from pET14b-RFB4 using oligonucleotides OL2318 introducing a NcoI site and OL2320 (Table 12). The WT PE or the lead amino acid substituted PE genes are amplified from pET14b-K-WT PE or pET14b-K-244 PE, respectively, using oligonucleotides OL2321 and OL2322 introducing a N-terminal 8xHis-EDLIK and a XhoI site (Table 12). Both scFv and PE genes are fused by performing a PCR with oligonucleotides OL2318 and OL2322 (Table 12). The resulting full-length fragments are subcloned into pET14b-K using NcoI and XhoI restriction enzymes. Plasmids are transformed into B21 (DE3) *E. coli* (EMD Millipore, Feltham, UK) and clones are inoculated into 2TY+Amp and grown overnight at 37° C. Two ml of overnight culture is inoculated into 350 ml 2TY+Amp media in a 1 L flask and grown to $OD_{600\text{ nm}}=1$ before addition of IPTG (Sigma) to 1 mM (final concentration). Cultures are grown overnight at 30 degrees C. overnight and centrifuged at 10000 rpm. Bacterial pellets are frozen at -80 until ready to use.

[1288] Pellets are defrosted on ice, extracted with 10 ml B-PER® Bacterial Protein Extraction Reagent (PIERCE® Biotechnology, Rockford, Ill.; Thermo Scientific, Hemel Hempstead, UK) containing Lysozyme and DNaseI (both Thermo Scientific), and rotated for 1 hour at room tempera-

ture. Samples are then centrifuged at 10000 rpm and supernatants are discarded. Each pellet is resuspended in 5 ml B-PER containing Lysozyme and DNaseI as above and extracted for an additional 30 min at room temperature. After centrifugation, pellets are pooled and washed successively with Wash Buffer A (50 mM Tris-HCl pH 8.0, 100 mM NaCl, 1 mM EDTA, 0.5 M urea and 1.0% Triton X-100), Wash Buffer B (Buffer A but without urea), and twice with Wash Buffer C (Buffer A but without urea or Triton X-100). After final wash, insoluble pellets are stored at -80°C . cOmplete® mini-EDTA protease inhibitors (Roche Diagnostics Ltd.) are included at each step.

[1289] Pellets are resuspended in 10 ml Solubilisation Buffer (50 mM Tris-HCl pH 8.0, 100 mM NaCl, 8 M Urea and 1 mM DTT). $\text{OD}_{280\text{ nm}}$ is determined and the samples are diluted to approximately 1 mg/ml in Solubilisation Buffer. Protein samples are allowed to denature for 4 hours at room temperature and centrifuged at 10000 rpm to remove insoluble debris. 10 ml of each solubilised protein samples is injected into a pre-soaked 12 ml, 3K MWCO cut-off SLIDE-A-LYSER® Dialysis Cassette dialysis device (Thermo Scientific; PIERCE® Biotechnology, Inc., IL, USA), and dialyzed by placement overnight in a beaker containing 2.5 L, Refolding Buffer A (50 mM Tris HCl pH 8.0, 100 mM NaCl, 5 mM reduced glutathione, 1 mM oxidised glutathione, 0.1 M arginine, 4 M urea). Dialysis buffer is replaced with, in order, 2.5 L Refolding Buffer B (Buffer A with 2 M urea), 2.5 L Refolding Buffer C (Buffer A with 1 M urea) and 5 L Refolding Buffer D (Buffer A without urea or arginine) for a minimum of 4 hours at each step.

[1290] Each sample is recovered from the dialysis cassette, buffer exchanged into 50 mM 2-N-morpholino)ethanesulfonic acid (MES) pH 6.2 using PD10 desalting columns (GE Healthcare, Little Chalfont, UK) and loaded onto a 1 ml SP FF Anion Exchange column (GE Healthcare). Each column is washed with 50 mM MES pH 6.2 before eluting using a linear 0M to 1M NaCl gradient in 50 mM MES pH 6.2. Protein-containing fractions are pooled and run through a pre-equilibrated 16/60 Size Exclusion column (GE Healthcare) using 1xPBS as running buffer. Fractions containing the main protein peaks are collected, pooled and concentrated to approximately 1 ml, filter sterilized and quantified.

[1291] For cytotoxicity analysis, Raji cells (ATCC, CCL-86) are propagated in growth medium (RPMI-1640, 10% FBS, 1% Pen/Strep) and harvested during mid-log growth phase. Cells are diluted to 1×10^5 cells/ml in growth medium and 50 microliter aliquots are dispensed per well in white walled, clear bottom 96 well plates (CORNING® catalogue #3610, FISHER SCIENTIFIC®, Loughborough, UK). Each protein concentration (8x4-fold dilutions from 500 nanograms/ml) is tested in triplicate wells, and controls containing cells or growth medium only are also included. Test protein is diluted to 2x desired concentration in growth medium. 50 microliters of the test protein dilutions or controls are added to the Raji cells and plates are incubated 72 hrs in a humidified cell culture incubator (37°C , 5% CO_2). After incubation, plates are equilibrated at room temperature for 1.0 min. CELLTITER-GLO® (PROMEGA® catalogue #G7571) is prepared according to manufacturer's instructions and 100 microliters is added per well. Plates are incubated for 10 min before reading via a FLUOstar OPTIMA fluorescence plate reader (BMG Labtech Ltd., Aylesbury, UK) (also known as a fluorometer).

REFERENCES

- [1292] Aida Y. & Pabst M., *Journal of immunological Methods*, 132:191-195 (1990)
- [1293] Al-Dosari et al., *AAPS Journal*, 11 (4):671-681 (2009).
- [1294] Allen et al. (Editor), *Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems*, Lippincott Williams & Wilkins; 9th Ed. (2011).
- [1295] Andre et al., *Curr Gene Ther.*, 10(4):267-280 (2010).
- [1296] Antoniewski et al., *Mol. Cell Biol.* 14:4465 (1994).
- [1297] Ash et al. (Editor), *Handbook of Pharmaceutical Additives, Third Edition*, Synapse Information Resources, Inc.; 3rd Ed. (2007).
- [1298] Baker & Carr, *Current Drug Safety*, 5(4):1-6 (2010).
- [1299] Baker & Jones, *Curr. Opin. Drug. Disc. Dev.*, 10(2): 219-227 (2007).
- [1300] Bodles-Brakhop et al., *Mol. Ther.*, 17(4):585-592 (2009).
- [1301] Brent et al., *Cell*, 43:729-736 (1985)
- [1302] Bryson et al., *BioDrugs.*, 24(1):1-8 (2010).
- [1303] Campana D. et al., *J. Immunol.*, 134:1524-1530 (1985).
- [1304] Canadian Patent No. 2,012,311.
- [1305] Chaudhary et al., *Proc. Natl. Acad. Sci. USA*, 87:308-312 (1990).
- [1306] Cherbas et. al., *Genes Dev.* 5:120-131 (1991).
- [1307] Chester et. al., *Expert Rev. Clin. Immunol.*, 1(4): 549-559 (2006).
- [1308] Colosimo et al., *Biotechniques*, 29(2):314-8, 320-322 (2000).
- [1309] Crouch, et al., *J. Immunol. Methods*, 160(1):81-88 (1993).
- [1310] Curiel et al. *Hum. Gene Ther.* 3:147 (1992).
- [1311] D'Avino et al., *Mol. Cell. Endocrinol.* 113:1 (1995).
- [1312] Donnelly et al., *Drug Deliv.* 17(4):187-207 (2010),
- [1313] European Patent No. 234,994B1
- [1314] European Patent No. 461,809B1.
- [1315] Feigner et al., *Proc. Natl. Acad. Sci. USA*. 84:7413 (1987).
- [1316] Feigner et al., *Science* 337:387 (1989).
- [1317] Golzio et al., *Curr Gene Ther.* 10(4):256-266 (2010).
- [1318] Hansen et al., *J. Immunother.* 33(3):297-304 (2011).
- [1319] Harlow & Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 2nd ed., ISBN 978-087969314-5 (1988).
- [1320] Harlow & Lane, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, ISBN 0879695447 (1999),
- [1321] Hausman & Cooper, *The Cell: A Molecular Approach*, Washington, D.C: ASM Press, p. 51 (2004).
- [1322] Higuchi, *Using PCR to Engineer DNA*, PCR Technology: Principles and Applications for DNA Amplification, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70 (1989).
- [1323] Hochuli et al., *J Interferon Cytokine Res.* 17 Suppl 1:S15-21 (1997).
- [1324] Holgate & Baker, *Idrugs*, 12(4):233-237 (2009).
- [1325] Hu et al., *Int. J. Cancer*, 127(9):2222-2229 (2010).
- [1326] Hutchinson et al., *J. Biol. Chem.* 255:6551 (1978).
- [1327] Hutchinson et al., *Proc. Natl. Acad. Sci. USA* 83:710 (1986).
- [1328] Hwang et al. *Cell* 48:129-136 (1987).

- [1329] Itoi et al., *J. Neurosci.* 31(16):6132-6139 (2011).
- [1330] Jaber & Baker, *J. Pharm. Biomed. Anal.* 43(4):1256-61 (2007).
- [1331] Karzenowski et al., *BioTechniques* 39: 191-200 (2005).
- [1332] Klein et al., *Curr. Opin. Biotechnol.* 4(5):583-590 (1993).
- [1333] Kreitman et. al., *Leak. Lymphoma Suppl.* 2:82-86 (2011).
- [1334] Krieg et al., *Nucl. Acids Res.* 12, 7057-7070 (1984).
- [1335] Kuan et al., *Int. J. Cancer* 129(1): 1.1.1-21 (2011).
- [1336] Kyte & Doolittle, *J. Mol. Biol.* 157(1):105-132 (1982).
- [1337] Lim, et al., *Hematology* 10(3):255-9 (2005).
- [1338] Mackey et al., *Proc. Natl. Acad. Sci USA* 85:8027 (1988).
- [1339] Mansfield, E., et al., *Blood.* 90:2020-2026 (1997).
- [1340] Mareeva et al., *J. Immunol. Methods* 353(1-2):78-86 (2010).
- [1341] Miller et al., *Somat. Cell Mol. Genet.* 27(1-6): 115-34 (2002).
- [1342] Nagata et al., *Adv. Drug Deliv. Rev.* 61(11):977-985 (2009).
- [1343] Namaka, et al., *Curr Med Res Opin.* 22(2):223-39 (2006).
- [1344] Niles, et al., *Anal. Biochem.*, 366, 197-206 (2007).
- [1345] O'Neil, "The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals," 14th Ed. (2006).
- [1346] Oliphant et al., *Gene* 44:177 (1986).
- [1347] Olsson et al., *J. Appl. Biochem* 5, 347-445 (1983).
- [1348] Onda et al., *Proc. Natl. Acad. Sci. USA* 105(32): 11311-11316 (2008).
- [1349] Onda et al., *Proc. Natl. Acad. Sci. USA* 108(14): 5742-5747 (2011).
- [1350] Pastan et al., *Leukemia and Lymphoma* 52(S2):87-90 (2011).
- [1351] Pastan et al., *Science* 254:1173-1173-1177 (1991).
- [1352] Pathak et al., *Biotechnol. J.* 4(11): 1559-1572 (2009).
- [1353] PDR Network, *Physicians' Desk Reference* 2011," PDR Network (2010).
- [1354] PDR Network, *Physicians' Desk Reference* 2012," PDR Network (2011).
- [1355] Pelham et al., *Eur. J. Biochem.* 67, 247-56 (1976).
- [1356] Perry et al., *Drugs R D.* 9(6):385-96 (2008).
- [1357] Pichon et al., *Curr Opin Biotechnol.* 21 (5):640-645 (2010).
- [1358] PROMEGA® Technical Bulletin #TB126 (Revised December 2010).
- [1359] PROMEGA® Technical Bulletin #TB288 (Revised June 2009).
- [1360] PROMEGA® Technical Bulletin #TB288 (Revised August 2011).
- [1361] PROMEGA® Technical Bulletin #TB306 (Revived May 2009).
- [1362] PROMEGA® Technical Bulletin #TB359 (Revived May 2009).
- [1363] PROMEGA® Technical Bulletin #TB359 (Revised October 2011).
- [1364] PROMEGA® Technical Bulletin #TM045 (Revised May 2011).
- [1365] PROMEGA® Technical Bulletin #TM051 (Revised March 2009).
- [1366] PROMEGA® Technical Bulletin #TM051 (Revised September 2011).
- [1367] Rochlitz et al., "Gene therapy of cancer," *Swiss Med. Wkly.*, 131(1-2):4-9 (2001).
- [1368] Rowe et al., (Editor), "Handbook of Pharmaceutical Excipients," Pharmaceutical Press, 6th Ed. (2009).
- [1369] Sadowski et al., *Nature* 335:563 (1988).
- [1370] Schellekens et al., *J. Interferon Cytokine Res.* 17(Suppl. 1), S5-S8 (1997).
- [1371] Seetharam et al., *Jour. Biol. Chem.* 266:17376-17381 (1991).
- [1372] Shapira et al., *Gastroenterology* 140(3):935-946 (2011).
- [1373] Siegall et al., *Mol. Cell. Biol.* 10(6):2443-2447 (1990).
- [1374] Siegall et al., *Biochemistry*, 30:7154-7159 (1991).
- [1375] Squirrell et al., *A Practical Guide to Industrial Uses of ATP Luminescence in Rapid Microbiology*, Cara Technology Ltd., Lingfield, pp. 107-113 (1997).
- [1376] Stish et al., *Br. J. Cancer* 101(7): 114-1123 (2009).
- [1377] Theuer et al., *J. Biol. Chem.*, 267(24):16872-16877 (1992).
- [1378] Thomas et al., *Clin. Cancer Res.* 10:7079-7087 (2004).
- [1379] U.S. Patent Application Publication No. 2005/0228016.
- [1380] U.S. Patent Application Publication No. 2006/0100416.
- [1381] U.S. Patent Application Publication No. 2009/0123441.
- [1382] U.S. Patent Application Publication No. 2009/0142341.
- [1383] U.S. Patent Application Publication. No. 2009/0298175.
- [1384] U.S. Pat. No. 4,892,827.
- [1385] U.S. Pat. No. 4,985,461.
- [1386] U.S. Pat. No. 5,117,057.
- [1387] U.S. Pat. No. 5,225,443.
- [1388] U.S. Pat. No. 5,324,637.
- [1389] U.S. Pat. No. 5,378,726.
- [1390] U.S. Pat. No. 5,459,127.
- [1391] U.S. Pat. No. 5,492,817.
- [1392] U.S. Pat. No. 5,530,028.
- [1393] U.S. Pat. No. 5,580,859.
- [1394] U.S. Pat. No. 5,583,024.
- [1395] U.S. Pat. No. 5,589,466.
- [1396] U.S. Pat. No. 5,641,641 .
- [1397] U.S. Pat. No. 5,650,289 .
- [1398] U.S. Pat. No. 5,674,713 .
- [1399] U.S. Pat. No. 5,693,622.
- [1400] U.S. Pat. No. 5,700,673.
- [1401] U.S. Pat. No. 5,821,238.
- [1402] U.S. Pat. No. 6,013,836.
- [1403] U.S. Pat. No. 6,258,603.
- [1404] U.S. Pat. No. 6,602,677.
- [1405] U.S. Pat. No. 6,982,152.
- [1406] U.S. Pat. No. 7,083,911,
- [1407] U.S. Pat. No. 7,241,584.
- [1408] U.S. Pat. No. 7,282,348.
- [1409] U.S. Pat. No. 7,304,161.
- [1410] U.S. Pat. No. 7,304,162.
- [1411] U.S. Pat. No. 7,375,093
- [1412] U.S. Pat. No. 7,452,663.
- [1413] U.S. Pat. No. 7,456,315.

- [1414] U.S. Pat. No. 7,531,326.
 [1415] U.S. Pat. No. 7,563,879.
 [1416] U.S. Pat. No. 7,700,310.
 [1417] U.S. Pat. No. 7,732,128.
 [1418] U.S. Pat. No. 7,741,067.
 [1419] U.S. Pat. No. 7,750,136.
 [1420] U.S. Pat. No. 7,919,269.
 [1421] U.S. Pat. No. 7,935,510.
 [1422] U.S. Pat. No. 8,076,517.
 [1423] U.S. Pat. No. 8,105,825.
 [1424] U.S. Pat. No. 8,854,044.
 [1425] Ulmer et al., *Science* 259:1745 (1.993).
 [1426] University of the Sciences in Philadelphia (Editor),
 "Remington: The Science and Practice of Pharmacy," Lip-
 pincott Williams & Wilkins, 21st Ed. (2005).
 [1427] Uskoković et al., *J. Biomed. Mater. Res. B App.*
Biomater 96(1):152-191 (2011).
 [1428] Wells, *Cell Biol Toxicol.* 26(1):21-28 (2010).
 [1429] Weldon & Pastan, *FEBS Journal* 278(23):4683-
 4700 (2011)
 [1430] WO 1995/018863 (PCT/FR1995/000022).
 [1431] WO 1995/021931 (PCT/FR1995/000098).
 [1432] WO 1996/017823 (PCT/FR1995/001595).
 [1433] WO 1996/025508 (PCT/FR1996/000248).
 [1434] WO 2001/070816 (PCT/US2001/090500).
 [1435] WO 2002/029075 (PCT/US2001/030608).
 [1436] WO 2002/066612 (PCT/US2002/005090).
 [1437] WO 2002/066613 (PCT/US2002/005090).
 [1438] WO 2002/066614 (PCT/US/2002/005706).
 [1439] WO 2002/066615 (PCT/US2002/005708).
 [1440] WO 2003/027266 (PCT/US/2002/05234).
 [1441] WO 2003/027289 (PCT/US2002/005026).
 [1442] WO 2005/108617 (PCT/US2005/015089).
 [1443] WO 2008/073154 (PCT/US2007/0116747).
 [1444] WO 2008/153801 (PCT/US2008/006757).
 [1445] WO 2009/025866 (PCT/US 2008/010040).
 [1446] WO 2009/045370 (PCT/US2008/011270).
 [1447] WO 2009/048560 (PCT/US2008/011563).
 [1448] WO 2010/042189 (PCT/US2009/005510).
 [1449] WO 2011/119773 (PCT/US2011/029682).
 [1450] Wu et al., *J. Biol. Chem.* 263:14621 (1988).
 [1451] Wu et al., *J. Biol. Chem.* 267:963 (1992).
 [1452] Wu et al., *J. Biol. Chem.* 262:4429 (1987).
 [1453] Xiong et al., *Pharmazie* 66(3):158-64 (2011).
 [1454] Yao et al., *Cell* 71:63 (1992).
 [1455] Yao et al., *Nature* 366:476 (1993).
 [1456] Zielinski et al., *J Immunother.* 32(8):817-825 (Oc-
 tober-2009).
 [1457] Zoller et al., *DNA* 3:479 (1984).

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 286

<210> SEQ ID NO 1

<211> LENGTH: 347

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(347)

<223> OTHER INFORMATION: *Pseudomonas aeruginosa* PE38

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (3)..(114)

<223> OTHER INFORMATION: Domain II (cytosolic translocation domain)

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (115)..(133)

<223> OTHER INFORMATION: Partial Domain IB

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (134)..(347)

<223> OTHER INFORMATION: Domain III (cytotoxic domain)

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (343)..(347)

<223> OTHER INFORMATION: Alternative carboxy-terminal tail (amino acids..REDLK)

<400> SEQUENCE: 1

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
 1 5 10 15

Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
 20 25 30

Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
 35 40 45

Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
 50 55 60

-continued

Ala	Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	65	70	75	80
Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	85	90	95	
Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	100	105	110	
Ala	Ser	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	115	120	125	
Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	130	135	140	
Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	145	150	155	160
Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	165	170	175	
Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	180	185	190	
Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	195	200	205	
Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	210	215	220	
Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	225	230	235	240
Ser	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu	245	250	255	
Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	260	265	270	
Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	275	280	285	
Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	290	295	300	
Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	305	310	315	320
Ile	Pro	Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	325	330	335	
Gln	Pro	Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys						340	345		

<210> SEQ ID NO 2
 <211> LENGTH: 359
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Gly-Ser-PE38 polypeptide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(12)
 <223> OTHER INFORMATION: artificial linker sequence
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (13)..(359)
 <223> OTHER INFORMATION: Pseudomonas aeruginosa PE38
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (15)..(126)
 <223> OTHER INFORMATION: Domain II (cytosolic translocation domain)
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (127)..(145)
 <223> OTHER INFORMATION: Partial Domain IB

-continued

```

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (146)..(359)
<223> OTHER INFORMATION: Domain III (cytotoxic domain)
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (355)..(359)
<223> OTHER INFORMATION: Alternative carboxy-terminal tail (amino
acids..REDLK)

<400> SEQUENCE: 2

Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Gly Ser Pro Glu Gly Gly
1          5          10          15
Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro Leu Glu
20          25          30
Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu Glu Gln
35          40          45
Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala Ala Arg
50          55          60
Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu Ala Ser
65          70          75          80
Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln Pro Glu
85          90          95
Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu Arg Phe
100         105         110
Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Ser Gly Pro
115         120         125
Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr Gly Ala
130         135         140
Glu Phe Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser Thr Arg Gly Thr
145         150         155         160
Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu
165         170         175
Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu Glu Ala
180         185         190
Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu
195         200         205
Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala
210         215         220
Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg
225         230         235         240
Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly
245         250         255
Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu
260         265         270
Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile
275         280         285
Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp
290         295         300
Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp
305         310         315         320
Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys
325         330         335
Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys
340         345         350

```

-continued

Pro Pro Arg Glu Asp Leu Lys
355

<210> SEQ ID NO 3
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 2x((Gx5)S) linker sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: artificial linker sequence

<400> SEQUENCE: 3

Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Gly Ser
1 5 10

<210> SEQ ID NO 4
<211> LENGTH: 347
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(347)
<223> OTHER INFORMATION: Variant of Pseudomonas aeruginosa PE38 in
SEQ ID NO:1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (114)..(114)
<223> OTHER INFORMATION: Ser-to-Asn change compared to SEQ ID NO:1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (141)..(141)
<223> OTHER INFORMATION: Ile-to-Val change compared to SEQ ID NO:1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (249)..(249)
<223> OTHER INFORMATION: Gly-to-Ser change compared to SEQ ID NO:1

<400> SEQUENCE: 4

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1 5 10 15
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20 25 30
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35 40 45
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50 55 60
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65 70 75 80
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
85 90 95
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100 105 110
Ala Asn Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
115 120 125
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser
130 135 140
Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
145 150 155 160
Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr

-continued

165					170					175						
Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	
180					185					190						
Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	
195					200					205						
Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	
210					215					220						
Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	
225					230					235					240	
Ser	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	
245					250					255						
Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	
260					265					270						
Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr		
275					280					285						
Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	
290					295					300						
Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	
305					310					315					320	
Ile	Pro	Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	
325					330					335						
Gln	Pro	Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys						
340					345											

<210> SEQ ID NO 5
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(9)
 <223> OTHER INFORMATION: Epitope 1

<400> SEQUENCE: 5

Ile Ser Phe Ser Thr Arg Gly Thr Gln
 1 5

<210> SEQ ID NO 6
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(9)
 <223> OTHER INFORMATION: Epitope 2

<400> SEQUENCE: 6

Gly Thr Gln Asn Trp Thr Val Glu Arg
 1 5

<210> SEQ ID NO 7
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(9)
 <223> OTHER INFORMATION: Epitope 3

<400> SEQUENCE: 7

-continued

Ile Val Phe Gly Gly Val Arg Ala Arg
1 5

<210> SEQ ID NO 8
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(9)
<223> OTHER INFORMATION: Epitope 4

<400> SEQUENCE: 8

Ala Arg Ser Gln Asp Leu Asp Ala Ile
1 5

<210> SEQ ID NO 9
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(9)
<223> OTHER INFORMATION: Epitope 5

<400> SEQUENCE: 9

Leu Arg Val Tyr Val Pro Arg Ser Ser
1 5

<210> SEQ ID NO 10
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(9)
<223> OTHER INFORMATION: Epitope 6

<400> SEQUENCE: 10

Ile Pro Asp Lys Glu Gln Ala Ile Ser
1 5

<210> SEQ ID NO 11
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Gly-Ser linker plus PE38 sequence
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 1

<400> SEQUENCE: 11

Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Gly Ser Pro Glu Gly
1 5 10 15

<210> SEQ ID NO 12
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Gly-Ser linker plus PE38 sequence
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 2

-continued

<400> SEQUENCE: 12

Gly Gly Ser Gly Gly Gly Gly Gly Ser Pro Glu Gly Gly Ser Leu
1 5 10 15

<210> SEQ ID NO 13

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Gly-Ser linker plus PE38 sequence

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 3

<400> SEQUENCE: 13

Gly Gly Gly Gly Gly Ser Pro Glu Gly Gly Ser Leu Ala Ala Leu
1 5 10 15

<210> SEQ ID NO 14

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Gly-Ser linker plus PE38 sequence

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 4

<400> SEQUENCE: 14

Gly Gly Ser Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His
1 5 10 15

<210> SEQ ID NO 15

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 5

<400> SEQUENCE: 15

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys
1 5 10 15

<210> SEQ ID NO 16

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 6

<400> SEQUENCE: 16

Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro
1 5 10 15

<210> SEQ ID NO 17

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

-continued

<223> OTHER INFORMATION: Peptide 7

<400> SEQUENCE: 17

Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr
1				5						10				15

<210> SEQ ID NO 18

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 8

<400> SEQUENCE: 18

Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe	Thr	Arg
1				5					10					15

<210> SEQ ID NO 19

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 9

<400> SEQUENCE: 19

Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln
1				5					10					15

<210> SEQ ID NO 20

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 10

<400> SEQUENCE: 20

His	Leu	Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly
1				5					10					15

<210> SEQ ID NO 21

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 11

<400> SEQUENCE: 21

Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln
1				5					10					15

<210> SEQ ID NO 22

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 12

-continued

<400> SEQUENCE: 22

Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu Glu Gln
1 5 10 15

<210> SEQ ID NO 23

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 13

<400> SEQUENCE: 23

His Arg Gln Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr
1 5 10 15

<210> SEQ ID NO 24

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 14

<400> SEQUENCE: 24

Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln
1 5 10 15

<210> SEQ ID NO 25

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 15

<400> SEQUENCE: 25

Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val
1 5 10 15

<210> SEQ ID NO 26

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 16

<400> SEQUENCE: 26

Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
1 5 10 15

<210> SEQ ID NO 27

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 17

<400> SEQUENCE: 27

-continued

Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala Ala
1 5 10 15

<210> SEQ ID NO 28
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 18

<400> SEQUENCE: 28

Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser
1 5 10 15

<210> SEQ ID NO 29
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 19

<400> SEQUENCE: 29

Arg Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln
1 5 10 15

<210> SEQ ID NO 30
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 20

<400> SEQUENCE: 30

Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln
1 5 10 15

<210> SEQ ID NO 31
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 21

<400> SEQUENCE: 31

Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg
1 5 10 15

<210> SEQ ID NO 32
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 22

<400> SEQUENCE: 32

Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu
1 5 10 15

-continued

<210> SEQ ID NO 33
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 23

<400> SEQUENCE: 33

Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu Ala Ser Pro
1 5 10 15

<210> SEQ ID NO 34
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 24

<400> SEQUENCE: 34

Val Asp Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly
1 5 10 15

<210> SEQ ID NO 35
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 25

<400> SEQUENCE: 35

Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
1 5 10 15

<210> SEQ ID NO 36
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 26

<400> SEQUENCE: 36

Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala
1 5 10 15

<210> SEQ ID NO 37
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 27

<400> SEQUENCE: 37

Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu
1 5 10 15

-continued

<210> SEQ ID NO 38
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 28

<400> SEQUENCE: 38

Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln Pro Glu
1 5 10 15

<210> SEQ ID NO 39
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 29

<400> SEQUENCE: 39

Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg
1 5 10 15

<210> SEQ ID NO 40
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 30

<400> SEQUENCE: 40

Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu
1 5 10 15

<210> SEQ ID NO 41
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 31

<400> SEQUENCE: 41

Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala
1 5 10 15

<210> SEQ ID NO 42
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 32

<400> SEQUENCE: 42

Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Glu
1 5 10 15

<210> SEQ ID NO 43
<211> LENGTH: 15

-continued

<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 33

<400> SEQUENCE: 43

Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu Arg
1 5 10 15

<210> SEQ ID NO 44
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 34

<400> SEQUENCE: 44

Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg
1 5 10 15

<210> SEQ ID NO 45
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 35

<400> SEQUENCE: 45

Thr Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr
1 5 10 15

<210> SEQ ID NO 46
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 36

<400> SEQUENCE: 46

Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp
1 5 10 15

<210> SEQ ID NO 47
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 37

<400> SEQUENCE: 47

Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly
1 5 10 15

<210> SEQ ID NO 48
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa

-continued

<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 38

<400> SEQUENCE: 48

Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Ser
1 5 10 15

<210> SEQ ID NO 49
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 39

<400> SEQUENCE: 49

Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala
1 5 10 15

<210> SEQ ID NO 50
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 40

<400> SEQUENCE: 50

Gly Asn Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly
1 5 10 15

<210> SEQ ID NO 51
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 41

<400> SEQUENCE: 51

Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
1 5 10 15

<210> SEQ ID NO 52
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 42

<400> SEQUENCE: 52

Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg
1 5 10 15

<210> SEQ ID NO 53
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE

-continued

<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 43

<400> SEQUENCE: 53

Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro
1 5 10 15

<210> SEQ ID NO 54
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 44

<400> SEQUENCE: 54

Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr Gly Ala
1 5 10 15

<210> SEQ ID NO 55
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 45

<400> SEQUENCE: 55

Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu
1 5 10 15

<210> SEQ ID NO 56
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 46

<400> SEQUENCE: 56

Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly
1 5 10 15

<210> SEQ ID NO 57
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 47

<400> SEQUENCE: 57

Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ile
1 5 10 15

<210> SEQ ID NO 58
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 48

-continued

<400> SEQUENCE: 58

Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser
1			5					10					15	

<210> SEQ ID NO 59

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 49

<400> SEQUENCE: 59

Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly
1			5					10					15	

<210> SEQ ID NO 60

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 50

<400> SEQUENCE: 60

Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
1			5					10					15	

<210> SEQ ID NO 61

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 51

<400> SEQUENCE: 61

Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val
1			5					10					15	

<210> SEQ ID NO 62

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 52

<400> SEQUENCE: 62

Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu
1			5					10					15	

<210> SEQ ID NO 63

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 53

<400> SEQUENCE: 63

-continued

Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala
1 5 10 15

<210> SEQ ID NO 64
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 54

<400> SEQUENCE: 64

Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln
1 5 10 15

<210> SEQ ID NO 65
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 55

<400> SEQUENCE: 65

Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu
1 5 10 15

<210> SEQ ID NO 66
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 56

<400> SEQUENCE: 66

Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr
1 5 10 15

<210> SEQ ID NO 67
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 57

<400> SEQUENCE: 67

Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
1 5 10 15

<210> SEQ ID NO 68
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 58

<400> SEQUENCE: 68

His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His

-continued

1	5	10	15
---	---	----	----

<210> SEQ ID NO 69
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 59

<400> SEQUENCE: 69

Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe
1				5					10				15	

<210> SEQ ID NO 70
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 60

<400> SEQUENCE: 70

Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala
1				5					10				15	

<210> SEQ ID NO 71
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 61

<400> SEQUENCE: 71

Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser
1				5					10				15	

<210> SEQ ID NO 72
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 62

<400> SEQUENCE: 72

Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe
1				5					10				15	

<210> SEQ ID NO 73
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 63

<400> SEQUENCE: 73

Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val
1				5					10				15	

-continued

<210> SEQ ID NO 74
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 64

<400> SEQUENCE: 74

Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
1 5 10 15

<210> SEQ ID NO 75
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 65

<400> SEQUENCE: 75

Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp
1 5 10 15

<210> SEQ ID NO 76
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 66

<400> SEQUENCE: 76

Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala
1 5 10 15

<210> SEQ ID NO 77
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 67

<400> SEQUENCE: 77

Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg
1 5 10 15

<210> SEQ ID NO 78
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 68

<400> SEQUENCE: 78

Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr
1 5 10 15

<210> SEQ ID NO 79

-continued

<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 69

<400> SEQUENCE: 79

Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly
1 5 10 15

<210> SEQ ID NO 80
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 70

<400> SEQUENCE: 80

Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala
1 5 10 15

<210> SEQ ID NO 81
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 71

<400> SEQUENCE: 81

Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr
1 5 10 15

<210> SEQ ID NO 82
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 72

<400> SEQUENCE: 82

Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala
1 5 10 15

<210> SEQ ID NO 83
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 73

<400> SEQUENCE: 83

Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
1 5 10 15

<210> SEQ ID NO 84
<211> LENGTH: 15
<212> TYPE: PRT

-continued

<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 74

<400> SEQUENCE: 84

Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp
1 5 10 15

<210> SEQ ID NO 85
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 75

<400> SEQUENCE: 85

Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly
1 5 10 15

<210> SEQ ID NO 86
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 76

<400> SEQUENCE: 86

Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg
1 5 10 15

<210> SEQ ID NO 87
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 77

<400> SEQUENCE: 87

Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala
1 5 10 15

<210> SEQ ID NO 88
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 78

<400> SEQUENCE: 88

Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg
1 5 10 15

<210> SEQ ID NO 89
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:

-continued

<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 79

<400> SEQUENCE: 89

Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val
1 5 10 15

<210> SEQ ID NO 90
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 80

<400> SEQUENCE: 90

Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
1 5 10 15

<210> SEQ ID NO 91
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 81

<400> SEQUENCE: 91

Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro
1 5 10 15

<210> SEQ ID NO 92
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 82

<400> SEQUENCE: 92

Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr
1 5 10 15

<210> SEQ ID NO 93
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 83

<400> SEQUENCE: 93

Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly
1 5 10 15

<210> SEQ ID NO 94
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)

-continued

<223> OTHER INFORMATION: Peptide 84

<400> SEQUENCE: 94

Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu
1 5 10 15

<210> SEQ ID NO 95

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 85

<400> SEQUENCE: 95

Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro
1 5 10 15

<210> SEQ ID NO 96

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 86

<400> SEQUENCE: 96

Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala
1 5 10 15

<210> SEQ ID NO 97

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 87

<400> SEQUENCE: 97

Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val
1 5 10 15

<210> SEQ ID NO 98

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 88

<400> SEQUENCE: 98

Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu
1 5 10 15

<210> SEQ ID NO 99

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 89

-continued

<400> SEQUENCE: 99

Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His
1 5 10 15

<210> SEQ ID NO 100

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 90

<400> SEQUENCE: 100

Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro
1 5 10 15

<210> SEQ ID NO 101

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 91

<400> SEQUENCE: 101

Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu
1 5 10 15

<210> SEQ ID NO 102

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 92

<400> SEQUENCE: 102

Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile
1 5 10 15

<210> SEQ ID NO 103

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 93

<400> SEQUENCE: 103

Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro
1 5 10 15

<210> SEQ ID NO 104

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: PEPTIDE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Peptide 94

<400> SEQUENCE: 104

-continued

Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu
1 5 10 15

<210> SEQ ID NO 105
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 95

<400> SEQUENCE: 105

Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg
1 5 10 15

<210> SEQ ID NO 106
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 96

<400> SEQUENCE: 106

Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr
1 5 10 15

<210> SEQ ID NO 107
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 97

<400> SEQUENCE: 107

Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly
1 5 10 15

<210> SEQ ID NO 108
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 98

<400> SEQUENCE: 108

Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu
1 5 10 15

<210> SEQ ID NO 109
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 99

<400> SEQUENCE: 109

Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg
1 5 10 15

-continued

<210> SEQ ID NO 110
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 100

<400> SEQUENCE: 110

Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val
1 5 10 15

<210> SEQ ID NO 111
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 101

<400> SEQUENCE: 111

Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser
1 5 10 15

<210> SEQ ID NO 112
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 102

<400> SEQUENCE: 112

Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro
1 5 10 15

<210> SEQ ID NO 113
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 103

<400> SEQUENCE: 113

Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro
1 5 10 15

<210> SEQ ID NO 114
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 104

<400> SEQUENCE: 114

Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val
1 5 10 15

-continued

<210> SEQ ID NO 115
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 105

<400> SEQUENCE: 115

Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp
1 5 10 15

<210> SEQ ID NO 116
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 106

<400> SEQUENCE: 116

Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro
1 5 10 15

<210> SEQ ID NO 117
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 107

<400> SEQUENCE: 117

Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile
1 5 10 15

<210> SEQ ID NO 118
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 108

<400> SEQUENCE: 118

Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys
1 5 10 15

<210> SEQ ID NO 119
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 109

<400> SEQUENCE: 119

Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala
1 5 10 15

<210> SEQ ID NO 120
<211> LENGTH: 15

-continued

<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 110

<400> SEQUENCE: 120

Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala
1 5 10 15

<210> SEQ ID NO 121
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 111

<400> SEQUENCE: 121

Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp
1 5 10 15

<210> SEQ ID NO 122
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 112

<400> SEQUENCE: 122

Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
1 5 10 15

<210> SEQ ID NO 123
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 113

<400> SEQUENCE: 123

Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly
1 5 10 15

<210> SEQ ID NO 124
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 114

<400> SEQUENCE: 124

Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
1 5 10 15

<210> SEQ ID NO 125
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa

-continued

<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 115

<400> SEQUENCE: 125

Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp
1 5 10 15

<210> SEQ ID NO 126
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(14)
<223> OTHER INFORMATION: Peptide 116

<400> SEQUENCE: 126

Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
1 5 10

<210> SEQ ID NO 127
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 117

<400> SEQUENCE: 127

Ile Thr Gly Pro Glu Glu Gly Gly Arg Leu Asp Thr Ile Leu
1 5 10 15

<210> SEQ ID NO 128
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 118

<400> SEQUENCE: 128

Pro Glu Glu Glu Gly Gly Arg Leu Asp Thr Ile Leu Gly Trp Pro
1 5 10 15

<210> SEQ ID NO 129
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 119

<400> SEQUENCE: 129

Glu Gly Gly Arg Leu Asp Thr Ile Leu Gly Trp Pro Leu Ala Glu
1 5 10 15

<210> SEQ ID NO 130
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE

-continued

```

<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Peptide 120

<400> SEQUENCE: 130

Arg Leu Asp Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val
1           5           10           15

<210> SEQ ID NO 131
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Hypothetical Peptide 121

<400> SEQUENCE: 131

Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg
1           5           10           15

<210> SEQ ID NO 132
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Hypothetical Peptide 122

<400> SEQUENCE: 132

Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile
1           5           10           15

<210> SEQ ID NO 133
<211> LENGTH: 613
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(613)
<223> OTHER INFORMATION: Pseudomonas Exotoxin A variant

<400> SEQUENCE: 133

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
1           5           10           15

Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
20          25          30

Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
35          40          45

Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
50          55          60

Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
65          70          75          80

Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
85          90          95

Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
100         105         110

Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
115         120         125

Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
130         135         140

```

-continued

Leu	Ala	Arg	Asp	Ala	Thr	Phe	Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu
145					150					155					160
Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met
				165					170					175	
Ala	Gln	Thr	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser
			180					185					190		
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr
	195						200					205			
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile
	210					215					220				
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys
225					230					235					240
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu
				245					250					255	
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
			260					265					270		
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly
		275					280					285			
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
	290					295					300				
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
305					310					315					320
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
				325					330					335	
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
			340					345					350		
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Ala	Asp	Val	Val
		355					360					365			
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp
	370					375					380				
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
385					390					395					400
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
			405						410					415	
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg
			420					425					430		
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln
		435					440					445			
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala
	450					455					460				
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly
465					470					475					480
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly
				485					490					495	
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr
			500					505					510		
Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu
		515					520					525			
Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly
	530					535					540				
Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu
545					550					555					560

-continued

Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg
 565 570 575

Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln
 580 585 590

Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
 595 600 605

Arg Glu Asp Leu Lys
 610

<210> SEQ ID NO 134
 <211> LENGTH: 649
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(649)
 <223> OTHER INFORMATION: Pseudomonas Exotoxin A variant

<400> SEQUENCE: 134

Asn Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala Ala Gly Glu Cys
 1 5 10 15

Ala Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro
 20 25 30

Thr Gly Ala Glu Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala
 35 40 45

Lys Ala Cys Val Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met
 50 55 60

Ser Val Asp Pro Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His
 65 70 75 80

Tyr Ser Met Val Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile
 85 90 95

Asp Asn Ala Leu Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu
 100 105 110

Gly Gly Val Glu Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln
 115 120 125

Ala Arg Gly Ser Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu
 130 135 140

Lys Pro Ser Asn Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn
 145 150 155 160

Gln Leu Ser His Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu
 165 170 175

Leu Leu Ala Lys Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His
 180 185 190

Glu Ser Asn Glu Met Gln Pro Thr Leu Ala Ile Ser His Ala Gly Val
 195 200 205

Ser Val Val Met Ala Gln Thr Gln Pro Arg Arg Glu Lys Arg Trp Ser
 210 215 220

Glu Trp Ala Ser Gly Lys Val Leu Cys Leu Leu Asp Pro Leu Asp Gly
 225 230 235 240

Val Tyr Asn Tyr Leu Ala Gln Gln Arg Cys Asn Leu Asp Asp Thr Trp
 245 250 255

Glu Gly Lys Ile Tyr Arg Val Leu Ala Gly Asn Pro Ala Lys His Asp
 260 265 270

Leu Asp Ile Lys Pro Thr Val Ile Ser His Arg Leu His Phe Pro Glu

-continued

275					280					285					
Gly	Gly	Ser	Leu	Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro
290					295					300					
Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu
305					310					315					320
Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala
			325						330					335	
Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu
		340						345					350		
Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln
		355					360					365			
Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu
	370					375					380				
Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Ser
385					390					395					400
Ala	Asp	Val	Val	Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala
			405						410					415	
Gly	Pro	Ala	Asp	Asn	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr
		420					425						430		
Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg
	435					440						445			
Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln
	450					455					460				
Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu
465					470					475					480
Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln
			485						490					495	
Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala
		500					505						510		
Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg
		515					520					525			
Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu
	530					535					540				
Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala
545					550					555					560
Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp
			565						570					575	
Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu
		580						585					590		
Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro
	595					600						605			
Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro
	610					615					620				
Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro
625					630					635					640
Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys							
			645												

<210> SEQ ID NO 135

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

-continued

<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(5)
<223> OTHER INFORMATION: Alternative carboxy terminal tail

<400> SEQUENCE: 135

Arg Glu Asp Leu Lys
1 5

<210> SEQ ID NO 136
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(4)
<223> OTHER INFORMATION: Alternative carboxy terminal tail

<400> SEQUENCE: 136

Arg Glu Asp Leu
1

<210> SEQ ID NO 137
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(4)
<223> OTHER INFORMATION: Alternative carboxy terminal tail

<400> SEQUENCE: 137

Lys Asp Glu Leu
1

<210> SEQ ID NO 138
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: alternative amino terminal portion of Domain IB

<400> SEQUENCE: 138

Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala Ala Gly Glu Cys Ala
1 5 10 15

<210> SEQ ID NO 139
<211> LENGTH: 35
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(35)
<223> OTHER INFORMATION: Domain IB

<400> SEQUENCE: 139

Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala Ala Gly Glu Cys Ala
1 5 10 15

Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr
20 25 30

Gly Ala Glu
35

<210> SEQ ID NO 140

-continued

```

<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Response element for ecdysone receptor
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 140

rrgggttcant gacacyy                                     17

<210> SEQ ID NO 141
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Ecdysone receptor response element
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (7)..(8)
<223> OTHER INFORMATION: nn at positions 7-8 is to indicate any number
      of spacer nucleotides

<400> SEQUENCE: 141

agggtcannag gtca                                         14

<210> SEQ ID NO 142
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: ecdysone receptor response element
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: ecdysone receptor response element

<400> SEQUENCE: 142

gggttgtaatg aattt                                       15

<210> SEQ ID NO 143
<211> LENGTH: 638
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(638)
<223> OTHER INFORMATION: Pseudomonas Exotoxin A variant

<400> SEQUENCE: 143

Met His Leu Ile Pro His Trp Ile Pro Leu Val Ala Ser Leu Gly Leu
1          5          10          15

Leu Ala Gly Gly Ser Ser Ala Ser Ala Ala Glu Glu Ala Phe Asp Leu
20          25          30

Trp Asn Glu Cys Ala Lys Ala Cys Val Leu Asp Leu Lys Asp Gly Val
35          40          45

Arg Ser Ser Arg Met Ser Val Asp Pro Ala Ile Ala Asp Thr Asn Gly
50          55          60

Gln Gly Val Leu His Tyr Ser Met Val Leu Glu Gly Gly Asn Asp Ala
65          70          75          80

Leu Lys Leu Ala Ile Asp Asn Ala Leu Ser Ile Thr Ser Asn Gly Leu
85          90          95

```

-continued

Thr	Ile	Arg	Leu	Glu	Gly	Gly	Val	Glu	Pro	Asn	Lys	Pro	Val	Arg	Tyr
			100					105					110		
Ser	Tyr	Thr	Arg	Gln	Ala	Arg	Gly	Ser	Trp	Ser	Leu	Asn	Trp	Leu	Val
		115					120					125			
Pro	Ile	Gly	His	Glu	Lys	Pro	Ser	Asn	Ile	Lys	Val	Phe	Ile	His	Glu
		130				135					140				
Leu	Asn	Ala	Gly	Asn	Gln	Leu	Ser	His	Met	Ser	Pro	Ile	Tyr	Thr	Ile
145				150					155						160
Glu	Met	Gly	Asp	Glu	Leu	Leu	Ala	Lys	Leu	Ala	Arg	Asp	Ala	Thr	Phe
			165					170						175	
Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu	Met	Gln	Pro	Thr	Leu	Ala	Ile
		180					185						190		
Ser	His	Ala	Gly	Val	Ser	Val	Val	Met	Ala	Gln	Ala	Gln	Pro	Arg	Arg
		195					200					205			
Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser	Gly	Lys	Val	Leu	Cys	Leu	Leu
	210					215					220				
Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr	Leu	Ala	Gln	Gln	Arg	Cys	Asn
225				230					235						240
Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile	Tyr	Arg	Val	Leu	Ala	Gly	Asn
			245					250						255	
Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys	Pro	Thr	Val	Ile	Ser	His	Arg
		260					265						270		
Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu	Ala	Ala	Leu	Thr	Ala	His	Gln
		275					280					285			
Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg
	290					295					300				
Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val
305				310					315						320
Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp	Gln	Val
			325					330						335	
Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu
		340					345						350		
Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala
		355				360						365			
Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu
	370				375						380				
Ala	Gly	Ala	Ala	Ser	Ala	Asp	Val	Val	Ser	Leu	Thr	Cys	Pro	Val	Ala
385				390					395						400
Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu
			405					410					415		
Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile
		420					425						430		
Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu
		435					440					445			
Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr
	450				455						460				
His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val
465				470					475						480
Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile
			485					490					495		
Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro
			500					505					510		

-continued

```

Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val
  515                      520                      525

Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala
  530                      535                      540

Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu
545                      550                      555                      560

Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg
  565                      570                      575

Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile
  580                      585                      590

Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp
  595                      600                      605

Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp
  610                      615                      620

Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
625                      630                      635

```

```

<210> SEQ ID NO 144
<211> LENGTH: 613
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(613)
<223> OTHER INFORMATION: PE variant

```

```

<400> SEQUENCE: 144

```

```

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
  1              5              10              15

Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
  20              25              30

Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
  35              40              45

Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
  50              55              60

Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
  65              70              75              80

Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
  85              90              95

Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
  100             105             110

Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
  115             120             125

Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
  130             135             140

Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His Glu Ser Asn Glu
  145             150             155             160

Met Gln Pro Thr Leu Ala Ile Ser His Ala Gly Val Ser Val Val Met
  165             170             175

Ala Gln Ala Gln Pro Arg Arg Glu Lys Arg Trp Ser Glu Trp Ala Ser
  180             185             190

Gly Lys Val Leu Cys Leu Leu Asp Pro Leu Asp Gly Val Tyr Asn Tyr
  195             200             205

Leu Ala Gln Gln Arg Cys Asn Leu Asp Asp Thr Trp Glu Gly Lys Ile

```

[illegible]

-continued

<210> SEQ ID NO 145
<211> LENGTH: 318
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(318)
<223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 145

Met Trp Glu Gln Leu Glu Gln Ser Gly Tyr Pro Val Gln Arg Leu Val
1 5 10 15

Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val
20 25 30

Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu
35 40 45

Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala
50 55 60

Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu
65 70 75 80

Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu
85 90 95

Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ile
100 105 110

Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu
115 120 125

Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr
130 135 140

His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val
145 150 155 160

Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile
165 170 175

Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro
180 185 190

Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val
195 200 205

Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala
210 215 220

Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu
225 230 235 240

Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg
245 250 255

Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile
260 265 270

Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp
275 280 285

Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp
290 295 300

Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
305 310 315

<210> SEQ ID NO 146
<211> LENGTH: 613
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*

-continued

<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(613)
<223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 146

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
1 5 10 15
Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
20 25 30
Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
35 40 45
Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
50 55 60
Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
65 70 75 80
Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
85 90 95
Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
100 105 110
Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
115 120 125
Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
130 135 140
Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His Glu Ser Asn Glu
145 150 155 160
Met Gln Pro Thr Leu Ala Ile Ser His Ala Gly Val Ser Val Val Met
165 170 175
Ala Gln Ala Gln Pro Arg Arg Glu Lys Arg Trp Ser Glu Trp Ala Ser
180 185 190
Gly Lys Val Leu Cys Leu Leu Asp Pro Leu Asp Gly Val Tyr Asn Tyr
195 200 205
Leu Ala Gln Gln Arg Cys Asn Leu Asp Asp Thr Trp Glu Gly Lys Ile
210 215 220
Tyr Arg Val Leu Ala Gly Asn Pro Ala Lys His Asp Leu Asp Ile Lys
225 230 235 240
Pro Thr Val Ile Ser His Arg Leu His Phe Pro Glu Gly Gly Ser Leu
245 250 255
Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro Leu Glu Thr Phe
260 265 270
Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly
275 280 285
Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser
290 295 300
Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly
305 310 315 320
Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala
325 330 335
Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg
340 345 350
Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Ser Ala Asp Val Val
355 360 365
Ser Leu Thr Cys Pro Val Ala Ala Gly Glu Cys Ala Gly Pro Ala Asp

-continued

370	375	380
Ser Gly Asp Ala Leu	Leu Glu Arg Asn Tyr	Pro Thr Gly Ala Glu Phe
385	390	395 400
Leu Gly Asp Gly Gly	Asp Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn	
	405	410 415
Trp Thr Val Glu Arg	Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg	
	420	425 430
Gly Tyr Val Phe Val	Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln	
	435	440 445
Ser Ile Val Phe Gly	Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala	
	450	455 460
Ile Trp Arg Gly Phe	Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly	
	465	470 475 480
Tyr Ala Gln Asp Gln	Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly	
	485	490 495
Ala Leu Leu Arg Val	Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr	
	500	505 510
Arg Thr Gly Leu Thr	Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu	
	515	520 525
Arg Leu Ile Gly His	Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly	
	530	535 540
Pro Glu Glu Glu Gly	Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu	
	545	550 555 560
Ala Glu Arg Thr Val	Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg	
	565	570 575
Asn Val Gly Gly Asp	Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln	
	580	585 590
Ala Ile Ser Ala Leu	Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro	
	595	600 605
Arg Glu Asp Leu Lys		
	610	
<210> SEQ ID NO 147		
<211> LENGTH: 613		
<212> TYPE: PRT		
<213> ORGANISM: Pseudomonas aeruginosa		
<220> FEATURE:		
<221> NAME/KEY: VARIANT		
<222> LOCATION: (1)..(613)		
<400> SEQUENCE: 147		
Ala Glu Glu Ala Phe	Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val	
1	5	10 15
Leu Asp Leu Lys Asp	Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro	
	20	25 30
Ala Ile Ala Asp Thr	Asn Gly Gln Gly Val Leu His Tyr Ser Met Val	
	35	40 45
Leu Glu Gly Gly Asn	Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu	
	50	55 60
Ser Ile Thr Ser Asp	Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu	
	65	70 75 80
Pro Asn Lys Pro Val	Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser	
	85	90 95
Trp Ser Leu Asn Trp	Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn	

-continued

100						105						110					
Ile	Lys	Val	Phe	Ile	His	Glu	Leu	Asn	Ala	Gly	Asn	Gln	Leu	Ser	His		
	115						120					125					
Met	Ser	Pro	Ile	Tyr	Thr	Ile	Glu	Met	Gly	Asp	Glu	Leu	Leu	Ala	Lys		
	130					135					140						
Leu	Ala	Arg	Asp	Ala	Thr	Phe	Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu		
145					150					155					160		
Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met		
				165					170					175			
Ala	Gln	Ala	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser		
				180				185					190				
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr		
		195					200				205						
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile		
	210					215					220						
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys		
225					230					235					240		
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu		
				245					250					255			
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe		
			260					265					270				
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly		
	275						280					285					
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser		
	290					295					300						
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly		
305					310					315					320		
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala		
			325						330					335			
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg		
			340					345					350				
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Ser	Ala	Asp	Val	Val		
		355					360					365					
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp		
	370					375					380						
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe		
385					390					395					400		
Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn		
			405						410					415			
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg		
			420					425					430				
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln		
		435					440					445					
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala		
	450					455					460						
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly		
465					470					475					480		
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly		
			485					490						495			
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr		
			500					505					510				

-continued

Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu
515 520 525

Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly
530 535 540

Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu
545 550 555 560

Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg
565 570 575

Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Gln Glu Gln
580 585 590

Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Gln Pro Pro
595 600 605

Arg Glu Asp Leu Arg
610

<210> SEQ ID NO 148
 <211> LENGTH: 346
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(346)
 <223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 148

Met Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His Leu
1 5 10 15

Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln
20 25 30

Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu
35 40 45

Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala
50 55 60

Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu
65 70 75 80

Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser
85 90 95

Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala
100 105 110

Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro
115 120 125

Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser Thr
130 135 140

Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg
145 150 155 160

Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe
165 170 175

Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser
180 185 190

Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro
195 200 205

Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly
210 215 220

Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser
225 230 235 240

-continued

Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala
 245 250 255
 Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu
 260 265 270
 Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile
 275 280 285
 Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile
 290 295 300
 Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile
 305 310 315 320
 Pro Asp Gln Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln
 325 330 335
 Pro Gly Gln Pro Pro Arg Glu Asp Leu Arg
 340 345

<210> SEQ ID NO 149
 <211> LENGTH: 613
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(613)
 <223> OTHER INFORMATION: PE variant
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(613)
 <223> OTHER INFORMATION: To use for fusion to ligand at c-terminus,
 delete AA 1-252 and 365-380; all of domain IB and portion of
 domain II (AA 350-394 can be deleted and replaced with GGGGS
 linker sequence.

<400> SEQUENCE: 149

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
 1 5 10 15
 Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
 20 25 30
 Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
 35 40 45
 Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
 50 55 60
 Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
 65 70 75 80
 Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
 85 90 95
 Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
 100 105 110
 Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
 115 120 125
 Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
 130 135 140
 Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His Glu Ser Asn Glu
 145 150 155 160
 Met Gln Pro Thr Leu Ala Ile Ser His Ala Gly Val Ser Val Val Met
 165 170 175
 Ala Gln Ala Gln Pro Arg Arg Glu Lys Arg Trp Ser Glu Trp Ala Ser
 180 185 190

-continued

Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr
	195						200					205			
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile
	210					215					220				
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys
225					230					235					240
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu
				245					250					255	
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
			260					265					270		
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly
	275						280					285			
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
	290					295					300				
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
305					310					315					320
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
				325					330					335	
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
			340					345					350		
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Ser	Ala	Asp	Val	Val
	355						360					365			
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp
	370					375					380				
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
385					390					395					400
Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
				405					410					415	
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg
		420						425					430		
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln
	435						440					445			
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala
	450					455					460				
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly
465					470					475					480
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly
				485					490					495	
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr
			500					505					510		
Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu
		515					520					525			
Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly
	530					535					540				
Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu
545					550					555					560
Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg
				565					570					575	
Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Lys	Glu	Gln
			580					585					590		
Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro	Pro
	595						600					605			

-continued

Arg Glu Asp Leu Lys
610

<210> SEQ ID NO 150
<211> LENGTH: 321
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(321)
<223> OTHER INFORMATION: PE variant with deletions for use as fusion
protein to ligand C-terminus

<400> SEQUENCE: 150

Gly Gly Gly Gly Ser Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln
1 5 10 15
Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg
20 25 30
Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val
35 40 45
Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val
50 55 60
Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu
65 70 75 80
Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala
85 90 95
Ala Ala Glu Ser Glu Arg Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly
100 105 110
Gly Asp Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu
115 120 125
Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe
130 135 140
Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe
145 150 155 160
Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly
165 170 175
Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp
180 185 190
Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg
195 200 205
Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu
210 215 220
Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly
225 230 235 240
His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu
245 250 255
Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr
260 265 270
Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly
275 280 285
Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala
290 295 300
Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu
305 310 315 320

-continued

Lys

<210> SEQ ID NO 151
<211> LENGTH: 321
<212> TYPE: PRT
<213> ORGANISM: *Pseudomonas aeruginosa*
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(321)
<223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 151

Gly Gly Gly Gly Ser Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln
1 5 10 15
Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg
20 25 30
Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val
35 40 45
Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val
50 55 60
Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu
65 70 75 80
Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala
85 90 95
Ala Ala Glu Ser Glu Arg Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly
100 105 110
Gly Asp Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu
115 120 125
Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe
130 135 140
Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe
145 150 155 160
Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly
165 170 175
Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp
180 185 190
Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg
195 200 205
Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu
210 215 220
Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly
225 230 235 240
His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu
245 250 255
Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr
260 265 270
Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly
275 280 285
Asp Leu Asp Pro Ser Ser Ile Pro Asp Gln Glu Gln Ala Ile Ser Ala
290 295 300
Leu Pro Asp Tyr Ala Ser Gln Pro Gly Gln Pro Pro Arg Glu Asp Leu
305 310 315 320
Arg

-continued

```

<210> SEQ ID NO 152
<211> LENGTH: 613
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(613)
<223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 152

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
1          5          10          15
Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
20          25          30
Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
35          40          45
Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
50          55          60
Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
65          70          75          80
Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
85          90          95
Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
100         105         110
Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
115         120         125
Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
130         135         140
Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His Glu Ser Asn Glu
145         150         155         160
Met Gln Pro Thr Leu Ala Ile Ser His Ala Gly Val Ser Val Val Met
165         170         175
Ala Gln Ala Gln Pro Arg Arg Glu Lys Arg Trp Ser Glu Trp Ala Ser
180         185         190
Gly Lys Val Leu Cys Leu Leu Asp Pro Leu Asp Gly Val Tyr Asn Tyr
195         200         205
Leu Ala Gln Gln Arg Cys Asn Leu Asp Asp Thr Trp Glu Gly Lys Ile
210         215         220
Tyr Arg Val Leu Ala Gly Asn Pro Ala Lys His Asp Leu Asp Ile Lys
225         230         235         240
Pro Thr Val Ile Ser His Arg Leu His Phe Pro Glu Gly Gly Ser Leu
245         250         255
Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro Leu Glu Thr Phe
260         265         270
Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly
275         280         285
Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser
290         295         300
Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly
305         310         315         320
Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala
325         330         335
Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg
340         345         350

```

-continued

Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Ser Ala Asp Val Val
 355 360 365
 Ser Leu Thr Cys Pro Val Ala Ala Gly Glu Cys Ala Gly Pro Ala Asp
 370 375 380
 Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe
 385 390 395 400
 Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn
 405 410 415
 Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg
 420 425 430
 Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln
 435 440 445
 Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala
 450 455 460
 Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly
 465 470 475 480
 Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly
 485 490 495
 Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr
 500 505 510
 Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu
 515 520 525
 Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly
 530 535 540
 Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu
 545 550 555 560
 Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg
 565 570 575
 Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln
 580 585 590
 Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
 595 600 605
 Arg Glu Asp Leu Lys
 610

<210> SEQ ID NO 153
 <211> LENGTH: 363
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(363)
 <223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 153

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
 1 5 10 15
 Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
 20 25 30
 Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
 35 40 45
 Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
 50 55 60
 Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
 65 70 75 80

-continued

Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
 85 90 95
 Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
 100 105 110
 Ala Ser Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala Ala Gly Glu
 115 120 125
 Cys Ala Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
 130 135 140
 Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser
 145 150 155 160
 Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
 165 170 175
 Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
 180 185 190
 Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
 195 200 205
 Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
 210 215 220
 Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
 225 230 235 240
 Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
 245 250 255
 Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
 260 265 270
 Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
 275 280 285
 Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr
 290 295 300
 Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
 305 310 315 320
 Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
 325 330 335
 Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
 340 345 350
 Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 355 360

<210> SEQ ID NO 154
 <211> LENGTH: 318
 <212> TYPE: PRT
 <213> ORGANISM: *Pseudomonas aeruginosa*
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(318)
 <223> OTHER INFORMATION: PE37 variant beginning with initiation
 methionine

 <400> SEQUENCE: 154

Met Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val
 1 5 10 15
 Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val
 20 25 30
 Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu
 35 40 45

-continued

Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala
50						55					60				
Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu
65					70					75					80
Ala	Gly	Ala	Ala	Asn	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu
				85					90					95	
Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Val
			100					105						110	
Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu
			115				120					125			
Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr
			130				135					140			
His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val
145					150					155					160
Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile
				165					170					175	
Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro
			180					185					190		
Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val
			195				200					205			
Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Ser	Leu	Thr	Leu	Ala
						215					220				
Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu
225					230					235					240
Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg
				245					250					255	
Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile
			260					265					270		
Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp
			275				280					285			
Pro	Ser	Ser	Ile	Pro	Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp
			290			295					300				
Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys		
305					310					315					

<210> SEQ ID NO 155

<211> LENGTH: 346

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (1)..(346)

<223> OTHER INFORMATION: PE variant beginning with methionine

<400> SEQUENCE: 155

Met	Gly	Gly	Ser	Leu	Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu
1				5					10				15		
Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln
			20					25					30		
Leu	Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu
			35				40					45			
Ala	Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala
			50			55				60					
Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu
65					70				75						80

-continued

Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser
 85 90 95
 Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala
 100 105 110
 Asn Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro
 115 120 125
 Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser Thr
 130 135 140
 Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg
 145 150 155 160
 Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe
 165 170 175
 Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser
 180 185 190
 Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro
 195 200 205
 Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly
 210 215 220
 Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser
 225 230 235 240
 Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu Ala
 245 250 255
 Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu
 260 265 270
 Asp Ala Ile Thr Gly Pro Glu Glu Gly Gly Arg Leu Glu Thr Ile
 275 280 285
 Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile
 290 295 300
 Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile
 305 310 315 320
 Pro Asp Gln Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln
 325 330 335
 Pro Gly Gln Pro Pro Arg Glu Asp Leu Arg
 340 345

<210> SEQ ID NO 156

<211> LENGTH: 613

<212> TYPE: PRT

<213> ORGANISM: *Pseudomonas aeruginosa*

<400> SEQUENCE: 156

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
 1 5 10 15
 Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
 20 25 30
 Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
 35 40 45
 Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
 50 55 60
 Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
 65 70 75 80
 Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
 85 90 95

-continued

Trp	Ser	Leu	Asn	Trp	Leu	Val	Pro	Ile	Gly	His	Glu	Lys	Pro	Ser	Asn
		100						105					110		
Ile	Lys	Val	Phe	Ile	His	Glu	Leu	Asn	Ala	Gly	Asn	Gln	Leu	Ser	His
		115					120					125			
Met	Ser	Pro	Ile	Tyr	Thr	Ile	Glu	Met	Gly	Asp	Glu	Leu	Leu	Ala	Lys
	130					135					140				
Leu	Ala	Arg	Asp	Ala	Thr	Phe	Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu
145					150					155					160
Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met
			165						170					175	
Ala	Gln	Ala	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser
			180					185					190		
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr
		195					200					205			
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile
	210					215					220				
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys
225					230					235					240
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu
			245						250					255	
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
			260					265					270		
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly
		275					280					285			
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
	290					295					300				
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
305					310					315					320
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
			325						330					335	
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
			340					345					350		
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Ala	Asp	Val	Val
		355					360					365			
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp
	370					375					380				
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
385					390					395					400
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
			405						410					415	
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg
			420					425					430		
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln
		435					440					445			
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala
	450					455					460				
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly
465					470					475					480
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly
			485						490					495	
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr

-continued

500					505					510					
Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu
515					520					525					
Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly
530					535					540					
Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu
545					550					555					
Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg
565					570					575					
Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Gln	Glu	Gln
580					585					590					
Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Gln	Pro	Pro
595					600					605					
Arg	Glu	Asp	Leu	Arg											
610															

<210> SEQ ID NO 157
 <211> LENGTH: 224
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(224)
 <223> OTHER INFORMATION: PE variant fusion-ready sequeunce with Gly-Ser linker at amino terminus for joining to C-terminal end of heterologous protein
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(5)
 <223> OTHER INFORMATION: Gly-Ser linker

<400> SEQUENCE: 157

Gly	Gly	Gly	Gly	Ser	Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly
1				5					10					15	
Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg
			20				25						30		
Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val
		35				40						45			
Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly
	50					55				60					
Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe
65					70				75					80	
Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln
			85						90					95	
Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val
		100						105					110		
Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Ser	Leu	Thr
		115					120					125			
Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His
		130				135						140			
Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly
145					150					155				160	
Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val
			165					170						175	
Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp
			180					185					190		

-continued

Leu Asp Pro Ser Ser Ile Pro Asp Gln Glu Gln Ala Ile Ser Ala Leu
195 200 205

Pro Asp Tyr Ala Ser Gln Pro Gly Gln Pro Pro Arg Glu Asp Leu Arg
210 215 220

<210> SEQ ID NO 158
 <211> LENGTH: 334
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(334)
 <223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 158

Met Trp Glu Gln Leu Glu Gln Ser Gly Tyr Pro Val Gln Arg Leu Val
1 5 10 15

Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val
20 25 30

Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu
35 40 45

Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala
50 55 60

Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu
65 70 75 80

Ala Gly Ala Ala Asn Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala
85 90 95

Ala Gly Glu Cys Ala Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu
100 105 110

Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val
115 120 125

Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu
130 135 140

Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr
145 150 155 160

His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val
165 170 175

Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile
180 185 190

Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro
195 200 205

Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val
210 215 220

Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala
225 230 235 240

Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu
245 250 255

Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg
260 265 270

Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile
275 280 285

Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp
290 295 300

Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp
305 310 315 320

-continued

Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 325 330

<210> SEQ ID NO 159
 <211> LENGTH: 318
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(318)
 <223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 159

Met Trp Glu Gln Leu Glu Gln Ser Gly Tyr Pro Val Gln Arg Leu Val
 1 5 10 15

Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val
 20 25 30

Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu
 35 40 45

Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala
 50 55 60

Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu
 65 70 75 80

Ala Gly Ala Ala Asn Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu
 85 90 95

Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val
 100 105 110

Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu
 115 120 125

Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr
 130 135 140

His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val
 145 150 155 160

Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile
 165 170 175

Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro
 180 185 190

Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val
 195 200 205

Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala
 210 215 220

Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu
 225 230 235 240

Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg
 245 250 255

Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile
 260 265 270

Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp
 275 280 285

Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp
 290 295 300

Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 305 310 315

-continued

```

<210> SEQ ID NO 160
<211> LENGTH: 363
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(363)
<223> OTHER INFORMATION: PE variant

<400> SEQUENCE: 160

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1      5      10     15
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20     25     30
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35     40     45
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50     55     60
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65     70     75     80
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
85     90     95
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100    105    110
Ala Asn Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala Ala Gly Glu
115    120    125
Cys Ala Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
130    135    140
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser
145    150    155    160
Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
165    170    175
Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
180    185    190
Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
195    200    205
Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
210    215    220
Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
225    230    235    240
Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
245    250    255
Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu
260    265    270
Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
275    280    285
Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr
290    295    300
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
305    310    315    320
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
325    330    335
Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
340    345    350

```

-continued

Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 355 360

<210> SEQ ID NO 161
 <211> LENGTH: 635
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas aeruginosa
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(635)
 <223> OTHER INFORMATION: PE variant from GenBank Accession Number
 YP_792118

<400> SEQUENCE: 161

Met His Leu Ile Pro His Trp Ile Pro Leu Val Ala Ser Leu Gly Leu
 1 5 10 15

Leu Ala Gly Gly Ser Phe Ala Ser Ala Ala Glu Glu Ala Phe Asp Leu
 20 25 30

Trp Asn Glu Cys Ala Lys Ala Cys Val Leu Asp Leu Lys Asp Gly Val
 35 40 45

Arg Ser Ser Arg Met Ser Val Asp Pro Ala Ile Ala Asp Thr Asn Gly
 50 55 60

Gln Gly Val Leu His Tyr Ser Met Val Leu Glu Gly Gly Asn Asp Ala
 65 70 75 80

Leu Lys Leu Ala Ile Asp Asn Ala Leu Ser Ile Thr Ser Asp Gly Leu
 85 90 95

Thr Ile Arg Leu Glu Gly Gly Val Glu Pro Asn Lys Pro Val Arg Tyr
 100 105 110

Ser Tyr Thr Arg Gln Ala Arg Gly Ser Trp Ser Leu Asn Trp Leu Val
 115 120 125

Pro Ile Gly His Glu Lys Pro Ser Asn Ile Lys Val Phe Ile His Glu
 130 135 140

Leu Asn Ala Gly Asn Gln Leu Ser His Met Ser Pro Ile Tyr Thr Ile
 145 150 155 160

Glu Met Gly Asp Glu Leu Leu Ala Lys Leu Ala Arg Asp Ala Thr Phe
 165 170 175

Phe Val Arg Ala His Glu Ser Asn Glu Met Gln Pro Thr Leu Ala Ile
 180 185 190

Ser His Ala Gly Val Ser Val Val Met Ala Gln Thr Gln Pro Arg Arg
 195 200 205

Glu Lys Arg Trp Ser Glu Trp Ala Ser Gly Lys Val Leu Cys Leu Leu
 210 215 220

Asp Pro Leu Asp Gly Val Tyr Asn Tyr Leu Ala Gln Gln Arg Cys Asn
 225 230 235 240

Leu Asp Asp Thr Trp Glu Gly Lys Ile Tyr Arg Val Leu Ala Gly Asn
 245 250 255

Pro Ala Lys His Asp Leu Asp Ile Lys Pro Thr Val Ile Ser His Arg
 260 265 270

Leu His Phe Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln
 275 280 285

Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg
 290 295 300

Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val
 305 310 315 320

Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val

-continued

325					330					335					
Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu
			340					345					350		
Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala
			355					360					365		
Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu
			370					375					380		
Ala	Ser	Ala	Asp	Val	Val	Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu
				390											400
Cys	Ala	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr
				405					410						415
Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser
				420					425						430
Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His
				435					440						445
Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr
				450					455						460
Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg
				465					470						480
Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp
				485					490						495
Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg
				500					505						510
Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser
				515					520						525
Ser	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu
				530					535						540
Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg
				545					550						555
Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	
				565					570						575
Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala
				580					585						590
Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser
				595					600						605
Ile	Pro	Asp	Gln	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser
				610					615						620
Gln	Pro	Gly	Lys	Pro	Ser	Arg	Glu	Asp	Leu	Lys					
				625					630						635

<210> SEQ ID NO 162

<211> LENGTH: 613

<212> TYPE: PRT

<213> ORGANISM: Pseudomonas aeruginosa

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (1)..(613)

<223> OTHER INFORMATION: PE variant from Genbank Accession number 1IKQ_A

<400> SEQUENCE: 162

Ala	Glu	Glu	Ala	Phe	Asp	Leu	Trp	Asn	Glu	Cys	Ala	Lys	Ala	Cys	Val
1				5					10					15	

Leu	Asp	Leu	Lys	Asp	Gly	Val	Arg	Ser	Ser	Arg	Met	Ser	Val	Asp	Pro
			20					25						30	

-continued

Ala	Ile	Ala	Asp	Thr	Asn	Gly	Gln	Gly	Val	Leu	His	Tyr	Ser	Met	Val
	35						40					45			
Leu	Glu	Gly	Gly	Asn	Asp	Ala	Leu	Lys	Leu	Ala	Ile	Asp	Asn	Ala	Leu
	50					55					60				
Ser	Ile	Thr	Ser	Asp	Gly	Leu	Thr	Ile	Arg	Leu	Glu	Gly	Gly	Val	Glu
65					70					75				80	
Pro	Asn	Lys	Pro	Val	Arg	Tyr	Ser	Tyr	Thr	Arg	Gln	Ala	Arg	Gly	Ser
				85					90					95	
Trp	Ser	Leu	Asn	Trp	Leu	Val	Pro	Ile	Gly	His	Glu	Lys	Pro	Ser	Asn
			100					105					110		
Ile	Lys	Val	Phe	Ile	His	Glu	Leu	Asn	Ala	Gly	Asn	Gln	Leu	Ser	His
		115					120					125			
Met	Ser	Pro	Ile	Tyr	Thr	Ile	Glu	Met	Gly	Asp	Glu	Leu	Leu	Ala	Lys
	130					135					140				
Leu	Ala	Arg	Asp	Ala	Thr	Phe	Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu
145					150					155					160
Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met
				165					170					175	
Ala	Gln	Ala	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser
			180					185					190		
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr
	195						200					205			
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile
	210					215					220				
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys
225					230					235					240
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu
				245					250					255	
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
			260					265					270		
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly
		275					280					285			
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
	290					295					300				
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
305					310					315					320
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
				325					330					335	
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
			340					345					350		
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Ala	Asp	Val	Val
		355					360					365			
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp
	370					375					380				
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
385					390					395					400
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
				405					410					415	
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg
			420					425					430		
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln
	435						440					445			

-continued

```

Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala
 450                455                460

Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly
465                470                475                480

Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly
                485                490                495

Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr
                500                505                510

Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu
                515                520                525

Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly
                530                535                540

Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu
545                550                555                560

Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg
                565                570                575

Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln
                580                585                590

Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
                595                600                605

Arg Glu Asp Leu Lys
 610

```

```

<210> SEQ ID NO 163
<211> LENGTH: 613
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(613)
<223> OTHER INFORMATION: PE variant from Genbank Accession number 1IKP_A

```

```

<400> SEQUENCE: 163

```

```

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
 1                5                10                15

Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
                20                25                30

Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
                35                40                45

Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
                50                55                60

Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
65                70                75                80

Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
                85                90                95

Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
                100               105               110

Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
                115                120                125

Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
                130                135                140

Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His Glu Ser Asn Glu
145                150                155                160

Met Gln Pro Thr Leu Ala Ile Ser His Ala Gly Val Ser Val Val Met

```

-continued

165								170					175				
Ala	Gln	Ala	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser		
			180					185					190				
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Gln	Leu	Asp	Gly	Val	Tyr	Asn	Tyr		
		195					200					205					
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile		
	210					215					220						
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys		
225					230					235				240			
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu		
				245					250					255			
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe		
			260					265					270				
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Ala	Glu	Gln	Leu	Glu	Gln	Cys	Gly		
	275						280					285					
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser		
	290					295					300						
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly		
305					310					315				320			
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala		
			325						330					335			
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg		
			340					345					350				
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Ala	Asp	Val	Val		
		355					360					365					
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp		
	370					375					380						
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe		
385					390					395				400			
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn		
			405						410					415			
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg		
			420					425					430				
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln		
		435					440					445					
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala		
	450					455					460						
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly		
465					470				475					480			
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly		
			485					490						495			
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr		
			500					505					510				
Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu		
		515					520					525					
Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly		
	530					535					540						
Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu		
545					550					555				560			
Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg		
			565						570					575			

```
<210> SEQ ID NO 164
<211> LENGTH: 517
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: IL2-PE fusion protein
<220> FEATURE:
<221> NAME/KEY: SIGNAL
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: Predicted signal peptide sequence
```

Met 1	Tyr	Arg	Met	Gln 5	Leu	Leu	Ser	Cys	Ile 10	Ala	Leu	Ser	Leu	Ala 15	Leu
Val	Thr	Asn	Ser 20	Ala	Pro	Thr	Ser	Ser 25	Ser	Thr	Lys	Lys	Thr 30	Gln	Leu
Gln	Leu	Glu 35	His	Leu	Leu	Leu	Asp 40	Leu	Gln	Met	Ile 45	Leu	Asn	Gly	Ile
Asn 50	Asn	Tyr	Lys	Asn	Pro	Lys 55	Leu	Thr	Arg	Met 60	Leu	Thr	Phe	Lys	Phe
Tyr 65	Met	Pro	Lys	Lys	Ala 70	Thr	Glu	Leu	Lys	His 75	Leu	Gln	Cys	Leu	Glu 80
Glu	Glu	Leu	Lys 85	Pro	Leu	Glu	Glu	Val 90	Leu	Asn	Leu	Ala	Gln	Ser 95	Lys
Asn	Phe	His	Leu 100	Arg	Pro	Arg	Asp 105	Leu	Ile	Ser	Asn	Ile 110	Asn	Val	Ile
Val	Leu	Glu 115	Leu	Lys	Gly	Ser	Glu 120	Thr	Thr	Phe	Met	Cys 125	Glu	Tyr	Ala
Asp 130	Glu	Thr	Ala	Thr	Ile	Val 135	Glu	Phe	Leu	Asn	Arg 140	Trp	Ile	Thr	Phe
Cys 145	Gln	Ser	Ile	Ile	Ser 150	Thr	Leu	Thr	Ile	Pro 155	Glu	Gly	Gly	Ser	Leu 160
Ala	Ala	Leu	Thr 165	Ala	His	Gln	Ala	Cys 170	His	Leu	Pro	Leu	Glu	Thr 175	Phe
Thr	Arg	His	Arg 180	Gln	Pro	Arg	Gly	Trp 185	Glu	Gln	Leu	Glu	Gln 190	Cys	Gly
Tyr	Pro	Val 195	Gln	Arg	Leu	Val	Ala 200	Leu	Tyr	Leu	Ala	Ala 205	Arg	Leu	Ser
Trp 210	Asn	Gln	Val	Asp	Gln 215	Val	Ile	Arg	Asn	Ala	Leu 220	Ala	Ser	Pro	Gly
Ser 225	Gly	Gly	Asp	Leu	Gly 230	Glu	Ala	Ile	Arg	Glu 235	Gln	Pro	Glu	Gln	Ala 240
Arg	Leu	Ala	Leu 245	Thr	Leu	Ala	Ala	Ala 250	Glu	Ser	Glu	Arg	Phe 255	Val	Arg
Gln	Gly	Thr	Gly 260	Asn	Asp	Glu	Ala	Gly 265	Ala	Ala	Ser	Ala	Asp 270	Val	Val
Ser	Leu	Thr 275	Cys	Pro	Val	Ala	Ala 280	Gly	Glu	Cys	Ala	Gly 285	Pro	Ala	Asp

-continued

```

Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe
 290                295                300

Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn
305                310                315                320

Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg
                325                330                335

Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln
                340                345                350

Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala
                355                360                365

Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly
                370                375                380

Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly
385                390                395                400

Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr
                405                410                415

Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu
                420                425                430

Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly
                435                440                445

Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu
                450                455                460

Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg
465                470                475                480

Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln
                485                490                495

Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
                500                505                510

Arg Glu Asp Leu Lys
 515

<210> SEQ ID NO 165
<211> LENGTH: 506
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: IL2-PE fusion protein with Gly-Ser linker
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1) .. (506)
<223> OTHER INFORMATION: IL2-PE fusion protein with Gly-Ser linker
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (154) .. (159)
<223> OTHER INFORMATION: Gly-Ser linker

<400> SEQUENCE: 165

Met Tyr Arg Met Gln Leu Leu Ser Cys Ile Ala Leu Ser Leu Ala Leu
 1                5                10                15

Val Thr Asn Ser Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu
                20                25                30

Gln Leu Glu His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile
                35                40                45

Asn Asn Tyr Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe
                50                55                60

Tyr Met Pro Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu
65                70                75                80

```

-continued

Glu	Glu	Leu	Lys	Pro	Leu	Glu	Glu	Val	Leu	Asn	Leu	Ala	Gln	Ser	Lys	85	90	95	
Asn	Phe	His	Leu	Arg	Pro	Arg	Asp	Leu	Ile	Ser	Asn	Ile	Asn	Val	Ile	100	105	110	
Val	Leu	Glu	Leu	Lys	Gly	Ser	Glu	Thr	Thr	Phe	Met	Cys	Glu	Tyr	Ala	115	120	125	
Asp	Glu	Thr	Ala	Thr	Ile	Val	Glu	Phe	Leu	Asn	Arg	Trp	Ile	Thr	Phe	130	135	140	
Cys	Gln	Ser	Ile	Ile	Ser	Thr	Leu	Thr	Gly	Gly	Gly	Gly	Gly	Ser	Pro	145	150	155	160
Glu	Gly	Gly	Ser	Leu	Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	165	170	175	
Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	180	185	190	
Leu	Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	195	200	205	
Ala	Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	210	215	220	
Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	225	230	235	240
Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	245	250	255	
Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	260	265	270	
Ser	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	275	280	285	
Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	290	295	300	
Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	305	310	315	320
Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	325	330	335	
Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	340	345	350	
Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	355	360	365	
Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	370	375	380	
Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	385	390	395	400
Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	405	410	415	
Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	420	425	430	
Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile		435	440	445	
Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	450	455	460	
Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	465	470	475	480
Pro	Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln				

-continued

	485	490	495
--	-----	-----	-----

Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 500 505

<210> SEQ ID NO 166
 <211> LENGTH: 517
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: IL2-PE fusion sequence
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(517)
 <223> OTHER INFORMATION: IL2-PE fusion protein
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: predicted signal peptide sequence
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (21)..(153)
 <223> OTHER INFORMATION: IL2 sequence
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (21)..(153)
 <223> OTHER INFORMATION: IL2 sequence
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (154)..(154)
 <223> OTHER INFORMATION: Linking isoleucine amino acid
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (154)..(154)
 <223> OTHER INFORMATION: Linking isoleucine amino acid
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (155)..(517)
 <223> OTHER INFORMATION: PE variant sequence

<400> SEQUENCE: 166

Met	Tyr	Arg	Met	Gln	Leu	Leu	Ser	Cys	Ile	Ala	Leu	Ser	Leu	Ala	Leu
1				5					10					15	
Val	Thr	Asn	Ser	Ala	Pro	Thr	Ser	Ser	Ser	Thr	Lys	Lys	Thr	Gln	Leu
			20					25					30		
Gln	Leu	Glu	His	Leu	Leu	Leu	Asp	Leu	Gln	Met	Ile	Leu	Asn	Gly	Ile
		35					40					45			
Asn	Asn	Tyr	Lys	Asn	Pro	Lys	Leu	Thr	Arg	Met	Leu	Thr	Phe	Lys	Phe
	50					55				60					
Tyr	Met	Pro	Lys	Lys	Ala	Thr	Glu	Leu	Lys	His	Leu	Gln	Cys	Leu	Glu
65					70					75				80	
Glu	Glu	Leu	Lys	Pro	Leu	Glu	Glu	Val	Leu	Asn	Leu	Ala	Gln	Ser	Lys
			85					90						95	
Asn	Phe	His	Leu	Arg	Pro	Arg	Asp	Leu	Ile	Ser	Asn	Ile	Asn	Val	Ile
			100					105					110		
Val	Leu	Glu	Leu	Lys	Gly	Ser	Glu	Thr	Thr	Phe	Met	Cys	Glu	Tyr	Ala
		115					120					125			
Asp	Glu	Thr	Ala	Thr	Ile	Val	Glu	Phe	Leu	Asn	Arg	Trp	Ile	Thr	Phe
	130					135					140				
Cys	Gln	Ser	Ile	Ile	Ser	Thr	Leu	Thr	Ile	Pro	Glu	Gly	Gly	Ser	Leu
145					150					155					160
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
			165						170					175	
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly

-continued

180					185					190					
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
		195					200					205			
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
	210					215					220				
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
	225					230					235				240
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
			245						250					255	
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Ala	Asp	Val	Val
			260					265					270		
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp
	275					280					285				
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
	290					295					300				
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
	305					310					315				320
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg
			325						330					335	
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln
			340					345					350		
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala
	355					360					365				
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly
	370					375					380				
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly
	385					390					395				400
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr
			405					410						415	
Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu
			420					425					430		
Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly
	435					440					445				
Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu
	450					455					460				
Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg
	465					470					475				480
Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Lys	Glu	Gln
			485					490						495	
Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro	Pro
			500					505					510		
Arg	Glu	Asp	Leu	Lys											
			515												

<210> SEQ ID NO 167
 <211> LENGTH: 630
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(630)
 <223> OTHER INFORMATION: Mesothelin sequence from GenBank accession
 NP_037536
 <400> SEQUENCE: 167

-continued

Met	Ala	Leu	Pro	Thr	Ala	Arg	Pro	Leu	Leu	Gly	Ser	Cys	Gly	Thr	Pro
1				5					10					15	
Ala	Leu	Gly	Ser	Leu	Leu	Phe	Leu	Leu	Phe	Ser	Leu	Gly	Trp	Val	Gln
		20						25					30		
Pro	Ser	Arg	Thr	Leu	Ala	Gly	Glu	Thr	Gly	Gln	Glu	Ala	Ala	Pro	Leu
		35					40					45			
Asp	Gly	Val	Leu	Ala	Asn	Pro	Pro	Asn	Ile	Ser	Ser	Leu	Ser	Pro	Arg
	50				55					60					
Gln	Leu	Leu	Gly	Phe	Pro	Cys	Ala	Glu	Val	Ser	Gly	Leu	Ser	Thr	Glu
65				70						75				80	
Arg	Val	Arg	Glu	Leu	Ala	Val	Ala	Leu	Ala	Gln	Lys	Asn	Val	Lys	Leu
			85					90						95	
Ser	Thr	Glu	Gln	Leu	Arg	Cys	Leu	Ala	His	Arg	Leu	Ser	Glu	Pro	Pro
		100						105					110		
Glu	Asp	Leu	Asp	Ala	Leu	Pro	Leu	Asp	Leu	Leu	Leu	Phe	Leu	Asn	Pro
	115					120						125			
Asp	Ala	Phe	Ser	Gly	Pro	Gln	Ala	Cys	Thr	Arg	Phe	Phe	Ser	Arg	Ile
	130					135					140				
Thr	Lys	Ala	Asn	Val	Asp	Leu	Leu	Pro	Arg	Gly	Ala	Pro	Glu	Arg	Gln
145				150						155					160
Arg	Leu	Leu	Pro	Ala	Ala	Leu	Ala	Cys	Trp	Gly	Val	Arg	Gly	Ser	Leu
			165					170						175	
Leu	Ser	Glu	Ala	Asp	Val	Arg	Ala	Leu	Gly	Gly	Leu	Ala	Cys	Asp	Leu
		180						185					190		
Pro	Gly	Arg	Phe	Val	Ala	Glu	Ser	Ala	Glu	Val	Leu	Leu	Pro	Arg	Leu
		195					200						205		
Val	Ser	Cys	Pro	Gly	Pro	Leu	Asp	Gln	Asp	Gln	Gln	Glu	Ala	Ala	Arg
	210					215					220				
Ala	Ala	Leu	Gln	Gly	Gly	Gly	Pro	Pro	Tyr	Gly	Pro	Pro	Ser	Thr	Trp
225				230						235					240
Ser	Val	Ser	Thr	Met	Asp	Ala	Leu	Arg	Gly	Leu	Leu	Pro	Val	Leu	Gly
			245					250						255	
Gln	Pro	Ile	Ile	Arg	Ser	Ile	Pro	Gln	Gly	Ile	Val	Ala	Ala	Trp	Arg
		260						265					270		
Gln	Arg	Ser	Ser	Arg	Asp	Pro	Ser	Trp	Arg	Gln	Pro	Glu	Arg	Thr	Ile
	275					280						285			
Leu	Arg	Pro	Arg	Phe	Arg	Arg	Glu	Val	Glu	Lys	Thr	Ala	Cys	Pro	Ser
	290					295					300				
Gly	Lys	Lys	Ala	Arg	Glu	Ile	Asp	Glu	Ser	Leu	Ile	Phe	Tyr	Lys	Lys
305				310						315					320
Trp	Glu	Leu	Glu	Ala	Cys	Val	Asp	Ala	Ala	Leu	Leu	Ala	Thr	Gln	Met
			325					330						335	
Asp	Arg	Val	Asn	Ala	Ile	Pro	Phe	Thr	Tyr	Glu	Gln	Leu	Asp	Val	Leu
		340						345					350		
Lys	His	Lys	Leu	Asp	Glu	Leu	Tyr	Pro	Gln	Gly	Tyr	Pro	Glu	Ser	Val
		355					360					365			
Ile	Gln	His	Leu	Gly	Tyr	Leu	Phe	Leu	Lys	Met	Ser	Pro	Glu	Asp	Ile
	370					375					380				
Arg	Lys	Trp	Asn	Val	Thr	Ser	Leu	Glu	Thr	Leu	Lys	Ala	Leu	Leu	Glu
385				390						395					400
Val	Asn	Lys	Gly	His	Glu	Met	Ser	Pro	Gln	Ala	Pro	Arg	Arg	Pro	Leu

-continued

405					410					415					
Pro	Gln	Val	Ala	Thr	Leu	Ile	Asp	Arg	Phe	Val	Lys	Gly	Arg	Gly	Gln
		420						425					430		
Leu	Asp	Lys	Asp	Thr	Leu	Asp	Thr	Leu	Thr	Ala	Phe	Tyr	Pro	Gly	Tyr
	435					440					445				
Leu	Cys	Ser	Leu	Ser	Pro	Glu	Glu	Leu	Ser	Ser	Val	Pro	Pro	Ser	Ser
	450					455					460				
Ile	Trp	Ala	Val	Arg	Pro	Gln	Asp	Leu	Asp	Thr	Cys	Asp	Pro	Arg	Gln
465					470					475					480
Leu	Asp	Val	Leu	Tyr	Pro	Lys	Ala	Arg	Leu	Ala	Phe	Gln	Asn	Met	Asn
		485						490						495	
Gly	Ser	Glu	Tyr	Phe	Val	Lys	Ile	Gln	Ser	Phe	Leu	Gly	Gly	Ala	Pro
		500						505					510		
Thr	Glu	Asp	Leu	Lys	Ala	Leu	Ser	Gln	Gln	Asn	Val	Ser	Met	Asp	Leu
	515						520					525			
Ala	Thr	Phe	Met	Lys	Leu	Arg	Thr	Asp	Ala	Val	Leu	Pro	Leu	Thr	Val
	530					535					540				
Ala	Glu	Val	Gln	Lys	Leu	Leu	Gly	Pro	His	Val	Glu	Gly	Leu	Lys	Ala
545					550					555					560
Glu	Glu	Arg	His	Arg	Pro	Val	Arg	Asp	Trp	Ile	Leu	Arg	Gln	Arg	Gln
		565						570					575		
Asp	Asp	Leu	Asp	Thr	Leu	Gly	Leu	Gly	Leu	Gln	Gly	Gly	Ile	Pro	Asn
	580					585							590		
Gly	Tyr	Leu	Val	Leu	Asp	Leu	Ser	Met	Gln	Glu	Ala	Leu	Ser	Gly	Thr
	595					600					605				
Pro	Cys	Leu	Leu	Gly	Pro	Gly	Pro	Val	Leu	Thr	Val	Leu	Ala	Leu	Leu
	610					615					620				
Leu	Ala	Ser	Thr	Leu	Ala										
625					630										

<210> SEQ ID NO 168
 <211> LENGTH: 80
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(80)
 <223> OTHER INFORMATION: CD24 sequence from Genbank accession number
 AAH64619

<400> SEQUENCE: 168

Met	Gly	Arg	Ala	Met	Val	Ala	Arg	Leu	Gly	Leu	Gly	Leu	Leu	Leu	Leu
1				5					10				15		
Ala	Leu	Leu	Leu	Pro	Thr	Gln	Ile	Tyr	Ser	Ser	Glu	Thr	Thr	Thr	Gly
		20					25					30			
Thr	Ser	Ser	Asn	Ser	Ser	Gln	Ser	Thr	Ser	Asn	Ser	Gly	Leu	Ala	Pro
	35					40						45			
Asn	Pro	Thr	Asn	Ala	Thr	Thr	Lys	Ala	Ala	Gly	Gly	Ala	Leu	Gln	Ser
	50					55					60				
Thr	Ala	Ser	Leu	Phe	Val	Val	Ser	Leu	Ser	Leu	Leu	His	Leu	Tyr	Ser
65					70					75					80

<210> SEQ ID NO 169
 <211> LENGTH: 40
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(40)
<223> OTHER INFORMATION: CD22 sequence from Genbank accession number
BAA36576

<400> SEQUENCE: 169

Val Arg Arg Ala Pro Leu Ser Glu Gly Pro His Ser Leu Gly Cys Tyr
1 5 10 15
Asn Pro Met Met Glu Asp Gly Ile Ser Tyr Thr Thr Leu Arg Phe Pro
20 25 30
Glu Met Asn Ile Pro Arg Thr Gly
35 40

<210> SEQ ID NO 170
<211> LENGTH: 272
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(272)
<223> OTHER INFORMATION: CD25 sequence from Genbank accession number
NP_000408

<400> SEQUENCE: 170

Met Asp Ser Tyr Leu Leu Met Trp Gly Leu Leu Thr Phe Ile Met Val
1 5 10 15
Pro Gly Cys Gln Ala Glu Leu Cys Asp Asp Asp Pro Pro Glu Ile Pro
20 25 30
His Ala Thr Phe Lys Ala Met Ala Tyr Lys Glu Gly Thr Met Leu Asn
35 40 45
Cys Glu Cys Lys Arg Gly Phe Arg Arg Ile Lys Ser Gly Ser Leu Tyr
50 55 60
Met Leu Cys Thr Gly Asn Ser Ser His Ser Ser Trp Asp Asn Gln Cys
65 70 75 80
Gln Cys Thr Ser Ser Ala Thr Arg Asn Thr Thr Lys Gln Val Thr Pro
85 90 95
Gln Pro Glu Glu Gln Lys Glu Arg Lys Thr Thr Glu Met Gln Ser Pro
100 105 110
Met Gln Pro Val Asp Gln Ala Ser Leu Pro Gly His Cys Arg Glu Pro
115 120 125
Pro Pro Trp Glu Asn Glu Ala Thr Glu Arg Ile Tyr His Phe Val Val
130 135 140
Gly Gln Met Val Tyr Tyr Gln Cys Val Gln Gly Tyr Arg Ala Leu His
145 150 155 160
Arg Gly Pro Ala Glu Ser Val Cys Lys Met Thr His Gly Lys Thr Arg
165 170 175
Trp Thr Gln Pro Gln Leu Ile Cys Thr Gly Glu Met Glu Thr Ser Gln
180 185 190
Phe Pro Gly Glu Glu Lys Pro Gln Ala Ser Pro Glu Gly Arg Pro Glu
195 200 205
Ser Glu Thr Ser Cys Leu Val Thr Thr Thr Asp Phe Gln Ile Gln Thr
210 215 220
Glu Met Ala Ala Thr Met Glu Thr Ser Ile Phe Thr Thr Glu Tyr Gln
225 230 235 240
Val Ala Val Ala Gly Cys Val Phe Leu Leu Ile Ser Val Leu Leu Leu
245 250 255

-continued

Ser Gly Leu Thr Trp Gln Arg Arg Gln Arg Lys Ser Arg Arg Thr Ile
260 265 270

<210> SEQ ID NO 171
 <211> LENGTH: 361
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(361)
 <223> OTHER INFORMATION: CD174 sequence from Genbank accession number
 NP_000140

<400> SEQUENCE: 171

Met Asp Pro Leu Gly Ala Ala Lys Pro Gln Trp Pro Trp Arg Arg Cys
1 5 10 15
 Leu Ala Ala Leu Leu Phe Gln Leu Leu Val Ala Val Cys Phe Phe Ser
20 25 30
 Tyr Leu Arg Val Ser Arg Asp Asp Ala Thr Gly Ser Pro Arg Ala Pro
35 40 45
 Ser Gly Ser Ser Arg Gln Asp Thr Thr Pro Thr Arg Pro Thr Leu Leu
50 55 60
 Ile Leu Leu Trp Thr Trp Pro Phe His Ile Pro Val Ala Leu Ser Arg
65 70 75 80
 Cys Ser Glu Met Val Pro Gly Thr Ala Asp Cys His Ile Thr Ala Asp
85 90 95
 Arg Lys Val Tyr Pro Gln Ala Asp Thr Val Ile Val His His Trp Asp
100 105 110
 Ile Met Ser Asn Pro Lys Ser Arg Leu Pro Pro Ser Pro Arg Pro Gln
115 120 125
 Gly Gln Arg Trp Ile Trp Phe Asn Leu Glu Pro Pro Pro Asn Cys Gln
130 135 140
 His Leu Glu Ala Leu Asp Arg Tyr Phe Asn Leu Thr Met Ser Tyr Arg
145 150 155 160
 Ser Asp Ser Asp Ile Phe Thr Pro Tyr Gly Trp Leu Glu Pro Trp Ser
165 170 175
 Gly Gln Pro Ala His Pro Pro Leu Asn Leu Ser Ala Lys Thr Glu Leu
180 185 190
 Val Ala Trp Ala Val Ser Asn Trp Lys Pro Asp Ser Ala Arg Val Arg
195 200 205
 Tyr Tyr Gln Ser Leu Gln Ala His Leu Lys Val Asp Val Tyr Gly Arg
210 215 220
 Ser His Lys Pro Leu Pro Lys Gly Thr Met Met Glu Thr Leu Ser Arg
225 230 235 240
 Tyr Lys Phe Tyr Leu Ala Phe Glu Asn Ser Leu His Pro Asp Tyr Ile
245 250 255
 Thr Glu Lys Leu Trp Arg Asn Ala Leu Glu Ala Trp Ala Val Pro Val
260 265 270
 Val Leu Gly Pro Ser Arg Ser Asn Tyr Glu Arg Phe Leu Pro Pro Asp
275 280 285
 Ala Phe Ile His Val Asp Asp Phe Gln Ser Pro Lys Asp Leu Ala Arg
290 295 300
 Tyr Leu Gln Glu Leu Asp Lys Asp His Ala Arg Tyr Leu Ser Tyr Phe
305 310 315 320

-continued

```

Arg Trp Arg Glu Thr Leu Arg Pro Arg Ser Phe Ser Trp Ala Leu Asp
      325              330              335

Phe Cys Lys Ala Cys Trp Lys Leu Gln Gln Glu Ser Arg Tyr Gln Thr
      340              345              350

Val Arg Ser Ile Ala Ala Trp Phe Thr
      355              360

<210> SEQ ID NO 172
<211> LENGTH: 420
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(420)
<223> OTHER INFORMATION: TPBG sequence from Genbank accession number
      CAA09930

<400> SEQUENCE: 172

Met Pro Gly Gly Cys Ser Arg Gly Pro Ala Ala Gly Asp Gly Arg Leu
 1              5              10              15

Arg Leu Ala Arg Leu Ala Leu Val Leu Leu Gly Trp Val Ser Ser Ser
      20              25              30

Ser Pro Thr Ser Ser Ala Ser Ser Phe Ser Ser Ser Ala Pro Phe Leu
      35              40              45

Ala Ser Ala Val Ser Ala Gln Pro Pro Leu Pro Asp Gln Cys Pro Ala
      50              55              60

Leu Cys Glu Cys Ser Glu Ala Ala Arg Thr Val Lys Cys Val Asn Arg
65              70              75              80

Asn Leu Thr Glu Val Pro Thr Asp Leu Pro Ala Tyr Val Arg Asn Leu
      85              90              95

Phe Leu Thr Gly Asn Gln Leu Ala Val Leu Pro Ala Gly Ala Phe Ala
      100             105             110

Arg Arg Pro Pro Leu Ala Glu Leu Ala Ala Leu Asn Leu Ser Gly Ser
      115             120             125

Arg Leu Asp Glu Val Arg Ala Gly Ala Phe Glu His Leu Pro Ser Leu
      130             135             140

Arg Gln Leu Asp Leu Ser His Asn Pro Leu Ala Asp Leu Ser Pro Phe
      145             150             155             160

Ala Phe Ser Gly Ser Asn Ala Ser Val Ser Ala Pro Ser Pro Leu Val
      165             170             175

Glu Leu Ile Leu Asn His Ile Val Pro Pro Glu Asp Glu Arg Gln Asn
      180             185             190

Arg Ser Phe Glu Gly Met Val Val Ala Ala Leu Leu Ala Gly Arg Ala
      195             200             205

Leu Gln Gly Leu Arg Arg Leu Glu Leu Ala Ser Asn His Phe Leu Tyr
      210             215             220

Leu Pro Arg Asp Val Leu Ala Gln Leu Pro Ser Leu Arg His Leu Asp
      225             230             235             240

Leu Ser Asn Asn Ser Leu Val Ser Leu Thr Tyr Val Ser Phe Arg Asn
      245             250             255

Leu Thr His Leu Glu Ser Leu His Leu Glu Asp Asn Ala Leu Lys Val
      260             265             270

Leu His Asn Gly Thr Leu Ala Glu Leu Gln Gly Leu Pro His Ile Arg
      275             280             285

Val Phe Leu Asp Asn Asn Pro Trp Val Cys Asp Cys His Met Ala Asp

```

-continued

290	295	300
Met Val Thr Trp Leu Lys Glu Thr Glu Val Val Gln Gly Lys Asp Arg		
305	310	315
Leu Thr Cys Ala Tyr Pro Glu Lys Met Arg Asn Arg Val Leu Leu Glu		
	325	330
Leu Asn Ser Ala Asp Leu Asp Cys Asp Pro Ile Leu Pro Pro Ser Leu		
	340	345
Gln Thr Ser Tyr Val Phe Leu Gly Ile Val Leu Ala Leu Ile Gly Ala		
	355	360
Ile Phe Leu Leu Val Leu Tyr Leu Asn Arg Lys Gly Ile Lys Lys Trp		
	370	375
Met His Asn Ile Arg Asp Ala Cys Arg Asp His Met Glu Gly Tyr His		
	385	390
Tyr Arg Tyr Glu Ile Asn Ala Asp Pro Arg Leu Thr Asn Leu Ser Ser		
	405	410
Asn Ser Asp Val		
	420	

<210> SEQ ID NO 173
 <211> LENGTH: 848
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(848)
 <223> OTHER INFORMATION: CD56 sequence from Genbank accession number NP_000606

<400> SEQUENCE: 173

Met Leu Gln Thr Lys Asp Leu Ile Trp Thr Leu Phe Phe Leu Gly Thr		
1	5	10
Ala Val Ser Leu Gln Val Asp Ile Val Pro Ser Gln Gly Glu Ile Ser		
	20	25
Val Gly Glu Ser Lys Phe Phe Leu Cys Gln Val Ala Gly Asp Ala Lys		
	35	40
Asp Lys Asp Ile Ser Trp Phe Ser Pro Asn Gly Glu Lys Leu Thr Pro		
	50	55
Asn Gln Gln Arg Ile Ser Val Val Trp Asn Asp Asp Ser Ser Ser Thr		
	65	70
Leu Thr Ile Tyr Asn Ala Asn Ile Asp Asp Ala Gly Ile Tyr Lys Cys		
	85	90
Val Val Thr Gly Glu Asp Gly Ser Glu Ser Glu Ala Thr Val Asn Val		
	100	105
Lys Ile Phe Gln Lys Leu Met Phe Lys Asn Ala Pro Thr Pro Gln Glu		
	115	120
Phe Arg Glu Gly Glu Asp Ala Val Ile Val Cys Asp Val Val Ser Ser		
	130	135
Leu Pro Pro Thr Ile Ile Trp Lys His Lys Gly Arg Asp Val Ile Leu		
	145	150
Lys Lys Asp Val Arg Phe Ile Val Leu Ser Asn Asn Tyr Leu Gln Ile		
	165	170
Arg Gly Ile Lys Lys Thr Asp Glu Gly Thr Tyr Arg Cys Glu Gly Arg		
	180	185
Ile Leu Ala Arg Gly Glu Ile Asn Phe Lys Asp Ile Gln Val Ile Val		
	195	200
		205

-continued

Asn Val Pro Pro Thr Ile Gln Ala Arg Gln Asn Ile Val Asn Ala Thr	
210 215 220	
Ala Asn Leu Gly Gln Ser Val Thr Leu Val Cys Asp Ala Glu Gly Phe	
225 230 235 240	
Pro Glu Pro Thr Met Ser Trp Thr Lys Asp Gly Glu Gln Ile Glu Gln	
245 250 255	
Glu Glu Asp Asp Glu Lys Tyr Ile Phe Ser Asp Asp Ser Ser Gln Leu	
260 265 270	
Thr Ile Lys Lys Val Asp Lys Asn Asp Glu Ala Glu Tyr Ile Cys Ile	
275 280 285	
Ala Glu Asn Lys Ala Gly Glu Gln Asp Ala Thr Ile His Leu Lys Val	
290 295 300	
Phe Ala Lys Pro Lys Ile Thr Tyr Val Glu Asn Gln Thr Ala Met Glu	
305 310 315 320	
Leu Glu Glu Gln Val Thr Leu Thr Cys Glu Ala Ser Gly Asp Pro Ile	
325 330 335	
Pro Ser Ile Thr Trp Arg Thr Ser Thr Arg Asn Ile Ser Ser Glu Glu	
340 345 350	
Lys Thr Leu Asp Gly His Met Val Val Arg Ser His Ala Arg Val Ser	
355 360 365	
Ser Leu Thr Leu Lys Ser Ile Gln Tyr Thr Asp Ala Gly Glu Tyr Ile	
370 375 380	
Cys Thr Ala Ser Asn Thr Ile Gly Gln Asp Ser Gln Ser Met Tyr Leu	
385 390 395 400	
Glu Val Gln Tyr Ala Pro Lys Leu Gln Gly Pro Val Ala Val Tyr Thr	
405 410 415	
Trp Glu Gly Asn Gln Val Asn Ile Thr Cys Glu Val Phe Ala Tyr Pro	
420 425 430	
Ser Ala Thr Ile Ser Trp Phe Arg Asp Gly Gln Leu Leu Pro Ser Ser	
435 440 445	
Asn Tyr Ser Asn Ile Lys Ile Tyr Asn Thr Pro Ser Ala Ser Tyr Leu	
450 455 460	
Glu Val Thr Pro Asp Ser Glu Asn Asp Phe Gly Asn Tyr Asn Cys Thr	
465 470 475 480	
Ala Val Asn Arg Ile Gly Gln Glu Ser Leu Glu Phe Ile Leu Val Gln	
485 490 495	
Ala Asp Thr Pro Ser Ser Pro Ser Ile Asp Gln Val Glu Pro Tyr Ser	
500 505 510	
Ser Thr Ala Gln Val Gln Phe Asp Glu Pro Glu Ala Thr Gly Gly Val	
515 520 525	
Pro Ile Leu Lys Tyr Lys Ala Glu Trp Arg Ala Val Gly Glu Glu Val	
530 535 540	
Trp His Ser Lys Trp Tyr Asp Ala Lys Glu Ala Ser Met Glu Gly Ile	
545 550 555 560	
Val Thr Ile Val Gly Leu Lys Pro Glu Thr Thr Tyr Ala Val Arg Leu	
565 570 575	
Ala Ala Leu Asn Gly Lys Gly Leu Gly Glu Ile Ser Ala Ala Ser Glu	
580 585 590	
Phe Lys Thr Gln Pro Val Gln Gly Glu Pro Ser Ala Pro Lys Leu Glu	
595 600 605	
Gly Gln Met Gly Glu Asp Gly Asn Ser Ile Lys Val Asn Leu Ile Lys	

-continued

610	615	620
Gln Asp Asp Gly Gly Ser Pro Ile Arg His Tyr Leu Val Arg Tyr Arg 625 630 635 640		
Ala Leu Ser Ser Glu Trp Lys Pro Glu Ile Arg Leu Pro Ser Gly Ser 645 650 655		
Asp His Val Met Leu Lys Ser Leu Asp Trp Asn Ala Glu Tyr Glu Val 660 665 670		
Tyr Val Val Ala Glu Asn Gln Gln Gly Lys Ser Lys Ala Ala His Phe 675 680 685		
Val Phe Arg Thr Ser Ala Gln Pro Thr Ala Ile Pro Ala Asn Gly Ser 690 695 700		
Pro Thr Ser Gly Leu Ser Thr Gly Ala Ile Val Gly Ile Leu Ile Val 705 710 715 720		
Ile Phe Val Leu Leu Leu Val Val Val Asp Ile Thr Cys Tyr Phe Leu 725 730 735		
Asn Lys Cys Gly Leu Phe Met Cys Ile Ala Val Asn Leu Cys Gly Lys 740 745 750		
Ala Gly Pro Gly Ala Lys Gly Lys Asp Met Glu Glu Gly Lys Ala Ala 755 760 765		
Phe Ser Lys Asp Glu Ser Lys Glu Pro Ile Val Glu Val Arg Thr Glu 770 775 780		
Glu Glu Arg Thr Pro Asn His Asp Gly Gly Lys His Thr Glu Pro Asn 785 790 795 800		
Glu Thr Thr Pro Leu Thr Glu Pro Glu Lys Gly Pro Val Glu Ala Lys 805 810 815		
Pro Glu Cys Gln Glu Thr Glu Thr Lys Pro Ala Pro Ala Glu Val Lys 820 825 830		
Thr Val Pro Asn Asp Ala Thr Gln Thr Lys Glu Asn Glu Ser Lys Ala 835 840 845		

<210> SEQ ID NO 174
 <211> LENGTH: 275
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(275)
 <223> OTHER INFORMATION: C-type lectin-like molecule-1 from Genbank
 accession number AAT11783

<400> SEQUENCE: 174

Met Trp Ile Asp Phe Phe Thr Tyr Ser Ser Met Ser Glu Glu Val Thr 1 5 10 15
Tyr Ala Asp Leu Gln Phe Gln Asn Ser Ser Glu Met Glu Lys Ile Pro 20 25 30
Glu Ile Gly Lys Phe Gly Glu Lys Ala Pro Pro Ala Pro Ser His Val 35 40 45
Trp Arg Pro Ala Ala Leu Phe Leu Thr Leu Leu Cys Leu Leu Leu Leu 50 55 60
Ile Gly Leu Gly Val Leu Ala Ser Met Phe His Val Thr Leu Lys Ile 65 70 75 80
Glu Met Lys Lys Met Asn Lys Leu Gln Asn Ile Ser Glu Glu Leu Gln 85 90 95
Arg Asn Ile Ser Leu Gln Leu Met Ser Asn Met Asn Ile Ser Asn Lys 100 105 110

-continued

```

Ile Arg Asn Leu Ser Thr Thr Leu Gln Thr Ile Ala Thr Lys Leu Cys
    115                      120                      125

Arg Glu Leu Tyr Ser Lys Glu Gln Glu His Lys Cys Lys Pro Cys Pro
    130                      135                      140

Arg Arg Trp Ile Trp His Lys Asp Ser Cys Tyr Phe Leu Ser Asp Asp
    145                      150                      155                      160

Val Gln Thr Trp Gln Glu Ser Lys Met Ala Cys Ala Ala Gln Asn Ala
    165                      170                      175

Ser Leu Leu Lys Ile Asn Asn Lys Asn Ala Leu Glu Phe Ile Lys Ser
    180                      185                      190

Gln Ser Arg Ser Tyr Asp Tyr Trp Leu Gly Leu Ser Pro Glu Glu Asp
    195                      200                      205

Ser Thr Arg Gly Met Arg Val Asp Asn Ile Ile Asn Ser Ser Ala Trp
    210                      215                      220

Val Ile Arg Asn Ala Pro Asp Leu Asn Asn Met Tyr Cys Gly Tyr Ile
    225                      230                      235                      240

Asn Arg Leu Tyr Val Gln Tyr Tyr His Cys Thr Tyr Lys Gln Arg Met
    245                      250                      255

Ile Cys Glu Lys Met Ala Asn Pro Val Gln Leu Gly Ser Thr Tyr Phe
    260                      265                      270

Arg Glu Ala
    275

```

```

<210> SEQ ID NO 175
<211> LENGTH: 613
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)..(613)
<223> OTHER INFORMATION: PE variant from Genbank Accession number GI
    17943391, version 1IKQ_A

```

```

<400> SEQUENCE: 175

```

```

Ala Glu Glu Ala Phe Asp Leu Trp Asn Glu Cys Ala Lys Ala Cys Val
  1                      5                      10                      15

Leu Asp Leu Lys Asp Gly Val Arg Ser Ser Arg Met Ser Val Asp Pro
    20                      25                      30

Ala Ile Ala Asp Thr Asn Gly Gln Gly Val Leu His Tyr Ser Met Val
    35                      40                      45

Leu Glu Gly Gly Asn Asp Ala Leu Lys Leu Ala Ile Asp Asn Ala Leu
    50                      55                      60

Ser Ile Thr Ser Asp Gly Leu Thr Ile Arg Leu Glu Gly Gly Val Glu
    65                      70                      75                      80

Pro Asn Lys Pro Val Arg Tyr Ser Tyr Thr Arg Gln Ala Arg Gly Ser
    85                      90                      95

Trp Ser Leu Asn Trp Leu Val Pro Ile Gly His Glu Lys Pro Ser Asn
    100                     105                     110

Ile Lys Val Phe Ile His Glu Leu Asn Ala Gly Asn Gln Leu Ser His
    115                     120                     125

Met Ser Pro Ile Tyr Thr Ile Glu Met Gly Asp Glu Leu Leu Ala Lys
    130                     135                     140

Leu Ala Arg Asp Ala Thr Phe Phe Val Arg Ala His Glu Ser Asn Glu
    145                     150                     155                     160

```

-continued

Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met
				165						170					175
Ala	Gln	Ala	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser
				180					185					190	
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr
		195						200				205			
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile
		210						215				220			
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys
225						230					235				240
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu
				245						250					255
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
				260						265				270	
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly
		275						280					285		
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
		290						295				300			
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
305						310					315				320
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
				325						330					335
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
				340					345					350	
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Ala	Asp	Val	Val
		355						360					365		
Ser	Leu	Thr	Cys	Pro	Val	Ala	Ala	Gly	Glu	Cys	Ala	Gly	Pro	Ala	Asp
		370						375				380			
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
385						390					395				400
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
				405						410					415
Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg
			420						425					430	
Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln
		435						440					445		
Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala
		450					455					460			
Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly
465						470					475				480
Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly
				485						490				495	
Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr
				500					505					510	
Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu
			515						520				525		
Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly
		530					535					540			
Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro	Leu
545						550					555				560
Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg
				565						570					575

-continued

Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln
 580 585 590

Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
 595 600 605

Arg Glu Asp Leu Lys
 610

<210> SEQ ID NO 176
 <211> LENGTH: 10
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Modified Kozak consensus sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(10)
 <223> OTHER INFORMATION: consensus sequence

<400> SEQUENCE: 176

gccaccatgg

10

<210> SEQ ID NO 177
 <211> LENGTH: 347
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pIEX02-228 PE-A amino acid substitution mutant
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (141)..(141)
 <223> OTHER INFORMATION: Amino acid change Ile-141 to Ala-141
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (152)..(152)
 <223> OTHER INFORMATION: Amino acid change Thr-152 to Arg-152
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (197)..(197)
 <223> OTHER INFORMATION: Amino acid change Asp-197 to Lys-197
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (241)..(241)
 <223> OTHER INFORMATION: Amino acid change Ser-241 to Thr-241
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (326)..(326)
 <223> OTHER INFORMATION: Amino acid change Gln-326 to Glu-326

<400> SEQUENCE: 177

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
 1 5 10 15

Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
 20 25 30

Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
 35 40 45

Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
 50 55 60

Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
 65 70 75 80

Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
 85 90 95

Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
 100 105 110

-continued

Ala	Ser	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr
		115					120					125			
Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ala	Ser	Phe	Ser
		130				135					140				
Thr	Arg	Gly	Thr	Gln	Asn	Trp	Arg	Val	Glu	Arg	Leu	Leu	Gln	Ala	His
145				150					155					160	
Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr
			165					170					175		
Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg
		180					185					190			
Ser	Gln	Asp	Leu	Lys	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp
		195				200					205				
Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg
	210					215					220				
Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser
225				230					235					240	
Thr	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu
			245					250						255	
Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg
		260					265					270			
Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr
	275					280					285				
Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala
	290				295					300					
Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser
305				310				315						320	
Ile	Pro	Asp	Lys	Glu	Glu	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser
			325					330					335		
Gln	Pro	Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys					
		340					345								

<210> SEQ ID NO 178
 <211> LENGTH: 347
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pIEX02-244 PE-A amino acid substitution mutant
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (141)..(141)
 <223> OTHER INFORMATION: Amino acid change Ile-141 to Thr-141
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (152)..(152)
 <223> OTHER INFORMATION: Amino acid change Thr-152 to Ala-152
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (192)..(192)
 <223> OTHER INFORMATION: Amino acid change Arg-192 to Ala-192
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (197)..(197)
 <223> OTHER INFORMATION: Amino acid change Asp-197 to Lys-197
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (241)..(241)
 <223> OTHER INFORMATION: Amino acid change Ser-241 to Thr-241
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (326)..(326)
 <223> OTHER INFORMATION: Amino acid change Gln-326 to Glu-326

-continued

<400> SEQUENCE: 178

```

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1      5      10      15
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20     25     30
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35     40     45
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50     55     60
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65     70     75     80
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Glu
85     90     95
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100    105    110
Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
115    120    125
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Thr Ser Phe Ser
130    135    140
Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg Leu Leu Gln Ala His
145    150    155    160
Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
165    170    175
Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Ala
180    185    190
Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
195    200    205
Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
210    215    220
Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
225    230    235    240
Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
245    250    255
Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
260    265    270
Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr
275    280    285
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
290    295    300
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
305    310    315    320
Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
325    330    335
Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
340    345

```

<210> SEQ ID NO 179

<211> LENGTH: 347

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pIEX02-246 PE-A amino acid substitution mutant

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

-continued

```

<222> LOCATION: (141)..(141)
<223> OTHER INFORMATION: Amino acid change Ile-141 to Ala-141
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (152)..(152)
<223> OTHER INFORMATION: Amino acid change Thr-152 to Ala-152
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (192)..(192)
<223> OTHER INFORMATION: Amino acid change Arg-192 to Ala-192
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (197)..(197)
<223> OTHER INFORMATION: Amino acid change Asp-197 to Lys-197
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (241)..(241)
<223> OTHER INFORMATION: Amino acid change Ser-241 to Thr-241
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (326)..(326)
<223> OTHER INFORMATION: Amino acid change Gln-326 to Glu-326

<400> SEQUENCE: 179

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1          5          10          15

Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20          25          30

Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35          40          45

Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50          55          60

Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65          70          75          80

Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
85          90          95

Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100         105         110

Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
115         120         125

Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ala Ser Phe Ser
130         135         140

Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg Leu Leu Gln Ala His
145         150         155         160

Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
165         170         175

Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Ala
180         185         190

Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
195         200         205

Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
210         215         220

Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
225         230         235         240

Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
245         250         255

Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
260         265         270

Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr

```

-continued

275	280	285
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala 290 295 300		
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser 305 310 315 320		
Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser 325 330 335		
Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys 340 345		
<210> SEQ ID NO 180		
<211> LENGTH: 347		
<212> TYPE: PRT		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: PE38 NULL MUTANT (E287D)		
<220> FEATURE:		
<221> NAME/KEY: MISC_FEATURE		
<222> LOCATION: (287)..(287)		
<223> OTHER INFORMATION: Amino acid change Glu-287 to Asp-287		
<400> SEQUENCE: 180		
Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His 1 5 10 15		
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu 20 25 30		
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr 35 40 45		
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn 50 55 60		
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg 65 70 75 80		
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu 85 90 95		
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala 100 105 110		
Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr 115 120 125		
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser 130 135 140		
Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His 145 150 155 160		
Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr 165 170 175		
Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg 180 185 190		
Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp 195 200 205		
Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg 210 215 220		
Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser 225 230 235 240		
Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu 245 250 255		
Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg		

-continued

260	265	270
Leu Asp Ala Ile Thr Gly Pro Glu Glu Gly Gly Arg Leu Asp Thr		
275	280	285
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala		
290	295	300
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser		
305	310	315
Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser		
325	330	335
Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys		
340	345	

<210> SEQ ID NO 181
 <211> LENGTH: 347
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PE VARIANT 238 (pIEX02-228 with E287D)
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (141)..(141)
 <223> OTHER INFORMATION: Amino acid change Ile-141 to Ala-141
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (152)..(152)
 <223> OTHER INFORMATION: Amino acid change Thr-152 to Arg-152
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (197)..(197)
 <223> OTHER INFORMATION: Amino acid change Asp-197 to Lys-197
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (241)..(241)
 <223> OTHER INFORMATION: Amino acid change Ser-241 to Thr-241
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (287)..(287)
 <223> OTHER INFORMATION: Amino acid change Glu-287 to Asp-287
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (326)..(326)
 <223> OTHER INFORMATION: Amino acid change Gln-326 to Glu-326

 <400> SEQUENCE: 181

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1 5 10 15
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20 25 30
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35 40 45
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50 55 60
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65 70 75 80
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
85 90 95
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100 105 110
Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
115 120 125
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ala Ser Phe Ser
130 135 140

-continued

Thr Arg Gly Thr Gln Asn Trp Arg Val Glu Arg Leu Leu Gln Ala His
 145 150 155 160
 Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
 165 170 175
 Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
 180 185 190
 Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
 195 200 205
 Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
 210 215 220
 Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
 225 230 235 240
 Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
 245 250 255
 Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
 260 265 270
 Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Asp Thr
 275 280 285
 Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
 290 295 300
 Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
 305 310 315 320
 Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
 325 330 335
 Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 340 345

<210> SEQ ID NO 182
 <211> LENGTH: 347
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PE VARIANT 245 (pIEX02-244 with E287D)
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (141)..(141)
 <223> OTHER INFORMATION: Amino acid change Ile-141 to Thr-141
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (152)..(152)
 <223> OTHER INFORMATION: Amino acid change Thr-152 to Ala-152
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (192)..(192)
 <223> OTHER INFORMATION: Amino acid change Arg-192 to Ala-192
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (197)..(197)
 <223> OTHER INFORMATION: Amino acid change Asp-197 to Lys-197
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (241)..(241)
 <223> OTHER INFORMATION: Amino acid change Ser-241 to Thr-241
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (287)..(287)
 <223> OTHER INFORMATION: Amino acid change Glu-287 to Asp-287
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (326)..(326)
 <223> OTHER INFORMATION: Amino acid change Gln-326 to Glu-326
 <400> SEQUENCE: 182

-continued

```

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1          5          10          15
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20          25          30
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35          40          45
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50          55          60
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65          70          75          80
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
85          90          95
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100         105         110
Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
115         120         125
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Thr Ser Phe Ser
130         135         140
Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg Leu Leu Gln Ala His
145         150         155         160
Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
165         170         175
Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Ala
180         185         190
Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
195         200         205
Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
210         215         220
Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
225         230         235         240
Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
245         250         255
Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
260         265         270
Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Asp Thr
275         280         285
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
290         295         300
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
305         310         315         320
Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
325         330         335
Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
340         345

```

<210> SEQ ID NO 183

<211> LENGTH: 347

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PE VARIANT 247(pIEX02-246 with E287D)

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (141) .. (141)

-continued

```

<223> OTHER INFORMATION: Amino acid change Ile-141 to Ala-141
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (152)..(152)
<223> OTHER INFORMATION: Amino acid change Thr-152 to Ala-152
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (192)..(192)
<223> OTHER INFORMATION: Amino acid change Arg-192 to Ala-192
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (197)..(197)
<223> OTHER INFORMATION: Amino acid change Asp-197 to Lys-197
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (241)..(241)
<223> OTHER INFORMATION: Amino acid change Ser-241 to Thr-241
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (287)..(287)
<223> OTHER INFORMATION: Amino acid change Glu-287 to Asp-287
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (326)..(326)
<223> OTHER INFORMATION: Amino acid change Gln-326 to Glu-326

<400> SEQUENCE: 183

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1          5          10          15

Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
20          25          30

Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
35          40          45

Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
50          55          60

Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65          70          75          80

Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
85          90          95

Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
100         105         110

Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
115         120         125

Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ala Ser Phe Ser
130         135         140

Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg Leu Leu Gln Ala His
145         150         155         160

Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
165         170         175

Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Ala
180         185         190

Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
195         200         205

Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
210         215         220

Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
225         230         235         240

Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
245         250         255

Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg

```

-continued

260	265	270
Leu Asp Ala Ile Thr Gly Pro Glu Glu Gly Gly Arg Leu Asp Thr 275 280 285		
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala 290 295 300		
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser 305 310 315 320		
Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser 325 330 335		
Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys 340 345		
 <210> SEQ ID NO 184 <211> LENGTH: 611 <212> TYPE: PRT <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: FUSION RFB4-WT PE38-HIS8-EDLK <400> SEQUENCE: 184		
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly 1 5 10 15		
Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ile Tyr 20 25 30		
Asp Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val 35 40 45		
Ala Tyr Ile Ser Ser Gly Gly Gly Thr Thr Tyr Tyr Pro Asp Thr Val 50 55 60		
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr 65 70 75 80		
Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys 85 90 95		
Ala Arg His Ser Gly Tyr Gly Ser Ser Tyr Gly Val Leu Phe Ala Tyr 100 105 110		
Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Gly Gly Gly Gly Ser 115 120 125		
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile Gln Met Thr Gln 130 135 140		
Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly Asp Arg Val Thr Ile Ser 145 150 155 160		
Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln 165 170 175		
Lys Pro Asp Gly Thr Val Lys Leu Leu Ile Tyr Tyr Thr Ser Ile Leu 180 185 190		
His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp 195 200 205		
Tyr Ser Leu Thr Ile Ser Asn Leu Glu Gln Glu Asp Phe Ala Thr Tyr 210 215 220		
Phe Cys Gln Gln Gly Asn Thr Leu Pro Trp Thr Phe Gly Gly Gly Thr 225 230 235 240		
Lys Leu Glu Ile Lys Ser Ser Gly Leu Val Pro Arg Gly Ser His Met 245 250 255		
Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His 260 265 270		

-continued

Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
 275 280 285
 Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
 290 295 300
 Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
 305 310 315 320
 Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
 325 330 335
 Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
 340 345 350
 Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
 355 360 365
 Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
 370 375 380
 Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Ile Ser Phe Ser
 385 390 395 400
 Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
 405 410 415
 Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
 420 425 430
 Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
 435 440 445
 Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
 450 455 460
 Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
 465 470 475 480
 Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
 485 490 495
 Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu
 500 505 510
 Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
 515 520 525
 Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr
 530 535 540
 Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
 545 550 555 560
 Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
 565 570 575
 Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
 580 585 590
 Gln Pro Gly Lys Pro Pro Arg His His His His His His His Glu
 595 600 605
 Asp Leu Lys
 610

<210> SEQ ID NO 185

<211> LENGTH: 614

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: FUSION RFB4-HIS6-WT PE38-EDLK

<400> SEQUENCE: 185

-continued

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	1	5	10	15
Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile	Tyr	20	25	30	
Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Arg	Leu	Glu	Trp	Val	35	40	45	
Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	50	55	60	
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu	Tyr	65	70	75	80
Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	85	90	95	
Ala	Arg	His	Ser	Gly	Tyr	Gly	Ser	Ser	Tyr	Gly	Val	Leu	Phe	Ala	Tyr	100	105	110	
Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	115	120	125	
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile	Gln	Met	Thr	Gln	130	135	140	
Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu	Gly	Asp	Arg	Val	Thr	Ile	Ser	145	150	155	160
Cys	Arg	Ala	Ser	Gln	Asp	Ile	Ser	Asn	Tyr	Leu	Asn	Trp	Tyr	Gln	Gln	165	170	175	
Lys	Pro	Asp	Gly	Thr	Val	Lys	Leu	Leu	Ile	Tyr	Tyr	Thr	Ser	Ile	Leu	180	185	190	
His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	195	200	205	
Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu	Gln	Glu	Asp	Phe	Ala	Thr	Tyr	210	215	220	
Phe	Cys	Gln	Gln	Gly	Asn	Thr	Leu	Pro	Trp	Thr	Phe	Gly	Gly	Gly	Thr	225	230	235	240
Lys	Leu	Glu	Ile	Lys	Ala	His	Gly	Gly	Ser	His	His	His	His	His	His	245	250	255	
Ser	Ser	Gly	Leu	Val	Pro	Arg	Gly	Ser	His	Met	Pro	Glu	Gly	Gly	Ser	260	265	270	
Leu	Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	275	280	285	
Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	290	295	300	
Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	305	310	315	320
Ser	Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	325	330	335	
Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	340	345	350	
Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	355	360	365	
Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Ser	Gly	Pro	Ala	370	375	380	
Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	385	390	395	400
Phe	Leu	Gly	Asp	Gly	Gly	Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	405	410	415	

-continued

Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu
 420 425 430
 Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala
 435 440 445
 Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp
 450 455 460
 Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr
 465 470 475 480
 Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn
 485 490 495
 Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe
 500 505 510
 Tyr Arg Thr Gly Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val
 515 520 525
 Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr
 530 535 540
 Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro
 545 550 555 560
 Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro
 565 570 575
 Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu
 580 585 590
 Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro
 595 600 605
 Pro Arg Glu Asp Leu Lys
 610

<210> SEQ ID NO 186
 <211> LENGTH: 611
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: FUSION RFB4-VARIANT 244-HIS8-EDLK

<400> SEQUENCE: 186

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ile Tyr
 20 25 30
 Asp Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
 35 40 45
 Ala Tyr Ile Ser Ser Gly Gly Gly Thr Thr Tyr Tyr Pro Asp Thr Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
 85 90 95
 Ala Arg His Ser Gly Tyr Gly Ser Ser Tyr Gly Val Leu Phe Ala Tyr
 100 105 110
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Gly Gly Gly Gly Ser
 115 120 125
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile Gln Met Thr Gln
 130 135 140

-continued

Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu	Gly	Asp	Arg	Val	Thr	Ile	Ser	145	150	155	160
Cys	Arg	Ala	Ser	Gln	Asp	Ile	Ser	Asn	Tyr	Leu	Asn	Trp	Tyr	Gln	Gln	165	170	175	
Lys	Pro	Asp	Gly	Thr	Val	Lys	Leu	Leu	Ile	Tyr	Tyr	Thr	Ser	Ile	Leu	180	185	190	
His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	195	200	205	
Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu	Gln	Glu	Asp	Phe	Ala	Thr	Tyr	210	215	220	
Phe	Cys	Gln	Gln	Gly	Asn	Thr	Leu	Pro	Trp	Thr	Phe	Gly	Gly	Gly	Thr	225	230	235	240
Lys	Leu	Glu	Ile	Lys	Ser	Ser	Gly	Leu	Val	Pro	Arg	Gly	Ser	His	Met	245	250	255	
Pro	Glu	Gly	Gly	Ser	Leu	Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	260	265	270	
Leu	Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	275	280	285	
Gln	Leu	Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	290	295	300	
Leu	Ala	Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	305	310	315	320
Ala	Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	325	330	335	
Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	340	345	350	
Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	355	360	365	
Ala	Ser	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	370	375	380	
Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Thr	Ser	Phe	Ser	385	390	395	400
Thr	Arg	Gly	Thr	Gln	Asn	Trp	Ala	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	405	410	415	
Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	420	425	430	
Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Ala	435	440	445	
Ser	Gln	Asp	Leu	Lys	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	450	455	460	
Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	465	470	475	480
Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	485	490	495	
Thr	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu	500	505	510	
Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	515	520	525	
Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	530	535	540	
Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	545	550	555	560

-continued

Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
 565 570 575

Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
 580 585 590

Gln Pro Gly Lys Pro Pro Arg His His His His His His His His Glu
 595 600 605

Asp Leu Lys
 610

<210> SEQ ID NO 187

<211> LENGTH: 614

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: FUSION RPB4-HIS6-WT PE38-EDLK

<400> SEQUENCE: 187

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ile Tyr
 20 25 30

Asp Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
 35 40 45

Ala Tyr Ile Ser Ser Gly Gly Gly Thr Thr Tyr Tyr Pro Asp Thr Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
 85 90 95

Ala Arg His Ser Gly Tyr Gly Ser Ser Tyr Gly Val Leu Phe Ala Tyr
 100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Gly Gly Gly Ser
 115 120 125

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile Gln Met Thr Gln
 130 135 140

Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly Asp Arg Val Thr Ile Ser
 145 150 155 160

Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln
 165 170 175

Lys Pro Asp Gly Thr Val Lys Leu Leu Ile Tyr Tyr Thr Ser Ile Leu
 180 185 190

His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp
 195 200 205

Tyr Ser Leu Thr Ile Ser Asn Leu Glu Gln Glu Asp Phe Ala Thr Tyr
 210 215 220

Phe Cys Gln Gln Gly Asn Thr Leu Pro Trp Thr Phe Gly Gly Gly Thr
 225 230 235 240

Lys Leu Glu Ile Lys Ala His Gly Gly Ser His His His His His His
 245 250 255

Ser Ser Gly Leu Val Pro Arg Gly Ser His Met Pro Glu Gly Gly Ser
 260 265 270

Leu Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro Leu Glu Thr
 275 280 285

-continued

Phe	Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys
290						295					300				
Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu
305					310					315					320
Ser	Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro
			325						330					335	
Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln
			340					345					350		
Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val
		355					360					365			
Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Ser	Gly	Pro	Ala
	370					375						380			
Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu
385					390					395					400
Phe	Leu	Gly	Asp	Gly	Gly	Asp	Thr	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln
			405						410					415	
Asn	Trp	Ala	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu
			420					425					430		
Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala
		435					440					445			
Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Ala	Ser	Gln	Asp	Leu	Lys
	450					455					460				
Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr
465					470					475					480
Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn
			485						490					495	
Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Thr	Leu	Pro	Gly	Phe
			500					505					510		
Tyr	Arg	Thr	Gly	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val
		515					520					525			
Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr
	530					535					540				
Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro
545					550					555					560
Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro
			565						570					575	
Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Lys	Glu
			580					585					590		
Glu	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro
	595						600					605			
Pro	Arg	Glu	Asp	Leu	Lys										
	610														

<210> SEQ ID NO 188

<211> LENGTH: 245

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RPB4 SCFV

<400> SEQUENCE: 188

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile	Tyr
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

-continued

20					25					30					
Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Arg	Leu	Glu	Trp	Val
		35					40					45			
Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	
	50					55				60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu	Tyr
65					70					75				80	
Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys
			85						90					95	
Ala	Arg	His	Ser	Gly	Tyr	Gly	Ser	Ser	Tyr	Gly	Val	Leu	Phe	Ala	Tyr
			100					105					110		
Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser
	115						120					125			
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Ser	Asp	Ile	Gln	Met	Thr	Gln	
	130					135					140				
Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu	Gly	Asp	Arg	Val	Thr	Ile	Ser
145					150					155					160
Cys	Arg	Ala	Ser	Gln	Asp	Ile	Ser	Asn	Tyr	Leu	Asn	Trp	Tyr	Gln	Gln
			165						170					175	
Lys	Pro	Asp	Gly	Thr	Val	Lys	Leu	Leu	Ile	Tyr	Tyr	Thr	Ser	Ile	Leu
			180					185					190		
His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp
		195					200					205			
Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu	Gln	Glu	Asp	Phe	Ala	Thr	Tyr
		210				215					220				
Phe	Cys	Gln	Gln	Gly	Asn	Thr	Leu	Pro	Trp	Thr	Phe	Gly	Gly	Gly	Thr
225					230					235					240
Lys	Leu	Glu	Ile	Lys											
				245											

<210> SEQ ID NO 189

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PE38 WT as expressed by pIEX02-001 (pET14b-K PE38)

<400> SEQUENCE: 189

```

atgggcagca gccatcatca tcatcatcac agcagcggcc tggtgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg    120
gaaaccttca ccggtcaccg tcagccgcgt ggttggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgaa ccaagtagat    240
cagggtcatcc gtaacgcgct ggcaagcccg ggttcgggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacaggc tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc agggtagcgg taacgacgaa gctggcgctg cgagcgggtcc tgcgactcc    420
ggtagcgtc tgctggaacg taactaccg accggtgcag aatttctggg tgatggcggc    480
gatattcttt tttctaccg cggcaccag aactggaccg ttgaacgtct gctgcaggcg    540
caccgtcaac tggaagaacg cggttacgtc ttcgtaggtt accacggtac ctctctggaa    600
gctgctcagt ctatcgtgtt cggtggtgta cgtgctcgta gccaggacct ggatgccatc    660

```

-continued

```

tggcgtggct tctacattgc gggatgatccg gccctggcct atggttatgc acaggatcag 720
gagccagacg ctctgtggtcg tatccgtaac ggcgtctctgc tgcgcgtgta cgtaccgcgc 780
agtcctctgc cgggttttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggc 840
gaagtggaac gcctgatccg ccatccgctg ccaactgcgtc tggacgctat cactggtcct 900
gaagaagagg gtggtcgcct ggagactatc ctgggttggc cgctggctga acgcactgta 960
gtaatcccg cgcgatccc aacggatccg cgcaatgttg gtggcgatct ggaccaagc 1020
tctatcccg ataaagaaca ggctatttct gccctgccgg actacgcctc ccagccgggt 1080
aaaccgccgc gtgaggacct gaagtaa 1107

```

<210> SEQ ID NO 190

<211> LENGTH: 368

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PE38 WT as expressed by pIEX02-001 (pET14b-K PE38)

<400> SEQUENCE: 190

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1           5           10           15

Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
20          25          30

His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
35          40          45

Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
50          55          60

Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
65          70          75          80

Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
85          90          95

Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
100         105         110

Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
115         120         125

Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
130         135         140

Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly
145         150         155         160

Asp Ile Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg
165         170         175

Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
180         185         190

Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly
195         200         205

Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe
210         215         220

Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
225         230         235         240

Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val
245         250         255

Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr
260         265         270

```

-continued

Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His
 275 280 285

Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly
 290 295 300

Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val
 305 310 315 320

Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp
 325 330 335

Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu
 340 345 350

Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 355 360 365

<210> SEQ ID NO 191
 <211> LENGTH: 1107
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pIEX02-001 E229D (pET14b-K PE38 E299D)

<400> SEQUENCE: 191

```

atgggcagca gccatcatca tcatcatcac agcagcggcc tgggtgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg    120
gaaaccttca cccgtcaccc tcagccgcgt gggtgggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgga ccaagtagat    240
caggctcatcc gtaacgcgct ggcaagcccg ggttcggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacaggc tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc agggtagcgg taacgacgaa gctggcgctg cgagcgggtc tgcgactcc    420
ggtgacgctc tcttggaacg taactaccg accggtgcag aatttctggg tgatggcggc    480
gatattctct tttctaccg cggcaccag aactggaccg ttgaacgtct gctgcaggcg    540
caccgtcaac tggaagaacg cggttacgtc ttcgtaggtt accacggtac ctctctggaa    600
gctgctcagt ctatcgtgtt cgggtggcgt cgtgctcgta gccaggacct ggatgccatc    660
tggtggtggt tctacattgc ggggtgatcc gccctggcct atggttatgc acaggatcag    720
gagccagacg ctctggtgct tatccgtaac ggcgctctgc tgcgcgtgta cgtaccgcgc    780
agctccctgc cgggtttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggt    840
gaagtggaa cgcctgatcg ccatccgctg cactgcgctc tggacgctat cactggtcct    900
gaagaagagg gtggtcgctt ggacactatc ctgggttggc cgctggctga acgcactgta    960
gtaatcccg cgcgatccc aacggatccg cgcaatgttg gtggcgatct ggacccaagc   1020
tctatcccg ataaagaaca ggctatttct gccctgccgg actacgcctc ccagccgggt   1080
aaaccgccgc gtgaggacct gaagtaa                                1107

```

<210> SEQ ID NO 192
 <211> LENGTH: 368
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pIEX02-001 E229D (pET14b-K PE38 E299D)

<400> SEQUENCE: 192

-continued

Met	Gly	Ser	Ser	His	His	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro
1				5						10				15	
Arg	Gly	Ser	His	Met	Pro	Glu	Gly	Gly	Ser	Leu	Ala	Ala	Leu	Thr	Ala
			20					25					30		
His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe	Thr	Arg	His	Arg	Gln
		35						40				45			
Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg
	50					55					60				
Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp
65					70					75					80
Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu
				85					90					95	
Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr
			100					105					110		
Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn
		115					120					125			
Asp	Glu	Ala	Gly	Ala	Ala	Ser	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu
	130						135				140				
Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly
145					150					155					160
Asp	Ile	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg
				165					170					175	
Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val
			180					185					190		
Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly
		195					200					205			
Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe
	210					215					220				
Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln
225					230					235					240
Glu	Pro	Asp	Ala	Arg	Gly	Arg	Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val
				245					250					255	
Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe	Tyr	Arg	Thr	Gly	Leu	Thr
			260					265					270		
Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His
		275					280					285			
Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly
	290					295					300				
Gly	Arg	Leu	Asp	Thr	Ile	Leu	Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val
305					310					315					320
Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp
				325					330					335	
Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu
			340					345					350		
Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys
		355					360					365			

<210> SEQ ID NO 193

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PE38 mut228 as expressed by pIEX02-228
 (pET14b-K PE38 S253T D209K I153A T164R Q338E)

-continued

<400> SEQUENCE: 193

```

atgggcagca gccatcatca tcacatcac agcagcggcc tggcgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg    120
gaaaccttca cccgtcacgc tcagccgcgt ggttgggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgaa ccaagtagat    240
cagggtcatcc gtaacgcgct ggcaagcccg ggttcgggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacagcg tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc aggggtacggg taacgacgaa gctggcgctg cgagcgggtcc tgccgactcc    420
ggtgacgctc tgctggaacg taactacccg accggtgcag aatttctggg tgatggcggc    480
gatgcctctt tttctaccgg cggcaccagc aactggagag ttgaacgtct gctgcaggcg    540
caccgtcaac tgaagaacg cggttacgtc ttcgtagggt accacgggtac ctctctggaa    600
gctgctcagt ctatcgtgtt cggtggcgta cgtgctcgta gccaggacct gaaggccatc    660
tggcgtggct tctacattgc ggggtgatccg gccctggcct atggttatgc acaggatcag    720
gagccagacg ctcggtggtc tatccgtaac ggcgctctgc tgcgcgtgta cgtaccgcgc    780
agcaccctgc cgggtttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggt    840
gaagtggaac gcctgatccg ccacccgctg ccaactgcgc tggacgctat cactggtcct    900
gaagaagagg gtggtgcgct ggagactatc ctgggttggc cgctggctga acgcactgta    960
gtaatcccg tccgcatccc aacggatccg cgcaatgttg gtggcgatct ggaccaagc    1020
tctatcccg ataaagaaga ggctatttct gccctgccgg actacgcctc ccagccgggt    1080
aaaccgccc gtaggacct gaagtaa                                1107

```

<210> SEQ ID NO 194

<211> LENGTH: 368

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PE38 mut228 as expressed by pLEX02-228
(pET14b-K PE38 S253T D209K I153A T164R Q338E)

<400> SEQUENCE: 194

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1          5          10          15

Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
20        25        30

His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
35        40        45

Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
50        55        60

Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
65        70        75        80

Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
85        90        95

Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
100       105       110

Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
115       120       125

Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu

```

-continued

130	135	140
Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly		
145	150	155 160
Asp Ala Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Arg Val Glu Arg		
	165	170 175
Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val		
	180	185 190
Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly		
	195	200 205
Gly Val Arg Ala Arg Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe		
	210	215 220
Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln		
	225	230 235 240
Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val		
	245	250 255
Tyr Val Pro Arg Ser Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr		
	260	265 270
Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His		
	275	280 285
Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly		
	290	295 300
Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val		
	305	310 315 320
Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp		
	325	330 335
Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu		
	340	345 350
Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys		
	355	360 365

<210> SEQ ID NO 195

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pIEX02-228 E299D (pET14b-K PE38 S253T D209K I153A T164R Q338E E299D)

<400> SEQUENCE: 195

```

atgggcagca gccatcatca tcatcatcac agcagcggcc tggtgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg    120
gaaaccttca ccgcgcaccg tcagccgcgt ggttggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgaa ccaagtagat    240
cagggtcatcc gtaacgcgct ggcaagcccg ggttcgggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacaggc tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc agggtagcgg taacgacgaa gctggcgctg cgagcgggtcc tgcgcactcc    420
ggtgacgctc tgctggaacg taactaccg accggtgcag aatttctggg tgatggcggc    480
gatgcctctt tttctaccg cggcaccag aactggagag ttgaacgtct gctgcaggcg    540
caccgtcaac tggaagaacg cggttacgtc ttcgtaggtt accacggtac ctctctggaa    600
gctgctcagt ctatcgtggt cggtggtgta cgtgctcgta gccaggacct gaaggccatc    660

```

-continued

```

tggcgtggct tctacattgc gggatgatccg gccctggcct atggttatgc acaggatcag 720
gagccagacg ctctgtggtcg tatccgtaac ggcgtctctgc tgcgcgtgta cgtaccgcgc 780
agcacctcgc cgggttttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggc 840
gaagtggaaac gcctgatccg ccatccgctg ccaactgcgtc tggacgctat cactggtcct 900
gaagaagagg gtggtgcgct ggacactatc ctgggttggc cgctggctga acgcaactga 960
gtaatcccg cgcgatccc aacggatccg cgcaatgttg gtggcgatct ggaccaagc 1020
tctatcccg ataaagaaga ggctatttct gccctgccgg actacgcctc ccagccgggt 1080
aaaccgccgc gtgaggacct gaagtaa 1107

```

<210> SEQ ID NO 196

<211> LENGTH: 368

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pIEX02-228 E299D (pET14b-K PE38 S253T D209K I153A T164R Q338E E299D)

<400> SEQUENCE: 196

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1           5           10           15

Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
          20           25           30

His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
          35           40           45

Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
          50           55           60

Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
65           70           75           80

Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
          85           90           95

Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
          100          105          110

Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
          115          120          125

Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
          130          135          140

Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly
145          150          155          160

Asp Ala Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Arg Val Glu Arg
          165          170          175

Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
          180          185          190

Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly
          195          200          205

Gly Val Arg Ala Arg Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe
          210          215          220

Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
225          230          235          240

Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val
          245          250          255

Tyr Val Pro Arg Ser Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr
          260          265          270

```

-continued

Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His
275 280 285

Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly
290 295 300

Gly Arg Leu Asp Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val
305 310 315 320

Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp
325 330 335

Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu
340 345 350

Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
355 360 365

<210> SEQ ID NO 197

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pIEX02-244 (pET14b-K PE38 S253T D209K R204A I153T T164A Q338E)

<400> SEQUENCE: 197

```
atgggcagca gccatcatca tcatcatcac agcagcggcc tggtgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg    120
gaaaccttca cccgtcaccg tcagccgcgt ggttggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgaa ccaagtagat    240
caggctcatcc gtaacgcgct ggcaagcccg ggttcgggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacagcg tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc aggggtacgg taacgacgaa gctggcgctg cgagcggtcc tgccgactcc    420
ggtgacgctc tgctggaacg taactaccgg accggtgcag aatttctggg tgatggcggc    480
gatacctctt tttctaccgg cggcaccag aactgggccc tgaaactct gctgcaggcg    540
caccgtcaac tgaagaacg cgggtacgtc ttcgtaggtt accacggtac ctctcggaa    600
gctgctcagt ctatcgtgtt cgggtggcgt cgtgctgcca gccaggacct gaaggccatc    660
tggcgtggct tctacattgc ggggtatccg gccctggcct atggttatgc acaggatcag    720
gagccagacg ctctggtgct tatccgtaac ggcgctctgc tgcgctgta cgtaccgcgc    780
agcaccctgc cgggtttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggt    840
gaagtggaac gcctgatcgg ccacccgctg ccaactgcgtc tggacgctat cactggtcct    900
gaagaagagg gtggtgcgct ggagactatc ctgggttggc cgctggctga acgcactgta    960
gtaatcccg tccgcatccc aacggatccg cgcaatgttg gtggcgatct ggaccaagc    1020
tctatcccg ataaagaaga ggctatttct gccctgccgg actacgcctc ccagccgggt    1080
aaaccgccc gtgaggacct gaagtaa                                     1107
```

<210> SEQ ID NO 198

<211> LENGTH: 368

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pIEX02-244 (pET14b-K PE38 S253T D209K R204A I153T T164A Q338E)

-continued

<400> SEQUENCE: 198

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
 1          5          10          15
Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
 20          25          30
His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
 35          40          45
Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
 50          55          60
Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
 65          70          75          80
Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
 85          90          95
Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
100          105          110
Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
115          120          125
Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
130          135          140
Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly
145          150          155          160
Asp Thr Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg
165          170          175
Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
180          185          190
Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly
195          200          205
Gly Val Arg Ala Ala Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe
210          215          220
Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
225          230          235          240
Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val
245          250          255
Tyr Val Pro Arg Ser Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr
260          265          270
Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His
275          280          285
Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly
290          295          300
Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val
305          310          315          320
Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp
325          330          335
Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu
340          345          350
Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
355          360          365

```

<210> SEQ ID NO 199

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: pIEX02-244 E299D (pET14b-K PE38 S253T D209K R204A I153T T164A Q338E E299D)

<400> SEQUENCE: 199

```

atgggcagca gccatcatca tcatcatcac agcagcggcc tggcgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg    120
gaaaccttca cccgtcaccg tcagccgcgt ggttgggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgaa ccaagtagat    240
caggctatcc gtaacgcgct ggcaagcccg ggttcgggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacaggc tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc agggtagcgg taacgacgaa gctggcgctg cgagcggtcc tgcgactcc    420
ggtgacgctc tcttggaacg taactaccg accggtgcag aatttctggg tgatggcggc    480
gatacctctt tttctaccg cggcaccag aactgggccc tgaaactct gctgcaggcg    540
caccgtcaac tggaagaacg cgggtacgtc ttcgtaggtt accacggtag ctctctggaa    600
gctgctcagt ctatcgtgtt cgggtggcgt cgtgctgcca gccaggacct gaaggccatc    660
tggcgtggct tctacattgc gggtgatccg gccctggcct atggttatgc acaggatcag    720
gagccagacg ctctgtggtc tatccgtaac ggcgctctgc tgcgctgta cgtaccgcgc    780
agcacctgc cgggtttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggt    840
gaagtggaac gcctgatcgg ccatccgctg ccaactgcgtc tggacgctat cactggctct    900
gaagaagagg gtggtcgctt ggacactatc ctgggttggc cgctggctga acgcactgta    960
gtaatcccg cgcgatccc aacggatccg cgcaatgttg gtggcgatct ggaccaagc    1020
tctatcccg ataaagaaga ggctatttct gccctgccgg actacgcctc ccagccgggt    1080
aaaccgccgc gtgaggacct gaagtaa                                1107

```

<210> SEQ ID NO 200

<211> LENGTH: 368

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pIEX02-244 E299D (pET14b-K PE38 S253T D209K R204A I153T T164A Q338E E299D)

<400> SEQUENCE: 200

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1           5           10           15

Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
20          25          30

His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
35          40          45

Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
50          55          60

Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
65          70          75          80

Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
85          90          95

Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
100         105         110

Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
115         120         125

```

-continued

Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
 130 135 140
 Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly
 145 150 155 160
 Asp Thr Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg
 165 170 175
 Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
 180 185 190
 Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly
 195 200 205
 Gly Val Arg Ala Ala Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe
 210 215 220
 Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
 225 230 235 240
 Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val
 245 250 255
 Tyr Val Pro Arg Ser Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr
 260 265 270
 Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His
 275 280 285
 Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly
 290 295 300
 Gly Arg Leu Asp Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val
 305 310 315 320
 Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp
 325 330 335
 Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu
 340 345 350
 Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 355 360 365

<210> SEQ ID NO 201

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

 <223> OTHER INFORMATION: pIEX02-246 (pET14b-K PE38 S253T D209K R204
 I153A Q338E T164A)

<400> SEQUENCE: 201

```

atgggcagca gccatcatca tcatcatcac agcagcggcc tggcgccgcg cggcagccat    60
atgccagaag gcggtagcct ggccgctctg accgcacatc aggccgtgtca cctgccgctg    120
gaaaccttca cccgtaccgc tcagccgcgt gggtgggaac aactggaaca atgtggctat    180
ccggtacagc gtctggtggc gctgtacctg gctgctcgtc tgtcttgga ccaagtagat    240
cagggtcatcc gtaacgcgct ggcaagcccg ggttccggtg gtgatctggg tgaagctatc    300
cgtgaacaac cggaacaggc tcgtctggcg ctgaccctgg cggcagcgga atctgaacgt    360
tttgtgcgcc agggtagcgg taacgacgaa gctggcgctg cgagcgggtc tgccgactcc    420
gggtgacgctc tgctggaacg taactaccgc accggtgcag aatttctggg tgatggcggc    480
gatgcctctt tttctaccgc cggcaaccag aactgggccc ttgaacgtct gctgcaggcg    540
caccgtcaac tggaagaacg cggttacgtc ttcgtaggtt accacggtac cttoctggaa    600

```

-continued

```

gctgctcagt ctatcgtgtt cgggtggcgta cgtgctgcc a gccaggacct gaaggccatc 660
tggcgtggct tctacattgc gggatgatccg gccctggcct atggttatgc acaggatcag 720
gagccagacg ctctgtggtcg tatccgtaac ggcgctctgc tgcgcgtgta cgtaccgcgc 780
agcacccctgc cgggttttcta tcgtactggc ctgaccctgg ctgcgccgga agcagccggt 840
gaagtggaaac gctgatccg ccatccgctg ccaactgcgtc tggacgctat cactggtcct 900
gaagaagagg gtggtcgcct ggagactatc ctgggttggc cgctggctga acgcaactgta 960
gtaatcccg cgcgatccc aacggatccg cgcaatgttg gtggcgatct ggaccaagc 1020
tctatcccg ataaagaaga ggctatttct gccctgccg actacgcctc ccagccgggt 1080
aaaccgccc gtgaggacct gaagtaa 1107

```

```

<210> SEQ ID NO 202
<211> LENGTH: 368
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pIEX02-246 (pET14b-K PE38 S253T D209K R204
I153A Q338E T164A)

```

```

<400> SEQUENCE: 202

```

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1          5          10          15
Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
20        25        30
His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
35        40        45
Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
50        55        60
Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
65        70        75        80
Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
85        90        95
Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
100       105       110
Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
115       120       125
Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
130       135       140
Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly
145       150       155       160
Asp Ala Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg
165       170       175
Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
180       185       190
Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly
195       200       205
Gly Val Arg Ala Ala Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe
210       215       220
Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
225       230       235       240
Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val
245       250       255

```

```
<210> SEQ ID NO 203
<211> LENGTH: 1107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pIEX02-246 E299D (pET14b-K PE38 S253T D209K
R204 I153A Q338E T164A E299D)
```

atgggcagca gccatcatac tcatacatcac agcagcggcc tgggtgccgc cgcgagccat	60
atgccagaag gcggttagcct ggccgctctg accgcacatc aggcctgtca cctgccgctg	120
gaaaccttca cccgtcaccc tcagccgcgt ggttggaac aactggaaca atgtggctat	180
ccggtacagc gtctggtggc gctgtacctg gctgctctgc tgtcttgga ccaagtagat	240
cagggtcatcc gtaacgcgct ggcaagcccg ggttcgggtg gtgatctggg tgaagctatc	300
cgtgaacaac cggaacaggc tcgtctggcg ctgacctcgg cggcagcgga atctgaacgt	360
tttgtgcgcc agggtagcgg taacagcagaa gctggcgctg cgagcggctc tgccgactcc	420
ggtgacgctc tctggaacg taactaccgc accggtgcag aatttctggg tgatggcggc	480
gatgcctctt tttctaccgc cggcacccag aactgggcgg ttgaacgtct gctgcaggcg	540
caccgtcaac tggaagaacg cggttacgct ttcgtagggt accacggtac cttcctggaa	600
gctgctcagt ctatcgtgtt cgggtggcgt cgtgctgcc gccaggacct gaaggccatc	660
tggcgtggct tctacattgc ggggtgatcc gccctggcct atggttatgc acaggatcag	720
gagccagacg ctcggtggtg tatccgtaac ggcgctctgc tgcgctgtga cgtaccgcgc	780
agcacccctgc cgggtttcta tcgtactggc ctgacctcgg ctgcgccgga agcagccggt	840
gaagtggaac gctgatccg ccataccgct ccaactgcgc tggacgctat cactggtcct	900
gaagaagagg gtggtcgctt ggacactatc ctgggttggc cgctggctga acgcactgta	960
gtaatcccg tccgcatccc aacggatccg cgcaatgttg gtggcgatct ggacccaagc	1020
tctatcccg ataaagaaga ggctatttct gccctgccgg actacgcctc ccagccgggt	1080
aaaccgccgc qtgaqgacgt qaaqtaa	1107

```
<210> SEQ ID NO 204
<211> LENGTH: 368
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pIEX02-246 E299D (pET14b-K PE38 S253T D209K
```

-continued

R204 I153A Q338E T164A E299D)

<400> SEQUENCE: 204

```

Met Gly Ser Ser His His His His His His Ser Ser Gly Leu Val Pro
 1          5          10          15

Arg Gly Ser His Met Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala
 20          25          30

His Gln Ala Cys His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln
 35          40          45

Pro Arg Gly Trp Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg
 50          55          60

Leu Val Ala Leu Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp
 65          70          75          80

Gln Val Ile Arg Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu
 85          90          95

Gly Glu Ala Ile Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr
100          105          110

Leu Ala Ala Ala Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn
115          120          125

Asp Glu Ala Gly Ala Ala Ser Gly Pro Ala Asp Ser Gly Asp Ala Leu
130          135          140

Leu Glu Arg Asn Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly
145          150          155          160

Asp Ala Ser Phe Ser Thr Arg Gly Thr Gln Asn Trp Ala Val Glu Arg
165          170          175

Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val
180          185          190

Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly
195          200          205

Gly Val Arg Ala Ala Ser Gln Asp Leu Lys Ala Ile Trp Arg Gly Phe
210          215          220

Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln
225          230          235          240

Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val
245          250          255

Tyr Val Pro Arg Ser Thr Leu Pro Gly Phe Tyr Arg Thr Gly Leu Thr
260          265          270

Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His
275          280          285

Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly
290          295          300

Gly Arg Leu Asp Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val
305          310          315          320

Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp
325          330          335

Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Glu Ala Ile Ser Ala Leu
340          345          350

Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
355          360          365

```

<210> SEQ ID NO 205

<211> LENGTH: 24

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 001 M13 FOR

<400> SEQUENCE: 205

cgccagggtt ttcccagtca cgac 24

<210> SEQ ID NO 206
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 002 M13 REV

<400> SEQUENCE: 206

agcggataac aatttcacac agga 24

<210> SEQ ID NO 207
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2043 RFB4 VH5' PCR primer sequence

<400> SEQUENCE: 207

gaagtgcagc tgggtggag 18

<210> SEQ ID NO 208
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2044 RFB4 VH3' PCR primer sequence

<400> SEQUENCE: 208

cagagccacc tccgcctgaa ccgcctccac ctgaggagac agtgaccag 49

<210> SEQ ID NO 209
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2045 RFB4 VK 5' PCR Primer Sequence

<400> SEQUENCE: 209

caggcggagg tggctctggc ggtggcggat cggatatcca gatgaccag 50

<210> SEQ ID NO 210
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2046 RFB4 VK 3' PCR Primer sequence

<400> SEQUENCE: 210

tttgatctcc agcttggtg 19

<210> SEQ ID NO 211
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2047 RFB4 Pull through Primer (FOR)

<400> SEQUENCE: 211

-continued

cccagccggc catggcggaa gtgcagctgg tggag 35

<210> SEQ ID NO 212
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2048 RFB4 Pull through Primer (REV)

<400> SEQUENCE: 212

ggtgctcgag tgcggccgcc cgtttgatct ccagcttggt g 41

<210> SEQ ID NO 213
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2097 IEX02 GroEL/ES REV

<400> SEQUENCE: 213

aaccgcccgg ccgttcttct ccgtgttgcc cggaaagcc 39

<210> SEQ ID NO 214
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2098 IEX02 GroEL/ES FOR

<400> SEQUENCE: 214

gggcaaaagc ttgttcttgt ttgagtcac tcatgg 36

<210> SEQ ID NO 215
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2154 IEX02 PE38 FOR, introducing NdeI

<400> SEQUENCE: 215

attgtccata tgccagaagg cggtagcctg gc 32

<210> SEQ ID NO 216
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2161 IEX02 PE38 REV, introducing XhoI

<400> SEQUENCE: 216

atcctcgagt tacttcaggt cctcacgcgg cg 32

<210> SEQ ID NO 217
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2162 IEX02 PE38 NM E229D FOR

<400> SEQUENCE: 217

gggtggtcgc ctggacacta tcctgggttg 30

<210> SEQ ID NO 218

-continued

<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2163 IEX02 PE38 NM E229D REV

<400> SEQUENCE: 218

caaccaggga tagtgtccag gcgaccaccc 30

<210> SEQ ID NO 219
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2164 IEX02 PE38 S253N

<400> SEQUENCE: 219

cagtagcgata gaaaccgggc agattgctgc gcggtacgta 40

<210> SEQ ID NO 220
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2165 IEX02 PE38 S253K

<400> SEQUENCE: 220

cagtagcgata gaaaccgggc agcttgctgc gcggtacgta 40

<210> SEQ ID NO 221
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2166 IEX02 PE38 S253P

<400> SEQUENCE: 221

cagtagcgata gaaaccgggc agagggtctgc gcggtacgta 40

<210> SEQ ID NO 222
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2167 IEX02 PE38 S253T

<400> SEQUENCE: 222

cagtagcgata gaaaccgggc aggggtctgc gcggtacgta 40

<210> SEQ ID NO 223
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2168 IEX02 PE38 Q206R

<400> SEQUENCE: 223

gtacgtgctc gtagcagaga cctggatgcc atc 33

<210> SEQ ID NO 224
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2169 IEX02 PE38 Q206R

-continued

<400> SEQUENCE: 224

gatggcatcc aggtctctgc tacgagcagc tac

33

<210> SEQ ID NO 225

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2170 IEX02 PE38 D209K

<400> SEQUENCE: 225

cgtagccagg acctgaaggc catctggcgt ggc

33

<210> SEQ ID NO 226

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2171 IEX02 PE38 D209K

<400> SEQUENCE: 226

gccacgccag atggccttca ggtcctggct acg

33

<210> SEQ ID NO 227

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2183 IEX02 PE38 I196A FOR, to pair with
OL2161

<400> SEQUENCE: 227

gaagctgctc agtctgccgt gttcggtggc gt

32

<210> SEQ ID NO 228

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2184 IEX02 PE38 I196A REV, to pair with
OL2268

<400> SEQUENCE: 228

acgccaccga acacggcaga ctgagcagct tc

32

<210> SEQ ID NO 229

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2185 IEX02 PE38 I196N FOR, to pair with
OL2161

<400> SEQUENCE: 229

gaagctgctc agtctaactgt gttcggtggc gt

32

<210> SEQ ID NO 230

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2186 IEX02 PE38 I196N REV, to pair with
OL2268

-continued

<400> SEQUENCE: 230

acgccaccga acacgttaga ctgagcagct tc

32

<210> SEQ ID NO 231

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2187 IEX02 to introduce I153A FOR

<400> SEQUENCE: 231

ggtgatggcg gcgatgcctc tttttctacc cgc

33

<210> SEQ ID NO 232

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2188 IEX02 to introduce I153A REV

<400> SEQUENCE: 232

gcgggtagaa aaagaggcat cgccgccatc acc

33

<210> SEQ ID NO 233

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2189 IEX02 to introduce I153T FOR

<400> SEQUENCE: 233

ggtgatggcg gcgatacctc tttttctacc cgc

33

<210> SEQ ID NO 234

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2190 IEX02 to introduce I153T REV

<400> SEQUENCE: 234

gcgggtagaa aaagaggat cgccgccatc acc

33

<210> SEQ ID NO 235

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2191 IEX02 to introduce I153H FOR

<400> SEQUENCE: 235

ggtgatggcg gcgatcactc tttttctacc cgc

33

<210> SEQ ID NO 236

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2192 IEX02 to introduce I153H REV

<400> SEQUENCE: 236

gcgggtagaa aaagagtgat cgccgccatc acc

33

-continued

<210> SEQ ID NO 237
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2193 IEX02 to introduce T164R FOR

<400> SEQUENCE: 237

gcacccagaa ctggagagtt gaacgtctgc tg 32

<210> SEQ ID NO 238
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2194 IEX02 to introduce T164R REV

<400> SEQUENCE: 238

cagcagacgt tcaactctcc agttctgggt gc 32

<210> SEQ ID NO 239
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2216 IEX02 Linker to optimize Kozak in
pET14b, to anneal with OL2217

<400> SEQUENCE: 239

catggtggct ctccttctta aagttaaca aaattat 39

<210> SEQ ID NO 240
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2217 IEX02 Linker to optimize Kozak in
pET14b, to anneal with OL2216

<400> SEQUENCE: 240

ctagaaataa ttttgtttaa cttaagaag gagagccac 39

<210> SEQ ID NO 241
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2268 IEX02 outside FOR spans over XbaI
site (pET14b) - to be paired with OL2161

<400> SEQUENCE: 241

atctccctct agaaataatt ttgtttaact ttaagaag 38

<210> SEQ ID NO 242
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2279 IEX02 FOR oligo to remove TM to be
paired with OL2161

<400> SEQUENCE: 242

gaagctgctc agtctatcgt gtccggtggc gt 32

<210> SEQ ID NO 243

-continued

<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2280 IEX02 REV oligo to remove TM to be
paired with OL2268

<400> SEQUENCE: 243

acgccaccga acacgataga ctgagcagct tc 32

<210> SEQ ID NO 244
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2281 IEX02 A201 REV, ONLY for templates
having Q206

<400> SEQUENCE: 244

ctctgctacg agcacgggcg ccaccgaaca cg 32

<210> SEQ ID NO 245
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2282 IEX02 A201 FOR, ONLY for templates
having Q206

<400> SEQUENCE: 245

cgtgttcggt ggcgcccgtg ctctagcag ag 32

<210> SEQ ID NO 246
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2283 IEX02 A204 REV, ONLY for templates
having Q206

<400> SEQUENCE: 246

catccaggtc tctgctggca gcacgtacgc cac 33

<210> SEQ ID NO 247
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2284 IEX02 A204 FOR, ONLY for templates
having Q206

<400> SEQUENCE: 247

gtggcgtagc tgctgccagc agagacctgg atg 33

<210> SEQ ID NO 248
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2285 IEX02 Q204 REV, ONLY for templates
having Q206

<400> SEQUENCE: 248

catccaggtc tctgctctga gcacgtacgc cac 33

-continued

<210> SEQ ID NO 249
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2286 IEX02 Q204 FOR, ONLY for templates
having Q206

<400> SEQUENCE: 249

gtggcgtagc tgctcagagc agagacctgg atg 33

<210> SEQ ID NO 250
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2287 IEX02 A158 REV

<400> SEQUENCE: 250

ccagttctgg gtgccggcgg tagaaaaaga g 31

<210> SEQ ID NO 251
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2288 IEX02 A158 FOR

<400> SEQUENCE: 251

ctctttttct accgccggca cccagaactg g 31

<210> SEQ ID NO 252
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2289 IEX02 Q158 REV

<400> SEQUENCE: 252

ccagttctgg gtgccctggg tagaaaaaga gatatc 36

<210> SEQ ID NO 253
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2290 IEX02 Q158 FOR

<400> SEQUENCE: 253

gatattctct tttctaccca gggcaccag aactgg 36

<210> SEQ ID NO 254
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2291 IEX02 S159 REV

<400> SEQUENCE: 254

gtccagttct gggtaggagcg ggtagaaaaa gagatatc 38

<210> SEQ ID NO 255
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: OL 2292 IEX02 S159 FOR

<400> SEQUENCE: 255

gatatactctt tttctaccgg ctccaccag aactggac 38

<210> SEQ ID NO 256
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2293 IEX02 T159 REV

<400> SEQUENCE: 256

accaccaga actggaccgt tgaac 25

<210> SEQ ID NO 257
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2294 IEX02 T159 FOR

<400> SEQUENCE: 257

ccagttctgg gtggtgcggg tagaaaaaga g 31

<210> SEQ ID NO 258
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2295 IEX02 generic REV oligo for mutations
at 333

<400> SEQUENCE: 258

gagcttgggt ccagatcgcc acc 23

<210> SEQ ID NO 259
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2296 IEX02 A333 FOR

<400> SEQUENCE: 259

ctggacccaa gctctgcccc ggataaagaa c 31

<210> SEQ ID NO 260
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2297 IEX02 N333 FOR

<400> SEQUENCE: 260

ctggacccaa gctctaaccg ggataaag 28

<210> SEQ ID NO 261
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2298 IEX02 T333 FOR

<400> SEQUENCE: 261

-continued

ctggacccaa gctctacccc ggataaag 28

<210> SEQ ID NO 262
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2299 IEX02 Q333 FOR

<400> SEQUENCE: 262

ctggacccaa gctctcagcc ggataaagaa c 31

<210> SEQ ID NO 263
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2300 IEX02 H333 FOR

<400> SEQUENCE: 263

ctggacccaa gctctcacccc ggataaag 28

<210> SEQ ID NO 264
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2301 IEX02 N338 FOR

<400> SEQUENCE: 264

ctggacccaa gctctatccc ggataaagaa aacgctattt ctgccctg 48

<210> SEQ ID NO 265
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2302 IEX02 E338 FOR

<400> SEQUENCE: 265

ctggacccaa gctctatccc ggataaagaa gaggctattt ctgccc 46

<210> SEQ ID NO 266
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2303 IEX02 V201A REV

<400> SEQUENCE: 266

ctgggtacga gcacggggcgc caccgaac 28

<210> SEQ ID NO 267
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2304 IEX02 V201A FOR

<400> SEQUENCE: 267

gttcggtggc gcccggtgctc gtagccag 28

<210> SEQ ID NO 268

-continued

<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2305 IEX02 R204A REV, ONLY for templates
having D209K

<400> SEQUENCE: 268

cttcaggtcc tggctggcag cacgtacgcc 30

<210> SEQ ID NO 269
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2306 IEX02 R204A FOR, ONLY for templates
having D209K

<400> SEQUENCE: 269

ggcgtagctg ctgccagcca ggacctgaag 30

<210> SEQ ID NO 270
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2307 IEX02 R204Q REV, ONLY for templates
having D209K

<400> SEQUENCE: 270

cttcaggtcc tggctctgag cacgtacgc 29

<210> SEQ ID NO 271
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2308 IEX02 R204Q FOR, ONLY for templates
having D209K

<400> SEQUENCE: 271

gcgtacgtgc tcagagccag gacctgaag 29

<210> SEQ ID NO 272
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2309 IEX02 Q161N REV

<400> SEQUENCE: 272

gttcaacggt ccagttgttg gtgccgctgg tag 33

<210> SEQ ID NO 273
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2310 IEX02 Q161N FOR

<400> SEQUENCE: 273

ctacccgcgg caccaacaac tggaccgttg aac 33

<210> SEQ ID NO 274
<211> LENGTH: 33

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2311 IEX02 Q161T REV

<400> SEQUENCE: 274

gttcaacggt ccagttggtg gtgccgagg tag 33

<210> SEQ ID NO 275
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2312 IEX02 Q161T FOR

<400> SEQUENCE: 275

ctacccgagg caccaccaac tggacgttg aac 33

<210> SEQ ID NO 276
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2313 IEX02 T164A REV

<400> SEQUENCE: 276

cagcagacgt tcaacggccc agttctgggt g 31

<210> SEQ ID NO 277
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2314 IEX02 T164A FOR

<400> SEQUENCE: 277

caccagaac tgggaggtg aacgtctgt g 31

<210> SEQ ID NO 278
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2315 IEX02 N162A REV

<400> SEQUENCE: 278

gacgttcaac ggtccaggcc tgggtgccgc ggg 33

<210> SEQ ID NO 279
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2316 IEX02 N162A FOR

<400> SEQUENCE: 279

ccgcggcac ccaggcctgg accgttgaac gtc 33

<210> SEQ ID NO 280
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OL 2318 IEX02 FOR RFB4 (NcoI)

-continued

<400> SEQUENCE: 280

attgccacca tggcggaggt gc 22

<210> SEQ ID NO 281

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2320 IEX02 REV to create RFB4 for RFB4-PE38-8xHis to pair with OL2318

<400> SEQUENCE: 281

caccaggccg ctgcttttga tctccagctt g 31

<210> SEQ ID NO 282

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2321 IEX02 FOR to create RFB4-PE38-8xHis to pair with OL2322

<400> SEQUENCE: 282

caagctggag atcaaaagca gcggcctggt g 31

<210> SEQ ID NO 283

<211> LENGTH: 66

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2322 IEX02 REV introducing 8xHis C-terminus of PE, introducing XhoI

<400> SEQUENCE: 283

cgattctcga gttacttcag gtctctgtgg tgggtggtgat gatgatgatg acgcggcggt 60

ttaccc 66

<210> SEQ ID NO 284

<211> LENGTH: 44

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2323 IEX02 FOR to create RFB4-6xHis PE38 fusions (pIEX02-302 and pIEX02-304) to pair with OL2161

<400> SEQUENCE: 284

caagctggag atcaaagctc atgggggcag ccatcatcat catc 44

<210> SEQ ID NO 285

<211> LENGTH: 44

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: OL 2324 IEX02 REV to create RFB4-6xHis PE38 fusions (pIEX02-302 and pIEX02-304) to pair with OL2318

<400> SEQUENCE: 285

gatgatgatg atggctgccc ccatgagctt tgatctccag cttg 44

<210> SEQ ID NO 286

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

```

<223> OTHER INFORMATION: Modified Kozak sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: r at position 7 is a purine (adenine or
                        guanine)

<400> SEQUENCE: 286

gccgcgcrccca ugg

```

13

1. A polypeptide having at least one *Pseudomonas* exotoxin A (PE-A) biological activity, wherein said polypeptide comprises one or more amino acid substitutions compared to a wild-type PE-A polypeptide, wherein said one or more amino acid substitutions is a substitution of a different amino acid at one or more positions corresponding to amino acid residues in the polypeptide of SEQ ID NO:1, wherein said substitution positions are selected from the group consisting of:

- a) isoleucine (I) at position 141;
- b) arginine (R) at position 146;
- c) glycine (G) at position 147;
- d) glutamine (Q) at position 149;
- e) asparagine (N) at position 150;
- f) threonine (T) at position 152;
- g) valine (V) at position 189;
- h) arginine (R) at position 192;
- i) glutamine (Q) at position 194;
- j) aspartic acid (D) at position 197;
- k) serine (S) at position 241;
- l) isoleucine (I) at position 321; and
- m) glutamine (Q) at position 326.

2. The polypeptide of claim 1, wherein said one or more amino acid substitutions is a conservative amino acid substitution.

3. The polypeptide of claim 1, wherein said one or more amino acid substitutions is selected from the group consisting of:

- a) isoleucine (I) at position 141 is substituted with alanine (A), threonine (T), or histidine (H);
- b) arginine (R) at position 146 is substituted with glutamine (Q) or alanine (A);
- c) glycine (G) at position 147 is substituted with serine (S);
- d) glutamine (Q) at position 149 is substituted with threonine (T);
- e) asparagine (N) at position 150 is substituted with alanine (A);
- f) threonine (T) at position 152 is substituted with alanine (A) or arginine (R);
- g) valine (V) at 189 is substituted with alanine (A);
- h) arginine (R) at position 192 is substituted with alanine (A) or glutamine (Q);
- i) glutamine (Q) at position 194 is substituted with arginine (R);
- j) aspartic acid (D) at position 197 is substituted with lysine (K);
- k) serine (S) at position 241 is substituted with threonine (T), asparagine (N), lysine (K), or proline (P);

l) isoleucine (I) at position 321 is substituted with alanine (A), asparagine (N), histidine (H), threonine (T), or glutamine (Q); and

m) glutamine (Q) at position 326 is substituted with glutamic acid (E).

4. The polypeptide of claim 1, wherein said polypeptide comprises:

- a) a substitution for isoleucine (I) at position 141, a substitution for threonine (T) at position 152, a substitution for arginine (R) at position 192, a substitution for aspartic acid (D) at position 197, a substitution for serine (S) at position 241, and a substitution for glutamine (Q) at position 326;
- b) a substitution of threonine (T) or alanine (A) for isoleucine (I) at position 141, a substitution alanine (A) or arginine (R) for threonine (T) at position 152, a substitution of alanine (A) for arginine (R) at position 192, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326;
- c) a substitution of threonine (T) for isoleucine (I) at position 141, a substitution alanine (A) for threonine (T) at position 152, a substitution of alanine (A) for arginine (R) at position 192, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (B) for glutamine (Q) at position 326;
- d) a substitution of alanine (A) for isoleucine (I) at position 141, a substitution alanine (A) for threonine (T) at position 152, a substitution of alanine (A) for arginine (R) at position 192, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326;
- e) a substitution of alanine (A) or threonine (T) for isoleucine (I) at position 141, a substitution of arginine (R) or alanine (A) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326;
- f) a substitution of alanine (A) for isoleucine (I) at position 141, a substitution of arginine (R) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326;
- g) a substitution of alanine (A) for isoleucine (I) at position 141, a substitution of alanine (A) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for

serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326; or

- h) a substitution of threonine (T) for isoleucine (I) at position 141, a substitution of alanine (A) for threonine (T) at position 152, a substitution of lysine (K) for aspartic acid (D) at position 197, a substitution of threonine (T) for serine (S) at position 241, and a substitution of glutamic acid (E) for glutamine (Q) at position 326.

5-7. (canceled)

8. The polypeptide of claim 1, wherein said polypeptide comprises:

- a) a substitution for isoleucine (I) at position 141, a substitution for threonine (T) at position 152, a substitution for aspartic acid (D) at position 197, a substitution for serine (S) at position 241, and a substitution for glutamine (Q) at position 326; or
- b) a substitution for isoleucine (I) at position 141, a substitution for threonine (T) at position 152, a substitution for arginine (R) at position 192, a substitution for aspartic acid (D) at position 197, and a substitution for serine (S) at position 241.

9-13. (canceled)

14. The polypeptide of claim 1, wherein the at least one *Pseudomonas* exotoxin A (PE-A) biological activity comprises the ability to inhibit in vitro transcription/translation compared to a corresponding wild-type or non-substituted PE-A polypeptide, wherein said ability to inhibit in vitro transcription/translation is in an amount selected from the group consisting of:

- a) at least 5% inhibition;
- b) at least 10% inhibition;
- c) at least 15% inhibition;
- d) at least 20% inhibition;
- e) at least 25% inhibition;
- f) at least 30% inhibition;
- g) at least 40% inhibition;
- h) at least 50% inhibition;
- i) at least 60% inhibition;
- j) at least 70% inhibition;
- k) at least 80% inhibition;
- l) at least 90% inhibition;
- m) at least 100% inhibition;
- n) about 100% inhibition; and
- o) 100% inhibition.

15. The polypeptide of claim 1, comprising a number of amino acid substitutions selected from the group consisting of:

- a) 1 amino acid substitution;
- b) 2 amino acid substitutions;
- c) 3 amino acid substitutions;
- d) 4 amino acid substitutions;
- e) 5 amino acid substitutions; and
- f) 6 amino acid substitutions.

16-17. (canceled)

18. The polypeptide of claim 1, wherein the last five or six amino acids in said polypeptide comprise one or more amino acid sequences selected from the group consisting of:

- a) Arg-Glu-Asp-Leu-Lys;
- b) Arg-Glu-Asp-Leu;
- c) Lys-Asp-Glu-Leu;
- d) Glu-Asp-Leu-Lys; and
- e) a dimer, trimer, pentamer, hexamer, septamer, or octamer of a), b), or c), or any combination thereof.

19. The polypeptide of claim 1, wherein said polypeptide has one or more biological activities selected from the group consisting of:

- a) eukaryotic cell killing activity (cell cytotoxicity);
- b) inhibits translation elongation factor EF-2 biological activity;
- c) induces or catalyzes ADP-ribosylation of EF-2; and
- d) inhibits protein synthesis.

20. The polypeptide of claim 1, wherein said one or more amino acid substitutions prevent or reduce host immunogenic responses compared to the same polypeptide without the corresponding said one or more amino acid substitutions.

21. A polypeptide comprising a biologically active fragment of the polypeptide of claim 1.

22. (canceled)

23. The polypeptide of claim 1, wherein said one or more amino acid substitutions prevent or reduce host immunogenic responses compared to a polypeptide comprising an amino acid sequence selected from the group consisting of:

- a) SEQ ID NO:1;
- b) SEQ ID NO:4;
- c) SEQ ID NO:133; and
- d) SEQ ID NO: 134.

24. The polypeptide of claim 1, wherein said polypeptide is a fusion protein comprising a different polypeptide.

25-34. (canceled)

35. A polynucleotide encoding the polypeptide of claim 1.

36. An expression vector comprising the polynucleotide of claim 35.

37. A host cell comprising the expression vector of claim 36.

38. A method of producing the polypeptide of claim 1, wherein said method comprises:

- a) obtaining a host cell comprising a polynucleotide encoding said polypeptide; and
- b) exposing said host cell to conditions wherein said polypeptide is produced.

39. A method of producing the polypeptide of claim 1, wherein said method comprises use of an expression system comprising:

- A) a first polynucleotide encoding a first hybrid polypeptide comprising:
 - i) a first ligand binding domain; and
 - ii) a DNA-binding domain;
- B) a second polynucleotide encoding a second hybrid polypeptide comprising:
 - i) a second ligand binding domain; and
 - ii) a transactivation domain; and

C) a third polynucleotide encoding the polypeptide of claim 1, wherein said third polynucleotide is operably associated with a response element capable of being bound by the DNA-binding domain of said first hybrid polypeptide;

wherein the first ligand binding domain and the second ligand binding domain are capable of ligand-induced dimerization,

wherein expression of the polypeptide of claim 1 is modulated by a ligand which induces dimerization of said first and said second ligand binding domains, and

wherein the polypeptide of claim 1 is produced by allowing said ligand to contact said first and said second ligand binding domains.

40-42. (canceled)

43. A method of treating a disease or disorder comprising administering to a subject in need thereof the polypeptide of claim **1**, the host cell of

44. A method of treating a disease or disorder comprising delivering to a subject in need thereof a polypeptide produced by the method of claim **39**, wherein said method comprises administration of the ligand to said subject.

45-47. (canceled)

48. A pharmaceutical composition comprising the polypeptide of claim **1**.

49-51. (canceled)

52. An *Pseudomonas* exotoxin A (PE-A) polypeptide, wherein said polypeptide comprises a mutation at a position corresponding to amino acid position 184 in SEQ ID NO:1 (or position 196 in SEQ ID NO:2) wherein an isoleucine at position 184 (or position 196 in SEQ id NO:2) is substituted with a different amino acid.

53. The polypeptide of claim **52**, wherein said polypeptide does not have PE-A biological activity.

54. A method for assaying the immunogenicity of a mutated form of *Pseudomonas* exotoxin A (PE-A), wherein said method comprises:

a) contacting immune cells with a mutated form of PE-A; and

b) assaying immune cell stimulation,

wherein said mutated form of PE-A comprises a mutation at a position corresponding to amino acid position 184 in SEQ ID NO:1 (or position 196 in SEQ ID NO:2) wherein an isoleucine at position 184 (or position 196 in SEQ ID NO:2) is substituted with a different amino acid, and wherein said mutated form of PE-A also comprises one or more additional amino acid substitutions compared to a wild-type form of PE-A.

55-56. (canceled)

* * * * *