



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2193705 C 2003/12/02

(11)(21) **2 193 705**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1996/05/20

(87) Date publication PCT/PCT Publication Date: 1996/11/21

(45) Date de délivrance/Issue Date: 2003/12/02

(85) Entrée phase nationale/National Entry: 1996/12/20

(86) N° demande PCT/PCT Application No.: US 1996/007297

(87) N° publication PCT/PCT Publication No.: 1996/036882

(30) Priorité/Priority: 1995/05/19 (08/444,895) US

(51) Cl.Int.⁶/Int.Cl.⁶ G01N 33/68, G01N 33/52, G01N 1/30

(72) Inventeurs/Inventors:

HAUGLAND, RICHARD P., US;
SINGER, VICTORIA L., US;
JONES, LAURIE JEANNE, US;
STEINBERG, THOMAS H., US

(73) Propriétaire/Owner:

MOLECULAR PROBES, INC., US

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : COLORATION DE PROTEINES PAR COLORANT A BASE DE MEROCYANINE

(54) Title: MEROCYANINE DYE PROTEIN STAINS

(57) Abrégé/Abstract:

This invention describes the use of merocyanine dyes, including styryl dyes, for detecting and quantifying poly(amino acids) by absorbance or fluorescence. Poly(amino acids) are detected in solution, in electrophoretic gels, or on solid supports.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :

G01N 33/68, 1/30

A1

(11) International Publication Number:

WO 96/36882

(43) International Publication Date: 21 November 1996 (21.11.96)

(21) International Application Number: PCT/US96/07297

(22) International Filing Date: 20 May 1996 (20.05.96)

(30) Priority Data:

08/444,895

19 May 1995 (19.05.95)

US

(71) Applicant: MOLECULAR PROBES, INC. [US/US]; 4849
Pitchford Avenue, Eugene, OR 97402 (US).

(72) Inventors: HAUGLAND, Richard, P.; 2233 Lasater, Eugene, OR 97402 (US). SINGER, Victoria, L.; 910 East 23rd Avenue, Eugene, OR 97405 (US). JONES, Laurie, Jeanne; 26629 Bennett Boulevard, Monroe, OR 97456 (US). STEINBERG, Thomas, H.; 1830 Washington Street, Eugene, OR 97401 (US).

(74) Agents: HELFENSTEIN, Allegra, J. et al.; Molecular Probes, Inc., P.O. Box 22010, Eugene, OR 97402 (US).


(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

2193705

(54) Title: MEROCYANINE DYE PROTEIN STAINS



(57) Abstract

This invention describes the use of merocyanine dyes, including styryl dyes, for detecting and quantifying poly(amino acids) by absorbance or fluorescence. Poly(amino acids) are detected in solution, in electrophoretic gels, or on solid supports.

MEROCYANINE DYE PROTEIN STAINS

FIELD OF THE INVENTION

5 The invention relates to staining of poly(amino acids), including peptides, polypeptides and proteins in solution, in gels and on solid supports, using merocyanine dyes.

BACKGROUND

10 Detection and analysis of poly(amino acids) is of great importance in a multitude of diverse activities, ranging from basic research to enzyme production, forensics analysis and diagnostics.

 Several methods have been utilized to detect proteins or other poly(amino acids) in
15 electrophoresis gels, including colorimetric methods like COOMASSIE Brilliant Blue (CBB) staining and silver staining, as well as fluorescent staining, e.g. Nile red. The invention possesses many advantages over known methods for staining poly(amino acids) on gels: Staining is rapid, simple, and relatively insensitive to poly(amino acid) composition. Visualization is possible without destaining, and the stained bands remain detectable for days. The dyes used in the current
20 method are readily soluble and stable in aqueous staining solutions. In addition, the dyes exhibit large Stokes shifts. Finally, the present invention allows the detection of as little as 1 ng of poly(amino acid) per band, which is comparable to silver staining, with less hazard and expense, and is more than an order of magnitude better than CBB or Nile red staining.

25 The invention can also be used to detect poly(amino acids) on filter membranes or other solid supports, or in solution. The use of this invention for staining poly(amino acids) in solution can also be used to quantitate poly(amino acids) with greater sensitivity than other known methods, including absorbance-based Lowry, Bradford, and bicinchoninic acid (BCA) methods, and fluorescence-based methods (e.g. Nile red), and with a greater dynamic range (Table 3).

30

 The preparation and characterization of merocyanine dyes useful for the present invention is well documented. A number of styryl merocyanine dyes (referred to as RH dyes) have been previously prepared for measuring electric potentials in cell membranes. A variety of other merocyanine dyes have been described for use in the photographic industry, without reference to

their fluorescence properties. The present invention, which describes the staining and detection of poly(amino acids) outside of the cellular milieu, is neither anticipated nor obvious from previous references.

5 DESCRIPTION OF DRAWINGS

Figure 1: Normalized excitation and emission spectra in the presence of poly(amino acid).

Fluorescence spectra were obtained of 1 μ M of Dye 801, 150 μ g/mL bovine serum albumin (BSA) and 0.05% SDS in 10 mM Tris-HCl, pH 7.5.

10

Figure 2: Fluorescence enhancement upon addition of poly(amino acid). Spectra were obtained of 1 μ M Dye 801 and 0.05% SDS in 10 mM Tris-HCl, pH 7.5, in the absence and presence of 500 μ g/mL BSA.

15 Figure 3: Correlation between poly(amino acid) concentration and fluorescence emission for Dye 307 as in Table 3, note 4, using 485 nm excitation and 590 nm emission.

Figure 4: Correlation between band fluorescence and total poly(amino acids) present in a gel band for Dye 801, as in Example 9.

20

SUMMARY OF THE INVENTION AND DESCRIPTION OF PREFERRED EMBODIMENTS

Poly(amino acids) are stained according to the present invention by incubating the poly(amino acids) with one or more merocyanine dyes. As used herein, a poly(amino acid) is any
25 homopolymer or heteropolymer of amino acids, including peptides, polypeptides, and proteins. The merocyanine dyes associate with the amino acid polymers either directly, or in the presence of a detergent, to yield both a strong colorimetric absorption and a strong fluorescence emission. Any poly(amino acid) thereby labeled can be detected with high sensitivity either in solution, or on a solid or semisolid support.

30

This invention provides a method of detecting a poly(amino acid), comprising the steps of: a) combining a sample mixture that is thought to contain a poly(amino acid) with a staining mixture that contains one or more merocyanine dyes to form a combined mixture; b) incubating the combined mixture for a time sufficient for the dye in the staining mixture to associate with the poly(amino acid) in the sample mixture to form a dye-poly(amino acid) complex that gives a detectable optical response upon illumination; c) illuminating said dye-poly(amino acid) complex; and, d) observing said optical response, providing that the staining mixture does not comprise a single merocyanine dye that is a 4'-hydroxy-3',5'-dihalostryryl merocyanine dye having a pKa value that changes in the presence of a poly(amino acid). The merocyanine dye may be a merocyanine dye having the formula Q-B-M as described below.

This invention also provides a method of detecting a poly(amino acid), comprising the steps of: a) combining a sample mixture that is thought to contain a poly(amino acid) with a staining mixture that contains one or more styryl dyes having the formula Q-B-M as described below, to form a combined mixture; b) incubating the combined mixture for a time sufficient for the dye in the staining mixture to associate with the poly(amino acid) in the sample mixture to form a dye-poly(amino acid) complex that gives a detectable optical response upon illumination; c) illuminating said dye-poly(amino acid) complex; and, d) observing said detectable optical response.

This invention also provides a combination consisting essentially of a detergent, a poly(amino acid), and one or more styryl dyes having the formula Q-B-M as described below, in a cell-free aqueous solution where said detergent is present at a concentration less than the critical micelle concentration for that detergent.

This invention also provides a kit for labelling the poly(amino acids) in a sample, comprising: a) one or more merocyanine dyes; and b) instructions for combining said merocyanine dye or dyes with a sample containing or thought to contain poly(amino acids), such that the dye or dyes form a detectable label in conjunction with poly(amino acids) in the sample, providing that the kit does not comprise a single merocyanine dye that is a 4'-hydroxy-3',5'-dihalostryryl merocyanine dye having a pKa value that changes in the presence of a poly(amino acid). The merocyanine dye may be a merocyanine dye having the formula Q-B-M as described below.


Selection of Dyes

Merocyanine dyes comprise a quarternary nitrogen heterocycle linked to an electron pair-donating moiety by an alkylene or polyalkylene bridge. A wide variety of electron pair-donating

groups are known that stabilize the formally positive charge of the quaternary nitrogen heterocycle by resonance. Suitable electron pair-donating groups include dialkylaminophenyl, dialkylaminonaphthyl, electron-rich heterocycles and acyclic moieties containing electron pair-donating groups.

5


Preferred merocyanine dyes have the general formula




- 10 wherein Q is a quaternized nitrogen heterocycle where the quaternizing group is a TAIL group, B is a covalent bridge that is an ethenyl or polyethenyl moiety, and M is an aromatic substituent or activated methylene substituent.

- The covalent bridge, B, has the formula $-(CR^7=CR^8)_n-$, where R^7 and R^8 are independently
 15 H, C_1-C_6 alkyl or phenyl. Alternatively, R^7 and R^8 taken in combination complete a 5- or 6-membered saturated ring. The subscript n has a value of 1-3. Higher values of n shift the excitation and emission to longer wavelengths. Typically, R^7 and R^8 are both H. Preferably $n = 1$ or 2.

- 20 The quaternized nitrogen heterocycle Q is typically a substituted or unsubstituted pyridinium, quinolinium, benzoquinolinium or benzazolum moiety having the formula




25 or



or the formula

5




where the ring substituents R^1 , R^2 , R^3 and R^4 are optionally and independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls. The ring substituents R^5 and R^6 are optionally and independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls, or phenyl. Alternatively, R^5 and R^6 , when taken in combination, form a fused 6-membered aromatic ring (yielding a quinolinium moiety) that is optionally and independently substituted one or more times by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls. The quinolinium ring is optionally substituted by an additional fused 6-membered aromatic ring (yielding a naphtho-substituted pyridinium, or a benzoquinoline), that is also optionally and independently substituted one or more times by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls. Typically, R^5 and R^6 are H, or form a substituted or unsubstituted benzo moiety. Preferably R^5 and R^6 , taken in combination, form a fused 6-membered substituted or unsubstituted benzo moiety yielding a quinolinium ring system.

In the benzazole ring Q1, the ring fragment X is optionally -S-, -O-, -NR⁷-, or -CR⁷R⁸-, wherein R⁷ and R⁸ are as defined previously. Typically R⁷ and R⁸ methyls. Preferably, X is O or S, more preferably X is S.


The quaternizing moiety, TAIL, is attached to the nitrogen atom of Q through a carbon atom and contains a total of 1-22 non-hydrogen atoms that are C, O, N or S, such that within TAIL each heteroatom is separated from any adjacent heteroatoms by at least two carbon atoms. TAIL is composed of bonds that are selected from the group consisting of carbon-carbon bonds (C-C), ether bonds (C-O-C), thioether bonds (C-S-C) or amine bonds (C-NR⁹-C). Any carbon atom in TAIL is optionally further substituted by hydroxy, carboxy, sulfo, amino, or ammonium. Any amine, amino or ammonium in TAIL is optionally substituted by an R⁹ that is a C₂-C₆ alkyl that is optionally further substituted by hydroxy, carboxy, sulfo, amino, amino substituted by 1-2 C₁-C₆ alkyls or ammonium substituted by 1-3 C₁-C₆ alkyls. Alternatively, the nitrogen atoms of TAIL form either one or two saturated 5- or 6-membered rings in combination with other C or N atoms in TAIL, such that the resulting rings are pyrrolidines, piperidines, piperazines or morpholines.

In one embodiment, the TAIL moiety includes at least one dialkylamino or a trialkylammonium substituent, where the alkyl substituents are methyl or ethyl. Preferably, TAIL is -CH₃ or CH₂CH₃, or is a C₃-C₂₂ alkyl that is optionally substituted one or more times by hydroxy, carboxy, sulfo, amino, amino substituted by 1-2 C₁-C₆ alkyls, or ammonium substituted by 1-3 C₁-C₆ alkyls. By "sulfo" is meant sulfonic acid (-SO₃H) or the common alkali metal salts of sulfonic acid. More preferably, TAIL is a C₃-C₁₂ alkyl that is linear and saturated, and substituted at its free terminus by hydroxy, carboxy, sulfo, amino, amino substituted by 1-2 C₁-C₆ alkyls, or ammonium substituted by 1-3 C₁-C₆ alkyls. Yet more preferably, TAIL is a C₃-C₄ alkyl that is substituted once by sulfo or carboxy.

The moiety M has the formula



or













where R^{11} and R^{12} are independently H, F, Cl, or $-CH_3$; typically H. R^{13} and R^{14} are independently C_1 - C_{18} alkyls that are linear, branched, saturated or unsaturated, and are optionally substituted one or more times by F, hydroxy or C_1 - C_6 alkoxy. Alternatively, R^{13} and R^{14} , when taken in combination, form a 5- or 6-membered saturated ring that optionally contains an oxygen heteroatom. In another embodiment of the dyes, R^{11} taken in combination with R^{13} and R^{10} taken in combination with R^{12} are independently $-(CH_2)_2-$ or $-(CH_2)_3-$, forming 5- or 6-membered rings. Preferably R^{13} and R^{14} are each linear alkyls, which may be the same or different, each having 4-8 carbon atoms, more preferably each having 5-7 carbon atoms.

Alternatively, M has a formula as depicted in Table 1, where R is H, phenyl, sulfophenyl, C_1 - C_6 alkyl or C_1 - C_6 alkyl substituted by carboxy. Such dyes have been described, for example, by Brooker et al., J. AM. CHEM. SOC. 73, 5326 (1951).

Table 1: Selected electron pair-donating moieties

5-pyrazolone	2-thiohydantoin
2,4-thiazolidinedione	2-imino-4-oxazolone

 2-imino-4-thiazolidone	 5(4H)-isoxazolone
 2-thio-2,4-oxazolidinedione	 barbituric acid
 5-thiopyrazolone	 2-thiobarbituric acid
 5(4H)-oxazolone	 3-thio-2,4-thiazolidinedione
 cyanoacetate	 malonate

For all dyes, any net positive or negative charges possessed by the dye are balanced by counterion(s), Ψ . Any counterion currently used in conjunction with biomolecules is suitable.

15 Preferred counterions include chloride, iodide, perchlorate, various sulfonates, alkali metal ions, alkaline earth metal ions, transition metal ions, ammonium or substituted ammonium ions.

Table 2: Selected dyes useful for the present invention

Dye no.	Dye Structure
101	
102	
103	
104	
201	
301	
302	
303	
304	
305	
306	
307	

308	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_8\text{H}_{17})_2$
309	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-(\text{CH}=\text{CH})_2-\text{C}_6\text{H}_4-\text{N}(\text{C}_4\text{H}_9)_2$
310	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-(\text{CH}=\text{CH})_2-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
311	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-(\text{CH}=\text{CH})_3-\text{C}_6\text{H}_4-\text{N}(\text{C}_4\text{H}_9)_2$
312	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{C}_6\text{H}_2(\text{Cl})_2-\text{N}(\text{C}_6\text{H}_{13})_2$
313	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{CH}_2(\text{CF}_2)_3\text{CF}_3)_2$
401	$\text{O}_3\text{S}-(\text{CH}_2)_3-\text{N}^+\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{C}_{10}\text{H}_6-\text{N}(\text{C}_4\text{H}_9)_2$
402	$\text{O}_3\text{S}-(\text{CH}_2)_3-\text{N}^+\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{C}_{10}\text{H}_6-\text{N}(\text{C}_8\text{H}_{17})_2$
501	$\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+\text{C}_6\text{H}_4-\text{CH}=\text{CH}-\text{C}_{10}\text{H}_6\text{N}$
601	$\text{C}_6\text{H}_4-\text{N}^+(\text{CH}_2)_4-\text{SO}_3^--\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
701	$\text{C}_{10}\text{H}_6-\text{N}^+(\text{CH}_2)_4-\text{SO}_3^--\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
702	$\text{C}_6\text{H}_4-\text{N}^+(\text{CH}_2)_3-\text{SO}_3^--\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_6\text{H}_{13})_2$

5

801	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
802	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_4\text{H}_9)_2$
803	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_6\text{H}_{13})_2$
804	$(\text{CH}_3)_2\text{N}-(\text{CH}_2)_3-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$ Ψ^-
805	$^-\text{OOC}-(\text{CH}_2)_2-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
806	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_3(\text{OCH}_3) \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
807	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_3(\text{N}(\text{C}_2\text{H}_5)_2) \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
808	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_5\text{H}_{11})_2$
809	$^-\text{O}_3\text{S}-(\text{CH}_2)_4-\text{N}^+ \begin{array}{c} \text{C}_6\text{H}_5 \\ \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array} -\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{N}(\text{C}_6\text{H}_{13})_2$

901	
902	
903	
904	
905	
906	

The synthesis of merocyanine dyes, including styryl dyes, is well documented. For example, the preparation of selected styryl dyes (RH dyes) is described by Rina Hildesheim (Grinvald et al., BIOPHYS. J. 39, 301 (1982), Leslie Loew (Loew et al., J. ORG. CHEM. 49, 2546 (1984), and others. Other merocyanine dyes are described by Brooker et al. (J. AM. CHEM. SOC. 73, 5326 (1951).

Many dyes useful for the invention are available from Molecular Probes, Inc. (Eugene,

Oregon).

The quaternizing moiety ("TAIL") is typically obtained by quaternization of a methyl-substituted pyridine, quinoline or other nitrogen heterocycle by an alkylating agent that already
5 contains other side chain substituents, or that is further reacted to yield the side chain substituents as described by Hildesheim (Grinvald et al. *supra*). In particular, quaternization with propane sultone or butane sultone is convenient and yields preferred dyes. Other substituents are usually added either before or after synthesis of the initial merocyanine dye via an intermediate haloalkyl quaternized heterocycle. However, addition of TAIL is also achieved subsequent to initial dye
10 synthesis, which usually requires an azastilbene or fused analog thereof, as described by Loew et al. (*supra*).

The styryl dyes of the present method are typically prepared by conversion of an aniline derivative that contains the desired R^1 , R^2 , R^3 , and R^4 substituents to a benzaldehyde,
15 cinnamaldehyde or pentadienal derivative using methods well known in the art. The aldehyde derivative is then condensed with a quaternized pyridinium, quinolinium or benzazolium salt to give the useful dye.

Method of Use

20

A staining mixture comprising one or more of the merocyanine dyes described above are combined with a sample mixture that is thought to contain poly(amino acids) for a sufficient time to form a dye-poly(amino acid) complex that gives a detectable colorimetric or fluorescent optical response upon illumination. The complex optionally contains detergent molecules, as described
25 above. The merocyanine dyes present in the complex interact non-covalently with either the poly(amino acid) itself, or with said detergent molecules present in the complex. The sample mixture or the combined mixture is optionally heated, typically to $> 90^\circ\text{C}$. Additional steps are optionally and independently used, in any combination, to provide for separation or purification of the poly(amino acids), for enhancing the detection of the poly(amino acids), or for quantification
30 of the poly(amino acids).

Sample Mixture

The sample mixture is an aqueous or mostly aqueous solution, or is a solid, paste,

emulsion or other solution that is combined with a mostly aqueous solution in the course of labeling. The sample mixture contains or is suspected to contain poly(amino acids). Where the sample mixture is an aqueous solution, the concentration of poly(amino acids) is typically 10 ng/mL - 50 μ g/mL, more preferably 50 ng/mL - 5 μ g/mL. For electrophoretic gels, the
5 concentration of poly(amino acids) is typically 1 ng/band - 4 μ g/band.

Poly(amino acids) suitable for the invention include both synthetic and naturally occurring poly(amino acids). The poly(amino acids) optionally incorporate non-peptide regions including lipid, phosphate, and/or carbohydrate regions. The poly(amino acids) are optionally relatively
10 homogeneous or heterogeneous mixtures of poly(amino acids), or are multi-subunit complexes. The poly(amino acids) in the sample mixture are optionally covalently or non-covalently bound to a solid surface such as glass, plastic, or semiconductor material; or are unbound.

The poly(amino acids) are optionally unmodified, or have been treated with a reagent so as
15 to enhance or decrease the mobility of the poly(amino acid) in an electrophoretic gel, such as by complexing with the peptide (to decrease migration), by cleaving selected peptide bonds (to increase migration of the resulting fragments), by changing the relative charge on the protein (as by phosphorylation or dephosphorylation) or by covalent coupling of a constituent such as occurs during glycosylation. Such interactions are detected by the change in electrophoretic mobility of
20 treated poly(amino acids), relative to untreated poly(amino acids).

Individual amino acids have been stained with these dyes, but typically poly(amino acids) are peptides of mw > 500 daltons (more typically > 800 daltons) or are proteins (Tables 3 and 5). Large poly(amino acids) (> 200,000 daltons) are not well resolved in gels. Smaller poly(amino
25 acids) (< 1000 daltons) are difficult to detect on gels and filters, but are readily detected in solution. In one embodiment of the invention, the poly(amino acids) present are a mixture of poly(amino acids) of different molecular weights (e.g. molecular weight standards).

The sample mixture optionally contains discrete biological ingredients other than the
30 desired poly(amino acids), including other poly(amino acids), amino acids, nucleic acids, carbohydrates, and lipids, which may or may not be removed in this method. In one embodiment, the poly(amino acids) are separated from each other or from other ingredients in the sample by mobility, by size or by binding affinity in the course of the method. Typically intact or fragmented biological membranes, liposomes or detergent micelles in the sample mixture are removed,

destroyed or dispersed below the concentration at which they assemble into micelles (critical micelle concentration or CMC) prior to or in the course of labeling with this method. Typically, the sample mixture is essentially cell-free.

5 Staining mixture

The staining mixture is combined with the sample mixture so as to facilitate contact between any dye and any poly(amino acids) present in the combined mixture. The staining mixture is typically prepared by dissolving the dye in an organic solvent, such as DMSO, DMF or methanol; and then diluted with an aqueous solution according to the assay being performed. For solution assays, the dye is diluted into an aqueous solution, preferably buffered, that optionally contains a detergent. For staining on gels or membranes, the dyes are diluted into water or buffer, optionally including acetic acid (typically 5%-7%).

For visible color detection, dye concentrations in the staining mixture are typically between 1 μM and 100 μM , preferably between about 5 μM and about 20 μM ; more preferably at least 10-15 μM or higher. For fluorescence detection, dye concentrations are typically greater than 0.10 μM and less than 10 μM ; preferably greater than about 0.50 μM and less than or equal to about 5 μM ; more preferably 1-3 μM . Concentrations below and above these values result in detectable staining for certain poly(amino acids), depending on the sensitivity of the detection method.

A particular dye is generally selected using one or more of the following criteria: sensitivity to poly(amino acids), dynamic range, photostability, staining time, and insensitivity to the presence of nucleic acids. The sensitivity and dynamic range of the dyes is determined using the procedures of Examples 1 and 7. Preferred dyes have a sensitivity of 1-2 ng or less of poly(amino acid) per band in electrophoretic gels, or 10-30 ng or less of poly(amino acid) per mL of solution. Preferably, the dyes have a dynamic range of about 3 or more orders of magnitude of poly(amino acid) concentration for solution assays. Preferred dyes, in an aqueous solution in the absence of poly(amino acids), possess a quantum yield < 0.05 , more preferably < 0.01 . Preferably, when combined with poly(amino acids), the dyes exhibit a fluorescence enhancement that is > 100 -fold, more preferably > 300 -fold relative to the dyes in the absence of poly(amino acids). Preferred dyes, when combined with poly(amino acids), have an absorption maximum within 10 nm of 488 nm, 514 nm, 543 nm, or 633 nm.

Detergent

Except where non-denaturing gels are being used, the invention optionally includes a detergent that is added simultaneously with or as part of the sample mixture or the staining mixture, or is added to the combined mixture as described below. Preferably, the detergent is combined with the sample mixture before the staining mixture is added. The detergent is any amphiphilic surface active agent or surfactant that serves to coat the poly(amino acids), i.e. non-covalently associate with the poly(amino acid). Useful detergents include non-ionic, cationic, anionic, amphoteric and fluorinated surfactants; many are commercially available. Any detergent utilized in protein gel electrophoresis is preferred. Typically, the detergent is an anionic detergent, preferably an alkyl sulfate or alkyl sulfonate salt (typically 6-18 carbons); more preferably, sodium dodecyl sulfate (SDS), sodium octadecyl sulfate, or sodium decyl sulfate; most preferably, SDS.

Preferably any detergent in the sample mixture, staining mixture, or combined mixture is present below the CMC for that detergent, in order to avoid poly(amino acid)-free micelle formation. The CMC is a function of the detergent being used and the ionic strength of the solution. For SDS solutions at moderate ionic strength, the CMC is about 0.1% of the solution by weight. Typically, for prestaining the sample mixture before electrophoretic separation of denatured poly(amino acids) using SDS, the concentration of the detergent is about 1-5% by weight, more typically about 2%. Where the combined mixture is used for a solution assay, the concentration of SDS in the combined mixture is typically less than 1% by weight, preferably 0.05-0.1% by weight. For non-ionic detergents such as PLURONIC and TRITON, the concentration of detergent in the combined mixture is preferably less than about 0.05% by weight.

Combined Mixture

In one aspect of the invention, the combined mixture is a solution (e.g. Example 7; Table 3); typically, an aqueous solution (preferably buffered) that consists essentially of poly(amino acids), one or more merocyanine dyes, particularly preferred dyes, and a detergent. The aqueous solution is optionally used as a separation medium, such as within a sedimentation gradient (e.g. a sucrose gradient) or when performing capillary electrophoresis. The sample mixture or combined mixture is optionally heated and cooled before the detection or quantification of poly(amino acids).

Table 3: Properties of selected dyes in solution assays

Dye No.	Absorbance (nm) ¹		Fluorescence (nm) ² dye + protein		Fluorescence Enhancement ³	Dynamic Range Limits ⁴ (μ g protein)	
	Free Dye	Dye+ Protein	Excitation	Emission		Lower	Upper
5	101	448	447	469	583	1.1	10
	102	509	508	526	624	2	0.05
	104	463	468	469	570	12.7	0.05
	106	440	437	448	578	1.1	50
10	201	480	507	494	635	142	0.05
	301	473	474	473	576	23.5	0.5
	302	479	476	469	576	188	0.05
	303	475	476	472	569	363	0.05
	304	464	473	470	569	533	0.05
	305	465	465	471	568	12.3	0.5
15	306	472	472	469	567	307	0.05
	307	461	472	469	569	443	0.05
	308	458	470	470	568	202	0.05
	309	507	507	483	629	283	0.05
	310	507	507	483	628	356	0.05
	311	479	508	504	669	158	0.25
20	401	472	476	470	611	679	0.25
	402	455	467	472	597	20.2	0.05
	501	506	506	496	589	76.2	0.25
	601	471	472	545	584	154	0.05
25	701	508	513	526	594	534	0.25
	801	538	543	547	631	1285	0.25
	802	542	543	554	623	351	0.05
	803	507	537	544	630	2238	0.25
	901	506	506	560	570	66.2	2.5

¹ Absorbance maxima are determined using 1 μ M dye + 0.05% SDS +/- 150 μ g/mL BSA (bovine serum albumin) in a 2.0 mL volume.

² Fluorescence excitation and emission spectra are measured using 1 μ M dye + 150 μ g/mL BSA + 0.05% SDS in 2.0 mL volume.

³ Fluorescence enhancements are calculated using 1 μ M dye +/- 500 μ g/mL BSA in 2.0 mL at fluorescence excitation and emission maxima.

⁴ Dynamic range limits are determined by titrating 1 μ M dye with 0.05 to 50 μ g protein in a 96-well microplate in a 200 μ L volume. Three protein solutions are titrated separately for each dye: lysozyme, BSA and a protein mixture that contains equal weight concentrations of IgG, avidin, streptavidin, cellulase and ovalbumin. The lower limit of detection is the amount of protein required to produce fluorescence 10% above background. The upper limit is where the linearity of signal with respect to protein concentration starts to degrade for the protein type producing signal saturation at the lowest concentration.

In another aspect of the invention, the merocyanine dyes are optionally used to prestain poly(amino acids) prior to separation; or are present as a component of the mobile phase during separation (e.g. Example 12) in gel or capillary electrophoresis. Alternatively, separated poly(amino acids) in electrophoretic (denaturing or non-denaturing) gels are post-stained using the staining mixture, or are transferred to a filter membrane or blot or other solid or semi-solid matrix before being combined with the staining mixture (e.g. Examples 1, 2, 3, and 5). Typically, polyacrylamide or agarose gels are used for electrophoresis. Alternatively, the gel is an iso-electric focusing gel, a gradient gel, a two-dimensional gel, or the gel is used for gel-mobility-shift analysis. The sample mixture or combined mixture is optionally heated before being applied to denaturing gels. The sensitivities of selected dyes of the present invention when used to post-stain electrophoretic gels are listed in Table 4.

Table 4: Labeling sensitivity in gels of selected dyes¹

Dye No.	Color ²	Sensitivity ³ (ng/band)	Background Staining
Coomassie Brilliant Blue	blue*	30-60 (low)	very high: requires destaining
Nile red	red/orange*	30-60 (low)	low/med
Silver staining	tan to black*	variable 2-10 (high)	varies, low to very high even with destaining
101	faint greenish	low	med
102	light orange	med	med

	103	red/orange	low	low/med
	104	yellow	med	low/med
	201	red/orange	med	low/med
	301	barely visible	low	low
5	302	greenish/pink	low	low
	303	pink/yellow to orange	med	med
	304	bright orange	high	low/med
	305	light orange	low	med
	306	orange	med	low
10	307	bright orange	high	low/med
	308	orange	high	med
	309	pink/orange	med	low
	310	red/orange	high	low/med
	311	light orange	low	low/med
15	401	orange	med	med
	402	red/orange	high	low
	501	orange	med	low/med
	601	red/orange	high	low
	701	red/orange	high	low
20	801	red	high	low
	803	red	high	low
	802	red	high	low
	901	orange	med	med/high

¹ All electrophoretic gels are SDS acrylamide mini-gels (Bio-Rad)

25 ² Color refers to the color of the fluorescence emission of the stained protein bands, except where marked with * where color refers to colorimetric appearance under white light.

30 ³ Sensitivity refers to the lower limit of protein detection, per band, using a 4X dilution series of BSA (using the procedure of Example 1). The sensitivity is defined to be the band containing the smallest amount of protein that is easily detectable by eye in a black and white Polaroid photograph taken at an optimal exposure. "Low" sensitivity means a detection limit comparable to that found with CBB and Nile red; "high" sensitivity means a detection limit comparable to that found with silver staining; "medium" sensitivity is intermediate between the two ranges.

The sensitivity of the present method for a variety of poly(amino acids) was compared to that of silver staining and CBB staining molecular weights (Table 5).

Table 5: Sensitivity of detection as a function of poly(amino acid) composition

Protein	Molecular Weight (daltons)	Sensitivity Dye 304 (ng)	Sensitivity Silver Staining (ng)	Sensitivity CBB Staining (ng)
myosin	200,000	1	1	8-16
β -galactosidase	116,250	1	2-4	8-16
phosphorylase B	97,400	2-4	1	30-60
fructose-6-phosphate kinase	85,200	1-2	2-4	31-62
bovine serum albumin	66,200	2-4	2-4	16-30
bovine cytochrome C oxidase (COX) subunit I	56,900	1-2	26-52	105
glutamate dehydrogenase	55,400	1-2	2-4	16
IgG (heavy chain)	50,000	2	4	34-69
ovalbumin	45,000	1-2	1-2	16-30
Protein A	41,000	1	1-2	16
aldolase	39,200	1-2	1	31-62
carbonic anhydrase	31,000	2	4-8	16-30
COX subunit II	29,900	1-2	2-4	27-54
triose phosphate isomerase	26,600	1	1	16-31
COX subunit III	26,000	0.8	0.8-1.5	48
Protein G	26,000	2	16	32-64
IgG (light chain)	25,000	14	56	166
soybean trypsin inhibitor	21,500	1-2	2-4	16-30
histone H1	20,000	10	80	80-160
β -bungarotoxin subunit	20,000	1.25-2.5	1.25	41
COX subunit IV	17,100	0.5-1	4-8	16-32
avidin	4 x 16,500	2	2-4	16
streptavidin	4 x 15,000	2	16	32-64

5	histone H3	15,700	7.5	15	30-60
	NEUTRALITE avidin	15,000	1-2	62	62
	lysozyme	14,400	1-2	4	16-30
	histone H2A/H2B	13,700	7	14-27	54
	pancreatic RNase A	13,700	4-8	16-30	125
10	COX subunit Va	12,400	1	45	16-32
	histone H4	11,200	3	22-45	45
	COX subunit Vb	10,700	1	4-8	16-32
	COX subunit VIa	9,400	1	4-8	16-32
	α -bungarotoxin	8,000	4-8	2-4	31-62
	β -bungarotoxin subunit	7,000	3-6	—	175-350
	aprotinin	6,500	2	4-8	16-30
	human insulin	6,000	8	8	125-250

15 Using the present invention, destaining of stained gels is generally not necessary for either colorimetric or fluorescent detection. At very high dye concentrations, destaining is optionally used to improve visible color detection in gels. Gels are optionally washed briefly after staining to prevent transfer of dye to other surfaces.

20 Additional reagents

 The method of the present invention optionally further comprises addition of an additional reagent, simultaneously or sequentially, to the sample mixture, the staining mixture, or the combined mixture. The additional reagent incorporates a means for producing a detectable
25 response, including the change in, or appearance of, color, fluorescence, reflectance, pH, chemiluminescence, infrared spectra, magnetic properties, radioactivity, light scattering, x-ray scattering, or production of an electron-rich substrate. An additional reagent is optionally a detection reagent that colocalizes with poly(amino acids) in general or with specific poly(amino acids); or is a probe for a specific component of the sample mixture (e.g. a nucleic acid stain; a
30 detectable member of a specific binding pair); or changes the mobility of the poly(amino acid) as described above. In one embodiment of the invention, the additional reagent is one or more additional merocyanine dyes, including preferred embodiments described above. Where the

additional merocyanine dye(s) have overlapping spectral characteristics such that energy transfer occurs, the labeled poly(amino acids) exhibit an extended Stokes shift. Alternatively, the additional reagent is another protein stain (such as CBB or silver stain) such that labeling of the poly(amino acids) is enhanced by the colocalization of staining.

5

Illumination and Observation

The combined mixture of dye and sample is illuminated to give an optical response that is an absorption of visible light (colorimetric response), or is a fluorescence emission (fluorescence response), or both. Illumination is by a light source capable of exciting the dye-poly(amino acid) complex, typically at or near the wavelength of maximum absorption of the dye-poly(amino acid) complex, such as an ultraviolet (254-370 nm) or visible (490-550 nm) wavelength emission lamp, an arc lamp, a laser, or even sunlight or ordinary room light. Preferably, the sample is excited with a wavelength within 20 nm of the maximum absorption of the dye-poly(amino acid) complex.

15

Preferably, the dye-poly(amino acid) complexes possess an absorption maximum between 480 and 650 nm, more preferably between 488 and 550 nm, most preferably matching the wavelength of a laser illumination source. The complexes also preferably excite in the UV at or near 300 nm.

20

The optical response is detected qualitatively, or optionally quantitatively, by means that include visual inspection, CCD cameras, video cameras, photographic film, or the use of instrumentation such as laser scanning devices, fluorometers, photodiodes, quantum counters, epifluorescence microscopes, scanning microscopes, flow cytometers, fluorescence microplate readers, or by means for amplifying the signal such as photomultiplier tubes. Recording the optical response using POLAROID film results in enhanced sensitivity of signal.

25

Quantification of the poly(amino acid) is typically performed by comparison of the optical response to a prepared standard or to a calibration curve, obtained from a standard dilution of a known concentration of a poly(amino acid) or poly(amino acid) mixture. Alternatively, the standard curve is generated by comparison with a reference dye or dyed particle that has been standardized versus the target dye-poly(amino acid) complex.

30

Due to the simplicity of use of the instant dyes, said dyes are particularly useful in the

formulation of a kit for the labeling of poly(amino acids), comprising one or more merocyanine dyes (preferably in a stock solution) and instructions for the use of the dye, optionally including a reference standard. In one embodiment, the instructions specify that the sample mixture and staining mixtures are applied, either simultaneously or sequentially to a solid or semi-solid support, the resulting color or fluorescence of which is then compared to a calibrated standard to determine the presence, and optionally the concentration, of poly(amino acid) present in the sample mixture. The support, dye solution, calibration standard and instructions for use, in combination, comprise a "dipstick" protein assay kit, allowing the rapid and convenient determination of protein concentration.

10

The examples below are given so as to illustrate the practice of this invention. They are not intended to limit or define the entire scope of this invention.

EXAMPLES

15

Example 1. Detection of proteins in SDS-polyacrylamide gels:

The pure protein or mixture of proteins is prepared in Loading Buffer (50 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol, and 0.015% bromophenol blue). Dithiothreitol is added to each sample, to 0.1 M. The samples are heated 4-5 minutes at 90-95 °C and loaded onto a 15% Tris-glycine polyacrylamide mini- or full-sized gel containing 0.05%-0.1% SDS, with a 4% stacking gel. The gel is electrophoresed under standard conditions, in Tris-glycine buffer containing 0.05%-0.1% SDS. The resulting gel is transferred to a staining dish containing 1-3 µM Dye 304 in 7.5% acetic acid (in water). The staining solution is gently agitated for 45-60 minutes. After 10 minutes, the protein bands are readily apparent, but sensitivity improves over about 30-40 minutes. The gel is removed from the staining dish, rinsed briefly in 7.5% acetic acid and transferred to a UV-transilluminator. The gel is photographed using 300 nm transillumination and black and white POLAROID 667 print film with a Wratten 9 gelatin filter, or using a CCD camera with UV illumination near 300 nm. The stained bands exhibit bright orange fluorescence. Under normal room lights, the stained bands appear as orange bands. Other dyes of the invention (See Table 3) yield stained gels that possess bands having visible coloring from yellow to purple and fluorescence emission from green to red.

30

Example 2. Detection of proteins in non-denaturing polyacrylamide gels:

A protein dilution series is prepared in Native Gel Loading Buffer (125 mM Tris-HCl, pH 6.8, 10% glycerol and 0.015% bromophenol blue). The samples are loaded onto a Tris-HCl non-denaturing polyacrylamide gel, and the gel is electrophoresed under standard conditions. The electrophoresed gel is stained and photographed as in Example 1. Staining sensitivity under these conditions is typically somewhat protein-selective.

Example 3. Detection of peptides in Tris-Tricine gels:

A dilution series of a short peptide is prepared in Loading Buffer. To each sample is added dithiothreitol to 0.1 M, the samples are heated 4-5 minutes at 90-95 °C then loaded onto 16.5% Tris-tricine gel. The gel is electrophoresed under standard conditions. The gel is stained and the protein bands are visualized as in Example 1. Polypeptides as small as 20 amino acids are readily detected using this method. Peptides such as a tetramer of the heptapeptide repeat found at the C-terminus of eukaryotic RNA polymerase II, tryptic peptides of trypsin and the small subunit of β -bungarotoxin, which are not stained with either CBB or silver staining, are readily detectable.

Example 4. Detection of proteins in 2-dimensional gels:

A mouse cytosolic extract (i.e., a whole cell lysate) is prepared and separated on a 2-dimensional gel under standard conditions. The gel is stained and visualized as in Example 1. An identical gel is stained in parallel with silver using a commercially available kit (Bio-Rad). The gel stained using the present invention exhibits more protein spots than the silver stained gel, indicating that the present invention is less protein-selective than silver staining. The signal intensity obtained with the instant method is more directly proportional to protein concentration than that obtained with silver staining. The sensitivity of staining using the present invention is superior to that of silver staining for detection of low molecular weight proteins (Table 3).

Example 5. Detection of proteins on filter membranes:

The protein is diluted in TBS (20 mM Tris-HCl, pH 7.5, 500 mM NaCl), then applied to a nylon or nitrocellulose filter membrane. The membrane is washed with TBS, then incubated with 2 μ M Dye 803 in 7.5% acetic acid or TBS. Alternatively, the proteins are first separated by gel electrophoresis and transferred to a nylon or nitrocellulose filter membrane using standard procedures. The blot is stained as above. The blot is illuminated at 300-365 nm. Protein spots or

bands are fluorescent. Blots are photographed as in Example 1. Stained proteins appear as faint white spots or white bands. Staining under these conditions is somewhat protein selective. Proteins are optionally detected using laser illumination sources.

5 Example 6. Western blotting of stained proteins:

A dilution series of bovine heart cytochrome oxidase complex (COX) is prepared in water. Samples are loaded onto a 12% polyacrylamide gel and the gel is electrophoresed under standard conditions. The resulting gel is cut in half, and one half is stained with Dye 801 in transfer buffer
10 (20% MeOH, 25 mM Tris-glycine, pH 8.3, 192 mM glycine) and visualized as in Example 1 above. The proteins are electrophoretically transferred from both halves of the gel to a filter membrane, according to standard methods. The blot is blocked, probed with mouse monoclonal antibodies directed against specific COX subunits, and the resulting signals are visualized using nitroblue tetrazolium and 5-bromo-4-chloro-3-indolyl phosphate, using standard procedures. The
15 two halves of the blot show equal sensitivity for detection of the COX subunits, indicating that staining proteins with the invention does not interfere with later immunological detection.

Example 7. Detection of proteins in solution:

20 A dilution series of proteins is prepared in 10 mM Tris-HCl, pH 7.5, containing 0.05% SDS. An equal volume of 2 μ M Dye 307 is added to each sample. The samples are incubated 15 minutes at room temperature. The fluorescence intensity of each sample is measured using 490 nm excitation, using a fluorometer or fluorescence microplate reader. The fluorescence of Dye 307 in buffer alone (control) is subtracted from that of the protein-containing samples. Protein
25 concentrations are determined by comparing the net fluorescence with that of a standard, such as bovine serum albumin, or protein mixture (Figure 3).

Example 8. Detection of proteins in gels in the presence of nucleic acids:

30 An extract from *E. coli* is prepared using standard methods, by suspending a bacterial cell pellet directly in Loading Buffer and heating it as in Example 1. The resulting sample is pipeted up and down repeatedly, through a fine bore syringe, to shear large DNA molecules. The sample is diluted serially and loaded onto a denaturing SDS gel in two duplicate dilution series and electrophoresed, as in Example 1. The electrophoresed gel is cut in half. One half is stained with

Dye 304 according to Example 1. The other half is stained with silver according to standard methods. The silver-stained half shows staining of bands that correspond to nucleic acids as well as protein. The half stained with Dye 304 exhibits only those bands that result from protein staining.

5

Example 9. Detection of proteins in gels using a laser-excited gel scanner:

Gels are prepared and stained as in Examples 1, 2, 3 or 4. The stained gels are scanned using a laser-excited gel scanner. Stained proteins display fluorescence emission having a
10 dynamic range of at least 3 orders of magnitude in protein concentration, with a sensitivity of a single ng per protein band (Figure 4, Table 2). Stained gels can also be analyzed using visible light lasers, or other scanners with light sources that overlap with the excitation spectra for the dye used.

15 Example 10. Detection of proteins spotted on a TLC plate or filter membrane:

Dilution series of BSA, lysozyme and an equi-mass mixture of myosin, β -galactosidase, phosphorylase B, bovine serum albumin, ovalbumin, carbonic anhydrase, trypsin inhibitor, lysozyme and aprotinin are prepared in 10 mM Tris-HCl, pH 7.5, containing 0.05% SDS. An
20 equal volume of Dye 803 in the same buffer is added to each and the samples are incubated for 15 minutes. A 1-5 μ L aliquot of each mixture is spotted onto a thin layer chromatography plate or a nylon or nitrocellulose filter membrane. Protein-containing spots are detected colorimetrically under room light, or are detected by fluorescence following ultraviolet or visible illumination. The color intensity or fluorescence intensity of a given spot is indicative of the amount of protein
25 present in that spot. The amount of protein in an unknown sample is determined by comparing either the colorimetric or fluorescence intensity with that of a standard of known concentration.

Example 11. Staining protein gels with dye in the running buffer:

30 Proteins are loaded and run on standard SDS gels, using standard methods, except that the running buffer contains 0.05% SDS and 1-3 μ M Dye 801. The stained gels are photographed directly after electrophoresis (as in Example 1) or are destained in 7.5% acetic acid for 20-50 minutes prior to photography. The sensitivity obtained is about the same as obtained by staining gels after electrophoresis. Migration of protein bands can be monitored through the glass plates

that support the gel, during electrophoresis.

Example 12. Prestaining of proteins prior to electrophoresis:

5 Proteins are diluted in Loading Buffer, heated to 90-95 °C for 4-5 minutes then cooled to room temperature. Dye 304 is added to 10 mM, and the samples are loaded onto a 12% polyacrylamide gel. The gel is electrophoresed and visualized using ultraviolet illumination as in Example 1, or as in Example 9, or with a CCD camera. The sensitivity of this method is somewhat less than in Examples 1 and 11.

10

It is to be understood that, while the foregoing invention has been described in detail by way of illustration and example, numerous modifications, substitutions, and alterations are possible without departing from the spirit and scope of the invention as described in the following claims.

15


5 What is claimed is:

1. A method of detecting a poly(amino acid), comprising the steps of:
 10 a) combining a sample mixture that is thought to contain a poly(amino acid) with a staining mixture that contains one or more merocyanine dyes to form a combined mixture; wherein each merocyanine dye independently has the formula

Q-B-M


15

where Q is



(Q1)


20 or



(Q2)

or

25



(Q3)

5 where

R^1 , R^2 , R^3 and R^4 are independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls;


10 R^5 and R^6 are independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls, or phenyl; or R^5 and R^6 taken in combination form a fused 6-membered aromatic ring that is unsubstituted or independently substituted one or more times by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2
15 C_1 - C_6 alkyls, or said fused 6-membered aromatic ring is unsubstituted or substituted by an additional fused 6-membered aromatic ring that is unsubstituted or independently substituted by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls;

20 X is -S-, -O-, -NR⁷-, or -CR⁷R⁸-, wherein R^7 and R^8 are independently H, C_1 - C_6 alkyl or phenyl; or R^7 and R^8 taken in combination complete a 5- or 6-membered saturated ring;

TAIL is attached to Q through a carbon atom and contains 1-22 non-hydrogen
25 atoms, that are C, O, N or S, such that each heteroatom is separated from any adjacent heteroatoms by two or more carbon atoms; and further such that TAIL is composed of carbon-carbon bonds (C-C), ether bonds (C-O-C), thioether bonds (C-S-C) or amine bonds (C-NR⁹-C); where any carbon atom in TAIL is unsubstituted or substituted by hydroxy, carboxy, sulfo, amino or
30 ammonium; and where any amine bond, amino or ammonium in TAIL is unsubstituted or substituted by hydroxy, carboxy, sulfo, amino, amino substituted by C_1 - C_6 alkyls, or ammonium substituted by 1-3 C_1 - C_6 alkyls, or said N atoms form either one or two saturated 5- or 6-membered rings in combination with additional C or N atoms in TAIL;

35

B is




5

where R^7 and R^8 are as defined previously;


$n = 1, 2$ or 3 ;

10 and M is



(M1)

15 or



(M2)

where


20

R^{11} and R^{12} are independently H, F, Cl, or $-CH_3$;

25

R^{13} and R^{14} are independently C_1 - C_{18} alkyl that is linear, branched, saturated or unsaturated, and is unsubstituted or substituted one or more times by F, hydroxy or C_1 - C_6 alkoxy; or R^{13} and R^{14} taken in combination form a 5- or 6-membered saturated ring containing 0 or 1 oxygen heteroatoms; or R^{13} taken in combination with R^{11} and R^{14} taken in combination with R^{12} independently are $-(CH_2)_2-$ or $-(CH_2)_3-$;

30 or M is



5

where R is independently H, phenyl, sulfophenyl, C1-C6 alkyl, or C1-C6 alkyl substituted by carboxy;

- 10 b) incubating the combined mixture for a time sufficient for the dye in the staining mixture to associate with the poly(amino acid) in the sample mixture to form a dye-poly(amino acid) complex that gives a detectable optical response upon illumination;
- c) illuminating said dye-poly(amino acid) complex; and
- d) observing said detectable optical response, providing that the staining mixture does

- 5 not consist of a single merocyanine dye that is a 4'-hydroxy-3',5'-dialhalostyryl merocyanine dye having a pKa value that changes in the presences of a poly(amino acid).


2. A method of detecting a poly(amino acid), comprising the steps of:

- 10 a) combining a sample mixture that is thought to contain a poly(amino acid) with a staining mixture that contains one or more styryl dyes having the formula

Q-B-M


where Q is

15



(Q1)


or



20

(Q2)

or



25

(Q3)

where

5 R^1 , R^2 , R^3 and R^4 are independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls;


R^5 and R^6 are independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls, or phenyl; or R^5 and
 10 R^6 taken in combination form a fused 6-membered aromatic ring that is unsubstituted or independently substituted one or more times by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls, or said fused 6-membered aromatic ring is unsubstituted or substituted by an additional fused 6-membered aromatic ring that is
 15 unsubstituted or independently substituted by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls;

X is -S-, -O-, -NR⁷-, or -CR⁷R⁸-, wherein R^7 and R^8 are independently H, C_1 - C_6 alkyl or phenyl; or R^7 and R^8 taken in combination complete a 5- or 6-
 20 membered saturated ring;

TAIL is attached to Q through a carbon atom and contains 1-22 non-hydrogen atoms, that are C, O, N or S, such that each heteroatom is separated from any adjacent heteroatoms by two or more carbon atoms; and further such that
 25 TAIL is composed of carbon-carbon bonds (C-C), ether bonds (C-O-C), thioether bonds (C-S-C) or amine bonds (C-NR⁹-C); where any carbon atom in TAIL is unsubstituted or substituted by hydroxy, carboxy, sulfo, amino or ammonium; and where any amine bond, amino or ammonium in TAIL is unsubstituted or substituted by hydroxy, carboxy, sulfo, amino, amino
 30 substituted by C_1 - C_6 alkyls, or ammonium substituted by 1-3 C_1 - C_6 alkyls, or said N atoms form either one or two saturated 5- or 6-membered rings in combination with additional C or N atoms in TAIL;

B is

35




where R^7 and R^8 are as defined previously;

5

$n = 1, 2 \text{ or } 3;$

and M is


10



(M1)

or

15



(M2)

where

20

R^{11} and R^{12} are independently H, F, Cl, or $-\text{CH}_3$;

25

R^{13} and R^{14} are independently $\text{C}_1\text{-C}_{18}$ alkyl that is linear, branched, saturated or unsaturated, and is unsubstituted or substituted one or more times by F, hydroxy or $\text{C}_1\text{-C}_6$ alkoxy; or R^{13} and R^{14} taken in combination form a 5- or 6-membered saturated ring containing 0 or 1 oxygen heteroatoms; or R^{13} taken in combination with R^{11} and R^{14} taken in combination with R^{12} independently are $-(\text{CH}_2)_2-$ or $-(\text{CH}_2)_3-$;

to form a combined mixture;

30

- b) incubating the combined mixture for a time sufficient for the dye in the staining mixture to associate with the poly(amino acid) in the sample mixture


- 5 to form a dye-poly(amino acid) complex that gives a detectable optical response upon illumination;
- c) illuminating said dye-poly(amino acid) complex; and
- d) observing said detectable optical response.

10 3. A method, as in Claim 1 or 2, wherein for one dye, M has the formula M1.

4. A method, as in Claim 1 or 2, wherein for one dye, M has the formula M2.


5. A method, as in Claim 1 or 2, wherein one dye has the formula

15




6. A method, as in Claim 1 or 2, wherein one dye has the formula

20




or



25

7. A method, as in Claim 1 or 2, wherein one dye has the formula



5

- where each R^{10} is independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls, or any two adjacent R^{10} substituents, when taken in combination, form a fused 6-membered aromatic ring that is optionally and independently substituted by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls.

8. A method, as in Claim 1 or 2, wherein for one dye,

15

TAIL is $-CH_3$, or CH_2CH_3 ; or TAIL is a C_3 - C_{22} alkyl chain that is linear or branched, saturated or unsaturated, that is unsubstituted or substituted one or more times by hydroxy, carboxy, sulfo, amino, amino substituted by 1-2 C_1 - C_6 alkyls, or ammonium substituted by 1-3 C_1 - C_6 alkyls.

20

9. A method, as in Claim 1 or 2, wherein for one dye,

M has the formula M1 or M2;

X is O or S;

B is $-(CH=CH)_n-$;

25

R^{11} and R^{12} are each H;

R^{13} and R^{14} are independently C_4 - C_8 alkyl; and

TAIL is sulfopropyl, sulfobutyl, aminopropyl or aminobutyl, the amines of which are substituted 1-3 times by any combination of H, methyl or ethyl.

30

10. A method, as in Claim 9, wherein for one dye,

M has the formula M1 or M2;

- 5 R^1 , R^2 , R^3 , and R^4 are each H;
 R^5 and R^6 are each H; and

said dye is present in the combined mixture at a concentration of $0.10\ \mu\text{M}$ - $10\ \mu\text{M}$
 for a fluorescent optical response or $10\ \mu\text{M}$ - $100\ \mu\text{M}$ for colorimetric optical
 response; and said poly(amino acid) has a molecular weight of 500-200,000 daltons.

10

11. A method, as in any one of Claims 1-10, further comprising removing,
 destroying, or dispersing below the critical micelle concentration any biological
 membranes that are present in the sample mixture.

- 15 12. A method, as in any one of Claims 1-10, further comprising adding a
 detergent to the sample mixture, staining mixture or combined mixture.

13. A method, as in any one of Claims 1-12, wherein said detectable
 optical response is a fluorescent response.

20

14. A method, as in any one of Claims 1-13, further comprising
 quantitating said poly(amino acid) by measuring said detectable optical response and
 comparing said measurement with a standard.

- 25 15. A method, as in Claim 2, wherein for one dye, Q is Q1.

16. A method, as in Claim 2, wherein for one dye, Q is Q2.

17. A method, as in Claim 2, wherein for one dye, Q is Q3.

30

18. A method, as in any one of Claims 1-17, further comprising
 electrophoretically separating the sample mixture before, after, or while it is
 combined with the staining mixture.

- 35 19. A method, as in any one of Claims 1-18, further transferring the
 sample mixture to a solid or semi-solid matrix before or after combining with the
 staining mixture.

5 20. A method, as in any one of Claims 1-19, further comprising the step
of heating the sample mixture prior to combining with the staining mixture or heating
the combined mixture.

21. A combination consisting essentially of:

10

a) one or more styryl dyes as defined any one of Claims 2-10;

b) a detergent; and

15 c) a poly(amino acid),


in a cell-free aqueous solution where said detergent is present at a concentration less
than the critical micelle concentration for that detergent.

20 22. A kit for staining the poly(amino acids) in a sample, comprising:

a) one or more merocyanine dyes wherein each merocyanine dye independently has
the formula


25 Q-B-M

where Q is




30

or



or



where

R^1 , R^2 , R^3 and R^4 are independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls;


R^5 and R^6 are independently H, Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, or amino substituted by 1-2 C_1 - C_6 alkyls, or phenyl; or R^5 and R^6 taken in combination form a fused 6-membered aromatic ring that is unsubstituted or independently substituted one or more times by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls, or said fused 6-membered aromatic ring is unsubstituted or substituted by an additional fused 6-membered aromatic ring that is unsubstituted or independently substituted by Cl, F, C_1 - C_6 alkyl, C_1 - C_6 perfluoroalkyl, C_1 - C_6 alkoxy, amino, amino substituted by 1-2 C_1 - C_6 alkyls;

X is -S-, -O-, -NR⁷-, or -CR⁷R⁸-, wherein R^7 and R^8 are independently H, C_1 - C_6 alkyl or phenyl; or R^7 and R^8 taken in combination complete a 5- or 6-membered saturated ring;

TAIL is attached to Q through a carbon atom and contains 1-22 non-hydrogen atoms, that are C, O, N or S, such that each heteroatom is separated from any

5 adjacent heteroatoms by two or more carbon atoms; and further such that
 TAIL is composed of carbon-carbon bonds (C-C), ether bonds (C-O-C),
 thioether bonds (C-S-C) or amine bonds (C-NR⁹-C); where any carbon atom
 in TAIL is unsubstituted or substituted by hydroxy, carboxy, sulfo, amino or
 ammonium; and where any amine bond, amino or ammonium in TAIL is
 10 unsubstituted or substituted by hydroxy, carboxy, sulfo, amino, amino
 substituted by C₁-C₆ alkyls, or ammonium substituted by 1-3 C₁-C₆ alkyls, or
 said N atoms form either one or two saturated 5- or 6-membered rings in
 combination with additional C or N atoms in TAIL;

15 B is




where R⁷ and R⁸ are as defined previously;

20

n = 1, 2 or 3;


and M is



25

(M1)

or



30

(M2)


where


5

R^{11} and R^{12} are independently H, F, Cl, or $-CH_3$;

R^{13} and R^{14} are independently C_1 - C_{18} alkyl that is linear, branched, saturated or unsaturated, and is unsubstituted or substituted one or more times by F, hydroxy or C_1 - C_6 alkoxy; or R^{13} and R^{14} taken in combination form a 5- or 6-membered saturated ring containing 0 or 1 oxygen heteroatoms; or R^{13} taken in combination with R^{11} and R^{14} taken in combination with R^{12} independently are $-(CH_2)_2-$ or $-(CH_2)_3-$;

15 or M is





5

where R is independently H, phenyl, sulfophenyl, C1-C6 alkyl, or C1-C6 alkyl substituted by carboxy;

- b) instructions for combining said merocyanine dye or dyes with a sample containing or thought to contain poly(amino acids), such that the dye or dyes form a detectable label in conjunction with poly(amino acids) in the sample, providing that the kit does not consist of a merocyanine dye that is 4'-hydroxy-3',5'-dihalostryryl merocyanine dye having a pKa value that changes in the presence of a poly(amino acid).

15

23. A kit, as in Claim 22, where said instructions further specify subjecting said sample to electrophoresis before, during or after being combined with said dye or dyes.


20

24. A method, as claimed in Claim 10, wherein said merocyanine dye is present in the combined mixture at a concentration of 0.10 μM - 10 μM for a fluorescent optical response or 10 μM - 100 μM for colorimetric optical response; said poly(amino acid) has a molecular weight of 500-200,000 daltons; and said combined mixture optionally further comprises a detergent that is an alkyl sulfate or alkyl sulfonate salt that is present in a concentration of less than 0.1% by weight.

30


5

25. A kit, as in Claim 22 or 23, wherein each of said merocyanine dyes has the formula



10

or



15 wherein

R⁵ and R⁶ are optionally and independently H, Cl, F, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxy, amino, or amino substituted by 1-2 C₁-C₆ alkyls; or R⁵ and R⁶ taken in combination form a fused 6-membered aromatic ring that is optionally substituted or independently substituted one or more times by Cl, F, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxy, amino, or amino substituted by 1-2 C₁-C₆ alkyls;

TAIL is -CH₃, or CH₂CH₃; or TAIL is a C₃-C₂₂ alkyl chain that is linear or branched, saturated or unsaturated, that is unsubstituted or substituted one or more times by hydroxy, carboxy, sulfo, amino, amino substituted by 1-2 C₁-C₆ alkyls, or ammonium substituted by 1-3 C₁-C₆ alkyls;

n = 1 or 2; and

30

R¹³ and R¹⁴ are independently C₁-C₁₈ alkyl that is linear, branched, saturated or

5 unsaturated, and is unsubstituted or substituted one or more times by F.

Figure 1




Figure 2




Figure 3




Figure 4

