



US 20140037745A1

(19) **United States**

(12) **Patent Application Publication**
Liang et al.

(10) **Pub. No.: US 2014/0037745 A1**

(43) **Pub. Date: Feb. 6, 2014**

(54) **CONTROLLED RELEASE COMPOSITIONS
OF GAMMA-HYDROXYBUTYRATE**

continuation of application No. 11/239,638, filed on
Sep. 30, 2005, now Pat. No. 8,193,211.

(71) Applicant: **Supernus Pharmaceuticals, Inc.,**
Rockville, MD (US)

(60) Provisional application No. 60/614,622, filed on Sep.
30, 2004.

(72) Inventors: **Likan Liang**, Boyds, MD (US); **Niraj
Shah**, Owings Mills, MD (US);
Padmanabh P. Bhatt, Rockville, MD
(US); **Scott Ibrahim**, Owings Mills, MD
(US)

Publication Classification

(51) **Int. Cl.**
A61K 9/50 (2006.01)
A61K 31/19 (2006.01)

(73) Assignee: **Supernus Pharmaceuticals, Inc.,**
Rockville, MD (US)

(52) **U.S. Cl.**
CPC **A61K 9/5047** (2013.01); **A61K 31/19**
(2013.01); **A61K 9/5042** (2013.01); **A61K**
9/5031 (2013.01); **A61K 9/5073** (2013.01)
USPC **424/495**; 514/557; 424/490; 424/494;
424/497

(21) Appl. No.: **14/035,021**

(22) Filed: **Sep. 24, 2013**

(57) **ABSTRACT**

Related U.S. Application Data

(63) Continuation of application No. 13/476,369, filed on
May 21, 2012, now Pat. No. 8,598,191, which is a

The present invention is directed to oral pulse-release phar-
maceutical dosage form containing an immediate release
component of gamma-hydroxybutyric acid, and one or more
delayed/controlled release components of gamma-hydroxy-
butyric acid.




Figure 1




Figure 2




Figure 3

Figure 4. Dissolution profile of an immediate release core at pH 1.1




Figure 5. Dissolution profile of an Opadry AMB-coated immediate release core at pH 1.1





Figure 6. Dissolution profile of an EC-coated immediate release core at pH 1.1






Figure 7

CONTROLLED RELEASE COMPOSITIONS OF GAMMA-HYDROXYBUTYRATE

FIELD OF THE INVENTION

[0001] The present invention is directed to pulse-released formulations of oxybate, or gamma-hydroxybutyric acid, salts, which reduce the number of dosages typically required for treatment. For instance, in the treatment of narcolepsy, a twice-nightly dosage regimen can be reduced to a single dose with the compositions of the present invention.

BACKGROUND OF THE INVENTION

[0002] Sodium gamma-hydroxybutyrate (GHB or sodium oxybate) is a naturally occurring metabolite of many mammalian tissues (Fishbein et al, J. Biol. Chem. 239:357-61 (1964), Mamelak, Neurosci Biobehav Rev. 13(4):187-98 (1989), Nelson et al, J. Neurochem., 37:1345-48 (1981)) and has broad indications including narcolepsy, cataplexy, sleep paralysis, alcoholism, chronic schizophrenia, catatonic schizophrenia, atypical psychoses, chronic brain syndrome, neurosis, drug addiction and withdrawal, Parkinson's disease and other neuropharmacological illnesses, hypertension, ischemia, circulatory collapse, radiation exposure, cancer, myocardial infarction, anesthesia induction, sedation, growth hormone production, heightened sexual desire, anorectic effects, euphoria, smooth muscle relaxation, muscle mass production, and sleep.

[0003] Currently, sodium gamma-hydroxybutyrate is prescribed for patients with narcolepsy (Xyrem®, Orphan Medical) as a twice-nightly solution. Patients take an initial dose of sodium gamma-hydroxybutyrate around bedtime and must wake up four hours later to take a second dose. Such a dose regimen is rather inconvenient.

[0004] Other dosage forms of sodium gamma-hydroxybutyrate have also been disclosed. For example, U.S. Pat. No. 5,594,030 discloses controlled release pharmaceutical compositions of gamma hydroxybutyric acid salts consisting of a nucleus in the form of granulates or tablets which comprises GHB and a cellulosic matrix, wherein the drug substance is released within 7 to 8 hours.

[0005] Sodium gamma-hydroxybutyrate is highly soluble, hygroscopic, and strongly alkaline, and the therapeutic dose is normally very high. For example, a daily dose of 4.5 to 9 grams of Xyrem® is prescribed to narcolepsy patients. These characteristics of sodium gamma-hydroxybutyrate have some significant effects on coated particles or tablets comprising GHB. The high solubility of sodium gamma-hydroxybutyrate likely leads to drug migration into the coating layer during the coating process, and dissolves rapidly when the coated articles encounter water or bodily fluids, creating "pores" that allow leakage of the drug from the coated articles. Further, when sodium gamma-hydroxybutyrate penetrates/diffuses into the coating film, it may interfere with the coating material itself. For example, penetrated/diffused sodium gamma-hydroxybutyrate may act as a strong base which reacts with pH sensitive coating polymers, such as Eudragit L30-D55 for instance, weakening the coating layer and lowering the coating efficiency.

[0006] Further, the absorption of sodium gamma-hydroxybutyrate seems to be capacity-limited (Palatini et al, Eur. J. Clin Pharmacol. (1993) 45:353-356), but it has been unclear whether the absorption of this drug is region-specific, which would affect the oral delivery of GHB.

[0007] Therefore, a need exists in the art for a more convenient dosing regimen, an effective dosage form of controlled release of gamma-hydroxybutyric acid salts and an efficient way to deliver gamma-hydroxybutyric acid salts to an animal in the gastrointestinal tract. The current invention satisfies these needs.

SUMMARY OF THE INVENTION

[0008] It is an object of the present invention to provide a convenient and effective dosage form of GHB, whereby the number of dosages can be reduced.

[0009] It is another object of the present invention to provide compositions of GHB that have a reduced likelihood of drug migration from the dosage form.

[0010] The present invention takes into account the surprising discovery by the present inventors that the oral absorption of sodium gamma-hydroxybutyrate is region specific in animals, and that the absorption is higher in the upper GI tract than in the lower GI tract.

[0011] The present invention is also directed to methods and compositions for the targeting of the upper GI tract of an animal for improved absorption of sodium gamma-hydroxybutyrate.

[0012] The current invention provides methods and compositions for convenient administration of multiple doses of one or more gamma-hydroxybutyric acid salts to an animal. It provides a convenient once nightly or once daily dose regimen for the oral delivery of one or more gamma-hydroxybutyric acid salts to an animal. With the compositions of the present invention, a patient does not need to wake up at night to take a second dose then go back to sleep.

[0013] The current invention also provides methods and compositions for the effective delayed/controlled release of multiple (i.e., more than one) doses of one or more gamma-hydroxybutyric acid salts. The current invention provides methods and compositions to improve the gastro-stability of delayed/controlled release particulates (e.g. beads, granules, minitabs or pellets) containing gamma-hydroxybutyric acid salts.

[0014] The current invention further provides methods and compositions for the effective delivery of multiple doses of gamma-hydroxybutyric acid salts to one or more specific regions in the gastrointestinal tract of an animal. It provides methods and compositions for the targeting of the upper GI tract of an animal to improve the effectiveness of the absorption of gamma-hydroxybutyric acid salts from the delayed/controlled release particles.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] FIG. 1. Dissolution profile of a colon-targeting delayed release prototype with a neutralizing agent in the barrier coat.

[0016] FIG. 2. Dissolution profile of a colon-targeting delayed release prototype without a neutralizing agent in the barrier coat.

[0017] FIG. 3. Dissolution profile of a duodenum-targeting delayed release prototype without a neutralizing agent in the barrier coat.

[0018] FIG. 4. Dissolution profile of an immediate release core of the present invention.

[0019] FIG. 5. Dissolution profile of an Opadry AMB-coated immediate release core of the present invention.

[0020] FIG. 6. Dissolution profile of an ethylcellulose-coated immediate release core of the present invention.

[0021] FIG. 7. Dog pharmacokinetic profiles—demonstrating region of absorption.

DETAILED DESCRIPTION OF THE INVENTION

[0022] The current invention provides methods and compositions for convenient administration of multiple (i.e. more than one, “pulsed”) doses of one or more gamma-hydroxybutyric acid salts to an animal.

[0023] It also provides methods and compositions for the effective delayed/controlled release of multiple doses of one or more gamma-hydroxybutyric acid salts.

[0024] The current invention provides methods and compositions to improve the gastro-stability of the delayed/controlled release particles containing gamma-hydroxybutyric acid salts.

[0025] The current invention further provides methods and compositions for the effective delivery of multiple doses of gamma-hydroxybutyric acid salts to one or more specific regions in the gastrointestinal tract of an animal for effective absorption.

[0026] Specifically, at the essence of the present invention is a dosage form comprising one or more pH sensitive delayed/controlled release particles (e.g. beads, granules, minitabs or pellets), wherein each of the pH sensitive delayed/controlled release particles is composed of an immediate release core comprising one or more gamma-hydroxybutyric acid salts and one or more pharmaceutically acceptable excipients, one or more barrier coats surrounding such core (with or without a neutralizing agent), a pH sensitive enteric release coat around said barrier coat, and optionally an overcoat.

[0027] The dosage forms of the current invention comprise an immediate release component in the form of a solid, a semi-solid or a liquid, comprising one or more gamma-hydroxybutyric acid salts and optionally one or more pharmaceutically acceptable excipients, wherein the immediate release component is present together with (or separated contained from) one or more pH sensitive delayed/controlled release particles.

[0028] The dosage forms thus provide, which administered together or sequentially, multiple release pulses of gamma-hydroxybutyric acid salts targeting multiple regions in the gastrointestinal tract of an animal for improved absorption.

[0029] In one of the preferred embodiments, the composition comprises multiple delayed release pellets or beads (used interchangeably herein) and an immediate release component. In a most preferred embodiment, the dosage form comprises a liquid immediate release component, and two delayed/controlled release pellets/beads.

[0030] Each of the pH sensitive delayed/controlled release particles in the current invention is designed to release its contents at a specific region in the gastrointestinal tract of an animal. The one or more pH sensitive delayed/controlled release particles releases the contents at one or more corresponding regions in the gastrointestinal tract of an animal.

[0031] The immediate release component, in the form of a solid, a semi-solid or a liquid, of the current invention releases its contents immediately for absorption upon oral administration. Preferably, due to the high dosage of GHB, the immediate release component is a liquid.

[0032] Combining the immediate release component and one or more pH sensitive delayed/controlled release particles

of the current invention can constitute a complete once-nightly or once-daily dose. The term “combining” as used herein means supplying and consuming all components (1) simultaneously in the same presentation or dosage form, or (2) simultaneously in different presentations or dosage forms, or (3) sequentially in the same presentation or dosage forms, or (4) sequentially in different presentations or dosage forms.

[0033] For example, an immediate release component in the form of particles and one or more pH sensitive delayed/controlled release particles are supplied as pre-mixed doses, and are consumed simultaneously at the time of dosing. Or, an immediate release component in the form of particles and one or more pH sensitive delayed/controlled release particles are supplied in separated parts, and are consumed simultaneously at the time of dosing. Alternatively, an immediate release component in the form of a powder and one or more pH sensitive delayed/controlled release particles are supplied in separate parts, and are consumed simultaneously at the time of dosing. In another embodiment, an immediate release component in the form of a solution and one or more pH sensitive delayed/controlled release particles are supplied in separate parts, and are consumed simultaneously at the time of dosing. Or, an immediate release component in the form of a solution and one or more pH sensitive delayed/controlled release particles are supplied in separated parts, and are consumed sequentially at the time of dosing. Other permutations would be apparent in those skilled in the art.

[0034] In one embodiment of the present invention, the delayed/controlled release component(s) is/are administered prior to the immediate release component, which can be administered from several minutes to about a half hour or more later (for practical reason, likely no more than about an hour later because the patient will become somewhat sleepy from the first dose). Thus, in its most basic form, the present invention is directed to the delayed/controlled release component(s), which have utility as a separately administrable dosage form. These components can be supplied as a separate entity, and preferable used in conjunction with an immediate release dosage form as is currently marketed.

[0035] Multiple (i.e. more than one) delayed releases can be achieved by combining multiple pH sensitive delayed/controlled release particles targeting certain sites of the GI tract of an animal. For example, an immediate release component can be combined with two pH sensitive delayed/controlled release particles that are released at two different sites in the GI tract to provide an immediate release and two other delayed release pulses.

[0036] An immediate release component can be combined with one type of pH sensitive delayed/controlled release particles to provide two pulses of gamma-hydroxybutyric acid salts, which can conveniently replace the nightly multi-dose regimen of the existing commercial product. In this case, a patient does not need to wake up and take a second dose during the night, as described earlier.

[0037] Preferably, an immediate release component is combined with one or more pH sensitive delayed/controlled release particles to provide multiple releases in a period of time. Preferably, an immediate release component is combined with one or more pH sensitive delayed/controlled release particles targeted to the upper GI tract of an animal. The inventors discovered that the absorption of sodium gamma-hydroxybutyrate in the GI tract of an animal is site specific, and that the absorption of sodium gamma-hydroxybutyrate in the upper GI tract is higher than in the lower GI

tract. The aforementioned combination therefore provides an initial dose and one or more delayed doses of gamma-hydroxybutyric acid salts, thereby providing an effective and convenient dose regimen for treating a patient.

[0038] More preferably, an immediate release component is combined with a single type of pH sensitive delayed/controlled release particles targeted to the duodenum or the jejunum of an animal to provide a two-pulse regiment to treat a patient.

[0039] The dose ratio of the immediate release component to one or more pH sensitive delayed/controlled release particles is dictated by the type of therapy and readily determined by the clinician, using currently available dosages as a reference. For example, the immediate release dose can be equivalent of, higher than, or lower than, the one or more delayed release doses.

[0040] It is contemplated that the delayed release dose amount, which is used to replace the second nightly dose (currently as a solution) in the current treatment of narcolepsy patients, can be the same as the immediate release dose amount, although the bioavailability is lower further along the GI tract, or even at a reduced dose amount, since the patients do not need to wake up and take a separate second nightly dose then go back to sleep.

[0041] It is also contemplated that the immediate release component can be at a slightly higher than normal dose, and the delayed release dose can be at a normal dose or at a reduced dose.

[0042] It is also contemplated that an immediate release component can be combined with one or more pH sensitive delayed/controlled release particles that are at reduced doses. For example, an immediate release dose can be combined with 0.7 equivalent dose of a duodenum-targeting delayed release component and 0.2 equivalent dose of a colon-targeting delayed release component to give a broader time coverage.

[0043] The immediate release component and one or more pH sensitive delayed/controlled release particles of the current invention can be administered to an animal directly, or mixed/sprinkled with fluids, soft foods (i.e. yogurt, apple-sauce), or pharmaceutically acceptable carriers. For example, an immediate release component in the form of a solution can be mixed with juice and the pH sensitive delayed/controlled release particles can be combined with foods (such as yogurts) for administration. Or, an immediate release component in the form of particles and the pH sensitive delayed/controlled release particles can be sprinkled with drinkable yogurt for dosing.

[0044] The Immediate Release Component

[0045] The dosage forms of the current invention comprise an immediate release component in the form of a solid, a semi-solid or a liquid. It can be a particle, a bead, a pellet, a granulate, a powder, a tablet, a minitab, a capsule, a caplet, a lozenge, a hard shell or soft shell capsule, a sachet, a cachet, a solid dispersion, a solid solution, a suspension, an emulsion, a lotion, a solution, a liquid drop, an elixir, a syrup, a tincture, a liquid spray, an aerosol, a gel, an ointment, a cream, or the like.

[0046] The immediate release component can be present together with one or more of the pH sensitive delayed/controlled release particles described herein, or separated from the pH sensitive delayed/controlled release particles.

[0047] For example, the immediate release component can be in the form of particles that are pre-mixed with the pH

sensitive delayed/controlled release particles. Or the two components can be provided as separate parts, possibly in a kit, wherein both components can be consumed together, or separately in a sequential manner.

[0048] In another example, the immediate release component can be in the form of a powder that is pre-mixed with the pH sensitive delayed/controlled release particles prior to ingestion. In this embodiment, the immediate release component is a powder comprising up to 100% of one or more gamma-hydroxybutyric acid salts and optionally one or more pharmaceutically acceptable excipients. Such a powder can be taken as is, or preferably is stirred into a drink or food along with the delayed/controlled release beads/pellets/minitabs.

[0049] In another preferred embodiment, the immediate release component is an aqueous solution (like the current Xyrem® product) of one or more gamma-hydroxybutyric acid salts stabilized with antioxidants, stabilizers, preservatives and neutralizing agents.

[0050] In yet another example, which is preferred because of the very high dosage needed for this drug, the immediate release component can be in the form of a solution that is provided separately from the pH sensitive delayed/controlled release particles, possibly in a kit form. The immediate release component is an aqueous solution (like the current Xyrem® product) of one or more gamma-hydroxybutyric acid salts stabilized with antioxidants, stabilizers, preservatives and neutralizing agents. Preferably, the delayed release particles are mixed with the liquid and then ingested.

[0051] The immediate release component of the current invention comprises one or more gamma-hydroxybutyric acid salts and optionally one or more pharmaceutically acceptable excipients, wherein the gamma-hydroxybutyric acid salts are selected from gamma-hydroxybutyric acid sodium salt, gamma-hydroxybutyric acid potassium salt, gamma-hydroxybutyric acid tetraammonium salt, or any other pharmaceutically acceptable salt forms of gamma-hydroxybutyric acid.

[0052] The immediate release component comprises from about 20% to about 100% by weight of one or more gamma-hydroxybutyric acid salts and optionally one or more pharmaceutically acceptable excipients.

[0053] The pharmaceutically acceptable excipients in the immediate release component are those known in the art as suitable for use in solid, semi-solid or liquid dosage forms, including but not limited to, binders, lubricants, anti-adherents, glidants, granulating aids, fillers, disintegrants, antioxidants, stabilizers, preservatives, neutralizing agents, buffering agents, tonics, moisture absorbents, colorants, flavorants, sweeteners, sugars, and taste-masking agents, suspending agents, thickening agents, gelling agents, solvents, solubilizers, surfactants, absorption enhancers, emulsifying agents, and combinations thereof.

[0054] The total amount of these pharmaceutically acceptable excipients in the immediate release component is from about 0% to about 80% by weight.

[0055] Examples of these pharmaceutically acceptable excipients in the immediate release component of the current invention include, but are not limited to, binders/fillers: microcrystalline cellulose, silicified microcrystalline cellulose, polyvinylpyrrolidone, hydroxypropyl cellulose, starch, pregelatinized starch, starch paste, lactose, mannitol, sorbitol, xylitol, sucrose, calcium phosphate, calcium carbonate, ethylcellulose, methylcellulose, and Acacia; lubricants/anti-adherents/glidants/granulating aids: talc, sodium lauryl

fumarate, fumed silicon dioxide, colloidal silica, titanium dioxide, kaolin, magnesium stearate, calcium stearate, stearic acid, hydrogenated vegetable oils, and sodium lauryl sulfate; disintegrants: sodium starch glycolate, croscarmellose sodium, cross-linked polyvinylpyrrolidone, and alginic acid; antioxidants/stabilizers/preservatives: riboflavin, tocopherol, vitamin E TPGS, BHT, BHA, cysteine and derivatives, ascorbates, sorbates, benzoates, propionates, bicarbonates, thiosulfates, metabisulfites, EDTA, carrageen, gums and benzyl alcohol; neutralizing agents: acids such as malic acid, citric acid, tartaric acid, ascorbic acid, oleic acid, capric acid, caprylic acid, benzoic acid, polyacids, acidic ionic resins, and other acidic excipients; suspending agents/thickening agents/gelling agents: mineral oils, vegetable oils, silicon dioxide, various gums such as xanthan gum, locust bean gum, gum Arabic, alginates, Carbopols, polyvinyl alcohols, carrageenan, gelatin, starches; or mixtures thereof.

[0056] Preferably, if the immediate release component is a solid pellet, bead or minitab or the like, that component is also used as the immediate release core of the pH sensitive delayed/controlled release particles by coating them using materials and methods similar to the barrier coats or the overcoat as described herein.

[0057] Delayed/Controlled Release Particles

[0058] The immediate release core of the pH sensitive delayed/controlled release particles (i.e., beads, pellets, minitabs, granulate, etc.) of the current invention comprises from about 20% to about 99% of one or more gamma-hydroxybutyric acid salts by weight of the core and one or more pharmaceutically acceptable excipients, wherein the gamma-hydroxybutyric acid salts are selected from gamma-hydroxybutyric acid sodium salt, gamma-hydroxybutyric acid potassium salt, gamma-hydroxybutyric acid tetraammonium salt, or any other pharmaceutically acceptable salt forms of gamma-hydroxybutyric acid, or combinations thereof.

[0059] One or more pharmaceutically acceptable excipients in the immediate release core of the pH sensitive delayed/controlled release particles of the current invention are excipients known in the art as suitable for use in particulates, including but not limited to binders, lubricants, anti-adherents, glidants, granulating aids, fillers, disintegrants, antioxidants, stabilizers, preservatives, neutralizing agents, buffering agents, moisture absorbents, colorants, flavorants and task-masking agents.

[0060] The total amount of these pharmaceutically acceptable excipients in the immediate release core is from about 1% to about 80% by weight of the core.

[0061] Examples of these pharmaceutically acceptable excipients in the immediate release core of the current invention include, but are not limited to, binders/fillers: microcrystalline cellulose, silicified microcrystalline cellulose, polyvinylpyrrolidone, hydroxypropyl cellulose, starch, pregelatinized starch, starch paste, lactose, mannitol, sorbitol, xylitol, sucrose, calcium phosphate, calcium carbonate, ethylcellulose, methylcellulose, and Acacia; lubricants/anti-adherents/glidants/granulating aids: talc, sodium lauryl fumarate, fumed silicon dioxide, colloidal silica, titanium dioxide, kaolin, magnesium stearate, calcium stearate, stearic acid, hydrogenated vegetable oils, and sodium lauryl sulfate;

[0062] disintegrants: sodium starch glycolate, croscarmellose sodium, cross-linked polyvinylpyrrolidone and alginic acid; antioxidants/stabilizers/preservatives: riboflavin, tocopherol, vitamin E TPGS, BHT, BHA, cysteine and derivatives, ascorbates, sorbates, benzoates, propionates, bicarbon-

ates, thiosulfates, metabisulfites, EDTA, carrageen, gums and benzyl alcohol; neutralizing agents: acids such as malic acid, citric acid, tartaric acid, ascorbic acid, oleic acid, capric acid, caprylic acid, benzoic acid, polyacids, acidic ionic resins, and other acidic excipients; or mixtures thereof.

[0063] Preferably, the immediate release core of the current invention comprises one or more excipients selected from binders, lubricants, anti-adherents, glidants and neutralizing agents.

[0064] The lubricants/anti-adherents/glidants may be selected from talc, sodium lauryl fumarate, fumed silicon dioxide, magnesium stearate and stearic acid, for instance. Preferably, the lubricants/anti-adherents/glidants are selected from one or both of talc and magnesium stearate.

[0065] In a preferred embodiment, the amount of talc in the immediate release core of the current invention about 1% to about 25% by weight of the core. More preferably, this amount is from about 5% to about 15% by weight of the core.

[0066] If magnesium stearate is used in the core it is present in an amount of from about 0% to about 10% by weight of the core. More preferably, this amount is from about 0.1% to about 5% by weight of the core.

[0067] Preferably, the binders/fillers in the immediate release core are selected from microcrystalline cellulose, silicified microcrystalline cellulose, polyvinylpyrrolidone, and hydroxypropyl cellulose.

[0068] Preferably, the immediate release core comprises microcrystalline cellulose or silicified microcrystalline cellulose at about 1% to about 80% by weight of the core. More preferably, the immediate release core comprises microcrystalline cellulose or silicified microcrystalline cellulose at about 3% to about 40% by weight of the core.

[0069] Preferably, the immediate release core comprises a neutralizing agent. The uptake of gamma-hydroxybutyric acid salts may be affected by the environmental pH and the ionization state of the salts. Preferably, the immediate release core contains a neutralizing agent to modulate the ionization state of the salt for better absorption in the gastrointestinal tract.

[0070] The immediate release cores in the pH sensitive delayed/controlled release particles of the current invention are made by techniques and equipment known in the art, for example dry blending, milling, dry granulation, wet granulation, pelletization, direct pelletization, extrusion, melt-extrusion, spheronization, drug layering, compaction, compression. Solvents can be used to facilitate the preparation of the immediate release core. These solvents can be removed partially or completely during the preparation of the core. Suitable solvents include, but are not limited to, water, alcohols, ketones and combinations thereof. For example, water and/or alcohols can be used during wet granulation and spheronization, or during direct pelletization, or during drug layering, and the solvents can be removed thereafter.

[0071] Barrier Coat(s)

[0072] One or more barrier coats applied to the pH sensitive delayed/controlled release particles of the current invention provides a barrier, and a neutralization zone when a neutralizing agent is used, between the immediate release core and the enteric coat, and functions to prevent gamma-hydroxybutyric acid salts from entering into or interfering with the enteric coat. The barrier coats can optionally act also as a controlled release coat to control the rate of release of gamma-hydroxybutyric acid salts from the immediate release core.

[0073] The barrier coats in the current invention provide a barrier and optionally a neutralization zone between the immediate release core and the enteric coat to prevent the alkaline gamma-hydroxybutyric acid salts from migrating into and interfering with the pH sensitive enteric coat. If the highly water-soluble and strongly alkaline gamma-hydroxybutyric acid salts migrate into the enteric coat, they not only create channels in the enteric coat which act as pore formers, but also react with the functional groups of the coat materials and weaken the enteric coat. By controlling the thickness and/or the permeability of the barrier coats, the migration of gamma-hydroxybutyric acid salts can be minimized. Further, neutralizing agents, mainly acidifiers, can be used in the barrier coat to neutralize gamma-hydroxybutyric acid salts in the barrier layer thus preventing these alkaline salts from reacting with the enteric coat material.

[0074] Moreover, the barrier coats can optionally act as a controlled release coat to control the rate of release of gamma-hydroxybutyric acid salts from the immediate release core, allowing for site specific and controlled release of gamma-hydroxybutyric acid salts in the GI tract of an animal.

[0075] Suitable coating materials for the barrier coats in the current invention include, but are not limited to, cellulose polymers such as ethylcellulose, methylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, cellulose acetate, cellulose acetate phthalate, polyvinyl alcohol, or other water-based or solvent-based coating materials.

[0076] Suitable neutralizing agents in the barrier coats of the current invention include, but are not limited to, acids such as malic acid, citric acid, tartaric acid, ascorbic acid, oleic acid, capric acid, caprylic acid, benzoic acid, polyacids (a polymer with multiple carboxylic acid functional groups or side chains, e.g. polymethacrylic acid, or molecules with multiple acid functional groups, e.g. EDTA, ethylenediaminetetraacetic acid), acidic ionic resins, and other acidic excipients, and are used in amounts sufficient to neutralize any migrating gamma-hydroxybutyric acid salts. Preferably, the amount of neutralizing agent in the barrier coat is at about 0.01% to about 10% mol/mol of the gamma-hydroxybutyric acid salts in the core. More preferably, this amount is at about 1% to about 5% mol/mol of the salts.

[0077] The barrier coats in the current invention can further comprise other additives known in the art, such as pore formers, plasticizers, anti-adherents, glidants, and antifoam agents. Pore formers suitable for use in the barrier coats of the invention are organic or inorganic agents, and include materials that can be dissolved, extracted or leached from the coating in the environment of use. Examples of the pore formers include, but are not limited to, organic compounds such as saccharides including sucrose, glucose, fructose, mannitol, mannose, galactose, sorbitol, pullulan, dextran; polymers soluble in the environment of use such as water-soluble hydrophilic polymers, hydroxyalkylcelluloses, carboxyalkylcelluloses, hydroxypropylmethylcellulose, cellulose ethers, acrylic resins, polyvinylpyrrolidone, cross-linked polyvinylpyrrolidone, polyethylene oxide, Carbowaxes, Carbopol, and the like, diols, polyols, polyhydric alcohols, polyalkylene glycols, polyethylene glycols, polypropylene glycols, or block polymers thereof, polyglycols, poly(a-w) alkylene diols; inorganic compounds such as alkali metal salts, lithium carbonate, sodium chloride, sodium bromide, potassium chloride, potassium sulfate, potassium phosphate, sodium acetate, sodium citrate, suitable calcium salts, and the like. The amount of pore formers used in the barrier coats

varies depending on the functions of the barrier coats. For example, if the pH sensitive delayed/controlled release particles are intended for immediate release after entering the targeted site in the GI tract, high amounts of pore formers (e.g. as high as about 50% by weight of the barrier coat) can be used. If the pH sensitive delayed/controlled release particles are for controlled release after entering the targeted site in the GI tract, little or no pore formers are used (e.g. no more than about 25% by weight of the barrier coat).

[0078] The rate of release of gamma-hydroxybutyric acid salts in the pH sensitive delayed/controlled release particles can also be controlled by varying the thickness and/or types of the barrier coats, with or without the use of pore formers. For example, when ethylcellulose is used together with PVP K30 (5%) as the pore former, or when ethylcellulose is used with or without a water-insoluble plasticizer and without the use of any pore formers, or when ethylcellulose is used with a water-soluble plasticizer such as triethyl citrate, the barrier coat can be between about % to about 20% weight gain on the particles in order to obtain different controlled release profiles. Or, when Opadry AMB is used as the barrier coat, the barrier coat can be from about 2% to about 10% weight gain on the particles, in order to obtain an immediate release profile.

[0079] The barrier coats can also be multiple coats of different coating materials. For example, the barrier coats can have an Opadry AMB initial barrier coat, and an ethylcellulose secondary barrier coat surrounding the initial coat, and optionally an Opadry tertiary barrier coat surrounding the secondary coat.

[0080] The barrier coats can be water-based coatings, or organic solvent-based coatings. Preferably, the barrier coat is organic solvent-based coating such as an alcohol or alcohol-water or ketone based coating.

[0081] Furthermore, the barrier coats of the current invention can provide moisture protection for hygroscopic gamma-hydroxybutyric acid salts inside the barrier coats.

The pH Sensitive Enteric Coat

[0082] The pH sensitive enteric release coat of the current invention enables targeted delivery of the particles to a specific region in the GI tract. It also provides a time delay in the release of the gamma-hydroxybutyric acid salts from the pH sensitive delayed/controlled release particles of the current invention. Combinations of more than one of these pH sensitive delayed/controlled release particles in a dosage form will provide multiple doses of gamma-hydroxybutyric acid salts delivered to multiple sites in the GI tract with multiple delay time periods or pulses. When combined with any controlled release characteristics of the barrier coats, the compositions of the current invention provide a wide spectrum of combined site specific, delayed and controlled release profiles for oral delivery of gamma-hydroxybutyric acid salts to an animal.

[0083] Materials suitable for use in the pH sensitive enteric coat of the current invention are pH sensitive coating materials known in the art. The pH sensitive coating materials include, but are not limited to, methacrylate-based coating materials such as polymers of methacrylic acid and methacrylates (e.g. Eudragit L 100-55, Eudragit L 30-D55, Eudragit L 100, Eudragit S 100, Eudragit FS 30 D), cellulose-based coating materials such as cellulose acetate phthalate, carboxymethyl ethylcellulose, cellulose acetate trimellitate, hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate succinate, Shellac-based coating mate-

rials such as Emcoat 120N and Marcoat 125, and other enteric coating polymers such as polyvinyl acetate phthalate.

[0084] Other additives such as solvents, plasticizers (e.g. PEG, triethyl citrate, dibutyl sebate), anti-tack agents (e.g. talc), anti-foam agents, colorants, fillers/extenders, flavorants, surfactants (e.g. sodium lauryl sulfate), bases, buffers, and other suitable additives known in the art can also be used together with the pH sensitive enteric coating materials.

[0085] The coating can be organic solvent-based, or aqueous-based, or organic solvent/aqueous based.

[0086] Preferably, the pH sensitive delayed/controlled release particles are prepared by coating the barrier-coated immediate release core with an appropriate pH sensitive coating material targeting to a specific region in the GI tract of an animal. The weight gain of the pH sensitive enteric coating is from about 10% to about 70% of the final enteric-coated particle weight. Preferably, the weight gain of this coating is from about 20% to about 60% of the final enteric-coated particles. More preferably, the weight gain of this coating is about 30% to about 50% of the final enteric-coated particle weight.

[0087] The pH sensitive enteric release coat can target both the upper part and the lower part of the GI tract of an animal. The pH sensitive enteric coat releases/dissolves in one of the stomach, the duodenum, the jejunum, the ileum, or the colon of an animal. Suitable pH sensitive enteric coating materials targeting each of these regions in humans are known in the art, such as Eudragit E 100 or Eudragit E PO (stomach), Eudragit L 30 D-55 and Eudragit L 100-55 (duodenum), Eudragit L 12.5 and Eudragit L 100 (jejunum), Eudragit S 100 (ileum), and Eudragit FS 30 D (colon).

[0088] Preferably, the pH sensitive enteric release coat releases/dissolves in the upper GI tract of an animal, which will allow for better absorption of the drug. In a more pre-

coats targeting the desired regions of the GI tract of an animal. Combinations of various particles with different pH sensitive enteric coats thus provide multiple pulses of gamma-hydroxybutyric acid salts with various delayed release times.

Overcoats

[0091] Optionally, the immediate and/or delayed/controlled release solid dosage forms of the current invention can be coated with an overcoat. The overcoat can be a moisture barrier coat, a protection coat, a seal coat, a taste-masking coat, a flavor coat, a polish coat, a color coat, or any other cosmetic coats. Suitable coating materials for such an overcoat are known in the art, including, but are not limited to, cellulosic polymers such as hydroxypropylmethylcellulose, hydroxypropylcellulose, microcrystalline cellulose carrageenan, and ethylcellulose.

[0092] Other additives known in the art can also be used in the overcoat, such as solvents, plasticizers (e.g. PEG, triethyl citrate, dibutyl sebate), anti-tack agents (e.g. Talc), anti-foam agents, colorants, fillers/extenders, flavorants, and surfactants (e.g. sodium lauryl sulfate).

[0093] The invention now will be described with respect to the following examples; however, the scope of the present invention is not intended to be limited thereby.

EXAMPLES

Example 1

Compositions of the Immediate Release Core and/or the Immediate Release Component

[0094]

TABLE 1

Ingredients	PD0231-25	PD0231-24A	PD0231-24B	PD0231-24C	PD0231-19	PD0231-17A	PD0231-16	PD0231-15A	PD0231-12	PD0231-10A
Sodium gamma-hydroxybutyrate	80	80	84	80	80	80	80	90	40	80
Avicel PH101	10	15	10	10	20	10	15	10	58	—
Talc	9	5	5	9	—	—	—	—	—	—
Magnesium stearate	1	—	1	1	—	—	—	—	—	—
SMCC 50	—	—	—	—	—	—	—	—	—	15
Emcompress	—	—	—	—	—	10	5	—	—	—
HPMC E5	—	—	—	—	—	—	—	—	2	—
PVP K30	—	—	—	—	—	—	—	—	—	5
Lactose	—	—	—	—	—	—	—	—	—	—
Water	10*	11.3*	11.3*	5.5*	11.3*	15*	15*	12*	10.5*	9*
Ethanol	—	—	—	—	—	—	—	—	—	9*

ferred embodiment, the pH sensitive enteric coat releases/dissolves in the duodenum or the jejunum of an animal.

[0089] Optionally, acidifiers or bases can be added to the pH enteric coating materials to adjust the target release/dissolution pH or region in the GI tract of an animal. Further, acidifiers in the pH sensitive enteric coat can also counteract the alkaline effect from any migrating gamma-hydroxybutyric acid salts. Suitable acidifiers are organic acids or inorganic acids, acidic excipients, and the aforementioned neutralizing agents.

[0090] The delay in release of gamma-hydroxybutyric acid salts from the particles of the current invention can be achieved by selecting different pH sensitive enteric release

TABLE 2

Ingredients	PD0231-10B	PD0231-10C	PD0231-9B	PD0231-8A
Sodium gamma-hydroxybutyrate	70	65	80	83
Avicel PH101	—	—	—	—
Talc	—	—	—	—
Magnesium stearate	—	—	—	—
SMCC 50	28	35	17	17
Emcompress	—	—	—	—
HPMC E5	2	—	1	—
PVP K30	—	—	—	—

TABLE 2-continued

Ingredients	PD0231- 10B	PD0231- 10C	PD0231- 9B	PD0231- 8A
Lactose	—	—	2	—
Water	20*	18*	10.8	9.5
Ethanol	—	—	—	—

Number in parts by weight.

*Removed partially or completely during preparation

Example 2

Preparation of the Immediate Release Core

[0095] The immediate release core can be made by techniques or processes or equipments known in the art, including but are not limited to dry blending, milling, dry granulation, wet granulation, pelletization, direct pelletization, extrusion, melt-extrusion, spheronization, drug layering, as exemplified by the following preparations:

[0096] Dry powders of sodium gamma-hydroxybutyric acid, Avicel PH101, Talc and magnesium stearate were screened and mixed briefly, then charged into a high shear granulator. Water was added to the mixture during the granulation. The granulates were extruded through a screen with a desirable pore size then spheronized to yield pellets. The pellets were dried in an oven for a sufficient time, for example overnight, then screened.

Example 3

Moisture Protection Coat of the Immediate Release Core

[0097] Opadry AMB coating solution:

Opadry AMB	25 g
Deionized water	475 g

[0098] Uncoated pellets from Example 2 (600 g) were charged into a fluid bed coater. The Opadry AMB coating solution was sprayed onto the pellets with a product temperature at 39° C. until 3% weight gain was reached to yield an Opadry AMB-coated immediate release core.

Example 4

Barrier Coats of the Immediate Release Core

[0099] Ethylcellulose coating solution:

Ethylcellulose	73.9 g
PVP K90	1.72 g
Triethyl citrate	8.1 g
Isopropyl alcohol	1000 g
Ethyl alcohol	1000 g

[0100] Uncoated pellets from Example 2 (600 g) were charged into a fluid bed coater. The ethylcellulose coating solution was sprayed onto the pellets with a product temperature at 35° C. until 3%, 6% or 9.2% weight gain was reached to yield the EC-coated immediate release core.

[0101] For a slower release core, PVP K90 is used at lower levels or can be omitted.

Example 5

Neutralizing Agent-Containing Barrier Coats of the Immediate Release Core

[0102] Neutralizing agent-containing barrier coat solution:

Opadry White	30 g
Malic acid	30 g
Deionized water	540 g

[0103] The 3% Opadry AMB-coated pellets (Example 3) were further coated with the neutralizing agent-containing barrier coat solution to 10% weight gain. An additional coat of Opadry AMB was also applied to some of the resultant pellets.

Example 6

pH Sensitive Enteric Release Coatings

[0104]

Enteric coating solution 1 (Duodenum):	
Eudragit L 30 D-55	840 g
Triethyl citrate	12 g
Talc	24 g
Deionized water	324 g

Enteric coating solution 2 (Jejunum):	
Eudragit L100	390 g
Talc	24 g
Triethyl citrate	34 g
Isopropyl alcohol	2460 g
Acetone	377 g
Deionized water	390 g

Enteric coating solution 3 (Colon):	
Eudragit FS 30 D	540 g
Triethyl citrate	9 g
Talc	45 g
Deionized water	304 g

[0105] (a) The ethylcellulose (EC)-coated immediate release core from Example 4 was further coated with enteric coating solution (1) to 40%, 45% or 50% weight gains to yield the duodenum-targeting particles.

[0106] (b) The EC-coated immediate release core from Example 4 was further coated with enteric coating solution (2) to 40%, 45%, 50%, or 60% weight gain to yield the jejunum-targeting particles.

[0107] (c) The Opadry AMB coated immediate release core from Example 3 was further coated with enteric coating solution (3) to 40%, 45% or 50% weight gain to yield the colon-targeting particles.

[0108] (d) The core coated with neutralizing agent-containing barrier coats from Example 5 was coated with an additional coat with enteric coating solution (3) to 40%, 45% or 50% weight gain to yield the colon-targeting particles.

Example 7

Dissolution Profiles of Various Prototypes

Gastro-Stability Improvement by the Neutralizing Agent in the Barrier Coats

Delayed/Controlled Release Prototypes

[0109] Colon-targeting prototype having a neutralizing agent (malic acid) in the barrier coat (PD0231-26B-50) does not release any sodium gamma-hydroxybutyrate at pH 1.1 and pH 6.0 for up to 3 hours (FIG. 1), whereas the one without the neutralizing agent in the barrier coat (PD0231-31F-50) releases 3% at pH 1.1 in 2 hours and 12% at pH 6.0 in 1 hour (FIG. 2). The neutralizing agent in the barrier coats thus improves the gastro-stability of the prototypes significantly.

[0110] Immediate Release Prototypes

[0111] Immediate release core (PD031-44), Opadry AMB-coated immediate release core (PD0231-27A) and an EC—barrier coated immediate release core (PD0231-38E) all showed an immediate release profile at pH 1.1.

Example 8

Canine PK Study

[0112] Four prototypes were used in the cross-over dog PK study, including an immediate release core (as the immediate release component in the current invention, IR) (see Ex. 2), an Eudragit L 30 D-55 coated delayed release prototype (DR2) (see Ex. 6a), an Eudragit FS 30 D coated delayed release prototype (DR1-no acid) (see Ex. 6c) and an Eudragit FS 30 D coated delayed release prototype with malic acid as the neutralizing agent in the barrier coats (DR1-with acid) (see Ex. 6d). A total of 6 dogs (3 males and 3 females) were given two oral capsules of one of the prototypes containing 1 g of sodium gamma-hydroxybutyrate per capsule. There was a minimum of a 2-day washout between each dose. Blood was collected at the following time points: 0 (pre-dose), 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 10, 12, and 14 Hrs post dose (for a total of 312 samples). Plasma samples were analyzed using a verified LC/MS/MS method. Relative bioavailability was determined by comparing the AUC from the delayed release prototype group to the AUC of the immediate release prototype group.

[0113] The results show that the lower in the GI, the lower the bioavailability (BA); i.e., absorption is higher at upper GI. The immediate release component has the highest BA, so GHB may be absorbed better in its acid form. The BAs for the delayed release components with or without a neutralizer in the barrier coat do not vary much, so the neutralizer helps the coating—in turn the gastro-stability—but does not affect the BA. See Table 3 and FIG. 7.

TABLE 3

Mean GHB Concentrations (ug/mL)				
Time Point (Hr)	Period			
	1	2	3	4
	DR1-w/Acid	DR1-No Acid	IR	DR2
0	0.00	0.00	0.00	0.00
0.5	0.00	0.00	116.04	0.00
1	0.00	4.76	248.27	1.53
2	4.99	11.62	195.51	32.52
3	26.31	31.88	117.56	100.99
4	35.14	38.26	47.21	100.57
5	29.18	34.77	8.74	54.99
6	21.09	27.83	0.00	23.42
7	11.25	9.13	0.00	7.52
8	8.67	2.53	0.00	0.34
10	1.43	3.03	0.00	0.00
12	0.98	0.67	0.00	0.00
14	0.43	0.00	0.00	0.00
Tmax (Hr)	4.2	5.2	1.2	3.7
Cmax (ug/mL)	38.77	58.44	249.5	112.7
AUClast	134.3	162.6	601.0	318.4
Rel BA	22%	27%	100%	53%

1.-56. (canceled)

57. An oral pharmaceutical dosage form, comprising (a) an immediate release component comprising gamma-hydroxybutyric acid (GHB), and (b) a delayed release component comprising GHB.

58. The oral dosage form of claim 57, wherein said delayed release component comprises particles containing GHB in a core, which core is immediately surrounded by a barrier coat to control the migration of GHB from the core, which in turn is surrounded by an enteric release coat that releases GHB at a predetermined pH after ingestion.

59. The oral dosage form of claim 58, wherein said barrier coat contains a neutralizing agent or agents selected from the group consisting of malic acid, citric acid, tartaric acid, ascorbic acid, oleic acid, capric acid, caprylic acid, benzoic acid, a polyacid, and acidic ionic resins.

60. The oral dosage form of claim 59, wherein the neutralizing agent(s) are used in amounts sufficient to neutralize any migrating gamma-hydroxybutyric acid salts.

61. The oral dosage form of claim 60, wherein said neutralizing agent(s) are used in an amount of about 0.01% to about 10% mol/mol of the GHB.

62. The oral dosage form of claim 61, wherein the amount is from about 1% to about 5% mol/mol of the GHB.

63. The oral dosage form of claim 58, wherein the barrier coat is composed of materials selected from ethylcellulose, methylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, cellulose acetate, cellulose acetate phthalate, polyvinyl alcohol, or other water-based or solvent-based coating materials.

64. The oral dosage form of claim 58, wherein the core is surrounded by more than one barrier coat.

65. The oral dosage form of claim 64, wherein a first barrier coat comprises Opadry AMB, a second barrier coat comprises ethylcellulose, and a third barrier coat comprises Opadry.

66. The oral dosage form of claim 58, wherein the enteric release coat is a pH sensitive material, which releases GHB at a predetermined pH in the gastrointestinal tract.

67. The oral dosage form of claim 66, wherein the pH sensitive material comprises methacrylate-based coating materials, cellulose-based coating materials, shellac-based

coating materials such as Emcoat 120N and Marcoat 125, polyvinyl acetate phthalate, or combinations thereof.

68. The oral dosage form of claim **67**, wherein the methacrylate-based coating materials are polymers of methacrylic acid and methacrylates and are selected from Eudragit E 100, Eudragit E PO, Eudragit 1, 12.5, Eudragit L 100-55, Eudragit L 30-D55, Eudragit L 100, Eudragit S 100, and Eudragit FS 30D.

69. The oral dosage form of claim **67**, wherein the cellulose-based coating materials are selected from cellulose acetate phthalate, carboxymethyl ethylcellulose, cellulose acetate trimellitate, hydroxypropylmethylcellulose phthalate, and hydroxypropylmethylcellulose acetate succinate.

70. The oral dosage form of claim **58**, wherein the core further comprises one or more excipients selected from binders, lubricants, anti-adherents, glidants and neutralizing agents.

71. The oral dosage form of claim **66**, wherein the pH sensitive enteric release coat allows the release of GLIB in the upper gastrointestinal tract.

72. The oral dosage form of claim **71**, wherein the enteric release coat comprises Eudragit L 30 D-55, Eudragit L 100-55, Eudragit L 12.5, Eudragit L 100, or combinations thereof.

73. The oral dosage form of claim **66**, wherein the enteric release coat further comprises acidic materials that counteract the alkaline effects of GHB.

74. The oral dosage form of claim **57**, wherein the immediate release component is a solid, a semi-solid or a liquid.

75. The oral dosage form of claim **57**, comprising two delayed release components, each of which releases GHB in different regions of the gastrointestinal tract.

76. The oral dosage form of claim **75**, wherein one of the delayed release components releases GHB in the duodenum, and the other delayed release component releases GHB in the jejunum.

77. The oral dosage form of claim **76**, wherein the delayed release component that releases GHB in the duodenum comprises a GHB containing core surrounded by a barrier coat, which in turn is surrounded by an enteric coating comprising Eudragit L 30 D-55 or Eudragit L 100-55, and the delayed release component that releases GLIB in the jejunum comprises a GLIB containing core surrounded by a barrier coat, which in turn is surrounded by a coating comprising Eudragit L 12.5 or Eudragit L 100.

78. An oral pharmaceutical composition, comprising a delayed release component comprising gamma-hydroxybutyric acid (GHB).

79. A method for the treatment of a subject in need of the effects of GHB, comprising administering an effective amount of the oral dosage form of claim **57** to the subject.

80. The method of claim **79**, wherein the subject is a human.

81. The method of claim **79** for the treatment of a condition selected from narcolepsy, cataplexy, sleep paralysis, alcoholism, chronic schizophrenia, catatonic schizophrenia, atypical psychoses, chronic brain syndrome, neurosis, drug addiction and withdrawal, Parkinson's disease, neuropharmacological illnesses, hypertension, ischemia, circulatory collapse, radiation exposure, cancer, and myocardial infarction.

* * * * *