



(86) Date de dépôt PCT/PCT Filing Date: 2001/05/17
(87) Date publication PCT/PCT Publication Date: 2001/11/22
(45) Date de délivrance/Issue Date: 2013/01/15
(85) Entrée phase nationale/National Entry: 2002/11/15
(86) N° demande PCT/PCT Application No.: EP 2001/005702
(87) N° publication PCT/PCT Publication No.: 2001/087931
(30) Priorité/Priority: 2000/05/18 (US09/573,872)

(51) Cl.Int./Int.Cl. *C12N 15/31* (2006.01),
A01N 63/00 (2006.01), *C07K 14/195* (2006.01),
C12N 15/62 (2006.01), *C12N 15/82* (2006.01)

(72) Inventeurs/Inventors:
BOETS, ANNEMIE, BE;
ARNAUT, GRETA, BE;
DAMME, NICOLE, BE;
VAN RIE, JEROEN, BE

(73) Propriétaire/Owner:
BAYER CROPS SCIENCE NV, BE

(74) Agent: SMART & BIGGAR

(54) Titre : PROTEINES SECRETEES INSECTICIDES DE BREVIBACILLUS LATEROSPORUS
(54) Title: INSECTICIDAL SECRETED PROTEINS OF BREVIBACILLUS LATEROSPORUS

(57) **Abrégé/Abstract:**

Bacterial insecticidal proteins and equivalents thereof were isolated. These proteins and the DNA sequences encoding them are useful to make insecticidal compositions or transgenic plants to protect plants from damage by insects, particularly coleopteran insects.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 November 2001 (22.11.2001)

PCT

(10) International Publication Number
WO 01/87931 A3

(51) International Patent Classification⁷: **C12N 15/31**,
15/62, C07K 14/195, A01N 63/00

patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(21) International Application Number: PCT/EP01/05702

(22) International Filing Date: 17 May 2001 (17.05.2001)

Declarations under Rule 4.17:

(25) Filing Language: English

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB,
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP,
KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL,
SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE,
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR),
OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML,
MR, NE, SN, TD, TG)

(26) Publication Language: English

(30) Priority Data:
09/573,872 18 May 2000 (18.05.2000) US

— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations

(71) Applicant: **AVENTIS CROPSCIENCE N.V.** [BE/BE];
Jozef Plateaustraat 22, B-9000 Gent (BE).

(72) Inventors: **BOETS, Annemie**; Schonenberg 6, B-9620
Velzeke (BE). **ARNAUT, Greta**; Aalterseweg 40, B-9910
Knesselare (BE). **DAMME, Nicole**; Colijnstraat 6, B-9770
Kruishoutem (BE). **VAN RIE, Jeroen**; Gravin Johannalaan
10, B-9900 Eeklo (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

Published:

— with international search report

(88) Date of publication of the international search report:
10 May 2002

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: BACTERIAL INSECTICIDAL PROTEINS

(57) Abstract: Bacterial insecticidal proteins and equivalents thereof were isolated. These proteins and the DNA sequences encod-
ing them are useful to make insecticidal compositions or transgenic plants to protect plants from damage by insects, particularly
coleopteran insects.



WO 01/87931 A3

75749-27

INSECTICIDAL SECRETED PROTEINS OF BREVIBACILLUS LATEROSPORUS

Background of the Invention

5 1. Field of the Invention

The present invention relates to new insecticidal secreted proteins ("ISPs") isolated from a bacterial strain, preferably a Brevibacillus species strain, most preferably a Brevibacillus laterosporus species strain which are insecticidal when ingested in combination with an ISP-complimentary protein such as another ISP protein of this invention, and to DNA sequences
10 encoding such proteins. These proteins are useful to prevent or minimize insect damage, particularly of corn rootworms, to plants in a field.

The present invention also relates to plants, particularly corn plants, that are rendered
15 insecticidal, preferably to coleopteran insects, particularly to Diabrotica spp., Leptinotarsa spp. and Anthonomus species insects, by the expression of the ISP proteins of this invention in cells of said plants.

The present invention also relates to a method for controlling damage by Diabrotica spp.,
20 Leptinotarsa spp. or Anthonomus species insects, preferably Diabrotica spp. insect pests, particularly corn rootworms, by having the ISP proteins of the invention, particularly the proteins with the amino acid sequence of any one of SEQ ID No. 2, 4, 8 or 10, or insecticidally-effective fragments thereof, ingested by said insects.

2. Description of the Prior Art

Some of the most destructive pests are found among the Diabroticine beetles. In North America, the three important species of corn rootworms, Diabrotica virgifera (the Western corn rootworm), Diabrotica barberi (the Northern corn rootworm) and Diabrotica undecimpunctata howardi (the Southern corn rootworm) are considered to be the most expensive insect pests to control (Metcalf, 1986, Foreword in "Methods for the Study of Pest Diabrotica", pp. vii-xv, eds. Krysan, J.L. and Miller, T.A., Springer-Verlag, New York). Diabrotica virgifera and Diabrotica barberi are considered the most serious insect pests of corn in the major corn-producing states of the United States and Canada (Levine and Oloumi-Sadeghi, 1991, Annu. Rev. Entomol. 36, 229-55). The larvae feed on the roots and thus cause direct damage to corn growth and corn yields. Costs for soil insecticides to control larval damage to the root systems of corn and aerial sprays to reduce beetle damage to corn silks, when combined with crop losses, can approach one billion dollars annually (Metcalf, 1986, supra). Recently, in some US states it was discovered that the crop rotation program of planting soybeans after corn lost its effect as corn rootworms have adapted to this situation.

Bacterial strains and/or genes with toxicity to corn rootworm have been described in US patents 6,023,013; 6,015,553; 6,001,637; 5,906,818; and 5,645,831. Also, PCT publications WO 00/09697, WO 99/57282, WO 98/18932, WO 97/40162, and WO 00/26378 relate to toxins and genes obtainable from Bacillus or other bacterial spp., some of which are described to have toxicity to corn rootworm. WO 98/44137, WO 94/21795 and WO 96/10083 relate to pesticidal Bacillus strains, characterized by pesticidal proteins and auxiliary proteins produced during vegetative growth, some of which are described to have toxicity to corn rootworm. US patent 5,055,293 describes a method to control corn rootworms by inoculating soil with parasporal-inclusion forming species of Bacillus laterosporus.

75749-27

Orlova et al. (1998, Applied Environmental Microbiol. 64, 2723) showed insecticidal activity to mosquitoes associated with protein crystals in crystal-forming strains of *Bacillus laterosporus*.

Objects and Summary of the Invention

5 An object of the present invention is to provide novel proteins and DNA sequences encoding such proteins with significant toxicity to insects, preferably *Diabrotica* spp. insects, particularly corn rootworm.

Specific aspects of the invention include:

10 - a protein comprising the amino acid sequence of the smallest active toxin of the protein of SEQ ID NO: 2, wherein said smallest active toxin is: a. a protein comprising an amino acid sequence with at least 97% sequence identity to a fragment of SEQ ID NO: 2, said fragment consisting of the amino acid sequence from amino acid position 38 to amino acid position 871 of SEQ ID NO: 2; and b. insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination
15 with a protein fragment of SEQ ID NO: 4 from amino acid position 51 to amino acid position 457;

20 - a protein comprising the amino acid sequence of the smallest active toxin of the protein of SEQ ID NO: 4, wherein said smallest active toxin is: a. a protein comprising an amino acid sequence with at least 97% sequence identity to a fragment of SEQ ID NO: 4, said fragment consisting of the amino acid sequence from amino acid position 51 to amino acid position 457 of SEQ ID NO: 4; and b. insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with a protein fragment of SEQ ID NO: 2 from amino acid position 38 to amino acid position 871;

25 - an insecticidal protein comprising a sequence with at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 2 from amino acid

475749-27

position 38 to amino acid position 768 or from amino acid position 38 to amino acid position 871;

5 - a protein comprising the amino acid sequence of SEQ ID NO: 2 from amino acid position 38 to amino acid position 768 or from amino acid position 38 to amino acid position 871;

- an insecticidal protein comprising a sequence with at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 4 from amino acid position 51 to amino acid position 449 or from amino acid position 51 to amino acid position 457;

10 - a protein comprising the amino acid sequence of SEQ ID NO: 4 from amino acid position 51 to amino acid position 449 or from amino acid position 51 to amino acid position 457;

15 - a protein comprising an amino acid sequence with at least 97% sequence identity to the amino acid sequence of SEQ ID NO: 2, wherein said protein is insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with the protein of SEQ ID NO: 4 from amino acid position 51 to amino acid position 457; and

20 - a protein comprising an amino acid sequence with at least 97% sequence identity to the amino acid sequence of SEQ ID NO: 4, wherein said protein is insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with the protein of SEQ ID NO: 2 from amino acid position 38 to amino acid position 871.

25 In one embodiment of the invention, a protein is provided comprising the amino acid sequence of the smallest active toxin of the protein of SEQ ID NO: 2, wherein said smallest active toxin is:

a) a fragment of the protein of SEQ ID NO: 2, and

*75749-27

b) insecticidal to Diabrotica virgifera larvae when ingested by said larvae in combination with the protein of SEQ ID NO: 4 from amino acid position 51 to 457. Also provided is a protein comprising the amino acid sequence of the smallest active toxin of the protein of SEQ ID NO: 4, wherein said smallest active
5 toxin is:

a) a fragment of the protein of SEQ ID NO: 4, and

b) insecticidal to Diabrotica virgifera larvae when ingested by said insect in combination with the protein of SEQ ID NO: 2 from amino acid position 38 to 871.

Particularly preferred is a protein characterized by an amino acid
10 sequence comprising the sequence of SEQ ID NO: 2 from amino acid position 38 to amino acid position 768 or 781, preferably the protein characterized by the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 10; and a protein characterized by an amino acid sequence comprising the sequence of SEQ ID NO: 4 from amino acid position 51 to amino acid position 449 or 457, preferably a protein characterized by
15 the amino acid sequence of SEQ ID NO: 4 or SEQ ID NO: 8.

A further object of the invention is a protein comprising the amino acid sequence of the protease-digestion fragment of the protein encoded by the isp1A DNA deposited at the BCCM-LMBP under accession number LMBP 4009, which protease-digestion fragment is insecticidal to Diabrotica virgifera upon combined application with the protein of SEQ ID. No. 4 from amino acid position 51 to amino acid position 457; and a protein comprising the amino acid sequence of the protease-digestion fragment of the protein encoded by the isp2A DNA deposited at the BCCM-LMBP under accession number LMBP 4009, which protease-digestion fragment is insecticidal to Diabrotica virgifera upon combined application with the protein of SEQ ID No. 2 from amino acid position 38 to amino acid position 871; particularly wherein said protease-digestion fragment is obtainable by treatment with coleopteran gut juice.

Also provided in accordance with this invention is a DNA sequence encoding the above proteins, particularly a DNA comprising an artificial DNA sequence having a different codon usage compared to the naturally occurring DNA sequence but encoding the same protein sequence, preferably contained in a chimeric gene operably linked to a plant-expressible promoter region (i.e., a promoter region which is suitable for expression in plant cells, this can be from bacterial, viral or plant origin or can be artificially made); particularly a promoter region which is preferentially active in root tissue.

In one embodiment of the invention, the promoter in said chimeric gene comprises the DNA sequence of SEQ ID No. 5 or 6 or a DNA hybridizing thereto under stringent hybridization conditions.

In a further embodiment of this invention, the chimeric gene further comprises a signal peptide for secretion from the cell or for targeting to a cellular organelle, particularly a chloroplast transit peptide.

Also provided is a plant cell, a plant or a seed, comprising any of these chimeric genes integrated in their cells, particularly a combination of the chimeric gene encoding the ISP1A, or an insecticidally effective fragment thereof, and the chimeric gene encoding the ISP2A protein,
5 or an insecticidally effective fragment thereof; particularly a corn cell, plant or seed.

In another embodiment of this invention, a micro-organism transformed to contain any of the above DNA sequences is provided.

10 Also provided is a process for controlling insects, particularly a process for rendering a plant resistant to coleopteran insects, comprising expressing any of the ISP proteins of this invention in cells of a plant, and regenerating transformed plants from said cells which are resistant to insects. In such process, the insect is preferably selected from the group consisting of:
rootworms, weevils, potato beetles, Diabrotica species, Anthonomus spp., Leptinotarsa spp.,
15 Agelastica alni, Hypera postica, Hypera brunneipennis, Haltica tombacina, Anthonomus grandis, Tenebrio molitor, Triboleum castaneum, Dicladispa armigera, Trichispa serica, Oulema oryzae, Colaspis brunnea, Lissorhoptrus oryzophilus, Phyllotreta cruciferae, Phyllotreta striolata, Psylliodes punctulata, Entomoscelis americana, Meligethes aeneus, Ceutorynchus sp., Psylliodes chrysocephala, Phyllotreta undulata, Leptinotarsa decemlineata, Diabrotica
20 undecimpunctata undecimpunctata, Diabrotica undecimpunctata howardi, Diabrotica barberi, and Diabrotica virgifera.

Yet another object of the present invention is to provide a method for rendering plants insecticidal against Coleoptera and a method for controlling Coleoptera, comprising planting,
25 sowing or growing in a field plants transformed with DNA sequences encoding the ISP proteins of the invention, particularly corn plants. In an embodiment of this invention, the ISP proteins

are combined with other insecticidal proteins or protein combinations, preferably corn rootworm-toxic proteins.

Also provided in accordance with this invention are ISP1A or ISP2A equivalents, preferably from a Brevibacillus laterosporus strain, particularly from a Brevibacillus laterosporus strain not forming crystalline inclusions. Such equivalents preferably have molecular weights of about 95 to about 100 kD for ISP1A equivalents, and about 45 to about 50 kD for ISP2A equivalents, as determined in standard 8-10 % SDS-PAGE gel electrophoresis using appropriate molecular weight markers.

Detailed Description of the Preferred Embodiments of the Invention

In accordance with this invention, new bacterial toxins and DNA sequences encoding them have been isolated and characterized. The new proteins were designated ISP1A and ISP2A, and the DNA sequences encoding them isp1A and isp2A.

In accordance with this invention "ISP1A protein" refers to any protein comprising the smallest protein fragment of the amino acid sequence of SEQ ID No. 2 which retains insecticidal activity, particularly to coleopteran insects, more particularly to corn rootworm, cotton boll weevil and Colorado potato beetle, preferably to Diabrotica spp., especially to Southern, Western and Northern corn rootworm, particularly to Diabrotica virgifera, upon combined ingestion by an insect with a suitable ISP-complimentary protein, particularly the mature ISP2A protein. This includes any protein with an amino acid sequence comprising the amino acid sequence from the amino acid at position 32 or 38, preferably 38, to an amino acid at a position from amino acid position 768 to amino acid position 871 in SEQ ID No. 2, particularly any protein with an amino acid sequence comprising at least the amino

acid sequence of SEQ ID No. 2 from amino acid position 38 to amino acid position 768. This also includes the protein obtained from the amino acid sequence of SEQ ID No. 2 by cleaving off part or all of the N-terminal signal peptide sequence or the protein wherein the signal peptide sequence has been replaced by another, e.g. a plant, targeting peptide, a methionine amino acid or a methionine-alanine dipeptide. Specifically, this includes the protein comprising an N-terminal prokaryotic or eucaryotic, e.g. bacterial or plant, signal peptide for secretion or targeting. This also includes hybrids or chimeric proteins comprising the above described smallest toxic protein fragment, e.g., a hybrid between an ISP1A and ISP2A protein of this invention. Further, included in the designation "ISP1A", as used herein, are also protease-resistant fragments of the ISP1A protein retaining insecticidal activity obtainable by treatment with insect gut juice, preferably coleopteran gut juice, particularly coleopteran gut proteases, preferably corn rootworm proteases, e.g., cysteine proteinases, serine proteinases, trypsin, chymotrypsin or trypsin-like proteases. Particularly, the coleopteran is selected from the group consisting of: corn rootworm, cotton boll weevil and Colorado potato beetle, Diabrotica spp., Diabrotica virgifera, Diabrotica barberi, Diabrotica undecimpunctata, Leptinotarsa decemlineata, and Anthonomus grandis. In a preferred embodiment of this invention, an ISP1A protein according to this invention is not insecticidal when ingested in isolation without providing simultaneously or sequentially, an ISP complimentary protein such as an ISP2A protein.

In accordance with this invention "ISP2A protein" refers to any protein comprising the smallest protein fragment of the amino acid sequence of SEQ ID No. 4 which retains insecticidal activity, particularly to coleopteran insects, more particularly to corn rootworm, cotton boll weevil and Colorado potato beetle, preferably to Diabrotica spp., especially to Diabrotica virgifera, Diabrotica barberi, Diabrotica undecimpunctata, upon combined ingestion by an insect with a suitable ISP-complimentary protein, particularly the mature

ISP1A protein. This includes any protein with an amino acid sequence comprising the amino acid sequence from the amino acid at position 43 or 51, preferably 51, to the amino acid at a position from amino acid position 449 to position 457 in SEQ ID No. 4, particularly any protein with an amino acid sequence comprising at least the amino acid sequence of SEQ ID

5 No. 4 from amino acid position 51 to position 449. This also includes the protein obtained from the amino acid sequence of SEQ ID No. 4 by cleaving off part or all of the N-terminal signal peptide sequence of the protein, and the protein wherein the signal peptide has been replaced by another, e.g., a plant, targeting peptide, a methionine amino acid or a methionine-alanine dipeptide. Specifically, this includes the protein comprising an N-

10 terminal prokaryotic or eucaryotic, e.g. bacterial or plant, signal peptide for secretion or targeting. This also includes hybrids or chimeric proteins comprising the above described smallest toxic protein fragment, e.g., a hybrid between an ISP1A and ISP2A protein of this invention. Further, included in the designation "ISP2A", as used herein, are also protease-resistant fragments of the ISP2A protein retaining insecticidal activity obtainable by

15 treatment with insect gut juice, preferably coleopteran gut juice, particularly coleopteran gut proteases, e.g., cysteine proteinases, serine proteinases, trypsin, chymotrypsin or trypsin-like proteases. In a preferred embodiment of this invention, an ISP2A protein according to this invention is not insecticidal when ingested in isolation without providing simultaneously or sequentially, an ISP complimentary protein such as an ISP1A protein.

20 An "ISP-complimentary protein", as used herein, refers to a protein, including but not limited to the mature ISP1A or ISP2A protein, which in combination with one of the ISP proteins of this invention, is insecticidal upon ingestion by an insect, particularly a coleopteran insect, preferably a corn rootworm, a cotton boll weevil or Colorado potato beetle, more particularly

25 Diabrotica virgifera, Diabrotica barberi, Diabrotica undecimpunctata or Anthonomus grandis. Particularly, also VIP proteins and active fragments thereof, particularly the mature VIP

proteins with their signal sequences cleaved off, as described in WO 98/44137, WO 94/21795 and WO 96/10083 are ISP-complimentary proteins in accordance with this invention. An ISP-complimentary protein to the ISP1A protein is ideally the ISP2A protein or the VIP2Aa or VIP2Ab protein or active fragments thereof (such as the mature proteins with the signal sequences removed) as described in US patent 5,990,383, or any bacterial secreted protein which has a sequence identity of at least 50 %, preferably at least 75 %, particularly at least 85 %, to any one of the ISP2 or VIP2 proteins, and which is insecticidal when ingested by an insect, preferably a coleopteran insect, particularly a corn rootworm, in combination with the mature ISP1A protein. An ISP-complimentary protein to the ISP2A protein is ideally the mature ISP1A protein, the VIP1Aa or VIP1Ab protein or active fragments thereof (such as the mature protein with the signal peptide removed) as described in US patent 5,990,383, or any bacterial secreted protein which has a sequence identity of at least 50 %, preferably at least 75 %, particularly at least 85 %, to any one of the ISP1 or VIP1 proteins, and which is insecticidal when ingested by an insect, preferably a coleopteran insect, particularly a corn rootworm, in combination with the mature ISP2A protein. For the avoidance of doubt, an ISP-complimentary protein and an ISP protein are always different proteins.

In a preferred embodiment of this invention, the ISP proteins of this invention, or their equivalents, when used in isolation, i.e., without any of the complementary ISP proteins present, do not result in any significant insecticidal activity, preferably to corn rootworm larvae, particularly to Diabrotica virgifera, when tested in a surface contamination assay on standard insect diet, and this at a concentration wherein the proteins (each applied in that concentration) in combination result in 100% mortality, preferably to corn rootworm larvae, particularly to Diabrotica virgifera. Particularly, the ISP1 proteins of this invention give no significant mortality (i.e., no difference with the controls using a buffer solution alone) to

Diabrotica virgifera larvae when only an ISP1 protein is applied at a concentration of 70ng/cm² in a surface contamination assay using standard corn rootworm diet, and the ISP2 proteins of this invention give no significant mortality (i.e., no difference with the controls using a buffer solution alone) to Diabrotica virgifera larvae when only an ISP2 protein is applied at a concentration of 36 ng/cm² in a surface contamination assay using standard corn rootworm diet, while ISP1 and ISP2 proteins applied together in these concentrations in the same type of assay give 100 % mortality to these larvae.

As used herein, "ISP1A equivalent" or "ISP2A equivalent" refers to a protein with the same or substantially the same toxicity to a target insect as the ISP1A or ISP2A protein, respectively, when applied to such target insect, preferably when ingested by such insect, in a binary combination with an ISP-complimentary protein, and with substantially the same amino acid sequence as the ISP1A or ISP2A protein, respectively. Also included in the definition of ISP1A or ISP2A equivalents are bacterial proteins of respectively about 45 to about 50 kD and about 95 to about 100 kD molecular weight as determined by standard 8-10 % SDS-PAGE gel electrophoresis, preferably from a Brevibacillus laterosporus strain, particularly from a Brevibacillus laterosporus strain not forming crystalline inclusions, which proteins in combination but not in isolation, have significant insecticidal activity to corn rootworm larvae.

The use of the terms "in combination", when referring to the application of an ISP protein (or its equivalent) and an ISP-complimentary protein to a target insect to get an insecticidal effect, includes the simultaneous application (i.e., in the same feed, cells or tissue and applied or ingested by an insect at the same moment) and the separate, sequential application (i.e. one is provided after the other but not applied or ingested at the same time) of an ISP protein (or its equivalent) and an ISP-complimentary protein of this invention, as long as these proteins are found together in the insect gut at one moment in time. The

ISP1A protein of this invention could thus be expressed in a certain type of cells or a certain zone in the roots of a plant, while the ISP2A protein of this invention could be expressed in another kind of cells or another zone in the roots of the same plant, so that the proteins will only interact once root material is ingested. Also included herein is the expression of an ISP
5 protein or its equivalent in roots of a plant, particularly a corn plant, and the expression of an ISP-complimentary protein in root-associated bacteria such as rhizobacteria strains (or vice versa).

"The same toxicity to a target insect", with respect to an ISP protein and an ISP equivalent
10 protein as used herein, means that the mean mortality of the ISP protein, in the presence of a suitable ISP-complimentary protein, is not significantly different from the mean mortality of the ISP equivalent, also in the presence of the same suitable ISP-complimentary protein. Particularly, this refers to the situation wherein the 95% fiducial limits of the LC50 of the ISP protein (when tested in the presence of a suitable ISP-complimentary protein) overlap with
15 the 95% fiducial limits of the LC50 of the ISP equivalent (when tested in the presence of a suitable ISP-complimentary protein).

"Substantially the same toxicity to a target insect", as used herein with respect to an ISP protein and an ISP equivalent protein, refers to levels of mean mortality of such proteins to a
20 target insect which are significantly different between the ISP and the ISP equivalent protein, but which are still within a range of insecticidal activity which is useful to control or kill the relevant target insect, preferably when such protein is expressed in a plant. In a preferred embodiment of this invention, proteins have substantially the same toxicity to a target insect when their LC50 values for such a target insect in the same (replicated) in vitro assay
25 conducted under the same assay conditions differ from each other by a factor 2 to 100, preferably 2 to 50, particularly 2 to 20, most preferably 2 to 10.

Also, functionally analogous amino acids can be used to replace certain amino acids in an ISP1A or ISP2A protein to obtain ISP1A or ISP2A equivalents. For example, one or more amino acids within the sequence can be substituted by other amino acids of a similar polarity which act as a functional equivalent, resulting in a silent alteration with respect to functionality of the protein. Substitutes for an amino acid within the sequence can be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Proteins wherein conservative amino acids replacements are made using amino acids of the above indicated same class which retain substantially the same insecticidal activity to a target insect compared to the original protein, are included herein as equivalents of the ISP proteins of this invention.

A protein with "substantially the same amino acid sequence" to an ISP1A protein, as used herein, refers to a protein with at least 90 %, particularly at least 95 %, preferably at least 97 % sequence identity with the ISP1A protein, wherein the percentage sequence identity is determined by using the blosum62 scoring matrix in the GAP program of the Wisconsin package of GCG (Madison, Wisconsin, USA) version 10.0 (GCG defaults used). "Sequence identity", as used throughout this application, when related to proteins, refers to the percentage of identical amino acids using this specified analysis. The "sequence identity", as used herein, when related to DNA sequences, is determined by using the nwsgapdna scoring matrix in the GAP program of the Wisconsin package of GCG (Madison, Wisconsin, USA) version 10.0 (GCG defaults used).

"ISP protein" or "ISP protein of this invention", as used herein, refers to any one of the new proteins of this invention and identified herein as ISP1A or ISP2A protein or their equivalents. An "ISP protoxin" refers to the full length ISP1A or ISP2A protein as it is encoded by the naturally-occurring bacterial DNA sequence, including the signal peptide. An "ISP toxin" refers to an insecticidal fragment thereof, particularly the smallest toxic fragment thereof or the mature protein with the signal peptide removed. A "mature" ISP, as used herein, refers to the ISP protein of this invention as secreted by its native bacterial host cell which is insecticidally active in combination with an ISP-complimentary protein (without the N-terminal bacterial signal peptide sequence). A mature ISP protein can have a complete native C-terminal or can have a C-terminal truncation.

Also included in this invention as an ISP protein is a first protein with an apparent molecular weight of its mature form of about 95 to about 100 kD, particularly about 100 kD, which is secreted by a bacterium, preferably a bacterium which is not B. thuringiensis or B. cereus, particularly a bacterium which is Brevibacillus laterosporus, or insecticidally effective fragments of said first protein, characterized by a significant insecticidal activity when combined with a second protein with an apparent molecular weight of its mature form of about 45 to about 50 kD, preferably about 45 kD, which is secreted by a bacterium, preferably a bacterium which is not B. thuringiensis or B. cereus, particularly a bacterium which is Brevibacillus laterosporus, or insecticidally effective fragments of said second protein, wherein the combination of said first and said second protein, or their insecticidal fragments, is significantly insecticidal to larvae of the Colorado potato beetle, the Western corn rootworm, the Southern corn rootworm, and the Northern corn rootworm, when ingested by said insects, and wherein said protein combination is not significantly insecticidal to Ostrinia nubilalis, Spodoptera frugiperda, Heliothis virescens, Helicoverpa zea and Sesamia nonagroides, when ingested by said insects, and wherein said

first protein alone has no significant insecticidal activity when ingested by an insect. Also included herein is the second protein as described above which is insecticidal when ingested by an insect in combination with the above described first protein.

5 "Apparent molecular weight", as used herein, is the molecular weight as evidenced by SDS-PAGE analysis upon comparison with molecular weight standards. As is appreciated by the skilled person, this molecular weight is an approximate determination and can have a range of variation of about 10-15 % with respect to the actual molecular weight as determined based on the amino acid sequence.

10

As used herein, the terms "isp1A DNA" or "isp2A DNA" refer to any DNA sequence encoding the ISP1A or ISP2A protein, respectively, as defined above. This includes naturally occurring, artificial or synthetic DNA sequences encoding the newly isolated proteins or their fragments as defined above, particularly DNA sequences with a modified codon usage adapted to more closely match the codon usage of a plant. Examples of artificial DNA sequences encoding an ISP1A and ISP2A protein are shown in SEQ ID Nos. 9 and 7, respectively (the ISP proteins encoded by these DNA sequences are defined herein as ISP1A-1 and ISP2A-1). Also included herein are DNA sequences encoding insecticidal proteins which are similar enough to the DNA sequence of SEQ ID No. 1, 3, 7 or 9 so that they can (i.e., have the ability to) hybridize to these DNA sequences under stringent hybridization conditions. "Stringent hybridization conditions", as used herein, refers particularly to the following conditions wherein hybridization is still obtained: immobilizing a first DNA sequence on filters, and prehybridizing the filters for either 1 to 2 hours in 50 % formamide, 5 % SSPE, 2x Denhardt's reagent and 0.1 % SDS at 42 ° C or 1 to 2 hours in 6x SSC, 2xDenhardt's reagent and 0.1 % SDS at 68 °C. The denatured radiolabeled second DNA is then added directly to the prehybridization fluid and incubation is carried out for 16 to

15

20

25

24 hours at one of the above selected temperatures. After incubation, the filters are then washed for 20 minutes at room temperature in 1x SSC, 0.1 % SDS, followed by three washes of 20 minutes each at 68 °C in 0.2x SSC and 0.1 % SDS. An autoradiograph is established by exposing the filters for 24 to 48 hours to X-ray film (Kodak XAR-2 or equivalent) at -70 °C with an intensifying screen. Of course, equivalent conditions and parameters can be used in this process while still retaining the desired stringent hybridization conditions. As used herein, stringent hybridization preferably occurs between DNA sequences with at least 90 to 95 %, preferably at least 97 %, particularly 99%, sequence identity.

Also included in the definition of "isp1 DNA" are all DNA sequences encoding a protein with a sequence identity of at least 90 %, preferably at least 95 %, particularly at least 97 %, most preferably at least 99 % with the protein of SEQ ID. No. 2 and which has substantially the same, preferably the same, insecticidal activity of the protein of SEQ ID No. 2, wherein said protein sequence identity is determined by using the blosum62 scoring matrix in the GAP program of the Wisconsin package of GCG (Madison, Wisconsin, USA) version 10.0 (GCG defaults used). Included in the definition of "isp2 DNA" are all DNA sequences encoding a protein with a sequence identity of at least 90 %, preferably at least 95 %, particularly at least 97 %, most preferably at least 99 %, with the protein of SEQ ID. No. 4 and which has substantially the same, preferably the same, insecticidal activity of the protein of SEQ ID No. 4, wherein said protein sequence identity is determined by using the blosum62 scoring matrix in the GAP program of the Wisconsin package of GCG (Madison, Wisconsin, USA) version 10.0 (GCG defaults used).

An "isp gene" or "isp DNA", as used herein, is a DNA sequence encoding an ISP protein in accordance with this invention, referring to any one of the isp1A or isp2A DNA sequences defined above.

- 5 An "isp DNA equivalent" or an "isp gene equivalent", as used herein, is a DNA encoding an ISP equivalent protein as defined above.

The terms "DNA/protein comprising the sequence X" and "DNA/protein with the sequence comprising sequence X", as used herein, refer to a DNA or protein including or containing at
10 least the sequence X in their nucleotide or amino acid sequence, so that other nucleotide or amino acid sequences can be included at the 5' (or N-terminal) and/or 3' (or C-terminal) end, e.g., an N-terminal transit or signal peptide. The term "comprising", as used herein, is open-ended language in the meaning of "including", meaning that other elements than those specifically recited can also be present. The term "consisting of", as used herein, is closed-
15 ended language, i.e., only those elements specifically recited are present. The term "DNA encoding a protein comprising sequence X", as used herein, refers to a DNA comprising a coding sequence which after transcription and translation results in a protein containing at least amino acid sequence X. A DNA encoding a protein need not be a naturally-occurring DNA, and can be a semi-synthetic, fully synthetic or artificial DNA and can include introns
20 and 5' and/or 3' flanking regions. The term "nucleotide sequence", as used herein, refers to the sequence of a DNA or RNA molecule, which can be in single- or double-stranded form.

The term "gene", as used herein refers to a DNA coding region flanked by 5' and/or 3' regulatory sequences allowing an RNA to be transcribed which can be translated to a
25 protein, typically comprising at least a promoter region. A "chimeric gene", when referring to an isp DNA of this invention, refers to an isp DNA sequence having 5' and/or 3' regulatory

sequences different from the naturally-occurring bacterial 5' and/or 3' regulatory sequences which drive the expression of the ISP protein in its native host cell.

"Insecticidal activity" or "insecticidally effective", when referring to an ISP protein of the invention or its equivalents, as used herein, means the capacity of an ISP protein to kill insects above the levels found in control treatment under the same assay conditions, upon the ingestion of such protein by an insect, preferably a coleopteran insect, particularly a corn rootworm, especially Diabrotica virgifera, in combination with an ISP-complimentary protein, such as a second ISP protein. Preferably the second ISP protein is the other protein encoded by the same bacterial operon as the first ISP or an insecticidally-effective fragment or equivalent thereof, yielding optimal insect mortality upon ingestion of the combined proteins. "Insect-controlling amounts" of an ISP protein, as used herein, refers to an amount of ISP protein which is sufficient to limit damage on a plant by insects feeding on such plant to commercially acceptable levels, e.g. by killing the insects or by inhibiting the insect development or growth in such a manner that they provide less damage to a plant and plant yield is not significantly adversely affected, when such ISP protein is provided with an ISP-complimentary protein, such as another ISP protein, preferably the other protein encoded by the same operon as the first ISP, or an insecticidally effective fragment thereof. "Insecticidally-effective ISP fragment", as used herein, refers to a fragment of an ISP protein of this invention which retains insecticidal activity when provided in combination with an ISP-complimentary protein, such as another ISP protein, preferably the other ISP encoded by the same operon as the first ISP. In the above definitions related to insecticidal activity, the insect preferably is a larva in any of the larval stages. Throughout this application, reference can be made to an "isp DNA and insecticidally-effective fragments or equivalents thereof", and in that case the insecticidal activity obviously refers to the activity of the protein encoded by the DNA and not to the insecticidal activity of the DNA itself.

In accordance with this invention, the ISP proteins of this invention and their equivalents, particularly the mature ISP1A and ISP2A proteins, were found to have no significant insecticidal activity to lepidopteran insects selected from the group consisting of: Heliothis virescens, Helicoverpa zea, Manduca sexta, Helicoverpa armigera, Spodoptera littoralis,
5 Spodoptera frugiperda, Sesamia nonagroides, and Ostrinia nubilalis.

In accordance with this invention, target insects susceptible to the ISP proteins of the invention or their equivalents are contacted with these proteins in insect-controlling amounts, preferably insecticidal amounts, by expression in a transgenic plant of DNA sequences
10 encoding such ISP proteins. Said target insects will only be affected by the insecticidal proteins when they ingest plant tissue. Thus, in another object of the present invention a method is provided for rendering plants insecticidal against Coleoptera and a method for controlling Coleoptera, comprising planting, sowing or growing in a field plants transformed with
15 DNA sequences encoding the ISP proteins of the invention, particularly corn plants.

The signal peptide of the ISP proteins of the invention can be removed or modified according to procedures known in the art, see, e.g., published PCT patent application WO 96/10083, or they can be replaced by another peptide such as a chloroplast transit peptide (e.g., Van Den
20 Broeck et al., 1985, Nature 313, 358, or preferably the modified chloroplast transit peptide of US patent 5, 510,471) causing transport of the protein to the chloroplasts, by a secretory signal peptide or a peptide targeting the protein to other plastids, mitochondria, the ER, or another organelle, or it can be replaced by a methionine amino acid or by a methionine-alanine dipeptide. Signal sequences for targeting to intracellular organelles or for secretion
25 outside the plant cell or to the cell wall are found in naturally targeted or secreted proteins, preferably those described by Klösgen et al. (1989, Mol. Gen. Genet. 217, 155-161), Klösgen

and Weil (1991, Mol. Gen. Genet. 225, 297-304), Neuhaus & Rogers (1998, Plant Mol. Biol. 38, 127-144), Bih et al. (1999, J. Biol. Chem. 274, 22884-22894), Morris et al. (1999, Biochem. Biophys. Res. Commun. 255, 328-333), Hesse et al. (1989, EMBO J. 8 2453-2461), Tavladoraki et al. (1998, FEBS Lett. 426, 62-66), Terashima et al. (1999, Appl. Microbiol. Biotechnol. 52, 516-523), Park et al. (1997, J. Biol. Chem. 272, 6876-6881), Shcherban et al. (1995, Proc. Natl. Acad. Sci USA 92, 9245-9249), all of which are incorporated herein by reference, particularly the signal peptide sequences from targeted or secreted proteins of corn. Although a DNA sequence encoding such a plant signal peptide can be inserted in the chimeric gene encoding the ISP1A and in the chimeric gene encoding the ISP2A protein for expression in plants, in one embodiment of this invention, a DNA sequence encoding such a signal peptide is only inserted in the chimeric ISP2A gene and the ISP1A chimeric gene lacks a signal peptide, or has a methionine amino acid or a methionine-alanine dipeptide instead of the signal peptide. In a preferred embodiment of this invention, the proteins are secreted from the roots as described by Gleba et al. (1999, Proc. Natl. Acad. Sci. USA 25, 5973-5977) and Borisjuk et al., 1999 (Nat. Biotechn. 17, 466-469), using a strong root-preferred promoter, particularly a strong root-preferred promoter from corn, and an N-terminal signal peptide for secretion from the root cells, preferably from corn, particularly the signal peptide of a secreted protein, preferably a corn secreted protein, selected from the group of: expansin, alpha-amylase, polyamine oxidase, cytokinin oxidase, endoxylanase.

Plants included in the scope of this invention are corn, cotton, soybean, peas, beans, lentils, potato, tomato, tobacco, lettuce, Brassica species plants, sugarcane, rice, oilseed rape, mustard, asparagus, wheat, barley, coffee, tea, vines, rubber plant, beet, turf grasses, sorghum, oats, rye, onions, carrots, leek, cucumber, squash, melon, sunflower, particularly any plant which is susceptible to damage by coleopteran insects, preferably corn rootworm, potato beetles or weevils, particularly insects of the Diabrotica or Leptinotarsa species, most

preferably any insect selected from the group consisting of: Diabrotica virgifera, Diabrotica barberi, Diabrotica undecimpunctata, Leptinotarsa decemlineata and Anthonomus grandis, most preferably corn plants.

- 5 "Corn", as used herein, refers to all plants of the species Zea mays, and any seeds, roots or grain, or other materials containing or directly produced from corn cells, of any variety of Zea mays, including but not limited to field corn, sweet corn, hybrid corn, white corn and dent corn, whether hybrid or inbred lines. Preferably, the corn plants used in this invention are suitable parent lines for producing hybrid corn, and most preferably they already carry an
- 10 endogenous or transgenic insect-resistance giving them protection from the major corn lepidopteran insect pests including but not limited to Ostrinia nubilalis.

The ISP1A and ISP2A proteins of this invention can be isolated in a conventional manner from the E. coli strain, deposited under the provisions of the Budapest Treaty on January 11,

15 2000 at the BCCM-LMBP (Belgian Coordinated Collections of Microorganisms - Laboratorium voor Moleculaire Biologie - Plasmidencollectie, University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium) under accession number LMBP 4009, or more preferably, they can be isolated from the supernatant of a Bacillus strain, preferably a crystal-minus Bacillus thuringiensis strain, transformed with a plasmid containing the

20 approximately 7 kb XbaI-EcoRI fragment of the plasmid pUCIB120/ISP as deposited under accession number LMBP 4009. The ISP proteins can be used to prepare specific monoclonal or polyclonal antibodies in a conventional manner (Höfte et al., 1988, Appl. Environm. Microbiol. 54, 2010). The ISP proteins can be treated with a protease such as cysteine proteases, to obtain the protease-resistant fragments of the ISP proteins.

The DNA sequences encoding the ISP proteins can be isolated in a conventional manner from the deposited strain or can be synthesized based on the encoded amino acid sequence.

5 DNA sequences encoding other ISP proteins in accordance with this invention can be identified by digesting total DNA from isolated bacterial strains with restriction enzymes; size fractionating the DNA fragments, so produced, into DNA fractions of 5 to 10 Kb, preferably 7 to 10 Kb; ligating these fractions to cloning vectors; screening the E. coli, transformed with the cloning vectors, with a DNA probe that was constructed from a region of known ISP
10 protein genes or with a DNA probe based on specific PCR fragments generated from isp DNA using primers corresponding to certain regions within known ISP protein genes. Such "other ISP proteins" isolated in accordance with this invention, like the ISP proteins of this invention, are characterized by any or all, preferably all, of the following characteristics: a) their significant insecticidal activity to larvae of the Southern corn rootworm, Diabrotica undecimpunctata, and to larvae of the Colorado potato beetle, Leptinotarsa decemlineata,
15 upon ingestion of such other ISP protein with an ISP-complimentary protein, preferably the mature ISP1A or ISP2A protein of this invention; b) their lack of significant insecticidal activity to larvae of lepidopteran insects, preferably Ostrinia nubilalis, upon ingestion of such other ISP protein with an ISP-complimentary protein, preferably the mature ISP1A or ISP2A
20 proteins of this invention; c) their apparent molecular weight of about 100 kD or about 45 kD of the mature forms (without signal peptide), d) their occurrence in the supernatant of a bacterial strain culture which is not Bacillus thuringiensis or Bacillus cereus, and e) by their lack of significant toxicity to corn rootworms, preferably Diabrotica virgifera, upon ingestion in the absence of an ISP complimentary protein.

25

Of course, any other DNA sequence differing in its codon usage but encoding the same protein or a similar protein with substantially the same insecticidal activity, can be constructed, depending on the particular purpose. It has been described in some prokaryotic and eucaryotic expression systems that changing the codon usage to that of the host cell is desired for gene expression in foreign hosts (Bennetzen & Hall, 1982, J. Biol. Chem. 257, 3026; Itakura, 1977, Science 198, 1056-1063). Codon usage tables are available in the literature (Wada et al., 1990, Nucl. Acids Res. 18, 2367-1411; Murray et al., 1989, Nucleic Acids Research 17, 477-498) and in the major DNA sequence databases. Accordingly, synthetic DNA sequences can be constructed so that the same or substantially the same proteins are produced. It is evident that several DNA sequences can be devised once the amino acid sequence of the ISP proteins of this invention is known. Such other DNA sequences include synthetic or semi-synthetic DNA sequences that have been changed in order to inactivate certain sites in the gene, e.g. by selectively inactivating certain cryptic regulatory or processing elements present in the native sequence, or by adapting the overall codon usage to that of a more related host organism, preferably that of the host organism in which expression is desired. Synthetic DNA sequences could also be made following the procedures described in EP 0 385 962, EP 0 618 967, or EP 0 682 115.

Small modifications to a DNA sequence such as described above can be routinely made by PCR-mediated mutagenesis (Ho et al., 1989, Gene 77, 51-59; White et al., 1989, Trends in Genet. 5, 185-189). New synthetic or semi-synthetic genes can be made by automated DNA synthesis and ligation of the resulting DNA fragments.

To prevent or delay the development of resistance by coleopteran insects, particularly corn rootworm, cotton boll weevil or Colorado potato beetle, preferably Leptinotarsa decemlineata, Diabrotica barberi, Diabrotica undecimpunctata, Anthonomus grandis or

Diabrotica virgifera, to transgenic hosts, particularly plants, expressing ISP proteins of this invention or their equivalents, it is preferred to also express in the same host, preferably a transgenic plant, another protein or another protein complex, which has a different mode of action, and a high toxicity to the same insect targeted by the first toxin or toxin complex

5 when produced in a transgenic host, preferably a plant. Suitable candidates to be combined with the ISP1A and ISP2A of the invention include the mature VIP1Aa protein when combined with the mature VIP2Aa or VIP2Ab protein of PCT publication WO 96/10083 in case these VIP proteins have a different mode of action compared to the ISP proteins; the corn rootworm toxins of *Photorhabdus* or *Xenorhabdus* spp., e.g., the insecticidal proteins of

10 *Photorhabdus luminescens* W-14 (Guo et al., 1999, J. Biol. Chem. 274, 9836-9842); the CryET70 protein of WO 00/26378; the insecticidal proteins produced by Bt strains PS80JJ1, PS149B1 and PS167H2 as described in WO 97/40162, particularly the about 14 kD and about 44 kD proteins of Bt strain PS149B1; the Cry3Bb protein of US patent 6,023,013; protease inhibitors such as the N2 and R1 cysteine proteinase inhibitors of soybean (Zhao et al., 1996,

15 Plant Physiol. 111, 1299-1306) or oryzastatine such as rice cystatin (Genbank entry S49967), corn cystatin (Genbank entries D38130, D10622, D63342) such as the corn cystatin expressed in plants as described by Irie et al. (1996, Plant Mol. Biol. 30, 149-157). Also included herein are all equivalents and variants, such as truncated proteins retaining insecticidal activity, of any of the above proteins.

20 Such combined expression can be achieved by transformation of a plant already transformed to express a corn rootworm toxic protein or protein complex, or by crossing plants transformed to express different corn rootworm toxic proteins. Alternatively, expression of the ISP proteins of the invention can be induced in roots upon feeding of corn rootworm larvae on root tissue, e.g., by using a wound-induced promoter region, preferably a wound-induced root-preferred

25 promoter region.

The 5 to 10, preferably 7 to 10, Kb fragments, prepared from total DNA of the isp genes of the invention, can be ligated in suitable expression vectors and transformed in E. coli, and the clones can then be screened by conventional colony immunoprobng methods (French et al., 1986, Anal.Biochem. 156, 417-423) for expression of the toxin with monoclonal or polyclonal antibodies raised against the ISP proteins. Also, the 5 to 10 Kb fragments; prepared from total DNA of the bacterial strains of the invention, can be ligated in suitable Bt shuttle vectors (Lereclus et al., 1992, Bio/Technology 10, 418) and transformed in a crystal minus Bt-mutant. The clones are then screened for production of ISP proteins (by SDS-PAGE, Western blot and/or insect assay).

The genes encoding the ISP proteins of this invention can be sequenced in a conventional manner (Maxam and Gilbert, 1980, Methods in Enzymol. 65, 499-560; Sanger, 1977, Proc. Natl. Acad. Sci. USA 74, 5463-5467) to obtain the DNA sequence. Sequence comparisons indicated that the genes are different from previously described genes encoding proteins secreted during the vegetative growth phase of Bacillus or other bacterial species and Bacillus thuringiensis crystal proteins with activity against Coleoptera (Crickmore, et al., 1998, Microbiology and Molecular Biology Reviews Vol 62: 807-813; WO 98/44137, WO 94/21795, WO 96/10083, WO 00/09697, WO 99/57282, and WO 97/46105).

In order to express all or an insecticidally effective part of the DNA sequence encoding an ISP protein of this invention in E. coli, in other bacterial strains or in plants, suitable restriction sites can be introduced, flanking the DNA sequence. This can be done by site-directed mutagenesis, using well-known procedures (e.g., Stanssens et al., 1989, Nucleic Acids Research 12, 4441-4454; White et al., 1989, supra). For expression in plants, the isp DNA of the invention, or its equivalent, is usually contained in a chimeric gene, flanked by 5' and 3' regulatory sequences including a plant-expressible promoter region and 3'

transcription termination and polyadenylation sequence active in plant cells. In order to obtain improved expression in plants, the codon usage of the isp DNA or its equivalent of this invention can be modified to form an equivalent, modified or artificial gene or gene part, or the isp DNA or insecticidally-effective parts thereof can be inserted in the chloroplast
5 genome and expressed there using a chloroplast-active promoter (e.g., Mc Bride et al., 1995, Bio/Technology 13, 362). For obtaining enhanced expression in monocot plants such as corn, a monocot intron also can be added to the chimeric gene, and the isp DNA sequence or its insecticidal part can be further changed in a translationally neutral manner, to modify possibly inhibiting DNA sequences present in the gene part by means of site-directed intron
10 insertion and/or by introducing changes to the codon usage, e.g., adapting the codon usage to that most preferred by the specific plant (Murray et al., 1989, supra) without changing significantly the encoded amino acid sequence.

The insecticidally effective isp DNA or its equivalent, preferably the isp chimeric gene,
15 encoding an ISP protein, can be stably inserted in a conventional manner into the nuclear genome of a single plant cell, and the so-transformed plant cell can be used in a conventional manner to produce a transformed plant that is insect-resistant. In this regard, a disarmed Ti-plasmid, containing the insecticidally effective isp gene part, in Agrobacterium tumefaciens can be used to transform the plant cell, and thereafter, a transformed plant can
20 be regenerated from the transformed plant cell using the procedures described, for example, in EP 0 116 718, EP 0 270 822, PCT publication WO 84/02913 and published European Patent application ("EP") 0 242 246 and in Gould et al. (1991, Plant Physiol. 95, 426-434). Preferred Ti-plasmid vectors each contain the insecticidally effective isp chimeric gene sequence between the border sequences, or at least located to the left of the right border
25 sequence, of the T-DNA of the Ti-plasmid. Of course, other types of vectors can be used to transform the plant cell, using procedures such as direct gene transfer (as described, for

example in EP 0 233 247), pollen mediated transformation (as described, for example in EP 0 270 356, PCT publication WO 85/01856, and US Patent 4,684,611), plant RNA virus-mediated transformation (as described, for example in EP 0 067 553 and US Patent 4,407,956), liposome-mediated transformation (as described, for example in US Patent 4,536,475), and other methods such as the methods for transforming certain lines of corn (Fromm et al., 1990, Bio/Technology 8, 833-839; Gordon-Kamm et al., 1990, The Plant Cell 2, 603-618, US Patent 5,767,367) and rice (Shimamoto et al., 1989, Nature 338, 274-276; Datta et al., 1990, Bio/Technology 8, 736-740) and the recently described method for transforming monocots generally (PCT publication WO 92/09696).

Different conventional procedures can be followed to obtain combined expression of two ISP proteins in transgenic plants as summarized below:

I. Chimeric gene constructs whereby two ISP DNA sequences and a marker gene are transferred to the plant genome as a single piece of DNA and lead to the insertion in a single locus in the genome.

Ia. The genes can be engineered in different transcriptional units each under control of a distinct promoter.

To express two ISP proteins and a marker protein as separate transcriptional units, several promoter fragments directing expression in plant cells can be used as described above. All combinations of the promoters mentioned above in the chimaeric constructs for one ISP gene are possible. The ISP coding region in each chimeric gene of this invention can be the intact isp gene or preferably an insecticidally-effective part of the intact isp gene. The individual chimeric genes are cloned in the same plasmid vector according to standard procedures (e.g., EP 0 193 259).

Ib. Two genes (e.g., either an isp and a marker gene or two isp genes) or more can be combined in the same transcriptional unit.

To express two isp genes in the same transcriptional unit, the following cases can be distinguished:

In a first case, hybrid genes in which the coding region of one gene is fused in frame
5 with the coding region of another gene can be placed under the control of a single promoter. Fusions can be made between either an isp and a marker gene or between two isp genes.

Also, between each gene encoding an ISP or an insecticidally-effective fragment thereof, a gene fragment encoding a protease (e.g., trypsin, cysteine or serine protease)-
10 sensitive protein part could be included, such as a gene fragment encoding a part of the ISPs which is removed or cleaved upon activation by the midgut enzymes of the target insect species. Alternatively, between each gene encoding an ISP or an insecticidally-effective fragment thereof, a gene fragment can be included encoding a peptide of about 16-
20 amino acids which has the capability to mediate cleavage at its own C-terminus by an
15 enzyme-independent reaction (Halpin et al., 1999, Plant J. 17, 453-459, US Patent 5,846,767), or a gene fragment encoding a linker peptide sequence which allows the production of a recombinant cell with significant toxicity to the target insects.

In a second case, the coding regions of the two respective isp genes can be combined in
20 dicistronic units placed under the control of a promoter. The coding regions of the two isp genes are placed after each other with an intergenic sequence of defined length. A single messenger RNA molecule is generated, leading to the translation into two separate gene products. Based on a modified scanning model (Kozak, 1987, Mol. Cell. Biol. 7, 3438-3445), the concept of reinitiation of translation has been accepted provided that a termination codon
25 in frame with the upstream ATG precedes the downstream ATG. Experimental data also

demonstrated that the plant translational machinery is able to synthesize several polypeptides from a polycistronic mRNA (Angenon et al., 1989, Mol. Cell Biol. 9, 5676-5684).

Based on the mechanism of internal initiation of translation (Jackson and Kaminski, 1995, RNA 1, 985-1000) initiation of translation of the second gene occurs by binding of the 43S pre-initiation complex to a specific intergenic sequence (internal ribosome entry sequence; IRES). Experimental data also demonstrated that the plant translational machinery is able to synthesize several polypeptides from a polycistronic mRNA containing intergenic IRES-elements (Hefferon et al., 1997, J Gen Virol 78, 3051-3059; Skulachev et al., 1999, Virology 263, 139-154; PCT patent publication WO 98/54342).

II. A chimeric construct with one isp gene that is transferred to the genome of a plant already transformed with one isp gene :

Several genes can be introduced into a plant cell during sequential transformation steps (retransformation), provided that an alternative system to select transformants is available for the second round of transformation, or provided that the selectable marker gene is excised from the plant genome using DNA recombination technology (e.g., published PCT applications WO 94/17176 and WO 91/09957). This retransformation leads to the combined expression of isp genes which are introduced at multiple loci in the genome. Preferably, two different selectable marker genes are used in the two consecutive transformation steps. The first marker is used for selection of transformed cells in the first transformation, while the second marker is used for selection of transformants in the second round of transformation. Sequential transformation steps using kanamycin and hygromycin have been described, for example by Sandler et al. (1988, Plant Mol. Biol. 11, 301-310) and Delauney et al. (1988, Proc. Natl. Acad. Sci. U.S.A. 85, 4300-4304).

III. Chimeric constructs with isp genes, that are separately transferred to the nuclear genome of separate plants in independent transformation events and are subsequently combined in a single plant genome through crosses.

5

The first plant should be a plant transformed with a first isp gene or an F1 plant derived thereof through selfing (preferably an F1 plant which is homozygous for the isp gene). The second plant should be a plant transformed with a second isp gene or an F1 plant derived thereof through selfing (preferably an F1 plant which is homozygous for the second isp gene). Selection methods can be applied to the plants obtained from this cross in order to select those plants having the two isp genes present in their genome (e.g., Southern blotting) and expressing the two ISPs (e.g., separate ELISA detection of the immunologically different ISPs). This is a useful strategy to produce hybrid varieties from two parental lines, each transformed with a different isp gene, as well as to produce inbred lines containing two different isp genes through crossing of two independent transformants (or their F1 selfed offspring) from the same inbred line.

IV. Chimeric constructs with one or more isp genes separately transferred to the genome of a single plant in the same transformation experiment leading to the insertion of the respective chimeric genes at the same or at multiple loci.

Cotransformation involves the simultaneous transformation of a plant with two different expression vectors, one containing a first isp gene, the second containing a second isp gene. Along with each isp gene, a different marker gene can be used, and selection can be made with the two markers simultaneously. Alternatively, a single marker can be used, and a sufficiently large number of selected plants can be screened in order to find those plants

having the two isp genes (e.g., by Southern blotting) and expressing the two proteins (e.g., by means of ELISA). Cotransformation with more than one T-DNA can be accomplished by using simultaneously two different strains of *Agrobacterium*, each with a different Ti-plasmid (Depicker et al., 1985, Mol. Gen. Genet. 201, 477-484) or with one strain of *Agrobacterium* containing two T-DNAs on separate plasmids (de Framond et al., 1986, Mol. Gen. Genet. 202, 125-131). Direct gene transfer, using a mixture of two plasmids or fragments thereof, can also be employed to cotransform plant cells with a selectable and a non-selectable gene (Schocher et al., 1986, Bio/technology 4, 1093-1096).

The resulting transformed plant can be used in a conventional plant breeding scheme to produce more transformed plants with the same characteristics or to introduce the insecticidally effective isp gene in other varieties of the same or related plant species. Seeds, which are obtained from the transformed plants, contain the insecticidally effective isp gene as a stable genomic insert. Cells of the transformed plant can be cultured in a conventional manner to produce the insecticidally effective portion of the ISP protein which can be recovered for use in conventional insecticide compositions against insects. Of course, the above possibilities of combined production of ISP proteins in plants is as well applicable to active fragments of the ISP proteins or the ISP equivalents of this invention.

The insecticidally effective isp gene is inserted in a plant cell genome so that the inserted gene is downstream (i.e., 3') of, and operably linked to, a promoter which can direct the expression of the gene part in the plant cell. This is preferably accomplished by inserting the isp chimeric gene in the plant cell genome, particularly in the nuclear or chloroplast genome. Preferred promoters include: the strong constitutive 35S promoters (the "35S promoters") of the cauliflower mosaic virus (CaMV) of isolates CM 1841 (Gardner et al., 1981, Nucl. Acids Res. 9, 2871-2887), CabbB-S (Franck et al., 1980, Cell 21, 285-294) and CabbB-JI (Hull and Howell, 1987, Virology 86, 482-493); promoters from the ubiquitin family (e.g., the maize

ubiquitin promoter of Christensen et al., 1992, Plant Mol. Biol. 18, 675-689; see also Cornejo et al., 1993, Plant Mol. Biol. 23, 567-581), the gos2 promoter (de Pater et al., 1992, Plant J. 2, 837-844), the emu promoter (Last et al., 1990, Theor. Appl. Genet. 81, 581-588), rice actin promoters such as the promoter described by Zhang et al. (1991, The Plant Cell 3, 1155-1165); and the TR1' promoter and the TR2' promoter (the "TR1' promoter" and "TR2' promoter", respectively) which drive the expression of the 1' and 2' genes, respectively, of the T-DNA (Velten et al., 1984, EMBO J 3, 2723-2730). Alternatively, a promoter can be utilized which is not constitutive but rather is specific for one or more tissues or organs of the plant (e.g., leaves and/or roots) whereby the inserted isp gene is expressed only in cells of the specific tissue(s) or organ(s). In a preferred embodiment of the invention, the insecticidally effective isp gene is selectively expressed in the roots of a plant, preferably corn, by placing the insecticidally effective gene part under the control of a root-preferred promoter (i.e., a promoter which is most active in root tissue, preferably a promoter with no or little transcription in non-root tissue such as pollen, leaves and stem, more particularly a promoter only resulting in detectable transcription in root tissue). Root-preferred promoters need not be exclusively active in root tissue. Root-preferred promoters in accordance with this invention include but are not limited to: promoters located immediately upstream of a DNA sequence corresponding to a root-specific cDNA, preferably from corn; the promoter of US patent 5,837,876, published PCT patent applications WO 00/29594, WO 00/73474, WO 01/00833, or US Patent 6,008,436; the ZRP2 promoter of US Patent 5,633,363 and Held et al. (1997, Plant Mol. Biol. 35, 367-375, Genbank accession number U38790); the promoter of US Patent 5,817,502; the promoter described by de Framond (1991, FEBS 290: 103-106; EP 0 452 269), the root-specific promoter of the peroxidase gene POX1 from wheat (Hertig et al., 1991, Plant Molec. Biol. 16, 171-174), the promoter of US patent 5,837,848, the promoter of PCT publication WO 015662, the promoter of Goddemeier et al. (1998, Plant Mol. Biol. 36, 799-802). Other root-specific or root-preferred promoters which may be useful in this invention

75749-27

include the promoters described by Hirel et al., 1992, Plant Mol. Biol. 20, 207; Keller and Baumgartner, 1991, The Plant Cell 3, 1051-1061; Sanger et al., 1990, Plant Mol. Biol. 14, 433-443; Miao et al., 1991, The Plant Cell 3, 11-22; Bogusz et al., 1990, The Plant Cell, 2, 633-641; Leach and Aoyagi, 1991, Plant Science 79, 69-76; Teeri et al., 1989, EMBO Journal 8, 343-
5 350.

Expression in leaves of a plant can be achieved by using the promoter of the ribulose-1,5-bisphosphate carboxylase small subunit gene of the plant itself or of another plant such as pea as disclosed in US Patent 5,254,799. Another alternative is to use a
10 promoter whose expression is inducible (e.g., by temperature or chemical factors). Also, for corn plants, particularly at the time adult corn rootworms are present in corn rootworm fields, expression in pollen is preferred, by using promoter regions resulting in expression in corn pollen, e.g. the promoters described in published PCT applications WO 93/25695 and WO
01/12799.

15

The insecticidally effective isp gene is inserted in the plant genome so that the inserted gene part is upstream (i.e., 5') of suitable 3' end transcription regulation signals (i.e., transcript formation and polyadenylation signals).

20 This is preferably accomplished by inserting the isp chimeric gene in the plant cell genome. The choice of the 3' regulatory sequence in the chimeric gene of the invention is not critical. In some cases good expression can be obtained when no such region is present in the chimeric gene, since a DNA sequence at the insertion site can act as a 3' regulatory sequence. Preferred polyadenylation and transcript formation signals include those of the
25 CaMV 35S (Mogen et al., 1990, The Plant Cell 2, 1261-1272), the octopine synthase gene (Gielen et al., 1984, EMBO J 3, 835-845), the nopaline synthase gene (Depicker et al., 1982,

J. Mol. Appl. Genet. 1, p. 561), and the T-DNA gene 7 (Velten and Schell, 1985, Nucleic Acids Research 13, 6981-6998), which act as 3'-untranslated DNA sequences in transformed plant cells.

- 5 The insecticidally effective isp gene can optionally be inserted in the plant genome as a hybrid gene (US Patent 5,254,799; Vaeck et al., 1987, Nature 327, 33-37) under the control of the same promoter as a selectable marker gene, such as the neo gene (EP 0 242 236) encoding kanamycin resistance, so that the plant expresses a fusion protein.
- 10 All or part of the isp gene, can also be used to transform other bacteria, such as a B. thuringiensis strain which has insecticidal activity against Lepidoptera or Coleoptera, or root-colonizing bacteria, e.g. root-colonizing Pseudomonas putida (Vilchez et al., J. Bacteriol. 182, 91-99). Thereby, a transformed bacterial strain can be produced which is useful for combatting insects and which can affect a wide spectrum of lepidopteran and coleopteran
- 15 insect pests. Transformation of bacteria, such as bacteria of the genus Agrobacterium, Bacillus or Escherichia, with all or part of the isp gene of this invention or its equivalent, incorporated in a suitable cloning vehicle, can be carried out in a conventional manner, such as heat shock transformation (Bergmans et al., 1981, J. Bacteriol 146, 564-570) or conventional electroporation techniques as described in Mahillon et al. (1989, FEMS
- 20 Microbiol. Letters 60, 205-210) and in PCT Patent publication WO 90/06999.

Transformed bacterial strains, such as Bacillus species strains, preferably Bacillus thuringiensis strains, containing the isp gene of this invention can be fermented by conventional methods (Dulmage, 1981, "Production of Bacteria for Biological Control of

25 Insects" in Biological Control in Crop Production, Ed. Paparizas, D.C., Osmun Publishers, Totowa, N.J., USA, pp. 129-141; Bernhard and Utz, 1993, "Production of Bacillus

thuringiensis insecticides for experimental and commercial uses", In Bacillus thuringiensis, An Environmental Biopesticide: Theory and Practice, pp.255-267, eds. Entwistle, P.F., Cory, J.S., Bailey, M.J. and Higgs, S., John Wiley and Sons, New York) to provide high yields of cells.

5

An insecticidal, particularly anti-coleopteran, preferably anti-corn rootworm composition of this invention can be formulated in a conventional manner using the microorganisms transformed with the isp gene, or preferably their respective ISP proteins or insecticidally effective portions thereof as an active ingredient, together with suitable carriers, diluents, emulsifiers and/or dispersants (e.g., as described by Bernhard and Utz, 1993, supra). This insecticide composition can be formulated as a wettable powder, pellets, granules or dust or as a liquid formulation with aqueous or non-aqueous solvents as a foam, gel, suspension, concentrate, etc. Known microorganisms include cells of Pseudomonas or other bacteria which serve to encapsulate the proteins in a stable environment prior to application to the insects. Also included in the invention is a product comprising the ISP1A and ISP2A protein of the invention as a combined preparation for simultaneous, separate or sequential use to protect corn plants against corn rootworms, particularly such product is an insecticidal composition or a transgenic corn plant.

20 A method for controlling insects, particularly Coleoptera, in accordance with this invention can comprise applying (e.g., spraying), to a locus (area) to be protected, an insecticidal amount of the ISP proteins or host cells transformed with the isp gene of this invention. The locus to be protected can include, for example, the habitat of the insect pests or growing vegetation or an area where vegetation is to be grown.

25

To obtain the ISP protein, cells of the recombinant hosts expressing the ISP protein can be grown in a conventional manner on a suitable culture medium and the protein can then be obtained from the medium using conventional means. The ISP protein can then be separated and purified by standard techniques such as chromatography, extraction,
5 electrophoresis, or the like. The protease-resistant toxin form can then be obtained by protease, e.g. cysteine or serine protease, digestion of the protein.

Bio-assays for testing the insecticidal efficacy are well known in the art. A method for corn rootworm testing is described in Marrone et al. (1985, J. Econ. Entomol. 78, 290-293), but
10 many other methods are available to test insects on artificial diet or on transgenic plants, e.g. US Patent 5,990,383. In a suitable bio-assay the proper controls are included to check for background mortality.

The following Examples illustrate the invention, and are not provided to limit the invention or
15 the protection sought. Many variants or equivalents of these Examples can be made or designed by a person of ordinary skill in the art without departing from the teachings of the invention and the common knowledge. The sequence listing referred to in this application (which forms part of this application and is attached thereto), is as follows:

20 Sequence Listing:

SEQ ID No. 1 - amino acid and DNA sequence of ISP1A protein and DNA

SEQ ID No. 2 - amino acid sequence of ISP1A protein.

SEQ ID No. 3 - amino acid and DNA sequence of ISP2A protein and DNA.

SEQ ID No. 4 - amino acid sequence ISP2A protein.

25 SEQ ID No. 5 - DNA sequence of the short zrp2 promoter fragment (n= any nucleotide)

SEQ ID No. 6 - DNA sequence of the long zrp2 promoter fragment (n=any nucleotide)

SEQ ID No. 7 - DNA sequence encoding ISP2A-1 protein

SEQ ID No. 8 - amino acid sequence of ISP2A-1 protein

SEQ ID No. 9 - DNA sequence encoding ISP1A-1 protein fragment

SEQ ID No. 10 - amino acid sequence of ISP1A-1 protein

5

Unless otherwise stated in the Examples, all procedures for making and manipulating recombinant DNA are carried out by the standard procedures described in Sambrook et al., Molecular Cloning - A Laboratory Manual, Second Ed., Cold Spring Harbor Laboratory Press, NY (1989), and in Volumes 1 and 2 of Ausubel et al. (1994) Current Protocols in Molecular Biology, Current Protocols, USA. Standard materials and methods for plant molecular biology work are described in Plant Molecular Biology Labfax (1993) by R.R.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications (UK). Procedures for PCR technology can be found in "PCR protocols: a guide to methods and applications", Edited by M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White (Academic Press, Inc., 1990).

10
15

EXAMPLES

Example 1: Characterization of the IB120-A strain

20

A bacterial strain, named herein the IB120-A strain, was found in a field in the state of Morelos, Mexico.

The strain was grown overnight on LB agar plates (LB medium with 1.5 % agar added; LB medium: 10 g/l trypton, 10 g/l NaCl, 5 g/l yeast extract, pH 7.3) at 28°C. For small scale cultures, 20 ml TB medium (Terrific Broth: 12 g/l tryptone, 24 g/l yeast extract, 3.8 g/l KH₂PO₄, 12.5 g/l K₂HPO₄, 5 ml/l glycerol, pH 7.1) was inoculated and grown for 65 hours at 28°C on a

25

rotating platform having about 70 rotations per minute. After 65 hours, a protease inhibitor mixture was added to the culture. This cocktail has the following ingredients (volumes given are those required to add to one 20 ml culture): 200µl PMSF (100mM), 200µl of a mixture of benzamidine.HCl (100mM) and epsilon-amino-n-caproic acid (500 mM), 400µl EGTA (0.5M),
5 40µl antipain (0.5mg/ml) / leupeptin (0.5mg/ml) and 20µl beta-mercapto ethanol (14M). The culture medium to which the protease inhibitor mixture had been added, was then centrifuged for 20 minutes at 3000 rpm. In some cases, the supernatant was concentrated about 4 times using Centriprep YM-10 Centrifugal Filter Devices (Millipore Cat. No. 4305). For long term storage, a loop of sporulated cells was added to 0.5ml of 25% or 50% glycerol and after
10 vortexing, stored at -70°C. Sporulated cells were obtained by growth of the strain on LB agar plates until sporulation (as apparent under the light microscope).

After cultivating on LB agar plates of single cell colonies, microscopical analysis of the IB120-A strain culture showed the presence of rod-shaped motile single vegetative cells and swollen
15 sporangia containing an ovale spore. No parasporal crystals were detected in cultures of the IB120-A strain.

Supernatant of the IB120-A strain showed strong toxicity to Diabrotica virgifera larvae (hereinafter termed "Dv") in surface contamination assays on artificial diet (diet as described by
20 Sutter et al. (1971, J. Econ. Entomol. 64, 65-67)). The assays were done at 24 °C (+/- 1°C) in 24 multiwell Costar plates at 50 microliter toxin solution per well (2cm²), 6 wells were analyzed with 4 L1 (first instar) larvae per well for each sample (scoring after 7 days).

Screening of supernatant harvested at 7, 24, 30, 48 and 120 hours after initiation of the culture
25 showed the strongest toxicity to Dv at 48 hours, with no significant toxicity found at 7 hours. At

48 hours, mortality in the 50-60 % range was still found when the supernatant (at total protein concentration of 214 microgr/ml) was diluted at 1/1024.

Loss of activity of IB120-A supernatant harvested 48, 65 and 144 hours after culture initiation (in dilutions of 1/1 to 1/8) upon heat treatment and retention of activity after ammonium sulphate precipitation indicated that the toxic compound is likely a protein.

Preliminary characterization of the IB120-A strain by PCR primers for gyrase B genes (Yamada et al., 1999, Appl. Environm. Microbiol. 65, 1483-1490) suggested that it was not a Bacillus strain of the subspecies thuringiensis, anthracis or cereus. Also, no amplification products were obtained using PCR primers specifically characterizing the 16S RNA sequences of the genus Paenibacillus which are also recognizing the species Bacillus lentimorbus and Bacillus popilliae (Pettersson et al., 1999, Int J Syst Bacteriol. 49(2), 531-540). Growth in NB (nutrient broth: 3 g/l bacto beef extract, 5 g/l bacto peptone, pH 6.8) medium indicated that the new strain was not a Bacillus larvae species. Thus, this IB120-A strain seems to be different from previously isolated Bacillus strains known to produce secreted insecticidal proteins which are not contained in crystals.

Based on the rod-like shape and the aerobic growth, the strain could be a Bacillus species strain. With a set of general and specific primers for cry3 Bacillus thuringiensis genes, no amplification products were found when analyzing strain IB120-A.

Detailed analysis of the IB 120-A strain using standard microbial identification techniques, including fatty acid analysis and API 50CHB tests combined with API 20E tests, showed that this strain is a Brevibacillus laterosporus species strain.

Example 2. Isolation and characterization of isp1 and isp2 DNAs/proteins:

In order to isolate the genes responsible for the toxicity of IB120-A, total DNA from this strain was prepared and partially digested with Sau3A. The digested DNA was size fractionated on a sucrose gradient and fragments ranging from 7 kb to 10 kb were ligated to the BamH1-digested and TsAP (thermosensitive alkaline phosphatase)- treated cloning vector pUC19 (Yannisch-Perron et al, 1985, Gene 33, 103-119.). The ligation mixture was electroporated in E. coli XL1-Blue cells. Transformants were plated on LB-triacillin plates containing Xgal and IPTG and white colonies were selected to be used in filter hybridization experiments. Recombinant E. coli clones containing the vector were then screened with a DIG labeled probe which was prepared as follow. First, a PCR was performed using as template cells from strain IB120-A. The resulting amplification product was gel-purified and used as template in a secondary PCR reaction using DIG- labeled dNTPs and the same primers as in the first PCR reaction. An appropriate amount of this amplification product was used in hybridization reactions.

Following the identification of a positive colony containing a plasmid harboring the full length isp genes, the sequence of these genes was determined using the dye terminator labeling method and a Perkin Elmer ABI Prism-377 DNA sequencer. The sequences of the 2 open reading frames found in the cloned DNA fragment of a positive colony are shown in SEQ ID No. 1 and 3. These DNA sequences were found to be organized in an operon and encode novel proteins, ISP1A (SEQ ID No. 2) and ISP2A (SEQ ID No. 4) which have been found to be the causal agents of the high insecticidal activity observed. A positive colony containing the pUC-derived plasmid with the genes responsible for toxicity (in plasmid pUCIB120/ISP) has been deposited under the provisions of the Budapest treaty in E. coli XL1Blue as LMBP 4009 on January 11, 2000.

A plasmid preparation was made from the positive colony and this plasmid was cut using XbaI and EcoRI (resulting in an about 7 Kb fragment) and ligated in the shuttle vector pSL40 which had been cut using the same restriction enzymes. This yielded plasmid pSLIB120/ISP. The plasmid pSLIB120/ISP was then transferred into a crystal-minus B. thuringiensis strain.

5 Supernatant from this recombinant Bt strain was obtained as follows.

The strain was grown overnight on LB agar plates containing erythromycin (20 µg/ml) at 28°C. For small scale cultures, 20 ml TB medium containing erythromycin (20 µg/ml) was inoculated and grown for 65 hours at 28°C on a rotating platform having about 70 rotations per minute.

10 After 65 hours, a protease inhibitor mixture was added to the culture. This cocktail has the following ingredients (volumes given are those required to add to one 20 ml culture): 200µl PMSF (100mM), 200µl of a mixture of benzamidine.HCl (100mM) / epsilon-amino-n-caproic acid (500 mM), 400µl EGTA (0.5M), 40µl antipain (0.5 mg/ml) / leupeptin (0.5 mg/ml) and 20µl beta-mercapto ethanol (14M). The culture medium to which the protease inhibitor mixture had
15 been added, was then centrifuged for 20 minutes at 3000 rpm. In some cases, the supernatant was concentrated about 4 times using centriprep YM-10 Centrifugal Filter Devices (Millipore, Cat. No. 4305).

Insecticidal activity of the supernatant at 48 hours after culture initiation of the recombinant Bt
20 strain containing the pSLIB120/ISP plasmid using the Dv surface contamination assay described above, showed that the supernatant still had significant mortality in the 60 % range at a 1/1024 dilution, while control mortality (the controls included supernatant of the untransformed Bt strain) was 0 %. This shows that the isolated DNA encodes an insecticidal ingredient of the IB120-A strain, and upon transfer to another bacterium is also expressed and secreted in the
25 culture medium. Analysis of toxicity of another independent Bt-crystal-minus strain transformed

with the pSLIB120/ISP plasmid confirmed the high insecticidal activity of the supernatants compared to that of the untransformed control.

The LC₅₀ value of the supernatant of the Bt strain expressing the two ISP proteins was found to be 3.5 ng/cm² after 4 days using the above-described surface contamination assay with Dv larvae (based on total supernatant protein concentration). Detailed analysis of the toxicity of supernatant produced by the recombinant Bt strain expressing the ISP1A and ISP2A proteins to selected coleopteran insects is reported in Table 1 below. The ISP proteins of the invention were found to have significant insecticidal activity to Western, Northern and Southern corn rootworm larvae, and also to larvae of the Colorado potato beetle and the cotton boll weevil.

The mature ISP1 and ISP2 proteins were purified to apparent homogeneity by ammonium sulphate precipitation followed by passage over different chromatographical columns. The chromatographical analysis suggest that the ISP1A and ISP2A proteins are present in the supernatant in equimolar ratio. The ISP1A protein was found to be of rather hydrophobic nature, while ISP2A was of rather hydrophilic nature. Amino-terminal sequence determination of the mature ISP1A and ISP2A proteins produced in the recombinant crystal-minus Bt strain showed that amino acid position 38 in SEQ ID No. 2 for ISP1A and amino acid position 51 in SEQ ID No. 4 for ISP2A are the N-terminal amino acids of the active proteins present in supernatant of such transformed Bt strain. The N-terminus for ISP1A was found to be IATTTQASKD, that for ISP2A was found to be LVKTTNNTED, which matches fully with the amino acid sequence of the DNA sequences isolated.

The apparent molecular weight of the pure mature ISP1A protein was determined to be about 100 kD in 8% SDS-PAGE gel electrophoresis, that of the pure mature ISP2A protein was determined to be about 45 kD protein in 10 % SDS-PAGE gel electrophoresis, using molecular weight markers of 37, 50, 75, 100, and 150 kD.

The activity of the mature ISP1A and ISP2A proteins as produced by the recombinant strain was evaluated against several insects. The results against selected coleopteran insects are shown in Table 1 below.

5 **Table 1: Coleopteran activity of ISP1A-ISP2A :**

Insect	Stage	LC50 (µg/ml)	LC90 (µg/ml)
		(95% CL)	(95% CL)
Dv	L1	0.437	1.689
		(0.321-0.563)	(1.250-2.597)
	L2	3.84	117.4
		(-)	(-)
	L3	21.674*	531.76
		(5.012-100.432)	(110.688-86864)
Db	L1	0.213	0.890
		(0.116-0.338)	(0.542-2.006)
Du	L1	4.91	30.06
		(1.65-13.26)	(11.52-329.72)
Ld	L1+2d	0.037	1.068
		(-)	(-)
Ag	L1	207.1	8759.2
		(84.3-654.7)	(1865.8-620175.8)

Legend to Table 1:

Dv: Diabrotica virgifera, Western corn rootworm; Db: Diabrotica barberi, Northern corn rootworm; Du: Diabrotica undecimpunctata howardi, Southern corn rootworm; Ld: Leptinotarsa
 10 decemlineata, Colorado potato beetle; Ag: Anthonomus grandis, cotton boll weevil; L1: first larval stage; L2: second larval stage; L3: third larval stage; L1+2d: 2d after egg hatch; *: 90%CL (CL=Confidence Limits, LC50: total supernatant protein concentration when 50% of the insects are killed).

Bioassays used: Dv, Db, Du: surface contamination assay on artificial diet (see above for
 15 description) at ratio of 25 µl/cm², scoring after 7 days; Ld: diptest with potato foliage (cut potato foliage dipped in toxin solution, allowed to dry and put in a petri dish (9 cm diam.) with 10 larvae; after 1 day untreated foliage was added to the petridish, scoring was after 5 days); Ag:

artificial diet incorporation test : diet as described by Moore and Whisnant, Handbook of Insect Rearing, Vol. I, Pritah Singh and R.F. Moore; Elsevier, Amsterdam, 1985, p. 217; assay: incorporation of 2 ml toxin solution per 25 ml diet, in 24 multiwell Costar plates (about 1 ml/well), 1 egg per well, 20 wells per sample, scoring after 14 days).

- 5 In the assays reported in Table 1, controls (untreated food, food supplied with supernatant of a non-transformed Bt-crystal-minus strain or with water) did not show any mortality above 20 % for any of the above insects (a variation from 5 to 20 % control mortality was found according to larval stage and insect species (the 20 % value is for the third larval stage of Dv)).
- 10 The LC50 values obtained for Diabrotica virgifera larvae using the bio-assay as described above, when an equimolar combination of ISP1A and ISP2A protein was applied after purification of these proteins from the supernatans of the recombinant Bt strain, were not significantly different from those obtained above.
- 15 Bio-assays of a mixture of the mature ISP1A and ISP2A protein in standard surface contamination assays against selected lepidopteran insects (Ostrinia nubilialis, Heliothis virescens, Helicoverpa zea, Spodoptera frugiperda, Sesamia nonagrioides, Spodoptera littoralis, Helicoverpa armigera, and Manduca sexta) showed that the ISP proteins of this invention do not cause any mortality in these insects above control levels. This evidences the
- 20 coleopteran-specific nature of the proteins of this invention. Preliminary bio-assays with two aphid species also showed that a mixture of the ISP1A and ISP2A proteins did not cause any significant mortality in these insects.

To determine which fragment of the isp genes are sufficient to encode an ISP protein complex

- 25 toxic to Diabrotica larvae, a series of C-terminally truncated ISP proteins were made by inserting a stopcodon in the ORF of the isp genes at different positions (and thus making

different 3' gene truncations). The stopcodons were inserted at random positions using the Genome Priming System (GPS, New England Biolabs, catalog #7100). Using this system, a 1.7 kb transposon (containing stopcodons in all three reading frames and 2 sequences complementary to specific primers) is inserted randomly but only once in the 'target' DNA. The position of the insertion, and therefore, the truncation can then be estimated by PCR screening using a gene-specific primer and a transposon-specific primer or can be precisely determined by sequencing. A set of constructs with a transposon at different positions in one of the isp genes was then transformed individually in a crystal minus Bt strain and the susceptibility of Western corn rootworm larvae to the supernatant of each recombinant Bt strain was tested using the assay described above.

For truncation of the isp2 gene, a fragment containing the gene was cut out of pSLIB120/ISP using BstBI and PmlI. Following the GPS reaction, this fragment was then ligated into the pSLIB120/ISP vector, cut with the same enzymes. Recombinant E. coli colonies were PCR-screened in order to estimate the position of the transposon insertion site. A set of 10 clones with the transposon at different positions in the second half (3' half) of the isp2 gene was selected. Plasmid DNA of these clones was transformed in a dcm and dam methylase negative E. coli strain GM2163. Plasmid DNA isolated from this strain was then transformed in a crystal-minus Bt strain. Supernatant was prepared and bioassayed, using supernatant from the Bt strain transformed with pSLIB120/ISP as a positive control and supernatant from the crystal-minus Bt strain as a negative control. For truncation of the isp1 gene, a fragment was cut out of pSLIB120/ISP using PmlI and EcoRI. The same methodology was followed as for the truncation of the isp2 gene.

The results of the transposon insertion analysis indicate that truncation of relatively large segments of the isp2A gene at the 3' end abolishes toxicity, indicating that no or only a

relatively short C-terminal truncation is allowed in the ISP2A protein. After sequencing, the smallest C-terminally truncated ISP2A toxin fragment retaining toxicity in this study when tested in combination with the ISP1A protein was found to end at amino acid position 449 in SEQ ID No. 4.

5

The results of these analyses indicate that a larger truncation of the ISP1A protein at the 3' end is possible: sequence determination showed that the toxic fragment with the largest C-terminal truncation ended at amino acid position 768 in SEQ ID No. 2. Thus, approximately 300 bp of the ISP1A coding region can be deleted at the 3' side without losing toxicity.

10

The above study also indicates that insecticidal activity is reduced to the background level when no functional ISP1A is produced, confirming the binary nature of the ISP proteins of this invention.

15 The ISP2A protein from amino acid position 51 to amino acid position 449 in SEQ ID No. 4 and the ISP1A protein from amino acid position 38 to amino acid position 768 in SEQ ID No. 2 are thus useful insecticidal fragments of the ISP proteins.

Furthermore, a chimeric gene containing a DNA fusion of the ISP1A and ISP2A genes was
20 expressed in a crystal-minus Bacillus thuringiensis berliner 1715 strain. A DNA sequence encoding an Arg-Lys-linker (RK RK RK) or a DNA sequence encoding a Gly-linker (GGGGGG) was provided between the open reading frames encoding the ISP2A and the ISP1A proteins (ISP2A still has its N-terminal signal peptide present, while ISP1A lacks its signal peptide), so that a translational fusion protein is produced; i.e. a fusion protein in which the specified linker
25 connects the ISP2A protein (as the N-terminal fusion partner) to the ISP1A protein (as the C-terminal fusion partner). The recombinant Bt strain proved to cause mortality to Diabrotica

virgifera which was significantly higher than that of the untransformed control strain (i.e., the crystal-minus Bt 1715 strain). This shows the possibility of producing the ISP proteins of the invention as a fusion protein in a recombinant host cell by using a single chimeric gene. Obviously, as explained in the description of the invention, multiple different approaches can be used to achieve the production of a fusion protein in the recombinant cell, e.g. cells of a plant.

Example 3: production of ISP proteins in plants:

To obtain optimal expression in plants, the isp1A and isp2A native DNA sequences are modified to produce modified DNA sequences encoding the ISP1A and ISP2A proteins. The optimized DNA sequences encoding the ISP1A-1 and ISP2A-1 proteins with their signal peptides replaced by a methionine and alanine amino acid are shown in SEQ ID No. 9 and 7, respectively.

These coding regions are inserted in a chimeric gene operably linked to suitable regulatory sequences, e.g., a promoter based on the short or long *zrp2* promoter sequences of SEQ ID No. 5 or 6 using standard techniques, and appropriate 3' transcription termination and polyadenylation sequences. In some constructs, the above chimeric genes contain a DNA sequence encoding the optimized chloroplast transit peptide (in place of the N-terminal bacterial signal peptide) as described in US Patent 5,510,471 (or Reissued US Patent RE 036449) resulting in targeting to the plastids or a signal peptide causing targeting to the cell wall. Other signal peptides effective in plant cells can similarly be used, preferably the signal peptide encoded by the alpha-amylase 3 gene of rice (Sutliff et al., 1991, Plant Molec. Biol. 16, 579-591), the signal peptide encoded by the ferredoxin NADP+ oxidoreductase gene from spinach (Oelmueller et al., 1993, Mol. Gen. Genet. 237, 261-272), or the signal peptide

encoded by the proteinase inhibitor II gene of potato (Keil et al., 1986, Nucl. Acids Res. 14, 5641-5650).

Corn cells are stably transformed with the above isp1A and isp2A chimeric genes using
5 Agrobacterium-mediated transformation (Ishida et al., 1996, Nature Biotechnology 14, 745-750; and U.S. Patent No. 5,591,616), protoplast transformation as described in US patent 5,792,936, or by electroporation using wounded and enzyme-degraded embryogenic callus, as described in WO 92/09696 or US Patent 5,641,664 (all of these above references are incorporated herein by reference). Corn plants are regenerated from the transformed cells and are selected for
10 optimal expression levels in roots and for adequate overall agronomic characteristics.

The thus obtained transgenic corn plants producing the above ISP proteins of the invention cause significant mortality to Diabrotica virgifera larvae attempting to feed on such plants.

In accordance with this invention, stably transformed corn plants are also obtained containing
15 several DNA constructs, so that the corn plant can express one or both the ISP proteins of the invention in its cells under control of the suitable regulatory control regions, together with an appropriate marker gene, preferably a herbicide resistance gene. In these DNA constructs, a preferred 3' transcription termination and polyadenylation sequence is a 215 bp fragment containing the 3' transcription termination and polyadenylation sequences obtained from
20 Cauliflower Mosaic virus (Oka et al., 1981, J. Mol. Biol. 147, 217-226). In some constructs, a 5' untranslated leader sequence can be added, preferably those chosen from the leader sequences of a cab22 gene from Petunia (Harpster et al., 1988, Mol. Gen Genetics 212, 182-190) or the zrp2 gene (Held et al, 1997, Plant Molecular Biology 35, 367-375, Genbank accession number U38790). Preferred promoters in the constructs are promoter-effective parts
25 of the 35S promoter (e.g., Odell et al., 1985, Nature 313, 810-812) or zrp2 promoters (Held et al, 1997, Plant Molecular Biology 35, 367-375, Genbank accession number U38790). The

constructs can also contain a signal peptide effective in plant cells besides the above optimized chloroplast transit peptide, namely the signal peptide of the alpha-amylase 3 gene of rice (Sutliff et al., 1991, Plant Molec. Biol. 16, 579-591), the signal peptide of the ferredoxin NADP+ oxidoreductase from spinach (Oelmueller et al., 1993, Mol. Gen. Genet. 237, 261-272), or the
 5 signal peptide from the proteinase inhibitor II of potato (Keil et al., 1986, Nucl. Acids Res. 14, 5641-5650). Transformations of plants with the ISP chimeric genes using a selection from the above preferred elements using techniques available to the person of ordinary skill in the art can be used to achieve the desired goal of the invention.

Preferably the plants also contain a gene encoding a protein conferring resistance to
 10 glufosinate or glyphosate, for which a different promoter and/or leader sequence is used, e.g. the *gos2* promoter and/or the *gos2* leader (de Pater et al., 1992, Plant J. 2, 837-844).

Needless to say, this invention is not limited to the above corn plants, transformation methods nor to the particular ISP proteins or DNA sequences used. Rather, the invention also includes
 15 any variant or equivalent of the ISP protein retaining insecticidal activity, such as a protein having substantially the amino acid sequence of the ISP proteins of the invention and having substantially the insecticidal activity of the ISP proteins. Also, any plant which is susceptible to damage by coleopteran insects, particularly corn rootworms, weevils or potato beetles, especially Diabrotica virgifera or Leptinotarsa decemlineata, preferably an insect selected from
 20 the group consisting of: Agelastica alni, Hypera postica, Hypera brunneipennis, Haltica tombacina, Anthonomus grandis, Tenebrio molitor, Triboleum castaneum, Dicladispa armigera, Trichispa serica, Oulema oryzae, Colaspis brunnea, Lissorhynchus oryzae, Phyllotreta cruciferae, Phyllotreta striolata, Psylliodes punctulata, Entomoscelis americana, Meligethes aeneus, Ceutorynchus sp., Psylliodes chrysocephala, Phyllotreta undulata, Leptinotarsa
 25 decemlineata, Diabrotica undecimpunctata undecimpunctata, Diabrotica undecimpunctata

howardi, Diabrotica barberi and Diabrotica virgifera, is included in the invention as a preferred target for transformation with a DNA encoding an ISP protein.

SEQUENCE LISTING

<110> Bayer Bioscience N.V.

<120> Insecticidal Secreted Proteins of Brevibacillus Laterosporus

<130> CRWISPW01

<140> CA 2,409,031

<141> 2001-05-17

<150> US 09/573,872

<151> 2000-05-18

<160> 10

<170> PatentIn version 3.0

<210> 1

<211> 2616

<212> DNA

<213> Brevibacillus laterosporus

<220>

<221> CDS

<222> (1)..(2613)

<400> 1

atg	aaa	tat	atg	aaa	aaa	gga	tta	tct	agt	ggt	gta	ata	ggg	acg	ttg	48
Met	Lys	Tyr	Met	Lys	Lys	Gly	Leu	Ser	Ser	Val	Val	Ile	Gly	Thr	Leu	
1				5				10						15		

ttc	gct	tct	atg	ttt	ttg	aat	ggg	aat	gta	aat	gct	ggt	tac	gcg	aac	96
Phe	Ala	Ser	Met	Phe	Leu	Asn	Gly	Asn	Val	Asn	Ala	Val	Tyr	Ala	Asn	
			20				25						30			

agt	aaa	aca	aat	cag	att	gct	aca	aca	act	cag	gca	agc	aaa	gac	aac	144
Ser	Lys	Thr	Asn	Gln	Ile	Ala	Thr	Thr	Thr	Gln	Ala	Ser	Lys	Asp	Asn	
		35				40						45				

caa	ata	gat	cga	gaa	gga	cta	ctt	ggt	tat	tac	ttt	aaa	gga	aaa	gat	192
Gln	Ile	Asp	Arg	Glu	Gly	Leu	Leu	Gly	Tyr	Tyr	Phe	Lys	Gly	Lys	Asp	
	50					55					60					

ttt	aat	gat	ctt	acc	ttg	ttt	gca	ccg	aca	cgt	gat	aat	act	ctt	att	240
Phe	Asn	Asp	Leu	Thr	Leu	Phe	Ala	Pro	Thr	Arg	Asp	Asn	Thr	Leu	Ile	
65					70					75					80	

tat	gac	caa	caa	aca	gca	aat	aca	cta	gta	gat	caa	aag	cat	caa	gaa	288
Tyr	Asp	Gln	Gln	Thr	Ala	Asn	Thr	Leu	Val	Asp	Gln	Lys	His	Gln	Glu	
				85				90						95		

tat	cat	tct	att	cgc	tgg	att	gga	ttg	att	cag	agt	agt	gca	aca	gga	336
Tyr	His	Ser	Ile	Arg	Trp	Ile	Gly	Leu	Ile	Gln	Ser	Ser	Ala	Thr	Gly	
			100					105							110	

gat ttc aca ttt aaa ttg tca gat gat gaa aat gcc atc att gaa ttg Asp Phe Thr Phe Lys Leu Ser Asp Asp Glu Asn Ala Ile Ile Glu Leu 115 120 125	384
gat ggg aaa gtt att tct gaa aaa ggt aac aat aaa caa agt gtt cat Asp Gly Lys Val Ile Ser Glu Lys Gly Asn Asn Lys Gln Ser Val His 130 135 140	432
tta gaa aaa gga cag ttg gtg caa ata aaa att gag tac caa tca gac Leu Glu Lys Gly Gln Leu Val Gln Ile Lys Ile Glu Tyr Gln Ser Asp 145 150 155 160	480
gat gca tta cat ata gat aat aaa att ttt aaa gag ctt aag cta ttc Asp Ala Leu His Ile Asp Asn Lys Ile Phe Lys Glu Leu Lys Leu Phe 165 170 175	528
aag ata gat agt caa aat cac tct caa caa gtt caa caa gat gaa ctg Lys Ile Asp Ser Gln Asn His Ser Gln Gln Val Gln Gln Asp Glu Leu 180 185 190	576
aga aac cct gag ttt aat aag aaa gaa acg caa gta ttc tta aag aaa Arg Asn Pro Glu Phe Asn Lys Lys Glu Thr Gln Val Phe Leu Lys Lys 195 200 205	624
gca tcg aaa aca aat ctt ttt aca caa aaa aca aaa aga gac att gat Ala Ser Lys Thr Asn Leu Phe Thr Gln Lys Thr Lys Arg Asp Ile Asp 210 215 220	672
gaa gat acg gat aca gat gga gat tct atc cct gat gtt tgg gaa gaa Glu Asp Thr Asp Thr Asp Gly Asp Ser Ile Pro Asp Val Trp Glu Glu 225 230 235 240	720
aac ggg tat acc att caa aac aaa gtc gca gtc aaa tgg gat gat tcg Asn Gly Tyr Thr Ile Gln Asn Lys Val Ala Val Lys Trp Asp Asp Ser 245 250 255	768
tta gca agt aaa ggg tat caa aaa ttt act tct aat cca cta gaa gca Leu Ala Ser Lys Gly Tyr Gln Lys Phe Thr Ser Asn Pro Leu Glu Ala 260 265 270	816
cac aca gtt gga gat ccc tat agt gat tat gaa aaa gct gca aga gat His Thr Val Gly Asp Pro Tyr Ser Asp Tyr Glu Lys Ala Ala Arg Asp 275 280 285	864
atg ccc tta tcg aat gca aaa gaa act ttt aat cct ctg gtt gcc gcc Met Pro Leu Ser Asn Ala Lys Glu Thr Phe Asn Pro Leu Val Ala Ala 290 295 300	912
ttt cca tca gta aat gtt agt tta gaa aag gtg att tta tcc aaa aat Phe Pro Ser Val Asn Val Ser Leu Glu Lys Val Ile Leu Ser Lys Asn 305 310 315 320	960
gaa gac ctt tcc cat agc gtt gaa agc agt caa tct acc aat tgg tct Glu Asp Leu Ser His Ser Val Glu Ser Ser Gln Ser Thr Asn Trp Ser 325 330 335	1008

tat acc aat act gaa ggc gtt aac gtc aat gcc gga tgg tca ggc tta Tyr Thr Asn Thr Glu Gly Val Asn Val Asn Ala Gly Trp Ser Gly Leu 340 345 350	1056
gga cct agt ttt gga gtt tct gtt aac tat caa cat agt gaa act gta Gly Pro Ser Phe Gly Val Ser Val Asn Tyr Gln His Ser Glu Thr Val 355 360 365	1104
gca aat gaa tgg ggt tct gcg acg aat gat ggc aca cat ata aat gga Ala Asn Glu Trp Gly Ser Ala Thr Asn Asp Gly Thr His Ile Asn Gly 370 375 380	1152
gcg gaa tct gct tat tta aac gca aat gtt cgc tat aat aac gtt ggg Ala Glu Ser Ala Tyr Leu Asn Ala Asn Val Arg Tyr Asn Asn Val Gly 385 390 395 400	1200
aca gga gca att tat gaa acg aaa cca aca acg agt ttt att ctt gat Thr Gly Ala Ile Tyr Glu Thr Lys Pro Thr Thr Ser Phe Ile Leu Asp 405 410 415	1248
gga aca aca att gga acg att aaa gca aaa gaa aat aca aca gct tta Gly Thr Thr Ile Gly Thr Ile Lys Ala Lys Glu Asn Thr Thr Ala Leu 420 425 430	1296
act att tta ccg gac caa agc tat cca gag aaa ggg aaa aac gga atc Thr Ile Leu Pro Asp Gln Ser Tyr Pro Glu Lys Gly Lys Asn Gly Ile 435 440 445	1344
gca att aac aca atg gat gat ttt aac tct cgc cca att cca tta aat Ala Ile Asn Thr Met Asp Asp Phe Asn Ser Arg Pro Ile Pro Leu Asn 450 455 460	1392
aaa gag caa cta aat act tat tta tct aat aaa aaa cca atc cta ctt Lys Glu Gln Leu Asn Thr Tyr Leu Ser Asn Lys Lys Pro Ile Leu Leu 465 470 475 480	1440
gaa aca gat caa gta gaa gga aaa tac gcc ata aag gat acc aat ggg Glu Thr Asp Gln Val Glu Gly Lys Tyr Ala Ile Lys Asp Thr Asn Gly 485 490 495	1488
aat att aca ata gct gga gat tgg aat ggt ata aca gat gaa att tct Asn Ile Thr Ile Ala Gly Asp Trp Asn Gly Ile Thr Asp Glu Ile Ser 500 505 510	1536
gct aaa acg gcc tct att att gta gat aat gga aat caa atg tca gaa Ala Lys Thr Ala Ser Ile Ile Val Asp Asn Gly Asn Gln Met Ser Glu 515 520 525	1584
aag aga gtt gca gcg aag gat tat aca aat cca gag gat aaa act cct Lys Arg Val Ala Ala Lys Asp Tyr Thr Asn Pro Glu Asp Lys Thr Pro 530 535 540	1632
aat tta tct gta aaa gaa gct cta aag tta gct tat cca gat gaa att Asn Leu Ser Val Lys Glu Ala Leu Lys Leu Ala Tyr Pro Asp Glu Ile 545 550 555 560	1680

gag gaa aaa gat ggt tta tta ttt tat aat gac caa cct att ttt gaa	1728
Glu Glu Lys Asp Gly Leu Leu Phe Tyr Asn Asp Gln Pro Ile Phe Glu	
565 570 575	
gca tct gta caa agt tat gtt gac gaa tat aca gct aaa caa att aga	1776
Ala Ser Val Gln Ser Tyr Val Asp Glu Tyr Thr Ala Lys Gln Ile Arg	
580 585 590	
aaa cag tta aat gat agt act ggt agc ttc aaa gat gtt aag aat tta	1824
Lys Gln Leu Asn Asp Ser Thr Gly Ser Phe Lys Asp Val Lys Asn Leu	
595 600 605	
tat gat gtg aaa tta gaa ccc aaa atg aat ttc aca ata aaa act agc	1872
Tyr Asp Val Lys Leu Glu Pro Lys Met Asn Phe Thr Ile Lys Thr Ser	
610 615 620	
act tta tat gat gga gga gaa tct gac aac aca aaa ata gga aat tgg	1920
Thr Leu Tyr Asp Gly Gly Glu Ser Asp Asn Thr Lys Ile Gly Asn Trp	
625 630 635 640	
tac tat act tat gtt gtc aac gga gga aat acg ggt aaa aaa caa tac	1968
Tyr Tyr Thr Tyr Val Val Asn Gly Gly Asn Thr Gly Lys Lys Gln Tyr	
645 650 655	
cgt tca gct aat aaa ggt gcc ttt act gag ctg tca aca gaa tca aag	2016
Arg Ser Ala Asn Lys Gly Ala Phe Thr Glu Leu Ser Thr Glu Ser Lys	
660 665 670	
aat aaa ttg aaa aaa aat ata gat tac tac gta agc cta tat atg aag	2064
Asn Lys Leu Lys Lys Asn Ile Asp Tyr Tyr Val Ser Leu Tyr Met Lys	
675 680 685	
gct gac tca aag gtt tca gtt gat ata gaa ata gac gga aaa cag gag	2112
Ala Asp Ser Lys Val Ser Val Asp Ile Glu Ile Asp Gly Lys Gln Glu	
690 695 700	
tca att gta aca gat aat ata acc tta gat cac gta ggt tac caa aga	2160
Ser Ile Val Thr Asp Asn Ile Thr Leu Asp His Val Gly Tyr Gln Arg	
705 710 715 720	
ata aac atc cta gtc ccc aat ctg gaa gga aac gaa ata aat act att	2208
Ile Asn Ile Leu Val Pro Asn Leu Glu Gly Asn Glu Ile Asn Thr Ile	
725 730 735	
tct att aaa ggt gac gga caa acc aat gtt tat tgg gat gat gtc tcc	2256
Ser Ile Lys Gly Asp Gly Gln Thr Asn Val Tyr Trp Asp Asp Val Ser	
740 745 750	
ttt gtc gaa gtg gga gca gaa gaa att gaa tat aaa gat cca gtt ccc	2304
Phe Val Glu Val Gly Ala Glu Glu Ile Glu Tyr Lys Asp Pro Val Pro	
755 760 765	
caa ttt gac att ata gaa gga gat ttt gat ttc ttt ggt gat cca ttg	2352
Gln Phe Asp Ile Ile Glu Gly Asp Phe Asp Phe Phe Gly Asp Pro Leu	
770 775 780	

gcg gta aaa tat cat gat gca acg tat ttt ata gat agt cct ttg att 2400
 Ala Val Lys Tyr His Asp Ala Thr Tyr Phe Ile Asp Ser Pro Leu Ile
 785 790 795 800

aca caa act cct gga act ttc tcc ttt act tat aaa gtg att ggg gaa 2448
 Thr Gln Thr Pro Gly Thr Phe Ser Phe Thr Tyr Lys Val Ile Gly Glu
 805 810 815

caa acg aag aca gta tta gat tcg gga tct ggt aaa aac gca aat cga 2496
 Gln Thr Lys Thr Val Leu Asp Ser Gly Ser Gly Lys Asn Ala Asn Arg
 820 825 830

atc aac cta gat ttt aaa aat gta aaa tca gat cgt tca ttc tta tat 2544
 Ile Asn Leu Asp Phe Lys Asn Val Lys Ser Asp Arg Ser Phe Leu Tyr
 835 840 845

aca tta tca tgt aaa gat gat tta tgg gga agc act cgc aca gca gtt 2592
 Thr Leu Ser Cys Lys Asp Asp Leu Trp Gly Ser Thr Arg Thr Ala Val
 850 855 860

gtt aga att ttt gct gta gat taa 2616
 Val Arg Ile Phe Ala Val Asp
 865 870

<210> 2
 <211> 871
 <212> PRT
 <213> Brevibacillus laterosporus

<400> 2
 Met Lys Tyr Met Lys Lys Gly Leu Ser Ser Val Val Ile Gly Thr Leu
 1 5 10 15
 Phe Ala Ser Met Phe Leu Asn Gly Asn Val Asn Ala Val Tyr Ala Asn
 20 25 30
 Ser Lys Thr Asn Gln Ile Ala Thr Thr Thr Gln Ala Ser Lys Asp Asn
 35 40 45
 Gln Ile Asp Arg Glu Gly Leu Leu Gly Tyr Tyr Phe Lys Gly Lys Asp
 50 55 60
 Phe Asn Asp Leu Thr Leu Phe Ala Pro Thr Arg Asp Asn Thr Leu Ile
 65 70 75 80
 Tyr Asp Gln Gln Thr Ala Asn Thr Leu Val Asp Gln Lys His Gln Glu
 85 90 95
 Tyr His Ser Ile Arg Trp Ile Gly Leu Ile Gln Ser Ser Ala Thr Gly
 100 105 110
 Asp Phe Thr Phe Lys Leu Ser Asp Asp Glu Asn Ala Ile Ile Glu Leu
 115 120 125
 Asp Gly Lys Val Ile Ser Glu Lys Gly Asn Asn Lys Gln Ser Val His
 130 135 140
 Leu Glu Lys Gly Gln Leu Val Gln Ile Lys Ile Glu Tyr Gln Ser Asp
 145 150 155 160
 Asp Ala Leu His Ile Asp Asn Lys Ile Phe Lys Glu Leu Lys Leu Phe
 165 170 175
 Lys Ile Asp Ser Gln Asn His Ser Gln Gln Val Gln Gln Asp Glu Leu
 180 185 190
 Arg Asn Pro Glu Phe Asn Lys Lys Glu Thr Gln Val Phe Leu Lys Lys
 195 200 205

Ala	Ser	Lys	Thr	Asn	Leu	Phe	Thr	Gln	Lys	Thr	Lys	Arg	Asp	Ile	Asp
210						215					220				
Glu	Asp	Thr	Asp	Thr	Asp	Gly	Asp	Ser	Ile	Pro	Asp	Val	Trp	Glu	Glu
225					230					235					240
Asn	Gly	Tyr	Thr	Ile	Gln	Asn	Lys	Val	Ala	Val	Lys	Trp	Asp	Asp	Ser
				245					250					255	
Leu	Ala	Ser	Lys	Gly	Tyr	Gln	Lys	Phe	Thr	Ser	Asn	Pro	Leu	Glu	Ala
			260					265					270		
His	Thr	Val	Gly	Asp	Pro	Tyr	Ser	Asp	Tyr	Glu	Lys	Ala	Ala	Arg	Asp
		275					280					285			
Met	Pro	Leu	Ser	Asn	Ala	Lys	Glu	Thr	Phe	Asn	Pro	Leu	Val	Ala	Ala
	290					295					300				
Phe	Pro	Ser	Val	Asn	Val	Ser	Leu	Glu	Lys	Val	Ile	Leu	Ser	Lys	Asn
305					310					315					320
Glu	Asp	Leu	Ser	His	Ser	Val	Glu	Ser	Ser	Gln	Ser	Thr	Asn	Trp	Ser
				325					330					335	
Tyr	Thr	Asn	Thr	Glu	Gly	Val	Asn	Val	Asn	Ala	Gly	Trp	Ser	Gly	Leu
			340					345						350	
Gly	Pro	Ser	Phe	Gly	Val	Ser	Val	Asn	Tyr	Gln	His	Ser	Glu	Thr	Val
		355					360					365			
Ala	Asn	Glu	Trp	Gly	Ser	Ala	Thr	Asn	Asp	Gly	Thr	His	Ile	Asn	Gly
	370					375					380				
Ala	Glu	Ser	Ala	Tyr	Leu	Asn	Ala	Asn	Val	Arg	Tyr	Asn	Asn	Val	Gly
385					390					395					400
Thr	Gly	Ala	Ile	Tyr	Glu	Thr	Lys	Pro	Thr	Thr	Ser	Phe	Ile	Leu	Asp
				405					410					415	
Gly	Thr	Thr	Ile	Gly	Thr	Ile	Lys	Ala	Lys	Glu	Asn	Thr	Thr	Ala	Leu
			420					425						430	
Thr	Ile	Leu	Pro	Asp	Gln	Ser	Tyr	Pro	Glu	Lys	Gly	Lys	Asn	Gly	Ile
		435					440					445			
Ala	Ile	Asn	Thr	Met	Asp	Asp	Phe	Asn	Ser	Arg	Pro	Ile	Pro	Leu	Asn
	450					455					460				
Lys	Glu	Gln	Leu	Asn	Thr	Tyr	Leu	Ser	Asn	Lys	Lys	Pro	Ile	Leu	Leu
465					470					475					480
Glu	Thr	Asp	Gln	Val	Glu	Gly	Lys	Tyr	Ala	Ile	Lys	Asp	Thr	Asn	Gly
				485					490					495	
Asn	Ile	Thr	Ile	Ala	Gly	Asp	Trp	Asn	Gly	Ile	Thr	Asp	Glu	Ile	Ser
			500					505					510		
Ala	Lys	Thr	Ala	Ser	Ile	Ile	Val	Asp	Asn	Gly	Asn	Gln	Met	Ser	Glu
		515					520					525			
Lys	Arg	Val	Ala	Ala	Lys	Asp	Tyr	Thr	Asn	Pro	Glu	Asp	Lys	Thr	Pro
	530					535					540				
Asn	Leu	Ser	Val	Lys	Glu	Ala	Leu	Lys	Leu	Ala	Tyr	Pro	Asp	Glu	Ile
545					550					555					560
Glu	Glu	Lys	Asp	Gly	Leu	Leu	Phe	Tyr	Asn	Asp	Gln	Pro	Ile	Phe	Glu
				565					570					575	
Ala	Ser	Val	Gln	Ser	Tyr	Val	Asp	Glu	Tyr	Thr	Ala	Lys	Gln	Ile	Arg
			580					585					590		
Lys	Gln	Leu	Asn	Asp	Ser	Thr	Gly	Ser	Phe	Lys	Asp	Val	Lys	Asn	Leu
		595					600					605			
Tyr	Asp	Val	Lys	Leu	Glu	Pro	Lys	Met	Asn	Phe	Thr	Ile	Lys	Thr	Ser
	610					615						620			
Thr	Leu	Tyr	Asp	Gly	Gly	Glu	Ser	Asp	Asn	Thr	Lys	Ile	Gly	Asn	Trp
625					630					635					640
Tyr	Tyr	Thr	Tyr	Val	Val	Asn	Gly	Gly	Asn	Thr	Gly	Lys	Lys	Gln	Tyr
				645					650					655	

Arg Ser Ala Asn Lys Gly Ala Phe Thr Glu Leu Ser Thr Glu Ser Lys
 660 665 670
 Asn Lys Leu Lys Lys Asn Ile Asp Tyr Tyr Val Ser Leu Tyr Met Lys
 675 680 685
 Ala Asp Ser Lys Val Ser Val Asp Ile Glu Ile Asp Gly Lys Gln Glu
 690 695 700
 Ser Ile Val Thr Asp Asn Ile Thr Leu Asp His Val Gly Tyr Gln Arg
 705 710 715 720
 Ile Asn Ile Leu Val Pro Asn Leu Glu Gly Asn Glu Ile Asn Thr Ile
 725 730 735
 Ser Ile Lys Gly Asp Gly Gln Thr Asn Val Tyr Trp Asp Asp Val Ser
 740 745 750
 Phe Val Glu Val Gly Ala Glu Glu Ile Glu Tyr Lys Asp Pro Val Pro
 755 760 765
 Gln Phe Asp Ile Ile Glu Gly Asp Phe Asp Phe Phe Gly Asp Pro Leu
 770 775 780
 Ala Val Lys Tyr His Asp Ala Thr Tyr Phe Ile Asp Ser Pro Leu Ile
 785 790 795 800
 Thr Gln Thr Pro Gly Thr Phe Ser Phe Thr Tyr Lys Val Ile Gly Glu
 805 810 815
 Gln Thr Lys Thr Val Leu Asp Ser Gly Ser Gly Lys Asn Ala Asn Arg
 820 825 830
 Ile Asn Leu Asp Phe Lys Asn Val Lys Ser Asp Arg Ser Phe Leu Tyr
 835 840 845
 Thr Leu Ser Cys Lys Asp Asp Leu Trp Gly Ser Thr Arg Thr Ala Val
 850 855 860
 Val Arg Ile Phe Ala Val Asp
 865 870

<210> 3
 <211> 1374
 <212> DNA
 <213> Brevibacillus laterosporus

<220>
 <221> CDS
 <222> (1)..(1371)

<400> 3
 atg att gtg att att ttc aca aac gta aaa gga ggg aat gag ttg aaa 48
 Met Ile Val Ile Ile Phe Thr Asn Val Lys Gly Gly Asn Glu Leu Lys
 1 5 10 15
 aag aat ttt tat aag aat ctt att tgt atg tct gct tta ttg tta gcc 96
 Lys Asn Phe Tyr Lys Asn Leu Ile Cys Met Ser Ala Leu Leu Leu Ala
 20 25 30
 atg cct ata tca agc aac gtt acg tac gct tac ggt agt gag aag gtt 144
 Met Pro Ile Ser Ser Asn Val Thr Tyr Ala Tyr Gly Ser Glu Lys Val
 35 40 45
 gat tat tta gta aaa acg act aac aat aca gag gat ttt aaa gag gat 192
 Asp Tyr Leu Val Lys Thr Thr Asn Asn Thr Glu Asp Phe Lys Glu Asp
 50 55 60

aag gaa aaa gcc aaa gaa tgg ggg aaa gaa aaa gag aaa gag tgg aaa Lys Glu Lys Ala Lys Glu Trp Gly Lys Glu Lys Glu Lys Glu Trp Lys 65 70 75 80	240
cta act gtt act gaa aaa aca agg atg aat aat ttt tta gat aat aaa Leu Thr Val Thr Glu Lys Thr Arg Met Asn Asn Phe Leu Asp Asn Lys 85 90 95	288
aat gat ata aaa aaa aat tat aaa gaa att act ttt tct atg gca ggt Asn Asp Ile Lys Lys Asn Tyr Lys Glu Ile Thr Phe Ser Met Ala Gly 100 105 110	336
tca ttt gaa gat gaa ata aaa gat tta aaa gag att gat aaa atg ttt Ser Phe Glu Asp Glu Ile Lys Asp Leu Lys Glu Ile Asp Lys Met Phe 115 120 125	384
gat aaa gcc aat cta tca agt tct att gtc acc tat aaa aat gtg gag Asp Lys Ala Asn Leu Ser Ser Ser Ile Val Thr Tyr Lys Asn Val Glu 130 135 140	432
ccc tca acg att gga ttt aac aaa cct tta aca gaa ggt aat act att Pro Ser Thr Ile Gly Phe Asn Lys Pro Leu Thr Glu Gly Asn Thr Ile 145 150 155 160	480
aat act gat gta caa gcc cag ttt aaa gaa caa ttt tta gga aaa gat Asn Thr Asp Val Gln Ala Gln Phe Lys Glu Gln Phe Leu Gly Lys Asp 165 170 175	528
att aag ttt gat agt tat ctt gac act cac tta act gca caa aat gtt Ile Lys Phe Asp Ser Tyr Leu Asp Thr His Leu Thr Ala Gln Asn Val 180 185 190	576
tct agt aaa gaa aga att att tta caa gtt aca gtg cca agt ggt aaa Ser Ser Lys Glu Arg Ile Ile Leu Gln Val Thr Val Pro Ser Gly Lys 195 200 205	624
gga tct act att cca aca aaa gca ggt gta att tta aat aat aat gag Gly Ser Thr Ile Pro Thr Lys Ala Gly Val Ile Leu Asn Asn Asn Glu 210 215 220	672
tat aaa atg cta att gat aat ggc tat gta ctc cat gtg gat aat ata Tyr Lys Met Leu Ile Asp Asn Gly Tyr Val Leu His Val Asp Asn Ile 225 230 235 240	720
tcg aaa gta gta aaa aaa ggt tat gaa tgt tta caa att caa gga acg Ser Lys Val Val Lys Lys Gly Tyr Glu Cys Leu Gln Ile Gln Gly Thr 245 250 255	768
cta aaa aag agt ctc gat ttt aaa aat gat att aat gct gag gct cat Leu Lys Lys Ser Leu Asp Phe Lys Asn Asp Ile Asn Ala Glu Ala His 260 265 270	816
cgt tgg ggt atg aaa aat tat gaa gga tgg gct aaa aat tta aca gat Arg Trp Gly Met Lys Asn Tyr Glu Gly Trp Ala Lys Asn Leu Thr Asp 275 280 285	864

cct caa agg gaa gct tta gat ggg tat gct aga caa gat tat aaa caa 912
 Pro Gln Arg Glu Ala Leu Asp Gly Tyr Ala Arg Gln Asp Tyr Lys Gln
 290 295 300

ata aat gat tat tta cga aat caa ggt gga agt gga aat gaa aaa cta 960
 Ile Asn Asp Tyr Leu Arg Asn Gln Gly Gly Ser Gly Asn Glu Lys Leu
 305 310 315 320

gat aca caa ata aaa aat att tct gaa gca tta gaa aag cag cca ata 1008
 Asp Thr Gln Ile Lys Asn Ile Ser Glu Ala Leu Glu Lys Gln Pro Ile
 325 330 335

cca gaa aat att act gtg tat aga tgg tgt gga atg gcg gaa ttt ggt 1056
 Pro Glu Asn Ile Thr Val Tyr Arg Trp Cys Gly Met Ala Glu Phe Gly
 340 345 350

tat caa att agt gat cct tta cct tct tta aaa gaa atg gaa gaa aaa 1104
 Tyr Gln Ile Ser Asp Pro Leu Pro Ser Leu Lys Glu Met Glu Glu Lys
 355 360 365

ttt tta aat aca atg aaa gaa gat aag gga tat atg agt act agt ttg 1152
 Phe Leu Asn Thr Met Lys Glu Asp Lys Gly Tyr Met Ser Thr Ser Leu
 370 375 380

tca agt gaa cgt ctt tct gca ttt ggt tcg aga aaa ttc att tta aga 1200
 Ser Ser Glu Arg Leu Ser Ala Phe Gly Ser Arg Lys Phe Ile Leu Arg
 385 390 395 400

tta caa gtt cct aaa gga agc aca ggg gca tat tta agc gct att ggg 1248
 Leu Gln Val Pro Lys Gly Ser Thr Gly Ala Tyr Leu Ser Ala Ile Gly
 405 410 415

gga ttt gca agt gaa aaa gaa atc ctt att gat aaa gat agt aac tat 1296
 Gly Phe Ala Ser Glu Lys Glu Ile Leu Ile Asp Lys Asp Ser Asn Tyr
 420 425 430

cat att gat aaa ata aca gag gta gtc att aaa ggt gtt aag cga tat 1344
 His Ile Asp Lys Ile Thr Glu Val Val Ile Lys Gly Val Lys Arg Tyr
 435 440 445

gta gtt gat gca acg tta tta aca aaa taa 1374
 Val Val Asp Ala Thr Leu Leu Thr Lys
 450 455

<210> 4
 <211> 457
 <212> PRT
 <213> Brevibacillus laterosporus

<400> 4
 Met Ile Val Ile Ile Phe Thr Asn Val Lys Gly Gly Asn Glu Leu Lys
 1 5 10 15
 Lys Asn Phe Tyr Lys Asn Leu Ile Cys Met Ser Ala Leu Leu Leu Ala
 20 25 30
 Met Pro Ile Ser Ser Asn Val Thr Tyr Ala Tyr Gly Ser Glu Lys Val
 35 40 45

Asp	Tyr	Leu	Val	Lys	Thr	Thr	Asn	Asn	Thr	Glu	Asp	Phe	Lys	Glu	Asp
50						55					60				
Lys	Glu	Lys	Ala	Lys	Glu	Trp	Gly	Lys	Glu	Lys	Glu	Lys	Glu	Trp	Lys
65					70					75					80
Leu	Thr	Val	Thr	Glu	Lys	Thr	Arg	Met	Asn	Asn	Phe	Leu	Asp	Asn	Lys
				85					90					95	
Asn	Asp	Ile	Lys	Lys	Asn	Tyr	Lys	Glu	Ile	Thr	Phe	Ser	Met	Ala	Gly
			100					105					110		
Ser	Phe	Glu	Asp	Glu	Ile	Lys	Asp	Leu	Lys	Glu	Ile	Asp	Lys	Met	Phe
		115					120					125			
Asp	Lys	Ala	Asn	Leu	Ser	Ser	Ser	Ile	Val	Thr	Tyr	Lys	Asn	Val	Glu
130						135					140				
Pro	Ser	Thr	Ile	Gly	Phe	Asn	Lys	Pro	Leu	Thr	Glu	Gly	Asn	Thr	Ile
145					150						155				160
Asn	Thr	Asp	Val	Gln	Ala	Gln	Phe	Lys	Glu	Gln	Phe	Leu	Gly	Lys	Asp
			165						170					175	
Ile	Lys	Phe	Asp	Ser	Tyr	Leu	Asp	Thr	His	Leu	Thr	Ala	Gln	Asn	Val
			180					185					190		
Ser	Ser	Lys	Glu	Arg	Ile	Ile	Leu	Gln	Val	Thr	Val	Pro	Ser	Gly	Lys
		195					200					205			
Gly	Ser	Thr	Ile	Pro	Thr	Lys	Ala	Gly	Val	Ile	Leu	Asn	Asn	Asn	Glu
210						215					220				
Tyr	Lys	Met	Leu	Ile	Asp	Asn	Gly	Tyr	Val	Leu	His	Val	Asp	Asn	Ile
225					230					235					240
Ser	Lys	Val	Val	Lys	Lys	Gly	Tyr	Glu	Cys	Leu	Gln	Ile	Gln	Gly	Thr
			245					250						255	
Leu	Lys	Lys	Ser	Leu	Asp	Phe	Lys	Asn	Asp	Ile	Asn	Ala	Glu	Ala	His
			260					265					270		
Arg	Trp	Gly	Met	Lys	Asn	Tyr	Glu	Gly	Trp	Ala	Lys	Asn	Leu	Thr	Asp
		275					280					285			
Pro	Gln	Arg	Glu	Ala	Leu	Asp	Gly	Tyr	Ala	Arg	Gln	Asp	Tyr	Lys	Gln
290						295					300				
Ile	Asn	Asp	Tyr	Leu	Arg	Asn	Gln	Gly	Gly	Ser	Gly	Asn	Glu	Lys	Leu
305					310					315					320
Asp	Thr	Gln	Ile	Lys	Asn	Ile	Ser	Glu	Ala	Leu	Glu	Lys	Gln	Pro	Ile
			325						330					335	
Pro	Glu	Asn	Ile	Thr	Val	Tyr	Arg	Trp	Cys	Gly	Met	Ala	Glu	Phe	Gly
			340					345					350		
Tyr	Gln	Ile	Ser	Asp	Pro	Leu	Pro	Ser	Leu	Lys	Glu	Met	Glu	Glu	Lys
		355					360					365			
Phe	Leu	Asn	Thr	Met	Lys	Glu	Asp	Lys	Gly	Tyr	Met	Ser	Thr	Ser	Leu
370						375					380				
Ser	Ser	Glu	Arg	Leu	Ser	Ala	Phe	Gly	Ser	Arg	Lys	Phe	Ile	Leu	Arg
385					390					395					400
Leu	Gln	Val	Pro	Lys	Gly	Ser	Thr	Gly	Ala	Tyr	Leu	Ser	Ala	Ile	Gly
			405						410					415	
Gly	Phe	Ala	Ser	Glu	Lys	Glu	Ile	Leu	Ile	Asp	Lys	Asp	Ser	Asn	Tyr
			420					425					430		
His	Ile	Asp	Lys	Ile	Thr	Glu	Val	Val	Ile	Lys	Gly	Val	Lys	Arg	Tyr
		435					440					445			
Val	Val	Asp	Ala	Thr	Leu	Leu	Thr	Lys							
450						455									

<210> 5

<211> 2658

<212> DNA
 <213> Zea mays

<220>
 <221> misc_feature
 <222> (21)..()
 <223> Nucleotide at position 21 is "n" wherein "n" = any nucleotide

<400> 5
 cctaaggccg atttcgtcct naggcaggcc caaaggaagg aagtacttca gtggatcaag 60
 atgttgatgt tccctgatgg gtatgcagct aacctgagta ggtgggggtga acttatctac 120
 tctgtgagtc ttagggatga agagtcatga ctccacata tggattgaac agattcttct 180
 ctgtgcatgg acaatctggg gcggcatcca acaacctca tggatcgccc ggccaatcgc 240
 cgcaccagtc catccgccc cctcgatgag acttatgttc ttagtggtga gacttcagaa 300
 cttattgata atgctgtatt ggatacttat gtttggttgc gatacttatg tgagaacttg 360
 agacttatga gacttatgtt cttgatactt atgtttgtgt tgagaacttg gatatttatg 420
 tttgtgttgg atacttatgt ctgtgatgat atatgtgatg tatatatgtg atgtatatgt 480
 gacatatgtg atgtatatgt ggtatctttt gtttggttgg atggaataga gaaagcaaag 540
 aaaaatgtgt atactggtca ctttgctgag tgtaacactc ggcaaaaagg tgctttgccg 600
 agtggttaggg ccatagcact cggtagagaa ccaatactta ggcaccggta aagctttttt 660
 ggcgagtgtt gtggccctgg cactcagctt tgccgagtgc ctacagagc actcgacaaa 720
 gaacctgaca aatggaccgg ctggtaaact ctttaccgag tgcaggtcag tagacactcg 780
 gcaaaggtaa cttctttgcc gagtgccgct tagaacattt gacaaagggg catctccgtt 840
 acccgggtgc gtgacggccg cttttctttg ccgagtgcct gatagaaagt actcggcaaa 900
 gaagtcgttg ccaatgtatt gtccgctgag gtctctttgt caagtattac actcggcaaa 960
 gactgtgccg agtggttttc agactttgcc gagtggttta agcactcagc aaagcgctcg 1020
 atttcggtag tgacggttgt ttggcaatag taaaatccag ccctctcccg tggggaaaaa 1080
 actggttaga tctggctcgt ggctaagatt ctctttcttc ctttgtaaa aaaagagaag 1140
 aaaaaaaaaa cgactgtcac ggtgccttgt ctggtaatga tcgcgcggtc ggctctgtcc 1200
 taaccgtaaa gatggacggg agctgatgat agcgtgacct ccaataaac aacaaggcg 1260
 tgttccccgc ggtcgaatat tttaagggcc actgattagg tgcgggtgaa tacatcaact 1320
 tcacgaacat catctgatct gatctgattt ggtctgatat gatctgggta gtcatttctg 1380
 caatgagcat ctatcagggtg aaccaattaa tattgatgac attatgagtt cgaagatata 1440
 ctctaaagtg ttatctaaat acagaagaca ttcgttcggt ctttgccctat aactctaaaa 1500
 ggcttgtaac accctcattc atcctctata tacgaagact ctctcctatc atttttatcg 1560
 atttattttt tttatattta gacaatggaa ttaaatagaa ctaaaatata tataagatga 1620
 tatctgagga cccgagatgg taatggggac tcgatcctcg attctccacg gagaattcct 1680
 ctaggatata ggtaatttgt cccacagagg attgaaacgg ggtaatttgg tccccatgtg 1740
 cccgtccgcg gaacttctct tgatctaaat tagtctatct ccatgttaaa actatactaa 1800
 aaattttaata cacagtctat tataaaatag caaactaaat tctaaagttg atgcatcttg 1860
 taattttaaa tctggtttgt tcaagttata ttcatttgat ataataaatt tgaatttgac 1920
 tcttaatatc gtattttttc ctaacgggga cggattctcc acggggataa attccatgat 1980
 acagatggga tgaaagaaaa atctcccgtg tgaacttttg caggaatggg gatgggccag 2040
 agaaattttc tccctgcggg gacggggggag ccatatcctc ggtggagaat ttcccattat 2100
 catccttatt tgtggtacat atatatgcat aatctttttt ttttgactga catgtgggaa 2160
 agtatcccat ctcaatagta gaaaatcttg ggaacggtag gatcgaacac aaagatcagc 2220
 tagcttgtaa tcaccgagcc atatagctag agggtaatag atcatgaatc aaatgttttt 2280
 ttcataaatt attaaggctc taaattatct ttaatttaaa aataaataaa aatatagttc 2340
 gattcttaca ttttatagtg taaaacttta aagtctatta ttaccctac ttattgagtt 2400
 atggttcagt tcttgctgac ggagagtaat gagatataga ataaggtagc ctatagaata 2460
 aagaatcttt ctctgaaaag tctgacgtac gtaaataaga tataataaaa aaaatacaaa 2520
 gagaagcgct ggactggaga tgctcctata tgccggcaatg cctgtgctta taaatagcca 2580
 cctcggtcgg caaggacatg aacggcggac gcagtgtgca tgcatacaag agcaacaaga 2640
 tactggcgca gaggagca 2658

<210> 6
 <211> 4325
 <212> DNA
 <213> Zea Mays

<220>
 <221> misc_feature
 <222> (372)..(1688)
 <223> Nucleotides at positions 372, 525, 586, 714, 737, 747, 754, 785,
 and 1688 are "n" wherein "n" = any nucleotid

<400> 6
 gcggccgcgg aattaaccct cactaaaggg aacgaattcg gatccctgtg gagaaatttt 60
 tacgtcgcg g gatgggatg gggagttatt cccctgtagg aaatgggtga cgcctaagag 120
 ggaggggtgaa gtaggacttc taaaactttc actaaactag gccacaaata attccctaga 180
 gcaaaaccta tgcaaatagt caaactagaa tgtgcaaacc aagttttgtc taagtgttgc 240
 tatctctacc gcaatggcta agtttcaatc tacactatat aagtatgaat acaagaatga 300
 aacttaaata cttaataata atgcggaaac ttaaagagca aggtagagat gcaaattctc 360
 gtggatgacg cntgcatttt tatcgaggta tccggaacca cgcaagggtc cgactaatcc 420
 tcattgggtgc ccctacgcaa agggaagccc acgcgagggc caagcacctc ggtagagtaa 480
 ctcttataga gagccgtggg ctttctccac gcgcaagtgg tgctntgctt tcagctctc 540
 tcagaccctc cccgtgtgtc ccactatcga gcttccggct gaaaangcca tgggcctcgt 600
 tccctccggg acacgggtggc ggccgtgaca caaatgcggg tatcacgggt ctcgcaagac 660
 tctcaccccc actggggaca aatttcaatg gctcgcacaa gagccgaggg gttnatgggt 720
 tatctaattc cactcanctt aactagnatt catntaaagc aagcgctaga gcggtctaac 780
 taacntaagc acttcacaaa gcacctacgc ttaatcaccg agtgattcta tttagcactt 840
 ggggtgcaaga gcacttgaga atgtctacta tatgccttgc tatgtctctt gggctcccaa 900
 acttggaat ggccgggttg tggtgtattt atagcccca acacaaaact agccgttgga 960
 ggaagctgct gctttttgtg gtgcaccgga cagtcgggtg gggtcaccag acagtccgac 1020
 gccctgtcc ggtgccctg tccgatgcgc ctagtgtgtg ggtctgtcag cgtaggtgac 1080
 cgttgccgcg caggcttttt gcaccggaca gtccgggtgg ctccctcga cagtgccacc 1140
 tggagctagc cgttagggct actgttctg gtgcaccgga cagtagtcg gtgtcttgt 1200
 ttggacagtc cgacttgtgg caacacttct tcttttcttg gactttactt gatcttcatg 1260
 atgtctctt ttgaggtgtt gctttcctaa gtgccttggg ccaagtaact tatcatcctg 1320
 tgaactacaa acacaaatag tagcaaacac attagtcac aggttatgtt gatcatcaaa 1380
 taccaaaatc tattaagcca aatggcccag ggtccatttt ccttacatcc ccgacgaaga 1440
 attctccgtt gccatcccta tctgtgtacg cactactgga atccgggtct ttgctgagta 1500
 ccgcactcgg caaagtccta ctctcggtaa cgatgccttt tgccgagagc aggactctcg 1560
 gcacaggaat aactcggcg aaggcggggt ctccgcaaag gccgttagcc accgtccaaa 1620
 gctgacggtc gttacctatg ccgagtggtg gaaagatatt gtgaaggcct aaggccgatt 1680
 tcgtcctnag cagggcccaa aggaaggaag tacttcagtg gatcaagatg ttgatgttcc 1740
 ctgatgggta tgcagctaac ctgagtaggt ggggtgaact tatctactct gtgagcttta 1800
 gggatgaaga gtcatgactt ccacatatgg attgaacaga ttcttctctg tgcattggaca 1860
 atctggggcg gcatccaaca acctcatgg atcggccggc caatcgccgc accagtccat 1920
 ccgccacct cgatgagact tatgttctta gtgttgagac ttcagaactt attgataatg 1980
 ctgtattgga tacttatgtt tgtgttcgat acttatgtga gaacttgaga cttatgagac 2040
 ttatgttctt gatacttatg tttgtgttga gaacttgat atttatgttt gtgttgata 2100
 cttatgtctg tgatgatata tgtgatgtat atatgtgat tatatgtgac atatgtgatg 2160
 tatatgtgg atcttttgtt tgtttggat gaatagagaa agcaaataaa aatgtgtata 2220
 ctggtcactt tgtcgagtgt aacactcggc aaaaagggtc tttgccgagt gttagggcca 2280
 tagcactcgg tagagaacca atacttaggc accggtaaag cttttttggc gagtgttgtg 2340
 gccctggcac tcagctttgc cgagtgcctc acagagcact cgacaaagaa cctgacaaat 2400
 ggaccgcgtg gtaaatacct taccgagtgc aggtcagtag aactcggca aaggtaactt 2460
 ctttgccgag tgccgcttag aacatttgac aaagggtcat ctccgttacc cgggtgtcgtg 2520
 acggccgctt ttctttgccg agtgctgat agaaagtact cggcaaagaa gtcgttgcca 2580
 atgtattgtt cgtgaggtc tctttgtcaa gtattacact cggcaaagac tgtgccgagt 2640
 gtttttcaga ctttgccgag tggtttaagc actcagcaaa gcgtctgatt tcggtagtag 2700

```

cggttgtttg gcaatagtaa aatccagccc tctcccgtgg ggaaaaaact ggtaggatct 2760
ggctcgtggc taagattctc tttcttccct ttgtaaaaaa agagaagaaa aaaaaaacga 2820
ctgtcacggg gccttgtctg gtaatgatcg cgcggtcggc tctgtcctaa cccgtaagat 2880
ggacggggagc tgatgatagc gtgacctcca aataaacaac aagggcgtgt tccccgcggg 2940
cgaatatatt aagggccact gattagggtgc ggttgaatac atcaacttca cgaacatcat 3000
ctgatctgat ctgatttggt ctgatatgat ctgggtagtc atttctgcaa tgagcatcta 3060
tcagggtgaac caattaatat tgatgacatt atgagttcga agatatactc taaagtgtta 3120
tctaaataca gaagacattc gttcgttctt tgcctataac tctaaaaggc ttgtaacacc 3180
ctcattcatc ctctatatac gaagactctc tcctatcatt tttatcgatt tatttttttt 3240
atattttagac aatggaatta aatagaacta aaatatatat aagatgatat ctgaggaccc 3300
gagatggtaa tggggactcg atcctcgatt ctccacggag aattcctcta ggatataggt 3360
aatttgtccc cacgaggatt gaaacggggg aatttgggtc ccatgtgccc gtcccgcgaa 3420
cttctcttga tctaaattag tctatttcca tgttaaaact atactaaaaa tttaatacac 3480
agtctattat aaaatagcaa actaaattct aaagttgatg catcttgtaa ttttaaattc 3540
ggtttgttca agtttatatt atttgatata ataaatttga atttgactct taatatcgta 3600
ttttttccta acggggacgg attctccacg gggataaatt ccatgatata gatgggatga 3660
aagaaaaatc tcccgatga acttttgcag gaatggggat gggccagaga aattttctcc 3720
ctgcgggggac gggggagcca tatcctcggt ggagaatttc ccattatcat ccttatttgt 3780
ggtacatata tatgcataat cttttttttt tgactgacat gtgggaaagt atcccatctc 3840
aatagtagaa aatcttggga acggtaggat cgaacacaaa gatcagctag cttgtaatca 3900
ccgagccata tagctagagg gtaatagatc atgaatcaaa tgtttttttc ataaattatt 3960
aaggctctaa attattttta atttaaaaaa aaataaaaaa atagttcgat tcttacattt 4020
tatagtgtaa aacttttaaag tctattatta cccctactta ttgagttatg gttcagttct 4080
tgtcgacgga gagtaatgag atatagaata aggtacccta tagaataaag aatctttctc 4140
tgaaaagtct gacgtacgta aataagatat aataaaaaaa atacaaagag aagcgctgga 4200
ctggagatgc tcctatatgc ggcaatgcct gtgcttataa atagccacct cggtcggcaa 4260
ggacatgaac ggcggacgca gtgtgcatgc atacaagagc aacaagatac tggcgcagag 4320
gagca 4325

```

<210> 7

<211> 1239

<212> DNA

<213> Artificial Sequence

<220>

<223> modified DNA encoding ISP2A protein

<220>

<221> CDS

<222> (3)..(1229)

<400> 7

```

cc atg gca cta gtg aag acc acc aac aac acc gag gac ttc aag gag 47
Met Ala Leu Val Lys Thr Thr Asn Asn Thr Glu Asp Phe Lys Glu
1 5 10 15

gac aag gag aag gcc aag gag tgg ggc aag gag aag gag aag gag tgg 95
Asp Lys Glu Lys Ala Lys Glu Trp Gly Lys Glu Lys Glu Lys Glu Trp
20 25 30

aag ctg acc gtg acc gag aag acg cgt atg aat aat ttc ctg gac aac 143
Lys Leu Thr Val Thr Glu Lys Thr Arg Met Asn Asn Phe Leu Asp Asn
35 40 45

```

aag aac gac atc aag aag aac tac aag gag atc acc ttc agc atg gct	191
Lys Asn Asp Ile Lys Lys Asn Tyr Lys Glu Ile Thr Phe Ser Met Ala	
50 55 60	
ggc tcc ttc gag gac gag atc aag gac ctg aag gag atc gac aag atg	239
Gly Ser Phe Glu Asp Glu Ile Lys Asp Leu Lys Glu Ile Asp Lys Met	
65 70 75	
ttc gac aag gcc aac ctg tcc tcc tcc atc gtg acc tac aag aac gtg	287
Phe Asp Lys Ala Asn Leu Ser Ser Ser Ile Val Thr Tyr Lys Asn Val	
80 85 90 95	
gag cca tcc acc atc gga ttt aac aag cca ctg acc gag ggc aac acc	335
Glu Pro Ser Thr Ile Gly Phe Asn Lys Pro Leu Thr Glu Gly Asn Thr	
100 105 110	
atc aac acc gac gtg cag gcc cag ttc aag gag cag ttc ctg ggc aag	383
Ile Asn Thr Asp Val Gln Ala Gln Phe Lys Glu Gln Phe Leu Gly Lys	
115 120 125	
gac atc aag ttc gac tcc tac ctg gac acc cac ctg act gct cag aac	431
Asp Ile Lys Phe Asp Ser Tyr Leu Asp Thr His Leu Thr Ala Gln Asn	
130 135 140	
gtg tcc tcc aag gag agg atc atc ctc caa gtg acc gtg cca tcc ggc	479
Val Ser Ser Lys Glu Arg Ile Ile Leu Gln Val Thr Val Pro Ser Gly	
145 150 155	
aag ggc tcc acc atc ccg acc aag gct ggt gtg atc ctg aac aac aac	527
Lys Gly Ser Thr Ile Pro Thr Lys Ala Gly Val Ile Leu Asn Asn Asn	
160 165 170 175	
gag tac aag atg ctg atc gac aac ggc tac gtg ctg cac gtg gac aac	575
Glu Tyr Lys Met Leu Ile Asp Asn Gly Tyr Val Leu His Val Asp Asn	
180 185 190	
atc tcc aag gtg gtg aag aag ggt tac gag tgc ctg cag atc cag ggt	623
Ile Ser Lys Val Val Lys Lys Gly Tyr Glu Cys Leu Gln Ile Gln Gly	
195 200 205	
acc ctg aag aag tcc ctg gac ttc aag aac gac atc aac gcc gag gct	671
Thr Leu Lys Lys Ser Leu Asp Phe Lys Asn Asp Ile Asn Ala Glu Ala	
210 215 220	
cac agg tgg ggc atg aag aac tac gag ggt tgg gct aag aac ctg acc	719
His Arg Trp Gly Met Lys Asn Tyr Glu Gly Trp Ala Lys Asn Leu Thr	
225 230 235	
gac cca cag agg gag gcc ctg gac ggc tac gct agg cag gac tac aag	767
Asp Pro Gln Arg Glu Ala Leu Asp Gly Tyr Ala Arg Gln Asp Tyr Lys	
240 245 250 255	
cag atc aac gac tac ctg cgg aac cag ggt ggc tcc ggc aac gag aag	815
Gln Ile Asn Asp Tyr Leu Arg Asn Gln Gly Gly Ser Gly Asn Glu Lys	
260 265 270	

ctg gac acc cag atc aag aac atc tcc gag gct ctg gag aag cag ccg 863
 Leu Asp Thr Gln Ile Lys Asn Ile Ser Glu Ala Leu Glu Lys Gln Pro
 275 280 285

atc cca gag aac atc acc gtg tac agg tgg tgc ggt atg gcc gag ttc 911
 Ile Pro Glu Asn Ile Thr Val Tyr Arg Trp Cys Gly Met Ala Glu Phe
 290 295 300

ggt tac cag att tcc gac cca ctg cca tcc ctg aag gag atg gag gag 959
 Gly Tyr Gln Ile Ser Asp Pro Leu Pro Ser Leu Lys Glu Met Glu Glu
 305 310 315

aag ttc ctg aac acc atg aag gag gac aag ggt tac atg tcc acc tcc 1007
 Lys Phe Leu Asn Thr Met Lys Glu Asp Lys Gly Tyr Met Ser Thr Ser
 320 325 330 335

ctg tcc tcc gag agg ctg tcc gct ttc ggc tcc agg aag ttc atc ctg 1055
 Leu Ser Ser Glu Arg Leu Ser Ala Phe Gly Ser Arg Lys Phe Ile Leu
 340 345 350

agg ctg cag gtg cca aag ggt tcc act ggt gcc tac ctg tcc gct atc 1103
 Arg Leu Gln Val Pro Lys Gly Ser Thr Gly Ala Tyr Leu Ser Ala Ile
 355 360 365

ggt ggc ttc gct tcc gag aag gag atc ctg atc gac aag gac tcc aac 1151
 Gly Gly Phe Ala Ser Glu Lys Glu Ile Leu Ile Asp Lys Asp Ser Asn
 370 375 380

tac cac atc gac aag atc acc gag gtg gtg atc aag ggt gtg aag agg 1199
 Tyr His Ile Asp Lys Ile Thr Glu Val Val Ile Lys Gly Val Lys Arg
 385 390 395

tac gta gtg gac gct acc ctg ctg acc aag tgaggctagc 1239
 Tyr Val Val Asp Ala Thr Leu Leu Thr Lys
 400 405

<210> 8
 <211> 409
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> amino acid sequence of ISP2A-1 protein

<400> 8
 Met Ala Leu Val Lys Thr Thr Asn Asn Thr Glu Asp Phe Lys Glu Asp
 1 5 10 15
 Lys Glu Lys Ala Lys Glu Trp Gly Lys Glu Lys Glu Lys Glu Trp Lys
 20 25 30
 Leu Thr Val Thr Glu Lys Thr Arg Met Asn Asn Phe Leu Asp Asn Lys
 35 40 45
 Asn Asp Ile Lys Lys Asn Tyr Lys Glu Ile Thr Phe Ser Met Ala Gly
 50 55 60
 Ser Phe Glu Asp Glu Ile Lys Asp Leu Lys Glu Ile Asp Lys Met Phe
 65 70 75 80

Asp	Lys	Ala	Asn	Leu	Ser	Ser	Ser	Ile	Val	Thr	Tyr	Lys	Asn	Val	Glu	
				85					90					95		
Pro	Ser	Thr	Ile	Gly	Phe	Asn	Lys	Pro	Leu	Thr	Glu	Gly	Asn	Thr	Ile	
			100					105					110			
Asn	Thr	Asp	Val	Gln	Ala	Gln	Phe	Lys	Glu	Gln	Phe	Leu	Gly	Lys	Asp	
		115					120					125				
Ile	Lys	Phe	Asp	Ser	Tyr	Leu	Asp	Thr	His	Leu	Thr	Ala	Gln	Asn	Val	
	130					135					140					
Ser	Ser	Lys	Glu	Arg	Ile	Ile	Leu	Gln	Val	Thr	Val	Pro	Ser	Gly	Lys	
145					150					155					160	
Gly	Ser	Thr	Ile	Pro	Thr	Lys	Ala	Gly	Val	Ile	Leu	Asn	Asn	Asn	Glu	
			165					170						175		
Tyr	Lys	Met	Leu	Ile	Asp	Asn	Gly	Tyr	Val	Leu	His	Val	Asp	Asn	Ile	
		180					185						190			
Ser	Lys	Val	Val	Lys	Lys	Gly	Tyr	Glu	Cys	Leu	Gln	Ile	Gln	Gly	Thr	
	195					200						205				
Leu	Lys	Lys	Ser	Leu	Asp	Phe	Lys	Asn	Asp	Ile	Asn	Ala	Glu	Ala	His	
	210					215					220					
Arg	Trp	Gly	Met	Lys	Asn	Tyr	Glu	Gly	Trp	Ala	Lys	Asn	Leu	Thr	Asp	
225				230						235					240	
Pro	Gln	Arg	Glu	Ala	Leu	Asp	Gly	Tyr	Ala	Arg	Gln	Asp	Tyr	Lys	Gln	
			245						250					255		
Ile	Asn	Asp	Tyr	Leu	Arg	Asn	Gln	Gly	Gly	Ser	Gly	Asn	Glu	Lys	Leu	
		260					265						270			
Asp	Thr	Gln	Ile	Lys	Asn	Ile	Ser	Glu	Ala	Leu	Glu	Lys	Gln	Pro	Ile	
	275					280						285				
Pro	Glu	Asn	Ile	Thr	Val	Tyr	Arg	Trp	Cys	Gly	Met	Ala	Glu	Phe	Gly	
	290					295					300					
Tyr	Gln	Ile	Ser	Asp	Pro	Leu	Pro	Ser	Leu	Lys	Glu	Met	Glu	Glu	Lys	
305				310						315					320	
Phe	Leu	Asn	Thr	Met	Lys	Glu	Asp	Lys	Gly	Tyr	Met	Ser	Thr	Ser	Leu	
			325						330					335		
Ser	Ser	Glu	Arg	Leu	Ser	Ala	Phe	Gly	Ser	Arg	Lys	Phe	Ile	Leu	Arg	
		340						345					350			
Leu	Gln	Val	Pro	Lys	Gly	Ser	Thr	Gly	Ala	Tyr	Leu	Ser	Ala	Ile	Gly	
	355						360					365				
Gly	Phe	Ala	Ser	Glu	Lys	Glu	Ile	Leu	Ile	Asp	Lys	Asp	Ser	Asn	Tyr	
	370					375					380					
His	Ile	Asp	Lys	Ile	Thr	Glu	Val	Val	Ile	Lys	Gly	Val	Lys	Arg	Tyr	
385				390						395					400	
Val	Val	Asp	Ala	Thr	Leu	Leu	Thr	Lys								
			405													

<210> 9
 <211> 2520
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> modified DNA encoding ISP1A protein

<220>
 <221> CDS
 <222> (3)..(2510)

<400> 9

cc atg gct atc gcc acc acc acc cag gct tcg aag gac aac cag atc	47
Met Ala Ile Ala Thr Thr Thr Gln Ala Ser Lys Asp Asn Gln Ile	
1 5 10 15	
gac agg gag ggc ctg ctg ggc tac tac ttc aag ggc aag gac ttc aac	95
Asp Arg Glu Gly Leu Leu Gly Tyr Tyr Phe Lys Gly Lys Asp Phe Asn	
20 25 30	
gac ctg acc ctg ttc gct cca acc agg gac aac acc ctg atc tac gac	143
Asp Leu Thr Leu Phe Ala Pro Thr Arg Asp Asn Thr Leu Ile Tyr Asp	
35 40 45	
cag cag acc gct aac acc ctc gtg gac cag aag cac cag gag tac cac	191
Gln Gln Thr Ala Asn Thr Leu Val Asp Gln Lys His Gln Glu Tyr His	
50 55 60	
tcc atc cgc tgg atc ggc ctg atc cag tcc tcc gcc act ggt gac ttc	239
Ser Ile Arg Trp Ile Gly Leu Ile Gln Ser Ser Ala Thr Gly Asp Phe	
65 70 75	
acc ttc aag ctg tcc gac gac gag aac gcc atc atc gag ctg gac ggc	287
Thr Phe Lys Leu Ser Asp Asp Glu Asn Ala Ile Ile Glu Leu Asp Gly	
80 85 90 95	
aag gtg atc tcc gag aag ggc aac aac aag cag tcc gtg cac ctc gaa	335
Lys Val Ile Ser Glu Lys Gly Asn Asn Lys Gln Ser Val His Leu Glu	
100 105 110	
aag ggc cag ctg gtg cag atc aag atc gag tac cag tcc gac gac gcc	383
Lys Gly Gln Leu Val Gln Ile Lys Ile Glu Tyr Gln Ser Asp Asp Ala	
115 120 125	
ctg cac atc gac aac aag atc ttc aag gag ctg aag ctg ttc aag atc	431
Leu His Ile Asp Asn Lys Ile Phe Lys Glu Leu Lys Leu Phe Lys Ile	
130 135 140	
gac tcc cag aac cac tcc cag cag gtg cag cag gac gag ctg agg aac	479
Asp Ser Gln Asn His Ser Gln Gln Val Gln Gln Asp Glu Leu Arg Asn	
145 150 155	
cca gag ttc aac aag aag gag acc cag gtg ttc ctg aag aag gcc tcc	527
Pro Glu Phe Asn Lys Lys Glu Thr Gln Val Phe Leu Lys Lys Ala Ser	
160 165 170 175	
aag acc aac ctg ttc acc cag aag acc aag agg gac atc gac gag gac	575
Lys Thr Asn Leu Phe Thr Gln Lys Thr Lys Arg Asp Ile Asp Glu Asp	
180 185 190	
acc gac acc gac ggc gac tcc atc ccg gac gtg tgg gag gag aac ggc	623
Thr Asp Thr Asp Gly Asp Ser Ile Pro Asp Val Trp Glu Glu Asn Gly	
195 200 205	
tac acc atc cag aac aag gtg gcc gtg aag tgg gac gac tcc ctg gcc	671
Tyr Thr Ile Gln Asn Lys Val Ala Val Lys Trp Asp Asp Ser Leu Ala	
210 215 220	

tcc aag ggc tac cag aag ttc acc agc aac cca ctc gaa gcc cac acc Ser Lys Gly Tyr Gln Lys Phe Thr Ser Asn Pro Leu Glu Ala His Thr 225 230 235	719
gtg ggc gac cca tac tcc gac tac gag aag gct gct agg gac atg cca Val Gly Asp Pro Tyr Ser Asp Tyr Glu Lys Ala Ala Arg Asp Met Pro 240 245 250 255	767
ctg tcc aac gcc aag gag acc ttc aac cca ctg gtg gct gct ttc cca Leu Ser Asn Ala Lys Glu Thr Phe Asn Pro Leu Val Ala Ala Phe Pro 260 265 270	815
tcc gtg aac gtg tcc ctc gaa aag gtg atc ctg tcc aag aac gag gac Ser Val Asn Val Ser Leu Glu Lys Val Ile Leu Ser Lys Asn Glu Asp 275 280 285	863
ctg tcc cac tcc gtg gag tcc tcc cag tcc acc aac tgg tcc tac acc Leu Ser His Ser Val Glu Ser Ser Gln Ser Thr Asn Trp Ser Tyr Thr 290 295 300	911
aac acc gag ggc gtg aac gtg aac gct ggt tgg tcc ggt ctg ggt cca Asn Thr Glu Gly Val Asn Val Asn Ala Gly Trp Ser Gly Leu Gly Pro 305 310 315	959
tcc ttc ggc gtg tcc gtg aac tac cag cac tcc gag acc gtg gcc aac Ser Phe Gly Val Ser Val Asn Tyr Gln His Ser Glu Thr Val Ala Asn 320 325 330 335	1007
gag tgg ggc tcc gcc acc aac gac ggc acc cac atc aac ggt gct gag Glu Trp Gly Ser Ala Thr Asn Asp Gly Thr His Ile Asn Gly Ala Glu 340 345 350	1055
tcc gcc tac ctg aac gcc aac gtg agg tac aac aac gtg ggc acc ggt Ser Ala Tyr Leu Asn Ala Asn Val Arg Tyr Asn Asn Val Gly Thr Gly 355 360 365	1103
gct atc tac gag acc aag cca acc acc tcc ttc atc ctg gac ggc acc Ala Ile Tyr Glu Thr Lys Pro Thr Thr Ser Phe Ile Leu Asp Gly Thr 370 375 380	1151
acc atc ggc acc atc aag gcc aag gag aac acc acc gct ctg acc atc Thr Ile Gly Thr Ile Lys Ala Lys Glu Asn Thr Thr Ala Leu Thr Ile 385 390 395	1199
ctg cca gac cag tcc tac cca gag aag ggc aag aac ggc atc gcc atc Leu Pro Asp Gln Ser Tyr Pro Glu Lys Gly Lys Asn Gly Ile Ala Ile 400 405 410 415	1247
aac acg atg gac gac ttc aac tcc agg ccg atc cca ctg aac aag gag Asn Thr Met Asp Asp Phe Asn Ser Arg Pro Ile Pro Leu Asn Lys Glu 420 425 430	1295
cag ctg aac acc tac ctg tcc aac aag aag ccg atc ctg ctc gaa acc Gln Leu Asn Thr Tyr Leu Ser Asn Lys Lys Pro Ile Leu Leu Glu Thr 435 440 445	1343

gac cag gtg gag ggc aag tac gcc atc aag gac acc aac ggc aac atc Asp Gln Val Glu Gly Lys Tyr Ala Ile Lys Asp Thr Asn Gly Asn Ile 450 455 460	1391
acc atc gct ggt gac tgg aac ggc atc acc gac gag atc tcc gcc aag Thr Ile Ala Gly Asp Trp Asn Gly Ile Thr Asp Glu Ile Ser Ala Lys 465 470 475	1439
acc gcc agc atc atc gtc gac aac ggc aac cag atg tcc gag aag agg Thr Ala Ser Ile Ile Val Asp Asn Gly Asn Gln Met Ser Glu Lys Arg 480 485 490 495	1487
gtg gct gct aag gac tac acc aac cca gag gac aag acc cca aat tta Val Ala Ala Lys Asp Tyr Thr Asn Pro Glu Asp Lys Thr Pro Asn Leu 500 505 510	1535
tcc gtg aag gag gcc ctg aag ctg gcc tac cca gac gag atc gag gag Ser Val Lys Glu Ala Leu Lys Leu Ala Tyr Pro Asp Glu Ile Glu Glu 515 520 525	1583
aag gac ggc ctg ctg ttc tac aac gac cag ccg atc ttc gag gct tcc Lys Asp Gly Leu Leu Phe Tyr Asn Asp Gln Pro Ile Phe Glu Ala Ser 530 535 540	1631
gtg cag tcc tac gtg gac gag tac acc gct aag cag atc agg aag cag Val Gln Ser Tyr Val Asp Glu Tyr Thr Ala Lys Gln Ile Arg Lys Gln 545 550 555	1679
ctg aac gac tcc acc ggt tcc ttc aag gac gtg aag aac ctg tac gac Leu Asn Asp Ser Thr Gly Ser Phe Lys Asp Val Lys Asn Leu Tyr Asp 560 565 570 575	1727
gtg aag ctc gaa ccg aag atg aac ttc aca ata aag acc tcc acc ctg Val Lys Leu Glu Pro Lys Met Asn Phe Thr Ile Lys Thr Ser Thr Leu 580 585 590	1775
tac gac ggt ggt gag tcc gac aac acc aag atc ggc aac tgg tac tac Tyr Asp Gly Gly Glu Ser Asp Asn Thr Lys Ile Gly Asn Trp Tyr Tyr 595 600 605	1823
acc tac gtg gtg aac ggt ggt aac acc ggt aag aag cag tac agg tcc Thr Tyr Val Val Asn Gly Gly Asn Thr Gly Lys Lys Gln Tyr Arg Ser 610 615 620	1871
gct aac aag ggt gct ttc acc gag ctg tcc acc gag tcc aag aac aag Ala Asn Lys Gly Ala Phe Thr Glu Leu Ser Thr Glu Ser Lys Asn Lys 625 630 635	1919
ctg aag aag aac atc gac tac tac gtg tcc ctg tac atg aag gct gac Leu Lys Lys Asn Ile Asp Tyr Tyr Val Ser Leu Tyr Met Lys Ala Asp 640 645 650 655	1967
tcc aag gtg tcc gtg gac atc gag atc gac ggt aag cag gag tcc atc Ser Lys Val Ser Val Asp Ile Glu Ile Asp Gly Lys Gln Glu Ser Ile 660 665 670	2015

gtg acc gac aac atc acc ctg gac cac gtg ggt tac cag agg atc aac Val Thr Asp Asn Ile Thr Leu Asp His Val Gly Tyr Gln Arg Ile Asn 675 680 685	2063
atc ctg gtg cca aac ctg gag ggc aac gag atc aac acc atc tcc atc Ile Leu Val Pro Asn Leu Glu Gly Asn Glu Ile Asn Thr Ile Ser Ile 690 695 700	2111
aag ggt gac ggc cag acc aac gtg tac tgg gac gac gtg tcc ttc gtg Lys Gly Asp Gly Gln Thr Asn Val Tyr Trp Asp Asp Val Ser Phe Val 705 710 715	2159
gag gtg ggt gct gag gag atc gag tac aag gac cca gtg cca cag ttc Glu Val Gly Ala Glu Glu Ile Glu Tyr Lys Asp Pro Val Pro Gln Phe 720 725 730 735	2207
gac atc atc gag ggt gac ttc gac ttc ttc ggt gac cca ctg gcc gtg Asp Ile Ile Glu Gly Asp Phe Asp Phe Phe Gly Asp Pro Leu Ala Val 740 745 750	2255
aag tac cac gac gct acc tac ttc atc gac tcc cca ctg att acc cag Lys Tyr His Asp Ala Thr Tyr Phe Ile Asp Ser Pro Leu Ile Thr Gln 755 760 765	2303
acc cca ggt acc ttc tcc ttc acc tac aag gtg atc ggt gag cag acc Thr Pro Gly Thr Phe Ser Phe Thr Tyr Lys Val Ile Gly Glu Gln Thr 770 775 780	2351
aag acc gtg ctg gac tcc ggt tcc ggc aag aac gct aac agg atc aac Lys Thr Val Leu Asp Ser Gly Ser Gly Lys Asn Ala Asn Arg Ile Asn 785 790 795	2399
ctg gac ttc aag aac gtg aag tcc gac agg tcc ttc ctg tac acc ctg Leu Asp Phe Lys Asn Val Lys Ser Asp Arg Ser Phe Leu Tyr Thr Leu 800 805 810 815	2447
tcc tgc aag gac gac ctg tgg ggc tcc acc agg acc gct gtg gtg agg Ser Cys Lys Asp Asp Leu Trp Gly Ser Thr Arg Thr Ala Val Val Arg 820 825 830	2495
atc ttc gct gtg gac tgaggctagc Ile Phe Ala Val Asp 835	2520

<210> 10
 <211> 836
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> amino acid sequence of ISPlA-1 protein

<400> 10
 Met Ala Ile Ala Thr Thr Thr Gln Ala Ser Lys Asp Asn Gln Ile Asp
 1 5 10 15

Arg	Glu	Gly	Leu	Leu	Gly	Tyr	Tyr	Phe	Lys	Gly	Lys	Asp	Phe	Asn	Asp	
			20					25					30			
Leu	Thr	Leu	Phe	Ala	Pro	Thr	Arg	Asp	Asn	Thr	Leu	Ile	Tyr	Asp	Gln	
		35					40					45				
Gln	Thr	Ala	Asn	Thr	Leu	Val	Asp	Gln	Lys	His	Gln	Glu	Tyr	His	Ser	
	50					55					60					
Ile	Arg	Trp	Ile	Gly	Leu	Ile	Gln	Ser	Ser	Ala	Thr	Gly	Asp	Phe	Thr	
65					70					75					80	
Phe	Lys	Leu	Ser	Asp	Asp	Glu	Asn	Ala	Ile	Ile	Glu	Leu	Asp	Gly	Lys	
				85					90					95		
Val	Ile	Ser	Glu	Lys	Gly	Asn	Asn	Lys	Gln	Ser	Val	His	Leu	Glu	Lys	
			100					105					110			
Gly	Gln	Leu	Val	Gln	Ile	Lys	Ile	Glu	Tyr	Gln	Ser	Asp	Asp	Ala	Leu	
		115					120					125				
His	Ile	Asp	Asn	Lys	Ile	Phe	Lys	Glu	Leu	Lys	Leu	Phe	Lys	Ile	Asp	
	130					135					140					
Ser	Gln	Asn	His	Ser	Gln	Gln	Val	Gln	Gln	Asp	Glu	Leu	Arg	Asn	Pro	
145					150					155					160	
Glu	Phe	Asn	Lys	Lys	Glu	Thr	Gln	Val	Phe	Leu	Lys	Lys	Ala	Ser	Lys	
				165					170					175		
Thr	Asn	Leu	Phe	Thr	Gln	Lys	Thr	Lys	Arg	Asp	Ile	Asp	Glu	Asp	Thr	
			180					185					190			
Asp	Thr	Asp	Gly	Asp	Ser	Ile	Pro	Asp	Val	Trp	Glu	Glu	Asn	Gly	Tyr	
		195					200					205				
Thr	Ile	Gln	Asn	Lys	Val	Ala	Val	Lys	Trp	Asp	Asp	Ser	Leu	Ala	Ser	
	210					215					220					
Lys	Gly	Tyr	Gln	Lys	Phe	Thr	Ser	Asn	Pro	Leu	Glu	Ala	His	Thr	Val	
225					230					235					240	
Gly	Asp	Pro	Tyr	Ser	Asp	Tyr	Glu	Lys	Ala	Ala	Arg	Asp	Met	Pro	Leu	
				245					250					255		
Ser	Asn	Ala	Lys	Glu	Thr	Phe	Asn	Pro	Leu	Val	Ala	Ala	Phe	Pro	Ser	
			260					265					270			
Val	Asn	Val	Ser	Leu	Glu	Lys	Val	Ile	Leu	Ser	Lys	Asn	Glu	Asp	Leu	
	275						280					285				
Ser	His	Ser	Val	Glu	Ser	Ser	Gln	Ser	Thr	Asn	Trp	Ser	Tyr	Thr	Asn	
	290					295					300					
Thr	Glu	Gly	Val	Asn	Val	Asn	Ala	Gly	Trp	Ser	Gly	Leu	Gly	Pro	Ser	
305					310					315					320	
Phe	Gly	Val	Ser	Val	Asn	Tyr	Gln	His	Ser	Glu	Thr	Val	Ala	Asn	Glu	
				325					330					335		
Trp	Gly	Ser	Ala	Thr	Asn	Asp	Gly	Thr	His	Ile	Asn	Gly	Ala	Glu	Ser	
			340					345					350			
Ala	Tyr	Leu	Asn	Ala	Asn	Val	Arg	Tyr	Asn	Asn	Val	Gly	Thr	Gly	Ala	
		355					360					365				
Ile	Tyr	Glu	Thr	Lys	Pro	Thr	Thr	Ser	Phe	Ile	Leu	Asp	Gly	Thr	Thr	
	370					375					380					
Ile	Gly	Thr	Ile	Lys	Ala	Lys	Glu	Asn	Thr	Thr	Ala	Leu	Thr	Ile	Leu	
385					390					395					400	
Pro	Asp	Gln	Ser	Tyr	Pro	Glu	Lys	Gly	Lys	Asn	Gly	Ile	Ala	Ile	Asn	
				405					410					415		
Thr	Met	Asp	Asp	Phe	Asn	Ser	Arg	Pro	Ile	Pro	Leu	Asn	Lys	Glu	Gln	
			420					425					430			
Leu	Asn	Thr	Tyr	Leu	Ser	Asn	Lys	Lys	Pro	Ile	Leu	Leu	Glu	Thr	Asp	
		435					440					445				
Gln	Val	Glu	Gly	Lys	Tyr	Ala	Ile	Lys	Asp	Thr	Asn	Gly	Asn	Ile	Thr	
	450					455					460					

Ile	Ala	Gly	Asp	Trp	Asn	Gly	Ile	Thr	Asp	Glu	Ile	Ser	Ala	Lys	Thr	465	470	475	480
Ala	Ser	Ile	Ile	Val	Asp	Asn	Gly	Asn	Gln	Met	Ser	Glu	Lys	Arg	Val	485	490		495
Ala	Ala	Lys	Asp	Tyr	Thr	Asn	Pro	Glu	Asp	Lys	Thr	Pro	Asn	Leu	Ser	500	505		510
Val	Lys	Glu	Ala	Leu	Lys	Leu	Ala	Tyr	Pro	Asp	Glu	Ile	Glu	Glu	Lys	515	520		525
Asp	Gly	Leu	Leu	Phe	Tyr	Asn	Asp	Gln	Pro	Ile	Phe	Glu	Ala	Ser	Val	530	535		540
Gln	Ser	Tyr	Val	Asp	Glu	Tyr	Thr	Ala	Lys	Gln	Ile	Arg	Lys	Gln	Leu	545	550		555
Asn	Asp	Ser	Thr	Gly	Ser	Phe	Lys	Asp	Val	Lys	Asn	Leu	Tyr	Asp	Val	565	570		575
Lys	Leu	Glu	Pro	Lys	Met	Asn	Phe	Thr	Ile	Lys	Thr	Ser	Thr	Leu	Tyr	580	585		590
Asp	Gly	Gly	Glu	Ser	Asp	Asn	Thr	Lys	Ile	Gly	Asn	Trp	Tyr	Tyr	Thr	595	600		605
Tyr	Val	Val	Asn	Gly	Gly	Asn	Thr	Gly	Lys	Lys	Gln	Tyr	Arg	Ser	Ala	610	615		620
Asn	Lys	Gly	Ala	Phe	Thr	Glu	Leu	Ser	Thr	Glu	Ser	Lys	Asn	Lys	Leu	625	630		635
Lys	Lys	Asn	Ile	Asp	Tyr	Tyr	Val	Ser	Leu	Tyr	Met	Lys	Ala	Asp	Ser	645	650		655
Lys	Val	Ser	Val	Asp	Ile	Glu	Ile	Asp	Gly	Lys	Gln	Glu	Ser	Ile	Val	660	665		670
Thr	Asp	Asn	Ile	Thr	Leu	Asp	His	Val	Gly	Tyr	Gln	Arg	Ile	Asn	Ile	675	680		685
Leu	Val	Pro	Asn	Leu	Glu	Gly	Asn	Glu	Ile	Asn	Thr	Ile	Ser	Ile	Lys	690	695		700
Gly	Asp	Gly	Gln	Thr	Asn	Val	Tyr	Trp	Asp	Asp	Val	Ser	Phe	Val	Glu	705	710		715
Val	Gly	Ala	Glu	Glu	Ile	Glu	Tyr	Lys	Asp	Pro	Val	Pro	Gln	Phe	Asp	725	730		735
Ile	Ile	Glu	Gly	Asp	Phe	Asp	Phe	Phe	Gly	Asp	Pro	Leu	Ala	Val	Lys	740	745		750
Tyr	His	Asp	Ala	Thr	Tyr	Phe	Ile	Asp	Ser	Pro	Leu	Ile	Thr	Gln	Thr	755	760		765
Pro	Gly	Thr	Phe	Ser	Phe	Thr	Tyr	Lys	Val	Ile	Gly	Glu	Gln	Thr	Lys	770	775		780
Thr	Val	Leu	Asp	Ser	Gly	Ser	Gly	Lys	Asn	Ala	Asn	Arg	Ile	Asn	Leu	785	790		795
Asp	Phe	Lys	Asn	Val	Lys	Ser	Asp	Arg	Ser	Phe	Leu	Tyr	Thr	Leu	Ser	805	810		815
Cys	Lys	Asp	Asp	Leu	Trp	Gly	Ser	Thr	Arg	Thr	Ala	Val	Val	Arg	Ile	820	825		830
Phe	Ala	Val	Asp													835			

*75749-27

CLAIMS:

1. A protein comprising the amino acid sequence of the smallest active toxin of the protein of SEQ ID NO: 2, wherein said smallest active toxin is:
 - a. a protein comprising an amino acid sequence with at least 97%
5 sequence identity to a fragment of SEQ ID NO: 2, said fragment consisting of the amino acid sequence from amino acid position 38 to amino acid position 871 of SEQ ID NO: 2; and
 - b. insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with a protein fragment of SEQ ID NO: 4 from amino acid
10 position 51 to amino acid position 457.
2. A protein comprising the amino acid sequence of the smallest active toxin of the protein of SEQ ID NO: 4, wherein said smallest active toxin is:
 - a. a protein comprising an amino acid sequence with at least 97%
sequence identity to a fragment of SEQ ID NO: 4, said fragment consisting of the
15 amino acid sequence from amino acid position 51 to amino acid position 457 of SEQ ID NO: 4; and
 - b. insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with a protein fragment of SEQ ID NO: 2 from amino acid position 38 to amino acid position 871.
- 20 3. An insecticidal protein comprising a sequence with at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 2 from amino acid position 38 to amino acid position 768 or from amino acid position 38 to amino acid position 871.
4. A protein comprising the amino acid sequence of SEQ ID NO: 2 from
25 amino acid position 38 to amino acid position 768 or from amino acid position 38 to amino acid position 871.

•75749-27

5. The protein according to claim 3 or 4 comprising the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 10.
6. An insecticidal protein comprising a sequence with at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 4 from amino acid position 51 to amino acid position 449 or from amino acid position 51 to amino acid position 457.
7. A protein comprising the amino acid sequence of SEQ ID NO: 4 from amino acid position 51 to amino acid position 449 or from amino acid position 51 to amino acid position 457.
8. The protein according to claim 6 or 7 comprising the amino acid sequence of SEQ ID NO: 4 or SEQ ID NO: 8.
9. A protein comprising an amino acid sequence with at least 97% sequence identity to the amino acid sequence of SEQ ID NO: 2, wherein said protein is insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with the protein of SEQ ID NO: 4 from amino acid position 51 to amino acid position 457.
10. A protein comprising an amino acid sequence with at least 97% sequence identity to the amino acid sequence of SEQ ID NO: 4, wherein said protein is insecticidal to *Diabrotica virgifera* larvae when ingested by said larvae in combination with the protein of SEQ ID NO: 2 from amino acid position 38 to amino acid position 871.
11. A DNA molecule encoding the protein as defined in any one of claims 1, 3, 4, 5 or 9.
12. A DNA molecule encoding the protein as defined in any one of claims 2, 6, 7, 8 or 10.

-75749-27

13. The DNA molecule of claim 11, comprising an artificial DNA sequence having a different codon usage compared to the naturally occurring DNA sequence but encoding the same protein sequence.

14. The DNA molecule of claim 12, comprising an artificial DNA sequence
5 having a different codon usage compared to the naturally occurring DNA sequence but encoding the same protein sequence.

15. A chimeric gene comprising the DNA molecule as defined in claim 13 and a plant-expressible promoter region.

16. A chimeric gene comprising the DNA molecule as defined in claim 14
10 and a plant-expressible promoter region.

17. The chimeric gene of claim 15 wherein said promoter region is active in root tissue.

18. The chimeric gene of claim 16, wherein said promoter region is active in root tissue.

15 19. The chimeric gene of claim 15, wherein said promoter region comprises the DNA sequence of SEQ ID NO: 5 or 6.

20. The chimeric gene of claim 16, wherein said promoter region comprises the DNA sequence of SEQ ID NO: 5 or 6.

21. The chimeric gene of claim 15 wherein said chimeric gene further
20 comprises a signal peptide for secretion from a cell or for targeting to a cellular organelle.

22. The chimeric gene of claim 16 wherein said chimeric gene further comprises a signal peptide for secretion from a cell or for targeting to a cellular organelle.

75749-27

23. The chimeric gene of claim 21 wherein said signal peptide is a chloroplast transit peptide.
24. The chimeric gene of claim 22 wherein said signal peptide is a chloroplast transit peptide.
- 5 25. A plant cell comprising the chimeric gene as defined in any one of claims 15, 17, 19, 21 or 23.
26. A plant cell comprising the chimeric gene as defined in any one of claims 16, 18, 20, 22 or 24.
- 10 27. A plant cell transformed to contain the chimeric gene as defined in any one of claims 15, 17, 19, 21 or 23, and the chimeric gene as defined in any one of claims 16, 18, 20, 22 or 24.
28. A plant cell comprising the protein as defined in any one of claims 1, 3, 4, 5 or 9, and the protein as defined in any one of claims 2, 6, 7, 8 or 10.
- 15 29. Use of a plant or a seed to control coleopteran insects, said plant or seed comprising the chimeric gene as defined in any one of claims 15, 17, 19, 21 or 23 integrated in its cells.
30. Use of a plant or a seed to control coleopteran insects, said plant or seed comprising the chimeric gene as defined in any one of claims 16, 18, 20, 22, or 24.
- 20 31. The use of claim 29 wherein said plant or seed is a corn plant or seed.
32. The use of claim 30 wherein said plant or seed is a corn plant or seed.
33. A microorganism, transformed to contain the DNA molecule as defined in claim 11 or the DNA molecule as defined in claim 12.

-75749-27

34. A process for controlling coleopteran insects, comprising expressing the protein of any one of claims 1, 3, 4, 5 or 9 in a cell of a plant, which cell also expresses the protein of any one of claims 2, 6, 7, 8 or 10.

35. A process for rendering a plant resistant to coleopteran insects, said process comprising transforming a plant cell with a first chimeric gene and a second chimeric gene, and regenerating a transformed plant from said cell, which plant is resistant to said coleopteran insects, wherein said first chimeric gene is the chimeric gene as defined in any one of claims 15, 17, 19, 21 or 23; and said second chimeric gene is the chimeric gene as defined in any one of claims 16, 18, 20, 22 or 24.

36. A process for controlling coleopteran insect pests, said process comprising planting, sowing or growing in a field, a plant transformed with a first chimeric gene and a second chimeric gene, wherein said first chimeric gene is the chimeric gene as defined in any one of claims 15, 17, 19, 21, or 23; and said second chimeric gene is the chimeric gene as defined in any one of claims 16, 18, 20, 22, or 24.

37. The process of claim 36, wherein said plant is a corn plant.

38. The process of any one of claims 34 to 36, wherein said insects are selected from the group consisting of rootworms, weevils, potato beetles, *Diabrotica* species, *Anthonomus* spp., *Leptinotarsa* spp., *Agelastica alni*, *Hypera postica*, *Hypera brunneipennis*, *Haltica tombacina*, *Anthonomus grandis*, *Tenebrio molitor*, *Triboleum castaneum*, *Dicladispa armigera*, *Trichispa serica*, *Oulema oryzae*, *Colaspis brunnea*, *Lissorhynchus oryzophilus*, *Phyllotreta cruciferae*, *Phyllotreta striolata*, *Psylliodes punctulata*, *Entomoscelis americana*, *Meligethes aeneus*, *Ceutorynchus* sp., *Psylliodes chrysocephala*, *Phyllotreta undulata*, *Leptinotarsa decemlineata*, *Diabrotica undecimpunctata undecimpunctata*, *Diabrotica undecimpunctata howardi*, *Diabrotica barberi*, and *Diabrotica virgifera*.

~75749-27

39. A product, comprising the protein as defined in any one of claims 1, 3, 4, 5 or 9 and the protein as defined in any one of claims 2, 6, 7, 8, or 10 to protect corn plants against corn rootworms.

40. The product of claim 39 which is a corn plant or an insecticidal
5 composition to protect corn plants against corn rootworms.