

(19) **DANMARK**

(10) **DK/EP 1620133 T3**



(12)

**Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 48/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-02-29**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2015-12-09**
- (86) Europæisk ansøgning nr.: **04760694.2**
- (86) Europæisk indleveringsdag: **2004-04-30**
- (87) Den europæiske ansøgnings publiceringsdag: **2006-02-01**
- (86) International ansøgning nr.: **US2004013685**
- (87) Internationalt publikationsnr.: **WO2004098648**
- (30) Prioritet: **2003-05-01 US 467195 P** **2003-06-04 US 475726 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **GENZYME CORPORATION, 500 Kendall Street, Cambridge, MA 02142, USA**
- (72) Opfinder: **DODGE, James, 1 Gates Circle, Worcester, MA 01603, USA**
PASSINI, Marco, A., 14 Nightingale Drive, Shrewsbury, Massachusetts 01545, USA
STEWART, Gregory R., 4735 Xene Lane South, Plymouth, Minnesota 55446, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **GENTERAPI TIL NEUROMETABOLISKE SYGDOMME**
- (56) Fremdragne publikationer:
PASSINI MA ET.AL.: "Distribution of a lysosomal enzyme in the adult brain by axonal transport and by cells of the rostral migratory stream" THE JOURNAL OF NEUROSCIENCE, vol. 22, no. 15, 1 August 2002 (2002-08-01), pages 6437-6446, XP002294218

DESCRIPTION

[0001] This application claims priority to U.S. patent application No. 60/467,195, filed on May 1, 2003, and U.S. patent application No. 60/475,726, filed on June 4, 2003.

Field of the Invention

[0002] The present disclosure relates to compositions and methods for treating disorders affecting the central nervous system (CNS). The present disclosure further relates to compositions comprising viral vectors such as adeno-associated virus (AAV) vectors, and methods of administration thereof.

Background of the Invention

[0003] A group of metabolic disorders known as lysosomal storage diseases (LSD) includes over forty genetic disorders, many of which involve genetic defects in various lysosomal hydrolases. Representative lysosomal storage diseases and the associated defective enzymes are listed in Table 1.

Table 1

<i>Lysosomal storage disease</i>	<i>Defective enzyme</i>
Aspartylglucosaminuria	Aspartylglucosaminidase
Fabry	α -Galactosidase A
Infantile Batten Disease* (CNL1)	Palmitoyl Protein Thioesterase
Classic Late Infantile Batten Disease* (CNL2)	Tripeptidyl Peptidase
Juvenile Batten Disease* (CNL3)	Lysosomal Transmembrane Protein
Batten, other forms* (CNL4-CNL8)	Multiple gene products
Cystinosis	Cysteine transporter
Farber	Acid ceramidase
Fucosidosis	Acid α -L-fucosidase
Galactosidosialidosis	Protective protein/cathepsin A
Gaucher types 1, 2*, and 3*	Acid β -glucosidase, or glucocerebrosidase
G _{M1} gangliosidosis*	Acid β -galactosidase
Hunter*	Iduronate-2-sulfatase
Hurler-Scheie*	α -L-Iduronidase
Krabbe*	Galactocerebrosidase
α -Mannosidosis*	Acid α -mannosidase
β -Mannosidosis*	Acid β -mannosidase
Maroteaux-Lamy	Arylsulfatase B
Metachromatic leukodystrophy*	Arylsulfatase A
Morquio A	N-Acetylgalactosamine-6-sulfate sulfatase
Morquio B	Acid β -galactosidase
Mucopolipidosis II/III*	N-Acetylglucosamine-1-phosphotransferase
Niemann-Pick A*, B	Acid sphingomyelinase
Niemann-Pick C*	NPC-1
Pompe*	Acid α -glucosidase
Sandhoff*	β -Hexosaminidase B
Sarfilippo A*	Heparan N-sulfatase

Lysosomal storage disease	Defective enzyme
Sanfilippo B*	α -N-Acetylglucosaminidase
Sanfilippo C*	Acetyl-CoA: α -glucosaminide N-acetyltransferase
Sanfilippo D*	N-Acetylglucosamine-6-sulfate sulfatase
Schindler Disease*	α -N-Acetylgalactosaminidase
Schindler-Kanzaki	α -N-Acetylgalactosaminidase
Sialidosis	α -Neuramidase
Sly*	β -Glucuronidase
Tay-Sachs*	β -Hexosaminidase A
Wolman*	Acid Lipase
* CNS involvement	

[0004] The hallmark feature of LSD is the abnormal accumulation of metabolites in the lysosomes which leads to the formation of large numbers of distended lysosomes in the perikaryon. A major challenge to treating LSD (as opposed to treating a liver-specific enzymopathy) is the need to reverse lysosomal storage pathology in multiple separate tissues. Some LSDs can be effectively treated by intravenous infusion of the missing enzyme, known as enzyme replacement therapy (ERT). For example, Gaucher type 1 patients have only visceral disease and respond favorably to ERT with recombinant glucocerebrosidase (Cerezyme®, Genzyme Corp.). However, patients with metabolic disease that affects the CNS (e.g., type 2 or 3 Gaucher disease) do not respond to intravenous ERT because the replacement enzyme is prevented from entering the brain by the blood brain barrier (BBB). Furthermore, attempts to introduce a replacement enzyme into the brain by direct injection have been unsuccessful in part due to enzyme cytotoxicity at high local concentrations (unpublished observations) and limited parenchymal diffusion rates in the brain (Pardridge, Peptide Drug Delivery to the Brain, Raven Press, 1991).

[0005] Gene therapy is an emerging treatment modality for disorders affecting the CNS, including LSDs. In this approach, restoration of the normal metabolic pathway and reversal of pathology occurs by transducing affected cells with a vector carrying a healthy version of the gene. CNS gene therapy has been facilitated by the development of viral vectors capable of effectively infecting post-mitotic neurons. For a review of viral vectors for gene delivery to the CNS, see Davidson et al. (2003) *Nature Rev.*, 4:353-364. Adeno-associated virus (AAV) vectors are considered optimal for CNS gene therapy because they are substantially non-toxic, non-immunogenic, neurotropic and can sustain long-term expression in the CNS (Kaplitt et al. (1994) *Nat. Genet.*, 8:148-154; Bartlett et al. (1998) *Hum. Gene Ther.*, 9:1181-1186; and Passini et al. (2002) *J. Neurosci.*, 22:6437-6446).

[0006] A therapeutic transgene product, e.g., an enzyme, can be secreted by transduced cells and subsequently taken up by other cells, in which it then alleviates pathology. This process is known as cross-correction (Neufeld et al. (1970) *Science*, 169:141-146). However, the correction of pathology, such as storage pathology in the context of LSD, is typically confined to the immediate vicinity of the injection site because of limited parenchymal diffusion of the injected vector and the secreted transgene product (Taylor et al. (1997) *Nat. Med.*, 3:771-774; Skorupa et al. (1999) *Exp. Neurol.*, 160:17-27). Thus, neuropathology affecting multiple brain regions requires widespread vector delivery, using multiple spatially distributed injections, especially in a large brain such as human. This significantly increases the risk of brain damage. In addition, some regions of the brain may be difficult to access surgically. Thus, other modes of vector transport within the CNS, besides diffusion, would be beneficial.

[0007] When administered at axonal endings, some viruses are internalized and transported retrogradely along the axon to the nucleus. Neurons in one brain region are interconnected by axons to distal brain regions thereby providing a transport system for vector delivery. Studies with adenovirus, HSV, and pseudo-rabies virus have utilized trafficking properties of these viruses to deliver genes to distal structures within the brain (Soudas et al. (2001) *FASEB J.*, 15:2283-2285; Breakefield et al. (1991) *New Biol.*, 3:203-218; and deFalco et al. (2001) *Science*, 291:2608-2613). However, drawbacks associated with these viral vectors, e.g., immunogenicity and/or cell toxicity, limit their utility.

[0008] Several groups have reported that the transduction of the brain by AAV serotype 2 (AAV2) is limited to the intracardial injection site (Kaplitt et al. (1994) *Nat. Genet.*, 8:148-154; Passini et al. (2002) *J. Neurosci.*, 22:6437-6446; and Chamberlin et al. (1998) *Brain Res.*, 793:169-175). One recent report suggests that retrograde axonal transport of AAV2 can also occur in select circuits of the normal rat brain (Kaspar et al. (2002) *Mol. Ther.*, 5:50-56). However, it is not known what specific parameters were responsible for the observed axonal transport, and whether sufficient and effective axonal transport would occur in a diseased neuron that is in a state of cellular dysfunction. Indeed, lesions observed in LSD neurons have been reported to interfere with or

even block axonal transport (reviewed in Walkley (1998) Brain Pathol., 8:175-193), suggesting that disease-compromised neurons would not support trafficking of AAV along their axons.

[0009] Therefore, there is a need in the art to develop new therapeutic methods for treating metabolic disorders that affect the CNS.

SUMMARY OF THE INVENTION

[0010] The present disclosure provides methods and compositions for treating or preventing metabolic disorders, such as lysosomal storage diseases (LSD), that are characterized by or associated with a risk of diminution of CNS function.

[0011] The present disclosure further provides methods for minimally invasive targeted delivery of a transgene to select regions in the brain of an affected subject.

[0012] Additional advantages of the present disclosure, including advantages of the present invention, will be set forth in part in the following description, and in part will be understood from the description, or may be learned by practice of the present disclosure, including the present invention.

[0013] In the experiments leading to the present disclosure, including the present invention, acid sphingomyelinase (ASM) knockout mice, a model of Niemann-Pick Type A disease, were administered an AAV2 vector carrying the human ASM gene (AAV-ASM) by a single intracranial injection into one hemisphere of the brain. The present disclosure, including the present invention, is based, in part, on the discovery and demonstration that the injection of high titer AAV-ASM into the diseased brain results in AAV-ASM expression within multiple distal sites in a pattern consistent with the topographical organization of the projection neurons that innervate the injection site. The present disclosure, including the present invention, is further based, in part, on the discovery and demonstration of extensive correction of lysosomal storage pathology at the injection site and all distal sites to which AAV-ASM was transported and where ASM was expressed.

[0014] Accordingly, in one aspect, the present disclosure provides methods for treating neurometabolic disorders in mammals. The populations treated by the methods of the present disclosure include, but are not limited to, patients having or at risk for developing a LSD, such as disorders listed in Table 1, particularly, if such disease affects the CNS. In an illustrative instance, the disease is Niemann-Pick A disease.

[0015] In one aspect, the disclosed methods include administering to the CNS of an afflicted subject an AAV viral vector carrying a transgene encoding a therapeutic product and allowing the transgene to be expressed within the CNS distally from the administration site at a therapeutic level. In an illustrative instance, the administration is accomplished by direct intraparenchymal injection of a high titer AAV vector solution into the diseased brain. Thereafter the transgene is expressed distally, contralaterally or ipsilaterally, to the administration site at a therapeutic level at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm from the administration site.

[0016] In another aspect, the present disclosure also provides a method of delivering a recombinant AAV genome to the nucleus of a disease-compromised neuron in vivo. In some instances, the cellular pathology exhibited by the neuron is that of a lysosomal storage disease such as disorders listed in Table 1. In an illustrative instance, the disease is Niemann-Pick A disease. The method of delivering a recombinant AAV genome to the nucleus of a disease-compromised neuron comprises contacting an axonal ending of the disease-compromised neuron with a composition comprising an AAV viral particle comprising the recombinant AAV genome and allowing the viral particle to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron. The concentration of the vector in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). In certain instances, the neuron is a projection neuron and/or the distance of the axonal ending to the nucleus of the neuron is at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm.

[0017] In another aspect, the present disclosure provides a method of delivering a therapeutic transgene product to a target cell of the CNS, which is a neuron or a glial cell, in a mammal afflicted with a neurometabolic disorder, e.g., a LSD that affects the CNS. The method includes contacting an axonal ending of a neuron with a composition containing an AAV vector carrying at least a part of a gene encoding a therapeutic transgene product, allowing the viral particle to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron; allowing the therapeutic transgene product to be expressed and secreted by the neuron, and allowing the target cell to uptake the therapeutic transgene product, wherein the

therapeutic transgene product thereby alleviates pathology in the target cell. In certain instances, the concentration of the vector in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml).

[0018] In the methods of the present disclosure, the therapeutic transgene encodes a biologically active molecule, expression of which in the CNS results in at least partial correction of neuropathology. In some instances, the therapeutic transgene product is a lysosomal hydrolase. In an illustrative instance, the lysosomal hydrolase is ASM. In other instances, the therapeutic transgene is a metalloendopeptidase, e.g., neprilysin.

[0019] On the basis of the disclosure that is contained herein, a first aspect of the present invention provides a composition comprising an AAV vector comprising a transgene encoding a lysosomal hydrolase, for use in treating a lysosomal storage disease in a mammal, wherein the composition is to be administered to a first site in the CNS of the mammal and the concentration of the vector in the composition is at least 5×10^{12} gp/ml such that the lysosomal hydrolase is expressed in a therapeutically effective amount in a second site in the CNS which contains neurons that project to the first site and which is distal to, and at least 2 mm from, said first site.

[0020] In a further aspect, the invention provides for the use of an AAV vector comprising a recombinant AAV genome comprising a transgene encoding a lysosomal hydrolase for the preparation of a composition for treating a lysosomal storage disease in a mammal, wherein the composition is to be contacted with an axonal ending of a disease-compromised neuron such that the AAV vector is endocytosed and retrogradely transported intracellularly along the axon of the neuron to the nucleus of the neuron, and wherein the neuron is a projection neuron and the distance from the axonal ending to the nucleus is at least 2mm.

[0021] Additional aspects and embodiments of the present invention are set forth in the appended claims.

[0022] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention as claimed.

BRIEF DESCRIPTION OF THE FIGURES

[0023]

Figure 1A depicts a representation of a cross-section of the ASMKO mouse brain, at 5 or 15 weeks following a 2 μ l injection of high titer (9.3×10^{12} gp/ml) AAV-ASM into the hippocampus. The site of injection is shown by a vertical line; ASM mRNA expression, as detected by in situ hybridization, is represented by the smaller circles; and ASM protein expression, as detected by immunohistochemical staining, is represented by the larger shaded circles. The expression pattern resulted in an extensive area of reversal of pathology (represented by the light shading) in the hippocampus and cortical regions in both hemispheres of the brain.

Figure 1B depicts the axonal transport of AAV to distal regions of the mouse brain following a high titer AAV injection into the hippocampus as described for Figure 1A. Injection into the hippocampus (10) resulted in axonal transport of the viral vector via the intrahippocampal circuit to the contralateral hippocampus (20) and via the entorhinodentate circuit to the entorhinal cortex (30). The site of injection is shown by a vertical line.

Figure 1C is a schematic diagram showing the connections of the intrahippocampal and entorhinodentate circuits of the mouse brain. Injection into the hippocampus (10) results in infection and transduction of cell bodies located in the cornu ammonis area 3 (CA3) and in the dentate granule cell layer (G). In addition, a subset of the injected AAV vector infects the axonal endings of the projection neurons innervating the injection site, undergoes retrograde axonal transport, and delivers the transgene to the CA3 field (CA3) and hilus (H) in the contralateral part of the hippocampus (20), and ipsilaterally in the entorhinal cortex (30).

Figure 2A depicts a representation of a cross-section of the ASMKO mouse brain, at 5 or 15 weeks following an intrahippocampal injection of high titer AAV-ASM as described in Figure 1A. ASM mRNA expression, as detected by in situ hybridization, is represented by the smaller circles; and ASM protein expression, as detected by immunohistochemical staining, is represented by the larger shaded circles. The injection resulted in ASM mRNA and protein to be detected in the septum. This expression pattern resulted in an extensive area of reversal of pathology (represented by the light shading).

Figure 2B depicts the axonal transport of AAV to distal regions of the mouse brain, following a high titer injection into the hippocampus as described in Figure 1A. Injection into the hippocampus (10) resulted in axonal transport of the viral vector via the

septohippocampal circuit from the injection site (represented by a vertical line) to the septum (40).

Figure 2C is a schematic diagram showing the connections of the septohippocampal circuit. Injection into the hippocampus resulted in transduction to cell bodies located in the CA3 field (11). In addition, a subset of the AAV vector infects the axonal endings of the projection neurons innervating the injection site, undergoes retrograde axonal transport, and delivers the transgene to the medial septum (40).

Figure 3 depicts the axonal transport of AAV in the nigrostriatal circuit, following a high titer injection of AAV into the striatum (50) of the mouse brain. Axonal transport of AAV occurs from the injection site (represented by a vertical line) to the substantia nigra (60).

Figure 4 depicts the axonal transport of AAV in the medullocerebellar circuit, following a high titer injection of AAV-ASM into the cerebellum (70) of the ASMKO mouse brain. Axonal transport of AAV2 occurs from the injection site (represented by a vertical line) to the medulla oblongata (80).

Figure 5 depicts axonal transport of AAV in the intrahippocampal, nigrostriatal, and entorhinodentate circuits following high-titer injection of AAV7-ASM in the ipsilateral hippocampus (10). Transduced cells were detected, as determined by in situ hybridization, along the entire rostral-caudal axis of the contralateral hippocampus (90), medial septum (40), and entorhinal cortex (100) after AAV7-ASM injection of the ipsilateral hippocampus (represented by a vertical line).

DETAILED DESCRIPTION OF THE INVENTION

[0024] In order that the present disclosure, including the present invention, may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

[0025] The term "transgene" refers to a polynucleotide that is introduced into a cell and is capable of being expressed under appropriate conditions and confers a desired property to a cell into which it was introduced, or otherwise leads to a desired therapeutic outcome.

[0026] The terms "genome particles (gp)," or "genome equivalents," as used in reference to a viral titer, refer to the number of virions containing the recombinant AAV DNA genome, regardless of infectivity or functionality. The number of genome particles in a particular vector preparation can be measured by procedures such as described in the Examples herein, or for example, in Clark et al. (1999) *Hum. Gene Ther.*, 10:1031-1039; Veldwijk et al. (2002) *Mol. Ther.*, 6:272-278.

[0027] The terms "infection unit (iu)," "infectious particle," or "replication unit," as used in reference to a viral titer, refer to the number of infectious and replication-competent recombinant AAV vector particles as measured by the infectious center assay, also known as replication center assay, as described, for example, in McLaughlin et al. (1988) *J. Virol.*, 62:1963-1973.

[0028] The term "transducing unit (tu)" as used in reference to a viral titer, refers to the number of infectious recombinant AAV vector particles that result in the production of a functional transgene product as measured in functional assays such as described in Examples herein, or for example, in Xiao et al. (1997) *Exp. Neurobiol.*, 144:113-124; or in Fisher et al. (1996) *J. Virol.*, 70:520-532 (LFU assay).

[0029] The terms "therapeutic," "therapeutically effective amount," and their cognates refer to that amount of a compound that results in prevention or delay of onset or amelioration of symptoms of neurometabolic disorder in a subject or an attainment of a desired biological outcome, such as correction of neuropathology, e.g., cellular pathology associated with a lysosomal storage disease such as that described herein or in Walkley (1998) *Brain Pathol.*, 8:175-193. The term "therapeutic correction" refers to that degree of correction that results in prevention or delay of onset or amelioration of symptoms of neurometabolic disorder in a subject. The effective amount can be determined by methods well-known in the art and as described in the subsequent sections.

Methods and Compositions

[0030] In the experiments leading to the present disclosure, including the present invention, acid sphingomyelinase (ASM) knockout mice (ASMKO) were used. ASMKO mice have been previously described and are an accepted model of types A and B Niemann-Pick disease (Horinouchi et al. (1995) *Nat. Genetics*, 10:288-293; Jin et al. (2002) *J. Clin. Invest.*, 109:1183-1191; and

Otterbach (1995) *Cell*, 81:1053-1061). Niemann-Pick disease (NPD) is classified as a lysosomal storage disease and is an inherited neurometabolic disorder characterized by a genetic deficiency in acid sphingomyelinase (ASM; sphingomyelin cholinephosphohydrolase, EC 3.1.3.12). The lack of functional ASM protein results in the accumulation of sphingomyelin substrate within the lysosomes of neurons and glia throughout the brain. This leads to the formation of large numbers of distended lysosomes in the perikaryon, which is a hallmark feature and the primary cellular phenotype of type A NPD. The presence of distended lysosomes correlates with the loss of normal cellular function and a progressive neurodegenerative course that leads to death of the affected individual in early childhood (*The Metabolic and Molecular Bases of Inherited Diseases*, eds. Scriver et al., McGraw-Hill, New York, 2001, pp. 3589-3610). Secondary cellular phenotypes (e.g., additional metabolic abnormalities) are also associated with this disease, notably the high level accumulation of cholesterol in the lysosomal compartment. Sphingomyelin has strong affinity for cholesterol, which results in the sequestering of large amounts of cholesterol in the lysosomes of ASMKO mice and human patients (Leventhal et al. (2001) *J. Biol. Chem.*, 276:44976-44983; Slotte (1997) *Subcell. Biochem.*, 28:277-293; and Viana et al. (1990) *J. Med. Genet.*, 27:499-504.)

[0031] The present disclosure, including the present invention, is based, in part, on the discovery and demonstration that an intrahippocampal injection of high titer AAV-ASM into the diseased brains of ASMKO mice results in expression of ASM mRNA and protein distally from the injection site in a pattern consistent with the topographical organization of the projection neurons that innervate the injection site. In addition to robust expression at the site of injection, ASM mRNA and protein are also detected in several distal regions outside of the ipsilateral (injected) hippocampus, specifically, in the contralateral hippocampal dentate gyrus and CA3, and the medial septum and entorhinal cortex. The present disclosure, including the present invention, is further based, in part, on the discovery and demonstration of the extensive correction of lysosomal storage pathology at the distal sites thereby allowing a larger volume of correction via a smaller number of injection sites.

[0032] Accordingly, in one aspect, the present disclosure provides methods for treating neurometabolic disorders in mammals. The populations treated by the methods of the present disclosure include, but are not limited to, patients having or at risk for developing a neurometabolic disorder, e.g., a LSD, such as diseases listed in Table 1, particularly, if such a disease affects the CNS. In an illustrative instance, the disease is type A Niemann-Pick disease. In certain instances, neurometabolic disorders may exclude Alzheimer's, Parkinson, Huntington, Tay Sachs, Lesch-Nyan, and Creutzfeldt-Jakob diseases. However, methods of the present disclosure utilizing a metalloendopeptidase as a therapeutic transgene, are specifically directed to the treatment of Alzheimer's disease and amyloid-related disorders.

[0033] In some instances, the method of treating a neurometabolic disorder comprises administration of a high titer AAV vector carrying a therapeutic transgene so that the transgene product is expressed at a therapeutic level in a second site within the CNS distal to the first site. In some instances, the viral titer of the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). In further instances, the administration is accomplished by direct intraparenchymal injection of a high titer AAV vector solution into the diseased brain, thereafter the transgene is expressed distally, contralaterally or ipsilaterally, to the administration site at a therapeutic level at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm from the administration site.

[0034] The distance between the first and the second sites is defined as the minimal distance region between the site of administration (first site) and the boundary of the detectable transduction of the distal site (second site) as measured using procedures known in the art or as described in the Examples, e.g., in situ hybridization. Some neurons in the CNS of larger mammals may span large distances by virtue of their axonal projections. For example, in humans, some axons may span a distance of 1000 mm or greater. Thus, in various methods of the present disclosure, AAV can be axonally transported along the entire length of the axon at such a distance to reach and transduce the parent cell body.

[0035] A site of vector administration within the CNS is chosen based on the desired target region of neuropathology and the topology of brain circuits involved so long as an administration site and the target region have axonal connections. The target region can be defined, for example, using 3-D stereotaxic coordinates. In some instances of the present disclosure, e.g., in some embodiments of the present invention, the administration site is chosen so that at least 0.1, 0.5, 1, 5, or 10 % of the total amount of vector injected is delivered distally at the target region of at least 1, 200, 500, or 1000 mm³. An administration site may be localized in a region innervated by projection neurons connecting distal regions of the brain. For example, the substantia nigra and ventral tegmental area send dense projections to the caudate and putamen (collectively known as the striatum). Neurons within the substantia nigra and ventral tegmentum can be targeted for transduction by retrograde transport of AAV following injection into the striatum. As another example, the hippocampus receives well-defined, predictable axonal projections from other regions of the brain. Other administration sites may be localized, for example, in the spinal cord, brainstem (medulla and pons), mesencephalon, cerebellum, diencephalon (thalamus, hypothalamus), telencephalon (corpus striatum, cerebral cortex, or within the cortex, the occipital, temporal, parietal or frontal lobes), or combinations thereof.

[0036] For identification of structures in the human brain, see, e.g., *The Human Brain: Surface, Three-Dimensional Sectional Anatomy With MRI, and Blood Supply*, 2nd ed., eds. Deuteron et al., Springer Vela, 1999; *Atlas of the Human Brain*, eds. Mai et al., Academic Press; 1997; and *Co-Planar Sterotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging*, eds. Tamarack et al., Thyme Medical Pub., 1988. For identification of structures in the mouse brain, see, e.g., *The Mouse Brain in Sterotaxic Coordinates*, 2nd ed., Academic Press, 2000. If desired, the human brain structure can be correlated to similar structures in the brain of another mammal. For example, most mammals, including humans and rodents, show a similar topographical organization of the entorhinal-hippocampus projections, with neurons in the lateral part of both the lateral and medial entorhinal cortex projecting to the dorsal part or septal pole of the hippocampus, whereas the projection to the ventral hippocampus originates primarily from neurons in medial parts of the entorhinal cortex (*Principles of Neural Science*, 4th ed., eds Kandel et al., McGraw-Hill, 1991; *The Rat Nervous System*, 2nd ed., ed. Paxinos, Academic Press, 1995). Furthermore, layer II cells of the entorhinal cortex project to the dentate gyrus, and they terminate in the outer two-thirds of the molecular layer of the dentate gyrus. The axons from layer III cells project bilaterally to the cornu ammonis areas CA1 and CA3 of the hippocampus, terminating in the stratum lacunose molecular layer.

[0037] The second (target) site can be located any region of the CNS, including the brain and the spinal cord, that contains neurons that project to the first (administration) site. In some instances of the present disclosure, e.g., in some embodiments of the present invention, the second site is in a region of the CNS chosen from the substantia nigra, the medulla oblongata, or the spinal cord.

[0038] To deliver the vector specifically to a particular region of the central nervous system, especially to a particular region of the brain, it may be administered by stereotaxic microinjection. For example, on the day of surgery, patients will have the stereotaxic frame base fixed in place (screwed into the skull). The brain with stereotaxic frame base (MRI-compatible with fiducial markings) will be imaged using high resolution MRI. The MRI images will then be transferred to a computer that runs stereotaxic software. A series of coronal, sagittal and axial images will be used to determine the target site of vector injection, and trajectory. The software directly translates the trajectory into 3-dimensional coordinates appropriate for the stereotaxic frame. Burr holes are drilled above the entry site and the stereotaxic apparatus localized with the needle implanted at the given depth. The vector in a pharmaceutically acceptable carrier will then be injected. The AAV vector is then administered by direct injection to the primary target site and retrogradely transported to distal target sites via axons. Additional routes of administration may be used, e.g., superficial cortical application under direct visualization, or other non-stereotaxic application.

[0039] The total volume of material to be administered, and the total number of vector particles to be administered, will be determined by those skilled in the art based upon known aspects of gene therapy. Therapeutic effectiveness and safety can be tested in an appropriate animal model. For example, a variety of well-characterized animal models exist for LSDs, e.g., as described herein or in Watson et al. (2001) *Methods Mol. Med.*, 76:383-403; or Jeyakumar et al. (2002) *Neuropath. Appl. Neurobiol.*, 28:343-357.

[0040] In experimental mice, the total volume of injected AAV solution is for example, between 1 to 5 μ l. For other mammals, including the human brain, volumes and delivery rates are appropriately scaled. For example, it has been demonstrated that volumes of 150 μ l can be safely injected in the primate brain (Janson et al. (2002) *Hum. Gene Ther.*, 13:1391-1412). Treatment may consist of a single injection per target site, or may be repeated along the injection tract, if necessary. Multiple injection sites can be used. For example, in some instances of the present disclosure, in addition to the first administration site, a composition comprising AAV carrying a transgene is administered to another site which can be contralateral or ipsilateral to the first administration site.

[0041] In another aspect, the present disclosure provides a method of delivering a recombinant AAV genome via retrograde axonal transport to the nucleus of a disease-compromised neuron in vivo. In some instances, the cellular pathology exhibited by a neuron is that of a LSD such as listed in Table 1 (see, e.g., Walkley (1998) *Brain Pathol.*, 8:175-193). In an illustrative instance, the disease is Niemann-Pick A disease. The method of delivering a recombinant AAV genome to the nucleus of a disease-compromised neuron comprises contacting an axonal ending of a disease-compromised neuron with a composition comprising an AAV viral particle comprising the recombinant AAV genome and allowing the viral particle to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron, wherein the concentration of AAV genomes in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). In certain instances, the neuron is a projection neuron and/or the distance from the axonal ending to the nucleus of the neuron is at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm. In various instances, the AAV genome is transported along the entire length of the axon, at distances varying depending on the axon length. In humans, these distances may be as much as 1000 mm or greater.

[0042] In another aspect, the present disclosure provides a method of delivering a transgene product to a target cell of the CNS, which is a neuron or a glial cell, in a mammal afflicted with a neurometabolic disorder, e.g., a LSD as listed in Table 1. The method comprises contacting an axonal ending of a neuron with a composition comprising an AAV vector carrying at least a part of a gene encoding a therapeutic transgene product; allowing the viral particles to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron; allowing the transgene product to be expressed and secreted by the neuron; and allowing a second cell to uptake the transgene product, wherein the transgene product thereby alleviates pathology in the second cell. In some instances, the concentration of the AAV vector in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). For example, lysosomal hydrolases can be secreted by transduced cells and subsequently taken up by another cell via mannose-6-phosphate receptor-mediated endocytosis, the second cell being transduced or non-transduced (Sando et al. (1977) *Cell*, 12:619-627; Taylor et al. (1997) *Nat. Med.*, 3:771-774; Miranda et al. (2000) *Gene Ther.*, 7:1768-1776; and Jin et al. (2002) *J. Clin. Invest.*, 109:1183-1191).

[0043] In the methods of the present disclosure, AAV of any serotype can be used so long as the vector is capable of undergoing retrograde axonal transport in a disease-compromised brain. The serotype of the viral vector used in certain instances of the present disclosure, and in certain embodiments of the invention, is selected from the group consisting from AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, and AAV8 (see, e.g., Gao et al. (2002) *PNAS*, 99:11854-11859; and *Viral Vectors for Gene Therapy: Methods and Protocols*, ed. Machida, Humana Press, 2003). Other serotype besides those listed herein can be used. However, AAV5 may be specifically excluded from the methods of the present disclosure utilizing a metalloendopeptidase, e.g., neprilysin, as a therapeutic transgene.

[0044] AAV vectors are derived from single-stranded (ss) DNA parvoviruses that are nonpathogenic for mammals (reviewed in Muzyska (1992) *Curr. Top. Microb. Immunol.*, 158:97-129). Briefly, AAV-based vectors have the rep and cap viral genes that account for 96% of the viral genome removed, leaving the two flanking 145-basepair (bp) inverted terminal repeats (ITRs), which are used to initiate viral DNA replication, packaging and integration. In the absence of helper virus, wild-type AAV integrates into the human host-cell genome with preferential site-specificity at chromosome 19q 13.3 or it may remain expressed episomally. A single AAV particle can accommodate up to 5 kb of ssDNA, therefore leaving about 4.5 kb for a transgene and regulatory elements, which is typically sufficient. However, trans-splicing systems as described, for example, in United States Patent No. 6,544,785, may nearly double this limit.

[0045] In an illustrative instance of the present disclosure, e.g., in an illustrative embodiment of the present invention, AAV is AAV2. Adeno-associated virus of many serotypes, especially AAV2, have been extensively studied and characterized as gene therapy vectors. Those skilled in the art will be familiar with the preparation of functional AAV-based gene therapy vectors. Numerous references to various methods of AAV production, purification and preparation for administration to human subjects can be found in the extensive body of published literature (see, e.g., *Viral Vectors for Gene Therapy: Methods and Protocols*, ed. Machida, Humana Press, 2003). Additionally, AAV-based gene therapy targeted to cells of the CNS has been described in United States Patent Nos. 6,180,613 and 6,503,888.

[0046] In certain methods of the present disclosure, the vector comprises a transgene operably linked to a promoter. The transgene encodes a biologically active molecule, expression of which in the CNS results in at least partial correction of neuropathology. In some instances, the transgene encodes a lysosomal hydrolase. In an illustrative instance, the lysosomal hydrolase is ASM. The genomic and functional cDNA sequences of human ASM have been published (see, e.g., United States Patent No. 5,773,278 and 6,541,218). Other lysosomal hydrolases can be used for appropriate diseases, for example, as listed in Table 1.

[0047] The present disclosure further provides methods of treating Alzheimer's disease in mammals, including humans. In such methods, the transgene encodes a metalloendopeptidase. The metalloendopeptidase can be, for example, the amyloid-beta degrading enzyme neprilysin (EC 3.4.24.11; sequence accession number, e.g., P08473 (SWISS-PROT)), the insulin-degrading enzyme insulysin (EC 3.4.24.56; sequence accession number, e.g., P14735 (SWISS-PROT)), or thimet oligopeptidase (EC 3.4.24.15; sequence accession number, e.g., P52888 (SWISS-PROT)).

[0048] The level of transgene expression in eukaryotic cells is largely determined by the transcriptional promoter within the transgene expression cassette. Promoters that show long-term activity and are tissue- and even cell-specific are used in some instances of the present disclosure, e.g., in some embodiments of the present invention. Nonlimiting examples of promoters include, but are not limited to, the cytomegalovirus (CMV) promoter (Kapliitt et al. (1994) *Nat. Genet.*, 8:148-154), CMV/human β -globin promoter (Mandel et al. (1998) *J. Neurosci.*, 18:4271-4284), GFAP promoter (Xu et al. (2001) *Gene Ther.*, 8:1323-1332),

the 1.8-kb neuron-specific enolase (NSE) promoter (Klein et al. (1998) *Exp. Neurol.*, 150:183-194), chicken beta actin (CBA) promoter (Miyazaki (1989) *Gene*, 79:269-277) and the β -glucuronidase (GUSB) promoter (Shiple et al. (1991) *Genetics*, 10:1009-1018). To prolong expression, other regulatory elements may additionally be operably linked to the transgene, such as, e.g., the Woodchuck Hepatitis Virus Post-Regulatory Element (WPRE) (Donello et al. (1998) *J. Virol.*, 72, 5085-5092) or the bovine growth hormone (BGH) polyadenylation site.

[0049] For some CNS gene therapy applications, it will be necessary to control transcriptional activity. To this end, pharmacological regulation of gene expression with AAV vectors can be obtained by including various regulatory elements and drug-responsive promoters as described, for example, in Haberma et al. (1998) *Gene Ther.*, 5:1604-16011; and Ye et al. (1995) *Science*, 283:88-91.

[0050] High titer AAV preparations can be produced using techniques known in the art, e.g., as described in United States Patent No. 5,658,776 and *Viral Vectors for Gene Therapy: Methods and Protocols*, ed. Machida, Humana Press, 2003.

[0051] The following examples provide illustrative embodiments of the invention. One of ordinary skill in the art will recognize the numerous modifications and variations that may be performed. The examples do not in any way limit the invention.

EXAMPLES

Titration of Recombinant Vectors

[0052] AAV vector titers were measured according to genome copy number (genome particles per milliliter). Genome particle concentrations were based on Taqman® PCR of the vector DNA as previously reported (Clark et al. (1999) *Hum. Gene Ther.*, 10:1031-1039; Veldwijk et al. (2002) *Mol. Ther.*, 6:272-278). Briefly, purified AAV-ASM was treated with capsid digestion buffer (50mM Tris-HCl pH 8.0, 1.0 mM EDTA, 0.5% SDS, 1.0 mg/ml proteinase K) at 50°C for 1 hour to release vector DNA. DNA samples were put through a polymerase chain reaction (PCR) with primers that anneal to specific sequences in the vector DNA, such as the promoter region, transgene, or the poly A sequence. The PCR results were then quantified by a Real-time Taqman® software, such as that provided by the Perkin Elmer-Applied Biosystems (Foster City, CA) Prism 7700 Sequence Detector System.

[0053] Vectors carrying an assayable marker gene such as the β -galactosidase or green fluorescent protein gene (GFP) can be titered using an infectivity assay. Susceptible cells (e.g., HeLa, or COS cells) are transduced with the AAV and an assay is performed to determine gene expression such as staining of β -galactosidase vector-transduced cells with X-gal (5-bromo-4chloro-3-indolyl- β -D-galactopyranoside) or fluorescence microscopy for GFP-transduced cells. For example, the assay is performed as follows: 4×10^4 HeLa cells are plated in each well of a 24-well culture plate using normal growth media. After attachment, i.e., about 24 hours later, the cells are infected with Ad type 5 at a multiplicity of infection (MOI) of 10 and transduced with serial dilutions of the packaged vector and incubated at 37°C. One to three days later, before extensive cytopathic effects are observed, the appropriate assay is performed on the cells (e.g., X-gal staining or fluorescence microscopy). If a reporter gene such as β -galactosidase is used, the cells are fixed in 2% paraformaldehyde, 0.5% glutaraldehyde and stained for β -galactosidase activity using X-gal. Vector dilutions that give well-separated cells are counted. Each positive cell represents 1 transduction unit (tu) of vector.

Correction of LSD pathology in the mouse brain

[0054] Ten-week old ASMKO mice contain significant NPD pathology in the central nervous system. Identification of homozygous recessive mutants was verified by PCR. Sixteen 10-week old ASMKO mice were anesthetized with isoflurane and mounted on a stereotaxic frame, an incision was made to expose the underlying skull, and a single drill hole was made over one hemisphere of each mouse. Two microliters of high titer (9.3×10^{12} gp/ml) AAV2-CMV-ASM (Targeted Genetics, Seattle, WA) were injected into the hippocampus at a final stereotaxic coordinate of 2.0 mm rostral of bregma, 1.5 mm right of midline, and 2.0 mm ventral to pial surface. This hippocampal coordinates ensured that the AAV2 vector was exposed to neurons of the dentate gyrus and of the cornu ammonis area 3 (CA3), as well as to axonal endings of projection neurons of the contralateral hippocampus, medial septum and entorhinal cortex. The injections were performed at a rate of 0.2 μ l/minute, and a total of 1.86×10^{10} genomic particles was administered into each brain. The mice were then sacrificed at either 5 (n = 8) or 15 (n = 8) weeks postinjection (pi). Eight brains (n = 4 each at 5 and 15 weeks pi) were analyzed for ASM mRNA and protein distribution, and for the reduction of the

supraphysiologic levels of cholesterol in the lysosomes. The remaining 8 brains (n=4 each at 5 and 15 weeks pi) were processed for histopathology to analyze the correction of accumulated and distended lysosomes, which is the most direct and accurate method for determining reversal of storage pathology for LSDs.

[0055] Robust transduction was detected in the injected (ipsilateral) hippocampus at 5 and 15 weeks pi. The granule cell layer and hilus of the dentate gyrus, and the pyramidal and oriens cell layers of CA3 were extensively transduced by the AAV2 vector. This impressive pattern of transduction extended to other regions of the ipsilateral hippocampus, such as the subiculum and cornu ammonis area 1 (CA1) and 2 (CA2). Immunofluorescence with an anti-human ASM monoclonal antibody confirmed the presence of ASM protein in many cells. The overall protein pattern was similar to the mRNA pattern, with some additional localized spread of protein.

[0056] Human ASM mRNA and protein were also detected in regions outside of the ipsilateral hippocampus at both time points. The contralateral dentate gyrus and CA3, and the medial septum and entorhinal cortex were positive for in situ hybridization and immunofluorescence (Figures 1A and 2A). The pattern of transduction at these distal sites was consistent with the topographical organization of the projection neurons that innervate the injection site (Figures 1B and 2B). This demonstrated that AAV2 underwent retrograde axonal transport in the intrahippocampal, septohippocampal and entorhinodentate circuits of ASMKO brains, and that the viral vector was targeted to the nucleus following transport up the axons (Figures 1C and 2C). The pattern of transduction does not support parenchymal diffusion as the reason for AAV2 transport to these distal sites. If such diffusion had occurred, structures between the injected and distal sites would have been exposed to migrating virus. But these intermediate structures were negative for in situ hybridization. For example the striatum, which possesses a strong natural tropism for AAV2, was negative for gene transfer despite being in the direct path between the hippocampus and medial septum. Thus, gene transfer to the distal sites must have arisen by retrograde axonal transport, which indicates that an affected projection neuron can function as an effective highway transport system for AAV2 movement through a diseased brain.

[0057] The ability of ASM to reverse the cholesterol abnormalities in the ASMKO brain additionally was investigated. Filipin is an autofluorescent molecule isolated from *Streptomyces filipinensis* that binds to cholesterol complexes (Leventhal et al. (2001) J. Biol. Chem., 276:44976-44983; and Sarna et al. (2001) Eur. J. Neurosci., 13:1-9). Uninjected ASMKO brains had high levels of filipin staining due to these abundant cholesterol complexes, whereas normal mouse brains produced no filipin staining.

[0058] Injection of AAV2-CMV-ASM resulted in the complete loss of filipin staining throughout the entire ipsilateral and contralateral hippocampus, septum and entorhinal cortex at 5 and 15 weeks pi of ASMKO mice (Figures 1A and 2A). This was in stark contrast to uninjected age-matched ASMKO controls, where high levels of filipin staining were detected in these same structures. The loss of filipin staining in AAV2-injected brains demonstrates that a secondary cellular phenotype (e.g. metabolic defect) of ASM disease was corrected. This strongly suggests that ASM protein was targeted to the lysosome and interacted with the sphingomyelin-cholesterol complex. This interaction likely resulted in the release of cholesterol from sphingomyelin, and subsequent entry of cholesterol into its normal biological pathways (such as degradation or translocation to the plasma membrane (Leventhal et al. (2001) J. Biol. Chem., 276:44976-44983)).

[0059] The loss of filipin staining was observed in all cell layers and subfields of the intrahippocampal, septohippocampal and entorhinodentate circuits. The area of cholesterol correction was far greater and more extensive than the ASM protein pattern. This indicates that, following retrograde axonal transport of AAV2, projection neurons may have functioned as "enzyme pumps" and secreted ASM protein into the surrounding tissue. Significantly, only a small amount of ASM is needed to have a therapeutic effect on cholesterol accumulation within ASMKO-affected cells, an amount below the detection limit of the immunofluorescent protocol.

[0060] Whether axonal transport of the AAV2-ASM vector results in the correction of the primary cellular phenotype of NPD also was evaluated. One-micron-thick histopathology brain sections demonstrated a remarkable reduction of accumulated and distended lysosomal pathology in AAV2-CMV-ASM-injected brains at 5 weeks pi (Table 2). Reversal of pathology resulting in partial or complete restoration of normal cellular architecture occurred in all regions of the ipsilateral and the contralateral hemispheres of the hippocampus. The medial septum and the entorhinal cortex also showed a substantial reduction in storage lesions. Similar to the filipin data, the number of cells corrected was greater and more widespread than the ASM protein pattern. Reversal of pathology was evident within regions known to project to the hippocampus including the intrahippocampal, septohippocampal and entorhinodentate circuits. Overall, the volume of correction was 30-35 mm³ or more in the contralateral hippocampus, 5-8 mm³ or more in the ipsilateral entorhinal cortex, 1-2 mm³ or more in the contralateral entorhinal cortex, and 2-3 mm³ or more in the medial septum. This further supports that axonal transport of the viral vector was responsible for this therapeutic effect, because nearby structures (that do not contribute to these circuits) would have been corrected if viral distribution was mediated merely by diffusion (see, "ipsilateral striatum" and "contralateral striatum" in Table 2).

[0061] To demonstrate that reversal of pathology was specific to ASM, an additional group of ASMKO mice was injected with a control vector carrying a reporter gene, AAV2-CMV-β-gal (n = 2 each at 5 and 15 weeks pi), and processed for histopathology. In all four brains, cells remained inundated with distended lysosomes, and contained other abnormalities such as cytoplasmic swelling and disorganized cellular layers.

Table 2

<i>Brain Region</i>	<i>Untreated</i>	<i>AAV2-ASM Treated</i>
Ipsilateral Hippocampus		
CA1 Field	++++	+
CA3 Field	++++	+
Dentate Granule Cell Layer	++++	+
Hilus	++++	+
Contralateral Hippocampus		
CA1 Field	++++	+
CA3 Field	++++	+
Dentate Granule Cell Layer	++++	+
Hilus	++++	+
Ipsilateral Entorhinal Cortex	++++	++
Contralateral Entorhinal Cortex	++++	+++
Ipsilateral Medial Septum	++++	++
Contralateral Medial Septum	++++	++
Ipsilateral Striatum	++++	++++
Contralateral Striatum	++++	++++
++++ high level of pathology in virtually all cells		
+++ pathology in many cells, correction is observed in some cells		
++ pathology in some cells, correction is observed in many cells		
+ little or no pathology is in most cells, virtually every cell is corrected		

[0062] Thus, in accordance with the present invention, a single injection of high titer AAV2 vector is sufficient to transfer the ASM gene to structures that innervate the ASMKO affected hippocampus. The number of structures positive for AAV2 vector was greater than that demonstrated by a recent study in the normal rat hippocampus, which showed axonal transport only in the entorhino-dentate circuit (Kaspar et al. (2002) Mol. Ther., 5:50-56). The results described herein demonstrate that axonal transport can occur in projection neurons afflicted with storage pathology, and that this mode of transport results in the clearance of storage pathology in proximal structures and multiple regions distal to the injection site. We also demonstrate that axonal transport is not limited to only those circuits associated with the hippocampus. Retrograde axonal transport occurred in the nigrostriatal (Figure 3) and in the medullocerebellar (Figure 4) circuits. This demonstrates that axonal transport of AAV2 in diseased-compromised neurons is a general property of the viral vector.

[0063] A similar study was performed with AAV1-ASM at the concentrations of $1-4 \times 10^{13}$ gp/ml and AAV7-ASM at the concentration of 8.4×10^{12} gp/ml. While AAV1 did not exhibit detectable retrograde axonal transport, AAV7 did undergo retrograde axonal transport, similar to AAV2, and produced correction of LSD pathology in distal regions (see Figure 5).

Treatment of Alzheimer's disease

[0064] Alzheimer's disease (AD) is a neurodegenerative disorder characterized by the accumulation of amyloid β -peptide ($A\beta$) due to decreased $A\beta$ catabolism. As $A\beta$ accumulates, it aggregates into extracellular plaques, causing impairment of synaptic function and loss of neurons. The pathology leads to dementia, loss of coordination, and death.

[0065] Neprilysin is a 97 kD membrane-bound zinc metalloendopeptidase that is the rate-limiting enzyme in the normal degradation of $A\beta$. Introduction of neprilysin may decelerate the progression of the disease by removing $A\beta$ pools before aggregation. Indeed, neprilysin was shown to degrade oligomeric forms of $A\beta$ thereby removing existing plaques in an animal model of AD (Kanemitsu et al. (2003) *Neurosci. Lett.*, 350:113-116). Neprilysin knockout mice exhibit high levels of $A\beta$ (Iwata et al. (2001) *J. Neurosci.*, 24:991-998.). Neprilysin inhibitors, such as thiorphan and phosphoramidon, increase $A\beta$ levels in mouse brain (Iwata et al. (2000) *Nat. Med.*, 6:143-150). Additionally, decreased neprilysin mRNA levels were found in areas of high amyloid plaque burden in human brains, further demonstrating the link between neprilysin and AD (Yasojima et al. (2001) *Neurosci. Lett.*, 297:97-100).

[0066] The areas of brain most affected by AD are the hippocampus, cortex, cerebellum, striatum and thalamus (see, e.g., Iwata et al. (2001) *supra*; Yasojima et al. (2001) *supra*). These are the same areas of the brain that show efficient retrograde axonal transport with AAV.

[0067] Accordingly, AAV can be used to deliver therapeutic transgenes to regions of high plaque burden by direct injection and subsequent translocation of virus through brain circuits to our target sites. Viral vector-mediated gene transfer of neprilysin was effective in treating mouse models of AD (Marr et al. (2003) *J. Neurosci.*, 23:1992-1996; Marr et al. (2004) *J. Mol. Neurosci.*, 22:5-11; Iwata et al. (2004) *J. Neurosci.*, 24:991-998). A recent report showed that AAV5-neprilysin removed $A\beta$ from the pre-synaptic terminals of the hippocampus in neprilysin-deficient mice, decelerating plaque formation at the synapses (Iwata et al. (2004) *supra*). In this report, neprilysin was found in the contralateral hippocampus but whether this is attributable to retrograde transport of virus or anterograde transport of expressed protein remains unknown.

[0068] The specification is most thoroughly understood in light of the teachings of the references cited within the specification. The embodiments within the specification provide an illustration of embodiments of the invention and should not be construed to limit the scope of the invention. The skilled artisan readily recognizes that many other embodiments are encompassed by the invention. To the extent the publications, patents, and biological sequences cited in this disclosure contradict or are inconsistent with the present specification, the present specification will supercede any such material. The citation of any references herein is not an admission that such references are prior art to the present invention.

[0069] Unless otherwise indicated, all numbers expressing quantities of ingredients, cell culture, treatment conditions, and so forth used in the specification, including claims, are to be understood as being modified in all instances by the term "about." Accordingly, unless otherwise indicated to the contrary, the numerical parameters are approximations and may vary depending upon the desired properties sought to be obtained by the present invention. Unless otherwise indicated, the term "at least" preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US46719503P \[0001\]](#)
- [US47572603P \[0001\]](#)

- [US6544785B \[0044\]](#)
- [US6160613E \[0045\]](#)
- [US6503888E \[0045\]](#)
- [US5773276A \[0046\]](#)
- [US6541219E \[0046\]](#)
- [US658776A \[0050\]](#)

Non-patent literature cited in the description

- **PARDRIDGE** Peptide Drug Delivery to the Brain Raven Press 19910000 [0004]
- **DAVIDSON et al.** Nature Rev., 2003, vol. 4, 353-364 [0005]
- **KAPLITT et al.** Nat. Genet., 1994, vol. 8, 148-154 [0005] [0008] [0048]
- **BARTLETT et al.** Hum. Gene Ther., 1998, vol. 9, 1181-1186 [0005]
- **PASSINI et al.** J. Neurosci., 2002, vol. 22, 6437-6446 [0005]
- **NEUFELD et al.** Science, 1970, vol. 169, 141-146 [0006]
- **TAYLOR et al.** Nat. Med., 1997, vol. 3, 771-774 [0006]
- **SKORUPA et al.** Exp. Neurol., 1999, vol. 160, 17-27 [0006]
- **SOUDAS et al.** FASEB J., 2001, vol. 15, 2283-2285 [0007]
- **BREAKEYFIELD et al.** New Biol., 1991, vol. 3, 203-218 [0007]
- **DEFALCO et al.** Science, 2001, vol. 291, 2608-2613 [0007]
- **PASSINI et al.** J. Neurosci., 2002, vol. 22, 6437-6446 [0009]
- **CHAMBERLIN et al.** Brain Res., 1998, vol. 793, 169-175 [0008]
- **KASPAR et al.** Mol. Ther., 2002, vol. 5, 50-56 [0008] [0062]
- **WALKLEY** Brain Pathol., 1998, vol. 8, 175-193 [0008] [0029] [0041]
- **CLARK et al.** Hum. Gene Ther., 1999, 1031-1039 [0026]
- **VELDWIJK et al.** Mol. Ther., 2002, vol. 6, 272-278 [0026] [0052]
- **MCLAUGHLIN et al.** J. Virol., 1988, vol. 62, 1963-1973 [0027]
- **XIAO et al.** Exp. Neurobiol., 1997, vol. 144, 113-124 [0028]
- **FISHER et al.** J. Virol., 1996, vol. 70, 520-532 [0028]
- **HORINOUCHE et al.** Nat. Genetics, 1995, vol. 10, 288-293 [0030]
- **JIN et al.** J. Clin. Invest., 2002, vol. 109, 1183-1191 [0030] [0042]
- **OTTERBACH** Cell, 1995, vol. 81, 1053-1061 [0030]
- The Metabolic and Molecular Bases of Inherited Diseases McGraw-Hill 200100003589-3610 [0030]
- **LEVENTHAL et al.** J. Biol. Chem., 2001, vol. 276, 44976-44983 [0030] [0058]
- **SLOTTE** Subcell. Biochem., 1997, vol. 28, 277-293 [0030]
- **VIANA ET LAJ.** Med. Genet., 1990, vol. 27, 499-504 [0030]
- The Human Brain: Surface, Three-Dimensional Sectional Anatomy With MRI, and Blood Supply Springer Vela 19990000 [0036]
- Atlas of the Human Brain Academic Press 19970000 [0036]
- Co-Planar Sterotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging Thyme Medical Pub. 19880000 [0036]
- The Mouse Brain in Sterotaxic Coordinates Academic Press 20000000 [0036]
- Principles of Neural Science McGraw-Hill 19910000 [0036]
- The Rat Nervous System Academic Press 19950000 [0036]
- **WATSON et al.** Methods Mol. Med., 2001, vol. 76, 383-403 [0039]
- **JEYAKUMAR et al.** Neuropath. Appl. Neurobiol., 2002, vol. 28, 343-357 [0039]
- **JANSON et al.** Hum. Gene Ther., 2002, vol. 13, 1391-1412 [0040]
- **SANDO et al.** Cell, 1977, vol. 12, 619-627 [0042]
- **TAYLOR et al.** Nat. Med., 1997, vol. 3, 771-774 [0042]
- **MIRANDA et al.** Gene Ther., 2000, vol. 7, 1768-1776 [0042]
- **GAO et al.** PNAS, 2002, vol. 99, 11854-11859 [0043]
- Viral Vectors for Gene Therapy: Methods and Protocols Humana Press 20030000 [0043] [0045] [0050]
- **MUZYSCKA** Curr. Top. Microb. Immunol., 1992, vol. 158, 97-129 [0044]
- **MANDEL et al.** J. Neurosci., 1998, vol. 18, 4271-4284 [0048]

- XU et al. Gene Ther., 2001, vol. 8, 1323-1332 [0048]
- KLEIN et al. Exp. Neurol., 1998, vol. 150, 183-194 [0048]
- MIYAZAKI Gene, 1989, vol. 79, 269-277 [0048]
- SHIPLEY et al. Genetics, 1991, vol. 10, 1009-1018 [0048]
- DONELLO et al. J. Virol., 1998, vol. 72, 5085-5092 [0048]
- HABERMA et al. Gene Ther., 1998, vol. 5, 1604-16011 [0049]
- YE et al. Science, 1995, vol. 283, 88-91 [0049]
- CLARK et al. Hum. Gene Ther., 1999, vol. 10, 1031-1039 [0052]
- LEVENTHAL et al. J. Biol. Chem., 2001, 44976-44983 [0057]
- SARNA et al. Eur. J. Neurosci., 2001, vol. 13, 1-9 [0057]
- KANEMITSU et al. Neurosci. Lett., 2003, vol. 350, 113-116 [0065]
- IWATA et al. J. Neurosci., 2001, vol. 24, 991-998 [0065]
- IWATA et al. Nat. Med., 2000, vol. 6, 143-150 [0065]
- YASOJIMA et al. Neurosci. Lett., 2001, vol. 297, 97-100 [0065]
- MARR et al. J. Neurosci., 2003, vol. 23, 1992-1996 [0067]
- MARR et al. J. Mol. Neurosci., 2004, vol. 22, 5-11 [0067]
- IWATA et al. J. Neurosci., 2004, vol. 24, 991-998 [0067]

Patentkrav

1. Sammensætning, som omfatter en AAV-vektor, der omfatter et transgen, hvilket transgen koder for en lysosomal
5 hydrolase, til anvendelse til behandling af en lysosomal ophobningssygdom hos et pattedyr, hvor sammensætningen skal administreres til et første sted i pattedyrets CNS, og koncentrationen af vektoren i sammensætningen er mindst 5×10^{12} gp/ml, således at den lysosomale hydrolase udtrykkes i en
10 terapeutisk effektiv mængde på et andet sted i CNS, der indeholder neuroner, som rager hen mod det første sted, og som er distale i forhold til, og mindst 2 mm fra, det første sted.
2. Sammensætning til anvendelse ifølge krav 1, hvor den
15 lysosomale ophobningssygdom er Niemann-Picks sygdom, type A.
3. Sammensætning til anvendelse ifølge krav 1, hvor pattedyret er et menneske.
- 20 4. Sammensætning til anvendelse ifølge krav 1, hvor sammensætningen skal administreres ved direkte injektion ind i hjernen.
5. Sammensætning til anvendelse ifølge krav 4, hvor det
25 andet sted er kontralateralt i forhold til det første sted.
6. Sammensætning til anvendelse ifølge krav 1, hvor det første sted forefindes i en region af CNS, som er valgt fra gruppen bestående af rygmærken, hjernestammen, hippocampus,
30 striatum, medulla, hjernebroen, mesencephalon, cerebellum, thalamus, hypothalamus, hjernebarken, occipitallappen, tindingelappen, parietallappen og frontallappen.
7. Sammensætning til anvendelse ifølge krav 4, hvor det
35 første sted forefindes i hippocampus, og det andet sted forefindes i en region af hjernen, som er valgt fra gruppen bestående af den kontralaterale gyrus dentatus og den kontralaterale CA3 og den mediale septum og den entorhinale

cortex.

8. Sammensætning til anvendelse ifølge krav 4, hvor det første sted forefindes i en region af hjernen, som er valgt fra gruppen bestående af striatum og cerebellum, og det andet sted forefindes i en region af hjernen, som er valgt fra gruppen bestående af substantia nigra og medulla oblongata.

9. Sammensætning til anvendelse ifølge krav 1, hvor sammensætningen skal administreres til et tredje sted i pattedyrets CNS.

10. Sammensætning til anvendelse ifølge krav 9, hvor det tredje sted er kontralateralt i forhold til det første sted.

11. Sammensætning til anvendelse ifølge krav 1, hvor den lysosomale hydrolase er aspartylglucosaminidase, α -galactosidase A, palmitoylproteinthioesterase, tripeptidylpeptidase, lysosomalt transmembranprotein, cysteintransportør, sur ceramidase, sur α -L-fucosidase, beskyttende protein/cathepsin A, sur β -glucosidase eller glucocerebrosidase, sur β -galactosidase, iduronat-2-sulfatase, α -L-iduronidase, galactocerebrosidase, sur α -mannosidase, sur β -mannosidase, arylsulfatase B, arylsulfatase A, N-acetylgalactosamin-6-sulfatsulfatase, N-acetylglucosamin-1-phosphotransferase, sur sphingomyelinase, NPC-1, sur α -glucosidase, β -hexosaminidase B, heparan N-sulfatase, α -N-acetylglucosaminidase, acetyl-CoA: α -glucosaminid-N-acetyltransferase, N-acetylglucosamin-6-sulfatsulfatase, α -N-acetylgalactosaminidase, α -neuramidase, β -glucuronidase, β -hexosaminidase A eller sur lipase.

12. Sammensætning til anvendelse ifølge krav 1, hvor den lysosomale hydrolase er sur sphingomyelinase.

13. Anvendelse af en AAV-vektor, der omfatter et rekombinant AAV-genom, hvilket genom omfatter et transgen, hvilket transgen koder for en lysosomal hydrolase, til fremstilling af

en sammensætning til behandling af en lysosomal ophobningssygdom hos et pattedyr, hvor sammensætningen skal bringes i kontakt med en axonende af en sygdomskompromitteret neuron, således at AAV-vektoren endocyteres og retrogradt
5 transporteres intracellulært langs neuronens axon til neuronens kerne, og den lysosomale hydrolase udtrykkes og secernerer af neuronen og optages af en målcelle, så patologi i målcellen derved mildnes, hvor koncentrationen af de rekombinante AAV-genomer i sammensætningen er mindst 5×10^{12}
10 gp/ml, og hvor neuronen er en projektionsneuron.

14. Anvendelse ifølge krav 13, hvor målcellen er en anden neuron eller en gliacelle.

15 15. Anvendelse ifølge krav 13, hvor afstanden fra axonenden til kernen er mindst 2 mm.

16. Anvendelse ifølge krav 13, hvor den lysosomale ophobningssygdom er Niemann-Picks sygdom, type A.

20

17. Anvendelse ifølge krav 13, hvor den lysosomale hydrolase er sur sphingomyelinase.

18. Sammensætning til anvendelse ifølge et hvilket som helst
25 af kravene 1-12, hvor nævnte AAV har en serotype, som er valgt fra gruppen bestående af AAV2, AAV3, AAV4, AAV5, AAV6, AAV7 og AAV8.

19. Anvendelse ifølge et hvilket som helst af kravene 13-17,
30 hvor nævnte AAV har en serotype, som er valgt fra gruppen bestående af AAV2, AAV3, AAV4, AAV5, AAV6, AAV7 og AAV8.

DRAWINGS




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 2C

