



(51) International Patent Classification:

C07K 14/54 (2006.01) A61K 38/20 (2006.01)
C07K 14/525 (2006.01) A61K 47/48 (2006.01)
C07K 16/18 (2006.01)

(21) International Application Number:

PCT/EP2012/062705

(22) International Filing Date:

29 June 2012 (29.06.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/539,131 26 September 2011 (26.09.2011) US

(71) Applicant (for all designated States except US): **PHILO-GEN S.P.A.** [IT/IT]; La Lizza 7, Siena, I-53100 Siena (IT).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **SCHWAGER, Kathrin** [CH/CH]; Stauffacherstrasse 35, CH-8004 Zurich (CH).(74) Agents: **KING, Hilary** et al.; Mewburn Ellis LLP, 33 Gutter Lane, London, Greater London EC2V 8AS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: IMMUNOCYTOKINE COMBINATION THERAPY




Figure 1B

(57) Abstract: Methods and compositions for treating tumours, especially skin tumours, by locally administering single doses of tumour necrosis factor alpha (TNF α) and interleukin-2 (IL2) at the tumour site, where the TNF α and IL2 are delivered as immunoconjugates comprising an antibody targeted to a splice isoform of an extracellular matrix component such as fibronectin.

Immunocytokine Combination Therapy

Field of the Invention

This invention relates to combination therapy for tumours in which a $\text{TNF}\alpha$ immunoconjugate and an IL2 immunoconjugate are administered directly to the tumour site.

5 Background

Tumour necrosis factor alpha ($\text{TNF}\alpha$) is a cytokine produced by many cell types, mainly activated monocytes and macrophages. It is expressed as a 26 kDa integral transmembrane precursor protein from which a mature protein of approximately 17 kDa is released by proteolytic cleavage. The soluble bioactive $\text{TNF}\alpha$ is a homotrimer that binds cell surface
10 receptors. $\text{TNF}\alpha$ has been shown to induce necrosis of solid tumours. It exerts its effects mainly on the endothelium of the tumour-associated vasculature, with increased permeability, upregulation of tissue factor, fibrin deposition and thrombosis, and massive destruction of the endothelial cells.

Interleukin-2 (IL2), a four α helix bundle cytokine produced by T helper 1 cells, plays an
15 essential role in the activation phases of both adaptive and innate immune responses. Although it is not believed to have a direct cytotoxic effect on cancer cells, it has been reported to induce tumour regression by stimulating a cell-mediated immune response.

Intratumoural injections of IL2 have been trialled in metastatic melanoma patients [1]. In that study, treatment was administered three times weekly for at least 2 weeks, and overall
20 69 % of patients were reported to achieve a complete response.

WO01/66298 described immunoconjugates comprising $\text{TNF}\alpha$ and IL2 respectively, fused to antibody L19. L19 specifically binds the ED-B domain of fibronectin isoform B-FN, which is one of the best known markers angiogenesis (US 10/382,107; WO01/62298). ED-B is an extra domain of 91 amino acids found in the B-FN isoform and is identical in mouse, rat,
25 rabbit, dog and man. B-FN accumulates around neovascular structures in aggressive tumours and other tissues undergoing angiogenesis, such as the endometrium in the proliferative phase and some ocular structures in pathological conditions, but is otherwise undetectable in normal adult tissues.

Carnemolla *et al.* [2] described enhancement of the antitumour properties of IL2 by its
30 targeted delivery to the tumour blood vessel extracellular matrix in an L19-IL2 immunoconjugate.

Christ *et al.* [3] described intratumoural administration of an IL2 immunoconjugate, a $\text{TNF}\alpha$ immunoconjugate, or antibody alone. The antibody used was anti-EGFR, which had an

anti-tumour effect. An anti-tumour immune response was reported following multiple injections of either fusion protein.

Borsi *et al.* [4] reported a study in which L19-TNF α and L19-IL2 immunoconjugates were administered intravenously to mice on days 7 and 10 following implantation of tumour cells. L19 was used to concentrate and maximise the anti-tumour effects of the systemically delivered cytokines. The combination of immunocytokines was reported to have a synergistic effect on tumour volume. Mice who received the combination treatment had markedly reduced tumour volume compared with those who received just one immunocytokine.

Summary of the Invention

Reported here are unexpected effects on tumours resulting from local administration of a combination of immunocytokines, TNF α -L19 and IL2-L19, at the tumour site. A single administration of these two immunocytokines promoted the complete eradication of large subcutaneous tumours in mice.

Mice received a single intratumoural injection of the TNF α -L19, a single intratumoural injection of IL2-L19, or the combination. No further treatments were given. Tumour volume was measured daily and it was observed that tumours in mice treated with the combination therapy rapidly reduced in size and appeared to be completely eliminated within a period of days and showed no regrowth. Compared with saline-treated control mice, mice treated with one cytokine alone also showed inhibition of tumour growth, but tumours in these mice nevertheless continued to increase slowly in size. These results show a synergistic effect of the combined cytokine therapy, and a remarkable therapeutic effect, in which tumours were eradicated following just a single dose of each cytokine.

Accordingly, a first aspect of the invention is a method of treating a tumour in a patient by injecting a single dose of a TNF α immunoconjugate and a single dose of an IL2 immunoconjugate at the tumour site.

The immunoconjugate comprises the cytokine linked to an antibody molecule, which targets the cytokine to the site of the lesion. The antibody molecule binds a splice isoform of an extracellular matrix component, which is selectively expressed by the extracellular matrix in tumour tissue. By combining this targeting effect with direct administration of the immunoconjugate to the tumour site, a very localised administration is achieved, which concentrates the effect of the cytokines at the tumour site and reduces side effects and toxicity associated with systemic use of the cytokines.

A number of splice isoforms of tumour extracellular matrix components are known, and antibody molecules targeting any such isoform may be used to selectively target the tumour. These include splice isoforms of fibronectin, such as B-FN. B-FN includes an extra domain ED-B, and antibody molecules of the invention are preferably targeted to this domain. A preferred

antibody molecule comprises the complementarity determining regions (CDRs) of antibody L19. These are, as illustrated in Figure 3:

VH CDR1	SFSMS	SEQ ID NO: 1
VH CDR 2	SISGSSGTTYADSVKG	SEQ ID NO: 2
VH CDR 3	PFPYFDY	SEQ ID NO: 3
VL CDR 1	RASQSVSSSFLA	SEQ ID NO: 4
VL CDR 2	YASSRAT	SEQ ID NO: 5
VL CDR 3	QQTGRIPPT	SEQ ID NO: 6

The TNF α immunoconjugate preferably comprises TNF α linked to an antibody molecule comprising the L19 CDRs. The IL2 immunoconjugate comprises IL2 linked to an antibody molecule, which may be an identical or different antibody molecule as the TNF α immunoconjugate. The antibody molecule in each immunoconjugate may bind the same extracellular matrix component, optionally the same splice isoform e.g. they may bind the same domain. Preferably, the IL2 immunoconjugate comprises IL2 linked to an antibody molecule comprising the L19 CDRs.

Preferably, the antibody molecule (of the TNF α and/or the IL2 immunoconjugate) comprises the L19 VH domain and/or the L19 VL domain. Amino acid sequences of the L19 VH and VL domains are SEQ ID NO: 7 and SEQ ID NO: 9 respectively (Figure 3).

Preferably the antibody molecule is a single chain Fv (scFv) or other antibody fragment of low molecular weight and/or lacking an Fc region. These properties assist with targeting and tissue penetration of the immunoconjugate at the tumour site. A preferred antibody molecule is scFv-L19, which is an scFv comprising an L19 VH domain and an L19 VL domain, wherein the VH and VL are conjoined in a single polypeptide chain by a peptide linker sequence. The VH domain contains VH CDR1, CDR2 and CDR3 sequences, and the VL domain contains VL CDR1, CDR2 and CDR3 sequences. The VH domain may have an amino acid sequence as set out in Figure 3 (SEQ ID NO: 7). The VL domain may have an amino acid sequence as set out in Figure 3 (SEQ ID NO: 9). The VH and VL domains are normally joined by a peptide linker such as the 12 residue linker shown in Figure 3 (SEQ ID NO: 8). Preferably, the scFv-L19 comprises or consists of the amino acid sequence shown in Figure 3 (SEQ ID NO: 10).

A molecular linker such as a peptide may be used to join the cytokine to the antibody molecule, facilitating expression of all or part of the immunocytokine as a fusion protein. Where the antibody molecule is also a single chain molecule, such as scFv, the entire immunocytokine polypeptide chain may conveniently be produced as a fusion protein. For the TNF α immunoconjugate, the fusion proteins are then assembled into trimers, allowing TNF α to adopt its normal trimeric form [4].

Optionally, the immunocytokine carries a detectable and/or functional label, such as a radioactive isotope. Radiolabelled L19, and its use in cancer therapy, has been described before.

It is generally convenient to provide the IL2 immunoconjugate and the TNF α immunoconjugate as separate molecules. They may be provided as a combined preparation, or as separate formulations to permit either simultaneous or sequential administration. The clinician can determine the most suitable manner of administering the single dose of each immunocytokine to the patient. For example, the method of treatment may comprise injecting the TNF α immunoconjugate and the IL2 immunoconjugate in separate injections, simultaneously or sequentially. Where sequential administration is used, the immunocytokines are preferably injected within 24 hours, 12 hours, 1 hour or more preferably within 30 minutes of each other. The two immunocytokines may be injected at the same point in the tumour site, or at different points. A combined injection of both immunocytokines may be administered. It may be preferable to administer a dose in multiple injections, for example to inject multiple locations across the tumour or around the tumour site, or to facilitate administration of a larger volume of immunocytokine.

The dose is an amount of cytokine, administered at one time, effective to treat the tumour in the combination therapy according to the invention. A single dose may be administered in a treatment period of 1 hour or less, preferably in a period of 30 minutes or less, e.g. 15, 10, 5 or 1 minute or less.

The quantity of TNF α or IL2 administered will depend on the size and nature of the tumour, among other factors. For example, the dose of an TNF α -scFv immunoconjugate may be in the range of 2 – 20 μ g, e.g. 5 – 10 μ g. The dose of IL2-scFv immunoconjugate may be in the range of 10 – 100 μ g, e.g. 20 – 40 μ g. Corresponding doses using other immunoconjugate formats may be straightforwardly calculated to administer an appropriate quantity of cytokine. These are examples only and, of course, different doses may be used. The clinician will determine a therapeutically effective amount for administration.

As reported here, a single dose of the TNF α immunoconjugate and a single dose of the IL2 immunoconjugate were sufficient for tumour therapy. Multiple doses were not required, and treatment of a tumour according to the present invention does not comprise repeating the combination therapy. In addition to the advantages this offers to patients, the single dose regimen provides a considerable advantage to clinicians and significant cost savings.

Accordingly, in treating a particular tumour, the method of the invention is not repeated. The method of treating the tumour may comprise:

(a) injecting a single dose of the TNF α immunoconjugate and a single dose of the IL2 immunoconjugate at the tumour site, and not repeating step (a).

The tumour is treated without any repeated administration of the combination of immunocytokines to the tumour site. As shown herein, a tumour may be treated without any subsequent injection of a TNF α immunoconjugate or an IL2 immunoconjugate. Indeed, the tumour may be treated without administering any further anti-cancer agent to the patient.

5 Optionally, the patient has not previously been given either TNF α or IL2 for the tumour, although in some cases a patient may have received previous therapy with only one of IL2, TNF α or an immunoconjugate including one of these cytokines, which did not achieve complete treatment of the tumour.

Accordingly, a method of the invention may comprise treating a tumour in a patient by
10 injecting a dose of the TNF α immunoconjugate and a dose of the IL2 immunoconjugate at the tumour site, wherein the tumour is treated without administering any subsequent dose of the TNF α immunoconjugate or the IL2 immunoconjugate to the tumour site.

Of course, the method of the invention may be used to treat multiple tumours in a patient, by performing the method on each tumour.

15 Other treatments that may be used in combination with the invention include the administration of suitable doses of pain relief drugs such as non-steroidal anti-inflammatory drugs (e.g. aspirin, paracetamol, ibuprofen or ketoprofen) or opiates such as morphine, or anti-emetics.

The immunocytokines are injected at the site of the tumour, preferably by intratumoural
20 injection. Peritumoural injection, e.g. local intradermal injection, is another suitable method for administering the immunocytokine locally to a tumour site.

The treated tumour may be a primary tumour or a metastatic tumour. The invention is particularly suited to treatment of skin tumours, e.g. malignant skin tumour, melanoma or carcinoma, since their location is amenable to direct local injection. Other tumours within the
25 body may also be treated, and injections may be guided to tumours within soft tissue or internal organs, e.g. by sonography [1]. The methods of the invention may also be used in a surgical context, where injection is performed before, during or after tumour surgery.

Treatment of a tumour according to the present invention may include complete eradication of the tumour. The disappearance of any evidence of vital tumour after stopping
30 injections represents complete treatment of the tumour. Disappearance of the tumour may be determined when the tumour has no discernable volume or is no longer visible. Treatment may comprise treatment to eradicate the tumour and prevent tumour regrowth.

A method of treating a tumour according to the present invention may comprise injecting
35 a single dose of the TNF α immunoconjugate and a single dose of the IL2 immunoconjugate at the tumour site, and observing disappearance of the tumour. Absence of tumour regrowth may also be observed.

Patients are preferably monitored during a followup period of at least one month, preferably at least six months or at least a year, after administration of the immunocytokine combination therapy. Disappearance of the tumour, and lack of tumour regrowth, may be observed in the followup period.

5 In the event of tumour recurrence after the followup period, or if other tumours develop, patients may receive a further treatment with immunocytokine combination therapy according to the invention, to remove the further tumour.

For example, a method according to the invention may comprise eradicating a tumour in a patient by injecting a single dose of the $\text{TNF}\alpha$ immunoconjugate and a single dose of the IL2
10 immunoconjugate at the tumour site, wherein the tumour disappears in the absence of further doses of the $\text{TNF}\alpha$ immunoconjugate and/or the IL2 immunoconjugate.

Further aspects of the invention relate to $\text{TNF}\alpha$ and IL2 immunoconjugates for use in any of the methods of the invention described herein. A composition comprising the $\text{TNF}\alpha$ immunoconjugate and/or the IL2 immunoconjugate may be provided for use in a method as
15 described. Compositions may further comprise additional components, such as pharmaceutically acceptable excipients. A composition may comprise the immunocytokines as separate formulations (e.g. separately packaged, optionally in a kit), or as a combined formulation. The formulation may be adapted for intratumoural administration. Use of the $\text{TNF}\alpha$ immunoconjugate and/or the IL2 immunoconjugate for the manufacture of a medicament for use
20 in a method as described herein is another aspect of the invention.

Nucleic acid molecules encoding immunoconjugates may be provided. The nucleic acids may be present in host cells. A method of producing the immunoconjugate may comprise by expressing the nucleic acid in cultured host cells, optionally followed by purifying the immunoconjugate from the host cell culture. The IL2 and $\text{TNF}\alpha$ immunoconjugates are
25 preferably produced in separate cell cultures. They may then be individually formulated as medicaments for administration as described.

Detailed Description

Certain aspects of the invention are as set out in the appended claims, which may be combined with any other part of the present disclosure.

30 An antibody molecule is an immunoglobulin whether natural or partly or wholly synthetically produced. The term also covers any polypeptide or protein comprising an antibody antigen-binding site. Thus, this term covers antibody fragments and derivatives, including any polypeptide comprising an antibody antigen-binding site, whether natural or wholly or partially synthetic. Chimeric molecules comprising an antibody antigen-binding site, or equivalent, fused
35 to another polypeptide are therefore included. Cloning and expression of chimeric antibodies is well known (EP0120694, EP0125023).

Further techniques available in the art of antibody engineering have made it possible to isolate human and humanised antibodies. For example, human hybridomas can be made as previously described [5]. Phage display is another established technique [5, WO92/01047]. Transgenic mice in which the mouse antibody genes are inactivated and functionally replaced with human antibody genes while leaving intact other components of the mouse immune system, can be used for isolating human antibodies [6].

Synthetic antibody molecules may be created by expression from genes generated by means of oligonucleotides synthesised and assembled within suitable expression vectors [7, 8].

It has been shown that fragments of a whole antibody can perform the function of binding antigens. Antibody fragments are preferred in conjugates of the invention owing to their small size and minimised interaction with other molecules and receptors (e.g. Fc receptor). Particularly preferred are single chain Fv molecules (scFv), wherein a VH domain and a VL domain are linked by a peptide linker which allows the two domains to associate to form an antigen binding site [9, 10]. scFv may be stabilised by the incorporation of disulphide bridges linking the VH and VL domains [11].

Another small antigen-binding antibody fragment is a dAb (domain antibody), namely the variable region of an antibody heavy or light chain [12]. VH dAbs occur naturally in camelids (e.g. camel, llama) and may be produced by immunising a camelid with a target antigen, isolating antigen-specific B cells and directly cloning dAb genes from individual B cells. dAbs are also producible in cell culture. Their small size, good solubility and temperature stability makes them particularly physiologically useful and suitable for selection and affinity maturation.

An antigen-binding site is the part of a molecule that specifically binds to and is complementary to all or part of the target antigen. In an antibody molecule it is referred to as the antibody antigen-binding site, and comprises the part of the antibody that specifically binds to and is complementary to all or part of the target antigen. Where an antigen is large, an antibody may only bind to a particular part of the antigen, which part is termed an epitope. An antibody antigen-binding site may be provided by one or more antibody variable domains. Preferably, an antibody antigen-binding site comprises an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH).

The term "specific" may be used to refer to the situation in which one member of a specific binding pair will not show any significant binding to molecules other than its specific binding partner(s). The term is also applicable where e.g. an antigen-binding site is specific for a particular epitope that is carried by a number of antigens, in which case the antibody carrying the antigen-binding site will be able to bind to the various antigens carrying the epitope.

In immunoconjugates of the invention, the antibody molecule binds an extracellular matrix component which is a marker of tumour growth. The extracellular matrix (ECM) is

remodelled during tumour growth, and alternative splice variants of ECM components may be selectively expressed at the site of the lesion.

One example is fibronectin. For example, the B-FN isoform of fibronectin contains an extra domain ED-B. An antibody molecule preferably binds specifically to ED-B of fibronectin isoform B-FN. The antibody molecule may comprise the L19 CDRs. For example, the antibody molecule may be an scFv having a VH domain with an amino acid sequence comprising VH CDR1, VH CDR2 and/or VH CDR3 of L19, and a VL domain with an amino acid sequence comprising VL CDR1, VL CDR2 and/or VL CDR3 of L19. An antibody molecule may comprise a VH domain having an amino acid sequence with at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% sequence identity with the amino acid sequence of the L19 VH domain as set out in SEQ ID NO: 7, and/or comprises a VL domain having an amino acid sequence with at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% sequence identity with the amino acid sequence of the L19 VL domain as set out in SEQ ID NO: 9. Preferably the antibody molecule is an scFv(L19) comprising an L19 VH domain (SEQ ID NO: 7) and an L19 VL domain (SEQ ID NO: 9). In a preferred embodiment, the antibody molecule is scFv(L19) having the amino acid sequence SEQ ID NO: 10 (Figure 3).

Modified forms of the L19 VH and/or VL domain may be employed in immunoconjugates of the invention, for example an antibody molecule may comprise the L19 VH or L19 VL domain in which 1, 2, 3, 4 or 5 amino acid substitutions have been made in a CDR and/or framework region, while retaining specific binding to fibronectin ED-B. Such amino acid substitutions are preferably conservative, e.g. substitution of one hydrophobic residue for another, one polar residue for another, arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine.

Another example is tenascin-C (TnC), which exists in various isoforms generated by alternative splicing. In neoplastic tissues TnC containing additional domains are more widely expressed than in normal tissues, especially isoforms containing domain C (cTN-C) (WO00/63699). Thus, an antibody molecule may bind a splice isoform of tenascin-C, e.g. it may bind domain C.

Nucleic acid molecules encoding the immunoconjugates and parts thereof also form part of the invention. The nucleic acid molecule may be a vector, e.g. a plasmid suitable for expression of the nucleotide sequence. Normally the nucleotide sequence is operably linked to a regulatory element such as a promoter for transcription.

The nucleic acid molecules may be contained in a host cell, which may be a cell co-transfected with the nucleic acid molecules or a daughter of such a cell. Cells, especially eukaryotic cells e.g. HEK and CHO cells, or bacterial cells e.g. *Escherichia coli*, containing the nucleic acid molecules also form part of the invention.

Immunoconjugates of the invention may be produced using recombinant techniques, for example by expressing all or part of the immunoconjugate as a fusion protein. Normally the

expression is performed in a host cell containing nucleic acid, as described above. Expression may therefore comprise culturing such a host cell. For TNF α fusion proteins, trimerisation of the subunits may occur in the cell or during purification of the fusion proteins from the cell.

Preferably the antibody molecule is conjugated with the cytokine by means of a peptide bond, e.g. within a fusion protein comprising the TNF α or IL2 and the antibody molecule or a polypeptide chain thereof. See WO01/66298 and Borsi *et al.* [4] for further information on preparation of immunoconjugates comprising TNF α or IL2. See Carnemolla *et al.* [2], Taniguchi *et al.* [13], Maeda *et al.* [14] or Devos *et al.* [15] for further IL2 sequence information useful in preparation of a fusion polypeptide comprising IL2.

TNF α used in immunoconjugates of the invention is preferably human TNF α . IL2 is preferably human IL-2. Antibody molecules are preferably human or humanised antibody molecules.

Also described is a method comprising formulating the immunoconjugate into a pharmaceutical composition. Generally this involves purifying the immunoconjugate and combining it with a physiologically acceptable carrier.

Compositions according to the present invention, and for use in accordance with the present invention, may comprise, in addition to active ingredient (immunoconjugate), a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. For injection at the tumour site, the immunoconjugate may be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability.

Brief Description of the Drawings

Figure 1A shows tumour volume over time in the experiment described in Example 1.

Figure 1B shows follow up data supplementary to Figure 1A.

Figure 2 shows average weight of the mice over time in the experiment described in Example 1.

Figure 3 shows the amino acid sequence of scFv(L19) (SEQ ID NO: 10). The VH and VL domains are shown separately (SEQ ID NO: 7 and SEQ ID NO: 9, respectively). The CDR1, 2 and 3 sequences in both the VH and VL domain are shown underlined. The VH and VL domains are separated by a 12 residue peptide linker sequence (SEQ ID NO: 8).

Experiments

Example 1 - Intratumoral Combination L19-IL2 and L19-muTNF

Procedure

Mouse strain: female, immunocompetent 10 weeks old 129SVE mice

5 20 Million F9 cells were implanted

Treatment was started at day 7 after tumour implantation

On day 7, mice received a single intratumoural injection of 30 µg L19-IL2, 7 µg L19-muTNF, the combination or PBS. The total volume injected was 90 µl.

No further injections were given.

10 Tumour volume was measured daily.

Results

Tumour volume measured from day 7 to day 12 (Figure 1A) – average results.

Tumours of mice who received only saline increased in size from about day 8 onwards.

15 Tumours of mice who received a single immunocytokine began to increase in size slowly from about day 10.

Tumours of mice who received the combination of immunocytokines decreased in size from day 9. By day 11 and 12, tumour volume was barely measurable. No tumour growth was observed in follow up measurements taken until day 20, when tumour volume was measured at zero.

Weight of the mice was also recorded over time (Figure 2).

20 Visual observations on day 10 after tumour implantation:

Large subcutaneous tumours had formed in the PBS-treated (control) mice (n = 4 mice).

Visible tumours were present in the mice treated with L19-IL2 and L19-TNF, but markedly smaller than the tumours in control mice, and tumours were barely visible or not visible in the mice treated with the combination (n = 5 mice in each group).

25

The results indicate that the single administration of the combination of immunocytokines was successful in achieving complete eradication of the treated tumour. It was observed that tumours did not regrow in the mice.

Further details

30

When tumours reached a size of 70 mm³ mice were randomly grouped and treatment was started. Mice received a single intratumoral injection of L19-IL2 (30 µg), L19-TNF (7 µg) or the combination in a volume of 90 µl PBS. The mice were monitored daily, and tumour volume was measured with a caliper, using the formula volume = length x width² x 0.5.

Figure 1B shows continuation of the data shown in Figure 1A. The mice treated with the combination remained without any measurable tumour in further follow up measurements until day 27. Five out of five mice in this group achieved clinical response.

References

- 1 Weide *et al. Cancer* 1 September 2010:4139-4146
- 2 Carnemolla *et al. Blood* 99:1659-1665 2002
- 3 Christ *et al. Clin Cancer Res* 7(5) :1385-97 2001
- 4 Borsi *et al. Blood* 102:4384-4392 2003
- 5 Kontermann, R & Dubel, S, *Antibody Engineering*, Springer-Verlag New York, LLC; 2001, ISBN: 3540413545
- 6 Mendez, M. et al. *Nature Genet*, 15(2): 146–156 1997
- 7 Knappik et al. *J. Mol. Biol.* (2000) 296, 57-86
- 8 Krebs et al. *Journal of Immunological Methods* 254 2001 67–84
- 9 Bird et al, *Science*, 242, 423-426, 1988
- 10 Huston et al, *PNAS USA*, 85, 5879-5883, 1988
- 11 Reiter, Y. et al, *Nature Biotech*, 14, 1239-1245, 1996
- 12 Holt et al *Trends in Biotechnology* 21, 484-490 2003
- 13 Taniguchi *et al. Nature* 302:305-310 1983
- 14 Maeda *et al. Biochem Biophys Res Comm* 115 :1040-1047 1983
- 15 Devos *et al. Nucl Acids Res* 11 :4307-4323 1983

What is claimed is:

1. A method of treating a tumour in a patient by injecting a single dose of a $\text{TNF}\alpha$ immunoconjugate and a single dose of an IL2 immunoconjugate at the tumour site, wherein
the $\text{TNF}\alpha$ immunoconjugate comprises $\text{TNF}\alpha$ linked to an antibody molecule that binds a splice isoform of an extracellular matrix component, and
the IL2 immunoconjugate comprises IL2 linked to an antibody molecule that binds a splice isoform of an extracellular matrix component.
2. A composition comprising a combination of a $\text{TNF}\alpha$ immunoconjugate and an IL2 immunoconjugate for use in a method of treating a tumour in a patient by injecting a single dose of the $\text{TNF}\alpha$ immunoconjugate and a single dose of the IL2 immunoconjugate at the tumour site, wherein
the $\text{TNF}\alpha$ immunoconjugate comprises $\text{TNF}\alpha$ linked to an antibody molecule that binds a splice isoform of an extracellular matrix component, and
the IL2 immunoconjugate comprises IL2 linked to an antibody molecule that binds a splice isoform of an extracellular matrix component.
3. A composition comprising a $\text{TNF}\alpha$ conjugate for use in a method of treating a tumour in a patient by injecting a single dose of the $\text{TNF}\alpha$ immunoconjugate and a single dose of an IL2 immunoconjugate at the tumour site, wherein
the $\text{TNF}\alpha$ immunoconjugate comprises $\text{TNF}\alpha$ linked to an antibody molecule that binds a splice isoform of an extracellular matrix component, and
the IL2 immunoconjugate comprises IL2 linked to an antibody molecule that binds a splice isoform of an extracellular matrix component.
4. A composition comprising an IL2 conjugate for use in a method of treating a tumour in a patient by injecting a single dose of the IL2 immunoconjugate and a single dose of a $\text{TNF}\alpha$ immunoconjugate at the tumour site, wherein
the $\text{TNF}\alpha$ immunoconjugate comprises $\text{TNF}\alpha$ linked to an antibody molecule that binds a splice isoform of an extracellular matrix component, and
the IL2 immunoconjugate comprises IL2 linked to an antibody molecule that binds a splice isoform of an extracellular matrix component..

5. A method according to claim 1, or a composition for use according to any of claims 2 to 4, wherein the antibody molecule in the TNF α immunoconjugate and the antibody molecule in the IL2 conjugate bind the same extracellular matrix component.
6. A method or composition for use according to any of the preceding claims, wherein the antibody molecule in the TNF α conjugate and the antibody molecule in the IL2 conjugate are identical antibody molecules.
7. A method or composition for use according to claim 5 or claim 6, wherein the antibody molecule binds a splice isoform of fibronectin.
8. A method or composition for use according to claim 7, wherein the antibody molecule binds B-FN.
9. A method or composition for use according to claim 8, wherein the antibody molecule comprises the L19 complementarity determining regions (CDRs),
wherein the L19 CDRs are:
- | | | |
|----------|------------------|---------------|
| VH CDR1 | SFSMS | SEQ ID NO: 1 |
| VH CDR 2 | SISGSSGTTYADSVKG | SEQ ID NO: 2 |
| VH CDR 3 | PFPYFDY | SEQ ID NO: 3 |
| VL CDR 1 | RASQSVSSSFLA | SEQ ID NO: 4 |
| VL CDR 2 | YASSRAT | SEQ ID NO: 5 |
| VL CDR 3 | QQTGRIPPT | SEQ ID NO: 6. |
10. A method or composition for use according to any of the preceding claims, wherein the antibody molecule comprises the L19 VH domain SEQ ID NO: 7 and the L19 VL domain SEQ ID NO: 9.
11. A method or a composition for use according to any of the preceding claims, wherein the antibody molecule is scFv.
12. A method or a composition for use according to claim 11, wherein the antibody molecule is scFv-L19 SEQ ID NO: 10.
13. A method or a composition for use according to any of the preceding claims, wherein one or both immunoconjugates carries a detectable label.

14. A method or a composition for use according to any of the preceding claims, wherein the injection is intratumoural injection.
15. A method or a composition for use according to any of the preceding claims, wherein the method comprises injecting the TNF α immunoconjugate and the IL2 immunoconjugate in separate injections.
16. A method or a composition for use according to any of the preceding claims, wherein the method comprises injecting the TNF α immunoconjugate and the IL2 immunoconjugate on the same day.
17. A method or a composition for use according to any of claims 1 to 14, wherein the method comprises injecting the TNF α immunoconjugate and the IL2 immunoconjugate in a combined injection.
18. A method or a composition for use according to any of the preceding claims, wherein the method comprises injecting the TNF α immunoconjugate and the IL2 immunoconjugate simultaneously.
19. A method or a composition for use according to any of claims 1 to 16, wherein the method comprises injecting the TNF α immunoconjugate and the IL2 immunoconjugate sequentially.
20. A method or a composition for use according to any of the preceding claims, wherein the dose of TNF α immunoconjugate is administered as a single injection.
21. A method or a composition for use according to any of the preceding claims, wherein the dose of IL2 immunoconjugate is administered as a single injection.
22. A method or a composition for use according to any of claims 11 to 21, comprising injecting a single dose of 10 – 100 μ g of an IL2-scFv immunoconjugate at the tumour site.
23. A method or a composition for use according to any of claims 11 to 22, comprising injecting a single dose of 2 – 20 μ g of a TNF α -scFv immunoconjugate at the tumour site.

24. A method or a composition for use according to any of the preceding claims, wherein the method comprises treating multiple tumours in the patient by injecting a single dose of the TNF α immunoconjugate and a single dose of the IL2 immunoconjugate at the site of each tumour.
25. A method or a composition for use according to any of the preceding claims, wherein the tumour is a primary tumour.
26. A method or a composition for use according to any of the preceding claims, wherein the tumour is a skin tumour.
27. A method or a composition for use according to any of the preceding claims, wherein the tumour is a carcinoma.
28. A method or a composition for use according to any of the preceding claims, wherein the patient is one who has not received previous treatment with IL-2 or TNF α .
29. A method or a composition for use according to any of the preceding claims, wherein treating the tumour comprises eradicating the tumour.




Figure 1A




Figure 1B




Figure 2

VH

CDR1
EVQLLESGGGLVQPGGSLRLSCAASGFTFSFMSWVRQAPGKGLEWVSSISGSSGTTYYADSVKGRFT

CDR2

CDR3
ISRDN SKNTLYLQMN SLRAEDTAVYYCAKFPFPYFDYWGQGLTVTVSS

LINKER
GDGSSGGGGAS

VL

CDR1
EIVLTQSPGTLSPGERATLSCRASQSVSSSF~~LA~~WYQQKPGQAPRLLIYYASSRAIGIPDRFSGS

CDR2

CDR3
GSGTDFTLTISRLEPEDFAVYYCQQIGRIPPTFGGQGTKVEIK

Figure 3

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/062705

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K14/54 C07K14/525 C07K16/18 A61K38/20 A61K47/48
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, EMBASE, BIOSIS, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01/62298 A2 (PHILOGEN S R L [IT]; ZARDI LUCIANO [IT]; NERI DARIO [CH]; CARNEMOLLA B) 30 August 2001 (2001-08-30) cited in the application the whole document	1-29
X	L. BORSI: "Selective targeted delivery of TNF to tumor blood vessels", BLOOD, vol. 102, no. 13, 15 December 2003 (2003-12-15), pages 4384-4392, XP55037714, ISSN: 0006-4971, DOI: 10.1182/blood-2003-04-1039 cited in the application the whole document	1-29
	----- -/--	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

12 September 2012

Date of mailing of the international search report

04/10/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Sirim, Pinar

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/062705

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ENRICA BALZA ET AL: "Therapy-induced antitumor vaccination in neuroblastomas by the combined targeting of IL-2 and TNF[alpha]", INTERNATIONAL JOURNAL OF CANCER, vol. 127, no. 1, 1 July 2010 (2010-07-01), pages 101-110, XP55037809, ISSN: 0020-7136, DOI: 10.1002/ijc.25018 the whole document	1-29
A	----- CHRIST ET AL: "Interleukin 2-antibody and tumor necrosis factor-antibody fusion proteins induce different antitumor immune responses in vivo.", CLINICAL CANCER RESEARCH, vol. 7, no. 5, 1 May 2001 (2001-05-01), pages 1385-1397, XP55037716, ISSN: 1078-0432 cited in the application the whole document	1-29
A	----- CARNEMOLLA BARBARA ET AL: "Enhancement of the antitumor properties of interleukin-2 by its targeted delivery to the tumor blood vessel extracellular matrix", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 99, no. 5, 1 March 2002 (2002-03-01), pages 1659-1665, XP002256864, ISSN: 0006-4971, DOI: 10.1182/BLOOD.V99.5.1659 cited in the application the whole document	1-29
A	----- HALIN C ET AL: "Synergistic therapeutic effects of a tumor targeting antibody fragment, fused to interleukin 12 and to tumor necrosis factor alpha", CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, US, vol. 63, no. 12, 15 June 2003 (2003-06-15), pages 3202-3210, XP002342601, ISSN: 0008-5472 the whole document -----	1-29

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/062705

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0162298	A2	30-08-2001	
		AT 333900 T	15-08-2006
		AT 526039 T	15-10-2011
		AU 3947001 A	03-09-2001
		CA 2399866 A1	30-08-2001
		DE 60121733 T2	09-08-2007
		DK 1257297 T3	20-11-2006
		DK 1719528 T3	09-01-2012
		EP 1257297 A2	20-11-2002
		EP 1719528 A2	08-11-2006
		ES 2269361 T3	01-04-2007
		ES 2373966 T3	10-02-2012
		JP 4917232 B2	18-04-2012
		JP 2003523406 A	05-08-2003
		JP 2012036216 A	23-02-2012
		PT 1257297 E	29-12-2006
		PT 1719528 E	06-01-2012
		US 2004013640 A1	22-01-2004
		US 2010316602 A1	16-12-2010
		WO 0162298 A2	30-08-2001
