



US 20030188330A1

(19)

United States

(12)

Patent Application Publication

(10)

Pub. No.: US 2003/0188330 A1

Heard

(43)

Pub. Date:

Oct. 2, 2003

(54) **GENES FOR MODIFYING PLANT TRAITS
XI**

(76) Inventor: **Jacqueline Heard**, San Mateo, CA
(US)

Correspondence Address:
MATTHEW KASER
Mendel Biotechnology, Inc.
21375 Cabot Blvd.
Hayward, CA 94545 (US)

(21) Appl. No.: **10/112,887**
(22) Filed: **Mar. 18, 2002**

Publication Classification

(51) **Int. Cl.⁷** **A01H 1/00**; C12N 15/82;
C07H 21/04; C07K 14/415
(52) **U.S. Cl.** **800/278**; 536/23.6; 530/370

(57) **ABSTRACT**

The invention relates to plant transcription factor polypeptides, polynucleotides that encode them, homologs and orthologs from a variety of plant species, and methods of using the polynucleotides and polypeptides to produce transgenic plants having advantageous properties compared to a reference plant. Sequence information related to these polynucleotides and polypeptides can also be used in bioinformatic search methods and is also disclosed.

GENES FOR MODIFYING PLANT TRAITS XI

FIELD OF THE INVENTION AND INTRODUCTION

[0001] This invention relates to the field of plant biology. More particularly, the present invention pertains to compositions and methods for phenotypically modifying a plant.

[0002] A plant's traits, such as its biochemical, developmental, or phenotypic characteristics, can be controlled through a number of cellular processes. One important way to manipulate that control is through transcription factors—proteins that influence the expression of a particular gene or sets of genes. Transgenic plants that comprise cells having altered levels of at least one selected transcription factor, for example, possess advantageous or desirable traits. Strategies for manipulating traits by altering a plant cell's transcription factor content can therefore result in plants and crops with commercially valuable properties. Applicants have identified polynucleotides encoding transcription factors, developed numerous transgenic plants using these polynucleotides, and have analyzed the plants for a variety of important traits. In so doing, applicants have identified important polynucleotide and polypeptide sequences for producing commercially valuable plants and crops as well as the methods for making them and using them. Other aspects and embodiments of the invention are described below and can be derived from the teachings of this disclosure as a whole.

BACKGROUND OF THE INVENTION

[0003] Transcription factors can modulate gene expression, either increasing or decreasing (inducing or repressing) the rate of transcription. This modulation results in differential levels of gene expression at various developmental stages, in different tissues and cell types, and in response to different exogenous (e.g., environmental) and endogenous stimuli throughout the life cycle of the organism.

[0004] Because transcription factors are key controlling elements of biological pathways, altering the expression levels of one or more transcription factors can change entire biological pathways in an organism. For example, manipulation of the levels of selected transcription factors may result in increased expression of economically useful proteins or metabolic chemicals in plants or to improve other agriculturally relevant characteristics. Conversely, blocked or reduced expression of a transcription factor may reduce biosynthesis of unwanted compounds or remove an undesirable trait. Therefore, manipulating transcription factor levels in a plant offers tremendous potential in agricultural biotechnology for modifying a plant's traits.

[0005] The present invention provides novel transcription factors useful for modifying a plant's phenotype in desirable ways.

SUMMARY OF THE INVENTION

[0006] In a first aspect, the invention relates to a recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a polypeptide comprising a polypeptide sequence selected from those of the Sequence Listing, SEQ ID NOs:2 and 4, or a complementary nucleotide sequence

thereof; (b) a nucleotide sequence encoding a polypeptide comprising a variant of a polypeptide of (a) having one or more, or between 1 and about 5, or between 1 and about 10, or between 1 and about 30, conservative amino acid substitutions; (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID NOs:1 and 3, or a complementary nucleotide sequence thereof; (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c); (e) a nucleotide sequence which hybridizes under stringent conditions over substantially the entire length of a nucleotide sequence of one or more of: (a), (b), (c), or (d); (f) a nucleotide sequence comprising at least 10 or 15, or at least about 20, or at least about 30 consecutive nucleotides of a sequence of any of (a)-(e), or at least 10 or 15, or at least about 20, or at least about 30 consecutive nucleotides outside of a region encoding a conserved domain of any of (a)-(e); (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide having a biological activity that modifies a plant's characteristic, functions as a transcription factor, or alters the level of transcription of a gene or transgene in a cell; (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g); (i) a nucleotide sequence having at least 60%, or at least 70%, or at least 80%, or at least 90 or at least 95% sequence identity to a nucleotide sequence of any of (a)-(g) or a 10 or 15 nucleotide, or at least about 20, or at least about 30 nucleotide region of a sequence of (a)-(g) that is outside of a region encoding a conserved domain; (j) a nucleotide sequence that encodes a polypeptide having at least 31% sequence identity to a polypeptide listed in the Sequence Listing or in Tables 6 and 7; (k) a nucleotide sequence which encodes a polypeptide having at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 95% sequence identity to a polypeptide listed in the Sequence Listing or in Tables 6 and 7; and (l) a nucleotide sequence that encodes a conserved domain of a polypeptide having at least 85%, or at least 90%, or at least 95%, or at least 98% sequence identity to a conserved domain of a polypeptide listed in the Sequence Listing or in Tables 6 and 7. The recombinant polynucleotide may further comprise a constitutive, inducible, or tissue-specific promoter operably linked to the nucleotide sequence. The invention also relates to compositions comprising at least two of the above-described polynucleotides.

[0007] In a second aspect, the invention comprises an isolated or recombinant polypeptide comprising a subsequence of at least about 10, or at least about 15, or at least about 20, or at least about 30 contiguous amino acids encoded by the recombinant or isolated polynucleotide described above, or comprising a subsequence of at least about 8, or at least about 12, or at least about 15, or at least about 20, or at least about 30 contiguous amino acids outside a conserved domain.

[0008] In a third aspect, the invention comprises an isolated or recombinant polynucleotide which encodes a polypeptide which is a paralog of the isolated polypeptide described in paragraph 6 above. In one aspect, the invention is an paralog which, when expressed in Arabidopsis, modifies a trait of the Arabidopsis plant.

[0009] In a fourth aspect, the invention comprises an isolated or recombinant polynucleotide which encodes a polypeptide which is an ortholog of the isolated polypeptide

described in paragraph 6 above. In one aspect, the invention is an ortholog which, when expressed in Arabidopsis, modifies a trait of the Arabidopsis plant.

[0010] In a fifth aspect, the invention comprises an isolated or recombinant polynucleotide which encodes a polypeptide which is a paralog of the isolated polypeptide described in paragraph 6 above. In one aspect, the invention is an paralog which, when expressed in Arabidopsis, modifies a trait of the Arabidopsis plant.

[0011] In another aspect, the invention comprises an isolated polypeptide which is an ortholog of the isolated polypeptide described in paragraph 6 above. In one aspect, the invention is an ortholog which, when expressed in Arabidopsis, modifies a trait of the Arabidopsis plant.

[0012] In yet another aspect, the invention comprises an isolated synthetic polypeptide which is a homolog of the isolated polypeptide described in paragraph 6 above. In one aspect, the invention is a synthetic polypeptide which, when expressed in Arabidopsis, modifies a trait of the Arabidopsis plant.

[0013] In another aspect, the invention is a transgenic plant comprising one or more of the above-described recombinant polynucleotides. In yet another aspect, the invention is a plant with altered expression levels of a polynucleotide described above or a plant with altered expression or activity levels of an above-described polypeptide. Further, the invention is a plant lacking a nucleotide sequence encoding a polypeptide described above or substantially lacking a polypeptide described above. The plant may be any plant, including, but not limited to, Arabidopsis, mustard, soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, pumpkin, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, vegetable brassicas, and mint or other labiates. In yet another aspect, the inventions is an isolated plant material of a plant, including, but not limited to, plant tissue, fruit, seed, plant cell, embryo, protoplast, pollen, and the like. In yet another aspect, the invention is a transgenic plant tissue culture of regenerable cells, including, but not limited to, embryos, meristematic cells, microspores, protoplast, pollen, and the like.

[0014] In a further aspect the invention provides a method of using the polynucleotide composition to breed progeny from a parent plant including crossing plants, producing seeds from transgenic plants, and methods of breeding using transgenic plants.

[0015] In a further aspect, the invention provides a progeny plant derived from a parental plant wherein said progeny plant exhibits at least three fold greater messenger RNA levels than said parental plant, wherein the messenger RNA encodes a DNA-binding protein which is capable of binding to a DNA regulatory sequence and inducing expression of a plant trait gene, wherein the progeny plant is characterized by a change in the plant trait compared to said parental plant. In yet a further aspect, the progeny plant exhibits at least ten fold greater messenger RNA levels compared to said parental plant. In yet a further aspect, the progeny plant exhibits at least fifty fold greater messenger RNA levels compared to said parental plant.

[0016] In a further aspect, the invention relates to a cloning or expression vector comprising the isolated or recombinant polynucleotide described above or cells comprising the cloning or expression vector.

[0017] In yet a further aspect, the invention relates to a composition produced by incubating a polynucleotide of the invention with a nuclease, a restriction enzyme, a polymerase; a polymerase and a primer; a cloning vector, or with a cell.

[0018] Furthermore, the invention relates to a method for producing a plant having a modified trait. The method comprises altering the expression of an isolated or recombinant polynucleotide of the invention or altering the expression or activity of a polypeptide of the invention in a plant to produce a modified plant, and selecting the modified plant for a modified trait. In one aspect, the plant is a monocot plant. In another aspect, the plant is a dicot plant. In another aspect the recombinant polynucleotide is from a dicot plant and the plant is a monocot plant. In yet another aspect the recombinant polynucleotide is from a monocot plant and the plant is a dicot plant. In yet another aspect the recombinant polynucleotide is from a monocot plant and the plant is a monocot plant. In yet another aspect the recombinant polynucleotide is from a dicot plant and the plant is a dicot plant.

[0019] In another aspect, the invention relates to a method of identifying a factor that is modulated by or interacts with a polypeptide encoded by a polynucleotide of the invention. The method comprises expressing a polypeptide encoded by the polynucleotide in a plant; and identifying at least one factor that is modulated by or interacts with the polypeptide. In one embodiment the method for identifying modulating or interacting factors is by detecting binding by the polypeptide to a promoter sequence, or by detecting interactions between an additional protein and the polypeptide in a yeast two hybrid system, or by detecting expression of a factor by hybridization to a microarray, subtractive hybridization, or differential display.

[0020] In yet another aspect, the invention is a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest. The method comprises placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of the invention and monitoring one or more of the expression level of the polynucleotide in the plant, the expression level of the polypeptide in the plant, and modulation of an activity of the polypeptide in the plant. In a farther aspect, the invention is a method of using a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest. The polynucleotide may be selected from the group comprising SEQ ID NOs:1 and 3, a variant or ortholog thereof. In the alternative, the polypeptide may be selected from the group comprising SEQ ID NOs:2 and 4, a variant or ortholog thereof. The method comprises placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of the invention and monitoring one or more of the expression level of the polynucleotide in the plant, the expression level of the polypeptide in the plant, and modulation of an activity of the polypeptide in the plant.

[0021] In yet another aspect, the invention relates to an integrated system, computer or computer readable medium comprising one or more character strings corresponding to a

polynucleotide of the invention, or to a polypeptide encoded by the polynucleotide. The integrated system, computer or computer readable medium may comprise a link between one or more sequence strings to a modified plant trait.

[0022] In yet another aspect, the invention is a method for identifying a sequence similar or homologous to one or more polynucleotides of the invention, or one or more polypeptides encoded by the polynucleotides. The method comprises providing a sequence database, and querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.

[0023] The method may further comprise of linking the one or more of the polynucleotides of the invention, or encoded polypeptides, to a modified plant phenotype.

[0024] Brief Description of the Sequence Listing and Tables

[0025] The Sequence Listing provides exemplary polynucleotide and polypeptide sequences of the invention. The traits associated with the use of the sequences are included in the "Examples" section of the invention disclosure.

[0026] Tables 1 through 5 are shown and described in the invention disclosure.

[0027] Table 6 lists a summary of orthologous and homologous sequences of the polynucleotide sequences and polypeptide sequences of the invention (SEQ ID NOS:1-4) identified using BLAST (TBLASTX program). The first column shows the polynucleotide sequence identifier (SEQ ID NO), the second column shows the transcription factor cDNA identifier (Gene ID), the third column shows the orthologous or homologous polynucleotide GenBank Accession Number (Test Sequence ID), the fourth column shows the orthologous or homologous polynucleotide sequence identifier (Test Sequence SEQ ID NO), the fifth column shows the calculated probability value that the sequence identity is due to chance (Smallest Sum Probability), and the sixth column shows the orthologous or homologous GenBank annotation (Test Sequence GenBank Annotation).

[0028] Table 7 lists orthologous and homologous sequences of the polynucleotide sequences and polypeptide sequences of the invention (SEQ ID NOS:1-4) identified using BLAST (TBLASTX program). The first column shows the polynucleotide sequence identifier (SEQ ID NO), the second column shows the transcription factor cDNA identifier (Gene ID), the third column shows the orthologous or homologous polynucleotide GenBank Accession Number (Test Sequence ID), the fourth column shows the orthologous or homologous polynucleotide sequence GenBank annotation (Test Sequence GenBank Annotation) identifier (Test Sequence SEQ ID NO), the fifth column shows the reading frame of the Test sequence which encodes the orthologous or homologous sequence (Reading Frame), the sixth column shows the calculated score value of the aligned sequences (High Score), the seventh column shows the calculated probability value that the sequence identity is due to chance (Smallest Sum Probability), and the eighth column shows the number of regions of the orthologous or homo-

gous Test Sequences which aligned with the sequence encoded by the transcription factor cDNA sequence GenBank annotation (N).

DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

[0029] In an important aspect, the present invention relates to polynucleotides and polypeptides, e.g. for modifying phenotypes of plants. Throughout this disclosure, various information sources are referred to and/or are specifically incorporated. The information sources include scientific journal articles, patent documents, textbooks, and world wide web browser-inactive page addresses, for example. While the reference to these information sources clearly indicates that they can be used by one of skill in the art, applicants specifically incorporate each and every one of the information sources cited herein, in their entirety, whether or not a specific mention of "incorporation by reference" is noted. The contents and teachings of each and every one of the information sources can be relied on and used to make and use embodiments of the invention.

[0030] It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a plant" includes a plurality of such plants, and a reference to "a stress" is a reference to one or more stresses and equivalents thereof known to those skilled in the art, and so forth.

[0031] The polynucleotides of the invention encode plant transcription factors. As one of ordinary skill in the art recognizes, transcription factors can be identified by the presence of a region or domain of structural similarity or identity to a specific consensus sequence or the presence of a specific consensus DNA-binding site or DNA-binding site motif (see, for example, Riechmann et al., *Science* 290: 2105-2110 (2000)). The plant transcription factors may belong to one of the following transcription factor families: the AP2 (APETALA2) domain transcription factor family (Riechmann and Meyerowitz (1998) *Biol. Chem.* 379:633-646); the MYB transcription factor family (Martin and Paz-Ares, (1997) *Trends Genet.* 13:67-73); the MADS domain transcription factor family (Riechmann and Meyerowitz (1997) *Biol. Chem.* 378:1079-1101); the WRKY protein family (Ishiguro and Nakamura (1994) *Mol. Gen. Genet.* 244:563-571); the ankyrin-repeat protein family (Zhang et al. (1992) *Plant Cell* 4:1575-1588); the zinc finger protein (Z) family (Klug and Schwabe (1995) *FASEB J.* 9: 597-604); the homeobox (HB) protein family (Duboule (1994) *Guidebook to the Homeobox Genes*, Oxford University Press); the CAAT-element binding proteins (Forsburg and Guarente (1989) *Genes Dev.* 3:1166-1178); the squamosa promoter binding proteins (SPB) (Klein et al. (1996) *Mol. Gen. Genet.* 1996 250:7-16); the NAM protein family (Souer et al. (1996) *Cell* 85:159-170); the IAA/AUX proteins (Rouse et al. (1998) *Science* 279:1371-1373); the HLH/MYC protein family (Littlewood et al. (1994) *Prot. Profile* 1:639-709); the DNA-binding protein (DBP) family (Tucker et al. (1994) *EMBO J.* 13:2994-3002); the bZIP family of transcription factors (Foster et al. (1994) *FASEB J.* 8:192-200); the Box P-binding protein (the BPF-1) family (da Costa e Silva et al. (1993) *Plant J.* 4:125-135); the high mobility group (HMG) family (Bustin and Reeves (1996) *Prog. Nucl. Acids Res. Mol. Biol.* 54:35-100); the scarecrow

(SCR) family (Di Laurenzio et al. (1996) *Cell* 86:423-433); the GF14 family (Wu et al. (1997) *Plant Physiol.* 114:1421-1431); the polycomb (PCOMB) family (Kennison (1995) *Annu. Rev. Genet.* 29:289-303); the teosinte branched (TEO) family (Luo et al. (1996) *Nature* 383:794-799); the ABI3 family (Giraudat et al. (1992) *Plant Cell* 4:1251-1261); the triple helix (TH) family (Dehesh et al. (1990) *Science* 250:1397-1399); the EIL family (Chao et al. (1997) *Cell* 89:1133-44); the AT-HOOK family (Reeves and Nissen (1990) *Journal of Biological Chemistry* 265:8573-8582); the SIFA family (Zhou et al. (1995) *Nucleic Acids Res.* 23:1165-1169); the bZIP2 family (Lu and Ferl (1995) *Plant Physiol.* 109:723); the YABBY family (Bowman et al. (1999) *Development* 126:2387-96); the PAZ family (Bohmert et al. (1998) *EMBO J.* 17:170-80); miscellaneous (MISC) transcription factors including the DPBF family (Kim et al. (1997) *Plant J.* 11: 1237-1251) and the SPF1 family (Ishiguro and Nakamura (1994) *Mol. Gen. Genet.* 244:563-571); the golden (GLD) family (Hall et al. (1998) *Plant Cell* 10:925-936); the TUBBY family (Boggin et al. (1999) *Science* 286:2119-2125); the heat shock family (Wu C (1995) *Annu Rev Cell Dev Biol* 11:441-469); the ENBP family (Christiansen et al (1996) *Plant Mol Biol* 32:809-821); the RING-zinc family (Jensen et al. (1998) *FEBS letters* 436:283-287); the PDBP family (Janik et al *Virology.* (1989)168:320-329); the PCF family (Cubas P, et al. *Plant J.* (1999)18:215-22); the SRS (SHI-related) family (Fridborg et al *Plant Cell* (1999) 11:1019-1032); the CPP (cysteine-rich polycomb-like) family (Cvitanich et al *Proc. Natl. Acad. Sci. U S A.* (2000) 97:8163-8168); the ARF (auxin response factor) family (Ulmasov, et al. (1999) *Proc. Natl. Acad. Sci. USA* 96: 5844-5849); the SWI/SNF family (Collingwood et al *J. Mol. End.* 23:255-275); the ACBF family (Seguin et al *Plant Mol Biol.* (1997) 35:281-291); PCGL (CG-1 like) family (*Plant Mol Biol.* (1994) 25:921-924); the ARID family (Vazquez et al *Development.* (1999) 126: 733-42); the Jumoni family (Balciunas et al *Trends Biochem Sci.* (2000) 25: 274-276); the bZIP-NIN family (Schauser et al *Nature.* (1999) 402: 191-195); the E2F family Kaelin et al (1992) *Cell* 70: 351-364); and the GRF-like family (Knaap et al (2000) *Plant Physiol.* 122: 695-704. As indicated by any part of the list above and as known in the art, transcription factors have been sometimes categorized by class, family, and sub-family according to their structural content and consensus DNA-binding site motif, for example. Many of the classes and many of the families and sub-families are listed here. However, the inclusion of one sub-family and not another, or the inclusion of one family and not another, does not mean that the invention does not encompass polynucleotides or polypeptides of a certain family or sub-family. The list provided here is merely an example of the types of transcription factors and the knowledge available concerning the consensus sequences and consensus DNA-binding site motifs that help define them as known to those of skill in the art (each of the references noted above are specifically incorporated herein by reference). A transcription factor may include, but is not limited to, any polypeptide that can activate or repress transcription of a single gene or a number of genes. This polypeptide group includes, but is not limited to, DNA-binding proteins, DNA-binding protein binding proteins, protein kinases, protein phosphatases, GTP-binding proteins, and receptors, and the like.

[0032] In addition to methods for modifying a plant phenotype by employing one or more polynucleotides and

polypeptides of the invention described herein, the polynucleotides and polypeptides of the invention have a variety of additional uses. These uses include their use in the recombinant production (i.e., expression) of proteins; as regulators of plant gene expression, as diagnostic probes for the presence of complementary or partially complementary nucleic acids (including for detection of natural coding nucleic acids); as substrates for further reactions, e.g., mutation reactions, PCR reactions, or the like; as substrates for cloning e.g., including digestion or ligation reactions; and for identifying exogenous or endogenous modulators of the transcription factors.

[0033] A "polynucleotide" is a nucleic acid sequence comprising a plurality of polymerized nucleotides, e.g., at least about 15 consecutive polymerized nucleotide, optionally at least about 30 consecutive nucleotides, at least about 50 consecutive nucleotides. In many instances, a polynucleotide comprises a nucleotide sequence encoding a polypeptide (or protein) or a domain or fragment thereof. Additionally, the polynucleotide may comprise a promoter, an intron, an enhancer region, a polyadenylation site, a translation initiation site, 5' or 3' untranslated regions, a reporter gene, a selectable marker, or the like. The polynucleotide can be single stranded or double stranded DNA or RNA. The polynucleotide optionally comprises modified bases or a modified backbone. The polynucleotide can be, e.g., genomic DNA or RNA, a transcript (such as an mRNA), a cDNA, a PCR product, a cloned DNA, a synthetic DNA or RNA, or the like. The polynucleotide can comprise a sequence in either sense or antisense orientations.

[0034] A "recombinant polynucleotide" is a polynucleotide that is not in its native state, e.g., the polynucleotide comprises a nucleotide sequence not found in nature, or the polynucleotide is in a context other than that in which it is naturally found, e.g., separated from nucleotide sequences with which it typically is in proximity in nature, or adjacent (or contiguous with) nucleotide sequences with which it typically is not in proximity. For example, the sequence at issue can be cloned into a vector, or otherwise recombined with one or more additional nucleic acid.

[0035] A "synthetic polynucleotide" is a polynucleotide not found in nature and encodes a polypeptide not found in nature. The encoded polypeptide comprises at least four consecutive amino acid residues of a polypeptide found in nature.

[0036] An "isolated polynucleotide" is a polynucleotide whether naturally occurring or recombinant, that is present outside the cell in which it is typically found in nature, whether purified or not. Optionally, an isolated polynucleotide is subject to one or more enrichment or purification procedures, e.g., cell lysis, extraction, centrifugation, precipitation, or the like.

[0037] A "polypeptide" is an amino acid sequence comprising a plurality of consecutive polymerized amino acid residues e.g., at least about 15 consecutive polymerized amino acid residues, optionally at least about 30 consecutive polymerized amino acid residues, at least about 50 consecutive polymerized amino acid residues. In many instances, a polypeptide comprises a polymerized amino acid residue sequence which is a transcription factor or a domain or portion or fragment thereof. Additionally, the polypeptide may comprise a localization domain, 2) an activation

domain, 3) a repression domain, 4) an oligomerization domain or 5) a DNA-binding domain, or the like. The polypeptide optionally comprises modified amino acid residues, naturally occurring amino acid residues not encoded by a codon, non-naturally occurring amino acid residues.

[0038] A “recombinant polypeptide” is a polypeptide produced by translation of a recombinant polynucleotide. A “synthetic polypeptide” is a polypeptide created by consecutive polymerization of isolated amino acid residues using methods well known in the art. An “isolated polypeptide,” whether a naturally occurring or a recombinant polypeptide, is more enriched in (or out of) a cell than the polypeptide in its natural state in a wild type cell, e.g., more than about 5% enriched, more than about 10% enriched, or more than about 20%, or more than about 50%, or more, enriched, i.e., alternatively denoted: 105%, 110%, 120%, 150% or more, enriched relative to wild type standardized at 100%. Such an enrichment is not the result of a natural response of a wild type plant. Alternatively, or additionally, the isolated polypeptide is separated from other cellular components with which it is typically associated, e.g., by any of the various protein purification methods herein.

[0039] A “synthetic polypeptide” is a polypeptide not found in nature and has activity of a polypeptide found in nature. The polypeptide comprises at least four consecutive amino acid residues of a polypeptide found in nature.

[0040] “Altered” nucleic acid sequences encoding polypeptide include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide encoding a polypeptide with at least one functional characteristic of the polypeptide. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding polypeptide, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding polypeptide. The encoded polypeptide protein may also be “altered”, and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent polypeptide. Deliberate amino acid substitutions may be made on the basis of similarity in residue side chain chemistry, including, but not limited to, polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological activity of polypeptide is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine. Alignments between different polypeptide sequences may be used to calculate “percentage sequence similarity”.

[0041] The term “plant” includes whole plants, shoot vegetative organs/structures (e.g., leaves, stems and tubers), roots, flowers and floral organs/structures (e.g., bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including embryo, endosperm, and seed coat) and fruit (the mature ovary), plant tissue (e.g., vascular tissue, ground tissue, and the like) and cells (e.g., guard cells, egg cells, and

the like), and progeny of same. The class of plants that can be used in the method of the invention is generally as broad as the class of higher and lower plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), gymnosperms, ferns, horsetails, psilophytes, lycophytes, bryophytes, and multicellular algae. (See for example, Daly et al. 2001 *Plant Physiology* 127:1328-1333; and also Tudge, C., *The Variety of Life*, Oxford University Press, New York, 2000, pp. 547-606.)

[0042] A “transgenic plant” refers to a plant that contains genetic material not found in a wild type plant of the same species, variety or cultivar. The genetic material may include a transgene, an insertional mutagenesis event (such as by transposon or T-DNA insertional mutagenesis), an activation tagging sequence, a mutated sequence, a homologous recombination event or a sequence modified by chimera-plasty. Typically, the foreign genetic material has been introduced into the plant by human manipulation, but any method can be used as one of skill in the art recognizes.

[0043] A transgenic plant may contain an expression vector or cassette. The expression cassette typically comprises a polypeptide-encoding sequence operably linked (i.e., under regulatory control of) to appropriate inducible or constitutive regulatory sequences that allow for the expression of polypeptide. The expression cassette can be introduced into a plant by transformation or by breeding after transformation of a parent plant. A plant refers to a whole plant as well as to a plant part, such as seed, fruit, leaf, or root, plant tissue, plant cells or any other plant material, e.g., a plant explant, as well as to progeny thereof, and to in vitro systems that mimic biochemical or cellular components or processes in a cell.

[0044] “Ectopic expression or altered expression” in reference to a polynucleotide indicates that the pattern of expression in, e.g., a transgenic plant or plant tissue, is different from the expression pattern in a wild type plant or a reference plant of the same species. The pattern of expression may also be compared with a reference expression pattern in a wild type plant of the same species. For example, the polynucleotide or polypeptide is expressed in a cell or tissue type other than a cell or tissue type in which the sequence is expressed in the wild type plant, or by expression at a time other than at the time the sequence is expressed in the wild type plant, or by a response to different inducible agents, such as hormones or environmental signals, or at different expression levels (either higher or lower) compared with those found in a wild type plant. The term also refers to altered expression patterns that are produced by lowering the levels of expression to below the detection level or completely abolishing expression. The resulting expression pattern can be transient or stable, constitutive or inducible. In reference to a polypeptide, the term “ectopic expression or altered expression” further may relate to altered activity levels resulting from the interactions of the polypeptides with exogenous or endogenous modulators or from interactions with factors or as a result of the chemical modification of the polypeptides.

[0045] A “fragment” or “domain,” with respect to a polypeptide, refers to a subsequence of the polypeptide. In some cases, the fragment or domain, is a subsequence of the polypeptide which performs at least one biological function

of the intact polypeptide in substantially the same manner, or to a similar extent, as does the intact polypeptide. For example, a polypeptide fragment can comprise a recognizable structural motif or functional domain such as a DNA-binding site or domain that binds to a DNA promoter region, an activation domain, or a domain for protein-protein interactions. Fragments can vary in size from as few as 6 amino acids to the full length of the intact polypeptide, but are preferably at least about 30 amino acids in length and more preferably at least about 60 amino acids in length. In reference to a polynucleotide sequence, "a fragment" refers to any subsequence of a polynucleotide, typically, of at least about 15 consecutive nucleotides, preferably at least about 30 nucleotides, more preferably at least about 50 nucleotides, of any of the sequences provided herein. [00451] A "conserved domain", with respect to a polypeptide, refers to a domain within a transcription factor family which exhibits a higher degree of sequence homology, such as at least 65% sequence identity including conservative substitutions, and preferably at least 80% sequence identity, and more preferably at least 85%, or at least about 86%, or at least about 87%, or at least about 88%, or at least about 90%, or at least about 95%, or at least about 98% amino acid residue sequence identity of a polypeptide of consecutive amino acid residues. A fragment or domain can be referred to as outside a consensus sequence or outside a consensus DNA-binding site that is known to exist or that exists for a particular transcription factor class, family, or sub-family. In this case, the fragment or domain will not include the exact amino acids of a consensus sequence or consensus DNA-binding site of a transcription factor class, family or sub-family, or the exact amino acids of a particular transcription factor consensus sequence or consensus DNA-binding site. Furthermore, a particular fragment, region, or domain of a polypeptide, or a polynucleotide encoding a polypeptide, can be "outside a conserved domain" if all the amino acids of the fragment, region, or domain fall outside of a defined conserved domain(s) for a polypeptide or protein. The conserved domains for each of polypeptides of SEQ ID NOs:2 and 4 are listed in Table 4 as described in Example VII. A comparison of the regions of the polypeptides in SEQ ID NOs:2 and 4 allows one of skill in the art to identify conserved domain(s) for any of the polypeptides listed or referred to in this disclosure, including those in Tables 6 and 7.

[0046] A "trait" refers to a physiological, morphological, biochemical, or physical characteristic of a plant or particular plant material or cell. In some instances, this characteristic is visible to the human eye, such as seed or plant size, or can be measured by biochemical techniques, such as detecting the protein, starch, or oil content of seed or leaves, or by the observation of the expression level of a gene or genes, e.g., by employing Northern analysis, RT-PCR, microarray gene expression assays, or reporter gene expression systems, or by agricultural observations such as stress tolerance, yield, or pathogen tolerance. Any technique can be used to measure the amount of, comparative level of, or difference in any selected chemical compound or macromolecule in the transgenic plants, however.

[0047] "Trait modification" refers to a detectable difference in a characteristic in a plant ectopically expressing a polynucleotide or polypeptide of the present invention relative to a plant not doing so, such as a wild type plant. In some cases, the trait modification can be evaluated quanti-

tatively. For example, the trait modification can entail at least about a 2% increase or decrease in an observed trait (difference), at least a 5% difference, at least about a 10% difference, at least about a 20% difference, at least about a 30%, at least about a 50%, at least about a 70%, or at least about a 100%, or an even greater difference compared with a wild type plant. It is known that there can be a natural variation in the modified trait. Therefore, the trait modification observed entails a change of the normal distribution of the trait in the plants compared with the distribution observed in wild type plant.

[0048] Trait modifications of particular interest include those to seed (such as embryo or endosperm), fruit, root, flower, leaf, stem, shoot, seedling or the like, including: enhanced tolerance to environmental conditions including freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved tolerance to pest infestations, including nematodes, mollicutes, parasitic higher plants or the like; decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals; improved growth under poor photoconditions (e.g., low light and/or short day length), or changes in expression levels of genes of interest. Other phenotype that can be modified relate to the production of plant metabolites, such as variations in the production of taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prennylipids (such as chlorophylls and carotenoids), glucosinolates, and terpenoids, enhanced or compositionally altered protein or oil production (especially in seeds), or modified sugar (insoluble or soluble) and/or starch composition. Physical plant characteristics that can be modified include cell development (such as the number of trichomes), fruit and seed size and number, yields of plant parts such as stems, leaves and roots, the stability of the seeds during storage, characteristics of the seed pod (e.g., susceptibility to shattering), root hair length and quantity, internode distances, or the quality of seed coat. Plant growth characteristics that can be modified include growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time, flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, as well as plant architecture characteristics such as apical dominance, branching patterns, number of organs, organ identity, organ shape or size.

[0049] Examples of plant trait modifications and how to measure and determine those plant traits or characteristics are provided in the invention disclosure and the "Examples" section, Table 5. The disclosures are intended to illustrate but not limit which plant trait or characteristic may be modified by the invention.

[0050] Polypeptides and Polynucleotides of the Invention

[0051] The present invention provides, among other things, transcription factors (TFs), and transcription factor homologue polypeptides, and isolated or recombinant polynucleotides encoding the polypeptides, or novel variant polypeptides or polynucleotides encoding novel variants of transcription factors derived from the specific sequences provided here. These polypeptides and polynucleotides may be employed to modify a plant's characteristic.

[0052] Exemplary polynucleotides encoding the polypeptides of the invention were identified in the *Arabidopsis*

thaliana GenBank database using publicly available sequence analysis programs and parameters. Sequences initially identified were then further characterized to identify sequences comprising specified sequence strings corresponding to sequence motifs present in families of known transcription factors. In addition, farther exemplary polynucleotides encoding the polypeptides of the invention were identified in the plant GenBank database using publicly available sequence analysis programs and parameters. Sequences initially identified were then further characterized to identify sequences comprising specified sequence strings corresponding to sequence motifs present in families of known transcription factors. Polynucleotide sequences meeting such criteria were confirmed as transcription factors.

[0053] Additional polynucleotides of the invention were identified by screening *Arabidopsis thaliana* and/or other plant cDNA libraries with probes corresponding to known transcription factors under low stringency hybridization conditions. Additional sequences, including full length coding sequences were subsequently recovered by the rapid amplification of cDNA ends (RACE) procedure, using a commercially available kit according to the manufacturer's instructions. Where necessary, multiple rounds of RACE are performed to isolate 5' and 3' ends. The full length cDNA was then recovered by a routine end-to-end polymerase chain reaction (PCR) using primers specific to the isolated 5' and 3' ends. Exemplary sequences are provided in the Sequence Listing.

[0054] The polynucleotides of the invention can be or were ectopically expressed in overexpressor or knockout plants and the changes in the characteristic(s) or trait(s) of the plants observed. Therefore, the polynucleotides and polypeptides can be employed to improve the characteristics of plants.

[0055] Producing Polypeptides

[0056] The polynucleotides of the invention include sequences that encode transcription factors and transcription factor homologue polypeptides and sequences complementary thereto, as well as unique fragments of coding sequence, or sequence complementary thereto. Such polynucleotides can be, e.g., DNA or RNA, e.g., mRNA, cRNA, synthetic RNA, genomic DNA, cDNA synthetic DNA, oligonucleotides, etc. The polynucleotides are either double-stranded or single-stranded, and include either, or both sense (i.e., coding) sequences and antisense (i.e., non-coding, complementary) sequences. The polynucleotides include the coding sequence of a transcription factor, or transcription factor homologue polypeptide, in isolation, in combination with additional coding sequences (e.g., a purification tag, a localization signal, as a fusion-protein, as a pre-protein, or the like), in combination with non-coding sequences (e.g., introns or inteins, regulatory elements such as promoters, enhancers, terminators, and the like), and/or in a vector or host environment in which the polynucleotide encoding a transcription factor or transcription factor homologue polypeptide is an endogenous or exogenous gene.

[0057] A variety of methods exist for producing the polynucleotides of the invention. Procedures for identifying and isolating DNA clones are well known to those of skill in the art, and are described in, e.g., Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology*

volume 152 Academic Press, Inc., San Diego, Calif. ("Berger"); Sambrook et al., *Molecular Cloning—A Laboratory Manual* (2nd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989 ("Sambrook") and *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2000) ("Ausubel").

[0058] Alternatively, polynucleotides of the invention, can be produced by a variety of in vitro amplification methods adapted to the present invention by appropriate selection of specific or degenerate primers. Examples of protocols sufficient to direct persons of skill through in vitro amplification methods, including the polymerase chain reaction (PCR) the ligase chain reaction (LCR), Qbeta-replicase amplification and other RNA polymerase mediated techniques (e.g., NASBA), e.g., for the production of the homologous nucleic acids of the invention are found in Berger, Sambrook, and Ausubel (all supra), as well as Mullis et al., (1987) *PCR Protocols A Guide to Methods and Applications* (Innis et al. eds) Academic Press Inc. San Diego, Calif. (1990) (Innis). Improved methods for cloning in vitro amplified nucleic acids are described in Wallace et al., U.S. Pat. No. 5,426,039. Improved methods for amplifying large nucleic acids by PCR are summarized in Cheng et al. (1994) *Nature* 369: 684-685 and the references cited therein, in which PCR amplicons of up to 40 kb are generated. One of skill will appreciate that essentially any RNA can be converted into a double stranded DNA suitable for restriction digestion, PCR expansion and sequencing using reverse transcriptase and a polymerase. See, e.g., Ausubel, Sambrook and Berger, all supra.

[0059] Alternatively, polynucleotides and oligonucleotides of the invention can be assembled from fragments produced by solid-phase synthesis methods. Typically, fragments of up to approximately 100 bases are individually synthesized and then enzymatically or chemically ligated to produce a desired sequence, e.g., a polynucleotide encoding all or part of a transcription factor. For example, chemical synthesis using the phosphoramidite method is described, e.g., by Beaucage et al. (1981) *Tetrahedron Letters* 22:1859-69; and Matthes et al. (1984) *EMBO J.* 3:801-5. According to such methods, oligonucleotides are synthesized, purified, annealed to their complementary strand, ligated and then optionally cloned into suitable vectors. And if so desired, the polynucleotides and polypeptides of the invention can be custom ordered from any of a number of commercial suppliers.

[0060] Homologous Sequences

[0061] Sequences homologous, i.e., that share significant sequence identity or similarity, to those provided in the Sequence Listing, derived from *Arabidopsis thaliana* or from other plants of choice are also an aspect of the invention. Homologous sequences can be derived from any plant including monocots and dicots and in particular agriculturally important plant species, including but not limited to, crops such as soybean, wheat, corn, potato, cotton, rice, oilseed rape (including canola), sunflower, alfalfa, sugarcane and turf; or fruits and vegetables, such as banana, blackberry, blueberry, strawberry, and raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers,

pineapple, pumpkin, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits (such as apple, peach, pear, cherry and plum) and vegetable brassicas (such as broccoli, cabbage, cauliflower, brussels sprouts and kohlrabi). Other crops, fruits and vegetables whose phenotype can be changed include barley, rye, millet, sorghum, currant, avocado, citrus fruits such as oranges, lemons, grapefruit and tangerines, artichoke, cherries, nuts such as the walnut and peanut, endive, leek, roots, such as arrowroot, beet, cassava, turnip, radish, yam, and sweet potato, and beans. The homologous sequences may also be derived from woody species, such as pine, poplar and eucalyptus, or mint or other labiates.

[0062] Transcription factors that are homologous to the listed sequences will typically share at least about 30% amino acid sequence identity, or at least about 30% amino acid sequence identity outside of a known consensus sequence or consensus DNA-binding site. More closely related transcription factors can share at least about 50%, about 60%, about 65%, about 70%, about 75% or about 80% or about 90% or about 95% or about 98% or more sequence identity with the listed sequences, or with the listed sequences but excluding or outside a known consensus sequence or consensus DNA-binding site, or with the listed sequences excluding one or all conserved domain. Factors that are most closely related to the listed sequences share, e.g., at least about 85%, about 90% or about 95% or more % sequence identity to the listed sequences, or to the listed sequences but excluding or outside a known consensus sequence or consensus DNA-binding site or outside one or all conserved domain. At the nucleotide level, the sequences will typically share at least about 40% nucleotide sequence identity, preferably at least about 50%, about 60%, about 70% or about 80% sequence identity, and more preferably about 85%, about 90%, about 95% or about 97% or more sequence identity to one or more of the listed sequences, or to a listed sequence but excluding or outside a known consensus sequence or consensus DNA-binding site, or outside one or all conserved domain. The degeneracy of the genetic code enables major variations in the nucleotide sequence of a polynucleotide while maintaining the amino acid sequence of the encoded protein. Conserved domains within a transcription factor family may exhibit a higher degree of sequence homology, such as at least 65% sequence identity including conservative substitutions, and preferably at least 80% sequence identity, and more preferably at least 85%, or at least about 86%, or at least about 87%, or at least about 88%, or at least about 90%, or at least about 95%, or at least about 98% sequence identity. Transcription factors that are homologous to the listed sequences should share at least 30%, or at least about 60%, or at least about 75%, or at least about 80%, or at least about 90%, or at least about 95% amino acid sequence identity over the entire length of the polypeptide or the homolog. In addition, transcription factors that are homologous to the listed sequences should share at least 30%, or at least about 60%, or at least about 75%, or at least about 80%, or at least about 90%, or at least about 95% amino acid sequence similarity over the entire length of the polypeptide or the homolog.

[0063] Identifying Orthologs And Paralogs

[0064] Several different methods are known by those of skill in the art for identifying and defining these functionally homologous sequences. Three general methods for defining

paralogs and orthologs are described; a paralog or ortholog may be identified by only one or more of the methods described below.

[0065] Orthologs and paralogs are evolutionarily related genes that have similar sequence and similar functions. Paralogs are related genes within a single species and are most likely a result of gene duplication, whereas orthologs are related genes in different species derived from a common ancestral molecule prior to speciation.

[0066] Within a single plant species, gene duplication may cause two copies of a particular gene, giving rise to two or more genes with similar sequence and similar function known as paralogs. A paralog is therefore a similar gene with a similar function within the same species. Paralogs typically cluster together or in the same clade (a group of similar genes) when a gene family phylogeny is analyzed using programs such as CLUSTAL (Thompson et al. (1994) *Nucleic Acids Res.* 22:4673-4680; Higgins et al. (1996) *Methods Enzymol.* 266:383-402). Groups of similar genes can also be identified using pair-wise BLAST analysis (Feng and Doolittle (1987) *J. Mol. Evol.* 25:351-360). For example, a clade of very similar MADS domain transcription factors from *Arabidopsis* all share a common function in flowering time (Ratcliffe et al. (2001) *Plant Physiol.* 126:122-132), and a group of very similar AP2 domain transcription factors from *Arabidopsis* are involved in tolerance of plants to freezing (Gilmour et al. (1998) *Plant J.* 16:433-442). Analysis of groups of similar genes with similar function that fall within one clade can yield sub-sequences that are particular to the clade. These sub-sequences, known as consensus sequences, can not only be used to define the sequences within each clade, but define the functions of these genes, since genes within each clade typically share the same function. (See also, for example, Mount, D. W. (2001) *Bioinformatics: Sequence and Genome Analysis* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. page 543.)

[0067] Speciation, the production of new species from a parental species, can also give rise to two or more genes with similar sequence and similar function. These genes, termed orthologs, often have an identical function within their host plants and are often interchangeable between species without losing function. Because plants have common ancestors, many genes in any plant species will have a corresponding orthologous gene in another plant species. Once a phylogenetic tree for a gene family of one species has been constructed using a program such as CLUSTAL (Thompson et al. (1994) *Nucleic Acids Res.* 22:4673-4680; Higgins et al. (1996) *Methods Enzymol.* 266:383-402), potential orthologous sequences can be placed into the phylogenetic tree and its relationship to genes from the species of interest can be determined. Once the ortholog pair has been identified, the function of the test ortholog can be determined by determining the function of the reference ortholog.

[0068] Orthologs can also be identified by pair-wise BLAST analysis by aligning a set of reference sequences against a set of test sequences. Test sequences with the closest match to a particular reference sequence, as determined by the P-value of the BLAST analysis, can be taken and individually aligned against the reference set of sequences. The individual test sequence will either best

match the particular reference sequence, in which case it is likely to be an ortholog, or not, in which case it may not be an ortholog.

[0069] A further way of identifying an ortholog is by identifying a consensus sequence within the candidate ortholog. Using pair-wise BLAST analysis, or programs such as CLUSTAL alignment program, sets of similar genes, or clades, can be identified. The particular sub-sequences which defining within a particular clade have in common to differentiate themselves can be derived from an alignment of those sequences. Orthologs would have the consensus sequence, or a sequence similar to the consensus sequence. Orthologs might also have a consensus sequence outside a conserved domain, which could be particular to that family of orthologous sequences.

[0070] Corresponding orthologs may bridge the monocot/dicot division of the plant kingdom and orthologous pairs of genes can be identified in rice and Arabidopsis, corn and Arabidopsis and Antirrhinum and corn. For example Peng et al showed that a mutant of the Arabidopsis gene termed *Gibberellin Insensitive* (GAI; mutant termed gai) encoded a transcription factor and which conferred a reduction in gibberellin responsiveness in the native plant (Peng et al. 1997 Genes and Development 11:3194-3205). In addition, Peng et al. subsequently showed that the Arabidopsis GAI protein has 62 % amino acid residue identity with the wheat Rht-D 1 a protein and 62% amino acid residue identity with the maize d8. Peng et al. showed that transgenic rice plants containing a mutant GAI allele give reduced response to gibberellin and are dwarfed, mimicking the dwarfed wheat variety from which the mutant Rht-D1a gene was isolated. Peng et al. taught that Arabidopsis GAI protein is an ortholog of the wheat Rht-D1a and maize d8 proteins. (Peng et al. 1999 Nature 400:256-261.)

[0071] In addition Fu et al. (2001 Plant Cell 13:1791-1802), Nandi et al. (2000 Curr. Biol. 10:215-218), Coupland (1995 Nature 377:482-483), and Weigel and Nilsson (1995 Nature 377:482-500) show that an Arabidopsis transcription factor expressed in an exogenous plant species elicits the same or very similar phenotypic response. Furthermore, Kater et al. (1998 Plant Cell 10:171-182), Mandel et al. (1992 Cell 71:133-143), and Suzuki et al. (2001 Plant J. 28:409-418) showed that a transcription factor expressed in another plant species elicits the same or very similar phenotypic response of the endogenous sequence, as often predicted in earlier studies of Arabidopsis transcription factors in Arabidopsis.

[0072] Identifying Polynucleotides Or Nucleic Acids By Hybridization

[0073] Polynucleotides homologous to the sequences illustrated in the Sequence Listing can be identified, e.g., by hybridization to each other under stringent or under highly stringent conditions. Single stranded polynucleotides hybridize when they associate based on a variety of well characterized physical-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. The stringency of a hybridization reflects the degree of sequence identity of the nucleic acids involved, such that the higher the stringency, the more similar are the two polynucleotide strands. Stringency is influenced by a variety of factors, including temperature, salt concentration and composition, organic and non-organic additives, solvents, etc. present in

both the hybridization and wash solutions and incubations (and number), as described in more detail in the references cited above.

[0074] An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is about 5° C. to 20° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Nucleic acid molecules that hybridize under stringent conditions will typically hybridize to a probe based on either the entire cDNA or selected portions, e.g., to a unique subsequence, of the cDNA under wash conditions of 0.2× SSC to 2.0× SSC, 0.1% SDS at 50-65° C. For example, high stringency is about 0.2× SSC, 0.1% SDS at 65° C. Ultra-high stringency will be the same conditions except the wash temperature is raised about 3 or about 5° C., and ultra-ultra-high stringency will be the same conditions except the wash temperature is raised about 6 or about 9° C. For identification of less closely related homologues washes can be performed at a lower temperature, e.g., 50° C. In general, stringency is increased by raising the wash temperature and/or decreasing the concentration of SSC, as known in the art.

[0075] As another example, stringent conditions can be selected such that an oligonucleotide that is perfectly complementary to the coding oligonucleotide hybridizes to the coding oligonucleotide with at least about a 5-10× higher signal to noise ratio than the ratio for hybridization of the perfectly complementary oligonucleotide to a nucleic acid encoding a transcription factor known as of the filing date of the application. Conditions can be selected such that a higher signal to noise ratio is observed in the particular assay which is used, e.g., about 15×, 25×, 35×, 50× or more. Accordingly, the subject nucleic acid hybridizes to the unique coding oligonucleotide with at least a 2× higher signal to noise ratio as compared to hybridization of the coding oligonucleotide to a nucleic acid encoding known polypeptide. Again, higher signal to noise ratios can be selected, e.g., about 5×, 10×, 25×, 35×, 50× or more. The particular signal will depend on the label used in the relevant assay, e.g., a fluorescent label, a colorimetric label, a radioactive label, or the like.

[0076] Alternatively, transcription factor homolog polypeptides can be obtained by screening an expression library using antibodies specific for one or more transcription factors. With the provision herein of the disclosed transcription factor, and transcription factor homologue nucleic acid sequences, the encoded polypeptide(s) can be expressed and purified in a heterologous expression system (e.g., *E. coli*) and used to raise antibodies (monoclonal or polyclonal) specific for the polypeptide(s) in question. Antibodies can also be raised against synthetic peptides derived from transcription factor, or transcription factor homologue, amino acid sequences. Methods of raising antibodies are well known in the art and are described in Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York. Such antibodies can then be used to screen an expression library produced from the plant from which it is desired to clone additional transcription factor

homologues, using the methods described above. The selected cDNAs can be confirmed by sequencing and enzymatic activity.

[0077] Sequence Variations

[0078] It will readily be appreciated by those of skill in the art, that any of a variety of polynucleotide sequences are capable of encoding the transcription factors and transcription factor homologue polypeptides of the invention. Due to the degeneracy of the genetic code, many different polynucleotides can encode identical and/or substantially similar polypeptides in addition to those sequences illustrated in the Sequence Listing.

[0079] For example, Table 1 illustrates, e.g., that the codons AGC, AGT, TCA, TCC, TCG, and TCT all encode the same amino acid: serine. Accordingly, at each position in the sequence where there is a codon encoding serine, any of the above trinucleotide sequences can be used without altering the encoded polypeptide.

TABLE 1			
Amino acid			Possible Codons
Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	TGC TGT
Aspartic acid	Asp	D	GAC GAT
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	TTC TTT
Glycine	Gly	G	GGA GGC GGG GGT
Histidine	His	H	CAC CAT
Isoleucine	Ile	I	ATA ATC ATT
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	TTA TTG CTA CTC CTG CTT
Methionine	Met	M	ATG
Asparagine	Asn	N	AAC AAT
Proline	Pro	P	CCA CCC CCG CCT
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGT
Serine	Ser	S	AGC AGT TCA TCC TCG TCT
Threonine	Thr	T	ACA ACC ACG ACT
Valine	Val	V	GTA GTC GTG GTT
Tryptophan	Trp	W	TGG
Tyrosine	Tyr	Y	TAC TAT

[0080] Sequence alterations that do not change the amino acid sequence encoded by the polynucleotide are termed "silent" variations. With the exception of the codons ATG and TGG, encoding methionine and tryptophan, respectively, any of the possible codons for the same amino acid can be substituted by a variety of techniques, e.g., site-directed mutagenesis, available in the art. Accordingly, any

and all such variations of a sequence selected from the above table are a feature of the invention.

[0081] In addition to silent variations, other conservative variations that alter one, or a few amino acids in the encoded polypeptide, can be made without altering the function of the polypeptide, these conservative variants are, likewise, a feature of the invention.

[0082] For example, substitutions, deletions and insertions introduced into the sequences provided in the Sequence Listing are also envisioned by the invention. Such sequence modifications can be engineered into a sequence by site-directed mutagenesis (Wu (ed.) *Meth. Enzymol.* (1993) vol. 217, Academic Press) or the other methods noted below. Amino acid substitutions are typically of single residues; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. In preferred embodiments, deletions or insertions are made in adjacent pairs, e.g., a deletion of two residues or insertion of two residues. Substitutions, deletions, insertions or any combination thereof can be combined to arrive at a sequence. The mutations that are made in the polynucleotide encoding the transcription factor should not place the sequence out of reading frame and should not create complementary regions that could produce secondary mRNA structure. Preferably, the polypeptide encoded by the DNA performs the desired function.

[0083] Conservative substitutions are those in which at least one residue in the amino acid sequence has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the Table 2 when it is desired to maintain the activity of the protein. Table 2 shows amino acids which can be substituted for an amino acid in a protein and which are typically regarded as conservative substitutions.

TABLE 2	
Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Gln	Asn
Cys	Ser
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu, Val
Leu	Ile; Val
Lys	Arg; Gln
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr; Gly
Thr	Ser; Val
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

[0084] Similar substitutions are those in which at least one residue in the amino acid sequence has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the Table 3 when it is desired to maintain the activity of the protein. Table 3 shows amino acids which can be substituted for an amino

acid in a protein and which are typically regarded as structural and functional substitutions. For example, a residue in column 1 of Table 3 may be substituted with residue in column 2; in addition, a residue in column 2 of Table 3 may be substituted with the residue of column 1.

TABLE 3

Residue	Similar Substitutions
Ala	Ser; Thr; Gly; Val; Leu; Ile
Arg	Lys; His; Gly
Asn	Gln; His; Gly; Ser; Thr
Asp	Glu; Ser; Thr
Gln	Asn; Ala
Cys	Ser; Gly
Glu	Asp
Gly	Pro; Arg
His	Asn; Gln; Tyr; Phe; Lys; Arg
Ile	Ala; Leu; Val; Gly; Met
Leu	Ala; Ile; Val; Gly; Met
Lys	Arg; His; Gln; Gly; Pro
Met	Leu; Ile; Phe
Phe	Met; Leu; Tyr; Trp; His; Val; Ala
Ser	Thr; Gly; Asp; Ala; Val; Ile; His
Thr	Ser; Val; Ala; Gly
Trp	Tyr; Phe; His
Tyr	Trp; Phe; His
Val	Ala; Ile; Leu; Gly; Thr; Ser; Glu

[0085] Substitutions that are less conservative than those in Table 2 or Table 3 can be selected by picking residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in protein properties will be those in which (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

[0086]

[0087] Further Modifying Sequences of the Invention—Mutation/Forced Evolution

[0088] In addition to generating silent or conservative substitutions as noted, above, the present invention optionally includes methods of modifying the sequences of the Sequence Listing. In the methods, nucleic acid or protein modification methods are used to alter the given sequences to produce new sequences and/or to chemically or enzymatically modify given sequences to change the properties of the nucleic acids or proteins. These sequences may be synthetic polynucleotides and synthetic polypeptides.

[0089] Thus, in one embodiment, given nucleic acid sequences are modified, e.g., according to standard mutagenesis or artificial evolution methods to produce modified sequences. For example, Ausubel, supra, provides additional details on mutagenesis methods. Artificial forced evolution methods are described, e.g., by Stemmer (1994)

Nature 370:389-391, and Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751, and U.S. Pat. Nos. 5,811,238, 5,837,500, and 6,242,568. Many other mutation and evolution methods are also available and expected to be within the skill of the practitioner. A synthetic polypeptide encoded by a synthetic polynucleotide may have as little as 20% amino acid residue sequence identity to the polypeptide encoded by the claimed polynucleotides and still modify a plant's trait or characteristic.

[0090] Similarly, chemical or enzymatic alteration of expressed nucleic acids and polypeptides can be performed by standard methods. For example, sequence can be modified by addition of lipids, sugars, peptides, organic or inorganic compounds, by the inclusion of modified nucleotides or amino acids, or the like. For example, protein modification techniques are illustrated in Ausubel, supra. Further details on chemical and enzymatic modifications can be found herein. These modification methods can be used to modify any given sequence, or to modify any sequence produced by the various mutation and artificial evolution modification methods noted herein.

[0091] Accordingly, the invention provides for modification of any given nucleic acid by mutation, evolution, chemical or enzymatic modification, or other available methods, as well as for the products produced by practicing such methods, e.g., using the sequences herein as a starting substrate for the various modification approaches.

[0092] For example, optimized coding sequence containing codons preferred by a particular prokaryotic or eukaryotic host can be used e.g., to increase the rate of translation or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, as compared with transcripts produced using a non-optimized sequence. Translation stop codons can also be modified to reflect host preference. For example, preferred stop codons for *S. cerevisiae* and mammals are TAA and TGA, respectively. The preferred stop codon for monocotyledonous plants is TGA, whereas insects and *E. coli* prefer to use TAA as the stop codon.

[0093] The polynucleotide sequences of the present invention can also be engineered in order to alter a coding sequence for a variety of reasons, including but not limited to, alterations which modify the sequence to facilitate cloning, processing and/or expression of the gene product. For example, alterations are optionally introduced using techniques which are well known in the art, e.g., site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to introduce splice sites, etc.

[0094] Furthermore, a fragment or domain derived from any of the polypeptides of the invention can be combined with domains derived from other transcription factors or synthetic domains to modify the biological activity of a transcription factor. For instance, a DNA-binding domain derived from a transcription factor of the invention can be combined with the activation domain of another transcription factor or with a synthetic activation domain. A transcription activation domain assists in initiating transcription from a DNA-binding site. Examples include the transcription activation region of VP16 or GAL4 (Moore et al. (1998) *Proc. Natl. Acad. Sci. USA* 95: 376-381; and Aoyama et al. (1995) *Plant Cell* 7:1773-1785), peptides derived from

bacterial sequences (Ma and Ptashne (1987) *Cell* 51: 113-119) and synthetic peptides (Giniger and Ptashne, (1987) *Nature* 330:670-672).

[0095] Expression and Modification of Polypeptides

[0096] Typically, polynucleotide sequences of the invention are incorporated into recombinant DNA (or RNA) molecules that direct expression of polypeptides of the invention in appropriate host cells, transgenic plants, in vitro translation systems, or the like. Due to the inherent degeneracy of the genetic code, nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence can be substituted for any listed sequence to provide for cloning and expressing the relevant homologue.

[0097] Vectors, Promoters, and Expression Systems

[0098] The present invention includes recombinant constructs comprising one or more of the nucleic acid sequences herein. The constructs typically comprise a vector, such as a plasmid, a cosmid, a phage, a virus (e.g., a plant virus), a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), or the like, into which a nucleic acid sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

[0099] General texts which describe molecular biological techniques useful herein, including the use and production of vectors, promoters and many other relevant topics, include Berger, Sambrook and Ausubel, supra. Any of the identified sequences can be incorporated into a cassette or vector, e.g., for expression in plants. A number of expression vectors suitable for stable transformation of plant cells or for the establishment of transgenic plants have been described including those described in Weissbach and Weissbach, (1989) *Methods for Plant Molecular Biology*, Academic Press, and Gelvin et al., (1990) *Plant Molecular Biology Manual*, Kluwer Academic Publishers. Specific examples include those derived from a Ti plasmid of *Agrobacterium tumefaciens*, as well as those disclosed by Herrera-Estrella et al. (1983) *Nature* 303: 209, Bevan (1984) *Nucl Acid Res.* 12: 8711-8721, Klee (1985) *Bio/Technology* 3: 637-642, for dicotyledonous plants.

[0100] Alternatively, non-Ti vectors can be used to transfer the DNA into monocotyledonous plants and cells by using free DNA delivery techniques. Such methods can involve, for example, the use of liposomes, electroporation, microprojectile bombardment, silicon carbide whiskers, and viruses. By using these methods transgenic plants such as wheat, rice (Christou (1991) *Bio/Technology* 9: 957-962) and corn (Gordon-Kamm (1990) *Plant Cell* 2: 603-618) can be produced. An immature embryo can also be a good target tissue for monocots for direct DNA delivery techniques by using the particle gun (Weeks et al. (1993) *Plant Physiol* 102: 1077-1084; Vasil (1993) *Bio/Technology* 10: 667-674; Wan and Lemeaux (1994) *Plant Physiol* 104: 37-48, and for *Agrobacterium*-mediated DNA transfer (Ishida et al. (1996) *Nature Biotech* 14: 745-750).

[0101] Typically, plant transformation vectors include one or more cloned plant coding sequence (genomic or cDNA)

under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant transformation vectors typically also contain a promoter (e.g., a regulatory region controlling inducible or constitutive, environmentally-or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, an RNA processing signal (such as intron splice sites), a transcription termination site, and/or a polyadenylation signal.

[0102] Examples of constitutive plant promoters which can be useful for expressing the TF sequence include: the cauliflower mosaic virus (CaMV) 35S promoter, which confers constitutive, high-level expression in most plant tissues (see, e.g., Odel et al. (1985) *Nature* 313:810); the nopaline synthase promoter (An et al. (1988) *Plant Physiol* 88:547); and the octopine synthase promoter (Fromm et al. (1989) *Plant Cell* 1: 977).

[0103] A variety of plant gene promoters that regulate gene expression in response to environmental, hormonal, chemical, developmental signals, and in a tissue-active manner can be used for expression of a TF sequence in plants. Choice of a promoter is based largely on the phenotype of interest and is determined by such factors as tissue (e.g., seed, fruit, root, pollen, vascular tissue, flower, carpel, etc.), inducibility (e.g., in response to wounding, heat, cold, drought, light, pathogens, etc.), timing, developmental stage, and the like. Numerous known promoters have been characterized and can favorably be employed to promote expression of a polynucleotide of the invention in a transgenic plant or cell of interest. For example, tissue specific promoters include: seed-specific promoters (such as the napin, phaseolin or DC3 promoter described in U.S. Pat. No. 5,773,697), fruit-specific promoters that are active during fruit ripening (such as the dru 1 promoter (U.S. Pat. No. 5,783,393), or the 2A11 promoter (U.S. Pat. No. 4,943,674) and the tomato polygalacturonase promoter (Bird et al. (1988) *Plant Mol Biol* 11:651), root-specific promoters, such as those disclosed in U.S. Pat. Nos. 5,618,988, 5,837,848 and 5,905,186, pollen-active promoters such as PTA29, PTA26 and PTA13 (U.S. Pat. No. 5,792,929), promoters active in vascular tissue (Ringli and Keller (1998) *Plant Mol Biol* 37:977-988), flower-specific (Kaiser et al. (1995) *Plant Mol Biol* 28:231-243), pollen (Baerson et al. (1994) *Plant Mol Biol* 26:1947-1959), carpels (Ohl et al. (1990) *Plant Cell* 2:837-848), pollen and ovules (Baerson et al. (1993) *Plant Mol Biol* 22:255-267), auxin-inducible promoters (such as that described in van der Kop et al. (1999) *Plant Mol Biol* 39:979-990 or Baumann et al. (1999) *Plant Cell* 11:323-334), cytokinin-inducible promoter (Guevara-Garcia (1998) *Plant Mol Biol* 38:743-753), promoters responsive to gibberellin (Shi et al. (1998) *Plant Mol Biol* 38:1053-1060, Willmott et al. (1998) 38:817-825) and the like. Additional promoters are those that elicit expression in response to heat (Ainley et al. (1993) *Plant Mol Biol* 22: 13-23), light (e.g., the pea rbcS-3A promoter, Kuhlmeier et al. (1989) *Plant Cell* 1:471, and the maize rbcS promoter, Schaffner and Sheen (1991) *Plant Cell* 3: 997); wounding (e.g., wun1, Siebertz et al. (1989) *Plant Cell* 1: 961); pathogens (such as the PR-1 promoter described in Buchel et al. (1999) *Plant Mol. Biol.* 40:387-396, and the PDF1.2 promoter described in Manners et al. (1998) *Plant Mol. Biol.* 38:1071-80), and chemicals such as methyl jasmonate or salicylic acid (Gatz et al. (1997) *Annu Rev Plant Physiol Plant Mol Biol* 48: 89-108). In addition, the timing of the expression can be

controlled by using promoters such as those acting at senescence (Gan and Amasino (1995) *Science* 270: 1986-1988); or late seed development (Odell et al. (1994) *Plant Physiol* 106:447-458).

[0104] Plant expression vectors can also include RNA processing signals that can be positioned within, upstream or downstream of the coding sequence. In addition, the expression vectors can include additional regulatory sequences from the 3'-untranslated region of plant genes, e.g., a 3' terminator region to increase mRNA stability of the mRNA, such as the PI-II terminator region of potato or the octopine or nopaline synthase 3' terminator regions.

[0105] Additional Expression Elements

[0106] Specific initiation signals can aid in efficient translation of coding sequences. These signals can include, e.g., the ATG initiation codon and adjacent sequences. In cases where a coding sequence, its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only coding sequence (e.g., a mature protein coding sequence), or a portion thereof, is inserted, exogenous transcriptional control signals including the ATG initiation codon can be separately provided. The initiation codon is provided in the correct reading frame to facilitate transcription. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

[0107] Expression Hosts

[0108] The present invention also relates to host cells which are transduced with vectors of the invention, and the production of polypeptides of the invention (including fragments thereof) by recombinant techniques. Host cells are genetically engineered (i.e., nucleic acids are introduced, e.g., transduced, transformed or transfected) with the vectors of this invention, which may be, for example, a cloning vector or an expression vector comprising the relevant nucleic acids herein. The vector is optionally a plasmid, a viral particle, a phage, a naked nucleic acid, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the relevant gene. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, Sambrook and Ausubel.

[0109] The host cell can be a eukaryotic cell, such as a yeast cell, or a plant cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Plant protoplasts are also suitable for some applications. For example, the DNA fragments are introduced into plant tissues, cultured plant cells or plant protoplasts by standard methods including electroporation (Fromm et al., (1985) *Proc. Natl. Acad. Sci. USA* 82, 5824, infection by viral vectors such as cauliflower mosaic virus (CaMV) (Hohn et al., (1982) *Molecular Biology of Plant Tumors*, (Academic Press, New York) pp. 549-560; U.S. Pat. No. 4,407,956), high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the

surface (Klein et al., (1987) *Nature* 327, 70-73), use of pollen as vector (WO 85/01856), or use of *Agrobacterium tumefaciens* or *A. rhizogenes* carrying a T-DNA plasmid in which DNA fragments are cloned. The T-DNA plasmid is transmitted to plant cells upon infection by *Agrobacterium tumefaciens*, and a portion is stably integrated into the plant genome (Horsch et al. (1984) *Science* 233:496-498; Fraley et al. (1983) *Proc. Natl. Acad. Sci. USA* 80, 4803).

[0110] The cell can include a nucleic acid of the invention which encodes a polypeptide, wherein the cells expresses a polypeptide of the invention. The cell can also include vector sequences, or the like. Furthermore, cells and transgenic plants which include any polypeptide or nucleic acid above or throughout this specification, e.g., produced by transduction of a vector of the invention, are an additional feature of the invention.

[0111] For long-term, high-yield production of recombinant proteins, stable expression can be used. Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The protein or fragment thereof produced by a recombinant cell may be secreted, membrane-bound, or contained intracellularly, depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides encoding mature proteins of the invention can be designed with signal sequences which direct secretion of the mature polypeptides through a prokaryotic or eukaryotic cell membrane.

[0112] Modified Amino Acids

[0113] Polypeptides of the invention may contain one or more modified amino acids. The presence of modified amino acids may be advantageous in, for example, increasing polypeptide half-life, reducing polypeptide antigenicity or toxicity, increasing polypeptide storage stability, or the like. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production or modified by synthetic or chemical means.

[0114] Non-limiting examples of a modified amino acid include incorporation or other use of acetylated amino acids, glycosylated amino acids, sulfated amino acids, prenylated (e.g., farnesylated, geranylgeranylated) amino acids, PEG modified (e.g., "PEGylated") amino acids, biotinylated amino acids, carboxylated amino acids, phosphorylated amino acids, etc. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature.

[0115] The modified amino acids may prevent or increase affinity of the polypeptide for another molecule, including, but not limited to, polynucleotide, proteins, carbohydrates, lipids and lipid derivatives, and other organic or synthetic compounds.

[0116] Identification of Additional Factors

[0117] A transcription factor provided by the present invention can also be used to identify additional endogenous or exogenous molecules that can affect a phenotype or trait of interest. On the one hand, such molecules include organic (small or large molecules) and/or inorganic compounds that affect expression of (i.e., regulate) a particular transcription

factor. Alternatively, such molecules include endogenous molecules that are acted upon either at a transcriptional level by a transcription factor of the invention to modify a phenotype as desired. For example, the transcription factors can be employed to identify one or more downstream gene with which is subject to a regulatory effect of the transcription factor. In one approach, a transcription factor or transcription factor homologue of the invention is expressed in a host cell, e.g., a transgenic plant cell, tissue or explant, and expression products, either RNA or protein, of likely or random targets are monitored, e.g., by hybridization to a microarray of nucleic acid probes corresponding to genes expressed in a tissue or cell type of interest, by two-dimensional gel electrophoresis of protein products, or by any other method known in the art for assessing expression of gene products at the level of RNA or protein. Alternatively, a transcription factor of the invention can be used to identify promoter sequences (i.e., binding sites) involved in the regulation of a downstream target. After identifying a promoter sequence, interactions between the transcription factor and the promoter sequence can be modified by changing specific nucleotides in the promoter sequence or specific amino acids in the transcription factor that interact with the promoter sequence to alter a plant trait. Typically, transcription factor DNA-binding sites are identified by gel shift assays. After identifying the promoter regions, the promoter region sequences can be employed in double-stranded DNA arrays to identify molecules that affect the interactions of the transcription factors with their promoters (Bulyk et al. (1999) *Nature Biotechnology* 17:573-577). A test promoter region element of a transcription factor gene may also be screened using a phage-display analysis and a phage library which comprises polynucleotides encoding any transcription factor to identify a transcription factor so encoded which binds to the test promoter region element. Such phage-display methods are well known in the art.

[0118] The identified transcription factors are also useful to identify proteins that modify the activity of the transcription factor. Such modification can occur by covalent modification, such as by phosphorylation, or by protein-protein (homo or-heteropolymer) interactions. Any method suitable for detecting protein-protein interactions can be employed. Among the methods that can be employed are co-immunoprecipitation, cross-linking and co-purification through gradients or chromatographic columns, and the two-hybrid yeast system.

[0119] The two-hybrid system detects protein interactions in vivo and is described in Chien, et al., (1991), *Proc. Natl. Acad. Sci. USA* 88, 9578-9582 and is commercially available from Clontech (Palo Alto, Calif.). In such a system, plasmids are constructed that encode two hybrid proteins: one consists of the DNA-binding domain of a transcription activator protein fused to the TF polypeptide and the other consists of the transcription activator protein's activation domain fused to an unknown protein that is encoded by a cDNA that has been recombined into the plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the cDNA library are transformed into a strain of the yeast *Saccharomyces cerevisiae* that contains a reporter gene (e.g., lacZ) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene. Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is

detected by an assay for the reporter gene product. Then, the library plasmids responsible for reporter gene expression are isolated and sequenced to identify the proteins encoded by the library plasmids. After identifying proteins that interact with the transcription factors, assays for compounds that interfere with the TF protein-protein interactions can be preformed.

[0120] Identification of Modulators

[0121] In addition to the intracellular molecules described above, extracellular molecules that alter activity or expression of a transcription factor, either directly or indirectly, can be identified. For example, the methods can entail first placing a candidate molecule in contact with a plant or plant cell. The molecule can be introduced by topical administration, such as spraying or soaking of a plant, and then the molecule's effect on the expression or activity of the TF polypeptide or the expression of the polynucleotide monitored. Changes in the expression of the TF polypeptide can be monitored by use of polyclonal or monoclonal antibodies, gel electrophoresis or the like. Changes in the expression of the corresponding polynucleotide sequence can be detected by use of microarrays, Northern, quantitative PCR, or any other technique for monitoring changes in mRNA expression. These techniques are exemplified in Ausubel et al. (eds) *Current Protocols in Molecular Biology*, John Wiley & Sons (1998, and supplements through 2001). Such changes in the expression levels can be correlated with modified plant traits and thus identified molecules can be useful for soaking or spraying on fruit, vegetable and grain crops to modify traits in plants.

[0122] Essentially any available composition can be tested for modulatory activity of expression or activity of any nucleic acid or polypeptide herein. Thus, available libraries of compounds such as chemicals, polypeptides, nucleic acids and the like can be tested for modulatory activity. Often, potential modulator compounds can be dissolved in aqueous or organic (e.g., DMSO-based) solutions for easy delivery to the cell or plant of interest in which the activity of the modulator is to be tested. Optionally, the assays are designed to screen large modulator composition libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtitre formats on microtitre plates in robotic assays).

[0123] In one embodiment, high throughput screening methods involve providing a combinatorial library containing a large number of potential compounds (potential modulator compounds). Such "combinatorial chemical libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as target compounds.

[0124] A combinatorial chemical library can be, e.g., a collection of diverse chemical compounds generated by chemical synthesis or biological synthesis. For example, a combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (e.g., in one example, amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound of a set length). Exemplary libraries include peptide libraries, nucleic acid libraries, antibody

libraries (see, e.g., Vaughn et al. (1996) *Nature Biotechnology*, 14(3):309-314 and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al. *Science* (1996) 274:1520-1522 and U.S. Pat. No. 5,593,853), peptide nucleic acid libraries (see, e.g., U.S. Pat. No. 5,539,083), and small organic molecule libraries (see, e.g., benzodiazepines, Baum *C&EN* January 18, page 33 (1993); isoprenoids, U.S. Pat. No. 5,569,588; thiazolidinones and metathiazanones, U.S. Pat. No. 5,549,974; pyrrolidines, U.S. Pat. Nos. 5,525,735 and 5,519,134; morpholino compounds, U.S. Pat. No. 5,506,337) and the like.

[0125] Preparation and screening of combinatorial or other libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Pat. No. 5,010,175, Furka, *Int. J. Pept. Prot. Res.* 37:487-493 (1991) and Houghton et al. *Nature* 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used.

[0126] In addition, as noted, compound screening equipment for high-throughput screening is generally available, e.g., using any of a number of well known robotic systems that have also been developed for solution phase chemistries useful in assay systems. These systems include automated workstations including an automated synthesis apparatus and robotic systems utilizing robotic arms. Any of the above devices are suitable for use with the present invention, e.g., for high-throughput screening of potential modulators. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art.

[0127] Indeed, entire high throughput screening systems are commercially available. These systems typically automate entire procedures including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. Similarly, microfluidic implementations of screening are also commercially available.

[0128] The manufacturers of such systems provide detailed protocols the various high throughput. Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like. The integrated systems herein, in addition to providing for sequence alignment and, optionally, synthesis of relevant nucleic acids, can include such screening apparatus to identify modulators that have an effect on one or more polynucleotides or polypeptides according to the present invention.

[0129] In some assays it is desirable to have positive controls to ensure that the components of the assays are working properly. At least two types of positive controls are appropriate. That is, known transcriptional activators or inhibitors can be incubated with cells/plants/etc. in one sample of the assay, and the resulting increase/decrease in transcription can be detected by measuring the resulting increase in RNA/protein expression, etc., according to the methods herein. It will be appreciated that modulators can also be combined with transcriptional activators or inhibitors to find modulators which inhibit transcriptional activation or transcriptional repression. Either expression of the nucleic

acids and proteins herein or any additional nucleic acids or proteins activated by the nucleic acids or proteins herein, or both, can be monitored.

[0130] In an embodiment, the invention provides a method for identifying compositions that modulate the activity or expression of a polynucleotide or polypeptide of the invention. For example, a test compound, whether a small or large molecule, is placed in contact with a cell, plant (or plant tissue or explant), or composition comprising the polynucleotide or polypeptide of interest and a resulting effect on the cell, plant, (or tissue or explant) or composition is evaluated by monitoring, either directly or indirectly, one or more of: expression level of the polynucleotide or polypeptide, activity (or modulation of the activity) of the polynucleotide or polypeptide. In some cases, an alteration in a plant phenotype can be detected following contact of a plant (or plant cell, or tissue or explant) with the putative modulator, e.g., by modulation of expression or activity of a polynucleotide or polypeptide of the invention. Modulation of expression or activity of a polynucleotide or polypeptide of the invention may also be caused by molecular elements in a signal transduction second messenger pathway and such modulation can affect similar elements in the same or another signal transduction second messenger pathway.

[0131] Subsequences

[0132] Also contemplated are uses of polynucleotides, also referred to herein as oligonucleotides, typically having at least 12 bases, preferably at least 15, more preferably at least 20, 30, or 50 bases, which hybridize under at least highly stringent (or ultra-high stringent or ultra-ultra- high stringent conditions) conditions to a polynucleotide sequence described above. The polynucleotides may be used as probes, primers, sense and antisense agents, and the like, according to methods as noted supra.

[0133] Subsequences of the polynucleotides of the invention, including polynucleotide fragments and oligonucleotides are useful as nucleic acid probes and primers. An oligonucleotide suitable for use as a probe or primer is at least about 15 nucleotides in length, more often at least about 18 nucleotides, often at least about 21 nucleotides, frequently at least about 30 nucleotides, or about 40 nucleotides, or more in length. A nucleic acid probe is useful in hybridization protocols, e.g., to identify additional polypeptide homologues of the invention, including protocols for microarray experiments. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other nucleic-acid amplification methods. See Sambrook and Ausubel, supra.

[0134] In addition, the invention includes an isolated or recombinant polypeptide including a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotides of the invention. For example, such polypeptides, or domains or fragments thereof, can be used as immunogens, e.g., to produce antibodies specific for the polypeptide sequence, or as probes for detecting a sequence of interest. A subsequence can range in size from about 15 amino acids in length up to and including the full length of the polypeptide.

[0135] Production of Transgenic Plants**[0136]** Modification of Traits

[0137] The polynucleotides of the invention are favorably employed to produce transgenic plants with various traits, or characteristics, that have been modified in a desirable manner, e.g., to improve the seed characteristics of a plant. For example, alteration of expression levels or patterns (e.g., spatial or temporal expression patterns) of one or more of the transcription factors (or transcription factor homologues) of the invention, as compared with the levels of the same protein found in a wild type plant, can be used to modify a plant's traits. An illustrative example of trait modification, improved characteristics, by altering expression levels of a particular transcription factor is described further in the Examples and the Sequence Listing.

[0138] Antisense and Cosuppression Approaches

[0139] In addition to expression of the nucleic acids of the invention as gene replacement or plant phenotype modification nucleic acids, the nucleic acids are also useful for sense and anti-sense suppression of expression, e.g., to down-regulate expression of a nucleic acid of the invention, e.g., as a further mechanism for modulating plant phenotype. That is, the nucleic acids of the invention, or subsequences or anti-sense sequences thereof, can be used to block expression of naturally occurring homologous nucleic acids. A variety of sense and anti-sense technologies are known in the art, e.g., as set forth in Lichtenstein and Nellen (1997) *Antisense Technology: A Practical Approach* IRL Press at Oxford University, Oxford, England. In general, sense or anti-sense sequences are introduced into a cell, where they are optionally amplified, e.g., by transcription. Such sequences include both simple oligonucleotide sequences and catalytic sequences such as ribozymes.

[0140] For example, a reduction or elimination of expression (i.e., a "knock-out") of a transcription factor or transcription factor homologue polypeptide in a transgenic plant, e.g., to modify a plant trait, can be obtained by introducing an antisense construct corresponding to the polypeptide of interest as a cDNA. For antisense suppression, the transcription factor or homologue cDNA is arranged in reverse orientation (with respect to the coding sequence) relative to the promoter sequence in the expression vector. The introduced sequence need not be the full length cDNA or gene, and need not be identical to the cDNA or gene found in the plant type to be transformed. Typically, the antisense sequence need only be capable of hybridizing to the target gene or RNA of interest. Thus, where the introduced sequence is of shorter length, a higher degree of homology to the endogenous transcription factor sequence will be needed for effective antisense suppression. While antisense sequences of various lengths can be utilized, preferably, the introduced antisense sequence in the vector will be at least 30 nucleotides in length, and improved antisense suppression will typically be observed as the length of the antisense sequence increases. Preferably, the length of the antisense sequence in the vector will be greater than 100 nucleotides. Transcription of an antisense construct as described results in the production of RNA molecules that are the reverse complement of mRNA molecules transcribed from the endogenous transcription factor gene in the plant cell.

[0141] Suppression of endogenous transcription factor gene expression can also be achieved using a ribozyme.

Ribozymes are RNA molecules that possess highly specific endoribonuclease activity. The production and use of ribozymes are disclosed in U.S. Pat. No. 4,987,071 and U.S. Pat. No. 5,543,508. Synthetic ribozyme sequences including antisense RNAs can be used to confer RNA cleaving activity on the antisense RNA, such that endogenous mRNA molecules that hybridize to the antisense RNA are cleaved, which in turn leads to an enhanced antisense inhibition of endogenous gene expression.

[0142] Vectors in which RNA encoded by a transcription factor or transcription factor homologue cDNA is over-expressed can also be used to obtain co-suppression of a corresponding endogenous gene, e.g., in the manner described in U.S. Pat. No. 5,231,020 to Jorgensen. Such co-suppression (also termed sense suppression) does not require that the entire transcription factor cDNA be introduced into the plant cells, nor does it require that the introduced sequence be exactly identical to the endogenous transcription factor gene of interest. However, as with antisense suppression, the suppressive efficiency will be enhanced as specificity of hybridization is increased, e.g., as the introduced sequence is lengthened, and/or as the sequence similarity between the introduced sequence and the endogenous transcription factor gene is increased.

[0143] Vectors expressing an untranslatable form of the transcription factor mRNA, e.g., sequences comprising one or more stop codon, or nonsense mutation) can also be used to suppress expression of an endogenous transcription factor, thereby reducing or eliminating its activity and modifying one or more traits. Methods for producing such constructs are described in U.S. Pat. No. 5,583,021. Preferably, such constructs are made by introducing a premature stop codon into the transcription factor gene. Alternatively, a plant trait can be modified by gene silencing using double-strand RNA (Sharp (1999) *Genes and Development* 13: 139-141).

[0144] Another method for abolishing the expression of a gene is by insertion mutagenesis using the T-DNA of *Agrobacterium tumefaciens*. After generating the insertion mutants, the mutants can be screened to identify those containing the insertion in a transcription factor or transcription factor homologue gene. Plants containing a single transgene insertion event at the desired gene can be crossed to generate homozygous plants for the mutation (Koncz et al. (1992) *Methods in Arabidopsis Research*. World Scientific).

[0145] Alternatively, a plant phenotype can be altered by eliminating an endogenous gene, such as a transcription factor or transcription factor homologue, e.g., by homologous recombination (Kempin et al. (1997) *Nature* 389:802).

[0146] A plant trait can also be modified by using the Cre-lox system (for example, as described in U.S. Pat. No. 5,658,772). A plant genome can be modified to include first and second lox sites that are then contacted with a Cre recombinase. If the lox sites are in the same orientation, the intervening DNA sequence between the two sites is excised. If the lox sites are in the opposite orientation, the intervening sequence is inverted.

[0147] The polynucleotides and polypeptides of this invention can also be expressed in a plant in the absence of an expression cassette by manipulating the activity or

expression level of the endogenous gene by other means. For example, by ectopically expressing a gene by T-DNA activation tagging (Ichikawa et al. (1997) *Nature* 390 698-701; Kakimoto et al. (1996) *Science* 274: 982-985). This method entails transforming a plant with a gene tag containing multiple transcriptional enhancers and once the tag has inserted into the genome, expression of a flanking gene coding sequence becomes deregulated. In another example, the transcriptional machinery in a plant can be modified so as to increase transcription levels of a polynucleotide of the invention (See, e.g., PCT Publications WO 96/06166 and WO 98/53057 which describe the modification of the DNA-binding specificity of zinc finger proteins by changing particular amino acids in the DNA-binding motif).

[0148] The transgenic plant can also include the machinery necessary for expressing or altering the activity of a polypeptide encoded by an endogenous gene, for example by altering the phosphorylation state of the polypeptide to maintain it in an activated state.

[0149] Transgenic plants (or plant cells, or plant explants, or plant tissues) incorporating the polynucleotides of the invention and/or expressing the polypeptides of the invention can be produced by a variety of well established techniques as described above. Following construction of a vector, most typically an expression cassette, including a polynucleotide, e.g., encoding a transcription factor or transcription factor homologue, of the invention, standard techniques can be used to introduce the polynucleotide into a plant, a plant cell, a plant explant or a plant tissue of interest. Optionally, the plant cell, explant or tissue can be regenerated to produce a transgenic plant.

[0150] The plant can be any higher plant, including gymnosperms, monocotyledonous and dicotyledonous plants. Suitable protocols are available for Leguminosae (alfalfa, soybean, clover, etc.), Umbelliferae (carrot, celery, parsnip), Cruciferae (cabbage, radish, rapeseed, broccoli, etc.), Curcubitaceae (melons and cucumber), Gramineae (wheat, corn, rice, barley, millet, etc.), Solanaceae (potato, tomato, tobacco, peppers, etc.), and various other crops. See protocols described in Ammirato et al. (1984) *Handbook of Plant Cell Culture—Crop Species*, Macmillan Publ. Co. Shimamoto et al. (1989) *Nature* 338:274-276; Fronim et al. (1990) *Bio/Technology* 8:833-839; and Vasil et al. (1990) *Bio/Technology* 8:429-434.

[0151] Transformation and regeneration of both monocotyledonous and dicotyledonous plant cells is now routine, and the selection of the most appropriate transformation technique will be determined by the practitioner. The choice of method will vary with the type of plant to be transformed; those skilled in the art will recognize the suitability of particular methods for given plant types. Suitable methods can include, but are not limited to: electroporation of plant protoplasts; liposome-mediated transformation; polyethylene glycol (PEG) mediated transformation; transformation using viruses; micro-injection of plant cells; micro-projectile bombardment of plant cells; vacuum infiltration; and *Agrobacterium tumefaciens* mediated transformation. Transformation means introducing a nucleotide sequence into a plant in a manner to cause stable or transient expression of the sequence.

[0152] Successful examples of the modification of plant characteristics by transformation with cloned sequences

which serve to illustrate the current knowledge in this field of technology, and which are herein incorporated by reference, include: U.S. Pat. Nos. 5,571,706; 5,677,175; 5,510,471; 5,750,386; 5,597,945; 5,589,615; 5,750,871; 5,268,526; 5,780,708; 5,538,880; 5,773,269; 5,736,369 and 5,610,042.

[0153] Following transformation, plants are preferably selected using a dominant selectable marker incorporated into the transformation vector. Typically, such a marker will confer antibiotic or herbicide resistance on the transformed plants, and selection of transformants can be accomplished by exposing the plants to appropriate concentrations of the antibiotic or herbicide.

[0154] After transformed plants are selected and grown to maturity, those plants showing a modified trait are identified. The modified trait can be any of those traits described above. Additionally, to confirm that the modified trait is due to changes in expression levels or activity of the polypeptide or polynucleotide of the invention can be determined by analyzing mRNA expression using Northern blots, RT-PCR or microarrays, or protein expression using immunoblots or Western blots or gel shift assays.

[0155] Integrated Systems—Sequence Identity

[0156] Additionally, the present invention may be an integrated system, computer or computer readable medium that comprises an instruction set for determining the identity of one or more sequences in a database. In addition, the instruction set can be used to generate or identify sequences that meet any specified criteria. Furthermore, the instruction set may be used to associate or link certain functional benefits, such improved characteristics, with one or more identified sequence.

[0157] For example, the instruction set can include, e.g., a sequence comparison or other alignment program, e.g., an available program such as, for example, the Wisconsin Package Version 10.0, such as BLAST, FASTA, PILEUP, FINDPATTERNS or the like (GCG, Madison, Wis.). Public sequence databases such as GenBank, EMBL, Swiss-Prot and PIR or private sequence databases such as PhytoSeq (Incyte Pharmaceuticals, Palo Alto, Calif.) can be searched.

[0158] Alignment of sequences for comparison can be conducted by the local homology algorithm of Smith and Waterman (1981) *Adv. Mol. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. U.S.A.* 85: 2444, by computerized implementations of these algorithms. After alignment, sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a comparison window to identify and compare local regions of sequence similarity. The comparison window can be a segment of at least about 20 contiguous positions, usually about 50 to about 200, more usually about 100 to about 150 contiguous positions. A description of the method is provided in Ausubel et al., *supra*.

[0159] A variety of methods for determining sequence relationships can be used, including manual alignment and computer assisted sequence alignment and analysis. This later approach is a preferred approach in the present invention, due to the increased throughput afforded by computer

assisted methods. As noted above, a variety of computer programs for performing sequence alignment are available, or can be produced by one of skill.

[0160] One example algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al. *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available, e.g., through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length *W* in the query sequence, which either match or satisfy some positive-valued threshold score *T* when aligned with a word of the same length in a database sequence. *T* is referred to as the neighborhood word score threshold (Altschul et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters *M* (reward score for a pair of matching residues; always >0) and *N* (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity *X* from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters *W*, *T*, and *X* determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (*W*) of 11, an expectation (*E*) of 10, a cutoff of 100, *M*=5, *N*=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (*W*) of 3, an expectation (*E*) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915). Unless otherwise indicated, "sequence identity" here refers to the % sequence identity generated from a tblastx using the NCBI version of the algorithm at the default settings using gapped alignments with the filter "off" (<http://www.ncbi.nlm.nih.gov/>). Additionally, BLASTX and TBLASTX programs may be used.

[0161] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5878). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (*P(N)*), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence (and, therefore, in this context, homologous) if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, or less than about 0.01, and or even less than about 0.001. An additional example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters.

[0162] The integrated system, or computer typically includes a user input interface allowing a user to selectively view one or more sequence records corresponding to the one or more character strings, as well as an instruction set which aligns the one or more character strings with each other or with an additional character string to identify one or more region of sequence similarity. The system may include a link of one or more character strings with a particular phenotype or gene function. Typically, the system includes a user readable output element which displays an alignment produced by the alignment instruction set.

[0163] The methods of this invention can be implemented in a localized or distributed computing environment. In a distributed environment, the methods may implemented on a single computer comprising multiple processors or on a multiplicity of computers. The computers can be linked, e.g. through a common bus, but more preferably the computer(s) are nodes on a network. The network can be a generalized or a dedicated local or wide-area network and, in certain preferred embodiments, the computers may be components of an intra-net or an internet.

[0164] Thus, the invention provides methods for identifying a sequence similar or homologous to one or more polynucleotides as noted herein, or one or more target polypeptides encoded by the polynucleotides, or otherwise noted herein and may include linking or associating a given plant phenotype or gene function with a sequence. In the methods, a sequence database is provided (locally or across an inter or intra net) and a query is made against the sequence database using the relevant sequences herein and associated plant phenotypes or gene functions.

[0165] Any sequence herein can be entered into the database, before or after querying the database. This provides for both expansion of the database and, if done before the querying step, for insertion of control sequences into the database. The control sequences can be detected by the query to ensure the general integrity of both the database and the query. As noted, the query can be performed using a web browser based interface. For example, the database can be a centralized public database such as those noted herein, and the querying can be done from a remote terminal or computer across an internet or intranet.

EXAMPLES

[0166] The following examples are intended to illustrate but not limit the present invention.

Example I

Full Length Gene Identification and Cloning

[0167] Putative transcription factor sequences (genomic or ESTs) related to known transcription factors were identified in the *Arabidopsis thaliana* GenBank database using the tblastn sequence analysis program using default parameters and a P-value cutoff threshold of -4 or -5 or lower, depending on the length of the query sequence. Putative transcription factor sequence hits were then screened to identify those containing particular sequence strings. If the sequence hits contained such sequence strings, the sequences were confirmed as transcription factors.

[0168] Alternatively, *Arabidopsis thaliana* cDNA libraries derived from different tissues or treatments, or genomic

libraries were screened to identify novel members of a transcription family using a low stringency hybridization approach. Probes were synthesized using gene specific primers in a standard PCR reaction (annealing temperature 60° C.) and labeled with ³²P dCTP using the High Prime DNA Labeling Kit (Boehringer Mannheim). Purified radiolabelled probes were added to filters immersed in Church hybridization medium (0.5 M NaPO₄ pH 7.0, 7% SDS, 1% w/v bovine serum albumin) and hybridized overnight at 60° C. with shaking. Filters were washed two times for 45 to 60 minutes with 1×SSC, 1% SDS at 60° C.

[0169] To identify additional sequence 5' or 3' of a partial cDNA sequence in a cDNA library, 5' and 3' rapid amplification of cDNA ends (RACE) was performed using the Marathon™ cDNA amplification kit (Clontech, Palo Alto, Calif.). Generally, the method entailed first isolating poly(A) mRNA, performing first and second strand cDNA synthesis to generate double stranded cDNA, blunting cDNA ends, followed by ligation of the Marathon™ Adaptor to the cDNA to form a library of adaptor-ligated ds cDNA.

[0170] Gene-specific primers were designed to be used along with adaptor specific primers for both 5' and 3' RACE reactions. Nested primers, rather than single primers, were used to increase PCR specificity. Using 5' and 3' RACE reactions, 5' and 3' RACE fragments were obtained, sequenced and cloned. The process can be repeated until 5' and 3' ends of the full-length gene were identified. Then the full-length cDNA was generated by PCR using primers specific to 5' and 3' ends of the gene by end-to-end PCR.

Example II

Construction of Expression Vectors

[0171] The sequence was amplified from a genomic or cDNA library using primers specific to sequences upstream and downstream of the coding region. The expression vector was pMEN20 or pMEN65, which are both derived from pMON316 (Sanders et al, (1987) *Nucleic Acids Research* 15:1543-58) and contain the CaMV 35S promoter to express transgenes. To clone the sequence into the vector, both pMEN20 and the amplified DNA fragment were digested separately with SalI and NotI restriction enzymes at 37° C. for 2 hours. The digestion products were subject to electrophoresis in a 0.8% agarose gel and visualized by ethidium bromide staining. The DNA fragments containing the sequence and the linearized plasmid were excised and purified by using a Qiaquick gel extraction kit (Qiagen, Calif.). The fragments of interest were ligated at a ratio of 3:1 (vector to insert). Ligation reactions using T4 DNA ligase (New England Biolabs, MA) were carried out at 16° C. for 16 hours. The ligated DNAs were transformed into competent cells of the *E. coli* strain DH5alpha by using the heat shock method. The transformations were plated on LB plates containing 50 mg/l kanamycin (Sigma, Mo.).

[0172] Individual colonies were grown overnight in five milliliters of LB broth containing 50 mg/l kanamycin at 37° C. Plasmid DNA was purified by using Qiaquick Mini Prep kits (Qiagen, Calif.).

Example III

Transformation of *Agrobacterium* with the Expression Vector

[0173] After the plasmid vector containing the gene was constructed, the vector was used to transform *Agrobacte-*

rium tumefaciens cells expressing the gene products. The stock of *Agrobacterium tumefaciens* cells for transformation were made as described by Nagel et al. (1990) *FEMS Microbiol Letts.* 67: 325-328. *Agrobacterium* strain ABI was grown in 250 ml LB medium (Sigma) overnight at 28° C. with shaking until an absorbance (A_{600}) of 0.5-1.0 was reached. Cells were harvested by centrifugation at 4,000×g for 15 min at 4° C. Cells were then resuspended in 250 μ l chilled buffer (1 mM HEPES, pH adjusted to 7.0 with KOH). Cells were centrifuged again as described above and resuspended in 125 μ l chilled buffer. Cells were then centrifuged and resuspended two more times in the same HEPES buffer as described above at a volume of 100 μ l and 750 μ l, respectively. Resuspended cells were then distributed into 40 μ l aliquots, quickly frozen in liquid nitrogen, and stored at -80° C.

[0174] *Agrobacterium* cells were transformed with plasmids prepared as described above following the protocol described by Nagel et al. For each DNA construct to be transformed, 50-100 ng DNA (generally resuspended in 10 mM Tris-HCl, 1 mM EDTA, pH 8.0) was mixed with 40 μ l of *Agrobacterium* cells. The DNA/cell mixture was then transferred to a chilled cuvette with a 2 mm electrode gap and subject to a 2.5 kV charge dissipated at 25 μ F and 200 μ F using a Gene Pulser II apparatus (Bio-Rad). After electroporation, cells were immediately resuspended in 1.0 ml LB and allowed to recover without antibiotic selection for 2-4 hours at 28° C. in a shaking incubator. After recovery, cells were plated onto selective medium of LB broth containing 100 μ g/ml spectinomycin (Sigma) and incubated for 24-48 hours at 28° C. Single colonies were then picked and inoculated in fresh medium. The presence of the plasmid construct was verified by PCR amplification and sequence analysis.

Example IV

Transformation of *Arabidopsis* Plants With *Agrobacterium tumefaciens* With Expression Vector

[0175] After transformation of *Agrobacterium tumefaciens* with plasmid vectors containing the gene, single *Agrobacterium* colonies were identified, propagated, and used to transform *Arabidopsis* plants. Briefly, 500 ml cultures of LB medium containing 50 mg/l kanamycin were inoculated with the colonies and grown at 28° C. with shaking for 2 days until an absorbance (A_{600}) of >2.0 is reached. Cells were then harvested by centrifugation at 4,000×g for 10 min, and resuspended in infiltration medium ($\frac{1}{2}$ ×Murashige and Skoog salts (Sigma), 1×Gamborg's B-5 vitamins (Sigma), 5.0% (w/v) sucrose (Sigma), 0.044 μ M benzylamino purine (Sigma), 200 μ l/l Silwet L-77 (Lehle Seeds) until an absorbance (A_{600}) of 0.8 was reached.

[0176] Prior to transformation, *Arabidopsis thaliana* seeds (ecotype Columbia) were sown at a density of ~10 plants per 4" pot onto Pro-Mix BX potting medium (Hummert International) covered with fiberglass mesh (18 mm×16 mm). Plants were grown under continuous illumination (50-75 μ E/m²/sec) at 22-23° C. with 65-70% relative humidity. After about 4 weeks, primary inflorescence stems (bolts) are cut off to encourage growth of multiple secondary bolts. After flowering of the mature secondary bolts, plants were prepared for transformation by removal of all siliques and opened flowers.

[0177] The pots were then immersed upside down in the mixture of *Agrobacterium* infiltration medium as described above for 30 sec, and placed on their sides to allow draining into a 1'x2' flat surface covered with plastic wrap. After 24 h, the plastic wrap was removed and pots are turned upright. The immersion procedure was repeated one week later, for a total of two immersions per pot. Seeds were then collected from each transformation pot and analyzed following the protocol described below.

Example V

Identification of Arabidopsis Primary Transformants

[0178] Seeds collected from the transformation pots were sterilized essentially as follows. Seeds were dispersed into in a solution containing 0.1% (v/v) Triton X-100 (Sigma) and sterile H₂O and washed by shaking the suspension for 20 min. The wash solution was then drained and replaced with fresh wash solution to wash the seeds for 20 min with shaking. After removal of the second wash solution, a solution containing 0.1% (v/v) Triton X-100 and 70% ethanol (Equistar) was added to the seeds and the suspension was shaken for 5 min. After removal of the ethanol/detergent solution, a solution containing 0.1% (v/v) Triton X-100 and 30% (v/v) bleach (Clorox) was added to the seeds, and the suspension was shaken for 10 min. After removal of the bleach/detergent solution, seeds were then washed five times in sterile distilled H₂O. The seeds were stored in the last wash water at 4° C. for 2 days in the dark before being plated onto antibiotic selection medium (1xMurashige and Skoog salts (pH adjusted to 5.7 with 1M KOH), 1xGamborg's B-5 vitamins, 0.9% phytagar (Life Technologies), and 50 mg/l kanamycin). Seeds were germinated under continuous illumination (50-75 $\mu\text{E}/\text{m}^2/\text{sec}$) at 22-23° C. After 7-10 days of growth under these conditions, kanamycin resistant primary transformants (T₁ generation) were visible and obtained. These seedlings were transferred first to fresh selection plates where the seedlings continued to grow for 3-5 more days, and then to soil (Pro-Mix BX potting medium).

[0179] Primary transformants were crossed and progeny seeds (T₂) collected; kanamycin resistant seedlings were selected and analyzed. The expression levels of the recombinant polynucleotides in the transformants varies from about a 5% expression level increase to a least a 100% expression level increase. Similar observations are made with respect to polypeptide level expression.

Example VI

Identification of Arabidopsis Plants with Transcription Factor Gene Knockouts

[0180] The screening of insertion mutagenized Arabidopsis collections for null mutants in a known target gene was essentially as described in Krysan et al (1999) *Plant Cell* 11:2283-2290. Briefly, gene-specific primers, nested by 5-250 base pairs to each other, were designed from the 5' and 3' regions of a known target gene. Similarly, nested sets of primers were also created specific to each of the T-DNA or transposon ends (the "right" and "left" borders). All possible combinations of gene specific and T-DNA/transposon primers were used to detect by PCR an insertion event within or close to the target gene. The amplified DNA fragments were then sequenced which allows the precise determination of

the T-DNA/transposon insertion point relative to the target gene. Insertion events within the coding or intervening sequence of the genes were deconvoluted from a pool comprising a plurality of insertion events to a single unique mutant plant for functional characterization. The method is described in more detail in Yu and Adam, U.S. application Ser. No. 09/177,733 filed Oct. 23, 1998.

Example VII

Identification of Modified Phenotypes in Overexpression or Gene Knockout Plants

[0181] Experiments were performed to identify those transformants or knockouts that exhibited modified biochemical characteristics. Among the biochemicals that were assayed were insoluble sugars, such as arabinose, fucose, galactose, mannose, rhamnose or xylose or the like; prenol lipids, such as lutein, beta-carotene, xanthophyll-1, xanthophyll-2, chlorophylls A or B, or alpha-, delta- or gamma-tocopherol or the like; fatty acids, such as 16:0 (palmitic acid), 16:1 (palmitoleic acid), 18:0 (stearic acid), 18:1 (oleic acid), 18:2 (linoleic acid), 20:0, 18:3 (linolenic acid), 20:1 (eicosenoic acid), 20:2, 22:1 (erucic acid) or the like; waxes, such as by altering the levels of C29, C31, or C33 alkanes; sterols, such as brassicasterol, campesterol, stigmastanol, sitosterol or stigmastanol or the like, glucosinolates, protein or oil levels.

[0182] Fatty acids were measured using two methods depending on whether the tissue was from leaves or seeds. For leaves, lipids were extracted and esterified with hot methanolic H₂SO₄ and partitioned into hexane from methanolic brine. For seed fatty acids, seeds were pulverized and extracted in methanol:heptane:toluene:2,2-dimethoxypropane:H₂SO₄ (39:34:20:5:2) for 90 minutes at 80° C. After cooling to room temperature the upper phase, containing the seed fatty acid esters, was subjected to GC analysis. Fatty acid esters from both seed and leaf tissues were analyzed with a Supelco SP-2330 column.

[0183] Glucosinolates were purified from seeds or leaves by first heating the tissue at 95° C. for 10 minutes. Preheated ethanol:water (50:50) is and after heating at 95° C. for a further 10 minutes, the extraction solvent is applied to a DEAE Sephadex column which had been previously equilibrated with 0.5 M pyridine acetate. Desulfoglucosinolates were eluted with 300 μl water and analyzed by reverse phase HPLC monitoring at 226 nm.

[0184] For wax alkanes, samples were extracted using an identical method as fatty acids and extracts were analyzed on a HP 5890 GC coupled with a 5973 MSD. Samples were chromatographed on a J&W DB35 mass spectrometer (J&W Scientific).

[0185] To measure prenol lipids levels, seeds or leaves were pulverized with 1 to 2% pyrogallol as an antioxidant. For seeds, extracted samples were filtered and a portion removed for tocopherol and carotenoid/chlorophyll analysis by HPLC. The remaining material was saponified for sterol determination. For leaves, an aliquot was removed and diluted with methanol and chlorophyll A, chlorophyll B, and total carotenoids measured by spectrophotometry by determining absorbance at 665.2 nm, 652.5 nm, and 470 nm. An aliquot was removed for tocopherol and carotenoid/chlorophyll composition by HPLC using a Waters uBondapak C18

column (4.6 mm×150 mm). The remaining methanolic solution was saponified with 10% KOH at 80° C. for one hour. The samples were cooled and diluted with a mixture of methanol and water. A solution of 2% methylene chloride in hexane was mixed in and the samples were centrifuged. The aqueous methanol phase was again re-extracted 2% methylene chloride in hexane and, after centrifugation, the two upper phases were combined and evaporated. 2% methylene chloride in hexane was added to the tubes and the samples were then extracted with one ml of water. The upper phase was removed, dried, and resuspended in 400 µl of 2% methylene chloride in hexane and analyzed by gas chromatography using a 50 m DB-5 ms (0.25 mm ID, 0.25 µm phase, J&W Scientific).

[0186] Insoluble sugar levels were measured by the method essentially described by Reiter et al., *Plant Journal* 12:335-345. This method analyzes the neutral sugar composition of cell wall polymers found in *Arabidopsis* leaves. Soluble sugars were separated from sugar polymers by extracting leaves with hot 70% ethanol. The remaining residue containing the insoluble polysaccharides was then acid hydrolyzed with allose added as an internal standard. Sugar monomers generated by the hydrolysis were then reduced to the corresponding alditols by treatment with NaBH₄, then were acetylated to generate the volatile alditol acetates which were then analyzed by GC-FID. Identity of the peaks was determined by comparing the retention times of known sugars converted to the corresponding alditol acetates with the retention times of peaks from wild-type plant extracts. Alditol acetates were analyzed on a Supelco SP-2330 capillary column (30 m×250 µm×0.2 µm) using a temperature program beginning at 180° C. for 2 minutes followed by an increase to 220° C. in 4 minutes. After holding at 220° C. for 10 minutes, the oven temperature is increased to 240° C. in 2 minutes and held at this temperature for 10 minutes and brought back to room temperature.

[0187] To identify plants with alterations in total seed oil or protein content, 150 mg of seeds from T2 progeny plants were subjected to analysis by Near Infrared Reflectance (NIR) using a Foss NirSystems Model 6500 with a spinning cup transport system.

[0188] Experiments were performed to identify those transformants or knockouts that exhibited an improved pathogen tolerance. For such studies, the transformants were exposed to biotrophic fungal pathogens, such as *Erysiphe orontii*, and necrotrophic flingal pathogens, such as *Fusarium oxysporum*. *Fusarium oxysporum* isolates cause vascular wilts and damping off of various annual vegetables, perennials and weeds (Mauch-Mani and Slusarenko (1994) *Molecular Plant-Microbe Interactions* 7: 378-383). For *Fusarium oxysporum* experiments, plants grown on petri dishes were sprayed with a fresh spore suspension of *F. oxysporum*. The spore suspension was prepared as follows: A plug of fungal hyphae from a plate culture was placed on a fresh potato dextrose agar plate and allowed to spread for one week. 5 ml sterile water was then added to the plate, swirled, and pipetted into 50 ml Armstrong Fusarium medium. Spores were grown overnight in Fusarium medium and then sprayed onto plants using a Preval paint sprayer. Plant tissue was harvested and frozen in liquid nitrogen 48 hours post infection.

[0189] *Erysiphe orontii* is a causal agent of powdery mildew. For *Erysiphe orontii* experiments, plants were

grown approximately 4 weeks in a greenhouse under 12 hour light (20° C., ~30% relative humidity (rh)). Individual leaves were infected with *E. orontii* spores from infected plants using a camel's hair brush, and the plants were transferred to a Percival growth chamber (20° C., 80% rh.). Plant tissue was harvested and frozen in liquid nitrogen 7 days post infection.

[0190] *Botrytis cinerea* is a necrotrophic pathogen. *Botrytis cinerea* was grown on potato dextrose agar in the light. Aspore culture was made by spreading 10 ml of sterile water on the ftingus plate, swirling and transferring spores to 10 ml of sterile water. The spore inoculum (approx. 105 spores/ml) was used to spray 10 day-old seedlings grown under sterile conditions on MS (minus sucrose) media. Symptoms were evaluated every day up to approximately 1 week.

[0191] Infection with bacterial pathogens *Pseudomonas syringae* pv maculicola strain 4326 and pv maculicola strain 4326 was performed by hand inoculation at two doses. Two inoculation doses allows the differentiation between plants with enhanced susceptibility and plants with enhanced resistance to the pathogen. Plants were grown for 3 weeks in the greenhouse, then transferred to the growth chamber for the remainder of their growth. Psm ES4326 was hand inoculated with 1 ml syringe on 3 fully-expanded leaves per plant (4½ wk old), using at least 9 plants per overexpressing line at two inoculation doses, OD=0.005 and OD=0.0005. Disease scoring occurred at day 3 post-inoculation with pictures of the plants and leaves taken in parallel.

[0192] In some instances, expression patterns of the pathogen-induced genes (such as defense genes) was monitored by microarray experiments. cDNAs were generated by PCR and resuspended at a final concentration of ~100 ng/µl in 3×SSC or 150 mM Na-phosphate (Eisen and Brown (1999) *Meth. in Enzymol.* 303:179-205). The cDNAs were spotted on microscope glass slides coated with polylysine. The prepared cDNAs were aliquoted into 384 well plates and spotted on the slides using an x-y-z gantry (OnmiGrid) purchased from GeneMachines (Menlo Park, Calif.) outfitted with quill type pins purchased from Telechem International (Sunnyvale, Calif.). After spotting, the arrays were cured for a minimum of one week at room temperature, rehydrated and blocked following the protocol recommended by Eisen and Brown (1999). [Sample total RNA (10 µg) samples were labeled using fluorescent Cy3 and Cy5 dyes. Labeled samples were resuspended in 4×SSC/0.03% SDS/4 µg salmon sperm DNA/2 µg tRNA/50 mM Na-pyrophosphate, heated for 95° C. for 2.5 minutes, spun down and placed on the array. The array was then covered with a glass coverslip and placed in a sealed chamber. The chamber was then kept in a water bath at 62° C. overnight. The arrays were washed as described in Eisen and Brown (1999) and scanned on a General Scanning 3000 laser scanner. The resulting files are subsequently quantified using Imagen software purchased from BioDiscovery (Los Angeles, Calif.).

[0193] Experiments were performed to identify those transformants or knockouts that exhibited an improved environmental stress tolerance. For such studies, the transformants were exposed to a variety of environmental stresses. Plants were exposed to chilling stress (6 hour exposure to 4-8° C.), heat stress (6 hour exposure to 32-37° C.), high salt stress (6 hour exposure to 200 mM NaCl),

drought stress (168 hours after removing water from trays), osmotic stress (6 hour exposure to 3 M mannitol), or nutrient limitation (nitrogen, phosphate, and potassium) (Nitrogen: all components of MS medium remained constant except N was reduced to 20 mg/l of NH₄ NO₃, or Phosphate: All components of MS medium except KH₂ PO₄, which was replaced by K₂SO₄, Potassium: All components of MS medium except removal of KNO₃ and KH₂PO₄, which were replaced by NaH₄PO₄).

[0194] Experiments were performed to identify those transformants or knockouts that exhibited a modified structure and development characteristics. For such studies, the transformants were observed by eye to identify novel structural or developmental characteristics associated with the ectopic expression of the polynucleotides or polypeptides of the invention.

[0195] Experiments were performed to identify those transformants or knockouts that exhibited modified sugar-sensing. For such studies, seeds from transformants were germinated on media containing 5% glucose or 9.4%

polynucleotide and polypeptide sequences of the invention. Table 4 includes, from left to right for each sequence: the first column shows the polynucleotide SEQ ID NO; the second column shows the polypeptide SEQ ID NO encoded by the polynucleotide; the third column shows the Mendel Gene ID No., GID; the fourth column, CDS, shows the start and stop nucleotide positions of the encoded polypeptide, respectively, with respect to the polynucleotide co-ordinates; the fifth column shows the amino acid residue positions of the conserved domain in amino acid (AA) co-ordinates; and the sixth column shows if the polynucleotide was knocked out (KO) or overexpressed (OE) in a transgenic plant. Table 5 includes, from left to right for each sequence: the first column shows the nucleotide SEQ ID NO.; the second column shows the polypeptide SEQ ID NO encoded by the polynucleotide; the third column shows the Mendel Gene ID No., GID; the fourth column shows if the polynucleotide was knocked out (KO) or overexpressed (OE) in a transgenic plant; and the fifth column shows the trait resulting from the knock out or overexpression of the polynucleotide in the transgenic plant.

TABLE 4

SEQ ID NO (polynucleotide)	SEQ ID NO (polypeptide)	GID	CDS	Conserved domain in AA coordinates	Knockout or overexpressor
1	2	G481	103 . . . 528	20–109	OE
3	4	G1466	16 . . . 1278	154–420	OE

sucrose which normally partially restrict hypocotyl elongation. Plants with altered sugar sensing may have either longer or shorter hypocotyls than normal plants when grown on this media. Additionally, other plant traits may be varied such as root mass.

[0196] Flowering time was measured by the number of rosette leaves present when a visible inflorescence of approximately 3 cm is apparent Rosette and total leaf number on the progeny stem are tightly correlated with the timing of flowering (Koomneef et al (1991) *Mol. Gen. Genet* 229:57-66. The vernalization response was measured. For vernalization treatments, seeds were sown to MS agar plates, sealed with micropore tape, and placed in a 4° C. cold room with low light levels for 6-8 weeks. The plates were then transferred to the growth rooms alongside plates containing freshly sown non-vernalized controls. Rosette leaves were counted when a visible inflorescence of approximately 3 cm was apparent.

[0197] Modified phenotypes observed for particular over-expressor or knockout plants are provided in Table 4 and Table 5. For a particular overexpressor that shows a less beneficial characteristic, it may be more useful to select a plant with a decreased expression of the particular transcription factor. For a particular knockout that shows a less beneficial characteristic, it may be more useful to select a plant with an increased expression of the particular transcription factor.

[0198] The sequences of the Sequence Listing or those discloses here can be used to prepare transgenic plants and plants with altered traits. The specific transgenic plants listed below are produced from the sequences of the Sequence Listing, as noted. Table 4 and Table 5 provide exemplary

[0199]

TABLE 5

SEQ ID NO (poly- nucle- otide)	SEQ ID NO (poly- peptide) GID	Observed phenotype or trait	Overexpressor or knockout
1	2	G481	Germination assay: OE
			High sucrose, osmotic stress
3	4	G1466	Seed composition assay: OE
			High seed oil and protein content

[0200] G481: Better Germination on High Sucrose

[0201] Seed of plants overexpressing sequence G481 (SEQ ID NOs: 1 and 2) showed slightly better germination when grown on high sucrose medium (5% glucose or 9.4% sucrose). The plants showed longer radicle and more cotyledon expansion. In one line (line 8) analysis of leaf insoluble sugars showed a slight increase in decrease in Rhamnose levels (16.9% vs. approximately 11.4% in wild-type plants) and a decrease in Arabinose and Xylose levels (15.3% and 12.8% vs. approximately 19.0% and 21% in wild-type plants, respectively).

[0202] In wild-type plants, G481 was predominantly expressed in flower and silique, and to a lesser extent, in rosette, embryo, and germinating seed.

[0203] The potential utility of G481 includes a possible role in sugar sensing, a plant mechanism that has been shown to be involved in the following: 1) altering storage compound accumulation (oil and/or protein) in seeds which could impact yield and seed quality, and 2) altering photo-

synthetic rate which could also impact yield in vegetative tissues as well as seed. Sugars are key regulatory molecules that affect diverse processes in higher plants including germination, growth, flowering, senescence, sugar metabolism and photosynthesis. Sucrose is the major transport form of photosynthate and its flux through cells has been shown to affect gene expression and alter storage compound accumulation in seeds (source-sink relationships).

[0204] The enhanced germination phenotype of transgenic plants overexpressing G481 under a condition of osmotic stress (such as high concentrations of sucrose) suggests the gene could also be used to improve plant tolerance to water deficit related conditions such as drought stress, salt stress, and freezing stress. Thus G481 could be used to engineer plants with enhanced stress tolerance that could ultimately impact survivability and yield.

[0205] G1466: Increased Seed Oil; Decreased Protein Content

[0206] Seed of plants overexpressing sequences G1466 (SEQ ID NOs:3 and 4) was subjected to NIR analysis and an increase in seed oil content compared with seed from wild-type plants was identified (39% vs. approximately 35% in wild-type plants). In addition, a slight decrease in seed protein content compared with seed from wild-type plants was identified (20.8% vs. approximately 23% in wild-type plants).

[0207] In wild-type plants, G1466 was expressed in all tissues examined.

[0208] Therefore, G1466 could be used to modify high value seed quality traits such protein, oil and carbohydrates content and composition in any plant in which the expression of this gene is altered. Altering the amount of seed oil, protein or carbohydrate could effectively increase the yield and quality. Altering the composition of seeds would improve feed quality by altering availability of energy and phosphorus, and to improve the amino acid balance of grain meal or to improve oil quality for human food and industrial uses. Altering seed composition with G1466 could also improved grain quality for the wet milling industry as well as provide a means for the production of novel polymers and chemicals.

Example VIII

Identification of Homologous Sequences

[0209] Homologous sequences from Arabidopsis and plant species other than Arabidopsis were identified using database sequence search tools, such as the Basic Local Alignment Search Tool (BLAST) (Altschul et al. (1990) *J. Mol. Biol.* 215:403-410; and Altschul et al. (1997) *Nucl. Acid Res.* 25: 3389-3402). The tblastx sequence analysis programs were employed using the BLOSUM-62 scoring matrix (Henikoff, S. and Henikoff, J. G. (1992) *Proc. Natl. Acad. Sci.* 89: 10915-10919).

[0210] Identified orthologs and homologs of Arabidopsis sequences are provided in Tables 6 and 7. The percent sequence identity among these sequences can be as low as 47%, or even lower sequence identity. Additionally, the entire NCBI GenBank database was filtered for sequences from all plants except *Arabidopsis thaliana* by selecting all entries in the NCBI GenBank database associated with NCBI taxonomic ID 33090 (Viridiplantae; all plants) and excluding entries associated with taxonomic ID 3701 (*Arabidopsis thaliana*). These sequences are compared to sequences representing genes of SEQ IDs NOs:2 and 4 using the Washington University TBLASTX algorithm (version 2.0a19MP) at the default settings using gapped alignments with the filter "off". For each gene of SEQ IDs NOs:2 and 4 individual comparisons were ordered by probability score (P-value), where the score reflects the probability that a particular alignment occurred by chance. For example, a score of 3.6e-40 is 3.6×10^{-40} . In addition to P-values, comparisons were also scored by percentage identity. Percentage identity reflects the degree to which two segments of DNA or protein are identical over a particular length. The identified homologous polynucleotide and polypeptide sequences and homologues of the Arabidopsis polynucleotides and polypeptides may be orthologs of the Arabidopsis polynucleotides and polypeptides.

[0211] As shown in Table 6, polynucleotide and polypeptide sequences which were identified as orthologous and homologous of SEQ ID NOs: 1 and 2 were found in *Gossypium arboreum*, *Glycine max*, *Zea mays*, *Gossypium hirsutum*, *Medicago truncatula*, *Lycopersicon esculentum*, *Solanum tuberosum*, *Triticum aestivum*, *Hordeum vulgare*, *Triticum monococcum*, *Oryza sativa*, *Vernonia galamensis*, *Argemone mexicana*, and *Triticum aestivum*.

[0212] As shown in Table 6, polynucleotide and polypeptide sequences which were identified as orthologous and homologous of SEQ ID NOs:3 and 4 were found in *Brassica oleracea*, *Lycopersicon esculentum*, *Lycopersicon pennellii*, *Lotus japonicus*, *Oryza sativa*, *Solanum tuberosum*, *Medicago truncatula*, *Glycine max*, *Euphorbia esula*, *Gossypium arboreum*, *Antirrhinum hispanicum*, *Oryza sativa* (japonica cultivar-group), and *Triticum aestivum*. Additional orthologous and homologous polynucleotides and polypeptide sequences from other plant species are shown in Table 7.

[0213] All references, publications, patent documents, web pages, and other documents cited or mentioned herein are hereby incorporated by reference in their entirety for all purposes. Although the invention has been described with reference to specific embodiments and examples, it should be understood that one of ordinary skill can make various modifications without departing from the spirit of the invention. The scope of the invention is not limited to the specific embodiments and examples provided.

TABLE 6

SEQ ID NO	Gene ID	Test Sequence ID	Smallest Sum		Test Sequence Species	Test Sequence GenBank Annotation
			Probability	Test Sequence		
1	G481	BG440251	9.00E-42	[<i>Gossypium arboreum</i>]	GA_Ea0006K20f	<i>Gossypium arboreum</i> 7-10 d
1	G481	BM887558	3.90E-41	[<i>Glycine max</i>]	sam40c09.y1	Gm-c1068 <i>Glycine max</i> cDNA clone SOY

TABLE 6-continued

SEQ ID NO	Gene ID	Test Sequence ID	Smallest Sum Probability	Test Sequence Species	Test Sequence GenBank Annotation
1	G481	ZMNFYB	1.70E-40	[<i>Zea mays</i>]	<i>Z. mays</i> mRNA for CAAT-box DNA binding protein subun
1	G481	AI728916	2.40E-40	[<i>Gossypium hirsutum</i>]	BNLGH12022 Six-day Cotton fiber Gossypi
1	G481	AW775623	3.80E-40	[<i>Medicago truncatula</i>]	EST334688 DSIL <i>Medicago truncatula</i> cDNA
1	G481	AW738727	9.80E-40	[<i>Lycopersicon esculentum</i>]	EST340154 tomato flower buds, anthe
1	G481	BG599785	1.70E-38	[<i>Solanum tuberosum</i>]	EST504680 cSTS <i>Solanum tuberosum</i> cDNA clo
1	G481	BE413647	2.50E-38	[<i>Triticum aestivum</i>]	SCU001.E10.R990714 ITEC SCU Wheat Endospe
1	G481	BF065056	5.80E-38	[<i>Hordeum vulgare</i>]	HV_CE0022M01f <i>Hordeum vulgare</i> seedling gre
1	G481	BG314203	1.00E-37	[<i>Triticum monococcum</i>]	WHE2460_E10_I20ZS <i>Triticum monococcum</i> i
1	G481	gi22380	1.10E-45	[<i>Zea mays</i>]	CAAT-box DNA binding protein subunit B (NF-YB).
1	G481	gi15408794	2.60E-30	[<i>Oryza sativa</i>]	putative CCAAT-binding transcription factor
1	G481	gi16902054	1.00E-28	[<i>Vernonia galamensis</i>]	CCAAT-box binding factor HAP3 B domai
1	G481	gi16902050	2.70E-28	[<i>Glycine max</i>]	CCAAT-box binding factor HAP3 B domain.
1	G481	gi16902056	4.30E-28	[<i>Argemone mexicana</i>]	CCAAT-box binding factor HAP3 B domain.
1	G481	gi16902058	1.50E-23	[<i>Triticum aestivum</i>]	CCAAT-box binding factor HAP3 B domain.
3	G1466	BH596941	1.40E-65	[<i>Brassica oleracea</i>]	BOHFG41TR BOHF <i>Brassica oleracea</i> genomic
3	G1466	BE462774	0.00014	[<i>Lycopersicon esculentum</i>]	EST325108 tomato flower buds 0-3 mm
3	G1466	AW399721	0.00049	[<i>Lycopersicon pennellii</i>]	EST310221 <i>L. pennellii</i> trichome, Cor
3	G1466	AV413010	0.00054	[<i>Lotus japonicus</i>]	AV413010 <i>Lotus japonicus</i> young plants (two-
3	G1466	OSIN00182	0.0029	[<i>Oryza sativa</i>]	chromosome 4 clone
3	G1466	BG599234	0.0047	[<i>Solanum tuberosum</i>]	OSINBa0086006, *** SEQUENC
3	G1466	BG580068	0.0095	[<i>Medicago truncatula</i>]	EST504134 cSTS <i>Solanum tuberosum</i> cDNA clo
3	G1466	BI316339	0.026	[<i>Glycine max</i>]	EST481790 GVN <i>Medicago truncatula</i> cDNA
3	G1466	BG409415	0.026	[<i>Euphorbia esula</i>]	saf02a08.y1 Gm-c1065 <i>Glycine max</i> cDNA
3	G1466	BF278686	0.11	[<i>Gossypium arboreum</i>]	clone GEN
3	G1466	gi8096405	2.30E-12	[<i>Oryza sativa</i>]	00787 leafy spurge Lambda HybriZAP 2.1 two-
3	G1466	gi13161526	3.40E-08	[<i>Antirrhinum hispanicum</i>]	GA_Eb0035E21f <i>Gossypium arboreum</i> 7-10 d
3	G1466	gi18844804	0.083	[<i>Oryza sativa</i> (japonica cultivar-group)]	Similar to <i>Arabidopsis thaliana</i> chromosome 1
3	G1466	gi9858770	0.83	[<i>Lycopersicon esculentum</i>]	S locus F-box (SLF)-S2 protein.
3	G1466	gi6505722	1	[<i>Triticum aestivum</i>]	contains ESTs D151
					BAC19.2.
					ribosomal RNA apurinic site specifcly

[0214]

TABLE 7

SEQ ID NO	Gene ID	Test Sequence ID	Test Sequence GenBank Annotation	Reading Frame	High Score	Smallest Sum Probability	N
1	G481	BG440251	BG440251 GA_Ea0006K20f <i>Gossypium arboreum</i> 7 . . .	3	263	9.00E-42	2
1	G481	BM887558	BM887558 sam40c09.y1 Gm-c1068 <i>Glycine max</i> cD . . .	3	268	3.90E-41	2
1	G481	BG362898	BG362898 sac13e07.y1 Gm-c1040 <i>Glycine max</i> cD . . .	3	263	4.90E-41	2
1	G481	AW395227	AW395227 sh45e04.y1 Gm-c1017 <i>Glycine max</i> cDN . . .	1	264	9.10E-41	2
1	G481	BM525962	BM525962 sak74b11.y1 Gm-c1036 <i>Glycine max</i> cD . . .	3	264	9.20E-41	2
1	G481	BI972318	BI972318 sag90a01.y1 Gm-c1084 <i>Glycine max</i> cD . . .	3	264	9.70E-41	2
1	G481	BG363233	BG363233 sac11h11.y1 Gm-c1040 <i>Glycine max</i> cD . . .	3	264	9.90E-41	2
1	G481	BE021941	BE021941 sm64d05.y1 Gm-c1028 <i>Glycine max</i> cDN . . .	1	264	1.00E-40	2
1	G481	BI316766	BI316766 saf73a12.y1 Gm-c1078 <i>Glycine max</i> cD . . .	3	264	1.10E-40	2
1	G481	BE803572	BE803572 sr60e11.y1 Gm-c1052 <i>Glycine max</i> cDN . . .	2	264	1.10E-40	2
1	G481	ZMNFYB	X59714 <i>Z. mays</i> mRNA for CAAT-box DNA binding . . .	2	262	1.70E-40	2
1	G481	BF071234	BF071234 st06h05.y1 Gm-c1065 <i>Glycine max</i> cDN . . .	3	258	2.00E-40	2
1	G481	AI728916	AI728916 BNLGH12022 Six-day Cotton fiber Go . . .	3	261	2.40E-40	2
1	G481	BF597252	BF597252 su96c06.y1 Gm-c1056 <i>Glycine max</i> cDN . . .	2	264	2.90E-40	2
1	G481	AW597630	AW597630 sj96g06.y1 Gm-c1023 <i>Glycine max</i> cDN . . .	2	259	3.40E-40	2
1	G481	AW775623	AW775623 EST334688 DSIL <i>Medicago truncatula</i> . . .	2	259	3.80E-40	2
1	G481	AW733618	AW733618 sk75h06.y1 Gm-c1016 <i>Glycine max</i> cDN . . .	3	257	6.00E-40	2
1	G481	AW738727	AW738727 EST340154 tomato flower buds, anthe . . .	1	249	9.80E-40	2
1	G481	BG642751	BG642751 EST510945 tomato shoot/meristem Lyc . . .	1	249	9.90E-40	2
1	G481	BE441135	BE441135 EST408405 tomato developing/immatur . . .	1	249	9.90E-40	2
1	G481	AW621652	AW621652 EST312450 tomato root during/after . . .	1	249	1.10E-39	2
1	G481	AI900024	AI900024 sb97g11.y1 Gm-c1012 <i>Glycine max</i> cDN . . .	1	264	3.20E-39	2
1	G481	BG445358	BG445358 GA_Ea0027N18f <i>Gossypium arboreum</i> 7 . . .	2	243	1.40E-38	2
1	G481	BG599785	BG599785 EST504680 cSTS <i>Solanum tuberosum</i> cD . . .	2	254	1.70E-38	2
1	G481	BG350430	BG350430 091D09 Mature tuber lambda ZAP Sola . . .	3	254	1.70E-38	2
1	G481	BE413647	BE413647 SCU001.E10.R990714 ITEC SCU Wheat E . . .	3	248	2.50E-38	2
1	G481	BE516510	BE516510 WHE611_D10_H19ZA Wheat ABA-treated . . .	2	248	3.00E-38	2
1	G481	BF065056	BF065056 HV_CE0022M01f <i>Hordeum vulgare</i> seed . . .	3	247	5.80E-38	2

TABLE 7-continued

SEQ ID	Test	Reading	High	Smallest Sum	
NO	Gene ID	Sequence ID	Test Sequence	GenBank	Annotation
1	G481	BG314203	BG314203 WHE2460_E10_I20ZS	Triticum monococc	...
1	G481	AI725612	AI725612 BNLGHI12445 Six-day	Cotton fiber Go	...
1	G481	AL387357	AL387357 MtBC42A04F1 MtBC	Medicago truncatul	...
1	G481	AW907348	AW907348 EST343471 potato stolon,	Cornell Un	...
1	G481	BG274786	BG274786 WHE2234_C03_E06ZS	Aegilops speltoid	...
1	G481	AW459387	AW459387 sh23f03.y1 Gm-c1016	Glycine max cDN	...
1	G481	BE804236	BE804236 sr77b04.y1 Gm-c1052	Glycine max cDN	...
1	G481	BE210041	BE210041 so38b01.y1 Gm-c1039	Glycine max cDN	...
1	G481	AW980494	AW980494 EST391647 GVN	Medicago truncatula c	...
1	G481	BG263362	BG263362 WHE2341_B02_C03ZS	Wheat pre-anthesi	...
1	G481	BH532457	BH532457 BOGLV08TR BOGL	Brassica oleracea ge	...
1	G481	BG847452	BG847452 1024017D03.y1 C.	reinhardtii CC-169	...
1	G481	BG857007	BG857007 1024049D01.y1 C.	reinhardtii CC-169	...
1	G481	BG858372	BG858372 1024057C11.y1 C.	reinhardtii CC-169	...
1	G481	BG850689	BG850689 1024029A11.y2 C.	reinhardtii CC-169	...
1	G481	BI718232	BI718232 1031024F10.y1 C.	reinhardtii CC-169	...
1	G481	BI719728	BI719728 1031045D08.y1 C.	reinhardtii CC-169	...
1	G481	BI875221	BI875221 963122G10.y1 C.	reinhardtii CC-1690	...
1	G481	BE496857	BE496857 WHE0761_D09_H17ZS	Wheat heat-stress	...
1	G481	BF651151	BF651151 NF101H10EC1F1090	Elicited cell cult	...
1	G481	BE441739	BE441739 925009A11.x1 C.	reinhardtii CC-2290	...
1	G481	BG846124	BG846124 1024012C11.y1 C.	reinhardtii CC-169	...
1	G481	AX288144	AX288144 Sequence 15 from Patent	WO0177311	...
1	G481	AW570530	AW570530 sj63c01.y1 Gm-c1033	Glycine max cDN	...
1	G481	AX180950	AX180950 Sequence 1 from Patent	WO0145493, 8	...
1	G481	BF585526	BF585526 FM1_23_E09.g1_A003	Floral-Induced M	...
1	G481	BI271802	BI271802 NF013D06FL1F1057	Developing flower	...
1	G481	BF270944	BF270944 GA_Eb0010B11f	Gossypium arboreum 7	...
1	G481	AI731275	AI731275 BNLGHI9078 Six-day	Cotton fiber Gos	...
1	G481	BI967397	BI967397 GM830001B20E03 Gm-r1083	Glycine max	...
1	G481	BF585616	BF585616 FM1_23_E09.b1_A003	Floral-Induced M	...
1	G481	BF263449	BF263449 HV_CeA0006M10f	Hordeum vulgare seed	...
1	G481	BF263455	BF263455 HV_CeA0006M16f	Hordeum vulgare seed	...
1	G481	BF459554	BF459554 061A04 Mature tuber	lambda ZAP Sola	...
1	G481	BF460267	BF460267 073E08 Mature tuber	lambda ZAP Sola	...
1	G481	BG594268	BG594268 EST492946 cSTS	Solanum tuberosum cD	...
1	G481	BI469382	BI469382 sai11b10.y1 Gm-c1053	Glycine max cD	...
1	G481	AI731250	AI731250 BNLGHI9010 Six-day	Cotton fiber Gos	...
1	G481	BG850688	BG850688 1024029A11.y1 C.	reinhardtii CC-169	...
1	G481	AW132359	AW132359 se03b02.y1 Gm-c1013	Glycine max cDN	...
1	G481	BM269434	BM269434 MEST409-G11.univ	ISUM5-RN Zea mays	...
1	G481	AW035570	AW035570 EST281308 tomato	callus, TAMU Lycop	...
1	G481	BE418716	BE418716 SCL074.B01R990724	ITEC SCL Wheat Le	...
1	G481	AW648378	AW648378 EST326832 tomato	germinating seedli	...
1	G481	AV424305	AV424305 AV424305 Lotus	japonicus young plan	...
1	G481	BF715909	BF715909 saa11e08.y1 Gm-c1058	Glycine max cD	...
1	G481	BI423967	BI423967 sah64c11.y1 Gm-c1049	Glycine max cD	...
1	G481	BG890447	BG890447 EST516298 cSTD	Solanum tuberosum cD	...
1	G481	AI495007	AI495007 sa89f03.y1 Gm-c1004	Glycine max cDN	...
1	G481	AW760103	AW760103 sl58b03.y1 Gm-c1027	Glycine max cDN	...
1	G481	BF517889	BF517889 NXSI_029_D01_F	NXSI (Nsf Xylem Side	...
1	G481	BE060015	BE060015 sn39h06.y1 Gm-c1027	Glycine max cDN	...
1	G481	AW625817	AW625817 EST319724 tomato	radicle, 5 d post-	...
1	G481	AW397727	AW397727 sg83f04.y1 Gm-c1026	Glycine max cDN	...
1	G481	BI207873	BI207873 EST525913 cTOS	Lycopersicon esculen	...
1	G481	AV632945	AV632945 AV632945 Chlamydomonas	reinhardtii	...
1	G481	AW931634	AW931634 EST357477 tomato	fruit mature green	...
1	G481	BM888735	BM888735 952068E04.y1 952	—BMS tissue from	...
1	G481	AI486503	AI486503 EST244824 tomato	ovary, TAMU Lycope	...
1	G481	BE641101	BE641101 Cri2_2_E11_SP6	Ceratopteris Spore L	...
1	G481	BI953657	BI953657 HVSME0013M03f	Hordeum vulgare gree	...
1	G481	BM341536	BM341536 MEST336-C11.T3	ISUM5-RN Zea mays cD	...
1	G481	AX180957	AX180957 Sequence 8 from Patent	WO0145493, 8	...
1	G481	AW201996	AW201996 sf09g11.y1 Gm-c1027	Glycine max cDN	...
1	G481	BG135204	BG135204 EST468096 tomato	crown gall Lycoper	...
1	G481	BI406257	BI406257 158C12 Mature tuber	lambda ZAP Sola	...
1	G481	BM341107	BM341107 MEST330-D11.T3	ISUM5-RN Zea mays cD	...
1	G481	AI782351	AI782351 EST263230 tomato	susceptible, Corne	...
1	G481	BM331836	BM331836 MEST171-B11.T3	ISUM5-RN Zea mays cD	...
1	G481	BM268414	BM268414 MEST395-C12.univ	ISUM5-RN Zea mays	...
1	G481	BM337630	BM337630 MEST215-B12.T3	ISUM5-RN Zea mays cD	...
1	G481	BM349646	BM349646 MEST253-D11.T3	ISUM5-RN Zea mays cD	...
1	G481	BE356560	BE356560 DG1_126_D05.b1_A002	Dark Grown 1 (D	...

TABLE 7-continued

SEQ ID	Test	Reading	High	Smallest Sum	
NO	Gene ID	Frame	Score	Probability	N
	Sequence ID	Test Sequence	GenBank	Annotation	
1	G481	AV632044	AV632044	AV632044	<i>Chlamydomonas reinhardtii</i> . . .
1	G481	BE054369	BE054369	GA_Ea0002A05f	<i>Gossypium arboreum</i> 7 . . .
1	G481	AC108500	AC108500	<i>Oryza sativa</i> chromosome 5 clone OJ1 . . .	
1	G481	BI129814	BI129814	G095P88Y	<i>Populus cambium</i> cDNA libra . . .
1	G481	BG318871	BG318871	NXPV_020_H08_F	NXPV (Nsf Xylem Plan . . .
1	G481	AV420653	AV420653	AV420653	<i>Lotus japonicus</i> young plan . . .
1	G481	AW348165	AW348165	GM210001A21D7	Gm-r1021 <i>Glycine max</i> . . .
1	G481	AI442376	AI442376	sa26b07.y1	Gm-c1004 <i>Glycine max</i> cDN . . .
1	G481	AI442765	AI442765	sa26b07.x1	Gm-c1004 <i>Glycine max</i> cDN . . .
1	G481	BE604847	BE604847	WHE1713-1716_D19_D19ZS	Wheat heat s . . .
1	G481	AW043377	AW043377	ST32F09	Pine TriplEx shoot tip libr . . .
1	G481	AP003271	AP003271	<i>Oryza sativa</i> genomic DNA, chromosom . . .	
1	G481	BG832836	BG832836	NXPV_081_C10_F	NXPV (Nsf Xylem Plan . . .
1	G481	AP004366	AP004366	<i>Oryza sativa</i> chromosome 1 clone P04 . . .	
1	G481	BE603222	BE603222	HVSMEh0102J16f	<i>Hordeum vulgare</i> 5-45 . . .
1	G481	AW754604	AW754604	PC04B12	Pine TriplEx pollen cone li . . .
1	G481	AW756413	AW756413	sl21a12.y1	Gm-c1036 <i>Glycine max</i> cDN . . .
1	G481	AW432980	AW432980	si03a01.y1	Gm-c1029 <i>Glycine max</i> cDN . . .
1	G481	BG551755	BG551755	sad42f11.y1	Gm-c1075 <i>Glycine max</i> cD . . .
1	G481	BF595304	BF595304	su76f03.y1	Gm-c1055 <i>Glycine max</i> cDN . . .
1	G481	BM308208	BM308208	sak43a12.y1	Gm-c1036 <i>Glycine max</i> cD . . .
1	G481	AC104284	AC104284	<i>Oryza sativa</i> chromosome 5 clone OJ1 . . .	
1	G481	BM528842	BM528842	sak69b03.y1	Gm-c1036 <i>Glycine max</i> cD . . .
1	G481	BI268123	BI268123	NF116D11IN1F1094	Insect herbivory M . . .
1	G481	AW981720	AW981720	PC15H07	Pine TriplEx pollen cone li . . .
1	G481	AA660543	AA660543	00429 MtRHE	<i>Medicago truncatula</i> cDN . . .
1	G481	BE726750	BE726750	894093C12.y3	<i>C. reinhardtii</i> CC-1690 . . .
1	G481	BI309186	BI309186	EST530596	GPOD <i>Medicago truncatula</i> . . .
1	G481	AW931376	AW931376	EST357219	tomato fruit mature green . . .
1	G481	BF636140	BF636140	NF060H09DT1F1079	Drought <i>Medicago t</i> . . .
1	G481	BG526135	BG526135	57-6	<i>Stevia</i> field grown leaf cDNA S . . .
1	G481	BI311277	BI311277	EST5313027	GESD <i>Medicago truncatula</i> . . .
1	G481	BI721770	BI721770	1031057H04.y1	<i>C. reinhardtii</i> CC-169 . . .
1	G481	AW200790	AW200790	se93e11.y1	Gm-c1027 <i>Glycine max</i> cDN . . .
1	G481	AY058919	AY058919	<i>Vernonia galamensis</i> CCAAT-box bindi . . .	
1	G481	BM109471	BM109471	EST557007	potato roots <i>Solanum tube</i> . . .
1	G481	AY058917	AY058917	<i>Glycine max</i> clone se2.11d12	CCAAT-b . . .
1	G481	AY058920	AY058920	<i>Argemone mexicana</i> CCAAT-box binding . . .	
1	G481	AU088581	AU088581	Rice callus	<i>Oryza sativa</i> c . . .
1	G481	AW720671	AW720671	LjNEST6a3rc	<i>Lotus japonicus</i> nodule . . .
1	G481	BI419749	BI419749	LjNEST14e12r	<i>Lotus japonicus</i> nodule . . .
1	G481	AW719547	AW719547	LjNEST6a3r	<i>Lotus japonicus</i> nodule I . . .
1	G481	BM134935	BM134935	WHE0460_A02_A03ZS	Wheat Fusarium gr . . .
1	G481	AI965590	AI965590	sc74b05.y1	Gm-c1018 <i>Glycine max</i> cDN . . .
1	G481	BI875522	BI875522	963125B06.y1	<i>C. reinhardtii</i> CC-1690 . . .
1	G481	BI531782	BI531782	1024116E03.y1	<i>C. reinhardtii</i> CC-169 . . .
1	G481	AW688588	AW688588	NF009C11ST1F1000	Developing stem Me . . .
1	G481	BG368375	BG368375	HVSMEi0018C01f	<i>Hordeum vulgare</i> 20 D . . .
1	G481	BF270164	BF270164	GA_Eb0007A21f	<i>Gossypium arboreum</i> 7 . . .
1	G481	BH472297	BH472297	BOGF90TF	BOGJ <i>Brassica oleracea</i> ge . . .
1	G481	AY058918	AY058918	<i>Glycine max</i> clone ses2w.pk0015.a4	C . . .
1	G481	BF291752	BF291752	WHE2205_F04_K07ZS	<i>Aegilops speltoid</i> . . .
1	G481	BF169598	BF169598	NXCI_125_B04_F	NXCI (Nsf Xylem Comp . . .
1	G481	BH470962	BH470962	BOGNF35TF	BOGN <i>Brassica oleracea</i> ge . . .
1	G481	BI952722	BI952722	HVSME0007119f	<i>Hordeum vulgare</i> gree . . .
1	G481	AL506199	AL506199	<i>Hordeum vulgare</i> Barke deve . . .	
1	G481	BH659234	BH659234	BOMDK68TR BO_2_3_KB	<i>Brassica olerac</i> . . .
1	G481	BE802539	BE802539	sr32f02.y1	Gm-c1050 <i>Glycine max</i> cDN . . .
1	G481	BE196056	BE196056	HVSMEh0091D23f	<i>Hordeum vulgare</i> 5-45 . . .
1	G481	AL509098	AL509098	<i>Hordeum vulgare</i> Barke deve . . .	
1	G481	AP003266	AP003266	<i>Oryza sativa</i> genomic DNA, chromosom . . .	
1	G481	AF410176	AF410176	<i>Zea mays</i> leafy cotyledon1 (Lec1) mR . . .	
1	G481	AX365282	AX365282	Sequence 18 from Patent WO0206499 . . .	
1	G481	AY058921	AY058921	<i>Triticum aestivum</i> CCAAT-box binding . . .	
1	G481	BG662094	BG662094	Ljirnp38-110-g8	Ljirnp Lambda Hy . . .
1	G481	BH645253	BH645253	BOMFL56TR BO_2_3_KB	<i>Brassica olerac</i> . . .
1	G481	AP004179	AP004179	<i>Oryza sativa</i> chromosome 2 clone OJ1 . . .	
1	G481	BM500534	BM500534	PAC000000000627	Pioneer AF-1 array . . .
1	G481	BH701005	BH701005	BOMMD16TR BO_2_3_KB	<i>Brassica olerac</i> . . .
1	G481	BH678940	BH678940	BOMIF09TF BO_2_3_KB	<i>Brassica olerac</i> . . .
1	G481	AP004791	AP004791	<i>Oryza sativa</i> (japonica cultivar-gro . . .	
1	G481	BE121888	BE121888	894015G05.y1	<i>C. reinhardtii</i> CC-1690 . . .
1	G481	AX288136	AX288136	Sequence 7 from Patent WO0177311. 1 . . .	

TABLE 7-continued

SEQ ID	Test	Reading	High	Smallest Sum	
NO	Gene ID	Frame	Score	Probability	N
	Sequence ID	Test Sequence	GenBank	Annotation	
1	G481	AV411210	AV411210	AV411210	<i>Lotus japonicus</i> young plan . . .
1	G481	AV425835	AV425835	AV425835	<i>Lotus japonicus</i> young plan . . .
1	G481	BI206716	EST524756	cTOS	<i>Lycopersicon esculen</i> . . .
1	G481	BF645376	BF645376	NF040B05EC1F1044	Elicited cell cult . . .
1	G481	BM347760	BM347760	MEST281-G09.T3	ISUM5-RN <i>Zea mays</i> cD . . .
1	G481	AW648379	AW648379	EST326833	tomato germinating seedli . . .
1	G481	BI176409	BI176409	EST521199	P. infestans-challenged l . . .
1	G481	BM109406	BM109406	EST556942	potato roots Solanum tube . . .
1	G481	BM348480	BM348480	MEST291-E08.T3	ISUM5-RN <i>Zea mays</i> cD . . .
1	G481	BI531808	BI531808	1024116G03.y1	<i>C. reinhardtii</i> CC-169 . . .
1	G481	BM158109	BM158109	NXLV_029_E11_F	NXLV (Nsf Xylem Late . . .
1	G481	AI966550	AI966550	sc51h01.y1	Gm-c1015 <i>Glycine max</i> cDN . . .
1	G481	BM341073	BM341073	MEST329-H08.T3	ISUM5-RN <i>Zea mays</i> cD . . .
1	G481	BM335521	BM335521	MEST162-H08.T3	ISUM5-RN <i>Zea mays</i> cD . . .
1	G481	AV422691	AV422691	AV422691	<i>Lotus japonicus</i> young plan . . .
1	G481	BG039303	BG039303	NXSL_097_E11_F	NXSL (Nsf Xylem Side . . .
1	G481	BF068031	BF068031	st86h12.y1	Gm-c1054 <i>Glycine max</i> cDN . . .
1	G481	BF777951	BF777951	NXSL_079_003_F	NXSL (Nsf Xylem Side . . .
1	G481	BI720257	BI720257	1031048F06.y1	<i>C. reinhardtii</i> CC-169 . . .
1	G481	BM441686	BM441686	EBed07_SQ001_E05_R	IGF Barley EBed0 . . .
1	G481	AX365284	AX365284	Sequence 20	from Patent WO0206499. . .
1	G481	BH153040	BH153040	Gm_ISb001_083_P16R	ISU Soybean BAC . . .
1	G481	BG589029	BG589029	EST490838	MHRP- <i>Medicago truncatula</i> . . .
1	G481	BG350792	BG350792	098C07	Mature tuber lambda ZAP Sola . . .
1	G481	BM094268	BM094268	sah27d01.y1	Gm-c1036 <i>Glycine max</i> cD . . .
1	G481	BM271333	BM271333	sak08b06.y1	Gm-c1075 <i>Glycine max</i> cD . . .
1	G481	AW693654	AW693654	NF066H08ST1F1000	Developing stem Me . . .
1	G481	C19290	C19290	C19290	Rice panicle at ripening stage . . .
1	G481	AL388746	AL388746	MtBC50E04F1	MtBC <i>Medicago truncatul</i> . . .
1	G481	BI068503	BI068503	C022P78U	Populus strain T89 leaves . . .
1	G481	BH591716	BH591716	BOGTY51TR	BOGT <i>Brassica oleracea</i> ge . . .
1	G481	BM333505	BM333505	MEST156-F03.T3	ISUM5-RN <i>Zea mays</i> cD . . .
1	G481	BH645120	BH645120	BOHZM09TF	BO_2_3_KB <i>Brassica olerac</i> . . .
1	G481	BM112643	BM112643	EST560179	potato roots Solanum tube . . .
1	G481	BE356637	BE356637	DG1_126_D05.g1	A002 Dark Grown 1 (D . . .
1	G481	BH580896	BH580896	BOGLY95TR	BOGL <i>Brassica oleracea</i> ge . . .
1	G481	BE400220	BE400220	AWB001.E09F000328	ITEC AWB Wheat Me . . .
1	G481	AW981721	AW981721	PC15H08	Pine TriplEx pollen cone li . . .
1	G481	BI325183	BI325183	baa05b02.x1	Cassava EYC library1 Ma . . .
1	G481	BE822946	BE822946	GM700019A20C12	Gm-r1070 <i>Glycine max</i> . . .
1	G481	BI271659	BI271659	NF026A03FL1F1020	Developing flower . . .
1	G481	AMTAM4	X59057	<i>A. majus</i> transposable element	Tam4 DNA . . .
1	G481	BG653330	BG653330	sad87a07.y1	Gm-c1055 <i>Glycine max</i> cD . . .
1	G481	BG644353	BG644353	EST505972	KV3 <i>Medicago truncatula</i> c . . .
1	G481	BG650027	BG650027	sad90h01.y1	Gm-c1055 <i>Glycine max</i> cD . . .
1	G481	AW734322	AW734322	sk81f11.y1	Gm-c1016 <i>Glycine max</i> cDN . . .
1	G481	BG046421	BG046421	saa63e06.y1	Gm-c1060 <i>Glycine max</i> cD . . .
1	G481	BG726061	BG726061	sac06d09.y1	Gm-c1055 <i>Glycine max</i> CD . . .
1	G481	BF070629	BF070629	st23b12.y1	Gm-c1065 <i>Glycine max</i> cDN . . .
1	G481	AI443631	AI443631	sa42e05.y1	Gm-c1004 <i>Glycine max</i> cDN . . .
1	G481	BF009637	BF009637	ss81h09.y1	Gm-c1064 <i>Glycine max</i> cDN . . .
1	G481	BF627138	BF627138	HVSMEb0004A20f	<i>Hordeum vulgare</i> seed . . .
1	G481	AV420244	AV420244	AV420244	<i>Lotus japonicus</i> young plan . . .
1	G481	BH637694	BH637694	1008018B12.2EL_x1	1008 —RescueMu G . . .
1	G481	AI967494	AI967494	Ljirmpst03-196-b3	Ljirnp Lambda Hy . . .
1	G481	AV416827	AV416827	AV416827	<i>Lotus japonicus</i> young plan . . .
1	G481	BG131195	BG131195	EST464087	tomato crown gall Lycoper . . .
1	G481	BF052943	BF052943	EST438173	potato leaves and petiole . . .
1	G481	AV409183	AV409183	AV409183	<i>Lotus japonicus</i> young plan . . .
1	G481	AW776198	AW776198	EST335263	DSIL <i>Medicago truncatula</i> . . .
1	G481	AL368819	AL368819	MtBA27A03F1	MtBA <i>Medicago truncatul</i> . . .
1	G481	C99407	C99407	C99407	Rice panicle at ripening stage . . .
1	G481	BH448006	BH448006	BOGBH37TR	BOGB <i>Brassica oleracea</i> ge . . .
1	G481	AW185273	AW185273	se89c10.y1	Gm-c1023 <i>Glycine max</i> cDN . . .
1	G481	AW423569	AW423569	sh68f08.y1	Gm-c1015 <i>Glycine max</i> cDN . . .
1	G481	BG507868	BG507868	sac82d02.y1	Gm-c1072 <i>Glycine max</i> cD . . .
1	G481	C19737	C19737	C19737	Rice panicle at ripening stage . . .
1	G481	AW719575	AW719575	LjNEST6a11r	<i>Lotus japonicus</i> nodule . . .
1	G481	AW706867	AW706867	sk07d05.y1	Gm-c1023 <i>Glycine max</i> cDN . . .
1	G481	AI460665	AI460665	sa71g05.y1	Gm-c1004 <i>Glycine max</i> cDN . . .
1	G481	AW759521	AW759521	sl44d11.y1	Gm-c1027 <i>Glycine max</i> cDN . . .
1	G481	BF595796	BF595796	sv04f12.y1	Gm-c1056 <i>Glycine max</i> cDN . . .
1	G481	BM178052	BM178052	saj68e01.y1	Gm-c1072 <i>Glycine max</i> cD . . .

TABLE 7-continued

SEQ ID	Test	Reading	High	Smallest Sum	
NO	Gene ID	Frame	Score	Probability	N
	Sequence ID	Test Sequence	GenBank	Annotation	
1	G481	BI424397	BI424397	saf34c08.y4 Gm-c1077 <i>Glycine max</i> cD . . .	1 110 0.16 1
1	G481	BG044257	BG044257	saa25h07.y1 Gm-c1059 <i>Glycine max</i> cD . . .	2 110 0.17 1
1	G481	AW234956	AW234956	sf21b08.y1 Gm-c1028 <i>Glycine max</i> cDN . . .	1 110 0.17 1
1	G481	AW759820	AW759820	sl54e01.y1 Gm-c1027 <i>Glycine max</i> cDN . . .	3 110 0.18 1
1	G481	AI959799	AI959799	sc94d03.y1 Gm-c1019 <i>Glycine max</i> cDN . . .	1 110 0.18 1
1	G481	AW201988	AW201988	sf09f10.y1 Gm-c1027 <i>Glycine max</i> cDN . . .	3 109 0.2 1
1	G481	AW164642	AW164642	se74f06.y1 Gm-c1023 <i>Glycine max</i> cDN . . .	3 110 0.2 1
1	G481	AT002114	AT002114	AT002114 Flower bud cDNA Brassica r . . .	2 98 0.23 1
1	G481	BF113032	BF113032	EST440542 tomato breaker fruit Lyco . . .	1 107 0.25 1
1	G481	BE659989	BE659989	1010 GmaxSC <i>Glycine max</i> cDNA, mRNA . . .	2 110 0.26 1
1	G481	OSA300218	AJ300218	<i>Oryza sativa</i> nf-yb1 gene and nf-YB1 . . .	2 111 0.27 1
1	G481	BG597547	BG597547	EST496225 cSTS <i>Solanum tuberosum</i> cD . . .	1 107 0.32 1
1	G481	BF644204	BF644204	NF060B12EC1F1101 Elicited cell cult . . .	2 105 0.33 1
1	G481	AF464906	AF464906	<i>Glycine max</i> repressor protein (Dr1) . . .	1 110 0.34 1
1	G481	BE436801	BE436801	EST407919 tomato breaker fruit, TIG . . .	2 107 0.36 1
1	G481	BE659987	BE659987	7-F12 GmaxSC <i>Glycine max</i> cDNA, mRNA . . .	1 109 0.36 1
1	G481	BG451060	BG451060	NF098C04DT1F1024 Drought Medicago t . . .	3 105 0.42 1
1	G481	BM436739	BM436739	WA009B06_53061 An expressed sequen . . .	3 107 0.45 1
1	G481	BF273545	BF273545	GA_Eb0018J12f <i>Gossypium arboreum</i> 7 . . .	1 104 0.5 1
1	G481	BM411335	BM411335	EST585662 tomato breaker fruit Lyco . . .	3 107 0.5 1
1	G481	AW830697	AW830697	sm06h11.y1 Gm-c1027 <i>Glycine max</i> cDN . . .	2 91 0.54 1
1	G481	BE202566	BE202566	EST392975 KV1 <i>Medicago truncatula</i> c . . .	2 105 0.55 1
1	G481	BG240158	BG240158	OV1_18_F02.b1_A002 Ovary 1 (OV1) So . . .	3 101 0.56 1
1	G481	BM817060	BM817060	HC01C02_T3.ab1 HC <i>Hordeum vulgare</i> c . . .	1 103 0.58 1
1	G481	BE998485	BE998485	EST430208 GVSN <i>Medicago truncatula</i> . . .	3 105 0.58 1
1	G481	BE022456	BE022456	sm74b08.y1 Gm-c1015 <i>Glycine max</i> cDN . . .	2 103 0.6 1
1	G481	BF647976	BF647976	NF013H04EC1F1042 Elicited cell cult . . .	1 105 0.63 1
1	G481	BG102166	BG102166	RHIZ2_21_F06.b1_A003 Rhizome2 (RHIZ . . .	3 101 0.63 1
1	G481	AU084707	AU084707	AU084707 <i>Cryptomeria japonica</i> inner . . .	1 101 0.68 1
1	G481	BG648823	BG648823	EST510442 HOGA <i>Medicago truncatula</i> . . .	1 105 0.7 1
1	G481	BG648909	BG648909	EST510528 HOGA <i>Medicago truncatula</i> . . .	2 105 0.72 1
1	G481	BM380524	BM380524	MEST521-B07.univ ISUM6 <i>Zea mays</i> cDN . . .	-3 103 0.75 1
1	G481	BG593107	BG593107	EST491785 cSTS <i>Solanum tuberosum</i> cD . . .	3 103 0.81 1
1	G481	BG052069	BG052069	RHIZ2_5_F11.b1_A003 Rhizome2 (RHIZ2 . . .	3 101 0.9 1
1	G481	BE497740	BE497740	WHE0956_G06_M12ZS Wheat pre-anthesi . . .	2 100 0.92 1
1	G481	BG873649	BG873649	MEST8-E10.T7-1 ISUM3-TL <i>Zea mays</i> cD . . .	3 101 0.94 1
1	G481	AW432997	AW432997	si03b08.y1 Gm-c1029 <i>Glycine max</i> cDN . . .	2 88 0.96 1
1	G481	AW064635	AW064635	ST33H06 Pine TriplEx shoot tip libr . . .	2 100 0.96 1
1	G481	AX365283	AX365283	Sequence 19 from Patent WO0206499 . . .	1 101 0.97 1
1	G481	BG648613	BG648613	EST510232 HOGA <i>Medicago truncatula</i> . . .	3 99 0.993 1
1	G481	BE640725	BE640725	Cri2_1_E07_SP6 Ceratopteris Spore L . . .	2 100 0.995 1
1	G481	AF464902	AF464902	<i>Oryza sativa</i> repressor protein (Dr1) . . .	2 101 0.996 1
1	G481	AF464903	AF464903	<i>Triticum aestivum</i> repressor protein . . .	2 100 0.999 1
1	G481	BE449790	BE449790	EST361228 tomato root, plants pre-a . . .	3 97 0.9991 1
1	G481	BI206380	BI206380	EST524420 cTOS Lycopersicon esculen . . .	1 97 0.9996 1
1	G481	GI-22380	CAAT-box	DNA binding protein subunit B (N . . .	1 262 1.10E-45 3
1	G481	GI-115840	CBFA_MAIZE	CCAAT-BINDING TRANSCRIPTION FAC . . .	1 262 1.10E-45 3
1	G481	GI-7443522	S22820	transcription factor NF-Y, CCAAT-bi . . .	1 262 1.10E-45 3
1	G481	GI-15408794	putative	CCAAT-binding transcription facto . . .	1 194 2.60E-30 2
1	G481	GI-16902054	CCAAT-box	binding factor HAP3 B domain [Ve . . .	1 187 1.00E-28 2
1	G481	GI-16902050	CCAAT-box	binding factor HAP3 B domain [Gl . . .	1 188 2.70E-28 2
1	G481	GI-16902056	CCAAT-box	binding factor HAP3 B domain [Ar . . .	1 189 4.30E-28 2
1	G481	GI-15321716	AF410176_1	leafy cotyledon1 [<i>Zea mays</i>] . . .	1 179 1.20E-26 3
1	G481	GI-16902052	CCAAT-box	binding factor HAP3 B domain [Gl . . .	1 178 2.90E-26 2
1	G481	GI-15408793	hypothetical	protein~similar to CCAAT-bind . . .	1 180 2.70E-24 2
1	G481	GI-16902058	CCAAT-box	binding factor HAP3 B domain [Tr . . .	1 175 1.50E-23 2
1	G481	GI-18481628	AF464906_1	repressor protein [<i>Glycine max</i>] . . .	1 110 2.10E-09 2
1	G481	GI-13928060	NF-YB1	protein [<i>Oryza sativa</i>] . . .	1 111 0.00069 1
1	G481	GI-18481620	AF464902_1	repressor protein [<i>Oryza sativa</i>] . . .	1 101 0.037 1
1	G481	GI-18481622	AF464903_1	repressor protein [<i>Triticum aes</i> . . .	1 100 0.052 1
3	G1466	BH596941	BH596941	BOHFG41TR BOHF <i>Brassica oleracea</i> ge . . .	1 557 1.40E-65 2
3	G1466	BH497171	BH497171	BOHKV94TF BOHK <i>Brassica oleracea</i> ge . . .	-1 531 7.70E-49 1
3	G1466	BH515525	BH515525	BOGZE84TR BOGZ <i>Brassica oleracea</i> ge . . .	-1 518 1.60E-47 1
3	G1466	BH685857	BH685857	BOHWL93TF BO_2_3_KB <i>Brassica olerac</i> . . .	1 409 6.90E-36 1
3	G1466	BH556937	BH556937	BOHCT42TF BOHC <i>Brassica oleracea</i> ge . . .	3 374 3.20E-32 1
3	G1466	BH716964	BH716964	BOMNM26TR BO_2_3_KB <i>Brassica olerac</i> . . .	-2 371 6.40E-32 1
3	G1466	BH556945	BH556945	BOHCT42TR BOHC <i>Brassica oleracea</i> ge . . .	-3 205 3.40E-24 2
3	G1466	BH586673	BH586673	BOHPK25TF BOHP <i>Brassica oleracea</i> ge . . .	-1 293 1.80E-23 1
3	G1466	BH479514	BH479514	BOHEU46TF BOHE <i>Brassica oleracea</i> ge . . .	-3 293 2.50E-23 1
3	G1466	BH442040	BH442040	BOHJW94TR BOHJ <i>Brassica oleracea</i> ge . . .	2 284 2.00E-22 1
3	G1466	BH468201	BH468201	BOHAW05TF BOHA <i>Brassica oleracea</i> ge . . .	3 263 6.80E-20 1
3	G1466	BH554599	BH554599	BOHES28TF BOHE <i>Brassica oleracea</i> ge . . .	-2 126 4.40E-17 3

TABLE 7-continued

SEQ ID	Test	Reading	High	Smallest Sum	
NO	Gene ID	Sequence ID	Test Sequence	GenBank	Annotation
			Frame	Score	Probability
					N
3	G1466	BH443975	BH443975	BOGYY14TR	BOGY <i>Brassica oleracea</i> ge . . .
3	G1466	BH478506	BH478506	BOGJO28TR	BOGJ <i>Brassica oleracea</i> ge . . .
3	G1466	BH248066	BH248066	BOGAU25TR	BOGA <i>Brassica oleracea</i> ge . . .
3	G1466	BH455609	BH455609	BOHPV23TF	BOHP <i>Brassica oleracea</i> ge . . .
3	G1466	BH453095	BH453095	BOGVR55TF	BOGV <i>Brassica oleracea</i> ge . . .
3	G1466	BH536461	BH536461	BOHJM41TR	BOHM <i>Brassica oleracea</i> ge . . .
3	G1466	BE462774	BE462774	EST325108	tomato flower buds 0-3 mm . . .
3	G1466	AW399721	AW399721	EST310221	<i>L. pennellii</i> trichome, Co . . .
3	G1466	AV413010	AV413010	AV413010	<i>Lotus japonicus</i> young plan . . .
3	G1466	AW032605	AW032605	EST276164	tomato callus, TAMU Lycop . . .
3	G1466	BI927221	BI927221	EST547110	tomato flower, 3-8 mm b . . .
3	G1466	OSIN00182	AL662981	<i>Oryza sativa</i> chromosome 4 clone	OSJ . . .
3	G1466	BG662160	BG662160	Ljirnp39-183-g3	Ljirnp Lambda Hy . . .
3	G1466	BG599234	BG599234	EST504134	cSTS <i>Solanum tuberosum</i> cD . . .
3	G1466	BG133465	BG133465	EST466357	tomato crown gall Lycoper . . .
3	G1466	BG596415	BG596415	EST495093	cSTS <i>Solanum tuberosum</i> cD . . .
3	G1466	BI922297	BI922297	EST542201	tomato callus Lycopersico . . .
3	G1466	BG097828	BG097828	EST462347	potato leaves and petiole . . .
3	G1466	BI178485	BI178485	EST519430	cSTE <i>Solanum tuberosum</i> cD . . .
3	G1466	BG580068	BG580068	EST481790	GVN <i>Medicago truncatula</i> c . . .
3	G1466	AW932502	AW932502	EST358345	tomato fruit mature green . . .
3	G1466	BH700517	BH700517	BOMEG26TR	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	AV414925	AV414925	AV414925	<i>Lotus japonicus</i> young plan . . .
3	G1466	BG887252	BG887252	EST513103	cSTD <i>Solanum tuberosum</i> cD . . .
3	G1466	BH649717	BH649717	BOHTZ83TF	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	AW691672	AW691672	NF047G10ST1F1000	Developing stem Me . . .
3	G1466	BI316339	BI316339	saf02a08.y1	Gm-c1065 <i>Glycine max</i> cD . . .
3	G1466	BG409415	BG409415	00787	leafy spurge Lambda HybriZAP . . .
3	G1466	AP003837	AP003837	<i>Oryza sativa</i> chromosome 7 clone	OJ1 . . .
3	G1466	AW928976	AW928976	EST337860	tomato flower buds 8 mm t . . .
3	G1466	OSIN00167	AL662965	<i>Oryza sativa</i> chromosome 4 clone	OSI . . .
3	G1466	BH660321	BH660321	BOMGE76TR	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	AW395660	AW395660	sg73f12.y1	Gm-c1007 <i>Glycine max</i> cDN . . .
3	G1466	BE459691	BE459691	EST414983	tomato developing/immatur . . .
3	G1466	BH558685	BH558685	BOGEH17TR	BOGE <i>Brassica oleracea</i> ge . . .
3	G1466	BH486360	BH486360	BOHLY92TF	BOHL <i>Brassica oleracea</i> ge . . .
3	G1466	AW218077	AW218077	EST296792	tomato flower buds, anthe . . .
3	G1466	AW650052	AW650052	EST328506	tomato germinating seedli . . .
3	G1466	BH698615	BH698615	BOHUQ33TR	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	BG596140	BG596140	EST494818	cSTS <i>Solanum tuberosum</i> cD . . .
3	G1466	BE205043	BE205043	EST397719	KV0 <i>Medicago truncatula</i> c . . .
3	G1466	BE919377	BE919377	EST423230	potato leaves and petiole . . .
3	G1466	BH437084	BH437084	BOHKZ27TF	BOHK <i>Brassica oleracea</i> ge . . .
3	G1466	BF278686	BF278686	GA_Eb0035E21f	<i>Gossypium arboreum</i> 7 . . .
3	G1466	BI933949	BI933949	EST553838	tomato flower, anthesis L . . .
3	G1466	BF096307	BF096307	EST360356	tomato nutrient deficient . . .
3	G1466	BH737844	BH737844	BOMAX76TF	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	BI929823	BI929823	EST549712	tomato flower, 3-8 mm b . . .
3	G1466	BH513097	BH513097	BOGEX21TR	BOGE <i>Brassica oleracea</i> ge . . .
3	G1466	BE203214	BE203214	EST403236	KV1 <i>Medicago truncatula</i> c . . .
3	G1466	BE202448	BE202448	EST392897	KV1 <i>Medicago truncatula</i> c . . .
3	G1466	BF644564	BF644564	NF016D06EC1F1057	Elicited cell cult . . .
3	G1466	BI420528	BI420528	LjNEST58c10r	<i>Lotus japonicus</i> nodule . . .
3	G1466	BF010496	BE010496	NXCI_084_G01_F	NXCI (Nsf Xylem Comp . . .
3	G1466	BI308763	BI308763	EST530173	GPOD <i>Medicago truncatula</i> . . .
3	G1466	BH469525	BH469525	BOGSA70TF	BOGS <i>Brassica oleracea</i> ge . . .
3	G1466	BE442810	BE442810	WHE1106_C02_E04ZS	Wheat etiolated s . . .
3	G1466	BH569775	BH569775	BOGMV91TF	BOGM <i>Brassica oleracea</i> ge . . .
3	G1466	BH723965	BH723965	BOMHF56TF	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	BH475440	BH475440	BOHNA92TF	BOHN <i>Brassica oleracea</i> ge . . .
3	G1466	BH730453	BH730453	BOHVD24TR	BO_2_3_KB <i>Brassica olerac</i> . . .
3	G1466	AC082644	AC082644	<i>Oryza sativa</i> chromosome 3 BAC	OSJNB . . .
3	G1466	AZ124264	AZ124264	T223049b	<i>Medicago truncatula</i> BAC li . . .
3	G1466	BI922762	BI922762	EST542666	tomato callus Lycopersico . . .
3	G1466	AI443156	AI443156	sa50f01.y1	Gm-c1004 <i>Glycine max</i> cDN . . .
3	G1466	BE249557	BE249557	NF022C06LF1F1049	Developing leaf Me . . .
3	G1466	AL379504	AL379504	MtBB45G03F1	MtBB <i>Medicago truncatul</i> . . .
3	G1466	BF646352	BF646352	NF068G05EC1F1039	Elicited cell cult . . .
3	G1466	BH498652	BH498652	BOGZR84TR	BOGZ <i>Brassica oleracea</i> ge . . .
3	G1466	BM442042	BM442042	EBan01_SQ002_A06_R	IGF Barley EBan0 . . .
3	G1466	BG447666	BG447666	NF019E03ST1F1000	Developing stem Me . . .
3	G1466	AV427033	AV427033	AV427033	<i>Lotus japonicus</i> young plan . . .
3	G1466	AW216996	AW216996	EST295710	tomato callus, TAMU Lycop . . .

TABLE 7-continued

SEQ ID	Test	Reading	High	Smallest Sum				
NO	Gene ID	Sequence ID	Test Sequence	GenBank Annotation	Frame	Score	Probability	N
3	G1466	AW648500	AW648500	EST326954 tomato germinating seedli . . .	3	102	0.55	1
3	G1466	AW649521	AW649521	EST327975 tomato germinating seedli . . .	1	102	0.59	1
3	G1466	BF187212	BF187212	EST443499 potato stolon, Cornell Un . . .	3	102	0.62	1
3	G1466	BE202443	BE202443	EST392892 KV1 <i>Medicago truncatula</i> c . . .	3	104	0.65	1
3	G1466	BE344294	BE344294	EST409456 potato stolon, Cornell Un . . .	3	102	0.66	1
3	G1466	C72509	C72509	C72509 Rice panicle at flowering stag . . .	2	89	0.67	1
3	G1466	BH743977	BH743977	gt29g06.b1 BoBuds01 Brassica olerac . . .	2	101	0.68	1
3	G1466	AI967453	AI967453	Ljirnp02-129-d2 Ljirnp Lambda Hy . . .	1	99	0.7	1
3	G1466	BI419994	BI419994	LjNEST42f9r <i>Lotus japonicus</i> nodule . . .	2	102	0.7	1
3	G1466	BG646874	BG646874	EST508493 HOGA <i>Medicago truncatula</i> . . .	2	104	0.7	1
3	G1466	AW694435	AW694435	NF076B06ST1F1048 Developing stem Me . . .	1	102	0.75	1
3	G1466	AP004055	AP004055	<i>Oryza sativa</i> chromosome 2 clone OJ1 . . .	1	107	0.77	1
3	G1466	AP004144	AP004144	<i>Oryza sativa</i> chromosome 2 clone OJ1 . . .	-3	107	0.77	1
3	G1466	OSJN00115	AL606997	<i>Oryza sativa</i> chromosome 4 clone OSJ . . .	-1	107	0.77	1
3	G1466	AP002071	AP002071	<i>Oryza sativa</i> genomic DNA, chromosom . . .	-1	107	0.77	1
3	G1466	BE095245	BE095245	00306 leafy spurge Lambda HybriZAP . . .	1	88	0.79	1
3	G1466	BM526605	BM526605	sal43e01.y1 Gm-c1059 <i>Glycine max</i> cD . . .	2	100	0.8	1
3	G1466	BI933721	BI933721	EST553610 tomato flower, anthesis L . . .	3	102	0.86	1
3	G1466	BI272408	BI272408	NF020H09FL1F1079 Developing flower . . .	3	102	0.87	1
3	G1466	BI933639	BI933639	EST553528 tomato flower, anthesis L . . .	3	102	0.89	1
3	G1466	BM108916	BM108916	EST556452 potato roots Solanum tube . . .	1	102	0.89	1
3	G1466	BH581102	BH581102	BOHKD50TF BOHK <i>Brassica oleracea</i> ge . . .	3	95	0.9	1
3	G1466	BH692394	BH692394	BOMKE64TF BO_2_3_KB Brassica olerac . . .	3	99	0.92	1
3	G1466	BG131783	BG131783	EST464675 tomato crown gall Lycoper . . .	3	97	0.93	1
3	G1466	AI896339	AI896339	EST265782 tomato callus, TAMU Lycop . . .	2	94	0.93	1
3	G1466	BG404995	BG404995	sac46g02.y1 Gm-c1062 <i>Glycine max</i> cD . . .	1	97	0.95	1
3	G1466	AP004042	AP004042	<i>Oryza sativa</i> chromosome 8 clone OJ1 . . .	1	104	0.95	1
3	G1466	AP004708	AP004708	<i>Oryza sativa</i> chromosome 8 clone P07 . . .	-3	104	0.95	1
3	G1466	AP004274	AP004274	<i>Oryza sativa</i> chromosome 7 clone P04 . . .	-1	104	0.95	1
3	G1466	BE919578	BE919578	EST423347 potato leaves and petiole . . .	1	100	0.95	1
3	G1466	BF634570	BF634570	NF061B09DT1F1076 Drought <i>Medicago t</i> . . .	3	100	0.96	1
3	G1466	BI420155	BI420155	LjNEST53f7r <i>Lotus japonicus</i> nodule . . .	3	96	0.97	1
3	G1466	AQ917137	AQ917137	T233170b <i>Medicago truncatula</i> BAC li . . .	-1	99	0.97	1
3	G1466	AW720029	AW720029	LjNEST15c9r <i>Lotus japonicus</i> nodule . . .	2	96	0.98	1
3	G1466	AW568479	AW568479	si59c09.y1 Gm-r1030 <i>Glycine max</i> cDN . . .	3	97	0.98	1
3	G1466	BI420626	BI420626	LjNEST59e5r <i>Lotus japonicus</i> nodule . . .	3	96	0.99	1
3	G1466	BE433506	BE433506	EST400035 tomato breaker fruit, TIG . . .	2	97	0.99	1
3	G1466	BI785473	BI785473	sai41d06.y1 Gm-c1065 <i>Glycine max</i> cD . . .	3	97	0.991	1
3	G1466	BH737087	BH737087	BOMGM48TR BO_2_3_KB Brassica olerac . . .	3	98	0.991	1
3	G1466	BM324672	BM324672	PIC1_34_B01.b1_A002 Pathogen-infect . . .	3	98	0.992	1
3	G1466	BH425892	BH425892	BOGDN33TF BOGD <i>Brassica oleracea</i> ge . . .	-3	98	0.994	1
3	G1466	BH645340	BH645340	BOHWO42TF BO_2_3_KB Brassica olerac . . .	-3	97	0.995	1
3	G1466	AW687534	AW687534	NF010F07RT1F1062 Developing root Me . . .	3	96	0.997	1
3	G1466	BF176934	BF176934	EM1_4_E11.b1_A002 Embryo 1 (EM1) So . . .	3	96	0.997	1
3	G1466	BI419241	BI419241	LjNEST44a5r <i>Lotus japonicus</i> nodule . . .	3	96	0.998	1
3	G1466	AL380183	AL380183	MtBB50H09F1 MtBB <i>Medicago truncatul</i> . . .	1	94	0.998	1
3	G1466	AC079128	AC079128	<i>Oryza sativa</i> chromosome 10 clone OS . . .	3	101	0.998	1
3	G1466	BF070938	BF070938	st85e05.y1 Gm-c1054 <i>Glycine max</i> cDN . . .	3	95	0.999	1
3	G1466	AW928966	AW928966	EST337850 tomato flower buds 8 mm t . . .	3	86	0.9995	1
3	G1466	BH429646	BH429646	BOGSF37TF BOGS <i>Brassica oleracea</i> ge . . .	-3	97	0.9997	1
3	G1466	BH560749	BH560749	BOGPC01TR BOGP <i>Brassica oleracea</i> ge . . .	-1	97	0.9997	1
3	G1466	AP004150	AP004150	<i>Oryza sativa</i> chromosome 2 clone OJ1 . . .	-1	100	0.9998	1
3	G1466	AP004077	AP004077	<i>Oryza sativa</i> chromosome 2 clone OJ1 . . .	1	100	0.9998	1
3	G1466	BG130763	BG130763	EST463655 tomato crown gall Lycoper . . .	3	96	0.9998	1
3	G1466	BH484808	BH484808	BOGQC69TR BOGQ <i>Brassica oleracea</i> ge . . .	1	97	0.9998	1
3	G1466	BF651138	BF651138	NF101G05EC1F1038 Elicited cell cult . . .	1	96	0.9999	1
3	G1466	BF650226	BF650226	NF090F10EC1F1089 Elicited cell cult . . .	3	96	0.9999	1
3	G1466	BF647598	BF647598	NF012A06EC1F1039 Elicited cell cult . . .	3	96	0.9999	1
3	G1466	BG447988	BG447988	NF103H10EC1F1090 Elicited cell cult . . .	3	96	0.9999	1
3	G1466	BH480188	BH480188	BOGZW54TR BOGZ <i>Brassica oleracea</i> ge . . .	3	97	0.99991	1
3	G1466	BF519054	BF519054	EST456514 DSIL <i>Medicago truncatula</i> . . .	3	96	0.99994	1
3	G1466	GI-8096405		Similar to <i>Arabidopsis thaliana</i> chromosome . . .	-3	137	2.30E-12	2
3	G1466	GI-8096413		hypothetical protein [<i>Oryza sativa</i>] . . .	-3	135	2.20E-11	2
3	G1466	GI-8096416		hypothetical protein [<i>Oryza sativa</i>] . . .	-3	111	9.40E-09	3
3	G1466	GI-13161526		S locus F-box (SLF)-S2 protein [Antirrhinu . . .	-3	92	3.40E-08	3
3	G1466	GI-13161540		SLF-S2 protein [Antirrhinum <i>hispanicum</i>] . . .	-3	92	3.40E-08	3
3	G1466	GI-13161528		S locus F-box (SLF)-S2-like protein [Antir . . .	-1	87	1.20E-07	3
3	G1466	GI-8096415		hypothetical protein [<i>Oryza sativa</i>] . . .	-1	102	3.60E-07	2
3	G1466	GI-18854994		AC087599_5 unknown protein [<i>Oryza sativa</i>] . . .	-1	90	1.10E-05	3
3	G1466	GI-14028986		AC079128_10 Unknown protein [<i>Oryza sativa</i>] . . .	-1	101	4.60E-05	2
3	G1466	GI-18854992		AC087599_3 putative transposase [<i>Oryza sat</i> . . .	-3	93	0.0014	3
3	G1466	GI-12039340		AC082644_9 hypothetical protein [<i>Oryza sat</i> . . .	-1	113	0.0029	1
3	G1466	GI-14018043		AC079936_2 Hypothetical protein [<i>Oryza sat</i> . . .	-1	87	0.012	2

TABLE 7-continued

SEQ ID	Test				Reading	High	Smallest Sum	
NO	Gene ID	Sequence ID	Test Sequence	GenBank Annotation	Frame	Score	Probability	N
3	G1466	GI-15451623	AC091734_4	Hypothetical protein [<i>Oryza sat</i> . . .	-1	87	0.012	2
3	G1466	GI-15528755		hypothetical protein [<i>Oryza sativa</i>]	-1	96	0.012	2
3	G1466	GI-8096406		hypothetical protein [<i>Oryza sativa</i>]	-3	98	0.013	2
3	G1466	GI-8096410		hypothetical protein [<i>Oryza sativa</i>]	-1	103	0.022	1
3	G1466	GI-18844804		contains ESTs D15126(C0122),C97919(C0122)~. . .	-1	99	0.083	1
3	G1466	GI-18449949	AC099733_6	Unknown protein [<i>Oryza sativa</i>]	-1	81	0.099	2
3	G1466	GI-18464016	AC090873_9	Hypothetical protein [<i>Oryza sat</i> . . .	-3	69	0.23	2
3	G1466	GI-19224986	AC077693_1	putative transposase protein, 5 . . .	-3	93	0.24	2
3	G1466	GI-12039332	AC082644_1	hypothetical protein [<i>Oryza sat</i> . . .	-1	91	0.5	1
3	G1466	GI-18461280		similar to <i>Oryza sativa</i> chromosome 1, P045 . . .	-3	72	0.61	2
3	G1466	GI-9858770	AF273333_2	BAC19.2 [<i>Lycopersicon esculentum</i>]	-1	70	0.83	2
3	G1466	GI-12039337	AC082644_6	hypothetical protein [<i>Oryza sat</i> . . .	-1	87	0.85	1
3	G1466	GI-18642684	AC074283_6	Hypothetical protein [<i>Oryza sat</i> . . .	-1	79	0.9	2
3	G1466	GI-18642688	AC074283_7	Unknown protein [<i>Oryza sativa</i>]	-1	75	0.97	1
3	G1466	GI-18087878	AC087182_15	hypothetical protein [<i>Oryza sa</i> . . .	-1	87	0.98	1
3	G1466	GI-18873858	AC079874_27	hypothetical protein [<i>Oryza sa</i> . . .	-3	81	0.994	1
3	G1466	GI-6505722		ribosomal RNA apurinic site specific lyase	-1	81	0.99993	1
3	G1466	GI-6513849		ribosomal RNA apurinic site specific lyase	-1	81	0.99993	1

[0215]

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 832

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (103)...(528)

<223> OTHER INFORMATION: G481

<400> SEQUENCE: 1

gagcgttttcg tagaaaaatt cgatttctct aaagccctaa aactaaaacg actatcccca 60

attccaagtt ctagggtttc catcttcccc aatctagtat aa atg gcg gat acg 114

Met Ala Asp Thr

1

cct tcg agc cca gct gga gat ggc gga gaa agc ggc ggt tcc gtt agg 162

Pro Ser Ser Pro Ala Gly Asp Gly Gly Glu Ser Gly Gly Ser Val Arg

5 10 15 20

gag cag gat cga tac ctt cct ata gct aat atc agc agg atc atg aag 210

Glu Gln Asp Arg Tyr Leu Pro Ile Ala Asn Ile Ser Arg Ile Met Lys

25 30 35

aaa gcg ttg cct cct aat ggt aag att gga aaa gat gct aag gat aca 258

Lys Ala Leu Pro Pro Asn Gly Lys Ile Gly Lys Asp Ala Lys Asp Thr

40 45 50

gtt cag gaa tgc gtc tct gag ttc atc agc ttc atc act agc gag gcc 306

Val Gln Glu Cys Val Ser Glu Phe Ile Ser Phe Ile Thr Ser Glu Ala

55 60 65

agt gat aag tgt caa aaa gag aaa agg aaa act gtg aat ggt gat gat 354

Ser Asp Lys Cys Gln Lys Glu Lys Arg Lys Thr Val Asn Gly Asp Asp

70 75 80

ttg ttg tgg gca atg gca aca tta gga ttt gag gat tac ctg gaa cct 402

Leu Leu Trp Ala Met Ala Thr Leu Gly Phe Glu Asp Tyr Leu Glu Pro

85 90 95 100

-continued

cta aag ata tac cta gcg agg tac agg gag ttg gag ggt gat aat aag	450
Leu Lys Ile Tyr Leu Ala Arg Tyr Arg Glu Leu Glu Gly Asp Asn Lys	
105 110 115	
 gga tca gga aag agt gga gat gga tca aat aga gat gct ggt ggc ggt	498
Gly Ser Gly Lys Ser Gly Asp Gly Ser Asn Arg Asp Ala Gly Gly Gly	
120 125 130	
 gtt tct ggt gaa gaa atg ccg agc tgg taa aagaagttgc aagtagtgat	548
Val Ser Gly Glu Glu Met Pro Ser Trp *	
135 140	
 taagaacaat cgccaaatga tcaagggaaa ttagagatca gtgagttggt tatagttgag	608
ctgatcgaca actatttcgg gtttactctc aatttcggtt atgttagttt gaacgtttgg	668
tttattgttt ccggtttagt tggttgtatt taaagatttc tctgttagat gttgagaaca	728
cttgaatgaa ggaaaaattt gtccacatcc tgttggttatt ttcgattcac tttcggaatt	788
tcatagctaa tttattctca tttaatacca aatccttaaa ttaa	832

<210> SEQ ID NO 2
 <211> LENGTH: 141
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 2

Met Ala Asp Thr Pro Ser Ser Pro Ala Gly Asp Gly Gly Glu Ser Gly	
1 5 10 15	
Gly Ser Val Arg Glu Gln Asp Arg Tyr Leu Pro Ile Ala Asn Ile Ser	
20 25 30	
Arg Ile Met Lys Lys Ala Leu Pro Pro Asn Gly Lys Ile Gly Lys Asp	
35 40 45	
Ala Lys Asp Thr Val Gln Glu Cys Val Ser Glu Phe Ile Ser Phe Ile	
50 55 60	
Thr Ser Glu Ala Ser Asp Lys Cys Gln Lys Glu Lys Arg Lys Thr Val	
65 70 75 80	
Asn Gly Asp Asp Leu Leu Trp Ala Met Ala Thr Leu Gly Phe Glu Asp	
85 90 95	
Tyr Leu Glu Pro Leu Lys Ile Tyr Leu Ala Arg Tyr Arg Glu Leu Glu	
100 105 110	
Gly Asp Asn Lys Gly Ser Gly Lys Ser Gly Asp Gly Ser Asn Arg Asp	
115 120 125	
Ala Gly Gly Gly Val Ser Gly Glu Glu Met Pro Ser Trp	
130 135 140	

<210> SEQ ID NO 3
 <211> LENGTH: 1342
 <212> TYPE: DNA
 <213> ORGANISM: Arabidopsis thaliana
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (16)...(1278)
 <223> OTHER INFORMATION: G1466

<400> SEQUENCE: 3

caacaagctt caaaa atg ggg gaa gaa gaa gag aac cct aat tcg atc gac	51
Met Gly Glu Glu Glu Glu Asn Pro Asn Ser Ile Asp	
1 5 10	
 att ctt ccc gag cta ctt gaa gaa gtt ctc ctt aga ttg ccc acg aaa	99
Ile Leu Pro Glu Leu Leu Glu Glu Val Leu Leu Arg Leu Pro Thr Lys	
15 20 25	

-continued

tcg atc ctc aaa tgc aga atc gtc tca aaa caa tgg agg tca ctc ctg	147
Ser Ile Leu Lys Cys Arg Ile Val Ser Lys Gln Trp Arg Ser Leu Leu	
30 35 40	
gaa tcg agt agg ttc gcg gag agg cat atg agt ctt caa aac agc cgc	195
Glu Ser Ser Arg Phe Ala Glu Arg His Met Ser Leu Gln Asn Ser Arg	
45 50 55 60	
cgg aga atc tta gct gct tac aac tgc gac tgc gcc gga cgg agg aag	243
Arg Arg Ile Leu Ala Ala Tyr Asn Cys Asp Cys Gly Gly Arg Arg Lys	
65 70 75	
ctc cta ccc gag tca cgg ttt gaa ggg gac gaa gag att gtc tat ctg	291
Leu Leu Pro Glu Ser Arg Phe Glu Gly Asp Glu Glu Ile Val Tyr Leu	
80 85 90	
cac tgc gac gcc tca cga ccc tcg atg act tgc caa ggt gtg atc tgc	339
His Cys Asp Ala Ser Arg Pro Ser Met Thr Cys Gln Gly Val Ile Cys	
95 100 105	
ttc ccc gag caa gat tgg atc atc gtt ttg aac cca tcg act agc caa	387
Phe Pro Glu Gln Asp Trp Ile Ile Val Leu Asn Pro Ser Thr Ser Gln	
110 115 120	
ctt cgg cga ttc cct tcc gcc ttg aac cat aac tgc aga ttt aga att	435
Leu Arg Arg Phe Pro Ser Gly Leu Asn His Asn Cys Arg Phe Arg Ile	
125 130 135 140	
gga tta tgg aag act ttc tct ccg gga aac tgg gta atg ggg ttt ggt	483
Gly Leu Trp Lys Thr Phe Ser Pro Gly Asn Trp Val Met Gly Phe Gly	
145 150 155	
aga gac aaa gtg aat ggg agg tat aaa gtg gtg agg atg tct ttt gct	531
Arg Asp Lys Val Asn Gly Arg Tyr Lys Val Val Arg Met Ser Phe Ala	
160 165 170	
ttc tgg aga gtt agg caa gag gag cct gtg gtg gaa tgt ggt gtt ctt	579
Phe Trp Arg Val Arg Gln Glu Glu Pro Val Val Glu Cys Gly Val Leu	
175 180 185	
gat gtt gat act ggt gaa tgg cgg aag ctg agt cca cct cct tat gtg	627
Asp Val Asp Thr Gly Glu Trp Arg Lys Leu Ser Pro Pro Pro Tyr Val	
190 195 200	
gtc aat gtg gga agc aaa tcg gta tgc gtg aat gga tct atc tac tgg	675
Val Asn Val Gly Ser Lys Ser Val Cys Val Asn Gly Ser Ile Tyr Trp	
205 210 215 220	
tta cac att cag acg gtt tac aga ata cta gcc ttg gat ctt cac aaa	723
Leu His Ile Gln Thr Val Tyr Arg Ile Leu Ala Leu Asp Leu His Lys	
225 230 235	
caa gag ttt cat aaa gtc cca gtg ccg cct acg cag atc act gtg gac	771
Gln Glu Phe His Lys Val Pro Val Pro Pro Thr Gln Ile Thr Val Asp	
240 245 250	
act cag atg gtg aac ctt gag gac cgt ctc gta ctt gct ata acc aga	819
Thr Gln Met Val Asn Leu Glu Asp Arg Leu Val Leu Ala Ile Thr Arg	
255 260 265	
gtt agc cct gaa tgg ata cta gag gta tgg gcc atg gat aca tac aaa	867
Val Ser Pro Glu Trp Ile Leu Glu Val Trp Gly Met Asp Thr Tyr Lys	
270 275 280	
gaa aaa tgg agc aag act tac tcc ata agt ttg gat cat aga gtt gtt	915
Glu Lys Trp Ser Lys Thr Tyr Ser Ile Ser Leu Asp His Arg Val Val	
285 290 295 300	
tcc tgg cga agg cag aaa agg tgg ttc acg ccc gtg gca gtt tct aag	963
Ser Trp Arg Arg Gln Lys Arg Trp Phe Thr Pro Val Ala Val Ser Lys	
305 310 315	
caa gcg aat ctt gtc ttc tat gac aat aag aag agg cta ttc aaa tat	1011
Gln Ala Asn Leu Val Phe Tyr Asp Asn Lys Lys Arg Leu Phe Lys Tyr	
320 325 330	

-continued

tat cca gtg aaa gat gag att cgt tgt ctc tcc tta gac att tgt gtt	1059
Tyr Pro Val Lys Asp Glu Ile Arg Cys Leu Ser Leu Asp Ile Cys Val	
335 340 345	
ctg tct cct tac gtg gaa aac ttg gtc cct ctt ccg tta aag cca agc	1107
Leu Ser Pro Tyr Val Glu Asn Leu Val Pro Leu Pro Leu Lys Pro Ser	
350 355 360	
cat ccg cat cct act ccg aaa aat tca gat ttt gaa atg agg ata tca	1155
His Pro His Pro Thr Pro Lys Asn Ser Asp Phe Glu Met Arg Ile Ser	
365 370 375 380	
aga tgc cgc ttg ttt tcg acg cca ggt tct tgg ata tcc aaa att ttg	1203
Arg Cys Arg Leu Phe Ser Thr Pro Gly Ser Trp Ile Ser Lys Ile Leu	
385 390 395	
aaa tgg aat gtt atg act cta gag att ttg ttt acc tct cta gca ata	1251
Lys Trp Asn Val Met Thr Leu Glu Ile Leu Phe Thr Ser Leu Ala Ile	
400 405 410	
gtt ggt tat ata tgc tta cct ctc tag gagttattta tcttgtttca	1298
Val Gly Tyr Ile Cys Leu Pro Leu *	
415 420	
aaatattagt tggttatata tgctttcgag atcttttctga taaa	1342

<210> SEQ ID NO 4
 <211> LENGTH: 420
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana
 <400> SEQUENCE: 4

Met Gly Glu Glu Glu Glu Asn Pro Asn Ser Ile Asp Ile Leu Pro Glu	
1 5 10 15	
Leu Leu Glu Glu Val Leu Leu Arg Leu Pro Thr Lys Ser Ile Leu Lys	
20 25 30	
Cys Arg Ile Val Ser Lys Gln Trp Arg Ser Leu Leu Glu Ser Ser Arg	
35 40 45	
Phe Ala Glu Arg His Met Ser Leu Gln Asn Ser Arg Arg Arg Ile Leu	
50 55 60	
Ala Ala Tyr Asn Cys Asp Cys Gly Gly Arg Arg Lys Leu Leu Pro Glu	
65 70 75 80	
Ser Arg Phe Glu Gly Asp Glu Glu Ile Val Tyr Leu His Cys Asp Ala	
85 90 95	
Ser Arg Pro Ser Met Thr Cys Gln Gly Val Ile Cys Phe Pro Glu Gln	
100 105 110	
Asp Trp Ile Ile Val Leu Asn Pro Ser Thr Ser Gln Leu Arg Arg Phe	
115 120 125	
Pro Ser Gly Leu Asn His Asn Cys Arg Phe Arg Ile Gly Leu Trp Lys	
130 135 140	
Thr Phe Ser Pro Gly Asn Trp Val Met Gly Phe Gly Arg Asp Lys Val	
145 150 155 160	
Asn Gly Arg Tyr Lys Val Val Arg Met Ser Phe Ala Phe Trp Arg Val	
165 170 175	
Arg Gln Glu Glu Pro Val Val Glu Cys Gly Val Leu Asp Val Asp Thr	
180 185 190	
Gly Glu Trp Arg Lys Leu Ser Pro Pro Pro Tyr Val Val Asn Val Gly	
195 200 205	
Ser Lys Ser Val Cys Val Asn Gly Ser Ile Tyr Trp Leu His Ile Gln	
210 215 220	

-continued

Thr	Val	Tyr	Arg	Ile	Leu	Ala	Leu	Asp	Leu	His	Lys	Gln	Glu	Phe	His
225					230					235					240
Lys	Val	Pro	Val	Pro	Pro	Thr	Gln	Ile	Thr	Val	Asp	Thr	Gln	Met	Val
			245						250					255	
Asn	Leu	Glu	Asp	Arg	Leu	Val	Leu	Ala	Ile	Thr	Arg	Val	Ser	Pro	Glu
			260					265					270		
Trp	Ile	Leu	Glu	Val	Trp	Gly	Met	Asp	Thr	Tyr	Lys	Glu	Lys	Trp	Ser
	275						280					285			
Lys	Thr	Tyr	Ser	Ile	Ser	Leu	Asp	His	Arg	Val	Val	Ser	Trp	Arg	Arg
	290					295					300				
Gln	Lys	Arg	Trp	Phe	Thr	Pro	Val	Ala	Val	Ser	Lys	Gln	Ala	Asn	Leu
305					310					315					320
Val	Phe	Tyr	Asp	Asn	Lys	Lys	Arg	Leu	Phe	Lys	Tyr	Tyr	Pro	Val	Lys
			325						330					335	
Asp	Glu	Ile	Arg	Cys	Leu	Ser	Leu	Asp	Ile	Cys	Val	Leu	Ser	Pro	Tyr
			340					345					350		
Val	Glu	Asn	Leu	Val	Pro	Leu	Pro	Leu	Lys	Pro	Ser	His	Pro	His	Pro
	355					360						365			
Thr	Pro	Lys	Asn	Ser	Asp	Phe	Glu	Met	Arg	Ile	Ser	Arg	Cys	Arg	Leu
	370					375					380				
Phe	Ser	Thr	Pro	Gly	Ser	Trp	Ile	Ser	Lys	Ile	Leu	Lys	Trp	Asn	Val
385					390					395					400
Met	Thr	Leu	Glu	Ile	Leu	Phe	Thr	Ser	Leu	Ala	Ile	Val	Gly	Tyr	Ile
			405						410					415	
Cys	Leu	Pro	Leu												
			420												

What is claimed is:

1. A transgenic plant comprising a recombinant polynucleotide having a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from those of SEQ ID NOs:2, 4, or a complementary nucleotide sequence thereof;
- (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);
- (c) one of SEQ ID NOs: 1, 3, or a complementary nucleotide sequence thereof;
- (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
- (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
- (f) a nucleotide sequence comprising at least 15 consecutive nucleotides outside of a conserved domain of any of (a)-(e);
- (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide that modifies a plant's trait;

- (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
- (i) a nucleotide sequence having at least 60% sequence identity to a nucleotide sequence of any of (a)-(g);
- (j) a nucleotide sequence having at least 95% sequence identity to a nucleotide sequence of any of (a)-(g);
- (k) a nucleotide sequence which encodes a polypeptide having at least 31% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (l) a nucleotide sequence which encodes a polypeptide having at least 60% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (m) a nucleotide sequence which encodes a polypeptide having at least 75% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (n) a nucleotide sequence which encodes a polypeptide having at least 95% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (o) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least 86% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;

- (p) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least about 90% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (q) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least about 95% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (r) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least about 98% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (s) a nucleotide sequence which encodes a polypeptide having at least 31% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4;
- (t) a nucleotide sequence which encodes a polypeptide having at least 60% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4;
- (u) a nucleotide sequence which encodes a polypeptide having at least 75% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4; and
- (v) a nucleotide sequence which encodes a polypeptide having at least 95% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4;

wherein the plant possesses an altered trait as compared to a wild type plant, or the plant exhibits an altered phenotype as compared to a wild type plant, or the plant expresses an altered level of one or more genes associated with a plant trait as compared to a wild type plant.

2. The transgenic plant of claim 1, further comprising a constitutive, inducible, or tissue-specific promoter operably linked to said nucleotide sequence.

3. The transgenic plant of claim 1, wherein the plant is selected from the group consisting of: soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, pumpkin, spinach, squash, sweet corn, tobacco, tomato, watermelon, mint and other labiates, rosaceous fruits, and vegetable brassicas.

4. An isolated or recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from those of SEQ ID NOs:2, 4, or a complementary nucleotide sequence thereof;
- (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);
- (c) one of SEQ ID NOs: 1, 3, or a complementary nucleotide sequence thereof;
- (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
- (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);

- (f) a nucleotide sequence comprising at least 15 consecutive nucleotides outside of a conserved domain of any of (a)-(e);
- (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide that modifies a plant's trait;
- (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
- (i) a nucleotide sequence having at least 60% sequence identity to a nucleotide sequence of any of (a)-(g);
- (j) a nucleotide sequence having at least 95% sequence identity to a nucleotide sequence of any of (a)-(g);
- (k) a nucleotide sequence which encodes a polypeptide having at least 31% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (l) a nucleotide sequence which encodes a polypeptide having at least 60% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (m) a nucleotide sequence which encodes a polypeptide having at least 75% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (n) a nucleotide sequence which encodes a polypeptide having at least 95% sequence identity outside of a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (o) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least 86% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (p) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least about 90% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (q) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least about 95% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (r) a nucleotide sequence encoding a polypeptide having an amino acid domain with at least about 98% sequence identity to a conserved domain of a polypeptide of SEQ ID NOs:2 or 4;
- (s) a nucleotide sequence which encodes a polypeptide having at least 31% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4;
- (t) a nucleotide sequence which encodes a polypeptide having at least 60% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4;
- (u) a nucleotide sequence which encodes a polypeptide having at least 75% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4; and
- (v) a nucleotide sequence which encodes a polypeptide having at least 95% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4.

5. The isolated or recombinant polynucleotide of claim 4, further comprising a constitutive, inducible, or tissue-specific promoter operably linked to the nucleotide sequence.

6. An isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4.

7. A method of using the isolated or recombinant polynucleotide of claim 4 for producing a plant having a modified trait, the method comprising selecting a polynucleotide that encodes a polypeptide, inserting the polynucleotide into an expression vector, introducing the vector into a plant or a cell of a plant to overexpress the polypeptide, thereby producing a modified plant, and selecting for a modified trait.

8. The transgenic plant of claim 1, wherein the trait is selected from the group consisting of: enhanced tolerance to freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved tolerance to pest infestations, decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals; improved growth under poor photoconditions.

9. The transgenic plant of claim 1, wherein the trait is an alteration in the level of one or more of the compounds selected from the group consisting of: taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prenyllipids, glucosinolates, and terpenoids.

10. The transgenic plant of claim 1, wherein the trait is an alteration in one or more physical characteristics selected from the group consisting of: the number of trichomes, fruit and seed size and number, yield of stems, leaves, or roots, stability of seeds during storage, susceptibility of the seed to shattering, root hair length and quantity, internode distances, or the quality of seed coat.

11. The transgenic plant of claim 1, wherein the trait is an alteration in a plant growth characteristic selected from the group consisting of: growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time, flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, apical dominance, branching patterns, number of organs, organ identity, and organ shape or size.

12. The transgenic plant of claim 1, wherein the trait is an alteration in one or more characteristics selected from the group consisting of protein or oil production, seed protein or oil production, insoluble sugar level, soluble sugar level, and starch composition.

13. The method of claim 7, wherein the trait is selected from the group consisting of: enhanced tolerance to freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved tolerance to pest infestations, decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals, improved growth under poor photoconditions.

14. The method of claim 7, wherein the trait is an alteration in the level of one or more of the compounds selected from the group consisting of: taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prenyllipids, glucosinolates, and terpenoids.

15. The method of claim 7, wherein the trait is an alteration in one or more physical characteristics selected from the group consisting of: the number of trichomes, fruit and seed size and number, yield of stems, leaves, or roots, stability of seeds during storage, susceptibility of the seed to shattering, root hair length and quantity, internode distances, or the quality of seed coat.

16. The method of claim 7, wherein the trait is an alteration in a plant growth characteristic selected from the group consisting of: growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time, flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, apical dominance, branching patterns, number of organs, organ identity, and organ shape or size.

17. The method of claim 7, wherein the trait is an alteration in one or more characteristics selected from the group consisting of protein or oil production, seed protein or oil production, insoluble sugar level, soluble sugar level, and starch composition.

18. A plant produced by the method of claim 13.

19. A plant produced by the method of claim 14.

20. A plant produced by the method of claim 15.

21. A plant produced by the method of claim 16.

22. A plant produced by the method of claim 17.

23. A method of using the isolated or recombinant polynucleotide of claim 4 for producing a plant having a modified trait, the method comprising selecting a polynucleotide that when expressed produces an antisense nucleic acid sequence, inserting the polynucleotide into an expression vector, introducing the vector into a plant or a cell of a plant to express the antisense nucleic acid, thereby producing a modified plant, and selecting for a modified trait.

24. The method of claim 23, wherein the trait is selected from the group consisting of: enhanced tolerance to freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved tolerance to pest infestations, decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals, improved growth under poor photoconditions.

25. The method of claim 23, wherein the trait is an alteration in the level of one or more of the compounds selected from the group consisting of: taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prenyllipids, glucosinolates, and terpenoids.

26. The method of claim 23, wherein the trait is an alteration in one or more physical characteristics selected from the group consisting of: the number of trichomes, fruit and seed size and number, yield of stems, leaves, or roots, stability of seeds during storage, susceptibility of the seed to shattering, root hair length and quantity, internode distances, or the quality of seed coat.

27. The method of claim 23, wherein the trait is an alteration in a plant growth characteristic selected from the group consisting of: growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time, flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, apical dominance, branching patterns, number of organs, organ identity, and organ shape or size.

28. The method of claim 23, wherein the trait is an alteration in one or more characteristics selected from the

group consisting of protein or oil production, seed protein or oil production, insoluble sugar level, soluble sugar level, and starch composition.

29. A plant produced by the method of claim 24.

30. A plant produced by the method of claim 25.

31. A plant produced by the method of claim 26.

32. A plant produced by the method of claim 27.

33. A plant produced by the method of claim 28.

34. An isolated or recombinant polypeptide comprising a subsequence of at least about 10 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4, wherein the contiguous amino acids are outside of a conserved domain.

35. An isolated or recombinant polypeptide comprising a subsequence of at least about 20 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4, wherein the contiguous amino acids are outside of a conserved domain.

36. An isolated or recombinant polypeptide comprising a subsequence of at least about 30 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4, wherein the contiguous amino acids are outside of a conserved domain.

37. An isolated or recombinant polypeptide comprising a subsequence of at least about 10 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4, wherein the contiguous amino acids are within a conserved domain.

38. An isolated or recombinant polypeptide comprising a subsequence of at least about 20 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4, wherein the contiguous amino acids are within a conserved domain.

39. An isolated or recombinant polypeptide comprising a subsequence of at least about 30 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4, wherein the contiguous amino acids are within a conserved domain.

40. A polypeptide having at least 31% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4, or the length of the polypeptide itself.

41. A polypeptide having at least 60% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4, or the length of the polypeptide itself.

42. A polypeptide having at least 75% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4, or the length of the polypeptide itself.

43. A polypeptide having at least 95% sequence identity over the entire length of a polypeptide of SEQ ID NOs:2 or 4, or the length of the polypeptide itself.

* * * * *