

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017299673 B2**

- (54) Title
Anti- GPRC5D antibodies, bispecific antigen binding molecules that bind GPRC5D and CD3, and uses thereof
- (51) International Patent Classification(s)
C07K 16/28 (2006.01) **A61P 35/02** (2006.01)
A61K 39/395 (2006.01)
- (21) Application No: **2017299673** (22) Date of Filing: **2017.07.20**
- (87) WIPO No: **WO18/017786**
- (30) Priority Data
- | | | |
|-------------------|-------------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 62/364,811 | 2016.07.20 | US |
- (43) Publication Date: **2018.01.25**
(44) Accepted Journal Date: **2024.08.01**
- (71) Applicant(s)
Janssen Biotech, Inc.
- (72) Inventor(s)
Attar, Ricardo;Chin, Diana;Edavettal, Suzanne;Gaudet, Francois;Li, Yingzhe;Luistro, Leopoldo;Majewski, Nathan;Mendonca, Mark;Pillarisetti, Kondandaram;Teplyakov, Alexey;Tornetta, Mark
- (74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU
- (56) Related Art
WO 2016/090329 A2
WO 2016/036937 A1



(51) International Patent Classification:

C07K 16/28 (2006.01) *A61P 35/02* (2006.01)
A61K 39/395 (2006.01)

(21) International Application Number:

PCT/US2017/042982

(22) International Filing Date:

20 July 2017 (20.07.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/364,811 20 July 2016 (20.07.2016) US

(71) Applicant: **JANSSEN PHARMACEUTICA NV**
[BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).


(72) Inventor; and

(71) Applicant: **ATTAR, Ricardo** [US/US]; 1400 McKean Road, Spring House, Pennsylvania 19477 (US).

(72) Inventors: **CHIN, Diana**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **EDAVETTAL, Suzanne**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **GAUDET, Francois**; 3210 Merryfield Row, La Jolla, 92121 (UA). **LI, Yingzhe**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **LUISTRO, Leopoldo**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **MAJEWSKI, Nathan**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **MEN-DONCA, Mark**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **PILLARISETTI, Kondandaram**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **TEPLYAKOV, Alexey**; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). **TORNETTA, Mark**;

(54) Title: ANTI- GPRC5D ANTIBODIES, BISPECIFIC ANTIGEN BINDING MOLECULES THAT BIND GPRC5D AND CD3, AND USES THEREOF

Figure 1



(57) Abstract: Provided herein are antibodies that specifically bind to GPRC5D. Also described are related polynucleotides capable of encoding the provided GPRC5D-specific antibodies or antigen-binding fragments, cells expressing the provided antibodies or antigen-binding fragments, as well as associated vectors and detectably labeled antibodies or antigen-binding fragments. In addition, methods of using the provided antibodies are described. For example, the provided antibodies may be used to diagnose, treat, or monitor GPRC5D-expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with GPRC5D-expressing cancer and thus may be amenable to treatment with a GPRC5D-specific anti-cancer therapeutic, such as the multispecific antibodies against GPRC5D and CD3 described herein.



1400 McKean Road, Spring House, Pennsylvania 19477
(US).

(74) **Agent:** SHIRTZ, Joseph F. et al.; JOHNSON & JOHNSON, One Johnson & Johnson Plaza, New Brunswick, New Jersey 08933 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

(88) **Date of publication of the international search report:**

01 March 2018 (01.03.2018)

**ANTI- GPRC5D ANTIBODIES, BISPECIFIC ANTIGEN BINDING MOLECULES
THAT BIND GPRC5D AND CD3, AND USES THEREOF**

5 Cross-reference to Related Applications

This application claims the benefit of United States Provisional Application Serial Number 62/364,811 filed 20 July 2016. The entire content of the aforementioned application is incorporated herein by reference in its entirety.

10 Sequence Listing

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on June 28, 2017, is named PRD3422USNP_SL.txt and is 57,673 bytes in size.

Technical Field

15 The disclosure provided herein relates to monoclonal antibodies that specifically bind G protein-coupled receptor class C group 5 member D (GPRC5D), multispecific antibodies specifically bind GPRC5D and cluster determinant 3 (CD3), and methods of producing and using the described antibodies.

20 Background

Multiple myeloma (MM) is the second most common hematological malignancy and constitutes 2% of all cancer deaths. MM is a heterogenous disease and caused by mostly by chromosome translocations inter alia t(11 ; 14),t(4; 14),t(8;14),del(13),del(17) (Drach et al., (1998) Blood 92(3):802-809; Gertz et al., (2005) Blood 106(8):2837-2840; Facon et al., (2001) **25** Blood 97(6): 1566-1571). MM-affected patients may experience a variety of disease-related symptoms due to, bone marrow infiltration, bone destruction, renal failure, immunodeficiency, and the psychosocial burden of a cancer diagnosis. As of 2006, the 5-year relative survival rate for MM was approximately 34% highlighting that MM is a difficult-to-treat disease where there are currently no curative options.

G-protein coupled receptor, class C, group 5, member D (GPRC5D) is an orphan, atypical, class C GPCR first identified in 2001 (Bräuner-Osborne et al. *Biochim Biophys Acta*. 1518(3):237-248, 2001). GPRC5D and other group 5 GPCRs have unusually short amino-terminal domains for class C receptors, and are therefore, predicted to be conformationally similar to class A receptors. In this regard they are unique, with sequence homology to class C GPCRs and predicted structural topology comparable to class A receptors. Functional consequence of GPRC5D activation has not been described and the ligand remains unknown. The gene has three exons and is located on chromosome 12p13.3 in humans. GPRC5D receptor is highly conserved among various species and shares 92% identity with cynomolgus monkey GPRC5D.

GPRC5D mRNA is predominantly expressed in all malignant plasma cells from MM patients (Atamaniuk JA et al. *Eur J Clin Invest* 42(9) 953-960; 2012; Frigyesi-blood and Cohen, et al. *Hematology* 18(6): 348-35; 2013). GPRC5D expression is variable among the patients and correlate well with plasma cell burden and genetic aberrations such as Rb-1 deletion (Atamaniuk JA et al. *Eur J Clin Invest* 42(9) 953-960; 2012).

This exclusive expression of GPRC5D on the plasma-cell lineage designates it as an ideal target for antimyeloma antibodies.

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

Unless the context clearly requires otherwise, throughout the description and the claims, the words “comprise”, “comprising”, and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of “including, but not limited to”.

Summary

In one aspect, the present disclosure provides an isolated antibody, or an antigen-binding fragment thereof, that binds specifically to GPRC5D comprising: a heavy chain complementary

determining region 1 (CDR1) comprising the amino acid sequence of SEQ ID NO: 4, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 8, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 12; and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 15, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 18, and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 21

In another aspect, the present disclosure provides an isolated cell expressing an antibody, or antigen-binding fragment of the invention.

In another aspect, the present disclosure provides an isolated GPRC5D x CD3 bispecific antibody comprising: a) a first heavy chain (HC1); b) a second heavy chain (HC2); c) a first light chain (LC1); and d) a second light chain (LC2), wherein the HC1 and the LC1 pair to form a first antigen-binding site that specifically binds CD3, and the HC2 and the LC2 pair to form a second antigen-binding site that specifically binds GPRC5D, wherein the HC1 comprises a VH region comprising the amino acid sequence of SEQ ID NO: 25 and the LC1 comprises a VL region comprising the amino acid sequence of SEQ ID NO: 26, and wherein the HC2 comprises a VH region comprising the amino acid sequence of SEQ ID NO: 55 and the LC2 comprises a VL region comprising the amino acid sequence of SEQ ID NO: 58.

In another aspect, the present disclosure provides an isolated cell expressing the GPRC5D x CD3 bispecific antibody, of the invention.

In another aspect, the present disclosure provides a method for treating a subject having cancer, said method comprising: administering a therapeutically effective amount of a GPRC5D x CD3 bispecific antibody the invention to a patient in need thereof for a time sufficient to treat the cancer.

In another aspect, the present disclosure provides a method for inhibiting growth or proliferation of cancer cells, said method comprising: administering a therapeutically effective amount of a GPRC5D x CD3 bispecific antibody of the invention to inhibit the growth or proliferation of cancer cells.

In another aspect, the present disclosure provides a method of redirecting a T cell to a GPRC5D-expressing cancer cell, said method comprising: administering a therapeutically effective amount of a GPRC5D x CD3 bispecific antibody of the invention to redirect a T cell to a cancer.

In another aspect, the present disclosure provides a pharmaceutical composition comprising a GPRC5D x CD3 bispecific antibody of the invention and a pharmaceutically acceptable carrier.

In another aspect, the present disclosure provides a method for generating a GPRC5D x CD3 bispecific antibody of the invention by culturing a cell of the invention.

In another aspect, the present disclosure provides one or more isolated synthetic polynucleotides encoding the HC1, the HC2, the LC1, and the LC2 of the isolated GPRC5D x CD3 bispecific antibody of the invention.

In another aspect, the present disclosure provides a kit comprising a GPRC5D x CD3 bispecific antibody as described herein and/or one or more polynucleotide(s) as described herein and packaging for the same.

In another aspect, the present disclosure provides an isolated antibody that binds specifically to GPRC5D comprising: a heavy chain comprising a variable heavy (VH) chain region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain comprising a variable light (VL) chain region comprising the amino acid sequence of SEQ ID NO: 58, wherein the antibody is an IgG4 isotype.

In another aspect, the present disclosure provides an isolated GPRC5D x CD3 bispecific antibody comprising: a GPRC5D binding portion comprising: a heavy chain comprising a variable heavy (VH) chain region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain comprising a variable light (VL) chain region comprising the amino acid sequence of SEQ ID NO: 58; and a CD3 binding portion comprising a heavy chain comprising SEQ ID NO: 99 and a light chain comprising SEQ ID NO: 100, wherein the antibody is an IgG4 isotype.

In another aspect, the present disclosure provides use of a GPRC5D x CD3 bispecific antibody of the invention in the manufacture of a medicament for the treatment of a subject having cancer.

Provided herein are antibodies that specifically bind to GPRC5D and antigen-binding fragments thereof. Also described are related polynucleotides capable of encoding the provided GPRC5D -specific antibodies and antigen-binding fragments, cells expressing the provided antibodies and antigen-binding fragments, as well as associated vectors and detectably labeled antibodies and antigen-binding fragments. In addition, methods of using the provided antibodies and antigen-binding fragments are described. For example, the GPRC5D -specific antibodies

and antigen-binding fragments may be used to diagnose or monitor GPRC5D -expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with GPRC5D -expressing cancer and thus may be amenable to treatment with a GPRC5D -specific anti-cancer therapeutic, such as the multispecific antibodies against GPRC5D and CD3 described herein.

Further provided herein are multispecific antibodies that immunospecifically bind to GPRC5D and CD3 and multispecific antigen-binding fragments thereof. Also described are related polynucleotides capable of encoding the provided GPRC5D x CD3-multispecific antibodies, cells expressing the provided antibodies, as well as associated vectors and detectably labeled multispecific antibodies. In addition, methods of using the provided multispecific antibodies are described. For example, the GPRC5D x CD3-multispecific antibodies may be used to diagnose or monitor GPRC5D -expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with GPRC5D -expressing cancer and thus may be amenable to treatment with a GPRC5D -specific anti-cancer therapeutic, such as the GPRC5D x CD3-multispecific antibodies described herein.

GPRC5D -Specific Antibodies

Described herein are isolated antibodies and antigen-binding fragments specific for GPRC5D. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments bind human GPRC5D. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments bind human GPRC5D and cynomolgus monkey GPRC5D. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments bind to one or more residues of a polypeptide having the amino acid sequence of SEQ ID NO. 22. This GPRC5D -specific antibody or antigen-binding fragment may induce ADCC *in vitro* with an EC₅₀ of 28 nM or less.

Table 1 provides a summary of examples of some GPRC5D-specific antibodies described herein:

Table 1. CDR sequences of mAbs generated against human GPRC5D

ID	HC-CDR1	HC-CDR2	HC-CDR3	LC-CDR1	LC-CDR2	LC-CDR3
GC5B81	SYAIS (SEQ ID NO 1)	GIPIFGTANYAQKFQG (SEQ ID NO 5)	ESRWRGYKLD (SEQ ID NO 9)	RASQSISSYLN (SEQ ID NO 13)	AASSLQS (SEQ ID NO 16)	QQSYSTPLT (SEQ ID NO 19)
GC5B465 GC5B597	NYWMS (SEQ ID NO 2)	GISYSGGSKYYASSVKG (SEQ ID NO 6)	AAFDGRRRAVRLD (SEQ ID NO 10)	RASQSISSYLN (SEQ ID NO 13)	AASSLQS (SEQ ID NO 16)	QQSYSTPLT (SEQ ID NO 19)
GC5B483 GC5B599	SYFIG (SEQ ID NO 3)	IIYPGKSDTRYSPSFQG (SEQ ID NO 7)	VYSFGGRHKALFDY (SEQ ID NO 11)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B596	GYTMN (SEQ ID NO 4)	LINPYNSDTNYAQLQG (SEQ ID NO 8)	VALRVALDY (SEQ ID NO 12)	KASQNVATHVG (SEQ ID NO 15)	SASYRYS (SEQ ID NO 18)	QQYNRYPYT (SEQ ID NO 21)
GC5B382	DYGMH	AIKYSGGSTYYADSVKG	RAESGPGLDY	KSSQSVLYSSNNKNYLA	WASTRES	QQYYSTPLT

	(SEQ ID NO 61)	(SEQ ID NO 67)	(SEQ ID NO 72)	(SEQ ID NO 98)	(SEQ ID NO 78)	(SEQ ID NO 80)
GC5B379	NYWMS (SEQ ID NO 2)	GISYSGGSKYYADSVKG (SEQ ID NO 28)	AAWDFGRRVRLDY (SEQ ID NO 30)	RASQSISSYLN (SEQ ID NO 13)	AASSLQS (SEQ ID NO 16)	QQSYSTPLT (SEQ ID NO 19)
GC5B373	SYWIG (SEQ ID NO 27)	IYPGDS DTRYSPSFQG (SEQ ID NO 29)	IGFYGRSFRIDY (SEQ ID NO 73)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B376	SYWIG (SEQ ID NO 27)	IYPGDS DTRYSPSFQG (SEQ ID NO 29)	VYSFGGRHKALFDY (SEQ ID NO 11)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B385	GYAMS (SEQ ID NO 62)	AISGSGGSTYYADSVKG (SEQ ID NO 68)	VDRSFGRSRYTLDY (SEQ ID NO 74)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B370 GC5B598	SYGIS (SEQ ID NO 63)	GIPIFGNINYAQKFQG (SEQ ID NO 69)	VSRRFKRFAYYFDY (SEQ ID NO 75)	KSSQSVLYSSNNKNYLA (SEQ ID NO 98)	WASTRES (SEQ ID NO 78)	QQYYSTPLT (SEQ ID NO 80)
GC5B602	GYSFTGYTMN (SEQ ID NO 64)	LINPYNGDTN (SEQ ID NO 70)	VALRVALDY (SEQ ID NO 12)	KASQNVATHVG (SEQ ID NO 15)	SASYRYS (SEQ ID NO 18)	QQYNRYPYT (SEQ ID NO 21)
GC5B603	SYAMS (SEQ ID NO 65)	AISGSGGSTYYADSVKG (SEQ ID NO 68)	SNFLPVVFDY (SEQ ID NO 76)	RASQSVRKSLA (SEQ ID NO 95)	TASNRAT (SEQ ID NO 79)	QQYFRAPIT (SEQ ID NO 81)
GC5B601	GFSLTSYNVH (SEQ ID NO 66)	VIWAGGSTNYNSALMS (SEQ ID NO 71)	DGIRLRFAY (SEQ ID NO 77)	KASQNVATHVG (SEQ ID NO 15)	SASYRYS (SEQ ID NO 18)	QQYNRYPYT (SEQ ID NO 21)

In some embodiments are provided a GPRC5D -specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1. In some embodiments are provided a GPRC5D-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1 and a light chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1. In some embodiments described herein, the GPRC5D-specific antibody or antigen-binding fragment thereof competes for binding to GPRC5D with an antibody or antigen-binding fragment thereof comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1 and a light chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1.

The IgG class is divided in four isotypes: IgG1, IgG2, IgG3 and IgG4 in humans. They share more than 95% homology in the amino acid sequences of the Fc regions but show major differences in the amino acid composition and structure of the hinge region. The Fc region mediates effector functions, such as antibody-dependent cellular cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC). In ADCC, the Fc region of an antibody binds to Fc receptors (FcγRs) on the surface of immune effector cells such as natural killers and macrophages, leading to the phagocytosis or lysis of the targeted cells. In CDC, the antibodies kill the targeted cells by triggering the complement cascade at the cell surface. The antibodies

described herein include antibodies with the described features of the variable domains in combination with any of the IgG isotypes, including modified versions in which the Fc sequence has been modified to effect different effector functions.

For many applications of therapeutic antibodies, Fc-mediated effector functions are not part of the mechanism of action. These Fc-mediated effector functions can be detrimental and potentially pose a safety risk by causing off-mechanism toxicity. Modifying effector functions can be achieved by engineering the Fc regions to reduce their binding to FcγRs or the complement factors. The binding of IgG to the activating (FcγRI, FcγRIIa, FcγRIIIa and FcγRIIIb) and inhibitory (FcγRIIb) FcγRs or the first component of complement (C1q) depends on residues located in the hinge region and the CH2 domain. Mutations have been introduced in IgG1, IgG2 and IgG4 to reduce or silence Fc functionalities. The antibodies described herein may include these modifications.

In one embodiment, the antibody comprises an Fc region with one or more of the following properties: (a) reduced effector function when compared to the parent Fc; (b) reduced affinity to FcγRI, FcγRIIa, FcγRIIb, FcγRIIIb and/or FcγRIIIa, (c) reduced affinity to FcγRI (d) reduced affinity to FcγRIIa (e) reduced affinity to FcγRIIb, (f) reduced affinity to FcγRIIIb or (g) reduced affinity to FcγRIIIa.

In some embodiments, the antibodies or antigen-binding fragments are IgG, or derivatives thereof, e.g., IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the antibody has an IgG1 isotype, the antibody contains L234A, L235A, and/or K409R substitution(s) in its Fc region. In some embodiments wherein the antibody has an IgG4 isotype, the antibody contains S228P, L234A, and L235A substitutions in its Fc region. The antibodies described herein may include these modifications.

In addition to the described GPRC5D-specific antibodies and antigen-binding fragments, also provided are polynucleotide sequences capable of encoding the described antibodies and antigen-binding fragments. Vectors comprising the described polynucleotides are also provided, as are cells expressing the GPRC5D-specific antibodies or antigen-binding fragments provided herein. Also described are cells capable of expressing the disclosed vectors. These cells may be mammalian cells (such as 293F cells, CHO cells), insect cells (such as Sf7 cells), yeast cells, plant cells, or bacteria cells (such as E. coli). The described antibodies may also be produced by hybridoma cells.

Methods of using GPRC5D-Specific Antibodies

Methods of using the described GPRC5D-specific antibodies or antigen-binding fragments are also disclosed. Particular antibodies for use in the methods discussed in this section include those with the set of CDRs described for antibodies in Table 1. For example, these antibodies or antigen-binding fragments may be useful in treating cancer, by interfering with GPRC5D-receptor interactions or where the antibody is conjugated to a toxin, so targeting the toxin to the GPRC5D-expressing cancer. Further, these antibodies or antigen-binding fragments may be useful for detecting the presence of GPRC5D in a biological sample, such as blood or serum; for quantifying the amount of GPRC5D in a biological sample, such as blood or serum; for diagnosing GPRC5D-expressing cancer; determining a method of treating a subject afflicted with cancer; or monitoring the progression of GPRC5D-expressing cancer in a subject. In some embodiments, GPRC5D-expressing cancer may be a lymphoma, such as multiple myeloma (MM). The described methods may be carried out before the subject receives treatment for GPRC5D-expressing cancer, such as treatment with a multispecific antibody against GPRC5D and CD3. Furthermore, the described methods may be carried out after the subject receives treatment for GPRC5D-expressing cancer, such as treatment with a multispecific antibody against GPRC5D and CD3 described herein.

The described methods of detecting GPRC5D in a biological sample include exposing the biological sample to one or more of the GPRC5D-specific antibodies or antigen-binding fragments described herein.

The described methods of diagnosing GPRC5D-expressing cancer in a subject also involve exposing the biological sample to one or more of the GPRC5D-specific antibodies or antigen-binding fragments described herein; however, the methods also include quantifying the amount of GPRC5D present in the sample; comparing the amount of GPRC5D present in the sample to a known standard or reference sample; and determining whether the subject's GPRC5D levels fall within the levels of GPRC5D associated with cancer.

Also described herein are methods of monitoring GPRC5D-expressing cancer in a subject. The described methods include exposing the biological sample to one or more of the GPRC5D-specific antibodies or antigen-binding fragments described herein; quantifying the amount of GPRC5D present in the sample that is bound by the antibody, or antigen-binding

fragment thereof; comparing the amount of GPRC5D present in the sample to either a known standard or reference sample or the amount of GPRC5D in a similar sample previously obtained from the subject; and determining whether the subject's GPRC5D levels are indicative of cancer progression, regression or stable disease based on the difference in the amount of GPRC5D in the compared samples.

The samples obtained, or derived from, subjects are biological samples such as urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated, tissues, surgically resected tumor tissue, biopsies, fine needle aspiration samples, or histological preparations.

The described GPRC5D-specific antibodies or antigen-binding fragments may be labeled for use with the described methods, or other methods known to those skilled in the art. For example, the antibodies described herein, or antigen-binding fragments thereof, may be labeled with a radiolabel, a fluorescent label, an epitope tag, biotin, a chromophore label, an ECL label, an enzyme, ruthenium, ¹¹¹In-DOTA, ¹¹¹In- diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase and beta-galactosidase, or poly-histidine or similar such labels known in the art.

GPRC5D-Specific Antibody Kits

Described herein are kits including the disclosed GPRC5D-specific antibodies or antigen-binding fragments thereof. The described kits may be used to carry out the methods of using the GPRC5D-specific antibodies or antigen-binding fragments provided herein, or other methods known to those skilled in the art. In some embodiments the described kits may include the antibodies or antigen-binding fragments described herein and reagents for use in detecting the presence of GPRC5D in a biological sample. Accordingly, the described kits may include one or more of the antibodies, or an antigen-binding fragment(s) thereof, described herein and a vessel for containing the antibody or fragment when not in use, instructions for use of the antibody or fragment, the antibody or fragment affixed to a solid support, and/or detectably labeled forms of the antibody or fragment, as described herein.

GPRC5D x CD3-Multispecific Antibodies

The redirection of T-lymphocytes to MM cells expressing GPRC5D via the TCR/CD3 complex represents an attractive alternative approach. The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha (α)/beta (β) or TCR gamma (γ)/delta (δ) heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma (γ), delta (δ), epsilon (ϵ), zeta (ζ), and eta (η). Human CD3 ϵ is described under UniProt P07766 (CD3E_HUMAN). An anti CD3 ϵ antibody described in the state of the art is SP34 (Yang SJ, The Journal of Immunology (1986) 137: 1097-1100). SP34 reacts with both primate and human CD3. SP34 is available from Pharmingen. A further anti CD3 antibody described in the state of the art is UCHT-1 (see WO2000041474). A further anti CD3 antibody described in the state of the art is BC-3 (Fred Hutchinson Cancer Research Institute; used in Phase I/II trials of GvHD, Anasetti et al., Transplantation 54: 844 (1992)). SP34 differs from UCHT-1 and BC-3 in that SP-34 recognizes an epitope present on solely the ϵ chain of CD3 (see Salmeron et al., (1991) J. Immunol. 147: 3047) whereas UCHT-1 and BC-3 recognize an epitope contributed by both the ϵ and γ chains. The sequence of an antibody with the same sequence as of antibody SP34 is mentioned in WO2008119565, WO2008119566, WO2008119567, WO2010037836, WO2010037837 and WO2010037838. A sequence which is 96% identical to VH of antibody SP34 is mentioned in US8236308 (WO2007042261).

Described herein are isolated multispecific antibodies that bind GPRC5D and CD3 ("GPRC5D x CD3 multispecific antibodies") and multispecific antigen-binding fragments thereof. In some embodiments an isolated antibody, or an antigen-binding fragment thereof, that binds immunospecifically to GPRC5D is provided.

In some embodiments, the GPRC5D-specific arm of the multispecific antibody binds human GPRC5D and cynomolgus monkey GPRC5D. In some embodiments, the GPRC5D-specific arm of the GPRC5D x CD3-multispecific antibodies or antigen-binding fragments binds the extracellular domain of human GPRC5D. In preferred embodiments, the GPRC5D x CD3 multispecific antibody or antigen-binding fragment is a bispecific antibody or antigen-binding fragment. In some embodiments, an isolated GPRC5D x CD3 bispecific antibody comprising: a) a first heavy chain (HC1); b) a second heavy chain (HC2); c) a first light chain (LC1); and d) a second light chain (LC2), wherein the HC1 and the LC1 pair to form a first antigen-binding site that immunospecifically binds GPRC5D, and the HC2 and the LC2 pair to form a second antigen-binding site that immunospecifically binds CD3, or a GPRC5D x CD3-bispecific binding

fragment thereof is provided. In another embodiment, an isolated cell expressing the antibody or bispecific binding fragment is provided. In some embodiments, the GPRC5D-binding arm (or “GPRC5D-specific arm”) of the GPRC5D x CD3 multispecific antibody is derived from a GPRC5D antibody described herein (for example, from an antibody having the CDR sequences listed in Table 1).

In some embodiments, the GPRC5D-specific arm of the GPRC5D x CD3-multispecific antibodies or antigen-binding fragments are IgG, or derivatives thereof. In some embodiments, the CD3-binding arm (or “CD3-specific arm”) of the GPRC5D x CD3 multispecific antibody is derived from the mouse monoclonal antibody SP34, a mouse IgG3/lambda isotype. (K.R. Abhinandan and A. C. Martin, 2008. Mol. Immunol. 45, 3832-3839). In some embodiments, the CD3-binding arm of the GPRC5D x CD3 multispecific antibody comprises one VH domain and one VL domain selected from Table 2.

Table 2. Heavy chains and light chains of the CD3-specific antibody and antigen-binding fragment.

VH	VL
CD3B219 (SEQ ID NO: 99): EVQLVESGGGLVQPGGSLRLSCAASGFTFN TYAMNWRQAPGKGLEWVARIRSKYNNYAT YYAASVKGRFTISRDDSKNSLYLQMNSLKTE DTAVYYCARHGNFGNSYVSWFAYWGQGT LTVSSASTKGPSVFPLAPCSRSTSESTAALG CLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGKTYTCNV DHKPSNTKVDKRVESKYGPPCPPCPAPEAAGGP SVFLFPPKPKDTLMISRTPEVTCVVDVSQED PEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPS SIEKTISKAKGQPREPQVYTLPPSQEEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYK TTPPVLDSDGSFLLYSKLTVDKSRWQEGNVF SCSVMHEALHNHYTQKSLSLSLGK	CD3B219 (SEQ ID NO: 100): QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYA NWWQQKPGQAPRGLIGGTNKRAPGTPARFSGSLL GGKAALTLSGVQPEDEAEYYCALWYSNLWFGG GTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVC LISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSN NKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVE KTVAPTECS

The IgG class is divided in four isotypes: IgG1, IgG2, IgG3 and IgG4 in humans. They share more than 95% homology in the amino acid sequences of the Fc regions but show major differences in the amino acid composition and structure of the hinge region. The Fc region mediates effector functions, such as antibody-dependent cellular cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC). In ADCC, the Fc region of an antibody binds to Fc receptors (FcγRs) on the surface of immune effector cells such as natural killers and macrophages, leading to the phagocytosis or lysis of the targeted cells. In CDC, the antibodies kill the targeted cells by triggering the complement cascade at the cell surface.

For many applications of therapeutic antibodies, Fc-mediated effector functions are not part of the mechanism of action. These Fc-mediated effector functions can be detrimental and potentially pose a safety risk by causing off-mechanism toxicity. Modifying effector functions can be achieved by engineering the Fc regions to reduce their binding to FcγRs or the complement factors. The binding of IgG to the activating (FcγRI, FcγRIIa, FcγRIIIa and FcγRIIIb) and inhibitory (FcγRIIb) FcγRs or the first component of complement (C1q) depends on residues located in the hinge region and the CH2 domain. Mutations have been introduced in IgG1, IgG2 and IgG4 to reduce or silence Fc functionalities.

In one embodiment, the antibody comprises an Fc region with one or more of the following properties: (a) reduced effector function when compared to the parent Fc; (b) reduced affinity to FcγRI, FcγRIIa, FcγRIIb, FcγRIIIb and/or FcγRIIIa, (c) reduced affinity to FcγRI (d) reduced affinity to FcγRIIa (e) reduced affinity to FcγRIIb, (f) reduced affinity to FcγRIIIb or (g) reduced affinity to FcγRIIIa.

In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG, or a derivative thereof. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG1, or a derivative thereof. In some embodiments, for example, the Fc region of the CD3-specific IgG1 antibody from which the CD3-binding arm is derived comprises L234A, L235A, and F405L substitutions in its Fc region. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG4, or a derivative thereof. In some embodiments, for example, the Fc region of the CD3-specific IgG4 antibody from which the CD3-binding arm is derived comprises S228P, L234A, L235A, F405L, and

R409K substitutions in its Fc region. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived binds CD3 ϵ on primary human T cells and/or primary cynomolgus T cells. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived activates primary human CD4⁺ T cells and/or primary cynomolgus CD4⁺ T cells.

In addition to the described GPRC5D x CD3-multispecific antibodies, also provided are polynucleotide sequences capable of encoding the described GPRC5D x CD3-multispecific antibodies. In some embodiments, an isolated synthetic polynucleotide encoding the HC1, the HC2, the LC1 or the LC2 of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment is provided. Vectors comprising the described polynucleotides are also provided, as are cells expressing the GPRC5D x CD3-multispecific antibodies provided herein. Also described are cells capable of expressing the disclosed vectors. These cells may be mammalian cells (such as 293F cells, CHO cells), insect cells (such as Sf7 cells), yeast cells, plant cells, or bacteria cells (such as E. coli). The described antibodies may also be produced by hybridoma cells. In some embodiments, methods for generating the GPRC5D x CD3 bispecific antibody or bispecific binding fragment by culturing cells is provided.

Further provided herein are pharmaceutical compositions comprising the GPRC5D x CD3 multispecific antibodies or antigen-binding fragments and a pharmaceutically acceptable carrier.

Methods of using GPRC5D x CD3-Multispecific Antibodies

Methods of using the described GPRC5D x CD3-multispecific antibodies and multispecific antigen-binding fragments thereof are also disclosed. For example, the GPRC5D x CD3-multispecific antibodies and multispecific antigen-binding fragments thereof may be useful in the treatment of a GPRC5D-expressing cancer in a subject in need thereof. In some embodiments, the GPRC5D-expressing cancer is a lymphoma, such as multiple myeloma.

The described methods of treating GPRC5D-expressing cancer in a subject in need thereof include administering to the subject a therapeutically effective amount of a described GPRC5D x CD3-multispecific antibody or multispecific antigen-binding fragment thereof. In some embodiments, the subject is a mammal, preferably a human. In preferred embodiments are

provided methods for treating a subject having cancer by administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody or bispecific antigen-binding fragment to a patient in need thereof for a time sufficient to treat the cancer.

Further provided herein are methods for inhibiting growth or proliferation of cancer cells by administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment to inhibit the growth or proliferation of cancer cells.

Also provided herein are methods of redirecting a T cell to a GPRC5D-expressing cancer cell by administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment to redirect a T cell to a cancer.

GPRC5D x CD3-Specific Antibody Kits

Described herein are kits including the disclosed GPRC5D x CD3-multispecific antibodies. The described kits may be used to carry out the methods of using the GPRC5D x CD3-multispecific antibodies provided herein, or other methods known to those skilled in the art.

In some embodiments the described kits may include the antibodies described herein and reagents for use in treating a GPRC5D-expressing cancer. Accordingly, the described kits may include one or more of the multispecific antibodies, or a multispecific antigen-binding fragment(s) thereof, described herein and a vessel for containing the antibody or fragment when not in use, and/or instructions for use of the antibody or fragment, the antibody or fragment affixed to a solid support, and/or detectably labeled forms of the antibody or fragment, as described herein.

Brief Description of the Drawings

Fig. 1: Concentration dependent binding profile of selected anti-GPRC5D mAbs against human GPRC5D and non-transfected HEK 293 cells. Three mAbs, GC5B36, GC5B168, and GC5B205 were also observed to bind to the non-transfected (GPRC5D null) HEK293 cells.

Fig. 2. Dose dependent binding of anti-GPCR5D x CD3 bispecific antibodies to human GPRC5D HEK293F cells.

Fig. 3. The binding profile of the bispecific antibodies on MM1R and H929 cells was compared with the overexpressed human GPCR5D HEK293 cells and non-transfected HEK293 cells using FACS.

5 **Fig. 4A and 4B.** T-cell mediated cytotoxicity and T cell activation of anti-GPCR5D x CD3 antibodies against human GPCR5D-expressing cells.

Fig. 5A and 5B T-cell mediated cytotoxicity and T cell activation of anti-GPCR5D x CD3 antibodies against cyno GPCR5D-expressing cells.

10

Fig. 6A and 6B. GPCR5DxCD3 Abs efficiently kill H929 cells in prophylactic NSG mice model. GCDB32 (6A) and GCDB35 (6B) resulted in complete tumor growth inhibition at the 10 ug doses, with GCDB32 also capable of 100% tumor growth inhibition at the 1 ug dose.

15 **Fig. 7A and 7B.** Comparison of potency in the absence (7A) and presence (7B) of Fc block.

Fig. 8A-8D. Binding of GCDB32 GCDB35, GCDB40 and GCDB43 to Fcγ receptors.

Fig. 9A and 9B. FACS binding assessment of hybridoma derived mAbs.

20

Fig. 10A and 10B. T-cell mediated cytotoxicity of GPCR5D x CD3 bispecific antibodies against H929 cells.

25 **Fig. 11A-11E.** GPCR5D x CD3 bispecific antibody binding to GPCR5D positive (H929, MM1R, LP1, OPM2) and negative cell lines (NALM6).

Fig. 12A-12D. GPCR5DxCD3 bispecific Abs are potent tumor inhibitors in vivo. All GPCR5D x CD3 bispecific Abs (GCDB32 shown in 12A, GCDB53 shown in 12B, GCDB61 shown in 12C, and GCDB72 shown in 12D) completely inhibited multiple myeloma cell (H929) tumor growth at the 10 ug and 1 ug doses. Differentiation was observed at the 0.1 ug dose with GCDB72 observed to have 80% tumor growth inhibition

30

Fig. 13: GPRC5D⁺ MM1.R cell lines were stained for 60 minutes with various concentrations of lead antibodies to measure the surface binding profiles (n=3). Phycoerythrin labelled human IgG4Fc was used as a secondary antibody to capture the signal (Southern Biotech, clone HP6025). Binding is expressed as normalized geometrical mean fluorescence intensity as determined by FACS. Graphing and fitting of data were done in GraphPad Prism 6 using nonlinear regression with variable slope (four parameters) and least square fit method.

Fig. 14A and 14B. GPRC5D⁺ cell lines were stained for 60 minutes with various concentrations of FAB6300 and GC5M481 antibodies to measure the surface binding profiles. Phycoerythrin labelled human IgG4Fc was used as a secondary antibody to capture the signal (Southern Biotech, clone HP6025). Binding is expressed as histograms (black line for the isotype and the red line denotes for the specific GPRC5D antibody (Fig. 14A). Fig. 14B shows the binding pattern of the lead molecules on GPRC5D⁺ multiple myeloma cell lines. Shaded, dotted line indicates isotype control and the solid line indicate lead molecule binding.

Fig. 15A and 15B. Frozen bone marrow-derived mononuclear cells from two different MM patients were used to assess GPRC5D x CD3 bispecific antibody binding, compared to IgG4 isotype control, plasma cell cytotoxicity and T-cell activation. For the cytotoxicity assay, T cells from the normal healthy donor were exogenously added to patient BM MNC samples and incubated with four lead molecules for 48 hours. GPRC5D x CD3 bispecific antibody binds to plasma cells in a dose dependent manner to all donor samples and the mean fluorescence intensities were recorded on the Y-axis. Note the loss of live plasma cells (CD138⁺) and the concomitant upregulation of CD25 on T cells in response to GPRC5D x CD3 bispecific antibody treatment.

Fig. 16. NSG mice were subcutaneously implanted with MM.1S human Multiple Myeloma cells on Day 0. Human PBMC were intravenously inoculated on study day 7. PBS, GCDB72 (0.1 µg, 1 µg, 10 µg and 50 µg/animal (equivalent to 0.005, 0.05, 0.5 and 2.5 mg/kg, respectively)) and Null control antibodies were intravenously dosed on Days 15, 18, 22, 24, 29, 32 and 36. Subcutaneous tumors were measured twice weekly and the results were presented as the mean

tumor volume, expressed $\text{mm}^3 \pm \text{SEM}$ of each group. GCDB72 antibody treatment when dosed at 1 μg (0.05 mg/kg) significantly inhibited sc tumor growth compared to PBS (TGI=64%, $p \leq 0.0001$). GCDB72 doses of 10 $\mu\text{g}/\text{animal}$ (0.5 mg/kg) and 50 $\mu\text{g}/\text{animal}$ (2.5 mg/kg) completely regressed tumor growth ($p \leq 0.0001$). The Null control antibodies had negligible or no effect. Statistical significance was evaluated using a 2-way ANOVA with multiple comparisons using Tukey's multiple comparisons test using Graph Pad Prism software (version 6). Differences between groups were considered significant when the probability value (p) was ≤ 0.05 .

Fig. 17. NSG were subcutaneously implanted with MM.1S human Multiple Myeloma cells on Day 0. Human PBMC were intravenously inoculated on study day 7. PBS, GCDB72 (0.1 μg , 1 μg , 10 μg and 50 $\mu\text{g}/\text{animal}$ (equivalent to 0.005, 0.05, 0.5 and 2.5 mg/kg, respectively)) and Null control antibodies were intravenously dosed on Days 15, 18, 22, 24, 29, 32 and 36. Body weight is presented as absolute body weight from start of treatment to the end of study

Detailed Description of Illustrative Embodiments

Definitions

Various terms relating to aspects of the description are used throughout the specification and claims. Such terms are to be given their ordinary meaning in the art unless otherwise indicated. Other specifically defined terms are to be construed in a manner consistent with the definitions provided herein.

As used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to "a cell" includes a combination of two or more cells, and the like.

The term "about" as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of up to $\pm 10\%$ from the specified value, as such variations are appropriate to perform the disclosed methods. Unless otherwise indicated, all numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." Accordingly, unless indicated

to the contrary, the numerical parameters set forth in the following specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least
5 be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain
10 errors necessarily resulting from the standard deviation found in their respective testing measurements.

“Isolated” means a biological component (such as a nucleic acid, peptide or protein) has been substantially separated, produced apart from, or purified away from other biological components of the organism in which the component naturally occurs, i.e., other chromosomal
15 and extrachromosomal DNA and RNA, and proteins. Nucleic acids, peptides and proteins that have been “isolated” thus include nucleic acids and proteins purified by standard purification methods. “Isolated” nucleic acids, peptides and proteins can be part of a composition and still be isolated if such composition is not part of the native environment of the nucleic acid, peptide, or protein. The term also embraces nucleic acids, peptides and proteins prepared by recombinant
20 expression in a host cell as well as chemically synthesized nucleic acids. An “isolated” antibody or antigen-binding fragment, as used herein, is intended to refer to an antibody or antigen-binding fragment which is substantially free of other antibodies or antigen-binding fragments having different antigenic specificities (for instance, an isolated antibody that specifically binds to GPRC5D is substantially free of antibodies that specifically bind antigens other than
25 GPRC5D). An isolated antibody that specifically binds to an epitope, isoform or variant of GPRC5D may, however, have cross-reactivity to other related antigens, for instance from other species (such as GPRC5D species homologs).

“Polynucleotide,” synonymously referred to as “nucleic acid molecule,” “nucleotides” or “nucleic acids,” refers to any polyribonucleotide or polydeoxyribonucleotide, which may be
30 unmodified RNA or DNA or modified RNA or DNA. “Polynucleotides” include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-

stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both
5 RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, “polynucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found
10 in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short nucleic acid chains, often referred to as oligonucleotides.

The meaning of “substantially the same” can differ depending on the context in which the term is used. Because of the natural sequence variation likely to exist among heavy and light
15 chains and the genes encoding them, one would expect to find some level of variation within the amino acid sequences or the genes encoding the antibodies or antigen-binding fragments described herein, with little or no impact on their unique binding properties (e.g., specificity and affinity). Such an expectation is due in part to the degeneracy of the genetic code, as well as to the evolutionary success of conservative amino acid sequence variations, which do not
20 appreciably alter the nature of the encoded protein. Accordingly, in the context of nucleic acid sequences, “substantially the same” means at least 65% identity between two or more sequences. Preferably, the term refers to at least 70% identity between two or more sequences, more preferably at least 75% identity, more preferably at least 80% identity, more preferably at least 85% identity, more preferably at least 90% identity, more preferably at least 91% identity, more
25 preferably at least 92% identity, more preferably at least 93% identity, more preferably at least 94% identity, more preferably at least 95% identity, more preferably at least 96% identity, more preferably at least 97% identity, more preferably at least 98% identity, and more preferably at least 99% or greater identity. The percent identity between two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical
30 positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The percent

identity between two nucleotide or amino acid sequences may e.g. be determined using the algorithm of E. Meyers and W. Miller, Comput. Appl. Biosci 4, 11-17 (1988) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences may be determined using the Needleman and Wunsch, J. Mol. Biol. 48, 444-453 (1970) algorithm.

The degree of variation that may occur within the amino acid sequence of a protein without having a substantial effect on protein function is much lower than that of a nucleic acid sequence, since the same degeneracy principles do not apply to amino acid sequences.

Accordingly, in the context of an antibody or antigen-binding fragment, “substantially the same” means antibodies or antigen-binding fragments having 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to the antibodies or antigen-binding fragments described. Other embodiments include GPRC5D specific antibodies, or antigen-binding fragments, that have framework, scaffold, or other non-binding regions that do not share significant identity with the antibodies and antigen-binding fragments described herein, but do incorporate one or more CDRs or other sequences needed to confer binding that are 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to such sequences described herein. A “vector” is a replicon, such as plasmid, phage, cosmid, or virus in which another nucleic acid segment may be operably inserted so as to bring about the replication or expression of the segment.

A “clone” is a population of cells derived from a single cell or common ancestor by mitosis. A “cell line” is a clone of a primary cell that is capable of stable growth in vitro for many generations. In some examples provided herein, cells are transformed by transfecting the cells with DNA.

The terms “express” and “produce” are used synonymously herein, and refer to the biosynthesis of a gene product. These terms encompass the transcription of a gene into RNA. These terms also encompass translation of RNA into one or more polypeptides, and further encompass all naturally occurring post-transcriptional and post-translational modifications. The expression or production of an antibody or antigen-binding fragment thereof may be within the cytoplasm of the cell, or into the extracellular milieu such as the growth medium of a cell culture.

The terms “treating” or “treatment” refer to any success or indicia of success in the attenuation or amelioration of an injury, pathology or condition, including any objective or subjective parameter such as abatement, remission, diminishing of symptoms or making the condition more tolerable to the patient, slowing in the rate of degeneration or decline, making the final point of degeneration less debilitating, improving a subject’s physical or mental well-being, or prolonging the length of survival. The treatment may be assessed by objective or subjective parameters; including the results of a physical examination, neurological examination, or psychiatric evaluations.

An “effective amount” or “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result. A therapeutically effective amount of a GPRC5D x CD3 antibody may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects.

“Antibody” refers to all isotypes of immunoglobulins (IgG, IgA, IgE, IgM, IgD, and IgY) including various monomeric, polymeric and chimeric forms, unless otherwise specified. Specifically encompassed by the term “antibody” are polyclonal antibodies, monoclonal antibodies (mAbs), and antibody-like polypeptides, such as chimeric antibodies and humanized antibodies.

“Antigen-binding fragments” are any proteinaceous structure that may exhibit binding affinity for a particular antigen. Antigen-binding fragments include those provided by any known technique, such as enzymatic cleavage, peptide synthesis, and recombinant techniques. Some antigen-binding fragments are composed of portions of intact antibodies that retain antigen-binding specificity of the parent antibody molecule. For example, antigen-binding fragments may comprise at least one variable region (either a heavy chain or light chain variable region) or one or more CDRs of an antibody known to bind a particular antigen. Examples of suitable antigen-binding fragments include, without limitation diabodies and single-chain molecules as well as Fab, F(ab’)2, Fc, Fabc, and Fv molecules, single chain (Sc) antibodies, individual antibody light chains, individual antibody heavy chains, chimeric fusions between antibody chains or CDRs and other proteins, protein scaffolds, heavy chain monomers or dimers,

light chain monomers or dimers, dimers consisting of one heavy and one light chain, a monovalent fragment consisting of the VL, VH, CL and CH1 domains, or a monovalent antibody as described in WO2007059782, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge region, a Fd fragment consisting essentially of the V.sub.H and C.sub.H1 domains; a Fv fragment consisting essentially of the VL and VH domains of a single arm of an antibody, a dAb fragment (Ward et al., Nature 341, 544-546 (1989)), which consists essentially of a VH domain and also called domain antibodies (Holt et al; Trends Biotechnol. 2003 Nov.; 21(11):484-90); camelid or nanobodies (Revets et al; Expert Opin Biol Ther. 2005 Jan.; 5(1):111-24); an isolated complementarity determining region (CDR), and the like. All antibody isotypes may be used to produce antigen-binding fragments. Additionally, antigen-binding fragments may include non-antibody proteinaceous frameworks that may successfully incorporate polypeptide segments in an orientation that confers affinity for a given antigen of interest, such as protein scaffolds. Antigen-binding fragments may be recombinantly produced or produced by enzymatic or chemical cleavage of intact antibodies. The phrase “an antibody or antigen-binding fragment thereof” may be used to denote that a given antigen-binding fragment incorporates one or more amino acid segments of the antibody referred to in the phrase.

The terms “CDR”, and its plural “CDRs”, refer to a complementarity determining region (CDR) of which three make up the binding character of a light chain variable region (CDRL1, CDRL2 and CDRL3) and three make up the binding character of a heavy chain variable region (CDRH1, CDRH2 and CDRH3). CDRs contribute to the functional activity of an antibody molecule and are separated by amino acid sequences that comprise scaffolding or framework regions. The exact definitional CDR boundaries and lengths are subject to different classification and numbering systems. CDRs may therefore be referred to by Kabat, Chothia, contact or any other boundary definitions. Despite differing boundaries, each of these systems has some degree of overlap in what constitutes the so called “hypervariable regions” within the variable sequences. CDR definitions according to these systems may therefore differ in length and boundary areas with respect to the adjacent framework region. See for example Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th ed. NIH Publication No. 91-3242 (1991); Chothia et al., “Canonical Structures For the Hypervariable Regions of Immunoglobulins,” *J. Mol. Biol.* 196:901 (1987); and MacCallum et al., “Antibody-Antigen

Interactions: Contact Analysis and Binding Site Topography,” *J. Mol. Biol.* 262:732 (1996)), each of which is hereby incorporated by reference in its entirety.

Typically, CDRs form a loop structure that can be classified as a canonical structure. The term “canonical structure” refers to the main chain conformation that is adopted by the antigen binding (CDR) loops. From comparative structural studies, it has been found that five of the six antigen binding loops have only a limited repertoire of available conformations. Each canonical structure can be characterized by the torsion angles of the polypeptide backbone. Correspondent loops between antibodies may, therefore, have very similar three dimensional structures, despite high amino acid sequence variability in most parts of the loops (Chothia et al., “Canonical Structures For the Hypervariable Regions of Immunoglobulins,” *J. Mol. Biol.* 196:901 (1987); Chothia et al., “Conformations of Immunoglobulin Hypervariable Regions,” *I* 342:877 (1989); Martin and Thornton, “Structural Families in Loops of Homologous Proteins: Automatic Classification, Modelling and Application to Antibodies,” *J. Mol. Biol.* 263:800 (1996), each of which is incorporated by reference in its entirety). Furthermore, there is a relationship between the adopted loop structure and the amino acid sequences surrounding it. The conformation of a particular canonical class is determined by the length of the loop and the amino acid residues residing at key positions within the loop, as well as within the conserved framework (i.e., outside of the loop). Assignment to a particular canonical class can therefore be made based on the presence of these key amino acid residues.

The term “polypeptide” is used interchangeably with the term “protein” and in its broadest sense refers to a compound of two or more subunit amino acids, amino acid analogs or peptidomimetics. The subunits may be linked by peptide bonds. In another embodiment, the subunit may be linked by other bonds, e.g., ester, ether, etc. As used herein the term “amino acid” refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D and L optical isomers, amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is commonly called a polypeptide or a protein.

“Specifically binds” or “binds specifically” or derivatives thereof when used in the context of antibodies, or antibody fragments, represents binding via domains encoded by immunoglobulin genes or fragments of immunoglobulin genes to one or more epitopes of a protein of interest, without preferentially binding other molecules in a sample containing a mixed

population of molecules. Typically, an antibody binds to a cognate antigen with a K_d of less than about 1×10^{-8} M, as measured by a surface plasmon resonance assay or a cell-binding assay. Phrases such as “[antigen]-specific” antibody (e.g., GPRC5D-specific antibody) are meant to convey that the recited antibody specifically binds the recited antigen.

5 Polynucleotide,” synonymously referred to as “nucleic acid molecule,” “nucleotides” or “nucleic acids,” refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. “Polynucleotides” include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. 10 “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, “polynucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short nucleic acid chains, often referred to as 15 oligonucleotides.

A “vector” is a replicon, such as plasmid, phage, cosmid, or virus in which another nucleic acid segment may be operably inserted so as to bring about the replication or expression of the segment.

As used herein, the term “host cell” can be any type of cell, e.g., a primary cell, a cell in 25 culture, or a cell from a cell line. In specific embodiments, the term “host cell” refers to a cell transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule, e.g., due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome. The terms 30 “expression” and “production” are used synonymously herein, and refer to the biosynthesis of a gene product. These terms encompass the transcription of a gene into RNA. These terms also

encompass translation of RNA into one or more polypeptides, and further encompass all naturally occurring post-transcriptional and post-translational modifications. The expression or production of an antibody or antigen-binding fragment thereof may be within the cytoplasm of the cell, or into the extracellular milieu such as the growth medium of a cell culture. The meaning of “substantially the same” can differ depending on the context in which the term is used. Because of the natural sequence variation likely to exist among heavy and light chains and the genes encoding them, one would expect to find some level of variation within the amino acid sequences or the genes encoding the antibodies or antigen-binding fragments described herein, with little or no impact on their unique binding properties (e.g., specificity and affinity). Such an expectation is due in part to the degeneracy of the genetic code, as well as to the evolutionary success of conservative amino acid sequence variations, which do not appreciably alter the nature of the encoded protein. Accordingly, in the context of nucleic acid sequences, “substantially the same” means at least 65% identity between two or more sequences. Preferably, the term refers to at least 70% identity between two or more sequences, more preferably at least 75% identity, more preferably at least 80% identity, more preferably at least 85% identity, more preferably at least 90% identity, more preferably at least 91% identity, more preferably at least 92% identity, more preferably at least 93% identity, more preferably at least 94% identity, more preferably at least 95% identity, more preferably at least 96% identity, more preferably at least 97% identity, more preferably at least 98% identity, and more preferably at least 99% or greater identity. The percent identity between two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The percent identity between two nucleotide or amino acid sequences may e.g. be determined using the algorithm of E. Meyers and W. Miller, *Comput. Appl. Biosci* 4, 11-17 (1988) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences may be determined using the Needleman and Wunsch, *J. Mol. Biol.* 48, 444-453 (1970) algorithm.

The degree of variation that may occur within the amino acid sequence of a protein without having a substantial effect on protein function is much lower than that of a nucleic acid

sequence, since the same degeneracy principles do not apply to amino acid sequences.

Accordingly, in the context of an antibody or antigen-binding fragment, “substantially the same” means antibodies or antigen-binding fragments having 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to the antibodies or antigen-binding fragments described. Other
5 embodiments include GPRC5D-specific antibodies, or antigen-binding fragments, that have framework, scaffold, or other non-binding regions that do not share significant identity with the antibodies and antigen-binding fragments described herein, but do incorporate one or more CDRs or other sequences needed to confer binding that are 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to such sequences described herein.

10 The term “subject” refers to human and non-human animals, including all vertebrates, e.g., mammals and non-mammals, such as non-human primates, mice, rabbits, sheep, dogs, cats, horses, cows, chickens, amphibians, and reptiles. In many embodiments of the described methods, the subject is a human.

15 The term “redirect” or “redirecting” as used herein refers to the ability of the GPRC5D x CD3 antibody to traffic the activity of T cells effectively, from its inherent cognate specificity toward reactivity against GPRC5D-expressing cells.

The term “sample” as used herein refers to a collection of similar fluids, cells, or tissues (e.g., surgically resected tumor tissue, biopsies, including fine needle aspiration), isolated from a
20 subject, as well as fluids, cells, or tissues present within a subject. In some embodiments the sample is a biological fluid. Biological fluids are typically liquids at physiological temperatures and may include naturally occurring fluids present in, withdrawn from, expressed or otherwise extracted from a subject or biological source. Certain biological fluids derive from particular tissues, organs or localized regions and certain other biological fluids may be more globally or
25 systemically situated in a subject or biological source. Examples of biological fluids include blood, serum and serosal fluids, plasma, lymph, urine, saliva, cystic fluid, tear drops, feces, sputum, mucosal secretions of the secretory tissues and organs, vaginal secretions, ascites fluids such as those associated with non-solid tumors, fluids of the pleural, pericardial, peritoneal, abdominal and other body cavities, fluids collected by bronchial lavage and the like. Biological
30 fluids may also include liquid solutions contacted with a subject or biological source, for example, cell and organ culture medium including cell or organ conditioned medium, lavage

fluids and the like. The term "sample," as used herein, encompasses materials removed from a subject or materials present in a subject.

A "known standard" may be a solution having a known amount or concentration of GPRC5D, where the solution may be a naturally occurring solution, such as a sample from a patient known to have early, moderate, late, progressive, or static cancer, or the solution may be a synthetic solution such as buffered water having a known amount of GPRC5D diluted therein. The known standards, described herein may include GPRC5D isolated from a subject, recombinant or purified GPRC5D protein, or a value of GPRC5D concentration associated with a disease condition.

As used herein, the terms "G-protein coupled receptor family C group 5 member D" and "GPRC5D" specifically include the human GPRC5D protein, for example as described in GenBank Accession No. BC069341, NCBI Reference Sequence: NP_061124.1 and UniProtKB/Swiss-Prot Accession No. Q9NZD1 (see also Brauner-Osborne, H. et al. 2001, *Biochim. Biophys. Acta* 1518, 237-248).

The term "CD3" refers to the human CD3 protein multi-subunit complex. The CD3 protein multi-subunit complex is composed to 6 distinctive polypeptide chains. These include a CD3 γ chain (SwissProt P09693), a CD3 δ chain (SwissProt P04234), two CD3 ϵ chains (SwissProt P07766), and one CD3 ζ chain homodimer (SwissProt 20963), and which is associated with the T cell receptor α and β chain. The term "CD3" includes any CD3 variant, isoform and species homolog which is naturally expressed by cells (including T cells) or can be expressed on cells transfected with genes or cDNA encoding those polypeptides, unless noted.

A "GPRC5D x CD3 antibody" is a multispecific antibody, optionally a bispecific antibody, which comprises two different antigen-binding regions, one of which binds specifically to the antigen GPRC5D and one of which binds specifically to CD3. A multispecific antibody can be a bispecific antibody, diabody, or similar molecule (see for instance *PNAS USA* 90(14), 6444-8 (1993) for a description of diabodies). The bispecific antibodies, diabodies, and the like, provided herein may bind any suitable target in addition to a portion of GPRC5D. The term "bispecific antibody" is to be understood as an antibody having two different antigen-binding regions defined by different antibody sequences. This can be understood as different target binding but includes as well binding to different epitopes in one target.

A “reference sample” is a sample that may be compared against another sample, such as a test sample, to allow for characterization of the compared sample. The reference sample will have some characterized property that serves as the basis for comparison with the test sample. For instance, a reference sample may be used as a benchmark for GPRC5D levels that are indicative of a subject having cancer. The reference sample does not necessarily have to be analyzed in parallel with the test sample, thus in some instances the reference sample may be a numerical value or range previously determined to characterize a given condition, such as GPRC5D levels that are indicative of cancer in a subject. The term also includes samples used for comparative purposes that are known to be associated with a physiologic state or disease condition, such as GPRC5D-expressing cancer, but that have an unknown amount of GPRC5D.

The term “progression,” as used in the context of progression of GPRC5D-expressing cancer, includes the change of a cancer from a less severe to a more severe state. This may include an increase in the number or severity of tumors, the degree of metastasis, the speed with which the cancer is growing or spreading, and the like. For example, “the progression of colon cancer” includes the progression of such a cancer from a less severe to a more severe state, such as the progression from stage I to stage II, from stage II to stage III, etc.

The term “regression,” as used in the context of regression of GPRC5D-expressing cancer, includes the change of a cancer from a more severe to a less severe state. This could include a decrease in the number or severity of tumors, the degree of metastasis, the speed with which the cancer is growing or spreading, and the like. For example, “the regression of colon cancer” includes the regression of such a cancer from a more severe to a less severe state, such as the progression from stage III to stage II, from stage II to stage I, etc.

The term “stable” as used in the context of stable GPRC5D-expressing cancer, is intended to describe a disease condition that is not, or has not, changed significantly enough over a clinically relevant period of time to be considered a progressing cancer or a regressing cancer.

The embodiments described herein are not limited to particular methods, reagents, compounds, compositions or biological systems, which can, of course, vary.

GPRC5D-Specific Antibodies and Antigen-Binding Fragments

Described herein are isolated monoclonal antibodies or antigen-binding fragments that specifically bind GPRC5D. The general structure of an antibody molecule comprises an antigen

binding domain, which includes heavy and light chains, and the Fc domain, which serves a variety of functions, including complement fixation and binding antibody receptors.

The described GPRC5D-specific antibodies or antigen-binding fragments include all isotypes, IgA, IgD, IgE, IgG and IgM, and synthetic multimers of the four-chain

immunoglobulin structure. The described antibodies or antigen-binding fragments also include the IgY isotype generally found in hen or turkey serum and hen or turkey egg yolk.

The GPRC5D-specific antibodies and antigen-binding fragments may be derived from any species by recombinant means. For example, the antibodies or antigen-binding fragments may be mouse, rat, goat, horse, swine, bovine, chicken, rabbit, camelid, donkey, human, or chimeric versions thereof. For use in administration to humans, non-human derived antibodies or antigen-binding fragments may be genetically or structurally altered to be less antigenic upon administration to a human patient.

In some embodiments, the antibodies or antigen-binding fragments are chimeric. As used herein, the term “chimeric” refers to an antibody, or antigen-binding fragment thereof, having at least some portion of at least one variable domain derived from the antibody amino acid sequence of a non-human mammal, a rodent, or a reptile, while the remaining portions of the antibody, or antigen-binding fragment thereof, are derived from a human.

In some embodiments, the antibodies are humanized antibodies. Humanized antibodies may be chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin sequence. The humanized antibody may include at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

The antibodies or antigen-binding fragments described herein can occur in a variety of forms, but will include one or more of the antibody CDRs shown in Table 1.

Described herein are isolated antibodies and antigen-binding fragments that immunospecifically bind to GPRC5D. In some embodiments, the GPRC5D-specific antibodies or antigen-binding fragments are human IgG, or derivatives thereof. While the GPRC5D-specific antibodies or antigen-binding fragments exemplified herein are human, the antibodies or antigen-binding fragments exemplified may be chimerized.

In some embodiments are provided a GPRC5D-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1. In some embodiments are provided a GPRC5D-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1 and a light chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 1, a heavy chain CDR2 comprising SEQ ID NO: 5, and a heavy chain CDR3 comprising SEQ ID NO: 9. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 1, a heavy chain CDR2 comprising SEQ ID NO: 5, a heavy chain CDR3 comprising SEQ ID NO: 9, a light chain CDR1 comprising SEQ ID NO: 13, a light chain CDR2 comprising SEQ ID NO: 16, and a light chain CDR3 comprising SEQ ID NO: 19. This GPRC5D-specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 52. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 52 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 56. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 2, a heavy chain CDR2 comprising SEQ ID NO: 6, and a heavy chain CDR3 comprising SEQ ID NO: 10. In some embodiments, the

GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 2, a heavy chain CDR2 comprising SEQ ID NO: 6, a heavy chain CDR3 comprising SEQ ID NO: 10, a light chain CDR1 comprising SEQ ID NO: 13, a light chain CDR2 comprising SEQ ID NO: 16, and a light chain CDR3 comprising SEQ ID NO: 19.

5 This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 53. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, 10 SEQ ID NO: 53 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 56. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments 15 comprise a heavy chain CDR1 comprising SEQ ID NO: 3, a heavy chain CDR2 comprising SEQ ID NO: 7, and a heavy chain CDR3 comprising SEQ ID NO: 11. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 3, a heavy chain CDR2 comprising SEQ ID NO: 7, a heavy chain CDR3 comprising SEQ ID NO: 11, a light chain CDR1 comprising SEQ ID NO: 14, a light 20 chain CDR2 comprising SEQ ID NO: 17, and a light chain CDR3 comprising SEQ ID NO: 20. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 54. In some embodiments, the GPRC5D-specific antibodies and antigen-binding 25 fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 54 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 57. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

30 In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 4, a heavy chain CDR2 comprising SEQ

ID NO: 8, and a heavy chain CDR3 comprising SEQ ID NO: 12. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 4, a heavy chain CDR2 comprising SEQ ID NO: 8, a heavy chain CDR3 comprising SEQ ID NO: 12, a light chain CDR1 comprising SEQ ID NO: 15, a light chain CDR2 comprising SEQ ID NO: 18, and a light chain CDR3 comprising SEQ ID NO: 21. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 55. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 55 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 58. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 61, a heavy chain CDR2 comprising SEQ ID NO: 67, and a heavy chain CDR3 comprising SEQ ID NO: 72. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 61, a heavy chain CDR2 comprising SEQ ID NO: 67, a heavy chain CDR3 comprising SEQ ID NO: 72, a light chain CDR1 comprising SEQ ID NO: 13, a light chain CDR2 comprising SEQ ID NO: 78, and a light chain CDR3 comprising SEQ ID NO: 80. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 82. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 82 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 92. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 2, a heavy chain CDR2 comprising SEQ ID NO: 28, and a heavy chain CDR3 comprising SEQ ID NO: 30. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 2, a heavy chain CDR2 comprising SEQ ID NO: 28, a heavy chain CDR3 comprising SEQ ID NO: 30, a light chain CDR1 comprising SEQ ID NO: 13, a light chain CDR2 comprising SEQ ID NO: 16, and a light chain CDR3 comprising SEQ ID NO: 19. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 83. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 83 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 56. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 27, a heavy chain CDR2 comprising SEQ ID NO: 29, and a heavy chain CDR3 comprising SEQ ID NO: 73. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 27, a heavy chain CDR2 comprising SEQ ID NO: 29, a heavy chain CDR3 comprising SEQ ID NO: 73, a light chain CDR1 comprising SEQ ID NO: 14, a light chain CDR2 comprising SEQ ID NO: 17, and a light chain CDR3 comprising SEQ ID NO: 20. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 84. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 84 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 57. The heavy chain variable domain and light chain variable domain of antibodies

discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 27, a heavy chain CDR2 comprising
5 SEQ ID NO: 29, and a heavy chain CDR3 comprising SEQ ID NO: 11. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 27, a heavy chain CDR2 comprising SEQ ID NO: 29 a heavy chain CDR3 comprising SEQ ID NO: 11, a light chain CDR1 comprising SEQ ID NO: 14, a light chain CDR2 comprising SEQ ID NO: 17, and a light chain CDR3 comprising SEQ ID NO: 20.
10 This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 85. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ
15 ID NO: 85 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 57. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments
20 comprise a heavy chain CDR1 comprising SEQ ID NO: 62, a heavy chain CDR2 comprising SEQ ID NO: 68, and a heavy chain CDR3 comprising SEQ ID NO: 74. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 62, a heavy chain CDR2 comprising SEQ ID NO: 68, a heavy chain CDR3 comprising SEQ ID NO: 74, a light chain CDR1 comprising SEQ ID NO: 14, a light
25 chain CDR2 comprising SEQ ID NO: 17, and a light chain CDR3 comprising SEQ ID NO: 20. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 86. In some embodiments, the GPRC5D-specific antibodies and antigen-binding
30 fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 86 and a light chain variable domain substantially the same as, or identical to, SEQ ID

NO: 57. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

5 In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 63, a heavy chain CDR2 comprising SEQ ID NO: 69, and a heavy chain CDR3 comprising SEQ ID NO: 75. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 63, a heavy chain CDR2 comprising SEQ ID NO: 69, a heavy chain CDR3 comprising SEQ ID NO: 75, a light chain CDR1 comprising SEQ ID NO: 13, a light
10 chain CDR2 comprising SEQ ID NO: 78, and a light chain CDR3 comprising SEQ ID NO: 80. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 87. In some embodiments, the GPRC5D-specific antibodies and antigen-binding
15 fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 87 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 92. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

20 In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 64, a heavy chain CDR2 comprising SEQ ID NO: 70, and a heavy chain CDR3 comprising SEQ ID NO: 12. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 64, a heavy chain CDR2 comprising SEQ ID NO: 70, a heavy chain
25 CDR3 comprising SEQ ID NO: 12, a light chain CDR1 comprising SEQ ID NO: 15, a light chain CDR2 comprising SEQ ID NO: 18, and a light chain CDR3 comprising SEQ ID NO: 21. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to,
30 SEQ ID NO: 88. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ

ID NO: 88 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 58. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

5 In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 65, a heavy chain CDR2 comprising SEQ ID NO: 68, and a heavy chain CDR3 comprising SEQ ID NO: 76. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 65, a heavy chain CDR2 comprising SEQ ID NO: 68, a heavy chain
10 CDR3 comprising SEQ ID NO: 76, a light chain CDR1 comprising SEQ ID NO: 95, a light chain CDR2 comprising SEQ ID NO: 79, and a light chain CDR3 comprising SEQ ID NO: 81. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to,
15 SEQ ID NO: 89. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 89 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 93. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is
20 an anti-GPRC5D arm.

In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 66, a heavy chain CDR2 comprising SEQ ID NO: 71, and a heavy chain CDR3 comprising SEQ ID NO: 77. In some embodiments, the GPRC5D-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1
25 comprising SEQ ID NO: 66, a heavy chain CDR2 comprising SEQ ID NO: 71, a heavy chain CDR3 comprising SEQ ID NO: 77, a light chain CDR1 comprising SEQ ID NO: 15, a light chain CDR2 comprising SEQ ID NO: 18, and a light chain CDR3 comprising SEQ ID NO: 21. This GPRC5D -specific antibody or antigen-binding fragment may comprise human framework sequences. In some embodiments, the GPRC5D -specific antibodies and antigen-binding
30 fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 91. In some embodiments, the GPRC5D-specific antibodies and antigen-binding

fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 91 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 94. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-GPRC5D arm.

In some embodiments, the antibodies or antigen-binding fragments are IgG, or derivatives thereof, e.g., IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the antibody is of IgG1 isotype, the antibody comprises an IgG1 Fc region (SEQ ID NO. 60).

SEQ ID NO. 60

10 ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS
GLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGG
PSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY
NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTKAKKGQPREPQVYTLPPSR
EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDK
15 SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

In some embodiments wherein the antibody is of IgG4 isotype, the antibody contains S228P, L234A, and L235A substitutions in its Fc region (SEQ ID NO. 59).

SEQ ID NO. 59

20 ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS
GLYSLSSVVTVPSSSLGTQTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEAAGGPS
VFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNS
TYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTKAKKGQPREPQVYTLPPSQEE
MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSR
25 WQEGNVFSCSVMHEALHNHYTQKSLSLSLGK

The specific antibodies defined by CDR and/or variable domain sequence discussed in the above paragraphs may include these modifications.

Also disclosed are isolated polynucleotides that encode the antibodies or antigen-binding fragments that immunospecifically bind to GPRC5D. The isolated polynucleotides capable of encoding the variable domain segments provided herein may be included on the same, or different, vectors to produce antibodies or antigen-binding fragments. An exemplary polynucleotide sequence that encodes a GPRC5D antibody is shown below:

35 Heavy Chain Sequence (SEQ ID NO: 96):

atggcctgggtctggaccctgctgtcctgatggccgctgccagagcatccaggcccaggtgcagctggtgcagagcggcg
ccgaggtgaagaagcccgccgagcgtgaaggtgagctgcaaggccagcggctacagcttcaccggctacaccatgaactgggtgcg

gcaggccccggccaggcctggagtggatgggcctgatcaacccctacaacagcgacaccaactacgcccagaagctgcaggggccg
 ggtgacctgaccaccgacaccagcaccagcaccgctacatggagctgcggagcctgcggagcgacgacaccgctgtactactgc
 ccccggtggccctgcgggtggccctggactactggggccaggccacctggtgaccgtgagcagcgccctccaccaaggcccatcc
 gtctccccctggcgccctgtctcaggagcacctccgagagcacagccgacctgggtgcctgggtcaaggactactccccgaaccggtg
 5 acggtgtcgtggaactcaggcgccctgaccagcggcgtgcacacctccccggtgtcctacagtcctcaggactctactccctcagcagcg
 tggtagaccgtgccctccagcagcttgggcacgaaaacctacacctgcaacgtagatcacaagcccagcaacaccaagggtggacaagaga
 gttgagtcctaatatggtcccccattgccaccatgccagcacctgaggccgcccgggggaccatcagttctctgttcccccaaaacca
 aggacactctcatgatctccggacccctgaggtcacgtgcgtgggtgggacgtgagccagggaagaccccaggtccagttcaactggt
 acgtggatggcgtggaggtgcataatgccaagacaaagccgcccggaggagcagttcaacagcacgtaccgtgtggtcagcgtctcacc
 10 gtctctgaccaggactggctgaacggcaaggagtacaagtgaagggtctccaacaaaggcctcccgtctccatcgagaaaaccatctcc
 aaagccaaagggcagccccgagagccacaggtgtacacctgcccccatcccaggaggagatgaccaagaaccaggtcagcctgacc
 tgctgtgtaaaaggcttctaccccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgctcc
 cgtgtggactccgacggctcttcttctctacagcaggctaaccgtggacaagagcaggtggcaggagggaatgtcttctcatgctcc
 gtgatgcatgaggctctgcacaaccactacacacagaagagcctctccctgtctctgggtaaatga

Light Chain Sequence (SEQ ID NO:90):

atgcgggtgtcggccagctgctgggactgctgctgtgcttccctggcgccagatgcgacatccagatgaccagagccccagcagc
 ctgagcgcagcgtggcgaccgggtgaccatcacctgcaaggccagccagaacgtggccaccacgtgggctggtaccagcagaag
 cccggcaaggcccccaagcggctgatctacagcgcagctaccgggtacagcggcgtgcccagccgggtcagcggcagcggcagcggc
 20 accgagttcacctgaccatcagcaacctgcagcccaggacttcgccactactactgccagcagtacaaccgggtaccctacaccttcg
 gccagggcaccaagctggagatcaagcgtacgggtgacctgtcttcatcttcccgcctatgatgagcagttgaaatctggaact
 gcctctgttgtgtcctgtgaataacttctatcccagagaggccaaagtacagtggaaaggtggataacgccctccaatcgggtaactccca
 ggagagtgctcacagagcaggacagcaaggacagcacctacagcctcagcagcaccctgacgtgagcaaagcagactacgagaaaca
 caaagtctacgcctgcgaagtcacccatcaggcgctgagctcgccgtcacaagagcttcaacaggggagagtgttga

Polynucleotides encoding recombinant antigen-binding proteins also are within the scope of the disclosure. In some embodiments, the polynucleotides described (and the peptides they encode) include a leader sequence. Any leader sequence known in the art may be employed.

The leader sequence may include, but is not limited to, a restriction site or a translation start site.

The GPRC5D-specific antibodies or antigen-binding fragments described herein include variants having single or multiple amino acid substitutions, deletions, or additions that retain the biological properties (e.g., binding affinity or immune effector activity) of the described GPRC5D-specific antibodies or antigen-binding fragments. In the context of the present invention the following notations are, unless otherwise indicated, used to describe a mutation; i) substitution of an amino acid in a given position is written as e.g. S228P which means a substitution of a Serine in position 228 with a Proline; and ii) for specific variants the specific three or one letter codes are used, including the codes Xaa and X to indicate any amino acid residue. Thus, the substitution of Serine for Arginine in position 228 is designated as: S228P, or the substitution of any amino acid residue for Serine in position 228 is designated as S228P. In

case of deletion of Serine in position 228 it is indicated by S228*. The skilled person may produce variants having single or multiple amino acid substitutions, deletions, or additions.

These variants may include: (a) variants in which one or more amino acid residues are substituted with conservative or nonconservative amino acids, (b) variants in which one or more amino acids are added to or deleted from the polypeptide, (c) variants in which one or more amino acids include a substituent group, and (d) variants in which the polypeptide is fused with another peptide or polypeptide such as a fusion partner, a protein tag or other chemical moiety, that may confer useful properties to the polypeptide, such as, for example, an epitope for an antibody, a polyhistidine sequence, a biotin moiety and the like. Antibodies or antigen-binding fragments described herein may include variants in which amino acid residues from one species are substituted for the corresponding residue in another species, either at the conserved or nonconserved positions. In other embodiments, amino acid residues at nonconserved positions are substituted with conservative or nonconservative residues. The techniques for obtaining these variants, including genetic (deletions, mutations, etc.), chemical, and enzymatic techniques, are known to persons having ordinary skill in the art.

The GPRC5D-specific antibodies or antigen-binding fragments described herein may embody several antibody isotypes, such as IgM, IgD, IgG, IgA and IgE. In some embodiments the antibody isotype is IgG1, IgG2, IgG3, or IgG4 isotype, preferably IgG1 or IgG4 isotype. Antibody or antigen-binding fragment thereof specificity is largely determined by the amino acid sequence, and arrangement, of the CDRs. Therefore, the CDRs of one isotype may be transferred to another isotype without altering antigen specificity. Alternatively, techniques have been established to cause hybridomas to switch from producing one antibody isotype to another (isotype switching) without altering antigen specificity. Accordingly, such antibody isotypes are within the scope of the described antibodies or antigen-binding fragments.

Also provided are vectors comprising the polynucleotides described herein. The vectors can be expression vectors. Recombinant expression vectors containing a sequence encoding a polypeptide of interest are thus contemplated as within the scope of this disclosure. The expression vector may contain one or more additional sequences such as but not limited to regulatory sequences (e.g., promoter, enhancer), a selection marker, and a polyadenylation signal. Vectors for transforming a wide variety of host cells are well known and include, but are not limited to, plasmids, phagemids, cosmids, baculoviruses, bacmids, bacterial artificial

chromosomes (BACs), yeast artificial chromosomes (YACs), as well as other bacterial, yeast and viral vectors.

Recombinant expression vectors within the scope of the description include synthetic, genomic, or cDNA-derived nucleic acid fragments that encode at least one recombinant protein which may be operably linked to suitable regulatory elements. Such regulatory elements may include a transcriptional promoter, sequences encoding suitable mRNA ribosomal binding sites, and sequences that control the termination of transcription and translation. Expression vectors, especially mammalian expression vectors, may also include one or more nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, other 5' or 3' flanking nontranscribed sequences, 5' or 3' nontranslated sequences (such as necessary ribosome binding sites), a polyadenylation site, splice donor and acceptor sites, or transcriptional termination sequences. An origin of replication that confers the ability to replicate in a host may also be incorporated.

The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. Exemplary vectors may be constructed as described by Okayama and Berg, 3 *Mol. Cell. Biol.* 280 (1983).

In some embodiments, the antibody- or antigen-binding fragment-coding sequence is placed under control of a powerful constitutive promoter, such as the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPRT), adenosine deaminase, pyruvate kinase, beta-actin, human myosin, human hemoglobin, human muscle creatine, and others. In addition, many viral promoters function constitutively in eukaryotic cells and are suitable for use with the described embodiments. Such viral promoters include without limitation, Cytomegalovirus (CMV) immediate early promoter, the early and late promoters of SV40, the Mouse Mammary Tumor Virus (MMTV) promoter, the long terminal repeats (LTRs) of Moloney leukemia virus, Human Immunodeficiency Virus (HIV), Epstein Barr Virus (EBV), Rous Sarcoma Virus (RSV), and other retroviruses, and the thymidine kinase promoter of Herpes Simplex Virus. In one embodiment, the GPRC5D-specific antibody or antigen-binding fragment thereof coding sequence is placed under control of an inducible promoter such as the metallothionein promoter, tetracycline-inducible promoter, doxycycline-inducible promoter, promoters that contain one or more interferon-stimulated response elements (ISRE) such as protein kinase R 2',5'-oligoadenylate synthetases, Mx genes, ADAR1, and the like.

Vectors described herein may contain one or more Internal Ribosome Entry Site(s) (IRES). Inclusion of an IRES sequence into fusion vectors may be beneficial for enhancing expression of some proteins. In some embodiments the vector system will include one or more polyadenylation sites (e.g., SV40), which may be upstream or downstream of any of the
5 aforementioned nucleic acid sequences. Vector components may be contiguously linked, or arranged in a manner that provides optimal spacing for expressing the gene products (i.e., by the introduction of “spacer” nucleotides between the ORFs), or positioned in another way. Regulatory elements, such as the IRES motif, may also be arranged to provide optimal spacing for expression.

10 The vectors may comprise selection markers, which are well known in the art. Selection markers include positive and negative selection markers, for example, antibiotic resistance genes (e.g., neomycin resistance gene, a hygromycin resistance gene, a kanamycin resistance gene, a tetracycline resistance gene, a penicillin resistance gene, a puromycin resistance gene, a blasticidin resistance gene), glutamate synthase genes, HSV-TK, HSV-TK derivatives for
15 ganciclovir selection, or bacterial purine nucleoside phosphorylase gene for 6-methylpurine selection (Gadi et al., 7 *Gene Ther.* 1738-1743 (2000)). A nucleic acid sequence encoding a selection marker or the cloning site may be upstream or downstream of a nucleic acid sequence encoding a polypeptide of interest or cloning site.

20 The vectors described herein may be used to transform various cells with the genes encoding the described antibodies or antigen-binding fragments. For example, the vectors may be used to generate GPRC5D-specific antibody or antigen-binding fragment-producing cells. Thus, another aspect features host cells transformed with vectors comprising a nucleic acid sequence encoding an antibody or antigen-binding fragment thereof that specifically binds GPRC5D, such as the antibodies or antigen-binding fragments described and exemplified herein.

25 Numerous techniques are known in the art for the introduction of foreign genes into cells and may be used to construct the recombinant cells for purposes of carrying out the described methods, in accordance with the various embodiments described and exemplified herein. The technique used should provide for the stable transfer of the heterologous gene sequence to the host cell, such that the heterologous gene sequence is heritable and expressible by the cell
30 progeny, and so that the necessary development and physiological functions of the recipient cells are not disrupted. Techniques which may be used include but are not limited to chromosome

transfer (e.g., cell fusion, chromosome mediated gene transfer, micro cell mediated gene transfer), physical methods (e.g., transfection, spheroplast fusion, microinjection, electroporation, liposome carrier), viral vector transfer (e.g., recombinant DNA viruses, recombinant RNA viruses) and the like (described in Cline, 29 *Pharmac. Ther.* 69-92 (1985)).

- 5 Calcium phosphate precipitation and polyethylene glycol (PEG)-induced fusion of bacterial protoplasts with mammalian cells may also be used to transform cells.

Cells suitable for use in the expression of the GPRC5D-specific antibodies or antigen-binding fragments described herein are preferably eukaryotic cells, more preferably cells of plant, rodent, or human origin, for example but not limited to NSO, CHO, CHOK1, perC.6, Tk-ts13, BHK, HEK293 cells, COS-7, T98G, CV-1/EBNA, L cells, C127, 3T3, HeLa, NS1, Sp2/0 myeloma cells, and BHK cell lines, among others. In addition, expression of antibodies may be accomplished using hybridoma cells. Methods for producing hybridomas are well established in the art.

Cells transformed with expression vectors described herein may be selected or screened for recombinant expression of the antibodies or antigen-binding fragments described herein. Recombinant-positive cells are expanded and screened for subclones exhibiting a desired phenotype, such as high level expression, enhanced growth properties, or the ability to yield proteins with desired biochemical characteristics, for example, due to protein modification or altered post-translational modifications. These phenotypes may be due to inherent properties of a given subclone or to mutation. Mutations may be effected through the use of chemicals, UV-wavelength light, radiation, viruses, insertional mutagens, inhibition of DNA mismatch repair, or a combination of such methods.

Methods of using GPRC5D-specific antibodies for treatment

25 Provided herein are GPRC5D-specific antibodies or antigen-binding fragments thereof for use in therapy. In particular, these antibodies or antigen-binding fragments may be useful in treating cancer, such as GPRC5D-expressing cancer. Accordingly, the invention provides a method of treating cancer comprising administering an antibody as described herein, such as GPRC5D-specific antibodies or antigen-binding fragments. For example, the use may be by interfering with GPRC5D-receptor interactions or where the antibody is conjugated to a toxin, so targeting the toxin to the GPRC5D-expressing cancer. In some embodiments GPRC5D-

expressing cancer includes lymphoma, such as multiple myeloma (MM). The antibodies for use in these methods include those described herein above, for example a GPRC5D-specific antibody or antigen-binding fragment with the features set out in Table 1, for example the CDRs or variable domain sequences, and in the further discussion of these antibodies.

5 In some embodiments described herein, immune effector properties of the GPRC5D-specific antibodies may be enhanced or silenced through Fc modifications by techniques known to those skilled in the art. For example, Fc effector functions such as Clq binding, complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cell-mediated phagocytosis (ADCP), down regulation of cell surface
10 receptors (e.g., B cell receptor; BCR), etc. may be provided and/or controlled by modifying residues in the Fc responsible for these activities.

"Antibody-dependent cell-mediated cytotoxicity" or "ADCC" refers to a cell-mediated reaction in which non-specific cytotoxic cells that express Fc receptors (FcRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and
15 subsequently cause lysis of the target cell.

The ability of monoclonal antibodies to induce ADCC can be enhanced by engineering their oligosaccharide component. Human IgG1 or IgG3 are N-glycosylated at Asn297 with the majority of the glycans in the well-known biantennary G0, G0F, G1, G1F, G2 or G2F forms. Antibodies produced by non-engineered CHO cells typically have a glycan fucose content of
20 about at least 85%. The removal of the core fucose from the biantennary complex-type oligosaccharides attached to the Fc regions enhances the ADCC of antibodies via improved Fc.gamma.RIIIa binding without altering antigen binding or CDC activity. Such mAbs can be achieved using different methods reported to lead to the successful expression of relatively high defucosylated antibodies bearing the biantennary complex-type of Fc oligosaccharides such as
25 control of culture osmolality (Konno et al., Cytotechnology 64:249-65, 2012), application of a variant CHO line Lec13 as the host cell line (Shields et al., J Biol Chem 277:26733-26740, 2002), application of a variant CHO line EB66 as the host cell line (Olivier et al., MAbs; 2(4), 2010; Epub ahead of print; PMID:20562582), application of a rat hybridoma cell line YB2/0 as the host cell line (Shinkawa et al., J Biol Chem 278:3466-3473, 2003), introduction of small
30 interfering RNA specifically against the .alpha. 1,6-fucosyltransferase (FUT8) gene (Mori et al., Biotechnol Bioeng 88:901-908, 2004), or coexpression of β -1,4-N-acetylglucosaminyltransferase

III and golgi α -mannosidase II or a potent alpha-mannosidase I inhibitor, kifunensine (Ferrara et al., J Biol Chem 281:5032-5036, 2006, Ferrara et al., Biotechnol Bioeng 93:851-861, 2006; Xhou et al., Biotechnol Bioeng 99:652-65, 2008).

In some embodiments described herein, ADCC elicited by the GPRC5D antibodies may also be enhanced by certain substitutions in the antibody Fc. Exemplary substitutions are for example substitutions at amino acid positions 256, 290, 298, 312, 356, 330, 333, 334, 360, 378 or 430 (residue numbering according to the EU index) as described in U.S. Pat. No. 6,737,056.

Methods of detecting GPRC5D

Provided herein are methods for detecting GPRC5D in a biological sample by contacting the sample with an antibody, or antigen-binding fragment thereof, described herein. As described herein, the sample may be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like. In some embodiments the described methods include detecting GPRC5D in a biological sample by contacting the sample with any of the GPRC5D-specific antibodies or antigen-binding fragments thereof described herein.

In some embodiments the sample may be contacted with more than one of the GPRC5D-specific antibodies or antigen-binding fragments described herein. For example, a sample may be contacted with a first GPRC5D-specific antibody, or antigen-binding fragment thereof, and then contacted with a second GPRC5D-specific antibody, or antigen-binding fragment thereof, wherein the first antibody or antigen-binding fragment and the second antibody or antigen-binding fragment are not the same antibody or antigen-binding fragment. In some embodiments, the first antibody, or antigen-binding fragment thereof, may be affixed to a surface, such as a multiwell plate, chip, or similar substrate prior to contacting the sample. In other embodiments the first antibody, or antigen-binding fragment thereof, may not be affixed, or attached, to anything at all prior to contacting the sample.

The described GPRC5D-specific antibodies and antigen-binding fragments may be detectably labeled. In some embodiments labeled antibodies and antigen-binding fragments may facilitate the detection GPRC5D via the methods described herein. Many such labels are readily known to those skilled in the art. For example, suitable labels include, but should not be

considered limited to, radiolabels, fluorescent labels, epitope tags, biotin, chromophore labels, ECL labels, or enzymes. More specifically, the described labels include ruthenium, ¹¹¹In-DOTA, ¹¹¹In- diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase and beta-galactosidase, poly-histidine (HIS tag), acridine dyes, cyanine dyes, fluorone dyes, oxazin dyes, phenanthridine dyes, rhodamine dyes, Alexafluor® dyes, and the like.

The described GPRC5D-specific antibodies and antigen-binding fragments may be used in a variety of assays to detect GPRC5D in a biological sample. Some suitable assays include, but should not be considered limited to, western blot analysis, radioimmunoassay, surface plasmon resonance, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

In some embodiments described herein detection of GPRC5D-expressing cancer cells in a subject may be used to determine that the subject may be treated with a therapeutic agent directed against GPRC5D.

GPRC5D is present at detectable levels in blood and serum samples. Thus, provided herein are methods for detecting GPRC5D in a sample derived from blood, such as a serum sample, by contacting the sample with an antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D. The blood sample, or a derivative thereof, may be diluted, fractionated, or otherwise processed to yield a sample upon which the described method may be performed. In some embodiments, GPRC5D may be detected in a blood sample, or a derivative thereof, by any number of assays known in the art, such as, but not limited to, western blot analysis, radioimmunoassay, surface plasmon resonance, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

Methods for Diagnosing Cancer

Provided herein are methods for diagnosing GPRC5D-expressing cancer in a subject. In some embodiments GPRC5D-expressing cancer include lymphomas, such as multiple myeloma (MM). In some embodiments, as described above, detecting GPRC5D in a biological sample, such as a blood sample or a serum sample, provides the ability to diagnose cancer in the subject

from whom the sample was obtained. Alternatively, in some embodiments other samples such as a histological sample, a fine needle aspirate sample, resected tumor tissue, circulating cells, circulating tumor cells, and the like, may also be used to assess whether the subject from whom the sample was obtained has cancer. In some embodiments, it may already be known that the subject from whom the sample was obtained has cancer, but the type of cancer afflicting the subject may not yet have been diagnosed or a preliminary diagnosis may be unclear, thus detecting GPRC5D in a biological sample obtained from the subject can allow for, or clarify, diagnosis of the cancer. For example, a subject may be known to have cancer, but it may not be known, or may be unclear, whether the subject's cancer is GPRC5D-expressing.

In some embodiments the described methods involve assessing whether a subject is afflicted with GPRC5D-expressing cancer by determining the amount of GPRC5D that is present in a biological sample derived from the subject; and comparing the observed amount of GPRC5D with the amount of GPRC5D in a control, or reference, sample, wherein a difference between the amount of GPRC5D in the sample derived from the subject and the amount of GPRC5D in the control, or reference, sample is an indication that the subject is afflicted with a GPRC5D-expressing cancer. In another embodiment the amount of GPRC5D observed in a biological sample obtained from a subject may be compared to levels of GPRC5D known to be associated with certain forms or stages of cancer, to determine the form or stage of the subject's cancer. In some embodiments the amount of GPRC5D in the sample derived from the subject is assessed by contacting the sample with an antibody, or an antigen-binding fragment thereof, that immunospecifically binds GPRC5D, such as the GPRC5D-specific antibodies described herein. The sample assessed for the presence of GPRC5D may be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like. In some embodiments GPRC5D-expressing cancer includes hematological cancer, such as multiple myeloma (MM). In some embodiments the subject is a human.

In some embodiments the method of diagnosing a GPRC5D-expressing cancer will involve: contacting a biological sample of a subject with a GPRC5D-specific antibody, or an antigen-binding fragment thereof (such as those derivable from the antibodies and fragments provided in Table 1), quantifying the amount of GPRC5D present in the sample that is bound by

the antibody or antigen-binding fragment thereof, comparing the amount of GPRC5D present in the sample to a known standard or reference sample; and determining whether the subject's GPRC5D levels fall within the levels of GPRC5D associated with cancer. In an additional embodiment, the diagnostic method can be followed with an additional step of administering or
5 prescribing a cancer-specific treatment. In another embodiment, the diagnostic method can be followed with an additional step of transmitting the results of the determination to facilitate treatment of the cancer. In some embodiments the cancer-specific treatment may be directed against GPRC5D-expressing cancers, such as the GPRC5D x CD3 multispecific antibodies described herein.

10 In some embodiments the described methods involve assessing whether a subject is afflicted with GPRC5D-expressing cancer by determining the amount of GPRC5D present in a blood or serum sample obtained from the subject; and comparing the observed amount of GPRC5D with the amount of GPRC5D in a control, or reference, sample, wherein a difference between the amount of GPRC5D in the sample derived from the subject and the amount of
15 GPRC5D in the control, or reference, sample is an indication that the subject is afflicted with a GPRC5D-expressing cancer.

In some embodiments the control, or reference, sample may be derived from a subject that is not afflicted with GPRC5D-expressing cancer. In some embodiments the control, or reference, sample may be derived from a subject that is afflicted with GPRC5D-expressing
20 cancer. In some embodiments where the control, or reference, sample is derived from a subject that is not afflicted with GPRC5D-expressing cancer, an observed increase in the amount of GPRC5D present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is afflicted with GPRC5D-expressing cancer. In some embodiments where the control sample is derived from a subject that is not afflicted with
25 GPRC5D-expressing cancer, an observed decrease or similarity in the amount of GPRC5D present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is not afflicted with GPRC5D-expressing cancer. In some embodiments where the control or reference sample is derived from a subject that is afflicted with GPRC5D-expressing cancer, an observed similarity in the amount of GPRC5D
30 present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is afflicted with GPRC5D-expressing cancer. In some

embodiments where the control or reference sample is derived from a subject that is afflicted with GPRC5D-expressing cancer, an observed decrease in the amount of GPRC5D present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is not afflicted with GPRC5D-expressing cancer.

5 In some embodiments the amount of GPRC5D in the sample derived from the subject is assessed by contacting the sample with an antibody, or an antigen-binding fragment thereof, that specifically binds GPRC5D, such as the antibodies described herein. The sample assessed for the presence of GPRC5D may be derived from a blood sample, a serum sample, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and
10 the like.

 In various aspects, the amount of GPRC5D is determined by contacting the sample with an antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D. In some embodiments, the sample may be contacted by more than one type of antibody, or antigen-
15 binding fragment thereof, that specifically binds GPRC5D. In some embodiments, the sample may be contacted by a first antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D and then contacted by a second antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D. GPRC5D-specific antibodies or antigen-binding fragments such as those described herein may be used in this capacity.

20 Various combinations of the GPRC5D-specific antibodies and antigen-binding fragments can be used to provide a “first” and “second” antibody or antigen-binding fragment to carry out the described diagnostic methods. In some embodiments GPRC5D-expressing cancer includes lymphomas, such as multiple myeloma (MM).

 In certain embodiments, the amount of GPRC5D is determined by western blot analysis,
25 radioimmunoassay, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

 In various embodiments of the described diagnostic methods a control or reference sample is used. This sample may be a positive or negative assay control that ensures the assay
30 used is working properly; for example, an assay control of this nature might be commonly used for immunohistochemistry assays. Alternatively, the sample may be a standardized reference for

the amount of GPRC5D in a biological sample from a healthy subject. In some embodiments, the observed GPRC5D levels of the tested subject may be compared with GPRC5D levels observed in samples from subjects known to have GPRC5D-expressing cancer. In some embodiments, the control subject may be afflicted with a particular cancer of interest. In some
5 embodiments, the control subject is known to have early stage cancer, which may or may not be GPRC5D-expressing cancer. In some embodiments, the control subject is known to have intermediate stage cancer, which may or may not be GPRC5D-expressing cancer. In some embodiments, the control subject is known to have late stage, which may or may not be GPRC5D-expressing cancer.

10 **Methods for Monitoring Cancer**

Provided herein are methods for monitoring GPRC5D-expressing cancer in a subject. In some embodiments GPRC5D-expressing cancer includes lymphomas, such as multiple myeloma (MM). In some embodiments the described methods involve assessing whether GPRC5D-
15 expressing cancer is progressing, regressing, or remaining stable by determining the amount of GPRC5D that is present in a test sample derived from the subject; and comparing the observed amount of GPRC5D with the amount of GPRC5D in a biological sample obtained, in a similar manner, from the subject at an earlier point in time, wherein a difference between the amount of GPRC5D in the test sample and the earlier sample provides an indication of whether the cancer
20 is progressing, regressing, or remaining stable. In this regard, a test sample with an increased amount of GPRC5D, relative to the amount observed for the earlier sample, may indicate progression of a GPRC5D-expressing cancer. Conversely, a test sample with a decreased amount of GPRC5D, relative to the amount observed for the earlier sample, may indicate regression of a GPRC5D-expressing cancer.

25 Accordingly, a test sample with an insignificant difference in the amount of GPRC5D, relative to the amount observed for the earlier sample, may indicate a state of stable disease for a GPRC5D-expressing cancer. In some embodiments the amount of GPRC5D in a biological sample derived from the subject is assessed by contacting the sample with an antibody, or an antibody fragment thereof, that specifically binds GPRC5D, such as the antibodies described
30 herein. The sample assessed for the presence of GPRC5D may be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue

associated (i.e., free cells), tissues (e.g., surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like. In some embodiments the subject is a human.

In some embodiments the methods of monitoring a GPRC5D-expressing cancer will
5 involve: contacting a biological sample of a subject with a GPRC5D-specific antibody, or antigen-binding fragment thereof (such as those derivable from the antibodies and fragments provided in Table 1), quantifying the amount of GPRC5D present in the sample, comparing the amount of GPRC5D present in the sample to the amount of GPRC5D determined to be in a biological sample obtained, in a similar manner, from the same subject at an earlier point in time;
10 and determining whether the subject's GPRC5D level has changed over time. A test sample with an increased amount of GPRC5D, relative to the amount observed for the earlier sample, may indicate progression of cancer. Conversely, a test sample with a decreased amount of GPRC5D, relative to the amount observed for the earlier sample, may indicate regression of a GPRC5D-expressing cancer. Accordingly, a test sample with an insignificant difference in the amount of
15 GPRC5D, relative to the amount observed for the earlier sample, may indicate a state of stable disease for a GPRC5D-expressing cancer. In some embodiments, the GPRC5D levels of the sample may be compared to a known standard or a reference sample, alone or in addition to the GPRC5D levels observed for a sample assessed at an earlier point in time. In an additional embodiment, the diagnostic method can be followed with an additional step of administering a
20 cancer-specific treatment. In some embodiments the cancer-specific treatment may be directed against GPRC5D-expressing cancers.

In various aspects, the amount of GPRC5D is determined by contacting the sample with an antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D. In some
25 embodiments, the sample may be contacted by more than one type of antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D. In some embodiments, the sample may be contacted by a first antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D and then contacted by a second antibody, or antigen-binding fragment thereof, that specifically binds GPRC5D. Antibodies such as those described herein may be used in this capacity.

30 Various combinations of the antibodies and antigen-binding fragments described in Table 1 can be used to provide a "first" and "second" antibody or antigen-binding fragment to carry out

the described monitoring methods. In some embodiments GPRC5D-expressing cancer includes a hematological cancer, such as multiple myeloma (MM).

In certain embodiments, the amount of GPRC5D is determined by western blot analysis, radioimmunoassay, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

Kits for Detecting GPRC5D

Provided herein are kits for detecting GPRC5D in a biological sample. These kits include one or more of the GPRC5D-specific antibodies described herein, or an antigen-binding fragment thereof, and instructions for use of the kit.

The provided GPRC5D-specific antibody, or antigen-binding fragment, may be in solution; lyophilized; affixed to a substrate, carrier, or plate; or detectably labeled.

The described kits may also include additional components useful for performing the methods described herein. By way of example, the kits may comprise means for obtaining a sample from a subject, a control or reference sample, *e.g.*, a sample from a subject having slowly progressing cancer and/or a subject not having cancer, one or more sample compartments, and/or instructional material which describes performance of a method of the invention and tissue specific controls or standards.

The means for determining the level of GPRC5D can further include, for example, buffers or other reagents for use in an assay for determining the level of GPRC5D. The instructions can be, for example, printed instructions for performing the assay and/or instructions for evaluating the level of expression of GPRC5D.

The described kits may also include means for isolating a sample from a subject. These means can comprise one or more items of equipment or reagents that can be used to obtain a fluid or tissue from a subject. The means for obtaining a sample from a subject may also comprise means for isolating blood components, such as serum, from a blood sample. Preferably, the kit is designed for use with a human subject.

Multispecific Antibodies

The binding domains of the anti-GPRC5D antibodies described herein recognize cells expressing GPRC5D on their surface. As noted above, GPRC5D expression can be indicative of a cancerous cell. More specific targeting to particular subsets of cells can be achieved by making bispecific molecules, such as antibodies or antibody fragments, which bind to GPRC5D and to another target, such as CD3 and BCMA. This is achieved by making a molecule which comprises a first region binding to GPRC5D and a second binding region binding to the other target antigen. The antigen-binding regions can take any form that allows specific recognition of the target, for example the binding region may be or may include a heavy chain variable domain, an Fv (combination of a heavy chain variable domain and a light chain variable domain), a binding domain based on a fibronectin type III domain (such as from fibronectin, or based on a consensus of the type III domains from fibronectin, or from tenascin or based on a consensus of the type III domains from tenascin, such as the Centyrin molecules from Janssen Biotech, Inc., see *e.g.* WO2010/051274 and WO2010/093627). Accordingly, bispecific molecules comprising two different antigen-binding regions which bind GPRC5D and another antigen, respectively, are provided.

Some of the multispecific antibodies described herein comprise two different antigen-binding regions which bind GPRC5D and CD3, respectively. In preferred embodiments, multispecific antibodies that bind GPRC5D and CD3 (GPRC5D x CD3-multispecific antibodies) and multispecific antigen-binding fragments thereof are provided. In some embodiments, the GPRC5D x CD3-multispecific antibody comprises a first heavy chain (HC1) and a first light chain (LC1) that pair to form a first antigen-binding site that specifically binds GPRC5D and a second heavy chain (HC2) and a second light chain (LC2) that pair to form a second antigen-binding site that specifically binds CD3. In preferred embodiments, the GPRC5D x CD3-multispecific antibody is a bispecific antibody comprising a GPRC5D-specific arm comprising a first heavy chain (HC1) and a first light chain (LC1) that pair to form a first antigen-binding site that specifically binds CD3 and a CD3-specific arm comprising second heavy chain (HC2) and a second light chain (LC2) that pair to form a second antigen-binding site that specifically binds GPRC5D. In some embodiments, the bispecific antibodies of the invention include antibodies having a full length antibody structure. "Full length antibody" as used herein refers to an antibody having two full length antibody heavy chains and two full length antibody light chains. A full length antibody heavy chain (HC) includes heavy chain variable and constant domains

VH, CH1, CH2, and CH3. A full length antibody light chain (LC) includes light chain variable and constant domains VL and CL. The full length antibody may be lacking the C-terminal lysine (K) in either one or both heavy chains. The term "Fab-arm" or "half molecule" refers to one heavy chain-light chain pair that specifically binds an antigen. In some embodiments, one of the antigen-binding domains is a non-antibody based binding domain, *e.g.* a binding domain of based on a fibronectin type 3 domain, *e.g.* Centyrin.

The GPRC5D-binding arm of the multispecific antibodies provided herein may be derived from any of the GPRC5D-specific antibodies described above. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises a heavy chain CDR1, CDR2, and CDR3 derived from an antibody clone as described in Table 1. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises heavy chain CDR1, CDR2, and CDR3 and light chain CDR1, CDR2, and CDR3 derived from an antibody clone as described in Table 1. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises heavy chain CDR1, CDR2, and CDR3 of clone GC5B81, GC5B465, GS5B483, GC5B596, GC5B382, GC5B379, GC5B373, GC5B376, GC5B385, GC5B370, GC5B602, GC5B603, GC5B599, GC5B601, GC5B598, or GC5B597.

In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises heavy chain CDR1, CDR2, and CDR3 and light chain CDR1, CDR2, and CDR3 of clone GC5B81, GC5B465, GS5B483 or GC5B596. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises a heavy chain variable domain derived from an antibody clone as described in Table 1. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises heavy chain variable domain and light chain variable domain derived from an antibody clone as described in Table 1. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises heavy chain variable domain of clone GC5B81, GC5B465, GS5B483 or GC5B596. In some exemplary embodiments of such GPRC5D-binding arms, the first antigen-binding region which binds GPRC5D comprises heavy chain variable domain and light chain variable domain of clone GC5B81, GC5B465, GS5B483, GC5B596, GC5B382, GC5B379,

GC5B373, GC5B376, GC5B385, GC5B370, GC5B602, GC5B603, GC5B599, GC5B601, GC5B598, or GC5B597. .

5 Table 3 provides a listing of GPRC5D x CD3 bi-specific antibodies having one heavy and light chain pair specific for GPRC5D and another heavy and light chain pair specific for CD3, where the particular antibody ID is listed to describe the antigen-specific antibody arms used to produce the described embodiment.

Table 3:

GPRC5D-specific arm = Ab ID	CD3-specific arm = Ab ID
GC5B81	CD3B219
GC5B465	CD3B219
GC5B483	CD3B219
GC5B596	CD3B219
GC5B382	CD3B219
GC5B379	CD3B219
GC5B373	CD3B219
GC5B376	CD3B219
GC5B385	CD3B219
GC5B370	CD3B219
GC5B602	CD3B219
GC5B603	CD3B219
GC5B599	CD3B219
GC5B601	CD3B219
GC5B598	CD3B219
GC5B597	CD3B219

10 In some embodiments of the bispecific antibodies, the GPRC5D-binding arm binds also binds cynomolgus GPRC5D, preferably the extracellular domain thereof.

In some embodiments, the GPRC5D-binding arm of the multispecific antibody is IgG, or a derivative thereof, e.g., IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the GPRC5D-binding arm has an IgG4 isotype, it contains S228P, L234A, and L235A
15 substitution(s) in its Fc region.

In some embodiments of the bispecific antibodies, the second antigen-binding arm binds human CD3. In some preferred embodiments, the CD3-specific arm of the GPRC5D x CD3 bispecific antibody is derived from a CD3-specific antibody that binds and activates human

primary T cells and/or cynomolgus monkey primary T cells. In some embodiments, the CD3-binding arm binds to an epitope at the N-terminus of CD3ε. In some embodiments, the CD3-binding arm contacts an epitope including the six N-terminal amino acids of CD3ε. In some embodiments, the CD3-specific binding arm of the bispecific antibody is derived from the mouse monoclonal antibody SP34, a mouse IgG3/lambda isotype. In some embodiments, the CD3-binding arm comprises the CDRs of antibody SP34. Such CD3-binding arms may bind to CD3 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. The CD3-specific binding arm may be a humanized version of an arm of mouse monoclonal antibody SP34. Human framework adaptation (HFA) may be used to humanize the anti-CD3 antibody from which the CD3-specific arm is derived. In some embodiments of the bispecific antibodies, the CD3-binding arm comprises a heavy chain and light chain pair selected from Table 2.

In some embodiments, the CD3-binding arm is IgG, or a derivative thereof. In some embodiments, the CD3-binding arm is IgG1, IgG2, IgG3, or IgG4. In some embodiments where in the CD3-binding arm has an IgG4 isotype, it contains S228P, L234A, L235A, F405L, and R409K substitution(s) in its Fc region. In some embodiments, the antibodies or antigen-binding fragments bind CD3ε on primary human T cells. In some embodiments, the antibodies or antigen-binding fragments bind CD3ε on primary cynomolgus T cells. In some embodiments, the antibodies or antigen-binding fragments bind CD3ε on primary human and cynomolgus T cells. In some embodiments, the antibodies or antigen-binding fragments activate primary human CD3+ T cells. In some embodiments, the antibodies or antigen-binding fragments activate primary cynomolgus CD4+ T cells.

In some embodiments are provided a GPRC5D x CD3 bispecific antibody having a GPRC5D-binding arm comprising a heavy chain of antibody clone GC5B81, GC5B465, GS5B483, GC5B596, GC5B382, GC5B379, GC5B373, GC5B376, GC5B385, GC5B370, GC5B602, GC5B603, GC5B599, GC5B601, GC5B598, or GC5B597. . In some embodiments are provided a GPRC5D x CD3 bispecific antibody having a GPRC5D-binding arm comprising a heavy chain and light chain of antibody clone GC5B81, GC5B465, GS5B483, GC5B596, GC5B382, GC5B379, GC5B373, GC5B376, GC5B385, GC5B370, GC5B602, GC5B603, GC5B599, GC5B601, GC5B598, or GC5B597. In some embodiments are provided a GPRC5D x CD3 bispecific antibody having a CD3-binding arm comprising a heavy chain of antibody

clone CD3B219. In some embodiments are provided a GPRC5D x CD3 bispecific antibody having a CD3-binding arm comprising a heavy chain and light chain of antibody clone CD3B219. In some embodiments are provided a GPRC5D x CD3 bispecific antibody having a GPRC5D-binding arm comprising a heavy chain of antibody clone GC5B81, GC5B465, GS5B483, GC5B596, GC5B382, GC5B379, GC5B373, GC5B376, GC5B385, GC5B370, GC5B602, GC5B603, GC5B599, GC5B601, GC5B598, or GC5B597 and a CD3-binding arm comprising a heavy chain of antibody clone CD3B219. In some embodiments are provided a GPRC5D x CD3 bispecific antibody having a GPRC5D-binding arm comprising a heavy chain and light chain of antibody clone GC5B81, GC5B465, GS5B483, GC5B596, GC5B382, GC5B379, GC5B373, GC5B376, GC5B385, GC5B370, GC5B602, GC5B603, GC5B599, GC5B601, GC5B598, or GC5B597 and a CD3-binding arm comprising a heavy chain and light chain of antibody clone CD3B219.

An exemplary GPRC5D x CD3 bispecific antibody is provided in Table 23.

Different formats of bispecific antibodies have been described and were recently reviewed by Chames and Baty (2009) Curr Opin Drug Disc Dev 12: 276.

In some embodiments, the bispecific antibody of the present invention is a diabody, a cross-body, or a bispecific antibody obtained via a controlled Fab arm exchange as those described in the present invention.

In some embodiments, the bispecific antibodies include IgG-like molecules with complementary CH3 domains to force heterodimerisation; recombinant IgG-like dual targeting molecules, wherein the two sides of the molecule each contain the Fab fragment or part of the Fab fragment of at least two different antibodies; IgG fusion molecules, wherein full length IgG antibodies are fused to an extra Fab fragment or parts of Fab fragment; Fc fusion molecules, wherein single chain Fv molecules or stabilized diabodies are fused to heavy-chain constant-domains, Fc-regions or parts thereof; Fab fusion molecules, wherein different Fab-fragments are fused together; ScFv- and diabody-based and heavy chain antibodies (e.g., domain antibodies, nanobodies) wherein different single chain Fv molecules or different diabodies or different heavy-chain antibodies (e.g. domain antibodies, nanobodies) are fused to each other or to another protein or carrier molecule.

In some embodiments, IgG-like molecules with complementary CH3 domains molecules include the Triomab/Quadroma (Trion Pharma/Fresenius Biotech), the Knobs-into-Holes

(Genentech), CrossMAbs (Roche) and the electrostatically-matched (Amgen), the LUZ-Y (Genentech), the Strand Exchange Engineered Domain body (SEEDbody)(EMD Serono), the Biclonic (Merus) and the DuoBody (Genmab A/S).

5 In some embodiments, recombinant IgG-like dual targeting molecules include Dual Targeting (DT)-Ig (GSK/Domantis), Two-in-one Antibody (Genentech), Cross-linked Mabs (Karmanos Cancer Center), mAb2 (F-Star) and CovX-body (CovX/Pfizer).

In some embodiments, IgG fusion molecules include Dual Variable Domain (DVD)-Ig (Abbott), IgG-like Bispecific (InnClone/Eli Lilly), Ts2Ab (MedImmune/AZ) and BsAb (Zymogenetics), HERCULES (Biogen Idec) and TvAb (Roche).

10 In some embodiments, Fc fusion molecules include to ScFv/Fc Fusions (Academic Institution), SCORPION (Emergent BioSolutions/Trubion, Zymogenetics/BMS), Dual Affinity Retargeting Technology (Fc-DART) (MacroGenics) and Dual(ScFv).sub.2-Fab (National Research Center for Antibody Medicine--China).

In some embodiments, Fab fusion bispecific antibodies include F(ab)₂
 15 (Medarex/AMGEN), Dual-Action or Bis-Fab (Genentech), Dock-and-Lock (DNL) (ImmunoMedics), Bivalent Bispecific (Biotechnol) and Fab-Fv (UCB-Celltech). ScFv-, diabody-based and domain antibodies include but are not limited to Bispecific T Cell Engager (BiTE) (Micromet), Tandem Diabody (Tandab) (Affimed), Dual Affinity Retargeting Technology (DART) (MacroGenics), Single-chain Diabody (Academic), TCR-like Antibodies (AIT,
 20 ReceptorLogics), Human Serum Albumin ScFv Fusion (Merrimack) and COMBODY (Epigen Biotech), dual targeting nanobodies (Ablynx), dual targeting heavy chain only domain antibodies.

Full length bispecific antibodies of the invention may be generated for example using Fab arm exchange (or half molecule exchange) between two mono specific bivalent antibodies by
 25 introducing substitutions at the heavy chain CH3 interface in each half molecule to favor heterodimer formation of two antibody half molecules having distinct specificity either in vitro in cell-free environment or using co-expression. The Fab arm exchange reaction is the result of a disulfide-bond isomerization reaction and dissociation-association of CH3 domains. The heavy-chain disulfide bonds in the hinge regions of the parent mono specific antibodies are reduced.
 30 The resulting free cysteines of one of the parent monospecific antibodies form an inter heavy-chain disulfide bond with cysteine residues of a second parent mono specific antibody molecule

and simultaneously CH3 domains of the parent antibodies release and reform by dissociation-association. The CH3 domains of the Fab arms may be engineered to favor heterodimerization over homodimerization. The resulting product is a bispecific antibody having two Fab arms or half molecules which each bind a distinct epitope, i.e. an epitope on GPRC5D and an epitope on CD3.

"Homodimerization" as used herein refers to an interaction of two heavy chains having identical CH3 amino acid sequences. "Homodimer" as used herein refers to an antibody having two heavy chains with identical CH3 amino acid sequences.

"Heterodimerization" as used herein refers to an interaction of two heavy chains having non-identical CH3 amino acid sequences. "Heterodimer" as used herein refers to an antibody having two heavy chains with non-identical CH3 amino acid sequences.

The "knob-in-hole" strategy (see, e.g., PCT Int'l. Publ. No. WO 2006/028936) may be used to generate full length bispecific antibodies. Briefly, selected amino acids forming the interface of the CH3 domains in human IgG can be mutated at positions affecting CH3 domain interactions to promote heterodimer formation. An amino acid with a small side chain (hole) is introduced into a heavy chain of an antibody specifically binding a first antigen and an amino acid with a large side chain (knob) is introduced into a heavy chain of an antibody specifically binding a second antigen. After co-expression of the two antibodies, a heterodimer is formed as a result of the preferential interaction of the heavy chain with a "hole" with the heavy chain with a "knob". Exemplary CH3 substitution pairs forming a knob and a hole are (expressed as modified position in the first CH3 domain of the first heavy chain/modified position in the second CH3 domain of the second heavy chain): T366Y/F405A, T366W/F405W, F405W/Y407A, T394W/Y407T, T394S/Y407A, T366W/T394S, F405W/T394S and T366W/T366S_L368A_Y407V.

Other strategies such as promoting heavy chain heterodimerization using electrostatic interactions by substituting positively charged residues at one CH3 surface and negatively charged residues at a second CH3 surface may be used, as described in US Pat. Publ. No. US2010/0015133; US Pat. Publ. No. US2009/0182127; US Pat. Publ. No. US2010/028637 or US Pat. Publ. No. US2011/0123532. In other strategies, heterodimerization may be promoted by the following substitutions (expressed as modified position in the first CH3 domain of the first heavy chain/modified position in the second CH3 domain of the second heavy chain):

L351Y_F405AY407V/T394W, T366I_K392M_T394W/F405A_Y407V,
T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F,
L351Y_Y407A/T366V_K409F_Y407A/T366A_K409F, or T350V_L351Y_F405A
Y407V/T350V_T366L_K392L_T394W as described in U.S. Pat. Publ. No. US2012/0149876 or
5 U.S. Pat. Publ. No. US2013/0195849.

In addition to methods described above, bispecific antibodies of the invention may be
generated in vitro in a cell-free environment by introducing asymmetrical mutations in the CH3
regions of two mono specific homodimeric antibodies and forming the bispecific heterodimeric
antibody from two parent monospecific homodimeric antibodies in reducing conditions to allow
10 disulfide bond isomerization according to methods described in Intl. Pat. Publ. No.

W02011/131746. In the methods, the first monospecific bivalent antibody (e.g., anti- GPRC5D
antibody) and the second monospecific bivalent antibody (e.g., anti-CD3 antibody) are
engineered to have certain substitutions at the CH3 domain that promotes heterodimer stability;
the antibodies are incubated together under reducing conditions sufficient to allow the cysteines
15 in the hinge region to undergo disulfide bond isomerization; thereby generating the bispecific
antibody by Fab arm exchange. The incubation conditions may optimally be restored to non-
reducing conditions. Exemplary reducing agents that may be used are 2-mercaptoethylamine (2-
MEA), dithiothreitol (DTT), dithioerythritol (DTE), glutathione, tris (2-carboxyethyl) phosphine
(TCEP), L-cysteine and beta-mercaptoethanol, preferably a reducing agent selected from the
20 group consisting of: 2-mercaptoethylamine, dithiothreitol and tris (2-carboxyethyl) phosphine.
For example, incubation for at least 90 min at a temperature of at least 20° C in the presence of at
least 25 mM 2-MEA or in the presence of at least 0.5 mM dithiothreitol at a pH from 5-8, for
example at pH of 7.0 or at pH of 7.4 may be used.

In addition to the described GPRC5D x CD3-multispecific antibodies, also provided are
25 polynucleotide sequences capable of encoding the described GPRC5D x CD3-multispecific
antibodies. Vectors comprising the described polynucleotides are also provided, as are cells
expressing the GPRC5D x CD3-multispecific antibodies provided herein. Also described are
cells capable of expressing the disclosed vectors. These cells may be mammalian cells (such as
293F cells, CHO cells), insect cells (such as Sf7 cells), yeast cells, plant cells, or bacteria cells
30 (such as *E. coli*). The described antibodies may also be produced by hybridoma cells.

Therapeutic composition and methods of treatment using multispecific antibodies and multispecific antigen-binding fragments thereof

The GPRC5D bispecific antibodies discussed above, for example the GPRC5D x CD3 bispecific antibodies discussed above, are useful in therapy. In particular, the GPRC5D bispecific antibodies are useful in treating cancer. Also provided herein are therapeutic compositions for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a multispecific antibody or multispecific antigen-binding fragment described herein and a pharmaceutically acceptable carrier. In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof. In one embodiment said pharmaceutical composition is for the treatment of a GPRC5D-expressing cancer, including (but not limited to) the following: GPRC5D-expressing B cell cancers, such as multiple myeloma (MM); and other cancers yet to be determined in which GPRC5D is expressed. Particular bispecific antibodies that may be used to treat cancer, such as hematological cancer, including the specific cancers discussed above, include antibodies GC5B81, GC5B465, GS5B483 or GC5B596.

The pharmaceutical compositions provided herein comprise: a) an effective amount of a multispecific antibody or antibody fragment of the present invention, and b) a pharmaceutically acceptable carrier, which may be inert or physiologically active. In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof. As used herein, the term "pharmaceutically acceptable carriers" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, and the like that are physiologically compatible. Examples of suitable carriers, diluents and/or excipients include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, and the like, as well as any combination thereof. In many cases, it will be preferable to include isotonic agents, such as sugars, polyalcohols, or sodium chloride in the composition. In particular, relevant examples of suitable carrier include: (1) Dulbecco's phosphate buffered saline, pH about 7.4, containing or not containing about 1 mg/mL to 25 mg/mL human serum albumin,

(2) 0.9% saline (0.9% w/v sodium chloride (NaCl)), and (3) 5% (w/v) dextrose; and may also contain an antioxidant such as tryptamine and a stabilizing agent such as Tween 20 ®.

The compositions herein may also contain a further therapeutic agent, as necessary for the particular disorder being treated. Preferably, the multispecific antibody or antibody fragment and the supplementary active compound will have complementary activities that do not adversely affect each other. In a preferred embodiment, the further therapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2. In a preferred embodiment, the further therapeutic agent is a chemotherapeutic agent.

The compositions of the invention may be in a variety of forms. These include for example liquid, semi-solid, and solid dosage forms, but the preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions. The preferred mode of administration is parenteral (e.g. intravenous, intramuscular, intraperitoneal, subcutaneous). In a preferred embodiment, the compositions of the invention are administered intravenously as a bolus or by continuous infusion over a period of time. In another preferred embodiment, they are injected by intramuscular, subcutaneous, intra-articular, intrasynovial, intratumoral, peritumoral, intralesional, or perilesional routes, to exert local as well as systemic therapeutic effects.

Sterile compositions for parenteral administration can be prepared by incorporating the antibody, antibody fragment or antibody conjugate of the present invention in the required amount in the appropriate solvent, followed by sterilization by microfiltration. As solvent or vehicle, there may be used water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, and the like, as well as combination thereof. In many cases, it will be preferable to include isotonic agents, such as sugars, polyalcohol's, or sodium chloride in the composition. These compositions may also contain adjuvants, in particular wetting, isotonizing, emulsifying, dispersing and stabilizing agents. Sterile compositions for parenteral administration may also be prepared in the form of sterile solid compositions which may be dissolved at the time of use in sterile water or any other injectable sterile medium.

The multispecific antibody or antibody fragment may also be orally administered. As solid compositions for oral administration, tablets, pills, powders (gelatin capsules, sachets) or granules may be used. In these compositions, the active ingredient according to the invention is mixed with one or more inert diluents, such as starch, cellulose, sucrose, lactose or silica, under

an argon stream. These compositions may also comprise substances other than diluents, for example one or more lubricants such as magnesium stearate or talc, a coloring, a coating (sugar-coated tablet) or a glaze.

As liquid compositions for oral administration, there may be used pharmaceutically acceptable solutions, suspensions, emulsions, syrups and elixirs containing inert diluents such as water, ethanol, glycerol, vegetable oils or paraffin oil. These compositions may comprise substances other than diluents, for example wetting, sweetening, thickening, flavoring or stabilizing products.

The doses depend on the desired effect, the duration of the treatment and the route of administration used; they are generally between 5 mg and 1000 mg per day orally for an adult with unit doses ranging from 1 mg to 250 mg of active substance. In general, the doctor will determine the appropriate dosage depending on the age, weight and any other factors specific to the subject to be treated.

Also provided herein are methods for killing a GPRC5D + cell by administering to a patient in need thereof a multispecific antibody which binds said GPRC5D and is able to recruit T cells to kill said GPRC5D + cell (i.e., T cell redirection). Any of the multispecific antibodies or antibody fragments of the invention may be used therapeutically. For example, in one embodiment the GPRC5D x CD3-multispecific antibody may be used therapeutically to treat cancer in a subject.

In a preferred embodiment, multispecific antibodies or antibody fragments of the invention are used for the treatment of a hyperproliferative disorder in a mammal. In a more preferred embodiment, one of the pharmaceutical compositions disclosed above, and which contains a multispecific antibody or antibody fragment of the invention, is used for the treatment of a hyperproliferative disorder in a mammal. In one embodiment, the disorder is a cancer. In particular, the cancer is a GPRC5D-expressing cancer, including (but not limited to) the following: GPRC5D-expressing B-cell cancers, such as multiple myeloma (MM); and other cancers yet to be determined in which GPRC5D is expressed. In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof.

Accordingly, the pharmaceutical compositions of the invention are useful in the treatment or prevention of a variety of cancers, including (but not limited to) the following: a GPRC5D-expressing cancer, including (but not limited to) the following: GPRC5D-expressing B/Plasma cell cancers, such as acute multiple myeloma (MM) or premalignant myelomas such as MGUS (Monoclonal Gammopathy of Undetermined Significance) and SMM (Smoldering Multiple myeloma) and plasmacytoma; and other cancers yet to be determined in which GPRC5D is expressed.

Similarly, further provided herein is a method for inhibiting the growth of selected cell populations comprising contacting GPRC5D-expressing target cells, or tissue containing such target cells, with an effective amount of a multispecific antibody or antibody fragment of the present invention, either alone or in combination with other cytotoxic or therapeutic agents, in the presence of a peripheral blood mononuclear cell (PBMC). In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof. In a preferred embodiment, the further therapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2. In a preferred embodiment, the further therapeutic agent is a chemotherapeutic agent. The method for inhibiting the growth of selected cell populations can be practiced *in vitro*, *in vivo*, or *ex vivo*.

Examples of *in vitro* uses include treatments of autologous bone marrow prior to their transplant into the same patient in order to kill diseased or malignant cells; treatments of bone marrow prior to its transplantation in order to kill competent T cells and prevent graft-versus-host-disease (GVHD); treatments of cell cultures in order to kill all cells except for desired variants that do not express the target antigen; or to kill variants that express undesired antigen. The conditions of non-clinical *in vitro* use are readily determined by one of ordinary skill in the art.

Examples of clinical *ex vivo* use are to remove tumor cells from bone marrow prior to autologous transplantation in cancer treatment. Treatment can be carried out as follows. Bone marrow is harvested from the patient or other individual and then incubated in medium containing serum to which is added the cytotoxic agent of the invention. Concentrations range from about 10 uM to 1 uM, for about 30 min to about 48 hr at about 37 °C. The exact conditions

of concentration and time of incubation, i.e., the dose, are readily determined by one of ordinary skill in the art. After incubation the bone marrow cells are washed with medium containing serum and returned to the patient by i.v. infusion according to known methods. In circumstances where the patient receives other treatment such as a course of ablative chemotherapy or total-body irradiation between the time of harvest of the marrow and reinfusion of the treated cells, the treated marrow cells are stored frozen in liquid nitrogen using standard medical equipment.

For clinical *in vivo* use, a therapeutically effective amount of the multispecific antibody or antigen-binding fragment is administered to a subject in need thereof. For example, the GPRC5D x CD3-multispecific antibodies and multispecific antigen-binding fragments thereof may be useful in the treatment of a GPRC5D-expressing cancer in a subject in need thereof. In some embodiments, the GPRC5D-expressing cancer is a B-cell cancer, such as multiple myeloma (MM). In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof. In some embodiments, the subject is a mammal, preferably a human. In some embodiments, the multispecific antibody or antigen-binding fragment will be administered as a solution that has been tested for sterility.

Dosage regimens in the above methods of treatment and uses are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage.

The efficient dosages and the dosage regimens for the multispecific antibodies and fragments depend on the disease or condition to be treated and may be determined by one skilled in the art. An exemplary, non-limiting range for a therapeutically effective amount of a compound of the present invention is about 0.001-10 mg/kg, such as about 0.001-5 mg/kg, for example about 0.001-2 mg/kg, such as about 0.001-1 mg/kg, for instance about 0.001, about 0.01, about 0.1, about 1 or about 10 mg/kg.

A physician or veterinarian having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the

physician or veterinarian could start doses of the multispecific antibody or fragment employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a bispecific antibody of the present invention will be that amount
5 of the compound which is the lowest dose effective to produce a therapeutic effect.

Administration may e.g. be parenteral, such as intravenous, intramuscular or subcutaneous. In one embodiment, the multispecific antibody or fragment may be administered by infusion in a weekly dosage of calculated by mg/m^2 . Such dosages can, for example, be based on the mg/kg dosages provided above according to the following: dose (mg/kg) $\times 70$: 1.8. Such administration
10 may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hr, such as of from 2 to 12 hr. In one embodiment, the multispecific antibody or fragment may be administered by slow continuous infusion over a long period, such as more than 24 hours, in order to reduce toxic side effects.

In one embodiment, the multispecific antibody or fragment may be administered in a
15 weekly dosage of calculated as a fixed dose for up to eight times, such as from four to six times when given once a week. Such regimen may be repeated one or more times as necessary, for example, after six months or twelve months. Such fixed dosages can, for example, be based on the mg/kg dosages provided above, with a body weight estimate of 70 kg. The dosage may be determined or adjusted by measuring the amount of bispecific antibody of the present invention
20 in the blood upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the GPRC5D antigen binding region of the multispecific antibodies of the present invention.

In one embodiment, the multispecific antibody or fragment may be administered by maintenance therapy, such as, e.g., once a week for a period of six months or more.

25 A multispecific antibody or fragment may also be administered prophylactically in order to reduce the risk of developing cancer, delay the onset of the occurrence of an event in cancer progression, and/or reduce the risk of recurrence when a cancer is in remission.

The multispecific antibodies and fragments thereof as described herein may also be administered in combination therapy, i.e., combined with other therapeutic agents relevant for
30 the disease or condition to be treated. Accordingly, in one embodiment, the antibody-containing medicament is for combination with one or more further therapeutic agent, such as a

chemotherapeutic agent. In some embodiments, the other therapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2. Such combined administration may be simultaneous, separate or sequential, in any order. For simultaneous administration the agents may be administered as one composition or as separate compositions, as appropriate.

5 In one embodiment, a method for treating a disorder involving cells expressing GPRC5D in a subject, which method comprises administration of a therapeutically effective amount of a multispecific antibody or fragment, such as a GPRC5D x CD3 bispecific antibody described herein, and radiotherapy to a subject in need thereof is provided. In one embodiment is provided a method for treating or preventing cancer, which method comprises administration of a
10 therapeutically effective amount of a multispecific antibody or fragment, such as a GPRC5D x CD3 antibody described herein, and radiotherapy to a subject in need thereof. Radiotherapy may comprise radiation or associated administration of radiopharmaceuticals to a patient is provided. The source of radiation may be either external or internal to the patient being treated (radiation treatment may, for example, be in the form of external beam radiation therapy (EBRT) or
15 brachytherapy (BT)). Radioactive elements that may be used in practicing such methods include, e.g., radium, cesium-137, iridium-192, americium-241, gold-198, cobalt-57, copper-67, technetium-99, iodide-123, iodide-131, and indium-111.

Kits

20 Also provided herein are includes kits, e.g., comprising a described multispecific antibody or antigen-binding fragment thereof and instructions for the use of the antibody or fragments for killing of particular cell types. In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as
25 described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof. The instructions may include directions for using the multispecific antibody or antigen-binding fragment thereof in vitro, in vivo or ex vivo.

Typically, the kit will have a compartment containing the multispecific antibody or antigen-binding fragment thereof. The multispecific antibody or antigen-binding fragment
30 thereof may be in a lyophilized form, liquid form, or other form amendable to being included in a kit. The kit may also contain additional elements needed to practice the method described on

the instructions in the kit, such a sterilized solution for reconstituting a lyophilized powder, additional agents for combining with the multispecific antibody or antigen-binding fragment thereof prior to administering to a patient, and tools that aid in administering the multispecific antibody or antigen-binding fragment thereof to a patient.

5

Diagnostic Uses

The multispecific antibodies and fragments described herein may also be used for diagnostic purposes. Thus, also provided are diagnostic compositions comprising a multispecific antibody or fragments as defined herein, and to its use. In preferred embodiments, the multispecific antibody is a GPRC5D x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a GPRC5D x CD3-bispecific antibody as described herein, or a GPRC5D x CD3-bispecific antigen-binding fragment thereof. In one embodiment, the present invention provides a kit for diagnosis of cancer comprising a container comprising a bispecific GPRC5D x CD3 antibody, and one or more reagents for detecting binding of the antibody to GPRC5D. Reagents may include, for example, fluorescent tags, enzymatic tags, or other detectable tags. The reagents may also include secondary or tertiary antibodies or reagents for enzymatic reactions, wherein the enzymatic reactions produce a product that may be visualized. For example, the multispecific antibodies described herein, or antigen-binding fragments thereof, may be labeled with a radiolabel, a fluorescent label, an epitope tag, biotin, a chromophore label, an ECL label, an enzyme, ruthenium, ¹¹¹In-DOTA, ¹¹¹In- diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase and beta-galactosidase, or poly-histidine or similar such labels known in the art.

EMBODIMENTS

The disclosure provided herein also provides the following non-limiting embodiments.

1. An isolated antibody, or an antigen-binding fragment thereof, that binds specifically to GPRC5D comprising:

a. a heavy chain complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 1, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 5, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 9;

5 b. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 2, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 6, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 10;

c. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 3, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 7, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 11;

10 d. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 4, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 8, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 12;

e. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 61, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 67, and a heavy chain CDR3
15 having the amino acid sequence of SEQ ID NO: 72;

f. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 2, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 28, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 30;

20 g. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 27, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 29, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 73;

h. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 27, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 29, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 11;

25 i. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 62, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 68, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 74;

j. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 63, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 69, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 75;

5 k. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 64, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 70, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 12;

l. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 65, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 68, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 76; or

10 m. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 66, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 71, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 77.

2. The isolated antibody, or an antigen-binding fragment of embodiment 1, wherein

15 a. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 1, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 5, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 9 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 13, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 16, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 19;

20 b. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 2, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 6, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 10 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 13, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 16, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 19;

25 c. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 3, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 7, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 11 further comprises a

light chain CDR1 having the amino acid sequence of SEQ ID NO: 14, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 17, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 20;

5 d. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 4, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 8, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 12 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 15, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 18, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 21;

10 e. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 61, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 67, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 72 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 13, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 78, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 80;

f. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 2, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 28, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 30 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 13, a light chain CDR2
20 having the amino acid sequence of SEQ ID NO: 6, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 19;

g. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 27, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 29, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 73 further
25 comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 14, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 17, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 20;

h. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 27, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 29,

and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 11 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 14, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 17, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 20;

5 i. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 62, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 68, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 74 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 14, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 17, and a light chain CDR3 having the
10 amino acid sequence of SEQ ID NO: 20;

j. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 63, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 69, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 75 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 13, a light chain
15 CDR2 having the amino acid sequence of SEQ ID NO: 78, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 80;

k. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 61, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 67, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 72 further
20 comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 13, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 78, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 80;

l. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 65, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 68, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 76 further
25 comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 95, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 79, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 81; or

m. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 66, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 71, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 77 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 15, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 18, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 21.

3. An isolated antibody, or an antigen-binding fragment thereof, that binds specifically to GPRC5D, and comprises a variable heavy (VH) chain region selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 82, 83, 84, 85, 86, 87, 88, 89, or 91.

4. The antibody of embodiment 3, wherein the antibody or antigen binding fragment thereof comprises a variable light (VL) chain region selected from the group consisting of SEQ ID NOs: 56, 57, 58, 92, 93, or 94.

5. The antibody of embodiment 3, wherein the antibody or antigen binding fragment thereof comprises a VH region selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 82, 83, 84, 85, 86, 87, 88, 89, or 91 and a VL region selected from the group consisting of SEQ ID NOs: 56, 57, 58, 92, 93, or 94.

6. The antibody of embodiment 5, wherein the VH chain region comprises SEQ ID NOs: 52, 53, or 83 paired with a VL chain region comprising SEQ ID NO: 56.

7. The antibody of embodiment 5, wherein the VH chain region comprises SEQ ID NOs: 54, 84, 85, 86, or 90 paired with a VL chain region comprising SEQ ID NO: 57.

8. The antibody of embodiment 5, wherein the VH chain region comprises SEQ ID NOs: 55 or 88 paired with a VL chain region comprising SEQ ID NO: 58.

9. The antibody of embodiment 5, wherein the VH chain region comprises SEQ ID NOs: 82 or 87 paired with a VL chain region comprising SEQ ID NO: 92.

10. The antibody of embodiment 5, wherein the VH chain region comprises SEQ ID NO: 89 paired with a VL chain region comprising SEQ ID NO: 93.

11. The antibody of embodiment 5, wherein the VH chain region comprises SEQ ID NO: 91 paired with a VL chain region comprising SEQ ID NO: 94.
12. The antibody or antigen-binding fragment of any one of embodiments 1 to 11 wherein the antibody or antigen-binding fragment thereof binds to a polypeptide having the amino acid
5 sequence of SEQ ID NO: 22.
13. The antibody or antigen-binding fragment of any one of embodiments 1 to 12 wherein the antibody or antigen-binding fragment is a human antibody or antigen-binding fragment.
14. The antibody or antigen-binding fragment of any one of embodiments 1 to 13 wherein the antibody or antigen-binding fragment is recombinant.
- 10 15. The antigen binding fragment of any one of embodiments 1 to 14 wherein the antigen binding fragment is a Fab fragment, a Fab2 fragment, or a single chain antibody.
16. The antibody or antigen-binding fragment of any one of embodiments 1 to 15 wherein the antibody or antigen-binding fragment thereof are of IgG1, IgG2, IgG3, or IgG4 isotype.
17. The antibody or antigen-binding fragment of any of embodiments 1 to 9 is an IgG1 or an
15 IgG4 isotype.
18. The antibody of embodiment 17 wherein the IgG1 has a K409R substitution in its Fc region.
19. The antibody of embodiment 17 wherein the IgG1 has an F405L substitution in its Fc region.
- 20 20. The antibody of embodiment 20 wherein the IgG4 has an F405L substitution and an R409K substitution in its Fc region.
21. The antibody of embodiment 16 further comprising an S228P substitution, an L234A substitution, and an L235A substitution in its Fc region.
22. The antibody or antigen-binding fragment of any one of embodiments 1 to 14 wherein the
25 antibody or antigen-binding fragment thereof specifically binds human GPRC5D and cross reacts to cynomolgus monkey GPRC5D.

23. The antibody or antigen-binding fragment of any one of embodiment 17 wherein the antibody or antigen-binding fragment induces ADCC *in vitro* with an EC₅₀ of less than about 28 nM.

24. An isolated cell expressing the antibody or antigen-binding fragment of any one of
5 embodiments 1 to 11.

25. The cell of embodiment 24 wherein the cell is a hybridoma.

26. The cell of embodiment 24 wherein the antibody is recombinantly produced.

27. An isolated GPRC5D x CD3 bispecific antibody comprising:

a) a first heavy chain (HC1);

10 b) a second heavy chain (HC2);

c) a first light chain (LC1); and

d) a second light chain (LC2),

wherein the HC1 and the LC1 pair to form a first antigen-binding site that specifically binds CD3, and the HC2 and the LC2 pair to form a second antigen-binding site that specifically binds
15 GPRC5D, or a GPRC5D x CD3 -bispecific binding fragment thereof.

28. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 27 wherein HC1 comprises SEQ ID NO: 25 and LC1 comprises SEQ ID NO: 26.

29. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 52 and LC2 comprises SEQ ID NO: 56.

20 30. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 53 and LC2 comprises SEQ ID NO: 56.

31. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 54 and LC2 comprises SEQ ID NO: 57.

25 32. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 55 and LC2 comprises SEQ ID NO: 58.

33. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 82 and LC2 comprises SEQ ID NO: 92.
34. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 83 and LC2 comprises SEQ ID NO: 56.
- 5 35. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 84 and LC2 comprises SEQ ID NO: 57.
36. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 85 and LC2 comprises SEQ ID NO: 57.
37. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 10 28 wherein HC2 comprises SEQ ID NO: 86 and LC2 comprises SEQ ID NO: 57.
38. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 87 and LC2 comprises SEQ ID NO: 92.
39. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 88 and LC2 comprises SEQ ID NO: 58.
- 15 40. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 89 and LC2 comprises SEQ ID NO: 93.
41. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 28 wherein HC2 comprises SEQ ID NO: 91 and LC2 comprises SEQ ID NO: 94.
42. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 20 28 wherein HC2 comprises SEQ ID NO: 85 and LC2 comprises SEQ ID NO: 57.
43. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42 wherein the antibody or bispecific binding fragment are of IgG1, IgG2, IgG3, or IgG4 isotype.
44. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 25 43 wherein the antibody or bispecific binding fragment is IgG4 isotype.

45. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiments 27 to 42 wherein the antibody or bispecific binding fragment thereof binds GPRC5D on the surface of human myeloma cells.
46. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiments 27 to 42 wherein the antibody or bispecific binding fragment thereof binds GPRC5D on the surface of human multiple myeloma cells.
47. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 27 to 42 wherein the antibody or bispecific binding fragment induces human T cell activation in vitro with an EC₅₀ of less than about 0.22 nM.
48. The GPRC5D x CD3 bispecific antibody or bispecific binding fragment of embodiment 27 to 42 wherein the antibody or bispecific binding fragment induces T-cell dependent cytotoxicity of GPRC5D-expressing cells in vitro with an EC₅₀ of less than about 0.89 nM.
49. An isolated cell expressing the antibody or bispecific binding fragment of any one of embodiments 27 to 42.
50. The cell of embodiment 49 wherein the cell is a hybridoma.
51. The cell of embodiment 49 wherein the antibody or bispecific binding fragment is recombinantly produced.
52. A method for treating a subject having cancer, said method comprising:
administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42 to a patient in need thereof for a time sufficient to treat the cancer.
53. A method for inhibiting growth or proliferation of cancer cells, said method comprising:
administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42 to inhibit the growth or proliferation of cancer cells.
54. A method of redirecting a T cell to a GPRC5D-expressing cancer cell, said method comprising:

administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42 to redirect a T cell to a cancer.

55. The method of embodiment 52, 53, or 54 wherein the cancer is a hematological cancer.

5 56. The method of embodiment 55 wherein the hematological cancer is a GPRC5D-expressing B cell cancer.

57. The method of embodiment 56 wherein the GPRC5D-expressing B cell cancer is multiple myeloma.

58. The method of embodiment 52 comprising administering a second therapeutic agent.

10 59. The method of embodiment 58 wherein the second therapeutic agent is a chemotherapeutic agent or a targeted anti-cancer therapy.

60. The method of embodiment 59 wherein the chemotherapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2.

15 61. The method of embodiment 59 wherein the second therapeutic agent is administered to said subject simultaneously with, sequentially, or separately from the bispecific antibody.

62. A pharmaceutical composition comprising the GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42 and a pharmaceutically acceptable carrier.

20 63. A method for generating the GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42 by culturing the cell of any one of embodiments 49 to 51.

64. An isolated synthetic polynucleotide encoding the HC1, the HC2, the LC1 or the LC2 of the GPRC5D x CD3 bispecific antibody or bispecific binding fragment of any one of embodiments 27 to 42.

25 65. A kit comprising the GPRC5D x CD3 bispecific antibody or bispecific binding fragment as defined in any one of embodiments 27 to 42 and/or a polynucleotide as defined in embodiment 63 and packaging for the same.

EXAMPLES

The following examples are provided to supplement the prior disclosure and to provide a better understanding of the subject matter described herein. These examples should not be considered to limit the described subject matter. It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be apparent to persons skilled in the art and are to be included within, and can be made without departing from, the true scope of the invention.

EXAMPLE 1: ANTIGENS

Due to difficulties in production of recombinant GPRC5D antigens, transfected cell lines displaying GPRC5D [human (SEQ ID NO: 22), cyno (SEQ ID NO: 23), and murine (SEQ ID NO: 24)] were generated using standard methods to be used as whole cell antigens for antibody generation and characterization studies (Table 4).

Table 4. GPRC5D-expressing cell lines

Protein	Cell Line	Promoter	Resistance
Human GPRC5D	HEK293T	CMV	Neomycin
Cyno GPRC5D	HEK293F	CMV	Blasticidin

EXAMPLE 2: GENERATION OF GPRC5D ANTIBODIES USING PHAGE DISPLAY

Two distinct approaches were followed in the generation of GPRC5D antibodies by phage: standard cell panning (negative selection) and FACS cell phage panning (competitive selection).

Standard Cell Phage Panning (negative selection):

In-house de novo phage libraries have been described in detail (Shi et al (2010) J. Mol. Biol. 397:385-396; Int. Pat. Publ. No. WO09/085462). These libraries were built on three human VH germline genes (IGHV1-69, 3-23, 5-51) and four human VL germline genes (A27, B3, L6, O12) designed to have high diversity in CDR-H3. Three de novo phage libraries (DNP00004 -

169HC /LC mix, DNP00005 -323HC /LC mix and DNP00006 -551HC / LC mix) displaying Fab variants on phage coat protein pIX were panned against GPRC5D-expressing HEK293 G5 stable cells (target cells) for round 1, 3, 5 and for rounds 2 and 4 the previous round of amplified Fab-pIX phage were applied to HEK293 background cells (the negative selection) (see Table 5).

- 5 Round 1 de novo Fab-pIX phage bound to the target cells were recovered for overnight amplification. The round 1 selected phage was applied to the background cells for round 2 where the unbound Fab-pIX phage were recovered for overnight negatively selected phage amplification. Another set of positive and negative rounds of panning were performed, rounds 3 and 4. The final round was performed with the last round of negatively selected amplified phage
10 split into two panning samples one for target cells and one for background cell panning.

Table 5: Standard cell phage panning flowchart-negative selection

Panning Round	Cells used as antigen
1	Human GPRC5D HEK293
2	HEK293
3	Human GPRC5D HEK293
4	HEK293
5 (split)	Human GPRC5D HEK293
	HEK293

15

Standard Cell Phage Panning (background selection):

- The Fab-pIX phage display libraries were added to HEK293 cells following similar environments and incubation times as performed with the standard cell panning procedure previously mentioned (Table 6). After three rounds the DNA corresponding to the HEK293
20 bound Fab-pIX phage was used to produce PCR amplicons for NGS. These NGS results would be used in an additional dynamic subtractive analysis within the NGS2.0 software to help in distinguishing possible target specific Fab candidates.

Table 6: Standard cell phage panning flowchart-background selection

Panning Round	Cells used as antigen
1	HEK293
2	HEK293
3	HEK293

FACS cell phage panning (competitive selection)

Three rounds of panning were performed by way of applying Fab-pIX phage to a mixture of target cells and background cells at the same time (Table 7). For rounds 1 and 2 the target cells with Fab-pIX phage attached were sorted by use of the GFP signal. To capture the bound Fab-pIX phage from the sorted cells an acid cell lysis was applied followed by an E. coli infection. The round 2 amplified Fab-pIX phage and cell mixture were sorted into two populations in the final round, gated for GFP target cells and gated for non-GFP background cells. The bound Fab-pIX phage for both cell populations was captured by acid lysis and E.coli infection

Table 7: FACS cell phage panning flowchart-competitive selection

Panning Round	Cells used as antigen
1	Human GPRC5D_GFP & HEK293
2	Human GPRC5D_GFP & HEK293
3	Human GPRC5D_GFP & HEK293

Next Generation Sequencing (NGS) of phage panning

The overnight growth under glucose repression was done for the six of the last round panning samples. These cultures were used to make mini-prep DNA, Qiagen QIAspin DNA kit. The six DNA samples were used as PCR template to generate amplicons that measured from HCDR1 to HCDR3. The six amplicons were gel purified and pooled by keeping version 2.1, 3.0 separate for the standard cell panning and completely pooled for FACS cell panning. These pooled gel purified amplicons were provided to Genewiz NGS services to process by MiSeq 1x300 technology. Files were delivered by Genewiz and uploaded into a local server (nas2.0). Within this server the NGS2.0 software application was used to load, read, and analyze the

sequence files. The top 88 sequences according to copy number (>50) and ratio of target cell (+) to background cell (-) sequences (ratio $>5:1$) were selected for IgG conversion. As only variable heavy chain sequence is determined by NGS, the full heavy chain constructs had to be electronically constructed into the appropriate frameworks. In addition, since only heavy chain sequence was known each of the candidates was paired with the four parental light chains (A27, B3, L6, O12). The final conversion for the candidates was done as human IgG4PAA.

ELISA screen and Standard Sequencing

From the same DNA prep used for NGS, a restriction enzyme digest and self-ligation was done to excise the gene pIX to enable soluble Fab expression. Ninety-two colonies, picked for each of the three panning samples were assessed for Fab expression by ELISA and were sequenced to determine both heavy chain and light chain by Sanger method. The final candidates with diversified sequence within the LCDRs were cloned into mammalian expressions plasmids.

EXAMPLE 4: INITIAL CHARACTERIZATION OF GPRC5D ANTIBODIES OBTAINED THROUGH PHAGE DISPLAY TECHNOLOGY

GPRC5D Binding:

Whole cell phage panning was completed using the human GPRC5D cells as antigen, as described above. Following NGS analysis, only heavy chain sequence was known for each of the candidates. Consequently, each heavy chain had to be paired with the 4 parental light chains (A27, B3, L6, O12), resulting in 348 mAbs from 87 Hc sequences identified using NGS. These mAbs were initially evaluated for binding to cyno GPRC5D using FACS. The cyno GPRC5D cell line was selected for initial screening to maximize the potential binding signal as the cyno GPRC5D cell line had a higher expression level than did the human GPRC5D cell line. Briefly, FACS screening was performed by normalizing protein concentration to 1 $\mu\text{g/ml}$ and 100 μl of protein was mixed with 200,000 cells per well. The mAb was allowed to incubate with cells for 1 hour at 4°C . The cells were then washed three times with PBS and 0.2% FBS. An anti-human secondary mAb conjugated with PE (Jackson cat # 709-116-149) was then added as the detection reagent. The cells and secondary antibody were incubated for 1 hour at 4°C . The cells were then

washed three times with PBS and 0.2% FBS. The cells were washed again using PBS and 0.2% FBS and then analyzed on FACSarray.

A large number of the hits were observed to bind to cyno GPRC5D, including 15 mAbs with MFI greater than 100,000, 23 mAbs with MFI less than 100,000 but greater than 10,000 and 63 mAbs with MFI less than 10,000 but greater than 1000. (Table 8).

Table 8: FACS binding data for NGS derived mAbs to cyno GPRC5D. Heavy chain sequences identified from NGS analysis were paired with each parental Lc as shown. GC5M29 was used as the control (PH9L3 light chain). mAbs highlighted have an MFI >1000

Hc ID	GPRC5D mAb [PH9L1 (A27)]	MFI	GPRC5D mAb [PH9L2 (B3)]	MFI	GPRC5D mAb [PH9L3 (L6)]	MFI	GPRC5D mAb [PH9L4 (O12)]	MFI
GC5H18	GC5B22	198	GC5B23	230	GC5B24	246	GC5B25	31324
GC5H17	GC5B66	242	GC5B67	272	GC5B68	2682	GC5B69	17416
GC5H20	GC5B114	199	GC5B115	266	GC5B116	253	GC5B117	5037
GC5H15	GC5B158	14407	GC5B159	139766	GC5B160	84111	GC5B161	7856
GC5H13	GC5B202	89948	GC5B203	9173	GC5B204	15203	GC5B205	52663
GC5H23	GC5B242	242	GC5B243	116641	GC5B244	249	GC5B245	469
GC5H14	GC5B282	487	GC5B283	341	GC5B284	249	GC5B285	87116
GC5H22	GC5B326	195	GC5B237	265	GC5B328	248	GC5B329	292
GC5H24	GC5B26	194	GC5B27	265	GC5B28	229	GC5B29	590
GC5H19	GC5B70	214	GC5B71	229	GC5B72	2013	GC5B73	3827
GC5H33	GC5B118	178	GC5B119	222	GC5B120	880	GC5B121	43924
GC5H36	GC5B162	86773	GC5B163	21970	GC5B164	122043	GC5B165	1870
GC5H31	GC5B206	467	GC5B207	3636	GC5B208	481	GC5B209	1409
GC5H29	GC5B246	202	GC5B247	3037	GC5B248	2835	GC5B249	1373
GC5H34	GC5B286	239	GC5B287	163990	GC5B288	259	GC5B289	271
GC5H35	GC5B330	202	GC5B331	552	GC5B332	191978	GC5B333	1591
GC5H26	GC5B30	204	GC5B31	230	GC5B32	16553	GC5B33	270
GC5H27	GC5B74	238	GC5B75	365	GC5B76	829	GC5B77	317
GC5H25	GC5B122	153	GC5B123	237	GC5B124	6026	GC5B125	6197
GC5H30	GC5B166	1510	GC5B167	6266	GC5B168	32482	GC5B169	327
GC5H47	GC5B210	1730	GC5B211	284	GC5B212	278	GC5B213	727
GC5H42	GC5B250	3969	GC5B251	213233	GC5B252	63295	GC5B253	574
GC5H37	GC5B290	8839	GC5B291	201024	GC5B292	110940	GC5B293	305
GC5H45	GC5B334	160	GC5B335	1434	GC5B336	3228	GC5B337	309
GC5H38	GC5B34	157	GC5B35	283	GC5B36	14936	GC5B37	297
GC5H40	GC5B78	162	GC5B79	309	GC5B80	428	GC5B81	74140
GC5H43	GC5B126	157	GC5B127	265	GC5B128	268	GC5B129	1283
GC5H49	GC5B170	1014	GC5B171	51995	GC5B172	1434	GC5B173	332

GC5H62	GC5B214	2565	GC5B215	274	GC5B216	263	GC5B217	1589
GC5H60	GC5B254	161	GC5B255	5788	GC5B256	460	GC5B257	308
GC5H94	GC5B294	190	GC5B295	2312	GC5B296	745	GC5B298	261
GC5H69	GC5B338	188	GC5B339	260	GC5B340	1798	GC5B341	290
GC5H85	GC5B38	10809	GC5B39	1112	GC5B40	110654	GC5B41	4379
GC5H96	GC5B82	165	GC5B83	272	GC5B84	1547	GC5B85	3563
GC5H97	GC5B130	174	GC5B131	293	GC5B132	306	GC5B133	343
GC5H76	GC5B174	219	GC5B175	3631	GC5B176	244	GC5B177	324
GC5H68	GC5B218	21420	GC5B219	310	GC5B220	777	GC5B221	762
GC5H79	GC5B256	714	GC5B257	1146	GC5B258	456	GC5B259	1129
GC5H71	GC5B298	185	GC5B299	1370	GC5B300	260	GC5B301	292
GC5H93	GC5B342	164	GC5B343	247	GC5B344	401	GC5B345	283
GC5H21	GC5B42	219	GC5B43	307	GC5B44	2694	GC5B45	1022
GC5H39	GC5B86	192	GC5B87	344	GC5B88	470	GC5B89	41027
GC5H50	GC5B134	764	GC5B135	340	GC5B136	600	GC5B137	2087
GC5H28	GC5B178	750	GC5B179	6708	GC5B180	1005	GC5B181	535
GC5H53	GC5B222	284	GC5B223	264	GC5B224	228	GC5B225	605
GC5H51	GC5B262	159	GC5B263	1013	GC5B267	238	GC5B268	309
GC5H64	GC5B302	184	GC5B303	329	GC5B304	240	GC5B305	1571
GC5H16	GC5B346	182	GC5B347	259	GC5B348	332	GC5B349	552
GC5H65	GC5B46	204	GC5B47	327	GC5B48	1320	GC5B49	392
GC5H32	GC5B90	335	GC5B91	340	GC5B92	1848	GC5B93	1780
GC5H54	GC5B138	193	GC5B139	354	GC5B140	323	GC5B141	292
GC5H99	GC5B182	196	GC5B183	6421	GC5B184	309	GC5B185	280
GC5H52	GC5B226	204	GC5B227	298	GC5B228	247	GC5B229	263
GC5H48	GC5B266	268	GC5B267	2160	GC5B268	235	GC5B269	281
GC5H44	GC5B306	204	GC5B307	313	GC5B308	302	GC5B309	621
GC5H46	GC5B350	198	GC5B351	287	GC5B352	4300	GC5B353	1708
GC5H56	GC5B50	229	GC5B51	36679	GC5B52	386	GC5B53	446
GC5H55	GC5B94	255	GC5B95	609	GC5B96	411	GC5B97	350
GC5H98	GC5B142	194	GC5B143	407	GC5B144	284	GC5B145	483
GC5H61	GC5B186	543	GC5B187	1132	GC5B188	376	GC5B189	333
GC5H92	GC5B320	297	GC5B321	294	GC5B322	280	GC5B323	386
GC5H5	GC5B17	243	GC5B18	301	GC5B19	1048	GC5B20	384
GC5H59	GC5B98	272	GC5B99	388	GC5B100	661	GC5B101	528
GC5H66	GC5B354	223	GC5B355	852	GC5B356	374	GC5B357	500
GC5H80	GC5B54	299	GC5B55	415	GC5B56	422	GC5B57	487
GC5H87	GC5B102	383	GC5B103	460	GC5B104	3130	GC5B105	456
GC5H67	GC5B146	325	GC5B147	370	GC5B148	383	GC5B149	487
GC5H73	GC5B190	236	GC5B191	2556	GC5B192	409	GC5B193	387
GC5H86	GC5B230	17885	GC5B231	543	GC5B232	494	GC5B233	620
GC5H70	GC5B270	390	GC5B271	660	GC5B272	389	GC5B273	450

GC5H90	GC5B314	219	GC5B315	334	GC5B316	363	GC5B317	403
GC5H95	GC5B358	211	GC5B359	344	GC5B360	414	GC5B361	449
GC5H74	GC5B58	294	GC5B59	448	GC5B60	477	GC5B61	548
GC5H78	GC5B106	267	GC5B107	440	GC5B108	474	GC5B109	497
GC5H77	GC5B150	312	GC5B151	452	GC5B152	412	GC5B153	709
GC5H91	GC5B194	296	GC5B195	448	GC5B196	491	GC5B197	483
GC5H57	GC5B234	128355	GC5B235	195348	GC5B236	134008	GC5B237	11255
GC5H41	GC5B274	313	GC5B275	7431	GC5B278	435	GC5B279	542
GC5H58	GC5B318	2608	GC5B319	5791	GC5B320	49616	GC5B321	4823
GC5H63	GC5B362	313	GC5B363	392	GC5B364	461	GC5B365	3613
GC5H72	GC5B62	326	GC5B63	463	GC5B64	485	GC5B65	694
GC5H75	GC5B110	208487	GC5B111	692	GC5B112	140269	GC5B113	104136
GC5H81	GC5B154	351	GC5B155	412	GC5B156	440	GC5B157	555
GC5H82	GC5B198	285	GC5B199	593	GC5B200	499	GC5B201	485
GC5H83	GC5B238	3698	GC5B239	4370	GC5B240	4676	GC5B241	825
GC5H88	GC5B278	3659	GC5B279	481	GC5B280	425	GC5B281	633
GC5H89	GC5B332	299	GC5B333	561	GC5B334	4467	GC5B335	494
GC5H84	GC5B366	544	GC5B367	474	GC5B368	6313	GC5B369	26116

The 40 mAbs with the highest binding affinity were selected for additional characterization, which consisted of repeating the binding study to cyno GPCR5D cells, as well as, assessment of binding to human GPCR5D expressing cells using FACS (Table 9). These data were analyzed to select 17 mAbs for purification and GPCR5D x CD3 bispecific antibody generation (highlighted). Factors used to select mAbs for purification and GPCR5D x CD3 bispecific antibody generation include specificity of binding to human GPCR5D, cross-reactivity to cyno GPCR5D and diversity in Hc sequence. For example, GC5H36 was paired with three distinct light chains and analyzed for binding (GC5B162, GC5B163, GC5B164). Only GC5B164 was advanced as this mAb was observed to have the higher MFI to human GPCR5D than GC5B162 or GC5B163.

Table 9: FACS binding data for NGS derived mAbs to cyno and human GPCR5D expressing HEK293F cells. Non-transfected HEK293F cells were used to evaluate binding specificity for GPCR5D. TF7M1636 was used as the isotype control. mAbs highlighted were selected for further analysis.

Protein ID	HC Peptide ID	cyno GPCR5D MFI	human GPCR5D MFI	293F MFI
GC5B38	GC5H85	6,864	280	215
GC5B110	GC5H75	142,277	2,543	235

GC5B158	GC5H15	4,923	285	235
GC5B162	GC5H36	75,139	2,981	230
GC5B202	GC5H13	30,470	2,152	256
GC5B218	GC5H68	10,451	277	263
GC5B230	GC5H86	19,103	364	290
GC5B234	GC5H57	134,258	1,657	261
GC5B51	GC5H56	22,661	667	480
GC5B159	GC5H15	129,370	1,987	258
GC5B163	GC5H36	7,605	840	310
GC5B171	GC5H49	24,133	1,144	1,321
GC5B235	GC5H57	158,011	2,643	254
GC5B243	GC5H23	67,848	20,458	221
GC5B251	GC5H42	74,510	11,418	235
GC5B281	GC5H34	109,418	5,317	570
GC5B291	GC5H37	77,798	10,427	880
GC5B32	GC5H26	3,597	14,656	358
GC5B36	GC5H38	5,353	2,002	352
GC5B40	GC5H85	67,279	4,510	310
GC5B112	GC5H75	104,875	470	433
GC5B160	GC5H15	40,881	635	276
GC5B164	GC5H36	78,672	12,664	343
GC5B168	GC5H30	41,779	28,885	1,238
GC5B204	GC5H13	5,090	390	212
GC5B236	GC5H57	78,546	1,442	192
GC5B252	GC5H42	22,010	1,498	204
GC5B292	GC5H37	63,366	1,872	480
GC5B320	GC5H58	63,345	22,891	231
GC5B332	GC5H35	106,515	13,348	253
GC5B25	GC5H18	22,079	705	431
GC5B69	GC5H17	1,593	851	284
GC5B81	GC5H40	62,756	9,986	288
GC5B89	GC5H39	31,613	1,068	251
GC5B113	GC5H75	66,475	550	323
GC5B121	GC5H33	35,493	3,061	299
GC5B205	GC5H13	63,610	25,199	260
GC5B237	GC5H57	2,579	267	248
GC5B285	GC5H14	81,405	19,870	225
GC5B369	GC5H84	16,241	559	252
	TF7M1636	956	2,621	1,538
	human 2nd	209	316	228
	anti hGPCR5D	9,001	379	70
	mouse anti isotype	60	82	65

unstained	44	50	33
-----------	----	----	----

The concentration dependent binding profile of each of the selected mAbs against human GPRC5D expressing HEK293 cells and non-transfected HEK293 cells was determined using FACS (Fig. 1). All mAbs were observed to bind to human GPRC5D in a dose dependent manner. Three mAbs, GC5B36, GC5B168, and GC5B205 were also observed to bind to the non-transfected (GPRC5D null) HEK293 cells and were deprioritized due to this non-specific interaction with the cells.

The remaining 14 mAbs were selected for bispecific antibody generation with an anti-CD3 arm CD3B219 and anti-RSV null arm B23M46 (Table 10).

Table 10: The relationship between GPRC5D mAb IDs and bispecific IDs.

GPRC5D mAb Protein ID	x CD3 bispecific ID	x B23Null bispecific ID
GC5B320	GCDB44	GCDB30
GC5B243	GCDB40	GCDB26
GC5B285	GCDB43	GCDB29
GC5B332	GCDB45	GCDB31
GC5B164	GCDB35	GCDB21
GC5B251	GCDB41	GCDB27
GC5B81	GCDB32	GCDB18
GC5B235	GCDB38	GCDB24
GC5B110	GCDB34	GCDB20
GC5B202	GCDB36	GCDB22
GC5B234	GCDB37	GCDB23
GC5B252	GCDB42	GCDB28
GC5B236	GCDB39	GCDB25
GC5B89	GCDB33	GCDB19

One bispecific antibody, GCDB38, precipitated during recombination and another, GCDB36, contained >10% aggregate. All other bispecific antibodies passed the standard release criteria and the null arm bispecific antibodies were analyzed for concentration dependent binding to human GPRC5D expressing HEK293 cells (Fig. 2).

All bispecific antibodies were observed to bind in a dose dependent manner. GC5B320 was included as a binding comparator to understand binding differences between bivalent mAbs

and monovalent bispecific antibodies. GCDB44 was included as a binding comparator to understand binding differences between anti-CD3 bispecific antibodies and anti-RSV null arm mAbs. The expected decrease in apparent binding affinity was observed when comparing the mAb GC5B320 to bispecific antibodies GCDB30 and GCDB44. In addition, the GCDB44 (anti-CD3 bispecific Ab) was observed to have slightly higher binding affinity to the anti-human GPRC5D cells compared with the GCDB30 (anti-RSV null arm bispecific Ab), demonstrating that the anti-CD3 arm is positively impacted binding to this cell line.

The panel of bispecific antibodies was also profiled for binding against MM1R and H929 cells, which endogenously express GPRC5D (Fig. 3). The binding profile of the bispecific antibodies on MM1R and H929 cells was compared with the overexpressed human GPCR5D HEK293 cells and non-transfected HEK293 cells using FACS. The bispecific antibodies were observed to bind to GPRC5D as expressed endogenously on MM1R and H929 cells with a range of affinity. The highest binding was observed to the human GPRC5D HEK cells, which as an overexpressed stable cell line, has a much higher receptor density than either MM1R or H929 cells. GCDB37, GCDB38, GCDB39 and GCDB41 were not advanced due to the low binding affinity observed for H929 and MM1R cells.

***In vitro* T-Cell Dependent Cytotoxicity**

The panel of bispecific antibodies were then profiled for potency in a T-cell mediated cytotoxicity assay using H929 and MM1R target cells (Fig. 4 A and B, Table 11). Briefly, target cells (H929, MM1R, OPM2, LP-1 and Daudi or HEK parent and HEK+GPRC5D cells) were counted and 10 million cells were centrifuged at 1350 rpm for 3 minutes and cell pellets were resuspended in 1 ml of diluted CFSE solution (CellTrace CFSE proliferation stain was reconstituted in 18 μ l of sterile DMSO and 1 μ l of the solution was diluted in 10 ml of sterile PBS) and incubated at room temperature for 8 minutes in dark. After the incubation, 1 ml of HI FBS was added to cell suspension to quench the surplus CFSE. Cells were washed twice in RPMI-1640 with 10% FBS. After reconstitution in 10 ml of RPMI, cells were counted and cell viability was recoded in a spreadsheet. Cells were diluted to 2.2×10^5 /ml and incubated at 37°C until use.

Pan T cells from normal donors were thawed in 37°C water bath, and cells were then centrifuged at 1350 rpm at 4°C for 3 minutes. The supernatants were discarded and reconstituted

in culture medium at 1.1×10^6 /ml concentration. 2×10^5 target cells were added to wells of a 96-well U-bottom plate, followed by Fc blocker (to final concentration of 2mg/ml). All cell lines were incubated at room temperature for 10 minutes to block Fc receptor activity. 1×10^5 T cells were added to the wells (5:1 Effector:Target ratio). After target and T cells were mixed, 20 μ l of

5 GPRC5D x CD3 bispecific Ab dilutions were added to each well. GPRC5D x CD3 bispecific antibodies were diluted to 800 μ g/ml (10X) in PBS. The titration was prepared in 4-fold serial dilutions in PBS in a 96-well U-bottom plate. The last column was left as PBS alone (vehicle control). The plates were incubated at 37°C with 5% CO_2 for 48 hours.

Two days later (48 hours), the plates were centrifuged and 100 μ l of supernatants were

10 stored at -80°C for cytokine release assay. Cells were washed in 200 μ l of PBS and incubated in 50 μ l of near-IR Live/Dead stain (1:200 dilution) and anti-CD25 PE antibody (1:50 dilution) for 20 minutes at room temperature. Then, the cells were washed once in 200 μ l of FACS buffer and finally reconstituted in 150 μ l of FACS buffer. Cells were analyzed using FACSCanto II and FlowJo 7.6 for target cytotoxicity (% target) and T cell activation CD25+ (% live T cells).

15 Graphing and fitting of data were done in GraphPad Prism 6 using nonlinear regression with variable slope (four parameters) function using least squares method.

Table 11: Average EC_{50} calculated from T-cell mediated cytotoxicity assessment of GPRC5D x CD3 bispecific antibodies using H929 and MM1R target cells.

GPRC5D x CD3 EC_{50} (nM)			
Bispecific ID	H929 (n= 5)	MM1R (n = 3)	Ranking (Average) H929/MM1R
GCDB32	0.39 ± 0.3	0.12 ± 0.1	2/1
GCDB33	4.29 ± 1.28	0.8 ± 0.18	7/6
GCDB34	2.13 ± 0.93	0.77 ± 0.24	5/5
GCDB35	1.39 ± 0.84	0.74 ± 0.32	4/4
GCDB36	5.2 ± 0.93	4.22 ± 0.55	8/9
GCDB40	1.2 ± 0.92	0.52 ± 0.38	3/3
GCDB41	NA	2.58 ± 0.15	10/8
GCDB43	0.36 ± 0.41	0.21 ± 0.29	1/2

GCDB44	10.29 ± 2.65	4.41 ± 0.29	9/10
GCDB45	2.25 ± 1	0.91 ± 0.93	6/7

All bispecific antibodies are active in T-cell mediated H929 cell killing, with a range of potencies observed (Table 11). Similar rank order was observed for both cell lines. However a lower EC₅₀ was observed in MM1R cells. Interestingly, binding affinity was not necessarily
 5 correlative of potency in T-cell mediated cytotoxicity assay. For example, GCDB44 was the highest affinity binding bispecific antibody but the least potent in the cytotoxicity assay. While GCDB43, with similar, albeit slightly lower, binding affinity was the most potent in the cytotoxicity assay.

To assess the functional cross-reactivity with cyno GPRC5D, the panel of bispecific
 10 antibodies was then profiled for T-cell mediated cytotoxicity and T cell activation using cyno GPRC5D HEK293 cells (Fig. 5A and B). All bispecific antibodies were active in this assay though a range of potency was observed against the cyno GPRC5D+ expressing cells.

In vivo efficacy

A benchmarking *in vivo* study was then completed to understand the *in vivo* potency of
 15 these GPRC5D X CD3 bispecific antibodies. GCDB32 and GCDB35 (Fig. 5 A & B) were selected to test in a H929 prophylactic tumor model. H929 cells were implanted into NSG mice, one week following injection of human PBMCs. Treatment of the bispecific antibodies was initiated at the same time that the H929 cells were implanted and continued every 2 or 3 days
 20 (q2d or q3d) at 10 ug, 1 ug, and 0.1 ug / animal doses for a total of five treatments. Ten mice were used in each group and PBS included as the vehicle control. Treatment was stopped at day 11 and the study was terminated on day 25 (Fig. 6 A and B) or day 26 (Fig. 12 A-D). All the GPRC5D x CD3 antibodies tested in this prophylactic model showed 100% tumor growth inhibition at the 10 and 1 ug/ animal dose except for GCDB35 which showed round 80 % tumor
 25 growth inhibition at 1 ug/animal dose. At tenfold lower dose (0.1ug/animal) these bispecific antibodies showed varying degree of efficacy ranging from 10 to 80 % tumor growth inhibition.

In vitro T-Cell Dependent Cytotoxicity in presence of Fc blocker

To gain an understanding of the specificity of the GPRC5D targeting arm, the panel of GPRC5D x CD3 bispecific antibodies was then assessed in a T-cell redirection cytotoxicity assay in the presence of Fc block. This experiment was critical to understanding specificity as the target cells for bispecific antibodies are B-cells expressing Fcγ receptors capability of interacting with the Fc portion of the bispecific antibody in the *in vitro* assay. A shift in potency was observed in the T-cell mediated cytotoxicity assay for a number of the bispecific antibodies, with the greatest shift observed for GCDB40 and GCDB34 (Fig. 7A-7B).

Direct measurement of binding interactions between the four most potent bispecific antibodies (GCDB32, GCDB35, GCDB40, and GCDB43) and Fcγ receptors was then completed (Fig. 8A-8D). One round of Alpha Screen analysis was carried out for each of the Fcγ receptors and bispecific antibodies listed above. All samples were assayed in duplicate. On FcγRI, the four bispecific antibodies behave like B21M hIgG4 PAA. That is, they are no more competitive than matched isotype control. Similar differences were also observed between the hIgG1 WT control and the four bispecific antibodies on FcγRIIIa, with GCD43 most like the IgG4PAA isotype control and the other bispecific antibodies having slightly higher affinity for FcγRIIIa. On both FcγRIIa and FcγRIIb, the four bispecific antibodies compete in the following order: GCDB40 > GCDB32 > GCDB43 > GCDB35. GCDB40 is the most competitive or binds with highest affinity to FcγRIIa and FcγRIIb. In fact GCDB40 competes on FcγRIIa and FcγRIIb to the same extent as hIgG1 WT, corroborating the observed shift in potency observed in the T-cell mediated cytotoxicity assay when Fc block was included. Due to the unexpected interactions with FcγRIIa and FcγRIIb, GCDB40 was not advanced.

Competition Binning Assay

Anti-GPRC5D mAbs were assessed for competition binding with each other to the human GPCR5D cells. Briefly, cells were plated at 50,000 cells/well in 50uL of media and allowed to settle for 90 minutes at 37C. Wells were then blocked using 3% BSA for 1 hour at room temperature. mAbs were labeled with ruthenium (II) tris-bipyridine, N-hydroxysuccinimide (Ru-label) following standard procedures. In a separate 96 well plate, 5uM of competitor mAb was incubated with 50nM of Ru-labeled mAb. The blocking solution was removed from the cell plate and 25uL of mAb solution added. Plates were incubated for 1 hour at room temperature with shaking. After washing the plates thrice with PBS, 150uL of MSD

read buffer (without surfactant) was added and the binding of the Ru-labeled antibody detected using a MSD plate reader.

All mAbs belong to the same competition group, with only GC5B420 and GC5B421 not fully competed by GC5B81, GC5B285, and/or GC5B332 (Blocked < 70%) (Table 12). It is hypothesized that simultaneous binding of two mAbs to GPRC5D may be sterically impossible given the small size of the extracellular domain of GPRC5D compared to the size of a mAb.

Table 12: Competition binding epitope binning of anti-GPRC5D mAbs. Anti-GPRC5D mAbs were assessed for competition binding with each other to the human GPCR5D cells (+/- = Blocked < 70%).

mAb ID	GC5B81	GC5B164	GC5B285	GC5B332	GC5B420	GC5B421
GC5B81	+	+	+	+	+/-	+/-
GC5B164	+	+	+	+	+	+
GC5B285	+	+	+	+	+	+/-
GC5B332	+	+	+	+	+/-	+
GC5B420	+	+	+	+	+	+
GC5B421	+	+	+	+	+	+
GC5B243	+	+	+	+	+	+

EXAMPLE 4: GENERATION OF GPRC5D ANTIBODIES USING HYBRIDOMA TECHNOLOGY

Three Balb/c mice were immunized intradermally-base of tail with pCMV6-neo (CMV promoter) plasmid DNA expressing full length human GPRC5D on days 0, 10, and 20. Mice received final intraperitoneal and intravenous immunizations on day 59 with Rat Basophilic Leukemia (RBL) cells over-expressing full length human GPRC5D. On day 63, lymph nodes and spleens were harvested, enrichment of B cells was performed, and the cells were used to generate ~3500 mAb secreting hybridomas.

EXAMPLE 5: INITIAL CHARACTERIZATION OF GPRC5D ANTIBODIES OBTAINED THROUGH HYBRIDOMA TECHNOLOGY

GPRC5D Binding

The hybridomas were screened using FACS for binding to both RBL and human GPRC5D expressing RBL cells. The ratio of background adjusted MFI of each mAb binding to the GPRC5D_RBL cells relative to the non-transfected RBL cells was calculated and any sample with a binding ratio greater than 3 were considered potentially positive. Ninety nine hybridomas had a ratio greater than 3 and were advanced for v-region cloning. Thirty one mAbs sequences were identified, synthesized, expressed and purified. Two of the 31 mAbs exhibited binding to RBL_GPRC5D cells above background (Fig. 9A and 8B). GCDB390 and GCDB396 were selected for expression, purification and bispecific antibody generation with anti-CD3 arm CD3B219 to generate the bispecific antibodies GCDB46 and GCDB47, respectively.

T-Cell Dependent Cytotoxicity

GCDB46 and GCDB47 were assessed in the T-cell mediated cytotoxicity assay (Fig. 10A and 10B). Both bispecific antibodies were observed to be potent, with reported EC₅₀ of 0.67 and 0.1 nM respectively. On the basis of these data both mAbs were advanced to lead optimization along with the three phage derived mAbs.

EXAMPLE 6: HIT EVALUATION, SELECTION AND OPTIMIZATION

Five GPRC5D bispecific antibodies were selected for lead optimization (GCDB32, GCDB43, GCDB35, GCDB46, GCDB47) based on binding, function, cross-reactivity and selectivity data summarized in Table 13.

Table 13: Lead optimization data for GPRC5D x CD3 bispecific antibodies. Bispecific antibodies were assessed for binding, function, cross-reactivity and selectivity.

GPRC5D x CD3 ID	Selectivity	T cell mediated Target cell killing			
	Binding to GPRC5A, B, C	H929 EC ₅₀ ± SD (nM)	MM1R EC ₅₀ ± SD (nM)	H929 No Fc vs Fc Block	Cyno GPRC5D EC ₅₀ (nM)
GCDB32	No Binding	0.43 ± 0.29	0.12 ± 0.01	2.4	0.19
GCDB43	No Binding	0.39 ± 0.07	0.54	2.8	0.95
GCDB40	No Binding	1.2 ± 0.83	0.52 ± 0.31	5.1	10.61
GCDB35	Low	1.39 ± 0.75	0.74 ± 0.26	2.7	4.65
GCDB34	No Binding	2.01 ± 0.74	0.77 ± 0.09	6.3	1.14
GCDB45	No Binding	2.36 ± 0.84	0.91 ± 0.76	2.2	2.82
GCDB33	No Binding	5.21 ± 1.21	0.8 ± 0.15	1.5	0.2
GCDB36	Low	5.57 ± 0.47	4.22 ± 0.29		3.35
GCDB46	No Binding	0.67			
GCDB47	No Binding	0.1			

GCDB48	No Binding	0.17			Not Active
--------	------------	------	--	--	------------

Lead optimization was aimed at addressing potential post-translation modification (PTM) sequence risks for GCDB32 (GC5B81 parent mAb), GCDB43 (GC5B285 parent mAb) and GCDB35 (GC5B164 parent mAb) as outlined in Table 14.

5

Table 14: PTM mitigation sequence potential liabilities for Phage derived hits. The Hc CDR sequences are shown with potential PTM sequence liabilities are underlined. (SEQ ID NOs for each listed sequence are provided in parenthesis)

GPCR5D ID	GPCR5D Hc ID	Hc CDR1	Hc CDR2	HcCDR3
GC5B81	GC5H40	SYAIS (1)	GIPIFGTANYAQKFQG (5)	ESR <u>W</u> RGYKLD (9)
GC5B285	GC5H14	NYWMS (2)	GISYSGGSKYYA <u>D</u> SVKG (28)	AA <u>W</u> DFGRRVRLDY (30)
GC5B164	GC5H36	SY <u>W</u> IG (27)	IIYPGD <u>S</u> DRYSPSFQG (29)	VYSFGGRHKALFDY (11)

10

All of the PTM variants assayed for GC5B81 had a substantial reduction in binding affinity, demonstrating the criticality of this residue (HC W102) to the paratope (Table 15).

Table 15: CDR sequences and binding data of the GC5B81 PTM library. The site of mutation is underlined. Binding was classified as MFI >10,000 = ++; MFI >1,000 = +; MFI <1,000 (SEQ ID NOs for each listed sequence are provided in parenthesis)

15

GPCR5D ID	Heavy Chain ID	Mutation	Hc CDR1	Hc CDR2	Hc CDR3	Human GPCR5D (FACS)
GC5B427	GC5H199	W102Y	SYAIS (1)	GIPIFGTANYAQKFQG (5)	ESR <u>Y</u> RGYKLDY (31)	-
GC5B428	GC5H198	W102V	SYAIS (1)	GIPIFGTANYAQKFQG (5)	ESR <u>V</u> RGYKLDY (32)	-
GC5B430	GC5H196	W102G	SYAIS (1)	GIPIFGTANYAQKFQG (5)	ESR <u>G</u> RGYKLDY (33)	-
GC5B431	GC5H195	W102A	SYAIS (1)	GIPIFGTANYAQKFQG (5)	ESR <u>A</u> RGYKLDY (34)	-
GC5B429	GC5H197	W102F	SYAIS (1)	GIPIFGTANYAQKFQG (5)	ESR <u>F</u> RGYKLDY (35)	-

Binding studies identified a number of mutations for both GC5B285 and GC5B164 which retained binding to human GPCR5D (Tables 16 and 17).

20

Table 16: CDR sequences and binding data of the GC5B164 PTM library. Site of mutation is underlined. Binding was classified as MFI >10,000 = ++; MFI >1,000 = +; MFI <1,000. (SEQ ID NOs for each listed sequence are provided in parenthesis)

GPRC5D ID	Heavy Chain ID	Mutation	Hc CDR1	Hc CDR2	Hc CDR3	Human GPR5D Binding (FACS)
GC5B471	GC5H278	D55A, W33Y	SY <u>Y</u> IG (36)	IYPG <u>A</u> SDTRYSPSFQG (40)	VYSFGGRHKALFDY (11)	++
GC5B472	GC5H277	D55A, W33V	SY <u>V</u> IG (37)	IYPG <u>A</u> SDTRYSPSFQG (40)	VYSFGGRHKALFDY (11)	+
GC5B473	GC5H276	D55A, W33F	SY <u>F</u> IG (3)	IYPG <u>A</u> SDTRYSPSFQG (40)	VYSFGGRHKALFDY (11)	++
GC5B474	GC5H275	D55A, W33G	SY <u>G</u> IG (38)	IYPG <u>A</u> SDTRYSPSFQG (40)	VYSFGGRHKALFDY (11)	+
GC5B475	GC5H274	D55A, W33A	SY <u>A</u> IG (39)	IYPG <u>A</u> SDTRYSPSFQG (40)	VYSFGGRHKALFDY (11)	+
GC5B476	GC5H273	D55S, W33Y	SY <u>Y</u> IG (36)	IYPG <u>S</u> SDTRYSPSFQG (41)	VYSFGGRHKALFDY (11)	++
GC5B477	GC5H272	D55S, W33V	SY <u>V</u> IG (37)	IYPG <u>S</u> SDTRYSPSFQG (41)	VYSFGGRHKALFDY (11)	+
GC5B478	GC5H271	D55S, W33F	SY <u>F</u> IG (3)	IYPG <u>S</u> SDTRYSPSFQG (41)	VYSFGGRHKALFDY (11)	++
GC5B479	GC5H270	D55S, W33G	SY <u>G</u> IG (38)	IYPG <u>S</u> SDTRYSPSFQG (41)	VYSFGGRHKALFDY (11)	+
GC5B480	GC5H269	D55S, W33A	SY <u>A</u> IG (39)	IYPG <u>S</u> SDTRYSPSFQG (41)	VYSFGGRHKALFDY (11)	+
GC5B481	GC5H268	D55K, W33Y	SY <u>Y</u> IG (36)	IYPG <u>K</u> SDTRYSPSFQG (7)	VYSFGGRHKALFDY (11)	++
GC5B482	GC5H267	D55K, W33V	SY <u>V</u> IG (37)	IYPG <u>K</u> SDTRYSPSFQG (7)	VYSFGGRHKALFDY (11)	++
GC5B483	GC5H266	D55K, W33F	SY <u>F</u> IG (3)	IYPG <u>K</u> SDTRYSPSFQG (7)	VYSFGGRHKALFDY (11)	++
GC5B484	GC5H265	D55K, W33G	SY <u>G</u> IG (38)	IYPG <u>K</u> SDTRYSPSFQG (7)	VYSFGGRHKALFDY (11)	+
GC5B485	GC5H264	D55K, W33A	SY <u>A</u> IG (39)	IYPG <u>K</u> SDTRYSPSFQG (7)	VYSFGGRHKALFDY (11)	+
GC5B486	GC5H263	D55E, W33Y	SY <u>Y</u> IG (36)	IYPG <u>E</u> SDTRYSPSFQG (42)	VYSFGGRHKALFDY (11)	++
GC5B487	GC5H262	D55E, W33V	SY <u>V</u> IG (37)	IYPG <u>E</u> SDTRYSPSFQG (42)	VYSFGGRHKALFDY (11)	+
GC5B488	GC5H261	D55E, W33F	SY <u>F</u> IG (3)	IYPG <u>E</u> SDTRYSPSFQG (42)	VYSFGGRHKALFDY (11)	++
GC5B489	GC5H260	D55E, W33G	SY <u>G</u> IG (38)	IYPG <u>E</u> SDTRYSPSFQG (42)	VYSFGGRHKALFDY (11)	-
GC5B490	GC5H259	D55E, W33A	SY <u>A</u> IG (39)	IYPG <u>E</u> SDTRYSPSFQG (42)	VYSFGGRHKALFDY (11)	+
GC5B491	GC5H258	D55Y, W33Y	SY <u>Y</u> IG (36)	IYPG <u>Y</u> SDTRYSPSFQG (43)	VYSFGGRHKALFDY (11)	++
GC5B492	GC5H257	D55Y, W33V	SY <u>V</u> IG (37)	IYPG <u>Y</u> SDTRYSPSFQG (43)	VYSFGGRHKALFDY (11)	++
GC5B493	GC5H256	D55Y, W33F	SY <u>F</u> IG (3)	IYPG <u>Y</u> SDTRYSPSFQG (43)	VYSFGGRHKALFDY (11)	++
GC5B494	GC5H255	D55Y, W33G	SY <u>G</u> IG (38)	IYPG <u>Y</u> SDTRYSPSFQG (43)	VYSFGGRHKALFDY (11)	+
GC5B495	GC5H254	D55Y, W33A	SY <u>A</u> IG (39)	IYPG <u>Y</u> SDTRYSPSFQG (43)	VYSFGGRHKALFDY (11)	+

GC5B496	GC5H253	W33Y	SY <u>Y</u> IG (36)	IIPGDS <u>D</u> TRYSPSFQG (29)	VYSFGGRHKALFDY (11)	++
GC5B497	GC5H252	W33V	SY <u>V</u> IG (37)	IIPGDS <u>D</u> TRYSPSFQG (29)	VYSFGGRHKALFDY (11)	++
GC5B498	GC5H251	W33F	SY <u>F</u> IG (3)	IIPGDS <u>D</u> TRYSPSFQG (29)	VYSFGGRHKALFDY (11)	++
GC5B499	GC5H250	W33G	SY <u>G</u> IG (38)	IIPGDS <u>D</u> TRYSPSFQG (29)	VYSFGGRHKALFDY (11)	-
GC5B500	GC5H249	W33A	SY <u>A</u> IG (39)	IIPGDS <u>D</u> TRYSPSFQG (29)	VYSFGGRHKALFDY (11)	-
GC5B501	GC5H248	D55A	SYWIG (27)	IIPG <u>A</u> SDTRYSPSFQG (40)	VYSFGGRHKALFDY (11)	++
GC5B502	GC5H247	D55S	SYWIG (27)	IIPG <u>S</u> SDTRYSPSFQG (41)	VYSFGGRHKALFDY (11)	++
GC5B503	GC5H246	D55K	SYWIG (27)	IIPG <u>K</u> SDTRYSPSFQG (7)	VYSFGGRHKALFDY (11)	++
GC5B504	GC5H245	D55E	SYWIG (27)	IIPG <u>E</u> SDTRYSPSFQG (42)	VYSFGGRHKALFDY (11)	++
GC5B505	GC5H244	D55Y	SYWIG (27)	IIPG <u>Y</u> SDTRYSPSFQG (43)	VYSFGGRHKALFDY (11)	++

Table 17: CDR sequences and binding data of the GC5B285 PTM library. Site of mutation is underlined. Binding was classified as MFI >10,000 = ++; MFI >1,000 = +; MFI <1,000.

5 (SEQ ID NOs for each listed sequence are provided in parenthesis)

GPRC5D ID	Heavy Chain ID	Mutation	Hc CDR1	Hc CDR2	Hc CDR3	Human GPRC5D (FACS)
GC5B463	GC5H234	D62A,W101Y	NYWMS (2)	GISYSGGSKYYA <u>A</u> SVKG (44)	AA <u>Y</u> DFGRRAVRLDY (48)	++
GC5B432	GC5H228	D62S,W101V	NYWMS (2)	GISYSGGSKYYA <u>S</u> SVKG (6)	AA <u>Y</u> DFGRRAVRLDY (49)	++
GC5B465	GC5H227	D62S,W101F	NYWMS (2)	GISYSGGSKYYA <u>S</u> SVKG (6)	AA <u>F</u> DFGRRAVRLDY (97)	++
GC5B433	GC5H223	D62K,W101V	NYWMS (2)	GISYSGGSKYYA <u>K</u> SVKG (45)	AA <u>Y</u> DFGRRAVRLDY (49)	++
GC5B434	GC5H222	D62K,W101F	NYWMS (2)	GISYSGGSKYYA <u>K</u> SVKG (45)	AA <u>F</u> DFGRRAVRLDY (97)	++
GC5B435	GC5H219	D62E,W101Y	NYWMS (2)	GISYSGGSKYYA <u>E</u> SVKG (46)	AA <u>Y</u> DFGRRAVRLDY (48)	+
GC5B436	GC5H217	D62E,W101F	NYWMS (2)	GISYSGGSKYYA <u>E</u> SVKG (46)	AA <u>F</u> DFGRRAVRLDY (97)	+
GC5B461	GC5H216	D62E,W101G	NYWMS (2)	GISYSGGSKYYA <u>E</u> SVKG (46)	AA <u>G</u> DFGRRAVRLDY (50)	-
GC5B462	GC5H215	D62E,W101A	NYWMS (2)	GISYSGGSKYYA <u>E</u> SVKG (46)	AA <u>A</u> DFGRRAVRLDY (51)	++
GC5B437	GC5H214	D62Y,W101Y	NYWMS (2)	GISYSGGSKYYA <u>Y</u> SVKG (47)	AA <u>Y</u> DFGRRAVRLDY (48)	+
GC5B438	GC5H213	D62Y,W101V	NYWMS (2)	GISYSGGSKYYA <u>Y</u> SVKG (47)	AA <u>Y</u> DFGRRAVRLDY (49)	+
GC5B439	GC5H212	D62Y,W101F	NYWMS (2)	GISYSGGSKYYA <u>Y</u> SVKG (47)	AA <u>F</u> DFGRRAVRLDY (97)	+
GC5B440	GC5H211	D62Y,W101G	NYWMS (2)	GISYSGGSKYYA <u>Y</u> SVKG (47)	AA <u>G</u> DFGRRAVRLDY (50)	-
GC5B441	GC5H210	D62Y,W101A	NYWMS (2)	GISYSGGSKYYA <u>Y</u> SVKG (47)	AA <u>A</u> DFGRRAVRLDY (51)	+

GC5B442	GC5H209	W101Y	NYWM S (2)	GISYSGGSKYYADSVK G (28)	AAVDFGRRVRLDY (48)	+
GC5B443	GC5H208	W101V	NYWM S (2)	GISYSGGSKYYADSVK G (28)	AAVDFGRRVRLDY (49)	+
GC5B444	GC5H207	W101F	NYWM S (2)	GISYSGGSKYYADSVK G (28)	AAEDFGRRVRLDY (97)	+
GC5B464	GC5H206	W101G	NYWM S (2)	GISYSGGSKYYADSVK G (28)	AAGDFGRRVRLDY (50)	+
GC5B445	GC5H205	W101A	NYWM S (2)	GISYSGGSKYYADSVK G (28)	AAADFGRRAVRLDY (51)	+
GC5B446	GC5H204	D62A	NYWM S (2)	GISYSGGSKYYAASVK G (44)	AAWDFGRRVRLD Y (30)	+
GC5B447	GC5H203	D62S	NYWM S (2)	GISYSGGSKYYA ~~S~~ SVKG (6)	AAWDFGRRVRLD Y (30)	+
GC5B448	GC5H202	D62K	NYWM S (2)	GISYSGGSKYYA ~~K~~ SVK G (45)	AAWDFGRRVRLD Y (30)	+
GC5B449	GC5H201	D62E	NYWM S (2)	GISYSGGSKYYA ~~E~~ SVKG (46)	AAWDFGRRVRLD Y (30)	+
GC5B450	GC5H200	D62Y	NYWM S (2)	GISYSGGSKYYA ~~Y~~ SVK G (47)	AAWDFGRRVRLD Y (30)	+

On the basis of this binding data, selected mAbs were generated as GPRC5D x CD3 bispecific antibodies and assessed for T-cell mediated cytotoxicity of H929 cells (Table 18).

5 **Table 18:** Functional Activity of selected GPRC5D PTM variants for GCDB164 and GCDB285

Parent GPRC5D mAb ID	GPRC5D Protein AA ID	GPRC5D x CD3 ID	H929 EC ₅₀ (nM)	Selected as Lead
GC5B285	GC5B432	GCDB50	1.3	
	GC5B433	GCDB51	0.81	
	GC5B434	GCDB52	2.81	
	GC5B465	GCDB53	0.24	XX
	GC5B463	GCDB54	3.09	
GC5B164	GC5B471	GCDB57	3.87	
	GC5B476	GCDB58	1.24	
	GC5B478	GCDB59	1.54	
	GC5B481	GCDB60	2.94	
	GC5B483	GCDB61	1.51	XX
	GC5B493	GCDB62	3.1	

A range of potency was observed in the T-cell mediated cytotoxicity assay that was not necessarily predicted by the observed binding affinity. For example, GC5B465 and GC5B463 bound with similar affinity to human GPRC5D differ only in the sequence of 2 amino acids (Table 17) and were observed to have a 12.5 fold difference in potency as GPRC5D x CD3 bispecific Abs (Table 18). On the basis of the functional data GC5B465 and GC5B483 were

selected as optimized sequences for GC5B285 (GCDB43 as CD3 bispecific) and GC5B164 (GCDB35 as CD3 bispecific).

Human framework adaptation was also completed for the murine hybridoma derived GPCR5D mAbs (GC5B390 and GC5B396, or GCDB46 and 47 as CD3 bispecific, respectively).

- 5 Binding studies identified a number of frameworks for GC5B396 and one framework for GC5B390 which retained binding to human GPCR5D (Table 19).

Table 19: Binding and functional data for the human framework adaptation of hybridoma derived anti-GPCR5D mAb libraries. Binding was classified as MFI >10,000 = +++; MFI >5,000 = ++; MFI > 1,000 = +; MFI <1,000.

GPRC5D Parent mAb ID	GPRC5D mAb AA ID	Heavy Chain ID	Light Chain ID	Human GPRC5D (FACS)	GPRC5D x CD3 ID	H929 EC50 (nM)
GPRC5D B396	GC5B541	GC5H241	GC5L58	+		
	GC5B540	GC5H240	GC5L58	+++	GCDB69	0.58
	GC5B539	GC5H242	GC5L58	++		
	GC5B538	GC5H243	GC5L58	++		
	GC5B537	GC5H241	GC5L57	+		
	GC5B536	GC5H240	GC5L57	+++	GCDB68	2.51
	GC5B535	GC5H242	GC5L57	+		
	GC5B534	GC5H243	GC5L57	++		
	GC5B533	GC5H241	GC5L56	+		
	GC5B532	GC5H240	GC5L56	+++	GCDB67	0.61
	GC5B531	GC5H242	GC5L56	++		
	GC5B530	GC5H243	GC5L56	+++	GCDB66	1.41
	GC5B529	GC5H241	GC5L55	++		
	GC5B528	GC5H240	GC5L55	+++	GCDB65	1.01
	GC5B527	GC5H242	GC5L55	-		
	GC5B526	GC5H243	GC5L55	++	GCDB64	1.5
GPRC5D B390	GC5B525	GC5H279	GC5L53	-		
	GC5B524	GC5H237	GC5L53	-		
	GC5B523	GC5H238	GC5L53	-		
	GC5B522	GC5H236	GC5L53	-		
	GC5B521	GC5H279	GC5L52	-		
	GC5B520	GC5H237	GC5L52	-		
	GC5B519	GC5H238	GC5L52	-		
	GC5B518	GC5H236	GC5L52	-		
	GC5B517	GC5H279	GC5L51	+		
	GC5B516	GC5H237	GC5L51	+		
	GC5B515	GC5H238	GC5L51	++	GCDB63	0.52
	GC5B514	GC5H236	GC5L51	+		
	GC5B513	GC5H279	GC5L50	-		
	GC5B512	GC5H237	GC5L50	+		
	GC5B511	GC5H238	GC5L50	+		

	GC5B510	GC5H236	GC5L50	+		
	GC5B509	GC5H279	GC5L49	-		
	GC5B508	GC5H237	GC5L49	+		
	GC5B507	GC5H238	GC5L49	+		
	GC5B506	GC5H236	GC5L49	+		

On the basis of the binding data several anti-GPCR5D mAbs were generated as CD3 bispecific antibodies and assessed for T-cell mediated cytotoxicity of H929 cells (Table 18). Functional analysis identified GCDB63, GCDB67, and GCDB69 as potent fully humanized GPRC5D x CD3 bispecific antibodies. On the basis of these data, the corresponding anti-GPCR5D mAbs, GC5B515, GC5B532, and GC5B540 were selected as the fully humanized sequences for GC5B390 and GC5B391.

Additional lead optimization of the fully humanized sequences was then completed aimed at addressing potential post-translation modification sequence risks for GC5B515, GC5B532, and GCDB540. A G56S mutation was generated in the heavy chain sequence to remove a potential deamidation risk for GC5B515 (Table 20).

Table 20: Binding and Functional Activity of selected GPRC5D PTM variants for GC5B532, GC5B540 and GCDB515.

GPRC5D Parent mAb ID	GPRC5D mAb AA ID	Mutation	Human GPRC5D (FACS)	GPRC5D x CD3 ID	H929 EC50 (nM)
GC5B540	GC5B590	M64K	+++		
	GC5B592	G99A	+		
	GC5B594	M64K, G99A	+		
GC5 B532	GC5B591	M64K	+++		
	GC5B593	G99A	+		
	GC5B595	M64K, G99A	+		
GC5 B515	GC5B596	G56S	++	GCDB72	0.15

The heavy chains for GC5B532 and GC5B540 contained both a potential isomerization and an oxidation risk. M64K and G99A mutations were generated to ameliorate this risk (Table 20). All of the variants tested with the G99A mutation had a substantial reduction in binding affinity, while the M64K and G56A variants were not impacted. Based on the binding data alone GC5B596 was progressed to functional assessment and demonstrated potency as a CD3 bispecific (GCDB72) in T-cell mediated cytotoxicity assays.

Thus, four GPRC5D bispecific mAbs were selected for additional characterization: GCDB32, GCDB53, GCDB61, and GCDB72. Depicted below in Tables 21 and 22 are the CDR and heavy and light chain sequences of the GPRC5D mAbs used for the generation of the bispecific molecules.

5 **Table 21.** CDR sequences of 4 GPRC5D mAb candidates that showed binding against human and cyno GPRC5D and that were functional when generated as CD3 bispecifics.

ID	HC-CDR1	HC-CDR2	HC-CDR3	LC-CDR1	LC-CDR2	LC-CDR3
GC5B81	SYAIS (SEQ ID NO 1)	GIPIFGTANYAQKFQG (SEQ ID NO 5)	ESRWRGYKLD (SEQ ID NO 9)	RASQSISSYLN (SEQ ID NO 13)	AASSLQS (SEQ ID NO 16)	QQSYSTPLT (SEQ ID NO 19)
GC5B465	NYWMS (SEQ ID NO 2)	GISYSGGSKYYASSVKG (SEQ ID NO 6)	AAFDGRRRAVRD (SEQ ID NO 10)	RASQSISSYLN (SEQ ID NO 13)	AASSLQS (SEQ ID NO 16)	QQSYSTPLT (SEQ ID NO 19)
GC5B483	SYFIG (SEQ ID NO 3)	IIYPGKSDTRYSPSFQG (SEQ ID NO 7)	VYSFGGRHKALFDY (SEQ ID NO 11)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B596	GYTMN (SEQ ID NO 4)	LINPYNSDTNYAQKLQG (SEQ ID NO 8)	VALRVALDY (SEQ ID NO 12)	KASQNVATHVG (SEQ ID NO 15)	SASYRYS (SEQ ID NO 18)	QQYNRYPYT (SEQ ID NO 21)

Table 22: Heavy and light chain variable region sequences of 4 GPRC5D mAb candidates that showed binding against human and cyno GPRC5D and that were functional when generated as CD3 bispecifics.

mAb AA ID	VH Amino Acid Sequence	SEQ ID NO	VL Amino Acid Sequence	SEQ ID NO
GC5B81	QVQLVQSGAEVKKPGSSV KVCKASGGTFSSYAISW VRQAPGGGLEWMGGIPI FGTANYAQKFQGRVTITA DESTSTAYMELSSLRSED TAVYYCARESRWRGYKLD YWGGQGLTVTVSS	52	DIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQ QKPGKAPKLLIYAASSLQ SGVPSRFSGSGSGTDFLT ISSLPEDFATYYCQQSYS TPLTFGQGTKVEIK	56
GC5B465	EVQLLESGGGLVQPGGSL RLSCAASGFTFSNYWMS WVRQAPGKGLEWVSGIS YSGGSKYYASSVKGRFTI SRDNSKNTLYLQMNSLRA EDTAVYYCAKAAFDGR RAVRDLYWGQGLTVTVS	53	DIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQ QKPGKAPKLLIYAASSLQ SGVPSRFSGSGSGTDFLT ISSLPEDFATYYCQQSYS TPLTFGQGTKVEIK	56

	S			
GC5B483	EVQLVQSGAEVKKPGESL KISCKGSGYSFTSYFIGWV RQMPGKGLEWMGIIYPG KSDTRYSPSFQGGVTISAD KSISTAYLQWSSLKASDT AMYYCARVYSFGGRHKA LFDYWGQGTLVTVSS	54	EIVLTQSPATLSLSPGERA TLSCRASQSVSSYLAWYQ QKPGQAPRLLIYDASNRA TGIPARFSGSGSGTDFTLTI SSLEPEDFAVYYCQQRSN WPLTFGQGGTKVEIK	57
GC5B596	QVQLVQSGAEVKKPGAS VKVSCKASGYSFTGYTM NWVRQAPGQGLEWMGLI NPYNSTNYAQKLQGRV TMTTDTSTSTAYMELRSL RSDDTAVYYCARVALRV ALDYWGQGTLVTVSS	55	DIQMTQSPSSLSASVGRD VTITCKASQNVATHVGW YQQKPGKAPKRLIYSASY RYSQVPSRFSGSGSGTEFT LTISNLPEDFATYYCQQ YNRYPTYTFGQGGTKLEIK	58

EXAMPLE 7: PREPARATION OF GPRC5D AND CD3 ANTIBODIES IN A BISPECIFIC FORMAT IN IgG4 S228P, L234A, L235A

The four monospecific GPRC5D antibodies (see table 21) were expressed as IgG4, having Fc substitutions S228P, L234A, and L235A or S228P, L234A, L235A, F405L, and R409K (CD3 arm) (numbering according to EU index). A monospecific anti-CD3 antibody CD3B219 was also generated comprising the VH and VL regions having the heavy chain of SEQ ID NO: 25 and the light chain of SEQ ID NO: 26 and IgG4 constant region with S228P, L234A, L235A, F405L, and R409K substitutions.

The monospecific antibodies were purified using standard methods using a Protein A column (HiTrap MabSelect SuRe column). After elution, the pools were dialyzed into D-PBS, pH 7.2.

Bispecific GPRC5D x CD3 antibodies were generated by combining a monospecific CD3 mAb and a monospecific GPRC5D mAb in *in-vitro* Fab arm exchange (as described in WO2011/131746). Briefly, at about 1-20 mg/mL at a molar ratio of 1.08:1 of anti-GPRC5D/anti-CD3 antibody in PBS, pH 7-7.4 and 75 mM 2-mercaptoethanolamine (2-MEA) was mixed together and incubated at 25-37 °C for 2-6 hours, followed by removal of the 2-MEA via dialysis, diafiltration, tangential flow filtration and/or spin cell filtration using standard methods.

Heavy and Light chains for the GPRC5D x CD3 bispecific Abs are shown below in Table 23.

Table 23. Heavy and Light Chain Sequences for bispecific Abs IgG4-PAA

Ab		Amino Acid Sequence
GCDB32	Heavy chain 1 CD3B219 (SEQ ID NO:25)	EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQ APGKGLEWVARIRSKYNNYATYYAASVKGRFTISRDDSK NSLYLQMNSLKTEDTAVYYCARHGNFGNSYVSWFAYWG QGTLVTVSS
	Light Chain 1 CD3B219 (SEQ ID NO:26)	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQ QKPGQAPRGLIGGTNKRAPGTPARFSGSLLGGKAALTLTG VQPEDEAEYYCALWYSNLWVFGGGTKLTVL
	Heavy chain 2 GC5B81 (SEQ ID NO:52)	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQ APGQGLEWMGGIPIFGTANYAQKFQGRVTITADESTSTA YMELSSLRSED TAVYYCARESRWRGYKLDYWGQGTLVT VSS
	Light Chain 2 GC5B81 (SEQ ID NO:56)	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPG KAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDF ATYYCQQSYSTPLTFGQGTKVEIK
GCDB53	Heavy chain 1 CD3B219 (SEQ ID NO:25)	EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQ APGKGLEWVARIRSKYNNYATYYAASVKGRFTISRDDSK NSLYLQMNSLKTEDTAVYYCARHGNFGNSYVSWFAYWG QGTLVTVSS
	Light Chain 1 CD3B219 (SEQ ID NO:26)	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQ QKPGQAPRGLIGGTNKRAPGTPARFSGSLLGGKAALTLTG VQPEDEAEYYCALWYSNLWVFGGGTKLTVL
	Heavy chain 2 GC5B465 (SEQ ID NO:53)	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYWMSWVRQ APGKGLEWVSGISYSGGSKYYASSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAKAAFDGRRAVRLDYWGQGT LTVSS
	Light Chain 2	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPG KAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDF

	GC5B465 (SEQ ID NO:56)	ATYYCQQSYSTPLTFGQGTKVEIK
GCDB61	Heavy chain 1 CD3B219 (SEQ ID NO:25)	EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQ APGKGLEWVARIRSKYNNYATYYAASVKGRFTISRDDSK NSLYLQMNSLKTEDTAVYYCARHGNFGNSYVSWFAYWG QGTLVTVSS
	Light Chain 1 CD3B219 (SEQ ID NO:26)	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQ QKPGQAPRGLIGGTNKRAPGTPARFSGSLLGGKAALTLTG VQPEDEAEYYCALWYSNLWVFGGGTKLTVL
	Heavy chain 2 GC5B483 (SEQ ID NO:54)	EVQLVQSGAEVKKPAGESLKISCKGSGYSFTSYFIGWVRQM PGKGLEWMGIIYPGKSDTRYSPSFQGGVTISADKSISTAYL QWSSLKASDTAMYYCARVYSFGGRHKALFDYWGGQTLV TVSS
	Light Chain 2 GC5B483 (SEQ ID NO:57)	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPG QAPRLLIYDASNRAITGIPARFSGSGSGTDFTLTISSLEPEDF AVYYCQQRSNWPLTFGQGTKVEIK
GCDB72	Heavy chain 1 CD3B219 (SEQ ID NO:25)	EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQ APGKGLEWVARIRSKYNNYATYYAASVKGRFTISRDDSK NSLYLQMNSLKTEDTAVYYCARHGNFGNSYVSWFAYWG QGTLVTVSS
	Light Chain 1 CD3B219 (SEQ ID NO:26)	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQ QKPGQAPRGLIGGTNKRAPGTPARFSGSLLGGKAALTLTG VQPEDEAEYYCALWYSNLWVFGGGTKLTVL
	Heavy chain 2 GC5B596 (SEQ ID NO:55)	QVQLVQSGAEVKKPGASVKVSCKASGYSTGYTMNWVR QAPGGGLEWMGLINPYNSDTNYAQKLQGRVTMTTDTSTS TAYMELRSLRSDDTAVYYCARVALRVALDYWGQGLVT VSS
	Light Chain 2 GC5B596 (SEQ ID NO:58)	DIQMTQSPSSLSASVGDRTITCKASQNVATHVGVYQQK PGKAPKRLIYSASYRSGVPSRFSGSGSGTEFTLTISNLPQ DFATYYCQQYNRYPYTFGQGTKLEIK

EXAMPLE 8: FUNCTIONAL CHARACTERIZATION OF GCDB32, GCDB53, GCDB61, AND GCDB72

GCDB32, GCDB53, GCDB61, and GCDB72 were assessed for binding to murine GPRC5D (Table 24). All four bispecific antibodies bound to murine GPRC5D, with a range of binding affinities observed.

Table 24: Binding of anti-GPCR5D x CD3 antibodies to murine GPCR5D.

GPRC5D x CD3 ID	Murine GPRC5D FACS binding (MFI)
GCDB32	1151
GCDB53	2658
GCDB61	3552
GCDB72	481

Cross-reactivity with cyno GPRC5D was also assessed using the T-cell redirection cytotoxicity assay of over-expressed human and cyno GPRC5D cell lines (Table 25). One GPRC5D x CD3 bispecific antibody was equipotent against human and cyno GPRC5D (GCDB32), while the other bispecifics tested were less potent at inducing cytotoxicity of cyno GPRC5D than human GPRC5D.

Table 25: Functional activity of lead GPRC5D x CD3 antibodies against human and cyno GPRC5D-expressing HEK cells.

GPRC5D x CD3 ID	Human GPRC5D HEK cells (EC ₅₀ nM)	Cyno GPRC5D HEK cells (EC ₅₀ nM)
GCDB32	0.04	0.07
GCDB53	0.08	0.36
GCDB61	0.03	1.22
GCDB72	0.03	3.41

Additional characterization was aimed at understanding in vitro binding (Fig. 11A-11E) and potency (Table 26).

Table 26: T-cell mediated cytotoxicity of several human GPRC5D-expressing B cell lines by lead GPRC5D x CD3 antibodies.

GPRC5D x CD3 ID	H929 cells EC ₅₀ nM (Rank order)	MM1R EC ₅₀ nM (Rank order)	OPM2 EC ₅₀ nM (Rank order)	LP-1 EC ₅₀ nM (Rank Order)
GCDB32	0.45 (3)	0.04 (1)	0.98 (2)	1.02 (2)
GCDB53	0.69 (4)	0.28 (4)	2.83 (4)	1.58 (3)
GCDB61	0.39 (2)	0.06 (3)	1.46 (3)	1.67 (4)
GCDB72	0.22 (1)	0.04 (1)	0.77 (1)	0.7 (1)

5 While a range of binding affinities were observed, with GCDB61 the strongest binder and GCDB72 & GCDB32 the weakest binders, the *in vitro* potencies were very similar for the panel of bispecific antibodies. However, based on rank order analysis GCDB72 was the most potent across the various B cell lines analyzed. *Ex vivo* binding and potency experiments using patient derived MNCs yielded larger similar results to the *in vitro* assays (Table 27 and Fig. 15A and
10 15B).

Table 27. GPCR5D x CD3 bispecific antibody binding, T-cell mediated cytotoxicity, and T-cell activation of MM patient derived MNCs.

Patient	GPRC5D x CD3 AA ID	Binding EC ₅₀ nM	Cytotoxicity EC ₅₀ nM	T cell activation EC ₅₀ nM
MM303BM	GCDB32	16.1	0.36	0.18
	GCDB53	3.98	0.42	0.17
	GCDB61	5.53	0.89	0.22
	GCDB72	27.5	0.30	0.12
MM305BM	GCDB32	70.4	0.26	0.16
	GCDB53	10.2	0.26	0.19
	GCDB61	6.93	0.6	0.22
	GCDB72	65.9	0.28	0.12

15 GCDB61 had the highest binding affinity to patient MNCs while GCDB72 & GCDB32 were the weakest binders. Again, even though differences in binding affinity were observed, all bispecific antibodies demonstrated sub-nanomolar efficacy in T-cell redirected cytotoxicity assays. The molecules were virtually indistinguishable on the basis of *in vitro* and *ex vivo* potency however, *in vivo* data provided differentiation (Fig. 12A-12D).
20

H929 cells were implanted into NSG mice, one week following injection of human PBMCs. Treatment of the bispecific antibodies was initiated at the same time that the H929 cells

were implanted and continued every 2 or 3 days (q2d or q3d) at 10 ug, 1 ug, and 0.1 ug / animal doses for a total of five treatments. Ten mice were used in each group and PBS included as the vehicle control. Treatment was stopped at day 11 and the study was terminated on day 26 (Fig. 12 A-D). All the GPRC5D x CD3 bispecific molecules tested in this prophylactic model showed 100% tumor growth inhibition at the 10 and 1 ug/animal dose. Differentiation was observed at the lowest dose, 0.1ug/animal, with GCDB72 demonstrating superiority to the other bispecific antibodies tested with 80% tumor growth inhibition observed.

EXAMPLE 9: GPRC5D ANTIBODY BINDING PROFILE ON GPRC5D⁺ MM1.R CELL LINE

The binding affinities of the GPRC5D antibodies on GPRC5D⁺ human MM cell line (MM1.R, purchased from ATCC (American Type Culture Collection)) were measured using FACS. Fig. 13 shows that all the lead antibodies bound to GPRC5D expressing MM.1R cells in a dose dependent manner with EC₅₀ values ranging from 0.10 nM to 135 nm all of which except GC5B602 are significantly lower than the values of commercial antibody FAB6300 (R& D Systems Cat No. FAB6300A, Clone No. 571961) which yielded an EC₅₀ value of 121.7 nM.

GPRC5D⁺ MM1.R cell lines were stained for 60 minutes with various concentrations of lead antibodies to measure the surface binding profiles (n=3). Phycoerythrin labelled human IgG4Fc was used as a secondary antibody to capture the signal (Southern Biotech, clone HP6025, Cat # 9200-09). Binding is expressed as normalized geometrical mean fluorescence intensity as determined by FACS. Graphing and fitting of data were done in GraphPad Prism 6 using nonlinear regression with variable slope (four parameters) and least square fit method.

In addition, the GPRC5D mAb GC5M481 binding profile was assessed in comparison with the commercial antibody using three GPRC5D⁺ (JIM3,OPM-2, and MM.1R; Cell lines purchased from ATCC) multiple myeloma cell lines (Fig. 14A). Further, the GPRC5D mAb (GC5M481) was profiled for cyno cross-reactivity using a cyno-GPRC5D expressing Daudi cells which showed strong binding compared to the parental cells (Fig. 14A). Also, five GPRC5D x CD3 bispecific antibodies (GCDB32, GCDB48, GCDB53, GCDB61 and GCDB72) when evaluated for the binding potential using GPRC5D⁺ (JIM3, OPM-2 and MM1.R) cell lines (Fig. 14B) showed a significant binding as evident by the shift in the histogram (black solid trace) compared to the isotype control (dotted, gray filled trace).

EXAMPLE 10: ANTI-TUMOR EFFICACY OF GCDB72 AGAINST SUBCUTANEOUS MM.1S HUMAN MULTIPLE MYELOMA XENOGRAFTS IN PBMC-HUMANIZED NSG MICE.

5 This *in vivo* study was performed to determine the efficacy of GCDB72 against established MM.1S human multiple myeloma (MM) xenografts in PBMC humanized NSG mice. Female NSG mice of similar weight and age were subcutaneously (sc) implanted with MM.1S human MM cells (1×10^7 cells in 200 μ L PBS per mouse) on the right dorsal hind flank on study day 0. On day 7 post tumor cell implantation, 1×10^7 human PBMC (in 200 μ L PBS) were
10 injected intravenously via the lateral tail vein. Treatments were initiated on day 15 when mean tumor volume was approximately $72 - 78 \text{ mm}^3$, each mouse received intravenous (iv) administration of PBS or GCDB72 DuoBody antibody at 0.1 μ g (0.005 mg/kg), 1 μ g (0.05 mg/kg), 10 μ g (0.5 mg/kg) and 50 μ g (2.5 mg/kg). Null DuoBody controls, CD3 x Null and Null x GPRC5D, were each dosed at 10 μ g per mouse. Treatments were administered
15 approximately every three days (q3d) for a total of seven doses. Robust anti-tumor efficacy was observed with the two high doses (10 μ g and 50 μ g) of GCDB72 where MM.1S sc tumors completely regressed in 100% (10 of 10 per group) of the animals by the end of the study (Fig. 16). Moreover, the 1 μ g per mouse dose significantly inhibited tumor growth by 65% ($p \leq 0.0001$) compared to PBS treated tumors, while the 0.1 μ g dose had little effect (TGI=19.3%, $p=0.0023$).
20 The effect of CD3 x Null was not considered efficacious (TGI=28%, $p \leq 0.0001$) and Null x GPRC5D had negligible effect of 3.1% TGI, $p=0.9971$. TGI was determined on day 36 when there at least 80% viable animals per group. Significant body weight loss and/or mortality began to manifest post day 36 due to GVHD (Fig. 17). The study was terminated on day 43 when there were 60% or less animals remaining in the groups.

25

EXAMPLE 11: ANTIBODY-DEPENDENT CELL-MEDIATED CYTOTOXICITY (ADCC) OF GPRC5D ANTIBODIES

A panel of anti-human GPRC5D mAbs were generated as IgG1 mAbs. In addition, a new panel of anti-human GPRC5D mAbs were generated as described in Example 2. Depicted below
30 in Tables 28 and 29 are the CDR and heavy and light chain variable region sequences of the new GPRC5D mAbs. These new antibodies were used for the generation of bispecific CD3 molecules

as described in Example 7 and also incorporated as IgG1 mAbs for ADCC activity assessment.

Table 28. CDR sequences of new panel of GPRC5D antibodies

ID	HC-CDR1	HC-CDR2	HC-CDR3	LC-CDR1	LC-CDR2	LC-CDR3
GC5B382	DYGMH (SEQ ID NO 61)	AIKYSGGSTYYADSVKG (SEQ ID NO 67)	RAESGPGLDY (SEQ ID NO 72)	KSSQSVLYSSNNKNYLA (SEQ ID NO 98)	WASTRES (SEQ ID NO 78)	QYYSTPLT (SEQ ID NO 80)
GC5B379	NYWMS (SEQ ID NO 2)	GISYSGGSKYYADSVKG (SEQ ID NO 28)	AAWDFGRRVRLDY (SEQ ID NO 30)	RASQSISSYLN (SEQ ID NO 13)	AASSLQS (SEQ ID NO 16)	QSYSTPLT (SEQ ID NO 19)
GC5B373	SYWIG (SEQ ID NO 27)	IYPGDS TRYSPSFQG (SEQ ID NO 29)	IGFYGRSFRIFDY (SEQ ID NO 73)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B376	SYWIG (SEQ ID NO 27)	IYPGDS TRYSPSFQG (SEQ ID NO 29)	VYSFGGRHKALFDY (SEQ ID NO 11)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B385	GYAMS (SEQ ID NO 62)	AISGSGGSTYYADSVKG (SEQ ID NO 68)	VDRSFGRSRYTLDY (SEQ ID NO 74)	RASQSVSSYLA (SEQ ID NO 14)	DASNRAT (SEQ ID NO 17)	QQRSNWPLT (SEQ ID NO 20)
GC5B370 GC5B597	SYGIS (SEQ ID NO 63)	GIPIFGNINYAQKFQG (SEQ ID NO 69)	VSRFRKRFAYYFDY (SEQ ID NO 75)	KSSQSVLYSSNNKNYLA (SEQ ID NO 98)	WASTRES (SEQ ID NO 78)	QYYSTPLT (SEQ ID NO 80)
GC5B602	GYSFTGYTMN (SEQ ID NO 64)	LINPYNGDTN (SEQ ID NO 70)	VALRVALDY (SEQ ID NO 12)	KASQNVATHVG (SEQ ID NO 15)	SASYRYS (SEQ ID NO 18)	QYNYRYPYT (SEQ ID NO 21)
GC5B603	SYAMS (SEQ ID NO 65)	AISGSGGSTYYADSVKG (SEQ ID NO 68)	SNFLPVVFDY (SEQ ID NO 76)	RASQSVRKSLA (SEQ ID NO 95)	TASNRAT (SEQ ID NO 79)	QYFRAPIT (SEQ ID NO 81)
GC5B601	GFSLTSYNVH (SEQ ID NO 66)	VIWAGGSTNYNSALMS (SEQ ID NO 71)	DGIRLRFAY (SEQ ID NO 77)	KASQNVATHVG (SEQ ID NO 15)	SASYRYS (SEQ ID NO 18)	QYNYRYPYT (SEQ ID NO 21)

5

Table 29. Heavy and Light chain variable region sequences of new panel of GPRC5D antibodies

mAb AA ID	VH Amino Acid Sequence	SEQ ID NO	VL Amino Acid Sequence	SEQ ID NO
-----------	------------------------	-----------	------------------------	-----------

GC5B382	EVQLLES GGGLVQPGGSL RLSCAASGFTFSDYGMH WVRQAPGKGLEWVSAIK YSGGSTYYADSVKGRFTI SRDNSKNTLYLQMNSLRA EDTAVYYCAKRAESGPGL DYWGQGTLVTVSS	82	DIVMTQSPDSLAVSLGER ATINCKSSQSVLYSSNNK NYLAWYQQKPGQPPKLLI YWASTRESGVDPDRFSGSG SGTDFTLTISSLQAEDVAV YYCQQYYSTPLTFGQGTK VEIK	92
GC5B379	EVQLLES GGGLVQPGGSL RLSCAASGFTFSNYWMS WVRQAPGKGLEWVSGIS YSGGSKYYADSVKGRFTI SRDNSKNTLYLQMNSLRA EDTAVYYCAKAAWDFGR RAVRDYGWQGTLVTVSS	83	DIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQ QKPGKAPKLLIYAASSLQ SGVPSRFSGSGSGTDFTLT ISSLPEDFATYYCQQSYS TPLTFGQGTKVEIK	56
GC5B373	EVQLVQSGAEVKKPGESL KISCKGSGYSFTSYWIGW VRQMPGKGLEWMGIYP GDS DTRYSPSFQGQVTISA DKSISTAYLQWSSLKASD TAMYYCARIGFYGRSFRIF DYWGQGTLVTVSS	84	EIVLTQSPATLSLSPGERA TLSCRASQSVSSYLAWYQ QKPGQAPRLLIYDASNRA TGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYCQQRSN WPLTFGQGTKVEIK	57
GC5B376	EVQLVQSGAEVKKPGESL KISCKGSGYSFTSYWIGW VRQMPGKGLEWMGIYP GDS DTRYSPSFQGQVTISA DKSISTAYLQWSSLKASD TAMYYCARVYSFGGRHK ALFDYWGQGTLVTVSS	85	EIVLTQSPATLSLSPGERA TLSCRASQSVSSYLAWYQ QKPGQAPRLLIYDASNRA TGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYCQQRSN WPLTFGQGTKVEIK	57
GC5B385	EVQLLES GGGLVQPGGSL RLSCAASGFTFSGYAMSW VRQAPGKGLEWVSAISGS GGSTYYADSVKGRFTISR DNSKNTLYLQMNSLRAE DTAVYYCAKVDRSFRGRS RYTL DYGWQGTLVTVSS	86	EIVLTQSPATLSLSPGERA TLSCRASQSVSSYLAWYQ QKPGQAPRLLIYDASNRA TGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYCQQRSN WPLTFGQGTKVEIK	57

GC5B370 GC5B597	QVQLVQSGAEVKKPGSSV KVSCASGGTFSSYGISW VRQAPGQGLEWMGGIPI FGNINYAQKFQGRVTITA DESTSTAYMELSSLRSED AVYYCARVSRFRKFAY YFDYWGQGTLVTVSS	87	DIVMTQSPDSLAVSLGER ATINCKSSQSVLYSSNNK NYLAWYQQKPGQPPKLLI YWASTRESGVDPDRFSGSG SGTDFTLTISLQAEDVAV YYCQQYYSTPLTFGQGTK VEIK	92
GC5B602	QVQLVQSGAEVKKPGAS VKVSCASGYSFTGYTM NWVRQAPGQGLEWMGLI NPYNGDTNYAQKLQGRV TMTTDTSTSTAYMELRSL RSDDTAVYYCARVALRV ALDYWGQGTLVTVSS	88	DIQMTQSPSSLSASVGDR VTITCKASQNVATHVGW YQKPGKAPKRLIYSASY RYSQVPSRFSGSGSGTEFT LTISNLQPEDFATYYCQQ YNRYPTYFGQGTKLEIK	58
GC5B603	EVQLLES GGGLVQPGGSL RLSCAASGFTFSSYAMSW VRQAPGKGLEWVSAISGS GGSTYYADSVKGRFTISR DNSKNTLYLQMNSLRAE DTAVYYCAKSNFLPVVFD YWGQGTLVTVSS	89	EIVLTQSPATLSLSPGERA TLSCRASQSVRKSLAWYQ QKPGQAPRLLIYTASNRA TGIPARFSGSGSGTDFTLT ISLEPEDFAVYYCQQYFR APITFGQGTKVEIKK	93
GC5B601	QVTLKESGPVLVKPTETL TLCTVSGFSLTSYNVHW IRQPPGKALEWLAVIWAG GSTNYNSALMSRLTISKD TSKSQVVLMTNMRAED TATYYCARDGIRLRFAYW QGQTLVTVSS	91	EIVMTQSPATLSVSPGERA TLSCASQNVATHVGWY QKPGQAPRLLIYSASYR YSGIPARFSGSGSGTEFTL TISLQSEDFAVYYCQQY NRYPPTYFGQGTKLEIK	94

ADCC activity against H929 cells (Tables 30 and 31). Briefly, multiple myeloma cells were labeled with Calcein-AM for 30 minutes at room temperature and re-suspend at 0.2×10^6 /ml in RPMI + 10% HI FBS after two PBS washes. PBMC's were thawed and resuspended after PBS wash at 3×10^6 cells per ml in RPMI growth media. 10000 or 50000 target cells were mixed with 100000 or 2500000 PBMC'S in presence of the antibody and incubated for 3 hours in CO2 incubator at 37°C . After the incubation plate was centrifuges at 200 g for 4 minutes and 100 ul of supernatant was transferred into a new 96 well plate and the fluorescence intensity was measured at 485/535 nM. RFU values were plotted to calculate the percent lysis

Table 30: Antibody dependent cytotoxicity of H929 cells by anti-GPRC5D mAbs with IgG1 Fc.

GPRC5D IgG4 Protein ID	GPRC5D IgG1 Protein ID	H929 Cytotoxicity EC ₅₀ (pM)	% Lysis
GC5B243	GC5B382	1346.5	29.14
GC5B285	GC5B379	87.4	17.3
GC5B332	GC5B373	244.2	32.94
GC5B164	GC5B376	22.0	32.21
GC5B320	GC5B385	944.0	29.73
GC5B251	GC5B370	24.2	8
GC5B515	GC5B602	27721.7	20
GC5B420	GC5B603	3.4	16
GC5B483	GC5B599	7.4	14
GC5B540	GC5B601	169.6	14
GC5B465	GC5B598	2.0	12
GC5B251	GC5B597	705.1	10

5 **Table 31:** Comparison of Antibody dependent cytotoxicity and T-cell mediated cytotoxicity of H929 cells.

GPRC5D IgG1 Protein ID	GPRC5D x CD3 DuoBody ID	ADCC	CD3 Redirection
		H929 Cytotoxicity EC ₅₀ (pM)	T-cell Mediated H929 Cytotoxicity EC ₅₀ (nM)
GC5B382	GCDB40	1346.5	1.2 ± 0.83
GC5B379	GCDB43	87.4	0.39 ± 0.07
GC5B373	GCDB45	244.2	2.36 ± 0.84
GC5B376	GCDB35	22.0	1.39 ± 0.75
GC5B385	GCDB44	944.0	>10
GC5B370	GCDB41	24.2	>10
GC5B602	GCDB72	27721.7	0.15
GC5B603	GCDB48	3.4	0.17
GC5B599	GCDB61	7.4	1.51
GC5B601	GCDB69	169.6	0.58
GC5B598	GCDB53	2.0	0.24

10 A range of potency was observed ranging from 2 pM to 27.7 nM. Binding affinity was not necessarily predictive of efficacy in the ADCC assay. For example, GC5B382 and GC5B379 had similar binding affinity to human GPRC5D cells but a 15x difference in cytotoxicity against

H929 cells in the ADCC assay. Similarly, cytotoxic inducement as a GPRC5D x CD3 bispecific was not predictive of potency in the ADCC assay as exemplified by GC5B370 and GC5B602. When formatted as a CD3 bispecific GC5B602 (GCDB63) had sub-nanomolar potency against H929 cells while GC5B370 as a CD3 bispecific (GCDB41) was essentially inactive. The same v-

regions, when formatted as IgG1 mAbs resulted in the opposite observation in the ADCC assay, with GC5B370 observed as more potent (by ~1100x fold) over GC5B602.

Brief Description of the Sequence Listing

SEQ ID NO:	Type	Species	Description	Sequence
1	PRT	human	GC5B81, GC5B427, GC5B428, GC5B430, GC5B431, AND GC5B429- HCDR1	SYAIS
2	PRT	human	GC5B379, GC5B598,	NYWMS

			GC5B465, GC5B285, GC5B463, GC5B432, GC5B433, GC5B434, GC5B435, GC5B436, GC5B461, GC5B462, GC5B437, GC5B438, GC5B439, GC5B440, GC5B441, GC5B442, GC5B443, GC5B444, GC5B464, GC5B445, GC5B446, GC5B447, GC5B448, GC5B449, and GC5B450 - HCDR1	
3	PRT	human	GC5B483, GC5B473, GC5B478, GC5B488, GC5B493, and GC5B498 - HCDR1	SYFIG
4	PRT	human	GC5B596- HCDR1	GYTMN
5	PRT	human	GC5B81, GC5B427, GC5B428, GC5B430, GC5B431, and GC5B429- HCDR2	GIPIFGTANYAQKFQG
6	PRT	human	GC5B598,	GISYSGGSKYYASSVKG

			GC5B465, GC5B432, AND GC5B447- HCDR2	
7	PRT	human	GC5B599, GC5B483, GC5B481, GC5B482, GC5B484, GC5B485, AND GC5B503- HCDR2	IIYPGKSDTRYSPSFQG
8	PRT	human	GC5B596- HCDR2	LINPYNSDTNYAQKLQG
9	PRT	human	GCB581- HCDR3	ESRWRGYKLD
10	PRT	human	GC5B598, GCB5465- HCDR3	AAFDFGRRAVRLD
11	PRT	human	GC5B376, GC5B599, GC5B483, GC5B164, GC5B471, GC5B472, GC5B473, GC5B474, GC5B475, GC5B476, GC5B477, GC5B478, GC5B479, GC5B480, GC5B481, GC5B482, GC5B484, GC5B485, GC5B486, GC5B487, GC5B488, GC5B489, GC5B490, GC5B491, GC5B492,	VYSFGGRHKALFDY

			GC5B493, GC5B494, GC5B495, GC5B496, GC5B497, GC5B498, GC5B499, GC5B500, GC5B501, GC5B502, GC5B503, GC5B504, AND GC5B505 -HCDR3	
12	PRT	human	GC5B602, GCB596- HCDR3	VALRVALDY
13	PRT	human	GC5B382, GC5B379, GC5B370, GC5B598, GC5B597, GC5B81, GC5B465- LCDR1	RASQSISSYLN
14	PRT	human	GC5B373, GC5B376, GC5B385, GC5B599, GC5B483- LCDR1	RASQSVSSYLA
15	PRT	human	GC5B605, GC5B601, GC5B596- LCDR1	KASQNVATHVG
16	PRT	human	GC5B81, GC5B465, GC5B379, GC5B598- LCDR2	AASSLQS
17	PRT	human	GC5B373, GC5B376, GC5B385, GC5B599, GC5B483-	DASNRAT

			LCDR2	
18	PRT	human	GC5B602, GC5B601, GC5B596- LCDR2	SASYRYS
19	PRT	human	GC5B379, GC5B598, GC5B81, GC5B465- LCDR3	QQSYSTPLT
20	PRT	human	GC5B373, GC5B376, GC5B385, GC5B599, GC5B483- LCDR3	QQRSNWPLT
21	PRT	human	GC5B602, GC5B601, GC5B596- LCDR3	QQYNRYPYT
22	PRT	human	GPRC5D	MYKDCIESTGDYFLLCDAEGPWGIILES LAILGIVVTILLLLAFLFLMRKIQDCSQW NVLPTQLLFLLSV LGLFGLAFAFIIELNQQTAPVRYFLFGVL FALCFSCLLAHASNLVKLVRGCVSFSW TTILCIAIGCSLLQ IIATEYVTLIMTRGMMFVNMTPCQLNV DFVVLLVYVLFMLALTFFVSKATFCGP CENWKQHGRILFITV LFSIIIWVWISMILLRGNPQFQRQPQWD DPVVCIALVTNAWVFLLLYIVPELCILY RSCRQECPLQGNAC PVTAYQHSFQVENQELSRARDSDGAEE DVALTSYGTPIQPQTVDPTQECFIPQAK LSPQQDAGGV
23	PRT	cyno	GPRC5D	MYKDCIESTGDYFLPCDSEGPWGIVLES LAILGIVVTILLLLAFLFLMRKIQDCSQW NVLPTQLLFLLSV LGLFGLAFAFIIQLNQQTAPVRYFLFGV LFALCFSCLLAHASNLVKLVRGRVSFS WTTILCIAIGCSLLQVIIAIEYVTLIMTRG MMFVHMTPYQLNVDFVVLLVYVLFML ALTFFVSKATFCGPCENWKQHGRILFIT VLFSSIIIWVWISMILLRGNPQFQRQPQW DDPVVCIALVTNA WVFLLLYIVPELCILYRSCRQECPSQGH

				ACPVTA YQ R S F Q V E N Q E L S R A R D S D G A E E D V A L T S F G T P I Q P Q T V D P T Q E C F I P R A K L S P Q Q D A G V
24	PRT	mouse	GPRC5D	MYEDCVKSTEDYYLFC DNEG PWAIVLE SLAVIGIVVTILLLLAFLFLMRKVQDCS QWNVLPTQFLFLLAV LGLFGLTFAFIIQLNHQTAPVRYFLFGVL FAICFSCLLAHASNLVKLVRGRVSFCWT TILFIAIGVSLQ TIIAIEYVTLIMTRGLMFEHMTPTYQLNV DFVCLLIYVLFLMALTFFVSKATFCGPC ENWKQHGR LIFATV LVSIHVVVWISMLLRGNPQLQRQPHW DDAVICIGLV TNAWVFLLIYIPELSILYR SCRQECPTQGNVC QVPVYQRSFRMDTQEPTRARDSDGAQE DVALTAYGTPIQLQSADPSREYLIPSATL SPQQDAGL
25	PRT	human	CD3B219- HEAVY CHAIN	EVQLVESGGGLVQPGGSLRLSCAASGF TFNTYAMNWVRQAPGKGLEWVARIRS KYN NYATYYAASVKGRFTISRDDSKNS LYLQMNSLKTEDTAVYYCARHGNFGN SYVSWFAYWGQGT LVT VSS
26	PRT	human	CD3B219- LIGHT CHAIN	QTVVTQEPSLTVSPGGTVTLTCRSSTGA VTTSNYANWVQQKPGQAPRGLIGGTN KRAPGTPARFSGSLLGGKAALTL SGVQ PEDEAEYYCALWYSNLWVFGGGTKLT VL
27	PRT	human	GC5B373, GC5B376, GC5B164, GC5B501, GC5B502, GC5B503, GC5B504, AND GC5B505 - HCDR1	SYWIG
28	PRT	human	GC5B379, GC5B285, GC5B442, GC5B443, GC5B444, GC5B464,	GISYSGGSKYYADSVKG

			AND GC5B445- HCDR2	
29	PRT	human	GC5B373, GC5B376, GC5B164, GC5B496, GC5B497, GC5B498, GC5B499, AND GC5B500- HCDR2	IIYPGD <u>S</u> DTRYSPSFQG
30	PRT	human	GC5B379, GC5B285, GC5B446, GC5B447, GC5B448, GC5B449, AND GC5B450- HCDR3	AAWDFGRRAVRLDY
31	PRT	human	GC5B427- HCDR3	ESRYRGYKLDY
32	PRT	human	GC5B428- HCDR3	ESRVRGYKLDY
33	PRT	human	GC5B430- HCDR3	ESRGRGYKLDY
34	PRT	human	GC5B431- HCDR3	ESRARGYKLDY
35	PRT	human	GC5B429 HCDR3- VH	ESRFRGYKLDY
36	PRT	human	GC5B471, GC5B476, GC5B486, GC5B481, GC5B491, and GC5B496, -HCDR1	SY <u>Y</u> IG
37	PRT	human	GC5B497, GC5B472, GC5B477, GC5B482, GC5B487,	SY <u>V</u> IG

			AND GC5B492- HCDR1	
38	PRT	human	GC5B474, GC5B479, GC5B484, GC5B489, GC5B494, AND GC5B499- HCDR1	SY <u>G</u> IG
39	PRT	human	GC5B475, GC5B480, GC5B485, GC5B490, GC5B495, AND GC5B500- HCDR1	SY <u>A</u> IG
40	PRT	human	GC5B471, GC5B472, GC5B473, GC5B474, GC5B475, AND GC5B501- HCDR2	IIYPG <u>A</u> SDTRYSPSFQG
41	PRT	human	GC5B476, GC5B477, GC5B478, GC5B479, GC5B480, AND GC5B502 - HCDR2	IIYPG <u>S</u> SDTRYSPSFQG
42	PRT	human	GC5B486, GC5B487, GC5B488, GC5B489, GC5B490, AND GC5B504- HCDR2	IIYPG <u>E</u> SDTRYSPSFQG
43	PRT	human	GC5B491, GC5B492, GC5B493,	IIYPG <u>Y</u> SDTRYSPSFQG

			GC5B494, GC5B495, AND GC5B505- HCDR2	
44	PRT	human	GC5B463 AND GC5B446- HCDR2	GISYSGGSKYYA <u>A</u> SVKG
45	PRT	human	GC5B433, GC5B434, AND GC5B448- HCDR2	GISYSGGSKYYA <u>K</u> SVKG
46	PRT	human	GC5B435, GC5B436, GC5B461, GC5B462, AND GC5B449- HCDR2	GISYSGGSKYYA <u>E</u> SVKG
47	PRT	human	GC5B437, GC5B438, GC5B439, GC5B440, GC5B441, AND GC5B450- HCDR2	GISYSGGSKYYA <u>Y</u> SVKG
48	PRT	human	GC5B463, GC5B435, GC5B437, AND GC5B442- HCDR3	AA <u>Y</u> DFGRRVRLDY
49	PRT	human	GC5B432, GC5B433, GC5B438, AND GC5B443- HCDR3	AA <u>Y</u> DFGRRVRLDY
50	PRT	human	GC5B461, GC5B440, AND GC5B464- HCDR3	AA <u>G</u> DFGRRVRLDY

51	PRT	human	GC5B462, GC5B441, AND GC5B445- HCDR3	AAADFGRRAVRLDY
52	PRT	human	GC5B81- VH	QVQLVQSGAEVKKPGSSVKVSCKASGG TFSSYAIWVRQAPGQGLEWMGGIPIF GTANYAQKFQGRVTITADESTSTAYME LSSLRSED ^T AVYYCARESRWRGYKLDY WGQGTLVTVSS
53	PRT	human	GC5B598, GC5B465- VH	EVQLLES ^G GGGLVQPGGSLRLSCAASGFT FSNYWMSWVRQAPGKGLEWVSGISYS GGSKYYASSVKGRFTISRDN ^S KN ^T LYL QMNSLRAED ^T AVYYCAKAAFD ^G RRRA VRLDYWGQGTLVTVSS
54	PRT	human	GC5B599, GC5B483- VH	EVQLVQSGAEVKKPGESLKISCKGSGYS FTSYFIGWVRQMPGKGLEWMGIIYPGK SDTRYSPSFQGGVTISADKSISTAYLQW SSLKASDTAMY ^C CARVYSFGGRHKALF DYWGQGTLVTVSS
55	PRT	human	GC5B596- VH	QVQLVQSGAEVKKPGASVKVSCKASG YSFTGYTMNWVRQAPGQGLEWMGLIN PYN ^S DTNYAQKLQGRVTMTTDTSTSTA YME ^L RLSRSD ^T AVYYCARVALRVAL DYWGQGTLVTVSS
56	PRT	human	GC5B379, GC5B598, GC5B81 and GC5465- VL	DIQMTQSPSSLSASVGDRVTITCRASQSI SSYLNWYQQKPGKAPKLLIYAASSLQS GVPSRFSGSGSGTDFTLTIS ^S LQPEDFAT YYCQQSYSTPLTFGQGTKVEIK
57	PRT	human	GC5B373, GC5B376, GC5B385, GC5B599, GC5B483- VL	EIVLTQSPATLSLSPGERATLSCRASQSV SSYLAWYQQKPGQAPRLIYDASN ^R AT GIPARFSGSGSGTDFTLTIS ^S LEPEDFAV YYCQQRSNWPLTFGQGTKVEIK
58	PRT	human	GC5B596, GC5B602- VL	DIQMTQSPSSLSASVGDRVTITCKASQN VATHVGWYQQKPGKAPKRLIYSASYR YSGVPSRFSGSGSGTEFTLTIS ^N LQPEDF ATYYCQQYNRY ^P YTFGQGTKLEIK
59	PRT	human	IgG4PAA	ASTKGPSVFPLAPCSRSTSESTAALGCL VKDYFPEPVT ^S WNSGALTSGVHTFPA VLQSSGLYSLSSV ^T VPSSSLG ^T KTYTC NVDHKPSNTKVDKRVESKYGPPCPPCP APEAAGGPSVFLFPPKPKDTLMISRTPE

				VTCVVVDVSQEDPEVQFNWYVDGVEV HNAKTKPREEQFNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKGLPSSIEKTISKA KGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKT TPPVLDSDGSFFLYSRLTVDKSRWQEG NVFSCSVMHEALHNHYTQKSLSLGLK
60	PRT	human	IgG1	ASTKGPSVFPLAPSSKSTSGGTAALGCL VKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSSVVTVPSSSLGTQTYIC NVNHKPSNTKVDKKVEPKSCDKTHTCP PCPAPELLGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPREEQYNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENN YKTPPVLDSDGSFFLYSKLTVDKSRW QQGNVFSCSVMHEALHNHYTQKSLSL PGK
61	PRT	human	GC5B382- HCDR1	DYGMH
62	PRT	human	GC5B385- HCDR1	GYAMS
63	PRT	human	GC5B370, GC5B597- HCDR1	SYGIS
64	PRT	human	GC5B602- HCDR1	GYSFTGYTMN
65	PRT	human	GC5B603- HCDR1	SYAMS
66	PRT	human	GC5B601- HCDR1	GFSLTSYNVH
67	PRT	human	GC5B382- HCDR2	AIKYSGGSTYYADSVKG
68	PRT	human	GC5B603, GC5B385- HCDR2	AISGSGGSTYYADSVKG
69	PRT	human	GC5B370, GC5B597- HCDR2	GIPIFGNINYAQKFQG
70	PRT	human	GC5B602- HCDR2	LINPYNGDTN
71	PRT	human	GC5B601- HCDR2	VIWAGGSTNYNSALMS

72	PRT	human	GC5B382-HCDR3	RAESGPGLDY
73	PRT	human	GC5B373-HCDR3	IGFYGRSFRIFDY
74	PRT	human	GC5B385-HCDR3	VDRSFGRSRYTLDY
75	PRT	human	GC5B370, GC5B597-HCDR3	VSRRFKRFAYYFDY
76	PRT	human	GC5B603-HCDR3	SNFLPVVFDY
77	PRT	human	GC5B601-HCDR3	DGIRLRFAY
78	PRT	human	GC5B382, GC5B370, GC5B597-LCDR2	WASTRES
79	PRT	human	GC5B603-LCDR2	TASNRAT
80	PRT	human	GC5B382, GC5B370, GC5B597-LCDR3	QQYYSTPLT
81	PRT	human	GC5B603-LCDR3	QQYFRAPIT
82	PRT	human	GC5B382-VH	EVQLLES GGGLVQPGGSLRLSCAASGFT FSDYGMHWVRQAPGKGLEWVSAIKYS GGSTYYADSVKGRFTISRDN SKNTLYL QMNSLRAEDTAVYYCAKRAESGPGLD YWGQGT LTVSS
83	PRT	human	GC5B379-VH	EVQLLES GGGLVQPGGSLRLSCAASGFT FSNYWMSWVRQAPGKGLEWVSGISYS GGSKYYADSVKGRFTISRDN SKNTLYL QMNSLRAEDTAVYYCAKAAWDFGRR AVRLDYWGQGT LTVSS
84	PRT	human	GC5B376-VH	EVQLVQSGAEVKKPGESLKISCKGSGYS FTSYWIGWVRQMPGKGLEWMGIIYPG DSDTRYSPSFQGQVTISADKSISTAYLQ WSSLKASDTAMY YCARIGFYGRSFRIF DYWGQGT LTVSS
85	PRT	human	GC5B376-VH	EVQLVQSGAEVKKPGESLKISCKGSGYS FTSYWIGWVRQMPGKGLEWMGIIYPG DSDTRYSPSFQGQVTISADKSISTAYLQ WSSLKASDTAMY YCARVYSFGGRHKA

				LFDYWGGQGLVTVSS
86	PRT	human	GC5B385-VH	EVQLLES GGGLVQP GGSRLRLSCAASGFT FSGYAMSWVRQAPGKGLEWVSAISGS GGSTYYADSVKGRFTISRDN SKNTLYL QMNSLRAEDTAVYYCAKVDRSFGRSR YTLDYWGGQGLVTVSS
87	PRT	human	GC5B370, GC5B597-VH	QVQLVQSGAEVKKPGSSVKV SCKASGG TFSSYGISWVRQAPGQGLEWMGGIIPF GNINYAQKFQGRVTITADESTSTAYME LSSLRSED TAVYYCARVSRRFKRFAYY FDYWGGQGLVTVSS
88	PRT	human	GC5B602-VH	QVQLVQSGAEVKKPGASVKV SCKASG YSFTGYTMNWVRQAPGQGLEWMGLIN PYNGDTN YAQKLQGRVTMTTDTSTSTA YMELRSLRSDDTAVYYCARVALRVAL DYWGQGLVTVSS
89	PRT	human	GC5B603-VH	EVQLLES GGGLVQP GGSRLRLSCAASGFT FSSYAMSWVRQAPGKGLEWVSAISGSG GSTYYADSVKGRFTISRDN SKNTLYLQ MNSLRAEDTAVYYCAKSNFLPVVFDY WGQGLVTVSS
90	DNA	human	GC5B596-light chain	Atgcgggtgctggcccagctgctgggactgctgctgctgc ttccctggcgccagatgcgacatccagatgaccagagcccc agcagcctgagcgccagcgtggcgaccgggtgaccatca cctgcaaggccagccagaacgtggccaccacgtgggctg gtaccagcagaagcccggcaaggccccaagcggctgatct acagcgccagctaccggtacagcggcgtgccagccggttc agcggcagcggcagcggcaccgagttaccctgaccatca gcaacctgcagcccaggacttcgccacctactactgccagc agtacaaccggtaccctacaccttcggccagggcaccaag ctggagatcaagcgtacggtggctgcaccatctgtcttcatt cccgccatctgatgacagttgaaatctggaactgcctctgttg tgtgcctgctgaataacttctatcccagagaggccaaagtaca gtggaaggtggataacgccctccaatcgggtaactcccagga gagtgtcacagagcaggacagcaaggacagcacctacagc ctcagcagcaccctgacgctgagcaaagcagactacgagaa acacaaagtctacgcctgcgaagtcacccatcagggcctgag ctcgcccgtcacaagagcttcaacaggggagagtgttga
91	PRT	human	GC5B601-VH	QVTLKESGPVLVKPTETLTLTCTVSGFS LTSYNVHWIRQPPGKALEWLAVIWAG GSTNYNSALMSRLTISKDTSKSQVVL MTNMRAEDTATYYCARDGIRLRFAYW

				GQGTLVTVSS
92	PRT	human	GC5B382, GC5B597, GC5B370 - VL	DIVMTQSPDSLAVSLGERATINCKSSQS VLYSSNNKNYLAWYQQKPGQPPKLLIY WASTRESGVPDRFSGSGSGTDFTLT LQAEDVAVYYCQQYYSTPLTFGQGTK VEIK
93	PRT	human	GC5B603- VL	EIVLTQSPATLSLSPGERATLSCRASQSV RKSLAWYQQKPGQAPRLLIYTASNRA T GIPARFSGSGSGTDFTLT TISSLEPEDFAV YYCQQYFRAPITFGQGTKVEIKK
94	PRT	human	GC5B601- VL	EIVMTQSPATLSVSPGERATLSCKASQN VATHVGWYQQKPGQAPRLLIYSASYRY SGIPARFSGSGSGTEFTLT TISSLQSEDF AVYYCQQYNRYPYTFGQGTKLEIK
95	PRT	human	GC5B603- LCDRI	RASQSVRKSLA
96	DNA	human	GC5B596- heavy chain	atggcctgggtctggaccctgctgttctgatggccgtgccc agagcatccaggcccagggtgcagctggtgcagagcggcgc cgaggtgaagaagcccggcgccagcgtgaaggtgagctgc aaggccagcggctacagcttcaccggctacaccatgaactg ggtgcggcaggccccggccaggcctggagtggatggggc ctgatcaacccctacaacagcgcacaccaactacgcccagaa gctgcagggccgggtgacctgaccaccgacaccagcacc agcaccgcctacatggagctgcggagcctgcggagcgcacg acaccgccgtgtactactgcgccgggtggcctgcgggtg gccctggactactggggccagggcaccctggtgaccgtgag cagcgcctccaccaaggggcccatccgtcttccccctggcgcc ctgctccaggagcacctccgagagcacagccgcccctgggt gcctggtcaaggactacttccccgaaccggtgacggtgctgt ggaactcaggcgccctgaccagcggcgtgcacaccttcccg gctgtctacagtccctcaggactctactccctcagcagcgtgg tgaccgtgccctccagcagcttgggcacgaaaacctacacct gcaacgtagatcacaagcccagcaacaccaaggtggacaa gagagttgagtccaaatatggtcccccattgcccaccatgcc agcacctgaggccgcccgggggaccatcagtcttctgttccc ccaaaacccaaggacactctcatgatctcccgacccctga ggtcacgtgcgtggtggtggacgtgagccaggaagaccccg aggtccagttcaactggtacgtggatggcgtggaggtgcata atgccaagacaaaagccgcccaggagcagttcaacagcac gtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactg gctgaacggcaaggagtacaagtgaaggtctccaacaaag gcctcccgtcctccatcgagaaaaccatctccaaagccaaag ggcagccccgagagccacaggtgtacacctgcccccatcc caggaggagatgaccaagaaccagggtcagcctgacctgcct ggtcaaaggcttctaccccagcgacatcgccgtggagtggg

				agagcaatgggcagccggagaacaactacaagaccacgcc tccgtgctggactccgacggctcttcttctctacagcagg ctaaccgtggacaagagcaggtggcaggaggggaatgttt ctcatgtccgtgatgcatgaggctctgcacaaccactacaca cagaagagcctctccctgtctctgggtaaatga
--	--	--	--	--

We Claim:

1. An isolated antibody, or an antigen-binding fragment thereof, that binds specifically to GPRC5D comprising:

a heavy chain complementary determining region 1 (CDR1) comprising the amino acid sequence of SEQ ID NO: 4, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 8, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 12; and

a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 15, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 18, and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 21.

2. The isolated antibody, or antigen-binding fragment thereof, of claim 1, comprising a variable heavy chain (VH) region comprising the amino acid sequence of SEQ ID NO: 55, and a variable light chain (VL) region comprising the amino acid sequence of SEQ ID NO: 58.

3. The isolated antibody, or antigen-binding fragment thereof, of claim 2, further comprising the amino acid sequence of SEQ ID NO: 59.

4. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 3, wherein the isolated antibody, or antigen-binding fragment thereof, binds to a polypeptide comprising the amino acid sequence of SEQ ID NO: 22.

5. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 4, wherein the antibody or antigen-binding fragment thereof is a human antibody or antigen-binding fragment thereof.

6. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 5, wherein the antibody or antigen-binding fragment thereof is recombinant.

7. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 6, wherein the antigen-binding fragment is a Fab fragment, a Fab2 fragment, or a single chain antibody.

8. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 7, wherein the antibody, or antigen-binding fragment thereof, are of IgG1, IgG2, IgG3, or IgG4 isotype.
9. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 8, wherein the antibody, or antigen binding fragment thereof, is of IgG1 or an IgG4 isotype.
10. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 9, wherein the antibody, or antigen-binding fragment thereof, specifically binds human GPRC5D and cross reacts to cynomolgus monkey GPRC5D.
11. The isolated antibody, or antigen-binding fragment thereof, of any one of claims 1 to 9 wherein the antibody or antigen-binding fragment thereof induces ADCC *in vitro* with an EC₅₀ of less than about 28 nM.
12. An isolated cell expressing the antibody, or antigen-binding fragment thereof, of any one of claims 1 to 11.
13. The isolated cell of claim 12, wherein the cell is a hybridoma.
14. The isolated cell of claim 12 or 13, wherein the antibody, or antigen-binding fragment thereof, is recombinantly produced.
15. An isolated GPRC5D x CD3 bispecific antibody comprising:
 - a) a first heavy chain (HC1);
 - b) a second heavy chain (HC2);
 - c) a first light chain (LC1); and
 - d) a second light chain (LC2),

wherein the HC1 and the LC1 pair to form a first antigen-binding site that specifically binds CD3, and the HC2 and the LC2 pair to form a second antigen-binding site that specifically binds GPRC5D,

wherein the HC1 comprises a VH region comprising the amino acid sequence of SEQ ID NO: 25 and the LC1 comprises a VL region comprising the amino acid sequence of SEQ ID NO: 26, and

wherein the HC2 comprises a VH region comprising the amino acid sequence of SEQ ID NO: 55 and the LC2 comprises a VL region comprising the amino acid sequence of SEQ ID NO: 58.

16. The GPRC5D x CD3 bispecific antibody, of claim 15, wherein the GPRC5D binding portion of the bispecific antibody further comprises the amino acid sequence of SEQ ID NO: 59.

17. The GPRC5D x CD3 bispecific antibody, of claim 15 or claim 16, wherein the CD3 binding portion of the bispecific antibody further comprises the amino acid sequence of SEQ ID NO: 59 comprising a F405L and a R409K substitution.

18. The GPRC5D x CD3 bispecific antibody, of any one of claims 15 to 17 wherein the antibody is of IgG1, IgG2, IgG3, or IgG4 isotype.

19. The GPRC5D x CD3 bispecific antibody, of claim 18 wherein the antibody or bispecific binding fragment thereof is of IgG4 isotype.

20. The GPRC5D x CD3 bispecific antibody, of any one of claims 15 to 19, wherein the antibody binds GPRC5D on the surface of human myeloma cells.

21. The GPRC5D x CD3 bispecific antibody, of any one of claims 15 to 20, wherein the antibody binds GPRC5D on the surface of human multiple myeloma cells.

22. The GPRC5D x CD3 bispecific antibody, of any one of claims 15 to 21 wherein the antibody induces human T cell activation *in vitro* with an EC₅₀ of less than about 0.22 nM.

23. The GPRC5D x CD3 bispecific antibody, of any one of claims 15 to 22, wherein the antibody induces T-cell dependent cytotoxicity of GPRC5D-expressing cells *in vitro* with an EC₅₀ of less than about 0.89 nM.

24. An isolated cell expressing the GPRC5D x CD3 bispecific antibody, of any one of claims 15 to 23.

25. The isolated cell of claim 24 wherein the cell is a hybridoma.

26. The isolated cell of claim 24 or claim 25, wherein the antibody is recombinantly produced.
27. A method for treating a subject having cancer, said method comprising:
administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23 to a patient in need thereof for a time sufficient to treat the cancer.
28. A method for inhibiting growth or proliferation of cancer cells, said method comprising:
administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23 to inhibit the growth or proliferation of cancer cells.
29. A method of redirecting a T cell to a GPRC5D-expressing cancer cell, said method comprising:
administering a therapeutically effective amount of the GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23 to redirect a T cell to a cancer.
30. The method of claim 27, 28, or 29 wherein the cancer is a hematological cancer.
31. The method of claim 30 wherein the hematological cancer is a GPRC5D-expressing B cell cancer.
32. The method of claim 31 wherein the GPRC5D-expressing B cell cancer is multiple myeloma.
33. The method of claim 27 comprising administering a second therapeutic agent.
34. The method of claim 33 wherein the second therapeutic agent is a chemotherapeutic agent or a targeted anti-cancer therapy.
35. The method of claim 34 wherein the chemotherapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2.
36. The method of claim 34 wherein the second therapeutic agent is administered to said subject simultaneously with, sequentially, or separately from the bispecific antibody.

37. A pharmaceutical composition comprising the GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23 and a pharmaceutically acceptable carrier.
38. A method for generating the GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23 by culturing the cell of any one of claims 24 to 26.
39. One or more isolated synthetic polynucleotides encoding the HC1, the HC2, the LC1, and the LC2 of the isolated GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23.
40. A kit comprising the GPRC5D x CD3 bispecific antibody as defined in any one of claims 15 to 23 and/or one or more polynucleotide(s) as defined in claim 39 and packaging for the same.
41. An isolated antibody that binds specifically to GPRC5D comprising:
 - a heavy chain comprising a variable heavy (VH) chain region comprising the amino acid sequence of SEQ ID NO: 55, and
 - a light chain comprising a variable light (VL) chain region comprising the amino acid sequence of SEQ ID NO: 58,wherein the antibody is an IgG4 isotype.
42. The isolated antibody of claim 41, wherein the heavy chain further comprises the amino acid sequence of SEQ ID NO: 59.
43. An isolated GPRC5D x CD3 bispecific antibody comprising:
 - a GPRC5D binding portion comprising:
 - a heavy chain comprising a variable heavy (VH) chain region comprising the amino acid sequence of SEQ ID NO: 55, and
 - a light chain comprising a variable light (VL) chain region comprising the amino acid sequence of SEQ ID NO: 58; and
 - a CD3 binding portion comprising a heavy chain comprising SEQ ID NO: 99 and a light chain comprising SEQ ID NO: 100,wherein the antibody is an IgG4 isotype.
44. The isolated GPRC5D x CD3 bispecific antibody of claim 43, wherein the GPRC5D heavy chain further comprises the amino acid sequence of SEQ ID NO: 59.

45. Use of the GPRC5D x CD3 bispecific antibody of any one of claims 15 to 23 in the manufacture of a medicament for the treatment of a subject having cancer.
46. A GPRC5D x CD3 bispecific antibody generated by the method of claim 38.

Figure 1




Figure 2




Figure 3




Figure 4A




Figure 4B




Figure 5A




Figure 5B

293F-GPRC5D(Cyno) cells, no Fc blocker (48hrs)

T cell donor: M6521(n=1)




Figure 6A




Figure 6B




Figure 7A




Figure 7B





Figure 8A
Human FcγRI






Figure 8C
Human FcγRIIb




Figure 8D
Human FcγRIIIa




Figure 9A




Figure 9B




Figure 10A




Figure 10B




Figure 11A




Figure 11B




Figure 11C




Figure 11D




Figure 11E




Figure 12A




Figure 12B




Figure 12C




Figure 12D





Figure 13



470 GC5M470.GC5B470.002

610 GC5B598 (GC5M610)

611 GC5B599 (GC5M611)

614 GC5B602 (GC5M614)

615 GC5B603 (GC5M615)

Antibody concentrations:

μg/mL	0	0.0003	0.001	0.005	0.020	0.078	0.313	1.250	5.000	20.000	80.000
nM	0	0.0020	0.008	0.032	0.130	0.520	2.078	8.313	33.250	133.000	532.000

Figure 14A

FAB6300




Figure 14A (Contd.)

GC5M481




Figure 14B




Figure 14B (Contd.)




Figure 14B (Contd.)




Figure 15A




Figure 15B




Figure 16





Figure 17



PRD3422WOPCT_SL.TXT
SEQUENCE LISTING

<110> JANSSEN PHARMACEUTICA NV

<120> ANTI- GPRC5D ANTIBODIES, BISPECIFIC ANTIGEN BINDING MOLECULES
THAT BIND GPRC5D AND CD3, AND USES THEREOF

<130> PRD3422WOPCT

<140>

<141>

<150> 62/364,811

<151> 2016-07-20

<160> 100

<170> PatentIn version 3.5

<210> 1

<211> 5

<212> PRT

<213> Homo sapiens

<400> 1

Ser Tyr Ala Ile Ser
1 5

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<400> 2

Asn Tyr Trp Met Ser
1 5

<210> 3

<211> 5

<212> PRT

<213> Homo sapiens

<400> 3

Ser Tyr Phe Ile Gly
1 5

<210> 4

<211> 5

<212> PRT

<213> Homo sapiens

<400> 4

Gly Tyr Thr Met Asn
1 5

<210> 5

<211> 17

<212> PRT

<213> Homo sapiens

PRD3422WOPCT_SL.TXT

<400> 5
 Gly Ile Ile Pro Ile Phe Gly Thr Ala Asn Tyr Ala Gln Lys Phe Gln
 1 5 10 15

Gly

<210> 6
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 6
 Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Ser Ser Val Lys
 1 5 10 15

Gly

<210> 7
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 7
 Ile Ile Tyr Pro Gly Lys Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln
 1 5 10 15

Gly

<210> 8
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 8
 Leu Ile Asn Pro Tyr Asn Ser Asp Thr Asn Tyr Ala Gln Lys Leu Gln
 1 5 10 15

Gly

<210> 9
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 9
 Glu Ser Arg Trp Arg Gly Tyr Lys Leu Asp
 1 5 10

<210> 10

<211> 13
<212> PRT
<213> Homo sapiens

<400> 10
Ala Ala Phe Asp Phe Gly Arg Arg Ala Val Arg Leu Asp
1 5 10

<210> 11
<211> 14
<212> PRT
<213> Homo sapiens

<400> 11
Val Tyr Ser Phe Gly Gly Arg His Lys Ala Leu Phe Asp Tyr
1 5 10

<210> 12
<211> 9
<212> PRT
<213> Homo sapiens

<400> 12
Val Ala Leu Arg Val Ala Leu Asp Tyr
1 5

<210> 13
<211> 11
<212> PRT
<213> Homo sapiens

<400> 13
Arg Ala Ser Gln Ser Ile Ser Ser Tyr Leu Asn
1 5 10

<210> 14
<211> 11
<212> PRT
<213> Homo sapiens

<400> 14
Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
1 5 10

<210> 15
<211> 11
<212> PRT
<213> Homo sapiens

<400> 15
Lys Ala Ser Gln Asn Val Ala Thr His Val Gly
1 5 10

<210> 16
<211> 7
<212> PRT
<213> Homo sapiens

<400> 16
Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 17
<211> 7
<212> PRT
<213> Homo sapiens

<400> 17
Asp Ala Ser Asn Arg Ala Thr
1 5

<210> 18
<211> 7
<212> PRT
<213> Homo sapiens

<400> 18
Ser Ala Ser Tyr Arg Tyr Ser
1 5

<210> 19
<211> 9
<212> PRT
<213> Homo sapiens

<400> 19
Gln Gln Ser Tyr Ser Thr Pro Leu Thr
1 5

<210> 20
<211> 9
<212> PRT
<213> Homo sapiens

<400> 20
Gln Gln Arg Ser Asn Trp Pro Leu Thr
1 5

<210> 21
<211> 9
<212> PRT
<213> Homo sapiens

<400> 21
Gln Gln Tyr Asn Arg Tyr Pro Tyr Thr
1 5

<210> 22
<211> 345
<212> PRT
<213> Homo sapiens

<400> 22
Met Tyr Lys Asp Cys Ile Glu Ser Thr Gly Asp Tyr Phe Leu Leu Cys

```

1              5              10              15
Asp Ala Glu Gly Pro Trp Gly Ile Ile Leu Glu Ser Leu Ala Ile Leu
      20      25      30
Gly Ile Val Val Thr Ile Leu Leu Leu Ala Phe Leu Phe Leu Met
      35      40      45
Arg Lys Ile Gln Asp Cys Ser Gln Trp Asn Val Leu Pro Thr Gln Leu
      50      55      60
Leu Phe Leu Leu Ser Val Leu Gly Leu Phe Gly Leu Ala Phe Ala Phe
      65      70      75      80
Ile Ile Glu Leu Asn Gln Gln Thr Ala Pro Val Arg Tyr Phe Leu Phe
      85      90      95
Gly Val Leu Phe Ala Leu Cys Phe Ser Cys Leu Leu Ala His Ala Ser
      100      105      110
Asn Leu Val Lys Leu Val Arg Gly Cys Val Ser Phe Ser Trp Thr Thr
      115      120      125
Ile Leu Cys Ile Ala Ile Gly Cys Ser Leu Leu Gln Ile Ile Ile Ala
      130      135      140
Thr Glu Tyr Val Thr Leu Ile Met Thr Arg Gly Met Met Phe Val Asn
      145      150      155      160
Met Thr Pro Cys Gln Leu Asn Val Asp Phe Val Val Leu Leu Val Tyr
      165      170      175
Val Leu Phe Leu Met Ala Leu Thr Phe Phe Val Ser Lys Ala Thr Phe
      180      185      190
Cys Gly Pro Cys Glu Asn Trp Lys Gln His Gly Arg Leu Ile Phe Ile
      195      200      205
Thr Val Leu Phe Ser Ile Ile Ile Trp Val Val Trp Ile Ser Met Leu
      210      215      220
Leu Arg Gly Asn Pro Gln Phe Gln Arg Gln Pro Gln Trp Asp Asp Pro
      225      230      235      240
Val Val Cys Ile Ala Leu Val Thr Asn Ala Trp Val Phe Leu Leu Leu
      245      250      255

```

Tyr Ile Val Pro Glu Leu Cys Ile Leu Tyr Arg Ser Cys Arg Gln Glu
260 265 270

Cys Pro Leu Gln Gly Asn Ala Cys Pro Val Thr Ala Tyr Gln His Ser
275 280 285

Phe Gln Val Glu Asn Gln Glu Leu Ser Arg Ala Arg Asp Ser Asp Gly
290 295 300

Ala Glu Glu Asp Val Ala Leu Thr Ser Tyr Gly Thr Pro Ile Gln Pro
305 310 315 320

Gln Thr Val Asp Pro Thr Gln Glu Cys Phe Ile Pro Gln Ala Lys Leu
325 330 335

Ser Pro Gln Gln Asp Ala Gly Gly Val
340 345

<210> 23

<211> 344

<212> PRT

<213> Macaca fascicularis

<400> 23

Met Tyr Lys Asp Cys Ile Glu Ser Thr Gly Asp Tyr Phe Leu Pro Cys
1 5 10 15

Asp Ser Glu Gly Pro Trp Gly Ile Val Leu Glu Ser Leu Ala Ile Leu
20 25 30

Gly Ile Val Val Thr Ile Leu Leu Leu Leu Ala Phe Leu Phe Leu Met
35 40 45

Arg Lys Ile Gln Asp Cys Ser Gln Trp Asn Val Leu Pro Thr Gln Leu
50 55 60

Leu Phe Leu Leu Ser Val Leu Gly Leu Phe Gly Leu Ala Phe Ala Phe
65 70 75 80

Ile Ile Gln Leu Asn Gln Gln Thr Ala Pro Val Arg Tyr Phe Leu Phe
85 90 95

Gly Val Leu Phe Ala Leu Cys Phe Ser Cys Leu Leu Ala His Ala Ser
100 105 110

Asn Leu Val Lys Leu Val Arg Gly Arg Val Ser Phe Ser Trp Thr Thr
115 120 125

Ile Leu Cys Ile Ala Ile Gly Cys Ser Leu Leu Gln Val Ile Ile Ala

130

135

Ile Glu Tyr Val Thr Leu Ile Met Thr Arg Gly Met Met Phe Val His
145 150 155 160

Met Thr Pro Tyr Gln Leu Asn Val Asp Phe Val Val Leu Leu Val Tyr
165 170 175

Val Leu Phe Leu Met Ala Leu Thr Phe Phe Val Ser Lys Ala Thr Phe
180 185 190

Cys Gly Pro Cys Glu Asn Trp Lys Gln His Gly Arg Leu Ile Phe Ile
195 200 205

Thr Val Leu Phe Ser Ile Ile Ile Trp Val Val Trp Ile Ser Met Leu
210 215 220

Leu Arg Gly Asn Pro Gln Phe Gln Arg Gln Pro Gln Trp Asp Asp Pro
225 230 235 240

Val Val Cys Ile Ala Leu Val Thr Asn Ala Trp Val Phe Leu Leu Leu
245 250 255

Tyr Ile Val Pro Glu Leu Cys Ile Leu Tyr Arg Ser Cys Arg Gln Glu
260 265 270

Cys Pro Ser Gln Gly His Ala Cys Pro Val Thr Ala Tyr Gln Arg Ser
275 280 285

Phe Gln Val Glu Asn Gln Glu Leu Ser Arg Ala Arg Asp Ser Asp Gly
290 295 300

Ala Glu Glu Asp Val Ala Leu Thr Ser Phe Gly Thr Pro Ile Gln Pro
305 310 315 320

Gln Thr Val Asp Pro Thr Gln Glu Cys Phe Ile Pro Arg Ala Lys Leu
325 330 335

Ser Pro Gln Gln Asp Ala Gly Val
340

<210> 24
<211> 344
<212> PRT
<213> Mus sp.

<400> 24
Met Tyr Glu Asp Cys Val Lys Ser Thr Glu Asp Tyr Tyr Leu Phe Cys
1 5 10 15

PRD3422WOPCT_SL.TXT

Asp Asn Glu Gly Pro Trp Ala Ile Val Leu Glu Ser Leu Ala Val Ile
20 25 30

Gly Ile Val Val Thr Ile Leu Leu Leu Ala Phe Leu Phe Leu Met
35 40 45

Arg Lys Val Gln Asp Cys Ser Gln Trp Asn Val Leu Pro Thr Gln Phe
50 55 60

Leu Phe Leu Leu Ala Val Leu Gly Leu Phe Gly Leu Thr Phe Ala Phe
65 70 75 80

Ile Ile Gln Leu Asn His Gln Thr Ala Pro Val Arg Tyr Phe Leu Phe
85 90 95

Gly Val Leu Phe Ala Ile Cys Phe Ser Cys Leu Leu Ala His Ala Ser
100 105 110

Asn Leu Val Lys Leu Val Arg Gly Arg Val Ser Phe Cys Trp Thr Thr
115 120 125

Ile Leu Phe Ile Ala Ile Gly Val Ser Leu Leu Gln Thr Ile Ile Ala
130 135 140

Ile Glu Tyr Val Thr Leu Ile Met Thr Arg Gly Leu Met Phe Glu His
145 150 155 160

Met Thr Pro Tyr Gln Leu Asn Val Asp Phe Val Cys Leu Leu Ile Tyr
165 170 175

Val Leu Phe Leu Met Ala Leu Thr Phe Phe Val Ser Lys Ala Thr Phe
180 185 190

Cys Gly Pro Cys Glu Asn Trp Lys Gln His Gly Arg Leu Ile Phe Ala
195 200 205

Thr Val Leu Val Ser Ile Ile Ile Trp Val Val Trp Ile Ser Met Leu
210 215 220

Leu Arg Gly Asn Pro Gln Leu Gln Arg Gln Pro His Trp Asp Asp Ala
225 230 235 240

Val Ile Cys Ile Gly Leu Val Thr Asn Ala Trp Val Phe Leu Leu Ile
245 250 255

Tyr Ile Ile Pro Glu Leu Ser Ile Leu Tyr Arg Ser Cys Arg Gln Glu

260

265

270

Cys Pro Thr Gln Gly Asn Val Cys Gln Val Pro Val Tyr Gln Arg Ser
 275 280 285

Phe Arg Met Asp Thr Gln Glu Pro Thr Arg Ala Arg Asp Ser Asp Gly
 290 295 300

Ala Gln Glu Asp Val Ala Leu Thr Ala Tyr Gly Thr Pro Ile Gln Leu
 305 310 315 320

Gln Ser Ala Asp Pro Ser Arg Glu Tyr Leu Ile Pro Ser Ala Thr Leu
 325 330 335

Ser Pro Gln Gln Asp Ala Gly Leu
 340

<210> 25
 <211> 125
 <212> PRT
 <213> Homo sapiens

<400> 25
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Thr Tyr
 20 25 30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Arg Ile Arg Ser Lys Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Ala
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Ser
 65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95

Tyr Cys Ala Arg His Gly Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe
 100 105 110

Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120 125

<210> 26
 <211> 109

PRD3422WOPCT_SL.TXT

<212> PRT

<213> Homo sapiens

<400> 26

Gln Thr Val Val Thr Gln Glu Pro Ser Leu Thr Val Ser Pro Gly Gly
1 5 10 15

Thr Val Thr Leu Thr Cys Arg Ser Ser Thr Gly Ala Val Thr Thr Ser
20 25 30

Asn Tyr Ala Asn Trp Val Gln Gln Lys Pro Gly Gln Ala Pro Arg Gly
35 40 45

Leu Ile Gly Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe
50 55 60

Ser Gly Ser Leu Leu Gly Gly Lys Ala Ala Leu Thr Leu Ser Gly Val
65 70 75 80

Gln Pro Glu Asp Glu Ala Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn
85 90 95

Leu Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu
100 105

<210> 27

<211> 5

<212> PRT

<213> Homo sapiens

<400> 27

Ser Tyr Trp Ile Gly
1 5

<210> 28

<211> 17

<212> PRT

<213> Homo sapiens

<400> 28

Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 29

<211> 17

<212> PRT

<213> Homo sapiens

<400> 29

Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln

1 5 10 15

Gly

<210> 30
<211> 14
<212> PRT
<213> Homo sapiens

<400> 30
Ala Ala Trp Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
1 5 10

<210> 31
<211> 11
<212> PRT
<213> Homo sapiens

<400> 31
Glu Ser Arg Tyr Arg Gly Tyr Lys Leu Asp Tyr
1 5 10

<210> 32
<211> 11
<212> PRT
<213> Homo sapiens

<400> 32
Glu Ser Arg Val Arg Gly Tyr Lys Leu Asp Tyr
1 5 10

<210> 33
<211> 11
<212> PRT
<213> Homo sapiens

<400> 33
Glu Ser Arg Gly Arg Gly Tyr Lys Leu Asp Tyr
1 5 10

<210> 34
<211> 11
<212> PRT
<213> Homo sapiens

<400> 34
Glu Ser Arg Ala Arg Gly Tyr Lys Leu Asp Tyr
1 5 10

<210> 35
<211> 11
<212> PRT
<213> Homo sapiens

<400> 35

Glu Ser Arg Phe Arg Gly Tyr Lys Leu Asp Tyr
 1 5 10

<210> 36
 <211> 5
 <212> PRT
 <213> Homo sapiens

<400> 36
 Ser Tyr Tyr Ile Gly
 1 5

<210> 37
 <211> 5
 <212> PRT
 <213> Homo sapiens

<400> 37
 Ser Tyr Val Ile Gly
 1 5

<210> 38
 <211> 5
 <212> PRT
 <213> Homo sapiens

<400> 38
 Ser Tyr Gly Ile Gly
 1 5

<210> 39
 <211> 5
 <212> PRT
 <213> Homo sapiens

<400> 39
 Ser Tyr Ala Ile Gly
 1 5

<210> 40
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 40
 Ile Ile Tyr Pro Gly Ala Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln
 1 5 10 15

Gly

<210> 41
 <211> 17
 <212> PRT
 <213> Homo sapiens

PRD3422WOPCT_SL.TXT

<400> 41

Ile Ile Tyr Pro Gly Ser Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln
1 5 10 15

Gly

<210> 42

<211> 17

<212> PRT

<213> Homo sapiens

<400> 42

Ile Ile Tyr Pro Gly Glu Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln
1 5 10 15

Gly

<210> 43

<211> 17

<212> PRT

<213> Homo sapiens

<400> 43

Ile Ile Tyr Pro Gly Tyr Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln
1 5 10 15

Gly

<210> 44

<211> 17

<212> PRT

<213> Homo sapiens

<400> 44

Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Ala Ser Val Lys
1 5 10 15

Gly

<210> 45

<211> 17

<212> PRT

<213> Homo sapiens

<400> 45

Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Lys Ser Val Lys
1 5 10 15

Gly

<210> 46
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 46
 Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Glu Ser Val Lys
 1 5 10 15

Gly

<210> 47
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 47
 Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Tyr Ser Val Lys
 1 5 10 15

Gly

<210> 48
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 48
 Ala Ala Tyr Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
 1 5 10

<210> 49
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 49
 Ala Ala Val Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
 1 5 10

<210> 50
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 50
 Ala Ala Gly Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
 1 5 10

<210> 51
 <211> 14
 <212> PRT

<213> Homo sapiens

<400> 51

Ala Ala Ala Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
 1 5 10

<210> 52

<211> 120

<212> PRT

<213> Homo sapiens

<400> 52

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Ser Ser Tyr
 20 25 30

Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Gly Ile Ile Pro Ile Phe Gly Thr Ala Asn Tyr Ala Gln Lys Phe
 50 55 60

Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Glu Ser Arg Trp Arg Gly Tyr Lys Leu Asp Tyr Trp Gly Gln
 100 105 110

Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 53

<211> 123

<212> PRT

<213> Homo sapiens

<400> 53

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

PRD3422WOPCT_SL.TXT

Ser Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Ser Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Ala Ala Phe Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 54
<211> 123
<212> PRT
<213> Homo sapiens

<400> 54
Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr
20 25 30

Phe Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
35 40 45

Gly Ile Ile Tyr Pro Gly Lys Ser Asp Thr Arg Tyr Ser Pro Ser Phe
50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ala Arg Val Tyr Ser Phe Gly Gly Arg His Lys Ala Leu Phe Asp Tyr
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 55
<211> 118
<212> PRT
<213> Homo sapiens

PRD3422WOPCT_SL.TXT

<400> 55

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Gly Tyr
20 25 30

Thr Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Ser Asp Thr Asn Tyr Ala Gln Lys Leu
50 55 60

Gln Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Val Ala Leu Arg Val Ala Leu Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser
115

<210> 56

<211> 107

<212> PRT

<213> Homo sapiens

<400> 56

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr Pro Leu
85 90 95

PRD3422WOPCT_SL.TXT
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 57
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 57
 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Leu
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 58
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 58
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asn Val Ala Thr His
 20 25 30

Val Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
 35 40 45

Tyr Ser Ala Ser Tyr Arg Tyr Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Asn Leu Gln Pro
 65 70 75 80

PRD3422WOPCT_SL.TXT

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Arg Tyr Pro Tyr
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 59
<211> 327
<212> PRT
<213> Homo sapiens

<400> 59
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
100 105 110

Glu Ala Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
115 120 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
130 135 140

Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
145 150 155 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
165 170 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
180 185 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
 195 200 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 210 215 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
 225 230 235 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 245 250 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285

Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320

Leu Ser Leu Ser Leu Gly Lys
 325

<210> 60
 <211> 330
 <212> PRT
 <213> Homo sapiens

<400> 60
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85

90

95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
 225 230 235 240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325 330

<210> 61
<211> 5
<212> PRT
<213> Homo sapiens

<400> 61
Asp Tyr Gly Met His
1 5

<210> 62
<211> 5
<212> PRT
<213> Homo sapiens

<400> 62
Gly Tyr Ala Met Ser
1 5

<210> 63
<211> 5
<212> PRT
<213> Homo sapiens

<400> 63
Ser Tyr Gly Ile Ser
1 5

<210> 64
<211> 10
<212> PRT
<213> Homo sapiens

<400> 64
Gly Tyr Ser Phe Thr Gly Tyr Thr Met Asn
1 5 10

<210> 65
<211> 5
<212> PRT
<213> Homo sapiens

<400> 65
Ser Tyr Ala Met Ser
1 5

<210> 66
<211> 10
<212> PRT
<213> Homo sapiens

<400> 66
Gly Phe Ser Leu Thr Ser Tyr Asn Val His
1 5 10

<210> 67
<211> 17
<212> PRT

<213> Homo sapiens

<400> 67

Ala Ile Lys Tyr Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 68

<211> 17

<212> PRT

<213> Homo sapiens

<400> 68

Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 69

<211> 17

<212> PRT

<213> Homo sapiens

<400> 69

Gly Ile Ile Pro Ile Phe Gly Asn Ile Asn Tyr Ala Gln Lys Phe Gln
1 5 10 15

Gly

<210> 70

<211> 10

<212> PRT

<213> Homo sapiens

<400> 70

Leu Ile Asn Pro Tyr Asn Gly Asp Thr Asn
1 5 10

<210> 71

<211> 16

<212> PRT

<213> Homo sapiens

<400> 71

Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Ser Ala Leu Met Ser
1 5 10 15

<210> 72

<211> 10

<212> PRT

<213> Homo sapiens

<400> 72

Arg Ala Glu Ser Gly Pro Gly Leu Asp Tyr
1 5 10

<210> 73

<211> 13

<212> PRT

<213> Homo sapiens

<400> 73

Ile Gly Phe Tyr Gly Arg Ser Phe Arg Ile Phe Asp Tyr
1 5 10

<210> 74

<211> 14

<212> PRT

<213> Homo sapiens

<400> 74

Val Asp Arg Ser Phe Gly Arg Ser Arg Tyr Thr Leu Asp Tyr
1 5 10

<210> 75

<211> 14

<212> PRT

<213> Homo sapiens

<400> 75

Val Ser Arg Arg Phe Lys Arg Phe Ala Tyr Tyr Phe Asp Tyr
1 5 10

<210> 76

<211> 10

<212> PRT

<213> Homo sapiens

<400> 76

Ser Asn Phe Leu Pro Val Val Phe Asp Tyr
1 5 10

<210> 77

<211> 9

<212> PRT

<213> Homo sapiens

<400> 77

Asp Gly Ile Arg Leu Arg Phe Ala Tyr
1 5

<210> 78

<211> 7

<212> PRT

<213> Homo sapiens

<400> 78

Trp Ala Ser Thr Arg Glu Ser

1

5

<210> 79

<211> 7

<212> PRT

<213> Homo sapiens

<400> 79

Thr Ala Ser Asn Arg Ala Thr

1

5

<210> 80

<211> 9

<212> PRT

<213> Homo sapiens

<400> 80

Gln Gln Tyr Tyr Ser Thr Pro Leu Thr

1

5

<210> 81

<211> 9

<212> PRT

<213> Homo sapiens

<400> 81

Gln Gln Tyr Phe Arg Ala Pro Ile Thr

1

5

<210> 82

<211> 119

<212> PRT

<213> Homo sapiens

<400> 82

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1

5

10

15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr

20

25

30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35

40

45

Ser Ala Ile Lys Tyr Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val

50

55

60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr

65

70

75

80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

PRD3422WOPCT_SL.TXT
 Ala Lys Arg Ala Glu Ser Gly Pro Gly Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Leu Val Thr Val Ser Ser
 115

<210> 83
 <211> 123
 <212> PRT
 <213> Homo sapiens

<400> 83
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Gly Ile Ser Tyr Ser Gly Gly Ser Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys Ala Ala Trp Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
 100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 84
 <211> 122
 <212> PRT
 <213> Homo sapiens

<400> 84
 Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
 1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr
 20 25 30

Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
 35 40 45

PRD3422WOPCT_SL.TXT

Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe
50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ala Arg Ile Gly Phe Tyr Gly Arg Ser Phe Arg Ile Phe Asp Tyr Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 85
<211> 123
<212> PRT
<213> Homo sapiens

<400> 85
Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr
20 25 30

Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
35 40 45

Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe
50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ala Arg Val Tyr Ser Phe Gly Gly Arg His Lys Ala Leu Phe Asp Tyr
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 86
<211> 123
<212> PRT
<213> Homo sapiens

PRD3422WOPCT_SL.TXT

<400> 86

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr
20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Val Asp Arg Ser Phe Gly Arg Ser Arg Tyr Thr Leu Asp Tyr
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 87

<211> 123

<212> PRT

<213> Homo sapiens

<400> 87

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Ser Ser Tyr
20 25 30

Gly Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Gly Ile Ile Pro Ile Phe Gly Asn Ile Asn Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

PRD3422WOPCT_SL.TXT

Ala Arg Val Ser Arg Arg Phe Lys Arg Phe Ala Tyr Tyr Phe Asp Tyr
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 88
<211> 118
<212> PRT
<213> Homo sapiens

<400> 88
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Gly Tyr
20 25 30

Thr Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Thr Asn Tyr Ala Gln Lys Leu
50 55 60

Gln Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Val Ala Leu Arg Val Ala Leu Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser
115

<210> 89
<211> 119
<212> PRT
<213> Homo sapiens

<400> 89
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

PRD3422WOPCT_SL.TXT

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Ser Asn Phe Leu Pro Val Val Phe Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ser
115

<210> 90
<211> 705
<212> DNA
<213> Homo sapiens

<400> 90
atgcgggtgc tggcccagct gctgggactg ctgctgctgt gcttccctgg cgccagatgc 60
gacatccaga tgaccagag ccccagcagc ctgagcgcca gcgtgggcga ccgggtgacc 120
atcacctgca aggccagcca gaacgtggcc acccacgtgg gctggtacca gcagaagccc 180
ggcaaggccc ccaagcggct gatctacagc gccagctacc ggtacagcgg cgtgccagc 240
cggttcagcg gcagcggcag cggcaccgag ttcaccctga ccatcagcaa cctgcagccc 300
gaggacttcg ccacctacta ctgccagcag tacaaccggt acccctacac cttcggccag 360
ggcaccaagc tggagatcaa gcgtacggtg gctgcaccat ctgtcttcat cttcccgcca 420
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taacttctat 480
cccagagagg ccaaagtaca gtggaagggtg gataacgccc tccaatcggg taactcccag 540
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg 600
ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc 660
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gttga 705

<210> 91
<211> 117
<212> PRT
<213> Homo sapiens

<400> 91
Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30

Asn Val His Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu
 35 40 45

Ala Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Ser Ala Leu Met
 50 55 60

Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val Val Leu
 65 70 75 80

Thr Met Thr Asn Met Arg Ala Glu Asp Thr Ala Thr Tyr Tyr Cys Ala
 85 90 95

Arg Asp Gly Ile Arg Leu Arg Phe Ala Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> 92
 <211> 113
 <212> PRT
 <213> Homo sapiens

<400> 92
 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
 20 25 30

Ser Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
 85 90 95

Tyr Tyr Ser Thr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110

Lys

PRD3422WOPCT_SL.TXT

<210> 93
 <211> 108
 <212> PRT
 <213> Homo sapiens

<400> 93
 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Arg Lys Ser
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Tyr Thr Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Phe Arg Ala Pro Ile
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Lys
 100 105

<210> 94
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 94
 Glu Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Lys Ala Ser Gln Asn Val Ala Thr His
 20 25 30
 Val Gly Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Tyr Ser Ala Ser Tyr Arg Tyr Ser Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Ser
 65 70 75 80

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

```
<210> 95
<211> 11
<212> PRT
<213> Homo sapiens
```

<400> 95
Arg Ala Ser Gln Ser Val Arg Lys Ser Leu Ala
1 5 10

```
<210> 96
<211> 1395
<212> DNA
<213> Homo sapiens
```

<400> 96						
atggcctggg	tctggaccct	gctgttcctg	atggccgctg	cccagagcat	ccaggcccag	60
gtgcagctgg	tgcagagcgg	cgccgaggtg	aagaagcccg	gcgccagcgt	gaaggtgagc	120
tgcaaggcca	gcggttacag	cttcaccggc	tacaccatga	actgggtgcg	gcaggccccc	180
ggccagggcc	tggagtggat	gggcctgata	aaccctaca	acagcgacac	caactacgcc	240
cagaagctgc	agggccgggt	gaccatgacc	accgacacca	gcaccagcac	cgcctacatg	300
gagctgcgga	gcctgcggag	cgacgacacc	gccgtgtact	actgcgcccg	ggtggccctg	360
cgggtggccc	tggactactg	gggccagggc	accctgggtga	ccgtgagcag	cgcctccacc	420
aagggcccat	ccgtcttccc	cctggcgccc	tgtctccagga	gcacctccga	gagcacagcc	480
gccctgggct	gcctgggtcaa	ggactacttc	cccgaaccgg	tgacgggtgtc	gtggaactca	540
ggcgccctga	ccagcggcgt	gcacaccttc	ccggctgtcc	tacagtccctc	aggactctac	600
tccctcagca	gcgtggtgac	cgtgccctcc	agcagcttgg	gcacgaaaac	ctacacctgc	660
aacgtagatc	acaagcccag	caacaccaag	gtggacaaga	gagttgagtc	caaatatggt	720
cccccattgc	caccatgccc	agcacctgag	gccgccgggg	gaccatcagt	cttcctgttc	780
cccccaaac	ccaaggacac	tctcatgata	tcccggaccc	ctgaggtcac	gtgcgtgggtg	840
gtggacgtga	gccaggaaga	ccccgaggtc	cagttcaact	ggtacgtgga	tggcgtggag	900
gtgcataatg	ccaagacaaa	gccgcggggag	gagcagttca	acagcacgta	ccgtgtggtc	960
agcgtcctca	ccgtcctgca	ccaggactgg	ctgaacggca	aggagtacaa	gtgcaaggtc	1020
tccaacaaag	gcctcccgtc	ctccatcgag	aaaaccatct	ccaaagccaa	agggcagccc	1080
cgagagccac	aggtgtacac	cctgccccca	tcccaggagg	agatgaccaa	gaaccaggtc	1140
agcctgacct	gcctgggtcaa	aggcttctac	cccagcgaca	tgcgccgtgga	gtgggagagc	1200

PRD3422WOPCT_SL.TXT

aatgggcagc cggagaacaa ctacaagacc acgcctcccg tgctggactc cgacggctcc 1260
 ttcttcctct acagcaggct aaccgtggac aagagcaggt ggcaggaggg gaatgtcttc 1320
 tcatgctccg tgatgcatga ggctctgcac aaccactaca cacagaagag cctctccctg 1380
 tctctgggta aatga 1395

<210> 97
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 97
 Ala Ala Phe Asp Phe Gly Arg Arg Ala Val Arg Leu Asp Tyr
 1 5 10

<210> 98
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 98
 Lys Ser Ser Gln Ser Val Leu Tyr Ser Ser Asn Asn Lys Asn Tyr Leu
 1 5 10 15

Ala

<210> 99
 <211> 452
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 99
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Thr Tyr
 20 25 30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Arg Ile Arg Ser Lys Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Ala
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Ser
 65 70 75 80

PRD3422WOPCT_SL.TXT

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95
 Tyr Cys Ala Arg His Gly Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe
 100 105 110
 Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr
 115 120 125
 Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser
 130 135 140
 Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu
 145 150 155 160
 Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His
 165 170 175
 Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser
 180 185 190
 Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys
 195 200 205
 Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu
 210 215 220
 Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala
 225 230 235 240
 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 245 250 255
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
 260 265 270
 Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu
 275 280 285
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr
 290 295 300
 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
 305 310 315 320
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser

325

335

Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
340 345 350

Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val
355 360 365

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
370 375 380

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
385 390 395 400

Pro Val Leu Asp Ser Asp Gly Ser Phe Leu Leu Tyr Ser Lys Leu Thr
405 410 415

Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val
420 425 430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
435 440 445

Ser Leu Gly Lys
450

<210> 100

<211> 215

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 100

Gln Thr Val Val Thr Gln Glu Pro Ser Leu Thr Val Ser Pro Gly Gly
1 5 10 15

Thr Val Thr Leu Thr Cys Arg Ser Ser Thr Gly Ala Val Thr Thr Ser
20 25 30

Asn Tyr Ala Asn Trp Val Gln Gln Lys Pro Gly Gln Ala Pro Arg Gly
35 40 45

Leu Ile Gly Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe
50 55 60

Ser Gly Ser Leu Leu Gly Gly Lys Ala Ala Leu Thr Leu Ser Gly Val

PRD3422W0PCT_SL.TXT

65

70

75

80

Gln Pro Glu Asp Glu Ala Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn
85 90 95

Leu Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln Pro
100 105 110

Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu
115 120 125

Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro
130 135 140

Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys Ala
145 150 155 160

Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala
165 170 175

Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Arg
180 185 190

Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr
195 200 205

Val Ala Pro Thr Glu Cys Ser
210 215