

(54) Title  
**PROBES TO MYCOBACTERIUM AVIUM MYCOBACTERIUM INTRACELLULARE AND MYCOBACTERIUM PARATUBERCULOSIS**

International Patent Classification(s)  
(51)<sup>5</sup> **C12N 015/11 C12Q 001/68**

(21) Application No. : **50305/93** (22) Application Date : **26.10.93**

(30) Priority Data

(31) Number (32) Date (33) Country  
**973342 06.11.92 US UNITED STATES OF AMERICA**

(43) Publication Date : **19.05.94**

(44) Publication Date of Accepted Application : **23.01.97**

(71) Applicant(s)  
**BECTON DICKINSON AND COMPANY**

(72) Inventor(s)  
**COLLEEN MARIE NYCZ; JAMES LEE SCHRAM; DARYL DEE SHANK; GLENN PHILANDER VONK**

(74) Attorney or Agent  
**F B RICE & CO , 28A Montague Street, BALMAIN NSW 2041**

(56) Prior Art Documents  
**AU 642140**  
**AU 663180**  
**WO 93/09251**

(57) Claim

1. A probe having a sequence consisting essentially of Seq ID No: 1, Seq ID No: 2, Seq ID No: 3, Seq ID No: 4, Seq ID No: 5, Seq ID No: 6, Seq ID No: 7, Seq ID No: 8, Seq ID No: 9, Seq ID No: 10, Seq ID No: 11, Seq ID No: 12, Seq No: 13, Seq ID No: 14, or Seq ID No: 15, and the modified backbone, modified nucleotide, labelled forms, and ribonucleic acid forms thereof.

**AUSTRALIA**

**Patents Act 1990**

**BECTON DICKINSON AND COMPANY**

ORIGINAL  
COMPLETE SPECIFICATION  
STANDARD PATENT

Invention Title:

**"Probes to mycobacterium avium mycobacterium  
intracellulare and mycobacterium paratuberculosis"**

The following statement is a full description of this invention including the best method of performing it known to us:-

FIELD OF THE INVENTION

5       The invention is in the field of molecular biology. In particular, the invention relates to diagnostics involving nucleic acid probes. More particularly, the invention relates to nucleic acid probes for detecting, amplifying, and isolating Mycobacterium avium, Mycobacterium intracellulare, and Mycobacterium paratuberculosis.

10       BACKGROUND OF THE INVENTION

15       Health care providers are encountering a significant increase in cases of mycobacterial infections. Many of these new cases are related to the AIDS epidemic. Physicians rely on clinical microbiologists to assist them in diagnosing mycobacterial infection. The diagnosis of such infections is, however, largely dependent on acid-fast staining and cultivation of the organism, followed by biochemical assays. These are time consuming processes. Hence, there is a

continuing need for new, and particularly rapid, methods of diagnosing mycobacterial infections.

SUMMARY OF THE INVENTION

5 The present invention provides nucleic acid probes and methods that are definitive, rapid, and cost effective for detecting, amplifying, and isolating Mycobacterium avium, Mycobacterium intracellulare, and Mycobacterium paratuberculosis. The invention also provides methods for isolating sequences of M. avium, M. intracellulare, and M. paratuberculosis.  
10 In addition, probes of the invention are useful for amplification of sequences to which they hybridize.

Probes of the invention can also be used to construct amino acid sequences for which they encode and antibodies raised to recognize the sequence.

15 Probes of the invention consist essentially of the nucleic acid sequences in Sequence ID No: 1-15. The probes can locate either M. avium, M. intracellulare or M. paratuberculosis or any combination thereof. Since the probes of the invention are unique, there is no cross-hybridization to nucleic acid from  
20 other microorganisms.

The probe sequences described in the invention are derived

from plasmid pMav29 originally identified by Fries et al.,  
Molec. Cell Probes 4:87 (1990). Plasmid pMav29 had  
cross-reactivity to M. intracellulare and M. avium, but also  
strong cross-reactivity to M. bovis and lesser cross-reactivity  
5 to M. tuberculosis. The probes described in this invention  
offer substantial improvement over that described by Fries et  
al. since specific cross-reactivity is observed only for M.  
intracellulare, M. avium, and M. paratuberculosis.

Consequently, these probes are useful for clinical diagnosis of  
10 M. intracellulare and M. avium complex infection which  
currently comprise approximately 20% of pathogenic isolates in  
mycobacteriology laboratories (Good, R. E. et al.(1982) J.  
Infec. Dis. 146:829-33). Diagnosis is becoming important in  
management of HIV infection where an estimated 30-50% develop  
15 M. intracellulare or M. avium co-infection (Morris, S. C. et  
al. (1991) J. Clin. Microbiol 29:2715-19). The probes of the  
invention are also useful for diagnosis of M. paratuberculosis  
infection implicated in Crohn's disease (Wayne, L. G. et al.  
(1992) Clin Microbiol.Rev. 5:1-25) and identified as the  
20 etiological agent of Johne's (veterinary) disease (Barclay et  
al. (1985) J. Bacteriology 164:896-903).

#### DETAILED DESCRIPTION

The present invention provides a probe having a sequence  
consisting essentially of Seq ID No:1-15, and the modified

backbone (includes modified sugar and phosphates groups (e.g. thiophosphates)), modified nucleotide, labelled forms, and ribonucleic acid forms thereof.

5 A method of amplifying Mycobacterium avium, Mycobacterium intracellulare, and Mycobacterium paratuberculosis which comprises the use of one or more nucleic acid sequences selected from the sequences consisting essentially of Seq ID No:1 through 15, and ribonucleic acid forms thereof are also provided.

10 Also provided is a method of amplifying Mycobacterium avium, Mycobacterium intracellulare and Mycobacterium paratuberculosis and detecting the amplified product which comprises the use of one or more sequences selected from the sequences consisting essentially of Seq ID No:1 - 15, and the modified backbone, modified nucleotide, labelled forms, and  
15 ribonucleic acid forms thereof.

A kit comprising a nucleic acid sequence selected from the sequences consisting essentially of Seq ID No:1 - 15, and the modified backbone, modified nucleotide, labelled forms, and  
20 ribonucleic acid forms thereof is also provided.

Table 1 sets forth probes of the invention and where they hybridize to regions of the pMav29 clone identified in Molec. Cell. Probes 4:87 (1992) (left to right is 5' → 3'):

TABLE 1

	<u>Probe Sequence</u>	<u>Probe Sequence Position in pMav29*</u>
5	(Seq ID No: 1)  TGGCCAACT GTGGGCGCAG GCCTGCGAGT GGGAACCGGT GACTCCAAA ACCTTGCGGC TCTTACAAGT CGGTGGCGCC AAGCTGGAGC CCGAGGA	72-168
10	(Seq ID No: 2)  TTGAATAGTC GGTACTTGT TGA CTCTCG GGCTCCA	156-168
15	(Seq ID No: 3)  TTGAAGTAAC CGACTATTGT TGA CTGGCCA AACTGTG	72-84
	(Seq ID No: 4)  TTGAATAGTA GGTAAGTTGT TGACACTTGT AAGAGCC	129-141
20	(Seq ID No: 5)  TTGAAGTAAC CGACTATTGT TGA CTGCGAG TGGAAC	95-107
	(Seq ID No: 6)  GGGAACCGGT GACTC	102-116
25	(Seq ID No: 7)  CAAAAACCTT GCGGC	117-131
	(Seq ID No: 8)  AACCGGTGAC TCCA	105-118

(Seq ID No: 9)

AAAACCTTGC GGC

119-131

(Seq ID No: 10)

AACCGGTGAC TCCA

105-118

5

(Seq ID No: 11)

GCGAGTGGGA ACCGGTGACT CCAAAAACCT  
TGCGGCTCTT ACAAGTC

96-142

(Seq ID No: 12)

ACCGGTGACT CCAAAAACCT TGCGGCTCTT  
ACAAGTCGGT GGCGCCAAG

106-154

10

(Seq ID No: 13)

GATCCCAGCC CCGAGGCGGC CTTCGCCAGT  
CGACCGCCAT GGCCTCACGG

1-50

(Seq ID No: 14)

GGGAACCGGT GACTCCAAAA ACCTTGCGGC  
TCTTACAAGT CGGTGGCGCC

102-151

15

(Seq ID No: 15)

GGCCAAACTG TGGGCGCAGG CCTGCGAGTG  
GGAACCGGTG ACTCCAAAAA CCTTGCGGCT  
CTTACAAGTC GGTGGCGCCA AGCTGGAGCC  
CGAGGACGCC CCGCCTG

73-180

20

\* Numbering according to pMav29 sequence having cytidine insert  
(See Example 1).



Probes of the invention are useful for amplifying nucleic acid sequences which hybridize to the probes. These nucleic acid sequences could then be cloned and used to express protein or an amino acid sequence based on the nucleic acid sequence or  
5 can be chemically synthesized. This amino acid sequence can then be used as an immunogen to make antibodies. Due to the degeneracy of the genetic code (many amino acids are selected by more than one codon) variations in the probe sequence will result in the same antibody specificity. Such antibodies can  
10 be polyclonal or monoclonal.

Probes of the invention can be deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). DNA is composed of nucleotides based on purines (adenine and guanine) and pyrimidines (cytosine and thymine), with adenine base pairing  
15 (complementary) to thymine and guanine base pairing to cytosine. RNA is similar to DNA, except for the sugar differences (ribose instead of deoxyribose) and the base uracil, which is present instead of thymine. Also, modified nucleosides can be used in constructing probes of the  
20 invention. Modified nucleosides are typically used to obtain a probe capable of withstanding more stringent hybridization conditions, capable of easier detection, and the like. Preferably the probes are DNA.

The present invention can detect the presence of M. avium,  
25 M. intracellulare, M. paratuberculosis, all three, or any combination thereof, in a variety of samples. Samples can

include clinical specimens such as fecal material, blood, sputum, saliva, urine, plaque samples, tissue samples, fixed tissues, and tissue culture monolayers obtained by standard techniques such as lavage, scraping, or biopsy. The location  
5 or format of the Mycobacterium is not key to the invention, the ability to specifically detect, amplify, or isolate M. avium, M. intracellulare, M. paratuberculosis, or combination thereof, is key to the invention.

Samples obtained for use with probes of the invention can  
10 be utilized directly or amplified before using probes of the invention. A variety of amplification methods are available. For example, polymerase chain reaction (PCR), PCR Technology, H.A. Erlich, Ed. (Stockton Press, New York, NY, 1989), transcription-based amplification system (TAS), Proc. Natl. Acad. Sci. USA 86:1173 (1989), ligation amplification  
15 reaction (LAR), Genomics 4:560 (1989), ligase based amplification system (LAS), Gene 89:117 (1990), Strand displacement amplification (SDA) Proc. Natl. Acad. Sci. USA 89:392 (1992) and Nucleic Acids Res., 20:7 (1992), and Q B replicase, Infect. Dis. 162:13 (1990). The goal of any  
20 sample preparation is to eliminate false positives and improve sensitivity. Such a goal is obtained by taking into account the way samples are prepared, and the selection of a medium or substance in which the sample is prepared.

25 Probes of the invention can be prepared in a variety of ways and therefore are not limited to any particular

preparation means. Suitable means for preparing the probes of the invention include using replication vectors and cloning the probe as a library, followed by appropriate screening procedures, and growing the vector in a suitable host.

5 Purification and isolation will result in the probe being separated from the vector with select restriction enzymes. Preferably the probes are synthesized using commercially available methods and equipment. For example, the solid phase phosphoramidite method can be used to produce the probes of the  
10 invention. C. Caruthers, et al., Cold Spring Harbour Symp. Quant. biol., 47:411-418 (1982), and Adams, et al., J. Am. Chem. Soc., 105:601 (1983).

Typically, synthesis of nucleic acid probes utilize phosphoramidite derivatives that are appropriately protected  
15 for synthesis. See S. P. Adams et al., J. Am. Chem. Soc., 105:661 (1983), L. J. McBride and M. H. Caruthers, Tetrahedron Lett., 24:245 (1983), and N. D. Sinha et al., Nucleic Acids Res., 12:4539 (1984). Methods for incorporating phosphoramidites into a nucleic acid strand  
20 include the use of solid phase, solution phase, triesters, and H-phosphonate intermediates as generally illustrated by Froehler et al., Nuc. Acids Res., 14:5399 (1986), Narang et al., Methods Enz., 68:90 (1979) and Ogilvie, K.K. et al., Proc., Natl. Acad. Sci. U.S.A., 85:5764 (1988).

Probes of the invention can be conventionally synthesized by the modified phosphotriester method using fully protected deoxyribonucleotide building blocks. Such synthetic methods can be carried out in substantial accordance with the procedure of Itakura et al., (1977), Science, 198:1056 and Crea et al., (1978), Proc. Nat. Acad. Sci. U.S.A., 75:575, Hsiung et al., (1983), Nucleic Acid Research, 11:3227, and Narang et al., (1980), Methods in Enzymology, 68:90. In addition to manual procedures, the probes can be synthesized using automated synthesizers, such as the Systec 1450A or ABI 380A Synthesizers.

Probes of the invention can be utilized with naturally occurring sugar-phosphate backbones as well as modified backbones including phosphorothioates, dithionates, alkyl phosphonates and  $\alpha$ -nucleotides. Modified sugar-phosphate backbones are generally illustrated by Miller and T'so, Ann. Reports Med. Chem., 23:295 (1988) and Moran et al., Nuc. Acids Res., 14:5019 (1987).

Use of probes in detection methods include Northern blots (RNA detection), Southern blots (DNA detection), western blots (protein detection), dot blots (DNA, RNA, or protein detection), and Slot blots (DNA, RNA or protein). Other detection methods include kits containing probes on a dipstick setup and the like.

To detect hybrid molecules formed from using the probes of the invention, typically a detectable marker is added to one of

the probes. Probes can be labelled by several methods. Probes can be radiolabelled and detected by autoradiography. Such labels for autoradiography include  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , and  $^{32}\text{P}$ . Typically the choice of radioactive isotopes depends on research preferences involving ease of synthesis, stability, and half lives of the isotopes. Other detectable markers include ligands, fluorophores, chemiluminescent agents, electrochemical labels, time-resolved fluorophores, enzymes, and antibodies. Other detectable markers for use with probes of the invention include biotin, radionucleotides, enzyme inhibitors, co-enzymes, luciferins, paramagnetic metals, spin labels, and monoclonal antibodies. The choice of label dictates the manner in which the label is bound to the probe.

Radioactive nucleotides can be incorporated into probes of the invention by several means. Such means include nick translation of double-stranded probes, copying single-stranded M13 plasmids having specific inserts with the Klenow fragment of DNA polymerase I of E. coli or other such DNA polymerase in the presence of radioactive dNTP, transcribing cDNA from RNA templates using reverse transcriptase in the presence of radioactive dNTP, transcribing RNA from vectors containing strong promoters such as SP6 promoters or T7 promoters using SP6 or T7 RNA polymerase in the presence of radioactive rNTP, tailing the 3' ends of probes with radioactive nucleotides using terminal transferase, and by phosphorylation of the 5' ends of probes using gamma  $^{32}\text{P}$  ATP and polynucleotide kinase.

Non-radioactive probes of the invention can be labelled by indirect means. For example, a ligand molecule can be covalently bound to a probe of the invention. The ligand can then bind to an anti-ligand molecule which is either inherently detectable or covalently bound to a signal system such as a detectable enzyme, a fluorescent compound, or chemiluminescent compound.

Probes of the invention can also be conjugated directly to signal generating compounds by such means as conjugation with an enzyme. Suitable enzymes for labels include hydrolases, particularly phosphatases, esterases, glycosidases, and oxidoreductases such as peroxidases. Fluorescent compounds include fluorescein and derivatives thereof, rhodamine and derivatives thereof, dansyl, umbelliferone, and the like.

Various hybridization conditions can be employed when using probes of the invention. The particular hybridization technique is not essential to the invention. Hybridization techniques are generally described in "Nucleic Acid Hybridization, A Practical Approach", edited by Hanes, B.D. and Higgins, S.J., IRL Press, 1985, Gall and Pardue (1969), Proc. Natl. Acad. Sci., USA, 62:378-383, John Burnsteil and Jones (1969) Nature 223:582-587, and Southern, E., J. Mol. Biol. 98:503 (1975). With routine experimentation the conditions which permit satisfactory hybridization are easily obtained.

The correct hybridization complex can be detected in accordance with the label used with the probes. Therefore, the choice of label will guide the choice of detection methods. For example, when the label is a hapten or antigen, the hybridization complex can be detected using antibodies. Typically, the antibody will be attached to a fluorescent or enzyme molecule which can be detected. Tijssen, P., "Practice in Theory of Enzyme Immunoassay, Laboratory Techniques in Biochemistry and Molecular Biology", Burdon, R. H., Van Knippenberg, P.H., Eds., Elsevier, 1985, PP.9-20. When the label is radioactive, the hybridization complex is exposed to X-ray film. Where the label is fluorescent, the sample is detected by irradiation with light of a particular wavelength. Enzyme labels are detected by incubation with appropriate substrate.

In addition to using probes of the invention for detecting sequences of mycobacteria, the probes of the invention can be used for isolating sequences of mycobacteria. In many diagnostic applications, merely detecting the presence of a sequence of mycobacteria in a sample is all that is desired. However, there are situations where it is desirable to isolate a sequence of mycobacteria. For example, a sequence can be isolated as a preliminary step to separate extraneous DNA and then the specific isolated DNA sequence is amplified using DNA amplification techniques.

Probes of the invention are also useful for amplification of mycobacteria sequences. For example, probes of the invention can be used as primers in a variety of amplification protocols such as polymerase chain reaction, transcription-based amplification system, ligation amplification reaction, ligase based amplification system and QB replicase (referenced above).

The probes of the invention can be conveniently provided in the form of a kit. The kits can also comprise probes in which detection can also be specific, for example, the probes can be designed for fluorescence, radioactive, and chemiluminescence detection.

The following examples illustrate the specific embodiments of the invention described in this document. As would be apparent to skilled artisans, various changes and modifications are possible and are contemplated within the scope of the invention described.



## MATERIALS AND METHODS

### Preparation of Capture Oligodeoxynucleotide Probe

The oligodeoxynucleotide capture probes were synthesized as described below. First, the capture oligomer was prepared using a DNA synthesizer (Model 380B, Applied Biosystems, Foster City, CA), and Biotin On™ reagent (Clontech, Palo Alto, CA) such that there were three biotin molecules (BBB) at the 5' terminus of the oligodeoxynucleotide. Purification was accomplished by reverse phase High Pressure Liquid Chromatography (HPLC, Brownlee Lab Aquapore RP 300 Column - 220 x 4.6 mm, C8 column 7 particle, 300 A pore size) with a UV monitor at 254 nm and a gradient of 14 to 44% Buffer B over one hour (Buffer B: 0.1M Triethylamine-Acetate pH 7 with 50% Acetonitrile; Buffer A: 0.1M Triethylamine-Acetate, pH 7) and a flow rate of 1 ml/minute.

The oligodeoxynucleotide detector probes were synthesized using a DNA synthesizer (Model 380B, Applied Biosystems, Foster City, CA) and a 3'-amino-modifier column (Glenn Research, Sterling, VA). This yielded an oligodeoxynucleotide with a 3' amine terminus required for subsequent conjugation with alkaline phosphatase. Calf intestine alkaline phosphatase (AP, enzyme immunoassay grade, Boehringer Mannheim, Indianapolis, IN) was dialyzed overnight at 4°C against 50 mM potassium

phosphate pH 7.5 and subsequently centrifuged to remove aggregates. The AP (4 mL, 10 mg/mL) was combined with a solution (40 uL) of succinimidyl-4-(p-maleimidophenyl)butyrate (SMPB, obtained from Pierce, Rockford, MD, 50 mM) dissolved in N,N'-dimethylformamide (DMF, Aldrich, Milwaukee, WI) and allowed to react in the dark at room temperature for 30 minutes. The derivatized AP was purified using a NAP-25 column (Pharmacia, Piscataway, NJ) previously equilibrated with 50 mM potassium phosphate pH 7.5 (degassed and purged with N<sub>2</sub>). The absorbances of the NAP-25 column fractions were read at 260 and 280 nm using an extinction coefficient of 0.75 ml/umole cm<sup>-1</sup>. Typically, about 170 nmoles of derivatized AP were obtained and stored on ice (less than 2 hours) until conjugation with the derivatized oligodeoxynucleotide.

The 3' amino-oligodeoxynucleotide (98.4 ul of 508.2 uM, 50 nmoles) was diluted in 13.4 ul of 1M potassium phosphate (pH 7.2) and mixed with 27 ul of a solution of n-succinimidyl-3-(2-pyridyldithio)propionate (50 mM, SPDP, Pierce, Rockford, IL) diluted in DMF. This mixture was incubated in the dark for 1 hour at room temperature. A solution of dithiothreitol (DTT, 1M) in 50 mM potassium phosphate (pH 7.5) was added to the oligodeoxynucleotide/DMF mixture to a final concentration of 0.1M DTT and allowed to incubate for 15 minutes at room temperature. Excess DTT and SPDP were separated from the derivatized oligodeoxynucleotide

by elution over a NAP-25 column with 50 mM potassium phosphate (pH 7.5). Within 10 minutes of purification, the reduced oligodeoxynucleotide was mixed with the SMPB derivatized AP. Rapid mixture of the reduced oligomer and the SMPB derivatized AP is necessary to prevent reoxidation of the thiolated oligomer. The resulting solution was incubated 2-4 hours at room temperature, then overnight at 4°C and quenched by addition of 1/100th the original volume of 50 mM beta-mercaptoethanol in 50 mM potassium phosphate (pH 7.5).

10 The crude conjugate was concentrated using a Centriprep 30™ centrifugal concentrator (Amicon, Danvers, MA) to approximately 2 ml. This material was further purified by HPLC using a DEAE-5PW column (7.5 mm x 7.5 cm), a gradient of 0 to 66% Buffer B (Buffer B: 20 mM Tris, 1M NaCl pH 7.5, Buffer A: 20 mM Tris pH 7.5) and a flow rate of 1 ml/minute. Absorbance was monitored at 254 nm. Fractions with  $A_{260}/A_{280}$  equal to 1 correspond to conjugate and were pooled. The protein concentration of the conjugated oligodeoxynucleotide was then determined (BCA Protein Assay Kit, Pierce, Rockford, IL).

20 The activities of the alkaline phosphate (AP) detector oligodeoxynucleotide probes were determined as follows. The conjugate was diluted to 5 ug/ml in 50 mM Tris-HCl, 100 mM NaCl, 1 mM  $MgCl_2$ , 1 mg/ml BSA, pH 7.5. The substrate, 4-nitrophenylphosphate (pNPP, 5mM), was dissolved in 1 M


diethanolamine, 1 mM  $MgCl_2$ , pH 9.8. AP activity was assayed as follows at 25°C. The conjugate (5  $\mu$ l) was diluted into 2 ml of the substrate solution and the change in absorbance monitored at 405 nm using a Hewlet Packard 8452 spectrophotometer.

5

The reaction rates were calculated from the linear region of the kinetic plots using the extinction coefficient of p-nitrophenol at 405 nm ( $18500 M^{-1}cm^{-1}$ ). The specific activity of the AP detector oligodeoxynucleotide probes were determined to be 850-1300  $\mu$ mole/minute/mg.

10

The purified AP detector oligodeoxynucleotide probe was diluted to 2  $\mu$ M in 20 mM Tris, 1M NaCl, 0.05% sodium azide, 50  $\mu$ g/ml sonicated salmon sperm DNA, pH 7.5, and stored thereafter at 4°C.



EXAMPLES

EXAMPLE 1

Development of Mycobacterium avium/Mycobacterium intracellulare  
Probe Sequences for Strand Displacement Amplification

5           The target sequence for the invention was derived from the  
clone pMav29 as disclosed by Fries, J. M. W., et al. (1990)  
10           Molec. Cell. Probes 4:87-105. The specificity of this clone  
was tested against genomic DNA from a number of mycobacteria  
(pre-digested with the restriction endonuclease PstI) using  
Southern blot analysis (Southern, E.M. (1975) J. Mol. Biol.  
15           98:503-517). Clone pMav29 was cross reactive to both M.  
avium and M. intracellulare, with lesser cross reactivity to M.  
tuberculosis and M. bovis. A similar clone, pMav22 was found  
to hybridize primarily to M. avium with lessor cross reactivity  
to M. kansasii and M. intracellulare. Fries et. al. continued  
work on pMav22 and developed a probe sequence specific for M.  
20           avium.

          In order to identify a region of pMav29 specific for only  
M. avium and M. intracellulare, four 50 mer  
oligodeoxynucleotides were synthesized spanning the first 150  
bases of pMav29 and a region from position 281-320 having  
relatively low GC/GCAT content. The oligomers were

5'-<sup>32</sup>P-labelled using T4 polynucleotide kinase and used to probe Southern blots of PstI digests of various genomic DNA preparations (Table I). The blots were visualized by autoradiography using X-ray film. The film was placed next to the blot, then developed using standard procedures. Since the position of each preparation of genomic DNA on the blot is known, it is possible to determine the cross-reactivity of the 5'-<sup>32</sup>P-labelled oligomers by the position of the dark spot (s) on the autoradiogram. That is, spots appear only where specific hybridization of 5'-<sup>32</sup>P-labelled oligomers to membrane bound genomic DNA occurred. The results are presented in Table 2.

TABLE 1

Species Test for Cross reactivity in Southern Blot Analysis

M. tuberculosis  
M. scrofulaceum  
M. kansasii  
M. intracellulare  
M. gordonae  
M. fortuitum  
M. chelonae  
M. bovis  
M. avium

Escherichia coli  
Neisseria gonorrhoeae  
Murine DNA isolated from McCoy cell tissue culture  
DNA isolated from salmon sperm

TABLE 2

Specificity of pMav29 Oligomers in Southern Blot Analysis  
(Numbering is according to pMav29 sequence described by Fries  
et al.)

5	Oligomer Position in pMav29	Mycobacteria Species Cross Reactivity
	1-50	<u>M. avium</u> , <u>M. intracellulare</u>
	51-100	<u>M. avium</u> , <u>M. intracellulare</u> , <u>M. kansasii</u> , <u>M. gordonae</u> , <u>M. fortuitum</u> , <u>M. scrofulaceum</u>
10	101-150	<u>M. avium</u> , <u>M. intracellulare</u>
	72-167	<u>M. avium</u> , <u>M. intracellulare</u>
	1-377 (pMav29)	<u>M. avium</u> , <u>M. intracellulare</u> , <u>M. kansasii</u> , <u>M. bovis</u> , <u>M. tuberculosis</u>
15		

Initial Strand Displacement Amplification (SDA) experiments indicated that the region 72-167 of pMav29 would be amenable to amplification. Specificity of this region was determined by

cloning/sequencing a Polymerase Chain Reaction (PCR)  
amplification product (position 72-167) obtained from M. avium  
genomic DNA. The sequence is in agreement with that published  
by Fries et. al. with the exception of a cytosine (C) insert  
5 between positions 93 and 94 of pMav29. The specificity of the  
clone having the C insert was rechecked as described above in  
Southern blot hybridization to PstI digested Mycobacterial  
DNA. This region (72-168, 97 bp) retains the original  
specificity of pMav29 (positions 72-167) for M. avium and M.  
10 intracellulare with negligible cross reactivity to other  
species (Table 2).

#### EXAMPLE 2

Strand Displacement Amplification of Mycobacterium  
avium-intracellulare complex

15 As previously described, Southern Blot analysis of pMav29  
indicated cross reactivity to M. tuberculosis, M. kansasii and M.  
bovis as well as M. avium and M. intracellulare (Table 2) Example  
1.

20 Southern blot analysis of subsequences derived from the  
pMav29 clone indicated regions of specificity for Mycobacterium  
avium and Mycobacterium intracellulare. Based on this  
information, several other factors were considered: 1) G+C



content should be close to 50%, 2) secondary structure should be minimal, 3) the target region should be large enough to accommodate the detector and capture probe.

Several regions of specificity based on Southern blot analysis were evaluated for G+C content. Bases 1-50 have specificity for M. avium and M. intracellulare but have a G+C content of 74%. Bases 105-153 (49 mer) and bases 95-141 (47 mer) showed G+C content of 55% and 53% respectively, which is close to the acceptable range.

Computer analysis of the secondary structure for these two potential target sequences was carried out. From this analysis it was noted that for the 49 mer target sequence, a hairpin due to a palindromic sequence could form between bases 144 and 151. The 47 mer target sequence showed no obvious secondary structure. However, one potential drawback for the 47 mer target was the relatively small number of nucleotides available for binding of detector and capture probes as discussed in following Example 3. With this in mind, SDA primer oligos were prepared for amplification of the 47 mer target.

Several primer sets and combinations were synthesized and tested for sensitivity. The best set of primers were chosen and tested under various conditions. Conditions that were evaluated included addition of glycerol or 1-methyl-2-pyrrolidinone. Sensitivity of the assay was increased by the addition of glycerol. Assay sensitivity was

increased even more by using 3% 1-methyl-2-pyrrolidinone instead of 5-10% glycerol. Assembled reaction conditions contained the following:

#### Reaction Mix

5 50mM potassium phosphate pH 7,6  
100 ug/ml bovine serum albumin  
1mM dithiothreitol  
1mM dCTP  
1mM dGTP  
10 1mM dATP alpha thio  
1mM TTP  
6mM magnesium chloride  
3% (v/v) 1-methyl-2-pyrrolidinone

#### Enzymes

15 5 units exonuclease free klenow (United States Biochemicals,  
Cleveland, OH)  
150 units HincII

#### Primer Mix

20 .05 uM 5'TTGAATAGTCGGTTACTTGTTGACTCCTCGGGCTCCA3'  
(Seq ID No: 2)  
.05 uM 5'TTGAAGTAACCGACTATTGTTGACTGGCCAACTGTG3'  
(Seq ID No: 3)  
0.5 uM 5'TTGAATAGTAGGTAAGTTGTTGACACTTGTAAGAGCC3'  
(Seq ID No: 4)

0.5 uM

5'TTGAAGTAACCGACTATTGTTGACTGCGAGTGGGAAC3'  
(Seq ID No: 5)

Two SDA primers (Seq ID No:2 and 4) bind to the anticoding strand and two SDA primers (Seq ID No:3 and 5) bind to the coding strand of the target genomic DNA sequence (Seq ID No:1).

All concentrations are final reagent concentrations in a sample volume of 50 ul. Target DNA was isolated from killed bacterial cells of M. avium and M. intracellulare. A series of stock target M. avium and M. intracellulare DNA solutions were assembled by dilution in the presence of 0.02ug/uL of human DNA.

The SDA assay procedure consists of the following steps:

Add reaction mix to primer mix and target

Heat to 95°C for 2 minutes

Cool to 37°C for 5 minutes

Add enzymes and incubate at 37°C for 2 hours

Stop reaction by heating to 95°C for 3 minutes

Detect amplified product

This procedure results in the detection of as few as 10 genomes of either M. avium or M. intracellulare.

### EXAMPLE 3

The following procedure is provided to illustrate the method by which oligomers were identified for assay detection of

Mycobacterium avium and Mycobacterium intracellulare DNA.

First the sequence pMAV-29 was screened as described above in Southern blot hybridizations to determine appropriate specificity for M. avium/intracellulare. Second, conditions were developed which permit amplification of target DNA using SDA (also described above). Once amplification using SDA is established, a central region of amplified target exists for binding of assay oligomers. These oligomers must be designed to minimize competition with SDA primers. Also, the secondary structure of these oligomers must be minimized, including homo dimerization and heterodimerization. Finally, the melting temperature (T<sub>m</sub>) must be adjusted to give high specificity for the target-oligomer duplex, but not so as to reduce sensitivity of detection under the expected hybridization conditions. A number of methods are well known in the art for assessment of oligomer hybridization (secondary structure, dimerization, T<sub>m</sub>). Especially useful are software packages such as Oligo™ (National Biosciences, Plymouth, MN) or PC-Gene™ (Intelligenetics Inc., Geneva, Switzerland).

Design of oligomers for the detection of M. avium/intracellulare SDA products followed the above mentioned strategy. The internal region of the SDA product to be used for detection was 21 nucleotides long. In order to meet the needed specificity and T<sub>m</sub> for the oligomers in the assay, the detection oligomers needed to overlap with the SDA oligomers.

Competition was kept to a minimum while satisfying the requirements for hybridization. Detection oligomers were designed to be of 13-15 nucleotides in length.

Homodimerization was kept between 2-8 bases, with the ideal conditions being 6-8 bases or less. Heterodimerization was held between 2-6 bases with the ideal being 3-5 bases or less. Any secondary structure (i.e., hairpin loops), was acceptable if the calculated Gibb's Free Energy was greater than -2.0

kcal/mole. In these examples, the Gibb's Free energy was generally between -2 and 3 kcal/mole for the most stable secondary structure as determined by computational methods (Oligo or PC-Gene software). Melting temperature was dictated by oligomer concentration, %GC of the oligomer, and sodium chloride (NaCl) concentration and oligomer length. Oligomer

concentration may range between 1 - 40 nM with the ideal being 5 - 20 nM. The NaCl concentration may range between 1 - 1000 mM with the ideal being 5 - 500mM NaCl. The %GC (G+C/G+C+A+T) may range from 40 - 80% with the preferred being 45 - 70%. The above factors may be adjusted to yield a calculated melting temperature of 20 - 70°C, most preferably greater than 25°C.

EXAMPLE 4

Preparation of Coated Microtiter Plates

Biotinylated bovine serum albumin (biotin\*BSA) (Pierce, Rockford, IL) was diluted to 1 ug/well in 0.3 M Glycine pH 9.6 (BRL, Bethesda, MD., prepared using autoclaved water) and was pipetted into each well (200 ul/well) (Dynatech, Chantilly, VA), and incubated at 4°C overnight. The plates were washed twice (375 ul/wash) using FTA hemagglutination buffer (Becton Dickinson Microbiology Systems, Cockeysville, MD) pH 7.2 prepared using autoclaved water. Streptavidin (5 ug/well) in hemagglutination buffer was added to the biotin\*BSA coated microtiter wells (100 ul/well). Plates were covered and incubated for 1 hour at 37°C. Unbound streptavidin was discarded by inversion and blocking buffer (300 ul/well) (hemagglutination buffer pH 7.2, 0.05% w/v bovine serum albumin, Sigma Chemical Co., St. Louis, Mo.) was added. The plates were covered and incubated (30 min, 37°C), and the blocking buffer discarded by inversion. Plates were washed twice with hemagglutination buffer (375 ul/well), then once using hemagglutination buffer with 2% w/v trehalose (375 ul/well) (Fluka, Ronkonkoma, N.Y.). Plates were dried for approximately 4 hours under vacuum below 0.5 Torr at 25°C, sealed in mylar pouches with desiccant, and stored overnight at room temperature prior to use. The plates were stored thereafter at 2-8°C.

The assay uses 2 oligomers (probes), for detection of the SDA amplified target DNA. A biotinylated (BBB) capture probe for binding to the target DNA and the streptavidin on the microtiter plate. A second detector probe is conjugated with alkaline phosphatase (AP) and binds to the target DNA. The AP provides an opportunity for detection by means of colorimetry, fluorimetry, or chemiluminescence. This assay used the following probes for the detection of specific Mycobacterium avium and Mycobacterium intracellulare DNA amplified by SDA:

Probe set 3.

5' BBB-A ACC GGT GAC TCC A 3' (Seq ID No: 10)

5' AAA ACC TTG CGG C-AP 3' (Seq ID No: 9)

B: biotin

AP: Alkaline Phosphatase

### Microtiter Plate Assay Procedure

SDA reactions (50 ul) containing 1000, 100, 10, 1, 0 genome  
copies/10ul were prepared as follows: 35ul of SDA reaction +  
140ul of SDA dilution buffer ( 6.25mM  $\text{KH}_2\text{PO}_4$ , 12.5ug/ml BSA,  
5 0.75mM  $\text{MgCl}_2$ , 0.125mM DTT, 0.375% glycerol, 0.375% NMP pH 7.6  
+ 125ug/ml sheared salmon sperm DNA (Sigma) were added to sterile  
siliconized tubes. Diluted SDA reactions were heated to 95°C for  
2 minutes to denature the DNA. Tubes were cooled for 5 minutes  
at room temperature and then 50ul of the  
10 denatured SDA reaction was added to each well. Immediately  
thereafter, 50ul/well of hybridization mix (1M sodium  
phosphate, 0.2% BSA, 40 nM capture probe, 10nM detector probe)  
was added. The plate was covered and incubated for 45 minutes  
at 37°C. Three stringency washes (300 ul/well) (10mM sodium  
15 phosphate pH 7, 0.1% w/v bovine serum albumin, 0.05% v/v  
Nonidet P-40) were performed at room temperature. Each wash  
was allowed to remain in the microtiter wells for 1 minute  
before removing. Lumiphos™ 530 (100 ul/well, Lumigen Inc.,  
20 Detroit, MI) substrate was added, and the plates were covered  
and incubated for 30 minutes at 37°C. Luminescence was read on  
a microtiter plate luminometer (Labsystems, Research Triangle  
Park, NC) at 37°C, using a 2 second/well integration time.



# RESULTS

Using the above procedure, the assay detected 1-10 genomic copies of M. avium or M. intracellulare DNA (see Table 3 and Table 4).

5

TABLE 3

Specific Signal - Relative Light Units (RLU)

<u>M. intracellulare genome copies</u>	<u>RLU's*</u>
1	58.82
10	174.09
100	9871.9
1000	7334.26

\* Average of triplicate points; Background subtracted:  
zero=55.4 RLU

EXAMPLE 5

Using the above example procedure, probes with a greater amount of competition with the SDA primers were compared to the previously mentioned capture and detector probes. The additional probes had the following sequences: Probe set 1

5' BBB-G GGA ACC GGT GAC TC 3' (Seq ID No: 6)

5' CAA AAA CCT TGC GGC-AP 3' (Seq ID No: 7)

B: Biotin

AP: Alkaline Phosphatase

RESULTS

The results show the significance of competition on the performance of the assay. While both sets of probes, 3 and 1, had minimal detectable levels of 1-10 genomic copies of M. avium, the effect of competition was seen by the decrease in sensitivity of the assay using probe set 1 with respect to that obtained using probe set 3 (Table 4). That is, probe set 1 exhibits reduced sensitivity due to increased competition with SDA primer oligodeoxynucleotides. Increased sensitivity provides for superior quantitation of amplified target and enhanced detection limits.

TABLE 4

	<u>M. avium genome copies</u>	<u>*RLU's Probe Set 1</u>	<u>*RLU's Probe Set 3</u>
	1	199.75	371.25
	10	297.00	620.25
5	100	468.05	1337.60
	1000	3358.50	9115.60

\* Average of duplicate points; Background subtracted: zero for Set 1, 122 RLU; zero for set 3, 341.4 RLU.

SUMMARY

10       Methods are presented which teach how to design  
oligonucleotide reagents for detection of SDA amplified DNA  
from less than 10 copies of M. avium/intracellulare.

15       Although the invention has been described with respect to  
specific modifications, the details thereof are not to be  
construed as limitations, for it will be apparent that various  
equivalents, changes and modifications may be resorted to  
without departing from the spirit and scope thereof, and it is  
understood that such equivalent embodiments are to be included  
therein.

EXAMPLE 6

Genomic DNA was isolated from organisms obtained from American Type Culture Collection, Brown University, the Trudeau Microbiological Collection, Laboratories Center Disease Control (Canada) and the Raleigh Veterans Administration Hospital.

5 These specimens of DNA were then screened using aliquots of  $10^6$  copies of purified genomic DNA and performing SDA and the microwell assay as described in substantial accordance with examples 2 and 4, above. The results were calibrated using assay results for synthesized target DNA and the apparent number of genomic copies was estimated for each specimen. In the cases of M. intracellulare, M. avium and M.

10 paratuberculosis the number of copies was estimated to be above 1000. This is greater than the upper limit of the assay method. All other results were less than 17 estimated copies indicating a relative specificity (genomic copies added/genomic copies estimated) of greater than  $10^6$  (Table 5). These results indicate high specificity for M. intracellulare, M. avium and M. paratuberculosis.

TABLE 5

	DNA Tested	# Genomic Copies Added	SDA/Assay Result, Genome equiv.
	M. avium	10 <sup>6</sup>	+, > 1,000
5	M. intracellulare	10 <sup>6</sup>	+, > 1,000
	M. paratuberculosis	10 <sup>6</sup>	+, > 1,000
	M. africanum	10 <sup>6</sup>	-, <1
	M. bovis	10 <sup>6</sup>	-, <1
	M. bovis BCG	10 <sup>6</sup>	-, <1
10	M. chelonae	10 <sup>6</sup>	-, <1
	M. fortuitum	10 <sup>6</sup>	-, <1
	M. gordonae	10 <sup>6</sup>	-, <1
	M. kansasii	10 <sup>6</sup>	-, <1
	M. microti	10 <sup>6</sup>	-, <1
15	M. tuberculosis	10 <sup>6</sup>	-, <1
	M. scrofulaceum	10 <sup>6</sup>	-, <1
	M. xenopi	10 <sup>6</sup>	-, <1
	B. pertussis	10 <sup>6</sup>	-, <1
	C. diphtheriae	10 <sup>6</sup>	-, <1
20	R. rhodochrous	10 <sup>6</sup>	-, <1
	E. coli	10 <sup>6</sup>	-, 10
	K. pneumoniae	10 <sup>6</sup>	-, 6
	N. lactamica	10 <sup>6</sup>	-, 10
	P. asteroides	10 <sup>6</sup>	-, 5
25	P. aeruginosa	10 <sup>6</sup>	-, 6
	S. aureus	10 <sup>6</sup>	-, 5
	S. boydii	10 <sup>6</sup>	-, 6
	S. pneumoniae	10 <sup>6</sup>	-, 17
	S. pyogenes	10 <sup>6</sup>	-, 7

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Vonk, Glenn P.  
Nycz, Colleen M.  
Schram, James L.  
Shank, Daryl D.

(ii) TITLE OF INVENTION: Probes to Mycobacterium Avium,  
Mycobacterium Intracellurae, and Mycobacterium Paratuberculosis

(iii) NUMBER OF SEQUENCES: 15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Richard J. Rodrick  
(B) STREET: 1 Becton Drive  
(C) CITY: Franklin Lakes  
(D) STATE: New Jersey  
(E) COUNTRY: U.S.A.  
(F) ZIP: 07417

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Stierwalt, Brian K.  
(B) REGISTRATION NUMBER: 33,213  
(C) REFERENCE/DOCKET NUMBER: P-2522

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 201-847-7116  
(B) TELEFAX: 201-848-9228

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 97 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TGGCCAAACT GTGGGCGCAG GCCTGCGAGT GGGAACCGGT GACTCCAAAA	50
ACCTTGCGGC TCTTACAAGT CCGTGGCGCC AAGCTGGAGC CCGAGGA	97

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TTGAATAGTC GGTACTTGT TGACTCCTCG GGCTCCA	37
---	----

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TTGAAGTAAC CGACTATTGT TGACTGGCCA AACTGTG	37
--	----

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TTGAATAGTA GGTAAGTTGT TGACACTTGT AAGAGCC

37

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TTGAAGTAAC CGACTATTGT TGACTGCGAG TGGGAAC

37

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGGAACCGGT GACTC

15

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)



(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CAAAAACCTT GCGGC

15

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 15 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AACCGGTGAC TCCA

14

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 13 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

AAAACCTTGC GGC

13

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 14 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

AACCGGTGAC TCCA

14

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 47 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GCGAGTGGGA ACCGGTGACT CCAAAAACCT TCGGCTCTT ACAAGTC

47

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:


(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)



(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AACCGGTGAC TCCAAAAACC TTGCGGCTCT TACAAGTCGG TGGCGCCAAG 50

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 50 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GATCCCAGCC CCGAGGCGGC CTCGCCAGT CGACCGCCAT GCGGTCACGG 50

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 50 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGGAACCGGT GACTCCAAAA ACCTTGCGGC TCTTACAAGT CGGTGGCGCC 50


(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 106 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GGCCAAACTG TGGGCGCAGG CCTGCGAGTG GGAACCGGTG ACTCCAAAAA	50
CCTTGCGGGCT CTTACAAGTC GGTGGCGCCA AGCTGGAGCC CGAGGACGCC	100
CCGCCTG	106



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A probe having a sequence consisting essentially of Seq ID No: 1, Seq ID No: 2, Seq ID No: 3, Seq ID No: 4, Seq ID No: 5, Seq ID No: 6, Seq ID No: 7, Seq ID No: 8, Seq ID No: 9, Seq ID No: 10, Seq ID No: 11, Seq ID No: 12, Seq No: 13, Seq ID No: 14, or Seq ID No: 15, and the modified backbone, modified nucleotide, labelled forms, and ribonucleic acid forms thereof.
2. The probe of claim 1 which is the ribonucleic acid form thereof.
3. A method of amplifying *Mycobacterium avium*, *Mycobacterium intracellulare*, or *Mycobacterium paratuberculosis* with the use of one or more nucleic acid sequences selected from the sequences consisting essentially of Seq ID No: 1, Seq ID No: 2, Seq ID No: 3, Seq ID No: 4, Seq ID No: 5, Seq ID No: 6, Seq ID No: 7, Seq ID No: 8, Seq ID No: 9, Seq ID No: 10, Seq ID No: 11, Seq ID No: 12, Seq No: 13, Seq ID No: 14, or Seq ID No: 15, and the modified backbone, modified nucleotide, labelled forms, and ribonucleic acid forms thereof.
4. The method of claim 3 in which the nucleic acid sequence is the deoxyribonucleic acid form thereof.
5. A method of detecting *Mycobacterium avium*, *Mycobacterium intracellulare*, or *Mycobacterium paratuberculosis* which comprises the use of one or more nucleic acid sequences selected from the sequences consisting essentially of Seq ID No: 1, Seq ID No: 2, Seq ID No: 3, Seq ID No: 4, Seq ID No: 5, Seq ID No: 6, Seq ID No: 7, Seq ID No: 8, Seq ID No: 9, Seq ID No: 10, Seq ID No: 11, Seq ID No: 12, Seq No: 13, Seq ID No: 14, or Seq ID No: 15, and the modified backbone, modified nucleotide, labelled forms, and ribonucleic acid forms thereof.
6. The method of claim 5 in which the nucleic acid sequence is the deoxyribonucleic acid form thereof.
7. A method of amplifying *Mycobacterium avium*, *Mycobacterium intracellulare*, or *Mycobacterium paratuberculosis* and detecting the amplified product which comprises the use of one or more sequences selected from the sequences consisting essentially of Seq ID No: 1, Seq ID No: 2, Seq ID No: 3, Seq ID No: 4, Seq ID No: 5, Seq ID No: 6, Seq ID No: 7, Seq ID No: 8, Seq ID No: 9,

Seq ID No: 10, Seq ID No: 11, Seq ID No: 12, Seq No: 13, Seq ID No: 14, or Seq ID No: 15, and the modified backbone, modified nucleotide, labelled forms, and ribonucleic acid forms thereof.

8. The method of claim 7 in which the nucleic acid sequence is the deoxyribonucleic acid form thereof.

9. A kit comprising a nucleic acid sequence selected from the sequences consisting essentially of Seq ID No: 1, Seq ID No: 2, Seq ID No: 3, Seq ID No: 4, Seq ID No: 5, Seq ID No: 6, Seq ID No: 7, Seq ID No: 8, Seq ID No: 9, Seq ID No: 10, Seq ID No: 11, Seq ID No: 12, Seq No: 13, Seq ID No: 14, or Seq ID No: 15, and the modified backbone, modified nucleotide, labelled forms, and ribonucleic acid forms thereof.

10. The kit of claim 9 in which the nucleic acid sequence is the deoxyribonucleic acid form thereof.

DATED THIS 25 DAY OF OCTOBER 1993

BECTON DICKINSON AND COMPANY

Patent Attorneys for the

Applicant:-

F.B.RICE & CO.

# ABSTRACT

The invention provides methods and nucleic acid probes for detecting, amplifying, and isolating sequences of Mycobacterium avium, Mycobacterium intracellulare, and Mycobacterium paratuberculosis.

