(12) STANDARD PATENT (11) Application No. **AU 2016298317 B2**
(19) AUSTRALIAN PATENT OFFICE

(54) Title
Targeted oligonucleotides

(51) International Patent Classification(s)
C07H 19/173 (2006.01) ***C07H 19/19*** (2006.01)
C07H 19/09 (2006.01) ***C07H 21/00*** (2006.01)

(21) Application No: **2016298317** (22) Date of Filing: **2016.07.28**

(87) WIPO No: **WO17/019918**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/198,110	**2015.07.28**	**US**
62/305,536	**2016.03.09**	**US**
62/198,051	**2015.07.28**	**US**
62/220,652	**2015.09.18**	**US**
62/269,671	**2015.12.18**	**US**
62/239,226	**2015.10.08**	**US**

(43) Publication Date: **2017.02.02**
(44) Accepted Journal Date: **2021.02.18**

(71) Applicant(s)
Caris Science, Inc.

(72) Inventor(s)
O'Neill, Heather;Mayer, Gunter;Miglarese, Mark;Spetzler, David

(74) Agent / Attorney
Pizzeys Patent and Trade Mark Attorneys Pty Ltd, PO Box 291, WODEN, ACT, 2606, AU

(56) Related Art
JP 2013094067 A
WO 2011147193 A1
"Pseudorca crassidens DNA, BAC clone: QGB1-271M02_F, 5'end.", EMBL, (2013-12-12), Database accession no. DE820782, URL: EBI

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
2 February 2017 (02.02.2017)

WIPO | PCT

(10) International Publication Number
WO 2017/019918 A1

(51) International Patent Classification:
C07H 19/173 (2006.01) *C07H 19/19* (2006.01)
C07H 21/00 (2006.01) *C07H 19/09* (2006.01)

(21) International Application Number:
PCT/US2016/044595

(22) International Filing Date:
28 July 2016 (28.07.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

62/198,051	28 July 2015 (28.07.2015)	US
62/198,110	28 July 2015 (28.07.2015)	US
62/220,652	18 September 2015 (18.09.2015)	US
62/239,226	8 October 2015 (08.10.2015)	US
62/269,671	18 December 2015 (18.12.2015)	US
62/305,536	9 March 2016 (09.03.2016)	US

(71) Applicant: **CARIS SCIENCE, INC.** [US/US]; 6655 N. Macarthur Blvd., Irving, TX 75039 (US).

(72) Inventors: **O'NEILL, Heather**; 1252 W Laguna Azul Ave, Mesa, AZ 85202 (US). **MAYER, Gunter**; Martin-legros Strasse 59, 53123 Bonn (DE). **MIGLARESE, Mark**; 1620 East Silverwood Drive, Phoenix, AZ 85048 (US). **SPETZLER, David**; 6045 N. 40th St., Paradise Valley, AZ 85253 (US).

(74) Agent: **AKHAVAN, Ramin**; Caris Science, Inc., 6655 N. Macarthur Blvd., Irving, TX 75039 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

— *with sequence listing part of description (Rule 5.2(a))*

WO 2017/019918 A1

(54) Title: TARGETED OLIGONUCLEOTIDES

(57) Abstract: Methods and compositions are provided for oligonucleotides that bind targets of interest. The targets include cells and microvesicles, such as those derived from various diseases. The oligonucleotides can be used for diagnostic and therapeutic purposes. The target of the oligonucleotides can be a target such as PARP1, HIST1H1B, HIST1H1D, NCL, FBL, SFPQ, RPL12, ACTB, HIST1H4A, SSBP1, NONO, H2AFJ, and DDX21, or a complex, subunit or fragment thereof.

TARGETED OLIGONUCLEOTIDES

CROSS REFERENCE

This application claims the benefit of U.S. Provisional Patent Application Nos. 62/198,051, filed July 28, 2015; 62/198,110, filed July 28, 2015; 62/220,652, filed September 18, 2015; 62/239,226, filed October 8, 2015; 62/269,671, filed December 18, 2015; and 62/305,536, filed March 9, 2016; all of which applications are incorporated herein by reference in their entirety. This application is related to International Patent Application Nos. PCT/US2016/021632, filed March 9, 2016; and PCT/US2016/040157, filed June 29, 2016; both of which applications are incorporated herein by reference in their entirety.

SEQUENCE LISTING SUBMITTED VIA EFS-WEB

[0001] The entire content of the following electronic submission of the sequence listing via the USPTO EFS-WEB server, as authorized and set forth in MPEP §1730 II.B.2(a), is incorporated herein by reference in its entirety for all purposes. The sequence listing is within the electronically filed text file that is identified as follows:

[0002] File Name: 826601_SeqListing_ST25.txt

[0003] Date of Creation: July 28, 2016

[0004] Size (bytes): 897,071 bytes

BACKGROUND OF THE INVENTION

[0005] The invention relates generally to the field of aptamers capable of binding to microvesicle surface antigens, which are useful as therapeutics in and diagnostics of cancer and/or other diseases or disorders in which microvesicles implicated. The invention further relates to materials and methods for the administration of aptamers capable of binding to microvesicles. The microvesicles may be derived from cells indicative of cancer, including without limitation a breast cancer.

[0006] Aptamers are oligomeric nucleic acid molecules having specific binding affinity to molecules, which may be through interactions other than classic Watson-Crick base pairing. Unless otherwise specified, an "aptamer" as the term is used herein can refer to nucleic acid molecules that can be used to characterize a phenotype, regardless of manner of target recognition. Unless other specified, the terms "aptamer," "oligonucleotide," "polynucleotide," or the like may be used interchangeably herein.

[0007] Aptamers, like peptides generated by phage display or monoclonal antibodies ("mAbs"), are capable of specifically binding to selected targets and modulating the target's activity, e.g., through binding aptamers may block their target's ability to function. Created by an in vitro selection process from pools of random sequence oligonucleotides, aptamers have been generated for over 100 proteins including growth factors, transcription factors, enzymes, immunoglobulins, and receptors. A typical aptamer is 10-15 kDa in size (30-45 nucleotides), binds its target with sub-nanomolar affinity, and discriminates against closely related targets (e.g., aptamers will typically not bind other proteins from the same gene family). A

series of structural studies have shown that aptamers are capable of using the same types of binding interactions (e.g., hydrogen bonding, electrostatic complementarity, hydrophobic contacts, steric exclusion) that drive affinity and specificity in antibody-antigen complexes.

[0008] Aptamers have a number of desirable characteristics for use as therapeutics and diagnostics including high specificity and affinity, biological efficacy, and excellent pharmacokinetic properties. In addition, they offer specific competitive advantages over antibodies and other protein biologics, for example:

[0009] Speed and control. Aptamers are produced by an entirely in vitro process, allowing for the rapid generation of initial leads, including therapeutic leads. In vitro selection allows the specificity and affinity of the aptamer to be tightly controlled and allows the generation of leads, including leads against both toxic and non-immunogenic targets.

[0010] Toxicity and Immunogenicity. Aptamers as a class have demonstrated little or no toxicity or immunogenicity. In chronic dosing of rats or woodchucks with high levels of aptamer (10 mg/kg daily for 90 days), no toxicity is observed by any clinical, cellular, or biochemical measure. Whereas the efficacy of many monoclonal antibodies can be severely limited by immune response to antibodies themselves, it is extremely difficult to elicit antibodies to aptamers most likely because aptamers cannot be presented by T-cells via the MHC and the immune response is generally trained not to recognize nucleic acid fragments.

[0011] Administration. Whereas most currently approved antibody therapeutics are administered by intravenous infusion (typically over 2-4 hours), aptamers can be administered by subcutaneous injection (aptamer bioavailability via subcutaneous administration is >80% in monkey studies (Tucker et al., J. Chromatography B. 732: 203-212, 1999)). This difference is primarily due to the comparatively low solubility and thus large volumes necessary for most therapeutic mAbs. With good solubility (>150 mg/mL) and comparatively low molecular weight (aptamer: 10-50 kDa; antibody: 150 kDa), a weekly dose of aptamer may be delivered by injection in a volume of less than 0.5 mL. In addition, the small size of aptamers allows them to penetrate into areas of conformational constrictions that do not allow for antibodies or antibody fragments to penetrate, presenting yet another advantage of aptamer-based therapeutics or prophylaxis.

[0012] Scalability and cost. Aptamers are chemically synthesized and are readily scaled as needed to meet production demand for diagnostic or therapeutic applications. Whereas difficulties in scaling production are currently limiting the availability of some biologics and the capital cost of a large-scale protein production plant is enormous, a single large-scale oligonucleotide synthesizer can produce upwards of 100 kg/year and requires a relatively modest initial investment. The current cost of goods for aptamer synthesis at the kilogram scale is estimated at $100/g, comparable to that for highly optimized antibodies.

[0013] Stability. Aptamers are chemically robust. They are intrinsically adapted to regain activity following exposure to factors such as heat and denaturants and can be stored for extended periods (>1 yr) at room temperature as lyophilized powders.

INCORPORATION BY REFERENCE

[0014] All publications, patents and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.

SUMMARY OF THE INVENTION

[0015] Compositions and methods of the invention provide aptamers that bind biomarkers of interest. In various embodiments, oligonucleotide probes of the invention are used in diagnostic, prognostic or theranostic processes to screen a biological sample for the presence or levels of biomarkers, including without limitation surface antigens, determined to provide a relevant readout. The diagnosis may be related to a disease or disorder, e.g., a cancer. In other embodiments, oligonucleotide probes of the invention are chemically modified or comprised within a pharmaceutical composition for therapeutic or medical imaging applications.

[0016] In an aspect, the invention provides an oligonucleotide comprising a sequence selected from any one of SEQ ID NOs. 4357-4368 or 4372-4407. In a preferred embodiment, the oligonucleotide comprises a sequence according to SEQ ID NO. 4357, i.e., the sequence of aptamer 10.36. The invention further provides an oligonucleotide having a substitution in aptamer 10.36 such as in SEQ ID NOs. 4372-4407. The substitution can be chosen to such that the aptamer retains or improves upon desired such as target recognition and G quadruplex structure. In a related aspect, the invention provides an oligonucleotide comprising a sequence selected from any one of SEQ ID NOs. 4357-4368, and a 5' region with sequence 5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131), a 3' region with sequence 5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132), or both.

[0017] The oligonucleotide of the invention can be capable of binding to a target in **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61**, or a subcomponent thereof. In some embodiments, the oligonucleotide is capable of binding to Ramos cells. The oligonucleotide can be capable of binding to a protein selected from the group consisting of PARP1, HIST1H1B, HIST1H1D, NCL, FBL, SFPQ, RPL12, ACTB, HIST1H4A, SSBP1, NONO, H2AFJ, and DDX21, or a complex, subunit or fragment thereof. In some embodiment, the oligonucleotide is capable of binding to cells comprising surface nucleolin.

[0018] The invention further provides an oligonucleotide comprising a nucleic acid sequence or a portion thereof that is at least 50, 55, 60, 65, 70, 75, 80, 85, 86, 86, 88, 89, 90, 95, 96, 97, 98, 99 or 100 percent homologous to an oligonucleotide sequence described above.

[0019] In another aspect, the invention provides a plurality of oligonucleotides comprising at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000,

3000, 4000, 5000, 6000, 7000, 8000, 9000, or at least 10000 different oligonucleotide sequences described above.

[0020] The oligonucleotide or the plurality of oligonucleotides provided by the invention may comprise a DNA, RNA, 2'-O-methyl or phosphorothioate backbone, or any combination thereof. The oligonucleotide or the plurality of oligonucleotides may comprise at least one of DNA, RNA, PNA, LNA, UNA, and any combination thereof.

[0021] In some embodiments, the oligonucleotide or the plurality of oligonucleotides comprises at least one functional modification selected from the group consisting of biotinylation, a non-naturally occurring nucleotide, a deletion, an insertion, an addition, and a chemical modification. The chemical modification can be chosen to modulate desired properties such as stability, capture, detection, or binding efficiency. In some embodiments, the chemical modification comprises at least one of C18, polyethylene glycol (PEG), PEG4, PEG6, PEG8, and PEG12. The oligonucleotide or plurality of oligonucleotides can be labeled. The oligonucleotide or plurality of oligonucleotides can be attached to a nanoparticle, liposome, gold, magnetic label, fluorescent label, light emitting particle, or radioactive label. The liposome or particle can incorporate desired entities such as chemotherapeutic agents or detectable labels.

[0022] In an aspect, the invention provides an isolated oligonucleotide or plurality of oligonucleotides having a sequence as described above. In a related aspect, the invention provides a composition comprising such isolated oligonucleotide or plurality of oligonucleotides.

[0023] In some embodiments, the isolated oligonucleotide or at least one member of the plurality of oligonucleotides is capable of binding to Ramos cell. In some embodiments, the isolated oligonucleotide or plurality of oligonucleotides is capable of binding to a protein selected from the group consisting of PARP1, HIST1H1B, HIST1H1D, NCL, FBL, SFPQ, RPL12, ACTB, HIST1H4A, SSBP1, NONO, H2AFJ, and DDX21, or a complex, subunit or fragment thereof. In some embodiments, the isolated oligonucleotide or plurality of oligonucleotides is capable of binding to a cell surface nucleolin complex.

[0024] The isolated oligonucleotide or plurality of oligonucleotides can by capable of inhibiting nucleolin activity. The isolated oligonucleotide or plurality of oligonucleotides can be capable of modulating cell proliferation. In some embodiments, the isolated oligonucleotide or plurality of oligonucleotides is capable of inducing apoptosis. The cell proliferation can be neoplastic or dysplastic growth. The cell proliferation can be that of cancer cells such as disclosed herein, including without limitation that of lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer. In certain embodiments, the cell proliferation is that of leukemia, lymphoma or renal carcinoma cells.

[0025] The isolated oligonucleotide or plurality of oligonucleotides may bind to a cell surface nucleolin, a complex comprising nucleolin, or another cell surface protein from **Table 61**. Such bound outer complex may mediate cellular internalization of the complex. Such binding may also interfere with nucleolin function in the nucleus, cytoplasm, or membrane.

[0026] In an aspect, the invention provides a method comprising synthesizing the at least one oligonucleotide or the plurality of oligonucleotides provided above. Techniques for synthesizing oligonucleotides are disclosed herein or are known in the art.

[0027] In another aspect, the invention provides a method comprising contacting a biological sample with the at least one oligonucleotide, the plurality of oligonucleotides, or composition as described above. The method can further comprise detecting a presence or level of a protein in **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61** in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. In some embodiments, the method further comprises detecting a presence or level of a protein in **Table 61** in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. In still other embodiments, the method comprises detecting a presence or level of a nucleolin protein or complex thereof in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. Relatedly, the method may further comprise detecting a presence or level of a cell population in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. For example, the cells may display a protein in in **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61** on their surface. The cell population can be any desired population, including without limitation neoplastic, malignant, tumor, hyperplastic, or dysplastic cells. In some embodiments, the cell population comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells.

[0028] The detecting step of the method may comprise detecting the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. The presence or level of oligonucleotide serves as a proxy for the level of oligonucleotide's target. The oligonucleotides can be detecting using any desired technique such as described herein or known in the art, including without limitation at least one of sequencing, amplification, hybridization, gel electrophoresis, chromatography, and any combination thereof. Any useful sequencing method can be employed, including without limitation at least one of next generation sequencing, dye termination sequencing, pyrosequencing, and any combination thereof. In some embodiments, the detecting comprises transmission electron microscopy (TEM) of immunogold labeled oligonucleotides. In some embodiments, the detecting comprises confocal microscopy of fluor labeled oligonucleotides.

[0029] The detecting step of the method may comprise detecting the protein or cell using techniques described herein or known in the art for detecting proteins, including without limitation at least one of an immunoassay, enzyme immunoassay (EIA), enzyme-linked immunosorbent assay (ELISA), enzyme-linked oligonucleotide assay (ELONA), affinity isolation, immunoprecipitation, Western blot, gel electrophoresis, microscopy or flow cytometry.

[0030] In some embodiments of the method, the detected protein is associated with a microvesicle population. The method may further comprise isolating the microvesicle population prior to the contacting

with the oligonucleotides, after the contacting, or both. The isolating may be in whole or in part. For example, the microvesicle population may be partially isolated from other components in the sample before or after contacting the sample with the oligonucleotide or plurality of oligonucleotides. The invention may use any appropriate techniques to isolate microvesicles. Various techniques of isolating microvesicles are disclosed herein or known in the art, including without limitation affinity purification, filtration, concentration, polymer precipitation, PEG precipitation, ultracentrifugation, a molecular crowding reagent, affinity selection, chromatography, or any combination thereof.

[0031] Any desired biological sample can be contacted with the oligonucleotide or plurality of oligonucleotides according to the invention. In various embodiments, the biological sample comprises a bodily fluid, tissue sample or cell culture. Any desired tissue sample can be contacted. In some embodiments, the tissue sample comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer tissue. Similarly, any desired cell culture sample can be contacted. In certain embodiments, the cell culture comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells. Any appropriate bodily fluid can be contacted, including without limitation peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair oil, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, or umbilical cord blood. In certain preferred embodiments, the bodily fluid comprises whole blood or a derivative or fraction thereof, such as sera or plasma. The bodily fluid may comprise cancer cells, including without limitation lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells.

[0032] The biological sample may be spiked with a purified or recombinant protein. In some embodiments, such protein is selected from **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61**, or complexes, subunits or fragments thereof.

[0033] As desired, the method of detecting the presence or level of the at least one oligonucleotide, the plurality of oligonucleotides, or composition bound to a target can be used to characterize a phenotype. The phenotype can be any appropriate phenotype, including without limitation a disease or disorder. In such cases, the characterizing may include providing, or assisting in providing, at least one of diagnostic, prognostic and theranostic information for the disease or disorder. Characterizing the phenotype may comprise comparing the presence or level to a reference. Any appropriate reference level can be used. For

example, the reference can be the presence or level determined in a sample from at least one individual without the phenotype or from at least one individual with a different phenotype. As a further example, if the phenotype is a disease or disorder, the reference level may be the presence or level determined in a sample from at least one individual without the disease or disorder, or with a different state of the disease or disorder (*e.g.*, in remission, different stage or grade, different prognosis, metastatic versus local, etc).

[0034] As noted, the sample can be from a subject suspected of having or being predisposed to a disease or disorder. The disease or disorder can be any disease or disorder that can be assessed by the subject method. For example, the disease or disorder may be a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. In an embodiment, the cancer comprises an acute lymphoblastic leukemia; acute myeloid leukemia; adrenocortical carcinoma; AIDS-related cancers; AIDS-related lymphoma; anal cancer; appendix cancer; astrocytomas; atypical teratoid/rhabdoid tumor; basal cell carcinoma; bladder cancer; brain stem glioma; brain tumor (including brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, astrocytomas, craniopharyngioma, ependymoblastoma, ependymoma, medulloblastoma, medulloepithelioma, pineal parenchymal tumors of intermediate differentiation, supratentorial primitive neuroectodermal tumors and pineoblastoma); breast cancer; bronchial tumors; Burkitt lymphoma; cancer of unknown primary site; carcinoid tumor; carcinoma of unknown primary site; central nervous system atypical teratoid/rhabdoid tumor; central nervous system embryonal tumors; cervical cancer; childhood cancers; chordoma; chronic lymphocytic leukemia; chronic myelogenous leukemia; chronic myeloproliferative disorders; colon cancer; colorectal cancer; craniopharyngioma; cutaneous T-cell lymphoma; endocrine pancreas islet cell tumors; endometrial cancer; ependymoblastoma; ependymoma; esophageal cancer; esthesioneuroblastoma; Ewing sarcoma; extracranial germ cell tumor; extragonadal germ cell tumor; extrahepatic bile duct cancer; gallbladder cancer; gastric (stomach) cancer; gastrointestinal carcinoid tumor; gastrointestinal stromal cell tumor; gastrointestinal stromal tumor (GIST); gestational trophoblastic tumor; glioma; hairy cell leukemia; head and neck cancer; heart cancer; Hodgkin lymphoma; hypopharyngeal cancer; intraocular melanoma; islet cell tumors; Kaposi sarcoma; kidney cancer; Langerhans cell histiocytosis; laryngeal cancer; lip cancer; liver cancer; lung cancer; malignant fibrous histiocytoma bone cancer; medulloblastoma; medulloepithelioma; melanoma; Merkel cell carcinoma; Merkel cell skin carcinoma; mesothelioma; metastatic squamous neck cancer with occult primary; mouth cancer; multiple endocrine neoplasia syndromes; multiple myeloma; multiple myeloma/plasma cell neoplasm; mycosis fungoides; myelodysplastic syndromes; myeloproliferative neoplasms; nasal cavity cancer; nasopharyngeal cancer; neuroblastoma; Non-Hodgkin lymphoma; nonmelanoma skin cancer; non-small cell lung cancer; oral cancer; oral cavity cancer; oropharyngeal cancer; osteosarcoma; other brain and spinal cord tumors; ovarian cancer; ovarian epithelial cancer; ovarian germ cell tumor; ovarian low malignant potential tumor; pancreatic cancer; papillomatosis; paranasal sinus cancer; parathyroid cancer; pelvic cancer; penile cancer; pharyngeal cancer; pineal parenchymal tumors of intermediate

differentiation; pineoblastoma; pituitary tumor; plasma cell neoplasm/multiple myeloma; pleuropulmonary blastoma; primary central nervous system (CNS) lymphoma; primary hepatocellular liver cancer; prostate cancer; rectal cancer; renal cancer; renal cell (kidney) cancer; renal cell cancer; respiratory tract cancer; retinoblastoma; rhabdomyosarcoma; salivary gland cancer; Sézary syndrome; small cell lung cancer; small intestine cancer; soft tissue sarcoma; squamous cell carcinoma; squamous neck cancer; stomach (gastric) cancer; supratentorial primitive neuroectodermal tumors; T-cell lymphoma; testicular cancer; throat cancer; thymic carcinoma; thymoma; thyroid cancer; transitional cell cancer; transitional cell cancer of the renal pelvis and ureter; trophoblastic tumor; ureter cancer; urethral cancer; uterine cancer; uterine sarcoma; vaginal cancer; vulvar cancer; Waldenström macroglobulinemia; or Wilm's tumor. The premalignant condition can be Barrett's Esophagus. The autoimmune disease can be inflammatory bowel disease (IBD), Crohn's disease (CD), ulcerative colitis (UC), pelvic inflammation, vasculitis, psoriasis, diabetes, autoimmune hepatitis, multiple sclerosis, myasthenia gravis, Type I diabetes, rheumatoid arthritis, psoriasis, systemic lupus erythematosis (SLE), Hashimoto's Thyroiditis, Grave's disease, Ankylosing Spondylitis Sjogrens Disease, CREST syndrome, Scleroderma, Rheumatic Disease, organ rejection, Primary Sclerosing Cholangitis, or sepsis. The cardiovascular disease can be atherosclerosis, congestive heart failure, vulnerable plaque, stroke, ischemia, high blood pressure, stenosis, vessel occlusion or a thrombotic event. The neurological disease can be Multiple Sclerosis (MS), Parkinson's Disease (PD), Alzheimer's Disease (AD), schizophrenia, bipolar disorder, depression, autism, Prion Disease, Pick's disease, dementia, Huntington disease (HD), Down syndrome, cerebrovascular disease, Rasmussen's encephalitis, viral meningitis, neurospsychiatric systemic lupus erythematosus (NPSLE), amyotrophic lateral sclerosis, Creutzfeldt-Jacob disease, Gerstmann-Straussler-Scheinker disease, transmissible spongiform encephalopathy, ischemic reperfusion damage (e.g. stroke), brain trauma, microbial infection, or chronic fatigue syndrome. The pain can be fibromyalgia, chronic neuropathic pain, or peripheral neuropathic pain. The infectious disease can be a bacterial infection, viral infection, yeast infection, Whipple's Disease, Prion Disease, cirrhosis, methicillin-resistant staphylococcus aureus, HIV, Hepatitis C virus (HCV), Epstein Barr virus, Helicobacter pylori, hepatitis, syphilis, meningitis, malaria, tuberculosis, or influenza.

[0035] As further described herein, the invention provides a kit comprising a reagent for carrying out the method. Similarly, the invention provides for the use of a reagent for carrying out the method. The reagent can be any useful reagent for carrying out the method. For example, the reagent can be the at least one oligonucleotide or the plurality of oligonucleotides, one or more primer for amplification or sequencing of such oligonucleotides, at least one binding agent to at least one protein, a binding buffer with or without $MgCl_2$, a sample processing reagent, a microvesicle isolation reagent, a cell isolation reagent, a detection reagent, a secondary detection reagent, a wash buffer, an elution buffer, a solid support, and any combination thereof. The microvesicle isolation reagent may comprise at least one of a concentrator unit, a filtration unit, a polymer, PEG, a size exclusion column, a binding agent to a microvesicle antigen, and any combination thereof; and/or the detection or secondary detection agent comprises streptavidin-horse

radish peroxide (HRP), a streptavidin-conjugated fluorophore, a streptavidin-conjugated quantum dot, and any combination thereof.

[0036] In an aspect, the invention provides a method of imaging a cell or tissue, comprising contacting the cell or tissue with at least one oligonucleotide or plurality of oligonucleotides as described above, and detecting the at least one oligonucleotide or the plurality of oligonucleotides in contact with at least one cell or tissue. In some embodiments, the at least one oligonucleotide or the plurality of oligonucleotides is labeled, e.g., in order to facilitate detection or medical imaging. The oligonucleotide or plurality of oligonucleotides can be attached to a nanoparticle, liposome, gold, magnetic label, fluorescent label, light emitting particle, radioactive label, or other useful label such as disclosed herein or known in the art. The oligonucleotides can be administered to a subject prior to the detecting. The at least one cell or tissue can comprise cells displaying nucleolin or another protein from **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61** on their surface. In some embodiments, the at least one cell or tissue comprises neoplastic, malignant, tumor, hyperplastic, or dysplastic cells. For example, the at least one cell or tissue may comprise lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells.

[0037] In an aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of the at least one oligonucleotide or the plurality of oligonucleotides of the invention, or a salt thereof, and a pharmaceutically acceptable carrier, diluent, or both. In some embodiments, the oligonucleotides are attached to a toxin or chemotherapeutic agent. In some embodiments, the oligonucleotides are attached to a liposome or nanoparticle. The liposome or nanoparticle may comprise a toxin or chemotherapeutic agent. In such embodiments, the at least one oligonucleotide or the plurality of oligonucleotides can be used for targeted delivery of the toxin, chemotherapeutic agent, liposome or nanoparticle to a desired target cell or tissue.

[0038] In a related aspect, the invention provides a method of treating or ameliorating a disease or disorder in a subject in need thereof, comprising administering such pharmaceutical composition to the subject. In another related aspect, the invention provides a method of inducing cytotoxicity in a subject, comprising administering such pharmaceutical to the subject. The pharmaceutical composition can be administered in any useful format. In various embodiments, the administering comprises at least one of intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, by inhalation, topical administration, or any combination thereof. The carrier or diluent can be any useful carrier or diluent, as described herein or known in the art. As desired, the pharmaceutical composition can be administered in combination with additional known chemotherapeutic agents.

[0039] In an aspect, the invention provides a multipartite (chimeric) construct that comprises a first segment that binds to a first target and a second segment that binds to a second target. As desired, the first segment, the second segment, or both, comprises SEQ ID NO. 4357, or a region that is at least 50, 55, 60,

65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99 or 100 percent homologous thereto. For example, the first segment can be according to any of SEQ ID NOs. 4372-4407.

[0040] In an embodiment, the first target comprises a protein selected from any of **Table 50**, **51**, **52**, **60** or **61**. For example, the first target can be a protein selected from the group consisting of PARP1, HIST1H1B, HIST1H1D, NCL, FBL, SFPQ, RPL12, ACTB, HIST1H4A, SSBP1, NONO, H2AFJ, and DDX21, or a complex, subunit or fragment thereof. In some embodiments, the first target comprises nucleolin or a complex thereof. The first segment may be capable of binding to Ramos cells.

[0041] As described herein, the invention contemplates various configuration of the multipartite construct. *See, e.g.,* **FIGs. 23A-D** herein and related discussion herein. The construct may further comprise a first oligonucleotide primer region and/or a second oligonucleotide primer region surrounding the first segment.

[0042] In some embodiments, the second target of the multipartite construct of the invention comprises an immunomodulatory molecule. For example, the immunomodulatory molecule may be selected from at least one of a member of the innate immune system, a member of the complement system, C1q, C1r, C1s, C1, C3a, C3b, C3d, C5a, C2, C4, and any combination thereof. In some embodiments, the second target comprises C1q or a subunit thereof, e.g., the A, B or C subunit. The second target may comprise any number of post-translational modifications. For example, when the immunomodulatory molecule comprises C1q A, the A subunit may have at least one modification selected from **Table 46**.

[0043] As noted above, the invention contemplates various configuration of the multipartite construct. In various embodiments of the invention, the second segment comprises an antibody or oligonucleotide. And in preferred embodiments, the second segment comprises an oligonucleotide. Such a second segment may further comprise a first oligonucleotide primer region and/or a second oligonucleotide primer region surrounding the second segment. The second segment may comprise an oligonucleotide as provided by the invention. For example, the oligonucleotide can be as described above, including without limitation a sequence selected from any one of SEQ ID NOs. 137-969 and 1072-4325, or a region that is at least 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99 or 100 percent homologous thereto.

[0044] As desired, the multipartite construct of the invention can include a linker region between the first segment and second segment. The linker region can be chosen to achieve any desired purpose, such as modulate the distance between the first and second targets, e.g., to relieve steric hindrance or to bring the targets into close proximity. The linker can also provide certain functionalities, including without limitation an immunostimulatory sequence, an anti-proliferative sequence, a pro-apoptotic sequence, or some combination thereof. In an embodiment, the linker region comprises one or more CpG motif. In another embodiment, the linker comprises a polyG sequence. In still other emobdiments, the multipartite construct of the invention comprises an immunostimulating moiety, a membrane disruptive moiety, or both.

[0045] The multipartite construct of the invention can be modified to comprise at least one oligonucleotide chemical modification. Various useful modifications are disclosed herein or known in the

art. In some embodiments, the modification is selected from the group consisting: of a chemical substitution at a sugar position; a chemical substitution at a phosphate position; a chemical substitution at a base position of the nucleic acid; and any combination thereof. The modification can be selected from the group consisting of: incorporation of a modified nucleotide, 3' capping, conjugation to an amine linker, conjugation to a high molecular weight, non-immunogenic compound, conjugation to a lipophilic compound, conjugation to a drug, conjugation to a cytotoxic moiety, labeling with a radioisotope, and any combination thereof. In a preferred embodiment, the non-immunogenic, high molecular weight compound is polyalkylene glycol, such as polyethylene glycol.

[0046] In preferred embodiments, the multipartite construct comprises an oligonucleotide polymer. Such a construct can be flanked by a first oligonucleotide primer region and a second oligonucleotide primer region. In various embodiments, the second segment comprises an oligonucleotide provided by the invention, e.g., an oligonucleotide that can bind C1q or a subunit thereof. For example, the multipartite construct may comprise an oligonucleotide having a sequence according to any one of SEQ ID NO. 4358-4368, or that is at least 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99 or 100 percent homologous thereto.

[0047] In a related aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of the multipartite construct described herein, or a salt thereof, and a pharmaceutically acceptable carrier, diluent or both. In a related aspect, the invention provides a method of treating or ameliorating a disease or disorder in a subject in need thereof, comprising administering such pharmaceutical composition to the subject. The pharmaceutical composition can be administered in any useful format. In various embodiments, the administering comprises at least one of intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, by inhalation, topical administration, or any combination thereof. The carrier or diluent can be any useful carrier or diluent, as described herein or known in the art.

[0048] In another related aspect, the invention provides a method of inducing killing of a cell, comprising contacting the cell with a multipartite construct described herein. In some embodiments, the cell that is killed comprises a disease or disorder.

[0049] As described above, the multipartite construct of the invention can be administered to a subject to treat a disease or disorder, or to induce cell killing. In various embodiments, the disease or disorder comprises a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. The cancer can include without limitation one of acute lymphoblastic leukemia; acute myeloid leukemia; adrenocortical carcinoma; AIDS-related cancers; AIDS-related lymphoma; anal cancer; appendix cancer; astrocytomas; atypical teratoid/rhabdoid tumor; basal cell carcinoma; bladder cancer; brain stem glioma; brain tumor (including brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, astrocytomas, craniopharyngioma, ependymoblastoma, ependymoma, medulloblastoma, medulloepithelioma, pineal parenchymal tumors of

intermediate differentiation, supratentorial primitive neuroectodermal tumors and pineoblastoma); breast cancer; bronchial tumors; Burkitt lymphoma; cancer of unknown primary site; carcinoid tumor; carcinoma of unknown primary site; central nervous system atypical teratoid/rhabdoid tumor; central nervous system embryonal tumors; cervical cancer; childhood cancers; chordoma; chronic lymphocytic leukemia; chronic myelogenous leukemia; chronic myeloproliferative disorders; colon cancer; colorectal cancer; craniopharyngioma; cutaneous T-cell lymphoma; endocrine pancreas islet cell tumors; endometrial cancer; ependymoblastoma; ependymoma; esophageal cancer; esthesioneuroblastoma; Ewing sarcoma; extracranial germ cell tumor; extragonadal germ cell tumor; extrahepatic bile duct cancer; gallbladder cancer; gastric (stomach) cancer; gastrointestinal carcinoid tumor; gastrointestinal stromal cell tumor; gastrointestinal stromal tumor (GIST); gestational trophoblastic tumor; glioma; hairy cell leukemia; head and neck cancer; heart cancer; Hodgkin lymphoma; hypopharyngeal cancer; intraocular melanoma; islet cell tumors; Kaposi sarcoma; kidney cancer; Langerhans cell histiocytosis; laryngeal cancer; lip cancer; liver cancer; lung cancer; malignant fibrous histiocytoma bone cancer; medulloblastoma; medulloepithelioma; melanoma; Merkel cell carcinoma; Merkel cell skin carcinoma; mesothelioma; metastatic squamous neck cancer with occult primary; mouth cancer; multiple endocrine neoplasia syndromes; multiple myeloma; multiple myeloma/plasma cell neoplasm; mycosis fungoides; myelodysplastic syndromes; myeloproliferative neoplasms; nasal cavity cancer; nasopharyngeal cancer; neuroblastoma; Non-Hodgkin lymphoma; nonmelanoma skin cancer; non-small cell lung cancer; oral cancer; oral cavity cancer; oropharyngeal cancer; osteosarcoma; other brain and spinal cord tumors; ovarian cancer; ovarian epithelial cancer; ovarian germ cell tumor; ovarian low malignant potential tumor; pancreatic cancer; papillomatosis; paranasal sinus cancer; parathyroid cancer; pelvic cancer; penile cancer; pharyngeal cancer; pineal parenchymal tumors of intermediate differentiation; pineoblastoma; pituitary tumor; plasma cell neoplasm/multiple myeloma; pleuropulmonary blastoma; primary central nervous system (CNS) lymphoma; primary hepatocellular liver cancer; prostate cancer; rectal cancer; renal cancer; renal cell (kidney) cancer; renal cell cancer; respiratory tract cancer; retinoblastoma; rhabdomyosarcoma; salivary gland cancer; Sézary syndrome; small cell lung cancer; small intestine cancer; soft tissue sarcoma; squamous cell carcinoma; squamous neck cancer; stomach (gastric) cancer; supratentorial primitive neuroectodermal tumors; T-cell lymphoma; testicular cancer; throat cancer; thymic carcinoma; thymoma; thyroid cancer; transitional cell cancer; transitional cell cancer of the renal pelvis and ureter; trophoblastic tumor; ureter cancer; urethral cancer; uterine cancer; uterine sarcoma; vaginal cancer; vulvar cancer; Waldenström macroglobulinemia; or Wilm's tumor. The premalignant condition can include without limitation Barrett's Esophagus. The autoimmune disease can include without limitation one of inflammatory bowel disease (IBD), Crohn's disease (CD), ulcerative colitis (UC), pelvic inflammation, vasculitis, psoriasis, diabetes, autoimmune hepatitis, multiple sclerosis, myasthenia gravis, Type I diabetes, rheumatoid arthritis, psoriasis, systemic lupus erythematosis (SLE), Hashimoto's Thyroiditis, Grave's disease, Ankylosing Spondylitis Sjogrens Disease, CREST syndrome, Scleroderma, Rheumatic Disease, organ rejection, Primary Sclerosing Cholangitis, or sepsis. The

cardiovascular disease can include without limitation one of atherosclerosis, congestive heart failure, vulnerable plaque, stroke, ischemia, high blood pressure, stenosis, vessel occlusion or a thrombotic event. The neurological disease can include without limitation one of Multiple Sclerosis (MS), Parkinson's Disease (PD), Alzheimer's Disease (AD), schizophrenia, bipolar disorder, depression, autism, Prion Disease, Pick's disease, dementia, Huntington disease (HD), Down's syndrome, cerebrovascular disease, Rasmussen's encephalitis, viral meningitis, neurospsychiatric systemic lupus erythematosus (NPSLE), amyotrophic lateral sclerosis, Creutzfeldt-Jacob disease, Gerstmann-Straussler-Scheinker disease, transmissible spongiform encephalopathy, ischemic reperfusion damage (e.g. stroke), brain trauma, microbial infection, or chronic fatigue syndrome. The pain can include without limitation one of fibromyalgia, chronic neuropathic pain, or peripheral neuropathic pain. The infectious disease can include without limitation one of a bacterial infection, viral infection, yeast infection, Whipple's Disease, Prion Disease, cirrhosis, methicillin-resistant staphylococcus aureus, HIV, Hepatitis C virus (HCV), Epstein Barr virus, Helicobacter pylori, hepatitis, syphilis, meningitis, malaria, tuberculosis, or influenza.

[0050] In an aspect, the invention provides a kit comprising a multipartite construct or a pharmaceutical composition provided by the invention, e.g., as described above. In a related aspect, the invention provides a kit comprising a reagent for carrying out the methods making use of such multipartite construct or a pharmaceutical composition. Similarly, the invention provides use of a reagent for carrying out the methods making use of such multipartite construct or a pharmaceutical composition. The invention also provides use of a reagent for the manufacture of a kit or reagent for carrying out the methods making use of such multipartite construct or a pharmaceutical composition. The invention further contemplates use of a reagent for the manufacture of a medicament for carrying out the method methods making use of such multipartite construct or a pharmaceutical composition. In such kits or uses, the reagent may comprise a multipartite construct or a pharmaceutical composition provided by the invention, e.g., as described above.

BRIEF DESCRIPTION OF THE DRAWINGS

[0051] **FIG. 1** illustrates a competitive assay selection strategy: the random pool of aptamer (the library) is incubated with the target protein, in this case, EpCAM. After washing and elution from the target, the eluted aptamers are again added to the target and allowed to bind. The antibody is then added to the reaction, competing with the aptamers at the epitope of the antibody. The aptamers displaced by the antibody are then collected.

[0052] **FIGs. 2A-2F** illustrate methods of assessing biomarkers such as microvesicle surface antigens. **FIG. 2A** is a schematic of a planar substrate coated with a capture agent, such as an aptamer or antibody, which captures vesicles expressing the target antigen of the capture agent. The capture agent may bind a protein expressed on the surface of vesicles shed from diseased cells ("disease vesicle"). The detection agent, which may also be an aptamer or antibody, carries a detectable label, here a fluorescent signal. The detection agent binds to the captured vesicle and provides a detectable signal via its fluorescent label. The detection agent can detect an antigen that is generally associated with vesicles, or is associated with a cell-

of-origin or a disease, e.g., a cancer. **FIG. 2B** is a schematic of a particle bead conjugated with a capture agent, which captures vesicles expressing the target antigen of the capture agent. The capture agent may bind a protein expressed on the surface of vesicles shed from diseased cells ("disease vesicle"). The detection agent, which may also be an aptamer or antibody, carries a detectable label, here a fluorescent signal. The detection agent binds to the captured vesicle and provides a detectable signal via its fluorescent label. The detection agent can detect an antigen that is generally associated with vesicles, or is associated with a cell-of-origin or a disease, e.g., a cancer. **FIG. 2C** is an example of a screening scheme that can be performed by using different combinations of capture and detection agents to the indicated biomarkers. The biomarker combinations can be detected using assays as shown in **FIGs. 2A-2B**. **FIGs. 2D-2E** present illustrative schemes for capturing and detecting vesicles to characterize a phenotype. **FIG. 2F** presents illustrative schemes for assessing vesicle payload to characterize a phenotype.

[0053] **FIGs. 3A-B** illustrates a non-limiting example of an aptamer nucleotide sequence and its secondary structure. **FIG. 3A** illustrates a secondary structure of a 32-mer oligonucleotide, Aptamer 4, with sequence 5'-CCCCCCGAATCACATGACTTGGGCGGGGGTCG (SEQ ID NO. 1). In the figure, the sequence is shown with 6 thymine nucleotides added to the end, which can act as a spacer to attach a biotin molecule. This particular oligo has a high binding affinity to the target, EpCAM (*see* **Table 5**). Additional candidate EpCAM binders are identified by modeling the entire database of sequenced oligos to the secondary structure of this oligo. **FIG. 3B** illustrates another 32-mer oligo with sequence 5'-ACCGGATAGCGGTTGGAGGCGTGCTCCACTCG (SEQ ID NO. 2) that has a different secondary structure than the aptamer in **FIG. 3A**. This aptamer is also shown with a 6-thymine tail.

[0054] **FIG. 4** illustrates a process for producing a target-specific set of aptamers using a cell subtraction method, wherein the target is a biomarker associated with a specific disease. In Step 1, a random pool of oligonucleotides are contacted with a biological sample from a normal patient. In Step 2, the oligos that did not bind in Step 1 are added to a biological sample isolated from diseased patients. The bound oligos from this step are then eluted, captured via their biotin linkage and then combined again with normal biological sample. The unbound oligos are then added again to disease-derived biological sample and isolated. This process can be repeated iteratively. The final eluted aptamers are tested against patient samples to measure the sensitivity and specificity of the set. Biological samples can include blood, including plasma or serum, or other components of the circulatory system, such as microvesicles.

[0055] **FIG. 5** illustrates results from a binding assay showing the binding affinity of an exemplary aptamer (Aptamer ID BTX176881 (SEQ ID NO: 3)) to the target EpCAM protein at various target concentrations. The aptamer to be tested is fixed to a substrate using a biotin tail and is incubated with various concentrations of target (125, 250 and 500 nM). The test is performed on a surface plasmon resonance machine (SPR). The SPR machine detects association and disassociation of the aptamer and the target. Target is applied until the association and disassociation events are equal, resulting in a plateau of the curve. The equations describing the curve at each concentration can then be used to calculate the K_D of the aptamer (*see* **Table 5**).

[0056] **FIGs. 6A-D** illustrate the use of an anti-EpCAM aptamer (Aptamer 4; SEQ ID NO. 1) to detect a microvesicle population. Vesicles in patient plasma samples were captured using bead-conjugated antibodies to the indicated microvesicle surface antigens: A) EGFR; B) PBP; C) EpCAM; D) KLK2. Fluorescently labeled Aptamer 4 was used as a detector in the microbead assay. The figure shows average median fluorescence values (MFI values) for three cancer (C1-C3) and three normal samples (N1-N3) in each plot. In each plot, the samples from left to right are ordered as: C1, C2, C3, N1, N2, N3.

[0057] **FIG. 7A** illustrates the sequence of EPCAM aptamer CAR003 (SEQ ID NO. 4). **FIG. 7B** illustrates the optimal secondary structure of CAR003 with a minimum free energy (ΔG) of -30.00 kcal/mol. For purposes of illustration, the aptamer is shown as an RNA aptamer (SEQ ID NO. 5) corresponding to the DNA sequence in **FIG. 7A**. **FIG. 7C** illustrates aptamer pool purification. The figure comprises an FPLC chromatogram with all product and fractions assigned in pools after checking quality on gel. **FIG. 7D** illustrates a SYBR GOLD stained gel with different FPLC fractions of CAR003 aptamer after synthesis. Different fractions were combined in pools based on amount of un-finished chains in order high to low (pool 1 – pool 3). The pools 1-3 correspond to those indicated in **FIG. 7C**. **FIG. 7E-F** illustrate binding of CAR003 to EPCAM protein in 25 mM HEPES with PBS-BN (**FIG. 7E**) or in 25 mM HEPES with 1 mM $MgCl_2$ (**FIG. 7F**). **FIG. 7G** illustrates CAR003 binding to EpCAM in the indicated salts with and without addition of bovine serum albumin (BSA). **FIG. 7H** illustrates the effect of denaturing on CAR003 binding to EPCAM protein. In each group of four bars, the aptamer is from left to right: Aptamer 4, CAR003 Pool 1, CAR003 Pool 2, CAR003 Pool 3. **FIG. 7I** illustrates titration of aptamers against EPCAM recombinant protein (constant input 5 μg). **FIG. 7J** illustrates a Western blot with CAR003 aptamer versus EPCAM his-tagged protein, BSA, and HSA (5 μg each). The gel was blocked 0.5% F127 and probed with ~ 50 μg/ml CAR003 biotinylated aptamer, fraction 3. The blot was visualized with NeutrAvidin-HRP followed by SuperSignal West Femto Chemiluminescent Substrate.

[0058] **FIGs. 8A-8D** illustrates methods to attach microvesicles to a substrate. **FIG. 8A** illustrates direct conjugation of a carboxylated microsphere to a vesicle surface antigen. **FIG. 8B** illustrates anchoring of a microvesicle to a microsphere via a biotin functionalized lipid anchor. **FIG. 8C** illustrates antibody binding to a vesicle surface antigen, wherein the antibody is conjugated to a carboxylated microsphere. **FIG. 8D** illustrates aptamer binding to a vesicle surface antigen, wherein the aptamer is conjugated to a carboxylated microsphere.

[0059] **FIG. 9** comprises a schematic for identifying a target of a selected aptamer, such as an aptamer selected by the process of the invention. The figure shows a binding agent **902**, here an aptamer for purposes of illustration, tethered to a substrate **901**. The binding agent **902** can be covalently attached to substrate **901**. The binding agent **902** may also be non-covalently attached. For example, binding agent **902** can comprise a label which can be attracted to the substrate, such as a biotin group which can form a complex with an avidin/streptavidin molecule that is covalently attached to the substrate. The binding agent **902** binds to a surface antigen **903** of microvesicle **904**. In the step signified by arrow (i), the microvesicle is disrupted while leaving the complex between the binding agent **902** and surface antigen

903 intact. Disrupted microvesicle **905** is removed, e.g., via washing or buffer exchange, in the step signified by arrow (ii). In the step signified by arrow (iii), the surface antigen **903** is released from the binding agent **902**. The surface antigen **903** can be analyzed to determine its identity.

[0060] **FIGs. 10A-10C** illustrate binding of selected aptamers against microbeads conjugated to various input sample. The aptamers were selected from an aptamer library as binding to microbeads conjugated to breast cancer-derived microvesicles. Experimental details are in the Examples herein. Each plot shows a different aptamer. The Y-axis indicates level of binding. In each group of samples, binding of 9 purified aptamer candidates is shown. The input sample is indicated on the X axis from left to right as follows: 1) Cancer Exosome: aptamer binding to microbeads conjugated to microvesicles isolated from plasma samples from breast cancer patients; 2) Cancer Non-exosome: aptamer binding to microbeads conjugated to plasma samples from breast cancer patients after removal of microvesicles by ultracentrifugation; 3) Non-Cancer Exosome: aptamer binding to microbeads conjugated to microvesicles isolated from plasma samples from normal (i.e., non-breast cancer) patients; 4) Non-Cancer Non-Exosome: aptamer binding to microbeads conjugated to plasma samples from breast cancer patients after removal of microvesicles by ultracentrifugation.

[0061] **FIGs. 11A-11B** illustrate enriching a naïve aptamer library for aptamers that differentiate between breast cancer and non-cancer microvesicles in plasma samples.

[0062] **FIGs. 12A-12G** illustrate using an oligonucleotide probe library to differentiate cancer and non-cancer samples.

[0063] **FIG. 13** shows protein targets of oligonucleotide probes run on a silver stained SDS-PAGE gel.

[0064] **FIGs. 14A-G** illustrate use of oligonucleotides that differentiate microvesicles in breast cancer plasma from normal controls.

[0065] **FIG. 15** shows a heatmap of clusters of oligonucleotides enriched against aggressive versus non-aggressive breast cancer plams samples.

[0066] **FIGs. 16A-B** illustrate a model generated using a training (**FIG. 16A**) and test (**FIG. 16B**) set from a round of cross validation. The AUC for the test set was 0.803. Another exemplary round of cross-validation is shown in **FIGs. 16C-D** with training (**FIG. 16C**) and test (**FIG. 16D**) sets. The AUC for the test set was 0.678.

[0067] **FIGs. 17A-E** show a photo-cleavable Biotin mediated purification of an oligonucleotide library.

[0068] **FIGs. 18A-C** illustrate SUPRA (SsDNA by Unequal length PRimer Asymmetric PCR), a protocol for single stranded DNA (ssDNA) oligonucleotide library preparation.

[0069] **FIGs. 19A-C** illustrate use of aptamers in methods of characterizing a phenotype. **FIG. 19A** is a schematic **1900** showing an assay configuration that can be used to detect and/or quantify a target of interest. In the figure, capture aptamer **1902** is attached to substrate **1901**. Target of interest **1903** is bound by capture aptamer **1902**. Detection aptamer **1904** is also bound to target of interest **1903**. Detection aptamer **1904** carries label **1905** which can be detected to identify target captured to substrate **1901** via capture aptamer **1902**. **FIG. 19B** is a schematic **1910** showing use of an aptamer pool to characterize a

phenotype. A pool of aptamers to a target of interest is provided **1911**. The pool is contacted with a test sample to be characterized **1912**. The mixture is washed to remove unbound aptamers. The remaining aptamers are disassociated and collected **1913**. The collected aptamers are identified **1914** and the identity of the retained aptamers is used to characterize the phenotype **1915**. **FIG. 19C** is a schematic **1920** showing an implementation of the method in **FIG. 19B**. A pool of aptamers identified as binding a microvesicle population is provided **1919**. The input sample comprises microvesicles that are isolated from a test sample **1920**. The pool is contacted with the isolated microvesicles to be characterized **1923**. The mixture is washed to remove unbound aptamers and the remaining aptamers are disassociated and collected **1925**. The collected aptamers are identified and the identity of the retained aptamers is used to characterize the phenotype **1926**.

[0070] **FIGs. 20A-I** illustrate development and use of an oligonucleotide probe library to distinguish biological sample types.

[0071] **FIGs. 21A-C** illustrate enriching a naïve oligonucleotide library with balanced design for oligonucleotides that differentiate between breast cancer and non-cancer microvesicles derived from plasma samples.

[0072] **FIGs. 22A-D** shows characterization of breast cancer samples as cancer or non-cancer using two different but related oligonucleotide probe libraries.

[0073] **FIGs. 23A-E** illustrate oligonucleotide constructs that recognize immunomodulatory (IMD) targets.

[0074] **FIGs. 24A-C** illustrate identification of oligonucleotides that recognize C1q and other targets.

[0075] **FIGs. 25A-G** illustrate identification of targets of aptamer 10.36 (SEQ ID NO. 4357).

[0076] **FIGs. 26A-E** illustrate cell killing by aptamer 10.36 (SEQ ID NO. 4357).

DETAILED DESCRIPTION OF THE INVENTION

[0077] The details of one or more embodiments of the invention are set forth in the accompanying description below. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. Other features, objects, and advantages of the invention will be apparent from the description. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the case of conflict, the present Specification will control.

[0078] Disclosed herein are compositions and methods that can be used to assess a biomarker profile, which can include a presence or level of one or more biomarkers. The compositions and methods of the invention comprise the use of oligonucleotide probes (aptamers) that bind microvesicle surface antigens or a functional fragment thereof. The antigens typically comprise proteins or polypeptides but can be any useful component displayed on a microvesicle surface including nucleic acids, lipids and/or

carbohydrates. In general, the oligonucleotides disclosed are synthetic nucleic acid molecules, including DNA and RNA, and variations thereof. Unless otherwise specified, the oligonucleotide probes can be synthesized in DNA or RNA format or as hybrid molecules as desired. The methods disclosed comprise diagnostic processes and techniques using one or more aptamer of the invention, to determine the level or presence of relevant microvesicle surface antigens or a functional fragment thereof. Alternatively, an oligonucleotide probe of the invention can also be used as a binding agent to capture, isolate, or enrich, a cell, cell fragment, vesicle or any other fragment or complex that comprises the antigen or functional fragments thereof.

[0079] The compositions and methods of the invention comprise individual oligonucleotides that are identified for use in assessing a biomarker profile. The invention further discloses compositions and methods of oligonucleotide pools that can be used to detect a biomarker profile in a given sample.

[0080] Oligonucleotide probes and sequences disclosed in the compositions and methods of the invention may be identified herein in the form of DNA or RNA. Unless otherwise specified, one of skill in the art will appreciate that an oligonucleotide may generally be synthesized as either form of nucleic acid and carry various chemical modifications and remain within the scope of the invention. The term aptamer may be used in the art to refer to a single oligonucleotide that binds specifically to a target of interest through mechanisms other than Watson crick base pairing, similar to binding of a monoclonal antibody to a particular antigen. Within the scope of this disclosure and unless stated explicitly or otherwise implicit in context, the terms aptamer, oligonucleotide and oligonucleotide probe, and variations thereof, may be used interchangeably to refer to an oligonucleotide capable of distinguishing biological entities of interest (e.g, biomarkers) whether or not the specific entity has been identified or whether the precise mode of binding has been determined.

[0081] An oligonucleotide probe of the invention can also be used to provide *in vitro* or *in vivo* detection or imaging, to provide any appropriate diagnostic readout (e.g., diagnostic, prognostic or theranostic).

[0082] Separately, an oligonucleotide probe of the invention can also be used for treatment or as a therapeutic to specifically target a cell, tissue or organ.

Aptamers

[0083] SELEX. A suitable method for generating an aptamer is with the process entitled "Systematic Evolution of Ligands by Exponential Enrichment" ("SELEX") generally described in, e.g., U.S. patent application Ser. No. 07/536,428, filed Jun. 11, 1990, now abandoned, U.S. Pat. No. 5,475,096 entitled "Nucleic Acid Ligands", and U.S. Pat. No. 5,270,163 (see also WO 91/19813) entitled "Nucleic Acid Ligands". Each SELEX-identified nucleic acid ligand, i.e., each aptamer, is a specific ligand of a given target compound or molecule. The SELEX process is based on the unique insight that nucleic acids have sufficient capacity for forming a variety of two- and three-dimensional structures and sufficient chemical versatility available within their monomers to act as ligands (i.e., form specific binding pairs) with virtually any chemical compound, whether monomeric or polymeric. Molecules of any size or composition can serve as targets.

[0084] SELEX relies as a starting point upon a large library or pool of single stranded oligonucleotides comprising randomized sequences. The oligonucleotides can be modified or unmodified DNA, RNA, or DNA/RNA hybrids. In some examples, the pool comprises 100% random or partially random oligonucleotides. In other examples, the pool comprises random or partially random oligonucleotides containing at least one fixed and/or conserved sequence incorporated within randomized sequence. In other examples, the pool comprises random or partially random oligonucleotides containing at least one fixed and/or conserved sequence at its 5' and/or 3' end which may comprise a sequence shared by all the molecules of the oligonucleotide pool. Fixed sequences are sequences such as hybridization sites for PCR primers, promoter sequences for RNA polymerases (e.g., T3, T4, T7, and SP6), restriction sites, or homopolymeric sequences, such as poly A or poly T tracts, catalytic cores, sites for selective binding to affinity columns, and other sequences to facilitate cloning and/or sequencing of an oligonucleotide of interest. Conserved sequences are sequences, other than the previously described fixed sequences, shared by a number of aptamers that bind to the same target.

[0085] The oligonucleotides of the pool preferably include a randomized sequence portion as well as fixed sequences necessary for efficient amplification. Typically the oligonucleotides of the starting pool contain fixed 5' and 3' terminal sequences which flank an internal region of 30-50 random nucleotides. The randomized nucleotides can be produced in a number of ways including chemical synthesis and size selection from randomly cleaved cellular nucleic acids. Sequence variation in test nucleic acids can also be introduced or increased by mutagenesis before or during the selection/amplification iterations.

[0086] The random sequence portion of the oligonucleotide can be of any length and can comprise ribonucleotides and/or deoxyribonucleotides and can include modified or non-natural nucleotides or nucleotide analogs. See, e.g. U.S. Pat. No. 5,958,691; U.S. Pat. No. 5,660,985; U.S. Pat. No. 5,958,691; U.S. Pat. No. 5,698,687; U.S. Pat. No. 5,817,635; U.S. Pat. No. 5,672,695, and PCT Publication WO 92/07065. Random oligonucleotides can be synthesized from phosphodiester-linked nucleotides using solid phase oligonucleotide synthesis techniques well known in the art. See, e.g., Froehler et al., Nucl. Acid Res. 14:5399-5467 (1986) and Froehler et al., Tet. Lett. 27:5575-5578 (1986). Random oligonucleotides can also be synthesized using solution phase methods such as triester synthesis methods. See, e.g., Sood et al., Nucl. Acid Res. 4:2557 (1977) and Hirose et al., Tet. Lett., 28:2449 (1978). Typical syntheses carried out on automated DNA synthesis equipment yield 10^{14}-10^{16} individual molecules, a number sufficient for most SELEX experiments. Sufficiently large regions of random sequence in the sequence design increases the likelihood that each synthesized molecule is likely to represent a unique sequence.

[0087] The starting library of oligonucleotides may be generated by automated chemical synthesis on a DNA synthesizer. To synthesize randomized sequences, mixtures of all four nucleotides are added at each nucleotide addition step during the synthesis process, allowing for random incorporation of nucleotides. As stated above, in one embodiment, random oligonucleotides comprise entirely random sequences; however, in other embodiments, random oligonucleotides can comprise stretches of nonrandom or

partially random sequences. Partially random sequences can be created by adding the four nucleotides in different molar ratios at each addition step.

[0088] The starting library of oligonucleotides may be for example, RNA, DNA, or RNA/DNA hybrid. In those instances where an RNA library is to be used as the starting library it is typically generated by transcribing a DNA library in vitro using T7 RNA polymerase or modified T7 RNA polymerases and purified. The library is then mixed with the target under conditions favorable for binding and subjected to step-wise iterations of binding, partitioning and amplification, using the same general selection scheme, to achieve virtually any desired criterion of binding affinity and selectivity. More specifically, starting with a mixture containing the starting pool of nucleic acids, the SELEX method includes steps of: (a) contacting the mixture with the target under conditions favorable for binding; (b) partitioning unbound nucleic acids from those nucleic acids which have bound specifically to target molecules; (c) dissociating the nucleic acid-target complexes; (d) amplifying the nucleic acids dissociated from the nucleic acid-target complexes to yield a ligand-enriched mixture of nucleic acids; and (e) reiterating the steps of binding, partitioning, dissociating and amplifying through as many cycles as desired to yield highly specific, high affinity nucleic acid ligands to the target molecule. In those instances where RNA aptamers are being selected, the SELEX method further comprises the steps of: (i) reverse transcribing the nucleic acids dissociated from the nucleic acid-target complexes before amplification in step (d); and (ii) transcribing the amplified nucleic acids from step (d) before restarting the process.

[0089] Within a nucleic acid mixture containing a large number of possible sequences and structures, there is a wide range of binding affinities for a given target. A nucleic acid mixture comprising, for example, a 20 nucleotide randomized segment can have 4^{20} candidate possibilities. Those which have the higher affinity constants for the target are most likely to bind to the target. After partitioning, dissociation and amplification, a second nucleic acid mixture is generated, enriched for the higher binding affinity candidates. Additional rounds of selection progressively favor better ligands until the resulting nucleic acid mixture is predominantly composed of only one or a few sequences. These can then be cloned, sequenced and individually tested for binding affinity as pure ligands or aptamers.

[0090] Cycles of selection and amplification are repeated until a desired goal is achieved. In the most general case, selection/amplification is continued until no significant improvement in binding strength is achieved on repetition of the cycle. The method is typically used to sample approximately 10^{14} different nucleic acid species but may be used to sample as many as about 10^{18} different nucleic acid species. Generally, nucleic acid aptamer molecules are selected in a 5 to 20 cycle procedure. In one embodiment, heterogeneity is introduced only in the initial selection stages and does not occur throughout the replicating process.

[0091] In one embodiment of SELEX, the selection process is so efficient at isolating those nucleic acid ligands that bind most strongly to the selected target, that only one cycle of selection and amplification is required. Such an efficient selection may occur, for example, in a chromatographic-type process wherein

the ability of nucleic acids to associate with targets bound on a column operates in such a manner that the column is sufficiently able to allow separation and isolation of the highest affinity nucleic acid ligands.

[0092] In many cases, it is not necessarily desirable to perform the iterative steps of SELEX until a single nucleic acid ligand is identified. The target-specific nucleic acid ligand solution may include a family of nucleic acid structures or motifs that have a number of conserved sequences and a number of sequences which can be substituted or added without significantly affecting the affinity of the nucleic acid ligands to the target. By terminating the SELEX process prior to completion, it is possible to determine the sequence of a number of members of the nucleic acid ligand solution family. The invention provides for the identification of aptamer pools and uses thereof that jointly can be used to characterize a test sample. For example, the aptamer pools can be identified through rounds of positive and negative selection to identify microvesicle indicative of a disease or condition. The invention further provides use of such aptamer pools to detect and/or quantify such microvesicles in a sample, thereby allowing a diagnosis, prognosis or theranosis to be provided.

[0093] A variety of nucleic acid primary, secondary and tertiary structures are known to exist. The structures or motifs that have been shown most commonly to be involved in non-Watson-Crick type interactions are referred to as hairpin loops, symmetric and asymmetric bulges, pseudoknots and myriad combinations of the same. Almost all known cases of such motifs suggest that they can be formed in a nucleic acid sequence of no more than 30 nucleotides. For this reason, it is often preferred that SELEX procedures with contiguous randomized segments be initiated with nucleic acid sequences containing a randomized segment of between about 20 to about 50 nucleotides and in some embodiments, about 30 to about 40 nucleotides. In one example, the 5'-fixed:random:3'-fixed sequence comprises a random sequence of about 30 to about 50 nucleotides.

[0094] The core SELEX method has been modified to achieve a number of specific objectives. For example, U.S. Pat. No. 5,707,796 describes the use of SELEX in conjunction with gel electrophoresis to select nucleic acid molecules with specific structural characteristics, such as bent DNA. U.S. Pat. No. 5,763,177 describes SELEX based methods for selecting nucleic acid ligands containing photoreactive groups capable of binding and/or photocrosslinking to and/or photoinactivating a target molecule. U.S. Pat. No. 5,567,588 and U.S. Pat. No. 5,861,254 describe SELEX based methods which achieve highly efficient partitioning between oligonucleotides having high and low affinity for a target molecule. U.S. Pat. No. 5,496,938 describes methods for obtaining improved nucleic acid ligands after the SELEX process has been performed. U.S. Pat. No. 5,705,337 describes methods for covalently linking a ligand to its target.

[0095] SELEX can also be used to obtain nucleic acid ligands that bind to more than one site on the target molecule, and to obtain nucleic acid ligands that include non-nucleic acid species that bind to specific sites on the target. SELEX provides means for isolating and identifying nucleic acid ligands which bind to any envisionable target, including large and small biomolecules such as nucleic acid-binding proteins and proteins not known to bind nucleic acids as part of their biological function as well

as lipids, cofactors and other small molecules. For example, U.S. Pat. No. 5,580,737 discloses nucleic acid sequences identified through SELEX which are capable of binding with high affinity to caffeine and the closely related analog, theophylline.

[0096] Counter-SELEX is a method for improving the specificity of nucleic acid ligands to a target molecule by eliminating nucleic acid ligand sequences with cross-reactivity to one or more non-target molecules. Counter-SELEX is comprised of the steps of: (a) preparing a candidate mixture of nucleic acids; (b) contacting the candidate mixture with the target, wherein nucleic acids having an increased affinity to the target relative to the candidate mixture may be partitioned from the remainder of the candidate mixture; (c) partitioning the increased affinity nucleic acids from the remainder of the candidate mixture; (d) dissociating the increased affinity nucleic acids from the target; e) contacting the increased affinity nucleic acids with one or more non-target molecules such that nucleic acid ligands with specific affinity for the non-target molecule(s) are removed; and (f) amplifying the nucleic acids with specific affinity only to the target molecule to yield a mixture of nucleic acids enriched for nucleic acid sequences with a relatively higher affinity and specificity for binding to the target molecule. As described above for SELEX, cycles of selection and amplification are repeated until a desired goal is achieved.

[0097] One potential problem encountered in the use of nucleic acids as therapeutics and vaccines is that oligonucleotides in their phosphodiester form may be quickly degraded in body fluids by intracellular and extracellular enzymes such as endonucleases and exonucleases before the desired effect is manifest. The SELEX method thus encompasses the identification of high-affinity nucleic acid ligands containing modified nucleotides conferring improved characteristics on the ligand, such as improved in vivo stability or improved delivery characteristics. Examples of such modifications include chemical substitutions at the ribose and/or phosphate and/or base positions. SELEX identified nucleic acid ligands containing modified nucleotides are described, e.g., in U.S. Pat. No. 5,660,985, which describes oligonucleotides containing nucleotide derivatives chemically modified at the 2' position of ribose, 5' position of pyrimidines, and 8' position of purines, U.S. Pat. No. 5,756,703 which describes oligonucleotides containing various 2'-modified pyrimidines, and U.S. Pat. No. 5,580,737 which describes highly specific nucleic acid ligands containing one or more nucleotides modified with 2'-amino (2'--NH_2), 2'-fluoro (2'-F), and/or 2'-O-methyl (2'-OMe) substituents.

[0098] Modifications of the nucleic acid ligands contemplated in this invention include, but are not limited to, those which provide other chemical groups that incorporate additional charge, polarizability, hydrophobicity, hydrogen bonding, electrostatic interaction, and fluxionality to the nucleic acid ligand bases or to the nucleic acid ligand as a whole. Modifications to generate oligonucleotide populations which are resistant to nucleases can also include one or more substitute internucleotide linkages, altered sugars, altered bases, or combinations thereof. Such modifications include, but are not limited to, 2'-position sugar modifications, 5-position pyrimidine modifications, 8-position purine modifications, modifications at exocyclic amines, substitution of 4-thiouridine, substitution of 5-bromo or 5-iodo-uracil; backbone modifications, phosphorothioate or allyl phosphate modifications, methylations, and unusual

base-pairing combinations such as the isobases isocytidine and isoguanosine. Modifications can also include 3' and 5' modifications such as capping.

[0099] In one embodiment, oligonucleotides are provided in which the P(O)O group is replaced by P(O)S ("thioate"), P(S)S ("dithioate"), P(O)NR_2 ("amidate"), P(O)R, P(O)OR', CO or CH_2 ("formacetal") or 3'-amine (--NH--CH_2--CH_2--), wherein each R or R' is independently H or substituted or unsubstituted alkyl. Linkage groups can be attached to adjacent nucleotides through an --O--, --N--, or --S-- linkage. Not all linkages in the oligonucleotide are required to be identical. As used herein, the term phosphorothioate encompasses one or more non-bridging oxygen atoms in a phosphodiester bond replaced by one or more sulfur atoms.

[00100] In further embodiments, the oligonucleotides comprise modified sugar groups, for example, one or more of the hydroxyl groups is replaced with halogen, aliphatic groups, or functionalized as ethers or amines. In one embodiment, the 2'-position of the furanose residue is substituted by any of an O-methyl, O-alkyl, O-allyl, S-alkyl, S-allyl, or halo group. Methods of synthesis of 2'-modified sugars are described, e.g., in Sproat, et al., Nucl. Acid Res. 19:733-738 (1991); Cotten, et al., Nucl. Acid Res. 19:2629-2635 (1991); and Hobbs, et al., Biochemistry 12:5138-5145 (1973). Other modifications are known to one of ordinary skill in the art. Such modifications may be pre-SELEX process modifications or post-SELEX process modifications (modification of previously identified unmodified ligands) or may be made by incorporation into the SELEX process.

[00101] Pre-SELEX process modifications or those made by incorporation into the SELEX process yield nucleic acid ligands with both specificity for their SELEX target and improved stability, e.g., in vivo stability. Post-SELEX process modifications made to nucleic acid ligands may result in improved stability, e.g., in vivo stability without adversely affecting the binding capacity of the nucleic acid ligand.

[00102] The SELEX method encompasses combining selected oligonucleotides with other selected oligonucleotides and non-oligonucleotide functional units as described in U.S. Pat. No. 5,637,459 and U.S. Pat. No. 5,683,867. The SELEX method further encompasses combining selected nucleic acid ligands with lipophilic or non-immunogenic high molecular weight compounds in a diagnostic or therapeutic complex, as described, e.g., in U.S. Pat. No. 6,011,020, U.S. Pat. No. 6,051,698, and PCT Publication No. WO 98/18480. These patents and applications teach the combination of a broad array of shapes and other properties, with the efficient amplification and replication properties of oligonucleotides, and with the desirable properties of other molecules.

[00103] The identification of nucleic acid ligands to small, flexible peptides via the SELEX method has also been explored. Small peptides have flexible structures and usually exist in solution in an equilibrium of multiple conformers, and thus it was initially thought that binding affinities may be limited by the conformational entropy lost upon binding a flexible peptide. However, the feasibility of identifying nucleic acid ligands to small peptides in solution was demonstrated in U.S. Pat. No. 5,648,214. In this patent, high affinity RNA nucleic acid ligands to substance P, an 11 amino acid peptide, were identified.

[00104] The aptamers with specificity and binding affinity to the target(s) of the present invention can be selected by the SELEX N process as described herein. As part of the SELEX process, the sequences selected to bind to the target are then optionally minimized to determine the minimal sequence having the desired binding affinity. The selected sequences and/or the minimized sequences are optionally optimized by performing random or directed mutagenesis of the sequence to increase binding affinity or alternatively to determine which positions in the sequence are essential for binding activity. Additionally, selections can be performed with sequences incorporating modified nucleotides to stabilize the aptamer molecules against degradation in vivo.

[00105] ***2' Modified SELEX***

[00106] For an aptamer to be suitable for use as a therapeutic, it is preferably inexpensive to synthesize, and safe and stable in vivo. Wild-type RNA and DNA aptamers are typically not stable is vivo because of their susceptibility to degradation by nucleases. Resistance to nuclease degradation can be greatly increased by the incorporation of modifying groups at the 2'-position.

[00107] Fluoro and amino groups have been successfully incorporated into oligonucleotide pools from which aptamers have been subsequently selected. However, these modifications greatly increase the cost of synthesis of the resultant aptamer, and may introduce safety concerns in some cases because of the possibility that the modified nucleotides could be recycled into host DNA by degradation of the modified oligonucleotides and subsequent use of the nucleotides as substrates for DNA synthesis.

[00108] Aptamers that contain 2'-O-methyl ("2'-OMe") nucleotides, as provided herein, may overcome one or more potential drawbacks. Oligonucleotides containing 2'-OMe nucleotides are nuclease-resistant and inexpensive to synthesize. Although 2'-OMe nucleotides are ubiquitous in biological systems, natural polymerases do not accept 2'-OMe NTPs as substrates under physiological conditions, thus there are no safety concerns over the recycling of 2'-OMe nucleotides into host DNA. The SELEX method used to generate 2'-modified aptamers is described, e.g., in U.S. Provisional Patent Application Ser. No. 60/430,761, filed Dec. 3, 2002, U.S. Provisional Patent Application Ser. No. 60/487,474, filed Jul. 15, 2003, U.S. Provisional Patent Application Ser. No. 60/517,039, filed Nov. 4, 2003, U.S. patent application Ser. No. 10/729,581, filed Dec. 3, 2003, and U.S. patent application Ser. No. 10/873,856, filed Jun. 21, 2004, entitled "Method for in vitro Selection of 2'-O-methyl substituted Nucleic Acids", each of which is herein incorporated by reference in its entirety.

METHODS

[00109] ***Biomarker Detection and Diagnotics***

[00110] The aptamers of the invention can be used in various methods to assess presence or level of biomarkers in a biological sample, e.g., biological entities of interest such as proteins, nucleic acids, or microvesicles. The aptamer functions as a binding agent to assess presence or level of the cognate target molecule. Therefore, in various embodiments of the invention directed to diagnostics, prognostics or theranostics, one or more aptamers of the invention are configured in a ligand-target based assay, where one or more aptamer of the invention is contacted with a selected biological sample, where the or more

aptamer associates with or binds to its target molecules. Aptamers of the invention are used to identify candidate biosignatures based on the biological samples assessed and biomarkers detected. In further embodiments, aptamers may themselves provide a biosignature for a particular condition or disease. A biosignature refers to a biomarker profile of a biological sample comprising a presence, level or other characteristic that can be assessed (including without limitation a sequence, mutation, rearrangement, translocation, deletion, epigenetic modification, methylation, post-translational modification, allele, activity, complex partners, stability, half life, and the like) of one or more biomarker of interest. Biosignatures can be used to evaluate diagnostic and/or prognostic criteria such as presence of disease, disease staging, disease monitoring, disease stratification, or surveillance for detection, metastasis or recurrence or progression of disease. For example, methods of the invention using aptamers against microvesicle surface antigen are useful for correlating a biosignature comprising microvesicle antigens to a selected condition or disease. A biosignature can also be used clinically in making decisions concerning treatment modalities including therapeutic intervention. A biosignature can further be used clinically to make treatment decisions, including whether to perform surgery or what treatment standards should be used along with surgery (e.g., either pre-surgery or post-surgery). As an illustrative example, a biosignature of circulating biomarkers that indicates an aggressive form of cancer may call for a more aggressive surgical procedure and/or more aggressive therapeutic regimen to treat the patient.

[00111] A biosignature can be used in any methods disclosed herein, e.g., to assess whether a subject is afflicted with disease, is at risk for developing disease or to assess the stage or progression of the disease. For example, a biosignature can be used to assess whether a subject has prostate cancer, colon cancer, or other cancer as described herein. Furthermore, a biosignature can be used to determine a stage of a disease or condition, such as colon cancer. The biosignature/biomarker profile comprising a microvesicle can include assessment of payload within the microvesicle. For example, one or more aptamer of the invention can be used to capture a microvesicle population, thereby providing readout of microvesicle antigens, and then the payload content within the captured microvesicles can be assessed, thereby providing further biomarker readout of the payload content.

[00112] A biosignature for characterizing a phenotype may comprise any number of useful criteria. As described further below, the term "phenotype" as used herein can mean any trait or characteristic that is attributed to a biosignature / biomarker profile. A phenotype can be detected or identified in part or in whole using the compositions and/or methods of the invention. In some embodiments, at least one criterion is used for each biomarker. In some embodiments, at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90 or at least 100 criteria are used. For example, for the characterizing of a cancer, a number of different criteria can be used when the subject is diagnosed with a cancer: 1) if the amount of microRNA in a sample from a subject is higher than a reference value; 2) if the amount of a microRNA within cell type specific vesicles (i.e. vesicles derived from a specific tissue or organ) is higher than a reference value; or 3) if the amount of microRNA within vesicles with one or more cancer specific biomarkers is higher than a reference value. Similar rules can apply if the amount of microRNA is less

than or the same as the reference. The method can further include a quality control measure, such that the results are provided for the subject if the samples meet the quality control measure. In some embodiments, if the criteria are met but the quality control is questionable, the subject is reassessed.

[00113] *Theranostics*

[00114] A biosignature can be used in therapy related diagnostics to provide tests useful to diagnose a disease or choose the correct treatment regimen, such as provide a theranosis. Theranostics includes diagnostic testing that provides the ability to affect therapy or treatment of a diseased state. Theranostics testing provides a theranosis in a similar manner that diagnostics or prognostic testing provides a diagnosis or prognosis, respectively. As used herein, theranostics encompasses any desired form of therapy related testing, including predictive medicine, personalized medicine, integrated medicine, pharmacodiagnostics and Dx/Rx partnering. Therapy related tests can be used to predict and assess drug response in individual subjects, i.e., to provide personalized medicine. Predicting a drug response can be determining whether a subject is a likely responder or a likely non-responder to a candidate therapeutic agent, e.g., before the subject has been exposed or otherwise treated with the treatment. Assessing a drug response can be monitoring a response to a drug, e.g., monitoring the subject's improvement or lack thereof over a time course after initiating the treatment. Therapy related tests are useful to select a subject for treatment who is particularly likely to benefit from the treatment or to provide an early and objective indication of treatment efficacy in an individual subject. Thus, a biosignature as disclosed herein may indicate that treatment should be altered to select a more promising treatment, thereby avoiding the great expense of delaying beneficial treatment and avoiding the financial and morbidity costs of administering an ineffective drug(s).

[00115] The compositions and methods of the invention can be used to identify or detect a biosignature associated with a variety of diseases and disorders, which include, but are not limited to cardiovascular disease, cancer, infectious diseases, sepsis, neurological diseases, central nervous system related diseases, endovascular related diseases, and autoimmune related diseases. Therapy related diagnostics (i.e., theranostics) are also useful in clinical diagnosis and management of many such diseases and disorders. Therapy related diagnostics also aid in the prediction of drug toxicity, drug resistance or drug response. Therapy related tests may be developed in any suitable diagnostic testing format, which include, but are not limited to, e.g., immunohistochemical tests, clinical chemistry, immunoassay, cell-based technologies, nucleic acid tests or body imaging methods. Therapy related tests can further include but are not limited to, testing that aids in the determination of therapy, testing that monitors for therapeutic toxicity, or response to therapy testing. Thus, a biosignature can be used to predict or monitor a subject's response to a treatment. A biosignature can be determined at different time points for a subject after initiating, removing, or altering a particular treatment.

[00116] In some embodiments, the compositions and methods of the invention provide for a determination or prediction as to whether a subject is responding to a treatment is made based on a change in the amount of one or more components of a biosignature (i.e., the microRNA, vesicles and/or biomarkers of interest),

an amount of one or more components of a particular biosignature, or the biosignature detected for the components. In another embodiment, a subject's condition is monitored by determining a biosignature at different time points. The progression, regression, or recurrence of a condition is determined. Response to therapy can also be measured over a time course. Thus, the invention provides a method of monitoring a status of a disease or other medical condition in a subject, comprising isolating or detecting a biosignature from a biological sample from the subject, detecting the overall amount of the components of a particular biosignature, or detecting the biosignature of one or more components (such as the presence, absence, or expression level of a biomarker). The biosignatures are used to monitor the status of the disease or condition.

[00117] One or more novel biosignatures of a vesicle can also be identified. For example, one or more vesicles can be isolated from a subject that responds to a drug treatment or treatment regimen and compared to a reference, such as another subject that does not respond to the drug treatment or treatment regimen. Differences between the biosignatures can be determined and used to identify other subjects as responders or non-responders to a particular drug or treatment regimen.

[00118] In some embodiments, a biosignature is used to determine whether a particular disease or condition is resistant to a drug, in which case a physician need not waste valuable time with such drug treatment. To obtain early validation of a drug choice or treatment regimen, a biosignature is determined for a sample obtained from a subject. The biosignature is used to assess whether the particular subject's disease has the biomarker associated with drug resistance. Such a determination enables doctors to devote critical time as well as the patient's financial resources to effective treatments.

[00119] Biosignatures can be used in the theranosis of a cancer, such as identifying whether a subject suffering from cancer is a likely responder or non-responder to a particular cancer treatment. The subject methods can be used to theranose cancers including those listed herein, e.g., in the "Phenotypes" section below. These include without limitation lung cancer, non-small cell lung cancer small cell lung cancer (including small cell carcinoma (oat cell cancer), mixed small cell/large cell carcinoma, and combined small cell carcinoma), colon cancer, breast cancer, prostate cancer, liver cancer, pancreatic cancer, brain cancer, kidney cancer, ovarian cancer, stomach cancer, melanoma, bone cancer, gastric cancer, breast cancer, glioma, glioblastoma, hepatocellular carcinoma, papillary renal carcinoma, head and neck squamous cell carcinoma, leukemia, lymphoma, myeloma, or other solid tumors.

[00120] A biosignature of circulating biomarkers, including markers associated with a component present in a biological sample (e.g., cell, cell-fragment, cell-derived extracellular vesicle), in a sample from a subject suffering from a cancer can be used select a candidate treatment for the subject. The biosignature can be determined according to the methods of the invention presented herein. In some embodiments, the candidate treatment comprises a standard of care for the cancer. The treatment can be a cancer treatment such as radiation, surgery, chemotherapy or a combination thereof. The cancer treatment can be a therapeutic such as anti-cancer agents and chemotherapeutic regimens. Further drug associations and rules that are used in embodiments of the invention are found in PCT/US2007/69286, filed May 18, 2007; PCT/

US2009/60630, filed October 14, 2009; PCT/ 2010/000407, filed February 11, 2010; PCT/US12/41393, filed June 7, 2012; PCT/US2013/073184, filed December 4, 2013; PCT/US2010/54366, filed October 27, 2010; PCT/US11/67527, filed December 28, 2011; PCT/US15/13618, filed January 29, 2015; and PCT/US16/20657, filed March 3, 2016.

Biomarker Detection

[00121] The compositions and methods of the invention can be used to assess any useful biomarkers in a biological sample for charactering a phenotype associated with the sample. Such biomarkers include all sorts of biological entities such as proteins, nucleic acids, lipids, carbohydrates, complexes of any thereof, and microvesicles. Various molecules associated with a microvesicle surface or enclosed within the microvesicle (referred to herein as "payload") can serve as biomarkers. The microvesicles themselves can also be used as biomarkers.

[00122] The aptamers of the invention can be used to assess levels or presence of a microvesicle population. See, e.g., **FIGs. 19B-C**. The aptamers of the invention can also be used to assess levels or presence of their specific target molecule. See, e.g., **FIG. 19A**. In addition, aptamers of the invention are used to capture or isolated a component present in a biological sample that has the aptamer's target molecule present. For example, if a given microvesicle surface antigen is present on a cell, cell fragment or cell-derived extracellular vesicle. A binding agent to the biomarker, including without limitation an aptamer provided by the invention, may be used to capture or isolate the cell, cell fragment or cell-derived extracellular vesicles. See, e.g., **FIGs. 2A-B**, **19A**. Such captured or isolated entities may be further characterized to assess additional surface antigens or internal "payload" molecules present (i.e., nucleic acid molecules, lipids, sugars, polypeptides or functional fragments thereof, or anything else present in the cellular milieu that may be used as a biomarker), where one or more biomarkers provide a biosignature to assess a desired phenotype, such a s disease or condition. See, e.g., **FIG. 2F**. Therefore, aptamers of the invention are used not only to assess one or more microvesicle surface antigen of interest but are also used to separate a component present in a biological sample, where the components themselves can be further assessed to identify a candidate biosignature.

[00123] The methods of the invention can comprise multiplex analysis of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 50, 75 or 100 different biomarkers. For example, an assay of a heterogeneous population of vesicles can be performed with a plurality of particles that are differentially labeled. There can be at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 50, 75 or 100 differentially labeled particles. The particles may be externally labeled, such as with a tag, or they may be intrinsically labeled. Each differentially labeled particle can be coupled to a capture agent, such as a binding agent, for a vesicle, resulting in capture of a vesicle. The multiple capture agents can be selected to characterize a phenotype of interest, including capture agents against general vesicle biomarkers, cell-of-origin specific biomarkers, and disease biomarkers. One or more biomarkers of the captured vesicle can then be detected by a plurality of binding agents. The binding agent can be directly labeled to facilitate detection. Alternatively, the binding agent is labeled by a secondary agent. For example, the

binding agent may be an antibody for a biomarker on the vesicle, wherein the binding agent is linked to biotin. A secondary agent comprises streptavidin linked to a reporter and can be added to detect the biomarker. In some embodiments, the captured vesicle is assayed for at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 50, 75 or 100 different biomarkers. For example, multiple detectors, i.e., detection of multiple biomarkers of a captured vesicle or population of vesicles, can increase the signal obtained, permitted increased sensitivity, specificity, or both, and the use of smaller amounts of samples. Detection can be with more than one biomarker, including without limitation more than one vesicle marker such as in any of **Tables 3-4**, and **Tables 18-25**.

[00124] An immunoassay based method (e.g., sandwich assay) can be used to detect a biomarker of a vesicle. An example includes ELISA. A binding agent can be bound to a well. For example, a binding agent such as an aptamer or antibody to an antigen of a vesicle can be attached to a well. A biomarker on the captured vesicle can be detected based on the methods described herein. **FIG. 2A** shows an illustrative schematic for a sandwich-type of immunoassay. The capture agent can be against a vesicle antigen of interest, e.g., a general vesicle biomarker, a cell-of-origin marker, or a disease marker. In the figure, the captured vesicles are detected using fluorescently labeled binding agent (detection agent) against vesicle antigens of interest. Multiple capture binding agents can be used, e.g., in distinguishable addresses on an array or different wells of an immunoassay plate. The detection binding agents can be against the same antigen as the capture binding agent, or can be directed against other markers. The capture binding agent can be any useful binding agent, e.g., tethered aptamers, antibodies or lectins, and/or the detector antibodies can be similarly substituted, e.g., with detectable (e.g., labeled) aptamers, antibodies, lectins or other binding proteins or entities. In an embodiment, one or more capture agents to a general vesicle biomarker, a cell-of-origin marker, and/or a disease marker are used along with detection agents against general vesicle biomarker, such as tetraspanin molecules including without limitation one or more of CD9, CD63 and CD81, or other markers in **Table 3** herein. Examples of microvesicle surface antigens are disclosed herein, *e.g.* in **Tables 3- 4** and **18-25**, or are known in the art, and examples useful in methods and compositions of the invention are disclosed of International Patent Application Nos. PCT/US2009/62880, filed October 30, 2009; PCT/US2009/006095, filed November 12, 2009; PCT/US2011/26750, filed March 1, 2011; PCT/US2011/031479, filed April 6, 2011; PCT/US11/48327, filed August 18, 2011; PCT/US2008/71235, filed July 25, 2008; PCT/US10/58461, filed November 30, 2010; PCT/US2011/21160, filed January 13, 2011; PCT/US2013/030302, filed March 11, 2013; PCT/US12/25741, filed February 17, 2012; PCT/2008/76109, filed September 12, 2008; PCT/US12/42519, filed June 14, 2012; PCT/US12/50030, filed August 8, 2012; PCT/US12/49615, filed August 3, 2012; PCT/US12/41387, filed June 7, 2012; PCT/US2013/072019, filed November 26, 2013; PCT/US2014/039858, filed May 28, 2013; PCT/IB2013/003092, filed October 23, 2013; PCT/US13/76611, filed December 19, 2013; PCT/US14/53306, filed August 28, 2014; and PCT/US15/62184, filed November 23, 2015; each of which applications is incorporated herein by reference in its entirety.

[00125] **FIG. 2D** presents an illustrative schematic for analyzing vesicles according to the methods of the invention. Capture agents are used to capture vesicles, detectors are used to detect the captured vesicles, and the level or presence of the captured and detected microvesicles is used to characterize a phenotype. Capture agents, detectors and characterizing phenotypes can be any of those described herein. For example, capture agents include antibodies or aptamers tethered to a substrate that recognize a vesicle antigen of interest, detectors include labeled antibodies or aptamers to a vesicle antigen of interest, and characterizing a phenotype includes a diagnosis, prognosis, or theranosis of a disease. In the scheme shown in **FIG. 2D i)**, a population of vesicles is captured with one or more capture agents against general vesicle biomarkers (**200**). The captured vesicles are then labeled with detectors against cell-of-origin biomarkers (**201**) and/or disease specific biomarkers (**202**). If only cell-of-origin detectors are used (**201**), the biosignature used to characterize the phenotype (**203**) can include the general vesicle markers (**200**) and the cell-of-origin biomarkers (**201**). If only disease detectors are used (**202**), the biosignature used to characterize the phenotype (**203**) can include the general vesicle markers (**200**) and the disease biomarkers (**202**). Alternately, detectors are used to detect both cell-of-origin biomarkers (**201**) and disease specific biomarkers (**202**). In this case, the biosignature used to characterize the phenotype (**203**) can include the general vesicle markers (**200**), the cell-of-origin biomarkers (**201**) and the disease biomarkers (**202**). The biomarkers combinations are selected to characterize the phenotype of interest and can be selected from the biomarkers and phenotypes described herein, e.g., in **Tables 1**, **3-4** and **18-25**.

[00126] In the scheme shown in **FIG. 2D ii)**, a population of vesicles is captured with one or more capture agents against cell-of-origin biomarkers (**210**) and/or disease biomarkers (**211**). The captured vesicles are then detected using detectors against general vesicle biomarkers (**212**). If only cell-of-origin capture agents are used (**210**), the biosignature used to characterize the phenotype (**213**) can include the cell-of-origin biomarkers (**210**) and the general vesicle markers (**212**). If only disease biomarker capture agents are used (**211**), the biosignature used to characterize the phenotype (**213**) can include the disease biomarkers (**211**) and the general vesicle biomarkers (**212**). Alternately, capture agents to one or more cell-of-origin biomarkers (**210**) and one or more disease specific biomarkers (**211**) are used to capture vesicles. In this case, the biosignature used to characterize the phenotype (**213**) can include the cell-of-origin biomarkers (**210**), the disease biomarkers (**211**), and the general vesicle markers (**213**). The biomarkers combinations are selected to characterize the phenotype of interest and can be selected from the biomarkers and phenotypes described herein.

[00127] The methods of the invention comprise capture and detection of microvesicles of interest using any combination of useful biomarkers. For example, a microvesicle population can be captured using one or more binding agent to any desired combination of cell of origin, disease specific, or general vesicle markers. The captured microvesicles can then be detected using one or more binding agent to any desired combination of cell of origin, disease specific, or general vesicle markers. **FIG. 2E** represents a flow diagram of such configurations. Any one or more of a cell-of-origin biomarker (**240**), disease biomarkers (**241**), and general vesicle biomarker (**242**) is used to capture a microvesicle population. Thereafter, any

one or more of a cell-of-origin biomarker (**243**), disease biomarkers (**244**), and general vesicle biomarker (**245**) is used to detect the captured microvesicle population. The biosignature of captured and detected microvesicles is then used to characterize a phenotype. The biomarkers combinations are selected to characterize the phenotype of interest and can be selected from the biomarkers and phenotypes described herein.

[00128] A microvesicle payload molecule can be assessed as a member of a biosignature panel. A payload molecule comprises any of the biological entities contained within a cell, cell fragment or vesicle membrane. These entities include without limitation nucleic acids, e.g., mRNA, microRNA, or DNA fragments; protein, e.g., soluble and membrane associated proteins; carbohydrates; lipids; metabolites; and various small molecules, e.g., hormones. The payload can be part of the cellular milieu that is encapsulated as a vesicle is formed in the cellular environment. In some embodiments of the invention, the payload is analyzed in addition to detecting vesicle surface antigens. Specific populations of vesicles can be captured as described above then the payload in the captured vesicles can be used to characterize a phenotype. For example, vesicles captured on a substrate can be further isolated to assess the payload therein. Alternately, the vesicles in a sample are detected and sorted without capture. The vesicles so detected can be further isolated to assess the payload therein. In an embodiment, vesicle populations are sorted by flow cytometry and the payload in the sorted vesicles is analyzed. In the scheme shown in **FIG. 2F iv)**, a population of vesicles is captured and/or detected (**220**) using one or more of cell-of-origin biomarkers (**220**), disease biomarkers (**221**), and/or general vesicle markers (**222**). The payload of the isolated vesicles is assessed (**223**). A biosignature detected within the payload can be used to characterize a phenotype (**224**). In a non-limiting example, a vesicle population can be analyzed in a plasma sample from a patient using antibodies against one or more vesicle antigens of interest. The antibodies can be capture antibodies which are tethered to a substrate to isolate a desired vesicle population. Alternately, the antibodies can be directly labeled and the labeled vesicles isolated by sorting with flow cytometry. The presence or level of microRNA or mRNA extracted from the isolated vesicle population can be used to detect a biosignature. The biosignature is then used to diagnose, prognose or theranose the patient.

[00129] In other embodiments, vesicle or cellular payload is analyzed in a population (e.g., cells or vesicles) without first capturing or detected subpopulations of vesicles. For example, a cellular or extracellular vesicle population can be generally isolated from a sample using centrifugation, filtration, chromatography, or other techniques as described herein and known in the art. The payload of such sample components can be analyzed thereafter to detect a biosignature and characterize a phenotype. In the scheme shown in **FIG. 2F v)**, a population of vesicles is isolated (**230**) and the payload of the isolated vesicles is assessed (**231**). A biosignature detected within the payload can be used to characterize a phenotype (**232**). In a non-limiting example, a vesicle population is isolated from a plasma sample from a patient using size exclusion and membrane filtration. The presence or level of microRNA or mRNA extracted from the vesicle population is used to detect a biosignature. The biosignature is then used to diagnose, prognose or theranose the patient.

[00130] The biomarkers used to detect a vesicle population can be selected to detect a microvesicle population of interest, e.g., a population of vesicles that provides a diagnosis, prognosis or theranosis of a selected condition or disease, including but not limited to a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. See Section "Phenotypes" herein for more detail. In an embodiment, the biomarkers are selected from the group consisting of EpCam (epithelial cell adhesion molecule), CD9 (tetraspanin CD9 molecule), PCSA (prostate cell specific antigen, *see* Rokhlin *et al.*, 5E10: a prostate-specific surface-reactive monoclonal antibody. Cancer Lett. 1998 **131**:129-36), CD63 (tetraspanin CD63 molecule), CD81 (tetraspanin CD81 molecule), PSMA (FOLH1, folate hydrolase (prostate-specific membrane antigen) 1), B7H3 (CD276 molecule), PSCA (prostate stem cell antigen), ICAM (intercellular adhesion molecule), STEAP (STEAP1, six transmembrane epithelial antigen of the prostate 1), KLK2 (kallikrein-related peptidase 2), SSX2 (synovial sarcoma, X breakpoint 2), SSX4 (synovial sarcoma, X breakpoint 4), PBP (prostatic binding protein), SPDEF (SAM pointed domain containing ets transcription factor), EGFR (epidermal growth factor receptor), and a combination thereof. One or more of these markers can provide a biosignature for a specific condition, such as to detect a cancer, including without limitation a carcinoma, a prostate cancer, a breast cancer, a lung cancer, a colorectal cancer, an ovarian cancer, melanoma, a brain cancer, or other type of cancer as disclosed herein. In an embodiment, a binding agent to one or more of these markers is used to capture a microvesicle population, and an aptamer of the invention is used to assist in detection of the capture vesicles as described herein. In other embodiments, an aptamer of the invention is used to capture a microvesicle population, and a binding agent to one or more of these markers is used to assist in detection of the capture vesicles as described herein. The binding agents can be any useful binding agent as disclosed herein or known in the art, e.g., antibodies or aptamers.

[00131] The methods of characterizing a phenotype can employ a combination of techniques to assess a component or population of components present in a biological sample of interest. For example, an aptamer of the invention can be used to assess a single cell, or a single extracellular vesicle or a population of cells or population of vesicles. A sample may be split into various aliquots, where each is analyzed separately. For example, protein content of one or more aliquot is determined and microRNA content of one or more other aliquot is determined. The protein content and microRNA content can be combined to characterize a phenotype. In another embodiment, a component present in a biological sample of interest is isolated and the payload therein is assessed (e.g., capture a population of subpopulation of vesicles using an aptamer of the invention and further assess nucleic acid or proteins present in the isolated vesicles).

[00132] In one embodiment, a population of vesicles with a given surface marker can be isolated by using a binding agent to a microvesicle surface marker. See, e.g., **FIGs. 2A, 2B, 21A**. The binding agent can be an aptamer that was identified to target the microvesicle surface marker using to the methods of the invention. The isolated vesicles is assessed for additional biomarkers such as surface content or payload,

which can be contemporaneous to detection of the aptamer-specific target or the assessment of additional biomarkers can be before or subsequent to aptamer-specific target detection.

[00133] A biosignature can be detected qualitatively or quantitatively by detecting a presence, level or concentration of a circulating biomarker, e.g., a microRNA, protein, vesicle or other biomarker, as disclosed herein. These biosignature components can be detected using a number of techniques known to those of skill in the art. For example, a biomarker can be detected by microarray analysis, polymerase chain reaction (PCR) (including PCR-based methods such as real time polymerase chain reaction (RT-PCR), quantitative real time polymerase chain reaction (Q-PCR/qPCR) and the like), hybridization with allele-specific probes, enzymatic mutation detection, ligation chain reaction (LCR), oligonucleotide ligation assay (OLA), flow-cytometric heteroduplex analysis, chemical cleavage of mismatches, mass spectrometry, nucleic acid sequencing, single strand conformation polymorphism (SSCP), denaturing gradient gel electrophoresis (DGGE), temperature gradient gel electrophoresis (TGGE), restriction fragment polymorphisms, serial analysis of gene expression (SAGE), or combinations thereof. A biomarker, such as a nucleic acid, can be amplified prior to detection. A biomarker can also be detected by immunoassay, immunoblot, immunoprecipitation, enzyme-linked immunosorbent assay (ELISA; EIA), radioimmunoassay (RIA), flow cytometry, or electron microscopy (EM).

[00134] Biosignatures can be detected using aptamers of the invention that function as either as capture agents and detection agents, as described herein. A capture agent can comprise an antibody, aptamer or other entity which recognizes a biomarker and can be used for capturing the biomarker. Biomarkers that can be captured include circulating biomarkers, e.g., a protein, nucleic acid, lipid or biological complex in solution in a bodily fluid. Similarly, the capture agent can be used for capturing a vesicle. A detection agent can comprise an antibody or other entity which recognizes a biomarker and can be used for detecting the biomarker vesicle, or which recognizes a vesicle and is useful for detecting a vesicle. In some embodiments, the detection agent is labeled and the label is detected, thereby detecting the biomarker or vesicle. The detection agent can be a binding agent, e.g., an antibody or aptamer. In other embodiments, the detection agent comprises a small molecule such as a membrane protein labeling agent. See, e.g., the membrane protein labeling agents disclosed in Alroy et al., US. Patent Publication US 2005/0158708. In an embodiment, vesicles are isolated or captured as described herein, and one or more membrane protein labeling agent is used to detect the vesicles. In many cases, the antigen or other vesicle-moiety that is recognized by the capture and detection agents are interchangeable.

[00135] In a non-limiting embodiment, a vesicle having a cell-of-origin specific antigen on its surface and a cancer-specific antigen on its surface, is captured using a binding agent that is specific to a cells-specific antigen, e.g., by tethering the capture antibody or aptamer to a substrate, and then the vesicle is detected using a binding agent to a disease-specific antigen, e.g., by labeling the binding agent used for detection with a fluorescent dye and detecting the fluorescent radiation emitted by the dye.

[00136] It will be apparent to one of skill in the art that where the target molecule for a binding agent (such as an aptamer of the invention) is informative as to assessing a condition or disease, the same

binding agent can be used to both capture a component comprising the target molecule (e.g., microvesicle surface antigen of interest) and also be modified to comprise a detectable label so as to detect the target molecule, e.g., binding $agent_1$-antigen-binding $agent_2$*, wherein the * signifies a detectable label; binding $agent_1$ and binding $agent_2$ may be the same binding agent or a different binding agent (e.g., same aptamer or different aptamer). In addition, binding $agent_1$ and binding $agent_2$ can be selected from wholly different categories of binding agents (e.g., antibody, aptamer, synthetic antibody, peptide-nucleic acid molecule, or any molecule that is configured to specifically bind to or associate with its target molecule). Such binding molecules can be selected solely based on their binding specificity for a target molecule. Examples of additional biomarkers that can be incorporated into the methods and compositions of the invention are known in the art, such as those disclosed in International Patent Publication Nos. WO/2012/174282 (Int'l Appl. PCT/US2012/042519 filed June 14, 2012) and WO/2013/020995 (Int'l Appl. PCT/US2012/050030 filed August 8, 2013). The detectable signal can itself be associated with a nucleic acid molecule that hybridizes with a stretch of nucleic acids present in each oligonucleotide comprising a probing library. The stretch can be the same or different as to one or more oligonucleotides in a library. The detectable signal can comprise fluorescence agents, including color-coded barcodes which are known, such as in U.S. Patent Application Pub. No. 20140371088, 2013017837, and 20120258870.

[00137] Techniques of detecting biomarkers or capturing sample components using an aptamer of the invention include the use of a planar substrate such as an array (e.g., biochip or microarray), with molecules immobilized to the substrate as capture agents that facilitate the detection of a particular biosignature. The array can be provided as part of a kit for assaying one or more biomarkers. Additional examples of binding agents described above and useful in the compositions and methods of the invention are disclosed in International Patent Publication No. WO/2011/127219, entitled "Circulating Biomarkers for Disease" and filed April 6, 2011, which application is incorporated by reference in its entirety herein. Aptamers of the invention can be included in an array for detection and diagnosis of diseases including presymptomatic diseases. In some embodiments, an array comprises a custom array comprising biomolecules selected to specifically identify biomarkers of interest. Customized arrays can be modified to detect biomarkers that increase statistical performance, e.g., additional biomolecules that identifies a biosignature which lead to improved cross-validated error rates in multivariate prediction models (e.g., logistic regression, discriminant analysis, or regression tree models). In some embodiments, customized array(s) are constructed to study the biology of a disease, condition or syndrome and profile biosignatures in defined physiological states. Markers for inclusion on the customized array be chosen based upon statistical criteria, e.g., having a desired level of statistical significance in differentiating between phenotypes or physiological states. In some embodiments, standard significance of p-value = 0.05 is chosen to exclude or include biomolecules on the microarray. The p-values can be corrected for multiple comparisons. As an illustrative example, nucleic acids extracted from samples from a subject with or without a disease can be hybridized to a high density microarray that binds to thousands of gene

sequences. Nucleic acids whose levels are significantly different between the samples with or without the disease can be selected as biomarkers to distinguish samples as having the disease or not. A customized array can be constructed to detect the selected biomarkers. In some embodiments, customized arrays comprise low density microarrays, which refer to arrays with lower number of addressable binding agents, e.g., tens or hundreds instead of thousands. Low density arrays can be formed on a substrate. In some embodiments, customizable low density arrays use PCR amplification in plate wells, e.g., TaqMan® Gene Expression Assays (Applied Biosystems by Life Technologies Corporation, Carlsbad, CA).

[00138] An aptamer of the invention or other useful binding agent may be linked directly or indirectly to a solid surface or substrate. See, e.g., **FIGs. 2A-2B, 9, 21A**. A solid surface or substrate can be any physically separable solid to which a binding agent can be directly or indirectly attached including, but not limited to, surfaces provided by microarrays and wells, particles such as beads, columns, optical fibers, wipes, glass and modified or functionalized glass, quartz, mica, diazotized membranes (paper or nylon), polyformaldehyde, cellulose, cellulose acetate, paper, ceramics, metals, metalloids, semiconductive materials, quantum dots, coated beads or particles, other chromatographic materials, magnetic particles; plastics (including acrylics, polystyrene, copolymers of styrene or other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon material, etc.), polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, plastics, ceramics, conducting polymers (including polymers such as polypyrole and polyindole); micro or nanostructured surfaces such as nucleic acid tiling arrays, nanotube, nanowire, or nanoparticulate decorated surfaces; or porous surfaces or gels such as methacrylates, acrylamides, sugar polymers, cellulose, silicates, or other fibrous or stranded polymers. In addition, as is known the art, the substrate may be coated using passive or chemically-derivatized coatings with any number of materials, including polymers, such as dextrans, acrylamides, gelatins or agarose. Such coatings can facilitate the use of the array with a biological sample.

[00139] As provided in the examples, below, an aptamer or other useful binding agent can be conjugated to a detectable entity or label.

[00140] Appropriate labels include without limitation a magnetic label, a fluorescent moiety, an enzyme, a chemiluminescent probe, a metal particle, a non-metal colloidal particle, a polymeric dye particle, a pigment molecule, a pigment particle, an electrochemically active species, semiconductor nanocrystal or other nanoparticles including quantum dots or gold particles, fluorophores, quantum dots, or radioactive labels. Protein labels include green fluorescent protein (GFP) and variants thereof (e.g., cyan fluorescent protein and yellow fluorescent protein); and luminescent proteins such as luciferase, as described below. Radioactive labels include without limitation radioisotopes (radionuclides), such as ^{3}H, ^{11}C, ^{14}C, ^{18}F, ^{32}P, ^{35}S, ^{64}Cu, ^{68}Ga, ^{86}Y, ^{99}Tc, ^{111}In, ^{123}I, ^{124}I, ^{125}I, ^{131}I, ^{133}Xe, ^{177}Lu, ^{211}At, or ^{213}Bi. Fluorescent labels include without limitation a rare earth chelate (e.g., europium chelate), rhodamine; fluorescein types including without limitation FITC, 5-carboxyfluorescein, 6-carboxy fluorescein; a rhodamine type including without limitation TAMRA; dansyl; Lissamine; cyanines; phycoerythrins; Texas Red; Cy3, Cy5, dapoxyl, NBD,

Cascade Yellow, dansyl, PyMPO, pyrene, 7-diethylaminocoumarin-3-carboxylic acid and other coumarin derivatives, Marina Blue™, Pacific Blue™, Cascade Blue™, 2-anthracenesulfonyl, PyMPO, 3,4,9,10-perylene-tetracarboxylic acid, 2,7-difluorofluorescein (Oregon Green™ 488-X), 5-carboxyfluorescein, Texas Red™-X, Alexa Fluor 430, 5-carboxytetramethylrhodamine (5-TAMRA), 6-carboxytetramethylrhodamine (6-TAMRA), BODIPY FL, bimane, and Alexa Fluor 350, 405, 488, 500, 514, 532, 546, 555, 568, 594, 610, 633, 647, 660, 680, 700, and 750, and derivatives thereof, among many others. See, e.g., "The Handbook--A Guide to Fluorescent Probes and Labeling Technologies," Tenth Edition, available on the internet at probes (dot) invitrogen (dot) com/handbook. The fluorescent label can be one or more of FAM, dRHO, 5-FAM, 6FAM, dR6G, JOE, HEX, VIC, TET, dTAMRA, TAMRA, NED, dROX, PET, BHQ, Gold540 and LIZ.

[00141] Using conventional techniques, an aptamer can be directly or indirectly labeled, e.g., the label is attached to the aptamer through biotin-streptavidin (e.g., synthesize a biotinylated aptamer, which is then capable of binding a streptavidin molecule that is itself conjugated to a detectable label; non-limiting example is streptavidin, phycoerythrin conjugated (SAPE)). Methods for chemical coupling using multiple step procedures include biotinylation, coupling of trinitrophenol (TNP) or digoxigenin using for example succinimide esters of these compounds. Biotinylation can be accomplished by, for example, the use of D-biotinyl-N-hydroxysuccinimide. Succinimide groups react effectively with amino groups at pH values above 7, and preferentially between about pH 8.0 and about pH 8.5. Alternatively, an aptamer is not labeled, but is later contacted with a second antibody that is labeled after the first antibody is bound to an antigen of interest.

[00142] Various enzyme-substrate labels may also be used in conjunction with a composition or method of the invention. Such enzyme-substrate labels are available commercially (e.g., U.S. Pat. No. 4,275,149). The enzyme generally catalyzes a chemical alteration of a chromogenic substrate that can be measured using various techniques. For example, the enzyme may catalyze a color change in a substrate, which can be measured spectrophotometrically. Alternatively, the enzyme may alter the fluorescence or chemiluminescence of the substrate. Examples of enzymatic labels include luciferases (e.g., firefly luciferase and bacterial luciferase; U.S. Pat. No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, malate dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRP), alkaline phosphatase (AP), β-galactosidase, glucoamylase, lysozyme, saccharide oxidases (e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like. Examples of enzyme-substrate combinations include, but are not limited to, horseradish peroxidase (HRP) with hydrogen peroxidase as a substrate, wherein the hydrogen peroxidase oxidizes a dye precursor (e.g., orthophenylene diamine (OPD) or 3,3',5,5'-tetramethylbenzidine hydrochloride (TMB)); alkaline phosphatase (AP) with para-nitrophenyl phosphate as chromogenic substrate; and β-D-galactosidase (β-D-Gal) with a chromogenic substrate (e.g., p-nitrophenyl- β-D-galactosidase) or fluorogenic substrate 4-methylumbelliferyl-β-D-galactosidase.

[00143] Aptamer(s) can be linked to a substrate such as a planar substrate. A planar array generally contains addressable locations (e.g., pads, addresses, or micro-locations) of biomolecules in an array format. The size of the array will depend on the composition and end use of the array. Arrays can be made containing from 2 different molecules to many thousands. Generally, the array comprises from two to as many as 100,000 or more molecules, depending on the end use of the array and the method of manufacture. A microarray for use with the invention comprises at least one biomolecule that identifies or captures a biomarker present in a biosignature of interest, e.g., a microRNA or other biomolecule or vesicle that makes up the biosignature. In some arrays, multiple substrates are used, either of different or identical compositions. Accordingly, planar arrays may comprise a plurality of smaller substrates.

[00144] The present invention can make use of many types of arrays for detecting a biomarker, e.g., a biomarker associated with a biosignature of interest. Useful arrays or microarrays include without limitation DNA microarrays, such as cDNA microarrays, oligonucleotide microarrays and SNP microarrays, microRNA arrays, protein microarrays, antibody microarrays, tissue microarrays, cellular microarrays (also called transfection microarrays), chemical compound microarrays, and carbohydrate arrays (glycoarrays). These arrays are described in more detail above. In some embodiments, microarrays comprise biochips that provide high-density immobilized arrays of recognition molecules (e.g., aptamers or antibodies), where biomarker binding is monitored indirectly (e.g., via fluorescence).

[00145] An array or microarray that can be used to detect one or more biomarkers of a biosignature and comprising one or more aptamers of the invention can be made according to the methods described in U.S. Pat. Nos. 6,329,209; 6,365,418; 6,406,921; 6,475,808; and 6,475,809, and U.S. Patent Application Ser. No. 10/884,269, each of which is herein incorporated by reference in its entirety. Custom arrays to detect specific selections of sets of biomarkers described herein can be made using the methods described in these patents. Commercially available microarrays can also be used to carry out the methods of the invention, including without limitation those from Affymetrix (Santa Clara, CA), Illumina (San Diego, CA), Agilent (Santa Clara, CA), Exiqon (Denmark), or Invitrogen (Carlsbad, CA). Custom and/or commercial arrays include arrays for detection proteins, nucleic acids, and other biological molecules and entities (e.g., cells, vesicles, virii) as described herein.

[00146] In some embodiments, multiple capture molecules are disposed on an array, e.g., proteins, peptides or additional nucleic acid molecules. In certain embodiments, the proteins are immobilized using methods and materials that minimize the denaturing of the proteins, that minimize alterations in the activity of the proteins, or that minimize interactions between the protein and the surface on which they are immobilized. The capture molecules can comprise one or more aptamer of the invention. In one embodiment, an array is constructed for the hybridization of a pool of aptamers. The array can then be used to identify pool members that bind a sample, thereby facilitating characterization of a phenotype. See **FIGs. 19B-19C** and related disclosure for further details.

[00147] Array surfaces useful may be of any desired shape, form, or size. Non-limiting examples of surfaces include chips, continuous surfaces, curved surfaces, flexible surfaces, films, plates, sheets, or

tubes. Surfaces can have areas ranging from approximately a square micron to approximately 500 cm^2. The area, length, and width of surfaces may be varied according to the requirements of the assay to be performed. Considerations may include, for example, ease of handling, limitations of the material(s) of which the surface is formed, requirements of detection systems, requirements of deposition systems (e.g., arrayers), or the like.

[00148] In certain embodiments, it is desirable to employ a physical means for separating groups or arrays of binding islands or immobilized biomolecules: such physical separation facilitates exposure of different groups or arrays to different solutions of interest. Therefore, in certain embodiments, arrays are situated within microwell plates having any number of wells. In such embodiments, the bottoms of the wells may serve as surfaces for the formation of arrays, or arrays may be formed on other surfaces and then placed into wells. In certain embodiments, such as where a surface without wells is used, binding islands may be formed or molecules may be immobilized on a surface and a gasket having holes spatially arranged so that they correspond to the islands or biomolecules may be placed on the surface. Such a gasket is preferably liquid tight. A gasket may be placed on a surface at any time during the process of making the array and may be removed if separation of groups or arrays is no longer desired.

[00149] In some embodiments, the immobilized molecules can bind to one or more biomarkers or vesicles present in a biological sample contacting the immobilized molecules. In some embodiments, the immobilized molecules modify or are modified by molecules present in the one or more vesicles contacting the immobilized molecules. Contacting the sample typically comprises overlaying the sample upon the array.

[00150] Modifications or binding of molecules in solution or immobilized on an array can be detected using detection techniques known in the art. Examples of such techniques include immunological techniques such as competitive binding assays and sandwich assays; fluorescence detection using instruments such as confocal scanners, confocal microscopes, or CCD-based systems and techniques such as fluorescence, fluorescence polarization (FP), fluorescence resonant energy transfer (FRET), total internal reflection fluorescence (TIRF), fluorescence correlation spectroscopy (FCS); colorimetric/spectrometric techniques; surface plasmon resonance, by which changes in mass of materials adsorbed at surfaces are measured; techniques using radioisotopes, including conventional radioisotope binding and scintillation proximity assays (SPA); mass spectroscopy, such as matrix-assisted laser desorption/ionization mass spectroscopy (MALDI) and MALDI-time of flight (TOF) mass spectroscopy; ellipsometry, which is an optical method of measuring thickness of protein films; quartz crystal microbalance (QCM), a very sensitive method for measuring mass of materials adsorbing to surfaces; scanning probe microscopies, such as atomic force microscopy (AFM), scanning force microscopy (SFM) or scanning electron microscopy (SEM); and techniques such as electrochemical, impedance, acoustic, microwave, and IR/Raman detection. See, e.g., *Mere L, et al.,"Miniaturized FRET assays and microfluidics: key components for ultra-high-throughput screening," Drug Discovery Today 4(8):363-369 (1999),* and references cited therein*; Lakowicz J R, Principles of Fluorescence Spectroscopy, 2nd Edition,*

Plenum Press (1999), or Jain KK: Integrative Omics, Pharmacoproteomics, and Human Body Fluids. In: Thongboonkerd V, ed., ed. Proteomics of Human Body Fluids: Principles, Methods and Applications. Volume 1: Totowa, N.J.: Humana Press, 2007, each of which is herein incorporated by reference in its entirety.

[00151] Microarray technology can be combined with mass spectroscopy (MS) analysis and other tools. Electrospray interface to a mass spectrometer can be integrated with a capillary in a microfluidics device. For example, one commercially available system contains eTag reporters that are fluorescent labels with unique and well-defined electrophoretic mobilities; each label is coupled to biological or chemical probes via cleavable linkages. The distinct mobility address of each eTag reporter allows mixtures of these tags to be rapidly deconvoluted and quantitated by capillary electrophoresis. This system allows concurrent gene expression, protein expression, and protein function analyses from the same sample *Jain KK: Integrative Omics, Pharmacoproteomics, and Human Body Fluids. In: Thongboonkerd V, ed., ed. Proteomics of Human Body Fluids: Principles, Methods and Applications. Volume 1: Totowa, N.J.: Humana Press, 2007, which* is herein incorporated by reference in its entirety.

[00152] A biochip can include components for a microfluidic or nanofluidic assay. A microfluidic device can be used for isolating or analyzing biomarkers, such as determining a biosignature. Microfluidic systems allow for the miniaturization and compartmentalization of one or more processes for isolating, capturing or detecting a vesicle, detecting a microRNA, detecting a circulating biomarker, detecting a biosignature, and other processes. The microfluidic devices can use one or more detection reagents in at least one aspect of the system, and such a detection reagent can be used to detect one or more biomarkers. In one embodiment, the device detects a biomarker on an isolated or bound vesicle. Various probes, antibodies, proteins, or other binding agents can be used to detect a biomarker within the microfluidic system. The detection agents may be immobilized in different compartments of the microfluidic device or be entered into a hybridization or detection reaction through various channels of the device.

[00153] A vesicle in a microfluidic device can be lysed and its contents detected within the microfluidic device, such as proteins or nucleic acids, e.g., DNA or RNA such as miRNA or mRNA. The nucleic acid may be amplified prior to detection, or directly detected, within the microfluidic device. Thus microfluidic system can also be used for multiplexing detection of various biomarkers. In an embodiment, vesicles are captured within the microfluidic device, the captured vesicles are lysed, and a biosignature of microRNA from the vesicle payload is determined. The biosignature can further comprise the capture agent used to capture the vesicle.

[00154] Novel nanofabrication techniques are opening up the possibilities for biosensing applications that rely on fabrication of high-density, precision arrays, e.g., nucleotide-based chips and protein arrays otherwise known as heterogeneous nanoarrays. Nanofluidics allows a further reduction in the quantity of fluid analyte in a microchip to nanoliter levels, and the chips used here are referred to as nanochips. See, e.g., *Unger M et al., Biotechniques 1999; 27(5):1008-14, Kartalov EP et al., Biotechniques 2006; 40(1):85-90*, each of which are herein incorporated by reference in their entireties. Commercially

available nanochips currently provide simple one step assays such as total cholesterol, total protein or glucose assays that can be run by combining sample and reagents, mixing and monitoring of the reaction. Gel-free analytical approaches based on liquid chromatography (LC) and nanoLC separations (*Cutillas et al. Proteomics, 2005;5:101-112 and Cutillas et al., Mol Cell Proteomics 2005;4:1038-1051*, each of which is herein incorporated by reference in its entirety) can be used in combination with the nanochips.

[00155] An array suitable for identifying a disease, condition, syndrome or physiological status can be included in a kit. A kit can include, an aptamer of the invention, including as non-limiting examples, one or more reagents useful for preparing molecules for immobilization onto binding islands or areas of an array, reagents useful for detecting binding of a vesicle to immobilized molecules, and instructions for use.

[00156] Further provided herein is a rapid detection device that facilitates the detection of a particular biosignature in a biological sample. The device can integrate biological sample preparation with polymerase chain reaction (PCR) on a chip. The device can facilitate the detection of a particular biosignature of a vesicle in a biological sample, and an example is provided as described in *Pipper et al., Angewandte Chemie, 47(21), p. 3900-3904 (2008)*, which is herein incorporated by reference in its entirety. A biosignature can be incorporated using micro-/nano-electrochemical system (MEMS/NEMS) sensors and oral fluid for diagnostic applications as described in *Li et al., Adv Dent Res 18(1): 3-5 (2005)*, which is herein incorporated by reference in its entirety.

[00157] ***Particle arrays***

[00158] As an alternative to planar arrays, assays using particles, such as bead based assays are also capable of use with an aptamer of the invention. Aptamers are easily conjugated with commercially available beads. See, e.g., Srinivas *et al.* Anal. Chem. 2011 Oct. 21, *Aptamer functionalized Microgel Particles for Protein Detection*; See also, review article on aptamers as therapeutic and diagnostic agents, Brody and Gold, Rev. Mol. Biotech. 2000, 74:5-13.

[00159] Multiparametric assays or other high throughput detection assays using bead coatings with cognate ligands and reporter molecules with specific activities consistent with high sensitivity automation can be used. In a bead based assay system, a binding agent for a biomarker or vesicle, such as a capture agent (e.g. capture antibody), can be immobilized on an addressable microsphere. Each binding agent for each individual binding assay can be coupled to a distinct type of microsphere (i.e., microbead) and the assay reaction takes place on the surface of the microsphere, such as depicted in **FIG. 2B**. A binding agent for a vesicle can be a capture antibody coupled to a bead. Dyed microspheres with discrete fluorescence intensities are loaded separately with their appropriate binding agent or capture probes. The different bead sets carrying different binding agents can be pooled as desired to generate custom bead arrays. Bead arrays are then incubated with the sample in a single reaction vessel to perform the assay.

[00160] Bead-based assays can also be used with one or more aptamers of the invention. A bead substrate can provide a platform for attaching one or more binding agents, including aptamer(s). For multiplexing, multiple different bead sets (e.g., Illumina, Luminex) can have different binding agents (specific to

different target molecules). For example, a bead can be conjugated to an aptamer of the invention used to detect the presence (quantitatively or qualitatively) of an antigen of interest, or it can also be used to isolate a component present in a selected biological sample (e.g., cell, cell-fragment or vesicle comprising the target molecule to which the aptamer is configured to bind or associate). Any molecule of organic origin can be successfully conjugated to a polystyrene bead through use of commercially available kits.

[00161] One or more aptamers of the invention can be used with any bead based substrate, including but not limited to magnetic capture method, fluorescence activated cell sorting (FACS) or laser cytometry. Magnetic capture methods can include, but are not limited to, the use of magnetically activated cell sorter (MACS) microbeads or magnetic columns. Examples of bead or particle based methods that can be modified to use an aptamer of the invention include methods and bead systems described in U.S. Patent Nos. 4,551,435, 4,795,698, 4,925,788, 5,108,933, 5,186,827, 5,200,084 or 5,158,871; 7,399,632; 8,124,015; 8,008,019; 7,955,802; 7,445,844; 7,274,316; 6,773,812; 6,623,526; 6,599,331; 6,057,107; 5,736,330; International Patent Publication No. WO/2012/174282; WO/1993/022684.

[00162] *Flow Cytometry*

[00163] Isolation or detection of circulating biomarkers, e.g., protein antigens, from a biological sample, or of the biomarker-comprising cells, cell fragments or vesicles may also be achieved using an aptamer of the invention in a cytometry process. As a non-limiting example, aptamers of the invention can be used in an assay comprising using a particle such as a bead or microsphere The invention provides aptamers as binding agents, which may be conjugated to the particle. Flow cytometry can be used for sorting microscopic particles suspended in a stream of fluid. As particles pass through they can be selectively charged and on their exit can be deflected into separate paths of flow. It is therefore possible to separate populations from an original mix, such as a biological sample, with a high degree of accuracy and speed. Flow cytometry allows simultaneous multiparametric analysis of the physical and/or chemical characteristics of single cells flowing through an optical/electronic detection apparatus. A beam of light, usually laser light, of a single frequency (color) is directed onto a hydrodynamically focused stream of fluid. A number of detectors are aimed at the point where the stream passes through the light beam; one in line with the light beam (Forward Scatter or FSC) and several perpendicular to it (Side Scatter or SSC) and one or more fluorescent detectors.

[00164] Each suspended particle passing through the beam scatters the light in some way, and fluorescent chemicals in the particle may be excited into emitting light at a lower frequency than the light source. This combination of scattered and fluorescent light is picked up by the detectors, and by analyzing fluctuations in brightness at each detector (one for each fluorescent emission peak), it is possible to deduce various facts about the physical and chemical structure of each individual particle. FSC correlates with the cell size and SSC depends on the inner complexity of the particle, such as shape of the nucleus, the amount and type of cytoplasmic granules or the membrane roughness. Some flow cytometers have eliminated the need for fluorescence and use only light scatter for measurement.

[00165] Flow cytometers can analyze several thousand particles every second in "real time" and can actively separate out and isolate particles having specified properties. They offer high-throughput automated quantification, and separation, of the set parameters for a high number of single cells during each analysis session. Flow cytometers can have multiple lasers and fluorescence detectors, allowing multiple labels to be used to more precisely specify a target population by their phenotype. Thus, a flow cytometer, such as a multicolor flow cytometer, can be used to detect one or more vesicles with multiple fluorescent labels or colors. In some embodiments, the flow cytometer can also sort or isolate different vesicle populations, such as by size or by different markers.

[00166] The flow cytometer may have one or more lasers, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more lasers. In some embodiments, the flow cytometer can detect more than one color or fluorescent label, such as at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 different colors or fluorescent labels. For example, the flow cytometer can have at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 fluorescence detectors.

[00167] Examples of commercially available flow cytometers that can be used to detect or analyze one or more vesicles, to sort or separate different populations of vesicles, include, but are not limited to the MoFlo™ XDP Cell Sorter (Beckman Coulter, Brea, CA), MoFlo™ Legacy Cell Sorter (Beckman Coulter, Brea, CA), BD FACSAria™ Cell Sorter (BD Biosciences, San Jose, CA), BD™ LSRII (BD Biosciences, San Jose, CA), and BD FACSCalibur™ (BD Biosciences, San Jose, CA). Use of multicolor or multi-fluor cytometers can be used in multiplex analysis of vesicles, as further described below. In some embodiments, the flow cytometer can sort, and thereby collect or sort more than one population of vesicles based one or more characteristics. For example, two populations of vesicles differ in size, such that the vesicles within each population have a similar size range and can be differentially detected or sorted. In another embodiment, two different populations of vesicles are differentially labeled.

[00168] The data resulting from flow-cytometers can be plotted in 1 dimension to produce histograms or seen in 2 dimensions as dot plots or in 3 dimensions with newer software. The regions on these plots can be sequentially separated by a series of subset extractions which are termed gates. Specific gating protocols exist for diagnostic and clinical purposes especially in relation to hematology. The plots are often made on logarithmic scales. Because different fluorescent dye's emission spectra overlap, signals at the detectors have to be compensated electronically as well as computationally. Fluorophores for labeling biomarkers may include those described in *Ormerod, Flow Cytometry 2nd ed., Springer-Verlag, New York (1999),* and in *Nida et al., Gynecologic Oncology 2005;4 889-894* which is incorporated herein by reference. In a multiplexed assay, including but not limited to a flow cytometry assay, one or more different target molecules can be assessed, wherein at least one of the target molecules is a microvesicle surface antigen assessed using an aptamer of the invention.

[00169] ***Microfluidics***

[00170] One or more aptamer of the invention can be disposed on any useful planar or bead substrate. In one aspect of the invention one or more aptamer of the invention is disposed on a microfluidic device,

thereby facilitating assessing, characterizing or isolating a component of a biological sample comprising a polypeptide antigen of interest or a functional fragment thereof. For example, the circulating antigen or a cell, cell fragment or cell-derived vesicles comprising the antigen can be assessed using one or more aptamers of the invention (alternatively along with additional binding agents). Microfluidic devices, which may also be referred to as "lab-on-a-chip" systems, biomedical micro-electro-mechanical systems (bioMEMs), or multicomponent integrated systems, can be used for isolating and analyzing a vesicle. Such systems miniaturize and compartmentalize processes that allow for binding of vesicles, detection of biosignatures, and other processes.

[00171] A microfluidic device can also be used for isolation of a vesicle through size differential or affinity selection. For example, a microfluidic device can use one more channels for isolating a vesicle from a biological sample based on size or by using one or more binding agents for isolating a vesicle from a biological sample. A biological sample can be introduced into one or more microfluidic channels, which selectively allows the passage of a vesicle. The selection can be based on a property of the vesicle, such as the size, shape, deformability, or biosignature of the vesicle.

[00172] In one embodiment, a heterogeneous population of vesicles can be introduced into a microfluidic device, and one or more different homogeneous populations of vesicles can be obtained. For example, different channels can have different size selections or binding agents to select for different vesicle populations. Thus, a microfluidic device can isolate a plurality of vesicles wherein at least a subset of the plurality of vesicles comprises a different biosignature from another subset of the plurality of vesicles. For example, the microfluidic device can isolate at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, or 100 different subsets of vesicles, wherein each subset of vesicles comprises a different biosignature.

[00173] In some embodiments, the microfluidic device can comprise one or more channels that permit further enrichment or selection of a vesicle. A population of vesicles that has been enriched after passage through a first channel can be introduced into a second channel, which allows the passage of the desired vesicle or vesicle population to be further enriched, such as through one or more binding agents present in the second channel.

[00174] Array-based assays and bead-based assays can be used with microfluidic device. For example, the binding agent can be coupled to beads and the binding reaction between the beads and vesicle can be performed in a microfluidic device. Multiplexing can also be performed using a microfluidic device. Different compartments can comprise different binding agents for different populations of vesicles, where each population is of a different cell-of-origin specific vesicle population. In one embodiment, each population has a different biosignature. The hybridization reaction between the microsphere and vesicle can be performed in a microfluidic device and the reaction mixture can be delivered to a detection device. The detection device, such as a dual or multiple laser detection system can be part of the microfluidic system and can use a laser to identify each bead or microsphere by its color-coding, and another laser can detect the hybridization signal associated with each bead.

[00175] Any appropriate microfluidic device can be used in the methods of the invention. Examples of microfluidic devices that may be used, or adapted for use with vesicles, include but are not limited to those described in U.S. Pat. Nos. 7,591,936, 7,581,429, 7,579,136, 7,575,722, 7,568,399, 7,552,741, 7,544,506, 7,541,578, 7,518,726, 7,488,596, 7,485,214, 7,467,928, 7,452,713, 7,452,509, 7,449,096, 7,431,887, 7,422,725, 7,422,669, 7,419,822, 7,419,639, 7,413,709, 7,411,184, 7,402,229, 7,390,463, 7,381,471, 7,357,864, 7,351,592, 7,351,380, 7,338,637, 7,329,391, 7,323,140, 7,261,824, 7,258,837, 7,253,003, 7,238,324, 7,238,255, 7,233,865, 7,229,538, 7,201,881, 7,195,986, 7,189,581, 7,189,580, 7,189,368, 7,141,978, 7,138,062, 7,135,147, 7,125,711, 7,118,910, 7,118,661, 7,640,947, 7,666,361, 7,704,735; and International Patent Publication WO 2010/072410; each of which patents or applications are incorporated herein by reference in their entirety. Another example for use with methods disclosed herein is described in *Chen et al., "Microfluidic isolation and transcriptome analysis of serum vesicles," Lab on a Chip, Dec. 8, 2009 DOI: 10.1039/b916199f.*

[00176] Other microfluidic devices for use with the invention include devices comprising elastomeric layers, valves and pumps, including without limitation those disclosed in U.S. Patent Nos. 5,376,252, 6,408,878, 6,645,432, 6,719,868, 6,793,753, 6,899,137, 6,929,030, 7,040,338, 7,118,910, 7,144,616, 7,216,671, 7,250,128, 7,494,555, 7,501,245, 7,601,270, 7,691,333, 7,754,010, 7,837,946; U.S. Patent Application Nos. 2003/0061687, 2005/0084421, 2005/0112882, 2005/0129581, 2005/0145496, 2005/0201901, 2005/0214173, 2005/0252773, 2006/0006067; and EP Patent Nos. 0527905 and 1065378; each of which application is herein incorporated by reference. In some instances, much or all of the devices are composed of elastomeric material. Certain devices are designed to conduct thermal cycling reactions (e.g., PCR) with devices that include one or more elastomeric valves to regulate solution flow through the device. The devices can comprise arrays of reaction sites thereby allowing a plurality of reactions to be performed. Thus, the devices can be used to assess circulating microRNAs in a multiplex fashion, including microRNAs isolated from vesicles. In an embodiment, the microfluidic device comprises (a) a first plurality of flow channels formed in an elastomeric substrate; (b) a second plurality of flow channels formed in the elastomeric substrate that intersect the first plurality of flow channels to define an array of reaction sites, each reaction site located at an intersection of one of the first and second flow channels; (c) a plurality of isolation valves disposed along the first and second plurality of flow channels and spaced between the reaction sites that can be actuated to isolate a solution within each of the reaction sites from solutions at other reaction sites, wherein the isolation valves comprise one or more control channels that each overlay and intersect one or more of the flow channels; and (d) means for simultaneously actuating the valves for isolating the reaction sites from each other. Various modifications to the basic structure of the device are envisioned within the scope of the invention. MicroRNAs can be detected in each of the reaction sites by using PCR methods. For example, the method can comprise the steps of the steps of: (i) providing a microfluidic device, the microfluidic device comprising: a first fluidic channel having a first end and a second end in fluid communication with each other through the channel; a plurality of flow channels, each flow channel terminating at a terminal wall; wherein each flow channel

branches from and is in fluid communication with the first fluidic channel, wherein an aqueous fluid that enters one of the flow channels from the first fluidic channel can flow out of the flow channel only through the first fluidic channel; and, an inlet in fluid communication with the first fluidic channel, the inlet for introducing a sample fluid; wherein each flow channel is associated with a valve that when closed isolates one end of the flow channel from the first fluidic channel, whereby an isolated reaction site is formed between the valve and the terminal wall; a control channel; wherein each the valve is a deflectable membrane which is deflected into the flow channel associated with the valve when an actuating force is applied to the control channel, thereby closing the valve; and wherein when the actuating force is applied to the control channel a valve in each of the flow channels is closed, so as to produce the isolated reaction site in each flow channel; (ii) introducing the sample fluid into the inlet, the sample fluid filling the flow channels; (iii) actuating the valve to separate the sample fluid into the separate portions within the flow channels; (iv) amplifying the nucleic acid in the sample fluid; (v) analyzing the portions of the sample fluid to determine whether the amplifying produced the reaction. The sample fluid can contain an amplifiable nucleic acid target, e.g., a microRNA, and the conditions can be polymerase chain reaction (PCR) conditions, so that the reaction results in a PCR product being formed.

[00177] The microfluidic device can have one or more binding agents attached to a surface in a channel, or present in a channel. For example, the microchannel can have one or more capture agents, such as a capture agent for a tissue related antigen in **Table 4**, one or more general microvesicle antigen in **Table 3** or a cell-of-origin or cancer related antigen in **Table 4**, including without limitation EpCam, CD9, CD63, CD81, B7H3, ICAM, STEAP, KLK2, SSX2, SSX4, PBP, SPDEF, and EGFR. The capture agent may be an aptamer selected by the methods of the invention. The surface of the channel can also be contacted with a blocking aptamer. In one embodiment, a microchannel surface is treated with avidin and a capture agent, such as an antibody, that is biotinylated can be injected into the channel to bind the avidin. In other embodiments, the capture agents are present in chambers or other components of a microfluidic device. The capture agents can also be attached to beads that can be manipulated to move through the microfluidic channels. In one embodiment, the capture agents are attached to magnetic beads. The beads can be manipulated using magnets.

[00178] A biological sample can be flowed into the microfluidic device, or a microchannel, at rates such as at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, or 50 μl per minute, such as between about 1-50, 5-40, 5-30, 3-20 or 5-15 μl per minute. One or more vesicles can be captured and directly detected in the microfluidic device. Alternatively, the captured vesicle may be released and exit the microfluidic device prior to analysis. In another embodiment, one or more captured vesicles are lysed in the microchannel and the lysate can be analyzed, e.g., to examine payload within the vesicles. Lysis buffer can be flowed through the channel and lyse the captured vesicles. For example, the lysis buffer can be flowed into the device or microchannel at rates such as at least about a, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 26, 27, 28, 29, 30, 35, 40, 45, or 50 μl per minute, such as between about 1-50, 5-40, 10-30, 5-30 or 10-35 μl per minute. The lysate can be collected

and analyzed, such as performing RT-PCR, PCR, mass spectrometry, Western blotting, or other assays, to detect one or more biomarkers of the vesicle.

Phenotypes

[00179] Disclosed herein are products and processes for characterizing a phenotype using the methods and compositions of the invention. The term "phenotype" as used herein can mean any trait or characteristic that is attributed to a biomarker profile that is identified using in part or in whole the compositions and/or methods of the invention. For example, a phenotype can be a diagnostic, prognostic or theranostic determination based on a characterized biomarker profile for a sample obtained from a subject. A phenotype can be any observable characteristic or trait of, such as a disease or condition, a stage of a disease or condition, susceptibility to a disease or condition, prognosis of a disease stage or condition, a physiological state, or response / potential response to therapeutics. A phenotype can result from a subject's genetic makeup as well as the influence of environmental factors and the interactions between the two, as well as from epigenetic modifications to nucleic acid sequences.

[00180] A phenotype in a subject can be characterized by obtaining a biological sample from a subject and analyzing the sample using the compositions and/or methods of the invention. For example, characterizing a phenotype for a subject or individual can include detecting a disease or condition (including pre-symptomatic early stage detecting), determining a prognosis, diagnosis, or theranosis of a disease or condition, or determining the stage or progression of a disease or condition. Characterizing a phenotype can include identifying appropriate treatments or treatment efficacy for specific diseases, conditions, disease stages and condition stages, predictions and likelihood analysis of disease progression, particularly disease recurrence, metastatic spread or disease relapse. A phenotype can also be a clinically distinct type or subtype of a condition or disease, such as a cancer or tumor. Phenotype determination can also be a determination of a physiological condition, or an assessment of organ distress or organ rejection, such as post-transplantation. The compositions and methods described herein allow assessment of a subject on an individual basis, which can provide benefits of more efficient and economical decisions in treatment.

[00181] In an aspect, the invention relates to the analysis of biomarkers such as microvesicles to provide a diagnosis, prognosis, and/or theranosis of a disease or condition. Theranostics includes diagnostic testing that provides the ability to affect therapy or treatment of a disease or disease state. Theranostics testing provides a theranosis in a similar manner that diagnostics or prognostic testing provides a diagnosis or prognosis, respectively. As used herein, theranostics encompasses any desired form of therapy related testing, including predictive medicine, personalized medicine, integrated medicine, pharmacodiagnostics and Dx/Rx partnering. Therapy related tests can be used to predict and assess drug response in individual subjects, i.e., to provide personalized medicine. Predicting a drug response can be determining whether a subject is a likely responder or a likely non-responder to a candidate therapeutic agent, e.g., before the subject has been exposed or otherwise treated with the treatment. Assessing a drug response can be monitoring a response to a drug, e.g., monitoring the subject's improvement or lack thereof over a time course after initiating the treatment. Therapy related tests are useful to select a subject for treatment who

is particularly likely to benefit from the treatment or to provide an early and objective indication of treatment efficacy in an individual subject. Thus, analysis using the compositions and methods of the invention may indicate that treatment should be altered to select a more promising treatment, thereby avoiding the great expense of delaying beneficial treatment and avoiding the financial and morbidity costs of administering an ineffective drug(s).

[00182] Thus, the compositions and methods of the invention may help predict whether a subject is likely to respond to a treatment for a disease or disorder. Characterizating a phenotype includes predicting the responder / non-responder status of the subject, wherein a responder responds to a treatment for a disease and a non-responder does not respond to the treatment. Biomarkers such as microvesicles can be analyzed in the subject and compared against that of previous subjects that were known to respond or not to a treatment. If the biomarker profile in the subject more closely aligns with that of previous subjects that were known to respond to the treatment, the subject can be characterized, or predicted, as a responder to the treatment. Similarly, if the biomarker profile in the subject more closely aligns with that of previous subjects that did not respond to the treatment, the subject can be characterized, or predicted as a non-responder to the treatment. The treatment can be for any appropriate disease, disorder or other condition, including without limitation those disclosed herein.

[00183] In some embodiments, the phenotype comprises a disease or condition such as those listed in **Tables 1** or **16**. For example, the phenotype can comprise detecting the presence of or likelihood of developing a tumor, neoplasm, or cancer, or characterizing the tumor, neoplasm, or cancer (e.g., stage, grade, aggressiveness, likelihood of metastatis or recurrence, etc). Cancers that can be detected or assessed by methods or compositions described herein include, but are not limited to, breast cancer, ovarian cancer, lung cancer, colon cancer, hyperplastic polyp, adenoma, colorectal cancer, high grade dysplasia, low grade dysplasia, prostatic hyperplasia, prostate cancer, melanoma, pancreatic cancer, brain cancer (such as a glioblastoma), hematological malignancy, hepatocellular carcinoma, cervical cancer, endometrial cancer, head and neck cancer, esophageal cancer, gastrointestinal stromal tumor (GIST), renal cell carcinoma (RCC) or gastric cancer. The colorectal cancer can be CRC Dukes B or Dukes C-D. The hematological malignancy can be B-Cell Chronic Lymphocytic Leukemia, B-Cell Lymphoma-DLBCL, B-Cell Lymphoma-DLBCL-germinal center-like, B-Cell Lymphoma-DLBCL-activated B-cell-like, and Burkitt's lymphoma.

[00184] The phenotype can be a premalignant condition, such as actinic keratosis, atrophic gastritis, leukoplakia, erythroplasia, Lymphomatoid Granulomatosis, preleukemia, fibrosis, cervical dysplasia, uterine cervical dysplasia, xeroderma pigmentosum, Barrett's Esophagus, colorectal polyp, or other abnormal tissue growth or lesion that is likely to develop into a malignant tumor. Transformative viral infections such as HIV and HPV also present phenotypes that can be assessed according to the invention.

[00185] A cancer characterized by the methods of the invention can comprise, without limitation, a carcinoma, a sarcoma, a lymphoma or leukemia, a germ cell tumor, a blastoma, or other cancers. Carcinomas include without limitation epithelial neoplasms, squamous cell neoplasms squamous cell

carcinoma, basal cell neoplasms basal cell carcinoma, transitional cell papillomas and carcinomas, adenomas and adenocarcinomas (glands), adenoma, adenocarcinoma, linitis plastica insulinoma, glucagonoma, gastrinoma, vipoma, cholangiocarcinoma, hepatocellular carcinoma, adenoid cystic carcinoma, carcinoid tumor of appendix, prolactinoma, oncocytoma, hurthle cell adenoma, renal cell carcinoma, grawitz tumor, multiple endocrine adenomas, endometrioid adenoma, adnexal and skin appendage neoplasms, mucoepidermoid neoplasms, cystic, mucinous and serous neoplasms, cystadenoma, pseudomyxoma peritonei, ductal, lobular and medullary neoplasms, acinar cell neoplasms, complex epithelial neoplasms, warthin's tumor, thymoma, specialized gonadal neoplasms, sex cord stromal tumor, thecoma, granulosa cell tumor, arrhenoblastoma, sertoli leydig cell tumor, glomus tumors, paraganglioma, pheochromocytoma, glomus tumor, nevi and melanomas, melanocytic nevus, malignant melanoma, melanoma, nodular melanoma, dysplastic nevus, lentigo maligna melanoma, superficial spreading melanoma, and malignant acral lentiginous melanoma. Sarcoma includes without limitation Askin's tumor, botryodies, chondrosarcoma, Ewing's sarcoma, malignant hemangio endothelioma, malignant schwannoma, osteosarcoma, soft tissue sarcomas including: alveolar soft part sarcoma, angiosarcoma, cystosarcoma phyllodes, dermatofibrosarcoma, desmoid tumor, desmoplastic small round cell tumor, epithelioid sarcoma, extraskeletal chondrosarcoma, extraskeletal osteosarcoma, fibrosarcoma, hemangiopericytoma, hemangiosarcoma, kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, lymphosarcoma, malignant fibrous histiocytoma, neurofibrosarcoma, rhabdomyosarcoma, and synovialsarcoma. Lymphoma and leukemia include without limitation chronic lymphocytic leukemia/small lymphocytic lymphoma, B-cell prolymphocytic leukemia, lymphoplasmacytic lymphoma (such as waldenström macroglobulinemia), splenic marginal zone lymphoma, plasma cell myeloma, plasmacytoma, monoclonal immunoglobulin deposition diseases, heavy chain diseases, extranodal marginal zone B cell lymphoma, also called malt lymphoma, nodal marginal zone B cell lymphoma (nmzl), follicular lymphoma, mantle cell lymphoma, diffuse large B cell lymphoma, mediastinal (thymic) large B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, burkitt lymphoma/leukemia, T cell prolymphocytic leukemia, T cell large granular lymphocytic leukemia, aggressive NK cell leukemia, adult T cell leukemia/lymphoma, extranodal NK/T cell lymphoma, nasal type, enteropathy-type T cell lymphoma, hepatosplenic T cell lymphoma, blastic NK cell lymphoma, mycosis fungoides / sezary syndrome, primary cutaneous CD30-positive T cell lymphoproliferative disorders, primary cutaneous anaplastic large cell lymphoma, lymphomatoid papulosis, angioimmunoblastic T cell lymphoma, peripheral T cell lymphoma, unspecified, anaplastic large cell lymphoma, classical hodgkin lymphomas (nodular sclerosis, mixed cellularity, lymphocyte-rich, lymphocyte depleted or not depleted), and nodular lymphocyte-predominant hodgkin lymphoma. Germ cell tumors include without limitation germinoma, dysgerminoma, seminoma, nongerminomatous germ cell tumor, embryonal carcinoma, endodermal sinus turmor, choriocarcinoma, teratoma, polyembryoma, and gonadoblastoma. Blastoma includes without limitation nephroblastoma, medulloblastoma, and retinoblastoma. Other cancers include without limitation labial carcinoma, larynx carcinoma,

hypopharynx carcinoma, tongue carcinoma, salivary gland carcinoma, gastric carcinoma, adenocarcinoma, thyroid cancer (medullary and papillary thyroid carcinoma), renal carcinoma, kidney parenchyma carcinoma, cervix carcinoma, uterine corpus carcinoma, endometrium carcinoma, chorion carcinoma, testis carcinoma, urinary carcinoma, melanoma, brain tumors such as glioblastoma, astrocytoma, meningioma, medulloblastoma and peripheral neuroectodermal tumors, gall bladder carcinoma, bronchial carcinoma, multiple myeloma, basalioma, teratoma, retinoblastoma, choroidea melanoma, seminoma, rhabdomyosarcoma, craniopharyngeoma, osteosarcoma, chondrosarcoma, myosarcoma, liposarcoma, fibrosarcoma, Ewing sarcoma, and plasmocytoma.

[00186] In a further embodiment, the cancer under analysis may be a lung cancer including non-small cell lung cancer and small cell lung cancer (including small cell carcinoma (oat cell cancer), mixed small cell/large cell carcinoma, and combined small cell carcinoma), colon cancer, breast cancer, prostate cancer, liver cancer, pancreas cancer, brain cancer, kidney cancer, ovarian cancer, stomach cancer, skin cancer, bone cancer, gastric cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, hepatocellular carcinoma, papillary renal carcinoma, head and neck squamous cell carcinoma, leukemia, lymphoma, myeloma, or a solid tumor.

[00187] In embodiments, the cancer comprises an acute lymphoblastic leukemia; acute myeloid leukemia; adrenocortical carcinoma; AIDS-related cancers; AIDS-related lymphoma; anal cancer; appendix cancer; astrocytomas; atypical teratoid/rhabdoid tumor; basal cell carcinoma; bladder cancer; brain stem glioma; brain tumor (including brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, astrocytomas, craniopharyngioma, ependymoblastoma, ependymoma, medulloblastoma, medulloepithelioma, pineal parenchymal tumors of intermediate differentiation, supratentorial primitive neuroectodermal tumors and pineoblastoma); breast cancer; bronchial tumors; Burkitt lymphoma; cancer of unknown primary site; carcinoid tumor; carcinoma of unknown primary site; central nervous system atypical teratoid/rhabdoid tumor; central nervous system embryonal tumors; cervical cancer; childhood cancers; chordoma; chronic lymphocytic leukemia; chronic myelogenous leukemia; chronic myeloproliferative disorders; colon cancer; colorectal cancer; craniopharyngioma; cutaneous T-cell lymphoma; endocrine pancreas islet cell tumors; endometrial cancer; ependymoblastoma; ependymoma; esophageal cancer; esthesioneuroblastoma; Ewing sarcoma; extracranial germ cell tumor; extragonadal germ cell tumor; extrahepatic bile duct cancer; gallbladder cancer; gastric (stomach) cancer; gastrointestinal carcinoid tumor; gastrointestinal stromal cell tumor; gastrointestinal stromal tumor (GIST); gestational trophoblastic tumor; glioma; hairy cell leukemia; head and neck cancer; heart cancer; Hodgkin lymphoma; hypopharyngeal cancer; intraocular melanoma; islet cell tumors; Kaposi sarcoma; kidney cancer; Langerhans cell histiocytosis; laryngeal cancer; lip cancer; liver cancer; malignant fibrous histiocytoma bone cancer; medulloblastoma; medulloepithelioma; melanoma; Merkel cell carcinoma; Merkel cell skin carcinoma; mesothelioma; metastatic squamous neck cancer with occult primary; mouth cancer; multiple endocrine neoplasia syndromes; multiple myeloma; multiple myeloma/plasma cell neoplasm; mycosis fungoides; myelodysplastic syndromes;

myeloproliferative neoplasms; nasal cavity cancer; nasopharyngeal cancer; neuroblastoma; Non-Hodgkin lymphoma; nonmelanoma skin cancer; non-small cell lung cancer; oral cancer; oral cavity cancer; oropharyngeal cancer; osteosarcoma; other brain and spinal cord tumors; ovarian cancer; ovarian epithelial cancer; ovarian germ cell tumor; ovarian low malignant potential tumor; pancreatic cancer; papillomatosis; paranasal sinus cancer; parathyroid cancer; pelvic cancer; penile cancer; pharyngeal cancer; pineal parenchymal tumors of intermediate differentiation; pineoblastoma; pituitary tumor; plasma cell neoplasm/multiple myeloma; pleuropulmonary blastoma; primary central nervous system (CNS) lymphoma; primary hepatocellular liver cancer; prostate cancer; rectal cancer; renal cancer; renal cell (kidney) cancer; renal cell cancer; respiratory tract cancer; retinoblastoma; rhabdomyosarcoma; salivary gland cancer; Sézary syndrome; small cell lung cancer; small intestine cancer; soft tissue sarcoma; squamous cell carcinoma; squamous neck cancer; stomach (gastric) cancer; supratentorial primitive neuroectodermal tumors; T-cell lymphoma; testicular cancer; throat cancer; thymic carcinoma; thymoma; thyroid cancer; transitional cell cancer; transitional cell cancer of the renal pelvis and ureter; trophoblastic tumor; ureter cancer; urethral cancer; uterine cancer; uterine sarcoma; vaginal cancer; vulvar cancer; Waldenström macroglobulinemia; or Wilm's tumor. The methods of the invention can be used to characterize these and other cancers. Thus, characterizing a phenotype can be providing a diagnosis, prognosis or theranosis of one of the cancers disclosed herein.

[00188] In some embodiments, the cancer comprises an acute myeloid leukemia (AML), breast carcinoma, cholangiocarcinoma, colorectal adenocarcinoma, extrahepatic bile duct adenocarcinoma, female genital tract malignancy, gastric adenocarcinoma, gastroesophageal adenocarcinoma, gastrointestinal stromal tumors (GIST), glioblastoma, head and neck squamous carcinoma, leukemia, liver hepatocellular carcinoma, low grade glioma, lung bronchioloalveolar carcinoma (BAC), lung non-small cell lung cancer (NSCLC), lung small cell cancer (SCLC), lymphoma, male genital tract malignancy, malignant solitary fibrous tumor of the pleura (MSFT), melanoma, multiple myeloma, neuroendocrine tumor, nodal diffuse large B-cell lymphoma, non epithelial ovarian cancer (non-EOC), ovarian surface epithelial carcinoma, pancreatic adenocarcinoma, pituitary carcinomas, oligodendroglioma, prostatic adenocarcinoma, retroperitoneal or peritoneal carcinoma, retroperitoneal or peritoneal sarcoma, small intestinal malignancy, soft tissue tumor, thymic carcinoma, thyroid carcinoma, or uveal melanoma. The methods of the invention can be used to characterize these and other cancers. Thus, characterizing a phenotype can be providing a diagnosis, prognosis or theranosis of one of the cancers disclosed herein.

[00189] The phenotype can also be an inflammatory disease, immune disease, or autoimmune disease. For example, the disease may be inflammatory bowel disease (IBD), Crohn's disease (CD), ulcerative colitis (UC), pelvic inflammation, vasculitis, psoriasis, diabetes, autoimmune hepatitis, Multiple Sclerosis, Myasthenia Gravis, Type I diabetes, Rheumatoid Arthritis, Psoriasis, Systemic Lupus Erythematosis (SLE), Hashimoto's Thyroiditis, Grave's disease, Ankylosing Spondylitis Sjogrens Disease, CREST syndrome, Scleroderma, Rheumatic Disease, organ rejection, Primary Sclerosing Cholangitis, or sepsis.

[00190] The phenotype can also comprise a cardiovascular disease, such as atherosclerosis, congestive heart failure, vulnerable plaque, stroke, or ischemia. The cardiovascular disease or condition can be high blood pressure, stenosis, vessel occlusion or a thrombotic event.

[00191] The phenotype can also comprise a neurological disease, such as Multiple Sclerosis (MS), Parkinson's Disease (PD), Alzheimer's Disease (AD), schizophrenia, bipolar disorder, depression, autism, Prion Disease, Pick's disease, dementia, Huntington disease (HD), Down's syndrome, cerebrovascular disease, Rasmussen's encephalitis, viral meningitis, neurospsychiatric systemic lupus erythematosus (NPSLE), amyotrophic lateral sclerosis, Creutzfeldt-Jacob disease, Gerstmann-Straussler-Scheinker disease, transmissible spongiform encephalopathy, ischemic reperfusion damage (e.g. stroke), brain trauma, microbial infection, or chronic fatigue syndrome. The phenotype may also be a condition such as fibromyalgia, chronic neuropathic pain, or peripheral neuropathic pain.

[00192] The phenotype may also comprise an infectious disease, such as a bacterial, viral or yeast infection. For example, the disease or condition may be Whipple's Disease, Prion Disease, cirrhosis, methicillin-resistant staphylococcus aureus, HIV, hepatitis, syphilis, meningitis, malaria, tuberculosis, or influenza. Viral proteins, such as HIV or HCV-like particles can be assessed in a vesicle, to characterize a viral condition.

[00193] The phenotype can also comprise a perinatal or pregnancy related condition (e.g. preeclampsia or preterm birth), metabolic disease or condition, such as a metabolic disease or condition associated with iron metabolism. For example, hepcidin can be assayed in a vesicle to characterize an iron deficiency. The metabolic disease or condition can also be diabetes, inflammation, or a perinatal condition.

[00194] The compositions and methods of the invention can be used to characterize these and other diseases and disorders that can be assessed via biomarkers. Thus, characterizing a phenotype can be providing a diagnosis, prognosis or theranosis of one of the diseases and disorders disclosed herein.

Subject

[00195] One or more phenotypes of a subject can be determined by analyzing one or more vesicles, such as vesicles, in a biological sample obtained from the subject. A subject or patient can include, but is not limited to, mammals such as bovine, avian, canine, equine, feline, ovine, porcine, or primate animals (including humans and non-human primates). A subject can also include a mammal of importance due to being endangered, such as a Siberian tiger; or economic importance, such as an animal raised on a farm for consumption by humans, or an animal of social importance to humans, such as an animal kept as a pet or in a zoo. Examples of such animals include, but are not limited to, carnivores such as cats and dogs; swine including pigs, hogs and wild boars; ruminants or ungulates such as cattle, oxen, sheep, giraffes, deer, goats, bison, camels or horses. Also included are birds that are endangered or kept in zoos, as well as fowl and more particularly domesticated fowl, i.e. poultry, such as turkeys and chickens, ducks, geese, guinea fowl. Also included are domesticated swine and horses (including race horses). In addition, any animal species connected to commercial activities are also included such as those animals connected to

agriculture and aquaculture and other activities in which disease monitoring, diagnosis, and therapy selection are routine practice in husbandry for economic productivity and/or safety of the food chain.

[00196] The subject can have a pre-existing disease or condition, such as cancer. Alternatively, the subject may not have any known pre-existing condition. The subject may also be non-responsive to an existing or past treatment, such as a treatment for cancer.

Samples

[00197] A sample used and/or assessed via the compositions and methods of the invention includes any relevant biological sample that can be used for biomarker assessment, including without limitation sections of tissues such as biopsy or tissue removed during surgical or other procedures, bodily fluids, autopsy samples, frozen sections taken for histological purposes, and cell cultures. Such samples include blood and blood fractions or products (e.g., serum, buffy coat, plasma, platelets, red blood cells, and the like), sputum, malignant effusion, cheek cells tissue, cultured cells (e.g., primary cultures, explants, and transformed cells), stool, urine, other biological or bodily fluids (e.g., prostatic fluid, gastric fluid, intestinal fluid, renal fluid, lung fluid, cerebrospinal fluid, and the like), etc. The sample can comprise biological material that is a fresh frozen & formalin fixed paraffin embedded (FFPE) block, formalin-fixed paraffin embedded, or is within an RNA preservative + formalin fixative. More than one sample of more than one type can be used for each patient.

[00198] The sample used in the methods described herein can be a formalin fixed paraffin embedded (FFPE) sample. The FFPE sample can be one or more of fixed tissue, unstained slides, bone marrow core or clot, core needle biopsy, malignant fluids and fine needle aspirate (FNA). In an embodiment, the fixed tissue comprises a tumor containing formalin fixed paraffin embedded (FFPE) block from a surgery or biopsy. In another embodiment, the unstained slides comprise unstained, charged, unbaked slides from a paraffin block. In another embodiment, bone marrow core or clot comprises a decalcified core. A formalin fixed core and/or clot can be paraffin-embedded. In still another embodiment, the core needle biopsy comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more, e.g., 3-4, paraffin embedded biopsy samples. An 18 gauge needle biopsy can be used. The malignant fluid can comprise a sufficient volume of fresh pleural/ascitic fluid to produce a 5x5x2mm cell pellet. The fluid can be formalin fixed in a paraffin block. In an embodiment, the core needle biopsy comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more, e.g., 4-6, paraffin embedded aspirates.

[00199] A sample may be processed according to techniques understood by those in the art. A sample can be without limitation fresh, frozen or fixed cells or tissue. In some embodiments, a sample comprises formalin-fixed paraffin-embedded (FFPE) tissue, fresh tissue or fresh frozen (FF) tissue. A sample can comprise cultured cells, including primary or immortalized cell lines derived from a subject sample. A sample can also refer to an extract from a sample from a subject. For example, a sample can comprise DNA, RNA or protein extracted from a tissue or a bodily fluid. Many techniques and commercial kits are available for such purposes. The fresh sample from the individual can be treated with an agent to preserve RNA prior to further processing, e.g., cell lysis and extraction. Samples can include frozen samples

collected for other purposes. Samples can be associated with relevant information such as age, gender, and clinical symptoms present in the subject; source of the sample; and methods of collection and storage of the sample. A sample is typically obtained from a subject.

[00200] A biopsy comprises the process of removing a tissue sample for diagnostic or prognostic evaluation, and to the tissue specimen itself. Any biopsy technique known in the art can be applied to the molecular profiling methods of the present invention. The biopsy technique applied can depend on the tissue type to be evaluated (e.g., colon, prostate, kidney, bladder, lymph node, liver, bone marrow, blood cell, lung, breast, etc.), the size and type of the tumor (e.g., solid or suspended, blood or ascites), among other factors. Representative biopsy techniques include, but are not limited to, excisional biopsy, incisional biopsy, needle biopsy, surgical biopsy, and bone marrow biopsy. An "excisional biopsy" refers to the removal of an entire tumor mass with a small margin of normal tissue surrounding it. An "incisional biopsy" refers to the removal of a wedge of tissue that includes a cross-sectional diameter of the tumor. Molecular profiling can use a "core-needle biopsy" of the tumor mass, or a "fine-needle aspiration biopsy" which generally obtains a suspension of cells from within the tumor mass. Biopsy techniques are discussed, for example, in Harrison's Principles of Internal Medicine, Kasper, et al., eds., 16th ed., 2005, Chapter 70, and throughout Part V.

[00201] Standard molecular biology techniques known in the art and not specifically described are generally followed as in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York (1989), and as in Ausubel et al., Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, Md. (1989) and as in Perbal, A Practical Guide to Molecular Cloning, John Wiley & Sons, New York (1988), and as in Watson et al., Recombinant DNA, Scientific American Books, New York and in Birren et al (eds) Genome Analysis: A Laboratory Manual Series, Vols. 1-4 Cold Spring Harbor Laboratory Press, New York (1998) and methodology as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057 and incorporated herein by reference. Polymerase chain reaction (PCR) can be carried out generally as in PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego, Calif. (1990).

[00202] The biological sample assessed using the compositions and methods of the invention can be any useful bodily or biological fluid, including but not limited to peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen (including prostatic fluid), Cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates or other lavage fluids, cells, cell culture, or a cell culture supernatant. A biological sample may also include the blastocyl cavity, umbilical cord blood, or maternal circulation which may be of fetal or maternal origin. The biological sample may also be a cell culture, tissue sample or biopsy from which vesicles and other circulating biomarkers may be obtained. For example, cells of

interest can be cultured and vesicles isolated from the culture. In various embodiments, biomarkers or more particularly biosignatures disclosed herein can be assessed directly from such biological samples (e.g., identification of presence or levels of nucleic acid or polypeptide biomarkers or functional fragments thereof) using various methods, such as extraction of nucleic acid molecules from blood, plasma, serum or any of the foregoing biological samples, use of protein or antibody arrays to identify polypeptide (or functional fragment) biomarker(s), as well as other array, sequencing, PCR and proteomic techniques known in the art for identification and assessment of nucleic acid and polypeptide molecules. In addition, one or more components present in such samples can be first isolated or enriched and further processed to assess the presence or levels of selected biomarkers, to assess a given biosignature (e.g., isolated microvesicles prior to profiling for protein and/or nucleic acid biomarkers).

[00203] Table 1 presents a non-limiting listing of diseases, conditions, or biological states and corresponding biological samples that may be used for analysis according to the methods of the invention.

Table 1: Examples of Biological Samples for Various Diseases, Conditions, or Biological States

Illustrative Disease, Condition or Biological State	**Illustrative Biological Samples**
Cancers/neoplasms affecting the following tissue types/bodily systems: breast, lung, ovarian, colon, rectal, prostate, pancreatic, brain, bone, connective tissue, glands, skin, lymph, nervous system, endocrine, germ cell, genitourinary, hematologic/blood, bone marrow, muscle, eye, esophageal, fat tissue, thyroid, pituitary, spinal cord, bile duct, heart, gall bladder, bladder, testes, cervical, endometrial, renal, ovarian, digestive/gastrointestinal, stomach, head and neck, liver, leukemia, respiratory/thorasic, cancers of unknown primary (CUP)	Tumor, blood, serum, plasma, cerebrospinal fluid (CSF), urine, sputum, ascites, synovial fluid, semen, nipple aspirates, saliva, bronchoalveolar lavage fluid, tears, oropharyngeal washes, feces, peritoneal fluids, pleural effusion, sweat, tears, aqueous humor, pericardial fluid, lymph, chyme, chyle, bile, stool water, amniotic fluid, breast milk, pancreatic juice, cerumen, Cowper's fluid or pre-ejaculatory fluid, female ejaculate, interstitial fluid, menses, mucus, pus, sebum, vaginal lubrication, vomit
Neurodegenerative/neurological disorders: Parkinson's disease, Alzheimer's Disease and multiple sclerosis, Schizophrenia, and bipolar disorder, spasticity disorders, epilepsy	Blood, serum, plasma, CSF, urine
Cardiovascular Disease: atherosclerosis, cardiomyopathy, endocarditis, vunerable plaques, infection	Blood, serum, plasma, CSF, urine
Stroke: ischemic, intracerebral hemorrhage, subarachnoid hemorrhage, transient ischemic attacks (TIA)	Blood, serum, plasma, CSF, urine
Pain disorders: peripheral neuropathic pain and chronic neuropathic pain, and fibromyalgia,	Blood, serum, plasma, CSF, urine
Autoimmune disease: systemic and localized diseases, rheumatic disease, Lupus, Sjogren's syndrome	Blood, serum, plasma, CSF, urine, synovial fluid
Digestive system abnormalities: Barrett's esophagus, irritable bowel syndrome, ulcerative colitis, Crohn's disease, Diverticulosis and Diverticulitis, Celiac Disease	Blood, serum, plasma, CSF, urine
Endocrine disorders: diabetes mellitus, various forms of Thyroiditis, adrenal disorders, pituitary	Blood, serum, plasma, CSF, urine

disorders	
Diseases and disorders of the skin: psoriasis	Blood, serum, plasma, CSF, urine, synovial fluid, tears
Urological disorders: benign prostatic hypertrophy (BPH), polycystic kidney disease, interstitial cystitis	Blood, serum, plasma, urine
Hepatic disease/injury: Cirrhosis, induced hepatotoxicity (due to exposure to natural or synthetic chemical sources)	Blood, serum, plasma, urine
Kidney disease/injury: acute, sub-acute, chronic conditions, Podocyte injury, focal segmental glomerulosclerosis	Blood, serum, plasma, urine
Endometriosis	Blood, serum, plasma, urine, vaginal fluids
Osteoporosis	Blood, serum, plasma, urine, synovial fluid
Pancreatitis	Blood, serum, plasma, urine, pancreatic juice
Asthma	Blood, serum, plasma, urine, sputum, bronchiolar lavage fluid
Allergies	Blood, serum, plasma, urine, sputum, bronchiolar lavage fluid
Prion-related diseases	Blood, serum, plasma, CSF, urine
Viral Infections: HIV/AIDS	Blood, serum, plasma, urine
Sepsis	Blood, serum, plasma, urine, tears, nasal lavage
Organ rejection/transplantation	Blood, serum, plasma, urine, various lavage fluids
Differentiating conditions: adenoma versus hyperplastic polyp, irritable bowel syndrome (IBS) versus normal, classifying Dukes stages A, B, C, and/or D of colon cancer, adenoma with low-grade hyperplasia versus high-grade hyperplasia, adenoma versus normal, colorectal cancer versus normal, IBS versus. ulcerative colitis (UC) versus Crohn's disease (CD),	Blood, serum, plasma, urine, sputum, feces, colonic lavage fluid
Pregnancy related physiological states, conditions, or affiliated diseases: genetic risk, adverse pregnancy outcomes	Maternal serum, plasma, amniotic fluid, cord blood

[00204] The methods of the invention can be used to characterize a phenotype using a blood sample or blood derivative. Blood derivatives include plasma and serum. Blood plasma is the liquid component of whole blood, and makes up approximately 55% of the total blood volume. It is composed primarily of water with small amounts of minerals, salts, ions, nutrients, and proteins in solution. In whole blood, red blood cells, leukocytes, and platelets are suspended within the plasma. Blood serum refers to blood plasma without fibrinogen or other clotting factors (i.e., whole blood minus both the cells and the clotting factors).

[00205] The biological sample may be obtained through a third party, such as a party not performing the analysis of the biomarkers, whether direct assessment of a biological sample or by profiling one or more vesicles obtained from the biological sample. For example, the sample may be obtained through a clinician, physician, or other health care manager of a subject from which the sample is derived. Alternatively, the biological sample may obtained by the same party analyzing the vesicle. In addition,

biological samples be assayed, are archived (e.g., frozen) or ortherwise stored in under preservative conditions.

[00206] Furthermore, a biological sample can comprise a vesicle or cell membrane fragment that is derived from a cell of origin and available extracellularly in a subject's biological fluid or extracellular milieu.

[00207] Methods of the invention can include assessing one or more vesicles, including assessing vesicle populations. A vesicle, as used herein, is a membrane vesicle that is shed from cells. Vesicles or membrane vesicles include without limitation: circulating microvesicles (cMVs), microvesicle, exosome, nanovesicle, dexosome, bleb, blebby, prostasome, microparticle, intralumenal vesicle, membrane fragment, intralumenal endosomal vesicle, endosomal-like vesicle, exocytosis vehicle, endosome vesicle, endosomal vesicle, apoptotic body, multivesicular body, secretory vesicle, phospholipid vesicle, liposomal vesicle, argosome, texasome, secresome, tolerosome, melanosome, oncosome, or exocytosed vehicle. Furthermore, although vesicles may be produced by different cellular processes, the methods of the invention are not limited to or reliant on any one mechanism, insofar as such vesicles are present in a biological sample and are capable of being characterized by the methods disclosed herein. Unless otherwise specified, methods that make use of a species of vesicle can be applied to other types of vesicles. Vesicles comprise spherical structures with a lipid bilayer similar to cell membranes which surrounds an inner compartment which can contain soluble components, sometimes referred to as the payload. In some embodiments, the methods of the invention make use of exosomes, which are small secreted vesicles of about 40–100 nm in diameter. For a review of membrane vesicles, including types and characterizations, see *Thery et al., Nat Rev Immunol. 2009 Aug;9(8):581-93.* Some properties of different types of vesicles include those in **Table 2**:

Table 2: Vesicle Properties

Feature	Exosomes	Microvesicles	Ectosomes	Membrane particles	Exosome-like vesicles	Apoptotic vesicles
Size	50-100 nm	100-1,000 nm	50-200 nm	50-80 nm	20-50 nm	50-500 nm
Density in sucrose	1.13-1.19 g/ml			1.04-1.07 g/ml	1.1 g/ml	1.16-1.28 g/ml
EM appearance	Cup shape	Irregular shape, electron dense	Bilamellar round structures	Round	Irregular shape	Heterogeneous
Sedimentation	100,000 g	10,000 g	160,000-200,000 g	100,000-200,000 g	175,000 g	1,200 g, 10,000 g, 100,000 g
Lipid composition	Enriched in cholesterol, sphingomyelin and ceramide; contains lipid rafts; expose PPS	Expose PPS	Enriched in cholesterol and diacylglycerol; expose PPS		No lipid rafts	
Major protein	Tetraspanins	Integrins,	CR1 and	CD133; no	TNFRI	Histones

markers	(e.g., CD63, CD9), Alix, TSG101	selectins and CD40 ligand	proteolytic enzymes; no CD63	CD63		
Intracellular origin	Internal compartments (endosomes)	Plasma membrane	Plasma membrane	Plasma membrane		

Abbreviations: phosphatidylserine (PPS); electron microscopy (EM)

[00208] Vesicles include shed membrane bound particles, or "microparticles," that are derived from either the plasma membrane or an internal membrane. Vesicles can be released into the extracellular environment from cells. Cells releasing vesicles include without limitation cells that originate from, or are derived from, the ectoderm, endoderm, or mesoderm. The cells may have undergone genetic, environmental, and/or any other variations or alterations. For example, the cell can be tumor cells. A vesicle can reflect any changes in the source cell, and thereby reflect changes in the originating cells, e.g., cells having various genetic mutations. In one mechanism, a vesicle is generated intracellularly when a segment of the cell membrane spontaneously invaginates and is ultimately exocytosed (see for example, *Keller et al., Immunol. Lett. 107 (2): 102–8 (2006)*). Vesicles also include cell-derived structures bounded by a lipid bilayer membrane arising from both herniated evagination (blebbing) separation and sealing of portions of the plasma membrane or from the export of any intracellular membrane-bounded vesicular structure containing various membrane-associated proteins of tumor origin, including surface-bound molecules derived from the host circulation that bind selectively to the tumor-derived proteins together with molecules contained in the vesicle lumen, including but not limited to tumor-derived microRNAs or intracellular proteins. Blebs and blebbing are further described in *Charras et al., Nature Reviews Molecular and Cell Biology, Vol. 9, No. 11, p. 730-736 (2008)*. A vesicle shed into circulation or bodily fluids from tumor cells may be referred to as a "circulating tumor-derived vesicle." When such vesicle is an exosome, it may be referred to as a circulating-tumor derived exosome (CTE). In some instances, a vesicle can be derived from a specific cell of origin. CTE, as with a cell-of-origin specific vesicle, typically have one or more unique biomarkers that permit isolation of the CTE or cell-of-origin specific vesicle, e.g., from a bodily fluid and sometimes in a specific manner. For example, a cell or tissue specific markers are used to identify the cell of origin. Examples of such cell or tissue specific markers are disclosed herein and can further be accessed in the Tissue-specific Gene Expression and Regulation (TiGER) Database, available at bioinfo.wilmer.jhu.edu/tiger/; Liu et al. (2008) TiGER: a database for tissue-specific gene expression and regulation. BMC Bioinformatics. 9:271; TissueDistributionDBs, available at genome.dkfz-heidelberg.de/menu/tissue_db/index.html.

[00209] A vesicle can have a diameter of greater than about 10 nm, 20 nm, or 30 nm. A vesicle can have a diameter of greater than 40 nm, 50 nm, 100 nm, 200 nm, 500 nm, 1000 nm, 1500 nm, 2000 nm or greater than 10,000 nm. A vesicle can have a diameter of about 20-2000 nm, about 20-1500 nm, about 30-1000 nm, about 30-800 nm, about 30-200 nm, or about 30-100 nm. In some embodiments, the vesicle has a diameter of less than 10,000 nm, 2000 nm, 1500 nm, 1000 nm, 800 nm, 500 nm, 200 nm, 100 nm, 50 nm, 40 nm, 30 nm, 20 nm or less than 10 nm. As used herein the term "about" in reference to a numerical

value means that variations of 10% above or below the numerical value are within the range ascribed to the specified value. Typical sizes for various types of vesicles are shown in **Table 2**. Vesicles can be assessed to measure the diameter of a single vesicle or any number of vesicles. For example, the range of diameters of a vesicle population or an average diameter of a vesicle population can be determined. Vesicle diameter can be assessed using methods known in the art, e.g., imaging technologies such as electron microscopy. In an embodiment, a diameter of one or more vesicles is determined using optical particle detection. See, e.g., U.S. Patent 7,751,053, entitled "Optical Detection and Analysis of Particles" and issued July 6, 2010; and U.S. Patent 7,399,600, entitled "Optical Detection and Analysis of Particles" and issued July 15, 2010.

[00210] In some embodiments, the methods of the invention comprise assessing vesicles directly such as in a biological sample without prior isolation, purification, or concentration from the biological sample. For example, the amount of vesicles in the sample can by itself provide a biosignature that provides a diagnostic, prognostic or theranostic determination. Alternatively, the vesicle in the sample may be isolated, captured, purified, or concentrated from a sample prior to analysis. As noted, isolation, capture or purification as used herein comprises partial isolation, partial capture or partial purification apart from other components in the sample. Vesicle isolation can be performed using various techniques as described herein, e.g., chromatography, filtration, centrifugation, flow cytometry, affinity capture (e.g., to a planar surface or bead), and/or using microfluidics. **FIGs. 19B-C** present an overview of a method of the invention for assessing microvesicles using an aptamer pool.

[00211] Vesicles such as exosomes can be assessed to provide a phenotypic characterization by comparing vesicle characteristics to a reference. In some embodiments, surface antigens on a vesicle are assessed. The surface antigens can provide an indication of the anatomical origin and/or cellular of the vesicles and other phenotypic information, e.g., tumor status. For example, wherein vesicles found in a patient sample, e.g., a bodily fluid such as blood, serum or plasma, are assessed for surface antigens indicative of colorectal origin and the presence of cancer. The surface antigens may comprise any informative biological entity that can be detected on the vesicle membrane surface, including without limitation surface proteins, lipids, carbohydrates, and other membrane components. For example, positive detection of colon derived vesicles expressing tumor antigens can indicate that the patient has colorectal cancer. As such, methods of the invention can be used to characterize any disease or condition associated with an anatomical or cellular origin, by assessing, for example, disease-specific and cell-specific biomarkers of one or more vesicles obtained from a subject.

[00212] In another embodiment, the methods of the invention comprise assessing one or more vesicle payload to provide a phenotypic characterization. The payload with a vesicle comprises any informative biological entity that can be detected as encapsulated within the vesicle, including without limitation proteins and nucleic acids, e.g., genomic or cDNA, mRNA, or functional fragments thereof, as well as microRNAs (miRs). In addition, methods of the invention are directed to detecting vesicle surface antigens (in addition or exclusive to vesicle payload) to provide a phenotypic characterization. For

example, vesicles can be characterized by using binding agents (e.g., antibodies or aptamers) that are specific to vesicle surface antigens, and the bound vesicles can be further assessed to identify one or more payload components disclosed therein. As described herein, the levels of vesicles with surface antigens of interest or with payload of interest can be compared to a reference to characterize a phenotype. For example, overexpression in a sample of cancer-related surface antigens or vesicle payload, e.g., a tumor associated mRNA or microRNA, as compared to a reference, can indicate the presence of cancer in the sample. The biomarkers assessed can be present or absent, increased or reduced based on the selection of the desired target sample and comparison of the target sample to the desired reference sample. Non-limiting examples of target samples include: disease; treated/not-treated; different time points, such as a in a longitudinal study; and non-limiting examples of reference sample: non-disease; normal; different time points; and sensitive or resistant to candidate treatment(s).

Microvesicle Isolation and Analysis

[00213] ***Sample Processing***

[00214] A vesicle or a population of vesicles may be isolated, purified, concentrated or otherwise enriched prior to and/or during analysis. Unless otherwise specified, the terms "purified," "isolated," or similar as used herein in reference to vesicles or biomarker components are intended to include partial or complete purification or isolation of such components from a cell or organism. Analysis of a vesicle can include quantitiating the amount one or more vesicle populations of a biological sample. For example, a heterogeneous population of vesicles can be quantitated, or a homogeneous population of vesicles, such as a population of vesicles with a particular biomarker profile, a particular biosignature, or derived from a particular cell type can be isolated from a heterogeneous population of vesicles and quantitated. Analysis of a vesicle can also include detecting, quantitatively or qualitatively, one or more particular biomarker profile or biosignature of a vesicle, as described herein.

[00215] A vesicle can be stored and archived, such as in a bio-fluid bank and retrieved for analysis as desired. A vesicle may also be isolated from a biological sample that has been previously harvested and stored from a living or deceased subject. In addition, a vesicle may be isolated from a biological sample which has been collected as described in *King et al., Breast Cancer Res 7(5): 198-204 (2005)*. A vesicle can be isolated from an archived or stored sample. Alternatively, a vesicle may be isolated from a biological sample and analyzed without storing or archiving of the sample. Furthermore, a third party may obtain or store the biological sample, or obtain or store the vesicle for analysis.

[00216] An enriched population of vesicles can be obtained from a biological sample. For example, vesicles may be concentrated or isolated from a biological sample using size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof.

[00217] Size exclusion chromatography, such as gel permeation columns, centrifugation or density gradient centrifugation, and filtration methods can be used. For example, a vesicle can be isolated by differential centrifugation, anion exchange and/or gel permeation chromatography (for example, as

described in US Patent Nos. 6,899,863 and 6,812,023), sucrose density gradients, organelle electrophoresis (for example, as described in U.S. Patent No. 7,198,923), magnetic activated cell sorting (MACS), or with a nanomembrane ultrafiltration concentrator. Various combinations of isolation or concentration methods can be used.

[00218] Highly abundant proteins, such as albumin and immunoglobulin in blood samples, may hinder isolation of vesicles from a biological sample. For example, a vesicle can be isolated from a biological sample using a system that uses multiple antibodies that are specific to the most abundant proteins found in a biological sample, such as blood. Such a system can remove up to several proteins at once, thus unveiling the lower abundance species such as cell-of-origin specific vesicles. This type of system can be used for isolation of vesicles from biological samples such as blood, cerebrospinal fluid or urine. The isolation of vesicles from a biological sample may also be enhanced by high abundant protein removal methods as described in *Chromy et al. J Proteome Res 2004; 3:1120-1127*. In another embodiment, the isolation of vesicles from a biological sample may also be enhanced by removing serum proteins using glycopeptide capture as described in *Zhang et al, Mol Cell Proteomics 2005;4:144-155*. In addition, vesicles from a biological sample such as urine may be isolated by differential centrifugation followed by contact with antibodies directed to cytoplasmic or anti-cytoplasmic epitopes as described in *Pisitkun et al., Proc Natl Acad Sci U S A, 2004;101:13368-13373*.

[00219] Plasma contains a large variety of proteins including albumin, immunoglobulins, and clotting proteins such as fibrinogen. About 60% of plasma protein comprises the protein albumin (e.g., human serum albumin or HSA), which contributes to osmotic pressure of plasma to assist in the transport of lipids and steroid hormones. Globulins make up about 35% of plasma proteins and are used in the transport of ions, hormones and lipids assisting in immune function. About 4% of plasma protein comprises fibrinogen which is essential in the clotting of blood and can be converted into the insoluble protein fibrin. Other types of blood proteins include: Prealbumin, Alpha 1 antitrypsin, Alpha 1 acid glycoprotein, Alpha 1 fetoprotein, Haptoglobin, Alpha 2 macroglobulin, Ceruloplasmin, Transferrin, complement proteins C3 and C4, Beta 2 microglobulin, Beta lipoprotein, Gamma globulin proteins, C-reactive protein (CRP), Lipoproteins (chylomicrons, VLDL, LDL, HDL), other globulins (types alpha, beta and gamma), Prothrombin and Mannose-binding lectin (MBL). Any of these proteins, including classes of proteins, or derivatives thereof (such as fibrin which is derived from the cleavage of fibrinogen) can be selectively depleted from a biological sample prior to further analysis performed on the sample. Without being bound by theory, removal of such background proteins may facilitate more sensitive, accurate, or precise detection of the biomarkers of interest in the sample.

[00220] Abundant proteins in blood or blood derivatives (e.g., plasma or serum) include without limitation albumin, IgG, transferrin, fibrinogen, IgA, α_2-Macroglobulin, IgM, α_1-Antitrypsin, complement C3, haptoglobulin, apolipoprotein A1, apolipoprotein A3, apolipoprotein B, α_1-Acid Glycoprotein, ceruloplasmin, complement C4, C1q, IgD, prealbumin (transthyretin), and plasminogen. Such proteins can be depleted using commercially available columns and kits. Examples of such columns comprise the

Multiple Affinity Removal System from Agilent Technologies (Santa Clara, CA). This system include various cartridges designed to deplete different protein profiles, including the following cartridges with performance characteristics according to the manufacturer: Human 14, which eliminates approximately 94% of total protein (albumin, IgG, antitrypsin, IgA, transferrin, haptoglobin, fibrinogen, alpha2-macroglobulin, alpha1-acid glycoprotein (orosomucoid), IgM, apolipoprotein AI, apolipoprotein AII, complement C3 and transthyretin); Human 7, which eliminates approximately 85 - 90% of total protein (albumin, IgG, IgA, transferrin, haptoglobin, antitrypsin, and fibrinogen); Human 6, which eliminates approximately 85 - 90% of total protein (albumin, IgG, IgA, transferrin, haptoglobin, and antitrypsin); Human Albumin/IgG, which eliminates approximately 69% of total protein (albumin and IgG); and Human Albumin, which eliminates approximately 50-55% of total protein (albumin). The ProteoPrep® 20 Plasma Immunodepletion Kit from Sigma-Aldrich is intended to specifically remove the 20 most abundant proteins from human plasma or serum, which is about remove 97-98% of the total protein mass in plasma or serum (Sigma-Aldrich, St. Louis, MO). According to the manufacturer, the ProteoPrep® 20 removes: albumin, IgG, transferrin, fibrinogen, IgA, α_2- Macroglobulin, IgM, α_1- Antitrypsin, complement C3, haptoglobulin, apolipoprotein A1, A3 and B; α_1- Acid Glycoprotein, ceruloplasmin, complement C4, C1q; IgD, prealbumin, and plasminogen. Sigma-Aldrich also manufactures ProteoPrep® columns to remove albumin (HSA) and immunoglobulins (IgG). The ProteomeLab IgY-12 High Capacity Proteome Partitioning kits from Beckman Coulter (Fullerton, CA) are specifically designed to remove twelve highly abundant proteins (Albumin, IgG, Transferrin, Fibrinogen, IgA, α2-macroglobulin, IgM, α1-Antitrypsin, Haptoglobin, Orosomucoid, Apolipoprotein A-I, Apolipoprotein A-II) from the human biological fluids such as serum and plasma. Generally, such systems rely on immunodepletion to remove the target proteins, e.g., using small ligands and/or full antibodies. The PureProteome™ Human Albumin/Immunoglobulin Depletion Kit from Millipore (EMD Millipore Corporation, Billerica, MA, USA) is a magnetic bead based kit that enables high depletion efficiency (typically >99%) of Albumin and all Immunoglobulins (i.e., IgG, IgA, IgM, IgE and IgD) from human serum or plasma samples. The ProteoExtract® Albumin/IgG Removal Kit, also from Millipore, is designed to deplete >80% of albumin and IgG from body fluid samples. Other similar protein depletion products include without limitation the following: Aurum™ Affi-Gels Blue mini kit (Bio-Rad, Hercules, CA, USA); Vivapure® anti-HSA/IgG kit (Sartorius Stedim Biotech, Goettingen, Germany), Qproteome albumin/IgG depletion kit (Qiagen, Hilden, Germany); Seppro® MIXED12-LC20 column (GenWay Biotech, San Diego, CA, USA); Abundant Serum Protein Depletion Kit (Norgen Biotek Corp., Ontario, Canada); GBC Human Albumin/IgG/Transferrin 3 in 1 Depletion Column/Kit (Good Biotech Corp., Taiwan). These systems and similar systems can be used to remove abundant proteins from a biological sample, thereby improving the ability to detect low abundance circulating biomarkers such as proteins and vesicles.

[00221] Thromboplastin is a plasma protein aiding blood coagulation through conversion of prothrombin to thrombin. Thrombin in turn acts as a serine protease that converts soluble fibrinogen into insoluble strands of fibrin, as well as catalyzing many other coagulation-related reactions. Thus, thromboplastin is a

protein that can be used to facilitate precipitation of fibrinogen/fibrin (blood clotting factors) out of plasma. In addition to or as an alternative to immunoaffinity protein removal, a blood sample can be treated with thromboplastin to deplete fibrinogen/fibrin. Thromboplastin removal can be performed in addition to or as an alternative to immunoaffinity protein removal as described above using methods known in the art. Precipitation of other proteins and/or other sample particulate can also improve detection of circulating biomarkers such as vesicles in a sample. For example, ammonium sulfate treatment as known in the art can be used to precipitate immunoglobulins and other highly abundant proteins.

[00222] In an embodiment, the invention provides a method of detecting a presence or level of one or more circulating biomarker such as a microvesicle in a biological sample, comprising: (a) providing a biological sample comprising or suspected to comprise the one or more circulating biomarker; (b) selectively depleting one or more abundant protein from the biological sample provided in step (a); (c) performing affinity selection of the one or more circulating biomarker from the sample depleted in step (b), thereby detecting the presence or level of one or more circulating biomarker. The biological sample may comprise a bodily fluid, e.g., peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, umbilical cord blood, or a derivative of any thereof. In some embodiments, the biological sample comprises peripheral blood, serum or plasma. Illustrative protocols and results from selectively depleting one or more abundant protein from blood plasma prior to vesicle detection can be found in **Example 40** of International Patent Publication No. WO/2014/082083, filed November 26, 2013, which patent publication is incorporated by reference herein in its entirety.

[00223] An abundant protein may comprise a protein in the sample that is present in the sample at a high enough concentration to potentially interfere with downstream processing or analysis. Typically, an abundant protein is not the target of any further analysis of the sample. The abundant protein may constitute at least 10^{-5}, 10^{-4}, 10^{-3}, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98 or at least 99% of the total protein mass in the sample. In some embodiments, the abundant protein is present at less than 10^{-5}% of the total protein mass in the sample, e.g., in the case of a rare target of interest. As described herein, in the case of blood or a derivative thereof, the one or more abundant protein may comprise one or more of albumin, IgG, transferrin, fibrinogen, fibrin, IgA, α2-Marcroglobulin, IgM, α1-Antitrypsin, complement C3, haptoglobulin, apolipoprotein A1, A3 and B; α1-Acid Glycoprotein, ceruloplasmin, complement C4, C1q, IgD, prealbumin (transthyretin), plasminogen, a derivative of any thereof, and a combination thereof. The one or more abundant protein in blood or a blood derivative may also comprise one or more of Albumin,

Immunoglobulins, Fibrinogen, Prealbumin, Alpha 1 antitrypsin, Alpha 1 acid glycoprotein, Alpha 1 fetoprotein, Haptoglobin, Alpha 2 macroglobulin, Ceruloplasmin, Transferrin, complement proteins C3 and C4, Beta 2 microglobulin, Beta lipoprotein, Gamma globulin proteins, C-reactive protein (CRP), Lipoproteins (chylomicrons, VLDL, LDL, HDL), other globulins (types alpha, beta and gamma), Prothrombin, Mannose-binding lectin (MBL), a derivative of any thereof, and a combination thereof.

[00224] In some embodiments, selectively depleting the one or more abundant protein comprises contacting the biological sample with thromboplastin to initiate precipitation of fibrin. The one or more abundant protein may also be depleted by immunoaffinity, precipitation, or a combination thereof. For example, the sample can be treated with thromboplastin to precipitate fibrin, and then the sample may be passed through a column to remove HSA, IgG, and other abundant proteins as desired.

[00225] "Selectively depleting" the one or more abundant protein comprises depleting the abundant protein from the sample at a higher percentage than depletion another entity in the sample, such as another protein or microvesicle, including a target of interest for downstream processing or analysis. Selectively depleting the one or more abundant protein may comprise depleting the abundant protein at a 1.1-fold, 1.2-fold, 1.3-fold, 1.4-fold, 1.5-fold, 1.6-fold, 1.7-fold, 1.8-fold, 1.9-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 11-fold, 12-fold, 13-fold, 14-fold, 15-fold, 16-fold, 17-fold, 18-fold, 19-fold, 20-fold, 25-fold, 30-fold, 40-fold, 50-fold, 60-fold, 70-fold, 80-fold, 90-fold, 100-fold, 200-fold, 300-fold, 400-fold, 500-fold, 600-fold, 700-fold, 800-fold, 900-fold, 1000-fold, 10^4-fold, 10^5-fold, 10^6-fold, 10^7-fold, 10^8-fold, 10^9-fold, 10^{10}-fold, 10^{11}-fold, 10^{12}-fold, 10^{13}-fold, 10^{14}-fold, 10^{15}-fold, 10^{16}-fold, 10^{17}-fold, 10^{18}-fold, 10^{19}-fold, 10^{20}-fold, or higher rate than another entity in the sample, such as another protein or microvesicle, including a target of interest for downstream processing or analysis. In an embodiment, there is little to no observable depletion of the target of interest as compared to the depletion of the abundant protein. In some embodiments, selectively depleting the one or more abundant protein from the biological sample comprises depleting at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the one or more abundant protein.

[00226] Removal of highly abundant proteins and other non-desired entities can further be facilitated with a non-stringent size exclusion step. For example, the sample can be processed using a high molecular weight cutoff size exclusion step to preferentially enrich high molecular weight vesicles apart from lower molecular weight proteins and other entities. In some embodiments, a sample is processed with a column (e.g., a gel filtration column) or filter having a molecular weight cutoff (MWCO) of 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10000, or greater than 10000 kiloDaltons (kDa). In an embodiment, a 700 kDa filtration column is used. In such a step, the vesicles will be retained or flow more slowly than the column or filter than the lower molecular weight entities. Such columns and filters are known in the art.

[00227] Isolation or enrichment of a vesicle from a biological sample can also be enhanced by use of sonication (for example, by applying ultrasound), detergents, other membrane-activating agents, or any

combination thereof. For example, ultrasonic energy can be applied to a potential tumor site, and without being bound by theory, release of vesicles from a tissue can be increased, allowing an enriched population of vesicles that can be analyzed or assessed from a biological sample using one or more methods disclosed herein.

[00228] With methods of detecting circulating biomarkers as described here, e.g., antibody affinity isolation, the consistency of the results can be optimized as desired using various concentration or isolation procedures. Such steps can include agitation such as shaking or vortexing, different isolation techniques such as polymer based isolation, e.g., with PEG, and concentration to different levels during filtration or other steps. It will be understood by those in the art that such treatments can be applied at various stages of testing the vesicle containing sample. In one embodiment, the sample itself, e.g., a bodily fluid such as plasma or serum, is vortexed. In some embodiments, the sample is vortexed after one or more sample treatment step, e.g., vesicle isolation, has occurred. Agitation can occur at some or all appropriate sample treatment steps as desired. Additives can be introduced at the various steps to improve the process, e.g., to control aggregation or degradation of the biomarkers of interest.

[00229] The results can also be optimized as desireable by treating the sample with various agents. Such agents include additives to control aggregation and/or additives to adjust pH or ionic strength. Additives that control aggregation include blocking agents such as bovine serum albumin (BSA), milk or StabilGuard® (a BSA-free blocking agent; Product code SG02, Surmodics, Eden Prairie, MN), chaotropic agents such as guanidium hydro chloride, and detergents or surfactants. Useful ionic detergents include sodium dodecyl sulfate (SDS, sodium lauryl sulfate (SLS)), sodium laureth sulfate (SLS, sodium lauryl ether sulfate (SLES)), ammonium lauryl sulfate (ALS), cetrimonium bromide, cetrimonium chloride, cetrimonium stearate, and the like. Useful non-ionic (zwitterionic) detergents include polyoxyethylene glycols, polysorbate 20 (also known as Tween 20), other polysorbates (e.g., 40, 60, 65, 80, etc), Triton-X (e.g., X100, X114), 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS), CHAPSO, deoxycholic acid, sodium deoxycholate, NP-40, glycosides, octyl-thio-glucosides, maltosides, and the like. In some embodiments, Pluronic F-68, a surfactant shown to reduce platelet aggregation, is used to treat samples containing vesicles during isolation and/or detection. F68 can be used from a 0.1% to 10% concentration, e.g., a 1%, 2.5% or 5% concentration. The pH and/or ionic strength of the solution can be adjusted with various acids, bases, buffers or salts, including without limitation sodium chloride (NaCl), phosphate-buffered saline (PBS), tris-buffered saline (TBS), sodium phosphate, potassium chloride, potassium phosphate, sodium citrate and saline-sodium citrate (SSC) buffer. In some embodiments, NaCl is added at a concentration of 0.1% to 10%, e.g., 1%, 2.5% or 5% final concentration. In some embodiments, Tween 20 is added to 0.005 to 2% concentration, e.g., 0.05%, 0.25% or 0.5 % final concentration. Blocking agents for use with the invention comprise inert proteins, e.g., milk proteins, non-fat dry milk protein, albumin, BSA, casein, or serum such as newborn calf serum (NBCS), goat serum, rabbit serum or salmon serum. The proteins can be added at a 0.1% to 10% concentration, e.g., 1%, 2%, 3%, 3.5%, 4%, 5%, 6%, 7%, 8%, 9% or 10% concentration. In some embodiments, BSA is added to 0.1%

to 10% concentration, e.g., 1%, 2%, 3%, 3.5%, 4%, 5%, 6%, 7%, 8%, 9% or 10% concentration. In an embodiment, the sample is treated according to the methodology presented in U.S. Patent Application 11/632946, filed July 13, 2005, which application is incorporated herein by reference in its entirety. Commercially available blockers may be used, such as SuperBlock, StartingBlock, Protein-Free from Pierce (a division of Thermo Fisher Scientific, Rockford, IL). In some embodiments, SSC/detergent (e.g., 20X SSC with 0.5% Tween 20 or 0.1% Triton-X 100) is added to 0.1% to 10% concentration, e.g., at 1.0% or 5.0% concentration.

[00230] The methods of detecting vesicles and other circulating biomarkers can be optimized as desired with various combinations of protocols and treatments as described herein. A detection protocol can be optimized by various combinations of agitation, isolation methods, and additives. In some embodiments, the patient sample is vortexed before and after isolation steps, and the sample is treated with blocking agents including BSA and/or F68. Such treatments may reduce the formation of large aggregates or protein or other biological debris and thus provide a more consistent detection reading.

[00231] ***Filtration and Ultrafiltration***

[00232] A vesicle can be isolated from a biological sample by filtering a biological sample from a subject through a filtration module and collecting from the filtration module a retentate comprising the vesicle, thereby isolating the vesicle from the biological sample. The method can comprise filtering a biological sample from a subject through a filtration module comprising a filter (also referred to herein as a selection membrane); and collecting from the filtration module a retentate comprising the vesicle, thereby isolating the vesicle from the biological sample. For example, in one embodiment, the filter retains molecules greater than about 100 kiloDaltons. In such cases, microvesicles are generally found within the retentate of the filtration process whereas smaller entities such as proteins, protein complexes, nucleic acids, etc, pass through into the filtrate.

[00233] The method can be used when determining a biosignature of one or more microvesicle. The method can also further comprise contacting the retentate from the filtration to a plurality of substrates, wherein each substrate is coupled to one or more capture agents, and each subset of the plurality of substrates comprises a different capture agent or combination of capture agents than another subset of the plurality of substrates.

[00234] Also provided herein is a method of determining a biosignature of a vesicle in a sample comprising: filtering a biological sample from a subject with a disorder through a filtration module, collecting from the filtration module a retentate comprising one or more vesicles, and determining a biosignature of the one or more vesicles. In one embodiment, the filtration module comprises a filter that retains molecules greater than about 100 or 150 kiloDaltons.

[00235] The method disclosed herein can further comprise characterizing a phenotype in a subject by filtering a biological sample from a subject through a filtration module, collecting from the filtration module a retentate comprising one or more vesicles; detecting a biosignature of the one or more vesicles; and characterizing a phenotype in the subject based on the biosignature, wherein characterizing is with at

least 70% sensitivity. In some embodiments, characterizing comprises determining an amount of one or more vesicle having the biosignature. Furthermore, the characterizing can be from about 80% to 100% sensitivity.

[00236] Also provided herein is a method for multiplex analysis of a plurality of vesicles. In some embodiments, the method comprises filtering a biological sample from a subject through a filtration module; collecting from the filtration module a retentate comprising the plurality of vesicles, applying the plurality of vesicles to a plurality of capture agents, wherein the plurality of capture agents is coupled to a plurality of substrates, and each subset of the plurality of substrates is differentially labeled from another subset of the plurality of substrates; capturing at least a subset of the plurality of vesicles; and determining a biosignature for at least a subset of the captured vesicles. In one embodiment, each substrate is coupled to one or more capture agents, and each subset of the plurality of substrates comprises a different capture agent or combination of capture agents as compared to another subset of the plurality of substrates. In some embodiments, at least a subset of the plurality of substrates is intrinsically labeled, such as comprising one or more labels. The substrate can be a particle or bead, or any combination thereof. In some embodiments, the filter retains molecules greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10000, or greater than 10000 kiloDaltons (kDa). In one embodiment, the filtration module comprises a filter that retains molecules greater than about 100 or 150 kiloDaltons. In one embodiment, the filtration module comprises a filter that retains molecules greater than about 9, 20, 100 or 150 kiloDaltons. In still another embodiment, the filtration module comprises a filter that retains molecules greater than about 7000 kDa.

[00237] In some embodiments, the method for multiplex analysis of a plurality of vesicles comprises filtering a biological sample from a subject through a filtration module, wherein the filtration module comprises a filter that retains molecules greater than about 100 kiloDaltons; collecting from the filtration module a retentate comprising the plurality of vesicles; applying the plurality of vesicles to a plurality of capture agents, wherein the plurality of capture agents is coupled to a microarray; capturing at least a subset of the plurality of vesicles on the microarray; and determining a biosignature for at least a subset of the captured vesicles. In some embodiments, the filter retains molecules greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10000, or greater than 10000 kiloDaltons (kDa). In one embodiment, the filtration module comprises a filter that retains molecules greater than about 100 or 150 kiloDaltons. In one embodiment, the filtration module comprises a filter that retains molecules greater than about 9, 20, 100 or 150 kiloDaltons. In still another embodiment, the filtration module comprises a filter that retains molecules greater than about 7000 kDa.

[00238] The biological sample can be clarified prior to isolation by filtration. Clarification comprises selective removal of cellular debris and other undesirable materials. For example, cellular debris and other components that may interfere with detection of the circulating biomarkers can be removed. The clarification can be by low-speed centrifugation, such as at about 5,000x g, 4,000x g, 3,000x g, 2,000x g, 1,000x g, or less. The supernatant, or clarified biological sample, containing the vesicle can then be collected and filtered to isolate the vesicle from the clarified biological sample. In some embodiments, the biological sample is not clarified prior to isolation of a vesicle by filtration.

[00239] In some embodiments, isolation of a vesicle from a sample does not use high-speed centrifugation, such as ultracentrifugation. For example, isolation may not require the use of centrifugal speeds, such as about 100,000x g or more. In some embodiments, isolation of a vesicle from a sample uses speeds of less than 50,000 x g, 40,000 x g, 30,000 x g, 20,000 x g, 15,000 x g, 12,000 x g, or 10,000 x g.

[00240] Any number of applicable filter configurations can be used to filter a sample of interest. In some embodiments, the filtration module used to isolate the circulating biomarkers from the biological sample is a fiber-based filtration cartridge. For example, the fiber can be a hollow polymeric fiber, such as a polypropylene hollow fiber. A biological sample can be introduced into the filtration module by pumping the sample fluid, such as a biological fluid as disclosed herein, into the module with a pump device, such as a peristaltic pump. The pump flow rate can vary, such as at about 0.25, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, or 10 mL/minute. The flow rate can be adjusted given the configuration, e.g., size and throughput, of the filtration module.

[00241] The filtration module can be a membrane filtration module. For example, the membrane filtration module can comprise a filter disc membrane, such as a hydrophilic polyvinylidene difluoride (PVDF) filter disc membrane housed in a stirred cell apparatus (e.g., comprising a magnetic stirrer). In some embodiments, the sample moves through the filter as a result of a pressure gradient established on either side of the filter membrane.

[00242] The filter can comprise a material having low hydrophobic absorptivity and/or high hydrophilic properties. For example, the filter can have an average pore size for vesicle retention and permeation of most proteins as well as a surface that is hydrophilic, thereby limiting protein adsorption. For example, the filter can comprise a material selected from the group consisting of polypropylene, PVDF, polyethylene, polyfluoroethylene, cellulose, secondary cellulose acetate, polyvinylalcohol, and ethylenevinyl alcohol (EVAL®, Kuraray Co., Okayama, Japan). Additional materials that can be used in a filter include, but are not limited to, polysulfone and polyethersulfone.

[00243] The filtration module can have a filter that retains molecules greater than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, or 900 kiloDaltons (kDa), such as a filter that has a MWCO (molecular weight cut off) of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, or 900 kDa, respectively. In

embodiments, the filtration module has a MWCO of 1000 kDa, 1500 kDa, 2000 kDa, 2500 kDa, 3000 kDa, 3500 kDa, 4000 kDa, 4500 kDa, 5000 kDa, 5500 kDa, 6000 kDa, 6500 kDa, 7000 kDa, 7500 kDa, 8000 kDa, 8500 kDa, 9000 kDa, 9500 kDa, 10000 kDa, or greater than 10000 kDa. Ultrafiltration membranes with a range of MWCO of 9 kDa, 20 kDa and/or 150 kDa can be used. In some embodiments, the filter within the filtration module has an average pore diameter of about 0.01 μm to about 0.15 μm, and in some embodiments from about 0.05 μm to about 0.12 μm. In some embodiments, the filter has an average pore diameter of about 0.06 μm, 0.07 μm, 0.08 μm, 0.09 μm, 0.1 μm, 0.11 μm or 0.2 μm.

[00244] The filtration module can be a commerically available column, such as a column typically used for concentrating proteins or for isolating proteins (e.g., ultrafiltration). Examples include, but are not limited to, columns from Millpore (Billerica, MA), such as Amicon® centrifugal filters, or from Pierce® (Rockford, IL), such as Pierce Concentrator filter devices. Useful columns from Pierce include disposable ultrafiltration centrifugal devices with a MWCO of 9 kDa, 20 kDa and/or 150 kDa. These concentrators consist of a high-performance regenerated cellulose membrane welded to a conical device. The filters can be as described in U.S. Patents 6,269,957 or 6,357,601, both of which applications are incorporated by reference in their entirety herein.

[00245] The retentate comprising the isolated vesicle can be collected from the filtration module. The retentate can be collected by flushing the retentate from the filter. Selection of a filter composition having hydrophilic surface properties, thereby limiting protein adsorption, can be used, without being bound by theory, for easier collection of the retentate and minimize use of harsh or time-consuming collection techniques.

[00246] The collected retentate can then be used subsequent analysis, such as assessing a biosignature of one or more vesicles in the retentate, as further described herein. The analysis can be directly performed on the collected retentate. Alternatively, the collected retentate can be further concentrated or purified, prior to analysis of one or more vesicles. For example, the retentate can be further concentrated or vesicles further isolated from the retentate using size exclusion chromatography, density gradient centrifugation, differential centrifugation, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof, such as described herein. In some embodiments, the retentate can undergo another step of filtration. Alternatively, prior to isolation of a vesicle using a filter, the vesicle is concentrated or isolated using techniques including without limitation size exclusion chromatography, density gradient centrifugation, differential centrifugation, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof.

[00247] Combinations of filters can be used for concentrating and isolating biomarkers. For example, the biological sample may first be filtered through a filter having a porosity or pore size of between about 0.01 μm to about 10 μm, e.g., 0.01 μm to about 2 μm or about 0.05 μm to about 1.5 μm, and then the sample is filtered. For example, prior to filtering a biological sample through a filtration module with a filter that retains molecules greater than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1000,

1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10000, or greater than 10000 kiloDaltons (kDa), such as a filter that has a MWCO (molecular weight cut off) of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10000, or greater than 10000 kDa, respectively, the biological sample may first be filtered through a filter having a porosity or pore size of between about 0.01 μm to about 10 μm, e.g., 0.01 μm to about 2 μm or about 0.05 μm to about 1.5 μm. In some embodiments, the filter has a pore size of about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9 or 2.0, 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0 or 10.0 μm. The filter may be a syringe filter. Thus, in one embodiment, the method comprises filtering the biological sample through a filter, such as a syringe filter, wherein the syringe filter has a porosity of greater than about 1 μm, prior to filtering the sample through a filtration module comprising a filter that retains molecules greater than about 100 or 150 kiloDaltons. In an embodiment, the filter is 1.2 μM filter and the filtration is followed by passage of the sample through a 7 ml or 20 ml concentrator column with a 150 kDa cutoff. Multiple concentrator columns may be used, e.g., in series. For example, a 7000 MWCO filtration unit can be used before a 150 MWCO unit.

[00248] The filtration module can be a component of a microfluidic device. Microfluidic devices, which may also be referred to as "lab-on-a-chip" systems, biomedical micro-electro-mechanical systems (bioMEMs), or multicomponent integrated systems, can be used for isolating, and analyzing, vesicles. Such systems miniaturize and compartmentalize processes that allow for binding of vesicles, detection of biomarkers, and other processes, such as further described herein.

[00249] The filtration module and assessment can be as described in Grant, R., et al., A filtration-based protocol to isolate human Plasma Membrane-derived Vesicles and exosomes from blood plasma, *J Immunol Methods* (2011) **371**:143-51 (Epub 2011 Jun 30), which reference is incorporated herein by reference in its entirety.

[00250] A microfluidic device can also be used for isolation of a vesicle by comprising a filtration module. For example, a microfluidic device can use one more channels for isolating a vesicle from a biological sample based on size from a biological sample. A biological sample can be introduced into one or more microfluidic channels, which selectively allows the passage of vesicles. The microfluidic device can further comprise binding agents, or more than one filtration module to select vesicles based on a property of the vesicles, for example, size, shape, deformability, biomarker profile, or biosignature.

[00251] The retentate from a filtration step can be further processed before assessment of microvesicles or other biomarkers therein. In an embodiment, the retentate is diluted prior to biomarker assessment, e.g., with an appropriate diluent such as a biologically compatible buffer. In some cases, the retentate is serially diluted. In an aspect, the invention provides a method for detecting a microvesicle population from a biological sample comprising: a) concentrating the biological sample using a selection membrane having a pore size of from 0.01 μm to about 10 μm, or a molecular weight cut off (MWCO) from about 1 kDa to

10000 kDa; b) diluting a retentate from the concentration step into one or more aliquots; and c) contacting each of the one or more aliquots of retentate with one or more binding agent specific to a molecule of at least one microvesicle in the microvesicle population. In a related aspect, the invention provides a method for detecting a microvesicle population from a biological sample comprising: a) concentrating the biological sample using a selection membrane having a pore size of from 0.01 μm to about 10 μm, or a molecular weight cut off (MWCO) from about 1 kDa to 10000 kDa; and b) contacting one or more aliquots of the retentate from the concentrating step with one or more binding agent specific to a molecule of at least one microvesicle in the microvesicle population.

[00252] The selection membrane can be sized to retain the desired biomarkers in the retentate or to allow the desired biomarkers to pass through the filter into the filtrate. The filter membrane can be chosen to have a certain pore size or MWCO value. The selection membrane can have a pore size of about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9 or 2.0, 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0 or 10.0 μm. The selection membrane can also have a MWCO of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000 or 10000 kDa.

[00253] The retentate can be separated and/or diluted into any number of desired aliquots. For example, multiple aliquots without any dilution or the same dilution can be used to determine reproducibility. In another example, multiple aliquots at different dilutions can be used to construct a concentration curve. In an embodiment, the retentate is separated and/or diluted into at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 200, 250, 300, 350 or 400 aliquots. The aliquots can be at a same dilution or at different dilutions.

[00254] A dilution factor is the ratio of the final volume of a mixture (the mixture of the diluents and aliquot) divided by the initial volume of the aliquot. The retentate can be diluted into one or more aliquots at a dilution factor of about 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000 and/or 100000. For example, the retentate can be diluted into one or more aliquot at a dilution factor of about 500.

[00255] To estimate a concentration or form a curve, the retentate can be diluted into multiple aliquots. In an embodiment of the method, the retentate is diluted into one or more aliquots at a dilution factor of about 100, 250, 500, 1000, 10000 and 100000. As desired, the method can further comprise detecting an amount of microvesicles in each aliquot of retentate, e.g., that formed a complex with the one or more binding agent. The curve can be used to determine a linear range of the amount of microvesicles in each aliquot detected versus dilution factor. A concentration of the detected microvesicles for the biological sample can be determined using the amount of microvesicles determined in one or more aliquot within the

linear range. The concentration can be compared to a reference concentration, e.g., in order to characterize a phenotype as described herein.

[00256] The invention also provides a related method comprising filtering a biological sample from a subject through a filtration module and collecting a filtrate comprising the vesicle, thereby isolating the vesicle from the biological sample. In such cases cells and other large entities can be retained in the retentate while microvesicles pass through into the filtrate. It will be appreciated that strategies to retain and filter microvesicles can be used in concert. For example, a sample can be filtered with a selection membrane that allows microvesicles to pass through, thereby isolating the microvesicles from large particles (cells, complexes, etc). The filtrate comprising the microvesicle can then be filtered using a selection membrane that retains microvesicles, thereby isolating the microvesicles from smaller particles (proteins, nucleic acids, etc). The isolated microvesicles can be further assessed according to the methods of the invention, e.g., to characterize a phenotype.

[00257] ***Precipitation***

[00258] Vesicles can be isolated using a polymeric precipitation method. The method can be in combination with or in place of the other isolation methods described herein. In one embodiment, the sample containing the vesicles is contacted with a formulation of polyethylene glycol (PEG). The polymeric formulation is incubated with the vesicle containing sample then precipitated by centrifugation. The PEG can bind to the vesicles and can be treated to specifically capture vesicles by addition of a capture moiety, e.g., a pegylated-binding protein such as an antibody. One of skill will appreciate that other polymers in addition to PEG can be used, e.g., PEG derivatives including methoxypolyethylene glycols, poly (ethylene oxide), and various polymers of formula $HO\text{-}CH_2\text{-}(CH_2\text{-}O\text{-}CH_2\text{-})n\text{-}CH_2\text{-}OH$ having different molecular weights. The efficiency of isolation may depend on various factors including the length of the polymer chains and concentration of polymer used. In preferred embodiments, PEG4000 or PEG 8000 may be used at a concentration of 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, or 10%, e.g., 4% or 8%.

[00259] In some embodiments of the invention, the vesicles are concentrated from a sample using the polymer precipitation method and the isolated vesicles are further separated using another approach. The second step can be used to identify a subpopulation of vesicles, e.g., that display certain biomarkers. The second separation step can comprise size exclusion, a binding agent, an antibody capture step, microbeads, as described herein.

[00260] In an embodiment, vesicles are isolated according to the ExoQuick™ and ExoQuick-TC™ kits from System Biosciences, Mountain View, CA USA. These kits use a polymer-based precipitation method to pellet vesicles. Similarly, the vesicles can be isolated using the Total Exosome Isolation (from Serum) or Total Exosome Isolation (from Cell Culture Media) kits from Invitrogen / Life Technologies (Carlsbad, CA USA). The Total Exosome Isolation reagent forces less-soluble components such as vesicles out of solution, allowing them to be collected by a short, low-speed centrifugation. The reagent is

added to the biological sample, and the solution is incubated overnight at 2 °C to 8 °C. The precipitated vesicles are recovered by standard centrifugation.

[00261] *Binding Agents*

[00262] Binding agents (also referred to as binding reagents) include agents that are capable of binding a target biomarker. A binding agent can be specific for the target biomarker, meaning the agent is capable of binding a target biomarker. The target can be any useful biomarker disclosed herein, such as a biomarker on the vesicle surface. In some embodiments, the target is a single molecule, such as a single protein, so that the binding agent is specific to the single protein. In other embodiments, the target can be a group of molecules, such as a family or proteins having a similar epitope or moiety, so that the binding agent is specific to the family or group of proteins. The group of molecules can also be a class of molecules, such as protein, DNA or RNA. The binding agent can be a capture agent used to capture a vesicle by binding a component or biomarker of a vesicle. In some embodiments, a capture agent comprises an antibody or fragment thereof, or an aptamer, that binds to an antigen on a vesicle. The capture agent can be optionally coupled to a substrate and used to isolate a vesicle, as further described herein.

[00263] A binding agent is an agent that binds to a circulating biomarker, such as a vesicle or a component of a vesicle. The binding agent can be used as a capture agent and/or a detection agent. A capture agent can bind and capture a circulating biomarker, such as by binding a component or biomarker of a vesicle. For example, the capture agent can be a capture antibody or capture antigen that binds to an antigen on a vesicle. A detection agent can bind to a circulating biomarker thereby facilitating detection of the biomarker. For example, a capture agent comprising an antibody or aptamer that is sequestered to a substrate can be used to capture a vesicle in a sample, and a detection agent comprising an antibody or aptamer that carries a label can be used to detect the captured vesicle via detection of the detection agent's label. In some embodiments, a vesicle is assessed using capture and detection agents that recognize the same vesicle biomarkers. For example, a vesicle population can be captured using a tetraspanin such as by using an anti-CD9 antibody bound to a substrate, and the captured vesicles can be detected using a fluorescently labeled anti-CD9 antibody to label the captured vesicles. In other embodiments, a vesicle is assessed using capture and detection agents that recognize different vesicle biomarkers. For example, a vesicle population can be captured using a cell-specific marker such as by using an anti-PCSA antibody bound to a substrate, and the captured vesicles can be detected using a fluorescently labeled anti-CD9 antibody to label the captured vesicles. Similarly, the vesicle population can be captured using a general vesicle marker such as by using an anti-CD9 antibody bound to a substate, and the captured vesicles can be detected using a fluorescently labeled antibody to a cell-specific or disease specific marker to label the captured vesicles.

[00264] The biomarkers recognized by the binding agent are sometimes referred to herein as an antigen. Unless otherwise specified, antigen as used herein is meant to encompass any entity that is capable of being bound by a binding agent, regardless of the type of binding agent or the immunogenicity of the

biomarker. The antigen further encompasses a functional fragment thereof. For example, an antigen can encompass a protein biomarker capable of being bound by a binding agent, including a fragment of the protein that is capable of being bound by a binding agent.

[00265] In one embodiment, a vesicle is captured using a capture agent that binds to a biomarker on a vesicle. The capture agent can be coupled to a substrate and used to isolate a vesicle, as further described herein. In one embodiment, a capture agent is used for affinity capture or isolation of a vesicle present in a substance or sample.

[00266] A binding agent can be used after a vesicle is concentrated or isolated from a biological sample. For example, a vesicle can first be isolated from a biological sample before a vesicle with a specific biosignature is isolated or detected. The vesicle with a specific biosignature can be isolated or detected using a binding agent for the biomarker. A vesicle with the specific biomarker can be isolated or detected from a heterogeneous population of vesicles. Alternatively, a binding agent may be used on a biological sample comprising vesicles without a prior isolation or concentration step. For example, a binding agent is used to isolate or detect a vesicle with a specific biosignature directly from a biological sample.

[00267] A binding agent can be a nucleic acid, protein, or other molecule that can bind to a component of a vesicle. The binding agent can comprise DNA, RNA, monoclonal antibodies, polyclonal antibodies, Fabs, Fab', single chain antibodies, synthetic antibodies, aptamers (DNA/RNA), peptoids, zDNA, peptide nucleic acids (PNAs), locked nucleic acids (LNAs), unlocked nucleic acid (UNA), lectins, synthetic or naturally occurring chemical compounds (including but not limited to drugs, labeling reagents), dendrimers, or a combination thereof. For example, the binding agent can be a capture antibody. In embodiments of the invention, the binding agent comprises a membrane protein labeling agent. See, e.g., the membrane protein labeling agents disclosed in Alroy et al., US. Patent Publication US 2005/0158708. In an embodiment, vesicles are isolated or captured as described herein, and one or more membrane protein labeling agent is used to detect the vesicles.

[00268] In some instances, a single binding agent can be employed to isolate or detect a vesicle. In other instances, a combination of different binding agents may be employed to isolate or detect a vesicle. For example, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 50, 75 or 100 different binding agents may be used to isolate or detect a vesicle from a biological sample. Furthermore, the one or more different binding agents for a vesicle can form a biosignature of a vesicle, as further described below.

[00269] Different binding agents can also be used for multiplexing. For example, isolation or detection of more than one population of vesicles can be performed by isolating or detecting each vesicle population with a different binding agent. Different binding agents can be bound to different particles, wherein the different particles are labeled. In another embodiment, an array comprising different binding agents can be used for multiplex analysis, wherein the different binding agents are differentially labeled or can be ascertained based on the location of the binding agent on the array. Multiplexing can be accomplished up to the resolution capability of the labels or detection method, such as described below. The binding agents

can be used to detect the vesicles, such as for detecting cell-of-origin specific vesicles. A binding agent or multiple binding agents can themselves form a binding agent profile that provides a biosignature for a vesicle. One or more binding agents can be selected from Fig. 2 of International Patent Publication No. WO/2011/127219, entitled "Circulating Biomarkers for Disease" and filed April 6, 2011, which application is incorporated by reference in its entirety herein. For example, if a vesicle population is detected or isolated using two, three, four or more binding agents in a differential detection or isolation of a vesicle from a heterogeneous population of vesicles, the particular binding agent profile for the vesicle population provides a biosignature for the particular vesicle population. The vesicle can be detected using any number of binding agents in a multiplex fashion. Thus, the binding agent can also be used to form a biosignature for a vesicle. The biosignature can be used to characterize a phenotype.

[00270] The binding agent can be a lectin. Lectins are proteins that bind selectively to polysaccharides and glycoproteins and are widely distributed in plants and animals. For example, lectins such as those derived from Galanthus nivalis in the form of Galanthus nivalis agglutinin ("GNA"), Narcissus pseudonarcissus in the form of Narcissus pseudonarcissus agglutinin ("NPA") and the blue green algae Nostoc ellipsosporum called "cyanovirin" (*Boyd et al. Antimicrob Agents Chemother 41(7): 1521 1530, 1997; Hammar et al. Ann N Y Acad Sci 724: 166 169, 1994; Kaku et al. Arch Biochem Biophys 279(2): 298 304, 1990*) can be used to isolate a vesicle. These lectins can bind to glycoproteins having a high mannose content (*Chervenak et al. Biochemistry 34(16): 5685 5695, 1995*). High mannose glycoprotein refers to glycoproteins having mannose-mannose linkages in the form of α-1→3 or α-1→6 mannose-mannose linkages.

[00271] The binding agent can be an agent that binds one or more lectins. Lectin capture can be applied to the isolation of the biomarker cathepsin D since it is a glycosylated protein capable of binding the lectins Galanthus nivalis agglutinin (GNA) and concanavalin A (ConA).

[00272] Methods and devices for using lectins to capture vesicles are described in International Patent Publications WO/2011/066589, entitled "METHODS AND SYSTEMS FOR ISOLATING, STORING, AND ANALYZING VESICLES" and filed November 30, 2010; WO/2010/065765, entitled "AFFINITY CAPTURE OF CIRCULATING BIOMARKERS" and filed December 3, 2009; WO/2010/141862, entitled "METHODS AND MATERIALS FOR ISOLATING EXOSOMES" and filed June 4, 2010; and WO/2007/103572, entitled "EXTRACORPOREAL REMOVAL OF MICROVESICULAR PARTICLES" and filed March 9, 2007, each of which applications is incorporated by reference herein in its entirety.

[00273] The binding agent can be an antibody. For example, a vesicle may be isolated using one or more antibodies specific for one or more antigens present on the vesicle. For example, a vesicle can have CD63 on its surface, and an antibody, or capture antibody, for CD63 can be used to isolate the vesicle. Alternatively, a vesicle derived from a tumor cell can express EpCam, the vesicle can be isolated using an antibody for EpCam and CD63. Other antibodies for isolating vesicles can include an antibody, or capture antibody, to CD9, PSCA, TNFR, CD63, B7H3, MFG-E8, EpCam, Rab, CD81, STEAP, PCSA, PSMA, or 5T4. Other antibodies for isolating vesicles can include an antibody, or capture antibody, to DR3, STEAP,

epha2, TMEM211, MFG-E8, Tissue Factor (TF), unc93A, A33, CD24, NGAL, EpCam, MUC17, TROP2, or TETS.

[00274] In some embodiments, the capture agent is an antibody to CD9, CD63, CD81, PSMA, PCSA, B7H3, EpCam, PSCA, ICAM, STEAP, or EGFR. The capture agent can also be used to identify a biomarker of a vesicle. For example, a capture agent such as an antibody to CD9 would identify CD9 as a biomarker of the vesicle. In some embodiments, a plurality of capture agents can be used, such as in multiplex analysis. The plurality of captures agents can comprise binding agents to one or more of: CD9, CD63, CD81, PSMA, PCSA, B7H3, EpCam, PSCA, ICAM, STEAP, and EGFR. In some embodiments, the plurality of capture agents comprise binding agents to CD9, CD63, CD81, PSMA, PCSA, B7H3, MFG-E8, and/or EpCam. In yet other embodiments, the plurality of capture agents comprises binding agents to CD9, CD63, CD81, PSMA, PCSA, B7H3, EpCam, PSCA, ICAM, STEAP, and/or EGFR. The plurality of capture agents comprises binding agents to TMEM211, MFG-E8, Tissue Factor (TF), and/or CD24.

[00275] The antibodies referenced herein can be immunoglobulin molecules or immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that specifically binds an antigen and synthetic antibodies. The immunoglobulin molecules can be of any class (e.g., IgG, IgE, IgM, IgD or IgA) or subclass of immunoglobulin molecule. Antibodies include, but are not limited to, polyclonal, monoclonal, bispecific, synthetic, humanized and chimeric antibodies, single chain antibodies, Fab fragments and $F(ab')_2$ fragments, Fv or Fv' portions, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, or epitope-binding fragments of any of the above. An antibody, or generally any molecule, "binds specifically" to an antigen (or other molecule) if the antibody binds preferentially to the antigen, and, e.g., has less than about 30%, 20%, 10%, 5% or 1% cross-reactivity with another molecule.

[00276] The binding agent can also be a polypeptide or peptide. Polypeptide is used in its broadest sense and may include a sequence of subunit amino acids, amino acid analogs, or peptidomimetics. The subunits may be linked by peptide bonds. The polypeptides may be naturally occurring, processed forms of naturally occurring polypeptides (such as by enzymatic digestion), chemically synthesized or recombinantly expressed. The polypeptides for use in the methods of the present invention may be chemically synthesized using standard techniques. The polypeptides may comprise D-amino acids (which are resistant to L- amino acid-specific proteases), a combination of D- and L-amino acids, β amino acids, or various other designer or non-naturally occurring amino acids (e.g., β-methyl amino acids, Cα- methyl amino acids, and Nα-methyl amino acids, etc.) to convey special properties. Synthetic amino acids may include ornithine for lysine, and norleucine for leucine or isoleucine. In addition, the polypeptides can have peptidomimetic bonds, such as ester bonds, to prepare polypeptides with novel properties. For example, a polypeptide may be generated that incorporates a reduced peptide bond, i.e., R_1-CH_2-NH-R_2, where R_1 and R_2 are amino acid residues or sequences. A reduced peptide bond may be introduced as a dipeptide subunit. Such a polypeptide would be resistant to protease activity, and would possess an

extended half- live in vivo. Polypeptides can also include peptoids (N-substituted glycines), in which the side chains are appended to nitrogen atoms along the molecule's backbone, rather than to the α-carbons, as in amino acids. Polypeptides and peptides are intended to be used interchangeably throughout this application, i.e. where the term peptide is used, it may also include polypeptides and where the term polypeptides is used, it may also include peptides. The term "protein" is also intended to be used interchangeably throughout this application with the terms "polypeptides" and "peptides" unless otherwise specified.

[00277] A vesicle may be isolated, captured or detected using a binding agent. The binding agent can be an agent that binds a vesicle "housekeeping protein," or general vesicle biomarker. The biomarker can be CD63, CD9, CD81, CD82, CD37, CD53, Rab-5b, Annexin V, MFG-E8 or other commonly observed vesicle markers include those listed in **Table 3**. Furthermore, any of the markers disclosed herein or in **Table 3** can be selected in identifying a candidate biosignature for a disease or condition, where the one or more selected biomarkers have a direct or indirect role or function in mechanisms involved in the disease or condition.

[00278] The binding agent can also be an agent that binds to a vesicle derived from a specific cell type, such as a tumor cell (e.g. binding agent for Tissue factor, EpCam, B7H3, RAGE or CD24) or a specific cell-of-origin. The binding agent used to isolate or detect a vesicle can be a binding agent for an antigen selected from **Fig. 1** of International Patent Publication No. WO/2011/127219, entitled "Circulating Biomarkers for Disease" and filed April 6, 2011, which application is incorporated by reference in its entirety herein. The binding agent for a vesicle can also be selected from those listed in **Fig. 2** of International Patent Publication No. WO/2011/127219. The binding agent can be for an antigen such as a tetraspanin, MFG-E8, Annexin V, 5T4, B7H3, caveolin, CD63, CD9, E-Cadherin, Tissue factor, MFG-E8, TMEM211, CD24, PSCA, PCSA, PSMA, Rab-5B, STEAP, TNFR1, CD81, EpCam, CD59, CD81, ICAM, EGFR, or CD66. A binding agent for a platelet can be a glycoprotein such as GpIa-IIa, GpIIb-IIIa, GpIIIb, GpIb, or GpIX. A binding agent can be for an antigen comprisine one or more of CD9, Erb2, Erb4, CD81, Erb3, MUC16, CD63, DLL4, HLA-Drpe, B7H3, IFNAR, 5T4, PCSA, MICB, PSMA, MFG-E8, Muc1, PSA, Muc2, Unc93a, VEGFR2, EpCAM, VEGF A, TMPRSS2, RAGE, PSCA, CD40, Muc17, IL-17-RA, and CD80. For example, the binding agent can be one or more of CD9, CD63, CD81, B7H3, PCSA, MFG-E8, MUC2, EpCam, RAGE and Muc17. One or more binding agents, such as one or more binding agents for two or more of the antigens, can be used for isolating or detecting a vesicle. The binding agent used can be selected based on the desire of isolating or detecting a vesicle derived from a particular cell type or cell-of-origin specific vesicle. The binding agent can be to one or more vesicle marker in **Table 4**.

[00279] A binding agent can also be linked directly or indirectly to a solid surface or substrate. A solid surface or substrate can be any physically separable solid to which a binding agent can be directly or indirectly attached including, but not limited to, surfaces provided by microarrays and wells, particles such as beads, columns, optical fibers, wipes, glass and modified or functionalized glass, quartz, mica,

diazotized membranes (paper or nylon), polyformaldehyde, cellulose, cellulose acetate, paper, ceramics, metals, metalloids, semiconductive materials, quantum dots, coated beads or particles, other chromatographic materials, magnetic particles; plastics (including acrylics, polystyrene, copolymers of styrene or other materials, polypropylene, polyethylene, polybutylene, polyurethanes, polytetrafluoroethylene (PTFE, Teflon®), etc.), polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, plastics, ceramics, conducting polymers (including polymers such as polypyrole and polyindole); micro or nanostructured surfaces such as nucleic acid tiling arrays, nanotube, nanowire, or nanoparticulate decorated surfaces; or porous surfaces or gels such as methacrylates, acrylamides, sugar polymers, cellulose, silicates, or other fibrous or stranded polymers. In addition, as is known the art, the substrate may be coated using passive or chemically-derivatized coatings with any number of materials, including polymers, such as dextrans, acrylamides, gelatins or agarose. Such coatings can facilitate the use of the array with a biological sample.

[00280] For example, an antibody used to isolate a vesicle can be bound to a solid substrate such as a well, such as commercially available plates (e.g. from Nunc, Milan Italy). Each well can be coated with the antibody. In some embodiments, the antibody used to isolate a vesicle is bound to a solid substrate such as an array. The array can have a predetermined spatial arrangement of molecule interactions, binding islands, biomolecules, zones, domains or spatial arrangements of binding islands or binding agents deposited within discrete boundaries. Further, the term array may be used herein to refer to multiple arrays arranged on a surface, such as would be the case where a surface bore multiple copies of an array. Such surfaces bearing multiple arrays may also be referred to as multiple arrays or repeating arrays.

[00281] Arrays typically contain addressable moieties that can detect the presense of an entity, e.g., a vesicle in the sample via a binding event. An array may be referred to as a microarray. Arrays or microarrays include without limitation DNA microarrays, such as cDNA microarrays, oligonucleotide microarrays and SNP microarrays, microRNA arrays, protein microarrays, antibody microarrays, tissue microarrays, cellular microarrays (also called transfection microarrays), chemical compound microarrays, and carbohydrate arrays (glycoarrays). DNA arrays typically comprise addressable nucleotide sequences that can bind to sequences present in a sample. MicroRNA arrays, e.g., the MMChips array from the University of Louisville or commercial systems from Agilent, can be used to detect microRNAs. Protein microarrays can be used to identify protein–protein interactions, including without limitation identifying substrates of protein kinases, transcription factor protein-activation, or to identify the targets of biologically active small molecules. Protein arrays may comprise an array of different protein molecules, commonly antibodies, or nucleotide sequences that bind to proteins of interest. In a non-limiting example, a protein array can be used to detect vesicles having certain proteins on their surface. Antibody arrays comprise antibodies spotted onto the protein chip that are used as capture molecules to detect proteins or other biological materials from a sample, e.g., from cell or tissue lysate solutions. For example, antibody arrays can be used to detect vesicle-associated biomarkers from bodily fluids, e.g., serum or urine. Tissue

microarrays comprise separate tissue cores assembled in array fashion to allow multiplex histological analysis. Cellular microarrays, also called transfection microarrays, comprise various capture agents, such as antibodies, proteins, or lipids, which can interact with cells to facilitate their capture on addressable locations. Cellular arrays can also be used to capture vesicles due to the similarity between a vesicle and cellular membrane. Chemical compound microarrays comprise arrays of chemical compounds and can be used to detect protein or other biological materials that bind the compounds. Carbohydrate arrays (glycoarrays) comprise arrays of carbohydrates and can detect, e.g., protein that bind sugar moieties. One of skill will appreciate that similar technologies or improvements can be used according to the methods of the invention.

[00282] A binding agent can also be bound to particles such as beads or microspheres. For example, an antibody specific for a component of a vesicle can be bound to a particle, and the antibody-bound particle is used to isolate a vesicle from a biological sample. In some embodiments, the microspheres may be magnetic or fluorescently labeled. In addition, a binding agent for isolating vesicles can be a solid substrate itself. For example, latex beads, such as aldehyde/sulfate beads (Interfacial Dynamics, Portland, OR) can be used.

[00283] A binding agent bound to a magnetic bead can also be used to isolate a vesicle. For example, a biological sample such as serum from a patient can be collected for colon cancer screening. The sample can be incubated with anti-CCSA-3 (Colon Cancer–Specific Antigen) coupled to magnetic microbeads. A low-density microcolumn can be placed in the magnetic field of a MACS Separator and the column is then washed with a buffer solution such as Tris-buffered saline. The magnetic immune complexes can then be applied to the column and unbound, non-specific material can be discarded. The CCSA-3 selected vesicle can be recovered by removing the column from the separator and placing it on a collection tube. A buffer can be added to the column and the magnetically labeled vesicle can be released by applying the plunger supplied with the column. The isolated vesicle can be diluted in IgG elution buffer and the complex can then be centrifuged to separate the microbeads from the vesicle. The pelleted isolated cell-of-origin specific vesicle can be resuspended in buffer such as phosphate-buffered saline and quantitated. Alternatively, due to the strong adhesion force between the antibody captured cell-of-origin specific vesicle and the magnetic microbeads, a proteolytic enzyme such as trypsin can be used for the release of captured vesicles without the need for centrifugation. The proteolytic enzyme can be incubated with the antibody captured cell-of-origin specific vesicles for at least a time sufficient to release the vesicles.

[00284] A binding agent, such as an antibody, for isolating vesicles is preferably contacted with the biological sample comprising the vesicles of interest for at least a time sufficient for the binding agent to bind to a component of the vesicle. For example, an antibody may be contacted with a biological sample for various intervals ranging from seconds days, including but not limited to, about 10 minutes, 30 minutes, 1 hour, 3 hours, 5 hours, 7 hours, 10 hours, 15 hours, 1 day, 3 days, 7 days or 10 days.

[00285] A binding agent, such as an antibody specific to an antigen listed in **Fig. 1** of International Patent Publication No. WO/2011/127219, entitled "Circulating Biomarkers for Disease" and filed April 6, 2011,

which application is incorporated by reference in its entirety herein, or a binding agent listed in **Fig. 2** of International Patent Publication No. WO/2011/127219, can be labeled to facilitate detection. Appropriate labels include without limitation a magnetic label, a fluorescent moiety, an enzyme, a chemiluminescent probe, a metal particle, a non-metal colloidal particle, a polymeric dye particle, a pigment molecule, a pigment particle, an electrochemically active species, semiconductor nanocrystal or other nanoparticles including quantum dots or gold particles, fluorophores, quantum dots, or radioactive labels. Various protein, radioactive, fluorescent, enzymatic, and other labels are described further above.

[00286] A binding agent can be directly or indirectly labeled, e.g., the label is attached to the antibody through biotin-streptavidin. Alternatively, an antibody is not labeled, but is later contacted with a second antibody that is labeled after the first antibody is bound to an antigen of interest.

[00287] Depending on the method of isolation or detection used, the binding agent may be linked to a solid surface or substrate, such as arrays, particles, wells and other substrates described above. Methods for direct chemical coupling of antibodies, to the cell surface are known in the art, and may include, for example, coupling using glutaraldehyde or maleimide activated antibodies. Methods for chemical coupling using multiple step procedures include biotinylation, coupling of trinitrophenol (TNP) or digoxigenin using for example succinimide esters of these compounds. Biotinylation can be accomplished by, for example, the use of D-biotinyl-N-hydroxysuccinimide. Succinimide groups react effectively with amino groups at pH values above 7, and preferentially between about pH 8.0 and about pH 8.5. Biotinylation can be accomplished by, for example, treating the cells with dithiothreitol followed by the addition of biotin maleimide.

[00288] ***Particle-based Assays***

[00289] As an alternative to planar arrays, assays using particles or microspheres, such as bead based assays, are capable of use with a binding agent. For example, antibodies or aptamers are easily conjugated with commercially available beads. *See, e.g.,* Fan *et al.*, Illumina universal bead arrays. Methods Enzymol. 2006 410:57-73; Srinivas *et al.* Anal. Chem. 2011 Oct. 21, *Aptamer functionalized Microgel Particles for Protein Detection*; See also, review article on aptamers as therapeutic and diagnostic agents, Brody and Gold, Rev. Mol. Biotech. 2000, 74:5-13.

[00290] Multiparametric assays or other high throughput detection assays using bead coatings with cognate ligands and reporter molecules with specific activities consistent with high sensitivity automation can be used. In a bead based assay system, a binding agent for a biomarker or vesicle, such as a capture agent (e.g. capture antibody), can be immobilized on an addressable microsphere. Each binding agent for each individual binding assay can be coupled to a distinct type of microsphere (i.e., microbead) and the assay reaction takes place on the surface of the microsphere, such as depicted in **FIG. 2B**. A binding agent for a vesicle can be a capture antibody or aptamer coupled to a bead. Dyed microspheres with discrete fluorescence intensities are loaded separately with their appropriate binding agent or capture probes. The different bead sets carrying different binding agents can be pooled as desired to generate custom bead arrays. Bead arrays are then incubated with the sample in a single reaction vessel to perform the assay.

[00291] Various particle/bead substrates and systems useful for the methods of the invention are described further above.

[00292] *Flow Cytometry*

[00293] In various embodiments of the invention, flow cytometry, which is described in further detail above, is used to assess a microvesicle population in a biological sample. If desired, the microvesicle population can be sorted from other particles (e.g., cell debris, protein aggregates, etc) in a sample by labeling the vesicles using one or more general vesicle marker. The general vesicle marker can be a marker in **Table 3**. Commonly used vesicle markers include tetraspanins such as CD9, CD63 and/or CD81. Vesicles comprising one or more tetraspanin are sometimes refered to as "Tet+" herein to indicate that the vesicles are tetraspanin-positive. The sorted microvesicles can be further assessed using methodology described herein. E.g., surface antigens on the sorted microvesicles can be detected using flow or other methods. In some embodiments, payload within the sorted microvesicles is assessed. As an illustrative example, a population of microvesicles is contacted with a labeled binding agent to a surface antigen of interest, the contacted microvesicles are sorted using flow cytometry, and payload with the microvesicles is assessed. The payload may be polypeptides, nucleic acids (e.g., mRNA or microRNA) or other biological entities as desired. Such assessment is used to characterize a phenotype as described herein, e.g., to diagnose, prognose or theranose a cancer.

[00294] In an embodiment, flow sorting is used to distinguish microvesicle populations from other biological complexes. In a non-limiting example, Ago2+/Tet+ and Ago2+/Tet- particles are detected using flow methodology to separate Ago2+ vesicles from vesicle-free Ago2+ complexes, respectively.

[00295] *Multiplexing*

[00296] Multiplex experiments comprise experiments that can simultaneously measure multiple analytes in a single assay. Vesicles and associated biomarkers can be assessed in a multiplex fashion. Different binding agents can be used for multiplexing different circulating biomarkers, e.g., microRNA, protein, or vesicle populations. Different biomarkers, e.g., different vesicle populations, can be isolated or detected using different binding agents. Each population in a biological sample can be labeled with a different signaling label, such as a fluorophore, quantum dot, or radioactive label, such as described above. The label can be directly conjugated to a binding agent or indirectly used to detect a binding agent that binds a vesicle. The number of populations detected in a multiplexing assay is dependent on the resolution capability of the labels and the summation of signals, as more than two differentially labeled vesicle populations that bind two or more affinity elements can produce summed signals.

[00297] Multiplexing of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 50, 75 or 100 different circulating biomarkers may be performed. For example, one population of vesicles specific to a cell-of-origin can be assayed along with a second population of vesicles specific to a different cell-of-origin, where each population is labeled with a different label. Alternatively, a population of vesicles with a particular biomarker or biosignature can be assayed along with a second population of vesicles with a

different biomarker or biosignature. In some cases, hundreds or thousands of vesicles are assessed in a single assay.

[00298] In one embodiment, multiplex analysis is performed by applying a plurality of vesicles comprising more than one population of vesicles to a plurality of substrates, such as beads. Each bead is coupled to one or more capture agents. The plurality of beads is divided into subsets, where beads with the same capture agent or combination of capture agents form a subset of beads, such that each subset of beads has a different capture agent or combination of capture agents than another subset of beads. The beads can then be used to capture vesicles that comprise a component that binds to the capture agent. The different subsets can be used to capture different populations of vesicles. The captured vesicles can then be analyzed by detecting one or more biomarkers.

[00299] Flow cytometry can be used in combination with a particle-based or bead based assay. Multiparametric immunoassays or other high throughput detection assays using bead coatings with cognate ligands and reporter molecules with specific activities consistent with high sensitivity automation can be used. For example, beads in each subset can be differentially labeled from another subset. In a particle based assay system, a binding agent or capture agent for a vesicle, such as a capture antibody, can be immobilized on addressable beads or microspheres. Each binding agent for each individual binding assay (such as an immunoassay when the binding agent is an antibody) can be coupled to a distinct type of microsphere (i.e., microbead) and the binding assay reaction takes place on the surface of the microspheres. Microspheres can be distinguished by different labels, for example, a microsphere with a specific capture agent would have a different signaling label as compared to another microsphere with a different capture agent. For example, microspheres can be dyed with discrete fluorescence intensities such that the fluorescence intensity of a microsphere with a specific binding agent is different than that of another microsphere with a different binding agent. Biomarkers bound by different capture agents can be differentially detected using different labels.

[00300] A microsphere can be labeled or dyed with at least 2 different labels or dyes. In some embodiments, the microsphere is labeled with at least 3, 4, 5, 6, 7, 8, 9, or 10 different labels. Different microspheres in a plurality of microspheres can have more than one label or dye, wherein various subsets of the microspheres have various ratios and combinations of the labels or dyes permitting detection of different microspheres with different binding agents. For example, the various ratios and combinations of labels and dyes can permit different fluorescent intensities. Alternatively, the various ratios and combinations maybe used to generate different detection patters to identify the binding agent. The microspheres can be labeled or dyed externally or may have intrinsic fluorescence or signaling labels. Beads can be loaded separately with their appropriate binding agents and thus, different vesicle populations can be isolated based on the different binding agents on the differentially labeled microspheres to which the different binding agents are coupled.

[00301] In another embodiment, multiplex analysis can be performed using a planar substrate, wherein the substrate comprises a plurality of capture agents. The plurality of capture agents can capture one or more

populations of vesicles, and one or more biomarkers of the captured vesicles detected. The planar substrate can be a microarray or other substrate as further described herein.

[00302] *Binding Agents*

[00303] A vesicle may be isolated or detected using a binding agent for a novel component of a vesicle, such as an antibody for a novel antigen specific to a vesicle of interest. Novel antigens that are specific to a vesicle of interest may be isolated or identified using different test compounds of known composition bound to a substrate, such as an array or a plurality of particles, which can allow a large amount of chemical/structural space to be adequately sampled using only a small fraction of the space. The novel antigen identified can also serve as a biomarker for the vesicle. For example, a novel antigen identified for a cell-of-origin specific vesicle can be a useful biomarker.

[00304] The term "agent" or "reagent" as used in respect to contacting a sample can mean any entity designed to bind, hybridize, associate with or otherwise detect or facilitate detection of a target molecule, including target polypeptides, peptides, nucleic acid molecules, leptins, lipids, or any other biological entity that can be detected as described herein or as known in the art. Examples of such agents/reagents are well known in the art, and include but are not limited to universal or specific nucleic acid primers, nucleic acid probes, antibodies, aptamers, peptoid, peptide nucleic acid, locked nucleic acid, lectin, dendrimer, chemical compound, or other entities described herein or known in the art.

[00305] A binding agent can be identified by screening either a homogeneous or heterogeneous vesicle population against test compounds. Since the composition of each test compound on the substrate surface is known, this constitutes a screen for affinity elements. For example, a test compound array comprises test compounds at specific locations on the substrate addressable locations, and can be used to identify one or more binding agents for a vesicle. The test compounds can all be unrelated or related based on minor variations of a core sequence or structure. The different test compounds may include variants of a given test compound (such as polypeptide isoforms), test compounds that are structurally or compositionally unrelated, or a combination thereof.

[00306] A test compound can be a peptoid, polysaccharide, organic compound, inorganic compound, polymer, lipids, nucleic acid, polypeptide, antibody, protein, polysaccharide, or other compound. The test compound can be natural or synthetic. The test compound can comprise or consist of linear or branched heteropolymeric compounds based on any of a number of linkages or combinations of linkages (e.g., amide, ester, ether, thiol, radical additions, metal coordination, etc.), dendritic structures, circular structures, cavity structures or other structures with multiple nearby sites of attachment that serve as scaffolds upon which specific additions are made. Thes test compound can be spotted on a substrate or synthesized in situ, using standard methods in the art. In addition, the test compound can be spotted or synthesized in situ in combinations in order to detect useful interactions, such as cooperative binding.

[00307] The test compound can be a polypeptide with known amino acid sequence, thus, detection of a test compound binding with a vesicle can lead to identification of a polypeptide of known amino sequence that can be used as a binding agent. For example, a homogenous population of vesicles can be applied to a

spotted array on a slide containing between a few and 1,000,000 test polypeptides having a length of variable amino acids. The polypeptides can be attached to the surface through the C-terminus. The sequence of the polypeptides can be generated randomly from 19 amino acids, excluding cysteine. The binding reaction can include a non-specific competitor, such as excess bacterial proteins labeled with another dye such that the specificity ratio for each polypeptide binding target can be determined. The polypeptides with the highest specificity and binding can be selected. The identity of the polypeptide on each spot is known, and thus can be readily identified. Once the novel antigens specific to the homogeneous vesicle population, such as a cell-of-origin specific vesicle is identified, such cell-of-origin specific vesicles may subsequently be isolated using such antigens in methods described hereafter.

[00308] An array can also be used for identifying an antibody as a binding agent for a vesicle. Test antibodies can be attached to an array and screened against a heterogeneous population of vesicles to identify antibodies that can be used to isolate or identify a vesicle. A homogeneous population of vesicles such as cell-of-origin specific vesicles can also be screened with an antibody array. Other than identifying antibodies to isolate or detect a homogeneous population of vesicles, one or more protein biomarkers specific to the homogenous population can be identified. Commercially available platforms with test antibodies pre-selected or custom selection of test antibodies attached to the array can be used. For example, an antibody array from Full Moon Biosystems can be screened using prostate cancer cell derived vesicles identifying antibodies to Bcl-XL, ERCC1, Keratin 15, CD81/TAPA-1, CD9, Epithelial Specific Antigen (ESA), and Mast Cell Chymase as binding agents, and the proteins identified can be used as biomarkers for the vesicles. The biomarker can be present or absent, underexpressed or overexpressed, mutated, or modified in or on a vesicle and used in characterizing a condition.

[00309] An antibody or synthetic antibody to be used as a binding agent can also be identified through a peptide array. Another method is the use of synthetic antibody generation through antibody phage display. M13 bacteriophage libraries of antibodies (e.g. Fabs) are displayed on the surfaces of phage particles as fusions to a coat protein. Each phage particle displays a unique antibody and also encapsulates a vector that contains the encoding DNA. Highly diverse libraries can be constructed and represented as phage pools, which can be used in antibody selection for binding to immobilized antigens. Antigen-binding phages are retained by the immobilized antigen, and the nonbinding phages are removed by washing. The retained phage pool can be amplified by infection of an *Escherichia coli* host and the amplified pool can be used for additional rounds of selection to eventually obtain a population that is dominated by antigen-binding clones. At this stage, individual phase clones can be isolated and subjected to DNA sequencing to decode the sequences of the displayed antibodies. Through the use of phase display and other methods known in the art, high affinity designer antibodies for vesicles can be generated.

[00310] Bead-based assays can also be used to identify novel binding agents to isolate or detect a vesicle. A test antibody or peptide can be conjugated to a particle. For example, a bead can be conjugated to an antibody or peptide and used to detect and quantify the proteins expressed on the surface of a population of vesicles in order to discover and specifically select for novel antibodies that can target vesicles from

specific tissue or tumor types. Any molecule of organic origin can be successfully conjugated to a polystyrene bead through use of a commercially available kit according to manufacturer's instructions. Each bead set can be colored a certain detectable wavelength and each can be linked to a known antibody or peptide which can be used to specifically measure which beads are linked to exosomal proteins matching the epitope of previously conjugated antibodies or peptides. The beads can be dyed with discrete fluorescence intensities such that each bead with a different intensity has a different binding agent as described above.

[00311] For example, a purified vesicle preparation can be diluted in assay buffer to an appropriate concentration according to empirically determined dynamic range of assay. A sufficient volume of coupled beads can be prepared and approximately 1 µl of the antibody-coupled beads can be aliqouted into a well and adjusted to a final volume of approximately 50 µl. Once the antibody-conjugated beads have been added to a vacuum compatible plate, the beads can be washed to ensure proper binding conditions. An appropriate volume of vesicle preparation can then be added to each well being tested and the mixture incubated, such as for 15-18 hours. A sufficient volume of detection antibodies using detection antibody diluent solution can be prepared and incubated with the mixture for 1 hour or more. The beads can then be washed before the addition of detection antibody (biotin expressing) mixture composed of streptavidin phycoereythin. The beads can then be washed and vacuum aspirated several times before analysis on a suspension array system using software provided with an instrument. The identity of antigens that can be used to selectively extract the vesicles can then be elucidated from the analysis.

[00312] Assays using imaging systems can be used to detect and quantify proteins expressed on the surface of a vesicle in order to discover and specifically select for and enrich vesicles from specific tissue, cell or tumor types. Antibodies, peptides or cells conjugated to multiple well multiplex carbon coated plates can be used. Simultaneous measurement of many analytes in a well can be achieved through the use of capture antibodies arrayed on the patterned carbon working surface. Analytes can then be detected with antibodies labeled with reagents in electrode wells with an enhanced electro-chemiluminescent plate. Any molecule of organic origin can be successfully conjugated to the carbon coated plate. Proteins expressed on the surface of vesicles can be identified from this assay and can be used as targets to specifically select for and enrich vesicles from specific tissue or tumor types.

[00313] The binding agent can also be an aptamer to a specific target. The term "specific" as used herein in regards to a binding agent can mean that an agent has a greater affinity for its target than other targets, typically with a much great affinity, but does not require that the binding agent is absolutely specific for its target.

[00314] ***Microfluidics***

[00315] The methods for isolating or identifying vesicles can be used in combination with microfluidic devices. The methods of isolating or detecting a vesicle, such as described herien, can be performed using a microfluidic device. Microfluidic devices, which may also be referred to as "lab-on-a-chip" systems,

biomedical micro-electro-mechanical systems (bioMEMs), or multicomponent integrated systems, can be used for isolating and analyzing a vesicle. Such systems miniaturize and compartmentalize processes that allow for binding of vesicles, detection of biosignatures, and other processes.

[00316] A microfluidic device can also be used for isolation of a vesicle through size differential or affinity selection. For example, a microfluidic device can use one more channels for isolating a vesicle from a biological sample based on size or by using one or more binding agents for isolating a vesicle from a biological sample. A biological sample can be introduced into one or more microfluidic channels, which selectively allows the passage of a vesicle. The selection can be based on a property of the vesicle, such as the size, shape, deformability, or biosignature of the vesicle.

[00317] In one embodiment, a heterogeneous population of vesicles can be introduced into a microfluidic device, and one or more different homogeneous populations of vesicles can be obtained. For example, different channels can have different size selections or binding agents to select for different vesicle populations. Thus, a microfluidic device can isolate a plurality of vesicles wherein at least a subset of the plurality of vesicles comprises a different biosignature from another subset of the plurality of vesicles. For example, the microfluidic device can isolate at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, or 100 different subsets of vesicles, wherein each subset of vesicles comprises a different biosignature.

[00318] In some embodiments, the microfluidic device can comprise one or more channels that permit further enrichment or selection of a vesicle. A population of vesicles that has been enriched after passage through a first channel can be introduced into a second channel, which allows the passage of the desired vesicle or vesicle population to be further enriched, such as through one or more binding agents present in the second channel.

[00319] Array-based assays and bead-based assays can be used with microfluidic device. For example, the binding agent can be coupled to beads and the binding reaction between the beads and vesicle can be performed in a microfluidic device. Multiplexing can also be performed using a microfluidic device. Different compartments can comprise different binding agents for different populations of vesicles, where each population is of a different cell-of-origin specific vesicle population. In one embodiment, each population has a different biosignature. The hybridization reaction between the microsphere and vesicle can be performed in a microfluidic device and the reaction mixture can be delivered to a detection device. The detection device, such as a dual or multiple laser detection system can be part of the microfluidic system and can use a laser to identify each bead or microsphere by its color-coding, and another laser can detect the hybridization signal associated with each bead.

[00320] Various microfluidic devices and methods are described above.

[00321] ***Combined Isolation Methodology***

[00322] One of skill will appreciate that various methods of sample treatment and isolating and concentrating circulating biomarkers such as vesicles can be combined as desired. For example, a biological sample can be treated to prevent aggregation, remove undesired particulate and/or deplete

highly abundant proteins. The steps used can be chosen to optimize downstream analysis steps. Next, biomarkers such as vesicles can be isolated, e.g., by chromotography, centrifugation, density gradient, filtration, precipitation, or affinity techniques. Any number of the later steps can be combined, e.g., a sample could be subjected to one or more of chromotography, centrifugation, density gradient, filtration and precipitation in order to isolate or concentrate all or most microvesicles. In a subsequent step, affinity techniques, e.g., using binding agents to one or more target of interest, can be used to isolate or identify microvesicles carrying desired biomarker profiles. Microfluidic systems can be employed to perform some or all of these steps.

[00323] An exemplary yet non-limiting isolation scheme for isolating and analysis of microvesicles includes the following: Plasma or serum collection -> highly abundant protein removal -> ultrafiltration -> nanomembrane concentration -> flow cytometry or particle-based assay.

[00324] Using the methods disclosed herein or known in the art, circulating biomarkers such as vesicles can be isolated or concentrated by at least about 2-fold, 3-fold, 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 12-fold, 15-fold, 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, 50-fold, 55-fold, 60-fold, 65-fold, 70-fold, 75-fold, 80-fold, 90-fold, 95-fold, 100-fold, 110-fold, 120-fold, 125-fold, 130-fold, 140-fold, 150-fold, 160-fold, 170-fold, 175-fold, 180-fold, 190-fold, 200-fold, 205-fold, 250-fold, 275-fold, 300-fold, 325-fold, 350-fold, 375-fold, 400-fold, 425-fold, 450-fold, 475-fold, 500-fold, 525-fold, 550-fold, 575-fold, 600-fold, 625-fold, 650-fold, 675-fold, 700-fold, 725-fold, 750-fold, 775-fold, 800-fold, 825-fold, 850-fold, 875-fold, 900-fold, 925-fold, 950-fold, 975-fold, 1000-fold, 1500-fold, 2000-fold, 2500-fold, 3000-fold, 4000-fold, 5000-fold, 6000-fold, 7000-fold, 8000-fold, 9000-fold, or at least 10,000-fold. In some embodiments, the vesicles are isolated or concentrated concentrated by at least 1 order of magnitude, 2 orders of magnitude, 3 orders of magnitude, 4 orders of magnitude, 5 orders of magnitude, 6 orders of magnitude, 7 orders of magnitude, 8 orders of magnitude, 9 orders of magnitude, or 10 orders of magnitude or more.

[00325] Once concentrated or isolated, the circulating biomarkers can be assessed, e.g., in order to characterize a phenotype as described herein. In some embodiments, the concentration or isolation steps themselves shed light on the phenotype of interest. For example, affinity methods can detect the presence or level of specific biomarkers of interest.

[00326] The various isolation and detection systems described herein can be used to isolate or detect circulating biomarkers such as vesicles that are informative for diagnosis, prognosis, disease stratification, theranosis, prediction of responder / non-responder status, disease monitoring, treatment monitoring and the like as related to such diseases and disorders. Combinations of the isolation techniques are within the scope of the invention. In a non-limiting example, a sample can be run through a chromatography column to isolate vesicles based on a property such as size of electrophoretic motility, and the vesicles can then be passed through a microfluidic device. Binding agents can be used before, during or after these steps.

[00327] The methods and compositions of the invention can be used with microvesicles isolated or detected using such methods as described herein. In various non-limiting examples: an aptamer provided

by the methods of the invention can be used as a capture and/or detector agent for a biomarker such as a protein or microvesicle; a sample such as a bodily fluid can be contacted with an oligonucleotide probe library of the invention before microvesicles in the sample are isolated using one or more technique described herein (e.g., chromatography, centrifugation, flow cytometry, filtration, affinity isolation, polymer precipitation, etc); microvesicles in a sample are isolated using one or more technique described herein (e.g., chromatography, centrifugation, flow cytometry, filtration, affinity isolation, polymer precipitation, etc) before contacting the microvesicles with an aptamer or oligonucleotide probe library of the invention. Contaminants such as highly abundant proteins can be removed in whole or in part at any appropriate step in such processes. These and various other useful iterations of such techniques for assessment of microvesicles and other biomarkers are contemplated by the invention.

Biomarkers

[00328] As described herein, the methods and compositions of the invention can be used in assays to detect the presence or level of one or more biomarker of interest. The biomarker can be any useful biomarker disclosed herein or known to those of skill in the art. In an embodiment, the biomarker comprises a protein or polypeptide. As used herein, "protein," "polypeptide" and "peptide" are used interchangeably unless stated otherwise. The biomarker can be a nucleic acid, including DNA, RNA, and various subspecies of any thereof as disclosed herein or known in the art. The biomarker can comprise a lipid. The biomarker can comprise a carbohydrate. The biomarker can also be a complex, e.g., a complex comprising protein, nucleic acids, lipids and/or carbohydrates. In some embodiments, the biomarker comprises a microvesicle. In an embodiment, the invention provides a method wherein a pool of aptamers is used to assess the presence and/or level of a population of microvesicles of interest without knowing the precise microvesicle antigen targeted by each member of the pool. See, e.g., **FIGs. 19B-C**. In other cases, biomarkers associated with microvesicles are assessed according to the methods of the invention. See, e.g., **FIGs. 2A-F**; **FIG. 19A**.

[00329] A biosignature comprising more than one biomarker can comprise one type of biomarker or multiple types of biomarkers. As a non-limiting example, a biosignature can comprise multiple proteins, multiple nucleic acids, multiple lipids, multiple carbohydrates, multiple biomarker complexes, multiple microvesicles, or a combination of any thereof. For example, the biosignature may comprise one or more microvesicle, one or more protein, and one or more microRNA, wherein the one or more protein and/or one or more microRNA is optionally in association with the microvesicle as a surface antigen and/or payload, as appropriate.

[00330] In some embodiments, vesicles are detected using vesicle surface antigens. A commonly expressed vesicle surface antigen can be referred to as a "housekeeping protein," or general vesicle biomarker. The biomarker can be CD63, CD9, CD81, CD82, CD37, CD53, Rab-5b, Annexin V or MFG-E8. Tetraspanins, a family of membrane proteins with four transmembrane domains, can be used as general vesicle biomarkers. The tetraspanins include CD151, CD53, CD37, CD82, CD81, CD9 and CD63. There have been over 30 tetraspanins identified in mammals, including the TSPAN1 (TSP-1), TSPAN2

(TSP-2), TSPAN3 (TSP-3), TSPAN4 (TSP-4, NAG-2), TSPAN5 (TSP-5), TSPAN6 (TSP-6), TSPAN7 (CD231, TALLA-1, A15), TSPAN8 (CO-029), TSPAN9 (NET-5), TSPAN10 (Oculospanin), TSPAN11 (CD151-like), TSPAN12 (NET-2), TSPAN13 (NET-6), TSPAN14, TSPAN15 (NET-7), TSPAN16 (TM4-B), TSPAN17, TSPAN18, TSPAN19, TSPAN20 (UP1b, UPK1B), TSPAN21 (UP1a, UPK1A), TSPAN22 (RDS, PRPH2), TSPAN23 (ROM1), TSPAN24 (CD151), TSPAN25 (CD53), TSPAN26 (CD37), TSPAN27 (CD82), TSPAN28 (CD81), TSPAN29 (CD9), TSPAN30 (CD63), TSPAN31 (SAS), TSPAN32 (TSSC6), TSPAN33, and TSPAN34. Other commonly observed vesicle markers include those listed in **Table 3**. One or more of these proteins can be useful biomarkers for the characterizing a phenotype using the subject methods and compositions.

Table 3: Proteins Observed in Vesicles from Multiple Cell Types

Class	Protein
Antigen Presentation	MHC class I, MHC class II, Integrins, Alpha 4 beta 1, Alpha M beta 2, Beta 2
Immunoglobulin family	ICAM1/CD54, P-selection
Cell-surface peptidases	Dipeptidylpeptidase IV/CD26, Aminopeptidase n/CD13
Tetraspanins	CD151, CD53, CD37, CD82, CD81, CD9 and CD63
Heat-shock proteins	Hsp70, Hsp84/90
Cytoskeletal proteins	Actin, Actin-binding proteins, Tubulin
Membrane transport and fusion	Annexin I, Annexin II, Annexin IV, Annexin V, Annexin VI, RAB7/RAP1B/RADGDI
Signal transduction	Gi2alpha/14-3-3, CBL/LCK
Abundant membrane proteins	CD63, GAPDH, CD9, CD81, ANXA2, ENO1, SDCBP, MSN, MFGE8, EZR, GK, ANXA1, LAMP2, DPP4, TSG101, HSPA1A, GDI2, CLTC, LAMP1, Cd86, ANPEP, TFRC, SLC3A2, RDX, RAP1B, RAB5C, RAB5B, MYH9, ICAM1, FN1, RAB11B, PIGR, LGALS3, ITGB1, EHD1, CLIC1, ATP1A1, ARF1, RAP1A, P4HB, MUC1, KRT10, HLA-A, FLOT1, CD59, C1orf58, BASP1, TACSTD1, STOM
Other Transmembrane Proteins	Cadherins: CDH1, CDH2, CDH12, CDH3, Deomoglein, DSG1, DSG2, DSG3, DSG4, Desmocollin, DSC1, DSC2, DSC3, Protocadherins, PCDH1, PCDH10, PCDH11x, PCDH11y, PCDH12, FAT, FAT2, FAT4, PCDH15, PCDH17, PCDH18, PCDH19; PCDH20; PCDH7, PCDH8, PCDH9, PCDHA1, PCDHA10, PCDHA11, PCDHA12, PCDHA13, PCDHA2, PCDHA3, PCDHA4, PCDHA5, PCDHA6, PCDHA7, PCDHA8, PCDHA9, PCDHAC1, PCDHAC2, PCDHB1, PCDHB10, PCDHB11, PCDHB12, PCDHB13, PCDHB14, PCDHB15, PCDHB16, PCDHB17, PCDHB18, PCDHB2, PCDHB3, PCDHB4, PCDHB5, PCDHB6, PCDHB7, PCDHB8, PCDHB9, PCDHGA1, PCDHGA10, PCDHGA11, PCDHGA12, PCDHGA2; PCDHGA3, PCDHGA4, PCDHGA5, PCDHGA6, PCDHGA7, PCDHGA8, PCDHGA9, PCDHGB1, PCDHGB2, PCDHGB3, PCDHGB4, PCDHGB5, PCDHGB6, PCDHGB7, PCDHGC3, PCDHGC4, PCDHGC5, CDH9 (cadherin 9, type 2 (T1-cadherin)), CDH10 (cadherin 10, type 2 (T2-cadherin)), CDH5 (VE-cadherin (vascular endothelial)), CDH6 (K-cadherin (kidney)), CDH7 (cadherin 7, type 2), CDH8 (cadherin 8, type 2), CDH11 (OB-cadherin (osteoblast)), CDH13 (T-cadherin - H-cadherin (heart)), CDH15 (M-cadherin (myotubule)), CDH16 (KSP-cadherin), CDH17 (LI cadherin (liver-intestine)), CDH18 (cadherin 18, type 2), CDH19 (cadherin 19, type 2), CDH20 (cadherin 20, type 2), CDH23 (cadherin 23, (neurosensory epithelium)), CDH10, CDH11, CDH13, CDH15, CDH16, CDH17, CDH18, CDH19, CDH22, CDH23, CDH24,

	CDH26, CDH28, CDH4, CDH5, CDH6, CDH7, CDH8, CDH9, CELSR1, CELSR2, CELSR3, CLSTN1, CLSTN2, CLSTN3, DCHS1, DCHS2, LOC389118, PCLKC, RESDA1, RET

[00331] Any of the types of biomarkers or specific biomarkers described herein can be used and/or assessed via the subject methods and compositions, e.g., to identify a useful biosignature. Exemplary biomarkers include without limitation those in **Table 4**. The markers can be detected as protein, RNA or DNA as appropriate, which can be circulating freely or in a complex with other biological molecules. As appropriate, the markers in **Table 4** can also be used for capture and/or detection of vesicles for characterizing phenotypes as disclosed herein. In some cases, multiple capture and/or detectors are used to enhance the characterization. The markers can be detected as protein or as mRNA, which can be circulating freely or in a complex with other biological molecules. See, e.g., **FIGs. 2D-E**. The markers can be detected as vesicle surface antigens and/or vesicle payload. The "Illustrative Class" indicates indications for which the markers are known markers. Those of skill will appreciate that the markers can also be used in alternate settings in certain instances. For example, a marker which can be used to characterize one type disease may also be used to characterize another disease as appropriate. Consider a non-limiting example of a tumor marker which can be used as a biomarker for tumors from various lineages. The biomarker references in **Tables 3** and **4** are those commonly used in the art. Gene aliases and descriptions can be found using a variety of online databases, including GeneCards® (www.genecards.org), HUGO Gene Nomenclature (www.genenames.org), Entrez Gene (www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene), UniProtKB/Swiss-Prot (www.uniprot.org), UniProtKB/TrEMBL (www.uniprot.org), OMIM (www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM), GeneLoc (genecards.weizmann.ac.il/geneloc/), and Ensembl (www.ensembl.org). Generally, gene symbols and names below correspond to those approved by HUGO, and protein names are those recommended by UniProtKB/Swiss-Prot. Common alternatives are provided as well. In some cases, biomarkers are referred to by Ensembl reference numbers, which are of the form "ENSG" followed by a number, e.g., ENSG00000005893 which corresponds to LAMP2. In **Table 4**, solely for sake of brevity, "E." is sometimes used to represent "ENSG00000". For example, "E.005893 represents "ENSG00000005893." Where a protein name indicates a precursor, the mature protein is also implied. Throughout the application, gene and protein symbols may be used interchangeably and the meaning can be derived from context as necessary.

Table 4: Illustrative Biomarkers

Illustrative Class	Biomarkers
Drug associated targets and prognostic markers	ABCC1, ABCG2, ACE2, ADA, ADH1C, ADH4, AGT, AR, AREG, ASNS, BCL2, BCRP, BDCA1, beta III tubulin, BIRC5, B-RAF, BRCA1, BRCA2, CA2, caveolin, CD20, CD25, CD33, CD52, CDA, CDKN2A, CDKN1A, CDKN1B, CDK2, CDW52, CES2, CK 14, CK 17, CK 5/6, c-KIT, c-Met, c-Myc, COX-2, Cyclin D1, DCK, DHFR, DNMT1, DNMT3A, DNMT3B, E-Cadherin, ECGF1, EGFR, EML4-ALK fusion, EPHA2, Epiregulin, ER, ERBR2, ERCC1, ERCC3, EREG, ESR1, FLT1, folate receptor, FOLR1, FOLR2, FSHB, FSHPRH1, FSHR, FYN, GART, GNA11, GNAQ, GNRH1, GNRHR1, GSTP1, HCK, HDAC1, hENT-1, Her2/Neu, HGF, HIF1A, HIG1, HSP90, HSP90AA1, HSPCA, IGF-1R, IGFRBP, IGFRBP3, IGFRBP4, IGFRBP5, IL13RA1, IL2RA, KDR, Ki67, KIT, K-RAS,

	LCK, LTB, Lymphotoxin Beta Receptor, LYN, MET, MGMT, MLH1, MMR, MRP1, MS4A1, MSH2, MSH5, Myc, NFKB1, NFKB2, NFKBIA, NRAS, ODC1, OGFR, p16, p21, p27, p53, p95, PARP-1, PDGFC, PDGFR, PDGFRA, PDGFRB, PGP, PGR, PI3K, POLA, POLA1, PPARG, PPARGC1, PR, PTEN, PTGS2, PTPN12, RAF1, RARA, ROS1, RRM1, RRM2, RRM2B, RXRB, RXRG, SIK2, SPARC, SRC, SSTR1, SSTR2, SSTR3, SSTR4, SSTR5, Survivin, TK1, TLE3, TNF, TOP1, TOP2A, TOP2B, TS, TUBB3, TXN, TXNRD1, TYMS, VDR, VEGF, VEGFA, VEGFC, VHL, YES1, ZAP70
Drug associated targets and prognostic markers	ABL1, STK11, FGFR2, ERBB4, SMARCB1, CDKN2A, CTNNB1, FGFR1, FLT3, NOTCH1, NPM1, SRC, SMAD4, FBXW7, PTEN, TP53, AKT1, ALK, APC, CDH1, C-Met, HRAS, IDH1, JAK2, MPL, PDGFRA, SMO, VHL, ATM, CSF1R, FGFR3, GNAS, ERBB2, HNF1A, JAK3, KDR, MLH1, PTPN11, RB1, RET, c-Kit, EGFR, PIK3CA, NRAS, GNA11, GNAQ, KRAS, BRAF
Drug associated targets and prognostic markers	ALK, AR, BRAF, cKIT, cMET, EGFR, ER, ERCC1, GNA11, HER2, IDH1, KRAS, MGMT, MGMT promoter methylation, NRAS, PDGFRA, Pgp, PIK3CA, PR, PTEN, ROS1, RRM1, SPARC, TLE3, TOP2A, TOPO1, TS, TUBB3, VHL
Drug associated targets	ABL1, AKT1, ALK, APC, AR, ATM, BRAF, BRAF, BRCA1, BRCA2, CDH1, cKIT, cMET, CSF1R, CTNNB1, EGFR, EGFR (H-score), EGFRvIII, ER, ERBB2 (HER2), ERBB4, ERCC1, FBXW7, FGFR1, FGFR2, FLT3, GNA11, GNAQ, GNAS, HER2, HNF1A, HRAS, IDH1, IDH2, JAK2, JAK3, KDR (VEGFR2), KRAS, MGMT, MGMT Promoter Methylation, microsatellite instability (MSI), MLH1, MPL, MSH2, MSH6, NOTCH1, NPM1, NRAS, PD-1, PDGFRA, PD-L1, Pgp, PIK3CA, PMS2, PR, PTEN, PTPN11, RB1, RET, ROS1, RRM1, SMAD4, SMARCB1, SMO, SPARC, STK11, TLE3, TOP2A, TOPO1, TP53, TS, TUBB3, VHL
Drug associated targets	1p19q co-deletion, ABL1, AKT1, ALK, APC, AR, ARAF, ATM, BAP1, BRAF, BRCA1, BRCA2, CDH1, CHEK1, CHEK2, cKIT, cMET, CSF1R, CTNNB1, DDR2, EGFR, EGFRvIII, ER, ERBB2 (HER2), ERBB3, ERBB4, ERCC1, FBXW7, FGFR1, FGFR2, FLT3, GNA11, GNAQ, GNAS, H3K36me3, HER2, HNF1A, HRAS, IDH1, IDH2, JAK2, JAK3, KDR (VEGFR2), KRAS, MDMT, MGMT, MGMT Methylation, Microsatellite instability, MLH1, MPL, MSH2, MSH6, NF1, NOTCH1, NPM1, NRAS, NY-ESO-1, PD-1, PDGFRA, PD-L1, Pgp, PIK3CA, PMS2, PR, PTEN, PTPN11, RAF1, RB1, RET, ROS1, ROS1, RRM1, SMAD4, SMARCB1, SMO, SPARC, STK11, TLE3, TOP2A, TOPO1, TP53, TRKA, TS, TUBB3, VHL, WT1
Drug associated targets	ABL1, AKT1, ALK, APC, AR, ATM, BRAF, BRAF, BRCA1, BRCA2, CDH1, cKIT, cMET, CSF1R, CTNNB1, EGFR, EGFR (H-score), EGFRvIII, ER, ERBB2 (HER2), ERBB4, ERCC1, FBXW7, FGFR1, FGFR2, FLT3, GNA11, GNAQ, GNAS, HER2, HNF1A, HRAS, IDH1, IDH2, JAK2, JAK3, KDR (VEGFR2), KRAS, MGMT, MGMT Promoter Methylation, microsatellite instability (MSI), MLH1, MPL, MSH2, MSH6, NOTCH1, NPM1, NRAS, PD-1, PDGFRA, PD-L1, Pgp, PIK3CA, PMS2, PR, PTEN, PTPN11, RB1, RET, ROS1, RRM1, SMAD4, SMARCB1, SMO, SPARC, STK11, TLE3, TOP2A, TOPO1, TP53, TS, TUBB3, VHL
Drug associated targets	1p19q, ALK, ALK (2p23), Androgen Receptor, BRCA, cMET, EGFR, EGFR, EGFRvIII, ER, ERCC1, Her2, Her2/Neu, MGMT, MGMT Promoter Methylation, microsatellite instability (MSI), MLH1, MSH2, MSH6, PD-1, PD-L1, PMS2, PR, PTEN, ROS1, RRM1, TLE3, TOP2A, TOP2A, TOPO1, TS, TUBB3
Drug associated targets	TOP2A, Chromosome 17 alteration, PBRM1 (PB1/BAF180), BAP1, SETD2 (ANTI-HISTONE H3), MDM2, Chromosome 12 alteration, ALK, CTLA4, CD3, NY-ESO-1, MAGE-A, TP, EGFR
5-aminosalicyclic acid (5-ASA) efficacy	μ-protocadherin, KLF4, CEBPα
Cancer treatment associated markers	AR, AREG (Amphiregulin), BRAF, BRCA1, cKIT, cMET, EGFR, EGFR w/T790M, EML4-ALK, ER, ERBB3, ERBB4, ERCC1, EREG, GNA11, GNAQ, hENT-1, Her2, Her2 Exon 20 insert, IGF1R, Ki67, KRAS, MGMT, MGMT methylation, MSH2, MSI, NRAS, PGP (MDR1), PIK3CA, PR, PTEN, ROS1, ROS1 translocation, RRM1, SPARC, TLE3, TOPO1, TOPO2A, TS, TUBB3, VEGFR2
Cancer treatment associated markers	AR, AREG, BRAF, BRCA1, cKIT, cMET, EGFR, EGFR w/T790M, EML4-ALK, ER, ERBB3, ERBB4, ERCC1, EREG, GNA11, GNAQ, Her2, Her2 Exon 20 insert, IGFR1, Ki67, KRAS, MGMT-Me, MSH2, MSI, NRAS, PGP (MDR-1), PIK3CA, PR, PTEN, ROS1 translocation, RRM1, SPARC, TLE3, TOPO1, TOPO2A, TS, TUBB3, VEGFR2
Colon cancer treatment	AREG, BRAF, EGFR, EML4-ALK, ERCC1, EREG, KRAS, MSI, NRAS, PIK3CA, PTEN, TS, VEGFR2

associated markers	
Colon cancer treatment associated markers	AREG, BRAF, EGFR, EML4-ALK, ERCC1, EREG, KRAS, MSI, NRAS, PIK3CA, PTEN, TS, VEGFR2
Melanoma treatment associated markers	BRAF, cKIT, ERBB3, ERBB4, ERCC1, GNA11, GNAQ, MGMT, MGMT methylation, NRAS, PIK3CA, TUBB3, VEGFR2
Melanoma treatment associated markers	BRAF, cKIT, ERBB3, ERBB4, ERCC1, GNA11, GNAQ, MGMT-Me, NRAS, PIK3CA, TUBB3, VEGFR2
Ovarian cancer treatment associated markers	BRCA1, cMET, EML4-ALK, ER, ERBB3, ERCC1, hENT-1, HER2, IGF1R, PGP(MDR1), PIK3CA, PR, PTEN, RRM1, TLE3, TOPO1, TOPO2A, TS
Ovarian cancer treatment associated markers	BRCA1, cMET, EML4-ALK (translocation), ER, ERBB3, ERCC1, HER2, PIK3CA, PR, PTEN, RRM1, TLE3, TS
Breast cancer treatment associated markers	BRAF, BRCA1, EGFR, EGFR T790M, EML4-ALK, ER, ERBB3, ERCC1, HER2, Ki67, PGP (MDR1), PIK3CA, PR, PTEN, ROS1, ROS1 translocation, RRM1, TLE3, TOPO1, TOPO2A, TS
Breast cancer treatment associated markers	BRAF, BRCA1, EGFR w/T790M, EML4-ALK, ER, ERBB3, ERCC1, HER2, Ki67, KRAS, PIK3CA, PR, PTEN, ROS1 translocation, RRM1, TLE3, TOPO1, TOPO2A, TS
NSCLC cancer treatment associated markers	BRAF, BRCA1, cMET, EGFR, EGFR w/T790M, EML4-ALK, ERCC1, Her2 Exon 20 insert, KRAS, MSH2, PIK3CA, PTEN, ROS1 (trans), RRM1, TLE3, TS, VEGFR2
NSCLC cancer treatment associated markers	BRAF, cMET, EGFR, EGFR w/T790M, EML4-ALK, ERCC1, Her2 Exon 20 insert, KRAS, MSH2, PIK3CA, PTEN, ROS1 translocation, RRM1, TLE3, TS
Mutated in cancers	AKT1, ALK, APC, ATM, BRAF, CDH1, CDKN2A, c-Kit, C-Met, CSF1R, CTNNB1, EGFR, ERBB2, ERBB4, FBXW7, FGFR1, FGFR2, FGFR3, FLT3, GNA11, GNAQ, GNAS, HNF1A, HRAS, IDH1, JAK2, JAK3, KDR, KRAS, MLH1, MPL, NOTCH1, NPM1, NRAS, PDGFRA, PIK3CA, PTEN, PTPN11, RB1, RET, SMAD4, SMARCB1, SMO, SRC, STK11, TP53, VHL
Mutated in cancers	ALK, BRAF, BRCA1, BRCA2, EGFR, ERRB2, GNA11, GNAQ, IDH1, IDH2, KIT, KRAS, MET, NRAS, PDGFRA, PIK3CA, PTEN, RET, SRC, TP53
Mutated in cancers	AKT1, HRAS, GNAS, MEK1, MEK2, ERK1, ERK2, ERBB3, CDKN2A, PDGFRB, IFG1R, FGFR1, FGFR2, FGFR3, ERBB4, SMO, DDR2, GRB1, PTCH, SHH, PD1, UGT1A1, BIM, ESR1, MLL, AR, CDK4, SMAD4
Mutated in cancers	ABL, APC, ATM, CDH1, CSFR1, CTNNB1, FBXW7, FLT3, HNF1A, JAK2, JAK3, KDR, MLH1, MPL, NOTCH1, NPM1, PTPN11, RB1, SMARCB1, STK11, VHL
Mutated in cancers	ABL1, AKT1, AKT2, AKT3, ALK, APC, AR, ARAF, ARFRP1, ARID1A, ARID2, ASXL1, ATM, ATR, ATRX, AURKA, AURKB, AXL, BAP1, BARD1, BCL2, BCL2L2, BCL6, BCOR, BCORL1, BLM, BRAF, BRCA1, BRCA2, BRIP1, BTK, CARD11, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD79A, CD79B, CDC73, CDH1, CDK12, CDK4, CDK6, CDK8, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CEBPA, CHEK1, CHEK2, CIC, CREBBP, CRKL, CRLF2, CSF1R, CTCF, CTNNA1, CTNNB1, DAXX, DDR2, DNMT3A, DOT1L, EGFR, EMSY (C11orf30), EP300, EPHA3, EPHA5, EPHB1, ERBB2, ERBB3, ERBB4, ERG, ESR1, EZH2, FAM123B (WTX), FAM46C, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCL, FBXW7, FGF10, FGF14, FGF19, FGF23, FGF3, FGF4, FGF6, FGFR1, FGFR2, FGFR3, FGFR4, FLT1, FLT3, FLT4, FOXL2, GATA1, GATA2, GATA3, GID4 (C17orf39), GNA11, GNA13, GNAQ, GNAS, GPR124, GRIN2A, GSK3B, HGF, HRAS, IDH1, IDH2, IGF1R, IKBKE, IKZF1, IL7R, INHBA, IRF4, IRS2, JAK1, JAK2, JAK3, JUN, KAT6A (MYST3), KDM5A, KDM5C, KDM6A, KDR, KEAP1, KIT, KLHL6, KRAS, LRP1B, MAP2K1, MAP2K2, MAP2K4, MAP3K1, MCL1, MDM2, MDM4, MED12, MEF2B, MEN1, MET, MITF, MLH1, MLL, MLL2, MPL, MRE11A, MSH2, MSH6, MTOR, MUTYH, MYC, MYCL1, MYCN, MYD88, NF1, NF2, NFE2L2, NFKBIA, NKX2-1, NOTCH1, NOTCH2, NPM1, NRAS, NTRK1, NTRK2, NTRK3, NUP93, PAK3, PALB2, PAX5, PBRM1, PDGFRA, PDGFRB, PDK1, PIK3CA, PIK3CG, PIK3R1, PIK3R2, PPP2R1A, PRDM1, PRKAR1A, PRKDC, PTCH1, PTEN, PTPN11, RAD50, RAD51, RAF1, RARA, RB1, RET, RICTOR, RNF43, RPTOR, RUNX1, SETD2, SF3B1, SMAD2, SMAD4, SMARCA4, SMARCB1, SMO, SOCS1, SOX10, SOX2, SPEN,

	SPOP, SRC, STAG2, STAT4, STK11, SUFU, TET2, TGFBR2, TNFAIP3, TNFRSF14, TOP1, TP53, TSC1, TSC2, TSHR, VHL, WISP3, WT1, XPO1, ZNF217, ZNF703
Gene rearrangement in cancer	ALK, BCR, BCL2, BRAF, EGFR, ETV1, ETV4, ETV5, ETV6, EWSR1, MLL, MYC, NTRK1, PDGFRA, RAF1, RARA, RET, ROS1, TMPRSS2
Cancer Related	ABL1, ACE2, ADA, ADH1C, ADH4, AGT, AKT1, AKT2, AKT3, ALK, APC, AR, ARAF, AREG, ARFRP1, ARID1A, ARID2, ASNS, ASXL1, ATM, ATR, ATRX, AURKA, AURKB, AXL, BAP1, BARD1, BCL2, BCL2L2, BCL6, BCOR, BCORL1, BCR, BIRC5 (survivin), BLM, BRAF, BRCA1, BRCA2, BRIP1, BTK, CA2, CARD11, CAV, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD33, CD52 (CDW52), CD79A, CD79B, CDC73, CDH1, CDK12, CDK2, CDK4, CDK6, CDK8, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CEBPA, CES2, CHEK1, CHEK2, CIC, CREBBP, CRKL, CRLF2, CSF1R, CTCF, CTNNA1, CTNNB1, DAXX, DCK, DDR2, DHFR, DNMT1, DNMT3A, DNMT3B, DOT1L, EGFR, EMSY (C11orf30), EP300, EPHA2, EPHA3, EPHA5, EPHB1, ERBB2, ERBB3, ERBB4, ERBR2 (typo?), ERCC3, EREG, ERG, ESR1, ETV1, ETV4, ETV5, ETV6, EWSR1, EZH2, FAM123B (WTX), FAM46C, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCL, FBXW7, FGF10, FGF14, FGF19, FGF23, FGF3, FGF4, FGF6, FGFR1, FGFR2, FGFR3, FGFR4, FLT1, FLT3, FLT4, FOLR1, FOLR2, FOXL2, FSHB, FSHPRH1, FSHR, GART, GATA1, GATA2, GATA3, GID4 (C17orf39), GNA11, GNA13, GNAQ, GNAS, GNRH1, GNRHR1, GPR124, GRIN2A, GSK3B, GSTP1, HDAC1, HGF, HIG1, HNF1A, HRAS, HSPCA (HSP90), IDH1, IDH2, IGF1R, IKBKE, IKZF1, IL13RA1, IL2, IL2RA (CD25), IL7R, INHBA, IRF4, IRS2, JAK1, JAK2, JAK3, JUN, KAT6A (MYST3), KDM5A, KDM5C, KDM6A, KDR (VEGFR2), KEAP1, KIT, KLHL6, KRAS, LCK, LRP1B, LTB, LTBR, MAP2K1, MAP2K2, MAP2K4, MAP3K1, MAPK, MCL1, MDM2, MDM4, MED12, MEF2B, MEN1, MET, MGMT, MITF, MLH1, MLL, MLL2, MPL, MRE11A, MS4A1 (CD20), MSH2, MSH6, MTAP, MTOR, MUTYH, MYC, MYCL1, MYCN, MYD88, NF1, NF2, NFE2L2, NFKB1, NFKB2, NFKBIA, NGF, NKX2-1, NOTCH1, NOTCH2, NPM1, NRAS, NTRK1, NTRK2, NTRK3, NUP93, ODC1, OGFR, PAK3, PALB2, PAX5, PBRM1, PDGFC, PDGFRA, PDGFRB, PDK1, PGP, PGR (PR), PIK3CA, PIK3CG, PIK3R1, PIK3R2, POLA, PPARG, PPARGC1, PPP2R1A, PRDM1, PRKAR1A, PRKDC, PTCH1, PTEN, PTPN11, RAD50, RAD51, RAF1, RARA, RB1, RET, RICTOR, RNF43, ROS1, RPTOR, RRM1, RRM2, RRM2B, RUNX1, RXR, RXRB, RXRG, SETD2, SF3B1, SMAD2, SMAD4, SMARCA4, SMARCB1, SMO, SOCS1, SOX10, SOX2, SPARC, SPEN, SPOP, SRC, SST, SSTR1, SSTR2, SSTR3, SSTR4, SSTR5, STAG2, STAT4, STK11, SUFU, TET2, TGFBR2, TK1, TLE3, TMPRSS2, TNF, TNFAIP3, TNFRSF14, TOP1, TOP2, TOP2A, TOP2B, TP53, TS, TSC1, TSC2, TSHR, TUBB3, TXN, TYMP, VDR, VEGF (VEGFA), VEGFC, VHL, WISP3, WT1, XDH, XPO1, YES1, ZAP70, ZNF217, ZNF703
Cancer Related	5T4, ABI1, ABL1, ABL2, ACKR3, ACSL3, ACSL6, ACVR1B, ACVR2A, AFF1, AFF3, AFF4, AKAP9, AKT1, AKT2, AKT3, ALDH2, ALK, AMER1, ANG1/ANGPT1/TM7SF2, ANG2/ANGPT2/VPS51, APC, AR, ARAF, ARFRP1, ARHGAP26, ARHGEF12, ARID1A, ARID1B, ARID2, ARNT, ASPSCR1, ASXL1, ATF1, ATIC, ATM, ATP1A1, ATP2B3, ATR, ATRX, AURKA, AURKB, AXIN1, AXL, BAP1, BARD1, BBC3, BCL10, BCL11A, BCL11B, BCL2, BCL2L1, BCL2L11, BCL2L2, BCL3, BCL6, BCL7A, BCL9, BCOR, BCORL1, BCR, BIRC3, BLM, BMPR1A, BRAF, BRCA1, BRCA2, BRD3, BRD4, BRIP1, BTG1, BTK, BUB1B, c-KIT, C11orf30, c15orf21, C15orf65, C2orf44, CA6, CACNA1D, CALR, CAMTA1, CANT1, CARD11, CARS, CASC5, CASP8, CBFA2T3, CBFB, CBL, CBLB, CBLC, CCDC6, CCNB1IP1, CCND1, CCND2, CCND3, CCNE1, CD110, CD123, CD137, CD19, CD20, CD274, CD27L, CD38, CD4, CD74, CD79A, CD79B, CDC73, CDH1, CDH11, CDK12, CDK4, CDK6, CDK7, CDK8, CDK9, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CDX2, CEBPA, CHCHD7, CHD2, CHD4, CHEK1, CHEK2, CHIC2, Chk1, CHN1, CIC, CIITA, CLP1, CLTC, CLTCL1, CNBP, CNOT3, CNTRL, COL1A1, COPB1, CoREST, COX6C, CRAF, CREB1, CREB3L1, CREB3L2, CREBBP, CRKL, CRLF2, CRTC1, CRTC3, CSF1R, CSF3R, CTCF, CTLA4, CTNNA1, CTNNB1, CUL3, CXCR4, CYLD, CYP17A1, CYP2D6, DAXX, DDB2, DDIT3, DDR1, DDR2, DDX10, DDX5, DDX6, DEK, DICER1, DLL-4, DNAPK, DNM2, DNMT3A, DOT1L, EBF1, ECT2L, EGFR, EIF4A2, ELF4, ELK4, ELL, ELN, EML4, EP300, EPHA3, EPHA5, EPHA7, EPHA8, EPHB1, EPHB2, EPS15, ERBB2, ERBB3, ERBB4, ERC1, ERCC1, ERCC2, ERCC3, ERCC4, ERCC5, ERG, ERRFI1, ESR1, ETBR, ETV1, ETV4, ETV5, ETV6, EWSR1, EXT1, EXT2, EZH2, EZR, FAK, FAM46C, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCL, FAS, FAT1, FBXO11, FBXW7, FCRL4,

	FEV, FGF10, FGF14, FGF19, FGF2, FGF23, FGF3, FGF4, FGF6, FGFR1, FGFR1OP, FGFR2, FGFR3, FGFR4, FH, FHIT, FIP1L1, FKBP12, FLCN, FLI1, FLT1, FLT3, FLT4, FNBP1, FOXA1, FOXL2, FOXO1, FOXO3, FOXO4, FOXP1, FRS2, FSTL3, FUBP1, FUS, GABRA6, GAS7, GATA1, GATA2, GATA3, GATA4, GATA6, GID4, GITR, GLI1, GMPS, GNA11, GNA13, GNAQ, GNAS, GNRH1, GOLGA5, GOPC, GPC3, GPHN, GPR124, GRIN2A, GRM3, GSK3B, GUCY2C, H3F3A, H3F3B, HCK, HERPUD1, HEY1, HGF, HIP1, HIST1H3B, HIST1H4I, HLF, HMGA1, HMGA2, HMT, HNF1A, HNRNPA2B1, HOOK3, HOXA11, HOXA13, HOXA9, HOXC11, HOXC13, HOXD11, HOXD13, HRAS, HSD3B1, HSP90AA1, HSP90AB1, IAP, IDH1, IDH2, IGF1R, IGF2, IKBKE, IKZF1, IL2, IL21R, IL6, IL6ST, IL7R, INHBA, INPP4B, IRF2, IRF4, IRS2, ITGAV, ITGB1, ITK, JAK1, JAK2, JAK3, JAZF1, JUN, KAT6A, KAT6B, KCNJ5, KDM5A, KDM5C, KDM6A, KDR, KDSR, KEAP1, KEL, KIAA1549, KIF5B, KIR3DL1, KLF4, KLHL6, KLK2, KMT2A, KMT2C, KMT2D, KRAS, KTN1, LASP1, LCK, LCP1, LGALS3, LGR5, LHFP, LIFR, LMO1, LMO2, LOXL2, LPP, LRIG3, LRP1B, LSD1, LYL1, LYN, LZTR1, MAF, MAFB, MAGI2, MALT1, MAML2, MAP2K1, MAP2K2, MAP2K4, MAP3K1, MAPK1, MAPK11, MAX, MCL1, MDM2, MDM4, MDS2, MECOM, MED12, MEF2B, MEK1, MEK2, MEN1, MET, MITF, MKL1, MLF1, MLH1, MLLT1, MLLT10, MLLT11, MLLT3, MLLT4, MLLT6, MMP9, MN1, MNX1, MPL, MPS1, MRE11A, MS4A1, MSH2, MSH6, MSI2, MSN, MST1R, MTCP1, MTOR, MUC1, MUC16, MUTYH, MYB, MYC, MYCL, MYCN, MYD88, MYH11, MYH9, NACA, NAE1, NBN, NCKIPSD, NCOA1, NCOA2, NCOA4, NDRG1, NF1, NF2, NFE2L2, NFIB, NFKB2, NFKBIA, NIN, NKX2-1, NONO, NOTCH1, NOTCH2, NOTCH3, NPM1, NR4A3, NRAS, NSD1, NT5C2, NTRK1, NTRK2, NTRK3, NUMA1, NUP214, NUP93, NUP98, NUTM1, NUTM2B, OLIG2, OMD, P2RY8, PAFAH1B2, PAK3, PALB2, PARK2, PARP1, PATZ1, PAX3, PAX5, PAX7, PAX8, PBRM1, PBX1, PCM1, PCSK7, PDCD1, PDCD1LG2, PDE4DIP, PDGFB, PDGFRA, PDGFRB, PDK1, PER1, PHF6, PHOX2B, PICALM, PIK3C2B, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PIK3R1, PIK3R2, PIM1, PKC, PLAG1, PLCG2, PML, PMS1, PMS2, POLD1, POLE, POT1, POU2AF1, POU5F1, PPARG, PPP2R1A, PRCC, PRDM1, PRDM16, PREX2, PRF1, PRKAR1A, PRKCI, PRKDC, PRLR, PRRX1, PRSS8, PSIP1, PTCH1, PTEN, PTK2, PTPN11, PTPRC, PTPRD, QKI, RABEP1, RAC1, RAD21, RAD50, RAD51, RAD51B, RAF1, RALGDS, RANBP17, RANBP2, RANKL, RAP1GDS1, RARA, RB1, RBM10, RBM15, RECQL4, REL, RET, RHOH, RICTOR, RMI2, RNF213, RNF43, ROS1, RPL10, RPL20, RPL5, RPN1, RPS6KB1, RPTOR, RUNX1, RUNx1T1, SBDS, SDC4, SDHA, SDHAF2, SDHB, SDHC, SDHD, SEPT5, SEPT6, SEPT9, SET, SETBP1, SETD2, SF3B1, SFPQ, SH2B3, SH3GL1, SLAMF7, SLC34A2, SLC45A3, SLIT2, SMAD2, SMAD3, SMAD4, SMARCA4, SMARCB1, SMARCE1, SMO, SNCAIP, SNX29, SOCS1, SOX10, SOX2, SOX9, SPECC1, SPEN, SPOP, SPTA1, SRC, SRGAP3, SRSF2, SRSF3, SS18, SS18L1, SSX1, SSX2, SSX4, STAG2, STAT3, STAT4, STAT5B, STEAP1, STIL, STK11, SUFU, SUZ12, SYK, TAF1, TAF15, TAL1, TAL2, TBL1XR1, TBX3, TCEA1, TCF12, TCF3, TCF7L2, TCL1A, TERC, TERT, TET1, TET2, TFE3, TFEB, TFG, TFPT, TFRC, TGFB1, TGFBR2, THRAP3, TIE2, TLX1, TLX3, TMPRSS2, TNFAIP3, TNFRSF14, TNFRSF17, TOP1, TOP2A, TP53, TPM3, TPM4, TPR, TRAF7, TRIM26, TRIM27, TRIM33, TRIP11, TRRAP, TSC1, TSC2, TSHR, TTL, U2AF1, UBA1, UBR5, USP6, VEGFA, VEGFB, VEGFR, VHL, VTI1A, WAS, WEE1, WHSC1, WHSC1L1, WIF1, WISP3, WNT11, WNT2B, WNT3, WNT3A, WNT4, WNT5A, WNT6, WNT7B, WRN, WT1, WWTR1, XPA, XPC, XPO1, YWHAE, ZAK, ZBTB16, ZBTB2, ZMYM2, ZNF217, ZNF331, ZNF384, ZNF521, ZNF703, ZRSR2
Cancer Related	ABL2, ACSL3, ACSL6, AFF1, AFF3, AFF4, AKAP9, AKT3, ALDH2, APC, ARFRP1, ARHGAP26, ARHGEF12, ARID2, ARNT, ASPSCR1, ASXL1, ATF1, ATIC, ATM, ATP1A1, ATR, AURKA, AXIN1, AXL, BAP1, BARD1, BCL10, BCL11A, BCL2L11, BCL3, BCL6, BCL7A, BCL9, BCR, BIRC3, BLM, BMPR1A, BRAF, BRCA1, BRCA2, BRIP1, BUB1B, C11orf30, C2orf44, CACNA1D, CALR, CAMTA1, CANT1, CARD11, CARS, CASC5, CASP8, CBFA2T3, CBFB, CBL, CBLB, CCDC6, CCNB1IP1, CCND2, CD274, CD74, CD79A, CDC73, CDH11, CDKN1B, CDX2, CHEK1, CHEK2, CHIC2, CHN1, CIC, CIITA, CLP1, CLTC, CLTCL1, CNBP, CNTRL, COPB1, CREB1, CREB3L1, CREB3L2, CRTC1, CRTC3, CSF1R, CSF3R, CTCF, CTLA4, CTNNA1, CTNNB1, CYLD, CYP2D6, DAXX, DDR2, DDX10, DDX5, DDX6, DEK, DICER1, DOT1L, EBF1, ECT2L, ELK4, ELL, EML4, EPHA3, EPHA5, EPHB1, EPS15, ERBB3, ERBB4, ERC1, ERCC2, ERCC3, ERCC4, ERCC5, ERG, ESR1, ETV1, ETV5, ETV6, EWSR1, EXT1, EXT2, EZR, FANCA, FANCC, FANCD2, FANCE, FANCG, FANCL, FAS, FBXO11, FBXW7, FCRL4, FGF14, FGF19, FGF23, FGF6, FGFR1OP, FGFR4, FH, FHIT, FIP1L1, FLCN, FLI1, FLT1,

	FLT3, FLT4, FNBP1, FOXA1, FOXO1, FOXP1, FUBP1, FUS, GAS7, GID4, GMPS, GNA13, GNAQ, GNAS, GOLGA5, GOPC, GPHN, GPR124, GRIN2A, GSK3B, H3F3A, H3F3B, HERPUD1, HGF, HIP1, HMGA1, HMGA2, HNRNPA2B1, HOOK3, HSP90AA1, HSP90AB1, IDH1, IDH2, IGF1R, IKZF1, IL2, IL21R, IL6ST, IL7R, IRF4, ITK, JAK1, JAK2, JAK3, JAZF1, KDM5A, KEAP1, KIAA1549, KIF5B, KIT, KLHL6, KMT2A, KMT2C, KMT2D, KRAS, KTN1, LCK, LCP1, LGR5, LHFP, LIFR, LPP, LRIG3, LRP1B, LYL1, MAF, MALT1, MAML2, MAP2K2, MAP2K4, MAP3K1, MDM4, MDS2, MEF2B, MEN1, MITF, MLF1, MLH1, MLLT1, MLLT10, MLLT3, MLLT4, MLLT6, MNX1, MRE11A, MSH2, MSH6, MSI2, MTOR, MYB, MYCN, MYD88, MYH11, MYH9, NACA, NCKIPSD, NCOA1, NCOA2, NCOA4, NF1, NFE2L2, NFIB, NFKB2, NIN, NOTCH2, NPM1, NR4A3, NSD1, NT5C2, NTRK2, NTRK3, NUP214, NUP93, NUP98, NUTM1, PALB2, PAX3, PAX5, PAX7, PBRM1, PBX1, PCM1, PCSK7, PDCD1, PDCD1LG2, PDGFB, PDGFRA, PDGFRB, PDK1, PER1, PICALM, PIK3CA, PIK3R1, PIK3R2, PIM1, PML, PMS2, POLE, POT1, POU2AF1, PPARG, PRCC, PRDM1, PRDM16, PRKAR1A, PRRX1, PSIP1, PTCH1, PTEN, PTPN11, PTPRC, RABEP1, RAC1, RAD50, RAD51, RAD51B, RAF1, RALGDS, RANBP17, RAP1GDS1, RARA, RBM15, REL, RET, RMI2, RNF43, RPL20, RPL5, RPN1, RPTOR, RUNX1, RUNX1T1, SBDS, SDC4, SDHAF2, SDHB, SDHC, SDHD, 8-Sep, SET, SETBP1, SETD2, SF3B1, SH2B3, SH3GL1, SLC34A2, SMAD2, SMAD4, SMARCB1, SMARCE1, SMO, SNX29, SOX10, SPECC1, SPEN, SRGAP3, SRSF2, SRSF3, SS18, SS18L1, STAT3, STAT4, STAT5B, STIL, STK11, SUFU, SUZ12, SYK, TAF15, TCF12, TCF3, TCF7L2, TET1, TET2, TFEB, TFG, TFRC, TGFBR2, TLX1, TNFAIP3, TNFRSF14, TNFRSF17, TP53, TPM3, TPM4, TPR, TRAF7, TRIM26, TRIM27, TRIM33, TRIP11, TRRAP, TSC1, TSC2, TSHR, TTL, U2AF1, USP6, VEGFA, VEGFB, VTI1A, WHSC1, WHSC1L1, WIF1, WISP3, WRN, WWTR1, XPA, XPC, XPO1, YWHAE, ZMYM2, ZNF217, ZNF331, ZNF384, ZNF521, ZNF703
Gene fusions and mutations in cancer	AKT3, ALK, ARHGAP26, AXL, BRAF, BRD3/4, EGFR, ERG, ESR1, ETV1/4/5/6, EWSR1, FGFR1, FGFR2, FGFR3, FGR, INSR, MAML2, MAST1/2, MET, MSMB, MUSK, MYB, NOTCH1/2, NRG1, NTRK1/2/3, NUMBL, NUTM1, PDGFRA/B, PIK3CA, PKN1, PPARG, PRKCA/B, RAF1, RELA, RET, ROS1, RSPO2/3, TERT, TFE3, TFEB, THADA, TMPRSS2
Gene fusions and mutations in cancer	ABL1 fusion to (ETV6, NUP214, RCSD1, RANBP2, SNX2, or ZMIZ1); ABL2 fusion to (PAG1 or RCSD1); CSF1R fusion to (SSBP2); PDGFRB fusion to (EBF1, SSBP2, TNIP1 or ZEB2); CRLF2 fusion to (P2RY8); JAK2 fusion to (ATF7IP, BCR, ETV6, PAX5, PPFIBP1, SSBP2, STRN3, TERF2, or TPR); EPOR fusion to (IGH or IGK); IL2RB fusion to (MYH9); NTRK3 fusion to (ETV6); PTK2B fusion to (KDM6A or STAG2); TSLP fusion to (IQGAP2); TYK2 fusion to (MYB)
Cytohesions	cytohesin-1 (CYTH1), cytohesin-2 (CYTH2; ARNO), cytohesin-3 (CYTH3; Grp1; ARNO3), cytohesin-4 (CYTH4)
Cancer/Angio	Erb 2, Erb 3, Erb 4, UNC93a, B7H3, MUC1, MUC2, MUC16, MUC17, 5T4, RAGE, VEGF A, VEGFR2, FLT1, DLL4, Epcam
Tissue (Breast)	BIG H3, GCDFP-15, PR(B), GPR 30, CYFRA 21, BRCA 1, BRCA 2, ESR 1, ESR2
Tissue (Prostate)	PSMA, PCSA, PSCA, PSA, TMPRSS2
Inflammation/Immune	MFG-E8, IFNAR, CD40, CD80, MICB, HLA-DRb, IL-17-Ra
Common vesicle markers	HSPA8, CD63, Actb, GAPDH, CD9, CD81, ANXA2, HSP90AA1, ENO1, YWHAZ, PDCD6IP, CFL1, SDCBP, PKN2, MSN, MFGE8, EZR, YWHAG, PGK1, EEF1A1, PPIA, GLC1F, GK, ANXA6, ANXA1, ALDOA, ACTG1, TPI1, LAMP2, HSP90AB1, DPP4, YWHAB, TSG101, PFN1, LDHB, HSPA1B, HSPA1A, GSTP1, GNAI2, GDI2, CLTC, ANXA5, YWHAQ, TUBA1A, THBS1, PRDX1, LDHA, LAMP1, CLU, CD86
Common vesicle membrane markers	CD63, GAPDH, CD9, CD81, ANXA2, ENO1, SDCBP, MSN, MFGE8, EZR, GK, ANXA1, LAMP2, DPP4, TSG101, HSPA1A, GDI2, CLTC, LAMP1, CD86, ANPEP, TFRC, SLC3A2, RDX, RAP1B, RAB5C, RAB5B, MYH9, ICAM1, FN1, RAB11B, PIGR, LGALS3, ITGB1, EHD1, CLIC1, ATP1A1, ARF1, RAP1A, P4HB, MUC1, KRT10, HLA-A, FLOT1, CD59, C1orf58, BASP1, TACSTD1, STOM
Common vesicle markers	MHC class I, MHC class II, Integrins, Alpha 4 beta 1, Alpha M beta 2, Beta 2, ICAM1/CD54, P-selection, Dipeptidylpeptidase IV/CD26, Aminopeptidase n/CD13, CD151, CD53, CD37, CD82, CD81, CD9, CD63, Hsp70, Hsp84/90 Actin, Actin-binding proteins, Tubulin, Annexin I, Annexin II, Annexin IV, Annexin V, Annexin VI, RAB7/RAP1B/RADGDI, Gi2alpha/14-3-3, CBL/LCK, CD63, GAPDH, CD9, CD81, ANXA2, ENO1, SDCBP, MSN, MFGE8, EZR, GK, ANXA1, LAMP2, DPP4, TSG101, HSPA1A, GDI2, CLTC, LAMP1, Cd86, ANPEP, TFRC, SLC3A2, RDX, RAP1B,

	RAB5C, RAB5B, MYH9, ICAM1, FN1, RAB11B, PIGR, LGALS3, ITGB1, EHD1, CLIC1, ATP1A1, ARF1, RAP1A, P4HB, MUC1, KRT10, HLA-A, FLOT1, CD59, C1orf58, BASP1, TACSTD1, STOM
Vesicle markers	A33, a33 n15, AFP, ALA, ALIX, ALP, AnnexinV, APC, ASCA, ASPH (246-260), ASPH (666-680), ASPH (A-10), ASPH (D01P), ASPH (D03), ASPH (G-20), ASPH (H-300), AURKA, AURKB, B7H3, B7H4, BCA-225, BCNP, BDNF, BRCA, CA125 (MUC16), CA-19-9, C-Bir, CD1.1, CD10, CD174 (Lewis y), CD24, CD44, CD46, CD59 (MEM-43), CD63, CD66e CEA, CD73, CD81, CD9, CDA, CDAC1 1a2, CEA, C-Erb2, C-erbB2, CRMP-2, CRP, CXCL12, CYFRA21-1, DLL4, DR3, EGFR, Epcam, EphA2, EphA2 (H-77), ER, ErbB4, EZH2, FASL, FRT, FRT c.f23, GDF15, GPCR, GPR30, Gro-alpha, HAP, HBD 1, HBD2, HER 3 (ErbB3), HSP, HSP70, hVEGFR2, iC3b, IL 6 Unc, IL-1B, IL6 Unc, IL6R, IL8, IL-8, INSIG-2, KLK2, L1CAM, LAMN, LDH, MACC-1, MAPK4, MART-1, MCP-1, M-CSF, MFG-E8, MIC1, MIF, MIS RII, MMG, MMP26, MMP7, MMP9, MS4A1, MUC1, MUC1 seq1, MUC1 seq11A, MUC17, MUC2, Ncam, NGAL, NPGP/NPFF2, OPG, OPN, p53, p53, PA2G4, PBP, PCSA, PDGFRB, PGP9.5, PIM1, PR (B), PRL, PSA, PSMA, PSME3, PTEN, R5-CD9 Tube 1, Reg IV, RUNX2, SCRN1, seprase, SERPINB3, SPARC, SPB, SPDEF, SRVN, STAT 3, STEAP1, TF (FL-295), TFF3, TGM2, TIMP-1, TIMP1, TIMP2, TMEM211, TMPRSS2, TNF-alpha, Trail-R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, VEGF A, YPSMA-1
Vesicle markers	NSE, TRIM29, CD63, CD151, ASPH, LAMP2, TSPAN1, SNAIL, CD45, CKS1, NSE, FSHR, OPN, FTH1, PGP9, ANNEXIN 1, SPD, CD81, EPCAM, PTH1R, CEA, CYTO 7, CCL2, SPA, KRAS, TWIST1, AURKB, MMP9, P27, MMP1, HLA, HIF, CEACAM, CENPH, BTUB, INTG b4, EGFR, NACC1, CYTO 18, NAP2, CYTO 19, ANNEXIN V, TGM2, ERB2, BRCA1, B7H3, SFTPC, PNT, NCAM, MS4A1, P53, INGA3, MUC2, SPA, OPN, CD63, CD9, MUC1, UNCR3, PAN ADH, HCG, TIMP, PSMA, GPCR, RACK1, PSCA, VEGF, BMP2, CD81, CRP, PRO GRP, B7H3, MUC1, M2PK, CD9, PCSA, PSMA
Vesicle markers	TFF3, MS4A1, EphA2, GAL3, EGFR, N-gal, PCSA, CD63, MUC1, TGM2, CD81, DR3, MACC-1, TrKB, CD24, TIMP-1, A33, CD66 CEA, PRL, MMP9, MMP7, TMEM211, SCRN1, TROP2, TWEAK, CDACC1, UNC93A, APC, C-Erb, CD10, BDNF, FRT, GPR30, P53, SPR, OPN, MUC2, GRO-1, tsg 101, GDF15
Vesicle markers	CD9, Erb2, Erb4, CD81, Erb3, MUC16, CD63, DLL4, HLA-Drpe, B7H3, IFNAR, 5T4, PCSA, MICB, PSMA, MFG-E8, Muc1, PSA, Muc2, Unc93a, VEGFR2, EpCAM, VEGF A, TMPRSS2, RAGE, PSCA, CD40, Muc17, IL-17-RA, CD80
Benign Prostate Hyperplasia (BPH)	BCMA, CEACAM-1, HVEM, IL-1 R4, IL-10 Rb, Trappin-2, p53, hsa-miR-329, hsa-miR-30a, hsa-miR-335, hsa-miR-152, hsa-miR-151-5p, hsa-miR-200a, hsa-miR-145, hsa-miR-29a, hsa-miR-106b, hsa-miR-595, hsa-miR-142-5p, hsa-miR-99a, hsa-miR-20b, hsa-miR-373, hsa-miR-502-5p, hsa-miR-29b, hsa-miR-142-3p, hsa-miR-663, hsa-miR-423-5p, hsa-miR-15a, hsa-miR-888, hsa-miR-361-3p, hsa-miR-365, hsa-miR-10b, hsa-miR-199a-3p, hsa-miR-181a, hsa-miR-19a, hsa-miR-125b, hsa-miR-760, hsa-miR-7a, hsa-miR-671-5p, hsa-miR-7c, hsa-miR-1979, hsa-miR-103
Metastatic Prostate Cancer	hsa-miR-100, hsa-miR-1236, hsa-miR-1296, hsa-miR-141, hsa-miR-146b-5p, hsa-miR-17*, hsa-miR-181a, hsa-miR-200b, hsa-miR-20a*, hsa-miR-23a*, hsa-miR-331-3p, hsa-miR-375, hsa-miR-452, hsa-miR-572, hsa-miR-574-3p, hsa-miR-577, hsa-miR-582-3p, hsa-miR-937, miR-10a, miR-134, miR-141, miR-200b, miR-30a, miR-32, miR-375, miR-495, miR-564, miR-570, miR-574-3p, miR-885-3p
Metastatic Prostate Cancer	hsa-miR-200b, hsa-miR-375, hsa-miR-141, hsa-miR-331-3p, hsa-miR-181a, hsa-miR-574-3p
Prostate Cancer	hsa-miR-574-3p, hsa-miR-141, hsa-miR-432, hsa-miR-326, hsa-miR-2110, hsa-miR-181a-2*, hsa-miR-107, hsa-miR-301a, hsa-miR-484, hsa-miR-625*
Metastatic Prostate Cancer	hsa-miR-582-3p, hsa-miR-20a*, hsa-miR-375, hsa-miR-200b, hsa-miR-379, hsa-miR-572, hsa-miR-513a-5p, hsa-miR-577, hsa-miR-23a*, hsa-miR-1236, hsa-miR-609, hsa-miR-17*, hsa-miR-130b, hsa-miR-619, hsa-miR-624*, hsa-miR-198
Metastatic Prostate Cancer	FOX01A, SOX9, CLNS1A, PTGDS, XPO1, LETMD1, RAD23B, ABCC3, APC, CHES1, EDNRA, FRZB, HSPG2, TMPRSS2_ETV1 fusion
Prostate Cancer	hsa-let-7b, hsa-miR-107, hsa-miR-1205, hsa-miR-1270, hsa-miR-130b, hsa-miR-141, hsa-miR-143, hsa-miR-148b*, hsa-miR-150, hsa-miR-154*, hsa-miR-181a*, hsa-miR-181a-2*, hsa-miR-18a*, hsa-miR-19b-1*, hsa-miR-204, hsa-miR-2110, hsa-miR-215, hsa-miR-217, hsa-miR-219-2-3p, hsa-miR-23b*, hsa-miR-299-5p, hsa-miR-301a, hsa-miR-301a, hsa-miR-326, hsa-miR-331-3p, hsa-miR-365*, hsa-miR-373*, hsa-miR-424, hsa-miR-424*, hsa-miR-432, hsa-miR-450a, hsa-miR-451, hsa-miR-484, hsa-miR-497, hsa-miR-517*, hsa-miR-517a, hsa-miR-518f, hsa-miR-574-3p, hsa-miR-595, hsa-miR-617, hsa-miR-625*, hsa-miR-628-5p,

	hsa-miR-629, hsa-miR-634, hsa-miR-769-5p, hsa-miR-93, hsa-miR-96
Prostate Cancer	CD9, PSMA, PCSA, CD63, CD81, B7H3, IL 6, OPG-13, IL6R, PA2G4, EZH2, RUNX2, SERPINB3, EpCam
Prostate Cancer	A33, a33 n15, AFP, ALA, ALIX, ALP, AnnexinV, APC, ASCA, ASPH (246-260), ASPH (666-680), ASPH (A-10), ASPH (D01P), ASPH (D03), ASPH (G-20), ASPH (H-300), AURKA, AURKB, B7H3, B7H4, BCA-225, BCNP, BDNF, BRCA, CA125 (MUC16), CA-19-9, C-Bir, CD1.1, CD10, CD174 (Lewis y), CD24, CD44, CD46, CD59 (MEM-43), CD63, CD66e CEA, CD73, CD81, CD9, CDA, CDAC1 1a2, CEA, C-Erb2, C-erbB2, CRMP-2, CRP, CXCL12, CYFRA21-1, DLL4, DR3, EGFR, Epcam, EphA2, EphA2 (H-77), ER, ErbB4, EZH2, FASL, FRT, FRT c.f23, GDF15, GPCR, GPR30, Gro-alpha, HAP, HBD 1, HBD2, HER 3 (ErbB3), HSP, HSP70, hVEGFR2, iC3b, IL 6 Unc, IL-1B, IL6 Unc, IL6R, IL8, IL-8, INSIG-2, KLK2, L1CAM, LAMN, LDH, MACC-1, MAPK4, MART-1, MCP-1, M-CSF, MFG-E8, MIC1, MIF, MIS RII, MMG, MMP26, MMP7, MMP9, MS4A1, MUC1, MUC1 seq1, MUC1 seq11A, MUC17, MUC2, Ncam, NGAL, NPGP/NPFF2, OPG, OPN, p53, p53, PA2G4, PBP, PCSA, PDGFRB, PGP9.5, PIM1, PR (B), PRL, PSA, PSMA, PSME3, PTEN, R5-CD9 Tube 1, Reg IV, RUNX2, SCRN1, seprase, SERPINB3, SPARC, SPB, SPDEF, SRVN, STAT 3, STEAP1, TF (FL-295), TFF3, TGM2, TIMP-1, TIMP1, TIMP2, TMEM211, TMPRSS2, TNF-alpha, Trail-R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, VEGF A, YPSMA-1
Prostate Cancer Vesicle Markers	5T4, ACTG1, ADAM10, ADAM15, ALDOA, ANXA2, ANXA6, APOA1, ATP1A1, BASP1, C1orf58, C20orf114, C8B, CAPZA1, CAV1, CD151, CD2AP, CD59, CD9, CD9, CFL1, CFP, CHMP4B, CLTC, COTL1, CTNND1, CTSB, CTSZ, CYCS, DPP4, EEF1A1, EHD1, ENO1, F11R, F2, F5, FAM125A, FNBP1L, FOLH1, GAPDH, GLB1, GPX3, HIST1H1C, HIST1H2AB, HSP90AB1, HSPA1B, HSPA8, IGSF8, ITGB1, ITIH3, JUP, LDHA, LDHB, LUM, LYZ, MFGE8, MGAM, MMP9, MYH2, MYL6B, NME1, NME2, PABPC1, PABPC4, PACSIN2, PCBP2, PDCD6IP, PRDX2, PSA, PSMA, PSMA1, PSMA2, PSMA4, PSMA6, PSMA7, PSMB1, PSMB2, PSMB3, PSMB4, PSMB5, PSMB6, PSMB8, PTGFRN, RPS27A, SDCBP, SERINC5, SH3GL1, SLC3A2, SMPDL3B, SNX9, TACSTD1, TCN2, THBS1, TPI1, TSG101, TUBB, VDAC2, VPS37B, YWHAG, YWHAQ, YWHAZ
Prostate Cancer Vesicle Markers	FLNA, DCRN, HER 3 (ErbB3), VCAN, CD9, GAL3, CDADC1, GM-CSF, EGFR, RANK, CSA, PSMA, ChickenIgY, B7H3, PCSA, CD63, CD3, MUC1, TGM2, CD81, S100-A4, MFG-E8, Integrin, NK-2R(C-21), PSA, CD24, TIMP-1, IL6 Unc, PBP, PIM1, CA-19-9, Trail-R4, MMP9, PRL, EphA2, TWEAK, NY-ESO-1, Mammaglobin, UNC93A, A33, AURKB, CD41, XAGE-1, SPDEF, AMACR, seprase/FAP, NGAL, CXCL12, FRT, CD66e CEA, SIM2 (C-15), C-Bir, STEAP, PSIP1/LEDGF, MUC17, hVEGFR2, ERG, MUC2, ADAM10, ASPH (A-10), CA125, Gro-alpha, Tsg 101, SSX2, Trail-R4
Prostate Cancer Vesicle Markers	NT5E (CD73), A33, ABL2, ADAM10, AFP, ALA, ALIX, ALPL, AMACR, Apo J/CLU, ASCA, ASPH (A-10), ASPH (D01P), AURKB, B7H3, B7H4, BCNP, BDNF, CA125 (MUC16), CA-19-9, C-Bir (Flagellin), CD10, CD151, CD24, CD3, CD41, CD44, CD46, CD59(MEM-43), CD63, CD66e CEA, CD81, CD9, CDA, CDADC1, C-erbB2, CRMP-2, CRP, CSA, CXCL12, CXCR3, CYFRA21-1, DCRN, DDX-1, DLL4, EGFR, EpCAM, EphA2, ERG, EZH2, FASL, FLNA, FRT, GAL3, GATA2, GM-CSF, Gro-alpha, HAP, HER3 (ErbB3), HSP70, HSPB1, hVEGFR2, iC3b, IL-1B, IL6 R, IL6 Unc, IL7 R alpha/CD127, IL8, INSIG-2, Integrin, KLK2, Label, LAMN, Mammaglobin, M-CSF, MFG-E8, MIF, MIS RII, MMP7, MMP9, MS4A1, MUC1, MUC17, MUC2, Ncam, NDUFB7, NGAL, NK-2R(C-21), NY-ESO-1, p53, PBP, PCSA, PDGFRB, PIM1, PRL, PSA, PSIP1/LEDGF, PSMA, RAGE, RANK, Reg IV, RUNX2, S100-A4, seprase/FAP, SERPINB3, SIM2 (C-15), SPARC, SPC, SPDEF, SPP1, SSX2, SSX4, STEAP, STEAP4, TFF3, TGM2, TIMP-1, TMEM211, Trail-R2, Trail-R4, TrKB (poly), Trop2, Tsg 101, TWEAK, UNC93A, VCAN, VEGF A, wnt-5a(C-16), XAGE, XAGE-1
Prostate Vesicle Membrane	ADAM 9, ADAM10, AGR2, ALDOA, ALIX, ANXA1, ANXA2, ANXA4, ARF6, ATP1A3, B7H3, BCHE, BCL2L14 (Bcl G), BCNP1, BDKRB2, BDNFCAV1-Caveolin1, CCR2 (CC chemokine receptor 2, CD192), CCR5 (CC chemokine receptor 5), CCT2 (TCP1-beta), CD10, CD151, CD166/ALCAM, CD24, CD283/TLR3, CD41, CD46, CD49d (Integrin alpha 4, ITGA4), CD63, CD81, CD9, CD90/THY1, CDH1, CDH2, CDKN1A cyclin-dependent kinase inhibitor (p21), CGA gene (coding for the alpha subunit of glycoprotein hormones), CLDN3- Claudin3, COX2 (PTGS2), CSE1L (Cellular Apoptosis Susceptibility), CXCR3, Cytokeratin 18, Eag1 (KCNH1), EDIL3 (del-1), EDNRB - Endothelial Receptor Type B, EGFR, EpoR, EZH2 (enhancer of Zeste Homolog2), EZR, FABP5, Farnesyltransferase/geranylgeranyl diphosphate synthase 1 (GGPS1), Fatty acid synthase (FASN), FTL (light and heavy), GAL3, GDF15-Growth Differentiation Factor 15, GloI, GM-

	CSF, GSTP1, H3F3A, HGF (hepatocyte growth factor), hK2 / Kif2a, HSP90AA1, HSPA1A / HSP70-1, HSPB1, IGFBP-2, IGFBP-3, IL1alpha, IL-6, IQGAP1, ITGAL (Integrin alpha L chain), Ki67, KLK1, KLK10, KLK11, KLK12, KLK13, KLK14, KLK15, KLK4, KLK5, KLK6, KLK7, KLK8, KLK9, Lamp-2, LDH-A, LGALS3BP, LGALS8, MMP 1, MMP 2, MMP 25, MMP 3, MMP10, MMP-14/MT1-MMP, MMP7, MTA1nAnS, Nav1.7, NKX3-1, Notch1, NRP1 / CD304, PAP (ACPP), PGP, PhIP, PIP3 / BPNT1, PKM2, PKP1 (plakophilin1), PKP3 (plakophilin3), Plasma chromogranin-A (CgA), PRDX2, Prostate secretory protein (PSP94) / β-Microseminoprotein (MSP) / IGBF, PSAP, PSMA, PSMA1, PTENPTPN13/PTPL1, RPL19, seprase/FAPSET, SLC3A2 / CD98, SRVN, STEAP1, Syndecan / CD138, TGFB, TGM2, TIMP-1TLR4 (CD284), TLR9 (CD289), TMPRSS1 / hepsin, TMPRSS2, TNFR1, TNFα, Transferrin receptor/CD71/TRFR, Trop2 (TACSTD2), TWEAK uPA (urokinase plasminoge activator) degrades extracellular matrix, uPAR (uPA receptor) / CD87, VEGFR1, VEGFR2
Prostate Vesicle Markers	ADAM 34, ADAM 9, AGR2, ALDOA, ANXA1, ANXA 11, ANXA4, ANXA 7, ANXA2, ARF6, ATP1A1, ATP1A2, ATP1A3, BCHE, BCL2L14 (Bcl G), BDKRB2, CA215, CAV1-Caveolin1, CCR2 (CC chemokine receptor 2, CD192), CCR5 (CC chemokine receptor 5), CCT2 (TCP1-beta), CD166/ALCAM, CD49b (Integrin alpha 2, ITGA4), CD90/THY1, CDH1, CDH2, CDKN1A cyclin-dependent kinase inhibitor (p21), CGA gene (coding for the alpha subunit of glycoprotein hormones), CHMP4B, CLDN3- Claudin3, CLSTN1 (Calsyntenin-1), COX2 (PTGS2), CSE1L (Cellular Apoptosis Susceptibility), Cytokeratin 18, Eag1 (KCNH1) (plasma membrane-K+-voltage gated channel), EDIL3 (del-1), EDNRB-Endothelial Receptor Type B, Endoglin/CD105, ENOX2 – Ecto-NOX disulphide Thiol exchanger 2, EPCA-2 Early prostate cancer antigen2, EpoR, EZH2 (enhancer of Zeste Homolog2), EZR, FABP5, Farnesyltransferase/geranylgeranyl diphosphate synthase 1 (GGPS1), Fatty acid synthase (FASN, plasma membrane protein), FTL (light and heavy), GDF15-Growth Differentiation Factor 15, GloI, GSTP1, H3F3A, HGF (hepatocyte growth factor), hK2 (KLK2), HSP90AA1, HSPA1A / HSP70-1, IGFBP-2, IGFBP-3, IL1alpha, IL-6, IQGAP1, ITGAL (Integrin alpha L chain), Ki67, KLK1, KLK10, KLK11, KLK12, KLK13, KLK14, KLK15, KLK4, KLK5, KLK6, KLK7, KLK8, KLK9, Lamp-2, LDH-A, LGALS3BP, LGALS8, MFAP5, MMP 1, MMP 2, MMP 24, MMP 25, MMP 3, MMP10, MMP-14/MT1-MMP, MTA1, nAnS, Nav1.7, NCAM2 – Neural cell Adhesion molecule 2, NGEP/D-TMPP/IPCA-5/ANO7, NKX3-1, Notch1, NRP1 / CD304, PGP, PAP (ACPP), PCA3- Prostate cancer antigen 3, Pdia3/ERp57, PhIP, phosphatidylethanolamine (PE), PIP3, PKP1 (plakophilin1), PKP3 (plakophilin3), Plasma chromogranin-A (CgA), PRDX2, Prostate secretory protein (PSP94) / β-Microseminoprotein (MSP) / IGBF, PSAP, PSMA1, PTEN, PTGFRN, PTPN13/PTPL1, PKM2, RPL19, SCA-1 / ATXN1, SERINC5/TPO1, SET, SLC3A2 / CD98, STEAP1, STEAP-3, SRVN, Syndecan / CD138, TGFB, Tissue Polypeptide Specific antigen TPS, TLR4 (CD284), TLR9 (CD289), TMPRSS1 / hepsin, TMPRSS2, TNFR1, TNFα, CD283/TLR3, Transferrin receptor/CD71/TRFR, uPA (urokinase plasminoge activator), uPAR (uPA receptor) / CD87, VEGFR1, VEGFR2
Prostate Cancer Treatment	hsa-miR-1974, hsa-miR-27b, hsa-miR-103, hsa-miR-146a, hsa-miR-22, hsa-miR-382, hsa-miR-23a, hsa-miR-376c, hsa-miR-335, hsa-miR-142-5p, hsa-miR-221, hsa-miR-142-3p, hsa-miR-151-3p, hsa-miR-21, hsa-miR-16
Prostate Cancer	let-7d, miR-148a, miR-195, miR-25, miR-26b, miR-329, miR-376c, miR-574-3p, miR-888, miR-9, miR1204, miR-16-2*, miR-497, miR-588, miR-614, miR-765, miR92b*, miR-938, let-7f-2*, miR-300, miR-523, miR-525-5p, miR-1182, miR-1244, miR-520d-3p, miR-379, let-7b, miR-125a-3p, miR-1296, miR-134, miR-149, miR-150, miR-187, miR-32, miR-324-3p, miR-324-5p, miR-342-3p, miR-378, miR-378*, miR-384, miR-451, miR-455-3p, miR-485-3p, miR-487a, miR-490-3p, miR-502-5p, miR-548a-5p, miR-550, miR-562, miR-593, miR-593*, miR-595, miR-602, miR-603, miR-654-5p, miR-877*, miR-886-5p, miR-125a-5p, miR-140-3p, miR-192, miR-196a, miR-2110, miR-212, miR-222, miR-224*, miR-30b*, miR-499-3p, miR-505*
Prostate (PCSA+ cMVs)	miR-182, miR-663, miR-155, mirR-125a-5p, miR-548a-5p, miR-628-5p, miR-517*, miR-450a, miR-920, hsa-miR-619, miR-1913, miR-224*, miR-502-5p, miR-888, miR-376a, miR-542-5p, miR-30b*, miR-1179
Prostate Cancer	miR-183-96-182 cluster (miRs-183, 96 and 182), metal ion transporter such as hZIP1, SLC39A1, SLC39A2, SLC39A3, SLC39A4, SLC39A5, SLC39A6, SLC39A7, SLC39A8, SLC39A9, SLC39A10, SLC39A11, SLC39A12, SLC39A13, SLC39A14
Prostate Cancer	RAD23B, FBP1, TNFRSF1A, CCNG2, NOTCH3, ETV1, BID, SIM2, LETMD1, ANXA1, miR-519d, miR-647
Prostate Cancer	RAD23B, FBP1, TNFRSF1A, NOTCH3, ETV1, BID, SIM2, ANXA1, BCL2

Prostate Cancer	ANPEP, ABL1, PSCA, EFNA1, HSPB1, INMT, TRIP13
Prostate Cancer	E2F3, c-met, pRB, EZH2, e-cad, CAXII, CAIX, HIF-1α, Jagged, PIM-1, hepsin, RECK, Clusterin, MMP9, MTSP-1, MMP24, MMP15, IGFBP-2, IGFBP-3, E2F4, caveolin, EF-1A, Kallikrein 2, Kallikrein 3, PSGR
Prostate Cancer	A2ML1, BAX, C10orf47, C1orf162, CSDA, EIFC3, ETFB, GABARAPL2, GUK1, GZMH, HIST1H3B, HLA-A, HSP90AA1, NRGN, PRDX5, PTMA, RABAC1, RABAGAP1L, RPL22, SAP18, SEPW1, SOX1
Prostate Cancer	NY-ESO-1, SSX-2, SSX-4, XAGE-lb, AMACR, p90 autoantigen, LEDGF
Prostate Cancer	A33, ABL2, ADAM10, AFP, ALA, ALIX, ALPL, ApoJ/CLU, ASCA, ASPH(A-10), ASPH(D01P), AURKB, B7H3, B7H3, B7H4, BCNP, BDNF, CA125(MUC16), CA-19-9, C-Bir, CD10, CD151, CD24, CD41, CD44, CD46, CD59(MEM-43), CD63, CD63, CD66eCEA, CD81, CD81, CD9, CD9, CDA, CDADC1, CRMP-2, CRP, CXCL12, CXCR3, CYFRA21-1, DDX-1, DLL4, DLL4, EGFR, Epcam, EphA2, ErbB2, ERG, EZH2, FASL, FLNA, FRT, GAL3, GATA2, GM-CSF, Gro-alpha, HAP, HER3(ErbB3), HSP70, HSPB1, hVEGFR2, iC3b, IL-1B, IL6R, IL6Unc, IL7Ralpha/CD127, IL8, INSIG-2, Integrin, KLK2, LAMN, Mammoglobin, M-CSF, MFG-E8, MIF, MISRII, MMP7, MMP9, MUC1, Muc1, MUC17, MUC2, Ncam, NDUFB7, NGAL, NK-2R(C-21), NT5E (CD73), p53, PBP, PCSA, PCSA, PDGFRB, PIM1, PRL, PSA, PSA, PSMA, PSMA, RAGE, RANK, RegIV, RUNX2, S100-A4, seprase/FAP, SERPINB3, SIM2(C-15), SPARC, SPC, SPDEF, SPP1, STEAP, STEAP4, TFF3, TGM2, TIMP-1, TMEM211, Trail-R2, Trail-R4, TrKB(poly), Trop2, Tsg101, TWEAK, UNC93A, VEGFA, wnt-5a(C-16)
Prostate Vesicles	CD9, CD63, CD81, PCSA, MUC2, MFG-E8
Prostate Cancer	miR-148a, miR-329, miR-9, miR-378*, miR-25, miR-614, miR-518c*, miR-378, miR-765, let-7f-2*, miR-574-3p, miR-497, miR-32, miR-379, miR-520g, miR-542-5p, miR-342-3p, miR-1206, miR-663, miR-222
Prostate Cancer	hsa-miR-877*, hsa-miR-593, hsa-miR-595, hsa-miR-300, hsa-miR-324-5p, hsa-miR-548a-5p, hsa-miR-329, hsa-miR-550, hsa-miR-886-5p, hsa-miR-603, hsa-miR-490-3p, hsa-miR-938, hsa-miR-149, hsa-miR-150, hsa-miR-1296, hsa-miR-384, hsa-miR-487a, hsa-miRPlus-C1089, hsa-miR-485-3p, hsa-miR-525-5p
Prostate Cancer	hsa-miR-451, hsa-miR-223, hsa-miR-593*, hsa-miR-1974, hsa-miR-486-5p, hsa-miR-19b, hsa-miR-320b, hsa-miR-92a, hsa-miR-21, hsa-miR-675*, hsa-miR-16, hsa-miR-876-5p, hsa-miR-144, hsa-miR-126, hsa-miR-137, hsa-miR-1913, hsa-miR-29b-1*, hsa-miR-15a, hsa-miR-93, hsa-miR-1266
Inflammatory Disease	miR-588, miR-1258, miR-16-2*, miR-938, miR-526b, miR-92b*, let-7d, miR-378*, miR-124, miR-376c, miR-26b, miR-1204, miR-574-3p, miR-195, miR-499-3p, miR-2110, miR-888
Prostate Cancer	A33, ADAM10, AMACR, ASPH (A-10), AURKB, B7H3, CA125, CA-19-9, C-Bir, CD24, CD3, CD41, CD63, CD66e CEA, CD81, CD9, CDADC1, CSA, CXCL12, DCRN, EGFR, EphA2, ERG, FLNA, FRT, GAL3, GM-CSF, Gro-alpha, HER 3 (ErbB3), hVEGFR2, IL6 Unc, Integrin, Mammaglobin, MFG-E8, MMP9, MUC1, MUC17, MUC2, NGAL, NK-2R(C-21), NY-ESO-1, PBP, PCSA, PIM1, PRL, PSA, PSIP1/LEDGF, PSMA, RANK, S100-A4, seprase/FAP, SIM2 (C-15), SPDEF, SSX2, STEAP, TGM2, TIMP-1, Trail-R4, Tsg 101, TWEAK, UNC93A, VCAN, XAGE-1
Prostate Cancer	A33, ADAM10, ALIX, AMACR, ASCA, ASPH (A-10), AURKB, B7H3, BCNP, CA125, CA-19-9, C-Bir (Flagellin), CD24, CD3, CD41, CD63, CD66e CEA, CD81, CD9, CDADC1, CRP, CSA, CXCL12, CYFRA21-1, DCRN, EGFR, EpCAM, EphA2, ERG, FLNA, GAL3, GATA2, GM-CSF, Gro alpha, HER3 (ErbB3), HSP70, hVEGFR2, iC3b, IL-1B, IL6 Unc, IL8, Integrin, KLK2, Mammaglobin, MFG-E8, MMP7, MMP9, MS4A1, MUC1, MUC17, MUC2, NGAL, NK-2R(C-21), NY-ESO-1, p53, PBP, PCSA, PIM1, PRL, PSA, PSMA, RANK, RUNX2, S100-A4, seprase/FAP, SERPINB3, SIM2 (C-15), SPC, SPDEF, SSX2, SSX4, STEAP, TGM2, TIMP-1, TRAIL R2, Trail-R4, Tsg 101, TWEAK, VCAN, VEGF A, XAGE
Prostate Vesicles	EpCam, CD81, PCSA, MUC2, MFG-E8
Prostate Vesicles	CD9, CD63, CD81, MMP7, EpCAM
Prostate Cancer	let-7d, miR-148a, miR-195, miR-25, miR-26b, miR-329, miR-376c, miR-574-3p, miR-888, miR-9, miR1204, miR-16-2*, miR-497, miR-588, miR-614, miR-765, miR92b*, miR-938, let-7f-2*, miR-300, miR-523, miR-525-5p, miR-1182, miR-1244, miR-520d-3p, miR-379, let-7b, miR-125a-3p, miR-1296, miR-134, miR-149, miR-150, miR-187, miR-32, miR-324-3p, miR-324-5p, miR-342-3p, miR-378, miR-378*, miR-384, miR-451, miR-455-3p, miR-485-3p, miR-487a, miR-490-3p, miR-502-5p, miR-548a-5p, miR-550, miR-562, miR-593, miR-593*, miR-595, miR-602, miR-603, miR-654-5p, miR-877*, miR-886-5p, miR-125a-5p,

	miR-140-3p, miR-192, miR-196a, miR-2110, miR-212, miR-222, miR-224*, miR-30b*, miR-499-3p, miR-505*
Prostate Cancer	STAT3, EZH2, p53, MACC1, SPDEF, RUNX2, YB-1, AURKA, AURKB
Prostate Cancer (Ensembl ENSG identifiers)	E.001036, E.001497, E.001561, E.002330, E.003402, E.003756, E.004838, E.005471, E.005882, E.005893, E.006210, E.006453, E.006625, E.006695, E.006756, E.007264, E.007952, E.008118, E.008196, E.009694, E.009830, E.010244, E.010256, E.010278, E.010539, E.010810, E.011052, E.011114, E.011143, E.011304, E.011451, E.012061, E.012779, E.014216, E.014257, E.015133, E.015171, E.015479, E.015676, E.016402, E.018189, E.018699, E.020922, E.022976, E.023909, E.026508, E.026559, E.029363, E.029725, E.030582, E.033030, E.035141, E.036257, E.036448, E.038002, E.039068, E.039560, E.041353, E.044115, E.047410, E.047597, E.048544, E.048828, E.049239, E.049246, E.049883, E.051596, E.051620, E.052795, E.053108, E.054118, E.054938, E.056097, E.057252, E.057608, E.058729, E.059122, E.059378, E.059691, E.060339, E.060688, E.061794, E.061918, E.062485, E.063241, E.063244, E.064201, E.064489, E.064655, E.064886, E.065054, E.065057, E.065308, E.065427, E.065457, E.065485, E.065526, E.065548, E.065978, E.066455, E.066557, E.067248, E.067369, E.067704, E.068724, E.068885, E.069535, E.069712, E.069849, E.069869, E.069956, E.070501, E.070785, E.070814, E.071246, E.071626, E.071859, E.072042, E.072071, E.072110, E.072506, E.073050, E.073350, E.073584, E.073756, E.074047, E.074071, E.074964, E.075131, E.075239, E.075624, E.075651, E.075711, E.075856, E.075886, E.076043, E.076248, E.076554, E.076864, E.077097, E.077147, E.077312, E.077514, E.077522, E.078269, E.078295, E.078808, E.078902, E.079246, E.079313, E.079785, E.080572, E.080823, E.081087, E.081138, E.081181, E.081721, E.081842, E.082212, E.082258, E.082556, E.083093, E.083720, E.084234, E.084463, E.085224, E.085733, E.086062, E.086205, E.086717, E.087087, E.087301, E.088888, E.088899, E.088930, E.088992, E.089048, E.089127, E.089154, E.089177, E.089248, E.089280, E.089902, E.090013, E.090060, E.090565, E.090612, E.090615, E.090674, E.090861, E.090889, E.091140, E.091483, E.091542, E.091732, E.092020, E.092199, E.092421, E.092621, E.092820, E.092871, E.092978, E.093010, E.094755, E.095139, E.095380, E.095485, E.095627, E.096060, E.096384, E.099331, E.099715, E.099783, E.099785, E.099800, E.099821, E.099899, E.099917, E.099956, E.100023, E.100056, E.100065, E.100084, E.100142, E.100191, E.100216, E.100242, E.100271, E.100284, E.100299, E.100311, E.100348, E.100359, E.100393, E.100399, E.100401, E.100412, E.100442, E.100575, E.100577, E.100583, E.100601, E.100603, E.100612, E.100632, E.100714, E.100739, E.100796, E.100802, E.100815, E.100823, E.100836, E.100883, E.101057, E.101126, E.101152, E.101222, E.101246, E.101265, E.101365, E.101439, E.101557, E.101639, E.101654, E.101811, E.101812, E.101901, E.102030, E.102054, E.102103, E.102158, E.102174, E.102241, E.102290, E.102316, E.102362, E.102384, E.102710, E.102780, E.102904, E.103035, E.103067, E.103175, E.103194, E.103449, E.103479, E.103591, E.103599, E.103855, E.103978, E.104064, E.104067, E.104131, E.104164, E.104177, E.104228, E.104331, E.104365, E.104419, E.104442, E.104611, E.104626, E.104723, E.104760, E.104805, E.104812, E.104823, E.104824, E.105127, E.105220, E.105221, E.105281, E.105379, E.105402, E.105404, E.105409, E.105419, E.105428, E.105486, E.105514, E.105518, E.105618, E.105705, E.105723, E.105939, E.105948, E.106049, E.106078, E.106128, E.106153, E.106346, E.106392, E.106554, E.106565, E.106603, E.106633, E.107104, E.107164, E.107404, E.107485, E.107551, E.107581, E.107623, E.107798, E.107816, E.107833, E.107890, E.107897, E.107968, E.108296, E.108312, E.108375, E.108387, E.108405, E.108417, E.108465, E.108561, E.108582, E.108639, E.108641, E.108848, E.108883, E.108953, E.109062, E.109184, E.109572, E.109625, E.109758, E.109790, E.109814, E.109846, E.109956, E.110063, E.110066, E.110104, E.110107, E.110321, E.110328, E.110921, E.110955, E.111057, E.111218, E.111261, E.111335, E.111540, E.111605, E.111647, E.111785, E.111790, E.111801, E.111907, E.112039, E.112081, E.112096, E.112110, E.112144, E.112232, E.112234, E.112473, E.112578, E.112584, E.112715, E.112941, E.113013, E.113163, E.113282, E.113368, E.113441, E.113448, E.113522, E.113580, E.113645, E.113719, E.113739, E.113790, E.114054, E.114127, E.114302, E.114331, E.114388, E.114491, E.114861, E.114867, E.115053, E.115221, E.115234, E.115239, E.115241, E.115257, E.115339, E.115540, E.115541, E.115561, E.115604, E.115648, E.115738, E.115758, E.116044, E.116096, E.116127, E.116254, E.116288, E.116455, E.116478, E.116604, E.116649, E.116726, E.116754, E.116833, E.117298, E.117308, E.117335, E.117362, E.117411, E.117425, E.117448, E.117480, E.117592, E.117593, E.117614, E.117676, E.117713, E.117748, E.117751,

	E.117877, E.118181, E.118197, E.118260, E.118292, E.118513, E.118523, E.118640, E.118898, E.119121, E.119138, E.119318, E.119321, E.119335, E.119383, E.119421, E.119636, E.119681, E.119711, E.119820, E.119888, E.119906, E.120159, E.120328, E.120337, E.120370, E.120656, E.120733, E.120837, E.120868, E.120915, E.120948, E.121022, E.121057, E.121068, E.121104, E.121390, E.121671, E.121690, E.121749, E.121774, E.121879, E.121892, E.121903, E.121940, E.121957, E.122025, E.122033, E.122126, E.122507, E.122566, E.122705, E.122733, E.122870, E.122884, E.122952, E.123066, E.123080, E.123143, E.123154, E.123178, E.123416, E.123427, E.123595, E.123901, E.123908, E.123983, E.123992, E.124143, E.124164, E.124181, E.124193, E.124216, E.124232, E.124529, E.124562, E.124570, E.124693, E.124749, E.124767, E.124788, E.124795, E.124831, E.124942, E.125246, E.125257, E.125304, E.125352, E.125375, E.125445, E.125492, E.125676, E.125753, E.125798, E.125844, E.125868, E.125901, E.125944, E.125995, E.126062, E.126267, E.126653, E.126773, E.126777, E.126814, E.126858, E.126883, E.126934, E.126945, E.126952, E.127022, E.127328, E.127329, E.127399, E.127415, E.127554, E.127616, E.127720, E.127824, E.127884, E.127914, E.127946, E.127948, E.128050, E.128311, E.128342, E.128609, E.128626, E.128683, E.128708, E.128881, E.129315, E.129351, E.129355, E.129514, E.129636, E.129657, E.129757, E.129810, E.129990, E.130175, E.130177, E.130193, E.130255, E.130299, E.130305, E.130338, E.130340, E.130402, E.130413, E.130612, E.130713, E.130764, E.130770, E.130810, E.130826, E.130935, E.131351, E.131467, E.131473, E.131771, E.131773, E.132002, E.132275, E.132323, E.132382, E.132475, E.132481, E.132589, E.132646, E.132716, E.132881, E.133313, E.133315, E.133687, E.133835, E.133863, E.133874, E.133961, E.134077, E.134138, E.134207, E.134248, E.134308, E.134444, E.134452, E.134548, E.134684, E.134759, E.134809, E.134851, E.134955, E.135052, E.135297, E.135298, E.135387, E.135390, E.135476, E.135486, E.135525, E.135597, E.135679, E.135740, E.135829, E.135842, E.135870, E.135900, E.135914, E.135926, E.135940, E.135999, E.136044, E.136068, E.136152, E.136169, E.136280, E.136371, E.136383, E.136450, E.136521, E.136527, E.136574, E.136710, E.136750, E.136807, E.136874, E.136875, E.136930, E.136933, E.136935, E.137055, E.137124, E.137312, E.137409, E.137497, E.137513, E.137558, E.137601, E.137727, E.137776, E.137806, E.137814, E.137815, E.137948, E.137955, E.138028, E.138031, E.138041, E.138050, E.138061, E.138069, E.138073, E.138095, E.138160, E.138294, E.138347, E.138363, E.138385, E.138587, E.138594, E.138621, E.138674, E.138756, E.138757, E.138760, E.138772, E.138796, E.139211, E.139405, E.139428, E.139517, E.139613, E.139626, E.139684, E.139697, E.139874, E.140263, E.140265, E.140326, E.140350, E.140374, E.140382, E.140451, E.140481, E.140497, E.140632, E.140678, E.140694, E.140743, E.140932, E.141002, E.141012, E.141258, E.141378, E.141425, E.141429, E.141522, E.141543, E.141639, E.141744, E.141873, E.141994, E.142025, E.142208, E.142515, E.142606, E.142698, E.142765, E.142864, E.142875, E.143013, E.143294, E.143321, E.143353, E.143374, E.143375, E.143390, E.143578, E.143614, E.143621, E.143633, E.143771, E.143797, E.143816, E.143889, E.143924, E.143933, E.143947, E.144136, E.144224, E.144306, E.144381, E.144410, E.144485, E.144566, E.144671, E.144741, E.144935, E.145020, E.145632, E.145741, E.145833, E.145888, E.145907, E.145908, E.145919, E.145990, E.146067, E.146070, E.146281, E.146433, E.146457, E.146535, E.146701, E.146856, E.146966, E.147044, E.147127, E.147130, E.147133, E.147140, E.147231, E.147257, E.147403, E.147475, E.147548, E.147697, E.147724, E.148158, E.148396, E.148488, E.148672, E.148737, E.148835, E.149182, E.149218, E.149311, E.149480, E.149548, E.149646, E.150051, E.150593, E.150961, E.150991, E.151092, E.151093, E.151247, E.151304, E.151491, E.151690, E.151715, E.151726, E.151779, E.151806, E.152086, E.152207, E.152234, E.152291, E.152359, E.152377, E.152409, E.152422, E.152582, E.152763, E.152818, E.152942, E.153113, E.153140, E.153391, E.153904, E.153936, E.154099, E.154127, E.154380, E.154639, E.154723, E.154781, E.154832, E.154864, E.154889, E.154957, E.155368, E.155380, E.155508, E.155660, E.155714, E.155959, E.155980, E.156006, E.156194, E.156282, E.156304, E.156467, E.156515, E.156603, E.156650, E.156735, E.156976, E.157064, E.157103, E.157502, E.157510, E.157538, E.157551, E.157637, E.157764, E.157827, E.157992, E.158042, E.158290, E.158321, E.158485, E.158545, E.158604, E.158669, E.158715, E.158747, E.158813, E.158863, E.158901, E.158941, E.158987, E.159147, E.159184, E.159348, E.159363, E.159387, E.159423, E.159658, E.159692, E.159761, E.159921, E.160049, E.160226, E.160285, E.160294, E.160633, E.160685, E.160691, E.160789, E.160862, E.160867, E.160948, E.160972, E.161202, E.161267, E.161649, E.161692, E.161714, E.161813, E.161939, E.162069, E.162298, E.162385, E.162437, E.162490,

	E.162613, E.162641, E.162694, E.162910, E.162975, E.163041, E.163064, E.163110, E.163257, E.163468, E.163492, E.163530, E.163576, E.163629, E.163644, E.163749, E.163755, E.163781, E.163825, E.163913, E.163923, E.163930, E.163932, E.164045, E.164051, E.164053, E.164163, E.164244, E.164270, E.164300, E.164309, E.164442, E.164488, E.164520, E.164597, E.164749, E.164754, E.164828, E.164916, E.164919, E.164924, E.165084, E.165119, E.165138, E.165215, E.165259, E.165264, E.165280, E.165359, E.165410, E.165496, E.165637, E.165646, E.165661, E.165688, E.165695, E.165699, E.165792, E.165807, E.165813, E.165898, E.165923, E.165934, E.166263, E.166266, E.166329, E.166337, E.166341, E.166484, E.166526, E.166596, E.166598, E.166710, E.166747, E.166833, E.166860, E.166946, E.166971, E.167004, E.167085, E.167110, E.167113, E.167258, E.167513, E.167552, E.167553, E.167604, E.167635, E.167642, E.167658, E.167699, E.167744, E.167751, E.167766, E.167772, E.167799, E.167815, E.167969, E.167978, E.167987, E.167996, E.168014, E.168036, E.168066, E.168071, E.168148, E.168298, E.168393, E.168575, E.168653, E.168746, E.168763, E.168769, E.168803, E.168916, E.169087, E.169093, E.169122, E.169189, E.169213, E.169242, E.169410, E.169418, E.169562, E.169592, E.169612, E.169710, E.169763, E.169789, E.169807, E.169826, E.169957, E.170017, E.170027, E.170037, E.170088, E.170144, E.170275, E.170310, E.170315, E.170348, E.170374, E.170381, E.170396, E.170421, E.170430, E.170445, E.170549, E.170632, E.170703, E.170743, E.170837, E.170854, E.170906, E.170927, E.170954, E.170959, E.171121, E.171155, E.171180, E.171202, E.171262, E.171302, E.171345, E.171428, E.171488, E.171490, E.171492, E.171540, E.171643, E.171680, E.171723, E.171793, E.171861, E.171953, E.172115, E.172283, E.172345, E.172346, E.172466, E.172590, E.172594, E.172653, E.172717, E.172725, E.172733, E.172831, E.172867, E.172893, E.172939, E.173039, E.173230, E.173366, E.173473, E.173540, E.173585, E.173599, E.173714, E.173726, E.173805, E.173809, E.173826, E.173889, E.173898, E.173905, E.174021, E.174100, E.174332, E.174842, E.174996, E.175063, E.175110, E.175166, E.175175, E.175182, E.175198, E.175203, E.175216, E.175220, E.175334, E.175416, E.175602, E.175866, E.175946, E.176102, E.176105, E.176155, E.176171, E.176371, E.176515, E.176900, E.176971, E.176978, E.176994, E.177156, E.177239, E.177354, E.177409, E.177425, E.177459, E.177542, E.177548, E.177565, E.177595, E.177628, E.177674, E.177679, E.177694, E.177697, E.177731, E.177752, E.177951, E.178026, E.178078, E.178104, E.178163, E.178175, E.178187, E.178234, E.178381, E.178473, E.178741, E.178828, E.178950, E.179091, E.179115, E.179119, E.179348, E.179388, E.179776, E.179796, E.179869, E.179912, E.179981, E.180035, E.180198, E.180287, E.180318, E.180667, E.180869, E.180979, E.180998, E.181072, E.181163, E.181222, E.181234, E.181513, E.181523, E.181610, E.181773, E.181873, E.181885, E.181924, E.182013, E.182054, E.182217, E.182271, E.182318, E.182319, E.182512, E.182732, E.182795, E.182872, E.182890, E.182944, E.183048, E.183092, E.183098, E.183128, E.183207, E.183292, E.183431, E.183520, E.183684, E.183723, E.183785, E.183831, E.183856, E.184007, E.184047, E.184113, E.184156, E.184254, E.184363, E.184378, E.184470, E.184481, E.184508, E.184634, E.184661, E.184697, E.184708, E.184735, E.184840, E.184916, E.185043, E.185049, E.185122, E.185219, E.185359, E.185499, E.185554, E.185591, E.185619, E.185736, E.185860, E.185896, E.185945, E.185972, E.186198, E.186205, E.186376, E.186472, E.186575, E.186591, E.186660, E.186814, E.186834, E.186868, E.186889, E.187097, E.187323, E.187492, E.187634, E.187764, E.187792, E.187823, E.187837, E.187840, E.188021, E.188171, E.188186, E.188739, E.188771, E.188846, E.189060, E.189091, E.189143, E.189144, E.189221, E.189283, E.196236, E.196419, E.196436, E.196497, E.196504, E.196526, E.196591, E.196700, E.196743, E.196796, E.196812, E.196872, E.196975, E.196993, E.197081, E.197157, E.197217, E.197223, E.197299, E.197323, E.197353, E.197451, E.197479, E.197746, E.197779, E.197813, E.197837, E.197857, E.197872, E.197969, E.197976, E.198001, E.198033, E.198040, E.198087, E.198131, E.198156, E.198168, E.198205, E.198216, E.198231, E.198265, E.198366, E.198431, E.198455, E.198563, E.198586, E.198589, E.198712, E.198721, E.198732, E.198783, E.198793, E.198804, E.198807, E.198824, E.198841, E.198951, E.203301, E.203795, E.203813, E.203837, E.203879, E.203908, E.204231, E.204316, E.204389, E.204406, E.204560, E.204574
Prostate Markers (Ensembl ENSG identifiers)	E.005893 (LAMP2), E.006756 (ARSD), E.010539 (ZNF200), E.014257 (ACPP), E.015133 (CCDC88C), E.018699 (TTC27), E.044115 (CTNNA1), E.048828 (FAM120A), E.051620 (HEBP2), E.056097 (ZFR), E.060339 (CCAR1), E.063241 (ISOC2), E.064489 (MEF2BNB-MEF2B), E.064886 (CHI3L2), E.066455 (GOLGA5), E.069535 (MAOB), E.072042

	(RDH11), E.072071 (LPHN1), E.074047 (GLI2), E.076248 (UNG), E.076554 (TPD52), E.077147 (TM9SF3), E.077312 (SNRPA), E.081842 (PCDHA6), E.086717 (PPEF1), E.088888 (MAVS), E.088930 (XRN2), E.089902 (RCOR1), E.090612 (ZNF268), E.092199 (HNRNPC), E.095380 (NANS), E.099783 (HNRNPM), E.100191 (SLC5A4), E.100216 (TOMM22), E.100242 (SUN2), E.100284 (TOM1), E.100401 (RANGAP1), E.100412 (ACO2), E.100836 (PABPN1), E.102054 (RBBP7), E.102103 (PQBP1), E.103599 (IQCH), E.103978 (TMEM87A), E.104177 (MYEF2), E.104228 (TRIM35), E.105428 (ZNRF4), E.105518 (TMEM205), E.106603 (C7orf44; COA1), E.108405 (P2RX1), E.111057 (KRT18), E.111218 (PRMT8), E.112081 (SRSF3), E.112144 (ICK), E.113013 (HSPA9), E.113368 (LMNB1), E.115221 (ITGB6), E.116096 (SPR), E.116754 (SRSF11), E.118197 (DDX59), E.118898 (PPL), E.119121 (TRPM6), E.119711 (ALDH6A1), E.120656 (TAF12), E.121671 (CRY2), E.121774 (KHDRBS1), E.122126 (OCRL), E.122566 (HNRNPA2B1), E.123901 (GPR83), E.124562 (SNRPC), E.124788 (ATXN1), E.124795 (DEK), E.125246 (CLYBL), E.126883 (NUP214), E.127616 (SMARCA4), E.127884 (ECHS1), E.128050 (PAICS), E.129351 (ILF3), E.129757 (CDKN1C), E.130338 (TULP4), E.130612 (CYP2G1P), E.131351 (HAUS8), E.131467 (PSME3), E.133315 (MACROD1), E.134452 (FBXO18), E.134851 (TMEM165), E.135940 (COX5B), E.136169 (SETDB2), E.136807 (CDK9), E.137727 (ARHGAP20), E.138031 (ADCY3), E.138050 (THUMPD2), E.138069 (RAB1A), E.138594 (TMOD3), E.138760 (SCARB2), E.138796 (HADH), E.139613 (SMARCC2), E.139684 (ESD), E.140263 (SORD), E.140350 (ANP32A), E.140632 (GLYR1), E.142765 (SYTL1), E.143621 (ILF2), E.143933 (CALM2), E.144410 (CPO), E.147127 (RAB41), E.151304 (SRFBP1), E.151806 (GUF1), E.152207 (CYSLTR2), E.152234 (ATP5A1), E.152291 (TGOLN2), E.154723 (ATP5J), E.156467 (UQCRB), E.159387 (IRX6), E.159761 (C16orf86), E.161813 (LARP4), E.162613 (FUBP1), E.162694 (EXTL2), E.165264 (NDUFB6), E.167113 (COQ4), E.167513 (CDT1), E.167772 (ANGPTL4), E.167978 (SRRM2), E.168916 (ZNF608), E.169763 (PRYP3), E.169789 (PRY), E.169807 (PRY2), E.170017 (ALCAM), E.170144 (HNRNPA3), E.170310 (STX8), E.170954 (ZNF415), E.170959 (DCDC5), E.171302 (CANT1), E.171643 (S100Z), E.172283 (PRYP4), E.172590 (MRPL52), E.172867 (KRT2), E.173366 (TLR9), E.173599 (PC), E.177595 (PIDD), E.178473 (UCN3), E.179981 (TSHZ1), E.181163 (NPM1), E.182319 (Tyrosine-protein kinase SgK223), E.182795 (C1orf116), E.182944 (EWSR1), E.183092 (BEGAIN), E.183098 (GPC6), E.184254 (ALDH1A3), E.185619 (PCGF3), E.186889 (TMEM17), E.187837 (HIST1H1C), E.188771 (C11orf34), E.189060 (H1F0), E.196419 (XRCC6), E.196436 (NPIPL2), E.196504 (PRPF40A), E.196796, E.196993, E.197451 (HNRNPAB), E.197746 (PSAP), E.198131 (ZNF544), E.198156, E.198732 (SMOC1), E.198793 (MTOR), E.039068 (CDH1), E.173230 (GOLGB1), E.124193 (SRSF6), E.140497 (SCAMP2), E.168393 (DTYMK), E.184708 (EIF4ENIF1), E.124164 (VAPB), E.125753 (VASP), E.118260 (CREB1), E.135052 (GOLM1), E.010244 (ZNF207), E.010278 (CD9), E.047597 (XK), E.049246 (PER3), E.069849 (ATP1B3), E.072506 (HSD17B10), E.081138 (CDH7), E.099785 (MARCH2), E.104331 (IMPAD1), E.104812 (GYS1), E.120868 (APAF1), E.123908 (EIF2C2), E.125492 (BARHL1), E.127328 (RAB3IP), E.127329 (PTPRB), E.129514 (FOXA1), E.129657 (SEC14L1), E.129990 (SYT5), E.132881 (RSG1), E.136521 (NDUFB5), E.138347 (MYPN), E.141429 (GALNT1), E.144566 (RAB5A), E.151715 (TMEM45B), E.152582 (SPEF2), E.154957 (ZNF18), E.162385 (MAGOH), E.165410 (CFL2), E.168298 (HIST1H1E), E.169418 (NPR1), E.178187 (ZNF454), E.178741 (COX5A), E.179115 (FARSA), E.182732 (RGS6), E.183431 (SF3A3), E.185049 (WHSC2), E.196236 (XPNPEP3), E.197217 (ENTPD4), E.197813, E.203301, E.116833 (NR5A2), E.121057 (AKAP1), E.005471 (ABCB4), E.071859 (FAM50A), E.084234 (APLP2), E.101222 (SPEF1), E.103175 (WFDC1), E.103449 (SALL1), E.104805 (NUCB1), E.105514 (RAB3D), E.107816 (LZTS2), E.108375 (RNF43), E.109790 (KLHL5), E.112039 (FANCE), E.112715 (VEGFA), E.121690 (DEPDC7), E.125352 (RNF113A), E.134548 (C12orf39), E.136152 (COG3), E.143816 (WNT9A), E.147130 (ZMYM3), E.148396 (SEC16A), E.151092 (NGLY1), E.151779 (NBAS), E.155508 (CNOT8), E.163755 (HPS3), E.166526 (ZNF3), E.172733 (PURG), E.176371 (ZSCAN2), E.177674 (AGTRAP), E.181773 (GPR3), E.183048 (SLC25A10; MRPL12 SLC25A10), E.186376 (ZNF75D), E.187323 (DCC), E.198712 (MT-CO2), E.203908 (C6orf221; KHDC3L), E.001497 (LAS1L), E.009694 (ODZ1), E.080572 (CXorf41; PIH1D3), E.083093 (PALB2), E.089048 (ESF1), E.100065 (CARD10), E.100739 (BDKRB1), E.102904 (TSNAXIP1), E.104824 (HNRNPL), E.107404 (DVL1), E.110066 (SUV420H1), E.120328 (PCDHB12), E.121903 (ZSCAN20), E.122025 (FLT3), E.136930 (PSMB7), E.142025 (DMRTC2), E.144136 (SLC20A1), E.146535 (GNA12), E.147140 (NONO), E.153391 (INO80C), E.164919 (COX6C), E.171540 (OTP), E.177951 (BET1L), E.179796 (LRRC3B), E.197479

	(PCDHB11), E.198804 (MT-CO1), E.086205 (FOLH1), E.100632 (ERH), E.100796 (SMEK1), E.104760 (FGL1), E.114302 (PRKAR2A), E.130299 (GTPBP3), E.133961 (NUMB), E.144485 (HES6), E.167085 (PHB), E.167635 (ZNF146), E.177239 (MAN1B1), E.184481 (FOXO4), E.188171 (ZNF626), E.189221 (MAOA), E.157637 (SLC38A10), E.100883 (SRP54), E.105618 (PRPF31), E.119421 (NDUFA8), E.170837 (GPR27), E.168148 (HIST3H3), E.135525 (MAP7), E.174996 (KLC2), E.018189 (RUFY3), E.183520 (UTP11L), E.173905 (GOLIM4), E.165280 (VCP), E.022976 (ZNF839), E.059691 (PET112), E.063244 (U2AF2), E.075651 (PLD1), E.089177 (KIF16B), E.089280 (FUS), E.094755 (GABRP), E.096060 (FKBP5), E.100023 (PPIL2), E.100359 (SGSM3), E.100612 (DHRS7), E.104131 (EIF3J), E.104419 (NDRG1), E.105409 (ATP1A3), E.107623 (GDF10), E.111335 (OAS2), E.113522 (RAD50), E.115053 (NCL), E.120837 (NFYB), E.122733 (KIAA1045), E.123178 (SPRYD7), E.124181 (PLCG1), E.126858 (RHOT1), E.128609 (NDUFA5), E.128683 (GAD1), E.130255 (RPL36), E.133874 (RNF122), E.135387 (CAPRIN1), E.135999 (EPC2), E.136383 (ALPK3), E.139405 (C12orf52), E.141012 (GALNS), E.143924 (EML4), E.144671 (SLC22A14), E.145741 (BTF3), E.145907 (G3BP1), E.149311 (ATM), E.153113 (CAST), E.157538 (DSCR3), E.157992 (KRTCAP3), E.158901 (WFDC8), E.165259 (HDX), E.169410 (PTPN9), E.170421 (KRT8), E.171155 (C1GALT1C1), E.172831 (CES2), E.173726 (TOMM20), E.176515, E.177565 (TBL1XR1), E.177628 (GBA), E.179091 (CYC1), E.189091 (SF3B3), E.197299 (BLM), E.197872 (FAM49A), E.198205 (ZXDA), E.198455 (ZXDB), E.082212 (ME2), E.109956 (B3GAT1), E.169710 (FASN), E.011304 (PTBP1), E.057252 (SOAT1), E.059378 (PARP12), E.082258 (CCNT2), E.087301 (TXNDC16), E.100575 (TIMM9), E.101152 (DNAJC5), E.101812 (H2BFM), E.102384 (CENPI), E.108641 (B9D1), E.119138 (KLF9), E.119820 (YIPF4), E.125995 (ROMO1), E.132323 (ILKAP), E.134809 (TIMM10), E.134955 (SLC37A2), E.135476 (ESPL1), E.136527 (TRA2B), E.137776 (SLTM), E.139211 (AMIGO2), E.139428 (MMAB), E.139874 (SSTR1), E.143321 (HDGF), E.164244 (PRRC1), E.164270 (HTR4), E.165119 (HNRNPK), E.165637 (VDAC2), E.165661 (QSOX2), E.167258 (CDK12), E.167815 (PRDX2), E.168014 (C2CD3), E.168653 (NDUFS5), E.168769 (TET2), E.169242 (EFNA1), E.175334 (BANF1), E.175416 (CLTB), E.177156 (TALDO1), E.180035 (ZNF48), E.186591 (UBE2H), E.187097 (ENTPD5), E.188739 (RBM34), E.196497 (IPO4), E.197323 (TRIM33), E.197857 (ZNF44), E.197976 (AKAP17A), E.064201 (TSPAN32), E.088992 (TESC), E.092421 (SEMA6A), E.100601 (ALKBH1), E.101557 (USP14), E.103035 (PSMD7), E.106128 (GHRHR), E.115541 (HSPE1), E.121390 (PSPC1), E.124216 (SNAI1), E.130713 (EXOSC2), E.132002 (DNAJB1), E.139697 (SBNO1), E.140481 (CCDC33), E.143013 (LMO4), E.145020 (AMT), E.145990 (GFOD1), E.146070 (PLA2G7), E.164924 (YWHAZ), E.165807 (PPP1R36), E.167751 (KLK2), E.169213 (RAB3B), E.170906 (NDUFA3), E.172725 (CORO1B), E.174332 (GLIS1), E.181924 (CHCHD8), E.183128 (CALHM3), E.204560 (DHX16), E.204574 (ABCF1), E.146701 (MDH2), E.198366 (HIST1H3A), E.081181 (ARG2), E.185896 (LAMP1), E.077514 (POLD3), E.099800 (TIMM13), E.100299 (ARSA), E.105419 (MEIS3), E.108417 (KRT37), E.113739 (STC2), E.125868 (DSTN), E.145908 (ZNF300), E.168575 (SLC20A2), E.182271 (TMIGD1), E.197223 (C1D), E.186834 (HEXIM1), E.001561 (ENPP4), E.011451 (WIZ), E.053108 (FSTL4), E.064655 (EYA2), E.065308 (TRAM2), E.075131 (TIPIN), E.081087 (OSTM1), E.092020 (PPP2R3C), E.096384 (HSP90AB1), E.100348 (TXN2), E.100577 (GSTZ1), E.100802 (C14orf93), E.101365 (IDH3B), E.101654 (RNMT), E.103067 (ESRP2), E.104064 (GABPB1), E.104823 (ECH1), E.106565 (TMEM176B), E.108561 (C1QBP), E.115257 (PCSK4), E.116127 (ALMS1), E.117411 (B4GALT2), E.119335 (SET), E.120337 (TNFSF18), E.122033 (MTIF3), E.122507 (BBS9), E.122870 (BICC1), E.130177 (CDC16), E.130193 (C8orf55; THEM6), E.130413 (STK33), E.130770 (ATPIF1), E.133687 (TMTC1), E.136874 (STX17), E.137409 (MTCH1), E.139626 (ITGB7), E.141744 (PNMT), E.145888 (GLRA1), E.146067 (FAM193B), E.146433 (TMEM181), E.149480 (MTA2), E.152377 (SPOCK1), E.152763 (WDR78), E.156976 (EIF4A2), E.157827 (FMNL2), E.158485 (CD1B), E.158863 (FAM160B2), E.161202 (DVL3), E.161714 (PLCD3), E.163064 (EN1), E.163468 (CCT3), E.164309 (CMYA5), E.164916 (FOXK1), E.165215 (CLDN3), E.167658 (EEF2), E.170549 (IRX1), E.171680 (PLEKHG5), E.178234 (GALNT11), E.179869 (ABCA13), E.179912 (R3HDM2), E.180869 (C1orf180), E.180979 (LRRC57), E.182872 (RBM10), E.183207 (RUVBL2), E.184113 (CLDN5), E.185972 (CCIN), E.189144 (ZNF573), E.197353 (LYPD2), E.197779 (ZNF81), E.198807 (PAX9), E.100442 (FKBP3), E.111790 (FGFR1OP2), E.136044 (APPL2), E.061794 (MRPS35), E.065427 (KARS), E.068885 (IFT80), E.104164 (PLDN; BLOC1S6), E.105127 (AKAP8), E.123066 (MED13L), E.124831 (LRRFIP1), E.125304 (TM9SF2), E.126934 (MAP2K2), E.130305 (NSUN5), E.135298 (BAI3), E.135900 (MRPL44), E.136371 (MTHFS), E.136574

	(GATA4), E.140326 (CDAN1), E.141378 (PTRH2), E.141543 (EIF4A3), E.150961 (SEC24D), E.155368 (DBI), E.161649 (CD300LG), E.161692 (DBF4B), E.162437 (RAVER2), E.163257 (DCAF16), E.163576 (EFHB), E.163781 (TOPBP1), E.163913 (IFT122), E.164597 (COG5), E.165359 (DDX26B), E.165646 (SLC18A2), E.169592 (INO80E), E.169957 (ZNF768), E.171492 (LRRC8D), E.171793 (CTPS; CTPS1), E.171953 (ATPAF2), E.175182 (FAM131A), E.177354 (C10orf71), E.181610 (MRPS23), E.181873 (IBA57), E.187792 (ZNF70), E.187823 (ZCCHC16), E.196872 (C2orf55; KIAA1211L), E.198168 (SVIP), E.160633 (SAFB), E.177697 (CD151), E.181072 (CHRM2), E.012779 (ALOX5), E.065054 (SLC9A3R2), E.074071 (MRPS34), E.100815 (TRIP11), E.102030 (NAA10), E.106153 (CHCHD2), E.126814 (TRMT5), E.126952 (NXF5), E.136450 (SRSF1), E.136710 (CCDC115), E.137124 (ALDH1B1), E.143353 (LYPLAL1), E.162490 (C1orf187; DRAXIN), E.167799 (NUDT8), E.171490 (RSL1D1), E.173826 (KCNH6), E.173898 (SPTBN2), E.176900 (OR51T1), E.181513 (ACBD4), E.185554 (NXF2), E.185945 (NXF2B), E.108848 (LUC7L3), E.029363 (BCLAF1), E.038002 (AGA), E.108312 (UBTF), E.166341 (DCHS1), E.054118 (THRAP3), E.135679 (MDM2), E.166860 (ZBTB39), E.183684 (THOC4; ALYREF), E.004838 (ZMYND10), E.007264 (MATK), E.020922 (MRE11A), E.041353 (RAB27B), E.052795 (FNIP2), E.075711 (DLG1), E.087087 (SRRT), E.090060 (PAPOLA), E.095139 (ARCN1), E.099715 (PCDH11Y), E.100271 (TTLL1), E.101057 (MYBL2), E.101265 (RASSF2), E.101901 (ALG13), E.102290 (PCDH11X), E.103194 (USP10), E.106554 (CHCHD3), E.107833 (NPM3), E.110063 (DCPS), E.111540 (RAB5B), E.113448 (PDE4D), E.115339 (GALNT3), E.116254 (CHD5), E.117425 (PTCH2), E.117614 (SYF2), E.118181 (RPS25), E.118292 (C1orf54), E.119318 (RAD23B), E.121022 (COPS5), E.121104 (FAM117A), E.123427 (METTL21B), E.125676 (THOC2), E.132275 (RRP8), E.137513 (NARS2), E.138028 (CGREF1), E.139517 (LNX2), E.143614 (GATAD2B), E.143889 (HNRPLL), E.145833 (DDX46), E.147403 (RPL10), E.148158 (SNX30), E.151690 (MFSD6), E.153904 (DDAH1), E.154781 (C3orf19), E.156650 (KAT6B), E.158669 (AGPAT6), E.159363 (ATP13A2), E.163530 (DPPA2), E.164749 (HNF4G), E.165496 (RPL10L), E.165688 (PMPCA), E.165792 (METTL17), E.166598 (HSP90B1), E.168036 (CTNNB1), E.168746 (C20orf62), E.170381 (SEMA3E), E.171180 (OR2M4), E.171202 (TMEM126A), E.172594 (SMPDL3A), E.172653 (C17orf66), E.173540 (GMPPB), E.173585 (CCR9), E.173809 (TDRD12), E.175166 (PSMD2), E.177694 (NAALADL2), E.178026 (FAM211B; C22orf36), E.184363 (PKP3), E.187634 (SAMD11), E.203837 (PNLIPRP3), E.169122 (FAM110B), E.197969 (VPS13A), E.136068 (FLNB), E.075856 (SART3), E.081721 (DUSP12), E.102158 (MAGT1), E.102174 (PHEX), E.102316 (MAGED2), E.104723 (TUSC3), E.105939 (ZC3HAV1), E.108883 (EFTUD2), E.110328 (GALNTL4), E.111785 (RIC8B), E.113163 (COL4A3BP), E.115604 (IL18R1), E.117362 (APH1A), E.117480 (FAAH), E.124767 (GLO1), E.126267 (COX6B1), E.130175 (PRKCSH), E.135926 (TMBIM1), E.138674 (SEC31A), E.140451 (PIF1), E.143797 (MBOAT2), E.149646 (C20orf152), E.157064 (NMNAT2), E.160294 (MCM3AP), E.165084 (C8orf34), E.166946 (CCNDBP1), E.170348 (TMED10), E.170703 (TTLL6), E.175198 (PCCA), E.180287 (PLD5), E.183292 (MIR5096), E.187492 (CDHR4), E.188846 (RPL14), E.015479 (MATR3), E.100823 (APEX1), E.090615 (GOLGA3), E.086062 (B4GALT1), E.138385 (SSB), E.140265 (ZSCAN29), E.140932 (CMTM2), E.167969 (ECI1), E.135486 (HNRNPA1), E.137497 (NUMA1), E.181523 (SGSH), E.099956 (SMARCB1), E.049883 (PTCD2), E.082556 (OPRK1), E.090674 (MCOLN1), E.107164 (FUBP3), E.108582 (CPD), E.109758 (HGFAC), E.111605 (CPSF6), E.115239 (ASB3), E.121892 (PDS5A), E.125844 (RRBP1), E.130826 (DKC1), E.132481 (TRIM47), E.135390 (ATP5G2), E.136875 (PRPF4), E.138621 (PPCDC), E.145632 (PLK2), E.150051 (MKX), E.153140 (CETN3), E.154127 (UBASH3B), E.156194 (PPEF2), E.163825 (RTP3), E.164053 (ATRIP), E.164442 (CITED2), E.168066 (SF1), E.170430 (MGMT), E.175602 (CCDC85B), E.177752 (YIPF7), E.182512 (GLRX5), E.188186 (C7orf59), E.198721 (ECI2), E.204389 (HSPA1A), E.010256 (UQCRC1), E.076043 (REXO2), E.102362 (SYTL4), E.161939 (C17orf49), E.173039 (RELA), E.014216 (CAPN1), E.054938 (CHRDL2), E.065526 (SPEN), E.070501 (POLB), E.078808 (SDF4), E.083720 (OXCT1), E.100084 (HIRA), E.101246 (ARFRP1), E.102241 (HTATSF1), E.103591 (AAGAB), E.104626 (ERI1), E.105221 (AKT2), E.105402 (NAPA), E.105705 (SUGP1), E.106346 (USP42), E.108639 (SYNGR2), E.110107 (PRPF19), E.112473 (SLC39A7), E.113282 (CLINT1), E.115234 (SNX17), E.115561 (CHMP3), E.119906 (FAM178A), E.120733 (KDM3B), E.125375 (ATP5S), E.125798 (FOXA2), E.127415 (IDUA), E.129810 (SGOL1), E.132382 (MYBBP1A), E.133313 (CNDP2), E.134077 (THUMPD3), E.134248 (HBXIP), E.135597 (REPS1), E.137814 (HAUS2), E.138041 (SMEK2), E.140382 (HMG20A), E.143578 (CREB3L4), E.144224 (UBXN4),

	E.144306 (SCRN3), E.144741 (SLC25A26), E.145919 (BOD1), E.146281 (PM20D2), E.152359 (POC5), E.152409 (JMY), E.154889 (MPPE1), E.157551 (KCNJ15), E.157764 (BRAF), E.158987 (RAPGEF6), E.162069 (CCDC64B), E.162910 (MRPL55), E.163749 (CCDC158), E.164045 (CDC25A), E.164300 (SERINC5), E.165898 (ISCA2), E.167987 (VPS37C), E.168763 (CNNM3), E.170374 (SP7), E.171488 (LRRC8C), E.178381 (ZFAND2A), E.180998 (GPR137C), E.182318 (ZSCAN22), E.198040 (ZNF84), E.198216 (CACNA1E), E.198265 (HELZ), E.198586 (TLK1), E.203795 (FAM24A), E.204231 (RXRB), E.123992 (DNPEP), E.184634 (MED12), E.181885 (CLDN7), E.186660 (ZFP91), E.126777 (KTN1), E.080823 (MOK), E.101811 (CSTF2), E.124570 (SERPINB6), E.148835 (TAF5), E.158715 (SLC45A3), E.110955 (ATP5B), E.127022 (CANX), E.142208 (AKT1), E.128881 (TTBK2), E.147231 (CXorf57), E.006210 (CX3CL1), E.009830 (POMT2), E.011114 (BTBD7), E.065057 (NTHL1), E.068724 (TTC7A), E.073584 (SMARCE1), E.079785 (DDX1), E.084463 (WBP11), E.091140 (DLD), E.099821 (POLRMT), E.101126 (ADNP), E.104442 (ARMC1), E.105486 (LIG1), E.110921 (MVK), E.113441 (LNPEP), E.115758 (ODC1), E.116726 (PRAMEF12), E.119681 (LTBP2), E.136933 (RABEPK), E.137815 (RTF1), E.138095 (LRPPRC), E.138294 (MSMB), E.141873 (SLC39A3), E.142698 (C1orf94), E.143390 (RFX5), E.148488 (ST8SIA6), E.148737 (TCF7L2), E.151491 (EPS8), E.152422 (XRCC4), E.154832 (CXXC1), E.158321 (AUTS2), E.159147 (DONSON), E.160285 (LSS), E.160862 (AZGP1), E.160948 (VPS28), E.160972 (PPP1R16A), E.165934 (CPSF2), E.167604 (NFKBID), E.167766 (ZNF83), E.168803 (ADAL), E.169612 (FAM103A1), E.171262 (FAM98B), E.172893 (DHCR7), E.173889 (PHC3), E.176971 (FIBIN), E.177548 (RABEP2), E.179119 (SPTY2D1), E.184378 (ACTRT3), E.184508 (HDDC3), E.185043 (CIB1), E.186814 (ZSCAN30), E.186868 (MAPT), E.196812 (ZSCAN16), E.198563 (DDX39B), E.124529 (HIST1H4B), E.141002 (TCF25), E.174100 (MRPL45), E.109814 (UGDH), E.138756 (BMP2K), E.065457 (ADAT1), E.105948 (TTC26), E.109184 (DCUN1D4), E.125257 (ABCC4), E.126062 (TMEM115), E.142515 (KLK3), E.144381 (HSPD1), E.166710 (B2M), E.198824 (CHAMP1), E.078902 (TOLLIP), E.099331 (MYO9B), E.102710 (FAM48A), E.107485 (GATA3), E.120948 (TARDBP), E.187764 (SEMA4D), E.103855 (CD276), E.117751 (PPP1R8), E.173714 (WFIKKN2), E.172115 (CYCS), E.005882 (PDK2), E.007952 (NOX1), E.008118 (CAMK1G), E.012061 (ERCC1), E.015171 (ZMYND11), E.036257 (CUL3), E.057608 (GDI2), E.058729 (RIOK2), E.071246 (VASH1), E.073050 (XRCC1), E.073350 (LLGL2), E.079246 (XRCC5), E.085733 (CTTN), E.091542 (ALKBH5), E.091732 (ZC3HC1), E.092621 (PHGDH), E.099899 (TRMT2A), E.099917 (MED15), E.101439 (CST3), E.103479 (RBL2), E.104611 (SH2D4A), E.105281 (SLC1A5), E.106392 (C1GALT1), E.107104 (KANK1), E.107798 (LIPA), E.108296 (CWC25), E.109572 (CLCN3), E.112110 (MRPL18), E.113790 (EHHADH), E.115648 (MLPH), E.117308 (GALE), E.117335 (CD46), E.118513 (MYB), E.118640 (VAMP8), E.119321 (FKBP15), E.122705 (CLTA), E.123983 (ACSL3), E.124232 (RBPJL), E.125901 (MRPS26), E.127399 (LRRC61), E.127554 (GFER), E.128708 (HAT1), E.129355 (CDKN2D), E.130340 (SNX9), E.130935 (NOL11), E.131771 (PPP1R1B), E.133863 (TEX15), E.134207 (SYT6), E.136935 (GOLGA1), E.141425 (RPRD1A), E.143374 (TARS2), E.143771 (CNIH4), E.146966 (DENND2A), E.148672 (GLUD1), E.150593 (PDCD4), E.153936 (HS2ST1), E.154099 (DNAAF1), E.156006 (NAT2), E.156282 (CLDN17), E.158545 (ZC3H18), E.158604 (TMED4), E.158813 (EDA), E.159184 (HOXB13), E.161267 (BDH1), E.163492 (CCDC141), E.163629 (PTPN13), E.164163 (ABCE1), E.164520 (RAET1E), E.165138 (ANKS6), E.165923 (AGBL2), E.166484 (MAPK7), E.166747 (AP1G1), E.166971 (AKTIP), E.167744 (NTF4), E.168071 (CCDC88B), E.169087 (HSPBAP1), E.170396 (ZNF804A), E.170445 (HARS), E.170632 (ARMC10), E.170743 (SYT9), E.171428 (NAT1), E.172346 (CSDC2), E.173805 (HAP1), E.175175 (PPM1E), E.175203 (DCTN2), E.177542 (SLC25A22), E.177679 (SRRM3), E.178828 (RNF186), E.182013 (PNMAL1), E.182054 (IDH2), E.182890 (GLUD2), E.184156 (KCNQ3), E.184697 (CLDN6), E.184735 (DDX53), E.184840 (TMED9), E.185219 (ZNF445), E.186198 (SLC51B), E.186205 (MOSC1; MARC1), E.189143 (CLDN4), E.196700 (ZNF512B), E.196743 (GM2A), E.198087 (CD2AP), E.198951 (NAGA), E.204406 (MBD5), E.002330 (BAD), E.105404 (RABAC1), E.114127 (XRN1), E.117713 (ARID1A), E.123143 (PKN1), E.130764 (LRRC47), E.131773 (KHDRBS3), E.137806 (NDUFAF1), E.142864 (SERBP1), E.158747 (NBL1), E.175063 (UBE2C), E.178104 (PDE4DIP), E.186472 (PCLO), E.069956 (MAPK6), E.112941 (PAPD7), E.116604 (MEF2D), E.142875 (PRKACB), E.147133 (TAF1), E.157510 (AFAP1L1), E.006625 (GGCT), E.155980 (KIF5A), E.134444 (KIAA1468), E.107968 (MAP3K8), E.117592 (PRDX6), E.123154 (WDR83), E.135297 (MTO1), E.135829 (DHX9), E.149548 (CCDC15), E.152086 (TUBA3E), E.167553 (TUBA1C),

	E.169826 (CSGALNACT2), E.171121 (KCNMB3), E.198033 (TUBA3C), E.147724 (FAM135B), E.170854 (MINA), E.006695 (COX10), E.067369 (TP53BP1), E.089248 (ERP29), E.112096 (SOD2), E.138073 (PREB), E.146856 (AGBL3), E.159423 (ALDH4A1), E.171345 (KRT19), E.172345 (STARD5), E.111647 (UHRF1BP1L), E.117877 (CD3EAP), E.155714 (PDZD9), E.156603 (MED19), E.075886 (TUBA3D), E.167699 (GLOD4), E.121749 (TBC1D15), E.090861 (AARS), E.093010 (COMT), E.117676 (RPS6KA1), E.157502 (MUM1L1), E.159921 (GNE), E.169562 (GJB1), E.179776 (CDH5), E.071626 (DAZAP1), E.085224 (ATRX), E.116478 (HDAC1), E.117298 (ECE1), E.176171 (BNIP3), E.177425 (PAWR), E.179348 (GATA2), E.187840 (EIF4EBP1), E.033030 (ZCCHC8), E.049239 (H6PD), E.060688 (SNRNP40), E.075239 (ACAT1), E.095627 (TDRD1), E.109625 (CPZ), E.113719 (ERGIC1), E.126773 (C14orf135; PCNXL4), E.149218 (ENDOD1), E.162975 (KCNF1), E.183785 (TUBA8), E.198589 (LRBA), E.105379 (ETFB), E.011052 (NME2), E.011143 (MKS1), E.048544 (MRPS10), E.062485 (CS), E.114054 (PCCB), E.138587 (MNS1), E.155959 (VBP1), E.181222 (POLR2A), E.183723 (CMTM4), E.184661 (CDCA2), E.204316 (MRPL38), E.140694 (PARN), E.035141 (FAM136A), E.095485 (CWF19L1), E.115540 (MOB4), E.123595 (RAB9A), E.140678 (ITGAX), E.141258 (SGSM2), E.158941 (KIAA1967), E.169189 (NSMCE1), E.198431 (TXNRD1), E.016402 (IL20RA), E.112234 (FBXL4), E.125445 (MRPS7), E.128342 (LIF), E.164051 (CCDC51), E.175866 (BAIAP2), E.102780 (DGKH), E.203813 (HIST1H3H), E.198231 (DDX42), E.030582 (GRN), E.106049 (HIBADH), E.130810 (PPAN), E.132475 (H3F3B), E.158290 (CUL4B), E.166266 (CUL5), E.026559 (KCNG1), E.059122 (FLYWCH1), E.107897 (ACBD5), E.121068 (TBX2), E.125944 (HNRNPR), E.134308 (YWHAQ), E.137558 (PI15), E.137601 (NEK1), E.147548 (WHSC1L1), E.149182 (ARFGAP2), E.159658 (KIAA0494), E.165699 (TSC1), E.170927 (PKHD1), E.186575 (NF2), E.188021 (UBQLN2), E.167552 (TUBA1A), E.003756 (RBM5), E.134138 (MEIS2), E.008196 (TFAP2B), E.079313 (REXO1), E.089127 (OAS1), E.106078 (COBL), E.113645 (WWC1), E.116288 (PARK7), E.121940 (CLCC1), E.136280 (CCM2), E.141639 (MAPK4), E.147475 (ERLIN2), E.155660 (PDIA4), E.162298 (SYVN1), E.176978 (DPP7), E.176994 (SMCR8), E.178175 (ZNF366), E.196591 (HDAC2), E.127824 (TUBA4A), E.163932 (PRKCD), E.143375 (CGN), E.076864 (RAP1GAP), E.138772 (ANXA3), E.163041 (H3F3A), E.165813 (C10orf118), E.166337 (TAF10), E.178078 (STAP2), E.184007 (PTP4A2), E.167004 (PDIA3), E.039560 (RAI14), E.119636 (C14orf45), E.140374 (ETFA), E.143633 (C1orf131), E.144935 (TRPC1), E.156735 (BAG4), E.159348 (CYB5R1), E.170275 (CRTAP), E.172717 (FAM71D), E.172939 (OXSR1), E.176105 (YES1), E.078295 (ADCY2), E.119888 (EPCAM), E.141522 (ARHGDIA), E.184047 (DIABLO), E.109062 (SLC9A3R1), E.170037 (CNTROB), E.066557 (LRRC40), E.074964 (ARHGEF10L), E.078269 (SYNJ2), E.090013 (BLVRB), E.100142 (POLR2F), E.100399 (CHADL), E.104365 (IKBKB), E.111261 (MANSC1), E.111907 (TPD52L1), E.112578 (BYSL), E.121957 (GPSM2), E.122884 (P4HA1), E.124693 (HIST1H3B), E.126653 (NSRP1), E.130402 (ACTN4), E.138757 (G3BP2), E.150991 (UBC), E.164828 (SUN1), E.175216 (CKAP5), E.176155 (CCDC57), E.177459 (C8orf47), E.183856 (IQGAP3), E.185122 (HSF1), E.122952 (ZWINT), E.151093 (OXSM), E.067704 (IARS2), E.088899 (ProSAP-interacting protein 1), E.091483 (FH), E.114388 (NPRL2), E.114861 (FOXP1), E.135914 (HTR2B), E.197837 (HIST4H4), E.127720 (C12orf26; METTL25), E.123416 (TUBA1B), E.047410 (TPR), E.117748 (RPA2), E.133835 (HSD17B4), E.067248 (DHX29), E.121879 (PIK3CA), E.132589 (FLOT2), E.136750 (GAD2), E.160789 (LMNA), E.166329, E.170088 (TMEM192), E.175946 (KLHL38), E.178163 (ZNF518B), E.182217 (HIST2H4B), E.184470 (TXNRD2), E.110321 (EIF4G2), E.171861 (RNMTL1), E.065978 (YBX1), E.115738 (ID2), E.143294 (PRCC), E.158042 (MRPL17), E.169093 (ASMTL), E.090565 (RAB11FIP3), E.185591 (SP1), E.156304 (SCAF4), E.092978 (GPATCH2), E.100056 (DGCR14), E.100583 (SAMD15), E.105723 (GSK3A), E.107551 (RASSF4), E.107581 (EIF3A), E.107890 (ANKRD26), E.110104 (CCDC86), E.112584 (FAM120B), E.113580 (NR3C1), E.114491 (UMPS), E.137312 (FLOT1), E.137955 (RABGGTB), E.141994 (DUS3L), E.147044 (CASK), E.152818 (UTRN), E.180667 (YOD1), E.184916 (JAG2), E.196526 (AFAP1), E.198783 (ZNF830), E.108465 (CDK5RAP3), E.156515 (HK1), E.036448 (MYOM2), E.061918 (GUCY1B3), E.070785 (EIF2B3), E.116044 (NFE2L2), E.128311 (TST), E.131473 (ACLY), E.132716 (DCAF8), E.138363 (ATIC), E.166596 (WDR16), E.170027 (YWHAG), E.174021 (GNG5), E.203879 (GDI1), E.160049 (DFFA), E.010810 (FYN), E.051596 (THOC3), E.006453 (BAI1-associated protein 2-like 1), E.126945 (HNRNPH2), E.165695 (AK8), E.069869 (NEDD4), E.111801 (BTN3A3), E.112232 (KHDRBS2), E.128626 (MRPS12), E.129636 (ITFG1), E.137948 (BRDT), E.147257 (GPC3), E.155380 (SLC16A1), E.159692 (CTBP1), E.166833 (NAV2), E.172466

	(ZNF24), E.175110 (MRPS22), E.176102 (CSTF3), E.179388 (EGR3), E.185359 (HGS), E.198001 (IRAK4), E.100603 (SNW1), E.162641 (AKNAD1), E.069712 (KIAA1107), E.073756 (PTGS2), E.077522 (ACTN2), E.101639 (CEP192), E.106633 (GCK), E.115241 (PPM1G), E.116649 (SRM), E.120370 (GORAB), E.124143 (ARHGAP40), E.127948 (POR), E.129315 (CCNT1), E.132646 (PCNA), E.135740 (SLC9A5), E.151726 (ACSL1), E.154380 (ENAH), E.157103 (SLC6A1), E.163930 (BAP1), E.164488 (DACT2), E.164754 (RAD21), E.175220 (ARHGAP1), E.180318 (ALX1), E.181234 (TMEM132C), E.197081 (IGF2R), E.092871 (RFFL), E.163644 (PPM1K), E.171723 (GPHN), E.108953 (YWHAE), E.072110 (ACTN1), E.077097 (TOP2B), E.090889 (KIF4A), E.114331 (ACAP2), E.114867 (EIF4G1), E.117593 (DARS2), E.118523 (CTGF), E.120915 (EPHX2), E.134759 (ELP2), E.138061 (CYP1B1), E.140743 (CDR2), E.151247 (EIF4E), E.152942 (RAD17), E.160685 (ZBTB7B), E.163923 (RPL39L), E.167642 (SPINT2), E.167996 (FTH1), E.185736 (ADARB2), E.198841 (KTI12), E.185860 (C1orf110), E.160226 (C21orf2), E.070814 (TCOF1), E.124749 (COL21A1), E.154639 (CXADR), E.065485 (PDIA5), E.023909 (GCLM), E.100714 (MTHFD1), E.108387 (SEPT4), E.160867 (FGFR4), E.134684 (YARS), E.123080 (CDKN2C), E.065548 (ZC3H15), E.116455 (WDR77), E.117448 (AKR1A1), E.100393 (EP300), E.138160 (KIF11), E.166263 (STXBP4), E.173473 (SMARCC1), E.124942 (AHNAK), E.174842 (GLMN), E.180198 (RCC1), E.185499 (MUC1), E.143947 (RPS27A), E.170315 (UBB), E.003402 (CFLAR), E.137055 (PLAA), E.142606 (MMEL1), E.147697 (GSDMC), E.163110 (PDLIM5), E.135842 (FAM129A), E.160691 (SHC1), E.197157 (SND1), E.029725 (RABEP1), E.127946 (HIP1), E.001036 (FUCA2), E.109846 (CRYAB), E.183831 (ANKRD45), E.189283 (FHIT), E.092820 (EZR), E.104067 (TJP1), E.120159 (C9orf82; CAAP1), E.154864 (PIEZO2), E.196975 (ANXA4), E.105220 (GPI), E.127914 (AKAP9), E.135870 (RC3H1), E.026508 (CD44), E.089154 (GCN1L1), E.100311 (PDGFB), E.119383 (PPP2R4), E.075624 (ACTB), E.177409 (SAMD9L), E.177731 (FLII), E.015676 (NUDCD3), E.146457 (WTAP), E.178950 (GAK), E.167110 (GOLGA2)
Prostate vesicle	LAMP2, ACPP, CTNNA1, HEBP2, ISOC2, HNRNPC, HNRNPM, TOMM22, TOM1, ACO2, KRT18, HSPA9, LMNB1, SPR, PPL, ALDH6A1, HNRNPA2B1, ATXN1, SMARCA4, ECHS1, PAICS, ILF3, PSME3, COX5B, RAB1A, SCARB2, HADH, ESD, SORD, ILF2, CALM2, ATP5A1, TGOLN2, ANGPTL4, ALCAM, KRT2, PC, NPM1, C1orf116, GPC6, ALDH1A3, HIST1H1C, XRCC6, HNRNPAB, PSAP, CDH1, SCAMP2, VASP, CD9, ATP1B3, HSD17B10, APAF1, EIF2C2, RAB5A, CFL2, FARSA, XPNPEP3, ENTPD4, APLP2, NUCB1, RAB3D, VEGFA, HPS3, TSNAXIP1, HNRNPL, PSMB7, GNA12, NONO, FOLH1, PRKAR2A, PHB, HIST3H3, MAP7, VCP, U2AF2, FUS, FKBP5, NDRG1, ATP1A3, NCL, RPL36, KRT8, C1GALT1C1, FASN, PTBP1, TXNDC16, DNAJC5, SLC37A2, HNRNPK, VDAC2, PRDX2, TALDO1, USP14, PSMD7, HSPE1, DNAJB1, YWHAZ, RAB3B, CORO1B, MDH2, HIST1H3A, LAMP1, STC2, DSTN, SLC20A2, ENPP4, WIZ, HSP90AB1, IDH3B, ECH1, C1QBP, SET, TNFSF18, ITGB7, SPOCK1, EIF4A2, CCT3, CLDN3, EEF2, LRRC57, RUVBL2, CLDN5, APPL2, TM9SF2, EIF4A3, DBI, DBF4B, SVIP, CD151, ALOX5, SLC9A3R2, RAB27B, DLG1, ARCN1, CHCHD3, RAB5B, RPS25, RPL10, DDAH1, HSP90B1, CTNNB1, PSMD2, PKP3, FLNB, EFTUD2, GLO1, PRKCSH, TMBIM1, SEC31A, TMED10, RPL14, MATR3, APEX1, B4GALT1, HNRNPA1, CPD, HSPA1A, CAPN1, CHRDL2, SPEN, SDF4, NAPA, SYNGR2, CHMP3, CNDP2, CCDC64B, SERINC5, VPS37C, DNPEP, CLDN7, KTN1, SERPINB6, ATP5B, CANX, AKT1, TTBK2, DDX1, DLD, LNPEP, LTBP2, LRPPRC, EPS8, AZGP1, VPS28, DHCR7, CIB1, DDX39B, HIST1H4B, UGDH, HSPD1, B2M, TOLLIP, CD276, CYCS, CUL3, GDI2, LLGL2, XRCC5, CTTN, PHGDH, CST3, RBL2, SLC1A5, CD46, VAMP8, CLTA, ACSL3, MRPS26, SNX9, GLUD1, TMED4, PTPN13, AP1G1, SYT9, DCTN2, IDH2, GLUD2, TMED9, CLDN4, GM2A, CD2AP, MBD5, SERBP1, NBL1, PRKACB, GGCT, PRDX6, DHX9, TUBA3E, TUBA1C, TUBA3C, ERP29, SOD2, KRT19, TUBA3D, AARS, COMT, MUM1L1, CDH5, ECE1, ACAT1, ENDOD1, TUBA8, ETFB, NME2, CS, VBP1, RAB9A, TXNRD1, LIF, BAIAP2, HIST1H3H, GRN, HIBADH, H3F3B, CUL4B, HNRNPR, YWHAQ, PKHD1, TUBA1A, PARK7, ERLIN2, PDIA4, TUBA4A, PRKCD, ANXA3, H3F3A, PTP4A2, PDIA3, ETFA, CYB5R1, CRTAP, OXSR1, YES1, EPCAM, ARHGDIA, DIABLO, SLC9A3R1, BLVRB, P4HA1, HIST1H3B, ACTN4, UBC, FH, HIST4H4, TUBA1B, HSD17B4, PIK3CA, FLOT2, LMNA, TMEM192, HIST2H4B, YBX1, EIF3A, FLOT1, UTRN, HK1, ACLY, ATIC, YWHAG, GNG5, GDI1, HNRNPH2, NEDD4, BTN3A3, SLC16A1, HGS, ACTN2, SRM, PCNA, ACSL1, RAD21, ARHGAP1, IGF2R, YWHAE, ACTN1, EIF4G1, EPHX2, EIF4E, FTH1, CXADR, MTHFD1, AKR1A1, STXBP4, AHNAK, MUC1, RPS27A, UBB, PDLIM5, FAM129A, SND1, FUCA2, CRYAB, EZR, TJP1, ANXA4, GPI, AKAP9, CD44,

	GCN1L1, ACTB, FLII, NUDCD3
Prostate Cancer vesicles	EGFR, GLUD2, ANXA3, APLP2, BclG, Cofilin 2 /cfL2, DCTN-50 / DCTN2, DDAH1, ESD, FARSLA, GITRL, PRKCSH, SLC20A2, Synaptogyrin 2 /SYNGR2, TM9SF2, Calnexin, TOMM22, NDRG1, RPL10, RPL14, USP14, VDAC2, LLGL2, CD63, CD81, uPAR / CD87, ADAM 9, BDKRB2, CCR5, CCT2 (TCP1-beta), PSMA, PSMA1, HSPB1, VAMP8, Rab1A, B4GALT1, Aspartyl Aminopeptidase /Dnpep, ATPase Na+/K+ beta 3/ATP1B3, BDNF, ATPB, beta 2 Microglobulin, Calmodulin 2 /CALM2, CD9, XRCC5 / Ku80, SMARCA4, TOM1, Cytochrome C, Hsp10 / HSPE1, COX2 / PTGS2, Claudin 4 /CLDN4, Cytokeratin 8, Cortactin/CTTN, DBF4B /DRF1, ECH1, ECHS1, GOLPH2, ETS1, DIP13B /appl2, EZH2 / KMT6, GSTP1, hK2 / Kif2a, IQGAP1, KLK13, Lamp-2, GM2A, Hsp40/DNAJB1, HADH/HADHSC, Hsp90B, Nucleophosmin, p130 /RBL2, PHGDH, RAB3B, ANXA1, PSMD7, PTBP1, Rab5a, SCARB2, Stanniocalcin 2 /STC2, TGN46 /TGOLN2, TSNAXIP1, ANXA2, CD46, KLK14, IL1alpha, hnRNP C1 + C2, hnRNP A1, hnRNP A2B1, Claudin 5, CORO1B, Integrin beta 7, CD41, CD49d, CDH2, COX5b, IDH2, ME1, PhIP, ALDOA, EDNRB/EDN3, MTA1, NKX3-1, TMPRSS2, CD10, CD24, CDH1, ADAM10, B7H3, CD276, CHRDL2, SPOCK1, VEGFA, BCHE, CD151, CD166/ALCAM, CSE1L, GPC6, CXCR3, GAL3, GDF15, IGFBP-2, HGF, KLK12, ITGAL, KLK7, KLK9, MMP 2, MMP 25, MMP10, TNFRI, Notch1, PAP - same as ACPP, PTPN13/PTPL1, seprase/FAP, TNFR1, TWEAK, VEGFR2, E-Cadherin, Hsp60, CLDN3- Claudin3, KLK6, KLK8, EDIL3 (del-1), APE1, MMP 1, MMP3, nAnS, PSP94 / MSP / IGBF, PSAP, RPL19, SET, TGFB, TGM2, TIMP-1, TNFRII, MDH2, PKP1, Cystatin C, Trop2 / TACSTD2, CCR2 / CD192, hnRNP M1-M4, CDKN1A, CGA, Cytokeratin 18, EpoR, GGPS1, FTL (light and heavy), GM-CSF, HSP90AA1, IDH3B, MKI67/Ki67, LTBP2, KLK1, KLK4, KLK5, LDH-A, Nav1.7/SCN9A, NRP1 / CD304, PIP3 / BPNT1, PKP3, CgA, PRDX2, SRVN, ATPase Na+/K+ alpha 3/ATP1A3, SLC3A2 / CD98, U2AF2, TLR4 (CD284), TMPRSS1, TNFα, uPA, GloI, ALIX, PKM2, FABP5, CAV1, TLR9 / CD289, ANXA4, PLEKHC1 / Kindlin-2, CD71 / TRFR, MBD5, SPEN/ RBM15, LGALS8, SLC9A3R2, ENTPD4, ANGPTL4, p97 / VCP, TBX5, PTEN, Prohibitin, LSP1, HOXB13, DDX1, AKT1, ARF6, EZR, H3F3A, CIB1, Ku70 (XRCC6), KLK11, TMBIM6, SYT9, APAF1, CLDN7, MATR3, CD90/THY1, Tollip, NOTCH4, 14-3-3 zeta/beta, ATP5A1, DLG1, GRP94, FKBP5/FKBP51, LAMP1, LGALS3BP, GDI2, HSPA1A, NCL, KLK15, Cytokeratin basic, EDN-3, AGR2, KLK10, BRG1, FUS, Histone H4, hnRNP L, Catenin Alpha 1, hnRNP K (F45)*, MMP7*, DBI*, beta catenin, CTH, CTNND2, Ataxin 1, Proteasome 20S beta 7, ADE2, EZH2, GSTP1, Lamin B1, Coatomer Subunit Delta, ERAB, Mortalin, PKM2, IGFBP-3, CTNND1 / delta 1-catenin / p120-catenin, PKA R2, NONO, Sorbitol Dehydrogenase, Aconitase 2, VASP, Lipoamide Dehydrogenase, AP1G1, GOLPH2, ALDH6A1, AZGP1, Ago2, CNDP2, Nucleobindin-1, SerpinB6, RUVBL2, Proteasome 19S 10B, SH3PX1, SPR, Destrin, MDM4, FLNB, FASN, PSME
Prostate Cancer vesicles	14-3-3 zeta/beta, Aconitase 2, ADAM 9, ADAM10, ADE2, AFM, Ago2, AGR2, AKT1, ALDH1A3, ALDH6A1, ALDOA, ALIX, ANGPTL4, ANXA1, ANXA2, ANXA3, ANXA3, ANXA4, AP1G1, APAF1, APE1, APLP2, APLP2, ARF6, Aspartyl Aminopeptidase /Dnpep, Ataxin 1, ATP5A1, ATPase Na+/K+ alpha 3/ATP1A3, ATPase Na+/K+ beta 3/ATP1B3, ATPase Na+/K+ beta 3/ATP1B3, ATPB, AZGP1, B4GALT1, B7H3, BCHE, BclG, BDKRB2, BDNF, BDNF, beta 2 Microglobulin, beta catenin, BRG1, CALM2, Calmodulin 2 /CALM2, Calnexin, Calpain 1, Catenin Alpha 1, CAV1, CCR2 / CD192, CCR5, CCT2 (TCP1-beta), CD10, CD151, CD166/ALCAM, CD24, CD276, CD41, CD46, CD49d, CD63, CD71 / TRFR, CD81, CD9, CD9, CD90/THY1, CDH1, CDH2, CDKN1A, CGA, CgA, CHRDL2, CIB1, CIB1, Claudin 4 /CLDN4, Claudin 5, CLDN3, CLDN3- Claudin3, CLDN4, CLDN7, CNDP2, Coatomer Subunit Delta, Cofilin 2 /cfL2, CORO1B, Cortactin/CTTN, COX2 / PTGS2, COX5b, CSE1L, CTH, CTNND1 / delta 1-catenin / p120-catenin, CTNND2, CXCR3, CYCS, Cystatin C, Cytochrome C, Cytokeratin 18, Cytokeratin 8, Cytokeratin basic, DBF4B /DRF1, DBI*, DCTN-50 / DCTN2, DDAH1, DDAH1, DDX1, Destrin, DIP13B /appl2, DIP13B /appl2, DLG1, Dnpep, E-Cadherin, ECH1, ECHS1, ECHS1, EDIL3 (del-1), EDN-3, EDNRB/EDN3, EGFR, EIF4A3, ENTPD4, EpoR, EpoR, ERAB, ESD, ESD, ETS1, ETS1, ETS-2, EZH2, EZH2 / KMT6, EZR, FABP5, FARSLA, FASN, FKBP5/FKBP51, FLNB, FTL (light and heavy), FUS, GAL3, gamma-catenin, GDF15, GDI2, GGPS1, GGPS1, GITRL, GloI, GLUD2, GM2A, GM-CSF, GOLM1/GOLPH2 Mab; clone 3B10, GOLPH2, GOLPH2, GPC6, GRP94, GSTP1, GSTP1, H3F3A, HADH/HADHSC, HGF, HIST1H3A, Histone H4, hK2 / Kif2a, hnRNP A1, hnRNP A2B1, hnRNP C1 + C2, hnRNP K (F45)*, hnRNP L, hnRNP M1-M4, HOXB13, Hsp10 / HSPE1, Hsp40/DNAJB1, Hsp60, HSP90AA1, Hsp90B, HSPA1A, HSPB1, IDH2, IDH3B,

	IDH3B, IGFBP-2, IGFBP-3, IgG1, IgG2A, IgG2B, IL1alpha, IL1alpha, Integrin beta 7, IQGAP1, ITGAL, KLHL12/C3IP1, KLK1, KLK10, KLK11, KLK12, KLK13, KLK14, KLK15, KLK4, KLK5, KLK6, KLK7, KLK8, KLK9, Ku70 (XRCC6), Lamin B1, LAMP1, Lamp-2, LDH-A, LGALS3BP, LGALS8, Lipoamide Dehydrogenase, LLGL2, LSP1, LSP1, LTBP2, MATR3, MBD5, MDH2, MDM4, ME1, MKI67/Ki67, MMP 1, MMP 2, MMP 25, MMP10, MMP-14/MT1-MMP, MMP3, MMP7*, Mortalin, MTA1, nAnS, nAnS, Nav1.7/SCN9A, NCL, NDRG1, NKX3-1, NONO, Notch1, NOTCH4, NRP1 / CD304, Nucleobindin-1, Nucleophosmin, p130 /RBL2, p97 / VCP, PAP - same as ACPP, PHGDH, PhIP, PIP3 / BPNT1, PKA R2, PKM2, PKM2, PKP1, PKP3, PLEKHC1 / Kindlin-2, PRDX2, PRKCSH, Prohibitin, Proteasome 19S 10B, Proteasome 20S beta 7, PSAP, PSMA, PSMA1, PSMA1, PSMD7, PSMD7, PSME3, PSP94 / MSP / IGBF, PTBP1, PTEN, PTPN13/PTPL1, Rab1A, RAB3B, Rab5a, Rad51b, RPL10, RPL10, RPL14, RPL14, RPL19, RUVBL2, SCARB2, seprase/FAP, SerpinB6, SET, SH3PX1, SLC20A2, SLC3A2 / CD98, SLC9A3R2, SMARCA4, Sorbitol Dehydrogenase, SPEN/ RBM15, SPOCK1, SPR, SRVN, Stanniocalcin 2 /STC2, STEAP1, Synaptogyrin 2 /SYNGR2, Syndecan, SYNGR2, SYT9, TAF1B / GRHL1, TBX5, TGFB, TGM2, TGN46 /TGOLN2, TIMP-1, TLR3, TLR4 (CD284), TLR9 / CD289, TM9SF2, TMBIM6, TMPRSS1, TMPRSS2, TNFR1, TNFRI, TNFRII, TNFSF18 / GITRL, TNFα, TNFα, Tollip, TOM1, TOMM22, Trop2 / TACSTD2, TSNAXIP1, TWEAK, U2AF2, uPA, uPAR / CD87, USP14, USP14, VAMP8, VASP, VDAC2, VEGFA, VEGFR1/FLT1, VEGFR2, VPS28, XRCC5 / Ku80, XRCC5 / Ku80
Prostate Vesicles / General Vesicles	EpCAM/TROP-1, HSA, Fibrinogen, GAPDH, Cholesterol Oxidase, MMP7, Complement Factor D/Adipsin, E-Cadherin, Transferrin Antibody, eNOS, IgM, CD9, Apolipoprotein B (Apo B), Ep-CAM, TBG, Kallekerin 3, IgA, IgG, Annexin V, IgG, Pyruvate Carboxylase, trypsin, AFP, TNF RI/TNFRSF1A, Aptamer CAR023, Aptamer CAR024, Aptamer CAR025, Aptamer CAR026
Ribonucleoprotein complexes & vesicles	GW182, Ago2, miR-let-7a, miR-16, miR-22, miR-148a, miR-451, miR-92a, CD9, CD63, CD81
Prostate Cancer vesicles	PCSA, Muc2, Adam10
Prostate Cancer vesicles	Alkaline Phosphatase (AP), CD63, MyoD1, Neuron Specific Enolase, MAP1B, CNPase, Prohibitin, CD45RO, Heat Shock Protein 27, Collagen II, Laminin B1/b1, Gai1, CDw75, bcl-XL, Laminin-s, Ferritin, CD21, ADP-ribosylation Factor (ARF-6)
Prostate Cancer vesicles	CD56/NCAM-1, Heat Shock Protein 27/hsp27, CD45RO, MAP1B, MyoD1, CD45/T200/LCA, CD3zeta, Laminin-s, bcl-XL, Rad18, Gai1, Thymidylate Synthase, Alkaline Phosphatase (AP), CD63, MMP-16 / MT3-MMP, Cyclin C, Neuron Specific Enolase, SIRP a1, Laminin B1/b1, Amyloid Beta (APP), SODD (Silencer of Death Domain), CDC37, Gab-1, E2F-2, CD6, Mast Cell Chymase, Gamma Glutamylcysteine Synthetase (GCS)
Prostate Cancer vesicles	EpCAM, MMP7, PCSA, BCNP, ADAM10, KLK2, SPDEF, CD81, MFGE8, IL-8
Prostate Cancer vesicles	EpCAM, KLK2, PBP, SPDEF, SSX2, SSX4
Prostate Cancer vesicles	ADAM-10, BCNP, CD9, EGFR, EpCam, IL1B, KLK2, MMP7, p53, PBP, PCSA, SERPINB3, SPDEF, SSX2, SSX4
Androgen Receptor (AR) pathway members in cMVs	GTF2F1, CTNNB1, PTEN, APPL1, GAPDH, CDC37, PNRC1, AES, UXT, RAN, PA2G4, JUN, BAG1, UBE2I, HDAC1, COX5B, NCOR2, STUB1, HIPK3, PXN, NCOA4
EGFR1 pathway members in cMVs	RALBP1, SH3BGRL, RBBP7, REPS1, SNRPD2, CEBPB, APPL1, MAP3K3, EEF1A1, GRB2, RAC1, SNCA, MAP2K3, CEBPA, CDC42, SH3KBP1, CBL, PTPN6, YWHAB, FOXO1, JAK1, KRT8, RALGDS, SMAD2, VAV1, NDUFA13, PRKCB1, MYC, JUN, RFXANK, HDAC1, HIST3H3, PEBP1, PXN, TNIP1, PKN2
TNF-alpha pathway members in cMVs	BCL3, SMARCE1, RPS11, CDC37, RPL6, RPL8, PAPOLA, PSMC1, CASP3, AKT2, MAP3K7IP2, POLR2L, TRADD, SMARCA4, HIST3H3, GNB2L1, PSMD1, PEBP1, HSPB1, TNIP1, RPS13, ZFAND5, YWHAQ, COMMD1, COPS3, POLR1D, SMARCC2, MAP3K3, BIRC3, UBE2D2, HDAC2, CASP8, MCM7, PSMD7, YWHAG, NFKBIA, CAST, YWHAB, G3BP2, PSMD13, FBL, RELB, YWHAZ, SKP1, UBE2D3, PDCD2, HSP90AA1, HDAC1, KPNA2, RPL30, GTF2I, PFDN2
Colorectal cancer	CD9, EGFR, NGAL, CD81, STEAP, CD24, A33, CD66E, EPHA2, Ferritin, GPR30, GPR110, MMP9, OPN, p53, TMEM211, TROP2, TGM2, TIMP, EGFR, DR3, UNC93A, MUC17, EpCAM, MUC1, MUC2, TSG101, CD63, B7H3

Colorectal cancer	DR3, STEAP, epha2, TMEM211, unc93A, A33, CD24, NGAL, EpCam, MUC17, TROP2, TETS
Colorectal cancer	A33, AFP, ALIX, ALX4, ANCA, APC, ASCA, AURKA, AURKB, B7H3, BANK1, BCNP, BDNF, CA-19-9, CCSA-2, CCSA-3&4, CD10, CD24, CD44, CD63, CD66 CEA, CD66e CEA, CD81, CD9, CDA, C-Erb2, CRMP-2, CRP, CRTN, CXCL12, CYFRA21-1, DcR3, DLL4, DR3, EGFR, Epcam, EphA2, FASL, FRT, GAL3, GDF15, GPCR (GPR110), GPR30, GRO-1, HBD 1, HBD2, HNP1-3, IL-1B, IL8, IMP3, L1CAM, LAMN, MACC-1, MGC20553, MCP-1, M-CSF, MIC1, MIF, MMP7, MMP9, MS4A1, MUC1, MUC17, MUC2, Ncam, NGAL, NNMT, OPN, p53, PCSA, PDGFRB, PRL, PSMA, PSME3, Reg IV, SCRN1, Sept-9, SPARC, SPON2, SPR, SRVN, TFF3, TGM2, TIMP-1, TMEM211, TNF-alpha, TPA, TPS, Trail-R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, VEGFA
Colorectal cancer	miR 92, miR 21, miR 9, miR 491
Colorectal cancer	miR-127-3p, miR-92a, miR-486-3p, miR-378
Colorectal cancer	TMEM211, MUC1, CD24 and/or GPR110 (GPCR 110)
Colorectal cancer	hsa-miR-376c, hsa-miR-215, hsa-miR-652, hsa-miR-582-5p, hsa-miR-324-5p, hsa-miR-1296, hsa-miR-28-5p, hsa-miR-190, hsa-miR-590-5p, hsa-miR-202, hsa-miR-195
Colorectal cancer vesicle markers	A26C1A, A26C1B, A2M, ACAA2, ACE, ACOT7, ACP1, ACTA1, ACTA2, ACTB, ACTBL2, ACTBL3, ACTC1, ACTG1, ACTG2, ACTN1, ACTN2, ACTN4, ACTR3, ADAM10, ADSL, AGR2, AGR3, AGRN, AHCY, AHNAK, AKR1B10, ALB, ALDH16A1, ALDH1A1, ALDOA, ANXA1, ANXA11, ANXA2, ANXA2P2, ANXA4, ANXA5, ANXA6, AP2A1, AP2A2, APOA1, ARF1, ARF3, ARF4, ARF5, ARF6, ARHGDIA, ARPC3, ARPC5L, ARRDC1, ARVCF, ASCC3L1, ASNS, ATP1A1, ATP1A2, ATP1A3, ATP1B1, ATP4A, ATP5A1, ATP5B, ATP5I, ATP5L, ATP5O, ATP6AP2, B2M, BAIAP2, BAIAP2L1, BRI3BP, BSG, BUB3, C1orf58, C5orf32, CAD, CALM1, CALM2, CALM3, CAND1, CANX, CAPZA1, CBR1, CBR3, CCT2, CCT3, CCT4, CCT5, CCT6A, CCT7, CCT8, CD44, CD46, CD55, CD59, CD63, CD81, CD82, CD9, CDC42, CDH1, CDH17, CEACAM5, CFL1, CFL2, CHMP1A, CHMP2A, CHMP4B, CKB, CLDN3, CLDN4, CLDN7, CLIC1, CLIC4, CLSTN1, CLTC, CLTCL1, CLU, COL12A1, COPB1, COPB2, CORO1C, COX4I1, COX5B, CRYZ, CSPG4, CSRP1, CST3, CTNNA1, CTNNB1, CTNND1, CTTN, CYFIP1, DCD, DERA, DIP2A, DIP2B, DIP2C, DMBT1, DPEP1, DPP4, DYNC1H1, EDIL3, EEF1A1, EEF1A2, EEF1AL3, EEF1G, EEF2, EFNB1, EGFR, EHD1, EHD4, EIF3EIP, EIF3I, EIF4A1, EIF4A2, ENO1, ENO2, ENO3, EPHA2, EPHA5, EPHB1, EPHB2, EPHB3, EPHB4, EPPK1, ESD, EZR, F11R, F5, F7, FAM125A, FAM125B, FAM129B, FASLG, FASN, FAT, FCGBP, FER1L3, FKBP1A, FLNA, FLNB, FLOT1, FLOT2, G6PD, GAPDH, GARS, GCN1L1, GDI2, GK, GMDS, GNA13, GNAI2, GNAI3, GNAS, GNB1, GNB2, GNB2L1, GNB3, GNB4, GNG12, GOLGA7, GPA33, GPI, GPRC5A, GSN, GSTP1, H2AFJ, HADHA, hCG_1757335, HEPH, HIST1H2AB, HIST1H2AE, HIST1H2AJ, HIST1H2AK, HIST1H4A, HIST1H4B, HIST1H4C, HIST1H4D, HIST1H4E, HIST1H4F, HIST1H4H, HIST1H4I, HIST1H4J, HIST1H4K, HIST1H4L, HIST2H2AC, HIST2H4A, HIST2H4B, HIST3H2A, HIST4H4, HLA-A, HLA-A29.1, HLA-B, HLA-C, HLA-E, HLA-H, HNRNPA2B1, HNRNPH2, HPCAL1, HRAS, HSD17B4, HSP90AA1, HSP90AA2, HSP90AA4P, HSP90AB1, HSP90AB2P, HSP90AB3P, HSP90B1, HSPA1A, HSPA1B, HSPA1L, HSPA2, HSPA4, HSPA5, HSPA6, HSPA7, HSPA8, HSPA9, HSPD1, HSPE1, HSPG2, HYOU1, IDH1, IFITM1, IFITM2, IFITM3, IGH@, IGHG1, IGHG2, IGHG3, IGHG4, IGHM, IGHV4-31, IGK@, IGKC, IGKV1-5, IGKV2-24, IGKV3-20, IGSF3, IGSF8, IQGAP1, IQGAP2, ITGA2, ITGA3, ITGA6, ITGAV, ITGB1, ITGB4, JUP, KIAA0174, KIAA1199, KPNB1, KRAS, KRT1, KRT10, KRT13, KRT14, KRT15, KRT16, KRT17, KRT18, KRT19, KRT2, KRT20, KRT24, KRT25, KRT27, KRT28, KRT3, KRT4, KRT5, KRT6A, KRT6B, KRT6C, KRT7, KRT75, KRT76, KRT77, KRT79, KRT8, KRT9, LAMA5, LAMP1, LDHA, LDHB, LFNG, LGALS3, LGALS3BP, LGALS4, LIMA1, LIN7A, LIN7C, LOC100128936, LOC100130553, LOC100133382, LOC100133739, LOC284889, LOC388524, LOC388720, LOC442497, LOC653269, LRP4, LRPPRC, LRSAM1, LSR, LYZ, MAN1A1, MAP4K4, MARCKS, MARCKSL1, METRNL, MFGE8, MICA, MIF, MINK1, MITD1, MMP7, MOBKL1A, MSN, MTCH2, MUC13, MYADM, MYH10, MYH11, MYH14, MYH9, MYL6, MYL6B, MYO1C, MYO1D, NARS, NCALD, NCSTN, NEDD4, NEDD4L, NME1, NME2, NOTCH1, NQO1, NRAS, P4HB, PCBP1, PCNA, PCSK9, PDCD6, PDCD6IP, PDIA3, PDXK, PEBP1, PFN1, PGK1, PHB, PHB2, PKM2, PLEC1, PLEKHB2, PLSCR3, PLXNA1, PLXNB2, PPIA, PPIB, PPP2R1A, PRDX1, PRDX2, PRDX3, PRDX5, PRDX6, PRKAR2A, PRKDC, PRSS23, PSMA2, PSMC6, PSMD11, PSMD3, PSME3, PTGFRN, PTPRF, PYGB, QPCT, QSOX1, RAB10, RAB11A, RAB11B, RAB13, RAB14, RAB15, RAB1A, RAB1B, RAB2A, RAB33B, RAB35, RAB43,

	RAB4B, RAB5A, RAB5B, RAB5C, RAB6A, RAB6B, RAB7A, RAB8A, RAB8B, RAC1, RAC3, RALA, RALB, RAN, RANP1, RAP1A, RAP1B, RAP2A, RAP2B, RAP2C, RDX, REG4, RHOA, RHOC, RHOG, ROCK2, RP11-631M21.2, RPL10A, RPL12, RPL6, RPL8, RPLP0, RPLP0-like, RPLP1, RPLP2, RPN1, RPS13, RPS14, RPS15A, RPS16, RPS18, RPS20, RPS21, RPS27A, RPS3, RPS4X, RPS4Y1, RPS4Y2, RPS7, RPS8, RPSA, RPSAP15, RRAS, RRAS2, RUVBL1, RUVBL2, S100A10, S100A11, S100A14, S100A16, S100A6, S100P, SDC1, SDC4, SDCBP, SDCBP2, SERINC1, SERINC5, SERPINA1, SERPINF1, SETD4, SFN, SLC12A2, SLC12A7, SLC16A1, SLC1A5, SLC25A4, SLC25A5, SLC25A6, SLC29A1, SLC2A1, SLC3A2, SLC44A1, SLC7A5, SLC9A3R1, SMPDL3B, SNAP23, SND1, SOD1, SORT1, SPTAN1, SPTBN1, SSBP1, SSR4, TACSTD1, TAGLN2, TBCA, TCEB1, TCP1, TF, TFRC, THBS1, TJP2, TKT, TMED2, TNFSF10, TNIK, TNKS1BP1, TNPO3, TOLLIP, TOMM22, TPI1, TPM1, TRAP1, TSG101, TSPAN1, TSPAN14, TSPAN15, TSPAN6, TSPAN8, TSTA3, TTYH3, TUBA1A, TUBA1B, TUBA1C, TUBA3C, TUBA3D, TUBA3E, TUBA4A, TUBA4B, TUBA8, TUBB, TUBB2A, TUBB2B, TUBB2C, TUBB3, TUBB4, TUBB4Q, TUBB6, TUFM, TXN, UBA1, UBA52, UBB, UBC, UBE2N, UBE2V2, UGDH, UQCRC2, VAMP1, VAMP3, VAMP8, VCP, VIL1, VPS25, VPS28, VPS35, VPS36, VPS37B, VPS37C, WDR1, YWHAB, YWHAE, YWHAG, YWHAH, YWHAQ, YWHAZ
Colorectal Cancer	hsa-miR-16, hsa-miR-25, hsa-miR-125b, hsa-miR-451, hsa-miR-200c, hsa-miR-140-3p, hsa-miR-658, hsa-miR-370, hsa-miR-1296, hsa-miR-636, hsa-miR-502-5p
Breast cancer	miR-21, miR-155, miR-206, miR-122a, miR-210, miR-21, miR-155, miR-206, miR-122a, miR-210, let-7, miR-10b, miR-125a, miR-125b, miR-145, miR-143, miR-145, miR-1b
Breast cancer	GAS5
Breast cancer	ER, PR, HER2, MUC1, EGFR, KRAS, B-Raf, CYP2D6, hsp70, MART-1, TRP, HER2, hsp70, MART-1, TRP, HER2, ER, PR, Class III b-tubulin, VEGFA, ETV6-NTRK3, BCA-225, hsp70, MART1, ER, VEGFA, Class III b-tubulin, HER2/neu (e.g., for Her2+ breast cancer), GPR30, ErbB4 (JM) isoform, MPR8, MISIIR, CD9, EphA2, EGFR, B7H3, PSM, PCSA, CD63, STEAP, CD81, ICAM1, A33, DR3, CD66e, MFG-E8, TROP-2, Mammaglobin, Hepsin, NPGP/NPFF2, PSCA, 5T4, NGAL, EpCam, neurokinin receptor-1 (NK-1 or NK-1R), NK-2, Pai-1, CD45, CD10, HER2/ERBB2, AGTR1, NPY1R, MUC1, ESA, CD133, GPR30, BCA225, CD24, CA15.3 (MUC1 secreted), CA27.29 (MUC1 secreted), NMDAR1, NMDAR2, MAGEA, CTAG1B, NY-ESO-1, SPB, SPC, NSE, PGP9.5, progesterone receptor (PR) or its isoform (PR(A) or PR(B)), P2RX7, NDUFB7, NSE, GAL3, osteopontin, CHI3L1, IC3b, mesothelin, SPA, AQP5, GPCR, hCEA-CAM, PTP IA-2, CABYR, TMEM211, ADAM28, UNC93A, MUC17, MUC2, IL10R-beta, BCMA, HVEM/TNFRSF14, Trappin-2, Elafin, ST2/IL1 R4, TNFRF14, CEACAM1, TPA1, LAMP, WF, WH1000, PECAM, BSA, TNFR
Breast cancer	CD9, MIS Rii, ER, CD63, MUC1, HER3, STAT3, VEGFA, BCA, CA125, CD24, EPCAM, ERB B4
Breast cancer	CD10, NPGP/NPFF2, HER2/ERBB2, AGTR1, NPY1R, neurokinin receptor-1 (NK-1 or NK-1R), NK-2, MUC1, ESA, CD133, GPR30, BCA225, CD24, CA15.3 (MUC1 secreted), CA27.29 (MUC1 secreted), NMDAR1, NMDAR2, MAGEA, CTAG1B, NY-ESO-1
Breast cancer	SPB, SPC, NSE, PGP9.5, CD9, P2RX7, NDUFB7, NSE, GAL3, osteopontin, CHI3L1, EGFR, B7H3, IC3b, MUC1, mesothelin, SPA, PCSA, CD63, STEAP, AQP5, CD81, DR3, PSM, GPCR, EphA2, hCEA-CAM, PTP IA-2, CABYR, TMEM211, ADAM28, UNC93A, A33, CD24, CD10, NGAL, EpCam, MUC17, TROP-2, MUC2, IL10R-beta, BCMA, HVEM/TNFRSF14, Trappin-2 Elafin, ST2/IL1 R4, TNFRF14, CEACAM1, TPA1, LAMP, WF, WH1000, PECAM, BSA, TNFR
Breast cancer	BRCA, MUC-1, MUC 16, CD24, ErbB4, ErbB2 (HER2), ErbB3, HSP70, Mammaglobin, PR, PR(B), VEGFA
Breast cancer	CD9, HSP70, Gal3, MIS, EGFR, ER, ICB3, CD63, B7H4, MUC1, DLL4, CD81, ERB3, VEGF, BCA225, BRCA, CA125, CD174, CD24, ERB2, NGAL, GPR30, CYFRA21, CD31, cMET, MUC2, ERBB4
Breast cancer	CD9, EphA2, EGFR, B7H3, PSMA, PCSA, CD63, STEAP, CD81, STEAP1, ICAM1 (CD54), PSMA, A33, DR3, CD66e, MFG-8e, TMEM211, TROP-2, EGFR, Mammoglobin, Hepsin, NPGP/NPFF2, PSCA, 5T4, NGAL, NK-2, EpCam, NK-1R, PSMA, 5T4, PAI-1, CD45
Breast cancer	PGP9.5, CD9, HSP70, gal3-b2c10, EGFR, iC3b, PSMA, PCSA, CD63, MUC1, DLL4, CD81, B7-H3, HER 3 (ErbB3), MART-1, PSA, VEGF A, TIMP-1, GPCR GPR110, EphA2, MMP9, mmp7, TMEM211, UNC93a, BRCA, CA125 (MUC16), Mammaglobin, CD174 (Lewis y), CD66e CEA, CD24 c.sn3, C-erbB2, CD10, NGAL, epcam, CEA

	(carcinoembryonic Antigen), GPR30, CYFRA21-1, OPN, MUC17, hVEGFR2, MUC2, NCAM, ASPH, ErbB4, SPB, SPC, CD9, MS4A1, EphA2, MIS RII, HER2 (ErbB2), ER, PR (B), MRP8, CD63, B7H4, TGM2, CD81, DR3, STAT 3, MACC-1, TrKB, IL 6 Unc, OPG - 13, IL6R, EZH2, SCRN1, TWEAK, SERPINB3, CDAC1, BCA-225, DR3, A33, NPGP/NPFF2, TIMP1, BDNF, FRT, Ferritin heavy chain, seprase, p53, LDH, HSP, ost, p53, CXCL12, HAP, CRP, Gro-alpha, Tsg 101, GDF15
Breast cancer	CD9, HSP70, Gal3, MIS (RII), EGFR, ER, ICB3, CD63, B7H4, MUC1, CD81, ERB3, MART1, STAT3, VEGF, BCA225, BRCA, CA125, CD174, CD24, ERB2, NGAL, GPR30, CYFRA21, CD31, cMET, MUC2, ERB4, TMEM211
Breast Cancer	5T4 (trophoblast), ADAM10, AGER/RAGE, APC, APP (β-amyloid), ASPH (A-10), B7H3 (CD276), BACE1, BAI3, BRCA1, BDNF, BIRC2, C1GALT1, CA125 (MUC16), Calmodulin 1, CCL2 (MCP-1), CD9, CD10, CD127 (IL7R), CD174, CD24, CD44, CD63, CD81, CEA, CRMP-2, CXCR3, CXCR4, CXCR6, CYFRA 21, derlin 1, DLL4, DPP6, E-CAD, EpCaM, EphA2 (H-77), ER(1) ESR1 α, ER(2) ESR2 β, Erb B4, Erbb2, erb3 (Erb-B3), PA2G4, FRT (FLT1), Gal3, GPR30 (G-coupled ER1), HAP1, HER3, HSP-27, HSP70, IC3b, IL8, insig, junction plakoglobin, Keratin 15, KRAS, Mammaglobin, MART1, MCT2, MFGE8, MMP9, MRP8, Muc1, MUC17, MUC2, NCAM, NG2 (CSPG4), Ngal, NHE-3, NT5E (CD73), ODC1, OPG, OPN, p53, PARK7, PCSA, PGP9.5 (PARK5), PR(B), PSA, PSMA, RAGE, STXBP4, Survivin, TFF3 (secreted), TIMP1, TIMP2, TMEM211, TRAF4 (scaffolding), TRAIL-R2 (death Receptor 5), TrkB, Tsg 101, UNC93a, VEGF A, VEGFR2, YB-1, VEGFR1, GCDPF-15 (PIP), BigH3 (TGFb1-induced protein), 5HT2B (serotonin receptor 2B), BRCA2, BACE 1, CDH1-cadherin
Breast Cancer	AK5.2, ATP6V1B1, CRABP1
Breast Cancer	DST.3, GATA3, KRT81
Breast Cancer	AK5.2, ATP6V1B1, CRABP1, DST.3, ELF5, GATA3, KRT81, LALBA, OXTR, RASL10A, SERHL, TFAP2A.1, TFAP2A.3, TFAP2C, VTCN1
Breast Cancer	TRAP; Renal Cell Carcinoma; Filamin; 14.3.3, Pan; Prohibitin; c-fos; Ang-2; GSTmu; Ang-1; FHIT; Rad51; Inhibin alpha; Cadherin-P; 14.3.3 gamma; p18INK4c; P504S; XRCC2; Caspase 5; CREB-Binding Protein; Estrogen Receptor; IL17; Claudin 2; Keratin 8; GAPDH; CD1; Keratin, LMW; Gamma Glutamylcysteine Synthetase(GCS)/Glutamate-cysteine Ligase; a-B-Crystallin; Pax-5; MMP-19; APC; IL-3; Keratin 8 (phospho-specific Ser73); TGF-beta 2; ITK; Oct-2/; DJ-1; B7-H2; Plasma Cell Marker; Rad18; Estriol; Chk1; Prolactin Receptor; Laminin Receptor; Histone H1; CD45RO; GnRH Receptor; IP10/CRG2; Actin, Muscle Specific; S100; Dystrophin; Tubulin-a; CD3zeta; CDC37; GABA a Receptor 1; MMP-7 (Matrilysin); Heregulin; Caspase 3; CD56/NCAM-1; Gastrin 1; SREBP-1 (Sterol Regulatory Element Binding Protein-1); MLH1; PGP9.5; Factor VIII Related Antigen; ADP-ribosylation Factor (ARF-6); MHC II (HLA-DR) Ia; Survivin; CD23; G-CSF; CD2; Calretinin; Neuron Specific Enolase; CD165; Calponin; CD95 / Fas; Urocortin; Heat Shock Protein 27/hsp27; Topo II beta; Insulin Receptor; Keratin 5/8; sm; Actin, skeletal muscle; CA19-9; GluR1; GRIP1; CD79a mb-1; TdT; HRP; CD94; CCK-8; Thymidine Phosphorylase; CD57; Alkaline Phosphatase (AP); CD59 / MACIF / MIRL / Protectin; GLUT-1; alpha-1-antitrypsin; Presenillin; Mucin 3 (MUC3); pS2; 14-3-3 beta; MMP-13 (Collagenase-3); Fli-1; mGluR5; Mast Cell Chymase; Laminin B1/b1; Neurofilament (160kDa); CNPase; Amylin Peptide; Gai1; CD6; alpha-1-antichymotrypsin; E2F-2; MyoD1
Ductal carcinoma in situ (DCIS)	Laminin B1/b1; E2F-2; TdT; Apolipoprotein D; Granulocyte; Alkaline Phosphatase (AP); Heat Shock Protein 27/hsp27; CD95 / Fas; pS2; Estriol; GLUT-1; Fibronectin; CD6; CCK-8; sm; Factor VIII Related Antigen; CD57; Plasminogen; CD71 / Transferrin Receptor; Keratin 5/8; Thymidine Phosphorylase; CD45/T200/LCA; Epithelial Specific Antigen; Macrophage; CD10; MyoD1; Gai1; bcl-XL; hPL; Caspase 3; Actin, skeletal muscle; IP10/CRG2; GnRH Receptor; p35nck5a; ADP-ribosylation Factor (ARF-6); Cdk4 ; alpha-1-antitrypsin; IL17; Neuron Specific Enolase; CD56/NCAM-1; Prolactin Receptor; Cdk7; CD79a mb-1; Collagen IV; CD94; Myeloid Specific Marker; Keratin 10; Pax-5; IgM (m-Heavy Chain); CD45RO; CA19-9; Mucin 2; Glucagon; Mast Cell Chymase; MLH1; CD1; CNPase; Parkin; MHC II (HLA-DR) Ia; B7-H2; Chk1; Lambda Light Chain; MHC II (HLA-DP and DR); Myogenin; MMP-7 (Matrilysin); Topo II beta; CD53; Keratin 19; Rad18; Ret Oncoprotein; MHC II (HLA-DP); E3-binding protein (ARM1); Progesterone Receptor; Keratin 8; IgG; IgA; Tubulin; Insulin Receptor Substrate-1; Keratin 15; DR3; IL-3; Keratin 10/13; Cyclin D3; MHC I (HLA25 and HLA-Aw32); Calmodulin; Neurofilament (160kDa)
Ductal carcinoma in situ (DCIS) v. other Breast cancer	Macrophage; Fibronectin; Granulocyte; Keratin 19; Cyclin D3; CD45/T200/LCA; EGFR; Thrombospondin; CD81/TAPA-1; Ruv C; Plasminogen; Collagen IV; Laminin B1/b1; CD10; TdT; Filamin; bcl-XL; 14.3.3 gamma; 14.3.3, Pan; p170; Apolipoprotein D; CD71 /

	Transferrin Receptor; FHIT
Breast cancer	5HT2B, 5T4 (trophoblast), ACO2, ACSL3, ACTN4, ADAM10, AGR2, AGR3, ALCAM, ALDH6A1, ANGPTL4, ANO9, AP1G1, APC, APEX1, APLP2, APP (_-amyloid), ARCN1, ARHGAP35, ARL3, ASAH1, ASPH (A-10), ATP1B1, ATP1B3, ATP5I, ATP5O, ATXN1, B7H3, BACE1, BAI3, BAIAP2, BCA-200, BDNF, BigH3, BIRC2, BLVRB, BRCA, BST2, C1GALT1, C1GALT1C1, C20orf3, CA125, CACYBP, Calmodulin, CAPN1, CAPNS1, CCDC64B, CCL2 (MCP-1), CCT3, CD10(BD), CD127 (IL7R), CD174, CD24, CD44, CD80, CD86, CDH1, CDH5, CEA, CFL2, CHCHD3, CHMP3, CHRDL2, CIB1, CKAP4, COPA, COX5B, CRABP2, CRIP1, CRISPLD1, CRMP-2, CRTAP, CTLA4, CUL3, CXCR3, CXCR4, CXCR6, CYB5B, CYB5R1, CYCS, CYFRA 21, DBI, DDX23, DDX39B, derlin 1, DHCR7, DHX9, DLD, DLL4, DNAJB1, DPP6, DSTN, eCadherin, EEF1D, EEF2, EFTUD2, EIF4A2, EIF4A3, EpCaM, EphA2, ER(1) ESR1 _, ER(2) ESR2 _, Erb B4, Erb2, erb3 (Erb-B3?), ERLIN2, ESD, FARSA, FASN, FEN1, FKBP5, FLNB, FOXP3, FUS, Gal3, GCDPF-15, GCNT2, GNA12, GNG5, GNPTG, GPC6, GPD2, GPER (GPR30), GSPT1, H3F3B, H3F3C, HADH, HAP1, HER3, HIST1H1C, HIST1H2AB, HIST1H3A, HIST1H3C, HIST1H3D, HIST1H3E, HIST1H3F, HIST1H3G, HIST1H3H, HIST1H3I, HIST1H3J, HIST2H2BF, HIST2H3A, HIST2H3C, HIST2H3D, HIST3H3, HMGB1, HNRNPA2B1, HNRNPAB, HNRNPC, HNRNPD, HNRNPH2, HNRNPK, HNRNPL, HNRNPM, HNRNPU, HPS3, HSP-27, HSP70, HSP90B1, HSPA1A, HSPA2, HSPA9, HSPE1, IC3b, IDE, IDH3B, IDO1, IFI30, IL1RL2, IL7, IL8, ILF2, ILF3, IQCG, ISOC2, IST1, ITGA7, ITGB7, junction plakoglobin, Keratin 15, KRAS, KRT19, KRT2, KRT7, KRT8, KRT9, KTN1, LAMP1, LMNA, LMNB1, LNPEP, LRPPRC, LRRC57, Mammaglobin, MAN1A1, MAN1A2, MART1, MATR3, MBD5, MCT2, MDH2, MFGE8, MFGE8, MGP, MMP9, MRP8, MUC1, MUC17, MUC2, MYO5B, MYOF, NAPA, NCAM, NCL, NG2 (CSPG4), Ngal, NHE-3, NME2, NONO, NPM1, NQO1, NT5E (CD73), ODC1, OPG, OPN (SC), OS9, p53, PACSIN3, PAICS, PARK7, PARVA, PC, PCNA, PCSA, PD-1, PD-L1, PD-L2, PGP9.5, PHB, PHB2, PIK3C2B, PKP3, PPL, PR(B)?, PRDX2, PRKCB, PRKCD, PRKDC, PSA, PSAP, PSMA, PSMB7, PSMD2, PSME3, PYCARD, RAB1A, RAB3D, RAB7A, RAGE, RBL2, RNPEP, RPL14, RPL27, RPL36, RPS25, RPS4X, RPS4Y1, RPS4Y2, RUVBL2, SET, SHMT2, SLAIN1, SLC39A14, SLC9A3R2, SMARCA4, SNRPD2, SNRPD3, SNX33, SNX9, SPEN, SPR, SQSTM1, SSBP1, ST3GAL1, STXBP4, SUB1, SUCLG2, Survivin, SYT9, TFF3 (secreted), TGOLN2, THBS1, TIMP1, TIMP2, TMED10, TMED4, TMED9, TMEM211, TOM1, TRAF4 (scaffolding), TRAIL-R2, TRAP1, TrkB, Tsg 101, TXNDC16, U2AF2, UEVLD, UFC1, UNC93a, USP14, VASP, VCP, VDAC1, VEGFA, VEGFR1, VEGFR2, VPS37C, WIZ, XRCC5, XRCC6, YB-1, YWHAZ
Lung cancer	Pgrmc1 (progesterone receptor membrane component 1)/sigma-2 receptor, STEAP, EZH2
Lung cancer	Prohibitin, CD23, Amylin Peptide, HRP, Rad51, Pax-5, Oct-3/, GLUT-1, PSCA, Thrombospondin, FHIT, a-B-Crystallin, LewisA, Vacular Endothelial Growth Factor(VEGF), Hepatocyte Factor Homologue-4, Flt-4, GluR6/7, Prostate Apoptosis Response Protein-4, GluR1, Fli-1, Urocortin, S100A4, 14-3-3 beta, P504S, HDAC1, PGP9.5, DJ-1, COX2, MMP-19, Actin, skeletal muscle, Claudin 3, Cadherin-P, Collagen IX, p27Kip1, Cathepsin D, CD30 (Reed-Sternberg Cell Marker), Ubiquitin, FSH-b, TrxR2, CCK-8, Cyclin C, CD138, TGF-beta 2, Adrenocorticotrophic Hormone, PPAR-gamma, Bcl-6, GLUT-3, IGF-I, mRANKL, Fas-ligand, Filamin, Calretinin, O ct-1, Parathyroid Hormone, Claudin 5, Claudin 4, Raf-1 (Phospho-specific), CDC14A Phosphatase, Mitochondria, APC, Gastrin 1, Ku (p80), Gai1, XPA, Maltose Binding Protein, Melanoma (gp100), Phosphotyrosine, Amyloid A, CXCR4 / Fusin, Hepatic Nuclear Factor-3B, Caspase 1, HPV 16-E7, Axonal Growth Cones, Lck, Ornithine Decarboxylase, Gamma Glutamylcysteine Synthetase(GCS)/Glutamate-cysteine Ligase, ERCC1, Calmodulin, Caspase 7 (Mch 3), CD137 (4-1BB), Nitric Oxide Synthase, brain (bNOS), E2F-2, IL-10R, L-Plastin, CD18, Vimentin, CD50/ICAM-3, Superoxide Dismutase, Adenovirus Type 5 E1A, PHAS-I, Progesterone Receptor (phospho-specific) - Serine 294, MHC II (HLA-DQ), XPG, ER Ca+2 ATPase2, Laminin-s, E3-binding protein (ARM1), CD45RO, CD1, Cdk2, MMP-10 (Stromilysin-2), sm, Surfactant Protein B (Pro), Apolipoprotein D, CD46, Keratin 8 (phospho-specific Ser73), PCNA, PLAP, CD20, Syk, LH, Keratin 19, ADP-ribosylation Factor (ARF-6), Int-2 Oncoprotein, Luciferase, AIF (Apoptosis Inducing Factor), Grb2, bcl-X, CD16, Paxillin, MHC II (HLA-DP and DR), B-Cell, p21WAF1, MHC II (HLA-DR), Tyrosinase, E2F-1, Pds1, Calponin, Notch, CD26/DPP IV, SV40 Large T Antigen, Ku (p70/p80), Perforin, XPF, SIM Ag (SIMA-4D3), Cdk1/p34cdc2, Neuron Specific Enolase, b-2-Microglobulin, DNA Polymerase Beta, Thyroid Hormone Receptor, Human, Alkaline Phosphatase (AP), Plasma Cell Marker, Heat Shock Protein 70/hsp70, TRP75 / gp75, SRF

	(Serum Response Factor), Laminin B1/b1, Mast Cell Chymase, Caldesmon, CEA / CD66e, CD24, Retinoid X Receptor (hRXR), CD45/T200/LCA, Rabies Virus, Cytochrome c, DR3, bcl-XL, Fascin, CD71 / Transferrin Receptor
Lung Cancer	miR-497
Lung Cancer	Pgrmc1
Ovarian Cancer	CA-125, CA 19-9, c-reactive protein, CD95(also called Fas, Fas antigen, Fas receptor, FasR, TNFRSF6, APT1 or APO-1), FAP-1, miR-200 microRNAs, EGFR, EGFRvIII, apolipoprotein AI, apolipoprotein CIII, myoglobin, tenascin C, MSH6, claudin-3, claudin-4, caveolin-1, coagulation factor III, CD9, CD36, CD37, CD53, CD63, CD81, CD136, CD147, Hsp70, Hsp90, Rab13, Desmocollin-1, EMP-2, CK7, CK20, GCDF15, CD82, Rab-5b, Annexin V, MFG-E8, HLA-DR. MiR-200 microRNAs (miR-200a, miR-200b, miR-200c), miR-141, miR-429, JNK, Jun
Prostate Cancer v normal	AQP2, BMP5, C16orf86, CXCL13, DST, ERCC1, GNAO1, KLHL5, MAP4K1, NELL2, PENK, PGF, POU3F1, PRSS21, SCML1, SEMG1, SMARCD3, SNAI2, TAF1C, TNNT3
Prostate Cancer v Breast Cancer	ADRB2, ARG2, C22orf32, CYorf14, EIF1AY, FEV, KLK2, KLK4, LRRC26, MAOA, NLGN4Y, PNPLA7, PVRL3, SIM2, SLC30A4, SLC45A3, STX19, TRIM36, TRPM8
Prostate Cancer v Colorectal Cancer	ADRB2, BAIAP2L2, C19orf33, CDX1, CEACAM6, EEF1A2, ERN2, FAM110B, FOXA2, KLK2, KLK4, LOC389816, LRRC26, MIPOL1, SLC45A3, SPDEF, TRIM31, TRIM36, ZNF613
Prostate Cancer v Lung Cancer	ASTN2, CAB39L, CRIP1, FAM110B, FEV, GSTP1, KLK2, KLK4, LOC389816, LRRC26, MUC1, PNPLA7, SIM2, SLC45A3, SPDEF, TRIM36, TRPV6, ZNF613
Prostate Cancer	miRs-26a+b, miR-15, miR-16, miR-195, miR-497, miR-424, miR-206, miR-342-5p, miR-186, miR-1271, miR-600, miR-216b, miR-519 family, miR-203
Integrins	ITGA1 (CD49a, VLA1), ITGA2 (CD49b, VLA2), ITGA3 (CD49c, VLA3), ITGA4 (CD49d, VLA4), ITGA5 (CD49e, VLA5), ITGA6 (CD49f, VLA6), ITGA7 (FLJ25220), ITGA8, ITGA9 (RLC), ITGA10, ITGA11 (HsT18964), ITGAD (CD11D, FLJ39841), ITGAE (CD103, HUMINAE), ITGAL (CD11a, LFA1A), ITGAM (CD11b, MAC-1), ITGAV (CD51, VNRA, MSK8), ITGAW, ITGAX (CD11c), ITGB1 (CD29, FNRB, MSK12, MDF20), ITGB2 (CD18, LFA-1, MAC-1, MFI7), ITGB3 (CD61, GP3A, GPIIIa), ITGB4 (CD104), ITGB5 (FLJ26658), ITGB6, ITGB7, ITGB8
Glycoprotein	GpIa-IIa, GpIIb-IIIa, GpIIIb, GpIb, GpIX
Transcription factors	STAT3, EZH2, p53, MACC1, SPDEF, RUNX2, YB-1
Kinases	AURKA, AURKB
Disease Markers	6Ckine, Adiponectin, Adrenocorticotropic Hormone, Agouti-Related Protein, Aldose Reductase, Alpha-1-Antichymotrypsin, Alpha-1-Antitrypsin, Alpha-1-Microglobulin, Alpha-2-Macroglobulin, Alpha-Fetoprotein, Amphiregulin, Angiogenin, Angiopoietin-2, Angiotensin-Converting Enzyme, Angiotensinogen, Annexin A1, Apolipoprotein A-I, Apolipoprotein A-II, Apolipoprotein A-IV, Apolipoprotein B, Apolipoprotein C-I, Apolipoprotein C-III, Apolipoprotein D, Apolipoprotein E, Apolipoprotein H, Apolipoprotein(a), AXL Receptor Tyrosine Kinase, B cell-activating Factor, B Lymphocyte Chemoattractant, Bcl-2-like protein 2, Beta-2-Microglobulin, Betacellulin, Bone Morphogenetic Protein 6, Brain-Derived Neurotrophic Factor, Calbindin, Calcitonin, Cancer Antigen 125, Cancer Antigen 15-3, Cancer Antigen 19-9, Cancer Antigen 72-4, Carcinoembryonic Antigen, Cathepsin D, CD 40 antigen, CD40 Ligand, CD5 Antigen-like, Cellular Fibronectin, Chemokine CC-4, Chromogranin-A, Ciliary Neurotrophic Factor, Clusterin, Collagen IV, Complement C3, Complement Factor H, Connective Tissue Growth Factor, Cortisol, C-Peptide, C-Reactive Protein, Creatine Kinase-MB, Cystatin-C, Endoglin, Endostatin, Endothelin-1, EN-RAGE, Eotaxin-1, Eotaxin-2, Eotaxin-3, Epidermal Growth Factor, Epiregulin, Epithelial cell adhesion molecule, Epithelial-Derived Neutrophil-Activating Protein 78, Erythropoietin, E-Selectin, Ezrin, Factor VII, Fas Ligand, FASLG Receptor, Fatty Acid-Binding Protein (adipocyte), Fatty Acid-Binding Protein (heart), Fatty Acid-Binding Protein (liver), Ferritin, Fetuin-A, Fibrinogen, Fibroblast Growth Factor 4, Fibroblast Growth Factor basic, Fibulin-1C, Follicle-Stimulating Hormone, Galectin-3, Gelsolin, Glucagon, Glucagon-like Peptide 1, Glucose-6-phosphate Isomerase, Glutamate-Cysteine Ligase Regulatory subunit, Glutathione S-Transferase alpha, Glutathione S-Transferase Mu 1, Granulocyte Colony-Stimulating Factor, Granulocyte-Macrophage Colony-Stimulating Factor, Growth Hormone, Growth-Regulated alpha protein, Haptoglobin, HE4, Heat Shock Protein 60, Heparin-Binding EGF-Like Growth Factor, Hepatocyte Growth Factor, Hepatocyte Growth Factor Receptor, Hepsin, Human Chorionic Gonadotropin beta, Human Epidermal Growth Factor Receptor 2, Immunoglobulin A, Immunoglobulin E,

	Immunoglobulin M, Insulin, Insulin-like Growth Factor I, Insulin-like Growth Factor-Binding Protein 1, Insulin-like Growth Factor-Binding Protein 2, Insulin-like Growth Factor-Binding Protein 3, Insulin-like Growth Factor Binding Protein 4, Insulin-like Growth Factor Binding Protein 5, Insulin-like Growth Factor Binding Protein 6, Intercellular Adhesion Molecule 1, Interferon gamma, Interferon gamma Induced Protein 10, Interferon-inducible T-cell alpha chemoattractant, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 Receptor antagonist, Interleukin-2, Interleukin-2 Receptor alpha, Interleukin-3, Interleukin-4, Interleukin-5, Interleukin-6, Interleukin-6 Receptor, Interleukin-6 Receptor subunit beta, Interleukin-7, Interleukin-8, Interleukin-10, Interleukin-11, Interleukin-12 Subunit p40, Interleukin-12 Subunit p70, Interleukin-13, Interleukin-15, Interleukin-16, Interleukin-25, Kallikrein 5, Kallikrein-7, Kidney Injury Molecule-1, Lactoylglutathione lyase, Latency-Associated Peptide of Transforming Growth Factor beta 1, Lectin-Like Oxidized LDL Receptor 1, Leptin, Luteinizing Hormone, Lymphotactin, Macrophage Colony-Stimulating Factor 1, Macrophage Inflammatory Protein-1 alpha, Macrophage Inflammatory Protein-1 beta, Macrophage Inflammatory Protein-3 alpha, Macrophage inflammatory protein 3 beta, Macrophage Migration Inhibitory Factor, Macrophage-Derived Chemokine, Macrophage-Stimulating Protein, Malondialdehyde-Modified Low-Density Lipoprotein, Maspin, Matrix Metalloproteinase-1, Matrix Metalloproteinase-2, Matrix Metalloproteinase-3, Matrix Metalloproteinase-7, Matrix Metalloproteinase-9, Matrix Metalloproteinase-9, Matrix Metalloproteinase-10, Mesothelin, MHC class I chain-related protein A, Monocyte Chemotactic Protein 1, Monocyte Chemotactic Protein 2, Monocyte Chemotactic Protein 3, Monocyte Chemotactic Protein 4, Monokine Induced by Gamma Interferon, Myeloid Progenitor Inhibitory Factor 1, Myeloperoxidase, Myoglobin, Nerve Growth Factor beta, Neuronal Cell Adhesion Molecule, Neuron-Specific Enolase, Neuropilin-1, Neutrophil Gelatinase-Associated Lipocalin, NT-proBNP, Nucleoside diphosphate kinase B, Osteopontin, Osteoprotegerin, Pancreatic Polypeptide, Pepsinogen I, Peptide YY, Peroxiredoxin-4, Phosphoserine Aminotransferase, Placenta Growth Factor, Plasminogen Activator Inhibitor 1, Platelet-Derived Growth Factor BB, Pregnancy-Associated Plasma Protein A, Progesterone, Proinsulin (inc. Total or Intact), Prolactin, Prostasin, Prostate-Specific Antigen (inc. Free PSA), Prostatic Acid Phosphatase, Protein S100-A4, Protein S100-A6, Pulmonary and Activation-Regulated Chemokine, Receptor for advanced glycosylation end products, Receptor tyrosine-protein kinase erbB-3, Resistin, S100 calcium-binding protein B, Secretin, Serotransferrin, Serum Amyloid P-Component, Serum Glutamic Oxaloacetic Transaminase, Sex Hormone-Binding Globulin, Sortilin, Squamous Cell Carcinoma Antigen-1, Stem Cell Factor, Stromal cell-derived Factor-1, Superoxide Dismutase 1 (soluble), T Lymphocyte-Secreted Protein I-309, Tamm-Horsfall Urinary Glycoprotein, T-Cell-Specific Protein RANTES, Tenascin-C, Testosterone, Tetranectin, Thrombomodulin, Thrombopoietin, Thrombospondin-1, Thyroglobulin, Thyroid-Stimulating Hormone, Thyroxine-Binding Globulin, Tissue Factor, Tissue Inhibitor of Metalloproteinases 1, Tissue type Plasminogen activator, TNF-Related Apoptosis-Inducing Ligand Receptor 3, Transforming Growth Factor alpha, Transforming Growth Factor beta-3, Transthyretin, Trefoil Factor 3, Tumor Necrosis Factor alpha, Tumor Necrosis Factor beta, Tumor Necrosis Factor Receptor I, Tumor necrosis Factor Receptor 2, Tyrosine kinase with Ig and EGF homology domains 2, Urokinase-type Plasminogen Activator, Urokinase-type plasminogen activator Receptor, Vascular Cell Adhesion Molecule-1, Vascular Endothelial Growth Factor, Vascular endothelial growth Factor B, Vascular Endothelial Growth Factor C, Vascular endothelial growth Factor D, Vascular Endothelial Growth Factor Receptor 1, Vascular Endothelial Growth Factor Receptor 2, Vascular endothelial growth Factor Receptor 3, Vitamin K-Dependent Protein S, Vitronectin, von Willebrand Factor, YKL-40
Disease Markers	Adiponectin, Adrenocorticotropic Hormone, Agouti-Related Protein, Alpha-1-Antichymotrypsin, Alpha-1-Antitrypsin, Alpha-1-Microglobulin, Alpha-2-Macroglobulin, Alpha-Fetoprotein, Amphiregulin, Angiopoietin-2, Angiotensin-Converting Enzyme, Angiotensinogen, Apolipoprotein A-I, Apolipoprotein A-II, Apolipoprotein A-IV, Apolipoprotein B, Apolipoprotein C-I, Apolipoprotein C-III, Apolipoprotein D, Apolipoprotein E, Apolipoprotein H, Apolipoprotein(a), AXL Receptor Tyrosine Kinase, B Lymphocyte Chemoattractant, Beta-2-Microglobulin, Betacellulin, Bone Morphogenetic Protein 6, Brain-Derived Neurotrophic Factor, Calbindin, Calcitonin, Cancer Antigen 125, Cancer Antigen 19-9, Carcinoembryonic Antigen, CD 40 antigen, CD40 Ligand, CD5 Antigen-like, Chemokine CC-4, Chromogranin-A, Ciliary Neurotrophic Factor, Clusterin, Complement C3, Complement Factor H, Connective Tissue Growth Factor, Cortisol, C-Peptide, C-Reactive Protein, Creatine Kinase-MB, Cystatin-C, Endothelin-1, EN-RAGE,

	Eotaxin-1, Eotaxin-3, Epidermal Growth Factor, Epiregulin, Epithelial-Derived Neutrophil-Activating Protein 78, Erythropoietin, E-Selectin, Factor VII, Fas Ligand, FASLG Receptor, Fatty Acid-Binding Protein (heart), Ferritin, Fetuin-A, Fibrinogen, Fibroblast Growth Factor 4, Fibroblast Growth Factor basic, Follicle-Stimulating Hormone, Glucagon, Glucagon-like Peptide 1, Glutathione S-Transferase alpha, Granulocyte Colony-Stimulating Factor, Granulocyte-Macrophage Colony-Stimulating Factor, Growth Hormone, Growth-Regulated alpha protein, Haptoglobin, Heat Shock Protein 60, Heparin-Binding EGF-Like Growth Factor, Hepatocyte Growth Factor, Immunoglobulin A, Immunoglobulin E, Immunoglobulin M, Insulin, Insulin-like Growth Factor I, Insulin-like Growth Factor-Binding Protein 2, Intercellular Adhesion Molecule 1, Interferon gamma, Interferon gamma Induced Protein 10, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 Receptor antagonist, Interleukin-2, Interleukin-3, Interleukin-4, Interleukin-5, Interleukin-6, Interleukin-6 Receptor, Interleukin-7, Interleukin-8, Interleukin-10, Interleukin-11, Interleukin-12 Subunit p40, Interleukin-12 Subunit p70, Interleukin-13, Interleukin-15, Interleukin-16, Interleukin-25, Kidney Injury Molecule-1, Lectin-Like Oxidized LDL Receptor 1, Leptin, Luteinizing Hormone, Lymphotactin, Macrophage Colony-Stimulating Factor 1, Macrophage Inflammatory Protein-1 alpha, Macrophage Inflammatory Protein-1 beta, Macrophage Inflammatory Protein-3 alpha, Macrophage Migration Inhibitory Factor, Macrophage-Derived Chemokine, Malondialdehyde-Modified Low-Density Lipoprotein, Matrix Metalloproteinase-1, Matrix Metalloproteinase-2, Matrix Metalloproteinase-3, Matrix Metalloproteinase-7, Matrix Metalloproteinase-9, Matrix Metalloproteinase-9, Matrix Metalloproteinase-10, Monocyte Chemotactic Protein 1, Monocyte Chemotactic Protein 2, Monocyte Chemotactic Protein 3, Monocyte Chemotactic Protein 4, Monokine Induced by Gamma Interferon, Myeloid Progenitor Inhibitory Factor 1, Myeloperoxidase, Myoglobin, Nerve Growth Factor beta, Neuronal Cell Adhesion Molecule, Neutrophil Gelatinase-Associated Lipocalin, NT-proBNP, Osteopontin, Pancreatic Polypeptide, Peptide YY, Placenta Growth Factor, Plasminogen Activator Inhibitor 1, Platelet-Derived Growth Factor BB, Pregnancy-Associated Plasma Protein A, Progesterone, Proinsulin (inc. Intact or Total), Prolactin, Prostate-Specific Antigen (inc. Free PSA), Prostatic Acid Phosphatase, Pulmonary and Activation-Regulated Chemokine, Receptor for advanced glycosylation end products, Resistin, S100 calcium-binding protein B, Secretin, Serotransferrin, Serum Amyloid P-Component, Serum Glutamic Oxaloacetic Transaminase, Sex Hormone-Binding Globulin, Sortilin, Stem Cell Factor, Superoxide Dismutase 1 (soluble), T Lymphocyte-Secreted Protein I-309, Tamm-Horsfall Urinary Glycoprotein, T-Cell-Specific Protein RANTES, Tenascin-C, Testosterone, Thrombomodulin, Thrombopoietin, Thrombospondin-1, Thyroid-Stimulating Hormone, Thyroxine-Binding Globulin, Tissue Factor, Tissue Inhibitor of Metalloproteinases 1, TNF-Related Apoptosis-Inducing Ligand Receptor 3, Transforming Growth Factor alpha, Transforming Growth Factor beta-3, Transthyretin, Trefoil Factor 3, Tumor Necrosis Factor alpha, Tumor Necrosis Factor beta, Tumor necrosis Factor Receptor 2, Vascular Cell Adhesion Molecule-1, Vascular Endothelial Growth Factor, Vitamin K-Dependent Protein S, Vitronectin, von Willebrand Factor
Oncology	6Ckine, Aldose Reductase, Alpha-Fetoprotein, Amphiregulin, Angiogenin, Annexin A1, B cell-activating Factor, B Lymphocyte Chemoattractant, Bcl-2-like protein 2, Betacellulin, Cancer Antigen 125, Cancer Antigen 15-3, Cancer Antigen 19-9, Cancer Antigen 72-4, Carcinoembryonic Antigen, Cathepsin D, Cellular Fibronectin, Collagen IV, Endoglin, Endostatin, Eotaxin-2, Epidermal Growth Factor, Epiregulin, Epithelial cell adhesion molecule, Ezrin, Fatty Acid-Binding Protein (adipocyte), Fatty Acid-Binding Protein (liver), Fibroblast Growth Factor basic, Fibulin-1C, Galectin-3, Gelsolin, Glucose-6-phosphate Isomerase, Glutamate-Cysteine Ligase Regulatory subunit, Glutathione S-Transferase Mu 1, HE4, Heparin-Binding EGF-Like Growth Factor, Hepatocyte Growth Factor, Hepatocyte Growth Factor Receptor, Hepsin, Human Chorionic Gonadotropin beta, Human Epidermal Growth Factor Receptor 2, Insulin-like Growth Factor-Binding Protein 1, Insulin-like Growth Factor-Binding Protein 2, Insulin-like Growth Factor-Binding Protein 3, Insulin-like Growth Factor Binding Protein 4, Insulin-like Growth Factor Binding Protein 5, Insulin-like Growth Factor Binding Protein 6, Interferon gamma Induced Protein 10, Interferon-inducible T-cell alpha chemoattractant, Interleukin-2 Receptor alpha, Interleukin-6, Interleukin-6 Receptor subunit beta, Kallikrein 5, Kallikrein-7, Lactoylglutathione lyase, Latency-Associated Peptide of Transforming Growth Factor beta 1, Leptin, Macrophage inflammatory protein 3 beta, Macrophage Migration Inhibitory Factor, Macrophage-Stimulating Protein, Maspin, Matrix Metalloproteinase-2, Mesothelin, MHC class I chain-related protein A, Monocyte Chemotactic Protein 1, Monokine Induced by Gamma Interferon, Neuron-Specific Enolase,

	Neuropilin-1, Neutrophil Gelatinase-Associated Lipocalin, Nucleoside diphosphate kinase B, Osteopontin, Osteoprotegerin, Pepsinogen I, Peroxiredoxin-4, Phosphoserine Aminotransferase, Placenta Growth Factor, Platelet-Derived Growth Factor BB, Prostasin, Protein S100-A4, Protein S100-A6, Receptor tyrosine-protein kinase erbB-3, Squamous Cell Carcinoma Antigen-1, Stromal cell-derived Factor-1, Tenascin-C, Tetranectin, Thyroglobulin, Tissue type Plasminogen activator, Transforming Growth Factor alpha, Tumor Necrosis Factor Receptor I, Tyrosine kinase with Ig and EGF homology domains 2, Urokinase-type Plasminogen Activator, Urokinase-type plasminogen activator Receptor, Vascular Endothelial Growth Factor, Vascular endothelial growth Factor B, Vascular Endothelial Growth Factor C, Vascular endothelial growth Factor D, Vascular Endothelial Growth Factor Receptor 1, Vascular Endothelial Growth Factor Receptor 2, Vascular endothelial growth Factor Receptor 3, YKL-40
Disease	Adiponectin, Alpha-1-Antitrypsin, Alpha-2-Macroglobulin, Alpha-Fetoprotein, Apolipoprotein A-I, Apolipoprotein C-III, Apolipoprotein H, Apolipoprotein(a), Beta-2-Microglobulin, Brain-Derived Neurotrophic Factor, Calcitonin, Cancer Antigen 125, Cancer Antigen 19-9, Carcinoembryonic Antigen, CD 40 antigen, CD40 Ligand, Complement C3, C-Reactive Protein, Creatine Kinase-MB, Endothelin-1, EN-RAGE, Eotaxin-1, Epidermal Growth Factor, Epithelial-Derived Neutrophil-Activating Protein 78, Erythropoietin, Factor VII, Fatty Acid-Binding Protein (heart), Ferritin, Fibrinogen, Fibroblast Growth Factor basic, Granulocyte Colony-Stimulating Factor, Granulocyte-Macrophage Colony-Stimulating Factor, Growth Hormone, Haptoglobin, Immunoglobulin A, Immunoglobulin E, Immunoglobulin M, Insulin, Insulin-like Growth Factor I, Intercellular Adhesion Molecule 1, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 Receptor antagonist, Interleukin-2, Interleukin-3, Interleukin-4, Interleukin-5, Interleukin-6, Interleukin-7, Interleukin-8, Interleukin-10, Interleukin-12 Subunit p40, Interleukin-12 Subunit p70, Interleukin-13, Interleukin-15, Interleukin-16, Leptin, Lymphotactin, Macrophage Inflammatory Protein-1 alpha, Macrophage Inflammatory Protein-1 beta, Macrophage-Derived Chemokine, Matrix Metalloproteinase-2, Matrix Metalloproteinase-3, Matrix Metalloproteinase-9, Monocyte Chemotactic Protein 1, Myeloperoxidase, Myoglobin, Plasminogen Activator Inhibitor 1, Pregnancy-Associated Plasma Protein A, Prostate-Specific Antigen (inc. Free PSA), Prostatic Acid Phosphatase, Serum Amyloid P-Component, Serum Glutamic Oxaloacetic Transaminase, Sex Hormone-Binding Globulin, Stem Cell Factor, T-Cell-Specific Protein RANTES, Thrombopoietin, Thyroid-Stimulating Hormone, Thyroxine-Binding Globulin, Tissue Factor, Tissue Inhibitor of Metalloproteinases 1, Tumor Necrosis Factor alpha, Tumor Necrosis Factor beta, Tumor Necrosis Factor Receptor 2, Vascular Cell Adhesion Molecule-1, Vascular Endothelial Growth Factor, von Willebrand Factor
Neurological	Alpha-1-Antitrypsin, Apolipoprotein A-I, Apolipoprotein A-II, Apolipoprotein B, Apolipoprotein C-I, Apolipoprotein H, Beta-2-Microglobulin, Betacellulin, Brain-Derived Neurotrophic Factor, Calbindin, Cancer Antigen 125, Carcinoembryonic Antigen, CD5 Antigen-like, Complement C3, Connective Tissue Growth Factor, Cortisol, Endothelin-1, Epidermal Growth Factor Receptor, Ferritin, Fetuin-A, Follicle-Stimulating Hormone, Haptoglobin, Immunoglobulin A, Immunoglobulin M, Intercellular Adhesion Molecule 1, Interleukin-6 Receptor, Interleukin-7, Interleukin-10, Interleukin-11, Interleukin-17, Kidney Injury Molecule-1, Luteinizing Hormone, Macrophage-Derived Chemokine, Macrophage Migration Inhibitory Factor, Macrophage Inflammatory Protein-1 alpha, Matrix Metalloproteinase-2, Monocyte Chemotactic Protein 2, Peptide YY, Prolactin, Prostatic Acid Phosphatase, Serotransferrin, Serum Amyloid P-Component, Sortilin, Testosterone, Thrombopoietin, Thyroid-Stimulating Hormone, Tissue Inhibitor of Metalloproteinases 1, TNF-Related Apoptosis-Inducing Ligand Receptor 3, Tumor necrosis Factor Receptor 2, Vascular Endothelial Growth Factor, Vitronectin
Cardiovascular	Adiponectin, Apolipoprotein A-I, Apolipoprotein B, Apolipoprotein C-III, Apolipoprotein D, Apolipoprotein E, Apolipoprotein H, Apolipoprotein(a), Clusterin, C-Reactive Protein, Cystatin-C, EN-RAGE, E-Selectin, Fatty Acid-Binding Protein (heart), Ferritin, Fibrinogen, Haptoglobin, Immunoglobulin M, Intercellular Adhesion Molecule 1, Interleukin-6, Interleukin-8, Lectin-Like Oxidized LDL Receptor 1, Leptin, Macrophage Inflammatory Protein-1 alpha, Macrophage Inflammatory Protein-1 beta, Malondialdehyde-Modified Low-Density Lipoprotein, Matrix Metalloproteinase-1, Matrix Metalloproteinase-10, Matrix Metalloproteinase-2, Matrix Metalloproteinase-3, Matrix Metalloproteinase-7, Matrix Metalloproteinase-9, Monocyte Chemotactic Protein 1, Myeloperoxidase, Myoglobin, NT-proBNP, Osteopontin, Plasminogen Activator Inhibitor 1, P-Selectin, Receptor for advanced

	glycosylation end products, Serum Amyloid P-Component, Sex Hormone-Binding Globulin, T-Cell-Specific Protein RANTES, Thrombomodulin, Thyroxine-Binding Globulin, Tissue Inhibitor of Metalloproteinases 1, Tumor Necrosis Factor alpha, Tumor necrosis Factor Receptor 2, Vascular Cell Adhesion Molecule-1, von Willebrand Factor
Inflammatory	Alpha-1-Antitrypsin, Alpha-2-Macroglobulin, Beta-2-Microglobulin, Brain-Derived Neurotrophic Factor, Complement C3, C-Reactive Protein, Eotaxin-1, Factor VII, Ferritin, Fibrinogen, Granulocyte-Macrophage Colony-Stimulating Factor, Haptoglobin, Intercellular Adhesion Molecule 1, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 Receptor antagonist, Interleukin-2, Interleukin-3, Interleukin-4, Interleukin-5, Interleukin-6, Interleukin-7, Interleukin-8, Interleukin-10, Interleukin-12 Subunit p40, Interleukin-12 Subunit p70, Interleukin-15, Interleukin-17, Interleukin-23, Macrophage Inflammatory Protein-1 alpha, Macrophage Inflammatory Protein-1 beta, Matrix Metalloproteinase-2, Matrix Metalloproteinase-3, Matrix Metalloproteinase-9, Monocyte Chemotactic Protein 1, Stem Cell Factor, T-Cell-Specific Protein RANTES, Tissue Inhibitor of Metalloproteinases 1, Tumor Necrosis Factor alpha, Tumor Necrosis Factor beta, Tumor necrosis Factor Receptor 2, Vascular Cell Adhesion Molecule-1, Vascular Endothelial Growth Factor, Vitamin D-Binding Protein, von Willebrand Factor
Metabolic	Adiponectin, Adrenocorticotropic Hormone, Angiotensin-Converting Enzyme, Angiotensinogen, Complement C3 alpha des arg, Cortisol, Follicle-Stimulating Hormone, Galanin, Glucagon, Glucagon-like Peptide 1, Insulin, Insulin-like Growth Factor I, Leptin, Luteinizing Hormone, Pancreatic Polypeptide, Peptide YY, Progesterone, Prolactin, Resistin, Secretin, Testosterone
Kidney	Alpha-1-Microglobulin, Beta-2-Microglobulin, Calbindin, Clusterin, Connective Tissue Growth Factor, Creatinine, Cystatin-C, Glutathione S-Transferase alpha, Kidney Injury Molecule-1, Microalbumin, Neutrophil Gelatinase-Associated Lipocalin, Osteopontin, Tamm-Horsfall Urinary Glycoprotein, Tissue Inhibitor of Metalloproteinases 1, Trefoil Factor 3, Vascular Endothelial Growth Factor
Cytokines	Granulocyte-Macrophage Colony-Stimulating Factor, Interferon gamma, Interleukin-2, Interleukin-3, Interleukin-4, Interleukin-5, Interleukin-6, Interleukin-7, Interleukin-8, Interleukin-10, Macrophage Inflammatory Protein-1 alpha, Macrophage Inflammatory Protein-1 beta, Matrix Metalloproteinase-2, Monocyte Chemotactic Protein 1, Tumor Necrosis Factor alpha, Tumor Necrosis Factor beta, Brain-Derived Neurotrophic Factor, Eotaxin-1, Intercellular Adhesion Molecule 1, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 Receptor antagonist, Interleukin-12 Subunit p40, Interleukin-12 Subunit p70, Interleukin-15, Interleukin-17, Interleukin-23, Matrix Metalloproteinase-3, Stem Cell Factor, Vascular Endothelial Growth Factor
Protein	14.3.3 gamma, 14.3.3 (Pan), 14-3-3 beta, 6-Histidine, a-B-Crystallin, Acinus, Actin beta, Actin (Muscle Specific), Actin (Pan), Actin (skeletal muscle), Activin Receptor Type II, Adenovirus, Adenovirus Fiber, Adenovirus Type 2 E1A, Adenovirus Type 5 E1A, ADP-ribosylation Factor (ARF-6), Adrenocorticotrophic Hormone, AIF (Apoptosis Inducing Factor), Alkaline Phosphatase (AP), Alpha Fetoprotein (AFP), Alpha Lactalbumin, alpha-1-antichymotrypsin, alpha-1-antitrypsin, Amphiregulin, Amylin Peptide, Amyloid A, Amyloid A4 Protein Precursor, Amyloid Beta (APP), Androgen Receptor, Ang-1, Ang-2, APC, APC11, APC2, Apolipoprotein D, A-Raf, ARC, Ask1 / MAPKKK5, ATM, Axonal Growth Cones, b Galactosidase, b-2-Microglobulin, B7-H2, BAG-1, Bak, Bax, B-Cell, B-cell Linker Protein (BLNK), Bcl10 / CIPER / CLAP / mE10, bcl-2a, Bcl-6, bcl-X, bcl-XL, Bim (BOD), Biotin, Bonzo / STRL33 / TYMSTR, Bovine Serum Albumin, BRCA2 (aa 1323-1346), BrdU, Bromodeoxyuridine (BrdU), CA125, CA19-9, c-Abl, Cadherin (Pan), Cadherin-E, Cadherin-P, Calcitonin, Calcium Pump ATPase, Caldesmon, Calmodulin, Calponin, Calretinin, Casein, Caspase 1, Caspase 2, Caspase 3, Caspase 5, Caspase 6 (Mch 2), Caspase 7 (Mch 3), Caspase 8 (FLICE), Caspase 9, Catenin alpha, Catenin beta, Catenin gamma, Cathepsin D, CCK-8, CD1, CD10, CD100/Leukocyte Semaphorin, CD105, CD106 / VCAM, CD115/c-fms/CSF-1R/M-CSFR, CD137 (4-1BB), CD138, CD14, CD15, CD155/PVR (Polio Virus Receptor), CD16, CD165, CD18, CD1a, CD1b, CD2, CD20, CD21, CD23, CD231, CD24, CD25/IL-2 Receptor a, CD26/DPP IV, CD29, CD30 (Reed-Sternberg Cell Marker), CD32/Fcg Receptor II, CD35/CR1, CD36GPIIIb/GPIV, CD3zeta, CD4, CD40, CD42b, CD43, CD45/T200/LCA, CD45RB, CD45RO, CD46, CD5, CD50/ICAM-3, CD53, CD54/ICAM-1, CD56/NCAM-1, CD57, CD59 / MACIF / MIRL / Protectin, CD6, CD61 / Platelet Glycoprotein IIIA, CD63, CD68, CD71 / Transferrin Receptor, CD79a mb-1, CD79b, CD8, CD81/TAPA-1, CD84, CD9, CD94, CD95 / Fas, CD98, CDC14A Phosphatase, CDC25C, CDC34, CDC37, CDC47, CDC6, cdh1, Cdk1/p34cdc2, Cdk2, Cdk3, Cdk4, Cdk5,

	Cdk7, Cdk8, CDw17, CDw60, CDw75, CDw78, CEA / CD66e, c-erbB-2/HER-2/neu Ab-1 (21N), c-erbB-4/HER-4, c-fos, Chk1, Chorionic Gonadotropin beta (hCG-beta), Chromogranin A, CIDE-A, CIDE-B, CITED1, c-jun, Clathrin, claudin 11, Claudin 2, Claudin 3, Claudin 4, Claudin 5, CLAUDIN 7, Claudin-1, CNPase, Collagen II, Collagen IV, Collagen IX, Collagen VII, Connexin 43, COX2, CREB, CREB-Binding Protein, Cryptococcus neoformans, c-Src, Cullin-1 (CUL-1), Cullin-2 (CUL-2), Cullin-3 (CUL-3), CXCR4 / Fusin, Cyclin B1, Cyclin C, Cyclin D1, Cyclin D3, Cyclin E, Cyclin E2, Cystic Fibrosis Transmembrane Regulator, Cytochrome c, D4-GDI, Daxx, DcR1, DcR2 / TRAIL-R4 / TRUNDD, Desmin, DFF40 (DNA Fragmentation Factor 40) / CAD, DFF45 / ICAD, DJ-1, DNA Ligase I, DNA Polymerase Beta, DNA Polymerase Gamma, DNA Primase (p49), DNA Primase (p58), DNA-PKcs, DP-2, DR3, DR5, Dysferlin, Dystrophin, E2F-1, E2F-2, E2F-3, E2F-4, E2F-5, E3-binding protein (ARM1), EGFR, EMA/CA15-3/MUC-1, Endostatin, Epithelial Membrane Antigen (EMA / CA15-3 / MUC-1), Epithelial Specific Antigen, ER beta, ER Ca+2 ATPase2, ERCC1, Erk1, ERK2, Estradiol, Estriol, Estrogen Receptor, Exo1, Ezrin/p81/80K/Cytovillin, F.VIII/VWF, Factor VIII Related Antigen, FADD (FAS-Associated death domain-containing protein), Fascin, Fas-ligand, Ferritin, FGF-1, FGF-2, FHIT, Fibrillin-1, Fibronectin, Filaggrin, Filamin, FITC, Fli-1, FLIP, Flk-1 / KDR / VEGFR2, Flt-1 / VEGFR1, Flt-4, Fra2, FSH, FSH-b, Fyn, Ga0, Gab-1, GABA a Receptor 1, GAD65, Gai1, Gamma Glutamyl Transferase (gGT), Gamma Glutamylcysteine Synthetase(GCS)/Glutamate-cysteine Ligase, GAPDH, Gastrin 1, GCDFP-15, G-CSF, GFAP, Glicentin, Glucagon, Glucose-Regulated Protein 94, GluR 2/3, GluR1, GluR4, GluR6/7, GLUT-1, GLUT-3, Glycogen Synthase Kinase 3b (GSK3b), Glycophorin A, GM-CSF, GnRH Receptor, Golgi Complex, Granulocyte, Granzyme B, Grb2, Green Fluorescent Protein (GFP), GRIP1, Growth Hormone (hGH), GSK-3, GST, GSTmu, H.Pylori, HDAC1, HDJ-2/DNAJ, Heat Shock Factor 1, Heat Shock Factor 2, Heat Shock Protein 27/hsp27, Heat Shock Protein 60/hsp60, Heat Shock Protein 70/hsp70, Heat Shock Protein 75/hsp75, Heat Shock Protein 90a/hsp86, Heat Shock Protein 90b/hsp84, Helicobacter pylori, Heparan Sulfate Proteoglycan, Hepatic Nuclear Factor-3B, Hepatocyte, Hepatocyte Factor Homologue-4, Hepatocyte Growth Factor, Heregulin, HIF-1a, Histone H1, hPL, HPV 16, HPV 16-E7, HRP, Human Sodium Iodide Symporter (hNIS), I-FLICE / CASPER, IFN gamma, IgA, IGF-1R, IGF-I, IgG, IgM (m-Heavy Chain), I-Kappa-B Kinase b (IKKb), IL-1 alpha, IL-1 beta, IL-10, IL-10R, IL17, IL-2, IL-3, IL-30, IL-4, IL-5, IL-6, IL-8, Inhibin alpha, Insulin, Insulin Receptor, Insulin Receptor Substrate-1, Int-2 Oncoprotein, Integrin beta5, Interferon-a(II), Interferon-g, Involucrin, IP10/CRG2, IPO-38 Proliferation Marker, IRAK, ITK, JNK Activating kinase (JKK1), Kappa Light Chain, Keratin 10, Keratin 10/13, Keratin 14, Keratin 15, Keratin 16, Keratin 18, Keratin 19, Keratin 20, Keratin 5/6/18, Keratin 5/8, Keratin 8, Keratin 8 (phospho-specific Ser73), Keratin 8/18, Keratin (LMW), Keratin (Multi), Keratin (Pan), Ki67, Ku (p70/p80), Ku (p80), L1 Cell Adhesion Molecule, Lambda Light Chain, Laminin B1/b1, Laminin B2/g1, Laminin Receptor, Laminin-s, Lck, Lck (p56lck), Leukotriene (C4, D4, E4), LewisA, LewisB, LH, L-Plastin, LRP / MVP, Luciferase, Macrophage, MADD, MAGE-1, Maltose Binding Protein, MAP1B, MAP2a,b, MART-1/Melan-A, Mast Cell Chymase, Mcl-1, MCM2, MCM5, MDM2, Medroxyprogesterone Acetate (MPA), Mek1, Mek2, Mek6, Mekk-1, Melanoma (gp100), mGluR1, mGluR5, MGMT, MHC I (HLA25 and HLA-Aw32), MHC I (HLA-A), MHC I (HLA-A,B,C), MHC I (HLA-B), MHC II (HLA-DP and DR), MHC II (HLA-DP), MHC II (HLA-DQ), MHC II (HLA-DR), MHC II (HLA-DR) Ia, Microphthalmia, Milk Fat Globule Membrane Protein, Mitochondria, MLH1, MMP-1 (Collagenase-I), MMP-10 (Stromilysin-2), MMP-11 (Stromelysin-3), MMP-13 (Collagenase-3), MMP-14 / MT1-MMP, MMP-15 / MT2-MMP, MMP-16 / MT3-MMP, MMP-19, MMP-2 (72kDa Collagenase IV), MMP-23, MMP-7 (Matrilysin), MMP-9 (92kDa Collagenase IV), Moesin, mRANKL, Muc-1, Mucin 2, Mucin 3 (MUC3), Mucin 5AC, MyD88, Myelin / Oligodendrocyte, Myeloid Specific Marker, Myeloperoxidase, MyoD1, Myogenin, Myoglobin, Myosin Smooth Muscle Heavy Chain, Nck, Negative Control for Mouse IgG1, Negative Control for Mouse IgG2a, Negative Control for Mouse IgG3, Negative Control for Mouse IgM, Negative Control for Rabbit IgG, Neurofilament, Neurofilament (160kDa), Neurofilament (200kDa), Neurofilament (68kDa), Neuron Specific Enolase, Neutrophil Elastase, NF kappa B / p50, NF kappa B / p65 (Rel A), NGF-Receptor (p75NGFR), brain Nitric Oxide Synthase (bNOS), endothelial Nitric Oxide Synthase (eNOS), nm23, NOS-i, NOS-u, Notch, Nucleophosmin (NPM), NuMA, O ct-1, Oct-2/, Oct-3/, Ornithine Decarboxylase, Osteopontin, p130, p130cas, p14ARF, p15INK4b, p16INK4a, p170, p170 / MDR-1, p18INK4c, p19ARF, p19Skp1, p21WAF1, p27Kip1, p300 / CBP, p35nck5a, P504S, p53, p57Kip2 Ab-7, p63 (p53 Family Member), p73, p73a, p73a/b, p95VAV, Parathyroid Hormone, Parathyroid Hormone Receptor Type 1, Parkin, PARP,

	PARP (Poly ADP-Ribose Polymerase), Pax-5, Paxillin, PCNA, PCTAIRE2, PDGF, PDGFR alpha, PDGFR beta, Pds1, Perforin, PGP9.5, PHAS-I, PHAS-II, Phospho-Ser/Thr/Tyr, Phosphotyrosine, PLAP, Plasma Cell Marker, Plasminogen, PLC gamma 1, PMP-22, Pneumocystis jiroveci, PPAR-gamma, PR3 (Proteinase 3), Presenillin, Progesterone, Progesterone Receptor, Progesterone Receptor (phospho-specific) - Serine 190, Progesterone Receptor (phospho-specific) - Serine 294, Prohibitin, Prolactin, Prolactin Receptor, Prostate Apoptosis Response Protein-4, Prostate Specific Acid Phosphatase, Prostate Specific Antigen, pS2, PSCA, Rabies Virus, RAD1, Rad51, Raf1, Raf-1 (Phospho-specific), RAIDD, Ras, Rad18, Renal Cell Carcinoma, Ret Oncoprotein, Retinoblastoma, Retinoblastoma (Rb) (Phospho-specific Serine608), Retinoic Acid Receptor (b), Retinoid X Receptor (hRXR), Retinol Binding Protein, Rhodopsin (Opsin), ROC, RPA/p32, RPA/p70, Ruv A, Ruv B, Ruv C, S100, S100A4, S100A6, SHP-1, SIM Ag (SIMA-4D3), SIRP a1, sm, SODD (Silencer of Death Domain), Somatostatin Receptor-I, SRC1 (Steroid Receptor Coactivator-1) Ab-1, SREBP-1 (Sterol Regulatory Element Binding Protein-1), SRF (Serum Response Factor), Stat-1, Stat3, Stat5, Stat5a, Stat5b, Stat6, Streptavidin, Superoxide Dismutase, Surfactant Protein A, Surfactant Protein B, Surfactant Protein B (Pro), Survivin, SV40 Large T Antigen, Syk, Synaptophysin, Synuclein, Synuclein beta, Synuclein pan, TACE (TNF-alpha converting enzyme) / ADAM17, TAG-72, tau, TdT, Tenascin, Testosterone, TGF beta 3, TGF-beta 2, Thomsen-Friedenreich Antigen, Thrombospondin, Thymidine Phosphorylase, Thymidylate Synthase, Thymine Glycols, Thyroglobulin, Thyroid Hormone Receptor beta, Thyroid Hormone Receptor, Thyroid Stimulating Hormone (TSH), TID-1, TIMP-1, TIMP-2, TNF alpha, TNFa, TNR-R2, Topo II beta, Topoisomerase IIa, Toxoplasma Gondii, TR2, TRADD, Transforming Growth Factor a, Transglutaminase II, TRAP, Tropomyosin, TRP75 / gp75, TrxR2, TTF-1, Tubulin, Tubulin-a, Tubulin-b, Tyrosinase, Ubiquitin, UCP3, uPA, Urocortin, Vacular Endothelial Growth Factor(VEGF), Vimentin, Vinculin, Vitamin D Receptor (VDR), von Hippel-Lindau Protein, Wnt-1, Xanthine Oxidase, XPA, XPF, XPG, XRCC1, XRCC2, ZAP-70, Zip kinase
Known Cancer Genes	ABL1, ABL2, ACSL3, AF15Q14, AF1Q, AF3p21, AF5q31, AKAP9, AKT1, AKT2, ALDH2, ALK, ALO17, APC, ARHGEF12, ARHH, ARID1A, ARID2, ARNT, ASPSCR1, ASXL1, ATF1, ATIC, ATM, ATRX, BAP1, BCL10, BCL11A, BCL11B, BCL2, BCL3, BCL5, BCL6, BCL7A, BCL9, BCOR, BCR, BHD, BIRC3, BLM, BMPR1A, BRAF, BRCA1, BRCA2, BRD3, BRD4, BRIP1, BTG1, BUB1B, C12orf9, C15orf21, C15orf55, C16orf75, CANT1, CARD11, CARS, CBFA2T1, CBFA2T3, CBFB, CBL, CBLB, CBLC, CCNB1IP1, CCND1, CCND2, CCND3, CCNE1, CD273, CD274, CD74, CD79A, CD79B, CDH1, CDH11, CDK12, CDK4, CDK6, CDKN2A, CDKN2a(p14), CDKN2C, CDX2, CEBPA, CEP1, CHCHD7, CHEK2, CHIC2, CHN1, CIC, CIITA, CLTC, CLTCL1, CMKOR1, COL1A1, COPEB, COX6C, CREB1, CREB3L1, CREB3L2, CREBBP, CRLF2, CRTC3, CTNNB1, CYLD, D10S170, DAXX, DDB2, DDIT3, DDX10, DDX5, DDX6, DEK, DICER1, DNMT3A, DUX4, EBF1, EGFR, EIF4A2, ELF4, ELK4, ELKS, ELL, ELN, EML4, EP300, EPS15, ERBB2, ERCC2, ERCC3, ERCC4, ERCC5, ERG, ETV1, ETV4, ETV5, ETV6, EVI1, EWSR1, EXT1, EXT2, EZH2, FACL6, FAM22A, FAM22B, FAM46C, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FBXO11, FBXW7, FCGR2B, FEV, FGFR1, FGFR1OP, FGFR2, FGFR3, FH, FHIT, FIP1L1, FLI1, FLJ27352, FLT3, FNBP1, FOXL2, FOXO1A, FOXO3A, FOXP1, FSTL3, FUBP1, FUS, FVT1, GAS7, GATA1, GATA2, GATA3, GMPS, GNA11, GNAQ, GNAS, GOLGA5, GOPC, GPC3, GPHN, GRAF, HCMOGT-1, HEAB, HERPUD1, HEY1, HIP1, HIST1H4I, HLF, HLXB9, HMGA1, HMGA2, HNRNPA2B1, HOOK3, HOXA11, HOXA13, HOXA9, HOXC11, HOXC13, HOXD11, HOXD13, HRAS, HRPT2, HSPCA, HSPCB, IDH1, IDH2, IGH@, IGK@, IGL@, IKZF1, IL2, IL21R, IL6ST, IL7R, IRF4, IRTA1, ITK, JAK1, JAK2, JAK3, JAZF1, JUN, KDM5A, KDM5C, KDM6A, KDR, KIAA1549, KIT, KLK2, KRAS, KTN1, LAF4, LASP1, LCK, LCP1, LCX, LHFP, LIFR, LMO1, LMO2, LPP, LYL1, MADH4, MAF, MAFB, MALT1, MAML2, MAP2K4, MDM2, MDM4, MDS1, MDS2, MECT1, MED12, MEN1, MET, MITF, MKL1, MLF1, MLH1, MLL, MLL2, MLL3, MLLT1, MLLT10, MLLT2, MLLT3, MLLT4, MLLT6, MLLT7, MN1, MPL, MSF, MSH2, MSH6, MSI2, MSN, MTCP1, MUC1, MUTYH, MYB, MYC, MYCL1, MYCN, MYD88, MYH11, MYH9, MYST4, NACA, NBS1, NCOA1, NCOA2, NCOA4, NDRG1, NF1, NF2, NFE2L2, NFIB, NFKB2, NIN, NKX2-1, NONO, NOTCH1, NOTCH2, NPM1, NR4A3, NRAS, NSD1, NTRK1, NTRK3, NUMA1, NUP214, NUP98, OLIG2, OMD, P2RY8, PAFAH1B2, PALB2, PAX3, PAX5, PAX7, PAX8, PBRM1, PBX1, PCM1, PCSK7, PDE4DIP, PDGFB, PDGFRA, PDGFRB, PER1, PHOX2B, PICALM, PIK3CA, PIK3R1, PIM1, PLAG1, PML, PMS1, PMS2, PMX1, PNUTL1, POU2AF1, POU5F1, PPARG, PPP2R1A, PRCC, PRDM1,

	PRDM16, PRF1, PRKAR1A, PRO1073, PSIP2, PTCH, PTEN, PTPN11, RAB5EP, RAD51L1, RAF1, RALGDS, RANBP17, RAP1GDS1, RARA, RB1, RBM15, RECQL4, REL, RET, ROS1, RPL22, RPN1, RUNDC2A, RUNX1, RUNXBP2, SBDS, SDH5, SDHB, SDHC, SDHD, SEPT6, SET, SETD2, SF3B1, SFPQ, SFRS3, SH3GL1, SIL, SLC45A3, SMARCA4, SMARCB1, SMO, SOCS1, SOX2, SRGAP3, SRSF2, SS18, SS18L1, SSH3BP1, SSX1, SSX2, SSX4, STK11, STL, SUFU, SUZ12, SYK, TAF15, TAL1, TAL2, TCEA1, TCF1, TCF12, TCF3, TCF7L2, TCL1A, TCL6, TET2, TFE3, TFEB, TFG, TFPT, TFRC, THRAP3, TIF1, TLX1, TLX3, TMPRSS2, TNFAIP3, TNFRSF14, TNFRSF17, TNFRSF6, TOP1, TP53, TPM3, TPM4, TPR, TRA@, TRB@, TRD@, TRIM27, TRIM33, TRIP11, TSC1, TSC2, TSHR, TTL, U2AF1, USP6, VHL, VTI1A, WAS, WHSC1, WHSC1L1, WIF1, WRN, WT1, WTX, XPA, XPC, XPO1, YWHAE, ZNF145, ZNF198, ZNF278, ZNF331, ZNF384, ZNF521, ZNF9, ZRSR2
Known Cancer Genes	AR, androgen receptor; ARPC1A, actin-related protein complex 2/3 subunit A; AURKA, Aurora kinase A; BAG4, BCl-2 associated anthogene 4; BCl2l2, BCl-2 like 2; BIRC2, Baculovirus IAP repeat containing protein 2; CACNA1E, calcium channel voltage dependent alpha-1E subunit; CCNE1, cyclin E1; CDK4, cyclin dependent kinase 4; CHD1L, chromodomain helicase DNA binding domain 1-like; CKS1B, CDC28 protein kinase 1B; COPS3, COP9 subunit 3; DCUN1D1, DCN1 domain containing protein 1; DYRK2, dual specificity tyrosine phosphorylation regulated kinase 2; EEF1A2, eukaryotic elongation transcription factor 1 alpha 2; EGFR, epidermal growth factor receptor; FADD, Fas-associated via death domain; FGFR1, fibroblast growth factor receptor 1, GATA6, GATA binding protein 6; GPC5, glypican 5; GRB7, growth factor receptor bound protein 7; MAP3K5, mitogen activated protein kinase kinase kinase 5; MED29, mediator complex subunit 5; MITF, microphthalmia associated transcription factor; MTDH, metadherin; NCOA3, nuclear receptor coactivator 3; NKX2-1, NK2 homeobox 1; PAK1, p21/CDC42/RAC1-activated kinase 1; PAX9, paired box gene 9; PIK3CA, phosphatidylinositol-3 kinase catalytic a; PLA2G10, phopholipase A2, group X; PPM1D, protein phosphatase magnesium-dependent 1D; PTK6, protein tyrosine kinase 6; PRKCI, protein kinase C iota; RPS6KB1, ribosomal protein s6 kinase 70kDa; SKP2, s-phase kinase associated protein; SMURF1, sMAD specific E3 ubiquitin protein ligase 1; SHH, sonic hedgehog homologue; STARD3, sTAR-related lipid transfer domain containing protein 3; YWHAQ, tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, zeta isoform; ZNF217, zinc finger protein 217
Mitotic Related Cancer Genes	Aurora kinase A (AURKA); Aurora kinase B (AURKB); Baculoviral IAP repeat-containing 5, survivin (BIRC5); Budding uninhibited by benzimidazoles 1 homolog (BUB1); Budding uninhibited by benzimidazoles 1 homolog beta, BUBR1 (BUB1B); Budding uninhibited by benzimidazoles 3 homolog (BUB3); CDC28 protein kinase regulatory subunit 1B (CKS1B); CDC28 protein kinase regulatory subunit 2 (CKS2); Cell division cycle 2 (CDC2)/CDK1 Cell division cycle 20 homolog (CDC20); Cell division cycle-associated 8, borealin (CDCA8); Centromere protein F, mitosin (CENPF); Centrosomal protein 110 kDa (CEP110); Checkpoint with forkhead and ring finger domains (CHFR); Cyclin B1 (CCNB1); Cyclin B2 (CCNB2); Cytoskeleton-associated protein 5 (CKAP5/ch-TOG); Microtubule-associated protein RP/ EB family member 1. End-binding protein 1, EB1 (MAPRE1); Epithelial cell transforming sequence 2 oncogene (ECT2); Extra spindle poles like 1, separase (ESPL1); Forkhead box M1 (FOXM1); H2A histone family, member X (H2AFX); Kinesin family member 4A (KIF4A); Kinetochore-associated 1 (KNTC1/ROD); Kinetochore-associated 2; highly expressed in cancer 1 (KNTC2/HEC1); Large tumor suppressor, homolog 1 (LATS1); Large tumor suppressor, homolog 2 (LATS2); Mitotic arrest deficient-like 1; MAD1 (MAD1L1); Mitotic arrest deficient-like 2; MAD2 (MAD2L1); Mps1 protein kinase (TTK); Never in mitosis gene a-related kinase 2 (NEK2); Ninein, GSK3b interacting protein (NIN); Non-SMC condensin I complex, subunit D2 (NCAPD2/CNAP1); Non-SMC condensin I complex, subunit H (NACPH/CAPH); Nuclear mitotic apparatus protein 1 (NUMA1); Nucleophosmin (nucleolar phosphoprotein B23, numatrin); (NPM1); Nucleoporin (NUP98); Pericentriolar material 1 (PCM1); Pituitary tumor-transforming 1, securin (PTTG1); Polo-like kinase 1 (PLK1); Polo-like kinase 4 (PLK4/SAK); Protein (peptidylprolyl cis/trans isomerase) NIMA-interacting 1 (PIN1); Protein regulator of cytokinesis 1 (PRC1); RAD21 homolog (RAD21); Ras association (RalGDS/AF-6); domain family 1 (RASSF1); Stromal antigen 1 (STAG1); Synuclein-c, breast cancer-specific protein 1 (SNCG, BCSG1); Targeting protein for Xklp2 (TPX2); Transforming, acidic coiled-coil containing protein 3 (TACC3); Ubiquitin-conjugating enzyme E2C (UBE2C); Ubiquitin-conjugating enzyme E2I (UBE2I/UBC9); ZW10 interactor, (ZWINT); ZW10, kinetochore-associated homolog

	(ZW10); Zwilch, kinetochore-associated homolog (ZWILCH)
Ribonucleoprotein complexes	Argonaute family member, Ago1, Ago2, Ago3, Ago4, GW182 (TNRC6A), TNRC6B, TNRC6C, HNRNPA2B1, HNRPAB, ILF2, NCL (Nucleolin), NPM1 (Nucleophosmin), RPL10A, RPL5, RPLP1, RPS12, RPS19, SNRPG, TROVE2, apolipoprotein, apolipoprotein A, apo A-I, apo A-II, apo A-IV, apo A-V, apolipoprotein B, apo B48, apo B100, apolipoprotein C, apo C-I, apo C-II, apo C-III, apo C-IV, apolipoprotein D (ApoD), apolipoprotein E (ApoE), apolipoprotein H (ApoH), apolipoprotein L, APOL1, APOL2, APOL3, APOL4, APOL5, APOL6, APOLD1
Cytokine Receptors	4-1BB, ALCAM, B7-1, BCMA, CD14, CD30, CD40 Ligand, CEACAM-1, DR6, Dtk, Endoglin, ErbB3, E-Selectin, Fas, Flt-3L, GITR, HVEM, ICAM-3, IL-1 R4, IL-1 RI, IL-10 Rbeta, IL-17R, IL-2Rgamma, IL-21R, LIMPII, Lipocalin-2, L-Selectin, LYVE-1, MICA, MICB, NRG1-beta1, PDGF Rbeta, PECAM-1, RAGE, TIM-1, TRAIL R3, Trappin-2, uPAR, VCAM-1, XEDAR
Prostate and colorectal cancer vesicles	ErbB3, RAGE, Trail R3
Colorectal cancer vesicles	IL-1 alpha, CA125, Filamin, Amyloid A
Colorectal cancer v adenoma vesicles	Involucrin, CD57, Prohibitin, Thrombospondin, Laminin B1/b1, Filamin, 14.3.3 gamma, 14.3.3 Pan
Colorectal adenoma vesicles	Involucrin, Prohibitin, Laminin B1/b1, IL-3, Filamin, 14.3.3 gamma, 14.3.3 Pan, MMP-15 / MT2-MMP, hPL, Ubiquitin, and mRANKL
Brain cancer vesicles	Prohibitin, CD57, Filamin, CD18, b-2-Microglobulin, IL-2, IL-3, CD16, p170, Keratin 19, Pds1, Glicentin, SRF (Serum Response Factor), E3-binding protein (ARM1), Collagen II, SRC1 (Steroid Receptor Coactivator-1) Ab-1, Caldesmon, GFAP, TRP75 / gp75, alpha-1-antichymotrypsin, Hepatic Nuclear Factor-3B, PLAP, Tyrosinase, NF kappa B / p50, Melanoma (gp100), Cyclin E, 6-Histidine, Mucin 3 (MUC3), TdT, CD21, XPA, Superoxide Dismutase, Glycogen Synthase Kinase 3b (GSK3b), CD54/ICAM-1, Thrombospondin, Gai1, CD79a mb-1, IL-1 beta, Cytochrome c, RAD1, bcl-X, CD50/ICAM-3, Neurofilament, Alkaline Phosphatase (AP), ER Ca+2 ATPase2, PCNA, F.VIII/VWF, SV40 Large T Antigen, Paxillin, Fascin, CD165, GRIP1, Cdk8, Nucleophosmin (NPM), alpha-1-antitrypsin, CD32/Fcg Receptor II, Keratin 8 (phospho-specific Ser73), DR5, CD46, TID-1, MHC II (HLA-DQ), Plasma Cell Marker, DR3, Calmodulin, AIF (Apoptosis Inducing Factor), DNA Polymerase Beta, Vitamin D Receptor (VDR), Bcl10 / CIPER / CLAP / mE10, Neuron Specific Enolase, CXCR4 / Fusin, Neurofilament (68kDa), PDGFR, beta, Growth Hormone (hGH), Mast Cell Chymase, Ret Oncoprotein, and Phosphotyrosine
Melanoma vesicles	Caspase 5, Thrombospondin, Filamin, Ferritin, 14.3.3 gamma, 14.3.3 Pan, CD71 / Transferrin Receptor, and Prostate Apoptosis Response Protein-4
Head and neck cancer vesicles	14.3.3 Pan, Filamin, 14.3.3 gamma, CD71 / Transferrin Receptor, CD30, Cdk5, CD138, Thymidine Phosphorylase, Ruv 5, Thrombospondin, CD1, Von Hippel-Lindau Protein, CD46, Rad51, Ferritin, c-Abl, Actin, Muscle Specific, LewisB
Membrane proteins	carbonic anhydrase IX, B7, CCCL19, CCCL21, CSAp, HER-2/neu, BrE3, CD1, CD1a, CD2, CD3, CD4, CD5, CD8, CD11A, CD14, CD15, CD16, CD18, CD19, CD20, CD21, CD22, CD23, CD25, CD29, CD30, CD32b, CD33, CD37, CD38, CD40, CD40L, CD44, CD45, CD46, CD52, CD54, CD55, CD59, CD64, CD67, CD70, CD74, CD79a, CD80, CD83, CD95, CD126, CD133, CD138, CD147, CD154, CEACAM5, CEACAM-6, alpha-fetoprotein (AFP), VEGF, ED-B fibronectin, EGP-1, EGP-2, EGF receptor (ErbB1), ErbB2, ErbB3, Factor H, FHL-1, Flt-3, folate receptor, Ga 733,GROB, HMGB-1, hypoxia inducible factor (HIF), HM1.24, HER-2/neu, insulin-like growth factor (ILGF), IFN-γ, IFN-α, IL-β, IL-2R, IL-4R, IL-6R, IL-13R, IL-15R, IL-17R, IL-18R, IL-2, IL-6, IL-8, IL-12, IL-15, IL-17, IL-18, IL-25, IP-10, IGF-1R, Ia, HM1.24, gangliosides, HCG, HLA-DR, CD66a-d, MAGE, mCRP, MCP-1, MIP-1A, MIP-1B, macrophage migration-inhibitory factor (MIF), MUC1, MUC2, MUC3, MUC4, MUC5, placental growth factor (P1GF), PSA (prostate-specific antigen), PSMA, PSMA dimer, PAM4 antigen, NCA-95, NCA-90, A3, A33, Ep-CAM, KS-1, Le(y), mesothelin, S100, tenascin, TAC, Tn antigen, Thomas-Friedenreich antigens, tumor necrosis antigens, tumor angiogenesis antigens, TNF-α, TRAIL receptor (R1 and R2), VEGFR, RANTES, T101, cancer stem cell antigens, complement factors C3, C3a, C3b, C5a, C5
Cluster of Differentiation (CD) proteins	CD1, CD2, CD3, CD4, CD5, CD6, CD7, CD8, CD9, CD10, CD11a, CD11b, CD11c, CD12w, CD13, CD14, CD15, CD16, CDw17, CD18, CD19, CD20, CD21, CD22, CD23, CD24, CD25, CD26, CD27, CD28, CD29, CD30, CD31, CD32, CD33, CD34, CD35, CD36, CD37, CD38, CD39, CD40, CD41, CD42, CD43, CD44, CD45, CD46, CD47, CD48,

	CD49a, CD49b, CD49c, CD49d, CD49e, CD49f, CD53, CD54, CD55, CD56, CD57, CD58, CD59, CD61, CD62E, CD62L, CD62P, CD63, CD68, CD69, CD71, CD72, CD73, CD74, CD80, CD81, CD82, CD83, CD86, CD87, CD88, CD89, CD90, CD91, CD95, CD96, CD100, CD103, CD105, CD106, CD107, CD107a, CD107b, CD109, CD117, CD120, CD127, CD133, CD134, CD135, CD138, CD141, CD142, CD143, CD144, CD147, CD151, CD152, CD154, CD156, CD158, CD163, CD165, CD166, CD168, CD184, CDw186, CD195, CD197, CD209, CD202a, CD220, CD221, CD235a, CD271, CD303, CD304, CD309, CD326
Interleukin (IL) proteins	IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8 or CXCL8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, IL-21, IL-22, IL-23, IL-24, IL-25, IL-26, IL-27, IL-28, IL-29, IL-30, IL-31, IL-32, IL-33, IL-35, IL-36
IL receptors	CD121a/IL1R1, CD121b/IL1R2, CD25/IL2RA, CD122/IL2RB, CD132/IL2RG, CD123/IL3RA, CD131/IL3RB, CD124/IL4R, CD132/IL2RG, CD125/IL5RA, CD131/IL3RB, CD126/IL6RA, CD130/IR6RB, CD127/IL7RA, CD132/IL2RG, CXCR1/IL8RA, CXCR2/IL8RB/CD128, CD129/IL9R, CD210/IL10RA, CDW210B/IL10RB, IL11RA, CD212/IL12RB1, IR12RB2, IL13R, IL15RA, CD4, CDw217/IL17RA, IL17RB, CDw218a/IL18R1, IL20R, IL20R, IL21R, IL20R, IL23R, IL20R, LY6E, IL20R1, IL27RA, IL28R, IL31RA
Mucin (MUC) proteins	MUC1, MUC2, MUC3A, MUC3B, MUC4, MUC5AC, MUC5B, MUC6, MUC7, MUC8, MUC12, MUC13, MUC15, MUC16, MUC17, MUC19, and MUC20
MUC1 isoforms	mucin-1 isoform 2 precursor or mature form (NP_001018016.1), mucin-1 isoform 3 precursor or mature form (NP_001018017.1), mucin-1 isoform 5 precursor or mature form (NP_001037855.1), mucin-1 isoform 6 precursor or mature form (NP_001037856.1), mucin-1 isoform 7 precursor or mature form (NP_001037857.1), mucin-1 isoform 8 precursor or mature form (NP_001037858.1), mucin-1 isoform 9 precursor or mature form (NP_001191214.1), mucin-1 isoform 10 precursor or mature form (NP_001191215.1), mucin-1 isoform 11 precursor or mature form (NP_001191216.1), mucin-1 isoform 12 precursor or mature form (NP_001191217.1), mucin-1 isoform 13 precursor or mature form (NP_001191218.1), mucin-1 isoform 14 precursor or mature form (NP_001191219.1), mucin-1 isoform 15 precursor or mature form (NP_001191200.1), mucin-1 isoform 16 precursor or mature form (NP_001191201.1), mucin-1 isoform 17 precursor or mature form (NP_001191202.1), mucin-1 isoform 18 precursor or mature form (NP_001191203.1), mucin-1 isoform 19 precursor or mature form (NP_001191204.1), mucin-1 isoform 20 precursor or mature form (NP_001191205.1), mucin-1 isoform 21 precursor or mature form (NP_001191206.1), mucin-1 isoform 1 precursor or mature form (NP_002447.4), ENSP00000357380, ENSP00000357377, ENSP00000389098, ENSP00000357374, ENSP00000357381, ENSP00000339690, ENSP00000342814, ENSP00000357383, ENSP00000357375, ENSP00000338983, ENSP00000343482, ENSP00000406633, ENSP00000388172, ENSP00000357378, P15941-1, P15941-2, P15941-3, P15941-4, P15941-5, P15941-6, P15941-7, P15941-8, P15941-9, P15941-10, secreted isoform, membrane bound isoform, CA 27.29 (BR 27.29), CA 15-3, PAM4 reactive antigen, underglycosylated isoform, unglycosylated isoform, CanAg antigen
MUC1 interacting proteins	ABL1, SRC, CTNND1, ERBB2, GSK3B, JUP, PRKCD, APC, GALNT1, GALNT10, GALNT12, JUN, LCK, OSGEP, ZAP70, CTNNB1, EGFR, SOS1, ERBB3, ERBB4, GRB2, ESR1, GALNT2, GALNT4, LYN, TP53, C1GALT1, C1GALT1C1, GALNT3, GALNT6, GCNT1, GCNT4, MUC12, MUC13, MUC15, MUC17, MUC19, MUC2, MUC20, MUC3A, MUC4, MUC5B, MUC6, MUC7, MUCL1, ST3GAL1, ST3GAL3, ST3GAL4, ST6GALNAC2, B3GNT2, B3GNT3, B3GNT4, B3GNT5, B3GNT7, B4GALT5, GALNT11, GALNT13, GALNT14, GALNT5, GALNT8, GALNT9, ST3GAL2, ST6GAL1, ST6GALNAC4, GALNT15, MYOD1, SIGLEC1, IKBKB, TNFRSF1A, IKBKG, MUC1
Tumor markers	Alphafetoprotein (AFP), Carcinoembryonic antigen (CEA), CA-125, MUC-1, Epithelial tumor antigen (ETA), Tyrosinase, Melanoma-associated antigen (MAGE), p53
Tumor markers	Alpha fetoprotein (AFP), CA15-3, CA27-29, CA19-9, CA-125, Calretinin, Carcinoembryonic antigen, CD34, CD99, CD117, Chromogranin, Cytokeratin (various types), Desmin, Epithelial membrane protein (EMA), Factor VIII, CD31 FL1, Glial fibrillary acidic protein (GFAP), Gross cystic disease fluid protein (GCDFP-15), HMB-45, Human chorionic gonadotropin (hCG), immunoglobulin, inhibin, keratin (various types), PTPRC (CD45), lymphocyte marker (various types, MART-1 (Melan-A), Myo D1, muscle-specific actin (MSA), neurofilament, neuron-specific enolase (NSE), placental alkaline phosphatase (PLAP), prostate-specific antigen, S100 protein, smooth muscle actin (SMA), synaptophysin, thyroglobulin, thyroid transcription factor-1, Tumor M2-PK, vimentin

Cell adhesion molecule (CAMs)	Immunoglobulin superfamily CAMs (IgSF CAMs), N-CAM (Myelin protein zero), ICAM (1, 5), VCAM-1, PE-CAM, L1-CAM, Nectin (PVRL1, PVRL2, PVRL3), Integrins, LFA-1 (CD11a+CD18), Integrin alphaXbeta2 (CD11c+CD18), Macrophage-1 antigen (CD11b+CD18), VLA-4 (CD49d+CD29), Glycoprotein IIb/IIIa (ITGA2B+ITGB3), Cadherins, CDH1, CDH2, CDH3, Desmosomal, Desmoglein (DSG1, DSG2, DSG3, DSG4), Desmocollin (DSC1, DSC2, DSC3), Protocadherin, PCDH1, T-cadherin, CDH4, CDH5, CDH6, CDH8, CDH11, CDH12, CDH15, CDH16, CDH17, CDH9, CDH10, Selectins, E-selectin, L-selectin, P-selectin, Lymphocyte homing receptor: CD44, L-selectin, integrin (VLA-4, LFA-1), Carcinoembryonic antigen (CEA), CD22, CD24, CD44, CD146, CD164
Annexins	ANXA1; ANXA10; ANXA11; ANXA13; ANXA2; ANXA3; ANXA4; ANXA5; ANXA6; ANXA7; ANXA8; ANXA8L1; ANXA8L2; ANXA9
Cadherins ("calcium-dependent adhesion")	CDH1, CDH2, CDH12, CDH3, Deomoglein, DSG1, DSG2, DSG3, DSG4, Desmocollin, DSC1, DSC2, DSC3, Protocadherins, PCDH1, PCDH10, PCDH11x, PCDH11y, PCDH12, FAT, FAT2, FAT4, PCDH15, PCDH17, PCDH18, PCDH19; PCDH20; PCDH7, PCDH8, PCDH9, PCDHA1, PCDHA10, PCDHA11, PCDHA12, PCDHA13, PCDHA2, PCDHA3, PCDHA4, PCDHA5, PCDHA6, PCDHA7, PCDHA8, PCDHA9, PCDHAC1, PCDHAC2, PCDHB1, PCDHB10, PCDHB11, PCDHB12, PCDHB13, PCDHB14, PCDHB15, PCDHB16, PCDHB17, PCDHB18, PCDHB2, PCDHB3, PCDHB4, PCDHB5, PCDHB6, PCDHB7, PCDHB8, PCDHB9, PCDHGA1, PCDHGA10, PCDHGA11, PCDHGA12, PCDHGA2; PCDHGA3, PCDHGA4, PCDHGA5, PCDHGA6, PCDHGA7, PCDHGA8, PCDHGA9, PCDHGB1, PCDHGB2, PCDHGB3, PCDHGB4, PCDHGB5, PCDHGB6, PCDHGB7, PCDHGC3, PCDHGC4, PCDHGC5, CDH9 (cadherin 9, type 2 (T1-cadherin)), CDH10 (cadherin 10, type 2 (T2-cadherin)), CDH5 (VE-cadherin (vascular endothelial)), CDH6 (K-cadherin (kidney)), CDH7 (cadherin 7, type 2), CDH8 (cadherin 8, type 2), CDH11 (OB-cadherin (osteoblast)), CDH13 (T-cadherin - H-cadherin (heart)), CDH15 (M-cadherin (myotubule)), CDH16 (KSP-cadherin), CDH17 (LI cadherin (liver-intestine)), CDH18 (cadherin 18, type 2), CDH19 (cadherin 19, type 2), CDH20 (cadherin 20, type 2), CDH23 (cadherin 23, (neurosensory epithelium)), CDH10, CDH11, CDH13, CDH15, CDH16, CDH17, CDH18, CDH19, CDH20, CDH22, CDH23, CDH24, CDH26, CDH28, CDH4, CDH5, CDH6, CDH7, CDH8, CDH9, CELSR1, CELSR2, CELSR3, CLSTN1, CLSTN2, CLSTN3, DCHS1, DCHS2, LOC389118, PCLKC, RESDA1, RET
ECAD (CDH1) downregulators	SNAI1/SNAIL, ZFHX1B/SIP1, SNAI2/SLUG, TWIST1, DeltaEF1
ECAD upregulators	AML1, p300, HNF3
ECAD interacting proteins	ACADVL, ACTG1, ACTN1, ACTN4, ACTR3, ADAM10, ADAM9, AJAP1, ANAPC1, ANAPC11, ANAPC4, ANAPC7, ANK2, ANP32B, APC2, ARHGAP32, ARPC2, ARVCF, BOC, C1QBP, CA9, CASP3, CASP8, CAV1, CBLL1, CCNB1, CCND1, CCT6A, CDC16, CDC23, CDC26, CDC27, CDC42, CDH2, CDH3, CDK5R1, CDON, CDR2, CFTR, CREBBP, CSE1L, CSNK2A1, CTNNA1, CTNNB1, CTNND1, CTNND2, DNAJA1, DRG1, EGFR, EP300, ERBB2, ERBB2IP, ERG, EZR, FER, FGFR1, FOXM1, FRMD5, FYN, GBAS, GNA12, GNA13, GNB2L1, GSK3B, HDAC1, HDAC2, HSP90AA1, HSPA1A, HSPA1B, HSPD1, IGHA1, IQGAP1, IRS1, ITGAE, ITGB7, JUP, KIFC3, KLRG1, KRT1, KRT9, LIMA1, LMNA, MAD2L2, MAGI1, MAK, MDM2, MET, MYO6, MYO7A, NDRG1, NEDD9, NIPSNAP1, NKD2, PHLPP1, PIP5K1C, PKD1, PKP4, PLEKHA7, POLR2E, PPP1CA, PRKD1, PSEN1, PTPN1, PTPN14, PTPRF, PTPRM, PTPRQ, PTTG1, PVR, PVRL1, RAB8B, RRM2, SCRIB, SET, SIX1, SKI, SKP2, SRC, TACC3, TAS2R13, TGM2, TJP1, TK1, TNS3, TTK, UBC, USP9X, VCL, VEZT, XRCC5, YAP1, YES1, ZC3HC1
Epithelial-mesenchymal transition (EMT)	SERPINA3, ACTN1, AGR2, AKAP12, ALCAM, AP1M2, AXL, BSPRY, CCL2, CDH1, CDH2, CEP170, CLDN3, CLDN4, CNN3, CYP4X1, DNMT3A, DSG3, DSP, EFNB2, EHF, ELF3, ELF5, ERBB3, ETV5, FLRT3, FOSB, FOSLl, FOXCl, FX YD 5, GPDIL, HMGAl, HMGA2, HOPX, IFI16, IGFBP2, IHH, IKBIP, IL-11, IL-18, IL6, IL8, ITGA5, ITGB3, LAMBl, LCN2, MAP7, MB, MMP7, MMP9, MPZL2, MSLN, MTA3, MTSS1, OCLN, PCOLCE2, PECAM1, PLAUR, PLXNB1, PPL, PPP1R9A, RASSF8, SCNN1A, SERPINB2, SERPINE1, SFRP1, SH3YL1, SLC27A2, SMAD7, SNAI1, SNAI2, SPARC, SPDEF, SRPX, STAT5A, TBX2, TJP3, TMEM125, TMEM45B, TWIST1, VCAN, VIM, VWF, XBP1, YBX1, ZBTB10, ZEB1, ZEB2
Vesicle Associated	ALB, C3, A2M, TF, APOB, KRT1, KRT10, FGA, IGHG1, SERPINA1, FGB, KRT2, HP, IGHG3, IGHA1, SERPINA3, C4A, IGKC, C4B, CP, IGHM, FGG, KRT9, IGHG2, FN1, CFH, SERPINC1, C4A, APOA1, GC, Ig mu heavy chain disease protein, IGHG4, HPX,

	IGHA2, IGLC2, ITIH1, KNG1, ITIH4, ITIH2, AGT, PLG, APOA4, KRT14, CFB, IGLC1, ITIH4, ORM1, ITIH4, AHSG, A1BG, IGLL5, SERPING1, Ig kappa chain V-I region DEE, APOE, Ig kappa chain V-I region OU, ORM2, AFM, Ig heavy chain V-III region BUT, C4BPA, KRT6A, SERPINF1, APCS, APOH, CLU, KRT5, Ig heavy chain V-III region BRO, Ig heavy chain V-III region GAL, HRG, Ig heavy chain V-III region CAM, VTN, SERPIND1, TTR, PON1, Ig heavy chain V-III region TIL, C1QC, SERPINA7, Ig kappa chain V-I region CAR, Ig kappa chain V-IV region Len, AMBP, KRT13, Ig kappa chain V-III region SIE, SERPINF2, Ig heavy chain V-III region VH26, C5, F2, IGKV4-1, C7, Ig kappa chain V-I region EU, Ig kappa chain V-III region NG9 (Fragment), GSN, LPA, LYZ, Ig kappa chain V-III region HAH, Ig lambda chain V-III region LOI, SERPINA6, AZGP1, C1S, CFHR1, C9, HRNR, APOL1, C1QB, Ig kappa chain V-I region Ni, Ig heavy chain V-III region WEA, Ig kappa chain V-II region TEW, SERPINA4, DCD, LRG1, GSN, RBP4, SMC3, PRSS3, IGJ, C6, SEPP1, HBA1, Ig kappa chain V-III region CLL, ABCF1, APOD, SERPINA5, PDE4D, C2, C8A, C1R, CD5L, CFHR2, FLG2, HBB, CFI, Ig kappa chain V-II region MIL, Ig heavy chain V-II region NEWM, C8G, Ig lambda chain V-III region SH, PGLYRP2, SBSN, Ig lambda chain V-I region WAH, Ig lambda chain V-IV region Hil, SAA4, F10, MASP1, SHROOM3, F13A1, Ig lambda chain V region 4A, GIT2, KLKB1, ATRN, Ig heavy chain V-I region HG3, ITIH3, CDK10, APOA2, Ig heavy chain V-II region OU, Ig heavy chain V-I region V35, UTF1, MAP1B, PAPLN, Ig kappa chain V-I region Lay, RNF207, VPS13D, CRYGN, HMCN1, SLC27A6, FN1, VWF, C8B, LGALS3BP, HP, PROS1, ECM1, HPR, LBP, HABP2, FCN2, KRT77, APOM, Ig kappa chain V-I region WEA, GC, PLA2G7, Ig kappa chain V-I region Scw, CFP, APOM, MASP1, IGKV1-5, F12, SERPINA1, F13B, FCN2, PCYOX1, C4BPB, LCAT, KRT73, Ig heavy chain V-III region GA, Ig kappa chain V-III region VG (Fragment), MBL2, EEF2, MAP3K6, EPHA5, APOC4, CAMP, SERPINA10, FCGBP, PCSK9, CPB2, CFHR5, SAFB2, C2CD4C, F5, NUP153, XYLT1, EP300, BMP8A, N4BP2, KRT4, KRT16, Ig kappa chain V-III region B6, KRT86, KRT85, ANXA1, KRT78, SPRR2E, CLU, CRNN, ARHGEF17, SPRR3, FN1, ARHGAP30, ACTG2, SFTPA1, CDC5L, FN1, IGLC7, FLG, SERPINA1, Ig heavy chain V-III region TUR, JUP, DSP, KNG1, KPRP, LCE1C, Ig heavy chain V-II region ARH-77, Ig kappa chain V-III region POM, FBLN1, C1QA, FCN3, Ig lambda chain V-IV region Bau, Ig lambda chain V-VI region WLT, UPF3A, SERPINF2, XIRP2, CFB, SERPINA3, DSG1, TTN, LRRCC1, MYO15A, ANKRD28, Ig heavy chain V-III region HIL, KIT, DNMT1, PLXND1, Ig kappa chain V-I region Mev, IGHD, RCBTB1, BCO1, KRT6B, KRT13, Ig kappa chain V-II region RPMI 6410, Ig kappa chain V-IV region B17, ACTB, FN1, SARDH, GK, EMC4, MED30, PIGR, HSPB1, DSP, VEPH1, SNX27, LRRC53, SIGLEC16, F9, Ig heavy chain V-III region TRO, APOC3, TOP2A, FLYWCH1, ACTL10
Vesicle Associated	KRT6A, DSP, KRT6B, ACTB, FLG, IVL, SFN, KRT77, LMNA, KRT15, LGALS7, HSPA8, EPPK1, HSPA1A, DSG1, GSN, HIST1H2BK, EEF1A1, RPLP2, KRT74, YWHAB, PKP1, JUP, HNRNPA1, HSP90AA1, HIST1H2AH, GAPDH, HIST1H1E, HSPB1, CALML5, DCD, YWHAQ, VCP, AHNAK, SFPQ, PLEC, SERBP1, P4HB, PPL, Ig lambda chain V-IV region Hil, EIF3B, HSPA5, C3, TUBB4A, IGHG1, RPS3A, PPIA, SPTBN2, PDIA3, KRT80, DBNL, RPL29, RPL3, ANXA2P2, TPI1, RDX, H1F0, PGAM2, IGLC2, EVPL, ENO1, HNRNPA2B1, RPL7A, MYL6, ANXA1, TRIM29, RPS19, POF1B, RPL6, MORC2, RTN4, CA/CK E, LYZ, ZDBF2, IGKC, Ig heavy chain V-III region TIL, C4BPA, ACTB, LCE1C, IGHG3, SHOX2, KRT17, KRT77, KRT80, PIGR, KNG1, DSG1, DSP, SHROOM3, FGA, KPRP, DUSP27, LCE1C, SARDH, LYZ, SHISA5, HSP90AB1, EEF1A1, FGB, SHROOM3, IGLC2, KRT85, BMP8A, LCE2B, KRT6A, IGKC, S100A9, EEF1A1, C3, DCD, S100A8, LCE1C, ALB, IGLC2, S100A9, HSP90AB1, ACTB, KRT5, Ig kappa chain V-II region MIL, HRNR, IGHG1, HIST1H4A, DEFA1, LYZ, C3, SHROOM3, Ig kappa chain V-IV region STH (Fragment), Ig lambda chain V-I region HA, IGHA2, SARDH, H3F3C, LTF, TF
Vesicle Associated	C3, A2M, APOB, IGKC, C4A, C4B, FGB, ALB, CFH, IGHG1, FGA, FN1, PLG, IGHM, FGG, TF, C5, CP, IGHG2, IGLC2, Ig mu heavy chain disease protein, ITIH1, PZP, IGHG3, IGLL5, HP, C4BPA, ITIH2, IGHA1, KRT1, KRT10, APOE, Ig kappa chain V-I region DEE, AMBP, F2, C7, C6, ITIH4, CFB, IGHG4, APOH, APOA1, CD5L, C1R, HPR, Ig kappa chain V-I region Scw, IGHA2, CFHR1, KRT2, Ig kappa chain V-III region SIE, HRG, Ig heavy chain V-III region BRO, C1QB, GC, Ig heavy chain V-III region TIL, Ig kappa chain V-III region NG9 (Fragment), Ig heavy chain V-III region BUT, Ig heavy chain V-III region TUR, C9, SERPIND1, Ig kappa chain V-I region WEA, Ig kappa chain V-I region Ni, Ig kappa chain V-IV region Len, Ig kappa chain V-I region EU, Ig kappa chain V-II region TEW, Ig heavy chain V-III region GAL, KNG1, VTN, C8B, Ig lambda chain V-III region

	LOI, Ig heavy chain V-II region NEWM, APCS, KLKB1, CFI, PROS1, LPA, KRT9, SERPINA1, Ig lambda chain V-III region SH, C8A, Ig kappa chain V-III region B6, Ig lambda chain V-IV region Hil, Ig kappa chain V-III region CLL, C1S, FCN3, SERPINC1, Ig kappa chain V-I region Mev, IGHD, C1QC, HPX, C8G, IGKV1-5, Ig kappa chain V-I region Wes, Ig heavy chain V-III region WEA, A1BG, GSN, FBLN1, HBB, ITIH3, F12, SERPINA3, APOC3, Ig kappa chain V-I region BAN, Ig kappa chain V-III region VH (Fragment), F13B, IGKV4-1, SERPINF2, CLU, HIST1H1D, PON1, IGJ, Ig kappa chain V-III region POM, Ig heavy chain V-III region CAM, Ig heavy chain V-III region BUR, Ig kappa chain V-III region VG (Fragment), APOD, Ig lambda chain V-IV region MOL, Ig heavy chain V-III region GAR, FCGBP, APOM, F13A1, Ig heavy chain V-I region HG3, C1QA, Ig lambda chain V-VI region WLT, C2, C4BPB, CFP, SERPINA4, SAA4, SERPINF1, LGALS3BP, HABP2, RCBTB1, APOL1, KCNQ2, F9, Ig heavy chain V-III region TRO, Ig heavy chain V-III region HIL, Ig heavy chain V-II region OU, APOA2, F11, Ig lambda chain V-I region WAH, Ig lambda chain V region 4A, Ig kappa chain V-II region RPMI 6410, Ig kappa chain V-III region IARC/BL41, KRT5, IGLL1, Ig heavy chain V-I region V35, HBA1, ADIPOQ, PGLYRP2, UPF3A, BCO1, ARFGAP3, SARDH, SERPINA1, KNG1, Ig kappa chain V-I region Kue, Ig kappa chain V-I region Lay, Ig kappa chain V-I region OU, Ig kappa chain V-II region MIL, Ig heavy chain V-III region VH26, Ig heavy chain V-III region GA, FN1, TTR, SERPING1, APOA4, PRSS1, ANXA6, CFTR, LBP, FBLN1, SPAG17, PDLIM2, ARHGEF17, IGLC7, AGRN, AGT, RBP4, AHSG, Ig kappa chain V-III region GOL, SERPINA5, GSN, Ig kappa chain V-III region HAH, CFHR2, GIT2, INCENP
Vesicle Associated	MUC5B, FABP5, HPX, CP, SPRR2E, SPRR2D, PDE4D, GC, CPD, CD14, LAP3, AFM, FCN2, DMBT1, LIFR, SNX27, LCN1, ARFIP1, APOH, KLKB1, XP32, H2AFV, KRT75, KRT6C, KRT83, KRT76, KRT33B, KRT72, KRT31, KRT73, DSG1, LCE1C, LCE1A, CFB, CFH, SERPINA1, Ig kappa chain V-I region EU, Ig kappa chain V-II region MIL, Ig lambda chain V-IV region Bau, Ig heavy chain V-III region GAL, IGLC6, ACTG2

[00332] Examples of additional biomarkers that can be incorporated into the methods and compositions of the invention include without limitation those disclosed in International Patent Application Nos. PCT/US2012/042519 (WO 2012/174282), filed June 14, 2012 and PCT/US2012/050030 (WO 2013/022995), filed August 8, 2012.

[00333] In various embodiments of the invention, the biomarkers or biosignature used to detect or assess any of the conditions or diseases disclosed herein can comprise one or more biomarkers in one of several different categories of markers, wherein the categories include without limitation one or more of: 1) disease specific biomarkers; 2) cell- or tissue-specific biomarkers; 3) vesicle-specific markers (e.g., general vesicle biomarkers); 4. angiogenesis-specific biomarkers; and 5) immunomodulatory biomarkers. Examples of all such markers are disclosed herein and known to a person having ordinary skill in the art. Furthermore, a biomarker known in the art that is characterized to have a role in a particular disease or condition can be adapted for use as a target in compositions and methods of the invention. In further embodiments, such biomarkers that are associated with vesicles can be all vesicle surface markers, or a combination of vesicle surface markers and vesicle payload markers (i.e., molecules enclosed by a vesicle). The biomarkers assessed can be from a combination of sources. For example, a disease or disorder may be detected or characterized by assessing a combination of proteins, nucleic acids, vesicles, circulating biomarkers, biomarkers from a tissue sample, and the like. In addition, as noted herein, the biological sample assessed can be any biological fluid, or can comprise individual components present within such biological fluid (e.g., vesicles, nucleic acids, proteins, or complexes thereof).

[00334] EpCAM is a pan-epithelial differentiation antigen that is expressed on many tumor cells. It is intricately linked with the Cadherin-Catenin pathway and hence the fundamental WNT pathway responsible for intracellular signalling and polarity. It has been used as an immunotherapeutic target in the treatment of gastrointestinal, urological and other carcinomas. (Chaudry MA, Sales K, Ruf P, Lindhofer H, Winslet MC (April 2007). *Br. J. Cancer* **96** (7): 1013–9.). It is expressed in undifferentiated pluripotent stem cells. EpCAM is a member of a family that includes at least two type I membrane proteins and functions as a homotypic calcium-independent cell adhesion molecule. Mutations in this gene result in congenital tufting enteropathy. EpCAM has been observed on the surface of microvesicles derived from cancer cell of various lineages. EpCAM is used as an exemplary surface antigen in various examples herein. One of skill will appreciate that various embodiments and examples using EpCAM can be applied to other microvesicle surface antigens as well.

Oligonucleotide Probe Methods

[00335] Nucleic acid sequences fold into secondary and tertiary motifs particular to their nucleotide sequence. These motifs position the positive and negative charges on the nucleic acid sequences in locations that enable the sequences to bind to specific locations on target molecules, e.g., proteins and other amino acid sequences. These binding sequences are known in the field as aptamers. Due to the trillions of possible unique nucleotide sequences in even a relatively short stretch of nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 nucleotides), a large variety of motifs can be generated, resulting in aptamers for almost any desired protein or other target.

[00336] Aptamers are created by randomly generating oligonucleotides of a specific length, typically 20-80 base pairs long, e.g., 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 or 80 base pairs. These random oligonucleotides are then incubated with the protein target of interest. After several wash steps, the oligonucleotides that bind to the target are collected and amplified. The amplified aptamers are then added to the target and the process is repeated, often 15-20 times. A common version of this process known to those of skill in the art as the SELEX method.

[00337] The end result comprises one or more aptamer with high affinity to the target. The invention provides further processing of such resulting aptamers that can be use to provide desirable characteristics: 1) competitive binding assays to identify aptamers to a desired epitope; 2) motif analysis to identify high affinity binding aptamers *in silico*; and 3) microvesicle-based aptamer selection assays to identify aptamers that can be used to detect a particular disease. The methods are described in more detail below and further in the Examples.

[00338] The invention further contemplates aptamer sequences that are highly homologous to the sequences that are discovered by the methods of the invention. "High homology" typically refers to a homology of 40% or higher, preferably 60% or higher, more preferably 80% or higher, even more

preferably 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or higher between a polynucleotide sequence sequence and a reference sequence. In an embodiment, the reference sequence comprises the sequence of one or more aptamer provided herein. Percent homologies (also referred to as percent identity) are typically carried out between two optimally aligned sequences. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences and comparison can be conducted, e.g., using the algorithm in "Wilbur and Lipman, Proc Natl Acad Sci USA 80: 726-30 (1983)". Homology calculations can also be performed using BLAST, which can be found on the NCBI server at: www.ncbi.nlm.nih.gov/BLAST/ (Altschul S F, et al, Nucleic Acids Res. 1997; 25(17):3389-402; Altschul S F, et al, J Mol. Biol. 1990; 215(3):403-10). In the case of an isolated polynucleotide which is longer than or equivalent in length to the reference sequence, e.g., a sequence identified by the methods herein, the comparison is made with the full length of the reference sequence. Where the isolated polynucleotide is shorter than the reference sequence, e.g., shorter than a sequence identified by the methods herein, the comparison is made to a segment of the reference sequence of the same length (excluding any loop required by the homology calculation).

[00339] The invention further contemplates aptamer sequences that are functional fragments of the sequences that are discovered by the methods of the invention. In the context of an aptamer sequence, a "functional fragment" of the aptamer sequence may comprise a subsequence that binds to the same target as the full length sequence. In some instances, a candidate aptamer sequence is from a member of a library that contains a 5' leader sequences and/or a 3' tail sequence. Such leader sequences or tail sequences may serve to facilitate primer binding for amplification or capture, etc. In these embodiments, the functional fragment of the full length sequence may comprise the subsequence of the candidate aptamer sequence absent the leader and/or tail sequences.

[00340] ***Competitive Antibody Addition***

[00341] Known aptamer production methods may involve eluting all bound aptamers from the target sequence. In some cases, this is not sufficient to identify the desired aptamer sequence. For example, when trying to replace an antibody in an assay, it may be desirable to only collect aptamers that bind to the specific epitope of the antibody being replaced. The invention provides a method comprising addition of an antibody that is to be replaced to the aptamer/target reaction in order to allow for the selective collection of aptamers which bind to the antibody epitope. In an embodiment, the method comprises incubating a reaction mixture comprising randomly generated oligonucleotides with a target of interest, removing unbound aptamers from the reaction mixture that do not bind the target, adding an antibody to the reaction mixture that binds to that epitope of interest, and collecting the aptamers that are displaced by the antibody. The target can be a protein. See, e.g., **FIG. 1**, which illustrates the method for identifying an aptamer to a specific epitope of EpCam.

[00342] ***Motif Analysis***

[00343] In most aptamer experiments, multiple aptamer sequences are identified that bind to the target. These aptamers will have various binding affinities. It can be time consuming and laborious to generate

quantities of these many aptamers sufficient to assess the affinities of each. To identify large numbers of aptamers with the highest affinities without physically screening large subsets, the invention provides a method comprising the analysis of the two dimensional structure of one or more high affinity aptamers to the target of interest. In an embodiment, the method comprises screening the database for aptamers that have similar two-dimensional structures, or motifs, but not necessarily similar primary sequences. In an embodiment, the method comprises identifying a high affinity aptamer using traditional methods such as disclosed herein or known in the art (e.g. surface plasmon resonance binding assay, see **FIG. 5**), approximating the two-dimensional structure of the high affinity aptamer, and identifying aptamers from a pool of sequences that are predicted to have a similar two-dimensional structure to the high affinity aptamer. The method thereby provides a pool of candidates that also bind the target of interest. The two-dimensional structure of an oligo can be predicting using methods known in the art, e.g., via free energy (ΔG) calculations performed using a commercially available software program such as Vienna or mFold, for example as described in Mathews, D., Sabina, J., Zucker, M. & Turner, H. Expanded sequence dependence of thermodynamic parameters provides robust prediction of RNA secondary structure. J. Mol. Biol. 288, 911-940 (1999); Hofacker et al., Monatshefte f. Chemie 125: 167-188 (1994); and Hofacker, I. L. Vienna RNA secondary structure server. Nucleic Acids Res. 31, 3429-3431 (2003), the contents of which are incorporated herein by reference in their entirety. See **FIGs. 3A-3B**.The pool of sequences can be sequenced from a pool of randomly generated aptamer candidates using a high-throughput sequencing platform, such as the Ion Torrent platform from Life Technologies. Identifying aptamers from a pool of sequences that are predicted to have a similar two-dimensional structure to the high affinity aptamer may comprise loading the resulting sequences into the software program of choice to identify members of the pool of sequences with similar two-dimensional structures as the high affinity aptamer. The affinities of the pool of sequences can then be determined in situ, e.g., surface plasmon resonance binding assay or the like.

[00344] ***Aptamer Subtraction Methods***

[00345] In order to develop an assay to detect a disease, for example, cancer, one typically screens a large population of known biomarkers from normal and diseased patients in order to identify markers that correlate with disease. This process only works if discriminating markers are already described. In order to address this problem, the invention provides a method comprising subtracting out non-discriminating aptamers from a large pool of aptamers by incubating them initially with non-target microvesicles or cells. The non-target cells can be normal cells or microvesicles shed therefrom. The aptamers that did not bind to the normal microvesicles or cells are then incubated with diseased microvesicles or cells. The aptamers that bind to the diseased microvesicles or cells but that did not bind to the normal cells are then possible candidates for an assay to detect the disease. This process is independent of knowing the existence of a particular marker in the diseased sample.

[00346] Subtraction methods can be used to identify aptamers that preferentially recognize a desired population of targets. In an embodiment, the subtraction method is used to identify aptamers that

preferentially recognize target from a diseased target population over a control (e.g., normal or non-diseased) population. The diseased target population may be a population of vesicles from a diseased individual or individuals, whereas the control population comprises vesicles from a non-diseased individual or individuals. The disease can be a cancer or other disease disclosed herein or known in the art. Accordingly, the method provides aptamers that preferentially identify disease targets versus control targets.

[00347] Circulating microvesicles can be isolated from control samples, e.g., plasma from "normal" individuals that are absent a disease of interest, such as an absence of cancer. Vesicles in the sample are isolated using a method disclosed herein or as known in the art. For example, vesicles can be isolated from the plasma by one of the following methods: filtration, ultrafiltration, nanomembrane ultrafiltration, the ExoQuick reagent (System Biosciences, Inc., Mountain View, CA), centrifugation, ultracentrifugation, using a molecular crowding reagent (e.g., TEXIS from Life Technologies), polymer precipitation (e.g., polyethylene glycol (PEG)), affinity isolation, affinity selection, immunoprecipitation, chromatography, size exclusion, or a combination of any of these methods. The microvesicles isolated in each case will be a mixture of vesicle types and will be various sizes although ultracentrifugation methods may have more tendencies to produce exosomal-sized vesicles. Randomly generated oligonucleotide libraries (e.g., produced as described in the **Examples** herein) are incubated with the isolated normal vesicles. The aptamers that do not bind to these vesicles are isolated, e.g., by spinning down the vesicles and collecting the supernatant containing the non-binding aptamers. These non-binding aptamers are then contacted with vesicles isolated from diseased patients (e.g., using the same methods as described above) to allow the aptamers to recognize the disease vesicles. Next, aptamers that are bound to the diseased vesicles are collected. In an embodiment, the vesicles are isolated then lysed using a chaotropic agent (e.g., SDS or a similar detergent), and the aptamers are then captured by running the lysis mixture over an affinity column. The affinity column may comprise streptavidin beads in the case of biotin conjugated aptamer pools. The isolated aptamers are the amplified. The process can then then repeated, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 or more times.

[00348] In one aspect of the invention, an aptamer profile is identified that can be used to characterize a biological sample of interest. In an embodiment, a pool of randomly generated oligonucleotides, e.g., at least 10, 10^2, 10^3, 10^4, 10^5, 10^6, 10^7, 10^8, 10^9, 10^{10}, 10^{11}, 10^{12}, 10^{13}, 10^{14}, 10^{15}, 10^{16}, 10^{17}, 10^{18}, 10^{19} or at least 10^{20} oligonucleotides, is contacted with a biological component or target of interest from a control population. The oligonucleotides that do not bind the biological component or target of interest from the control population are isolated and then contacted with a biological component or target of interest from a test population. The oligonucleotides that bind the biological component or target of interest from the test population are retained. The retained oligonucleotides can be used to repeat the process by contacting the retained oligonucleotides with the biological component or target of interest from the control population, isolating the retained oligonucleotides that do not bind the biological component or target of interest from the control population, and again contacting these isolated oligonucleotides with the biological component

or target of interest from the test population and isolating the binding oligonucleotides. The "component" or "target" can be anything that is present in sample to which the oligonucleotides are capable of binding (e.g., polypeptides, peptide, nucleic acid molecules, carbodyhrates, lipids, etc.). The process can be repeated any number of desired iterations, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 or more times. The resulting oligonucleotides comprise aptamers that can differentially detect the test population versus the control. These aptamers provide an aptamer profile, which comprises a biosignature that is determined using one or more aptamer, e.g., a biosignature comprising a presense or level of the component or target which is detected using the one or more aptamer.

[00349] An exemplary process is illustrated in **FIG. 4**, which demonstrates the method to identify aptamer that preferentially recognize cancer vesicles using vesicles from normal (non-cancer) individuals as a control. In the figure, exosomes are exemplified but one of skill will appreciate that other microvesicles can be used in the same manner. The resulting aptamers can provide a profile that can differentially detect the cancer vesicles from the normal vesicles. One of skill will appreciate that the same steps can be used to derive an aptamer profile to characterize any disease or condition of interest.

[00350] In an embodiment, the invention provides an isolated polynucleotide that encodes a polypeptide, or a fragment thereof, identified by the methods above. The invention further provides an isolated polynucleotide having a nucleotide sequence that is at least 60% identical to the nucleotide sequence identified by the methods above. More preferably, the isolated nucleic acid molecule is at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more, identical to the nucleotide sequence identified by the methods above. In the case of an isolated polynucleotide which is longer than or equivalent in length to the reference sequence, e.g., a sequence identified by the methods above, the comparison is made with the full length of the reference sequence. Where the isolated polynucleotide is shorter than the reference sequence, e.g., shorter than a sequence identified by the methods above, the comparison is made to a segment of the reference sequence of the same length (excluding any loop required by the homology calculation).

[00351] In a related aspect, the invention provides a method of characterizing a biological phenotype using an aptamer profile. The aptamer profile can be determined using the method above. The aptamer profile can be determined for a test sample and compared to a control aptamer profile. The phenotype may be a disease or disorder such as a cancer. Characterizing the phenotype can include without limitation providing a diagnosis, prognosis, or theranosis. Thus, the aptamer profile can provide a diagnostic, prognostic and/or theranostic readout for the subject from whom the test sample is obtained.

[00352] In another embodiment, an aptamer profile is determined for a test sample by contacting a pool of aptamer molecules to the test sample, contacting the same pool of aptamers to a control sample, and identifying one or more aptamer molecules that differentially bind a component or target in the test sample but not in the control sample (or vice versa). A "component" or "target" as used in the context of the biological test sample or control sample can be anything that is present in sample to which the aptamers are capable of binding (e.g., polypeptides, peptide, nucleic acid molecules, carbodyhrates, lipids,

etc.). For example, if a sample is a plasma or serum sample, the aptamer molecules may bind a polypeptide biomarker that is solely expressed or differentially expressed (over- or underexpressed) in a disease state as compared to a non-diseased subject. Comparison of the aptamer profile in the test sample as compared to the control sample may be based on qualitative and quantitative measure of aptamer binding (e.g., binding versus no binding, or level of binding in test sample versus different level of binding in the reference control sample).

[00353] In an aspect, the invention provides a method of identifying a target-specific aptamer profile, comprising contacting a biological test sample with a pool of aptamer molecules, contacting the pool to a control biological sample, identifying one or more aptamers that bind to a component in said test sample but not to the control sample, thereby identifying an aptamer profile for said biological test sample. In an embodiment, a pool of aptamers is selected against a disease sample and compared to a reference sample, the aptamers in a subset that bind to a component(s) in the disease sample but not in the reference sample can be sequenced using conventional sequencing techniques to identify the subset that bind, thereby identifying an aptamer profile for the particular disease sample. In this way, the aptamer profile provides an individualized platform for detecting disease in other samples that are screened. Furthermore, by selecting an appropriate reference or control sample, the aptamer profile can provide a diagnostic, prognostic and/or theranostic readout for the subject from whom the test sample is obtained.

[00354] In a related aspect, the invention provides a method of selecting a pool of aptamers, comprising: (a) contacting a biological control sample with a pool of oligonucleotides; (b) isolating a first subset of the pool of oligonucleotides that do not bind the biological control sample; (c) contacting the biological test sample with the first subset of the pool of oligonucleotides; and (d) isolating a second subset of the pool of oligonucleotides that bind the biological test sample, thereby selecting the pool of aptamers. The pool of oligonucleotides may comprise any number of desired sequences, e.g., at least 10, 10^2, 10^3, 10^4, 10^5, 10^6, 10^7, 10^8, 10^9, 10^{10}, 10^{11}, 10^{12}, 10^{13}, 10^{14}, 10^{15}, 10^{16}, 10^{17}, 10^{18}, 10^{19} or at least 10^{20} oligonucleotides may be present in the starting pool. Steps (a)-(d) may be repeated to further hone the pool of aptamers. In an embodiment, these steps are repeated at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or at least 20 times.

[00355] As described herein, the biological test sample and biological control sample may comprise microvesicles. In an embodiment, the biological test sample and optionally biological control sample comprise a bodily fluid. The bodily fluid may comprise without limitation peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, Cowper's fluid, pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair, tears, cyst fluid, pleural fluid, peritoneal fluid, malignant fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates or other lavage fluids. Tthe biological test sample and optionally biological control may also comprise a tumor sample, e.g., cells from a tumor or tumor tissue.

In other embodiments, the biological test sample and optionally biological control sample comprise a cell culture medium. In embodiments, the biological test sample comprises a diseased sample and the biological control sample comprises a non-diseased sample. Accordingly, the pool of aptamers may be used to provide a diagnostic, prognostic and/or theranostic readout a disease.

[00356] As noted, the invention can be used to assess microvesicles. Microvesicles are powerful biomarkers because the vesicles provide one biological entity that comprises multiple pieces of information. For example as described, a vesicle can have multiple surface antigens, each of which provides complementary information. Consider a cancer marker and a tissue specific marker. If both markers are individually present in a sample, e.g., both are circulating proteins or nucleic acids, it may not be ascertainable whether the cancer marker and the tissue specific marker are derived from the same anatomical locale. However, if both the cancer marker and the tissue specific marker are surface antigens on a single microvesicle, the vesicle itself links the two markers and provides an indication of a disease (via the cancer marker) and origin of the disease (via the tissue specific marker). Furthermore, the vesicle can have any number of surface antigens and also payload that can be assessed. Accordingly, the invention provides a method for identifying binding agents comprising contacting a plurality of extracellular microvesicles with a randomly generated library of binding agents, identifying a subset of the library of binding agents that have an affinity to one or more components of the extracellular microvesicles. The binding agents may comprise aptamers, antibodies, and/or any other useful type of binding agent disclosed herein or known in the art.

[00357] In a related aspect, the invention provides a method for identifying a plurality of target ligands comprising, (a) contacting a reference microvesicle population with a plurality of ligands that are capable of binding one or more microvesicle surface markers, (b) isolating a plurality of reference ligands, wherein the plurality of reference ligands comprise a subset of the plurality of ligands that do not have an affinity for the reference microvesicle population; (c) contacting one or more test microvesicle with the plurality of reference ligands; and (d) identifying a subset of ligands from the plurality of reference ligands that form complexes with a surface marker on the one or more test microvesicle, thereby identifying the plurality of target ligands. The term "ligand" can refer a molecule, or a molecular group, that binds to another chemical entity to form a larger complex. Accordingly, a binding agent comprises a ligand. The plurality of ligands may comprise aptamers, antibodies and/or other useful binding agents described herein or known in the art.

[00358] The invention further provides kits comprising one or more reagent to carry out the methods above. In an embodiment, the one or more reagent comprises a library of potential binding agents that comprises one or more of an aptamer, antibody, and other useful binding agents described herein or known in the art.

[00359] ***Negative and Positive Aptamer Selection***

[00360] Aptamers can be used in various biological assays, including numerous types of assays which rely on a binding agent. For example, aptamers can be used instead of or along side antibodies in immune-

based assays. The invention provides an aptamer screening method that identifies aptamers that do not bind to any surfaces (substrates, tubes, filters, beads, other antigens, etc.) throughout the assay steps and bind specifically to an antigen of interest. The assay relies on negative selection to remove aptamers that bind non-target antigen components of the final assay. The negative selection is followed by positive selection to identify aptamers that bind the desired antigen.

[00361] In an aspect, the invention provides a method of identifying an aptamer specific to a target of interest, comprising (a) contacting a pool of candidate aptamers with one or more assay components, wherein the assay components do not comprise the target of interest; (b) recovering the members of the pool of candidate aptamers that do not bind to the one or more assay components in (a); (c) contacting the members of the pool of candidate aptamers recovered in (b) with the target of interest in the presence of one or more confounding target; and (d) recovering a candidate aptamer that binds to the target of interest in step (c), thereby identifying the aptamer specific to the target of interest. In the method, steps (a) and (b) provide negative selection to remove aptamers that bind non-target entities. Conversely, steps (c) and (d) provide positive selection by identifying aptamers that bind the target of interest but not other confounding targets, e.g., other antigens that may be present in a biological sample which comprises the target of interest. The pool of candidate aptamers may comprise at least 10, 10^2, 10^3, 10^4, 10^5, 10^6, 10^7, 10^8, 10^9, 10^{10}, 10^{11}, 10^{12}, 10^{13}, 10^{14}, 10^{15}, 10^{16}, 10^{17}, 10^{18}, 10^{19} or at least 10^{20} nucleic acid sequences. One illustrative approach for performing the method is provided in **Example 7**.

[00362] In some embodiments, steps (a)-(b) are optional. In other embodiments, steps (a)-(b) are repeated at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or at least 20 times before positive selection in step (c) is performed. The positive selection can also be performed in multiple rounds. Steps (c)-(d) can be repeated at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or at least 20 times before identifying the aptamer specific to the target of interest. Multiple rounds may provide improved stringency of selection.

[00363] In some embodiments, the one or more assay components contacted with the aptamer pool during negative selection comprise one or more of a substrate, a bead, a planar array, a column, a tube, a well, or a filter. One of skill will appreciate that the assay components can include any substance that may be part of a biological assay.

[00364] The target of interest can be any appropriate entity that can be detected when recognized by an aptamer. In an embodiment, the target of interest comprises a protein or polypeptide. As used herein, "protein," "polypeptide" and "peptide" are used interchangeably unless stated otherwise. The target of interest can be a nucleic acid, including DNA, RNA, and various subspecies of any thereof as disclosed herein or known in the art. The target of interest can comprise a lipid. The target of interest can comprise a carbohydrate. The target of interest can also be a complex, e.g., a complex comprising protein, nucleic acids, lipids and/or carbohydrates. In some embodiments, the target of interest comprises a microvesicle. In such cases, the aptamer can be a binding agent to a microvesicle surface antigen, e.g., a protein. General microvesicle surface antigens include tetraspanin, CD9, CD63, CD81, CD63, CD9, CD81, CD82,

CD37, CD53, Rab-5b, Annexin V, and MFG-E8. Additional general microvesicle surface antigens are provided in **Table 3** herein.

[00365] The microvesicle surface antigen can also be a biomarker of a disease or disorder. In such cases, the aptamer may be used to provide a diagnosis, prognosis or theranosis of the disease or disorder. For example, the one or more protein may comprise one or more of PSMA, PCSA, B7H3, EpCam, ADAM-10, BCNP, EGFR, IL1B, KLK2, MMP7, p53, PBP, SERPINB3, SPDEF, SSX2, and SSX4. These markers can be used detect a prostate cancer. Additional microvesicle surface antigens are provided in **Tables 3-4** herein.

[00366] The one or more confounding target can be an antigen other than the target of interest. For example, a confounding target can be another entity that may be present in a sample to be assayed. As a non-limiting example, consider that the sample to be assessed is a plasma sample from an individual. The target of interest may be a protein, e.g., a microvesicle surface antigen, which is present in the sample. In this case, a confounding target could be selected from any other antigen that is likely to be present in the plasma sample. Accordingly, the positive selection should provide candidate aptamers that recognize the target of interest but have minimal, if any, interactions with the confounding targets. In some embodiments, the target of interest and the one or more confounding target comprise the same type of biological entity, e.g., all protein, all nucleic acid, all carbohydrate, or all lipids. As a non-limiting example, the target of interest can be a protein selected from the group consisting of SSX4, SSX2, PBP, KLK2, SPDEF, and EpCAM, and the one or more confounding target comprises the other members of this group. In other embodiments, the target of interest and the one or more confounding target comprise different types of biological entities, e.g., any combination of protein, nucleic acid, carbohydrate, and lipids. The one or more confounding targets may also comprise different types of biological entities, e.g., any combination of protein, nucleic acid, carbohydrate, and lipids.

[00367] In an embodiment, the invention provides an isolated polynucleotide, or a fragment thereof, identified by the methods above. The invention further provides an isolated polynucleotide having a nucleotide sequence that is at least 60% identical to the nucleotide sequence identified by the methods above. More preferably, the isolated nucleic acid molecule is at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more, identical to the nucleotide sequence identified by the methods above. In the case of an isolated polynucleotide which is longer than or equivalent in length to the reference sequence, e.g., a sequence identified by the methods above, the comparison is made with the full length of the reference sequence. Where the isolated polynucleotide is shorter than the reference sequence, e.g., shorter than a sequence identified by the methods above, the comparison is made to a segment of the reference sequence of the same length (excluding any loop required by the homology calculation).

[00368] In a related aspect, the invention provides a method of selecting a group of aptamers, comprising: (a) contacting a pool of aptamers to a population of microvesicles from a first sample; (b) enriching a subpool of aptamers that show affinity to the population of microvesicles from the first sample; (c)

contacting the subpool to a second population of microvesicles from a second sample; and (d) depleting a second subpool of aptamers that show affinity to the second population of microvesicles from the second sample, thereby selecting the group of aptamers that have preferential affinity for the population of microvesicles from the first sample.

[00369] The first sample and/or second sample may comprise a biological fluid such as disclosed herein. For example, the biological fluid may include without limitation blood, a blood derivative, plasma, serum or urine. The first sample and/or second sample may also be derived from a cell culture.

[00370] In an embodiment, the first sample comprises a cancer sample and the second sample comprises a control sample, such as a non-cancer sample. The first sample and/or and the second sample may each comprise a pooled sample. For example, the first sample and/or second sample can comprise bodily fluid from 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100 or more than 100 individuals. In such cases, the members of a pool may be chosen to represent a desired phenotype. In a non-limiting example, the members of the first sample pool may be from patients with a cancer and the members of the second sample pool may be from non-cancer controls.

[00371] Steps (a)-(d) can be repeated a desired number of times in order to further enrich the pool in aptamers that have preferential affinity for the population of microvesicles from the first sample. For example, steps (a)-(d) can be repeated 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more than 20 times. The output from step (d) can be used as the input to repeated step (a). In embodiment, the first sample and/or second sample are replaced with a different sample before repeating steps (a)-(d). In a non-limiting example, members of a first sample pool may be from patients with a cancer and members of a second sample pool may be from non-cancer controls. During subsequent repetitions of steps (a)-(d), the first sample pool may comprise samples from different cancer patients than in the prior round/s. Similarly, the second sample pool may comprise samples from different controls than in the prior round/s.

[00372] In still another related aspect, the invention provides a method of enriching a plurality of oligonucleotides, comprising: (a) contacting a first microvesicle population with the plurality of oligonucleotides; (b) fractionating the first microvesicle population contacted in step (a) and recovering members of the plurality of oligonucleotides that fractionated with the first microvesicle population; (c) contacting the recovering members of the plurality of oligonucleotides from step (b) with a second microvesicle population; (d) fractionating the second microvesicle population contacted in step (c) and recovering members of the plurality of oligonucleotides that did not fractionate with the second microvesicle population; (e) contacting the recovering members of the plurality of oligonucleotides from step (d) with a third microvesicle population; and (f) fractionating the third microvesicle population contacted in step (a) and recovering members of the plurality of oligonucleotides that fractionated with the third microvesicle population; thereby enriching the plurality of oligonucleotides. The first and third microvesicle populations may have a first phenotype while the second microvesicle population has a second phenotype. Thus, positive selection occurs for the microvesicle populations associated with the

first phenotype and negative selection occurs for the microvesicle populations associated with the second phenotype. An example of such selection schemes is described in **Example 18** herein, wherein the first phenotype comprises biopsy-positive breast cancer and the second phenotype comprises non-breast cancer (biopsy-negative or healthy).

[00373] In some embodiments, the first phenotype comprises a medical condition, disease or disorder and the second phenotype comprises a healthy state or a different state of the medical condition, disease or disorder. The first phenotype can be a healthy state and the second phenotype comprises a medical condition, disease or disorder. The medical condition, disease or disorder can be any detectable medical condition, disease or disorder, including without limitation a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. Various types of such conditions are disclosed herein. See, e.g., Section "Phenotypes" herein.

[00374] Any useful method to isolate microvesicles in whole or in part can be used to fractionate the samples. See, e.g., Section "Microvesicle Isolation and Analysis" herein. In an embodiment, the fractionating comprises ultracentrifugation in step (b) and polymer precipitation in steps (d) and (f). The polymer can be polyethylene glycol (PEG). Any appropriate form of PEG may be used. For example, the PEG may be PEG 8000. The PEG may be used at any appropriate concentration. For example, the PEG can be used at a concentration of 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14% or 15% to isolate the microvesicles. In some embodiments, the PEG is used at a concentration of 6%.

[00375] The contacting can be performed in the presence of a competitor, which may reduce non-specific binding events. Any useful competitor can be used. In an embodiment, the competitor comprises at least one of salmon sperm DNA, tRNA, dextran sulfate and carboxymethyl dextran. As desired, different competitors or competitor concentrations can be used at different contacting steps.

[00376] The method can be repeated to achieve a desired enrichment. In an embodiment, steps (a)-(f) are repeated at least once. These steps can be repeated 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more than 20 times as desired. At the same time, each of the contacting steps can be repeated as desired. In some embodiments, the method further comprises: (i) repeating steps (a)-(b) at least once prior to step (c), wherein the recovered members of the plurality of oligonucleotides that fractionated with the first microvesicle population in step (b) are used as the input plurality of oligonucleotides for the repetition of step (a); (ii) repeating steps (c)-(d) at least once prior to step (e), wherein the recovered members of the plurality of oligonucleotides that did not fractionate with the second microvesicle population in step (d) are used as the input plurality of oligonucleotides for the repetition of step (c); and/or (iii) repeating steps (e)-(f) at least once, wherein the recovered members of the plurality of oligonucleotides that fractionated with the third microvesicle population in step (f) are used as the input plurality of oligonucleotides for the repetition of step (e). Repetitions (i)-(iii) can be repeated any desired number of times, e.g., (i)-(iii) can be repeated 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,

14, 15, 16, 17, 18, 19, 20 or more than 20 times. In an embodiment, (i)-(iii) each comprise three repetitions.

[00377] The method may further comprise identifying the members of the selected group of aptamers or oligonucleotides, e.g., by DNA sequencing. The sequencing may be performed by Next Generation sequencing as desired and after or before any desired step in the method.

[00378] The method may also comprise identifying the targets of the selected group of aptamers/oligonucleotides. Methods to identify such targets are disclosed herein.

[00379] ***Oligonucleotide Probe Target Identification***

[00380] The methods and kits above can be used to identify binding agents that differentiate between two biomarker populations. The invention further provides methods of identifying the targets of binding agent. For example, the methods may further comprise identifying a surface marker of a target microvesicle that is recognized by the binding agent.

[00381] In an embodiment, the invention provides a method of identifying a target of a binding agent comprising: (a) contacting the binding agent with the target to bind the target with the binding agent, wherein the target comprises a surface antigen of a microvesicle; (b) disrupting the microvesicle under conditions which do not disrupt the binding of the target with the binding agent; (c) isolating the complex between the target and the binding agent; and (d) identifying the target bound by the binding agent. The binding agent can be a binding agent identified by the methods above, e.g., an aptamer, ligand, antibody, or other useful binding agent that can differentiate between two populations of biomarkers.

[00382] An illustrative schematic for carrying on the method is shown in **FIG. 9**. The figure shows a binding agent **902**, here an oligonucleotide probe or aptamer for purposes of illustration, tethered to a substrate **901**. The binding agent **902** can be covalently attached to substrate **901**. The binding agent **902** may also be non-covalently attached. For example, binding agent **902** can comprise a label which can be attracted to the substrate, such as a biotin group which can form a complex with an avidin/streptavidin molecule that is covalently attached to the substrate. This can allow a complex to be formed between the aptamer and the microvesicle while in solution, followed by capture of the aptamer using the biotin label. The binding agent **902** binds to a surface antigen **903** of microvesicle **904**. In the step signified by arrow (i), the microvesicle is disrupted while leaving the complex between the binding agent **902** and surface antigen **903** intact. Disrupted microvesicle **905** is removed, e.g., via washing or buffer exchange, in the step signified by arrow (ii). In the step signified by arrow (iii), the surface antigen **903** is released from the binding agent **902**. The surface antigen **903** can be analyzed to determine its identity using methods disclosed herein and/or known in the art. The target of the method can be any useful biological entity associated with a microvesicle. For example, the target may comprise a protein, nucleic acid, lipid or carbohydrate, or other biological entity disclosed herein or known in the art.

[00383] In some embodiments of the method, the target is cross-linked to the binding agent prior disrupting the microvesicle. Without being bound by theory, this step may assist in maintaining the complex between the binding agent and the target while the vesicle is disrupted. Any useful method of

crosslinking disclosed herein or known in the art can be used. In embodiments, the cross-linking comprises photocrosslinking, an imidoester crosslinker, dimethyl suberimidate, an N-Hydroxysuccinimide-ester crosslinker, bissulfosuccinimidyl suberate (BS3), an aldehyde, acrolein, crotonaldehyde, formaldehyde, a carbodiimide crosslinker, N,N'-dicyclohexylcarbodiimide (DDC), N,N'-diisopropylcarbodiimide (DIC), 1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC or EDAC), Succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC), a Sulfosuccinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate (Sulfo-SMCC), a Sulfo-N-hydroxysuccinimidyl-2-(6-[biotinamido]-2-(p-azido benzamido)-hexanoamido) ethyl-1,3'-dithioproprionate (Sulfo-SBED), 2-[N2-(4-Azido-2,3,5,6-tetrafluorobenzoyl)-N6-(6-biotin-amidocaproyl)-L-lysinyl]ethyl methanethiosulfonate (Mts-Atf-Biotin; available from Thermo Fisher Scientific Inc, Rockford IL.), 2-{N2-[N6-(4-Azido-2,3,5,6-tetrafluorobenzoyl-6-amino-caproyl)-N6-(6-biotinamidocaproyl)-L-lysinylamido]}ethyl methanethiosultonate (Mts-Atf-LC-Biotin; available from Thermo Fisher Scientific Inc), a photoreactive amino acid (e.g., L-Photo-Leucine and L-Photo-Methionine, see, e.g., Suchanek, M., et al. (2005). Photo-leucine and photo-methionine allow identification of protein-protein interactions. Nat. Methods 2:261-267), an N-Hydroxysuccinimide (NHS) crosslinker, an NHS-Azide reagent (e.g., NHS-Azide, NHS-PEG4-Azide, NHS-PEG12-Azide; each available from Thermo Fisher Scientific, Inc.), an NHS-Phosphine reagent (e.g., NHS-Phosphine, Sulfo-NHS-Phosphine; each available from Thermo Fisher Scientific, Inc.), or any combination or modification thereof.

[00384] A variety of methods can be used to disrupt the microvesicle. For example, the vesicle membrane can be disrupted using mechanical forces, chemical agents, or a combination thereof. In embodiments, disrupting the microvesicle comprises use of one or more of a detergent, a surfactant, a solvent, an enzyme, or any useful combination thereof. The enzyme may comprise one or more of lysozyme, lysostaphin, zymolase, cellulase, mutanolysin, a glycanase, a protease, and mannase. The detergent or surfactant may comprise one or more of a octylthioglucoside (OTG), octyl beta-glucoside (OG), a nonionic detergent, Triton X, Tween 20, a fatty alcohol, a cetyl alcohol, a stearyl alcohol, cetostearyl alcohol, an oleyl alcohol, a polyoxyethylene glycol alkyl ether (Brij), octaethylene glycol monododecyl ether, pentaethylene glycol monododecyl ether, a polyoxypropylene glycol alkyl ether, a glucoside alkyl ether, decyl glucoside, lauryl glucoside, octyl glucoside, a polyoxyethylene glycol octylphenol ethers, a polyoxyethylene glycol alkylphenol ether, nonoxynol-9, a glycerol alkyl ester, glyceryl laurate, a polyoxyethylene glycol sorbitan alkyl esters, polysorbate, a sorbitan alkyl ester, cocamide MEA, cocamide DEA, dodecyldimethylamine oxide, a block copolymers of polyethylene glycol and polypropylene glycol, poloxamers, polyethoxylated tallow amine (POEA), a zwitterionic detergent, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS), a linear alkylbenzene sulfonate (LAS), a alkyl phenol ethoxylate (APE), cocamidopropyl hydroxysultaine, a betaine, cocamidopropyl betaine, lecithin, an ionic detergent, sodium dodecyl sulfate (SDS), cetrimonium bromide (CTAB), cetyl trimethylammonium chloride (CTAC), octenidine dihydrochloride, cetylpyridinium chloride (CPC), benzalkonium chloride (BAC), benzethonium chloride (BZT), 5-Bromo-5-nitro-1,3-dioxane,

dimethyldioctadecylammonium chloride, dioctadecyldimethylammonium bromide (DODAB), sodium deoxycholate, nonyl phenoxypolyethoxylethanol (Tergitol-type NP-40; NP-40), ammonium lauryl sulfate, sodium laureth sulfate (sodium lauryl ether sulfate (SLES)), sodium myreth sulfate, an alkyl carboxylate, sodium stearate, sodium lauroyl sarcosinate, a carboxylate-based fluorosurfactant, perfluorononanoate, perfluorooctanoate (PFOA or PFO), and a biosurfactant. Mechanical methods of disruption that can be used comprise without limitation mechanical shear, bead milling, homogenation, microfluidization, sonication, French Press, impingement, a colloid mill, decompression, osmotic shock, thermolysis, freeze-thaw, desiccation, or any combination thereof.

[00385] As shown in **FIG. 9**, the binding agent may be tethered to a substrate. The binding agent can be tethered before or after the complex between the binding agent and target is formed. The substrate can be any useful substrate such as disclosed herein or known in the art. In an embodiment, the substrate comprises a microsphere. In another embodiment, the substrate comprises a planar substrate. The binding agent can also be labeled. Isolating the complex between the target and the binding agent may comprise capturing the binding agent via the label. For example, the label can be a biotin label. In such cases, the binding agent can be attached to the substrate via a biotin-avidin binding event.

[00386] Methods of identifying the target after release from the binding agent will depend on the type of target of interest. For example, when the target comprises a protein, identifying the target may comprise use of mass spectrometry (MS), peptide mass fingerprinting (PMF; protein fingerprinting), sequencing, N-terminal amino acid analysis, C-terminal amino acid analysis, Edman degradation, chromatography, electrophoresis, two-dimensional gel electrophoresis (2D gel), antibody array, and immunoassay. Nucleic acids can be identified by sequencing.

[00387] One of skill will appreciate that the method can be used to identify any appropriate target, including those not associated with a vesicle. For example, with respect to the **FIG. 9**, all steps except for the step signified by arrow (i) (i.e., disrupting the microvesicle), could be performed for a circulating target such as a protein, nucleic acid, lipid, carbohydrate, or combination thereof. The target can be any useful target, including without limitation a cell, an organelle, a protein complex, a lipoprotein, a carbohydrate, a microvesicle, a virus, a membrane fragment, a small molecule, a heavy metal, a toxin, a drug, a nucleic acid, mRNA, microRNA, a protein-nucleic acid complex, and various combinations, fragments and/or complexes of any of these.

[00388] In an aspect, the invention provides a method of identifying at least one protein associated with at least one microvesicle in a biological sample, comprising: a) contacting the at least one microvesicle with an oligonucleotide probe library, b) isolating at least one protein bound by at least one member of the oligonucleotide probe library in step a); and c) identifying the at least one protein isolated in step b). The isolating can be performed using any useful method such as disclosed herein, e.g., by immunopreciption or capture to a substrate. Similarly, the identifying can be performed using any useful method such as disclosed herein, including without limitation use of mass spectrometry, 2-D gel electrophoresis or an antibody array. Examples of such methodology are presented herein in **Examples 23-25**.

[00389] The targets identified by the methods of the invention can be detected, e.g., using the oligonucleotide probes of the invention, for various purposes as desired. For example, an identified microvesicle surface antigen can then be used to detect a microvesicle. In an aspect, the invention provides a method of detecting at least one microvesicle in a biological sample comprising contacting the biological sample with at least one binding agent to at least one microvesicle surface antigen and detecting the at least one microvesicle recognized by the binding agent to the at least one protein. In an embodiment, the at least one microvesicle surface antigen is selected from **Tables 3-4** herein. The at least one microvesicle surface antigen can be a protein in any of **Tables 18-25**. See **Example 23**. The at least one binding agent may comprise any useful binding agent, including without limitation a nucleic acid, DNA molecule, RNA molecule, antibody, antibody fragment, aptamer, peptoid, zDNA, peptide nucleic acid (PNA), locked nucleic acid (LNA), unlocked nucleic acid (UNA), lectin, peptide, dendrimer, membrane protein labeling agent, chemical compound, or a combination thereof. In some embodiments, the at least one binding agent comprises at least one oligonucleotide, such as an oligonucleotide probe as provided herein.

[00390] The at least one binding agent can be used to capture and/or detect the at least one microvesicle. Methods of detecting biomarkers and microvesicle using binding agents are provided herein. See, e.g., **FIGs. 2A-B**, which figures describe sandwich assay formats. In some embodiments, the at least one binding agent used to capture the at least one microvesicle is bound to a substrate. Any useful substrate can be used, including without limitation a planar array, a column matrix, or a microbead. See, e.g., **FIGs. 2A-B**. In some embodiments, the at least one binding agent used to detect the at least one microvesicle is labeled. Various useful labels are provided herein or known in the art, including without limitation a magnetic label, a fluorescent moiety, an enzyme, a chemiluminescent probe, a metal particle, a non-metal colloidal particle, a polymeric dye particle, a pigment molecule, a pigment particle, an electrochemically active species, a semiconductor nanocrystal, a nanoparticle, a quantum dot, a gold particle, a fluorophore, or a radioactive label.

[00391] In an embodiment, the detecting is used to characterize a phenotype. The phenotype can be any appropriate phenotype of interest. In some embodiments, the phenotype is a disease or disorder. The characterizing may comprise providing diagnostic, prognostic and/or theranostic information for the disease or disorder. The characterizing may be performed by comparing a presence or level of the at least one microvesicle to a reference. The reference can be selected per the characterizing to be performed. For example, when the phenotype comprises a disease or disorder, the reference may comprise a presence or level of the at least one microvesicle in a sample from an individual or group of individuals without the disease or disorder. The comparing can be determining whether the presence or level of the microvesicle differs from that of the reference. In some embodiments, the detected at least one microvesicle is found at higher levels in a healthy sample as compared to a diseased sample. In another embodiment, the detected at least one microvesicle is found at higher levels in a diseased sample as compared to a healthy sample. When multiplex assays are performed, e.g., using a plurality of binding agents to different biomarkers,

some microvesicle antigens may be observed at a higher level in the biological samples as compared to the reference whereas other microvesicle antigens may be observed at a lower level in the biological samples as compared to the reference.

[00392] The method can be used to detect the at least one microvesicle in any appropriate biological sample. For example, the biological sample may comprise a bodily fluid, tissue sample or cell culture. The bodily fluid or tissue sample can be from a subject having or suspected of having a medical condition, a disease or a disorder. Thus, the method can be used to provide a diagnostic, prognostic, or theranostic read out for the subject. Any appropriate bodily fluid can be used, including without limitation peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair oil, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, or umbilical cord blood.

[00393] The method of the invention can be used to detect or characterize any appropriate disease or disorder of interest, including without limitation Breast Cancer, Alzheimer's disease, bronchial asthma, Transitional cell carcinoma of the bladder, Giant cellular osteoblastoclastoma, Brain Tumor, Colorectal adenocarcinoma, Chronic obstructive pulmonary disease (COPD), Squamous cell carcinoma of the cervix, acute myocardial infarction (AMI) / acute heart failure, Chron's Disease, diabetes mellitus type II, Esophageal carcinoma, Squamous cell carcinoma of the larynx, Acute and chronic leukemia of the bone marrow, Lung carcinoma, Malignant lymphoma, Multiple Sclerosis, Ovarian carcinoma, Parkinson disease, Prostate adenocarcinoma, psoriasis, Rheumatoid Arthritis, Renal cell carcinoma, Squamous cell carcinoma of skin, Adenocarcinoma of the stomach, carcinoma of the thyroid gland, Testicular cancer, ulcerative colitis, or Uterine adenocarcinoma.

[00394] In some embodiments, the disease or disorder comprises a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. The cancer can include without limitation one of acute lymphoblastic leukemia; acute myeloid leukemia; adrenocortical carcinoma; AIDS-related cancers; AIDS-related lymphoma; anal cancer; appendix cancer; astrocytomas; atypical teratoid/rhabdoid tumor; basal cell carcinoma; bladder cancer; brain stem glioma; brain tumor (including brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, astrocytomas, craniopharyngioma, ependymoblastoma, ependymoma, medulloblastoma, medulloepithelioma, pineal parenchymal tumors of intermediate differentiation, supratentorial primitive neuroectodermal tumors and pineoblastoma); breast cancer; bronchial tumors; Burkitt lymphoma; cancer of unknown primary site; carcinoid tumor; carcinoma of unknown primary site; central nervous system atypical teratoid/rhabdoid tumor; central nervous system embryonal tumors; cervical cancer; childhood cancers; chordoma; chronic lymphocytic leukemia; chronic myelogenous

leukemia; chronic myeloproliferative disorders; colon cancer; colorectal cancer; craniopharyngioma; cutaneous T-cell lymphoma; endocrine pancreas islet cell tumors; endometrial cancer; ependymoblastoma; ependymoma; esophageal cancer; esthesioneuroblastoma; Ewing sarcoma; extracranial germ cell tumor; extragonadal germ cell tumor; extrahepatic bile duct cancer; gallbladder cancer; gastric (stomach) cancer; gastrointestinal carcinoid tumor; gastrointestinal stromal cell tumor; gastrointestinal stromal tumor (GIST); gestational trophoblastic tumor; glioma; hairy cell leukemia; head and neck cancer; heart cancer; Hodgkin lymphoma; hypopharyngeal cancer; intraocular melanoma; islet cell tumors; Kaposi sarcoma; kidney cancer; Langerhans cell histiocytosis; laryngeal cancer; lip cancer; liver cancer; lung cancer; malignant fibrous histiocytoma bone cancer; medulloblastoma; medulloepithelioma; melanoma; Merkel cell carcinoma; Merkel cell skin carcinoma; mesothelioma; metastatic squamous neck cancer with occult primary; mouth cancer; multiple endocrine neoplasia syndromes; multiple myeloma; multiple myeloma/plasma cell neoplasm; mycosis fungoides; myelodysplastic syndromes; myeloproliferative neoplasms; nasal cavity cancer; nasopharyngeal cancer; neuroblastoma; Non-Hodgkin lymphoma; nonmelanoma skin cancer; non-small cell lung cancer; oral cancer; oral cavity cancer; oropharyngeal cancer; osteosarcoma; other brain and spinal cord tumors; ovarian cancer; ovarian epithelial cancer; ovarian germ cell tumor; ovarian low malignant potential tumor; pancreatic cancer; papillomatosis; paranasal sinus cancer; parathyroid cancer; pelvic cancer; penile cancer; pharyngeal cancer; pineal parenchymal tumors of intermediate differentiation; pineoblastoma; pituitary tumor; plasma cell neoplasm/multiple myeloma; pleuropulmonary blastoma; primary central nervous system (CNS) lymphoma; primary hepatocellular liver cancer; prostate cancer; rectal cancer; renal cancer; renal cell (kidney) cancer; renal cell cancer; respiratory tract cancer; retinoblastoma; rhabdomyosarcoma; salivary gland cancer; Sézary syndrome; small cell lung cancer; small intestine cancer; soft tissue sarcoma; squamous cell carcinoma; squamous neck cancer; stomach (gastric) cancer; supratentorial primitive neuroectodermal tumors; T-cell lymphoma; testicular cancer; throat cancer; thymic carcinoma; thymoma; thyroid cancer; transitional cell cancer; transitional cell cancer of the renal pelvis and ureter; trophoblastic tumor; ureter cancer; urethral cancer; uterine cancer; uterine sarcoma; vaginal cancer; vulvar cancer; Waldenström macroglobulinemia; or Wilm's tumor. The premalignant condition can include without limitation Barrett's Esophagus. The autoimmune disease can include without limitation one of inflammatory bowel disease (IBD), Crohn's disease (CD), ulcerative colitis (UC), pelvic inflammation, vasculitis, psoriasis, diabetes, autoimmune hepatitis, multiple sclerosis, myasthenia gravis, Type I diabetes, rheumatoid arthritis, psoriasis, systemic lupus erythematosis (SLE), Hashimoto's Thyroiditis, Grave's disease, Ankylosing Spondylitis Sjogrens Disease, CREST syndrome, Scleroderma, Rheumatic Disease, organ rejection, Primary Sclerosing Cholangitis, or sepsis. The cardiovascular disease can include without limitation one of atherosclerosis, congestive heart failure, vulnerable plaque, stroke, ischemia, high blood pressure, stenosis, vessel occlusion or a thrombotic event. The neurological disease can include without limitation one of Multiple Sclerosis (MS), Parkinson's Disease (PD), Alzheimer's Disease (AD), schizophrenia, bipolar disorder, depression, autism, Prion

Disease, Pick's disease, dementia, Huntington disease (HD), Down's syndrome, cerebrovascular disease, Rasmussen's encephalitis, viral meningitis, neurospsychiatric systemic lupus erythematosus (NPSLE), amyotrophic lateral sclerosis, Creutzfeldt-Jacob disease, Gerstmann-Straussler-Scheinker disease, transmissible spongiform encephalopathy, ischemic reperfusion damage (e.g. stroke), brain trauma, microbial infection, or chronic fatigue syndrome. The pain can include without limitation one of fibromyalgia, chronic neuropathic pain, or peripheral neuropathic pain. The infectious disease can include without limitation one of a bacterial infection, viral infection, yeast infection, Whipple's Disease, Prion Disease, cirrhosis, methicillin-resistant staphylococcus aureus, HIV, HCV, hepatitis, syphilis, meningitis, malaria, tuberculosis, or influenza. One of skill will appreciate that oligonucleotide probes or plurality of oligonucleotides or methods of the invention can be used to assess any number of these or other related diseases and disorders.

[00395] In a related aspect, the invention provides a kit comprising a reagent for carrying out the methods herein. In still another related aspect, the invention provides for use of a reagent for carrying out the methods. The reagent may comprise at least one binding agent to the at least one protein. The binding agent may be an oligonucleotide probe as provided herein.

[00396] ***Sample Characterization***

[00397] The aptamers of the invention can be used to characterize a biological sample. For example, an aptamer can be used to bind a biomarker in the sample. The presence or level of the bound biomarker can indicate a characteristic of the example, such as a diagnosis, prognosis or theranosis of a disease or disorder associated with the sample.

[00398] In an aspect, the invention provides an aptamer comprising a nucleic acid sequence that is at least about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99 or 100 percent homologous of any of: a) SEQ ID NOs. 11-24 or a sequence in **Table 8**; a) SEQ ID NOs. 25-44 or a sequence in **Table 11**; b) SEQ ID NOs. 47-130 or a sequence in **Table 12**; or c) a functional variation or fragment of any preceding sequence. A functional variation or fragment includes a sequence comprising modifications that is still capable of binding a target molecule, wherein the modifications comprise without limitation at least one of a deletion, insertion, point mutation, truncation or chemical modification. In a related aspect, the invention provides a method of characterizing a disease or disorder, comprising: (a) contacting a biological test sample with one or more aptamer of the invention, e.g., any of those in this paragraph or modifications thereof; (b) detecting a presence or level of a complex between the one or more aptamer and the target bound by the one or more aptamer in the biological test sample formed in step (a); (c) contacting a biological control sample with the one or more aptamer; (d) detecting a presence or level of a complex between the one or more aptamer and the target bound by the one or more aptamer in the biological control sample formed in step (c); and (e) comparing the presence or level detected in steps (b) and (d), thereby characterizing the disease or disorder.

[00399] The biological test sample and biological control sample can each comprise a tissue sample, a cell culture, or a biological fluid. In some embodiments, the biological test sample and biological control

sample comprise the same sample type, e.g., both are tissue samples or both are fluid samples. In other embodiments, different sample types may be used for the test and control samples. For example, the control sample may comprise an engineered or otherwise artificial sample.

[00400] The biological fluid may comprise a bodily fluid. The bodily fluid may include without limitation one or more of peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, or umbilical cord blood. In some embodiments, the bodily fluid comprises blood, serum or plasma.

[00401] The biological fluid may comprise microvesicles. For example, the biological fluid can be a tissue, cell culture, or bodily fluid which comprises microvesicles released from cells in the sample. The microvesicles can be circulating microvesicles.

[00402] The one or more aptamer can bind a target biomarker, e.g., a biomarker useful in characterizing the sample. The biomarker may comprise a polypeptide or fragment thereof, or other useful biomarker described herein or known in the art (lipid, carbohydrate, complex, nucleic acid, etc). In embodiments, the polypeptide or fragment thereof is soluble or membrane bound. Membrane bound polypeptides may comprise a cellular surface antigen or a microvesicle surface antigen. The biomarker can be a biomarker selected from **Table 3** or **Table 4**.

[00403] The characterizing can comprises a diagnosis, prognosis or theranosis of the disease or disorder. Various diseases and disorders can be characterized using the compositions and methods of the invention, including without limitation a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, a neurological disease or disorder, an infectious disease, and/or pain. See, e.g., section herein "Phenotypes" for further details. In embodiments, the disease or disorder comprises a proliferative or neoplastic disease or disorder. For example, the disease or disorder can be a cancer. In some embodiments, the cancer comprises a breast cancer, ovarian cancer, prostate cancer, lung cancer, colorectal cancer, melanoma, or brain cancer.

[00404] **FIG. 19A** is a schematic **1900** showing an assay configuration that can be used to detect and/or quantify a target of interest using one or more aptamer of the invention. Capture aptamer **1902** is attached to substrate **1901**. The substrate can be a planar substrate, well, microbead, or other useful substrate as disclosed herein or known in the art. Target of interest **1903** is bound by capture aptamer **1902**. The target of interest can be any appropriate entity that can be detected when recognized by an aptamer or other binding agent. The target of interest may comprise a protein or polypeptide, a nucleic acid, including DNA, RNA, and various subspecies thereof, a lipid, a carbohydrate, a complex, e.g., a complex comprising protein, nucleic acids, lipids and/or carbohydrates. In some embodiments, the target of interest comprises a microvesicle. The target of interest can be a microvesicle surface antigen. The target of

interest may be a biomarker, including a vesicle associated biomarker, in **Tables 3** or **4**. The microvesicle input can be isolated from a sample using various techniques as described herein, e.g., chromatography, filtration, centrifugation, flow cytometry, affinity capture (e.g., to a planar surface, column or bead), and/or using microfluidics. Detection aptamer **1904** is also bound to target of interest **1903**. Detection aptamer **1904** carries label **1905** which can be detected to identify target captured to substrate **1901** via capture aptamer **1902**. The label can be a fluorescent, radiolabel, enzyme, or other detectable label as disclosed herein. Either capture aptamer **1902** or detection aptamer **1904** can be substituted with another binding agent, e.g., an antibody. For example, the target may be captured with an antibody and detected with an aptamer, or vice versa. When the target of interest comprises a complex, the capture and detection agents (aptamer, antibody, etc) can recognize the same or different targets. For example, when the target is a microvesicle, the capture agent may recognize one microvesicle surface antigen while the detection agent recognizes another microvesicle surface antigen. Alternately, the capture and detection agents can recognize the same surface antigen.

[00405] The aptamers of the invention may be identified and/or used for various purposes in the form of DNA or RNA. Unless otherwise specified, one of skill in the art will appreciate that an aptamer may generally be synthesized in various forms of nucleic acid. The aptamers may also carry various chemical modifications and remain within the scope of the invention.

[00406] In some embodiments, an aptamer of the invention is modified to comprise at least one chemical modification. The modification may include without limitation a chemical substitution at a sugar position; a chemical substitution at a phosphate position; and a chemical substitution at a base position of the nucleic acid. In some embodiments, the modification is selected from the group consisting of: biotinylation, incorporation of a fluorescent label, incorporation of a modified nucleotide, a 2'-modified pyrimidine, 3' capping, conjugation to an amine linker, conjugation to a high molecular weight, non-immunogenic compound, conjugation to a lipophilic compound, conjugation to a drug, conjugation to a cytotoxic moiety, and labeling with a radioisotope, or other modification as disclosed herein. The position of the modification can be varied as desired. For example, the biotinylation, fluorescent label, or cytotoxic moiety can be conjugated to the 5' end of the aptamer. The biotinylation, fluorescent label, or cytotoxic moiety can also be conjugated to the 3' end of the aptamer.

[00407] In some embodiments, the cytotoxic moiety is encapsulated in a nanoparticle. The nanoparticle can be selected from the group consisting of: liposomes, dendrimers, and comb polymers. In other embodiments, the cytotoxic moiety comprises a small molecule cytotoxic moiety. The small molecule cytotoxic moiety can include without limtation vinblastine hydrazide, calicheamicin, vinca alkaloid, a cryptophycin, a tubulysin, dolastatin-10, dolastatin-15, auristatin E, rhizoxin, epothilone B, epithilone D, taxoids, maytansinoids and any variants and derivatives thereof. In still other embodiments, the cytotoxic moiety comprises a protein toxin. For example, the protein toxin can be selected from the group consisting of diphtheria toxin, ricin, abrin, gelonin, and Pseudomonas exotoxin A. Non-immunogenic, high molecular weight compounds for use with the invention include polyalkylene glycols, e.g., polyethylene

glycol. Appropriate radioisotopes include yttrium-90, indium-111, iodine-131, lutetium-177, copper-67, rhenium-186, rhenium-188, bismuth-212, bismuth-213, astatine-211, and actinium-225. The aptamer may be labeled with a gamma-emitting radioisotope.

[00408] In some embodiments of the invention, an active agent is conjugated to the aptamer. For example, the active agent may be a therapeutic agent or a diagnostic agent. The therapeutic agent may be selected from the group consisting of tyrosine kinase inhibitors, kinase inhibitors, biologically active agents, biological molecules, radionuclides, adriamycin, ansamycin antibiotics, asparaginase, bleomycin, busulphan, cisplatin, carboplatin, carmustine, capecotabine, chlorambucil, cytarabine, cyclophosphamide, camptothecin, dacarbazine, dactinomycin, daunorubicin, dexrazoxane, docetaxel, doxorubicin, etoposide, epothilones, floxuridine, fludarabine, fluorouracil, gemcitabine, hydroxyurea, idarubicin, ifosfamide, irinotecan, lomustine, mechlorethamine, mercaptopurine, melphalan, methotrexate, rapamycin (sirolimus), mitomycin, mitotane, mitoxantrone, nitrosurea, paclitaxel, pamidronate, pentostatin, plicamycin, procarbazine, rituximab, streptozocin, teniposide, thioguanine, thiotepa, taxanes, vinblastine, vincristine, vinorelbine, taxol, combretastatins, discodermolides, transplatinum, anti-vascular endothelial growth factor compounds ("anti-VEGFs"), anti-epidermal growth factor receptor compounds ("anti-EGFRs"), 5-fluorouracil and derivatives, radionuclides, polypeptide toxins, apoptosis inducers, therapy sensitizers, enzyme or active fragment thereof, and combinations thereof.

[00409] ***Oligonucleotide Pools to Characterize a Sample***

[00410] The complexity and heterogeneity present in biology challenges the understanding of biological systems and disease. Diversity exists at various levels, e.g., within and between cells, tissues, individuals and disease states. See, e.g., **FIG. 20A**. **FIG. 20B** overviews various biological entities that can be assessed to characterize such samples. As shown in the Figure, as one moves from assessing DNA, to RNA, to protein, and finally to protein complexes, the amount of diversity and complexity increases dramatically. The oligonucleotide probe library method of the invention can be used characterize complex biological sources, e.g., tissue samples, cells, circulating tumor cells, microvesicles, and complexes such as protein and proteolipid complexes.

[00411] Current methods to characterize biological samples may not adequately address such complexity and diversity. As shown in **FIG. 20C**, such current methods often have a trade off between measuring diversity and complexity. As an example, consider high throughput sequencing technology. Next generation approaches may query many 1000s of molecular targets in a single assay. However, such approaches only probe individual DNA and/or RNA molecules, and thus miss out on the great diversity of proteins and biological complexes. On the other hand, flow cytometry can probe biological complexes, but are limited to a small number of pre-defined ligands. For example, a single assay can probe a handful of differentially labeled antibodies to pre-defined targets.

[00412] The oligonucleotide probe library of the invention address the above challenges with current biological detection technologies. The size of the starting library can be adjusted to measure as many different entities as there are library members. In this Example, the initial untrained oligonucleotide

library has the potential to measure 10^{12} or more biological features. A larger and/or different library can be constructed as desired. The technology is adapted to find differences between samples without assumptions about what "should be different." For example, the probe library may distinguish based on individual proteins, protein modifications, protein complexes, lipids, nucleic acids, different folds or conformations, or whatever is there that distinguishes a sample of interest. Thus, the method provides an unbiased approach to identify differences in biological samples that can be used to identify different populations of interest.

[00413] In the context herein, the use of the oligonucleotide library probe to assess a sample may be referred to as Adaptive Dynamic Artificial Poly-ligand Targeting, or ADAPT™ (previously referred to as Topological Oligonucleotide Profiling: TOP™). Although as noted the terms aptamer and oligonucleotides are typically used interchangeable herein, some differences between "classic" individual aptamers and ADAPT probes are as follows. Individual aptamers may comprise individual oligonucleotides selected to bind to a known specific target in an antibody-like "key-in-lock" binding mode. They may be evaluated individually based on specificity and binding affinity to the intended target. However, ADAPT probes may comprise a library of oligonucleotides intended to produce multi-probe signatures. The ADAPT probes comprise numerous potential binding modalities (electrostatic, hydrophobic, Watson-Crick, multi-oligo complexes, etc.). The ADAPT probe signatures have the potential to identify heterogeneous patient subpopulations. For example, a single ADAPT probe library can be assembled to differentiate multiple disease states, as demonstrated herein. Unlike classic single aptamers, the binding targets may or may not be isolated or identified. It will be understood that screening methods that identify individual aptamers, e.g., SELEX, can also be used to enrich a naive library of oligonucleotides to identify a ADAPT probe library.

[00414] The general method of the invention is outlined in **FIG. 20D**. One input to the method comprises a randomized oligonucleotide library with the potential to measure 10^{12} or more biological features. As outlined in the figure, the method identifies a desired number (e.g., ~10^5-10^6) that are different between two input sample types. The randomized oligonucleotide library is contacted with a first and a second sample type, and oligonucleotides that bind to each sample are identified. The bound oligonucleotide populations are compared and oligonucleotides that specifically bind to one or the other biological input sample are retained for the oligonucleotide probe library, whereas oligonucleotides that bind both biological input samples are discarded. This trained oligonucleotide probe library can then be contacted with a new test sample and the identities of oligonucleotides that bind the test sample are determined. The test sample is characterized based on the profile of oligonucleotides that bound. See, e.g., **FIG. 20H**.

[00415] Extracellular vesicles provide an attractive vehicle to profile the biological complexity and diversity driven by many inter-related sources. There can be a great deal of heterogeneity between patient-to-patient microvesicle populations, or even in microvesicle populations from a single patient under different conditions (e.g., stress, diet, exercise, rest, disease, etc). Diversity of molecular phenotypes within microvesicle populations in various disease states, even after microvesicle isolation and sorting by

vesicle biomarkers, can present challenges identifying surface binding ligands. This situation is further complicated by vesicle surface-membrane protein complexes. The oligonucleotide probe library can be used to address such challenges and allow for characterization of biological phenotypes. The approach combines the power of diverse oligonucleotide libraries and high throuput (next-generation) sequencing technologies to probe the complexity of extracellular microvesicles. See **FIG. 20E**.

[00416] ADAPT™ profiling may provide quantitative measurements of dynamic events in addition to detection of presence/absence of various biomarkers in a sample. For example, the binding probes may detect protein complexes or other post-translation modifications, allowing for differentiation of samples with the same proteins but in different biological configurations. Such configurations are illustrated in **FIGs. 20F-G**. In **FIG. 20F**, microvesicles with various surface markers are shown from an example microvesicle sample population: Sample Population A. The indicated Bound Probing Oligonucleotides **2001** are contacted to two surface markers **2002** and **2003** in a given special relationship. Here, probes unique to these functional complexes and spatial relationships may be retained. In contrast, in microvesicle Sample Population B shown in **FIG. 20F**, the two surface markers **2002** and **2003** are found in disparate spacial relationship. Here, probes **2001** are not bound due to absence of the spatial relationship of the interacting components **2002** and **2003**.

[00417] An illustrative approach **2010** for using ADAPT profiling to assess a sample is shown in **FIG. 20H**. The probing library **2011** is mixed with sample **2012**. The sample can be as described herein, e.g., a bodily fluid from a subject having or suspected of having a disease. The probes are allowed to bind the sample **2020** and the microvesicles are pelleted **2015**. The supernatant **2014** comprising unbound oligonucleotides is discarded. Oligonucleotide probes bound to the pellet **2015** are eluted **2016** and sequenced **2017**. The profile **2018** generated by the bound oligonucleotide probes as determined by the sequening **2017** is used to characterize the sample **2012**. For example, the profile **2018** can be compared to a reference, e.g., to determine if the profile is similar or different from a reference profile indicative of a disease or healthy state, or other phenotypic characterization of interest. The comparison may indicate the presence of a disease, provide a diagnosis, prognosis or theranosis, or otherwise characterize a phenotype associated with the sample **2012**. **FIG. 20I** illustrates another schematic for using TOP™ profiling to characterize a phenotype. A patient sample such as a bodily fluid disclosed herein is collected **2021**. The sample is contacted with the ADAPT ™ library pool **2022**. Microvesicles (MVs) are isolated from the contacted sample **2023**, e.g., using ultracentrifugation, filtration, polymer precipitation or other appropriate technique or combination of techniques disclosed herein. Oligonucleotides that bound the isolated microvesicles are collected and identity is determined **2024**. The identity of the bound oligonucleotides can be determined by any useful technique such as sequencing, high throughput sequencing (e.g., NGS), amplification including without limitation qPCR, or hybridization such as to a planar or particle based array. The identity of the bound oligonucleotides is used to characterize the sample, e.g., as containing disease related microvesicles.

[00418] In an aspect, the invention provides a method of characterizing a sample by contacting the sample with a pool of different oligonucleotides (e.g., an aptamer pool), and determining the frequency at which various oligonucleotides in the pool bind the sample. For example, a pool of oligonucleotides is identified that preferentially bind to microvesicles from cancer patients as compared to non-cancer patients. A test sample, e.g., from a patient suspected of having the cancer, is collected and contacted with the pool of oligonucleotides. Oligonucleotides that bind the test sample are eluted from the test sample, collected and identified, and the composition of the bound oligonucleotides is compared to those known to bind cancer samples. Various sequencing, amplification and hybridization techinques can be used to identify the eluted oligonucleotides. For example, when a large pool of oligonucleotides is used, oligonucleotide identification can be performed by high throughput methods such as next generation sequencing or via hybridization. If the test sample is bound by the oligonucleotide pool in a similar manner (e.g., as determined by bioinformatics classification methods) to the microvesicles from cancer patients, then the test sample is indicative of cancer as well. Using this method, a pool of oligonucleotides that bind one or more microvesicle antigen can be used to characterize the sample without necessarily knowing the precise target of each member of the pool of oligonucleotides. **Examples 26-30** and others herein illustrate embodiments of the invention.

[00419] In an aspect, the invention provides a method for characterizing a condition for a test sample comprising: contacting a microvesicle sample with a plurality of oligonucleotide capable of binding one or more target(s) present in said microvesicle sample, identifying a set of oligonucleotides that form a complex with the sample wherein the set is predetermined to characterize a condition for the sample, thereby characterizing a condition for a sample.

[00420] In an related aspect, the invention provides a method for identifying a set of oligonucleotides associated with a test sample, comprising: (a) contacting a microvesicle sample with a plurality of oligonucleotides, isolating a set of oligonucleotides that form a complex with the microvesicle sample, (b) determining sequence and/or copy number for each of the oligonucleotides, thereby identifying a set of oligonucleotides associated with the test sample.

[00421] In still another related aspect, the invention provides a method of diagnosing a sample as cancerous or predisposed to be cancerous, comprising contacting a microvesicle sample with a plurality of oligonucleotides that are predetermined to preferentially form a complex with microvesicles from a cancer sample as compared to microvesicles from a non-cancer sample.

[00422] The oligonucleotides can be identified by sequencing, e.g., by dye termination (Sanger) sequencing or high throughput methods. High throughput methods can comprise techiques to rapidly sequence a large number of nucleic acids, including next generation techniques such as Massively parallel signature sequencing (MPSS; Polony sequencing; 454 pyrosequencing; Illumina (Solexa) sequencing; SOLiD sequencing; Ion Torrent semiconductor sequencing; DNA nanoball sequencing; Heliscope single molecule sequencing; Single molecule real time (SMRT) sequencing, or other methods such as Nanopore DNA sequencing; Tunnelling currents DNA sequencing; Sequencing by hybridization; Sequencing with

mass spectrometry; Microfluidic Sanger sequencing; Microscopy-based techniques; RNAP sequencing; In vitro virus high-throughput sequencing. The oligonucleotides may also be identified by hybridization techniques. For example, a microarray having addressable locals to hybridize and thereby detect the various members of the pool can be used. Alternately, detection can be based on one or more differentially labelled oligonucleotides that hybridize with various members of the oligonucleotide pool. The detectable signal of the label can be associated with a nucleic acid molecule that hybridizes with a stretch of nucleic acids present in various oligonucleotides. The stretch can be the same or different as to one or more oligonucleotides in a library. The detectable signal can comprise fluorescence agents, including color-coded barcodes which are known, such as in U.S. Patent Application Pub. No. 20140371088, 2013017837, and 20120258870. Other detectable labels (metals, radioisotopes, etc) can be used as desired.

[00423] The plurality or pool of oligonucleotides can comprise any desired number of oligonucleotides to allow characterization of the sample. In various embodiments, the pool comprises at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, or at least 10000 different oligonucleotide members.

[00424] The plurality of oligonucleotides can be pre-selected through one or more steps of positive or negative selection, wherein positive selection comprises selection of oligonucleotides against a sample having substantially similar characteristics compared to the test sample, and wherein negative selection comprises selection of oligonucleotides against a sample having substantially different characteristics compared to the test sample. Substantially similar characteristics mean that the samples used for positive selection are representative of the test sample in one or more characteristic of interest. For example, the samples used for positive selection can be from cancer patients or cell lines and the test sample can be a sample from a patient having or suspected to have a cancer. Substantially different characteristics mean that the samples used for negative selection differ from the test sample in one or more characteristic of interest. For example, the samples used for negative selection can be from individuals or cell lines that do not have cancer (e.g., "normal" or otherwise "control" samples) and the test sample can be a sample from a patient having or suspected to have a cancer. The cancer can be a breast cancer, ovarian cancer, prostate cancer, lung cancer, colorectal cancer, melanoma, brain cancer, or other cancer.

[00425] By selecting samples representative of the desired phenotypes to detect and/or distinguish, the characterizing can comprise a diagnosis, prognosis or theranosis for any number of diseases or disorders. Various diseases and disorders can be characterized using the compositions and methods of the invention, including without limitation a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, a neurological disease or disorder, an infectious disease, and/or pain. See, e.g., section herein "Phenotypes" for further details. In embodiments, the disease or disorder comprises a proliferative or neoplastic disease or disorder. For

example, the disease or disorder can be a cancer. In some embodiments, the cancer comprises a breast cancer, ovarian cancer, prostate cancer, lung cancer, colorectal cancer, melanoma, or brain cancer.

[00426] **FIG. 19B** is a schematic **1910** showing use of an oligonucleotide pool to characterize a phenotype of a sample, such as those listed above. A pool of oligonucleotides to a target of interst is provided **1911**. For example, the pool of oligonucleotides can be enriched to target one or more microvesicle. The members of the pool may bind different targets (e.g., a microvesicle surface antigen) or different epitopes of the same target present on the one or more microvesicle. The pool is contacted with a test sample to be characterized **1912**. For example, the test sample may be a biological sample from an individual having or suspected of having a given disease or disorder. The mixture is washed to remove unbound oligonucleotides. The remaining oligonucleotides are eluted or otherwise disassociated from the sample and collected **1913**. The collected oligonucleotides are identified, e.g., by sequencing or hybridization **1914**. The presence and/or copy number of the identified is used to characterize the phenotype **1915**. For example, the pool of oligonucleotides may be chosen as oligonucleotides that preferentially recognize microvesicles shed from cancer cells. The method can be employed to detect whether the sample retains oligonucleotides that bind the cancer-related microvesicles, thereby allowing the sample to be characterized as cancerous or not.

[00427] **FIG. 19C** is a schematic **1920** showing an implementation of the method in **FIG. 19B**. A pool of oligonucleotides identified as binding a microvesicle population is provided **1919**. The input sample comprises a test sample comprising microvesicles **1922**. For example, the test sample may be a biological sample from an individual having or suspected of having a given disease or disorder. The pool is contacted with the isolated microvesicles to be characterized **1923**. The microvesicle population can be isolated before or after the contacting **1923** from the sample using various techniques as described herein, e.g., chromatography, filtration, ultrafiltration, centrifugation, ultracentrifugation, flow cytometry, affinity capture (e.g., to a planar surface, column or bead), polymer precipitation, and/or using microfluidics. The mixture is washed to remove unbound oligonucleotides and the remaining oligonucleotides are eluted or otherwise disassociated from the sample and collected **1924**. The collected oligonucleotides are identified **1925** and the presence and/or copy number of the retained oligonucleotides is used to characterize the phenotype **1926** as above.

[00428] As noted, in embodiment of **FIG. 19C**, the pool of oligonucleotides **1919** is directly contacted with a biological sample that comprises or is expected to comprise microvesicles. Microvesicles are thereafter isolated from the sample and the mixture is washed to remove unbound oligonucleotides and the remaining oligonucleotides are disassociated and collected **1924**. The following steps are performed as above. As an example of this alternate configuration, a biological sample, e.g., a blood, serum or plasma sample, is directly contacted with the pool of oligonucleotides. Microvesicles are then isolated by various techniques disclosed herein, including without limitation ultracentrifugation, ultrafiltration, flow cytometry, affinity isolation, polymer precipitation, chromatography, various combinations thereof, or the like. Remaining oligonucleotides are then identified, e.g., by sequencing, hybridization or amplification.

[00429] In a related aspect, the invention provides a composition of matter comprising a plurality of oligonucleotides that can be used to carry out the methods comprising use of an oligonucleotide pool to characterize a phenotype. The plurality of oligonucleotides can comprise any of those described herein.

[00430] In a related aspect, the invention provides a method of performing high-throughput sequencing comprising: performing at least one (i) negative selection or (ii) one positive selection of a plurality of oligonucleotides with a microvesicle sample; obtaining a set of oliognucleotides to provide a negative binder subset or positive binder subset of the plurality of oligonucleotides, wherein the negative binder subset of the plurality of oligonucleotides does not bind the microvesicle sample and wherein the positive binder subset of the plurality of oligonucleotides does bind the microvesicle sample; contacting the negative binder subset or positive binder subset with a test sample; eluting oligonucleotides that bound to the test sample to provide a plurality of eluate oligonucleotides; and performing high-throughput sequencing of the plurality of eluate oligonucleotides to identify sequence and/or copy number of the members of the plurality of eluate oligonucleotides. Negative and positive selection of the plurality of oligonucleotides using microvesicle sample can be performed as disclosed herein. The oligonucleotide profile revealed by the sequence and/or copy number of the members of the plurality of eluate oligonucleotides can be used to characterize a phenotype of the test sample as described herein.

[00431] In a similar aspect, the invention provides a method for identifying oligonucleotides specific for a test sample. The method comprises: (a) enriching a plurality of oligonucleotides for a sample to provide a set of oligonucleotides predetermined to form a complex with a target sample; (b) contacting the plurality in (a) with a test sample to allow formation of complexes of oligonucleotides with test sample; (c) recovering oligonucleotides that formed complexes in (b) to provide a recovered subset of oligonucleotides; and (d) profiling the recovered subset of oligonucleotides by high-throughput sequencing or hybridization, thereby identifying oligonucleotides specific for a test sample. The test sample may comprise a plurality of microvesicles. The oligonucleotides may comprise RNA, DNA or both. In some embodiment, the method further comprises performing informatics analysis to identify a subset of oligonucleotides comprising sequence identity of at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or at least 99% to the oligonucleotides predetermined to form a complex with the target sample.

[00432] One of skill will appreciate that the method can be used to identify any appropriate target, including those not associated with a microvesicle. The target can be any useful target, including without limitation a cell, an organelle, a protein complex, a lipoprotein, a carbohydrate, a microvesicle, a virus, a membrane fragment, a small molecule, a heavy metal, a toxin, a drug, a nucleic acid (including without limitation microRNA (miR) and messenger RNA (mRNA)), a protein-nucleic acid complex, and various combinations, fragments and/or complexes of any of these. The target can, e.g., comprise a mixture of microvesicles and non-microvesicle entities.

[00433] In an aspect, the invention also provides a method comprising contacting an oligonucleotide or plurality of oligonucleotides with a sample and detecting the presence or level of binding of the

oligonucleotide or plurality of oligonucleotides to a target in the sample, wherein the oligonucleotide or plurality of oligonucleotides can be those provided by the invention above. The sample may comprise a biological sample, an organic sample, an inorganic sample, a tissue, a cell culture, a bodily fluid, blood, serum, a cell, a microvesicle, a protein complex, a lipid complex, a carbohydrate, or any combination, fraction or variation thereof. The target may comprise a cell, an organelle, a protein complex, a lipoprotein, a carbohydrate, a microvesicle, a membrane fragment, a small molecule, a heavy metal, a toxin, or a drug.

[00434] In a related aspect, the invention provides a method comprising: a) contacting a biological sample comprising microvesicles with an oligonucleotide probe library, wherein optionally the oligonucleotide probe library comprises an oligonucleotide or plurality of oligonucleotides those provided by the invention above; b) identifying oligonucleotides bound to at least a portion of the microvesicles; and c) characterizing the sample based on a profile of the identified oligonucleotides.

[00435] In another aspect, the invention provides a method comprising: a) contacting a sample with an oligonucleotide probe library comprising at least 10^6, 10^7, 10^8, 10^9, 10^{10}, 10^{11}, 10^{12}, 10^{13}, 10^{14}, 10^{15}, 10^{16}, 10^{17}, or at least 10^{18} different oligonucleotide sequences oligonucleotides to form a mixture in solution, wherein the oligonucleotides are capable of binding a plurality of entities in the sample to form complexes, wherein optionally the oligonucleotide probe library comprises an oligonucleotide or plurality of oligonucleotides as provided by the invention above; b) partitioning the complexes formed in step (a) from the mixture; and c) detecting oligonucleotides present in the complexes partitioned in step (b) to identify an oligonucleotide profile for the sample. In an embodiment, the detecting step comprises performing sequencing of all or some of the oligonucleotides in the complexes, amplification of all or some of the oligonucleotides in the complexes, and/or hybridization of all or some of the oligonucleotides in the complexes to an array. The array can be any useful array, such as a planar or particle-based array.

[00436] In still another aspect, the invention provides a method for generating an enriched oligonucleotide probe library comprising: a) contacting a first oligonucleotide library with a biological test sample and a biological control sample, wherein complexes are formed between biological entities present in the biological samples and a plurality of oligonucleotides present in the first oligonucleotide library; b) partitioning the complexes formed in step (a) and isolating the oligonucleotides in the complexes to produce a subset of oligonucleotides for each of the biological test sample and biological control sample; c) contacting the subsets of oligonucleotides in (b) with the biological test sample and biological control sample wherein complexes are formed between biological entities present in the biological samples and a second plurality of oligonucleotides present in the subsets of oligonucleotides to generate a second subset group of oligonucleotides; and d) optionally repeating steps b)-c), one, two, three or more times to produce a respective third, fourth, fifth or more subset group of oligonucleotides, thereby producing the enriched oligonucleotide probe library. In a related aspect, the invention provides a plurality of oligonucleotides comprising at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 300, 400, 500, 600, 700, 800, 900,

1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000, 200000, 300000, 400000, or 500000 different oligonucleotide sequences, wherein the plurality results from the method in this paragraph, wherein the library is capable of distinguishing a first phenotype from a second phenotype. In some embodiments, the first phenotype comprises a disease or disorder and the second phenotype comprises a healthy state; or wherein the first phenotype comprises a disease or disorder and the second phenotype comprises a different disease or disorder; or wherein the first phenotype comprises a stage or progression of a disease or disorder and the second phenotype comprises a different stage or progression of the same disease or disorder; or wherein the first phenotype comprises a positive response to a therapy and the second phenotype comprises a negative response to the same therapy.

[00437] In yet another aspect, the invention provides a method of characterizing a disease or disorder, comprising: a) contacting a biological test sample with the oligonucleotide or plurality of oligonucleotides provided by the invention; b) detecting a presence or level of complexes formed in step (a) between the oligonucleotide or plurality of oligonucleotides provided by the invention and a target in the biological test sample; and c) comparing the presence or level detected in step (b) to a reference level from a biological control sample, thereby characterizing the disease or disorder. The step of detecting may comprise performing sequencing of all or some of the oligonucleotides in the complexes, amplification of all or some of the oligonucleotides in the complexes, and/or hybridization of all or some of the oligonucleotides in the complexes to an array. The sequencing may be high-throughput or next generation sequencing.

[00438] In the methods of the invention, the biological test sample and biological control sample may each comprise a tissue sample, a cell culture, or a biological fluid. In some embodiments, the biological fluid comprises a bodily fluid. Useful bodily fluids within the method of the invention comprise peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, or umbilical cord blood. In some preferred embodiments, the bodily fluid comprises blood, serum or plasma. The biological fluid may comprise microvesicles. In such case, the complexes may be formed between the oligonucleotide or plurality of oligonucleotides and at least one of the microvesicles.

[00439] The biological test sample and biological control sample may further comprise isolated microvesicles, wherein optionally the microvesicles are isolated using at least one of chromatography, filtration, ultrafiltration, centrifugation, ultracentrifugation, flow cytometry, affinity capture (e.g., to a planar surface, column or bead), polymer precipitation, and using microfluidics. The vesicles can also be isolated after contact with the oligonucleotide or plurality of oligonucleotides.

[00440] In various embodiments of the methods of the invention, the oligonucleotide or plurality of oligonucleotides binds a polypeptide or fragment thereof. The polypeptide or fragment thereof can be soluble or membrane bound, wherein optionally the membrane comprises a microvesicle membrane. The membrane could also be from a cell or a fragment of a cell of vesicle. In some embodiments, the polypeptide or fragment thereof comprises a biomarker in **Table 3**, **Table 4** or any one of **Tables 18-25**. For example, the polypeptide or fragment thereof could be a general vesicle marker such as in **Table 3** or a tissue-related or disease-related marker such as in **Table 4**, or a vesicle associated biomarker provided in any one of **Tables 18-25**. The oligonucleotide or plurality of oligonucleotides may bind a microvesicle surface antigen in the biological sample. For example, the oligonucleotide or plurality of oligonucleotides can be enriched from a naïve library against microvesicles.

[00441] As noted above, the microvesicles may be isolated in whole or in part using polymer precipitation. In an embodiment, the polymer comprises polyethylene glycol (PEG). Any appropriate form of PEG may be used. For example, the PEG may be PEG 8000. The PEG may be used at any appropriate concentration. For example, the PEG can be used at a concentration of 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14% or 15% to isolate the microvesicles. In some embodiments, the PEG is used at a concentration of 6%.

[00442] The invention provides oligonucleotide probes that can be used to carry out the methods herein. See, e.g., **Examples 26-32**. In an aspect, the invention provides an oligonucleotide comprising a sequence according to any one of SEQ ID NOs 137-969 and 1072-4150. In a related aspect, the invention provides an oligonucleotide comprising a sequence according to any one of the SEQ ID NOs in **Table 40**. In another related aspect, the invention provides an oligonucleotide comprising a sequence according to any one of the SEQ ID NOs in the row "2000v1" in **Table 43**. In still another related aspect, the invention provides an oligonucleotide comprising a sequence according to any one of the SEQ ID NOs in the row "2000v2" in **Table 43**. In yet another related aspect, the invention provides an oligonucleotide comprising a sequence according to any one of the SEQ ID NOs in the row "Common" in **Table 43**.

[00443] The oligonucleotides of the invention can comprise flanking regions for various purposes, including without limitation amplification, capture, conjugation or spacing. For example, the invention provides an oligonucleotide comprising a sequence according to any one of the SEQ ID NOs above and further having a 5' region with sequence 5'-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131) and/or a 3' region with sequence 5'-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132).

[00444] The invention further provides oligonucleotides homologous to the SEQ ID NOs above. For example, the invention provides an oligonucleotide comprising a nucleic acid sequence or a portion thereof that is at least 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99 or 100 percent homologous to an oligonucleotide sequence of any one of the SEQ ID NOs above. The homologous sequences may comprise similar properties to the listed sequences, such as similar binding properties.

[00445] In an aspect, the invention provides a plurality of oligonucleotides comprising at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95,

100, 125, 150, 175, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, or at least 10000 different oligonucleotide sequences as described in the paragraphs above. For example, the invention provides a plurality of oligonucleotides comprising member sequences having at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, or all variable regions according to SEQ ID NOs 137-969 and 1072-4150.

[00446] The plurality of oligonucleotides can comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, or all SEQ ID NOs listed in **Table 40**. In an embodiment, the plurality of oligonucleotides comprises at least the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, or SEQ ID NOs listed in **Table 40**.

[00447] The plurality of oligonucleotides can also comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, or all SEQ ID NOs listed in row "2000v1" of **Table 43**. In an embodiment, the plurality of oligonucleotides comprises at least the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, or all SEQ ID NOs listed in row "2000v1" of **Table 43**.

[00448] The plurality of oligonucleotides can comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145 or all SEQ ID NOs listed in row "2000v2" of **Table 43**. In an embodiment, the plurality of oligonucleotides comprises at least the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145 or all SEQ ID NOs listed in row "2000v2" of **Table 43**.

[00449] The plurality of oligonucleotides can also comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or all variable regions listed in row "Common" of **Table 43**. In an embodiment, the plurality of oligonucleotides comprises at least the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or all SEQ ID NOs listed in row "Common" of **Table 43**.

[00450] The oligonucleotide or at least one member of the plurality of oligonucleotides can have least one functional modification selected from the group consisting of DNA, RNA, biotinylation, a non-naturally occurring nucleotides, a deletion, an insertion, an addition, and a chemical modification. Such modifications may provide additional or altered functions to the oligonucleotides, including without limitation capture, detection, stability, or binding properties.

[00451] Such oligonucleotides and plurality of oligonucleotides (pools) can be used to characterize a phenotype as described herein. In an aspect, the invention provides a method of characterizing a phenotype in a sample comprising: (a) contacting the sample with at least one oligonucleotide or plurality of oligonucleotides provided by the invention (see above); and (b) identifying a presence or level of a complex formed between the at least one oligonucleotide or plurality of oligonucleotides and the sample, wherein the presence or level is used to characterize the phenotype. Any useful technique for identifying

can be used according to the invention. In various embodiments, the identifying comprises sequencing, amplification, hybridization, gel electrophoresis or chromatography. In an embodiment, identifying by hybridization comprises contacting the sample with at least one labeled probe that is configured to hybridize with at least one oligonucleotide. The at least one labeled probe can be directly or indirectly attached to a label. Any useful label can be used, including without limitation a fluorescent or magnetic label. In another embodiment, identifying by sequencing comprises next generation sequencing, dye termination sequencing, and/or pyrosequencing.

[00452] In the methods of the invention, the complex formed between the at least one oligonucleotide or the plurality of oligonucleotides and the sample can be a complex formed between a microvesicle population in the sample and the at least one oligonucleotide or plurality of oligonucleotides. The microvesicle population can be isolated in whole or in part from other constituents in the sample before of after the contacting. In embodiments, the isolating uses affinity purification, filtration, polymer precipitation, PEG precipitation, ultracentrifugation, a molecular crowding reagent, affinity isolation, affinity selection, or any combination thereof.

[00453] In the methods of the invention, the phenotype can be any detectable phenotype. In some embodiments, the phenotype comprises a disease or disorder. In such cases, the characterizing can be a diagnosis, prognosis and/or theranosis for the disease or disorder. The theranosis can be any type of therapy-related such as described herein. The theranosis includes without limitation predicting a treatment efficacy or lack thereof, or monitoring a treatment efficacy.

[00454] The characterizing step of the methods of the invention may entail comparing the presence or level to a reference. Any useful reference can be used. In an embodiment wherein the phenotype comprises a disease or disorder, the reference can be the presence or level determined in a sample from an individual without a disease or disorder, or from an individual with a different state of the disease or disorder. In some embodiments, the comparison to the reference of at least one oligonucleotide comprising a sequence having a SEQ ID NO provided above indicates that the sample comprises a cancer sample or a non-cancer/normal sample.

[00455] The disease or disorder detected by the oligonucleotide, plurality of oligonucleotides, or methods provided here may comprise any appropriate disease or disorder of interest, including without limitation Breast Cancer, Alzheimer's disease, bronchial asthma, Transitional cell carcinoma of the bladder, Giant cellular osteoblastoclastoma, Brain Tumor, Colorectal adenocarcinoma, Chronic obstructive pulmonary disease (COPD), Squamous cell carcinoma of the cervix, acute myocardial infarction (AMI) / acute heart failure, Chron's Disease, diabetes mellitus type II, Esophageal carcinoma, Squamous cell carcinoma of the larynx, Acute and chronic leukemia of the bone marrow, Lung carcinoma, Malignant lymphoma, Multiple Sclerosis, Ovarian carcinoma, Parkinson disease, Prostate adenocarcinoma, psoriasis, Rheumatoid Arthritis, Renal cell carcinoma, Squamous cell carcinoma of skin, Adenocarcinoma of the stomach, carcinoma of the thyroid gland, Testicular cancer, ulcerative colitis, or Uterine adenocarcinoma.

[00456] In some embodiments, the disease or disorder comprises a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. The cancer can include without limitation one of acute lymphoblastic leukemia; acute myeloid leukemia; adrenocortical carcinoma; AIDS-related cancers; AIDS-related lymphoma; anal cancer; appendix cancer; astrocytomas; atypical teratoid/rhabdoid tumor; basal cell carcinoma; bladder cancer; brain stem glioma; brain tumor (including brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, astrocytomas, craniopharyngioma, ependymoblastoma, ependymoma, medulloblastoma, medulloepithelioma, pineal parenchymal tumors of intermediate differentiation, supratentorial primitive neuroectodermal tumors and pineoblastoma); breast cancer; bronchial tumors; Burkitt lymphoma; cancer of unknown primary site; carcinoid tumor; carcinoma of unknown primary site; central nervous system atypical teratoid/rhabdoid tumor; central nervous system embryonal tumors; cervical cancer; childhood cancers; chordoma; chronic lymphocytic leukemia; chronic myelogenous leukemia; chronic myeloproliferative disorders; colon cancer; colorectal cancer; craniopharyngioma; cutaneous T-cell lymphoma; endocrine pancreas islet cell tumors; endometrial cancer; ependymoblastoma; ependymoma; esophageal cancer; esthesioneuroblastoma; Ewing sarcoma; extracranial germ cell tumor; extragonadal germ cell tumor; extrahepatic bile duct cancer; gallbladder cancer; gastric (stomach) cancer; gastrointestinal carcinoid tumor; gastrointestinal stromal cell tumor; gastrointestinal stromal tumor (GIST); gestational trophoblastic tumor; glioma; hairy cell leukemia; head and neck cancer; heart cancer; Hodgkin lymphoma; hypopharyngeal cancer; intraocular melanoma; islet cell tumors; Kaposi sarcoma; kidney cancer; Langerhans cell histiocytosis; laryngeal cancer; lip cancer; liver cancer; lung cancer; malignant fibrous histiocytoma bone cancer; medulloblastoma; medulloepithelioma; melanoma; Merkel cell carcinoma; Merkel cell skin carcinoma; mesothelioma; metastatic squamous neck cancer with occult primary; mouth cancer; multiple endocrine neoplasia syndromes; multiple myeloma; multiple myeloma/plasma cell neoplasm; mycosis fungoides; myelodysplastic syndromes; myeloproliferative neoplasms; nasal cavity cancer; nasopharyngeal cancer; neuroblastoma; Non-Hodgkin lymphoma; nonmelanoma skin cancer; non-small cell lung cancer; oral cancer; oral cavity cancer; oropharyngeal cancer; osteosarcoma; other brain and spinal cord tumors; ovarian cancer; ovarian epithelial cancer; ovarian germ cell tumor; ovarian low malignant potential tumor; pancreatic cancer; papillomatosis; paranasal sinus cancer; parathyroid cancer; pelvic cancer; penile cancer; pharyngeal cancer; pineal parenchymal tumors of intermediate differentiation; pineoblastoma; pituitary tumor; plasma cell neoplasm/multiple myeloma; pleuropulmonary blastoma; primary central nervous system (CNS) lymphoma; primary hepatocellular liver cancer; prostate cancer; rectal cancer; renal cancer; renal cell (kidney) cancer; renal cell cancer; respiratory tract cancer; retinoblastoma; rhabdomyosarcoma; salivary gland cancer; Sézary syndrome; small cell lung cancer; small intestine cancer; soft tissue sarcoma; squamous cell carcinoma; squamous neck cancer; stomach (gastric) cancer; supratentorial primitive neuroectodermal tumors; T-cell lymphoma; testicular cancer; throat cancer;

thymic carcinoma; thymoma; thyroid cancer; transitional cell cancer; transitional cell cancer of the renal pelvis and ureter; trophoblastic tumor; ureter cancer; urethral cancer; uterine cancer; uterine sarcoma; vaginal cancer; vulvar cancer; Waldenström macroglobulinemia; or Wilm's tumor. The premalignant condition can include without limitation Barrett's Esophagus. The autoimmune disease can include without limitation one of inflammatory bowel disease (IBD), Crohn's disease (CD), ulcerative colitis (UC), pelvic inflammation, vasculitis, psoriasis, diabetes, autoimmune hepatitis, multiple sclerosis, myasthenia gravis, Type I diabetes, rheumatoid arthritis, psoriasis, systemic lupus erythematosis (SLE), Hashimoto's Thyroiditis, Grave's disease, Ankylosing Spondylitis Sjogrens Disease, CREST syndrome, Scleroderma, Rheumatic Disease, organ rejection, Primary Sclerosing Cholangitis, or sepsis. The cardiovascular disease can include without limitation one of atherosclerosis, congestive heart failure, vulnerable plaque, stroke, ischemia, high blood pressure, stenosis, vessel occlusion or a thrombotic event. The neurological disease can include without limitation one of Multiple Sclerosis (MS), Parkinson's Disease (PD), Alzheimer's Disease (AD), schizophrenia, bipolar disorder, depression, autism, Prion Disease, Pick's disease, dementia, Huntington disease (HD), Down's syndrome, cerebrovascular disease, Rasmussen's encephalitis, viral meningitis, neurospsychiatric systemic lupus erythematosus (NPSLE), amyotrophic lateral sclerosis, Creutzfeldt-Jacob disease, Gerstmann-Straussler-Scheinker disease, transmissible spongiform encephalopathy, ischemic reperfusion damage (e.g. stroke), brain trauma, microbial infection, or chronic fatigue syndrome. The pain can include without limitation one of fibromyalgia, chronic neuropathic pain, or peripheral neuropathic pain. The infectious disease can include without limitation one of a bacterial infection, viral infection, yeast infection, Whipple's Disease, Prion Disease, cirrhosis, methicillin-resistant staphylococcus aureus, HIV, HCV, hepatitis, syphilis, meningitis, malaria, tuberculosis, or influenza. One of skill will appreciate that the oligonucleotide or plurality of oligonucleotides or methods of the invention can be used to assess any number of these or other related diseases and disorders.

[00457] In some embodiments of the invention, the oligonucleotide or plurality of oligonucleotides and methods of use thereof are useful for characterizing certain diseases or disease states. As desired, a pool of oligonucleotides useful for characterizing various diseases is assembled to create a master pool that can be used to probe useful for characterizing the various diseases. One of skill will also appreciate that pools of oligonucleotides useful for characterizing specific diseases or disorders can be created as well. The sequences provided herein can also be modified as desired so long as the functional aspects are still maintained (e.g., binding to various targets or ability to characterize a phenotype). For example, the oligonucleotides may comprise DNA or RNA, incorporate various non-natural nucleotides, incorporate other chemical modifications, or comprise various deletions or insertions. Such modifications may facilitate synthesis, stability, delivery, labeling, etc, or may have little to no effect in practice. In some cases, some nucleotides in an oligonucleotide may be substituted while maintaining functional aspects of the oligonucleotide. Similarly, 5' and 3' flanking regions may be substituted. In still other cases, only a portion of an oligonucleotide may be determined to direct its functionality such that other portions can be

deleted or substituted. Numerous techniques to synthesize and modify nucleotides and polynucleotides are disclosed herein or are known in the art.

[00458] In an aspect, the invention provides a kit comprising a reagent for carrying out the methods of the invention provided herein. In a similar aspect, the invention contemplates use of a reagent for carrying out the methods of the invention provided herein. In embodiments, the reagent comprises an oligonucleotide or plurality of oligonucleotides. The oligonucleotide or plurality of oligonucleotides can be those provided herein. The reagent may comprise various other useful components including without limitation microRNA (miR) and messenger RNA (mRNA)), a protein-nucleic acid complex, and various combinations, fragments and/or complexes of any of these. Theone or more of: a) a reagent configured to isolate a microvesicle, optionally wherein the at least one reagent configured to isolate a microvesicle comprises a binding agent to a microvesicle antigen, a column, a substrate, a filtration unit, a polymer, polyethylene glycol, PEG4000, PEG8000, a particle or a bead; b) at least one oligonucleotide configured to act as a primer or probe in order to amplify, sequence, hybridize or detect the oligonucleotide or plurality of oligonucleotides; and c) a reagent configured to remove one or more abundant protein from a sample, wherein optionally the one or more abundant protein comprises at least one of albumin, immunoglobulin, fibrinogen and fibrin.

[00459] ***Recovery of oligonucleotide probes post-probing***

[00460] As described herein, the oligonucleotide probes of the invention can be used to probe a sample in order to characterize a phenotype. The methods may entail recovering the oligonucleotide probes that bound various biological entities in the sample in order to identify the bound probes. In an aspect, the invention provides a method of detecting at least one oligonucleotide in a sample, comprising: (a) providing the at least one oligonucleotide comprising a capture moiety; (b) contacting the sample with the at least one oligonucleotide provided in (a); (c) capturing the at least one oligonucleotide that formed a complex with a component in the sample in (b); and (d) identifying a presence or level of the at least one oligonucleotide captured in (c), wherein optionally the presence or level is used to characterize a phenotype. *See, e.g.,* **Example 33** and **FIGs. 17A-E**. The at least one oligonucleotide may be captured to a substrate, including without limitation a bead or planar substrate. The capture moiety can be any useful capture moiety, including without limitation a biotin moiety. The capture moiety can be cleavable, e.g., photocleavable or chemically cleavable. In an embodiment, the at least one oligonucleotide is captured to a substrate coupled to avidin or streptavidin. Such configuration is particularly useful when the capture moiety comprises a biotin moiety. In some embodiments, the captured at least one oligonucleotide is released from the substrate by irradiation prior to the identifying. Any useful irradiation, e.g., ultra violet (UV) light may be used. Any useful technique for identifying can be used according to the invention. In various embodiments, the identifying comprises sequencing, amplification, hybridization, gel electrophoresis or chromatography. In an embodiment, identifying by hybridization comprises contacting the sample with at least one labeled probe that is configured to hybridize with at least one oligonucleotide. The at least one labeled probe can be directly or indirectly attached to a label. Any useful label can be

used, including without limitation a fluorescent or magnetic label. In another embodiment, identifying by sequencing comprises next generation sequencing, dye termination sequencing, and/or pyrosequencing. The at least one oligonucleotide can be an oligonucleotide or plurality of oligonucleotides provided by the invention. See e.g., the oligonucleotides and plurality of oligonucleotides described above.

[00461] ***Single strand DNA (ssDNA) library preparation***

[00462] In an embodiment, the invention provides a nucleic acid molecule comprising a 5' leader region which is 5' of a variable region, which is 5' of a tail region, wherein the leader region comprises a lengthener region, a terminator region and a forward primer region, and the tail region comprises a reverse primer region. The nucleic acid molecule may be used for asymmetric or unequal length PCR applications as desired, e.g., to recover ssDNA. *See, e.g.,* **Example 34** and **FIGs. 18A-C**. The lengthener region can be any desired length. In some embodiments, the lengthener region comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides. The lengthener region may comprise a poly-A sequence. Similarly, the terminator region can be any desired length. In some embodiments, the terminator region comprises at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides. The terminator region may comprise a non-nucleotide terminator. For example, the non-nucleotide terminator can be a polymer such as triethylene glycol or the like. The forward primer region can be any desired length. In some embodiments, the forward primer region comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides. The variable region can be any desired length. In some embodiments, the variable region comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides. In some embodiments, the variable region binds a target molecule or complex through non-Watson-Crick base pairing. For example, the variable region may act as an aptamer and bind proteins or other entities. Finally, the reverse primer region can be any desired length. In some embodiments, the reverse primer region comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides.

[00463] In a related aspect, the invention provides a method of generating a single-stranded DNA (ssDNA) molecule comprising: a) providing a mixture comprising a nucleic acid molecule as described in the paragraph above, and forward and reverse primers configured to amplify the nucleic acid molecule from the forward primer region and reverse primer region, respectively; and b) performing asymmetric polymerase chain reaction (PCR) on the mixture in a) to favorably amplify the reverse strand of the nucleic acid molecule, wherein the forward and reverse primers in the mixture are at a ratio of at least about 1:5 (F/R) in favor of the reverse primers; thereby generating the ssDNA molecule. In an embodiment, the ratio is between about 1:20-1:50 (F/R) in favor of the reverse primers. For example, the ratio can be between about 1:37.5 (F/R) in favor of the reverse primers. The method may further comprise

isolating the amplified reverse strand of the nucleic acid molecule on a native gel. The method may also further comprise: c) denaturing the amplified nucleic acid molecules from b); and d) isolating the denatured reverse strand of the nucleic acid molecules from c). In an embodiment, the denatured reverse strand of the nucleic acid molecules is isolated on a denaturing gel. The mixture in a) can comprise additional components as desired. For example, the mixture may further comprise at least one of an enrichment buffer, non-target molecules, proteins, microvesicles, and polyethyleve glycol.

[00464] In a related aspect, the invention provides a kit comprising a reagent for carrying out the methods herein. In still another related aspect, the invention provides for use of a reagent for carrying out the methods. In various embodiements, the reagent comprises at least one of a buffer, a nucleic acid molecule described above, and forward and/or reverse primers configured to amplify the nucleic acid molecule.

[00465] ***Detecting Watson-Crick base pairing with an oligonucleotide probe***

[00466] The oligonucleotide probes provided by the invention can bind via non-Watson Crick base pairing. However, in some cases, the oligonucleotide probes provided by the invention can bind via Watson Crick base pairing. The oligonucleotide probe libraries of the invention, e.g., as described above, can query both types of binding events simultaneously. For example, some oligonucleotide probes may bind the microvesicle protein antigens in the classical aptamer sense, whereas other oligonucleotide probes may bind microvesicles via nucleic acids associated with the microvesicles, e.g., nucleic acid (including without limitation microRNA and mRNA) on the surface of the microvesicles or as payload. Such surface bound nucleic acids can be associated with proteins. For example, they may comprise Argonaute-microRNA complexes. The argonaute protein can be Ago1, Ago2, Ago3 and/or Ago4.

[00467] In addition to the oligonucleotide probe library approach described herein which relies on determining a sequence of the oligonucleotides (e.g., via sequencing, hybridization or amplification), assays can also be designed to detect Watson Crick base pairing. In some embodiments, these approaches rely on Ago2-mediated cleavage wherein an Ago2-microRNA complex can be used to detected using oligonucleotide probes. These approaches can use oligonucleotide probes to detect Ago2-microRNA complexes, which may or may not be associated with microvesicles. The detection can be used to characterize a phenotype as described herein. Such detection can be used along side the sequencing identification methods described herein (e.g., via sequencing, hybridization or amplification).For further details, see PCT/US15/62184, filed November 23, 2015, which application is incorporated by reference herein in its entirety.

Therapeutics

[00468] As used herein "therapeutically effective amount" refers to an amount of a composition that relieves (to some extent, as judged by a skilled medical practitioner) one or more symptoms of the disease or condition in a mammal. Additionally, by "therapeutically effective amount" of a composition is meant an amount that returns to normal, either partially or completely, physiological or biochemical parameters associated with or causative of a disease or condition. A clinician skilled in the art can determine the therapeutically effective amount of a composition in order to treat or prevent a particular disease

condition, or disorder when it is administered, such as intravenously, subcutaneously, intraperitoneally, orally, or through inhalation. The precise amount of the composition required to be therapeutically effective will depend upon numerous factors, e.g., such as the specific activity of the active agent, the delivery device employed, physical characteristics of the agent, purpose for the administration, in addition to many patient specific considerations. But a determination of a therapeutically effective amount is within the skill of an ordinarily skilled clinician upon the appreciation of the disclosure set forth herein.

[00469] The terms "treating," "treatment," "therapy," and "therapeutic treatment" as used herein refer to curative therapy, prophylactic therapy, or preventative therapy. An example of "preventative therapy" is the prevention or lessening the chance of a targeted disease (e.g., cancer or other proliferative disease) or related condition thereto. Those in need of treatment include those already with the disease or condition as well as those prone to have the disease or condition to be prevented. The terms "treating," "treatment," "therapy," and "therapeutic treatment" as used herein also describe the management and care of a mammal for the purpose of combating a disease, or related condition, and includes the administration of a composition to alleviate the symptoms, side effects, or other complications of the disease, condition. Therapeutic treatment for cancer includes, but is not limited to, surgery, chemotherapy, radiation therapy, gene therapy, and immunotherapy.

[00470] As used herein, the term "agent" or "drug" or "therapeutic agent" refers to a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials such as bacteria, plants, fungi, or animal (particularly mammalian) cells or tissues that are suspected of having therapeutic properties. The agent or drug can be purified, substantially purified or partially purified. An "agent" according to the present invention, also includes a radiation therapy agent or a "chemotherapuetic agent."

[00471] As used herein, the term "diagnostic agent" refers to any chemical used in the imaging of diseased tissue, such as, e.g., a tumor.

[00472] As used herein, the term "chemotherapuetic agent" refers to an agent with activity against cancer, neoplastic, and/or proliferative diseases, or that has ability to kill cancerous cells directly.

[00473] As used herein, "pharmaceutical formulations" include formulations for human and veterinary use with no significant adverse toxicological effect. "Pharmaceutically acceptable formulation" as used herein refers to a composition or formulation that allows for the effective distribution of the nucleic acid molecules of the instant invention in the physical location most suitable for their desired activity.

[00474] As used herein the term "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated.

[00475] *Aptamer-Toxin Conjugates as a Cancer Therapeutic*

[00476] Previous work has developed the concept of antibody-toxin conjugates ("immunoconjugates") as potential therapies for a range of indications, mostly directed at the treatment of cancer with a primary focus on hematological tumors. A variety of different payloads for targeted delivery have been tested in pre-clinical and clinical studies, including protein toxins, high potency small molecule cytotoxics, radioisotopes, and liposome-encapsulated drugs. While these efforts have successfully yielded three FDA-approved therapies for hematological tumors, immunoconjugates as a class (especially for solid tumors) have historically yielded disappointing results that have been attributable to multiple different properties of antibodies, including tendencies to develop neutralizing antibody responses to non-humanized antibodies, limited penetration in solid tumors, loss of target binding affinity as a result of toxin conjugation, and imbalances between antibody half-life and toxin conjugate half-life that limit the overall therapeutic index (reviewed by Reff and Heard, Critical Reviews in Oncology/Hematology, 40 (2001):25-35).

[00477] Aptamers are functionally similar to antibodies, except their absorption, distribution, metabolism, and excretion ("ADME") properties are intrinsically different and they generally lack many of the immune effector functions generally associated with antibodies (e.g., antibody-dependent cellular cytotoxicity, complement-dependent cytotoxicity). In comparing many of the properties of aptamers and antibodies previously described, several factors suggest that toxin-delivery via aptamers offers several concrete advantages over delivery with antibodies, ultimately affording them better potential as therapeutics. Several examples of the advantages of toxin-delivery via aptamers over antibodies are as follows:

[00478] 1) Aptamer-toxin conjugates are entirely chemically synthesized. Chemical synthesis provides more control over the nature of the conjugate. For example, the stoichiometry (ratio of toxins per aptamer) and site of attachment can be precisely defined. Different linker chemistries can be readily tested. The reversibility of aptamer folding means that loss of activity during conjugation is unlikely and provides more flexibility in adjusting conjugation conditions to maximize yields.

[00479] 2) Smaller size allows better tumor penetration. Poor penetration of antibodies into solid tumors is often cited as a factor limiting the efficacy of conjugate approaches. See Colcher, D., Goel, A., Pavlinkova, G., Beresford, G., Booth, B., Batra, S. K. (1999) "Effects of genetic engineering on the pharmacokinetics of antibodies," Q. J. Nucl. Med., 43: 132-139. Studies comparing the properties of unPEGylated anti-tenascin C aptamers with corresponding antibodies demonstrate efficient uptake into tumors (as defined by the tumor:blood ratio) and evidence that aptamer localized to the tumor is unexpectedly long-lived ($t_{1/2}$>12 hours) (Hicke, B. J., Stephens, A. W., "Escort aptamers: a delivery service for diagnosis and therapy", J. Clin. Invest., 106:923-928 (2000)).

[00480] 3) Tunable PK. Aptamer half-life/metabolism can be easily tuned to match properties of payload, optimizing the ability to deliver toxin to the tumor while minimizing systemic exposure. Appropriate modifications to the aptamer backbone and addition of high molecular weight PEGs should make it

possible to match the half-life of the aptamer to the intrinsic half-life of the conjugated toxin/linker, minimizing systemic exposure to non-functional toxin-bearing metabolites (expected if $t_{1/2}$(aptamer)<<$t_{1/2}$(toxin)) and reducing the likelihood that persisting unconjugated aptamer will functionally block uptake of conjugated aptamer (expected if $t_{1/2}$(aptamer)>>$t_{1/2}$ (toxin)).

[00481] 4) Relatively low material requirements. It is likely that dosing levels will be limited by toxicity intrinsic to the cytotoxic payload. As such, a single course of treatment will likely entail relatively small (<100 mg) quantities of aptamer, reducing the likelihood that the cost of oligonucleotide synthesis will be a barrier for aptamer-based therapies.

[00482] 5) Parenteral administration is preferred for this indication. There will be no special need to develop alternative formulations to drive patient/physician acceptance.

[00483] The invention provides a pharmaceutical composition comprising a therapeutically effective amount of an aptamer provided by the invention or a salt thereof, and a pharmaceutically acceptable carrier or diluent. The invention also provides a pharmaceutical composition comprising a therapeutically effective amount of the aptamer or a salt thereof, and a pharmaceutically acceptable carrier or diluent. Relatedly, the invention provides a method of treating or ameliorating a disease or disorder, comprising administering the pharmaceutical composition to a subject in need thereof. Administering a therapeutically effective amount of the composition to the subject may result in: (a) an enhancement of the delivery of the active agent to a disease site relative to delivery of the active agent alone; or (b) an enhancement of microvesicles clearance resulting in a decrease of at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% in a blood level of microvesicles targeted by the aptamer; or (c) an decrease in biological activity of microvesicles targeted by the aptamer of at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%. In an embodiment, the biological activity of microvesicles comprises immune suppression or transfer of genetic information. The disease or disorder can include without limitation those disclosed herein. For example, the disease or disorder may comprise a neoplastic, proliferative, or inflammatory, metabolic, cardiovascular, or neurological disease or disorder. See, e.g., section "Phenotypes."

[00484] ***Anti-target and multivalent oligonucleotides***

[00485] As noted above, the target of oligonucleotide probes can be identified. For example, when the target comprises a protein or protein complex (e.g., a nucleoprotein or lipoprotein), identifying the target may comprise use of mass spectrometry (MS), peptide mass fingerprinting (PMF; protein fingerprinting), sequencing, N-terminal amino acid analysis, C-terminal amino acid analysis, Edman degradation, chromatography, electrophoresis, two-dimensional gel electrophoresis (2D gel), antibody array, or immunoassay. Such approaches can be applied to identify a number of targets recognized by an oligonucleotide probe library. For example, an oligonucleotide probe library can be incubated with a sample of interest, bound members of the library captured, and the targets bound to the captured members identified. *See* **Example 23** herein for an example of such target identification using mass spectrometry.

[00486] The oligonucleotide aptamers to the various targets can be used for multiple purposes. In some embodiments, the aptamers are used as therapeutic agents. Immunotherapeutic approaches using antibodies that recognize foreign/misfolded antigens (*e,g.*, anti-CD20, anti-CD30, anti-CD33, anti-CD52, anti-EGFR, anti-nucleolin, anti-nucleophosmin, etc.) can selectively kill target cells via linked therapeutic agents or by stimulating the immune system through activation of cell-mediated cytotoxicity. Aptamers or oligonucleotides are an attractive immunotherapeutic alternative for various reasons such as low cost, small size, ease and speed of synthesis, stability and low immunogenicity. In an embodiment, immunotherapeutic agents are conjugated to disease specific target oligonucleotide or antibody (Ab) for targeted cell killing via recruitment of complement proteins and the downstream membrane attack complex. *See, e.g.,* Zhou and Rossi, Cell-type-specific, Aptamer-functionalized Agents for Targeted Disease Therapy, Mol Ther Nucleic Acids. 2014 Jun 17;3:e169. doi: 10.1038/mtna.2014.21; Pei et al., Clinical applications of nucleic acid aptamers in cancer, Mol Clin Oncol. 2014 May;2(3):341-348. Epub 2014 Feb 10. This approach can be applied to target diseased host cells such as cancer cells, gram negative bacteria, viral and/or parasitic infections, and the like.

[00487] In some embodiments, the invention provides a multipartite construct comprising a binding agent specific to a biological target with another binding agent specific to immunomodulatory entity. Examples of such constructs are shown in **FIG. 23A**. In Design 1 in the figure, the horizontal line indicates an oligonucleotide construct, which construct comprises a 5' primer **2301** (Primer 1), a variable region **2302** that can be an aptamer to a target of interest, a 3' primer **2303** (Primer 2), and an immunomodulatory domain region ("IMD") **2304**. The complete Design 1 construct can be used to bring a target of interest in proximity with an immunomodulatory agent. The primers can be designed for any desired purpose, e.g., amplification, capture, modification, direct or indirect labeling, and the like. In some embodiments, the target of the variable region is a disease marker and thus the construct is targeted to a disease cell or microvesicle. The immunomodulatory domain region can act as an immune stimulator or suppressor. Any appropriate immune stimulator or suppressor can be used, e.g., a small molecule, antibody or an aptamer. Thus, the construct can modulate the immune response at a target of interest, e.g., at a cell or microvesicle carrying the target. The basic construct can be modified as desired. For example, Design 2 in **FIG. 23A** shows the construct carrying a linker **2305** between Primer 2 **2303** and the IMD **2304**. Such linkers are explained further below and can be inserted between any components of the construct as desired. Linkers can provide a desired space between the regions of the construct and can be manipulated to influence other properties such as stability. Design 3 in **FIG. 23A** shows another example wherein the IMD **2304** is an oligonucleotide and the variable region **2302** and IMD **2304** lie between the primers **2301** and **2303**. One of skill will appreciate that one or more linker, such as **2305** of Design 2, can also be inserted into Design 3, e.g., between the variable region **2302** and IMD **2304**. One of skill will further appreciate that the ordering of the oligonucleotide segments from 5' to 3' can be modified, e.g., reversed. As a concrete example which will be described further below, **FIG. 23B** illustrates Design 1 and Design 2 from **FIG. 23A** wherein the variable region comprises an anti-CD20 oligonucleotide **2311** and the IMD comprises an

anti-C1q oligonucleotide **2312**, e.g., an oligonucleotide provided herein. *See, e.g.,* **Example 35**. This constructs of **FIG. 23B** can be used used to target a CD20+ cell population and stimulate C1q mediated cell killing.

[00488] As noted, the multipartite constructs may be synthesized and/or modified as desired. In some embodiments of the invention, the multipartite oligonucleotide construct is synthesized directly with or without a linker in between the oligonucleotide segments. *See, e.g.,* **FIG. 23A** Design 3, which can be generated directly via amplification by Primer 1 **2301** and Primer 2 **2303**. One or more linker can act as a spacer to create a desired spacing between the target of the variable region segment **2302** and the target of the IMD segment **2304**. The spacing can be determined via computer modeling or via experimentation due to steric hindrance or other considerations. Following the example of **FIG. 23B**, the type and size of the linker may be dependent upon steric hindrance between the CD20 target protein and the C1q protein/MAC complex.

[00489] The multipartite constructs can be generated against any appropriate target. The targets can include without limitation diseased cells, cancer cells, circulating tumor cells (CTCs), immune cells (e.g., B-cells, T-cells, macrophages, dendritic cells), microvesicles, bacteria, viruses or other parasites. The target can be large biological complexes, e.g., protein complexes, ribonucleoprotein complexes, lipid complexes, or a combination thereof. It will be understood that the specific target of the multipartite constructs can be a certain member of the foregoing macromolecular targets. For example, consider that the desired target of the multipartite construct is a cell or microvesicle. In such case, the multipartite construct can be directed to a specific biomarker, e.g., a surface antigen, of the cell or microvesicle. As a non-limiting example, the target of interest can be B-cells and the specifc target of the variable region of the multipartite construct can be CD20. CD20 is a cellular marker of B-cells targeted by the monoclonal antibodies (mAb) rituximab, obinutuzumab, ofatumumab, ibritumomab tiuxetan, and tositumomab, which are used as agents in the treatment of B-cell lymphomas and leukemias. As another non-limiting example, the target of interest can be cancer cells and the specifc target of the variable region of the multipartite construct can be c-MET. MET is a membrane receptor that is essential for embryonic development and wound healing. Abnormal MET activation in cancer correlates with poor prognosis, where aberrantly active MET triggers tumor growth, formation of new blood vessels (angiogenesis), and cancer spread to other organs (metastasis). MET has been observed to be deregulated in many types of human malignancies, including cancers of kidney, liver, stomach, breast, and brain. *See* **FIG. 23D** for illustration (discussed further below). Other biomarkers can be used as the specifc target as desired. For example, the biomarker can be selected from **Table 3**, **Table 4**, **Tables 18-26**, or **Table 45** herein. See **FIG. 23C**, which illustrates a construct of the invention **2331** having a segment that recognizes a biomarker **2332** ("Marker of Interest") on a cell or vesicle surface **2333** ("Membrane"), and another segment **2334** that attracts an immune response ("Complement"). The construct **2331** can be such as in **FIGs. 23A-B** or any other desired configuration. Binding of such a construct to a target can cause a complement cascade and induce apoptosis.

[00490] In some embodiments of the invention, the target biomarker is selected from the group consisting of CD19, CD20, CD21, CD22 (also known as LL2), CDIM, and Lym-1. The target biomarker can be a membrane associated protein. In embodiments, the membrane associated protein is selected from the group consisting of CD4, CD19, DC-SIGN/CD209, HIV envelope glycoprotein gp120, CCR5, EGFR/ErbB1, EGFR2/ErbB2/HER2, EGFR3/ErbB3, EGFR4/ErbB4, EGFRvIII, Transferrin Receptor, PSMA, VEGF, VEGF-2, CD25, CD11a, CD33, CD20, CD3, CD52, CEA, TAG-72, LDL receptor, insulin receptor, megalin receptor, LRP, mannose receptor, P63/CKAP4 receptor, arrestin, ASGP, CCK-B, HGFR, RON receptor, FGFR, ILR, AFP, CA125/MUC16, PDGFR, stem cell factor receptor, colony stimulating factor-1 receptor, integrins, TLR, BCR and BAFF-R. The target biomarker can also be a cellular receptor selected from the group consisting of: nucleolin, human epidermal growth factor receptor 2 (HER2), CD20, a transferrin receptor, an asialoglycoprotein receptor, a thyroid-stimulating hormone (TSH) receptor, a fibroblast growth factor (FGF) receptor, CD3, the interleukin 2 (IL-2) receptor, a growth hormone receptor, an insulin receptor, an acetylcholine receptor, an adrenergic receptor, a vascular endothelial growth factor (VEGF) receptor, a protein channel, cadherin, a desmosome, and a viral receptor. In various embodiments, the target biomarker is a cell surface molecule selected from the group consisting of IgM, IgD, IgG, IgA, IgE, CD19, CD20, CD21, CD22, CD24, CD40, CD72, CD79a, CD79b, CD1d, CD5, CD9, CD10, CD1d, CD23, CD27, CD38, CD48, CD80, CD86, CD138, CD148, and combinations thereof. The target biomarker can be a lymphocyte-directing target such as one or moreT-cell receptor motifs, T-cell α chains, T-cell β chains, T-cell γ chains, T-cell Δ chains, CCR7, CD3, CD4, CD5, CD7, CD8, CD11b, CD11c, CD16, CD19, CD20, CD21, CD22, CD25, CD28, CD34, CD35, CD40, CD45RA, CD45RO, CD52, CD56, CD62L, CD68, CD80, CD95, CD117, CD127, CD133, CD137 (4-1 BB), CD163, F4/80, IL-4Ra, Sca-1, CTLA-4, GITR, GARP, LAP, granzyme B, LFA-1, or transferrin receptor.

[00491] In some embodiments, the target biomarker comprises a growth factor, vascular endothelial growth factor (VEGF), TGF, TGFβ, PDGF, IGF, FGF, cytokine, lymphokine, hematopoietic factor, M-CSR, GM-CSF, TNF, interleukin, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-1 3, IL-14, IL-15, IL-16, IL-17, IL18, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, GM-CSF, thrombopoietin, stem cell factor, erythropoietin, hepatocyte growth factor/NK1, angiogenic factor, angiopoietin, Ang-1, Ang-2, Ang-4, Ang-Y, human angiopoietin-like polypeptide, angiogenin, morphogenic protein-1, bone morphogenic protein receptor, bone morphogenic protein receptor IA, bone morphogenic protein receptor IB, neurotrophic factor, chemotactic factor, CD proteins, CD3, CD4, CD8, CD19, CD20, erythropoietin, osteoinductive factors, immunotoxin, bone morphogenetic protein (BMP), interferon, interferon-alpha, interferon-beta, interferon-gamma, colony stimulating factor (CSF), M-CSF, GM-CSF, G-CSF, superoxide dismutase, T-cell receptor; surface membrane protein, decay accelerating factor, viral antigen, portion of the AIDS envelope, transport protein, homing receptor, addressin, regulatory protein, integrin, CD11a, CD11b, CD11c, CD18, ICAM, VLA-4, VCAM, tumor associated

antigen, HER2, HER3, HER4, nucleophosmin, a heterogeneous nuclear ribonucleoproteins (hnRNPs), fibrillarin; or fragments or variants thereof.

[00492] In still other embodiments, the target biomarker is selected from the group consisting of epidermal growth factor receptor, transferrin receptor, platelet-derived growth factor receptor, Erb-B2, CD 19, CD20, CD45, CD52, Ep-CAM, alpha ([alpha])-fetoprotein, carcinoembryonic antigen peptide-1, caspase-8, CDC27, CDK4, carcino-embryonic antigen, calcium-activated chloride channel-2, cyclophilin B, differentiation antigen melanoma, elongation factor 2, Ephrin type-A receptor 2, 3, Fibroblast growth factor-5, fibronectin, glycoprotein 250, G antigen, N-acetylglucosaminyltransferase V, glycoprotein 100 kD, helicase antigen, human epidermal receptor-2/neurological, heat shock protein 70-2 mutated, human signet ring tumor-2, human telomerase reverse transcriptase, intestinal carboxyl esterase, interleukin 13 receptor [alpha]2 chain, [beta]-D-galactosidase 2-[alpha]-L-fucosyltransferase, melanoma antigen, melanoma antigen recognized by T cells-1/Melanoma antigen A, melanocortin 1 receptor, macrophage colony-stimulating factor, mucin 1, 2, melanoma ubiquitous mutated 1, 2, 3, New York-esophageous 1, ocular albinism type 1 protein, O-linked N-acetyl glucosamine transferase gene, protein 15, promyelocytic leukemia/retinoic acid receptor [alpha], prostate-specific antigen, prostate-specific membrane antigen, receptor-type protein-tyrosinephosphatase kappa, renal antigen, renal ubiquitous 1, 2, sarcoma antigen, squamous antigen rejecting tumor 1, 2, 3, synovial sarcoma, Survivin-2B, synaptotagmin I/synovial sarcoma, X fusion protein, translocation Ets-family leukemia/acute myeloid leukemia 1, transforming growth factor [beta] receptor 2, triosephosphate isomerase, taxol resistant associated protein 3, testin-related gene, tyrosinase related protein 1, and tyrosinase related protein 2.

[00493] The target biomarker can be a cancer-associated or tumor associated antigen. The cancer-associated antigen may include without limitation one or more of human Her2/neu, Her1/EGF receptor (EGFR), HER2 (ERBB2), Her3, Her4, A33 antigen, B7H3, CD5, CD19, CD20, CD22, CD23 (IgE Receptor), C242 antigen, 5T4, IL-6, IL-13, vascular endothelial growth factor VEGF (e.g., VEGF-A), VEGFR-1, VEGFR-2, CD30, CD33, CD37, CD40, CD44, CD51, CD52, CD56, CD74, CD80, CD152, CD200, CD221, CCR4, HLA-DR, CTLA-4, N PC-1C, tenascin, vimentin, insulin-like growth factor 1 receptor (IGF-1R), alpha-fetoprotein, insulin-like growth factor 1 (IGF-1), carbonic anhydrase 9 (CA-IX), carcinoem bryonic antigen (CEA), integrin αvβ3, integrin α5βι, folate receptor 1, transmembrane glycoprotein NMB, fibroblast activation protein alpha (FAP), glypican 1, glypican 3, glycoprotein 75, TAG-72, MUC1, MUC16 (also known as CA-125), phosphatidylserine, prostate-specific membrane antigen (PMSA), NR-LU-13 antigen, TRAIL-R1, tumor necrosis factor receptor superfamily member 10b (TNFRSF10B or TRAIL-R2), SLAM family member 7 (SLAM F7), EGP40 pancarcinoma antigen, B-cell activating factor (BAFF), platelet- derived growth factor receptor, glycoprotein EpCAM (17-1A), Programmed Death-1 (PD1), Programmed Death Ligand 1 (PD-L1), protein disulfide isomerase (PDI), Phosphatase of Regenerating Liver 3 (PRL-3), prostatic acid phosphatase, Lewis-Y antigen, GD2 (a disialoganglioside expressed on tumors of neuroectodermal origin), or mesothelin. For example, the target can be one or more of human Her2/neu, Herl/EGFR, TNF-a, B7H3 antigen, CD20, VEGF, CD52, CD33,

CTLA-4, tenascin, alpha-4 (α4) integrin, IL-23, amyloid-β, Huntingtin, CD25, nerve growth factor (NGF), TrkA, and α-synuclein. In some embodiments, the target biomarker is a tumor antigen selected from the group consisting of PSMA, BRCA1, BRCA2, alpha-actinin-4, BCR-ABL fusion protein (b3a2), CASP-8, β-catenin, Cdc27, CDK4, dek-can fusion protein, Elongation factor 2, ETV6-AML1 fusion protein, LDLR-fucosyltransferase AS fusion protein, hsp70-2, KIAAO205, MART2, MUM-1f, MUM-2, MUM-3, neo-PAP, Myosin class I, OS-9g, pml-RAR alpha fusion protein, PTPRK, K-ras, N-ras, CEA, gp100/Pmel17, Kallikrein 4, mammaglobin-A, Melan-A/MART-1, PSA, TRP-1/gp75, TRP-2, tyrosinase, CPSF, EphA3, G250/MN/CAIX, HER-2/neu, Intestinal carboxyl esterase, alpha-fetoprotein, M-CSF, MUC1, p53, PRAME, RAGE-1, RU2AS, survivin, Telomerase, WT1, or CA125. In still other embodiments, the target biomarker is a tumor antigen selected from the group consisting of 4-1BB, 5T4, AGS-5, AGS-16, Angiopoietin 2, B7.1, B7.2, B7DC, B7H1, B7H2, B7H3, BT-062, BTLA, CAIX, Carcinoembryonic antigen, CTLA4, Cripto, ED-B, ErbBl, ErbB2, ErbB3, ErbB4, EGFL7, EpCAM, EphA2, EphA3, EphB2, EphB3, FAP, Fibronectin, Folate Receptor, Ganglioside GM3, GD2, glucocorticoid-induced tumor necrosis factor receptor (GITR), gp100, gpA33, GPNMB, ICOS, IGFIR, Integrin av, Integrin αvβ, KIR, LAG-3, Lewis Y, Mesothelin, c-MET, MN Carbonic anhydrase IX, MUC1, MUC16, Nectin-4, NKGD2, NOTCH, OX40, OX40L, PD-1, PDL1, PSCA, PSMA, RANKL, RORl, ROR2, SLC44A4, Syndecan-1, TACI, TAG-72, Tenascin, TIM3, TRAILRl, TRAILR2,VEGFR-1, VEGFR-2, VEGFR-3, and variants thereof. In still other embodiments, the target biomarker is a tumor-associated antigen selected from the group consisting of Lewis Y, Muc-1, erbB-2,-3 and-4, Ep-CAM, EGF-receptor (e.g., EGFR type I or EGFR type II), EGFR deletion neoepitope, CA19-9, Muc-1, LeY, TF-, Tn-and sTn-antigen, TAG-72, PSMA, STEAP, Cora antigen, CD7, CD19 and CD20, CD22, CD25, Ig-α and Ig-β, A33 and G250, CD30, MCSP and gp100, CD44-v6, MT-MMPs, (MIS) receptor type II, carboanhydrase 9, F19-antigen, Ly6, desmoglein 4, PSCA, Wue-1, GD2 and GD3 as well as TM4SF-antigens (CD63, L6, CO-29, SAS) and the alpha and/or gamma subunit of the fetal type acetylcholinreceptor (AChR). The target biomarker can be a cancer antigen selected from A33, BAGE, Bcl-2, β-catenin, CA125, CA19-9, CD5, CD19, CD20, CD21, CD22, CD33, CD37, CD45, CD123, CEA, c-Met, CS-1, cyclin B1, DAGE, EBNA, EGFR, ephrinB2, estrogen receptor, FAP, ferritin, folate-binding protein, GAGE, G250, GD-2, GM2, gp75, gp100 (Pmel 17), HER-2/neu, HPV E6, HPV E7, Ki-67, LRP, mesothelin, p53, PRAME, progesterone receptor, PSA, PSMA, MAGE, MART, mesothelin, MUC, MUM-1 -B, myc, NYESO-1, ras, RORI, survivin, tenascin, TSTA tyrosinase, VEGF, and WT1. The target biomarker can also be a tumor antigen selected from carcinoembryonic antigen (CEA), alpha-fetoprotein (AFP), prostate specific antigen (PSA), prostate specific membrane antigen (PSMA), CA- 125 (epithelial ovarian cancer), soluble Interleukin-2 (IL-2) receptor, RAGE-1, tyrosinase, MAGE-1, MAGE-2, NY-ESO-1, Melan- A/MART- 1, glycoprotein (gp) 75, gp100, beta-catenin, PRAME, MUM-1, ZFP161, Ubiquilin-1, HOX-B6, YB-1, Osteonectin, ILF3, or IGF-1. In some embodiments, the cancer-related antigen is one or more of CD2, CD4, CD19, CD20, CD22, CD23, CD30, CD33, CD37, CD40, CD44v6, CD52, CD56, CD70, CD74, CD79a, CD80, CD98, CD138, EGFR (Epidermal growth factor

receptor), VEGF (Vascular endothelial growth factor), VEGFRI (Vascular endothelial growth factor receptor I), PDGFR (Platelet-derived growth factor receptor), RANKL (Receptor activator of nuclear factor kappa-B ligand), GPNMB (Transmembrane glycoprotein Neuromedin B), EphA 2 (Ephrin type-A receptor 2), PSMA (Prostate-specific membrane antigen), Cripto (Cryptic family protein 1B), EpCAM (Epithelial cell adhesion molecule), CTLA 4 (Cytotoxic T-Lymphocyte Antigen 4), IGF- IR (Type 1 insulin-like growth factor receptor), GP3 (M13 bacteriophage), GP9 (Glycoprotein IX (platelet), CD42a, GP 40 (Glycoprotein 40kDa), GPC3 (glypican-3), GPC1 (glypican-1), TRAILR1 (Tumor necrosis factor-related apoptosis-inducing ligand receptor 1), TRAILRII (Tumor necrosis factor-related apoptosis-inducing ligand receptor II), FAS (Type II transmembrane protein), PS (phosphatidyl serine) lipid, Gal GalNac Gal N-linked, Muc1 (Mucin 1, cell surface associated, PEM), Muc18, CD146, A5B1 integrin ($\alpha 5\beta 1$), $\alpha 4\beta 1$ integrin, αv integrin (Vitronectin Receptor), Chondrolectin, CAIX (Carbonic anhydrase IX, gene G250/MN-encoded transmembrane protein), GD2 gangloside, GD3 gangloside, GM1 gangloside, Lewis Y, Mesothelin, HER2 (Human Epidermal Growth factor 2), HER3, HER4, FN14 (Fibroblast Growth Factor Inducible 14), CS1 (Cell surface glycoprotein, CD2 subset 1, CRACC, SLAMF7, CD319), 41BB CD137, SIP (Siah-1 Interacting Protein), CTGF (Connective tissue growth factor), HLADR (MHC class II cell surface receptor), PD-1 (Programmed Death 1, Type I membrane protein, PD-L1 (Programmed Death Ligand 1), PD-L2 (Programmed Death Ligand 2), IL-2 (Interleukin-2), IL-8 (Interleukin-8), IL-13 (Interleukin-13), PIGF (Phosphatidylinositol-glycan biosynthesis class F protein), NRP1 (Neuropilin-1), ICAM1, CD54, GC182 (Claudin 18.2), Claudin, HGF (Hepatocyte growth factor), CEA (Carcinoembryonic antigen), LTβR (lymphotoxin β receptor), Kappa Myeloma, Folate Receptor alpha, GRP78 (BIP, 78 kDa Glucose-regulated protein), A33 antigen, PSA (Prostate-specific antigen), CA 125 (Cancer antigen 125 or carbohydrate antigen 125), CA19.9, CA15.3, CA242, leptin, prolactin, osteopontin, IGF- II (Insulin-like growth factor 2), fascin, sPIgR (secreted chain of polymorphic immunoglobulin receptor), 14-3-3 protein eta, 5T4 oncofetal protein, ETA (epithelial tumor antigen), MAGE (Melanoma-associated antigen), MAPG (Melanoma-associated proteoglycan, NG2), vimentin, EPCA-1 (Early prostate cancer antigen-2), TAG-72 (Tumor-associated glycoprotein 72), factor VIII, Neprilysin (Membrane metallo-endopeptidase) and 17-1 A (Epithelial cell surface antigen 17-1A). The cancer antigen can be selected from the group consisting of carbonic anhydrase IX, alpha-fetoprotein, A3, antigen specific for A33 antibody, Ba 733, BrE3-antigen, CA125, CD1, CD1a, CD3, CD5, CD15, CD16, CD19, CD20, CD21, CD22, CD23, CD25, CD30, CD33, CD38, CD45, CD74, CD79a, CD80, CD138, colon-specific antigen-p (CSAp), CEA (CEACAM5), CEACAM6, CSAp, EGFR, EGP-1, EGP-2, Ep-CAM, Flt-1, Flt-3, folate receptor, HLA-DR, human chorionic gonadotropin (HCG) and its subunits, HER2/neu, hypoxia inducible factor (HIF-1), Ia, IL-2, IL-6, IL-8, insulin growth factor-1 (IGF-1), KC4-antigen, KS-1-antigen, KS1-4, Le-Y, macrophage inhibition factor (MIF), MAGE, MUC1, MUC2, MUC3, MUC4, MUC16, NCA66, NCA95, NCA90, antigen specific for PAM-4 antibody, placental growth factor, p53, prostatic acid phosphatase, PSA, PSMA, RS5, S100, TAC, TAG-72, tenascin, TRAIL

receptors, Tn antigen, Thomson-Friedenreich antigens, tumor necrosis antigens, VEGF, ED-B fibronectin, 17-1A-antigen, an angiogenesis marker, an oncogene marker and an oncogene product.

[00494] The tumor marker can be a generic tumor marker or be associated with certain tumor types, such as those originating from different anatomical origins. In an embodiment, the tumor marker can be chosen to correspond to a certain tumor type. For example, exexmplary tumor markers and associated tumor types include without limitation the following, listed as antigen (optional name) (cancer types): Alpha fetoprotein (AFP) (germ cell tumor, hepatocellular carcinoma); CA15-3 (breast cancer); CA27-29 (breast cancer); CA19-9 (mainly pancreatic cancer, but also colorectal cancer and other types of gastrointestinal cancer); CA-125 (ovarian cancer, endometrial cancer, fallopian tube cancer, lung cancer, breast cancer and gastrointestinal cancer); Calcitonin (medullary thyroid carcinoma); Calretinin (mesothelioma, sex cord-gonadal stromal tumour, adrenocortical carcinoma, synovial sarcoma); Carcinoembryonic antigen (gastrointestinal cancer, cervix cancer, lung cancer, ovarian cancer, breast cancer, urinary tract cancer); CD34 (hemangiopericytoma/solitary fibrous tumor, pleomorphic lipoma, gastrointestinal stromal tumor, dermatofibrosarcoma protuberans); CD99 (MIC2) (Ewing sarcoma, primitive neuroectodermal tumor, hemangiopericytoma/solitary fibrous tumor, synovial sarcoma, lymphoma, leukemia, sex cord-gonadal stromal tumour); CD117 (gastrointestinal stromal tumor, mastocytosis, seminoma); Chromogranin (neuroendocrine tumor); Chromosomes 3, 7, 17, and 9p21 (bladder cancer); Cytokeratin (various types) (various carcinoma, some types of sarcoma); Desmin (smooth muscle sarcoma, skeletal muscle sarcoma, endometrial stromal sarcoma); Epithelial membrane antigen (EMA) (many types of carcinoma, meningioma, some types of sarcoma); Factor VIII (CD31, FL1) (vascular sarcoma); Glial fibrillary acidic protein (GFAP) (glioma (astrocytoma, ependymoma)); Gross cystic disease fluid protein (GCDFP-15) (breast cancer, ovarian cancer, salivary gland cancer); HMB-45 (melanoma, PEComa (for example angiomyolipoma), clear cell carcinoma, adrenocortical carcinoma); Human chorionic gonadotropin (hCG) (gestational trophoblastic disease, germ cell tumor, choriocarcinoma); Immunoglobulin (lymphoma, leukemia); Inhibin (sex cord-gonadal stromal tumour, adrenocortical carcinoma, hemangioblastoma); keratin (various types) (carcinoma, some types of sarcoma); lymphocyte marker (various types, lymphoma, leukemia); MART-1 (Melan-A) (melanoma, steroid-producing tumors (adrenocortical carcinoma, gonadal tumor)); Myo D1 (rhabdomyosarcoma, small, round, blue cell tumour); muscle-specific actin (MSA) (myosarcoma (leiomyosarcoma, rhabdomyosarcoma); neurofilament (neuroendocrine tumor, small-cell carcinoma of the lung); neuron-specific enolase (NSE) (neuroendocrine tumor, small-cell carcinoma of the lung, breast cancer); placental alkaline phosphatase (PLAP) (seminoma, dysgerminoma, embryonal carcinoma); prostate-specific antigen (prostate); PTPRC (CD45) (lymphoma, leukemia, histiocytic tumor); S100 protein (melanoma, sarcoma (neurosarcoma, lipoma, chondrosarcoma), astrocytoma, gastrointestinal stromal tumor, salivary gland cancer, some types of adenocarcinoma, histiocytic tumor (dendritic cell, macrophage)); smooth muscle actin (SMA) (gastrointestinal stromal tumor, leiomyosarcoma, PEComa); synaptophysin (neuroendocrine tumor); thyroglobulin (thyroid cancer but not typically medullary thyroid cancer); thyroid transcription factor-1

(all types of thyroid cancer, lung cancer); Tumor M2-PK (colorectal cancer, Breast cancer, renal cell carcinoma, Lung cancer, Pancreatic cancer, Esophageal Cancer, Stomach Cancer, Cervical Cancer, Ovarian Cancer); Vimentin (sarcoma, renal cell carcinoma, endometrial cancer, lung carcinoma, lymphoma, leukemia, melanoma). Additional tumor types and associated biomarkers comprise the following, listed as tumor type (markers): Colorectal (M2-PK, CEA, CA 19-9, CA 125); Breast (CEA, CA 15-3, Cyfra 21-1); Ovary (CEA, CA 19-9, CA 125, AFP, BHCG); Uterine (CEA, CA 19-9, CA 125, Cyfra 21-1, SCC); Prostate (PSA); Testicle (AFP, BHCG); Pancreas/Stomach (CEA, CA 19-9, CA 72-4); Liver (CEA, AFP); Oesophagus (CEA, Cyfra 21-1); Thyroid (CEA, NSE); Lung (CEA, CA 19-9, CA 125, NSE, Cyfra 21-1); Bladder (CEA, Cyfra 21-1, TPA). One or more of these markers can be used as the target biomarker recognized by the variable region of the multipartite construct of the invention.

[00495] In some embodiments of the invention, the target biomarker recognized by the variable region comprises one or more of PDGF, IgE, IgE Fcε R1, PSMA, CD22, TNF-alpha, CTLA4, PD-1, PD-L1, PD-L2, FcRIIB, BTLA, TIM-3, CD11c, BAFF, B7-X, CD19, CD20, CD25, and CD33. The target biomarker can also be a protein comprising one or more of insulin-like growth factor 1 receptor (IGF1R), IGF2R, insulin-like growth factor (IGF), mesenchymal epithelial transition factor receptor (c-met), hepatocyte growth factor (HGF), epidermal growth factor receptor (EGFR), ErbB2, ErbB3, epidermal growth factor (EGF), heregulin, fibroblast growth factor receptor (FGFR), platelet-derived growth factor receptor (PDGFR), platelet-derived growth factor (PDGF), vascular endothelial growth factor receptor (VEGFR), vascular endothelial growth factor (VEGF), tumor necrosis factor receptor (TNFR), tumor necrosis factor alpha (TNF-a), folate receptor (FOLR), folate, transferrin receptor (TfR), mesothelia, Fc receptor, c-kit receptor, c-kit, a4 integrin, P-selectin, sphingosine-1-phosphate receptor-1 (S1PR), hyaluronate receptor, leukocyte function antigen-1 (LFA-1), CD4, CD11, CD18, CD20, CD25, CD27, CD52, CD70, CD80, CD85, CD95 (Fas receptor), CD106 (vascular cell adhesion molecule 1 (VCAM1)), CD166 (activated leukocyte cell adhesion molecule (ALCAM)), CD 178 (Fas ligand), CD253 (TNF-related apoptosis-inducing ligand (TRAIL)), inducible costimulator (ICOS) ligand, CCR2, CXCR3, CCR5, CXCL12 (stromal cell-derived factor 1 (SDF-1)), interleukin 1 (IL-1), cytotoxic T-lymphocyte antigen 4 (CTLA-4), MART-1, gp100, MAGE-1, ephrin (Eph) receptor, mucosal addressin cell adhesion molecule 1 (MAdCAM-1), carcinoembryonic antigen (CEA), LewisY, MUC-1, epithelial cell adhesion molecule (EpCAM), cancer antigen 125 (CA125), prostate specific membrane antigen (PSMA), TAG-72 antigen, and fragments thereof. In various embodiments, the target biomarker comprises one or more of PSMA, PSCA, e selectin, an ephrin, ephB2, cripto-1, TENB2 (TEMFF2), ERBB2 receptor (HER2), MUC1, CD44v6, CD6, CD19, CD20, CD22, CD23, CD25, CD30, CD33, CD56, IL-2 receptor, HLA-DR10 B subunit, EGFR, CA9, caveolin-1 and nucleolin.

[00496] The target biomarker can be a microvesicle antigen, such as a microvesicle antigen selected from any of **Table 3**, **Table 4**, **Tables 18-26**, or **Table 45**. For example, the target biomarker can be one or more microvesicle antigen selected from CD9, EphA2, EGFR, B7H3, PSMA, PCSA, CD63, STEAP, CD81, B7H3, STEAP1, ICAM1 (CD54), A33, DR3, CD66e, MFG-e8, Hepsin, TMEM211, TROP-2,

EGFR, Mammoglobin, Hepsin, NPGP/NPFF2, PSCA, 5T4, NGAL, NK-2, EpCam, NK-1R, 5T4, PAI-1, and CD45. The target biomarker can be one or more microvesicle antigen selected from SPB, SPC, NSE, PGP9.5, CD9, P2RX7, NDUFB7, NSE, Gal3, Osteopontin, CHI3L1, EGFR, B7H3, iC3b, MUC1, Mesothelin, SPA, TPA, PCSA, CD63, AQP5, DLL4, CD81, DR3, PSMA, GPCR 110 (GPRl 10), EPHA2, CEACAM, PTP, CABYR, TMEM211, ADAM28, UNC93a, A33, CD24, CD10, NGAL, EpCam, MUC17, TROP2 and MUC2. In some embodiments, the target biomarker comprises one or more microvesicle antigen selected from CD9, CD63, CD81, B7H3, PRO GRP, CYTO 18, FTH1, TGM2, CENPH, ANNEXIN I, ANNEXIN V, ERBB2, EGFR, CRP, VEGF, CYTO 19, CCL2, Osteopontin (OST19), Osteopontin (OST22), BTUB, CD45, TIMP, NACC1, MMP9, BRCAl, P27, NSE, M2PK, HCG, MUCl, CEA, CEACAM, CYTO 7, EPCAM, MS4A1, MUC1, MUC2, PGP9, SPA, SPA, SPD, P53, GPCR (GPR110), SFTPC, UNCR2, NSE, INGA3, INTO b4, MMP1, PNT, RACK1, NAP2, HLA, BMP2, PTH1R, PAN ADH, NCAM, CD151, CKS1, FSHR, HIF, KRAS, LAMP2, SNAIL, TRIM29, TSPAN1, TWIST1, ASPH and AURKB. In another embodiment, the target biomarker is selected from the group of proteins consisting of CD9, PSMA, PCSA, CD63, CD81, B7H3, IL 6, OPG-13, IL6R, PA2G4, EZH2, RUNX2, SERPINB3, and EpCam. In another embodiment, a target biomarker is selected from the group of proteins consisting of A33, a33 n15, AFP, ALA, ALIX, ALP, AnnexinV, APC, ASCA, ASPH (246-260), ASPH (666-680), ASPH (A-10), ASPH (D01P), ASPH (D03), ASPH (G-20), ASPH (H-300), AURKA, AURKB, B7H3, B7H4, BCA-225, BCNP1, BDNF, BRCA, CA125 (MUC16), CA-19-9, C-Bir, CD1.1, CD10, CD174 (Lewis y), CD24, CD44, CD46, CD59 (MEM-43), CD63, CD66e CEA, CD73, CD81, CD9, CDA, CDAC1 1a2, CEA, C-Erb2, C-erbB2, CRMP-2, CRP, CXCL12, CYFRA21-1, DLL4, DR3, EGFR, Epcam, EphA2, EphA2 (H-77), ER, ErbB4, EZH2, FASL, FRT, FRT c.f23, GDF15, GPCR, GPR30, Gro-alpha, HAP, HBD 1, HBD2, HER 3 (ErbB3), HSP, HSP70, hVEGFR2, iC3b, IL 6 Unc, IL-1B, IL6 Unc, IL6R, IL8, IL-8, INSIG-2, KLK2, L1CAM, LAMN, LDH, MACC-1, MAPK4, MART-1, MCP-1, M-CSF, MFG-E8, MIC1, MIF, MIS RII, MMG, MMP26, MMP7, MMP9, MS4A1, MUC1, MUC1 seq1, MUC1 seq11A, MUC17, MUC2, Ncam, NGAL, NPGP/NPFF2, OPG, OPN, p53, p53, PA2G4, PBP, PCSA, PDGFRB, PGP9.5, PIM1, PR (B), PRL, PSA, PSMA, PSME3, PTEN, R5-CD9 Tube 1, Reg IV, RUNX2, SCRN1, seprase, SERPINB3, SPARC, SPB, SPDEF, SRVN, STAT 3, STEAP1, TF (FL-295), TFF3, TGM2, TIMP-1, TIMP1, TIMP2, TMEM211, TMPRSS2, TNF-alpha, Trail-R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, VEGF A, and YPSMA-1. The target biomarker can be selected from the group of proteins consisting of 5T4, A33, ACTG1, ADAM10, ADAM15, AFP, ALA, ALDOA, ALIX, ALP, ALX4, ANCA, Annexin V, ANXA2, ANXA6, APC, APOA1, ASCA, ASPH, ATP1A1, AURKA, AURKB, B7H3, B7H4, BANK1, BASP1, BCA-225, BCNP1, BDNF, BRCA, C1orf58, C20orf114, C8B, CA125 (MUC16), CA-19-9, CAPZA1, CAV1, C-Bir, CCSA-2, CCSA-3&4, CD1.1, CD10, CD151, CD174 (Lewis y), CD24, CD2AP, CD37, CD44, CD46, CD53, CD59, CD63, CD66 CEA, CD73, CD81, CD82, CD9, CDA, CDAC1 1a2, CEA, C-Erbb2, CFL1, CFP, CHMP4B, CLTC, COTL1, CRMP-2, CRP, CRTN, CTNND1, CTSB, CTSZ, CXCL12, CYCS, CYFRA21-1, DcR3, DLL4, DPP4, DR3, EEF1A1, EGFR, EHD1, ENO1, EpCAM, EphA2, ER, ErbB4,

EZH2, F11R, F2, F5, FAM125A, FASL, Ferritin, FNBP1L, FOLH1, FRT, GAL3, GAPDH, GDF15, GLB1, GPCR (GPR110), GPR30, GPX3, GRO-1, Gro-alpha, HAP, HBD 1, HBD2, HER 3 (ErbB3), HIST1H1C, HIST1H2AB, HNP1-3, HSP, HSP70, HSP90AB1, HSPA1B, HSPA8, hVEGFR2, iC3b, ICAM, IGSF8, IL 6, IL-1B, IL6R, IL8, IMP3, INSIG-2, ITGB1, ITIH3, JUP, KLK2, L1CAM, LAMN, LDH, LDHA, LDHB, LUM, LYZ, MACC-1, MAPK4, MART-1, MCP-1, M-CSF, MFGE8, MGAM, MGC20553, MIC1, MIF, MIS RII, MMG, MMP26, MMP7, MMP9, MS4A1, MUC1, MUC17, MUC2, MYH2, MYL6B, Ncam, NGAL, NME1, NME2, NNMT, NPGP/NPFF2, OPG, OPG-13, OPN, p53, PA2G4, PABPC1, PABPC4, PACSIN2, PBP, PCBP2, PCSA, PDCD6IP, PDGFRB, PGP9.5, PIM1, PR (B), PRDX2, PRL, PSA, PSCA, PSMA, PSMA1, PSMA2, PSMA4, PSMA6, PSMA7, PSMB1, PSMB2, PSMB3, PSMB4, PSMB5, PSMB6, PSMB8, PSME3, PTEN, PTGFRN, Rab-5b, Reg IV, RPS27A, RUNX2, SCRN1, SDCBP, seprase, Sept-9, SERINC5, SERPINB3, SERPINB3, SH3GL1, SLC3A2, SMPDL3B, SNX9, SPARC, SPB, SPDEF, SPON2, SPR, SRVN, SSX2, SSX4, STAT 3, STEAP, STEAP1, TACSTD1, TCN2, tetraspanin, TF (FL-295), TFF3, TGM2, THBS1, TIMP, TIMP1, TIMP2, TMEM211, TMPRSS2, TNF-alpha, TPA, TPI1, TPS, Trail-R2, Trail-R4, TrKB, TROP2, TROP2, Tsg 101, TUBB, TWEAK, UNC93A, VDAC2, VEGF A, VPS37B, YPSMA-1, YWHAG, YWHAQ, and YWHAZ. In another embodiment, the target biomarker is selected from the group of proteins consisting of 5T4, ACTG1, ADAM10, ADAM15, ALDOA, ANXA2, ANXA6, APOA1, ATP1A1, BASP1, C1orf58, C20orf114, C8B, CAPZA1, CAV1, CD151, CD2AP, CD59, CD9, CD9, CFL1, CFP, CHMP4B, CLTC, COTL1, CTNND1, CTSB, CTSZ, CYCS, DPP4, EEF1A1, EHD1, ENO1, F11R, F2, F5, FAM125A, FNBP1L, FOLH1, GAPDH, GLB1, GPX3, HIST1H1C, HIST1H2AB, HSP90AB1, HSPA1B, HSPA8, IGSF8, ITGB1, ITIH3, JUP, LDHA, LDHB, LUM, LYZ, MFGE8, MGAM, MMP9, MYH2, MYL6B, NME1, NME2, PABPC1, PABPC4, PACSIN2, PCBP2, PDCD6IP, PRDX2, PSA, PSMA, PSMA1, PSMA2, PSMA4, PSMA6, PSMA7, PSMB1, PSMB2, PSMB3, PSMB4, PSMB5, PSMB6, PSMB8, PTGFRN, RPS27A, SDCBP, SERINC5, SH3GL1, SLC3A2, SMPDL3B, SNX9, TACSTD1, TCN2, THBS1, TPI1, TSG101, TUBB, VDAC2, VPS37B, YWHAG, YWHAQ, and YWHAZ. In another embodiment, the target biomarker is selected from the group of proteins consisting of CD9, CD63, CD81, PSMA, PCSA, B7H3 and EpCam. In another embodiment, the target biomarker is selected from the group of proteins consisting of a tetraspanin, CD9, CD63, CD81, CD63, CD9, CD81, CD82, CD37, CD53, Rab-5b, Annexin V, MFG-E8, Muc1, GPCR 110, TMEM211 and CD24 In another embodiment, the target biomarker is selected from the group of proteins consisting of A33, AFP, ALIX, ALX4, ANCA, APC, ASCA, AURKA, AURKB, B7H3, BANK1, BCNP1, BDNF, CA-19-9, CCSA-2, CCSA-3&4, CD10, CD24, CD44, CD63, CD66 CEA, CD66e CEA, CD81, CD9, CDA, C-Erb2, CRMP-2, CRP, CRTN, CXCL12, CYFRA21-1, DcR3, DLL4, DR3, EGFR, Epcam, EphA2, FASL, FRT, GAL3, GDF15, GPCR (GPR110), GPR30, GRO-1, HBD 1, HBD2, HNP1-3, IL-1B, IL8, IMP3, L1CAM, LAMN, MACC-1, MGC20553, MCP-1, M-CSF, MIC1, MIF, MMP7, MMP9, MS4A1, MUC1, MUC17, MUC2, Ncam, NGAL, NNMT, OPN, p53, PCSA, PDGFRB, PRL, PSMA, PSME3, Reg IV, SCRN1, Sept-9, SPARC, SPON2, SPR, SRVN, TFF3, TGM2, TIMP-1, TMEM211, TNF-alpha, TPA, TPS, Trail-

R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, and VEGFA. In another embodiment, the target biomarker is selected from the group of proteins consisting of CD9, EGFR, NGAL, CD81, STEAP, CD24, A33, CD66E, EPHA2, Ferritin, GPR30, GPR110, MMP9, OPN, p53, TMEM211, TROP2, TGM2, TIMP, EGFR, DR3, UNC93A, MUC17, EpCAM, MUC1, MUC2, TSG101, CD63, B7H3, CD24, and a tetraspanin. The target biomarker can be selected from the group of proteins consisting of 5HT2B, 5T4 (trophoblast), ACO2, ACSL3, ACTN4, ADAM10, AGR2, AGR3, ALCAM, ALDH6A1, ANGPTL4, ANO9, AP1G1, APC, APEX1, APLP2, APP (Amyloid precursor protein), ARCN1, ARHGAP35, ARL3, ASAH1, ASPH (A-10), ATP1B1, ATP1B3, ATP5I, ATP5O, ATXN1, B7H3, BACE1, BAI3, BAIAP2, BCA-200, BDNF, BigH3, BIRC2, BLVRB, BRCA, BST2, C1GALT1, C1GALT1C1, C20orf3, CA125, CACYBP, Calmodulin, CAPN1, CAPNS1, CCDC64B, CCL2 (MCP-1), CCT3, CD10(BD), CD127 (IL7R), CD174, CD24, CD44, CD80, CD86, CDH1, CDH5, CEA, CFL2, CHCHD3, CHMP3, CHRDL2, CIB1, CKAP4, COPA, COX5B, CRABP2, CRIP1, CRISPLD1, CRMP-2, CRTAP, CTLA4, CUL3, CXCR3, CXCR4, CXCR6, CYB5B, CYB5R1, CYCS, CYFRA 21, DBI, DDX23, DDX39B, derlin 1, DHCR7, DHX9, DLD, DLL4, DNAJB1, DPP6, DSTN, eCadherin, EEF1D, EEF2, EFTUD2, EIF4A2, EIF4A3, EpCaM, EphA2, ER(1) (ESR1), ER(2) (ESR2), Erb B4, Erbb2, erbb3 (Erb-B3), ERLIN2, ESD, FARSA, FASN, FEN1, FKBP5, FLNB, FOXP3, FUS, Gal3, GCDPF-15, GCNT2, GNA12, GNG5, GNPTG, GPC1, GPC2, GPC3, GPC4, GPC5, GPC6, GPD2, GPER (GPR30), GSPT1, H3F3B, H3F3C, HADH, HAP1, HER3, HIST1H1C, HIST1H2AB, HIST1H3A, HIST1H3C, HIST1H3D, HIST1H3E, HIST1H3F, HIST1H3G, HIST1H3H, HIST1H3I, HIST1H3J, HIST2H2BF, HIST2H3A, HIST2H3C, HIST2H3D, HIST3H3, HMGB1, HNRNPA2B1, HNRNPAB, HNRNPC, HNRNPD, HNRNPH2, HNRNPK, HNRNPL, HNRNPM, HNRNPU, HPS3, HSP-27, HSP70, HSP90B1, HSPA1A, HSPA2, HSPA9, HSPE1, IC3b, IDE, IDH3B, IDO1, IFI30, IL1RL2, IL7, IL8, ILF2, ILF3, IQCG, ISOC2, IST1, ITGA7, ITGB7, junction plakoglobin, Keratin 15, KRAS, KRT19, KRT2, KRT7, KRT8, KRT9, KTN1, LAMP1, LMNA, LMNB1, LNPEP, LRPPRC, LRRC57, Mammaglobin, MAN1A1, MAN1A2, MART1, MATR3, MBD5, MCT2, MDH2, MFGE8, MFGE8, MGP, MMP9, MRP8, MUC1, MUC17, MUC2, MYO5B, MYOF, NAPA, NCAM, NCL, NG2 (CSPG4), Ngal, NHE-3, NME2, NONO, NPM1, NQO1, NT5E (CD73), ODC1, OPG, OPN (SC), OS9, p53, PACSIN3, PAICS, PARK7, PARVA, PC, PCNA, PCSA, PD-1, PD-L1, PD-L2, PGP9.5, PHB, PHB2, PIK3C2B, PKP3, PPL, PR(B), PRDX2, PRKCB, PRKCD, PRKDC, PSA, PSAP, PSMA, PSMB7, PSMD2, PSME3, PYCARD, RAB1A, RAB3D, RAB7A, RAGE, RBL2, RNPEP, RPL14, RPL27, RPL36, RPS25, RPS4X, RPS4Y1, RPS4Y2, RUVBL2, SET, SHMT2, SLAIN1, SLC39A14, SLC9A3R2, SMARCA4, SNRPD2, SNRPD3, SNX33, SNX9, SPEN, SPR, SQSTM1, SSBP1, ST3GAL1, STXBP4, SUB1, SUCLG2, Survivin, SYT9, TFF3 (secreted), TGOLN2, THBS1, TIMP1, TIMP2, TMED10, TMED4, TMED9, TMEM211, TOM1, TRAF4 (scaffolding), TRAIL-R2, TRAP1, TrkB, Tsg 101, TXNDC16, U2AF2, UEVLD, UFC1, UNC93a, USP14, VASP, VCP, VDAC1, VEGFA, VEGFR1, VEGFR2, VPS37C, WIZ, XRCC5, XRCC6, YB-1, YWHAZ, or any combination thereof. In other embodiments, the target biomarker is selected from the group consisting of p53, p63, p73, mdm-2, procathepsin-D, B23, C23, PLAP, CA125, MUC-1, HER2,

NY-ESO-1, SCP1, SSX-1, SSX-2, SSX-4, HSP27, HSP60, HSP90, GRP78, TAG72, HoxA7, HoxB7, EpCAM, ras, mesothelin, survivin, EGFK, MUC-1, or c-myc. The microvesicle antigen can be from any of **Tables 18-26** or **Table 45**.

[00497] One of skill will appreciate that the above biomarker listings are not intended to be mutually exclusive. For example, a single target biomarker can have one or more of the following attributes: cancer/tumor antigen, cell antigen, microvesicle antigen, membrane antigen, and any combination thereof. In some embodiments, the target biomarker will have all of these attributes.

[00498] As noted above, the IDM domain can be constructed to illicit a complement mediated immune response that can induce apoptosis. Such IDM can include but are not limited to C1q, C1r, C1s, C1, C3a, C3b, C3d, C5a, C2, C4, and cytokines. The IDM region may comprise an oligonucleotide sequence including without limitation Toll-Like Receptor (TLR) agonists like CpG sequences which are immunostimulatory and/or polyG sequences which can be anti-proliferative or pro-apoptotic. The moiety can be vaccine like moiety or antigen that stimulates an immune response. In an embodiment, the immune stimulating moiety comprises a superantigen. In some embodiments, the superantigen can be selected from the group consisting of staphylococcal enterotoxins (SEs), a Streptococcus pyogenes exotoxin (SPE), a Staphylococcus aureus toxic shock-syndrome toxin (TSST-1), a streptococcal mitogenic exotoxin (SME), a streptococcal superantigen (SSA), a hepatitis surface antigen, or a combination thereof. Other bacterial antigens that can be used with the invention comprise bacterial antigens such as Freund's complete adjuvant, Freund's incomplete adjuvant, monophosphoryl-lipid A/trehalose dicorynomycolate (Ribi's adjuvant), BCG (Calmette-Guerin Bacillus; Mycobacterium bovis), and Corynebacterium parvum. The immune stimulating moiety can also be a non-specific immunostimulant, such as an adjuvant or other non-specific immunostimulator. Useful adjuvants comprise without limitation aluminium salts, alum, aluminium phosphate, aluminium hydroxide, squalene, oils, MF59, and AS03 ("Adjuvant System 03"). The adjuvant can be selected from the group consisting of Cationic liposome-DNA complex JVRS-100, aluminum hydroxide vaccine adjuvant, aluminum phosphate vaccine adjuvant, aluminum potassium sulfate adjuvant, Alhydrogel, ISCOM(s)™, Freund's Complete Adjuvant, Freund's Incomplete Adjuvant, CpG DNA Vaccine Adjuvant, Cholera toxin, Cholera toxin B subunit, Liposomes, Saponin Vaccine Adjuvant, DDA Adjuvant, Squalene-based Adjuvants, Etx B subunit Adjuvant, IL-12 Vaccine Adjuvant, LTK63 Vaccine Mutant Adjuvant, TiterMax Gold Adjuvant, Ribi Vaccine Adjuvant, Montanide ISA 720 Adjuvant, Corynebacterium-derived P40 Vaccine Adjuvant, MPL™ Adjuvant, AS04, AS02, Lipopolysaccharide Vaccine Adjuvant, Muramyl Dipeptide Adjuvant, CRL1005, Killed Corynebacterium parvum Vaccine Adjuvant, Montanide ISA 51, Bordetella pertussis component Vaccine Adjuvant, Cationic Liposomal Vaccine Adjuvant, Adamantylamide Dipeptide Vaccine Adjuvant, Arlacel A, VSA-3 Adjuvant, Aluminum vaccine adjuvant, Polygen Vaccine Adjuvant, Adjumer™, Algal Glucan, Bay R1005, Theramide®, Stearyl Tyrosine, Specol, Algammulin, Avridine®, Calcium Phosphate Gel, CTA1-DD gene fusion protein, DOC/Alum Complex, Gamma Inulin, Gerbu Adjuvant, GM-CSF, GMDP, Recombinant hIFN-gamma/Interferon-g, Interleukin-1β, Interleukin-2, Interleukin-7, Sclavo peptide,

Rehydragel LV, Rehydragel HPA, Loxoribine, MF59, MTP-PE Liposomes, Murametide, Murapalmitine, D-Murapalmitine, NAGO, Non-Ionic Surfactant Vesicles, PMMA, Protein Cochleates, QS-21, SPT (Antigen Formulation), nanoemulsion vaccine adjuvant, AS03, Quil-A vaccine adjuvant, RC529 vaccine adjuvant, LTR192G Vaccine Adjuvant, E. coli heat-labile toxin, LT, amorphous aluminum hydroxyphosphate sulfate adjuvant, Calcium phosphate vaccine adjuvant, Montanide Incomplete Seppic Adjuvant, Imiquimod, Resiquimod, AF03, Flagellin, Poly(I:C), ISCOMATRIX®, Abisco-100 vaccine adjuvant, Albumin-heparin microparticles vaccine adjuvant, AS-2 vaccine adjuvant, B7-2 vaccine adjuvant, DHEA vaccine adjuvant, Immunoliposomes Containing Antibodies to Costimulatory Molecules, SAF-1, Sendai Proteoliposomes, Sendai-containing Lipid Matrices, Threonyl muramyl dipeptide (TMDP), Ty Particles vaccine adjuvant, Bupivacaine vaccine adjuvant, DL-PGL (Polyester poly (DL-lactide-co-glycolide)) vaccine adjuvant, IL-15 vaccine adjuvant, LTK72 vaccine adjuvant, MPL-SE vaccine adjuvant, non-toxic mutant E112K of Cholera Toxin mCT-E112K, and Matrix-S. Additional adjuvants that can be used with the multipartite constructs of the invention can be identified using the Vaxjo database. See Sayers S, Ulysse G, Xiang Z, and He Y. Vaxjo: a web-based vaccine adjuvant database and its application for analysis of vaccine adjuvants and their uses in vaccine development. Journal of Biomedicine and Biotechnology. 2012;2012:831486. Epub 2012 Mar 13. PMID: 22505817; www.violinet.org/vaxjo/. Other useful non-specific immunostimulators comprise histamine, interferon, transfer factor, tuftsin, interleukin-1, female sex hormones, prolactin, growth hormone vitamin D, deoxycholic acid (DCA), tetrachlorodecaoxide (TCDO), and imiquimod or resiquimod, which are drugs that activate immune cells through the toll-like receptor 7. A multipartite construct can be created that comprises more than one immunomodulating moiety, e.g., using segments that span CpG sequences which are immunostimulatory with complement directed segments that can stimulate apoptosis.

[00499] ***Personalized multipartite constructs***

[00500] The oligonucleotide probe libraries of the invention can be used to construct personalized multipartite constructs. As an example, an oligonucleotide probe library can be used to probe a sample from a patient. The biomarker targets are identified for library members that preferentially recognize diseased cells, e.g., cancer cells, from the individual. Such methodology is described herein. See, e.g., **Example 23**. The variable region of one or more oligonucleotide probe to the patient's diseased cells is used to synthesize a multipartite construct of the invention. Such construct can be used as a personalized treatment for the patient. **FIG. 23E** illustrates a flow chart of this approach. An oligonucleotide probe library **2351** is contacted with patient sample **2352**. The sample can be any useful patient sample, including without limitation tissue such as biopsy or tissue removed during surgical or other procedures, bodily fluids, frozen sections taken for histological purposes, cell cultures, and various embodiments, fractions, and components of any thereof as described herein. See, e.g., section entitled "Samples" above. Oligonucleotide probes that bind the patient sample are identified **2353**. Methodology for identifying oligonucleotide probe members that bind a biological sample is described herein, and includes without limitation sequence analysis such as next generation sequencing, amplification and/or hybridization

approaches. In an embodiment, flow sorting is used to separate diseased cells from the contacted patient sample and sorted and bound oligonucleotides are identified **2353**. In an optional step, the biomarkers recognized by the oligonucleotide probe binders are identified **2354**. This step can be optional as the target biomarker of a disease-specific oligonucleotide probe may not be necessary to synthesize a multipartite construct specific for the patient's disease. However, the identification of the target may assist in design of the multipartite construct. For example, identification of the target may allow computer modeling of the construct's *in vivo* interactions or help to select the most promising probe from multiple candidates, e.g., by avoiding toxicity associated with non-disease specific targets. The oligonucleotide probe binders are used to create a multipartite construct of the invention **2355**. The multipartite construct can be as shown in **FIG. 23A** or variants thereof. In an embodiment, the variable region of an oligonucleotide probe binder is used as the variable region as shown in **FIG. 23A**. The multipartite construct can be administered to the patient **2356**, thereby providing a personalized therapy for the patient. Further as noted in the figure, constructs comprising the variable region of an oligonucleotide probe binder can be used for other purposes, such as disease monitoring **2357**. As an example of such methodology, the oligonucleotide probe binder can be used to detect the presence or absence of disease markers in the patient over time such as in an immunoassay format.

[00501] ***Anti-C1q oligonucleotides***

[00502] The complement system is a part of the immune system that enhances (complements) the ability of antibodies and phagocytic cells to clear microbes and damaged cells from an organism. It is part of the innate immune system, which is not adaptable and does not change over the course of an individual's lifetime. However, it can be recruited and brought into action by the adaptive immune system. Complement activation or fixation can stimulate phagocytes to clear foreign and damaged material, induce inflammation to attract additional phagocytes, and activate the cell-killing membrane attack complex. The "classical" complement pathway is triggered by activation of the C1-complex, which occurs when C1q binds to IgM or IgG complexed with antigens. The C1-complex is composed of 1 molecule of C1q, 2 molecules of C1r and 2 molecules of C1s, or $C1qr_2s_2$. Such immunoglobulin-mediated binding of the complement uses the ability of the immunoglobulin system to detect and bind to non-self antigens. C1q can also directly identify various structures and ligands on microbial surfaces and apoptotic cells, and binds additional self proteins including C-reactive protein (CRP), HIV-1, phosphatidylserine (PS), HTLV-1, and others. Because the complement system has the potential to be extremely damaging to host tissues, its activation must be tightly regulated. The classical pathway is inhibited by C1-inhibitor, which binds to C1 to prevent its activation. C1q also performs a number of non-complement functions, including without limitation such diverse functions as clearance of bacterial pathogens, induction of angiogenesis during wound healing, tolerance induction, anti-inflammatory responses and inhibiting T cell response. As a result of these diverse functions, complement and C1q play a role in diverse diseases and disorders, including without limitation autoimmune settings, pregnancy disorders, pathogen infection, aggregated proteins leading to neurodegenerative diseases, inflammation, and cancer. Deficiencies have been

associated with autoimmune disease (e.g., systemic lupus erythematosus), pathogen infection and cancer. However, the tumor microenvironment may also hijack C1q to promote cell adhesion, migration and proliferation. S*ee, e.g.*, Kouser et al., Emerging and Novel Functions of Complement Protein C1q, Front Immunol. 2015; 6: 317. Published online 2015 Jun 29; Son et al., Fundamental role of C1q in autoimmunity and inflammation, Immunol Res. 2015 Dec; 63(1-3): 101–106; Ghebrehiwet et al., The C1q Family of Proteins: Insights into the Emerging Non-Traditional Functions, Front Immunol. 2012; 3: 52; Nayak et al., Complement and non-complement activating functions of C1q: a prototypical innate immune molecule. Innate Immun. 2012 Apr;18(2):350-63.

[00503] C1q is a Ca2+ dependent hexameric complex comprised of 18 polypeptide chains, 6 of three different subunits (C1q A chain (P02745), C1q B chain (P02746), and C1q C chain (P02747)), that binds C1r and C1s to form the C1 complex, the first component in classical pathway of complement. C1q globular heads form a pattern recognition complex that binds to various targets, including without limitation clustered antigen-antibody Fc immune complexes (e.g., IgG, IgM), C-reactive protein (CRP), abnormal proteins (e.g., prion and beta-amyloid), apoptotic and secondary necrotic cells, phosphatidylserine and the surface of a subpopulation of microparticles in human plasma. Recognition of IgG and IgM on a cell surface can induce a complement cascade and lead to apoptosis. *See, e.g.,* Kishore et al., C1q and tumor necrosis factor superfamily: modularity and versatility, TRENDS in Immunology 25 (2004) 551-561; Nayak et al., Complement and non-complement activating functions of C1q: a prototypical innate immune molecule, Innate Immunity 18 (2012) 350-363. Aptamer-biotin-C1q protein conjugates have been used to induce complement mediated cell death. *See, e.g.,* Bruno, Aptamer–biotin–streptavidin–C1q complexes can trigger the classical complement pathway to kill cancer cells, In Vitro Cell Dev Biol —Animal (2010) 46:107–113.

[00504] C1q globular heads has been shown to bind DNA and recognize apoptotic cells. *See, e.g.,* Païdassi et al., The lectin-like activity of human C1q and its implication in DNA and apoptotic cell recognition, FEBS Letters 582 (2008) 3111-3116; Navratil et al., The globular heads of C1q specifically recognize surface blebs of apoptotic vascular endothelial cells, J Immunol 166 (2001) 3231-3239. DNA binds C1qA and activates the complement cascade without interfering with the ability of C1q to bind antibody Fc regions. *See, e.g.,* Jiang et al., DNA binds and activates complement via residues 14-26 of the human C1q A chain, J Biol Chem 267 (1992) 25597-25601; Garlatti, et al. Cutting edge: C1q binds deoxyribose and heparan sulfate through neighboring sites of its recognition domain, J Immunol 185 (2010) 808-812.

[00505] C1q protein quantification has been used for disease monitoring and monoclonal antibody (mAb) production. For example, C1q mAb is used to coat ELISA plates to capture and quantitate immune complexes in clinical samples. Various companies sell diagnostic kits for immune complex detection and quantitation which are based on the ability of C1q to bind well to immune complexes, but to not bind significantly to monomeric immunoglobulins. Because the DNA recognition domain of C1q does not overlap with the Fc-recognition domain, a DNA based ELISA may further allow a more accurate quantitation of immune complex detection.

[00506] **Example 35** herein presents identification of an anti-C1q oligonucleotide aptamers and describes various uses thereof. The aptamers to C1q were identified via oligonucleotide probe analysis of plasma microvesicles followed by identification of oligonucleotide probe targets using gel electrophoresis and mass spectrometry analysis.

[00507] Anti-C1q aptamers of the invention can be used for multiple purposes. As described above, the invention provides a multipartite construct having a disease specific target oligonucleotide or antibody (Ab) that can recognize a target of interest and an immunomodulatory region. In an embodiment of the invention, the immunomodulatory region comprises the C1q aptamer. Such construct can act as an immunotherapeutic agent for targeted cell killing via recruitment of complement proteins and the downstream membrane attack complex (MAC). By linking the C1q aptamer segment to another segment that specifically binds to a target of interest (e.g., a biomarker present on a cell or microvesicle of interest), the construct can bring C1q into proximity of a target. See **FIG. 23C**, which illustrates a construct **2331** having a segment that recognizes a Marker of Interest **2332** on a Membrane **2333**, and another segment that attracts the Complement system **2334**. Such binding can cause a complement cascade and induce complement mediated cell killing. This approach can be applied in multiple setting, e.g., to recognize cancer cells, gram negative bacteria, and/or viral and/or parasitic infections. For example, an anti-CD20 specific oligonucleotide can be linked with an anti-C1q specific oligonucleotide. The linkage to create the oligonucleotide - oligonucleotide construct can include but is not limited to direct synthesis with a spacer between the two oligonucleotide recognition sites. Different biomarkers can be used as the target of interest, thereby directing the complement cascade to the various targets as desired. The spacer type and size can be configured based on steric hindrance between the target protein and the C1q protein/MAC complex. As noted above, the target specific oligonucleotides/Abs can be chosen to specically recognize various targets of interest, including but not limited to cancer cells, circulating tumor cells, immune cells (e.g., B-cells, T-cells, neutrophils, macrophage, dendritic cells) microvesicles, bacteria, viruses or parasites. In addition to C1q, the target of the complement specific oligonucleotide segment can include without limitation C1r, C1s, C1, C3a, C3b, C3d, C5a, C2, C4, and cytokines.

[00508] The multipartite construct of the invention can comprise a linear molecule, a circular molecule, and/or adopt various secondary structures. **FIG. 23D** illustrates a construct **2340** having a C1q recruiting domain **2341** (nucleotides 37-77) and a C-met targeting domain **2342** (nucleotides 1-36). As desired, the C1q recruiting domain **2341** can comprise an anti-C1q oligonucleotide sequence of the invention. *See e.g.,* **Example 35**. As shown in the illustration, the C1q recruiting domain **2341** comprises a single stranded hairpin and a complementary base pairing region, and the C-met targeting domain **2342** comprises a complementary base pairing region and a region having a more complex secondary structure. Such structures can be estimated using available software programs such as Vienna or mfold (available at mfold.rit.albany.edu). Such structural estimates can also be used to design derivatives of the sequences, e.g., by substituting, adding or deleting nucleotides in order to increase or decrease melting temperature,

facilitate additions of non-natural nucleotide analogs, direct chemical modification, and/or manipulate structure or other parameters.

[00509] The invention further provides a method of molecular profiling of patient specific autoantigens by identifying autoantigens bound to complement 1 (C1) in plasma. The invention also provides immunoassays that detect levels of C1q protein. Such assays can be any applicable immunoassay format using the anti-C1q oligonucleotide of the invention, including without limitation an oligonucleotide based ELISA, Western analysis, flow cytometry, or affinity isolation. The immunoassay can be applied to various settings, including without limitation: 1) monitor cancer patient specific immune responses before, during and after administration of immunosuppressing drugs for optimal treatment with chemotherapeutic agents; 2) monitor immune responses in patients with autoimmune disorders in response to administration of immunosuppressing drugs such as TNF blockers; 3) detect levels of C1q and/or anti-C1q autoantibodies in patients with systemic lupus erythematosus (SLE); 4) quantitatative C1q assay for mAb biosimilar production to satisfy the EMA biosimilar antibody guidance measures; 5) a WHO secondary test as a companion test to mAb based ELISAs; 6) as a marker for apoptosis/secondary necrosis; and 7) a C1q test for research purposes.

[00510] The anti-C1q oligonucleotides of the invention can undergo various modifications such as described herein or known in the art. For example, modifications can be made to alter desired characteristics, including without limitation in vivo stability, specificity, affinity, avidity or nuclease susceptibility. Alterations to the half life may improve stability in vivo or may reduce stability to limit in vivo toxicity. Such alterations can include mutations, truncations or extensions. The 5' and/or 3' ends of the multipartite oligonucleotide constructs can be protected or deprotected to modulate stability as well. Modifications to improve in vivo stability, specificity, affinity, avidity or nuclease susceptibility or alter the half life to influence in vivo toxicity may be at the 5' or 3' end and include but are not limited to the following: locked nucleic acid (LNA) incorporation, unlocked nucleic acid (UNA) incorporation, phosphorothioate backbone instead of phosphodiester backbone, amino modifiers (i.e. C6-dT), dye conjugates (Cy dues, Fluorophores, etc), Biotinylation, PEG linkers, Click chemistry linkers, dideoxynucleotide end blockers, inverted end bases, cholesterol TEG or other lipid based labels. *See, e.g.,* Campbell, MA and Wengel, J (2011). Locked vs. unlocked nucleic acids (LNA vs. UNA): contrasting structures work towards common therapeutic goals. Chem Soc Rev 40: 5680–5689; and Wahlestedt, C, Salmi, P, Good, L, Kela, J, Johnsson, T, Hökfelt, T et al. (2000). Potent and nontoxic antisense oligonucleotides containing locked nucleic acids. Proc Natl Acad Sci USA 97: 5633–5638; which publications are incorporated by reference herein in their entirety.

[00511] ***Aptamer 10.36***

[00512] We reported that aptamer 10.36 (5'-CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG; SEQ ID NO. 4357) forms a G-quadruplex structure and is taken up by Burkitt's Lymphoma (Ramos) cells via a clathrin-mediated endocytotic pathway and that this aptamer is taken up by Ramos cells to a greater extent than other lymphoma derived cell lines (e.g., Jurkat, Raji, or P12). Opazo F, *et al.* (2015). Modular

Assembly of Cell-targeting Devices Based on an Uncommon G-quadruplex Aptamer Molecular Therapy. Nucleic Acids 4, e251. Aptamer 10.36 comprises a central G rich region surrounded by flanking complementary strands. See Opazo et al.

[00513] In an aspect, the invention provides an oligonucleotide comprising a sequence selected from any one of SEQ ID NOs. 4357-4368 or 4372-4407. In a preferred embodiment, the oligonucleotide comprises a sequence according to SEQ ID NO. 4357, i.e., the sequence of aptamer 10.36. The invention further provides an oligonucleotide having a substitution in aptamer 10.36 such as in SEQ ID NOs. 4372-4407. The sequence can comprise the central G rich region of 10.36 (i.e., nucleotides 9-25 of SEQ ID NO. 4357) surrounded by complementary flanking regions. The flanking regions can be any useful length, e.g., at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or at least 10 nucleotides in length. See, e.g., Opazo et al. The aptamer sequence may also comprise additions and deletions. For example, at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or at least 10 nucleotides may be inserted between the G rich region and the flanking regions as desired. Alternately, nucleotides may be deleted between the G rich region and the flanking regions as desired. Substitutions, additions and deletions in the sequence can be chosen such that the aptamer retains or improves upon desired such as stability, target recognition and G quadruplex structure. In a related aspect, the invention provides an oligonucleotide comprising a sequence selected from any one of SEQ ID NOs. 4357-4368, and a 5' region with sequence 5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131), a 3' region with sequence 5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132), or both.

[00514] The oligonucleotide of the invention can be capable of binding to a target in **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61**, or a subcomponent thereof. In some embodiments, the oligonucleotide is capable of binding to Ramos cells. The oligonucleotide can be capable of binding to a protein selected from the group consisting of PARP1, HIST1H1B, HIST1H1D, NCL, FBL, SFPQ, RPL12, ACTB, HIST1H4A, SSBP1, NONO, H2AFJ, and DDX21, or a complex, subunit or fragment thereof. In some embodiment, the oligonucleotide is capable of binding to cells comprising surface nucleolin.

[00515] The invention further provides an oligonucleotide comprising a nucleic acid sequence or a portion thereof that is at least 50, 55, 60, 65, 70, 75, 80, 85, 86, 86, 88, 89, 90, 95, 96, 97, 98, 99 or 100 percent homologous to an oligonucleotide sequence described above.

[00516] In another aspect, the invention provides a plurality of oligonucleotides comprising at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, or at least 10000 different oligonucleotide sequences described above.

[00517] The oligonucleotide or the plurality of oligonucleotides provided by the invention may comprise a DNA, RNA, 2'-O-methyl or phosphorothioate backbone, or any combination thereof. The oligonucleotide or the plurality of oligonucleotides may comprise at least one of DNA, RNA, PNA, LNA, UNA, and any combination thereof.

[00518] In some embodiments, the oligonucleotide or the plurality of oligonucleotides comprises at least one functional modification selected from the group consisting of biotinylation, a non-naturally occurring nucleotide, a deletion, an insertion, an addition, and a chemical modification. The chemical modification can be chosen to modulate desired properties such as stability, capture, detection, or binding efficiency. In some embodiments, the chemical modification comprises at least one of C18, polyethylene glycol (PEG), PEG4, PEG6, PEG8, and PEG12. The oligonucleotide or plurality of oligonucleotides can be labeled. The oligonucleotide or plurality of oligonucleotides can be attached to a nanoparticle, liposome, gold, magnetic label, fluorescent label, light emitting particle, or radioactive label. The liposome or particle can incorporate desired entities such as chemotherapeutic agents or detectable labels. Other useful modifications are disclosed herein.

[00519] In an aspect, the invention provides an isolated oligonucleotide or plurality of oligonucleotides having a sequence as described above. In a related aspect, the invention provides a composition comprising such isolated oligonucleotide or plurality of oligonucleotides.

[00520] In some embodiments, the isolated oligonucleotide or at least one member of the plurality of oligonucleotides is capable of binding to Ramos cell. In some embodiments, the isolated oligonucleotide or plurality of oligonucleotides is capable of binding to a protein selected from the group consisting of PARP1, HIST1H1B, HIST1H1D, NCL, FBL, SFPQ, RPL12, ACTB, HIST1H4A, SSBP1, NONO, H2AFJ, and DDX21, or a complex, subunit or fragment thereof. In some embodiments, the isolated oligonucleotide or plurality of oligonucleotides is capable of binding to a cell surface nucleolin complex.

[00521] The isolated oligonucleotide or plurality of oligonucleotides can by capable of inhibiting nucleolin activity. The isolated oligonucleotide or plurality of oligonucleotides can be capable of modulating cell proliferation. In some embodiments, the isolated oligonucleotide or plurality of oligonucleotides is capable of inducing apoptosis. The cell proliferation can be neoplastic or dysplastic growth. The cell proliferation can be that of cancer cells such as disclosed herein, including without limitation that of lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer. In certain embodiments, the cell proliferation is that of leukemia, lymphoma or renal carcinoma cells.

[00522] The isolated oligonucleotide or plurality of oligonucleotides may bind to a cell surface nucleolin, a complex comprising nucleolin, or another cell surface protein from **Table 61**. Such bound outer complex may mediate cellular internalization of the complex. Such binding may also interfere with nucleolin function in the nucleus, cytoplasm, or membrane.

[00523] In an aspect, the invention provides a method comprising synthesizing the at least one oligonucleotide or the plurality of oligonucleotides provided above. Techniques for synthesizing oligonucleotides are disclosed herein or are known in the art.

[00524] In another aspect, the invention provides a method comprising contacting a biological sample with the at least one oligonucleotide, the plurality of oligonucleotides, or composition as described above.

The method can further comprise detecting a presence or level of a protein in **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61** in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. In some embodiments, the method further comprises detecting a presence or level of a protein in **Table 61** in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. In still other embodiments, the method comprises detecting a presence or level of a nucleolin protein or complex thereof in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. Relatedly, the method may further comprise detecting a presence or level of a cell population in the biological sample that is bound by the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. For example, the cells may display a protein in in **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61** on their surface. The cell population can be any desired population, including without limitation neoplastic, malignant, tumor, hyperplastic, or dysplastic cells. In some embodiments, the cell population comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells.

[00525] The detecting step of the method may comprise detecting the at least one oligonucleotide or at least one member of the plurality of oligonucleotides. The presence or level of oligonucleotide serves as a proxy for the level of oligonucleotide's target. The oligonucleotides can be detecting using any desired technique such as described herein or known in the art, including without limitation at least one of sequencing, amplification, hybridization, gel electrophoresis, chromatography, and any combination thereof. Any useful sequencing method can be employed, including without limitation at least one of next generation sequencing, dye termination sequencing, pyrosequencing, and any combination thereof. In some embodiments, the detecting comprises transmission electron microscopy (TEM) of immunogold labeled oligonucleotides. In some embodiments, the detecting comprises confocal microscopy of fluor labeled oligonucleotides.

[00526] The detecting step of the method may comprise detecting the protein or cell using techniques described herein or known in the art for detecting proteins, including without limitation at least one of an immunoassay, enzyme immunoassay (EIA), enzyme-linked immunosorbent assay (ELISA), enzyme-linked oligonucleotide assay (ELONA), affinity isolation, immunoprecipitation, Western blot, gel electrophoresis, microscopy or flow cytometry.

[00527] In some embodiments of the method, the detected protein is associated with a microvesicle population. The method may further comprise isolating the microvesicle population prior to the contacting with the oligonucleotides, after the contacting, or both. The isolating may be in whole or in part. For example, the microvesicle population may be partially isolated from other components in the sample before or after contacting the sample with the oligonucleotide or plurality of oligonucleotides. The invention may use any appropriate techniques to isolate microvesicles. Various techniques of isolating microvesicles are disclosed herein or known in the art, including without limitation affinity purification,

filtration, concentration, polymer precipitation, PEG precipitation, ultracentrifugation, a molecular crowding reagent, affinity selection, chromatography, or any combination thereof.

[00528] Any desired biological sample can be contacted with the oligonucleotide or plurality of oligonucleotides according to the invention. In various embodiments, the biological sample comprises a bodily fluid, tissue sample or cell culture. Any desired tissue or cell culture sample can be contacted. In some embodiments, the tissue or cell culture sample comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells. Similarly, any appropriate bodily fluid can be contacted, such as those disclosed herein. In certain preferred embodiments, the bodily fluid comprises whole blood or a derivative or fraction thereof, such as sera or plasma. The bodily fluid may comprise cancer cells, including without limitation lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells.

[00529] The biological sample may be spiked with a purified or recombinant protein (or both). In some embodiments, such protein is selected from **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61**, or complexes, subunits or fragments thereof. For example, the spiking can be used as a control in an assay.

[00530] As desired, the method of detecting the presence or level of the at least one oligonucleotide, the plurality of oligonucleotides, or composition bound to a target can be used to characterize a phenotype. The phenotype can be any appropriate phenotype, including without limitation a disease or disorder. In such cases, the characterizing may include providing, or assisting in providing, at least one of diagnostic, prognostic and theranostic information for the disease or disorder. Characterizing the phenotype may comprise comparing the presence or level to a reference. Any appropriate reference level can be used. For example, the reference can be the presence or level determined in a sample from at least one individual without the phenotype or from at least one individual with a different phenotype. As a further example, if the phenotype is a disease or disorder, the reference level may be the presence or level determined in a sample from at least one individual without the disease or disorder, or with a different state of the disease or disorder (*e.g.*, in remission, different stage or grade, different prognosis, metastatic versus local, etc).

[00531] As noted, the sample can be from a subject suspected of having or being predisposed to a disease or disorder. The disease or disorder can be any disease or disorder that can be assessed by the subject method. For example, the disease or disorder may be a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain. In certain embodiments, the cancer is a cancer disclosed herein (see, e.g., Section "Phenotypes"), including without limitation lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer.

[00532] As further described herein, the invention provides a kit comprising a reagent for carrying out the method. Similarly, the invention provides for the use of a reagent for carrying out the method. The reagent can be any useful reagent for carrying out the method. For example, the reagent can be the at least one oligonucleotide or the plurality of oligonucleotides, one or more primer for amplification or sequencing of such oligonucleotides, at least one binding agent to at least one protein, a binding buffer with or without $MgCl_2$, a sample processing reagent, a microvesicle isolation reagent, a cell isolation reagent, a detection reagent, a secondary detection reagent, a wash buffer, an elution buffer, a solid support, and any combination thereof. The microvesicle isolation reagent may comprise at least one of a concentrator unit, a filtration unit, a polymer, PEG, a size exclusion column, a binding agent to a microvesicle antigen, and any combination thereof; and/or the detection or secondary detection agent comprises streptavidin-horse radish peroxide (HRP), a streptavidin-conjugated fluorophore, a streptavidin-conjugated quantum dot, and any combination thereof.

[00533] The G quadruplex aptamer AS1411 has been shown to decrease viability of a variety of cancer cell types and has undergone phase II clinical trials. *See, e.g.,* Bates, P et al. (2009). Discovery and development of the G-rich oligonucleotide AS1411 as a novel treatment for cancer. Experimental and Molecular Pathology 86:151–164; Rosenberg J, et al. (2014). A phase II trial of the nucleolin-targeted DNA aptamer AS1411 in metastatic refractory renal cell carcinoma. Invest New Drugs 32:178-187, which references are incorporated by reference herein in their entirety. AS1411 is believed to bind the protein nucleolin, a nucleolar phosphoprotein which is overexpressed on the surface of certain cancer cells. AS1411 bound to surface nucleolin may be internalized and prevent nuclear nucleiln from stabilizing BCL2 mRNA. Reduced levels of BCL2 may then lead to apoptosis. AS1411 has been used to create nucleolin targeted liposomes and nanoparticles for drug delivery to cancer cells. *See, e.g.,* Wu et al. Nucleolin Targeting AS1411 Modified Protein Nanoparticle for Antitumor Drugs Delivery, Mol. Pharmaceutics, 2013, 10 (10), pp 3555–3563; Li et al., Nucleolin-targeting liposomes guided by aptamer AS1411 for the delivery of siRNA for the treatment of malignant melanomas, Biomaterials 35: 3840–3850 (2014); Ai et al., Multifunctional AS1411-functionalized fluorescent gold nanoparticles for targeted cancer cell imaging and efficient photodynamic therapy, Talanta 118:54-60 (2014); Zhang et al. Nucleolin targeting AS1411 aptamer modified pH-sensitive micelles for enhanced delivery and antitumor efficacy of paclitaxel. Nano Research 2015 8, 201-218. AS1411 has also been used as an imaging agent. See, e.g., Li et al., Aptamer imaging with Cu-64 labeled AS1411: Preliminary assessment in lung cancer, Nuc Med Biol, 41:179-185 (2014).

[00534] In an aspect, the invention provides a method of imaging a cell or tissue, comprising contacting the cell or tissue with at least one oligonucleotide or plurality of oligonucleotides as described above, e.g., aptamer 10.36, and detecting the oligonucleotides in contact with at least one cell or tissue. In some embodiments, the oligonucleotides are labeled, e.g., in order to facilitate detection or medical imaging. The oligonucleotides can be attached to a nanoparticle, liposome, gold, magnetic label, fluorescent label, light emitting particle, radioactive label, or other useful label such as disclosed herein or known in the art.

The oligonucleotides can be administered to a subject prior to the detecting. The cell or tissue can comprise cells displaying nucleolin or another protein from **Table 50**, **Table 52**, **Table 53**, **Table 60** or **Table 61** on their surface. In some embodiments, the cell or tissue comprises neoplastic, malignant, tumor, hyperplastic, or dysplastic cells. For example, the cell or tissue may comprise lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, non-small cell lung cancer, or other cancer cells such as described herein.

[00535] In an aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of the 10.36 oligonucleotide aptamers, or a salt thereof, and a pharmaceutically acceptable carrier, diluent, or both. In some embodiments, the oligonucleotides are attached to a toxin or chemotherapeutic agent. In some embodiments, the oligonucleotides are attached to a liposome or nanoparticle. The liposome or nanoparticle may comprise a toxin or chemotherapeutic agent. In such embodiments, the at least one oligonucleotide or the plurality of oligonucleotides can be used for targeted delivery of the toxin, chemotherapeutic agent, liposome or nanoparticle to a desired target cell or tissue.

[00536] In a related aspect, the invention provides a method of treating or ameliorating a disease or disorder in a subject in need thereof, comprising administering such pharmaceutical composition to the subject. In another related aspect, the invention provides a method of inducing cytotoxicity in a subject, comprising administering such pharmaceutical to the subject. The pharmaceutical composition can be administered in any useful format. In various embodiments, the administering comprises at least one of intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, by inhalation, topical administration, or any combination thereof. The carrier or diluent can be any useful carrier or diluent, as described herein or known in the art. As desired, the pharmaceutical composition can be administered in combination with additional known chemotherapeutic agents such as described herein or known in the art, e.g., cyclophosphamide, etoposide, doxorubicin, methotrexate, vincristine, procabazine, prednisone, dexamethasone, tamoxifen citrate, carboplatin, cisplatin, oxaliplatin, 5-fluorouracil, camptothecin, zoledronic acid, Ibandronate or mytomicin.

[00537] ***Modifications***

[00538] Modifications to the one or more oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, or any combination thereof, can be made to alter desired characteristics, including without limitation *in vivo* stability, specificity, affinity, avidity or nuclease susceptibility. Alterations to the half life may improve stability in vivo or may reduce stability to limit *in vivo* toxicity. Such alterations can include mutations, truncations or extensions. The 5' and/or 3' ends of the multipartite oligonucleotide constructs can be protected or deprotected to modulate stability as well. Modifications to improve *in vivo* stability, specificity, affinity, avidity or nuclease susceptibility or alter the half life to influence *in vivo* toxicity may be at the 5' or 3' end and include but are not limited to the following: locked nucleic acid (LNA) incorporation, unlocked nucleic acid (UNA) incorporation,

phosphorothioate backbone instead of phosphodiester backbone, amino modifiers (i.e. C6-dT), dye conjugates (Cy dues, Fluorophores, etc), Biotinylation, PEG linkers, Click chemistry linkers, dideoxynucleotide end blockers, inverted end bases, cholesterol TEG or other lipid based labels.

[00539] Linkage options for segments of the oligonucleotide of the invention can be on the 5' or 3' end of an oligonucleotide or to a primary amine, sulfhydryl or carboxyl group of an antibody and include but are not limited to the following: Biotin-target oligonucleotide /Ab, streptavidin-complement oligonucleotide or vice versa, amino modified-target Ab/ oligonucleotide, thiol/carboxy-complement oligonucleotide or vice versa, Click chemistry-target Ab/ oligonucleotide, corresponding Click chemistry partner-complement oligonucleotide or vice versa. The linkages may be covalent or non-covalent and may include but are not limited to monovalent, multivalent (i.e. bi, tri or tetra-valent) assembly, to a DNA scaffold (i.e. DNA origami structure), drug/chemotherapeutic agent, nanoparticle, microparticle or a micelle or liposome.

[00540] A linker region can comprise a spacer with homo- or multifunctional reactive groups that can vary in length and type. These include but are not limited to the following: spacer C18, PEG4, PEG6, PEG8, and PEG12.

[00541] The multipartite oligonucleotide of the invention can further comprise additional elements to add desired biological effects. For example, the oligonucleotide of the invention may comprise a membrane disruptive moiety. The oligonucleotide of the invention may also be conjugated to one or more chemical moiety that provides such effects. For example, the oligonucleotide of the invention may be conjugated to a detergent-like moiety to disrupt the membrane of a target cell or microvesicle. Useful ionic detergents include sodium dodecyl sulfate (SDS, sodium lauryl sulfate (SLS)), sodium laureth sulfate (SLS, sodium lauryl ether sulfate (SLES)), ammonium lauryl sulfate (ALS), cetrimonium bromide, cetrimonium chloride, cetrimonium stearate, and the like. Useful non-ionic (zwitterionic) detergents include polyoxyethylene glycols, polysorbate 20 (also known as Tween 20), other polysorbates (e.g., 40, 60, 65, 80, etc), Triton-X (e.g., X100, X114), 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS), CHAPSO, deoxycholic acid, sodium deoxycholate, NP-40, glycosides, octyl-thio-glucosides, maltosides, and the like. One of skill will appreciate that functional fragments, such as membrance disruptive moieties, can be covalently or non-covalently attached to the oligonucleotide of the invention.

[00542] Oligonucleotide segments, including those of a multipartite construct, can include any desireable base modification known in the art. In certain embodiments, oligonucleotide segments are 10 to 50 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies oligonucleotides of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length, or any range derivable there within.

[00543] In certain embodiments, a multipartite construct comprises a chimeric oligonucleotide that contains two or more chemically distinct regions, each made up of at least one nucleotide. Such chimeras can be referred to using terms such as multipartite, multivalent, or the like. The oligonucleotides portions

may contain at least one region of modified nucleotides that confers one or more beneficial properties, e.g., increased nuclease resistance, bioavailability, increased binding affinity for the target. Chimeric nucleic acids of the invention may be formed as composite structures of two or more oligonucleotides, two or more types of oligonucleotides (e.g., both DNA and RNA segments), modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics. Such compounds have also been referred to in the art as hybrids. Representative United States patents that teach the preparation of such hybrid structures comprise, but are not limited to, US patent nos: 5,013,830; 5,149,797; 5, 220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, each of which is herein incorporated by reference in its entirety.

[00544] In certain embodiments, an oligonucleotide of the invention comprises at least one nucleotide modified at the 2' position of the sugar, including without limitation a 2'-0-alkyl, 2'-0-alkyl-0-alkyl or 2'-fluoro-modified nucleotide. In other embodiments, RNA modifications include 2'- fluoro, 2'-amino and 2' O-methyl modifications on the ribose of pyrimidines, a basic residue or an inverted base at the 3' end of the RNA. Such modifications are routinely incorporated into oligonucleotides and these oligonucleotides have been shown to have higher target binding affinity in some cases than 2'-deoxyoligonucleotides against a given target.

[00545] A number of nucleotide and nucleoside modifications have been shown to make an oligonucleotide more resistant to nuclease digestion, thereby prolonging *in vivo* half- life. Specific examples of modified oligonucleotides include those comprising backbones comprising, for example, phosphorothioates, phosphotriesters, methyl phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. The constructs of the invention can comprise oligonucleotides with phosphorothioate backbones and/or heteroatom backbones, e.g., CH2 -NH-0-CH2, CH,~N(CH3)~0~CH2 (known as a methylene(methylimino) or MMI backbone], CH2 -O-N (CH3)-CH2, CH2 -N (CH3)-N (CH3)-CH2 and O-N (CH3)- CH2 -CH2 backbones, wherein the native phosphodiester backbone is represented as O- P— O- CH,); amide backbones (De Mesmaeker et ah, 1995); morpholino backbone structures (Summerton and Weller, U.S. Pat. No. 5,034,506); peptide nucleic acid (PNA) backbone (wherein the phosphodiester backbone of the oligonucleotide is replaced with a polyamide backbone, the nucleotides being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone (Nielsen, et al., 1991), each of which is herein incorporated by reference in its entirety. Phosphorus- containing linkages include, but are not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3 '-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3 '-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3*-5* to 5*-3* or 2*-5* to 5*-2*; see U.S. Patent Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5, 177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302;

5,286,717; 5,321, 131; 5,399,676; 5,405,939; 5,453,496; 5,455, 233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563, 253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference in its entirety. Morpholino-based oligomeric compounds are known in the art described in Braasch & Corey, Biochemistry vol. 41, no. 14, 2002, pages 4503 - 4510; Genesis vol. 30, 2001, page 3; Heasman, J. Dev. Biol. vol. 243, 2002, pages 209 - 214; Nasevicius et al. Nat. Genet. vol. 26, 2000, pages 216 - 220; Lacerra et al. Proc. Natl. Acad. Sci. vol. 97, 2000, pages 9591 - 9596 and U.S. Pat. No. 5,034,506, issued Jul. 23, 1991, each of which is herein incorporated by reference in its entirety. Cyclohexenyl nucleic acid oligonucleotide mimetics are described in Wang et al., J. Am. Chem. Soc. Vol. 122, 2000, pages 8595 - 8602, the contents of which is incorporated herein in its entirety. An oligonucleotide of the invention can comprise at least such modification as desired.

[00546] Modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that can be formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These comprise those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH2 component parts; see U.S. Patent Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216, 141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596, 086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, each of which is herein incorporated by reference in its entirety. An oligonucleotide of the invention can comprise at least such modification as desired.

[00547] In certain embodiments, an oligonucleotide of the invention comprises one or more substituted sugar moieties, e.g., one of the following at the 2' position: OH, SH, SCH_3, F, OCN, OCH_3 OCH_3, OCH_3 $O(CH_2)n$ CH_3, $O(CH_2)n$ NH_2 or $O(CH_2)n$ CH_3 where n is from 1 to about 10; Ci to CIO lower alkyl, alkoxyalkoxy, substituted lower alkyl, alkaryl or aralkyl; CI; Br; CN; CF_3; OCF_3; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; $SOCH_3$; SO_2 CH_3; ONO_2; N O_2; N_3; NH_2; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacokinetic/pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy [2'-0- CH2CH2OCH3, also known as 2'-0-(2-methoxyethyl)]. Other preferred modifications include 2*-methoxy (2*-0-CH3), 2*-propoxy (2*-OCH2 CH2CH3) and 2*-fiuoro (2*-F). Similar modifications may also be made at other positions on the oligonucleotide, e.g., the 3' position of the sugar on the 3' terminal nucleotide and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyls in place of the pentofuranosyl group.

[00548] In certain embodiments, an oligonucleotide of the invention comprises one or more base modifications and/or substitutions. As used herein, "unmodified" or "natural" bases include adenine (A), guanine (G), thymine (T), cytosine (C) and uracil (U). Modified bases include, without limitation, bases found only infrequently or transiently in natural nucleic acids, e.g., hypoxanthine, 6-methyladenine, 5 -Me pyrimidines, particularly 5-methylcytosine (also referred to as 5-methyl-2' deoxy cytosine and often referred to in the art as 5-Me-C), 5- hydroxymethylcytosine (HMC), glycosyl HMC and gentobiosyl HMC, as well as synthetic bases, e.g., 2-aminoadenine, 2-(methylamino)adenine, 2-(imidazolylalkyl)adenine, 2- (aminoalklyamino)adenine or other heterosubstituted alkyladenines, 2-thiouracil, 2- thiothymine, 5-bromouracil, 5-hydroxymethyluracil, 8-azaguanine, 7-deazaguanine, N6 (6-aminohexyl)adenine and 2,6-diaminopurine (Kornberg, 1980; Gebeyehu, et ah, 1987). A "universal" base known in the art, e.g., inosine, can also be included. 5-Me-C substitutions can also be included. These have been shown to increase nucleic acid duplex stability by 0.6- 1.20C. See, e.g., Sanghvi et al., 'Antisense Research & Applications', 1993, CRC PRESS pages 276 - 278. Further suitable modified bases are described in U.S. Patent Nos. 3,687,808, as well as 4,845,205; 5,130,302; 5,134,066; 5,175, 273; 5, 367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,596,091; 5,614,617; 5,750,692, and 5,681,941, each of which is herein incorporated by reference.

[00549] It is not necessary for all positions in a given oligonucleotide to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single oligonucleotide or even at within a single nucleoside within an oligonucleotide.

[00550] In certain embodiments, both a sugar and an internucleoside linkage, i.e., the backbone, of one or more nucleotide units within an oligonucleotide of the invention are replaced with novel groups. The base can be maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to retain hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, for example, an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative patents that teach the preparation of PNA compounds comprise, but are not limited to, U.S. Patent Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al. Science vol. 254, 1991, page 1497, which is herein incorporated by reference.

[00551] In certain embodiments, the oligonucleotide of the invention is linked (covalently or non-covalently) to one or more moieties or conjugates that enhance activity, cellular distribution, or localization. Such moieties include, without limitation, lipid moieties such as a cholesterol moiety (Letsinger et al. Proc. Natl. Acad. Sci. Usa. vol. 86, 1989, pages 6553 - 6556), cholic acid (Manoharan et al. Bioorg. Med. Chem. Let. vol. 4, 1994, pages 1053 - 1060), a thioether, e.g., hexyl-S- tritylthiol (Manoharan et al. Ann. N. Y. Acad. Sci. Vol. 660, 1992, pages 306 - 309; Manoharan et al. Bioorg. Med. Chem. Let. vol. 3, 1993, pages 2765 - 2770), a thiocholesterol (Oberhauser et al. Nucl. Acids Res. vol. 20,

1992, pages 533 - 538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Kabanov et al. Febs Lett. vol. 259, 1990, pages 327 - 330; Svinarchuk et al. Biochimie. vol. 75, 1993, pages 49 - 54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1 ,2-di-O-hexadecyl- rac- glycero-3-H-phosphonate (Manoharan et al. Tetrahedron Lett. vol. 36, 1995, pages 3651 - 3654; Shea et al. Nucl. Acids Res. vol. 18, 1990, pages 3777 - 3783), a polyamine or a polyethylene glycol chain (Mancharan et al. Nucleosides & Nucleotides vol. 14, 1995, pages 969 - 973), or adamantane acetic acid (Manoharan et al. Tetrahedron Lett. vol. 36, 1995, pages 3651 - 3654), a palmityl moiety (Mishra et al. Biochim. Biophys. Acta vol. 1264, 1995, pages 229 - 237), or an octadecylamine or hexylamino- carbonyl-t oxycholesterol moiety (Crooke et al. J. Pharmacol. Exp. Ther. vol. 277, 1996, pages 923 - 937), each of which is herein incorporated by reference in its entirety. See also U.S. Patent Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241,5,391,723; 5,416,203,5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference in its entirety.

[00552] The oligonucleotide of the invention can be modified to incorporate a wide variety of modified nucleotides as desired. For example, the construct may be synthesized entirely of modified nucleotides or with a subset of modified nucleotides. The modifications can be the same or different. Some or all nucleotides may be modified, and those that are modified may contain the same modification. For example, all nucleotides containing the same base may have one type of modification, while nucleotides containing other bases may have different types of modification. All purine nucleotides may have one type of modification (or are unmodified), while all pyrimidine nucleotides have another, different type of modification (or are unmodified). Thus, the construct may comprise any combination of desired modifications, including for example, ribonucleotides (2'-OH), deoxyribonucleotides (2'-deoxy), 2'-amino nucleotides (2'-NH2), 2'- fluoro nucleotides (2'-F) and 2'-0-methyl (2'-OMe) nucleotides.

[00553] In some embodiments, the oligonucleotide of the invention is synthesized using a transcription mixture containing modified nucleotides in order to generate a modified construct. For example, a transcription mixture may contain only 2'-OMe A, G, C and U and/or T triphosphates (2'-OMe ATP, 2'-OMe UTP and/or 2*-OMe TTP, 2*-OMe CTP and 2*-OMe GTP), referred to as an MNA or mRmY mixture. Oligonucleotides generated therefrom are referred to as MNA oligonucleotides or mRmY oligonucleotides and contain only 2'-0-methyl nucleotides. A transcription mixture containing all 2'-OH nucleotides is referred to as an "rN" mixture, and oligonucleotides generated therefrom are referred to as "rN", "rRrY" or RNA oligonucleotides. A transcription mixture containing all deoxy nucleotides is referred to as a "dN" mixture, and oligonucleotides generated therefrom are referred to as "dN", "dRdY" or DNA oligonucleotides. Aternatively, a subset of nucleotides (e.g., C, U and /or T) may comprise a first

modified nucleotides (e.g, 2'-OMe) nucleotides and the remainder (e.g., A and G) comprise a second modified nucleotide (e.g., 2'-OH or 2'-F). For example, a transcription mixture containing 2'-F U and 2'-OMe A, G and C is referred to as a "fUmV" mixture, and oligonucleotides generated therefrom are referred to as "fUmV" oligonucleotides. A transcription mixture containing 2'-F A and G, and 2'-OMe C and U and/or T is referred to as an "fRmY" mixture, and oligonucleotides generated therefrom are referred to as "fRmY" oligonucleotides. A transcription mixture containing 2'-F A and 2'-OMe C, G and U and/or T is referred to as "fAmB" mixture, and oligonucleotides generated therefrom are referred to as "fAmB" oligonucleotides.

[00554] One of skill in the art can improve pre-identified aptamer segments (e.g., variable regions or immunomodulatory regions that comprise an aptamer to a biomarker target or other entity) using various process modifications. Examples of such process modifications include, but are not limited to, truncation, deletion, substitution, or modification of a sugar or base or internucleotide linkage, capping, and PEGylation. In addition, the sequence requirements of an aptamer may be explored through doped reselections or aptamer medicinal chemistry. Doped reselections are carried out using a synthetic, degenerate pool that has been designed based on the aptamer of interest. The level of degeneracy usually varies from about 70-85% from the aptamer of interest. In general, sequences with neutral mutations are identified through the doped reselection process. Aptamer medicinal chemistry is an aptamer improvement technique in which sets of variant aptamers are chemically synthesized. These variants are then compared to each other and to the parent aptamer. Aptamer medicinal chemistry is used to explore the local, rather than global, introduction of substituents. For example, the following modifications may be introduced: modifications at a sugar, base, and/or internucleotide linkage, such as 2'-deoxy, 2'-ribo, or 2'-0-methyl purines or pyrimidines, phosphorothioate linkages may be introduced between nucleotides, a cap may be introduced at the 5' or 3' end of the aptamer (such as 3' inverted dT cap) to block degradation by exonucleases, or a polyethylene glycol (PEG) element may be added to the aptamer to increase the half-life of the aptamer in the subject.

[00555] Additional compositions comprising an oligonucleotide of the invention and uses thereof are further described below.

[00556] ***Pharmaceutical Compositions***

[00557] In an aspect, the invention provides pharmaceutical compositions comprising one or more oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, as described above, or any combination thereof. The invention further provides methods of administering such compositions.

[00558] The term "condition," as used herein means an interruption, cessation, or disorder of a bodily function, system, or organ. Representative conditions include, but are not limited to, diseases such as cancer, inflammation, diabetes, and organ failure.

[00559] The phrase "treating," "treatment of," and the like include the amelioration or cessation of a specified condition.

[00560] The phrase "preventing," "prevention of," and the like include the avoidance of the onset of a condition.

[00561] The term "salt," as used herein, means two compounds that are not covalently bound but are chemically bound by ionic interactions.

[00562] The term "pharmaceutically acceptable," as used herein, when referring to a component of a pharmaceutical composition means that the component, when administered to an animal, does not have undue adverse effects such as excessive toxicity, irritation, or allergic response commensurate with a reasonable benefit/risk ratio. Accordingly, the term "pharmaceutically acceptable organic solvent," as used herein, means an organic solvent that when administered to an animal does not have undue adverse effects such as excessive toxicity, irritation, or allergic response commensurate with a reasonable benefit/risk ratio. Preferably, the pharmaceutically acceptable organic solvent is a solvent that is generally recognized as safe ("GRAS") by the United States Food and Drug Administration ("FDA"). Similarly, the term "pharmaceutically acceptable organic base," as used herein, means an organic base that when administered to an animal does not have undue adverse effects such as excessive toxicity, irritation, or allergic response commensurate with a reasonable benefit/risk ratio.

[00563] The phrase "injectable" or "injectable composition," as used herein, means a composition that can be drawn into a syringe and injected subcutaneously, intraperitoneally, or intramuscularly into an animal without causing adverse effects due to the presence of solid material in the composition. Solid materials include, but are not limited to, crystals, gummy masses, and gels. Typically, a formulation or composition is considered to be injectable when no more than about 15%, preferably no more than about 10%, more preferably no more than about 5%, even more preferably no more than about 2%, and most preferably no more than about 1% of the formulation is retained on a 0.22 μm filter when the formulation is filtered through the filter at 98° F. There are, however, some compositions of the invention, which are gels, that can be easily dispensed from a syringe but will be retained on a 0.22 μm filter. In one embodiment, the term "injectable," as used herein, includes these gel compositions. In one embodiment, the term "injectable," as used herein, further includes compositions that when warmed to a temperature of up to about 40° C. and then filtered through a 0.22 μm filter, no more than about 15%, preferably no more than about 10%, more preferably no more than about 5%, even more preferably no more than about 2%, and most preferably no more than about 1% of the formulation is retained on the filter. In one embodiment, an example of an injectable pharmaceutical composition is a solution of a pharmaceutically active compound (for example, one or more oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, as described above, or any combination thereof) in a pharmaceutically acceptable solvent. One of skill will appreciate that injectable solutions have inherent properties, e.g., sterility, pharmaceutically acceptable excipients and free of harmful measures of pyrogens or similar contaminants.

[00564] The term "solution," as used herein, means a uniformly dispersed mixture at the molecular or ionic level of one or more substances (solute), in one or more other substances (solvent), typically a liquid.

[00565] The term "suspension," as used herein, means solid particles that are evenly dispersed in a solvent, which can be aqueous or non-aqueous.

[00566] The term "animal," as used herein, includes, but is not limited to, humans, canines, felines, equines, bovines, ovines, porcines, amphibians, reptiles, and avians. Representative animals include, but are not limited to a cow, a horse, a sheep, a pig, an ungulate, a chimpanzee, a monkey, a baboon, a chicken, a turkey, a mouse, a rabbit, a rat, a guinea pig, a dog, a cat, and a human. In one embodiment, the animal is a mammal. In one embodiment, the animal is a human. In one embodiment, the animal is a non-human. In one embodiment, the animal is a canine, a feline, an equine, a bovine, an ovine, or a porcine.

[00567] The phrase "drug depot," as used herein means a precipitate, which includes one or more oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, as described above, or any combination thereof, formed within the body of a treated animal that releases the oligonucleotide over time to provide a pharmaceutically effective amount of the oligonucleotide.

[00568] The phrase "substantially free of," as used herein, means less than about 2 percent by weight. For example, the phrase "a pharmaceutical composition substantially free of water" means that the amount of water in the pharmaceutical composition is less than about 2 percent by weight of the pharmaceutical composition.

[00569] The term "effective amount," as used herein, means an amount sufficient to treat or prevent a condition in an animal.

[00570] The nucleotides that make up the oligonucleotide of the invention can be modified to, for example, improve their stability, i.e., improve their in vivo half-life, and/or to reduce their rate of excretion when administered to an animal. The term "modified" encompasses nucleotides with a covalently modified base and/or sugar. For example, modified nucleotides include nucleotides having sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3′ position and other than a phosphate group at the 5′ position. Modified nucleotides may also include 2′ substituted sugars such as 2′-O-methyl-; 2′-O-alkyl; 2′-O-allyl; 2′-S-alkyl; 2′-S-allyl; 2′-fluoro-; 2′-halo or 2′-azido-ribose; carbocyclic sugar analogues; α-anomeric sugars; and epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, and sedoheptulose.

[00571] Modified nucleotides are known in the art and include, but are not limited to, alkylated purines and/or pyrimidines; acylated purines and/or pyrimidines; or other heterocycles. These classes of pyrimidines and purines are known in the art and include, pseudoisocytosine; N4,N4-ethanocytosine; 8-hydroxy-N6-methyladenine; 4-acetylcytosine, 5-(carboxyhydroxylmethyl) uracil; 5-fluorouracil; 5-bromouracil; 5-carboxymethylaminomethyl-2-thiouracil; 5-carboxymethylaminomethyl uracil; dihydrouracil; inosine; N6-isopentyl-adenine; 1-methyladenine; 1-methylpseudouracil; 1-methylguanine;

2,2-dimethylguanine; 2-methyladenine; 2-methylguanine; 3-methylcytosine; 5-methylcytosine; N6-methyladenine; 7-methylguanine; 5-methylaminomethyl uracil; 5-methoxy amino methyl-2-thiouracil; β-D-mannosylqueosine; 5-methoxycarbonylmethyluracil; 5-methoxyuracil; 2 methylthio-N6-isopentenyladenine; uracil-5-oxyacetic acid methyl ester; psueouracil; 2-thiocytosine; 5-methyl-2 thiouracil, 2-thiouracil; 4-thiouracil; 5-methyluracil; N-uracil-5-oxyacetic acid methylester; uracil 5-oxyacetic acid; queosine; 2-thiocytosine; 5-propyluracil; 5-propylcytosine; 5-ethyluracil; 5-ethylcytosine; 5-butyluracil; 5-pentyluracil; 5-pentylcytosine; and 2,6,-diaminopurine; methylpsuedouracil; 1-methylguanine; and 1-methylcytosine.

[00572] An oligonucleotide of the invention can also be modified by replacing one or more phosphodiester linkages with alternative linking groups. Alternative linking groups include, but are not limited to embodiments wherein P(O)O is replaced by P(O)S, P(S)S, P(O)NR2, P(O)R, P(O)OR′, CO, or CH2, wherein each R or R′ is independently H or a substituted or unsubstituted C1-C20 alkyl. A preferred set of R substitutions for the P(O)NR2 group are hydrogen and methoxyethyl. Linking groups are typically attached to each adjacent nucleotide through an —O— bond, but may be modified to include —N— or —S— bonds. Not all linkages in an oligomer need to be identical.

[00573] The oligonucleotide of the invention can also be modified by conjugation to a polymer, for example, to reduce the rate of excretion when administered to an animal. For example, the oligonucleotide can be "PEGylated," i.e., conjugated to polyethylene glycol ("PEG"). In one embodiment, the PEG has an average molecular weight ranging from about 20 kD to 80 kD. Methods to conjugate an oligonucleotide with a polymer, such PEG, are known to those skilled in the art (See, e.g., Greg T. Hermanson, Bioconjugate Techniques, Academic Press, 1966).

[00574] The oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, as described above, or any combination thereof, can be used in the pharmaceutical compositions disclosed herein or known in the art.

[00575] In one embodiment, the pharmaceutical composition further comprises a solvent.

[00576] In one embodiment, the solvent comprises water.

[00577] In one embodiment, the solvent comprises a pharmaceutically acceptable organic solvent. Any useful and pharmaceutically acceptable organic solvents can be used in the compositions of the invention.

[00578] In one embodiment, the pharmaceutical composition is a solution of the salt in the pharmaceutically acceptable organic solvent.

[00579] In one embodiment, the pharmaceutical composition comprises a pharmaceutically acceptable organic solvent and further comprises a phospholipid, a sphingomyelin, or phosphatidyl choline. Without wishing to be bound by theory, it is believed that the phospholipid, sphingomyelin, or phosphatidyl choline facilitates formation of a precipitate when the pharmaceutical composition is injected into water and can also facilitate controlled release of the oligonucleotide from the resulting precipitate. Typically, the phospholipid, sphingomyelin, or phosphatidyl choline is present in an amount ranging from greater than 0 to 10 percent by weight of the pharmaceutical composition. In one embodiment, the phospholipid,

sphingomyelin, or phosphatidyl choline is present in an amount ranging from about 0.1 to 10 percent by weight of the pharmaceutical composition. In one embodiment, the phospholipid, sphingomyelin, or phosphatidyl choline is present in an amount ranging from about 1 to 7.5 percent by weight of the pharmaceutical composition. In one embodiment, the phospholipid, sphingomyelin, or phosphatidyl choline is present in an amount ranging from about 1.5 to 5 percent by weight of the pharmaceutical composition. In one embodiment, the phospholipid, sphingomyelin, or phosphatidyl choline is present in an amount ranging from about 2 to 4 percent by weight of the pharmaceutical composition.

[00580] The pharmaceutical compositions can optionally comprise one or more additional excipients or additives to provide a dosage form suitable for administration to an animal. When administered to an animal, the oligonucleotide containing pharmaceutical compositions are typically administered as a component of a composition that comprises a pharmaceutically acceptable carrier or excipient so as to provide the form for proper administration to the animal. Suitable pharmaceutical excipients are described in Remington's Pharmaceutical Sciences 1447-1676 (Alfonso R. Gennaro ed., 19th ed. 1995), incorporated herein by reference. The pharmaceutical compositions can take the form of solutions, suspensions, emulsion, tablets, pills, pellets, capsules, capsules containing liquids, powders, suppositories, emulsions, aerosols, sprays, suspensions, or any other form suitable for use.

[00581] In one embodiment, the pharmaceutical compositions are formulated for intravenous or parenteral administration. Typically, compositions for intravenous or parenteral administration comprise a suitable sterile solvent, which may be an isotonic aqueous buffer or pharmaceutically acceptable organic solvent. Where necessary, the compositions can also include a solubilizing agent. Compositions for intravenous administration can optionally include a local anesthetic such as lidocaine to lessen pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where oligonucleotide-containing pharmaceutical compositions are to be administered by infusion, they can be dispensed, for example, with an infusion bottle containing, for example, sterile pharmaceutical grade water or saline. Where the pharmaceutical compositions are administered by injection, an ampoule of sterile water for injection, saline, or other solvent such as a pharmaceutically acceptable organic solvent can be provided so that the ingredients can be mixed prior to administration.

[00582] In another embodiment, the pharmaceutical compositions are formulated in accordance with routine procedures as a composition adapted for oral administration. Compositions for oral delivery can be in the form of tablets, lozenges, aqueous or oily suspensions, granules, powders, emulsions, capsules, syrups, or elixirs, for example. Oral compositions can include standard excipients such as mannitol, lactose, starch, magnesium stearate, sodium saccharin, cellulose, and magnesium carbonate. Typically, the excipients are of pharmaceutical grade. Orally administered compositions can also contain one or more agents, for example, sweetening agents such as fructose, aspartame or saccharin; flavoring agents such as peppermint, oil of wintergreen, or cherry; coloring agents; and preserving agents, to provide a

pharmaceutically palatable preparation. Moreover, when in tablet or pill form, the compositions can be coated to delay disintegration and absorption in the gastrointestinal tract thereby providing a sustained action over an extended period of time. Selectively permeable membranes surrounding an osmotically active driving compound are also suitable for orally administered compositions. A time-delay material such as glycerol monostearate or glycerol stearate can also be used.

[00583] The pharmaceutical compositions further comprising a solvent can optionally comprise a suitable amount of a pharmaceutically acceptable preservative, if desired, so as to provide additional protection against microbial growth. Examples of preservatives useful in the pharmaceutical compositions of the invention include, but are not limited to, potassium sorbate, methylparaben, propylparaben, benzoic acid and its salts, other esters of parahydroxybenzoic acid such as butylparaben, alcohols such as ethyl or benzyl alcohol, phenolic compounds such as phenol, or quaternary compounds such as benzalkonium chlorides (e.g., benzethonium chloride).

[00584] In one embodiment, the pharmaceutical compositions of the invention optionally contain a suitable amount of a pharmaceutically acceptable polymer. The polymer can increase the viscosity of the pharmaceutical composition. Suitable polymers for use in the compositions and methods of the invention include, but are not limited to, hydroxypropylcellulose, hydoxypropylmethylcellulose (HPMC), chitosan, polyacrylic acid, and polymethacrylic acid.

[00585] Typically, the polymer is present in an amount ranging from greater than 0 to 10 percent by weight of the pharmaceutical composition. In one embodiment, the polymer is present in an amount ranging from about 0.1 to 10 percent by weight of the pharmaceutical composition. In one embodiment, the polymer is present in an amount ranging from about 1 to 7.5 percent by weight of the pharmaceutical composition. In one embodiment, the polymer is present in an amount ranging from about 1.5 to 5 percent by weight of the pharmaceutical composition. In one embodiment, the polymer is present in an amount ranging from about 2 to 4 percent by weight of the pharmaceutical composition. In one embodiment, the pharmaceutical compositions of the invention are substantially free of polymers.

[00586] In one embodiment, any additional components added to the pharmaceutical compositions of the invention are designated as GRAS by the FDA for use or consumption by animals. In one embodiment, any additional components added to the pharmaceutical compositions of the invention are designated as GRAS by the FDA for use or consumption by humans.

[00587] The components of the pharmaceutical composition (the solvents and any other optional components) are preferably biocompatible and non-toxic and, over time, are simply absorbed and/or metabolized by the body.

[00588] As described above, the pharmaceutical compositions of the invention can further comprise a solvent.

[00589] In one embodiment, the solvent comprises water.

[00590] In one embodiment, the solvent comprises a pharmaceutically acceptable organic solvent.

[00591] In an embodiment, the oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, as described above, or any combination thereof, are available as the salt of a metal cation, for example, as the potassium or sodium salt. These salts, however, may have low solubility in aqueous solvents and/or organic solvents, typically, less than about 25 mg/mL. The pharmaceutical compositions of the invention comprising (i) an amino acid ester or amino acid amide and (ii) a protonated oligonucleotide, however, may be significantly more soluble in aqueous solvents and/or organic solvents. Without wishing to be bound by theory, it is believed that the amino acid ester or amino acid amide and the protonated oligonucleotide form a salt, such as illustrated above, and the salt is soluble in aqueous and/or organic solvents.

[00592] Similarly, without wishing to be bound by theory, it is believed that the pharmaceutical compositions comprising (i) an oligonucleotide of the invention; (ii) a divalent metal cation; and (iii) optionally a carboxylate, a phospholipid, a phosphatidyl choline, or a sphingomyelin form a salt, such as illustrated above, and the salt is soluble in aqueous and/or organic solvents.

[00593] In one embodiment, the concentration of the oligonucleotide of the invention in the solvent is greater than about 2 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide of the invention in the solvent is greater than about 5 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is greater than about 7.5 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is greater than about 10 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is greater than about 12 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is greater than about 15 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is ranges from about 2 percent to 5 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is ranges from about 2 percent to 7.5 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent ranges from about 2 percent to 10 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is ranges from about 2 percent to 12 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is ranges from about 2 percent to 15 percent by weight of the pharmaceutical composition. In one embodiment, the concentration of the oligonucleotide in the solvent is ranges from about 2 percent to 20 percent by weight of the pharmaceutical composition.

[00594] Any pharmaceutically acceptable organic solvent can be used in the pharmaceutical compositions of the invention. Representative, pharmaceutically acceptable organic solvents include, but are not limited to, pyrrolidone, N-methyl-2-pyrrolidone, polyethylene glycol, propylene glycol (i.e., 1,3-propylene glycol), glycerol formal, isosorbid dimethyl ether, ethanol, dimethyl sulfoxide, tetraglycol,

tetrahydrofurfuryl alcohol, triacetin, propylene carbonate, dimethyl acetamide, dimethyl formamide, dimethyl sulfoxide, and combinations thereof.

[00595] In one embodiment, the pharmaceutically acceptable organic solvent is a water soluble solvent. A representative pharmaceutically acceptable water soluble organic solvents is triacetin.

[00596] In one embodiment, the pharmaceutically acceptable organic solvent is a water miscible solvent. Representative pharmaceutically acceptable water miscible organic solvents include, but are not limited to, glycerol formal, polyethylene glycol, and propylene glycol.

[00597] In one embodiment, the pharmaceutically acceptable organic solvent comprises pyrrolidone. In one embodiment, the pharmaceutically acceptable organic solvent is pyrrolidone substantially free of another organic solvent.

[00598] In one embodiment, the pharmaceutically acceptable organic solvent comprises N-methyl-2-pyrrolidone. In one embodiment, the pharmaceutically acceptable organic solvent is N-methyl-2-pyrrolidone substantially free of another organic solvent.

[00599] In one embodiment, the pharmaceutically acceptable organic solvent comprises polyethylene glycol. In one embodiment, the pharmaceutically acceptable organic solvent is polyethylene glycol substantially free of another organic solvent.

[00600] In one embodiment, the pharmaceutically acceptable organic solvent comprises propylene glycol. In one embodiment, the pharmaceutically acceptable organic solvent is propylene glycol substantially free of another organic solvent.

[00601] In one embodiment, the pharmaceutically acceptable organic solvent comprises glycerol formal. In one embodiment, the pharmaceutically acceptable organic solvent is glycerol formal substantially free of another organic solvent.

[00602] In one embodiment, the pharmaceutically acceptable organic solvent comprises isosorbid dimethyl ether. In one embodiment, the pharmaceutically acceptable organic solvent is isosorbid dimethyl ether substantially free of another organic solvent.

[00603] In one embodiment, the pharmaceutically acceptable organic solvent comprises ethanol. In one embodiment, the pharmaceutically acceptable organic solvent is ethanol substantially free of another organic solvent.

[00604] In one embodiment, the pharmaceutically acceptable organic solvent comprises dimethyl sulfoxide. In one embodiment, the pharmaceutically acceptable organic solvent is dimethyl sulfoxide substantially free of another organic solvent.

[00605] In one embodiment, the pharmaceutically acceptable organic solvent comprises tetraglycol. In one embodiment, the pharmaceutically acceptable organic solvent is tetraglycol substantially free of another organic solvent.

[00606] In one embodiment, the pharmaceutically acceptable organic solvent comprises tetrahydrofurfuryl alcohol. In one embodiment, the pharmaceutically acceptable organic solvent is tetrahydrofurfuryl alcohol substantially free of another organic solvent.

[00607] In one embodiment, the pharmaceutically acceptable organic solvent comprises triacetin. In one embodiment, the pharmaceutically acceptable organic solvent is triacetin substantially free of another organic solvent.

[00608] In one embodiment, the pharmaceutically acceptable organic solvent comprises propylene carbonate. In one embodiment, the pharmaceutically acceptable organic solvent is propylene carbonate substantially free of another organic solvent.

[00609] In one embodiment, the pharmaceutically acceptable organic solvent comprises dimethyl acetamide. In one embodiment, the pharmaceutically acceptable organic solvent is dimethyl acetamide substantially free of another organic solvent.

[00610] In one embodiment, the pharmaceutically acceptable organic solvent comprises dimethyl formamide. In one embodiment, the pharmaceutically acceptable organic solvent is dimethyl formamide substantially free of another organic solvent.

[00611] In one embodiment, the pharmaceutically acceptable organic solvent comprises at least two pharmaceutically acceptable organic solvents.

[00612] In one embodiment, the pharmaceutically acceptable organic solvent comprises N-methyl-2-pyrrolidone and glycerol formal. In one embodiment, the pharmaceutically acceptable organic solvent is N-methyl-2-pyrrolidone and glycerol formal. In one embodiment, the ratio of N-methyl-2-pyrrolidone to glycerol formal ranges from about 90:10 to 10:90.

[00613] In one embodiment, the pharmaceutically acceptable organic solvent comprises propylene glycol and glycerol formal. In one embodiment, the pharmaceutically acceptable organic solvent is propylene glycol and glycerol formal. In one embodiment, the ratio of propylene glycol to glycerol formal ranges from about 90:10 to 10:90.

[00614] In one embodiment, the pharmaceutically acceptable organic solvent is a solvent that is recognized as GRAS by the FDA for administration or consumption by animals. In one embodiment, the pharmaceutically acceptable organic solvent is a solvent that is recognized as GRAS by the FDA for administration or consumption by humans.

[00615] In one embodiment, the pharmaceutically acceptable organic solvent is substantially free of water. In one embodiment, the pharmaceutically acceptable organic solvent contains less than about 1 percent by weight of water. In one embodiment, the pharmaceutically acceptable organic solvent contains less about 0.5 percent by weight of water. In one embodiment, the pharmaceutically acceptable organic solvent contains less about 0.2 percent by weight of water. Pharmaceutically acceptable organic solvents that are substantially free of water are advantageous since they are not conducive to bacterial growth. Accordingly, it is typically not necessary to include a preservative in pharmaceutical compositions that are substantially free of water. Another advantage of pharmaceutical compositions that use a pharmaceutically acceptable organic solvent, preferably substantially free of water, as the solvent is that hydrolysis of the oligonucleotide is minimized. Typically, the more water present in the solvent the more readily the oligonucleotide can be hydrolyzed. Accordingly, oligonucleotide containing pharmaceutical

compositions that use a pharmaceutically acceptable organic solvent as the solvent can be more stable than oligonucleotide containing pharmaceutical compositions that use water as the solvent.

[00616] In one embodiment, comprising a pharmaceutically acceptable organic solvent, the pharmaceutical composition is injectable.

[00617] In one embodiment, the injectable pharmaceutical compositions are of sufficiently low viscosity that they can be easily drawn into a 20 gauge and needle and then easily expelled from the 20 gauge needle. Typically, the viscosity of the injectable pharmaceutical compositions are less than about 1,200 cps. In one embodiment, the viscosity of the injectable pharmaceutical compositions are less than about 1,000 cps. In one embodiment, the viscosity of the injectable pharmaceutical compositions are less than about 800 cps. In one embodiment, the viscosity of the injectable pharmaceutical compositions are less than about 500 cps. Injectable pharmaceutical compositions having a viscosity greater than about 1,200 cps and even greater than about 2,000 cps (for example gels) are also within the scope of the invention provided that the compositions can be expelled through an 18 to 24 gauge needle.

[00618] In one embodiment, comprising a pharmaceutically acceptable organic solvent, the pharmaceutical composition is injectable and does not form a precipitate when injected into water.

[00619] In one embodiment, comprising a pharmaceutically acceptable organic solvent, the pharmaceutical composition is injectable and forms a precipitate when injected into water. Without wishing to be bound by theory, it is believed, for pharmaceutical compositions that comprise a protonated oligonucleotide and an amino acid ester or amide, that the α-amino group of the amino acid ester or amino acid amide is protonated by the oligonucleotide to form a salt, such as illustrated above, which is soluble in the pharmaceutically acceptable organic solvent but insoluble in water. Similarly, when the pharmaceutical composition comprises (i) an oligonucleotide; (ii) a divalent metal cation; and (iii) optionally a carboxylate, a phospholipid, a phosphatidyl choline, or a sphingomyelin, it is believed that the components of the composition form a salt, such as illustrated above, which is soluble in the pharmaceutically acceptable organic solvent but insoluble in water. Accordingly, when the pharmaceutical compositions are injected into an animal, at least a portion of the pharmaceutical composition precipitates at the injection site to provide a drug depot. Without wishing to be bound by theory, it is believed that when the pharmaceutically compositions are injected into an animal, the pharmaceutically acceptable organic solvent diffuses away from the injection site and aqueous bodily fluids diffuse towards the injection site, resulting in an increase in concentration of water at the injection site, that causes at least a portion of the composition to precipitate and form a drug depot. The precipitate can take the form of a solid, a crystal, a gummy mass, or a gel. The precipitate, however, provides a depot of the oligonucleotide at the injection site that releases the oligonucleotide over time. The components of the pharmaceutical composition, i.e., the amino acid ester or amino acid amide, the pharmaceutically acceptable organic solvent, and any other components are biocompatible and non-toxic and, over time, are simply absorbed and/or metabolized by the body.

[00620] In one embodiment, comprising a pharmaceutically acceptable organic solvent, the pharmaceutical composition is injectable and forms liposomal or micellar structures when injected into water (typically about 500 μL are injected into about 4 mL of water). The formation of liposomal or micellar structures are most often formed when the pharmaceutical composition includes a phospholipid. Without wishing to be bound by theory, it is believed that the oligonucleotide in the form of a salt, which can be a salt formed with an amino acid ester or amide or can be a salt with a divalent metal cation and optionally a carboxylate, a phospholipid, a phosphatidyl choline, or a sphingomyelin, that is trapped within the liposomal or micellar structure. Without wishing to be bound by theory, it is believed that when these pharmaceutically compositions are injected into an animal, the liposomal or micellar structures release the oligonucleotide over time.

[00621] In one embodiment, the pharmaceutical composition further comprising a pharmaceutically acceptable organic solvent is a suspension of solid particles in the pharmaceutically acceptable organic solvent. Without wishing to be bound by theory, it is believed that the solid particles comprise a salt formed between the amino acid ester or amino acid amide and the protonated oligonucleotide wherein the acidic phosphate groups of the oligonucleotide protonates the amino group of the amino acid ester or amino acid amide, such as illustrated above, or comprises a salt formed between the oligonucleotide; divalent metal cation; and optional carboxylate, phospholipid, phosphatidyl choline, or sphingomyelin, as illustrated above. Pharmaceutical compositions that are suspensions can also form drug depots when injected into an animal.

[00622] By varying the lipophilicity and/or molecular weight of the amino acid ester or amino acid amide it is possible to vary the properties of pharmaceutical compositions that include these components and further comprise an organic solvent. The lipophilicity and/or molecular weight of the amino acid ester or amino acid amide can be varied by varying the amino acid and/or the alcohol (or amine) used to form the amino acid ester (or amino acid amide). For example, the lipophilicity and/or molecular weight of the amino acid ester can be varied by varying the R1 hydrocarbon group of the amino acid ester. Typically, increasing the molecular weight of R1 increase the lipophilicity of the amino acid ester. Similarly, the lipophilicity and/or molecular weight of the amino acid amide can be varied by varying the R3 or R4 groups of the amino acid amide.

[00623] For example, by varying the lipophilicity and/or molecular weight of the amino acid ester or amino acid amide it is possible to vary the solubility of the oligonucleotide of the invention in water, to vary the solubility of the oligonucleotide in the organic solvent, vary the viscosity of the pharmaceutical composition comprising a solvent, and vary the ease at which the pharmaceutical composition can be drawn into a 20 gauge needle and then expelled from the 20 gauge needle.

[00624] Furthermore, by varying the lipophilicity and/or molecular weight of the amino acid ester or amino acid amide (i.e., by varying R1 of the amino acid ester or R3 and R4 of the amino acid amide) it is possible to control whether the pharmaceutical composition that further comprises an organic solvent will form a precipitate when injected into water. Although different oligonucleotides exhibit different

solubility and behavior, generally the higher the molecular weight of the amino acid ester or amino acid amide, the more likely it is that the salt of the protonated oligonucleotide and the amino acid ester of the amide will form a precipitate when injected into water. Typically, when R1 of the amino acid ester is a hydrocarbon of about C16 or higher the pharmaceutical composition will form a precipitate when injected into water and when R1 of the amino acid ester is a hydrocarbon of about C12 or less the pharmaceutical composition will not form a precipitate when injected into water. Indeed, with amino acid esters wherein R1 is a hydrocarbon of about C12 or less, the salt of the protonated oligonucleotide and the amino acid ester is, in many cases, soluble in water. Similarly, with amino acid amides, if the combined number of carbons in R3 and R4 is 16 or more the pharmaceutical composition will typically form a precipitate when injected into water and if the combined number of carbons in R3 and R4 is 12 or less the pharmaceutical composition will not form a precipitate when injected into water. Whether or not a pharmaceutical composition that further comprises a pharmaceutically acceptable organic solvent will form a precipitate when injected into water can readily be determined by injecting about 0.05 mL of the pharmaceutical composition into about 4 mL of water at about 98° F. and determining how much material is retained on a 0.22 μm filter after the composition is mixed with water and filtered. Typically, a formulation or composition is considered to be injectable when no more than 10% of the formulation is retained on the filter. In one embodiment, no more than 5% of the formulation is retained on the filter. In one embodiment, no more than 2% of the formulation is retained on the filter. In one embodiment, no more than 1% of the formulation is retained on the filter.

[00625] Similarly, in pharmaceutical compositions that comprise a protonated oligonucleotide and a diester or diamide of aspartic or glutamic acid, it is possible to vary the properties of pharmaceutical compositions by varying the amount and/or lipophilicity and/or molecular weight of the diester or diamide of aspartic or glutamic acid. Similarly, in pharmaceutical compositions that comprise an oligonucleotide; a divalent metal cation; and a carboxylate, a phospholipid, a phosphatidyl choline, or a sphingomyelin, it is possible to vary the properties of pharmaceutical compositions by varying the amount and/or lipophilicity and/or molecular weight of the carboxylate, phospholipid, phosphatidyl choline, or sphingomyelin.

[00626] Further, when the pharmaceutical compositions that further comprises an organic solvent form a depot when administered to an animal, it is also possible to vary the rate at which the oligonucleotide is released from the drug depot by varying the lipophilicity and/or molecular weight of the amino acid ester or amino acid amide. Generally, the more lipophilic the amino acid ester or amino acid amide, the more slowly the oligonucleotide is released from the depot. Similarly, when the pharmaceutical compositions that further comprises an organic solvent and also further comprise a carboxylate, phospholipid, phosphatidyl choline, sphingomyelin, or a diester or diamide of aspartic or glutamic acid and form a depot when administered to an animal, it is possible to vary the rate at which the oligonucleotide is released from the drug depot by varying the amount and/or lipophilicity and/or molecular weight of the

carboxylate, phospholipid, phosphatidyl choline, sphingomyelin, or the diester or diamide of aspartic or glutamic acid.

[00627] Release rates from a precipitate can be measured injecting about 50 μL of the pharmaceutical composition into about 4 mL of deionized water in a centrifuge tube. The time that the pharmaceutical composition is injected into the water is recorded as T=0. After a specified amount of time, T, the sample is cooled to about −9° C. and spun on a centrifuge at about 13,000 rpm for about 20 min. The resulting supernatant is then analyzed by HPLC to determine the amount of oligonucleotide present in the aqueous solution. The amount of oligonucleotide in the pellet resulting from the centrifugation can also be determined by collecting the pellet, dissolving the pellet in about 10 μL of methanol, and analyzing the methanol solution by HPLC to determine the amount of oligonucleotide in the precipitate. The amount of oligonucleotide in the aqueous solution and the amount of oligonucleotide in the precipitate are determined by comparing the peak area for the HPLC peak corresponding to the oligonucleotide against a standard curve of oligonucleotide peak area against concentration of oligonucleotide. Suitable HPLC conditions can be readily determined by one of ordinary skill in the art.

[00628] ***Methods of Treatment***

[00629] The pharmaceutical compositions of the invention are useful in human medicine and veterinary medicine. Accordingly, the invention further relates to a method of treating or preventing a condition in an animal comprising administering to the animal an effective amount of the pharmaceutical composition of the invention.

[00630] In one embodiment, the invention relates to methods of treating a condition in an animal comprising administering to an animal in need thereof an effective amount of a pharmaceutical composition of the invention.

[00631] In one embodiment, the invention relates to methods of preventing a condition in an animal comprising administering to an animal in need thereof an effective amount of a pharmaceutical composition of the invention.

[00632] Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, by inhalation, or topical. The mode of administration is left to the discretion of the practitioner. In some embodiments, administration will result in the release of the oligonucleotide of the invention, *e.g.*, a multipartite construct, an anti-C1Q oligonucleotide, a 10.36 oligonucleotide, as described above, or any combination thereof, into the bloodstream.

[00633] In one embodiment, the method of treating or preventing a condition in an animal comprises administering to the animal in need thereof an effective amount of an oligonucleotide by parenterally administering the pharmaceutical composition of the invention. In one embodiment, the pharmaceutical compositions are administered by infusion or bolus injection. In one embodiment, the pharmaceutical composition is administered subcutaneously.

[00634] In one embodiment, the method of treating or preventing a condition in an animal comprises administering to the animal in need thereof an effective amount of an oligonucleotide by orally administering the pharmaceutical composition of the invention. In one embodiment, the composition is in the form of a capsule or tablet.

[00635] The pharmaceutical compositions can also be administered by any other convenient route, for example, topically, by absorption through epithelial or mucocutaneous linings (e.g., oral, rectal, and intestinal mucosa, etc.).

[00636] The pharmaceutical compositions can be administered systemically or locally.

[00637] The pharmaceutical compositions can be administered together with another biologically active agent.

[00638] In one embodiment, the animal is a mammal.

[00639] In one embodiment the animal is a human.

[00640] In one embodiment, the animal is a non-human animal.

[00641] In one embodiment, the animal is a canine, a feline, an equine, a bovine, an ovine, or a porcine.

[00642] The effective amount administered to the animal depends on a variety of factors including, but not limited to the type of animal being treated, the condition being treated, the severity of the condition, and the specific multipartite construct being administered. A treating physician can determine an effective amount of the pharmaceutical composition to treat a condition in an animal.

[00643] In one embodiment, the multipartite construct comprises an anti-EpCAM aptamer segment. For example, the target of interest comprises EpCAM. In another embodiment, the target is selected from the group of proteins consisting of a EGFR, PBP, EpCAM, and KLK2. In another embodiment, the target is selected from the group of proteins consisting of a tetraspanin, EpCam, CD9, PCSA, CD63, CD81, PSMA, B7H3, PSCA, ICAM, STEAP, KLK2, SSX2, SSX4, PBP, SPDEF, and EGFR. In another embodiment, the target is selected from the group of proteins consisting of CD9, PSMA, PCSA, CD63, CD81, B7H3, IL 6, OPG-13, IL6R, PA2G4, EZH2, RUNX2, SERPINB3, and EpCam. In another embodiment, a target is selected from the group of proteins consisting of A33, a33 n15, AFP, ALA, ALIX, ALP, AnnexinV, APC, ASCA, ASPH (246-260), ASPH (666-680), ASPH (A-10), ASPH (D01P), ASPH (D03), ASPH (G-20), ASPH (H-300), AURKA, AURKB, B7H3, B7H4, BCA-225, BCNP1, BDNF, BRCA, CA125 (MUC16), CA-19-9, C-Bir, CD1.1, CD10, CD174 (Lewis y), CD24, CD44, CD46, CD59 (MEM-43), CD63, CD66e CEA, CD73, CD81, CD9, CDA, CDAC1 1a2, CEA, C-Erb2, C-erbB2, CRMP-2, CRP, CXCL12, CYFRA21-1, DLL4, DR3, EGFR, Epcam, EphA2, EphA2 (H-77), ER, ErbB4, EZH2, FASL, FRT, FRT c.f23, GDF15, GPCR, GPR30, Gro-alpha, HAP, HBD 1, HBD2, HER 3 (ErbB3), HSP, HSP70, hVEGFR2, iC3b, IL 6 Unc, IL-1B, IL6 Unc, IL6R, IL8, IL-8, INSIG-2, KLK2, L1CAM, LAMN, LDH, MACC-1, MAPK4, MART-1, MCP-1, M-CSF, MFG-E8, MIC1, MIF, MIS RII, MMG, MMP26, MMP7, MMP9, MS4A1, MUC1, MUC1 seq1, MUC1 seq11A, MUC17, MUC2, Ncam, NGAL, NPGP/NPFF2, OPG, OPN, p53, p53, PA2G4, PBP, PCSA, PDGFRB, PGP9.5, PIM1, PR (B), PRL, PSA, PSMA, PSME3, PTEN, R5-CD9 Tube 1, Reg IV, RUNX2, SCRN1, seprase, SERPINB3,

SPARC, SPB, SPDEF, SRVN, STAT 3, STEAP1, TF (FL-295), TFF3, TGM2, TIMP-1, TIMP1, TIMP2, TMEM211, TMPRSS2, TNF-alpha, Trail-R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, VEGF A, and YPSMA-1. In another embodiment, the target is selected from the group of proteins consisting of 5T4, ACTG1, ADAM10, ADAM15, ALDOA, ANXA2, ANXA6, APOA1, ATP1A1, BASP1, C1orf58, C20orf114, C8B, CAPZA1, CAV1, CD151, CD2AP, CD59, CD9, CD9, CFL1, CFP, CHMP4B, CLTC, COTL1, CTNND1, CTSB, CTSZ, CYCS, DPP4, EEF1A1, EHD1, ENO1, F11R, F2, F5, FAM125A, FNBP1L, FOLH1, GAPDH, GLB1, GPX3, HIST1H1C, HIST1H2AB, HSP90AB1, HSPA1B, HSPA8, IGSF8, ITGB1, ITIH3, JUP, LDHA, LDHB, LUM, LYZ, MFGE8, MGAM, MMP9, MYH2, MYL6B, NME1, NME2, PABPC1, PABPC4, PACSIN2, PCBP2, PDCD6IP, PRDX2, PSA, PSMA, PSMA1, PSMA2, PSMA4, PSMA6, PSMA7, PSMB1, PSMB2, PSMB3, PSMB4, PSMB5, PSMB6, PSMB8, PTGFRN, RPS27A, SDCBP, SERINC5, SH3GL1, SLC3A2, SMPDL3B, SNX9, TACSTD1, TCN2, THBS1, TPI1, TSG101, TUBB, VDAC2, VPS37B, YWHAG, YWHAQ, and YWHAZ. In another embodiment, the target is selected from the group of proteins consisting of CD9, CD63, CD81, PSMA, PCSA, B7H3 and EpCam. CD9, CD63, CD81, PSMA, PCSA, B7H3 and EpCam. In another embodiment, the target is selected from the group of proteins consisting of a tetraspanin, CD9, CD63, CD81, CD63, CD9, CD81, CD82, CD37, CD53, Rab-5b, Annexin V, MFG-E8,. Muc1, GPCR 110, TMEM211 and CD24 In another embodiment, the target is selected from the group of proteins consisting of A33, AFP, ALIX, ALX4, ANCA, APC, ASCA, AURKA, AURKB, B7H3, BANK1, BCNP1, BDNF, CA-19-9, CCSA-2, CCSA-3&4, CD10, CD24, CD44, CD63, CD66 CEA, CD66e CEA, CD81, CD9, CDA, C-Erb2, CRMP-2, CRP, CRTN, CXCL12, CYFRA21-1, DcR3, DLL4, DR3, EGFR, Epcam, EphA2, FASL, FRT, GAL3, GDF15, GPCR (GPR110), GPR30, GRO-1, HBD 1, HBD2, HNP1-3, IL-1B, IL8, IMP3, L1CAM, LAMN, MACC-1, MGC20553, MCP-1, M-CSF, MIC1, MIF, MMP7, MMP9, MS4A1, MUC1, MUC17, MUC2, Ncam, NGAL, NNMT, OPN, p53, PCSA, PDGFRB, PRL, PSMA, PSME3, Reg IV, SCRN1, Sept-9, SPARC, SPON2, SPR, SRVN, TFF3, TGM2, TIMP-1, TMEM211, TNF-alpha, TPA, TPS, Trail-R2, Trail-R4, TrKB, TROP2, Tsg 101, TWEAK, UNC93A, and VEGFA. In another embodiment, the target is selected from the group of proteins consisting of CD9, EGFR, NGAL, CD81, STEAP, CD24, A33, CD66E, EPHA2, Ferritin, GPR30, GPR110, MMP9, OPN, p53, TMEM211, TROP2, TGM2, TIMP, EGFR, DR3, UNC93A, MUC17, EpCAM, MUC1, MUC2, TSG101, CD63, B7H3, CD24, and a tetraspanin.

[00644] The immunosuppressive target can be a tumor-derived protein found on cMVs and/or cancer cells, including without limitation TGF-β, CD39, CD73, IL10, FasL or TRAIL.

[00645] In one embodiment, the multipartite construct can inhibit angiogenesis. In one embodiment, the multipartite construct can inhibit angiogenesis and the disease being treated is cancer. In one embodiment, the aptamer can inhibit angiogenesis and the disease being treated is a solid tumor.

The multipartite construct can be a multipartite construct that inhibits a neoplastic growth or a cancer. In embodiments, the cancer comprises an acute lymphoblastic leukemia; acute myeloid leukemia; adrenocortical carcinoma; AIDS-related cancers; AIDS-related lymphoma; anal cancer; appendix cancer;

astrocytomas; atypical teratoid/rhabdoid tumor; basal cell carcinoma; bladder cancer; brain stem glioma; brain tumor (including brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, astrocytomas, craniopharyngioma, ependymoblastoma, ependymoma, medulloblastoma, medulloepithelioma, pineal parenchymal tumors of intermediate differentiation, supratentorial primitive neuroectodermal tumors and pineoblastoma); breast cancer; bronchial tumors; Burkitt lymphoma; cancer of unknown primary site; carcinoid tumor; carcinoma of unknown primary site; central nervous system atypical teratoid/rhabdoid tumor; central nervous system embryonal tumors; cervical cancer; childhood cancers; chordoma; chronic lymphocytic leukemia; chronic myelogenous leukemia; chronic myeloproliferative disorders; colon cancer; colorectal cancer; craniopharyngioma; cutaneous T-cell lymphoma; endocrine pancreas islet cell tumors; endometrial cancer; ependymoblastoma; ependymoma; esophageal cancer; esthesioneuroblastoma; Ewing sarcoma; extracranial germ cell tumor; extragonadal germ cell tumor; extrahepatic bile duct cancer; gallbladder cancer; gastric (stomach) cancer; gastrointestinal carcinoid tumor; gastrointestinal stromal cell tumor; gastrointestinal stromal tumor (GIST); gestational trophoblastic tumor; glioma; hairy cell leukemia; head and neck cancer; heart cancer; Hodgkin lymphoma; hypopharyngeal cancer; intraocular melanoma; islet cell tumors; Kaposi sarcoma; kidney cancer; Langerhans cell histiocytosis; laryngeal cancer; lip cancer; liver cancer; malignant fibrous histiocytoma bone cancer; medulloblastoma; medulloepithelioma; melanoma; Merkel cell carcinoma; Merkel cell skin carcinoma; mesothelioma; metastatic squamous neck cancer with occult primary; mouth cancer; multiple endocrine neoplasia syndromes; multiple myeloma; multiple myeloma/plasma cell neoplasm; mycosis fungoides; myelodysplastic syndromes; myeloproliferative neoplasms; nasal cavity cancer; nasopharyngeal cancer; neuroblastoma; Non-Hodgkin lymphoma; nonmelanoma skin cancer; non-small cell lung cancer; oral cancer; oral cavity cancer; oropharyngeal cancer; osteosarcoma; other brain and spinal cord tumors; ovarian cancer; ovarian epithelial cancer; ovarian germ cell tumor; ovarian low malignant potential tumor; pancreatic cancer; papillomatosis; paranasal sinus cancer; parathyroid cancer; pelvic cancer; penile cancer; pharyngeal cancer; pineal parenchymal tumors of intermediate differentiation; pineoblastoma; pituitary tumor; plasma cell neoplasm/multiple myeloma; pleuropulmonary blastoma; primary central nervous system (CNS) lymphoma; primary hepatocellular liver cancer; prostate cancer; rectal cancer; renal cancer; renal cell (kidney) cancer; renal cell cancer; respiratory tract cancer; retinoblastoma; rhabdomyosarcoma; salivary gland cancer; Sézary syndrome; small cell lung cancer; small intestine cancer; soft tissue sarcoma; squamous cell carcinoma; squamous neck cancer; stomach (gastric) cancer; supratentorial primitive neuroectodermal tumors; T-cell lymphoma; testicular cancer; throat cancer; thymic carcinoma; thymoma; thyroid cancer; transitional cell cancer; transitional cell cancer of the renal pelvis and ureter; trophoblastic tumor; ureter cancer; urethral cancer; uterine cancer; uterine sarcoma; vaginal cancer; vulvar cancer; Waldenström macroglobulinemia; or Wilm's tumor. The compositions and methods of the invention can be used to treat these and other cancers.

Kits

[00646] The invention also provides a kit comprising one or more reagent to carry out the methods of the invention. For example, the one or more reagent can be the one or more aptamer, a buffer, blocker, enzyme, or combination thereof. The one or more reagent may comprise any useful reagents for carrying out the subject methods, including without limitation aptamer libraries, substrates such as microbeads or planar arrays or wells, reagents for biomarker and/or microvesicle isolation (e.g., via chromatography, filtration, ultrafiltration, centrifugation, ultracentrifugation, flow cytometry, affinity capture (e.g., to a planar surface, column or bead), polymer precipitation, and/or using microfluidics), aptamers directed to specific targets, aptamer pools that facilitate detection of a biomarker/microvesicle population, reagents such as primers for nucleic acid sequencing or amplification, arrays for nucleic acid hybridization, detectable labels, solvents or buffers and the like, various linkers, various assay components, blockers, and the like. The one or more reagent may also comprise various compositions provided by the invention. In an embodiment, the one or more reagent comprises one or more aptamer of the invention. The one or more reagent can comprise a substrate, such as a planar substate, column or bead. The kit can contain instructions to carry out various assays using the one or more reagent.

[00647] In an embodiment, the kit comprises an oligonucleotide probe or composition provided herein. The kit can be configured to carry out the methods provided herein. For example, the kit can include an aptamer of the invention, a substrate, or both an aptamer of the invention and a substrate.

[00648] In an embodiment, the kit is configured to carry out an assay. For example, the kit can contain one or more reagent and instructions for detecting the presence or level of a biological entity in a biological sample. In such cases, the kit can include one or more binding agent to a biological entity of interest. The one or more binding agent can be bound to a substrate.

[00649] In an embodiment, the kit comprises a set of oligonucleotides that provide a particular oligonucleotide profile for a biological sample. An oligonucleotide profile can include, without limitation, a profile that can be used to characterize a particular disease or disorder. For example, the disease or disorder can be a proliferative disease or disorder, including without limitation a cancer. In some embodiments, the cancer comprises a breast cancer.

EXAMPLES

Example 1: Identification of DNA oligonucleotides that bind a target

[00650] The target is affixed to a solid substrate, such as a glass slide or a magnetic bead. For a magnetic bead preparation, beads are incubated with a concentration of target protein ranging from 0.1 to 1 mg/ml. The target protein is conjugated to the beads according to a chemistry provided by the particular bead manufacturer. Typically, this involves coupling via an N-hydroxysuccinimide (NHS) functional group process. Unoccupied NHS groups are rendered inactive following conjugation with the target.

[00651] Randomly generated oligonucleotides (oligos) of a certain length, such as 32 base pairs long, are added to a container holding the stabilized target. Each oligo contains 6 thymine nucleotides (a "thymine tail") at either the 5 or 3 prime end, along with a single molecule of biotin conjugated to the thymine tail.

Additional molecules of biotin could be added. Each oligo is also manufactured with a short stretch of nucleotides on each end (5-10 base pairs long) corresponding to amplification primers for PCR ("primer tails"). The sequences are shown absent the thymine tails or primer tails.

[00652] The oligonucleotides are incubated with the target at a specified temperature and time in phosphate-buffered saline (PBS) at 37 degrees Celsius in 500 microliter reaction volume.

[00653] The target/oligo combination is washed 1-10 times with buffer to remove unbound oligo. The number of washes increases with each repetition of the process (as noted below).

[00654] The oligos bound to the target are eluted using a buffer containing a chaotropic agent such as 7 M urea or 1% SDS and collected using the biotin tag. The oligos are amplified using the polymerase chain reaction using primers specific to 5' and 3' sequences added to the randomized region of the oligos. The amplified oligos are added to the target again for another round of selection. This process is repeated as desired to observe binding enrichment.

Example 2: Competitive assay

[00655] The process is performed as in **Example 1** above, except that a known ligand to the target, such as an antibody, is used to elute the bound oligo species (as opposed to or in addition to the chaotropic agent). In this case, anti-EpCAM antibody from Santa Cruz Biotechnology, Inc. was used to elute the aptamers from the target EpCAM.

Example 3: Screening and Affinity Analysis

[00656] All aptamers generated from the binding assays described above are sequenced using a high-throughput sequencing platform, such as the Ion Torrent from Life Technologies:

[00657] Library Preparation - Aptamers were pooled after ligating barcodes and adapter sequences (Life Technologies) according to manufacturer protocols. In brief, equimolar pools of the aptamers were made using the following steps: Analyzed an aliquot of each library with a Bioanalyzer™ instrument and Agilent DNA 1000 Kit or Agilent High Sensitivity Kit, as appropriate for the final library concentration. The molar concentration (nmol/L) of each amplicon library was detetrmined using the commercially available software (Agilent).

[00658] An equimolar pool of the library was prepared at the highest possible concentration.

[00659] The combined concentration of the pooled library stock was calculated.

[00660] The template dilution factor of the library pool was determined using the following equation: Template Dilution Factor = (Library pool concentration [pM])/26 pM).

[00661] Template Preparation – Using a freshly diluted library, the aptamer pool resulting from binding assays provided above were sequenced using conventional sequencing protocols. High throughput (NextGen) sequencing methods can be used as desired.

[00662] Twenty aptamers were selected based on direct or competitive assays assessing binding to EpCAM (as described above).

[00663] Affinity Measurements – These twenty aptamers were then tested for binding affinity using an *in vitro* binding platform. SPR can be used for this step, e.g., a Biacore SPR machine using the T200 control software, as follows:

[00664] Dilute the antigen to a concentration of 32 nM.

[00665] Prepare necessary dilutions for kinetics, starting at 32nM prepare two-fold dilutions of antigen down to 0.5nM.

[00666] The Biacore 200 control software is programmed with the following conditions: Solution: HBS-EP+ Buffer; Number of cycles: 3; Contact time: 120s; Flow rate: 30µl/min; Dissociation time: 300s; Solution: Glycine-HCl pH 2.5; Contact time: 120s; Flow rate: 20µl/min; Stabilization period: 0s.The binding affinities of these aptamers are then measured using the SPR assay above, or an alternate in vitro assay assessing the aptamer for a desired function.

[00667] **FIG. 5** shows the SPR data for aptamer BTX176881 (SEQ ID NO: 3). The figure comprises an association and dissociation graph of 1:1 fitting model of the biotinylated aptamers to EpCAM protein at the indicated concentrations (nM). **Table 5** shows the calculated K_d values from the SPR measurements that are illustrated in **FIG. 5**. In addition, **Table 5** shows the SPR data and calculated K_d values for BTX187269 (SEQ ID NO: 6) and Aptamer 4 (SEQ ID NO. 1).

Table 5: Calculated K_D values from SPR measurements

Immobilized aptamer	Analyte	Conc (nM)	Response	K_d (nM)	Full R^2	Full Chi^2
BTX176881 (SEQ ID No: 3)	EpCAM protein	500	0.2434	8.40	0.989322	0.179008
		250	0.136	8.40	0.989322	0.179008
		100	0.0776	8.40	0.989322	0.179008
BTX187269 (SEQ ID NO: 6)	EpCAM protein	500	0.2575	7.12	0.990323	0.215697
		250	0.1584	7.12	0.990323	0.215697
		100	0.0551	7.12	0.990323	0.215697
Aptamer 4 (SEQ ID NO. 1)	EpCAM protein	500	0.2742	10.10	0.986276	0.299279
		250	0.1618	10.10	0.986276	0.299279
		100	0.0809	10.10	0.986276	0.299279

[00668] *K_d, R^2 and Chi^2 values by Global fitting for single reference method.

Example 4: Motif analysis

[00669] The process of **Example 3** is followed to identity a high affinity aptamer to a target of interest. Once a high affinity aptamer is identified, its sequence is then analyzed using a software program to estimate its two-dimensional folding structure. Well-known sequence alignment programs and algorithms for motif identification can be used to identify sequence motifs and reduce the dimensionality of even large data sets of sequences. Further, software programs such as Vienna and mfold are well-known to those skilled in the art of aptamer selection and can be used to further group sequences based on secondary structure motifs (shared shapes). See **FIG. 3A** and **FIG. 3B** for example structure predictions. Shared secondary structure of course, does not guarantee identical three-dimensional structure. Therefore "wet-lab" validation of aptamers is still useful as no one set of *in silico* tools has yet been able to fully predict the optimal aptamer among a set of aptamer candidates.

Example 5: Microvesicle-based aptamer subtraction assay

[00670] Circulating microvesicles are isolated from normal plasma (e.g., from individuals without cancer) using one of the following methods: 1) Isolation using the ExoQuick reagent according to manufacturer's protocol; 2) Ultracentrifugation comprising spin at 50,000 to 150,000g for 1 to 20 hours then resuspending the pellet in PBS; 3) Isolation using the TEXIS reagent from Life Technologies according to manufacturer's protocol; and 4) filtration methodology. The filtration method is described in more detail as follows:

[00671] Place syringe and filter (1.2 μm Acrodisc Syringe Filter Versapor Membrane Non-Pyrogenic Ref: 4190, Pall Life Sciences) on open 7 ml 150K MWCO column (Pierce concentrators, 150K MWCO (molecular weight cut off) 7 ml. Part number: 89922). Fill open end of syringe with 5.2 ml of filtered 1X PBS prepared in sterile molecular grade water.

[00672] Pipette patient plasma (900-1000 μl) into the PBS in the syringe, pipette mix twice

[00673] Filter the plasma into the 7 ml 150K MWCO column.

[00674] Centrifuge 7 ml 150K MWCO columns at 2000 x g at 20°C (16°C to 24°C) for 1 hour.

[00675] After 1 hour spin, pour the flow-through into 10% bleach to be discarded.

[00676] Visually inspect sample volume. If plasma concentrate is above the 8.5 ml graduation on the concentrator tube, continue to spin plasma sample at 10 minute increments at 2000 x g at 20°C (16°C to 24°C) checking volume after each spin until plasma concentrate is between 8.0 and 8.5 mls.

[00677] Pipette mix slowly on the column a minimum of 6 times and adjust pipette to determine plasma concentrate volume. If volume is between 100 μl and Target Volume, transfer plasma concentrate to previously labeled co-polymer 1.5 ml tube. If volume is still greater than Target Volume, repeat the above centrifugation step.

[00678] Pour ~45 mls of filtered 1X PBS prepared in sterile molecular grade water into 50 ml conical tube for use in the next step.

[00679] Add the appropriate amount of filtered 1X PBS to reconstitute the sample to the Target Volume.

[00680] The microvesicles produced using any of the isolation methods will comprise a mixture of vesicle types and will be various sizes with the possible exception of ultracentrifugation methods, which may favor isolating exosome size particles.

[00681] Randomly generated oligonucleotides (produced as described in **Example 1** above) are incubated with the isolated normal vesicles in PBS overnight at room temperature or at 4 degrees Celsius.

[00682] The aptamers that do not bind to these vesicles are isolated by spinning down the vesicles at 50,000 to 150,000 X g for 1 to 20 hours and collecting the supernatant.

[00683] The aptamer oligonucleotides are collected from the supernatant by running the mixture over a column containing streptavidin-coated beads. These aptamers are then added to vesicles isolated from diseased patients (using the same methods as above) and incubated overnight in PBS at room temperature or 4 degrees Celsius.

[00684] The vesicles are then spun at 50,000 to 150,000 X g for 1 to 20 hours and the supernatant is discarded. The vesicles are resuspended in PBS and lysed using SDS or some similar detergent.

[00685] The aptamers are then captured by running the lysis mixture over a column of streptavidin-coated beads. The isolated aptamers are then subjected to a round of PCR to amplify the products.

[00686] The process is then repeated for a set number of times, e.g., 5 times. The remaining aptamer pool has been depleted of aptamers that recognize microvesicles found in "normal" plasma. Accordingly, this method can be used to enrich the pool in aptamers that recognize cancer vesicles. See **FIG. 4**.

Example 6: Detection of Microvesicles using anti-EpCAM aptamers

[00687] Aptamers can be used as binding agents to detect a biomarker. In this Example, aptamers are used as binding agents to detect EpCAM protein associated with microvesicles.

[00688] **FIGs. 6A-D** illustrate the use of an anti-EpCAM aptamer (Aptamer 4; SEQ ID NO. 1) to detect a microvesicle population in plasma samples. Plasma samples were obtained from three men with prostate cancer and three men without prostate cancer (referred to as controls or normals). Antibodies to the following microvesicle surface protein antigens of interest were conjugated to microbeads (Luminex Corp, Austin, TX): **FIG. 6A**) EGFR (epidermal growth factor receptor); **FIG. 6B**) PBP (prostatic binding protein; also known as PEBP1 (phosphatidylethanolamine binding protein 1)); **FIG. 6C**) EpCAM (epithelial cell adhesion molecule); and **FIG. 6D**) KLK2 (kallikrein-related peptidase 2). Microvesicles in the plasma samples were captured using the bead-conjugated antibodies. Fluorescently labeled Aptamer 4 was used as a detector in the microbead assay. **FIGs. 6A-D** show the average median fluorescence values (MFI values) detected for the bead-captured and Aptamer 4 detected microvesicles. Each plot individually shows the three cancer (C1-C3) and three normal samples (N1-N3). These data show that, on average, the prostate cancer samples have higher levels of microvesicles containing the target proteins than the normals.

Example 7: Negative and Positive Selection of Aptamers

[00689] Aptamers can be used in various biological assays, including numerous types of assays which rely on a binding agent. For example, aptamers can be used instead of antibodies in immune-based assays. This Example provides an aptamer screening method that identifies aptamers that do not bind to any surfaces (substrates, tubes, filters, beads, other antigens, etc.) throughout the assay steps and bind specifically to an antigen of interest. The assay relies on negative selection to remove aptamers that bind non-target antigen components of the final assay. The negative selection is followed by positive selection to identify aptamers that bind the desired antigen.

[00690] Preliminary experiments were done with five DNA aptamer libraries with 10^{15} sequences each and variable lengths (60, 65, 70, 75, 80-mers) were pre-amplified and strand separated so that forward strand (non-biotinylated) serves as an aptamer. Multiple rounds of negative selection and positive selection were performed. Before each round, the recovered aptamer products were PCR amplified and strand separated using standard methodology. Selections were performed as follows:

[00691] Negative selection

[00692] 1. Prepare bead negative Selection Mix: Incubate 1200 non-magnetic beads with standard blocking agent for 20 min.

[00693] 2. Add 50 µl of aptamer library (5 libraries total) to a PCR strip tube with 4.5 µl of each bead mixture. Incubate for 2 h at 37°C with agitation at 550 rpm.

[00694] 3. Pre-wet filter plate (1.2 µm, Millipore) with PBS-BN buffer. Add 150 µl PBS-BN.

[00695] 4. Transfer samples from the PCR strip tubes to the filter plate, incubate for 1 h at room temperature with agitation at 550 rpm.

[00696] 5. Collect flow-through from filter plate into a collection (NBS) plate using a vacuum manifold.

[00697] 6. Concentrate and clean samples to remove excess materials as desired.

[00698] The negative selection process is repeated up to 6-7 times.

[00699] Positive selection

[00700] Before starting, conjugate the protein biomarkers of interest (here, SSX4, SSX2, PBP, KLK2, SPDEF) to desired non-magnetic microbeads using conditions known in the art. The recombinant purified starting material included: SPDEF recombinant protein from Novus Biologicals (Littleton, CO, USA), catalog number H00025803-P01; KLK2 recombinant protein from Novus, catalog number H00003817-P02; SSX2 recombinant protein from Novus, catalog number H00006757-P01; PBP recombinant protein from Fitzgerald Industries International (Action, MA, USA), catalog number 30R-1382; SSX4 recombinant protein from GenWay Biotech, Inc. (San Diego, CA, USA), catalog number GWB-E219AC.

[00701] 1. Bead blocking: Incubate a desired number of each bead (8400 x number of aptamer libraries (5) x an overage factor of (1.2)) with a starting block for 20 min.

[00702] 2. Mix 50 µl of each aptamer library sample to PCR strip tubes add 2.3 µl of bead sample with particular antigen. Incubate for 2 h at 37°C with agitation at 550 rpm.

[00703] 3. Pre-wet filter plate (1.2 µm, Millipore) with PBS-BN buffer. Add 150 µl PBS-BN.

[00704] 4. Transfer samples from the PCR strip tubes to the filter plate, incubate for 1 h at room temperature with agitation at 550 rpm.

[00705] 5. Wash 3x with PBS-BN, add 50 µl of PBS and collect samples from the top of the filter to the 1.5 ml tubes.

[00706] The positive selection is repeated up to 16 times. Certain rounds of positive selection have additional steps to treat the recovered RNA (i.e., remaining aptamer candidates) as follows:

[00707] Round 8 of positive selection was modified as follows:

[00708] 1. After the third wash (PBS-BN) 25 µl of sample were collected from the top of the filter into 1.5 ml tubes.

[00709] 2. The filter plate was incubated at 45°C for ~10 min and washed immediately using vacuum. The plate was washed three more times with PBS-BN.

[00710] 3. 50 µl of PBS were added to the plate and step 2 was repeated.

[00711] 4. After the last wash, 25 µl of PBS was added to the wells. The samples were mixed well and collected from the top of the filter into 1.5 ml tubes.

[00712] Round 9 of positive selection was modified as follows:

[00713] 1. After the final wash in step 5), 5 µg/ml Streptavidin-PE was added to the aptamer mixture and incubated for 30 min at room temperature with agitation at 550 rpm.

[00714] 2. Samples on filter plate were washed 3x with PBS-BN (+ additional 500 mM NaCl).

[00715] 3. One additional wash with regular PBS-BN was performed.

[00716] 4. 50 µl of PBS was added to the samples followed by collection as above into 1.5 ml tubes.

[00717] 5. Samples stored at -20°C.

[00718] Round 14 of positive selection was modified as follows:

[00719] Before start this round, the antigens of interest (SSX4, SSX2, PBP, KLK2, SPDEF) were conjugated to carboxylated magnetic beads using methods known in the art.

[00720] 1. Bead blocking: take desired number of each non-magnetic bead (3000 x number of aptamer libraries (5) x an overage factor of 1.2), add starting block (3:1, blocking per 1200 beads), make 5 mixes of 4 antigens and supplement each with different target antigen conjugated to magnetic beads (see **Table 6** below, wherein the antigens are conjugated to non-magnetic beads except as indicated), incubate 20 min.

Table 6: Bead blocking mixtures

Blocking Mix	Non-magnetic bead antigens	Magnetic bead antigens
1	SSX4 + PBP + KLK2 + SPDEF	SSX2
2	SSX2 + PBP + KLK2 + SPDEF	SSX4
3	SSX2 + SSX4 + KLK2 + SPDEF	PBP
4	SSX2 + SSX4 + PBP + SPDEF	KLK2
5	SSX2 + SSX4 + PBP + KLK2	SPDEF

[00721] 2. Add 50 µl of aptamer libraries to PCR strip tubes, add bead mixtures with target antigen on magnetic beads to the tubes with pre-selected corresponding aptamer library and incubate for 2 h at 37°C with agitation at 550 rpm.

[00722] 3. Pre-wet filter plate with PBS-BN buffer, add 150 µl PBS-BN.

[00723] 4. Transfer samples from PCR strip tubes to filter plate, incubate 1h room temperature with agitation at 550 rpm.

[00724] 5. After last (standard) wash, add 5µg/ml Streptavidin-PE, incubate for 30 min room temperature with agitation at 550 rpm;

[00725] 6. Wash 3x with PBS-BN (+ additional 500 mM NaCl).

[00726] 7. Perform one additional wash with regular PBS-BN.

[00727] 8. 50 µl of PBS was added to the samples followed by collection as above into 1.5 ml tubes.

[00728] 9. Remove the magnetic beads using a magnetic stand, and replace with fresh PBS buffer.

[00729] 10. Samples stored at -20°C for subsequent DNA extraction and strand separation.

[00730] Optional steps implemented in the later round of positive selection are intended to increase stringency of aptamer binding (e.g., increased heat or salt concentration).

Example 8: Discovery and Characterization of anti-EpCAM aptamers

[00731] In this Example, an aptamer to EpCAM identified using the technique in the Example above is characterized. After selection for a pool of EpCAM binding aptamers as described above, the aptamer library was sequenced using the Ion Torrent standard protocol (Life Technologies, Carlsbad, CA). Lead candidates were selected as those having (a) high abundant motifs across all read sequences with full expected length product and (b) strong secondary structure (**FIG. 7B**).

[00732] Aptamers wre selected for EPCAM protein conjugated to MicroPlex beads in competition with SSX4, SSX2, PBP, KLK2, and SPDEF recombinant proteins. A portion of the aptamers was selected in initial rounds against EpCAM that was attached to an Fc tag, and after round 8 the selection was switched to EPCAM with a Histidine tag. Another portion of the aptamers was selected in initial rounds against EpCAM that was attached to a Histidine tag, and after round 8 the selection was switched to EPCAM with an Fc tag. Methods of using Fc and histidine tags for protein purification and capture are known to those of skill in the art.

[00733] Aptamer Characterization

[00734] CAR003 is an aptamer candidate identified using the above methodology. As an RNA aptamer, CAR003 with alternate tail sequence has the following RNA sequence (SEQ ID NO. 5):

[00735] 5′-auccagagug acgcagcagu cuuuucugau ggacacgugg uggucuagua ucacuaagcc accgugucca-3′

[00736] CAR003 was further characterized. EpCAM aptamer CAR003 is modified as desired by attachment of a biotin moiety on the 5'end or 3' end. The biotin can be used to bind the aptamer using a streptavidin-biotin system, e.g., for labeling, capture and/or anchoring. **FIG. 7B** illustrates the optimal secondary structure of CAR003 with a minimum free energy (ΔG) of -30.00 kcal/mol. For purposes of illustration, the aptamer is shown as an RNA aptamer (SEQ ID NO. 5) corresponding to the CAR003 DNA sequence (SEQ ID NO. 4).

[00737] Synthesis and purification. The selected CAR003 aptamer was re-synthesized using AKTA OligoPilot 100 Synthesizer (GE Healthcare Life Sciences Corp., Piscataway, NJ) with a 3'Biotin and final detritylation. The product was purified with anion exchange chromatography by FPLC. Several fractions after FPLC were combined as shown as the indicated Pools 1-3 in **FIG. 7C**. The figure comprises an FPLC chromatogram with all product and fractions assigned in pools after checking quality on gel. **FIG. 7D** illustrates a SYBR GOLD stained gel with different FPLC fractions of CAR003 aptamer after synthesis. Different fractions were combined in pools based on amount of unfinished chains in order high to low (pool 1 – pool 3). The pools 1-3 correspond to those indicated in **FIG. 7C**.

[00738] CAR003 aptamer characterization. Purified CAR003 aptamer was tested for binding to recombinant EPCAM protein with a polyhistidine tag ("His tagged") using the following internally developed assay. Anti-His tag conjugated beads were mixed with EPCAM-His tagged protein. The

aptamer to be tested was labeled with streptavidin-phycoerythrin (SA-PE). The EpCAM-beads and SA-PE labeled aptamers were mixed. Binding was determined as median flourescent value in a bead assay as described herein. MFI values (**FIG. 7E-F**) increase with increased binding of the SA-PE labeled aptamer to the recombinant EpCAM. **FIG. 7E-F** illustrate binding of CAR003 to EPCAM protein in 25 mM HEPES with PBS-BN (PBS, 1% BSA, 0.05% Azide, pH 7.4) (**FIG. 7E**) or in 25 mM HEPES with 1 mM $MgCl_2$ (**FIG. 7F**). EPCAM aptamer Aptamer 4 (see above) was used for comparison. As shown in the figures, CAR003 pool 3 more efficiently binds its target in the presence of $MgCl_2$ (**FIG. 7F**) than in the presence of BSA (**FIG. 7E**).

[00739] To understand its performance further, CAR003 binding was tested in the presence of both BSA and $MgCl_2$ in various buffers. **FIG. 7G** illustrates CAR003 binding to EpCAM in the indicated salts with and without addition of bovine serum albumin (BSA). Again, CAR003 binding to EpCAM is more efficient when BSA is not present. Additionally, 150 mM NaCl was tested but did not appear to improve CAR003 performance over $MgCl_2$.

[00740] Another factor which might influence performance of aptamer is denaturing with different salt compositions. **FIG. 7H** illustrates the effect of denaturing on CAR003 binding to EPCAM protein. As seen from the chart, denaturing of the aptamer has a postive effect on CAR003 binding to EpCAM similar as the effect on CAR003 from $MgCl_2$. However, denaturing in the presence of $MgCl_2$ may not synergistically improve binding of CAR003 to EpCAM. Interestingly, CAR003 appeared more stable compared to control Aptamer 4 in the conditions tested.

[00741] CAR003 affinity to EpCAM in the bead assay environment was assessed in the same assay as above with aptamer titrated across a constant input of antigen. **FIG. 7I** illustrates titration of aptamers against EPCAM recombinant protein (constant input 5 μg). Under the conditions tested, Aptamer 4 had a higher affinity to EPCAM protein compared to CAR003 as suggested from saturation level starting at 5 μg of aptamer input.

[00742] In order to evaluate specificity of CAR003, it was tested using Western Blot against EPCAM recombinant protein, and controls comprising bovine serum albumin (BSA) and human serum albumin (HSA). **FIG. 7J** illustrates a Western blot with CAR003 aptamer versus EPCAM his-tagged protein, BSA, and HSA (5 μg each). The gel was blocked 0.5% F127 and probed with ~50 μg/ml CAR003 biotinylated aptamer, fraction 3. The blot was visualized with NeutrAvidin-HRP followed by SuperSignal West Femto Chemiluminescent Substrate. The Western blot probed with CAR003 aptamer showed a clear preference of the aptamer to EPCAM protein over the albumins.

[00743] CAR003 test with plasma samples. Plasma samples from five prostate cancer and five normal subjects were tested with CAR003 to detect microvesicles using bead-conjugated proteins to capture the microvesicles and SA-PE labeled aptamer to detect the vesicles. SA-PE labeled Aptamer 4 detector was used as control. Fold changes of Cancer over Normal are shown in **Table 7**. The fold changes are shown without normalization ("Raw") or with normalization to a negative control. The vesicles were captured with bead conjugated antibodies to SSX4, PBP, SPDEF, EPCAM, KLK2 and SSX2 as indicated.

Table 7: CAR003 to detect microvesicles

		SSX4	PBP	SPDEF	EPCAM	KLK2	SSX2
Raw	**Standard protocol**	0.87	0.39	0.71	0.63	0.93	0.87
	Incubation in presence of 1mM MgCl2 and absence of PBS-BN	0.77	0.39	0.69	0.6	0.91	0.81
	Aptamer 4 control (standard protocol)	0.78	0.67	0.81	0.72	1.19	0.79
Normalized to Negative control	**Standard protocol**	1.49	0.84	1.13	1.17	1.5	1.38
	Incubation in presence of 1mM MgCl2 and absence of PBS-BN	1.27	0.83	1.08	1.1	1.46	1.29
	Aptamer 4 control (standard protocol)	1.18	0.96	1.11	1.04	1.82	1.1

[00744] Under the conditions tested, the samples detected with CAR003 had lower MFI values as compared to detection with Aptamer 4, whereas CAR003 had a better signal-to-noise ratio and showed better separation between cancer and normal samples with SSX4, SPDEF, EPCAM and SSX2 capturing markers.

[00745] Control Aptamer

[00746] The characteristics of the aptamers (size, stability, binding affinity and specificity, etc) can be compared against control aptamers specific to EpCAM or other targets. For example, the aptamers are compared to the anti-VEGF aptamer 5' biotin-CA ATT GGG CCC GTC CGT ATG GTG GGT (SEQ ID NO. 7) as described in Kaur and Yung, 2012.

[00747] References:

1) Müller, J., et al. "Selection of high affinity DNA-aptamer for activated protein C using capillary electrophoresis." *Research in Pharmaceutical Sciences* 7.5 (2012): S987.

2) Cerchia, L., and V. de Franciscis. "Nucleic Acid Aptamers Against Protein Kinases." *Current medicinal chemistry* 18.27 (2011): 4152-4158.

3) Wu, Jie, et al. "Identification, Characterization and Application of a G-Quadruplex Structured DNA Aptamer against Cancer Biomarker Protein Anterior Gradient Homolog 2." *PloS ONE* 7.9 (2012): e46393

4) Mitkevich, Olga V., et al. "DNA aptamers detecting generic amyloid epitopes." *Prion* 6.4 (2012): 400-406.

5) Kaur H, Yung L-YL (2012) Probing High Affinity Sequences of DNA Aptamer against $VEGF_{165}$. PLoS ONE 7(2): e31196. doi:10.1371/journal.pone.0031196.

Example 9: Aptamer Target Identification

[00748] In this Example, aptamers conjugated to microspheres are used to assist in determining the target of two aptamers identified by library screening methods as described above. The general approach is shown in **FIG. 9**. The approach is used to verify the targets of CAR003, an aptamer identified by library screening to recognize EpCAM. See decription above for CAR003. In this approach, the sequence of CAR003 is randomly rearranged before linkage to the microspheres. The microspheres are used as controls to bind to targets that are similar but not identical to the intended target molecule.

[00749] The protocol used is as follows:

[00750] 1) The candidate aptamers (here, CAR003) and negative control aptamers (here, randomly arranged CAR003) are synthesized with modifications to allow capture (here, the aptamers are biotinylated) and crosslinking (here, using the Sulfo-SBED Biotin Label Transfer Reagent and Kit, Catalog Number 33073 from Thermo Fisher Scientific Inc., Rockford, IL, to allow photocrosslinking).

[00751] 2) Each of the aptamers is individually mixed with microvesicles having the target of interest (here, BrCa cell line microvesicles).

[00752] 3) After incubation to allow the aptamers to bind target, ultraviolet light is applied to the mixtures to trigger crosslinking of the aptamers with the microvesicle targets.

[00753] 4) The microvesicles are lysed, thereby releasing the crosslinked aptamer-target complex into solution.

[00754] 5) The crosslinked aptamer-target complexes are captured from solution using a streptavidin coated substrate.

[00755] 6) The crosslinked aptamer-target complexes for each aptamer are run individually on SDS-PAGE gel electrophoresis. The captured protein targets are visualized with Coomasie Blue staining.

[00756] 7) The crosslinking and binding steps may be promiscuous so that multiple bands including the intended target but also random proteins will appear on each of the gels. The intended target will be found in a band that appears on the gel with the candidate aptamer (here, CAR003) but not the related negative control aptamers (here, randomly arranged CAR003). The bands corresponding to the target are excised from the gel.

[00757] 8) Mass spectrometry (MS) is used to identify the aptamer target from the excised bands.

Example 10: Aptamers to Breast Cancer (BrCa) derived microvesicles

[00758] In this Example, an aptamer library is screened to identify aptamers that distinguish between microvesicles circulating in the blood of breast cancer patients and microvesicles circulating in the blood of healthy, control individuals (i.e., without breast cancer).

[00759] Microvesicles were isolated from plasma of a pool of 60 breast cancer patients (BrCa+). Microvesicles were also isolated from pool of 60 non-cancer samples (BrCa-). Microvesicles were

isolated from the plasma using ultracentrifugation (120,000 x g). Microvesicles were in the pellet from the ultracentrifugation. The supernatant from the ultracentrifugation was saved to use as a control. The microvesicles from both sample types were conjugated to MagPlex beads (Luminex Corp, Austin TX). Optionally, the isolated microvesicles are incubated with anti-HSA/IgG/Fibrinogen beads to remove these highly abundant blood proteins. However, the conjugation step can be optimized to favor conjugation of the microvesicles such that removal of highly abundant proteins may be less of an issue.

[00760] The aptamer library used consisted of a 2'F SUL1 RNA aptamer library. The sequence is 5'-GGGAGGACGAUGCGG-N40-CAGACGACUCGCUGAGGAUCCGAGA-3' (SEQ ID NO. 8). The aptamer library consists of three sections: Forward primer – 15 nucleotides, variable region – 40 nucleotides; reverse primer – 25 nucleotides. All pyrimidines (C and U) were 2'Fluoro modified.

[00761] The aptamer library was incubated with either the cancer or control microvesicle-conjugated beads. Thirteen rounds of positive selection for aptamers that bind the microvesicles were performed in parallel for both types of samples. See **Example 11** below for detailed protocol of the positive selection steps. Negative selection was not performed.

[00762] A number of representative sequences obtained from these procedures are shown in **Table 8**. The sequences in the table were identified in the aptamer pools from selection against BrCa microvesicles but were not in the aptamer pools selected against non-cancer samples. In **Table 8**, the sequences are shown 5' to 3' from left to right, wherein each complete sequence consists of a 5' leader sequence 5'-GGGAGGACGAUGCGG (SEQ ID NO. 9) followed by the indicated Variable Sequence followed by the 3' tail sequence 5'-CAGACGACUCGCUGAGGAUCCGAGA (SEQ ID NO. 10). Each sequence is derived from a library having a leader and tail (see description above) with a variable sequence between. It is understood that the nucleotide sequences that are disclosed in **Table 8** can also be modified to the extent that resulting modifications result in an aptamer having about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, and 99 percent homology to the disclosed sequence and retain the functionality of binding to microvesicle antigens or functional fragments thereof.

Table 8: BrCa microvesicle aptamer candidate sequences

ID	SEQ ID NO.	Variable Sequence
BRCA_APT1_RNA	11	CGCGUCUUCCCCGCAUUGCCGCAAUUGCCAUACAUUAAUA
BRCA_APT2_RNA	13	GUCCGGAACGCCUCGAUCCUCGCAUAAUAUGAUACGUCUG
BRCA_APT3_RNA	15	GUCCAUGGUACGCCUCGAUUCCGCCCAUACAUGCAUGUAA
BRCA_APT4_RNA	17	CACUAUCCGUUUGUCCGUCCUCUUGUGGUAUUGCGCAUGC
BRCA_APT5_RNA	19	UCUUCCAUCUGGUCGCGAUACAGAAUACGAUUAACAUAAA
BRCA_APT6_RNA	21	GAUCACGCUGCCCUUUGUUUAAGGCCUUUAUACAAACGCA
BRCA_APT7	23	UAUUCGCCAGUCACAUCAACUAUGAUGACGCUUGACUGGA

_RNA		

[00763] Each sequence in **Table 8** is synthesized in two variants for further investigation: 5' biotinylated and 3' biotinylated. This provides aptamer variants that can be captured at the 5' end or the 3' end as desired. The aptamers are further synthesized with each pyrimidine (C and U) 2'Fluoro modified.

[00764] The DNA sequence corresponding to each RNA sequence in **Table 8** is provided in the sequence listing, where the DNA sequence directly follows its corresponding RNA sequence. For example SEQ ID NO. 12 is the DNA sequence corresponding to RNA sequence SEQ ID NO. 11, etc. The DNA forms of the aptamers are synthesized for further characterization as well.

[00765] The aptamers above were identified using positive selection for aptamers which recognize BrCa and non-BrCa microvesicles conjugated to microspheres.

Example 11: Aptamer library selection protocol

[00766] This Example provides the protocol for SUL1 RNA library selection performed in the Example above. The protocol can be followed for other aptamer libraries and sample input as desired.

[00767] Preparation

[00768] The working space is cleaned with 80 % EtOH before working.

[00769] Beads are MagPlex beads (Luminex Corp., Austin, TX). Other beads can be substituted as desired.

[00770] Buffers/Reagents to Prepare:

- MilliQ water
- 100mM $MgCl_2$
- 5x Transcription Buffer (200mM Tris pH 7.9)
- 1x PBS
- 1x PBS with 3mM $MgCl_2$
- 10x PBS
- Selection buffer (1x PBS with 0.1% BSA and 3mM $MgCl_2$)

[00771] Before starting with selection, remove the bead storage buffer, and wash beads with 1x PBS w/ 3mM $MgCl_2$ 1 times (200uL total in all 4 tubes). 200,000 beads per selection are used.

[00772] Binding 2'F SUL1 RNA pool to microvesicle coated magnetic beads

[00773] Abbreviations: TK- Transcription; NTC- No template control.

[00774] Steps:

1. **1st Round:** Mix 1nmol purified 2'F SUL1 RNA with 20 µl of resuspended beads (conjugated with microvesicle). 10uL of 10x PBS +1% BSA, 3µl 100 mM $MgCl_2$, and 47uL H_2O. This gives a final concentration of 1x PBS, 0.1% BSA, 3mM $MgCl_2$.

    1.1 The addition of $MgCl_2$ in this step gives a concentration of 3mM $MgCl_2$. This is the binding concentration for the entire process.

    1.2 **Following Rounds:** Mix 20 µl of transcription product (15 mM $MgCl_2$ inside) with 20 µl of washed microvesicle coated beads, plus 9uL 10x PBS with 1% BSA, 51uL H_2O. No additional $MgCl_2$ is needed because the $MgCl_2$in the diluted transcription product (TK) provides a final concentration of 3mM $MgCl_2$.

2. Incubate for 30 min at 37 °C, shake at 1000rpm, and pipet mix every 10 minutes.
3. Wash the beads:
   3.1 **One washing cycle comprises**:
      3.1.1 Remove the beads from the magnet
      3.1.2 Resuspend beads in 100µl 1x PBS +3mM $MgCl_2$ off the magnet.
      3.1.3 Incubate sample for 30 seconds off of the magnet.
      3.1.4 Place the sample back onto the magnet, and wait until the beads are on the side.
      3.1.5 Remove and discard the supernatant.
      3.1.6 Resuspend in 100µl 1x PBS +3 mM $MgCl_2$ + 0.1% BSA off of the magnet.
      3.1.7 Incubate sample for 3 minutes off of the magnet.
      3.1.8 Place the sample back onto the magnet, and wait until the beads are on the side.
      3.1.9 Remove and discard the supernatant
   3.2 **1st Round:** Place bead mixture on a magnet and remove the supernatant. Wash once with 100 µl 1x PBS +3 mM $MgCl_2$ +0.1% BSA (by pipette mixing the beads), and discard buffer.
   3.3 **Following Rounds:** Increase the washing steps every second round by one more washing step up to 3 washing steps.
4. Add 55 µl MilliQ water to the bead sample.
5. Elute the RNA by incubating the bead sample for 5 min at 80 °C
   5.1 Check if there is 50 µl, if not spin the sample down to spin down the condensed water off the top.
   5.2 Transfer the supernatant to a new vial. Work quickly to avoid the strands rebinding the beads.
      5.2.1 Use 50 µl eluate for the following RT-PCR and store the rest at -20°C

[00775] **RT-PCR of recovered aptamer candidates**

[00776] Tips

- The rest of the RT-PCR sample and the TK-PCR sample is stored at -20 °C
- RNA can be stored at 4°C for ~1h
- RT-PCR product can be stored overnight at 4°C
- Proceed to the next selection cycle for optimal RNA quality immediately after transcription.
- Avoid vortexing RNA
- Mix on ice
- Use 0.5ml PCR tubes
- Every RT-PCR should have a no-template control (NTC) with water instead of template
- Do not freeze-thaw DTT more than one time

6. Prepare a Master Mix before the first round per **Table 9**, check it with 0.5 pmol RNA and store aliquots of 48 µl at -20°C until usage.

Table 9: RT-PCR master mix

Reagent	Volume (µl)/reaction	Final concentration
5x Colorless GoTaq Flexi Buffer Promega cat# M890A	20	1x
5 x first strand buffer (Invitrogen) lot# 1300427	4	0.2 x
100 mM DTT	2	2 mM
100 µM SUL1 F primer	1	1 µM
100 µM SUL1 R primer	1	1 µM
100 mM $MgCl_2$	1.5	1.5 mM
25 mM (**each**) dNTPs	1.2	300 µM
MilliQ water	17.3	
Total	48	

7. Add 50 µl MilliQ water as negative control (NTC) (pipette this first) **or** 50 µl selection eluate. Pipet mix.
8. Incubate at 65 °C for 5 min.
9. After cooling to 4°C, add:
   9.1 1 µl Superscript II Reverse Transcriptase (Invitrogen, cat# 18064) (200 U/µl)
   9.2 1 µl GoTaqFlexi DNA polymerase (5 units/µl) Promega cat# M8305.

PCR –Program (SARTPCR)

a) 10 min 54 °C
   (This step is only for reverse transcriptase, should more rounds be needed, do not repeat step A.)
b) 1 min 95 °C
c) 1 min 60 °C
d) 1 min 72 °C

10. Cycle steps b-d for
    10.1 **1st round** b-d 4 cycles. Run 5 µL PCR products on a 4% agarose gel.
        10.1.1 Subsequent rounds: The amount of RNA is decreased after the first round, leading to an increase in required PCR-cycles. To determine the number of cycles needed each time, check the band intensity from the agarose gel from the previous round of selection. Use that number of cycles to start the next round of RT-PCR. Note: Always check results on an agarose gel.
            10.1.1.1 Agarose gel results: product band should be seen at the target length. The band intensity should be about the same as the 50bp ladder band (if not a little less intense). If the band is not intense enough (barely visible), cycle an appropriate amount more and re-check on an agarose gel.

[00777] Transcription

[00778] All mixing performed on ice. Prepare transcription Master Mix per **Table 10** and store aliquots of 85.7 µl at -20°C until use.

11. Verify pH of stock 200 mM Tris pH 7.9 before use. A change in pH over time may cause problems with the transcription.

Table 10: Transcription (TK) Master Mix for SUL1 library

Reagent	Volume (µl) for one reaction	Volume (µl) for 20 reactions	Final concentration
5x Transcription buffer (200 mMTris, pH 7.9)	20	400	1x
100 mM DTT	5	100	5 mM
100mM ATP	1	20	1 mM
100mM GTP	1	20	1 mM
100mM 2'F-dUTP	3	60	3 mM
100mM 2'F-dCTP	3	60	3 mM
100 mM $MgCl_2$	15	300	15 mM
MilliQ water	37	740	
Total volume	85	1700 µl	

12. Add 10 µl RT-PCR product to the mastermix.
13. Add 1 µl RNasin (40 units/µl)

13.1 Promega Recombinant RNasin Ribonuclease Inhibitor cat# N2515/N2511
14. Add 4 µl T7 Y639F mutant polymerase (25U/µl use: 100U total per reaction)
15. Perform the reaction for 30 min at 37 °C
16. Use the transcription-product directly for the next selection round. If the next step is not feasible, freeze transcription product at -20C.

[00779] Subsequent Rounds

[00780] Repeat the bead incubation, the RT-PCR and transcription as often as needed. Try to have similar band intensity of the RT-PCR product for the sample in all rounds as noted above.

[00781] Binding Assay

[00782] A binding assay is performed after desired rounds of selection to determine to assess non-specific binding of cancer selected aptamers to control beads (conjugated to supernatant from plasma ultracentrifugation, see above) and likewise for non-cancer control samples. Binding assays can also be performed to assess binding of selected aptamers against the intended target microvesicles.

[00783] Cherenkov protocol: Performed using ^{32}P radioactively labeled aptamer library.

[00784] Final concentration of selection buffer: 1x PBS+ 3mM $MgCl_2$ + 0.01% BSA pH 7.4

[00785] Wash buffer: 1x PBS + 3mM $MgCl_2$ pH 7.4

1. Remove microvesicle samples from -80°C freezer and thaw.
2. Place beads on magnet (200,000 per sample experiment), remove bead storage buffer.
3. Wash 1x 200µL for 1 minute each with 1x PBS, 3mM $MgCl_2$ buffer. Pool beads to make 200,000 in one tube.
4. Resuspend beads in 70µL of the selection buffer. (10µl of 10x PBS, 1% BSA + 3µL 100mM $MgCl_2$ +57µL H_2O per sample).
5. Add 30µL radioactively labeled RNA aptamer library to their respective sample.
6. Incubate shaking at 1000 rpm at 37°C for 30 min.
7. Place samples on a magnet.
8. Remove and save supernatant.
9. Wash beads with 200µL wash buffer 1x PBS 3mM $MgCl_2$ pH 7.4, incubating off magnet for 3 min.
10. Place samples on the magnet, remove and save wash solution.
11. Repeat steps 9, 10.
12. Add 100µL water to the sample, pipette mix.
13. Heat at 80°C for 5 minutes.
14. Place samples on a magnet, remove supernatant, and save.
15. Resuspend beads in 100µL water.
16. Measure radioactivity of every fraction using scintillation counter.
17. Analyze amount of background binding present.

[00786] Negative Selection

[00787] As desired, a negative selection step is added prior to incubating the aptamer library with the beads conjugated to the target microvesicles (i.e., procedure "**Binding 2'F SUL1 RNA pool to microvesicle coated magnetic beads**" above). The negative selection can be performed using beads conjugated to the supernatant or the input samples (e.g., plasma) after microvesicles are filtered or sedimented from the sample (referred to as "no microvesicle coated beads," "microvesicle depleted samples," or similar). The steps are:

[00788] 1) Start with aptamer library product from the desired round after transcription as described above. Wash the beads before start: remove storage buffer, wash beads with 200µL wash buffer, then replace buffer as stated below:

[00789] 2) Negative selection step: Add and pipet mix 20 µl of transcription product (15 mM $MgCl_2$) with freshly washed 'no microvesicle' coated beads with 10µL 10x PBS with 1% BSA, 70µL H_2O. No additional $MgCl_2$ is needed because the $MgCl_2$ in the diluted transcription product (TK) provides a final concentration of 3mM $MgCl_2$.

[00790] 3) Incubate for 30 min at 37 °C, shake at 1000rpm.

[00791] 4) Remove supernatant and add it to the positive selection beads (directly), which are washed microvesicle coated beads.

[00792] Continue with positive selection incubation. See **Binding 2'F SUL1 RNA pool to microvesicle coated magnetic beads** above, starting at step 2. Additional steps through transcription are as detailed above.

Example 12: Additional Aptamers to Breast Cancer (BrCa) derived microvesicles

[00793] In this Example, an aptamer library is screened to identify aptamers that distinguish between microvesicles circulating in the blood of breast cancer patients and microvesicles circulating in the blood of healthy, control individuals (i.e., without breast cancer). The procedure used the same samples and aptamer library as in **Example 10** above. The procedure in this Example differs in that negative selection was performed before each positive selection starting after the third round of positive selection.

[00794] Negative selection serves to remove aptamers that bind soluble/abundant/non-informative and common proteins for cancer and non-cancer proteins. Negative selection include performing negative selection on the aptamer candidates selected against BrCa+ microvesicles as follows: (i) using microbeads conjugated to the supernatant from the BrCa+ plasma ultracentrifugation step (which should not contain microvesicles); (ii) using microbeads conjugated to the supernatant from the BrCa- plasma ultracentrifugation step (which should not contain microvesicles); (iii) using microbeads conjugated to BrCa- microvesicles. Negative selection can also be performed on the aptamer candidates selected against BrCa- microvesicles as follows: (i) using microbeads conjugated to the supernatant from the BrCa+ plasma ultracentrifugation step (which should not contain microvesicles); (ii) using microbeads conjugated to the supernatant from the BrCa- plasma ultracentrifugation step (which should not contain microvesicles); (iii) using microbeads conjugated to BrCa+ microvesicles. Negative selection rounds are performed between rounds of positive selection as described herein.

[00795] Microvesicles were isolated from plasma of a pool of 60 breast cancer patients (BrCa+). Microvesicles were also isolated from pool of 60 non-cancer samples (BrCa-). Microvesicles were isolated from the plasma using ultracentrifugation (120,000 x g). Microvesicles were in the pellet from the ultracentrifugation. The supernatant from the ultracentrifugation was saved to use as a control. The microvesicles from both sample types were conjugated to MagPlex beads (Luminex Corp, Austin TX).

[00796] The aptamer library used consisted of a 2'F SUL1 RNA aptamer library. The sequence is 5'-GGGAGGACGAUGCGG-N40-CAGACGACUCGCUGAGGAUCCGAGA-3' (SEQ ID NO. 8). The aptamer library consists of three sections: Forward primer – 15 nucleotides, variable region – 40 nucleotides; reverse primer – 25 nucleotides. All pyrimidines (C and U) were 2'Fluoro modified.

[00797] The aptamer library was incubated with either the cancer or control microvesicle-conjugated beads. Nine rounds of positive selection for aptamers that bind the microvesicles were performed in parallel for both types of samples. Negative selection against beads conjugated to the input plasma supernatant after ultracentrifugation before positive selection in rounds 4-9. See **Example 11** above for detailed protocol of the positive selection and negative selection steps.

[00798] The aptamers that were retained from the above positive selection were sequenced using Next Generation sequencing technology consisting of Ion Torrent NGS (Life Technologies, Inc., Carlsbad, CA). The MiSeq system may be used also (Illumina, Inc., San Diego, CA). The sequences are compared to identify aptamers that are found in the cancer samples and not the control samples, and vice versa. Such aptamers provide candidates that can be used to distinguish between BrCa and non-BrCa samples.

[00799] The sequencing data was analyzed according to the following procedure:

[00800] Step 1: Sequences were ranked according to frequencies in entire aptamer pool recovered in round 9 after negative selection against beads conjugated to microvesicle-depleted cancer plasma followed by positive selection against beads conjugated to cancer microvesicles.

[00801] Step 2: Fold changes were calculated between noted sample in Step 1 and: (i) same sample after additional negative selection against microvesicle depleted cancer plasma; (ii) same sample after additional negative selection against non-cancer microvesicles; (iii) same sample after additional negative selection against microvesicles depleted non-cancer plasma.

[00802] Step 3: Sequences were ranked based on fold changes calculated in Step 2 to identify sequences which are abundant or deficient in aptamer pool selected for breast cancer derived microvesicles.

[00803] Step 4: Possible mutant sequences (e.g., due to PCR or other errors) were removed based on results of consolidation analysis.

[00804] Step 5: Sequences were identified with fold changes greater than 3 and minimum frequency 50 in all three variants (i, ii and iii in step 2).

[00805] The same selection schemes as in steps 1-5 were performed for aptamers selected against beads conjugated to non-cancer microvesicles.

[00806] A number of representative sequences obtained from these procedures are shown in **Table 11**. The sequences in the table were identified in the aptamer pools from selection against BrCa microvesicles but were not in the aptamer pools selected against non-cancer samples. In **Table 11**, the sequences are shown 5' to 3' from left to right, wherein each complete sequence consists of a 5' leader sequence 5'-GGGAGGACGAUGCGG (SEQ ID NO. 9) followed by the indicated Variable Sequence followed by the 3' tail sequence 5'-CAGACGACUCGCUGAGGAUCCGAGA (SEQ ID NO. 10). Each sequence is derived from a library having a leader and tail (see description above) with a variable sequence between. It is understood

that the nucleotide sequences that are disclosed in **Table 11** can also be modified to the extent that resulting modifications result in an aptamer having about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, and 99 percent homology to the disclosed sequence and retain the functionality of binding to microvesicle antigens or functional fragments thereof.

Table 11: BrCa microvesicle aptamer candidate sequences

ID (Figure)	SEQ ID NO.	Variable Sequence
BCE8 **(FIG. 10Ai)**	25	UACCGCCUCAUCAUCGGACACGACGUGUAUCAGUUGGCUG
BCE9 **(FIG. 10Aii)**	26	GUUCUCGCCUCUGUCCUCAUGGUUCGAACCGGUAUGCAUG
BCE10 **(FIG. 10Aiii)**	27	GCGGUUUCUUCUCCUGACUACAUGAGAUUAAUAAACGCGC
BCE11 **(FIG. 10Aiv)**	28	CCGCCUCGAACACUGACGUCGUGGAACCUUCGAUUGCUAG
BCE12 **(FIG. 10Av)**	29	AAUCACAGUAAUUCUGCCCCUCUGAUGAAACCGGUUACUU
BCE13 **(FIG. 10Avi)**	30	CUUAGUGAUUUCGCCGCCCCUCUGUUUAGUGGCCAUUGGA
BCE14 **(FIG. 10Avii)**	31	ACACUAUUCCGGUAAGUCAUCGUUUAACCGUUUGUUGCAA
BCE15 **(FIG. 10Aviii)**	32	UGCGCAACGCCUUGAUUCACUCCUACAGUGUGUCUAUAGA
BCE16 **(FIG. 10Aix)**	33	AAUGUUAAGCUUACAUACGCCUGGGUCACUCUUUGUUCUG
BCE17 **(FIG. 10Bi)**	34	GUAAAUAUUCACGUUGAAUCGCCUUGCUCCUCUUAGUCUG
BCE18 **(FIG. 10Bii)**	35	CCGCCUCGGAUCGUUCCCAAUGGUGGUACCCCUAUUAAUG
BCE19 **(FIG. 10Biii)**	36	UGUAGAUCGUUCUUAUCCGCCUCGGUCUUCCCCAGGUUAA
BCE20 **(FIG. 10Biv)**	37	AUCGUCGGGCCCCUUUUAUGAAACUUACAUGAAAGCGCAC
BCE21 **(FIG. 10Bv)**	38	UAAGAGUGCACAGUACUGCCUCGAUCCUCCAUGGCUUAAG
BCE22 **(FIG. 10Bvi)**	39	GAAUUAGUACUGACGGCCGCCUUGAUCCUCCGUUAGUCUG
BCE23 **(FIG. 10Bvii)**	40	GCCCGCCUCCGAAGCCCUCCUAAGUGCACUUUAAACCGCG
BCE24 **(FIG. 10Bviii)**	41	CCGCCUGGGAUCACUCUCUACGCGUAUAAAUGCUCUGUCA
BCE25 **(FIG. 10Bix)**	42	AGUCUGACCCUGUUAUGGACUACCAUAUCAGAAAGGUACU
BCE26 **(FIG. 10Ci)**	43	GGUGAUCCUCCCCCCCGCCUCGAAGAUUUGUGCACAUAUC
BCE27 **(FIG. 10Cii)**	44	GCUACCAUCGUCUAGUGAGUCACCCUUAGUUCAUCAAGGC

[00807] Each sequence in **Table 11** is synthesized in two variants for further investigation: 5' biotinylated and 3' biotinylated. This provides aptamer variants that can be captured at the 5' end or the 3' end as desired. The aptamers are further synthesized with each pyrimidine (C and U) 2'Fluoro modified. The

aptamers may also be synthesized as the DNA sequence corresponding to each RNA sequence in **Table 11**. The aptamer libraries can also be filtered based on predicted secondary sequence, free energy, and other parameters as described herein.

[00808] **FIGs. 10A-C** show binding of the aptamers in **Table 11** against microbeads conjugated to various input samples. The aptamer is indicated above each plot and the plot for each aptamer is indicated in the ID column in **Table 11**. The input sample is indicated on the X axis from left to right as follows: 1) Cancer Exosome: aptamer binding to microbeads conjugated to microvesicles isolated from plasma samples from breast cancer patients; 2) Cancer Non-exosome: aptamer binding to microbeads conjugated to plasma samples from breast cancer patients after removal of microvesicles by ultracentrifugation; 3) Non-Cancer Exosome: aptamer binding to microbeads conjugated to microvesicles isolated from plasma samples from normal (i.e., non-breast cancer) patients; 4) Non-Cancer Non-Exosome: aptamer binding to microbeads conjugated to plasma samples from breast cancer patients after removal of microvesicles by ultracentrifugation. As shown in **FIGs. 10A-C**, the aptamers were each able to distinguish between the cancer microvesicle samples versus the supernatant control samples and the non-cancer microvesicles. Further, all sequences in **Table 11** were observed as binding more abundantly to cancer derived microvesicles as compared to non-cancer derived microvesicles with the exception of BCE10 and BCE14, which were observed as binding more abundantly to non-cancer derived microvesicles as compared to cancer derived microvesicles.

[00809] Based on the comparisons performed in this Example, aptamers that bind different starting input are obtained, including: 1) aptamers that preferentially bind cancer-derived microvesicles over non-cancer derived microvesicles; 2) aptamers that preferentially bind non-cancer-derived microvesicles over cancer derived microvesicles; 3) aptamers that bind both non-cancer-derived microvesicles and cancer derived microvesicles (e.g., "universal" binders); and 4) aptamers that bind plasma components that have been depleted of microvesicles.

[00810] The aptamer libraries in this Example are further subjected to four rounds of additional negative and positive selection. The positive selection is performed as described in this Example. The negative selection rounds are performed using the beads conjugated to non-cancer microvesicles as negative selection for aptamers obtained by positive selection against beads conjugated to cancer microvesicles. Similarly, the negative selection rounds are performed using the beads conjugated to cancer microvesicles as negative selection for aptamers obtained by positive selection against beads conjugated to non-cancer microvesicles.

Example 13: Disease Diagnosis

[00811] This example illustrates the use of oligonucleotide probes of the present invention to diagnose a proliferative disease.

[00812] A suitable quantity of an oligonucleotide or pool of oligonucleotides that bind a BrCa-derived population of microvesicles, such as identified in **Example 10** or **Example 12** or various Examples below, is synthesized via chemical means known in the art. The oligonucleotides are conjugated to a

diagnostic agent suitable for detection, such as a fluorescent moiety, using a conjugation method known in the art.

[00813] The composition is applied to microvesicles isolated from blood samples taken from a test cohort of patients suffering from a proliferative disease associated with the overexpression of microvesicles, e.g. breast cancer. The composition is likewise applied to microvesicles isolated from blood samples taken from a negative control cohort, not suffering from a proliferative disease.

[00814] The use of appropriate detection techniques (e.g., microbead assay or flow cytometry) on the test cohort samples indicates the presence of disease, while the same techniques applied to the control cohort samples indicate the absence of disease.

[00815] The results show that the oligonucleotides of the present invention are useful in diagnosing proliferative diseases.

Example 14: Theranostics

[00816] This example illustrates the use of oligonucleotide probes of the present invention to provide a theranosis for a drug for treating a proliferative disease.

[00817] A suitable quantity of an oligonucleotide or pool of oligonucleotides that bind a BrCa-derived population of microvesicles, such as identified in **Example 10** or **Example 12** or various Examples below, is synthesized via chemical means known in the art. The probes are conjugated to an agent suitable for detection, such as a fluorescent moiety, using a method known in the art such as conjugation. The oligonucleotide probe or panel of oligonucleotide probes are within a suitable composition, such as a buffered solution.

[00818] Treatment selection. The composition is applied to microvesicles isolated from blood samples taken from a test cohort of patients suffering from a proliferative disease, e.g. breast cancer, that responded to a certain treatment, e.g., trautuzamab. The composition is likewise applied to microvesicles isolated from blood samples taken from a control cohort consisting of patients suffering from the same proliferative disease that did not respond to the treatment. The use of appropriate detection techniques (e.g., microbead assay or flow cytometry) on the test cohort samples indicates that probes which bind the samples are useful for identifying patients that will respond to the treatment, while the same techniques applied to the control cohort samples identifies probes useful for identifying patients that will not respond to the treatment.

[00819] Treatment monitoring. In another setting, the composition is applied to microvesicles isolated from blood samples taken from a test cohort of patients suffering from a proliferative disease, e.g. breast cancer, prior to or during a course of treatment, such as surgery, radiotherapy and/or chemotherapy. The composition is then applied to microvesicles isolated from blood samples taken from the patients over a time course. The use of appropriate detection techniques (e.g., microbead assay or flow cytometry) on the test cohort samples indicates whether the detected population of disease-related microvesicles increases, decreases, or remains steady in concentration over time during the course of treatment. An increase in the population of disease-related microvesicles post-treatment may indicate that the treatment is ineffective

whereas a dcrease in the population of disease-related microvesicles post-treatment may indicate that the treatment has a beneficial effect.

[00820] The results show that the oligonucleotide probes of the present invention are useful in theranosing proliferative diseases.

Example 15: Therapeutic Oligonucleotide Probes

[00821] This example illustrates the use of oligonucleotide probes of the present invention to treat a proliferative disease.

[00822] A suitable quantity of an oligonucleotide or pool of oligonucleotides that bind a BrCa-derived population of microvesicles, such as identified in **Example 10** or **Example 12** or various Examples below, is synthesized via chemical means known in the art. The oligonucleotides are conjugated to a chemotherapeutic agent, such as Doxil, using a conjugation method known in the art. The conjugate is formulated in an aqueous composition.

[00823] The composition is administered intravenously, in one or more doses, to a test cohort of mice suffering from a proliferative disease associated with the overexpression of the microvesicles, e.g. a breast cancer model. A control cohort, not suffering from a proliferative disease is administered the identical composition intravenously, according to a corresponding dosage regimen.

[00824] Pathological analysis of tumor samples and/or mouse survival indicates that mortality and/or morbidity are improved in the test cohort over the control cohort.

[00825] The results show that the oligonucleotides of the present invention are useful in treating proliferative diseases.

[00826] Useful oligonucleotides can be used to treat proliferative diseases in other organisms, e.g., a human.

Example 16: Oligonucleotide – Sequencing Detection Method

[00827] This example illustrates the use of an oligonucleotide pool to detect microvesicles that are indicative of a phenotype of interest. The method makes use of a pool of oligonucleotides that have been enriched against a target of interest that is indicative of a phenotype of interest. The method in this Example allows efficient use of a library of oligonucleotides to preferentially recognize a target entity.

[00828] For purposes of illustration, the method is described in the Example with a microvesicle target from a bodily fluid sample. One of skill will appreciate that the method can be extended to other types of target entity (e.g., cells, proteins, various other biological complexes), sample (e.g., tissue, cell culture, biopsy, other bodily fluids) and other phenotypes (other cancers, other diseases, etc) by enriching an aptamer library against the desired input samples.

[00829] General workflow:

[00830] 1) Obtain sample (plasma, serum, urine or any other biological sample) of patients with unknown medical etymology and pre-treating them accordingly to ensure availability of the target of interest (see below). Where the target of interest is a microvesicle population, the microvesicles can be isolated and optionally tethered to a solid support such as a microbead.

[00831] 2) Expose pre-treated sample to an oligonucleotide pool carrying certain specificity against target of interest. As described herein, an oligonucleotide pool carrying certain specificity against the target of interest can be enriched using various selection schemes, e.g., using non-cancer microvesicles for negative selection and cancer microvesicles for positive selection as described above. DNA or RNA oligonucleotides can be used as desired.

[00832] 3) Contact oligonucleotide library with the sample.

[00833] 4) Elute any oligonucleotides bound to the target.

[00834] 5) Sequence the eluted oligonucleotides. Next generation sequencing methods can be used.

[00835] 6) Analyze oligonucleotide profile from the sequencing. A profile of oligonucleotides known to bind the target of interest indicates the presence of the target within the input sample. The profile can be used to characterize the sample, e.g., as cancer or non-cancer.

[00836] Protocol variations:

[00837] Various configurations of the assay can be performed. Four exemplary protocols are presented for the purposes of the oligonucleotide-sequencing assay. Samples can be any appropriate biological sample. The protocols can be modified as desired. For example, the microvesicles can be isolated using alternate techniques instead or or in addition to ultracentrifugation. Such techniques can be disclosed herein, e.g., polymer precipitation (e.g., PEG), column chromatography, and/or affinity isolation.

[00838] Protocol 1:

[00839] Ultracentrifugation of 1-5 ml bodily fluid samples (e.g., plasma/serum/urine) (120K x g, no sucrose) with two washes of the precipitate to isolate microvesicles.

[00840] Measure total protein concentration of recovered sample containing the isolated microvesicles.

[00841] Conjugate the isolated microvesicles to magnetic beads (for example MagPlex beads (Luminex Corp. Austin TX)).

[00842] Incubate conjugated microvesicles with oligonucleotide pool of interest.

[00843] Wash unbound oligonucleotides by retaining beads using magnet.

[00844] Elute oligonucleotides bound to the microvesicles.

[00845] Amplify and purify the eluted oligonucleotides.

[00846] Oligonucleotide sequencing (for example, Next generation methods; Ion Torrent: fusion PCR, emulsion PCR, sequencing).

[00847] Assess oligonucleotide profile.

[00848] Protocol 2:

[00849] This alternate protocol does not include a microvesicle isolation step, microvesicles conjugation to the beads, or separate partitioning step. This may present non-specific binding of the oligonucleotides against the input sample.

[00850] Remove cells/debris from bodily fluid sample and dilute sample with PBS containing $MgCl_2$ (2mM).

[00851] Pre-mix sample prepared above with oligonucleotide library.

[00852] Ultracentrifugation of oligonucleotide/sample mixture (120K x g, no sucrose). Wash precipitated microvesicles.

[00853] Recover precipitate and elute oligonucleotides bound to microvesicles.

[00854] Amplify and purify the eluted oligonucleotides.

[00855] Oligonucleotide sequencing (for example, Next generation methods; Ion Torrent: fusion PCR, emulsion PCR, sequencing).

[00856] Assess oligonucleotide profile.

[00857] Protocol 3:

[00858] This protocol uses filtration instead of ultracentrifugation and should require less time and sample volume.

[00859] Remove cells/debris from bodily fluid sample and dilute it with PBS containing $MgCl_2$ (2mM).

[00860] Pre-mix sample prepared above with oligonucleotide library.

[00861] Load sample into filter (i.e., 150K or 300K MWCO filter or any other that can eliminate unbound or unwanted oligonucleotides). Centrifuge sample to concentrate. Concentrated sample should contain microvesicles.

[00862] Wash concentrate. Variant 1: Dilute concentrate with buffer specified above to the original volume and repeat centrifugation. Variant 2: Dilute concentrate with buffer specified above to the original volume and transfer concentrate to new filter unit and centrifuge. Repeat twice.

[00863] Recover concentrate and elute oligonucleotides bound to microvesicles.

[00864] Amplify and purify the eluted oligonucleotides.

[00865] Oligonucleotide sequencing (for example, Next generation methods; Ion Torrent: fusion PCR, emulsion PCR, sequencing).

[00866] Assess oligonucleotide profile.

[00867] Protocol 4:

[00868] Ultracentrifugation of 1-5 ml bodily fluid sample (120K x g, no sucrose) with 2 washes of the precipitate to isolate microvesicles.

[00869] Pre-mix microvesicles with oligonucleotide pool.

[00870] Load sample into 300K MWCO filter unite and centrifuge (2000xg). Concentration rate is ~3x.

[00871] Wash concentrate. Variant 1: Dilute concentrate with buffer specified above to the original volume and centrifuge. Repeat twice. Variant 2: Dilute concentrate with buffer specified above to the original volume and transfer concentrate to new filter unit and centrifuge. Repeat twice

[00872] Recover concentrate and elute oligonucleotides bound to microvesicles.

[00873] Amplify and purify the eluted oligonucleotides.

[00874] Oligonucleotide sequencing (for example, Next generation methods; Ion Torrent: fusion PCR, emulsion PCR, sequencing).

[00875] Assess oligonucleotide profile.

[00876] In alterations of the above protocols, polymer precipitation is used to isolate microvesicles from the patient samples. For example, the oligonucleotides are added to the sample and then PEG4000 or PEG8000 at 4% or 8% concentration is used to precipitate and thereby isolate microvesicles. Elution, recovery and sequence analysis continues as above.

Example 17: Plasma/Serum probing with an Oligonucleotide Probe Library

[00877] The following protocol is used to probe a plasma or serum sample using an oligonucleotide probe library.

[00878] Input oligonucleotide library:

[00879] Use 2 ng input of oligonucleotide library per sample.

[00880] Input oligonucleotide library is a mixture of two libraries, cancer and non-cancer enriched, concentration is 16.3 ng/ul.

[00881] Dilute to 0.2ng/ul working stock using Aptamer Buffer (3mM $MgCl_2$ in 1X PBS)

[00882] Add 10ul from working stock (equal to 2 ng library) to each optiseal tube

[00883] Materials:

[00884] PBS, Hyclone SH30256.01, LN: AYG165629, bottle# 8237, exp. 7/2015

[00885] Round Bottom Centrifuge Tubes, Beckman 326820, LN:P91207

[00886] OptiSeal Centrifuge tubes and plugs, polyallomer Konical, Beckman 361621, lot# Z10804SCA

[00887] Ultracentrifuge rotor: 50.4 TI

[00888] Ultracentrifuge rotor: 50.4 TI, Beckman Caris ID# 0478

[00889] Protocol:

[00890] 1 Pre-chill tabletop centrifuge, ultracentrifuge, buckets, and rotor at 4°C.

[00891] 2 Thaw plasma or serum samples

[00892] 3 Dilute 1ml of samples with 1:2 with Aptamer Buffer (3mM $MgCl_2$ in 1X PBS)

[00893] 4 Spin at 2000xg, 30 min, 4°C to remove debris (tabletop centrifuge)

[00894] 5 Transfer supernatants for all samples to a round bottom conical

[00895] 6 Spin at 12,000xg, 45 min, 4°C in ultracentrifuge to remove additional debris.

[00896] 7 Transfer supernatant about 1.8ml for all samples into new OptiSeal bell top tubes (uniquely marked).

[00897] 8 Add 2ng (in 10 ul) of DNA Probing library to each optiseal tube

[00898] 9 QS to 4.5 ml with Aptamer Buffer

[00899] 10 Fix caps onto the OptiSeal bell top tubes

[00900] 11 Apply Parafilm around caps to prevent leakage

[00901] 12 Incubate plasma and oligonucleotide probe library for 1 hour at room temperature with rotation

[00902] 13 Remove parafilm (but not caps)

[00903] 14 Place correct spacer on top of each plugged tube

[00904] 15 Mark pellet area on the tubes, insure this marking is facing outwards from center.

[00905] 16 Spin tubes at 120,000 x g, 2hr, 4°C (inner row, 33,400 rpm) to pellet microvesicles.

[00906] 17 Check marking is still pointed away from center.

[00907] 18 Completely remove supernatant from pellet, by collecting liquid from opposite side of pellet marker and using a 10 ml syringe barrel and 21G2 needle

[00908] 19 Discard supernatant in appropriate biohazard waste container

[00909] 20 Add 1 ml of 3 mM MgCl2 diluted with 1X PBS

[00910] 21 Gentle vortex, 1600rpm for 5 sec and incubate 5 min at RT.

[00911] 22 QS to ~4.5 mL with 3 mM Mg Cl2 diluted with 1X PBS

[00912] 23 Fix caps onto the OptiSeal bell top tubes.

[00913] 24 Place correct spacer on top of each plugged tube.

[00914] 25 Mark pellet area on the tubes, insure this marking is facing outwards from center.

[00915] 26 Spin tubes at 120,000 x g, 70 min, 4°C (inner row 33,400 rpm) to pellet microvesicles

[00916] 27 Check marking in still pointed away from center.

[00917] 28 Completely remove supernatant from pellet, by collecting liquid from opposite side of pellet marker and using a 10 ml syringe barrel and 21G2 needle

[00918] 29 Discard supernatant in appropriate biohazard waste container

[00919] 30 Add 1 ml of 3 mM MgCl2 diluted with 1X PBS

[00920] 31 Gentle vortex, 1600rpm for 5 sec and incubate 5 min at RT.

[00921] 32 QS to ~4.5 mL with 3 mM Mg Cl2 diluted with 1X PBS

[00922] 33 Fix caps onto the OptiSeal bell top tubes.

[00923] 34 Place correct spacer on top of each plugged tube.

[00924] 35 Mark pellet area on the tubes, insure this marking is facing outwards from center.

[00925] 36 Spin tubes at 120,000 x g, 70 min, 4°C (inner row 33,400 rpm) to pellet microvesicles

[00926] 37 Check marking is still pointed away from center.

[00927] 38 Save an aliquot of the supernatant (100ul into a 1.5ml tube)

[00928] 39 Completely remove supernatant from pellet, by collecting liquid from opposite side of pellet marker and using a 10 ml syringe barrel and 21G2 needle

[00929] 40 Add 50 ul of Rnase-free water to the side of the pellet

[00930] 41 Leave for 15min incubation on bench top

[00931] 42 Cut top off tubes using clean scissors.

[00932] 43 Resuspend pellet, pipette up and down on the pellet side

[00933] 44 Measure the volume, make a note on the volume in order to normalize all samples

[00934] 45 Transfer the measured resuspended eluted microvesicles with bound oligonucleotides to a Rnase free 1.5ml Eppendorf tube

[00935] 46 Normalize all samples to 100ul to keep it even across samples and between experiments.

[00936] Next Generation Sequencing Sample Preparation:

[00937] I) Use 50 ul of sample from above, resuspended in 100 ul H2O and containing microvesicle/oligo complexes, as template in Transposon PCR, 14 cycles.

[00938] II) AMPure transposon PCR product, use entire recovery for indexing PCR, 10 cycles.

[00939] III) Check indexing PCR product on gel, proceed with AMPure if band is visible. Add 3 cylces if band is invisible, check on gel. After purification quantify product with QuBit and proceed with denaturing and dilting for loading on HiSeq flow cell (Illumina Inc., San Diego, CA).

[00940] IV) 5 samples will be multiplexed per one flow cell. 10 samples per HiSeq.

Example 18: Enrichment of Oligonucleotides to Breast Cancer (BrCa) and non-BrCa derived microvesicles

[00941] In this Example, a naive oligonucleotide probe library is screened to enrich oligonucleotides that identify microvesicles circulating in the blood of breast cancer patients and microvesicles circulating in the blood of healthy, control individuals (i.e., without breast cancer). The procedure in this Example used the 9-round selection scheme shown in **FIG. 11A**.

[00942] In **FIGs. 11A-11B**, the following abreviations are used: CE- Cancer exosome samples; NCE- Non-cancer exosome samples; CNE- Cancer no exosome samples; NCNE- Non-cancer no exosome samples; MFI- Median Fluorescence Intensity; GC- GC content group analysis; Heat- heat elution; NaOH- sodium hydroxide elution. The terms exosome and microvesicle are used interchangeably.

[00943] As indicated in **FIG. 11A**, 0.1 mg/mL salmon sperm DNA was added as a competitor in selection rounds 7-9. **FIG. 11B** shows the enrichment of aptamers that bind to microvesicles (CE and NCE). Binding was determined by fluorescently labeling aptamer candidates and detecting binding to bead-capture microvesicles using a microbead assay format. At the end of round 9, only aptamers having GC content between ~30% and 70% were selected to generate panels to diffentiate cancer and non-cancer samples. In addition, several groups of aptamers were selected using the following criteria: 1) 12 most commonly observed; 2) 12 most commonly observed having > 2-fold change in binding observed between cancer and normal samples; 3) 30 most commonly observed having > 2-fold change in binding observed between cancer and normal samples, >10 read counts in the library determined by high throughput sequencing, heat elution after selection round 9; and 4) 30 most commonly observed having > 2-fold change in binding observed between cancer and normal samples, >10 read counts in the library determined by high throughput sequencing, NaOH elution after selection round 9.

[00944] Retained sequences are shown in **Table 12** with selection criteria 1-4 indicated. In **Table 12**, the sequences are shown 5' to 3' from left to right, wherein each complete sequence consists of a 5' leader sequence 5'- GCTGTGTGACTCCTGCAA (SEQ ID NO. 45) followed by the indicated Variable Sequence followed by the 3' tail sequence 5'-GCAGCTGTATCTTGTCTCC (SEQ ID NO. 46).

Table 12: BrCa microvesicle oligonucleotide sequences

SEQ ID NO.	Selection Criteria	Variable Sequence
47	3	GTCCGCTTGGGGGTGGGTATCGAAAATTCGCGTTCTGGGGCGG
48	3	CTTCGCTCGTTTCAATGTTCAGTTCTGGTCTTTAGGTCTTTAG
49	3	TAGGAGGATAATATGTATCCGGTGCATCGGCCTGACTTTCCCC
50	3	TAGGAGGGCTGAAAACTTGCTTCTTCCCGGTGCAGTTATAACA
51	3	TAGGAGGATAATATGTATCCGATGCATGGGCCCGACTTTCCCC

52	3	TGCGTTTTCCTTGGGGTTCAGGGTAGGATGGGGGTGGAGGTGG
53	3	GGGGGGCGGGTGGGCTGGTAAAAGGTGATGCGGGGGTATTGTT
54	3	TAGGAGGATAATATGTATCCGGTGCATCGGCCCAACTTTCCCC
55	3	CCAATCTCGGAAGGTTTAAATAAGGTGGTCTTTAGGTCTTTAG
56	3	TACGTTAAGTGGGTCGGGAGGGGGGAATAGGGGGTTTGGTTGG
57	3	GTGAGGTCGGTGGGGCGGGGGATGGTGGGGGGTCGTTTACATT
58	3	CGATATTGGGGGGGGGTTGGCGGGCTATTTTCCGGGTGGGTG
59	3	TCGGTTCCTGTTAATTCGCTGGTGGTTGGTGGGGTGGCGGATG
60	3	GCGGGGAGGTGGGGGTGGGTGGAGGGTTGGTTACTCTCTACT
61	3	CGGGAATGGGAGGGTGGGGGTGGTGGCCGGGTCGTGTTATACC
62	3	TAGGAGGATAATATGTATCCGGTGCATCGGTCCGACTTTCCCC
63	3	TATAGTTGGGCGTAGGCGGGGGGGGGTGGTTGGGAGGTCCAAG
64	3	TAGGAGGATAATATGTATCCGGTGCATCGGCCCGACCTTCCCC
65	3	TCGATCGTCCTCAGGATCTCGTGGTTCAAATCATAAAGATTAT
66	3	TGCTGTCTGGGCGGGGGCGGTCTTGTGGTTTCTTTGGGGGGGG
67	3	GCGAGACAGGAGGGTGGTCTTATACGTGGGGGGGGGTGGTTGG
68	3	TAGGAGGGCTGAAGACTTGCTTCTTCCCGGTGCAGTTATAACA
69	3	CGTGTGGGGGGTGGGTTGGGCTCGGGTTGTTATCAGTTCCATG
70	3	CGATATTGGGGGGGGGGTTGGCGGGCTATTTTCCGGGTGGGTG
71	3	TAGGAGGATAATATGTATCCGGTGCATCGGCCCGACTTTCCTC
72	3	TAGGAGGATAATATGTATCCGGTGCATCGGCTCGACTTTCCCC
73	3	CTTCGCTCGTTTCAATGTTTAGTTCTGGTCTTTAGGTCTTTAG
74	3	GGACATGGGTTGGGTCGGGAGGGGGTGGTCGGTTGGGCAGTAA
75	3	TAGGAGGATAATATGTATACGATGCATCGGCCCGACTTTCCCC
76	3	TAGGAGGATAATATGTATCAGATGCATCGGCCCGACTTTCCCC
77	4	TGGATGTATGGGGTCTCGGGGTGGGAGGGTTCAACTTATCTGG
78	4	GTCCGCTTGGGGGTGGGTATCGAAAATTCGCGTTCTGGGGCGG
79	4	TCTTGTACAAATAGGAGGGAAGGGGGTTTTGGGAGGTGGGTGG
80	4	CTTCGCTCGTTTCAATGTTCAGTTCTGGTCTTTAGGTCTTTAG
81	4	CTGATGTTAGTAGGTCGGGGTCCGGTTGGGGGTTGGGTTGAGG
82	4	TGGCTCGTGGACGTGGTGGTGGCGGGTCGTGGGGGTGGGTAGG
83	4	TAGGAGGATAATATGTATCCGATGCATCGGCACGACTTTCCCC
84	4	CGGGGAGGGGGGGTCGGGGTATTTATTGTGTATGTTTTTTGTG
85	4	TGACTGTATCTTGGGGCGGGTTCTGGGGGGGGTGTATTGTTCA
86	4	TAGGAGGATAATATGTATCCGATGCATCGGCCCGACTTTCACC
87	4	TCATGTCTAGGGGGGGGAAGTCGGTTTGGGTGGGTACTCTGTG
88	4	CTTGTTTCGCTTTGGGTTTGGGCGGGTGGGTCAATTCCTGTTG
89	4	GGTGGGGGCCCTCGGTACTGTGGCGGGGTGGGTGGGTTTAGTG
90	4	TGCGTTTTCCTTGGGGTTCAGGGTAGGATGGGGGTGGAGGTGG
91	4	TGGTGGGGTGGGTCTGTGGGGTGGTTTTGTTCTTATCGGGGTT
92	4	TGGGTGGGTTACGGGTGGGTGTTGTTATCGCGCTGATCTGGTT
93	4	CGTGACGGTTTTAATCAGGGGGGGGACTCTAACATTTGGGTGG
94	4	TGCTCATATCGTGGGGGGATGGGGTTGCTAGTGGATGGGGTGG
95	4	GGGGGGCGGGTGGGCTGGTAAAAGGTGATGCGGGGGTATTGTT
96	4	TGATGTGGGGGGTGGGGTATAATACTTATGTTTGGGGTTTGGG
97	4	TTGGGTGGGTGGGAATGGGTATTTTTTCTTCGGGCGATGTTTG
98	4	TAGTTGGATACTGGATTTGGGAGGGATGGGGGGAGGAGGGTGG
99	4	CTGGAAGGTTGGGTGTGGGGCGGGGGGGAGGTCTCTCTATGT
100	4	TACGTTAAGTGGGTCGGGAGGGGGGAATAGGGGGTTTGGTTGG
101	4	TGCCGTGGGCCGGGGAGGGTGGGTTGGCGTCTCTGTTTCGATA
102	4	GGGGGGGCGGCCGGGGTGGACTGTGGCCCGTCTTCGTATTGTT
103	4	TGGTGGTGGCTGGGGGACGGGTATCCTGGAATTAGGGGTGGG
104	4	TCGGTTCATGTTGGGTGGGATCGGGGCGGGGTTTCTTCTCCTG
105	4	TAGGAGGATAATATGTATACGATGCATCGGCCCGACTTTCCCC

106	4	TCGTGGGTGTGTTCTGGGGGGCGTGGACGGGGGTTAATGGGCA
107	1	TAGGAGGATAATATGTATCCGATGCATCGGCCCGACTTTCCCC
108	1	TAAGAGTCTTAGGATGACGTCATTCGTCCGTCACGGTGCGGGA
109	1	GCCTGGCCAGATAGCAGTTACCTTACGGGATCTATATTCACCG
110	1	TGGCTCGTGGACGTGGTGGTGGCGGGTCGTGGGGGTGGGTAGG
111	1	TTCGTGTTCATCTGTTTATTGTTATTCACAATCCGTCTTTTGT
112	1	CGAATCCGTCGACCGCCCTCGTAACCATCCAACCCAGTCTCCT
113	1	CAGGAAAGTTACTTATCTTTTAGACGGTTATGTTCTCATTACT
114	1	ATCTGCTCGATACGTAGATAGTCTGCTCGATACAAAAGGGTGG
115	1	CGATGAGCGGGCTCCGATATTGTGTCGGCGTGACTACCTTGAT
116	1	GGTCGGCCTATTTATTGCGGCACGTTGTTTCTTGATGTCGCCC
117	1	TCGAATTCAGTTCTACATGTGTATTTCTTGCTCCGTTTCAGAA
118	1	TCGGTCCGGCAACAAACCCCACGCGCCTTCGATATGCCTGTCG
119	2	CCAATCTCGGAAGGTTTAAATAAGGTGGTCTTTAGGTCTTTAG
120	2	CGATATTGGGGGGGGGGTTGGCGGGCTATTTTCCGGGTGGGTG
121	2	CGGGAATGGGAGGGTGGGGGTGGTGGCCGGGTCGTGTTATACC
122	2	CGTGGTGGGTTCTTGGGTGGGGGGGGAGGGTTGGCGTAGTATA
123	2	GTGAGGTCGGTGGGGCGGGGGATGGTGGGGGGTCGTTTACATT
124	2	TACGTTAAGTGGGTCGGGAGGGGGGAATAGGGGGTTTGGTTGG
125	2	TAGGAGGATAATATGTATCCGGTGCATCGGCCCAACTTTCCCC
126	2	TAGGAGGGCTGAAAACTTGCTTCTTCCCGGTGCAGTTATAACA
127	2	TCGGTTCCTGTTAATTCGCTGGTGGTTGGTGGGGTGGCGGATG
128	2	TGCTGTCTGGGCGGGGGCGGTCTTGTGGTTTCTTTGGGGGGGG
129	2	AACATGGGGTTCTGATAAACCTGTGCCAAATACATTGTTGAGT
130	2	ATTCCTCCTTTGTAAGGGATGATATCCAACTCCTCTGCGTTCG

[00945] Individual aptamer candidates in **Table 12** are synthesized. The aptamers are labeled with a fluorescent label and used to detect binding to bead-capture microvesicles using a microbead assay format. The microbead results are used to identify aptamers and panels of aptamers that differentiate cancer and non-cancer plasma samples.

Example 19: Aptamer based precipitation of microvesicles in plasma

[00946] Aptamers can specifically recognize target proteins with nanomolar affinity and have some potential advantages compared to antibodies due to their chemical stability, ease of synthesis and overall reproducibility. Microvesicles were isolated from a rat glioma cell line model that expresses or does not express the human EGFR protein in order to optimize aptamer-microvesicle complex formation and isolation in plasma. Microvesicles spiked into human plasma were precipitated with the positive EGFR DNA or RNA aptamer but not with a reverse complement control sequence. There was no observed binding of the positive or negative aptamer to microvesicles that do not express EGFR. Binding of aptamers to microvesicles was confirmed with aptamer based ELISAs, EMSA assays, and flow cytometry. These results demonstrate aptamer-based precipitation of microvesicles from a complex biological sample.

Example 20: Quantitative proteomics of EGFR, EGFRviii, and EGFR negative microvesicles from a rat glioma cell line

[00947] In this Example, we performed quantitative proteomics on microvesicles isolated from a rat glioma cell line F98 (parental line), F98(EGFR) and F98(EGFRviii). Different lots of microvesicles purified from each cell line type were considered biological replicates and each biological replicate was analyzed in four technical replicates by tandem mass tag 6-plex (TMT) peptide labeling and LC-MS/MS. The general microvesicle markers CD81, CD9, CD63, Tsg101, and Alix were identified in all technical replicates. Multiple proteins were found to be up or down-regulated including a 7.5-fold increase in EGFR and a 2.6-fold increase in the EGFR binding partner HMCN2 in the F98[EGFR] compared with the parental line. These results indicate that expression of a single biomarker may affect global proteomic changes in microvesicles.

Example 21: Oligonucleotide probe library

[00948] This Example presents further development of the oligonucleotide probe library to detect biological entities such as described in **Example 18** above. In this Example, steps were taken to reduce the presence of double stranded oligonucleotides (dsDNA) when probing the patient samples. The data were also generated comparing the effects of 8% and 6% PEG used to precipitate microvesicles (and potentially other biological entities) from the patient samples.

[00949] Protocol:

[00950] 1) Pre-chill tabletop centrifuge at 4°C.

[00951] 2) Protease inhibition: dissolve 2 tablets of "cOmplete ULTRA MINI EDTA-free EASYpack" protease inhibitor in 1100 ul of H_2O (20x stock of protease inhibitor).

[00952] 3) Add 50 ul of protease inhibitor to the sample (on top of frozen plasma) and start thawing: 1 ml total ea.

[00953] 4) To remove cells/debris, spin samples at 10,000 x g, 20 min, 4°C. Collect 1 ml supernatant (SN).

[00954] 5) Mix 1 ml supernatant from step 4 with 1ml of 2xPBS 6 mM $MgCl_3$, collect 400 ul into 3 tubes (replicates A, B, C) and use it in step 6.

[00955] 6) Add competitor per **Table 13**: make dilutions in 1xPBS, 3mM $MgCl_2$, mix well, pour into trough, pipet using multichannel.

Table 13: Competitors

units	Type of Competitor	Stock Concentration	Intermediate stock concentration	Number of samples	Volume from stock to make intermediate stock, ul	Buffer to make intermediate stock	Final Volume, ul	Final Concentration
ng/ul	Salmon DNA	-	40	-	-	-	425.5	0.8

ng/ul	tRNA	-	40	-	-	-	425.5	0.8
x	S1	20	0.5	280	65.5	2555.6	425.5	0.01

[00956] 7) Incubate for 10 min, RT, end-over-end rotation

[00957] Pool of 6-3S and 8-3S oligonucleotide probing libraries is ready: 2.76 ng/ul (~185 ng). Save pool stock and dilutions. New pool can be made by mixing 171.2ul (500ng) of library 6-3S (2.92 ng/ul) with 190.8ul (500ng) of library 8-3S (2.62 ng/ul). Aliquot pooled library into 30 ul and store at -80C.

[00958] Add ssDNA oligonucleotide probing library to the final concentration 2.5 pg/ul for binding. Make dilutions in 1xPBS, 3mM $MgCl_2$.

Table 14: Probe library calculations

Original stock, ng/ul	Lib Name	Required working stock (ng/ul)	ul from original stock to make working	ul of buffer to make working	Final volume, ul	Number of samples	Volume per sample from working stock	Final concentration (pg/ul)
2.76	**Pooled library 6-3S/8-3S**	0.1	26.1	694.1	720.2	60	10.9	2.5

[00959] 8) Binding: Incubate for 1h at RT with rotation.

[00960] 9) Prepare polymer solution: 20% PEG8000 in 1x PBS 3mM MgCl2 (dilute 40% PEG8000 with 2xPBS with 6mM MgCl2). Add 20% PEG8000 to sample to the final concentration 6%. Invert few times to mix, incubate for 15 min at 4C

Table 15: PEG calculations

PEG MW	PEG stock, %	Final conc., %	Final volume, ul	Volume 20% PEG to add, ul	Volume of buffer to adjust final volume, ul	Sample volume before adding PEG	Total samples	Total 20% PEG needed, ml
8000	20	6	622.8	186.9	-0.4	436.4	60	11.2

[00961] 10) Spin at 10,000 x g for 5 min, RT.

[00962] 11) Remove SN, add 1ml 1xPBS, 3 mM MgCl2 and wash pellet by gentle invertion with 1ml aptamer buffer.

[00963] 12) Remove buffer, Re-suspend pellets in 100 ul H2O: incubate at RT for 10 min on mixmate 900rpm to re-suspend.

[00964] 13) Make sure each sample is re-suspended by pipeting after step 13. Make notes on hardly re-suspendable samples.

[00965] 14) 50 ul of re-suspended sample to indexing PCR -> next generation sequencing (NGS).

[00966] 15) Keep leftover at 4C

[00967] Technical Validation:

[00968] The current protocol was tested versus a protocol using 8% PEG8000 to precipitate microvesicles. The current protocol further comprises steps to reduce dsDNA in the oligonucleotide probing libraries.

[00969] **FIG. 12A** shows the within sample variance (black) between binding replicates and the between sample variance (grey). Black is on top of grey, thus any observable grey oligo is informative about differences in the biology of two paitent samples. This evaluation of Sources of Variance shows that the technical variances is significantly smaller than the biological variance.

[00970] **FIG. 12B** shows the impact of using a higher proportion of single stranded DNA and PEG 6% isolation (white bars) compared to when there is a higher amount of double stranded DNA and 8% PEG (grey). This data indicates that the protocol in this Example improves biological separation between patients.

[00971] The plots in **FIG. 12C** show the difference between an earlier protocol (PEG 8% with increased dsDNA) and a modified protocol of the Example (PEG 6% no dsDNA). The black is the scatter between replicates (independent binding events) and the grey is the difference between patients. This data shows that the signal to noise increased significantly using the newer protocol.

[00972] Patient testing:

[00973] The protocol above was used to test patient samples having the following characteristics:

Table 16: Patient characteristics

Sample Type	Description
Cancer	Mixed type carcinoma;Malignant;
Cancer	Invasive, predominant intraductal component (8500/3)
Cancer	Fibrocystic Changes;Invasive lobular carcinoma - 8520/3;Lobular carcinoma in situ - 8520/2;Benign;In situ and grade 3 intraepith;Malignant;Fat necrosis, periductal inflammation, malignant cellsFat necrosis;Inflammation;Benign;
Cancer	Invasive, predominant intraductal component (8500/3)
Cancer	Mucinous (colloid) adenocarcinoma (8480/3)
Cancer	Invasive lobular carcinoma - 8520/3;Microcalcifications;Benign;Malignant;
Cancer	Otherfibrocystic changeInvasive, NOS (8500/3)
Cancer	Invasive ductal carcinoma, not otherwise specified (NOS) - 8500/3;Malignant;
Cancer	Invasive ductal carcinoma, not otherwise specified (NOS) - 8500/3;Malignant;
Cancer	Intraductal carcinoma, non-infiltrating, NOS (in situ) (8500/2)
Cancer	Atypical lobular hyperplasia Otherfibrocystic changes, inter and intralobular fibrosis, apocrine metaplasia, columnar cell change, microcalcificationsInvasive, NOS (8500/3)
Cancer	FibroadenomaInvasive, NOS (8500/3)
Cancer	Ductal carcinoma in situ - 8500/2;Invasive ductal carcinoma, not otherwise specified (NOS) - 8500/3;Microcalcifications;Benign;In situ and grade 3 intraepith;Malignant;
Cancer	Ductal carcinoma in situ - 8500/2;Invasive lobular carcinoma - 8520/3;Lobular carcinoma in situ - 8520/2;In situ and grade 3 intraepith;Malignant;

Cancer	Ductal carcinoma in situ - 8500/2;Invasive ductal carcinoma, not otherwise specified (NOS) - 8500/3;Microcalcifications;Benign;In situ and grade 3 intraepith;Malignant;Focal Micropapillary Features, invasive ductal carcinoma with micropapillary features, invasive ductal carcinoma with mucinous and micropapillary featInvasive ductal carcinoma with micropapillary and mucinous features;Invasive micropapillary carcinoma - 8507/3;Malignant;
Cancer	Invasive, predominant intraductal component (8500/3)
Cancer	Invasive ductal carcinoma, not otherwise specified (NOS) - 8500/3;Malignant;
Cancer	Invasive, NOS (8500/3)
Cancer	Infiltrating duct and lobular carcinoma (8522/3)
Cancer	Invasive, predominant in situ component (8520/3)
Non-Cancer	Otherusual ductal hyperplasia, apocrine metaplasia, microcysts, elastosis
Non-Cancer	Otherstromal fibrosis, fibrous cyst wall
Non-Cancer	Otherfibrocystic change, stromal fibrosis, cyst formation, microcalcifications, apocrine metaplasia, sclerosing adenosis, usual ductal hyperplasia
Non-Cancer	Otherfibrocystic changes, apocrine metaplasia, cystic change, usual ductal hyperplasia
Non-Cancer	Otherfibrocystic change, microcalcifications
Non-Cancer	Fibroadenoma
Non-Cancer	Otherintraductal papilloma, sclerosis, microcalcifications, stromal fibrosis
Non-Cancer	Fibroadenoma
Non-Cancer	Otherfat necrosis
Non-Cancer	Otherstromal fibrosis, microcalcifications
Non-Cancer	Otherfibrocystic change, microcystic change, focal secretory features
Non-Cancer	Otherstromal fibrosis
Non-Cancer	Fibroadenoma Otheradenosis, columnar cell change/hyperplasia, usual ductal hyperplasia
Non-Cancer	OtherFNA - insufficient material for diagnosis
Non-Cancer	Otherintraductal papilloma
Non-Cancer	Otherfibrocystic changes, duct ectasia, usual ductal hyperplasia, apocrine metaplasia, microcalcifications

[00974] Microvesicles (and potentially other biological entities) were precipitated in blood (plasma) samples from the above patients using polymer precipitation with PEG as indicated above. The protocol was used to probe the samples with the oligonucleotide probe libraries. Sequences that bound the PEG precipitated samples were identified using next generation sequencing (NGS).

[00975] **FIG. 12D** shows scatter plots of a selection of results from testing the 40 patients listed previously. The spread in the data indicates that large numbers of oligos were detected that differed between samples. The number of significant oligos found is much greater than would be expected randomly as shown in **Table 17**. The table shows the number of oligonucleotides sorted by copy number detected and p-value. The d-# indicates the number copies of a sequence observed for the data in the rows.

Table 17: Expected versus observed sequences

	Total Number	P-0.1	P-0.05	P-0.01	P-0.005
d-50	83,632	47,020	30,843	5,934	2,471
d-100	52,647	29,106	19,446	3,893	1,615
d-200	28,753	14,681	9,880	2,189	914
d-500	10,155	4,342	2,927	725	315

d-50	100.0%	56.2%	36.9%	7.1%	3.0%
d-100	100.0%	55.3%	36.9%	7.4%	3.1%
d-200	100.0%	51.1%	34.4%	7.6%	3.2%
d-500	100.0%	42.8%	28.8%	7.1%	3.1%
Maximum expected		10.0%	5.0%	1.0%	0.5%

[00976] As a control, the cancer and non-cancer samples were randomly divided into two groups. Such randomization of the samples significantly reduced the number of oligos found that differentiate between sample groups. Indeed, there was a 50-fold increase in informative oligos between the cancer/non-cancer grouping versus random grouping. **FIG. 12E** shows data as in **Table 17** and indicates the number of observed informative oligos between the indicated sample groups.

[00977] **FIG. 12F** shows distinct groups of oligos that differentiate between cancer and non-cancer samples. The figure shows a heatmap of the 40 samples tested with oligos selected that had more than 500 copies and p-value less than 0.005. There are clear subpopulations emerging with a distinct non-cancer cohort at the top. The non-cancer samples have boxes around them on the left axis. **FIG. 12G** is similar and shows results with an additional 20 cancer and 20 non-cancer samples. As shown, analysis with the 80 samples provides the emergence of more distinct and larger clusters.

[00978] The data for the additional 80 samples was also used to compare the consistency of informative oligos identified in different screening experiments. Of the 315 informative oligos identified using the first set of 40 patients, 86% of them showed fold-change in a consistent manner when tested on the independent set of 40 patients.

Example 22: Enrichment of Oligonucleotide probes to Breast Cancer (BrCa) and non-BrCa derived microvesicles using a balanced library design

[00979] In this Example, a naïve ADAPT oligonucleotide library was screened to enrich oligonucleotides that identify microvesicles circulating in the blood of breast cancer patients and microvesicles circulating in the blood of healthy, control individuals (i.e., without breast cancer). The input library was the naïve F-TRin-35n-B 8-3s library, which comprises a 5' region (5' CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) followed by the random naïve aptamer sequences of 35 nucleotides and a 3' region (5' CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). The "balanced" design is described in **Example 23** of Int'l Patent Publication WO/2015/031694 (Appl. No. PCT/US2014/053306, filed August 28, 2014). The working library comprised approximately 2 x 10^{13} synthetic oligonucleotide sequences. The naïve library may be referred to as the "L0 Library" herein.

[00980] The L0 Library was enriched against fractionated plasma samples from breast cancer patients and from healthy (non-breast cancer) controls using the protocol shown in **FIG. 21A**. In Step 1, an aliquot of approximately 10^{11} sequences of PCR-amplified L0 was incubated with pooled blood-plasma from 59 breast cancer patients with positive biopsy (represented by "Source A" in **FIG. 21A**). In parallel, another aliquot of 10^{11} sequences was incubated with pooled blood-plasma from 30 patients with suspected breast cancer who proved negative on biopsy and 30 self declared healthy women (represented by "Source B" in **FIG. 21A**). In Step 2, microvesicles (extracellular vesicles, "EV") were precipitated using ultracentrifugation (UC) from both L0-samples. The EV-associated oligodeoxynucleotides (ODNs) were recovered from the respective pellets. In Step 3, a counter-selection step (Step 3) was carried out by incubation of each enriched library with plasma from the different cohorts to drive the selection pressure towards enrichment of ODNs specifically associated with each sample cohort. In this step, sequences contained in the EV pellets were discarded. In Step 4, a second positive selection was performed. In this step, the sequences contained in the respective supernatants (sn) from Step 3 were mixed with plasma from another aliquot of each positive control sample-population, and EVs were again isolated. EV-associated ODNs were recovered, representing two single-round libraries called library L1 for positive enrichment of cancer (positive biopsy) patients, and library L2 for the positive enrichment against control patients. In a final step, L1 and L2 were amplified by PCR, reverted to single stranded DNA (ssDNA), and mixed to yield library L3.

[00981] This enrichment scheme was iterated two times more using L3 as the input to further reduce the complexity of the profiling library to approximately 10^6 different sequences. In Step 2, UC was used for partitioning of microvesicles, which may increase the specificity for the EV fraction. In Steps 3 and 4, partitioning was performed using PEG-precipitation. This procedure enriches for ODNs specific for each biological source. Library L3 contains those ODNs that are associated with targets characteristic for EV-populations from both sources, i.e. ODNs acting as aptamers that bind to molecules preferentially expressed in each source. A total of biopsy-positive (n = 59), biopsy-negative (n = 30), and self-declared normal (n = 30) were used in the first round of L3 enrichment, while only the cancer and non-cancer samples were used in the subsequent rounds.

[00982] The enriched libraries were characterized using next-generation-sequencing (NGS) to measure copy numbers of sequences contained in each profiling library. NGS of L0 shows that the vast majority of sequences existed in low copy numbers, whereas libraries L1 and L2 showed significantly higher average counts per sequence (**FIG. 21B**) and a reduced amount of different sequences, with unaltered total valid reads, (**FIG. 21C**) consistent with an enrichment process.

Example 23: Analysis of ADAPT-identified biomarkers

[00983] As described herein, e.g., in the section entitled "Aptamer Target Identification," an unknown target recognized by an aptamer can be identified. In this Example, an oligonucleotide probe library (also referred to as Adaptive Dynamic Artificial Poly-ligand Targeting (ADAPT) libraries or Topographical Oligonucleotide Probe "TOP" libraries) was developed as described here and targets of the screened

oligonucleotides were determined. This Example used a ADAPT library generated by enriching microvesicles collected from the blood of breast cancer patients and normal controls (i.e., non-cancer individuals). The enrichment protocols are described herein in **Example 22**.

[00984] Materials & Methods

[00985] *SBED library conjugation*

[00986] A naïve F-TRin-35n-B 8-3s library was enriched against microvesicles from normal female plasma. The naïve unenriched library comprised a 5' region (5' CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) followed by the random naïve aptamer sequences of 35 nucleotides and a 3' region (5' CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). The naïve library may be referred to as the "L0 Library" herein and the enriched library referred to as the "L2 library." See **Example 22**. The screened library was PCR amplified with a C6-amine sense primer (C6 Amine-5' CTAGCATGACTGCAGTACGT 3' (SEQ ID NO. 131)) and a 5' phosphorylated anti-sense primer (5' Phos TCGTCGGCAGCGTCA (SEQ ID NO. 133)), the purified product was strand separated and conjugated with sulfo-SBED (Thermo Scientific) according to Vinkenborg et al. (Angew Chem Int Ed Engl. 2012, 51:9176-80) with the following modifications: The reaction was scaled down to 5µg C6-amine DNA library (8.6 µM) in 25mM HEPES-KOH, 0.1M NaCl, pH 8.3 and incubated with either 100-fold molar excess of sulfo-SBED or DMSO in a 21µL volume for 30 min at room temp in the dark. The SBED-conjugated library was immediately separated from the unconjugated library and free sulfo-SBED by injection onto a Waters X-BridgeTM OST C-18 column (4.6 mm x 50 mm) and fractionated by HPLC (Agilent 1260 Infinity) with a linear gradient Buffer A: 100 mM TEAA, pH7.0, 0% ACN to 100 mM TEAA, pH7.0, 25% ACN at 0.2ml/min, 65 °C. There SBED-conjugated fractions were desalted into water with Glen Gel-PakTM Cartridges and concentrated by speed-vac. SBED conjugation was confirmed by LC-MS and/or a dot blot with streptavidin-HRP detection.

[00987] *Binding reaction and cross-linking*

[00988] SBED library functionalization was tested by performing the ADAPT assay with SBED vs DMSO mock conjugated control C6-amine library and sequenced on a HiSeq 2500TM (Illumina Corp.). The aptamer precipitation was performed with forty-eight ADAPT reactions incubated for 1hr with end-over-end rotation at room temp with a 5ng input of SBED conjugated library per 200 µL of plasma (pre-spun to remove cellular debris at 10,000 xg for 20min, 4 °C) in 1X PBS, 3mM $MgCl_2$, 0.01mM dextran sulfate, 40 ng/µl salmon sperm DNA and 40 ng/µl yeast transfer RNA, and cOmplete ULTRA Mini EDTA-free TM protease inhibitors (Roche) equivalent to ~240ng library and 9.6 mls plasma. A duplicate set of 48 reactions was prepared with the DMSO control C6-amine library. Aptamer library-protein complexes were precipitated with incubation in 6% PEG8000 for 15 min at 4 °C then centrifuged at 10,000 xg for 5 min. Pellets were washed with 1ml 1x PBS, 3mM MgCl2 by gentle inversion to remove unbound aptamers. The washed pellets were resuspended in 100µL of water and subjected to photo-cross-linking at 365nm with a hand-held 3UV (254NM/302NM/365NM) lamp, 115 volts (Thermo Scientific) for 10 min on ice with 1-2 cm between the 96-well plate and lamp.

[00989] *Oligonucleotide precipitation*

[00990] Cross-linked reactions were subsequently pooled (~4.8ml) per library or 4.8 ml of 1X PBS (AP bead only control) and incubated with 10 µL of Prepared Dynabeads® MyOne™ Streptavidin C1 (10mg/ml) (Life Technologies) (pre-washed with 1X PBS, 0.01% Triton X-100) shaking for 1 hr at room temp. Beads were transferred to an eppendorf tube and lysed for 20 min with lysis buffer (50 mM Tris-HCl, 10mM MgCl2, 200mM NaCl, 0.5% Triton X-100, 5% glycerol, pH 7.5) on ice, washed 3 times with wash buffer 1 (10mM Tris-HCl, 1mM EDTA, 2M NaCl, 1% Triton X-100), followed by 2 times with wash buffer 2 (10mM Tris-HCl, 1mM EDTA, 2M NaCl, 0.01% Triton X-100) as described by Vinkenborg et al. (Angew Chem Int Ed Engl. 2012, 51:9176-80). Cross-linked proteins were eluted by boiling 15 min in 1X LDS sample buffer with reducing agent added (Life Technologies) and loaded on a 4-12% SDS-PAGE gradient gel (Life Technology). Proteins and DNA were detected with double staining with Imperial Blue Protein Stain (Thermo Scientific) followed by Prot-SIL2 TM silver stain kit (Sigma) used according to manufacturer's instructions in order to enhance sensitivity and reduce background.

[00991] *Protein identification*

[00992] Protein bands that appeared to differ between the cancer and normal were excised from the gradient gels and subjected to liquid chromatography-tandem mass spectrometry (LC-MS/MS).

[00993] Results

[00994] ADAPT protein targets were identified from bands cut from a silver stained SDS-PAGE gel (**FIG. 13**). Aptamer-SBED protein complexes (lane 3) or Aptamer-DMSO protein complexes (control-lane 4) were precipitated with 6% PEG8000, subjected to UV photo-cross-linking, and pulled-down with Streptavidin coated beads. Eluate was analyzed under reducing conditions by SDS-PAGE and silver staining. Aptamer library alone (5ng) (lane 1) was loaded as a control for migration of the library (second to bottom arrows) and an equal volume of eluate from a bead only sample (lane 4) was loaded as a streptavidin control to control for potential leaching of the streptavidin monomer (bottom arrow) under the harsh elution conditions. Upper arrows ("Targets") indicate specific or more predominant bands identified with the SBED-conjugated library vs. the mock DMSO treated control C6-amine library. Indicated target protein bands were cut out and sent for LC-MS/MS protein identification or indicated DNA library bands were eluted, reamplified and sequenced. The identified proteins are those that appeared as upregulated in the normal samples.

[00995] Tables 18-25 list human proteins that were identified in 8 bands excised from the silver stained gel. In all tables the proteins are those identified in the oligo-SBED protein complexes with proteins identified in the corresponding control lanes removed. The band numbers in the tables indicate different bands cut from the gel (**FIG. 13**). Accession numbers in the table are from the UniProt database (www.uniprot.org). "GN=" is followed by the gene name. Various protein classifications indicated in the **Tables 18-25** include Nucleic Acid Binding Proteins (NAB), Tumor suppressors (TS), cell adhesion/cytoskeletal (CA/CK) and abundant plasma proteins (ABP). In Table 18, the proteins listed

below the row "SBED associated" were identified by peptide fragments linked to SBED, indicating that the peptides are from near the cross-linking sites. Class is not reported for these proteins.

Table 18: Band 3

Accession number	Class	Protein name
P02538	CA/CK	Keratin, type II cytoskeletal 6A GN=KRT6A
P15924	CA/CK	Desmoplakin GN=DSP
P04259	CA/CK	Keratin, type II cytoskeletal 6B GN=KRT6B
P60709	CA/CK	Actin, cytoplasmic 1 GN=ACTB
P20930	CA/CK	Filaggrin GN=FLG
P07476	CA/CK	Involucrin GN=IVL
P31947	TS	14-3-3 protein sigma GN=SFN
Q7Z794	CA/CK	Keratin, type II cytoskeletal 1b GN=KRT77
P02545	NAB	Prelamin-A/C GN=LMNA
P19012	CA/CK	Keratin, type I cytoskeletal 15 GN=KRT15
P47929	CA/CK & TS	Galectin-7 GN=LGALS7
P11142		Heat shock cognate 71 kDa protein GN=HSPA8
P58107	NAB	Epiplakin GN=EPPK1
P08107		Heat shock 70 kDa protein 1A/1B GN=HSPA1A
Q02413	CA/CK	Desmoglein-1 GN=DSG1
P06396	CA/CK	Gelsolin GN=GSN
O60814	NAB	Histone H2B type 1-K GN=HIST1H2BK
P68104	NAB	Elongation factor 1-alpha 1 GN=EEF1A1
P05387	NAB	60S acidic ribosomal protein P2 GN=RPLP2
Q7RTS7	CA/CK	Keratin, type II cytoskeletal 74 GN=KRT74
P31946	TS	14-3-3 protein beta/alpha GN=YWHAB
Q13835	CA/CK	Plakophilin-1 GN=PKP1
P14923	CA/CK	Junction plakoglobin GN=JUP
P09651	NAB	Heterogeneous nuclear ribonucleoprotein A1 GN=HNRNPA1
P07900		Heat shock protein HSP 90-alpha GN=HSP90AA1
Q96KK5	NAB	Histone H2A type 1-H GN=HIST1H2AH
P04406-	CA/CK	Glyceraldehyde-3-phosphate dehydrogenase GN=GAPDH
P10412	NAB	Histone H1.4 GN=HIST1H1E
P04792		Heat shock protein beta-1 GN=HSPB1
Q9NZT1		Calmodulin-like protein 5 GN=CALML5
P81605		Dermcidin GN=DCD
P27348	TS	14-3-3 protein theta GN=YWHAQ
P55072	NAB	Transitional endoplasmic reticulum ATPase GN=VCP
Q09666	NAB	Neuroblast differentiation-associated protein AHNAK GN=AHNAK
P23246	NAB	Splicing factor, proline- and glutamine-rich GN=SFPQ
Q15149	CA/CK	Plectin GN=PLEC
Q8NC51	NAB	Plasminogen activator inhibitor 1 RNA-binding protein GN=SERBP1
P07237		Protein disulfide-isomerase GN=P4HB
O60437	CA/CK	Periplakin GN=PPL
P01717	ABP	Ig lambda chain V-IV region Hil
P55884	NAB	Eukaryotic translation initiation factor 3 subunit B GN=EIF3B
P11021		78 kDa glucose-regulated protein GN=HSPA5
P01024		Complement C3 GN=C3
P04350	CA/CK	Tubulin beta-4A chain GN=TUBB4A
P01857	ABP	Ig gamma-1 chain C region GN=IGHG1
P61247	NAB	40S ribosomal protein S3a GN=RPS3A
P62937		Peptidyl-prolyl cis-trans isomerase A GN=PPIA

O15020	CA/CK	Spectrin beta chain, non-erythrocytic 2 GN=SPTBN2
P30101		Protein disulfide-isomerase A3 GN=PDIA3
Q6KB66	CA/CK	Keratin, type II cytoskeletal 80 GN=KRT80
Q9UJU6	CA/CK	Drebrin-like protein GN=DBNL
P47914	NAB	60S ribosomal protein L29 GN=RPL29
P39023	NAB	60S ribosomal protein L3 GN=RPL3
A6NMY6	CA/CK	Putative annexin A2-like protein GN=ANXA2P2
P60174	CA/CK	Triosephosphate isomerase GN=TPI1
P35241	CA/CK	Radixin GN=RDX
P07305	NAB	Histone H1.0 GN=H1F0
P15259	CA/CK	Phosphoglycerate mutase 2 GN=PGAM2
P0CG05	ABP	Ig lambda-2 chain C regions GN=IGLC2
Q92817	CA/CK	Envoplakin GN=EVPL
P06733	NAB	MBP-1 of Alpha-enolase GN=ENO1
P22626	NAB	Heterogeneous nuclear ribonucleoproteins A2/B1 GN=HNRNPA2B1
P62424	NAB	60S ribosomal protein L7a GN=RPL7A
P60660	CA/CK	Myosin light polypeptide 6 GN=MYL6
P04083	NAB	Annexin A1 GN=ANXA1
Q14134	NAB	Tripartite motif-containing protein 29 GN=TRIM29
P39019	NAB	40S ribosomal protein S19 GN=RPS19
Q8WVV4	CA/CK	Protein POF1B GN=POF1B
Q02878	NAB	60S ribosomal protein L6 GN=RPL6
Q9Y6X9	NAB	MORC family CW-type zinc finger protein 2 GN=MORC2
Q9NQC3	NAB	Reticulon-4 GN=RTN4
Q5T753	CA/CK	Late cornified envelope protein 1E GN= CA/CK E
SBED associated		
P56202		Cathepsin W
P80188		Neutrophil gelatinase-associated lipocalin precursor
Q13017		Rho GTPase-activating protein 5
Q6UB98		Ankyrin repeat domain-containing protein 12
P54753		Ephrin type-B receptor 3
Q5JRS4		Olfactory receptor 10J3
P82279		Protein crumbs homolog 1
O00763		Acetyl-CoA carboxylase 2
P02533; P08779		Keratin, type 1 cytoskeletal 14, 16
P26012		Integrin beta-8
Q14766		Latent-transforming growth factor beta-binding protein 1

Table 19: Band 9

Accession number	Class	Protein name
P61626		Lysozyme C GN=LYZ
Q9HCK1	NAB	DBF4-type zinc finger-containing protein 2 GN=ZDBF2

Table 20: Band 1

Accession number	Class	Protein name
P01834	ABP	Ig kappa chain C region GN=IGKC
P01765	ABP	Ig heavy chain V-III region TIL
P04003	NAB	C4b-binding protein alpha chain GN=C4BPA
P60709	CA/CK	Actin, cytoplasmic 1 GN=ACTB

Q5T751	CA/CK	Late cornified envelope protein 1C GN=LCE1C

Table 21: Band 5

Accession number	Class	Protein name
P01860	ABP	Ig gamma-3 chain C region GN=IGHG3
O60902	NAB	Short stature homeobox protein 2 GN=SHOX2

Table 22: Band 7

Accession number	Class	Protein name
Q04695	CA/CK	Keratin, type I cytoskeletal 17 GN=KRT17
Q7Z794	CA/CK	Keratin, type II cytoskeletal 1b GN=KRT77
Q6KB66	CA/CK	Keratin, type II cytoskeletal 80 GN=KRT80
P01833		Polymeric immunoglobulin receptor GN=PIGR
P01042		Kininogen-1 GN=KNG1
Q02413	CA/CK	Desmoglein-1 GN=DSG1
P15924	CA/CK	Desmoplakin GN=DSP
Q8TF72		Protein Shroom3 GN=SHROOM3
P02671	ABP	Fibrinogen alpha chain GN=FGA
Q5T749	CA/CK	Keratinocyte proline-rich protein GN=KPRP
Q5VZP5		Inactive dual specificity phosphatase 27 GN=DUSP27
Q5T751	CA/CK	Late cornified envelope protein 1C GN=LCE1C
Q9UL12		Sarcosine dehydrogenase, mitochondrial GN=SARDH
P00698		Lysozyme C OS=Gallus gallus GN=LYZ
Q8N114		Protein shisa-5 GN=SHISA5

Table 23: Band 15

Accession number	Class	Protein name
P08238		Heat shock protein HSP 90-beta GN=HSP90AB1
P68104	NAB	Elongation factor 1-alpha 1 GN=EEF1A1
P02675	ABP	Fibrinogen beta chain GN=FGB
Q8TF72		Protein Shroom3 GN=SHROOM3
P0CG05	ABP	Ig lambda-2 chain C regions GN=IGLC2
P78386	CA/CK	Keratin, type II cuticular Hb5 GN=KRT85
Q7Z5Y6		Bone morphogenetic protein 8A GN=BMP8A
O14633	CA/CK	Late cornified envelope protein 2B GN=LCE2B

Table 24: Band 17

Accession number	Class	Protein name
P02538	CA/CK	Keratin, type II cytoskeletal 6A GN=KRT6A
P01834	ABP	Ig kappa chain C region GN=IGKC
P06702		Protein S100-A9 GN=S100A9
P68104	NAB	Elongation factor 1-alpha 1 GN=EEF1A1
P01024		Complement C3 GN=C3
P81605		Dermcidin GN=DCD
P05109		Protein S100-A8 GN=S100A8
Q5T751	CA/CK	Late cornified envelope protein 1C GN=LCE1C

Table 25: Band 19

Accession number	Class	Protein name
P02768	NAB	Serum albumin GN=ALB
P0CG05	ABP	Ig lambda-2 chain C regions GN=IGLC2
P06702		Protein S100-A9 GN=S100A9
P08238		Heat shock protein HSP 90-beta GN=HSP90AB1
P60709	CA/CK	Actin, cytoplasmic 1 GN=ACTB
P13647	CA/CK	Keratin, type II cytoskeletal 5 GN=KRT5
P01616	ABP	Ig kappa chain V-II region MIL
Q86YZ3	CA/CK	Hornerin GN=HRNR
P01857	ABP	Ig gamma-1 chain C region GN=IGHG1
P62805	NAB	Histone H4 GN=HIST1H4A
P59665		Neutrophil defensin 1 GN=DEFA1
P61626		Lysozyme C GN=LYZ
P01024	ABP	Complement C3 GN=C3
Q8TF72		Protein Shroom3 GN=SHROOM3
P83593	ABP	Ig kappa chain V-IV region STH (Fragment)
P01700	ABP	Ig lambda chain V-I region HA
P01877	ABP	Ig alpha-2 chain C region GN=IGHA2
Q9UL12		Sarcosine dehydrogenase, mitochondrial GN=SARDH
Q6NXT2	NAB	Histone H3.3C GN=H3F3C
P02788	NAB	Lactotransferrin GN=LTF
P02787	ABP	Serotransferrin GN=TF

[00996] Certain proteins were identified in multiple bands. For example, IGLC2 was identified in bands 3, 15 and 19 and SHROOM3 was identified in bands 7, 15, 19. This may be due to degradation products, isoforms or the like. These experiments identified 108 proteins (plus 2 lysozyme controls), comprising among others 34 Nucleic Acid Binding Proteins (NAB) where 7 of the 34 are putative tumor suppressors/repressors; 37 cell adhesion/cytoskeletal (CA/CK); and 14 abundant plasma proteins (ABP). All of the tumor suppressors/repressors are DNA/RNA binding proteins. Other proteins comprise chaperones, signaling molecules etc.

[00997] For comparison to the above pull-down results with the enriched L2 library, mass spectrometry was used to identify proteins in whole plasma samples ("neat plasma") (**Table 26**, row "Neat"), in PEG enriched plasma fractions (comprising a microvesicle-containing fraction) (**Table 26**, row "PEG"), and in unenriched L0 library pull-down (**Table 26**, row "Unenriched") using the same healthy (non-breast cancer) plasma samples as described above (column "Plasma," rows "Non-cancer" in the table). Descriptions in the table comprise the gene name followed by accession numbers from the UniProt database (www.uniprot.org) in parentheses. 81 proteins were pulled down by L2 (**Tables 18-25**) that were not detectable in the PEG-precipitated and neat plasma, indicating that these proteins became enriched due to their interaction with L2 oligonucleotide probes. Further details are found in Example 28 of International Patent Application PCT/US15/62184, filed November 23, 2015, which application is incorporated by reference herein in its entirety. The methods above were further repeated to identify proteins that were identified in cancer samples using pull-down with the L1 library (column "Plasma,"

rows "Cancer" in the table). With the proteins identified in the cancer plasma samples, the row "Non-SBED" and "SBED refer to proteins identified by peptide fragments that were unassociated with or associated with SBED, respectively.

Table 26: Proteins identified in plasma

Plasma	Sample	Description (Gene Name (Accession))
Non-cancer	Neat	ALB (P02768), C3 (P01024), APOB (P04114), TF (P02787), A2M (P01023), C4B (P0C0L5), C4A (P0C0L4), FGA (P02671), FGB (P02675), CP (P00450), IGKC (P01834), IGHG1 (P01857), HP (P00738), FGG (P02679), IGHG3 (P01860), CFH (P08603), GC (P02774), IGHG2 (P01859), SERPINA1 (P01009), ITIH4 (Q14624), SERPINC1 (P01008), IGHM (P01871), HPR (P00739), CFB (P00751), APOA1 (P02647), FN1 (P02751), IGLL5 (B9A064), PLG (P00747), A1BG (P04217), ITIH2 (P19823), SERPINA3 (P01011), KNG1 (P01042), IGHG4 (P01861), APOA4 (P06727), F2 (P00734), HPX (P02790), C4BPA (P04003), ITIH1 (P19827), IGHA1 (P01876), IGLC2 (P0CG05), SERPING1 (P05155), GSN (P06396), PON1 (P27169), AGT (P01019), APOH (P02749), HRG (P04196), APCS (P02743), AFM (P43652), IGHA2 (P01877), APOE (P02649), AHSG (P02765), SERPIND1 (P05546), C5 (P01031), AMBP (P02760), ORM1 (P02763), AZGP1 (P25311), KRT1 (P04264), CFI (P05156), HBB (P68871), CD5L (O43866), C9 (P02748), RBP4 (P02753), VTN (P04004), SERPINA6 (P08185), SERPINF2 (P08697), CLU (P10909), KRT2 (P35908), C1QC (P02747), TTR (P02766), Ig kappa chain V-II region TEW (P01617), Ig heavy chain V-III region BUT (P01767), Ig heavy chain V-III region TUR (P01779), APOA2 (P02652), ORM2 (P19652), CFHR1 (Q03591), PGLYRP2 (Q96PD5), LYZ (P00698), Ig kappa chain V-IV region Len (P01625), Ig heavy chain V-III region GAL (P01781), IGHD (P01880), C1QB (P02746), C8A (P07357), C8G (P07360), KRT10 (P13645), APOL1 (O14791), IGJ (P01591), KLKB1 (P03952), Ig lambda chain V region 4A (P04211), Ig kappa chain V-III region VG (Fragment) (P04433), HBA1 (P69905), APOM (O95445), C1R (P00736), F12 (P00748), Ig lambda chain V-III region SH (P01714), Ig lambda chain V-IV region Hil (P01717), Ig heavy chain V-III region WEA (P01763), Ig heavy chain V-II region NEWM (P01825), LRG1 (P02750), IGKV4-1 (P06312), PROS1 (P07225), PRSS1 (P07477), C7 (P10643), C6 (P13671), CPN2 (P22792), SERPINF1 (P36955), Ig lambda chain V-III region LOI (P80748), ABCF1 (Q8NE71), FCN3 (O75636), F10 (P00742), Ig heavy chain V-I region HG3 (P01743), Ig heavy chain V-III region HIL (P01771), Ig heavy chain V-II region OU (P01814), Ig kappa chain V-I region BAN (P04430), APOD (P05090), C2 (P06681), C8B (P07358), LPA (P08519), C1S (P09871), HIST1H1D (P16402), Ig heavy chain V-I region V35 (P23083), LGALS3BP (Q08380), ATF7IP (Q6VMQ6), RASSF6 (Q6ZTQ3), MTMR14 (Q8NCE2), DCHS1 (Q96JQ0), HIST1H2AH (Q96KK5), HMCN1 (Q96RW7), UPF3A (Q9H1J1), IGLC7 (A0M8Q6), UNC13B (O14795), APBA3 (O96018), GSN (P06396), C4A (P0C0L4), PZP (P20742), SERPINA4 (P29622), SHMT2 (P34897), KRT9 (P35527), SEPP1 (P49908), COG7 (P83436), ITIH3 (Q06033), TMEM198 (Q66K66), F13A1 (P00488), Ig kappa chain V-I region Lay (P01605), S100A6 (P06703), KRT5 (P13647), CDH2 (P19022), SAA4 (P35542), RXRG (P48443), ARFGEF3 (Q5TH69), IQCE (Q6IPM2), C19orf68 (Q86XI8), CELSR3 (Q9NYQ7), C1RL (Q9NZP8), SARDH (Q9UL12), MYH4 (Q9Y623)
Non-cancer	PEG	C3 (P01024), A2M (P01023), APOB (P04114), IGKC (P01834), C4A (P0C0L4), C4B (P0C0L5), FGB (P02675), ALB (P02768), CFH (P08603), IGHG1 (P01857), FGA (P02671), FN1 (P02751), PLG (P00747), IGHM (P01871), FGG (P02679), TF

		(P02787), C5 (P01031), CP (P00450), IGHG2 (P01859), IGLC2 (P0CG05), Ig mu heavy chain disease protein (P04220), ITIH1 (P19827), PZP (P20742), IGHG3 (P01860), IGLL5 (B9A064), HP (P00738), C4BPA (P04003), ITIH2 (P19823), IGHA1 (P01876), KRT1 (P04264), KRT10 (P13645), APOE (P02649), Ig kappa chain V-I region DEE (P01597), AMBP (P02760), F2 (P00734), C7 (P10643), C6 (P13671), ITIH4 (Q14624), CFB (P00751), IGHG4 (P01861), APOH (P02749), APOA1 (P02647), CD5L (O43866), C1R (P00736), HPR (P00739), Ig kappa chain V-I region Scw (P01609), IGHA2 (P01877), CFHR1 (Q03591), KRT2 (P35908), Ig kappa chain V-III region SIE (P01620), HRG (P04196), Ig heavy chain V-III region BRO (P01766), C1QB (P02746), GC (P02774), Ig heavy chain V-III region TIL (P01765), Ig kappa chain V-III region NG9 (Fragment) (P01621), Ig heavy chain V-III region BUT (P01767), Ig heavy chain V-III region TUR (P01779), C9 (P02748), SERPIND1 (P05546), Ig kappa chain V-I region WEA (P01610), Ig kappa chain V-I region Ni (P01613), Ig kappa chain V-IV region Len (P01625), Ig kappa chain V-I region EU (P01598), Ig kappa chain V-II region TEW (P01617), Ig heavy chain V-III region GAL (P01781), KNG1 (P01042), VTN (P04004), C8B (P07358), Ig lambda chain V-III region LOI (P80748), Ig heavy chain V-II region NEWM (P01825), APCS (P02743), KLKB1 (P03952), CFI (P05156), PROS1 (P07225), LPA (P08519), KRT9 (P35527), SERPINA1 (P01009), Ig lambda chain V-III region SH (P01714), C8A (P07357), Ig kappa chain V-III region B6 (P01619), Ig lambda chain V-IV region Hil (P01717), Ig kappa chain V-III region CLL (P04207), C1S (P09871), FCN3 (O75636), SERPINC1 (P01008), Ig kappa chain V-I region Mev (P01612), IGHD (P01880), C1QC (P02747), HPX (P02790), C8G (P07360), IGKV1-5 (P01602), Ig kappa chain V-I region Wes (P01611), Ig heavy chain V-III region WEA (P01763), A1BG (P04217), GSN (P06396), FBLN1 (P23142), HBB (P68871), ITIH3 (Q06033), F12 (P00748), SERPINA3 (P01011), APOC3 (P02656), Ig kappa chain V-I region BAN (P04430), Ig kappa chain V-III region VH (Fragment) (P04434), F13B (P05160), IGKV4-1 (P06312), SERPINF2 (P08697), CLU (P10909), HIST1H1D (P16402), PON1 (P27169), IGJ (P01591), Ig kappa chain V-III region POM (P01624), Ig heavy chain V-III region CAM (P01768), Ig heavy chain V-III region BUR (P01773), Ig kappa chain V-III region VG (Fragment) (P04433), APOD (P05090), Ig lambda chain V-IV region MOL (P06889), Ig heavy chain V-III region GAR (P80419), FCGBP (Q9Y6R7), APOM (O95445), F13A1 (P00488), Ig heavy chain V-I region HG3 (P01743), C1QA (P02745), Ig lambda chain V-VI region WLT (P06318), C2 (P06681), C4BPB (P20851), CFP (P27918), SERPINA4 (P29622), SAA4 (P35542), SERPINF1 (P36955), LGALS3BP (Q08380), HABP2 (Q14520), RCBTB1 (Q8NDN9), APOL1 (O14791), KCNQ2 (O43526), F9 (P00740), Ig heavy chain V-III region TRO (P01762), Ig heavy chain V-III region HIL (P01771), Ig heavy chain V-II region OU (P01814), APOA2 (P02652), F11 (P03951), Ig lambda chain V-I region WAH (P04208), Ig lambda chain V region 4A (P04211), Ig kappa chain V-II region RPMI 6410 (P06310), Ig kappa chain V-III region IARC/BL41 (P06311), KRT5 (P13647), IGLL1 (P15814), Ig heavy chain V-I region V35 (P23083), HBA1 (P69905), ADIPOQ (Q15848), PGLYRP2 (Q96PD5), UPF3A (Q9H1J1), BCO1 (Q9HAY6), ARFGAP3 (Q9NP61), SARDH (Q9UL12), SERPINA1 (P01009), KNG1 (P01042), Ig kappa chain V-I region Kue (P01604), Ig kappa chain V-I region Lay (P01605), Ig kappa chain V-I region OU (P01606), Ig kappa chain V-II region MIL (P01616), Ig heavy chain V-III region VH26 (P01764), Ig heavy chain V-III region GA (P01769), FN1 (P027510), TTR (P02766), SERPING1 (P05155), APOA4 (P06727), PRSS1 (P07477), ANXA6 (P08133), CFTR (P13569), LBP (P18428), FBLN1 (P23142), SPAG17 (Q6Q759), PDLIM2 (Q96JY6),

		ARHGEF17 (Q96PE2), IGLC7 (A0M8Q6), AGRN (O00468), AGT (P01019), RBP4 (P02753), AHSG (P02765), Ig kappa chain V-III region GOL (P04206), SERPINA5 (P05154), GSN (P06396), Ig kappa chain V-III region HAH (P18135), CFHR2 (P36980), GIT2 (Q141611), INCENP (Q9NQS7
Non-cancer	Unenriched	LYZ (P00698), IGHG1 (P01857), KRT14 (P02533), KRT6A (P02538), APOA1 (P02647), FGA (P02671), ALB (P02768), HSPB1 (P04792), COL1A2 (P08123), KRT16 (P08779), IGLC6 (P0CF74), KRT5 (P13647), DSP (P15924), DSP (P15924), LGALS7 (P47929), ACTG2 (P63267), DCD (P81605), DSG1 (Q02413), FLG2 (Q5D862), SBSN (Q6UWP8), KRT73 (Q86Y46), HRNR (Q86YZ3), KRT78 (Q8N1N4), SHROOM3 (Q8TF72), TREX2 (Q9BQ50), SPATA7 (Q9P0W8)
Cancer	Non-SBED	MUC5B (Q9HC84), FABP5 (Q01469), HPX (P02790), CP (P00450), SPRR2E (P22531), SPRR2D (P22532), PDE4D (Q08499-7), GC (P02774-2), CPD (O75976), CD14 (P08571), LAP3 (P28838-2), AFM (P43652), FCN2 (Q15485-2), DMBT1 (Q9UGM3-9), LIFR (P42702), SNX27 (Q96L92-3), LCN1 (P31025), ARFIP1 (P53367-2), APOH (P02749), KLKB1 (P03952), XP32 (Q5T750), H2AFV (Q71UI9-5), KRT75 (O95678), KRT6C (P48668), KRT83 (P78385), KRT76 (Q01546), KRT33B (Q14525), KRT72 (Q14CN4-3), KRT31 (Q15323), KRT73 (Q86Y46-2), DSG1 (Q02413-2), LCE1C (Q5T751), LCE1A (Q5T7P2), CFB (P00751), CFH (P08603), SERPINA1 (P01009-2), Ig kappa chain V-I region EU (P01598), Ig kappa chain V-II region MIL (P01616), Ig lambda chain V-IV region Bau (P01715), Ig heavy chain V-III region GAL (P01781), IGLC6 (P0CF74) and ACTG2 (P63267-2)
Cancer	SBED	Probable ubiquitin carboxyl-terminal hydrolase FAF-Y (O00507-1; O00507-2); Homeobox protein Hox-C12 (P31275); putative uncharacterized protein C18orf65 (Q6ZTR6); Piwi-like protein 2 (Q8TC59-2; Q8TC59-1); Probable ubiquitin carboxyl-terminal hydrolase FAF-X (Q93008-1; Q93008-3); Histone-lysine N-methyltransferase ASH1L (Q9NR48; Q9NR48-2); PNMA-like protein 2 (Q9ULN7-5)

[00998] The biomarkers in this Example can be used to detect microvesicles that are indicative of cancer or non-cancer samples.

Example 24: Identification of biomarkers through affinity enrichment with an enriched oligonucleotide library and mass spectrometry

[00999] This Example continues upon the Example above. Identification of protein-protein and nucleic acid-protein complexes by affinity purification mass spectrometry (AP-MS) can be hampered in samples comprising complex mixtures of biological components (e.g., bodily fluids including without limitation blood and derivatives thereof). For example, it may be desireable to detect low abundance protein and nucleic acid-protein complexes in a complex milieu comprising various components that may interact promiscuously with specific binding sites such as high abundance proteins that interact non-specifically with the affinity resin. AP-MS has been used previously to enrich for pre-identified targets of interest using individual DNA or RNA aptamers or specific nucleic acid binding domains. In this Example, an enriched oligonucleotide probing library was used as the affinity reagent. This approach combined with mass spectrometry enables the identification of differentially expressed biomarker from different disease states or cellular perturbations without relying on a priori knowledge of the targets of interest. Such

biomarker may comprise proteins, nucleic acids, miRNA, mRNA, carbohydrates, lipid targets, combinations thereof, or other components in a biological system.

[001000] The method comprises identification of an enriched oligonucleotide probe library according to the methods of the invention followed by target identification with affinity purification of the bound probing library and mass spectrometry. The members of the enriched oligonucleotide probing library comprise an affinity tag. A biological sample is probed with the oligonucleotide probe library, affinity purification of the oligonucleotide probe library via the affinity tag is performed which will accordingly purify biological entities in complex with various members of the probe library, and read-out of targets that purified with the members of the probe library is performed using liquid chromatography-tandem mass spectrometry (LC-MS/MS) for proteins or oligonucleotide targets (e.g., miRNA or mRNA) with next generation sequencing (NGS). Confirmation of protein targets is performed using quantitative mass spectrometry (MS), e.g., using MRM/SRM or SWATH based methods.

[001001] The method of the Example lends itself to various options. For example, any appropriate affinity tags can be used for affinity pull-down, including without limitation anti-sense oligonucleotides, biotin, polyhistidine, FLAG octapeptide (i.e., N-DYKDDDDK-C (SEQ ID NO. 134), where N stands for Amino-terminus and C stands for Carboxy terminus), 3X FLAG, Human influenza hemagglutinin (HA)-tag (i.e., N-YPYDVPDYA-C (SEQ ID NO. 135)), myc-tag (N-EQKLISEEDL-C (SEQ ID NO. 136)), other such as known in the art, and combinations thereof. Similarly, any appropriate enrichment support can be used in addition to the magnetic streptavidin beads exemplified herein, including without limitation other bead systems, agarose beads, planar arrays or column chromatography supports. It follows that the various supports can be coupled with the various affinity reagents appropriate for the oligonucleotide library, including without limitation streptavidin, avidin, anti-His tag antibodies, nickel, and the like. The different affinity tags and supports can be combined as desired. This Example used cross-linking but in certain cases such cross-linking is not necessary and may even be undesirable, e.g., to favor identification of high affinity complex formation. When cross-linking is desired, any appropriate cross-linkers can be used to carry out the invention, including BS2G, DSS, formaldehyde, and the like. Other appropriate cross-linkers and methods are described herein. See, e.g., Section "Aptamer Target Identification." Lysis buffers and wash stringencies can be varied, e.g, depending on whether complexes are cross-linked or not. Less stringent lysis/wash conditions may produce a wider array of potential protein complexes of interest whereas more stringent lysis/wash conditions may favor higher affinity oligo-target complexes and/or targets comprising specific proteins (e.g., by disassociating larger complexes bound to the oligos). One of skill will further appreciate that qualitative and/or quantitative LC-MS/MS may be used for target detection and verification. Similarly, metabolic labeling and label-free approaches may be used for quantitative MS, including without limitation spectral counting, SILAC, dimethyl labeling, TMT labeling, Targeted MS with SRM/MRM or SWATH, and the like.

[001002] References:

[001003] Vickenborg et al. "Aptamer based affinity labeling of proteins", Angew Chem Int. 51(36):9176-80 (2012).

[001004] Tacheny, M, Arnould, T., Renard, A. "Mass spectrometry-based identification of proteins interacting with nucleic acids", Journal of Proteomics 94; 89-109 (2013).

[001005] Faoro C and Ataide SF. "Ribonomic approaches to study the RNA-binding proteome.", FEBS Lett. 588(20):3649-64 (2014).

[001006] Budayeva HG, Cristea, IM, "A mass spectrometry view of stable and transient protein inteeractions." Adv Exp Med Biol. 806:263-82 (2014).

Example 25: Protocol for Affinity capture using oligonucleotide probing library

[001007] This Example presents a detailed protocol for the method of affinity capture using an oligonucleotide probing library presented in the Example above.

[001008] Protocol:

[001009] The oligonucleotide probe library comprises F-TRin-35n-B-8-3s described herein either desthiobiotin labeled or unlabeled library and binding to normal (i.e., non-cancer) female plasma. The oligonucleotide probe library is enriched against the plasma samples as described elsewhere (e.g., in **Example 21**). The plasma samples are processed separately against the desthiobiotin labeled or unlabeled oligonucleotide libraries. General parameters included the following:

[001010] 48 normal plasma samples are pooled for enrichment of each oligonucleotide library (Desthiobiotin or Unlabeled)

[001011] 200 µl input plasma per sample

[001012] Ultracentrifugation (UC) is used to pre-clear the samples

[001013] 5 ng of each aptamer library is added to each sample

[001014] Binding competitors for all library samples include 0.01mM dextran sulfate, 340ng for tRNA and 340 ng Salmon sperm DNA as described elsewhere herein

[001015] 6% PEG 8000 is used for precipitation of microvesicles within the samples

[001016] Affinity purification is performed with C1 Streptavidin beads (MyOne Strptavidin Beads C1- 65001, lot 2ml (10mg/ml))

[001017] Buffers:

[001018] Plasma dilution: 6 mM MgCl2 in 2X PBS

[001019] Pellet Wash Buffer: 1X PBS, 3mM MgCl2

[001020] PEG Ppt Buffer: 20% Peg8000 in 1X PBS, 3mM MgCl2

[001021] Bead Prep Buffer: 1XPBS containing 0.01% Triton X-100

[001022] Lysis Buffer: prepare a 2X stock solution consisting of 100mM Tris-HCl, 20mM MgCl2, 400mM NaCl, 1% Triton X-100, 10% glycerol, pH 7.5. Diluted to 1X with water 1:1 prior to using.

[001023] AP Wash buffer 1: 10mM Tris-HCl, 1mM EDTA, 2M NaCl, 1% Triton X-100, pH 7.5

[001024] AP wash buffer 2: 10mM Tris-HCL, 1mM EDTA, 2M NaCl, 0.01% Triton X-100, pH 7.5

[001025] Biotin Elution buffer 1: 5mM Biotin, 20mM Tris, 50mM NaCl, pH 7.5

[001026] 1X LDS, 1X Reducing buffer 2

[001027] Reagent/Instrument Prep:

[001028] Pre-chill Ultracentrifuge to 4 °C.

[001029] Protease inhibition: dissolve 2 tablets of "cOmplete ULTRA MINI EDTA-free EASYpack" protease inhibitor in 1100 µl of H2O (20x stock of protease inhibitor).

[001030] Plasma Preparation (for each of Desthiobiotin or Unlabeled oligonucleotide libraries):

[001031] 1. Add 50 µl of protease inhibitor to each ml of sample (on top of frozen plasma) in a room temperature (RT) water bath. Will use 22 mls of pooled plasma, so 1100 µl inhibitor.

[001032] 2. To remove cell/debris, spin samples at 7500 xg 20min, 4 °C in the Ultracentrifuge.

[001033] 3. Collect the supernatant, pool and measure volume & record.

[001034] 4. Add an equal volume of 2X PBS, 6mM $MgCl_2$ to the plasma.

[001035] 5. Label low-retention eppendorf tubes 1-96.

[001036] 6. Transfer 400µl of each sample to eppendorf tubes based on appropriate tube map

[001037] 7. Using an electronic P200, add competitors: 8.6 µl of 40ng/µl Salmon sperm DNA; 8.6 µl of 40ng/µl tRNA; 8.6µl of 0.5X S1.

[001038] 8. Incubate at RT with end over end rotation for 10 min.

[001039] 9. Add 10µL of appropriate oligo library, mix well. Save any leftover diluted library for gel control (see below).

[001040] 10. Incubate 1 hr at RT with end over end rotation.

[001041] 11. Using an electronic repeat P100, add 187µl of 20% PEG 8000 to sample for a final 6% concentration to the 435.5 µl of sample/oligo library. Invert a few times to mix and incubate for 15 min at 4 °C

[001042] 12. Spin each sample in table top centrifuge at 10,000 xg for 5 min.

[001043] 13. Remove supernatant and discard, add 1ml 1x PBS, 3mM $MgCl_2$ to pellet.

[001044] 14. Wash pellet by gentle inversion

[001045] 15. Remove buffer, re-suspend pellets in 100µl 1X PBS, 3mM $MgCl_2$: incubate at RT for 10 min on mixmate @ 900rpm to re-suspend. Make sure each sample is well re-suspended by pipetting.

[001046] 16. Pool all desthiobiotin library samples into one 50ml falcon tube, and the unlabeled library into another, total volume for each should be 4800µl.

[001047] 17. Take 10µL aliquot for the input into AP sample for gel (add 10 µL of 2x LDS buffer w/ 2X reducing agent.

[001048] Affinity Purification:

[001049] 18. Prepare 10µL of MyOne Strep-coated Magnetic beads per each condition into a 1.5 ml eppendorf tube and place on a magnetic bead rack. Have a Bead only control as well (n=3)

[001050] 19. Remove supernatant and wash 1X 500µl with Bead buffer.

[001051] 20. Discard supernatant

[001052] 21. Resuspend beads in an equal volume of 1X PBS, 3mM $MgCl_2$ (equal vol to what was taken out originally = 10µl)

[001053] 22. Add the 10µl of beads directly to the 4780µL from step 19. To Bead only control add PBS.

[001054] 23. Incubate samples with streptavidin beads 1hr RT on plate shaker (taped).

[001055] 24. Place on the large magnetic stand for 1 min and remove supernatant

[001056] 25. Add 1.5 mL of 1X lysis buffer to the samples (do 3 X 500µl with a good rinse of the 50mL falcon tube for each to collect all the beads) and transfer to a new set of eppendorf tubes.

[001057] 26. Incubate for 20 min on ice.

[001058] 27. Place tubes in magnetic bead rack, let equilibrate 1 min and remove the supernatant.

[001059] 28. Wash the beads with wash buffer #1 via vortexing. Resuspend well.

[001060] 29. Place tubes on magnetic bead rack, let equilibrate 1 min and remove the supernatant

[001061] 30. Wash 2 additional times as with wash buffer # 1 steps 27-29 (total 3 washes with wash buffer #1)

[001062] 31. Repeat steps 27-29 (2) additional times with wash buffer #2

[001063] 32. During the last wash transfer beads to a new eppendorf tube. (to reduce non-specific binding)

[001064] 33. Do one dry spin to make sure all residual wash buffer is removed.

[001065] 34. Add 10µl of Biotin Elution buffer 1 to beads

[001066] 35. Incubate for 15 minutes at 37°C.

[001067] 36. Place on magnetic stand for 1 min, collect sup and transfer to a new tube, add 10µL of 2X LDS, 2X Reducing agent to eluted sample. Save as Elution #1.

[001068] 37. Add 10µl of 1X LDS Sample Buffer, 1X Reducing buffer to magnetic beads.

[001069] 38. Boil the samples for 15 min at 90 °C. The boiling time is 15 minutes to essure the streptavidin on the beads unfolds and releases the biotinylated aptapmer-protein complex.

[001070] 39. Place samples on magnetic stand on ice and collect the eluted sample. This is Elution #2. Discard the beads.

[001071] 40. Gel 1 layout:

[001072] Lane 1: 5ng Desthiobiotin library

[001073] Lane 2: 1X LDS

[001074] Lane 3: Marker

[001075] Lane 4: Desthiobiotin Elution #1

[001076] Lane 5: Unlabeled Elution #1

[001077] Lane 6: Bead only Elution #1

[001078] Lane 7: Desthiobiotin Elution #2

[001079] Lane 8: Unlabeled Elution #2

[001080] Lane 9: Bead only Elution #2

[001081] Lane 10: Input for AP (saved from step 17)

[001082] Running Reducing SDS gel:

[001083] Prepare 1X MOPS SDS Running Buffer from 20X MOPS SDS Buffer

[001084] Use 10 or 12 well 4 -12 % Bis Tris gel

[001085] Peel off tape seal and place in the gel box. Insert spacer for second gel cassette if needed

[001086] Fill the inside/upper chamber with running buffer MOPS (1X) and 500ul Antioxidant

[001087] Remove the comb carefully, not disturbing the wells

[001088] Rinse the wells with the running buffer to remove the storage buffer which can interfere with sample running

[001089] Slowly load samples to each well carefully using L-20 tip

[001090] Fill the outer/lower chamber with approximately 600ml of running buffer MOPS (1X)

[001091] Place top portion of unit and secure correct electrodes

[001092] Run the gel to migrate proteins

[001093] 100 V constant for samples to move through stack (until all samples line up) for 15 min

[001094] Increase to 150 V constant for running (until visible sample buffer comes to bottom) for ~1 hr

[001095] At the end of the run, stop the power supply and remove the gel cassettes from cell

[001096] Disassemble the gel cassette by with gel knife.

[001097] Remove one side of cassette case. Trim off the gel foot and wells (avoid drying gel).

[001098] Transfer gel into container filled with Mili Q water and perform a quick wash.

[001099] Silver staining:

[001100] Materials:

[001101] ProteoSilver TMSilver Stain Kit, Sigma Catalog No. PROT-SIL1, Lot No. SLBJ0252V

[001102] Ethanol, Fisher Scientific Catalog No. BP2818-4, Lot No. 142224

[001103] Acetic acid, Acros organics Catalog No. 14893-0025, Lot No. B0520036

[001104] Water, Sigma Catalog No. W4502, Lot No. RNBD1581

[001105] Preparation:

[001106] 1. Fixing solution. Add 50 ml of ethanol and 10 ml of acetic acid to 40 ml of ultrapure water.

[001107] 2. 30% Ethanol solution. Add 30 ml of ethanol to 70 ml of ultrapure water.

[001108] 3. Sensitizer solution. Add 1 ml of ProteoSilver Sensitizer to 99 ml of ultrapure water.The prepared solution should be used within 2 hours. A precipitate may form in the ProteoSilver

Sensitizer. This precipitate will not affect the performance of the solution. Simply allow the precipitate to settle and remove 1 ml of the supernatant.

[001109] 4. Silver solution. Add 1 ml of ProteoSilver Silver Solution to 99 ml of ultrapure water. The prepared solution should be used within 2 hours.

[001110] 5. Developer solution. Add 5 ml ProteoSilver Developer 1 and 0.1 ml ProteoSilver Developer 2 to 95 ml of ultrapure water. The developer solution should be prepared immediately (<20 minutes) before use.

[001111] 6. All steps should be carried out in the hood and waste needs to be collected in toxic designated container.

[001112] Procedure

[001113] A. Direct Silver Staining

[001114] • All steps are carried out at room temperature on an orbital shaker at 60 to 70 rpm.

[001115] 1. Fixing - After electrophoresis of the proteins in the mini polyacrylamide gel, place the gel into a clean tray with 100 ml of the Fixing solution overnight in the hood. Cover tightly.

[001116] 2. Ethanol wash - Decant the Fixing solution and wash the gel for 10 minutes with 100 ml of the 30% Ethanol solution.

[001117] 3. Water wash – Decant the 30% Ethanol solution and wash the gel for 10 minutes with 200 ml of ultrapure water.

[001118] 4. Sensitization – Decant the water and incubate the gel for 10 minutes with 100 ml of the Sensitizer solution.

[001119] 5. Water wash – Decant the Sensitizer solution and wash the gel twice, each time for 10 minutes with 200 ml of ultrapure water.

[001120] 7. Silver equilibration – Decant the water and equilibrate the gel for 10 minutes with 100 ml of the Silver solution.

[001121] 8. Water wash – Decant the Silver solution and wash the gel for 1 to 1.5 minutes with 200 ml of ultrapure water.

[001122] 9. Gel development –Decant the water and develop the gel with 100 ml of the Developer solution. Development times of 3 to 7 minutes are sufficient to produce the desired staining intensity for most gels. Development times as long as 10 to 12 minutes may be required to detect bands or spots with very low protein concentrations (0.1 ng/mm2).

[001123] 10. Stop - Add 5 ml of the ProteoSilver Stop Solution to the developer solution to stop the developing reaction and incubate for 5 minutes. Bubbles of CO_2 gas will form in the mixture.

[001124] 11. Storage – Decant the Developer/Stop solution and wash the gel for 15 minutes with 200 ml of ultrapure water. Store the gel in fresh, ultrapure water and take picture for documentation.

[001125] Protein identification

[001126] Protein bands of interest were excised from the gradient gels and subjected to liquid chromatography-tandem mass spectrometry (LC-MS/MS) as above.

Example 26: Oligonucleotide probes: breast cancer versus non-cancer

[001127] This Example presents breast cancer oligonucleotide probes identified in a library enriched against balanced fractions pool of plasma-derived microvesicles from patients with 50% aggressive cancer. General methodology is as presented in **Example 21** above. The samples comprised pools of 30 each of breast cancer patient plasma and healthy plasma (i.e., non-cancer controls). A set of cancer specific aptamers was identified where each aptamer has a fold change exceeding two when compared to either the healthy plasma pool (normal and non-cancer) or process control (no negative selection enriched library).

[001128] Methodology

[001129] The F-TRin-35n-B 8-3s library as described herein was enriched against microvesicles from the plasma samples. See **Example 21** with the modifications noted below. The screened library comprised a 5' region (5' CTAGCATGACTGCAGTACGT 3' (SEQ ID NO. 131)) followed by the random naïve aptamer sequences and a 3' region (5' CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)).In the previous enrichment protocol in **Example 21**, positive, negative and positive selections were performed before each cycle of PCR to re-amplify the library. But in the current enrichment protocol, the aptamer library was purified with streptavidin beads after each selection and the beads were directly used for PCR amplification. Also as compared to prior experiments, the new sample pool for aptamer enrichment was balanced for different plasma fractions and collection vial for each of cancer, non-cancer and normal patients. Specifically, plasma was initially collected in 4 tubes from each patient, each of those 4 tubes were split into 3 aliquots, resulting in 12 aliquots from each patient. For example, a 1st tube out of the 4 results in aliquots 1A, 1B and 1C, the same split is repeated for the other three aliquots, which results in 12 aliquots from each patient (1A-C, 2A-C, 3A-C, 4A-C). Correspondingly, a pool of 60 patients consists of each variant repeated 5 times (5x12). Enrichment was done according to the selection methodology outlined above, where the library was PCR amplified after each binding round for 7 rounds, which include 3 positive selections against cancer-derived samples, 3 negative selections against controls, and a final positive selection. The enriched library was subjected to the probing test on cancer and healthy pools of plasma samples. There was a subset of 296 cancer specific aptamers, which have a relatively higher read count and fold change > 2 as compared to healthy pool and process control. The detailed protocol is as follows:

[001130] Equipment & Supplies

[001131] 1xPBS (HiClone): SH30256.01, Lot #: AZC186921, bottle#1476, exp. 05/2016. Supplemented with 3mM $MgCl_2$ (4227844, USB) in steps 6, 7, 11.

[001132] Table Top centrifuge: 0363

[001133] 20X S1: Aptamer Science TT 070214, LN: 14F-01-S1, exp. 2015-06

[001134] PEG8000- lot #SLBJ9928V cat # 91458, protease inhibitor Ref - 05892791, Water - ref 10977-015, lot # 1606173

[001135] 2X PBS+ 6mM $MgCl_2$ - PBS (Sigma) -SLBK2636V, Water - RNBD2918, $MgCl_2$ - 4227844 (USB). Used in steps: 5, 9.

[001136] Stock yeast tRNA (Ambion) - lot # 1406019. Salmon DNA (Invitrogen)- lot # 1617974

[001137] Starting solution comprises 5 ng Non-Enriched F-Trin-35n-B aptamer library, 300 ul of plasma, 0.01x S1 + 0.8ng/ul Salmon DNA/tRNA (competitor DNAs), 6% PEG8000 (to precipitate microvesicles); final volume 600 ul.

[001138] Round 1 (1st positive enrichment)

[001139] Step 1: Pre-chill tabletop centrifuge at 4°C.

[001140] Step 2: Protease inhibition: dissolve 1 tablet of "cOmplete ULTRA MINI EDTA-free EASYpack" protease inhibitor in 550 ul of H2O (20x stock of protease inhibitor).

[001141] Step 3: Add 50 ul of protease inhibitor to the sample (on top of frozen plasma) and start thawing: 1 ml total ea.

[001142] Step 4: Cell spin: To remove cells/debris, spin samples at 10,000 x g, 20 min, 4°C. Collect the entire volume of supernatant (SN) without disturbing the pellet.

[001143] Step 5: Mix SN from step 4 with equal volume of 2xPBS 6 mM $MgCl_2$, collect 600 ul ea into 2ml Fisher Low binding tubes for use in step 6. Store remaining sample at 4°C for the following rounds.

[001144] Step 6: Blocking: Add competitors in order: 1) Salmon DNA Stock; 2) tRNA; 3) S1. Make dilutions to desired concentration (see starting solution above) in 1xPBS, 3mM MgCl2, mix well. Incubate for 10 min at room temperature (RT), end-over-end rotation.

[001145] Step 7: Binding: Add ssDNA Probing library to the final concentration 12.5 pg/ul for binding. Make dilutions in 1xPBS, 3mM $MgCl_2$.

[001146] Step 8: Precipitation: Add buffer (20% PEG8000 in 1x PBS with 3mM $MgCl_2$) to sample to the final PEG concentration 6%.

[001147] Step 9: Spin at 10,000 x g for 5 min, RT.

[001148] Step 10: Wash: Remove SN, add 1ml 1xPBS, 3 mM MgCl2 and wash pellet by gentle invertion with 1ml aptamer buffer.

[001149] Step 11: Resuspention: Remove buffer, Re-suspend pellets in 200 ul H2O: incubate at RT for 10 min on mixmate 900rpm. Ensure each sample is re-suspended by pipeting after step 11.

[001150] Step 12: Purification: Aptamers elution from PEG/Protein pellet with Streptavidin beads (Dynabeads #65001: Dynabeads® MyOne™ Streptavidin C1):

[001151] -> Beads stock: 10 mg/ml; Capacity: >2,500 pmoles/mg;

[001152] -> 10 ul beads should bind 250 pmoles Biotin

[001153] -> 3 ul beads for each aptamer library (AL) sample can bind 75 pmoles Biotin

[001154] Library input in the protocol above is ~5ng, which is 0.17 pmol

[001155] 12.1) Pre-washing Streptavidin Magnetic Beads:

[001156] 12.1.1 Add 10 uL of Streptavidin Magnetic Beads into 1.5mL microcentrifuge tube (3ul x 2 samples + overage)

[001157] 12.1.2 Place the tube into a magnetic stand to collect the beads against the side of the tube. Remove and discard the supernatant.

[001158] 12.1.3 Add 0.5mL of Wash Buffer (1xPBS with 0.1% Tween 20) to the tube. Invert the tube several times or vortex gently to mix. Collect the beads with a magnetic stand, then remove and discard the supernatant.

[001159] 12.1.4 Wash once with 0.5ml 1xPBS, collect the beads with a magnetic stand, then remove and discard the supernatant.

[001160] 12.1.5 Add 35 ul 1xPBS to tubes. Aliquot into 10ul per well (using repeater pipet) in 1.5 ml Fisher low-binding tubes. Add 40 ul of 1xPBS.

[001161] 12.2) Incubate 100 ul of sample, recovered in step 12, at 50°C for 10 min (mixmate, 500 rpm) (to denature/remove protein/PEG from aptamer library)

[001162] 12.3) Sample Binding:

[001163] 12.3.6 Add heat denatured samples to beads (aliquoted in step 5).

[001164] 12.3.7 Incubate for 30min, 37°C, mixmate at 800 rpm. Spin down at 2000rpm for 20 sec.

[001165] 12.3.8 Collect the beads (with bound aptamers) using magnet and remove the supernatant by multichanel pipet.

[001166] 12.3.9 Take tubes off the magnet, add 300ul of 1xPBS, pipet well. Collect the beads with magnetic stand, then remove supernatant.

[001167] 12.3.10 Add 100ul H2O to resuspend beads by multichanel pipet.

[001168] 12.3.11 Block the beads with Biotin: add 7.5 ul from 10uM stock, incubate 15 min, RT, 800rpm.

[001169] 12.3.12 Collect the beads (with bound aptamers) using magnet and remove the supernatant by multichanel pipet. Take tubes off the magnet, add 200ul of 1xPBS, pipet well. Collect the beads with magnetic stand, then remove supernatant.

[001170] 12.3.13 Add 100ul H2O

[001171] Step 13: Use beads with bound aptamer library directly in re-AMP PCR (3x33ul)

[001172] Step 14: Agarose gel

[001173] SYBR Gold gel lot #: H204044-01

[001174] 2ul of sample + 8ul of loading buffer. Run for 10 cycles

[001175] If no dsDNA bands appear, run additional 3 cycles

[001176] Optional: if PCR product has non-specific bands, perform gel cut.

[001177] Step 15: dsDNA purification with Nucleospin column (NTI binding buffer):

[001178] 2x volume of buffer NTI per sample volume (600ul NTI to 300ul sample)

[001179] - combine 3 wells per column

[001180] - 5 min elution in 30 ul NE buffer, RT. Add 20 ul of NE after elution.

[001181] Step 16: Optional: Gel, SybrGold, Agarose: to verify product with correct size is observed

[001182] Step 17: Quantify dsDNA (QuBit, Life Technologies) (keep 5 ul of dsDNA as control for gel later)

[001183] Step 18: Lambda digestion at 37C for 2 h; heat inactivation at 80C for 10 min.

[001184] Step 19: ssDNA purification with Nucleospin column (NTC binding buffer).

[001185] - combine 3 samples per column

[001186] - elution: 5 miin in 30 ul NE buffer, RT

[001187] Step 20: Quantify dsDNA (QuBit, Life Technologies) ssDNA (5 ul)

[001188] Step 21: Gel ssDNA: load between 2ng and 10 ng

[001189] Round 2 (2d positive enrichment)

[001190] Repeat protocol starting from step 6 above, using diluted plasma samples stored in step 5. All steps are the same except for step 7. Input of library is 5 times less

[001191] Step 7: Binding: Add ssDNA Probing library to the final concentration 2.5 pg/ul for binding. Make dilutions in 1xPBS, 3mM MgCl2.

[001192] Steps 6-13: As above. Use entire 100 ul of re-suspended sample to re-amplify via PCR

[001193] Steps 14-20: ssDNA preparation as above.

[001194] Round 3 (3rd positive enrichment)

[001195] Repeat steps as shown for round 2

[001196] Round 4 (Negative enrichment)

[001197] Make sure to use correct negative samples (e.g., non-cancer sample in the case of positive cancer selection above, or vice versa)

[001198] Steps 1-9: Repeat as in Round 2 above (ssDNA input is 2.5pg/ul).

[001199] Step 10: Collect SN (~900ul) and discard pellet after PEG precipitation.

[001200] Step 11: Resuspention: Not needed in this step

[001201] Step 12: Purification: Aptamers elution from PEG/Protein pellet with Streptavidin beads (Dynabeads #65001: Dynabeads® MyOne™ Streptavidin C1):

[001202] -> Beads stock: 10 mg/ml; Capacity: >2,500 pmoles/mg;

[001203] -> 10 ul beads should bind 250 pmoles Biotin

[001204] -> 3 ul beads for each AL sample can bind 75 pmoles Biotin

[001205] Library input in the protocol above is ~5ng, which is 0.17 pmol

[001206] CHANGE FOR SN -> For each sample, split 900ul SN collected in step 10 above into two aliquots 450ul ea, consider 3ul of beads per each aliquot.

[001207] 12.1) Pre-washing Streptavidin Magnetic Beads:

[001208] 12.1.1 Add 30 uL of Streptavidin Magnetic Beads into 1.5mL microcentrifuge tube. Four samples make 8 aliquots (for beads treatment) x 3ul beads + overage.

[001209] 12.1.2 Place the tube into a magnetic stand to collect the beads against the side of the tube. Remove and discard the supernatant.

[001210] 12.1.3 Add 0.5mL of Wash Buffer (1xPBS with 0.1% Tween 20) to the tube. Invert the tube several times or vortex gently to mix. Collect the beads with a magnetic stand, then remove and discard the supernatant.

[001211] 12.1.4 Wash 1 time with 0.5ml 1xPBS, collect the beads with a magnetic stand, then remove and discard the supernatant.

[001212] 12.1.5 Add 105 ul 1xPBS to tubes. Aliquot into 10ul per well (using repeater pipet) in 1.5 ml Fisher low-binding tubes. Add 40 ul of 1xPBS.

[001213] 12.2) Incubate 900 ul of sample, recovered in step 10, at 50°C for 10 min (mixmate, 500 rpm) (to denature/remove protein/PEG from aptamer library)

[001214] 12.3) Sample Binding:

[001215] 12.3.6 Add 1/2 (450ul) heat denatured samples to beads (aliquoted in step 5).

[001216] 12.3.7 Incubate for 30min, 37°C, end-over-end rotation. Spin down at 2000rpm for 20 sec.

[001217] 12.3.8 Collect the beads (with bound aptamers) using magnet and remove the supernatant by multichanel pipet.

[001218] 12.3.9 Take tubes off the magnet, add 300ul of 1xPBS, pipet well. Collect the beads with magnetic stand, then remove supernatant.

[001219] 12.3.10 Add 100ul H2O to resuspend beads by multichanel pipet.

[001220] 12.3.11 Block the beads with Biotin: add 7.5 ul from 10uM stock, incubate 15 min, RT, 800rpm.

[001221] 12.3.12 Collect the beads (with bound aptamers) using magnet and remove the supernatant by multichanel pipet. Take tubes off the magnet, add 200ul of 1xPBS, pipet well. Collect the beads with magnetic stand, then remove supernatant.

[001222] 12.3.13 Add 100ul H2O per purification sample (total 200ul per negative enrichment sample)

[001223] 12.3.14 Split each sample into 3 aliquots (33ul each), this results into 6 wells per enrichment sample.

[001224] Step 13: Perform PCR

[001225] Rounds 5, 6: same as round 4

[001226] Round 7: repeats steps as shown for round 2

[001227] Results

[001228] **FIGs. 14A-B** are scatter plots showing high correlation of frequencies between replicates of cancer pool (**FIG. 14A**) and appearance of cancer specific subset of aptamers when cancer and healthy

pools are compared (circled region in **FIG. 14B**). **FIGs. 14C-F** are scatter plots showing the appearance of cancer specific subset of aptamers when the cancer pool profile of the enriched library (i.e., C-RN7) was compared to the cancer pool of the process control library (CP; **FIG. 14C**) and healthy pool profile of process control library (HP; **FIG. 14D**). We did not observe a similar subset when the healthy pool profile (C-RN7 v HP) of the enriched library was compared to both variants of profile of process control (**FIGs. 14E-F**, see points along x-axis).

[001229] Selections of aptamer sequences identified in the enrichment screening are shown in **Tables 27-29**. In these tables, each complete aptamer sequence is assembled from 5' to 3' as a 5' region, the variable region, and 3' region. The sequences are shown 5' to 3' from left to right, wherein each complete sequence consists of a 5' leader sequence 5'- CTAGCATGACTGCAGTACGT (SEQ ID NO. 131) followed by the indicated Variable Region sequence followed by the 3' tail sequence 5'- CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132).The top 50 cancer specific sequences are shown in **Table 27**, each having a relatively high copy number in these experiments and a cancer/normal copy number ratio of at least 2-fold higher in the cancers. **Table 28** comprises 25 sequences with mid-level copy number and a cancer/normal copy number difference of 0.8-1.2-fold. **Table 29** shows a selection of 25 more sequences with low copy numbers and cancer/normal copy number ratio of ~1.0 (i.e., no significant difference).

Table 27: Cancer-Specific Sequences

SEQ ID NO.	Variable Region
137	AACCCGCGTGATTGGGGTACTGCCCATGCGACTTT
138	AACTGAAGGCCAATTAAACTCCACTGTGCTAATCT
139	ACATGCCGGCGTTGATACTCTACTGTGCTCCATTATGA
140	AGACATGCCACGCCCTTTACATACTCCACTGTGCCA
141	AGCCACGCAGACCTATCTCTACTGTGCCAATGTT
142	AGGTACGTAAGGCTGCCATTACTCTCTACTGTGCCGA
143	AGTCTAGCGATCGACTTTTCTCTACGGTGCCCTTC
144	AGTGTGCGCATGAATCCATTACTCCACTGTGCCTTGA
145	AGTTGCAGACCCTTGTATTCTCTACCGTGCCCTATGA
146	CACTGCTAACTGATATTGTTGAACTCCACCGTGC
147	CATGCGACAAGCTTCCATTAACCATCCGCTATTTATGA
148	CATGCGGACCACGCTTTCACAAACTCATGATTCAAA
149	CATGCGTACGTCCTATTCTCTCTACTGTGCATTAA
150	CCTGCAATGCGGAAACCCTTCTACTCTACTGTGCC
151	CGCGAGCCAGGGTACTGCCTGCGGTCTAAACAATCA
152	CGTGCGACTTGTTCTCTACTGTGCCAATAAACCATA
153	GATAGCGCATATTCCTCTACTGTGCTACTTGTTAT
154	GCAAACCAATTAGTGTACCTCTACCGTGCCCAATGA
155	GCAACTACTCTCTACCGTGCCCAGTGATTTCTCCATGA
156	GCACCACATGTCAATCTCTACTGTGCTTATCTCATA
157	GCAGCACAAAAGTTACTCTCCTCTACCGTGCCCCTA
158	GCAGTTTGCGTGGATACTCTACTGTGCTGACCCTTA
159	GCATACATGAACTCTACCGTGCCCGTTGACTTGAAA
160	GCATAGCGGTTCACATTATTTTACTCTACCGTGCC
161	GCCACAGTATCGTTCTCGAGCTGGTTCCCGCCAT
162	GCGAACATCACTCTACCGTGCCATCTGAAATGACGTGA
163	GCGAACTTGTAACTCTACTGTGCTTATATTAAGGTTGA

164	GCGACCAATACTCTACTGTGCTGATACGGATTTTA
165	GCGACCACTCTATAACCTCAACCGTGCTCACTCCT
166	GCGCAAGTTGTCTCTACCGTGCCCCGAAAGTTTTGGA
167	GCGCACCACCTACATTGTCTCCACCGTGCTTTATT
168	GCGCACCCACGTGAACTCTACCGTGCCTATTTCCTA
169	GCGCACCGACCTCTACCGTGCCAAAATAGGTTATCTGA
170	GCGCCACCACACAAACCTTCCCTATCGAGGGAGAT
171	GCGCGGACTATATTACTCTACTGTGCCCTGCTTATGA
172	GCGGATAGGACAATAACTTACTCCACTGTGCCATC
173	GCGGTGACCTCTACTGTGCGCCCCAAGCCTTAGTT
174	GCGTACAGAACCTCTACCGTGCCCACTCACTTCCATGA
175	GCGTCCCCTCCCGGATGGTCCTTTCTCTACTGTGC
176	GCGTGCTAACGGTATGCAAGACGTATGCGATTTTC
177	GCTAGACCGATCCACCTCAAACCTCTACTGTGCCT
178	GCTAGCTTAGCTCTACCGTGCACATTCCGCTATTT
179	GGTGCAGGCAAGATATTTTACTCTACTGTGCATTT
180	GTAAATGTACATGCGTATCCTCACCTCTACTGTGC
181	TAACACGTCTTTCACTCTACTGTGCCCTTTATGCC
182	TAATGGCATGCGGACCTATCCTCTACCGTGCTCCTTGA
183	TATGCGATTTCTCTACCGTGCCAATATGCCTTGTT
184	TCTGCGATTCTACCGTTACTCTACCGTGCCACCAAA
185	TGCGCAGTCATTTCGCCATGTTCTCTACCGTGCCAA
186	TGTCAGGCGGTAGTACTCTCCACCGTGCCTATTGTTGA

Table 28: Moderate Control Sequences

SEQ ID NO.	Variable Region
187	ACATGCATACCCTACGAATCGTTCCATCACCATAC
188	AGTCAGTGCGCCCGCTATTTACGATCTCACTGTTC
189	AGTGCAGTCGCGATGGGAACTTCTTCTTTGCTTTA
190	ATGGCCATGCGAACCGAAACCTAGCCCATTTTCCTA
191	ATTGCGGTCATCCCCTTCCACGCTATACCACCAT
192	CACTGCAGGTCACTGGCGCCCTATTTCCTCACATT
193	CATGCCCGTAAACGCCTAATCTACCACCTTTTGCT
194	CATGCCGCCTTTCGACTTCCATATCCCAACACGCC
195	CATGCCGGTCATTCCATACTCAAACTCATTCACTC
196	CATGCGAGCTACCCTTATCATCTCATGTTTGCTTT
197	CATGCGATTGCGCCCCATTCCTTGCCTTATCTCAC
198	CATGCGCACATCGTACTCCGTAGCCCTAAATTATCA
199	CATGCGCCCTGTGCGTTCTCTACTACCCTAATCAT
200	CATGTACGCACACCCACTGATTCTTTACCACCCACTGA
201	GCAGGACACCCCCTCATTATTTGTTTCATCCTACG
202	GCATCAGGCGCACTTTCATTTGCTAATCGTTTTTT
203	GCCATGCGAGAGGGTTATTGTTGATCTCATGGGTT
204	GCGATAAAGCCCATGTAACCCTCTGAAAACCCGAT
205	GCGCCACGAGCTACCCGACTTCCGATTACTTTCTT
206	GCGGACATCGGGCACTTTATACAACCACCTTTTTG
207	GCTGCAGGCCGTCCCAAATATTCGCTCCCACACATA
208	GCTGGTCAGGCGTTCCACACTTCCTACCCGCTTTT
209	TATGCGCAGGACCACCTATTACACGACCTTATCCT
210	TCAACTGGGGTCAGTGCCAACGCTTACTTTCTTCT
211	TCAAGCAGTCATAGCCGGTTTCACGCTTTACTTAC

Table 29: Negative Control Sequences

SEQ ID NO.	Variable Region
212	AAAACACGCAAACTACCTGGAAACTTGACTTCTTT
213	AAAACACTCACGATACCTACCCTGGTCTTCACAAC
214	AAAACAGCAGGCATTTTCCTGTACTTTCGAATTCA
215	AAAACCACAGCTACCATTATCGCATTGGCCCTACT
216	AAAACCCAAGCACCGAACGAACTATCCCTTTTTCT
217	AAAACCCACCACGCGTCGTACGCATACCACCCTTT
218	AAAACCGGTCCTGCGCCTTCTTCTCCGCTTTTATT
219	CAGGTACTGTGGCAGGCGATCCATTGCTCTCTTTT
220	CAGGTCAGCAGTCCAGGCGATTACTTTCTCTTTTC
221	CAGTAACCGCTGTCTGTCATTTTTCAACATTCTGT
222	CAGTAACTGTAGTCGTCGAACCTTGCTTACAACCCA
223	CAGTAAGGAAGCTCATGCGGTACAGGTCTACCTCG
224	CAGTAAGTGCTGTCACTGCGCCACTTGTAAATACTTGA
225	CAGTAATGCGAAGCTAGTACGTTCTCTCTTATTTTA
226	CAGTACAAACATCGCGATTCTTTCCTTGATACTGT
227	GAGGGCGGCTCTAATGCATGCGCTTTCATCTTTGC
228	GAGGGCGGGCAGTCATTTCGATTATATCTGCCTACA
229	GAGGGCGGTCCAACGCGCAATGTCTTTTTTCTTCCGA
230	GAGGGCGTACAGGAGTCTTGCGGTCCACATTTTAT
231	GAGGGCTACTGAAGTTAATGGCATTCTTCTCTATC
232	GAGGGCTCATGCCAGCCGCTTTTCTTACCTTTATC
233	GAGGGCTGAACAAGTAGTGCGTACATTTTCCTTCCTGA
234	GAGGGCTGCAGGCATCCAATTTCCTCCTTGTCCCT
235	GAGGGGCCGTCCTTCGTACGGAATGTGCATGCGCT
236	GAGGGGCGGTCATTGCCGTTAGCCTCCTTTCGCCT

[001230] The data described above (i.e., **FIGs. 15A-F** and **Tables 27-29)** were obtained with 1 ng input of the specified oligonucleotide libraries used to probe the various samples. In addition, probing experiments were performed with lower titers of the C-R7N oligonucleotide library, specifically 0.1, 0.01, 0.001 and 0.0001 ng. A total of 826 unique oligonucleotide sequences were detected between all titers. An eighth enrichment round (round 8) was performed with four titers of the library C-R7N-1 (i.e., 1 ng input in round 7): 1, 0.1, 0.01 and 0.001 ng input. From each titer, sets of oligonucleotides that passed the same filters described above were obtained. The composite set of oligonucleotides was compared to the sets of oligonucleotides obtained from the titrations of the C-R7N library noted above. The variable regions of 733 oligonucleotides observed in at least two of the five sets of oligonucleotides are listed in rank order by occurrence in SEQ ID NOs. 237-969. These oligonucleotides were identified as oligonucleotide probes that selectively bind to cancer samples. As in **Table 27**, the oligonucleotides were synthesized with a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)) flanking the variable regions. The 733 cancer specific sequences which can be used to identify cancer samples along with those in **Table 27** above.

[001231] As noted, next generation sequencing technologies can be used to identify and quantify members of the oligonucleotide library that bind to a sample. We also designed probes that can be used to

identify and quantify members of the oligonucleotide library that bind to a sample without requiring a PCR step. This procedure relies on hybridization of the bound oligonucleotide with oligonucleotide-specific complementary probe. The complementary probe can directly or indirectly carry a tag that can be detected. For example, the oligonucleotide-specific complementary probe may carry a fluorescent tag. A general design is shown in **FIG. 14G**. In the figure, an individual member of the oligonucleotide library **1401** is bound by complementary probe **1402**. Complementary probe **1402** consists of three sections from 5' to 3': 1) probe part B **1403**; 2) probe complement **1404**; and 3) probe part A **1405**. The member of the oligonucleotide library **1401** is bound by complementary probe **1402** via base pair hybridization between the variable region of oligonucleotide library **1401** and the probe complement region **1404** of complementary probe **1402**. Tag probe **1406** which is fluorescently labeled (indicated by the circles in the figure) and capture probe **1407** hydridize to the probe part A **1405** and probe part B **1403** of complementary probe **1402**, respectively. These features allow for the capture of the entire complex (i.e., oligonucleotide library **1401**, complementary probe **1402**, tag probe **1406**, and capture probe **1407**) via the capture probe, combined with detection via the label of the tag probe **1406**.

[001232] For these experiments, the complementary probes were designed to work with tag probe and capture probe supplied as part of the nCounter nucleic acid detection system from NanoString Technologies, Inc. (Seattle, WA). The complementary probes for nCounter quantification were designed as reverse complement to the top 50 oligonucletide probe binders and 50 negative controls, described above. The complete sequences are shown in **Table 30**. In the table, the ID column is an identifier and the Type column indicates if the sequence is to a control oligonucleotide probe ("neutral") or to a cancer binding oligonucleotide probe ("binder"). Each complete sequence comprises from 5' to 3': 1) 5' Region (Probe Part B **1403**); 2) Probe Complement Region **1404**; and 3) 3' Region (Probe Part A **1405**), where region indentification numbers correspond to **FIG. 14G**. The probe complement region may be shorter than the variable sequence of the corresponding oligonucleotide probe in order to optimize binding parameters (e.g., specificity and melting point). The probe complement region may also be so optimized to bind against the variable sequence of the corresponding oligonucleotide probe and also to at least a portion of its 5' and 3' flanking sequences (e.g., as shown in **Tables 27-29** above). The oligonucleotides each have a 5' region **1403** consisting of the sequence (5′ – CGAAAGCCATGACCTCCGATCACTC (SEQ ID NO. 970)) 5' of the Probe Complement Region **1404** and a 3' region **1405** as shown in **Table 30** located 3' of the Probe Complement Region **1404**. Probe Part A **1405** shown in **FIG. 14G** corresponds to SEQ ID NO. 971.

Table 30: Complementary Probe Sequences

SEQ ID NO	Type	Probe Complement Region 1504	3' Region (Probe Part A 1505)
972	neutral	CAAGTTTCCAGGTAGTTTGCGTGTTTTA CGTAC	CCTCAAGACCTAAGCGACAGCGTG ACCTTGTTTCA
973	neutral	CAGGGTAGGTATCGTGAGTGTTTTACGT ACT	CATCCTCTTCTTTTCTTGGTGTTG AGAAGATGCTC
974	neutral	GAAAGTACAGGAAAATGCCTGCTGTTTT ACGTAC	CACAATTCTGCGGGTTAGCAGGAA GGTTAGGGAAC

975	neutral	GAGACAGAGTAGGGCCAATGCGATAATG	CTGTTGAGATTATTGAGCTTCATC ATGACCAGAAG
976	neutral	GAGACAGAGAAAAAGGGATAGTTCGTTC GGT	CAAAGACGCCTATCTTCCAGTTTG ATCGGGAAACT
977	neutral	GACGCGTGGTGGGTTTTACGTAC	CGAACCTAACTCCTCGCTACATTC CTATTGTTTTC
978	neutral	GCGCAGGACCGGTTTTACGTAC	CCAATTTGGTTTTACTCCCCTCGA TTATGCGGAGT
979	binder	GAGACAGAAAGTCGCATGGGCAGTA	CTTTCGGGTTATATCTATCATTTA CTTGACACCCT
980	binder	AGTGGAGTTTAATTGGCCTTCAGTTACG TACT	CAACAGCCACTTTTTTTCCAAATT TTGCAAGAGCC
981	neutral	GAACGATTCGTAGGGTATGCATGTACGT ACT	CACCGTGTGGACGGCAACTCAGAG ATAACGCATAT
982	binder	AGTATCAACGCCGGCATGTACGTAC	CCTGGAGTTTATGTATTGCCAACG AGTTTGTCTTT
983	binder	TAAAGGGCGTGGCATGTCTACGTAC	CAGATAAGGTTGTTATTGTGGAGG ATGTTACTACA
984	binder	GAGACAGAACATTGGCACAGTAGAGATA GGT	CTTCCTTCCTGTGTTCCAGCTACA AACTTAGAAAC
985	binder	TAATGGCAGCCTTACGTACCTACGTACT	CATAAAATTGGTTTTGCCTTTCAG CAATTCAACTT
986	neutral	CGGGCGCACTGACTACGTAC	CTGGTCAAGACTTGCATGAGGACC CGCAAATTCCT
987	binder	GAGACAGGAAGGGCACCGTAGA	CTTTCGTTGGGACGCTTGAAGCGC AAGTAGAAAAC
988	neutral	CCATCGCGACTGCACTACGTACT	CCAGCAGACCTGCAATATCAAAGT TATAAGCGCGT
989	binder	GAGACAGTCAAGGCACAGTGGAGTAATG	CCTGCCAATGCACTCGATCTTGTC ATTTTTTTGCG
990	binder	GGTAGAGAATACAAGGGTCTGCAACTAC GTAC	CAAACTGGAGAGAGAAGTGAAGAC GATTTAACCCA
991	neutral	CGGTTCGCATGGCCATACGTAC	CGATTGCTGCATTCCGCTCAACGC TTGAGGAAGTA
992	neutral	GAAGGGGATGACCGCAATACGTACT	CTGAGGCTGTTAAAGCTGTAGCAA CTCTTCCACGA
993	neutral	CCAGTGACCTGCAGTGACGTACT	CTAGGACGCAAATCACTTGAAGAA GTGAAAGCGAG
994	binder	TGGAGTTCAACAATATCAGTTAGCAGTG ACGTAC	CCACGCGATGACGTTCGTCAAGAG TCGCATAATCT
995	neutral	CCTGCCACAGTACCTGACGTACT	CATTTGGAATGATGTGTACTGGGA ATAAGACGACG
996	neutral	CTGGACTGCTGACCTGACGTACT	CACAAGAATCCCTGCTAGCTGAAG GAGGGTCAAAC
997	neutral	GAAAAATGACAGACAGCGGTTACTGACG TAC	CTTGACGTAGATTGCTATCAGGTT ACGATGACTGC
998	neutral	AGGTTCGACGACTACAGTTACTGACGTA C	CTTACAGATCGTGTGCTCATGACT TCCACAGACGT
999	neutral	CGCATGAGCTTCCTTACTGACGTACT	CTTGGAGGAGTTGATAGTGGTAAA ACAACATTAGC
1000	neutral	GAGACAGTCAAGTATTTACAAGTGGCGC AG	CCTACGTATATATCCAAGTGGTTA TGTCCGACGGC
1001	neutral	GAGACAGTAAAATAAGAGAGAACGTACT AGCTTCGCATTA	CAGCAAGAAGGAGTATGGAACTTA TAGCAAGAGAG
1002	neutral	GGAAAGAATCGCGATGTTTGTACTGACG TAC	CACCCCTCCAAACGCATTCTTATT GGCAAATGGAA

1003	neutral	GGCGTTTACGGGCATGACGTAC	CCCGAAGCAATACTGTCGTCACTC TGTATGTCCGT
1004	neutral	GTCGAAAGGCGGCATGACGTAC	CCGGGAATCGGCATTTCGCATTCT TAGGATCTAAA
1005	neutral	GAGACAGGAGTGAATGAGTTTGAGTATG GAATGAC	CCGATCTTCATAACGGACAAACTG AACGGGCCATT
1006	binder	GTTAATGGAAGCTTGTCGCATGACGTAC T	CGCTATGCAGACGAGCTGGCAGAG GAGAGAAATCA
1007	neutral	ATGATAAGGGTAGCTCGCATGACGTACT	CATTCGCAACCATGTGAAGTAATG TGAGCGTACTT
1008	neutral	GGCGCAATCGCATGACGTACT	CACCAGTTAGCGTGGCGTATACCA TGTTGTTAACA
1009	neutral	GAGACAGTGATAATTTAGGGCTACGGAG TACG	CCTGAATCAATAGAACAATATCAG TTATGGCGGTG
1010	neutral	GAGACAGATGATTAGGGTAGTAGAGAAC GCAC	CGGTTGTTAATATGACAGGCCGCT AAAGACGTTCT
1011	binder	GAGACAGTTTGAATCATGAGTTTGTGAA AGCGT	CCGTCTCAGATGAGTGGGTTAATC AATCAAGTATG
1012	binder	GAGAGAATAGGACGTACGCATGACGTAC T	CTGACACATTAGTAACGTCGGCAA GCACTTAGTCG
1013	neutral	TGGGTGTGCGTACATGACGTACT	CGTGAACCAGATTATGTATGGACG CGCAATAGATA
1014	binder	GGTTTCCGCATTGCAGGACGTAC	CATACGAAATTTGAGCAAGCAATT GAAGGCTTAGA
1015	binder	CCCTGGCTCGCGACGTAC	CTATCAGCTAATAGGGTCGGCTCA ACAGTGTATCC
1016	binder	CAGTAGAGAACAAGTCGCACGACGTAC	CTATCAATTCGTGACCCCGATCAT CCAGTCCAGAA
1017	neutral	GAGACAGGCAAAGATGAAAGCGCATG	CTTGAGCTCTAGGCCCAAAACGAC CTTAATGGTCA
1018	neutral	TGCCCGCCCTCACGTAC	CTAGCCCAGATCCTACGAGATGAG CTACGTAACTA
1019	neutral	GTTGGACCGCCCTCACGTAC	CAAATGCACTCTATATGGAGGGAG AGTAGCTGGAT
1020	neutral	ACTCCTGTACGCCCTCACGTAC	CCTGGTCTAGGTATCTAATTCGTG GGTCGGGTACT
1021	neutral	GCCATTAACTTCAGTAGCCCTCACGTAC	CATTAGCTCGGATGCTATCAGCTT GCGCCTATTAT
1022	neutral	GAGACAGGATAAAGGTAAGAAAAGCGGC TG	CACGATCTGTATTTTGCACCTTTC GCTATGCTGAG
1023	neutral	GAGACAGTCAGGAAGGAAAATGTACGCA C	CTGTGTCCGTCTATACGCATACTG GTCCACATATA
1024	neutral	GAGACAGAGGGACAAGGAGGAAATTGG	CATGTTGGAGTTAACGGAGACCCG CCATCGTTTAC
1025	neutral	GGACGGCCCCTCACGTAC	CGCTCATTTTGAACATACGATTGC GATTACGGAAA
1026	neutral	ATGACCGCCCCTCACGTACT	CCTATGCATCATGTGCCTCACTAG GACATCATGCT
1027	binder	CACAGTAGAGGAATATGCGCTATCACGT ACT	CCTAAATTGGGAAAAAAGGTTTTA GCTATTGATGG
1028	binder	GAGACAGTCATTGGGCACGGTAGA	CTTCAGTTAAAGGCTATCTTGCTC CGCTCGTTCTC
1029	binder	GCACGGTAGAGAGTAGTTGCACGTAC	CTTAAAGCTATCCACGAATGTCAA AAATGTGGTTT
1030	binder	GAGACAGTATGAGATAAGCACAGTAGAG ATTGACATGT	CCCGAATGTATAATGCTGACGTTC TTGCTTTTGGC

1031	binder	GAGACAGTAGGGGCACGGTAGAG	CCTATTGAAGCAATCCTCTCCCCA ATACTTAAAAA
1032	neutral	GGGGGTGTCCTGCACGTAC	CTACGGTTACCGTCTTTATAAGTG AACAAAACCGG
1033	binder	GTAGAGTATCCACGCAAACTGCACGTAC	CTCTGTGAACTGTCATCGGTCCGA TCAATTAGTCT
1034	binder	GCACGGTAGAGTTCATGTATGCACGTAC	CTCCCCTTTCCCAAGTAAATGTAC GGGAATTATCG
1035	binder	GAGACAGGGCACGGTAGAGTAAAATAAT GTGA	CGCTTTATTATGTGTTCGTCTAAC TCTGTTTCTGT
1036	neutral	AAAGTGCGCCTGATGCACGTAC	CCGAGTGCATGAGCTGTCTTTCAC ATGATACATCG
1037	binder	CTCGAGAACGATACTGTGGCACGTAC	CTATTTCTGTTCACGGATGAAGGC CTATATCAATG
1038	neutral	CCCTCTCGCATGGCACGTAC	CCATCCACTTTCATGGAAACAATA AGAGCAGGGAA
1039	binder	GAGACAGTCACGTCATTTCAGATGGCA	CACAAACTCACTACTACCAACAAC CTCACCAAAAA
1040	binder	GAGACAGTCAACCTTAATATAAGCACAG TAGAGTTACAAGTTC	CTCATGTCCTCTGTTAATCCAGCC TGAATATGCCA
1041	binder	GCACAGTAGAGTATTGGTCGCACGTAC	CAGAAATGTCACTCCCATGGTGGC TGATATAGAAA
1042	binder	GAGACAGAGGAGTGAGCACGGTT	CATGTCGAACCTTGGATAGGAGCG ACCGATTACGT
1043	neutral	GGTTACATGGGCTTTATCGCACGTACT	CTCAGGTTGTTACTTGAAGGGTTC AACACGAGCTC
1044	binder	GGTAGAGACAACTTGCGCACGTACT	CAGAAGATCAAAAAACGATCCCTG TCCATCAATAC
1045	binder	GAGACAGAATAAAGCACGGTGGAGACAA TG	CTTAGGCTACCAAATGAATTTAAA GCCAGCTGAAA
1046	binder	GAGACAGTAGGAAATAGGCACGGTAGAG TT	CCAATGCTTGCAGTATGTATCCTG ATCGTGCGTGC
1047	binder	AGGTCGGTGCGCACGTAC	CCTGCATTCTCATGGAAATGCAAT GGATTCATTCC
1048	binder	GTGTGGTGGCGCACGTACT	CCTGTTGCAGTATCACGTAAATAC CTACTTCGATA
1049	neutral	GAGACAGAAGAAAGTAATCGGAAGTCGG GTAG	CTAGCTGTTATGGCTATTGCTGAA ACAGCAAAATT
1050	binder	ACAGTAGAGTAATATAGTCCGCGCACGT AC	CCTTACGACTTCACTGCAATTGAC GATTCAGTTAA
1051	neutral	GAGACAGCAAAAAGGTGGTTGTATAAAG TGCC	CCTCATACCAATGTAAAGTATAGT TAACGCCCTGT
1052	binder	GAGACAGGATGGCACAGTGGAGTAAG	CATCTCCATGACTGCTTGAGCGGC TGGAGAATCTG
1053	binder	ACAGTAGAGGTCACCGCACGTAC	CTTTCGCCACCCATATAAACCCCA CTTCGTCCTCA
1054	binder	CGGTAGAGGTTCTGTACGCACGTAC	CAAGGCAGAGCAAATGTGACACTG TCTATCAGTAC
1055	binder	GGGAGGGGACGCACGTAC	CCTACATATATAGGAAAAGGGAAG GTAGAAGAGCT
1056	binder	GAGACAGGAAAATCGCATACGTCTTGCA TAC	CCTTCTGGAATTTCTTCCTTTGAT TTTGCCATTTT
1057	binder	GAGGTGGATCGGTCTAGCACGTAC	CTCCTAAGGTTGCTGATTTGGTTG TTGGAGACCCA
1058	binder	ACGGTAGAGCTAAGCTAGCACGTACT	CAAGGCCTAGCCTAAAGGTTCTTG CAGAGCAACAT

1059	neutral	CGGCCTGCAGCACGTACT	CCTAATTAGCTCTAGGAAACACAACCCCGGGATTT
1060	neutral	GAACGCCTGACCAGCACGTAC	CTTGAGTTATACGGAACTTCGCAAAAGTATTCCCT
1061	binder	GAGACAGAAATGCACAGTAGAGTAAAATATCTTGCCTG	CCATCCATCAACAACTGCTCCAACAGCCTTTCCAT
1062	binder	GAGACAGGCACAGTAGAGGTGAGGATA	CCGGCACAAGCAGACAAAATCAACATGGTCATTTA
1063	binder	GAGACAGGGCATAAAGGGCACAGTA	CGCTCACGTGATCTACCCTAGCTGACCGCTAATGA
1064	binder	GAGACAGTCAAGGAGCACGGTAGAG	CTCTAACCTGCATACATATGGCATTTAGTTGTTCA
1065	binder	GGCACGGTAGAGAAATCGCATAACGTAC	CCTTCTTGAAGACCTATGTAAAGAAACGGGTCACT
1066	neutral	GAGACAGAGGATAAGGTCGTGTAATAGGTGG	CCATACGCATGACTACATTACAACGGGCCAGGAAG
1067	neutral	GAGACAGAGAAGAAAGTAAGCGTTGGCAC	CAGGTCTCGATCTCGTACAAAACGACTATGACCAT
1068	neutral	CCGGCTATGACTGCTTGAACGTACT	CATGAACGTGTCGTGTTATGCAGCGGTATGTCGTG
1069	binder	GAGACAGTTTGGTGGCACGGTAGA	CGTCGTGTCTTAGACGACTGTGTGTGATTCTCGAG
1070	binder	GCGAAATGACTGCGCAACGTACT	CCACCGTTTTGCCAGTTCCACCAGTAACTGTAAGG
1071	binder	AGAGTACTACCGCCTGACAACGTACT	CGCCCTGGAACAACGGTTATATTCTCTGGCAGAAC

Example 27: Oligonucleotide probes: aggressive breast cancer, non-aggressive breast cancer, and non-cancer

[001233] This Example presents oligonucleotide probes identified in an oligonucleotide probe library enriched against plasma-derived microvesicles from patients with aggressive breast cancer, non-aggressive breast cancer, and non-cancer (breast biopsy-negative samples). The general methodology follows that in **Example 26** above. Oligonucleotides were screened using plasma samples from 60 individuals with breast cancer and 60 individuals without breast cancer.

[001234] A combination of ultracentrifugation and PEG based partitioning of microvesicles from each input plasma sample. The F-TRin-35n-B 8-3s library as described herein was enriched against microvesicles from pooled plasma samples (cancer or non-cancer pools) which were isolated using ultracentrifugation. Subsequent enrichments were peformed by isolating the microvesicles using PEG precitation. As in **Example 26**, the input oligonucleotide library was exposed to positive, negative and again positive targets sequentially and then PCR amplified. We performed two parallel tracks of initial enrichment against pooled samples, one track where pooled cancer samples were considered positive and non-cancer samples were negative, and another track where pooled non-cancer samples were considered positive and cancer samples were negative. The enriched libraries resulting from each track were pooled together before probing individual patient samples in the following experiments.

[001235] Oligonucleotides that distinguish aggressive and non-aggressive breast cancer

[001236] We first identified oligonucleotides that distinguished between aggressive and non-aggressive breast cancer. For these experiments, the aggressive breast cancer patient samples were considered as positive samples (31 samples) and non-aggressive breast cancer patient samples were considered as negative samples (29 samples). The enriched libraries from above were subjected to the oligonucleotide probing assay on the individual cancer plasma samples. A two-step normalization procedure was performed to normalize the sum of the copy numbers for all oligonucleotides associated with the PEG-precipitated pellet. The copy number of each recovered oligonucleotide was divided by the total number of sequence reads for each sample, then multiplied by the mean of totals of all samples. This data treatment accounts for the different number of sequence reads for each patient even though the same amount of DNA library was added for each patient-sample. The final value for each oligonucleotide for each patient is the average of normalized copy numbers from three probing replicates. We excluded oligonucleotides with an average read-count across all patients ≤100, and subsequently selected among the remaining oligonucleotides only those that show a >20% CV across the 60 patient samples. This process resulted in approximately 12000 remaining oligonucleotides.

[001237] We compared patient profiles using clustering analysis with one minus Pearson correlation coefficients ($1 - P_c$) as the distance measure. The resulting heat-map based on the number of oligonucleotide sequence counts allows for visualisation of similarities in oligonucleotide copy numbers among all patient samples. See **FIG. 15**. In the figure, the X-axis shows individual oligonucleotides and the Y-axis shows samples (aggressive or other (i.e., non-aggressive)). Oligonucleotides from the indicated Cluster 1 have relatively low counts for the majority of aggressive cancers. Oligonucleotides from the indicated Clusters 2 and 3 have relatively higher counts in a majority of aggressive cancer samples. It is noted that oligonucleotide probes that are observed at either higher or lower relative copy number in the advanced cancer samples versus non aggressive can be used as markers to distinguish these sample groupings. Oligonucleotides from the indicated Cluster 4 were more selective toward particular aggressive cancer samples, and were able to identify additional 12 advanced cancer samples that were missed by oligonucleotides in Clusters 1-3. Cluster 4 oligonucleotides may be useful for detecting particular subsets of aggressive cancers.

[001238] The distances between samples within each cluster are considerably smaller than the distance between the two clusters. As can be seen in **FIG. 15**, these two groups of samples have very different proportions of advanced cancer cases versus other; in cluster I, 15 of the 19 patients are advanced cancer cases (79%), while in cluster II, only 16 of 40 patients are advanced cases (40%). Fisher exact p-value equals 0.01 with an odds ratio of 4.9, indicating that the ratio of advanced to other cancer in these two clusters is statistically significant.

[001239] The SEQ ID NOs. of the variable regions from the oligonucleotides in Clusters 1-4 in **FIG. 15** are shown in **Table 31**. As in **Table 27**, the oligonucleotides were each synthesized with a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132))

flanking the variable region. The sequences can be used in the identification of aggressive breast cancer samples.

Table 31: Clusters 1-4 Variable Regions SEQ ID NOs

Cluster	SEQ ID NO
Cluster 4	1072-1121
Cluster 3	1122-1171
Cluster 2	1172-1221
Cluster 1	1222-1421

[001240] A selection of 50 negative control oligonucleotides was made from the same experiments as above. As criteria, we chose those oligonucleotides with the lowest variability (% cv) between biological samples when taking all 120 samples from the noted probing into account. The variable regions of the 50 negative controls are listed in SEQ ID NOs 1422-1471. As in **Table 27**, the oligonucleotides were synthesized with a 5' region consisting of the sequence (5'-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5'-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)) flanking the variable regions. These sequences can be used as negative controls when identifying aggressive breast cancer samples.

[001241] Oligonucleotides specific to non-cancer samples versus cancer samples

[001242] We also identified a set of oligonucleotides that were specific to the non-cancer plasma samples. In addition to probing the cancer samples noted above, the enriched oligonucleotide probe library was used to perform the probing test on the 60 individual non-cancer plasma samples. Each sample was probed as described above. The oligonucleotides recovered with each individual non-cancer and cancer plasma sample were identified using next-generation sequencing. The averaged frequencies of individual oligonucleotide sequence reads between non-cancer and cancer samples were used to calculate a fold-change of non-cancer versus cancer. Oligonucleotides with both read counts above 100 and fold change (non-cancer/cancer) greater or equal to 2 were identified as non-cancer specific oligonucleotide probes (SEQ ID NOs. 1472-1486). In addition, a selection of negative control oligonucleotides were selected as those with fold change less than or equal to 1 and read count below 20 (SEQ ID NOs 1487-1501).

[001243] The variable regions of the non-cancer specific oligonucleotides and negative control oligonucleotides are listed by rank order in SEQ ID NOs. 1472-1486 and SEQ ID NOs 1487-1501, respectively. As in **Table 27**, the oligonucleotides were synthesized with a 5' region consisting of the sequence (5'-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5'-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)) flanking the variable regions. The sequences can be used to identify non-cancer samples (SEQ ID NOs. 1472-1486) or used as negative controls for such experiments (SEQ ID NOs 1487-1501).

[001244] Oligonucleotides specific to cancer samples versus non-cancer

[001245] We then identified a set of oligonucleotides that were specific to the cancer plasma samples as compared to the non-cancer samples. Pooled cancer and non-cancer samples were probed as

described above. The oligonucleotides recovered with each individual cancer and non-cancer plasma sample were identified using next-generation sequencing. The averaged frequencies of individual oligonucleotide sequence reads between cancer and non-cancer samples were used to calculate a fold-change of cancer versus non-cancer. Oligonucleotides with both read counts above 10 and fold change (non-cancer/cancer) greater or equal to 1.5 were identified as cancer specific oligonucleotide probes (SEQ ID NOs. 1502-1539). In addition, a selection of negative control oligonucleotides were selected as those with fold change less than or equal to 1 and read count below 20. See SEQ ID NOs 1487-1501 as described above.

[001246] The variable regions of the cancer specific oligonucleotides are listed by rank order in SEQ ID NOs. 1502-1539. As in **Table 27**, the full length oligonucleotides were synthesized with a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)) flanking the variable regions. These sequences which can be used to identify cancer samples.

Example 28: Oligonucleotide probes: Aggressive Breast Cancer, Non-Aggressive Cancer and Non-cancer/Healthy blood plasma targets with UC and PEG

[001247] This Example builds on the work presented in **Example 27**. Based on the results of the probing Aggressive cancer pool (50% patients with aggressive breast cancer), Non-Aggressive cancer pool and Healthy Pool (50% biopsy confirmed non-cancer and 50% healthy non-cancer), we obtained aptamers specific for:

[001248] Aggressive cancer compared to non-aggressive cancer and to non-cancer/healthy samples.

[001249] Non-aggressive cancer compared to aggressive and to non-cancer/healthy samples.

[001250] Non-Cancer/Healthy samples compared to aggressive and non-aggressive samples.

[001251] A combination of ultracentrifugation and PEG based partitioning of microvesicles from each input plasma sample. This approach potentially provides a cleaner patient sample and thus provide higher specificity for the intended target. As in **Example 27**, the input oligonucleotide library was exposed to positive, negative and again positive targets sequentially and then PCR amplified. Enriched libraries were subjected to the oligonucleotide probing assay on the plasma samples using sample pools as follows: ACP – aggressive cancer pool; NACP – non-aggressive cancer pool; HP – healthy pool (mix of non-cancer and normal samples). Each sample pool was probed two naïve (i.e., untrained) oligonucleotide probe libraries (6-3S and 8-3S described hereing) with three replicates. Samples were sequenced in multiplex with 9 samples per flow cell. Indexing PCR was performed either with or without internal standard. The average number of sequence read counts between three replicates was considered for fold change and TTEST calculations. Sequences were identified that met the following criteria: fold change between pools >= 2; p-value =<0.05, % cv =< 20%, counts >=100.

[001252] The joint selection of oligonucleotides that passed the above criteria is shown in **Tables 32** and **33**. As in **Table 27**, the oligonucleotides each have a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-

CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Tables 32** and **33** list in rank order the SEQ ID NOs of the variable region that are positioned between the 5' and 3' regions. The differences are that the starting libraries were derived from a first round of enrichment against a cancer pool (**Table 32**) or a first round of enrichment against a non-cancer pool (**Table 33**). However, after the first round the aliquots were pooled for subsequent enrichment. The combined oligonucleotide probe library can be used to distinguish between aggressive breast cancer, non-aggressive breast cancer, and non-cancer plasma samples in a single assay according to the column "Comparison." In the Comparison column, sequences elevated in group 1 as compared to group 2 are shown as group 1 / group 2. The legend for this column is "ACP": Aggressive cancer pool; "HP": healthy & non-cancer pool; "NACP": Non-aggressive cancer pool. "IntSt" in the comparison row indicates that an internal standard was used during the enrichment. By way of illustration, the first two sets of comparisons in **Table 32** are labeled ACP/HP, which indicates that these oligonucleotides were observed at high copy numbers after probing the aggressive breast cancer sample pool versus the healthy/non-cancer sample pool. Continuing with this illustration, higher numbers of such oligonucleotide probes binding to a sample can indicate the presence of an aggressive breast cancer.

Table 32: Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.
ACP/HP (IntSt)	1540-1551
ACP/HP	1552-1585
HP/ACP (IntSt)	1586-1611
HP/ACP	1612-1650
NACP/HP (IntSt)	1651-1655
NACP/HP	1656-1733
HP/NACP (IntSt)	1734-1754
HP/NACP	1755-1789
NACP/ACP (IntSt)	1790-1793
ACP/NACP (IntSt)	1794-1805
ACP/NACP	1806-1815

Table 33: Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.
ACP/HP (IntSt)	1816-1822
ACP/HP	1823-2004
HP/ACP (IntSt)	2005-2054
HP/ACP	2055-2234
NACP/HP (IntSt)	2235-2242
NACP/HP	2243-2318
HP/NACP (IntSt)	2319-2356
HP/NACP	2357-2491
NACP/ACP (IntSt)	2492-2507
NACP/ACP	2508-2583
ACP/NACP (IntSt)	2584-2593
ACP/NACP	2594-2728

Example 29: Oligonucleotide probes: Non-Aggressive Cancer probes identified in libraries enriched with multiple protocols

[001253] This Example builds on the work presented in **Example 28**. Here, we identified a set of oligonucleotide probes specific for non-aggressive cancer samples. The oligonucleotide probes were identified based on probing using an aggressive cancer pool (ACP), non-aggressive cancer pool (NACP) and healthy pool (HP) of plasma samples with naïve oligonucleotide libraries. Compared to the work in **Example 28**, four different enrichment schemes were used, as further detailed below. For purposes of these experiments, non-aggressive (or non-advanced) cancer is defined as sample with tumor size not exceeding stage T1, presence of nodes N0, Metastasis M0 and positive estrogen/progesterone receptors.

[001254] In each case, the enriched libraries were subjected to the probing test on pooled plasma samples. The library enrichment was performed as described herein, see e.g., **Example 21**. The libraries were exposed to positive and negative targets and PCR amplified either after each binding or after series positive-negative-positive binding as described below. The positive and negative pools are indicated below for the different enrichment schemes. After each round of enrichment, next generation sequencing was used to determine a copy number of the recovered oligonucleotide probes. Averaged frequencies of individual oligonucleotide between the indicate pools were used to calculate the fold change. Oligonucleotide with counts above 50, fold change >=2 and %cv between replicates below 20% were considered as candidates and shown in the tables below.

[001255] Scheme 1

[001256] The positive target was the NACP plasma and the negative target was the HP plasma. The enrichment started with the naïve (non-enriched; NE) library. Microvesicles in the samples were isolated using ultracentrifugation in all stages. Each round of enrichment consisted of a complete set of positive-negative-positive binding events followed by PCR amplification of the enriched library. Two such rounds of binding were performed. The resulting library after the second enrichment (R2) is used for subsequent probing of the pooled samples (aggressive cancer, non-aggressive cancer and healthy). As in **Example 28** above, comparisons performed included ACP/HP, HP/ACP, ACP/NACP, NACP/ACP, HP/NACP, NACP/HP.

[001257] Resulting oligonucleotides identified in these selections are shown in **Table 34**. As in **Table 27**, the oligonucleotides each have a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Table 34** lists the SEQ ID NOs in rank order of the variable regions that are positioned between the 5' and 3' regions. The combined oligonucleotide probe library can be used to distinguish between aggressive breast cancer, non-aggressive breast cancer, and non-cancer plasma samples in a single assay according to the column "Comparison." In the Comparison column, sequences elevated in group 1 as compared to group 2 are shown as group 1 / group 2. The legend for this column is "ACP": Aggressive cancer pool; "HP": healthy & non-cancer pool; "NACP": Non-aggressive cancer pool. By way of illustration, the first two sets of comparisons in **Table**

34 are labeled ACP/HP, which indicates that these oligonucleotides were observed at high copy numbers after probing the aggressive breast cancer sample pool versus the healthy/non-cancer sample pool. Continuing with this illustration, higher numbers of such oligonucleotide probes binding to a sample can indicate the presence of an aggressive breast cancer.

Table 34: Enrichment Scheme 1 Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.
HP/ACP	2729-2737
ACP/HP	2738-2742
ACP/NACP	2743-2745
NACP/ACP	2746-2753
NACP/HP	2754
HP/NACP	2755-2763

[001258] Scheme 2

[001259] In this enrichment, the positive target is NACP plasma and the negative target is the HP plasma. The starting input library is the C-R8N-1 (*i.e.*, previously enriched for a cancer pool with 48% of aggressive cancer samples; see description of the C-RN7 library in **Example 26** with one additional round of positive selection performed). The microvesicle partitioning was performed using PEG precipitation. PCR amplification is done after each binding. The resulting library after this ninth (Round 9) round of enrichment is used for probing. Three different input criteria for used for round 9: 1) R9-2 – library input 0.1 ng; 2) R9-1c – library input is 1ng; 3) R9 – 5x input of competitor DNA. The R9-2 and R9-1c oligonucleotide libraries were then used for subsequent probing of the pooled samples (aggressive cancer, non-aggressive cancer and healthy). As above, comparisons performed included ACP/HP, HP/ACP, ACP/NACP, NACP/ACP, HP/NACP, NACP/HP.

[001260] Resulting oligonucleotides identified from the R9-1c and R9-2 oligonucleotide libraries in these selections are shown in **Table 35** and **Table 36**, respectively. As in **Table 27**, the oligonucleotides each have a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Tables 35** and **36** list the SEQ ID NOs in rank order of the variable regions that are positioned between the 5' and 3' regions. When combined, the oligonucleotide probe libraries can be used to distinguish between aggressive breast cancer, non-aggressive breast cancer, and non-cancer plasma samples in a single assay according to the column "Comparison." In the Comparison column, sequences elevated in group 1 as compared to group 2 are shown as group 1 / group 2. The legend for this column is "ACP": Aggressive cancer pool; "HP": healthy & non-cancer pool; "NACP": Non-aggressive cancer pool. By way of illustration, the first two sets of comparisons in **Table 35** are labeled ACP/HP, which indicates that these oligonucleotides were observed at high copy numbers after probing the aggressive breast cancer sample pool versus the healthy/non-cancer sample pool. Continuing with this illustration, higher numbers of such oligonucleotide probes binding to a sample can indicate the presence of an aggressive breast cancer.

Table 35: Enrichment Scheme 2 R9-1c Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.
ACP/HP	2764-2791
HP/ACP	2792-2793
ACP/NACP	2794-2815
NACP/ACP	2816-2830
HP vs NACP	2831-2845
NACP vs HP	2846-2858

Table 36: Enrichment Scheme 2 R9-2 Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.
ACP/HP	2859-2867
HP/ACP	2868-2892
ACP/NACP	2893-2904
NACP/ACP	2905-2938
HP/NACP	2939-2941
NACP/HP	2942-2950

[001261] Scheme 3

[001262] The positive target was the NACP plasma and the negative target was the HP plasma. The enrichment started with the naïve (non-enriched; NE) library. Microvesicles in the samples were isolated using PEG precipitation. Six rounds of enrichment were performed with PCR amplification performed after each round. The round alternated between a single selection of positive or negative selection (i.e., positive selection in rounds 1, 3, 5 and negative selection in rounds 2, 4, and 6). The resulting library after round 6 is used for probing of the pooled samples (aggressive cancer, non-aggressive cancer and healthy) with library input 0.1 ng. As above, comparisons performed included ACP/HP, HP/ACP, ACP/NACP, NACP/ACP, HP/NACP, NACP/HP.

[001263] Resulting oligonucleotides identified in these selections are shown in **Table 37**. As in **Table 27**, the oligonucleotides each have a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Table 37** lists the SEQ ID NOs in rank order of the variable regions that are positioned between the 5' and 3' regions. The combined oligonucleotide probe library can be used to distinguish between aggressive breast cancer, non-aggressive breast cancer, and non-cancer plasma samples in a single assay according to the column "Comparison." In the Comparison column, sequences elevated in group 1 as compared to group 2 are shown as group 1 / group 2. The legend for this column is "ACP": Aggressive cancer pool; "HP": healthy & non-cancer pool; "NACP": Non-aggressive cancer pool. By way of illustration, the first two sets of comparisons in **Table 37** are labeled ACP/HP, which indicates that these oligonucleotides were observed at high copy numbers after probing the aggressive breast cancer sample pool versus the healthy/non-cancer sample pool. Continuing with this illustration, higher numbers of such oligonucleotide probes binding to a sample can indicate the presence of an aggressive breast cancer.

Table 37: Enrichment Scheme 3 Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.
ACP/HP	2951-2960
HP/ACP	2961-2972
ACP/NACP	2973-2976
NACP/ACP	2977-2990
HP vs NACP	2991-3003
NACP/HP	3004-3017

[001264] Scheme 4

[001265] Starting aptamer library is C-R8N-3 (library previously enriched for Cancer pool with 48% of aggressive cancer samples and input 0.001ng). Microvesicles in the samples were isolated using PEG precipitation. PCR amplification is done after each binding. There was only negative target used (the NACP), which is unique for the selection scheme versus the three above. This enrichment was meant to further enrich C-R8N library toward aggressive cancer targets by removing oligonucleotides specific to non-aggressive targets. Four negative selection rounds were performed. Two different input criteria for used for probing after round 13 (R13): 1) R13-3 – library input 0.01 ng; 2) R13-4 – library input 0.001 ng. The R13-3 and R13-4 oligonucleotide libraries were then used for subsequent probing of the pooled samples (aggressive cancer, non-aggressive cancer and healthy). As above, comparisons performed included ACP/HP, HP/ACP, ACP/NACP, NACP/ACP, HP/NACP, NACP/HP.

[001266] The variable regions of the resulting oligonucleotides identified from the R13-3 and R13-4 oligonucleotide libraries in these selections are shown in **Table 38** and **Table 39**, respectively. As in **Table 27**, the oligonucleotides each have a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Tables 38** and **39** list the SEQ ID NOs in rank order of the variable regions that are positioned between the 5' and 3' regions. When combined, the oligonucleotide probe libraries can be used to distinguish between aggressive breast cancer, non-aggressive breast cancer, and non-cancer plasma samples in a single assay according to the column "Comparison." In the Comparison column, sequences elevated in group 1 as compared to group 2 are shown as group 1 / group 2. The legend for this column is "ACP": Aggressive cancer pool; "HP": healthy & non-cancer pool; "NACP": Non-aggressive cancer pool. By way of illustration, the first two sets of comparisons in **Table 38** are labeled ACP/HP, which indicates that these oligonucleotides were observed at high copy numbers after probing the aggressive breast cancer sample pool versus the healthy/non-cancer sample pool. Continuing with this illustration, higher numbers of such oligonucleotide probes binding to a sample can indicate the presence of an aggressive breast cancer.

Table 38: Enrichment Scheme 4 R13-3 Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.	Variable Region Sequence
ACP/HP	3018-3030	ACCATCGGGTCACGTAAAAGCGGTTCTAATTTCTAA
HP/ACP	3031-3043	ACGCAGTGCCGCCGAATCCTGATCCCCTAACTTTC
ACP/NACP	3044-3052	ACCACTGAACTCATGGTACTAATCACACACCAATT
NACP/ACP	3053-3061	ACTGACTGCATACATCCGCATATTAAACCCGTATT

HP/ACP	3062-3083	ACGCCTGTACGATGTCCTCGGTCTCATTTTATTTC
ACP/HP	3084-3096	ACTGCGGTCATCGAGTTACTGGCTATTGCCTGACC

Table 39: Enrichment Scheme 4 R13-4 Oligonucleotide Probe Comparison and Variable Regions

Comparison	SEQ ID NO.	Variable Region Sequence
ACP/HP	3097-3099	AATGCATTGCGTTATTTCAATTGAGGCGTCCTAAAA
HP/ACP	3100-3107	CATACGCACGGTCCTCTACTATAATTGTCCACCAA
ACP/NACP	3108-3117	AGCCCGTTGCTACGTTGCAGTAGGTGTAACTTCACTGA
NACP/ACP	3118-3130	AGCGGTCACCGCCTATGGTTATTCATTTTTCTCTTA
HP/NACP	3131-3141	AGCCCGTTGCTACGTTGCAGTAGGTGTAACTTCACTGA
NACP/HP	3142-3150	ACCACGTTGGTTATTGCCCTGCTTTGAAGTCTTACGA

Example 30: Use of an oligonucleotide probe library to characterize Breast Cancer samples

[001267] An oligonucleotide probe library comprising approximately 2000 different probe sequences was constructed and used to probe approximately 500 individual breast cancer and non-cancer samples. The probe sequences were derived from different screening experiments and are listed herein in SEQ ID NOs 137-969 and 1072-3150. The oligonucleotides listed in these tables were synthesized and pooled together. The samples were plasma samples from 212 breast cancer patients, 177 biospy confirmed non-cancer patients, and 117 normal control patients (self-reported as non-cancer). Experiments details are as provided above. The plasma samples were contacted with the oligonucleotide probe library and microvesicles were isolated using PEG precipitation. Oligonucleotides that were recovered with the microvesicles were isolated. Next Generation Sequencing (Illumina HiSeq) was used to identify the isolated sequences for each sample.

[001268] Analysis of significance of difference identified 18 aptamers with p-values below 0.01 when compared Cancer/Normal, 15 aptamers with p-values below 0.001 when compared cancer/Non-Cancer, 28 aptamers with p-values below 0.001 when compared Non-Cancer/Normal.

[001269] Multi-oligonucleotide panels were next contructed using a cross-validation approach. Briefly, 50 samples were randomly withheld from the sample cohort. The performance of individual oligonucleotides to distinguish the remaining cancers and non-cancer/normals was determined using logistic regression methodology. Additional oligonucleotides were added iteratively and performance was assessed using logistic regression until further performance improvements were no longer obtained with additional oligonucleotides. The approach generally led to panels of approximately 20-100 different probe sequences. The contructed panels were then used to classify the 50 withheld samples and diagnostic performance was assessed using Receiver Operating Curve (ROC) analysis and estimation of the Area Under the Curve (AUC).

[001270] In approximately 300 rounds of cross-validation, the average AUC was 0.6, thus showing that the average performance was statistically better than random (i.e., AUC of 0.5) and that the probe library could distinguish breast cancer and non-breast cancer/normal patient samples. AUC values as high as 0.8 were observed for particular cross validations. **FIGs. 16A-B** illustrate a model generated using a training (**FIG. 16A**) and test (**FIG. 16B**) set from a round of cross validation. The AUC was 0.803. The

variable regions of the sequences used to build this model are shown in **Table 40**. Another exemplary round of cross-validation is shown in **FIGs. 16C-D**. The AUC was 0.678.

[001271] The SEQ ID NOs. of the sequences used in the model in **FIGs. 16A-B** are listed in rank in **Table 40**. As in **Table 27**, the oligonucleotides were synthesized with a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)) flanking the variable regions.

Table 40: Oligonucleotide Probe Variable Regions

Rank Ordered SEQ ID NOs
215, 1286, 961, 1837, 780, 1319, 3032, 626, 2816, 1311, 364, 3102, 3115, 886, 414, 517, 599, 246, 416, 223, 507, 586, 1455, 1560, 1241, 2771, 1513, 2994, 2757, 461, 1917, 1178, 299, 1409, 959, 785, 322, 636, 1244, 665, 592, 823, 168, 1183, 3000, 182, 534, 1580, 2753, 2989, 1957, 2829, 1960, 856, 3149, 283, 1551, 1974, 605, 363, 266, 3140, 2242, 1306, 652, 634, 2763, 1270, 1728, 893, 1266, 1372, 1141, 1731, 1197, 1649

[001272] The data presented in this Example demonstrate that an oligonucleotide pool comprising members having the variable regions listed in SEQ ID NOs 137-969 and 1072-3150, e.g., a pool of probes having the variable regions listed in **Table 40**, can be used to distinguish plasma from individuals having breast cancer versus plasma from non-breast cancer individuals.

Example 31: Updated oligonucleotide pools to characterize Breast Cancer samples

[001273] The profiling of 500 clinical samples with an oligonucleotide probe library comprising 2000 oligonucleotides showed significant ability to distinguish breast cancer from non-cancer control samples. See **Example 30** above. When performing cross-validation in these experiments, it was observed that 85 out of 500 were misclassified at a higher rate compared to the other samples. We therefore identified another selection of oligonucleotide probe which were able to correctly classify the noted samples. These 85 samples were profiled with a naïve oligonucleotide probe library (6-3S/8-3S), and enriched on breast cancer and non-cancer plasma using methodology presented in the Examples above. The selected oligonucleotides were compared to a positive controls cohort, which comprised of the cancer and non-cancer samples that were consistently classified correctly within the experiments in **Example 30** above. Oligonucleotides were selected based on absolute number of copies sequenced of at least 50 and various criteria when comparing sample groups including: 1) copy number fold-change (fc) of at least 1.2; 2) effect size (es) above 0.6; 3) t-test (p-value <= 0.05); and 4) Kolmogorov-Smirnov test (ks) (p-value <= 0.05). Effect size is a population effect size calculated as (mean(group1) – mean(group2)) / standard deviation(group1 and group2).

[001274] The profiling data were normalized by dividing the count of each particular oligonucleotide by the total counts for particular sample and multiplying by the global mean across the entire experiment. These normalized values were used to calculate the statistical criteria specified above to compare the following samples type: Misclassified Cancer ("C"), misclassified non-cancer ("NC"), positive control Cancer ("C-P"), positive control Non-Cancer ("NC-P"), misclassified Normal ("N"). The

comparisons performed and numbers of resulting oligonucleotides are shown in **Table 41**. Negative controls are oligonucleotides that did not match any criteria and thus should not distinguish be any samples groups.

Table 41: Oligonucleotide probe candidates were selected from the following comparisons

Specificity	Total
C/NC-P* A** and C/NC-P B***	82
C/NC-P A	108
C/NC-P B	13
Negative control	60
C-P/NC A C-P/NC B	34
C-P/NC A	68
C-P/NC B	101
NC/C-P A NC/C-P B	7
NC/C-P A	196
NC-P/C A NC-P/C B	65
NC-P/C A	83
NC-P/C B	55
C+C-P/N	103
N/C+C-P	25
Total	**1000**

[001275] * "C/NC-P" in **Table 41** means that oligonucleotides were selected with specificity toward cancer ("C") while compared to Non-Cancer positive ("NC-P") control.

[001276] A** - selection criteria: counts > 50, fc ≥ 1.2, es > 0.6, ttest $p<0.05$,

[001277] B*** - selection criteria: counts >200, estimated sample size <71, fc >1.2, ttest/ks test $p<0.05$.

[001278] The sequences selected based on **Table 41** are shown in **Table 42**. As in **Table 27**, the oligonucleotides each have a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Table 42** lists in rank order the SEQ ID NOs of the variable regions that are positioned between the 5' and 3' regions. The column "Specificity" shows the specificity of the oligonucleotides based on the comparisons in **Table 41**.

Table 42: Oligonucleotide Probe Variable Regions

Specificity	SEQ ID NO.
C/NC-P A and C/NC-P B	3151-3232
C/NC-P A	3233-3340
C/NC-P B	3341-3353
Negative control	3354-3368
C-P/NC A C-P/NC B	3369-3402
C-P/NC A	3403-3470
C-P/NC B	3471-3571
Negative control	3572-3586
NC/C-P A NC/C-P B	3587-3593
NC/C-P A	3594-3789
Negative control	3790-3804
NC-P/C A NC-P/C B	3805-3869

NC-P/C A	3870-3952
NC-P/C B	3953-4007
Negative control	4008-4022
C/N	4023-4125
N/C	4126-4150

[001279] The 1000 selected oligonucleotides are synthesized as DNA with a 5' biotinylation. The synthesized oligonucleotides are pooled with 1000 of the most informative aptamers from the profiling performed in **Example 30** to create a further optimized oligonucleotide probing library that can be used to distinguish cancer and normal samples.

Example 32: Comparison of Oligonucleotide Pools

[001280] In this Example, the performance of the oligonucleotide pool from **Example 30** was compared to the performance of the oligonucleotide pool from **Example 31** to characterize breast cancer samples. In this Example, the oligonucleotide pool from **Example 30** is referred to as the 2000v1 and the oligonucleotide pool from **Example 30** is referred to as the 2000v2. A random forest approach was used to analyze the data, which data was obtained for the 2000v1 library in **Example 30**. The 2000v2 data was obtained via tested on a cohort of 858 plasma samples which comprised some samples described in **Example 30** with additional samples according to three groupings: 194 breast cancer positive, 382 breast cancer negative as confirmed by biopsy ("non-cancer"), and 282 self declared normals ("normal").

[001281] ROC curves are shown in **FIGs. 22A-D** for the various settings. AUC values are shown above the curve and are statistically significant from random (i.e., AUC = 0.5) in all cases. In **FIG. 22A**, the 2000v1 probe library was used to distinguish cancer samples versus biopsy confirmed non-cancer samples. In **FIG. 22B**, the 2000v2 probe library was used to distinguish the same sample groupings. A small improvement in AUC was observed with the 2000v2 probe library in this setting. In **FIG. 22C**, the 2000v1 probe library was used to distinguish cancer samples versus self-declared normal samples. In **FIG. 22D**, the 2000v2 probe library was used to distinguish the same sample groupings. A small decline in AUC was observed with the 2000v2 probe library in this setting. Highly stringent

[001282] The SEQ ID NOs. of the variable regions of informative oligonucleotides from the 2000v1 and 2000v2 libraries are listed in **Table 43**. As above, the oligonucleotides were synthesized with a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)) flanking the variable regions. The listed oligonucleotides were used in the Random Forest modeling to generate AUC values, *e.g.*, as shown in **FIGs. 22A-D**, and are listed in rank order. The row "Common" in **Table 43** shows overlapping informative sequences from the 2000v1 and 2000v2 settings.

Table 43: Informative oligonucleotide probes

Setting	SEQ ID NO.
2000v1	1102, 624, 3032, 597, 2831, 1923, 706, 752, 1188, 238, 605, 2809, 651, 611, 2830, 1163, 1187, 1106, 421, 843, 1186, 1100, 1235, 3140, 173, 1243, 289, 178, 3035, 1080, 890, 1225, 1995, 508, 183, 435, 2584, 2795, 1814, 1924, 3075, 939, 1183, 431, 938, 1731, 1095, 446, 623, 1732, 3024, 216, 626, 672, 457, 830, 1497, 777, 610, 3038, 1460, 665, 1079, 148, 1445,

	885, 1697, 1197, 1769, 965, 164, 838, 606, 3018, 493, 1830, 859, 2047, 1837, 404, 1693, 1201, 899, 264, 841, 1678, 1949, 628, 354, 1523, 3036, 495, 727, 1997, 676, 546, 1507, 615, 616, 3096, 1435, 1194, 1195, 698, 660, 1994, 1123, 290, 934, 428, 715, 751, 3064, 602, 345, 662, 1820, 586, 241, 929, 1136, 316, 803, 3095, 936, 1381, 537, 254, 1119, 1652, 1955, 1534, 2965, 496, 405, 757, 612, 145, 1248, 599, 942, 505, 907, 1353, 2786, 1509, 211, 1170, 736, 486, 815, 540, 1540, 1250, 3000, 2971, 348, 3108, 445, 1548, 1723, 888, 1284, 604, 858, 1372, 1083, 584, 767, 3135, 855, 2730, 805, 1557, 565, 1600, 920, 2330, 1977, 710, 1220, 1907, 441, 795, 1939, 1386, 908, 2986, 350, 552, 2953, 3134, 250, 765, 827, 1216, 526, 969, 1903, 693, 650, 1359, 789, 747, 826, 2002, 1291, 639, 1244, 573, 3119, 523, 594, 854, 898, 429, 1881, 569, 1538, 848, 738, 911, 669, 909, 329, 577, 2964, 272, 648, 1481, 1719, 816, 2981, 533, 2748, 1660, 3050, 755, 1706, 1504, 3059, 1276, 1547, 2732, 922, 2977, 2952, 1522, 1273, 1399, 1556, 2998, 853, 1256, 845, 625, 2005, 2987, 2985, 1327, 807, 923, 1500, 1331, 2497, 641, 1282, 1940, 629, 2970, 1150, 1915, 1862, 389, 1304, 2979, 3061, 1328, 2740, 819, 353, 396, 2983, 1164, 3029, 1566, 2041, 2972, 966
2000v2	4116, 171, 3280, 3401, 3879, 1175, 3291, 651, 3343, 592, 3110, 3682, 605, 3231, 3556, 389, 624, 399, 3173, 3483, 795, 3918, 211, 309, 3595, 4101, 3780, 1169, 803, 1392, 3332, 415, 3652, 1610, 3262, 163, 1477, 4044, 3319, 722, 3607, 3414, 564, 511, 826, 431, 4052, 3329, 692, 3153, 295, 1548, 635, 3900, 370, 3698, 342, 3545, 614, 178, 3699, 467, 3307, 770, 4055, 3912, 164, 3232, 4048, 2762, 3856, 2587, 421, 3683, 3000, 633, 3712, 3466, 3769, 3537, 4058, 4018, 2808, 4026, 3785, 4083, 3151, 1288, 4050, 3560, 3786, 1121, 3742, 3128, 4132, 1171, 3619, 3471, 566, 3751, 4110, 4091, 787, 1176, 3735, 538, 240, 3064, 2754, 3035, 199, 3777, 831, 1247, 1188, 3667, 3086, 3741, 3650, 1272, 3402, 3317, 3618, 1466, 1590, 3489, 3659, 1091, 3601, 4023, 1469, 1777, 2008, 4059, 1593, 4062, 205, 2838, 3087, 2835, 3661, 1661, 3551, 3612, 1322, 3629, 3789, 2017, 1325
Common	1188, 3064, 421, 624, 389, 803, 431, 826, 3035, 164, 178, 651, 3000, 1548, 211, 605, 795

[001283] The data presented in this Example demonstrate that an oligonucleotide pool comprising members having the variable regions listed in SEQ ID NOs 137-969 and 1072-4150, e.g., a pool of probes having the variable regions listed in **Table 43**, can be used to distinguish plasma from individuals having breast cancer versus plasma from non-breast cancer individuals.

Example 33: Photo-cleavable Biotin mediated purification of an oligonucleotide library

[001284] Tag probe and capture probe detection systems may have advantages in the read-out of oligonucleotide-based assays since such systems allow direct quantification of oligonucleotides without amplification. Amplification may introduce amplification bias, e.g., wherein some sequences are preferentially amplified versus others. One such system is the nCounter nucleic acid detection system from NanoString Technologies, Inc. (Seattle, WA). This system can be used to detect oligonucleotide probes of the invention. See, e.g., **FIG. 14G** and related discussion. Current applications of tag and capture probe detection systems include gene expression analysis, RNA:Protein analysis; single cell gene expression analysis; miRNA expression analysis, miRGE analysis, copy number variation analysis, lncRNA expression analysis; ChiP-String expression analysis. *See, e.g.,* www.nanostring.com/applications/.

[001285] In this Example, we improved using the Nanostring system to detect the oligonucleotide probe libraries of the invention. As described above, a pre-enriched oligonucleotide probe library is contacted with a biological sample such as plasma, and microvesicles bound by various oligonucleotide probes are precipitated, e.g., with PEG. Thus, the oligonucleotide probe library is present in the mix with

PEG/proteins. This mixture is compatible with PCR steps used for next-generation sequencing (NGS) preparation, but the Nanostring system may not detect probes in these mixtures. Without being bound by theory, this interference may be due to noted impurities with hybridization of the probes and other oligonucleotides used in purifications steps. We developed a protocol for purification of post-probing oligonucleotide probe library, which utilizes photo-cleavable (PC) Biotin (see, e.g., www.idtdna.com/site/Catalog/Modifications/Product/2291; references 1-4) and thus allows oligonucleotide probe detection using the Nanostring tag-capture system.

[001286] The steps of the purification of post-probing oligonucleotide probe library are shown in schematic **1700** in **FIG. 17A** and include the following: (i) modification of the 5'-end of each member of the oligonucleotide library with photocleavable-Biotin (PC-Biotin) **1701** via PCR with a PC-Biotinylated primer to create PC-Biotinylated aptamer library **1702**; (ii) probing of plasma samples using the PC-Biotinyltated oligonucleotide probe library followed by PEG precipitation as described herein, resulting in a pellet **1703** which contains, inter alia, bound oligonucleotide library members, proteins and microvesicles **1705**, and PEG; (iii) heat denaturing of the re-suspended pellet to break oligonucleotide/protein interactions and make PEG less viscous before mixing with streptavidin beads **1704**; (iv) after short incubation with streptavidin beads **1704** to allow library **1702** to bind to the beads **1706**, washing of the beads to remove impurities; (v) UV photo-cleavage **1707** of the spacer between the PC-Biotin and oligonucleotide molecules, capturing the beads **1708** using a magnet and collecting the released oligonucleotide library **1709** in the solution; and (vi) Nanostring **1710** sample preparation according to the manufacturer's instructions. In **FIG. 17A**, "PCB" refers to photocleavable biotin, oligonucleotide library members are refered to as "AL," "Str. Beads" refers to streptavidin beads, and "NS" refers to the Nanostring assay. The steps above are identified in the plot.

[001287] **FIG. 17B** show a 4% agarose gel with results of photocleavable biotin library cleavage test with Streptavidin protein. Lane 1 is a 50bp DNA ladder and lane 10 is a Control with no biotin library with Streptavidin protein. The gel shows that both biotinylated ("B") and photo-cleavable biotinylated ("PC-B") modified oligonucleotides were bound to Streptavidin, which resulted in slower mobility in gel (lanes 4 and 5) as compared to the unbound oligonucleotides (lanes 2 and 3). After exposure to 365 nm UV radiation, the PC-B oligonucleotides were photo-cleaved, which resulted in faster mobility in gel and returning to the original molecular weight (i.e., before mixing with Streptavidin protein) (lanes 7-9, where minutes UV exposure are indicated). In contrast, no change to the B oligonucleotides was observed after 10 min exposure to UV (lane 8). These results demonstrate that the photocleavable biotin library can be captured by streptavidin beads and released by UV-mediated cleavage.

[001288] To test the workflow, a pool of plasma from patients with breast cancer and another pool of plasma from healthy donors were probed with an oligonucleotide library compriseing 100 oligonucleotide probes, 50 of which were selected as cancer specific and other 50 were selected as equivalent binders to both types of samples (i.e., "neutral"). The Nanostring probe system **1720** was designed as in **FIG. 17C**. See **FIG. 14G** and description thereof for further details of the Nanostring probe

system. The "Nanostring Probe A" **1722** and "Nanostring Probe B" **1727** were designed in part as reverse complements to the oligonucleotide probe library members of the invention (labeled "Aptamer" **1724** in the figure). As can be seen in **FIGs. 17D-E**, relevant Nanostring counts were obtained for the photocleavable oligonucleotide library (min relevant count is 300). Furthermore, the oligonucleotide probes selected as binders ("Binder") for the cancer pool ("CP"), indeed show stronger binding to the cancer pool as compared to counts from healthy pool ("HP"). The binding ratio between cancers and normal was ~5.25. For oligonucleotide probes selected as neutral binders ("Neutral"), there was no relevant difference between the cancer pool and healthy pool.

[001289] The Nanostring protocol uses hybridization of target specific probes. We tested 100 different oligonucleotide probes, which are complementary to portion of each specific oligonucleotide, as well as universal Primer probe for Nanostring detection using the above procedure. Following the standard Nanostring detection protocol, all probes were mixed with recovered and purified library, hybridized overnight and efficiently detected using the Nanostring tag-capture probe approach. In contrast, without purification no quantification of oligonucleotides with Nanostring was observed (data not shown).

[001290] In this Example, we established a process for oligonucleotide probe library purification after probing test samples. This process is efficient, low cost, can be accomplished in short time frame, and can be used for any purpose when library needs to be purified from probing product directly and quickly, given no biotin labeling is needed in downstream applications.

[001291] References:

[001292] 1. Photocleavable biotin derivatives: a versatile approach for the isolation of biomolecules. Proc Natl Acad Sci U S A. 1995, 16, 7590-7594.

[001293] 2. Photocleavable biotin phosphoramidite for 5'-end-labeling, affinity purification and phosphorylation of synthetic oligonucleotides. Nucleic Acids Res. 1996, 2, 361-366.

[001294] 3. Photocleavable affinity tags for isolation and detection of biomolecules. Methods Enzymol. 1998, 291, 135-154.

[001295] 4. Matrix-assisted laser desorption/ionization mass spectrometry of DNA using photocleavable biotin. Biomol Eng. 1999, 16, 127-133.

Example 34: Single stranded DNA (ssDNA) oligonucleotide library preparation for library development

[001296] The preparation of high yield and high quality ssDNA libraries is a critical step in SELEX (Systematic Evolution of Ligands by EXponential enrichment) [1, 2] as well as in other biological applications, such as DNA chips and microarrays [3], and single-stranded conformation polymorphism technique (SSCP) [4]. The standard approach for preparing ssDNA libraries includes PCR amplification to first generate a double stranded (dsDNA) library, followed by ssDNA separation and purification. Several strategies of ssDNA preparation have been developed to date, each with advantages and disadvantages:

[001297] Lambda exonuclease digestion [2, 5-7]

[001298] The dsDNA standard PCR product is followed by Lambda exonuclease to digest the complementary strand and leave the target ssDNA. ssDNA purification is then performed to remove enzymes and unwanted buffer.

[001299] *Advantages*: Regular PCR amplification has high yield in generating dsDNA.

[001300] *Disadvantages*: The purity of final ssDNA is limited by enzyme digestion efficiency. Also dsDNA needs to be purified prior to digestion, together with post-digestion purification there will be two purifications, which results in substantial loss of input material. The digestion usually requires at least 2 hours. The digestion rate may not be consistent.

[001301] Asymmetric PCR [8, 9]

[001302] The procedure generates target ssDNA as the main product and less dsDNA products and non-target ssDNA. The band corresponding to the target ssDNA is cut from a native gel.

[001303] *Advantages*: The final ssDNA product potentially has high purity.

[001304] *Disadvantages*: Separation of strands is possible in the native gel, but the yield is typically low and the presence of non-target strand cannot be excluded. The yield cannot be increased on denaturing gel because the strands have the same length.

[001305] Biotin-streptavidin magnetic beads separation [10, 11]

[001306] The non-target PCR primer is biotinylated so final PCR products are Biotinylated-dsDNA, which can be captured by streptavidin magnetic beads and denatured to release the non-biotin labeled target ssDNA.

[001307] *Advantages*: The final ssDNA product has relatively high purity.

[001308] *Disadvantages*: In most cases, the input library needs to be biotinylated, but it may be difficult to replace or release the captured target strands from streptavidin beads. Post-denaturing purification is required to remove NaOH and/or acid used for neutralization.

[001309] Unequal primer length PCR [12]

[001310] The non-target PCR primer has a chemical modified spacer and a few extra nucleotides following. In the PCR reaction, the DNA polymerase will stop at the spacer, resulting in unequal length of PCR dsDNA product. Then target ssDNA can be cut from a denaturing PAGE gel.

[001311] *Advantages*: The final ssDNA product has high purity because the target ssDNA is not mixed with non-target strands.

[001312] *Disadvantages*: ssDNA cannot be seen on native gel. Requires time consuming denaturing PAGE gel. It may be difficult to denature some dsDNA library, which can limit the final yield.

[001313] Indirect purification method [13]

[001314] The indirect purification strategy combines Asymmetric PCR and Biotin-streptavidin magnetic beads separation. In short, regular PCR is used to generate sufficient template, then asymmetric PCR with excess of target primer and less biotinylated complementary primers, followed by biotin-streptavidin separation.

[001315] *Advantages*: May increase yield and purity of ssDNA product.

[001316] *Disadvantages*: It cannot produce biotinylated target ssDNA library. The process is relatively long and complicated and may be prone to generate mutants of the original sequence.

[001317] The invention provides methods of enriching oligonucleotide probe libraries against a target of interest. As the probes comprise ssDNA, the process may comprise PCR amplification then conversion back into ssDNA after each round of enrichment. In this Example, we developed a strategy for preparation of a ssDNA oligonucleotide library. The goals were to develop a process that is efficient and quick, while delivering high quality/purity ssDNA. We aimed to combine PCR and ssDNA prep in one step, remain efficient in the presence of selection buffer, target molecules, other sample components (e.g., highly abundant proteins for plasma samples) and other assay components (e.g., PEG precipitation solution that may be used to precipitate microvesicles). In addition, we desired the method to be able to generate ssDNA library with any modification, including without limitation Biotin.

[001318] We have used an optimized version of Lambda exonuclease digestion protocol for preparation of ssDNA oligonucleotide library. However, the digestion yield limits the overall recovery and is not consistent between different library preparations. In some cases, the ssDNA band is hardly visible on the gel following digestion. We have also observed incomplete digestion of dsDNA in the ssDNA product. In this Example, we developed an alternative protocol, termed "ssDNA by Unequal length PRimer Asymmetric PCR," or SUPRA. It lacks disadvantages from the known methods listed above, and provides high quality and yield up to 10x higher yield of ssDNA oligonucleotide library as compared to the previous methods. It is relatively fast and convenient technically, since target ssDNA can be distinguished from non-target DNA on a gel.

[001319] A schematic comparing standard PCR **1800** and unequal length PCR **1810** is shown in **FIG. 18A**. In regular PCR **1800**, a formard primer **1801** and reverse primer **1803** are hybridized with the reverse strand of an aptamer library **1802**. The PCR reaction is performed, thereby creating equal length forward **1804** and reverse strands **1802**. The strands are denatured in equal length single strands **1805**. In unequal length PCR **1801**, a formard primer **1811** having a lengthener segment and terminator segment and a reverse primer **1813** are hybridized with the reverse strand of an aptamer library **1812**. The PCR reaction is performed, thereby creating unequal length forward **1814** and reverse strands **1812**. The strands are denatured into unequal length single strands **1814** and **1812** that can be separated by size, e.g., on a denaturing gel.

[001320] The steps of SUPRA include: (i) Modification of regular non-target primer with two Isp9 (Internal Spacer 9; triethylene glycol spacer) as terminator and 32 extra nucleotides (e.g., poly-A) as lengthener. It is referred as Unequal-Forward-Double isp9 primer (UF-D9); (ii) Perform asymmetric PCR, by mixing DNA template, UF-D9 and regular target (reverse) primer at ratio that favors the reverse primer, e.g., 1:37.5. The PCR program has longer elongation step (e.g., 3 min instead of standard 1 min) and more cycles due to linear amplification mode (instead of exponential). The PCR product contains a majority of target ssDNA and small portion of dsDNA. (iii) Mix PCR reaction products 1:1 with

denaturing buffer (e.g., 180 mM NaOH and 6 mM EDTA) and denature samples by heating (e.g., 70°C for 10 min) and cooling (e.g., incubation on ice for 3 min); (iv) Run denatured products in denaturing buffer on an agarose gel stained with SybrGold. The non-target strand, which is longer due to the lengthener, will appear as upper band (if visible) and the target strand (strong lower band) is cut and purified. The process can include optional steps, including without limitation: (v) Weigh the gel pieces and purify ssDNA from the gel pieces (e.g., using the ssDNA Nucleospin kit or the like); (vi) quantification of the yield and native gel can be used to check the purity and yield of final product (e.g., using the ssDNA Qubit kit or the like).

[001321] The first step (i) uses a specific design of the forward primer with efficient terminator and lengthener, which creates non-target strand of unequal length. The DNA polymerase used to build the target strand will stop polymerization once it reaches the terminator, and the lengthener facilitates differentiation between the target and non-target strands. In the second step (ii), the ratio between the two primers is shifted toward the reverse primer, to produce a majority of target ssDNA. The ratio, however, should not limit double strand templates production to keep reaction running. **FIG. 18B** is a gel showing titration of forward and reverse primers input in asymmetrical PCR. The optimal condition, at which target strand is clearly visible, is in the range 1:20-1:50 F:R primers ratio. As shown in the figure, the ratio between two primers in asymmetric PCR can affect dsDNA and ssDNA amount in final products. The PCR thermocycler program is also adjusted to provide efficiency in the asymmetric PCR. In the third step (iii), a reliable denaturing method is used to separate target ssDNA to ensure the final yield and high purity.

[001322] As desired, the final step (vi) estimates the ratio of residual dsDNA, e.g., using ssDNA Qubit kit. In cases where the yield is not critical, the denaturing steps (iii and iv) can be skipped and the PCR products can be directly run on native gel. There will be a dsDNA band, but lower MW target ssDNA band can be distinguished and purified from gel. This is also a way to visualize the target band directly after PCR for a quality check or purification without denaturing. The purity of final product will be the same but yield will be lower.

[001323] A comparison of native versus denatured gel purification is shown in **FIG. 18C**. A post-probing oligonucleotide probe library was PCRed using unequal length primers mixed at a ratio of 1:38 (Forward/Reverse). In the figure, the left lane on each gel is a 50 bp molecular weight ladder and the lower band is the reverse primer. The positions of the dsDNA and ss DNA are indicated. A native gel showed the presence of both dsDNA and ssDNA (target strand) (**FIG. 18C, panel A**). Here, part of the target reverse strand is migrating in dsDNA. Thus, using the native gel, one can purify target ssDNA with moderate recovery. When a higher yield is desired, the PCR products can be run on denaturing agarose gel as described above. This approach provides maximal recovery wherein only target strand is visible, and can be cut from gel and purified (**FIG. 18C, panel B**). In this case, the reverse strand ssDNA, which is part of the dsDNA on native gel (**FIG. 18C, panel A**), is denatured and migrates together with other

free molecules of target ssDNA strand, while forward strand becomes invisible due to limited amplification.

[001324] Compared to standard asymmetric PCR, which has relatively low yield and does not allow to distinguish target and non-target strands on denaturing gel, SUPRA delivers different lengths of target and non-target that can be purified on both native gel and denaturing gels. Compared to unequal primer length PCR, which uses lengthy Urea-PAGE protocol and produces only dsDNA, SUPRA has less dsDNA and free target ssDNA can be cut even from native gel if yield is not critical.

[001325] SUPRA has been used in the oligonucleotide probe library enrichment methods provided by the invention. The method is robust. In the presence of enrichment buffer, target/non-target molecules, proteins, exosomes/microvesicles, PEG and other components, SUPRA provides high quality and quantity of the ssDNA oligonucleotide library.

[001326] References:

[001327] 1. Comparison of different methods for generation of single-stranded DNA for SELEX processes. Anal. Bioanal. Chem. 2012, 404, 835–842.

[001328] 2. Upgrading SELEX Technology by Using Lambda Exonuclease Diogestion for Single-Straded DNA Generation. Molecules 2010, 15, 1-11.

[001329] 3. Tang, K.; Fu, D.J.; Julien, D.; Braun, A.; Cantor, C.R.; Koster, H. Chip-based genotyping by mass spectrometry. Proc. Natl. Acad. Sci. USA 1999, 96, 10016–10020.

[001330] 4. Kuypers, A.W.; Linssen, P.C.; Willems, P.M.; Mensink, E.J. On-line melting of double-stranded DNA for analysis of single-stranded DNA using capillary electrophoresis. J. Chromatogr. B Biomed. Appl. 1996, 675, 205–211.

[001331] 5. Higuchi, R.G.; Ochman, H. Production of single-stranded DNA templates by exonuclease digestion following the polymerase chain reaction. Nucleic Acids Res. 1989, 17, 5865.

[001332] 6. Jones, L.A.; Clancy, L.E.; Rawlinson, W.D.; White, P.A. High-affinity aptamers to subtype 3a hepatitis C virus polymerase display genotypic specificity. Antimicrob. Agents Chemother. 2006, 50, 3019–3027.

[001333] 7. S. S. Oh, K. Ahmads, M. Cho, Y. Xiao, H. T. Soh, "Rapid, Efficient Aptamer Generation: Kinetic-Challenge Microfluidic SELEX," presented in the 12th Annual UC Systemwide Bioengineering Symposium, Jun. 13~15, 2011, Santa Barbara, U.S.A

[001334] 8. Gyllensten, U.B.; Erlich, H.A. Generation of single-stranded DNA by the polymerase chain reaction and its application to direct sequencing of the HLA-DQA locus. Proc. Natl. Acad. Sci. USA 1988, 85, 7652–7656.

[001335] 9. Wu, L.; Curran, J.F. An allosteric synthetic DNA. Nucleic Acids Res. 1999, 27, 1512–1516.

[001336] 10. Espelund, M.; Stacy, R.A.; Jakobsen, K.S. A simple method for generating single-stranded DNA probes labeled to high activities. Nucleic Acids Res. 1990, 18, 6157–6158.

[001337] 11. A. Paul, M. Avci-Adali, G. Ziemer, H.P. Wendel. Streptavidin-coated magnetic beads for DNA strand separation implicate a multitude of problems during cell-SELEX. Oligonucleotides 2009, 19, 243–254.

[001338] 12. Williams K., Bartel D. PCR product with strands of unequal length. Nucleic Acids Research, 1995, Vol. 23, No. 20.

[001339] 13. Indirect purification method provides high yield and quality ssDNA sublibrary for potential aptamer selection. Anal. Biochem. 2015, online available.

Example 35: Anti-C1q oligonucleotides

[001340] This Example presents identification of anti-C1q aptamers. In this Example, biotinylated forms of the oligonucleotides identified in **Example 27** were synthesized and immobilized to streptavidin beads and incubated with the PEG precipitated plasma fraction. Proteins that immunoprecipitated with oligonucleotides from SEQ ID NO. 1472-1486 were separated by SDS-PAGE. Specific bands were excised from the gel and proteins therein identified by in-gel trypsin digestions and LC-MS/MS such as in **Example 23** above.

[001341] Immunoprecipitation of specific oligonucleotides conjugated to magnetic beads was performed to identify oligonucleotide targets. Several oligonucleotides were identified that recognized C1q and subunits. The sequences of the variable regions of identified anti-C1q aptamers include 5′-ACTATAGAACAATCCACCGCTTTAAATCAACTATCTTA (SEQ ID NO. 1472), also known by the identifier 83S-B1; 5′-TACCCCTGACAATCCTCGCGCCGAGGCCTCCATCCTGA (SEQ ID NO. 1475), also known by the identifier 83S-B4; 5′-GATTTTAAAACCCTTGCACCTGATTGTGCCAATCCA (SEQ ID NO. 1477), also known by the identifier 83S-B6; and 5′-AACCCAATTCACATACACTCTCACCCCCACTAAACA (SEQ ID NO. 1482), also known by the identifier 83S-B11. Of note, the variable regions of 83S-B1 and 83S-B4 share the motif 5′-ACAATCC (SEQ ID NO. 4151), suggesting that this may comprise a consensus sequence that facilitates binding to C1q. Further as noted below, an oligonucleotide comprising the reverse complement of this putative consensus sequence did not bind C1q. Several variants of the anti-C1q aptamers were also identified in the experiments of **Example 27**. The sequences of the variable regions thereof are shown in **Table 44**. As above, the full length oligonucleotide probes in **Table 44** each had a 5' region consisting of the sequence (5′-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131)) and a 3' region consisting of the sequence (5′-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132)). **Table 44** shows the variable region that is positioned between the 5' and 3' regions of the sequences.

Table 44: Anti-C1q Variable Regions

Variable Region	SEQ ID NO.
83S-B1 (SEQ ID NO. 1472) and variants below	
ACTATAGAACAATCCACCGCTTTAAATCAACTATCTTA	1472
ACTATAGAACAATCCACCGCTTTAAATCAACTATCTGA	4152
ACTATAGAACAATCCACCGCTTTAAATCAACTATCTCA	4153
ACTATAGAACAATCCACCGCTTTAAATCAACTATCTAA	4154

ACTATAGAACAATCCACCGCTTTAAATAAACTATCTGA	4155
ACTATAGAACAATCCACCTCTTTAAATCAACTATCTGA	4156
ACTATAGAACAATCCACCGCTTTAAATCAACTATCCGA	4157
AATATAGAACAATCCACCGCTTTAAATCAACTATCTGA	4158
ACTATAGAACAATCCACCGCTTTAAATCAACTATCCTA	4159
ACTATAGAACAATCCACCTCTTTAAATCAACTATCTTA	4160
ACTATAGAACAATCCCCCGCTTTAAATCAACTATCTGA	4161
ACTATAGAACAATCCGCCGCTTTAAATCAACTATCTGA	4162
ACTATAGAACAATCCTCCGCTTTAAATCAACTATCTGA	4163
ACTATATAACAATCCACCGCTTTAAATCAACTATCTGA	4164
83S-B4 (SEQ ID NO. 1475) and variants below	
TACCCCTGACAATCCTCGCGCCGAGGCCTCCATCCTGA	1475
TACCCCTGACAATCCTCGCGCCGAGGCCTCCATCCTCA	1485
TACCCCTGACAATCCTCGCGCCGAGGCCTCCATCCTAA	4165
TACCCCTGACAATCCTCGCGCCGAGGCCTCCATCCTAA	4166
83S-B6 (SEQ ID NO. 1477) and variants below	
GATTTTAAAACCCTTGCACCTGATTGTGCCAATCCA	1477
GATTTTAAAACCCTTGCACCTGATTGTGCCAATCAA	4167
GATTTTAAAACCCTTGCACCTGATTGTGCCAATCGA	4168
GATTTTAAAACCCTTGCACCTGATTGTGCCAATCTA	4169
83S-B11 (SEQ ID NO. 1482) and variants below	
AACCCAATTCACATACACTCTCACCCCCACTAAACA	1482
AACCCAATTCACATACACTCTCACCCTCACTAAACA	4170
AACCCAATTCACATACACTCTCACCCTCACTAAAAA	4171
AACCCAATTCACATACACTCTCACCCTCACTAAAGA	4172
AACCCAATTCACATACACTCTCACCCTCACTAAATA	4173
AACCCAATTCACATACACTCTCACCCTCACTAACCA	4174
AACCCAATTCACATACACTCTCACCCTCACTAACAA	4175
AACCCAATTCACATACACTCTCTCCCTCACTAAACA	4176
AACCCAATTCACATACACTCTCCCCCTCACTAAACA	4177
AACCCAATTCACATACACTCTCGCCCTCACTAAACA	4178
AACCCAATTCACATACTCTCTCACCCTCACTAAACA	4179
AACCCAATTCACATAAACTCTCACCCTCACTAAACA	4180
AACCCAATCCACATACACTCTCACCCTCACTAAACA	4181
AACCCAATTCACATACACTCTCACCCTCACTAAACC	4182
AACCCAATTCACATACGCTCTCACCCTCACTAAACA	4183
AACCCAATTCACATACACTCTCTCCCTCACTAAAAA	4184
AACCCAATTCACATACCCTCTCACCCTCACTAAACA	4185
AACCCAATTCACATACACTCTCGCCCTCACTAAAAA	4186
AACCCAATTCACATACACTCTTACCCTCACTAAACA	4187
AACCCAATACACATACACTCTCACCCTCACTAAACA	4188
AACCCAATTCACATACACTCTCACCCTCTCTAAACA	4189
AACCCAATTCACATACACTCTCCCCCTCACTAAAAA	4190
AACCCAATTCACATACCCTCTCACCCTCACTAAAAA	4191
AACCCAATTCACATACACTCTCACCCTCACTAAACT	4192
AACCCAATTCACATACACCCTCACCCTCACTAAACA	4193
AACCCAATTCACATACTCTCTCACCCTCACTAAAAA	4194
AACCCAATGCACATACACTCTCACCCTCACTAAACA	4195
AGCCCAATTCACATACACTCTCACCCTCACTAAACA	4196
AACCCAATTCACATACACTCTAACCCTCACTAAACA	4197
AACCCAAATCACATACACTCTCACCCTCACTAAACA	4198
AACCCAATTCACATACACTCTCACCCTCACTAAAAT	4199
AACCCAATTCACATACACTCTCTCCCTCACTAAAGA	4200
AACCCAATTCTCATACACTCTCACCCTCACTAAACA	4201

ATCCCAATTCACATACACTCTCACCCTCACTAAACA	4202
AACCCAACTCACATACACTCTCACCCTCACTAAACA	4203
AACCCAATTCACATACACTCTCACCCTCACTAAAAC	4204
AACCCAATTCACATACACTCTCACCCTCACTAACTA	4205
AACCCAATTCACATACACTCTCGCCCTCACTAAAGA	4206
AACCCAATTCACATTCACTCTCACCCTCACTAAACA	4207
AACCAAATTCACATACACTCTCACCCTCACTAAACA	4208
AACCCAATTCACATAAACTCTCACCCTCACTAAAAA	4209
AACCCAATTCACATACACGCTCACCCTCACTAAACA	4210
AACCCAATTCACATACACTCTCACACTCACTAAACA	4211
AACCCAATTCACATACACTCTCACCATCACTAAACA	4212
AACCCAATTCACATACACTCTCACCCTCACTAACGA	4213
AACCCAATTCACATACACTCTCACCCTCGCTAAACA	4214
AACCCAATTCACATACACTCTCCCCCTCACTAAAGA	4215
AACCCAATTCGCATACACTCTCACCCTCACTAAACA	4216
AACCCAATTCACATACACTCACACCCTCACTAAACA	4217
AACCCAATTCACATACACTCCCACCCTCACTAAACA	4218
AACCCAATTCACATACACTCTCACCCTCACTAAACG	4219
AACCCAATTCACATACTCTCTCACCCTCACTAAAGA	4220
AACCCAATTCACATACTCTCTCACCCTCACTAAATA	4221
AACCCAGTTCACATACACTCTCACCCTCACTAAACA	4222
AAACCAATTCACATACACTCTCACCCTCACTAAACA	4223
AACACAATTCACATACACTCTCACCCTCACTAAACA	4224
AACCCAACTCACATACACTCTCACCCTCACTAAAAA	4225
AACCCAAGTCACATACACTCTCACCCTCACTAAACA	4226
AACCCAATCCACATACACTCTCACCCTCACTAAAAA	4227
AACCCAATTAACATACACTCTCACCCTCACTAAACA	4228
AACCCAATTCACAAACACTCTCACCCTCACTAAACA	4229
AACCCAATTCACACACACTCTCACCCCCACTAAACA	4230
AACCCAATTCACACACACTCTCACCCTCACTAAACA	4231
AACCCAATTCACACACACTCTCCCCCTCACTAAACA	4232
AACCCAATTCACAGACACTCTCACCCTCACTAAACA	4233
AACCCAATTCACATACAATCTCACCCTCACTAAACA	4234
AACCCAATTCACATACACACTCACCCTCACTAAACA	4235
AACCCAATTCACATACACCCTCACCCTCACTAAAAA	4236
AACCCAATTCACATACACTATCACCCTCACTAAAAA	4237
AACCCAATTCACATACACTATCACCCTCACTAAACA	4238
AACCCAATTCACATACACTCCCACCCTCACTAAAAA	4239
AACCCAATTCACATACACTCGCACCCTCACTAAACA	4240
AACCCAATTCACATACACTCTCAACCTCACTAAACA	4241
AACCCAATTCACATACACTCTCACACTCACTAAAAA	4242
AACCCAATTCACATACACTCTCACCATCCCTAAACA	4243
AACCCAATTCACATACACTCTCACCCACACTAAACA	4244
AACCCAATTCACATACACTCTCACCCGCACTAAACA	4245
AACCCAATTCACATACACTCTCACCCTAACTAAACA	4246
AACCCAATTCACATACACTCTCACCCTCAATAAACA	4247
AACCCAATTCACATACACTCTCACCCTCACCAAACA	4248
AACCCAATTCACATACACTCTCACCCTCACTAAAAG	4249
AACCCAATTCACATACACTCTCACCCTCACTAAAGC	4250
AACCCAATTCACATACACTCTCACCCTCACTAAAGT	4251
AACCCAATTCACATACACTCTCACCCTCACTAAATC	4252
AACCCAATTCACATACACTCTCACCCTCACTAAATT	4253
AACCCAATTCACATACACTCTCACCCTCACTAAGAA	4254
AACCCAATTCACATACACTCTCACCCTCACTAAGCA	4255

AACCCAATTCACATACACTCTCACCCTCACTAAGGA	4256
AACCCAATTCACATACACTCTCACCCTCACTAAGTA	4257
AACCCAATTCACATACACTCTCACCCTCACTAATAA	4258
AACCCAATTCACATACACTCTCACCCTCACTAATCA	4259
AACCCAATTCACATACACTCTCACCCTCACTAATGA	4260
AACCCAATTCACATACACTCTCACCCTCACTAATTA	4261
AACCCAATTCACATACACTCTCACCCTCACTACACA	4262
AACCCAATTCACATACACTCTCACCCTCACTAGACA	4263
AACCCAATTCACATACACTCTCACCCTCACTATACA	4264
AACCCAATTCACATACACTCTCACCCTCACTCAACA	4265
AACCCAATTCACATACACTCTCACCCTCACTGAACA	4266
AACCCAATTCACATACACTCTCACCCTCACTTAACA	4267
AACCCAATTCACATACACTCTCACCCTCATTAAACA	4268
AACCCAATTCACATACACTCTCACCCTCCCTAAAAA	4269
AACCCAATTCACATACACTCTCACCCTCCCTAAACA	4270
AACCCAATTCACATACACTCTCACCCTCCCTAAAGA	4271
AACCCAATTCACATACACTCTCACCCTCCCTAAATA	4272
AACCCAATTCACATACACTCTCACCCTCGCTAAAAA	4273
AACCCAATTCACATACACTCTCACCCTCTCTAAAAA	4274
AACCCAATTCACATACACTCTCACCCTGACTAAACA	4275
AACCCAATTCACATACACTCTCACCCTTACTAAACA	4276
AACCCAATTCACATACACTCTCACCTTCACTAAACA	4277
AACCCAATTCACATACACTCTCACCTTCTCTAAACA	4278
AACCCAATTCACATACACTCTCACGCTCACTAAACA	4279
AACCCAATTCACATACACTCTCACTCTCACTAAAAA	4280
AACCCAATTCACATACACTCTCACTCTCACTAAACA	4281
AACCCAATTCACATACACTCTCAGCCTCACTAAACA	4282
AACCCAATTCACATACACTCTCATCCTCACTAAACA	4283
AACCCAATTCACATACACTCTCCCCCTCACTAAATA	4284
AACCCAATTCACATACACTCTCGCCCTCACTAAATA	4285
AACCCAATTCACATACACTCTCTCCCTCACTAAATA	4286
AACCCAATTCACATACACTGTCACCCTCACTAAACA	4287
AACCCAATTCACATACACTTTCACCCTCACTAAACA	4288
AACCCAATTCACATACATTCTCACCCTCACTAAACA	4289
AACCCAATTCACATACCCTCTCACCCTCACTAAAGA	4290
AACCCAATTCACATACGCTCTCACCCTCACTAAAAA	4291
AACCCAATTCACATACGCTCTCACCCTCACTAAAGA	4292
AACCCAATTCACATATACTCTCACCCTCACTAAAAA	4293
AACCCAATTCACATATACTCTCACCCTCACTAAACA	4294
AACCCAATTCACATCCACTCTCACCCTCACTAAACA	4295
AACCCAATTCACATGCACTCTCACCCTCACTAAACA	4296
AACCCAATTCACCTACACTCTCACCCTCACTAAAAA	4297
AACCCAATTCACCTACACTCTCACCCTCACTAAACA	4298
AACCCAATTCACGTACACTCTCACCCTCACTAAACA	4299
AACCCAATTCACTTACACTCTCACCCTCACTAAACA	4300
AACCCAATTCAGATACACTCTCACCCTCACTAAACA	4301
AACCCAATTCATATACACTCTCACCCTCACTAAAAA	4302
AACCCAATTCATATACACTCTCACCCTCACTAAACA	4303
AACCCAATTCCCATACACTCTCACCCTCACTAAACA	4304
AACCCAATTTACATACACTCTCACCCTCACTAAACA	4305
AACCCACTTCACATACACTCTCACCCTCACTAAACA	4306
AACCCATTTCACATACACTCTCACCCTCACTAAACA	4307
AACCCCATTCACATACACTCTCACCCTCACTAAAAA	4308
AACCCCATTCACATACACTCTCACCCTCACTAAACA	4309

AACCCGATTCACATACACTCTCACCCTCACTAAACA	4310
AACCCTATTCACATACACTCTCACCCTCACTAAAAA	4311
AACCCTATTCACATACACTCTCACCCTCACTAAACA	4312
AACCGAATTCACATACACTCTCACCCTCACTAAACA	4313
AACCTAATTCACATACACTCTCACCCTCACTAAAAA	4314
AACCTAATTCACATACACTCTCACCCTCACTAAACA	4315
AACTCAATTCACATACACTCTCACCCTCACTAAACA	4316
AATCCAATTCACATACACTCTCACCCTCACTAAACA	4317
ACCCCAATTCACATACACTCTCACCCTCACTAAAAA	4318
ACCCCAATTCACATACACTCTCACCCTCACTAAACA	4319
AGCCCAATTCACATACACTCTCACCCTCACTAAAAA	4320
ATCCCAATTCACATACACTCTCACCCTCACTAAAAA	4321
CACCCAATTCACATACACTCTCACCCTCACTAAAAA	4322
CACCCAATTCACATACACTCTCACCCTCACTAAACA	4323
GACCCAATTCACATACACTCTCACCCTCACTAAACA	4324
TACCCAATTCACATACACTCTCACCCTCACTAAACA	4325

[001342] Specificity for C1q as a target was performed by immunoprecipitation of C1q depleted or spiked serum and plasma samples. SDS-PAGE analysis of the immunoprecipitation products demonstrated that bands corresponding to C1q subunits (i.e., A, B, C) were only present in the samples spiked with C1q. As a control, oligonucleotides comprising the reverse complements to SEQ ID NOs. 1472 and 1482 were synthesized. For example, the construct comprising the reverse complement to SEQ ID NOs. 1472 comprised the variable region sequence 5′-TTAAGATAGTTGATTTAAAGCGGTGGATTGTTCTATAGT (SEQ ID NO. 4326).

[001343] **FIG. 24A** shows a silver-stained reducing sodium dodecyl sulfate-polyacrylamide (SDS-PA) gel of immunoprecipitated proteins from plasma samples with the indicated oligonucleotides. In the figure, B1 comprises the variable region SEQ ID NO. 1472, B11 comprises the variable region SEQ ID NO. 1482, and L4 and L15 comprises variable regions without specific recognition of C1q. The individual biotinylated oligonucleotides were immobilized on streptavidin magnetic beads, contacted with PEG-precipitated plasma isolates from a pool of healthy donors, and the beads were isolated. Proteins that immunoprecipitated with the beads are shown on the gel. Controls included the reverse complements of B1 and B11 (lanes B1rc and B11rc, respectively) and a control using beads without oligonucleotides (Lane "NO"). Dashed arrows indicate pulled down proteins C1qA, C1qB, and C1qC. The dotted arrow indicates the heavy chain of IgM (IgMHC). The band labeled "SA" is streptavidin and the lane kDa is a protein ladder. As shown in **FIG. 24A**, specific bands indicated by the dashed arrows were observed in the B1 and B11 immunoprecipitation experiments. These bands were excised and analyzed by LC-MS/MS. This analysis revealed the complement protein complex C1q as the target of sequences B1 and B11, and the figure indicates the positions of C1q subunits A, B and C. In contrast, reverse complement versions of B1 and B11 (i.e., B1rc, B11rc) did not immunoprecipitate C1q. Also shown in the figure, IgM co-precipitated with C1q. Without being bound by theory, these findings are consistent with previous observations where IgM was found to remain complexed with C1q in sera that were precipitated with PEG. *See* Krieger, et al. Characterization of immune complexes detected by the 125I-C1q binding assay in

breast cancer. Clin Immunol Immunopathol 46, 14-23 (1988); which publication is incorporated by reference herein in its entirety.

[001344] To further investigate the interaction of aptamer B1 and B11 with C1q, we performed a filter retention assay. Radioactively labelled B1 and B11 were incubated with increasing concentrations of purified C1q and subsequently passed through a nitrocellulose membrane. The amount of oligonucleotides retained on membrane-immobilized C1q was then quantified by autoradiography. Results are shown in **FIG. 24B**. These experiments demonstrate that B1 and B11 exhibit concentration dependent binding to C1q, whereas B11rc did not. Similar results were obtained with B11 and B11rc in an Enzyme-Linked Oligonucleotide Assay (ELONA; *see* US Patent 5789163, which patent is incorporated herein in its entirety). See **FIG. 24C**. The figure shows ELONA analysis of B11 (circles) at indicated concentrations and a fixed C1q concentration (0.625 nM). As a control, the reverse complement of B11, B11RC, was used (squares). "No aptamer" control (triangles) shows low background binding of detector Streptavidin-HRP. B11 specifically binds C1q (estimated K_D around 40 nM). This binding behaviour indicates that B1 and B11 specifically interact with C1q with high affinity.

[001345] A second control comprised the reverse complement to the consensus motif (*i.e.*, SEQ ID NO. 4151). This construct comprised the variable region sequence 5′-TACTATAGATGGATTGTCCGCTTTAAATCAACTATCTTA (SEQ ID NO. 4327) and did not immunoprecipitate C1q from plasma.

[001346] **Table 45** lists proteins that were identified in the immunoprecipitation with B1 (*i.e.*, SEQ ID NO. 1472) using PSMs (peptide spectrum matches) sorted by area. The proteins in the list were observed only in the immunoprecipitation with B1 (i.e., not observed with negative controls) or at levels at least 2-fold over negative controls. The mass spectrometry results were obtained from a single band that was run a short distance in a gel in order to clean up the sample. Accession numbers in the table are from the UniProt database (www.uniprot.org). "GN=" is followed by the gene name. As seen in **Table 45**, C1q subunits A-C were prominent in the results. IgM was also observed at high levels. Without being bound by theory, this may be due to complexes between C1q and immunoglobulins as noted above. Complement C4b-binding protein alpha chain was also observed at 10-fold higher levels versus controls.

Table 45: Oligonucleotide probe targets

Accession number	Protein name
P02747	Complement C1q subcomponent subunit C GN=C1QC
P02746	Complement C1q subcomponent subunit B GN=C1QB
P01871	Ig mu chain C region GN=IGHM
P04220	Ig mu heavy chain disease protein
P02745	Complement C1q subcomponent subunit A GN=C1QA
P01834	Ig kappa chain C region GN=IGKC
P0CG05	Ig lambda-2 chain C regions GN=IGLC2
B9A064	Immunoglobulin lambda-like polypeptide 5 GN=IGLL5
P01857	Ig gamma-1 chain C region GN=IGHG1
P01860	Ig gamma-3 chain C region GN=IGHG3
P01859	Ig gamma-2 chain C region GN=IGHG2

P01766	Ig heavy chain V-III region BRO
O43866	CD5 antigen-like GN=CD5L
P01877	Ig alpha-2 chain C region GN=IGHA2
P01617	Ig kappa chain V-II region TEW
P01767	Ig heavy chain V-III region BUT
P06310	Ig kappa chain V-II region RPMI 6410
P01768	Ig heavy chain V-III region CAM
P01779	Ig heavy chain V-III region TUR
P01620	Ig kappa chain V-III region SIE
P01606	Ig kappa chain V-I region OU
P04208	Ig lambda chain V-I region WAH
P18135	Ig kappa chain V-III region HAH
P01825	Ig heavy chain V-II region NEWM
P01591	Immunoglobulin J chain GN=IGJ
P01598	Ig kappa chain V-I region EU
P01605	Ig kappa chain V-I region Lay
P04433	Ig kappa chain V-III region VG (Fragment)
P01610	Ig kappa chain V-I region WEA
Q8WZ74	Cortactin-binding protein 2 GN=CTTNBP2
P01781	Ig heavy chain V-III region GAL
P04003	C4b-binding protein alpha chain GN=C4BPA
P00739	Haptoglobin-related protein GN=HPR
P06312	Ig kappa chain V-IV region (Fragment) GN=IGKV4-1
P00738-2	Isoform 2 of Haptoglobin GN=HP
P01625	Ig kappa chain V-IV region Len
P01717	Ig lambda chain V-IV region Hil
Q04695	Keratin, type I cytoskeletal 17 GN=KRT17
Q96PE2	Rho guanine nucleotide exchange factor 17 GN=ARHGEF17
P04211	Ig lambda chain V region 4A
P05090	Apolipoprotein D GN=APOD
P34096	Ribonuclease 4 GN=RNASE4
P01714	Ig lambda chain V-III region SH
O14490-3	Isoform 3 of Disks large-associated protein 1 GN=DLGAP1
P04196	Histidine-rich glycoprotein GN=HRG
P35030-5	Isoform 5 of Trypsin-3 GN=PRSS3
P04430	Ig kappa chain V-I region BAN
P00748	Coagulation factor XII GN=F12
P06727	Apolipoprotein A-IV GN=APOA4
P02655	Apolipoprotein C-II GN=APOC2
P01611	Ig kappa chain V-I region Wes
Q569G3	Uncharacterized protein C5orf47 GN=C5orf47
Q9Y490	Talin-1 GN=TLN1
Q5D862	Filaggrin-2 GN=FLG2
P01876	Ig alpha-1 chain C region GN=IGHA1
P01703	Ig lambda chain V-I region NEWM
P02768	Serum albumin GN=ALB
P10909-3	Isoform 3 of Clusterin GN=CLU
P60709	Actin, cytoplasmic 1 GN=ACTB
P04114	Apolipoprotein B-100 GN=APOB
P0C0L4-2	Isoform 2 of Complement C4-A GN=C4A
P0C0L5	Complement C4-B GN=C4B
Q9H4B7	Tubulin beta-1 chain GN=TUBB1
P02675	Fibrinogen beta chain GN=FGB

P01024	Complement C3 GN=C3
P35579	Myosin-9 GN=MYH9
P21333-2	Isoform 2 of Filamin-A GN=FLNA
P02679-2	Isoform Gamma-A of Fibrinogen gamma chain GN=FGG
P00698	Lysozyme C OS=Gallus gallus GN=LYZ
P02671	Fibrinogen alpha chain GN=FGA
P02647	Apolipoprotein A-I GN=APOA1
P68363-2	Isoform 2 of Tubulin alpha-1B chain GN=TUBA1B
Q14624	Inter-alpha-trypsin inhibitor heavy chain H4 GN=ITIH4
P00747	Plasminogen GN=PLG
P02649	Apolipoprotein E GN=APOE
Q9Y2P0	Zinc finger protein 835 GN=ZNF835
Q96IY4	Carboxypeptidase B2 GN=CPB2
P08519	Apolipoprotein(a) GN=LPA
P04264	Keratin, type II cytoskeletal 1 GN=KRT1
P35908	Keratin, type II cytoskeletal 2 epidermal GN=KRT2
P02751-10	Isoform 10 of Fibronectin GN=FN1
P01023	Alpha-2-macroglobulin GN=A2M
P01009-2	Isoform 2 of Alpha-1-antitrypsin GN=SERPINA1
P05160	Coagulation factor XIII B chain GN=F13B
P04004	Vitronectin GN=VTN
Q86YZ3	Hornerin GN=HRNR
P13645	Keratin, type I cytoskeletal 10 GN=KRT10
P13647	Keratin, type II cytoskeletal 5 GN=KRT5
Q03591	Complement factor H-related protein 1 GN=CFHR1
P08603	Complement factor H GN=CFH
P35527	Keratin, type I cytoskeletal 9 GN=KRT9
P00488	Coagulation factor XIII A chain GN=F13A1

[001347] The oligonucleotide sequences in **Table 44** can be futher modified to assess the effects of various subsequences and structural conformation on the ability to bind C1q. Exemplary rationally designed modifications are shown in **Table 46**. For example, the modifications can be based on the secondary structure of the oligonucleotides as estimated using mFold. The sequences in the table may comprise a modification such as a 5' Biotin motif. The Name column indicates the sequence and/or modications thereof. For example, "83s B1 Full" (SEQ ID NO. 4328) consists of the 83s B1 sequence (SEQ ID NO. 1472) shown above with 5' and 3' regions (i.e., SEQ ID NOs. 131 and 132, respectively). Beneath are truncations ("TrXX-YY," where XX is the start position of the truncation and YY is the end position of the truncation as compared to the parent), substitutions (e.g., T(23)G indicating that T at position 23 of the parent sequence is modified to G), and combinations thereof. "RC" indicates reverse complement sequence, in this case for use as a negative control. The italicized and underlined sequences highlight the 5′-ACAATCC consensus sequence.

Table 46: Modified anti-C1q oligonucleotide probe targets

Name	Sequence	SEQ ID NO.
83s B1 Full	CTAGCATGACTGCAGTACGTACTATAGA*ACAATCC*ACCGCTTTAAATCA ACTATCTTACTGTCTCTTATACACATCTGACGCTGCCGACGA	4328
83s B1_Tr17-62	ACGTACTATAGAACAATCCACCGCTTTAAATCAACTATCTTACTGT	4329

83s B1_Tr19-59	GTACTATAGAACAATCCACCGCTTTAAATCAACTATCTTAC	4330
83s B1_T(23)G	CTAGCATGACTGCAGTACGTAC**G**ATAGAACAATCCACCGCTTTAAATCA ACTATCTTACTGTCTCTTATACACATCTGACGCTGCCGACGA	4331
83s B1_Tr17-62T(23)G	ACGTAC**G**ATAGAACAATCCACCGCTTTAAATCAACTATCTTACTGT	4332
83s B1_T(57)G	CTAGCATGACTGCAGTACGTACTATAGA*ACAATCCA*CCGCTTTAAATCA ACTATCT**G**ACTGTCTCTTATACACATCTGACGCTGCCGACGA	4333
83s B1_T(57)G_Tr24-76	ATAGA*ACAATCCA*CCGCTTTAAATCAACTATCT**G**ACTGTCTCTTATACA CATC	4334
83S-B11 Full	CTAGCATGACTGCAGTACGTAACCCAATTCACATACACTCTCACCCCCA CTAAACACTGTCTCTTATACACATCTGACGCTGCCGACGA	4335
B11_C(47)T	CTAGCATGACTGCAGTACGTAACC*CAAT**T**CA*CATACACTCTCACCC**T**CA CTAAACACTGTCTCTTATACACATCTGACGCTGCCGACGA	4336
B11_Tr19-35	GTAACCCAATTCACATAC	4337
B11 Tr_G18_19-35_36C	GGTAACCCAATTCACATACC	4338
83s B1RC (Negative Control)	CTAGCATGACTGCAGTACGTTAAGATAGTTGATTTAAAGCGGTGGATTG TTCTATAGTCTGTCTCTTATACACATCTGACGCTGCCGACGA	4339
83S-B4 Full	CTAGCATGACTGCAGTACGTTACCCCTG*ACAATCC*TCGCGCCGAGGCCT CCATCCTGACTGTCTCTTATACACATCTGACGCTGCCGACGA	4340
83s-B14 Full	CTAGCATGACTGCAGTACGTTACCCCTG*ACAATCC*TCGCGCCGAGGCCT CCATCCTCACTGTCTCTTATACACATCTGACGCTGCCGACGA	4341
83s-B6 Full	CTAGCATGACTGCAGTACGTGATTTTAAAACCCTTGCACCTGATTGTGC *CAATCCA*CTGTCTCTTATACACATCTGACGCTGCCGACGA	4342

[001348] In further analysis, C1q was immunoprecipitated from female plasma samples using the 83s B1 aptamer (or its negative control). For sequences, see **Table 46** above. C1q was eluted with TFA/Urea +/- acetonitrile then detected by either an in-solution trypsin digestion followed by mass spectrometry or SDS-PAGE stained for total protein as in **FIG. 24A** (gel not shown). Mass Spectrometry was performed using the Q Exactive HF Mass Spectrometer system from ThermoFisher Scientific (Waltham, MA) according to the manufacturer's instructions. Various post-translational modifications (PTM) have been observed for C1q A. For example, the UniProt database lists various oxidation sites (e.g., P45, K48, P54, P57, P73, P79, P85) and galactosyl sites (e.g., K48, K67). Other modifications were reported in Pflieger et al., Analysis of Human C1q by Combined Bottom-up and Top-down Mass Spectrometry, Mol Cell Proteomics. 2010 Apr; 9(4): 593–610, which publication is incorporated by reference herein in its entirety. **Table 47** lists PTMs observed in this study. The modifications in bold italics are previously unreported to our knowledge.

Table 47: Observed C1q A post-translational modifications

Position	Target	Modification	Classification	Sequence Motif	SEQ ID NO.
45	P	Oxidation	Post-translational	PGRRGRpGLKGEQ	4343
48	K	Galactosyl	O-linked glycosylation	RGRPGLkGEQGEP	4344
48	K	Oxidation	Post-translational	RGRPGLkGEQGEP	4345
54	P	Oxidation	Post-translational	KGEQGEpGAPGIR	4346
57	P	Oxidation	Post-translational	QGEPGApGIRTGI	4346

67	K	Galactosyl	O-linked glycosylation	TGIQGLkGDQGEP	4347
73	P	Oxidation	Post-translational	KGDQGEpGPSGNP	4348
75	P	Oxidation	Post-translational	DQGEPGpSGNPGK	4349
79	P	Oxidation	Post-translational	PGPSGNpGKVGYP	4350
81	***K***	***Galactosyl***	***O-linked glycosylation***	***PSGNPGkVGYPGP***	4351
85	P	Oxidation	Post-translational	PGKVGYpGPSGPL	4352
87	***P***	***Oxidation***	***Post-translational***	***KVGYPGpSGPLGA***	4353
106	***P***	***Oxidation***	***Post-translational***	***KGTKGSpGNIKDQ***	4354
126	M	Oxidation	Artefact	IRRNPPmGGNVVI	4355
205	M	Oxidation	Artefact	QVVSGGmVLQLQQ	4356

Example 36: C1q immunoassay and isolation

[001349] This Example illustrates C1q immunoassays using an anti-C1q oligonucleotide. A nucleic acid construct is synthesized comprising an anti-C1q oligonucleotide region, for example, as described in **Example 35** above. The anti-C1q construct comprises a 5' biotin modification to facilitate specific recognition by a desired moiety attached to streptavidin.

[001350] A labeled anti-C1q aptamer construct is contructed. The anti-C1q construct is contacted with fluorescently labeled streptavidin such as a streptavidin - Alexa Fluor® 488 conjugate from Thermo Fisher Scientific, Catalog number: S11223. This creates a fluorescently labeled anti-C1q construct which is used to detect C1q in various immunoassay formats. In one scenario, a biological sample known or suspected to contain C1q is contacted with an ELISA plate. The plate is washed and contacted with the fluorescently labeled anti-C1q construct. The fluorescent signal is read from the wells in the plate, thereby providing an indication of the presence or amount of C1q in the biological sample. In another scenario, a biological sample is directly contacted with the fluorescently labeled anti-C1q construct. The contacted sample is subjected to flow cytometry to detect fluorescent particles of the size of microvesicles, thereby providing an indication of the presence or amount of microvesicle associated C1q in the biological sample. Alternate labels such as disclosed herein or known in the art can be used in such formats.

[001351] Various modifications of the above scenarios are performed. For example, the anti-C1q aptamer is directly labeled with Alexa Fluor during the oligonucleotide synthesis process.

[001352] An immobilized anti-C1q aptamer is contructed. In one scenario, the anti-C1q aptamer construct is contacted with streptavidin conjugated beads. The beads are contacted with a biological sample known or suspected to contain C1q. The beads are precipitated (e.g., by centrifugation or magnetism) and washed. Proteins that precipitate with the beads are analyzed, thereby providing an indication of the presence or amount of C1q and associated proteins in the biological sample. In another scenario, the anti-C1q aptamer construct is contacted with streptavidin agarose resin, e.g., Pierce™ Streptavidin Agarose, Thermo Fisher Scientific Catalog number: 20347 or Pierce™ High Capacity Streptavidin Agarose Thermo Fisher Scientific Catalog number: 20357. The resins are placed in a spin column or chromatography column, respectively. The anti-C1q aptamer is contacted with the resin where it is bound by the streptavidin. A biological sample known or suspected to comprise C1q is allowed to pass through the resin. C1q and associated proteins in the biological sample are retained by the anti-C1q

aptamer within the resin and are then analyzed after elution. In either scenario, if desired, the anti-C1q aptamer is contacted with the biological sample in solution and then the sample is contacted with the beads or resin. C1q and associated proteins are analyzed as above. This modification allows the C1q and aptamer to bind freely in solution prior to aptamer immobilization.

[001353] Various modifications of the above scenarios are performed. For example, the anti-C1q aptamer is directly conjugated to a bead or other desired surface.

[001354] One of skill will appreciate that the anti-C1q aptamer construct can be used in any desired scenario where antibodies are conventionally used. *See, e.g.,* Toh et al., Aptamers as a replacement for antibodies in enzyme-linked immunosorbent assay. Biosens Bioelectron. 2015 Feb 15;64:392-403. doi: 10.1016/j.bios.2014.09.026. Epub 2014 Sep 16; Chen and Yang, Replacing antibodies with aptamers in lateral flow immunoassay. Biosens Bioelectron. 2015 Sep 15;71:230-42. doi: 10.1016/j.bios.2015.04.041. Epub 2015 Apr 14; Guthrie et al, Assays for cytokines using aptamers. Methods. 2006 Apr;38(4):324-30; Romig et al., Aptamer affinity chromatography: combinatorial chemistry applied to protein purification. J Chromatogr B Biomed Sci Appl. 1999 Aug 20;731(2):275-84.

Example 37: C1q Detection in Bodily Fluids

[001355] This Example describes using an anti-C1q oligonucleotide to detect C1q in bodily fluids. A bodily fluid such as blood or a derivative thereof, including without limitation sera or plasma, is obtained from a subject. An assay such as described in **Example 36** is used to detect C1q in the bodily fluid. As desired, such detection may assist in the diagnosis, prognosis or theranosis of a disease or disorder. *See, e.g.,* **Examples 13-14** herein.

[001356] In one scenario, the anti-C1q aptamer is used to capture and quantitate immune complexes in clinical samples based on the ability of C1q to preferentially bind immune complexes versus monomeric immunoglobulins.

Example 38: C1q Inhibition

[001357] This Example describes using an anti-C1q oligonucleotide to inhibit C1q mediated apoptosis.

[001358] Deficiency in the host C1q inhibitor protein, encoded by the SERPING1 gene (also known as C1NH and others), is associated with hereditary angioedema ("hereditary angioneurotic edema"; "HAE"), which comprises swelling due to leakage of fluid from blood vessels into connective tissue. In addition to facial swelling and/or abdominal pain, C1q inhibitor deficiency also predisposes to autoimmune diseases, e.g., lupus erythematosus, and age related macular degeneration. C1q inhibitors CINRYZE® (C1 esterase inhibitor [human]) and RUCONEST® (C1 esterase inhibitor [recombinant]) are approved by the US FDA for HAE symptoms. *See, e.g.,* Bernstein, Hereditary angioedema: a current state-of-the-art review, VIII: current status of emerging therapies. Ann. Allergy Asthma Immunol. 100 (1 Suppl 2): S41–6 (January 2008). A pharmaceutical composition comprising an anti-C1q oligonucleotide of the invention is administered to an HAE patient in sufficient dosage to ameliorate symptoms of HAE.

[001359] As the complement cascade damages cells, C1q inhibitors have potential functionality for diseases other than HAE. *See e.g.,* Caliezi et al, C1-Esterase inhibitor: an anti-inflammatory agent and its potential use in the treatment of diseases other than hereditary angioedema. Pharmacol. Rev. 52 (1): 91–112 (March 2000). During heart attack, lack of oxygen in heart cells causes necrosis in heart cells. Dying heart cells spill their contents in the extracellular environment, which triggers the complement cascade and may increase the damage for the surviving heart cells. A pharmaceutical composition comprising an anti-C1q oligonucleotide of the invention is administered to a heart attack victim in sufficient dosage to ameliorate complement-mediated heart damage. Similarly, a pharmaceutical composition comprising an anti-C1q oligonucleotide of the invention is administered to a patient following organ transplantation to ameliorate delayed graft function.

[001360] The complement cascade is a first defense against non-self cells and is also activated by malignant cells. Thus, complement appears to play a role in cancer surveillance by the immune system. Recently, however, it has been reported that the complement system can also play a role in promoting tumor growth, inflammation and angiogenesis. Thus, complement inhibition has been proposed for cancer treatment. *See, e.g.,* Pio et al, Complement inhibition: a promising concept for cancer treatment, Semin Immunol. 2013 Feb; 25(1): 54–64; Pio et al., The role of complement in tumor growth. Adv Exp Med Biol. 2014;772:229-62. It has further been reported that C1q may act in the tumor microenvironment to promote cancer independently of its complement function. Bulla *et al.* report that C1q may promote cancer cell adhesion, migration and proliferation as well as angiogenesis and metastasis. *See* Bulla et al., C1q acts in the tumour microenvironment as a cancer-promoting factor independently of complement activation. Nat Commun. 2016; 7: 10346. A pharmaceutical composition comprising an anti-C1q oligonucleotide of the invention is administered to a cancer victim in sufficient dosage to treat the cancer in the victim. As desired, the composition is administered in combination with traditional cancer therapeutics or in addition to immune therapies, e.g., cell-based tumor immunotherapies, or antitumor vaccines.

Example 39: Complement cascade initiation

[001361] This Example describes using an anti-C1q oligonucleotide to direct complement mediated cell killing.

[001362] Antibody immunotherapies may direct the killing of target cells through several mechanisms, including without limitation complement-dependent cytotoxicity (CDC), antibody-dependent cellular cytotoxicity (ADCC), apoptosis, and direct growth arrest. *See, e.g.,* Taylor and Lindorfer, The role of complement in mAb-based therapies of cancer. Methods. 2014 Jan 1;65(1):18-27. doi: 10.1016/j.ymeth.2013.07.027. Epub 2013 Jul 22; Rogers *et al.*, Complement in monoclonal antibody therapy of cancer. Immunol Res. 2014 Aug;59(1-3):203-10. doi: 10.1007/s12026-014-8542-z; Zhou *et al.*, The Role of Complement in the Mechanism of Action of Rituximab for B-Cell Lymphoma: Implications for Therapy, The Oncologist September 2008 vol. 13 no. 9 954-966; Di Gaetano *et al.*, Complement activation determines the therapeutic activity of rituximab *in vivo*. J Immunol. 2003 Aug 1;171(3):1581-7.

[001363] The Fc regions of membrane-bound therapeutic antibodies interact with the heterooligomeric Clq complex and activate the classical complement pathway to initiate CDC. A multipartite construct comprising an anti-C1q oligonucleotide of the invention is provided in a pharmaceutical composition. The pharmaceutical composition is administered to a cancer victim in sufficient dosage to treat the cancer in the victim. The construct comprises the anti-C1q oligonucleotide connected to an anti-tumor domain. *See, e.g.,* **FIGs. 23A-D** herein and related discussion. The anti-tumor domain can be to a general tumor antigen or specific to a tumor of a given origin. *See, e.g.,* Section "Anti-target and multivalent oligonucleotides" herein for illustrative targets. The anti-tumor domain may comprise the C10.36 aptamer provided herein (SEQ ID NO. 4357). As desired, the multipartite construct comprising an anti-C1q oligonucleotide is used to direct complement mediated cell killing where cancer cells and tumors have adapted to evade the complement system.

[001364] As noted herein, C1q may play a role in both preventing cancer, e.g., through complement mediated cell killing, and facilitating the growth of cancer, e.g., through cancer cell adhesion, migration and proliferation as well as angiogenesis and metastasis, perhaps in a non-complement fashion. Such disparate roles allow for combination treatment with oligonucleotides of the invention that both inhibit the roles of C1q in promoting cancer while targeting C1q mediated cell killing to cancer cells. A multipartite construct comprising an anti-C1q oligonucleotide of the invention is provided in a pharmaceutical composition. An anti-C1q oligonucleotide of the invention that inhibits non-complement functions of C1q is provided in a pharmaceutical composition. The pharmaceutical compositions are administered to a cancer victim in sufficient dosage to treat the cancer in the victim.

Example 40: Anti-lymphona oligonucleotide probes

[001365] This Example presents aptamer sequences that can be used to target cancer cells. For example, the aptamer can be used to target the C1q aptamers of the invention (see, e.g., **Table 44**) to cancer cells. This Example uses the C10 aptamer from Raddatz et al., Enrichment of cell-targeting and population-specific aptamers by fluorescence-activated cell sorting. Angew Chem Int Ed Engl. 2008;47(28):5190-3; which publication is incorporated by reference herein in its entirety. The C10 aptamer was found to recognize Burkitt Lymphoma cells. Here we have discovered a 36-mer sequence of Aptamer C10, the "C10.36" or "10.36" aptamer, that is responsible for binding Burkitt Lymphoma cells. We reported characterization of the structure of Aptamer 10.36 in Opazo *et al.*, Modular Assembly of Cell-targeting Devices Based on an Uncommon G-quadruplex Aptamer, Molecular Therapy—Nucleic Acids (2015) 4, e251; doi:10.1038/mtna.2015.25; published online 1 September 2015; which publication is incorporated by reference herein in its entirety.

[001366] The oligonucleotide sequences in **Table 48** comprise various combinations of C10.36 linked to anti-C1q aptamers of the invention (*i.e.*, B1 (SEQ ID NO. 1472) and B11(SEQ ID. NO. 1482) derivatives as indicated). These oligonucleotides can be futher modified to assess the effects of various subsequences and structural conformation on the ability to bind cancer cells. For example, the modifications can be based on the secondary structure of the oligonucleotides as estimated using mFold.

Exemplary rationally designed modifications are shown in **Table 48**. The sequences may further comprise desired modification such as a 5' Biotin motif. The "Name" column indicates the sequence and/or modications thereof. For example, "C10.36" is the functional truncation of the C10 aptamer as described. Beneath are truncations ("TrXX-YY," where XX is the start position of the truncation and YY is the end position of the truncation as compared to the parent), substitutions (e.g., T(57)G indicating that T at position 57 of the parent sequence is modified to G), and combinations thereof. The italicized and underlined sequences highlight the 5′-ACAATCC consensus sequence (SEQ ID NO. 4151) described above.

Table 48: C10.36 truncated oligonucleotide and derivatives

Name	Sequence	SEQ ID NO.
C10.36	CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4357
C10.36_B1_T(57)G_Tr24-76	CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAGATAGA*ACAATCC*ACCGCTTTAAATCAACTATCTGACTGTCTCTTATACACATC	4358
B1_T(57)G_Tr24-76_C10.36	ATAGA*ACAATCCA*CCGCTTTAAATCAACTATCT**G**ACTGTCTCTTATACACATCCTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4359
B1_C(29)T_T(57)G_Tr24-76_C10.36	ATAGAACAAT**T**CACCGCTTTAAATCAACTATCTGACTGTCTCTTATACACATCCTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4360
C10.36_83s B1_Tr17-62	CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAGACGTACTATAGA*ACAATCC*ACCGCTTTAAATCAACTATCTTACTGT	4361
83s B1_Tr17-62_C10.36	ACGTACTATAGAACAATCCACCGCTTTAAATCAACTATCTTACTGTCTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4362
C10.36_83s B1_Tr19-59	CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAGGTACTATAGA*ACAATCC*ACCGCTTTAAATCAACTATCTTAC	4363
B1_Tr19-59_C10.36_83s	GTACTATAGA*ACAATCC*ACCGCTTTAAATCAACTATCTTACCTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4364
C10.36_83s B1_Tr17-62T(23)G	CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAGACGTAC**G**ATAGA*ACAATCC*ACCGCTTTAAATCAACTATCTTACTGT	4365
83s B1_Tr17-62T(23)G_C10.36	ACGTAC**G**ATAGA*ACAATCC*ACCGCTTTAAATCAACTATCTTACTGTCTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4366
C10.36_83s_B11_Tr19-35	CTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAGGTAACCCAATTCACATAC	4367
83s_B11_Tr19-35_C10.36	GTAACCCAATTCACATACCTAACCCCGGGTGTGGTGGGTGGGCAGGGGGGTTAG	4368

Example 41: Aptamer 10.36 binds a ribonucleoprotein complex

[001367] We reported that aptamer 10.36 (SEQ ID NO. 4357) forms a G-quadruplex structure and is taken up by Burkitt's Lymphoma (Ramos) cells via a clathrin-mediated endocytotic pathway. Opazo *et al*. This aptamer is taken up by Ramos cells to a greater extent than other lymphoma derived cell lines (e.g., Jurkat, Raji, or P12). Opazo *et al*. In this Example, we report identification of the target of aptamer 10.36.

[001368] A single point mutant in aptamer 10.36, 10.36(G24A) (SEQ ID NO. 4369), disrupts the G-quadruplex structure and results in loss of binding to Ramos cells. Opazo *et al*. This point mutant was

used as a control in the aptamer precipitation experiments in this Example to control for non-specific binding to nucleic acid binding proteins.

[001369] **Preferential Binding to Ramos Cells:**

[001370] A titration of 10.36 and 10.36(G24A) demonstrated binding of 10.36 preferentially to Ramos cells and lack of binding to 10.36(G24A) to either Ramos or Jurkat cells. See **FIG. 25A**. Panels **i-ii** show results of flow cytometry with Ramos cells and panels **iii-iv** show results with Jurkat cells. Panels **i** and **iii** show results with aptamer 10.36 and panels **ii** and **iv** show results with control aptamer 10.36(G24A). To perform these experiments, aptamer stocks were diluted to 10 μM in Hank's Buffered Salt Solution (HBSS), denatured, and allowed to cool to room temperature for 15 minutes. They were further incubated with streptavidin, R-phycoerythrin conjugate (SAPE; at 1 SAPE molecule per 4 aptamer molecules) for 30 minutes at room temperature in the dark, then diluted in the appropriate media (RPMI medium (Gibco)-10% serum, 0.1 mg/mL sheared salmon sperm DNA (sssDNA), 0.1 mg/mL yeast tRNA, 0.01mM dextran sulfate, with or without prior heat inactivation of serum). The cells were washed, counted, and resuspended in media. They were then incubated with diluted aptamers and SAPE in media for 15 min at 37 °C, 5% CO_2. After incubation with aptamers, cells were washed twice with HBSS, then resuspended in HBSS for flow analysis on Apogee flow cytometry platform (Apogee Flow Systems Ltd, Middlesex, England).

[001371] As seen in **FIG. 25A**, SAPE by itself had the same binding pattern as unstained flow experiments under all conditions. Ramos cells had a clear positive population at all 10.36 concentrations with titration effect seen with increasing concentration of 10.36 (i.e., 200 nM to 800 nM, as indicated). See panel **i**. Jurkat cells were dimly positive at the highest concentration of C10.36 aptamer, 800μm. See panel **iii**. All cells were negative for C10.36(G24A) staining. See panels **ii** and **iv**. Slightly brighter staining of Ramos cells was observed in media comprising heat-inactivated serum. Data not shown.

[001372] **Target Identification**:

[001373] Targets of 10.36 were assessed by affinity purification of aptamers bound to Ramos or Jurkat cells with streptavidin Dynabeads® (ThermoFisher, Sci). Proteins eluted from the beads were separated by SDS-PAGE and detected by silver stain. Protocol used as follows:

[001374] 1. Pellet appropriate volume of cells by centrifuging for 5 minutes at room temperature (RT) at 500*xg*.

[001375] 2. Aspirate media.

[001376] 3. Resuspend cells in 500 μL binding buffer (see above for flow cytometry) to provide 2X10^5 cells per reaction.

[001377] 4. Aptamers: dilute aptamer to 10 μM stock and heat denature for 5 min at 95 °C; allow to cool 15 min on bench.

[001378] 5. Dilute aptamer stock to a working stock using binding buffer.

[001379] 6. Add 50μl of aptamer to each well of a 96 well deep plate.

[001380] 7. Add 50μl of cells to the above wells.

[001381] 8. Briefly tap/mix the samples.

[001382] 9. Incubate 15 minutes at 37 °C, 5% CO_2.

[001383] 10. Add 10µL of Dynabeads® MyOne™ Streptavidin C1 to all the wells (Thermo Fisher Scientific).

[001384] 11. Incubate beads with aptamer–cell solution for 30 mins at RT mixing.

[001385] 12. Discard supernatant via magnetic capture with a MagMax™ platform (Thermo Fisher Scientific).

[001386] 13. Wash beads twice with 200µL of 1X HBSS.

[001387] 14. Lyse cells with 200µl lysis buffer (50 mM Tris-HCl, 3 mM $MgCl_2$, 150 mM NaCl, 1% NP40, 0.5% sodium deoxycholate, pH 7.5) for 15 min at RT. Mix on Magmax.

[001388] 15. Wash again with with 200µL of 1X HBSS

[001389] 16. Elute sample with 15µL 0.3%TFA, 6M Urea heat 37 °C for 10min; transfer supernatant to fresh tube then add 5µL of 4X LDS, 2µL of 10X reducing agent

[001390] 17. To remaining beads at 22µL of 1X LDS, 1X reducing agent, boil for 15min at 95 °C, freeze at -20C

[001391] 18. Run 4-12% SDS-PAGE of TFA/Urea eluted samples

[001392] 19. Silver stain gels

[001393] 20. Cut out bands and process to in gel trypsin digestion

[001394] An exemplary gel is shown in **FIG. 25B**. Lanes in the Gel are shown in **Table 49**:

Table 49: Silver staining gel lanes

Lane	Cells	Aptamer
kDa	Molecular weight marker	
1	Ramos	10.36
2	Ramos	10.36(G24A)
3	Ramos	None
4	Jurkat	10.36
5	Jurkat	10.36(G24A)
6	Jurkat	None

[001395] As shown in **FIG. 25B**, few proteins bound to the G24A (lanes 2, 5) or no aptamer (lanes 3, 6) control beads in either Ramos or Jurkat cells. Two duplicate gels were run for 10 min, the entire lane was cut out in one band and subjected to in-gel trypsin digestion and LC-MS/MS. See, e.g., **Example 24**. Sixty-five of 81 identified proteins were identified as differentially binding to Ramos vs. Jurkat cells with at least a 2-fold change, suggesting these proteins are related to the difference in binding between Ramos and Jurkat cells. Of these, 51 were unique to 10.36 bound to Ramos cells. *See* **Table 50**. Accession numbers in the table are from the UniProt database (www.uniprot.org). "GN=" is followed by the gene name. Proteins in the table are sorted by level detected in Ramos cells, from high to low. The most abundant proteins included components of nucleolin containing complexes nucleolin (NCL),

nucleophosmin (NPM1), actin (ACTB), heterogeneous nuclear ribonucleoproteins (HNRNP) (Pinol-Roma, Hovanessian et al.) as well as other nucleoproteins. The table indicates the fold change in Ramos cells as compared to Jurkat cells. "n/d" indicates that the protein was observed in Ramos but not detected in Jurkat cells.

Table 50: Proteins differentially binding to Ramos vs. Jurkat cells

Accession Number	Protein Name	Fold Change (Ramos / Jurkat)
P19338	Nucleolin GN=NCL	3.0
P22626	Cluster of heterogeneous nuclear ribonucleoproteins A2/B1	2.0
P06748	Cluster of Nucleophosmin GN=NPM1	8.0
P60709	Cluster of Actin, cytoplasmic 1	5.3
P22087	rRNA 2'-O-methyltransferase fibrillarin	4.0
P09651	Cluster of Heterogeneous nuclear ribonucleoprotein A1	2.0
Q04837	Single-stranded DNA-binding protein, mitochondrial	5.5
P11142	Heat shock cognate 71 kDa protein	n/d
Q00839	Heterogeneous nuclear ribonucleoprotein U	8.0
P52272	Heterogeneous nuclear ribonucleoprotein M	7.0
Q07955	Serine/arginine-rich splicing factor 1	7.0
P55769	NHP2-like protein 1	n/d
Q12906	Interleukin enhancer-binding factor 3	n/d
P06576	ATP synthase subunit beta, mitochondrial	n/d
P31942	Cluster of Heterogeneous nuclear ribonucleoprotein H3	6.0
Q86V81	THO complex subunit 4	2.0
Q9NR30	Nucleolar RNA helicase 2	n/d
Q96PK6	RNA-binding protein 14	n/d
P37108	Signal recognition particle 14 kDa protein	n/d
P09874	Poly [ADP-ribose] polymerase 1	n/d
P17096	Cluster of High mobility group protein HMG-I/HMG-Y	n/d
Q99729	Heterogeneous nuclear ribonucleoprotein A/B	n/d
P11021	78 kDa glucose-regulated protein	n/d
P23246	splicing factor, proline- and glutamine-rich	n/d
P27694	Replication protein A 70 kDa DNA-binding subunit	n/d
Q13151	Heterogeneous nuclear ribonucleoprotein A0	n/d
P04792	Heat shock protein beta-1	n/d
P31943	Heterogeneous nuclear ribonucleoprotein H	n/d
P51991	Cluster of Heterogeneous nuclear ribonucleoprotein A3	4.0
P02647	Apolipoprotein A-I	n/d
O43390	heterogeneous nuclear ribonucleoprotein r	n/d
P08865	40S ribosomal protein SA	n/d
O14979	Cluster of Heterogeneous nuclear ribonucleoprotein D-like	n/d
Q15365	Cluster of Poly(RC)-binding protein 1	n/d
P26599	Polypyrimidine tract-binding protein 1	n/d
Q14624	Inter-alpha-trypsin inhibitor heavy chain H4	n/d

Q14103	heterogeneous nuclear ribonucleoprotein D0	n/d
P04406	glyceraldehyde-3-phosphate dehydrogenase	n/d
O60506	Heterogeneous nuclear ribonucleoprotein Q	n/d
Q15717	ELAV-like protein 1	n/d
Q8NFH5	Nucleoporin NUP53	n/d
Q9NX24	H/ACA ribonucleoprotein complex subunit 2	2.0
P01009	alpha-1-antitrypsin	n/d
Q14978	nucleolar and coiled-body phosphoprotein 1	n/d
P10809	60 kDa heat shock protein, mitochondrial	n/d
P14866	Heterogeneous nuclear ribonucleoprotein L	n/d
Q13243	Cluster of Serine/arginine-rich splicing factor 5	n/d
Q6RFH5	WD repeat-containing protein 74	n/d
Q9NZI8	Insulin-like growth factor 2 mRNA-binding protein 1	n/d
P46087	Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase	n/d
P18754	Regulator of chromosome condensation	n/d
P11940	Cluster of Polyadenylate-binding protein 1	n/d
P25705	ATP synthase subunit alpha, mitochondrial	n/d
Q15173	Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit beta isoform GN=PPP2R5B	n/d
Q9Y3B9	RRP15-like protein	n/d
Q9Y3Y2	Chromatin target of PRMT1 protein	n/d
P84103	Serine/arginine-rich splicing factor 3	n/d
P30154	Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A beta isoform GN=PPP2R1B	n/d
Q00577	Transcriptional activator protein Pur-alpha	n/d
1A31_HUMAN	HLA class I histocompatibility antigen, A-31 alpha chain GN=HLA-A	n/d
O00479	High mobility group nucleosome-binding domain-containing protein 4	n/d
P62263	40S ribosomal protein S14	n/d
Q16629	serine/arginine-rich splicing factor 7	n/d
P52907	F-actin-capping protein subunit alpha-1	n/d
Q12905	Interleukin enhancer-binding factor 2	n/d

[001396] Aptamer 10.36 is internalized by Ramos cells through a clathrin dependent mechanism at the cell surface. *See* Opazo *et al*. Both NCL and NPM1 have both been shown to be expressed on the cell surface of cancer cells and together regulate K-Ras signaling. See Gervin et al.; Inder et al. Ribonucleoprotein containing structures have been shown to accumulate on the cell surface of apoptotic cells. See Biggiogera et al. for review. To assess whether aptamer 10.36 could bind nucleolin at the cell-surface, aptamer precipitations were conducted in the presence of 80 μM Dynasore®, which inhibits clathrin mediated endocytosis. For methodology, see Opazo et al; Kirchhausen et al. To reduce viscosity of the cellular lysate, endonuclease HL-dsDNase, which specifically cleaves dsDNA but does not degrade the single stranded DNA aptamer (data not shown), was added to the lysis buffer at a final concentration of 4 Units per 100 μl. The number of total proteins identified by LC-MS/MS from gel bands run in

triplicate in the presence of endonuclease HL-dsDNase was much lower and was similar with and without Dynasore®. See **FIG. 25C**. In the figure, lanes are indicated in **Table 51**. Panels **i**, **ii** and **iii** are triplicate gels. Consistent with the flow cytometry results described above, higher levels of protein were detected with Ramos cells and aptamer 10.36 as opposed to Jurkat cells or control aptamer 10.36(G24A).

Table 51: Silver stained gel lanes

Lane	Cells	Aptamer	Dynasore
1	Mol Weight Ladder	n/a	n/a
2	Ramos	10.36	+
3	Ramos	10.36	-
4	Ramos	10.36(G24A)	+
5	Ramos	10.36(G24A)	-
6	Jurkat	10.36	+
7	Jurkat	10.36	-
8	Jurkat	10.36(G24A)	+
9	Jurkat	10.36(G24A)	-

[001397] Eight proteins were differentially bound to 10.36 and Ramos cells and included NCL, FBL, hnRNP C1/C2, SFPQ and DDX21. See **Table 52**. The last column in the table indicates the fold change in Ramos cells (R) as compared to Jurkat cells (J), in the presence (+) or absence (-) of Dynasore. "n/d" indicates that the protein was observed in Ramos but not detected in Jurkat cells. Dynasore inhibits internalization of 10.36 (*see* Opazo et al) but made minimal difference on the proteins we detected under our conditions. NCL, NMP1 and DDX21 have been shown to directly bind G-quadruplex DNA structures. See Arcovito et al.; Girvan et al.; Tosoni et al.; Brazda, et al.; Mendoza et al. SFPQ was originally described as a myoblast cell surface antigen and is a DNA- and RNA binding protein, involved in several nuclear processes. See Gower et al. DDX21 is a nucleolar helicase that can unfold Q-quadruplex structures. See Mendoza et al. To our knowledge this is the first time the translational control proteins SFPQ and DDX21 have been identified co-precipitating with cell surface nucleolin.

Table 52: Proteins differentially binding to Ramos vs. Jurkat cells

Accession Number	Protein Name	Fold Change (R / J) Dynasore + / -
P13645	Cluster of Keratin, type I cytoskeletal 10	7.6 / 1.2
P07910	Heterogeneous nuclear ribonucleoproteins C1/C2 (hnRNP C1/C2) GN=HNRNPC	1.3 / 4.9
P19338	Nucleolin GN=NCL	n/d / n/d
P23246	splicing factor, proline- and glutamine-rich (SFPQ)	2.6 / n/d
Q7L8J4	SH3 domain-binding protein 5-like GN=SH3BP5L	n/d / 4.1
P14923	Junction plakoglobin	n/d / n/d
P22087	rRNA 2'-O-methyltransferase fibrillarin GN=FBL	n/d / n/d
Q9NR30	Nucleolar RNA helicase 2 GN=DDX21	n/d / n/d

[001398] To increase the yield of the surface target proteins or protein complexes and limit endocytotic aptamer internalization, ~1µg of aptamer 10.36 and control aptamer 10.36(G24A) were immobilized to 1µm streptavidin Dynabeads® (ThermoFisher, Sci). Ramos or Jurkat cells were captured from the solution using the beads in the presence of Dynasore® and endonuclease HL-dsDNase. As above, differential detection by LC-MS/MS analysis of digests of triplicate gel bands (not shown) found ribonucleoproteins precipitated with the 10.36 aptamer specifically in Ramos cells. See **Table 53**. Accession numbers in the table are from the UniProt database (www.uniprot.org). "GN=" is followed by the gene name. Proteins in the table are sorted by level detected in Ramos cells, from high to low. The table indicates the fold change in Ramos cells as compared to Jurkat cells. "n/d" indicates that the protein was observed in Ramos but not detected in Jurkat cells.

Table 53: Proteins differentially binding to Ramos vs. Jurkat cells

Accession Number	Protein Name	Fold Change (Ramos / Jurkat)
P16403	Cluster of Histone H1.2	9.2
P19338	Nucleolin	4.4
P07910	Cluster of Heterogeneous nuclear ribonucleoproteins C1/C2	n/d
P23246	splicing factor, proline- and glutamine-rich	10.0
P22626	heterogeneous nuclear ribonucleoproteins A2/B1	14.0
P60709	Cluster of Actin, cytoplasmic 1	7.0
P09651	Heterogeneous nuclear ribonucleoprotein A1	13.5
P22087	rRNA 2'-O-methyltransferase fibrillarin	n/d
Q9NR30	Nucleolar RNA helicase 2	n/d
P55769	NHP2-like protein 1	n/d
Q13151	Heterogeneous nuclear ribonucleoprotein A0	n/d
P62633	Cellular nucleic acid-binding protein	n/d
P31942	Heterogeneous nuclear ribonucleoprotein H3	n/d
P61978	Heterogeneous nuclear ribonucleoprotein K	n/d
P09874	Poly [ADP-ribose] polymerase 1	n/d
P46087	Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase	n/d
P35268	60S ribosomal protein L22	n/d
Q01844	RNA-binding protein EWS	n/d
Q9UKM9	RNA-binding protein Raly	n/d
Q14978-1	nucleolar and coiled-body phosphoprotein 1	n/d
Q86V81	THO complex subunit 4	n/d
P62750	60S ribosomal protein L23a	n/d
P06748	Nucleophosmin	n/d
Q9NZI8	Insulin-like growth factor 2 mRNA-binding protein 1	n/d
P68431	Histone H3.1	n/d
Q9NX24	H/ACA ribonucleoprotein complex subunit 2	n/d
P27694	Replication protein A 70 kDa DNA-binding subunit	n/d
O43390	Cluster of heterogeneous nuclear ribonucleoprotein r	n/d

P67809	Nuclease-sensitive element-binding protein 1	n/d
Q96PK6	RNA-binding protein 14	n/d
Q12906	Interleukin enhancer-binding factor 3	n/d
P37108	Signal recognition particle 14 kDa protein	n/d
Q12905	Interleukin enhancer-binding factor 2	n/d
P11142	Heat shock cognate 71 kDa protein	n/d
Q00839	Heterogeneous nuclear ribonucleoprotein U	n/d
P50914	60S ribosomal protein L14	n/d
P11021	78 kDa glucose-regulated protein	n/d
P31943	Heterogeneous nuclear ribonucleoprotein H	n/d
Q8TEM1	Nuclear pore membrane glycoprotein 210	n/d

[001399] Four proteins were identified as differentially bound in all the experimental conditions tested as shown in **Tables 50**, **52** and **53**: NCL, FBL, SFPQ, and DDX21. Without being bound by theory, these data indicate that aptamer 10.36 binds a ribonucleoprotein complex comprising multiple proteins including without limitation nucleolin, fibrillarin, actin, SFPQ, hnRNPM, hnRNPC1/C2. To examine which proteins 10.36 binds directly as opposed to indirectly as part of a ribonucleoprotein complex, we increased the stringency of salt in the wash buffers from 133 mM to 1M NaCl in the presence of Dynasore. **FIG. 25D** shows gels run at the indicated salt concentrations (i.e., panel **i** = 250 mM; panel **ii** = 500 mM; panel **iii** = 1M). Lanes in the gel are shown in **Table 54**. Lane 1 is a molecular weight (MW) marker.

Table 54: Silver staining gel lanes

Lane	Cells	Aptamer
1	MW	
2	Ramos	10.36
3	Ramos	10.36(G24A)
4	Ramos	AS1411
5	Ramos	CRO26
6	Jurkat	10.36
7	Jurkat	10.36(G24A)
8	Jurkat	AS1411
9	Jurkat	CRO26

[001400] We compared aptamer 10.36 to aptamer AS1411 (SEQ ID NO. 4370), an anti-NCL homodimeric DNA aptamer that binds specifically to cancer cells, and CRO26 (SEQ ID NO. 4371), a biologically inactive control oligonucleotide. *See* Reyes-Reyes et al.; Soundararajan et al. See **FIG. 25D** and **Table 54**. AS1411 had no detectable binding to Ramos or Jurkat cells under our affinity purification experimental conditions. See **FIG. 25D**, lanes 4, 8. To assess whether immobilization of the aptamer disrupted its ability to homodimerize, binding was repeated in-solution. These experiments were carried out as follows:

[001401] 1. Pellet appropriate volume of cells by centrifuging for 5 minutes at RT, 500xg

[001402] 2. Aspirate media.

[001403] 3. Resuspend cells 500µL binding buffer (same as described above) to provide $1X10^5$ cells per reaction

[001404] 4. Aptamers: dilute aptamer to 10 µM stock and heat denature for 5 min at 95 °C; allow to cool 15 min on bench

[001405] 5. Dilute aptamer stock to a working stock using binding buffer containing Dynasore to inhibit endocytosis of aptamer

[001406] 6. Add 50µl of aptamer to each well of a 96 well deep plate

[001407] 7. Add 50µl of cells to the above wells

[001408] 8. Briefly tap/mix the samples

[001409] 9. Incubate 15 minutes at 37C, 5% CO_2

[001410] 10. Add 10µL of Dynabeads® MyOne™ Streptavidin C1 to all the wells

[001411] 11. Incubate beads with Aptamer–cell solution for 30 mins at RT mixing

[001412] 12. Discard supernatant

[001413] 13. Wash beads twice with 200 µL of 1X HBSS.

[001414] 14. Lyse cells with 200µl lysis buffer for 30min at rm temp Mix on Magmax

[001415] 15. Wash again with with 200 µL of 1X HBSS

[001416] 16. Elute sample with 1% Rapigest heat 37°C for 10 min

[001417] 17. Run one replicate for silver stain and one for anti-nucleolin western blot

[001418] 18. Run 4-12% SDS-PAGE gel

[001419] 19. Silver stain gel or proceed to western blot

[001420] 20. Process remaining replicates in solution digestion

[001421] **FIG. 25E**, panel i, shows that greater levels of protein precipitated with aptamers 10.36 (lane 2) and AS1411 (lane 4) in Ramos cells than controls (i.e., 10.36(G24A) (lane 4) or CRO26 (lane 5)) or with Jurkat cells. However, nucleolin was not detected bound to AS1411 in Ramos cells by Western blot with an anti-NCL antibody (lane 5) under our experimental conditions. See **FIG. 25E**, panel **ii**. Lanes in the figure are shown in **Tables 55** (panel **i**) and **56** (panel **ii**). Lane 1 in **Table 55** and Lane 2 in **Table 56** indicate molecular weight (MW) marker ladders at the weights indicated in kiloDaltons (kDa). In **Table 56**, lane 1 refers to 50 ng of recombinant purified nucleolin protein. The position of nucleolin is indicated by the arrow in **FIG. 25E**, panel **ii**.

Table 55: Silver staining gel

Lane	Cells	Aptamer
1	MW	
2	Ramos	10.36
3	Ramos	G24A
4	Ramos	AS1411
5	Ramos	CRO26

6	Jurkat	10.36
7	Jurkat	G24A
8	Jurkat	AS1411
9	Jurkat	CRO26

Table 56: Western blot with anti-nucleolin antibody (anti-NCL ab)

Lane	Cells	Aptamer
1	50 ng rec. NCL	
2	MW	
3	Ramos	10.36
4	Ramos	G24A
5	Ramos	AS1411
6	Ramos	CRO26

[001422] To further explore target binding of aptamer 10.36, the aptamer was conjugated to crosslinking agent SBED. See **Example 23** above. To perform these experiments, SBED labeled 10.36 aptamer or SBED labeled 10.36(G24A) control aptamer were incubated with Ramos cells. After incubation to allow binding of the aptamers to their targets, the aptamer-target complexes were photo-crosslinked with UV light. Cross-linked complexes were affinity purified and eluted proteins were washed under very stringent conditions (2M NaCl), eluted by boiling in LDS sample buffer, and run on SDS-PAGE for 10 min for clean up. The entire lane was extracted, digested with trypsin, and peptides were detected by LC-MS/MS and identified with Protein Discoverer (Thermo) with SBED as a dynamic modification. **Table 57** lists proteins identified with an SBED modification. Accession numbers in the table are from the UniProt database (www.uniprot.org). "GN=" is followed by the gene name. Proteins in the table are sorted by level detected in Ramos cells, from high to low. IL3RA (CD123) expression is correlated with leukemia and lymphomas and affects proliferation. See Gupta et al.; Zhang et al; Ehninger et al. JMJD1C family member JMJD2 has been shown to be required for expression of Il3RA and survival of acute myeloid leukemia cells. Agger et al. RanBP2 forms a fusion protein with ALK (UniProt Q9UM73) that has been observed in lymphomas. See Galietta et al; Vassileva et al. ALK also forms a fusion with nucleophosmin, which we identified along with nucleolin, fibrillarin, actin, SFPQ, hnRNPM, hnRNPC1/C2 and other proteins. See above.

Table 57: SBED-linked proteins binding to Ramos cells

Accession	Description
P26951-1	Interleukin-3 receptor subunit alpha GN=IL3RA
P49792	E3 SUMO-protein ligase RanBP2 GN=RANBP2
Q701N4	keratin-associated protein 5-2 GN=KRTAP5-2
Q9Y694-1	Solute carrier family 22 member 7 GN=SLC22A7
Q15652	probable JmjC domain-containing histone demethylation protein 2C GN=JMJD1C

[001423] We also assessed whether aptamer 10.36 would bind the prostate cancer cell line DU145 and breast cancer cell line MCF7. Aptamer pull-down experiments using bead captured aptamers was performed as above. We used ~500 ng of aptamer per 10 μL of beads per reaction and $2.5x10^5$ cells per reaction. Results were compared to aptamer AS1411. In addition, the aptamers were biotinylated on either the 5' or 3' ends to determine whether biotin placement affected aptamer binding. Results with Ramos cells and DU145 cells are shown in **FIG. 25F**. Lanes in the gel are shown in **Table 58**. As shown in the figure, 10.36 precipitated many proteins in Ramos cells, regardless of the position of the biotin (lanes 2 and 4), but did not pull down proteins in DU145 cells. Similar results to DU145 were observed with MCF7 cells (data not shown). Under our experimental conditions, AS1411 did not precipitate detectable protein in any case although AS1411 has previously been reported to kill DU145 cells. See Bates et al. Without being bound by theory, any differences may be due to experimental conditions such as dextran sulfate in our buffer.

Table 58: Lanes in silver stained gel

Lane	Cells	Aptamer	Biotin position
1	MW		
2	Ramos	10.36	5' Biotin
3	Ramos	AS1411	5' Biotin
4	Ramos	10.36	3' Biotin
5	Ramos	AS1411	3' Biotin
6	DU145	10.36	5' Biotin
7	DU145	AS1411	5' Biotin
8	DU145	10.36	3' Biotin
9	DU145	AS1411	3' Biotin

[001424] We performed pull-down experiments to identify proteins that precipitate with aptamer 10.36 in Ramos cells. Conditions were as above and included $1x10^5$ cells per reaction, both binding steps at 4 °C for 30 mins, lysis for 30 min, and additions of HL-dsDNAse to shear genomic DNA, various concentrations of non-biotinylated 10.36 competitor, and Dynasore to inhibit endocytosis. Macropinocytosis (and other forms of internalization) are inhibited at 4 °C. Proteins were run on triplicate silver stained gels as shown in **FIG. 25G**. Lanes in the gel are as in **Table 59**. Various concentrations of non-biotinylated 10.36 competitor are shown in **Table 59**. As shown in **FIG. 25G**, only the highest concentrations of competitor appeared to reduce binding, indicating that the binding of biotinylated 10.36 is not saturated under our experimental conditions.

Table 59: Lanes in silver stained gel

Lane	Cells	Aptamer	Competitor
1	MW		
2	Ramos	10.36	None
3	Ramos	10.36	1XNon-Bio 10.36

4	Ramos	10.36	10XNon-Bio 10.36
5	Ramos	10.36	100XNon-Bio 10.36

[001425] We performed differential detection by LC-MS/MS analysis of digests of different gel bands indicated in **FIG. 25G** for one gel (the "fractionated" gel; **FIG. 25G**) and for two otherwise identical gels used LC-MS/MS analysis of the entire gel lane (the "unfractionated" gels; not shown). See **Tables 60-61**. Accession numbers in the tables are from the UniProt database (www.uniprot.org). The column "Band" in **Table 60** indicates the band the gel shown in **FIG. 25G**. For each band, the proteins are listed in order of abundance. **Table 61** lists proteins identified in the two unfractionated gels. These proteins are also indicated by their appearance in the bands in **FIG. 25G**.

Table 60: Proteins precipititated with aptamer 10.36 in Ramos cells

Band	Accession	Gene ID	Description
1	P09874	PARP1	Poly [ADP-ribose] polymerase 1
	P19338	NCL	Nucleolin
	P46087-4	NOP2	Isoform 4 of Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase
	Q00839	HNRNPU	Heterogeneous nuclear ribonucleoprotein U
	Q13435	SF3B2	Splicing factor 3b subunit 2
2	P23246-1	SFPQ	**splicing factor, proline- and glutamine-rich**
	P19338	NCL	**Nucleolin**
	Q9NR30-1	DDX21	Nucleolar RNA helicase 2
	P46087-4	NOP2	Isoform 4 of Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase
	Q00839	HNRNPU	Heterogeneous nuclear ribonucleoprotein U
	P11387	TOP1	DNA topoisomerase 1
	Q14978-2	NOLC1	Isoform Beta of Nucleolar and coiled-body phosphoprotein 1
	P17480-1	UBTF	Nucleolar transcription factor 1
	Q12906-7	ILF3	Isoform 7 of Interleukin enhancer-binding factor 3
3	Q9NVP1	DDX18	ATP-dependent RNA helicase DDX18
	P11021	HSPA5	78 kDa glucose-regulated protein
	Q13769	THOC5	THO complex subunit 5 homolog
	Q08170	SRSF4	Serine/arginine-rich splicing factor 4
	P17844	DDX5	probable ATP-dependent RNA helicase DDX5
	P27694	RPA1	Replication protein A 70 kDa DNA-binding subunit
	P11940-1	PABPC1	Polyadenylate-binding protein 1
	O00567	NOP56	Nucleolar protein 56
4	P60709	ACTB	**Actin, cytoplasmic 1**
	P05109	S100A8	Protein S100-A8
	Q15233	NONO	Non-POU domain-containing octamer-binding protein
	P06702	S100A9	Protein S100-A9
	Q06830	PRDX1	peroxiredoxin-1
	P25311	AZGP1	Zinc-alpha-2-glycoprotein

	P38159-1	RBMX	RNA-binding motif protein, X chromosome
	P07910-1	HNRNPC	Heterogeneous nuclear ribonucleoproteins C1/C2
	P67809	YBX1	Nuclease-sensitive element-binding protein 1
	P36578	RPL4	60S ribosomal protein L4
	Q12905	ILF2	Interleukin enhancer-binding factor 2
	Q05639	EEF1A2	Elongation factor 1-alpha 2
	O60832-1	DKC1	H/ACA ribonucleoprotein complex subunit 4
	Q13601	KRR1	KRR1 small subunit processome component homolog
	Q15287-1	RNPS1	RNA-binding protein with serine-rich domain 1
	P61978-2	HNRNPK	Isoform 2 of Heterogeneous nuclear ribonucleoprotein K
	Q13247	SRSF6	Serine/arginine-rich splicing factor 6
	P42696	RBM34	RNA-binding protein 34
	Q969G3	SMARCE1	SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily E member 1
5	P22087	FBL	**rRNA 2'-O-methyltransferase fibrillarin**
	P06702	S100A9	**Protein S100-A9**
	P04406-1	GAPDH	glyceraldehyde-3-phosphate dehydrogenase
	P07910-2	HNRNPC	**Isoform C1 of Heterogeneous nuclear ribonucleoproteins C1/C2**
	Q9UKM9-1	RALY	RNA-binding protein Raly
	P09651-1	HNRNPA1	Heterogeneous nuclear ribonucleoprotein A1
	P06748	NPM1	Nucleophosmin OS=Homo sapiens GN=NPM1 PE=1 SV=2
	Q08188	TGM3	Protein-glutamine gamma-glutamyltransferase E
	P07355-2	ANXA2	Isoform 2 of Annexin A2
6	P16402	HIST1H1D	**Histone H1.3**
	P16401	HIST1H1B	**Histone H1.5**
	P62979	RPS27A	**Ubiquitin-40S ribosomal protein S27a**
	Q9BTM1-2	H2AFJ	Isoform 2 of Histone H2A.J
	P06702	S100A9	Protein S100-A9
	Q5D862	FLG2	Filaggrin-2
	P30050-1	RPL12	60S ribosomal protein L12
	P05109	S100A8	Protein S100-A8
	P31151	S100A7	Protein S100-A7
	Q06830	PRDX1	peroxiredoxin-1
	P04406-1	GAPDH	glyceraldehyde-3-phosphate dehydrogenase
	Q9NY12-1	GAR1	H/ACA ribonucleoprotein complex subunit 1
	P62633-1	CNBP	Cellular nucleic acid-binding protein
	P15927-3	RPA2	Isoform 3 of Replication protein A 32 kDa subunit
	P62913	RPL11	60S ribosomal protein L11
	Q86V81	ALYREF	THO complex subunit 4
	P62081	RPS7	40S ribosomal protein S7
	P62750	RPL23A	60S ribosomal protein L23a

	P14678-3		Isoform SM-B1 of Small nuclear ribonucleoprotein-associated proteins B and B'
	Q9BRL6-1	SRSF8	serine/arginine-rich splicing factor 8
	Q92522	H1FX	Histone H1x
	Q07020	RPL18	60S ribosomal protein L18
	P26373-1	RPL13	60S ribosomal protein L13
	Q92979	EMG1	Ribosomal RNA small subunit methyltransferase Nep1
	P09661	SNRPA1	U2 small nuclear ribonucleoprotein A'
7	Q04837	SSBP1	**Single-stranded DNA-binding protein, mitochondrial**
	P06702	S100A9	Protein S100-A9
	P62805	HIST1H4A; HIST1H4F; HIST1H4D; HIST1H4J; HIST2H4A; HIST2H4B; HIST1H4H; HIST1H4C; HIST4H4; HIST1H4E; HIST1H4I; HIST1H4B; HIST1H4K; HIST1H4L	histone H4
	P05109	S100A8	Protein S100-A8
	P62979	RPS27A	Ubiquitin-40S ribosomal protein S27a
	P31151	S100A7	Protein S100-A7
	Q5D862	FLG2	Filaggrin-2
	P04406-1	GAPDH	glyceraldehyde-3-phosphate dehydrogenase
	P25311	AZGP1	Zinc-alpha-2-glycoprotein
	P60709	ACTB	Actin, cytoplasmic 1
	P04908	HIST1H2AB; HIST1H2AE	**histone H2A type 1-B/E**
	P68431	HIST1H3F; HIST1H3C; HIST1H3D; HIST1H3G; HIST1H3H; HIST1H3B; HIST1H3A; HIST1H3E; HIST1H3I; HIST1H3J	**Histone H3.1**
	P55769	NHP2L1; SNU13	**NHP2-like protein 1**
	Q5QNW6-2	HIST2H2BF	**Isoform 2 of Histone H2B type 2-F**
	P53999	SUB1	Activated RNA polymerase II transcriptional coactivator p15
	P37108	SRP14	Signal recognition particle 14 kDa protein
	P06899	HIST1H2BJ	Histone H2B type 1-J
	P62888	RPL30	60S ribosomal protein L30
	P62899-2	RPL31	Isoform 2 of 60S ribosomal protein L31
	P61353	RPL27	60S ribosomal protein L27
	P62318	SNRPD3	small nuclear ribonucleoprotein sm d3
	P35268	RPL22	60S ribosomal protein L22
	P69905	HBA2; HBA1	Hemoglobin subunit alpha
	Q8IUE6	HIST2H2AB	Histone H2A type 2-B
	P62829	RPL23	60S ribosomal protein L23
	P62851	RPS25	40S ribosomal protein S25
	P62263	RPS14	40S ribosomal protein S14
	P62269	RPS18	40S ribosomal protein S18
	P60866-2	RPS20	Isoform 2 of 40S ribosomal protein S20
8	P62805	HIST1H4A; HIST1H4F; HIST1H4D;	**histone H4**

		HIST1H4J; HIST2H4A; HIST2H4B; HIST1H4H; HIST1H4C; HIST4H4; HIST1H4E; HIST1H4I; HIST1H4B; HIST1H4K; HIST1H4L	
	P31151	S100A7	Protein S100-A7
	Q5D862	FLG2	Filaggrin-2
	P49458	SRP9; SRP9P1	Signal recognition particle 9 kDa protein

Table 61: Proteins precipititated with aptamer 10.36 in Ramos cells

Accession	Gene ID	
P09874	PARP1	Poly [ADP-ribose] polymerase 1
P16401	HIST1H1B	Histone H1.5
P16402	HIST1H1D	Histone H1.3
P19338	NCL	Nucleolin
P22087	FBL	rRNA 2'-O-methyltransferase fibrillarin
P23246	SFPQ	splicing factor, proline- and glutamine-rich
P30050	RPL12	60S ribosomal protein L12
P60709	ACTB	Actin, cytoplasmic 1
P62805	HIST1H4A	histone H4
Q04837	SSBP1	Single-stranded DNA-binding protein, mitochondrial
Q15233	NONO	Non-POU domain-containing octamer-binding protein
Q9BTM1	H2AFJ	Histone H2A.J
Q9NR30	DDX21	Nucleolar RNA helicase 2

[001426] Taken together, aptamer 10.36 appears to bind a ribonucleoprotein complex on the surface of Ramos cells comprising multiple proteins including without limitation nucleolin, fibrillarin, actin, SFPQ, hnRNPM, and hnRNPC1/C2. See, e.g., **Table 61**. This binding appears to be specific to Ramos cells as much less protein was pulled down with 10.36 in Jurkat (human T lymphocyte cells from a T cell leukemia patient) or DU145 (prostate cancer cells from a brain metastasis). By silver staining gel, no proteins were observed pulled down in MCF7 (breast cancer) cells.

[001427] **Refererences**:

[001428] Opazo F, et al. (2015). Modular Assembly of Cell-targeting Devices Based on an Uncommon G-quadruplex Aptamer Molecular Therapy. Nucleic Acids 4, e251.

[001429] Mayer, G, et al. (2005). Light-induced formation of G-quadruplex DNA secondary structures. Chembiochem 6: 1966–1970.

[001430] Reyes-Reyes, EM, et al. (2010). A new paradigm for aptamer therapeutic AS1411 action: uptake by macropinocytosis and its stimulation by a nucleolin-dependent mechanism. Cancer Res 70: 8617–8629.

[001431] Pinol-Roma (1999). Association of Nonribosomal Nucleolar Proteins in Ribonucleoprotein Complexes during Interphase and Mitosis. Molecular Biology of the Cell: 10, 77–90.

[001432] Biggiogera, M, et al. (2004). Rearrangement of nuclear ribonucleoprotein (RNP)-containing structures during apoptosis and transcriptional arrest. Biology of the cell 96, 603-615.

[001433] Arcovito A, et al. (2014). Synergic role of nucleophosmin three-helix bundle and a flanking unstructured tail in the interaction with G-quadruplex DNA. J Biol Chem.;289:21230-41.

[001434] Mellgren R (2011). A new twist on plasma membrane repair. Communicative & Integrative Biology 4:2, 198-200.

[001435] Tosoni et al (2015). Nueclolin stabilizes G-quadruplex structures folded by the LTR promorter and silences HIV-1 viral transcription. Nucleic Acids Research 43:8884-97.

[001436] Girvan AC, et al. (2006). AGRO100 inhibits activation of nuclear factor-kappaB (NF-kappaB) by forming a complex with NF-kappaB essential modulator (NEMO) and nucleolin. Mol Cancer Ther. 5:1790-9.

[001437] Brázda V, et al. (2014). DNA and RNA Quadruplex-Binding Proteins. Int. J. Mol. Sci. 15, 17493-17517.

[001438] Inder KL, et al. (2010). Nucleophosmin and nucleolin regulate K-Ras signaling. Commun Integr Biol. 3:188-90.

[001439] Wang K, et al. 2010). Export of microRNAs and microRNA-protective protein by mammalian cells. Nucleic Acids Research 38, 7248–7259.

[001440] Hovanessian AG, et al. (2000). The cell-surface-expressed nucleolin is associated with the actin cytoskeleton. Exp Cell Res. 261:312-28.

[001441] Gower HJ, et al. (1989). Cloning and characterization of a myoblast cell surface antigen defined by 24.1D5 monoclonal antibody. Development 105:723-31.

[001442] Cai Y, et al. (2015). C1q protein binds to the apoptotic nucleolus and causes C1 protease degradation of nucleolar proteins. J Biol Chem. 290:22570-80.

[001443] Mendoza O, et al. (2016). G-quadruplexes and helicases. Nucleic Acids Res. 44:1989-2006.

[001444] Kirchhausen, T, et al. (2008). Use of dynasore, the small molecule inhibitor of dynamin, in the regulation of endocytosis. Methods Enzymol 438: 77–93.

[001445] Soundararajan, S. et al. (2009). Plasma Membrane Nucleolin Is a Receptor for the Anticancer Aptamer AS1411 in MV4-11 Leukemia Cells. Mol Pharmacol. Nov; 76(5): 984–991.

[001446] Gupta, SK et al. Gene copy number alteration profile and its clinical correlation in B-cell acute lymphoblastic leukemia. Leuk Lymphoma. 2016 Jun 24:1-10.

[001447] Zhang, W et al. Expressions of CD96 and CD123 in Bone Marrow Cells of Patients with Myelodysplastic Syndromes. Clin Lab. 2015;61(10):1429-34.

[001448] Ehninger A et al. Distribution and levels of cell surface expression of CD33 and CD123 in acute myeloid leukemia. Blood Cancer J. 2014 Jun 13;4:e218. doi: 10.1038/bcj.2014.39.

[001449] Agger K et al., Jmjd2/Kdm4 demethylases are required for expression of Il3ra and survival of acute myeloid leukemia cells. Genes Dev. 2016 Jun 1;30(11):1278-88.

[001450] Galietta A et al., NPM/ALK binds and phosphorylates the RNA/DNA-binding protein PSF in anaplastic large-cell lymphoma. Blood. 2007 Oct 1;110(7):2600-9.

[001451] Vassileva MT et al. SUMO modification of heterogeneous nuclear ribonucleoproteins. Mol Cell Biol. 2004 May;24(9):3623-32.

[001452] Bates, P et al. (2009). Discovery and development of the G-rich oligonucleotide AS1411 as a novel treatment for cancer. Experimental and Molecular Pathology 86:151–164.

Example 42: Cell killing mediated by Aptamer 10.36

[001453] This work expands on that in **Example 41** above. Here we explored cell killing mediated by aptamer 10.36. Aptamer AS1411 has been shown to decrease viability of a variety of cancer cell types and has undergone phase II clinical trials. See, e.g., Bates et al.; Rosenburg et al. Since 10.36 binds Ramos cells as determined by flow cytometry (*see, e.g.,* **FIG. 25A**) and pulls down nucleolin complexes in Ramos cells with a ~10-fold lesser concentration than AS1411, we tested a titration of 10.36 and AS1411 in cellular viability experiments on Ramos, Jurkat and DU145 cell lines.

[001454] To perform these experiments, aptamers were directly added at various concentrations to log phase cells and incubated at 37 °C with CO_2 for up to 6 days. In order to assess cell viability, ATP levels were quantitated with CellTiter-Glo® Luminescent Cell Viability Assay. (Promega, Madison, WI). The reagent was added at indicated time points, mixed to allow cell lysis and incubated for 10 mins at room temperature to allow signal stabilization. Luminescence was detected using a Synergy 2 microplate reader (BioTek Instruments, Inc., Winooski, VT). Experiments were performed in quadruplicate.

[001455] Representative results are shown in **FIGs. 26A-E**. **FIG. 26A** shows viability of Ramos cells after three days incubation of aptamer 10.36 and control 10.36(G24A) at the indicated concentrations of aptamer. Aptamer 10.36 killed over 50% of the cells at 1.0 μM and 10 μM whereas 10.36(G24A) had no effect. **FIG. 26B** shows similar results with the addition of aptamer AS1411 and control CRO26. In these experiments, 10.36 killed Ramos cells with ~5-fold more efficiency than AS1411. **FIGs. 26C-E** show results after 6 days incubation with aptamer in Ramos (**FIG. 26C**), Jurkat (**FIG. 26D**), and DU145 cells (**FIG. 26E**). After 6 days, killing of Ramos cells was similar between aptamers 10.36 and AS1411, whereas only AS1411 had an effect on the viability of Jurkat and DU145 cells. These data are consistent with findings in the Example above that proteins immunoprecipitated with aptamer 10.36 in Ramos cells but not Jurkat or DU145 cells.

[001456] Comparison of aptamer 10.36 with aptamer AS1411 reveals similarities and differences. Both aptamers are able to immunoprecipitate nucleolin-containing complexes. See **Tables 50** and **52-53** above; Bates et al. In addition, both form G-quadruplex structures. See Opazo et al.; Bates et al. Functionally, however, aptamer 10.36 kills Ramos cells more rapidly (day 2 – day 3) than AS1411 (day 4-6). See, e.g., **FIGs. 26B-C**. In addition, aptamer 10.36 kills Ramos cells at a lower concentration than AS1411 (as low as 200 nM). See, e.g., **FIGs. 26B-C**. However, aptamer 10.36 killed Ramos but not Jurkat or DU145 cells under our conditions, whereas aptamer AS1411 killed all cells under the same conditions. Compare **FIGs. 26C-E**. In accordance with the cell killing data, we found that aptamer 10.36 binds to Ramos cells with higher affinity than AS1411. *See, e.g.,* **FIGs. 25D-E** above and related discussion in the Example above. Finally, recognition of Ramos cells by aptamer 10.36 was impervious to

extreme conditions (i.e., 1M NaCl; 4 °C). See **Example 41**. Without being bound by theory, these data suggest that both aptamers recognize a similar target (e.g., nucleolin comprising protein complexes) but operate via different mechanism.

[001457] **Refererences (see also references in Example 41)**:

[001458] Rosenberg J, et al. (2014). A phase II trial of the nucleolin-targeted DNA aptamer AS1411 in metastatic refractory renal cell carcinoma. Invest New Drugs 32:178-187.

Example 43: Cell Growth Inhibition or Killing

[001459] The 10.36 oligonucleotide aptamer of the invention can be used for inhibiting the growth of neoplastic, hyperplastic, malignant or otherwise hyperproliferative cells. See, e.g., **Example 42**. This Example describes using the aptamer to treat a cancer.

[001460] A pharmaceutical composition comprising an anti-10.36 oligonucleotide of the invention is administered to a cancer victim in sufficient dosage (e.g., a therapeutically effective amount) to treat the cancer in the victim. As desired, the composition is administered in combination with traditional cancer therapeutics or in addition to immune therapies, e.g., cell-based tumor immunotherapies, or antitumor vaccines.

[001461] Relatedly, the 10.36 aptamer is used to target a liposome, nanoparticle or other chemotherapeutic agent to a hyperproliferative cell. See, e.g., Liao J et al., Cell-specific aptamers and their conjugation with nanomaterials for targeted drug delivery. Expert Opin Drug Deliv. 2015 Mar;12(3):493-506; Zhu H et al., Nucleic acid aptamer-mediated drug delivery for targeted cancer therapy. ChemMedChem. 2015 Jan;10(1):39-45; Khedri M, et al., Cancer immunotherapy via nucleic acid aptamers. Int Immunopharmacol. 2015 Dec;29(2):926-36. A pharmaceutical composition comprising an anti-10.36 oligonucleotide of the invention on the surface of a liposome is administered to a cancer victim in sufficient dosage (e.g., a therapeutically effective amount) to treat the cancer in the victim. As desired, the composition is administered in combination with traditional cancer therapeutics or in addition to immune therapies, e.g., cell-based tumor immunotherapies, or antitumor vaccines.

[001462] Relatedly, the 10.36 aptamer is used as the targeting domain of a chimeric, multi-part aptamer construct of the invention. A 10.36 region is connected to a segment which also leads to cell killing, such as an immunomodulatory domain. One non-limiting example comprises an anti-C1q oligonucleotide of the invention. See **Example 40**. A pharmaceutical composition comprising an anti-10.36 chimeric oligonucleotide is administered to a cancer victim in sufficient dosage (e.g., a therapeutically effective amount) to treat the cancer in the victim. As desired, the composition is administered in combination with traditional cancer therapeutics or in addition to immune therapies, e.g., cell-based tumor immunotherapies, or antitumor vaccines.

Example 44: Cell Imaging

[001463] This Example describes using a 10.36 oligonucleotide aptamer as an imaging agent.

[001464] The aptamer is combined with imaging agents including without limitation ananomaterial such as a magnetic nanomaterial, quantum dot, gold or radionuclide probe as desired. Sun and Zu.

Aptamers and their applications in nanomedicine. Small. 2015 May;11(20):2352-64; Dougherty CA et al., Applications of aptamers in targeted imaging: state of the art. Curr Top Med Chem. 2015;15(12):1138-52. The nanomaterial or other imaging agent is directly conjugated to the aptamer or encapsulated in a 10.36 targeted liposome or other nanoparticle. The construct can be configured to recognize surface nucleolin complexes and not internalize, thereby preferentially recognizing cancer cells. The aptamer targeted construct is administered to a patient and imaged to visualize the location of desired cells such as surface nucleolin positive cells.

Example 45: 10.36 immunoassay and isolation

[001465] This Example illustrates immunoassays using a 10.36 oligonucleotide. A nucleic acid construct is synthesized comprising oligonucleotide region corresponding to 10.36 (*i.e.*, SEQ ID NO. 4357). The oligonucleotide construct may comprise a biotin modification to facilitate specific recognition by a desired moiety attached to streptavidin. Alternate modifications and sequence variants that retain binding ability (e.g., do not disrupt the G quaruplex structure) may be used as desired. See, e.g., SEQ ID NOs. 4372-4407.

[001466] A labeled 10.36 aptamer is contructed. The 10.36 construct is contacted with fluorescently labeled streptavidin such as a streptavidin - Alexa Fluor® 488 conjugate from Thermo Fisher Scientific, Catalog number: S11223. This creates a fluorescently labeled 10.36 construct which is used to detect targets in various immunoassay formats. In one scenario, a biological sample known or suspected to contain a target of 10.36 (see, e.g., **Example 41**) is contacted with an ELISA plate. The plate is washed and contacted with the fluorescently labeled 10.36 construct. The fluorescent signal is read from the wells in the plate, thereby providing an indication of the presence or amount of target in the biological sample. In another scenario, a biological sample is directly contacted with the fluorescently labeled 10.36 construct. The contacted sample is subjected to flow cytometry to detect fluorescent particles of the size of cells, thereby providing an indication of the presence or amount of cells having surface displayed target in the biological sample. Alternate labels such as disclosed herein or known in the art can be used in such formats.

[001467] Various modifications of the above scenarios are performed. For example, the 10.36 aptamer is directly labeled with Alexa Fluor during the oligonucleotide synthesis process.

[001468] An immobilized 10.36 aptamer is contructed. In one scenario, the 10.36 construct is contacted with streptavidin conjugated beads. The beads are contacted with a biological sample known or suspected to contain a target of 10.36 (see, e.g., **Example 41**). The beads are precipitated (e.g., by centrifugation or magnetism) and washed. Proteins that precipitate with the beads are analyzed, thereby providing an indication of the presence or amount of target in the biological sample. In another scenario, the 10.36 aptamer construct is contacted with streptavidin agarose resin, e.g., Pierce™ Streptavidin Agarose, Thermo Fisher Scientific Catalog number: 20347 or Pierce™ High Capacity Streptavidin Agarose Thermo Fisher Scientific Catalog number: 20357. The resins are placed in a spin column or chromatography column, respectively. The aptamer is contacted with the resin where it is bound by the

streptavidin. A biological sample known or suspected to comprise a target of 10.36 is allowed to pass through the resin. Targets (proteins, complexes, cells, etc) in the biological sample are retained by the aptamer within the resin and are then analyzed after elution. In either scenario, if desired, the 10.36 aptamer is contacted with the biological sample in solution and then the sample is contacted with the beads or resin. This step allows the 10.36 aptamer and target to bind freely in solution prior to aptamer immobilization.

[001469] Various modifications of the above scenarios are performed. For example, the 10.36 aptamer is directly conjugated to a bead or other desired surface.

[001470] One of skill will appreciate that the 10.36 aptamer construct can be used in any desired scenario where antibodies are conventionally used. *See, e.g.,* Toh et al., Aptamers as a replacement for antibodies in enzyme-linked immunosorbent assay. Biosens Bioelectron. 2015 Feb 15;64:392-403. doi: 10.1016/j.bios.2014.09.026. Epub 2014 Sep 16; Chen and Yang, Replacing antibodies with aptamers in lateral flow immunoassay. Biosens Bioelectron. 2015 Sep 15;71:230-42. doi: 10.1016/j.bios.2015.04.041. Epub 2015 Apr 14; Guthrie et al, Assays for cytokines using aptamers. Methods. 2006 Apr;38(4):324-30; Romig et al., Aptamer affinity chromatography: combinatorial chemistry applied to protein purification. J Chromatogr B Biomed Sci Appl. 1999 Aug 20;731(2):275-84.

Example 46: Target Detection in Bodily Fluids

[001471] This Example describes using a 10.36 oligonucleotide aptamer to detect cancer cells in bodily fluids. A bodily fluid such as blood or a derivative thereof, including without limitation sera or plasma, is obtained from a subject. An assay such as described in **Example 45** is used to detect 10.36 bound to cells in the bodily fluid. As desired, such detection may assist in the diagnosis, prognosis or theranosis of a disease or disorder. *See, e.g.,* **Examples 13-14** herein.

[001472] Although preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

2016298317 25 Jan 2021

CLAIMS

What is claimed is:

1. An oligonucleotide having up to 80 nucleotides and comprising:

(i) a sequence according to SEQ ID NO. 4357; or

(ii) a nucleic acid sequence that has at least 90, 95, 96, 97, 98, or 99 percent identity to the oligonucleotide sequence according to SEQ ID NO. 4357

wherein the oligonucleotide is capable of binding to a Heterogeneous nuclear ribonucleoprotein U protein (HNRPU), or a complex comprising HNRPU, or to a cell expressing HNRPU on its surface.

2. An oligonucleotide of claim 1 consisting of the sequence of SEQ ID NO. 4357.

3. The oligonucleotide of claim 1, wherein the oligonucleotide comprises a 5' region with sequence 5'-CTAGCATGACTGCAGTACGT (SEQ ID NO. 131), a 3' region with sequence 5'-CTGTCTCTTATACACATCTGACGCTGCCGACGA (SEQ ID NO. 132), or both.

4. The oligonucleotide of any one of claims 1-3, wherein the oligonucleotide is capable of binding to Ramos cells.

5. The oligonucleotide of any one of claims 1-3, wherein the oligonucleotide is capable of binding to cells comprising surface nucleolin.

6. The oligonucleotide according to any preceding claim, wherein the oligonucleotide comprises a DNA, RNA, 2'-O-methyl or phosphorothioate backbone, or any combination thereof; and/or wherein the oligonucleotide comprises at least one of DNA, RNA, PNA, LNA, UNA, and any combination thereof; and/or wherein the oligonucleotide comprises at least one functional modification selected from the group consisting of biotinylation, a non-naturally occurring nucleotide, a deletion, an insertion, an addition, and a chemical modification, optionally wherein the chemical modification comprises at least one of C18, polyethylene glycol (PEG), PEG4, PEG6, PEG8, and PEG12.

2016298317 25 Jan 2021

7. The oligonucleotide according to any preceding claim, wherein the oligonucleotide is labeled, optionally wherein the oligonucleotide is attached to a nanoparticle, liposome, gold, magnetic label, fluorescent label, light emitting particle, or radioactive label.

8. A composition comprising an isolated oligonucleotide according to any preceding claim wherein the isolated oligonucleotide is capable of binding to Ramos cells; capable of binding to a cell surface nucleolin complex; capable of modulating cell proliferation, optionally wherein the cell proliferation is neoplastic or dysplastic growth or is that of leukemia, lymphoma or renal carcinoma cells; or capable of inducing apoptosis.

9. A method comprising synthesizing the oligonucleotide according to claims 1-7.

10. A method comprising contacting a biological sample with the oligonucleotide according to any one of claims 1-7, optionally further comprising detecting a presence or level of a cell population in the biological sample that is bound by the oligonucleotide, optionally wherein the cell population comprises neoplastic, malignant, tumor, hyperplastic, or dysplastic cells, or wherein the cell population comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells..

11. The method of claim 10, further comprising detecting a presence or level of a protein selected from the group consisting of Nucleolin; Cluster of heterogeneous nuclear ribonucleoproteins A2/B1; Cluster of Nucleophosmin; Cluster of Actin, cytoplasmic 1; rRNA 2'-O-methyltransferase fibrillarin; Cluster of Heterogeneous nuclear ribonucleoprotein Al; Single-stranded DNA-binding protein, mitochondrial; Heat shock cognate 71 kDa protein; Heterogeneous nuclear ribonucleoprotein M; Serine/arginine-rich splicing factor 1; NHP2-like protein 1; Interleukin enhancer-binding factor 3; ATP synthase subunit beta, mitochondrial; Cluster of Heterogeneous nuclear ribonucleoprotein H3; THO complex subunit 4; Nucleolar RNA helicase 2; RNA-binding protein 14; Signal recognition particle 14 kDa protein; Poly [ADP-ribose] polymerase 1; Cluster of High mobility group protein HMG-1/HMG-Y;

25 Jan 2021 2016298317

Heterogeneous nuclear ribonucleoprotein A/B; 78 kDa glucose-regulated protein; splicing factor, praline- and glutamine-rich; Replication protein A 70 kDa DNA-binding subunit; Heterogeneous nuclear ribonucleoprotein AO; Heat shock protein beta-I; Heterogeneous nuclear ribonucleoprotein H; Cluster of Heterogeneous nuclear ribonucleoprotein A3; Apolipoprotein A-I; heterogeneous nuclear ribonucleoprotein r; 40S ribosomal protein SA; Cluster of Heterogeneous nuclear ribonucleoprotein D-like; Cluster of Poly(RC)-binding protein 1; Polypyrimidine tract-binding protein 1; Inter-alpha-trypsin inhibitor heavy chain H4; heterogeneous nuclear ribonucleoprotein D O; glyceraldehyde-3-phosphate dehydrogenase; Heterogeneous nuclear ribonucleoprotein Q; ELAV-like protein 1; Nucleoporin NUP53; H/ACA ribonucleoprotein complex subunit 2; 9 alpha- 1-antitrypsin; nucleolar and coiled-body phosphoprotein 1; 60 kDa heat shock protein, mitochondrial; Heterogeneous nuclear ribonucleoprotein L; Cluster of Serine/arginine-rich splicing factor 5; WD repeat-containing protein 74; Insulin-like growth factor 2 mRNA-binding protein 1; Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase; Regulator of chromosome condensation; Cluster of Polyadenylate-binding protein 1; ATP synthase subunit alpha, mitochondrial; Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit beta isoform; RRP15-like protein; Chromatin target of PRMT1 protein; Serine/arginine-rich splicing factor 3; Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A beta isoform; Transcriptional activator protein Pur-alpha; HUMAN HLA class I histocompatibility antigen, A-3 1 alpha chain; High mobility group nucleosome-binding domain-containing protein 4; 40S ribosomal protein S14; serine/arginine-rich splicing factor 7; F-actin-capping protein subunit alpha- 1; Interleukin enhancer-binding factor; Cluster of Histone HI.2; Cluster of Heterogeneous nuclear ribonucleoproteins C1/C2; splicing factor, proline- and glutamine-rich; heterogeneous nuclear ribonucleoproteins A2/B1; Cluster of Actin, cytoplasmic 1; Heterogeneous nuclear ribonucleoprotein A1; rRNA 2'-0-methyltransferase fibrillarin; Nucleolar RNA helicase 2; NHP2-like protein 1; Heterogeneous nuclear ribonucleoprotein AO; Cellular nucleic acid-binding protein; Heterogeneous nuclear ribonucleoprotein H3; Heterogeneous nuclear ribonucleoprotein K; Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase; RNA-binding protein EWS; RNA-binding protein Raly; nucleolar and coiled-body phosphoprotein 1; THO complex subunit 4; 60S ribosomal protein L23a; Nucleophosmin; Insulin-like growth factor 2 mRNA-binding protein 1; Histone H3.1; H/ACA ribonucleoprotein complex subunit 2; Replication protein A 70 kDa DNA-binding

2016298317 25 Jan 2021

subunit; Cluster of heterogeneous nuclear ribonucleoprotein r; Nuclease -sensitive element-binding protein 1; RNA-binding protein 14; Interleukin enhancer-binding factor 3; Signal recognition particle 14 kDa protein; Interleukin enhancer-binding factor 2; Heat shock cognate 71kDa protein; 60S ribosomal protein L14; 78 kDa glucose-regulated protein; Heterogeneous nuclear ribonucleoprotein H; Nuclear pore membrane glycoprotein 210; Isoform 4 of Probable 28S rRNA (cytosine(4447)- C(5))-methyltransferase; Heterogeneous nuclear ribonucleoprotein U; Splicing factor 3b subunit 2; splicing factor, proline- and glutamine-rich; Nucleolar RNA helicase 2; Isoform 4 of Probable 28S rRNA (cytosine(4447)-C(5))-methyltransferase; DNA topoisomerase 1; phosphoprotein 1 Isoform Beta of Nucleolar and coiled-body; Nucleolar transcription factor 1; Isoform 7 of Interleukin enhancer-binding factor 3; ATP-dependent RNA helicase DDX18; 78 kDa glucose-regulated protein; THO complex subunit 5 homolog; Serine/arginine-rich splicing factor 4; probable ATP-dependent RNA helicase DDX5; Replication protein A 70 kDa DNA-binding subunit; Polyadenylate-binding protein 1; Nucleolar protein 56; Non-POU domain-containing octamer-binding protein; peroxiredoxin-1; Zinc-alpha-2-glycoprotein; RNA-binding motif protein, X chromosome; Heterogeneous nuclear ribonucleoproteins Cl/C2; Nuclease-sensitive element-binding protein 1; 60S ribosomal protein L4; Interleukin enhancer-binding factor 2; Elongation factor I -alpha 2; ACA ribonucleoprotein complex subunit 4; KRRl small subunit processome component homolog; RNA-binding protein with serine-rich domain 1; Isoform 2 of Heterogeneous nuclear ribonucleoprotein K; Serine/arginine-rich splicing factor 6; RNA-binding protein 34; SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily E member 1; rRNA 2'-O-methvltransferase fibrillarin; glyceraldehyde-3-phosphate dehydrogenase; Isoform Cl of Heterogeneous nuclear ribonucleoproteins Cl/C2; RNA-binding protein Ralv; Heterogeneous nuclear ribonucleoprotein Al; Nucleophosmin; Protein-glutamine gamma-glutamyltransferase E; Isoform 2 of Annexin A2; Histone Hl.3; Histone Hl.5; Ubiquitin-40S ribosomal protein S27a; Isoform 2 of Histone H2A.J; Protein S100-A9; ribosomal protein L12; Protein S100-A8; peroxiredoxin-1; glyceraldehyde-3-phosphate dehydrogenase; H/ACA ribonucleoprotein complex subunit 1; Cellular nucleic acid-binding protein; Isoform 3 of Replication protein A 32 kDa subunit; 60S ribosomal protein Ll 1; THO complex subunit 4; 40S ribosomal protein S7; 60S ribosomal protein L23a; Isoform SM-B1 of Small nuclear ribonucleoprotein-associated proteins B and B'; serine/arginine-rich splicing factor 8; Histone Hlx; 60S ribosomal protein L18;

2016298317 25 Jan 2021

60S ribosomal protein L13; Ribosomal RNA small subunit methyltransferase Nepl; U2 small nuclear ribonucleoprotein A'; Single-stranded DNA-binding protein, mitochondrial; histone H4; Ubiquitin-40S ribosomal protein S27a; Protein S100-A7; glyceraldehyde-3-phosphate dehydrogenase; Zinc-alpha-2-glycoprotein; histone H2A type 1-B/E; Histone H3.1; NHP2-like protein 1; Isoform 2 of Hi stone H2B type 2-F; Activated RNA polymerase II transcriptional coactivator p15; Signal recognition particle 14 kDa protein; Histone H2B type 1-J; 60S ribosomal protein L30; RPL31 Isoform 2 of 60S ribosomal protein L3l; 60S ribosomal protein L27; small nuclear ribonucleoprotein sm d3; 60S ribosomal protein L22; Hemoglobin subunit alpha; Histone H2A type 2-B; 60S ribosomal protein L23; 40S ribosomal protein S25; 40S ribosomal protein S14; 40S ribosomal protein S18 Isoform 2 of 40S ribosomal protein S20; Signal recognition particle 9 kDa protein; Protein S 100-A7; Filaggrin-2; Poly [ADP-ribose] polymerase 1(PARP1); Histone H1.5; Histone H1.3; Nucleolin; rRNA 2'-O-methyltransferase fibrillarin; splicing factor, praline- and glutamine-rich; 60S ribosomal protein L12; Actin, cytoplasmic 1; Single-stranded DNA-binding protein, mitochondrial; Non-POU domain-containing octamer-binding protein; Histone H2A.J; and Nucleolar RNA helicase 2 in the biological sample that is bound by the oligonucleotide.

12. The method of claim 10, further comprising detecting a presence or level of a protein selected from the group consisting of Poly [ADP-ribose] polymerase 1 (PARP1); Histone H1.5; Histone H1.3; Nucleolin; rRNA 2'-O-methyltransferase fibrillarin; splicing factor, praline- and glutamine-rich; 60S ribosomal protein L12; Actin, cytoplasmic 1; histone H4; Single-stranded DNA-binding protein, mitochondrial; Non-POU domain-containing octamer-binding protein; Histone H2A.J; and Nucleolar RNA helicase in the biological sample that is bound by the oligonucleotide.

13. The method of claim 10, wherein the biological sample comprises a bodily fluid, tissue sample or cell culture.

14. The method of claim 13, wherein the tissue sample comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung

2016298317 25 Jan 2021

cancer tissue; or wherein the cell culture comprises lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells; or wherein the bodily fluid comprises peripheral blood, sera, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, broncheoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, female ejaculate, sweat, fecal matter, hair oil, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, or umbilical cord blood, optionally wherein the bodily fluid comprises cells selected from the group consisting of lymphoma, leukemia, renal carcinoma, sarcoma, hemangiopericytoma, melanoma, abdominal cancer, gastric cancer, colon cancer, cervical cancer, prostate cancer, pancreatic cancer, breast cancer, or non-small cell lung cancer cells.

15. A pharmaceutical composition comprising a therapeutically effective amount of the oligonucleotide according to any one of claims 1-7, or a salt thereof, and a pharmaceutically acceptable carrier, diluent, or both; optionally wherein the oligonucleotide is attached to a toxin or chemotherapeutic agent, or wherein the oligonucleotide is attached to a liposome or nanoparticle, optionally wherein the liposome or nanoparticle comprises a toxin or chemotherapeutic agent.

16. The composition of claim 15 for use in treating or ameliorating a disease or disorder in a subject in need thereof.

17. The composition of claim 15 when used in a method of inducing cytotoxicity in a subject.

18. The composition of claim 15, wherein the composition is formulated for intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, inhalation, or topical administration.

2016298317 25 Jan 2021

19. The composition of any one of claims 16-17, wherein the subject has a cancer, a premalignant condition, an inflammatory disease, an immune disease, an autoimmune disease or disorder, a cardiovascular disease or disorder, neurological disease or disorder, infectious disease or pain.

Selection strategy
Aptamer library
EpCAM
Single round of binding with extensive washing "Rnd1 direct"
EpCAM
Additional round of binding, washing and competitive displacement with anti-EpCAM antibody "Rnd2 competitive"

FIG. 1


Substrate
Capture Agent
Capture Agent
Capture Agent
Disease Vesicle
Disease Vesicle
Detection Agent
Detection Agent
Fluorescence
Fluorescence
Fluorescence

FIG. 2A


Fluorescence
Detection Agent
Disease Vesicle
Capture Agent
Capture Agent
Capture Agent
Bead
Fluorescence
Detection Agent
Disease Vesicle
Capture Agent
Capture Agent
Capture Agent
Other Vesicle
Fluorescence
Detection Agent

FIG. 2B

Screening Scheme

5	X	20	=	100
Detection Antibodies		Capture Antibodies		Combinations Screened

CD63	CD9	Rab
CD9	PSCA	IgG
CD81	TNFR	CD81
B7H3	CD63 2X	STEAP
EpCam	B7H3	PCSA
	Rab IgG	PSMA
	MFG-E8	5T4
	EpCam 2X	CD24
	CD63	TMEM211

General vesicle biomarker antibodies: CD9, CD63, CD81
Cell of Origin biomarker antibodies: PSCA, MFG-E8, Rab, STEAP, PCSA, PSMA, 5T4, TMEM211
Cancer biomarker antibodies: EpCam, B7H3, CD24
Control antibodies: Rab IgG, IgG

FIG. 2C


i)
General vesicle biomarker/s
200
Cell-of-origin biomarker/s
201
Disease Biomarker/s
202
Characterize Phenotype
203
ii)
Cell-of-origin biomarker/s
210
Disease Biomarker/s
211
General vesicle biomarker/s
212
Characterize Phenotype
213
Capture
Detectors

FIG. 2D


iii)
Cell-of-origin biomarker/s
240
Disease Biomarker/s
241
General vesicle biomarker/s
242
Capture
Cell-of-origin biomarker/s
243
Disease Biomarker/s
244
General vesicle biomarker/s
245
Detectors
Characterize Phenotype
246

FIG. 2E


iv)
Cell-of-origin biomarker/s
220
Disease Biomarker/s
221
General vesicle biomarker/s
222
Analyze Payload
223
Characterize Phenotype
224
v)
Isolate vesicles
230
Analyze Payload
231
Characterize Phenotype
232

FIG. 2F



SEQ ID NO. 2

FIG. 3B



SEQ ID NO. 1

FIG. 3A


Random oligonucleotide pool (1x10^15)
Combine with exosomes from normal patients
1
Retain unbound oligonucleotides and amplify
Separate oligonucleotides from cancer exosomes
3
Combine with exosomes from cancer patients
2
Retain bound oligonucleotides and amplify
Cancer-specific oligonucleotide pool

FIG. 4


500
250
100
Loading Sample ID: EpCAM DIR1 aptamer_biotin - by Sample ID; Conc. (nM)
nm
0.24
0.21
0.18
0.15
0.12
0.09
0.06
0.03
0.00
0
200
400
600
800
1,000
1,200
Time (sec)
EpCAM protein, 500
EpCAM protein, 250
EpCAM protein, 123

FIG. 5


Aptamer 4 (SEQ ID NO. 1)
A)
EGFR capture
Average MFI
C1
C2
C3
N1
N2
N3
B)
PSP capture
Average MFI
C1
C2
C3
N1
N2
N3
C)
EpCAM capture
Average MFI
C1
C2
C3
N1
N2
N3
D)
KLK2 capture
Average MFI
C1
C2
C3
N1
N2
N3

FIG. 6

A) SEQ ID NO. 4

5′–ATCCAGAGTGACGCAGCAGTCTTTTCTGATGGACAC
GTGGTGGTCTAGTATCTGGACACGGTGGCTTAGT

B)



SEQ ID NO. 5

FIGs. 7A-B



FIG. 7C


Pool 3
Pool 2
Pool 1
C7
C8
C9
C10
C11
C12
D1
D2
D3
D4
M
M
D6
D8
D10
D12

FIG. 7D


Apt4 Control
Pool 1
Pool 2
Pool 3
MFI
600
500
400
300
200
100
0
0
5
10
15
20
EPCAM-His tagged, ug

FIG. 7E


Apt4 Control
Pool 1
Pool 2
Pool 3
MFI
1400
1200
1000
800
600
400
200
0
0
5
10
15
20
EPCAM-His tagged, ug

FIG. 7F


600
500
400
300
200
100
0
25 mM HEPES, 1 mM MgCl2
EPCAM, ug
0
1
5
25 mM HEPES, 150 mM NaCl
EPCAM, ug
0
1
5
25 mM HEPES, 1 mM MgCl2 in PBS-BN
EPCAM, ug
0
1
5
25 mM HEPES, 150 mM NaCl, PBS-BN
EPCAM, ug
0
1
5

FIG. 7G


CAR003 vs EPCAM antigen. Effect of denaturing vs MgCl2.
Apt4 Control
Pool 1
Pool 2
Pool 3
1600
1400
1200
1000
800
600
400
200
0
0
1
5
0
1
5
0
1
5
0
1
5
Antigen, ug
Antigen, ug
Antigen, ug
Antigen, ug
No denature, 1 mM MgCl2
75C, 5min, No MgCl2
75C, 5min, 1mM MgCl2

FIG. 7H


Apt4
Pool 1
Pool 2
Pool 3
MFI
1400
1200
1000
800
600
400
200
0
0
2
4
6
8
10
12
Aptamer, ug

FIG. 7I


BSA - Exp. MW 66 kDa
60
50
40
30
20
EPCAM His tagged Exp. MW 29 kDa Appears as 36 kDa due to glycosilation
M EPCAM BSA HSA M

FIG. 7J


cMV
substrate
A) Direct conjugation of cMVs to a substrate
cMV
Lipid chains
Streptavidin
substrate
B) Biotin functionalized lipid crosslinking of cMVs to a substrate

FIGs. 8A-B

C) Immuno-precipitation of cMVs with cMV specific antibodies immobilized to a substrate

D) Immuno-precipitation of cMVs with cMV specific aptamer immobilized to a substrate

FIGs. 8C-D



FIG. 9


i) BCE8
ii) BCE9
iii) BCE10
iv) BCE11
v) BCE12
vi) BCE13
vii) BCE14
viii) BCE15
ix) BCE16
Cancer...
Cancer non...
Non-cancer...
Non-Cancer...

FIG. 10A


i) BCE17
ii) BCE18
iii) BCE19
iiii) BCE20
v) BCE21
vi) BCE22
vii) BCE23
viii) BCE24
ix) BCE25
Cancer exosomes
Cancer non-...
Non-cancer...
Non-Cancer...

FIG. 10B


i) BCE26
60
50
40
30
20
10
0
Cancer exosomes
Cancer non-...
Non-cancer...
Non-Cancer...
ii) BCE27
60
50
40
30
20
10
0
Cancer exosomes
Cancer non-...
Non-cancer...
Non-Cancer...

FIG. 10C


Rounds 1 – 3
Rounds 4 – 6
Rounds 7 – 9
Heat → NaOH
EL
CE
NCE
SS
PCR
EL
CE
SN
Mix
100K CNE
+
100K NCNE
50K CNE
50K NCNE
100K NCE
D3 non-biotinylated library
0.1mg/mL salmon sperm DNA

FIG. 11A


Mean Fluorescence Intensity (MFI)
4500
4000
3500
3000
2500
2000
1500
1000
500
0
CE
NCE
CNE
NCNE
1
2
3
4
5
6
7
8
9
Round of Selection
No competitor DNA during selection
Competitor DNA during selection

FIG. 11B



FIG. 12A


Oligonucleotides detected
4000
3500
3000
2500
2000
1500
1000
500
0
Cancer
Normal
Current Protocol
Prior Protocol

FIG. 12B



FIG. 12C



FIG. 12D

Random

	P-0.005
d-50	49
d-100	27
d-200	17
d-500	5

	P-0.005
d-50	0.06%
d-100	0.05%
d-200	0.06%
d-500	0.05%
Random	0.50%

Cancer v Non-Cancer

	P-0.005
d-50	2471
d-100	1615
d-200	914
d-500	315

	P-0.005
d-50	2.95%
d-100	3.07%
d-200	3.18%
d-500	3.10%
Random	0.50%

FIG. 12E



FIG. 12F


Cluster of non-cancer
Cluster of non-cancer
Cluster of non-cancer

FIG. 12G


kDa
160
110
80
60
50
40
30
20
15
10
1
2
3
4
5
1
3
5
7
9
15
17
19
Targets
Oligos
Strept

FIG. 13



FIGs. 14A-B


C)
Control vs CP
C-R7N vs CP
0
1000
2000
3000
4000
D)
Control vs HP
C-R7N vs CP
E)
Control vs CP
C-R7N vs HP
F)
Control vs HP
C-R7N vs HP

FIGs. 14C-F


B
5'
Capture probe 1407
3'
Probe part B 1403 (SEQ ID NO. 131)
Complementary probe 1402
Probe complement 1404
Oligonucleotide probe 1401
Probe Part A 1405
3'
Tag probe 1406
5'-B

FIG. 14G


Cluster 4, 50
Cluster 2, 50
Cluster 3, 50
Cluster 1, 200
Color Key
Column Z-Score

FIG. 15


A)
ValidationSet-20-TrainSet-7-ROC
True Positive Rate
False Positive Rate
B)
ValidationSet-20-TestSet-7-ROC
True Positive Rate
False Positive Rate
AUC = 0.80

FIG. 16A-B


C)
ValidationSet-20-TrainSet-8-ROC
True Positive Rate
False Positive Rate
D)
ValidationSet-20-TestSet-8-ROC
AUC = 0.68
True Positive Rate
False Positive Rate

FIG. 16C-D


1700
1702
PCB-AL
i) Photocleavable biotin 1701
ii) Enrichment
1703
iii) Heat
Capture
Str. beads
1704
Purify
1705
1706
v)
UV 1707
1708
AL
1709
vi)
MS
1710

FIG. 17A



FIG. 17B


1720
1722
Nanostring Probe A
3'
5'
Reporter Tag
1721
Aptamer Probe
1723
5'
3'
Aptamer
1724
3'
5'
Primer Probe
1725
3'
5'
1727
Nanostring Probe B
5'
Capture Tag
1726
3'
Biotin
1728

FIG. 17C


D)
Nanostring 5hr Scan, Raw Count
5000
4500
4000
3500
3000
2500
2000
1500
1000
500
0
CP
HP
Binder
Neutral
E)
Nanostring 5hr Scan, Normalized Count
4500
4000
3500
3000
2500
2000
1500
1000
500
0
CP
HP
Binder
Neutral

FIGS. 17D-E


Regular PCR 1800
F-primer
1801
R (AL)
1802
R-primer 1803
PCR
F
1804
R (AL)
1802
Denature
F & R
1805
Denaturing gel
PCR
U-PCR
Unequal PCR 1810
Unequal-F-primer
1811
Lengthener
Terminator
R (AL)
1812
R-primer 1813
PCR
F (longer)
1814
R (AL)
1812
Denature
F (longer)
1814
R (AL)
1812

FIG. 18A


M 1 2 3 4 5 6 7 8 9
100 pmol R-primer
200 pmol R-primer
400 pmol R-primer
1000
500
250
125
62.5
31.25
15.625
7.8125
pmol F-primer

FIG. 18B



FIG. 18C



FIG. 19A


1910
Provide Aptamer Pool to Target of Interest
1911
Contact Pool with Sample
1912
Wash, Retain Aptamers that Bind Sample
1913
Identify Retained Aptamers
1914
Characterize Sample
1915

FIG. 19B


1920
Provide Aptamer Pool to Target of Interest
1919
Sample comprising Microvesicles
1922
Contact Pool with Microvesicles
1923
Wash, Retain Aptamers that Bind Microvesicles
1924
Characterize Sample
1926
Identify Retained Aptamers
1925

FIG. 19C


Person
•Inter-person heterogeneity
•Time of day
•Environment
Disease
•Disease phenotype
•Comorbidities
•Disease severity
Biological Complexity
Cellular
•Membrane blebs
•Golgi
•Nuclear
•Endocytosis
Tissue
•Different organs
•Different fluids

FIG. 20A


DNA
RNA
Protein
Protein Complexes
~20k genes
10^5 transcripts
10^6 isoforms
10^7-10^8 ???
MOLECULAR COMPLEXITY AND DIVERSITY

FIG. 20B


Molecular Target Diversity
Biological Complexity
Gap
Sequencing
Mass Spec
RNA Microarray
Antibody arrays
Microscopy
Immunoassays
Flow cytometry

FIG. 20C


Population Specific to Sample type 2 (Retain)
Population of biological features Common to Both Samples
Discard
Population Specific to Sample type 1 (Retain)

FIG. 20D


Sequencing
Target, e.g.,
Extracellular Vesicles
Oligonucleotide probe
library

FIG. 20E


2002
Bound Probing Oligonucleotides 2001
2003
Sample Population A
2002
2003

FIG. 20F


Unbound Probing Oligonucleotides 2001
Sample Population B
2002
2003
2002
2003

FIG. 20G


2010
Sample 2012
Probing Library 2011
Bind 2013
Supernatant 2014
Microvesicle pellet 2015
Elute 2016
Sequencing 2017
Profile 2018

FIG. 20H


2021
Patient Sample
2022
TOP library
2023
Isolate MVs
2024
NGS, PCR, Array
Other

FIG. 20I


Step 1
Incubation with plasma
EV-target-specific ODN association
Step 2
Positive selection
recovery of EVs with bound ODNs
Step 3
Counter selection
Removal of ODNs common to source A, B EVs
Step 4
Positive selection
Recovery of ODNs specific for source A, B EVs
Source A
Source B
Input library of 2.10^15 ssDNAs Library L0
10^11 ssDNAs
10^11 ssDNAs
Discard ODNs in supernatant
Eluted ODNs
ODNs in supernatant
Eluted ODNs
Library L1
Eluted ODNs
Library L2
Profiling-library L3

FIG. 21A



FIG. 21C



FIG. 21B


Cancer vs Non-Cancer
A)
0.616
Sensitivity
Specificity
2000v1
B)
0.637
Sensitivity
Specificity
2000v2

FIGs. 22A-B


Cancer vs Normal
C)
0.698
2000v1
D)
0.659
2000v2
Sensitivity
Specificity
0.0
0.2
0.4
0.6
0.8
1.0

FIGs. 22C-D


Design 1
Primer 1
2301
Variable Region
2302
Primer 2
2303
IMD 2304
Design 2
Primer 1
2301
Variable Region
2302
Primer 2
2303
Linker
2305
(e.g., polymer, nucleotide)
IMD 2304
Design 3
Primer 1
2301
Variable Region
2302
IMD
2304
Primer 2
2303

FIG. 23A


Design 1
Primer 1
2301
Anti-CD20
2311
Primer 2
2303
Anti-C1q
2312
Design 2
Primer 1
2301
Anti-CD20
2311
Primer 2
2303
Linker
2305
(e.g., polymer, nucleotide)
Anti-C1q
2312

FIG. 23B


Construct
2331
Marker
of
Interest
2332
Complement
2334
Membrane
2333

FIG. 23C


2340
C1Q recruiting domain 2341
C-met targeting domain 2342
71
40
1
35
5
30

FIG. 23D


Patient Sample
Patient Sample 2352
Identify probes that bind sample 2353
Identify Probe Targets 2354
Synthesize Construct 2355
Administer Construct 2356
Monitor Disease 2357
Oligonucleotide probe library 2351

FIG. 23E


kDA
B1
B11
L4
L15
B1rc
B11rc
NO
160
110
80
60
50
40
30
20
15
10
IgM$_{HC}$
C1QA, B
C1QC
SA

FIG. 24A


amount DNA bound [%]
0
1
2
3
4
5
B1RC
B1
B11
0
6.6
66.0
330.0
660.0
conc. C1Q [nM]

FIG. 24B


Absorbance [450-540 nm]
2.0
1.5
1.0
0.5
0.0
0
20
40
60
80
100
B11
B11RC
no aptamer
Aptamer conc. [nM]
[C1Q] = 0.625 nM

FIG. 24C


i)
ii)
iii)
iv)
Count 200
F2-488
4.82
SAPE only
Unstained
800 nM
400 nM
200 nM
800 nM
Ramos
Jurkat
10.36
10.36(G24A)

FIG. 25A



FIG. 25B



FIG. 25C



FIG. 25D


i)
kDa
260
160
110
80
60
40
30
20
15
10
1 2 3 4 5 6 7 8 9
Silver stain
iii)
kDa
80
60
50
40
30
20
1 2 3 4 5 6
Anti-NCL ab

FIG. 25E



FIG. 25F


1 2 3 4 5
260
160
110
80
60
50
40
30
20
15
1) PARP1, NCL
2) SFPQ, NCL, DDX21
3)
4) ACTB, NONO
5) FBL
6) HIST1HD, HIST1H1B, H2AFJ, RPL12
7) SSBP1, HIST1H4A, ACTB
8) HIST1H4A

FIG. 25G


G24A
10.36
Ramos
% Viability
150
100
50
0
0.00
0.01
0.10
1.00
10.00
Apt Conc µM

FIG. 26A


Ramos
CRO26
AS1411
G24A
10.36
% Viability
150
100
50
0
0
5
10
Apt Conc µM

FIG. 26B


CRO26
AS1411
G24A
10.36
Ramos
% Viability
150
100
50
0
0
5
10
Apt Conc µM

FIG. 26C


CRO26
AS1411
G24A
10.36
Jurkat
% Viability
150
100
50
0
0
5
10
Apt Conc µM

FIG. 26D


DU145
CRO26
AS1411
G24A
10.36
% Viability
150
100
50
0
0
5
10
Apt Conc µM

FIG. 26E

SEQUENCE LISTING

<110> CARIS SCIENCE, INC.

<120> TARGETED OLIGONUCLEOTIDES

<130> 37901-826.601

<150> US 62/198,051
<151> 2015-07-28

<150> US 62/198,110
<151> 2015-07-28

<150> US 62/220,652
<151> 2015-09-18

<150> US 62/239,206
<151> 2015-10-08

<150> US 62/269,671
<151> 2015-12-18

<150> US 62/305,536
<151> 2016-03-09

<160> 4407

<170> PatentIn version 3.5

<210> 1
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1
ccccccgaat cacatgactt gggcgggggt cg 32

<210> 2
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2
accggatagc ggttggaggc gtgctccact cg 32

<210> 3
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3
ccgcgcagat atacaacgta cctctgtgcg ca 32

<210> 4
<211> 70

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4
atccagagtg acgcagcagt cttttctgat ggacacgtgg tggtctagta tctggacacg 60

gtggcttagt 70

<210> 5
<211> 70
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 5
auccagagug acgcagcagu cuuuucugau ggacacgugg uggucuagua ucacuaagcc 60

accgugucca 70

<210> 6
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 6
ctgtgaggcg tactgcggtg agcctctcat ta 32

<210> 7
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 7
caattgggcc cgtccgtatg gtgggt 26

<210> 8
<211> 80
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<220>
<221> misc_feature
<222> (16)..(55)
<223> n is a, c, g, or u

<400> 8
gggaggacga ugcggnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnncagac 60

gacucgcuga ggauccgaga 80

<210> 9
<211> 15
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 9
gggaggacga ugcgg 15

<210> 10
<211> 25
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 10
cagacgacuc gcugaggauc cgaga 25

<210> 11
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 11
cgcgucuucc ccgcauugcc gcaauugcca uacauuaaua 40

<210> 12
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 12
cgcgtcttcc ccgcattgcc gcaattgcca tacattaata 40

<210> 13
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 13
guccggaacg ccucgauccu cgcauaauau gauacgucug 40

<210> 14
<211> 40
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 14
gtccggaacg cctcgatcct cgcataatat gatacgtctg 40

<210> 15
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 15
guccauggua cgccucgauu ccgcccauac augcauguaa 40

<210> 16
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 16
gtccatggta cgcctcgatt ccgcccatac atgcatgtaa 40

<210> 17
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 17
cacuauccgu uuguccgucc ucuuguggua uugcgcaugc 40

<210> 18
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 18
cactatccgt ttgtccgtcc tcttgtggta ttgcgcatgc 40

<210> 19
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 19
ucuuccaucu ggucgcgaua cagaauacga uuaacauaaa 40

<210> 20
<211> 40

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 20
tcttccatct ggtcgcgata cagaatacga ttaacataaa 40

<210> 21
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 21
gaucacgcug cccuuuguuu aaggccuuua uacaaacgca 40

<210> 22
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 22
gatcacgctg ccctttgttt aaggccttta tacaaacgca 40

<210> 23
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 23
uauucgccag ucacaucaac uaugaugacg cuugacugga 40

<210> 24
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 24
tattcgccag tcacatcaac tatgatgacg cttgactgga 40

<210> 25
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 25
uaccgccuca ucaucggaca cgacguguau caguuggcug 40

<210> 26
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 26
guucucgccu cuguccucau gguucgaacc gguaugcaug 40

<210> 27
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 27
gcgguuucuu cuccugacua caugagauua auaaacgcgc 40

<210> 28
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 28
ccgccucgaa cacugacguc guggaaccuu cgauugcuag 40

<210> 29
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 29
aaucacagua auucugcccc ucugaugaaa ccgguuacuu 40

<210> 30
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 30
cuuagugauu ucgccgcccc ucuguuuagu ggccauugga 40

<210> 31
<211> 40
<212> RNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 31
acacuauucc gguaagucau cguuuaaccg uuuguugcaa 40

<210> 32
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 32
ugcgcaacgc cuugauucac uccuacagug ugucuauaga 40

<210> 33
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 33
aauguuaagc uuacauacgc cugggucacu cuuuguucug 40

<210> 34
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 34
guaaauauuc acguugaauc gccuugcucc ucuuagucug 40

<210> 35
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 35
ccgccucgga ucguucccaa uggugguacc ccuauuaaug 40

<210> 36
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 36
uguagaucgu ucuuauccgc cucggucuuc cccagguuaa 40

<210> 37
<211> 40

<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 37
aucgucgggc cccuuuuaug aaacuuacau gaaagcgcac 40

<210> 38
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 38
uaagagugca caguacugcc ucgauccucc auggcuuaag 40

<210> 39
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 39
gaauuaguac ugacggccgc cuugauccuc cguuagucug 40

<210> 40
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 40
gcccgccucc gaagcccucc uaagugcacu uuaaaccgcg 40

<210> 41
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 41
ccgccuggga ucacucucua cgcguauaaa ugcucuguca 40

<210> 42
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 42
agucugaccc uguuauggac uaccauauca gaaagguacu 40

<210> 43
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 43
ggugauccuc ccccccgccu cgaagauuug ugcacauauc 40

<210> 44
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 44
gcuaccaucg ucuagugagu cacccuuagu ucaucaaggc 40

<210> 45
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 45
gctgtgtgac tcctgcaa 18

<210> 46
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 46
gcagctgtat cttgtctcc 19

<210> 47
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 47
gtccgcttgg gggtgggtat cgaaaattcg cgttctgggg cgg 43

<210> 48
<211> 43
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 48
cttcgctcgt ttcaatgttc agttctggtc tttaggtctt tag 43

<210> 49
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 49
taggaggata atatgtatcc ggtgcatcgg cctgactttc ccc 43

<210> 50
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 50
taggagggct gaaaacttgc ttcttcccgg tgcagttata aca 43

<210> 51
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 51
taggaggata atatgtatcc gatgcatggg cccgactttc ccc 43

<210> 52
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 52
tgcgttttcc ttggggttca gggtaggatg ggggtggagg tgg 43

<210> 53
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 53
gggggggcggg tgggctggta aaaggtgatg cgggggtatt gtt 43

<210> 54
<211> 43

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 54
taggaggata atatgtatcc ggtgcatcgg cccaactttc ccc 43

<210> 55
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 55
ccaatctcgg aaggtttaaa taaggtggtc tttaggtctt tag 43

<210> 56
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 56
tacgttaagt gggtcgggag gggggaatag ggggtttggt tgg 43

<210> 57
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 57
gtgaggtcgg tggggcgggg gatggtgggg ggtcgtttac att 43

<210> 58
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 58
cgatattggg ggggggttgg cgggctattt tccgggtggg tg 42

<210> 59
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 59
tcggttcctg ttaattcgct ggtggttggt ggggtggcgg atg 43

<210> 60
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 60
gcggggaggt gggggtgggt ggagggttgg ttactctcta ct 42

<210> 61
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 61
cgggaatggg agggtggggg tggtggccgg gtcgtgttat acc 43

<210> 62
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 62
taggaggata atatgtatcc ggtgcatcgg tccgactttc ccc 43

<210> 63
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 63
tatagttggg cgtaggcggg ggggggtggt tgggaggtcc aag 43

<210> 64
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 64
taggaggata atatgtatcc ggtgcatcgg cccgaccttc ccc 43

<210> 65
<211> 43
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 65
tcgatcgtcc tcaggatctc gtggttcaaa tcataaagat tat 43

<210> 66
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 66
tgctgtctgg gcgggggcgg tcttgtggtt tctttggggg ggg 43

<210> 67
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 67
gcgagacagg agggtggtct tatacgtggg ggggggtggt tgg 43

<210> 68
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 68
taggagggct gaagacttgc ttcttcccgg tgcagttata aca 43

<210> 69
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 69
cgtgtggggg gtgggttggg ctcgggttgt tatcagttcc atg 43

<210> 70
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 70
cgatattggg gggggggttg gcgggctatt ttccgggtgg gtg 43

<210> 71
<211> 43

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 71
taggaggata atatgtatcc ggtgcatcgg cccgactttc ctc 43

<210> 72
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 72
taggaggata atatgtatcc ggtgcatcgg ctcgactttc ccc 43

<210> 73
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 73
cttcgctcgt ttcaatgttt agttctggtc tttaggtctt tag 43

<210> 74
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 74
ggacatgggt tgggtcggga gggggtggtc ggttgggcag taa 43

<210> 75
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 75
taggaggata atatgtatac gatgcatcgg cccgactttc ccc 43

<210> 76
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 76
taggaggata atatgtatca gatgcatcgg cccgactttc ccc 43

<210> 77
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 77
tggatgtatg gggtctcggg gtgggagggt tcaacttatc tgg 43

<210> 78
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 78
gtccgcttgg gggtgggtat cgaaaattcg cgttctgggg cgg 43

<210> 79
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 79
tcttgtacaa ataggaggga agggggtttt gggaggtggg tgg 43

<210> 80
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 80
cttcgctcgt ttcaatgttc agttctggtc tttaggtctt tag 43

<210> 81
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 81
ctgatgttag taggtcgggg tccggttggg ggttgggttg agg 43

<210> 82
<211> 43
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 82
tggctcgtgg acgtggtggt ggcgggtcgt gggggtgggt agg 43

<210> 83
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 83
taggaggata atatgtatcc gatgcatcgg cacgactttc ccc 43

<210> 84
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 84
cggggagggg gggtcggggt atttattgtg tatgtttttt gtg 43

<210> 85
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 85
tgactgtatc ttggggcggg ttctgggggg ggtgtattgt tca 43

<210> 86
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 86
taggaggata atatgtatcc gatgcatcgg cccgactttc acc 43

<210> 87
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 87
tcatgtctag ggggggggaag tcggtttggg tgggtactct gtg 43

<210> 88
<211> 43

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 88
cttgtttcgc tttgggtttg ggcgggtggg tcaattcctg ttg 43

<210> 89
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 89
ggtgggggcc ctcggtactg tggcggggtg ggtgggttta gtg 43

<210> 90
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 90
tgcgttttcc ttggggttca gggtaggatg ggggtggagg tgg 43

<210> 91
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 91
tggtggggtg ggtctgtggg gtggttttgt tcttatcggg gtt 43

<210> 92
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 92
tgggtgggtt acgggtgggt gttgttatcg cgctgatctg gtt 43

<210> 93
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 93
cgtgacggtt ttaatcaggg gggggactct aacatttggg tgg 43

<210> 94
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 94
tgctcatatc gtgggggat ggggttgcta gtggatgggg tgg 43

<210> 95
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 95
gggggcggg tgggctggta aaaggtgatg cgggggtatt gtt 43

<210> 96
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 96
tgatgtgggg ggtggggtat aatacttatg tttggggttt ggg 43

<210> 97
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 97
ttgggtgggt gggaatgggt atttttttctt cgggcgatgt ttg 43

<210> 98
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 98
tagttggata ctggatttgg gagggatggg gggaggaggg tgg 43

<210> 99
<211> 42
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 99
ctggaaggtt gggtgtgggg cgggggggag gtctctctat gt 42

<210> 100
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 100
tacgttaagt gggtcgggag gggggaatag ggggtttggt tgg 43

<210> 101
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 101
tgccgtgggc cggggagggt gggttggcgt ctctgtttcg ata 43

<210> 102
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 102
ggggggggcgg ccggggtgga ctgtggcccg tcttcgtatt gtt 43

<210> 103
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 103
tggtggtggc tgggggacgg gtatcctgga attaggggtg gg 42

<210> 104
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 104
tcggttcatg ttgggtggga tcggggcggg gtttcttctc ctg 43

<210> 105
<211> 43

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 105
taggaggata atatgtatac gatgcatcgg cccgactttc ccc 43

<210> 106
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 106
tcgtgggtgt gttctggggg gcgtggacgg gggttaatgg gca 43

<210> 107
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 107
taggaggata atatgtatcc gatgcatcgg cccgactttc ccc 43

<210> 108
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 108
taagagtctt aggatgacgt cattcgtccg tcacggtgcg gga 43

<210> 109
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 109
gcctggccag atagcagtta ccttacggga tctatattca ccg 43

<210> 110
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 110
tggctcgtgg acgtggtggt ggcgggtcgt gggggtgggt agg 43

<210> 111
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 111
ttcgtgttca tctgtttatt gttattcaca atccgtcttt tgt 43

<210> 112
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 112
cgaatccgtc gaccgccctc gtaaccatcc aacccagtct cct 43

<210> 113
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 113
caggaaagtt acttatcttt tagacggtta tgttctcatt act 43

<210> 114
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 114
atctgctcga tacgtagata gtctgctcga tacaaaaggg tgg 43

<210> 115
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 115
cgatgagcgg gctccgatat tgtgtcggcg tgactacctt gat 43

<210> 116
<211> 43
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 116
ggtcggccta tttattgcgg cacgttgttt cttgatgtcg ccc 43

<210> 117
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 117
tcgaattcag ttctacatgt gtatttcttg ctccgtttca gaa 43

<210> 118
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 118
tcggtccggc aacaaacccc acgcgccttc gatatgcctg tcg 43

<210> 119
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 119
ccaatctcgg aaggtttaaa taaggtggtc tttaggtctt tag 43

<210> 120
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 120
cgatattggg gggggggttg gcgggctatt ttccgggtgg gtg 43

<210> 121
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 121
cgggaatggg agggtggggg tggtggccgg gtcgtgttat acc 43

<210> 122
<211> 43

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 122
cgtggtgggt tcttgggtgg gggggaggg ttggcgtagt ata 43

<210> 123
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 123
gtgaggtcgg tggggcgggg gatggtgggg ggtcgtttac att 43

<210> 124
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 124
tacgttaagt gggtcgggag gggggaatag ggggtttggt tgg 43

<210> 125
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 125
taggaggata atatgtatcc ggtgcatcgg cccaactttc ccc 43

<210> 126
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 126
taggagggct gaaaacttgc ttcttcccgg tgcagttata aca 43

<210> 127
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 127
tcggttcctg ttaattcgct ggtggttggt ggggtggcgg atg 43

<210> 128
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 128
tgctgtctgg gcgggggcgg tcttgtggtt tctttggggg ggg 43

<210> 129
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 129
aacatggggt tctgataaac ctgtgccaaa tacattgttg agt 43

<210> 130
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 130
attcctcctt tgtaagggat gatatccaac tcctctgcgt tcg 43

<210> 131
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 131
ctagcatgac tgcagtacgt 20

<210> 132
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 132
ctgtctctta tacacatctg acgctgccga cga 33

<210> 133
<211> 15
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 133
tcgtcggcag cgtca 15

<210> 134
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 134

Asp Tyr Lys Asp Asp Asp Asp Lys
1 5

<210> 135
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 135

Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
1 5

<210> 136
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 136

Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
1 5 10

<210> 137
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 137
aacccgcgtg attggggtac tgcccatgcg acttt 35

<210> 138
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 138
aactgaaggc caattaaact ccactgtgct aatct 35

<210> 139
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 139
acatgccggc gttgatactc tactgtgctc cattatga 38

<210> 140
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 140
agacatgcca cgccctttac atactccact gtgcca 36

<210> 141
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 141
agccacgcag acctatctct actgtgccaa tgtt 34

<210> 142
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 142
aggtacgtaa ggctgccatt actctctact gtgccga 37

<210> 143
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 143
agtctagcga tcgacttttc tctacggtgc ccttc 35

<210> 144
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 144
agtgtgcgca tgaatccatt actccactgt gccttga 37

<210> 145
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 145
agttgcagac ccttgtattc tctaccgtgc cctatga 37

<210> 146
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 146
cactgctaac tgatattgtt gaactccacc gtgc 34

<210> 147
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 147
catgcgacaa gcttccatta accatccgct atttatga 38

<210> 148
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 148
catgcggacc acgctttcac aaactcatga ttcaaa 36

<210> 149
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 149
catgcgtacg tcctattctc tctactgtgc attaa 35

<210> 150
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 150
cctgcaatgc ggaaaccctt ctactctact gtgcc 35

<210> 151
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 151
cgcgagccag ggtactgcct gcggtctaaa caatca 36

<210> 152
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 152
cgtgcgactt gttctctact gtgccaataa accata 36

<210> 153
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 153
gatagcgcat attcctctac tgtgctactt gttat 35

<210> 154
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 154
gcaaaccaat tagtgtacct ctaccgtgcc caatga 36

<210> 155
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 155
gcaactactc tctaccgtgc ccagtgattt ctccatga 38

<210> 156
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 156
gcaccacatg tcaatctcta ctgtgcttat ctcata 36

<210> 157
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 157
gcagcacaaa agttactctc ctctaccgtg ccccta 36

<210> 158
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 158
gcagtttgcg tggatactct actgtgctga ccctta 36

<210> 159
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 159
gcatacatga actctaccgt gcccgttgac ttgaaa 36

<210> 160
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 160
gcatagcggt tcacattatt ttactctacc gtgcc 35

<210> 161
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 161
gccacagtat cgttctcgag ctggttcccg ccat 34

<210> 162
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 162
gcgaacatca ctctaccgtg ccatctgaaa tgacgtga 38

<210> 163
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 163
gcgaacttgt aactctactg tgcttatatt aaggttga 38

<210> 164
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 164
gcgaccaata ctctactgtg ctgatacgga tttta 35

<210> 165
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 165
gcgaccactc tataacctca accgtgctca ctcct 35

<210> 166
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 166
gcgcaagttg tctctaccgt gccccgaaag ttttgga 37

<210> 167
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 167
gcgcaccacc tacattgtct ccaccgtgct ttatt 35

<210> 168
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 168
gcgcacccac gtgaactcta ccgtgcctat ttccta 36

<210> 169
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 169
gcgcaccgac ctctaccgtg ccaaaatagg ttatctga 38

<210> 170
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 170
gcgccaccac acaaaccttc cctatcgagg gagat 35

<210> 171
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 171
gcgcggacta tattactcta ctgtgccctg cttatga 37

<210> 172
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 172
gcggatagga caataactta ctccactgtg ccatc 35

<210> 173
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 173
gcggtgacct ctactgtgcg ccccaagcct tagtt 35

<210> 174
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 174
gcgtacagaa cctctaccgt gcccactcac ttccatga 38

<210> 175
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 175
gcgtcccctc ccggatggtc ctttctctac tgtgc 35

<210> 176
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 176
gcgtgctaac ggtatgcaag acgtatgcga ttttc 35

<210> 177
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 177
gctagaccga tccacctcaa acctctactg tgcct 35

<210> 178
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 178
gctagcttag ctctaccgtg cacattccgc tattt 35

<210> 179
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 179
ggtgcaggca agatatttta ctctactgtg cattt 35

<210> 180
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 180
gtaaatgtac atgcgtatcc tcacctctac tgtgc 35

<210> 181
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 181
taacacgtct ttcactctac tgtgcccttt atgcc 35

<210> 182
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 182
taatggcatg cggacctatc ctctaccgtg ctccttga 38

<210> 183
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 183
tatgcgattt ctctaccgtg ccaatatgcc ttgtt 35

<210> 184
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 184
tctgcgattc taccgttact ctaccgtgcc accaaa 36

<210> 185
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 185
tgcgcagtca tttcgccatg ttctctaccg tgccaa 36

<210> 186
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 186
tgtcaggcgg tagtactctc caccgtgcct attgttga 38

<210> 187
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 187
acatgcatac cctacgaatc gttccatcac catac 35

<210> 188
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 188
agtcagtgcg cccgctattt acgatctcac tgttc 35

<210> 189
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 189
agtgcagtcg cgatgggaac ttcttctttg cttta 35

<210> 190
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 190
atggccatgc gaaccgaaac ctagcccatt ttccta 36

<210> 191
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 191
attgcggtca tccccttcca cgctatacca ccat 34

<210> 192
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 192
cactgcaggt cactggcgcc ctatttcctc acatt 35

<210> 193
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 193
catgcccgta aacgcctaat ctaccacctt ttgct 35

<210> 194
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 194
catgccgcct ttcgacttcc atatcccaac acgcc 35

<210> 195
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 195
catgccggtc attccatact caaactcatt cactc 35

<210> 196
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 196
catgcgagct acccttatca tctcatgttt gcttt 35

<210> 197
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 197
catgcgattg cgccccattc cttgccttat ctcac 35

<210> 198
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 198
catgcgcaca tcgtactccg tagccctaaa ttatca 36

<210> 199
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 199
catgcgccct gtgcgttctc tactacccta atcat 35

<210> 200
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 200
catgtacgca cacccactga ttctttacca cccactga 38

<210> 201
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 201
gcaggacacc ccctcattat ttgtttcatc ctacg 35

<210> 202
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 202
gcatcaggcg cactttcatt tgctaatcgt ttttt 35

<210> 203
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 203
gccatgcgag agggttattg ttgatctcat gggtt 35

<210> 204
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 204
gcgataaagc ccatgtaacc ctctgaaaac ccgat 35

<210> 205
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 205
gcgccacgag ctacccgact tccgattact ttctt 35

<210> 206
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 206
gcggacatcg ggcactttat acaaccacct ttttg 35

<210> 207
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 207
gctgcaggcc gtcccaaata ttcgctccca cacata 36

<210> 208
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 208
gctggtcagg cgttccacac ttcctacccg ctttt 35

<210> 209
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 209
tatgcgcagg accacctatt acacgacctt atcct 35

<210> 210
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 210
tcaactgggg tcagtgccaa cgcttacttt cttct 35

<210> 211
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 211
tcaagcagtc atagccggtt tcacgcttta cttac 35

<210> 212
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 212
aaaacacgca aactacctgg aaacttgact tcttt 35

<210> 213
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 213
aaaacactca cgatacctac cctggtcttc acaac 35

<210> 214
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 214
aaaacagcag gcattttcct gtactttcga attca 35

<210> 215
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 215
aaaaccacag ctaccattat cgcattggcc ctact 35

<210> 216
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 216
aaaacccaag caccgaacga actatccctt tttct 35

<210> 217
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 217
aaaacccacc acgcgtcgta cgcataccac ccttt 35

<210> 218
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 218
aaaaccggtc ctgcgccttc ttctccgctt ttatt 35

<210> 219
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 219
caggtactgt ggcaggcgat ccattgctct ctttt 35

<210> 220
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 220
caggtcagca gtccaggcga ttactttctc ttttc 35

<210> 221
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 221
cagtaaccgc tgtctgtcat ttttcaacat tctgt 35

<210> 222
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 222
cagtaactgt agtcgtcgaa ccttgcttac aaccca 36

<210> 223
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 223
cagtaaggaa gctcatgcgg tacaggtcta cctcg 35

<210> 224
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 224
cagtaagtgc tgtcactgcg ccacttgtaa atacttga 38

<210> 225
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 225
cagtaatgcg aagctagtac gttctctctt atttta 36

<210> 226
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 226
cagtacaaac atcgcgattc tttccttgat actgt 35

<210> 227
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 227
gagggcggct ctaatgcatg cgctttcatc tttgc 35

<210> 228
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 228
gagggcgggc agtcatttcg attatatctg cctaca 36

<210> 229
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 229
gagggcggtc caacgcgcaa tgtctttttt cttccga 37

<210> 230
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 230
gagggcgtac aggagtcttg cggtccacat tttat 35

<210> 231
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 231
gagggctact gaagttaatg gcattcttct ctatc 35

<210> 232
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 232
gagggctcat gccagccgct tttcttacct ttatc 35

<210> 233
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 233
gagggctgaa caagtagtgc gtacattttc cttcctga 38

<210> 234
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 234
gagggctgca ggcatccaat ttcctccttg tccct 35

<210> 235
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 235
gaggggccgt ccttcgtacg gaatgtgcat gcgct 35

<210> 236
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 236
gaggggcggt cattgccgtt agcctccttt cgcct 35

<210> 237
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 237
gcgaaactcg ccatatactt aaacttgccc ttttgga 37

<210> 238
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 238
aactgcacgt aaccttaatc actccaccgt gccattga 38

<210> 239
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 239
ttcgccgtta agcatcgtgt catgcgcaac tgttaa 36

<210> 240
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 240
acgcggacct agcatatcgt agtgcgtggc acttt 35

<210> 241
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 241
tatgcgcgcc atcaccggaa ctaggtttgg gcttt 35

<210> 242
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 242
gagtacggaa tcgtgtcatc agctggatac ttttt 35

<210> 243
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 243
atgcactggg ctgaccggta ttgcgtattt ctctttga 38

<210> 244
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 244
tatgccccgt tccattcttc attcactact agccc 35

<210> 245
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 245
aagcacaggt actagcaggt tcgcactccc tgacta 36

<210> 246
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 246
cacgccccaa tatgccatac tctactgtgc ccctt 35

<210> 247
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 247
aatgcgcgct aaagaaggac ttgtcttgcc agccc 35

<210> 248
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 248
gaacggatcg tagtgaaaca ggactggagc ggccc 35

<210> 249
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 249
tcgtaggaca aaggcactgg ttgaatcttt tctat 35

<210> 250
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 250
tgcgcgctaa acactgcaag tctagccccg cattc 35

<210> 251
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 251
tcacctacca agtctccaat cattgcaccc tctac 35

<210> 252
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 252
gcctatcaat gtctctctac tgtgccggac ttttta 36

<210> 253
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 253
aggactgccc ccgcctgact ctactgtgcc atctca 36

<210> 254
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 254
ggtagctcca ccgtgccata ctcgatcttt gttac 35

<210> 255
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 255
ttccgcgacc taactccacc gtgccataat ccctttga 38

<210> 256
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 256
agacttaagc aaggtcgtac gtaggtagtc ccacca 36

<210> 257
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 257
gctaactgca ggccccctat ccgacctaaa tataca 36

<210> 258
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 258
atcgtagtac gaatgtcaag cgtacacgct ttccta 36

<210> 259
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 259
gcgttgatct ctaccgtgca ccgaacactc cccaca 36

<210> 260
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 260
gccgctatat caatcctcta ctgtgcgaat taacctga 38

<210> 261
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 261
ttgctgttat cgtcctctac tgtgccggta ttcct 35

<210> 262
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 262
gtcatgcgga agctttcctg ttttggtata atgtc 35

<210> 263
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 263
gcgaacagat ttctctacgg tgcttacaca tattta 36

<210> 264
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 264
actggaagga gtcctacctg cctgacgcgt gacgta 36

<210> 265
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 265
gccaatactc cactgtgctt gattttactt ttgaatga 38

<210> 266
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 266
tgacctacta agtcagcgca catcctgggc tttct 35

<210> 267
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 267
tgtctctact gtgcccaatt gttcggaaat aactt 35

<210> 268
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 268
aacctctgct gtgcgttaaa ctctaccgtg cctac 35

<210> 269
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 269
tacaacgcgt gcttggggtg acgtggtgcc tcttta 36

<210> 270
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 270
tagtcttgcc gtctatctct accgtgcctg gcccta 36

<210> 271
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 271
gcggaactcg ccccccttaca tcatttttt tggtt 35

<210> 272
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 272
tgcgtgctaa acggtctagc acatgcgggc ctcac 35

<210> 273
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 273
gctgcagtgc gactctggcc tccactgtgc taccttga 38

<210> 274
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 274
ggccacgact ggttccccaa ttcacaggta tacat 35

<210> 275
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 275
gcgatagcct taacttgcct tctgtgaact cgcct 35

<210> 276
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 276
tcacctacta ggtccatcca atacttattg tccca 35

<210> 277
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 277
catgcggcaa gcttttaact cagctttatc tcata 35

<210> 278
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 278
catgacattt tctccactgt gcttgtaatt agtattga 38

<210> 279
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 279
aactccaccg tgcccacttc ttgtttcata gtttaa 36

<210> 280
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 280
ccatgcgatt ttgtttcccc tctactgtgc ttatctga 38

<210> 281
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 281
tgttcacctg tccttcctct ctactgtgcc gttct 35

<210> 282
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 282
atgcgcgcga tcgaggtccc cgatggcttc gctcc 35

<210> 283
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 283
tccgcgtgta accttggcat gtacctagtg tccaca 36

<210> 284
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 284
catgccaagc tttcatttag gatttcgccc acatca 36

<210> 285
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 285
gcgtctgcgt cctagttttc tctaccgtgc ttttcga 37

<210> 286
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 286
ggctgctaga cctccaccgt gccctttctt tggtta 36

<210> 287
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 287
gagtcaggcc caggcttaca tatcactagc tttta 35

<210> 288
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 288
aatactctac tgtgcccccc gttttctcat cgatca 36

<210> 289
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 289
aggcatgcgc agagttaatt tttttctcta ctgtg 35

<210> 290
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 290
gacctgcagc gcacgcccga tatctctact gtgcca 36

<210> 291
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 291
ccggggctag gaaagctacc acgtcatgcg gaacta 36

<210> 292
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 292
catctcaaaa ctttcctcta ccgtgcaaac attta 35

<210> 293
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 293
gcgatcgaac ttgcctccta caatcatcaa ctagc 35

<210> 294
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 294
cagtcatccc gtggactctg tgctcgtatc tccattga 38

<210> 295
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 295
tcgtagtacg cggtgaaccc tcaatttacg ccccta 36

<210> 296
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 296
gcgtcgtcca actccactgt gcccatttta ggtctga 37

<210> 297
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 297
gcatatgcgg tgtttctact ctaccgtgca aaacc 35

<210> 298
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 298
gcggcaacac ttcctctact gtgcttcctg gcttaa 36

<210> 299
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 299
gattcggtcc ccttcttctc tactgtgcct tttct 35

<210> 300
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 300
ccgcgtgctt ggtgcgtgct gcaggaaatt ggaaa 35

<210> 301
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 301
gcgtgccaac ggactggagc tctattgatt gttcta 36

<210> 302
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 302
atgtacgtgc gtacctttta tattctctac tgtgc 35

<210> 303
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 303
agagcctggg cttgctagct ggatttggtc acgaaa 36

<210> 304
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 304
tactgcgcgc ctggacccgt gtgctagaca tgcgc 35

<210> 305
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 305
ccgtggtgcc ctgcgcgcta agaggctcac ggagta 36

<210> 306
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 306
gtaccggacc tagggactaa cgtactggca cctct 35

<210> 307
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 307
taccgcgagc ttgggaagtg ctgctacgtc tgccga 36

<210> 308
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 308
ggtatgcccg cttttactct actgtgcccg acata 35

<210> 309
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 309
gtgcaacact aagttgaccg gattgagtct agcgaa 36

<210> 310
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 310
gactgcaggg tgtcacctct accgtgccat gaggt 35

<210> 311
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 311
ttcgtagtac caatagtcag taatgccctt tcgcc 35

<210> 312
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 312
atgcactgta tgcgggctat gcgaatacct agttatga 38

<210> 313
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 313
cccggcgcgc tacgggacga ttatgagctc ttgcaa 36

<210> 314
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 314
catgcgcaca ggctttttaa tctccatgaa gtttc 35

<210> 315
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 315
gcgggcttac acgtactgga ggggttgcat gccattga 38

<210> 316
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 316
gtcgtagtac tcgcggtcaa gagattcctt atttc 35

<210> 317
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 317
cacgcgtgct tggcgacgtt cacttgcggg cctat 35

<210> 318
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 318
gccacctcac tccactgtgc tatcttatcg ccttca 36

<210> 319
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 319
cgtgtccttt gaccgtactg cgaaacgcct accaaa 36

<210> 320
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 320
gaatcgtaac aaactgcagg cctatctcta cttgt 35

<210> 321
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 321
gcaggaaacc tctactgtgc cttgtagtat actttga 37

<210> 322
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 322
gcagtccata gcgctcctgg tctcctcttt tgctttga 38

<210> 323
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 323
cgtgccactc aaatcctcct ctaccgtgct ttttt 35

<210> 324
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 324
tggagcacgc cgccaatgga cagtccacct tacct 35

<210> 325
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 325
gatgggcctg cgtgctaagg cgcgataacg ggtgt 35

<210> 326
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 326
cgcgggccag gtctgcatgc atgcgaggtc acacc 35

<210> 327
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 327
gcgtgcttac gaacaaggtg cgtgcccgat tctccga 37

<210> 328
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 328
tcagcggacc tagcgaacga ttgcggccac cactta 36

<210> 329
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 329
gccgcgcgtg ccagcataat cgtactatgt ctgcgtga 38

<210> 330
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 330
atcgtagaac tagctagtct gcaatttctg gttac 35

<210> 331
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 331
gccgcgtgcc tggccaaact gggtcaactt ttctc 35

<210> 332
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 332
tctatcgcgt gccagtgtac gtacggtgtc ttatttga 38

<210> 333
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 333
ttacgttccc ggactctact gtgcccctcc ctata 35

<210> 334
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 334
ccaaccgcgc gcctggggac tacgtgacat ctagc 35

<210> 335
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 335
tcgtagtact gctgctttgt gcgtccatac tcgtta 36

<210> 336
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 336
atcgtagtac tgcaactggg attattcaca cactaga 37

<210> 337
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 337
gaattccgcg cgtagtcgag tactagggaa atcgaa 36

<210> 338
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 338
catgcggacc atgattttta ttgagttttt gcttt 35

<210> 339
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 339
attcgagcgg tcttgcgggt atccctccac tgtgc 35

<210> 340
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 340
catgcgcatt actctggtca acttgccgaa tcgaaa 36

<210> 341
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 341
gggcgtatcg tagccacgta caagggttgc atattga 37

<210> 342
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 342
gcgagaagct ttcaatcctc aaccttacgc tttata 36

<210> 343
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 343
acgcctaagt tgcctctacc gtgccgagcc ctcca 35

<210> 344
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 344
tgcggagcta gcaacgaggg tagagtgcga attcc 35

<210> 345
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 345
gataagtcac ggacggtttc tctactgtgc ctttt 35

<210> 346
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 346
gcgtgccaac aacctgcgga caaaagaccc cctaaa 36

<210> 347
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 347
ggtcatgccg ctaggggcga actcctctga gcctgtga 38

<210> 348
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 348
cctcgcgcgc cagggacgtt gcaagttaaa ccatatga 38

<210> 349
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 349
gcaggcgggc caggcttaat acacgcttct actta 35

<210> 350
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 350
atcgtagtaa tgcagcaatt gcaatactcc tatt 34

<210> 351
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 351
accgaaccta gggacgcccc aggaatcaca atgtta 36

<210> 352
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 352
tactgctaac tttcactcta ccgtgcaaac cacca 35

<210> 353
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 353
cccggaagtg ggttcgagct gcatcgtgtc aggcc 35

<210> 354
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 354
tagtaacttc atgcccccag tggaactcct ctgaca 36

<210> 355
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 355
gtccacgcgt gcctgaggac acagccagga aactt 35

<210> 356
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 356
aatcgctggc cagaaagtta gcagtcgagg cccaaa 36

<210> 357
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 357
agggcgtgcc acctcgaaca cgcaggtaat ttttca 36

<210> 358
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 358
tcacctacca tgtccgagta acactcgccc ctcca 35

<210> 359
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 359
cgtggccggt ggtccgtact gccaaaatat ttaca 35

<210> 360
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 360
cgcgtgccag cggactttca ccatgcgata cgtgg 35

<210> 361
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 361
gagttaaagc tatctctacc gtgcccccta aagtta 36

<210> 362
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 362
tatcgcgcgc cagaagagta agcaagtcat gaccca 36

<210> 363
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 363
tgaccaaggg ttccgctgtg gggatacggt taaggga 37

<210> 364
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 364
agccggagct aggctagtgc gcaccagtca aaccta 36

<210> 365
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 365
tcgtagtatt gcaagggcac cttactccca ccatttga 38

<210> 366
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 366
gcgcagctta tcgccatccg caaatccaca cgctt 35

<210> 367
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 367
cccacgcgtg cctgtggatg gcgcactcgg gcataga 37

<210> 368
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 368
caaggagcac gtatgcactg tgcgagttgg tccgta 36

<210> 369
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 369
agttagtctc taccgtgccc attattccgt tctgaa 36

<210> 370
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 370
gcctgtgaac cttaaactct accgtgcctc tttgt 35

<210> 371
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 371
gctaaccaag tcatgggacc acgctctatt gggata 36

<210> 372
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 372
gcaggcatta ttagtgtctc taccgtgctc tggac 35

<210> 373
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 373
taccgcgcgc ctggtagttg cagtattcct ttgtttga 38

<210> 374
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 374
taacctacca ggccatgcgc aggactgtaa tgcca 35

<210> 375
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 375
gcgccacatc gcaagcttat cgacaccccg ctttca 36

<210> 376
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 376
cccggggcta gggatcgggt gcaatcattg aattt 35

<210> 377
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 377
gcgagcctag cccgtaactg gaaggccact actct 35

<210> 378
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 378
gacgggtgcg ggctagcaag ctttacctct attatga 37

<210> 379
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 379
gcgtgccaac gtgactgtgt cgcctttgct ctgta 35

<210> 380
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 380
atcgtaggac tgcgaaatca ttgcgggttt attgt 35

<210> 381
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 381
ttacctacca aggcacgccg ttttatataa ggtgg 35

<210> 382
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 382
ccgcgtgctt ggattgtgca gtcctccctc tgact 35

<210> 383
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 383
ttacctacca gtgcatggcg ataagtcttt cctata 36

<210> 384
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 384
gacgtctgca ggtgagcatt ggtctactta ttggt 35

<210> 385
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 385
cgcgatcgta gcagtacgga ggttcttcta gcatatga 38

<210> 386
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 386
cgcgaatcgt agccgacgtg cttgggcttg cgacc 35

<210> 387
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 387
aagcacggta cgcacagtcc gacttttgcc ttcaaa 36

<210> 388
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 388
tccggaacta ggagacgtac acgccgcaat gcccc 35

<210> 389
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 389
ccgcggacct agcgaggaca gcgcattgct agcta 35

<210> 390
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 390
agcccgcgtg cctgggagac gacctccgac ttgcg 35

<210> 391
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 391
gcgtgctaag gtactgtaca gtctacgcct atgtt 35

<210> 392
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 392
taccgcgata ctgcacgtgt gcgaccgctt ttccttga 38

<210> 393
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 393
tgacctacca aggagagttt catttgatct aaacta 36

<210> 394
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 394
cttagtgccc taagtctgcg tgactctact gtgcc 35

<210> 395
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 395
tgcaggcaga gcccgttaag ctttttatca actcca 36

<210> 396
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 396
gaggcgaatc gtagctccac gtggtgccat gcaga 35

<210> 397
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 397
catccgctag ttgcttttaa tgaccaccgc ttcca 35

<210> 398
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 398
aagcactgta caaactggat gcgtacactt cctcca 36

<210> 399
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 399
gcgtgccaag gtacggaagg tccatgcaac ggttt 35

<210> 400
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 400
cgtgtaccgc ttgtaccgta caaagtcacc ttgtta 36

<210> 401
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 401
catgcgcgct ccactggtct taattttcct ccatga 36

<210> 402
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 402
tctgcgtgct aagagcggcc ctgtagttgc gccttga 37

<210> 403
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 403
gcgtgccaac gattctgcgc tgggctagta ccatta 36

<210> 404
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 404
gcgcgctaac ggacaaatcc cctaacctcg ataaa 35

<210> 405
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 405
attctctacc gtgccacatt ttatcctctg caaac 35

<210> 406
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 406
tgtctggatg ccatgcgcca cgcttttctc actatga 37

<210> 407
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 407
atcgtagtgc tgcaggatct acctacatta ttctgga 37

<210> 408
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 408
gcagtctcat ctgtcttctc taccgtgctc gaatca 36

<210> 409
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 409
gtctccggcg atatctccac tgtgcccttc tctat 35

<210> 410
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 410
gcaagtcatt ccagtcctga ctctaccgtg cttac 35

<210> 411
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 411
gtgatggtcg tgcccgcttt atcttacatc tgctt 35

<210> 412
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 412
gcgtcgacga gatttttgac ctctaccgtg ccgac 35

<210> 413
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 413
tgcgtcaggc gtccgctttt accccaccat gcttt 35

<210> 414
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 414
atgacacgcg cgcctgctag tactgcgctc catgcga 37

<210> 415
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 415
ggtcaagcat tgcgactcta ccgtgcctca aactaa 36

<210> 416
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 416
cacgtactgt ttactctacc gtgcattaaa cgttt 35

<210> 417
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 417
agcaggccct tttcacatac tctctactgt gctcc 35

<210> 418
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 418
taatgcagga gtgtgccaag cttttatttc atgtc 35

<210> 419
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 419
catgcgaccg ctttacctac ataggattac tcttctga 38

<210> 420
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 420
ttgcggatct agcaacgaat ggaaggaaca cgtac 35

<210> 421
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 421
atcgtagtac ccgcgggaaa aaaccactta tcatc 35

<210> 422
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 422
tcgctgcacc gtgtccatgc ggctacacca attcg 35

<210> 423
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 423
gcagaatacc ttactctacc gtgccattaa tacat 35

<210> 424
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 424
gcgccgaata gcctctactg tgcgttaccc tggattga 38

<210> 425
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 425
gcaggccgta tcactctacc gtgcaaaaag tatac 35

<210> 426
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 426
tactgaggcg acaattctct accgtgcccc catttga 37

<210> 427
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 427
gccactagta tcctccactg tgcaaaatta caataga 37

<210> 428
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 428
ggtattgccg tgtagttcgt tctctactgt gcctttga 38

<210> 429
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 429
cttgcgagat tagctttatc acctgttttc cgctt 35

<210> 430
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 430
gcacctctac tgtgcccaca acgaattgga ctcac 35

<210> 431
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 431
ctccgcggtc atccactgct ccattcaact cgcct 35

<210> 432
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 432
aatctccacc gtgcactatc ttgtctcaaa gtttc 35

<210> 433
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 433
gctagctgaa ttcaccatat ctctactgtg cactaa 36

<210> 434
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 434
ctgcgtgcta agacgtacgg tgtcaacatc tctccga 37

<210> 435
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 435
gcgacgggac ttagcttata ttagatttcg cttct 35

<210> 436
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 436
catgcggaag cttactcatt tttattcgat taccaa 36

<210> 437
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 437
taacctacca agcaatcacc gttttggaaa tcatt 35

<210> 438
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 438
gcccagaaaa actctactgt gcgatgacct gttaaa 36

<210> 439
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 439
tccgcgtgcc tggaaaggta ctgtccactt gaacaa 36

<210> 440
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 440
gagtttgcag gcttggggac tctactgtgc aattttga 38

<210> 441
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 441
ttcgcctcca ccgtgcattt ctaactttat ttctta 36

<210> 442
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 442
ccctcggact gagcactgca tacaagaagc cttgc 35

<210> 443
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 443
cccgtgcctc cccttatctc taccgtgcct cctatga 37

<210> 444
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 444
actcagtcat cgtacgcttt ccgatatcgc tttata 36

<210> 445
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 445
tacgcgtgcc aggaactgta ccagtcttat gccgc 35

<210> 446
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 446
gcggatttgg tctctaccgt gcctactact agttaa 36

<210> 447
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 447
ggcacggaat gcatcgtgtc atggcgtctt tcccctga 38

<210> 448
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 448
gtgcaggcca cccagatctc tactgtgcct cttac 35

<210> 449
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 449
atcgtaggac tgcacatcgt agtttagtta ctcac 35

<210> 450
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 450
tcacctacaa atgtcaggca tatactactc tatta 35

<210> 451
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 451
gtacaagcag tttggcgtat atctctactg tgctga 36

<210> 452
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 452
gataacgttt gattactcca ctgtgccata ccctt 35

<210> 453
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 453
tgagcgatcg tagccaacgt actgcctttc ctattga 37

<210> 454
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 454
gcgtcgacca cgttaatatt taccctctac tgtgc 35

<210> 455
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 455
gcctggtcaa ctcgccgcaa tacttttacc catcc 35

<210> 456
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 456
gtcaccctaa ccgtattctc caccgtgcct tacttga 37

<210> 457
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 457
gatagaattg agctctactg tgcctatctg gttat 35

<210> 458
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 458
gtccaaatta gctctctact gtgcctatcc gttaa 35

<210> 459
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 459
gcgcacttgt tcttcttggt cgcggcgtgc cacgcga 37

<210> 460
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 460
gcggtcagag ctaattctgt caactcgccc tttcta 36

<210> 461
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 461
gagcggaact agcgatgcgt acaaggttaa tttcta 36

<210> 462
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 462
aacactctac tgtgcctcca cgatattttc atact 35

<210> 463
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 463
gtagtgatgc cacctcaact gtgcccttgt tttct 35

<210> 464
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 464
ttcctgcagt tctattctct actgtgccct ttgtta 36

<210> 465
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 465
gactgctctc atagtatgaa ctccactgtg cccttga 37

<210> 466
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 466
accatgcgcg caaccttctc taccgtgcat ctttttga 38

<210> 467
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 467
gcgttatacc tccactgtgc tcaattttca tctaaa 36

<210> 468
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 468
acgcgtgcct ggtttactgc agacgagccc cttcatga 38

<210> 469
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 469
gcacacacac tttactctac cgtgccattt tgaagtga 38

<210> 470
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 470
atctgcgggc atccatagat aatactctac tgtgctga 38

<210> 471
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 471
aaatcattcc tgtggtctct ctaccgtgcc aatat 35

<210> 472
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 472
gccataacct atctctactg tgctaaaccg gttaca 36

<210> 473
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 473
gcgactctac ggaaatttgg gatctctacc gtgcctga 38

<210> 474
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 474
acctctactg tgccgattcc attgttcttc cactc 35

<210> 475
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 475
gcgaccaagc cattccaaac tagccctgat ttcaa 35

<210> 476
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 476
ctgtacggct gtctgggcct agtattcccg ctttaa 36

<210> 477
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 477
gggaactgca ggttaaactc tactgtgcta tgttta 36

<210> 478
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 478
gcgtgggttc ctctaccgtg ccgaggttgt tcctaa 36

<210> 479
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 479
tcgcggtcag caattctcca ccgtgcgatt acctttga 38

<210> 480
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 480
tacacgtcac cttactctac tgtgccgtaa tcctca 36

<210> 481
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 481
atgcactgta cgcaggtcac gcgttttacc actactga 38

<210> 482
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 482
gttgcgtgct aacacacgtg actgcctatt tcctatga 38

<210> 483
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 483
gctactgcga taactctacc gtgccactcc cttgttga 38

<210> 484
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 484
tatccaacca ctagtaagcc tccaccgtgc cattta 36

<210> 485
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 485
gctagttacc agtaacctct accgtgcctc ttctgga 37

<210> 486
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 486
cggcgaactc ctctgtagca gtcgtgcggt tcataa 36

<210> 487
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 487
aactgcgtgc ctagtacgga cttgcaccct tgtgg 35

<210> 488
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 488
gcaggatccg ttctttcaac tcgcctattt ctgga 35

<210> 489
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 489
actgcgaagc cttgcatcca attactctac tgtgc 35

<210> 490
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 490
gcgataatct tgtcattact ctaccgtgcc gttta 35

<210> 491
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 491
atgccggcct attactacct aactctactg tgctta 36

<210> 492
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 492
gcgatcaagc gtatttcttc ttacctctac tgtgcga 37

<210> 493
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 493
gatgacacgc cctttccact ccactgtgcc tgttttga 38

<210> 494
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 494
tacatactcc actgtgccac ctttcccgac tttcca 36

<210> 495
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 495
catgcaagat tttgcgtaac ttaattaatt gcttt 35

<210> 496
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 496
gcactatatc cgtacctaaa cctccactgt gcttt 35

<210> 497
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 497
gcgctataac tccgatgtat tctccaccgt gccca 35

<210> 498
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 498
atcgtagtgc tgcaggaaac gagattgatg caccca 36

<210> 499
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 499
tgcgagtggg atttcgtccg acacctcacc atgca 35

<210> 500
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 500
agggtcatct agaacggttg acacgaaccg ttgct 35

<210> 501
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 501
gctaggaaag gacccaaatc tctactgtgc ctacc 35

<210> 502
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 502
gcgaggaata cctatctcta ctgtgccacc aaacg 35

<210> 503
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 503
gccgtttact ctaccgtgcc gtatcgattt ccatta 36

<210> 504
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 504
aggtcgtgcg tcccgttcaa ctagccctgt tctca 35

<210> 505
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 505
cattgcgttc tggctttcgt tcattttcgc tttac 35

<210> 506
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 506
gcactcatcg cggattatct ccaccgtgcc gacct 35

<210> 507
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 507
catgccagct ttaactttga gatctgttag ttttc 35

<210> 508
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 508
gccaggatct caactgtgcc attgcctttt tctgaa 36

<210> 509
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 509
ttacctccac cgtgccaatc actgtttgat cgccttga 38

<210> 510
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 510
taacctacca agtccaattt ggagtctcta cgatta 36

<210> 511
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 511
gcctcaactt gagtttctat cattctccac tgtgc 35

<210> 512
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 512
catagctgca tactctaccg tgcactcatt ccctaga 37

<210> 513
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 513
gacatcaact gcgaactccc ctgagtgtga tatcgga 37

<210> 514
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 514
tatctccacc gtgcccgtta ataatctgat ctgaa 35

<210> 515
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 515
tgcgtgccaa acggactcaa ggtaccatgc atcaca 36

<210> 516
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 516
acatagactg ccgagcgttc ttctctaccg tgccca 36

<210> 517
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 517
atgaccgagg tacccgacgg cctcatagga tccata 36

<210> 518
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 518
gcggttatgg ggttctcaac tgtgccaatc tccat 35

<210> 519
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 519
gggtcatgcg atttctttgt tcactcaact cgccc 35

<210> 520
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 520
gctgttgcgt ccatatctcc actgtgcttt tggttga 37

<210> 521
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 521
atgcgtgcca atacgacgtc aggccatgaa gccat 35

<210> 522
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 522
atgaactggt ggacctttat cctctactgt gcaaaa 36

<210> 523
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 523
ccgcgtgcct ggaccttgca cgacatgcgg caata 35

<210> 524
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 524
tctgcgggct agtatctaat gatctctacc gtgcttga 38

<210> 525
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 525
catgcggagg cttttaacac catgcattct cacc 34

<210> 526
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 526
ttacgtatct gagcgctagt accagtacct cgagtga 37

<210> 527
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 527
agtcaggccc gcttttattt ttgtggttat ttttg 35

<210> 528
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 528
cttgccctcc gacatgaact ctactgtgca acctatga 38

<210> 529
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 529
atcgcgtgcc agaacgtaac tcgcccatcg ac 32

<210> 530
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 530
tgacgcgtgc ttggcaacgt ggcaggacaa ttatt 35

<210> 531
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 531
accgcgtgat tggaagatac tgcgtcttcg atttaa 36

<210> 532
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 532
gtgtgctgac cttagcaacc gaactgaggc cctac 35

<210> 533
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 533
gcgcgccaag gtactgcgga tagcgcattg gacaca 36

<210> 534
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 534
taccgacgcg ctattttcta ctctactgtg ctcat 35

<210> 535
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 535
aaccgaacgt ccgataaact cgccctacta atact 35

<210> 536
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 536
accaggtcca ctttcggtga tagaccctgc acaca 35

<210> 537
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 537
taaattgccc gtttgttctc tactgtgcca ctccg 35

<210> 538
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 538
gccaccttga ctctgatcca actctactgt gccattga 38

<210> 539
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 539
cgtgccaccg ctggcttcct acttcgcttt aactatga 38

<210> 540
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 540
tgttctctac tgtgcccgat aacacgcttt tgttc 35

<210> 541
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 541
ctggtgctgc accgtgtcac ctatctttac gcatta 36

<210> 542
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 542
atcgtagtac gaagctagcc ataacctgtt atctca 36

<210> 543
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 543
gcgtgtttta tctctactgt gcgaacgttt ggaattga 38

<210> 544
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 544
agctaggggt atctctactg tgcgattctt ccgggtga 38

<210> 545
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 545
cgtgcccgct ttattctccc tggcttattg gtccca 36

<210> 546
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 546
gatgcgccgc tttacactct tgttcttacc cgctttga 38

<210> 547
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 547
ggggcatgta ccaaccgtta tttactctac tgtgc 35

<210> 548
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 548
gcatcgtgtc catgcccaaa tggttcgtct tttta 35

<210> 549
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 549
atgcactgga cgcgcatgtc gtttcatttc tccattga 38

<210> 550
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 550
aagcacgggc tgcattcgat tcaggtaaac aatt 34

<210> 551
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 551
acacccgcat gaggtatcct gtggatcgtg ggacgga 37

<210> 552
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 552
gccagctttc gttcatatac cactatcgct ttgta 35

<210> 553
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 553
acagtcataa cagactccgc tgcgtgctaa gcggatga 38

<210> 554
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 554
gcgcgctaac gggtatctgc aggtcctact agttta 36

<210> 555
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 555
atgcacctgt ctgtcagtcc ccgctaggcc atatt 35

<210> 556
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 556
ccatgcgccc ggttgagatt taacggatct cgctttga 38

<210> 557
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 557
gcacacctcg ctttaacact tgtaactacg cttca 35

<210> 558
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 558
tcacctacta aggactgata gctgcctaca ttgtt 35

<210> 559
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 559
ccacgagtct aggggcgtaa cgcagtaagg cataca 36

<210> 560
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 560
acatgcgtgc taaggtacta ctggtgtcga acctttga 38

<210> 561
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 561
atcgtagtac atacagtgcg tttcgtaaat tgcct 35

<210> 562
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 562
gcgccccgac tgccactagc ttttaattct cgctt 35

<210> 563
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 563
tgctagctgg cccatactca actgtgcgtt actcaa 36

<210> 564
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 564
actctgcatc cttcaactct actgtgcggt tattt 35

<210> 565
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 565
gctaactgca gggcacacaa tctcttaagg cttaatga 38

<210> 566
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 566
gcgtgggcct ggtctttact actaaacaaa ttggt 35

<210> 567
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 567
gatcgtaacg cactgagggt caggcattac ccgcc 35

<210> 568
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 568
gcccggggct aggggaaaga acctgccggt catggga 37

<210> 569
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 569
gcgacctagg gtaacgatcg tagttcaccc ataat 35

<210> 570
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 570
ttacctacca cagtctccgc ctcgttatcc ttatca 36

<210> 571
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 571
gccgcgtgcc tggccgtggt gcatcactaa gttgaa 36

<210> 572
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 572
ggcgcgcgct agccatacga actgcttgaa cattt 35

<210> 573
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 573
tgcgaatcgt agccatcgtc cactgctgat ttgcgga 37

<210> 574
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 574
gcgaccgatt cttctctcaa acgcttactt ctatga 36

<210> 575
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 575
gccaaatgat ctccactgtg cgaaatgggc ttttt 35

<210> 576
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 576
tgagcacaca atggacgtga caccaatctt tcttatga 38

<210> 577
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 577
tcatcacgcg ctagtgaacg tgttgtagca ccccta 36

<210> 578
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 578
tcatgtgcgt gccaactgac gaggtgcagg ctagcc 36

<210> 579
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 579
atcgtagctg ctgcagtccc ctgtcttaat atccctga 38

<210> 580
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 580
cgtatcgtag acgttacgga aggagataat gcgcaa 36

<210> 581
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 581
cttgccagct ttccttgtct tcgctttcat tacata 36

<210> 582
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 582
gctgtcattg aatctctact gtgcgatcca ctttttga 38

<210> 583
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 583
catgcgccag cttcttctga ttttatctat ctctatga 38

<210> 584
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 584
ccatgcccgc ttaagcttct tttctatgtt tctacg 36

<210> 585
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 585
tagcaggagt accctccacc gtgcacattg tttcc 35

<210> 586
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 586
catgcccgct ttgtttctcc gctttttggt attccga 37

<210> 587
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 587
atgcacgtcg aaaggcatgc ggtacaactc ctatga 36

<210> 588
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 588
agacagcagg catagcgggg aacctctacc gtgcc 35

<210> 589
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 589
aagtaacgcg ggttacaact ttttctccac cgtgc 35

<210> 590
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 590
acagtctcat gcccgctttt aatcataccg cttaca 36

<210> 591
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 591
gcgcccgacc attccctaag cttttgaccc cactatga 38

<210> 592
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 592
gcgacgacta tcgcttgttc ttgaatacct cgcttta 37

<210> 593
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 593
aaactctact gtgcctcagt ttcaaaccgg ttct 34

<210> 594
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 594
gtcgtagtac tgcgaaatgg gccatgcgca aactt 35

<210> 595
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 595
cgtgcggatt tctccaccgt gcttttgacc tctga 35

<210> 596
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 596
gactgcgacc tacacttatt tgctctactg tgccttga 38

<210> 597
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 597
catgcggctt ctaccaggtc aaactagccc caatt 35

<210> 598
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 598
aaccggagct aggtacgtgt ctgtcaggca tgaaga 36

<210> 599
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 599
atgagtccgc gtgcctggac ctttaccccc gacgaa 36

<210> 600
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 600
catgccgtac gcttattgga tacaccatcg ctttttga 38

<210> 601
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 601
gcgttggtgt tcctctctac tgtgcctgat tttttga 37

<210> 602
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 602
atcgtagtac tgcacgggtg ccacgtgtct tcccaga 37

<210> 603
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 603
aggcacgtac aggcatcgac cgctttcttc tttcc 35

<210> 604
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 604
gccatgcggg ataatattaa atcctctacc gtgcca 36

<210> 605
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 605
tctgccatgc ggctcattcc gaaactctac tgtgcga 37

<210> 606
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 606
ttttgcgatt cctctactgt gccttctggt tttgt 35

<210> 607
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 607
ctgcagcaat catgcaggct ttcatcgctt tcatc 35

<210> 608
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 608
tcacctacca agatctctcg cgccctatag tttct 35

<210> 609
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 609
caccgcgtgc caggggatcg gaatgatatc aactg 35

<210> 610
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 610
taaactacca agggcatgcc accatctact acatga 36

<210> 611
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 611
atcgtagtac tgaaaagcgg gatccagttc taatt 35

<210> 612
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 612
gaaggcaagc acccccctgc tcaactagcc aacaa 35

<210> 613
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 613
ctgccctcgt tctccaccgt gcccactttg taaactga 38

<210> 614
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 614
gatgcaaggg cccgttagta ctggggtcta attaaa 36

<210> 615
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 615
catgcgatgg tcttgcactt ggttagtaat ctccttga 38

<210> 616
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 616
gtgcgcgcta acacggagag tctcaactta acctg 35

<210> 617
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 617
aacaaattgc cgtggggacg tggtcatgcg gattta 36

<210> 618
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 618
gcgtggcacg aagctttcta tttttacttg attcca 36

<210> 619
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 619
caatgccagc ttacgtttat atcccaattc gcttt 35

<210> 620
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 620
aacggtcatg ccagcttttg gtacttagtt ctcct 35

<210> 621
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 621
gaacgcaggc atccgcatta gctttttcct tctatga 37

<210> 622
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 622
gcgccccgta cagcatcgtg tcatggacct acccta 36

<210> 623
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 623
tcgtagtaac tgcaggatcc actccactct catttga 37

<210> 624
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 624
atgcactgta cgagcgggat cgctagttta caaca 35

<210> 625
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 625
gcgcgctaac gtcactgcac atccagggcc ctatc 35

<210> 626
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 626
acccgatcgt agggaacgta ccgataacaa aattatga 38

<210> 627
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 627
actggtaggc atcgacctca agctttctca acatca 36

<210> 628
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 628
acacctacca ggtcacgcgt tcgaacttca cttaa 35

<210> 629
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 629
accgcgcgct agggacggga tgcttcggcc tgtta 35

<210> 630
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 630
taagcaacta gggtgctgtc atcgcgatct ctttt 35

<210> 631
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 631
atcccgggac tagggcgacg tgatgcatct ctcat 35

<210> 632
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 632
gacaaggtct gcccgcttcc cacactacgc ttaca 35

<210> 633
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 633
gcgcccgctt ctcataacta tggctttatt ctcaca 36

<210> 634
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 634
tgctccgggc ctaggagatt ctacgtacgc aatgca 36

<210> 635
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 635
gccagactcg ccccctattt ttttggaact cgccta 36

<210> 636
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 636
gcatccgtta aactcgcctc tcaaatcgaa tcccatga 38

<210> 637
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 637
ctcgcagcat gcgaaggctt tactcacaac tatcg 35

<210> 638
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 638
ccgtgcgaac gttatgcgac catgcgctct ctcttga 37

<210> 639
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 639
gtctagccgc ttttgttatg atattcttct ctttatga 38

<210> 640
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 640
gcgtgccaac gtaccttcct tatctggtga actaca 36

<210> 641
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 641
gtgcgcgcta acactcctct cggtcatttg ctgga 35

<210> 642
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 642
agcgcgtgat tgcacgtgac tgcacttcta atctt 35

<210> 643
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 643
tccgaaccta ggaagccgga tggaacatgc gccca 35

<210> 644
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 644
catcgcaagc tttaatacaa accctatcca ttgct 35

<210> 645
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 645
actgcgtgcc gagtacagga ctacagttaa tggcaa 36

<210> 646
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 646
ccatgcgtgc ctatggacgt ccccaaacat tgtct 35

<210> 647
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 647
tatcgcgggc aggcgattcg cttgtattta atccg 35

<210> 648
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 648
actgcaccgt gtcatggcga gagctcacct aattca 36

<210> 649
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 649
gacatgcaaa ggcccttaat tgactccact gtgcc 35

<210> 650
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 650
catttgcgtg cctaaatgcg gcgcgggctg gcgac 35

<210> 651
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 651
gctggccaat gtatctcaac cgtgccgaaa ttatt 35

<210> 652
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 652
tgctaggact agccggtttt ctattactct atttt 35

<210> 653
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 653
tccggagcta ggaactacgg cgatctctgc gcctaa 36

<210> 654
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 654
gcctagtgga caccccgatt cgacctttt tgctt 35

<210> 655
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 655
agtggaccaa cggatcgcgt gccaggccga tgggca 36

<210> 656
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 656
gataaccagt gtcatgcaaa cactatacat tcctta 36

<210> 657
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 657
actgcacaac attgattctg acaggttgac accaa 35

<210> 658
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 658
agtgcgcaat agtgcgacaa gcttctatcc atttttga 38

<210> 659
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 659
gctgacgttc gtagtccatg tacgggctgt catcaa 36

<210> 660
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 660
tcgtgcagta tcgcggtcat gtcacactcc cgtaca 36

<210> 661
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 661
agcacgatcg tagtgcgaat cgaacggggt ctgcga 36

<210> 662
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 662
gcgcgctaac gactgtagga atccaattct gaccga 36

<210> 663
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 663
agtgactgcg ggcactgcga tcgcttttga ttatttga 38

<210> 664
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 664
agccacccat cctcaaccgt gccataatcg cctcta 36

<210> 665
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 665
gcctctaccg tgcctgtgac ccaccagttt tttat 35

<210> 666
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 666
tcgtagtacg caggtggtca agcacgattt cttta 35

<210> 667
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 667
tcagtacgca ctagtacatt gccaagcttt ccacatga 38

<210> 668
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 668
aagcggaact agccaaggca agggaaatgc cctca 35

<210> 669
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 669
ggaccgaatc taggctgtac tgtgcacgca tcgcc 35

<210> 670
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 670
ttacctacct ggtcaggcat aacatttagt tcctc 35

<210> 671
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 671
aagaccgcgt gccaggtcaa cgtacgcaca tgcaatga 38

<210> 672
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 672
atcgtagtac tgtagggcct ctgtttcact caatatga 38

<210> 673
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 673
aggacaggcg ggttttgctg acgttgtcta tactg 35

<210> 674
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 674
tactccactg tgcatccgat cttcttcccc aactc 35

<210> 675
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 675
tctccactgt gcccgtaact tttcttttac tgtccga 37

<210> 676
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 676
ccgggcctag gaggactgca tgaccccttt aaccaa 36

<210> 677
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 677
ccgggcctag gacggttcca cagtcagcga atcct 35

<210> 678
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 678
aacgcgcgct tgggatgtat gcatcatcta gattt 35

<210> 679
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 679
cacgacgcac cttaccgctt tctatttttc cgctt 35

<210> 680
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 680
gcacttggcc gctttaacta ctacgctttg tcctc 35

<210> 681
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 681
cgtgtacgcc agtgccgtgc agtcaggcga tctatga 37

<210> 682
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 682
cttaggcaat gacgtgccct tatgcgactt cgccca 36

<210> 683
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 683
gcaccgtttc caccgcgcgt ggatcttccc aactttga 38

<210> 684
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 684
gcggacctag cttacggact gtaattaacg ttttt 35

<210> 685
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 685
atgcgcatcc cctctaccgt gcaattttcc atccctga 38

<210> 686
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 686
gacgcgcgcc agacagggaa ctgcacaggc atgcca 36

<210> 687
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 687
ccgagactag gacccctgtg ttcatgcgct atttc 35

<210> 688
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 688
gctaactgca ggcacagcct aagaaacctc atgta 35

<210> 689
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 689
tgcagtctag cccgcttaat gtattccttg attcttga 38

<210> 690
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 690
gcgtgctagc gtactgcatg gatactttgg tagct 35

<210> 691
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 691
tcgtagtaca gcagaccctc cctacctaca attggtga 38

<210> 692
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 692
gccgctgact gagttctctc tactgtgcgt atttt 35

<210> 693
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 693
gccagcagcc ctgaactcct ctgaagcatc gccgca 36

<210> 694
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 694
taggtctgca catgcgctag gcttttctta ccctg 35

<210> 695
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 695
gcgacgcccc acgctttgac ctaaacccat caatatga 38

<210> 696
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 696
cgctacagac atgcgcatct tgacttatat tcaattga 38

<210> 697
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 697
atgcactact gcgagcgagc acaggtttcc tcctt 35

<210> 698
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 698
aacgcgtgcc agagaagaac cgcggtctcg ccccatga 38

<210> 699
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 699
cattgcgaac tagctttctt ctccttctag ctttt 35

<210> 700
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 700
catgccctag aacattgctt taacactttc gcttt 35

<210> 701
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 701
tcatgcgatc aatcgctctg cgtgcctaga tcgattga 38

<210> 702
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 702
gcgaccgtgg cttcctacct ccaagctttc cgac 34

<210> 703
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 703
gcgtgctaac gtactagacc aaagcgagta acatca 36

<210> 704
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 704
tcacgcgcgc ttgggattac gtacaatcat caatgga 37

<210> 705
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 705
gcactacacc ctattcccgc gtaggtcccc gagcct 36

<210> 706
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 706
taactccact gtgcgatcat tgtttttctt ttttga 36

<210> 707
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 707
atcgtagtac tggtactcct gccctatttc gataa 35

<210> 708
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 708
acactctgca gcctagaact ctaccgtgct aattt 35

<210> 709
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 709
gtgcgcgcgc tagcatgcgt aatcatgtca tttaa 35

<210> 710
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 710
tcaatctcaa ctgtgcttcc ctggttttta gtttc 35

<210> 711
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 711
gtcctgcgtg ctaaggaccc taagcttcat gcgtatga 38

<210> 712
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 712
tcgccgccca aggatccctc tactgtgcca ctttaa 36

<210> 713
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 713
gcgtgcctac gtaacgttca tgcggatgga ccgat 35

<210> 714
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 714
cgtgccttgg ccggaccagt ccttttgatc actac 35

<210> 715
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 715
caggcatgac gatttggctt aacttccctt gctta 35

<210> 716
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 716
tatgcgtatc caatcgcttt tcaattccta tgctt 35

<210> 717
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 717
ccgcgtgcca ggcgtaatgt aggcaggccc tttat 35

<210> 718
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 718
gctaacagca gtcattgaat cctgtacgct ccaaa 35

<210> 719
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 719
atcgcctgcg ccaggactgg ggtgcagatc atgcg 35

<210> 720
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 720
ctgagcagtc accattcttt atctctactg tgctcga 37

<210> 721
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 721
acccacatcc tgcgtgctaa ggacggtggg gccgg 35

<210> 722
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 722
ccggacctag gaatcgactc aggcgtggta gaacaa 36

<210> 723
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 723
tcatggccac taaactcgcc ccataaaacc ttgtc 35

<210> 724
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 724
gactctactg tgcccgtttg ttgttcttcc cgaac 35

<210> 725
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 725
ggagccgggc ctgccacgta gctcggaaaa tgcgga 36

<210> 726
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 726
gatttcacta gctctctact gtgcccattt ctgttga 37

<210> 727
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 727
agcggatcta gccagtatgc actcatgtca ctttt 35

<210> 728
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 728
actgcacgtg cgcgcttaag ttttcgcttc tcttt 35

<210> 729
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 729
gtcgagggga actcctctga ggctcaacac atatctga 38

<210> 730
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 730
gcgatccggt taaactacac actctactgt gcgaa 35

<210> 731
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 731
gctaaccaag gaatgcgctg ccttagacag aattg 35

<210> 732
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 732
tcacctacca gggcgtggta ggtttttctc tctac 35

<210> 733
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 733
cctcgcgtgc cagggaccta gtgcaatcta tccgaga 37

<210> 734
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 734
ccgagcgaac gtacggaaag tgcgtacttt atctttga 38

<210> 735
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 735
agtacgtgac cactactttc cctcctttgt acttga 36

<210> 736
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 736
tcatgcggtt gattggtcta ctcatgtttc ttaattga 38

<210> 737
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 737
gcacaatccc acgcttccta ttataccacg attcta 36

<210> 738
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 738
acaacccggc gcgccacggg gtgatcgaca actgt 35

<210> 739
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 739
gatcgtccgc gtgcctggca acgatacccg acaag 35

<210> 740
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 740
gcgtgtctct ctactgtgcg cctcattctt atata 35

<210> 741
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 741
tcgtagtaca cgcagtcgac aaacaagcta gtatt 35

<210> 742
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 742
ttacctacca acgtaatgcg ctacaaccag agagta 36

<210> 743
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 743
ttagcgaaac tagcgtacga caaggtccaa tgtaa 35

<210> 744
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 744
tgggcggatc tagcaaaaac ggacgcctcc tgcgaa 36

<210> 745
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 745
aagcacggga cgcatgtcgt ggctacctga attct 35

<210> 746
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 746
gtgtgcaggc atgacgataa gctttctaac tcctatga 38

<210> 747
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 747
tctgcgcgcc tagaagtacg caatcaccac gcgac 35

<210> 748
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 748
atgtcgctgc aggtgcgtgc aagctttaat ctccttga 38

<210> 749
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 749
tccgggccta ggaacggtca ctagggaatg cccctga 37

<210> 750
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 750
acgtagtgca catcctagac tccactgtgc ctctta 36

<210> 751
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 751
tcatgcatag gcttatatga caccatcgct tcttt 35

<210> 752
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 752
actgccggct gcaagctttt aggtacactt gattca 36

<210> 753
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 753
cacgcgccgc tttttcgcct atacgctttt tattttga 38

<210> 754
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 754
tctgactgca cgcatgggac cgcttaccac taaccga 37

<210> 755
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 755
tgcgggacta gcataagttc cacatgcgaa gacaaa 36

<210> 756
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 756
gcgtgcctac gtacaggcgg atatgcgcgt ttccc 35

<210> 757
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 757
gcagtcgcgt atcgattcgc tttcttatct ggttatga 38

<210> 758
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 758
cgcaggcatg caggctttta gtttaattac tcctttga 38

<210> 759
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 759
gacactccag cactaaagct agtgtggtat cactta 36

<210> 760
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 760
cactgcgctt ctagtactac ctccactgtg ctttttga 38

<210> 761
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 761
acacatgccg cttaatattg attcgctttt atcat 35

<210> 762
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 762
gagtagccaa acgcgtgctt ggtggcgatc acgta 35

<210> 763
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 763
tactaaggaa agcggtaaaa cgtacactga aatct 35

<210> 764
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 764
ggtggtgcga aaaatgcggt gtgaactccc ctgaa 35

<210> 765
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 765
tatcgcgagc caggagacgt aaaaaggaaa tgcgg 35

<210> 766
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 766
ggccaccgac cactatgcat cgtgtcatga cgtaa 35

<210> 767
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 767
atccgatctc taccgtgcaa cccacccctt tgttta 36

<210> 768
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 768
gcggtcgatc ctctcaactg tgcacttttt actacga 37

<210> 769
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 769
attcgtagct caactctact gtgcattttt tgtttga 37

<210> 770
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 770
gccgctaaac tgtctccact gtgctacctt atcttga 37

<210> 771
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 771
cgcgtgcttg cgtactgtcc gcgtaatcac acaca 35

<210> 772
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 772
tctccactgt gccgtttcct ccgttcttac tcctaa 36

<210> 773
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 773
acctgcgtgc taaggaacgt agcattcgtg tgcga 35

<210> 774
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 774
acacgcgcgc ttgggactga actcagcaat gcacc 35

<210> 775
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 775
tcacctacta acgtgtgcgg ggtttttaca tcact 35

<210> 776
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 776
gatagccgct tagccaacac aaccgctttt gattca 36

<210> 777
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 777
gcggccttgc atcgattcaa ctcgccctat ttcaa 35

<210> 778
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 778
agcgcatcgt agccgtacag caatgtaata ttctc 35

<210> 779
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 779
gctaactgca ggccaccacg cttgccctgg ctacc 35

<210> 780
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 780
aagcactgtg ccgcagatat gcgatttttg tgatt 35

<210> 781
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 781
tgagccgcgc gcctggccac gacgattcat gtgattga 38

<210> 782
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 782
catgcgtaag cttatgtttc gattacgttc ccattga 37

<210> 783
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 783
gtggtccgcg cgaaccacca tgggtgcagg cgaat 35

<210> 784
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 784
tcacctacca gggaaatgtc cagccacccc gagct 35

<210> 785
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 785
gcagggcttg caggctttta attgtttcac tatcg 35

<210> 786
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 786
gctaagtgcg gtcactgccc caggagcttg atgcta 36

<210> 787
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 787
gtgcgggact agcacgcagt acagctttct gcgtc 35

<210> 788
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 788
gcgggactag cgagtttagc ctgatactac attga 35

<210> 789
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 789
gcggacgatt cctgttcctt tgcttatttc tctgaga 37

<210> 790
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 790
acacggacct agggtcgtac tgcaggaagt tgttc 35

<210> 791
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 791
actccactgt gcctctttaa cttctttcca tatatga 37

<210> 792
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 792
gcgtgcctac gacgccaggc actagtttag taaac 35

<210> 793
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 793
aagcggggag caacgtactg ccatttacta ttccatga 38

<210> 794
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 794
tttaacggaa cgacctgtgg ctgcacgttt ctcca 35

<210> 795
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 795
tttgtactga gcaagcgtac ataccgaagc cccgt 35

<210> 796
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 796
catgcgcgat tagcttgttc taccaccgct ttaca 35

<210> 797
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 797
catgtcgacc tctttgtaaa ctcgccctat tttcctga 38

<210> 798
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 798
cattgcgcgc tttcaataac ctccgcttat aacac 35

<210> 799
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 799
catctagagg ctcattcata ttaacacgct ttcct 35

<210> 800
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 800
tcgtagtgct gcaagtcctt agtaccttgg atcttga 37

<210> 801
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 801
atgcacggtt ctgctcaggc catcattttc ttcac 35

<210> 802
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 802
gcgcgcctac gttaatgcac ttcgtgacat cttcc 35

<210> 803
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 803
ccggtcgtag ggtgactgcg ggcataagct gggtt 35

<210> 804
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 804
tgcgcaggcc cgttgtactc gctcactgct agatt 35

<210> 805
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 805
aagcacggac aagcatgtat gcgatcacgc ccttta 36

<210> 806
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 806
gactgcggga ctagcaacgt actgtggctg gcgat 35

<210> 807
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 807
gcctccgtaa cctacccgcg tgcctggggg ttacc 35

<210> 808
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 808
gcagggaatc cagcgaccgt tcctctaccg tgcct 35

<210> 809
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 809
catgcggaag ctttctctgt tgatgatgaa ccctttga 38

<210> 810
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 810
actgcgctaa tgcccgctta acctccatga ttgtt 35

<210> 811
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 811
ggggtcgcgt gccagaccac gactcatgtc tgtgc 35

<210> 812
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 812
atcgtagaac tgcagacccg catggctacc tttgt 35

<210> 813
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 813
attgcggatc gtagctctaa ctgcgcatgc ggtaca 36

<210> 814
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 814
tacacaccgc tttactatcc tcaatttcgc tttcc 35

<210> 815
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 815
tgcctctact gtgccgatac cggtttcttt atatt 35

<210> 816
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 816
tcccgcgcgc tagggtaatg atccagcaga catgta 36

<210> 817
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 817
tcgtagtact cgcgcttccc gtatcacaat tttgaa 36

<210> 818
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 818
aggtcgcgcg ccagggacgt gctgcagaac cacctga 37

<210> 819
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 819
tctaccgcgc gcttggagga acgttccgga ctggca 36

<210> 820
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 820
gcaaaataat accgctacat tctctacaac tcgcc 35

<210> 821
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 821
gcgtggggtg cgcgctaacc acacttgacc agacta 36

<210> 822
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 822
cccggcgagc cacgggctgt acgagcaagt cacac 35

<210> 823
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 823
gcgaggacca tcccgcgtgc ctgggagtca ctttaa 36

<210> 824
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 824
atgcacggat gcggctagcg atctccccta aacactga 38

<210> 825
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 825
atgcactgta cgtagagtaa tgcggttcat tccttga 37

<210> 826
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 826
gatcgtaacg ctgcaaggtc actccgggaa aacca 35

<210> 827
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 827
cctgcgcgct aggggagata ctggttcgtg cgatttga 38

<210> 828
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 828
acgcgtgcta ggacggactg aacggtaaaa ctaag 35

<210> 829
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 829
actgcagagt atgcaagctt gaccaccata cgctttga 38

<210> 830
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 830
catgcgcagg ctttctcttt agttctcttc tggttga 37

<210> 831
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 831
tcaagcgaat ctagcggtac acctcgcagg ctcgc 35

<210> 832
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 832
catgcaagct tcgttcaaca tccaacctag ctgta 35

<210> 833
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 833
gcgtgccaac gttgcatcaa ctattttcat gaatt 35

<210> 834
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 834
gcgtgctgtg actccaccgt gccttatact ttgaatga 38

<210> 835
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 835
catgcgcaag cttcaacgac acaactaaac ctgcta 36

<210> 836
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 836
ggctgaccac gatgaagttt gcgcaagctt ttatta 36

<210> 837
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 837
aacgcgagcc tgtacggtac tgcccctatc ccaccga 37

<210> 838
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 838
gcgggcctac gactgcagga cccctgaccc attct 35

<210> 839
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 839
gctaactgcg tgtcatttcg gtttcctaca tccata 36

<210> 840
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 840
ccggatctag gagagtactg gctgcacatg cgacaa 36

<210> 841
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 841
tacgcgcgac taacattaat gcggtcttat catcca 36

<210> 842
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 842
cgccgcgcca gggtactgca ggaagcgcta cgacttga 38

<210> 843
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 843
gcagtcagca acgcttcttt atgtcatacc cgctt 35

<210> 844
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 844
catgcgaaag cttctcatac ttgatcacgt ttccc 35

<210> 845
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 845
cacgcggttc gtagcggaat gtactggtca atccta 36

<210> 846
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 846
ccggggctag gaacgggggc agtcagagcg accata 36

<210> 847
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 847
ccggcagtct tatctctact gtgccttctt taatc 35

<210> 848
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 848
gagttgcgag catgcggacg cttttgtttt gcttttga 38

<210> 849
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 849
tacgcaggag ggccatgacg ctgactatac gcttttga 38

<210> 850
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 850
ttctgcgcaa tttcaaccct ctactgtgcc caatttga 38

<210> 851
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 851
tcataccctc atgatcatgc agtcaccact cttag 35

<210> 852
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 852
gcggagctag cacggtactg cgcataaact atgattga 38

<210> 853
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 853
gaccggaact aggtacggct gcagagcgca accgc 35

<210> 854
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 854
tcgtagtcac cgcgggtcat ccactattaa caccttga 38

<210> 855
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 855
acgcgacagc tacgagtccc cgcggagtgc aacgc 35

<210> 856
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 856
tcacctacta gggaaggcca cacggcacgc acatta 36

<210> 857
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 857
aaccggaact aggtgtggga cctgaaggca tggacga 37

<210> 858
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 858
gtaaactgcg ctcttcttcg ccctcccctg tctcgga 37

<210> 859
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 859
tcccatcgcc ggacgctcca tatctttcgg atcct 35

<210> 860
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 860
tgcctctacc gtgccgtgtt gattttctcc attcta 36

<210> 861
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 861
gcgggccgtg actgtttacc actctactgt gcgattga 38

<210> 862
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 862
gcgatccatg acgaccaata tataccgctt tgaca 35

<210> 863
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 863
tgtactctct actgtgccct tcctgagtgt ttatca 36

<210> 864
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 864
taaggcgaaa ctagctgaag gtactgcgaa ttttta 36

<210> 865
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 865
atgcacggta cggaaggatt ctgccttccc atcacga 37

<210> 866
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 866
ggtcgcgtgc caggcacgga gatgcggtcc tttgca 36

<210> 867
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 867
aacctgaagg aaccgcatct ctaccgtgcc tatat 35

<210> 868
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 868
catgcaggct ttccaaccca ctttaatagc taact 35

<210> 869
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 869
tcatgccagg ctttttcttt gaattccgat ttaata 36

<210> 870
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 870
gcgttagatc tctaccgtgc tcaacctctt ctgtaa 36

<210> 871
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 871
gtgacgccgc tatctttaca actctactgt gcttta 36

<210> 872
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 872
tattgcgaac tcctctgagc tacggtacgt gtggca 36

<210> 873
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 873
tctatcaact cgccccaaat tacccgcccc acata 35

<210> 874
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 874
catgtctagg tctttcgtaa tcctctaccg tgcca 35

<210> 875
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 875
ctgcggtcag tgcccatccg ctcccaacta gcctta 36

<210> 876
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 876
ccactccacc gtgcccttct ataggttttc tcaactga 38

<210> 877
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 877
gccttcccac gtcgagtcag ttcgacgtcc ccgtt 35

<210> 878
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 878
tcatcctttt actcgaccgt gccccccctta gttcctga 38

<210> 879
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 879
taaaggccac gcatcccccg ttaaaacatc attaaa 36

<210> 880
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 880
ttcttctcca ctgtgcccca actcaatttt tgatta 36

<210> 881
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 881
tctccaccgt gccatcgaaa atttactctc tgctca 36

<210> 882
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 882
taacctacta aggcatgtcg actctatcat atgcttga 38

<210> 883
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 883
aaaggcatga catatatcct ctaccgtgcc cgccta 36

<210> 884
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 884
catgccagat ttcgcactag ttacctcctt agctta 36

<210> 885
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 885
catgccggag cgatttattc aatatcttct atgagtga 38

<210> 886
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 886
atcgtagtac aaaagggggt tatctcttgg tttcg 35

<210> 887
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 887
tagacatatc gttctggact ttttctccac tgtgc 35

<210> 888
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 888
atcgtagtac tgtggtccgc cactacccga actatga 37

<210> 889
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 889
gaagagcccc gcctgcttcc ttatccagat catgcga 37

<210> 890
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 890
gcgcgatacg acatctcctt agcttttgtt ttgct 35

<210> 891
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 891
gacgtcatgg ccaccctagc ttcatttctt gcttt 35

<210> 892
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 892
ccggtcgtag gactaccgcc aggcatatta ttgtttga 38

<210> 893
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 893
tgtggtagta cgcagtcagt cgctgtatcc cgctaa 36

<210> 894
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 894
tatgcgcccc tggtctatat tcggaaaacc ctacctga 38

<210> 895
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 895
ttaggtatca tgcggccacc tggtctaata ttttt 35

<210> 896
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 896
tcgtagtact cagggcccca ccaaatttct taaat 35

<210> 897
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 897
accgcgcgcc tggtatcctg ctatgtctag tccct 35

<210> 898
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 898
gatcgtagcg tacgtaggaa atccccgaaa caacg 35

<210> 899
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 899
cgtgccatgg ctttaaccgt atcttcgctt tttctga 37

<210> 900
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 900
gcgatcgtag cacggtaaac tggccccctt gcaata 36

<210> 901
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 901
acgcactgta taaagtctac gcgcccttca accata 36

<210> 902
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 902
cttacgcgtg cttggaacga tgtattgcgg ccaattga 38

<210> 903
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 903
tcgtaggact gcaaaccaca cgccttgttc gaaaaa 36

<210> 904
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 904
ccgggtctag gacggctctg ttgtctcctc cctgt 35

<210> 905
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 905
atcgtagcac tgcagggcgc ctagtttaca atggg 35

<210> 906
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 906
gacgtcatcg tgccttcttg gctgaccgta cagga 35

<210> 907
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 907
ttcatccgcg cgaatgaaac cgatctacat ccctatga 38

<210> 908
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 908
atcgtagtac catacccgc atgccgccaa aattt 35

<210> 909
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 909
aaacccgcga gcaggcattg aactcccctg aatacga 37

<210> 910
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 910
gcaatgacgt gcgttcgacc gattatctct actatga 37

<210> 911
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 911
aagcactgta cggaagtagg cccgccccca aattaa 36

<210> 912
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 912
tcgtagtatc tggaggtcgt gcgttgcact attac 35

<210> 913
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 913
gatcgtatac gtacgctagg tctctttcgc ttatca 36

<210> 914
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 914
tgggcaaccg gaactagtac agcccacgta cgttga 36

<210> 915
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 915
tcgcaaactc agcttcctct ctaccgtgca tgttaa 36

<210> 916
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 916
gcgcgctaac accctgcggt acaccaggaa ctccttga 38

<210> 917
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 917
ccggtcactg ctggagtgcg aaagctttct cttgt 35

<210> 918
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 918
gcagaaggaa atgcgcgcta attgacaacg ctcgaa 36

<210> 919
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 919
accgcgcgcc tgggaccgta atggatccta cataatga 38

<210> 920
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 920
gcgatcgtag ctaagggatc tactagggaa tgccca 36

<210> 921
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 921
ttacctacca gggatgcggc cccgaagcca ttctt 35

<210> 922
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 922
taggcaacta atgcagagac gcgacccacg acatttga 38

<210> 923
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 923
ctcctagcag gcgggccctg acaggcggtc gtagc 35

<210> 924
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 924
tcacctacca cggtcatctt tacttgagac aacaa 35

<210> 925
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 925
taagcaactc ctgccgtgcc attaaagctt ttata 35

<210> 926
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 926
taccgctgcc atgaatcatc tctgttaaac tagcca 36

<210> 927
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 927
gagtacgcga cgccttggtc tactactctt ccctgga 37

<210> 928
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 928
gcggtcgtag caacactgca tcctcatgat actta 35

<210> 929
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 929
tccgacgggg catcgtgtca ggcgcatgaa atttt 35

<210> 930
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 930
atcgtagtag tgccgcgcta gcctgacgca aaatt 35

<210> 931
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 931
aggcgtgcat gacgatttaa ttaaccaccg aaccatga 38

<210> 932
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 932
catcatacga ccgcgtgatt ggtgagttgt aacgga 36

<210> 933
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 933
catgcaagct tggattttct cacgtttcat tgtta 35

<210> 934
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 934
gcgcataggc tttttatatt tattctctat gcttt 35

<210> 935
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 935
gcccgattcc agaaaccacg cttttaccat ttcct 35

<210> 936
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 936
gctagcattg gcttttgttt atttgcttct atcttga 37

<210> 937
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 937
gcgctactag ccgatatatt tccgttccac tcttt 35

<210> 938
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 938
gactaaatac ctcactatct ccactgtgcc gttaaa 36

<210> 939
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 939
gcaaattctg ttcgatccac actattaact cgctt 35

<210> 940
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 940
gaatcatccg gccctaaact ccactgtgcc ctcttga 37

<210> 941
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 941
tgctaggtga tgaattaatt tggtctcttt ttggt 35

<210> 942
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 942
gcctacagga tttttgattg atcgatttcg cttct 35

<210> 943
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 943
catgcaattc ttcgcttcct attccttaag cttgta 36

<210> 944
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 944
gctaactgct tggcgtgcga gctgaaccaa cgctta 36

<210> 945
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 945
gccgaaaaca tcgctgttcg tcttatcaac tcgccc 36

<210> 946
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 946
atgcctgtcc aaaaactcta ccgtgccgtg ttacta 36

<210> 947
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 947
catgcgtcga ttcactatct tcgctttcaa ttgctga 37

<210> 948
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 948
gtcaaatgtc aatgggcacc acgcactgtt cgcgc 35

<210> 949
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 949
gcgacacgat tggtcaattc ctcgctttcc tctcta 36

<210> 950
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 950
catagggacg gaacgggcac ggtagctcct aacgta 36

<210> 951
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 951
taggcaacta ctaggtgatg cgtcaattac ctctta 36

<210> 952
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 952
tcacctacca caagatcgtg cccctgaatc acttgtga 38

<210> 953
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 953
agtcaggcat ggtagacgct tctacttcgc tttcta 36

<210> 954
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 954
taacctacca agtacatgac ctttcacttg atcttga 37

<210> 955
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 955
tatgccagct tctcatatat cccgctttct atcaatga 38

<210> 956
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 956
gaagccgcgc gccaagcaaa gggtactgcc cctct 35

<210> 957
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 957
atgcacggtg cggccggtca tgcaagacac ttcgta 36

<210> 958
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 958
ttcatttatc tccaccgtgc actccatgga tctta 35

<210> 959
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 959
gcaggcttcg aatcctccac tgtgctaata catgaa 36

<210> 960
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 960
atgcactgtt gtgatcggcc tgagcaggcg cccaa 35

<210> 961
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 961
aacccctcgc cgcgtgcctg gcgacggagt agtaatga 38

<210> 962
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 962
actccaccgt gcccttaat ctcatgtgcc acactga 37

<210> 963
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 963
gcgattatat taatttctct accgtgcgat cccaa 35

<210> 964
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 964
atgcatcagc gcgcttatta atcaatcgct tttgt 35

<210> 965
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 965
gcgtgccagc gtactcgcat ctcacagaac ccgt 34

<210> 966
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 966
caaccactat tcgcgtgcca gggtggggaa acgtt 35

<210> 967
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 967
tatgcaaccg ggactgagag tcctagcgct ttttta 36

<210> 968
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 968
gcccgctttg tcaccggaat taattaactc gcctca 36

<210> 969
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 969
tcacctacta gattcgtgcc agaatttcaa atttc 35

<210> 970
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 970
cgaaagccat gacctccgat cactc 25

<210> 971
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 971
cctcaagacc taagcgacag cgtgaccttg tttca 35

<210> 972
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 972
caagtttcca ggtagtttgc gtgttttacg taccctcaag acctaagcga cagcgtgacc 60

ttgtttca 68

<210> 973
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 973
cagggtaggt atcgtgagtg ttttacgtac tcatcctctt cttttcttgg tgttgagaag 60

atgctc 66

<210> 974
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 974
gaaagtacag gaaaatgcct gctgttttac gtaccacaat tctgcgggtt agcaggaagg 60

ttagggaac 69

<210> 975
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 975
gagacagagt agggccaatg cgataatgct gttgagatta ttgagcttca tcatgaccag 60

aag 63

<210> 976
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 976
gagacagaga aaaagggata gttcgttcgg tcaaagacgc ctatcttcca gtttgatcgg 60

gaaact 66

<210> 977
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 977
gacgcgtggt gggttttacg taccgaacct aactcctcgc tacattccta ttgttttc 58

<210> 978
<211> 57
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 978
gcgcaggacc ggttttacgt acccaatttg gttttactcc cctcgattat gcggagt 57

<210> 979
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 979
gagacagaaa gtcgcatggg cagtactttc gggttatatc tatcatttac ttgacaccct 60

<210> 980
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 980
agtggagttt aattggcctt cagttacgta ctcaacagcc actttttttc caaattttgc 60

aagagcc 67

<210> 981
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 981
gaacgattcg tagggtatgc atgtacgtac tcaccgtgtg gacggcaact cagagataac 60

gcatat 66

<210> 982
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 982
agtatcaacg ccggcatgta cgtaccctgg agtttatgta ttgccaacga gtttgtcttt 60

<210> 983
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 983
taaagggcgt ggcatgtcta cgtaccagat aaggttgtta ttgtggagga tgttactaca 60

<210> 984
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 984
gagacagaac attggcacag tagagatagg tcttccttcc tgtgttccag ctacaaactt 60

agaaac 66

<210> 985
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 985
taatggcagc cttacgtacc tacgtactca taaaattggt tttgcctttc agcaattcaa 60

ctt 63

<210> 986
<211> 55
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 986
cgggcgcact gactacgtac ctggtcaaga cttgcatgag gacccgcaaa ttcct 55

<210> 987
<211> 57

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 987
gagacaggaa gggcaccgta gactttcgtt gggacgcttg aagcgcaagt agaaaac 57

<210> 988
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 988
ccatcgcgac tgcactacgt actccagcag acctgcaata tcaaagttat aagcgcgt 58

<210> 989
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 989
gagacagtca aggcacagtg gagtaatgcc tgccaatgca ctcgatcttg tcatttttt 60

gcg 63

<210> 990
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 990
ggtagagaat acaagggtct gcaactacgt accaaactgg agagagaagt gaagacgatt 60

taaccca 67

<210> 991
<211> 57
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 991
cggttcgcat ggccatacgt accgattgct gcattccgct caacgcttga ggaagta 57

<210> 992
<211> 60
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 992
gaaggggatg accgcaatac gtactctgag gctgttaaag ctgtagcaac tcttccacga 60

<210> 993
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 993
ccagtgacct gcagtgacgt actctaggac gcaaatcact tgaagaagtg aaagcgag 58

<210> 994
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 994
tggagttcaa caatatcagt tagcagtgac gtacccacgc gatgacgttc gtcaagagtc 60

gcataatct 69

<210> 995
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 995
cctgccacag tacctgacgt actcatttgg aatgatgtgt actgggaata agacgacg 58

<210> 996
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 996
ctggactgct gacctgacgt actcacaaga atccctgcta gctgaaggag ggtcaaac 58

<210> 997
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 997
gaaaaatgac agacagcggt tactgacgta ccttgacgta gattgctatc aggttacgat 60

gactgc 66

<210> 998
<211> 64
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 998
aggttcgacg actacagtta ctgacgtacc ttacagatcg tgtgctcatg acttccacag 60

acgt 64

<210> 999
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 999
cgcatgagct tccttactga cgtactcttg gaggagttga tagtggtaaa acaacattag 60

c 61

<210> 1000
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1000
gagacagtca agtatttaca agtggcgcag cctacgtata tatccaagtg gttatgtccg 60

acggc 65

<210> 1001
<211> 75
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1001
gagacagtaa aataagagag aacgtactag cttcgcatta cagcaagaag gagtatggaa 60

cttatagcaa gagag 75

<210> 1002
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1002
ggaaagaatc gcgatgtttg tactgacgta ccacccctcc aaacgcattc ttattggcaa 60

atggaa 66

<210> 1003
<211> 57
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1003
ggcgtttacg ggcatgacgt accccgaagc aatactgtcg tcactctgta tgtccgt 57

<210> 1004
<211> 57
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1004
gtcgaaaggc ggcatgacgt acccgggaat cggcatttcg cattcttagg atctaaa 57

<210> 1005
<211> 70
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1005
gagacaggag tgaatgagtt tgagtatgga atgacccgat cttcataacg gacaaactga 60

acgggccatt 70

<210> 1006
<211> 64
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1006
gttaatggaa gcttgtcgca tgacgtactc gctatgcaga cgagctggca gaggagagaa 60

atca 64

<210> 1007
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1007
atgataaggg tagctcgcat gacgtactca ttcgcaacca tgtgaagtaa tgtgagcgta 60

ctt 63

<210> 1008
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1008
ggcgcaatcg catgacgtac tcaccagtta gcgtggcgta taccatgttg ttaaca 56

<210> 1009
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1009
gagacagtga taatttaggg ctacggagta cgcctgaatc aatagaacaa tatcagttat 60

ggcggtg 67

<210> 1010
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1010
gagacagatg attagggtag tagagaacgc accggttgtt aatatgacag gccgctaaag 60

acgttct 67

<210> 1011
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1011
gagacagttt gaatcatgag tttgtgaaag cgtccgtctc agatgagtgg gttaatcaat 60

caagtatg 68

<210> 1012
<211> 64
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1012
gagagaatag gacgtacgca tgacgtactc tgacacatta gtaacgtcgg caagcactta 60

gtcg 64

<210> 1013
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1013
tgggtgtgcg tacatgacgt actcgtgaac cagattatgt atggacgcgc aatagata 58

<210> 1014
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1014
ggtttccgca ttgcaggacg taccatacga aatttgagca agcaattgaa ggcttaga 58

<210> 1015
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1015
ccctggctcg cgacgtacct atcagctaat agggtcggct caacagtgta tcc 53

<210> 1016
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1016
cagtagagaa caagtcgcac gacgtaccta tcaattcgtg accccgatca tccagtccag 60

aa 62

<210> 1017
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1017
gagacaggca aagatgaaag cgcatgcttg agctctaggc ccaaaacgac cttaatggtc 60

a 61

<210> 1018
<211> 52

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1018
tgcccgccct cacgtaccta gcccagatcc tacgagatga gctacgtaac ta 52

<210> 1019
<211> 55
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1019
gttggaccgc cctcacgtac caaatgcact ctatatggag ggagagtagc tggat 55

<210> 1020
<211> 57
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1020
actcctgtac gccctcacgt accctggtct aggtatctaa ttcgtgggtc gggtact 57

<210> 1021
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1021
gccattaact tcagtagccc tcacgtacca ttagctcgga tgctatcagc ttgcgcctat 60

tat 63

<210> 1022
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1022
gagacaggat aaaggtaaga aaagcggctg cacgatctgt attttgcacc tttcgctatg 60

ctgag 65

<210> 1023
<211> 64
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide

<400> 1023
gagacagtca ggaaggaaaa tgtacgcacc tgtgtccgtc tatacgcata ctggtccaca 60

tata 64

<210> 1024
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1024
gagacagagg gacaaggagg aaattggcat gttggagtta acggagaccc gccatcgttt 60

ac 62

<210> 1025
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1025
ggacggcccc tcacgtaccg ctcattttga acatacgatt gcgattacgg aaa 53

<210> 1026
<211> 55
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1026
atgaccgccc ctcacgtact cctatgcatc atgtgcctca ctaggacatc atgct 55

<210> 1027
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1027
cacagtagag gaatatgcgc tatcacgtac tcctaaattg ggaaaaaagg ttttagctat 60

tgatgg 66

<210> 1028
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1028
gagacagtca ttgggcacgg tagacttcag ttaaaggcta tcttgctccg ctcgttctc 59

<210> 1029
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1029
gcacggtaga gagtagttgc acgtaccta aagctatcca cgaatgtcaa aaatgtggtt 60

t 61

<210> 1030
<211> 73
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1030
gagacagtat gagataagca cagtagagat tgacatgtcc cgaatgtata atgctgacgt 60

tcttgctttt ggc 73

<210> 1031
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1031
gagacagtag gggcacggta gagcctattg aagcaatcct ctccccaata cttaaaaa 58

<210> 1032
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1032
gggggtgtcc tgcacgtacc tacggttacc gtctttataa gtgaacaaaa ccgg 54

<210> 1033
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1033
gtagagtatc cacgcaaact gcacgtacct ctgtgaactg tcatcggtcc gatcaattag 60

tct 63

<210> 1034
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1034
gcacggtaga gttcatgtat gcacgtacct ccccttccc aagtaaatgt acgggaatta 60

tcg 63

<210> 1035
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1035
gagacagggc acggtagagt aaaataatgt gacgctttat tatgtgttcg tctaactctg 60

tttctgt 67

<210> 1036
<211> 57
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1036
aaagtgcgcc tgatgcacgt acccgagtgc atgagctgtc tttcacatga tacatcg 57

<210> 1037
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1037
ctcgagaacg atactgtggc acgtacctat ttctgttcac ggatgaaggc ctatatcaat 60

g 61

<210> 1038
<211> 55
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1038
ccctctcgca tggcacgtac ccatccactt tcatggaaac aataagagca gggaa 55

<210> 1039
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1039
gagacagtca cgtcatttca gatggcacac aaactcacta ctaccaacaa cctcaccaaa 60

aa 62

<210> 1040
<211> 78
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1040
gagacagtca accttaatat aagcacagta gagttacaag ttcctcatgt cctctgttaa 60

tccagcctga atatgcca 78

<210> 1041
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1041
gcacagtaga gtattggtcg cacgtaccag aaatgtcact cccatggtgg ctgatataga 60

aa 62

<210> 1042
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1042
gagacagagg agtgagcacg gttcatgtcg aaccttggat aggagcgacc gattacgt 58

<210> 1043
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1043
ggttacatgg gctttatcgc acgtactctc aggttgttac ttgaagggtt caacacgagc 60

tc 62

<210> 1044
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1044
ggtagagaca acttgcgcac gtactcagaa gatcaaaaaa cgatccctgt ccatcaatac 60

<210> 1045
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1045
gagacagaat aaagcacggt ggagacaatg cttaggctac caaatgaatt taaagccagc 60

tgaaa 65

<210> 1046
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1046
gagacagtag gaaataggca cggtagagtt ccaatgcttg cagtatgtat cctgatcgtg 60

cgtgc 65

<210> 1047
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1047
aggtcggtgc gcacgtaccc tgcattctca tggaaatgca atggattcat tcc 53

<210> 1048
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1048
gtgtggtggc gcacgtactc ctgttgcagt atcacgtaaa tacctacttc gata 54

<210> 1049
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1049
gagacagaag aaagtaatcg gaagtcgggt agctagctgt tatggctatt gctgaaacag 60

caaaatt 67

<210> 1050
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1050
acagtagagt aatatagtcc gcgcacgtac ccttacgact tcactgcaat tgacgattca 60

gttaa 65

<210> 1051
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1051
gagacagcaa aaaggtggtt gtataaagtg cccctcatac caatgtaaag tatagttaac 60

gccctgt 67

<210> 1052
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1052
gagacaggat ggcacagtgg agtaagcatc tccatgactg cttgagcggc tggagaatct 60

g 61

<210> 1053
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1053
acagtagagg tcaccgcacg tacctttcgc cacccatata aaccccactt cgtcctca 58

<210> 1054
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1054
cggtagaggt tctgtacgca cgtaccaagg cagagcaaat gtgacactgt ctatcagtac 60

<210> 1055
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1055
gggaggggac gcacgtaccc tacatatata ggaaaaggga aggtagaaga gct 53

<210> 1056
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1056
gagacaggaa aatcgcatac gtcttgcata cccttctgga atttcttcct ttgattttgc 60

catttt 66

<210> 1057
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1057
gaggtggatc ggtctagcac gtacctccta aggttgctga tttggttgtt ggagaccca 59

<210> 1058
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1058
acggtagagc taagctagca cgtactcaag gcctagccta aaggttcttg cagagcaaca 60

t 61

<210> 1059
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1059
cggcctgcag cacgtactcc taattagctc taggaaacac aaccccggga ttt 53

<210> 1060
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1060
gaacgcctga ccagcacgta ccttgagtta tacggaactt cgcaaaagta ttccct 56

<210> 1061
<211> 73
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1061
gagacagaaa tgcacagtag agtaaaatat cttgcctgcc atccatcaac aactgctcca 60

acagcctttc cat 73

<210> 1062
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1062
gagacaggca cagtagaggt gaggataccg gcacaagcag acaaaatcaa catggtcatt 60

ta 62

<210> 1063
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1063
gagacagggc ataaagggca cagtacgctc acgtgatcta ccctagctga ccgctaatga 60

<210> 1064
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1064
gagacagtca aggagcacgg tagagctcta acctgcatac atatggcatt tagttgttca 60

<210> 1065
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1065
ggcacggtag agaaatcgca taacgtaccc ttcttgaaga cctatgtaaa gaaacgggtc 60

act 63

<210> 1066
<211> 66
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1066
gagacagagg ataaggtcgt gtaataggtg gccatacgca tgactacatt acaacgggcc 60

aggaag 66

<210> 1067
<211> 64
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1067
gagacagaga agaaagtaag cgttggcacc aggtctcgat ctcgtacaaa acgactatga 60

ccat 64

<210> 1068
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1068
ccggctatga ctgcttgaac gtactcatga acgtgtcgtg ttatgcagcg gtatgtcgtg 60

<210> 1069
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1069
gagacagttt ggtggcacgg tagacgtcgt gtcttagacg actgtgtgtg attctcgag 59

<210> 1070
<211> 58

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1070
gcgaaatgac tgcgcaacgt actccaccgt tttgccagtt ccaccagtaa ctgtaagg 58

<210> 1071
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1071
agagtactac cgcctgacaa cgtactcgcc ctggaacaac ggttatattc tctggcagaa 60

c 61

<210> 1072
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1072
tcggacttga tctacacatt gacacattga taccctga 38

<210> 1073
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1073
cgtttgactt agacggactg ccttgatcac tgactga 37

<210> 1074
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1074
gagttctcct taaaatctga ccctgacaac gagtgga 37

<210> 1075
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1075
atactggagg tcatgattct gatcccgtga tattctga 38

<210> 1076
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1076
ggatacacgg gagtctgcgt tgacacattt tgatttga 38

<210> 1077
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1077
gtttgcaatt gcttaggata ctcaacctga ctccatga 38

<210> 1078
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1078
gcggtattta tgataatgac tcgtgacccc aatattga 38

<210> 1079
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1079
ggtcagctag taaagtttat ccgatgattt gattttga 38

<210> 1080
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1080
gcaggatagt catgcaaaat attcaacgat aattaga 37

<210> 1081
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1081
tgtgactata acgcactaag ccgattctga cgatatga 38

<210> 1082
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1082
ttttgtacct gcttcgataa ccaacagtta ggatttga 38

<210> 1083
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1083
gactatcaaa tgaccccgat cgatacgtag gacactga 38

<210> 1084
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1084
aactggccgt gtgtgccggt gggataactg ttacta 36

<210> 1085
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1085
gaggactagc cgatacttga taatgattat gacgta 36

<210> 1086
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1086
gtgaatgaca cacgtctgac cttggatcgc tgattga 37

<210> 1087
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1087
accgtggact tattgcttat attggactta gagcctga 38

<210> 1088
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1088
gaggacgtac gccttgatgt ttaactggat tactga 36

<210> 1089
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1089
cacccctttt gattatcctt attgatactt atatga 36

<210> 1090
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1090
tcccgtgatg tgggaacgta gatctttgac tgatttga 38

<210> 1091
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1091
tcgtaatgcc caataaccaa ctggttaagt agacctga 38

<210> 1092
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1092
gcgatcacaa tttaccggac cgatctccct agacaa 36

<210> 1093
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1093
gcgtacggaa cggaaccgac acaacctgaa taccta 36

<210> 1094
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1094
gcgtacggaa cggaaccgac acaacctgaa taccga 36

<210> 1095
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1095
gcttgaatcc acgacgaaat tacttaaaat aaccctga 38

<210> 1096
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1096
gccggggcgg agcctacgat ttatatcgcg attattga 38

<210> 1097
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1097
aagtgcacac tgcctcgatt tacatttcgg acctatga 38

<210> 1098
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1098
gcgtactgtg cgcgattccg ggatatatat tcgattga 38

<210> 1099
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1099
cacagacgtt ctcgttgata cccggacccg atatttga 38

<210> 1100
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1100
tccattttgg aacgtcgaat tactcggata ctaattga 38

<210> 1101
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1101
actcgacgta aaactatctg tgactcctaa tacattga 38

<210> 1102
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1102
accgtaactt gctagccttc cggatccctc caatctga 38

<210> 1103
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1103
ttcgtacaga tacccccgat accccgccac ctgattga 38

<210> 1104
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1104
atattaatgc cttgatccga ctccgacctc ttga 34

<210> 1105
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1105
ttgacgcgac gtactatttt gatcgtattc caatctga 38

<210> 1106
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1106
gcgccgccac ggtacttaaa aactcggact ttacttga 38

<210> 1107
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1107
cgtagtactg ctacatccga accgaccata aatactga 38

<210> 1108
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1108
gaggacgcag ttgaccggat ttccaccgat tatattga 38

<210> 1109
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1109
gaggactgaa ccatctcgga ccacccatgg atacttga 38

<210> 1110
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1110
tgcgcgagca aacgcgacta ttgaacctta gaacctga 38

<210> 1111
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1111
gctggtgacc cctcttgacc ttccttattg atttg 35

<210> 1112
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1112
gagtacctca ccgctttcga ttatcgttct gaccttga 38

<210> 1113
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1113
agccacttcc tgctttgaac cggactcctt atataa 36

<210> 1114
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1114
attgagcgta cgcatctacc ggattacctt acttttga 38

<210> 1115
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1115
gaacttacgt tactaacctg actctatctt acattga 37

<210> 1116
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1116
gcaccactat tgtgggaatt cctgatttta ctattga 37

<210> 1117
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1117
gcgtgcctgc ctggctctga ctccgattat tcattga 37

<210> 1118
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1118
agttccttac taacacaact tttggatatt taaattga 38

<210> 1119
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1119
actgcctttt cttactggga tacttattga ttattga 37

<210> 1120
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1120
atagtacact acctactgac tccctatatt atgaatga 38

<210> 1121
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1121
gcgatacaac ccttcatttc tgactaccga tatattga 38

<210> 1122
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1122
gcatctttct tcttaatgat ttaccgttct cacata 36

<210> 1123
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1123
accacgtgtt attctctgta tttcactttg tctcaa 36

<210> 1124
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1124
accgggaaca tttctgtcgt ttcatttcat cactaa 36

<210> 1125
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1125
agtttattgc tccttcccac tattgtctat ctcgca 36

<210> 1126
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1126
tcgcttcctt gtacttccta tacccttttc accaca 36

<210> 1127
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1127
actaactcgc ccactcttct catttggaac ctatc 35

<210> 1128
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1128
actccccctc agctatcata catcactcgc atctaa 36

<210> 1129
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1129
aatatatacg cgctacaaat cttttctcct actgatga 38

<210> 1130
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1130
actatactca catcctattt aattattggc ttaatga 37

<210> 1131
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1131
atcgatcctg cctaacctgg taaacaactc atttctga 38

<210> 1132
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1132
ttagctgctt cccagaccca agctattatt tttca 35

<210> 1133
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1133
tccctgcatc ttaacttcta tagggtttgc ttaaaa 36

<210> 1134
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1134
gggtcactcc gtaaggcctc tactccaaat tataaa 36

<210> 1135
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1135
aactgtacgg cgcaatttta ctctcaagta ccaaca 36

<210> 1136
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1136
tgcggacttt attcgcttga tgcctatcct gttat 35

<210> 1137
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1137
tccgccacat cgggttatac ttttgaacta gttaaa 36

<210> 1138
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1138
tgttccttcg agttccatcc caaagagtta attcatga 38

<210> 1139
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1139
tcctgcgact ctacggtata ttcctttaat cgaca 35

<210> 1140
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1140
agtaaaattc cggttcatcc acactgagct ttcaatga 38

<210> 1141
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1141
ttaactcctt gaagtacgtc ctaggccccc tcacca 36

<210> 1142
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1142
tggagtactg gtgccgaacg tttttgaacg ggttga 36

<210> 1143
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1143
gagtactata gagccaagac ctcatttagt cactctga 38

<210> 1144
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1144
ccgacccata acactttcgc ttcaagcatc taggca 36

<210> 1145
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1145
cgtccgcacc ccttttttcac cttgttctgc tatgca 36

<210> 1146
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1146
gagcctctgt aacccaatct tcaactgcag ggacc 35

<210> 1147
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1147
gagttctcca gctagatatt ccaatacgga accga 35

<210> 1148
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1148
gagtattgcg aagattgggt ttatcggttc taacatga 38

<210> 1149
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1149
aaacgctatt acggctgacc cggtgactcg ggattga 37

<210> 1150
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1150
accgactcca gtttcacggg aatattctca ccggga 36

<210> 1151
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1151
gttgtcggtt tccattagtt cgatatcttg ccgtgaga 38

<210> 1152
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1152
atgggaatca cgtagctgta acgataatcc atgggtga 38

<210> 1153
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1153
aggtactgac gattctttaa agtgactact ggtga 35

<210> 1154
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1154
ggcggtggaa gtccgaactg ccgagttaaa ttccctga 38

<210> 1155
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1155
tatcggtctt acgaaagccg gcacgtacgt ttctttga 38

<210> 1156
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1156
tgacggatga taggcatgac ctcgtagacc acatctga 38

<210> 1157
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1157
tttgtaccat ccttacaaat gagatcttgt tatgttga 38

<210> 1158
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1158
tacatgcgta ctataaaccc tattgtttac tggtaa 36

<210> 1159
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1159
gcaagtctag atcttcattc atttgttcat gtttt 35

<210> 1160
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1160
gagtttgcgg gaatcttccc cgaacaaggg gctaa 35

<210> 1161
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1161
tagaccgttg gaggtccggg gttggtgccc gatta 35

<210> 1162
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1162
acactcatac gagatctgat tatacacgcg ccttgga 37

<210> 1163
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1163
gtcgtcggga gaccacttat ctttttttcca tgccgga 37

<210> 1164
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1164
tcacctgacg ttacactcct ccgtacaaac gtgta 35

<210> 1165
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1165
tcctgattaa ttcatggtca caactcccgg tttcc 35

<210> 1166
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1166
tccttttgcc catgattatt ctcgatgtaa aaacga 36

<210> 1167
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1167
aacgctaata tctgccgatg ggaatcctcg attagga 37

<210> 1168
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1168
agggtgatat ttacgttatc ttcgaatttc tactgga 37

<210> 1169
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1169
tctcacttac aaagcgttta atcaacgcag ttctta 36

<210> 1170
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1170
ccgcaaactt aggggctaat aattttcgtt atccgga 37

<210> 1171
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1171
tgccagaaaa tcgaattttt atacgtaatg gcatgga 37

<210> 1172
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1172
atcgatcctg acgccgaaaa cttgacacga ctaaa 35

<210> 1173
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1173
atgctggtaa cttttgcaca tcttacttgg taaac 35

<210> 1174
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1174
gcgattactc agcacccccc tacttgtttt taatca 36

<210> 1175
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1175
actgcgggac tatctgaatt attggtatta taaca 35

<210> 1176
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1176
gagtactgaa ggaactggaa ccgagtattc ttgtca 36

<210> 1177
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1177
gagaactgcc tggtaatata ttcccaaacc tggaca 36

<210> 1178
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1178
gagtgcgtag gtcttacatc ctatcgaata tttaca 36

<210> 1179
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1179
ctcagacttt acacaatctt tctctccaca caagca 36

<210> 1180
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1180
gccactttca gaatctcgtg tcacacccat catca 35

<210> 1181
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1181
gcgttccccg tggacaatat cactttctta tcgtca 36

<210> 1182
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1182
tcctggtcta cgacttacat ctgatccgac ccaaa 35

<210> 1183
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1183
ggaccttatt tgttcgaccc ccgttatcac ctgaaa 36

<210> 1184
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1184
gaggactttt actgatcaac cccacctttc gaaaa 35

<210> 1185
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1185
gctactactt atccgactac tctaaaaaca actaca 36

<210> 1186
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1186
aactgtcatc cgccgactaa ctcaatctaa ctcaa 35

<210> 1187
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1187
aactgtcatc cgccgactaa ctcaatctaa ctcta 35

<210> 1188
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1188
aactgtcatc cgccgactaa ctcaatctaa ctcga 35

<210> 1189
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1189
tactccggcc caactatata cttctccaca ccaac 35

<210> 1190
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1190
tactccggcc caactatata cttctccaca ccatc 35

<210> 1191
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1191
tactccggcc caactatata cttctccaca ccagc 35

<210> 1192
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1192
tcctacctgc ccaatcccct tttgattaca tatat 35

<210> 1193
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1193
ccacgctttg gggacgctat tactatccct ttcaa 35

<210> 1194
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1194
aatgtactga taatcccttt tacactggtt actat 35

<210> 1195
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1195
ttagtcgtac ttttatctat attgggccta ataaa 35

<210> 1196
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1196
ttcgccacgt tcctaaatat tttacttatt actaaa 36

<210> 1197
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1197
ttccatgact ctgcctatac ttattatatc tcataa 36

<210> 1198
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1198
ttggagctac aatcaaactt catatctctt tctaaa 36

<210> 1199
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1199
tgcccaaaac tgcctttaga aacttattac attcaa 36

<210> 1200
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1200
tctgcgatcg ccatttatct cgacccttta cctaaa 36

<210> 1201
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1201
aacttttttg cactaacctg gacctcatat ataat 35

<210> 1202
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1202
aaccgctttt cacactacca ctcacgctct cata 34

<210> 1203
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1203
ccgccgtacc cacgctccaa ttattctcat cctaca 36

<210> 1204
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1204
tatctccgtc caacacatct ctggtcacca cctcca 36

<210> 1205
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1205
tttctccgct catcataact gacgccatcc actca 35

<210> 1206
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1206
acttcgctca ttcacttgcc aactcctgca caatc 35

<210> 1207
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1207
tccctcaccg ttgcccttat tccgtactca cttatga 37

<210> 1208
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1208
gtgcttctcc ttttgactcg cctcactctt ggtcca 36

<210> 1209
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1209
ttttctcctc actccgctcc gtgccttgac atcaaa 36

<210> 1210
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1210
taccttcttc aatatccgtc cgcttaacct ggattca 37

<210> 1211
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1211
cctcctctgc ctaaatacgt tttctccacc atcata 36

<210> 1212
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1212
actttttaa attttgttcc cacgtatcct taatta 36

<210> 1213
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1213
ttttccgggt ttcttctcca gtaggttgac ttttatga 38

<210> 1214
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1214
tgttttggcc tccacggttt agtacttatt ttttt 35

<210> 1215
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1215
gacttttttg ctcatatttt caccaatcaa ttcca 35

<210> 1216
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1216
tgagttctct tttatattgg tcgccacttt ttccaa 36

<210> 1217
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1217
ttccttcact tgggctttca aatcatttat aaacaa 36

<210> 1218
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1218
ttcttgactt cagttcgatt ttgttattct tagtttga 38

<210> 1219
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1219
gttaattcct tgctcgagtt tttccgctta ttccttga 38

<210> 1220
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1220
ggtacatttc gtccctacca ttggtttatt cttttga 37

<210> 1221
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1221
attatttttc gttctctcac ttgttttcga tcattga 37

<210> 1222
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1222
taattgagct tttgctccta acggcgagag ccgtt 35

<210> 1223
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1223
tgatatacgc gcgagtagaa catcaaccag tgagaa 36

<210> 1224
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1224
taagccgagc gcatgtaggt tcgttttcac gagagtga 38

<210> 1225
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1225
atgaacgcga gtgaggacgc tttagttaat tatgt 35

<210> 1226
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1226
actcggatag cggggccggg ggaggcctac acattga 37

<210> 1227
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1227
acgcagtccg taaattgata accgggaggt tcgagtga 38

<210> 1228
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1228
agtcccacga gagtggttct ctctagataa gagga 35

<210> 1229
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1229
gagtaccata acaaggcctg aagaaccttg caggtga 37

<210> 1230
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1230
gcggggtgtg aacccgctcg tatgtaacaa cggattga 38

<210> 1231
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1231
gagtacggtt acggtgagta accatttaga cgtaaga 37

<210> 1232
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1232
tgggatccgc gaggatctga tggggttctt atgtttga 38

<210> 1233
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1233
gagtacccgg gcagggagcc agttattatg ttttga 36

<210> 1234
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1234
gagtacctag atatctcaaa agatatttat cgttaa 36

<210> 1235
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1235
gagtactctg cgcaagtgtt gaactttgta gaaat 35

<210> 1236
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1236
gagtacttcc cgcggggtta tggaatgaat tggata 36

<210> 1237
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1237
taatgtctct tccgccgtag tacgagcggt tccgctga 38

<210> 1238
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1238
gagtacagcc gcccttgtac cgaggcctta ttaagtga 38

<210> 1239
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1239
ttagtttacc gtcaaaggcc tcataatgtt acaactga 38

<210> 1240
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1240
cgtagtattg cgggctcgtc tcattgcatt cctattga 38

<210> 1241
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1241
cgtgagacgt ggttgcccac cacagtgaat aaatttga 38

<210> 1242
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1242
gctccactat ttaaccgtga gttgttaacc gggggga 37

<210> 1243
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1243
gctccagtac cgctgtacga attatttaag aaaacga 37

<210> 1244
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1244
gcccacattt ttttcgagtg tgtaaggggt gagaaa 36

<210> 1245
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1245
gcttattccg gatgttgcct gcaattaggg gactcga 37

<210> 1246
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1246
gcttgtctcg cggttttgaa agtaatcttt aacgaga 37

<210> 1247
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1247
gcgcgaaagc tcaaataaaa ctaaagtaca acaaa 35

<210> 1248
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1248
gctaagctgg gagaaaatca caccctcgat cgtaaa 36

<210> 1249
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1249
ctcacattag ggctcggtaa atataacgat ctgacga 37

<210> 1250
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1250
actgtagcta ttatgaaagt ctgggttatt ggcct 35

<210> 1251
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1251
aatcgtatgg agcgttgaac aaatgaaccc gtagt 35

<210> 1252
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1252
agatatactt gaaaggtggt attatgaatg ttgggtga 38

<210> 1253
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1253
atggtgactc aattaaaatg tgaggtgggg gagctaga 38

<210> 1254
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1254
tcgtggaggg tattgacctt ctaatgatgg gggggga 37

<210> 1255
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1255
ggttcctatg aaggacacgg ggggggaggt atcaatga 38

<210> 1256
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1256
aaggggcggt agactttagg ggctgaatgg ggtggga 37

<210> 1257
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1257
gagtgcactg gatgaatttt gaggggggaa ggaccga 37

<210> 1258
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1258
tgcggtgggt ggtgggtggg aaataattat gtgggga 37

<210> 1259
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1259
taccagactg gaagtcaatg atcatggttg tctccga 37

<210> 1260
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1260
aacggtacaa catctgctag agaactgcaa cttcta 36

<210> 1261
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1261
actatatcca atgctatgcg cgcattcagg gtagaa 36

<210> 1262
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1262
agcatcggcc gatcgcctca atcacgtttt cgatt 35

<210> 1263
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1263
aggcttacaa tacgttaccc cttcttcgta gacggtga 38

<210> 1264
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1264
cttgacgtca catttgccta acactgtagg tcacaa 36

<210> 1265
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1265
tctggtagca catgagtttg gcattcctct ctgca 35

<210> 1266
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1266
tgttgtccgt atatattcct tgtgtgttaa ttccg 35

<210> 1267
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1267
gagcactata agtgtgtatc tatttgtaac tttct 35

<210> 1268
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1268
tgtggaggca ctcttgtatg attgatttcg tatct 35

<210> 1269
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1269
aatctaacgc atcaaatacg tttgggtagt cattaa 36

<210> 1270
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1270
tggtcaacgt tatcgtatcc tactgatcac taact 35

<210> 1271
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1271
ttgtcgcacc acatcatagg tgccacgatg tcccaa 36

<210> 1272
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1272
actgttacaa cagaagggtt agatcacata ctacaa 36

<210> 1273
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1273
actcgaggga actaggaact actacaacag tgtgt 35

<210> 1274
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1274
ccatgaaact acaacatatc aagtcgagtg tggtgtga 38

<210> 1275
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1275
cggccgacgt agctcaaatt ctacactatg gttgtga 37

<210> 1276
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1276
ggcagtacta tcaacttaca tgccaatatt cgatgtga 38

<210> 1277
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1277
gctaccagcg ttgcagtctg cggattctgt caaata 36

<210> 1278
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1278
cttactcaaa gtggcgtgta ggtcattagg ttgta 35

<210> 1279
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1279
cgtagtacta gcaacattag ggtcgtaaca taacta 36

<210> 1280
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1280
acgaccgtgg ggttagggcg catattcttt ggctta 36

<210> 1281
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1281
tactctcatg ggtatctttc tgaatcgaac gttgtga 37

<210> 1282
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1282
gcggtttaac ctcgaaagta agggcgatta ggcga 35

<210> 1283
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1283
gcgggttcaa attgaggagg aaagacgaaa gctct 35

<210> 1284
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1284
gatcgaatcg atactgcagc ggagtgacac gggtctaa 38

<210> 1285
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1285
tagtccgggt cgttttaaat tcagtttggc atttt 35

<210> 1286
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1286
aaccaggcat agttggccgc aggcaacccc atttt 35

<210> 1287
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1287
agtccaaaca cgaggttccg tcgcgagaac tttaa 35

<210> 1288
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1288
agtccggcaa ggggcgcaac ttatcatagt cctctga 37

<210> 1289
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1289
taggctcccg cctttagctt agtttcagat agcaa 35

<210> 1290
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1290
cgcggtattt tgcctcaaac tagtcgttaa tctat 35

<210> 1291
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1291
ttacttcggg agctaggctc acccctacat tagtgga 37

<210> 1292
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1292
agtgcaattg gatatatcag tgaggagttt caattga 37

<210> 1293
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1293
ccgaaatcgg ggtagattat ggggtcctgg aaatcga 37

<210> 1294
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1294
taagtttgta gaatttaacg tatggaacgg ggacgtga 38

<210> 1295
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1295
tgagcgagcg cctgatattc ggtactgcat atctc 35

<210> 1296
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1296
gatcgaatta ggggaattcg ttggaatcga cccata 36

<210> 1297
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1297
tctgtgcgaa gggtttagtc tcgggcgaaa cttaatga 38

<210> 1298
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1298
agggaagtcc tagttggagt gattttccgt gattctga 38

<210> 1299
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1299
aggaatggct agtctaagac aaactcctct gtttatga 38

<210> 1300
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1300
gattgtcacg gatgattctg gatttatata ggtacga 37

<210> 1301
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1301
attcatggga gcttgaccgg agttattgtt tgttgtga 38

<210> 1302
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1302
tatcgaactt agatgtgaca ctgtttttta gtcta 35

<210> 1303
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1303
agctacggct tatcgttggg gtccaagttc atgaaga 37

<210> 1304
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1304
agtctacccc ccaaattgtt ggtattccga gcataga 37

<210> 1305
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1305
tcgtcccaag ctctggactg gtaagttctt atgta 35

<210> 1306
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1306
tgctacagtt caacctcacg tttttgaact gcgtga 36

<210> 1307
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1307
cgtttggacg ggggtgtcct ttttttgat attacga 37

<210> 1308
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1308
ttcctcgtgt tacaggcact ttattagcct gtgggga 37

<210> 1309
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1309
cttggcccgt tgaccacgac ataatggtct tatata 36

<210> 1310
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1310
aggatattta ccctgcatct tttcgttacg gtatga 36

<210> 1311
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1311
actgtctgga agtcggaact ttcaccatta tgatga 36

<210> 1312
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1312
gaatggagga actttctcgt attcacgttt tgttgtga 38

<210> 1313
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1313
aaccgttagt gcaaagggga gtacggcctt tgttttga 38

<210> 1314
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1314
ctcgtacgca aacgctttga atagaacgtt tgtcttga 38

<210> 1315
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1315
agatacccgg tggttatctt aagggtgtga cttttga 37

<210> 1316
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1316
tagggttcag tatctggcga ttctgttgat gttcctga 38

<210> 1317
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1317
cgagctccct taagggtgca tcgggattca ttgtttga 38

<210> 1318
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1318
cacatccgga gaaggtgcgt ttatacatga atttctga 38

<210> 1319
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1319
aaggacgacg cggaacaatc tgtagctatg actcttga 38

<210> 1320
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1320
acttgttcct gaagtgggtt tgacttgcaa acgttga 37

<210> 1321
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1321
agtacacgcc cgttggtttc gttcggaaat tacttga 37

<210> 1322
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1322
tgaacatctt tgttattcaa cgggatcacg caacgga 37

<210> 1323
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1323
taatccgatg caatttgtat ttactgacaa tgcgta 36

<210> 1324
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1324
aactggagga tcccggaatt tatcctccca tacgaa 36

<210> 1325
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1325
gaattgcgga ctgacagata gatgcggtga cattgga 37

<210> 1326
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1326
tcgcactagt tctatgaggg aaactacagg aacga 35

<210> 1327
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1327
gattgacctg gaaggcaata gctggtcaac gactgga 37

<210> 1328
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1328
gcagtacggc acgagggata ctctgcccta tatatga 37

<210> 1329
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1329
gccatcatgg atatatgcac gtatggtgct tttagga 37

<210> 1330
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1330
tgcgaaactc tagatgatta tgattaaact tgtactga 38

<210> 1331
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1331
aactgacccc attgggtcac ccgtgatagt tatcgga 37

<210> 1332
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1332
gtactgctct aggagaaaat aaccgtatga aacgatga 38

<210> 1333
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1333
ttgcgtattt gccttagaac gcaataaagc atttgtga 38

<210> 1334
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1334
atctacactg gaactgacag ggggtagtag tatcctga 38

<210> 1335
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1335
tagggattga atctcctcta acattcgaca acaggtga 38

<210> 1336
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1336
agcgttcaac gcggttgagt gggtgccgaa ctatatga 38

<210> 1337
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1337
ggacctgatg ccatttagtg tggataatgt tattgtga 38

<210> 1338
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1338
tgatatctag gaaccatgcg gtaatggtcc gagta 35

<210> 1339
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1339
tgaaagccgt gactggtatt aacggcaaat aacggga 37

<210> 1340
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1340
tgcatgatac gggcttcaag gagcttctta ttcgta 36

<210> 1341
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1341
agacacattg tgtgcaacat ggaagtggtc tttagga 37

<210> 1342
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1342
gagttgctct atgtacttcc tgaggggtga tgtgatga 38

<210> 1343
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1343
cctgttatct atacttgcgt gagcaaatta ctttaga 37

<210> 1344
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1344
tggagtacgc gtgcgttaca gtccgaataa ggctttga 38

<210> 1345
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1345
cacagtcagg cgtaatgaga atgagcactt tggtttga 38

<210> 1346
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1346
tcatttacgt gagtaacctg gcttagtatc ggagta 36

<210> 1347
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1347
gatggaacgt agcgaaaccg ggtttacgcg tgatctga 38

<210> 1348
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1348
taattatcgt cgtgagatcg atactacggt tgtgcga 37

<210> 1349
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1349
gaggactggc attaccgtgg tcagctaaac tagac 35

<210> 1350
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1350
tgcagcatac ctaattgttc accgttggga aaggtga 37

<210> 1351
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1351
agctttgtaa tacgtgagtt tgggtaggtc atcgta 36

<210> 1352
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1352
cctgcgtcct aaacggataa cgaaaacgaa tactc 35

<210> 1353
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1353
ccctgcacga acgaaccgga ataaatcgat ggtga 35

<210> 1354
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1354
gcgcttttct caaccgacga ccatcgaata gaagaa 36

<210> 1355
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1355
ttaggctaag gaacgggacc cattcgatga aaatca 36

<210> 1356
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1356
tcggagtttg caatccggca attattagat tacaca 36

<210> 1357
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1357
accggaaatg aaccgtcgct tattccacga attag 35

<210> 1358
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1358
gagaattgcg gtaaatttcg gggcaatatc ctctc 35

<210> 1359
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1359
acgccccgaa tgatcggtgg gctgtatttt acaata 36

<210> 1360
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1360
gagtagtggg ctcacggggg cagtatatcg gatacga 37

<210> 1361
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1361
gagtgcgatg gcgtccacgg ggatttttcg gaataa 36

<210> 1362
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1362
accacgtaca aggaagttag gattcgggga ccgtga 36

<210> 1363
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1363
gaaagctcaa gctaatgaga atacgatcat acacc 35

<210> 1364
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1364
aggacgacga atttactcta ggggtgggct ggttaa 36

<210> 1365
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1365
gtacactggt gtgattgttg tggtggagtt ggtaaga 37

<210> 1366
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1366
gggtagtacg gggagcactg ggctagattg tgacta 36

<210> 1367
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1367
ggtggaacgg gggttgggat tgtaagggag tatgt 35

<210> 1368
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1368
acggtaaaaa gggcaatgct tggcgtctat cagacga 37

<210> 1369
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1369
gtgtggaccg gctagtaaag gctgttgggg taatt 35

<210> 1370
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1370
gcctatcggt ggatagtttg gtcattcatt ggagc 35

<210> 1371
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1371
actctccatg ggtggggctc ccgaaagaat ttctaga 37

<210> 1372
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1372
ttaaagccat caaaatctat ctaaagttgt agaagga 37

<210> 1373
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1373
gtctacataa tttgtatacg gtttaaaatc tcgagga 37

<210> 1374
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1374
tctcggagta gaatccacca gaatcacagg aattctga 38

<210> 1375
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1375
gaaaggtgcc aatcgccggt acacaattta ggaaa 35

<210> 1376
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1376
cataaacacc ttatgaattt atgtagaaat cgaaa 35

<210> 1377
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1377
gaaccgaagt agtagattta attatacata atgt 34

<210> 1378
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1378
gccgcaatac aaacgagagg cttctcccga ttaata 36

<210> 1379
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1379
acctgcggaa tatcgtgaga attgaaaaaa gcaaa 35

<210> 1380
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1380
taggcgaatc gaaaaaaagt aatcattatg ggttgtga 38

<210> 1381
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1381
ataccggacc aaggacgccg gggtactaag gactcga 37

<210> 1382
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1382
cgcataatag tcgacgtttt tttttgtcgg attcgga 37

<210> 1383
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1383
acaggaaggc caaagttttt ccaaggtcga ttgcctga 38

<210> 1384
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1384
gaaatcgttt ggtaatttag ttcggaaccc agggta 36

<210> 1385
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1385
aaaacgtctc atgcggcctt tcgagggcat atctc 35

<210> 1386
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1386
aaggttcgag aattctcact gcagatcatc cgtcatga 38

<210> 1387
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1387
agtgctctct cttgggggta gtcatccaga atagt 35

<210> 1388
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1388
taagtcccaa tttatttgaa gacgttgtaa tcgtc 35

<210> 1389
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1389
aaaacaggct aaaaaggtac atccccattt tgctta 36

<210> 1390
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1390
gtgcgcaagg tttagcgcat acgtagttga ataccga 37

<210> 1391
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1391
ttccaagtac ccaggtttag acctgggacg atagcga 37

<210> 1392
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1392
tttccaccaa ccctgatttg aaaaagacac gcaaa 35

<210> 1393
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1393
ctccgcccaa cgagagttct acgtaaagat caaac 35

<210> 1394
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1394
tgtcccgcgc ttgacatcat cctaagatta aagaaga 37

<210> 1395
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1395
attaccgtta atttaaagta cccacgggtt gttgg 35

<210> 1396
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1396
gcatcgtgga aagattctct tataatcaat tggggga 37

<210> 1397
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1397
acacacacgg ggtagatcac ataggcaaac gcttg 35

<210> 1398
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1398
acaaacgtac taagggcgaa gccttttact cgcgca 36

<210> 1399
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1399
cgtgcaaacc gctgtaagca ttcaaaccaa ctggatga 38

<210> 1400
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1400
acacaaacac aatctaggag ttaaatacag ttgga 35

<210> 1401
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1401
actttcacga tgttgggggg gcccaattac agatg 35

<210> 1402
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1402
acgtttagat taagttcgct tatgaatgtg ggatt 35

<210> 1403
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1403
attgccgttg atgttcgtga ttttattaag tggggga 37

<210> 1404
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1404
gttcttaggc ctgttgttac caggattttc gtgtaga 37

<210> 1405
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1405
agtccggata cctgtttgta tgcgaaaggg aattta 36

<210> 1406
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1406
gttaatgtca atctgaaccg acgtgagtat tcggc 35

<210> 1407
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1407
accacggttg ggttgcggct aacgtatagc cgatgga 37

<210> 1408
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1408
gtgccttggt tgtaattgga ctttatggct acctaa 36

<210> 1409
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1409
gatttgcgag tgtgaaacga attatagcac acgtctga 38

<210> 1410
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1410
atgctggtga accaaactca agcatgacat ttaag 35

<210> 1411
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1411
cctgcccacg aattgtatat gtaattgcac tatcga 36

<210> 1412
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1412
tctgcaggac ttttgattcg tgggttcaac ctcat 35

<210> 1413
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1413
gcacattatg tcacgtagag gcacagtcgg taatcga 37

<210> 1414
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1414
gggctacgtt atcccattca attcgaatta aatgta 36

<210> 1415
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1415
acattgccga attaattcgt tgtcgaggaa agacaa 36

<210> 1416
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1416
cccacgtgtt acgggtttta tcatgacatc actat 35

<210> 1417
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1417
catgcacgtt ttattgtgga aatgtaagct tgggc 35

<210> 1418
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1418
tgcaagggaa atacgttcct gtgttgtttc gatgaga 37

<210> 1419
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1419
tgggcagtct ctatttgtca ttacggtttt atgtc 35

<210> 1420
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1420
agcaggcaac agtatcagcc ctatttgtcg taaaatga 38

<210> 1421
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1421
ggctacggga aagatgtcac gtctgtttca tttgatga 38

<210> 1422
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1422
actacgtttt aatttcagta acccatccga tgctc 35

<210> 1423
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1423
aacaaacgcc aaccatatgg taggattcct ctttta 36

<210> 1424
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1424
tgataaccat cggtacctca ccgctcccaa ctctta 36

<210> 1425
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1425
cgaaccgacg aaatttccac accgctttgt atgtta 36

<210> 1426
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1426
tacctcatgt ttgtcactat ggaaatcatc ccgac 35

<210> 1427
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1427
agttatttct cgccacataa cactctatgg aaacta 36

<210> 1428
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1428
atctcggatt tcccgtaggt cactggcaca tttcta 36

<210> 1429
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1429
aattttccgc tgggacgaat ttctaaccct atttgga 37

<210> 1430
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1430
gaagcaccga atcacctggc atggcccttt atcta 35

<210> 1431
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1431
actgccgacg cacaacttgg gtggggctct aattt 35

<210> 1432
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1432
ttcgctgacc tcaccggaca tatcatttaa agcta 35

<210> 1433
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1433
tcccttatta tgcgaaacaa atatgtagtt cttac 35

<210> 1434
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1434
tctaaaactc acgctaggtg gtacaaccga ttcta 35

<210> 1435
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1435
gtagccaacg aagctcgcgt attctatttg ccctta 36

<210> 1436
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1436
acctcaccaa gtccttaagg attctgtatt cataaa 36

<210> 1437
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1437
agaatctggc tattgatccc cacacattgc gaaaa 35

<210> 1438
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1438
acctaccaat ctagatccgc ttttatggat tgctc 35

<210> 1439
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1439
gacttctaaa cacatggggt gcccgcaaca tattt 35

<210> 1440
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1440
tccagaacgt cagctctcca tacgcaatta attcatga 38

<210> 1441
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1441
gacggcctta taaggcactc ctcattcatt taata 35

<210> 1442
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1442
gggacctgca agcctatcaa cgggttatac atttgtga 38

<210> 1443
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1443
tacacttcca tttgatgcgc tcaccttgaa tctgta 36

<210> 1444
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1444
aatcgaactc attactttcc aatggacctg ggagaa 36

<210> 1445
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1445
agtttacgag ttacactaca acggacgaat ctttta 36

<210> 1446
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1446
aacttttctg gctttataca catacggttg ccatt 35

<210> 1447
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1447
agaactgggc cttggaatca cacacctaac tagtta 36

<210> 1448
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1448
attcgcacta ataacatgtt gctccattca acctaa 36

<210> 1449
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1449
tatctgctgt tttctctggt gccaataact acatc 35

<210> 1450
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1450
ttgcacataa ttttctgctt taaccaagcg aacccga 37

<210> 1451
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1451
taaaattctg aaaatacacg tgtctttcgc attac 35

<210> 1452
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1452
tcccgaacac atcccccttt taagggcttt gctaa 35

<210> 1453
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1453
aaccacacgg ggctctcatc gatacttcta gtgattga 38

<210> 1454
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1454
atctagaatt ccgtttgcgc gagtaacttc cctgt 35

<210> 1455
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1455
cccgcttatc gctagtacct catttagaat tggaaa 36

<210> 1456
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1456
acctgtaaat ttggctcact tttcctttac actgga 36

<210> 1457
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1457
aatagtatca cacctatctg gcttcctgta cccgt 35

<210> 1458
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1458
acccaccctt gagcaccatt aaatagtgat ttcgctga 38

<210> 1459
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1459
tcacgaaccc tttagctcta ttttgtttga cgcatga 37

<210> 1460
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1460
gaggaacgtt tctcccctca acatttaaat aaacaga 37

<210> 1461
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1461
atcctacgca taattctcga ggatctattt cgaaa 35

<210> 1462
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1462
tgcccgcaat tacccctaga tccacagctt ggttta 36

<210> 1463
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1463
ggctcttata caaaacattg ggttctcgac ctccctga 38

<210> 1464
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1464
gtgagctgct aatatttatc ctgaacttgc ttacga 36

<210> 1465
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1465
agttacctta caacgaggtt tcttgaatgg attct 35

<210> 1466
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1466
ttacaccatc cctacgcggt aatagcaact ctgtt 35

<210> 1467
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1467
gcaccacgca cctgtgggga ttcccctaatg tcttta 36

<210> 1468
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1468
ctatcggact aatacacgta aaaataactc tggtt 35

<210> 1469
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1469
ggcgtactta tcacaaacgg tggaaccaca cttct 35

<210> 1470
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1470
tattcttatc caacccacga aatcaattgt attttga 37

<210> 1471
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1471
agatccgaat ccgtaactcc aaaatattta tgcac 35

<210> 1472
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1472
actatagaac aatccaccgc tttaaatcaa ctatctta 38

<210> 1473
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1473
gccgttcttc tgcgacgaac acatcctccc ttaat 35

<210> 1474
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1474
gcgcgaggcc cgcgggctca tcacttttgt tatct 35

<210> 1475
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1475
taccctgac aatcctcgcg ccgaggcctc catcctga 38

<210> 1476
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1476
ttcggtctaa tttttagcgg ccaaattttc ttatca 36

<210> 1477
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1477
gattttaaaa cccttgcacc tgattgtgcc aatcca 36

<210> 1478
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1478
gcgcgaggcc cgcgggctca tcacttttgt tatat 35

<210> 1479
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1479
gccgttcttc tgcgacgaac acatcctccc ttact 35

<210> 1480
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1480
tgctcggccc tattaacacg tgccgtgtct tgttca 36

<210> 1481
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1481
tcgcgccgac cgtattacac caactctatg gactctaa 38

<210> 1482
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1482
aacccaattc acatacactc tcacccccac taaaca 36

<210> 1483
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1483
gccgttcttc tgcgacgaac acatcctccc ttatt 35

<210> 1484
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1484
gcgcgaggcc cgcgggctca tcacttttgt tatgt 35

<210> 1485
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1485
taccctgac aatcctcgcg ccgaggcctc catcctca 38

<210> 1486
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1486
ttccgcgcaa acgaatgact ctttgttgaa aaacga 36

<210> 1487
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1487
gcgtagtggc tgcttggcgt ttccgcatac taggca 36

<210> 1488
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1488
aaacctcggc taaccgccaa gggtagtgtc ccccgca 37

<210> 1489
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1489
atgagacgta gcaacactcg agacccttgg agcactca 38

<210> 1490
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1490
agctgccagg gttcgggacc ttaacggatg tcccaa 36

<210> 1491
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1491
tcacgagctt ccactacgtc tatccgcgta gacggaa 37

<210> 1492
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1492
gatcgacgat tcgatcatcc aacgtattta caccaga 37

<210> 1493
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1493
gcagccggcg accgggatgt cgtcgggact tgctta 36

<210> 1494
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1494
atgaaccatg gttacacctc taatgtagcc tcatgtca 38

<210> 1495
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1495
ccaccggtac ttattattcg ccatccacac ccagtta 37

<210> 1496
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1496
tagttacgcc actactctgt gggagaaacc cgaga 35

<210> 1497
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1497
atccacccaa aatcgatata agtgcgcccc caatctaa 38

<210> 1498
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1498
agggggggaaa gtcatcgggt ggctgtacta cttca 35

<210> 1499
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1499
gggggtcgta ctcccgagat ttgcgttacg gtgaaca 37

<210> 1500
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1500
gagtatcctc tggtcgttcg gtacagaaac ggtgtaa 37

<210> 1501
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1501
acatggcctt taaatctacg gagtactgtt gggggga 37

<210> 1502
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1502
gcgtagtggc tgcttggcgt ttccgcatac taggaa 36

<210> 1503
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1503
tccgataccg aacttcactc cgtgggcggg gttgaa 36

<210> 1504
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1504
gcagtgttat ttctcttggt ttctttgatt tcacttaa 38

<210> 1505
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1505
acatccagat acccttgact taagattgtg accactaa 38

<210> 1506
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1506
acaggttctg cgaccagaaa gttaggacta acctaga 37

<210> 1507
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1507
gcagtgttat ttctcttggt ttctttgatt tcacttca 38

<210> 1508
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1508
ccgcagtttt gttcccgggg gaatccggac tgaacga 37

<210> 1509
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1509
aggtccggtt cctatggact gcctgacccc gcgaaga 37

<210> 1510
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1510
ttgtaacgat ttgggagtcc agagactcgc tgttt 35

<210> 1511
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1511
tttgatttaa gtactcccgc gttcctccga ataccca 37

<210> 1512
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1512
gcagagcgca cgtgtactca atgggttgtt aattttaa 38

<210> 1513
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1513
gaactctgta ttattcctgt cattagccgt tggtaa 36

<210> 1514
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1514
gaggccctac cgtgtatttt gataatggat ttgtttaa 38

<210> 1515
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1515
tgtaatcacg ggtgcgaact cgagagaatt tgtgctaa 38

<210> 1516
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1516
ggccggtagt ccacgctttc gcgagatagt gactgtga 38

<210> 1517
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1517
aggtcctcag aacttgtgcg tgagcaaaga atgaag 36

<210> 1518
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1518
gagtatctgg gttcgagagc gcatttgccc gctgttga 38

<210> 1519
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1519
gtattaatct tcgctgtggg agagcgatta cgcca 35

<210> 1520
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1520
tatccttgga tcaccgatcg atgcgagagg tagtaa 36

<210> 1521
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1521
ttttctacgt gacggagagt ccagatcgaa gacaaaga 38

<210> 1522
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1522
acaatgcatc gtagcgggtc tcggagctcg ccttaa 36

<210> 1523
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1523
tacaaagaac ggaagctatt caaaaacaca cggagga 37

<210> 1524
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1524
ggacgaccgg atatagtaaa gtccccggct tctgt 35

<210> 1525
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1525
agccatcatg ttggcacggt cctttgatac cattgga 37

<210> 1526
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1526
attaccgtca acgggtcctg ggctccatac gaggca 36

<210> 1527
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1527
ctgctttagg tgtggtcgca ctcaaatgtg aatgg 35

<210> 1528
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1528
gcggggtaag aactcgatga ggtgagtgga tttgta 36

<210> 1529
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1529
aagggattgg aacttggatg ttttcagttg gcgca 35

<210> 1530
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1530
tgcctcggct agcatcacaa tagatggttt gcacttga 38

<210> 1531
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1531
gagacgaagg cacctttcg acgcgataag ctcgttga 38

<210> 1532
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1532
accgcgaggt cttttggaac agggcatatg aatcca 36

<210> 1533
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1533
acgtgaaact accgagcggt gcttttcgct gtagcga 37

<210> 1534
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1534
caacgcctga ctactgtaga gtgaacgctg gtggtga 37

<210> 1535
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1535
gcgaaggttt gggtaaaagg aaatagactt tgttgtga 38

<210> 1536
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1536
acaagtcgct gctcagcttc ctagggcagt actcaa 36

<210> 1537
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1537
aaacgggatc tgtcgcgagt acagtgttgc ccagta 36

<210> 1538
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1538
taaaggaaca gcgattcggg ctgttatttt aggtga 36

<210> 1539
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1539
tgtgggatta ttgtagttgt agtaaggtga ccggca 36

<210> 1540
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1540
cgctaaccgg aactaggttg caaagctatg gtggta 36

<210> 1541
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1541
cccttaggtt ccagtgaagt tatgcaactt catgta 36

<210> 1542
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1542
ctattccagc ttttttgatc ccgcttttga gagttga 37

<210> 1543
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1543
acaggacgtg cgcgctaacg ttctattagt tgtgctga 38

<210> 1544
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1544
gagtttcgtt ttatgtgttg gctttatttc cctcga 36

<210> 1545
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1545
tggcgttgta gccacgtccc tccgctagta gagaaa 36

<210> 1546
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1546
tcgtagtact ggtcttaatt tgttgtagtc ctaag 35

<210> 1547
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1547
tccacacgcg gcgagctacg cgactaaaat cacgta 36

<210> 1548
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1548
tatcaaattc ggtcgtagta tttgtcgccg taacgtga 38

<210> 1549
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1549
gctacgaagt ccaaacgtac gcacctgttt ggagta 36

<210> 1550
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1550
tccgggcacc atatctccac tgtgcgaatc actat 35

<210> 1551
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1551
tccgggcacc atatctccac tgtgcgaatc actct 35

<210> 1552
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1552
acaggacgtg cgcgctaacg ttctattagt tgtgctga 38

<210> 1553
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1553
agcaaacgat gggccatcat tcacgcttgg tattaa 36

<210> 1554
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1554
agggacgtga tatttcaatg ctgaatatac acacctga 38

<210> 1555
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1555
agtgattctc acttgttccc ggggttcaaa gatctga 37

<210> 1556
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1556
atcaagggag gatttttcgg tcggcctagg gacaca 36

<210> 1557
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1557
atacgcggcc taataataag atcgtcatac cccaa 35

<210> 1558
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1558
atctcacgat aaagctaacg gtgagagtat ctact 35

<210> 1559
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1559
ccgccggcga tgggggatcg cttattttat tgttcga 37

<210> 1560
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1560
cgggtacgtt ggatcgacac cgaccatctt gacgta 36

<210> 1561
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1561
cgctaaccgg aactaggttg caaagctatg gtggta 36

<210> 1562
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1562
cgagtgctgc gcgggttcct tggggtgctc caatttga 38

<210> 1563
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1563
cgtgtaaccc gcccaattcc atcgtcacga aattt 35

<210> 1564
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1564
ctattccagc ttttttgatc ccgcttttga gagttga 37

<210> 1565
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1565
cctcgaacct gcgcgggaga aggcgcttgc taatctga 38

<210> 1566
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1566
cttggtccct gctcggttga atgacgcaat gagtgtga 38

<210> 1567
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1567
ctcgtaggac tgaggtcgat agttatttca caccgtga 38

<210> 1568
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1568
gacacgacgt atctccttat tccctgctat aagtttga 38

<210> 1569
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1569
gacgttcttg caccgcgagg tctgcagcaa ttctaa 36

<210> 1570
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1570
gcaactattg ggccgtattc gtccacgggg cttgt 35

<210> 1571
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1571
gcgtagggtt ggttatgcat caattttccg ctaattga 38

<210> 1572
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1572
gccttcgggg tttattaagt tattacggat ctagta 36

<210> 1573
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1573
gcgcgtgagc tgcaccgaca tcacgccgat tctctga 37

<210> 1574
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1574
gctacgaagt ccaaacgtac gcacctgttt ggagta 36

<210> 1575
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1575
ggtcactcgt aggtctcccg catcgagaaa gccgta 36

<210> 1576
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1576
gtggatggca ctaatatccc cttctcgtta aagtgtga 38

<210> 1577
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1577
gtccaccggg attcgtatgg gtatttttac ccacc 35

<210> 1578
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1578
gttacggtat gcgaggtctc acgttgcgat cattttga 38

<210> 1579
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1579
gttcatcgtt cctatgtcac tatttcccta ctgctga 37

<210> 1580
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1580
tactaggggc cgtgagcctt gtgccccaag tagccga 37

<210> 1581
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1581
tatcaaattc ggtcgtagta tttgtcgccg taacgtga 38

<210> 1582
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1582
tccacacgcg gcgagctacg cgactaaaat cacgta 36

<210> 1583
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1583
tcgtagtact ggtcttaatt tgttgtagtc ctaag 35

<210> 1584
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1584
tcgtgcgctc ggttctacgc cactgtacga gacacga 37

<210> 1585
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1585
tgtacgaacc gggcctagga tagttgctac atgta 35

<210> 1586
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1586
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 1587
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1587
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 1588
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1588
actccatggg ccccccggtt tggaccccgg gttta 35

<210> 1589
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1589
gaaaatctgg cgacttagtt ttcgtttgat ccctta 36

<210> 1590
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1590
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 1591
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1591
aggataactg gtttttatct ctgctaattc actcg 35

<210> 1592
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1592
cgtggcctgt ttggagacgt actgtgtcag gcatttga 38

<210> 1593
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1593
accttgggat cctgtttaga ccgaatcggt gcgcta 36

<210> 1594
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1594
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 1595
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1595
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 1596
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1596
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 1597
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1597
agatcttttc ggttattggg gggaaccgag acgta 35

<210> 1598
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1598
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 1599
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1599
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 1600
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1600
gcctcgacga cagtgaccta catcgctcgt ttcgt 35

<210> 1601
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1601
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 1602
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1602
gttgccttgt accccatagc tggtaaggtc ctagta 36

<210> 1603
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1603
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 1604
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1604
tcgttatagc ctatatacac gcggttgact cgctga 36

<210> 1605
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1605
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 1606
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1606
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 1607
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1607
taattggccc gtttccaaat ttatagattc ctaca 35

<210> 1608
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1608
tccgacccat atcgacgata ctggcgtggc gctactga 38

<210> 1609
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1609
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 1610
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1610
gctggtatag ctgggtttct atcgtttatt ctcgta 36

<210> 1611
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1611
tttgcttcat ttgcacttcc cccgttattc cgctca 36

<210> 1612
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1612
aatttttact gcgtggaaag gattcgggaa tttgttga 38

<210> 1613
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1613
accttgggat cctgtttaga ccgaatcggt gcgcta 36

<210> 1614
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1614
aggataactg gtttttatct ctgctaattc actcg 35

<210> 1615
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1615
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 1616
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1616
aggagtacgc gttcatccag acacggaata ctttga 36

<210> 1617
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1617
agccttacgc catgccgaga gagcgctttg tttcga 36

<210> 1618
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1618
agggcttagt ttcctgattt tactacagct cgctc 35

<210> 1619
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1619
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 1620
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1620
agtcctgatg ctcaatgcgc gagtttcatt gagct 35

<210> 1621
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1621
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 1622
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1622
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 1623
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1623
cgtaagccgg tttcagccca cctggagatt cttca 35

<210> 1624
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1624
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 1625
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1625
cgtagttact cgtcatgggg tctttgttat gtggta 36

<210> 1626
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1626
gaaaatctgg cgacttagtt ttcgtttgat ccctta 36

<210> 1627
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1627
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 1628
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1628
gagtcccctt ctttgatgct ggggtgagct cctccga 37

<210> 1629
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1629
gatgatcgcg cctttcgttt gtgacggtta ccaggga 37

<210> 1630
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1630
gcctcgacga cagtgaccta catcgctcgt ttcgt 35

<210> 1631
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1631
ggcccccgct ccctagtttg agcgccctcc acgattga 38

<210> 1632
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1632
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 1633
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1633
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 1634
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1634
gttgccttgt accccatagc tggtaaggtc ctagta 36

<210> 1635
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1635
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 1636
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1636
tactggggaa gctccgggcg cttggtttcg ccacctga 38

<210> 1637
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1637
tcatttcgga ctgttctaac gcgagttatg aaggg 35

<210> 1638
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1638
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 1639
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1639
tcgaggggaa ttgtccacat gcatggtttc ataagga 37

<210> 1640
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1640
tccgacccat atcgacgata ctggcgtggc gctactga 38

<210> 1641
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1641
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 1642
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1642
tgttcttact aggcttagca gcctgttccc gttattga 38

<210> 1643
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1643
tgttttcaga ccagagtagg aattccgagg gcggtta 37

<210> 1644
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1644
tgttttgctg cttagagcgt aatcgtattg gctct 35

<210> 1645
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1645
ttcctaggat gaaacagaaa caccgattgg tccagga 37

<210> 1646
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1646
ttccggtttt gggcgactcc gtttgcgccc acgctga 37

<210> 1647
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1647
tttgcaagtt cgctacgtat atccactgtg aagcc 35

<210> 1648
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1648
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 1649
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1649
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 1650
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1650
ttttcgcatt aaaaacgggt gtccaccggt tggtgga 37

<210> 1651
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1651
cgcgcgagca actacactaa cggaaaaccc gttcaa 36

<210> 1652
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1652
ctactcgccg caacccaaca ctcagtcaaa cgatca 36

<210> 1653
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1653
ataacgcacg tttccataac tcataccact ggccttga 38

<210> 1654
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1654
gagagaagta tttgttaagc cacatagttt tcttg 35

<210> 1655
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1655
ccttactgca tggtttggta ccgctattgt ccccca 36

<210> 1656
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1656
aaaattatgc gccttaggga agacatacgg cctgtga 37

<210> 1657
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1657
aatgatcgga cttgcaggct caaccctact gatggga 37

<210> 1658
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1658
aaggcacgta cacactatca cctggggtgt gcctc 35

<210> 1659
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1659
aatagattaa agtgaacgtg agtgtttcca tatta 35

<210> 1660
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1660
aattcgtcac tcgttgtgtg tagccggtgc ccaattga 38

<210> 1661
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1661
acgggagatc ctttaagtac gtcgaagcca gcaata 36

<210> 1662
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1662
actccgttac gtgttggcgc tttcagttca gcacgga 37

<210> 1663
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1663
agacagtgga cttctcgtcg aaaatgcatc tctat 35

<210> 1664
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1664
agagaacgtt gcccttgagt gtttcttgag ccttgga 37

<210> 1665
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1665
agcgtgtatc atcactgatt tagttatccc gagcta 36

<210> 1666
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1666
agaggccagc ttagtgtggc caccgtgagt ttacta 36

<210> 1667
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1667
atatgtatac gaaccctcct aatcgcacta tcagt 35

<210> 1668
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1668
ataacgcacg tttccataac tcataccact ggccttga 38

<210> 1669
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1669
agtacctatc aaggattttg ccgaatccgt gccgca 36

<210> 1670
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1670
agttcactct aacttccgtg ccgaaggcgt tatcc 35

<210> 1671
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1671
cacatcgact gcgcgatcaa tattagggtg gacat 35

<210> 1672
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1672
attgcattct tcaggacgct tcgcccggga ttctt 35

<210> 1673
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1673
catccctgac gcgagttgga acatgagtac gtact 35

<210> 1674
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1674
caccgcttag tcttatttag ggtcgctaat ctggta 36

<210> 1675
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1675
caaaccgctg caaaaggatc cccgttatgg ctacat 36

<210> 1676
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1676
ccggaccatt aggtcttctt gttcaaatac tttgatga 38

<210> 1677
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1677
ccgagcttcg tttgggttca tttcgtccga ttcgca 36

<210> 1678
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1678
cctcctatta cagttccccg ggaatgaata gcaactga 38

<210> 1679
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1679
cgcctatacc gcactcgctt ggaacgacac aaagctga 38

<210> 1680
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1680
cgtggattcg cccacatgat gatggccggt atgtttga 38

<210> 1681
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1681
ccttttgtag gacgcccttt ctgcgcatcc ggtagga 37

<210> 1682
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1682
ccttaactac acgagagcat gtcaaggggt aaatgga 37

<210> 1683
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1683
ctactcgccg caacccaaca ctcagtcaaa cgatca 36

<210> 1684
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1684
cgagtgctgc gcgggttcct tggggtgctc caatttga 38

<210> 1685
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1685
cggagtacga agccgcttag attacctagc ccgtta 36

<210> 1686
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1686
cgagcgacgt ttgtttcaaa aatcactggg aatatga 37

<210> 1687
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1687
ctacatcgct ctccgtaatc gggcgattca tttca 35

<210> 1688
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1688
gaaaacgttg gaagttttta cgccgctctc agccgtga 38

<210> 1689
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1689
gaatttatgc aatccattag gaaaatgggt gtgcg 35

<210> 1690
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1690
ctgacagtct tccctggtct attgtccgca gatcga 36

<210> 1691
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1691
cttggtccct gctcggttga atgacgcaat gagtgtga 38

<210> 1692
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1692
gagggactgc acagcggtga tattctaaat gtacga 36

<210> 1693
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1693
gaccgggtag gcttctaggc gccccctcc ttacttga 38

<210> 1694
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1694
gaccaaagta tgtgtggtat tgatatgccg tgaac 35

<210> 1695
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1695
gagaaccgtg ctcaaggcag aattgtcacc gctcga 36

<210> 1696
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1696
gacgagtgac ttgcactacg aaggccctcg tcgcctga 38

<210> 1697
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1697
gagtttgtct tgccccgggt cgggacgttg tatccta 37

<210> 1698
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1698
gagtggctta ctttcattgg atccgtgagc cttgc 35

<210> 1699
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1699
gcaatacgct agataggctt agaaacggaa tacacga 37

<210> 1700
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1700
gatgtcgtca cggaagtaca tatggcgctg tagcc 35

<210> 1701
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1701
gccactggat ttaactgcat cccgtctctg aggct 35

<210> 1702
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1702
gcaactattg ggccgtattc gtccacgggg cttgt 35

<210> 1703
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1703
gcaatatggg atatgtgtaa ctcagtgaga caggaga 37

<210> 1704
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1704
gattacataa ctcgtacgtc gcttgtttta agtttga 37

<210> 1705
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1705
gcgtgctgct ttgtactacg gtctcatcgt tgtagga 37

<210> 1706
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1706
gccctggcac caaagtcgca caatcaatcg tgtgc 35

<210> 1707
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1707
gccgttggta ttctgtcgtc ttcctttgtt ctatca 36

<210> 1708
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1708
ggcactgtgc tgcgatcagg aaaagtcaca taccgga 37

<210> 1709
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1709
ggcccttgaa ttaacatgat gaatatcaca acgggtga 38

<210> 1710
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1710
gggtgtttcc tgcttatgac tacttttctt tacgca 36

<210> 1711
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1711
ggctcgtatt ttcccagtat cgtttccggt acgtc 35

<210> 1712
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1712
ggggatctcg ggcgatgaag cctagctatt gtact 35

<210> 1713
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1713
ggtcccgtat tccagatata ctaacggtta ccacc 35

<210> 1714
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1714
gtactgtccg tgtgaacggg tctctgcaga aattcga 37

<210> 1715
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1715
gtgatcggta ctgttgaact tgttgcaccg cccgctga 38

<210> 1716
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1716
gtgtcacatg gggtgaccgg attgaggaga ggatatga 38

<210> 1717
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1717
gtgtcctttc tgggttgatg catttacggt ttggtga 37

<210> 1718
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1718
gttcggttgg ttcccgcgct gaacaatatt acctgga 37

<210> 1719
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1719
gttgggaccc gattaaccca acagaattcg gaataa 36

<210> 1720
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1720
tacctgagcg ctggagctcg ttagtcggaa taggaa 36

<210> 1721
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1721
tacccgttca ccggatagga tgttcgtcct taatt 35

<210> 1722
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1722
taccgatcaa cccagcaagg cttgggtgga ttgaatga 38

<210> 1723
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1723
tccccccgtc gcttacggtg ttggacgggc tcatta 36

<210> 1724
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1724
tcttcataaa tgactaacgc ttcttagtga tgttga 36

<210> 1725
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1725
tgccatacct acccagagtg agacttctag cctgga 36

<210> 1726
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1726
tgaggtccct tttgtcgggg tatatccatg aggtttga 38

<210> 1727
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1727
tggcttggcc cgttcgttcg tgatcaagat tttttga 37

<210> 1728
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1728
tggtctgaaa ttcccgtgag gagatagatc atttctga 38

<210> 1729
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1729
tgtccaaatc actcattctc gtaaggttac cgtcc 35

<210> 1730
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1730
tgttctgtct tatgccatga agttcttata agtaga 36

<210> 1731
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1731
ttacccctgt aatgccctac atgatgttcg gttcatga 38

<210> 1732
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1732
ttgtaccagt gcgtatgttg tctcatcaat gtttt 35

<210> 1733
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1733
ttttagcaac gggagttgca gcggggttgt taaacga 37

<210> 1734
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1734
aggataactg gtttttatct ctgctaattc actcg 35

<210> 1735
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1735
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 1736
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1736
gaaaatctgg cgacttagtt ttcgtttgat ccctta 36

<210> 1737
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1737
aatttttact gcgtggaaag gattcgggaa tttgttga 38

<210> 1738
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1738
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 1739
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1739
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 1740
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1740
attccgtgag gtacgtaagg caaccagttt tattt 35

<210> 1741
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1741
accttgggat cctgtttaga ccgaatcggt gcgcta 36

<210> 1742
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1742
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 1743
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1743
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 1744
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1744
gcctcgacga cagtgaccta catcgctcgt ttcgt 35

<210> 1745
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1745
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 1746
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1746
tccgacccat atcgacgata ctggcgtggc gctactga 38

<210> 1747
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1747
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 1748
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1748
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 1749
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1749
tttgcttcat ttgcacttcc cccgttattc cgctca 36

<210> 1750
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1750
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 1751
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1751
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 1752
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1752
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 1753
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1753
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 1754
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1754
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 1755
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1755
accttgggat cctgtttaga ccgaatcggt gcgcta 36

<210> 1756
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1756
aggataactg gtttttatct ctgctaattc actcg 35

<210> 1757
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1757
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 1758
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1758
agccttacgc catgccgaga gagcgctttg tttcga 36

<210> 1759
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1759
aggagtacgc gttcatccag acacggaata ctttga 36

<210> 1760
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1760
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 1761
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1761
agtcctgatg ctcaatgcgc gagtttcatt gagct 35

<210> 1762
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1762
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 1763
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1763
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 1764
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1764
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 1765
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1765
gaaaatctgg cgacttagtt ttcgtttgat ccctta 36

<210> 1766
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1766
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 1767
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1767
gagcgactca tcccggggtt tggatcagcg ctccgga 37

<210> 1768
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1768
gagtcccctt ctttgatgct ggggtgagct cctccga 37

<210> 1769
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1769
ggtctcttgg tttagatccc gtcttttctg tggaaa 36

<210> 1770
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1770
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 1771
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1771
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 1772
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1772
gtccaggctc tatgcttagt acccgttcat ttgcttga 38

<210> 1773
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1773
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 1774
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1774
tactggggaa gctccgggcg cttggtttcg ccacctga 38

<210> 1775
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1775
tcatttcgga ctgttctaac gcgagttatg aaggg 35

<210> 1776
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1776
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 1777
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1777
tcctagaaat ctgtattgtt tctcaaggcc acctg 35

<210> 1778
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1778
tcgaggggaa ttgtccacat gcatggtttc ataagga 37

<210> 1779
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1779
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 1780
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1780
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 1781
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1781
tgttttatct ggatcacact agctattgta cggtt 35

<210> 1782
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1782
tgttttcaga ccagagtagg aattccgagg gcggtta 37

<210> 1783
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1783
ttcctaggat gaaacagaaa caccgattgg tccagga 37

<210> 1784
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1784
ttccggtttt gggcgactcc gtttgcgccc acgctga 37

<210> 1785
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1785
tttgcaagtt cgctacgtat atccactgtg aagcc 35

<210> 1786
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1786
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 1787
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1787
tttgagcgcc tcagtttgcg cgacttccag cccgtga 37

<210> 1788
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1788
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 1789
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1789
ttttcgcatt aaaaacgggt gtccaccggt tggtgga 37

<210> 1790
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1790
ctactcgccg caacccaaca ctcagtcaaa cgatca 36

<210> 1791
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1791
ataacgcacg tttccataac tcataccact ggccttga 38

<210> 1792
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1792
atgctatccc gtgaaacgag aacatacggt ttttttga 38

<210> 1793
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1793
gagagaagta tttgttaagc cacatagttt tcttg 35

<210> 1794
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1794
cgctaaccgg aactaggttg caaagctatg gtggta 36

<210> 1795
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1795
agggcaatct gtcccgccct cagctttttt atttaa 36

<210> 1796
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1796
agtgattctc acttgttccc ggggttcaaa gatctga 37

<210> 1797
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1797
acaggacgtg cgcgctaacg ttctattagt tgtgctga 38

<210> 1798
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1798
accgaaaacc cgttcccgcc gtactgcttc tatattga 38

<210> 1799
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1799
gctacgaagt ccaaacgtac gcacctgttt ggagta 36

<210> 1800
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1800
tccacacgcg gcgagctacg cgactaaaat cacgta 36

<210> 1801
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1801
tccgggcacc atatctccac tgtgcgaatc actat 35

<210> 1802
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1802
tccgggcacc atatctccac tgtgcgaatc actct 35

<210> 1803
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1803
tcgtagtact ggtcttaatt tgttgtagtc ctaag 35

<210> 1804
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1804
tttccgatta ttatgaccgc ttttattatt ggaggtga 38

<210> 1805
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1805
tcttggacgt gcgtgatccc cattttatat atgattga 38

<210> 1806
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1806
acaggacgtg cgcgctaacg ttctattagt tgtgctga 38

<210> 1807
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1807
agcaaacgat gggccatcat tcacgcttgg tattaa 36

<210> 1808
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1808
cgctaaccgg aactaggttg caaagctatg gtggta 36

<210> 1809
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1809
ctattccagc ttttttgatc ccgcttttga gagttga 37

<210> 1810
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1810
gctacgaagt ccaaacgtac gcacctgttt ggagta 36

<210> 1811
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1811
tccgggcacc atatctccac tgtgcgaatc actct 35

<210> 1812
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1812
tcttggacgt gcgtgatccc cattttatat atgattga 38

<210> 1813
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1813
tcgtagtact ggtcttaatt tgttgtagtc ctaag 35

<210> 1814
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1814
tgcccttagt ttccgtatga ccactttgat tcttttga 38

<210> 1815
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1815
tttccgatta ttatgaccgc ttttattatt ggaggtga 38

<210> 1816
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1816
attcgcatag cctatgggaa gggaattgtg aggggtga 38

<210> 1817
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1817
gaggttccct gtgggtcatg tcgcacgatt tttagtga 38

<210> 1818
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1818
acaggacgtg cgcgctaacg ttctattagt tgtgctga 38

<210> 1819
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1819
tatcaaattc ggtcgtagta tttgtcgccg taacgtga 38

<210> 1820
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1820
tgcccgcttt tggtttatat ggcttcttat ctttcga 37

<210> 1821
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1821
tttccgatta ttatgaccgc ttttattatt ggaggtga 38

<210> 1822
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1822
tccgggcacc atatctccac tgtgcgaatc actat 35

<210> 1823
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1823
aaacgtgggg gcgtcccagc cccatgtgtt atctctga 38

<210> 1824
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1824
aaccctctcc catcctctcg aaacactaga ttacctga 38

<210> 1825
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1825
aacatacgtc ccaagcgcat ctcgattcta tcttc 35

<210> 1826
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1826
aaccacgcct aaatggattc acttcctcaa aaattga 37

<210> 1827
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1827
aacgccgcac ttcgttaaaa gtctataaaa ctctaga 37

<210> 1828
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1828
aacgaagtat tataaccctt gtgtatcccg accatga 37

<210> 1829
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1829
aacaaagttc acccatacca caaccccaat tcgac 35

<210> 1830
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1830
aacggctggg aggggtagca ttttaactat atttctga 38

<210> 1831
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1831
aagtggtccg tttcggcaat aagtacaacg ggatctga 38

<210> 1832
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1832
aactctgcat aactaaccga cgccaactat tatcctga 38

<210> 1833
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1833
aatcaggctt gtttgctaga caaaataagc acgaaa 36

<210> 1834
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1834
aattgcgttt tccggttgac taaaaacacg gaatctga 38

<210> 1835
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1835
aataaacact tcgaccaaac taaggcctta tctcca 36

<210> 1836
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1836
aagtttcctg gtcgcccgct tctgaaaagt tagtttga 38

<210> 1837
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1837
aagagggcta tgatttagcc ggtgcctcct tgctttga 38

<210> 1838
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1838
aagccggtat cgcccgtaac attcccttcc tctagga 37

<210> 1839
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1839
aacttcaccg atacggttct tttctctaac ccgcttga 38

<210> 1840
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1840
aaggatagcc ataggctgat ggggactcaa caggtga 37

<210> 1841
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1841
aatcgtctta cgtgcccttg ataaatccca tactaa 36

<210> 1842
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1842
aattcgcgcg aattgaagag gcccatggcc cgctctga 38

<210> 1843
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1843
accatgaatc accgcgcctg ctactgcaac ttttgga 37

<210> 1844
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1844
acactggcat ttttcgaaac atagaatgaa tgggaa 36

<210> 1845
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1845
acacccacaa gctttaactt atcataactc gccccga 37

<210> 1846
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1846
acccaacgtt gagcgcaata ccccttcata atacc 35

<210> 1847
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1847
acctctctct gaaatttaac tctattctcc cattc 35

<210> 1848
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1848
accggactag atatttacca cattgcattt ccata 35

<210> 1849
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1849
acgctgaatc tggttcacaa cggggaacgt gttag 35

<210> 1850
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1850
acccgctgtg tgaaccaccg tgccttcgga cgtcatga 38

<210> 1851
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1851
accgctctct gaaatttatc tctattctcc cattc 35

<210> 1852
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1852
acggtacact cactttccta aatccctctt tatcta 36

<210> 1853
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1853
acgccccagc ctatttacct tggagtttct ttgactga 38

<210> 1854
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1854
acccggcgaa gacgttagct ctggcgctcc gttca 35

<210> 1855
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1855
actatcccag gtttacatta gtgtgccccg cattctga 38

<210> 1856
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1856
actgcttttt ttgcccaata tctcgagtaa ccccctga 38

<210> 1857
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1857
actgaacact ctacatgacc ataaacacct ctctt 35

<210> 1858
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1858
actcgggctt ccatcaagct tctgattggt gaggtga 37

<210> 1859
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1859
actgaacact ctacatgact gtaaacacct ctctt 35

<210> 1860
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1860
acgtttttct cgctcgtaat tttatcgtta atttctga 38

<210> 1861
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1861
actactcatc cccccgtctt aacaacgatc ccactga 37

<210> 1862
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1862
actacgcgct catcggccag agttccggtc aaccgga 37

<210> 1863
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1863
acttcaccca cacaataccc ttatcgataa ctcaaa 36

<210> 1864
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1864
acgttacgcc gggggactat aagcagttca gtagca 36

<210> 1865
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1865
agacctattc cccaatcctg aattacaaca tctcta 36

<210> 1866
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1866
agcacccgcc cgcttctatt cctcgatcct acact 35

<210> 1867
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1867
agccccagag tttatccggc ttggccggca attcc 35

<210> 1868
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1868
aggaatcgat cctggtggtt ttagctttga ccctt 35

<210> 1869
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1869
agacttagac tcaataactg gacccttac acaccgt 37

<210> 1870
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1870
agtgattctc acttgttccc ggggttcaaa gatctga 37

<210> 1871
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1871
atacacgctg cccatttcca ctcgtcaaac cactatga 38

<210> 1872
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1872
agttcgcgca acccttctc gcttgtgtgg cttaa 35

<210> 1873
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1873
atcaagccgg tgtaggcgct ccaggccttt cagcc 35

<210> 1874
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1874
agtcatccat tacctcaatg ttcatctatt ttcac 35

<210> 1875
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1875
atcctctacg ttccaaccac ttacttttat gccattga 38

<210> 1876
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1876
atcacactct aactgtaatc aatcatttcg aaccctga 38

<210> 1877
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1877
atcccttcca attccctata attactactc taatgtga 38

<210> 1878
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1878
atcccacttg ccatcttcgg tcctatcaaa tggtaga 37

<210> 1879
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1879
atcttaatgc cttgatccga ctcctacctc ttga 34

<210> 1880
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1880
atgccccccg cgcttttaac tcacacctgc tctgctga 38

<210> 1881
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1881
atgcagtcgt taacgtccat cgcgtgtgtc ttccg 35

<210> 1882
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1882
atccctttta ttgttaaact tcatagaaat catactga 38

<210> 1883
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1883
atcttcccgc cacttgtgcc ttatcattac cccccga 37

<210> 1884
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1884
atcccaattc acatacactc tcaccctcac taaaca 36

<210> 1885
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1885
atctacaccc tatcgaaact tcccatgaaa ctacaa 36

<210> 1886
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1886
attggtgccc gttgtgagga tcaagcaaca tgcta 35

<210> 1887
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1887
caaacgtacc ttactgcata caccttccct gttattga 38

<210> 1888
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1888
attgaacgag cgagaaatct aagcgccggc atctg 35

<210> 1889
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1889
atttcctaac attgctctct taactgttcc tatcc 35

<210> 1890
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1890
cacctggcca ttacactttc cttttgagtt ctctcga 37

<210> 1891
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1891
attttagcat acggctcctt ggtttcatca gctgttga 38

<210> 1892
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1892
attcgaattc tccacctgat aaccgccaca ttaaa 35

<210> 1893
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1893
atgtttgtcg gcctttgtgg gtcctttcgc ttccca 36

<210> 1894
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1894
ccgcaaccca actcgtattc taatctacta cacattga 38

<210> 1895
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1895
cccttaggtt ccagtgaagt tatgcaactt catgta 36

<210> 1896
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1896
ccccaaggac gcacgggggt acccgcatac tgtta 35

<210> 1897
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1897
ccagccggtc caggtggctg acccgcgaag gagcta 36

<210> 1898
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1898
cccgtcatga accgggaccc aatcttacct tttatga 37

<210> 1899
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1899
ccgcgcgtta gctacggatg gcattaaccc agttgga 37

<210> 1900
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1900
catggctctc atccgaagac gcgccttctt ctatttga 38

<210> 1901
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1901
ccgctggcgc tatttttttt ctatgctgtt ggagctga 38

<210> 1902
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1902
catggattta gtcgtcccta gccgaacttc tgttaa 36

<210> 1903
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1903
ccgcttaacg gggcacccaa ctaattcgat atctctga 38

<210> 1904
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1904
cggtccgtcg acccatgttt tttggctcta ttgcta 36

<210> 1905
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1905
cgctaaccgg aactaggttg caaagctatg gtggta 36

<210> 1906
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1906
cgctattatt actctccccg ggtaatatcc tatattga 38

<210> 1907
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1907
ccttgaaagg tgtaacgcac gttggatcgt cgtgaa 36

<210> 1908
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1908
cgacgaaacc ggacaccata cccttacgac ttttaa 36

<210> 1909
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1909
ccttaccccg agcgtgcccc cttcttaagt attcg 35

<210> 1910
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1910
cgttttatag tttcgatcta aggttcatta tatata 36

<210> 1911
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1911
cgcttaattc ttggttatat tatctaacta cgcaatga 38

<210> 1912
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1912
ctctaccgag tcccttacat atgtacccac ttccaga 37

<210> 1913
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1913
gaacccgagg gagggattct caatgtgcaa tgtatga 37

<210> 1914
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1914
cttcggggca ctaccttcgt ttttttgttc tactatga 38

<210> 1915
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1915
gaagccccat aaaagacgcc gaaatattgc atctgga 37

<210> 1916
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1916
gaaactaaac cgactgggtt aaattttatc atgcgga 37

<210> 1917
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1917
gagcttaaac ctctccgagc gcttatggct cgattga 37

<210> 1918
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1918
gaggaaacgc ggcagcctta taggccgatc tttca 35

<210> 1919
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1919
gaggtttaat aaataggcca tgcttattgg tccgctga 38

<210> 1920
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1920
gatcacgtcg cggattcccc cgtttcgcga gaggaga 37

<210> 1921
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1921
gagttccttt atttgcattt ttatcagatc aagttta 37

<210> 1922
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1922
gagtcctcta cttcatataa tctccagagg ggggga 36

<210> 1923
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1923
gagtttcgat attcgtctta aatccacccc tgtaa 35

<210> 1924
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1924
gagtacgatt tgaccactat cttcgtagat tccata 36

<210> 1925
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1925
gataacggga accaaatcgc gctgtagagg cctggga 37

<210> 1926
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1926
gatctacccg accctattaa ccttagtact tgacaa 36

<210> 1927
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1927
gatccctaat tgttttccta ttaatcgttt tcccatga 38

<210> 1928
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1928
gatcgctcgc caggcacact gccgtttcga gtatta 36

<210> 1929
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1929
gcacgacgaa cgagtccggg aggaccggct tcccca 36

<210> 1930
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1930
gatcttatcc ggcccacatc tttaacccca acact 35

<210> 1931
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1931
gccgtctcgg ggtaatcggc ctaatacgta gacat 35

<210> 1932
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1932
gccgcggggg gccggggagg tcccatcgag gttatga 37

<210> 1933
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1933
gctaaccttt gttatactag gactttctcc acaggtga 38

<210> 1934
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1934
ggggtttctt ggagcgcaat acggtttgtt tctga 35

<210> 1935
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1935
ggcgaaacta gcctggattg gagttagacc tttca 35

<210> 1936
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1936
gggggagttg catggttgca actctggagg ccctttga 38

<210> 1937
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1937
gggtttaacg tgcccacccg tcagggtagt cctacga 37

<210> 1938
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1938
gggtatgccg ttctgtccgg gccttatgcg catat 35

<210> 1939
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1939
gggacaatag aattaaactc gggcttaggt cctcgga 37

<210> 1940
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1940
ggggagatgg ctgggtcttg ataccggaaa cttgcga 37

<210> 1941
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1941
ggcttgtata cctattatgt taggcaggtg ccctttga 38

<210> 1942
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1942
gtggcccggt tgtgttgtca cggacgtgcg gtccca 36

<210> 1943
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1943
gtgttgtatg tcttcgcaat cctagccgac cgccaa 36

<210> 1944
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1944
gtgcgcgttg gtgggccgct tgtaactata ccctttga 38

<210> 1945
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1945
gttttggtgg ttcaaaccgg tcgcccactg gcatt 35

<210> 1946
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1946
tacatcccgg ctagatccgt gaggactggc ggtgtc 36

<210> 1947
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1947
taaaagaact cttttcgtca accatacctc aggcctga 38

<210> 1948
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1948
tagtccatat atagcttaaa ctgatgcggg ggttcga 37

<210> 1949
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1949
tacccttatc ggcaccccca cacttatggt aattc 35

<210> 1950
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1950
tacgggaatg ggtttggccc taaaagccag acacga 36

<210> 1951
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1951
tagctaacgg tcccaacaag catgttggga ttggca 36

<210> 1952
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1952
tacgccatga tacttacatc tagccacctt ttttttga 38

<210> 1953
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1953
tatatctggc caccggcggg ggggctggtg atctta 36

<210> 1954
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1954
tacctgagcg ctggagctcg ttagtcggaa taggaa 36

<210> 1955
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1955
tatcaaattc ggtcgtagta tttgtcgccg taacgtta 38

<210> 1956
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1956
tactagcgat catttgttcg ctttgggttg gttattga 38

<210> 1957
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1957
taggtggtcg tgcctacatt gactgtttcc atcca 35

<210> 1958
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1958
tactccggcc caactatata cttctccaca cctcc 35

<210> 1959
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1959
tatcacgact tcccaccttc ttccgccttt gtcccaa 37

<210> 1960
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1960
tcacagggcc tgaaggtagt agatattaaa cgctgga 37

<210> 1961
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1961
tcattatacg gatcccgcgc cgaacctggc cgact 35

<210> 1962
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1962
tcactttggt aaactattcc cccgtttctc tcctatga 38

<210> 1963
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1963
tcaatacgcg tccggaaatc cccctgtgca agctgga 37

<210> 1964
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1964
tcaatgcttg ctgctcgtag ctttcaaccg tggtgga 37

<210> 1965
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1965
tatcgtttgc catattccgc cagagaaaag tctcc 35

<210> 1966
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1966
tcctcgtggc ttaccggtat actcccccaa ctcattga 38

<210> 1967
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1967
tccctccccg ttgcccttat tccgtactca cttatga 37

<210> 1968
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1968
tcctgtctcc gctaggtaac atccgaatac tctcttga 38

<210> 1969
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1969
tcctcgacca tctacaaact ttgaagccac acccca 36

<210> 1970
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1970
tccgggcacc atatctccac tgtgcgaatc actct 35

<210> 1971
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1971
tccactcaat ccccggttcg tatgctctgt accattga 38

<210> 1972
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1972
tcccgtccca cctttacttt gctacttcac catcttga 38

<210> 1973
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1973
tccttgcata cctcggctct cctttctctc acatttga 38

<210> 1974
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1974
tctcgtgcgc gatttctgtt ctagggtagg agaggga 37

<210> 1975
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1975
tcggctcatc caaaattact tgcgggtttc ataaatga 38

<210> 1976
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1976
tctcgcacct ttattttctc cgctcttttg tttcca 36

<210> 1977
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1977
tcgctacaat tggccttttc tgtatttggg ttctatga 38

<210> 1978
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1978
tgcacgccga gcaaggcggt actctaggac ccacgga 37

<210> 1979
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1979
tgccgctttt accacttgac ttacgataag aaaactga 38

<210> 1980
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1980
tgagtttttc taaataccgc agcctcagcc acgcgga 37

<210> 1981
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1981
tgccttttt ttcaccggtt tgattccgct ttcg 34

<210> 1982
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1982
tgaaagccag gtattttggg gctccttttt cctcc 35

<210> 1983
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1983
tgacctacaa agggtttgaa ataaatcgct cgacgtga 38

<210> 1984
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1984
tggaaagtta gatttcctgg ggtctcgctt tgagt 35

<210> 1985
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1985
tgtacttact tttactcgat caccttataa tttcc 35

<210> 1986
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1986
tggcctattc tcgtacaatt tacttcttat catcctga 38

<210> 1987
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1987
ttagtacgtg ccgagccccg agccctagag acctgtga 38

<210> 1988
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1988
tgttctgtct tatgccatga agttcttata agtaga 36

<210> 1989
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1989
tgttagttcc actttccgat aatccgtctt cctta 35

<210> 1990
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1990
tgttaggcct ttatttacct tatcttaatt gtttta 36

<210> 1991
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1991
tgtttttaca cgcctacttt ttatctatct ccacatga 38

<210> 1992
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1992
ttccccttttt atttgcccgt aataatcgct tgacatga 38

<210> 1993
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1993
ttgaacctcg gtagagccgt aaggcctatc cggtttga 38

<210> 1994
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1994
ttcgaccctt ttccatactt cttctcacgc aaattga 37

<210> 1995
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1995
ttcttaatgt aaacttttgt tgatactgtg ccacc 35

<210> 1996
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1996
ttcccctcta ctcgcctatt ttcctcataa aattta 36

<210> 1997
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1997
ttcctccccc ccgtgcaatt ttagaggtcc cgtgg 35

<210> 1998
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1998
ttctcaccgt acatccctta cctatcccct tttat 35

<210> 1999
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 1999
tttccgatta ttatgaccgc ttttattatt ggaggtga 38

<210> 2000
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2000
tttgcgaatc tagctttcct gtatacatga caataa 36

<210> 2001
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2001
ttggtctatt atgctctata tgtgaagtgc gaaaa 35

<210> 2002
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2002
ttttagttcc actttccgat aatccgcctt cctta 35

<210> 2003
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2003
ttttagtata cccaccgcga tactctcagt cttgc 35

<210> 2004
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2004
ttttggcgtg agctccaggt agtggttgaa acccatga 38

<210> 2005
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2005
agttcgccgt tggaggctct ttcgacatca tgctaga 37

<210> 2006
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2006
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 2007
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2007
aaagcgatca cgtttctttg ggctgctgtc tagca 35

<210> 2008
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2008
ccattaacgg taagacttaa ttccgtttgg cgtgta 36

<210> 2009
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2009
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 2010
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2010
aatcgtactc acggtgacat ccttggtttg atggta 36

<210> 2011
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2011
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 2012
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2012
gagttttact gccgtgtcaa ttatcggtcc tgtgttga 38

<210> 2013
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2013
cgggacgaat cttagtttat tccattcgca agtgctga 38

<210> 2014
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2014
gatcataccg cgatatagac attgcaccgt ttcta 35

<210> 2015
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2015
aggataactg gtttttatct ctgctaattc actcg 35

<210> 2016
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2016
agggacgaat cttagtttat tccattcgca agtgctga 38

<210> 2017
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2017
atcgcttaag ttccgtaagg gttaaggttc gttattga 38

<210> 2018
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2018
gagtacgcgc aatgacaatc gacttgtttt cggtc 35

<210> 2019
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2019
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 2020
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2020
cctcagttta cgtactcatc ttgaatgggg gaaata 36

<210> 2021
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2021
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 2022
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2022
agatcttttc ggttattggg gggaaccgag acgta 35

<210> 2023
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2023
agattaggtt gctctgtccc gtgcttaagt atcgctga 38

<210> 2024
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2024
acattttccg catgggacgg aatgttcttc ttcattga 38

<210> 2025
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2025
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 2026
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2026
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 2027
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2027
tacctcatgt ttgtcactat ggaaatcatc ccgac 35

<210> 2028
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2028
tacctcatgt ttgtcactat ggaaatcatc ccgcc 35

<210> 2029
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2029
tgcttctgca tgcaacgtgc tctggactag ctgtc 35

<210> 2030
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2030
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 2031
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2031
ttttcgcatt aaaaacgggt gtccaccggt tggtgga 37

<210> 2032
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2032
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 2033
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2033
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 2034
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2034
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 2035
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2035
gttgccttgt accccatagc tggtaaggtc ctagta 36

<210> 2036
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2036
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 2037
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2037
ttggcctttg agttagtgcc tagatagcct gagtta 36

<210> 2038
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2038
tcattggttc cgatccgggg ctctggacct gacata 36

<210> 2039
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2039
tcgttatagc ctatatacac gcggttgact cgctga 36

<210> 2040
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2040
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 2041
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2041
gcggagcgct tatataatgg tgttgtaggt ccatta 36

<210> 2042
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2042
gtacgccgct gccaataatg tctgttgtta cattc 35

<210> 2043
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2043
tttaatgggg agctacgtct cgaatcgatt ttggta 36

<210> 2044
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2044
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 2045
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2045
ttcccttgtt cgagtttccc agcctggacg ccctg 35

<210> 2046
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2046
tccctgtcgg tccaaacgta agggcgggct aaaaca 36

<210> 2047
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2047
gtctggctgg aaaatcaggg ttgcctatgt ttcaaa 36

<210> 2048
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2048
tacattccgg gctgttcagc aatgtcgctc cagtta 36

<210> 2049
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2049
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 2050
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2050
tttgcaagtt cgctccgtat atccactgtg aagac 35

<210> 2051
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2051
tttgcaagtt cgctccgtat atccactgtg aagcc 35

<210> 2052
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2052
tcctctccgt ctgtgaagac gtgtgtggcg gaatta 36

<210> 2053
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2053
tgttttgctg cttagagcgt aatcgtattg gctct 35

<210> 2054
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2054
tttgcttcat ttgcacttcc cccgttattc cgctca 36

<210> 2055
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2055
aaccaaacct gtacccgaat aaatcatcca ttaacga 37

<210> 2056
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2056
aacccaattc acatactctc tcaccctcac taaaca 36

<210> 2057
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2057
aacaacctcg aatttgtata ggttcgcatc actctga 37

<210> 2058
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2058
aactgtcatc cgctgactaa ctcaatctaa ctcca 35

<210> 2059
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2059
acattaagga tttatcttca cccatttttg cctcttga 38

<210> 2060
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2060
acaaagtaac gtacctcagt tccggtactt acgggtga 38

<210> 2061
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2061
accccacctg cacaccttca tctcatctat tacat 35

<210> 2062
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2062
acattttccg catgggacgg aatgttcttc ttcattga 38

<210> 2063
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2063
acaattaata taaattcctt acatcaatcc tagacga 37

<210> 2064
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2064
acattcccgt ggtggtcctt acgggtgggc cagcctga 38

<210> 2065
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2065
aatttttact gcgtggaaag gattcgggaa tttgttga 38

<210> 2066
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2066
accggactgc acagggatcc cgataggagg gtgac 35

<210> 2067
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2067
accttgcaa tcaatccccc cttttacctg tacatga 37

<210> 2068
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2068
actcgccctt cacgacttta tccaaattta ctttca 36

<210> 2069
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2069
acgtctgcca aatctactcg cattcgatac acaca 35

<210> 2070
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2070
actactgggc gcactagcgc gaggctccac tgaat 35

<210> 2071
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2071
actcacattc ccacccccta ttgcctttaa atacctga 38

<210> 2072
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2072
actgcagcat caagccacta attggggctt gtatcga 37

<210> 2073
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2073
acggttggag cttgtgggaa actaggttgg gttcgga 37

<210> 2074
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2074
agatcttttc ggttattggg gggaaccgag acgta 35

<210> 2075
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2075
agcacccacc cgcttctatt cctcgatcat acact 35

<210> 2076
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2076
aggataactg gtttttatct ctgctaattc actcg 35

<210> 2077
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2077
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 2078
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2078
agattgctcc gcagtacggt gtagggcgat aaaaa 35

<210> 2079
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2079
agccgcccct ttagacagag gttacgactt gcgta 35

<210> 2080
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2080
actttggggc tttctcggtc ttcgcactat ggtactga 38

<210> 2081
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2081
agccttacgc catgccgaga gagcgctttg tttcga 36

<210> 2082
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2082
agaaccccat tctcattcca ccagttacta cgctta 36

<210> 2083
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2083
aggtatacaa atgtgtagtc tagtcctgcc gatcga 36

<210> 2084
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2084
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 2085
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2085
agggattcgc cctgtgtcac gctcaaactg tagtaa 36

<210> 2086
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2086
agtaatcccc ggcgtataat ttcagtcctg tggtttga 38

<210> 2087
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2087
agttgatgca gtgggacctg agattaggct tatga 35

<210> 2088
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2088
agtattcctc tggtagtatt tttccttaat tccgta 36

<210> 2089
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2089
agtcctgatg ctcaatgcgc gagtttcatt gagct 35

<210> 2090
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2090
agtacctctt gtgggtcttg gggctgccac tgcct 35

<210> 2091
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2091
atcaccgtcc ttatcataac cagttcgttt tttata 36

<210> 2092
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2092
atgcttccat ctctgttttt taacccctta ccgtta 36

<210> 2093
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2093
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 2094
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2094
caccctaatc accgctaacc ttaccataac aatttga 37

<210> 2095
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2095
cagccgctct agttacaggt ttactgcggc taaacga 37

<210> 2096
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2096
cacagtatcc ttgtacagat tttgatggtt gttgg 35

<210> 2097
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2097
catcggggta cgcttccttt aaaagatcct gtgcc 35

<210> 2098
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2098
attcatggtg ccgtgagaaa ataaccatgt acgcaa 36

<210> 2099
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2099
atttccatac cattgtaggc cttctcgtat cgtaa 35

<210> 2100
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2100
cccatagggc acttttttaca cccctcgcat actattga 38

<210> 2101
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2101
ccgtgatgca gatgaatgta cggacgtctg ccggtga 37

<210> 2102
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2102
cccaccgagg gcaggtggtg acgtttcccc agccttga 38

<210> 2103
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2103
cccagacaac taccgcggag tagcggaagc cgtactga 38

<210> 2104
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2104
ccgcgcatca cattgggctc tgggatgccc gctgca 36

<210> 2105
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2105
cctagaccac gtgtgtgtcc cgggtaggga agtct 35

<210> 2106
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2106
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 2107
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2107
ccattaacgg taagacttaa ttccgtttgg cgtgta 36

<210> 2108
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2108
ccccaccgcg cacaccccga actttacata tgcatga 37

<210> 2109
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2109
ccacacccat tttgactgaa cagcctatcc ggtactga 38

<210> 2110
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2110
cctgttcaat tcatcctcca ggttcgtgtt acccta 36

<210> 2111
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2111
cctttcggac tagcttctgg cccatttggc gccata 36

<210> 2112
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2112
ctacaattag aagctcggtt tcttctgcgg tgacg 35

<210> 2113
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2113
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 2114
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2114
cgcgggggcc tatatgctgt ttccaccgct agtta 35

<210> 2115
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2115
cgctggctgc acgcacgcct gggcgcactg cccta 35

<210> 2116
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2116
cgtataggtt aatacgtact ggagcccgga gggga 35

<210> 2117
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2117
cctgaatggg cggtccgcac ttcgttagta tcccaga 37

<210> 2118
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2118
ctcaacgtag tctgcggagt tactccagaa tgacg 35

<210> 2119
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2119
gaacttcgcg gtgactcgtg cccttcgttg tcctttga 38

<210> 2120
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2120
gaaccatcat tggcttcatt tagctattct acacc 35

<210> 2121
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2121
ctgtccttaa atggaaggtc attttggctc aggtctga 38

<210> 2122
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2122
gacacggggg caaaggtggt ttgtgtaccc cgtacga 37

<210> 2123
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2123
gaaggtcatc ggactgttct gtcaatcccc ctttgga 37

<210> 2124
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2124
gagctcaaat cggggactac tgcattgtct actcg 35

<210> 2125
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2125
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 2126
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2126
gacgccttga aagcgacgcc tgatactcgg gggcctga 38

<210> 2127
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2127
gagggcccca agtcaccgct gtagttctag gcccaga 37

<210> 2128
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2128
gacggtcgta ctggagcatc ctgagactgt tcgag 35

<210> 2129
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2129
gagcgactca tcccggggtt tggatcagcg ctccgga 37

<210> 2130
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2130
gagtacaggt ccggcgccct atcgagaaag gaaaa 35

<210> 2131
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2131
gagtactgcc atgaggccag aggtaagaag ccgaatga 38

<210> 2132
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2132
gagtacagtt ggattccgcc cttgtatctg tacga 35

<210> 2133
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2133
gagtacctac ccgtcgaata ttgagattcg ggaacga 37

<210> 2134
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2134
gagtctcata cccctgaata tgtttttatg ctttc 35

<210> 2135
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2135
gagtggctca accagggtcc agaattacaa aaaata 36

<210> 2136
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2136
gcaatgtccc ctgttttcta ctaactctga tcttga 36

<210> 2137
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2137
gcagttctaa aggtggaaga atccaaatga cccaga 36

<210> 2138
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2138
gcaccttctc ctttaattcc cagccaataa ccatc 35

<210> 2139
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2139
gatgtaacgg gtctgcgggg agtttgtcat gcggga 36

<210> 2140
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2140
gccatcatat tcattattcc atgcctcccc atccc 35

<210> 2141
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2141
gatcataccg cgatatagac attgcaccgt ttcta 35

<210> 2142
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2142
gcaatgaaca cctacacata cctgccttca cgatta 36

<210> 2143
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2143
gccctcactt caacgtttca taactctaca ttaata 36

<210> 2144
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2144
gcgccgatgg ggcatcggtc gcgacagtcc cgccgtga 38

<210> 2145
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2145
gcgttcccct tggacaatat cactttctta tcgtaa 36

<210> 2146
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2146
gcctcgacga cagtgaccta catcgctcgt ttcgt 35

<210> 2147
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2147
gcctctgcca agcaattcat cttaacgacc tctcta 36

<210> 2148
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2148
gccgccggag cacggctgtc tttatacgca ctgta 35

<210> 2149
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2149
gcggagcgct tatataatgg tgttgtaggt ccatta 36

<210> 2150
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2150
gcgtacatag atctaggctt tccatccagg atatgtga 38

<210> 2151
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2151
gcgccggcat actccttata ttccactatc cctgta 36

<210> 2152
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2152
gctaacgtac tctttttttc aacttcactg actattga 38

<210> 2153
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2153
gctcccgaat tgttttccta ttaatcgttt tcccatga 38

<210> 2154
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2154
gctgaacagg tcacggtccc ccccacacgt taacta 36

<210> 2155
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2155
gctatacgac tgcaagggtc tgaccgcagg gaattga 37

<210> 2156
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2156
gcttccgtat caagagtggc tcgcacccac agagta 36

<210> 2157
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2157
ggattacgta cccgactttt tttctggttt tcttt 35

<210> 2158
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2158
gggaccggac cctgtccttg gtcgtttgat ccacga 36

<210> 2159
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2159
ggtatttttt tgtgctaatc tcgccgaatc gtacctga 38

<210> 2160
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2160
ggctcaccga ctccttactt tgctggtact acctta 36

<210> 2161
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2161
gggaacgtaa ggttttggtg aacagagggt agtaaga 37

<210> 2162
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2162
ggtgcttgtc cctgtctcta cgtaactttt gaagttga 38

<210> 2163
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2163
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 2164
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2164
ggggtccttt tccatcacac atgtccagtt ccgta 35

<210> 2165
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2165
ggccttctta ccgactgggg cttcgggttt tacgt 35

<210> 2166
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2166
gtgggcccgc attgaaaccg cattcgtaga gcctgga 37

<210> 2167
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2167
gtacgccgct gccaataatg tctgttgtta cattc 35

<210> 2168
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2168
gtacttgccg ggttacgcag cgcgtccctc ttcgcga 37

<210> 2169
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2169
gtgagcttta cgtaatgcgg tagttcgtac taacgga 37

<210> 2170
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2170
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 2171
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2171
gtcgcctatc ccagtgcttc gaggggcgac gtgtctga 38

<210> 2172
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2172
gtgtacattt ccgcgcaagc tactcagttt tttttga 37

<210> 2173
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2173
gtttactcct gtaccgtgag gtgaagggaa agcaa 35

<210> 2174
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2174
tacattccgg gctgttcagc aatgtcgctc cagtta 36

<210> 2175
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2175
gttttttact atggttcatc taggattcca gtttaga 37

<210> 2176
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2176
gttctttcac ggacggcctc ctttccgatc tattca 36

<210> 2177
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2177
gttgccttgt accccatagc tggtaaggtc ctagta 36

<210> 2178
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2178
gttccgggct ctttgtgtag tggtgagcac ggcacga 37

<210> 2179
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2179
taaatgcccg gttcattgaa tcttcacaaa gatggga 37

<210> 2180
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2180
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 2181
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2181
tacaccatga tacttacatc tagcaacctt ttttttga 38

<210> 2182
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2182
taaccgttcg ctgggttcca gtcccttgtt aagcc 35

<210> 2183
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2183
tagtgttaaa cgtgagtgca cttaatcacc ctggta 36

<210> 2184
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2184
taccttgacc tgcatcggtt ctcttcataa tcctaa 36

<210> 2185
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2185
tagtcctgca ctgttttaca gttcttggta atatttga 38

<210> 2186
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2186
tacttcaaca cctcgctcgt tacataccaa tttct 35

<210> 2187
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2187
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 2188
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2188
tcatcgtcgc gtggccggta ggctgggaaa tcctg 35

<210> 2189
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2189
tcacggatgc gaatcaaatg aactacctaa gtccgga 37

<210> 2190
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2190
tcattggttc cgatccgggg ctctggacct gacata 36

<210> 2191
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2191
tcctagaaat ctgtattgtt tctcaaggcc acctg 35

<210> 2192
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2192
tccagccaca tggatgaaca gtgggccttt tcacgga 37

<210> 2193
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2193
tcgaggggaa ttgtccacat gcatggtttc ataagga 37

<210> 2194
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2194
tccttccgac gataacttac gtgtgacacg cgtactga 38

<210> 2195
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2195
tccgacccat atcgacgata ctggcgtggc gctactga 38

<210> 2196
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2196
tcccgagttt gtcgatctcg agtggggccc ttacctga 38

<210> 2197
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2197
tccctgtaac agggtctctc tctttatata ttctca 36

<210> 2198
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2198
tcgttcctcc tttcgggtcc gtgaggatcc ggaacga 37

<210> 2199
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2199
tcgttatagc ctatatacac gcggttgact cgctga 36

<210> 2200
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2200
tctcctttcc aacgggacgc tgtgttcccc tcagtga 37

<210> 2201
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2201
tctgcactac cgtgaggtat ggtgataggt ggcga 35

<210> 2202
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2202
tcgggccaga tcgcattcgt agttgctgtt tccgta 36

<210> 2203
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2203
tgcacttaat tccacttttc cacccttttc actttga 37

<210> 2204
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2204
tgcacataat tccaccttc cacccttttc actttga 37

<210> 2205
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2205
tgagcatctt cacacgcatg gctccacctc ataaatga 38

<210> 2206
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2206
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 2207
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2207
tggtctgcac acgcacgtcg cctcttgggg ttatgga 37

<210> 2208
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2208
tgtagttttg caatctatgg gctaggacca tgtcatga 38

<210> 2209
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2209
tgctgttctc ctggcgatag gagattagcc ggacgga 37

<210> 2210
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2210
tgctaagagg gctaaggcca cacacttaat aactta 36

<210> 2211
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2211
tgcttctgca tgcaacgtgc tctggactag ctgtc 35

<210> 2212
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2212
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 2213
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2213
tggagtacta gttccgagga cgtgagggaa gactcg 36

<210> 2214
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2214
tggtccctgg ttccacttca ttgagagcag agcat 35

<210> 2215
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2215
tgctcggatc ttcgattcgt gccgcctgac aaatca 36

<210> 2216
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2216
tgttttgctg cttagagcgt aatcgtattg gctct 35

<210> 2217
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2217
ttacgatgct caagggtccc gttcgttcac agatatga 38

<210> 2218
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2218
ttacaaaccc cggtgctcta aagttatcga tctcttga 38

<210> 2219
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2219
tgttgggggg ttagcacggg tttatgggtt gcttc 35

<210> 2220
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2220
ttgcagtgga gtaagccctg accgaggagc gacga 35

<210> 2221
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2221
ttctcgccta catctatctc aatattgtcc taccga 36

<210> 2222
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2222
ttgacatgcg atgtgcacag ttagttaatt tcaaca 36

<210> 2223
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2223
ttctatttac taaatacaac ctttgtagca gacaaga 37

<210> 2224
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2224
ttccacactt cactgtaccc actcttacca gccaca 36

<210> 2225
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2225
ttcctctagc ctccgactgg cgtgtgactc taaaga 36

<210> 2226
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2226
tttccgctat agtgcttctg gtacgacgcg tacgg 35

<210> 2227
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2227
tttatactca tacccgaact gttccacacc gcaactga 38

<210> 2228
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2228
tttattcgcc ctttgttgac ctacaccaca tcacttga 38

<210> 2229
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2229
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 2230
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2230
ttgtagtttg gccctttcga gggccgttgc ctggttga 38

<210> 2231
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2231
ttttgttcc ataggtacgg tctccggagg aatcgtga 38

<210> 2232
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2232
ttttgctaac gcgacggagg cgccgagagc tatgctga 38

<210> 2233
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2233
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 2234
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2234
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 2235
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2235
cgcgcgagca actacactaa cggaaaaccc gttcaa 36

<210> 2236
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2236
ctactcgccg caacccaaca ctcagtcaaa cgatca 36

<210> 2237
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2237
aacgcctaac tggacagaaa aatcgtacgc taccaga 37

<210> 2238
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2238
ataacgcacg tttccataac tcataccact ggccttga 38

<210> 2239
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2239
atgctatccc gtgaaacgag aacatacggt ttttttga 38

<210> 2240
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2240
tggactggtc gtggggtgtg aacgaatcca atgtg 35

<210> 2241
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2241
ttacccctgt aatgccctac atgatgttcg gttcatga 38

<210> 2242
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2242
tgcaccctac ccatttagat gtatcagcca cgggca 36

<210> 2243
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2243
aacccgtagt agcacaccct gtctaaccca aatcaa 36

<210> 2244
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2244
aaacgcgatc ttacatcatt acaatccctc aaccc 35

<210> 2245
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2245
aaatcccatc gagcccttaa cccagaccaa ctacg 35

<210> 2246
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2246
aagaacatga gactgaatat cgtacgcgag taaaatga 38

<210> 2247
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2247
aagagtatag gtccgaggga ggacctacta attac 35

<210> 2248
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2248
aatttcggtt acaccccgct ctctatcact atacgtga 38

<210> 2249
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2249
accactagag tcctcaagaa ctcatccacc cacctga 37

<210> 2250
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2250
acggtatttg ttttttcgat tgatggggaa gtata 35

<210> 2251
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2251
accggcgaat aacgccaaag cctgcgactg gccttga 37

<210> 2252
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2252
acgcttttaa aaccatacac tcgataacta taccc 35

<210> 2253
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2253
accctactat taacccatca aaagttcgct acccta 36

<210> 2254
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2254
acgcccatcg tttcgggcct ttgtccattt cataca 36

<210> 2255
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2255
accttccctt caacgggtgt attccggcgg gtagc 35

<210> 2256
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2256
acttcaccca cacaatacca tcatcgataa ctcaaa 36

<210> 2257
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2257
agcggctaat cactgaagta gccccaggtt ttgaa 35

<210> 2258
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2258
agcaggccca acggcctctg ggttagtatc cgcgtga 37

<210> 2259
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2259
agccgatcct tgtccccgtt gaaggccgga tcgaaa 36

<210> 2260
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2260
aggcatctca agcgctcgga ggtacgcact atttctga 38

<210> 2261
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2261
agagacggct tcaacgcggc tttcgggctc tcttc 35

<210> 2262
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2262
agtaccccct tatccctggg gcctcccggg gtacc 35

<210> 2263
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2263
atcaacacca ttccctatat ccaacactaa acgtaa 36

<210> 2264
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2264
agtttctcga caggggcact ctggccacgt ggctgga 37

<210> 2265
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2265
aggggatact cagaacccga tgtaaatctt aagaa 35

<210> 2266
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2266
atcttcccac cacttgtgcc ttatcattgc cccccga 37

<210> 2267
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2267
attcgaattc tccacctgat aaccgccaca ttaaa 35

<210> 2268
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2268
ccagctagac cagacgaatg tggaactgaa tctgc 35

<210> 2269
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2269
ccctaaaagc gacccactca gcggatttga cttgc 35

<210> 2270
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2270
ccaccccacc gagaggggct catggcatac gctga 35

<210> 2271
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2271
cccgggaaca cggtccgttc gtaggagaac gaccaa 36

<210> 2272
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2272
ccatgctctc cggaggccgt tttacggagg gagcgtga 38

<210> 2273
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2273
cgcgcgagca actacactaa cggaaaaccc gttcaa 36

<210> 2274
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2274
cgcttccgct acgtaccccc cgggcgttcg ccttgga 37

<210> 2275
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2275
ctcaaagcaa cttgctggtg ccagcaccga agtccga 37

<210> 2276
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2276
ccttacattt atgttgaggc ccccggtaat tattgtga 38

<210> 2277
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2277
ctggtaccct gcgattcggg tctcgacttg gttcc 35

<210> 2278
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2278
gaacgtacgc cttcctcgta ataattggaa ccttcta 37

<210> 2279
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2279
cttcgccatc ttccggtcgg cgaaaaacag gccac 35

<210> 2280
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2280
gaggaaacgc ggcagcctta taggccgatc tttca 35

<210> 2281
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2281
gaggacattg gaccgcccca tacgaacgcc actctga 37

<210> 2282
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2282
gagtacgact aattatctta acaacattta tgaaca 36

<210> 2283
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2283
gagtagttgc atcgaagcgg aggcttttcc ggatc 35

<210> 2284
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2284
gcactgtccc cggttttcta ctaactctga tcttga 36

<210> 2285
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2285
gcgtcctgca ctgttttaca gttcttggta atatttga 38

<210> 2286
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2286
ggcacacacc tagcctgaaa ggtcctttaa cattgga 37

<210> 2287
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2287
ggcccagtac tcacgctaga ccctcgaggg gactcga 37

<210> 2288
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2288
ggacccgtac cgacacctat agccttcccc cagag 35

<210> 2289
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2289
ggtctagcgg gtacaccttc tggggttcgg tttgtga 37

<210> 2290
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2290
gggacaatag aattaaactc gggcttaggt cctcgga 37

<210> 2291
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2291
gggaactcgc aaaatttctt gaccccaagg tacccga 37

<210> 2292
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2292
ggtccttgag catcaaatcg ctaaggggtc ccgcaga 37

<210> 2293
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2293
gtctggtttt tatattcaga gccctgtttg ggttaga 37

<210> 2294
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2294
taaggatagc gctcggcctt tattgatact tgttt 35

<210> 2295
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2295
taaccgcatc actttcactt gacactaata cctcatga 38

<210> 2296
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2296
taatatccct gctctcgaac acttcgcttt aaacta 36

<210> 2297
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2297
gttgggaccc gattaaccca acagaattcg gaataa 36

<210> 2298
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2298
gttccgctcc agggaacaga ggatcaggga gttcga 36

<210> 2299
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2299
tatatccgca tcccgaagca tagatctcaa gtactga 37

<210> 2300
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2300
tatacacccg aaaagcgggt tatcaaaaat tatcatga 38

<210> 2301
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2301
tactatcttc tagttcaaca gccacttatg accca 35

<210> 2302
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2302
tacgcccctt aacctctaac cagcgagggt agttc 35

<210> 2303
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2303
tcaacgaccg cgaggttgaa cgacgtgggg actgta 36

<210> 2304
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2304
tattatgttc acggccggag gggggccggg ggatgcga 38

<210> 2305
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2305
tcgcatacaa caaaattacg cctaaccata acacca 36

<210> 2306
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2306
tgaagtactc gtaagtttta tagtttttca ttctatga 38

<210> 2307
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2307
tgacaaaagt cccctgaggg gtggactcgt gaccca 36

<210> 2308
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2308
tgcgggtgca tgccgagtga gccgaacgac ccatc 35

<210> 2309
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2309
tggccggggt ttggccccag aaagatacgc cggggtga 38

<210> 2310
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2310
tggacctact acgcttgggc aacgcggggg ccccta 36

<210> 2311
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2311
tggacgattt tgccaccgtt agtacaaccc gggggtga 38

<210> 2312
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2312
tgtagcccag gagaaaccct accccaacgg acgcg 35

<210> 2313
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2313
tgtggcgagg gccgcctcca ggcatgaaat gtatca 36

<210> 2314
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2314
tgtttgcttc atagtaacgt tgagactgca taaactga 38

<210> 2315
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2315
tgttagttcc actttccgat aatccgcctc cctta 35

<210> 2316
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2316
ttctcgacgg ccttccaggc gggagacaaa gctgc 35

<210> 2317
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2317
ttcatggacg aaacgggcgg cggacattta ttcgctga 38

<210> 2318
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2318
tttgctccgg tgtttgaccc tcgcagtccg gagagtga 38

<210> 2319
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2319
aaagcgatca cgtttctttg ggctgctgtc tagca 35

<210> 2320
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2320
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 2321
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2321
aggataactg gtttttatct ctgctaattc actcg 35

<210> 2322
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2322
agggacgaat cttagtttat tccattcgca agtgctga 38

<210> 2323
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2323
aacgcgctgg tcttcgtgta ttgtttcgcg ctaagga 37

<210> 2324
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2324
aactatcatt catcttggtg tgatctacct ttgtt 35

<210> 2325
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2325
agttcgccgt tggaggctct ttcgacatca tgctaga 37

<210> 2326
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2326
cttctatata tctcattctt ccggtttatt caatca 36

<210> 2327
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2327
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 2328
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2328
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 2329
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2329
attccgtgag gtacgtaagg caaccagttt tattt 35

<210> 2330
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2330
acgcttcttt tacttagcta accattattc attgt 35

<210> 2331
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2331
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 2332
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2332
gagttttact gccgtgtcaa ttatcggtcc tgtgttga 38

<210> 2333
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2333
gatcataccg cgatatagac attgcaccgt ttcta 35

<210> 2334
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2334
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 2335
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2335
tacctcatgt ttgtcactat ggaaatcatc ccgcc 35

<210> 2336
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2336
tgttttgctg cttagagcgt aatcgtattg gctct 35

<210> 2337
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2337
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 2338
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2338
tcattggttc cgatccgggg ctctggacct gacata 36

<210> 2339
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2339
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 2340
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2340
ttcccttgtt cgagtttccc agcctggacg ccctg 35

<210> 2341
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2341
ttctgccggt ttttacggtc aaattctaga tagtttga 38

<210> 2342
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2342
ggtctcttgg tttagatccc gtcttttctg tggaaa 36

<210> 2343
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2343
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 2344
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2344
tcgttatagc ctatatacac gcggttgact cgctga 36

<210> 2345
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2345
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 2346
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2346
gtacgccgct gccaataatg tctgttgtta cattc 35

<210> 2347
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2347
tttaatgggg agctacgtct cgaatcgatt ttggta 36

<210> 2348
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2348
gtctggctgg aaaatcaggg ttgcctatgt ttcaaa 36

<210> 2349
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2349
tttgcttcat ttgcacttcc cccgttattc cgctca 36

<210> 2350
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2350
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 2351
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2351
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 2352
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2352
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 2353
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2353
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 2354
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2354
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 2355
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2355
tacattccgg gctgttcagc aatgtcgctc cagtta 36

<210> 2356
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2356
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 2357
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2357
aaatcccaac gagcccttaa cccagaccaa ctaca 35

<210> 2358
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2358
aaccaaacct gtacccgaat aaatcatcca ttaacga 37

<210> 2359
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2359
aacgtagtga ggtattccgc acccccttta ttcgtga 37

<210> 2360
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2360
aacccaattc acatacactc tcacccacac taaaca 36

<210> 2361
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2361
aaaaccgctc gcttaagctt tctttgactt tttgatga 38

<210> 2362
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2362
aatgcacaac cgcgctgaaa atcaccttta tcgaaa 36

<210> 2363
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2363
aattgatctt tttgtttacg aattcgtatt ccattga 37

<210> 2364
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2364
aatcaccaca aatactttcg cctatctcgc ttatctga 38

<210> 2365
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2365
aacttgtctt atgctctcca ccttacctat taacaa 36

<210> 2366
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2366
aactcctgat atcgtttctt cgtttccgcc aattcta 37

<210> 2367
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2367
accccacctg cacaccttca tctcatctat tacat 35

<210> 2368
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2368
acattttccg catgggacgg aatgttcttc ttcattga 38

<210> 2369
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2369
accgggaaca tttctgttgt ttcatttcat cactaa 36

<210> 2370
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2370
acgcttcttt tacttagcta accattattc attgt 35

<210> 2371
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2371
acccttgcaa tcaatccccc cttttacctg tacatga 37

<210> 2372
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2372
accgacgcac tctcctttct aaaatctatc tactt 35

<210> 2373
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2373
acgttaattt cccaaaaacg atcccacttt gaaca 35

<210> 2374
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2374
actcgccctt cacgacttta tcaaaatgta ctttca 36

<210> 2375
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2375
actcacattc ccacccccta ttgcctttaa atacctga 38

<210> 2376
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2376
agatcttttc ggttattggg gggaaccgag acgta 35

<210> 2377
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2377
agccttacgc catgccgaga gagcgctttg tttcga 36

<210> 2378
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2378
acttcgccca cacaatacca ttatcgataa ctcaaa 36

<210> 2379
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2379
aggataactg gtttttatct ctgctaattc actcg 35

<210> 2380
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2380
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 2381
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2381
agattaggtt gctctgtccc gtgcttaagt atcgctga 38

<210> 2382
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2382
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 2383
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2383
agtcctgatg ctcaatgcgc gagtttcatt gagct 35

<210> 2384
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2384
atatgcccgc tccctactca atatcttccc ataca 35

<210> 2385
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2385
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 2386
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2386
atgcattatc ccagctttaa acgctctacc caccgaa 37

<210> 2387
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2387
atgcttccat ctctgttttt taacccctta ccgtta 36

<210> 2388
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2388
atctgcttgt aagtccccgt gtatacctgt tcagt 35

<210> 2389
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2389
atcactggag tgatagtctg ggccctgaaa ctcct 35

<210> 2390
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2390
atcaccgtcc ttatcatacc cacttcgttt tttata 36

<210> 2391
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2391
attccgtgag gtacgtaagg caaccagttt tattt 35

<210> 2392
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2392
caccctaatc accgctaacc ttaccataac aatttga 37

<210> 2393
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2393
attcatggtg ccgtgagaaa ataaccatgt acgcaa 36

<210> 2394
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2394
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 2395
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2395
cccatagggc acttttttaca cccctcgcat actattga 38

<210> 2396
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2396
ccagcacact gacttttcct tacccacctc ctttc 35

<210> 2397
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2397
ccccgccatg gtttcaaatg gctgaagcta tctattga 38

<210> 2398
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2398
ccggtatctg cactcgtttg gaatcgacca acccctga 38

<210> 2399
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2399
ccgtctactg caacaggagg ccaccacggt cttccga 37

<210> 2400
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2400
cgctcttctt actttattca atatggtcat aaattga 37

<210> 2401
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2401
cgtttcttta tcccaaagtc attcgcatgc cctttga 37

<210> 2402
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2402
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 2403
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2403
cgcgcctcct tccccgatga tatgccactg ccgca 35

<210> 2404
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2404
cgcgggggcc tatatgctgt ttccaccgct agtta 35

<210> 2405
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2405
cgcttcttgg gcaggaggcg acacaggttc atattga 37

<210> 2406
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2406
gaaccatcat tggcttcatt tagctattct acacc 35

<210> 2407
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2407
ctgtcccttg taaggattat tcgtaggtat ttagtga 37

<210> 2408
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2408
ctgcggtcac ctgcatagta attaataata tacgtga 37

<210> 2409
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2409
cttcgctaaa tggctggggg aatcatactg ccggaga 37

<210> 2410
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2410
ctcctgcaga tcttggtacc tcttgttctt tcata 35

<210> 2411
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2411
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 2412
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2412
gaggactttt acttatcaac cccacctttc gaaca 35

<210> 2413
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2413
gagtacttga ctcttgtata ttttttctcc ccctt 35

<210> 2414
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2414
gatcccgttc ctttttaac gatctatttt cttttga 37

<210> 2415
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2415
gccatcatat tcattattcc atgcctcccc atccc 35

<210> 2416
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2416
gatcataccg cgatatagac attgcaccgt ttcta 35

<210> 2417
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2417
gcctcgacga cagtgaccta catcgctcgt ttcgt 35

<210> 2418
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2418
gccctcactt caacgtttca taactctaca ttaata 36

<210> 2419
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2419
gcctctgcca agcaattcat cttaacgacc tctcta 36

<210> 2420
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2420
gcgccggcat actccttata ttccactatc cctgta 36

<210> 2421
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2421
gcggaatagt ccacgggacg ctaagcgtgg agctcga 37

<210> 2422
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2422
gcgtccactc tatccttgct ttaatacatc gttaaa 36

<210> 2423
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2423
gcctttccag gttcttcttt tcgaccgttt atttttga 38

<210> 2424
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2424
ggccctggtg gcaccaagcc gcgagcccac tcaactga 38

<210> 2425
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2425
ggtgcccaac tggtgcttct ccttctttga gacgt 35

<210> 2426
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2426
gggagttttt atcctacaat gtaaattatc atacttga 38

<210> 2427
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2427
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 2428
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2428
ggtcccactc tgaccatcaa ttcacctaac ttgtaa 36

<210> 2429
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2429
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 2430
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2430
gtgggcccgc attgaaaccg cattcgtaga gcctgga 37

<210> 2431
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2431
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 2432
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2432
gtccaggctc tatgcttagt acccgttcat ttgcttga 38

<210> 2433
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2433
gtcgaaagcg gagcgactct ccgcacaacc ccaag 35

<210> 2434
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2434
gttgccttgt accccatagc tggtaaggtc ctagta 36

<210> 2435
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2435
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 2436
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2436
gttcagattt tgctttgtga cctggtgtgg ggtct 35

<210> 2437
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2437
gttccgggct ctttgtgtag tggtgagcac ggcacga 37

<210> 2438
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2438
tacattccgg gctgttcagc aatgtcgctc cagtta 36

<210> 2439
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2439
taattggccc gtttccaaat ttatagattc ctaca 35

<210> 2440
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2440
tactctagtc agaggatctc ttgcgatcaa tatgga 36

<210> 2441
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2441
tactgtcctt ttcaattctt gtcttggttt tatta 35

<210> 2442
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2442
tactcggccc tactacacga actcttttca aatca 35

<210> 2443
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2443
tacttcaaca cctcgctcgt tacataccaa tttct 35

<210> 2444
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2444
tattgtagcc tgtacggcgg ctgcacctcg tacgttga 38

<210> 2445
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2445
tcactttgcc cttttgtttc ttcattaatt ccttc 35

<210> 2446
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2446
tcacgctccc ggggttctat ctatttcgtt attcttga 38

<210> 2447
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2447
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 2448
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2448
tcattggttc cgatccgggg ctctggacct gacata 36

<210> 2449
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2449
tccgacccat atcgacgata ctggcgtggc gctactga 38

<210> 2450
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2450
tccatcatat tcattattcc atgcctaccc atccc 35

<210> 2451
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2451
tcgaggggaa ttgtccacat gcatggtttc ataagga 37

<210> 2452
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2452
tccccataat ctccgtcctc attgcttcca aaatttga 38

<210> 2453
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2453
tcccctcgcc actgacccat ctactcgcca aatcttga 38

<210> 2454
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2454
tcgggcaata ccctttccca tattatcttc tttcctta 38

<210> 2455
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2455
tctcgggatt ttccgcgact actcgagcgc tactca 36

<210> 2456
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2456
tcgcttcctt gtacttccta ttccctttc accaca 36

<210> 2457
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2457
tcgttatagc ctatatacac gcggttgact cgctga 36

<210> 2458
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2458
tcttcacgat atgcagtgaa aacaccaccc aactta 36

<210> 2459
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2459
tctgcttgat ggtgctttca gtttcttact tctacga 37

<210> 2460
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2460
tcttctgtaa ccattttgtt ccggtttttt tattt 35

<210> 2461
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2461
tgaagctccg aaggggtgca cttgccgaca gtagta 36

<210> 2462
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2462
tgacgtctcc tgttcggtta accctgtgtg cctgt 35

<210> 2463
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2463
tgcccttctt cgtttactct tatccacatt accttga 37

<210> 2464
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2464
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 2465
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2465
tgacatatcg ctcttacatc ccctcacaca atttca 36

<210> 2466
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2466
tgctccgagt accataaact aatttgaagc tctac 35

<210> 2467
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2467
tgctgttctc ctggcgatag gagattagcc ggacgga 37

<210> 2468
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2468
tggaccggtc aaaggatggc accggttcct ctcaca 36

<210> 2469
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2469
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 2470
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2470
tgctttgtac aacgttggtg tttggcattt gctgttga 38

<210> 2471
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2471
ttacaaaccc cggtgctcta aagttatcga tctcttga 38

<210> 2472
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2472
tgtgccactc gtgttcgaat ttctgttcct tatct 35

<210> 2473
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2473
tgttacgccc gtcctctata caccaattac gatatga 37

<210> 2474
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2474
tgttaggcct ttatttccat tatcttaatt gtttta 36

<210> 2475
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2475
tgttttgctg cttagagcgt aatcgtattg gctct 35

<210> 2476
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2476
ttctgccgga tacatttgat tcaataacta cttgta 36

<210> 2477
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2477
ttctgcttgc gagccgtcca gttagggccg ccacc 35

<210> 2478
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2478
ttctgccggt ttttacggtc aaattctaga tagtttga 38

<210> 2479
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2479
ttgcacgccc caatacacct aatctttatc cataca 36

<210> 2480
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2480
ttcgttccgg accacccatt tcctcaatca tatattga 38

<210> 2481
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2481
ttctgcgttt tacgatgctc gatgtctgtg acatgga 37

<210> 2482
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2482
ttgctccgac ctagtttact aaatttcaac cattga 36

<210> 2483
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2483
tttcattata ccccatcgac tttctctcta gttcca 36

<210> 2484
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2484
ttgtcccagg cagtttccac gagaacgcgt acggt 35

<210> 2485
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2485
tttggacgta tcactcccta accatttttt taccctga 38

<210> 2486
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2486
tttgcttcat ttgcacttcc cccgttattc cgctca 36

<210> 2487
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2487
tttcctgttt cttctttgtg tgtccatata atcga 35

<210> 2488
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2488
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 2489
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2489
tttaaacgta cgatattact tccgatccat acatta 36

<210> 2490
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2490
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 2491
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2491
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 2492
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2492
cgcgcgagca actacactaa cggaaaaccc gttcaa 36

<210> 2493
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2493
ctactcgccg caacccaaca ctcagtcaaa cgatca 36

<210> 2494
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2494
aacgcctaac tggacagaaa aatcgtacgc taccaga 37

<210> 2495
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2495
ataacgcacg tttccataac tcataccact ggccttga 38

<210> 2496
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2496
acaactatca aataaacctg cgccctctat gaatgga 37

<210> 2497
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2497
gaccaataga cccggccaag actggagatg aaagta 36

<210> 2498
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2498
atgctatccc gtgaaacgag aacatacggt ttttttga 38

<210> 2499
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2499
acgggagatc ctttaagtac gtcgaagcca gcaata 36

<210> 2500
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2500
acggtatttg ttttttcgat tgatggggaa gtaga 35

<210> 2501
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2501
ccacccgtac gaagccgcgt tttctgagtg cctattga 38

<210> 2502
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2502
ttacccctgt aatgccctac atgatgttcg gttcatga 38

<210> 2503
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2503
ggccaatttg aatgagggtt gtgaattatt taaga 35

<210> 2504
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2504
ggcctaacca tgggggtgg tgattattat gaaggga 37

<210> 2505
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2505
ttcatcgcgg ggccgaggta agcctaatct tccgta 36

<210> 2506
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2506
gtaactggta tcctaaaaga ataggttcgt accccga 37

<210> 2507
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2507
tgcaccctac ccatttagat gtatcagcca cgggca 36

<210> 2508
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2508
aacccgtagt agcacaccct gtctaaccca aatcaa 36

<210> 2509
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2509
aaacgcgatc ttacatcatt acaatccctc aaccc 35

<210> 2510
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2510
aaatcccatc gagcccttaa cccagaccaa ctacg 35

<210> 2511
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2511
aagaacatga gactgaatat cgtacgcgag taaaatga 38

<210> 2512
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2512
aagagtatag gtccgaggga ggacctacta attac 35

<210> 2513
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2513
aatttcggtt acaccccgct ctctatcact atacgtga 38

<210> 2514
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2514
accactagag tcctcaagaa ctcatccacc cacctga 37

<210> 2515
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2515
acggtatttg ttttttcgat tgatggggaa gtata 35

<210> 2516
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2516
accggcgaat aacgccaaag cctgcgactg gccttga 37

<210> 2517
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2517
acgcttttaa aaccatacac tcgataacta taccc 35

<210> 2518
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2518
accctactat taacccatca aaagttcgct acccta 36

<210> 2519
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2519
acgcccatcg tttcgggcct ttgtccattt cataca 36

<210> 2520
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2520
accttccctt caacgggtgt attccggcgg gtagc 35

<210> 2521
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2521
acttcaccca cacaatacca tcatcgataa ctcaaa 36

<210> 2522
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2522
agcggctaat cactgaagta gccccaggtt ttgaa 35

<210> 2523
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2523
agcaggccca acggcctctg ggttagtatc cgcgtga 37

<210> 2524
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2524
agccgatcct tgtccccgtt gaaggccgga tcgaaa 36

<210> 2525
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2525
aggcatctca agcgctcgga ggtacgcact atttctga 38

<210> 2526
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2526
agagacggct tcaacgcggc tttcgggctc tcttc 35

<210> 2527
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2527
agtaccccct tatccctggg gcctcccggg gtacc 35

<210> 2528
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2528
atcaacacca ttccctatat ccaacactaa acgtaa 36

<210> 2529
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2529
agtttctcga caggggcact ctggccacgt ggctgga 37

<210> 2530
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2530
aggggatact cagaacccga tgtaaatctt aagaa 35

<210> 2531
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2531
atcttcccac cacttgtgcc ttatcattgc cccccga 37

<210> 2532
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2532
attcgaattc tccacctgat aaccgccaca ttaaa 35

<210> 2533
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2533
ccagctagac cagacgaatg tggaactgaa tctgc 35

<210> 2534
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2534
ccctaaaagc gacccactca gcggatttga cttgc 35

<210> 2535
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2535
ccaccccacc gagaggggct catggcatac gctga 35

<210> 2536
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2536
cccgggaaca cggtccgttc gtaggagaac gaccaa 36

<210> 2537
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2537
ccatgctctc cggaggccgt tttacggagg gagcgtga 38

<210> 2538
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2538
cgcgcgagca actacactaa cggaaaaccc gttcaa 36

<210> 2539
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2539
cgcttccgct acgtaccccc cgggcgttcg ccttgga 37

<210> 2540
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2540
ctcaaagcaa cttgctggtg ccagcaccga agtccga 37

<210> 2541
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2541
ccttacattt atgttgaggc ccccggtaat tattgtga 38

<210> 2542
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2542
ctggtaccct gcgattcggg tctcgacttg gttcc 35

<210> 2543
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2543
gaacgtacgc cttcctcgta ataattggaa ccttcta 37

<210> 2544
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2544
cttcgccatc ttccggtcgg cgaaaaacag gccac 35

<210> 2545
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2545
gaggaaacgc ggcagcctta taggccgatc tttca 35

<210> 2546
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2546
gaggacattg gaccgcccca tacgaacgcc actctga 37

<210> 2547
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2547
gagtacgact aattatctta acaacattta tgaaca 36

<210> 2548
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2548
gagtagttgc atcgaagcgg aggcttttcc ggatc 35

<210> 2549
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2549
gcactgtccc cggttttcta ctaactctga tcttga 36

<210> 2550
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2550
gcgtcctgca ctgttttaca gttcttggta atatttga 38

<210> 2551
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2551
ggcacacacc tagcctgaaa ggtcctttaa cattgga 37

<210> 2552
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2552
ggcccagtac tcacgctaga ccctcgaggg gactcga 37

<210> 2553
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2553
ggacccgtac cgacacctat agccttcccc cagag 35

<210> 2554
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2554
ggtctagcgg gtacaccttc tggggttcgg tttgtga 37

<210> 2555
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2555
gggacaatag aattaaactc gggcttaggt cctcgga 37

<210> 2556
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2556
gggaactcgc aaaatttctt gaccccaagg tacccga 37

<210> 2557
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2557
ggtccttgag catcaaatcg ctaaggggt ccgcaga 37

<210> 2558
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2558
gtctggtttt tatattcaga gccctgtttg ggttaga 37

<210> 2559
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2559
taaggatagc gctcggcctt tattgatact tgttt 35

<210> 2560
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2560
taaccgcatc actttcactt gacactaata cctcatga 38

<210> 2561
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2561
taatatccct gctctcgaac acttcgcttt aaacta 36

<210> 2562
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2562
gttgggaccc gattaaccca acagaattcg gaataa 36

<210> 2563
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2563
gttccgctcc agggaacaga ggatcaggga gttcga 36

<210> 2564
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2564
tatatccgca tcccgaagca tagatctcaa gtactga 37

<210> 2565
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2565
tatacacccg aaaagcgggt tatcaaaaat tatcatga 38

<210> 2566
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2566
tactatcttc tagttcaaca gccacttatg accca 35

<210> 2567
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2567
tacgcccctt aacctctaac cagcgagggt agttc 35

<210> 2568
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2568
tcaacgaccg cgaggttgaa cgacgtgggg actgta 36

<210> 2569
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2569
tattatgttc acggccggag gggggccggg ggatgcga 38

<210> 2570
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2570
tcgcatacaa caaaattacg cctaaccata acacca 36

<210> 2571
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2571
tgaagtactc gtaagtttta tagtttttca ttctatga 38

<210> 2572
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2572
tgacaaaagt cccctgaggg gtggactcgt gaccca 36

<210> 2573
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2573
tgcgggtgca tgccgagtga gccgaacgac ccatc 35

<210> 2574
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2574
tggccggggt ttggccccag aaagatacgc cggggtga 38

<210> 2575
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2575
tggacctact acgcttgggc aacgcggggg ccccta 36

<210> 2576
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2576
tggacgattt tgccaccgtt agtacaaccc gggggtga 38

<210> 2577
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2577
tgtagcccag gagaaaccct accccaacgg acgcg 35

<210> 2578
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2578
tgtggcgagg gccgcctcca ggcatgaaat gtatca 36

<210> 2579
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2579
tgtttgcttc atagtaacgt tgagactgca taaactga 38

<210> 2580
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2580
tgttagttcc actttccgat aatccgcctc cctta 35

<210> 2581
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2581
ttctcgacgg ccttccaggc gggagacaaa gctgc 35

<210> 2582
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2582
ttcatggacg aaacgggcgg cggacattta ttcgctga 38

<210> 2583
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2583
tttgctccgg tgtttgaccc tcgcagtccg gagagtga 38

<210> 2584
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2584
atctatcgcc tatcgctaca acccgtattc gcttctta 38

<210> 2585
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2585
acaggacgtg cgcgctaacg ttctattagt tgtgctga 38

<210> 2586
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2586
attcgcatag cctatgggaa gggaattgtg aggggtga 38

<210> 2587
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2587
attctcttta ttttatcgct ttcatccagt gcatatga 38

<210> 2588
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2588
gaggttccct gtgggtcatg tcgcacgatt tttagtga 38

<210> 2589
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2589
ggcttcttat ttgcttgaat atccgcttcc tctct 35

<210> 2590
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2590
tccgggcacc atatctccac tgtgcgaatc actat 35

<210> 2591
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2591
tccgggcacc atatctccac tgtgcgaatc actct 35

<210> 2592
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2592
tttccgatta ttatgaccgc ttttattatt ggaggtga 38

<210> 2593
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2593
tgcccgcttt tggtttatat ggcttcttat ctttcga 37

<210> 2594
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2594
aaatcccaac gagcccttaa cccagaccaa ctaca 35

<210> 2595
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2595
aaccaaacct gtacccgaat aaatcatcca ttaacga 37

<210> 2596
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2596
aacgtagtga ggtattccgc acccccttta ttcgtga 37

<210> 2597
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2597
aacccaattc acatacactc tcacccacac taaaca 36

<210> 2598
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2598
aaaaccgctc gcttaagctt tctttgactt tttgatga 38

<210> 2599
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2599
aatgcacaac cgcgctgaaa atcaccttta tcgaaa 36

<210> 2600
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2600
aattgatctt tttgtttacg aattcgtatt ccattga 37

<210> 2601
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2601
aatcaccaca aatactttcg cctatctcgc ttatctga 38

<210> 2602
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2602
aacttgtctt atgctctcca ccttacctat taacaa 36

<210> 2603
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2603
aactcctgat atcgtttctt cgtttccgcc aattcta 37

<210> 2604
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2604
accccacctg cacaccttca tctcatctat tacat 35

<210> 2605
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2605
acattttccg catgggacgg aatgttcttc ttcattga 38

<210> 2606
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2606
accgggaaca tttctgttgt ttcatttcat cactaa 36

<210> 2607
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2607
acgcttcttt tacttagcta accattattc attgt 35

<210> 2608
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2608
acccttgcaa tcaatccccc cttttacctg tacatga 37

<210> 2609
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2609
accgacgcac tctcctttct aaaatctatc tactt 35

<210> 2610
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2610
acgttaattt cccaaaaacg atcccacttt gaaca 35

<210> 2611
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2611
actcgccctt cacgacttta tcaaaatgta ctttca 36

<210> 2612
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2612
actcacattc ccacccccta ttgcctttaa atacctga 38

<210> 2613
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2613
agatcttttc ggttattggg gggaaccgag acgta 35

<210> 2614
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2614
agccttacgc catgccgaga gagcgctttg tttcga 36

<210> 2615
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2615
acttcgccca cacaatacca ttatcgataa ctcaaa 36

<210> 2616
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2616
aggataactg gtttttatct ctgctaattc actcg 35

<210> 2617
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2617
agctcatgga cgcactggtg gcagtgcgac tactaa 36

<210> 2618
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2618
agattaggtt gctctgtccc gtgcttaagt atcgctga 38

<210> 2619
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2619
ataatgccag atctttggtt tatcttgcgc ttggta 36

<210> 2620
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2620
agtcctgatg ctcaatgcgc gagtttcatt gagct 35

<210> 2621
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2621
atatgcccgc tccctactca atatcttccc ataca 35

<210> 2622
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2622
atcgtagtca tagacccggc tttggtttcc gggta 35

<210> 2623
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2623
atgcattatc ccagctttaa acgctctacc caccgaa 37

<210> 2624
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2624
atgcttccat ctctgttttt taacccctta ccgtta 36

<210> 2625
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2625
atctgcttgt aagtccccgt gtatacctgt tcagt 35

<210> 2626
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2626
atcactggag tgatagtctg ggccctgaaa ctcct 35

<210> 2627
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2627
atcaccgtcc ttatcatacc cacttcgttt tttata 36

<210> 2628
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2628
attccgtgag gtacgtaagg caaccagttt tattt 35

<210> 2629
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2629
caccctaatc accgctaacc ttaccataac aatttga 37

<210> 2630
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2630
attcatggtg ccgtgagaaa ataaccatgt acgcaa 36

<210> 2631
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2631
ccattagctg agtttcggtg gacgcatgat agtttga 37

<210> 2632
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2632
cccatagggc acttttttaca cccctcgcat actattga 38

<210> 2633
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2633
ccagcacact gacttttcct tacccacctc ctttc 35

<210> 2634
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2634
ccccgccatg gtttcaaatg gctgaagcta tctattga 38

<210> 2635
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2635
ccggtatctg cactcgtttg gaatcgacca acccctga 38

<210> 2636
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2636
ccgtctactg caacaggagg ccaccacggt cttccga 37

<210> 2637
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2637
cgctcttctt actttattca atatggtcat aaattga 37

<210> 2638
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2638
cgtttcttta tcccaaagtc attcgcatgc cctttga 37

<210> 2639
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2639
ccttgagccg ccgggtttcg gtccttattc gtgaatga 38

<210> 2640
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2640
cgcgcctcct tccccgatga tatgccactg ccgca 35

<210> 2641
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2641
cgcgggggcc tatatgctgt ttccaccgct agtta 35

<210> 2642
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2642
cgcttcttgg gcaggaggcg acacaggttc atattga 37

<210> 2643
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2643
gaaccatcat tggcttcatt tagctattct acacc 35

<210> 2644
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2644
ctgtcccttg taaggattat tcgtaggtat ttagtga 37

<210> 2645
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2645
ctgcggtcac ctgcatagta attaataata tacgtga 37

<210> 2646
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2646
cttcgctaaa tggctggggg aatcatactg ccggaga 37

<210> 2647
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2647
ctcctgcaga tcttggtacc tcttgttctt tcata 35

<210> 2648
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2648
gacctgggct gtctttactt tgtttcgcgg ggctgga 37

<210> 2649
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2649
gaggactttt acttatcaac cccacctttc gaaca 35

<210> 2650
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2650
gagtacttga ctcttgtata ttttttctcc ccctt 35

<210> 2651
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2651
gatcccgttc ctttttaac gatctatttt cttttga 37

<210> 2652
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2652
gccatcatat tcattattcc atgcctcccc atccc 35

<210> 2653
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2653
gatcataccg cgatatagac attgcaccgt ttcta 35

<210> 2654
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2654
gcctcgacga cagtgaccta catcgctcgt ttcgt 35

<210> 2655
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2655
gccctcactt caacgtttca taactctaca ttaata 36

<210> 2656
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2656
gcctctgcca agcaattcat cttaacgacc tctcta 36

<210> 2657
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2657
gcgccggcat actccttata ttccactatc cctgta 36

<210> 2658
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2658
gcggaatagt ccacgggacg ctaagcgtgg agctcga 37

<210> 2659
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2659
gcgtccactc tatccttgct ttaatacatc gttaaa 36

<210> 2660
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2660
gcctttccag gttcttcttt tcgaccgttt atttttga 38

<210> 2661
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2661
ggccctggtg gcaccaagcc gcgagcccac tcaactga 38

<210> 2662
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2662
ggtgcccaac tggtgcttct ccttctttga gacgt 35

<210> 2663
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2663
gggagttttt atcctacaat gtaaattatc atacttga 38

<210> 2664
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2664
ggtctggggg gattcctacg gttccaacag cataatga 38

<210> 2665
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2665
ggtcccactc tgaccatcaa ttcacctaac ttgtaa 36

<210> 2666
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2666
ggctcttggt ttagccctta ttgtttatca ttcat 35

<210> 2667
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2667
gtgggcccgc attgaaaccg cattcgtaga gcctgga 37

<210> 2668
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2668
gtgcgttggt taccgatgga ggggacccgt agacta 36

<210> 2669
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2669
gtccaggctc tatgcttagt acccgttcat ttgcttga 38

<210> 2670
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2670
gtcgaaagcg gagcgactct ccgcacaacc ccaag 35

<210> 2671
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2671
gttgccttgt accccatagc tggtaaggtc ctagta 36

<210> 2672
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2672
gttgttcggg cgtcgcaaca cggtcccatt gtctta 36

<210> 2673
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2673
gttcagattt tgctttgtga cctggtgtgg ggtct 35

<210> 2674
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2674
gttccgggct ctttgtgtag tggtgagcac ggcacga 37

<210> 2675
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2675
tacattccgg gctgttcagc aatgtcgctc cagtta 36

<210> 2676
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2676
taattggccc gtttccaaat ttatagattc ctaca 35

<210> 2677
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2677
tactctagtc agaggatctc ttgcgatcaa tatgga 36

<210> 2678
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2678
tactgtcctt ttcaattctt gtcttggttt tatta 35

<210> 2679
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2679
tactcggccc tactacacga actcttttca aatca 35

<210> 2680
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2680
tacttcaaca cctcgctcgt tacataccaa tttct 35

<210> 2681
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2681
tattgtagcc tgtacggcgg ctgcacctcg tacgttga 38

<210> 2682
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2682
tcactttgcc cttttgtttc ttcattaatt ccttc 35

<210> 2683
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2683
tcacgctccc ggggttctat ctatttcgtt attcttga 38

<210> 2684
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2684
tcacccgacg gcaacaaggg catttgcttc atagta 36

<210> 2685
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2685
tcattggttc cgatccgggg ctctggacct gacata 36

<210> 2686
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2686
tccgacccat atcgacgata ctggcgtggc gctactga 38

<210> 2687
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2687
tccatcatat tcattattcc atgcctaccc atccc 35

<210> 2688
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2688
tcgaggggaa ttgtccacat gcatggtttc ataagga 37

<210> 2689
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2689
tccccataat ctccgtcctc attgcttcca aaatttga 38

<210> 2690
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2690
tcccctcgcc actgacccat ctactcgcca aatcttga 38

<210> 2691
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2691
tcgggcaata ccctttccca tattatcttc tttcctta 38

<210> 2692
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2692
tctcgggatt ttccgcgact actcgagcgc tactca 36

<210> 2693
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2693
tcgcttcctt gtacttccta ttccctttc accaca 36

<210> 2694
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2694
tcgttatagc ctatatacac gcggttgact cgctga 36

<210> 2695
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2695
tcttcacgat atgcagtgaa aacaccaccc aactta 36

<210> 2696
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2696
tctgcttgat ggtgctttca gtttcttact tctacga 37

<210> 2697
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2697
tcttctgtaa ccattttgtt ccggtttttt tattt 35

<210> 2698
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2698
tgaagctccg aaggggtgca cttgccgaca gtagta 36

<210> 2699
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2699
tgacgtctcc tgttcggtta accctgtgtg cctgt 35

<210> 2700
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2700
tgcccttctt cgtttactct tatccacatt accttga 37

<210> 2701
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2701
tcttgtcggg attttgctcc tttcatactc atggta 36

<210> 2702
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2702
tgacatatcg ctcttacatc ccctcacaca atttca 36

<210> 2703
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2703
tgctccgagt accataaact aatttgaagc tctac 35

<210> 2704
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2704
tgctgttctc ctggcgatag gagattagcc ggacgga 37

<210> 2705
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2705
tggaccggtc aaaggatggc accggttcct ctcaca 36

<210> 2706
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2706
tgcgtgagta acgcacaggg gaacactccg ttttgtga 38

<210> 2707
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2707
tgctttgtac aacgttggtg tttggcattt gctgttga 38

<210> 2708
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2708
ttacaaaccc cggtgctcta aagttatcga tctcttga 38

<210> 2709
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2709
tgtgccactc gtgttcgaat ttctgttcct tatct 35

<210> 2710
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2710
tgttacgccc gtcctctata caccaattac gatatga 37

<210> 2711
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2711
tgttaggcct ttatttccat tatcttaatt gtttta 36

<210> 2712
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2712
tgttttgctg cttagagcgt aatcgtattg gctct 35

<210> 2713
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2713
ttctgccgga tacatttgat tcaataacta cttgta 36

<210> 2714
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2714
ttctgcttgc gagccgtcca gttagggccg ccacc 35

<210> 2715
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2715
ttctgccggt ttttacggtc aaattctaga tagtttga 38

<210> 2716
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2716
ttgcacgccc caatacacct aatctttatc cataca 36

<210> 2717
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2717
ttcgttccgg accacccatt tcctcaatca tatattga 38

<210> 2718
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2718
ttctgcgttt tacgatgctc gatgtctgtg acatgga 37

<210> 2719
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2719
ttgctccgac ctagtttact aaatttcaac cattga 36

<210> 2720
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2720
tttcattata ccccatcgac tttctctcta gttcca 36

<210> 2721
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2721
ttgtcccagg cagtttccac gagaacgcgt acggt 35

<210> 2722
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2722
tttggacgta tcactcccta accatttttt taccctga 38

<210> 2723
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2723
tttgcttcat ttgcacttcc cccgttattc cgctca 36

<210> 2724
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2724
tttcctgttt cttctttgtg tgtccatata atcga 35

<210> 2725
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2725
ttgtagctgc tattgttttc tcgattgaga actta 35

<210> 2726
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2726
tttaaacgta cgatattact tccgatccat acatta 36

<210> 2727
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2727
ttttgttaaa caacgccggc aagggtcgtt gttacga 37

<210> 2728
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2728
ttttattggc tcgtttccta taatatgttg aatcc 35

<210> 2729
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2729
aattgtgaat cacaactcct ggtatatagg ctttga 36

<210> 2730
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2730
cggcatacga acctgtctca aatcctagta aggtaa 36

<210> 2731
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2731
agttgtgaaa cacaactcct ggtatatagg ctttga 36

<210> 2732
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2732
gagcttgata tcatcttcca cttcttcgct attcta 36

<210> 2733
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2733
ggtcataatg tgaatcatga atagtactaa tttactga 38

<210> 2734
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2734
tggcggctca cgtattctga aagttagtgg tgcgga 36

<210> 2735
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2735
tagctcatgg gttgcatcct caggataact ccgctga 37

<210> 2736
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2736
tcttattttc gcaagatctt ttatctaaat acaat 35

<210> 2737
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2737
tttcgaccat tgtgggattt ccggcggtcc aattatga 38

<210> 2738
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2738
actaggaaga acaggtacgt agggacacct gacagtga 38

<210> 2739
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2739
gacagctatc aacgaccgac aacggaaatt ggtgtga 37

<210> 2740
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2740
gggcgtccct tgcgttctct caacgaatgt gtcgcga 37

<210> 2741
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2741
tggcgggcgt gggtgttgcc gagacgtcag taccgtga 38

<210> 2742
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2742
tggagcggcg agttggaatc ctgatgtggg ctcatga 37

<210> 2743
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2743
atagataaat ggggaggtcg cgcattgaat gatggtga 38

<210> 2744
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2744
gccgtatgat taagcgatct ctggcatatg gcatctga 38

<210> 2745
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2745
tctcgaagtt ccgtttagcc ccaaggtcgc gggcgtga 38

<210> 2746
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2746
aaccgggtgt ccaaaccgta atttaatgtt tctggga 37

<210> 2747
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2747
aaccgggcgt ccaaaccgta atttaatgtt tctggga 37

<210> 2748
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2748
ataattcgcg tccggtcctt ggaaataaac ggcgt 35

<210> 2749
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2749
gggcggcctc gttttatcgg gaaaaggaac aaagt 35

<210> 2750
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2750
gggcggcctc gttttatcgg gaaaaggaac aatat 35

<210> 2751
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2751
gggcggcctc gttttatcgg gaaaaggaac aatct 35

<210> 2752
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2752
tggctgctgg acgcgtctta cagccgacca catcaa 36

<210> 2753
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2753
tagccctaat tgcgcttctt agcagggctc gtggttga 38

<210> 2754
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2754
tttaggcttt agggtacgac agcaataggt acaaa 35

<210> 2755
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2755
atagataaat ggggagagcg cgcattgaat gatggtga 38

<210> 2756
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2756
gagagaagta tttgttaagc cacatagttt tcata 35

<210> 2757
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2757
gagagaagta tttgttaagc cacatagttt tcatc 35

<210> 2758
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2758
gagcttgata tcatcttcca cttcttcgct attcta 36

<210> 2759
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2759
tcttatttta gcaagatctt ttatctaaat aaact 35

<210> 2760
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2760
tcttatttta gcaagatctt ttatctaaat acctt 35

<210> 2761
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2761
tcttatttta gcaagatctt ttatctaaat actaa 35

<210> 2762
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2762
tcttatttta gcaagatctt ttatctaaat actag 35

<210> 2763
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2763
tgggaggttc agaagtcttt ggccttgatc tgcagga 37

<210> 2764
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2764
accaccaaat ccccaaccac ttttccgctt atccca 36

<210> 2765
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2765
catgcagccc cacaccgctt ttatattgtt ttctc 35

<210> 2766
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2766
catgcatagg ctcactcctg gaacactacc tttat 35

<210> 2767
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2767
caacagtcat gtcaccctat tatttgcttt cattca 36

<210> 2768
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2768
catgccatat gtcccatcac ttacctacca tcccta 36

<210> 2769
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2769
catgcgatcc aatcctatag tccttggacc ttata 35

<210> 2770
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2770
catgcgatcc ctcacacaac tttcatccca atacc 35

<210> 2771
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2771
cttgcgatat taactttcca tcactttcct tcttt 35

<210> 2772
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2772
catgcgccga tacccatttc ccaattcctc ttcaa 35

<210> 2773
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2773
cactgccacg tcaaatgcct ctttcatttc cattt 35

<210> 2774
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2774
cactgcggtc gagcctccta attgttctca ccctta 36

<210> 2775
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2775
cgaacagtgg tacacgacgt agtgcactct ttttatga 38

<210> 2776
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2776
attgatctgc atgtgccatt ttcccacctt tgttt 35

<210> 2777
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2777
gcacatgcgt actctcatta acttcattac catcc 35

<210> 2778
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2778
gcgattaccc actcttgtaa actaatttgc ccatt 35

<210> 2779
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2779
gagacacgcc actctcacga cacgttcctt atttctga 38

<210> 2780
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2780
gataacgttt gattactcca ctgtgccata ccctt 35

<210> 2781
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2781
gccacaacac aattcttccg ccatcactgc ccaa 34

<210> 2782
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2782
gcctgaccac ccgtataatt taccgcctaa acctt 35

<210> 2783
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2783
gccacagtat cgttctcgag ctggttcccg ccat 34

<210> 2784
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2784
tacgactcat acttccattc cgcctaacct tttcca 36

<210> 2785
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2785
gcgtatcacc atggctcact tttcctatct ccttt 35

<210> 2786
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2786
ggccacgtaa cagctactca acgcgcaaca ctaac 35

<210> 2787
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2787
gcggccacac tttgccttct tattcgttaa ttccta 36

<210> 2788
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2788
tgctgtcaac gcgaagatct ttctctttta ttctt 35

<210> 2789
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2789
tatgcggcga acgccttcta tttcctggtt ctctt 35

<210> 2790
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2790
tatgccccac gataatttcc tctttttgca tattt 35

<210> 2791
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2791
ttacctacca cagtctccgc ctcgttatcc ttatca 36

<210> 2792
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2792
acaccacgcc tctgccggtg gacgtactgc ccatta 36

<210> 2793
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2793
gccacgtaga acacccatcc tgatccatac aatactga 38

<210> 2794
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2794
agcgcatcgt agccgtacag caatgtaata ttctc 35

<210> 2795
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2795
acgcacgtgc agtcgtatcc cgtttcttgc gcttta 36

<210> 2796
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2796
gacctagaag agggcgctta aaattgggag tggtt 35

<210> 2797
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2797
atgcactacg cagatgcaaa ctatctttcg ccttc 35

<210> 2798
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2798
cttgcgagat tagctttatc acctgttttc cgctt 35

<210> 2799
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2799
gatgtagctg ccacgagtgc gtatactcct ttatctga 38

<210> 2800
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2800
gcccagctgc ggtcacttgc ccagccattc caccaa 36

<210> 2801
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2801
gataacgttt gattactcca ctgtgccata ccctt 35

<210> 2802
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2802
gccacagtat cgttctcgag ctggttcccg ccat 34

<210> 2803
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2803
gccgagtcaa tcaagctcgg attacagtgt ttacc 35

<210> 2804
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2804
tactgcaggt acagagtatc ttcatccacc tccat 35

<210> 2805
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2805
ggtgccgttg aggtatacgg accgtgcagg ccatca 36

<210> 2806
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2806
taactccact gtgcgatcat tgtttttctt ttttga 36

<210> 2807
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2807
gtcgtgcgag gtcaagtact cctattctct ctaaa 35

<210> 2808
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2808
tggagggccg agtggtgggg gtttgagaag cattatga 38

<210> 2809
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2809
tcacgacgta caaaggttgg cgtacttacc ttactga 37

<210> 2810
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2810
tgcactgtag gtcatttcgt ttacttgcca ccatc 35

<210> 2811
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2811
tgtaacagca gtcgcgggga attattggct cttgaa 36

<210> 2812
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2812
tgtacaagtc agacgcgccc aaccctgttt acacta 36

<210> 2813
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2813
tgtacacctg caatgcgtgg tattactttg ttccttga 38

<210> 2814
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2814
tccacatgcc gttgaactct ttattcccct taccttga 38

<210> 2815
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2815
ttggtacccg atagaatcta gtttttgtag ccaag 35

<210> 2816
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2816
acggacatgc gagcaacaat gaacgtcttt ttttc 35

<210> 2817
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2817
aaccacgaag gtacggacac actttacata ccctta 36

<210> 2818
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2818
atgtacgcgg tacccatcac acactatgtg tcccttga 38

<210> 2819
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2819
agtcacacaa gcctccgatt cattcttcta aacat 35

<210> 2820
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2820
catgcgcagg tatcttgcta cactattctc ccgtttga 38

<210> 2821
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2821
atgcgcaccc gctatctacc cgtgacacac gcctta 36

<210> 2822
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2822
atctcgaacc accaccttcc atcccttatg ttgtc 35

<210> 2823
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2823
gcgaaccgga acttaatatc cttgtaaccg ctatc 35

<210> 2824
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2824
gcaacccgac ctcgctccaa ccgcttacga ctccta 36

<210> 2825
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2825
gcaccccaca tcccacgatc ccatggatac ccgcca 36

<210> 2826
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2826
gccaacctca ccaccggtac gccttatcgc taatta 36

<210> 2827
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2827
tcagtaccgc cattcccaaa aatttttccc caaca 35

<210> 2828
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2828
tgaagccagc tgacgtacat ccacatattc cctac 35

<210> 2829
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2829
tcaaccaact caccctcatt tactcgcata acatta 36

<210> 2830
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2830
tatgcccgcc tgactctctt attactggtt ccgac 35

<210> 2831
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2831
actgctggca ccgcaatcta cctaaagttt ccgtta 36

<210> 2832
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2832
accacgcact caggttcatg catcttgaac tttac 35

<210> 2833
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2833
gaccgtcgca ggccgtggct agcaggacgt tggcta 36

<210> 2834
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2834
gacgaacgtg ctgttggtat tttttgaact cttattga 38

<210> 2835
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2835
gacatcctgg aggtcctaac tgcggtctca cacttga 37

<210> 2836
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2836
atgccggggt cacgacgtta ccgcactcca tctat 35

<210> 2837
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2837
atcagcaacg cccagtggtg caagctttac ccttt 35

<210> 2838
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2838
gactaccgta cgtgcatggc ccccagttac acaata 36

<210> 2839
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2839
tacctcaacc gcgaactttg ctacgttttg taggg 35

<210> 2840
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2840
tacgctgtcg tgtgaggccc taccgctccc ctatc 35

<210> 2841
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2841
gtcacggaag cagttacgaa cgatgccctt ccatc 35

<210> 2842
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2842
tgtaacagca gtcgcgggga attattggct cttgaa 36

<210> 2843
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2843
ttgtgacgtc tcgtgcgcat taccggttaa tattt 35

<210> 2844
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2844
ttggtacccg atagaatcta gtttttgtag ccaag 35

<210> 2845
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2845
tttaacgtag tgagtcacca ttgatccttt aattt 35

<210> 2846
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2846
acccacgcct accctaatcc ttgtctgtta ttttc 35

<210> 2847
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2847
catgccgaat aatctctttc gttcctggtc ctttta 36

<210> 2848
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2848
catgccgcca aaccacactg ttccgtctta cttatga 37

<210> 2849
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2849
catgcgaaca acctttacca ttcactagca cccct 35

<210> 2850
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2850
catgcgacac gctctatcgc tttcacatct tcgcttga 38

<210> 2851
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2851
catgcgatcc ctcacacaac tttcatccca atacc 35

<210> 2852
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2852
catgcgcact ttcctcacat gataccctgc atgac 35

<210> 2853
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2853
catgcgcatc caatcacact ttttaatttc atttc 35

<210> 2854
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2854
atgcctagca gccaagcaat gcctttcatt tattc 35

<210> 2855
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2855
gccgcagcca tcatctacgc ttcacactca tcacca 36

<210> 2856
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2856
ggtacgcagg cgcgatttgt acatttttga tcctga 36

<210> 2857
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2857
tctaacgtac tatcatgtgc cacgccttca tttcctga 38

<210> 2858
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2858
tcaactgcca ccgaccccaa cccgcgaccc atcct 35

<210> 2859
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2859
caaggagcac gtatgcactg tgcgagttgg tccgta 36

<210> 2860
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2860
cgcgagccag ggtactgcct gcggtctaaa caataa 36

<210> 2861
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2861
cgcgagccag ggtactgcct gcggtctaaa caatca 36

<210> 2862
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2862
cgcgggccag gtctgcatgc atgcgaggtc acacc 35

<210> 2863
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2863
taacacgtct ttcactctac tgtgcccttt atgcc 35

<210> 2864
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2864
gcggagctag cacggtactg cgcataaact atgattga 38

<210> 2865
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2865
tctgcgattc taccgttact ctaccgtgcc accaaa 36

<210> 2866
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2866
tgggtgtcgt actggtacca aggagtgcga aatcta 36

<210> 2867
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2867
tgcgtgctaa acggtctagc acatgcgggc ctcac 35

<210> 2868
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2868
acgcgcataa accaatcact accgaccact tattc 35

<210> 2869
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2869
acgtcaggca taaattcttc aaactcttac tatca 35

<210> 2870
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2870
catgcccgga ccattattat tcgcttacca ttccca 36

<210> 2871
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2871
catgccgacc actgttcctt ttacctctta tacctga 37

<210> 2872
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2872
catgcgaaca gacgaatgat tcttacaacc gcaac 35

<210> 2873
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2873
cccgtccgca ggcatccacg tttatgcctt ttatc 35

<210> 2874
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2874
cacgcgaccc tcattaacat catctaaccc aatcca 36

<210> 2875
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2875
catgcgccaa ttttactgta catttctttt tttat 35

<210> 2876
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2876
catgcgcgcc gacctactcg cttaatctca cgttt 35

<210> 2877
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2877
catggcgacc ctcaatctca cattttacct cttct 35

<210> 2878
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2878
gacggactgc gtatcgcata gtttatcctt ttttt 35

<210> 2879
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2879
gcccaccata tcgcaatcat attcctcgcc ttgtt 35

<210> 2880
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2880
gcgcatgtca accttacctc tattggccct tattc 35

<210> 2881
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2881
gcatccagtc cataactcaa tctcacctgt aattc 35

<210> 2882
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2882
gccgcatacc cctgtaattt cgtaacctac accca 35

<210> 2883
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2883
gcgcaactcc gactatcact gccttaccaa tcac 34

<210> 2884
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2884
gcgacgcctc tgacctacat ataactttga tagtaa 36

<210> 2885
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2885
gcgacgcgtg aacaccctta ttctgtaatc ttttt 35

<210> 2886
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2886
gtactgtcaa cccgatcttt gtttgtattt atttt 35

<210> 2887
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2887
gcgcggtact cattatcctt atatcatctt acatttga 38

<210> 2888
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2888
tatgcgagcc catttttcac ctttgctatt tcttaga 37

<210> 2889
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2889
tcctccagca cattctacca cactaccaca ttcat 35

<210> 2890
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2890
tatgcggtgc cacttctcta ttacttctat tactt 35

<210> 2891
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2891
tcatgcagcc tagtatctat cctaccatat tattca 36

<210> 2892
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2892
tgacggacag tgcatgcgtt atacttattc tctat 35

<210> 2893
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2893
aagcctggcc gtggtgctgc agtttattcc gttagtga 38

<210> 2894
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2894
cgcgagccag ggtactgcct gcggtctaaa caataa 36

<210> 2895
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2895
cgcgagccag ggtactgcct gcggtctaaa caatca 36

<210> 2896
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2896
gcattcgcgt actgtggaca acctatatcc gggtaa 36

<210> 2897
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2897
gcaactatcg cgagcgcctt caagattagc gtgtca 36

<210> 2898
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2898
gcgcaccgac ctctaccgtg ccaaaatagg ttatctga 38

<210> 2899
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2899
gtgcggggcc acaggtgata gcaatgcatg ccctatga 38

<210> 2900
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2900
tcaccccaag cagtcattgc ctacaggagt ctggtga 37

<210> 2901
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2901
tgcagttagt gcaatgaata tctcggacac gcggaga 37

<210> 2902
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2902
tccgcagaca gccgggggtc aggactacgg ctacg 35

<210> 2903
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2903
ttacagctac tctgcgctcc aagtggacac tttacga 37

<210> 2904
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2904
ttagcatcac tgacttggat acatacagca tgtcga 36

<210> 2905
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2905
agcacgactg caatccacat ccgttcttat tcctt 35

<210> 2906
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2906
aatgcgcaca tctgtttttg taccttagaa tgtttga 37

<210> 2907
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2907
acgtcaggca taaattcttc aaactcttac tatca 35

<210> 2908
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2908
catgcaccac tcgcacattc caccacaatc tataca 36

<210> 2909
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2909
ctgccactag ccgcccatac tatccacacc ataat 35

<210> 2910
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2910
ccatgcgcgg tcacacatac cagtccttac cattt 35

<210> 2911
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2911
catgccgaca caccactttt ttccgaccga cttcc 35

<210> 2912
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2912
catgccgcct ttcgacttcc atatcccaac acgac 35

<210> 2913
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2913
catgccgcct ttcgacttcc atatcccaac acgcc 35

<210> 2914
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2914
catgccggtc attccatact caaactcatt cactc 35

<210> 2915
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2915
cttgccgatc ccgctcccta gttcccttgc ccact 35

<210> 2916
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2916
cacgcatgtc tgcgctcgta tcttttcacc ctatga 36

<210> 2917
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2917
cactgcggtc aacacccacc aacctccgac cttactga 38

<210> 2918
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2918
catgcgtacg ccactttcgc ctactctaac atatctga 38

<210> 2919
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2919
cagtctgcgc gacccttttct gttccttcat cccca 35

<210> 2920
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2920
cattgcgcct tctattttat tttccttata tttgt 35

<210> 2921
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2921
cattgcggcc attacccctc taccgttacc ctttttga 38

<210> 2922
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2922
gcacacacct gcggagcacc actccaaatt accca 35

<210> 2923
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2923
gcacactacg tttaccctgg cccttctaat tgttt 35

<210> 2924
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2924
gcactgccat tgatttcttc ccacttcaca ctctt 35

<210> 2925
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2925
gcgaacactg atcttaacta acgcccgccc aattta 36

<210> 2926
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2926
gcacccccaa tgaccgccat caaaccatat attcca 36

<210> 2927
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2927
gcgcaaccat cactataacc catttacctg ttttca 36

<210> 2928
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2928
gcgcaccaca cttgagttct ttaactttcc ctttatga 38

<210> 2929
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2929
gccgacgtac ctgctgggtc tctctattca ttttt 35

<210> 2930
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2930
tacgccgtcc accatcctct tctatatacc taacc 35

<210> 2931
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2931
tactacgaac catcccactc tttgcacctg ttttc 35

<210> 2932
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2932
tcatgcgcgg tacacatttc ttaatcactt ttct 34

<210> 2933
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2933
tgcgggcatt ggtgcaatct taacttattc tttat 35

<210> 2934
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2934
tggtagtgct gcatcaccga tcatcctttt acatt 35

<210> 2935
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2935
tccgcgactg ttcatccccc atcccaacat aacccga 37

<210> 2936
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2936
tgtgaacgtc actgcaccct aaactaatat tcccttga 38

<210> 2937
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2937
ttgacaactg ccactaccaa ctgaacccgc tactta 36

<210> 2938
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2938
ttgtcaagcg ctatccacca tttctctttt atctatga 38

<210> 2939
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2939
caatgcgacg aggcttctat gtcatgaact aagct 35

<210> 2940
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2940
ccgaacggtc gcaaagtggt gtagcctcta atccgga 37

<210> 2941
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2941
catggcgacc ctcaatctca cattttacct cttct 35

<210> 2942
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2942
accggaatgt acatgccctg gtttttgtta ttatttga 38

<210> 2943
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2943
accatgcgcc caccactatt ttttccactt cctta 35

<210> 2944
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2944
acggagtacc ctgcccacta tataacttaa accttga 37

<210> 2945
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2945
acgggcgtgt cgcactgtct gtcactggct tattc 35

<210> 2946
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2946
agacggtggg tgggtgggtg gtgctgcagt ctacc 35

<210> 2947
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2947
catgccgcac cccaacacgt accatttcat tattt 35

<210> 2948
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2948
catgcgggca tccttagctt aacctcaatt ccatta 36

<210> 2949
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2949
catggccatt ctaccctcca ttccattcct tgcct 35

<210> 2950
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2950
cgttcggtac gcagtcattt cactctatct tgttc 35

<210> 2951
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2951
aacccacgag ttgttggtcc acgcctcgtt atcacga 37

<210> 2952
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2952
aacccatgcc tgccccatgt gcattcaggc agagca 36

<210> 2953
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2953
accgccacag tcatcaaccc acctggccga cgacga 36

<210> 2954
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2954
gacaacgcaa ccgcaggctc atggagccaa tatgta 36

<210> 2955
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2955
gacaacacgt cgcatgatct taaaagagga cctagga 37

<210> 2956
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2956
gagtggtcgc cactcacaac gatgcggtat ccccttga 38

<210> 2957
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2957
gcgtggtggt ccgccggtgg tgagtcgatg aactgga 37

<210> 2958
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2958
gtgatcgata gcgagtgttg agctgtcgag agtgga 36

<210> 2959
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2959
tcacacaccg cgtggctcgg acgacaagta gtccta 36

<210> 2960
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2960
tgaacctgca cagctgttaa gaggtggctt acgta 35

<210> 2961
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2961
actgaccaca gctacgtaag caataaccga tcgtt 35

<210> 2962
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2962
aagagtacgg gcatgagtaa taagggagta gccgt 35

<210> 2963
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2963
agatcgcggt caatcggtgg gtagttagga tcggg 35

<210> 2964
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2964
gaacacacgt cgatcgccac tgagagcaga tgtca 35

<210> 2965
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2965
atcggaggcc gtatcaccat atcccaccac cactt 35

<210> 2966
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2966
cttattttag caagatcttt tatctaaata caat 34

<210> 2967
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2967
gacaaacgta gtgctgaaca tggcgttctg gtccaa 36

<210> 2968
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2968
ataccgggac aggcgacaga agggtgaacc aatgt 35

<210> 2969
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2969
gagtggacgt ggacgcgaaa cgggtcaggc ataacga 37

<210> 2970
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2970
taccacgtgg cgcctgacga cttacgcaac tgggga 36

<210> 2971
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2971
tacagccgac atgaacatcc cttaatccca ccctaa 36

<210> 2972
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2972
tgataccgcg caacgagact taccacgtca ggatgga 37

<210> 2973
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2973
atggtgggag caccgacgta ggaggcctat ggcgta 36

<210> 2974
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2974
gacaacacgt cgcatgatct taaaagagga cctagga 37

<210> 2975
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2975
gtgaacgtcg ctagaaacga tggtgtaaat agtcttga 38

<210> 2976
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2976
gtgatcgata gcgagtgttg agctgtcgag agtgga 36

<210> 2977
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2977
aacacgcaac acacaactct cggcaaatcc agcgcga 37

<210> 2978
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2978
acgacgcacc ctgccactat cgtagatact ccgacta 37

<210> 2979
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2979
gaccaatgga ccggggacta ctctgagaaa gcgcga 36

<210> 2980
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2980
agtggaacgc accgaagtta cctacctgag atggga 36

<210> 2981
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2981
gacgacgggg atcctgacaa ttcctgacta agcgcga 37

<210> 2982
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2982
gacccgccca gtaagcacac acacagccca aattt 35

<210> 2983
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2983
gagcggactc gaatagaact ccgcaacacc agcgcga 37

<210> 2984
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2984
gagacgaggc cgggatcatg ctagggatcg gacagga 37

<210> 2985
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2985
taccgacacc ctcgtgttcc caaacaagga gacgga 36

<210> 2986
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2986
ggatacgcac cttaaggagg gacgaataaa agcgcga 37

<210> 2987
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2987
tagcacagtg gtgcttacgg aacatcacca gcgcga 36

<210> 2988
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2988
tagtcggtcg gcctgccgat gcgtcaaggt agcgcga 37

<210> 2989
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2989
taggaatcac cggctcgggc gaggccggga agcgcga 37

<210> 2990
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2990
tgagccgcaa ccggcaaaca aggcttcaac agcgcga 37

<210> 2991
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2991
aacaccacac cgccgctgat ccggatggaa cctcga 36

<210> 2992
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2992
aagagtacgg gcatgagtaa taagggagta gccgt 35

<210> 2993
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2993
acacgtgccc caccaatatg cgataagact ttgtta 36

<210> 2994
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2994
gacaccgaag acgaggctaa agagagaagt gccata 36

<210> 2995
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2995
atcgccgcac acctgccact cactcttcca tcgct 35

<210> 2996
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2996
cttattttag caagatcttt tatctaaata caat 34

<210> 2997
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2997
gactaacgcc gcttatcctc aacgcctaac cttgtt 36

<210> 2998
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2998
gagtggtcgc ctgcaggtgc attccatcca gctccga 37

<210> 2999
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 2999
gtggaggctg caatacggtg caagagagct taggg 35

<210> 3000
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3000
gtcaccgcat gcgggtcaag ccctcaactt tacttga 37

<210> 3001
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3001
taagtcgcgc cccagtcagt gattgataca gatagga 37

<210> 3002
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3002
tcaacatgcg tgcaagtggc accgttctcg tttcga 36

<210> 3003
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3003
tcttatttag caagatcttt tatctaaata caat 34

<210> 3004
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3004
aacacgcaac acacaactct cggcaaatcc agcgcga 37

<210> 3005
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3005
acgacgcacc ctgccactat cgtagatact ccgacta 37

<210> 3006
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3006
gaccaatgga ccggggacta ctctgagaaa gcgcga 36

<210> 3007
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3007
gacgacgggg atcctgacaa ttcctgacta agcgcga 37

<210> 3008
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3008
gagcggactc gaatagaact ccgcaacacc agcgcga 37

<210> 3009
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3009
gagacgaggc cgggatcatg ctagggatcg gacagga 37

<210> 3010
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3010
tacgcgggtg gtcgacccca ctagtagtca tccata 36

<210> 3011
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3011
taccgacacc ctcgtgttcc caaacaagga gacgga 36

<210> 3012
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3012
ggatacgcac cttaaggagg gacgaataaa agcgcga 37

<210> 3013
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3013
tagcacagtg gtgcttacgg aacatcacca gcgcga 36

<210> 3014
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3014
tagtcggtcg gcctgccgat gcgtcaaggt agcgcga 37

<210> 3015
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3015
taggaatcac cggctcgggc gaggccggga agcgcga 37

<210> 3016
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3016
tccgaacggg acgtacacca agccgagctc tatcctga 38

<210> 3017
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3017
tgagccgcaa ccggcaaaca aggcttcaac agcgcga 37

<210> 3018
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3018
accatcgggt cacgtaaaag cggttctaat ttctaa 36

<210> 3019
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3019
acccacatcc tgcgtgctaa ggacggtggg gccgg 35

<210> 3020
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3020
gaacaggcgt tccctactga tttatctgtc cccata 36

<210> 3021
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3021
gaccatgcga gcggcaaagc acggaaaatg tttgaa 36

<210> 3022
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3022
gagtcatgtc cgcctttaac ccgacttaag aactttga 38

<210> 3023
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3023
gcaagatgaa gttggtactt tcctctctat tcttttga 38

<210> 3024
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3024
gccacaccgc ccgtacatac ttcctcacca acattga 37

<210> 3025
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3025
gcgcgcccac cagtatcttt tggtctagat aggca 35

<210> 3026
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3026
gggcccacgc caccgtttgt ccacaggaag gatac 35

<210> 3027
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3027
gtgcggtcat ggaattcttt taagcatgat aatacga 37

<210> 3028
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3028
ttgcaaagcg caccgtccga gccggtggca ccgaa 35

<210> 3029
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3029
ttcacgccac attcacggga actgagagtg acgacga 37

<210> 3030
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3030
tttgcggtca ttaacttgcg tgtaggacac cggtgga 37

<210> 3031
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3031
acgcagtgcc gccgaatcct gatcccctaa ctttc 35

<210> 3032
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3032
aatgcgtccg acctaactgt tgcgattctt ctttt 35

<210> 3033
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3033
acacacaagt ccatgcgggc ctcacttttc gcctatga 38

<210> 3034
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3034
agtcccaact gcagggttta tacctctggt tctctc 36

<210> 3035
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3035
gcacacgccc acattgtcct actagctatt ttatta 36

<210> 3036
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3036
gccgcccctg agtagttttc gtttttcttg tcctctga 38

<210> 3037
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3037
gcgatcgata tgttgcctat cctcttacga catct 35

<210> 3038
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3038
gcgttgtccg tgtccatcca tcttgtcatt tgttt 35

<210> 3039
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3039
taacgtacaa ccgggcgcac aggccattct tctac 35

<210> 3040
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3040
gttcggggtg catgtcgtga agccgctaac ctctt 35

<210> 3041
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3041
tgctcagtca agcgagtacc actacgccta ttccgtga 38

<210> 3042
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3042
tgctaaacgt actggggctt cctcagatct ttgcttga 38

<210> 3043
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3043
tgtacccccg ccgtaatcat ctatcttgtc cccgaa 36

<210> 3044
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3044
accactgaac tcatggtact aatcacacac caatt 35

<210> 3045
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3045
actgtacagt agtccacgaa catatcgctt tctatga 37

<210> 3046
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3046
accgcgcgct gcagtatatc ccacttcgtt ctatatga 38

<210> 3047
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3047
agtggatcac tgggcggagt gcgttaggaa gacgg 35

<210> 3048
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3048
gattacgcct taagtggtac tcagttatga ggaga 35

<210> 3049
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3049
gcgtaggtca tgcgacgtat aaccttccca caaatga 37

<210> 3050
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3050
tacaccaaaa agatcggtga ttaccatgca acaagga 37

<210> 3051
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3051
taccagacgt accgcacacc tttcatctgc ctaac 35

<210> 3052
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3052
tttgagactg cgggctgtgc gttgctttga tatcc 35

<210> 3053
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3053
actgactgca tacatccgca tattaaaccc gtatt 35

<210> 3054
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3054
acatctccag gattcccctt cagccatggc cctaca 36

<210> 3055
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3055
acgctcagct ggcatgtacc ccccctctct acacttga 38

<210> 3056
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3056
actcgcgtgt ccaacacgga ctgcagtagc aatcca 36

<210> 3057
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3057
gcaggtggtg caggttattc atgcgaaagg gtactga 37

<210> 3058
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3058
gcggtggaaa cgtaatcgga cactgcgcat gcgaca 36

<210> 3059
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3059
tgaacctgcg atggtccgtg cgaacccgat ctcaca 36

<210> 3060
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3060
tcacgtcccc aacaacccgg agtaattccc tgcca 35

<210> 3061
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3061
tgggtacaca tccggcatgc caataacgct taagt 35

<210> 3062
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3062
acgcctgtac gatgtcctcg gtctcatttt atttc 35

<210> 3063
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3063
aaacgtcaag ggaaagcgta gtctcctcga ccctt 35

<210> 3064
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3064
actgggaagt acgctgtcag gcatttgcta actaaa 36

<210> 3065
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3065
aatcattgca ggtatatgcg agcgaactac gattc 35

<210> 3066
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3066
atacgttccg gccaactgaa acggagttcg ggtct 35

<210> 3067
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3067
atgcatagct gcccttctta tccgtttcca actaca 36

<210> 3068
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3068
atgcgaaact gtaagcaccc ttccaccttg taccatga 38

<210> 3069
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3069
gacgccgaaa cactaagctt ttaacactta cgatatga 38

<210> 3070
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3070
attatcccag cgatcccgag tcccggcatt tactg 35

<210> 3071
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3071
catgcgcata tcgctccgtt cgttcctcaa tttcca 36

<210> 3072
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3072
gcccgaaccc gaaaaccgat cgcctgaatc ccaattga 38

<210> 3073
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3073
gcaggctcgc gtagcttcaa actagcagac cacgt 35

<210> 3074
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3074
gcacgggacg caagctagac tacacaccga ctttttga 38

<210> 3075
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3075
gctggcactg cgaatatcga tctacactat cccactga 38

<210> 3076
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3076
gctagcacac aggatccctc tttattagca cttcaa 36

<210> 3077
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3077
tatgcgaccc gttaaacccg tactcttcct aaact 35

<210> 3078
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3078
tagtcatgcg cgcatctcaa aatttacttc tcaaatga 38

<210> 3079
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3079
tctacactcc gctccttgac ctaatttaaa gaccttga 38

<210> 3080
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3080
tcgcggaaac ctgcccctgc atcgtagcca tccac 35

<210> 3081
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3081
ttcaccacca ccccttttgga ccctttttacc gagaga 36

<210> 3082
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3082
ttgcagacct catttacgca cttgggactt ctgcca 36

<210> 3083
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3083
ttggcacctc cactgcttgt tctcgattca tgtgcga 37

<210> 3084
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3084
actgcggtca tcgagttact ggctattgcc tgacc 35

<210> 3085
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3085
cccgcggtaa tagtgcgggc ggacgcggtc atgaca 36

<210> 3086
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3086
gaacgtagac tgactacggt ccacacgccc attgta 36

<210> 3087
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3087
ccaacgacag atatggacgt gctgcggggg caatc 35

<210> 3088
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3088
atgcgatgtc acgtgacgtg agtaaccgcc caccc 35

<210> 3089
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3089
gcccaccctt gccagcatgt atgcttgcat gtaac 35

<210> 3090
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3090
gccggtcatg cggtttgcac accactccat tctac 35

<210> 3091
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3091
gccaggccca ttccttgcgg gaggttaaca ccagca 36

<210> 3092
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3092
gcggtggaaa cgtaatcgga cactgcgcat gcgaca 36

<210> 3093
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3093
gtttcacccc ggctgggcta gttctgttaa tgcgaga 37

<210> 3094
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3094
tgtaccgcgg ttgcgactat ttccttactg cccata 36

<210> 3095
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3095
ttcgcgacga taactagttg catgtctgaa ttcac 35

<210> 3096
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3096
tgtcactgca cgatgagact ttcggcgaat atactga 37

<210> 3097
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3097
aatgcattgc gttatttcaa ttgaggcgtc ctaaaa 36

<210> 3098
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3098
gcactgccaa ctttcttctg gggactattt agctttga 38

<210> 3099
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3099
ggtaccaaaa gtactgccat gagagccact tcccaa 36

<210> 3100
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3100
catacgcacg gtcctctact ataattgtcc accaa 35

<210> 3101
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3101
atgctgcggt aatggagccg gttcttaaac tgcct 35

<210> 3102
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3102
agttactgac aggcgcgtct ttcgcaacac tttata 36

<210> 3103
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3103
gcacccgagc ctaatagtcc tctattcccc ttcca 35

<210> 3104
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3104
gcgagcagtc catggcccat cctcgactta acctca 36

<210> 3105
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3105
ggtagcaggc ataccctacg gaacagttct gtggca 36

<210> 3106
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3106
taggcacgtg catcaggcgt tctttgcccc tgttta 36

<210> 3107
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3107
tgtgcagtca gtgaacggtg acctagttat gattt 35

<210> 3108
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3108
agcccgttgc tacgttgcag taggtgtaac ttcactga 38

<210> 3109
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3109
actaactgca agcgcccttg gcccgatttg acctttga 38

<210> 3110
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3110
actaccgcaa actgcactgt ttaaaactcc ttgaa 35

<210> 3111
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3111
acgactatcg tacatgtgcg tagtttagat tccaa 35

<210> 3112
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3112
agttgaccac ggagtgtact gcaatggagt gtaccga 37

<210> 3113
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3113
gaacgagccc ggatgttagg atcctcgatt tgcgta 36

<210> 3114
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3114
agtgctgcca ctaagctaga aaagttttca tgccc 35

<210> 3115
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3115
atcgtaacgg actgctattt gtacggatcc ctataa 36

<210> 3116
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3116
gcagtctagc gcccgcttgt accacatcct cgctttga 38

<210> 3117
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3117
tgtactgagt gcacaacacg tgatgatgat atctt 35

<210> 3118
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3118
agcggtcacc gcctatggtt attcattttt ctctta 36

<210> 3119
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3119
actcagcgat gccctaactt ggacgttcgg ctact 35

<210> 3120
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3120
cacaccgacg aaactgtcct cgcctactac ctatc 35

<210> 3121
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3121
agtgaggcgt cacggctatt actccgtgct aattt 35

<210> 3122
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3122
catacgcacg gtcctctact ataattgtcc accaa 35

<210> 3123
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3123
gaaacggcca tgaacctttt aatctttcca ccttt 35

<210> 3124
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3124
gcgagaggat caccgtgagg cttcccaact atttta 36

<210> 3125
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3125
gcgaggcgcc ctccaaccca acccattcaa tggta 35

<210> 3126
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3126
gcgcagtccg tccacatttg gtctactgct ttcct 35

<210> 3127
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3127
gtgcgcctac atatatctcg gttacaacct ttttc 35

<210> 3128
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3128
tgcatcccac cactatgata tacctaatgc cctct 35

<210> 3129
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3129
tgcaggccgc atcacctctt gtgccggttc atcaaa 36

<210> 3130
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3130
tgcgaggcta gcttgttgcc actatttggt ccttatga 38

<210> 3131
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3131
agcccgttgc tacgttgcag taggtgtaac ttcactga 38

<210> 3132
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3132
agggcaacga acgcaggaat ccctcattcc tacgg 35

<210> 3133
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3133
agtgctgcca ctaagctaga aaagttttca tgccc 35

<210> 3134
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3134
atcctcgtta cgtgcacgga cttctcatgc gaact 35

<210> 3135
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3135
gcccaggttt gacgactgcc gtgcgctcgg tcttaa 36

<210> 3136
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3136
ggtcatggcg caacttcatc ttttcacacc tgattga 37

<210> 3137
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3137
ggtagcaggc ataccctacg gaacagttct gtggca 36

<210> 3138
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3138
tcccgcaatt acgcagaccg actacgattc caaaa 35

<210> 3139
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3139
tgtacgtagg taagcgatac acactgcacc cctactga 38

<210> 3140
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3140
tgacctctgc agtggccagc ccattcttta gcttc 35

<210> 3141
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3141
ttactgcagg tgcgatcgcg ttctccaaca tctct 35

<210> 3142
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3142
accacgttgg ttattgccct gctttgaagt cttacga 37

<210> 3143
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3143
aacacactgc gcctttactt taattttcga ttatta 36

<210> 3144
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3144
agcaccgacg tgataggact gtaatgccct cctct 35

<210> 3145
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3145
atgcgcgagt cataaggcgt ttcttttcac tggct 35

<210> 3146
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3146
atgcgacatg cgttacaata atccgcgcaa gccgc 35

<210> 3147
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3147
gagtgcgcgt acaatttgtc ctcttgccgc ttttt 35

<210> 3148
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3148
taagtcatgg gcctcatcta tacctgatcg atctc 35

<210> 3149
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3149
tcaggcggga tatgcgctca gggaagcggg gagcaa 36

<210> 3150
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3150
tgagctcgac ctgccgtcgt acaatttccc ttactga 37

<210> 3151
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3151
acttaccgct taatacactt gccggacctt gtgcttga 38

<210> 3152
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3152
actatagcca cctaacccag cactaaaccc acgca 35

<210> 3153
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3153
acccccccgc ttcgtgcact tcactcccga aatcc 35

<210> 3154
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3154
gccgcaccat aactttagcc tacccagtgc cctgc 35

<210> 3155
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3155
tgccaccaaa tacccaaccc tagagtagcc ttatta 36

<210> 3156
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3156
actctagcca cttgatccac cacgtcggtt cccacga 37

<210> 3157
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3157
ttagcgataa acacatcatc ccggacacgc cgtgt 35

<210> 3158
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3158
taccgtcgct caaaaccggc ctaccctaaa cccactga 38

<210> 3159
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3159
accgccgctc tactctgtca ccgaacgccc tttggga 37

<210> 3160
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3160
agcccgaccc gcttgcctaa acccacccat gacactga 38

<210> 3161
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3161
aggtacccac cgcaaagcac cagtaaccac aaccc 35

<210> 3162
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3162
gcatccgacc aatagaccca cacatctgta ccaaaga 37

<210> 3163
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3163
tcccgccgct accgtaagcc actccgaagt actca 35

<210> 3164
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3164
acgccagacc acccgatatc cccccaaatg gtaac 35

<210> 3165
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3165
aaacccactt cttacccggt aatctgccga ccgcatga 38

<210> 3166
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3166
acgcgtgggc tactaaactc catcaccccg tgccc 35

<210> 3167
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3167
actattccgc atcattgagc acccgaccga ccaaaa 36

<210> 3168
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3168
acacaaaggc tataaccgcc gactctccac cagcc 35

<210> 3169
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3169
accaacacca accttctacg ccgtaccctc tgcaaa 36

<210> 3170
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3170
aaaccgctat ttaccccgga ccacgttgac accga 35

<210> 3171
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3171
tacgtccacc ccgaatccta attcatacac ctaaaga 37

<210> 3172
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3172
aaatgccaca tcacttattc accccgcgca ccccga 36

<210> 3173
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3173
acaatcacct tacaccttac gacctgctta atagaga 37

<210> 3174
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3174
gttcgtgtac gtcgcattaa ccctaccata aaacc 35

<210> 3175
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3175
gcgtatcaaa aaacccaccg ctaactacca caaccga 37

<210> 3176
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3176
gtgaacccac ctgcgcccta cactattaaa cgagctga 38

<210> 3177
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3177
tcaacatgcc cacatacacc agacacacta tacac 35

<210> 3178
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3178
catgcaaccc gcacttaggc gcgagccaat tacca 35

<210> 3179
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3179
acccccaatc cttataccat aaacacatga ccggtga 37

<210> 3180
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3180
accggtcaac cccccacaga acatttcccg tcaccga 37

<210> 3181
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3181
acatgaatcc aactaaaacc cgctccgtgc cgaacga 37

<210> 3182
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3182
atcccccaaa cacaaacgcg agtttcccgg gtcca 35

<210> 3183
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3183
ctctgcgcga gtcaccctca cctagatccc tgatatga 38

<210> 3184
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3184
ctacccaacg gtcacaccca ccgcgtatcc actac 35

<210> 3185
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3185
ttacacccgt caactaacca tacccacgag agagc 35

<210> 3186
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3186
accccactac taccctctcg tccgcctaac tgtc 34

<210> 3187
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3187
accagtccat atcccgccat tgtgtcgttc gatcga 36

<210> 3188
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3188
gccacccccct accctaacga cccaatatga ctggtga 37

<210> 3189
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3189
caccaaatat caagtgcctc tctaaccttc tgacctga 38

<210> 3190
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3190
actgtcacta ccccacgcgc aaccaacaaa taccca 36

<210> 3191
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3191
agctcgacta aaccacccgt ataaaagcta tacccga 37

<210> 3192
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3192
gagcgcccta cttcaactcc caaaccacat gaccg 35

<210> 3193
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3193
aacaaagttc acccatccca caaccccaat tcgac 35

<210> 3194
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3194
cttctgaccc gtagaccaca acttcccgac ttcccga 37

<210> 3195
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3195
accccatacc attaactccg gattccccag tgctc 35

<210> 3196
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3196
agccacccac aaaacgcgca atcagtaaac ccacctga 38

<210> 3197
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3197
tcctaacgct gtcgacgcca ccacccggac ccact 35

<210> 3198
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3198
ccaatacccc cgcgcccaat cccacacgtt atcgta 36

<210> 3199
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3199
agcttaccca ctcccgaaag acaccctgac accgta 36

<210> 3200
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3200
atgacctaat ccaccacccc tgcacctcag tctaca 36

<210> 3201
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3201
atcccgcaca cctaaagcaa acgtggctta ggtcc 35

<210> 3202
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3202
atcctccgcc cgggcactaa ccagctcctg accacga 37

<210> 3203
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3203
actgcgccca tcagatacca aaacctgcaa accgc 35

<210> 3204
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3204
gccacccgcc tttagcccta cagatacaca cactaa 36

<210> 3205
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3205
acccccctacg ccggctgact aaaccgcaca ccctca 36

<210> 3206
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3206
gtttaccgta ccccccttgca ctgcccgacc aaccca 36

<210> 3207
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3207
actggaaccc caatcggagc gaagcctgac taccg 35

<210> 3208
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3208
tgccaaccta ccataccatc aattgacaga tacctga 37

<210> 3209
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3209
tcacccacgc catatactgc actccaaacc ccgctga 37

<210> 3210
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3210
aaactgttcc tgtgacaaaa ccctcaaccc atacca 36

<210> 3211
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3211
acgccaccta cccacctcac tatgaaatat actcgga 37

<210> 3212
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3212
gtccccttaa gtccacccat acacgttact gaccc 35

<210> 3213
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3213
gagaacccac cagcaccaat tcacaaacac ttacgga 37

<210> 3214
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3214
acaagcctaa cccgacccct agttctcacg ttgaca 36

<210> 3215
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3215
acaccctcca aagcacaaac ctacccatta gccaatga 38

<210> 3216
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3216
aagtacccca actcactgta ggtcactacg acaaca 36

<210> 3217
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3217
acgccgaccc gatattttag caaatatcac ctacc 35

<210> 3218
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3218
acgttaacac ccgcctccca cacccacttg acaggtga 38

<210> 3219
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3219
catcaagcag tcaacctgac tcatcccatg taccc 35

<210> 3220
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3220
acgcagttcc accacccatt gtcaccaaga tcatcga 37

<210> 3221
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3221
ctatgcaaca cccgccttcg cacgttcaca ctaccga 37

<210> 3222
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3222
agcatgacac ggcacctacc acccgaacga ccatc 35

<210> 3223
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3223
ttaataccga tttacacgct acccgtcccg ggaggga 37

<210> 3224
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3224
aatacgctcc agctcaccta accctactaa tcctcga 37

<210> 3225
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3225
acggaacaat cccccactat gctccacttc ttact 35

<210> 3226
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3226
gcacacaaac attccaccag gtgacaaccc cgaca 35

<210> 3227
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3227
atccgctctc acgacgccga ccaaacacat tccggga 37

<210> 3228
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3228
tctgcacgcg aagctccgcc gaccacaata caaac 35

<210> 3229
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3229
tgaaatcgac cacttgagag cgtacttaag cacccg 36

<210> 3230
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3230
aaccgaatcc caggacctaa ctcatcccat gaaa 34

<210> 3231
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3231
atcctcctct gacctcgatc tcccttacca aacgg 35

<210> 3232
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3232
tggtccaacc cccacgccgc catcactgga caacttga 38

<210> 3233
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3233
gcaccgcgcc tctaccctac ttacctaccc catcaa 36

<210> 3234
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3234
ggggcgcacg aaatcagaac aaccctaaaa ccccta 36

<210> 3235
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3235
gactacccgc tcactcaccc gtcccaaatt agatatga 38

<210> 3236
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3236
tacgccatca aacccgatac ctaaatttag cacac 35

<210> 3237
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3237
ttgcagaaca cgaccaccac cgctttagag ccacca 36

<210> 3238
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3238
taactcggtc ctgatgctct acccgtatca accct 35

<210> 3239
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3239
ccgcatccca cagaactcgg ctactcaaac cacta 35

<210> 3240
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3240
gcagcactcc acatcatcaa ccatccacaa acctc 35

<210> 3241
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3241
actacataga aggacccaaa catccctacc gcgct 35

<210> 3242
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3242
aacggcctaa cacttagaaa cctgactagc cctac 35

<210> 3243
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3243
gccccgccgg gctatccgcg acctagcctc cctct 35

<210> 3244
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3244
gagtaatgct acccacaacc cgcctaaacc atatca 36

<210> 3245
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3245
ctcactagac tcccaactaa ccttcttgct ccctaga 37

<210> 3246
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3246
gaaactcaaa cacactaaga tcctaatcat atccatga 38

<210> 3247
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3247
gtcgtagcac gctccttaca cgctttaatc ctagatga 38

<210> 3248
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3248
aacgaaacgg tctgctacca ccatccccac acacc 35

<210> 3249
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3249
ccccagagaa cccgcaccta gtcacacgac cactt 35

<210> 3250
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3250
ccgagtaccg tcattaatca actagccgac accac 35

<210> 3251
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3251
actgcccgat tgtcgctgga attccgctat ctccgga 37

<210> 3252
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3252
tgcgtcccca cgaaaccata gccaacccac cggaca 36

<210> 3253
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3253
accgcaggtc atttcccagt actaccgaca gacgca 36

<210> 3254
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3254
gcccttatgc aaaacccgcg gaccggagac catac 35

<210> 3255
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3255
acccataagc ctagaaccac cctatcacac gaagttga 38

<210> 3256
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3256
tagtccgagc acgcctataa aatgagccct aattctg 37

<210> 3257
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3257
cacaccactc gtagctgtct gtcccccgct ttagt 35

<210> 3258
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3258
cgttaccgta ctaatcccac ccaggcctat tcgtgga 37

<210> 3259
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3259
gcccggggcc cactagatcc agactcaaca ccgcca 36

<210> 3260
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3260
ccagcatatg ccctacccgc atccgatcaa ctat 34

<210> 3261
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3261
tccatatcaa accaccctaa ccagacgaat cgggc 35

<210> 3262
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3262
acctttgacc gcccatgaac tcaccggggg acggtga 37

<210> 3263
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3263
ccatcccgct ccctgtaacg gaatcactcc cctcc 35

<210> 3264
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3264
cctgcacacc gatacaatgc ctaaaaccgc atgtc 35

<210> 3265
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3265
gcctctgaat caacccgcta gcacttctag acttgga 37

<210> 3266
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3266
aacgtcaaaa actccgcgcc attaacccca accggga 37

<210> 3267
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3267
ttatctactt agcccgccct tactacagtc cgcagga 37

<210> 3268
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3268
ttcacgccga cccgaccctg ctttaaagag ttcacga 37

<210> 3269
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3269
agtcactcac cgcgcacgaa cgaaacccgc tgcac 35

<210> 3270
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3270
gtccgttctt gtcccgccaa accatagtaa tccgatga 38

<210> 3271
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3271
actctcaatc ctcaggctct gcccttccaa ccagatga 38

<210> 3272
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3272
aacccctatt ccgtggacac tcctatcatg cacgctga 38

<210> 3273
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3273
accaataacc aactatatac cttagatgac ttgaa 35

<210> 3274
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3274
gaccctcagt gcctacccca gtgcaaacaa ttctgga 37

<210> 3275
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3275
acaccgtgcc caaactatac ggaccacacc gactatga 38

<210> 3276
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3276
aacctgtctt ttcgctatcc cacatccgag agacca 36

<210> 3277
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3277
aaacccgcat ctgggaaccg gccccaactt ggccttga 38

<210> 3278
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3278
tccgccagta cgcatccaat atagtccgga ctatgga 37

<210> 3279
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3279
tgacacgaac gttatgcaat cccgactctg gacaaa 36

<210> 3280
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3280
gcctggcact atactgagtc catccgtaca tcccg 35

<210> 3281
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3281
aatcgtacag gtgtctcccc caccaccccg cgaaaa 36

<210> 3282
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3282
aacccgcacc ccacaactgc tacctactgg tgcata 36

<210> 3283
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3283
acaaccccgc accgttactg ccacctctac tcacg 35

<210> 3284
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3284
gaggcattat atcatcccgc atccaatata gaccatga 38

<210> 3285
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3285
gtcgcgttct acgccccaac cctagctccc ttccaga 37

<210> 3286
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3286
acccaagcaa gaatccgact atagctcacc tagaca 36

<210> 3287
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3287
tcttgcacct gcccgcctca gtagactacc cgctgga 37

<210> 3288
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3288
aagttactcc tccatctctg cacatggcac tagcatga 38

<210> 3289
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3289
atgcaggcaa ttaggacatt tcactctcgc ccgtga 36

<210> 3290
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3290
acccatattg atccccacca ccaccatcaa cgttctga 38

<210> 3291
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3291
aaagcccagt aactcccgga agattagtac cccccga 37

<210> 3292
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3292
acttaaggag atcctcccca ctacacatac ccaaaga 37

<210> 3293
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3293
cccaaatacc tattcccgcc agtccacgct tatgta 36

<210> 3294
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3294
aggcacctat tacgtccaac ccatcccact cggtc 35

<210> 3295
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3295
acccgccacg aaagcccgac taggtaatac gccgta 36

<210> 3296
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3296
atcgctccgt ttaaatccta aacccacagc cgcttga 37

<210> 3297
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3297
tgtcatccca cctaaacggg gaccatgcat ctaaatga 38

<210> 3298
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3298
gagcactgag tctgacccac accgcacagg ggacta 36

<210> 3299
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3299
ggtaatccgc ctccacactt gcgtgtccca tcatatga 38

<210> 3300
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3300
tgacaatgac ggacccgata ctaacccgca gtacttga 38

<210> 3301
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3301
ggagaaacgt acttgccatc caccccggac atacc 35

<210> 3302
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3302
acatttgccg cctattcggg cccgcctatc ctgtaga 37

<210> 3303
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3303
acgctcgccg ccttttccta gctgtatccc aatgatga 38

<210> 3304
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3304
cctcgccgct tctgccctac gagtacccca actaca 36

<210> 3305
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3305
acccgcccgc ttacctgaga tatcgagagc cccatga 37

<210> 3306
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3306
ttcccgtccg gtactgccct aatcataaag aaacg 35

<210> 3307
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3307
gtgggtcatc acttcccgca aatcccgatt cctacga 37

<210> 3308
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3308
acaccgcttt gctcggcgac ttatcccaaa tggta 35

<210> 3309
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3309
gcgtatcaaa aaacccaccg ctaactacaa caaccga 37

<210> 3310
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3310
ggcccgcctg ccaccgaaaa cgcgagtact tccaaga 37

<210> 3311
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3311
gccccagcgc ccacctcatt ccaaatccac gcaat 35

<210> 3312
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3312
ctcaggctgc aactacccag gttacccatc acttcga 37

<210> 3313
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3313
acgccgtttc tgtgctacca acccccctgtg ctact 35

<210> 3314
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3314
aaccaccaac tctacgcgag ttatcctgcc cgacga 36

<210> 3315
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3315
ccctaccacc cgaaaacttg aaatcacccg cgtata 36

<210> 3316
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3316
gacactgcac tctgtacccc aacctatgcc tctac 35

<210> 3317
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3317
ctactataat ccgctccacc aaaaaatagc ctcagtga 38

<210> 3318
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3318
aggtgtcaac tgccactaca tccacctgtt tccgatga 38

<210> 3319
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3319
tgtagtattg ccttccctct tccgctataa aatgatga 38

<210> 3320
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3320
gagaacgccc attcgataac caaaccaact tataaga 37

<210> 3321
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3321
aggtcatttc tactccgatt gccagtcacc gatcgtga 38

<210> 3322
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3322
acttccttgg cctggattct acccaataag aactca 36

<210> 3323
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3323
cttacaaatc cttgcctacg acccgaccac tggcc 35

<210> 3324
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3324
gtgcgcctac tgctcctacc tctgaccaca tgtgtga 37

<210> 3325
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3325
cccctgccta aaccggttag tgccttgccg ctccc 35

<210> 3326
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3326
actgagtgca tcccacgtac tttacaaatc cgactga 37

<210> 3327
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3327
aatccgatct taccgctatc tgataccgag aatta 35

<210> 3328
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3328
cccaccatta cgacctatac cccatgttac acgtatga 38

<210> 3329
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3329
caccgcctta gtcccgtaaa tactatgctc ctctg 35

<210> 3330
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3330
acagtataca tcacacccct cattttcgga gtgccga 37

<210> 3331
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3331
gcccagtacc ggtccacccc gcttccgtac taccctga 38

<210> 3332
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3332
gaacgtcacc gtatcccccg catgatcagg gcccca 36

<210> 3333
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3333
gcgcctacga ttgacaccgc cggaacatac ccacaga 37

<210> 3334
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3334
tattgcccca agatcgaccc aacctcaacg cccaatta 38

<210> 3335
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3335
tccactaccg gcgccgacgg gccgcatgta atttctga 38

<210> 3336
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3336
tcaccaaacg tcatccttca cccgagactg tagggga 37

<210> 3337
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3337
accgcctcta tcggaacttt gcttagatac tacgc 35

<210> 3338
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3338
ggacaaaaaa caccttagcc ccaaatcact accacta 37

<210> 3339
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3339
cgcgagccac acctccgcac atttaccctc gggatga 37

<210> 3340
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3340
ttgtggcttc acccatcaat ctcctctggg tctactga 38

<210> 3341
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3341
ctctgctgac tatacccatc ataacacgcg agccta 36

<210> 3342
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3342
taacagcccg acacacccga actactatgg acactga 37

<210> 3343
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3343
acaccgcatc tgccctcggt caccctacac atcattga 38

<210> 3344
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3344
actgcctaca ctacttcgtg cgaacgatag ctctt 35

<210> 3345
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3345
aattccgctc tcctgaacgc ttggagtccc attca 35

<210> 3346
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3346
tgctggtcat cacaaccgac gataacgctt aatat 35

<210> 3347
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3347
gtgacggaac ctactactgc actgaaccac cgccc 35

<210> 3348
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3348
gcccgcgatt cacgctccac ttaatccgat aacct 35

<210> 3349
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3349
gatatgatcc cacttcctag acacctatat tacctga 37

<210> 3350
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3350
cttatctgct gtcccttcg cttggatcac gattaa 36

<210> 3351
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3351
aatcaactct catgccgccc atcagtccac gttatga 37

<210> 3352
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3352
cgcttactcg tactcatcat cacgtgacat tcacta 36

<210> 3353
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3353
tccctatccc cccgccagta aaacccacat gacgga 36

<210> 3354
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3354
actccatatt cacgataacc cacagatccg ttaaaa 36

<210> 3355
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3355
tccatcccga ctaattttta gacctttgct acacaga 37

<210> 3356
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3356
gcgttcgttc tatactttct aacgttcttt acacaga 37

<210> 3357
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3357
taccggataa caacgccaat tttcattaat ttcaaga 37

<210> 3358
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3358
ggactacacg tgagcgctta ttctttgact ttacaga 37

<210> 3359
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3359
atcacgggag tgatagtctg ggccctgaaa ctact 35

<210> 3360
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3360
aaatcccatc gagcccttaa cccagaccaa ctcaa 35

<210> 3361
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3361
atcatcccgg cttttgagta aaccagcttt tcacc 35

<210> 3362
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3362
gagtacttga ctctggtata ttttttctcc ccatt 35

<210> 3363
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3363
actttacacc cgcgaacaat attacctccc cgaac 35

<210> 3364
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3364
aacctggcca ttacactttc cttttgagtt ctctaga 37

<210> 3365
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3365
gagtaggcct ctgcactccc ttctcgagat ccgtaa 36

<210> 3366
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3366
tcttggagga ctttttttg acggtttttt gtggt 35

<210> 3367
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3367
acgcacatct cattccgaat accttacgaa taacaga 37

<210> 3368
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3368
gacctataca tatccgcgca cataaactcg aactaga 37

<210> 3369
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3369
ggaccgtttt tgttttattg atgccgattg accgc 35

<210> 3370
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3370
atcttttgcc tgatcgcttc gaaattcgtt ccgcctga 38

<210> 3371
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3371
ttttgcatat acccttggtt cttgttccat ccgagga 37

<210> 3372
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3372
attttgcaa catttgaaat ttgggcattg ggaac 35

<210> 3373
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3373
gagcacttaa tttgtttgta tactgatggt tttggt 36

<210> 3374
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3374
cgtagtaccg gtctacttca ttgcccactt gtatttga 38

<210> 3375
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3375
atgtggtggc ctgaattgag tccttttttgt tttgctga 38

<210> 3376
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3376
acttaccctg gtgcactgaa agtgaacatt tgtttga 37

<210> 3377
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3377
ttgcaactat aggttgcatt ggtttttcga tacttga 37

<210> 3378
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3378
gacgctacta agtgtcgtgg gatttttttgc tctct 35

<210> 3379
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3379
gcctggatta ctttgtgaga gtttttgggg tttgc 35

<210> 3380
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3380
ggtatagcgg tcttgcaaac ttgcaaaatc ctctt 35

<210> 3381
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3381
gagtagttcc ggtcgtttac gggcttttgt ttatttga 38

<210> 3382
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3382
tgggactttg ccgcgaagct cttttctgtt atctatga 38

<210> 3383
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3383
gattcgtagc ctttaagttt tggattgttg cctct 35

<210> 3384
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3384
ggggccattt agtatgttgt ttctttttac gatgc 35

<210> 3385
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3385
aaatcgaaat cgggcctttt gaaacattgc cttagtga 38

<210> 3386
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3386
ggggctaatc gttcgaattt cgctgtgcca tttcttga 38

<210> 3387
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3387
tttcggatat taactttgcg acctcggggt ctatcga 37

<210> 3388
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3388
tgcattctct cggttagttc tgtctgtttt tgttttga 38

<210> 3389
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3389
ttcactctac ctctctcttg ctctgggctt atagtga 37

<210> 3390
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3390
ggtttcgaat tgttagttcc tttagtatga tgccttga 38

<210> 3391
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3391
gaattgcctt agttggaatt gatgttattt gtttttga 38

<210> 3392
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3392
atttcctgcg tgggttcgcc tcgaattgcc catcttga 38

<210> 3393
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3393
gcggagaagc tttgtatgtt ttccaccttg gttattga 38

<210> 3394
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3394
gaaccagctg ttgattttac cgccttgccc ttttc 35

<210> 3395
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3395
attttgcccg tgattcttat tttgtcgctg attgaga 37

<210> 3396
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3396
tgcccccggt gtttgcttta cctgtaagtt ttcgtga 37

<210> 3397
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3397
gctcaccgcg aatattttgc ttgggatttc tgggga 36

<210> 3398
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3398
agccttaatg aatgaaccat caacgaattg ctctt 35

<210> 3399
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3399
gtgacgattg cgaaattgac ttctcctctt ctgtatga 38

<210> 3400
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3400
gagttgaatg ttacccctct gcacaatatt ttagttga 38

<210> 3401
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3401
gcgcaccttg gggtttagtc atattacttg gttcttga 38

<210> 3402
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3402
ttttggccta actccgggtt tcgctttcct gcctgtga 38

<210> 3403
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3403
gagtgaggtt tgatttgcct attttcagag gggttc 36

<210> 3404
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3404
gacggtggtc ttgggggggg ttttgttttc cagtatga 38

<210> 3405
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3405
ttcaggtagg agcatttggt ggtcttcgct ttcttga 37

<210> 3406
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3406
tcgggtaggt tatccctgta tcacctttt tttgc 35

<210> 3407
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3407
tcactcatgc acttttgagc tttgttgctg aattttga 38

<210> 3408
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3408
acacgcgctt tggttattgt gcttttgctt caccgtga 38

<210> 3409
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3409
ttatatctgg ttgacttttt tttgcgggtt gcacc 35

<210> 3410
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3410
ggttgcttat gggtcgtacc atttgcatga ttctttga 38

<210> 3411
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3411
tgcccgtcct tggttttcac gtgatttgat tgtgcga 37

<210> 3412
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3412
tgcttctgta agcggagcta ttaataggta tctttga 37

<210> 3413
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3413
ttttccatgg ttgtttttgc ctcatcctct cctagtga 38

<210> 3414
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3414
agcaccccga tcgttagggg ttctgttaaa ttctttga 38

<210> 3415
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3415
atttcttgcc gttgccctgt ttttttgttt atgaaa 36

<210> 3416
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3416
ttatgcaatt ttcttccgct tggaccataa caggttga 38

<210> 3417
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3417
tctttggtcc gtgtaaataa ttttgcctgc ttttttga 38

<210> 3418
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3418
aacgcgtact gagcaataaa gcttttgccg tgttt 35

<210> 3419
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3419
agtgagtcga cgacgacgtt tctcgatatc ccccttga 38

<210> 3420
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3420
tttgttccat tttttatgca ttgttttcag tacgtga 37

<210> 3421
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3421
ttttagctct catttggttg aggattattc tgctttga 38

<210> 3422
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3422
ttccttaggg ggtttgacct catggtcctt cgtgtga 37

<210> 3423
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3423
ttacaatgat tttcgaaccg ggggttcgag ctatttga 38

<210> 3424
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3424
gttcgcttgg ccgtggactt cctatattaa atttg 35

<210> 3425
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3425
gccctttgc tttttgaggg atttaacatg cttgg 35

<210> 3426
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3426
ggaaatagaa ttgtaggatt ctctagtgct tttct 35

<210> 3427
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3427
gaccgtgttc ggaccggatg tccgacttct tcttga 36

<210> 3428
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3428
gagtaatcgt aaattttgcc ccgctggctc tggcttga 38

<210> 3429
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3429
ttactccttg acgattgtga gttctcttta gaagc 35

<210> 3430
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3430
gcacgggaag gtactgctcg tcatattaac ttggttga 38

<210> 3431
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3431
tccgtagatc cgccccaagc atgggttttt agtttga 37

<210> 3432
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3432
ttctatgcaa agctccaacg gtggagtggt tttcta 36

<210> 3433
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3433
tctttatcta gcttggtggg ctatgcaagc ccatttga 38

<210> 3434
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3434
atcccgtttt ttacctttgt gagcattatg ccgcttga 38

<210> 3435
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3435
gttttcgcga gttatttccg taatctgctc ttagtga 37

<210> 3436
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3436
aaattaaacc tttgcgcgtt ttggtgccct ttgaa 35

<210> 3437
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3437
ttttgccgtt ttccatttat ccgtgtaaat gattca 36

<210> 3438
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3438
gtggtaccat ttgactccaa cggtgattat tgctta 36

<210> 3439
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3439
gcggtcgttt tgacttgtct tcggtctcat tggattga 38

<210> 3440
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3440
gagtggaacg ttagctttac tttggtggga tcttga 36

<210> 3441
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3441
acggttcgca ctgttcaacg tggtttctcc ttgggga 37

<210> 3442
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3442
attgctgtgc ctgactattt gaacctttga ccgta 35

<210> 3443
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3443
attgttttta tttttgccat agggtgtgcc ttagta 36

<210> 3444
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3444
acgtcaacca tatgattagc tttgttgccc ttggaga 37

<210> 3445
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3445
atttcaaatt gctttgctgg acggtgctcc ttacttga 38

<210> 3446
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3446
tttgcatgaa gccattgacc ggattggatt cttta 35

<210> 3447
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3447
tcctggagct tctacgtaaa tttgatgcct tttgatga 38

<210> 3448
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3448
acagaggtac gttgattctc ctcccttgtt agcccta 37

<210> 3449
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3449
attgcatggg gattctgcct atgttggttc tgttttga 38

<210> 3450
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3450
gccacgtatt catcacatgg ggggatgtcc ttgtt 35

<210> 3451
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3451
aatgcatcga tgagccattg acttgtttgg tttgta 36

<210> 3452
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3452
attttgatca catgtgttgt gtaccatttg ctatt 35

<210> 3453
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3453
tgtttgttgc ccgctcggcg cagttaccct ttgct 35

<210> 3454
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3454
tgccacccgt gtttggggcc ctagtgtttt tttcgtga 38

<210> 3455
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3455
cgtcagtaat tgcagcatga caatttcctg cttgttga 38

<210> 3456
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3456
tgttccctga ttttgccact ggacctagta ggatttga 38

<210> 3457
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3457
catgccgaaa cgatggtttg gatcgtgctc ctctg 35

<210> 3458
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3458
ttcgctatcc gggtgatcgc tgtggaggta tttcttga 38

<210> 3459
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3459
tctaattttt tgcgtggttc tggcctatca aagtttga 38

<210> 3460
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3460
gcttccttct gggtgacggg tgatccccgt gcttgtga 38

<210> 3461
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3461
gagtacactc aattttagtt tggtgcatgc cctgca 36

<210> 3462
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3462
ttgctctgtt cttaccgccc agtgaacgcg tgttcga 37

<210> 3463
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3463
acgtgcacaa tgttggattg tttttgccct ttgtatga 38

<210> 3464
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3464
tatttgcccg attgtgaagg tgtagccctt gatctga 37

<210> 3465
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3465
tcgacttgcc agggactttt gcatccgcct tgatatga 38

<210> 3466
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3466
gcagggccgc ctgttaaggt tcctaatttg ccgtt 35

<210> 3467
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3467
ccgttggtgg gacacccctg atcttgcctc ttattga 37

<210> 3468
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3468
acccgatagc tttgaggttt gtttttctgt gcctt 35

<210> 3469
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3469
tctagcgttt gcttaccgca cactttctgt gaaaaa 36

<210> 3470
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3470
gagctttaga ggcgcttgag ggcgactttc ctgtttga 38

<210> 3471
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3471
aatagaatga tttgtccggt cccgtatttc ttgct 35

<210> 3472
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3472
ttcgtaacgt tgaccgtgtt gtttcgttta tgcctga 37

<210> 3473
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3473
agacccgtga ggatacattt ttgtttaatt tggcttga 38

<210> 3474
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3474
ttaacgccca agctcgggga ccgtgttctg tttttga 37

<210> 3475
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3475
gcatggtttt gttcgaatat ttccgaatca cgatttga 38

<210> 3476
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3476
ttgcaaatgg aacctatgcc ctgttttaat gacttga 37

<210> 3477
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3477
attttatcgg ttttgtacct gatttgcttt gattgga 37

<210> 3478
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3478
agtggacgcg ttacctcact tttgcttgaa tttgc 35

<210> 3479
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3479
atgtttgttg ttgttcctcc ccttatgact taatc 35

<210> 3480
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3480
tttgcctgtg aggtaaatct catagtatcc gtttttga 38

<210> 3481
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3481
ttttgcttcg tgtggtggtt aactttcgcg cttac 35

<210> 3482
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3482
tgatcaatct ggtccctctt ttttggagtc ttcac 35

<210> 3483
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3483
ttagcattcg ccttccaaac ccacgggttt gccttga 37

<210> 3484
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3484
acacggttgt ccctggatta gctctttgat ttggtga 37

<210> 3485
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3485
ttcactcgag gtttcggttt tataaatttc tgtacga 37

<210> 3486
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3486
cccaacttgg tttctgttta tgatctattc ggtgatga 38

<210> 3487
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3487
attgcccttg tttggtgtgt gtatttttct catgta 36

<210> 3488
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3488
actttctggg tttacttctg gtggtttcca tgtatga 37

<210> 3489
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3489
ttttgaccct gcgaaccgaa tgaccgtaat cgtta 35

<210> 3490
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3490
gattcttcaa ctcgggttga cacctgttcg ccttttga 38

<210> 3491
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3491
ggtggtcctt tgttgacgga atttatttgg ttctctga 38

<210> 3492
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3492
tatttcttcc gtgttttcca tactgtgatg atggaga 37

<210> 3493
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3493
tttggctctt tagcacacgg gtttctagtt ccgtt 35

<210> 3494
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3494
ggaagtactg tactccgaat cggatcggat ttttttga 38

<210> 3495
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3495
aataggaact tgccctcgct tttaattgtt gcctt 35

<210> 3496
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3496
ttttctgact tttggggtgg tctaaatata cttggtga 38

<210> 3497
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3497
aatttgcctc gtttttccgg gtcctgaagg gatct 35

<210> 3498
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3498
tcgacttaat gtcttattcg gagccctact tgaaa 35

<210> 3499
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3499
gccgctaacg ctgaaccgac gttgttaaat ccgtgtga 38

<210> 3500
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3500
ttccttgatg gggttcgcct ttttctgcca actattga 38

<210> 3501
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3501
gcaccgaaga gtacgtttcg accttggatt attgc 35

<210> 3502
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3502
tactgtccta ttccccgtag ttttttgtac ctgga 35

<210> 3503
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3503
ttgcgaactt ggttagcccg cctagagggt tgatttga 38

<210> 3504
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3504
gaatgtcacc ttgtggtagt gggtttctgg ttttga 36

<210> 3505
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3505
tgaaccactt agatccgttc cttaaccctg cttatga 37

<210> 3506
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3506
ttaatcccct ttcttctcgc ttatgaagaa actta 35

<210> 3507
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3507
gagtgcgggt tgatcacttg gtggggtatc ttcttga 37

<210> 3508
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3508
tcaacacgct gaccatatgt tacttgtgcc ctgct 35

<210> 3509
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3509
aattctctca gtttcgctcg tagcgagttt ttgcca 36

<210> 3510
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3510
aagttctttg ccctttcttt gctacctgaa taatatga 38

<210> 3511
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3511
tgcagtatgg acacagtggg tattttgttg tttggtga 38

<210> 3512
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3512
ttggagctgt actcccgctt tatatctgta tctcgtga 38

<210> 3513
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3513
actttaacga aactttgccg tatttcgttt gattc 35

<210> 3514
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3514
acattgtaaa tcccaattct gatggtttgt ttgtttga 38

<210> 3515
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3515
atccggtttg atcttcggag atgatatgaa atttgtga 38

<210> 3516
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3516
acggcgtgag ccatggaaaa gtaaattttg cccga 35

<210> 3517
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3517
atccaaaaca taacctttgc cgggggtatt ctctt 35

<210> 3518
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3518
ttgctcggtc aaatattgtt cgttgtttca acccgtga 38

<210> 3519
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3519
attgcatatc cgttttctca actgtaccgc ttgggtga 38

<210> 3520
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3520
acctggaccg gtgtcgaacc ggaaatctgc cctct 35

<210> 3521
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3521
catgcagtgg gtaactaaac gcttctgtat actggtga 38

<210> 3522
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3522
tcatgaagct tggcttctga aatccacacg atttt 35

<210> 3523
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3523
gcacccggtt ttgtttaact gcttgtagga ccacc 35

<210> 3524
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3524
ttttcgccgt gaggaatacc gttgggtcct ttgtt 35

<210> 3525
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3525
tcgttaaatt gggcttgcct gtgccttttt tcccgtga 38

<210> 3526
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3526
attgggggtc accttgggtt tggatgtttt ttgtttga 38

<210> 3527
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3527
ttgacgaaat tacccggtat tgtgcttata aattgga 37

<210> 3528
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3528
tgcgattgat tccttccgtg ccgttttgaa actatga 37

<210> 3529
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3529
tctgtgccaa taactctaaa tgccgcttag tttgaa 36

<210> 3530
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3530
atggtgatcg gtggtttgtt actctgtccg tttcttga 38

<210> 3531
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3531
gcccggttac gtaccttggg gtgtttgtct cttgc 35

<210> 3532
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3532
tgcgagtcct ctgaatctga tcctaaactt cttgc 35

<210> 3533
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3533
actccggcct tcttctgtat cggtttttt gctgttga 38

<210> 3534
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3534
tcgactcttg tgccatggtt tccttatctg gtatt 35

<210> 3535
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3535
ttcgatctgt gcgaacagat tttgttttgc cctgga 36

<210> 3536
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3536
ttcgtcgttt aaacacattt gtccgctatg cctta 35

<210> 3537
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3537
atggaagggg ttccatttcc tttatttggg ttttcga 37

<210> 3538
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3538
tgcttgggtg caatgcgaac gcatattaat ttgttga 37

<210> 3539
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3539
gcattgtgta acttcggttt ttttggaatt ttcgc 35

<210> 3540
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3540
gggtccaaac ggcttggcca ttttctctat ttctga 36

<210> 3541
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3541
gataatcttg ctccgtagaa acatacttac tctggtga 38

<210> 3542
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3542
atcgagtgtc tcccgtttta tctttgattg tttgc 35

<210> 3543
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3543
tctctgtatt cgtttttctg atatttggtt gtagaa 36

<210> 3544
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3544
tctcctgggt cccgttcggt tatattatgc ttgtga 36

<210> 3545
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3545
accgctccct tttgttcggg cttggccaat ttccttga 38

<210> 3546
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3546
aggccaattt tatagcggtg actcactttt gccct 35

<210> 3547
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3547
gggtgaggta ccttctccgt gtgattctcc gtttttga 38

<210> 3548
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3548
gaaacgtgat ataccgact tgttgcttac gaatttga 38

<210> 3549
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3549
ttgcctgaga gatgccgctt tgttcttcgt aatcta 36

<210> 3550
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3550
ttctctgacc cggttttttt acatatgtga cgttttga 38

<210> 3551
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3551
tgcacctcgt aaaaaaactc gggtctggaa tttcgtga 38

<210> 3552
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3552
cataaactct ccttcctgca ctcttccgga ttacttga 38

<210> 3553
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3553
acccgggttt gcatactgtt cttctgtcct tatga 35

<210> 3554
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3554
tctgaaacta gcatgagcct tactgacctg atgtttga 38

<210> 3555
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3555
tattgcccgc cactttctga atgatttgat ctgct 35

<210> 3556
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3556
gacgcaccaa attatgcaat acaaattctg cccttga 37

<210> 3557
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3557
gggttacgtt attctccggt accttgggtt taaaa 35

<210> 3558
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3558
gacaactccc gtgaggagtt tctgatttta tgttt 35

<210> 3559
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3559
ttactagccg gtggatctgc ccacgttcgt ttttttga 38

<210> 3560
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3560
gcacaaattg cccggtataa ttgttatacg ctatgtga 38

<210> 3561
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3561
taccgaggtc gagggatttt tgtttgtcct tatgtga 37

<210> 3562
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3562
aaaattgccg gcgccgtagc cttgttttca tcttta 36

<210> 3563
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3563
tccgcgggcc gtaatttggt tagattttct gtttgtga 38

<210> 3564
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3564
ttgcctcgct ttgaccagtg ttcctgtttc tccttga 37

<210> 3565
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3565
gggcggaccg cctggggttt tgaaccttac aatcttga 38

<210> 3566
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3566
gagtacgtga ccggtgaaaa atattgggat gtgtgtga 38

<210> 3567
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3567
tttgcgcgtg agcgttataa aattttccga ctaccga 37

<210> 3568
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3568
ggtcgccttg ccacaccgat aatcgttgaa ttttgtga 38

<210> 3569
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3569
tgccgctgga cactatatat ggtttctgaa atgcc 35

<210> 3570
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3570
gagaacgcac aagttgccac gaggtttatt tgccttga 38

<210> 3571
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3571
tgccctgtaa aacctcggac ggtttcctgt ttgttga 37

<210> 3572
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3572
tccatctaag attatctcgt atcacactat taaat 35

<210> 3573
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3573
tatcaaccca aactaaccgc tcaataatga attaaga 37

<210> 3574
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3574
agtcagtact aaactctcta aacatctgga tccaaa 36

<210> 3575
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3575
ttcgcaagtc caggatcgcg actaaatacc ttacaa 36

<210> 3576
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3576
tgcctcatcc cgaatagttt tgtcaaccca tccaca 36

<210> 3577
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3577
acactctacc aaatagccac tcgcaaacat ataaaga 37

<210> 3578
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3578
tcgtcactca ctcaatatcc ttggtaccta ctcaaa 36

<210> 3579
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3579
tgcaggcccc tgtgaacacc tttattgaat agcaca 36

<210> 3580
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3580
gcactgtgac ctaccacttt gacctctgtt taaac 35

<210> 3581
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3581
acctcgacca tctacaaact ttgaagccac accaaa 36

<210> 3582
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3582
taagcctgac tccatcgtct attatttttt ttaagc 36

<210> 3583
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3583
atcgccacga ccaaccacca cggatcccac ataaa 35

<210> 3584
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3584
tactcgacga aatagctaac aacggacccg ataaaa 36

<210> 3585
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3585
aaatcatccg tagtttatcc tttgtctcca tccaaga 37

<210> 3586
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3586
gggaacgtaa ctcaggtttt tttcatccca acaca 35

<210> 3587
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3587
aactcggcca cccgtagacg tatatccaat tcacttga 38

<210> 3588
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3588
ctgcagcttg atcttcgtca caacacatct attacga 37

<210> 3589
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3589
ccttccaatc tgcatcgcta acgcatacac cattatga 38

<210> 3590
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3590
gatctattca ccttagccat aattatagac cccaatga 38

<210> 3591
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3591
ccgtactgtt tacacatatc tatccacgta ttaca 35

<210> 3592
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3592
acgtctgcca aatctactcg cattcgatac acacc 35

<210> 3593
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3593
ctgtcgacat tccgctataa aattaccaac ccaat 35

<210> 3594
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3594
gagtacgcaa ttattatccg aagtcaacta tagcaa 36

<210> 3595
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3595
gagtaactag tcatcaattc tatcttcgca cactaa 36

<210> 3596
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3596
atgcgctact gcctagctcc cgctcgccca tataa 35

<210> 3597
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3597
acaatatcga tctccaaaac tatctcgtgt atctca 36

<210> 3598
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3598
acgtacccga ccccggcgtt aagccgtctt tgtactga 38

<210> 3599
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3599
gtcactcatt atatggttct acactacgat catca 35

<210> 3600
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3600
gactccgtcg ttcatttcac taacacgtat gccaga 36

<210> 3601
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3601
tagtacccac ttattactat cggagttcaa acataa 36

<210> 3602
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3602
tcgagcgcac agattcgttt tccaccgcga acttca 36

<210> 3603
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3603
gccacgtttc ctactttcgt tcgttcgcca cgtagga 37

<210> 3604
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3604
tcactgaccc acctctcgat ccccaattaa atatt 35

<210> 3605
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3605
cgcttcccga atcgagagat caagcgatca gccgta 36

<210> 3606
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3606
tcttacgtac gcttcgtaga caattattca tttctga 37

<210> 3607
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3607
tgctacacga aacctgtact gctagctatc tatcatga 38

<210> 3608
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3608
ctagctcttc agtcgcactc ctagtctcac cattca 36

<210> 3609
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3609
tgaccattag ttaactctat cttttgaaat aatct 35

<210> 3610
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3610
agcctcatag gcattccggt gtcacgcaaa cacata 36

<210> 3611
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3611
tgccgtcccg ggactcaatc cacgttttta ttttctca 38

<210> 3612
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3612
tgccgctaag gcagttacta cagggtctta ttacatga 38

<210> 3613
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3613
gacccttgt taaatgtatc cttgttaatt acgat 35

<210> 3614
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3614
caaccgtcct ctaatttcac gcctgtcggt cattca 36

<210> 3615
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3615
ctacaccact attcaaaacg aatatctcat catactga 38

<210> 3616
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3616
ttgtacctgc actgcctcaa gttattacat ctcata 36

<210> 3617
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3617
taggcgagag cctttgacag gttaccaggt ggtggtga 38

<210> 3618
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3618
agtgactgcg caaatcacta caacttccat ccaaaga 37

<210> 3619
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3619
gagtgactgc tttgacctat acacacaacc aacttaa 37

<210> 3620
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3620
atttaaccct aaatatttca acacacaaac accaca 36

<210> 3621
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3621
ctctggactg caacacagtg ccgttccgac cttcttga 38

<210> 3622
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3622
tacttagtct tcttcggatc gcgaccgtgg tcggta 36

<210> 3623
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3623
atatcgtacc tattttctaa tcttacgctg ttcac 35

<210> 3624
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3624
cccccgcacc acattattcc gatcatatta tccattga 38

<210> 3625
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3625
acttaatatt ccattctcaa ttacactatt cggtt 35

<210> 3626
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3626
ttcgtcactg ctacttttat attaagccca tagaatga 38

<210> 3627
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3627
cctcgcatcc gacttttaca atcacataat cacgta 36

<210> 3628
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3628
ttgaaagccc ctcaatggcc acctattttg cattca 36

<210> 3629
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3629
gagtactgtc tctattccac aagcataact acttca 36

<210> 3630
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3630
aggcgactgc cgcagtgccg ctgtccgttt catcatga 38

<210> 3631
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3631
aatgccaagt cttatcgcta gcaataatat tgaactga 38

<210> 3632
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3632
atcccctgtc tgcatgcagg cctatgaccg cactca 36

<210> 3633
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3633
gggggcggga gctacgcaag catgcctttg acattga 37

<210> 3634
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3634
taatttctct gcatctatct caaagcacca ctatctta 38

<210> 3635
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3635
acccatctac cgcattatta acgttataac cctaaa 36

<210> 3636
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3636
cctgcctatc gccccttcat attctcagat cagcaga 37

<210> 3637
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3637
ttctcgtctc ggggctaata cacatacaat tataca 36

<210> 3638
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3638
gctcattcta ttttagcata tcatccatct aatcg 35

<210> 3639
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3639
aatccgtcat gtattaaaca cttctatcac attcg 35

<210> 3640
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3640
actcttctga atcgaaacca atctcacttt cagggga 37

<210> 3641
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3641
ctaccatcac atcctatcat cgcatttagt cttgta 36

<210> 3642
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3642
ctaatcatct cttcaaattc aacggcgaac aaaac 35

<210> 3643
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3643
taagcacttc cgttttaatc tttacgacgc acccatga 38

<210> 3644
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3644
tactatcggc acttctacgg aggataccct cgatta 36

<210> 3645
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3645
tgagctgccg ctatggtgac cttctatccc attat 35

<210> 3646
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3646
ccttacagac ggtcaccgtc ttgcgggtta ttacaa 36

<210> 3647
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3647
ccagtcagtc tgttactgtc ttgttcccat ccctca 36

<210> 3648
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3648
cgtttccaca cgccgattca aatccctcat taacca 36

<210> 3649
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3649
acctaacatc ctatcaccgc ttatgtattg catca 35

<210> 3650
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3650
gcagtaggtt gtgatgctag ctaattatac tatac 35

<210> 3651
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3651
cccgccgcac gtccaatatt caaccatgac cttactga 38

<210> 3652
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3652
atcgccatca tcgctttatc tggtcgtggg aaccc 35

<210> 3653
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3653
tgtcatccgt tatttcattc accttctact atgtaa 36

<210> 3654
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3654
tactattatc atccacgcca tcagccgggt cttcgga 37

<210> 3655
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3655
ccttacatac atactccgca tacgtgccta tttaatga 38

<210> 3656
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3656
acaactatca aataaacctg cgccctctat taatgga 37

<210> 3657
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3657
gcaccatcgc cggactgctc atatccccgt tgtaa 35

<210> 3658
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3658
ctcgcagtaa tggtgcccac atgttacgtc tctttga 37

<210> 3659
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3659
ttagtactgt cattcgagga ttcgaatcta gcttta 36

<210> 3660
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3660
tgtaaacgta cgatattact tccgatccat acatca 36

<210> 3661
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3661
attcaagcaa tctagtaatt attctttcac tgccgga 37

<210> 3662
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3662
taccatctct attccacgtc cacgacacgc tgccatga 38

<210> 3663
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3663
tgttgacaat gccgctgtcc cgtctaccct tcgata 36

<210> 3664
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3664
catgtcgccg tcgcgcacag cctacatccc actta 35

<210> 3665
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3665
gagtacttct cataaacgaa atatcaaatt cgaat 35

<210> 3666
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3666
tgactgcgaa actaacgtat ccccatatat acaattga 38

<210> 3667
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3667
ttcagcgcgt tcttcacagc ggggtagcac tcatca 36

<210> 3668
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3668
tcagtacatc attaatcacc atatcacgca ttatc 35

<210> 3669
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3669
agtctactgc atcatctccc ttactccaag tagtaa 36

<210> 3670
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3670
gctcatactc agcactgcga attgccagtc cttcac 36

<210> 3671
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3671
gctgcatctc gtcactccag gtggttggta cattgtga 38

<210> 3672
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3672
gttagctaga acgaacgcac gtcttctatc tttta 35

<210> 3673
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3673
ttcgttattg ccttagttac actctattat aggtta 36

<210> 3674
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3674
tatcggacac ttactacatt accgtcgctt ataccga 37

<210> 3675
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3675
atcaacgtac tcgaacgcgt ccacacccca ccatctga 38

<210> 3676
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3676
gagctactct gaatctcatt gatatctaca ataaatga 38

<210> 3677
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3677
taataggacg gcgggctctt gccgatattt cctgta 36

<210> 3678
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3678
catcttatac agctcagcct ctattcccat acgtc 35

<210> 3679
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3679
ctgccggatc ttgcacgcac tgctattata tttcca 36

<210> 3680
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3680
actgcctaag gctatcgcat caatatgaac tcaac 35

<210> 3681
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3681
ctactgcacg acgcgagtcc ttatcaatat ctcacga 37

<210> 3682
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3682
gagtacttac gtaacgagtt attaatgtcc aaaat 35

<210> 3683
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3683
aaggactcaa tgccactata cgacttattc caatca 36

<210> 3684
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3684
cttttctagc tagcatttac acttttatac acaac 35

<210> 3685
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3685
tcgggttctc cctctcgcca cctaggttta aaacgga 37

<210> 3686
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3686
cgcatgcact gaacatatac acatctcgtc gctcctga 38

<210> 3687
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3687
ttacctctct acaccattta ccaacggcta cgccga 36

<210> 3688
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3688
gacgtcttct ttcacagcat gccgtaacag ttttctga 38

<210> 3689
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3689
ctacagacta actagttccc gaattctgcc acaac 35

<210> 3690
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3690
aatccacgtg cccaacataa agatctaact cacaaga 37

<210> 3691
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3691
tacgacgcct ctgttaatcc gtttttatat accaca 36

<210> 3692
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3692
ggcgggaatc acgccaaacc tctgtcacct atcgctga 38

<210> 3693
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3693
cttctgccac ttcatctcgt caaggttcat ttcgttga 38

<210> 3694
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3694
cgcacaacgt accggtccag ggaacacatt ttcgca 36

<210> 3695
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3695
ctctcaccag cacgtcccac acattttcac aaacctga 38

<210> 3696
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3696
tttctcgcgc ggcaggcatc cgttctaata taccatga 38

<210> 3697
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3697
accacaactt atcgcatctc attgattgga ctaata 36

<210> 3698
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3698
tcccctcagc catcgtggaa caactgcggt gtttc 35

<210> 3699
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3699
atacgtctgc gattatttag tggtctctca cacata 36

<210> 3700
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3700
cacgagtcgc cacacctcag ctgatgctca cctaca 36

<210> 3701
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3701
agctacgtac cgaatttcgt tcataatagg taatga 36

<210> 3702
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3702
acacattcgg ccatgtcatc ctgcacttca cgtga 35

<210> 3703
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3703
taaccatcga tcccaacaca tgtcagcacg gcacctga 38

<210> 3704
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3704
tacctagtcc ggccaccagt ctcccacaca ttacg 35

<210> 3705
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3705
atgtctgcgg accatctatc ggatcgccga acctt 35

<210> 3706
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3706
cgcacttacc atgcgtatcc tttatcggtc tgaacga 37

<210> 3707
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3707
gagtcccggt cccccgggtc catcatctac tccatga 37

<210> 3708
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3708
atctgctgtt gtctttccca ttagttatcc caatcga 37

<210> 3709
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3709
agccacacct aatagctcac atccggttac cgtcctga 38

<210> 3710
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3710
ccggctcaga gccacgctat atcttcacta ctaactga 38

<210> 3711
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3711
gcgaagtttc acttccagtt tctgacgccc ttcatga 37

<210> 3712
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3712
tcctaggctg ctcttcgccg cacaccggac ttcaca 36

<210> 3713
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3713
catgcaccgg tctcacggtc gcataataac tatcatta 38

<210> 3714
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3714
tgtcaatatc tgaatcggtg gatattggcg gaggttga 38

<210> 3715
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3715
gggtaccgtg cagactcaca gaacctaata tatagga 37

<210> 3716
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3716
tttaccctgc atcgatcgcg agacccacca atcatga 37

<210> 3717
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3717
tggcggccct tctaccgccc gtcaccacct ctatgga 37

<210> 3718
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3718
tacaccaata accatacaac tacatctaat acgtta 36

<210> 3719
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3719
gagtactcgt gatcttttac acacatctgc ctaaaga 37

<210> 3720
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3720
agcaccggaa tgcccgggcc cttgtacgcc tagtga 36

<210> 3721
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3721
ttaagtcatc tgcactagta cgccctatat taaaa 35

<210> 3722
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3722
agtctcgcag aaaaaatcta tttcccattt gactgga 37

<210> 3723
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3723
aggaactgcc taccatcgat ttttaatacc acacga 36

<210> 3724
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3724
tttcttctta tgagggggag gtgcggttcg gaataa 36

<210> 3725
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3725
gccaggattc ggccatttta cctatctcga tttaaa 36

<210> 3726
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3726
cgtgccgaac aactcacacg ctacaccttt atttttaa 38

<210> 3727
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3727
agactatcat aactccttca cgaaccttcc acataa 36

<210> 3728
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3728
tcaccgaccc tcagtacttc ccctgcgcat gactaga 37

<210> 3729
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3729
tctcactact acccactaaa atcggttgat actac 35

<210> 3730
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3730
acgcacgggt atgcatcatt gatggtttcc ggtac 35

<210> 3731
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3731
acgttcctgt acgccgcggg cggtattttc acaattga 38

<210> 3732
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3732
tcgccctatc aactccctgt attttcatcc ctcaatca 38

<210> 3733
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3733
cgcctacgat cacttccccg cgcgactgtc tcatc 35

<210> 3734
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3734
cacaacatat ctactctata ctagctcaat tcaccga 37

<210> 3735
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3735
cattgcgacc ttatcccacc acgaattgcc actaaa 36

<210> 3736
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3736
ttactatcca gtatcacacc gaccttccaa tccga 35

<210> 3737
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3737
agaacttgat cgtttctggg gtttgggaat catgtga 37

<210> 3738
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3738
gggcttgcaa acaacttcac gtatgatccg cacct 35

<210> 3739
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3739
aatcccgtga gggtcgtttc aagtttcaat aatcc 35

<210> 3740
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3740
ggcttacaca gccacctatc tttggccaat ttataga 37

<210> 3741
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3741
agcgccttag gcttccttct gcggctaacc atcata 36

<210> 3742
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3742
gaattagctc acttactatt tcattttgac cataa 35

<210> 3743
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3743
ggccttcccc ttcatccgcc tcaacgctgg ccacc 35

<210> 3744
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3744
aggcaggcga ctacggatca acatttctat acccta 36

<210> 3745
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3745
gttagtgcgt attgtaggct gacgaacgct accgta 36

<210> 3746
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3746
gcggcccggc gatcctctag tccaaccggt ctacta 36

<210> 3747
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3747
ggggggcac gtacgacatt aggtccttaa atcactca 38

<210> 3748
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3748
taaccgcgta cggactcaag ccacccctat taaaca 36

<210> 3749
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3749
gctatttgtt aagtccagtc gcaaatcacc actat 35

<210> 3750
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3750
ccgggcactc ggactttcac tctgtcagtt tctaca 36

<210> 3751
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3751
tcgcaatata tatctccgtg gtgtaccatc agtcaa 36

<210> 3752
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3752
ccctctactg gcttctacgt cgcacgactg ctacc 35

<210> 3753
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3753
aacctcacat gctggcgatc actagcaaac cacc 34

<210> 3754
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3754
aaactgcctt gaacaagttt cgatatcaca ttccatga 38

<210> 3755
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3755
cctcataacc gattaccaac gccatcaaaa tatag 35

<210> 3756
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3756
aacaaggcca aaatacgcat ataattctgg ttgaaa 36

<210> 3757
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3757
agtgcgaact cttattactg ctattaaaac tgtcc 35

<210> 3758
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3758
tcttgctacg tttgttcgca ccaaacttgt cctgca 36

<210> 3759
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3759
gttctcgcgc gcacatcaat tatatcacaa ataa 34

<210> 3760
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3760
atttacgttc gcttcatgga cgcggtcata ggacctga 38

<210> 3761
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3761
tcgcaccggc cctcaatacc atttacacat ccacgga 37

<210> 3762
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3762
ctcgccacga cttaattctc aattcctcac aactga 36

<210> 3763
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3763
aactgtacgg cgcaatttta ctctcaagta ccaaaa 36

<210> 3764
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3764
cactgtcaag ttttccattc cactatacat caatca 36

<210> 3765
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3765
ctttcatcat aacccgggga atttaccggc cgcctca 37

<210> 3766
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3766
ggtatttgct agatcttcaa ttacctcgcc gtatataa 38

<210> 3767
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3767
tgcctgcatt gctcgtggag gctcattgac tttca 35

<210> 3768
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3768
ctctgtcctt gctttctctc tcattccgat gactatga 38

<210> 3769
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3769
gatctccact tcaggcgcca ttctataccc attaatga 38

<210> 3770
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3770
tatcaatttt attaataccc tcattgtgct cctactga 38

<210> 3771
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3771
gatgcgagaa ccactatcca tcccacacgc cgatt 35

<210> 3772
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3772
agcactctgt caggtcgatc attatttcac aacaca 36

<210> 3773
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3773
gagttcggcc ctcgttgatc ctcgattaca aattaa 36

<210> 3774
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3774
gggggtacgt ccagcttatc atctcatacc attattga 38

<210> 3775
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3775
cgcacgtcac gagcacacca acactacttc accactga 38

<210> 3776
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3776
acatgtcgtc cacctatgtt gttaccatta tttat 35

<210> 3777
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3777
attcaatccc ctcctatggc ttgccccccg gttct 35

<210> 3778
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3778
cggtattcgt acgtctgcgc gcaacccatt ataaca 36

<210> 3779
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3779
tccatgcagc cacgctcagt tactactact agtgatga 38

<210> 3780
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3780
atcacatttc catcgatctc caaggttctc aaact 35

<210> 3781
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3781
gtgttcctca cactaatcgt tctactttct taact 35

<210> 3782
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3782
tcacactcca tagatctctc ggtagacgca gtccga 36

<210> 3783
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3783
ccattacaac actacctcta tatggtccta tccgca 36

<210> 3784
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3784
cgctatctct acgccaatta agtcgctcct ggtaca 36

<210> 3785
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3785
gcagatacca agtcattact ttattataca taacca 36

<210> 3786
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3786
attcgtactc tataaattcc aatggcgtac ccacaa 36

<210> 3787
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3787
tcagtacacc attaatcacc atatcacgca ttatc 35

<210> 3788
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3788
aaccgggtac ccacaccacc cgtaaccatc ccaattta 38

<210> 3789
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3789
aaacggacgc gcgaaactag ttggaacatt attcc 35

<210> 3790
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3790
aacaacgaaa taggaaatcc tctgctgaac caaat 35

<210> 3791
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3791
catgcccccc gagtttcgta gttattttcc taaac 35

<210> 3792
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3792
acgcaattca gaacatattc gcaactcttt caaat 35

<210> 3793
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3793
tttcgaattc tccacctgat acccgccaca ttaaa 35

<210> 3794
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3794
accctcctat taacccatca aaagttcgct accata 36

<210> 3795
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3795
atactccaca acgaacatcc aaactgttta catcaga 37

<210> 3796
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3796
gagtacctag ccgcaccggt cttaattttc ttaaaa 36

<210> 3797
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3797
aagctgcacg gatcttccat gtcccctcta ttaaa 35

<210> 3798
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3798
agttctatta tggctcctaa ctgttcttgt tttcaga 37

<210> 3799
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3799
acactcgcct agaagccacc acccatttct cataaa 36

<210> 3800
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3800
cctccaccat ccgaacatta actaattacc cgccaga 37

<210> 3801
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3801
tacttcctat agttacccac atctactaac caaac 35

<210> 3802
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3802
gagtactggc acgcgatacc ctaactgtac atcaca 36

<210> 3803
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3803
agccagcaag gctagcatca cataattctc atatc 35

<210> 3804
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3804
atgaatccgc actactaaaa caatttatcc ttataga 37

<210> 3805
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3805
tctgtgctgg ctttagaagg ttttctgttt tttccta 37

<210> 3806
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3806
acatagcact gctaggttta tctcattgtt tttaa 35

<210> 3807
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3807
ttctggcaat tggcctctca ctctctatgg ttttgtga 38

<210> 3808
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3808
ttttgattta ttctgcttgc tgtttgggct tcttt 35

<210> 3809
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3809
tatattcaca atcgattctt ttcacttcct cattttga 38

<210> 3810
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3810
tgtaggagtt gtctcttttt catattttct gtctt 35

<210> 3811
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3811
taggactggt taatcctccg aaattgtttt tctat 35

<210> 3812
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3812
ttctatcggg ggagttccta ctctggtttt tttcttga 38

<210> 3813
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3813
gcctgggtgg ttatttttt ccatatcgaa attta 35

<210> 3814
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3814
ttagttttct cactgtttcc cgttgactat tctcta 36

<210> 3815
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3815
ctagtgacca ctgcttcttt tggttatctt aatcttga 38

<210> 3816
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3816
gcctgggttg ttatttttt ccatatcgaa attta 35

<210> 3817
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3817
tgtatgattt tggtttttgt tttggtttca cgtggtga 38

<210> 3818
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3818
gcagtgttat ttctcttggt ttctttgatt tcacttga 38

<210> 3819
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3819
ttctatcggg ggagttccta ctcttgtttt tttcttga 38

<210> 3820
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3820
ggcctggttc tcgtacatct gtattttggt ttttctga 38

<210> 3821
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3821
gcacccgggg tttttatctg agttgttttt cattt 35

<210> 3822
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3822
tttatccatt gtccttggac tctcctccgt ttagaa 36

<210> 3823
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3823
ctgcgctggc caatctcatg ttttgttttt attaatga 38

<210> 3824
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3824
tgcttcttta atctgttttc tactattttg atataa 36

<210> 3825
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3825
gatcccgaat tgttttccta ttaatcgttt tcccatga 38

<210> 3826
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3826
tcccttggga ctctttggtt attaacattt ctctttga 38

<210> 3827
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3827
tctaactgct ggaccctcct gttttttatt agtcta 36

<210> 3828
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3828
gcagtacgga ggtttaagtt gtttctccta ttttttga 38

<210> 3829
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3829
atgaccgttt cttggaagtt tccgattttt tttgt 35

<210> 3830
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3830
taatgcattg ttcgcacctg gttttttaa tggtta 36

<210> 3831
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3831
gagcaataat tcctgtttct atctctacaa ccatttga 38

<210> 3832
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3832
actgacttaa tctaatttgt tttccattct catgaa 36

<210> 3833
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3833
gtgcacgggc gtttttaat ttaagtttct tgttttga 38

<210> 3834
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3834
ttatactatc caaatctcgt gtctttcctc attat 35

<210> 3835
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3835
gtcgtaattt gttctttata tctatttctt gttgta 36

<210> 3836
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3836
ccttggagtt gctggtaacc tgttgttttt tattga 36

<210> 3837
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3837
acgtgcctga taattatatg tatgtttctt gtttt 35

<210> 3838
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3838
atctttcact tagtctcgga gtttttttgtt tcctta 36

<210> 3839
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3839
attcgtttat ttttatttcc tgattggttt atttca 36

<210> 3840
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3840
ttcactggtt tttttctttc tgacctatag attgc 35

<210> 3841
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3841
ttgctaattt ctgggtttgt tacgtccttt tatata 36

<210> 3842
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3842
tacttttttg atttttgatc ttggcccacg aaataa 36

<210> 3843
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3843
tctctcttgt caatagaatt tttgtttgta tcttcga 37

<210> 3844
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3844
tgtgctcact atgtctctcc gttgtctgtt ttttta 36

<210> 3845
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3845
aatcaagcag ggttcaccat tgtcgttttt ttttta 36

<210> 3846
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3846
gatgtactgt tcgttttgtt ttgtcctttt catcatga 38

<210> 3847
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3847
cgcatactgc cgcatcgcat ttgttttatt gttgta 36

<210> 3848
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3848
ttgtggttgt tcatacctgc tatcttttt cgttt 35

<210> 3849
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3849
gagtgatgcg gaataaatcg ctttgttttg tttcctga 38

<210> 3850
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3850
ttagtaggta actttcactg ttttttttatc gtttt 35

<210> 3851
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3851
gggtctcttt ttctttaact cttttatcta aatttga 37

<210> 3852
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3852
tgacgttttt ttgactggtt tttctgaagt ggttttga 38

<210> 3853
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3853
gacgccaccg ttttttattt tagttgattt agata 35

<210> 3854
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3854
tcataaatca ctctatcttt gtttttcagt cttagtga 38

<210> 3855
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3855
ttacggatgc tttttttgt cctgagatta ctttt 35

<210> 3856
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3856
gactgtacga tggcttaatt gttttgtatt ttgtt 35

<210> 3857
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3857
ttattgtttc taccatgttt gttgcatatg ttttt 35

<210> 3858
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3858
tttttccgtt caactctact ttcgatatat tatggtga 38

<210> 3859
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3859
tagtctttcc gattgttttt gttatttata tgcctga 37

<210> 3860
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3860
gcggggcgcc ctcgatttgg ttttattttt ctcaca 36

<210> 3861
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3861
tgttactttt ttatcttgtt ctgccatctc taattga 37

<210> 3862
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3862
tatctatcac taagttttta tatctaaggt ccatta 36

<210> 3863
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3863
ggctttgttc attttcttac tctttgtttg tagtttga 38

<210> 3864
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3864
gccggtttat ttaccgagtt tttcatcttt tcttatga 38

<210> 3865
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3865
agggtactag cattttcgat tttgtgaaat tttctga 37

<210> 3866
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3866
atactgcagt agcacttgtc gtttttctcc aatattga 38

<210> 3867
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3867
ttcgttccta agaatttgcc atttgttttt tggtc 35

<210> 3868
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3868
tcactctatc cagcatgcga ggtgttattc tatttga 37

<210> 3869
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3869
agtcgataga ataatttttt ggttttctca ctcat 35

<210> 3870
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3870
tactgtcctt ttcaattcgt gtcttggttt tatta 35

<210> 3871
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3871
taaacgtggt actgcgcacg gttatttctt tctcttga 38

<210> 3872
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3872
tggggtactt gcctactttt tacctggttg ttatttga 38

<210> 3873
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3873
ttggggttaa cgttctgatc tcgttttctt attattga 38

<210> 3874
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3874
ggaatgcagc actcgaggtt ttctttgttt ttccca 36

<210> 3875
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3875
gcgcccaaat ccgttgcttt gttgttttgt acttttga 38

<210> 3876
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3876
agtagccttt ctctcgatct ttgtttgttt tttgatga 38

<210> 3877
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3877
actttcggtt gaccatcata tatctctttc ttacttga 38

<210> 3878
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3878
gtcgtaccaa tatcggatat cctggttttt ctgtc 35

<210> 3879
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3879
ttactgaacc tctccggccc tctatttttt gaagttga 38

<210> 3880
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3880
ttttccgttt aattcgatca ctttcattgt ctcca 35

<210> 3881
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3881
gcgttgacat gggattcttt gtctcttagt ctttt 35

<210> 3882
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3882
agttctatta tggctcctaa ctgttcttgt tttccga 37

<210> 3883
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3883
gctgcactct acacttatgg ttgtctatct tttactga 38

<210> 3884
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3884
tgaacgtagc agttttgaca tttagtatat ctttttga 38

<210> 3885
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3885
gcggtccgta tgggattttt atttcgttct tgtta 35

<210> 3886
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3886
ttttaggggt cctgtacatt ctatcctttt atcat 35

<210> 3887
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3887
ttgtccttcc ctgttagtct atcatctttc ctctgga 37

<210> 3888
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3888
acgacgcccc ttttgttgtt tttcttggta tcttc 35

<210> 3889
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3889
gagggccccc tatcatattg ttttttttatc ctccatga 38

<210> 3890
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3890
tgctgccgta acactgttta tttctattct tgatta 36

<210> 3891
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3891
tcgactctgg tttgtttttt tggtttatta cttcgga 37

<210> 3892
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3892
tgttcaaccg gtcccattgt tttttgtcgt aatct 35

<210> 3893
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3893
cacgccttcc ttttgggttg ttgatctcgt ttttt 35

<210> 3894
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3894
aaacgcattt ggctcaattg atcttattat ttttctga 38

<210> 3895
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3895
tttaccttgc cctggttacg aaatatttgt ctctca 36

<210> 3896
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3896
tataccatcg cttctcatac tggggtttat ttttttga 38

<210> 3897
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3897
taaatcgttc tcgtaacacc ttctatcttt tttatga 37

<210> 3898
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3898
atggatttgg ccttcactct atgttctcgt tttct 35

<210> 3899
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3899
catcgcatct cgaacgggcc ttttgtttta ttatttga 38

<210> 3900
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3900
gcctttacac atttctgttt tgataaagta attct 35

<210> 3901
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3901
gtgcgctatc taatgtcacc gtttttttcc tcaattga 38

<210> 3902
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3902
tgtcacgtac ctcgtgtttc ttttctgtta aacttga 37

<210> 3903
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3903
aatggaacaa actactttca ttttctttct ctcattga 38

<210> 3904
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3904
tttcccaaat ctagcttttt tcaataatcc atatctga 38

<210> 3905
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3905
ttccatctca ccaatgttgg tgactgttgc cattttga 38

<210> 3906
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3906
actcgtgttt ttttcgtttt gtgcccttcg attctta 37

<210> 3907
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3907
agtgcccgga ttagctggct acagtttttt tctct 35

<210> 3908
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3908
aaactccttt tttgtatgtc gccatttctt tctta 35

<210> 3909
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3909
tcttggcatg tcctcgtttt ttgatatctg ctttta 36

<210> 3910
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3910
tgtcgtagta gtgcgcattt gtttaactct ctatca 36

<210> 3911
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3911
tccactgttt acggttttaa attgtttcct tgccctga 38

<210> 3912
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3912
agtccatgcg agctgcatgt actctcggtt ttttatga 38

<210> 3913
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3913
gcgtgtttag ttaattcgtt ttttaccaat ggggtga 37

<210> 3914
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3914
agtttccctt ccgtttcttt ctcagtttaa ctctttga 38

<210> 3915
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3915
ggtgctccgg gcttgtaact ttatagtttt ttccttga 38

<210> 3916
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3916
gactctttga tctccttgaa cgtttttgtt atttc 35

<210> 3917
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3917
tcgtttccaa tgatttctcg atttacactc catcttga 38

<210> 3918
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3918
tttctgtcaa cactctcgaa catatattta ctccttga 38

<210> 3919
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3919
acgcggagcg gttgtttttt tttttgtaat ccacaga 37

<210> 3920
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3920
ttgttaactt ttcatctcat ctttttcgtt acgtga 36

<210> 3921
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3921
acgttggtgc aaacgttttt tgatttctag tattttga 38

<210> 3922
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3922
tatgccctct aattctgggc tttttttttct gttgttga 38

<210> 3923
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3923
gtttcatctt gttttattag atcttagatt ttttgtga 38

<210> 3924
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3924
gcaatgcgat ttgtttcgtg tttatcctat ctagctga 38

<210> 3925
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3925
tcgtgctttt ctggtagttg accacactgt ttttttga 38

<210> 3926
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3926
ttccatacgt atatttttt gaaattcgac ttttttga 38

<210> 3927
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3927
acgtttttga cctctagttc tgtagcattt tgttttga 38

<210> 3928
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3928
actgaacttt ttttgtttac gaatccatct attgttga 38

<210> 3929
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3929
gccgcatagg ggcaaaattg tttttctctc ttacttga 38

<210> 3930
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3930
accctggggg tttgtaaggc tctttttttcg taatttga 38

<210> 3931
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3931
agttacatac cgtttttcgt tacttgaatt caaat 35

<210> 3932
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3932
acgatcacca caataatgtc ttttttttcgt tttcta 36

<210> 3933
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3933
gggactgaca tccataatcg ggtttttttt ctagaa 36

<210> 3934
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3934
atcgggtgcg cccccgaatt tttccatgtt tttgttga 38

<210> 3935
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3935
ttttgttacc cattttggtc cctatcctca ctcatga 37

<210> 3936
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3936
agttggaccc gtacgattcc ctaactttca acctaa 36

<210> 3937
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3937
tttctggtct aattatttaa ccattctctt gcttta 36

<210> 3938
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3938
attggcgcct atactgcttt gtattgtttt cgatttga 38

<210> 3939
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3939
tcctacaagc aaacggtgaa gttgaatttt atttttga 38

<210> 3940
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3940
actgctgact gttttatttg acgataataa ttttttga 38

<210> 3941
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3941
tttggtcgta tattctattt ttggatgtga attac 35

<210> 3942
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3942
tacggtacat ggtggagtct cttttgataa ttttttga 38

<210> 3943
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3943
ttttgcaatg gtgattaatg cggttcgcct gtttttga 38

<210> 3944
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3944
aggttgcagt ccgggatctt tattgttaat attta 35

<210> 3945
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3945
gtttaggctg tcaatggcct ttgttttta ttttaa 36

<210> 3946
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3946
aggacttatt acattctctg tgttccttat ctttatga 38

<210> 3947
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3947
gctctttcct gtcttgttta tttcgttggt tttcc 35

<210> 3948
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3948
gggttacgta cccgactttt tttctggttt tctct 35

<210> 3949
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3949
tcgcaagcta atagctttat tttttctgtc tttcta 36

<210> 3950
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3950
ttactatgca tcgtttgaat ccgttgtttc tccactga 38

<210> 3951
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3951
tggagtttct gctgctcgac ttggtatcct gtttgtga 38

<210> 3952
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3952
gagtaccgtg tcaacgtttt tttttgtatt atcaatga 38

<210> 3953
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3953
tctattaacc cgtttacgtg ctcacctcgt ttttttga 38

<210> 3954
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3954
gcatgagctc atttgtttta tcttgtcgtt ttttgga 37

<210> 3955
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3955
ttatctctat actatgttgt ttctctattt accactga 38

<210> 3956
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3956
tagctcacga cgatttattt tctcatttct tatttga 37

<210> 3957
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3957
ttgcaaggca tccgccttga gttttttttc agttt 35

<210> 3958
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3958
gcgttgtttc gttatggggt ttcttctcac aatgga 36

<210> 3959
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3959
gcaaatctcc ttgtttatcc atctattcct cttagga 37

<210> 3960
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3960
gccgtgcttc tctttctttt tcggttaaat ttcttga 37

<210> 3961
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3961
gtaggccttt gatttctttc cttccgtttc ttttgtga 38

<210> 3962
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3962
atctcggtaa tctcatcatg ggggttgttt attttga 37

<210> 3963
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3963
gcaggccgtt cctcatattt gttgaattgg ttgttga 37

<210> 3964
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3964
accgaacttt tttcggattc ctcggggttt ttttttga 38

<210> 3965
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3965
tggagtactg ggcttgtccg taatttttat gtaaca 36

<210> 3966
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3966
tccatgtacg tcgatattga catctttatt tttta 35

<210> 3967
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3967
atgttaggcc acgattgttt tattgttgat ttttttga 38

<210> 3968
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3968
gcggctcgca ttgcttttta tttcgtagac tatcctga 38

<210> 3969
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3969
tctacctgga tggcgataat ttttttgat ctttttga 38

<210> 3970
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3970
gagtgcccat tagtaatttt tcttccactt tattt 35

<210> 3971
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3971
gagtttccct ttttattagt gccttttagt tgtttga 37

<210> 3972
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3972
ccataacact tggaggttct ttgtttgttt tgttttga 38

<210> 3973
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3973
ttatttcctc tcttaacgtt ttttactaac ctatatga 38

<210> 3974
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3974
ccgtccgctc ctttaatccg ttttgttttt ctgctga 37

<210> 3975
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3975
taattggctt ttttacttat atttcctcat ttgtt 35

<210> 3976
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3976
tactacaggc ctctttcgat tttccgggtg ttttt 35

<210> 3977
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3977
gttctgtggt tcgttttctt aatgttagta ctctca 36

<210> 3978
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3978
ggctgttttt tcttgtattt cggatttacc aatatga 37

<210> 3979
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3979
gggggactat ctactaggct ttcgtttttt cctct 35

<210> 3980
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3980
agcaattatg tgcgtttgtt tttaatttat gttca 35

<210> 3981
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3981
ttccacacgg tgttgaacat ttgttttatt gatttga 37

<210> 3982
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3982
ggtgtcgctt atttttatct ttgaattggt cataaa 36

<210> 3983
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3983
aacttttatt gatatacgca ttttttttcct gggata 36

<210> 3984
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3984
catggtcgtt tttctgtcct ctaaacttac tatccga 37

<210> 3985
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3985
tagtctatgg cttggctttc catggtttcc tttcttga 38

<210> 3986
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3986
agcactttcg tcatagtggt tattgttgta aattgtga 38

<210> 3987
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3987
ttaagcgtac gaccgagttt ttatatcata tcttga 36

<210> 3988
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3988
gctctctctg gtataggttt ttttttgtta tccaca 36

<210> 3989
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3989
tgcctaggta tatttcgaat tttcatagga ttttca 36

<210> 3990
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3990
ttctttttta tgatgatccg atcctatttt ctttcga 37

<210> 3991
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3991
ttgttttcct tgttatgttt aaactgtcaa aatcca 36

<210> 3992
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3992
tactgcagat attttaagtt tctgttaatc tttgtga 37

<210> 3993
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3993
ttctctcaaa atagtttttc tttccgcagg atgta 35

<210> 3994
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3994
ttttagcgta actggatttt tttcacgacg ttttctga 38

<210> 3995
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3995
catgcatggc tttctttcca tgttctttct atccatga 38

<210> 3996
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3996
acatgcgacc gttttcttga ctttcttcgt tattga 36

<210> 3997
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3997
agctcctgtt ttccatttca aatcacgtta atctatga 38

<210> 3998
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3998
acggcgcctc tcctgatttt ccatcttttc tttcctga 38

<210> 3999
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 3999
tcgatatctc ttgtttttgt taaattcgtc ttcaa 35

<210> 4000
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4000
ctctgtccca aatccgtatc catttttct tactctga 38

<210> 4001
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4001
gtgcttcgct ctctgagttt ttcatctgtc aattt 35

<210> 4002
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4002
aagcaacgct gacctttcag actttttta tttgttga 38

<210> 4003
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4003
actctttctg ggggtttttc ttcttataga tttcttga 38

<210> 4004
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4004
ttatggtaca ctctctattg atgtttttct atctta 36

<210> 4005
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4005
ccctcaatta attatgcttg tattttgttt ctatct 36

<210> 4006
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4006
ttctcgctgt ttgttttccc attatgttac gctattga 38

<210> 4007
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4007
gaggttttgc gattggattt tttggttgtt atcttga 37

<210> 4008
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4008
aacccgtagt agcacaccct gtcgaaccca aataaa 36

<210> 4009
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4009
aacacttcta ggttcgcaac cccttataac cctaaa 36

<210> 4010
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4010
actccgcaat attcctttta aatttaggtt gcaac 35

<210> 4011
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4011
acttcaacga gcgttgatag ctcggatttt atcaa 35

<210> 4012
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4012
tctgcataac aaactcgaca cgccctagcc ccaata 36

<210> 4013
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4013
gagtacctca tcgcatatcg agcgcacttt atacaga 37

<210> 4014
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4014
accccaatac cattgcacct gaggactcaa tcaat 35

<210> 4015
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4015
ttccaccgag ttcctttcca gccaatatag catata 36

<210> 4016
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4016
ctcattcgct accaaccctg tgatctcttc taaac 35

<210> 4017
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4017
aactgtcatc cgccgactaa ctcaatctaa ctaca 35

<210> 4018
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4018
gagtactggc acgcgatacc ctaactgtac atcaaa 36

<210> 4019
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4019
ggctgcactt atctgtaaca catcactatt ccccaga 37

<210> 4020
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4020
tatctcaact cttctatatc acatgaccat aataaa 36

<210> 4021
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4021
tgaacacgaa aaaactaagc attaaatcac acacc 35

<210> 4022
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4022
aattaaccaa caaacccata acgagtccct ttatt 35

<210> 4023
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4023
aatgtaactg gcaatttctt tctcactgat ttctc 35

<210> 4024
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4024
acaatactgc ctccacactt ccaccttctt ttgtca 36

<210> 4025
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4025
acactatcaa gcgcttcgca tttcctttct ccaat 35

<210> 4026
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4026
acccgtgcgg cttaacctta tcttctttat ttcat 35

<210> 4027
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4027
accgctgggg ctctttccga caatttcttc cattt 35

<210> 4028
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4028
acgcaaccgt agcaaatttc ttagtgtttt ttttt 35

<210> 4029
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4029
acgggtcacg ccttccactt tacttggttt tattttga 38

<210> 4030
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4030
agcagtcatt ttacttctgt tttccttaaa attcta 36

<210> 4031
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4031
agccagtcac aacttctcgt ttcgctctct ttttctga 38

<210> 4032
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4032
agctaccggg gtttaccttc ctttcttctc tcctctga 38

<210> 4033
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4033
agttggcttg tcatcatttt tttgcttatc ttctctga 38

<210> 4034
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4034
atccaaacga tcaccttggt ctctctttac ttttca 36

<210> 4035
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4035
atccttattt ctatctcatc ctcttgtcgc cgact 35

<210> 4036
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4036
atcgaaggta ctacgtaaat ttctcattct ctttt 35

<210> 4037
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4037
atggtctgtc atggcaccga ttacatttct ttctc 35

<210> 4038
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4038
atgtgctctt cgttatttct cttatttcta tcttt 35

<210> 4039
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4039
attacttctg ctccattttc ctttttttcat acatt 35

<210> 4040
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4040
attcctgcgt tgcgtatctt ccttcgctta ttttc 35

<210> 4041
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4041
ccccactatt tatccgcaca tcttttgttc tctttga 37

<210> 4042
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4042
cctccgcctt actttcttat ttctctttac gaatt 35

<210> 4043
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4043
cgaggaccga cttttccact tatctttgcc ctttt 35

<210> 4044
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4044
ctccatcgaa ctctatctct ttttcttaac aatccga 37

<210> 4045
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4045
gagtacctca caattttccc gcgacgccct ttttta 36

<210> 4046
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4046
gatcccgttc ctttttaac gctctatttt cttttga 37

<210> 4047
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4047
gcaaatgcct ttcattaccc attgcttttc tgaca 35

<210> 4048
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4048
gcaatgaaca cctacacata cctgcctaca cgatta 36

<210> 4049
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4049
gcaatttttc attcttaggg tttctctcac ttccca 36

<210> 4050
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4050
gcacatttct taacttttct gtttggattt accaatga 38

<210> 4051
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4051
gcacggctta cgctcttctt ttccgctatc tatttga 37

<210> 4052
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4052
gcagttcgcc tccactcact cttttccgat ttatca 36

<210> 4053
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4053
gcatcctccc gtgtcgctca cttttccttt gcttc 35

<210> 4054
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4054
gccaactttg aatgggctct catcctatct ctctttga 38

<210> 4055
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4055
gcccgcgctc gcatcctatt tcttcttttt ttgtaa 36

<210> 4056
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4056
gccctcgtac tcttctttgg aacttacttt cattt 35

<210> 4057
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4057
gccgcattat ttcgtttttt cttcactctg cacttga 37

<210> 4058
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4058
gcctatctta ctatatctcg tttactttcc ttatatga 38

<210> 4059
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4059
gcctgtacct gtttactttt atcaatcttg tgcttga 37

<210> 4060
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4060
gccttccgct tggtgttttt tttactttta gatcta 36

<210> 4061
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4061
gcgatggtct accttctttt aacaaatttt cgctc 35

<210> 4062
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4062
gcgccttacg ttaattcaca tatcccttta ccatctga 38

<210> 4063
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4063
gcggtctggg aatttctacc ctgcttctat ctttttga 38

<210> 4064
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4064
gctacgcctt actctctgat aatctccact ttccg 35

<210> 4065
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4065
gctatctgag ctcccgcttt ctttcgatta tatat 35

<210> 4066
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4066
ggcccgattt cctatgtcat ttggtttttt ctttc 35

<210> 4067
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4067
ggccgtcgca tacctagatt tctttatcct ttttc 35

<210> 4068
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4068
ggcgccaatt tccttctatc gcaattcatt tcctctga 38

<210> 4069
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4069
ggctaggccg ccttcttatc tttatgccct tgttaa 36

<210> 4070
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4070
ggcttcttat ttgcttgaat atccgcttcc tctat 35

<210> 4071
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4071
gggcacggtt ttttcttttc tggtcgctat ccctta 36

<210> 4072
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4072
ggtaccctga ttttttctcg cctattttaa tattt 35

<210> 4073
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4073
ggtccaatta ccctttcaa cttatcgcac cattt 35

<210> 4074
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4074
gtttcgtggt atgcgaacct tctattactt atttta 36

<210> 4075
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4075
gttttcttcc cggttaaccc tttatttttt ttggt 35

<210> 4076
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4076
tacactttct ctagttacta ctcaccttct ttctctga 38

<210> 4077
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4077
tagttacttt taatcaatac tctctatttt tttgc 35

<210> 4078
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4078
tatccttcgc attattcttt ctcatcggtc cttta 35

<210> 4079
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4079
tatctttatt gctctgggtg attgcctctt gtttt 35

<210> 4080
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4080
tatttcccac gctattcatc tattctcttt acctt 35

<210> 4081
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4081
tccactggct tggcattgcc tccttgatcc ttctta 36

<210> 4082
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4082
tccagcacgc aacctcagct tcatcttttt ctctt 35

<210> 4083
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4083
tccatttgta ctgtgcgcct aattttctac tcttt 35

<210> 4084
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4084
tccgacctgg ttatttttt gccttttct attat 35

<210> 4085
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4085
tcgcgatttc ccatgtctct atacttctta caccca 36

<210> 4086
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4086
tcgtacgaga tgcactcttt ctacttaagt tttct 35

<210> 4087
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4087
tcgtcattgc cttccttttt catgtttcat tttctga 37

<210> 4088
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4088
tcgtgatctc tatttggttt cttccacgca tttct 35

<210> 4089
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4089
tcgttctatc cgtttgtgca tcttccttct ttacca 36

<210> 4090
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4090
tctaggcatc tccgcttcct aaatttcttt ctgta 35

<210> 4091
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4091
tctgcccgca ctactaaatt tttatctctt cacta 35

<210> 4092
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4092
tctttagcat ctcatctttt ctgccctcac gcctta 36

<210> 4093
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4093
tgcacgagga ttacctttc tttttctcga ttctt 35

<210> 4094
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4094
tgcttatccc gggaatatcc tttgctctct tttcca 36

<210> 4095
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4095
tggctctcac gtctccattt tattttctat ttttctga 38

<210> 4096
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4096
tggggcttaa ccatctcttc tttctctttt accagga 37

<210> 4097
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4097
tggtactaac ctctatctca tcctaaaatc tcttta 36

<210> 4098
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4098
tgtcgacctt taccttattc atttgaaatc gcttt 35

<210> 4099
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4099
ttccttgcaa tttcctttct ttttattcat ccggca 36

<210> 4100
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4100
ttcgaccctt ttccatactt tttctcacgc aaattga 37

<210> 4101
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4101
ttctcgatcc tttgccgcat ttctcacttt tctgttga 38

<210> 4102
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4102
ttctcgtagc tacaaatcct ttaacacttc tcttca 36

<210> 4103
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4103
ttgcgacgcc tctactttca ccccactaaa atttctga 38

<210> 4104
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4104
ttgtacgttc gctttatttt gcattccgcc ttttt 35

<210> 4105
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4105
tttcctcgct atttctgtct ctttgccctt cttggga 37

<210> 4106
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4106
ttttcaggta cgaccccata ctttttttc attttga 37

<210> 4107
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4107
aacgccgtac agttgctgat ttcacccact ctttt 35

<210> 4108
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4108
actccttcta aaattacact tatatcaccc atacc 35

<210> 4109
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4109
agaaccgtcc ctgcattttt attggttatt ttttt 35

<210> 4110
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4110
cttgcatatt gttcttcctg tttgcttttt cttcttga 38

<210> 4111
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4111
gagtacccta ttacgctccc cccccattat ctctt 35

<210> 4112
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4112
gcaccttcta ttttcacccc gatttaccct ttgcctga 38

<210> 4113
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4113
gccgccttgg gtccattatt atctttatat ttttctga 38

<210> 4114
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4114
gccgcttgcc tttatttatc atacgctatt aatcttga 38

<210> 4115
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4115
gcctggttat ctttgaatag ctttcgtatt ccttt 35

<210> 4116
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4116
gccttgcatc ttcacttatg gtgtcttacc acttta 36

<210> 4117
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4117
gcgaggctat tatctctggc gctgttcgcc cactt 35

<210> 4118
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4118
gctagcatgc actcctttac ttcttctcat cacacga 37

<210> 4119
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4119
gttcaactta tcttcctttt tctttctggg actggga 37

<210> 4120
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4120
taccgctcca ctcttcttta ttttcggtat ctttc 35

<210> 4121
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4121
tctagccgtc actattcttc gtacgtttcc atcct 35

<210> 4122
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4122
tctcactact cgctcttttt ccttttctcc acttatga 38

<210> 4123
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4123
tcttacctaa tttcttactc tgccacttta cacgc 35

<210> 4124
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4124
tggtacgtac tcgcgccgcg gtctttttct tttcctga 38

<210> 4125
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4125
ttttacttac ccataatttc cttacattac tttata 36

<210> 4126
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4126
aaatagctcg cacgagtagc cccgatagaa acataa 36

<210> 4127
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4127
aatcaaatca tacaatagag accgaaatta caaccga 37

<210> 4128
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4128
acaggcttgg ttcaacgaaa atcttcagtt ccggaa 36

<210> 4129
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4129
accctggggt gtccgggata aaaatataca aatca 35

<210> 4130
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4130
agagaataaa atgggggcta gaagctaaaa atataa 36

<210> 4131
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4131
cgtgcacatt ctaacggaat cagtaccaca caccatga 38

<210> 4132
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4132
ctgcttaaaa ctatcccata atcctaatga ccgga 35

<210> 4133
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4133
gaatgattta caccacaatt agactactga tatcaa 36

<210> 4134
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4134
gagttgctgc aagggactcc ctgaacactt atgacga 37

<210> 4135
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4135
gcgccgaata gttcgacccg aaaccctaat gaatcga 37

<210> 4136
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4136
gggctgtgca aacacaatat ataaaaactc aacattga 38

<210> 4137
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4137
ggtctgcccc tgaccgactc aaccgattaa gaaca 35

<210> 4138
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4138
taaaacgaaa ctaggaaaga tacacctaaa aagaa 35

<210> 4139
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4139
tactaacaac gggtagaccg acgaataccc acaca 35

<210> 4140
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4140
tactggaggg ctacaacctg ctacatagat acacatga 38

<210> 4141
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4141
tataacgccc taacatattg actacgagag actggga 37

<210> 4142
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4142
tcaatatccc gtcctcgggg ctagttatag tctaaa 36

<210> 4143
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4143
tcccataata aaccctatca agtgtagaca aacatga 37

<210> 4144
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4144
tcgaacccgc cacaaaatac ttaagtcagt caacaga 37

<210> 4145
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4145
tcgcattccc gacaaaatat atactggact acttc 35

<210> 4146
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4146
tgaaccgtta tattaccaaa ttagagctta gaaata 36

<210> 4147
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4147
tgacatacta taaccggggc cattaataat acacatga 38

<210> 4148
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4148
tgcaacgtaa aacacacaaa ctctaatgat accgga 36

<210> 4149
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4149
tgcgacaaca gccattcggg aactacccga tcacca 36

<210> 4150
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4150
ttccggctat aatacaaaca tatcacccga cttgaa 36

<210> 4151
<211> 7
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4151
acaatcc 7

<210> 4152
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4152
actatagaac aatccaccgc tttaaatcaa ctatctga 38

<210> 4153
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4153
actatagaac aatccaccgc tttaaatcaa ctatctca 38

<210> 4154
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4154
actatagaac aatccaccgc tttaaatcaa ctatctaa 38

<210> 4155
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4155
actatagaac aatccaccgc tttaaataaa ctatctga 38

<210> 4156
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4156
actatagaac aatccacctc tttaaatcaa ctatctga 38

<210> 4157
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4157
actatagaac aatccaccgc tttaaatcaa ctatccga 38

<210> 4158
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4158
aatatagaac aatccaccgc tttaaatcaa ctatctga 38

<210> 4159
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4159
actatagaac aatccaccgc tttaaatcaa ctatccta 38

<210> 4160
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4160
actatagaac aatccacctc tttaaatcaa ctatctta 38

<210> 4161
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4161
actatagaac aatcccccgc tttaaatcaa ctatctga 38

<210> 4162
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4162
actatagaac aatccgccgc tttaaatcaa ctatctga 38

<210> 4163
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4163
actatagaac aatcctccgc tttaaatcaa ctatctga 38

<210> 4164
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4164
actatataac aatccaccgc tttaaatcaa ctatctga 38

<210> 4165
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4165
taccctgac aatcctcgcg ccgaggcctc catcctaa 38

<210> 4166
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4166
taccctgac aatcctcgcg ccgaggcctc catcctaa 38

<210> 4167
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4167
gattttaaaa cccttgcacc tgattgtgcc aatcaa 36

<210> 4168
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4168
gattttaaaa cccttgcacc tgattgtgcc aatcga 36

<210> 4169
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4169
gattttaaaa cccttgcacc tgattgtgcc aatcta 36

<210> 4170
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4170
aacccaattc acatacactc tcaccctcac taaaca 36

<210> 4171
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4171
aacccaattc acatacactc tcaccctcac taaaaa 36

<210> 4172
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4172
aacccaattc acatacactc tcaccctcac taaaga 36

<210> 4173
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4173
aacccaattc acatacactc tcaccctcac taaata 36

<210> 4174
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4174
aacccaattc acatacactc tcaccctcac taacca 36

<210> 4175
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4175
aacccaattc acatacactc tcaccctcac taacaa 36

<210> 4176
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4176
aacccaattc acatacactc tctccctcac taaaca 36

<210> 4177
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4177
aacccaattc acatacactc tccccctcac taaaca 36

<210> 4178
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4178
aacccaattc acatacactc tcgccctcac taaaca 36

<210> 4179
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4179
aacccaattc acatactctc tcaccctcac taaaca 36

<210> 4180
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4180
aacccaattc acataaactc tcaccctcac taaaca 36

<210> 4181
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4181
aacccaatcc acatacactc tcaccctcac taaaca 36

<210> 4182
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4182
aacccaattc acatacactc tcaccctcac taaacc 36

<210> 4183
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4183
aacccaattc acatacgctc tcaccctcac taaaca 36

<210> 4184
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4184
aacccaattc acatacactc tctccctcac taaaaa 36

<210> 4185
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4185
aacccaattc acataccctc tcaccctcac taaaca 36

<210> 4186
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4186
aacccaattc acatacactc tcgccctcac taaaaa 36

<210> 4187
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4187
aacccaattc acatacactc ttaccctcac taaaca 36

<210> 4188
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4188
aacccaatac acatacactc tcaccctcac taaaca 36

<210> 4189
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4189
aacccaattc acatacactc tcaccctctc taaaca 36

<210> 4190
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4190
aacccaattc acatacactc tccccctcac taaaaa 36

<210> 4191
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4191
aacccaattc acataccctc tcaccctcac taaaaa 36

<210> 4192
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4192
aacccaattc acatacactc tcaccctcac taaact 36

<210> 4193
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4193
aacccaattc acatacaccc tcaccctcac taaaca 36

<210> 4194
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4194
aacccaattc acatactctc tcaccctcac taaaaa 36

<210> 4195
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4195
aacccaatgc acatacactc tcaccctcac taaaca 36

<210> 4196
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4196
agcccaattc acatacactc tcaccctcac taaaca 36

<210> 4197
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4197
aacccaattc acatacactc taaccctcac taaaca 36

<210> 4198
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4198
aacccaaatc acatacactc tcaccctcac taaaca 36

<210> 4199
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4199
aacccaattc acatacactc tcaccctcac taaaat 36

<210> 4200
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4200
aacccaattc acatacactc tctccctcac taaaga 36

<210> 4201
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4201
aacccaattc tcatacactc tcaccctcac taaaca 36

<210> 4202
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4202
atcccaattc acatacactc tcaccctcac taaaca 36

<210> 4203
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4203
aacccaactc acatacactc tcaccctcac taaaca 36

<210> 4204
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4204
aacccaattc acatacactc tcaccctcac taaaac 36

<210> 4205
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4205
aacccaattc acatacactc tcaccctcac taacta 36

<210> 4206
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4206
aacccaattc acatacactc tcgccctcac taaaga 36

<210> 4207
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4207
aacccaattc acattcactc tcaccctcac taaaca 36

<210> 4208
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4208
aaccaaattc acatacactc tcaccctcac taaaca 36

<210> 4209
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4209
aacccaattc acataaactc tcaccctcac taaaaa 36

<210> 4210
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4210
aacccaattc acatacacgc tcaccctcac taaaca 36

<210> 4211
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4211
aacccaattc acatacactc tcacactcac taaaca 36

<210> 4212
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4212
aacccaattc acatacactc tcaccatcac taaaca 36

<210> 4213
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4213
aacccaattc acatacactc tcaccctcac taacga 36

<210> 4214
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4214
aacccaattc acatacactc tcaccctcgc taaaca 36

<210> 4215
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4215
aacccaattc acatacactc tccccctcac taaaga 36

<210> 4216
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4216
aacccaattc gcatacactc tcaccctcac taaaca 36

<210> 4217
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4217
aacccaattc acatacactc acaccctcac taaaca 36

<210> 4218
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4218
aacccaattc acatacactc ccaccctcac taaaca 36

<210> 4219
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4219
aacccaattc acatacactc tcaccctcac taaacg 36

<210> 4220
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4220
aacccaattc acatactctc tcaccctcac taaaga 36

<210> 4221
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4221
aacccaattc acatactctc tcaccctcac taaata 36

<210> 4222
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4222
aacccagttc acatacactc tcaccctcac taaaca 36

<210> 4223
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4223
aaaccaattc acatacactc tcaccctcac taaaca 36

<210> 4224
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4224
aacacaattc acatacactc tcaccctcac taaaca 36

<210> 4225
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4225
aacccaactc acatacactc tcaccctcac taaaaa 36

<210> 4226
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4226
aacccaagtc acatacactc tcaccctcac taaaca 36

<210> 4227
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4227
aacccaatcc acatacactc tcaccctcac taaaaa 36

<210> 4228
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4228
aacccaatta acatacactc tcaccctcac taaaca 36

<210> 4229
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4229
aacccaattc acaaacactc tcaccctcac taaaca 36

<210> 4230
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4230
aacccaattc acacacactc tcacccccac taaaca 36

<210> 4231
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4231
aacccaattc acacacactc tcaccctcac taaaca 36

<210> 4232
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4232
aacccaattc acacacactc tccccctcac taaaca 36

<210> 4233
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4233
aacccaattc acagacactc tcaccctcac taaaca 36

<210> 4234
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4234
aacccaattc acatacaatc tcaccctcac taaaca 36

<210> 4235
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4235
aacccaattc acatacacac tcaccctcac taaaca 36

<210> 4236
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4236
aacccaattc acatacaccc tcaccctcac taaaaa 36

<210> 4237
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4237
aacccaattc acatacacta tcaccctcac taaaaa 36

<210> 4238
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4238
aacccaattc acatacacta tcaccctcac taaaca 36

<210> 4239
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4239
aacccaattc acatacactc ccaccctcac taaaaa 36

<210> 4240
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4240
aacccaattc acatacactc gcaccctcac taaaca 36

<210> 4241
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4241
aacccaattc acatacactc tcaacctcac taaaca 36

<210> 4242
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4242
aacccaattc acatacactc tcacactcac taaaaa 36

<210> 4243
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4243
aacccaattc acatacactc tcaccatccc taaaca 36

<210> 4244
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4244
aacccaattc acatacactc tcacccacac taaaca 36

<210> 4245
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4245
aacccaattc acatacactc tcacccgcac taaaca 36

<210> 4246
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4246
aacccaattc acatacactc tcaccctaac taaaca 36

<210> 4247
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4247
aacccaattc acatacactc tcaccctcaa taaaca 36

<210> 4248
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4248
aacccaattc acatacactc tcaccctcac caaaca 36

<210> 4249
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4249
aacccaattc acatacactc tcaccctcac taaaag 36

<210> 4250
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4250
aacccaattc acatacactc tcaccctcac taaagc 36

<210> 4251
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4251
aacccaattc acatacactc tcaccctcac taaagt 36

<210> 4252
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4252
aacccaattc acatacactc tcaccctcac taaatc 36

<210> 4253
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4253
aacccaattc acatacactc tcaccctcac taaatt 36

<210> 4254
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4254
aacccaattc acatacactc tcaccctcac taagaa 36

<210> 4255
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4255
aacccaattc acatacactc tcaccctcac taagca 36

<210> 4256
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4256
aacccaattc acatacactc tcaccctcac taagga 36

<210> 4257
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4257
aacccaattc acatacactc tcaccctcac taagta 36

<210> 4258
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4258
aacccaattc acatacactc tcaccctcac taataa 36

<210> 4259
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4259
aacccaattc acatacactc tcaccctcac taatca 36

<210> 4260
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4260
aacccaattc acatacactc tcaccctcac taatga 36

<210> 4261
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4261
aacccaattc acatacactc tcaccctcac taatta 36

<210> 4262
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4262
aacccaattc acatacactc tcaccctcac tacaca 36

<210> 4263
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4263
aacccaattc acatacactc tcaccctcac tagaca 36

<210> 4264
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4264
aacccaattc acatacactc tcaccctcac tataca 36

<210> 4265
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4265
aacccaattc acatacactc tcaccctcac tcaaca 36

<210> 4266
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4266
aacccaattc acatacactc tcaccctcac tgaaca 36

<210> 4267
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4267
aacccaattc acatacactc tcaccctcac ttaaca 36

<210> 4268
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4268
aacccaattc acatacactc tcaccctcat taaaca 36

<210> 4269
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4269
aacccaattc acatacactc tcaccctccc taaaaa 36

<210> 4270
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4270
aacccaattc acatacactc tcaccctccc taaaca 36

<210> 4271
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4271
aacccaattc acatacactc tcaccctccc taaaga 36

<210> 4272
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4272
aacccaattc acatacactc tcaccctccc taaata 36

<210> 4273
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4273
aacccaattc acatacactc tcaccctcgc taaaaa 36

<210> 4274
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4274
aacccaattc acatacactc tcaccctctc taaaaa 36

<210> 4275
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4275
aacccaattc acatacactc tcaccctgac taaaca 36

<210> 4276
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4276
aacccaattc acatacactc tcacccttac taaaca 36

<210> 4277
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4277
aacccaattc acatacactc tcaccttcac taaaca 36

<210> 4278
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4278
aacccaattc acatacactc tcaccttctc taaaca 36

<210> 4279
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4279
aacccaattc acatacactc tcacgctcac taaaca 36

<210> 4280
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4280
aacccaattc acatacactc tcactctcac taaaaa 36

<210> 4281
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4281
aacccaattc acatacactc tcactctcac taaaca 36

<210> 4282
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4282
aacccaattc acatacactc tcagcctcac taaaca 36

<210> 4283
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4283
aacccaattc acatacactc tcatcctcac taaaca 36

<210> 4284
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4284
aacccaattc acatacactc tccccctcac taaata 36

<210> 4285
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4285
aacccaattc acatacactc tcgccctcac taaata 36

<210> 4286
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4286
aacccaattc acatacactc tctccctcac taaata 36

<210> 4287
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4287
aacccaattc acatacactg tcaccctcac taaaca 36

<210> 4288
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4288
aacccaattc acatacactt tcaccctcac taaaca 36

<210> 4289
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4289
aacccaattc acatacattc tcaccctcac taaaca 36

<210> 4290
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4290
aacccaattc acataccctc tcaccctcac taaaga 36

<210> 4291
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4291
aacccaattc acatacgctc tcaccctcac taaaaa 36

<210> 4292
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4292
aacccaattc acatacgctc tcaccctcac taaaga 36

<210> 4293
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4293
aacccaattc acatatactc tcaccctcac taaaaa 36

<210> 4294
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4294
aacccaattc acatatactc tcaccctcac taaaca 36

<210> 4295
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4295
aacccaattc acatccactc tcaccctcac taaaca 36

<210> 4296
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4296
aacccaattc acatgcactc tcaccctcac taaaca 36

<210> 4297
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4297
aacccaattc acctacactc tcaccctcac taaaaa 36

<210> 4298
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4298
aacccaattc acctacactc tcaccctcac taaaca 36

<210> 4299
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4299
aacccaattc acgtacactc tcaccctcac taaaca 36

<210> 4300
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4300
aacccaattc acttacactc tcaccctcac taaaca 36

<210> 4301
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4301
aacccaattc agatacactc tcaccctcac taaaca 36

<210> 4302
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4302
aacccaattc atatacactc tcaccctcac taaaaa 36

<210> 4303
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4303
aacccaattc atatacactc tcaccctcac taaaca 36

<210> 4304
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4304
aacccaattc ccatacactc tcaccctcac taaaca 36

<210> 4305
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4305
aacccaattt acatacactc tcaccctcac taaaca 36

<210> 4306
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4306
aacccacttc acatacactc tcaccctcac taaaca 36

<210> 4307
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4307
aacccatttc acatacactc tcaccctcac taaaca 36

<210> 4308
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4308
aaccccattc acatacactc tcaccctcac taaaaa 36

<210> 4309
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4309
aaccccattc acatacactc tcaccctcac taaaca 36

<210> 4310
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4310
aacccgattc acatacactc tcaccctcac taaaca 36

<210> 4311
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4311
aaccctattc acatacactc tcaccctcac taaaaa 36

<210> 4312
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4312
aaccctattc acatacactc tcaccctcac taaaca 36

<210> 4313
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4313
aaccgaattc acatacactc tcaccctcac taaaca 36

<210> 4314
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4314
aacctaattc acatacactc tcaccctcac taaaaa 36

<210> 4315
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4315
aacctaattc acatacactc tcaccctcac taaaca 36

<210> 4316
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4316
aactcaattc acatacactc tcaccctcac taaaca 36

<210> 4317
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4317
aatccaattc acatacactc tcaccctcac taaaca 36

<210> 4318
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4318
accccaattc acatacactc tcaccctcac taaaaa 36

<210> 4319
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4319
accccaattc acatacactc tcaccctcac taaaca 36

<210> 4320
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4320
agcccaattc acatacactc tcaccctcac taaaaa 36

<210> 4321
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4321
atcccaattc acatacactc tcaccctcac taaaaa 36

<210> 4322
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4322
cacccaattc acatacactc tcaccctcac taaaaa 36

<210> 4323
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4323
cacccaattc acatacactc tcaccctcac taaaca 36

<210> 4324
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4324
gacccaattc acatacactc tcaccctcac taaaca 36

<210> 4325
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4325
tacccaattc acatacactc tcaccctcac taaaca 36

<210> 4326
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4326
ttaagatagt tgatttaaag cggtggattg ttctatagt 39

<210> 4327
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4327
tactatagat ggattgtccg ctttaaatca actatctta 39

<210> 4328
<211> 91
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4328
ctagcatgac tgcagtacgt actatagaac aatccaccgc tttaaatcaa ctatcttact 60

gtctcttata cacatctgac gctgccgacg a 91

<210> 4329
<211> 46
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4329
acgtactata gaacaatcca ccgctttaaa tcaactatct tactgt 46

<210> 4330
<211> 41
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4330
gtactataga acaatccacc gctttaaatc aactatctta c 41

<210> 4331
<211> 91
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4331
ctagcatgac tgcagtacgt acgatagaac aatccaccgc tttaaatcaa ctatcttact 60

gtctcttata cacatctgac gctgccgacg a 91

<210> 4332
<211> 46
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4332
acgtacgata gaacaatcca ccgctttaaa tcaactatct tactgt 46

<210> 4333
<211> 91
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4333
ctagcatgac tgcagtacgt actatagaac aatccaccgc tttaaatcaa ctatctgact 60

gtctcttata cacatctgac gctgccgacg a 91

<210> 4334
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4334
atagaacaat ccaccgcttt aaatcaacta tctgactgtc tcttatacac atc 53

<210> 4335
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4335
ctagcatgac tgcagtacgt aacccaattc acatacactc tcacccccac taaacactgt 60

ctcttataca catctgacgc tgccgacga 89

<210> 4336
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4336
ctagcatgac tgcagtacgt aacccaattc acatacactc tcaccctcac taaacactgt 60

ctcttataca catctgacgc tgccgacga 89

<210> 4337
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4337
gtaacccaat tcacatac 18

<210> 4338
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4338
ggtaacccaa ttcacatacc 20

<210> 4339
<211> 91
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4339
ctagcatgac tgcagtacgt taagatagtt gatttaaagc ggtggattgt tctatagtct 60

gtctcttata cacatctgac gctgccgacg a 91

<210> 4340
<211> 91
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4340
ctagcatgac tgcagtacgt taccccctgac aatcctcgcg ccgaggcctc catcctgact 60

gtctcttata cacatctgac gctgccgacg a 91

<210> 4341
<211> 91
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4341
ctagcatgac tgcagtacgt taccccctgac aatcctcgcg ccgaggcctc catcctcact 60

gtctcttata cacatctgac gctgccgacg a 91

<210> 4342
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4342
ctagcatgac tgcagtacgt gattttaaaa cccttgcacc tgattgtgcc aatccactgt 60

ctcttataca catctgacgc tgccgacga 89

<210> 4343
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4343

Pro Gly Arg Arg Gly Arg Pro Gly Leu Lys Gly Glu Gln
1 5 10

<210> 4344
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4344

Arg Gly Arg Pro Gly Leu Lys Gly Glu Gln Gly Glu Pro
1 5 10

<210> 4345
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4345

Lys Gly Glu Gln Gly Glu Pro Gly Ala Pro Gly Ile Arg
1 5 10

<210> 4346
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4346

Gln Gly Glu Pro Gly Ala Pro Gly Ile Arg Thr Gly Ile
1 5 10

<210> 4347
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4347

Thr Gly Ile Gln Gly Leu Lys Gly Asp Gln Gly Glu Pro
1 5 10

<210> 4348
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4348

Lys Gly Asp Gln Gly Glu Pro Gly Pro Ser Gly Asn Pro
1 5 10

<210> 4349
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4349

Asp Gln Gly Glu Pro Gly Pro Ser Gly Asn Pro Gly Lys
1 5 10

<210> 4350

<211> 13
<212> PRT
<213> Homo sapiens

<400> 4350

Pro Gly Pro Ser Gly Asn Pro Gly Lys Val Gly Tyr Pro
1 5 10

<210> 4351
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4351

Pro Ser Gly Asn Pro Gly Lys Val Gly Tyr Pro Gly Pro
1 5 10

<210> 4352
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4352

Pro Gly Lys Val Gly Tyr Pro Gly Pro Ser Gly Pro Leu
1 5 10

<210> 4353
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4353

Lys Val Gly Tyr Pro Gly Pro Ser Gly Pro Leu Gly Ala
1 5 10

<210> 4354
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4354

Lys Gly Thr Lys Gly Ser Pro Gly Asn Ile Lys Asp Gln
1 5 10

<210> 4355
<211> 13
<212> PRT
<213> Homo sapiens

<400> 4355

Ile Arg Arg Asn Pro Pro Met Gly Gly Asn Val Val Ile
1 5 10

<210> 4356
<211> 13
<212> PRT

<213> Homo sapiens

<400> 4356

Gln Val Val Ser Gly Gly Met Val Leu Gln Leu Gln Gln
1 5 10

<210> 4357
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4357
ctaaccccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4358
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4358
ctaaccccgg gtgtggtggg tgggcagggg ggttagatag aacaatccac cgctttaaat 60

caactatctg actgtctctt atacacatc 89

<210> 4359
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4359
atagaacaat ccaccgcttt aaatcaacta tctgactgtc tcttatacac atcctaaccc 60

cgggtgtggt gggtgggcag gggggttag 89

<210> 4360
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4360
atagaacaat tcaccgcttt aaatcaacta tctgactgtc tcttatacac atcctaaccc 60

cgggtgtggt gggtgggcag gggggttag 89

<210> 4361
<211> 82
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 4361
ctaaccccgg gtgtggtggg tgggcagggg ggttagacgt actatagaac aatccaccgc 60

tttaaatcaa ctatcttact gt 82

<210> 4362
<211> 82
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4362
acgtactata gaacaatcca ccgctttaaa tcaactatct tactgtctaa ccccgggtgt 60

ggtgggtggg caggggggtt ag 82

<210> 4363
<211> 77
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4363
ctaaccccgg gtgtggtggg tgggcagggg ggttaggtac tatagaacaa tccaccgctt 60

taaatcaact atcttac 77

<210> 4364
<211> 77
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4364
gtactataga acaatccacc gctttaaatc aactatctta cctaaccccg ggtgtggtgg 60

gtgggcaggg gggttag 77

<210> 4365
<211> 82
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4365
ctaaccccgg gtgtggtggg tgggcagggg ggttagacgt acgatagaac aatccaccgc 60

tttaaatcaa ctatcttact gt 82

<210> 4366
<211> 82
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4366
acgtacgata gaacaatcca ccgctttaaa tcaactatct tactgtctaa ccccgggtgt 60

ggtgggtggg caggggggtt ag 82

<210> 4367
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4367
ctaaccccgg gtgtggtggg tgggcagggg ggttaggtaa cccaattcac atac 54

<210> 4368
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4368
gtaacccaat tcacatacct aaccccgggt gtggtgggtg ggcaggggggg ttag 54

<210> 4369
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4369
ctaaccccgg gtgtggtggg tggacagggg ggttag 36

<210> 4370
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide

<400> 4370
ggtggtggtg gttgtggtgg tggtgg 26

<210> 4371
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 4371
cctcctcctc cttctcctcc tcctcc 26

<210> 4372
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (1)..(1)
<223> n is a, c, g, or t

<400> 4372
ntaaccccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4373
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (2)..(2)
<223> n is a, c, g, or t

<400> 4373
cnaaccccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4374
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (3)..(3)
<223> n is a, c, g, or t

<400> 4374
ctnaccccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4375
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (4)..(4)

<223> n is a, c, g, or t

<400> 4375
ctanccccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4376
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (5)..(5)
<223> n is a, c, g, or t

<400> 4376
ctaancccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4377
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (6)..(6)
<223> n is a, c, g, or t

<400> 4377
ctaacnccgg gtgtggtggg tgggcagggg ggttag 36

<210> 4378
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (7)..(7)
<223> n is a, c, g, or t

<400> 4378
ctaaccncgg gtgtggtggg tgggcagggg ggttag 36

<210> 4379
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (8)..(8)
<223> n is a, c, g, or t

<400> 4379
ctaacccngg gtgtggtggg tgggcagggg ggttag 36

<210> 4380
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (9)..(9)
<223> n is a, c, g, or t

<400> 4380
ctaaccccng gtgtggtggg tgggcagggg ggttag 36

<210> 4381
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (10)..(10)
<223> n is a, c, g, or t

<400> 4381
ctaaccccgn gtgtggtggg tgggcagggg ggttag 36

<210> 4382
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (11)..(11)
<223> n is a, c, g, or t

<400> 4382
ctaaccccgg ntgtggtggg tgggcagggg ggttag 36

<210> 4383
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (12)..(12)
<223> n is a, c, g, or t

<400> 4383
ctaaccccgg gngtggtggg tgggcagggg ggttag 36

<210> 4384
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (13)..(13)
<223> n is a, c, g, or t

<400> 4384
ctaaccccgg gtntggtggg tgggcagggg ggttag 36

<210> 4385
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (14)..(14)
<223> n is a, c, g, or t

<400> 4385
ctaaccccgg gtgnggtggg tgggcagggg ggttag 36

<210> 4386
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (15)..(15)
<223> n is a, c, g, or t

<400> 4386
ctaaccccgg gtgtngtggg tgggcagggg ggttag 36

<210> 4387
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (16)..(16)
<223> n is a, c, g, or t

<400> 4387
ctaaccccgg gtgtgntggg tgggcagggg ggttag 36

<210> 4388
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (17)..(17)
<223> n is a, c, g, or t

<400> 4388
ctaaccccgg gtgtggnggg tgggcagggg ggttag 36

<210> 4389
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (18)..(18)
<223> n is a, c, g, or t

<400> 4389
ctaaccccgg gtgtggtngg tgggcagggg ggttag 36

<210> 4390
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (19)..(19)
<223> n is a, c, g, or t

<400> 4390
ctaaccccgg gtgtggtgng tgggcagggg ggttag 36

<210> 4391
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (20)..(20)
<223> n is a, c, g, or t

<400> 4391
ctaaccccgg gtgtggtggn tgggcagggg ggttag 36

<210> 4392
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (21)..(21)
<223> n is a, c, g, or t

<400> 4392
ctaaccccgg gtgtggtggg ngggcagggg ggttag 36

<210> 4393
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (22)..(22)
<223> n is a, c, g, or t

<400> 4393
ctaaccccgg gtgtggtggg tnggcagggg ggttag 36

<210> 4394
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>

<221> misc_feature
<222> (23)..(23)
<223> n is a, c, g, or t

<400> 4394
ctaaccccgg gtgtggtggg tgngcagggg ggttag 36

<210> 4395
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (24)..(24)
<223> n is a, c, g, or t

<400> 4395
ctaaccccgg gtgtggtggg tggncagggg ggttag 36

<210> 4396
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (25)..(25)
<223> n is a, c, g, or t

<400> 4396
ctaaccccgg gtgtggtggg tgggnagggg ggttag 36

<210> 4397
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (26)..(26)
<223> n is a, c, g, or t

<400> 4397
ctaaccccgg gtgtggtggg tgggcngggg ggttag 36

<210> 4398
<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (27)..(27)
<223> n is a, c, g, or t

<400> 4398
ctaaccccgg gtgtggtggg tgggcanggg ggttag 36

<210> 4399
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (28)..(28)
<223> n is a, c, g, or t

<400> 4399
ctaaccccgg gtgtggtggg tgggcagngg ggttag 36

<210> 4400
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (29)..(29)
<223> n is a, c, g, or t

<400> 4400
ctaaccccgg gtgtggtggg tgggcaggng ggttag 36

<210> 4401
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (30)..(30)
<223> n is a, c, g, or t

<400> 4401
ctaaccccgg gtgtggtggg tgggcagggn ggttag 36

<210> 4402
<211> 36

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (31)..(31)
<223> n is a, c, g, or t

<400> 4402
ctaaccccgg gtgtggtggg tgggcagggg ngttag 36

<210> 4403
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (32)..(32)
<223> n is a, c, g, or t

<400> 4403
ctaaccccgg gtgtggtggg tgggcagggg gnttag 36

<210> 4404
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (33)..(33)
<223> n is a, c, g, or t

<400> 4404
ctaaccccgg gtgtggtggg tgggcagggg ggntag 36

<210> 4405
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (34)..(34)
<223> n is a, c, g, or t

<400> 4405
ctaaccccgg gtgtggtggg tgggcagggg ggtnag 36

<210> 4406
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (35)..(35)
<223> n is a, c, g, or t

<400> 4406
ctaaccccgg gtgtggtggg tgggcagggg ggttng 36

<210> 4407
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<220>
<221> misc_feature
<222> (36)..(36)
<223> n is a, c, g, or t

<400> 4407
ctaaccccgg gtgtggtggg tgggcagggg ggttan 36