



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2139948 C 2001/12/04

(11)(21) **2 139 948**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1993/07/09
(87) Date publication PCT/PCT Publication Date: 1994/01/20
(45) Date de délivrance/Issue Date: 2001/12/04
(85) Entrée phase nationale/National Entry: 1995/01/10
(86) N° demande PCT/PCT Application No.: US 93/06479
(87) N° publication PCT/PCT Publication No.: WO 94/01139
(30) Priorité/Priority: 1992/07/13 (07/912,934) US

(51) Cl.Int.⁶/Int.Cl.⁶ A61K 48/00, A01K 67/027, C12N 5/10,
C12N 15/87, C12Q 1/02, G01N 33/48
(72) Inventeur/Inventor:
Ledley, Fred D., US
(73) Propriétaire/Owner:
BAYLOR COLLEGE OF MEDICINE, US
(74) Agent: SMART & BIGGAR

(54) Titre : APPLICATION AUX ARTICULATIONS D'UNE THERAPIE GENIQUE SOMATIQUE
(54) Title: TARGETING SOMATIC GENE THERAPY TO JOINTS

(57) **Abrégé/Abstract:**

A method of transfecting a cell in a structure of a joint is disclosed, wherein a DNA vector containing a nucleic acid cassette encoding, for example, a cell ablation agent or a therapeutic agent, is directly injected into the joint. The invention also includes a transfected synovial cell.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 48/00, 39/395, C12N 5/10 C12N 15/85, 15/90	A1	(11) International Publication Number: WO 94/01139 (43) International Publication Date: 20 January 1994 (20.01.94)
(21) International Application Number: PCT/US93/06479 (22) International Filing Date: 9 July 1993 (09.07.93) (30) Priority data: 07/912,934 13 July 1992 (13.07.92) US (71) Applicant: BAYLOR COLLEGE OF MEDICINE [US/ US]; One Baylor Plaza, Houston, TX 77030 (US). (72) Inventor: LEDLEY, Fred, D. ; 4911 Braesvalley, Houston, TX 77096 (US). (74) Agent: WARBURG, Richard, J.; Lyon & Lyon, 611 West Sixth Street, Suite 3400, Los Angeles, CA 90017 (US).		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: TARGETING SOMATIC GENE THERAPY TO JOINTS (57) Abstract A method of transfecting a cell in a structure of a joint is disclosed, wherein a DNA vector containing a nucleic acid cassette encoding, for example, a cell ablation agent or a therapeutic agent, is directly injected into the joint. The invention also includes a transfected synovial cell.		

-1-

TARGETING SOMATIC GENE THERAPY TO JOINTS

FIELD OF THE INVENTION

5 The present invention relates generally to the introduction of genetic material into the joints and cells comprising the joint structures for the purpose of somatic gene therapy. Further, it relates to methods for introducing novel genetic material into cells within the joint and achieving expression of gene products with therapeutic potential.

BACKGROUND OF INVENTION

10 Somatic gene therapy involves the treatment of genetic or acquired disease by the introduction of recombinant genes into somatic cells. Many different organs have been traditionally considered targets for somatic gene therapy including the bone
15 marrow, liver, endothelial cells, epithelial cells, fibroblasts, and muscle. In general, somatic gene therapy has traditionally been considered for those organs and cells which can be studied in vitro or ex vivo. One of the reasons for this focus is that many schemes for somatic gene therapy involve the harvest of cells from an organ
20 by surgical means, growing these cells in the laboratory, introducing genes into these cells using viral vectors, and then reimplanting these cells into the body using a cellular transplantation procedure. Broad applications of ex vivo schemes for gene therapy, particularly the transplantation of fibroblasts
25 into many sites, have been proposed.

More recently attention has turned to schemes for somatic gene therapy which involve in vivo delivery of recombinant genes to essential tissues by direct injection or targeted delivery. The present invention describes a novel approach to somatic gene therapy

2139948

-2-

5 involving the direct delivery of recombinant genes to joint spaces and cells comprising essential structures of the joint for the purposes of treating various forms of arthritis or other diseases of the joints. Major joints may be targeted for somatic gene therapy by injection of genetic material into the joint space. This therapy would be intended to prevent the often crippling and painful manifestations of inflammatory or degenerative arthritis as well as to enhance the process of repair or regeneration after illness, injury, or surgery.

10 SUMMARY OF INVENTION

An object of the present invention is the transformation of cells comprising essential structures of the joint.

15 An additional object of the present invention is a method of direct delivery of genetic material to the cells comprising essential structures of the joint.

A further object of the present invention is the in vivo introduction of genetic material into cells comprising essential structures of the joint.

20 Another object of the present invention is a method of injecting genetic material into the joint space for uptake by cells constituting structures of the joint.

25 An additional object of the present invention is the coupling of genetic materials to non-genetic materials termed vehicles which enhance the uptake, stability, and expression of genetic material into cells of the joint.

A further object of the present invention is a method for treating pathophysiological conditions resulting from inflammatory processes.

30 Another object of the present invention is a method for treating hypertrophy or inappropriate proliferation of cellular elements of the joint.

60724-2257

-3-

A further object of the present invention is a method for enhancing repair, regeneration, and recovery of essential structures comprising the joint after surgery or injury.

An additional object of the present invention is a
5 method for ablating certain cells in the joint.

A further object of the present invention is a method for generating animal models of arthritis.

In accomplishing the foregoing objects there is provided in accordance with one aspect of the present invention
10 a medicament for transfecting a cell in a structure of a joint comprising (a) a pharmaceutically acceptable carrier, diluent or vehicle and (b) an effective amount of a DNA vector adapted to the introduced directly into the joint. The vector is comprised of elements required for directing transcription of
15 the genetic material and expression of a specific gene product encoded by a nucleic acid cassette within the vector. The following elements are linked sequentially at appropriate distance for allowing functional expression: a promoter; a 5' mRNA leader sequence; an initiation site; a nucleic acid
20 cassette containing the sequence to be expressed; a 3' untranslated region; and a polyadenylation signal. One or more of these elements may be eliminated for specific applications.

In an embodiment of this invention the vector is comprised of DNA sequences which can be inserted into cells of
25 the joint and can transform these cells to produce novel gene products.

In a preferred embodiment, the nucleic acid cassette encodes a protein, polypeptide, or anti-sense RNA. The protein can be selected from the group including enzymes, proteins
30 constituents of the extracellular matrix, as well as cytokines, interleucins, growth factors, toxins, as well as receptors for

60724-2257

-4-

these ligands and also natural or genetically modified receptors for steroid hormones.

In an alternative embodiment, the nucleic acid cassette encodes a protein, polypeptide, or anti-sense RNA for the purpose of generating animal models of joint disease. The protein can be selected from the group including growth factors, tissue transplantation antigens (histocompatibility antigens), viral antigens, non-viral antigens, interferon, or toxins.

10 The pharmaceutically acceptable carriers, diluents and vehicles are generally well-known in the art.

In preferred embodiments the medicament contains the genetic material in conjunction with non-genetic material termed vehicles which comprise a solutions or suspensions. This non-genetic material can be selected from the group including sucrose, protamine, polybrene, spermidine, polylysine, other polycations, proteins, CaPO_4 precipitates, soluble or insoluble particles, or matrices for slow release of genetic material. The proteins may be selected from the group including lactoferrin, histone, natural or synthetic DNA binding proteins, natural or synthetic DNA binding compounds, viral proteins, non-viral proteins or any combinations of these. In addition, vehicles may be comprised of synthetic compounds which bind both to DNA and function as ligands for receptors on cells comprising structures of the joint.

In specific embodiments of the present invention, the nucleic acid cassette is transiently expressed by cells of the joints or is persistently expressed by cells of the joint as an episome. Alternatively, the nucleic acid cassette is stably expressed by cells of the joint.

60724-2257

-4a-

This approach can be used for the treatment of pathophysiological conditions resulting from inflammatory diseases affecting the joints, degenerative diseases of the joints, proliferative diseases of the joints, repair and
5 regeneration of essential joint structures as well as other acquired diseases of the joints. This approach can also be used to generate animal models of arthritis.

As well known in the art, the medicament may be put in a commercial package for practical use, transportation,
10 storage, etc. Such a commercial package usually carries a written matter which describes indications of the medicament or instructions about how to use the medicament.

Other and further objects, features and advantages will be apparent from the following descriptions of the
15 presently preferred embodiments in the invention which are given for the purpose of disclosure and when taken in conjunction with the accompanying drawings.

-5-

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows expression of E. coli β -galactosidase in synovial structures of the rabbit knee after DNA mediated gene transfer into the joint. Arrows indicate the regions of synovium stained with X-gal.

- A. Experimental animal after injection with CMV-B-gal.
- B. Control animal injected with different DNA construct.
- C. Close-up of experimental animal injected with CMV-B-gal.
- D. Close-up of control animal.

Figure 2 is a schematic of one proposed mechanism for DNA mediated gene transfer into synovial cells. Synovial cells form the boundary between the synovial fluid and adjacent tissues. The synovial fluid is synthesized by synovial cells and is constantly remodelled by pinocytosis. DNA injected into the joint can be taken up during the process of pinocytosis.

Figure 3 is a schematic of one proposed mechanism for DNA mediated gene transfer into inflammatory cells invading the joint. Packaging DNA in a form which is preferentially taken up by phagocytes (such as particulate DNA) can lead to preferential ingestion of DNA by phagocytosis. This can be used to alter gene expression in phagocytes or ablate these cells.

Figure 4 is a schematic of one proposed mechanism for targeting DNA to different cells such as chondrocytes and synovial cells as a complex with a vehicle that is capable of receptor mediated uptake, membrane fusion, or enhanced pinocytosis.

The drawings are not necessarily to scale and certain features of the invention may be exaggerated in scale and shown in schematic form in the interest of clarity and conciseness.

DETAILED DESCRIPTION OF THE INVENTION

It will be readily apparent to one skilled in the art that various substitutions and modification may be made to the invention

-6-

disclosed herein without departing from the scope and spirit of the invention.

5 The terms "structures comprising the joint" and "cells of the joint" refer to all of the cellular and non-cellular materials which comprise the joint and are involved in normal function of the joint or are present within the joint due to pathological conditions. These include materials associated with: the joint capsule such as, synovial membranes, synovial fluid, synovial cells; the cartilaginous components of the joint such as chondrocytes, extracellular matrix of cartilage; the bony structures such as bone, 10 periosteum of bones, periosteal cells, osteoblasts, osteoclasts; the immunological components such as inflammatory cells, lymphocytes, mast cells, monocytes, eosinophils; other cells like fibroblasts; or combinations thereof.

15 The term "transformed" as used herein refers to transient or permanent changes in the characteristics (expressed phenotype) of a cell by the mechanism of gene transfer. Genetic material is introduced into a cell in a form where it expresses a specific gene product or alters the expression or effect of endogenous gene products. 20

The term "transfection" as used herein refers to the process of introducing a DNA expression vector into a cell. Various methods of transfection are possible including microinjection, CaPO_4 precipitation, liposome fusion (e.g. lipofection) or use of a gene 25 gun.

The term "transient" as used herein relates to the introduction of genetic material into a cell to express specific proteins, peptides, or RNA etc. The introduced genetic material is not integrated into the host cell genome or replicated and is 30 accordingly eliminated from the cell over a period of time.

The term "stable" as used herein refers to the introduction of genetic material into the chromosome of the targeted cell where it integrates and becomes a permanent component of the genetic material

-7-

in that cell. Gene expression after stable transduction can permanently alter the characteristics of the cell leading to stable transformation.

5 The term "persistent" as used herein refers to the introduction of genes into the cell together with genetic elements which enable episomal (extrachromosomal) replication. This can lead to apparently stable transformation of the characteristics of the cell without the integration of the novel genetic material into the chromosome of the host cell.

10 The term "ligand" refers to a protein capable of binding to a receptor on a cell.

15 The term "nucleic acid cassette" as used herein refers to the genetic material of interest which can express a protein, or a peptide, or RNA to incorporate it transiently, permanently or episomally into the structures comprising the joint. The nucleic acid cassette is positionally and sequentially oriented in a vector with other necessary elements such that the nucleic acid in the cassette can be transcribed and, when necessary, translated in the cells of the joint.

20 The term "vector " as used herein refers to a construction comprised of genetic material designed to direct transformation of a targeted cell. A vector contains multiple genetic elements positionally and sequentially oriented with other necessary elements such that the nucleic acid in a nucleic acid cassette can be transcribed and when necessary translated in the transfected cells.
25 In the present invention the preferred vector comprises the following elements linked sequentially at appropriate distance for allowing functional expression: a promoter; a 5' mRNA leader sequence; an initiation site; a nucleic acid cassette, containing the sequence to be expressed; a 3' untranslated region; and a
30 polyadenylation signal.

The term "genetic material" as used herein refers to contiguous fragments of DNA or RNA. The genetic material which is

2139948

-8-

introduced into cells comprising essential structures of the joint according to the methods described herein can be any DNA or RNA. For example the nucleic acid can be: 1) normally found in cells of the joints, 2) normally found in cells of the joints but not expressed at physiologically appropriate levels, 3) normally found in cells of the joints but not expressed at optimal levels in certain pathological conditions, 4) novel fragments of genes normally expressed or not expressed in structures of the joint, 5) synthetic modifications of genes expressed or not expressed within structures the joints, 6) any other DNA which may be modified for expression in cells of the joint, and 7) any combination of the above.

The term "pharmacological dose" as used herein refers to a dose of vector and level of gene expression resulting from the action of the promotor on the nucleic acid cassette when introduced into the appropriate cell type which will produce sufficient protein, polypeptide, or antisense RNA to either (1) increase the level of protein production, (2) decrease or stop the production of a protein, (3) inhibit the action of a protein, (4) inhibit proliferation or accumulation of specific cell types, (5) induce proliferation or accumulation of specific cell types. As an example, if a protein is being produced which causes the accumulation of inflammatory cells within the joint, the expression of this protein can be inhibited, or the action of this protein be interfered by expression of appropriate regulatory proteins. The dose will depend on the protein being expressed, the promoter, uptake and action of the protein RNA. Given any set of parameters, one skilled in the art will be able to determine the dose.

The term "ablation agent" as used herein refers to an agent which is capable of destroying the cell in which it is present. Examples of ablation agents used in the present invention include diphtheria toxin and herpes thymidine kinase.

-9-

The term "vehicle" as used herein refers to non-genetic material combined with the vector in a solution or suspension which enhance the uptake, stability, and expression of genetic material into cells of the joint. Examples of a vehicle include: sucrose, protamine, polybrene, spermidine, polylysine, other polycations, proteins, CaPO_4 precipitates, soluble and insoluble particles, or matrices for slow release of genetic material. The proteins may be selected from the group including lactoferrin, histone, natural or synthetic DNA binding proteins, natural or synthetic DNA binding compounds, viral proteins, non-viral proteins or any combinations of these. In addition, vehicles may be comprised of synthetic compounds which bind both to DNA and function as ligands for receptors on cells comprising structures of the joint.

Examples of proteins or polypeptides which can be expressed by the vector in the transformed cells of the joint for the purposes of somatic gene therapy include enzymes, extracellular matrix elements, cytokines, interleucins, growth factors, toxins, as well as receptors for these ligands and also natural or genetically modified receptors for steroid hormones.

Examples of proteins or polypeptides which can be expressed by the vector in the transformed cells of the joint for the purposes of generating animal models of arthritis include growth factors, tissue transplantation antigens (histocompatibility antigens), viral antigens, non-viral antigens, interferon, or fragments of diphtheria toxin.

One specific embodiment of the present invention is a method of transfecting a cell in a structure of a joint comprising the step of introducing a DNA vector directly into the joint, said vector comprised of the following elements linked sequentially at appropriate distances for allowing functional gene expression: a promotor; a 5' mRNA leader sequence, an initiation site, a nucleic acid cassette containing the nucleic acid sequence to be expressed, a 3' untranslated region, and a polyadenylation signal. One skilled

2139948

-10-

in the art will readily recognize how to order and position these elements such that the nucleic acid sequence is expressed. Further, one skilled in the art will readily recognize that certain elements of this DNA vector may be eliminated for specific applications. In preferred embodiments, the DNA vector without the nucleic acid cassette will contain a restriction site for insertion of the specific cassette. Once the specific cassette is inserted, the restriction site may no longer be present.

In the present invention the nucleic acid cassette is activated with a promotor. The promotor is any of the wide variety of promotors known in the art. Examples of the promotors which are used in the present invention are the retroviral LTR promotor, cytomegalovirus promotor, dihydrofolate reductase promotor, viral promotors, or non-viral promotors. Alternatively, the promotor can be selected from those shown to specifically express in the select cell types which may be found associated with the structures of the joint such as synovial cells, fibroblasts, lymphocytes, periosteal cells, chondrocytes, osteoblasts, osteoclasts, or more than one of these cell types.

One skilled in the art will recognize that the selection of the promotor will depend on the vector, the nucleic acid cassette, the cell type to be targeted, and the desired biological effect. One skilled in the art will also recognize that in the selection of a promotor the parameters can include: achieving sufficiently high levels of gene expression to achieve a physiological effect, maintaining a critical level of gene expression, achieving temporal regulation of gene expression, achieving cell type specific expression, achieving pharmacological, endocrine, paracrine, or autocrine regulation of gene expression, and preventing inappropriate or undesirable levels of expression. Any given set of selection requirements will depend on the conditions but can be readily determined once the specific requirements are determined.

-11-

Those promoters which naturally occur in the cells comprising structures of the joint will be preferred.

Introduction of a vector into cells comprising structures of the joint can be used to treat a variety of pathological conditions. These conditions can result from the abnormal production of a protein or a polypeptide, for example too little or too much of a protein or polypeptide, the production of an abnormal protein or polypeptide, atopic production of a protein or peptide, or abnormal regulation of production of a protein or polypeptide. Introduction of a vector into cells comprising structures of the joint can also be used to provide proteins or peptides or genetic elements (DNA or RNA) with therapeutic actions. One skilled in the art will recognize that even complex pathophysiological events such as inflammatory diseases, degenerative, diseases, injury and regeneration can be understood as comprising a series of molecular interactions between proteins and can be treated according to the embodiment of this invention.

For example, in the treatment of a pathological condition the vector in a vehicle will be introduced into cells comprising structures of the joint by injecting a pharmacological dose of the vector and vehicle into a joint.

The preferred embodiments of this invention involves transient or persistent expression within the joint. This is preferable to stable expression since it enables adjustment of the level of expression in response to the evolution of the disease process.

Stable expression may presently be achieved by the use of viral vectors or the transplantation of cells which are stably transformed with viral or DNA vectors.

Specific diseases which can be treated by administration of vectors to cells within the joint include various arthritis, avascular necrosis, or injuries requiring repair and regeneration of structures comprising the joint. Various types of arthritis can be treated. A list of these various types are shown below. This list

-12-

is based on a classification of joint disorders: (after table 358-1, p. 2048 Harrison's Principles of Internal Medicine, Thorn, G. W., Adams, R. D., Braunwald, E., Isselbacher, K. J., Petersdorf, R. G., McGraw Hill Book Company, 1977)

- 5 I. Periarticular
 - A. Tendinitis
 - B. Bursitis
 - C. Fibrositis
 - D. Bone lesions
 - 10 E. Soft Tissue inflammation
- II. Articular
 - A. Cartilage and ligaments
 1. Degenerative joint disease
 2. Traumatic disorders
 - 15 3. Neuropathic arthropathy
 4. Metabolic disorders
 - B. Synovium
 1. synovial tumors
 2. pigmented villonodular synovitis
 - 20 3. hemorrhagic disorders
 4. septic disorders
 5. crystal induced disorders (gout)
 - a. Immune complex disease and vasculitis
 - b. Systemic lupus erythematosus
 - 25 c. rheumatoid arthritis
 - d. Reiter's syndrome
 - e. Psoriasis
 - f. ankylosing spondylitis
 - g. scleroderma
 - 30 h. arthritis of intestinal disease

The method of treating a pathophysiological condition or repairing and regenerating structures of the joint comprises the steps of injecting a joint of a human with a pharmacological dose of

-13-

a DNA vector in a vehicle, wherein the nucleic acid cassette in the vector encodes a protein, polypeptide or RNA. The vector is taken up by appropriate cells within the joint and expresses the protein, polypeptide or RNA. Specific embodiments of the invention include
5 treatment of disorders listed above.

Specific embodiments of the present invention include a method of ablation of inflammatory cells in a joint comprising the step of introducing a vector into the joint, said vector comprised of the following elements linked sequentially at appropriate distances for
10 allowing function expression: a promoter which is specifically expressed in leucocytes; a 5' mRNA leader sequence; an initiation site; a nucleic acid cassette containing the sequence for an ablation agent; a 3' untranslated region; and a polyadenylation signal; wherein the vector is targeted for selective uptake by
15 phagocytic cells in the joint. In the preferred embodiment, the vector is directly targeted by attaching the vector to a molecule which attaches to the surface of the phagocytic cell.

In specific embodiments of the present invention for the treatment of arthritis, a soluble receptor for cytokines can be
20 used. Examples of a soluble receptor include IL-1 or IL-6.

An alternative embodiment of the present invention includes a method of treating arthritis in humans comprising the step of injecting an inflamed joint of a human with a pharmacological dose of a DNA vector in a vehicle, wherein the nucleic acid cassette in
25 the vector encodes a sequence for a steroid receptor. This can be the normal receptor or it can be a genetically modified receptor.

When treating pathophysiological conditions or repairing or regenerating structures of the joint, it is found that a useful nucleic acid cassette encodes antisense RNA to prostaglandin
30 synthase.

Another embodiment of the present invention is a method of making an animal model for inflammatory arthritis comprising the step of injecting the joint of an animal with a functional DNA

-14-

vector in a vehicle. The nucleic acid cassette in the vector encodes a transplantation antigen, cell adhesion antigen, antigen from micro-organisms, viral antigen, synthetic antigen or recombinant antigen. Examples of the transplantation antigen include the histocompatibility antigens, Class I transplantation antigen, Class II transplantation antigen and other transplantation antigens, allogeneic transplantation antigen and xenogeneic transplantation antigen.

Another embodiment of the present invention is a transformed synovial cell comprised of nucleic acid incorporated into a synovial cell by gene transfer. In specific embodiments, the nucleic acid can be any of the genetic materials described above. Synovial cells are capable of expressing either protein, polypeptide or antisense RNA. Generally, synovial cells can express cytokines, interleucins, receptors for natural ligands, genetically modified receptors for natural ligands, inhibitors of natural ligands, steroid receptors, genetically modified steroid receptors, cell adhesion molecules, genetically modified adhesion molecules, enzymes affecting prostaglandin metabolism, enzymes involved in extracellular matrix, RNA molecules inhibiting production of transplantation antigens, cell adhesion molecules, receptors, or enzymes involved in prostaglandin metabolism. The gene in the transformed synovial cells can specifically encode a steroid receptor, IL-1, IL-6, IL-8, or soluble IL-1 receptor, the transplantation antigen or an antisense RNA to prostaglandin synthase.

Another embodiment of the present invention is a method of targeting a DNA vector to joints comprising the steps of identifying human antibodies from patients with autoimmune arthritis, cloning the antibodies to develop monoclonals with the same epitope binding determinants and coupling DNA vectors to the monoclonals; wherein the DNA/monoclonal complex is delivered to the cells of the joint. In the preferred embodiment the identified antibodies have the property of binding specifically to proteins on cells within the

-15-

joint. One skilled in the art recognizes that a variety of methods are available for identifying antibodies.

5 An alternative method of gene therapy is a method of introducing DNA into an extracellular space enclosed by a cellular membrane. For example, this method of gene therapy can be used in both humans and animals and it comprises the step of introducing a pharmacological dose of a DNA vector in a vehicle into an extracellular space in the human or animal. The extracellular space is enclosed by a cellular membrane. An example of this type of
10 treatment would be the introduction of a DNA vector into the synovium of the joint.

Another method for gene therapy in humans and animals is introduction of a pharmacological dose of DNA vector in a vehicle into a fluid space in the animal or human. This fluid space is
15 reabsorbed or remodelled by surrounding cells. Examples of this type of gene therapy would be the introduction of a DNA vector into a follicle of the thyroid, the synovium of the joint or the vitreous of the eye.

Another embodiment of the present invention is a method for
20 gene therapy in humans and animals, comprising the step of introducing a pharmacological dose of a DNA vector into a fluid of the animals or humans. The flow and removal of this fluid is by endocytosis or pinocytosis from the surrounding cells. Examples of this occurs in the thyroid follicles, synovium of the joint, and in
25 the vitreous of the eye.

Another embodiment of the present invention is a method of gene therapy in animals or humans comprising the step of introducing a pharmacological dose of DNA vector in a vehicle into a fluid space of the animal or human. The flow and removal of the fluid through
30 a membrane causes the fluid to be filtered and the cells of the membrane then uptake the filtered DNA. An example of this is in the kidney and in the central nervous system, including the spinal column and brain.

-16-

One skilled in the art will readily recognize that a variety of other tissues, fluid space and extracellular spaces may be used. As long as they meet the specific criteria of being surrounded by a cellular membrane or the uptake in removal of the fluid or the filtering of the fluid.

The following examples are offered by way of illustration and are not intended to limit the invention in any manner.

Example 1

Introduction of a Marker Gene into Cells of the Joint

A vector comprising the cytomegalovirus immediate early promotor, 5' untranslated sequences and an intron from SV40, 3' untranslated sequences from SV40 including the polyadenylation signal and a nucleic acid cassette containing the E. coli beta-galactosidase gene was combined with a vehicle containing 20% sucrose at pH 7.4. This solution was injected into the knee joint of rabbits using a transpatellar injection. As a control, an identical vector containing the E. coli chloramphenicol acetyltransferase gene in place of the beta-galactosidase gene was injected into joints on the opposite leg using identical methods. Three days after injection animals were sacrificed, the joint was disarticulated, and stained in a solution containing X-gal at pH 7.2. Under these conditions cells taking up and expressing the E. coli beta-galactosidase gene will be stained blue. Cells not expressing beta-galactosidase do not stain with this dye under these conditions.

The analysis showed that there was diffuse blue staining in regions of the joint representing the synovial cells of joint injected with the beta-galactosidase containing vector (figures 1A, 1C). No staining was observed over the tendons of bony surfaces of the joints. No blue staining was apparent in joints injected with the control vector (figures 1B, 1D).

-17-

These results demonstrate that cells of the joint are capable of taking up DNA vectors and are capable of expressing gene products encoded by the vector.

Example 2

5 Expression of anti-inflammatory factors
 to prevent inflammatory arthritis

10 The inflammatory attack on joints in animal models and human diseases may be mediated, in part, by secretion of cytokines such as IL-1 and IL-6 which stimulate the local inflammatory response. The inflammatory reaction may be modified by local secretion of soluble fragments of the receptors for these ligands. The complex between the ligand and the soluble receptor prevents the ligand from binding to the receptor which is normally resident on the surface of cells, thus preventing the stimulation of the inflammatory effect. Therapy
15 consists of the construction of a vector containing the soluble form of receptors for appropriate cytokines (for example IL-1 or IL-6) together with promoters capable of inducing high level expression in structures of the joint and a vehicle which enables efficient uptake of this vector. This DNA is injected into affected joints where the
20 secretion of an inhibitor for IL-1 such as a soluble IL-1 receptor or natural IL-1 inhibitor modifies the local inflammatory response and resulting arthritis.

 This method is useful in treating episodes of arthritis which characterize many "autoimmune" or "collagen vascular" diseases.
25 This method can also prevent disabling injury of large joints by inflammatory arthritis.

Example 3

Induction of "steroid response" by gene transfer of steroid
 receptors into cells of the joint

30 Current therapy for severe arthritis involves the administration of pharmacological agents including steroids to

2139948

-18-

depress the inflammatory response. Steroids can be administered systemically or locally by direct injection into the joint space.

5 Steroids normally function by binding to receptors within the cytoplasm of cells. Formation of the steroid-receptor complex changes the structure of the receptor so that it becomes capable of binding to specific sequences within the genome of the cell and altering the expression of specific genes. Genetic modifications of the steroid receptor can be made which enable this receptor to bind naturally occurring steroids with higher affinity, bind
10 pharmacological forms of steroid at higher affinity, or bind to synthetic steroids. Other modifications can be made to create steroid receptor which is "constitutively active" meaning that it is capable of binding to DNA and regulating gene expression in the absence of steroid altogether.

15 One approach to treating arthritis is to introduce a vector in which the nucleic acid cassette expresses a genetically modified steroid receptor into cells of the joint. Expression of a constitutively active steroid receptor within cells of the joint induces the therapeutic effects of steroids without the profound
20 systemic toxicity of these drugs. Of particular importance is the ability to target these genes differentially to specific cell types (for example synovial cells versus lymphocytes) to affect the activity of these cells.

25 Alternatively, steroid receptors which have a higher affinity for natural steroids can be introduced into the joint. These receptors exert an increased anti-inflammatory effect when stimulated by physiological concentrations of steroids or lower doses of pharmacologically administered steroids. Alternatively, constitution of a steroid receptor which binds a novel steroid
30 enables the use of drugs which would affect only cells taking up this receptor. These strategies obtain a therapeutic effect from steroids on arthritis without the profound systemic complications associated with these drugs.

-19-

Example 4

Inhibition of prostaglandin synthase

Drugs which inhibit the enzyme prostaglandin synthase are important agents in the treatment of arthritis. This is due, in part, to the important role of certain prostaglandin in stimulating the local immune response. Salicylates are widely used drugs but can be administered in limited doses which are often inadequate for severe forms of arthritis.

Gene transfer is used to inhibit the action of prostaglandin synthase specifically in affected joints by the expression of an antisense RNA for prostaglandin synthase. The complex formed between the antisense RNA and mRNA for prostaglandin synthase interferes with the proper processing and translation of this mRNA and lowers the levels of this enzyme in treated cells.

Alternatively, genes encoding enzymes which alter prostaglandin metabolism can be transferred into the joint. These have an important anti-inflammatory effect by altering the chemical composition or concentration of inflammatory prostaglandin.

Example 5

Generating an animal model of inflammatory arthritis

Inflammatory arthritis is thought to result from an autoimmune attack on cells comprising essential structures of the joint. The association of various forms of arthritis with certain tissue transplantation (Histocompatibility) antigens, suggests that these antigens, or dysregulation of these antigens, may play an important role in the genesis of these diseases.

Animal models of inflammatory arthritis can be generated by gene transfer of vectors capable of expressing heterogenic (not self) or xenogeneic (other species) transplantation antigens within the joint, thus mimicking the effect of dysregulation of tissue transplantation antigens. This induces an autoimmune attack not only on the transplantation antigens themselves, but on other

-20-

antigens within the joint which are presented to the immune system in an abnormal manner.

Alternatively, overproduction of interferon in tissues is thought to induce dysregulation of class I antigens and potentiate a inflammatory reaction. This can be modeled in animals by introduction of vectors expressing interferon into cells of the joint.

Example 6

Gene transfer to enhance repair or regeneration of joints

10 The regenerative capacity of the joint is limited by the fact that chondrocytes are not capable of remodelling and repairing cartilaginous tissues such as tendons and cartilage. Further, collagen which is produced in response to injury is of a different type -- lacking the tensile strength of normal collagen. Further, 15 the injury collagen is not remodeled effectively by available collagenase.

Gene transfer using promoters specific to chondrocytes (i.e., collagen promoters) is used to express different collagens or collagenase for the purpose of improving the restoration of function in the joints and prevent scar formation. Gene transfer into 20 fibroblasts or muscle cells in the environment of the joint is used to locally secrete growth factors such as IGF-1. The growth factors maintain muscle mass and proliferation and enhance the strength of the damaged joint.

25 Gene transfer for these purposes is affected by direct introduction of DNA into the joint space where it comes into contact with chondrocytes and synovial cells. further, the genes infiltrate into the environment of the joint where they are taken up by fibroblasts, myoblasts, and other constituents of periarticular tissue. Additionally, the gene gun can be used interoperatively or 30 via arthroscopy.

2139948

-21-

Example 7

Treatment of gouty arthritis by gene transfer

Gout is caused by the accumulation of uric acid in joints. This remains a common and painful disorder in the aging population despite medical management. Gene transfer into the joints is used to express products (for example enzymes) capable of degrading uric acid to non-toxic products. These products include urease or urate oxidase which are capable of metabolizing uric acid or urate binding globulins which render this compound more soluble and thus prevent crystalline formation within the joint.

Example 8

Persistent expression using episomal vectors

In each of the foregoing examples, transient expression of recombinant genes induces the desired biological response. In some diseases more persistent expression of recombinant genes is desirable. This is achieved by adding elements which enable extrachromosomal (episomal) replication of DNA to the structure of the vector. Vectors capable of episomal replication are maintained as extrachromosomal material and can replicate. These sequences will not be eliminated by simple degradation but will continue to be copied. Episomal vectors provide prolonged or persistent expression, though not necessarily stable or permanent, expression of recombinant genes in the joint. Persistent as opposed to stable expression is desirable to enable adjustments in the pharmacological dose of the recombinant gene product as the disease evolves over time.

Example 9

Gene delivery using the gene gun

The repair and regeneration of tissues comprising the joint following injury or surgery is enhanced by directly delivering recombinant genes to specific sites at the time of surgery. This is

-22-

done in conjunction with surgical reconstructions or arthroscopy using the "gene gun" (Yang, N.S. et al.; In vivo and in vitro gene transfer to mammalian somatic cells by particle bombardment; Proc. Natl. Acad. Sci. USA 87:9568-72 (1990)) to deliver particles coated with DNA to specific regions of the joint. The use of the gene gun and particle bound DNA enables constitution of gene expression at highly specific locations.

Example 10

Vehicles for gene delivery into cells of the joint

Initial experiments used DNA in solution for gene transfer into cells of the joint. This DNA is taken up by synovial cells during the process of these cells continually resorbing and remodeling the synovial fluid by secretion and pinocytosis. Gene delivery is enhanced by packaging DNA into lipophilic particles (for example lipofectin) which binds nonspecifically to hydrophobic membranes resulting in a fusion of the particle with the membrane and release of the DNA into the cytoplasm. Similarly, complexes of DNA and proteins which bind to receptors on the surface of cells in the joint enhances uptake and expression. Alternatively, particulate DNA complexed with CaPO_4 or polycations can be efficient substrates for phagocytosis by monocytes or other inflammatory cells.

Example 11

Ablation of inflammatory cells invading the joint.

Inflammatory cells invading the joint are ablated using vectors which contain the diphtheria toxin gene under the control of a promotor which is expressed only in leukocytes. These vectors are delivered to the joint as particles and are selectively taken up by phagocytotic cells. The uptake and expression of this vector results in expression of diphtheria toxin which is lethal to that specific cell. Alternatively, a vector which expressed herpes

thymidine kinase gene under the control of a specific promoter could be introduced into cells, and these cells are then ablated by administration of acytovin.

All patents and publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains.

One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The present examples of DNA vectors along with the methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

10

60724-2257

24

CLAIMS:

1. A medicament for in vivo transfection and expression of a nucleic acid sequence in a cell in structure of a joint, which comprises:

5 (a) a pharmaceutically acceptable carrier, diluent or vehicle and

(b) an effective amount of a non-viral DNA vector,

wherein the DNA vector comprises the following elements linked sequentially at appropriate distance for
10 allowing functional expression:

a promoter;

a 5' mRNA leader sequence;

an initiation site;

a nucleic acid sequence encoding a polypeptide that
15 affects a joint pathophysiology;

a 3' untranslated region; and

a polyadenylation signal.

2. The medicament of claim 1, which comprises a pharmaceutically acceptable the vehicle selected from the group
20 consisting of sucrose, protamine, polybrene, polylysine, polycations, proteins, CaPO_4 , spermidine, soluble or insoluble particles, and matrices for slow release.

3. A medicament for in vivo transfection and expression of a nucleic acid sequence in a cell in structure of a joint,
25 which comprises:

60724-2257

25

(a) a pharmaceutically acceptable carrier, diluent or vehicle, and

(b) an effective amount of a non-viral vector,

wherein the vector comprises the following elements
5 linked sequentially at appropriate distances for allowing functional expression:

a promoter that is specifically expressed in leukocytes;

a 5' mRNA leader sequence;

10 an initiation site;

a nucleic acid sequence encoding an agent for ablation of inflammatory cells in a joint;

a 3' untranslated region; and

a polyadenylation signal.

15 4. The medicament of claim 3, wherein the vector is selectively targeted for selective uptake by phagocytic cells in the joint by combining the vector with a molecule able to attach to a phagocytic cell surface.

5. The medicament of claim 3 or 4, wherein the ablation
20 agent is diphtheria toxin.

6. The medicament of claim 3, 4 or 5, wherein the vector is combined with a protein having trophic properties.

7. A medicament for direct injection into a mammalian joint in vivo which comprises:

25 (a) a pharmaceutically acceptable carrier, diluent or vehicle, and

60724-2257

26

(b) an effective amount of a non-viral DNA vector,

wherein the DNA vector comprises a nucleic acid cassette encoding a protein, polypeptide or RNA involved in a pathophysiologic condition of a mammalian joint.

5 8. The medicament of claim 7, wherein the pathophysiologic condition is selected from the group consisting of periarticular arthritis, articular arthritis of cartilage and ligaments and articular arthritis of synovium.

9. The medicament of claim 7 or 8, wherein the nucleic
10 acid cassette encodes a biological molecule selected from the group consisting of cytokines, interleukins, receptors for natural ligands, genetically modified receptors for natural ligands, inhibitors of natural ligands, steroid receptors, genetically modified steroid receptors, cell adhesion
15 molecules, genetically modified adhesion molecules, enzymes affecting prostaglandin metabolism, enzymes involved in extracellular matrix, RNA molecules inhibiting production of transplantation antigens, cell adhesion molecules, receptors involved in prostaglandin metabolism and enzymes involved in
20 prostaglandin metabolism.

10. The medicament of claim 9, wherein the biological molecule is selected from the group consisting of a steroid receptor, IL-1, IL-6, a transplantation antigen and an antisense RNA to prostaglandin synthase.

25 11. A medicament for injection into a joint of a mammal having a pathophysiologic condition of the joint, which comprises:

(a) a pharmaceutically acceptable carrier, diluent or vehicle, and

30 (b) an effective amount of a non-viral DNA vector,

60724-2257

27

wherein the DNA vector comprises a nucleic acid cassette encoding a protein or polypeptide, or RNA respective thereto, involved in repairing and regenerating a structure of the joint.

5 12. The medicament of claim 11, wherein the pathophysiologic condition is selected from the group consisting of periarticular arthritis, articular arthritis of cartilage and ligaments and articular arthritis of synovium.

10 13. The medicament of claim 11 or 12, wherein the protein or polypeptide, or RNA respective thereto, involved in repairing and regenerating a structure of the joint is selected from the group consisting of collagens, growth factors, extracellular matrix proteins, inhibitory growth factors, enzymes involved in modifying extracellular matrix and enzymes
15 involved in remodelling extracellular matrix.

14. A medicament for in vivo transfection and expression of a nucleic acid sequence in a cell in structure of a mammalian joint, which comprises:

(a) a pharmaceutically acceptable vehicle, and

20 (b) an effective amount of a non-viral DNA vector,

wherein a nucleic acid cassette in the vector encodes the sequence for a soluble form of a receptor, wherein the receptor is involved in a pathophysiology of arthritis in a mammal.

25 15. The medicament of claim 14, wherein the soluble receptor is a cytokine receptor.

16. The medicament of claim 15, wherein the cytokine receptor is specific for IL-1 or IL-6.

60724-2257

28

17. A method of making a mammalian model for inflammatory arthritis comprising the step of injecting a joint of a mammal with a non-viral DNA vector;

wherein a nucleic acid cassette in the vector encodes
5 a cytokine or an antigen selected from the group consisting of a transplantation antigen, cell adhesion antigen, antigen from a micro-organism, viral antigen, synthetic antigen and recombinant antigens.

18. The method of claim 17, wherein the transplantation
10 antigen is selected from the group consisting of histocompatibility antigen, allogeneic transplantation antigen, and xenogeneic transplantation antigen.

19. The method of claim 17 or 18, wherein the cytokine is an interferon.

15 20. A medicament for injection into a joint of a human having gouty arthritis which comprises:

(a) a pharmaceutically acceptable carrier, diluent or vehicle, and

(b) an effective amount of a non-viral DNA vector,

20 wherein the DNA vector comprises a nucleic acid cassette encoding and expressing a product capable of degrading uric acid.

21. The medicament of claim 20, wherein the cassette encodes the nucleic acid sequence for urease or urate oxidase.

25 22. A medicament for injection into an extracellular space in a mammal which comprises:

(a) a pharmaceutically acceptable carrier, diluent or vehicle, and

60724-2257

29

(b) an effective amount of a non-viral DNA vector,

wherein the extracellular space is enclosed by a cellular membrane and wherein the DNA vector comprises a nucleic acid cassette encoding a therapeutic protein,
5 polypeptide or RNA.

23. The medicament of claim 22, wherein the extracellular space is a synovium of a joint.

24. A medicament for injection into a fluid space in a mammal which comprises:

10 (a) a pharmaceutically acceptable carrier, diluent or vehicle, and

(b) an effective amount of a non-viral DNA vector,

wherein the fluid space is reabsorbed or remodeled by surrounding cells and wherein the DNA vector comprises a
15 nucleic acid cassette encoding a therapeutic protein, polypeptide or RNA.

25. The medicament of claim 24, wherein the fluid space is selected from the group consisting of a thyroid follicle, a joint synovium and a vitreous of an eye.

26. The medicament according to any one of claims 1 to 16 or any one of claims 20 to 25, which is contained in a commercial package carrying a written matter which describes
20 instructions for direct injection of the medicament into the joint.

60724-2257

30

27. The medicine according to claim 26, wherein the
written matter describes that the medicament is to be used for
treatment of a pathophysiological condition resulting from an
inflammatory disease affecting the joint, a degenerative
5 disease of the joint, a proliferative disease of the joint or
for repair or regeneration of an essential joint structure.

SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS

2139948

1/6




Figure 1A

SUBSTITUTE SHEET

2139948

2/6



Figure 1B

SUBSTITUTE SHEET

2139948

3/6



Figure 1C

2139948

4/6




Figure 1D

SUBSTITUTE SHEET

2139948

5/6




Figure 2




Figure 3

SUBSTITUTE SHEET

2139948

6/6




Figure 4

SUBSTITUTE SHEET