



(86) **Date de dépôt PCT/PCT Filing Date:** 2017/08/16  
 (87) **Date publication PCT/PCT Publication Date:** 2018/02/22  
 (45) **Date de délivrance/Issue Date:** 2024/01/23  
 (85) **Entrée phase nationale/National Entry:** 2019/02/01  
 (86) **N° demande PCT/PCT Application No.:** JP 2017/030003  
 (87) **N° publication PCT/PCT Publication No.:** 2018/034356  
 (30) **Priorité/Priority:** 2016/08/16 (JP2016-159743)

(51) **Cl.Int./Int.Cl. C07K 7/06** (2006.01),  
**A61K 38/03** (2006.01), **A61K 45/00** (2006.01),  
**A61K 49/04** (2006.01), **A61K 49/06** (2006.01),  
**A61K 49/14** (2006.01), **A61P 35/00** (2006.01),  
**C07K 14/00** (2006.01), **C07K 19/00** (2006.01),  
**C12N 15/09** (2006.01)

(72) **Inventeurs/Inventors:**  
 FUKUDA, MICHIKO, JP;  
 NONAKA, MOTOHIRO, JP

(73) **Propriétaire/Owner:**  
 NATIONAL INSTITUTE OF ADVANCED INDUSTRIAL  
 SCIENCE AND TECHNOLOGY, JP

(74) **Agent:** SMART & BIGGAR LP

(54) **Titre : PEPTIDE CIBLE DE TUMEUR MALIGNNE**  
 (54) **Title: MALIGNANT TUMOR TARGET PEPTIDE**

(57) **Abrégé/Abstract:**

The present invention provides a peptide comprising the amino acid sequence of any of the formulas (I)-(III) below: (I) an amino acid sequence of (X<sub>1</sub>)[D]P[D](X<sub>2</sub>)[D] wherein X<sub>1</sub> is W or F, X<sub>2</sub> is S or T, and each amino acid symbol immediately followed by symbol [D] is a D form of the amino acid, (II) an amino acid sequence of P[D]T[D](X)<sub>n</sub>F[D] wherein (X)<sub>n</sub> is any amino acid in the number of n selected independently of each other, n is an integer of 0-4, and the symbol [D] is as defined above, (III) an amino acid sequence that is a Retro-inverso of the amino acid sequence of any of the aforementioned (I) and the aforementioned (II); and a conjugate containing the peptide and a functional part.

## Abstract

The present invention provides a peptide comprising the amino acid sequence of any of the formulas (I)-(III) below:

5 (I) an amino acid sequence of (X1)[D]P[D](X2)[D] wherein X1 is W or F, X2 is S or T, and each amino acid symbol immediately followed by symbol [D] is a D form of the amino acid,

(II) an amino acid sequence of P[D]T[D](X)<sub>n</sub>F[D] wherein (X)<sub>n</sub> is any amino acid in the number of n selected independently of  
10 each other, n is an integer of 0-4, and the symbol [D] is as defined above,

(III) an amino acid sequence that is a Retro-inverso of the amino acid sequence of any of the aforementioned (I) and the aforementioned (II);

15 and a conjugate containing the peptide and a functional part.

## DESCRIPTION

Title of the Invention: MALIGNANT TUMOR TARGET PEPTIDE

[Technical Field]

[0001]

5       The present invention generally relates to the field of cancer biology, and more particularly to peptides that bind to annexin A1 and use thereof.

[Background Art]

[0002]

10       Among brain tumors, many tumors originating from meninges, brain or spinal nerve, such as meningiomas and schwannomas, are benign brain tumors and can be completely cured if they can be removed by surgery. In contrast, neuroepithelial tumors including glioma are basically malignant brain tumors.

15       Especially, grade 4 glioblastoma shows extremely poor prognosis (5-year survival rate is about 10%) even if radiation therapy and chemotherapy are performed after removal by craniotomy. A major reason for the lack of effective chemotherapy for brain tumor is the presence of a blood-brain barrier. On the other  
20       hand, temozolomide, which is a new drug effective for glioblastoma, was approved in Japan in 2006. This drug has a very small molecular weight (194 Da) and was clarified to be able to cross the blood-brain barrier by diffusion. However, the drug has only been able to extend life expectancy from 12  
25       months to 16 months. To dramatically improve the treatment results for malignant brain tumor in the future, therefore, it is essential to develop a therapeutic drug that accumulates in the brain tumor at an overwhelmingly high concentration and can actively cross the wall of vascular endothelium, rather than a  
30       passive method relying on the diffusion of small molecules as for the blood-brain barrier.

[0003]

      Recently, the economic burden on patients is becoming increasingly heavy due to the rising research and development  
35       expenses and the widespread use of antibody pharmaceuticals.

In the future, this trend is considered to be further accelerated by the sophistication of medical care such as diagnosis system by genetic information and the like. In addition, the impact of these soaring drug prices on national health care costs is also immeasurable. Furthermore, expensive antibody pharmaceuticals become inaccessible to underdeveloped countries, leading to further medical inequality on a global scale. If measures are to be taken for the above-mentioned problems from a longer-term perspective, it is necessary to explore the possibility of inexpensive biopharmaceuticals such as short-chain peptides in the future in addition to the pursuit of a superior drug seed.

[0004]

Fukuda, one of the present inventors, hosted a laboratory in Sanford Burnham Prebys Medical Discovery Institute in the United States for more than 30 years until she was assigned to the National Institute of Advanced Industrial Science and Technology in 2014. During that time, Fukuda succeeded for the first time in the world in inhibiting sugar chain-dependent cancer metastasis by using a peptide mimicking the structure of sugar chain (non-patent document 1). Furthermore, in the process of examining vascular endothelial receptors interacting with these sugar chain mimetic peptides, she found that a peptide named IF7 binds to annexin A1 (Anxa1) (non-patent document 2). Anxa1 was clarified by the group of Jan Schnitzer et al. to be the most specific of the currently known tumor vessel specific marker molecules and is expressed intracellularly in normal cells but strongly expressed on luminal surfaces adjacent to the bloodstream in tumor neovascular endothelial cells (non-patent document 3). Fukuda et al. clarified that a drug (IF7-SN38) obtained by binding an anticancer agent (SN38) to IF7 clears tumor in cancer-bearing mice at a low dose (non-patent document 2). While IF7 binds to the N-terminal region of mouse Anxa1, the amino acid sequence in that region is highly conserved between mouse and human.

Therefore, IF7-SN38 may also be useful in human. This finding was taken up in the "In this issue in PNAS" of the PNAS journal and the NIH introduced a special article. The finding further attracted considerable attention, including reports by media  
 5 around the world.

[Document List]

[non-patent documents]

[0005]

non-patent document 1: Fukuda et al., Cancer Research, 60: 450-  
 10 6, 2000

non-patent document 2: Hatakeyama et al., Proc. Natl. Acad. Sci USA, 108: 19587-92, 2011

non-patent document 3: Oh et al., Nature 429: 629-35, 2004

[SUMMARY OF THE INVENTION]

15 [Problems to be Solved by the Invention]

[0006]

When IF7 is administered intravenously to cancer-bearing mice, it reaches the blood vessels surrounding the tumor, after which IF7 is taken up into vesicles from the luminal side of  
 20 vascular endothelial cells and moves to the basal side, where it is released into the stroma where cancer cells are present. Also in mouse vascular endothelial F2 cells, IF7 bound to Anxal actively traversed tumor vascular endothelial cells by transcytosis. Therefore, it was suggested that IF7 has an  
 25 activity to cross the blood vessel-brain tumor barrier in the brain tumor. Surprisingly, intravenous injection of fluorescent-labeled IF7 into brain tumor model mice transplanted with glioma cells resulted in accumulation of fluorescence at a high concentration in the tumor site in the  
 30 brain. Fluorescence also crossed blood vessels and reached cancer cells in the brain stroma. Furthermore, subcutaneous tumor and brain tumor were created in the same mouse (dual tumor model), and the therapeutic effect of IF7-SN38 was tested. As a result, both brain tumor and subcutaneous tumor were  
 35 suppressed, and the effect thereof was higher in the brain

tumor than in the subcutaneous tumor. Similar results were obtained in B16 melanoma metastatic brain tumor model, and the results remained same even when the host mouse was changed to C57BL/6 or SCID strain. These facts indicate that DDS  
5 targeting Anxal not only efficiently cross the blood-brain tumor barrier but also provide excellent therapeutic effects in brain tumor.

[0007]

As described above, while IF7 has an excellent malignancy  
10 targeting activity, it has two problems for clinical development: low solubility and low stability. First, IF7 peptide is protease sensitive and easily degraded. In fact, in an experiment including intravenous administration of A488-IF7 to healthy mice, the fluorescence signal of the peripheral  
15 blood almost disappeared in about 1 hour. However, by tail vein administration to cancer-bearing mice, fluorescence of A488-IF7 recovered from the peripheral blood was significantly less than in healthy mice. This result suggests that most of A488-IF7 accumulates rapidly in the tumor while undergoing  
20 degradation. Also, IF7 is highly hydrophobic and causes difficulty in dosage form, for example, when a compound bound with an anticancer agent is administered intravenously, a surfactant needs to be added to avoid precipitation after dissolution in DMSO.

25 [0008]

Therefore, it is an object of the present invention to provide a novel peptide that binds to annexin A1 and use thereof capable of at least partially overcoming the above-mentioned problems associated with IF7.

30 [Means of Solving the Problems]

[0009]

To solve the above-mentioned problems, the present inventors conducted intensive studies and had an idea of exploring a D-form peptide capable of binding to annexin A1.  
35 To this end, the present inventors synthesized a D-form peptide

(D-MC16) in mirror image relation with 16-residue L-form peptide (L-MC16) having a modifying Cys residue added to the N-terminal 15 residues of Anxal known to bind to IF7, and screened using a T7 phage library to identify a plurality of 7-  
 5 amino acid L-form peptides that bind to D-MC16. They conducted further studies of a peptide called TIT7 peptide, which was particularly promising among the identified L-form peptides. As the inventors predicted, they confirmed that the D-form peptide (dTIT7 peptide), which is in mirror image relation with  
 10 the TIT7 peptide, binds to Anxal. Furthermore, the present inventors have demonstrated by in vivo experiments that dTIT7 peptides accumulate in the tumor site when administered to brain tumor model mice via the tail vein, that dTIT7 peptide (GA-dTIT7) bound with an anticancer agent (geldanamycin)  
 15 markedly inhibited tumor growth when intravenously administered to cancer-bearing mice, thus causing massive necrosis in the tumor site, and that oral administration of GA-dTIT7 suppressed tumor growth in both brain tumor model mice transplanted with C6 cells into the brain and brain metastasis model mice  
 20 transplanted with B16 cells into the brain, the tumor continued to reduce even after cessation of drug administration, and complete cure was finally achieved in some animals.

[0010]

The present inventors have conducted further studies  
 25 based on the above results, and have completed the present invention. The present invention is as follows.

[1] A peptide comprising the amino acid sequence of any of the formulas (I)-(III) below:

- (I) an amino acid sequence of (X1) [D]P[D] (X2) [D] wherein X1 is  
 30 W or F, X2 is S or T, and each amino acid symbol immediately followed by symbol [D] is a D form of the amino acid,  
 (II) an amino acid sequence of P[D]T[D] (X)<sub>n</sub> F[D] wherein (X)<sub>n</sub> is any amino acid in the number of n selected independently of each other, n is an integer of 0-4, and the symbol [D] is as  
 35 defined above,

(III) an amino acid sequence that is a Retro-inverso of the amino acid sequence of any of the aforementioned (I) and the aforementioned (II).

[2] A peptide comprising the amino acid sequence of any of the following (i)-(vii) (in the following sequences, symbol [D] is as defined above):

(i) an amino acid sequence of T[D] I[D] T[D] W[D] P[D] T[D] M[D],

(ii) an amino acid sequence of L[D] R[D] F[D] P[D] T[D] V[D] L[D],

(iii) an amino acid sequence of L[D] L[D] S[D] W[D] P[D] S[D] A[D],

(iv) an amino acid sequence of S[D] P[D] T[D] S[D] L[D] L[D] F[D],

(v) an amino acid sequence of M[D] P[D] T[D] L[D] T[D] F[D] R[D],

(vi) an amino acid sequence of any of the aforementioned (i)-(v), in which 1 or several amino acids are inserted, substituted or deleted, or these are combined,

(vii) an amino acid sequence that is a Retro-inverso of the amino acid sequence of any of the aforementioned (i)-(vi).

[3] The peptide of [2], comprising the amino acid sequence of the above-mentioned (i).

[4] The peptide of [2], consisting of the amino acid sequence of the above-mentioned (i).

[5] A conjugate comprising the peptide of any of [1] to [4], and one or more components.

[6] The conjugate of [5], wherein the one or more components comprise an anticancer agent.

[7] The conjugate of [6], wherein the above-mentioned anticancer agent is selected from the group consisting of an antimetabolite, an alkylating agent, an anticancer antibiotic, a microtubule inhibitor, a platinum preparation, a topoisomerase inhibitor, a molecular targeting agent, and an anti-angiogenic agent.

[8] The conjugate of [6] or [7], which is selected from the group consisting of  
enocitabine, capecitabine, capecitabine, carmofour, cladribine, gemcitabine, cytarabine, cytarabine ocfosphate, tegafur, tegafur/uracil,  
5 tegafur/Gimeracil/Oteracil potassium, doxifluridine, nelarabine, hydroxycarbamide, fluorouracil, fludarabine, pemetrexed, pentostatin, mercaptopurine, methotrexate; cyclophosphamide, ifosfamide, melphalan, busulfan, thiotepa, nimustine, ranimustine, dacarbazine, procarbazine, temozolomide,  
10 carmustine, streptozotocin, bendamustine; actinomycin D, aclarubicin, amrubicin, idarubicin, epirubicin, zinostatin stimalamer, daunorubicin, doxorubicin, pirarubicin, bleomycin, peplomycin, mitomycin C, mitoxantrone, liposomal doxorubicin;  
15 vinblastine, vincristine, vindesine, paclitaxel, docetaxel, oxaliplatin, carboplatin, cisplatin, nedaplatin; camptothecin, irinotecan, nogitecan, SN-38, doxorubicin, etoposide, levofloxacin, ciprofloxacin; legolafenib, cetuximab, panitumumab, ramsilmab, gefitinib,  
20 erlotinib, afatinib, crizotinib, alectinib, ceritinib, libertinib, trastuzumab, lapatinib, pertuzumab, sunitinib, sorafenib, axitinib, pazopanib, nivolumab, pembrolizumab, ipilimumab, vemurafenib, everolimus, temsirolimus, rituximab, bevacizumab, geldanamycin;  
25 angiostatin, endostatin, metastatin, anti-VEGF antibody and VEGFR-2 inhibitor.

[9] The conjugate of [5], wherein the one or more components comprise a detectable substance.

[10] The conjugate of [9], wherein the detectable substance  
30 enables detection of the conjugate in vivo by a means selected from the group consisting of X-ray photography, computed tomography (CT), nuclear magnetic resonance imaging (MRI), ultrasonography, scintigraphy, positron emission tomography (PET), endoscopy and laparoscopy.

35 [11] The conjugate of [9] or [10], wherein the detectable

substance is a radioisotope, an MRI enhancer, a radiopaque substance, a contrast agent or a fluorescent substance.

[12] The conjugate of any of [9] to [11], wherein the detectable substance is selected from the group consisting of  
 5 a radioactive nuclide selected from  $^{18}\text{F}$ ,  $^{51}\text{Mn}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{52}\text{Fe}$ ,  $^{55}\text{Co}$ ,  
 $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$ ,  $^{82\text{m}}\text{Rb}$ ,  $^{83}\text{Sr}$ ,  $^{86}\text{Y}$ ,  $^{89}\text{Zr}$ ,  $^{94\text{m}}\text{Tc}$ ,  $^{110}\text{In}$ ,  
 $^{120}\text{I}$ ,  $^{124}\text{I}$  and the like. Examples of the radioactive nuclide  
 useful for detection of  $\gamma$  ray include  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{67}\text{Cu}$ ,  
 $^{67}\text{Ga}$ ,  $^{75}\text{Se}$ ,  $^{97}\text{Ru}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{114\text{m}}\text{In}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{169}\text{Yb}$ ,  $^{197}\text{Hg}$   
 10 and  $^{201}\text{Tl}$ ;

a paramagnetic ion selected from chromium (III), manganese (II),  
 iron(III), iron(II), cobalt(II), nickel(II), copper(II),  
 neodymium (III), samarium (III), ytterbium (III), gadolinium  
 (III), vanadium (II), terbium (III), dysprosium (III), holmium  
 15 (III) and erbium (III),

lanthanum (III), gold (III), lead (II), bismuth (III);

an iodine compound, a barium compound, a gallium compound, a  
 thallium compound;

rhodamine, fluorescein, Cy dye, Alexa (registered trademark)  
 20 Fluor, phycoerythrin (PE), allophycocyanin (APC), derivatives  
 thereof and a near infrared fluorescent reagent.

[13] A composition comprising the peptide of any of [1] to [4]  
 or the conjugate of any of [5] to [12], and a pharmacologically  
 acceptable carrier.

25 [14] A composition for the treatment of cancer, comprising the  
 conjugate of any of [6] to [8], and a pharmacologically  
 acceptable carrier.

[15] A composition for testing for cancer, comprising the  
 conjugate of any of [9] to [12], and a pharmacologically  
 30 acceptable carrier.

[16] The composition of [14] or [15], wherein the cancer is a  
 solid cancer or a liquid cancer.

[17] The composition of [16], wherein the solid cancer is an  
 angiogenic solid cancer.

35 [18] The composition of [16] or [17], wherein the solid cancer

is cancer of the brain or nervous system, cancer of the head and neck, cancer of the digestive tract, cancer of the urinary or reproductive organ, cancer of the respiratory system, cancer of the breast, cancer of the skin, cancer of the bone, or cancer of the muscle.

[19] The composition of any of [16] to [18], wherein the solid cancer is brain tumor, spinal cord tumor, laryngeal cancer, oral cancer, salivary gland cancer, paranasal sinus cancer, thyroid cancer, stomach cancer, esophageal cancer, small intestine cancer, colon cancer, rectal cancer, anal cancer, liver cancer, biliary tract cancer, pancreatic cancer, kidney cancer, kidney cell cancer, bladder cancer, prostate cancer, renal pelvis and ureter cancer, gall bladder cancer, bile duct cancer, testis cancer, penile cancer, uterine cancer, endometrial cancer, uterine sarcoma, cervical cancer, vaginal cancer, vulvar cancer, ovarian cancer, fallopian tube cancer, lung cancer, breast cancer, malignant melanoma, osteosarcoma or rhabdomyosarcoma.

[20] The composition of [19], wherein the solid cancer is a benign or malignant brain tumor.

[21] The composition of [20], wherein the brain tumor is a primary brain tumor or a metastatic brain tumor.

[22] The composition of [20] or [21], wherein the brain tumor is meningioma, pituitary adenoma, schwannoma, astrocytoma, oligodendroglioma, anaplastic astrocytoma, anaplastic oligodendroglioma, anaplastic oligoastrocytoma or glioblastoma.

[23] The composition of [16], wherein the liquid cancer is B-cell lymphoma.

[0010A]

The present invention as claimed relates to:

[1] A peptide comprising the amino acid sequence T[D] I[D] T[D] W[D] P[D] T[D] M[D], wherein each amino acid symbol immediately

followed by symbol [D] is a D form of the amino acid;

[2] A peptide consisting of the amino acid sequence T[D] I[D] T[D] W[D] P[D] T[D] M[D], wherein each amino acid symbol immediately followed by symbol [D] is a D form of the amino acid;

[3] A conjugate comprising the peptide according to [1] or [2], and one or more components;

[4] The conjugate according to [3], wherein the one or more components comprise an anticancer agent;

[5] The conjugate according to [4], wherein the anticancer agent is selected from the group consisting of an antimetabolite, an alkylating agent, an anticancer antibiotic, a microtubule inhibitor, a platinum preparation, a topoisomerase inhibitor, a molecular targeting agent, and an anti-angiogenic agent;

[6] The conjugate according to [4] or [5], wherein the anticancer agent is selected from the group consisting of

enocitabine, capecitabine, carmofour, cladribine, gemcitabine, cytarabine, cytarabine ocfosphate, tegafur, tegafur/uracil, tegafur/Gimeracil/Oteracil potassium, doxifluridine, nelarabine, hydroxycarbamide, fluorouracil, fludarabine, pemetrexed, pentostatin, mercaptopurine, methotrexate;

cyclophosphamide, ifosfamide, melphalan, busulfan, thiotepa, nimustine, ranimustine, dacarbazine, procarbazine, temozolomide, carmustine, streptozotocin, bendamustine;

actinomycin D, aclarubicin, amrubicin, idarubicin, epirubicin, zinostatin stimalamer, daunorubicin, doxorubicin, pirarubicin, bleomycin, peplomycin, mitomycin C, mitoxantrone, liposomal doxorubicin;

vinblastine, vincristine, vindesine, paclitaxel, docetaxel, oxaliplatin, carboplatin, cisplatin, nedaplatin;

camptothecin, irinotecan, nogitecan, SN-38, etoposide, levofloxacin, ciprofloxacin;

legolafenib, cetuximab, panitumumab, ramsilmab, gefitinib, erlotinib, afatinib, crizotinib, alectinib, ceritinib, libertinib, trastuzumab, lapatinib, pertuzumab, sunitinib, sorafenib, axitinib, pazopanib, nivolumab, pembrolizumab, ipilimumab, vemrafenib, everolimus, temsirolimus, rituximab, bevacizumab, geldanamycin;  
angiostatin, endostatin, metastatin, anti-VEGF antibody and VEGFR-2 inhibitor;

[7] The conjugate according to [3], wherein the one or more components comprise a detectable substance;

[8] The conjugate according to [7], wherein the detectable substance enables detection of the conjugate in vivo by a means selected from the group consisting of X-ray photography, computed tomography (CT), nuclear magnetic resonance imaging (MRI), ultrasonography, scintigraphy, positron emission tomography (PET), endoscopy and laparoscopy;

[9] The conjugate according to [7] or [8], wherein the detectable substance is a radioisotope, an MRI enhancer, a radiopaque substance, a contrast agent or a fluorescent substance;

[10] The conjugate according to any one of [7] to [9], wherein the detectable substance is selected from the group consisting of a radioactive nuclide selected from  $^{18}\text{F}$ ,  $^{51}\text{Mn}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{52}\text{Fe}$ ,  $^{55}\text{Co}$ ,  $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$ ,  $^{82\text{m}}\text{Rb}$ ,  $^{83}\text{Sr}$ ,  $^{86}\text{Y}$ ,  $^{89}\text{Zr}$ ,  $^{94\text{m}}\text{Tc}$ ,  $^{110}\text{In}$ ,  $^{120}\text{I}$ ,  $^{124}\text{I}$ ,  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{67}\text{Cu}$ ,  $^{67}\text{Ga}$ ,  $^{75}\text{Se}$ ,  $^{97}\text{Ru}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{114\text{m}}\text{In}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{169}\text{Yb}$ ,  $^{197}\text{Hg}$  and  $^{201}\text{Tl}$ ;

a paramagnetic ion selected from chromium (III), manganese (II), iron(III), iron(II), cobalt(II), nickel(II), copper(II), neodymium (III), samarium (III), ytterbium (III), gadolinium (III), vanadium (II), terbium (III), dysprosium (III), holmium (III) and erbium (III), lanthanum (III), gold (III), lead (II), bismuth (III);

an iodine compound, a barium compound, a gallium compound, a thallium compound;

rhodamine, fluorescein, Cy dye, Alexa (registered trademark) Fluor, phycoerythrin (PE), allophycocyanin (APC), derivatives thereof and a near infrared fluorescent reagent;

[11] A composition comprising the peptide according to [1] or [2] or the conjugate according to any one of [3] to [10], and a pharmacologically acceptable carrier;

[12] A composition for use in the treatment of cancer, comprising the conjugate according to any one of [4] to [6], and a pharmacologically acceptable carrier;

[13] A composition for use in testing for cancer, comprising the conjugate according to any one of [7] to [10], and a pharmacologically acceptable carrier;

[14] The composition for use according to [12] or [13], wherein the cancer is a solid cancer or a liquid cancer;

[15] The composition for use according to [14], wherein the solid cancer is an angiogenic solid cancer;

[16] The composition for use according to [14] or [15], wherein the solid cancer is cancer of the brain or nervous system, cancer of the head and neck, cancer of the digestive tract, cancer of the urinary or reproductive organ, cancer of the respiratory system, cancer of the breast, cancer of the skin, cancer of the bone, or cancer of the muscle;

[17] The composition for use according to any one of [14] to [16], wherein the solid cancer is brain tumor, spinal cord tumor, laryngeal cancer, oral cancer, salivary gland cancer, paranasal sinus cancer, thyroid cancer, stomach cancer, esophageal cancer, small intestine cancer, colon cancer, rectal cancer, anal cancer, liver cancer, biliary tract cancer, pancreatic cancer, kidney cancer, kidney cell cancer, bladder cancer, prostate cancer,

renal pelvis and ureter cancer, gall bladder cancer, bile duct cancer, testis cancer, penile cancer, uterine cancer, endometrial cancer, uterine sarcoma, cervical cancer, vaginal cancer, vulvar cancer, ovarian cancer, fallopian tube cancer, lung cancer, breast cancer, malignant melanoma, osteosarcoma or rhabdomyosarcoma;

[18] The composition for use according to [17], wherein the solid cancer is a benign or malignant brain tumor;

[19] The composition for use according to [18], wherein the brain tumor is a primary brain tumor or a metastatic brain tumor;

[20] The composition for use according to [18] or [19], wherein the brain tumor is meningioma, pituitary adenoma, schwannoma, astrocytoma, oligodendroglioma, anaplastic astrocytoma, anaplastic oligodendroglioma, anaplastic oligoastrocytoma or glioblastoma;

[21] The composition for use according to [14], wherein the liquid cancer is B-cell lymphoma;

[22] Use of the conjugate according to any one of [4] to [6] for the treatment of cancer;

[23] The use according to [22], wherein the cancer is a solid cancer or a liquid cancer;

[24] The use according to [23], wherein the solid cancer is an angiogenic solid cancer;

[25] The use according to [23] or [24], wherein the solid cancer is cancer of the brain or nervous system, cancer of the head and neck, cancer of the digestive tract, cancer of the urinary or reproductive organ, cancer of the respiratory system, cancer of the breast, cancer of the skin, cancer of the bone, or cancer of the muscle;

[26] The use according to any one of [23] to [25], wherein the solid cancer is brain tumor, spinal cord tumor, laryngeal cancer,

oral cancer, salivary gland cancer, paranasal sinus cancer, thyroid cancer, stomach cancer, esophageal cancer, small intestine cancer, colon cancer, rectal cancer, anal cancer, liver cancer, biliary tract cancer, pancreatic cancer, kidney cancer, kidney cell cancer, bladder cancer, prostate cancer, renal pelvis and ureter cancer, gall bladder cancer, bile duct cancer, testis cancer, penile cancer, uterine cancer, endometrial cancer, uterine sarcoma, cervical cancer, vaginal cancer, vulvar cancer, ovarian cancer, fallopian tube cancer, lung cancer, breast cancer, malignant melanoma, osteosarcoma or rhabdomyosarcoma;

[27] The use according to [26], wherein the solid cancer is a benign or malignant brain tumor;

[28] The use according to [27], wherein the brain tumor is a primary brain tumor or a metastatic brain tumor;

[29] The use according to [27] or [28], wherein the brain tumor is meningioma, pituitary adenoma, schwannoma, astrocytoma, oligodendroglioma, anaplastic astrocytoma, anaplastic oligodendroglioma, anaplastic oligoastrocytoma or glioblastoma;

[30] The use according to [23], wherein the liquid cancer is B-cell lymphoma.

[Effect of the Invention]

[0011]

According to the present invention, a novel peptide that binds to annexin A1 is provided.

As evident in the case of erythropoietin and recent anti-PD-1 antibodies, clinically successful pharmaceuticals have a very high specificity of expression of the biomolecule targeted

by drugs. Since annexin A1 targeted by the peptide of the present invention is the most specific molecule among the currently known neovascular target markers of malignant tumor, an anticancer agent in which the peptide of the present  
5 invention is combined may exhibit a superior therapeutic effect compared to the existing anticancer agents.

In particular, the peptide-anticancer agent conjugate is not only capable of accumulating an anticancer agent in malignant tumor with high efficiency but is also protease  
10 resistant, thus allowing for a reduction in the number of effective doses and even lower doses. In addition, complete cure of the malignant tumor may be expected by the combined use of an anticancer agent and immunotherapy, since the immune system is considered to be preserved.

15 The characteristic feature of annexin A1 is that it is expressed on the blood side of neovascular endothelial cell formed in a tumor, and when bound on the blood side with a ligand such as the peptide of the present invention, the ligand is transported to the basal side by transcytosis and is  
20 actively released into the stroma where cancer cells are present. This property may also serve as a mechanism to actively cross the vascular brain barrier. The peptide of the present invention targeting annexin A1, unlike conventional angiogenesis inhibitors (Avastin) and temozolomide (described  
25 above), allows the treatment of malignant brain tumor by an innovative mechanism. That is, currently, chemotherapeutic agents that cross the blood-brain barrier, that accumulate in tumor tissue, and that can be stable in the body and administered orally have been developed, but a chemotherapeutic  
30 agent combining all of them does not exist except the peptide of the present invention (Fig. 10).

Since the peptide of the present invention can be combined with various anticancer agents, it can afford a wide range of physiological activity.

35 The present invention opens a breakthrough for the

treatment of brain tumors. In addition, ANXA1 is known to be expressed on vascular surfaces of various malignant tumors, and a dTIT7-bound anticancer agent is expected to be clinically applicable as a therapeutic agent for not only brain tumor but also a wide range of cancers.

In addition, since the peptide of the present invention permits easy chemical synthesis and modification, it is highly promising as a middle molecular biopharmaceutical replacing the antibody pharmaceuticals. In particular, since the peptide of the present invention can be a short chain peptide (e.g., 7 residues of amino acids), it can be produced economically by chemical synthesis.

Furthermore, the peptide of the present invention and a conjugate containing the peptide and other functional part have excellent stability and can also be used as pharmaceuticals for oral administration. Clinical trials are therefore also easy. After actually approved as a medical drug, it can be spread as a pharmaceutical even in the underdeveloped countries where medical facilities are scarce.

[0012]

In addition, a conjugate containing the peptide of the present invention and a detectable substance is useful in, for example, cancer diagnostic applications. The present inventors attempted PET testing of mice using IF7 in the past, but no tumor-specific images were obtained. The reason for the failure is considered to be that IF7 bound to a radioactive reagent was buried inside the compound due to the high hydrophobicity of IF7. The peptide of the present invention can be water soluble and thus can overcome this problem. The peptide of the present invention permits relatively easy modification of, for example, the N-terminal, can be labeled with reagents for 3D imaging such as PET, and can be developed as a diagnostic agent.

[Brief Description of the Drawings]

[0013]

Fig. 1 shows a brain tumor targeting effect of IF7. Brain tissue sections after intravenous administration of IF7-A488 to brain tumor model mice. A488 (left) and Hoechst (right, tumor site). RQ7 is an inverse sequence negative control of IF7. Bars; 5 mm.

Fig. 2 shows a fluorescence microscopic image of brain tumor tissue section from mouse treated with the fluorescently labeled IF7. An enlarged view of the brain tumor tissue section shown in Figure 1. Vascular endothelial cell was stained with CD31 specific antibody (red). IF7-A488 (green) passed through blood vessels to reach cancer cells in the stroma. RQ7 is an inverse sequence control of IF7. Bars; 40  $\mu$ m.

Fig. 3 explains a Dual Tumor Model Mouse. To investigate the influence on the vascular-brain tumor barrier (BBTB), luciferase (Luc)-expressing cancer cells were transplanted into the brain and under the skin of one mouse to create a tumor. When BBTB inhibits penetration of the anticancer drug administered via the tail vein into the brain tumor (BBTB+), the effects of the anticancer drug appear only in the subcutaneous tumor. On the other hand, when the anticancer drug penetrates the brain tumor (BBTB-), it is effective in both the brain and subcutaneous tumors.

Fig. 4 shows therapeutic effects of IF7-SN38 on brain tumor. C6-Luc cells (cells forcedly expressing luciferase in rat glioma) were transplanted into the brains and under the skin of immunodeficient SCID mouse to create tumor to give a dual tumor model. IF7-SN38 was administered daily via the tail vein, luminescence due to luciferase activity in the tumor was measured and viability of C6-Luc cells was determined. IF7-SN38 not only inhibited the growth of brain tumor, but also inhibited brain tumor more strongly than the subcutaneous tumor.

Fig. 5 shows quantitative analysis of fluorescence remaining in the blood of A488-IF7 administered intravenously to cancer bearing mice and healthy mice. A488-IF7 was

administered via the tail vein to mice with subcutaneous carcinoma of melanoma B16, and peripheral blood was collected from the ocular vein over time and fluorescence was measured. The recovered fluorescence decreased depending on the size of the tumor.

Fig. 6 explains screening for D-form peptide that binds to Anx1. (A) D-form peptide sequence identification method by mirror image screen. IF7 binds to Anx1 N terminal peptide ((1) L-form MC16) consisting of 16 amino acid residues. (2) D-form MC16 is synthesized and the sequence binding thereto is selected. (3) The obtained L-form peptide is synthesized with D-amino acid to form a D-form peptide that binds to Anx1. (B) Screening results of T7 phage library targeting D-MC16. Phage enrichment was seen for each round. (C) Sequence analysis of D-MC16 specific T7 phages by next-generation sequencer. (D) Enriched peptides and proportion thereof. (E) Confirmation of binding of TIT7 phage to D-MC16 (plaque formation experiment).

Fig. 7 shows the results of IRDye-dTIT7 administration experiment. Rat glioma cells (C6-Luc) forcedly expressing luciferase were transplanted into the brain of nude mouse to prepare a brain tumor model mouse. dTIT7 (IRDye-dTIT7, unpurified product) labeled with near-infrared fluorescent reagent (IRDye 800CW) was injected into the tail vein, and fluorescence was observed over time by IVIS Imager. Accumulation of fluorescence signals was observed in the brain tumor site and kidney after 24 hours. (A) Overall image over time (B) Enlarged photograph (C) Signal quantification by a software of IVIS imager.

Fig. 8 shows the structure of GA-dTIT7 condensate. An amino group is introduced into geldanamycin to bind a maleimide linker. Cysteine is added in advance to the C-terminal of dTIT7 to perform the synthesis. It is reacted with maleimide via the sulfhydryl group of the cysteine residue to give GA-dTIT7 condensate.

Fig. 9 shows the results of experiment including oral

administration of a drug of dTIT7 bound with an anticancer agent geldanamycin (GA) (GA-dTIT7) to a brain tumor model mouse every other day. Survival of brain tumor was quantified by luminescence by luciferase. GA-dWIP7 containing WIPTTMT  
 5 (peptide in which the order of D-amino acid sequence constituting dTIT7 was changed) bound with geldanamycin was administered as a control to a mouse. A graph (upper right) quantifying the viable cell signal (left) of the malignant tumor and the survival curve (lower right) are shown.

10 Fig. 10 conceptually explains the targeted treatment of malignant brain tumor with an IF7, dTIT7-bound anticancer agent. For convenience, the blood-brain barrier was considered to be present above the neck. General anticancer drugs do not have a tumor targeting ability. Temozolomide penetrates into brain  
 15 tumor to some extent but is not concentrated. Avastin acts on new blood vessels but does not act directly on tumor cell. In contrast, Anxal-bound peptide crosses the blood-brain tumor barrier and accumulates only in tumor at a high concentration.

Fig. 11 shows the results of intermolecular interactions  
 20 of Anxal with dLRF7, dSPT7, dMPT7 and dLLS7 peptides. All peptides tested showed positive binding to Anxal.

Fig. 12 shows the experiment results of administration of IRDye-dLRF7, IRDye-dSPT7, IRDye-dMPT7 and IRDye-dLLS7 to brain tumor model nude mice. All mice, except those injected with  
 25 dLLS7, showed strong signals at the brain tumor site, and three peptides of dLRF7, dMPT7 and dSPT7, other than dLLS7, showed a brain tumor targeting ability via the vasculature route.

[DETAILED DESCRIPTION OF THE INVENTION]

[0014]

30 1. Peptide

The present invention provides a peptide containing the amino acid sequence of any of (I)-(III) below.

(I) an amino acid sequence of (X1) [D]P[D] (X2) [D] wherein X1 is W or F, X2 is S or T,

35 (II) an amino acid sequence of P[D]T[D] (X)<sub>n</sub> F[D] wherein (X)<sub>n</sub>

is any amino acid in the number of n selected independently of each other, n is an integer of 0-4,  
 (III) an amino acid sequence that is a Retro-inverso of the amino acid sequence of any of the aforementioned (I) and the  
 5 aforementioned (II).

[0015]

The present invention also provides a peptide containing the amino acid sequence of any of the following (i)-(vii):

- (i) an amino acid sequence of T[D] I[D] T[D] W[D] P[D] T[D]  
 10 M[D],  
 (ii) an amino acid sequence of L[D] R[D] F[D] P[D] T[D] V[D] L[D],  
 (iii) an amino acid sequence of L[D] L[D] S[D] W[D] P[D] S[D] A[D],  
 15 (iv) an amino acid sequence of S[D] P[D] T[D] S[D] L[D] L[D] F[D],  
 (v) an amino acid sequence of M[D] P[D] T[D] L[D] T[D] F[D] R[D],  
 (vi) an amino acid sequence of any of the aforementioned (i)-  
 20 (v), in which 1 or several amino acids are inserted, substituted or deleted, or these are combined,  
 (vii) an amino acid sequence that is a Retro-inverso of the amino acid sequence of any of the aforementioned (i)-(vi).

[0016]

25 In the present specification, the amino acid sequence of the chain peptide is described according to the conventional manner of peptide indication with the N-terminal side on the left side and the C-terminal side on the right side. In addition, each amino acid symbol with symbol [D] immediately  
 30 following the amino acid sequence indicates a D form of the amino acid, and each amino acid symbol without symbol [D] immediately following the amino acid sequence indicates an L form of the amino acid, unless contrary to the context. In the present specification, a peptide containing the amino acid  
 35 sequence of any one of the above-mentioned (I)-(II) and a

peptide containing the amino acid sequence of any one of the above-mentioned (i)-(vii) are collectively referred to as the peptide of the present invention.

[0017]

5 The amino acid sequence of the above-mentioned (I) may be any of W[D] P[D] S[D], W[D] P[D] T[D], F[D] P[D] S[D] and F[D] P[D] T[D].

[0018]

In the amino acid sequence of the above-mentioned (II),  
10 the integer n is 0 to 4, preferably 2 to 3. X may be any amino acid selected independently of each other.

[0019]

The peptide of the present invention may consist of the amino acid sequence of any of the above-mentioned (I)-(III) and  
15 (i)-(vii), or may have one or more amino acids added to the N-terminal side and/or C-terminal side of these sequences.

[0020]

In the present specification, the peptide refers to one in which two or more amino acids are peptide-bonded. The  
20 length of the peptide of the present invention is not particularly limited. The peptide of the present invention may contain at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14 or at least 15 amino acids.  
25 The peptide of the present invention may consist of up to 50, 45, 40, 35, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8 or 7 amino acids. The peptide of the present invention may have a length of 3-10, 3-15, 3-20, 3-25, 3-30, 3-40, 3-50, 4-10, 4-15, 4-20, 4-25, 4-30, 4-40, 4-50, 5-10, 5-15, 5-20, 5-  
30 25, 5-30, 5-40, 5-50, 6-10, 6-15, 6-20, 6-25, 6-30, 6-40, 6-50, 7-10, 7-15, 7-20, 7-25, 7-30, 7-40, 7-50, 8-10, 8-15, 8-20, 8-25, 8-30, 8-40 or 8-50 amino acids. For example, the peptide of the present invention may have a length of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids.

35 [0021]

The peptide of the present invention may be composed of a combination of a form D amino acid and a form L amino acid. In more detail, all the amino acids constituting the peptide containing the amino acid sequence of any of the above-mentioned (I)-(II) and (i)-(vi) may be form D amino acids, may contain form L amino acid in addition to form D amino acids, and all the amino acids constituting the peptide containing the amino acid sequence of the above-mentioned (III) or (vii) may be form L amino acids, or may contain form D amino acid in addition to form L amino acids. The L-form amino acid may be a naturally occurring L-form amino acid and examples include glycine, alanine, leucine, proline, phenylalanine, tyrosine, methionine, serine, threonine, cysteine, aspartic acid, glutamic acid, asparagine, glutamine, lysine, arginine, hydroxylysine, histidine, tryptophan, valine, all of which are in the L form. The D-form amino acid includes, for example, an optical isomer of an L-form amino acid as described above. In the present specification, glycine, which is an amino acid that does not exhibit optical activity, can be read as an L-form and D-form amino acid unless contrary to the context.

[0022]

The peptide of the present invention may contain a modified or unusual amino acid such as those referred to under 37 C.F.R. 1.821-1.822 and the like. In one embodiment, the peptide of the present invention does not contain a modified or unusual amino acid. Examples of the modified or unusual amino acid include 3-aminoadipic acid,  $\beta$ -alanine, 2-aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisobutyric acid, 2-aminopimelic acid, 2,4-diaminobutyric acid, desmosine, 2,2'-diaminopimelic acid, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, allo-isoleucine, N-methylglycine, N-methylisoleucine, 6-N-methyllysine, N-methylvaline, norvaline, norleucine, ornithine and the like.

[0023]

The amino terminal and/or carboxy terminal of the peptide of the present invention may be modified. The modification of the amino terminal may be methylation (e.g.,  $-\text{NHCH}_3$  or  $-\text{N}(\text{CH}_3)_2$ ),  
5 acetylation (e.g., by acetic acid or its halogenated derivative thereof), or any protecting group such as benzyloxycarbonyl group, carboxylate functional group ( $\text{RCOO}-$ ) or sulfonyl functional group ( $\text{R-SO}_2-$ ) (wherein R is selected from alkyl, aryl, heteroaryl, and alkylaryl, and the like may be introduced.  
10 Modification of the carboxy terminal include amidation ( $-\text{CONH}_2$ ), esterification ( $-\text{COOR}$ ) and the like. Here, as R in the ester, for example, a  $\text{C}_{1-6}$  alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl; a  $\text{C}_{3-8}$  cycloalkyl group such as cyclopentyl, cyclohexyl; a  $\text{C}_{6-12}$  aryl group such as phenyl,  $\alpha$ -naphthyl; a  
15 phenyl- $\text{C}_{1-2}$  alkyl group such as benzyl, phenethyl; a  $\text{C}_{7-14}$  aralkyl group such as  $\alpha$ -naphthyl- $\text{C}_{1-2}$  alkyl group; a pivaloyloxymethyl group and the like are used.

[0024]

The peptide of the present invention may undergo various  
20 modifications other than that at the N-terminal or C-terminal. Chemical modification may be, for example, methylation, acetylation, phosphorylation, and the like. When the peptide of the present invention has a carboxyl group (or carboxylate) at a site other than the C-terminal, the carboxyl group may be  
25 amidated or esterified. As the ester in this case, for example, the C-terminal ester described above and the like are used. Alternatively, the substituents on the side chain of the amino acid in a molecule (e.g.,  $-\text{OH}$ ,  $-\text{SH}$ , amino group, imidazole group, indole group, guanidino group, etc.) may be protected  
30 with an appropriate protecting group (e.g.,  $\text{C}_{1-6}$  acyl group such as  $\text{C}_{1-6}$  alkanoyl group such as formyl, acetyl, etc.).

[0025]

The peptide of the present invention containing the amino acid sequence of the above-mentioned (vi) may contain a partial  
35 sequence consisting of four consecutive amino acids, a partial

sequence consisting of five consecutive amino acids, or a partial sequence consisting of six consecutive amino acids in the amino acid sequence of the above-mentioned (i)-(v). To be specific, for example, when the original amino acid sequence is the amino acid sequence of the above-mentioned (i), the peptide may contain the amino acid sequence of T[D] I[D] T[D] W[D], the amino acid sequence of I[D] T[D] W[D] P[D], the amino acid sequence of T[D] W[D] P[D] T[D], the amino acid sequence of W[D] P[D] T[D] M[D], the amino acid sequence of T[D] I[D] T[D] W[D] P[D], the amino acid sequence of I[D] T[D] W[D] P[D] T[D], the amino acid sequence of T[D] W[D] P[D] T[D] M[D], the amino acid sequence of T[D] I[D] T[D] W[D] P[D] T[D] or the amino acid sequence of I[D] T[D] W[D] P[D] T[D] M[D].

[0026]

The position of mutation (i.e., insertion, substitution, deletion, and combination thereof) in the amino acid sequence of the above-mentioned (vi) is not particularly limited. The mutation may consist of (a) insertion only, (b) substitution only, (c) deletion only, (d) insertion and substitution only, (e) insertion and deletion only, (f) substitution and deletion only, or (g) a combination of insertion, substitution and deletion. The number of mutation in the above-mentioned (a)-(g) is, for example, 1-5, preferably 1-4, more preferably 1-3, further more preferably 1 or 2, even more preferably 1.

[0027]

The amino acid inserted in the amino acid sequence of the above-mentioned (vi) may be any amino acid described above. The amino acid may be an L-form or D-form amino acid, for example, a naturally occurring L-form amino acid described above, or a D-form amino acid which is an optical isomer thereof. The amino acid may also undergo various chemical modifications described above.

[0028]

Preferably, the substitution in the amino acid sequence of the above-mentioned (vi) is a conservative amino acid

substitution. The "conservative amino acid substitution" is well known in the art. For example, conservative amino acid substitution can be defined as substitution between amino acids having similar properties of the side chain. Thus, the conservative amino acid substitution may be, for example, (1) substitution between aromatic amino acids (Phe, Trp, Tyr), (2) substitution between non-polar aliphatic amino acids (Gly, Ala, Val, Leu, Met, Ile, Pro), (3) substitution between non-charged polar amino acids (Ser, Thr, Cys, Asn, Gln), (4) substitution between basic amino acids (Lys, Arg, His), or (5) substitution between acidic amino acids (Asp, Glu). In the present specification, the conservative amino acid substitution may also include substitution of a form D amino acid with a form L amino acid which is an optical isomer thereof. Thus, the conservative amino acid substitution may also be substitution between two L-form amino acids, substitution between two D-form amino acids, or substitution between an L-form amino acid and a D-form amino acid within the groups of the above-mentioned (1)-(5).

20 [0029]

In one embodiment, the sequence of the above-mentioned (vi) may be encompassed in the sequence of the above-mentioned (I) or (II). In this case, the sequence of the above-mentioned (vii) is encompassed in the sequence of the above-mentioned (III). In such embodiments, the fourth amino acid may be W[D] or F[D] and the fifth amino acid may be P[D] and the sixth amino acid may be T[D] or S[D] in the sequence of the above-mentioned (i), the third amino acid may be W[D] or F[D] and the fourth amino acid may be P[D] and the fifth amino acid may be T[D] or S[D] in the sequence of the above-mentioned (ii), the fourth amino acid may be W[D] or F[D] and the fifth amino acid may be P[D] and the sixth amino acid may be T[D] or S[D] in the sequence of the above-mentioned (iii), the second amino acid may be P[D] and the third amino acid may be T[D] and the sixth amino acid may be F[D] in the sequence of the above-mentioned

(iv), and, the second amino acid may be P[D] and the third amino acid may be T[D] and the fifth amino acid may be F[D] in the sequence of the above-mentioned (v).

[0030]

5 Retro-inverso isomer of peptide refers to a peptide in which the chirality of each amino acid residue is reversed ("inverso") and the direction of the amino acid sequence is reversed ("Retro") relative to the original peptide. Retro-inverso isomer is known to exhibit a structure and function  
 10 similar to those of the original peptide (e.g., Acc. Chem. Res., 1993, 26(5), pp 266 273, and PLoS One. 2013 Dec 2; 8(12):e80390). Specifically, the amino acid sequence of the above-mentioned (vii) includes, for example, MTPWTIT (SEQ ID NO: 1) which is the Retro-inverso of the sequence of the above-  
 15 mentioned (i), LVTPFRL (SEQ ID NO: 2) which is the Retro-inverso of the sequence of the above-mentioned (ii), ASPWLL (SEQ ID NO: 3) which is the Retro-inverso of the sequence of the above-mentioned (iii), FLLSTPS (SEQ ID NO: 4) which is the Retro-inverso of the sequence of the above-mentioned (iv), and  
 20 RFTLTPM (SEQ ID NO: 5) which is the Retro-inverso of the sequence of the above-mentioned (v).

[0031]

The peptide of the present invention may contain two or more sequences selected from the above-mentioned (I)-(III) and  
 25 (i)-(vii). In one embodiment, the peptide of the present invention contains a tandem repeat of any of the sequences of the above-mentioned (I)-(III) and (i)-(vii) (i.e., a structure in which identical sequence portions are directly linked to each other). In another embodiment, the peptide of the present  
 30 invention contains a structure in which two or more different sequences of the above-mentioned (I)-(III) and (i)-(vii) are directly linked. Specific repeat structures include, for example, T[D]I[D]T[D]W[D]P[D]T[D]M[D]-  
 T[D]I[D]T[D]E[D]P[D]T[D]M[D], and T[D]I[D]T[D]-T[D]I[D]T[D].  
 35 Alternatively, the peptide of the present invention may

constitute a multivalent peptide with a dendrimer.

[0032]

The peptide of the present invention may be a free form or a salt form. Examples of the salt of the peptide of the present invention include a pharmaceutically acceptable acid addition salt and a base addition salt. Examples of the acid addition salt include salts with inorganic acids such as hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid and the like, and salts with organic acids such as acetic acid, malic acid, succinic acid, tartaric acid, citric acid and the like. Examples of the base addition salt include salts with alkali metals such as sodium, potassium and the like, salts with alkaline earth metals such as calcium, magnesium and the like, and salts with amines such as ammonium, triethylamine and the like.

[0033]

The peptide of the present invention may bind to annexin A1. Annexin A1 (Anx1) is a known protein belonging to the annexin family, also known as lipocortin 1. Genetic sequence and amino acid sequence of Anx1 are known for various species. For example, RefSeq No. NM-000700.2 (SEQ ID NO: 6) of mRNA, RefSeq No. NP-000691.1 (SEQ ID NO: 7) of protein for human Anx1; RefSeq No. NM-010730.2 (SEQ ID NO: 8) of mRNA, RefSeq No. NP-034860.2 [ARRAY ID NO:9] of protein for mouse Anx1 can be mentioned. The peptide of the present invention may bind to the N-terminal region of Anx1. More particularly, the peptide of the present invention may bind to a region consisting of 1st to 15th amino acids of SEQ ID NO: 7 in human Anx1 or a region consisting of 1st to 15th amino acids of SEQ ID NO: 9 in murine Anx1. The peptide of the present invention may have dissociative constant (Kd value) of preferably less than  $10^{-6}$  M, more preferably less than  $10^{-7}$  M, further more preferably less than  $5 \times 10^{-8}$  M when, for example, intermolecular interactions with mouse or human Anx1 are measured using the QCM (Quartz Crystal Microbalance) method. For the above-mentioned

measurement, human recombinant Anxal protein consisting of 346 amino acid residues, which is commercially available from ATGen Corporation (Seongnam-si, South Korea) can be used.

[0034]

5           The peptide of the present invention can be produced according to a known peptide synthesis method. The method of peptide synthesis may be any of, for example, a solid phase synthesis process and a liquid phase synthesis process. The object peptide can be produced by condensing a partial peptide  
10 or amino acid capable of constituting the peptide of the present invention and the remaining portion and, when the product has a protecting group, by eliminating the protecting group.

          Here, the condensation and elimination of a protecting  
15 group can be performed according to a method known per se, for example, the methods described in the following (1) - (8).

- (1) M. Bodanszky & M.A. Ondetti, Peptide Synthesis, Interscience Publishers, New York (1966)
- (2) Schroeder & Luebke, The Peptide, Academic Press, New York  
20 (1965)
- (3) Nobuo Izumiya, et al.: Peptide Gosei-no-Kiso to Jikken (Basics and experiments of peptide synthesis), published by Maruzen Co. (1975)
- (4) Haruaki Yajima and Shunpei Sakakibara: Seikagaku Jikken  
25 Koza (Biochemical Experiment) 1, Tanpakushitsu no Kagaku (Chemistry of Proteins) IV, 205 (1977)
- (5) Haruaki Yajima, ed.: Zoku Iyakuhiin no Kaihatsu (A sequel to Development of Pharmaceuticals), Vol. 14, Peptide Synthesis, published by Hirokawa Shoten.
- 30 (6) Stewart, J.M. & Young, J.D., "Solid phase peptide synthesis (2nd ed.)", Pierce Chemical Company, Rockford (1984)
- (7) Atherton, E. & Sheppard, R.C., "Solid Phase peptide synthesis: a practical approach", IRL Press, Oxford (1989)
- (8) "Fmoc Solid Phase Peptide Synthesis: A Practical Approach  
35 (Practical Approach Series)", Oxford University Press (2000)

[0035]

The thus-obtained peptide can be purified and isolated by a known purification method. Examples of the purification method include solvent extraction, distillation, column chromatography, liquid chromatography, recrystallization, combinations thereof and the like.

When the peptide obtained by the above-mentioned method is in a free form, the form can be converted to a suitable salt by a known method or a method analogous thereto; conversely, when the peptide is obtained in the form of a salt, the salt can be converted to a free form or other salt by a known method or a method analogous thereto.

[0036]

As stated above, annexin A1 is known to have the highest specificity among the currently known tumor vessel specific marker molecules, and is expressed intracellularly in normal cells, but strongly expressed on luminal surface adjacent to the blood stream in tumor neovascular endothelial cells (Oh et al., Nature 429: 629-35, 2004). Thus, the peptide of the present invention can selectively bind to angiogenic tumors in vivo. In addition, Anxal is expressed on the blood side of neovascular endothelial cells formed in tumors, and when bound on the blood side to ligands such as the present peptides, the ligands are transported to the basal side by transcytosis and are actively released into the stroma where cancerous cells are present. Thus, the peptide of the present invention can target malignancies in vivo. Thus, the peptide of the present invention is useful, for example, for targeting malignant tumors.

[0037]

## 2. Conjugate

The present invention also provides a conjugate in which one or more components are bound with the above-mentioned peptide of the present invention (hereinafter to be also referred to as the conjugate of the present invention).

[0038]

The component is not particularly limited as long as it can be linked to the peptide of the present invention, and may be suitable for administration to an animal (e.g., human), and  
5 may perform some function in the body of the animal. The component may be natural or non-natural. Examples of the component include, but are not limited to, biological materials (e.g., cell, phage, virus, etc.), oligonucleotides and nucleic acids (e.g., DNA, RNA, DNA/RNA chimera, etc.), peptide,  
10 polypeptide, protein, antibody, lipid, polysaccharide, small molecule compounds (e.g., organic or inorganic compounds of not more than 1000 Da), particles (e.g., gold particles, various nanoparticles, etc.), combinations thereof, and the like.

[0039]

15 The component may perform a given functionality at a target site in the body of an animal (e.g., human). The type of the functionality is not particularly limited. Since the conjugate of the present invention can be targeted to malignant tumor by the action of the part corresponding to the peptide of  
20 the present invention, preferred examples of the functionality include anti-cancer activity and provision of detectability. Thus, the component may be, for example, an anticancer agent or a detectable substance.

[0040]

25 (Anticancer agent)

In the present specification, the anticancer agent refers to a drug intended to suppress growth of a malignant tumor (cancer). The mechanism of the action of anticancer agent is not particularly limited. The anticancer agent may be an  
30 antimetabolite, an alkylating agent, an anticancer antibiotic, a microtubule inhibitor, a platinum preparation, a topoisomerase inhibitor, a molecular targeting agent, or the like. The conjugate of the present invention may contain two or more the same or different anticancer agents.

35 [0041]

The metabolic antagonist may be, for example, a folic acid antimetabolite, a dihydropteroate synthase inhibitor, a dihydrofolate reductase inhibitor (DHFR inhibitor), a pyrimidine metabolism inhibitor, a thymidylate synthase inhibitor, a purine metabolism inhibitor, an IMPDH inhibitor, a ribonucleotide reductase inhibitor, a ribonucleotide reductase inhibitor, a nucleotide analog, L-asparaginase and the like. Specific examples of the metabolic antagonist include enocitabine (SUNRABIN), capecitabine (Xeloda), carmofour (Mifuro1), cladribine (Leustatin), gemcitabine (Gemzar), cytarabine (kiloside), cytarabine ocfosphate (Starasid), tegafur (Achillon, AFTHOUR, Tefseal, Futrafur, Lunasin etc.), tegafur/uracil (UFT), tegafur/Gimeracil/Oteracil potassium (TS-1: T-S-One), doxifluridine (FURTURON), nelarabine (Arranon G), hydroxycarbamide (HYDREA), fluorouracil (5-FU, carzonal, Benton, Lunachol, Lunapon), fludarabine (fludara), pemetrexed (alimta), pentostatin (cofolin), mercaptopurine (leukerin), methotrexate (methotrexate) and the like.

[0042]

Specific examples of the alkylating agent include nitrogen mustard based alkylating agents such as cyclophosphamide (endoxane), ifosfamide (ifomide), melphalan (alckeran), busulfan, thiotepa (tespamine) and the like, nitrosourea-based alkylating agents such as nimustine (nidran), ranimustine (cymerin), dacarbazine (dacarbazine), procarbazine (procarbazine hydrochloride), temozolomide (Temodal), carmustine (Gliadel), streptozotocin (zanosar), bendamustine (treakisym) and the like.

[0043]

Specific examples of the anti-cancer antibiotic include actinomycin D (cosmegen), aclarubicin (aclacinone), amrubicin (Calsed), idarubicin (idamycin), epirubicin (epirubicin hydrochloride, Farmorubicin), zinostatin stimalamer (Smancs), daunorubicin (daunomycin), doxorubicin (adriacin), pirarubicin (Pinorubin, THERARUBICIN), bleomycin (Bleo), peplomycin (Peleo),

mitomycin C (Mitomycin), mitoxantrone (Novantrum), liposomal doxorubicin (Doxil) and the like.

[0044]

Examples of the microtubule inhibitor include vinca  
5 alkaloid microtubule polymerization inhibitors such as  
vinblastine (exal), vincristine (oncovin), vindesine (Fildesin)  
and the like, taxane type microtubule depolymerization  
inhibitors such as paclitaxel (taxol), docetaxel (taxotere) and  
the like.

10. [0045]

Examples of the platinum preparation include oxaliplatin  
(El Prat), carboplatin (Carboplatin, Carbomerck, Paraplatin),  
cisplatin (IA-call, Konaburi, Cisplatin etc.), nedaplatin  
(Aqupla) and the like.

15 [0046]

Examples of the topoisomerase inhibitor include type I  
topoisomerase inhibitors such as camptothecin and a derivative  
thereof (e.g., irinotecan (Campto), nogitecan (HYCAMTIN), SN-38  
and the like) and the like; type II topoisomerase inhibitors  
20 such as anthracycline drugs such as doxorubicin (adriacin) and  
the like, epipodophyllotoxin drugs such as etoposide (Lastet,  
VePesid) and the like, and quinolone drugs such as levofloxacin  
(cravit), ciprofloxacin (ciproxan) and the like.

[0047]

25 Examples of the molecular target drug include legolafenib  
(Stivarga), cetuximab (Erbix), panitumumab (Vectibix),  
ramsilimab (Cyramza), gefitinib (Iressa), erlotinib (Tarceva),  
afatinib (GIOTRIE), crizotinib (XALKORI), alectinib (ALECENSA),  
ceritinib, Lenvatinib (Lenvima), trastuzumab (Herceptin),  
30 lapatinib (Tykerb), pertuzumab (PERJETA), sunitinib (Sutent),  
sorafenib (Nexavar), axitinib (Inlyta), pazopanib (Votrient),  
Nivolumab (OPDIVO), pembrolizumab, ipilimumab (YERVOY),  
vemurafenib (ZELBORAF), everolimus (AFINITOR), temsirolimus  
(TORISEL), rituximab (Rituxan), bevacizumab (Avastin),  
35 geldanamycin and the like.

[0048]

The anticancer agent may also be an anti-angiogenic agent. The anti-angiogenic agent may be one that inhibits vascular endothelial growth factor (VEGF) or other angiogenic factor, or  
 5 a receptor thereof. Specific examples of the anti-angiogenic agent include angiostatin, endostatin, métastatin, anti VEGF antibody (e.g., Avastin), VEGFR-2 inhibitor (e.g., SU5416, SU6668), and the like.

[0049]

10 (Detectable substance)

In the present specification, the detectable substance refers to any substance that makes the conjugate of the present invention containing the substance detectable. Preferably, the detectable substance enables detection of the conjugate of the  
 15 present invention in vivo directly or indirectly using an appropriate visualization or imaging means. Examples of the visualization or imaging means include, but are not limited to, X-ray photography, computed tomography (CT), nuclear magnetic resonance imaging (MRI), ultrasonography, scintigraphy,  
 20 positron emission tomography (PET), endoscopy, laparoscopy, and the like. The detectable substance may be, for example, a radioisotope, an enhancing agent for MRI (e.g., paramagnetic ions), a radiopaque substance, a contrast agent, a fluorescent substance, or the like.

25 [0050]

Examples of the radioactive nuclide useful for PET include  $^{18}\text{F}$ ,  $^{51}\text{Mn}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{52}\text{Fe}$ ,  $^{55}\text{Co}$ ,  $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$ ,  $^{82\text{m}}\text{Rb}$ ,  $^{83}\text{Sr}$ ,  $^{86}\text{Y}$ ,  $^{89}\text{Zr}$ ,  $^{94\text{m}}\text{Tc}$ ,  $^{110}\text{In}$ ,  $^{120}\text{I}$ ,  $^{124}\text{I}$  and the like.  
 Examples of the radioactive nuclide useful for detection of  $\gamma$   
 30 ray include  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{67}\text{Cu}$ ,  $^{67}\text{Ga}$ ,  $^{75}\text{Se}$ ,  $^{97}\text{Ru}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{114\text{m}}\text{In}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{169}\text{Yb}$ ,  $^{197}\text{Hg}$ ,  $^{201}\text{Tl}$  and the like.

[0051]

Preferable examples of the paramagnetic ion include chromium (III), manganese (II), iron(III), iron(II), cobalt(II),  
 35 nickel(II), copper(II), neodymium (III), samarium (III),

ytterbium (III), gadolinium (III), vanadium (II), terbium (III), dysprosium (III), holmium (III), erbium (III) and the like, and gadolinium is particularly preferable. In addition, metals such as lanthanum (III), gold (III), lead (II), bismuth (III) and the like are also useful for X-ray imaging and the like.

[0052]

Examples of the radiopaque substance and contrast agent include iodine compounds (for example, organic iodic acids such as iodocarboxylic acid, iodoform, triiodophenol, tetraiodoethylene, etc.), barium compounds (for example, barium sulfate and the like), gallium compounds (such as gallium citrate), thallium compounds (such as thallium chloride) and the like.

[0053]

Examples of the fluorescent substances include rhodamine, fluorescein, Cy dye, Alexa (registered trademark) Fluor, phycoerythrin (PE), allophycocyanin (APC), derivatives thereof and the like. Near infrared fluorescent reagents such as indocyanine are also exemplified as preferable fluorescent substances.

[0054]

(Binding of peptide of the present invention with component)

The mode of binding of the peptide of the present invention with one or more components in the conjugate of the present invention is not particularly limited. The binding may be direct or indirect via a linker etc. The binding may be by covalent binding, non-covalent binding, or a combination thereof. One or more components may be directly or indirectly bonded at the N-terminal, C-terminal, or other position of the peptide of the present invention. The binding of a peptide with other component (or second peptide) is well known in the art, and the binding may be by any known means in the conjugate of the present invention.

[0055]

For example, when the binding is via a linker, known

crosslinkers such as NHS ester, imide ester, maleimide, carbodiimide, allyl azide, diaziline, isocyanide, psoralen and the like can be used. Depending on the crosslinker to be used, the peptide of the present invention may be modified as appropriate. For example, a cysteine can be added in advance to the C-terminal of the peptide of the present invention for binding with a maleimide linker.

[0056]

For linking radioactive metal or paramagnetic ion mentioned above to the peptide of the present invention, an appropriate chelating agent (e.g., ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), 4,7,10-tetraazacyclododecane-N-N',N'',N'''-tetraacetic acid (DOTA) and the like) and/or metallothionein and the like can be used. See, for example, Culali Aktolun et al. ed., "Nuclear Medicine Therapy: Principles and Clinical Applications", Springer, 2013 and the like.

[0057]

### 3. Composition

The present invention also provides a composition containing the peptide or conjugate of the present invention and a pharmacologically acceptable carrier (hereinafter to be also referred to as the composition of the present invention. The composition may be provided in a dosage form suitable for oral or parenteral administration.

[0058]

Examples of the composition for parenteral administration include injection, suppository and the like. The injection may include dosage forms such as intravenous injection, subcutaneous injection, intradermal injection, intramuscular injection, drip intravenous injection, and the like. Such injections can be prepared according to a known method. As a method for preparing an injection, for example, the peptide or conjugate of the present invention can be prepared by dissolving, suspending or emulsifying the peptide or conjugate

of the present invention in a sterile aqueous solution or oily solution generally used for injection. As an aqueous solution for injection, for example, physiological saline, an isotonic solution containing glucose and other adjuvant may be used and  
5 may be used in combination with suitable solubilizing agents such as alcohols (e.g., ethanol), polyalcohols (e.g., propylene glycol, polyethylene glycol), nonionic surfactants (e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)), and the like. As the oily solution,  
10 for example, sesame oil, soybean oil, or the like is used, and benzyl benzoate, benzyl alcohol, or the like may be used in combination as a solubilizing agent. The prepared injection solution is preferably filled in a suitable ampoule. A suppository to be used for rectal administration can be  
15 prepared by mixing the peptide or conjugate of the present invention with a conventional base for suppository.

[0059]

The composition for oral administration includes, for example, solid or liquid dosage forms, specifically tablet  
20 (including sugar-coated tablet, film-coated tablet), pill, granule, powder, capsule (including soft capsule), syrup, emulsion, suspension, and the like. Such compositions are produced by a known method and may contain carrier, diluent or excipient generally used in the field of preparations. As the  
25 carrier and excipient for tablets, for example, lactose, starch, sucrose, magnesium stearate are used.

[0060]

Each of the above-mentioned compositions may contain other active ingredient as long as blending with the above-  
30 mentioned peptide or conjugate does not cause an undesirable interaction.

[0061]

The above-mentioned parenteral or oral pharmaceutical compositions are conveniently formulated in a dosage unit form  
35 compatible with the dosage of the active ingredient. Examples

of the dosage form for such dosage unit include tablet, pill, capsule, injection (ampoule) and suppository. The content of the peptide or conjugate is generally preferably 1 to 500 mg per dosage unit form, preferably, in particular, 1 to 100 mg for injection, and 10 to 250 mg for other dosage forms.

[0062]

The composition of the present invention can target tumor, particularly angiogenic malignant tumor, and preferably can accumulate the peptide or conjugate in the tumor. Thus, the composition of the present invention containing an anticancer agent in the conjugate is useful for the treatment or prevention of the targeted malignant tumor. The composition of the present invention containing a detectable substance in the conjugate is also useful for the examination or diagnosis of malignant tumor.

[0063]

The malignant tumor (cancer) may be any kind of cancer, and may also be solid cancer or liquid cancer. As the solid cancer, those that express Anxal on the cell surface, and therefore, solid tumors that are angiogenic, are preferably mentioned. Examples of the solid cancer include cancer of the brain/nervous system (e.g., brain tumor, spinal cord tumor etc.), head and neck cancer (e.g., laryngeal cancer, oral cancer, salivary gland cancer, paranasal sinus cancer, thyroid cancer etc.) digestive organ cancer (e.g., stomach cancer, esophageal cancer, small intestine cancer, colon cancer, rectal cancer, anal cancer, liver cancer, biliary tract cancer, pancreatic cancer etc.), urinary or reproductive organ cancer (e.g., kidney cancer, kidney cell cancer, bladder cancer, prostate cancer, renal pelvis and ureter cancer, gall bladder cancer, bile duct cancer, testis cancer, penile cancer, uterine cancer, endometrial cancer, uterine sarcoma, cervical cancer, vaginal cancer, vulvar cancer, ovarian cancer, fallopian tube cancer and the like), respiratory system cancer (e.g., lung cancer (including small cell lung cancer, non-small cell lung

cancer, metastatic lung cancer), bronchial cancer and the like), breast cancer, skin cancer (e.g., malignant melanoma etc.), bone cancer (e.g., osteosarcoma etc.), muscle cancer (e.g., rhabdomyosarcoma etc.) and the like.

5 [0064]

As the liquid cancer, leukemia, malignant lymphoma, multiple myeloma, myelodysplastic syndrome and the like can be mentioned. As leukemia, acute myeloid leukemia, acute lymphocytic leukemia, chronic myeloid leukemia, chronic  
 10 lymphocytic leukemia and the like can be mentioned. The malignant lymphoma is classified into Hodgkin lymphoma and non-Hodgkin's lymphoma, and examples of non-Hodgkin's lymphoma include B cell lymphoma, adult T cell lymphoma, lymphoblastic lymphoma, diffuse large cell lymphoma, Burkitt lymphoma,  
 15 follicular lymphoma, MALT lymphoma, peripheral T cell lymphoma, mantle cell lymphoma and the like.

[0065]

Since the peptide or conjugate of the present invention can efficiently cross the blood-brain tumor barrier, brain  
 20 tumor can be recited as a particularly preferred target. The brain tumor may be a primary brain tumor or a metastatic brain tumor. The brain tumor may also be a benign brain tumor (e.g., meningioma, pituitary adenoma, schwannoma, etc.) or malignant brain tumor, preferably malignant brain tumor. Examples of the  
 25 malignant brain tumor include grade 2 brain tumor s such as astrocytoma, oligodendroglioma, grade 3 brain tumor s such as anaplastic astrocytoma, anaplastic oligodendroglioma, anaplastic oligoastrocytoma, and grade 4 brain tumor s such as glioblastoma.

30 [0066]

The composition of the present invention may be administered to an animal, particularly a mammal, expressing annexin A1. Examples of the mammal include, but are not limited to, laboratory animals such as rodents such as mouse,  
 35 rat, hamster, guinea pig, and rabbit and the like, domestic

animals such as pig, cow, goat, horse, sheep, mink and the like, pets such as dog, cat and the like, primate such as human, monkey, rhesus monkey, marmoset, orangutan, chimpanzee, and the like.

5 [0067]

The dosage of the composition of the present invention also varies depending on the purpose of administration, administration subject, target disease, condition, administration route, and the like. For example, when used for  
10 the treatment or prevention of cancer described above, it is convenient to administer the conjugate of the present invention containing an anticancer agent once a week intravenously or orally generally at 0.1 to 10 mg per kg body weight as a single dose. Alternatively, when used for testing or diagnosing  
15 cancer described above, it is convenient to administer the conjugate of the present invention containing a detectable substance intravenously or orally at generally about 0.1 to 10 mg/kg body weight prior to the testing.

[Examples]

20 [0068]

The present invention is described in more detail in the following by referring to examples and the like, but the present invention is not limited by the following examples and the like.

25 [0069]

(Preparation of experimental animal)

Rat glioma C6 cells were cultured in Dulbecco's Modified Eagles Medium added with 10% fetal bovine serum, high glucose and antibiotic. Lentivirus vector PGK-Luc was prepared at the  
30 Virus Core Facility of Sanford-Burnham-Prebys Medical Discovery Institute. C6 cell and B16 cell were infected with PGK-Luc lentivirus and luciferase positive cells were produced. Using a stereotactic frame, C6-Luc cells ( $4.8 \times 10^4$  cells suspended in 4  $\mu$ l PBS) were injected into the brain corpus striatum of  
35 C57BL/6 mouse. Two days later, the mouse was imaged for tumors

expressing luciferase. To this end, 100  $\mu$ l of luciferin (30 mg/ml PBS) was injected into the peritoneal cavity, the mouse was anesthetized with isoflurane gas (20 ml/min) together with oxygen (1 ml/min) and placed under a camera provided with  
5 Xenogen IVIS 200 imager. The photon number was measured for 1-10 seconds. For the dual tumor model, C6-Luc cells were similarly injected into the brain of NOD-SCID mouse and tumor growth was monitored by Xenogen imager.

When the brain tumor became detectable, C6-Luc cells  
10 ( $2 \times 10^5$  cells suspended in 100  $\mu$ l PBS) were subcutaneously injected into the dorsal flank of the same mouse. The photon number in the brain tumor and subcutaneous tumor was measured using Xenogen imager. A dual tumor model of the same strain C57BL/6 mouse was also prepared using B16-Luc cells. B16-Luc  
15 cells ( $5 \times 10^4$  cells) were injected into the brain of 8- to 10-week-old C57BL/6 female mouse as described above for C6-Luc cells. B16-Luc cells ( $2 \times 10^5$  cells) were subcutaneously injected when B16-Luc tumor in the brain became detectable.

[0070]

20 Examination Example 1: Administration experiment of IF7 to tumor-bearing mouse

IF7 (i.e., L form peptide having the amino acid sequence of IFLLWQR (SEQ ID NO: 10)) was intravenously administered to a tumor-bearing mouse. After reaching the blood vessels around  
25 the tumor, IF7 was taken into the vesicle from the luminal side of vascular endothelial cells, moved to the basal side and was released to the stroma where cancer cells were present.

Even in mouse vascular endothelial F2 cells, IF7 bound to Anxal actively transected tumor vascular endothelial cells by  
30 transcytosis. Therefore, it was suggested that IF7 has an activity to cross the vascular-brain tumor barrier in the brain tumor.

Surprisingly, when fluorescently labeled IF7 was intravenously injected into a brain tumor model mouse  
35 transplanted with glioma cells, accumulation of fluorescence at

a high concentration was observed in the brain tumor site (Figs. 1 and 2). In addition, fluorescence crossed the blood vessel and reached cancer cells in the brain stroma.

[0071]

5 Examination Example 2: Administration experiment of IF7-SN38 to dual tumor model

Using a mouse within which both subcutaneous tumor and brain tumor are present (dual tumor model), the treatment effect of IF7-SN38 was verified. The experiment was conducted  
10 as follows.

IF7C(RR)-SN38 was injected into the tumor-bearing mouse via the tail vein when the photon number of the brain tumor became  $>1.0 \times 10^6$  (this generally occurred 2 weeks later). IF7C(RR)-SN38 (14.2 mg or 6.63 mmol) was dissolved in 100  $\mu$ l of  
15 dimethyl sulfoxide (DMSO), 1  $\mu$ L of the solution (142 mg or 66.3 nmoles) was diluted with 50% ethanol solution (10  $\mu$ l) of Cremophore EL, and 90  $\mu$ l of PBS was further added. The IF7C(RR)-SN38 amount per injection was 142 mg/mouse or 7.1 mg/kg. The inverse sequence of IF7 or RQ7C(RR)-SN38 prepared  
20 similarly was used as a control. Irinotecan dissolved in PBS and PBS alone were used as a negative control.

As a result, both the brain tumor and the subcutaneous tumor were suppressed, and the effect thereof was higher in the brain tumor than in the subcutaneous tumor (Figs. 3 and 4).  
25 Similar results were obtained in the B16 melanoma metastatic brain tumor model, and the results did not change even when the host mouse was changed to C57BL/6 or SCID strain. These facts indicate that DDS targeting Anxal not only efficiently crosses the blood-brain tumor barrier, but also provides an excellent  
30 therapeutic effect in the brain tumor irrespective of the tumor cell type and mouse strain.

[0072]

Examination Example 3: Study of pharmacokinetics of IF7

As demonstrated in Examination Examples 1 and 2, IF7 has  
35 a superior malignant tumor targeting activity. In terms of

clinical development, however, there are problems in two points: poor solubility and low stability. That is, IF7 peptide is protease sensitive and is susceptible to degradation. Indeed, in an experiment including intravenous administration of A488-IF7 to healthy mouse, the fluorescence signal of peripheral blood almost disappeared in about 1 hour (Fig. 5). However, with tail vein administration to tumor-bearing mouse, fluorescence recovered from the peripheral blood of A488-IF7 was markedly lower than that in healthy mice. This result suggests that A488-IF7 is mostly accumulated rapidly in the tumor while being degraded. In addition, IF7 has high hydrophobicity and causes difficulty in dosage form such as the need for addition of a surfactant after dissolution in DMSO, in an attempt to avoid precipitation when a compound bound with an anticancer drug is intravenously administered.

[0073]

#### Example 1: Identification of dTIT7 peptide

The present inventors considered that a more superior therapeutic agent could be created once the above-mentioned weakness associated with IF7 was overcome, and started to develop a next generation peptide consisting of a novel sequence.

The technique to be the basis of this research is a method of screening a phage library that displays on the surface about 1 billion ( $20^7$  species) of random peptides consisting of 7 amino acid residues. All amino acids constituting living organisms are L-form amino acids except for some exceptions. Therefore, it is not possible to obtain the sequence of a D-form peptide that binds to the target by general phage library screening. Therefore, a mirror image screening using a T7 phage library (Funke et al., Mol. Biosystem. 2009, 783-6) was newly performed (Fig. 6).

Screening was performed based on the following findings and assumptions to obtain a phage having high binding affinity to Anx1. It has been clarified that IF7 interacts with the N-

terminal 15 residues of Anx1 (MAMVSEFLKQAWFIE; SEQ ID NO: 11),  
and that IF7 binds to L-peptide with 16 residues (L-MC16) which  
is Anx1 added with Cys residue for modification to the N-  
terminal 15 residues thereof (Sasai et al., unpublished).  
5 First, D-MC16 (all 16 bases are D-form) which is in mirror  
image relation with L-MC16 was synthesized and 7 amino acids  
(L-form TIT7 peptide) that bind to D-MC16 were identified by  
screening using T7 phage library. Based on the obtained  
sequence, TIT7 peptide (dTIT7) composed of D-form amino acids  
10 was synthesized. This dTIT7 peptide is expected to bind to  
Anx1.

When analyzing the peptide sequence displayed by the  
phage, it was a general practice to optimize the sequence by  
introducing various mutations into the peptide sequence.  
15 Currently, it is possible to evaluate and identify the optimal  
peptide sequences at once from the sequences of not less than  
10,000 candidate peptides by applying the obtained phage groups  
to the next generation DNA sequencer.

Analysis results revealed that the frequency of  
20 appearance of 7 amino acids starting with TIT (threonine-  
isoleucine-threonine) is high as a sequence binding to the N-  
terminal part of Anx1, and among them, the TIT7 sequence  
"TITWPTM" (SEQ ID NO: 12) is prominent. While the frequency of  
the partial sequence of TIT7 also appeared in the top ranking  
25 position, no sequence substitute of TIT7 was found. In  
addition, the peptide of TIT7 was composed of a novel sequence  
completely different from the sequence of IF7 (IFLLWQR; SEQ ID  
NO: 10).

The above results showed that straight chain peptide TIT7  
30 of seven amino acids that consists of all D-form amino acids  
was successfully obtained.

Furthermore, when the intermolecular interaction between  
dTIT7 and Anx1 was measured by the QCM method, the  
dissociation constant (Kd value) was  $4.57 \times 10^{-8}$  M.

35 As shown in Fig. 6D, peptides with sequences of LRFPTVL

(SEQ ID NO: 13), SPTSLLF (SEQ ID NO: 14), MPTLTFR (SEQ ID NO: 15), LKGLMRI (SEQ ID NO: 16) and LLSWPSA (SEQ ID NO: 17) frequently appeared in addition to the TIT7 sequence (SEQ ID NO: 12), and among the vast number of  $10^9$  species of random peptides, these six peptides occupied about 50% of the whole. [0074]

Example 2: Brain tumor targeting ability of dTIT7 peptide

dTIT7 (IRDye-dTIT7) labeled with near infrared fluorescent reagent IRDye 800CW was prepared. A brain tumor nude mouse model in which rat glioma was transplanted into the brain of the mouse was prepared and IRDye-dTIT7 was injected via the tail vein of the mouse. As a result, it was observed by in vivo imaging that IRDye-dTIT7 accumulates at the tumor site of a brain tumor model mouse (Fig. 7).

[0075]

Example 3: Antitumor effect of dTIT7-bound anti-cancer agent on brain tumor model

Next, the effect of dTIT7-anticancer agent conjugate on brain tumor model mouse was examined.

GA-dTIT7 in which an anticancer drug geldanamycin was bound to dTIT7 peptide (Fig. 8) is resistant to both protease and esterase and is considered to be more stable in vivo. GA-dTIT7 at a 1/10-fold (in terms of molar number) of the recommended dose was intravenously administered to a tumor-bearing mouse (melanoma B16 cells, subcutaneously administered) every 1 or 2 days. As a result, tumor growth was remarkably suppressed, and a large amount of necrosis was observed at the tumor site of mouse administered with GA-dTIT7 by histopathological observation. This effect strongly suggests the possibility that dTIT7 is superior to IF7 which required consecutive administration (Hatakeyama et al., Proc. Natl. Acad. Sci USA, 108: 19587-92, 2011).

More surprisingly, when GA-dTIT7 was orally administered to a brain tumor model transplanted with B16 cells into the brain, tumor regression was observed, the tumor continued to

decrease even after administration of the drug was stopped, and finally, complete cure was observed in 2 out of 4 mice (Fig. 9). Similar therapeutic effects were also observed in a brain tumor model using C6 cells. This suggests that GA-dTIT7 may lead to the complete cure of brain tumor as an anticancer agent that can be administered orally.

[0076]

Example 4: Intermolecular interaction between dLRF7, dSPT7, dMPT7 and dLLS7 peptides and Anxal

10 As for the peptides of the sequences of LRFPTVL (SEQ ID NO: 13), SPTSLLF (SEQ ID NO: 14), MPTLTFR (SEQ ID NO: 15) and LLSWPSA (SEQ ID NO: 17) with high emergence frequency besides the TIT7 sequence (SEQ ID NO: 12) described in Example 1, similar to the dTIT7 peptide, peptides (dLRF7, dSPT7, dMPT7 and 15 dLLS7 peptides) composed of D-form amino acids were synthesized and the intermolecular interaction with Anxal was measured for each of them by the following method.

The sensor chip was washed twice with piranha solution for 5 min and then immersed overnight in a solution of 0.9 mM 20 hydroxy-EG3-undecanethiol (Dojindo) and 0.1 mM amino-EG6-undecanethiol (Dojindo). After washing with water, the chip was treated with 1 mM GMBS (Dojindo) and incubated at room temperature for 1 hr. After washing away the remaining GMBS, 1 mM peptide solution was added onto the chip and the chip was 25 incubated for 30 min. The peptide-bound sensor chip was placed in a QCM apparatus (Single-Q, ASONE), and 0.5 ml of PBS was added to the well. Various doses of Anxal were injected into the reaction chamber and the binding affinity (kd value) for each peptide was determined.

30 As a result, all peptides tested showed positive binding to Anxal (Table 1 and Fig. 11).

[0077]

[Table 1]

name of synthetic peptide	kd value (M)
dLRF7	$3.119 \times 10^{-8}$
dSPT7	$9.029 \times 10^{-8}$
dMPT7	$2.982 \times 10^{-7}$
dLLS7	$5.418 \times 10^{-8}$

[0078]

5 Example 5: Brain tumor targeting ability of dLRF7, dSPT7, dMPT7 and dLLS7 peptides

DLRF7, dSPT7, dMPT7 and dLLS7 peptides which were extended by adding terminal cysteine were chemically synthesized. Then, each peptide was bound to near infrared  
 10 fluorescent dye IRDye 800CW maleimide (Li-Cor) via a cysteine residue. The resulting peptide conjugate was purified by HPLC and then lyophilized. Each peptide-IRDye conjugate was dissolved in DMSO and 6% glucose and intravenously injected into brain tumor model nude mouse at 100  $\mu$ l/mouse. After 24 hr,  
 15 the fluorescence image was measured by IVIS spectrum (Perkin Elmer).

As a result, all mice except those injected with dLLS7 showed a strong signal of IRDye at the brain tumor site, and three peptides dLRF7, dMPT7 and dSPT7 other than dLLS7 showed a  
 20 brain tumor targeting ability via the vasculature pathway (Fig. 12).

[0079]

This application is based on a patent application No. 2016-159743 filed in Japan (filing date: August 16, 2016).

25

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 85026686 Seq 04-02-2019 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

CLAIMS:

1. A peptide comprising the amino acid sequence T[D] I[D] T[D] W[D] P[D] T[D] M[D], wherein each amino acid symbol immediately followed by symbol [D] is a D form of the amino acid.
2. A peptide consisting of the amino acid sequence T[D] I[D] T[D] W[D] P[D] T[D] M[D], wherein each amino acid symbol immediately followed by symbol [D] is a D form of the amino acid.
3. A conjugate comprising the peptide according to claim 1 or 2, and one or more components.
4. The conjugate according to claim 3, wherein the one or more components comprise an anticancer agent.
5. The conjugate according to claim 4, wherein the anticancer agent is selected from the group consisting of an antimetabolite, an alkylating agent, an anticancer antibiotic, a microtubule inhibitor, a platinum preparation, a topoisomerase inhibitor, a molecular targeting agent, and an anti-angiogenic agent.
6. The conjugate according to claim 4 or 5, wherein the anticancer agent is selected from the group consisting of enocitabine, capecitabine, carmofour, cladribine, gemcitabine,

85026686

cytarabine, cytarabine ocfosphate, tegafur, tegafur/uracil, tegafur/Gimeracil/Oteracil potassium, doxifluridine, nelarabine, hydroxycarbamide, fluorouracil, fludarabine, pemetrexed, pentostatin, mercaptopurine, methotrexate;

cyclophosphamide, ifosfamide, melphalan, busulfan, thiotepa, nimustine, ranimustine, dacarbazine, procarbazine, temozolomide, carmustine, streptozotocin, bendamustine;

actinomycin D, aclarubicin, amrubicin, idarubicin, epirubicin, zinostatin stimalamer, daunorubicin, doxorubicin, pirarubicin, bleomycin, peplomycin, mitomycin C, mitoxantrone, liposomal doxorubicin;

vinblastine, vincristine, vindesine, paclitaxel, docetaxel, oxaliplatin, carboplatin, cisplatin, nedaplatin;

camptothecin, irinotecan, nogitecan, SN-38, etoposide, levofloxacin, ciprofloxacin;

legolafenib, cetuximab, panitumumab, ramsilmab, gefitinib, erlotinib, afatinib, crizotinib, alectinib, ceritinib, libertinib, trastuzumab, lapatinib, pertuzumab, sunitinib, sorafenib, axitinib, pazopanib, nivolumab, pembrolizumab, ipilimumab, vemurafenib, everolimus, temsirolimus, rituximab, bevacizumab, geldanamycin;

angiostatin, endostatin, metastatin, anti-VEGF antibody and VEGFR-2 inhibitor.

7. The conjugate according to claim 3, wherein the one or more components comprise a detectable substance.

8. The conjugate according to claim 7, wherein the detectable substance enables detection of the conjugate in vivo by a means selected from the group consisting of X-ray photography, computed tomography (CT), nuclear magnetic resonance imaging (MRI), ultrasonography, scintigraphy, positron emission tomography (PET), endoscopy and laparoscopy.

9. The conjugate according to claim 7 or 8, wherein the detectable substance is a radioisotope, an MRI enhancer, a radiopaque substance, a contrast agent or a fluorescent substance.

10. The conjugate according to any one of claims 7 to 9, wherein the detectable substance is selected from the group consisting of a radioactive nuclide selected from  $^{18}\text{F}$ ,  $^{51}\text{Mn}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{52}\text{Fe}$ ,  $^{55}\text{Co}$ ,  $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$ ,  $^{82\text{m}}\text{Rb}$ ,  $^{83}\text{Sr}$ ,  $^{86}\text{Y}$ ,  $^{89}\text{Zr}$ ,  $^{94\text{m}}\text{Tc}$ ,  $^{110}\text{In}$ ,  $^{120}\text{I}$ ,  $^{124}\text{I}$ ,  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{67}\text{Cu}$ ,  $^{67}\text{Ga}$ ,  $^{75}\text{Se}$ ,  $^{97}\text{Ru}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{114\text{m}}\text{In}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{169}\text{Yb}$ ,  $^{197}\text{Hg}$  and  $^{201}\text{Tl}$ ; a paramagnetic ion selected from chromium (III), manganese (II), iron(III), iron(II), cobalt(II), nickel(II), copper(II), neodymium (III), samarium (III), ytterbium (III), gadolinium

(III), vanadium (II), terbium (III), dysprosium (III), holmium (III) and erbium (III), lanthanum (III), gold (III), lead (II), bismuth (III);

an iodine compound, a barium compound, a gallium compound, a thallium compound;

rhodamine, fluorescein, Cy dye, Alexa (registered trademark) Fluor, phycoerythrin (PE), allophycocyanin (APC), derivatives thereof and a near infrared fluorescent reagent.

11. A composition comprising the peptide according to claim 1 or 2 or the conjugate according to any one of claims 3 to 10, and a pharmacologically acceptable carrier.

12. A composition for use in the treatment of cancer, comprising the conjugate according to any one of claims 4 to 6, and a pharmacologically acceptable carrier.

13. A composition for use in testing for cancer, comprising the conjugate according to any one of claims 7 to 10, and a pharmacologically acceptable carrier.

14. The composition for use according to claim 12 or 13, wherein the cancer is a solid cancer or a liquid cancer.

15. The composition for use according to claim 14, wherein the

solid cancer is an angiogenic solid cancer.

16. The composition for use according to claim 14 or 15, wherein the solid cancer is cancer of the brain or nervous system, cancer of the head and neck, cancer of the digestive tract, cancer of the urinary or reproductive organ, cancer of the respiratory system, cancer of the breast, cancer of the skin, cancer of the bone, or cancer of the muscle.

17. The composition for use according to any one of claims 14 to 16, wherein the solid cancer is brain tumor, spinal cord tumor, laryngeal cancer, oral cancer, salivary gland cancer, paranasal sinus cancer, thyroid cancer, stomach cancer, esophageal cancer, small intestine cancer, colon cancer, rectal cancer, anal cancer, liver cancer, biliary tract cancer, pancreatic cancer, kidney cancer, kidney cell cancer, bladder cancer, prostate cancer, renal pelvis and ureter cancer, gall bladder cancer, bile duct cancer, testis cancer, penile cancer, uterine cancer, endometrial cancer, uterine sarcoma, cervical cancer, vaginal cancer, vulvar cancer, ovarian cancer, fallopian tube cancer, lung cancer, breast cancer, malignant melanoma, osteosarcoma or rhabdomyosarcoma.

18. The composition for use according to claim 17, wherein the

solid cancer is a benign or malignant brain tumor.

19. The composition for use according to claim 18, wherein the brain tumor is a primary brain tumor or a metastatic brain tumor.

20. The composition for use according to claim 18 or 19, wherein the brain tumor is meningioma, pituitary adenoma, schwannoma, astrocytoma, oligodendroglioma, anaplastic astrocytoma, anaplastic oligodendroglioma, anaplastic oligoastrocytoma or glioblastoma.

21. The composition for use according to claim 14, wherein the liquid cancer is B-cell lymphoma.

22. Use of the conjugate according to any one of claims 4 to 6 for the treatment of cancer.

23. The use according to claim 22, wherein the cancer is a solid cancer or a liquid cancer.

24. The use according to claim 23, wherein the solid cancer is an angiogenic solid cancer.

25. The use according to claim 23 or 24, wherein the solid cancer is cancer of the brain or nervous system, cancer of the head and neck, cancer of the digestive tract, cancer of the urinary or

reproductive organ, cancer of the respiratory system, cancer of the breast, cancer of the skin, cancer of the bone, or cancer of the muscle.

26. The use according to any one of claims 23 to 25, wherein the solid cancer is brain tumor, spinal cord tumor, laryngeal cancer, oral cancer, salivary gland cancer, paranasal sinus cancer, thyroid cancer, stomach cancer, esophageal cancer, small intestine cancer, colon cancer, rectal cancer, anal cancer, liver cancer, biliary tract cancer, pancreatic cancer, kidney cancer, kidney cell cancer, bladder cancer, prostate cancer, renal pelvis and ureter cancer, gall bladder cancer, bile duct cancer, testis cancer, penile cancer, uterine cancer, endometrial cancer, uterine sarcoma, cervical cancer, vaginal cancer, vulvar cancer, ovarian cancer, fallopian tube cancer, lung cancer, breast cancer, malignant melanoma, osteosarcoma or rhabdomyosarcoma.

27. The use according to claim 26, wherein the solid cancer is a benign or malignant brain tumor.

28. The use according to claim 27, wherein the brain tumor is a primary brain tumor or a metastatic brain tumor.

29. The use according to claim 27 or 28, wherein the brain tumor is meningioma, pituitary adenoma, schwannoma, astrocytoma,

85026686

oligodendroglioma, anaplastic astrocytoma, anaplastic  
oligodendroglioma, anaplastic oligoastrocytoma or glioblastoma.

30. The use according to claim 23, wherein the liquid cancer is  
B-cell lymphoma.

Fig. 1






Fig. 2



5

Fig. 3



**Fig. 4**






Fig. 5



5

Fig. 6



**Fig. 7**






Fig. 9




Fig. 10




Fig. 11




Fig. 12

