

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 63/00, B09C 1/10, C02F 3/32	A1	(11) International Publication Number: WO 96/32016 (43) International Publication Date: 17 October 1996 (17.10.96)
(21) International Application Number: PCT/US96/04631 (22) International Filing Date: 4 April 1996 (04.04.96) (30) Priority Data: 08/421,209 13 April 1995 (13.04.95) US 60/006,662 13 November 1995 (13.11.95) US (71) Applicant: RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY [US/US]; Old Queens Somerset Street, New Brunswick, NJ 08901 (US). (72) Inventors: CHET, Ilan; Shikun Ezrahi, 70400 Nes Ziona (IL). SALT, David; 231 Montgomery Street, Highland Park, NJ 08904 (US). BLAYLOCK, Michael, J.; M5 Quincy Circle, Dayton, NJ 08810 (US). RASKIN, Ilya; 48 Alexandria Drive, Manapalan, NJ 07726 (US). (74) Agent: JARRELL, Brenda, Herschbach; Choate, Hall & Stewart, Exchange Place, 53 State Street, Boston, MA 02109 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS OF PROMOTING PHYTOREMEDIATION (57) Abstract Methods and compositions or enhancing metal uptake of plants, such as members of the family Brassicaceae, comprise treating the roots, plants, seeds, and/or soil in which the plants are grown, with metal-uptake altering microorganisms, preferably of the bacterial genus <i>Pseudomonas</i> and <i>Bacillus</i> .		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5 **METHODS OF PROMOTING PHYTOREMEDIATION**

BACKGROUND OF THE INVENTION

 Agricultural growers and researchers have developed methods of improving root crop yield, and it has been known for many years that some soil
10 microorganisms that are beneficial to plant growth may be added to the soil to either enhance production of plant hormones, produce antibiotics that kill harmful soil organisms, or aid in incorporation of inorganic mineral nutrients, such as iron and manganese, into the plant.

 In a commonly used method, root crop production is enhanced, and root crop
15 diseases, such as root rot, are inhibited by treating the seeds, seed pieces, or roots, or by inoculating the soil, with bacteria isolated from roots or tubers and selected for their ability to inhibit the growth of common plant pathogens. Such growth-promoting root and tuber-associated bacteria are subsequently introduced into the soil surrounding a root or tuber crop, usually at concentrations much higher than
20 naturally present. The method is particularly useful for introducing such bacteria into soils where they are not naturally present.

 Recently, researchers have developed methods of using crop plants to remove pollutants such as heavy metals from soils and waters. See, for example, U.S. Patent Number 5,393,426 and Raskin et al., Current Opinion in Biotechnology, 5: 285-290
25 (1994). These methods are, however, subject to the normal constraints imposed on conventional agronomic practices by bad weather, poor soil drainage, inadequate soil fertilization, and the like.

SUMMARY OF THE INVENTION

30 The present invention solves these problems and is based, in part, on our discovery that plant uptake of heavy metals can be altered (i.e. enhanced or inhibited) by the administration, directly to the soil/plant environment, of isolated microorganisms that are associated with plant roots. These microorganisms have the ability to tolerate heavy metals in their growth medium.

35 In one aspect, a method for altering uptake of a heavy metal by a plant includes contacting a plant with an isolated microorganism that is capable of altering

heavy metal uptake of the plant, and then incubating the plant and isolated microorganism for a time and under conditions sufficient for the isolated microorganism to alter heavy metal uptake of the plant. It is preferred that the isolated microorganism is obtained from an environment of use that contains a concentration of at least one heavy metal. The isolated microorganisms are obtained from the environment of use by extracting microorganisms from a rhizosphere portion of the plant; exposing the extracted microorganisms to selective growth conditions such that only those microorganisms capable of growing in the presence of heavy metals will proliferate; and purifying the selected microorganisms.

10 Preferred microorganisms for use in the present method include bacteria and fungi.

An additional method for altering heavy metal uptake of a plant includes obtaining a series of isolated microorganisms from a heavy metal-containing environment of use, preferably an environment that includes a rhizospheric portion of the plant. Next, one determines to what extent individual members of the isolated microorganisms alter heavy metal uptake of the plant. At least one member of the series of isolated microorganisms is chosen that is capable of altering heavy metal uptake by the plant and this member is re-introduced back into the environment of use. The environment of use may include a hydroponic environment.

15

A further embodiment of the invention is a composition for altering heavy metal uptake by a plant. The composition includes at least one isolated microorganism capable of altering heavy metal uptake of the plant in combination with an agronomically acceptable carrier. A preferred composition includes bacteria of the genus *Pseudomonas* or the genus *Bacillus* in a concentration from about 10^3 cells/ml to about 10^{10} cells/ml in a carrier that may be a seed of the plant, a wettable powder, a dust, or granules. A method of using the composition to alter heavy metal uptake of a plant includes applying an effective amount of the composition to a plant seed prior to planting of the seed, applying an effective amount of the composition to soil where a plant seed is to be planted, or applying an effective amount of the composition to roots of the plant.

20

25

30

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph illustrating total cadmium accumulation in shoots of 2-

week old B. juncea seedlings grown hydroponically in the presence of isolated bacterial strains, previously selected from the rhizosphere of B. juncea. "B.t.I" denotes Bacillus thuringiensis strain I; "B.t.II" denotes B. thuringiensis strain II; "P.p." denotes Pseudomonas putida; "P.c." denotes Pseudomonas chlororaphis; and
5 "C" denotes control treatment.

Figure 2 is a graph illustrating total cadmium removal from solution (bottom graph) and total cadmium shoot concentration in shoots of 2-week old B. juncea seedlings grown hydroponically in the presence of isolated bacterial strains (X-axis), previously selected from the rhizosphere of B. juncea. The designation "con" refers
10 to the control experiment. Strains "8", "10" and "19" refer to Bacillus thuringiensis strain I; Pseudomonas putida; and B. thuringiensis strain II, respectively.

Figure 3 illustrates the effect of isolated bacterial isolates on total lead accumulation during phytoextraction by Brassica juncea.

Figure 4 illustrate the effect of isolated bacterial isolates on total cadmium
15 accumulation during phytoextraction by Brassica juncea.

DETAILED DESCRIPTION OF THE INVENTION

The present invention pertains to methods of using isolated microorganisms selected from the rhizosphere of plants growing in environments containing amounts
20 of heavy metals. Specifically, we have discovered that microorganisms may be isolated from soils contaminated with heavy metals, and when the isolated microorganisms are applied to the plant under appropriate conditions during phytoremediation, heavy metal uptake of the plant is altered.

The term "heavy metal" refers to soluble or insoluble metals. The term
25 "soluble metals" preferably refers to soluble metal cationic or anionic species, which may be soluble in solution at environmentally relevant temperatures (e.g., greater than 0 degrees C to less than about 45 degrees C). "Insoluble metals" refers to metal ions that are substantially insoluble in solution at environmental relevant temperatures and also includes non-ionic, elemental forms of the metal. The present
30 method is useful for enhancing phytoremediation of metals selected from the commonly known heavy metals and radioactive metals such as zinc, cadmium,

chromium, lead, copper, silver, nickel, aluminum, manganese, arsenic, mercury and selenium, as well as radionuclides such as strontium, uranium, cesium, and others.

The term "alter" refers to the ability of the microorganisms to either increase or decrease the uptake of heavy metal by the plant, when compared to a plant
5 lacking the population of isolated microorganisms.

The term "microorganisms" means prokaryotic or eukaryotic unicellular or multicellular organisms, and includes bacteria, actinomycetes, fungi, protozoans, slime molds and blue-green algae. The exact nature and type of microorganism is not intended to limit the methods of the invention. The term "isolated
10 microorganism" means that the particular microorganism has been removed from its natural habitat, exposed to selective growth conditions, i.e., grown in the presence high concentrations of a heavy metal, and purified (see Example 1). Preferably, the term "isolated" refers to pure and mixed cultures of naturally occurring microorganisms that have been selected for their ability to grow and tolerate
15 concentrations of heavy metals, as well as naturally occurring microorganisms that have been genetically manipulated so that they contain RNA or DNA polymers, portions of genomic nucleic acid, cDNA, synthetic nucleic acid and/or polypeptides encoded by these nucleic acids that: (i) are not associated with all of a nucleic acid or polypeptide with which it is associated in nature; or (ii) are linked to a nucleic
20 acid, polypeptide, or other chemical moiety other than that to which it is linked in nature; or (iii) do not exist in nature. Methods of genetically manipulating bacteria are well known in the art (see, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, NY, 1989, incorporated herein by reference).

25 As but one example of the genetic manipulations useful in the present context, it is well known that many plants and microorganisms synthesize high-affinity metal transport systems (e.g., iron, zinc, copper, manganese) that utilize biosynthetic chelating agents called phytosiderophores and siderophores, respectively. Some microorganisms, such as *Escherichia coli* and *Streptomyces*
30 *pilosus*, may actually use other siderophores in addition to their own (Crowley et al., *Plant and Soil*, 130: 179-198, 1991). Thus, plant DNA sequences encoding upon

expression for one or more phytosiderophores can be introduced into isolated rhizosphere microorganisms in order to alter metal uptake of a plant.

The term "rhizosphere" is a term commonly used by persons having ordinary skill in the art and refers to that environment adjacent to the roots of a plant,
5 regardless of whether the plant is growing in soil or in water.

The term "phytoremediation" refers to the practice of using plants to remove pollutants from the environment. The term "phytoremediation" is generic to methods that may be used for plants growing in soils ("phytoextraction": see United States Patent 5,364,451, issued November 15, 1994, incorporated herein by
10 reference) or plants grown hydroponically in water ("rhizofiltration": see United States Patent 5,393,426, issued February 28, 1995, incorporated herein by reference).

The plants used in the preferred methods of the present invention may be either terrestrial plants or aquatic plants. The term "terrestrial plants" refers to
15 photosynthetic plants that normally grow in soils or sediments, but that may be grown hydroponically. The soils or sediments can include a variety of soil types having wide ranges of water and organic matter content. Terrestrial plants can therefore include crop-related plants and/or plants associated with environments such as wetlands, as well as portions of such plants such as excised roots or shoots.
20 "Aquatic plants" are those plants that spend their entire life cycle completely floating on, or submerged in, an aqueous solution, such as floating ferns (e.e., Azolla), duckweed (Lemna sp.), and water hyacinth (Eichhornia), as well as isolated plant cells or cell suspensions capable of metal uptake.

Plants particularly suitable for the present invention include turfgrasses and
25 members of the family Brassicaceae as well as the common sunflower, Helianthus annuus L. Exemplary turfgrasses include Colonial bentgrass, Kentucky bluegrass, perennial ryegrass, creeping bentgrass, a variety of fescues and lovegrasses, Bermudagrass, Buffalograss, centipedegrass, switch grass, Japanese lawngrass and coastal panicgrass. Members of the Brassicaceae include Brassica juncea and B.
30 oleracea. Other plants include spinach, sorghum, tobacco and corn. Plants may also include those plants that are selectively bred and/or genetically engineered.

Microorganisms useful in the present invention may be identified by first isolating a number of microbial strains from the vicinity of healthy roots, usually from the surfaces of such roots, by conventional microbiological methods (see, generally, Elad and Chet, 1987, "The Role of Competition for Nutrients in
5 Biocontrol of Pythium damping off by bacteria", *Phytopathology*, 77: 190-195; Ordentlich et al., 1988, "Rhizosphere Colonization by *Serratia marcescens* for control of *Sclerotium rolfii*, *Soil. Biol. Biochem.*, 19: 747-751, each of which is incorporated herein by reference).

Microbial strains are directly selected for an ability to grow in high
10 concentrations of heavy metals by successive rounds of supplementing microbial growth media with increasing concentrations of one or more heavy metals. Microbial colonies that are able to grow on heavy metal-contaminated media are reisolated on the same media and purified using conventional procedures. Microorganisms that have been subject to the selection pressure of heavy metal
15 exposure are further selected for an ability to alter uptake of heavy metals by plants under particular plant growth conditions (see Examples 2-4).

Exemplary bacteria useful for the invention may be from the genus *Bacillus* and *Pseudomonas*. We have discovered strains of *Pseudomonas putida*,
Pseudomonas chlororaphis, and *Bacillus thuringiensis* to be useful, although it will
20 be appreciated that the specific strain of bacteria is not intended to limit the applicability of the method.

In general, the methods of the present invention involve application of a single type of isolated microorganism, alone or in combination with one or more other isolated microorganisms, to seeds, seed pieces or roots of plants such as sunflower,
25 Brassicas, potatoes, sugar beets, radishes and the like. Concentrations of at least about 10^3 cells/ml can be used and preferably from about 10^5 to about 10^9 cells/ml can be used in combination with an agronomically acceptable carrier medium. A paste may be used to apply to the seeds at a concentration up to 10^{10} cells/ml. While it is preferred to apply the microorganisms directly to the seeds, or portions of plant
30 prior to planting, the microorganisms can also be used to colonize soil in the area of planting prior to planting. In particular, they may be used to recolonize soil which has been fumigated or pasteurized.

The microorganisms may be utilized effectively in diverse formulations, including agronomically acceptable adjuvants and carriers normally employed for facilitating the dispersion of active ingredients for agricultural applications, recognizing as known fact that the dosage, formulation, mode of application, and other variables may affect its activity in any given application. Thus, the previously described exemplary pseudomonads and bacilli may be formulated as a suspension or dispersion, in aqueous or nonaqueous media, as a dust, as a wettable powder, as an emulsifiable concentrate, as a granule, or as any of several other known types of formulations, depending on the desired mode of application. These compositions may be applied as sprays, dust, or granules to the seeds, seed pieces, roots, plants, soil, or plant situs against which activity is desired. Microorganisms may also be added directly to the soil, in potting mixtures as well as directly to aqueous hydroponic systems.

In order to prepare compositions in the form of dust, granules, water dispersible powders, aqueous dispersions, or emulsions and dispersions in organic liquids, the carrier or diluent agent in such formulations may be a finely divided solid, an organic liquid, water, a wetting agent, a dispersing agent, or emulsifying agent, or any suitable combination of these. Generally, when liquids and wettable powders are prepared, a conditioning agent comprising one or more surface-active agents or surfactants is present in amounts sufficient to render a given composition containing the active material, the microorganism, dispersible in water or in oil.

Microorganisms are obtained as described herein and cultured by standard procedures. To convert the desired microorganism to a form which will facilitate the preparation of the following described compositions, a slurry is prepared which is then dried into a primary, agronomically acceptable carrier such as vermiculite, whereby the microorganism is adsorbed onto the carrier, becomes the concentrate for preparing the desired composition. If desired, the slurry can be used as the concentrate for fungal antagonist compositions.

The surface active agent used in the invention can be a wetting, dispersing, or emulsifying agent which will assist dispersion of the effective composition. The surface active agent or surfactant can include such anionic, cationic, and nonionic agents as have heretofore been generally employed in plant control compositions of

similar types. Suitable surface-active agents are set forth, for example, in "Detergents and Emulsifiers", 1971 Annual by John W. McCutcheon, Inc.

In general, 1-10% by weight of the surface-active agent will be used in compositions of this invention and ordinarily the amount of surface-active agent will
5 range from 1-5%, but may even be less than 1% by weight.

Additional surface-active agents can be added to formulations to increase the ratio of surfactants to active ingredients up to as high as 5:1 by weight. Such compositions may have a greater biological effectiveness than can be expected when the components are used separately. When used at higher ratios, it is preferred that
10 the surfactant be present in the range of one-fifth to five parts surfactant for each one part of active agent. More specific formulations are disclosed in Examples 6 to 10, presented hereinafter.

EXAMPLES

15

EXAMPLE 1: Selection and Identification of Microorganisms

I. Selection of Microorganisms

Phytoextraction: Plants growing in heavy metal contaminated soil were collected from sites in New Jersey (Jersey City and Trenton NJ). The roots were cut and
20 washed in 1L tap water to remove the soil particles. The roots were then shaken in about 100 ml sterile water. Aseptic scrapings were also taken of the root surface and placed in about 2ml sterile water. Aliquots of the sterile water containing the scrapings were diluted further for use in selection media, below.

Bacteria were selected on Pseudomonas agar F (mfg by Difco, Inc.) or soil
25 extract agar and fungi were selected on Potato Dextrose agar (obtained from Difco, Inc.). All selection media were supplemented with 20 ug/ml of one or more of the following heavy metals: cadmium as CdSO_4 , chromium (VI) as $\text{K}_2\text{Cr}_2\text{O}_7$ and lead as $\text{Pb}(\text{NO}_3)_2$.

After bacterial colonies were observed, they were picked using conventional
30 sterile microbiological techniques and subject to further rounds of selection on Pseudomonas F agar supplemented with 1 ug/ml of one or more of the following heavy metals: cadmium as CdSO_4 , lead as $\text{Pb}(\text{NO}_3)_2$, chromium (VI) as $\text{K}_2\text{Cr}_2\text{O}_7$,

selenium as Na_2SeO_4 , manganese as MnCl_2 , nickel as $\text{Ni}(\text{NO}_3)_2$, and arsenic as NaAsO_2 . Growing bacteria were reisolated to check the purity of the colony by re-streaking the bacteria onto the same media.

5 After fungal colonies were observed, colonies were purified by taking single spores from the colony or hyphal tips using conventional microbiological sterile technique and growing them on potato dextrose agar containing heavy metal, as above.

Rhizofiltration: Plants growing in heavy metal contaminated water were collected
10 and the roots were cut and washed in 1L tap water to remove the soil particles. The roots were then shaken in about 100 ml sterile water. Aseptic scrapings were also taken of the root surface and placed in about 2ml sterile water. Aliquots of the sterile water containing the scrapings were diluted further for use in selection media, as described below.

15 Bacteria were selected on Pseudomonas agar F (mfg by Difco, Inc. address) or soil extract agar and fungi were selected on Potato Dextrose agar (obtained from Difco, Inc.).

All selection media were supplemented with 20 ug/ml of one or more of the following heavy metals: cadmium as CdSO_4 , chromium (VI) as $\text{K}_2\text{Cr}_2\text{O}_7$ and lead as
20 $\text{Pb}(\text{NO}_3)_2$.

After bacterial colonies were observed, they were picked using conventional sterile microbiological techniques and subject to further rounds of selection on Pseudomonas F agar supplemented with 1 ug/ml of one or more of the following heavy metals: cadmium as CdSO_4 , lead as $\text{Pb}(\text{NO}_3)_2$, chromium (VI) as $\text{K}_2\text{Cr}_2\text{O}_7$,
25 selenium as Na_2SeO_4 , manganese as MnCl_2 , nickel as $\text{Ni}(\text{NO}_3)_2$, and arsenic as NaAsO_2 . Growing bacteria were reisolated to check the purity of the colony by re-streaking the bacteria onto the same media.

After fungal colonies were observed, colonies were purified by taking single spores from the colony or hyphal tips using conventional microbiological sterile
30 technique and growing them on potato dextrose agar containing heavy metal, as above.

II. Identification of Selected Microorganisms

Identification selected microorganisms was carried out by comparing the distribution of fatty acids using gas chromatography according to the methods of Sasser, M., "Identification of Bacteria Through Fatty Acid Analysis", pp. 199-204 in
5 Methods in Phytobacteriology, (ed. Z. Klement, K. Rudolph, and D. Sands), 1990, Akademia Kiado Budapest, incorporated herein by reference.

EXAMPLE 2: Growth of Selected Microorganisms and Application to Plant

Phytoextraction: Bacteria and fungi were grown in liquid *Pseudomonas* F medium
10 and Potato Dextrose medium, respectively, in 250 ml Erlenmeyer flasks containing 100 ml medium with continuous shaking (about 135 rpm) for about 24 hours. Bacteria were collected by centrifugation at about 10,000 rpm, and fungi were collected by centrifugation at about 6,000 rpm. The respective pellets were washed twice by resuspension in 0.1M phosphate buffer (pH 7.0), followed by
15 centrifugation.

Microbial water suspensions were added to phytoextraction systems by irrigation up to the field capacity of the soil. Suspensions contained 10^8 bacteria/ml or 10^8 CFU (colony forming units)/ml of fungi.

20 *Rhizofiltration:* Bacteria were grown in liquid *Pseudomonas* F medium and collected, as above. Bacterial suspensions (about 10^8 bacteria/ml as a final concentration) were added to rhizofiltration systems by immersing hydroponically-grown roots in the suspension and then adding the rest of the suspension to the hydroponic solution itself.

25

EXAMPLE 3: Greenhouse Screening for Altered Phytoextraction

Strains of selected microorganisms are grown on appropriate media (e.g., *Pseudomonas* liquid for bacteria; potato dextrose broth for fungi) for 48 hr, centrifuged, washed and resuspended. The pellet is diluted to give suspensions of
30 approximately 10^8 - 10^9 bacteria or CFU/ml. Several uniform *Brassica juncea* seedlings are dipped for 5 min. in each suspension and planted in pots containing loamy sand field soil. Fresh weights of roots and shoots are taken from 4 week-old

plants. These experiments are repeated three times with strains that alter heavy metal uptake.

EXAMPLE 4: Screening for Altered Rhizofiltration

5 I. Methods

Alteration of Cadmium Uptake: Seeds of *Brassica juncea* were germinated and grown hydroponically with the roots growing in an aerated nutrient solution (a diluted and modified solution after Hoagland and Arnon, Univ. California Agricultural Experiment Station Circular 347- 1938), The modified Hoagland's
10 solution contained ammonium phosphate monobasic (0.25 mol m^{-3}); boric acid (12.5 mmol m^{-3}); calcium nitrate (0.7 mol m^{-3}); copper sulfate pentahydrate ($0.075 \text{ mmol m}^{-3}$); ferric tartrate (4.75 mmol m^{-3}); magnesium sulfate anhydrous (0.5 mol m^{-3}); manganese chloride tetrahydrate (2.25 mmol m^{-3}); potassium nitrate (1.5 mol m^{-3}); and zinc sulfate heptahydrate (0.2 mmol m^{-3}). After about 1 week,
15 bacterial or fungal suspensions were added to the roots of the seedlings, as described above, and plants allowed to grow for a further week. The 2 week old seedlings were then exposed to 1 micromolar cadmium containing ^{109}Cd as a tracer, in modified Hoagland's solution for about 24 hours. Shoots and roots were harvested, washed and oven dried. Dried plant material was then ground and analyzed for Cd
20 using a gamma counter with a solid NaI crystal scintillant.

Alteration of Lead Uptake: Seeds of *Brassica juncea* are germinated and grown hydroponically with the roots growing in an aerated Hoagland's nutrient solution as described above. After about 1 week, bacterial or fungal suspensions are added to
25 the roots of the seedlings, as described above, and plants allowed to grow for a further week. Shoots and roots are harvested, washed and oven dried. Plants are then ashed at 500°C and the solid material dissolved in a mixture of HCl and HNO_3 . Lead is then measured in the solubilized plant material using Direct Current Plasma Optical Emission Spectroscopy (Fisons Model SS-7).

30

II. Results and Discussion

Rhizofiltration: Figure 1 illustrates the ability of Bacillus and Pseudomonas strains to enhance rhizofiltration of cadmium in B. juncea seedlings as compared to a control treatment in which the roots contained only endogenous rhizosphere bacteria and did not have isolated Bacillus and Pseudomonas strains added back to the roots. Figure 2 is a series of graphs from another series of rhizofiltration experiments on B. juncea. Evidence of altered cadmium uptake is evidenced by the increased cadmium removal from solution (as compared to controls) and increased cadmium in the shoot (as compared to controls), when plotted against the isolated microbial strain applied to the plant. Most microbial isolates tended to increase accumulation of cadmium in plant shoots, when compared to the control treatments. Although variability in cadmium distribution within the plant shoot is to be expected, the results are not so variable as to mask the dramatic alteration in cadmium uptake due to some isolates.

Phytoextraction: Bacteria and fungi were grown in liquid Pseudomonas F medium and Potato Dextrose medium, respectively, in 250 ml Erlenmeyer flasks containing 100 ml medium with continuous shaking (about 135 rpm), as above. Microbial water suspensions were added to laboratory-scale phytoextraction systems by irrigation up to the field capacity of the soil. Suspensions contained 10^8 bacteria/ml or 10^8 CFU (colony forming units)/ml of fungi. Figures 3 and 4 illustrate the effect of isolated bacterial isolates on total Brassica juncea accumulation during phytoextraction of lead and cadmium, respectively. Strain numbers 8 and 10 are B. thuringiensis Strain I and P. putida, respectively. The "sterile control" consisted of sterile soil. The "control" consisted of soil containing only endogenous rhizosphere bacteria and did not have isolated Bacillus and Pseudomonas strains added back to the soil. All isolates generally effected a net enhancement of metal removal from soil.

EXAMPLE 5: Establishment of Microorganisms on Root Surfaces and in the Rhizospheres of Treated Plants

Microbial strains (e.g., B. thuringiensis strain I) are extracted, selected and purified from freshly dug roots of, for example, Brassica juncea using methods

- described herein. Roots are dipped for 5 minutes in suspensions containing approximately 10^9 CFU/ml of microbial strain. Roots are then placed on paper towels in the laboratory and sampled at various time intervals to determine the effect of air-drying on the survival of the inoculum. Laboratory temperature and relative humidity is about 24° C and 20% RH during the air-drying period. Population determinations are made by removing root tissue from treated roots and placing them in 100 ml sterile distilled water. Serial water dilutions are subsequently made and plated on growth medium. One piece of root tissue from each of three roots is sampled at each time interval and three dilution plates are made from each peice.
- 10 Plates are incubated for 24 hr at 28° C.

The ability of the microorganisms to survive on roots in field soil, or during hydroponic growth, is determined by previously described inoculation and sampling procedures. Generally, sampling of roots is made by excising one cm root sections from the tips and midsections of three roots per plant and washing them individually (along with soil particles in the case of phytoextraction experiments) with about 1 ml sterile distilled water. About 0.1 ml of these suspensions is plated on the appropriate medium. The population of microorganisms on roots of treated and non-treated roots may be followed over time using this procedure.

20 EXAMPLE 6: Dusts

Dusts are dense powder compositions which are intended for application in dry form. Dusts are characterized by their free- flowing and rapid settling properties so that they are not readily windborne to areas where their presence is not desired. They contain primarily an active ingredient and a dense, free-flowing solid extender. Their performance is sometimes aided by the inclusion of a wetting agent, and convenience in manufacture frequently demands the inclusion of an inert absorptive grinding aid.

The wettable powder as described above can also be used in the preparation of dusts. While such wettable powders can be used directly in dust form, it is more advantageous to dilute them by blending with the dense dust diluent. In this manner, dispersing agents, corrosion inhibitors, and antifoam agents may also be used as components of a dust.

Thus, the dust compositions of this invention can comprise from about 0.5 to 20.0 weight percent active ingredient, 5 to 25 weight percent filler, 0 to 1.0 weight percent wetting agent and from about 30 to 90 weight percent dense, free-flowing extender, as these terms are used herein. Such dust formulations can contain, in addition, minor amounts of dispersants, corrosion inhibitors, and antifoam agents derived from the wettable powders used to make the dust.

EXAMPLE 7: Wettable Powders

Wettable powders are water-dispersible compositions containing the active material, an inert solid extender, and one or more surfactants to provide rapid wetting and to prevent heavy flocculations when suspended in water. The inert extenders which are preferred for use in the wettable powders of this invention containing the 25 active compounds are of mineral origin.

Extenders suitable for the wettable powder formulations of this invention are the natural clays, diatomaceous earth and synthetic mineral fillers derived from silica and silicate. Most preferred fillers for this invention are kaolinites, attapulgite clay, montmorillonite clays, synthetic silica, synthetic magnesium silicate and calcium sulfate dihydrate.

Among the more preferred surfactants are the non-ionic and anionic types. They are most suitable for the preparation of dry, wettable products of this invention and dispersants. Occasionally, a liquid, nonionic compound which is primarily an emulsifier, may serve as both wetter and dispersant.

Most preferred wetting agents are alkylbenzene and alkylnaphthalene sulfonates, sulfated fatty alcohols, amines or acid amides, long chain esters of sodium isothionate, esters of sodium sulfosuccinate, sulfated or sulfonated vegetable oils, and ditertiary acetylenic glycols. Preferred dispersants are methyl cellulose, polyvinyl alcohol, lignin sulfonates, polymeric alkylnaphthalene sulfonates, sodium naphthalene sulfonates, polymethylene bisnaphthalene sulfonate and sodium-N-methyl-N-(long chain acid) taurates.

Wetting and dispersing agents in these preferred wettable powder compositions of the invention are usually present at concentrations of from about 0.5

weight percent to 5 weight percent. The inert extender then completes the formulation. Where needed, 0.1 weight percent of the extender may be replaced by a corrosion inhibitor or an antifoaming agent or both.

Thus, wettable powder formulations of the invention will contain from about
5 25 to 90 weight percent active material, from 0.5 to 2.0 percent wetting agent, from 0.25 to 5.0 weight percent dispersant, and from 9.25 to 74.25 weight percent inert extender, as these terms are described above.

When the wettable powder contains a corrosion inhibitor or an antifoaming agent or both, the corrosion inhibitor should not exceed about 1 percent of the
10 composition, and the antifoaming agent should not exceed about 0.5 percent by weight of the composition, both replacing equivalent amounts of the inert extender.

EXAMPLE 9: Emulsifiable Oils

Emulsifiable oils are usually solutions or suspensions of active material in
15 nonwater miscible solvents together with a surfactant and/or emulsifier. For compositions of this invention, emulsifiable oil compositions can be made by mixing the active ingredient with an organic solvent and surfactant. Suitable surfactants are those ionic or nonionic agents known to the art as emulsifying agents.

Emulsifying agents most suitable for the emulsifiable oil compositions of this
20 invention are long chain alkyl or mercaptan polyethoxy alcohols, alkylaryl polyethoxy alcohols, sorbitan fatty acid esters, polyoxyethylene ethers with sorbitan fatty acid esters, polyethylene glycol esters with fatty rosin acids, fatty alkylol amide condensates, calcium and amine salts of fatty alcohol sulfates, oil soluble petroleum sulfonates, or preferably mixtures of these emulsifying agents. Such emulsifying
25 agents should compromise from about 1 to 10 weight percent of the total composition. As described above, however, up to 5 parts of emulsifying agent for each part of active ingredient can be used.

Thus, emulsifiable oil compositions of the present invention can consist of from about 10 to 50 weight percent active ingredients, about 40 to 82 percent
30 solvents, and about 1 to 10 weight percent emulsifies, as these terms are defined and used above.

EXAMPLE 10: Granules

Granules are physically stable, particulate compositions containing mycelium, sclerotia, spores, or other microorganisms which adhere to, or are distributed through, a basic matrix of a coherent, inert carrier with microscopic dimensions. In order to aid leaching of the active ingredient from the granule, a surfactant can be present.

The inert carrier is preferably of mineral origin, and suitable carriers are natural clays, some pyrophyllites and vermiculite. Suitable wetting agents can be anionic or nonionic.

For the granule composition of this invention, most suitable carriers are of two types. The first are porous, absorptive pre-formed granules, such as preformed and screened granular attapulgite or heat expanded granular, screened vermiculite. On either of these, a suspension of the active agent can be sprayed and will be absorbed at concentrations up to 25 weight percent of the total weight. The second type are initially powdered kaolin clays, hydrated attapulgite or bentonite clays in the form of sodium, calcium or magnesium bentonites. Water-soluble salts such as sodium salts may also be present to aid in the disintegration of the granules in the presence of moisture. These ingredients are blended with the active components to give mixtures that are granulated, followed by drying to yield formulations with the active component distributed uniformly throughout the mass. Such granules can also be made with 25 to 30 weight percent active component but more frequently a concentration of about 10 weight percent is desired for optimum distribution. The granular compositions of this invention are believed to be most useful in a size range of 15-30 mesh.

The most suitable wetting agents for the granular compositions of this invention depend upon the type of granule used. When pre-formed granules are sprayed with active material in liquid form, the most suitable wetting agents are nonionic, liquid wetters miscible with the solvent. These are more generally known in the art as emulsifiers and comprise alkyl-aryl polyether alcohols, alkyl polyether alcohols, polyoxyethylene sorbitan fatty acid esters, polyethylene glycol esters with fatty or rosin acids, fatty alkylol amide concentrates, oil soluble

petroleum or vegetable oil sulfonates, of mixtures of these. Such agents will usually compromise up to about 5 weight percent of the total composition.

When the active ingredient is first mixed with a powdered carrier and subsequently granulated, liquid non-ionic wetters can still be used, but it is usually preferable to incorporate at the mixing stage, one of the solid, powdered anionic wetting agents such as those previously listed for the wettable powders. Such agents should compromise about 0 to 2 weight percent of the total composition.

Thus, the preferred granular formulations of this invention compromise about 5 to 30 weight percent active material, about 0 to 5 weight percent wetting agent, and about 65 to 90 percent inert mineral carrier, as these terms are used herein.

EQUIVALENTS

Those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents to the specific products and processes described herein. Such equivalents are considered to be within the scope of the invention and are intended to be covered by the following claims.

We claim:

1. A method for altering uptake of a heavy metal by a plant, comprising:
contacting a plant with an isolated microorganism that is capable of altering
5 heavy metal uptake of the plant; and
incubating the plant and isolated microorganism for a time and under
conditions sufficient for the isolated microorganism to alter heavy metal uptake of
the plant.
- 10 2. The method of claim 1, wherein the step of contacting comprises contacting
with an isolated microorganism obtained from an environment of use that contains a
concentration of at least one heavy metal.
3. The method of claim 2, wherein the step of incubating comprises incubating
15 the plant and isolated microorganism in the environment of use.
4. The method of claim 2, wherein the isolated microorganisms are obtained
from the environment of use by a method comprising:
extracting microorganisms from a rhizosphere portion of the plant;
20 exposing the extracted microorganisms to selective growth conditions such
that those microorganisms capable of growing in the presence of heavy metals will
proliferate; and
purifying the selected microorganisms.
- 25 5. The method of claim 3, wherein the step of incubating comprises incubating
in a soil environment.
6. The method of claim 3, wherein the step of incubating comprises incubating
in a hydroponic environment.
- 30 7. The method of claim 1, wherein the isolated microorganism is a
microorganism selected from the group consisting of bacteria and fungi.

8. The method of claim 7, wherein the isolated bacteria is a member of the bacterial genus *Pseudomonas* or a member of the bacterial genus *Bacillus*.
9. A method for altering heavy metal uptake of a plant, comprising the steps of:
- 5 obtaining a plurality of isolated microorganisms from a heavy metal-containing environment of use, wherein the environment of use includes a rhizospheric portion of the plant;
- determining to what extent individual members of said plurality of isolated microorganisms alter heavy metal uptake of the plant;
- 10 choosing at least one member of said plurality of isolated microorganisms that is capable of altering heavy metal uptake by the plant; and
- re-introducing said at least one member of said plurality of isolated microorganisms back into the environment of use.
- 15 10. The method of claim 9, wherein the step of obtaining comprises:
- extracting microorganisms from a rhizosphere portion of the plant;
- exposing the extracted microorganisms to selective growth conditions such that only those microorganisms capable of growing in the presence heavy metals will proliferate;
- 20 and
- purifying the selected microorganisms.
11. A composition for altering heavy metal uptake by a plant, comprising at least one isolated microorganism capable of altering heavy metal uptake of the plant in
- 25 combination with an agronomically acceptable carrier.
12. The composition of claim 11, wherein the isolated microorganism is a bacteria of the genus *Pseudomonas* or the genus *Bacillus*.
- 30 13. The composition of claim 11, wherein the isolated microorganism is present in a concentration from about 10^3 cells/ml to about 10^{10} cells/ml.

14. The composition of claim 11, wherein the agronomically acceptable carrier comprises a seed of the plant.
15. The composition of claim 11, in the form of a wettable powder.
- 5 16. The composition of claim 11 in a dust.
17. The composition of claim 11 in a form of granules.
- 10 18. A method of altering heavy metal uptake of a plant comprising applying an effective amount of the composition of claim 11 to a plant seed prior to planting of the seed.
- 15 19. A method of altering heavy metal uptake of a plant comprising applying an effective amount of the composition of claim 11 to soil where a plant seed is to be planted.
- 20 20. A method of altering heavy metal uptake of a plant comprising applying an effective amount of the composition of claim 11 to roots of the plant.
21. The method of claim 20, wherein the roots are grown in a hydroponic system.
22. The method of claim 1, wherein the step of contacting comprises contacting
25 with an isolated microorganism that is a genetically manipulated microorganism.




FIG. 1




FIG. 4

2 / 3




FIG.2

3/3




FIG. 3

INTERNATIONAL SEARCH REPORT

Inte onal Application No
PCT/US 96/04631

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A01N63/00 B09C1/10 C02F3/32

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A01N B09C C02F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	W0,A,94 29466 (PHYTOTECH INC) 22 December 1994	1,7,11, 20,22
Y	see page 20, line 32 - page 21, line 20 see page 1, line 8 - page 2, line 8 see page 3, line 21 - line 31 see page 9, line 21 - page 11, line 1 --- -/--	2-5, 8-10, 12-19

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

23 August 1996

Date of mailing of the international search report

04-09-1996

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

Int. onal Application No
PCT/US 96/04631

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOURNAL OF INDUSTRIAL MICROBIOLOGY, vol. 14, 3 April 1995, pages 208-212, XP000577876 B.MÜLLER ET AL.: "Leaching of zinc from an industrial filter dust with Penicillium, Pseudomonas and Corynebacterium: citric acid is the leaching agent rather than amino acids" see SUMMARY see page 208, column 1, paragraph 1 - column 2, paragraph 3 ---	2-6, 8-10, 12-19,21
Y	WO,A,94 29226 (PHYTOTECH INC) 22 December 1994 see the whole document ---	6,21
Y	DE,A,41 00 758 (SCHLUTTIG ALEXANDER DR) 23 July 1992 see claim 3 ---	2-6, 8-10, 12-19,21
X	CHEMICAL ABSTRACTS, vol. 122, no. 1, 2 January 1995 Columbus, Ohio, US; abstract no. 8909, M.K.BANKS ET AL.: "Effects of plants and soil microflora on leaching of zinc from mine tailings" XP002011478 see abstract & CHEMOSPHERE, vol. 29, no. 8, 1994, pages 1691-1699, ---	1,7,11, 13,20
P,X	BIO/TECHNOLOGY, vol. 13, May 1996, pages 468-474, XP000577130 D.E.SALT ET AL. : "Phytoremediation: A Novel Strategy for the Removal of Toxic Metals from the Environment Using Plants" see page 472, column 1, paragraph 2 ---	1-22
P,X	BIOLOGICAL ABSTRACTS, vol. 98, Philadelphia, PA, US; abstract no. 368177, I.WEISSENHORN ET AL.: "Arbuscular mycorrhizal contribution to heavy metal uptake by maize (Zea mays L.) in pot culture with contaminated soil." XP002011476 see abstract & MYCORRHIZA, vol. 5, no. 4, 12 May 1995, pages 245-251, ---	1-5,7, 9-11, 13-20
	-/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/04631

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	BIOLOGICAL ABSTRACTS, vol. 98, Philadelphia, PA, US; abstract no. 798373, J.A.ENTRY ET AL.: "Phytoremediation of soil contaminated with low concentrations of radionuclides" XP002011477 see abstract & WATER AIR AND SOIL POLLUTION, vol. 88, no. 1-2, 1996, pages 167-176, ---	1-5,7, 9-11, 13-20
P,X	ENVIRONMENTAL SCIENCE AND TECHNOLOGY, vol. 29, no. 5, 1 May 1995, pages 1239-1245, XP000505197 DUSHENKOV V ET AL: "RHIZOFILTRATION: THE USE OF PLANTS TO REMOVE HEAVY METALS FROM AQUEOUS STREAMS" see page 1245, column 1, paragraph 2 ---	6,21
A	DE,A,43 19 992 (KUEHN BERNDT ; KOENIG PETER (DE)) 22 December 1994 see column 5, line 21 - line 27 -----	1-22

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. nal Application No

PCT/US 96/04631

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9429466	22-12-94	US-A- 5364451 AU-B- 7097894 CA-A- 2163665 EP-A- 0701621	15-11-94 03-01-95 22-12-94 20-03-96
WO-A-9429226	22-12-94	US-A- 5393426 AU-B- 7049994 CA-A- 2163666 EP-A- 0701537	28-02-95 03-01-95 22-12-94 20-03-96
DE-A-4100758	23-07-92	NONE	
DE-A-4319992	22-12-94	WO-A- 9500264	05-01-95