



US 20070026397A1

(19) **United States**

(12) **Patent Application Publication**
Freskgard et al.

(10) **Pub. No.: US 2007/0026397 A1**

(43) **Pub. Date: Feb. 1, 2007**

(54) **METHOD FOR PRODUCING
SECOND-GENERATION LIBRARY**

(75) Inventors: **Per-Ola Freskgard**, Norrkoping (SE);
Alex Haahr Gouliaev, Veksoe
Sjaelland (DK); **Thomas Thisted**,
Frederikssund (DK); **Eva Kampmann
Olsen**, Herlev (DK)

Correspondence Address:
BROWDY AND NEIMARK, P.L.L.C.
624 NINTH STREET, NW
SUITE 300
WASHINGTON, DC 20001-5303 (US)

(73) Assignee: **Nuevolution A/S**, Copenhagen O (DK)

(21) Appl. No.: **10/546,538**

(22) PCT Filed: **Feb. 23, 2004**

(86) PCT No.: **PCT/DK04/00117**

§ 371(c)(1),
(2), (4) Date: **Jun. 26, 2006**

Related U.S. Application Data

(60) Provisional application No. 60/448,480, filed on Feb. 21, 2003. Provisional application No. 60/448,460, filed on Feb. 21, 2003. Provisional application No. 60/504,748, filed on Sep. 22, 2003.

(30) **Foreign Application Priority Data**

Feb. 21, 2003 (DK)..... PA 2003 00268
Feb. 21, 2003 (DK)..... PA 2003 00269
Sep. 18, 2003 (DK)..... PA 2003 01356


Publication Classification


(51) **Int. Cl.**
C40B 30/06 (2006.01)
C40B 40/08 (2006.01)
(52) **U.S. Cl.** **435/6**

(57) **ABSTRACT**

The present invention relates to a method for generating a second-generation library. In a first step, a library of encoded molecules associated with an identifier nucleic acid comprising codons identifying chemical entities that have participated in the formation of the encoded molecule is provided. In a second step, the library is partitioned and encoded molecules having a certain property are selected. Codons of identifiers of selected encoded molecules are subsequently identified, and a second-generation library is prepared using at least some of the chemical entities coded for by the identified codons. The new focused library may be used of another partition step to select encoded molecules with a certain property.

Fig. 1








Fig. 3






Fig. 5

Panel A



Panel B






Fig. 6

Fig. 7





Fig. 8








Fig. 9

Fig. 10











Fig. 13










Fig. 16




Fig. 17

METHOD FOR PRODUCING SECOND-GENERATION LIBRARY

[0001] Various patent and non-patent references cited in the present application are hereby incorporated by reference in their entirety.

TECHNICAL FIELD OF THE INVENTION

[0002] The invention relates to a method for producing a second-generation compound library with an improved desired property profile. In nature and artificial methods based on the natural system, the parent genotype is carried on to the off-spring and results in a phenotype in which the exact type and sequence of amino acids is retained, unless a mutation and/or recombination has occurred. The present method only retains the identity of chemical entities, e.g. amino acids, while the sequence wholly or partly is scrambled. The result is a focused second-generation library with lower diversity.

BACKGROUND OF THE INVENTION

[0003] The biological evolution is based on the survival of specific genotypes that encode phenotypes with the most suitable functionalities in a certain environment. In all living species DNA programs the genotype. DNA serves two important functions in the natural selection process. One function is obviously to encode for the type of nucleotides used and the other function is to encode for the specific order of nucleotide sequences in a nucleic acid sequence. The strategy used in nature, i.e. encoding for the exact type as well as the precise sequence of nucleotides, ensures an extremely similarity between the progeny and its parents. Thus, conserving almost the exact sequence and type of the nucleotides is absolutely essential in order to create off spring with a high functionality. The changes in the genotype from one generation to another, which allow for evolution, are determined by the random mutation rate and recombination between the two parent's genotypes.

[0004] The natural selection cannot afford too many changes in the DNA from one generation to the next in order to secure survival of the species. Therefore, nature has evolved sophisticated means to proofread the copying of the DNA from the parents to its progeny and secured that the characteristics of phenotype from one generation to the next is carried only by the DNA.

[0005] Within the art of selecting ligands from a library of encoded polypeptides associated with a corresponding identifier nucleic acid sequence, the method of nature is used. Thus, when more than a single library generation is needed, the identifier nucleic acid sequences (genotype) carries the information from one generation to the next.

[0006] WO 93/03172 A1 discloses a method for identifying a polypeptide ligand having a desired property in a polypeptide library. In a first step, a translatable mRNA mixture is provided, which is mixed with a mixture of ribosome complexes to form a translation product attached to the mRNA strand responsible for the formation thereof. In a second step the ribosome complexes binding to a target are partitioned from and remainder of the library. In a third step, an amplification of mRNA strands of the partitioned ribosome complexes, which has bound to the target follows. The amplified mRNA strands are used for the production of a

second generation library, which is subjected to a renewed contact with the target. The method is repeated a sufficient number of times until the size of the library has narrowed to a small pool of high affinity binders.

[0007] In WO98/31700 A1 a method for selecting a DNA molecule, which encodes for a desired protein, is disclosed. The method implies the initial presence of a pool of candidates RNA molecules, which subsequently is translated into a corresponding pool of RNA-protein fusions. Subsequently the mRNA-protein fusion products are subjected to a selection process, i.e. the fusion products are presented for a target molecule, and a new pool of complexes capable of binding to the target are partitioned. From the new pool of complexes, the mRNAs are recovered and amplified for use in a subsequent round of library generation. Xu, L. et al *Chemistry & Biology*, Vol. 9, 933-942, August 2002 discloses a practical embodiment in which a library of more than 10^{12} unique mRNA-protein fusion products through ten rounds of library generation and selection are used to identify a high affinity binding protein.

[0008] The preparation of libraries of synthetic molecules associated with a corresponding identifier nucleic acid sequence, and the selection of synthetic molecules from such libraries, have been the subject of various patent applications. When two or more generations of libraries are needed, the identifier nucleic acid sequence is used as carrier between an initial library and the next generation library.

[0009] Thus, in WO 00/23458A1 libraries of complexes comprising non-natural molecules attached to corresponding nucleic acid sequences are suggested. After a selection of the library has been conducted, the nucleic acid sequences of successful complexes are amplified by PCR and a new library is prepared from these nucleic acid sequences. The same method of carrying information from an initial library to the next library is applied in WO 02/074929A2 and WO 02/103008A2.

[0010] The present invention provides a new method for evolving encoded molecules. The method is based on the identification of chemical entities used in the synthesis of reaction products of successful complexes and the application, at least in part, of these chemical entities in the preparation of the next generation library. The utilization of preferable chemical entities and the exclusion of certain undesired chemical entities in the next library generation generally imply that the next generation library has a smaller size compared to the size of the initial library, thereby, at the same time, retaining the desirable encoded molecules in the library.

SUMMARY OF THE INVENTION

[0011] The present invention concerns a method for producing a composition of molecules with an improved desired property, said method comprising the steps of: providing an initial library comprising a plurality of different encoded molecules associated with a corresponding identifier nucleic acid sequence, wherein each encoded molecule comprises a reaction product of multiple chemical entities and the identifier nucleic acid sequence comprises codons identifying said chemical entities; subjecting the initial library to a condition partitioning members having encoded molecules displaying a predetermined property from the remainder of the initial library; identifying codons of the

identifier nucleic acid sequences of the partitioned members of the initial library; and preparing a second-generation library of encoded molecules using the chemical entities coded for by the codons of the partitioned members of the initial library or a part thereof.

[0012] The present invention relates to a novel approach to perform evolution of molecules with a desired property, said approach being different from the approach of nature and the prior art. The invention is based on the selecting of chemical entities, the counterpart of amino acids in Nature, instead of the precise sequence of chemical entities. This new approach is powerful in *ex vivo* conditions when high functionality of the off spring is not vital for success and when the number of chemical entities relative to the number of reactants used in each encoded molecule is high.

[0013] The method disclosed herein will be increasingly effective as the library size increases. This is due to the fact that more chemical entities is used when a library size is increased, when the number of reactions for the formation of the encoded is fixed and the fact that different chemical entities tend to be involved in encoded molecules having the desired property. The chemical entities, which are part of the final selected molecules, will be enriched in each round of selection. Finally, when the diversity has been extensively reduced, the enriched molecules are decoded from the identifier nucleic acid sequence comprising the codons of the chemical entities that have participated in the formation of the encoded molecule.

[0014] The strategy of performing enrichment of chemical entities instead of specific combinations of chemical entities more efficiently search the chemical space for all combinations of chemical entities that are eager to show a certain property, such as a binding ability towards a target. Thus, chemical entities having a certain impact on the formation of encoded molecules is allowed in a new library to recombine in each new library generation. In a certain aspect of the invention, the recombination is random, *i.e.* once a chemical entity has qualified as being of interest it is allowed in every position of the reaction sequence. In another aspect of the invention, the recombination is semi-random, *i.e.* once a chemical entity is qualified as being of interest it is used in a certain position in the reaction sequence of the encoded molecule. In still a further aspect of the invention, the amount of the chemical entity used in a subsequent library generation is dependent on the frequency and the amount of the partitioned library members.

[0015] The present invention may be of special interest when a group of chemical entities are selected from a larger pool of chemical entities in the formation of a first library. Selecting chemical entities resulting in encoded molecules having a certain property in a first library and spiking with remaining chemical entities of the pool allows for the formation of a second-generation library not necessarily of a smaller size but enriched in encoded molecules having a certain property.

[0016] The second-generation library may be formed of a reaction product of the chemical entities without attaching the reaction product to a nucleic acid. In an embodiment of such second-generation library the individual reaction products are formed in discrete reaction compartments in accordance with traditional combi-chem technology. In a certain aspect of the invention, the second-generation library is

prepared as the first generation library, *i.e.* the second-generation library comprises a plurality of different encoded molecules associated with a corresponding identifier nucleic acid sequence, wherein each encoded molecule comprises a reaction product of multiple chemical entities and the identifier nucleic acid sequence comprises codons identifying said chemical entities.

[0017] In a preferred aspect of the invention, it comprises subjecting the second-generation library to a condition partitioning members having encoded molecules displaying a predetermined property from the remainder of the second-generation library. The second-generation library may be partitioned as to the same property or a different property. Notably, the second-generation library can be screening against the same target or a different target.

[0018] After the partitioning of the second-generation library, the invention comprises the step of deducing the identity of the encoded molecule(s) using the identifier nucleic acid sequence, when present. Optionally, a third or further generation library may be formed and screened before the final deducing step is performed. In a certain embodiment, the decoding includes that the codons of the identifier nucleic acid sequence is decoded to establish the synthesis history of the encoded molecules. The synthesis history includes the identity of the chemical entities used and the point in time they enter the sequence of reactions resulting in the encoded molecule.

[0019] The encoded molecule is preferably a reaction product in which multiple chemical entity precursors have participated. The encoded molecule may have any chemical structure. Generally, the multiple chemical entities are precursors for a structural unit appearing in the encoded molecule. However, the chemical entities may also perform a chemical reaction with the nascent encoded molecule, which result in an altering or removal of chemical groups. In certain aspects of the invention, the encoded molecule is a scaffolded molecule, *i.e.* various chemical entities have reacted with a chemical core structure like steroid, benzodiazepine, retinol, camphor, ephedrine, penicillin, cannabinol, coumarin, oxazol, etc. In certain other aspects of the invention the encoded molecule is fully or partly a polymer. The polymer may be of a type which occurs naturally or may be a non-naturally occurring polymer. Nature only has the possibility of preparing α -polypeptides using the recognition of a codon of an mRNA strand by the anticodon of a charged tRNA. In some aspects of the invention, the encoded molecule is not a α -peptide. Notably, in some aspects of the invention, the chemical entities are reacted without enzymatic interaction to produce the encoded molecule.

[0020] The encoded molecule can be associated with the nucleic acid sequence identifier in any appropriate way. In a certain aspect of the invention, the encoded molecule associated with the corresponding identifier nucleic acid sequence is a bifunctional complex. The bifunctional complex may be formed by covalent or non-covalent attachment of the encoded molecule to the identifier nucleic acid sequence. In another aspect of the invention, an identifier nucleic acid sequence is physically a distinct entity separated from the encoded molecule, wherein the identifier identifies the spatial position of an encoded molecule, *e.g.* in

the same compartment in which an encoded molecule is formed a corresponding identifier oligonucleotide is generated.

[0021] The conditions partitioning complexes of interests from the remainder of the library may be chosen from a variety of possibilities. In one aspect the condition relates to physical parameters, so that complexes displaying a physical stability under e.g. certain temperature conditions, certain acidic conditions, certain radiation conditions etc. are selected from the library. In other aspects of the invention the condition for partitioning the desired complexes includes subjecting the initial library to a molecular target and partitioning complexes binding to this target. The molecular target may be any compound of interest. Exemplary targets are proteins, carbohydrates, polysaccharides, hormones, receptors, antibodies, viruses, antigens, cells, tissues etc. In certain aspects the target is immobilized on a solid support, such as column material and contacted with the candidate complexes in a fluid media followed by a partitioning of the complexes capable of binding to the target under the contacting conditions used. Typically the binding complexes are eluted from the column using increased stringency conditions.

[0022] The complexes as such or only the identifier part is harvested after the partitioning step. Usually the identifier nucleic acid sequences are amplified prior to the identification step. The amplification is suitably performed applying polymerase chain reaction (PCR). The amplified identifiers may be explicitly or implicitly identified. When the codons are identified explicitly, the sequence and identity of nucleotides in the codon is made known to the experimenter, whereas, when the codons of the identifiers are implicitly identified, the experimenter is not presented for the information.

[0023] Any suitable method for identifying codons may be used. In a certain aspect of the invention, traditional sequencing, e.g. by using a modification of the Sangers method or pyrosequencing methods, identifies the codons. In another aspect of the invention, the codons of the identifier nucleic acid sequences of the partitioned members of the initial library are identified by contacting said identifier nucleic acid sequences with a pool of nucleic acid fragments under conditions allowing for hybridisation.

[0024] The pool of nucleic acid fragments may be immobilized or in solution. In a certain aspect of the invention, the pool of nucleic acid fragments comprises a plurality of single stranded nucleic acid probes immobilized in discrete areas of a solid support, wherein the nucleic acid probes are capable of hybridising to a codon of the identifier nucleic acid sequence comprising codons. The nucleic acid probes may be positioned on a microarray, such that the identity of the codons is revealed by observing the discrete areas of the support in which a hybridisation event has occurred.

[0025] The nucleic acid probe can be directly hybridised to the identifier or the nucleic acid probe of the array is hybridised to an identifier nucleic acid sequence through an adapter oligonucleotide having a sequence complementing the probe as well as one or more codons of the identifier nucleic acid sequence. The probe may identify a single codon of an identifier or a probe of the array is capable of hybridising to two codons of the identifier nucleic acid sequence or a sequence complementary to said sequence.

The ability to hybridise two or more codons makes it possible to study the influences of neighbouring chemical entities on each other. In a certain aspect, a nucleic acid probe of the array is capable of hybridising to all codons of an identifier nucleic acid sequence. This latter option will fully decode the identity of the encoded molecule. Usually however, a fully decoding is only possible for a relative small library size, as it presupposes a nucleic acid probe for each member of the library.

[0026] When single codons are detected, useful information about a certain codon may be gathered by detecting the codon together with a framing sequence identifying the position in the reaction history of the chemical entity corresponding to said codon.

[0027] As an example, if a library of complexes is prepared from 100 chemical entities and the three reactions, i.e. each identifier comprises 4 codons, the library size is 10^8 . For most practical uses 10^8 is in the excess of what is possible to detect on an array, especially if multiple determinations for each identifier are considered necessary to obtain a high accuracy. However, an array of just 100 probes complementary to the 100 codons will reveal important information prior to or subsequent to a selection. In the event a framing sequence is detected together with the codon an array of 400 probes is needed.

[0028] A suitable method for identifying an hybridisation event is to use a label. Therefore, in a preferred embodiment, the existence of a hybridisation event is measured through labelling of the identifier nucleic acid sequence, or an amplification product thereof. When the label emits light, the hybridisation event is measured by the emission of light in a scanner. To reveal the relative abundance of each chemical entity in the library of encoded molecules, the relative intensity of light in each discrete spot is measured.

[0029] The measurement of a hybridisation event may be conducted by various methods known in the art. In the event the label emits lights, the presence or absence of a hybridisation event may be measured in a scanner, e.g. a confocal scanner. The scanner may be connected with computer software, which is able to quantify the amount of lights measured. The amount of light measured correlates with the amount of identifier annealed to the probes. Thus, it is possible to measure not only the presence or absence of one or more codons of an identifier; it is also possible to measure the relative amount of the codons in one or more identifiers.

[0030] After the complexes have been partitioned and the specific codons have been identified on the microarray, the information can be used to design optimized libraries including chemical entities based on both the selection data and the chemical structure. The microarray analysis will first of all detect which chemical entities pass the partitioning step. Secondly, the relative intensity on the microarray will reflect the relative binding affinity of the chemical entities. Finally, the structures of the chemical entities are directly identified due to the position of the probes on the array. For instance, chemical entities that are strongly selected in a partitioning process but possess some unfavourable chemical structure can be excluded in the next generation of library. Similarly, chemical entities that are weakly selected in a partitioning process but possess some favourable chemical structure can be included in the next generation of library. Thus, the next generation library design can be based both on a rational

choice of chemical entities with lead-like structures and the selection pressure detected on the microarray.

[0031] Another method of identifying codons includes that nucleic acid fragments are primer oligonucleotides, and the identification involves subjecting the hybridisation complex between the primer oligonucleotides and the identifier nucleic acid sequences to a condition allowing for an extension reaction to occur when the primer is sufficient complementary to a part of the identifier nucleic acid sequence, and evaluating based on measurement of the extension reaction, the presence, absence, or relative abundance of one or more codons.

[0032] The extension reaction requires a primer, a polymerase as well as a collection of deoxyribonucleotide triphosphates (abbreviated dNTP's herein) to proceed. An extension product may be obtained in the event the primer is sufficient complementary to an identifier oligonucleotide for a polymerase to recognise the double helix as a substrate. After binding of the polymerase to the double helix, the deoxyribonucleotide triphosphates (blend of DATP, dCTP, dGTP, and dTTP) are incorporated into the extension product using the identifier oligonucleotide as identifier. The conditions allowing for the extension reaction to occur usually includes a suitable buffer. The buffer may be any aqueous or organic solvent or mixture of solvents in which the polymerase has a sufficient activity. To facilitate the extension process the polymerase and the mixture of dNTP's are generally included in a buffer which is added to the identifier oligonucleotide and primer mixture. An exemplary kit comprising the polymerase and the nNTP's for performing the extension process comprises the following: 50 mM KCl; 10 mM Tris-HCl at pH 8.3; 1.5 mM MgCl₂; 0.001% (wt/vol) gelatin, 200 μM DATP; 200 μM DTTTP; 200 μM dCTP; 200 μM dGTP; and 2.5 units *Thermus aquaticus* (Taq) DNA polymerase I (U.S. Pat. No. 4,889,818) per 100 microliters (μl) of buffer.

[0033] The primer may be selected to be complementary to one or more codons or parts of such codons. The length of the primers may be determined by the length of the codons, however, the primers usually are at least about 11 nucleotides in length, more preferred at least 15 nucleotides in length to allow for an efficient extension by the polymerase. The presence or absence of one or more codons is indicated by the presence of or absence of an extension product. The extension product may be measured by any suitable method, such as size fractioning on an agarose gel and staining with ethidium bromide.

[0034] In a preferred embodiment the admixture of identifier oligonucleotide and primer is thermocycled to obtain a sufficient number of copies of the extension product. The thermocycling is typically carried out by repeatedly increasing and decreasing the temperature of the mixture within a temperature range whose lower limit is about 30 degrees Celsius (30° C.) to about 55° C. and whose upper limit is about 90° C. to about 100° C. The increasing and decreasing can be continuous, but is preferably phasic with time periods of relative temperature stability at each of temperatures favouring polynucleotide synthesis, denaturation and hybridization.

[0035] When a single complex is analysed in accordance with the present method, the result may be used to verify the presence or absence of a specific chemical entity during the

formation of the display molecule. The formation of an extension product is indicative of the presence of an oligonucleotide part complementary to the primer in the identifier oligonucleotide. Conversely, the absence of an extension product is indicative of the absence of an oligonucleotide part complementary to the primer in the identifier oligonucleotide. Selecting the sequence of the primer such that it is complementary to one or more codons will therefore provide information of the structure of the encoded molecule coded for by this codon(s).

[0036] In a preferred aspect of the invention, in the mixture of the identifier oligonucleotide and the primer oligonucleotide, a second primer complementary to a sequence of the extension product is included. The second primer is also termed reverse primer and ensures an exponential increase of the number of produced extension products. The method using a forward and reverse primer is well known to skilled person in the art and is generally referred to as polymerase chain reaction (abbreviated PCR) in the present application with claims. In one embodiment of the invention the reverse primer is annealed to a part of the extension product downstream, i.e. near the 3' end of the extension product, or a part complementing the coding part of the identifier oligonucleotide. In another embodiment, the first primer (forward primer) anneals to an upstream position of the identifier oligonucleotide, preferably before the coding part, and the reverse primer anneals to a sequence of the extension product complementing one or more codons or parts thereof.

[0037] The amplicons resulting from the PCR process may be stained during or following the reaction to ease the detection. A staining after the PCR process may be prepared with e.g. ethidium bromide or a similar staining agent. As an example, amplicons from the PCR process is run on an agarose gel and subsequently stained with ethidium bromide. Under UV illumination bands of amplicons becomes visible. It is possible to incorporate the staining agent in the agarose gel or to allow a solution of the staining agent to migrate through the gel. The amplicons may also be stained during the PCR process by an intercalating agent, like CYBR. In presence of the intercalating agent while the amplification proceeds it will incorporate in the double helix. The intercalation agent may then be made visible by irradiation by a suitable source.

[0038] The intensity of the staining is informative of the relative abundance of a specific amplicon. Thus, it is possible to quantify the occurrence of a codon in an identifier oligonucleotide. When a library of bifunctional complexes has been subjected to a selection the codons in the pool of identifier oligonucleotides which has been selected can be quantified using this method. As an example a sample of the selected identifier oligonucleotides is subjected to various PCR amplifications with different primers in separate compartments and the PCR product of each compartment is analysed by electrophoresis in the presence of ethidium bromide. The bands that appear can be quantified by a densitometric analysis after irradiation by ultraviolet light and the relative abundance of the codons can be measured.

[0039] Alternatively, the primers may be labelled with a suitable small molecule, like biotin or digoxigenin. A PCR-ELISA analysis may subsequently be performed based on the amplicons comprising the small molecule. A preferred method includes the application of a solid support covered

with streptavidin or avidin when biotin is used as label and anti-digoxigenin when digoxigenin is used as the label. Once captured, the amplicons can be detected using an enzyme-labelled avidin or anti-digoxigenin reporter molecule similar to a standard ELISA format.

[0040] To avoid laborious post-PCR handling steps required to evaluate the amplicons, it is in a certain embodiment preferred to measure the extension process “real time”. Several real time PCR processes has been developed and all the suitable real time PCR process available to the skilled person in the art can be used in the evaluating step of the present invention and are include in the present scope of protection. The PCR reactions discussed below are of particular interest.

[0041] The monitoring of accumulating amplicons in real time has been made possible by labelling of primers, probes, or amplicons with fluorogenic molecules. The real time PCR amplification is usually performed with a speed faster than the conventional PCR, mainly due to reduced cycles time and the use of sensitive methods for detection of emissions from the fluorogenic labels. The most commonly used fluorogenic oligoprobes rely upon fluorescent resonance energy transfer (FRET) between fluorogenic labels or between one fluorophore and a dark or “black-hole” nonfluorescent quencher (NFQ), which disperse energy as heat rather than fluorescence. FRET is a spectroscopic process by which energy is passed between molecules separated by 10-100 Å that have overlapping emission and absorption spectra. An advantage of many real time PCR methods is that they can be carried out in a closed system, i.e. a system which does not need to be opened to examine the result of the PCR. A closed system implies a reduced result turnaround, minimisation of the potential for carry-over contamination and the ability to closely scrutinise the assay's performance.

[0042] The real time PCR methods currently available to the skilled person can be classified into either amplicon sequence specific or non-specific methods. The basis for the non-specific detection methods is a DNA-binding fluorogenic molecule. Included in this class are the earliest and simplest approaches to real time PCR. Ethidium bromide, YO-PRO1, and SYBR® green 1 all fluorescence when associated with double stranded DNA which is exposed to a suitable wavelength of light. This approach requires the fluorescent agent to be present during the PCR process and provides for a real time detection of the fluorescent agent as it is incorporated into the double stranded helix.

[0043] The amplicons sequence specific methods includes, but are not limited to, the TaqMan®, hairpin, LightCycler®, Sunrise®, and Scorpions methods. The LightCycler® method also designated “HybProbes” make use of a pair of adjacent, fluorogenic hybridisation oligonucleotide probes. A first, usually the upstream oligoprobe is labelled with a 3' donor fluorophore and the second, usually the downstream probe is commonly labelled with either a Light cycler Red 640 or Red 705 acceptor fluorophore at the 5' terminus so that when both oligoprobes are hybridised the two fluorophores are located in close proximity, such as within 10 nm, of each other. The close proximity provides for the emission of a fluorescence when irradiated with a suitable light source, such a blue diode in case of the LightCycler®. The region for annealing of the probes may

be any suitable position that does not interfere with the primer annealing. In a suitable setup, the site for binding the probes are positioned downstream of the codon region on the identifier oligonucleotide. Alternatively, when a reverse primer is used, the region for annealing the probes may be at the 3' end of the strand complementing the identifier oligonucleotide. Another embodiment of the LightCycler method includes that the pair of oligonucleotide probes are annealed to one or more codons and primer sites exterior to the coding part of the identifier oligonucleotide are used for PCR amplification.

[0044] The TaqMan® method, also referred to as the 5' nuclease or hydrolysis method, requires an oligoprobe, which is attached to a reporter fluorophore, such as 6-carboxy-fluorescein, and a quencher fluorophore, such as 6-carboxy-tetramethyl-rhodamine, at each end. When in close proximity, i.e. annealed to an identifier oligonucleotide, or a sequence complementing the identifier oligonucleotide, the quencher will “hijack” the emissions that have resulted from the excitation of the reporter. As the polymerase progresses along the relevant strand, it displaces and the hydrolyses the oligoprobe via its 5'→3' endonuclease activity. Once the reporter is removed from the extinguishing influence of the quencher, it is able to release excitation energy at a wavelength that can be monitored by a suitable instrument, such as ABI Prism® 7700. The fractional cycle number at which the real-time fluorescence signal mirrors progression of the reaction above the background noise is normally used as an indicator of successful identifier oligonucleotide amplification. This threshold cycle (C_T) is defined as the PCR cycle in which the gain in fluorescence generated by the accumulating amplicons exceeds 10 standard deviations of the mean baseline fluorescence. The C_T is proportional to the number of identifier oligonucleotide copies present in the sample. The TaqMan probe is usually designed to hybridise at a position downstream of a primer binding site, be it a forward or a reverse primer. When the primer is designed to anneal to one or more codons of the identifier oligonucleotide, the presence of these one or more codons is indicated by the emittance of light. Furthermore, the quantity of the identifier oligonucleotides comprising the one or more codons may be measured by the C_T value.

[0045] The Hairpin method involves an oligoprobe, in which a fluorophore and a quencher are positioned at the termini. The labels are held in close proximity by distal stem regions of homologous base pairing deliberately designed to create a hairpin structure which result in quenching either by FRET or a direct energy transfer by a collisional mechanism due to the intimate proximity of the labels. When direct energy transfer by a collision mechanism is used the quencher is usually different from the FRET mechanism, and is suitably 4-(4'-dimethylamino-phenylazo)-benzene (DAB-CYL). In the presence of a complementary sequence, usually downstream of a primer, or within the bounds of the primer binding sites in case of more than one a single primer, the oligoprobe will hybridise, shifting into an open configuration. The fluorophore is now spatially removed from the quencher's influence and fluorescence emissions are monitored during each cycle. In a certain aspect, the hairpin probe may be designed to anneal to a codon in order to detect this codon if present on the identifier oligonucleotide. This embodiment may be suitable if codons only differs from each other with a single or a few nucleotides, because it is well-known that the occurrence of a mismatch

between a hairpin oligoprobe and its target sequence has a greater destabilising effect on the duplex than the introduction of an equivalent mismatch between the target oligonucleotide and a linear oligoprobe. This is probably because the hairpin structure provides a highly stable alternate conformation.

[0046] The Sunrise and Scorpion methods are similar in concept to the hairpin oligoprobe, except that the label becomes irreversible incorporated in to the PCR product. The Sunrise method involves a primer (commercially available as Amplifluor™ hairpin primers) comprising a 5' fluorophore and a quencher, e.g. DABCYL. The labels are separated by complementary stretches of sequence that create a stem when the sunrise primer is closed. At the 3' terminus is a target specific primer sequence. In a preferred embodiment the target sequence is a codon, optionally more codons. The sunrise primer's sequence is intended to be duplicated by the nascent complementary strand and, in this way, the stem is destabilised, the two fluorophores are held apart, usually between 15 and 25 nucleotides, and the fluorophore is free to emit its excitation energy for monitoring. The Scorpion primer resembles the sunrise primer, but deviate in having a moiety that blocks duplication on the signalling portion of the scorpion primer. The blocking moiety is typically hexethylene glycol. In addition to the difference in structure, the function of the scorpion primers differs slightly in that the 5' region of the oligonucleotide is designed to hybridise to a complementary region within the amplicons. In a certain embodiment the complementary region is a codon on the identifier oligonucleotide. The hybridisation forces the labels apart disrupting the hairpin and permitting emission in the same way as the hairpin probes.

[0047] After the selection has been performed the codon profile is indicative of the chemical entities that have been used in the synthesis of encoded molecules having a certain property, such as an affinity towards a target. In the event the selection has been sufficient effective it may be possible directly to deduce a part or the entire structure of encoded molecules with the desired property. Alternatively, it may be possible to deduce a structural unit appearing more frequently among the encoded molecules after the selection, which gives important information to the structure-activity-relationship (SAR). If the selection process has not narrowed the size of the library to a manageable number, the formation of a second-generation library is useful. In the formation of the second-generation library chemical entities, which have not been involved in the synthesis of encoded molecules that have been successful in the selection may be omitted, thus limiting the size of the new library and at the same time increasing the concentration of complexes with the requested property, e.g. the ability to bind to a target. The second-generation library may then be subjected to more stringent selection conditions to allow only the encoded molecules with a higher affinity to bind to the target. The second-generation library may also be generated using the chemical entities coded for in addition to certain chemical entities suspected of increasing the performance of the final encoded molecule. The indication of certain successful chemical entities may be obtained from the SAR. The use in a second-generation library of chemical entities, which have proved to be interesting for further investigation in a pre-

ceding library, may thus entail a shuffling with new chemical entities that may focus the second-generation library in a certain desired direction.

[0048] An Example of implicit identification of codons includes that the nucleic acid fragment is associated with a chemical entity precursor capable of being transferred to a recipient reactive group. The recipient reactive group may be a part of a chemical scaffold and the chemical entity precursor may add a structural unit to said scaffold. It is preferred that the nucleic acid fragment codes for the chemical entity. In some aspects of the present invention each member of the nucleic acid fragment pool comprises an anticodon, which identifies the chemical entity. When a plurality of chemical entities are present the anticodon is preferably unique, i.e. a unique correspondence between the chemical entities and the associated anticodons exists.

[0049] The identifier nucleic acid sequence comprises codons, which may be able to pair with one or more anticodons of the pool of nucleic acid fragments. The pairing between one or more codons of an identifier nucleic acid sequence and one or more anticodons is preferably specific, i.e. the one or more codons of the identifier nucleic acid sequence are only recognized by particular anticodons. The nucleic acid fragment containing more than one anticodon can encode for scaffold molecules where each anticodon encodes for specific chemical entities of that scaffold molecule. The specific pairing makes it possible implicitly to decode the codon of an identifier nucleic acid sequence. In the method according to the invention, non-specific pairing between codons and anticodons can be cleaved with an enzyme or chemically treated to break the double stranded nucleotides. The non-pairing region can be cleaved using enzymes that cleaves specifically nucleotide sequences with mismatches. Notably, the enzyme is selected from T4 endonuclease VII, T4 endonuclease I, CEL I, nuclease S1, or variants thereof. The cleavage is preferable used when more than one codon and anticodon is involved in pairing between the identifier nucleic acid sequence and the nucleic acid fragment.

[0050] The pool of nucleic acid fragments associated with a chemical entity may comprise anticodons complemented by codons of one or more identifier nucleic acid sequence as well as anticodons which are not complemented by codons on any identifier nucleic acid sequence. In other words, the amount of genetic information contained in the anticodons of the pool is larger than the amount of genetic information complemented by the codons.

[0051] The contacting of the one or more identifier nucleic acid sequences with the pool of nucleic acid fragments are usually conducted at conditions, which allow for hybridisation, i.e. conditions at which cognate nucleic acid sequences can anneal to each other. To facilitate the recovery of nucleic acid fragments, which have annealed to the identifier nucleic acid sequences, the identifier nucleic acid sequences are usually immobilized on a solid support. Examples of suitable solid supports include beads and column material, e.g. beads and column material associated with a second part of the affinity pair to bind identifier nucleic acid sequences attached to the first part of the molecular affinity pair. In certain aspects of the invention the solid support is associated with streptavidin and the identifier nucleic acid sequences are attached to biotin.

[0052] When the identifier nucleic acid sequences are immobilized on a solid support the pool of nucleic acid fragments is typically present in a mobile phase, i.e. dissolved in a liquid. The identifier nucleic acids will hybridise to these nucleic acid fragments in the pool which are sufficient complementary to a particular part of an identifier nucleic acid sequence for a binding to occur. Fragments not finding any complementing sequence will remain in the solution. In the event, the identifier nucleic acid sequences are segregated into codons and the fragments comprises anticodons, the anticodons which are able to anneal to a codons will be caught while fragments not having a cognate codon will be maintained in the mobile phase. When codons and anticodons are present in the method of the present invention, specific hybridisation implies that the tendency of an anticodon to cross-hybridise to another codon will be impede or avoided. To avoid cross-hybridisation, codons may be designed such that each codon is distinguished from all other codons by one, two or more mismatching nucleotides.

[0053] The mobile phase is subsequently separated from the solid phase e.g. by washing, and the enriched pool of fragments is recovered. The recovery of the nucleic acid fragments are usually done by subjecting the hybrid to denaturing conditions, i.e. conditions which separate the two strands. If the parent nucleic acid sequences are immobilized on beads, the separation of the fragments can be effected using denaturing conditions and centrifugation/spinning.

[0054] The enriched pool of nucleic acid fragments associated with a chemical entity may be used directly to prepare a next generation library of complexes, in which each member of the library comprises an encoded molecule and the nucleic acid sequence which codes for this molecule. In one embodiment of the invention, building blocks comprising a particular transferable chemical entity associated with an anticodon corresponding to the anticodons of the detected fragments are used in the generation of the next generation library. In another embodiment, additional building blocks are added having modified transferable chemical entities in order to improve on a certain property of the encoded molecule.

[0055] The complexes may be prepared by various known methods starting from the nucleic acid fragment comprising the anticodon and the chemical entity, as disclosed above. According to a particular method, the next generation library is formed by a) mixing under hybridisation conditions, nascent bifunctional complexes comprising a chemical entity or a reaction product of chemical entities, and an identifier nucleic acid sequence comprising codon(s) identifying said chemical entities, with the recovered nucleic acid fragments, said fragments comprising an oligonucleotide sufficient complementary to at least a part of the identifier nucleic acid sequence to allow for hybridisation, a transferable chemical entity and an anticodon identifying the chemical entity, to form hybridisation products; and b) transferring the chemical entities of the nucleic acid fragments to the nascent bifunctional complexes through a reaction involving a reactive group of the nascent bifunctional complex, in conjunction with a transfer of the genetic information of the anticodon.

[0056] Preferably, the above method for preparing the next generation library comprises the further step of c) separating

the components of the hybridisation product and recovering the complexes. If further chemical entities are intended to participate in the formation of the encoded molecule of the nascent complex, steps a) through c) are repeated as appropriate using the recovered complexes in step c) as the nascent bifunctional complexes in step a) of the next round.

[0057] The genetic information of the anticodon may be transferred to the nascent complex by a variety of methods. According to a first embodiment the genetic information of the anticodon is transferred by enzymatically extending the oligonucleotide identifier region to obtain a codon attached to the bifunctional complex having received the chemical entity. A second embodiment implies that genetic information of the anticodon is transferred to the nascent complexes by hybridisation to a cognate codon of the nascent complex.

[0058] According to the first embodiment, the enriched pool of fragments comprises an affinity oligonucleotide sufficient complementary to an identifier region of the nascent complex, said oligonucleotide being distinct from the anticodon. Accordingly, the oligonucleotide identifier region of the nascent complex anneals to the affinity oligonucleotide of the building block to form the hybridisation product, while the anticodon remains single stranded. Subsequently, the chemical entity is transferred to the recipient reactive group of the complex to form the encoded molecule prior to, simultaneously with, or subsequent to the enzymatic extension of the hybridisation product using the anticodon as identifier. Specific examples of suitable enzymes are polymerases and ligases, which requires dNTPs and oligonucleotides, respectively as substrates. The method for forming the complexes according to this first embodiment is the subject PCT/DK03/00739, the content thereof being incorporated herein by reference.

[0059] According to the second embodiment, the anticodon form part of the affinity oligonucleotide, i.e. the anticodon is a part of or the entire affinity oligonucleotide. Initially, a plurality of identifiers comprising different codons and/or different order of codons is provided. The identifiers are associated with a recipient reactive group, i.e. the reactive group may be covalently attached to the identifier or attached by hybridisation. Notably, a codon of the identifier may be used for the attachment of a building block harbouring the reactive group. The identifiers are subsequently contacted with the enriched pool of building blocks, i.e. nucleic acid fragments associated with a transferable chemical entity. The mixture of identifiers and building blocks are maintained at hybridisation conditions to anneal the anticodon of the building blocks to the cognate codon of the identifier. After or simultaneously with the annealing step, the chemical entity is transferred to the recipient reactive group of the identifier. The method for forming the complexes according to the second embodiment is the subject of various patent applications, including WO 02/103008, WO 02/074929, Danish patent application No. PA 2002 01347, and U.S. provisional patent application No. 60/409,968. The content of these patent applications are incorporated herein by reference in their entirety.

[0060] The new generation of library complexes may be used in a partition step, in which the library of complexes is subjected to a condition partitioning complexes displaying a predetermined property from the remainder of the next generation library, as explained above. Thus, using the

present method, it is possible to repeat the partitioning procedure a desired number of times using still more stringent conditions, until a single or a few encoded molecules are identified which display the desired property to a high extent. When the partitioning is based on an affinity assay, the library of encoded molecules are increasingly narrowed in size from one generation to the next and at the same time the high affinity binders are increased in concentration.

[0061] The outcome of a codon analysis will be dependent of the enrichment factor in the selection process. An efficient and specific selection will generate a large difference between the specific binders compared to the background. Still, there will be a large amount of molecules in the background that will reduce the possibility to obtain measurable differences between the binders and the background in the codon analysis procedure. If the enrichment factor (or too large library) is not good enough to distinguish a specific binder among the background binders, the signal in the codon analysis will probably not be detectable. However, there will be a continuing of binders that use a certain chemical entity in a certain position. These "non optimal" binders (a certain important chemical entity in one position and less important in the other position) will be many due to the diversity obtained when only one (or a few) positions are important in the selection process. Therefore, the sum of all molecules with a preferable chemical entity in a certain position will be larger than the sum of all molecules with a non-binding chemical entity, which will make the codon analysis easier.

[0062] This invention may involve an extensive analysis of all the chemical entities in a library and how they are involved in the binding to targets. This information can be used both to design new libraries and in the final process where the lead structures are produced and pre-clinical candidates are picked. The extensive data obtained in the codon analysis can for instance be used for selecting candidates with the appropriate specificity. This can be done if selection has been performed on a family of proteins where one of the members is the target.

[0063] The invention enables pharmacophore identification and transformation into small molecule drugs. In cases where peptide-like libraries is used, the peptide/peptidomimetic lead to small molecule conversion process is supported by medicinal chemistry and cheminformatics and guided by matching the pharmacophore derived from massive structure activity relationship (SAR) data information from the codon analysis. A "pharmacophore" is a description of the structural criteria a molecule must fulfil in order that it is active against a specified biological receptor. These criteria are usually the 3D spatial relationships of a set of chemical features, and sometimes include the steric boundaries, within which the molecule must fit. There is a set of software methods, which automatically infers such pharmacophores, given a SAR, in the absence of direct macromolecular structural data.

[0064] The extensive SAR information obtained using the codon analyses described in this invention can be combined with molecular modeling technologies to refine for example pharmacophore models and the plausible interactions between the potential binders and a target.

[0065] The codon analysis is also a valuable experimental tool for SAR on weak binders. The codon analysis measures

the abundance of chemical entities after a selection in all binding molecules. Thus, even weak binders, which there might be many of, is detected even though the detected codon is selected in many different combinations. The selection procedure can also be tuned to enrich predominantly for weak binders, which will simplify the codon analysis data.

[0066] This invention is also suitable for replacing the laborious task of extracting SAR information by hand with an automated process using suitable algorithm and software programs. The codon analysis (e.g. array or QPCR measurements) can be directly feed into a data handling software program that use both the codon abundances and structural data to generate SAR information and potential pharmacophore models.

[0067] The SAR information and potential pharmacophore models obtained from the codon analysis can be used to design focused libraries in an array format allowing massive and parallel testing. Thus, the selection procedure and codon analysis can be seen as a diversity reduction step to allow a complete test of potential binders in an array format.

[0068] Various methods for identifying the codons of the identifiers of step iii) are disclosed herein. When a pool of partitioned identifier nucleic acid sequences is subjected to the identification step it is normally not practically to decode a sufficient number of sequences comprising the entire "genome" of an encoded molecule to ensure that all interesting encoded molecules have been revealed. Therefore, a modified sequencing technique preferably identifies the codons in each position occurring with the highest frequency. The next generation library is then build using in each position the chemical entities occurring with the highest frequency.

[0069] In a certain embodiment of the invention, the codon identification step uses the entire population of identifier nucleic acid sequences in the analysis and informs the experimenter of the relative abundance of each codon in a certain position. The codon information may be obtained using microarray, QPCR, or any equivalent method for revealing the identity of codons. In contrary, sequencing a subset of identifier nucleic acid sequences only provides the experimenter with a limited insight as to the population of codons and the corresponding encoded molecules.

DETAILED DESCRIPTION OF THE INVENTION

[0070] Complex

[0071] The complex comprises an encoded molecule and an identifier oligonucleotide. The identifier comprises codons that identify the encoded molecule. Preferably, the identifier oligonucleotide identifies the encoded molecule uniquely, i.e. in a library of complexes a particular identifier is capable of distinguishing the molecule it is attached to from the rest of the molecules.

[0072] The encoded molecule and the identifier may be attached directly to each other or through a bridging moiety. In one aspect of the invention, the bridging moiety is a selectively cleavable linkage.

[0073] The identifier oligonucleotide may comprise two or more codons. In a preferred aspect the identifier oligonucleotide

otide comprises three or more codons. The sequence of each codon can be decoded utilizing the present method to identify reactants used in the formation of the encoded molecule. When the identifier comprises more than one codon, each member of a pool of chemical entities can be identified and the order of codons is informative of the synthesis step each member has been incorporated in.

[0074] In a certain embodiment, the same codon is used to code for several different chemical entities. In a subsequent identification step, the structure of the encoded molecule can be deduced taking advantage of the knowledge of different attachment chemistries, steric hindrance, deprotection of orthogonal protection groups, etc. In another embodiment, the same codon is used for a group of chemical entities having a common property, such as a lipophilic nature, a certain attachment chemistry etc. In a preferred embodiment, however, the codon is unique i.e. a similar combination of nucleotides does not appear on the identifier oligonucleotide coding for another chemical entity. In a practical approach, for a specific chemical entity, only a single combination of nucleotides is used. In some aspects of the invention, it may be advantageous to use several codons for the same chemical entity, much in the same way as Nature uses up to six different codons for a single amino acid. The two or more codons identifying the same chemical entity may carry further information related to different reaction conditions.

[0075] The sequence of the nucleotides in each codon may have any suitable length. The codon may be a single nucleotide or a plurality of nucleotides. In some aspects of the invention, it is preferred that each codon independently comprises four or more nucleotides, more preferred 4 to 30 nucleotides. In some aspects of the invention the lengths of the codons vary.

[0076] A certain codon may be distinguished from any other codon in the library by only a single nucleotide. However, to facilitate a subsequent decoding process and to increase the ability of the primer to discriminate between codons it is in general desired to have two or more mismatches between a particular codon and any other codon appearing on identifier oligonucleotide. As an example, if a codon length of 5 nucleotides is selected, more than 100 nucleotide combinations exist in which two or more mismatches appear. For a certain number of nucleotides in the codon, it is generally desired to optimize the number of mismatches between a particular codon relative to any other codon appearing in the library.

[0077] The identifier oligonucleotide will in general have at least two codons arranged in sequence, i.e. next to each other. Two neighbouring codons may be separated by a framing sequence. Depending on the encoded molecule formed, the identifier may comprise further codons, such as 3, 4, 5, or more codons. Each of the further codons may be separated by a suitable framing sequence. Preferably, all or at least a majority of the codons of the identifier are separated from a neighbouring codon by a framing sequence. The framing sequence may have any suitable number of nucleotides, e.g. 1 to 20. Alternatively, codons on the identifier may be designed with overlapping sequences.

[0078] The framing sequence, if present, may serve various purposes. In one setup of the invention, the framing sequence identifies the position of the codon. Usually, the

framing sequence either upstream or downstream of a codon comprises information which positions the chemical entity and the reaction conditions in the synthesis history of the encoded molecule. The framing sequence may also or in addition provide for a region of high affinity. The high affinity region may ensure that a hybridisation event with an anti-codon will occur in frame. Moreover, the framing sequence may adjust the annealing temperature to a desired level.

[0079] A framing sequence with high affinity can be provided by incorporation of one or more nucleobases forming three hydrogen bonds to a cognate nucleobase. Examples of nucleobases having this property are guanine and cytosine. Alternatively, or in addition, the framing sequence may be subjected to backbone modification. Several backbone modifications provides for higher affinity, such as 2'-O-methyl substitution of the ribose moiety, peptide nucleic acids (PNA), and 2'-O-methylene cyclisation of the ribose moiety, also referred to as LNA (Locked Nucleic Acid).

[0080] The sequence comprising a codon and an adjacent framing sequence has in a certain aspect of the invention a total length of 11 nucleotides or more, preferably 15 nucleotides or more. A primer may be designed to complementary to the codon sequence as well as the framing sequence. The presence of an extension reaction under conditions allowing for such reaction to occur is indicative of the presence of the chemical entity encoded in the codon as well as the position said chemical entity has in the entire synthesis history of the encoded molecule.

[0081] The identifier may comprise flanking regions around the coding section. The flanking regions can also serve as priming sites for amplification reactions, such as PCR or as binding region for oligonucleotide probe. The identifier may in certain embodiments comprise an affinity region having the property of being able to hybridise to a building block.

[0082] It is to be understood that when the term identifier oligonucleotide is used in the present description and claims, the identifier oligonucleotide may be in the sense or the anti-sense format, i.e. the identifier can be a sequence of codons which actually codes for the encoded molecule or can be a sequence complementary thereto. Moreover, the identifier may be single-stranded or double-stranded, as appropriate.

[0083] The encoded molecule part of the complex is generally of a structure expected of having an effect according to the property sought for, e.g. the encoded molecule has a binding affinity towards a target. When the target is of pharmaceutical importance, the encoded molecule is generally a possible drug candidate. The complex may be formed by tagging a library of different possible drug candidates with a tag, e.g. a nucleic acid tag identifying each possible drug candidate. In another embodiment of the invention, the molecule formed by a variety of reactants which have reacted with each other and/or a scaffold molecule. Optionally, this reaction product may be post-modified to obtain the final molecule displayed on the complex. The post-modification may involve the cleavage of one or more chemical bonds attaching the encoded molecule to the identifier in order more efficiently to display the encoded molecule.

[0084] The formation of an encoded molecule generally starts by a scaffold, i.e. a chemical unit having one or more

reactive groups capable of forming a connection to another reactive group positioned on a chemical entity, thereby generating an addition to the original scaffold. A second chemical entity may react with a reactive group also appearing on the original scaffold or a reactive group incorporated by the first chemical entity. Further chemical entities may be involved in the formation of the final reaction product. The formation of a connection between the chemical entity and the nascent encoded molecule may be mediated by a bridging molecule. As an example, if the nascent encoded molecule and the chemical entity both comprise an amine group a connection between these can be mediated by a dicarboxylic acid. A synthetic molecule is in general produced in vitro and may be a naturally occurring or an artificial substance. Usually, a synthetic molecule is not produced using the naturally translation system in an in vitro process.

[0085] The chemical entities that are precursors for structural additions or eliminations of the encoded molecule may be attached to a building block prior to the participation in the formation of the reaction product leading the final encoded molecule. Besides the chemical entity, the building block generally comprises an anti-codon. In some embodiments the building blocks also comprise an affinity region providing for affinity towards the nascent complex.

[0086] Thus, the chemical entities are suitably mediated to the nascent encoded molecule by a building block, which further comprises an anticodon. The anti-codon serves the function of transferring the genetic information of the building block in conjunction with the transfer of a chemical entity. The transfer of genetic information and chemical entity may occur in any order. The chemical entities are preferably reacted without enzymatic interaction in some aspects of the invention. Notably, the reaction of the chemical entities is preferably not mediated by ribosomes or enzymes having similar activity. In other aspects of the invention, enzymes are used to mediate the reaction between a chemical entity and a nascent encoded molecule.

[0087] According to certain aspects of the invention the genetic information of the anti-codon is transferred by specific hybridisation to a codon on a nucleic acid identifier. Another method for transferring the genetic information of the anti-codon to the nascent complex is to anneal an oligonucleotide complementary to the anti-codon and attach this oligonucleotide to the complex, e.g. by ligation. A still further method involves transferring the genetic information of the anti-codon to the nascent complex by an extension reaction using a polymerase and a mixture of dNTPs.

[0088] The chemical entity of the building block may in most cases be regarded as a precursor for the structural entity eventually incorporated into the encoded molecule. In other cases the chemical entity provides for the eliminations of chemical units of the nascent encoded molecule. Therefore, when it in the present application with claims is stated that a chemical entity is transferred to a nascent encoded molecule it is to be understood that not necessarily all the atoms of the original chemical entity is to be found in the eventually formed encoded molecule. Also, as a consequence of the reactions involved in the connection, the structure of the chemical entity can be changed when it appears on the nascent encoded molecule. Especially, the cleavage resulting in the release of the entity may generate a reactive group

which in a subsequent step can participate in the formation of a connection between a nascent complex and a chemical entity.

[0089] The chemical entity of the building block comprises at least one reactive group capable of participating in a reaction which results in a connection between the chemical entity of the building block and another chemical entity or a scaffold associated with the nascent complex. The number of reactive groups which appear on the chemical entity is suitably one to ten. A building block featuring only one reactive group is used i.a. in the end positions of polymers or scaffolds, whereas building blocks having two reactive groups are suitable for the formation of the body part of a polymer or scaffolds capable of being reacted further. One, two or more reactive groups intended for the formation of connections, are typically present on scaffolds. Non-limiting examples of scaffolds are opiates, steroids, benzodiazepines, hydantoines, and peptidylphosphonates.

[0090] The reactive group of the chemical entity may be capable of forming a direct connection to a reactive group of the nascent complex or the reactive group of the building block may be capable of forming a connection to a reactive group of the nascent complex through a bridging fill-in group. It is to be understood that not all the atoms of a reactive group are necessarily maintained in the connection formed. Rather, the reactive groups are to be regarded as precursors for the structure of the connection.

[0091] The subsequent cleavage step to release the chemical entity from the building block can be performed in any appropriate way. In an aspect of the invention the cleavage involves usage of a chemical reagent or an enzyme. The cleavage results in a transfer of the chemical entity to the nascent encoded molecule or in a transfer of the nascent encoded molecule to the chemical entity of the building block. In some cases it may be advantageous to introduce new chemical groups as a consequence of linker cleavage. The new chemical groups may be used for further reaction in a subsequent cycle, either directly or after having been activated. In other cases it is desirable that no trace of the linker remains after the cleavage.

[0092] In another aspect, the connection and the cleavage is conducted as a simultaneous reaction, i.e. either the chemical entity of the building block or the nascent encoded molecule is a leaving group of the reaction. In some aspects of the invention, it is appropriate to design the system such that the connection and the cleavage occur simultaneously because this will reduce the number of steps and the complexity. The simultaneous connection and cleavage can also be designed such that either no trace of the linker remains or such that a new chemical group for further reaction is introduced, as described above.

[0093] The attachment of the chemical entity to the building block, optionally via a suitable spacer can be at any entity available for attachment, e.g. the chemical entity can be attached to a nucleobase or the backbone. In general, it is preferred to attach the chemical entity at the phosphor of the internucleoside linkage or at the nucleobase. When the nucleobase is used for attachment of the chemical entity, the attachment point is usually at the 7 position of the purines or 7-deaza-purins or at the 5 position of pyrimidines. The nucleotide may be distanced from the reactive group of the chemical entity by a spacer moiety. The spacer may be

designed such that the conformational space sampled by the reactive group is optimized for a reaction with the reactive group of the nascent encoded molecule.

[0094] The encoded molecules may have any chemical structure. In a preferred aspect, the encoded molecule can be any compound that may be synthesized in a component-by-component fashion. In some aspects the synthetic molecule is a linear or branched polymer. In another aspect the synthetic molecule is a scaffolded molecule. The term "encoded molecule" also comprises naturally occurring molecules like α -polypeptides etc, however produced in vitro usually in the absence of enzymes, like ribosomes. In certain aspects, the synthetic molecule of the library is a non- α -polypeptide.

[0095] The encoded molecule may have any molecular weight. However, in order to be orally available, it is in this case preferred that the synthetic molecule has a molecular weight less than 2000 Daltons, preferably less than 1000 Dalton, and more preferred less than 500 Daltons.

[0096] The size of the library may vary considerably pending on the expected result of the inventive method. In some aspects, it may be sufficient that the library comprises two, three, or four different complexes. However, in most events, more than two different complexes are desired to obtain a higher diversity. In some aspects, the library comprises 1,000 or more different complexes, more preferred 1,000,000 or more different complexes. The upper limit for the size of the library is only restricted by the size of the vessel in which the library is comprised. It may be calculated that a vial may comprise up to 10^{14} different complexes.

[0097] Methods for Forming Libraries of Complexes

[0098] The encoded molecules associated with an identifier oligonucleotide having two or more codons that code for reactants that have reacted in the formation of the molecule part of the complex may be formed by a variety of processes. Generally, the preferred methods can be used for the formation of virtually any kind of encoded molecule. Suitable examples of processes include prior art methods disclosed in WO 93/20242, WO 93106121, WO 00/23458, WO 02/074929, and WO 02/103008, the content of which being incorporated herein by reference as well as methods of the present applicant not yet public available, including the methods disclosed in PCT/DK03/00739 filed 30 Oct. 2003, and DK PA 2003 00430 filed 20 Mar. 2003. Any of these methods may be used, and the entire content of the patent applications are included herein by reference.

[0099] Below five presently preferred embodiments are described. A first embodiment disclosed in more detail in WO 02/103008 is based on the use of a polymerase to incorporate unnatural nucleotides as building blocks. Initially, a plurality of identifier oligonucleotides is provided. Subsequently primers are annealed to each of the identifiers and a polymerase is extending the primer using nucleotide derivatives, which have appended chemical entities. Subsequent to or simultaneously with the incorporation of the nucleotide derivatives, the chemical entities are reacted to form a reaction product. The encoded molecule may be post-modified by cleaving some of the linking moieties to better present the encoded molecule.

[0100] Several possible reaction approaches for the chemical entities are apparent. First, the nucleotide deriva-

tives can be incorporated and the chemical entities subsequently polymerised. In the event the chemical entities each carry two reactive groups, the chemical entities can be attached to adjacent chemical entities by a reaction of these reactive groups. Exemplary of the reactive groups are amine and carboxylic acid, which upon reaction form an amide bond. Adjacent chemical entities can also be linked together using a linking or bridging moiety. Exemplary of this approach is the linking of two chemical entities each bearing an amine group by a bi-carboxylic acid. Yet another approach is the use of a reactive group between a chemical entity and the nucleotide building block, such as an ester or a hoister group. An adjacent building block having a reactive group such as an amine may cleave the interspaced reactive group to obtain a linkage to the chemical entity, e.g. by an amide linking group.

[0101] A second embodiment for obtainment of complexes disclosed in WO 02/103008 pertains to the use of hybridisation of building blocks to an identifier and reaction of chemical entities attached to the building blocks in order to obtain a reaction product. This approach comprises that identifiers are contacted with a plurality of building blocks, wherein each building block comprises an anti-codon and a chemical entity. The anti-codons are designed such that they recognise a sequence, i.e. a codon, on the identifier. Subsequent to the annealing of the anti-codon and the codon to each other a reaction of the chemical entity is effected.

[0102] The identifier may be associated with a scaffold. Building blocks bringing chemical entities in may be added sequentially or simultaneously and a reaction of the reactive group of the chemical entity may be effected at any time after the annealing of the building blocks to the identifier.

[0103] A third embodiment for the generation of a complex includes chemical or enzymatic ligation of building blocks when these are lined up on a identifier. Initially, identifiers are provided, each having one or more codons. The identifiers are contacted with building blocks comprising anti-codons linked to chemical entities. The two or more anti-codons annealed on an identifier are subsequently ligated to each other and a reaction of the chemical entities is effected to obtain a reaction product. The method is disclosed in more detail in DK PA 2003 00430 filed 20 Mar. 2003.

[0104] A fourth embodiment makes use of the extension by a polymerase of an affinity sequence of the nascent complex to transfer the anti-codon of a building block to the nascent complex. The method implies that a nascent complex comprising a scaffold and an affinity region is annealed to a building block comprising a region complementary to the affinity section. Subsequently, the anti-codon region of the building block is transferred to the nascent complex by a polymerase. The transfer of the chemical entity may be transferred prior to, simultaneously with or subsequent to the transfer of the anti-codon. This method is disclosed in detail in PCT/DK03100739.

[0105] A fifth embodiment also disclosed in PCT/DK03/00739 comprises reaction of a reactant with a reaction site on nascent bifunctional molecule and addition of a nucleic acid tag to the nascent bifunctional molecule using an enzyme, such as a ligase. When a library is formed, usually an array of compartments is used for reaction of reactants and enzymatic addition of tags with the nascent bifunctional molecule.

[0106] Thus, the codons are either pre-made into one or more identifiers before the encoded molecules are generated or the codons are transferred simultaneously with the formation of the encoded molecules.

[0107] After or simultaneously with the formation of the reaction product some of the linkers to the identifier may be cleaved, however, usually at least one linker is maintained to provide for the complex.


[0108] Nucleotides

[0109] The nucleotides used in the present invention may be linked together in a sequence of nucleotides, i.e. an oligonucleotide. Each nucleotide monomer is normally composed of two parts, namely a nucleobase moiety, and a backbone. The backbone may in some cases be subdivided into a sugar moiety and an internucleoside linker.

[0110] The nucleobase moiety may be selected among naturally occurring nucleobases as well as non-naturally occurring nucleobases. Thus, "nucleobase" includes not only the known purine and pyrimidine heterocycles, but also heterocyclic analogues and tautomers thereof. Illustrative examples of nucleobases are adenine, guanine, thymine, cytosine, uracil, purine, xanthine, diaminopurine, 8-oxo-N⁶-methyladenine, 7-deazaxanthine, 7-deazaguanine, N⁴,N⁴-ethanocytosin, N⁶,N⁶-ethano-2,6-diamino-purine, 5-methylcytosine, 5-(C³-C⁶)-alkynylcytosine, 5-fluorouracil, 5-bromouracil, pseudoisocytosine, 2-hydroxy-5-methyl-4-triazolopyridine, isocytosine, isoguanine, inosine and the "non-naturally occurring" nucleobases described in Benner et al., U.S. Pat. No. 5,432,272. The term "nucleobase" is intended to cover these examples as well as analogues and tautomers thereof. Especially interesting nucleobases are adenine, guanine, thymine, cytosine, 5-methylcytosine, and uracil, which are considered as the naturally occurring nucleobases in relation to therapeutic and diagnostic application in humans.


[0111] Examples of suitable specific pairs of nucleobases are shown below:

Natural Base Pairs




-continued



Synthetic Base Pairs




-continued



[0112] Suitable examples of backbone units are shown below (B denotes a nucleobase):



-continued



[0113] The sugar moiety of the backbone is suitably a pentose but may be the appropriate part of a PNA or a six-member ring. Suitable examples of possible pentoses include ribose, 2'-deoxyribose, 2'-O-methyl-ribose, 2'-flour-ribose, and 2',4'-O-methylene-ribose (LNA). Suitably the nucleobase is attached to the 1' position of the pentose entity.


[0114] An internucleoside linker connects the 3' end of preceding monomer to a 5' end of a succeeding monomer when the sugar moiety of the backbone is a pentose, like ribose or 2'-deoxyribose. The internucleoside linkage may be the natural occurring phosphodiester linkage or a derivative thereof. Examples of such derivatives include phosphorothioate, methylphosphonate, phosphoramidate, phosphorriester, and phosphodithioate. Furthermore, the internucleoside linker can be any of a number of non-phosphorous-containing linkers known in the art.

[0115] Preferred nucleic acid monomers include naturally occurring nucleosides forming part of the DNA as well as the RNA family connected through phosphodiester linkages. The members of the DNA family include deoxyadenosine,

deoxyguanosine, deoxythymidine, and deoxycytidine. The members of the RNA family include adenosine, guanosine, uridine, cytidine, and inosine. Inosine is a non-specific pairing nucleoside and may be used as universal base because inosine can pair nearly isoenergetically with A, T, and C. Other compounds having the same ability of non-specifically base-pairing with natural nucleobases have been formed. Suitable compounds which may be utilized in the present invention includes among others the compounds depicted below

EXAMPLES OF UNIVERSAL BASES

[0116]



[0117] Building Block

[0118] The chemical entities or reactants that are precursors for structural additions or eliminations of the encoded molecule may be attached to a building block prior to the participation in the formation of the reaction product leading to the final encoded molecule. Besides the chemical entity, the building block generally comprises an anti-codon.

[0119] The chemical entity of the building block comprises at least one reactive group capable of participating in a reaction, which results in a connection between the chemical entity of the building block and another chemical entity

or a scaffold associated with the nascent complex. The connection is facilitated by one or more reactive groups of the chemical entity. The number of reactive groups, which appear on the chemical entity, is suitably one to ten. A building block featuring only one reactive group is used i.a. in the end positions of polymers or scaffolds, whereas building blocks having two reactive groups are suitable for the formation of the body part of a polymer or scaffolds capable of being reacted further. One, two or more reactive groups intended for the formation of connections are typically present on scaffolds.

[0120] The reactive group of the building block may be capable of forming a direct connection to a reactive group of the nascent complex or the reactive group of the building block may be capable of forming a connection to a reactive group of the nascent complex through a bridging fill-in group. It is to be understood that not all the atoms of a reactive group are necessarily maintained in the connection formed. Rather, the reactive groups are to be regarded as precursors for the structure of the connection.

[0121] The subsequent cleavage step to release the chemical entity from the building block can be performed in any appropriate way. In an aspect of the invention the cleavage involves usage of a reagent or an enzyme. The cleavage results in a transfer of the chemical entity to the nascent encoded molecule or in a transfer of the nascent encoded molecule to the chemical entity of the building block. In some cases it may be advantageous to introduce new chemical groups as a consequence of linker cleavage. The new chemical groups may be used for further reaction in a subsequent cycle, either directly or after having been activated. In other cases it is desirable that no trace of the linker remains after the cleavage.

[0122] In another aspect, the connection and the cleavage are conducted as a simultaneous reaction, i.e. either the chemical entity of the building block or the nascent encoded molecule is a leaving group of the reaction. In general, it is preferred to design the system such that the connection and the cleavage occur simultaneously because this will reduce the number of steps and the complexity. The simultaneous connection and cleavage can also be designed such that either no trace of the linker remains or such that a new chemical group for further reaction is introduced, as described above.


[0123] The attachment of the chemical entity to the building block, optionally via a suitable spacer can be at any entity available for attachment, e.g. the chemical entity can be attached to a nucleobase or the backbone. In general, it is preferred to attach the chemical entity at the phosphor of the internucleoside linkage or at the nucleobase. When the nucleobase is used for attachment of the chemical entity, the attachment point is usually at the 7 position of the purines or 7-deaza-purins or at the 5 position of pyrimidines. The nucleotide may be distanced from the reactive group of the chemical entity by a spacer moiety. The spacer may be designed such that the conformational space sampled by the reactive group is optimized for a reaction with the reactive group of the nascent encoded molecule or reactive site.

[0124] The anticodon complements the codon of the identifier oligonucleotide sequence and generally comprises the same number of nucleotides as the codon. The anti-codon may be adjoined with a fixed sequence, such as a sequence complementing a framing sequence.

[0125] Various specific building blocks are envisaged. Building blocks of particular interest are shown below.


[0126] Building Blocks Transferring a Chemical Entity to a Recipient Nucleophilic Group

[0127] The building block indicated below is capable of transferring a chemical entity (CE) to a recipient nucleophilic group, typically an amine group. The bold lower horizontal line illustrates the building block comprising an anti-codon and the vertical line illustrates a spacer. The 5membered substituted N-hydroxysuccinimid (NHS) ring serves as an activator, i.e. a labile bond is formed between the oxygen atom connected to the NHS ring and the chemical entity. The labile bond may be cleaved by a nucleophilic group, e.g. positioned on a scaffold



[0128] The 5-membered substituted N-hydroxysuccinimid (NHS) ring serves as an activator, i.e. a labile bond is formed between the oxygen atom connected to the NHS ring and the chemical entity. The labile bond may be cleaved by a nucleophilic group, e.g. positioned on a scaffold, to transfer the chemical entity to the scaffold, thus converting the remainder of the fragment into a leaving group of the reaction. When the chemical entity is connected to the activator through a carbonyl group and the recipient group is an amine, the bond formed on the scaffold will be an amide bond. The above building block is the subject of WO03078627A2, the content of which is incorporated herein in their entirety by reference.

[0129] Another building block, which may form an amide bond, is




[0130] R may be absent or NO₂, CF₃, halogen, preferably Cl, Br, or I, and Z may be S or O. This type of building block is disclosed in WO03078626A2. The content of this patent application is incorporated herein in the entirety by reference.

[0131] A nucleophilic group can cleave the linkage between Z and the carbonyl group thereby transferring the chemical entity —(C=O)—CE' to said nucleophilic group.

[0132] Building Blocks Transferring a Chemical Entity to a Recipient Reactive Group Forming a C=C Bond


[0133] A building block as shown below is able to transfer the chemical entity to a recipient aldehyde group thereby forming a double bond between the carbon of the aldehyde and the chemical entity



[0134] The above building block is disclosed in WO03078445A2, the content of which being incorporated herein in the entirety by reference.


[0135] Building Blocks Transferring a Chemical Entity to a Recipient Reactive Group Forming a C—C Bond

[0136] The below building block is able to transfer the chemical entity to a recipient group thereby forming a single bond between the receiving moiety, e.g. a scaffold, and the chemical entity.



[0137] The above building block is disclosed in WO03078445A2, the content of which being incorporated herein in the entirety by reference.

[0138] Another building block capable of transferring a chemical entity to a receiving reactive group forming a single bond is



[0139] The receiving group may be a nucleophile, such as a group comprising a hetero atom, thereby forming a single bond between the chemical entity and the hetero atom, or the receiving group may be an electronegative carbon atom, thereby forming a C—C bond between the chemical entity and the scaffold. The above building block is disclosed in WO03078446A2, the content of which is incorporated herein by reference.

[0140] The chemical entity attached to any of the above building blocks may be selected from a large arsenal of chemical structures. Examples of chemical entities are H or entities selected among the group consisting of a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R⁴, 0-3 R⁵ and 0-3 R⁹ or C₁-C₃ alkylene-NR⁴₂, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O—NR⁴₂, C₁-C₂ alkylene-O—NR⁴C(O)R⁸, C₁-C₂ alkylene-O—NR⁴C(O)OR⁸ substituted with 0-3 R⁹.

[0141] where R⁴ is H or selected independently among the group consisting of C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R⁹ and

[0142] R⁵ is selected independently from —N₃, —CNO, —C(NO₂)NH₂, —NHOH, —NHNHR⁶, —C(O)R⁶, —SnR⁶₃, —B(OR⁶)₂, —P(O)(OR⁶)₂ or the group consisting of C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl said group being substituted with 0-2 R⁷,

[0143] where R⁶ is selected independently from H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-5 halogen atoms selected from —F, —Cl, —Br, and —I; and


[0144] R⁷ is independently selected from —NO₂, —COOR⁶, —COR⁶, —CN, —OSiR⁶₃, —OR⁶ and —NR⁶₂.

[0145] R⁸ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-3 substituents independently selected from —F, —Cl, —NO₂, —R³, —OR³, —SiR³₃

[0146] R⁹ is =O, —F, —Cl, —Br, —I, —CN, —NO₂, —OR⁶, —NR⁶₂, —NR⁶C(O)R⁸, —NR⁶C(O)OR⁸, —SR⁶, —S(O)R⁶, —S(O)₂R⁶, —COOR⁶, —C(O)NR⁶₂ and —S(O)₂NR⁶₂.

[0147] Cross-Link Cleavage Building Blocks

[0148] It may be advantageous to split the transfer of a chemical entity to a recipient reactive group into two separate steps, namely a cross-linking step and a cleavage step because each step can be optimized. A suitable building block for this two-step process is illustrated below:



[0149] Initially, a reactive group appearing on the chemical entity precursor (abbreviated FEP) reacts with a recipient reactive group, e.g. a reactive group appearing on a scaffold, thereby forming a cross-link. Subsequently, a cleavage is performed, usually by adding an aqueous oxidising agent such as I₂, Br₂, Cl₂, H⁺, or a Lewis acid. The cleavage results in a transfer of the group HZ-FEP- to the recipient moiety, such as a scaffold.

[0150] In the above formula

[0151] Z is O, S, NR⁴

[0152] Q is N, CR¹

[0153] P is a valence bond, O, S, NR⁴, or a group C₅₋₇arylene, C₁₋₆alkylene, C₁₋₆O-alkylene, C₁₋₆S-alkylene, NR¹-alkylene, C₁₋₆alkylene-O, C₁₋₆alkylene-S option said group being substituted with 0-3 R⁴, 0-3 R⁵ and 0-3 R⁹ or C₁-C₃ alkylene-NR⁴₂, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O—NR⁴₂, C₁-C₂ alkylene-O—NR⁴C(O)R⁸, C₁-C₂ alkylene-O—NR⁴C(O)OR⁸ substituted with 0-3 R⁹,

[0154] B is a group comprising D-E-F, in which

[0155] D is a valence bond or a group C₁₋₆alkylene, C₁₋₆alkenylene, C₁₋₆alkynylene, C₅₋₇arylene, or C₅₋₇heteroarylene, said group optionally being substituted with 1 to 4 group R¹¹,

[0156] E is, when present, a valence bond, O, S, NR⁴, or a group C₁₋₆alkylene, C₁₋₆alkenylene, C₁₋₆alkynylene, C₅₋₇arylene, or C₅₋₇heteroarylene, said group optionally being substituted with 1 to 4 group R¹¹,

[0157] F is, when present, a valence bond, O, S, or NR⁴,

[0158] A is a spacing group distancing the chemical structure from the complementing element, which may be a nucleic acid,

[0159] R¹, R², and R³ are independent of each other selected among the group consisting of H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R⁴, 0-3 R⁵ and 0-3 R⁹ or C₁-C₃ alkylene-NR⁴₂, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O—NR⁴₂, C₁-C₂ alkylene-O—NR⁴C(O)R⁸, C₁-C₂ alkylene-O—NR⁴C(O)OR⁸ substituted with 0-3 R⁹,

- [0160] FEP is a group selected among the group consisting of H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R⁴, 0-3 R⁵ and 0-3 R⁹ or C₁-C₃ alkylene-NR⁴, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O-NR⁴, C₁-C₂ alkylene-O-NR⁴C(O)R⁸, C₁-C₂ alkylene-O-NR⁴C(O)OR⁸ substituted with 0-3 R⁹,
- [0161] where R⁴ is H or selected independently among the group consisting of C₂-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R⁹ and
- [0162] R⁵ is selected independently from -N₃, -CNO, -C(NO₂)NH₂, -NHOH, -NHNHR⁶, -C(O)R⁶, -SnR⁶, -B(OR⁶)₂, -P(O)(OR⁶)₂ or the group consisting of C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl said group being substituted with 0-2 R⁷,
- [0163] where R⁶ is selected independently from H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-5 halogen atoms selected from -F, -Cl, -Br, and -I; and R⁷ is independently selected from -NO₂, -COOR⁶, -COR⁶, -CN, -OSiR⁶, -OR⁶ and -NR⁶.
- [0164] R⁸ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -NO₂, -R³, -OR³, -SiR³₃
- [0165] R⁹ is =O, -F, -Cl, -Br, -I, -CN, -NO₂, -OR⁶, -NR⁶, -NR⁶-C(O)R⁸, -NR⁶-C(O)OR⁸, -SR⁶, -S(O)R⁶, -S(O)₂R⁶, -COOR⁶, -C(O)NR⁶₂ and -S(O)₂NR⁶.
- [0166] In a preferred embodiment Z is O or S, P is a valence bond, Q is CH, B is CH₂, and R¹, R², and R³ is H. The bond between the carbonyl group and Z is cleavable with aqueous I₂.
- [0167] Partitioning Conditions
- [0168] The partition step may be referred to as a selection or a screen, as appropriate, and includes the screening of the library for encoded molecules having predetermined desirable characteristics. Predetermined desirable characteristics can include binding to a target, catalytically changing the target, chemically reacting with a target in a manner which alters/modifies the target or the functional activity of the target, and covalently attaching to the target as in a suicide inhibitor.
- [0169] The target can be any compound of interest. E.g. the target can be a protein, peptide, carbohydrate, polysaccharide, glycoprotein, hormone, receptor, antigen, antibody, virus, substrate, metabolite, transition state analogue, cofactor, inhibitor, drug, dye, nutrient, growth factor, cell, tissue, etc. without limitation. Particularly preferred targets include, but are not limited to, angiotensin converting enzyme, renin, cyclooxygenase, 5-lipoxygenase, IIL-10 converting enzyme, cytokine receptors, PDGF receptor, type II inosine monophosphate dehydrogenase, β-lactamases, integrin, and fungal cytochrome P-450. Targets can include, but are not limited to, bradykinin, neutrophil elastase, the HIV proteins,
- including tat, rev, gag, int, RT, nucleocapsid etc., VEGF, bFGF, TGFβ, KGF, PDGF, thrombin, theophylline, caffeine, substance P, IgE, sPLA2, red blood cells, glioblastomas, fibrin clots, PBMCs, hCG, lectins, selectins, cytokines, ICP4, complement proteins, etc.
- [0170] Encoded molecules having predetermined desirable characteristics can be partitioned away from the rest of the library while still attached to the identifier nucleic acid sequence by various methods known to one of ordinary skill in the art. In one embodiment of the invention the desirable products are partitioned away from the entire library without chemical degradation of the attached nucleic acid identifier such that the identifiers are amplifiable. The identifiers may then be amplified, either still attached to the desirable encoded molecule or after separation from the desirable encoded molecule.
- [0171] In a preferred embodiment, the desirable encoded molecule acts on the target without any interaction between the nucleic acid attached to the desirable encoded molecule and the target. In one embodiment, the bound complex-target aggregate can be partitioned from unbound complexes by a number of methods. The methods include nitrocellulose filter binding, column chromatography, filtration, affinity chromatography, centrifugation, and other well known methods.
- [0172] Briefly, the library of complexes is subjected to the partitioning step, which may include contact between the library and a column onto which the target is immobilised. Identifier nucleic acids associated with undesirable encoded molecules, i.e. encoded molecules not bound to the target under the stringency conditions used, will pass through the column. Additional undesirable encoded molecules (e.g. encoded molecules which cross-react with other targets) may be removed by counter-selection methods. Desirable complexes are bound to the column and can be eluted by changing the conditions of the column (e.g., salt, pH, surfactant, etc.) or the identifier.
- [0173] Additionally, encoded molecules which react with a target can be separated from those products that do not react with the target. In one example, a chemical compound which covalently attaches to the target (such as a suicide inhibitor) can be washed under very stringent conditions. The resulting complex can then be treated with proteinase, DNase or other suitable reagents to cleave a linker and liberate the nucleic acids which are associated with the desirable chemical compound. The liberated nucleic acids can be amplified.
- [0174] In another example, the predetermined characteristic of the desirable product is the ability of the product to transfer a chemical group (such as acyl transfer) to the target and thereby inactivate the target. One could have a product library where all of the products have a thioester chemical group. Upon contact with the target, the desirable products will transfer the chemical group to the target concomitantly changing the desirable product from a thioester to a thiol. Therefore, a partitioning method which would identify products that are now thiols (rather than thioesters) will enable the selection of the desirable products and amplification of the nucleic acid associated therewith.
- [0175] There are other partitioning and screening processes, which are compatible with this invention that are

known to one of ordinary skill in the art. In one embodiment, the products can be fractionated by a number of common methods and then each fraction is then assayed for activity. The fractionization methods can include size, pH, hydrophobicity, etc.

[0176] Inherent in the present method is the selection of encoded molecules on the basis of a desired function; this can be extended to the selection of molecules with a desired function and specificity. Specificity can be required during the selection process by first extracting identifier nucleic acid sequences of chemical compounds which are capable of interacting with a non-desired "target" (negative selection, or counter-selection), followed by positive selection with the desired target. As an example, inhibitors of fungal cytochrome P450 are known to cross-react to some extent with mammalian cytochrome P450 (resulting in serious side effects). Highly specific inhibitors of the fungal cytochrome could be selected from a library by first removing those products capable of interacting with the mammalian cytochrome, followed by retention of the remaining products which are capable of interacting with the fungal cytochrome.

BRIEF DESCRIPTION OF THE FIGURES

[0177] FIG. 1 illustrates the overall process of building block evolution.

[0178] FIG. 2 shows the distribution of codon in different positions in an output from a selection.

[0179] FIG. 3 shows the difference between identifier driven and building block driven evolution.

[0180] FIG. 4 shows a method for reducing the library diversity through codon analysis.

[0181] FIG. 5 discloses two embodiments of using a Taqman probe (5' nuclease probe) in the measurement of the presence or absence of a certain codon.

[0182] FIG. 6 shows a standard curve referred to in example 4.

[0183] FIG. 7 shows a result of example 4.

[0184] FIG. 8 discloses a result of example 4.

[0185] FIG. 9 discloses a scheme relating to combined structural information and codon abundances in library design.

[0186] FIG. 10 discloses a relationship between codon analysis and structural information.

[0187] FIG. 11 shows the detection of single codons of identifiers.

[0188] FIG. 12 shows the detection of codon pairs of identifiers.

[0189] FIG. 13 shows the detection of codon pairs at specific codon positions.

[0190] FIG. 14 shows the detection of single codons of identifiers after the separation of the individual codons.

[0191] FIG. 15 discloses a method for selecting from a library, complexes capable of binding to a target molecule.

[0192] FIG. 16 discloses a method for enriching specific nucleic acid fragments and the utility of these fragments for the generation of a new library.

[0193] FIG. 17 discloses a method for reducing the diversity of a library of complexes.

DETAILED DESCRIPTION OF THE FIGURES

[0194] FIG. 1A Shows the principle steps in BB evolution. An initial library of desired size is produced. This initial library is subjected to a selection process where encoded molecules that associate with a target of interest are enriched. The encoding identifier oligonucleotide is preferably amplified and the used in the codon analysis step. This step monitors the relative abundance of each codon in the selected library. The information obtained in this analysis is used to design a new enriched library, which contains the preferable chemical entities and their corresponding codons. This new library is then subjected to a new selection process to select for binders. This diversity reduction cycle can be repeated until the desirable result is obtained and the binders have been obtained.

[0195] FIG. 1B shows how the diversity of a library (n^4) is reduced by reducing the number of chemical entities (n) in the library. Thus, by removing chemical entities not involved in the encoded molecules partitioned, a reduction in library diversity can be obtained to allow the identification of binders.

[0196] The identifier oligonucleotide that encodes for the display molecule is composed of codons and associated with the encoded molecule, as shown in FIG. 2. These codons possess information about the chemical entities in the encoded molecule. Each of these codon positions can be analysed for the precise sequence, which will reflect which chemical entities that have been enrich for in the selection process. The relative amount can also be obtained by comparing the signal in the measuring procedure (e.g. QPCR and array analysis). Each codon position will have its own fingerprint on which chemical entities that the selected display molecules possess. These fingerprints in each position can subsequently be used to put together a new more focused library with a lower and more enrich diversity that can be subjected to another round of selection. This can then repeated until the preferable encoded molecules have been obtained.

[0197] FIG. 3 illustrates the main difference between identifier and chemical entity (CE) evolution. In both cases the initial selection starts on a library with certain diversity. After the first round of selection the encoding identifiers are amplified where the distribution is maintained. This distribution is then transferred to the next generation which is used in a new selection. Thus, the strongest binders that were enriched in the first round of selection will be present at a relatively higher concentration compared to the weaker binders and the background. In the CE-driven evolution the codon analysis is used to design a new library. In this example, the new library is constructed to contain all the chemical entities that were identified as a positive signal in the analysis. In other words, all the chemical entities that were not detected through the codon analysis were excluded in the new library. The new library is designed to have an equal amount of each selected chemical entity, which will generate all the possible display molecules at the same concentration. This will allow all binders to compete at the same concentration and potentially retain a more diverse set of binders in each round of selection. This is especially important for small molecules here not only the affinity is of interest.

[0198] FIG. 4. This illustrates the process where the diversity is reduced through the codon analysis. An initial library of 10^{10} (e.g. 317*317*317*317) library members is subjected to a selection. The enrich identifier oligonucleotides are amplified and used in the codon analysis. The codon analysis result is used to design a new 10^7 (e.g. 57*57*57*57) library where the enriched chemical entities are included. This new library is again subjected to a selection process. The identifier oligonucleotides are amplified and used for codon analysis. This new codon analysis results is again used to design a new 10^4 (e.g. 10*10*10*10) library where the enriched chemical entities are included. Finally a last selection step is performed in this reduced diversity library to identify the binders.

[0199] A preferred embodiment of the invention utilizing a universal Taqman probe is shown in FIG. 5. Four codons are shown (P1 through P4; bold pattern) along with flanking regions (light pattern). A universal Taqman probe anneals to a region adjacent to the codon region, but within the amplicon defined by the universal PCR primers Pr.1 and Pr. 2. These primers could be the same as used for amplification of the identifier oligonucleotides encoding binders after an enrichment process on a specific target. However, are minimal length identifiers preferred during the encoding process, the region involved in Taqman probe annealing could be appended to the library identifier oligonucleotides by e.g. overlap PCR, ligation, or by employing a long downstream PCR primer containing the necessary sequences. The added length corresponding to the region necessary for annealing of the Taqman probe would be from 20 to 40 nts depending on the type of TaqMan probe and T_A of the PCR primers. The Q-PCR reactions are preferably performed in a 96- or 384-well format on a real-time PCR thermocycling machine.

[0200] Panel A shows the detection of abundance of a specific codon sequence in position one. Similar primers are prepared for all codon sequences. For each codon sequence utilized to encode a specific BB in the library a Q-PCR reaction is performed with a primer oligonucleotide complementary to the codon sequence in question. A downstream universal reverse primer Pr. 2 is provided after the Taqman probe to provide for an exponential amplification of the PCR amplicon. The setup is most suited for cases where the codon constitutes a length corresponding to a length suitable for a PCR primer.

[0201] Panel B shows the detection of abundance of a specific codon sequence in a specific codon position using a primer, which is complementing a codon and a framing sequence. Similar primers are used for all the codons and framing sequences. For each codon sequence utilized to encode a specific BB at a specific codon position in the library a Q-PCR reaction is performed with an oligo complementary to the codon sequence in question as well as a short region up- or downstream of the codon region which ensures extension of the primer in a PCR reaction only when annealed to the codon sequence in that specific codon position. The number of specific primers and Q-PCR reactions needed to cover all codon sequences in all possible codon positions equals the number of codon sequences times the number of codon positions. Thus, monitoring the abundance of 96 different codon sequences in 4 different positions can be performed in a single run on four 96 wells micro titre plates (as shown in Panel B) or a single 384 well plate

on a suitable instrument This architecture allows for the decoding of a $8,5 \times 10^7$ library of different encoded molecules.

[0202] Quantification is performed relative to the amount of full-length PCR product obtained in a parallel control reaction on the same input material performed with the two external PCR primers Pr.1+Pr. 2. Theoretically, a similar rate of accumulation of this control amplicon compared to the accumulation of a product utilizing a single codon+sequence specific primer would indicate a 100% dominance of this particular sequence in the position in question.

[0203] Although the setups shown in Panel A and B employ a Taqman probe strategy, other detection systems (SYBR green, Molecular Beacons etc.) could be utilized. In theory, multiplex reactions employing up to 4 different fluorophores in the same reaction could increase throughput correspondingly.

[0204] An example of how a deconvolution process of a library of encoded molecules occurs is described in the following. Imagine that at the end of a selection scheme a pool of 3 ligand families (and the corresponding coding identifiers) are dominating the population and present at approx. the same concentration. Three different chemical entities are present in the first position of the encoded compounds, and each of these chemical entities are present in combination with one unique chemical entity out of 3 different chemical entities in position P2. Only one chemical entity in position 3 gives rise to active binders, whereas any of a 20% subset of chemical entities (e.g. determined by charge, size or other characteristics) is present in position 4. The outcome of the initial codon profile analysis would be: 3 codon sequences are equally dominating in position P1, 3 other codon sequences in position P2, 1 unique codon sequence is dominant in P3 whereas somewhat similarly increased levels of 20% of the codon sequences (background levels of the remaining 80% sequences) are seen in P4. In such cases it could be relevant to use an iterative Q-PCR ("IQ-PCR") strategy to perform a further deconvolution of a library after selection. Again with reference to the example above, by taking the PCR products from the 3 individual wells that contained primers giving the high yields in position P1, diluting the product appropriately and performing a second round of Q-PCR on each of these identifier oligonucleotides separately, it would be possible to deduce which codon sequence(s) is preferred in P2 when a given codon sequence is present in P1.

[0205] FIG. 9. This figure illustrates the possibility to combine structural information about the chemical entities and the relative abundance when designing a new more focused library. The structural information about the chemical entities can be used at least in two ways. First the similarities between the chemical entities in each position can be used to choose chemical entities to a new library. Secondly, the combination of the selected chemical entities can be analyzed to investigate possible pattern that generate potential ligands. This is especially useful if the binding site or the structure of a known ligand is known. Any type of structural analysis tool can be used that generate information about the structure of separate chemical entities or combination of chemical entities (the potential binders). By combining these three analysis approaches a more focused library can be generated that potentially will contain more

specific binders compare to background binders. This new focused library can be used in another round of selection to reduce the diversity. This procedure can be repeated until the desired binders have been identified.

[0206] FIG. 10. This figure shows how the combination of codon analysis and structural information can generate valuable information. This invention allows the performance of structure activity relationship analysis (SAR) where the relative abundance in the codon analysis will represent the activity parameter (e.g. IC_{50} values) in the SAR measurements. Pharmacophore models can be generated, focused libraries can be designed, certain follow up chemistry can be used and information in the hit to lead process can be used.

[0207] FIG. 11 shows an array detection system in which a single codon is detected. Initially a library of selected complexes (29), i.e. complexes comprised of the initial library, which display a certain property, is provided as disclosed above. The initial library of complexes is prepared from e.g. 100 codons and identifiers having 4 codons in sequence, which theoretical gives a library of 10^8 complexes. The selected complexes are subjected to amplification to amplify the identifiers of the selected complexes and the amplification products are added to an array (30). The array (30) comprises probes (32) complementary to each of the codons of the identifiers (31). At hybridisation conditions the PCR products of the identifiers are annealed to the cognate probes of the array and in a suitable scanner the spatial position of the annealed probes are detected to elucidate the codons (33) of the identifier. The quantity of each codon may be measured to find codons abundant in more than one identifier and/or codons leading to encoded molecules with high affinity. The information may be used for decoding of the encoded molecule of the complexes displaying the desired property or the information may be used for selection of building blocks, which is to be added in a next round of library formation.

[0208] FIG. 12 discloses an array detection system for establishing codons pairs, i.e. codons in the vicinity of each other. Initially (as shown in this example) a library of complexes is prepared from 100 different codons deposited on an identifier in a sequence of four, making the total amount of combinations possible 10^8 . The initial library is subjected to a condition in order to select a sub-library (29) displaying a desired property. The identifiers of the sub-library are amplified by a PCR reaction and the reaction product is added under hybridisation conditions to an array (34).

[0209] The array is designed with probes (35) capable of detecting two codons at a time. To cover all possible combinations of a library based on 100 different codons 10^4 probes are needed, which is practically feasible with the current technology. The detection of the codons may be conducted quantitatively, i.e. the relative abundance of each of the codon pairs may be determined. The detection on the array may be used to reconstruct the selected identifiers (36) as three overlapping codon pair detections depict the entire identifier. In the event the same codon pair appears on more than one identifier, the information on the relative abundance of each codon pair maybe used to decipher the sequence of codons of the selected identifiers as it can be assumed that each codon pair of the same identifier appears in the same amounts in the PCR products added to the array.

[0210] FIG. 13 discloses an array for detecting codon pairs at specific codon positions. Initially, a library of complexes comprising identifiers with framing sequences is provided. The framing sequence is specific for each position of the codons on the identifier. Four times more probes on the microarray is needed per each codon if the position of the codons also should be detected in the analysis which is practically feasible with current technology. The position is detected due to the framing sequences next to each codon. The initial library is subjected to a selection process to isolate complexes (37) having a desired property. The selected complexes are amplified by a PCR reaction and the reaction products are added to an array (38). The array comprises probes capable of detecting codon pairs as wells as the framing sequences (40) between the codons. The framing sequence determines the position of the codon in the reaction history, i.e. it is possible to deduct which chemical entity that reacted at which point in time of the synthesis history of the encoded molecule, thus making it possible to reconstruct the structure of the encoded molecule.

[0211] The detection of the codon pairs may be conducted quantitatively, i.e. the relative abundance of each of the codon pairs may be determined. The detection on the array may be used to reconstruct the selected identifiers (41) as three overlapping codon pair detections depict the entire identifier. In the event the same codon pair appears on more than one identifier, the information on the relative abundance of each codon pair maybe used to decipher the sequence of codons of the selected identifiers as it can be assumed that each codon pair of the same identifier appears in the same amounts in the PCR products added to the array.

[0212] FIG. 14 shows an array detection system in which a single codon is detected. Initially a library of selected complexes (42), i.e. complexes comprised of the initial library which display a certain property, is provided as disclosed above. The initial library of complexes is prepared from e.g. 100 codons and identifiers having 4 codons in sequence, which theoretical gives a library of 10^8 complexes. The selected complexes are subjected to amplification to amplify the identifiers of the selected complexes and the amplification products are treated with suitable reagents to cut between the individual codons (43). The individual codon is the applied to the array. The array (44) comprises probes (45) complementary to each of the codons of the identifiers (46). At hybridisation conditions the PCR products of the identifiers are annealed to the cognate probes of the array and in a suitable scanner the spatial position of the annealed probes are detected to elucidate the codons (47) of the identifier. The quantity of each codon may be measured to find codons abundant in more than one identifier and/or codons leading to encoded molecules with high affinity. The information may be used for decoding of the encoded molecule of the complexes displaying the desired property or the information may be used for selection of building blocks, which is to be added in a next round of library formation.

[0213] FIG. 15 discloses a method for selection of a suitable complex in several steps. In a first step the library of complexes 1 is provided. Each member of the library comprises an encoded molecule 2 composed of four chemical entities which is attached to an identifier oligonucleotide 3, which comprises four codons. The initial library shown comprises three complexes. In a second step the library of

complexes is incubated with immobilized target molecules 4. The encoded molecule having an affinity towards the target molecule is bound to the immobilized target whereas encoded molecules not having affinity towards the target under the conditions used remains in the liquid media. The complexes remaining in the liquid media are discarded by a washing process, while the bound complexes remain attached to the immobilized target molecules. The washing process is usually conducted using mild stringency conditions in the initial rounds of selection. In later stage selections the working stringency conditions are usually increased to allow only high affinity binders to remain attached to the target. Subsequent to the washing step the complexes having affinity towards the target molecule are recovered. The recovery process usually requires high stringency conditions to detach the encoded molecule from the immobilized target. The selected sub-library resulting from the elution is subjected to an amplification process. The amplification of the identifier nucleic acid sequence of the selected complexes is usually performed using the PCR method. Preferably, a modification of the PCR method is followed such that a biotin molecule is attached to one of the primers to obtain a handle for subsequent immobilization. The result of the amplification step is multiple copies of the identifier nucleic acid sequences, which codes for the encoded molecules which have survived the selection step.

[0214] FIG. 16 discloses an enrichment process of building blocks. The building blocks can be used for generation of a new library. Initially, identifier nucleic acid sequences are immobilized on solid support. In one aspect of the invention the identifier nucleic acid sequences are the product of the selection procedure described in FIG. 1. Each codon of the identifier nucleic acid sequence is identified with an uppercase letter, i.e. A, B, C, or D. The immobilized identifier acid sequences are contacted with the pool of building blocks under hybridisation conditions. Each of the building blocks are illustrated with a sequence complementary to a codon which may or may not be present on the identifier nucleic acid sequence. The complementary sequences are indicated with an apostrophe, e.g. A', B', etc. The transferable chemical entity of a building block is illustrated with a lowercase letter. The conditions providing for hybridisation of the complementing sequences of the pool of building blocks to the immobilised identifier nucleic acid sequence are preferably such that cognate nucleic acid sequences are hybridised to each other while sequences not recognizing any immobilized sequence remain in aqueous media. The immobilized sequences of the identifier nucleic acid sequences are thus used as bait in catching building blocks with complementing sequences. Following the incubation step, non-binding building blocks are removed by washing, whereby the part of the pool of building blocks not being able to find a complementing sequence is discarded. The building blocks attached to the immobilized nucleic acid sequences are detached using dehybridisation conditions. The diminished pool of building blocks may be used in a subsequent round for preparing a new library of complexes, in which the encoded molecule comprises a reaction product comprising additions from chemical entities attached to the enriched building blocks. Because the order of building blocks which have participated in the formation of the encoded molecules successful in the selection procedure, is not preserved by the method for enriching building blocks a scrambling of the encoded molecules may be

obtained in some of the methods described herein for obtaining a library of complexes. In some applications of the library it will be an advantage to have a scrambling of the building blocks because and increased diversity is obtained.

[0215] FIG. 17 discloses a method for reducing the diversity of the library of complexes resulting from the method described in FIG. 16. In some of the applications of the library the diversity induced by scrambling of the building blocks are not desired. In a first step the sequences complementary to the identifier acid sequences used in FIG. 16 are provided and immobilized on a suitable solid support. In one aspect of the invention the complementary sequence is obtained from the PCR product resulting from the method according to FIG. 15. Alternatively, the complementing sequence may be obtained by extending the identifier nucleic acid sequence using a suitable primer, optionally attached to a handle such as a biotin or dinitrophenol. In a second step the immobilized complementary sequence is incubated with the scrambled library under conditions, which provide for hybridisation between the complementary sequence and members of the library having affinity towards this sequence. Members of the library not having affinity to the complementary sequences remains in the media and is discarded, while members of the library being able to hybridise to the immobilized nucleic acid sequences is recovered. Occasionally, nucleic acids not perfectly matching with the complementary sequence immobilized on the solid support are caught. In one aspect of the invention the hybridisation products, prior to the recovery step, are treated with an enzyme capable of recognizing mismatching nucleotides and cleaving the double stranded helix in which they are situated. An example of an enzyme with this ability is T4 endonuclease VII. After the treatment with the enzyme, complexes displaying a hybridisation toward the immobilized sequence are eluted under dehybridisation conditions. Nucleotide sequences remaining from the cleavage by the enzyme will also be present in the new library, however, these sequences will not have any effect of a subsequent selection because no molecule is attached thereto.

EXAMPLES

Example 1

Enrichment of Nucleic Acid Fragments

[0216] A codon was included in the oligonucleotide sequence shown below. The codon is underlined and the boldface sequences represent the "framing" regions next to each codon. These framing regions can be used for specifying the position of each codon.

Biotin-AATTCCGGAAC**ATACTAGTCAAC**ATGA-3' (SEQ ID NO:1)

[0217] This identifier oligonucleotide was immobilized on streptavidin beads using standard protocols, i.e. 600 pmol identifier oligonucleotide with 5'-dT biotin in 50 µl 100 mM Mes pH 6.0 was mixed with 50 µl SA-magnetic beads (Roche). The mixture was washed 2-3 times with 100 mM MES pH 6.0 to remove non-bound identifier oligonucleotides. To reduce background binding, the oligos and beads were incubated at RT for 10 min on shaker, then incubated on ice for 10 min while rotating the tube. Finally, the sample was washed with 100 mM MES 4 times in 800 µl at 60° C.

[0218] In the case where a PCR product is immobilized, the complementing (non-sense) strand is removed using 10 mM NaOH. This will generate single-stranded DNA with the selected codons. The same procedure described in this example can be used for a collection of different identifier nucleic acid molecules that contain one or more codons. The codons in the identifier nucleic acid molecules can be the same or different determined from the enrichment performed on the initial library.


[0219] The immobilized identifier nucleic acid molecule was mixed with the pool of nucleic acid fragments shown below. This pool of fragments illustrates an original pool that was used for generating an initial library of complexes. Each fragment may possess in the 3'-end a specific chemical entity that is encoded by the codon sequence. These nucleic acid fragments contain a specific sequence in the codon region (underlined) while the framing region shown in boldface is identical among the fragments. Thus, the pool of fragments represents different codons in the same position of the identifier nucleic acid.

1. CGT GTG ATC GAA CTC GTG TG **GTATGATCAGTTG**TACT-5'
(SEQ ID NO:2)
2. CGT GTG ATC GAA CTC GTG TG **GTATCTAGTCGGT**TACT-5'
(SEQ ID NO:3)
3. CGT GTG ATC GAA CTC GTG TG **GTATTCGAGTGT**TTTA-5'
(SEQ ID NO:4)

-continued

4. CGT GTG ATC GAA CTC GTG TG **GTATAGCTCATGG**TACT-5'
(SEQ ID NO:5)


[0220] The nucleic acid fragments are mixed with the immobilized identifier nucleic acid using 600 pmol of each nucleic acid fragment mixed with the immobilized identifier nucleic acid molecules (100 mM MES pH 6.0, 150 mM NaCl). The mixture was incubated at 25° C. for 30 minutes in a shaker. The non-hybridized fragments were removed by 4 times washing in 800 µl 100 mM MES, 150 mM NaCl. This step should separate the complementing fragments (bound) encoding for the select chemical entities from the non-complementing fragments (non-bound) encoding for chemical entities that were not effective in the preceding selection process. The annealed fragments are eluted from the immobilized identifier nucleic acid molecules by re-suspending the beads in 25 µl 60° C. H₂O and incubating for 2 min at 60° C. The enriched fragments were purified on a micro-spin gel filtration column (BiRad). The eluted fragments were prepared for mass spectroscopy (MS) analysis by mixing in half volume of ion exchanger resin and incubating minimum 2 h at 25° C. on a shaker. After incubation the resin was removed by centrifugation and 15 µl of the supernatant was mixed with 7 µl of water, 2 µl of piperidine and imidazole (each 625 mM) and 24 µl acetonitrile. The sample was analysed using a Mass Spectroscopy instrument (Bruker Daltonics, Esquire 3000plus). The result for the MS analysis is shown below.



Component	Deconvoluted Mass	Molecule	Absolute Abundance	Relative Abundance
A.	11438.39	[M - H] -	112674	96.98 → MATCH
B	8663.73	[M - H] -	61549	53.06
C	6616.63	[M - H] -	41245	35.50

[0221] The mass from the correct complementary fragment (number 1) is obtained in the MS analysis (11438.39, expected 11439 Da) No masses for the other fragments (number 2-4) could not be found in the MS spectra (expected masses; 11415, 11430, 11424 Da). This result shows that the right fragment is strongly enriched and other fragments with the wrong codon sequences are removed. The enrichment is possible even when the "spacing" region (boldface) is


identical in each fragment. Two control experiments were also performed to validate the enrichment protocol. In the first experiment, the fragment with the correct codon sequence (number 1) was mixed with the immobilized identifier molecule as described above. The sample was washed and eluted also as described above and prepared for MS analysis. The result from the MS analysis is shown below.



Component	Deconvoluted Mass	Molecule	Absolute Abundance	Relative Abundance
A	11438.39	[M - H] -	127110	96.21 → MATCH
B	8663.61	[M - H] -	82800	62.67
C	6616.52	[M - H] -	51480	38.97

[0222] The result indicates that the fragment with the correct sequence (number 1) anneals to the immobilized identifier molecules and is eluted under the conditions used in this example. The expected mass (11439) correlate well with the experimental mass, 11438.39 Da.

[0223] In the other control experiment, a fragment with a wrong codon sequence (number 3) was allow to bind to the immobilized identifier molecule as described above. Again, the eluted sample was prepared and analysed with MS. The result is shown below.



Component	Deconvoluted Mass	Molecule	Absolute Abundance	Relative Abundance
A	8663.47	[M - H]-	89308	96.60 - <i>ExoSP1</i>
B	6616.52	[M - H]-	41963	45.39 - <i>PEIHI</i>
C	8678.68	[M - H]-	13651	14.77 - <i>Exot.</i>

[0224] In this experiment, no mass was found that corresponded to the expected mass (11430) of the tested building block (number 1). Again, this shows that fragments with an anticodon sequence different from the enriched codons in the identifier nucleic acid molecules are not captured using this approach.

[0225] The enriched fragments obtained using this strategy may then be used to generate a new library of encoded molecules. This new library will contain encoded molecules composed of the enriched chemical entities. Thus, the library size have been reduced due to the removal of chemical entities not involved in binding encoded molecules, and enriched in chemical entities that are highly represented in the encoded molecules which binds to the target molecule.

[0226] Example 1 shows the possibility of enriching for specific building block molecules, i.e. nucleic acid fragments associated with transferable chemical entities. The same procedure can be used for a larger pool of building block than four as used herein. The codon design will determine the maximum number of building blocks that can be used. The sequence in the codon region should be large enough to allow discrimination in the annealing step. Various conditions can be used to increase the stringency in the annealing step. Parameters such as temperature, salt, pH, formamide concentration, time and other conditions could be used.

Example 2

(Model): Multiple Codon Selection in a Library

[0227] This example describes the enrichment of building blocks using an identifier nucleic acid (identifier) molecule with multiple codons. These codons encode for a displayed molecule (DM) that are attached to the identifier molecule before the selection is performed. The library size is determined both by the number of different chemical entities and the total number of chemical entities. The identifier molecule shown below contains three codons. The codons, which codes for the displayed molecule are indicated with underlines and the region separating (framing region) the codons in boldface. The size of the codons can be varied dependent in the diversity need in the library and the optimal setup for chemical entity enrichment. The framing region can also be varied dependent on the discrimination needed to distinguish the precise position of a codon in the identifier molecule. The framing region will also be important for the generation of the library. This can be understood when the encoding is accomplished by extension of the encoding region as disclosed in DK PA 2002 01955 and U.S. No. 60/434,425, incorporated herein by reference. There need to be a perfect match in the 3'-end in order to get efficient extension with a polymerase or a ligase. The size of this spacing/framing region should be long enough to form a

complementing region to allow extension with a polymerase or ligase. Preferably, the spacing region should be between 3 and 6 nucleotides. The codon region together with the spacing region will also be useful when codons are to be identified using a micro array setup. The identifier molecule with the right codon sequences will hybridize to the array and be detected.

[0228] The sequence below represents an enriched identifier molecule attached to the displayed molecule (DM). This identifier molecule has been enriched due to the fact that the DM binds to the target molecule in the selection process. In practice, more than one enriched identifier molecules will be obtained when using a library of displayed molecules attached to its identifier sequence.

DM-GCACACTAGCTTGAGCACACTGACACATGGAGATCACATGCTTCGAC
AAATGCAGGACTCCCGCAGCTTTACGATCCCGCAGGTAACCGT

[0229] This identifier molecule is amplified with two primers (below) using a standard PCR reaction. For example, 500 nM of each primer, 2,5 units Taq polymerase, 0.2 mM of each NTP, in a PCR buffer (50 mM KCl, 10 mM Tris-Cl, 3 mM DTT, 1.5 mM MgCl₂, 0.1 mg/ml BSA). Run 25 cycles (94° C. melt for 30 seconds, 55° C. anneal for 45 seconds, 72° C. extension for 60 seconds).

B-GCACACTAGCTTGAGCACACTGACA-3'
 CGAAATGCTAGGGCGTCCATTGGCA-5'

[0230] This will amplify the identifier molecule from the selection process and add a biotin in the 5'-end of one of the strand (below). This amplified product is then immobilized on a solid support, streptavidin beads for example. This can be performed identical as describe in example 1.

[0231] When the identifier molecules have been immobilized and the excess has been removed by a washing step (as describe in example 1), the complementing non-sense stand is removed by incubating in 10 NaOH for about 2 min and washed with 100 mM Mes buffer, pH 6.0. This procedure will generate the strand shown below where the codon regions are exposed to allow hybridization with the complementing sequences.

B-GCACACTAGCTTGAGCACACTGACACATGGAGATCACATGCTTCGACA
ATGCAGGACTCCCGCAGCTTTACGATCCCGCAGGTAACCGT

[0232] The next step is to protect the complementing sequences outside the codons to prevent the binding of the building block to these sequences. This can be performed by adding "blocking" oligonucleotides that has a complementing sequence. This is shown below.

B-
GCACACTAGCTTGAGCACACTGACA [Ⓢ]CTTCGACAATGCAGGACTCCCGCAGCTTTACGATCCCCAGGTAACC
GT
CGTGTGATCGAACTCGTGTGACTGT CGAAATGCTAGGGCGTCCATT-
GGCA

[Ⓢ] indicates text missing or illegible when filed

[0233] Next, the pool of different building blocks is added and is allowed annealing to the codon region in the identifier region. The position of annealing is determined by the spacing region shown in boldface. The stringency is adjusted to only allow hybridization of the correct building block in the right position. This can be accomplished by mixing the right component together using various conditions. The condition can for example include the presence of salt, formamide and various buffers adjusted to suitable pH and temperature. Below is the correct building block that will anneal to the enriched identifier molecules. These building blocks is annealed and eluted as described in example 1.

CE-CGTGTGATCGAACTCGTGTGACTGTGTACCTCTAGTGT**TAC**

[0234] The next pool of building blocks is blocked with an oligonucleotide that also protects the first codon. This is necessary to prevent binding of the building blocks in that codon.

B-
GCACACTAGCTTGAGCACACTGACACATGGAGATCAC [Ⓢ]AGGACTCCCGCAGCTTTACGATCCCGCAGGTAACC
GT
CGTGTGATCGAACTCGTGTGACTGTGT**A**IIIIIIIIII CGAAATGCTAGGGCGTCCATT-
GGCA

[Ⓢ] indicates text missing or illegible when filed

[0235] Again, the library of building blocks is added to enrich for the selected codons. Below is the building block with the correct sequence. These building blocks is annealed and eluted as described in example 1.

CE--CGTGTGATCGAACTCGTGTGACTGTGT**A**IIIIIIIIII**TAC**GAACT
GTTACG

[0236] Finally, the identifier molecule is protected with a blocking oligo that expose only the last codon.

B--
GCACACTAGCTTGAGCACACTGACACATGGAGATCATGCTTCGACAA ~~CGAAATGCTAGGGCGTCCATT~~ [Ⓢ] ~~CGAAATGCTAGGGCGTCCATT~~ GCTTTACGATCCCGCAGGTAACC
GT
CGTGTGATCGAACTCGTGTGACTGTGT**A**IIIIIIIIII**TAC**IIIIIIIIII CGAAATGCTAGGGCGTCCATT-
GGCA

[Ⓢ] indicates text missing or illegible when filed

[0237] A new pool of building blocks is added and allowed hybridizing to the identifier molecule. These building blocks is annealed and eluted as described in example 1.

CE--CGTGTGATCGAACTCGTGTGACTGTGT**A**IIIIIIIIII**TAC**IIIIIIIIII
IIII**AC**GTCTCGAGGGCGT


[0238] The enrichment of each library of building blocks are performed in separate tubes in order to keep the libraries of building block separated. The enrichment is performed with building blocks loaded with chemical entities (CE).

Example 3

Template Versus Chemical Entity Evolution


[0239] The graph below illustrates the relationship between the number of chemical entities and the library size.

The example below is calculated on that the final encoded molecules contains four chemical entities that is individually encoded by the corresponding building block (n^4 , where n is the number of building blocks). The graph shows that the diversity decreases dramatically with the reduction of the total number of building blocks. If the number of different building can be reduced to about 20-30 (library size of $16 \cdot 10^3$ and $81 \cdot 10^4$, respectively) in the selection process, then the library size for the final round of selection is low enough for identification of the binding molecules.



[0240] When the same analysis is performed on a protein another situation is obtained. The example shown below is on a very small protein (50 amino acids in length). The diversity is enormous when all amino acids are included in the library. The size of the library is also decreasing with the total number of amino acids, but not to the same extent as

show above for a small molecule. Even when the different amino acids are reduced to 2, the library size is huge ($1.2 \cdot 10^{15}$). This shows that amino acid enrichment is impossible on protein. This is even more pronounced for mid-size protein which contains about 300 amino acids.



Example 4

Codon Analysis

[0241] This example illustrates one possibility to perform codon analysis on a whole population of different identifier oligonucleotides. The analysis can also be performed using array where the probe oligonucleotides (complementary to the codons) are immobilized in discreet areas and the signal is monitored dependent on the amount of identifiers oligonucleotides are hybridised in each specific area. Codon analysis can also be performed using standard sequencing using a polymerase extension step.

[0242] In FIG. 5, Four codons are shown (P1 through P4; bold pattern) along with flanking regions (light pattern). A universal Taqman probe anneals to a region adjacent to the codon region, but within the amplicon defined by the universal PCR primers Pr.1 and Pr. 2. These primers could be the same as used for amplification of the identifier oligonucleotides encoding binders after an enrichment process on a specific target. However, are minimal length identifiers preferred during the encoding process, the region involved in Taqman probe annealing could be appended to the library identifier oligonucleotides by e.g. overlap PCR, ligation, or by employing a long downstream PCR primer containing the necessary sequences. The added length corresponding to the region necessary for annealing of the Taqman probe would be from 20 to 40 nts depending on the type of TaqMan probe and T_A of the PCR primers. The Q-PCR reactions are preferably performed in a 96 or 384-well format on a real-time PCR thermocycling machine.

[0243] FIG. 5, panel A, shows the detection of abundance of a specific codon sequence in position one. Similar primers are prepared for all codon sequences. For each codon sequence utilized to encode a specific BB in the library a Q-PCR reaction is performed with a primer oligonucleotide complementary to the codon sequence in question. A downstream universal reverse primer Pr. 2 is provided after the Taqman probe to provide for an exponential amplification of the PCR amplicon. The setup is most suited for cases where the codon constitutes a length corresponding to a length suitable for a PCR primer.

[0244] FIG. 5, panel B shows the detection of abundance of a specific codon sequence in a specific codon position using a primer which is complementing a codon and a framing sequence. Similar primers are used for all the codons and framing sequences. For each codon sequence utilized to encode a specific BB at a specific codon position in the library a Q-PCR reaction is performed with an oligo complementary to the codon sequence in question as well as a short region up- or downstream of the codon region which ensures extension of the primer in a PCR reaction only when annealed to the codon sequence in that specific codon

position. The number of specific primers and Q-PCR reactions needed to cover all codon sequences in all possible codon positions equals the number of codon sequences times the number of codon positions. Thus, monitoring the abundance of 96 different codon sequences in 4 different positions can be performed in a single run on four 96 wells micro titre plates (as shown in FIG. 5, panel B) or a single 384 well plate on a suitable instrument. This architecture allows for the decoding of a 8.5×10^7 library of different encoded molecules.

[0245] Quantification is performed relative to the amount of full-length PCR product obtained in a parallel control reaction on the same input material performed with the two external PCR primers Pr.1+Pr. 2. Theoretically, a similar rate of accumulation of this control amplicon compared to the accumulation of a product utilizing a single codon+sequence specific primer would indicate a 100% dominance of this particular sequence in the position in question.

[0246] Although the setups shown in FIG. 5, panel A and B employ a Taqman probe strategy, other detection systems (SYBR green, Molecular Beacons etc.) could be utilized. In theory, multiplex reactions employing up to 4 different fluorophores in the same reaction could increase throughput correspondingly.

[0247] An example of how a deconvolution process of a library of encoded molecules occurs is described in the following. Imagine that at the end of a selection scheme a pool of 3 ligand families (and the corresponding coding identifiers) are dominating the population and present at approx. the same concentration. Three different chemical entities are present in the first position of the encoded compounds, and each of these chemical entities are present in combination with one unique chemical entity out of 3 different chemical entities in position P2. Only one chemical entity in position 3 gives rise to active binders, whereas any of a 20% subset of chemical entities (e.g. determined by charge, size or other characteristics) are present in position 4. The outcome of the initial codon profile analysis would be: 3 codon sequences are equally dominating in position P1, 3 other codon sequences in position P2, 1 unique codon sequence is dominant in P3 whereas somewhat similarly increased levels of 20% of the codon sequences (background levels of the remaining 80% sequences) are seen in P4. In such cases it could be relevant to use an iterative Q-PCR ("IQ-PCR") strategy to perform a further deconvolution of a library after selection. Again with reference to the example above, by taking the PCR products from the 3 individual wells that contained primers giving the high yields in position P1, diluting the product appropriately and performing a second round of Q-PCR on each of these identifier oligonucleotides separately, it would be possible to deduce which codon sequence(s) is preferred in P2 when a given codon sequence is present in P1.

Identifiers used for Q-PCR quantification

	P1	P2
5' -	<u>CAGCTTGGACACCACGTCATAC</u> TAGCTGCTAGAGATGTGGTGATATTAGTGTGTGACGATGGTACGCACA	
	GGAAGAAGACAGAAGACCTG	TCAGGAGTCGAGAACTGAAG
	TGTGTACGTC AACACGTCAG	

-continued

Identifiers used for Q-PCR quantification

TGTGGAACCTACCATCCAAGG
 CCATCCAACATCGTTGGAAG
 AACCTGTCCCTGTGAGACTG
 TCACGAAGCTGGATGATGAG
 TAGCATCGATCGAACGTAGG
 TCGAAGCTACTGTGAGATC

P3

P4

AGTACGAACTGCATCAGAGAGGACGAGCAGGACCTGGAACCTGGTGC*TTCCTCCACCAGTCTCTGAC-3'

CTCGACCACTGCAGGTGGAGCTCC
 CGTGCTTCCTCTGCTGCCACCACC
 CCTGGTGTGAGGTGAGCAGCAGC
 CTCGACGAGGTCCATCCTGGTCCG
 CGTGAGGAGCAGGTCCCTCTGTCC
 CCTGACACTGGTCTGGTCCGAGGC
 CCATCTCGACGACCTGCTCCTGGG
 CCACGAGTCTCCACTGTGTCAGG
 CCACTGAGCTGCTCCTCCAGGTGC

Oligos for Identifier synthesis:

FPv2: CAGCTTGGACACCACGTTCATAC
 RPv2: GTCAGAGACGTGGTGGAGGAA

Temp1-1: CAGCTTGGACACCACGTTCATACACTAGCTGCTAGAGATGTGGTATATTAGTGTGTGACGAT
 Temp1-2: CAGCTTGGACACCACGTTCATACGGAAGAAGACAGAAAGACCTGATATTAGTGTGTGACGAT
 Temp1-3: CAGCTTGGACACCACGTTCATACACTCAGGAGTCGAGAACGAAAGATATTAGTGTGTGACGAT
 Temp1-4: CAGCTTGGACACCACGTTCATACACTGTGTACGTCAACACGTCAGATATTAGTGTGTGACGAT
 Temp1-5: CAGCTTGGACACCACGTTCATACACTGTGGAACCTACCATCCAAGGATATTAGTGTGTGACGAT
 Temp1-6: CAGCTTGGACACCACGTTCATACCATCCAACATCGTTGGAAGATATTAGTGTGTGACGAT
 Temp1-7: CAGCTTGGACACCACGTTCATACAACTGTCTGTGAGATCTGATATTAGTGTGTGACGAT
 Temp1-8: CAGCTTGGACACCACGTTCATACACTCAGAAAGCTGGATGATGAGATATTAGTGTGTGACGAT
 Temp1-9: CAGCTTGGACACCACGTTCATACACTAGCATCGATCGAACGTAAGGATATTAGTGTGTGACGAT
 Temp1-10: CAGCTTGGACACCACGTTCATACACTCGAAGCTACTGTGAGATGATATTAGTGTGTGACGAT

Temp2: GTCCTCTCTGATGCACGTTCTGACTTGTGCGTACCATCGTCACACACTAATATC

Temp3-1: GAACGTGCATCAGAGAGGACGAGCAGGACCTGGAACCTGGTGCATTCAGCTTCTAGGAAGACT
 Temp3-2: GAACGTGCATCAGAGAGGACCTGACCACTGCAGGTGGAGCTCCAATTCAGCTTCTAGGAAGACT
 Temp3-3: GAACGTGCATCAGAGAGGACGTGCTTCTCTGCTGACACCACGAATTCAGCTTCTAGGAAGACT
 Temp3-4: GAACGTGCATCAGAGAGGACCTGGTGTGCGAGGTGAGCAGCAGCAATTCAGCTTCTAGGAAGACT
 Temp3-5: GAACGTGCATCAGAGAGGACCTGACGAGGTCCATCCTGGTGCATTCAGCTTCTAGGAAGACT
 Temp3-6: GAACGTGCATCAGAGAGGACGTGAGGAGCAGGTCTCTGTCGAATTCAGCTTCTAGGAAGACT
 Temp3-7: GAACGTGCATCAGAGAGGACCTGACACTGGTCTGGTTCGAGGCAATTCAGCTTCTAGGAAGACT
 Temp3-8: GAACGTGCATCAGAGAGGACCTCGACGACCTGCTCCTGGGAATTCAGCTTCTAGGAAGACT
 Temp3-9: GAACGTGCATCAGAGAGGACCTCAGAGTCTCCACTGGTCCAGGAATTCAGCTTCTAGGAAGACT
 Temp3-10: GAACGTGCATCAGAGAGGACCTGAGCTGCTCCTCAGGTGGAATTCAGCTTCTAGGAAGACT

Temp4: GTCAGAGACGTGGTGGAGGAACTTCTCCTAGAAGCTGGAATT

Taqman MGB probe binding region: * = AATTCAGCTTCTAGGAAGAC

[0248] Synthesis of Identifier Oligonucleotides:

[0249] The 10 identifier oligonucleotides were assembled in 10 separate 50 μ l PCR reactions each containing 0.05 pmol of the oligos Q-Temp1-X, Q-Temp2, Q-Temp3-X and Q-Temp4 (x=1 through 10) and 25 pmol of the external primers FPv2 and RPv2 with TA=53° C. The 160 bp products were gel-purified using QIAquick Gel Extraction Kit from QIAGEN (Cat. No. 28706) and quantified on spectrophotometer. As a control, 20 ng of each of the identifiers (as estimated from these measurements) were loaded on an agarose gel.

[0250] Preparation of Samples for Q-PCR:

[0251] Sample A Generated by mixing 20 ng from each identifier oligonucleotide prep. Volume was adjusted to 50 μ l. Concentration: 4 ng/ μ l=38.46 fmol/ μ l (160 bp \times 650

Da/bp=1.04 \times 10⁵ g/mol. 1 ng=9.615 fmol). Diluted to 10⁷ copies/5 μ l (0.00332 fmol/ μ l).

[0252] Sample B: 20 ng/20 μ l stocks of each identifier were prepared. The sample was mixed as follows:

[0253] 5 μ l undil. Identifier #10

[0254] 5 μ l 2 \times dil. Identifier #9

[0255] 5 μ l 4 \times dil. Identifier #8

[0256] 5 μ l 8 \times dil. Identifier #7

[0257] 5 μ l 16 \times dil. Identifier #6

[0258] 5 μ l 32 \times dil. Identifier #5

[0259] 5 μ l 64 \times dil. Identifier #4

[0260] 5 μ l 128 \times dil. Identifier #3

[0261] 5 µl 256× dil. Identifier #2

[0262] 5 µl 512× dil. Identifier #1

[0263] Concentration: 10 ng/50 µl=0.20 ng/µl=1.923 fmol/µl. Diluted 579.2-fold to 10⁷ copies/5 µl (0.00332 fmol/µl).

[0264] Standard curve: The samples for the standard curve was prepared by diluting Sample A 116.55-fold to 10⁹ copies/5 µl (0.33 fmol/µl) and subsequently performing a 10-fold serial dilution of this sample. 5 µl was used for each PCR reaction. The standard curve is shown in FIG. 2.

[0265] Q-PCR Reactions

[0266] For 5 ml premix (for one 96-well plate):

[0267] 2.5 ml Taqman Universal PCR Master Mix (Applied Biosystems; includes Taq polymerase, dNTPs and optimized Taq pol. buffer)

[0268] 450 µl RPv2 (10 pmol/ul)

[0269] 25 µl Taqman probe (6-FAM-TCCAGCTTCTAG-GAAGAC-MGBNFQ; 50 µM; Applied Biosystems)

[0270] 1075 µl H₂O

[0271] 40.5 µl premix was aliquoted into each well and 4.5 µl of relevant upstream PCR primer (FPv2 (for standard curve) or one of the codon specific primers listed below; 10 pmol/µl) and 5 µl sample (H₂O in wells for negative controls) was added. The codon-specific PCR primers were: (Tm calculations shown are from Vector NTI; matched to Tm for RPv2 (67.7° C.))

P1-1:	GTCATACTAGCTGCTAGAGATGTGGTGATA	66.8° C.
P1-2:	CATACGGAAGAAGACAGAAGACCTGATA	67.8° C.
P1-3:	TCATACTCAGGAGTCGAGAAGCTGAAGATA	67.6° C.
P1-4:	CATACTGTGTACGTCAACACGTCAGATA	67.4° C.
P1-5:	CATACTGTGGAAC TACCATCCAAGGATA	68.0° C.
P1-6:	CCATCCAACATCGTTGGAAGAT	67.8° C.
P1-7:	CATACAACCTGTCTGTGAGATCTGATA	67.7° C.
P1-8:	ATACTCACGAAGCTGGATGATGAGATA	67.3° C.
P1-9:	CATACTAGCATCGATCGAACGTAGGATA	68.1° C.
P1-10:	TCATACTCGAAGCTACTGTGAGATGATA	68.2° C.
P2-1:	ATATTAGTGTGTGACGATGGTACGCA	67.8° C.
P3-1:	ACAAGTACGAACGTGCATCAGAGA	67.7° C.
P4-1:	CGAGCAGGACCTGGAACCT	67.7° C.
P4-2:	TCGACCAGCTCAGGTGGA	68.3° C.
P4-3:	GCTTCTCTGCTGCACCA	66.7° C.
P4-4:	GGTGTGAGGTGAGCAGCA	69.1° C.
P4-5:	CGACGAGGTCATCCTGGT	68.6° C.
P4-6:	GTGAGGAGCAGGTCCTCCTGT	68.0° C.
P4-7:	CTGACACTGGTCGTGGTCTGA	68.8° C.
P4-8:	CATCTCGACGACCTGCTCCT	67.9° C.

-continued

P4-9: ACGAGGTCTCCACTGGTCCA 68.3° C.

P4-10: ACTGAGCTGCTCCTCCAGGT 66.5° C.

[0272] Thermocycling/measurement of fluorescence was performed on an Applied Biosystems ABI Prism 7900HT real-time instrument utilizing the standard cycling parameters:

[0273] 95° C. 10 min;

[0274] 40 cycles of

[0275] 95° C. 15 sec;

[0276] 60° C. 1 min

[0277] All samples were run in duplicate.

[0278] Results

[0279] FIG. 6 shows the standard curve calculated by the 7900HT system software. The log of the starting copy number was plotted against the measured C_T value. The relationship between C_T and starting copy number was linear in the range from 10 to 10⁹ identifier copies.

[0280] This standard curve was utilized by the system software to calculate the quantity in the “unknown” samples as shown below.

TABLE I

Sample A (Shown graphically in FIG. 7)			
Sample A: Equimolar ratios	Observed A	Observed B	Expected
FPv2	12539947.00	11977503.00	10000000
P1-1	445841.90	480382.03	1000000
P1-2	884840.70	847478.56	1000000
P1-3	1013073.56	948770.00	1000000
P1-4	764187.94	741304.40	1000000
P1-5	1352874.60	1275155.50	1000000
P1-6	1284075.60	1337928.50	1000000
P1-7	658161.80	747371.56	1000000
P1-8	742187.20	653874.00	1000000
P1-9	824587.75	705785.75	1000000
P1-10	813550.75	836037.90	1000000
P2-1	13145159.00	14482606.00	10000000
P3-1	13263911.00	12773780.00	10000000
P4-1	1430704.80	1472576.80	1000000
P4-2	2681652.00	2481824.80	1000000
P4-3	1933106.80	2085476.40	1000000
P4-4	1359684.40	1364621.40	1000000
P4-5	2206709.80	2065813.60	1000000
P4-6	1652718.10	1873777.20	1000000
P4-7	1468208.10	1416153.00	1000000
P4-8	1664467.50	1581067.00	1000000
P4-9	1462520.60	1594593.80	1000000
P4-10	2020088.20	1912277.40	1000000

[0281]

TABLE II

Sample B (Shown graphically in FIG. 8)			
Sample B: 2-fold dil.	Observed A	Observed B	Expected
FPv2	4.97E+06	5.05E+06	10000000
P1-1	9955.07	10899.97	9765.625
P1-2	12732.32	13469.12	19531.25
P1-3	25542.8	25419.85	39062.5
P1-4	34748.89	44070.81	78125
P1-5	110881.41	123734.13	156250
P1-6	163687.44	166220.5	312500
P1-7	156993.81	172005.64	625000
P1-8	343176.78	374809.13	1250000
P1-9	646619.44	576151	2500000
P1-10	1.49E+06	1.72E+06	5000000
P2-1	5.19E+06	5.37E+06	10000000
P3-1	5.29E+06	5.09E+06	10000000
P4-1	(no signal)	70223.8	9765.625
P4-2	42103.32	22733.17	19531.25
P4-3	54480.62	39663.62	39062.5
P4-4	51293.07	43950.9	78125
P4-5	137946.95	115027.34	156250
P4-6	174134.64	156442.55	312500
P4-7	316505.78	283856.84	625000
P4-8	737661.44	691296.75	1250000
P4-9	1.42E+06	1.45E+06	2500000
P4-10	3.72E+06	3.52E+06	5000000

[0282] The results of the experiments show the possibility of accurately quantification of identifier oligonucleotides down to or even below 10 copies with a 9 fold dynamic range, and reliable relative quantification of the tested codons in various positions in the identifier oligonucleotide.

Example 5

Codon Analysis

[0283] Another possibility to analyse codons in identifier oligonucleotides is to use array format with attached probe oligonucleotides.

[0284] Six adaptors with the different anti-codon sequences in all three positions were designed. All the adaptors contain a probe binding sequence (20 nucleotides) that allows discrete binding on the microarray. Probe design is known in the art. Adaptors harbouring one to three deletions in the spacing region were used as negative controls to ensure that only the framing region is responsible for the hybridization of the identifier. Thus, the negative controls contain another framing sequence. The identifier oligonucleotide harbours the complementing codon sequence and the position directing framing regions.

Adaptor oligonucleotides

3' CTCATCGGAAGGGCTCGTAACGCTGGGTTTGGGGCGCTGGGTTTGGGGCGTGGGTTTGGGGCGG-5'

3' TTTGGTAGCTGAGTGCCCTAGGCTGGGTTTGGGGCGCTGGGTTTGGGGCGTGGGTTTGGGGCGG-5'

3' TAACTGGTTTACGCCACGCGCTGGGTTTGGGGCGCTGGGTTTGGGGCGTGGGTTTGGGGCGG-5'

3' TAATTGAGCTGACGGCGCACGGCTGGGTTTGGGGCGCTGGGTTTGGGGCGTGGGTTTGGGGCGG-5'

-continued

3' TGTTGCTACTCTGGCCGAGGCTGGGTTTGGGGCTGGGTTTGGGGCTGGGTTTGGGGCGG-5'

3' ACGGGATAACAACGCAGCCTGGCTGGGTTTGGGGCTGGGTTTGGGGCTGGGTTTGGGGCGG-5'


[0285] Identifier Oligonucleotide

Biotin-5' GCCACCCAAACCCCG

[0286] GenFlex hybridisation and scanning. Prior to hybridization, the Adaptor mix (100 pM final concentration for each of the adaptor oligonucleotides) in a hybridization buffer (100 mM MES, 1 M NaCl, 20 mM EDTA, 0.01% Tween 20, 1x Denhardt's), was heated to 95° C. for 5 min and subsequently cooled and maintained at 40° C. for 5 min before loading onto the Affymetrix GenFlex probe array cartridge. The probe array was then incubated for 2 h at 45° C. at constant rotation (60 rpm). The remaining Adaptor mix was removed from the GenFlex cartridge, and replaced with the identifier in a hybridization buffer (100 mM MES, 1 M NaCl, 20 mM EDTA, 0.01% Tween 20, 1x Denhardt's). The identifier hybridisation mix was heated to 95° C. for 5 min and subsequently cooled and maintained at 40° C. for 5 min before loading onto the Affymetrix GenFlex probe array cartridge and hybridised for 2 h at 45° C. at constant rotation (60 rpm). The washing and staining procedure was performed in the Affymetrix Fluidics Station. The probe array was exposed to 2 washes in 6xSSPE-T at 25° C. followed by 12 washes in 0.5xSSPE-T at 40° C. The biotinylated Identifier oligonucleotide was stained with a streptavidin-phycoerythrin conjugate, final concentration 2 µg/µl (Molecular Probes, Eugene, Oreg.) in 6xSSPE-T for 10 min at 25° C. followed by 6 washes in 6xSSPE-T at 25° C.

[0287] The probe arrays were scanned at 560 nm using a confocal laser-scanning microscope with an argon ion laser as the excitation source (Hewlett Packard GeneArray Scanner G2500A). The readings from the quantitative scanning were analysed by the Affymetrix Gene Expression Analysis Software. The results are depicted in Scheme 1.

Scheme 1:



[0288] The Array analysis shows that the codons including the framing regions are able to distinguish between the different probe oligonucleotides. The designed probes will only detect codons with the correct framing region allowing distinguishing first of the right codon and secondly as to which position the codon is positioned. Only one deletion in

both framing regions reduces significantly the hybridization of the identifier. Thus, the framing sequence may be used to obtain information about the position of a specific codon and the point in the reaction history when a given reaction of a chemical entity has occurred.

[0289] The information obtained in this example using either QPCR or array codon analysis as example can be used to generate a new more focused library. The signal from the QPCR analysis or the array analysis can directly be used to combine preferable chemical entities.


Example 6

Generation of a Second-Generation Library

[0290] The information obtained from a codon analysis performed according to the principles described in Examples 4 or 5 can be utilized for assembly a new more focused library. Sequence information can also be used to design a second-generation library with reduced diversity. This example illustrates how sequence data can be utilized to make a more focused library with the enriched chemical entities. Identical strategy can be based on the codon analysis methods described in Examples 4 or 5.

[0291] A 700-member library was generated composing of $4 \times 25 \times 7$ chemical entities. The library generation protocol is described below with the sequence information and chemical entity structure.



[0292] General arrangement of each complex composed of display molecule and identifier oligonucleotide in the library generation:



[0293] Specific codons in each oligo (Ax, Bx, Cx) was used and can be designed by using a specific nucleotide sequence for each chemical entity. In this particular setup, two complementary oligonucleotides (e.g. oligo Ax and oligo ax) containing a particular codon are allow to hybridize before the ligation step. The ligation of each codon oligonucleotide in each position is ligated with that attachment of the encoded chemical entity.


[0294] Overview of the Library Generation Procedure:

[0295] First Round of Library Generation (Round A):




[0296] "Pnt" corresponds to pentenoyl—an amine protecting group. "R" can be any molecule fragment. The chemical used in library generation comprise a primary (shown) or a secondary amine.


[0297] Second Round of Library Generation (Round B):



[0298] Third Round of Library Generation (Round C):



-continued



[0299] General Procedure: Library Generation, Selection and Mismatch Subsequent Selection

[0300] First Round of Library Generation (Round A):

[0301] First oligonucleotides of the A series are each modified by adding to each type of oligo a small molecule building block (BBAX) to the 5' amine forming an amide bond. After this step the identifier is comprised of oligo Ax.

[0302] Second Round of Library Generation (Round B):

[0303] 4 nmol of a mixture of different modified A oligos are then split into a number tubes corresponding to the number of different building blocks to be used in round B. 190 pmol Oligo a and 2 μ l heering DNA is added to each tube and the DNA material in each tube is lyophilized. The lyophilized DNA is then redissolved in 50 μ l water and purified by spinning through Biospin P-6 columns (Biorad) equilibrated with water.

[0304] Addition of Building Block

[0305] The DNA material in each tube is again lyophilized and redissolved in 2 μ l 100 mM Naborate pH 8.0/100 mM sulfo N-hydroxy succinimide (sNHS). For each tube 10 μ l building block BBBX (100 mM in dimethyl sulfoxide [DMSO]) is preactivated by mixing with 10 μ l 1-Ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC) (90 mM in dimethylformamide [DMF]) and incubating at 30 $^{\circ}$ C. for 30 min. 3 μ l of this preactivated mixture is then mixed with the 2 μ l in each tube and allowed to react 45 min at 30 $^{\circ}$ C. Then an additional 3 μ l freshly preactivated BB is added and the reaction is allowed to proceed for 45 min at 30 $^{\circ}$ C. The resulting mixture is then purified by spinning through Bio-Rad P6 DG (Desalting gel).

[0306] Addition of Codon Oligonucleotide

[0307] The DNA material is then lyophilized and redissolved in 10 μ l water containing 200 pmol oligo Bx (eg. B1) and the corresponding oligo bx (eg. b1). This is done so that the codon in oligo Bx identifies the BBBX added to the DNA identifier. 10 units of T4 DNA ligase (Promega) and 1.2 μ l T4 DNA ligase buffer is then added to each tube and the mixture is incubated at 20 $^{\circ}$ C. for 1 hour. The DNAn identifier linked to the small molecules now comprises an Ax oligo with a Bx oligo ligated to its 3' end. The reactions are then pooled, an appropriate volume of water is allowed to evaporate and the remaining sample is purified by spinning through Biospin P-6 columns (Biorad) equilibrated with water.

[0308] Removal of Building Block Protecting Group

[0309] The pooled sample (~50 μ l) is adjusted to 10 mM Na-acetate (pH 5). 0.25 volumes of 25 mM Iodine in tetrahydrofuran/water (1:1) is added and the sample is incubate at 37 $^{\circ}$ C. for 2 h. The reaction is then quenched by addition of 2 μ l of 1M Na₂S₂O₃ and incubation at room temperature for 5 min. The complexes are then purified by spinning through Biospin P-6 columns (Biorad) equilibrated with water

[0310] To remove sulphonamide protecting groups, the sample is adjusted to 50 μ l 100 mM sodium borate pH 8.5 and 20 μ l 1 500 mM 4-methoxy thiophenol (in acetonitrile) is added and the reaction is incubated at 25 $^{\circ}$ C. overnight. Then the complexes are purified by spinning through Biospin P-6 columns (Biorad) equilibrated with water and then lyophilized.

[0311] Third Round of Library Generation (Round C):

[0312] The samples are dissolved in 175 μ l 100 mM Na-borate pH 8.0 and distributed into 25 wells (7 μ l/well). 2 μ l 100 mM BB_cx in water/DMSO and 1 μ l of 250 mM DMT-MM is added to each reaction and incubated at 30 $^{\circ}$ C. overnight. Water is added to 50 μ l and the reactions are then spin purified using Bio-Rad P6 DG (Desalting gel) and subsequently water is allowed to evaporate so that the final volume is 10 μ l.

[0313] Addition of Building Block

[0314] The DNA material is then lyophilized and redissolved in 10 μ l water containing 200 pmol oligo Cx (eg. C1) and the corresponding oligo cx (eg. c1). This is done so that the codon in oligo Cx corresponds to the BB_cx added to the DNA identifier. 10 units of T4 DNA ligase (Promega) and 1.2 μ l T4 DNA ligase buffer is then added to each tube and incubated at 20 $^{\circ}$ C. for 1 hour. The DNAn identifier linked to the small molecules now comprises and Ax oligo with a Bx ligated to its 3' end and a Cx oligo ligated to the 3' end of the Bx oligo. The reactions are then pooled, the pooled sample volume is reduced by evaporation and the sample is purified by spinning through Biospin P-6 columns (Biorad) equilibrated with water. The pooled sample (~50 μ l) is adjusted to 10 mM Na-acetate (pH 5). 0.25 volumes of 25 mM Iodine in tetrahydrofuran/water (1:1) is added and the sample is incubate at 37 $^{\circ}$ C. for 2 h. The reaction is then quenched by addition of 2 μ l of 1 M Na₂S₂O₃ and incubation at RT for 5 min. Then the DNA identifiers (carrying small molecules) are purified by spinning through Biospin P-6 columns (Biorad) equilibrated with water and then lyophilized.

[0315] Final Deprotection Step

[0316] Some building blocks contain methyl esters that are deprotected to acids by dissolving the pooled sample in 5 μ l 20 mM NaOH, heating to 80 $^{\circ}$ C. for 10 minutes and adding 5 μ l of 20 mM HCl.

[0317] Final Extension Step

[0318] To ensure that the DNA identifiers are double stranded prior to selection oligo d is extended along the identifier by adding to the sample 10 μ l of 5 \times sequenase EX-buffer [100 mM Hepes, pH 7.5, 50 mM MgCl₂, 750 mM NaCl] and 4000 pmol oligo d. Annealing is performed by heating to 80 $^{\circ}$ C. and cooling to 20 $^{\circ}$ C. To the sample is then

added 500 μ L dNTP, water to 50 μ L and 39 units of Sequenase version 2.0 (USB). The reaction is incubated at 37° C. for 1 hour.

[0319] Selection

[0320] This library is subjected to selection, whereby binders to the selection target are enriched.

[0321] Maxisorp ELISA wells (NUNC A/S, Denmark) were coated with each 100 μ L 2 μ g/mL integrin α V β 3 in PBS buffer [2.8 mM NaH₂PO₄, 7.2 mM Na₂HPO₄, 0.15 M NaCl, pH 7.2] overnight at 4° C. Then the integrin solution was substituted for 200 μ L blocking buffer [TBS, 0.05% Tween 20 (Sigma P-9416), 1% bovine serum albumin (Sigma A-7030), 1 mM MnCl₂] which was left on for 3 hours at room temperature. Then the wells were washed 10 times with blocking buffer and the encoded library was added to the wells after diluting it 100 times with blocking buffer. Following 2 hours incubation at room temperature the wells were washed 10 times with blocking buffer. After the final wash the wells were cleared of wash buffer and subsequently inverted and exposed to UV light at 300-350 nm for 30 seconds using a trans-illuminator set at 70% power. Then 100 μ L blocking buffer without Tween-20 was immediately added to each well, the wells were shaken for 30 seconds, and the solutions containing eluted identifiers were removed for PCR amplification.





[0322] Cloning

[0323] A TOPO-TA (Invitrogen) ligation reaction is assembled with 4 μ L PCR product, 1 μ L salt solution (Invitrogen) and 1 μ L vector. Water is added to 6 μ L. The reaction is then incubated at RT for 30 min. Heat-shock competent TOP10 *E. coli* cells are then thawed on ice and 5 μ L of the ligation reaction is added to the thawed cells. The cells are then incubated 30 min on ice, heatshocked in 42° C. water for 30 sec, and then put on ice again. 250 μ L of growth medium is added to the cells and they are incubated 1 h at 37° C. The medium containing cells is then spread on a growth plate containing 100 μ g/ml ampicillin and incubated at 37° C. for 16 hours.




[0324] Sequencing

[0325] Individual *E. coli* clones are then picked and transferred to PCR wells containing 50 μ L water. These 50 μ L were incubated at 94° C. for 5 minutes and used in a 20 μ L in a 25 μ L PCR reaction with 5 pmol of each TOPO primer M13 forward & M13 reverse and Ready-To-Go PCR beads (Amersham Biosciences). The following PCR profile is used: 94° C. 2 min, then 30 \times (94° C. 4 sec, 50° C. 30 sec, 72° C. 1 min) then 72° C. 10 min. Primers and nucleotides are then degraded by adding 1 μ L 1:1 EXO/SAP mixture (USB corp.) to 2 μ L PCR product and incubating at 37° C. for 15 min and then 80° C. for 15 min to heat-inactivate the enzymes. 5 pmol T7 primer is added and water is added to 12 μ L. Then 8 μ L DYEnamic ET cycle sequencing Terminator Mix (Applied biosystems) is added to each well. A thermocycling profile of 30 \times (95° C. 20 sec, 50° C. 15 sec, 60° C. 1 min) is then run. Then 10 μ L water is added to each well and sequencing reactions are purified using seq96 spinplates (Amersham Biosciences). Reactions are then run on a MegaBace capillary electrophoresis instrument (Molecular Dynamics) using injection parameters 2 kV, 60 sec and run parameters: 9 kV 45 min and analyzed using Contig Express software (Informax).

[0326] The chemical entities used in each position are shown below.

Position 1	
Building Block	Smiles
BB-A-000098	 <p>Structure 9</p>
BB-A-000112	 <p>Structure 123</p>
BB-A-000282	 <p>Structure 81</p>
BB-A-000283	 <p>Structure 82</p>


[0327]

Position 2	
BB-A-0099182	 <p>Structure 131</p>
BB-A-0000613	 <p>Structure 36</p>
BB-A-0000084	 <p>Structure 33</p>

-continued


Position 2

BB-A-0000832




Structure 14

BB-A-909683




Structure 70

BB-A-0011001




Structure 137

BB-A-0001183




Structure 137

BB-A-0007821




Structure 160

BB-A-0001182



Structure 114

BB-A-0001182




Structure 1H

-continued


Position 2

BB-A-0001614




Structure 295

BB-A-0001642




Structure 300

BB-A-0002382




Structure 166

BB-A-0063143




Structure 168

BB-A-0982162




Structure 168

BB-A-0002182




Structure 153

BB-A-0083172




Structure 453

-continued



[0328]



[0329] After the selection as described above, the codons in the identifier oligonucleotides were analysed. Before the analysis, the identifier oligonucleotides were amplified using the constant flanking regions and the amplified material was used in the identifier sequence analysis.

[0330] A sequence codon analysis of the selected codons showed a bias for specific chemical entities. They are listed in the table below. For instance, in position 1 chemical entity 98 was seen 47 times (out of 51 sequences, 92%, compare to 25% before the selection) and chemical entity 99 was seen 14 times (out of 51 sequences, 27%, compare to 4% before

selection) and chemical entity 53 was seen 35 times (out of 51 sequences, 68%, compare to 14% before selection).

[0331] The chemical entities listed in the table below can then be used to generate a new and more focused library.

Oligo(-s)	Count	pos 1	pos 2	pos 3
BB-A-000098	47	98		
BB-A-000282	4	282		
Ⓢ	Ⓢ		Ⓢ	
Ⓢ	Ⓢ		Ⓢ	
BBA0004242	6		424	
BBA0004182	5		418	
BBA0001101	2			
BBA0003172	2			
BBA0004212	2			
BBA0004232	2			
BBA000064	1			
BBA0001011	1			
BBA0003132	1			
BBA0003142	1			
BBA0003152	1			
Ⓢ	Ⓢ			Ⓢ
Ⓢ	Ⓢ			Ⓢ
BBA0001006	4			100
BBA0008512	2			
BBA0008312	1			

Ⓢ indicates text missing or illegible when filed

[0332] The new focused library with the selected chemical entities can be selected against the target and the outcome from the selection can be analysed. The most abundant binders will be the combination between the chemical entities 98-99-53 and the second most abundant binder is 98-158-53 as shown below.

Oligo(-s)	Count	pos 1	pos 2	pos 3
BB-A-000098 BBA000099 BBA0000531	11	98	Ⓢ	53
BB-A-000098 BBA0001582 BBA0000531	7	98	158	53
BB-A-000098 BBA0004242 BBA0000531	4	98	424	53
BB-A-000098 BBA0001582 BBA0001391	3	98	158	139
BB-A-000098 BBA0004182 BBA0000531	3	98	418	53
BB-A-000098 BBA000099 BBA0001391	2	98	Ⓢ	139
BB-A-000098 BBA0001582 BBA0001006	2	98	158	100

Ⓢ indicates text missing or illegible when filed

[0333] This example exemplifies the possibility to reduce the library diversity by using the enriched chemical entities in a new library and perform another round of selection on the chosen chemical entities.

Example 7

[0334] The following experiment illustrates the principle of chemical entity (also termed building block herein) evolution through multiple rounds of library generation and selection. The experiment is not intended to limit the scope of the current invention.

[0335] Libraries were assembled by the combination of building blocks (BB) each of which was encoded by an oligonucleotide (oligo). Some of the building blocks carried an amine functional group and a carboxylic acid functional group. The building block amine was protected by N-pentenoylation and deprotected by iodine treatment prior to the

reaction of the following building block. Oligonucleotide 1 (Oligo1) carried an amine functional group to allow reaction with the building block 1's carboxylic acid and oligonucleotides are optionally derivatized by phosphorylation to allow ligation. Oligonucleotide 3 (oligo3) also comprised a primer region for PCR amplification. EDC/NHS, EDC/sulfoNHS or DMTMM was used as coupling reagents.

[0336] The following scheme describes the split and mix assembly of the libraries:

[0337] i.) n times [BB1+Oligo1→BB1-Oligo1] in separate wells

[0338] * Optionally purify product

[0339] ii.) mix all n wells into one tube

[0340] iii.) split product of ii.) into m separate wells

[0341] iv.) m times [BB2+BB1-Oligo1+Oligo2→BB2-BB1-Oligo1-Oligo2] in separate wells

[0342] * Optionally purify product

[0343] v.) mix all m wells into one tube

[0344] vi.) split product of v.) into p separate wells

[0345] vii.) p times [BB3+BB2-BB1-Oligo1-Oligo2+Oligo3→BB3-BB2-BB1-Oligo1-Oligo2-Oligo3] in separate wells

[0346] * Optionally purify product

[0347] viii.) mix all p wells into one tube

[0348] ix.) Selection was performed and binders isolated


[0349] x.) PCR of DNA and sequencing

[0350] xi.) Analyse for building block abundancy and full sequence information



[0351] Building block abundances analysis may be done by QPCR or by sequencing full sequences and then analyzing for the abundance of individual building blocks.

[0352] The following types of building blocks were used, wherein R describes a group which is varied for different building blocks:


Building block types used in position 1, 2 and 3




Both (R) and (S) isomers were for some R-groups used



Both (R) and (S) isomers were for some R-groups used



[0353] The overall process leads to molecules of the following structure, where the oligonucleotide was double stranded.



[0354] The oligonucleotide was made double stranded by the use of double stranded Oligo's 1, 2 and 3 with an overhang to allow ligation of both strands.

[0355] Summary of the experimental outcome:

[0356] Two libraries of 61,875 members (Library 1 and 2) were generated as described in example 6 above and selected for binders of the Integrin $\alpha\beta 3$ receptor separately. The libraries were generated with 99 different building blocks in position 1, 25 different building blocks in position 2 and 25 different building blocks in position 3.

[0357] The identified sequences were then analyzed for the abundances of building blocks at each position in the sequence. The most abundant building blocks at each position from the two libraries 1 and 2 were then used again to generate a new and smaller library of 1,365 members, which was selected for binders of the Integrin $\alpha\beta 3$ receptor. The library was generated with 7 different building blocks in position 1, 13 different building blocks in position 2 and 15 different building blocks in position 3.

[0358] In the tables below, each of the building block numbers identify one specific building block or in two instances (library 1) a mixture of three different building blocks. The same numbers are used for each building block in all libraries, however the oligonucleotide used to identify each building block may not necessarily be the same between libraries to avoid potential problems of cross contamination.

[0359] The following tables describes the codon sequences and corresponding building blocks used. The codon is only indicated for one of the strands.

<u>Library 1, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
1	TGTC	BBA000092
2	CGAGC	BBA000354
3	GGATA	BBA000085
4	CGCTG	BBA000086
5	GTTAT	BBA000098

-continued

<u>Library 1, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
6	AGTGC	BBA000099
7	ACCTG	BBA000089
8	CTGGT	BBA000090
9	TAGGA	BBA000087
10	ACTCA	BBA000088
11	CTTAC	BBA000153
12	CGCAC	BBA000154
13	TCGCG	BBA000059
14	CGGAT	BBA000152
15	GAGAT	BBA000101
16	TGTAG	BBA000110
17	GTGTT	BBA000112
18	AGATG	BBA000113
19	ATCCT	BBA000114
20	TTGCT	BBA000286

-continued			-continued		
<u>Library 1, Position 1</u>			<u>Library 1, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID	Codon no.	Codon sequence ID	Building Block ID
21	ACGTA	BBA000123	56	TCTGC	BBA000196
22	ATCAC	BBA000124	57	ATAGG	BBA000197
23	TATCC	BBA000155	58	CTACC	BBA000198
24	GGAAG	BBA000156	59	AAGTG	BBA000201
25	CGGTC	BBA000158	60	TCCAA	BBA000202
26	TGCTT	BBA000159	61	GCTCT	BBA000203
27	TTAGC	BBA000160	62	GGAGT	BBA000204
28	GCTGA	BBA000161	63	AATCG	BBA000205
29	GAACG	BBA000162	64	AAGCT	BBA000206
30	CATGG	BBA000163	65	CCGAA	BBA000207
31	TGGTA	BBA000165	66	TTTGT	BBA000208
32	TCAAG	BBA000166	67	CCGTG	BBA000209
33	ATCGA	BBA000167	68	TTTCG	BBA000210
34	ATGCA	BBA000168	69	TGAGG	BBA000211
35	ACTAG	BBA000169	70	GTTGC	BBA000212
36	TACCT	BBA000170	71	AACTA	BBA000112
37	TACGA	BBA000171	72	AACTA	BBA000280
38	CTTCA	BBA000172	73	CCTCG	BBA000281
39	CTCTT	BBA000173	74	AGCAA	BBA000282
40	TCATC	BBA000174	75	TTCCA	BBA000313
41	ATTCC	BBA000175	76	AGACT	BBA000314
42	CGACG	BBA000176	77	AGGTT	BBA000315
43	CCTGT	BBA000177	78	GCGTC	BBA000316
44	CCTTC	BBA000178	79	AACGT	BBA000317
45	ACACC	BBA000179	80	CAAGA	BBA000287
46	TAACA	BBA000180	81	AGAGA	BBA000419
47	TAACA	BBA000098	82	GTA CT	BBA000420
48	CCAGG	BBA000181	83	TAGAG	BBA000421
49	ATGTC	BBA000182	84	ACGAT	BBA000422
50	GAGGA	BBA000183	85	GACCA	BBA000200
51	GGTCA	BBA000184	86	TCGTT	BBA000194
52	GACTT	BBA000185	87	GTCTC	B8A000427
53	GGTGG	BBA000186	88	CAGCA	BBA000428
54	CAACT	BBA000190	89	TAGTC	BBA000199
55	ATGAG	BBA000195	90	GGGTG	BBA000187

-continued

<u>Library 1, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
91	CTCAG	BBA000191
92	AGAAC	BBA000284
93	GCGAG	BBA000458
94	GATGT	BBA000459
95	TCACT	BBA000461
96	CGTCT	OBA000610
97	AGCTC	OBA000611
98	CACTC	OBA000609
99	CAGTT	OBA000615

[0360]

<u>Library 1, Position 2</u>		
Codon no.	Codon sequence ID	Building Block ID
1	AGTACGAACGTGCATCAGAG	BBA000098
2	TAGTCTCCTCCACTTCCATG	BBA000099
3	TACATCGTTCCAGACTACCG	BBA000085
4	TCCAGTGCAAGACTGAACAG	BBA000153
5	AGCATCACTACTCTGTCTGG	BBA000206
6	TCTTGTCAACCTTCCATGCG	BBA000200
7	AAGGACGTTCTAGTAGGTG	BBA000208
8	GGAACCATCAAGATCCTGAG	BBA000091
9	ATCTCTGACGAGATCCAAG	BBA000090
10	TCAAGGTGGTGGTGTACTG	BBA000092
11	TCGAAC TTGTTGCTTCCTCG	BBA000123
12	CTGAGTGTGTAGTACCAACG	BBA000156
13	ATCTTGTTGTTCTCCTGCG	BBA000163
14	TAGTAGCTTGGAGTAGACCG	BBA000197
15	TTCACTCCATGCAGCATGTG	BBA000083
16	ACGATGGTGATCGATCAACG	BBA000181
17	TTCAGTGCTTGAGCTACCTG	BBA000152
18	TTGACTCTTCTTGACCAG	BBA000088
19	TCAACCAACTGGTCTTTGGG	BBA000100

-continued

<u>Library 1, Position 2</u>		
Codon no.	Codon sequence ID	Building Block ID
20	TAGTACTCTACTGCTGCG	BBA00087 101 196
21	TACACCATGACTTGCAGACG	BBA00087 101 196
22	GCATCTTGAGTCGTTGAACG	BBA000059
23	GACTCATCTCACTGGAGTTG	BBA000124
24	TCCAGCTTCTAGGAAGACAG	BBA000160
25	CTTCTTGAGTGCCTAGCAG	BBA000201

[0361]

<u>Library 1, Position 3</u>		
Codon no.	Codon sequence ID	Building Block ID
1	CGAGCAGGACCTGGAACCTGGTGC	BBA000098
2	CTCGACCACTGCAGGTGGAGCTCC	BBA000099
3	CGTGCTTCTCTGCTGCACCACCG	BBA000085
4	CCTGGTGTGCGAGGTGAGCAGCAGC	BBA000153
5	CTCGACGAGGTCCATCCTGGTFCG	BBA000206
6	CGTGAGGAGCAGGTCCCTCCTGTCG	BBA000200
7	CCTGACACTGGTCTGTTGTCGAGGC	BBA000208
8	CCATCTCGACGACCTGTCTCTGGG	BBA000091
9	CCACGAGGTCTCCACTGGTCCAGG	BBA000090
10	CCACTGAGCTGTCTCCAGGTGG	BBA000092
11	CCTCCTGTCTGCAGTCCATCCG	BBA000123
12	CAGCACCTGGAGGTAGGACCACGG	BBA000156
13	CGACCAGACGAGGACCAGGTAGGC	BBA000163
14	CCAGGTTGAGGACCTCGTCAGCC	BBA000197
15	CGAGCACGAGGAGCACGTGTCCAG	BBA000100
16	CCACGTCCACAGGTGCACCAGGTG	BBA000181
17	CCTGGTGTCCACGACGTGCTTCG	BBA000152
18	CACGTGACGACCTGGTCAGGTGGG	BBA000088
19	CGTAGCTCGTGTGGTCTCTCTGG	BBA000101
20	CGACGACCACCTTGGACACCC	BBA000196
21	CCTACGTCGTGCTCACGTCTGCC	BBA000087
22	CGACGACGCTAGGAGGAGGTGGG	BBA000083
23	CTGGTGGAGCTGCACGACACAGC	BBA000059

-continued

<u>Library 1, Position 3</u>		
Codon no.	Codon sequence ID	Building Block ID
24	CAGGACTGGACGACGACCAGGTCG	BBA000124
25	CGATGCTGCAGACGACCAGCACCC	BBA000160

[0362]

<u>Library 2, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
1	TGTTC	BBA000092
2	CGAGC	BBA000354
3	GGATA	BBA000085
4	CGCTG	BBA000086
5	GTTAT	BBA000098
6	AGTGC	BBA000099
7	ACCTG	BBA000089
8	CTGGT	BBA000090
9	TAGGA	BBA000087
10	ACTCA	BBA000088
11	CTTAC	BBA000153
12	CGCAC	BBA000154
13	TCGCG	BBA000059
14	CGGAT	BBA000152
15	GAGAT	BBA000101
16	TGTAG	BBA000110
17	GTGTT	BBA000112
18	AGATG	BBA000113
19	ATCCT	BBA000114
20	TTGCT	BBA000286
21	ACGTA	BBA000123
22	ATCAC	BBA000124
23	TATCC	BBA000155
24	GGAAG	BBA000156
25	CGGTC	BBA000158
26	TGCTT	BBA000159

-continued

<u>Library 2, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
27	TTAGC	BBA000160
28	GCTGA	BBA000161
29	GAACG	BBA000162
30	CATGG	BBA000163
31	TGGTA	BBA000165
32	TCAAG	BBA000166
33	ATCGA	BBA000167
34	ATGCA	BBA000168
35	ACTAG	BBA000169
36	TACCT	BBA000170
37	TACGA	BBA000171
38	CTTCA	BBA000172
39	CTCTT	BBA000173
40	TCATC	BBA000174
41	ATTCC	BBA000175
42	CGACG	BBA000176
43	CCTGT	BBA000177
44	CCTTC	BBA000178
45	ACACC	BBA000179
46	TAACA	BBA000180
47	TAACA	BBA000098
48	CCAGG	BBA000181
49	ATGTC	BBA000182
50	GAGGA	BBA000183
51	GGTCA	BBA000184
52	GACTT	BBA000185
53	GGTGG	BBA000186
54	CAACT	BBA000190
55	ATGAG	BBA000195
56	TCTGC	BBA000196
57	ATAGG	BBA000197
58	CTACC	BBA000198
59	AAGTG	BBA000201
60	TCCAA	BBA000202
61	GCTCT	BBA000203

-continued

<u>Library 2, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
62	GGAGT	BBA000204
63	AATCG	BBA000205
64	AAGCT	BBA000206
65	CCGAA	BBA000207
66	TTTGT	BBA000208
67	CCGTG	BBA000209
68	TTTCG	BBA000210
69	TGAGG	BBA000211
70	GTTGC	BBA000212
71	AACTA	BBA000112
72	AACTA	BBA000280
73	CCTCG	BBA000281
74	AGCAA	BBA000282
75	TTCCA	BBA000313
76	AGACT	BBA000314
77	AGGTT	BBA000315
78	GCGTC	BBA000316
79	AACGT	BBA000317
80	CAAGA	BBA000287
81	AGAGA	BBA000419
82	GTACT	BBA000420
83	TAGAG	BBA000421
84	ACGAT	BBA000422
85	GACCA	BBA000200
86	TCGTT	BBA000194
87	GTCTC	BBA000427
88	CAGCA	BBA000428
89	TAGTC	BBA000199
90	GGGTG	BBA000187
91	CTCAG	BBA000191
92	AGAAC	BBA000284
93	GCGAG	BBA000458
94	GATGT	BBA000459
95	TCACT	BBA000461
96	CGTCT	OBA000610

-continued

<u>Library 2, Position 1</u>		
Codon no.	Codon sequence ID	Building Block ID
97	AGCTC	OBA000611
98	CACTC	OBA000609
99	CAGTT	OBA000615

[0363]

<u>Library 2, Position 2</u>		
Codon no.	Codon sequence ID	Building Block ID
1	AGTACGAACGTGCATCAGAG	BBA000059
2	TAGTCTCCTCCACTTCCATG	BBA000085
3	TACATCGTTCCAGACTACCG	BBA000098
4	TCCAGTGCAAGACTGAACAG	BBA000099
5	AGCATCACTACTCTGTCTGG	BBA000101
6	TCTTGTC AACCTTCCATGCG	BBA000110
7	AAGGACGTFCC TAGTAGGTG	BBA000113
8	GGAACCATCAAGATCCTGAG	BBA000114
9	ATCTCTGACGAGATCCAAG	BBA000123
10	TCAAGGTTGGTGGTACTG	BBA000124
11	TCGAACTTGTGTCTCCTCG	BBA000152
12	CTGAGTGTGTAGTACCAACG	BBA000158
13	ATCTTGGTGTCTCCTGCG	BBA000160
14	TAGTAGCTTGGAGTAGACCG	BBA000161
15	TTCATCCATGCAGCATGTG	BBA000167
16	ACGATGGTGATCGATCAACG	BBA000176
17	TTCAGTGCTTGAGCTACCTG	BBA000181
18	TTGACTCTTCTTGACCAG	BBA000313
19	TCAACCACTGGTTCTTGGG	BBA000314
20	TAGTACTCTACACTGCTGCG	BBA000315
21	TACACCATGACTTGCAGACG	BBA000316
22	GCATCTTGAGTCGTTGAACG	BBA000317
23	GACTCATCTCACTGGAGTTG	BBA000420
24	TCCAGCTTCTAGGAAGACAG	BBA000421
25	CTTCTTGAGTGCAC TAGCAG	BBA000422

[0364]

<u>Library 2, Position 3</u>		
Codon no.	Codon sequence ID	Building Block ID
1	CGAGCAGGACCTGGAACCTGGTGC	BBA000052
2	CTCGACCACTGCAGGTGGAGCTCC	BBA000053
3	CGTGCTTCCTCTGCTGCACCACCG	BBA000054
4	CCTGGTGTGCGAGGTGAGCAGCAGC	BBA000056
5	CTCGACGAGGTCCATCCTGGTCGC	BBA000057
6	CGTGAGGAGCAGGTCTCCTGTGCG	BBA000058
7	CCTGACACTGGTCGTGGTCGAGGC	BBA000062
8	CCATCTCGACGACCTGCTCCTGGG	BBA000139
9	CCACGAGGTCTCCACTGGTCCAGG	BBA000140
10	CCACTGAGCTGTCCCTCCAGGTGG	BBA000100
11	CCTCCTGTCTGCACGTCCATCCG	BBA000059
12	CAGCACCTGGAGGTAGGACCACGG	BBA000085
13	CGACCAGACGAGGACCAGGTAGGC	BBA000098
14	CCAGGTTTCGAGGACCTCGTCAGCC	BBA000099
15	CGAGCACGAGGAGCAGCTGTCCAG	BBA000101
16	CCACGTCCACAGGTGCACCAGGTG	BBA000110
17	CCTGGTGTCTCCACGACGTGCTTCG	BBA000113
18	CACGTGACGACCTGGTCAGGTGGG	BBA000114
19	CGTAGCTCGTGTGGTCTCCTGG	BBA000123
20	CGACGACCACCCTTGGACACCC	BBA000124

-continued

<u>Library 2, Position 3</u>		
Codon no.	Codon sequence ID	Building Block ID
21	CCTACGTCGTGCTCACGTCCTGCC	BBA000152
22	CGACGACAGCTAGGAGGAGGTGGG	BBA000158
23	CTGGTGGAGCTGCACGACACAGC	BBA000160
24	CAGGACTGGACGACGACCAGGTCTG	BBA000161
25	CGATGCTGCAGACGACCAGCACCC	BBA000167

[0365]

<u>Library 3, Position 1</u>			
Codon no.	Codon sequence ID	Building Block ID	More abundant in position 1 in library no.
1	TGTTC	BBA000092	1
2	ACTCA	BBA000088	1
3	CTTAC	BBA000153	1 and 2
4	CGGAT	BBA000152	1
5	ATTCC	BBA000175	1 and 2
6	GTCTC	BBA000427	1
7	ACAGT	BBA000098	1 and 2

[0366]

<u>Library 3, Position 2</u>			
Codon no.	Codon sequence ID	Building Block ID	More abundant in position 2 in library no.
1	6CACAAAGTACGAACGTGCATCAGAG	BBA000059	1
2	6CACATAGTCTCCTCCACTTCCATG	BBA000083	1
3	6CACATACATCGTTCAGACTACCG	BBA000085	2
4	6CACATCCAGTGCAAGACTGAACAG	BBA000088	1
5	6CACAAGCATCACTACTCTGTCTGG	BBA000090	1
6	6CACATCTTGTCAACCTTCCATGCG	BBA000099	1 and 2
7	6CACAAAGGACGTTCTAGTAGGTG	BBA000110	
8	6CACAGGAACCATCAAGATCCTGAG	BBA000114	2
9	6CACAATCTCTGACGAGATCCAAGG	BBA000152	2

-continued

<u>Library 3, Position 2</u>			
Codon no.	Codon sequence ID	Building Block ID	More abundant in position 2 in library no.
10	6CACATCAAGGTTGGTGGTGTACTG	BBA000160	2
11	6CACATCGAACTTGTGCTTCCTCG	BBA000200	1
12	6CACACTGAGTGTAGTACCAACG	BBA000201	1
13	6CACAATCTTGGTTGTTCTCCTGCG	BBA000422	2

[0367]

<u>Library 3, Position 3</u>			
Codon no.	Codon sequence ID	Building Block ID	More abundant in position 3 in library no.
1	6AGGACGAGCAGGACCTGGAACCTGGTGCCTCCACCACGTCTCCG	BBA000053	2
2	6AGGACTCGACCACTGCAGGTGGAGTCCGTTCCACCACGTCTCCG	BBA000085	1
3	6AGGACGTGCTTCTCTGCTGCACCACCGTTCCACCACGTCTCCG	BBA000087	1
4	6AGGACCTGGTGTGAGGTGAGCAGCAGCGTTCCACCACGTCTCCG	BBA000090	1
5	6AGGACTCGACGAGGTCCATCCTGGTGCCTCCACCACGTCTCCG	BBA000091	1
6	6AGGACGTGAGGAGCAGGTCCCTCTGTCGGTTCCACCACGTCTCCG	BBA000098	1
7	6AGGACCTGACACTGGTGCCTGAGGCGTTCCACCACGTCTCCG	BBA000100	1 and 2
8	6AGGACCATCTCGACGACCTGCTCCTGGGGTTCCACCACGTCTCCG	BBA000139	2
9	6AGGACCACGAGGTCTCCACTGGTCCAGGGTTCCACCACGTCTCCG	BBA000140	2
10	6AGGACCACCTGAGTGTCTCCAGGTGGTTCCACCACGTCTCCG	BBA000152	
11	6AGGACCTCCTGTCTGCACGTCCATCCGGTTCCACCACGTCTCCG	BBA000153	1
12	6AGGACAGCACCTGGAGGTAGGACCAGGGTTCCACCACGTCTCCG	BBA000161	
13	6AGGACGACCAGCAGGACCAGGTAGGCGTTCCACCACGTCTCCG	BBA000167	2
14	6AGGACCAGGTTGAGGACCTCGTCAGCCGTTCCACCACGTCTCCG	BBA000197	1
15	6AGGACGAGCAGGAGCAGTGTCCAGTTCCACCACGTCTCCG	BBA000200	1

[0368] A subset of the isolated sequences from the library post selection was analysed:

- (1) GGCAGCACAGTCGTCGCACATACATCGTTCCAGACTACCGAGGACCTGACACTGGTCTGTCGAGGCGTTTCCT
- (2) GGCAGCACAGTCGTCGCTACATGCTTGTCAACCTTCCATGCGAGTACCTTACACTGGTCTGTCGAGGCGTTTCCT
- (3) GGCAGCCGGAT423CGTCGCACATCTTGTCAACCTTCCATGCGAGGACCTGACACTGGTCTGTCGAGGCGTTTCCT
- (4) GGCAGCCTTACGTCGCACAATTCTCTGACAGAAATCCAACGGAGG

-continued

- ACCTGACACGTGCGTCTGGCTCGATGCGTTTCCTC
- (5) GGCAGCACAGTCGTCGCACATCATTGTCAAAACCTTCCATGCGAGGACCATCTCGACGACCTGCTCCTGGGGTTCCTC
 - (6) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGACCATCTCGACGACCTGCTCCTGGGGTTCCTC
 - (7) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGACCATCTCGACGACCTGCTCCTGGGGTTCCTC
 - (8) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGACCATCTCGACGACCTGCTCCTGGGGTTCCTC

-continued

- (9) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
CATCTCGACGACCTGCTCCTGGGGTTCCTC
- (10) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
CATCTCGACGACCTGCTCCTGGGGTTCCTC
- (11) GGCAGCACTAGATCGTCGCACATCTTGTCAACCTTCCATGCGAGG
ACCATCTTCGACTGANCTGCCTCCTGTGGGGTTCCTC
- (12) GGCAGCACAGATCGTCGCACATCTTGTCAACCTTCCATGCGAGGA
CCATCTCGACGANCTGCTCCTGGGGTTCCTC
- (13) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
CATCACGACTACCTTGGCTCCCTGGGGTTCCTC
- (14) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGACC
ATCTCGACGACCTGCTCCTGGGGCCCTC
- (15) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
CATCTCGACGACCTGCTCCTGGGGTTCCTC
- (16) GGCAGCCGGATCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
CATCTCGACGACCTGCTCCTGGGGTTCCTC
- (17) GGCAGCCGGATCGTCGCACATCTTGTCAACCTTCCATGCGAGGACC
ATCTCGACGACCTGCTCCTGGGGTTCCTC
- (18) GGCAGCCGGATCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
CATCTCGACGACCTGCTCCTGGGGTTCCTC
- (19) GGCAGCACAGTCGTCGCAATCCAGTCAAGACTGAACAGAGGACCA
TCTCGACGACCTGCTCCTGGGGTTCCTC
- (20) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTTCCATGCGAGG
ACGAGCAGGACCTGGAACCTGGTGCCTTCCTC
- (21) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGACC
AGCAGGACCTGGAACCTGGTGCCTTCCTC
- (22) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
GATGACGAGACCTGGAACCTGGTGCCTTCCTC
- (23) GGCAGCCGGATCGTCGCAGATCTTGGTNAANCTTCCATGCGAGGA
CGAGCATGAACCTGGAACCTGGTGCCTTCCTC
- (24) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGACC
AGGTCCTCACTGGTCCAGGGTTCCTC
- (25) GGCAGCACAGTCGTCGGCACATCTTGTCAACCTTCCATGCGAG
GACCAGGAGTCTCCACTGGTCCAGGGTTCCTC
- (26) GGCAGCCGGATCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
GACCAAGACGAGGACAGGTAGGCGTTCCTC
- (27) GGCAGCCGGAT423CGTCGCACATCTTGTCAACCTTCCATGCGAG
GACGTGATGGAGCAAGTCTCCTGTGCGTTCCTC
- (28) GGCAGCACAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGAC
ACGAGGTCTCCACTGGTCCAGGTTCCTC
- (29) GCCCAAACAAGTCGTCGCACATCTTGTCAACCTTCCATGCGAGGA
CCGAGNNGTAGCTGGANNCTCGGATGCGTTCCTC
- (30) GCAGCACAGATCGTCGCACATGCTTGTCAAGCCTTTCCATCGGGA
GGACCATCTACGAGCGAGCACTTGTGCTGCGTGGGGTTCCTC
- (31) GGCAGCCGGATCGTCGCACATCAATGGTTTGGTGGTGATACTGA
GGACCAGAGCTTACACTTGGTTCAGGGTTCCTC


[0369] These sequences could be translated into the following building block compositions:

Sequence no.	Position 1	Position 2	Position 3
1	BBA000098	BBA000085	BBA000100
2	BBA000098	BBA000099	BBA000100
3	BBA000152	BBA000099	BBA000100
4	BBA000153	BBA000152	BBA000100
5	BBA000098	BBA000099	BBA000139
6	BBA000098	BBA000099	BBA000139
7	BBA000098	BBA000099	BBA000139
8	BBA000098	BBA000099	BBA000139
9	BBA000098	BBA000099	BBA000139
10	BBA000098	BBA000099	BBA000139
11	BBA000098	BBA000099	BBA000139
12	BBA000098	BBA000099	BBA000139
13	BBA000098	BBA000099	BBA000139
14	BBA000098	BBA000099	BBA000139
15	BBA000098	BBA000099	BBA000139
16	BBA000152	BBA000099	BBA000139
17	BBA000152	BBA000099	BBA000139
18	BBA000152	BBA000099	BBA000139
19	BBA000098	BBA000088	BBA000139
20	BBA000098	BBA000099	BBA000053
21	BBA000098	BBA000099	BBA000053
22	BBA000098	BBA000099	BBA000053
23	BBA000152	BBA000099	BBA000053
24	BBA000152	BBA000099	BBA000140
25	BBA000098	BBA000099	BBA000140
26	BBA000152	BBA000099	BBA000167
27	BBA000152	BBA000099	BBA000098
28	BBA000098	BBA000099	BBA000200
29	BBA000098	BBA000099	—
30	BBA000098	BBA000099	—
31	BBA000152	BBA000160	—


[0370] In position 1 L-Asp (BBA00098) dominated. D-Asp was also found (BBA000152)

[0371] In position 2 Gly (BBA00099) dominated.


[0372] In position 3 building blocks carrying an amidine and no amine functionality was found to dominate:



BBA000139




BBA000053




BBA000100

-continued



BBA000140

[0373] The most abundant sequence was thereby found to correspond to the following structure:



BBA000139-BBA000099-BBA000098

[0374] The following 3 sequences

[0375] BBA000098-BBA000099-BBA000139

[0376] BBA000098-BBA000099-BBA000100

[0377] BBA000098-BBA000099-BBA000053

out of the 31 identified sequences were selected for further analysis using a standard ELISA assay and thereby-verified as binders of the $\alpha\beta 3$ Integrin receptor.

[0378] While the invention has been described with references to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention. All patent and literature references cited herein are hereby incorporated by reference in their entirety.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 455

<210> SEQ ID NO 1

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: biotin

<400> SEQUENCE: 1

aattcggaa catactagtc aacatga

27

<210> SEQ ID NO 2

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 2

tcatgttgac tagtatggtg tgctcaagct agtgtgc

37

-continued

<210> SEQ ID NO 3
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 3

tcattggctg atctatggtg tgctcaagct agtgtgc 37

<210> SEQ ID NO 4
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 4

tcatttgta gcttatggtg tgctcaagct agtgtgc 37

<210> SEQ ID NO 5
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 5

tcatggtact cgatatggtg tgctcaagct agtgtgc 37

<210> SEQ ID NO 6
<211> LENGTH: 89
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: displayed molecule

<400> SEQUENCE: 6

gcacactagc ttgagcacac tgacacatgg agatcacatg cttcgacaat gcaggactcc 60
cgcagcttta cgatcccga ggtaaccgt 89

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: biotin

<400> SEQUENCE: 7

gcacactagc ttgagcacac tgaca 25

<210> SEQ ID NO 8
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 8

acggttacct gcgggatcgt aaagc 25

<210> SEQ ID NO 9

<211> LENGTH: 89

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: biotin

<400> SEQUENCE: 9

gcacactagc ttgagcacac tgacacatgg agatcacatg cttcgacaat gcaggactcc 60

cgcagcttta cgatcccgca ggtaaccgt 89

<210> SEQ ID NO 10

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 10

tgtcagtggt ctcaagctag tgtgc 25

<210> SEQ ID NO 11

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 11

acggttacct gcgggatcgt aaagc 25

<210> SEQ ID NO 12

<211> LENGTH: 40

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (40)..(40)

<223> OTHER INFORMATION: chemical entity

<400> SEQUENCE: 12

catgtgatct ccatgtgtca gtgtgctcaa gctagtggtgc 40

<210> SEQ ID NO 13

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(9)

<223> OTHER INFORMATION: N represents inosine (I)

-continued

<400> SEQUENCE: 13

nnnnnnnna tgtgtcagtg tgctcaagct agtgtgc 37

<210> SEQ ID NO 14
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 14

acggttacct gcgggatcgt aaagc 25

<210> SEQ ID NO 15
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (52)..(52)
<223> OTHER INFORMATION: chemical entity
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (16)..(24)
<223> OTHER INFORMATION: N represents inosine (I)

<400> SEQUENCE: 15

gcattgtcga agcatnnnnn nnnnatgtgt cagtgtgctc aagctagtgt gc 52

<210> SEQ ID NO 16
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(9)
<223> OTHER INFORMATION: N represents inosine (I)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (13)..(21)
<223> OTHER INFORMATION: N represents inosine (I)

<400> SEQUENCE: 16

nnnnnnnnnc atnnnnnnnn natgtgtcag tgtgctcaag ctagtgtgc 49

<210> SEQ ID NO 17
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 17

acggttacct gcgggatcgt aaagc 25

<210> SEQ ID NO 18
<211> LENGTH: 64
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (16)..(24)
<223> OTHER INFORMATION: N represents inosine (I)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (64)..(64)
<223> OTHER INFORMATION: chemical entity
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (28)..(36)
<223> OTHER INFORMATION: N represents inosine (I)

<400> SEQUENCE: 18

tgcgggagtc ctgcannnnn nnnncatnnn nnnnnnatgt gtcagtgatgc tcaagctagt    60
gtgc                                                                    64

<210> SEQ ID NO 19
<211> LENGTH: 139
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 19

cagcctggac accacgtcat actagctgct agagatgtgg tgatattagt gtgtgacgat    60
ggtagcgaca agtacgaacg tgcacagag aggacgagca ggacctggaa cctggtgctt    120
cctccaccac gtctctgac                                                    139

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 20

ggaagaagac agaagacctg                                                    20

<210> SEQ ID NO 21
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 21

ctcgaccact gcaggtggag ctcc                                              24

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 22

tcaggagtcg agaactgaag                                                    20

<210> SEQ ID NO 23
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 23

ctgtgcttctct ctgctgcacc accg 24

<210> SEQ ID NO 24

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 24

tgtgtacgtc aacacgtcag 20

<210> SEQ ID NO 25

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 25

cctgggtgctc aggtgagcag cagc 24

<210> SEQ ID NO 26

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 26

tgtggaacta ccatccaagg 20

<210> SEQ ID NO 27

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 27

ctcgacgagg tccatcctgg tcgc 24

<210> SEQ ID NO 28

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 28

ccatccaaca tcggtggaag 20

<210> SEQ ID NO 29

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 29

-continued

cgtgaggagc aggtcctcct gtcg 24

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 30

aacctgtcct gtgagatctg 20

<210> SEQ ID NO 31
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 31

cctgacactg gtcgtggtcg aggc 24

<210> SEQ ID NO 32
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 32

tcacgaagct ggatgatgag 20

<210> SEQ ID NO 33
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 33

ccatctcgac gacctgctcc tggg 24

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 34

tagcatcgat cgaacgtagg 20

<210> SEQ ID NO 35
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 35

ccacgaggtc tccactggtc cagg 24

<210> SEQ ID NO 36

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 36

tcgaagctac tgtcgagatg 20

<210> SEQ ID NO 37
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 37

ccactgagct gtcctccag gtgg 24

<210> SEQ ID NO 38
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 38

cagcttggac accacgtcat ac 22

<210> SEQ ID NO 39
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 39

gtcagagacg tggaggagga a 21

<210> SEQ ID NO 40
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 40

cagcttggac accacgtcat actagctgct agagatgtgg tgatattagt gtgtgacgat 60

<210> SEQ ID NO 41
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 41

cagcttggac accacgtcat acggaagaag acagaagacc tgatattagt gtgtgacgat 60

<210> SEQ ID NO 42
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 42

cagcttggac accacgtcat actcaggagt cgagaactga agatattagt gtgtgacgat 60

<210> SEQ ID NO 43

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 43

cagcttggac accacgtcat actgtgtacg tcaacacgtc agatattagt gtgtgacgat 60

<210> SEQ ID NO 44

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 44

cagcttggac accacgtcat actgtggaac taccatccaa ggatattagt gtgtgacgat 60

<210> SEQ ID NO 45

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 45

cagcttggac accacgtcat accatccaa catcgttggga agatattagt gtgtgacgat 60

<210> SEQ ID NO 46

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 46

cagcttggac accacgtcat acaacctgtc ctgtgagatc tgatattagt gtgtgacgat 60

<210> SEQ ID NO 47

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 47

cagcttggac accacgtcat actcaagaa ctggatgatg agatattagt gtgtgacgat 60

<210> SEQ ID NO 48

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 48

-continued

cagcttgac accacgtcat actagcatcg atcgaacgta ggatattagt gtgtgacgat 60

<210> SEQ ID NO 49
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 49

cagcttgac accacgtcat actcgaagct actgtcgaga tgatattagt gtgtgacgat 60

<210> SEQ ID NO 50
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 50

gtctctctcg atgcacgttc gtacttgtgc gtaccatcgt cacacactaa tacc 54

<210> SEQ ID NO 51
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 51

gaacgtgcat cagagaggac gagcaggacc tggaacctgg tgcaattcca gcttctagga 60

agact 65

<210> SEQ ID NO 52
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 52

gaacgtgcat cagagaggac tcgaccactg caggtggagc tccaattcca gcttctagga 60

agact 65

<210> SEQ ID NO 53
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 53

gaacgtgcat cagagaggac gtgcttcctc tgctgcacca ccgaattcca gcttctagga 60

agact 65

<210> SEQ ID NO 54
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 54

gaacgtgcat cagagaggac ctggtgtcga ggtgagcagc agcaattcca gcttctagga 60
agact 65

<210> SEQ ID NO 55

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 55

gaacgtgcat cagagaggac tcgacgaggt ccatcctggt cgcaattcca gcttctagga 60
agact 65

<210> SEQ ID NO 56

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 56

gaacgtgcat cagagaggac gtgaggagca ggtcctcctg tcgaattcca gcttctagga 60
agact 65

<210> SEQ ID NO 57

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 57

gaacgtgcat cagagaggac ctgacactgg tcgtggtcga ggcaattcca gcttctagga 60
agact 65

<210> SEQ ID NO 58

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 58

gaacgtgcat cagagaggac catctogacg acctgctcct gggaattcca gcttctagga 60
agact 65

<210> SEQ ID NO 59

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 59

gaacgtgcat cagagaggac cactggtct ccaactggtcc aggaattcca gcttctagga 60
agact 65

-continued

<210> SEQ ID NO 60
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 60

gaacgtgcat cagagaggac cactgagctg ctcctccagg tggaattcca gcttctagga 60
agact 65

<210> SEQ ID NO 61
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 61

gtcagagacg tggtgaggga agtcttccta gaagctggaa tt 42

<210> SEQ ID NO 62
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: 6-FAM
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: MGBNFQ

<400> SEQUENCE: 62

tccagcttct aggaagac 18

<210> SEQ ID NO 63
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 63

gtcatactag ctgctagaga tgtggtgata 30

<210> SEQ ID NO 64
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 64

catacggaag aagacagaag acctgata 28

<210> SEQ ID NO 65
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 65

tcatactcag gagtcgagaa ctgaagata 29

<210> SEQ ID NO 66

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 66

catactgtgt acgtcaacac gtcagata 28

<210> SEQ ID NO 67

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 67

catactgtgg aactaccatc caaggata 28

<210> SEQ ID NO 68

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 68

ccatccaaca tcggtggaag at 22

<210> SEQ ID NO 69

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 69

catacaacct gtctgtgag atctgata 28

<210> SEQ ID NO 70

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 70

atactcacga agctggatga tgagata 27

<210> SEQ ID NO 71

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 71

-continued

catactagca tcgatcgaac gtaggata	28
<210> SEQ ID NO 72	
<211> LENGTH: 29	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: synthetic construct	
<400> SEQUENCE: 72	
tcatactcga agctactgtc gagatgata	29
<210> SEQ ID NO 73	
<211> LENGTH: 26	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: synthetic construct	
<400> SEQUENCE: 73	
atattagtgt gtgacgatgg tacgca	26
<210> SEQ ID NO 74	
<211> LENGTH: 24	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: synthetic construct	
<400> SEQUENCE: 74	
acaagtacga acgtgcatca gaga	24
<210> SEQ ID NO 75	
<211> LENGTH: 19	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: synthetic construct	
<400> SEQUENCE: 75	
cgagcaggac ctggaacct	19
<210> SEQ ID NO 76	
<211> LENGTH: 18	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: synthetic construct	
<400> SEQUENCE: 76	
tcgaccactg caggtgga	18
<210> SEQ ID NO 77	
<211> LENGTH: 18	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: synthetic construct	
<400> SEQUENCE: 77	
gcttcctctg ctgcacca	18
<210> SEQ ID NO 78	

-continued

<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 78

ggtgtcgagg tgagcagca 19

<210> SEQ ID NO 79
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 79

cgacgaggtc catcctggt 19

<210> SEQ ID NO 80
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 80

gtgaggagca ggtcctcctg t 21

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 81

ctgacactgg tcgtggtcga 20

<210> SEQ ID NO 82
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 82

catctcgacg acctgctcct 20

<210> SEQ ID NO 83
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 83

acgaggtctc cactggtcca 20

<210> SEQ ID NO 84
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 84

actgagctgc tctccaggt 20

<210> SEQ ID NO 85

<211> LENGTH: 62

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 85

ggcgggtttg ggtgcggggt ttgggtcggg ggtttgggtg gcaatgctcg ggaaggctac 60

tc 62

<210> SEQ ID NO 86

<211> LENGTH: 62

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 86

gcggggtttg ggtcgggggt ttgggtgctg ggtttgggtc ggatcccgtg agtcgatggt 60

tt 62

<210> SEQ ID NO 87

<211> LENGTH: 62

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 87

cgggggtttg ggtgcggggt ttgggtgctg ggtttgggtg cgcgcaccgc agtttgggtca 60

at 62

<210> SEQ ID NO 88

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 88

gcggggtttg ggtcgggggt tgggtgctg tttgggtcgg cacgcggcag tcgagttaat 60

<210> SEQ ID NO 89

<211> LENGTH: 57

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 89

gcggggtttg ggtcgggggt gggtcgggtt tgggtcggag cccggtetca tcgttgt 57

<210> SEQ ID NO 90

<211> LENGTH: 56

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 90
gcgggggttg ggtgggttg ggtgggttg ggtcggtccg acgcaacaat agggca 56

<210> SEQ ID NO 91
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: biotin

<400> SEQUENCE: 91
gccacccaaa cccccg 16

<210> SEQ ID NO 92
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 92
tgttc 5

<210> SEQ ID NO 93
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 93
cgagc 5

<210> SEQ ID NO 94
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 94
ggata 5

<210> SEQ ID NO 95
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 95
cgctg 5

<210> SEQ ID NO 96
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 96

gttat 5

<210> SEQ ID NO 97
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 97

agtgc 5

<210> SEQ ID NO 98
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 98

acctg 5

<210> SEQ ID NO 99
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 99

ctggt 5

<210> SEQ ID NO 100
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 100

tagga 5

<210> SEQ ID NO 101
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 101

actca 5

<210> SEQ ID NO 102
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 102

cttac 5

<210> SEQ ID NO 103

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 103

cgcac 5

<210> SEQ ID NO 104

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 104

tcgcg 5

<210> SEQ ID NO 105

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 105

cggat 5

<210> SEQ ID NO 106

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 106

gagat 5

<210> SEQ ID NO 107

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 107

tgtag 5

<210> SEQ ID NO 108

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 108

gtggtt 5

-continued

<210> SEQ ID NO 109
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 109

agatg 5

<210> SEQ ID NO 110
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 110

atcct 5

<210> SEQ ID NO 111
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 111

ttgct 5

<210> SEQ ID NO 112
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 112

acgta 5

<210> SEQ ID NO 113
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 113

atcac 5

<210> SEQ ID NO 114
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 114

tatcc 5

<210> SEQ ID NO 115
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 115

ggaag 5

<210> SEQ ID NO 116
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 116

cggtc 5

<210> SEQ ID NO 117
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 117

tgctt 5

<210> SEQ ID NO 118
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 118

ttagc 5

<210> SEQ ID NO 119
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 119

gctga 5

<210> SEQ ID NO 120
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 120

gaacg 5

<210> SEQ ID NO 121
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 121

catgg 5

<210> SEQ ID NO 122

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 122

tggtta 5

<210> SEQ ID NO 123

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 123

tcaag 5

<210> SEQ ID NO 124

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 124

atcga 5

<210> SEQ ID NO 125

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 125

atgca 5

<210> SEQ ID NO 126

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 126

actag 5

<210> SEQ ID NO 127

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 127

tacct 5

-continued

<210> SEQ ID NO 128
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 128

tacga 5

<210> SEQ ID NO 129
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 129

cttca 5

<210> SEQ ID NO 130
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 130

ctctt 5

<210> SEQ ID NO 131
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 131

tcatc 5

<210> SEQ ID NO 132
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 132

attcc 5

<210> SEQ ID NO 133
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 133

cgacg 5

<210> SEQ ID NO 134
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 134

cctgt 5

<210> SEQ ID NO 135
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 135

ccttc 5

<210> SEQ ID NO 136
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 136

acacc 5

<210> SEQ ID NO 137
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 137

taaca 5

<210> SEQ ID NO 138
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 138

taaca 5

<210> SEQ ID NO 139
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 139

ccagg 5

<210> SEQ ID NO 140
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 140

atgtc

5

<210> SEQ ID NO 141

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 141

gagga

5

<210> SEQ ID NO 142

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 142

ggtca

5

<210> SEQ ID NO 143

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 143

gactt

5

<210> SEQ ID NO 144

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 144

ggtgg

5

<210> SEQ ID NO 145

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 145

caact

5

<210> SEQ ID NO 146

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 146

atgag

5

-continued

<210> SEQ ID NO 147
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 147

tctgc 5

<210> SEQ ID NO 148
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 148

atagg 5

<210> SEQ ID NO 149
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 149

ctacc 5

<210> SEQ ID NO 150
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 150

aagtg 5

<210> SEQ ID NO 151
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 151

tccaa 5

<210> SEQ ID NO 152
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 152

gctct 5

<210> SEQ ID NO 153
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 153

ggagt 5

<210> SEQ ID NO 154
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 154

aatcg 5

<210> SEQ ID NO 155
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 155

aagct 5

<210> SEQ ID NO 156
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 156

ccgaa 5

<210> SEQ ID NO 157
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 157

tttgt 5

<210> SEQ ID NO 158
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 158

ccgtg 5

<210> SEQ ID NO 159
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 159

tttcg 5

<210> SEQ ID NO 160

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 160

tgagg 5

<210> SEQ ID NO 161

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 161

gttgc 5

<210> SEQ ID NO 162

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 162

aacta 5

<210> SEQ ID NO 163

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 163

aacta 5

<210> SEQ ID NO 164

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 164

cctcg 5

<210> SEQ ID NO 165

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 165

agcaa 5

-continued

<210> SEQ ID NO 166
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 166

ttcca 5

<210> SEQ ID NO 167
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 167

agact 5

<210> SEQ ID NO 168
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 168

aggtt 5

<210> SEQ ID NO 169
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 169

gcgtc 5

<210> SEQ ID NO 170
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 170

aacgt 5

<210> SEQ ID NO 171
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 171

caaga 5

<210> SEQ ID NO 172
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 172

agaga 5

<210> SEQ ID NO 173
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 173

gtact 5

<210> SEQ ID NO 174
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 174

tagag 5

<210> SEQ ID NO 175
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 175

acgat 5

<210> SEQ ID NO 176
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 176

gacca 5

<210> SEQ ID NO 177
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 177

tcggtt 5

<210> SEQ ID NO 178
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 178

gtctc 5

<210> SEQ ID NO 179

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 179

cagca 5

<210> SEQ ID NO 180

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 180

tagtc 5

<210> SEQ ID NO 181

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 181

gggtg 5

<210> SEQ ID NO 182

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 182

ctcag 5

<210> SEQ ID NO 183

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 183

agaac 5

<210> SEQ ID NO 184

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 184

gcgag 5

-continued

<210> SEQ ID NO 185
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 185

gatgt 5

<210> SEQ ID NO 186
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 186

tcact 5

<210> SEQ ID NO 187
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 187

cgtct 5

<210> SEQ ID NO 188
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 188

agctc 5

<210> SEQ ID NO 189
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 189

cactc 5

<210> SEQ ID NO 190
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 190

cagtt 5

<210> SEQ ID NO 191
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 191
agtacgaacg tgcacacagag 20

<210> SEQ ID NO 192
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 192
tagtctcctc cacttccatg 20

<210> SEQ ID NO 193
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 193
tacatcggtc cagactaccg 20

<210> SEQ ID NO 194
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 194
tccagtgcaa gactgaacag 20

<210> SEQ ID NO 195
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 195
agcatcacta ctctgtctgg 20

<210> SEQ ID NO 196
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 196
tcttgtcaac cttccatgcg 20

<210> SEQ ID NO 197
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 197
aaggacgttc ctagtaggtg 20

<210> SEQ ID NO 198
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 198
ggaaccatca agatcctgag 20

<210> SEQ ID NO 199
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 199
atctctgacg agatccaagg 20

<210> SEQ ID NO 200
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 200
tcaaggttg tggtgtactg 20

<210> SEQ ID NO 201
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 201
tcgaacttgt tgcttctcg 20

<210> SEQ ID NO 202
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 202
ctgagtgtgt agtaccaacg 20

<210> SEQ ID NO 203
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 203
atcttggttg ttctcctgag 20

-continued

<210> SEQ ID NO 204
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 204

tagtagcttg gagtagaccg 20

<210> SEQ ID NO 205
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 205

ttcactccat gcagcatgtg 20

<210> SEQ ID NO 206
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 206

acgatgtga tcgatcaacg 20

<210> SEQ ID NO 207
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 207

ttcagtgctt gagctacctg 20

<210> SEQ ID NO 208
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 208

ttggactcct cttgcaccag 20

<210> SEQ ID NO 209
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 209

tcaaccaact ggttcttggg 20

<210> SEQ ID NO 210
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 210
tagtactcta cactgctgcg 20

<210> SEQ ID NO 211
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 211
tacacatga cttgcagacg 20

<210> SEQ ID NO 212
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 212
gcatcttgag tcggtgacg 20

<210> SEQ ID NO 213
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 213
gactcatctc actggagttg 20

<210> SEQ ID NO 214
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 214
tccagcttct aggaagacag 20

<210> SEQ ID NO 215
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 215
cttcttgagt gcactagcag 20

<210> SEQ ID NO 216
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 216
cgagcaggac ctggaacctg gtgc 24

<210> SEQ ID NO 217
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 217
ctcgaccact gcaggtggag ctcc 24

<210> SEQ ID NO 218
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 218
cgtgcttctct ctgctgcacc accg 24

<210> SEQ ID NO 219
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 219
cctggtgtcg aggtgagcag cagc 24

<210> SEQ ID NO 220
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 220
ctcgacgagg tccatctctg tcgc 24

<210> SEQ ID NO 221
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 221
cgtgaggagc agtcctcct gtcg 24

<210> SEQ ID NO 222
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 222
cctgacactg gtcgtggtcg aggc 24

-continued

<210> SEQ ID NO 223
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 223

ccatctcgac gacctgctcc tggg 24

<210> SEQ ID NO 224
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 224

ccacgaggtc tccactggtc cagg 24

<210> SEQ ID NO 225
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 225

ccactgagct gctcctccag gtgg 24

<210> SEQ ID NO 226
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 226

cctcctgtcc tgcacgtcca tccg 24

<210> SEQ ID NO 227
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 227

cagcacctgg aggtaggacc acgg 24

<210> SEQ ID NO 228
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 228

cgaccagacg aggaccaggt aggc 24

<210> SEQ ID NO 229
<211> LENGTH: 24
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 229

ccaggttcga ggacctcgtc agcc 24

<210> SEQ ID NO 230
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 230

cgagcacgag gagcacgtgt ccag 24

<210> SEQ ID NO 231
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 231

ccacgtccac aggtgcacca ggtg 24

<210> SEQ ID NO 232
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 232

cctggtgctc cagcagctgc ttcg 24

<210> SEQ ID NO 233
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 233

cacgtgacga cctggtcagg tggg 24

<210> SEQ ID NO 234
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 234

cgtagctcgt gctggtcctc ctgg 24

<210> SEQ ID NO 235
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 235
cgacgaccac caccttgac accc 24

<210> SEQ ID NO 236
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 236
cctacgtcgt gctcacgtcc tgcc 24

<210> SEQ ID NO 237
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 237
cgacgacagc taggaggagg tggg 24

<210> SEQ ID NO 238
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 238
ctggtggagc tgcacgagca cagc 24

<210> SEQ ID NO 239
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 239
caggactgga cgacgaccag gtcg 24

<210> SEQ ID NO 240
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 240
cgatgctgca gacgaccagc accc 24

<210> SEQ ID NO 241
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 241
tgttc 5

-continued

<210> SEQ ID NO 242
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 242

cgagc 5

<210> SEQ ID NO 243
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 243

ggata 5

<210> SEQ ID NO 244
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 244

cgctg 5

<210> SEQ ID NO 245
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 245

gttat 5

<210> SEQ ID NO 246
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 246

agtgc 5

<210> SEQ ID NO 247
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 247

acctg 5

<210> SEQ ID NO 248
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 248

ctggt 5

<210> SEQ ID NO 249
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 249

tagga 5

<210> SEQ ID NO 250
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 250

actca 5

<210> SEQ ID NO 251
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 251

cttac 5

<210> SEQ ID NO 252
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 252

cgcac 5

<210> SEQ ID NO 253
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 253

tcgcg 5

<210> SEQ ID NO 254
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 254

cggat

5

<210> SEQ ID NO 255

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 255

gagat

5

<210> SEQ ID NO 256

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 256

tgtag

5

<210> SEQ ID NO 257

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 257

gtggt

5

<210> SEQ ID NO 258

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 258

agatg

5

<210> SEQ ID NO 259

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 259

atcct

5

<210> SEQ ID NO 260

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 260

ttgct

5

-continued

<210> SEQ ID NO 261
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 261

acgta 5

<210> SEQ ID NO 262
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 262

atcac 5

<210> SEQ ID NO 263
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 263

tatcc 5

<210> SEQ ID NO 264
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 264

ggaag 5

<210> SEQ ID NO 265
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 265

cggtc 5

<210> SEQ ID NO 266
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 266

tgctt 5

<210> SEQ ID NO 267
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 267

ttagc 5

<210> SEQ ID NO 268
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 268

gctga 5

<210> SEQ ID NO 269
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 269

gaacg 5

<210> SEQ ID NO 270
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 270

catgg 5

<210> SEQ ID NO 271
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 271

tggtta 5

<210> SEQ ID NO 272
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 272

tcaag 5

<210> SEQ ID NO 273
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 273

atcga 5

<210> SEQ ID NO 274

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 274

atgca 5

<210> SEQ ID NO 275

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 275

actag 5

<210> SEQ ID NO 276

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 276

tacct 5

<210> SEQ ID NO 277

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 277

tacga 5

<210> SEQ ID NO 278

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 278

cttca 5

<210> SEQ ID NO 279

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 279

ctctt 5

-continued

<210> SEQ ID NO 280
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 280

tcatc 5

<210> SEQ ID NO 281
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 281

attcc 5

<210> SEQ ID NO 282
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 282

cgacg 5

<210> SEQ ID NO 283
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 283

cctgt 5

<210> SEQ ID NO 284
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 284

ccttc 5

<210> SEQ ID NO 285
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 285

acacc 5

<210> SEQ ID NO 286
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 286

taaca 5

<210> SEQ ID NO 287
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 287

taaca 5

<210> SEQ ID NO 288
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 288

ccagg 5

<210> SEQ ID NO 289
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 289

atgtc 5

<210> SEQ ID NO 290
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 290

gagga 5

<210> SEQ ID NO 291
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 291

ggtca 5

<210> SEQ ID NO 292
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 292

gactt 5

<210> SEQ ID NO 293

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 293

ggtgg 5

<210> SEQ ID NO 294

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 294

caact 5

<210> SEQ ID NO 295

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 295

atgag 5

<210> SEQ ID NO 296

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 296

tctgc 5

<210> SEQ ID NO 297

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 297

atagg 5

<210> SEQ ID NO 298

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 298

ctacc 5

-continued

<210> SEQ ID NO 299
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 299

aagtg 5

<210> SEQ ID NO 300
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 300

tccaa 5

<210> SEQ ID NO 301
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 301

gctct 5

<210> SEQ ID NO 302
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 302

ggagt 5

<210> SEQ ID NO 303
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 303

aatcg 5

<210> SEQ ID NO 304
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 304

aagct 5

<210> SEQ ID NO 305
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 305

ccgaa 5

<210> SEQ ID NO 306
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 306

tttgt 5

<210> SEQ ID NO 307
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 307

ccgtg 5

<210> SEQ ID NO 308
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 308

tttcg 5

<210> SEQ ID NO 309
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 309

tgagg 5

<210> SEQ ID NO 310
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 310

gttgc 5

<210> SEQ ID NO 311
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 311

aacta 5

<210> SEQ ID NO 312

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 312

aacta 5

<210> SEQ ID NO 313

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 313

cctcg 5

<210> SEQ ID NO 314

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 314

agcaa 5

<210> SEQ ID NO 315

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 315

ttcca 5

<210> SEQ ID NO 316

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 316

agact 5

<210> SEQ ID NO 317

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 317

aggtt 5

-continued

<210> SEQ ID NO 318
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 318

gcgtc 5

<210> SEQ ID NO 319
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 319

aacgt 5

<210> SEQ ID NO 320
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 320

caaga 5

<210> SEQ ID NO 321
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 321

agaga 5

<210> SEQ ID NO 322
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 322

gtact 5

<210> SEQ ID NO 323
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 323

tagag 5

<210> SEQ ID NO 324
<211> LENGTH: 5
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 324

acgat 5

<210> SEQ ID NO 325
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 325

gacca 5

<210> SEQ ID NO 326
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 326

tcggt 5

<210> SEQ ID NO 327
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 327

gtctc 5

<210> SEQ ID NO 328
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 328

cagca 5

<210> SEQ ID NO 329
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 329

tagtc 5

<210> SEQ ID NO 330
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 330

gggtg

5

<210> SEQ ID NO 331

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 331

ctcag

5

<210> SEQ ID NO 332

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 332

agaac

5

<210> SEQ ID NO 333

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 333

gcgag

5

<210> SEQ ID NO 334

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 334

gatgt

5

<210> SEQ ID NO 335

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 335

tcact

5

<210> SEQ ID NO 336

<211> LENGTH: 5

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 336

cgtct

5

-continued

<210> SEQ ID NO 337
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 337

agctc 5

<210> SEQ ID NO 338
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 338

cactc 5

<210> SEQ ID NO 339
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 339

cagtt 5

<210> SEQ ID NO 340
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 340

agtacgaacg tgcacagag 20

<210> SEQ ID NO 341
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 341

tagtctctc cacttccatg 20

<210> SEQ ID NO 342
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 342

tacatcggtc cagactaccg 20

<210> SEQ ID NO 343
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 343

tccagtgcaa gactgaacag 20

<210> SEQ ID NO 344
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 344

agcatcacta ctctgtctgg 20

<210> SEQ ID NO 345
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 345

tcttgtcaac cttccatgcg 20

<210> SEQ ID NO 346
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 346

aaggacgttc ctagtaggtg 20

<210> SEQ ID NO 347
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 347

ggaaccatca agatcctgag 20

<210> SEQ ID NO 348
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 348

atctctgacg agatccaagg 20

<210> SEQ ID NO 349
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 349

tcaaggttg tggtgtactg 20

<210> SEQ ID NO 350

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 350

tcgaacttgt tgcttctcg 20

<210> SEQ ID NO 351

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 351

ctgagtgtgt agtaccacg 20

<210> SEQ ID NO 352

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 352

atcttggttg ttctcctcg 20

<210> SEQ ID NO 353

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 353

tagtagcttg gagtagaccg 20

<210> SEQ ID NO 354

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 354

ttcactccat gcagcatgtg 20

<210> SEQ ID NO 355

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 355

acgatgtga tcgatcaacg 20

-continued

<210> SEQ ID NO 356
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 356

ttcagtgctt gagctacctg 20

<210> SEQ ID NO 357
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 357

ttggactcct cttgcaccag 20

<210> SEQ ID NO 358
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 358

tcaaccaact ggttcttggg 20

<210> SEQ ID NO 359
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 359

tagtactcta cactgctgcg 20

<210> SEQ ID NO 360
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 360

tacacatga cttgcagacg 20

<210> SEQ ID NO 361
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 361

gcatcttgag tcggtgaacg 20

<210> SEQ ID NO 362
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 362
gactcatctc actggagttg 20

<210> SEQ ID NO 363
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 363
tccagcttct aggaagacag 20

<210> SEQ ID NO 364
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 364
cttcttgagt gcactagcag 20

<210> SEQ ID NO 365
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 365
cgagcaggac ctggaacctg gtgc 24

<210> SEQ ID NO 366
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 366
ctcgaccact gcaggtggag ctcc 24

<210> SEQ ID NO 367
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 367
cgtgcttcct ctgctgcacc accg 24

<210> SEQ ID NO 368
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 368
cctggtgtcg aggtgagcag cagc 24

<210> SEQ ID NO 369
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 369
ctcgacgagg tccatcctgg tcgc 24

<210> SEQ ID NO 370
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 370
cgtgaggagc aggtcctcct gtcg 24

<210> SEQ ID NO 371
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 371
cctgacactg gtcgtggtcg aggc 24

<210> SEQ ID NO 372
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 372
ccatctcgac gacctgctcc tggg 24

<210> SEQ ID NO 373
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 373
ccacgaggtc tccactggtc cagg 24

<210> SEQ ID NO 374
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 374
ccactgagct gctcctccag gtgg 24

-continued

<210> SEQ ID NO 375
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 375

cctcctgtcc tgcacgtcca tccg 24

<210> SEQ ID NO 376
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 376

cagcacctgg aggtaggacc acgg 24

<210> SEQ ID NO 377
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 377

cgaccagacg aggaccaggt aggc 24

<210> SEQ ID NO 378
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 378

ccaggttcga ggacctcgtc agcc 24

<210> SEQ ID NO 379
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 379

cgagcacgag gagcacgtgt ccag 24

<210> SEQ ID NO 380
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 380

ccacgtccac aggtgcacca ggtg 24

<210> SEQ ID NO 381
<211> LENGTH: 24
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 381

cctggtgctc cacgacgtgc ttcg 24

<210> SEQ ID NO 382
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 382

cacgtgacga cctggtcagg tggg 24

<210> SEQ ID NO 383
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 383

cgtagctcgt gctggtcctc ctgg 24

<210> SEQ ID NO 384
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 384

cgacgaccac caccttgac accc 24

<210> SEQ ID NO 385
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 385

cctacgtcgt gctcacgtcc tgcc 24

<210> SEQ ID NO 386
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 386

cgacgacagc taggaggagg tggg 24

<210> SEQ ID NO 387
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 387
ctggtggagc tgcacgagca cagc 24

<210> SEQ ID NO 388
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 388
caggactgga cgacgaccag gtcgs 25

<210> SEQ ID NO 389
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 389
cgatgctgca gacgaccagc accc 24

<210> SEQ ID NO 390
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 390
tgttc 5

<210> SEQ ID NO 391
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 391
actca 5

<210> SEQ ID NO 392
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 392
cttac 5

<210> SEQ ID NO 393
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 393
cggat 5

-continued

<210> SEQ ID NO 394
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 394

attcc 5

<210> SEQ ID NO 395
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 395

gtctc 5

<210> SEQ ID NO 396
<211> LENGTH: 5
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 396

acagt 5

<210> SEQ ID NO 397
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 397

cacaagtacg aacgtgcatc agag 24

<210> SEQ ID NO 398
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 398

cacatagtct cctccacttc catg 24

<210> SEQ ID NO 399
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 399

cacatacatc gttccagact accg 24

<210> SEQ ID NO 400
<211> LENGTH: 24
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 400
cacatccagt gcaagactga acag 24

<210> SEQ ID NO 401
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 401
cacaagcatc actactctgt ctgg 24

<210> SEQ ID NO 402
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 402
cacatcttgt caaccttcca tgcg 24

<210> SEQ ID NO 403
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 403
cacaaaggac gttcctagta ggtg 24

<210> SEQ ID NO 404
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 404
cacaggaacc atcaagatcc tgag 24

<210> SEQ ID NO 405
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 405
cacaatctct gacgagatcc aagg 24

<210> SEQ ID NO 406
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 406

cacatcaagg ttggtggtgt actg 24

<210> SEQ ID NO 407

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 407

cacatcgaac ttggtgcttc ctcg 24

<210> SEQ ID NO 408

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 408

cacactgagt gtgtagtacc aacg 24

<210> SEQ ID NO 409

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 409

cacaatcttg gttgttctcc tgcg 24

<210> SEQ ID NO 410

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 410

aggacgagca ggacctgga cctgggctgt tctccacca cgtctccg 48

<210> SEQ ID NO 411

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 411

aggactcgac cactgcaggt ggagctccgt tctccacca cgtctccg 48

<210> SEQ ID NO 412

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 412

aggacgtgct tctctgctg caccaccgt tctccacca cgtctccg 48

-continued

<210> SEQ ID NO 413
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 413

aggacctggt gtcgaggtga gcagcagcgt tctccacca cgtctccg 48

<210> SEQ ID NO 414
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 414

aggactcgac gaggtccatc ctggtcgcgt tctccacca cgtctccg 48

<210> SEQ ID NO 415
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 415

aggacgtgag gagcaggtcc tctgtcgcgt tctccacca cgtctccg 48

<210> SEQ ID NO 416
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 416

aggacctgac actggtcgtg gtcgagcgt tctccacca cgtctccg 48

<210> SEQ ID NO 417
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 417

aggaccatct cgacgacctg ctctggggt tctccacca cgtctccg 48

<210> SEQ ID NO 418
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 418

aggaccacga ggtctccact ggtccagggt tctccacca cgtctccg 48

<210> SEQ ID NO 419
<211> LENGTH: 48
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 419
aggaccactg agctgctcct ccaggtgggt tctccacca cgtctccg 48

<210> SEQ ID NO 420
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 420
aggacctcct gtctctgcag tccatccggt tctccacca cgtctccg 48

<210> SEQ ID NO 421
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 421
aggacagcac ctggaggtag gaccacgggt tctccacca cgtctccg 48

<210> SEQ ID NO 422
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 422
aggacgacca gacgaggacc aggtaggcgt tctccacca cgtctccg 48

<210> SEQ ID NO 423
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 423
aggaccaggt tcgaggacct cgtcagcgt tctccacca cgtctccg 48

<210> SEQ ID NO 424
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 424
aggacgagca cgaggagcac gtgtccaggt tctccacca cgtctccg 48

<210> SEQ ID NO 425
<211> LENGTH: 74
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

-continued

<400> SEQUENCE: 425

ggcagcacag tcgtcgcaca tacatcggtc cagactaccg aggacctgac actggtcgtg 60
gtcgaggcgt tcct 74

<210> SEQ ID NO 426

<211> LENGTH: 77

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 426

ggcagcacag tcgtcgctac atgcttgta accttccatg cgagtaacctt acactggctc 60
gtggtcgagg cgttcct 77

<210> SEQ ID NO 427

<211> LENGTH: 74

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 427

ggcagccgga tcgtcgcaca tcttgtaaac cttccatgag aggacctgac actggtcgtg 60
gtcgaggcgt tcct 74

<210> SEQ ID NO 428

<211> LENGTH: 80

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 428

ggcagcctta cgtcgcacaa ttctctgaca gaaatccaac ggaggacctg acacgtgcgt 60
cgtggctcga tgcgttcctc 80

<210> SEQ ID NO 429

<211> LENGTH: 78

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (74)..(74)

<223> OTHER INFORMATION: N

<400> SEQUENCE: 429

ggcagcacag tcgtcgcaca tcattgtaca aaccttccat gcgaggacca tctcgacgac 60
ctgctcctgg ggtncctc 78

<210> SEQ ID NO 430

<211> LENGTH: 75

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 430

-continued

 ggcagcacag tcgtcgca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60

ctcctggggg tctctc 75

<210> SEQ ID NO 431

<211> LENGTH: 75

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 431

ggcagcacag tcgtcgca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60

ctcctggggg tctctc 75

<210> SEQ ID NO 432

<211> LENGTH: 75

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 432

ggcagcacag tcgtcgca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60

ctcctggggg tctctc 75

<210> SEQ ID NO 433

<211> LENGTH: 75

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 433

ggcagcacag tcgtcgca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60

ctcctggggg tctctc 75

<210> SEQ ID NO 434

<211> LENGTH: 75

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 434

ggcagcacag tcgtcgca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60

ctcctggggg tctctc 75

<210> SEQ ID NO 435

<211> LENGTH: 82

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (61)..(61)

<223> OTHER INFORMATION: N

<400> SEQUENCE: 435

ggcagcacta gatcgtcgca catctgtca accttccatg cgaggaccat cttogactga 60

-continued

nctgcctcct gtgggettcc tc 82

 <210> SEQ ID NO 436
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (58)..(58)
 <223> OTHER INFORMATION: N

 <400> SEQUENCE: 436

 ggcagcacag atcgtcgcac atcttgtcaa cttccatgc gaggaccatc tcgacganct 60
 gctcctgggg ttcctc 76

 <210> SEQ ID NO 437
 <211> LENGTH: 78
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

 <400> SEQUENCE: 437

 ggcagcacag tcgtcgcaca tcttgtcaac cttccatgcg aggaccatca cgactacctt 60
 ggctccctgg ggttcctc 78

 <210> SEQ ID NO 438
 <211> LENGTH: 74
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

 <400> SEQUENCE: 438

 ggcagcacag tcgtcgcaca tcttgtcacc ttccatgcga ggaccatctc gacgacctgc 60
 tcctgggggt cctc 74

 <210> SEQ ID NO 439
 <211> LENGTH: 75
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

 <400> SEQUENCE: 439

 ggcagcacag tcgtcgcaca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60
 ctctgggggt tcctc 75

 <210> SEQ ID NO 440
 <211> LENGTH: 75
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

 <400> SEQUENCE: 440

 ggcagccgga tcgtcgcaca tcttgtcaac cttccatgcg aggaccatct cgacgacctg 60
 ctctgggggt tcctc 75

-continued

<210> SEQ ID NO 441
<211> LENGTH: 74
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 441

ggcagccgga tcgtcgaca tcttgcacc ttccatgca ggaccatctc gacgacctgc 60

tcctggggtt cctc 74

<210> SEQ ID NO 442
<211> LENGTH: 75
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 442

ggcagccgga tcgtcgaca tcttgcacc ttccatgca aggaccatctc cgacgacctg 60

ctctggggtt tcctc 75

<210> SEQ ID NO 443
<211> LENGTH: 68
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 443

ggcagcacag tcgtcgcaat ccagtcaaga ctgaacagag gaccatctcg acgacctgct 60

cctggggtt 68

<210> SEQ ID NO 444
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 444

ggcagcacag tcgtcgaca tcttgcacc cttttccatg cgaggacgag caggacctgg 60

aacctgggtc gttcctc 77

<210> SEQ ID NO 445
<211> LENGTH: 74
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 445

ggcagcacag tcgtcgaca tcttgcacc ttccatgca ggacgagcag gacctggaac 60

ctggtcggtt cctc 74

<210> SEQ ID NO 446
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 446

ggcagcacag tcgtcgcaca tcttgtaac cttccatgcg aggacgatgc aggacctgga 60

acctggtgcg ttcctc 76

<210> SEQ ID NO 447

<211> LENGTH: 76

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (28)..(28)

<223> OTHER INFORMATION: N

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (31)..(31)

<223> OTHER INFORMATION: N

<400> SEQUENCE: 447

ggcagccgga tcgtcgcaca tcttggtnaa ncttccatgc gaggacgagc atgaactgga 60

acctggtgcg ttcctc 76

<210> SEQ ID NO 448

<211> LENGTH: 72

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 448

ggcggatcgt cgcacatctt gtcaaccttc catgcgagga ccacgaggtc tccactggtc 60

caggggttcc tc 72

<210> SEQ ID NO 449

<211> LENGTH: 77

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 449

ggcagcacag tcgtcgcaa tctttgtca accttccatg cgaggaccac gaggtctcca 60

ctggtccagg gttcctc 77

<210> SEQ ID NO 450

<211> LENGTH: 75

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 450

ggcagccgga tcgtcgcaca tcttgtaac cttccatgcg aggacgacca agacgaggac 60

caggtaggcg ttcctc 75

<210> SEQ ID NO 451

<211> LENGTH: 76

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 451

ggcagccgga tcgctgcaca tcttgtcaac cttccatgcg aggacgtgat ggagcaagtc 60
ctcctgtcgg ttctctc 76

<210> SEQ ID NO 452
<211> LENGTH: 73
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 452

ggcagcacag tcgctgcaca tcttgtcaac cttccatgcg aggacacgag gtctccactg 60
gtccaggttc ctc 73

<210> SEQ ID NO 453
<211> LENGTH: 79
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc feature
<222> LOCATION: (51)..(53)
<223> OTHER INFORMATION: N
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (63)..(64)
<223> OTHER INFORMATION: N

<400> SEQUENCE: 453

gccccaaaca gtcgctgcac atcttgtcaa ccttccatgc gaggaccgag nngtagctg 60
gannctcgga tgcgttcct 79

<210> SEQ ID NO 454
<211> LENGTH: 84
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 454

gcagcacaga tcgctgcaca tgcttgtcaa gcctttccat cgcgaggacc atcctacgga 60
gcgagcactt gctgcctggg gttc 84

<210> SEQ ID NO 455
<211> LENGTH: 81
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 455

ggcagccgga tcgctgcaca tcaatggttt ggctggatgat actgaggacc acgacgtcta 60
cacttggttc cagggttcct c 81

1. A method for producing a composition of molecules with an improved desired property, comprising the steps of:

- i) providing an initial library comprising a plurality of different encoded molecules associated with a corresponding identifier nucleic acid sequence, wherein each encoded molecule comprises a reaction product of multiple chemical entities and the identifier nucleic acid sequence comprises codons identifying said chemical entities,
- ii) subjecting the initial library to a condition partitioning members having encoded molecules displaying a predetermined property from the remainder of the initial library,
- iii) identifying codons of the identifier nucleic acid sequences of the partitioned members of the initial library, and
- iv) preparing a second-generation library of encoded molecules using the chemical entities coded for by the codons of the partitioned members of the initial library or a part thereof.

2. The method according to claim 1, wherein the second-generation library comprises a plurality of different encoded molecules associated with a corresponding identifier nucleic acid sequence, wherein each encoded molecule comprises a reaction product of multiple chemical entities and the identifier nucleic acid sequence comprises codons identifying said chemical entities.

3. The method of claim 1, further comprising subjecting the second generation library to a condition partitioning members having encoded molecules displaying a predetermined property from the remainder of the second generation library.

4-6. (canceled)

7. The method according to claim 1, wherein the encoded molecule is covalently associated with the corresponding identifier nucleic acid sequence.

8. (canceled)

9. The method according to claim 1, wherein the chemical entities are reacted without enzymatic interaction to produce the encoded molecule.

10. The method according to claim 1, wherein some or all chemical entities are not naturally occurring α -amino acids or precursors thereof.

11. The method according to claim 1, wherein the encoded molecule is not an α -polypeptide.

12. The method according to claim 1, wherein each codon comprises 4 or more nucleotides.

13. The method according to claim 1, wherein the codons are separated by a framing sequence.

14. The method according to claim 13, wherein the framing sequence positions the reaction of a chemical entity in the synthesis history of the encoded molecule.

15. (canceled)

16. The method according to claim 1, wherein the identifier nucleic acid sequence comprises three or more codons.

17. The method according to claim 1, wherein the identifier nucleic acid sequence is amplifiable and comprises codons identifying chemical entities, which have participated in the formation of the encoded molecule.

18. (canceled)

19. The method according to claim 1, wherein the encoded molecule has a molecular weight less than 2000 Dalton, preferably less than 1000 Dalton, and more preferred less than 500 Dalton.

20-21. (canceled)

22. The method according to claim 1, wherein identifier nucleic acid sequence prior to step iii) is amplified.

23. (canceled)

24. The method according to claim 1, wherein the codons of the identifier nucleic acid sequences of the partitioned members of the initial library are identified by contacting said identifier nucleic acid sequences with a pool of nucleic acid fragments under conditions allowing for hybridisation.

25-33. (canceled)

34. The method according to claim 24, wherein nucleic acid fragments are primer oligonucleotides, and the identification involves subjecting the hybridisation complex between the primer oligonucleotides and the identifier nucleic acid sequences to a condition allowing for an extension reaction to occur when the primer is sufficient complementary to a part of the identifier nucleic acid sequence, and evaluating based on measurement of the extension reaction, the presence, absence, or relative abundance of one or more codons.

35-47. (canceled)

48. The method according to claim 24, wherein the nucleic acid fragment is associated with a chemical entity precursor capable of being transferred to a recipient reactive group.

49-59. (canceled)

60. The method according to claim 1, wherein the second generation library is formed by

a) mixing under hybridisation conditions, nascent bifunctional complexes comprising a chemical entity or a reaction product of chemical entities, and an identifier nucleic acid sequence comprising codon(s) identifying said chemical entities, with the recovered nucleic acid fragments, said fragments comprising an oligonucleotide sufficient complementary to at least a part of the identifier nucleic acid sequence to allow for hybridisation, a transferable chemical entity and an anticodon identifying the chemical entity, to form hybridisation products,

b) transferring the chemical entities of the nucleic acid fragments to the nascent bifunctional complexes through a reaction involving a reactive group of the nascent bifunctional complex, in conjunction with a transfer of the genetic information of the anticodon.

61-64. (canceled)

65. The method according to, wherein the second generation library are subjected to a partitioning according to step ii) of claim 1.

66. The method according to claim 1, wherein, prior to the partitioning, the second generation library of complexes are contacted with sequences complementary to the identifier nucleic acid sequences, and the complexes which have hybridised with the complementary sequences are recovered.

67-68. (canceled)

69. The method according to claim 1, wherein the second-generation library is prepared using chemical entities appearing in the initial library and chemical entities foreign to the initial library.

70-75. (canceled)