



HU000035739T2

(19) **HU**(11) Lajstromszám: **E 035 739**(13) **T2****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 14 738117**(22) A bejelentés napja: **2014. 01. 10.**(51) Int. Cl.: **A61K 51/00** (2006.01)**C07D403/14** (2006.01)**C07D413/14** (2006.01)**C07D401/14** (2006.01)(96) Az európai bejelentés bejelentési száma:
EP 20140738117(97) Az európai bejelentés közzétételi adatai:
EP 2943227 A1 **2014. 07. 17.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 14/011047(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2943227 B1 **2017. 09. 20.**

(87) A nemzetközi közzétételi szám:

WO 14110372

(30) Elsőbbségi adatok:

201361752350 P **2013. 01. 14.** **US****201361785788 P** **2013. 03. 14.** **US**

(73) Jogosult(ak):

Molecular Insight Pharmaceuticals, Inc., New York, New York 10007 (US)

(72) Feltaláló(k):

BABICH, John, W, New York, New York 10028 (US)
ZIMMERMAN, Craig, Topsfield, Massachusetts 01983 (US)**JOYAL, John, Melrose, Massachusetts 02176 (US)****LU, Genliang, Winchester, Massachusetts 01890 (US)**

(74) Képviselő:

Danubia Szabadalmi és Jogi Iroda Kft., Budapest

(54)

Triazin alapú radiogyógyszerek és radiológiai képalkotó ágensek

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 943 227 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
20.09.2017 Bulletin 2017/38

(51) Int Cl.:
A61K 51/00 ^(2006.01) **C07D 413/14** ^(2006.01)
C07D 401/14 ^(2006.01) **C07D 403/14** ^(2006.01)

(21) Application number: **14738117.2**

(86) International application number:
PCT/US2014/011047

(22) Date of filing: **10.01.2014**

(87) International publication number:
WO 2014/110372 (17.07.2014 Gazette 2014/29)

(54) **TRIAZINE BASED RADIOPHARMACEUTICALS AND RADIOIMAGING AGENTS**

RADIOPHARMAZEUTIKA AUF TRIAZINBASIS UND STRAHLUNGSABBILDUNGSMITTEL

PRODUITS RADIOPHARMACEUTIQUES À BASE DE TRIAZINE ET AGENTS DE RADIO-IMAGERIE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **14.01.2013 US 201361752350 P**
14.03.2013 US 201361785788 P

(43) Date of publication of application:
18.11.2015 Bulletin 2015/47

(73) Proprietor: **Molecular Insight Pharmaceuticals,
Inc.**
Tarrytown, NY 10591 (US)

(72) Inventors:
• **BABICH, John, W**
New York, New York 10028 (US)
• **ZIMMERMAN, Craig**
Topsfield, Massachusetts 01983 (US)
• **JOYAL, John**
Melrose, Massachusetts 02176 (US)

• **LU, Genliang**
Winchester, Massachusetts 01890 (US)

(74) Representative: **Nederlandsch Octrooibureau**
P.O. Box 29720
2502 LS The Hague (NL)

(56) References cited:
WO-A1-2013/022797 WO-A1-2013/166110
WO-A2-2008/058192 WO-A2-2010/065902
WO-A2-2012/074840 US-A- 5 514 505
US-A1- 2011 183 954 US-A1- 2012 009 121

• **BANERJEE SANGEETA R ET AL: "Synthesis and
evaluation of technetium-99m- and
rhenium-labeled inhibitors of the
prostate-specific membrane antigen (PSMA)",
JOURNAL OF MEDICINAL CHEMISTRY,
AMERICAN CHEMICAL SOCIETY, US, vol. 51, no.
15, 14 August 2008 (2008-08-14), pages
4504-4517, XP002577449, ISSN: 0022-2623, DOI:
10.1021/JM800111U [retrieved on 2008-07-19]**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**CROSS-REFERENCE TO RELATED APPLICATIONS**

- 5 **[0001]** The present application claims the benefit of U.S. Provisional Patent Application Nos. 61/752,350, filed on January 14, 2013, and 61/785,788, filed on March 14, 2013.

FIELD

- 10 **[0002]** The present technology relates generally to the field of radiopharmaceuticals and their use in nuclear medicine as tracers, imaging agents and for the treatment of various disease states.

BACKGROUND

- 15 **[0003]** Many tumors express unique proteins that are predictors of malignancy and a poor prognosis. The expression of such proteins on the surface of tumor cells offers a unique opportunity to use such proteins as markers for the diagnoses of a cancer condition, to evaluate the progression of a cancer condition and to use such proteins as targets for the delivery of a radiotherapeutic agent. Radioactive molecules that selectively bind to specific tumor cell surface proteins provide an attractive route for imaging and treating tumors under non-invasive conditions. In particular, the present invention provides radiolabeled ligands that specifically bind the prostate-specific membrane antigen (PSMA) protein, over expressed on many cancer cells, as agents for imaging or radiation based therapy of PSMA-expressing cancer cells.

- 20 **[0004]** With over a million men suffering from prostate cancer, it is estimated that the disease will strike one in six U.S. men between the ages of 60 and 80. There are more than 300,000 new cases of prostate cancer diagnosed each year and the mortality from the disease is second only to lung cancer. An estimated \$2 billion is currently spent worldwide on surgical, radiation and drugs as treatments for prostate cancer. There is presently no effective therapy for relapsing, metastatic, androgen-independent prostate cancer. New agents that enable rapid visualization of prostate cancer and specific targeting of this cancer tissue for therapeutic purposes are presently needed.

- 25 **[0005]** Human prostate-specific membrane antigen (PSMA), also known as folate hydrolase I (FOLH1), is a trans-membrane, 750 amino acid type II glycoprotein which is primarily expressed in the epithelium of normal human prostate tissue, but is upregulated in prostate cancer, including metastatic disease. PSMA is a unique exopeptidase with reactivity toward poly-gamma-glutamated folates, capable of sequentially removing the poly-gamma-glutamyl termini. Since PSMA is expressed by virtually all prostate cancers and its expression is further increased in poorly differentiated, metastatic and hormone-refractory carcinomas, it is a very attractive target for prostate imaging and therapy. Developing ligands that interact with PSMA and carry appropriate radionuclides, therefore, may provide a promising and novel approach for the detection, treatment and management of prostate cancer.

- 30 **[0006]** The radio-immunoconjugate form of the anti-PSMA monoclonal antibody (mAb) 7E11, known as the PROS-TASCINT scan, is currently being used to diagnose prostate cancer metastasis and recurrence. More recently, monoclonal antibodies that bind to the extracellular domain of PSMA and have a radionuclide were shown to accumulate in PSMA-positive prostate tumor models in animals. However, diagnosis and tumor detection using monoclonal antibodies has been limited by the low permeability of the monoclonal antibody in solid tumor. Tumor detection using low molecular weight radiopharmaceutical compounds, therefore, hold promise and are being explored as potential diagnostic and radiotherapeutic alternatives to radioconjugates of monoclonal antibodies.

- 35 **[0007]** The selective targeting of cancer cells with radiopharmaceuticals, either for imaging or therapeutic purposes is challenging. A variety of radionuclides are known to be useful for radio-imaging or cancer radiotherapy, including ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ¹⁷⁷Lu, ^{99m}Tc, ¹²³I and ¹³¹I. Recently it has been shown that some compounds containing a glutamate-urea-glutamate (GUG) or a glutamate-urea-lysine (GUL) recognition element linked to a radionuclide-complex exhibit high affinity for PSMA. Importantly, the present inventors found that the avidity of the GUL-radionuclide conjugate and GUG-radionuclide conjugate depends at least in part on the chemical nature and size of the linker or spacer joining the GUL or GUG group to the radionuclide complex.

- 40 **[0008]** WO 2008/058192 A2 discloses heterodimers of glutamic acid which have diagnostic and therapeutic properties, such as the treatment of prostate cancer and other diseases related to NAALADase inhibition.

- 45 **[0009]** The present invention focuses on GUL-radiocomplexes or GUG-radiocomplexes that have a one or more optionally substituted triazine groups as part of a linker conjugating the GUL or GUG groups to the radiocomplex. More specifically, the present invention explores the structure-function activity of such triazine-based linkers, for instance by exploring the relationship between binding affinity and linker length as well as the relationship between binding affinity and the position of the optionally substituted triazine moiety such as a piperazinyl-triazine-*p*-aminobenzyl group within the linker. Also described are methods for synthesizing the triazine based radiopharmaceuticals, as well as methods for characterization and for using the inventive GUL-radionuclide and GUG-radionuclide conjugates for the diagnosis and


treatment of cancer.

SUMMARY


[0010] The present invention relates to compounds having a PSMA targeting moiety linked to a radionuclide chelating group as well as radionuclide complexes of the inventive compounds. More specifically, the present technology is focused on the synthesis and use of compounds that conform to the general structure [PSMA recognition motif]-linker-[radionuclide chelating group] and radionuclide complexes of the inventive compounds. As further described below, the inventive compounds and their radionuclide complexes comprise a 1,3,5-triazine moiety within the linker. The incorporation of the 1,3,5-triazine group has advantages since it provides three sites of attachments for the PSMA recognition motif and radionuclide chelating group and also improves the pharmacokinetic properties of the inventive compounds and their radionuclide complexes.

[0011] The invention also provides pharmaceutically acceptable formulations of the inventive compounds and their radionuclide complexes. Such formulations are suitable for treating a variety of disease conditions including without limitation prostate cancer, breast cancer, colorectal cancer, brain cancer, lung cancer, liver cancer, endometrial cancer, bone cancer, ovarian cancer, testicular cancer, skin cancer, pancreatic cancer, uterine cancer, cervical cancer, bladder cancer, esophageal cancer, gastric cancer, head and neck cancers, or kidney cancer.

[0012] In one embodiment therefore, are provided compounds that conform to Formula I and to stereoisomers, tautomers, prodrugs, and pharmaceutically acceptable salts or esters thereof.




[0013] In Formula I, A is $(\text{CHR}^1)_m$ or $\text{C}(\text{O})$ and W is selected from the group consisting of $-\text{C}(\text{O})-(\text{CH}_2)_p-$; $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$; $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$; $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q-$; $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$; and $-(\text{CH}_2)_m-\text{NR}_a-(\text{CH}_2)_n-$. Substituent Y is selected from $-\text{NH}-$, $-\text{NR}^2-$, or



while X in Formula I is selected from $-(\text{C}_1-\text{C}_{10})\text{alkylene}-(\text{C}_3-\text{C}_{10})\text{arylene}$, $-(\text{C}_3-\text{C}_{10})\text{arylene}$, $-(\text{C}_3-\text{C}_{10})\text{arylene}-(\text{C}_1-\text{C}_{10})\text{alkylene}$, phenylene, $-(\text{C}_1-\text{C}_{10})\text{alkylene}-(\text{C}_3-\text{C}_{10})\text{cycloalkylene}$, $-(\text{C}_3-\text{C}_{10})\text{cycloalkylene}$, or $-(\text{C}_3-\text{C}_{10})\text{cycloalkylene}-(\text{C}_1-\text{C}_{10})\text{alkylene}$.


[0014] R^1 and R^2 in Formula I can each independently selected from H, $-(\text{C}_1-\text{C}_{10})\text{alkyl}$, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})\text{alkyl}$, benzyl, $-(\text{C}_3-\text{C}_{10})\text{cycloalkyl}$, or $-(\text{C}_3-\text{C}_{10})\text{aryl}$. For Formula I compounds, R^a and R^b are each independently selected from the group consisting of H, $-\text{OH}$, $-(\text{C}_1-\text{C}_{10})\text{alkyl}$, $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-\text{T}$, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})\text{alkyl}$, $-(\text{C}_1-\text{C}_{10})\text{alkylene}-\text{C}(\text{O})-$, $-(\text{C}_1-\text{C}_{10})\text{alkylene}-\text{C}(\text{O})-\text{Z}$, benzyl, $-(\text{C}_3-\text{C}_{10})\text{cycloalkyl}$, $-(\text{C}_3-\text{C}_{10})\text{aryl}-(\text{C}_1-\text{C}_{10})\text{alkylene}$, $-(\text{C}_3-\text{C}_{10})\text{aryl}$, halo- $-(\text{C}_1-\text{C}_{10})\text{alkyl}$, hydroxy- $-(\text{C}_1-\text{C}_{10})\text{alkyl}$, $-\text{NH}-(\text{C}_1-\text{C}_{10})\text{alkyl}$, and $-(\text{C}_1-\text{C}_{10})\text{alkylene}-\text{NR}^d\text{R}^e$, or R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heteroaryl or (C_3-C_6) -heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O.


[0015] Z in Formula I is selected from $-\text{OH}$, $-\text{O}(\text{C}_1-\text{C}_{10})\text{alkyl}$,




and substituent R^c can be selected from -OH, -O(C₁-C₁₀)alkyl, -Obenzyl, -O(C₃-C₁₀)cycloalkyl, -O(C₃-C₁₀)aryl, -O-(C₁-C₁₀)alkylene-(C₃-C₁₀)aryl, or -O-(C₁-C₁₀)alkylene-(C₃-C₁₀)cycloalkyl.

[0016] For Formula I compounds, R^3 is selected from H, halogen, -OH, -NH₂, -(CH₂)_p-COOH, or -(CH₂)_p-NH₂, substituent T is selected from -H, -OH, -COOH, or -NR^dR^e and R^d and R^e are each independently selected from H, bond, -OH, -(C₁-C₁₀)alkyl, or -(C₃-C₁₀)heteroaryl-(C₁-C₁₀)alkylene. Subscripts m, n, p, q, t and r in Formula I are each independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; and group D is selected from






[0017] Any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene in Formula 1 is optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of $-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ haloalkyl, $-(C_1-C_{10})$ aminoalkyl, $-(C_1-C_{10})$ alkylene-COOH, $-(C_1-C_{10})$ hydroxyalkyl, -OH, halogen, -NH₂, -COOH, -C(O)-(C₁-C₁₀)alkyl, $-(C_1-C_{10})$ alkylene-C(O)-, $-(C_1-C_{10})$ alkylene-C(O)-X, -NH-(C₁-C₁₀)alkyl, and $-(C_1-C_{10})$ alkylene-NR^dRe, and -NR^dRe. Pursuant to these definitions, for certain Formula I compounds, X is phenylene, r is 1 and D is




[0018] The present invention also provides compounds that conform to Formula II, to stereoisomers, tautomers, prodrugs, and pharmaceutically acceptable salts or esters thereof, and to their pharmaceutically acceptable formulations as therapeutics for treating various disease states associated uncontrolled proliferation of cells.




[0019] In Formula II, A is $(CHR^1)_m$ or C(O) and substituent W is selected from the group consisting of $-C(O)-(CH_2)_p-$, $-C(O)-[CH_2-CH_2-O]_n-$, $-[CH_2-CH_2-O]_n-(CH_2)_2-$, $-C(O)-[CH(R^3)]_q-$, $-(CH_2)_m-O-(CH_2)_n-$, $-(CH_2)_m-S-(CH_2)_n-$, $-(CH_2)_m-S(O)-(CH_2)_n-$, $-(CH_2)_m-S(O)_2-(CH_2)_n-$, and $-(CH_2)_m-NR_a-(CH_2)_n-$.

[0020] Group Y in Formula II is selected from -NH-, -NR²-,




or



while variables R^1 and R^2 are each independently selected from H, $-(C_1-C_{10})$ alkyl, $-C(O)-(C_1-C_{10})$ alkyl, benzyl, $-(C_3-C_{10})$ cycloalkyl, or $-(C_3-C_{10})$ aryl.

[0021] In Formula II, R^a and R^b are each independently selected from the group consisting of H, -OH, $-(C_1-C_{10})$ alkyl, $-[CH_2-CH_2-O]_n-(CH_2)_2-T$, $-C(O)-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ alkylene- $C(O)-$, $-(C_1-C_{10})$ alkylene- $C(O)-Z$, benzyl, $-(C_3-C_{10})$ cycloalkyl, $-(C_3-C_{10})$ aryl- (C_1-C_{10}) alkylene, $-(C_3-C_{10})$ aryl, halo- (C_1-C_{10}) alkyl, hydroxy- (C_1-C_{10}) alkyl, $-NH-(C_1-C_{10})$ alkyl, and $-(C_1-C_{10})$ alkylene- NR^dR^e . Alternatively, R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heteroaryl or (C_3-C_6) -heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O.

[0022] Z in Formula II is selected from -OH, $-O(C_1-C_{10})$ alkyl,




and substituent R^c is selected from -OH, $-O(C_1-C_{10})$ alkyl, -Obenzyl, $-O(C_3-C_{10})$ cycloalkyl, $-O(C_3-C_{10})$ aryl, $-O(C_1-C_{10})$ alkylene- (C_3-C_{10}) aryl, or $-O(C_1-C_{10})$ alkylene- $-(C_3-C_{10})$ cycloalkyl.


[0023] For Formula II compounds R^3 is selected from H, halogen, -OH, $-NH_2$, $-(CH_2)_p-COOH$, or $-(CH_2)_p-NH_2$, T is selected from -H, -OH, $-COOH$, or NR^dR^e and each of R^d and R^e are independently selected from H, bond, -OH, $-(C_1-C_{10})$ alkyl, or $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene.

[0024] Any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene in Formula II can be optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of $-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ haloalkyl, $-(C_1-C_{10})$ aminoalkyl, $-(C_1-C_{10})$ alkylene- $COOH$, $-(C_1-C_{10})$ hydroxyalkyl, $-NH_2$, $-COOH$, $-C(O)-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ alkylene- $C(O)-$, $-(C_1-C_{10})$ alkylene- $C(O)-X$, $-NH-(C_1-C_{10})$ alkyl, and $-(C_1-C_{10})$ alkylene- NR^dR^e , and NR^dR^e and subscripts m, n, p, q, t and x are each independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

[0025] For certain Formula II compounds A is $(CH_2)_m$, W is $-C(O)-(CH_2)_p-$ and Y is -NH- or




In one embodiment, A is $(CH_2)_2$, W is $-C(O)-(CH_2)_7-$ or $-C(O)-(CH_2)_{10}-$ and Y is




with R^a and R^b each independently being hydrogen or methyl and substituent R^c is -OH.

[0026] In one embodiment, R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heterocycloalkyl, for example, a group selected from piperidine, piperazine, morpholine, thiomorpholine, isothiazolidine, isoxazolidine, pyrrolidine, imidazolidine, thiazolidine, oxazolidine, or 4-(piperidin-4-yl)butanoic acid.

[0027] For certain other Formula II compounds, R^a is -H and R^b is



with groups R^d and R^e each independently being a $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene, such as




[0028] Also encompassed by the present technology are metal complexes comprising a radionuclide and a compound according to Formula I or Formula II. The radionuclide used is selected from the group consisting of ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{225}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , $^{117\text{m}}\text{Sn}$, ^{169}Yb , ^{90}Y , ^{96}Y , ^{89}Zr and ^{177}Lu .

[0029] The present invention also provides a pharmaceutically acceptable salt, stereoisomer, tautomer, or prodrug of a Formula I or a Formula II compound as well as the radionuclide complexes of Formula I or Formula II compounds.


[0030] Radionuclide complexes of Formula I or II compounds and their pharmaceutical formulations are useful for obtaining radiographic images or for treating a number of diseases and conditions, including but not limited to prostate cancer, breast cancer, colon cancer, brain cancer, lung cancer, liver cancer, endometrial cancer, bone cancer, ovarian cancer, or kidney cancer.

[0031] In one embodiment, the invention provides a method of obtaining a radiographic image of one or more tissues that express prostate-specific membrane antigen (PSMA) by (a) contacting one or more tissues that express PSMA with a metal complex comprising a radionuclide and a compound according to Formula III




or a pharmaceutically acceptable salt or solvate thereof; and (b) recording a radiographic image of the one or more tissues.

[0032] Pursuant to this methodology, variable G in Formula III is




L is selected from -NH-(C₁-C₁₀)alkylene-, -NH-(C₁-C₁₀)alkylene-C(O)-, -C(O)-(C₁-C₁₀)alkylene-, -C(O)-(C₁-C₁₀)alkylene-C(O)- or



and R^a and R^b are each independently selected from the group consisting of H, -OH, -(C₁-C₁₀)alkyl, -[CH₂-CH₂-O]_n-(CH₂)₂-T, -C(O)-(C₁-C₁₀)alkyl, -(C₁-C₁₀)alkylene-C(O)-, -(C₁-C₁₀)alkylene-C(O)-Z, benzyl, -(C₃-C₁₀)cycloalkyl, -(C₃-C₁₀)aryl-(C₁-C₁₀)alkylene, -(C₃-C₁₀)aryl, halo-(C₁-C₁₀)alkyl, hydroxy-(C₁-C₁₀)alkyl, -NH-(C₁-C₁₀)alkyl, and -(C₁-C₁₀)alkylene-NR^dRe-.

[0033] For certain Formula III compounds R^a and R^b together with the nitrogen to which they are bonded form a (C₃-C₆)-heteroaryl or (C₃-C₆)-heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O.

[0034] Substituent Z in Formula III is selected from -OH, -O(C₁-C₁₀)alkyl,



substituents R^d and R^e are each independently selected from H, bond, -OH, $-(C_1-C_{10})$ alkyl, or $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene and subscript n is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

[0035] Pursuant to one embodiment, as noted above, the invention provides a radionuclide complex of Formula I or Formula II compounds as therapeutics for treating a subject diagnosed with cancer for instance prostate cancer. Treatment according to the inventive methodology is effected by administering to a subject a therapeutically effective amount of a prostate-specific membrane antigen (PSMA) binding complex comprising a triazinylene linker and capable of being retained in a PSMA-expressing tumor tissue for a longer interval of time than non-PSMA expressing tissue.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036]

Figure 1 illustrates tissue biodistribution of the ^{177}Lu -complex of (2S)-2-(3-(1-carboxy-5-(11-(4-(4-(2-(2-carboxyethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid according to the present invention in LNCap Xenograft mice.

Figure 2 illustrates tissue biodistribution of the ^{177}Lu -complex of (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid according to the present invention in LNCap Xenograft mice.

Figure 3 illustrates tissue biodistribution of the ^{177}Lu -complex of (21S, 25S)-8,15,23-trioxo-1-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1thioxo-2,7,16,22,24-pentaazaheptacosane-21,25,27-tricarboxylic acid used as a control in LNCap Xenograft mice.

Figure 4 illustrates tissue biodistribution of the ^{177}Lu -complex of (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid according to the present invention in LNCap Xenograft mice.

Figure 5 illustrates in vivo inhibition of LNCaP tumor growth by ^{177}Lu -complex of (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

Figure 6 illustrates a radiographic image obtained by administering to a subject having prostate cancer a ^{68}Ga complex of (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

DETAILED DESCRIPTION

[0037] There are two categories of radiopharmaceuticals: (i) those with biological distribution determined strictly by blood flow, or perfusion, and targeting high capacity systems such as glomerular filtration, phagocytosis, hepatocyte clearance and bone absorption and (ii) those with distribution determined by specific enzymatic or receptor binding interactions, which are low-capacity sites. The radiopharmaceuticals according to Formula I or Formula II belong to the second category and are synthesized by conjugating the radionuclide coordination complex to a biologically active molecule selective for PSMA protein using a linker that has a triazine moiety.

[0038] The terms "linker," "spacer," "linker group" or "spacer group" are used interchangeably in this document and refer to a group that spans the distance between two other identified groups, or which "spaces" them apart. The linker

or spacer may be a bond, an organic group, or an inorganic group or atom.

[0039] In some embodiments, the linker or spacer is an optionally substituted (C₁-C₁₅)alkylene, a (C₂-C₁₅)alkenylene, a (C₂-C₁₅)alkynylene group, a -C(O)-(C₁-C₁₅)alkylene-, a -C(O)-(C₃-C₁₅)arylene-(C₁-C₁₅)alkylene-, -W-Y-(C₃-C₁₅)heteroarylene-NH-X-(CH₂)_r, or a-C(O)-(C₁-C₁₅)alkylene-Y-(C₃-C₁₅)heteroarylene-NH-X-, where the variables "W", "X" and "Y" are further described below. Illustrative substituent groups include without limitation carboxyl groups, carboxylate, hydroxyl groups, and amino (NR^aR^b) groups. For certain embodiments, the (C₁-C₁₅)alkylene group in the linker described above can be replaced by a (C₁-C₁₅)polyol, for example, a polyethylene glycol (PEG) moiety. Exemplary linker or spacer groups are illustrated without limitation throughout the specification and working examples.

[0040] For convenience, certain terms employed herein and within the appended claims are defined here.

[0041] As used herein, "about" will be understood by persons of ordinary skill in the art and will vary to some extent depending upon the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art, given the context in which it is used, "about" will mean up to plus or minus 10% of the particular term.

[0042] The embodiments, illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "comprising," "including," "containing," etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the claimed technology. Additionally, the phrase "consisting essentially of" will be understood to include those elements specifically recited and those additional elements that do not materially affect the basic and novel characteristics of the claimed technology. The phrase "consisting of" excludes any element not specified.

[0043] The use of the terms "a" and "an" and "the" and similar referents in the context of describing the elements (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context.

[0044] The terms "lipophilic group" and "lipophilic moiety" as used herein refer to a group, moiety or substituent that has a greater affinity for non-polar or non-aqueous environments versus polar or aqueous environments. For example, Merriam Webster's online dictionary defines "lipophilic" as "having an affinity for lipids (as fats)." Illustrative lipophilic moieties include aliphatic hydrocarbon radicals, e.g., alkyl radicals, aromatic hydrocarbon radicals, and long-chain acyl radicals; all of them have increasing lipophilicity as the number of constituent carbons increases. In general, addition of a lipophilic moiety to a particular compound will increase the compound's affinity for octanol in the standard octanol/water partition-coefficient-determination protocol; this protocol may be used to gauge a compound's relative hydrophobicity (lipophilicity) and hydrophilicity.

[0045] The term "ligand" refers to a species that interacts in some fashion with another species. In one example, a ligand may be a Lewis base that is capable of forming a coordinate bond with a Lewis Acid. In other examples, a ligand is a species, often organic, that forms a coordination complex with a metal ion. In biochemistry and pharmacology, a ligand is a substance (usually a small molecule), that forms a complex with a biomolecule to serve a biological purpose. In a narrower sense, a ligand is a signal triggering molecule, binding to a site on a target protein. The binding occurs by intermolecular forces, such as ionic bonds, hydrogen bonds and van der Waals forces.

[0046] The term "chelating agent" refers to a molecule, often an organic one, and often a Lewis base, having two or more unshared electron pairs available for donation to a metal ion. The metal ion is usually coordinated by two or more electron pairs to the chelating agent. The terms, "bidentate chelating agent", "tridentate chelating agent", and "tetradentate chelating agent" are art-recognized and refer to chelating agents having, respectively, two, three, and four electron pairs readily available for simultaneous donation to a metal ion coordinated by the chelating agent. Usually, the electron pairs of a chelating agent forms coordinate bonds with a single metal ion; however, in certain examples, a chelating agent may form coordinate bonds with more than one metal ion, with a variety of binding modes being possible.

[0047] The term "coordination" refers to an interaction in which one multi-electron pair donor coordinatively bonds (is "coordinated") to one metal ion.

[0048] The term radionuclide refers to an atom with an unstable nucleus, which is a nucleus characterized by excess energy available to be imparted either to a newly created radiation particle within the nucleus or to an atomic electron.

The radionuclide can undergo radioactive decay and in the process emit subatomic ionizing particles. Illustrative of subatomic ionizing particles without limitation are alpha (α) particles, beta (β) particle and gamma (γ) rays. Exemplary radionuclides include without limitation elements belonging to the lanthanide series, actinide series as well as radioisotopes of transition metals. Illustrative radionuclides may include, but are not limited to ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ⁶⁴Cu, ¹⁵³Gd, ¹⁵⁵Gd, ¹⁵⁷Gd, ⁵⁹Fe, ²²⁵Ac, ²¹²Bi, ²¹³Bi, ⁵⁵Co, ⁶⁷Cu, ¹⁶⁵Dy, ¹⁶⁶Ho, ¹⁹²Ir, ²²³Ra, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁰⁵Rh, ²¹²Pb, ²¹³Pb, ¹⁴⁹Tb, ²²⁷Th, ¹⁵³Sm, ⁸⁹Sr, ^{117m}Sn, ¹⁶⁹Yb, ⁹⁰Y, ⁸⁵Y, ⁸⁹Zr and ¹⁷⁷Lu. However, the term is not limited to these four radionuclides.

[0049] Fmoc is an abbreviation for the chemical group: fluorenylmethyloxycarbonyl.

[0050] The phrases "effective amount" or "therapeutically-effective amount" as used herein means that amount of a compound, material, or composition comprising a compound of the invention, or other active ingredient which is effective

for producing some desired therapeutic effect in at least a sub-population of cells in an animal at a reasonable benefit/risk ratio applicable to any medical treatment. A therapeutically effective amount with respect to a compound of the invention means that amount of therapeutic agent alone, or in combination with other therapies, that provides a therapeutic benefit in the treatment or prevention of a disease. Used in connection with a compound of the invention, the term can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease, or enhances the therapeutic efficacy of or synergies with another therapeutic agent.

[0051] As used herein, the terms "treating" or "treatment" is intended to encompass also diagnosis, prophylaxis, therapy and cure. The patient receiving this treatment is any animal in need, including primates, in particular humans, and other mammals such as equines, cattle, swine and sheep; and poultry and pets in general.

[0052] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0053] The phrase "pharmaceutically-acceptable carrier" as used herein means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, or solvent encapsulating material, involved in carrying or transporting the subject compound from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; and (22) other non-toxic compatible substances employed in pharmaceutical formulations.

[0054] A "pharmaceutically acceptable salt" is a pharmaceutically acceptable, organic or inorganic acid or base salt of a compound of the invention. Representative pharmaceutically acceptable salts include, e.g., alkali metal salts, alkali earth salts, ammonium salts, water-soluble and water-insoluble salts, such as the acetate, amsonate (4,4-diaminostilbene-2, 2'-disulfonate), benzenesulfonate, benzonate, bicarbonate, bisulfate, bitartrate, borate, bromide, butyrate, calcium, calcium edetate, camsylate, carbonate, chloride, citrate, clavulinate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexafluorophosphate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, 3-hydroxy-2-naphthoate, oleate, oxalate, palmitate, pamoate (1,1-methene-bis-2-hydroxy-3-naphthoate, einbonate), pantothenate, phosphate/diphosphate, picrate, polygalacturonate, propionate, p-toluenesulfonate, salicylate, stearate, subacetate, succinate, sulfate, sulfosalicylate, suramate, tannate, tartrate, teoclate, tosylate, triethiodide, and valerate salts. A pharmaceutically acceptable salt can have more than one charged atom in its structure. In this instance the pharmaceutically acceptable salt can have multiple counterions. Thus, a pharmaceutically acceptable salt can have one or more charged atoms and/or one or more counterions.

[0055] The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0056] The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

[0057] A "patient" includes an animal, such as a human, cow, horse, sheep, lamb, pig, chicken, turkey, quail, cat, dog, mouse, rat, rabbit or guinea pig. The animal can be a mammal such as a non-primate and a primate (e.g., monkey and human). In one embodiment, a patient is a human, such as a human infant, child, adolescent or adult.

[0058] The term "prodrug" refers to a precursor of a drug that is a compound which upon administration to a patient, must undergo chemical conversion by metabolic processes before becoming an active pharmacological agent. Illustrative prodrugs of compounds in accordance with Formula I are esters, preferably alkyl esters or fatty acid esters.

[0059] The term "heteroatom" refers to an atom of any element other than carbon or hydrogen. Illustrative heteroatoms include boron, nitrogen, oxygen, phosphorus, sulfur and selenium.

[0060] In general, "substituted" refers to an alkyl, alkylene, alkenyl, alkenylene, alkyne, alkynylene, aryl, arylene,

cycloalkyl, or cycloalkylene group, as defined below in which one or more bonds to a hydrogen atom contained therein are replaced by a bond to non-hydrogen or non-carbon atoms. Substituted groups also include groups in which one or more bonds to a carbon(s) or hydrogen(s) atom are replaced by one or more bonds, including double or triple bonds, to a heteroatom. Thus, a substituted group will be substituted with one or more substituents, unless otherwise specified.

In some embodiments, a substituted group is substituted with 1, 2, 3, 4, 5, or 6 substituents. Examples of substituent groups include: halogens (i.e., F, Cl, Br, and I); hydroxyls; alkoxy, alkenoxy, alkynoxy, aryloxy, aralkyloxy, heterocycloxy, and heterocyclylalkoxy groups; carbonyls (oxo); carboxyls; esters; urethanes; oximes; hydroxylamines; alkoxyamines; aralkoxyamines; thiols; sulfides; sulfoxides; sulfones; sulfonyls; sulfonamides; amines; N-oxides; hydrazines; hydrazides; hydrazones; azides; amides; ureas; amidines; guanidines; enamines; imides; isocyanates; isothiocyanates; cyanates; thiocyanates; imines; nitro groups; nitriles (i.e., CN), haloalkyl, aminoalkyl, hydroxyalkyl, cycloalkyl and the like.

[0061] Alkyl groups include straight chain and branched chain alkyl groups having from 1 to 12 carbon atoms, and typically from 1 to 10 carbons or, in some embodiments, from 1 to 8, 1 to 6, or 1 to 4 carbon atoms. Examples of straight chain alkyl groups include groups such as methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, and n-octyl groups. Examples of branched alkyl groups include, but are not limited to, isopropyl, isobutyl, sec-butyl, tert-butyl, neopentyl, isopentyl, and 2,2-dimethylpropyl groups. Alkyl groups may be substituted or unsubstituted. Unless the number of carbons is otherwise specified, "lower alkyl" refers to an alkyl group, as defined above, but having from one to about ten carbons, alternatively from one to about six carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths.

[0062] The terms "alkylene" and "substituted alkylene" refer to divalent alkyl and divalent substituted alkyl, respectively. Examples of alkylene include without limitation, ethylene ($-\text{CH}_2-\text{CH}_2-$). "Optionally substituted alkylene" refers to alkylene or substituted alkylene.

[0063] The term "alkylcarbonyl" or "alkylenecarbonyl" denote a $-(\text{C}_1-\text{C}_8)\text{alkyl}-\text{C}(\text{O})-$ or $-\text{C}(\text{O})-(\text{C}_1-\text{C}_8)\text{alkyl}-$ groups in which at least one of the methylenes in the C_1-C_8 alkyl group is replaced with a $\text{C}(\text{O})$ group. Representative examples include, but are not limited to, acetyl, propionyl, and $\text{CH}_3(\text{CH}_2)_2\text{C}(\text{O})-$ group, or $-\text{H}_2(\text{CH}_2)_2\text{C}(\text{O})-$.

[0064] The terms "cyclic alkyl" or "cycloalkyl" refers to a saturated or partially saturated non-aromatic cyclic alkyl groups of from 3 to 14 carbon atoms and no ring heteroatoms and having a single ring or multiple rings including fused and bridged ring systems. Cycloalkyl groups may be substituted or unsubstituted. Cycloalkyl or cyclic alkyl groups include mono-, bi- or tricyclic alkyl groups having from 3 to 14 carbon atoms in the ring(s), or, in some embodiments, 3 to 12, 3 to 10, 3 to 8, or 3 to 4, 5, 6 or 7 carbon atoms. Illustrative monocyclic cycloalkyl groups include, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl groups. Bi- and tricyclic ring systems include both bridged cycloalkyl groups and fused rings, such as, but not limited to, bicyclo[2.1.1]hexane, adamantyl, decalyl, and the like.

[0065] A "cycloalkylene" is a divalent saturated or partially saturated non-aromatic cyclo alkyl groups having 3 to 14 carbon atoms and no ring heteroatoms.

[0066] Alkenyl groups include straight and branched chain and cycloalkyl groups as defined above, except that at least one double bond exists between two carbon atoms. Thus, alkenyl groups have from 2 to about 12 carbon atoms in some embodiments, from 2 to 10 carbon atoms in other embodiments, and from 2 to 8 carbon atoms in other embodiments. Examples include, but are not limited to vinyl, allyl, $-\text{CH}=\text{CH}(\text{CH}_3)$, $-\text{CH}=\text{C}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{C}(\text{CH}_3)=\text{CH}(\text{CH}_3)$, $-\text{C}(\text{CH}_2\text{CH}_3)=\text{CH}_2$, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, and hexadienyl, among others. Alkenyl groups may be substituted or unsubstituted. Representative substituted alkenyl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di- or trisubstituted with substituents such as those listed above.

[0067] The term "alkenylene" refers to divalent alkene. Examples of alkenylene include without limitation, ethenylene ($-\text{CH}=\text{CH}-$) and all stereoisomeric and conformational isomeric forms thereof. "Substituted alkenylene" refers to divalent substituted alkene. "Optionally substituted alkenylene" refers to alkenylene or substituted alkenylene.

[0068] "Alkyne" or "alkynyl" refers to straight and branched chain unsaturated hydrocarbon having the indicated number of carbon atoms and at least one triple bond. Examples of a $(\text{C}_2-\text{C}_8)\text{alkynyl}$ group include, but are not limited to, acetylene, propyne, 1-butyne, 2-butyne, 1-pentyne, 2-pentyne, 1-hexyne, 2-hexyne, 3-hexyne, 1-heptyne, 2-heptyne, 3-heptyne, 1-octyne, 2-octyne, 3-octyne and 4-octyne. An alkynyl group can be unsubstituted or optionally substituted with one or more substituents as described herein below.

[0069] The term "alkynylene" refers to divalent alkyne. Examples of alkynylene include without limitation, ethynylene, propynylene. "Substituted alkynylene" refers to divalent substituted alkyne.

[0070] Aryl groups are cyclic aromatic hydrocarbons that do not contain heteroatoms. Aryl groups include monocyclic, bicyclic and polycyclic ring systems. Thus, aryl groups include, but are not limited to, phenyl, azulenyl, heptalenyl, biphenylenyl, indacenyl, fluorenyl, phenanthrenyl, triphenylenyl, pyrenyl, naphthacenyl, chrysenyl, biphenyl, anthracenyl, indenyl, indanyl, pentalenyl, and naphthyl groups. In some embodiments, aryl groups contain 6-14 carbons, and in others from 3 to 12 or even 3-10 carbon atoms in the ring portions of the groups. Aryl group includes both substituted and unsubstituted aryl groups. Substituted aryl groups may be mono-substituted or substituted more than once. For example,

monosubstituted aryl groups include, but are not limited to, 2-, 3-, 4-, 5-, or 6-substituted phenyl or naphthyl groups, which may be substituted with substituent groups such as those listed above.

[0071] "Arylene" denotes divalent aryl, and "substituted arylene" refers to divalent substituted aryl. "Optionally substituted arylene" refers to arylene or substituted arylene. Illustrative of the arylene group is phenylene.

[0072] "Heterocycloalkyl" means a saturated or unsaturated non-aromatic monocyclic, bicyclic, tricyclic or polycyclic ring system that has from 5 to 14 atoms in which from 1 to 3 carbon atoms in the ring are replaced by heteroatoms of O, S or N. A heterocycloalkyl is optionally fused with benzo or heteroaryl of 5-6 ring members, and includes oxidized S or N, such as sulfinyl, sulfonyl and N-oxide of a tertiary ring nitrogen. The point of attachment of the heterocycloalkyl ring is at a carbon or heteroatom such that a stable ring is retained. Examples of heterocycloalkyl groups include without limitation morpholino, tetrahydrofuranyl, dihydropyridinyl, piperidinyl, pyrrolidinyl, piperazinyl, dihydrobenzofuryl, and dihydroindolyl.

[0073] "Optionally substituted heterocycloalkyl" denotes heterocycloalkyl that is substituted with 1 to 3 substituents, e.g., 1, 2 or 3 substituents, attached at any available atom to produce a stable compound, wherein the substituents are as described herein.

[0074] The term "cycloalkyl" refer to monocyclic, bicyclic, tricyclic, or polycyclic, 3-to 14-membered ring systems, which are either saturated, unsaturated or aromatic. The cycloalkyl group may be attached via any atom. Cycloalkyl also contemplates fused rings wherein the cycloalkyl is fused to an aryl or heteroaryl ring as defined above. Representative examples of cycloalkyl include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. A cycloalkyl group can be unsubstituted or optionally substituted with one or more substituents as described herein below.

[0075] The term "cycloalkylene" refers to divalent cycloalkyl. The term "optionally substituted cycloalkylene" refers to cycloalkylene that is substituted with 1 to 3 substituents, e.g., 1, 2 or 3 substituents, attached at any available atom to produce a stable compound, wherein the substituents are as described herein.

[0076] The term "(C₃-C₁₄)aryl-(C₁-C₆)alkylene" refers to a divalent alkylene wherein one or more hydrogen atoms in the C₁-C₆ alkylene group is replaced by a (C₃-C₁₄)aryl group. Examples of (C₃-C₁₄)aryl-(C₁-C₆)alkylene groups include without limitation 1-phenylbutylene, phenyl-2-butylene, 1-phenyl-2-methylpropylene, phenylmethylen, phenylpropylene, and naphthylethylene.

[0077] The term "(C₁-C₁₀)alkylene-(C₃-C₁₄)arylene" refers to a divalent arylene in which one or more hydrogen atoms in the C₃-C₁₄ arylene is replaced by a (C₁-C₁₀)alkyl group and wherein one of the hydrogens of the alkyl group is replaced by another group. Examples of "(C₁-C₁₀)alkylene-(C₃-C₁₄)arylene groups include without limitation butylene-4-phenylene, propylene-2-phenylene, and 1-[2-methylpropylene] phenylene.

[0078] The term "(C₃-C₁₄)arylene-(C₁-C₁₀)alkylene" refers to a divalent alkylene in which one or more hydrogen atoms in the C₁-C₁₀ alkylene is replaced by a divalent (C₃-C₁₄)arylene group. Exemplary of "(C₃-C₁₄)arylene-(C₁-C₁₀)alkylene group include without limitation phenylene-4-butylene, phenylene-2-butylene, and phenylene-1-[2-methylpropylene].

[0079] Aralkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to an aryl group as defined above. In some embodiments, aralkyl groups contain 7 to 20 carbon atoms, 7 to 14 carbon atoms or 7 to 10 carbon atoms.

[0080] "Heterocyclyl" or heterocycloalkyl refers to non-aromatic ring compounds containing 3 or more ring members, of which one or more ring carbon atoms are replaced with a heteroatom such as, but not limited to, N, O, and S. In some embodiments, heterocyclyl groups include 3 to 20 ring members, whereas other such groups have 3 to 6, 3 to 10, 3 to 12, or 3 to 15 ring members. Heterocyclyl groups encompass unsaturated, partially saturated and saturated ring systems, such as, for example, imidazolyl, imidazolyl and imidazolidinyl groups. Heterocyclyl groups may be substituted or unsubstituted. Heterocyclyl groups include, but are not limited to, aziridinyl, azetidyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, thiazolidinyl, tetrahydrothiophenyl, tetrahydrofuranyl, dioxolyl, furanyl, thiophenyl, pyrrolyl, pyrrolinyl, imidazolyl, imidazolyl, pyrazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, piperidyl, piperazinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, tetrahydrothiopyranyl, oxathiane, dioxyl, dithianyl, pyranyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, dihydropyridyl, dihydrodithiyl, dihydrodithiyl, homopiperazinyl, quinuclidyl, indolyl, indolyl, isoindolyl, azaindolyl (pyrrolopyridyl), indazolyl, indolizyl, benzotriazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl, benzthiazolyl, benzoxadiazolyl, benzoxazinyl, benzodithiyl, benzoxathiyl, benzothiazinyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[1,3]dioxolyl, pyrazolopyridyl, imidazopyridyl (azabenzimidazolyl), triazolopyridyl, isoxazolopyridyl, purinyl, xanthinyl, adeninyl, guaninyl, quinolinyl, isoquinolinyl, quinolizyl, quinoxalinyl, quinazolyl, cinnolinyl, phthalazinyl, naphthyridinyl, pteridinyl, thianaphthalenyl, dihydrobenzothiazinyl, dihydrobenzofuranyl, dihydroindolyl, dihydrobenzodioxinyl, tetrahydroindolyl, tetrahydroindazolyl, tetrahydrobenzimidazolyl, tetrahydrobenzotriazolyl, tetrahydropyrrolopyridyl, tetrahydropyrazolopyridyl, tetrahydroimidazopyridyl, tetrahydrotriazolopyridyl, and tetrahydroquinolinyl groups. Heterocyclyl groups may be substituted or unsubstituted. Representative substituted heterocyclyl groups may be mono-substituted or substituted more than once, such as, but not limited to, pyridyl or morpholinyl groups, which are 2-, 3-, 4-, 5-, or 6-substituted, or disubstituted with various substituents such as those listed above.

[0081] Heteroaryl groups are aromatic ring compounds containing 5 or more ring members, of which, one or more

ring carbon atoms are replaced with heteroatom such as, but not limited to, N, O, and S. Heteroaryl groups may be substituted or unsubstituted. Heteroaryl groups include, but are not limited to, groups such as pyrrolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, thiophenyl, benzothiophenyl, furanyl, benzofuranyl, indolyl, azaindolyl (pyrrolopyridyl), indazolyl, benzimidazolyl, imidazopyridyl (azabenzimidazolyl), pyrazolopyridyl, triazolopyridyl, benzotriazolyl, benzoxazolyl, berizothiazolyl, benzothiadiazolyl, imidazopyridyl, isoxazolopyridyl, thianaphthalenyl, purinyl, xanthinyl, adeninyl, guaninyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinoxalinyl, and quinazolinyl groups.

[0082] The term "alkoxy" refers to an -O-alkyl group having the indicated number of carbon atoms. For example, a (C₁-C₁₀)alkoxy group includes -O-methyl (methoxy), -O-ethyl (ethoxy), -O-propyl (propoxy), -O-isopropyl (isopropoxy), -O-butyl (butoxy), -O-*sec*-butyl (*sec*-butoxy), -O-*tert*-butyl (*tert*-butoxy), -O-pentyl (pentoxy), -O-isopentyl (isopentoxy), -O-neopentyl (neopentoxy), -O-hexyl (hexyloxy), -O-isohexyl (isohexyloxy), and -O-neohexyl (neohexyloxy). Examples of cycloalkoxy groups include but are not limited to cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, and the like. Alkoxy groups may be substituted or unsubstituted.

[0083] The term "carbocycle" refers to an aromatic or non-aromatic ring in which each atom of the ring is carbon.

[0084] The term "nitro" refers to -NO₂.

[0085] The term "halogen" is art-recognized and refers to -F, -Cl, -Br or -I; the term "sulfhydryl" is art-recognized and refers to -SH; the term "hydroxyl" means -OH; and the term "sulfonyl" is art-recognized and refers to -SO₂-. "Halide" designates the corresponding anion of the halogens, and "pseudohalide" has the definition set forth on 560 of "Advanced Inorganic Chemistry" by Cotton and Wilkinson.

[0086] The term "amine or amino" refers to an -NR^dR^e group wherein R^d and R^e each independently refer to a hydrogen, (C₁-C₈)alkyl, aryl, heteroaryl, and heterocycloalkyl group. When R^d and R^e are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR^dR^e is meant to include 1-pyrrolidinyl, pyridinyl or a 4-morpholinyl ring.

[0087] The term "amido" is art recognized as an amino-substituted carbonyl and includes a moiety that may be represented by the general formula, $-C(O)NR^dR^e$ group wherein R^d and R^e are as defined above.

[0088] The term "nitrile or cyano" can be used interchangeably and refer to a -CN group which is bound to a carbon atom of a heteroaryl ring, aryl ring and a heterocycloalkyl ring.


[0089] The term "aminoalkyl," refers to an (C)-C₁₀alkyl group wherein one or more hydrogen atoms in the (C₁-C₁₀)alkyl group is replaced with a -NR^dR^e group, where R^d and R^e can be the same or different, for example, R^d and R^e each independently refer to a hydrogen, (C₁-C₈)alkyl, aryl, heteroaryl, heterocycloalkyl, (C₁-C₈)haloalkyl, and (C₁-C₁₀)hydroxyalkyl group. Examples of aminoalkyl groups include, but are not limited to, aminomethyl, aminoethyl, 4-aminobutyl and 3-aminobutyl.

[0090] The term "haloalkoxy," refers to an -O-(C₁-C₈)alkyl group wherein one or more hydrogen atoms in the C₁-C₈ alkyl group is replaced with a halogen atom, which can be the same or different. Examples of haloalkyl groups include, but are not limited to, difluoromethoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, 4-chlorobutoxy, 3-bromopropoxy, pentachloroethoxy, and 1,1,1-trifluoro-2-bromo-2-chloroethoxy.

[illegible]


[0092] A "hydroxyl" or "hydroxy" refers to an -OH group.





[0093] The terms "carboxyl" and "carboxylate" include such moieties as may be represented by the general formulas:



wherein E is a bond or represents O or S, and R^f and R^f individually is H, alkyl, alkenyl, aryl, or a pharmaceutically acceptable salt. Where E is O, and R^f is as defined above, the moiety is referred to herein as a carboxyl group, and particularly when R^f is a hydrogen, the formula represents a "carboxylic acid". In general, where the expressly shown oxygen is replaced by sulfur, the formula represents a "thiocarbonyl" group.

[0094] The substituent -CO₂H, may be replaced with bioisosteric replacements such as:





and the like, wherein R has the same definition as R' and R" as defined herein. See, e.g., THE PRACTICE OF MEDICINAL CHEMISTRY (Academic Press: New York, 1996), at page 203.


[0095] The terms "alkoxyl" or "alkoxy" refer to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxyl groups include methoxy, ethoxy, propoxy, butoxy, *tert*-butoxy and the like. An "ether" is two hydrocarbons covalently linked by an oxygen. "Ether" also encompasses polyethers where more than one ether group, or linkage, may be present in a given group. "Ether" also encompasses cyclic ethers, and crown ethers, where the ether linkage is within a cyclic group.

[0096] The term "(C₅-C₁₄)aryl-(C₁-C₁₀)alkylene" refers to a divalent alkylene wherein one or more hydrogen atoms in the C₁-C₁₀ alkylene group is replaced by a (C₃-C₁₄)aryl group. Examples of (C₃-C₁₄)aryl-(C₁-C₁₀)alkylene groups include without limitation 1-phenylbutylene, phenyl-2-butylene, 1-phenyl-2-methylpropylene, phenylmethylene, phenylpropylene, and naphthylethylene.


[0097] The term "(C₅-C₁₄)heteroaryl-(C₁-C₁₀)alkylene" refers to a divalent alkylene wherein one or more hydrogen atoms in the C₁-C₁₀ alkylene group is replaced a (C₃-C₁₄)heteroaryl group. Examples of (C₃-C₁₄)heteroaryl-(C₁-C₁₀)alkylene groups include without limitation 1-pyridylbutylene, quinoliny-2-butylene and 1-pyridyl-2-methylpropylene.

[0098] The term "-(C₅-C₁₄)heteroarylene-(C₁-C₁₀)alkylene-" refers to a divalent alkylene wherein one or more hydrogen atoms in the C₁-C₁₀ alkylene group is replaced a (C₃-C₁₄)heteroaryl group and wherein one of the hydrogens or one of the heteroatoms of the (C₃-C₁₄)heteroaryl group is bonded to another group, for example, a (C₁-C₁₀)alkyl group.

[0099] A "benzyl" is



while the term "benzylene" denotes a divalent benzyl moiety that is represented by the following structure



[0100] A halogen refers to chlorine, bromine, fluorine, or iodine.

[0101] The definition of each expression, e.g. alkyl, m, n, and the like, when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

[0102] The terms triflyl, tosyl, mesyl, and nonafllyl refer to trifluoromethanesulfonyl, *p*-toluenesulfonyl, methanesulfonyl, and nonafluorobutanesulfonyl groups, respectively. The terms triflate, tosylate, mesylate, and nonaflate are art-recognized and refer to trifluoromethanesulfonate ester, *p*-toluenesulfonate ester, methanesulfonate ester, and nonafluorobutanesulfonate ester functional groups and molecules that contain the groups, respectively. The abbreviations Me, Et, Ph, Tf, Nf, Ts, and Ms represent methyl, ethyl, phenyl, trifluoromethanesulfonyl, nonafluorobutanesulfonyl, *p*-toluenesulfonyl and methanesulfonyl, respectively. A more comprehensive list of the abbreviations utilized by organic chemists of

ordinary skill in the art appears in the first issue of each volume of the Journal of Organic Chemistry; this list is typically presented in a table entitled Standard List of Abbreviations.


[0103] Certain compounds contained in the compositions may exist in particular geometric or stereoisomeric forms. In addition, compounds may also be optically active. The compounds may also include cis- and trans-isomers, R- and S'-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. If, for instance, a particular enantiomer of compound is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

[0104] The phrase "protecting group" as used herein means temporary substituents which protect a potentially reactive functional group from undesired chemical transformations. Examples of such protecting groups include esters of carboxylic acids, silyl ethers of alcohols, and acetals and ketals of aldehydes and ketones, respectively. The field of protecting group chemistry has been reviewed (Greene, T.W.; Wuts, P.G.M. Protective Groups in Organic Synthesis, 3rd ed.; Wiley: New York, 1999).

[0105] Unless otherwise indicated, "stereoisomer" means one stereoisomer of a compound that is substantially free of other stereoisomers of that compound. Thus, a stereomerically pure compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, for example greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, or greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, or greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound.


[0106] If there is a discrepancy between a depicted structure and a name given that structure, then the depicted structure controls. Additionally, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.

[0107] As described above, the present invention relates to compounds according to Formula I.



[0108] For Formula I compounds variable A is $(\text{CHR}^1)_m$ or $\text{C}(\text{O})$ and W is selected from the group consisting of $\text{C}(\text{O})-(\text{CH}_2)_p$; $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n$; $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2$; $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q$; $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n$; $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n$; $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n$; $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n$; and $-(\text{CH}_2)_m-\text{NR}_a-(\text{CH}_2)_n$.


[0109] Variable Y in Formula I is selected from $-\text{NH}-$, $-\text{NR}^2-$, or



and X is group selected from $-(\text{C}_1-\text{C}_{10})\text{alkylene}$ - $(\text{C}_3-\text{C}_{10})\text{arylene}$, $-(\text{C}_3-\text{C}_{10})\text{arylene}$, $-(\text{C}_3-\text{C}_{10})\text{arylene}-(\text{C}_1-\text{C}_{10})\text{alkylene}$ -, phenylene, $-(\text{C}_1-\text{C}_{10})\text{alkylene}-(\text{C}_3-\text{C}_{10})\text{cycloalkylene}$, $-(\text{C}_3-\text{C}_{10})\text{cycloalkylene}$, or $-(\text{C}_3-\text{C}_{10})\text{cycloalkylene}-(\text{C}_1-\text{C}_{10})\text{alkylene}$ -. For certain Formula I compounds X is a $-(\text{C}_3-\text{C}_{10})\text{arylene}$, such as a phenylene group.

[0110] Substituent groups R^1 and R^2 in Formula I are each independently selected from H, $-(C_1-C_{10})$ alkyl, $-C(O)-(C_1-C_{10})$ alkyl, benzyl, $-(C_3-C_{10})$ cycloalkyl, or $-(C_3-C_{10})$ aryl, while groups R^a and R^b are each independently selected from the group consisting of H, $-OH$, $-(C_1-C_{10})$ alkyl, $-[CH_2-CH_2-O]_n-(CH_2)_2-T$, $-C(O)-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ alkylene- $C(O)-$, $-(C_1-C_{10})$ alkylene- $C(O)-Z$, benzyl, $-(C_3-C_{10})$ cycloalkyl, $-(C_3-C_{10})$ aryl- (C_1-C_{10}) alkylene, $-(C_3-C_{10})$ aryl, halo- (C_1-C_{10}) alkyl, hydroxy- (C_1-C_{10}) alkyl, $-NH-(C_1-C_{10})$ alkyl, and $-(C_1-C_{10})$ alkylene- NR^dR^e . For certain Formula I compounds R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heteroaryl or (C_3-C_6) -heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O.


[0111] Z in Formula I can be selected from $-OH$, $-O(C_1-C_{10})$ alkyl,




substituent R^c is selected from $-OH$, $-O(C_1-C_{10})$ alkyl, $-Obenzyl$, $-O(C_3-C_{10})$ cycloalkyl, $-O(C_3-C_{10})$ aryl, $-O-(C_1-C_{10})$ alkylene- (C_3-C_{10}) aryl, or $-O-(C_1-C_{10})$ alkylene- $-(C_3-C_{10})$ cycloalkyl and R^3 is selected from H, halogen, $-OH$, $-NH_2$, $-(CH_2)_p-COOH$, or $-(CH_2)_p-NH_2$.

[0112] In Formula I T is selected from $-H$, $-OH$, $-COOH$, or $-NR^dR^e$ and when T is $-NR^dR^e$, substituent groups R^d and R^e are each independently selected from H, bond, $-OH$, $-(C_1-C_{10})$ alkyl, or $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene;

[0113] Subscripts m, n, p, q, t and r are each independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; and the chelator group D is






[0114] For Formula I compounds any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene is optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of $-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ haloalkyl, $-(C_1-C_{10})$ aminoalkyl, $-(C_1-C_{10})$ alkylene-COOH, $-(C_1-C_{10})$ hydroxyalkyl, -OH, halogen, $-NH_2$, -COOH, $-C(O)-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ alkylene-C(O)-, $-(C_1-C_{10})$ alkylene-C(O)-X, $-NH-(C_1-C_{10})$ alkyl, and $-(C_1-C_{10})$ alkylene-NR^dRe, and -NR^dRe.


[0115] In one aspect for an inventive Formula I compound X is phenylene, subscript "r" is 1 and D is




the metal chelator DOTA. Pursuant to these qualifications is a Formula II compound as illustrated below. For certain Formula II compounds A is $(CHR^1)_m$, W is a $C(O)-(CH_2)_7-$ or $-C(O)-(CH_2)_{10}-$ group and Y is




[0116] In one embodiment, A is $(CHR^1)_m$ with R¹ being a hydrogen and m is 2. For certain Formula II compounds R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heterocycloalkyl selected from piperidine, piperazine, morpholine, thiomorpholine, isothiazolidine, isoxazolidine, pyrrolidine, imidazolidine, thiazolidine or oxazolidine. For some Formula II compounds R^a is -H and R^b is



with R^d and R^e each independently being a $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene, for example, R^d and R^e are each independently





[0117] An illustrative Formula II compound that comports with the above definition is illustrated below:



[0118] Other exemplary Formula I or Formula II compounds include without limitation compounds mentioned in Table 1 below. While some exemplary compounds are depicted with stereochemistry, it should be understood that the invention includes all possible stereoisomers, such as diastereomers, of the compounds.

Table 1





5

10

15

20

25






30

35


40

45

50




55




[0119] Pharmaceutically acceptable salts and/or solvates of the inventive Formula I and Formula II compounds illustrated above are also within the scope of the present invention. In some embodiments, the chelator group, for example, the DOTA group is not complexed with a radionuclide. When DOTA is uncomplexed the carboxylic acid groups of the DOTA group can be in the form of a free acid, or in the form of a salt. The free carboxylic acid groups can also be esterified to obtain the prodrug form of Formula I or Formula II compounds. Suitable ester prodrugs include various alkyl esters, including saturated and unsaturated C₈ to C₁₈ fatty acids.

[0120] The inventive compounds are glutamate-urea-lysine (GUL-) or glutamate-urea-glutamate (GUG) analogs in which a chelator group is conjugated to the GUL- or GUG-moiety via a linker.



[0121] As further discussed below, the length and chemical nature of the linker group is believed to influence the binding avidity of the inventive compounds to the target tissue. Thus, radionuclide complexes of Formula I or Formula II compounds having a piperazine-triazinyl-*p*-aminobenzyl-DOTA moiety within the linker were observed to concentrate to a greater extent in tumor tissue than non-tumor tissue, such as blood, heart, lungs, liver, spleen, stomach, large and small intestines, testes, skeletal muscle, bone, brain, and adipose tissue.





[0122] These compounds, moreover, were rapidly cleared by the kidneys. It was observed that over a period of 96 hours, the piperazine-triazinyl-*p*-aminobenzyl-DOTA containing compounds initially concentrated in the kidneys but at longer intervals of time were rapidly cleared by the kidneys. For example, Formula I or Formula II compounds concentrate to a greater extent in the kidneys than tumor at 4 hours post administration. However, the concentration of the inventive compounds in tumor did not change as a function of time. Thus, the tumor concentration of Formula I or Formula II compounds at 4 hours post administration is similar to their tumor concentration at 24 hours and 96 hours post administration.






[0123] Depending on whether the Formula I or Formula II compounds are to be used as radioimaging agents or radio pharmaceuticals, different radionuclides are complexed to the compounds. Illustrative of suitable radionuclides are those selected from the actinide series, lanthanide series and radionuclides of transition metals, for example, ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{225}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , $^{117\text{m}}\text{Sn}$, ^{169}Yb , ^{90}Y , ^{86}Y , ^{89}Zr and ^{177}Lu .

[0124] Illustrative of Formula I or Formula II compounds complexed to an exemplary radionuclide ^{177}Lu are those illustrated below in Table 2.











or a pharmaceutically acceptable salt or solvate thereof.


[0125] Figure 1 and Figure 2 illustrate results of a bio-distribution study of a GUL-[piperazine-triazinyl-*p*-aminobenzyl]-DOTA-¹⁷⁷Lu complexes according to Formula I or Formula II in LNCap xenograft mice, while Figure 3 illustrates results of a bio-distribution study of a GUL- [alkylene thiourea]-DOTA-¹⁷⁷Lu complex in LNCap xenograft mice.



(A)



(B)




(C)

[0126] As illustrated by the bar graphs in these figures, complexes (A), (B) and (C) concentrate in kidneys and tumor to a greater extent than other tissues. In fact, at 4 hours post administration, the observed concentration for each complex (A), (B) and (C) was greater in kidneys than in tumor. As illustrated by Figures 1-3, however, at 24 hours and 96 hours post administration the concentration of the inventive GUL- [piperazine-triazinyl-*p*-aminobenzyl]-DOTA-¹⁷⁷Lu complexes (A) and (B) in LNCap tumor cells remained unchanged while the concentration of complex (C) used as a control decreases in LNCAP tumor cells at these longer intervals of time.

[0127] These results were unexpected and suggest a greater ability for radionuclide complexes of Formula I or Formula II compounds to concentrate in tumor cells. Moreover, as illustrated in Figures 1 and 2, inventive complexes (A) and (B) are rapidly cleared from the kidneys. Because radionuclide complexes of Formula I or Formula II compounds concentrate in tumor and are rapidly cleared by the kidneys, radionuclide complexes of Formula I or Formula II compounds are candidate therapeutics for treating cancer, for example, prostate cancer.

[0128] Further confirmation that the inventive complexes concentrate in LNCaP tumors but are more rapidly cleared

from other tissues including kidneys post administration to LNCaP tumor bearing mice was obtained in a separate extended biodistribution study using the GUL- [piperazine-triazinyl-*p*-aminobenzyl]-DOTA-¹⁷⁷Lu complex (D), illustrated below.



[0129] As illustrated by the bar graph in Figure 4, the inventive complex concentrates to a greater extent in kidneys and tumor than other tissues at shorter time intervals post administration. For instance, there is a gradual increase in the concentration of complex (D) in kidneys and tumor as a function of time over the first eight hours post administration. At longer time intervals, for example between 24 hours to 96 hours however, the concentration of complex (D) in kidney decreases while there is no observable change in the concentration of complex (D) in tumor.







[0130] To further investigate the pharmacokinetics of tumor retention and renal clearance, the biodistribution study was extended to 3 weeks. Tissue analysis at 1 week post administration of complex (D) indicated no appreciable change in the intracellular concentration of this complex in LNCaP tumor cells. The intrarenal concentration at 1 week post administration of complex (D) is significantly lower than the intrarenal concentration of complex (D) at earlier time intervals, for example, within eight hours post administration.

[0131] At 3 weeks post administration, tissue analysis indicates a decrease in the intratumoral concentration of complex (D). However, the decrease in the concentration of the inventive complex in tumor is less compared to the decrease in the intrarenal concentration of complex (D). As mentioned above, the extended biodistribution study confirmed initial observations that within the same period of time there is a more rapid decrease in the concentration of complex (D) from the kidneys than tumor. Taken together, these results illustrate a greater affinity for the inventive radionuclide complexes that comport with Formula I or Formula II for tumor cells than non-tumor tissue, such as blood, heart, lungs, liver, spleen, stomach, large and small intestines, testes, skeletal muscle, bone, brain, and adipose tissue. Accordingly, Formula I and Formula II compounds are candidate therapeutic or imaging agents for selectively imaging LNCaP tumor cells.









[0132] The compounds of Formula I or Formula II were screened in a human prostate cancer cell competitive binding assay using PSMA positive (+), LNCaP cells against the known inhibitor of PSMA, (7S,14S,18S)-7-amino-1-(1-(carboxymethyl)-1H-imidazol-2-yl)-2-((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)-8,16-dioxo-2,9,15,17-tetraazaicosane-14,18,20-tricarboxylic acid (^{99m}Tc-MIP-1405), and IC₅₀ values were calculated.

[0133] Briefly, LNCaP human prostate cancer cells were obtained from American Type Culture Collection, Rockville, MD. LNCaP cells were maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum (FBS). Binding of the radiolabeled compound and competition with cold derivatives to LNCaP cells was performed according to published methods. Cells were plated in 12-well plates at approximately 4 x 10⁵ cells/well and incubated for 48 hours in a humidified incubator at 37 °C/5% carbon dioxide prior to addition of compound. Solutions of the Formula I or Formula II compounds were prepared and diluted in serum-free cell culture medium containing 0.5% bovine serum albumin (BSA) in combination with 3nM ^{99m}Tc-MIP-1405 (known inhibitor). Total binding was determined by incubating ^{99m}Tc-MIP-1405 without test compound. Plates were incubated at room temperature for 1 hour. Cells were removed from the plates and transferred to eppendorff tubes. Samples were microcentrifuged for 15 seconds at 10k x g. The medium was aspirated and the pellet was washed twice by dispersal in fresh assay medium followed by microcentrifugation. Cell binding of ^{99m}Tc-MIP-1405 was determined by counting the cell pellet in an automated gamma counter. Table 3 illustrates the IC₅₀ values of representative Formula II non-radioactive ¹⁷⁵Lu complexes.



Table 3

Complex	IC ₅₀ (nM)
	7.2
	11
	20
	6.7
	47
	1.3

(continued)

	Complex	IC ₅₀ (nM)
5		13
10		
15		10
20		
25		40
30		
35		129
40		
45		90
50		
55		121
		22
		20

(continued)

Complex	IC ₅₀ (nM)
	17
	6

[0134] As illustrated above, Formula I and Formula II compounds of the invention bind to PSMA expressed on the surface of prostate cancer cells with IC₅₀ values in the nanomolar range. The inventive compounds, therefore, are candidate radiotherapeutic agents for inhibiting the growth of prostate cancer tumor. Please note that in some structures depicted above and elsewhere in this disclosure there may or may not be dashed or solid lines showing putative interactions between certain functional groups and a metal radionuclide. These depictions are merely illustrative of possible bonding interactions, but by no means should they be interpreted as the only possible or actual metal-ligand interaction(s) present for the particular metal complex depicted. For example, it is possible, perhaps even probably, that one or more of the macrocyclic aza groups are contributing to the overall bonding interactions between the metal ion and the chelating ligand.

[0135] Figure 5 illustrates the *in vivo* efficacy of an exemplary lutetium complex of the invention to inhibit the growth of LNCaP tumors in mice. Arrest of tumor growth was determined by administering 450 μ Ci of the lutetium complex of (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid to each mouse in the study group. Mice in the control group were administered saline. Tumor volumes in the test and control group of mice were measured twice weekly. Tumor volumes of mice receiving the lutetium complex according to the invention, were significantly lower than the tumor volumes of mice in the control group.

[0136] In fact, as illustrated in Figure 5, LNCaP tumor volumes of mice in the test group were observed to decrease to values lower than the tumor volume at the start of the study upon administration of the inventive complex. In contrast, there was an increase in the volume of LNCaP tumors in mice receiving saline. These observations indicate that radionuclide complexes of the inventive Formula I and Formula II compounds are effective at arresting the growth of prostate cancer *in vivo*.

[0137] According to another embodiment of the invention, radiometal complexes of Formula I and Formula II compounds were used for imaging prostate cancer and accompanying metastasis in a subject. Briefly, ⁶⁸Ga was complexed to (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid and the resultant complex was administered to a subject having prostate cancer. The subject was then imaged at 1 hour and 3 hours post administration of the inventive ⁶⁸Ga complex using, for example, a ⁶⁸Ga-PSMA PET/CT scanner. As illustrated in Figure 6, PSMA specific lesions were detected in the lymph nodes and bone, in addition to the prostate tissue itself. Some imaging agent was also visible in the subject's bladder at hour 1, which was cleared by the 3-hour scan. The radiographic image in Figure 6 further indicates that the inventive complex accumulates in the lacrimal and salivary glands, kidney, liver, and urinary bladder. Overall, this imaging study supports the use of radiometal complexes of the inventive compounds as suitable agents for radioimaging of cancers, such as prostate cancer.

[0138] Because Formula I and Formula II compounds and their radionuclide complexes can have one or more chiral centers, the present invention encompasses both enantiomers, as well as all of the diastereoisomers. Moreover, both L and D-forms of the natural amino acids can be used for synthesizing the Formula I and Formula II compounds. That is, the present invention encompasses stereoisomers, tautomers, and prodrugs of Formula I and Formula II compounds and their radionuclide complexes.

[0139] As noted above, radionuclide complexes of Formula I or Formula II compounds may contain one or more radi-

onucleides which are suitable for use as radio-imaging agents or as radio-therapeutics for the treatment of diseases associated with the uncontrolled and rapid proliferation of cells, for example, PSMA expressing prostate cancer cells. Accordingly, in one embodiment, a pharmaceutical composition is provided including a complex that includes a metal and a compound of Formula I or Formula II, a salt, solvate, stereoisomer, or tautomer thereof, and a pharmaceutically acceptable carrier.


[0140] In general, metal complexes of a Formula I or a Formula II compound or pharmaceutical compositions thereof, may be administered orally, or *via* a parenteral route, usually by injection. Parenteral routes include, but are not limited to, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion. In some embodiments, the compound, or pharmaceutical composition thereof, is administered orally. Such compositions may take the form of tablets, pills, capsules, semisolids, powders, solutions, suspensions, elixirs, aerosols, or any other appropriate compositions.

[0141] According to another aspect, a pharmaceutical composition is provided, which is suitable for *in vivo* imaging and radiotherapy. Suitable pharmaceutical compositions may contain a radio imaging agent, or a radiotherapeutic agent that has a radionuclide either as an element, *i.e.* radioactive iodine, or a radioactive metal chelate complex of the compound of Formula I or Formula II in an amount sufficient for imaging, together with a pharmaceutically acceptable radiological vehicle. The radiological vehicle should be suitable for injection or aspiration, such as human serum albumin; aqueous buffer solutions, e.g., tris(hydromethyl) aminomethane (and its salts), phosphate, citrate, bicarbonate, etc; sterile water; physiological saline; and balanced ionic solutions containing chloride and or dicarbonate salts or normal blood plasma cations such as calcium, potassium, sodium, and magnesium.


[0142] The concentration of the imaging agent or the therapeutic agent in the radiological vehicle should be sufficient to provide satisfactory imaging. For example, when using an aqueous solution, the dosage is about 1.0 to 50 millicuries. The actual dose administered to a patient for imaging or therapeutic purposes, however, is determined by the physician administering treatment. The imaging agent or therapeutic agent should be administered so as to remain in the patient for about 1 to 24 hours, although both longer and shorter time periods are acceptable. Therefore, convenient ampoules containing 1 to 10 mL of aqueous solution may be prepared.

[0143] Imaging may be carried out in the normal manner, for example by injecting a sufficient amount of the imaging composition to provide adequate imaging and then scanning with a suitable machine, such as a gamma camera. In certain embodiments, a method of imaging a region in a patient, for example, imaging one or more tissues that express prostate-specific membrane antigen (PSMA) includes the steps of: (i) administering to a patient a diagnostically effective amount of a Formula 1, formula II or Formula III compound complexed with a radionuclide so as to contact the one or more tissues expressing PSMA with a radionuclide complex of a Formula I, Formula II or Formula III compound; and (ii) recording a radiographic image of the one or more tissues. In one embodiment the tissue imaged is a prostate tissue or a prostate cancer tissue. According to the inventive methodology, imaging can be carried out by administering to a patient a diagnostically effective amount of a Formula I compound complexed to a radionuclide, a Formula 11 compound complexed to a radionuclide or a Formula III compound complexed to a radionuclide, or a pharmaceutically acceptable salt or solvate of the inventive complexes.


[0144] In one embodiment, therefore, imaging is carried out using a radionuclide complex of a Formula III compound.



[0145] In Formula III, G is




is selected from -NH-(C₁-C₁₀)alkylene-, -NH-(C₁-C₁₀)alkylene-C(O)-, -C(O)-(C₁-C₁₀)alkylene-, -C(O)-(C₁-C₁₀)alkylene-C(O)- or



and variables R^a and R^b are each independently selected from the group consisting of H, -OH, -(C₁-C₁₀)alkyl, -[CH₂-CH₂-O]_n-(CH₂)₂-T, -C(O)-(C₁-C₁₀)alkyl, -(C₁-C₁₀)alkylene-C(O)-, -(C₁-C₁₀)alkylene-C(O)-Z, benzyl, -(C₃-C₁₀)cycloalkyl, -(C₃-C₁₀)aryl-(C₁-C₁₀)alkylene, -(C₃-C₁₀)aryl, halo-(C₁-C₁₀)alkyl, hydroxy-(C₁-C₁₀)alkyl, -NH-(C₁-C₁₀)alkyl, and -(C₁-C₁₀)alkylene-NR^dR^e-, or R^a and R^b together with the nitrogen to which they are bonded form a (C₃-C₆)-heteroaryl or (C₃-C₆)-heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O.

[0146] Z in Formula III is selected from -OH, -O(C₁-C₁₀)alkyl,



while R^d and R^e are each independently selected from H, bond, -OH, -(C₁-C₁₀)alkyl, or -(C₃-C₁₀)heteroaryl-(C₁-C₁₀)alkylene. Subscript "n" is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. For Formula III compounds, any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene is optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of -(C₁-C₁₀)alkyl, -(C₁-C₁₀)haloalkyl, -(C₁-C₁₀)aminoalkyl, -(C₁-C₁₀)alkylene-COOH, -(C₁-C₁₀)hydroxyalkyl, -NH₂-, -COOH, -C(O)-(C₁-C₁₀)alkyl, -(C₁-C₁₀)alkylene-C(O)-, -(C₁-C₁₀)alkylene-C(O)-X, -NH-(C₁-C₁₀)alkyl, and -(C₁-C₁₀)alkylene-NR^dR^e-, and -NR^dR^e.

[0147] The metal used to form the complex is a radionuclide selected from ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ⁶⁴Cu, ¹⁵³Gd, ¹⁵⁵Gd, ¹⁵⁷Gd, ⁵⁹Fe, ²²⁵Ac, ²¹²Bi, ²¹³Bi, ⁵⁵Co, ⁶⁷Cu, ¹⁶⁵Dy, ¹⁶⁶Ho, ¹⁹²Ir, ²²³Ra, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁰⁵Rh, ²¹²Pb, ²¹³Pb, ¹⁴⁹Tb, ²²⁷Th, ¹⁵³Sm, ⁸⁹Sr, ^{117m}Sn, ¹⁶⁹Yb, ⁹⁰Y, ⁸⁶Y, ⁸⁹Zr and ¹⁷⁷Lu.

[0148] The amount of a Formula I, Formula II or Formula III compound, or a formulation comprising a complex of a radiometal and a compound according to Formula I or Formula II, or its salt, solvate, stereoisomer, or tautomer that is administered to a patient depends on several physiological factors that are routinely used by the physician, including the nature of imaging to be carried out, tissue to be targeted for imaging and the body weight and medical history of the patient to be imaged.

[0149] Also described is a method for treating a patient diagnosed with cancer by administering to such a patient a therapeutically effective amount of a prostate-specific membrane antigen (PSMA) binding complex comprising a triazinylene linker. In one embodiment of this methodology, the prostate-specific membrane antigen (PSMA) binding complex comprising a triazinylene linker is a Formula I, Formula II or Formula III compound complexed to a radionuclide, or a pharmaceutically acceptable salt or solvate of the complex. Radionuclide complexes of Formula I, Formula II and Formula III compounds, as described above, are preferentially retained in PSMA-expressing tumor tissue than non-PSMA expressing tissues such as kidney, liver, spleen, heart, blood, lungs, muscle, bone, large intestine, small intestine, brain, or fat. In addition to prostate cancer, radionuclide complexes of Formula I or Formula II compounds are also candidate therapeutics for treating breast cancer, colon cancer, brain cancer, lung cancer, liver cancer or kidney cancer.

[0150] The present invention, thus generally described, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

EXAMPLES

General Protocol for Cell Culture

[0151] Human prostate cancer LNCaP cells were obtained from the American Type Culture Collection. Cell culture supplies were from Invitrogen unless otherwise noted. LNCaP cells were maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum (Hyclone), 4 mM L-glutamine, 1 mM sodium pyruvate, 10 mM hepes, 2.5 mg/mL D-glucose, and 50 µg/mL gentamicin in a humidified incubator at 37 °C/5% CO₂. Cells were removed from flasks for passage, inoculation of mice or for transfer to 12-well assay plates by incubating them with 0.25% trypsin/EDTA.

General Protocol for Competitive Binding

[0152] The ability of non-radioactive lutetium containing PSMA inhibitors to compete with ^{99m}Tc -((7S,14S,18S)-7-amino-1-(1-(carboxymethyl)-1*H*-imidazol-2-yl)-2-((1-(carboxymethyl)-1*H*-imidazol-2-yl)methyl)-8,16-dioxo-2,9,15,17-tetraazaicosane-14,18,20-tricarboxylic acid) for binding to PSMA in LNCaP cells was examined. LNCaP cells (4×10^5 cells/well in 12-well plates in triplicate) were incubated for 1 hour with 3nM of the ^{99m}Tc -complex in RPMI medium containing 0.5% BSA in the presence of 1-10,000 nM test compounds. Cells were removed to Eppendorf tubes by gently pipetting, washed twice with RPMI + 0.5% BSA and counted.

Mouse Studies

[0153] All animal studies were approved by the Institute for Animal Care and Use Committee in accordance with the guidelines set forth by the U.S. Public Health Service *Policy on Humane Care and Use of Laboratory Animals*. Mice were housed under standard conditions in approved facilities with 12 hour light/dark cycles and given food and water *ad libitum*. Male athymic NCr-nu/nu mice were purchased from Taconic. For inoculation in mice, LNCaP cells were resuspended at 10^7 cells/ml in a 1:1 mixture of cell culture medium:Matrigel (BD Biosciences). Each mouse was injected in the right flank with 0.25 ml of the cell suspension. Mice were used for tissue distribution studies when the tumors reached approximately 100-400 mm³.

Tissue Distribution

[0154] A quantitative analysis of the tissue distribution of ^{177}Lu -labeled compounds was performed in separate groups of male NCr-nu/nu mice bearing LNCaP cell xenografts. The compounds were administered via the tail vein as a bolus injection (approximately 10 μCi /mouse) in a constant volume of 0.05 mL. The animals ($n=5$ /time point) were euthanized by asphyxiation with carbon dioxide at the indicated time points after injection. Tissues, for example, blood, heart, lungs, liver, spleen, kidneys, stomach, large and small intestines (with contents), testes, skeletal muscle, bone, brain, adipose, and tumor were dissected, excised, weighed wet, and counted in an automated γ -counter. Tissue time-radioactivity levels were expressed as percent injected dose per gram of tissue (%ID/g).

In Vivo Efficacy

[0155] Mice bearing LNCaP xenografts having an average volume of ~100-500 mm³, were randomly assigned to a control group or a treatment group ($n = 10$ mice per group). Mice in the control group were administered saline while mice in the test group received 450 μCi /mouse of ^{177}Lu -complex of the inventive Formula I or Formula II compound. Each animal was administered the test article intravenously in a volume of 0.05 mL. Tumor dimensions were measured twice weekly with digital calipers and tumor volumes were calculated using the formula $(4/3 \times \pi \times \text{width}^2 \times \text{length})/6$. Measurements were made until tumor volumes in the vehicle group reached the maximum allowed by IACUC guidelines (1,500 mm³).

General Synthetic Methods.

[0156] General procedure for the synthesis of Formula I compounds and for complexation of a Formula I compound with a radionuclide are described. While a protocol for complexing lutetium to a Formula I compound is exemplified below, it is to be understood that a similar synthetic procedure can be followed for complexing other radionuclides. Therefore, while lutetium may specifically be shown in various examples described below, complexes with other radionuclides such as In, Y, Zr, Ga, Lu, Cu, Gd, Ac Fe, Bi Co, Dy Ho, Ir, Ra, Re, Rh, Sr or Sm are within the scope of the present invention. Additionally, it is to be understood that various isotopes of these elements may be complexed, for example, ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{225}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , ^{117m}Sn , ^{169}Yb , ^{90}Y , ^{86}Y , ^{89}Zr and ^{177}Lu .

General Experimental Conditions for the Formation of the lutetium Complexes

[0157] The lutetium complexes of Formula I compounds were conveniently isolated from the reactions that involve contacting commercially available LuCl_3 with a compound according to Formula I. Briefly, a 10^{-6}M - 10^{-4}M solution of the desired Formula I or Formula II compound in an equal volume mixture of 1:1 acetonitrile and phosphate buffer was contacted with LuCl_3 in a sealed vial. The reaction mixture was heated at 100 °C for 30 to 45 minutes. Upon cooling, the reaction was analyzed for completion and purity by reverse-phase high pressure liquid chromatography (RP-HPLC) and if required was purified using RP-HPLC or a C18 Sep Pak column. The overall average yield of the lutetium complexed

product following purification was in the range from about 20% to about 99%. The radiochemical purity (RCP), after HPLC purification, however, was consistently $\geq 95\%$.


[0158] Initial results demonstrated radiolabeling of a Formula I or a Formula II compound at concentrations as low as 10^{-6} M, the radiochemical yield (RCY) at this concentration of reagents was approximately $\leq 80\%$. To achieve a higher RCY, greater than 95%, the reaction temperature and concentration of reagents in the reaction mixture were increased to 10^{-4} M.

[0159] A similar synthetic strategy was used to incorporate other radionuclides. Moreover, the introduction of a radionuclide can be made either prior to deprotection of a Formula I or Formula II compound, or after deprotecting a Formula I compound.


Synthesis of Exemplary Triazine-piperazine Based Formula I, Formula II, or Formula III Compounds

[0160] Schemes A, B and C illustrate general synthetic protocols for exemplary Formula I compounds. Briefly, *p*-aminobenzyl DOTA is contacted with cyanuric chloride followed by reaction of the resultant product with an amine. The product thus formed is then contacted with a GUG- or GUL-linker-piperazine moiety to obtain a Formula I compound.


Scheme A.



Scheme B




Scheme C




Example 1: (2S)-2-(3-((1S)-1-carboxy-5-(8-((4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)amino)octanamido)pentanediolic acid lutetium complex.

[0161]



Step 1. (18S,22S)-tri-*tert*-butyl 1-(9*H*-fluoren-9-yl)-3,12,20-trioxo-2-oxa-4,13,19,21-tetraazatetracosane-18,22,24-tricarboxylate.


[0162]



[0163] A solution of (S)-di-*tert*-butyl 2-(3-((S)-6-amino-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (1.9677 g, 4.03 mmol), 8-(((9*H*-fluoren-9-yl)methoxy)carbonyl)amino)octanoic acid (1.84 g, 4.84 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI; 0.770 g, 4.03 mmol), HOBT (0.544 g, 4.03 mmol) and *N,N*-diisopropylethylamine (DIPEA; 2.0 mL) in DCE (100 mL) was stirred at room temperature for overnight. The solvent was evaporated to give a residue, which was purified by silica gel column chromatography (Biotage) using a mixture of DCM/MeOH as the eluent to give (18S,22S)-tri-*tert*-butyl 1-(9*H*-fluoren-9-yl)-3,12,20-trioxo-2-oxa-4,13,19,21-tetraazatetracosane-18,22,24-tricarboxylate (2.099 g, 61%) as a white solid. MS (ESI), 851.2 (M+H)⁺.

Step 2. (S)-di-*tert*-butyl 2-(3-((S)-6-(8-aminooctanamido)-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate.


[0164]



[0165] To a solution of (18S,22S)-tri-*tert*-butyl 1-(9*H*-fluoren-9-yl)-3,12,20-trioxo-2-oxa-4,13,19,21-tetraazatetracosane-18,22,24-tricarboxylate (1.983 mg, 2.333 mmol) in DMF (4.0 mL) was added piperidine (4.0 mL). The mixture was stirred at room temperature for 3 hrs following which the solvent was evaporated under reduce pressure to afford a residue, which was purified by column chromatography using a Biotage SP4 column and gradient elution using 100% DCM to a 1:1 mixture of DCM:methanol as the eluting solvent. The product (S)-di-*tert*-butyl-2-(3-((S)-6-(8-aminooctanamido)-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (1.039 mg, 71%), thus obtained was characterized using ¹H NMR and mass spectrometry. ¹H NMR (400 MHz, DMSO-*d*₆) 7.71 (t, *J* = 5.2 Hz, 1 H), 6.29 (d, *J* = 8.0 Hz, 1 H), 6.25 (d, *J* = 8.4 Hz, 1 H), 5.74 (brs, 2 H), 4.05-3.91 (m, 2 H), 3.01-2.88 (m, 2 H), 2.63 (t, *J* = 6.8 Hz, 2 H), 2.20-1.22 (m, 49 H); MS (ESI), 629.3 (M+H)⁺.

Step 3. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(dimethylamino)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid.

[0166]




[0167] To a solution of *p*-NH₂-Bn-DUTA-tetra(*t*-Bu-ester) (67.8 mg, 0.080 mmol) and cyanuric chloride (14.7 mg, 0.080 mmol) in DCM (4.0 mL) was added DIPEA (0.10 mL). This solution was stirred at room temperature for 3 hrs, following which the solvent was removed under a stream of nitrogen to give a residue. To a DMSO (4.0 mL) solution of the residue was added (S)-di-*tert*-butyl 2-(3-((S)-6-(8-aminooctanamido)-1-(*tert*-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (50.3 mg, 0.08 mmol) and K₂CO₃ (100mg). The suspension was stirred at room temperature for about 2 hrs and then a tetrahydrofuran solution of dimethylamine (0.3 mL, 2.0 M in THF) was added to the reaction mixture. After stirring at room temperature continuously for 16 hrs, the reaction mixture was lyophilized to afford the crude triazine intermediate. The crude product was deprotected by the addition of TFA (4.0 mL) and DCM (1.0 mL) and stirring the reaction mixture at room temperature for 4 hours. Removal of the solvent using a stream of nitrogen gas gave a residue, which was purified using Biotage SP4 via C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(dimethylamino)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (67 mg) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.83-7.60 (m, 3 H), 7.17 (d, *J* = 8.0 Hz, 2 H), 6.32 (d, *J* = 8.0 Hz, 1 H), 6.28 (d, *J* = 8.4 Hz, 1 H), 4.10-1.27 (m, 61 H); MS (ESI), 1091.4 (M+H)⁺.

Step 4. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(dimethylamino)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0168] To solid (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(dimethylamino)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (5.7 mg, 0.00522 mmol) was added LuCl₃ (1.46 mL of a 0.00357 mmol/mL, 0.00522 mmol) and acetonitrile (0.50 mL). The reaction mixture was heated at 95 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(dimethylamino)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex (6.2 mg) as a white solid. MS (ESI), 1263.0 (M+H)⁺.


Example 2. (S)-2-(3-((S)-1-Carboxy-5-(8-((4-(piperidin-1-yl)-6-((4-(((S)-1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)amino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0169]



Step 1. (2S)-2-(3-((S)-1-carboxy-5-(8-(4-(piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid.

[0170]



[0171] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL). This reaction mixture was stirred at room temperature for 2 hours following which the solvent was removed using a stream of nitrogen to give a residue. The residue thus obtained was dissolved in DMSO (4.0 mL) and (S)-di-*tert*-butyl 2-(3-((S)-6-(8-aminooctanamido)-1-(*tert*-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (31.4 mg, 0.05 mmol) and K₂CO₃ (100 mg) were added. The suspension was stirred at room temperature for 2 hrs and then piperidine (0.10 mL) was added. The reaction mixture was stirred at room temperature for an additional 14 hrs and then lyophilized to afford a triazine intermediate, which was deprotected by the addition of TFA (2.0 mL) in DCM (1.0 mL). Deprotection was carried out by stirring the reaction mixture at room temperature for 4 hours. Following deprotection, the solvent was removed using a stream of nitrogen to give a residue, which was purified by Biotage SP4 using C18 cartridge to give pure (2S)-2-(3-((S)-1-carboxy-5-(8-(4-(piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid


[0172] (25.8 mg) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) 7.75-7.60 (m, 3 H), 7.18 (d, *J* = 7.2 Hz, 2 H), 6.33 (d, *J* = 7.6 Hz, 1 H), 6.30 (d, *J* = 8.0 Hz, 1 H), 4.12-1.24 (m, 65 H); MS (ESI), 1131.2 (M+H)⁺.

Step 2. (2S)-2-(3-((S)-1-carboxy-5-(8-(4-(piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0173] To solid (2S)-2-(3-((S)-1-carboxy-5-(8-(4-(piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (9.2 mg, 0.00814 mmol) was added LuCl₃ (1.60 mL, of a 0.00513 mmol/mL, 0.0082 mmol) and acetonitrile (0.50 mL). The reaction mixture was heated at 95 °C for 1 hour and then lyophilized to give (2S)-2-(3-((S)-1-carboxy-5-(8-(4-(piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex (9.4 mg) as a white solid. MS (ESI), 1302.2 (M+H)⁺.


Example 3. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-morpholino-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0174]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-morpholino-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid.

[0175]




[0176] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL). The reaction was stirred at room temperature for 2 hours and the solvent was then removed using a stream of nitrogen to give a residue. The residue was dissolved in DMSO (4.0 mL) and (S)-di-*tert*-butyl 2-(3-((S)-6-(8-aminooctanamido)-1-(*tert*-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (31.4 mg, 0.05 mmol) and K₂CO₃ (100 mg) were then added to the DMSO solution. The suspension was stirred at room temperature for 2 hours following which morpholine (0.10 mL) was added and the reaction mixture was stirred at room temperature for an additional 14 hours. The reaction mixture was lyophilized to afford a triazine intermediate to which was added TFA (2.0 mL) and DCM (1.0 mL). This mixture was stirred at room temperature for 4 hours to effect deprotection following which the solvent was removed using a stream of nitrogen to give a residue of the crude product. Purification was effected using a Biotage SP4 and a C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-morpholino-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (29.8 mg) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) 7.75-7.65 (m, 3 H), 7.14 (m, 2 H), 6.55 (m, 2 H), 6.33 (d, *J* = 8.0 Hz, 1 H), 6.30 (d, *J* = 8.4 Hz, 1 H), 4.10-1.27 (m, 61 H); MS (ESI), 1133.2 (M+H)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-morpholino-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0177] To solid (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-morpholino-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (10.4 mg, 0.0092 mmol) was added LuCl₃ (1.80 mL, 0.00513 mmol/mL, 0.0092 mmol) and acetonitrile (0.50 mL). The reaction mixture was heated at 95 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-morpholino-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex (9.9 mg) as a white solid. MS (ESI), 1304.9 (M+H)⁺.


Example 4. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-((4-carboxy-1,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-6-(piperazin-1-yl)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0178]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-((4-carboxy-1,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-6-(piperazin-1-yl)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid.

[0179]




[0180] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL). The reaction was stirred at room temperature for 2 hrs. The solvent was then removed using a stream of nitrogen to give a residue, which was dissolved in DMSO (4.0 mL) and (S)-di-*tert*-butyl 2-(3-((S)-6-(8-aminooctanamido)-1-*tert*-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (31.43 mg, 0.05 mmol) and K₂CO₃ (100 mg) were then added to the DMSO solution. The resultant suspension was stirred at room temperature for 2 hrs following which piperazine (100 mg) was added and stirring was continued at room temperature for an additional 16 hrs. The crude reaction was then lyophilized and the triazine intermediate thus obtained was added deprotected using TFA (2.0 mL) and DCM (1.0 mL). Deprotection was carried out by stirring the mixture at room temperature overnight, following which the solvent was removed using a stream of nitrogen to give a residue of the crude product which was purified by Biotage SP4 using a C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-((4-carboxy-1,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-6-(piperazin-1-yl)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (18.9 mg) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) 8.85 (m, 2 H), 7.75-7.65 (m, 4 H), 7.16 (m, 2 H), 6.55 (m, 2 H), 6.32 (d, *J* = 8.8 Hz, 1 H), 6.29 (d, *J* = 8.4 Hz, 1 H), 4.11-1.23 (m, 61 H); MS (ESI), 1132.2 (M+H)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-((4-carboxy-1,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-6-(piperazin-1-yl)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0181] To solid (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-((4-carboxy-1,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-6-(piperazin-1-yl)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid (7.8 mg, 0.0069 mmol) was added LuCl₃ (1.80 mL of a 0.00385 mmol/mL, 0.0069 mmol) and acetonitrile (0.5 mL). The reaction mixture was heated at 95 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-((4-carboxy-1,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-6-(piperazin-1-yl)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex (8.3 mg) as a white solid. MS (ESI), 1303.6 (M+H)⁺.


Example 5. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0182]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido) pentyl)ureido)pentanedioic acid.

[0183]




[0184] DIPEA (0.10 mL) was added to a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocylics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) and mixture was stirred at room temperature for 2 hrs. The solvent was then removed using a stream of nitrogen to give a residue, which was dissolved in DMSO (4.0 mL). (S)-di-*tert*-butyl 2-(3-((S)-6-(8-amino-octanamido)-1-(*tert*-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (31.43 mg, 0.05 mmol) and K₂CO₃ (100 mg) were added to the DMSO solution and the resultant suspension was stirred at room temperature for 2 hrs following which 4-(piperidin-4-yl)butanoic acid (30 mg) was added. After stirring at room temperature for an additional 16 hours the reaction mixture was lyophilized to afford a triazine intermediate which was deprotected using TFA (2.0 mL) and DCM (1.0 mL). After stirring at room temperature overnight the solvent was removed using a stream of nitrogen to give a residue of the titled crude. Purification was effected using Biotage SP4 and a C18 cartridge to obtain (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido) pentyl)ureido)pentanedioic acid (18.8 mg) as a white solid. MS (ESI), 608.8 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido) pentyl)ureido)pentanedioic acid lutetium complex.

[0185] To solid (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido) pentyl)ureido)pentanedioic acid (7.4 mg, 0.006086 mmol) was added LuCl₃ (1.58 mL of a 0.00385 mmol/mL, 0.006086 mmol). The reaction mixture was heated at 95 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(8-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-(4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenylamino)-1,3,5-triazin-2-ylamino)octanamido) pentyl)ureido)pentanedioic acid lutetium complex (9.0 mg) as a white solid. MS (ESI), 1388.8 (M+H)⁺.


Example 6. ((2S,2'S)-2,2'-((((1S,1'S)-((8,8'-((6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl))bis(octanoyl))bis(azanediyl))bis(1-carboxypentane-5,1-diyl))bis(azanediyl))bis(carbonyl))bis(azanediyl))dipentanedioic acid lutetium complex.

[0186]



Step 1. ((2S,2'S)-2,2'-((((1S,1'S)-((8,8'-((6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl))bis(octanoyl))bis(azanediyl))bis(1-carboxypentane-5,1-diyl))bis(azanediyl))bis(carbonyl))bis(azanediyl))dipentanedioic acid.

[0187]



[0188] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL) and the mixture stirred at room temperature for 2 hours. Following stirring, the solvent was removed using a stream of nitrogen to give a residue. This residue was dissolved in DMSO (4.0 mL) and (S)-di-*tert*-butyl 2-(3-((S)-6-(8-amino-octanoyl)-1-(*tert*-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (62.8 mg, 0.10 mmol) and K₂CO₃ (100 mg) were added to the resultant DMSO solution. The suspension thus obtained was stirred at room temperature for 72 hours and then lyophilized to afford a triazine intermediate which was deprotected using TFA (4.0 mL) and DCM (1.0 mL). The TFA/DCM mixture was stirred at room temperature overnight following which the solvent was removed using a stream of nitrogen to afford the titled crude as a solid. The crude was purified by Biotage SP4 using C18 cartridge to give pure ((2S,2'S)-2,2'-((((1S,1'S)-((8,8'-((6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl))bis(octanoyl))bis(azanediyl))bis(1-carboxypentane-5,1-diyl))bis(azanediyl))bis(carbonyl))bis(azanediyl))dipentanedioic acid (10.0 mg) as a white solid. MS (ESI), 753.2 (M/2+H)⁺.


Step 2. ((2S,2'S)-2,2'-((((1S,1'S)-((8,8'-((6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl))bis(octanoyl))bis(azanediyl))bis(1-carboxypentane-5,1-diyl))bis(azanediyl))bis(carbonyl))bis(azanediyl))dipentanedioic acid lutetium complex.

[0189] To solid ((2S,2'S)-2,2'-((((1S,1'S)-((8,8'-((6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl))bis(octanoyl))bis(azanediyl))bis(1-carboxypentane-5,1-diyl))bis(azanediyl))bis(carbonyl))bis(azanediyl))dipentanedioic acid (8.5 mg, 0.005646 mmol) was added LuCl₃

(1.47 mL of a 0.00385 mmol/mL, 0.005646 mmol). The reaction mixture was heated at 70 °C for 1 hour and then lyophilized to give (2S,2'S)-2,2'-((((1S,1'S)-((8,8'-((6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl))bis(octanoyl))bis(azanediyl))bis(1-carboxypentane-5,1-diyl))bis(azanediyl))bis(carbonyl))bis(azanediyl)) dipentanedioic acid lutetium complex (8.6 mg) as a white solid. MS (ESI), 1678.0 (M+H)⁺.


Example 7. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0190]



Step 1. (S)-di-tert-butyl 2-(3-((S)-6-(11-(4-((benzyloxy)carbonyl)piperazin-1-yl)undecanamido)-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate.


[0191]



[0192] A solution of (S)-di-tert-butyl 2-(3-((S)-6-amino-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (1.023 g, 2.097 mmol), 11-(4-((benzyloxy)carbonyl)piperazin-1-yl)undecanoic acid (0.77 g, 1.9059 mmol), EDCI (0.40 g, 2.097 mmol), HOBt (0.27 g, 2.097 mmol) and DIPEA (1.0 mL) in dichloroethane (DCE; 25 mL) was stirred at room temperature overnight. The following day, the solvent was evaporated to give a residue, which was purified using Biotage column chromatography and a mixture of DCM/MeOH as the eluant to give (S)-di-tert-butyl 2-(3-((S)-6-(11-(4-((benzyloxy)carbonyl)piperazin-1-yl)undecanamido)-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (1.52 g, 91%) as a yellowish solid. MS (ESI), 874.3 (M+H)⁺.

Step 2. (S)-di-tert-butyl 2-(3-((S)-1-(tert-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate.

[0193]




[0194] To a solution of (S)-di-tert-butyl 2-(3-((S)-6-(11-(4-((benzyloxy)carbonyl)piperazin-1-yl)undecanamido)-1-(tert-butoxy)-1-oxohexan-2-yl)ureido)pentanedioate (1.50 g, 1.72 mmol) and ammonium formate (1.0 g) in ethanol (60 mL) was added palladium on carbon (300 mg). The reaction mixture was stirred at room temperature for overnight and filtered

through a pad of celite followed by washing of the celite pad using ethyl acetate (EtOAc). The solvent was removed under reduced pressure and the residue dissolved in dichloromethane (DCM). The DCM solution was washed using saturated sodium bicarbonate and then partitioned to separate the organic layer from the aqueous layer. Concentration of the organic layer under reduced pressure afforded the titled product as a yellowish solid (1.2345 g, 97 % yield). MS (ESI), 740.4 (M+H)⁺.

Step 3. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0195]




[0196] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL) and resultant mixture was stirred at room temperature for 2 hrs. Following stirring, the solvent was removed under a stream of nitrogen to give a residue, which was dissolved in DMSO (1.0 mL) prior to the addition of (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (34 mg, 0.05 mmol) and K₂CO₃ (50 mg). The resultant suspension was stirred at room temperature for 2 hrs, following which a tetrahydrofuran solution of dimethylamine (0.2 mL, 2.0 M in THF) was added. After additional stirring of the reaction mixture at room temperature for 16 hours, the reaction was lyophilized to afford the crude triazine intermediate. Deprotection of the crude using TFA (2.0 mL) and DCM (1.0 mL) was carried out at room temperature overnight. The following day, the solvent was removed under a stream of nitrogen to give a residue, which was purified by Biotage SP4 using C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (24 mg) as a white solid. MS (ESI), 601.2 (M/2+H)⁺.

Step 4. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0197] To solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (9.4 mg, 0.00783 mmol) was added LuCl₃ (1.02 mL of a 0.00770 mmol/mL, 0.00783 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (11.1 mg) as a white solid. MS (ESI), 1373.7 (M+H)⁺.


Example 8. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0198]



Step 1. ((2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0199]




[0200] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 ml) and the solution was stirred at room temperature for 2 hrs. The solvent was then removed using a stream of nitrogen to give a residue, which was dissolved in DMSO (1.0 mL) prior to the addition of piperidine (4.25 mg, 0.05 mmol). The resultant suspension was stirred at room temperature for 2 hours following which (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) and K₂CO₃ (50 mg) were added to the DMSO solution. After additional stirring at room temperature for 16 hours and the mixture was lyophilized to afford the crude triazine intermediate, which was deprotected using TFA (2.0 mL) and DCM (1.0 mL). Deprotection was carried out by stirring the crude at room temperature overnight and the following day the solvent was removed using a stream of nitrogen to give a residue which was purified by Biotage SP4 using a C18 cartridge to give ((2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (22 mg) as a white solid. MS (ESI), 621.2 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0201] To solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (12.4 mg, 0.01 mmol) was added LuCl₃ (1.30 mL of a 0.00770 mmol/mL, 0.01 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (14.0 mg) as a white solid. MS (ESI), 1413.7 (M+H)⁺.


Example 9. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(2-(2-(2-carboxyethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0202]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((2-(2-(2-carboxyethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0203]




[0204] To a DCM solution (2.0 mL) of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclus) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) was added DIPEA (0.10 mL). After stirring at room temperature for 2 hours the solvent was removed under a stream of nitrogen to give a residue, which was dissolved in DMSO (1.0 mL). Tert-butyl 3-(2-(2-aminoethoxy)ethoxy)propanoate (11.67 mg, 0.05 mmol) and K₂CO₃ (50 mg) were then added to the DMSO solution and the resultant suspension was stirred at room temperature for 2 hours. (S)-di-tert-butyl 2-(3-((S)-1-(tert-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) was then added. After stirring for 16 hours the reaction mixture was lyophilized to afford the crude triazine intermediate which was deprotected using TFA (2.0 mL) and DCM (1.0 mL). Deprotection was carried out by stirring the crude at room temperature overnight and the following day the solvent was removed using a stream of nitrogen to give a residue which was purified by Biotage SP4 using C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((2-(2-(2-carboxyethoxy)ethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (29.4 mg) as a white solid. MS (ESI), 667.2 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((2-(2-(2-carboxyethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0205] To solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((2-(2-(2-carboxyethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (13.1 mg, 0.01 mmol) was added LuCl₃ (1.30 mL of a 0.00770 mmol/mL, 0.01 mmol). The reaction mixture was heated at 90 °C for 1 hour and lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((2-(2-(2-carboxyethoxy)ethoxy)ethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (14.5 mg) as a white solid. MS (ESI), 1505.7 (M+H)⁺.


Example 10. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((26-carboxy-3,6,9,12,15,18,21,24-octaoxaheptacosyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0206]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((26-carboxy-3,6,9,12,15,18,21,24-octaoxaheptacosyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0207]



[0208] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL). The reaction was stirred at room temperature for 2 hrs and the solvent removed following stirring using a stream of nitrogen. The residue thus obtained was dissolved in DMSO (1.0 mL) and 1-amino-3,6,9,12,15,18,21,24-octaoxaheptacosan-27-oic acid (22.1 mg, 0.05 mmol) and K₂CO₃ (50 mg) were added to the DMSO solution. The resultant suspension was stirred at room temperature for 2 hrs following which (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) was then added. After stirring for an additional 16 hours at room temperature the crude reaction was lyophilized to afford the triazine intermediate, which was deprotected overnight at room temperature using TFA (2.0 mL) and DCM (1.0 mL). The crude product was purified by Biotage SP4 using a C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((26-carboxy-3,6,9,12,15,18,21,24-octaoxaheptacosyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (31.4 mg) as a white solid. MS (ESI), 799.3 (M/2+H)⁺.


Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((26-carboxy-3,6,9,12,15,18,21,24-octaoxaheptacosyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0209] LuCl₃ (0.69 mL of a 0.00770 mmol/mL, 0.00532 mmol) was added to solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((26-carboxy-3,6,9,12,15,18,21,24-octaoxaheptacosyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (8.5 mg, 0.00532 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-((26-carboxy-3,6,9,12,15,18,21,24-octaoxaheptacosyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (8.2 mg) as a white solid. MS (ESI), 885.2 (M/2+H)⁺.


Example 11. (2S)-2-(3-((1S)-5-(11-(4-(4-(((S)-5-bis((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)amino)-1-carboxypentyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex.

[0210]



Step 1. (2S)-2-(3-((1S)-5-(11-(4-(4-(((S)-5-bis((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)amino)-1-carboxypentyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid.

[0211]



[0212] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL). After stirring at room temperature for 2 hours the solvent was removed using a stream of nitrogen gas to give a residue. This residue was dissolved in DMSO (1.0 mL) and (S)-2-amino-6-bis((1-(2-(tert-butoxy)-2-oxoethyl)-1H-imidazol-2-yl)methyl)amino)hexanoic acid (26.7 mg, 0.05 mmol) and K₂CO₃ (50 mg) were then added. The resultant suspension was stirred at room temperature overnight. The following day (S)-di-tert-butyl 2-(3-((S)-1-(tert-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) was added and the reaction mixture was stirred at room temperature for an additional 24 hours. Lyophilization afforded the crude triazine intermediate which was deprotected overnight at room temperature using TFA (3.0 mL) and DCM (1.0 mL). The deprotected crude final product was purified by Biotage SP4 using a C18 cartridge to give (2S)-2-(3-((1S)-5-(11-(4-(4-(((S)-5-bis((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)amino)-1-carboxypentyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid (41.5 mg) as a


white solid. MS (ESI), 789.6 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-5-(11-(4-(4-(((S)-5-(bis((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)amino)-1-carboxypentyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex.

[0213] To solid (2S)-2-(3-((1S)-5-(11-(4-(4-(((S)-5-(bis((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)amino)-1-carboxypentyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid (16.3 mg, 0.0103 mmol) was added LuCl₃ (1.0 mL, 0.0103 mmol/mL, 0.0103 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-5-(11-(4-(4-(((S)-5-(bis((1-(carboxymethyl)-1H-imidazol-2-yl)methyl)amino)-1-carboxypentyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex (15.7 mg) as a white solid. MS (ESI), 875.6 (M/2+H)⁺.


Example 12. (2S)-2-(3-((1S)-5-(11-(4-(4-(bis(carboxymethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex.

[0214]



Step 1. (2S)-2-(3-((1S)-5-(11-(4-(4-(bis(carboxymethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid.

[0215]




[0216] To a DCM (2.0 mL) solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocylics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) was added DIPEA (0.10 mL) and resultant mixture was stirred at room temperature for 2 hrs. Removal of the solvent using a stream of nitrogen gave a residue which was dissolved in DMSO (1.0 mL) prior to the addition of di-*tert*-butyl 2,2'-azanediyl diacetate (24.5 mg, 0.10 mmol) and K₂CO₃ (50 mg) were added. The resultant suspension was stirred at room temperature for overnight and the following day (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxy-6-(1-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) was added and the stirring continued at room temperature for 24 hours. Lyophilization of this suspension afforded the triazine intermediate, which was deprotected at room temperature overnight using TFA (3.0 mL) and DCM (1.0 mL). The deprotected crude product was purified by Biotage SP4 using a C18 cartridge to give (2S)-2-(3-((1S)-5-(11-(4-(4-(bis(carboxymethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid (27.0 mg) as a white solid. MS (ESI), 645.2 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-5-(11-(4-(4-(bis(carboxymethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex.

[0217] LuCl_3 (0.89 mL of a 0.0103 mmol/mL, 0.00915 mmol) was added to solid reagent of (2S)-2-(3-((1S)-5-(11-(4-(4-(bis(carboxymethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid (11.8 mg, 0.00915 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-5-(11-(4-(4-(bis(carboxymethyl)amino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex (12.0 mg) as a white solid. MS (ESI), 731.2 ($M/2+H$)⁺.


Example 13. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(methylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0218]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(methylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0219]




[0220] To a DCM (2.0 mL) solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) was added DIPEA (0.10 mL) and the solution stirred at room temperature for 2 hours. After stirring the solvent was removed using a stream of nitrogen gas to give a residue. This residue was dissolved in DMSO (1.0 mL) and the solution was contacted with methanamine (0.10 mL, 2.0 M in THF) and K₂CO₃ (50 mg). The resultant suspension was stirred at room temperature for 4 hours. (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) was added then added to the DMSO solution and the reaction mixture was stirred at room temperature for an additional 24 hrs prior to lyophilization to afford the crude triazine intermediate. Deprotection using TFA (3.0 mL) and DCM (1.0 mL) at room temperature, overnight followed by removal of the solvent using a stream of nitrogen gave crude product which was purified by Biotage SP4 using a C18 cartridge to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(methylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (10.8 mg) as a white solid. MS (ESI), 594.2 ($M/2+H$)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(methylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0221] To solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(methylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid, (7.7 mg, 0.00649 mmol) was added LuCl_3 (0.63 mL of a 0.0103 mmol/mL, 0.00649 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(methylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (7.9 mg) as a white solid. MS (ESI), 680.2 ($M/2+H$)⁺.


Example 14. (2S)-2-(3-((1S)-5-(11-(4-(4-(4-(3-aminopropyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex.

[0222]



Step 1. (2S)-2-(3-((1S)-5-(11-(4-(4-(4-(3-aminopropyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid.

[0223]



[0224] To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics) (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL) and the solution was stirred at room temperature for 2 hours. After stirring the solvent was removed under a stream of nitrogen to give a residue which was dissolved in DMSO (1.0 mL) prior to the addition of (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) and K_2CO_3 (50 mg). The resultant suspension was stirred at room temperature for 2 hours and 3-(piperazin-1-yl)propan-1-amine (47 mg) was then added following which the reaction mixture was stirred for an additional 16 hours at room temperature. Lyophilization after 16 hours afforded the crude triazine intermediate which was deprotected at room temperature, overnight using TFA (2.0 mL) and DCM (1.0 mL). The deprotected product was purified by Biotage SP4 using a C18 cartridge to give (2S)-2-(3-((1S)-5-(11-(4-(4-(4-(3-aminopropyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phe-


nyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid (25 mg) as a white solid. MS (ESI), 650.3 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-5-(11-(4-(4-(4-(3-aminopropyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex.

[0225] To solid (2S)-2-(3-((1S)-5-(11-(4-(4-(4-(3-aminopropyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid (10.7 mg, 0.00824 mmol) was added LuCl₃ (0.80 mL of a 0.0103 mmol/mL, 0.00824 mmol). The reaction mixture was heated at 90 °C for 1 hour and then lyophilized to give (2S)-2-(3-((1S)-5-(11-(4-(4-(4-(3-aminopropyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)-1-carboxypentyl)ureido)pentanedioic acid lutetium complex (10.2 mg) as a white solid. MS (ESI), 736.2 (M/2+H)⁺.


Example 15. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(carboxymethyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0226]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(carboxymethyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0227]



[0228] To a DCM solution (2.0 mL) of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics), (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) was added DIPEA (0.10 mL) the resultant solution was stirred at room temperature for 2 hours. After stirring, the solvent was removed using a stream of nitrogen to give a residue which was dissolved in DMSO (1.0 mL) prior to the addition of (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) and K₂CO₃ (50 mg). The suspension thus obtained was stirred at room temperature for 2 hrs and *tert*-butyl 2-(piperazin-1-yl)acetate (50 mg) was then added to the reaction mixture and stirring was continued at room temperature for an additional 16 hours. Lyophilization of the reaction mixture at the end of 16 hours afforded a residue of the protected final product. This residue was contacted with TFA (2.0 mL) and DCM (1.0 mL) at room temperature overnight to cause removal of protecting groups, following which the solvent was removed under a stream of nitrogen to give crude deprotected product that was purified by Biotage SP4 using a C18 cartridge. The titled compound (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(carboxymethyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.


6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (14 mg) as obtained as a white solid. MS (ESI), 650.8 (M/2+H)⁺.

- 5 Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(carboxymethyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

10 **[0229]** To solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(carboxymethyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (6.0 mg, 0.00426 mmol) was added LuCl₃ (0.45 mL of a 0.0103 mmol/mL, 0.00462 mmol). The reaction mixture was heated at 90 °C for 1 hour and lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(carboxymethyl)piperazin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (5.6 mg) as a white solid. MS (ESI), 736.8 (M/2+H)⁺.


Example 16. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0230]



Step 1. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0231]



50 **[0232]** To a solution of *p*-NH₂-Bn-DOTA-tetra(*t*-Bu-ester) (Macrocyclics), (42.4 mg, 0.050 mmol) and cyanuric chloride (9.2 mg, 0.050 mmol) in DCM (2.0 mL) was added DIPEA (0.10 mL). Following stirring at room temperature for 2 hrs, the solvent was removed using a stream of nitrogen to give a residue which was dissolved in DMSO (1.0 mL) prior to the addition of (S)-di-*tert*-butyl 2-(3-((S)-1-(*tert*-butoxy)-1-oxo-6-(11-(piperazin-1-yl)undecanamido)hexan-2-yl)ureido)pentanedioate (37 mg, 0.05 mmol) and K₂CO₃ (50 mg). The suspension formed was stirred at room temperature for 2 hrs and 4-(piperidin-4-yl)butanoic acid (160 mg) was then added to the suspension. After continuous stirring at room temperature for 72 hrs, the reaction was stopped by lyophilization to afford the protected triazine compound. Deprotection at room temperature, overnight using TFA (4.0 mL) and DCM (1.0 mL), followed by purification using Biotage SP4 and


a C18 cartridge gave (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (15.3 mg) as a white solid. MS (ESI), 650.8 (M/2+H)⁺.

Step 2. (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex.

[0233] To solid (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid (6.9 mg, 0.00520 mmol) was added LuCl₃ (0.50 mL, 0.0103 mmol/mL, 0.00520 mmol). The reaction mixture was heated at 90 °C for 1 hour and lyophilized to give (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(4-(3-carboxypropyl)piperidin-1-yl)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid lutetium complex (7.9 mg) as a white solid. MS (ESI), 750.2 (M/2+H)⁺.

Example 17. ⁶⁸Ga Labeling of (2S)-2-(3-((1S)-1-carboxy-5-(11-(4-(4-(dimethylamino)-6-((4-((1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecan-2-yl)methyl)phenyl)amino)-1,3,5-triazin-2-yl)piperazin-1-yl)undecanamido)pentyl)ureido)pentanedioic acid.

[0234]



[0235] ⁶⁸Ga was synthesized using a gallium-68 generator (IDB Holland). A 1 mL fraction of the generator eluate (eluted using 0.6 M HCl suprapure) containing the highest ⁶⁸Ga activity was mixed with the reaction mixture that containing 2 µL of the target compound (10 mM solution in DMSO) and 10 µL of ascorbic acid (20% in water). The pH of the reaction mixture was adjusted to be in the pH range of 3.6 - 3.9 by the addition of approximately 290 µL of an aqueous solution of sodium acetate (2.5 M in water).

[0236] The mixture was heated at 90 °C for 10 minutes with stirring. A test sample of the reaction mixture was analyzed by HPLC to confirm complete complexation. The reaction mixture was then diluted with 2 mL saline (0.9% sodium chloride) and loaded onto a pre-conditioned Plea Cartridge (60 mg, Varian, Bond Elut Plexa). The cartridge was rinsed with 2 mL saline prior to elution of the desired complex using 0.5 mL ethanol. The eluent was passed through a sterile filter (Millipore, Millex-GV) fitted to a syringe followed by washing of the filter by passing 5 mL of saline and 200 µL of phosphate buffer.

[0237] The radio-labelled compound was analyzed by HPLC on a Chromolith Performance RP-18e column (100 x 3 mm Merck KGaA, Darmstadt, Germany) using a linear gradient from 0% to 100% acetonitrile in water (both containing 0.1% TFA) over 5 min. UV absorbance was detected at 214 nm. Under these conditions ⁶⁸Ga-MIP-1558 is eluted at about 2.25 min. The radiochemical yields ranged from 77% - 97%, average RCP = 87% (data corrected for radioactive decay).

EQUIVALENTS

[0238] While certain embodiments have been illustrated and described, it should be understood that changes and modifications can be made therein in accordance with ordinary skill in the art without departing from the technology in its broader aspects as defined in the following claims.

[0239] The present disclosure is not to be limited in terms of the particular embodiments described in this application. The present disclosure is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled. It is to be understood that this disclosure is not limited to particular methods, reagents, compounds compositions or biological systems, which can of course vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.


[0240] In addition, where features or aspects of the disclosure are described in terms of Markush groups, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group.

[0241] As will be understood by one skilled in the art, for any and all purposes, particularly in terms of providing a written description, all ranges disclosed herein also encompass any and all possible subranges and combinations of subranges thereof. Any listed range can be easily recognized as sufficiently describing and enabling the same range being broken down into at least equal halves, thirds, quarters, fifths, tenths, etc. As a non-limiting example, each range discussed herein can be readily broken down into a lower third, middle third and upper third, etc. As will also be understood by one skilled in the art all language such as "up to," "at least," "greater than," "less than," and the like, include the number recited and refer to ranges which can be subsequently broken down into subranges as discussed above. Finally, as will be understood by one skilled in the art, a range includes each individual member, including the first and last number listed for the range.

[0242] Other embodiments are set forth in the following claims.

Claims

1. A compound according to formula I




wherein:

A is $(\text{CHR}^1)_m$ or $\text{C}(\text{O})$;


W is $\text{C}(\text{O})-(\text{CH}_2)_p$ -, $-\text{C}(\text{O})[\text{CH}_2\text{CH}_2\text{O}]_n$ -, $-\text{[CH}_2\text{CH}_2\text{O}]_n(\text{CH}_2)_2$ -, $-\text{C}(\text{O})[\text{CH}(\text{R}^3)]_q$ -, $-(\text{CH}_2)_m\text{O}-(\text{CH}_2)_n$ -, $-(\text{CH}_2)_m\text{S}-(\text{CH}_2)_n$ -, $-(\text{CH}_2)_m\text{S}(\text{O})-(\text{CH}_2)_n$ -, $-(\text{CH}_2)_m\text{S}(\text{O})_2-(\text{CH}_2)_n$ -, or $-(\text{CH}_2)_m\text{NR}_a-(\text{CH}_2)_n$ -,

Y is $-\text{NH}-$, $-\text{NR}^2$ -, or



X is $-(\text{C}_1\text{-C}_{10})\text{alkylene}-(\text{C}_3\text{-C}_{10})\text{arylene}$ -, $-(\text{C}_3\text{-C}_{10})\text{arylene}$ -, $-(\text{C}_3\text{-C}_{10})\text{arylene}-(\text{C}_1\text{-C}_{10})\text{alkylene}$ -, phenylene, $-(\text{C}_1\text{-C}_{10})\text{alkylene}-(\text{C}_3\text{-C}_{10})\text{cycloalkylene}$ -, $-(\text{C}_3\text{-C}_{10})\text{cycloalkylene}$ -, or $-(\text{C}_3\text{-C}_{10})\text{cycloalkylene}-(\text{C}_1\text{-C}_{10})\text{alkylene}$ -, R^1 and R^2 are each independently H, $-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, $-\text{C}(\text{O})-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, benzyl, $-(\text{C}_3\text{-C}_{10})\text{cycloalkyl}$ -, or $-(\text{C}_3\text{-C}_{10})\text{aryl}$;

R^a and R^b are each independently H, $-\text{OH}$ -, $-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, $-\text{[CH}_2\text{CH}_2\text{O}]_n(\text{CH}_2)_2$ -, $-\text{C}(\text{O})-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, $-(\text{C}_1\text{-C}_{10})\text{alkylene}-\text{C}(\text{O})$ -, $-(\text{C}_1\text{-C}_{10})\text{alkylene}-\text{C}(\text{O})-\text{Z}$ -, benzyl, $-(\text{C}_3\text{-C}_{10})\text{cycloalkyl}$ -, $-(\text{C}_3\text{-C}_{10})\text{aryl}-(\text{C}_1\text{-C}_{10})\text{alkylene}$ -, $-(\text{C}_3\text{-C}_{10})\text{aryl}$ -, halo- $-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, hydroxy- $-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, $-\text{NH}-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -, or $-(\text{C}_1\text{-C}_{10})\text{alkylene}-\text{NR}^d\text{Re}$ -, or R^a and R^b together with the nitrogen to which they are bonded form a $(\text{C}_3\text{-C}_6)$ -heteroaryl or $(\text{C}_3\text{-C}_6)$ -heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O; Z is $-\text{OH}$ -, $-\text{O}-(\text{C}_1\text{-C}_{10})\text{alkyl}$ -,



R^c is -OH, -O(C₁-C₁₀)alkyl, -Obenzyl, -O(C₃-C₁₀)cycloalkyl, -O(C₃-C₁₀)aryl, -O-(C₁-C₁₀)alkylene-(C₃-C₁₀)aryl, or -O-(C₁-C₁₀)alkylene-(C₃-C₁₀)cycloalkyl,


R^3 is H, halogen, -OH, -NH₂, -(CH₂)_p-COOH, or -(CH₂)_p-NH₂;


T is -H, -OH, -COOH, or -NR^dR^e;

R^d and R^e are each independently H, bond, -OH, -(C₁-C₁₀)alkyl, or -(C₃-C₁₀)heteroaryl-(C₁-C₁₀)alkylene;

m, n, p, q, t and r are each independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; and


D is






wherein any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene is optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of $-(\text{C}_1\text{-C}_{10})\text{alkyl}$, $-(\text{C}_1\text{-C}_{10})\text{haloalkyl}$, $-(\text{C}_1\text{-C}_{10})\text{aminoalkyl}$, $-(\text{C}_1\text{-C}_{10})\text{alkylene-COOH}$, $-(\text{C}_1\text{-C}_{10})\text{hydroxyalkyl}$, $-\text{OH}$, halogen, $-\text{NH}_2$, $-\text{COOH}$, $-\text{C}(\text{O})-(\text{C}_1\text{-C}_{10})\text{alkyl}$, $-(\text{C}_1\text{-C}_{10})\text{alkylene-C}(\text{O})-$, $-(\text{C}_1\text{-C}_{10})\text{alkylene-C}(\text{O})-\text{X}$, $-\text{NH}-(\text{C}_1\text{-C}_{10})\text{alkyl}$, and $-(\text{C}_1\text{-C}_{10})\text{alkylene-NR}^d\text{R}^e$, and $-\text{NR}^d\text{R}^e$.

2. The compound of Claim 1, wherein X is phenylene, r is 1 and D is



3. The compound of Claim 2, wherein the compound is a compound according to Formula II




wherein:

A is $(\text{CHR}^1)_m$ or $\text{C}(\text{O})$;

W is selected from the group consisting of $-\text{C}(\text{O})-(\text{CH}_2)_p-$, $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$, $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$, $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q-$, $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$, $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$, $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$, $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$, and $-(\text{CH}_2)_m-\text{NR}^a-(\text{CH}_2)_n-$.

Y is selected from $-\text{NH}-$, $-\text{NR}^2-$,





R^1 and R^2 are each independently selected from H, $-(\text{C}_1\text{-C}_{10})\text{alkyl}$, $-\text{C}(\text{O})-(\text{C}_1\text{-C}_{10})\text{alkyl}$, benzyl, $-(\text{C}_3\text{-C}_{10})\text{cy-}$

cloalkyl, or $-(C_3-C_{10})$ aryl;

R^a and R^b are each independently selected from the group consisting of H, -OH, $-(C_1-C_{10})$ alkyl, $-[CH_2-CH_2-O]_n-(CH_2)_2-T$, $-C(O)-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ alkylene- $C(O)-$, $-(C_1-C_{10})$ alkylene- $C(O)-Z$, benzyl, $-(C_3-C_{10})$ cycloalkyl, $-(C_3-C_{10})$ aryl- (C_1-C_{10}) alkylene, $-(C_3-C_{10})$ aryl, halo- (C_1-C_{10}) alkyl, hydroxy- (C_1-C_{10}) alkyl, $-NH-(C_1-C_{10})$ alkyl, and $-(C_1-C_{10})$ alkylene- NR^dR^e , or R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heteroaryl or (C_3-C_6) -heterocycloalkyl that can further comprise one or more heteroatoms selected from N, S, or O;

Z is selected from -OH, $-O(C_1-C_{10})$ alkyl,



R^c is selected from -OH, $-O(C_1-C_{10})$ alkyl, $-Obenzyl$, $-O(C_3-C_{10})$ cycloalkyl, $-O(C_3-C_{10})$ aryl, $-O(C_1-C_{10})$ alkylene- $-(C_3-C_{10})$ aryl, or $-O(C_1-C_{10})$ alkylene- $-(C_3-C_{10})$ cycloalkyl,

R^3 is selected from H, halogen, -OH, $-NH_2$, $-(CH_2)_p-COOH$, or $-(CH_2)_p-NH_2$;

T is selected from -H, -OH, $-COOH$, or $-NR^dR^e$;

R^d and R^e are each independently selected from H, bond, -OH, $-(C_1-C_{10})$ alkyl, or $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene;


m, n, p, q, t and x are each independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

wherein any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene is optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of $-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ haloalkyl, $-(C_1-C_{10})$ aminoalkyl, $-(C_1-C_{10})$ alkylene- $COOH$, $-(C_1-C_{10})$ hydroxyalkyl, $-NH_2$, $-COOH$, $-C(O)-(C_1-C_{10})$ alkyl, $-(C_1-C_{10})$ alkylene- $C(O)-$, $-(C_1-C_{10})$ alkylene- $C(O)-X$, $-NH-(C_1-C_{10})$ alkyl, and $-(C_1-C_{10})$ alkylene- NR^dR^e , and $-NR^dR^e$.

4. The compound of Claim 3, wherein A is $(CHR^1)_m$ and W is $-C(O)-(CH_2)_p-$.

5. The compound of Claim 4, wherein R^1 is hydrogen and m is 2.

6. The compound of Claim 3, wherein Y is -NH- or




7. The compound of Claim 3, wherein R^a and R^b are each independently hydrogen or methyl and R^c is -OH.


8. The compound of Claim 3, wherein R^a and R^b together with the nitrogen to which they are bonded form a (C_3-C_6) -heterocycloalkyl selected from piperidine, piperazine, morpholine, thiomorpholine, isothiazolidine, isoxazolidine, pyrrolidine, imidazolidine, thiazolidine or oxazolidine.

9. The compound of Claim 8, wherein the (C_3-C_6) -heterocycloalkyl is piperidine or 4-(piperidin-4-yl)butanoic acid.


10. The compound according to claim 8, wherein R^d and R^e are each independently $-(C_3-C_{10})$ heteroaryl- (C_1-C_{10}) alkylene.

11. The compound of Claim 3, which is selected from the following compounds:






5




10




15

20



25

30



35

40

45

50

55

5

10

15

20

25

30


35


40

45

50

55






15


20

25



40


45




50

55

67



or



R^c is selected from -OH, -O(C₁-C₁₀)alkyl, -Obenzyl, -O(C₃-C₁₀)cycloalkyl, -O(C₃-C₁₀)aryl, -O-(C₁-C₁₀)alkylene-(C₃-C₁₀)aryl, or -O-(C₁-C₁₀)alkylene-(C₃-C₁₀)cycloalkyl,

R^3 is selected from H, halogen, -OH, -NH₂, -(CH₂)_p-COOH, or -(CH₂)_p-NH₂;

T is selected from -H, -OH, -COOH, or -NR^dR^e;


R^d and R^e are each independently selected from H, bond, -OH, -(C₁-C₁₀)alkyl, or -(C₃-C₁₀)heteroaryl-(C₁-C₁₀)alkylene;


m, n, p, q, t and x are each independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

wherein any alkyl, alkylene, aryl, arylene, heteroaryl, heteroarylene, cycloalkyl, cycloalkylene, heterocycloalkyl, or heterocycloalkylene is optionally substituted with 1, 2, or 3 substituent groups selected from the group consisting of -(C₁-C₁₀)alkyl, -(C₁-C₁₀)haloalkyl, -(C₁-C₁₀) aminoalkyl, -(C₁-C₁₀)alkylene-COOH, -(C₁-C₁₀)hydroxyalkyl, -NH₂, -COOH, -C(O)-(C₁-C₁₀)alkyl, -(C₁-C₁₀)alkylene-C(O)-, -(C₁-C₁₀)alkylene-C(O)-X, -NH-(C₁-C₁₀)alkyl, and -(C₁-C₁₀)alkylene-NR^dR^e-, and -NR^dR^e;


the radionuclide is selected from the group consisting of ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ⁶⁴Cu, ¹⁵³Gd, ¹⁵⁵Gd, ¹⁵⁷Gd, ⁵⁹Fe, ²²⁵Ac, ²¹²Bi, ²¹³Bi, ⁵⁵Co, ⁶⁷Cu, ¹⁶⁵Dy, ¹⁶⁶Ho, ¹⁹²Ir, ²²³Ra, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁰⁵Rh, ²¹²Pb, ²¹³Pb, ¹⁴⁹Tb, ²²⁷Th, ¹⁵³Sm, ⁸⁹Sr, ^{117m}Sn, ¹⁶⁹Yb, ⁹⁰Y, ⁸⁶Y, ⁸⁹Zr and ¹⁷⁷Lu.


14. The metal complex of Claim 13, which is:















or a pharmaceutically acceptable salt or solvate thereof.

15. The metal complex of Claim 13, which is:














or a pharmaceutically acceptable salt or solvate thereof.

16. A pharmaceutical composition comprising the compound of Claim 3 or the metal complex of Claim 13, or a pharmaceutically acceptable salt, solvate, or ester thereof; and a pharmaceutically acceptable carrier.
17. The compound of any one of claims 1 to 11, for use in a method of obtaining a radiographic image of one or more tissues that express prostate-specific membrane antigen (PSMA), said method comprising the following steps:
 - a. contacting one or more tissues that express PSMA with a metal complex comprising a radionuclide and said compound or a pharmaceutically acceptable salt, solvate, or ester thereof; and
 - b. obtaining a radiographic image of the one or more tissues.

18. The metal complex of any one of claims 12 to 15, for use in a method of obtaining a radiographic image of one or more tissues that express prostate-specific membrane antigen (PSMA), said method comprising the following steps:


- a. contacting one or more tissues that express PSMA with said metal complex or a pharmaceutically acceptable salt, solvate, or ester thereof; and
- b. obtaining a radiographic image of the one or more tissues.

19. The compound for use of claim 17 or the metal complex for use of claim 18, wherein the one or more tissues are prostate tissue or prostate cancer tissue.

20. The compound for use of claim 17 or the metal complex for use of claim 18, wherein the radionuclide is selected from the group consisting of ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{225}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , ^{117}mSn , ^{169}Yb , ^{90}Y , ^{86}Y , ^{89}Zr and ^{177}Lu :

Patentansprüche

1. Eine Verbindung gemäß der Formel I




wobei:

A $(\text{CHR}^1)_m$ oder C(O) ist;

W $-\text{C}(\text{O})-(\text{CH}_2)_p-$; $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$; $-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$; $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q-$; $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$; oder $-(\text{CH}_2)_m-\text{NR}_a-(\text{CH}_2)_n-$ ist,

Y $-\text{NH}-$, $-\text{NR}^2-$ oder




ist;

X $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en- $(\text{C}_3-\text{C}_{10})\text{Aryl}$ en-, $-(\text{C}_3-\text{C}_{10})\text{Aryl}$ en-, $-(\text{C}_3-\text{C}_{10})\text{Aryl}$ en- $(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en-, Phenylen-, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en- $(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$ en-, $-(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$ en-, oder $-(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$ en- $(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en- ist;

R^1 und R^2 jeweils unabhängig H-, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, Benzyl-, $-(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$ -, oder $-(\text{C}_3-\text{C}_{10})\text{Aryl}$ sind;

R^a und R^b jeweils unabhängig H-, $-\text{OH}$ -, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en- $\text{C}(\text{O})-$ -, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en- $\text{C}(\text{O})-\text{Z}$ -, Benzyl-, $-(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$ -, $-(\text{C}_3-\text{C}_{10})\text{Aryl}$ -($\text{C}_1-\text{C}_{10})\text{Alkyl}$ en-, $-(\text{C}_3-\text{C}_{10})\text{Aryl}$ -, Halogen- $(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, Hydroxy- $(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, $-\text{NH}-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -, oder $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$ en- NR^d - R^e - sind, oder R^a und R^b zusammen mit dem Stickstoff, an den sie gebunden sind, ein (C_3-C_6) -Heteroaryl oder (C_3-C_6) -Heterocycloalkyl bilden, die weiterhin ein oder mehrere Heteroatome ausgewählt aus N, S, oder O umfassen können;

Z $-\text{OH}$ -, $-\text{O}(\text{C}_1-\text{C}_{10})\text{Alkyl}$ -,



, oder

ist;

R^c -OH, -O(C₁-C₁₀)Alkyl, -OBenzyl, -O(C₃-C₁₀)Cycloalkyl, -O(C₃-C₁₀)Aryl, -O-(C₁-C₁₀)Alkyl-(C₃-C₁₀)Aryl, oder -O-(C₁-C₁₀)Alkyl-(C₃-C₁₀)Cycloalkyl ist;




R³ H, Halogen, -OH, -NH₂, -(CH₂)_p-COOH, oder -(CH₂)_p-NH₂ ist;


T -H, -OH -COOH oder -NR^dR^e ist;

R^d und R^e jeweils unabhängig H, Bindung, -OH, -(C₁-C₁₀)Alkyl, oder -(C₃-C₁₀)Heteroaryl-(C₁-C₁₀)Alkyl sind;

m, n, p, q, t und r jeweils unabhängig 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 oder 10 sind; und

D ist






H₃ [5-MeOsal]₃ TAME] , oder


Punkt der
Anbindung
an den Linker

wobei jedes Alkyl, Alkylen, Aryl, Arylen, Heteroaryl, Heteroarylen, Cycloalkyl, Cycloalkylen, Heterocycloalkyl, oder Heterocycloalkylen wahlweise substituiert ist mit 1, 2, oder 3 Substituentengruppen ausgewählt aus der Gruppe bestehend aus -(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Halogenalkyl, -(C₁-C₁₀)Aminoalkyl, -(C₁-C₁₀)Alkylen-CO-OH, -(C₁-C₁₀)Hydroxyalkyl, -OH, Halogen, -NH₂, -COOH, -C(O)-(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Alkylen-C(O)-, -(C₁-C₁₀)Alkylen-C(O)-X, -NH-(C₁-C₁₀)Alkyl, und -(C₁-C₁₀)Alkylen-NR^dRe, und -NR^dRe.

2. Die Verbindung aus Anspruch 1, wobei X Phenyl ist, r 1 ist und D ist



3. Die Verbindung aus Anspruch 2, wobei die Verbindung eine Verbindung gemäß der Formel II ist




II

wobei:

A (CHR¹)_m oder C(O) ist;

W ausgewählt ist aus der Gruppe bestehend aus -C(O)-(CH₂)_p-; -C(O)-[CH₂-CH₂-O]_n-, -[CH₂-CH₂-O]_n-(CH₂)₂-, -C(O)-[CH(R³)]_lq-, -(CH₂)_m-O-(CH₂)_n-, -(CH₂)_m-S-(CH₂)_n-, -(CH₂)_m-S(O)-(CH₂)_n-, -(CH₂)_m-S(O)₂-(CH₂)_n- und -(CH₂)_m-NR^a-(CH₂)_n-,

Y ausgewählt ist aus -NH-, -NR²-,




ist;


R¹ und R² jeweils unabhängig ausgewählt sind aus H, -(C₁-C₁₀)Alkyl, -C(O)-(C₁-C₁₀)Alkyl, Benzyl, -(C₃-C₁₀)Cycloalkyl, oder -(C₃-C₁₀)Aryl sind;

R^a und R^b jeweils unabhängig ausgewählt sind aus der Gruppe bestehend aus H, -OH, -(C₁-C₁₀)Alkyl, -[CH₂-CH₂-O]_n-(CH₂)₂-T, -C(O)-(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Alkylen-C(O)-, -(C₁-C₁₀)Alkylen-C(O)-Z, Benzyl, -(C₃-C₁₀)Cycloalkyl, -(C₃-C₁₀)Aryl-(C₁-C₁₀)Alkylen, -(C₃-C₁₀)Aryl, Halogen-(C₁-C₁₀)Alkyl, Hydroxy-(C₁-C₁₀)Alkyl, -NH-(C₁-C₁₀)Alkyl, oder -(C₁-C₁₀)Alkylen-NR^dR^e, oder R^a und R^b zusammen mit dem Stickstoff, an den sie gebunden sind, ein (C₃-C₆)-Heteroaryl oder (C₃-C₆)-Heterocycloalkyl bilden, die weiterhin ein oder mehrere Heteroatome ausgewählt aus N, S, oder O umfassen können;

Z ausgewählt ist aus -OH, -O(C₁-C₁₀)Alkyl,



oder



R^c ausgewählt ist aus -OH, -O(C₁-C₁₀)Alkyl, -OBenzyl, -O(C₃-C₁₀)Cycloalkyl, -O(C₃-C₁₀)Aryl, -O-(C₁-C₁₀)Alkylen-(C₃-C₁₀)Aryl, oder -O-(C₁-C₁₀)Alkylen-(C₃-C₁₀)Cycloalkyl,

R³ ausgewählt ist aus H, Halogen, -OH, -NH₂, -(CH₂)_p-COOH, oder -(CH₂)_p-NH₂;

T ausgewählt ist aus -H, -OH, -COOH oder -NR^dR^e;

R^d und R^e jeweils unabhängig ausgewählt sind aus H, Bindung, -OH, -(C₁-C₁₀)Alkyl, oder -(C₃-C₁₀)Heteroaryl-(C₁-C₁₀)Alkylen;


m, n, p, q, t und x jeweils unabhängig 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 oder 10 sind;

wobei jedes Alkyl, Alkylen, Aryl, Arylen, Heteroaryl, Heteroarylen, Cycloalkyl, Cycloalkylen, Heterocycloalkyl, oder Heterocycloalkylen wahlweise substituiert ist mit 1, 2, oder 3 Substituentengruppen ausgewählt aus der Gruppe bestehend aus -(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Halogenalkyl, -(C₁-C₁₀)Aminoalkyl, -(C₁-C₁₀)Alkylen-CO-OH, -(C₁-C₁₀)Hydroxyalkyl, -NH₂, -COOH, -C(O)-(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Alkylen-C(O)-, -(C₁-C₁₀)Alkylen-C(O)-X, -NH-(C₁-C₁₀)Alkyl, und -(C₁-C₁₀)Alkylen-NR^dR^e, und -NR^dR^e.

4. Die Verbindung aus Anspruch 3, wobei A (CHR¹)_m und W -C(O)-(CH₂)_p- ist.


5. Die Verbindung aus Anspruch 4, wobei R¹ Wasserstoff ist und m 2 ist.


6. Die Verbindung aus Anspruch 3, wobei Y -NH- oder




ist.


7. Die Verbindung aus Anspruch 3, wobei R^a und R^b jeweils unabhängig Wasserstoff oder Methyl sind und R^c -OH ist.
8. Die Verbindung aus Anspruch 3, wobei R^a und R^b zusammen mit dem Stickstoff, an den sie gebunden sind, ein (C₃-C₆)-Heterocycloalkyl bilden, das ausgewählt ist aus Piperidin, Piperazin, Morpholin, Thiomorpholin, Isothiazolidin, Isoxazolidin, Pyrrolidin, Imidazolidin, Thiazolidin oder Oxazolidin.
9. Die Verbindung aus Anspruch 8, wobei das (C₃-C₆)-Heterocycloalkyl Piperidin oder 4-(Piperidin-4-yl)butansäure ist.
10. Die Verbindung aus Anspruch 8, wobei R^d und R^e jeweils unabhängig (C₃-C₆)-Heteroaryl-(C₁-C₁₀)-Alkylen sind.
11. Die Verbindung aus Anspruch 3, welche ausgewählt ist aus den folgenden Verbindungen:








5




10




15

20



25

30



35

40

oder einem pharmazeutisch annehmbaren Salz, Solvat, oder Ester davon.


45

12. Ein Metallkomplex, umfassend ein Radionuklid und eine Verbindung nach Anspruch 1.

13. Der Metallkomplex aus Anspruch 12, wobei die Verbindung

50

55




ist, wobei:

A $(\text{CHR}^1)_m$ oder $\text{C}(\text{O})$ ist;

W ausgewählt ist aus der Gruppe bestehend aus $-\text{C}(\text{O})-(\text{CH}_2)_p-$, $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$, $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$, $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q-$, $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$, $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$, $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$, $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$, und $-(\text{CH}_2)_m-\text{NR}_a-(\text{CH}_2)_n-$,

Y $-\text{NH}-$, $-\text{NR}^2-$,




ist;


R^1 und R^2 jeweils unabhängig ausgewählt sind aus H, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, Benzyl, $-(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$, oder $-(\text{C}_3-\text{C}_{10})\text{Aryl}$;

R^a und R^b jeweils unabhängig ausgewählt sind aus der Gruppe bestehend aus H, $-\text{OH}$, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-\text{T}$, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}-\text{C}(\text{O})-$, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}-\text{C}(\text{O})-\text{Z}$, Benzyl, $-(\text{C}_3-\text{C}_{10})\text{Cycloalkyl}$, $-(\text{C}_3-\text{C}_{10})\text{Aryl}$, $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, $-(\text{C}_3-\text{C}_{10})\text{Aryl}$, Halogen- $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, Hydroxy- $-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, $-\text{NH}-(\text{C}_1-\text{C}_{10})\text{Alkyl}$, und $-(\text{C}_1-\text{C}_{10})\text{Alkyl}-\text{NR}^d\text{Re}$, oder R^a und R^b zusammen mit dem Stickstoff, an den sie gebunden sind, ein (C_3-C_6) -Heteroaryl oder (C_3-C_6) -Heterocycloalkyl bilden, die weiterhin ein oder mehrere Heteroatome ausgewählt aus N, S, oder O umfassen können;

Z ausgewählt ist aus $-\text{OH}$, $-\text{O}(\text{C}_1-\text{C}_{10})\text{Alkyl}$,





oder





R^c ausgewählt ist aus -OH, -O(C₁-C₁₀)Alkyl, -OBenzyl, -O(C₃-C₁₀)Cycloalkyl, -O(C₃-C₁₀)Aryl, -O-(C₁-C₁₀)Alkyl-(C₃-C₁₀)Aryl, oder -O-(C₁-C₁₀)Alkyl-(C₃-C₁₀)Cycloalkyl,
 R^3 ausgewählt ist aus H, Halogen, -OH, -NH₂, -(CH₂)_p-COOH, oder -(CH₂)_p-NH₂;
 T ausgewählt ist aus -H, -OH -COOH, oder -NR^dR^e;
 R^d und R^e jeweils unabhängig ausgewählt sind aus H, Bindung, -OH, -(C₁-C₁₀)Alkyl, oder -(C₃-C₁₀)Heteroaryl-(C₁-C₁₀)Alkyl;
 m, n, p, q, t und x jeweils unabhängig 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 oder 10 sind;
 wobei jedes Alkyl, Alkylen, Aryl, Arylen, Heteroaryl, Heteroarylen, Cycloalkyl, Cycloalkylen, Heterocycloalkyl, oder Heterocycloalkylen wahlweise substituiert ist mit 1, 2, oder 3 Substituentengruppen ausgewählt aus der Gruppe bestehend aus -(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Halogenalkyl, -(C₁-C₁₀)Aminoalkyl, -(C₁-C₁₀)Alkyl-CO-OH, -(C₁-C₁₀)Hydroxyalkyl, -NH₂, -COOH, -C(O)-(C₁-C₁₀)Alkyl, -(C₁-C₁₀)Alkyl-C(O)-, -(C₁-C₁₀)Alkyl-C(O)-X, -NH-(C₁-C₁₀)Alkyl, und -(C₁-C₁₀)Alkyl-NR^dR^e, und -NR^dR^e; und
 das Radionuklid ausgewählt ist aus der Gruppe bestehend aus ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ⁶⁴Cu, ¹⁵³Gd, ¹⁵⁵Gd, ¹⁵⁷Gd, ⁵⁹Fe, ²²⁵Ac, ²¹²Bi, ²¹³Bi, ⁵⁵Co, ⁶⁷Cu, ¹⁶⁵Dy, ¹⁶⁶Ho, ¹⁹²Ir, ²²³Ra, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁰⁵Rh, ²¹²Pb, ²¹³Pb, ¹⁴⁹Tb, ²²⁷Th, ¹⁵³Sm, ⁸⁹Sr, ^{117m}Sn, ¹⁶⁹Yb, ⁹⁰Y, ⁸⁶Y, ⁸⁹Zr und ¹⁷⁷Lu.


14. Der Metallkomplex nach Anspruch 13, welcher













oder ein pharmazeutisch annehmbares Salz oder Solvat davon ist.


15. Der Metallkomplex nach Anspruch 13, welcher











oder ein pharmazeutisch annehmbares Salz oder Solvat davon ist.

16. Eine pharmazeutische Zusammensetzung, umfassend die Verbindung nach Anspruch 3 oder den Metallkomplex nach Anspruch 13, oder ein pharmazeutisch annehmbares Salz, Solvat oder Ester davon; und einen pharmazeutisch annehmbaren Träger.
17. Die Verbindung nach einem der Ansprüche 1 bis 11 zur Verwendung in einem Verfahren zur Gewinnung eines Röntgenbildes eines oder mehrerer Gewebe, welche das Prostata-spezifische Membran-Antigen (PSMA) exprimieren, wobei das Verfahren die folgenden Schritte umfasst:
 - a. Kontaktieren eines oder mehrerer Gewebe, die PSMA exprimieren, mit einem Metallkomplex, der ein Radionuklid und die genannte Verbindung oder ein pharmazeutisch annehmbares Salz, Solvat oder Ester davon umfasst; und
 - b. Erhalten eines Röntgenbildes des einen oder der mehreren Gewebe.

18. Der Metallkomplex nach einem der Ansprüche 12 bis 15, zur Verwendung in einem Verfahren zur Gewinnung eines Röntgenbildes eines oder mehrerer Gewebe, welche das Prostata-spezifische Membran-Antigen (PSMA) exprimieren, wobei das Verfahren die folgenden Schritte umfasst:


- Kontaktieren eines oder mehrerer Gewebe, die PSMA exprimieren, mit dem Metallkomplex oder einem pharmazeutisch annehmbaren Salz, Solvat oder Ester davon; und
- Erhalten eines Röntgenbildes des einen oder der mehreren Gewebe.

19. Die Verbindung zur Verwendung nach Anspruch 17 oder der Metallkomplex zur Verwendung nach Anspruch 18, wobei das eine oder die mehreren Gewebe Prostatagewebe oder Prostatakrebsgewebe sind.

20. Die Verbindung zur Verwendung nach Anspruch 17 oder der Metallkomplex zur Verwendung nach Anspruch 18, wobei das Radionuklid ausgewählt ist aus der Gruppe bestehend aus ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{225}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , $^{117\text{m}}\text{Sn}$, ^{169}Yb , ^{90}Y , ^{86}Y , ^{89}Zr und ^{177}Lu .

Revendications

1. Composé selon la formule 1



dans lequel

A est $(\text{CHR}^1)_m$ ou $\text{C}(\text{O})$;

W est $-\text{C}(\text{O})-(\text{CH}_2)_p-$; $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$; $-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$; $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q-$; $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$; ou $-(\text{CH}_2)_m-\text{NR}_a-(\text{CH}_2)_n-$

Y est $-\text{NH}-$, $-\text{NR}^2-$, ou




X est-alkylène en $(\text{C}_1-\text{C}_{10})$ -arylène en $(\text{C}_3-\text{C}_{10})$, -arylène en $(\text{C}_3-\text{C}_{10})$, arylène en $(\text{C}_3-\text{C}_{10})$ -alkylène en $(\text{C}_1-\text{C}_{10})$ -, phenylène, -alkylène en $(\text{C}_1-\text{C}_{10})$ -cycloalkylène en $(\text{C}_3-\text{C}_{10})$ -,cycloalkylène en $(\text{C}_3-\text{C}_{10})$, ou -cycloalkylène en $(\text{C}_3-\text{C}_{10})$ -alkylène en $-(\text{C}_1-\text{C}_{10})$;


R^1 and R^2 sont chacun indépendamment H, -alkyle en $(\text{C}_1-\text{C}_{10})$, $-\text{C}(\text{O})$ -alkyle en $(\text{C}_1-\text{C}_{10})$, benzyle, -cycloalkyle en $-(\text{C}_3-\text{C}_{10})$ ou -aryle en $-(\text{C}_3-\text{C}_{10})$;

R^a et R^b sont chacun indépendamment H, $-\text{OH}$, -alkyle en $-(\text{C}_1-\text{C}_{10})$, $-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-\text{T}$, $-\text{C}(\text{O})$ -alkyle en $(\text{C}_1-\text{C}_{10})$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{C}(\text{O})-$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{C}(\text{O})-\text{Z}$, benzyle, -cycloalkyle en $(\text{C}_3-\text{C}_{10})$, -aryle en $-(\text{C}_3-\text{C}_{10})$ -alkylène en $-(\text{C}_1-\text{C}_{10})$, -aryle en $(\text{C}_3-\text{C}_{10})$, halogéno-alkyle en $(\text{C}_1-\text{C}_{10})$, hydroxy-alkyle en $(\text{C}_1-\text{C}_{10})$, $-\text{NH}-$ alkyle $(\text{C}_1-\text{C}_{10})$, ou -alkylène en $-(\text{C}_1-\text{C}_{10})-\text{NR}^d\text{Re}$, ou R^a et R^b ensemble avec l'azote auquel ils sont liés forment un hétéroaryle en (C_3-C_6) ou hétérocycloalkyle en (C_3-C_6) qui peut en outre comporter un ou plusieurs hétéroatomes choisis parmi N, S, or O ;

Z est $-\text{OH}$, $-\text{O}$ -alkyle en $(\text{C}_1-\text{C}_{10})$,



ou




R^c est -OH, -O-alkyle en (C_1-C_{10}) , O-benzyle, -O-cycloalkyle en (C_3-C_{10}) , -O-aryle en (C_3-C_{10}) , -O-alkylène en (C_1-C_{10}) -aryle en $-(C_3-C_{10})$, ou -O-alkylène en (C_1-C_{10}) -cycloalkyle en (C_3-C_{10}) ;

R^3 est H, halogène, -OH, -NH₂, $-(CH_2)_p-COOH$, ou $-(CH_2)_p-NH_2$;


T est -H, -OH, -COOH, ou $-NR^dR^e$;


R^d et R^e sont chacun indépendamment H, liaison, -OH, -alkyle en (C_1-C_{10}) , ou -hétéroaryle en (C_3-C_{10}) -alkylène en (C_1-C_{10}) ;

m, n, p, q, t et r sont chacun indépendamment 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, ou 10 ; et



D est







H3 [5-MeOsal)3 TAME], ou

dans lequel tout alkyle, alkylène, aryle, arylène, hétéroaryle, hétéroarylène, cycloalkyle, cycloalkylène, hétérocycloalkyle ou hétérocycloalkylène est éventuellement substitué avec 1, 2, ou 3 groupes substituants choisis dans le groupe comprenant -alkyle en (C₁-C₁₀), -halogénoalkyle en (C₁-C₁₀), -aminoalkyle en (C₁-C₁₀), -alkylène en (C₁-C₁₀)-COOH, -hydroxyalkyle en (C₁-C₁₀), -OH, halogène, -NH₂, -COOH, -C(O)-alkyle en (C₁-C₁₀), -alkylène en (C₁-C₁₀)-C(O)-, -alkylène en (C₁-C₁₀)-C(O)-X, -NH--alkyle en (C₁-C₁₀), et -alkylène en -(C₁-C₁₀)-NR^dR^e-, et -NR^dR^e.

2. Composé selon la revendication 1, dans lequel X est phénylène, r est 1 et D est



3. Composé selon la revendication 2, dans lequel le composé est un composé selon la Formule II




II

dans lequel :

A est $(\text{CHR}^1)_m$ ou $\text{C}(\text{O})$;

W est choisi parmi le groupe constitué de $-\text{C}(\text{O})-(\text{CH}_2)_p-$; $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$; $[\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-$;
 $-\text{C}(\text{O})-[\text{CH}(\text{R}^3)]_q-$; $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$; et
 $-(\text{CH}_2)_m-\text{NR}^a-(\text{CH}_2)_n-$;


Y est choisi parmi $-\text{NH}-$; $-\text{NR}^2-$;




R^1 et R^2 sont chacun indépendamment choisis parmi H, -alkyle en $(\text{C}_1-\text{C}_{10})$, $-\text{C}(\text{O})$ -alkyle en $(\text{C}_1-\text{C}_{10})$, benzyle, -cycloalkyle en $(\text{C}_3-\text{C}_{10})$ ou -aryle en $(\text{C}_3-\text{C}_{10})$;

R^a et R^b sont chacun indépendamment choisis dans le groupe comprenant H, -OH, -alkyle en $(\text{C}_1-\text{C}_{10})$, $[-\text{CH}_2-\text{CH}_2-\text{O}]_n-(\text{CH}_2)_2-\text{T}$, $-\text{C}(\text{O})$ -alkyle en $(\text{C}_1-\text{C}_{10})$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{C}(\text{O})-$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{C}(\text{O})-\text{Z}$, benzyle, -cycloalkyle en $(\text{C}_3-\text{C}_{10})$, -aryle en $-(\text{C}_3-\text{C}_{10})$ -alkylène en $-(\text{C}_1-\text{C}_{10})$, -aryle en $(\text{C}_3-\text{C}_{10})$, halo-alkyle en $(\text{C}_1-\text{C}_{10})$, hydroxy-alkyle en $(\text{C}_1-\text{C}_{10})$, $-\text{NH}-$ alkyle en $(\text{C}_1-\text{C}_{10})$ et -alkylène en $(\text{C}_1-\text{C}_{10})-\text{NR}^d\text{R}^e$, ou R^a et R^b avec l'azote auquel ils sont liés forment un hétéroaryle en (C_3-C_6) ou hétérocycloalkyle en (C_3-C_6) qui peut comporter en outre un ou plusieurs hétéroatomes choisis parmi N, S, or O ;

Z est choisi parmi -OH, alkyle en $-\text{O}(\text{C}_1-\text{C}_{10})$



ou



R^c est choisi parmi -OH, -O-alkyle en $(\text{C}_1-\text{C}_{10})$, -O-benzyle, -O-cycloalkyle en $(\text{C}_3-\text{C}_{10})$, -O-aryle en $(\text{C}_3-\text{C}_{10})$, -O-alkylène en $(\text{C}_1-\text{C}_{10})$ -aryle en $(\text{C}_3-\text{C}_{10})$, ou -O-alkylène en $(\text{C}_1-\text{C}_{10})$ -cycloalkyle en $-(\text{C}_3-\text{C}_{10})$;

R^3 est choisi parmi H, halogène, -OH, $-\text{NH}_2$, $-(\text{CH}_2)_p-\text{COOH}$, ou $-(\text{CH}_2)_p-\text{NH}_2$;

T est choisi parmi -H, -OH, $-\text{COOH}$, ou $-\text{NR}^d\text{R}^e$;

R^d et R^e sont chacun choisis indépendamment parmi H, liaison, -OH, -alkyle en $(\text{C}_1-\text{C}_{10})$, ou hétéroaryle en $(\text{C}_3-\text{C}_{10})$ -alkylène en $-(\text{C}_1-\text{C}_{10})$;


m, n, p, q, t et x sont chacun indépendamment 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, ou 10 ; et

dans lequel tout alkyle, alkylène, aryle, arylène, hétéroaryle, hétéroarylène, cycloalkyle, cycloalkylène, hétérocycloalkyle, ou hétérocycloalkylène est éventuellement substitué avec 1, 2, ou 3 groupes substituants choisis dans le groupe comprenant -alkyle en $(\text{C}_1-\text{C}_{10})$, -halogénoalkyle en $(\text{C}_1-\text{C}_{10})$, -aminoalkyle en $(\text{C}_1-\text{C}_{10})$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{COOH}$, -hydroxyalkyle en $(\text{C}_1-\text{C}_{10})$, $-\text{NH}_2$, $-\text{COOH}$, $-\text{C}(\text{O})$ -alkyle en $(\text{C}_1-\text{C}_{10})$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{C}(\text{O})-$, -alkylène en $(\text{C}_1-\text{C}_{10})-\text{C}(\text{O})-\text{X}$, $-\text{NH}-$ alkyle en $(\text{C}_1-\text{C}_{10})$, et -alkylène en $(\text{C}_1-\text{C}_{10})-\text{NR}^d\text{R}^e$, et $-\text{NR}^d\text{R}^e$.

4. Composé selon la revendication 3, dans lequel A est $(\text{CHR}^1)_m$ et W est $-\text{C}(\text{O})-(\text{CH}_2)_p-$.

5. Composé selon la revendication 4, dans lequel R^1 est hydrogène et m est 2.

5 6. Composé selon la revendication 3, dans lequel Y est -NH- ou



10

7. Composé selon la revendication 3, dans lequel R^a et R^b sont chacun indépendamment hydrogène ou méthyle, et R^c est -OH.

15

8. Composé selon la revendication 3, dans lequel R^a et R^b , conjointement avec l'azote auquel ils sont liés, forment un hétérocycloalkyle en (C_3-C_6) choisi parmi la pipéridine, la pipérazine, la morpholine, la thiomorpholine, fisothiazolidine, fisoazolidine, la pyrrolidine, l'imidazolidine, la thiazolidine ou foxazolidine.

20

9. Composé selon la revendication 8, dans lequel l'hétérocycloalkyle en (C_3-C_6) est de la pipéridine ou de l'acide 4-(pipéridine-4-yl)butanoïque.

10. Composé selon la revendication 8, dans lequel R^d et R^e sont chacun indépendamment -hétéroaryle en $(\text{C}_3-\text{C}_{10})$ -alkylène en $(\text{C}_1-\text{C}_{10})$.

25

11. Composé selon la revendication 3, qui est choisi parmi les composés suivants :

30


35


40

45

50


55





5


10




15

20

25




30




35

40




45

50




55

R^a et R^b sont chacun indépendamment choisis parmi H, -OH, -alkyle en $-(C_1-C_{10})$, $-[CH_2-CH_2-O]_n-(CH_2)_2-T$, -C(O)-alkyle en (C_1-C_{10}) , -alkylène en $(C_1-C_{10})-C(O)-$, -alkylène en $(C_1-C_{10})-C(O)-Z$, benzyle, -cycloalkyle en (C_3-C_{10}) , -aryle en $-(C_3-C_{10})$, -alkylène en $-(C_1-C_{10})$, -aryle en (C_3-C_{10}) , halogéno-alkyle en (C_1-C_{10}) , hydroxy-alkyle en (C_1-C_{10}) , -NH--alkyle (C_1-C_{10}) , ou -alkylène en $-(C_1-C_{10})-NR^dR^e$, ou R^a et R^b ensemble avec l'azote auquel ils sont liés forment un hétéroaryle en (C_3-C_6) ou hétérocycloalkyle en (C_3-C_6) qui peut en outre comporter un ou plusieurs hétéroatomes choisis parmi N, S, or O ;
Z est choisi parmi -OH, -O-alkyle en (C_1-C_{10}) ,



ou



R^c est choisi parmi -OH, -O-alkyle en (C_1-C_{10}) , O-benzyle, -O-cycloalkyle en (C_3-C_{10}) , -O-aryle en (C_3-C_{10}) , -O-alkylène en (C_1-C_{10}) -aryle en $-(C_3-C_{10})$, ou -O-alkylène en (C_1-C_{10}) -cycloalkyle en (C_3-C_{10}) ,

R^3 est choisi parmi H, halogène, -OH, -NH₂, $-(CH_2)_p-COOH$, ou $-(CH_2)_p-NH_2$;

T est choisi parmi -H, -OH, -COOH, ou $-NR^dR^e$;


R^d et R^e sont chacun choisis indépendamment parmi H, liaison, -OH, -alkyle en (C_1-C_{10}) , ou -hétéroaryle en (C_3-C_{10}) -alkylène en (C_1-C_{10}) ;


m, n, p, q, t et x sont chacun indépendamment 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, ou 10 ; et


dans lequel tout alkyle, alkylène, aryle, arylène, hétéroaryle, hétéroarylène, cycloalkyle, cycloalkylène, hétérocycloalkyle ou hétérocycloalkylène est éventuellement substitué avec 1, 2, ou 3 groupes substituants choisis dans le groupe comprenant -alkyle en (C_1-C_{10}) , -halogénoalkyle en (C_1-C_{10}) , -aminoalkyle en (C_1-C_{10}) , -alkylène en $(C_1-C_{10})-COOH$, -hydroxyalkyle en (C_1-C_{10}) , -NH₂, -COOH, -C(O)-alkyle en (C_1-C_{10}) , -alkylène en $(C_1-C_{10})-C(O)-$, -alkylène en $(C_1-C_{10})-C(O)-X$, -NH--alkyle en (C_1-C_{10}) , et -alkylène en $-(C_1-C_{10})-NR^dR^e$, et $-NR^dR^e$; et

le radionucléide est choisi parmi le groupe constitué de ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ⁶⁴Cu, ¹⁵³Gd, ¹⁵⁵Gd, ¹⁵⁷Gd, ⁵⁹Fe, ²²⁵Ac, ²¹²Bi, ²¹³Bi, ⁵⁵Co, ⁶⁷Cu, ¹⁶⁵Dy, ¹⁶⁶Ho, ¹⁹²Ir, ²²³Ra, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁰⁵Rh, ²¹²Pb, ²¹³Pb, ¹⁴⁹Tb, ²²⁷Th, ¹⁵³Sm, ⁸⁹Sr, ^{117m}Sn, ¹⁶⁹Yb, ⁹⁰Y, ⁸⁶Y, ⁸⁹Zr et ¹⁷⁷Lu.


14. Complexe métallique de la revendication 13, qui est :















ou un sel pharmaceutiquement acceptable ou un solvate de celui-ci.

15. Complexe métallique de la revendication 13, qui est :














ou un sel pharmaceutiquement acceptable ou un solvate de celui-ci.

16. Composition pharmaceutique comprenant le composé de la revendication 3 ou le complexe métallique de la revendication 13, ou un sel, solvate ou ester pharmaceutiquement acceptable de celui-ci ; et un support pharmaceutiquement acceptable.

17. Composé selon l'une quelconque des revendications 1 à 11, destiné à être utilisé dans une méthode d'obtention d'une image radiographique d'un ou plusieurs tissus qui expriment un antigène de membrane spécifique de la prostate (PSMA), ledit procédé comprenant les étapes suivantes :


- a. une mise en contact d'un ou plusieurs tissus qui expriment le PSMA avec un complexe métallique comprenant un radionucléide et ledit composé ou un sel, solvate ou ester pharmaceutiquement acceptable de celui-ci ; et
- b. l'obtention d'une image radiographique des un ou plusieurs tissus.

18. Complexe métallique selon l'une quelconque des revendications 12 à 15, destiné à être utilisé dans une méthode d'obtention d'une image radiographique d'un ou plusieurs tissus qui expriment un antigène de membrane spécifique de la prostate (PSMA), ledit procédé comprenant les étapes suivantes :

- a. une mise en contact d'un ou plusieurs tissus qui expriment le PSMA avec ledit complexe métallique ou un sel, solvate ou ester pharmaceutiquement acceptable de celui-ci ; et
- b. l'obtention d'une image radiographique des un ou plusieurs tissus.

19. Composé destiné à être utilisé selon la revendication 17 ou complexe métallique destiné à être utilisé selon la revendication 18, dans lequel les un ou plusieurs tissus sont des tissus de prostate ou du tissu de cancer de la prostate.

20. Composé destiné à être utilisé selon la revendication 17 ou complexe métallique destiné à être utilisé selon la revendication 18, dans lequel le radionucléide est choisi dans le groupe constitué de ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{225}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , $^{117\text{m}}\text{Sn}$, ^{169}Yb , ^{90}Y , ^{86}Y , ^{89}Zr et ^{177}Lu .



٧٤٥




FIG. 2





FIG. 3





 DEPARTMENT OF HEALTH AND HUMAN SERVICES




FIG. 5




FIG. 6

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61752350 A [0001]
- US 61785788 B [0001]
- WO 2008058192 A2 [0008]

Non-patent literature cited in the description

- THE PRACTICE OF MEDICINAL CHEMISTRY. Academic Press, 1996, 203 [0094]
- GREENE, T.W. ; WUTS, P.G.M. Protective Groups in Organic Synthesis. Wiley, 1999 [0104]


TRIAZIN ALAPÚ RADIOGYÓGYSZEREK ÉS RADIOLÓGIAI KÉPALKOTÓ ÁGENSEK

Szabadalmi igénypontok



SZTNH-100086540

I. (I) képletű vegyület



ahol:

A jelentése $(\text{CHR}^1)_m$ vagy $\text{C}(\text{O})$;

W jelentése $-\text{C}(\text{O})-(\text{CH}_2)_p-$; $-\text{C}(\text{O})[-\text{CH}_2-\text{CH}_2-\text{O}]_n-$; $-\{\text{CH}_2-\text{CH}_2-\text{O}\}_n-(\text{CH}_2)_2-$; $-\text{C}(\text{O})-[\text{CH}(\text{R}^1)]_q-$; $-(\text{CH}_2)_m-\text{O}-$; $-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})-(\text{CH}_2)_n-$; $-(\text{CH}_2)_m-\text{S}(\text{O})_2-(\text{CH}_2)_n-$ vagy $-(\text{CH}_2)_m-\text{NR}_a-(\text{CH}_2)_n-$;

Y jelentése $-\text{NH}-$; $-\text{NR}^2-$ vagy




X jelentése $-(\text{C}_1-\text{C}_{10})$ alkilén- $(\text{C}_3-\text{C}_{10})$ arilén-, $-(\text{C}_3-\text{C}_{10})$ arilén-, $-(\text{C}_3-\text{C}_{10})$ arilén- $(\text{C}_1-\text{C}_{10})$ alkilén-, fenilén-, $-(\text{C}_1-\text{C}_{10})$ alkilén- $(\text{C}_3-\text{C}_{10})$ cikloalkilén-, $-(\text{C}_3-\text{C}_{10})$ cikloalkilén vagy $-(\text{C}_3-\text{C}_{10})$ cikloalkilén- $(\text{C}_1-\text{C}_{10})$ alkilén-;

R^1 és R^2 jelentése egymástól függetlenül H, $-(\text{C}_1-\text{C}_{10})$ alkil-, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})$ alkil-, benzil-, $-(\text{C}_3-\text{C}_{10})$ cikloalkil vagy $-(\text{C}_3-\text{C}_{10})$ aril;

R^a és R^b jelentése egymástól függetlenül H, $-\text{OH}$, $-(\text{C}_1-\text{C}_{10})$ alkil-, $-\{\text{CH}_2-\text{CH}_2-\text{O}\}_n-(\text{CH}_2)_2-$, $-\text{C}(\text{O})-(\text{C}_1-\text{C}_{10})$ alkil-, $-(\text{C}_1-\text{C}_{10})$ alkilén- $\text{C}(\text{O})-$, $-(\text{C}_1-\text{C}_{10})$ alkilén- $\text{C}(\text{O})-\text{Z}$, benzil-, $-(\text{C}_3-\text{C}_{10})$ cikloalkil-, $-(\text{C}_3-\text{C}_{10})$ aril- $(\text{C}_1-\text{C}_{10})$ alkilén-, $-(\text{C}_3-\text{C}_{10})$ aril-, halo- $(\text{C}_1-\text{C}_{10})$ alkil-, hidroxi- $(\text{C}_1-\text{C}_{10})$ alkil-, $-\text{NH}-(\text{C}_1-\text{C}_{10})$ alkil vagy $-(\text{C}_1-\text{C}_{10})$ alkilén- NR^dR^e -, vagy R^a és R^b azzal a nitrogénatommal együtt, amelyhez kapcsolódnak, (C_3-C_6) -heteroaril vagy (C_3-C_6) -heterocikloalkil csoportot alkot, amely tartalmazhat továbbá egy vagy több, a következők közül választott heteroatomot: N, S vagy O;

Z jelentése $-\text{OH}$, $-\text{O}(\text{C}_1-\text{C}_{10})$ alkil,



R^c jelentése $-\text{OH}$, $-\text{O}(\text{C}_1-\text{C}_{10})$ alkil-, $-\text{Obenzil}$ -, $-\text{O}(\text{C}_3-\text{C}_{10})$ cikloalkil-, $-\text{O}(\text{C}_3-\text{C}_{10})$ aril-, $-\text{O}-(\text{C}_1-\text{C}_{10})$ alkilén- $(\text{C}_3-\text{C}_{10})$ aril vagy $-\text{O}-(\text{C}_1-\text{C}_{10})$ alkilén- $(\text{C}_3-\text{C}_{10})$ cikloalkil,


R^3 jelentése H, halogén, $-\text{OH}$, $-\text{NH}_2$, $-(\text{CH}_2)_6-\text{COOH}$ vagy $-(\text{CH}_2)_6-\text{NH}_2$;

T jelentése $-\text{H}$, $-\text{OH}$, $-\text{COOH}$ vagy $-\text{NR}^d\text{R}^e$;

R^d és R^e jelentése egymástól függetlenül H, kötés, $-\text{OH}$, $-(\text{C}_1-\text{C}_{10})$ alkil vagy $-(\text{C}_3-\text{C}_{10})$ heteroaril- $(\text{C}_1-\text{C}_{10})$ alkilén;


m, n, p, q, t és r értéke egymástól függetlenül 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 vagy 10; és

D jelentése



ahol bármely alkil, alkilén, aril, arilén, heteroaril, heteroarilén, cikloalkil, cikloalkilén, heterocikloalkil vagy heterocikloalkilén adott esetben szubsztituált 1, 2 vagy 3, a következők alkotta csoportból választott szubsztituenssel: $-(\text{C}_1\text{-C}_{10})$ alkil, $-(\text{C}_1\text{-C}_{10})$ haloalkil, $-(\text{C}_1\text{-C}_{10})$ aminoalkil, $-(\text{C}_1\text{-C}_{10})$ alkilén-COOH, $-(\text{C}_1\text{-C}_{10})$ hidroxialkil, -OH, halogén, $-\text{NH}_2$, $-\text{COOH}$, $-\text{C}(\text{O})-(\text{C}_1\text{-C}_{10})$ alkil, $-(\text{C}_1\text{-C}_{10})$ alkilén-C(O)-, $-(\text{C}_1\text{-C}_{10})$ alkilén-C(O)-X, $-\text{NH}-(\text{C}_1\text{-C}_{10})$ alkil és $-(\text{C}_1\text{-C}_{10})$ alkilén- NR^{dR^e} és $-\text{NR}^{\text{dR}^e}$.

2. Az 1. igénypont szerinti vegyület, ahol X jelentése fenilén, r értéke 1 és D jelentése



enssel: $-(C_1-C_{10})$ alkil, $-(C_1-C_{10})$ haloalkil, $-(C_1-C_{10})$ aminoalkil, $-(C_1-C_{10})$ alkilén-COOH, $-(C_1-C_{10})$ hidroxialkil, $-NH_2$, $-COOH$, $-C(O)-(C_1-C_{10})$ alkil, $-(C_1-C_{10})$ alkilén-C(O)-, $-(C_1-C_{10})$ alkilén-C(O)-X, $-NH-(C_1-C_{10})$ alkil és $-(C_1-C_{10})$ alkilén- NR^dR^e és $-NR^dR^e$.

4. A 3. igénypont szerinti vegyület, ahol A jelentése $(CHR^1)_m$ és W jelentése $-C(O)-(CH_2)_p-$.

5. A 4. igénypont szerinti vegyület, ahol R^1 jelentése hidrogén és m értéke 2.

6. A 3. igénypont szerinti vegyület, ahol Y jelentése $-NH-$ vagy




7. A 3. igénypont szerinti vegyület, ahol R^a és R^b jelentése egymástól függetlenül hidrogén vagy metil és R^c jelentése $-OH$.


8. A 3. igénypont szerinti vegyület, ahol R^a és R^b azzal a nitrogénatommal együtt, amelyhez kapcsolódnak, piperidin, piperazin, morfolin, tiomorfolin, izotiazolidin, izoxazolidin, pirrolidin, imidazolidin, tiazolidin vagy oxazolidin közül választott (C_3-C_6) -heterocikloalkil csoportot alkot.


9. A 8. igénypont szerinti vegyület, ahol a (C_1-C_6) -heterocikloalkil piperidin vagy 4-(piperidin-4-il)butánsav.


10. A 8. igénypont szerinti vegyület, ahol R^d és R^e jelentése egymástól függetlenül $-(C_3-C_{10})$ heteroaril- $-(C_1-C_{10})$ alkilén.

11. A 3. igénypont szerinti vegyület, amely a következő vegyületek közül választott:










vagy gyógyászatilag elfogadható sója, szolvátja vagy észtere.

12. Az 1. igénypont szerinti vegyületet és radionuklidot tartalmazó fémkomplex.

13. A 12. igénypont szerinti fémkomplex, ahol a vegyület




ahol:

A jelentése $(CHR^1)_m$ vagy $C(O)$;

W jelentése a következők alkotta csoportból választott: $-C(O)-(CH_2)_p-$; $-C(O)-[CH_2-CH_2-O]_n-$; $-[CH_2-CH_2-O]_n-$; $-(CH_2)_2-$; $-C(O)-[CH(R^3)]_q-$; $-(CH_2)_m-O-(CH_2)_n-$; $-(CH_2)_m-S-(CH_2)_n-$; $-(CH_2)_m-S(O)-(CH_2)_n-$; $-(CH_2)_n-S(O)_2-$; $-(CH_2)_n-$ és $-(CH_2)_m-NR^6-(CH_2)_n-$.


Y jelentése -NH-, -NR²- közül választott,



R¹ és R² jelentése egymástól függetlenül H, -(C₁-C₁₀)alkil, -C(O)-(C₁-C₁₀)alkil, benzil, -(C₃-C₁₀)cikloalkil vagy -(C₃-C₁₀)aril közül választott;

R^a és R^b jelentése egymástól függetlenül a következők alkotta csoportból választott: H, -OH, -(C₁-C₁₀)alkil, -[CH₂-CH₂-O]_n-(CH₂)_x-T, -C(O)-(C₁-C₁₀)alkil, -(C₁-C₁₀)alkilén-C(O)-, -(C₁-C₁₀)alkilén-C(O)-Z, benzil, -(C₃-C₁₀)cikloalkil, -(C₃-C₁₀)aril-(C₁-C₁₀)alkilén, -(C₃-C₁₀)aril, halo-(C₁-C₁₀)alkil, hidroxil-(C₁-C₁₀)alkil, -NH-(C₁-C₁₀)alkil és -(C₁-C₁₀)alkilén-NR^dR^e-, vagy R^a és R^b azzal a nitrogénatommal együtt, amelyhez kapcsolódnak (C₃-C₆)-heteroaril vagy (C₁-C₆)-heterocikloalkil csoportot alkot, amely tartalmazhat továbbá egy vagy több, a következők közül választott heteroatomot: N, S vagy O;

Z jelentése -OH, -O(C₁-C₁₀)alkil közül választott,



R^c jelentése -OH, -O(C₁-C₁₀)alkil, -Obenzil, -O(C₃-C₁₀)cikloalkil, -O(C₃-C₁₀)aril, -O-(C₁-C₁₀)alkilén-(C₃-C₁₀)aril vagy -O-(C₁-C₁₀)alkilén-(C₃-C₁₀)cikloalkil közül választott,

R³ jelentése H, halogén, -OH, -NH₂, -(CH₂)_p-COOH vagy -(CH₂)_p-NH₂ közül választott;

T jelentése -H, -OH, -COOH vagy -NR^dR^e közül választott;


R^d és R^e jelentése egymástól függetlenül H, kötés, -OH, -(C₁-C₁₀)alkil vagy -(C₃-C₁₀)heteroaril-(C₁-C₁₀)alkilén közül választott;


m, n, p, q, t és x értéke egymástól függetlenül 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 vagy 10;





ahol bármely alkil, alkilén, aril, arilén, heteroaril, heteroarilén, cikloalkil, cikloalkilén, heterocikloalkil vagy heterocikloalkilén adott esetben szubsztituált 1, 2 vagy 3, a következők alkotta csoportból választott szubsztituenssel: -(C₁-C₁₀)alkil, -(C₁-C₁₀)haloalkil, -(C₁-C₁₀) aminoalkil, -(C₁-C₁₀)alkilén-COOH, -(C₁-C₁₀)hidroxialkil, -NH₂, -COOH, -C(O)-(C₁-C₁₀)alkil, -(C₁-C₁₀)alkilén-C(O)-, -(C₁-C₁₀)alkilén-C(O)-X, -NH-(C₁-C₁₀)alkil és -(C₁-C₁₀)alkilén-NR^dR^e- és -NR^dR^e;




a radionuklid a következők alkotta csoportból választott: ¹¹¹In, ⁹⁰Y, ⁶⁸Ga, ⁶⁴Cu, ¹⁵³Gd, ¹⁵⁵Gd, ¹⁵⁷Gd, ⁵⁹Fe, ²²⁵Ac, ²¹²Bi, ²¹³Bi, ⁵⁵Co, ⁶⁷Cu, ¹⁶⁵Dy, ¹⁶⁶Ho, ¹⁹²Ir, ²²³Ra, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁰⁵Rh, ²¹²Pb, ²¹³Pb, ¹⁴⁹Tb, ²²⁷Th, ¹⁵³Sm, ⁸⁹Sr, ^{117m}Sn, ¹⁶⁹Yb, ⁹⁰Y, ⁸⁶Y, ⁸⁹Zr és ¹⁷⁷Lu.

14. A 13. igénypont szerinti fémkomplex, amely:












vagy gyógyászati lag elfogadható sója vagy szolvátja.

15. A 13. igénypont szerinti fémkomplex, amely:







b. az egy vagy több szövet radiográfiás képének előállítása.

18. A 12-15. igénypontok bármelyike szerinti fémkomplex egy vagy több, prosztataspecifikus membrán-antigént (PSMA) expresszáló szövet radiográfiás képének előállítására szolgáló eljárásban történő alkalmazásra, amely eljárás tartalmazza a következő lépéseket:

a. egy vagy több, PSMA-t expresszáló szövet érintkeztetése a fémkomplexszel vagy annak gyógyászatilag elfogadható sójával, szolvátjával vagy észterével; és

b. egy vagy több szövet radiográfiás képének előállítása.

19. A 17. igénypont szerinti vegyület vagy a 18. igénypont szerinti fémkomplex az ott meghatározott alkalmazásra, ahol az egy vagy több szövet prosztataszövet vagy prosztatarákszövet.

20. A 17. igénypont szerinti vegyület vagy a 18. igénypont szerinti fémkomplex az ott meghatározott alkalmazásra, ahol a radionuklid a következők alkotta csoportból választott: ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{59}Fe , ^{223}Ac , ^{212}Bi , ^{213}Bi , ^{55}Co , ^{67}Cu , ^{165}Dy , ^{166}Ho , ^{192}Ir , ^{223}Ra , ^{186}Re , ^{188}Re , ^{105}Rh , ^{212}Pb , ^{213}Pb , ^{149}Tb , ^{227}Th , ^{153}Sm , ^{89}Sr , ^{117}mSn , ^{169}Yb , ^{90}Y , ^{86}Y , ^{89}Zr és ^{177}Lu .