

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 July 2009 (30.07.2009)

PCT

(10) International Publication Number
WO 2009/094536 A1

(51) International Patent Classification:

C07D 311/82 (2006.01) **C07F 5/02** (2006.01)
C07D 471/06 (2006.01) **C09B 11/24** (2006.01)
C07F 9/28 (2006.01) **G01N 33/533** (2006.01)

(74) Agent: **BLODGETT, James**; Life Technologies Corporation c/o Intellevate, P.O. Box 52050, Minneapolis, MN 55402 (US).

(21) International Application Number:

PCT/US2009/031830

(22) International Filing Date: 23 January 2009 (23.01.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

61/023,191 24 January 2008 (24.01.2008) US

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **LIFE TECHNOLOGIES CORPORATION** [US/US]; 5791 Van Allen Way, Carlsbad, CA 92008 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CHEN, Aimei** [US/US]; 3853 Monroe Street, Eugene, OR 97405 (US). **GEE, Kyle, R.** [US/US]; 37699 Upper Camp Creek Road, Springfield, OR 97478 (US). **KANG, Hee Chol** [US/US]; 4980 Blanton Road, Eugene, OR 97405 (US).

Published:

— with international search report

(54) Title: FLUOROGENIC HYDRAZINE-SUBSTITUTED COMPOUNDS

(57) Abstract: The present disclosure is directed to fluorogenic schiff base-forming dyes capable of detecting analytes containing aldehyde and ketone groups. The dyes contain nucleophilic hydrazinyl appendages and are capable of binding and detecting analytes *in situ*.

WO 2009/094536 A1

FLUOROGENIC HYDRAZINE-SUBSTITUTED COMPOUNDS

RELATED APPLICATIONS

5

This application claims the benefit of U.S. Provisional Application No. 61/023,191, filed January 24, 2008, the disclosure of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

10

Schiff base-forming fluorogenic dyes containing a hydrazinyl substituent appended to a fluorophore are disclosed. The compounds are useful in the detection of analytes containing aldehyde and ketone groups.

BACKGROUND OF THE INVENTION

15

A myriad of biomarkers containing aldehyde and ketone moieties exist and can play an important role in the biological, forensic, medical and industrial sciences. In particular, aldehydes and ketones are known to be key end products in the degradation of a variety of biological molecules, such as, lipids, nucleic acids, carbohydrates and proteins. In a number of instances, these end products are a result of oxidative stress. In one example, malondialdehyde and 4-hydroxynonenal are markers for lipid peroxidation.

20

A number of reagents for the detection of aldehyde and ketone-containing moieties have been proposed, however, each with limited success. Among the most widely used of such reagents are dansyl hydrazine, fluorescein thiosemicarbazide, various biotin hydrazides, biotin hydroxylamine (ARP), and various aromatic amines (2-aminopyridine, 8-aminonaphthalene-1,3,6-disulfonic acid, 1-aminopyrene-3,6,8-trisulfonic acid, 2-aminoacridone). Unfortunately, these reagents require additional purification and/or secondary reagents.

25

Existing methods of labeling carbohydrates that utilize hydrazine, hydroxylamine and amine derivatization reagents have focused on labeling aldehydes present in, or introduced into, carbohydrates, particularly the so-called "reducing sugars". Aldehydes are typically introduced into carbohydrates by periodate oxidation. The adduct formed with the reducing sugar can be further stabilized by treatment with borohydride or a cyanoborohydride. The derivatization reaction typically proceeds or is followed by a separation technique such as chromatography, electrophoresis, precipitation, affinity isolation or other means before direct or indirect detection of the labeled product. Unlike the foregoing, which require purification and the use of a secondary detection reagent, the

35

reagents of the present invention permit rapid in-situ detection of aldehyde and ketone moieties upon contact.


U.S. Patent No. 6,967,251 (Haugland et al.) describes aniline-substituted quinazolinone compounds which can display fluorescent changes upon binding of aldehyde-containing compounds in a gel. However, the aniline-substituted quinazolinone core, which is important for binding of the analyte, is less fluorescent and fluorogenic than the compounds of the present invention. Accordingly, the present invention provides substantial advantages over the previously described aldehyde and ketone detector compounds.

A family of hydrazinyl substituted xanthene dyes have been previously described in U.S. Pat. No. 4,420,627 (Widiger). However, each of the xanthene moieties in Widiger are substituted to prevent analyte binding of the molecule. Particularly, the terminal nitrogen atom in the hydrazinyl moiety in Widiger is carbonylated, such that the hydrazinyl group is not nucleophilic and unable to bind an aldehyde or ketone moiety. Accordingly, the compounds are not functional in the analyte detection method of the present invention, which involve a quenched molecule up until the analyte binding event. Accordingly, the compounds in Widiger are functionally and structurally very different from those of the present invention.

SUMMARY OF THE INVENTION

The present invention provides hydrazinyl-substituted compounds capable of binding aldehyde and ketone-containing analytes in a solution, wherein upon binding of the analyte, the hydrazinyl-substituted compounds become highly fluorescent, thereby indicating the presence of the analyte.

One aspect of the invention provides a compound of Formula I:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;


R^1 and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^3 is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, anlyte substituted hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R^2 and R^3 , R^3 and R^4 , or R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.

Another aspect of the invention provides a compound of Formula II:



II

or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an anlyte;


Y is H or an anlyte;

L is -NH- or a covalent bond;

R¹ is selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and R¹¹ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl.

Another aspect of the invention provides a compound of Formula III:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;


wherein,

R¹ is selected from the group consisting of H, alkyl and substituted alkyl;


R², R⁴, R⁵, R⁶, R⁷, R⁸ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R¹¹ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl.


5 Another aspect of the invention provides a compound selected from the group consisting of:



Another embodiment of the invention provides a compound selected from the group consisting of:



Another aspect of the invention provides a compound of Formula V:



5 or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;

Z¹ a fluorophore and Z² is R⁸; or


Z¹ is R⁸ and Z² is a fluorophore;

10 R⁵, R⁶, R⁸ and R⁹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R¹⁴ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

20 X and R¹⁴ are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group.

Another aspect of the invention provides a compound of Formula VI:



VI

or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

5 G is a 5-6 membered fused aryl or heteroaryl group;

X is H or an anlyte;

R¹ is selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, anlyte substituted hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

25 R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.

Another aspect of the invention provides a composition comprising:

30 (a) an anlyte; and

(b) a compound as described herein.

Another aspect of the invention provides a method for determining the presence of an analyte of interest in a sample, wherein the method comprises:

contacting the sample with a compound described herein;

incubating the sample and the compound for a sufficient amount of time to form a complex between the compound and analyte;

illuminating the complex with an appropriate wavelength to form an illuminated complex; and

detecting emissions from the illuminated complex.


Another aspect of the invention provides a kit for detecting an analyte in a sample, wherein the kit comprises:

a compound as describe herein that binds the analyte; and

instructions for detecting the analyte.

Another aspect of the invention provides a compound comprising a (9-phenyl-3H-xanthen-3-ylidene)hydrazine, wherein the hydrazine group is unsubstituted or substituted with an analyte.

Another aspect of the invention provides a method of synthesizing a compound of Formula III, a tautomer thereof, a stereoisomer thereof, or a salt thereof:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;


wherein,

R^1 and R^{11} are each independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl;


5 R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

the method comprising:

contacting a compound of formula IV:



with R^1 -NH-NH- R^{12} and R^{11} -NH-NH- R^{12} , wherein R^{12} is a protecting group, thereby obtaining a compound of Formula III(a):



contacting R^{12} with a deprotecting reagent, thereby obtaining a compound of Formula III.

In another embodiment the SO_3^- group in formula III is a CO_2^- and the SO_2 group is a CO.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows emission spectra of 2-(6-hydrazinyl-3-hydrazono-3H-xanthen-9-yl)benzenesulfonate before and after contact with acetaldehyde. There was a significant fluorescent increase upon reaction with an aldehyde. Spectra were taken at pH 6 in a 50 mM phosphate buffer solution.

Figure 2 shows quantitation of aldehyde content in solution. 20 μ M of 2-(6-hydrazinyl-3-hydrazono-3H-xanthen-9-yl)benzenesulfonate was incubated with aldehyde for 15 min at room temperature at pH 6 in a 50 mM phosphate buffer solution.

Figure 3A shows the excitation maximum at 519 nm for un-reacted 2-(6-hydrazinyl-3-hydrazono-3H-xanthen-9-yl)benzenesulfonate. There were no fluorescent emissions detected from excitation at 519 nm.

Figure 3B shows excitation maximum at 548 nm for aldehyde-reacted 2-(6-hydrazinyl-3-hydrazono-3H-xanthen-9-yl)benzenesulfonate and a bright emission spectra at 581 nm for the same species. Spectra were taken at pH 6 in 50 mM phosphate buffer solution.

DETAILED DESCRIPTION OF THE INVENTION

Introduction:

The present invention provides compounds with a hydrazine appendage capable of binding aldehyde and ketone groups present on analytes of interest. The compounds of the present invention generally have a fluorescent core that is quenched in the unbound state. Once the hydrazine reacts with an aldehyde or ketone, a Schiff base (imine) is formed. The resultant compound is highly fluorescent, thereby providing an excellent method for detection of aldehydes and ketones in solution.

The compounds of the present invention provide distinct advantages over known aldehyde detection reagents, including being fluorogenic with a high quantum yield, unique emission and excitation wavelengths outside the range of endogenous particles, highly stable and high specificity for ketones and aldehydes, soluble in a variety of solutions, particularly aqueous solutions, and compatibility with biological applications. Additionally, reaction with the compounds of the present invention proceeds under milder conditions (rt, aqueous acetic acid) than existing assays. Furthermore, the Schiff base formation appears to be a near-instantaneous reaction, whereas existing assays require at least 1 hour reaction time at elevated temperatures. Additionally, the available wavelength range for the compounds is large and tunable.

Definitions:

Before describing the present invention in detail, it is to be understood that this invention is not limited to specific compositions or process steps, as such may vary. It should be noted that, as used in this specification and the appended claims, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. Thus, for example, reference to "a hydrazine group" includes a plurality of hydrazine groups and reference to "an analyte" includes a plurality of analytes and the like.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention is related. The following terms are defined for purposes of the invention as described herein.

"Alkyl" refers to monovalent saturated aliphatic hydrocarbyl groups having from 1 to 10 carbon atoms and preferably 1 to 6 carbon atoms. This term includes, by way of example, linear and branched hydrocarbyl groups such as methyl (CH_3 -), ethyl (CH_3CH_2 -), *n*-propyl ($\text{CH}_3\text{CH}_2\text{CH}_2$ -), isopropyl ($(\text{CH}_3)_2\text{CH}$ -), *n*-butyl ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2$ -), isobutyl ($(\text{CH}_3)_2\text{CHCH}_2$ -), *sec*-butyl ($(\text{CH}_3)(\text{CH}_3\text{CH}_2)\text{CH}$ -), *t*-butyl ($(\text{CH}_3)_3\text{C}$ -), *n*-pentyl ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ -), and neopentyl ($(\text{CH}_3)_3\text{CCH}_2$ -).

"Substituted alkyl" refers to an alkyl group having from 1 to 5, preferably 1 to 3, or more preferably 1 to 2 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, cycloalkenyl, substituted cycloalkenyl, cycloalkenyloxy, substituted cycloalkenyloxy, cycloalkenylthio, substituted

cycloalkenylthio, guanidino, substituted guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclylthio, substituted heterocyclylthio, nitro, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said
5 substituents are defined herein.

"Alkoxy" refers to the group –O-alkyl wherein alkyl is defined herein. Alkoxy includes, by way of example, methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, *t*-butoxy, *sec*-butoxy, and *n*-pentoxy.

10 "Substituted alkoxy" refers to the group –O-(substituted alkyl) wherein substituted alkyl is defined herein.

"Acyl" refers to the groups H-C(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, alkenyl-C(O)-, substituted alkenyl-C(O)-, alkynyl-C(O)-, substituted alkynyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, cycloalkenyl-C(O)-, substituted
15 cycloalkenyl-C(O)-, aryl-C(O)-, substituted aryl-C(O)-, heteroaryl-C(O)-, substituted heteroaryl-C(O)-, heterocyclic-C(O)-, and substituted heterocyclic-C(O)-, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined
20 herein. Acyl includes the "acetyl" group CH₃C(O)-.

"Acylamino" refers to the groups –NRC(O)alkyl, -NRC(O)substituted alkyl, -NRC(O)cycloalkyl, -NRC(O)substituted cycloalkyl, -NRC(O)cycloalkenyl, -NRC(O)substituted cycloalkenyl, -NRC(O)alkenyl, -NRC(O)substituted alkenyl, -NRC(O)alkynyl, -NRC(O)substituted alkynyl, -NRC(O)aryl, -NRC(O)substituted aryl,
25 -NRC(O)heteroaryl, -NRC(O)substituted heteroaryl, -NRC(O)heterocyclic, and -NRC(O)substituted heterocyclic wherein R is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined
30 herein.

"Acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, alkenyl-C(O)O-, substituted alkenyl-C(O)O-, alkynyl-C(O)O-, substituted alkynyl-C(O)O-, aryl-C(O)O-, substituted aryl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, cycloalkenyl-C(O)O-, substituted cycloalkenyl-C(O)O-, heteroaryl-C(O)O-, substituted
35 heteroaryl-C(O)O-, heterocyclic-C(O)O-, and substituted heterocyclic-C(O)O- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl,

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

"Amino" refers to the group -NH_2 .

5 "Substituted amino" refers to the group -NR'R'' where R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, -SO_2 -alkyl, -SO_2 -substituted alkyl, -SO_2 -alkenyl, -SO_2 -substituted alkenyl, -SO_2 -cycloalkyl, -SO_2 -substituted cycloalkyl, -SO_2 -cycloalkenyl, -SO_2 -substituted cycloalkenyl, -SO_2 -aryl, -SO_2 -substituted aryl, -SO_2 -heteroaryl, -SO_2 -substituted heteroaryl, -SO_2 -heterocyclic, and -SO_2 -substituted heterocyclic and wherein R' and R'' are optionally joined, together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, provided that R' and R'' are both not hydrogen, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein. When R' is hydrogen and R'' is alkyl, the substituted amino group is sometimes referred to herein as alkylamino. When R' and R'' are alkyl, the substituted amino group is sometimes referred to herein as dialkylamino. When referring to a monosubstituted amino, it is meant that either R' or R'' is hydrogen but not both. When referring to a disubstituted amino, it is meant that neither R' nor R'' are hydrogen.

"Aminocarbonyl" refers to the group -C(O)NR'R'' where R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminothiocabonyl" refers to the group -C(S)NR'R'' where R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonylamino" refers to the group -NRC(O)NR'R'' where R is hydrogen or alkyl and R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminothiocabonylamino" refers to the group -NRC(S)NR'R'' where R is hydrogen or alkyl and R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonyloxy" refers to the group -O-C(O)NR'R'' where R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

“Aminosulfonyl” refers to the group $-\text{SO}_2\text{NR}'\text{R}''$ where R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

“Aminosulfonyloxy” refers to the group $-\text{O}-\text{SO}_2\text{NR}'\text{R}''$ where R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

“Aminosulfonylamino” refers to the group $-\text{NR}-\text{SO}_2\text{NR}'\text{R}''$ where R is hydrogen or alkyl and R' and R'' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

“Amidino” refers to the group $-\text{C}(=\text{NR}''')\text{R}'\text{R}''$ where R' , R'' , and R''' are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R' and R'' are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted

heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

5 "Aryl" or "Ar" refers to a monovalent aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (*e.g.*, phenyl) or multiple condensed rings (*e.g.*, naphthyl or anthryl) which condensed rings may or may not be aromatic (*e.g.*, 2-benzoxazolinone, 2H-1,4-benzoxazin-3(4H)-one-7-yl, and the like) provided that the point of attachment is at an aromatic carbon atom. Preferred aryl groups include phenyl and
10 naphthyl.

"Substituted aryl" refers to aryl groups which are substituted with 1 to 5, preferably 1 to 3, or more preferably 1 to 2 substituents selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl,
15 aminothiocabonyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, cycloalkenyl, substituted
20 cycloalkenyl, cycloalkenyloxy, substituted cycloalkenyloxy, cycloalkenylthio, substituted cycloalkenylthio, guanidino, substituted guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclylthio, substituted heterocyclylthio, nitro, SO₃H, substituted
25 sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said substituents are defined herein.

"Aryloxy" refers to the group –O-aryl, where aryl is as defined herein, that includes, by way of example, phenoxy and naphthoxy.

30 "Substituted aryloxy" refers to the group –O-(substituted aryl) where substituted aryl is as defined herein.

"Arylthio" refers to the group –S-aryl, where aryl is as defined herein.

"Substituted arylthio" refers to the group –S-(substituted aryl), where substituted aryl is as defined herein.

35 "Alkenyl" refers to alkenyl groups having from 2 to 6 carbon atoms and preferably 2 to 4 carbon atoms and having at least 1 and preferably from 1 to 2 sites of

alkenyl unsaturation. Such groups are exemplified, for example, by vinyl, allyl, and but-3-en-1-yl.

"Substituted alkenyl" refers to alkenyl groups having from 1 to 3 substituents, and preferably 1 to 2 substituents, selected from the group consisting of alkoxy, substituted
5 alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, cycloalkenyl, substituted
10 cycloalkenyl, cycloalkenyloxy, substituted cycloalkenyloxy, cycloalkenylthio, substituted cycloalkenylthio, guanidino, substituted guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted
15 heterocyclyloxy, heterocyclylthio, substituted heterocyclylthio, nitro, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said substituents are defined herein and with the proviso that any hydroxy substitution is not attached to a vinyl (unsaturated) carbon atom.

"Alkynyl" refers to alkynyl groups having from 2 to 6 carbon atoms and preferably 2 to 3 carbon atoms and having at least 1 and preferably from 1 to 2 sites of
20 alkynyl unsaturation.

"Substituted alkynyl" refers to alkynyl groups having from 1 to 3 substituents, and preferably 1 to 2 substituents, selected from the group consisting of alkoxy, substituted
25 alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, cycloalkenyl, substituted
30 cycloalkenyl, cycloalkenyloxy, substituted cycloalkenyloxy, cycloalkenylthio, substituted cycloalkenylthio, guanidino, substituted guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclylthio, substituted heterocyclylthio, nitro, SO₃H, substituted
35 sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said

substituents are defined herein and with the proviso that any hydroxy substitution is not attached to an acetylenic carbon atom.

“Carbonyl” refers to the divalent group —C(O)— which is equivalent to —C(=O)— .

5 “Carboxyl” or “carboxy” refers to —COOH or salts thereof.

“Carboxyl ester” or “carboxy ester” refers to the groups —C(O)O—alkyl , $\text{—C(O)O—substituted alkyl}$, —C(O)O—alkenyl , $\text{—C(O)O—substituted alkenyl}$, —C(O)O—alkynyl , $\text{—C(O)O—substituted alkynyl}$, —C(O)O—aryl , $\text{—C(O)O—substituted aryl}$, —C(O)O—cycloalkyl , $\text{—C(O)O—substituted cycloalkyl}$, $\text{—C(O)O—cycloalkenyl}$, $\text{—C(O)O—substituted cycloalkenyl}$,
10 —C(O)O—heteroaryl , $\text{—C(O)O—substituted heteroaryl}$, $\text{—C(O)O—heterocyclic}$, and $\text{—C(O)O—substituted heterocyclic}$ wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

15 “(Carboxyl ester)amino” refers to the group —NR—C(O)O—alkyl , substituted —NR—C(O)O—alkyl , —NR—C(O)O—alkenyl , $\text{—NR—C(O)O—substituted alkenyl}$, —NR—C(O)O—alkynyl , $\text{—NR—C(O)O—substituted alkynyl}$, —NR—C(O)O—aryl , $\text{—NR—C(O)O—substituted aryl}$, $\text{—NR—C(O)O—cycloalkyl}$, $\text{—NR—C(O)O—substituted cycloalkyl}$, $\text{—NR—C(O)O—cycloalkenyl}$, $\text{—NR—C(O)O—substituted cycloalkenyl}$, $\text{—NR—C(O)O—heteroaryl}$, $\text{—NR—C(O)O—substituted heteroaryl}$, $\text{—NR—C(O)O—heterocyclic}$, and $\text{—NR—C(O)O—substituted heterocyclic}$ wherein R is alkyl or hydrogen, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

25 “(Carboxyl ester)oxy” refers to the group —O—C(O)O—alkyl , substituted —O—C(O)O—alkyl , —O—C(O)O—alkenyl , $\text{—O—C(O)O—substituted alkenyl}$, —O—C(O)O—alkynyl , $\text{—O—C(O)O—substituted alkynyl}$, —O—C(O)O—aryl , $\text{—O—C(O)O—substituted aryl}$, $\text{—O—C(O)O—cycloalkyl}$, $\text{—O—C(O)O—substituted cycloalkyl}$, $\text{—O—C(O)O—cycloalkenyl}$, $\text{—O—C(O)O—substituted cycloalkenyl}$, $\text{—O—C(O)O—heteroaryl}$, $\text{—O—C(O)O—substituted heteroaryl}$,
30 $\text{—O—C(O)O—heterocyclic}$, and $\text{—O—C(O)O—substituted heterocyclic}$ wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

“Cyano” refers to the group —CN .

"Cycloalkyl" refers to cyclic alkyl groups of from 3 to 10 carbon atoms having single or multiple cyclic rings including fused, bridged, and spiro ring systems. Examples of suitable cycloalkyl groups include, for instance, adamantyl, cyclopropyl, cyclobutyl, cyclopentyl, and cyclooctyl.

5 "Cycloalkenyl" refers to non-aromatic cyclic alkyl groups of from 3 to 10 carbon atoms having single or multiple cyclic rings and having at least one $>C=C<$ ring unsaturation and preferably from 1 to 2 sites of $>C=C<$ ring unsaturation.

"Substituted cycloalkyl" and "substituted cycloalkenyl" refers to a cycloalkyl or cycloalkenyl group having from 1 to 5 or preferably 1 to 3 substituents selected from the group consisting of oxo, thione, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, cycloalkenyl, substituted cycloalkenyl, cycloalkenyloxy, substituted cycloalkenyloxy, cycloalkenylthio, substituted cycloalkenylthio, guanidino, substituted guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclylthio, substituted heterocyclylthio, nitro, SO_3H , substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said substituents are defined herein.

"Cycloalkyloxy" refers to $-O$ -cycloalkyl.

25 "Substituted cycloalkyloxy refers to $-O$ -(substituted cycloalkyl).

"Cycloalkylthio" refers to $-S$ -cycloalkyl.

"Substituted cycloalkylthio" refers to $-S$ -(substituted cycloalkyl).

"Cycloalkenyloxy" refers to $-O$ -cycloalkenyl.

"Substituted cycloalkenyloxy refers to $-O$ -(substituted cycloalkenyl).

30 "Cycloalkenylthio" refers to $-S$ -cycloalkenyl.

"Substituted cycloalkenylthio" refers to $-S$ -(substituted cycloalkenyl).

"Guanidino" refers to the group $-NHC(=NH)NH_2$.

"Substituted guanidino" refers to $-NR^{13}C(=NR^{13})N(R^{13})_2$ where each R^{13} is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic

and two R¹³ groups attached to a common guanidino nitrogen atom are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, provided that at least one R¹³ is not hydrogen, and wherein said substituents are as defined herein.

5 "H" indicates hydrogen.

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo.

"Hydroxy" or "hydroxyl" refers to the group –OH.

"Heteroaryl" refers to an aromatic group of from 1 to 10 carbon atoms and 1 to 4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur within the
10 ring. Such heteroaryl groups can have a single ring (e.g., pyridinyl or furyl) or multiple condensed rings (e.g., indolizinyl or benzothienyl) wherein the condensed rings may or may not be aromatic and/or contain a heteroatom provided that the point of attachment is through an atom of the aromatic heteroaryl group. In one embodiment, the nitrogen and/or the sulfur ring atom(s) of the heteroaryl group are optionally oxidized to provide for the N-oxide
15 (N→O), sulfinyl, or sulfonyl moieties. Preferred heteroaryls include pyridinyl, pyrrolyl, indolyl, thiophenyl, and furanyl.

"Substituted heteroaryl" refers to heteroaryl groups that are substituted with from 1 to 5, preferably 1 to 3, or more preferably 1 to 2 substituents selected from the group consisting of the same group of substituents defined for substituted aryl.

20 "Heteroaryloxy" refers to –O-heteroaryl.

"Substituted heteroaryloxy" refers to the group –O-(substituted heteroaryl).

"Heteroarylthio" refers to the group –S-heteroaryl.

"Substituted heteroarylthio" refers to the group –S-(substituted heteroaryl).

"Heterocycle" or "heterocyclic" or "heterocycloalkyl" or "heterocyclyl" refers to
25 a saturated or unsaturated group having a single ring or multiple condensed rings, including fused bridged and spiro ring systems, from 1 to 10 carbon atoms and from 1 to 4 hetero atoms selected from the group consisting of nitrogen, sulfur or oxygen within the ring wherein, in fused ring systems, one or more the rings can be cycloalkyl, aryl or heteroaryl provided that the point of attachment is through the non-aromatic ring. In one embodiment,
30 the nitrogen and/or sulfur atom(s) of the heterocyclic group are optionally oxidized to provide for the N-oxide, sulfinyl, sulfonyl moieties.

"Substituted heterocyclic" or "substituted heterocycloalkyl" or "substituted heterocyclyl" refers to heterocyclyl groups that are substituted with from 1 to 5 or preferably 1 to 3 of the same substituents as defined for substituted cycloalkyl.

35 "Heterocyclioxy" refers to the group –O-heterocyclyl.

"Substituted heterocyclyloxy" refers to the group $-O-(\text{substituted heterocycyl})$.

"Heterocyclylthio" refers to the group $-S-\text{heterocycyl}$.

"Substituted heterocyclylthio" refers to the group $-S-(\text{substituted heterocycyl})$.

Examples of heterocycle and heteroaryls include, but are not limited to,

- 5 azetidine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, dihydroindole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline,
 10 phthalimide, 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholinyl, thiomorpholinyl (also referred to as thiamorpholinyl), 1,1-dioxothiomorpholinyl, piperidinyl, pyrrolidine, and tetrahydrofuranyl.


"Hydrazinyl" refers to the group $-NHNH_2$, $=NNH-$, or $=N^{(+)}HNNH_2$.

- 15 "Substituted hydrazinyl" refers to a hydrazinyl group, wherein a non-hydrogen atom, such as an alkyl group, is appended to one or both of the hydrazinyl amine groups. An example of substituted hydrazinyl is $-N(\text{alkyl})-NH_2$ or $=N^{+}(\text{alkyl})-NH_2$.

"Nitro" refers to the group $-NO_2$.

"Oxo" refers to the atom $(=O)$ or $(-O^-)$.

- 20 "Spirocyclyl" refers to divalent saturated cyclic group from 3 to 10 carbon atoms having a cycloalkyl or heterocyclyl ring with a spiro union (the union formed by a single atom which is the only common member of the rings) as exemplified by the following structure:



"Sulfonyl" refers to the divalent group $-S(O)_2-$.

- 25 "Substituted sulfonyl" refers to the group $-SO_2\text{-alkyl}$, $-SO_2\text{-substituted alkyl}$, $-SO_2\text{-alkenyl}$, $-SO_2\text{-substituted alkenyl}$, $-SO_2\text{-cycloalkyl}$, $-SO_2\text{-substituted cycloalkyl}$, $-SO_2\text{-cycloalkenyl}$, $-SO_2\text{-substituted cycloalkenyl}$, $-SO_2\text{-aryl}$, $-SO_2\text{-substituted aryl}$, $-SO_2\text{-heteroaryl}$, $-SO_2\text{-substituted heteroaryl}$, $-SO_2\text{-heterocyclic}$, $-SO_2\text{-substituted heterocyclic}$, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl,
 30 heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein. Substituted sulfonyl includes groups such as methyl- SO_2- , phenyl- SO_2- , and 4-methylphenyl- SO_2- .

"Sulfonyloxy" refers to the group $-\text{OSO}_2\text{-alkyl}$, $-\text{OSO}_2\text{-substituted alkyl}$, $-\text{OSO}_2\text{-alkenyl}$, $-\text{OSO}_2\text{-substituted alkenyl}$, $-\text{OSO}_2\text{-cycloalkyl}$, $-\text{OSO}_2\text{-substituted cycloalkyl}$, $-\text{OSO}_2\text{-cycloalkenyl}$, $-\text{OSO}_2\text{-substituted cycloalkenyl}$, $-\text{OSO}_2\text{-aryl}$, $-\text{OSO}_2\text{-substituted aryl}$, $-\text{OSO}_2\text{-heteroaryl}$, $-\text{OSO}_2\text{-substituted heteroaryl}$, $-\text{OSO}_2\text{-heterocyclic}$, $-\text{OSO}_2\text{-substituted heterocyclic}$, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Thioacyl" refers to the groups H-C(S)- , alkyl-C(S)- , substituted alkyl-C(S)- , alkenyl-C(S)- , substituted alkenyl-C(S)- , alkynyl-C(S)- , substituted alkynyl-C(S)- , cycloalkyl-C(S)- , substituted cycloalkyl-C(S)- , $\text{cycloalkenyl-C(S)-}$, substituted $\text{cycloalkenyl-C(S)-}$, aryl-C(S)- , substituted aryl-C(S)- , heteroaryl-C(S)- , substituted heteroaryl-C(S)- , $\text{heterocyclic-C(S)-}$, and substituted $\text{heterocyclic-C(S)-}$, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Thiol" refers to the group $-\text{SH}$.


"Thiocarbonyl" refers to the divalent group $-\text{C(S)-}$ which is equivalent to $-\text{C(=S)-}$.

"Thione" refers to the atom $(=\text{S})$.


"Alkylthio" refers to the group $-\text{S-alkyl}$ wherein alkyl is as defined herein.

"Substituted alkylthio" refers to the group $-\text{S-(substituted alkyl)}$ wherein substituted alkyl is as defined herein.


A dashed line projecting from a substituent, such as:



indicates the point of attachment to the base molecule. For a fused ring, dashed lines indicate portions of the base molecule where the fused ring is attached, such as:



wherein the full molecule could have the structure:



“Stereoisomer” or “stereoisomers” refer to compounds that differ in the
chirality of one or more stereocenters. Stereoisomers include enantiomers and
5 diastereomers.

“Tautomer” refers to alternate forms of a compound that differ in the position
of a proton, such as enol-keto and imine-enamine tautomers, or the tautomeric forms of
heteroaryl groups containing a ring atom attached to both a ring -NH- moiety and a ring =N-
moeity such as pyrazoles, imidazoles, benzimidazoles, triazoles, and tetrazoles.

10 “Patient,” “subject” or “individual” refers to mammals and includes humans
and non-human mammals, such as monkeys, dogs, cats, horses, cows, pigs or rats.

“Salt” refers to acceptable salts of a compound, which salts are derived from
a variety of organic and inorganic counter ions well known in the art and include, by way of
example only, sodium, potassium, calcium, magnesium, ammonium, and
15 tetraalkylammonium; and when the molecule contains a basic functionality, salts of organic
or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate,
maleate, and oxalate.

“Treating” or “treatment” of a disease in a patient refers to 1) preventing the
disease from occurring in a patient that is predisposed or does not yet display symptoms of
20 the disease; 2) inhibiting the disease or arresting its development; or 3) ameliorating or
causing regression of the disease.

The terms “protein” and “polypeptide” are used herein in a generic sense to
include polymers of amino acid residues of any length. The term “peptide” is used herein to
refer to polypeptides having less than 250 amino acid residues, typically less than 100 amino
25 acid residues. The terms apply to amino acid polymers in which one or more amino acid
residues are an artificial chemical analogue of a corresponding naturally occurring amino
acid, as well as to naturally occurring amino acid polymers.

The term “reactive group” as used herein refers to a group that is capable of
reacting with another chemical group to form a covalent bond, i.e. is covalently reactive
30 under suitable reaction conditions, and generally represents a point of attachment for
another substance. The reactive group is a moiety, such as carboxylic acid or succinimidyl
ester, on the compounds of the present invention that is capable of chemically reacting with

a functional group on a different compound to form a covalent linkage. Reactive groups generally include nucleophiles, electrophiles and photoactivatable groups.

Exemplary reactive groups include, but not limited to, olefins, acetylenes, alcohols, phenols, ethers, oxides, halides, aldehydes, ketones, carboxylic acids, esters, amides, cyanates, isocyanates, thiocyanates, isothiocyanates, amines, hydrazines, hydrazones, hydrazides, diazo, diazonium, nitro, nitriles, mercaptans, sulfides, disulfides, sulfoxides, sulfones, sulfonic acids, sulfinic acids, acetals, ketals, anhydrides, sulfates, sulfenic acids isonitriles, amidines, imides, imidates, nitrones, hydroxylamines, oximes, hydroxamic acids thiohydroxamic acids, allenes, ortho esters, sulfites, enamines, ynamines, ureas, pseudoureas, semicarbazides, carbodiimides, carbamates, imines, azides, azo compounds, azoxy compounds, and nitroso compounds. Reactive functional groups also include those used to prepare bioconjugates, *e.g.*, N-hydroxysuccinimide esters, maleimides and the like. Methods to prepare each of these functional groups are well known in the art and their application to or modification for a particular purpose is within the ability of one of skill in the art (see, for example, Sandler and Karo, eds., Organic Functional Group Preparations, Academic Press, San Diego, 1989).

The term "detectable response" as used herein refers to an occurrence of or a change in, a signal that is directly or indirectly detectable either by observation or by instrumentation. Typically, the detectable response is an optical response resulting in a change in the wavelength distribution patterns or intensity of absorbance or fluorescence or a change in light scatter, fluorescence lifetime, fluorescence polarization, or a combination of the above parameters.

The term "dye" as used herein refers to a compound that emits light to produce an observable detectable signal.

The term "fluorophore" or "fluorogenic" as used herein refers to a composition that demonstrates a change in fluorescence upon binding to a biological compound or analyte interest. Preferred fluorophores of the present invention include fluorescent dyes having a high quantum yield in aqueous media. Exemplary fluorophores include xanthene, indole, borapolyazaindacene, furan, and benzofuran, among others. The fluorophores of the present invention may be substituted to alter the solubility, spectral properties or physical properties of the fluorophore.

The term "carrier molecule" as used herein refers to a biological or a non-biological component that is covalently bonded to a compound of the present invention. Such components include, but are not limited to, an amino acid, a peptide, a protein, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid, a hapten, a

psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a polymeric microparticle, a biological cell, a virus and combinations thereof. "Covalently bonded" as used herein indicates a direct covalent linkage or through a number of atoms corresponding to a linker moiety.

5 The term "Linker" as used herein, refers to a single covalent bond or a series of stable covalent bonds incorporating 1-20 nonhydrogen atoms selected from the group consisting of C, N, O, S and P that covalently attach the fluorogenic or fluorescent compounds to another moiety such as a chemically reactive group or a biological and non-biological component. Exemplary linking members include a moiety that includes -C(O)NH-,
10 -C(O)O-, -NH-, -S-, -O-, and the like. A "cleavable linker" is a linker that has one or more cleavable groups that may be broken by the result of a reaction or condition. The term "cleavable group" refers to a moiety that allows for release of a portion, e.g., a fluorogenic or fluorescent moiety, of a conjugate from the remainder of the conjugate by cleaving a bond linking the released moiety to the remainder of the conjugate. Such cleavage is either
15 chemical in nature, or enzymatically mediated. Exemplary enzymatically cleavable groups include natural amino acids or peptide sequences that end with a natural amino acid.

 In addition to enzymatically cleavable groups, it is within the scope of the present invention to include one or more sites that are cleaved by the action of an agent other than an enzyme. Exemplary non-enzymatic cleavage agents include, but are not
20 limited to, acids, bases, light (e.g., nitrobenzyl derivatives, phenacyl groups, benzoin esters), and heat. Many cleavable groups are known in the art. See, for example, Jung *et al.*, *Biochem. Biophys. Acta*, **761**: 152-162 (1983); Joshi *et al.*, *J. Biol. Chem.*, **265**: 14518-14525 (1990); Zarling *et al.*, *J. Immunol.*, **124**: 913-920 (1980); Bouizar *et al.*, *Eur. J. Biochem.*, **155**: 141-147 (1986); Park *et al.*, *J. Biol. Chem.*, **261**: 205-210 (1986); Browning *et al.*, *J.*
25 *Immunol.*, **143**: 1859-1867 (1989). Moreover a broad range of cleavable, bifunctional (both homo- and hetero-bifunctional) spacer arms are commercially available.

 An exemplary cleavable group, an ester, is cleavable group that may be cleaved by a reagent, e.g. sodium hydroxide, resulting in a carboxylate-containing fragment and a hydroxyl-containing product.

30 The linker can be used to attach the compound to another component of a conjugate, such as a targeting moiety (e.g., antibody, ligand, non-covalent protein-binding group, etc.), an analyte, a biomolecule, a drug and the like.

 Unless indicated otherwise, the nomenclature of substituents that are not explicitly defined herein are arrived at by naming the terminal portion of the functionality

followed by the adjacent functionality toward the point of attachment. For example, the substituent "arylalkyloxycabonyl" refers to the group (aryl)-(alkyl)-O-C(O)-.

It is understood that in all substituted groups defined above, polymers arrived at by defining substituents with further substituents to themselves (e.g., substituted aryl
5 having a substituted aryl group as a substituent which is itself substituted with a substituted aryl group, which is further substituted by a substituted aryl group etc.) are not intended for inclusion herein. In such cases, the maximum number of such substitutions is three. For example, serial substitutions of substituted aryl groups with two other substituted aryl groups are limited to -substituted aryl-(substituted aryl)-substituted aryl.

10 Similarly, it is understood that the above definitions are not intended to include impermissible substitution patterns (e.g., a methyl substituted with 5 fluoro groups). Such impermissible substitution patterns are well known to the skilled artisan.

A simple, sensitive reagent for the selective detection of aldehydes/ketones is a useful tool for high-throughput screening systems in chemistry and biology. In particular,
15 aldehydes have been widely known as among the key end products from a degradation of a variety of biological molecules, e.g. lipids, nucleic acids, carbohydrates, proteins, induced by oxidative stress.


Malondialdehyde (MDA) is a major lipid peroxidation product. The present invention involves reaction of non-fluorescent (or virtually non-fluorescent) fluorophore-
20 hydrazines with MDA under mild conditions such that a fluorescent pyrazole is formed. Thus a fluorogenic assay for lipid peroxidation is presented that allows for a large wavelength range to be utilized.

Lipid peroxidation is a major indicator of oxidative stress. Researchers need to measure lipid peroxidation in many types of samples including drugs, food products, and
25 human and biological tissue. The extent of lipid peroxidation provides important information regarding free radical activity in disease states, and it also provides a measure of antioxidant activity of potential therapeutic compounds. Malondialdehyde (MDA) is a key lipid peroxidation product. The most commonly used assay of lipid peroxidation involves the reaction of MDA with 2 equivalents of thiobarbituric acid, aka the OXI-TEK TBARS Assay
30 Kit. The resulting oxonol product is detected by absorbance or fluorescence (e.g. Janero D., "Malondialdehyde and thiobarbituric acid reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury," *Free Radical Biology & Medicine*, **1998**, 9: 515-40).

Drawbacks to this method include: 1) samples must be heated in strong acetic acid for 1 hour at 95°C; 2) only a single wavelength set (530 nm excitation/550 nm emission) is
35 produced.

MDA is known to react with aryl hydrazines to form pyrazoles (Otteneider et al, "Reaction of Malondialdehyde-DNA adducts with Hydrazines- Development of a Facile Assay for Quantification of Malondialdehyde Equivalents in DNA," *Chem. Res. Toxicol.* **2002**, 15: 312-8).

- 5 The present invention describes a new class of fluorogenic dyes for the detection and quantitation of aldehyde/ketone functional groups. The compounds are capable of reacting with a variety of analytes, such as malondialdehyde (MDA) as shown below:




10

- Aniline versions of visible wavelength fluorophores such as fluoresceinamine, 5-amino-tetramethylrhodamine, and 8-aminophenyl-BODIPY are virtually non-fluorescent because of internal charge transfer (ICT). During the fluorophore excited state, the LUMO is filled by intramolecular transfer of electron density from the aniline nitrogen electron lone pair. This filling precludes relaxation to the ground state via fluorescence. Thus, even though the compounds absorb visible light, they do not fluoresce as they return to the ground state. These compounds can be made fluorescent by conversion into products in which the aniline nitrogen electron lone pair becomes unavailable for ICT. In one embodiment, the present invention involves reaction of
- 15



fluorophore-hydrazine combinations (non-fluorescent because of ICT) with MDA to form fluorescent pyrazole products. The pyrazole products are fluorescent because the electron lone pair on the “first” nitrogen atom of the hydrazine moiety becomes part of the aromaticity of the pyrazole, i.e. this electron lone pair is no longer available to quench fluorescence by ICT.

5



This principle can be applied to a number of other fluorophore hydrazine combinations, for example:

10



The aldehyde or ketone functional group is typically naturally present on the analyte prior to its conjugation to the reagent of the invention. Alternatively, the aldehyde or ketone functionality is formed on the target substance by chemical, light, heat, radiation, or enzymatic treatment prior to reaction with a reagent of the invention. In one aspect of the invention, the target substance is treated with an oxidizing condition, such as a chemical

15

oxidizing agent (for example a periodate, a strong acid, or ozone), oxidizing radiation, photolysis, or enzymatic oxidation.

Where they are not present, aldehydes and ketones are also introduced into molecules using extrinsic reagents that already contain an aldehyde or ketone. For instance, aldehydes are introduced at aliphatic amine sites with the reagents succinimidyl 4-formylbenzoate or succinimidyl 4-formylphenoxyacetate (Molecular Probes, Eugene OR). These reagents selectively modify proteins on the surface of live cells, and thereby permit the analysis of the topology of peptide and protein exposure on cells surfaces following, for instance, lysis and gel electrophoresis. Additionally, galactosides are enzymatically transferred to a target carbohydrate using UDP-galactose:N-acetylglucosamine galactosyltransferase and, following galactose oxidase-catalyzed oxidation to an aldehyde (as described by Shaper et al. J. SUPRAMOL. STRUCTURE 6, 291-299 (1977)), the target carbohydrate can be modified by a reagent of the invention. Glycoproteins such as horseradish peroxidase are oxidized to aldehydes and their conjugates subsequently used in various detection schemes according to the instant invention

The oligosaccharide components of cell surface glycoproteins play a role in the interactions that regulate many important biological processes, from cell-cell adhesion to signal transduction. Sialic acids are the most abundant terminal components of oligosaccharides on mammalian cell-surface glycoproteins and are synthesized from the six-carbon precursor N-acetylmannosamine. When cells in culture are incubated with N-levulinoyl-D-mannosamine, this ketone-containing monosaccharide serves as a substrate in the oligosaccharide synthesis pathway, resulting in ketone-tagged cell-surface oligosaccharides (as described in U.S. Pat. No. 6,075,134 to Bertozzi et al. (2000), incorporated by reference). If these tagged cells are then labeled with a reagent of the invention, they are readily identified or traced using either by imaging or flow cytometry.

The conjugated target is typically a peptide, a protein, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid, a sugar, a polysaccharide, a lipid, a lipopolysaccharide, a ganglioside, a drug, a hormone, or a ligand having a molecular weight less than 2,000 Daltons. Preferably, the conjugated target is a protein, a nucleic acid, a lipid, a lipopolysaccharide, a ganglioside, a drug, or a hormone.

The use of the invention to label aldehyde- and ketone-containing target substances comprises combining a reagent of the present invention with a sample that contains or is thought to contain a desired target, incubating the mixture of reagent and

sample for a time sufficient for the reagent to form a covalent conjugate with the target substance in the sample, such that the conjugate exhibits a detectable fluorescent signal.

5 The characteristics of the resulting reagent-target conjugate, including the presence, location, intensity, excitation and emission spectra, fluorescence polarization, fluorescence lifetime, photobleaching rate and other physical properties of the fluorescent signal can be used to detect, differentiate, sort, quantitate, and/or analyze aspects or portions of the sample. The reagents of the invention are optionally used in conjunction with one or more additional detection reagents (preferably having detectably different
10 fluorescence characteristics).

Selected Target Substances Containing Aldehydes or Ketones Formaldehyde
Acetone Benzaldehydes Reducing sugars and polysaccharides in ring-opened forms
Steroids Keto acids Aldehyde- or ketone-containing drugs Aldehyde- or ketone-containing
15 environmental pollutants Aldehyde- or ketone-containing organics Acid-treated deoxyribonucleic acids Oxidized sugars Oxidized polysaccharides Oxidized glycols Oxidized glycoproteins Oxidized glycolipids Oxidized glycosaminoglycans Oxidized ribonucleic acids Oxidized biological cells Oxidized N-terminal serine residues of proteins Oxidized N-terminal
20 threonine residues of proteins

Typically, when the reagent of the invention is used in the form of a staining solution, preferably an aqueous or aqueous miscible solution that is compatible with the sample and the intended use. For biological samples, where minimal perturbation of cell morphology or physiology is desired, the staining solution is selected accordingly. For
25 solution assays, the staining solution preferably does not perturb the native conformation of the target substance.

Although typically used in an aqueous or aqueous miscible solution, the staining solution is typically prepared by first dissolving the reagent in a water-miscible
30 organic solvent such as dimethylsulfoxide (DMSO), dimethylformamide (DMF), or a lower alcohol, such as methanol or ethanol. This stock solution is typically prepared at a concentration of greater than about 50-times that used in the final staining solution, then diluted one or more times with an aqueous solvent or a buffer solution such that the reagent is present in an effective amount. Typically, the reagent is first dissolved in 100% DMF, and
35 then diluted with buffer. The staining solution optionally further comprises additional formulation components, such as acids, buffering agents, inorganic salts, polar organic solvents, antioxidants, and ion chelators.

The pH of the staining solution is optionally modified by the inclusion of a buffering agent. Any buffering agent that is compatible with the target substance in the sample is suitable for inclusion in the staining solution. In a preferred embodiment the buffer is PBS. In another embodiment, the buffering agent is one of the so-called "Good's" buffers. "Good's" buffers include BES (N,N-bis[2-hydroxyethyl]-2-aminoethanesulfonic acid; 2-[bis(2-hydroxyethyl)amino]ethanesulfonic acid), BICINE (N,N-bis[2-hydroxyethyl]glycine), CAPS (3-[cyclohexylamino]-1-propanesulfonic acid), EPPS (N-[2-hydroxyethyl]piperazine-N'-[3-propanesulfonic acid]), HEPES ((N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid]), MES (2-[N-morpholino]ethanesulfonic acid), MOPS (3-[N-morpholino]propanesulfonic acid), PIPES (piperazine-N,N'-bis[2-ethanesulfonic acid]; 1,4-piperazinediethanesulfonic acid), TAPS (N-tris [hydroxymethyl]methyl-3-aminopropanesulfonic acid; ([2-hydroxy-1, 1-bis(hydroxymethyl)ethyl]amino-1-propanesulfonic acid), TES (N-tris[hydroxymethyl]methyl-2-aminoethanesulfonic acid; 2-([2-hydroxy-1,1-bis(hydroxymethyl)ethyl]amino)ethanesulfonic acid), or TRICINE (N-tris [hydroxymethyl]methylglycine; N-[2-hydroxy-1,1-bis(hydroxymethyl)ethyl]glycine).

Other preferred buffering agents include salts of formate, citrate, acetate, N-(2-hydroxyethyl)-N'-(2-sulfoethyl)piperazine, imidazole, N-(2-hydroxyethylpiperazine)-N'-2-ethanesulfonic acid, Tris(hydroxymethyl)aminomethane acetate, or Tris(hydroxymethyl)aminomethane hydrochloride. In a preferred embodiment, the buffering agent is MES, sodium acetate, or acetic acid, preferably acetic acid. The buffering agent or mixture of buffering agents is typically present in the staining solution at a concentration of 20 mM to 500 mM, preferably about 25 mM to about 100 mM. Where the buffering agent is acetic acid, it is preferably present in a concentration of about 1%-6%, more preferably at about 3%.

In a particularly advantageous formulation of the staining solution, the staining solution additionally comprises an inorganic salt. Advantageous inorganic salts produce staining formulations that exhibit low background signals when staining glycoproteins in electrophoretic gels. Particularly useful and inexpensive salts include sodium chloride, ammonium sulfate, magnesium chloride, magnesium acetate, zinc chloride, magnesium sulfate and magnesium glucuronate present in the staining solution at a concentration of 1-50%. In a preferred embodiment, the inorganic salt is sodium chloride or magnesium chloride, more preferably magnesium chloride.


An effective amount of reagent is the amount of reagent sufficient to give a detectable

fluorescence response in combination with the desired target. The reagent concentration in the solution must be sufficient both to contact the target in the sample and to combine with the target in an amount sufficient to give a signal, but too much reagent may cause problems with background fluorescence or speckling in gels. The optimal concentration and
 5 composition of the staining solution is determined by the nature of the sample (including physical, biological, biochemical and physiological properties), the nature of the reagent-target interaction (including the transport rate of the reagent to the site of the target), and the nature of the analysis being performed, and can be determined using standard procedures, similar to those described in examples below.

10 In another embodiment of the invention, the target substance contains a carboxylic acid or sulfonic acid functional group, therefore the functional group must first be activated before combining with a staining solution containing a reagent of the invention, depending upon the properties of the target substance. Typically carbodiimides, such as
 15 EDAC, or dicyclohexylcarbodiimide (DCC) are used to activate carboxylic acids, whereas sulfonic acids most often require formation of their sulfonyl chloride by standard means. The reagent adducts of carboxylic acids and sulfonic acids are typically used to characterize the target substance, or the conjugates are used as fluorescent tracers. Carboxylic acid and sulfonic acids do not form stable adducts when stained in gels or solutions, thus
 20 differentiating them from aldehydes and ketones.

Particular Aspects of the Invention:

One aspect of the invention provides a compound of Formula I:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
 wherein,
 X is H or an analyte;

R^1 and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and


R^3 is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, anlyte substituted hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R^2 and R^3 , R^3 and R^4 , or R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.

In another embodiment, R^1 is H. Alternatively, R^1 is alkyl or substituted alkyl. In a preferred embodiment R^{1a} is H. It is noted that R^{1a} is not a carbonyl group since the resultant amide would not be nucleophilic nor functional as a aldehyde/ketone reactive moiety. In another embodiment, R^3 hydrazinyl or substituted hydrazinyl. In another embodiment, R^3 is methoxy. In another embodiment, R^3 is hydroxy. In another embodiment, R^3 is amino.

30

In another embodiment of the invention, R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group or substituted fused heterocyclyl group of the formula:




R^{12} and R^{13} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

In another embodiment, R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are all H. In another embodiment, R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are all H. In another embodiment, R^9 is $-SO_3^-$. In another embodiment, R^9 is $-CO_2^-$. In another embodiment, X is H. In another embodiment, R^3 is methoxy and R^9 is $-CO_2^-$.

In another embodiment of the invention, X is an analyte. In another embodiment, the analyte is bound through an imine. In another embodiment, the analyte is selected from the group consisting of an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a drug, a lipid, and a synthetic polymer. In another embodiment, the analyte comprises a carbonyl group. In another embodiment, the analyte comprises an aldehyde group. In another embodiment, R^3 is analyte substituted hydrazinyl. In another embodiment, the analyte is bound through an imine. In another embodiment the analyte is malondialdehyde. In another embodiment the analyte is 4-hydroxynonenal.

Another particular aspect of the invention provides a salt of the compound.

Another aspect of the invention provides a compound of Formula II:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
wherein,

X is H or an analyte;

Y is H or an analyte;

L is -NH- or a covalent bond;


R¹ is selected from the group consisting of H, alkyl and substituted alkyl;

5 R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

10 R¹¹ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl.

In another more particular embodiment, R⁹ is SO₃⁻. In another embodiment, R⁹ is CO₂⁻. In another embodiment, R², R⁴, R⁵, R⁶, R⁷, R⁸ and R¹⁰ are H. In another embodiment, R¹ and R¹¹ are H. In another embodiment, R¹ is alkyl or substituted alkyl and R¹¹ is alkyl or substituted alkyl. In another embodiment thereof, R¹ and R¹¹ are -(CH₂)₃SO₃⁻. In another embodiment, X and Y are H. In another embodiment, X is an analyte and Y is an analyte. In another embodiment, L is -NH-. In another embodiment, L is a covalent bond.

Another aspect of the invention provides a compound of Formula III:



25

or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

R¹ is selected from the group consisting of H, alkyl and substituted alkyl;


R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and


R^{11} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl.

In a more particular embodiment thereof, R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are hydrogen. More particularly, R^1 and R^{11} are H. In another embodiment, R^1 is alkyl or substituted alkyl and R^{11} is alkyl or substituted alkyl. In another embodiment, R^1 and R^{11} are both $-(\text{CH}_2)_3\text{SO}_3^-$.


In another embodiment the SO_3^- group in formula III is a CO_2^- .

Another aspect of the invention provides a compound selected from the group consisting of:






Another embodiment of the invention provides a compound selected from the group consisting of:



Another embodiment thereof provides a salt or tautomer thereof.

Another aspect of the invention provides a compound of Formula V:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
wherein,

X is H or an analyte;

Z^1 a fluorophore and Z^2 is R^8 ; or

Z^1 is R^8 and Z^2 is a fluorophore;

R^5 , R^6 , R^8 and R^9 are each independently selected from the group consisting of H,


5 alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^{14} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

15 X and R^{14} are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group.

In another embodiment, one of Z^1 or Z^2 is selected from the group consisting of a xanthene, an indole and a borapolyazaindacine.

20 In a more particular embodiment, Z^1 or Z^2 is:



wherein,

R^2 , R^4 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

30 R^3 is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, analyte substituted hydrazinyl, alkyl,

substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R^2 and R^3 , R^3 and R^4 , or R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group;


R^{15} is selected from the group consisting of $=\text{O}$ and $=^{(+)}\text{NR}^{16}\text{R}^{17}$; and

R^{16} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, heteroaryl, and substituted heteroaryl; and

R^{17} is selected from the group consisting of alkyl, substituted alkyl, amino, and substituted amino.


In a more particular embodiment, R^{15} is $=\text{O}$. In another embodiment, R^{15} is $=^{(+)}\text{NR}^{16}\text{R}^{17}$. In another embodiment, R^{16} and R^{17} are methyl. In another embodiment, R^2 , R^4 and R^{10} are H. In another embodiment, R^3 is amino or substituted amino. In another embodiment, R^3 is hydroxy. In another embodiment, R^3 is $-\text{N}(\text{CH}_3)_2$.

In another embodiment, R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group or substituted fused heterocyclyl group of the following structure:



wherein, R^{12} and R^{13} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

In a more particular embodiment, Z¹ or Z² is:




R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³ are each independently selected from the group
 5 consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, -SO₃H, sulfonyl, substituted sulfonyl, sulfonyloxy,
 10 thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

In another embodiment, R¹⁹ and R²² are H. In another embodiment, R¹⁸, R²⁰, R²¹ and R²³ are methyl. In another embodiment, R¹⁸, R¹⁹, R²⁰, R²¹, R²² and R²³ are H.

15

In a more particular embodiment, Z¹ or Z² is:



R²⁴ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and
 20

R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, -SO₃H, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.
 25

In another embodiment thereof, R^{24} , R^{25} , R^{27} , R^{28} and R^{29} are H. In another embodiment, R^{26} is a carboxyl ester. In another embodiment, R^{26} is $-\text{CO}_2\text{CH}_3$. In another embodiment, R^5 , R^6 , R^8 and R^{14} are all H. In another embodiment, R^9 is $-\text{SO}_3^-$. In another embodiment, R^9 is $-\text{CO}_2^-$.

5


In a more particular embodiment, X is H. In another embodiment, X is an analyte. More particularly still, the analyte is bound through an imine. In another embodiment, the analyte is selected from the group consisting of an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a drug, a lipid, and a synthetic polymer. In another embodiment, the analyte comprises a carbonyl group. In another embodiment, the analyte comprises an aldehyde group. In another embodiment the analyte is malondialdehyde. In another embodiment the analyte is 4-hydroxynonenal.

10

15

In a more particular embodiment X and R^{14} are taken together to form a fused heteroaryl or heterocyclyl group. More particular still, X and R^{14} are taken together to form a pyrazole group.

Another aspect of the invention provides a compound of Formula VI:



VI

20

or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

G is a 5-6 membered fused aryl or heteroaryl group;

X is H or an analyte;

25

R^1 is selected from the group consisting of H, alkyl and substituted alkyl;

R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl


30

ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^3 is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, analyte substituted hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R^2 and R^3 , R^3 and R^4 , or R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.

In another embodiment, G is a 6-membered aryl group. More particularly, G is phenyl. In a more particular embodiment thereof, the compound has the formula VI(a):




VI(a)

wherein each of the variables are described herein.

In another embodiment thereof, R^1 and R^2 are taken together to form a phenyl group.

In another embodiment, R^1 is H. Alternatively, R^1 is alkyl or substituted alkyl. In another embodiment, R^3 hydrazinyl or substituted hydrazinyl. In another embodiment, R^3 is methoxy. In another embodiment, R^3 is hydroxy. In another embodiment, R^3 is amino.

In another embodiment of the invention, R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group or substituted fused heterocyclyl group of the formula:



R^{12} and R^{13} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

In another embodiment, R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are all H. In another embodiment, R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are all H. In another embodiment, R^9 is $-SO_3^-$. In another embodiment, R^9 is $-CO_2^-$. In another embodiment, X is H. In another embodiment, R^3 is methoxy and R^9 is $-CO_2^-$.


In another embodiment of the invention, X is an analyte. In another embodiment, the analyte is bound through an imine. In another embodiment, the analyte is selected from the group consisting of an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a drug, a lipid, and a synthetic polymer. In another embodiment, the analyte comprises a carbonyl group. In another embodiment, the analyte comprises an aldehyde group. In another embodiment, R^3 is analyte substituted hydrazinyl. In another embodiment, the analyte is bound through an imine. In another embodiment the analyte is malondialdehyde. In another embodiment the analyte is 4-hydroxynonenal.

Another aspect of the invention provides a composition comprising:

- (a) an analyte; and
- (b) a compound as described herein.

Another more particular aspect of the invention provides a composition comprising:

- (a) an analyte; and
 (b) a compound of Formula I:



- 5 or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
 wherein,
 X is H or an analyte;
 R¹ and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;
- 10 R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and
- 15 R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or
- 20 R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.
- 25

A more particular embodiment further comprises a buffer solution. Another embodiment provides an additional dye in solution. The additional dye may be another aldehyde/ketone sensitive dye or not. Additional dyes are well known in the art and include those described in The Handbook, A guide to Fluorescent Probes and Labeling Technologies, by R. Haugland, 10th Edition.

In another embodiment of the composition, the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus.

A more particular aspect of the composition provides any of the above embodiments, particularly those describing R¹-R¹⁰ and X.

Another aspect of the invention provides a method for determining the presence of an analyte of interest in a sample, wherein the method comprises:

contacting the sample with a compound described herein;


incubating the sample and the compound for a sufficient amount of time to form a complex between the compound and analyte;

illuminating the complex with an appropriate wavelength to form an illuminated complex; and

detecting emissions from the illuminated complex.

Another more particular aspect of the invention provides a method for determining the presence of an analyte of interest in a sample, wherein the method comprises:

contacting the sample with a compound having Formula I:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

5 X is H or an analyte;

R¹ and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group;

incubating the sample and the compound for a sufficient amount of time to form a complex between the compound and analyte;

illuminating the complex with an appropriate wavelength to form an illuminated complex; and

detecting emissions from the illuminated complex.

5

In another embodiment, the analyte and the compound are connected by a covalent bond. In another embodiment, the compound becomes fluorescent after formation of the complex. In another embodiment, the analyte is bound to the compound through an imine. In another embodiment, the analyte is selected from the group consisting of: an amino
10 acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus. In another embodiment, the analyte comprises a carbonyl group. In another embodiment, the analyte comprises an aldehyde group. In another embodiment
15 the analyte is malondialdehyde. In another embodiment the analyte is 4-hydroxynonenal.

In another embodiment, the sample comprises live cells, intracellular fluids, extracellular fluids, biological fluids, biological fermentation media, environmental sample, industrial samples, proteins, peptides, buffer solutions biological fluids or chemical reactors.

20 In another embodiment, the sample comprises blood cells, immune cells, cultured cells, muscle tissue, neurons, extracellular vesicles; vascular tissue, blood fluids, saliva, urine; water, soil, waste water, sea water; pharmaceuticals, foodstuffs or beverages. In another embodiment, the sample is immobilized on a polymeric membrane, within a polymeric gel, on a microparticle, on a microarray, on a silicon chip, on a glass slide, on a microwell plate,
25 and on a microfluidic chip.


Another aspect of the invention provides a kit for detecting an analyte in a sample, wherein the kit comprises:

a compound as describe herein that binds the analyte; and

30 instructions for detecting the analyte.

Another aspect of the invention provides a kit for detecting an analyte in a sample, wherein the kit comprises:

a compound of Formula I that binds the analyte:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

5 X is H or an analyte;

R¹ and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group; and instructions for detecting the analyte.


A more particular embodiment further comprises instructions for covalently bonding the compound to the analyte. A more particular embodiment further comprising one

or more of the following: a buffering agent, a purification medium, a vial comprising the analyte, or an organic solvent.

Another aspect of the invention provides a compound comprising a (9-phenyl-3H-xanthen-3-ylidene)hydrazine, wherein the hydrazine group is unsubstituted or substituted with an analyte. In another more particular embodiment thereof, (9-phenyl-3H-xanthen-3-ylidene)hydrazine is substituted with methoxy, amino, sulfonyl, carboxyl, sulfonylalkyl, alkyl, or a second hydrazinyl group. More particular still, the (9-phenyl-3H-xanthen-3-ylidene)hydrazine is (E)-(9-phenyl-3H-xanthene-6-yl-3-ylidene)bis(hydrazine). In another embodiment, the analyte comprises an aldehyde group.

Another aspect of the invention provides a composition comprising:

- (a) an analyte; and
- (b) a compound of Formula V:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
wherein,

X is H or an analyte;


Z is a fluorophore;

R^5 , R^6 , R^8 and R^9 are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and R^{14} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

X and R¹⁴ are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group.

Another more particular embodiment further comprises a buffer solution. In another embodiment, the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus.

Another aspect of the invention provides a method for determining the presence of an analyte of interest in a sample, wherein the method comprises: contacting the sample with compound of Formula V:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;

Z is a fluorophore;

R⁵, R⁶, R⁸ and R⁹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R¹⁴ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

X and R¹⁴ are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group;

incubating the sample and the compound for a sufficient amount of time to form a complex between the compound and analyte;

illuminating the complex with an appropriate wavelength to form an illuminated complex; and


detecting emissions from the illuminated complex.

In another embodiment, the analyte and the compound are connected by a covalent bond. In another embodiment, the compound becomes fluorescent after formation of the complex. In another embodiment, the analyte is bound to the compound through an imine. In another embodiment, the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus. In another embodiment, the analyte comprises a carbonyl group. In another embodiment, the analyte comprises an aldehyde group. In another embodiment the analyte is malondialdehyde. In another embodiment the analyte is 4-hydroxynonenal.

In another embodiment, the sample comprises live cells, intracellular fluids, extracellular fluids, biological fluids, biological fermentation media, environmental sample, industrial samples, proteins, peptides, buffer solutions biological fluids or chemical reactors. In another embodiment, the sample comprises blood cells, immune cells, cultured cells, muscle tissue, neurons, extracellular vesicles; vascular tissue, blood fluids, saliva, urine; water, soil, waste water, sea water; pharmaceuticals, foodstuffs or beverages. In another embodiment, the sample is immobilized on a polymeric membrane, within a polymeric gel, on a microparticle, on a microarray, on a silicon chip, on a glass slide, on a microwell plate, and on a microfluidic chip.

Another aspect of the invention provides a kit for detecting an analyte in a sample, wherein the kit comprises:

a compound of Formula V that binds the analyte:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

5 X is H;

Z is a fluorophore;

R⁵, R⁶, R⁸ and R⁹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

15 R¹⁴ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or


X and R¹⁴ are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group; and

20 instructions for detecting the analyte.

Another embodiment thereof further comprises instructions for covalently bonding the compound to the analyte.

25 Another embodiment thereof further comprises one or more of the following: a buffering agent, a purification medium, a vial comprising the analyte, or an organic solvent.

Another aspect of the invention provides a method of synthesizing a compound of Formula III, a tautomer thereof, a stereoisomer thereof, or a salt thereof:




or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,


- 5 R^1 and R^{11} are each independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl;

- R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;
- 15 the method comprising:

contacting a compound of formula IV:



- 20 with R^1 -NH-NH- R^{12} and R^{11} -NH-NH- R^{12} , wherein R^{12} is a protecting group, thereby obtaining a compound of Formula III(a):



III(a)

contacting R^{12} with a deprotecting reagent, thereby obtaining a compound of Formula III.

5

In another embodiment the SO_3^- group in formula III is a CO_2^- and the SO_2 group is a CO.

In another embodiment, the protecting group is Boc. In another embodiment, the deprotecting reagent comprises an acid. In another embodiment, the acid is trifluoroacetic acid (TFA).

10


Another embodiment further comprises a purifying step after obtaining the compound of Formula III(a). Another embodiment further comprises a purifying step after obtaining the compound of Formula III. In another embodiment, the purifying step comprises at least one of: column chromatography, trituration, recrystallization, filtration, or aqueous separation.

15

Another embodiment further comprises synthesizing a compound of Formula IV, comprising:

20

(c) contacting a compound of Formula IV(a):



IV(a)


with a chlorinating agent, thereby obtaining the compound of Formula IV.

In another embodiment, the chlorinating agent comprises PCl_5 or POCl_3 .

Another embodiment further comprises a purifying step after obtaining the compound of Formula IV. In another embodiment, the purifying step comprises at least one of: column chromatography, trituration, recrystallization, filtration, or aqueous separation.

One particular synthetic example is provided in Scheme 1:

10



15

Various ancillary materials will frequently be employed in an assay in accordance with the present invention. In an exemplary embodiment, buffers and/or stabilizers are present in the kit components. In another exemplary embodiment, the kits comprise indicator solutions or indicator "dipsticks", blotters, culture media, cuvettes, and the like. In yet another exemplary embodiment, the kits comprise indicator cartridges (where a kit component is bound to a solid support) for use in an automated detector. In another exemplary embodiment, the kit further comprises molecular weight markers, wherein said markers are selected from phosphorylated and non-phosphorylated polypeptides, calcium-binding and non-calcium binding polypeptides, sulfonated and non-sulfonated polypeptides, and sialylated and non-sialylated polypeptides. In another exemplary embodiment, the kit further comprises a member selected from a fixing solution, a detection reagent, a standard, a wash solution, and combinations thereof.

20

25

In an exemplary embodiment, the compounds of the invention comprise a reactive group which is a member selected from an acrylamide, an activated ester of a carboxylic acid, an acyl azide, an acyl nitrile, an aldehyde, an alkyl halide, an anhydride, an aniline, an aryl halide, an azide, an aziridine, a boronate, a carboxylic acid, a diazoalkane, a

haloacetamide, a halotriazine, a hydrazine, a hydrazide, an imido ester, an isocyanate, an isothiocyanate, a maleimide, a phosphoramidite, a reactive platinum complex, a sulfonyl halide, a thiol group, and a photoactivatable group.

These reactive groups can be covalently attached either during or after the synthesis of the dyes in order to provide reactive group-containing- dyes. In this way, reactive group-containing-dyes can be covalently attached to a wide variety of carrier molecules or solid supports that contain or are modified to contain functional groups with suitable reactivity, resulting in chemical attachment of the components. In an exemplary embodiment, the reactive group of a compound of the invention and the functional group of the carrier molecule of solid support comprise electrophiles and nucleophiles that can generate a covalent linkage between them. Alternatively, the reactive group comprises a photoactivatable group, which becomes chemically reactive only after illumination with light of an appropriate wavelength. Typically, the conjugation reaction between the reactive group and the carrier molecule/solid support results in one or more atoms of the reactive group being incorporated into a new linkage attaching the dye to the carrier molecule/solid support. Selected examples of functional groups and linkages are shown in Table 1, where the reaction of an electrophilic group and a nucleophilic group yields a covalent linkage.

Table 1: Examples of some routes to useful covalent linkages with electrophile and nucleophile reactive groups

Electrophilic Group	Nucleophilic Group	Resulting Covalent Linkage
activated esters*	amines/anilines	Carboxamides
acyl azides**	amines/anilines	Carboxamides
acyl halides	amines/anilines	Carboxamides
acyl halides	alcohols/phenols	Esters
acyl nitriles	alcohols/phenols	Esters
acyl nitriles	amines/anilines	Carboxamides
Aldehydes	amines/anilines	Imines
aldehydes or ketones	hydrazines	Hydrazones
aldehydes or ketones	hydroxylamines	Oximes
alkyl halides	amines/anilines	alkyl amines
alkyl halides	Carboxylic acids	Esters
alkyl halides	Thiols	Thioethers
alkyl halides	alcohols/phenols	Ethers
alkyl sulfonates	Thiols	Thioethers
alkyl sulfonates	Carboxylic acids	Esters

alkyl sulfonates	alcohols/phenols	Ethers
Anhydrides	alcohols/phenols	Esters
Anhydrides	amines/anilines	Carboxamides
aryl halides	Thiols	Thiophenols
aryl halides	Amines	aryl amines
Aziridines	Thiols	Thioethers
Boronates	Glycols	boronate esters
carboxylic acids	amines/anilines	Carboxamides
carboxylic acids	Alcohols	Esters
carboxylic acids	hydrazines	Hydrazides
Carbodiimides	Carboxylic acids	N-acylureas or anhydrides
Diazoalkanes	Carboxylic acids	Esters
Epoxides	Thiols	Thioethers
haloacetamides	Thiols	Thioethers
Halotriazines	amines/anilines	Aminotriazines
Halotriazines	alcohols/phenols	triazinyl ethers
imido esters	amines/anilines	Amidines
Isocyanates	amines/anilines	Ureas
Isocyanates	alcohols/phenols	Urethanes
isothiocyanates	amines/anilines	Thioureas
Maleimides	Thiols	Thioethers
phosphoramidites	Alcohols	phosphite esters
silyl halides	Alcohols	silyl ethers
sulfonate esters	amines/anilines	alkyl amines
sulfonate esters	Thiols	Thioethers
sulfonate esters	Carboxylic acids	Esters
sulfonate esters	Alcohols	Ethers
sulfonyl halides	amines/anilines	Sulfonamides
sulfonyl halides	phenols/alcohols	sulfonate esters

* Activated esters, as understood in the art, generally have the formula -CO Ω , where Ω is a good leaving group (e.g. oxysuccinimidyl (-OC₄H₄O₂) oxysulfosuccinimidyl (-OC₄H₃O₂-SO₃H), -1-oxybenzotriazolyl (-OC₆H₄N₃); or an aryloxy group or aryloxy substituted one or more times by electron withdrawing substituents such as nitro, fluoro, chloro, cyano, or trifluoromethyl, or combinations thereof, used to form activated aryl esters; or a carboxylic acid activated by a carbodiimide to form an anhydride or mixed anhydride -OCOR^a or -OCNR^aNHR^b, where R^a and R^b, which may be the same or different, are C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, or C₁-C₆ alkoxy; or cyclohexyl, 3-dimethylaminopropyl, or N-morpholinoethyl).

** Acyl azides can also rearrange to isocyanates

In one aspect, the compound comprises (in addition to the reactive hydrazine) at least one reactive group that selectively reacts with an amine group. This amine-reactive group is selected from the group consisting of succinimidyl ester, sulfonyl halide, tetrafluorophenyl ester and isothiocyanates. Thus, in one aspect, the present compounds form a covalent bond with an amine-containing molecule in a sample. In another aspect, the compound comprises at least one reactive group that selectively reacts with a thiol group. This thiol-reactive group is selected from the group consisting of maleimide, haloalkyl and haloacetamide (including any reactive groups disclosed in U.S. Patent Nos. 5,362,628; 5,352,803 and 5,573,904).

Choice of the reactive group used to attach the compound of the invention to the substance to be conjugated typically depends on the reactive or functional group on the substance to be conjugated and the type or length of covalent linkage desired. The types of functional groups typically present on the organic or inorganic substances (biomolecule or non-biomolecule) include, but are not limited to, amines, amides, thiols, alcohols, phenols, aldehydes, ketones, phosphates, imidazoles, hydrazines, hydroxylamines, disubstituted amines, halides, epoxides, silyl halides, carboxylate esters, sulfonate esters, purines, pyrimidines, carboxylic acids, olefinic bonds, or a combination of these groups. A single type of reactive site may be available on the substance (typical for polysaccharides or silica), or a variety of sites may occur (e.g., amines, thiols, alcohols, phenols), as is typical for proteins.

Typically, the reactive group will react with an amine, a thiol, an alcohol, an aldehyde, a ketone, or with silica. Preferably, reactive groups react with an amine or a thiol functional group, or with silica. In one embodiment, the reactive group is an acrylamide, an activated ester of a carboxylic acid, an acyl azide, an acyl nitrile, an aldehyde, an alkyl halide, a silyl halide, an anhydride, an aniline, an aryl halide, an azide, an aziridine, a boronate, a diazoalkane, a haloacetamide, a halotriazine, a hydrazine (including hydrazides), an imido ester, an isocyanate, an isothiocyanate, a maleimide, a phosphoramidite, a reactive platinum complex, a sulfonyl halide, or a thiol group. By "reactive platinum complex" is particularly meant chemically reactive platinum complexes such as described in U.S. Patent No. 5,714,327.

Where the reactive group is an activated ester of a carboxylic acid, such as a succinimidyl ester of a carboxylic acid, a sulfonyl halide, a tetrafluorophenyl ester or an isothiocyanates, the resulting compound is particularly useful for preparing conjugates of carrier molecules such as proteins, nucleotides, oligonucleotides, or haptens. Where the reactive group is a maleimide, haloalkyl or haloacetamide (including any reactive groups

disclosed in U.S. Patent Nos. 5,362,628; 5,352,803 and 5,573,904 (*supra*)) the resulting compound is particularly useful for conjugation to thiol-containing substances. Where the reactive group is a hydrazide, the resulting compound is particularly useful for conjugation to periodate-oxidized carbohydrates and glycoproteins, and in addition is an aldehyde-fixable polar tracer for cell microinjection. Where the reactive group is a silyl halide, the resulting compound is particularly useful for conjugation to silica surfaces, particularly where the silica surface is incorporated into a fiber optic probe subsequently used for remote ion detection or quantification.

In a particular aspect, the reactive group is a photoactivatable group such that the group is only converted to a reactive species after illumination with an appropriate wavelength. An appropriate wavelength is generally a UV wavelength that is less than 400 nm. This method provides for specific attachment to only the target molecules, either in solution or immobilized on a solid or semi-solid matrix. In this way, present dye compounds that comprise a photoactivatable reactive group associate with anionic proteins and can be covalently conjugated to the proteins. Photoactivatable reactive groups include, without limitation, benzophenones, aryl azides and diazirines.

Preferably, the reactive group is a photoactivatable group, succinimidyl ester of a carboxylic acid, a haloacetamide, haloalkyl, a hydrazine, an isothiocyanate, a maleimide group, an aliphatic amine, a silyl halide, a cadaverine or a psoralen. More preferably, the reactive group is a succinimidyl ester of a carboxylic acid, a maleimide, an iodoacetamide, or a silyl halide. In a particular embodiment the reactive group is a succinimidyl ester of a carboxylic acid, a sulfonyl halide, a tetrafluorophenyl ester, an isothiocyanates or a maleimide.

The selection of a covalent linkage to attach the reporter molecule to the carrier molecule or solid support typically depends on the chemically reactive group on the component to be conjugated. The discussion regarding reactive groups in the section immediately preceding is relevant here as well. In addition to the hydrazinyl appendage, exemplary reactive groups typically present on the biological or non-biological components include, but are not limited to, amines, thiols, alcohols, phenols, aldehydes, ketones, phosphates, imidazoles, hydrazines, hydroxylamines, disubstituted amines, halides, epoxides, sulfonate esters, purines, pyrimidines, carboxylic acids, or a combination of these groups. A single type of reactive site may be available on the component (typical for polysaccharides), or a variety of sites may occur (e.g. amines, thiols, alcohols, phenols), as is typical for proteins. A carrier molecule or solid support may be conjugated to more than one reporter molecule, which may be the same or different, or to a substance that is additionally modified by a hapten. Although some selectivity can be obtained by careful

control of the reaction conditions, selectivity of labeling is best obtained by selection of an appropriate reactive compound.

In another exemplary embodiment, the dye is covalently bound to a carrier molecule. If the compound has a reactive group, then the carrier molecule can alternatively
5 be linked to the compound through the reactive group. The reactive group may contain both a reactive functional moiety and a linker, or only the reactive functional moiety.

A variety of carrier molecules are useful in the present invention. Exemplary carrier molecules include antigens, steroids, vitamins, drugs, haptens, metabolites, toxins, environmental pollutants, amino acids, peptides, proteins, nucleic acids, nucleic acid
10 polymers, carbohydrates, lipids, and polymers.

In an exemplary embodiment, the carrier molecule comprises an amino acid, a peptide, a protein, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a polymeric microparticle, a biological cell, a virus and combinations thereof. In
15 another exemplary embodiment, the carrier molecule is selected from a hapten, a nucleotide, an oligonucleotide, a nucleic acid polymer, a protein, a peptide or a polysaccharide. In another exemplary embodiment, at least one member selected from R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} and R^{29} comprise a carrier molecule. In another exemplary
20 embodiment one of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} and R^{29} comprises a carrier group bound through a substituted alkyl group or reactive group, such as an –alkyl-succinamidyl group.

In an exemplary embodiment, the carrier molecule comprises an amino acid, a peptide, a protein, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a
25 nucleic acid, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a polymeric microparticle, a biological cell, a virus and combinations thereof. In another exemplary embodiment, the carrier molecule is selected from a hapten, a nucleotide, an oligonucleotide, a nucleic acid polymer, a protein, a peptide or a polysaccharide. In a preferred embodiment the carrier molecule is amino acid, a peptide, a
30 protein, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a tyramine, a synthetic polymer, a polymeric microparticle, a biological cell, cellular components, an ion chelating moiety, an enzymatic substrate or a virus. In another preferred embodiment, the carrier molecule is an antibody or fragment thereof, an antigen, an avidin or streptavidin, a biotin, a
35 dextran, an antibody binding protein, a fluorescent protein, agarose, and a non-biological

microparticle. Typically, the carrier molecule is an antibody, an antibody fragment, antibody-binding proteins, avidin, streptavidin, a toxin, a lectin, or a growth factor. Preferred haptens include biotin, digoxigenin and fluorophores.

Antibody binding proteins include, but are not limited to, protein A, protein G,
5 soluble Fc receptor, protein L, lectins, anti-IgG, anti-IgA, anti-IgM, anti-IgD, anti-IgE or a fragment thereof.

In an exemplary embodiment, the enzymatic substrate is selected from an amino acid, peptide, sugar, alcohol, alkanolic acid, 4-guanidinobenzoic acid, nucleic acid, lipid, sulfate, phosphate, $-\text{CH}_2\text{OCOalkyl}$ and combinations thereof. Thus, the enzyme
10 substrates can be cleaved by enzymes selected from the group consisting of peptidase, phosphatase, glycosidase, dealkylase, esterase, guanidinobenzotase, sulfatase, lipase, peroxidase, histone deacetylase, endoglycoceramidase, exonuclease, reductase and endonuclease.

In another exemplary embodiment, the carrier molecule is an amino acid
15 (including those that are protected or are substituted by phosphates, carbohydrates, or C_1 to C_{22} carboxylic acids), or a polymer of amino acids such as a peptide or protein. In a related embodiment, the carrier molecule contains at least five amino acids, more preferably 5 to 36 amino acids. Exemplary peptides include, but are not limited to, neuropeptides, cytokines, toxins, protease substrates, and protein kinase substrates. Other exemplary peptides may
20 function as organelle localization peptides, that is, peptides that serve to target the conjugated compound for localization within a particular cellular substructure by cellular transport mechanisms. Preferred protein carrier molecules include enzymes, antibodies, lectins, glycoproteins, histones, albumins, lipoproteins, avidin, streptavidin, protein A, protein G, phycobiliproteins and other fluorescent proteins, hormones, toxins and growth factors.
25 Typically, the protein carrier molecule is an antibody, an antibody fragment, avidin, streptavidin, a toxin, a lectin, or a growth factor. Exemplary haptens include biotin, digoxigenin and fluorophores.

In another exemplary embodiment, the carrier molecule comprises a nucleic acid base, nucleoside, nucleotide or a nucleic acid polymer, optionally containing an
30 additional linker or spacer for attachment of a fluorophore or other ligand, such as an alkynyl linkage (U.S. Pat. No. 5,047,519), an aminoalkyl linkage (U.S. Pat. No. 4,711,955) or other linkage. In another exemplary embodiment, the nucleotide carrier molecule is a nucleoside or a deoxynucleoside or a dideoxynucleoside.

Exemplary nucleic acid polymer carrier molecules are single- or multi-
35 stranded, natural or synthetic DNA or RNA oligonucleotides, or DNA/RNA hybrids, or

incorporating an unusual linker such as morpholine derivatized phosphates (AntiVirals, Inc., Corvallis OR), or peptide nucleic acids such as *N*-(2-aminoethyl)glycine units, where the nucleic acid contains fewer than 50 nucleotides, more typically fewer than 25 nucleotides.

5 In another exemplary embodiment, the carrier molecule comprises a carbohydrate or polyol that is typically a polysaccharide, such as dextran, FICOLL, heparin, glycogen, amylopectin, mannan, inulin, starch, agarose and cellulose, or is a polymer such as a poly(ethylene glycol). In a related embodiment, the polysaccharide carrier molecule includes dextran, agarose or FICOLL.

10 In another exemplary embodiment, the carrier molecule comprises a lipid (typically having 6–25 carbons), including glycolipids, phospholipids, and sphingolipids. Alternatively, the carrier molecule comprises a lipid vesicle, such as a liposome, or is a lipoprotein (see below). Some lipophilic substituents are useful for facilitating transport of the conjugated dye into cells or cellular organelles.

15 Alternatively, the carrier molecule is cells, cellular systems, cellular fragments, or subcellular particles. Examples of this type of conjugated material include virus particles, bacterial particles, virus components, biological cells (such as animal cells, plant cells, bacteria, or yeast), or cellular components. Examples of cellular components that can be labeled, or whose constituent molecules can be labeled, include but are not limited to lysosomes, endosomes, cytoplasm, nuclei, histones, mitochondria, Golgi apparatus,
20 endoplasmic reticulum and vacuoles.

In another embodiment the carrier molecule is a metal chelating moiety. While any chelator that binds a metal ion of interest and gives a change in its fluorescence properties is a suitable conjugate, preferred metal chelating moieties are crown ethers, including diaryldiaza crown ethers, as described in U.S. Pat. No. 5,405,975 to Kuhn et al.
25 (1995); derivatives of 1,2-bis-(2-aminophenoxyethane)-*N,N,N',N'*-tetraacetic acid (BAPTA), as described in U.S. Pat. No. 5,453,517 to Kuhn et al. (1995) (incorporated by reference) and U.S. Pat. No. 5,049,673 to Tsien et al. (1991); derivatives of 2-carboxymethoxy-aniline-*N,N*-diacetic acid (APTRA), as described by Ragu et al., *Am. J. Physiol.*, 256: C540 (1989); and pyridyl-based and phenanthroline metal ion chelators, as described in U.S. Pat. No.
30 5,648,270 to Kuhn et al. (1997).

Aldehyde/ketone-sensing conjugates of the invention are optionally prepared in chemically reactive forms and further conjugated to polymers such as dextrans to improve their utility as sensors as described in U.S. Pat. Nos. 5,405,975 and 5,453,517.

35 In another exemplary embodiment, the carrier molecule non-covalently associates with organic or inorganic materials. Exemplary embodiments of the carrier

molecule that possess a lipophilic substituent can be used to target lipid assemblies such as biological membranes or liposomes by non-covalent incorporation of the dye compound within the membrane, *e.g.*, for use as probes for membrane structure or for incorporation in liposomes, lipoproteins, films, plastics, lipophilic microspheres or similar materials.

- 5 In an exemplary embodiment, the carrier molecule comprises a specific binding pair member wherein the present compounds are conjugated to a specific binding pair member and are used to detect an analyte in a sample. Alternatively, the presence of the labeled specific binding pair member indicates the location of the complementary member of that specific binding pair; each specific binding pair member having an area on
- 10 the surface or in a cavity which specifically binds to, and is complementary with, a particular spatial and polar organization of the other. Exemplary binding pairs are set forth in Table 2.

Table 2. Representative Specific Binding Pairs

Antigen	Antibody
Biotin	avidin (or streptavidin or anti-biotin)
IgG*	protein A or protein G
Drug	drug receptor
Folate	folate binding protein
Toxin	toxin receptor
Carbohydrate	lectin or carbohydrate receptor
Peptide	peptide receptor
Protein	protein receptor
enzyme substrate	Enzyme
DNA (RNA)	cDNA (cRNA)†
Hormone	hormone receptor
Ion	Chelator
Antibody	antibody-binding proteins

* IgG is an immunoglobulin

- 15 † cDNA and cRNA are the complementary strands used for hybridization

Analytes:

- Analytes of interest are preferably bound through a hydrazinyl group on the dye or a reporter moiety bound to the dye. The analytes include: an amino acid, a peptide, a
- 20 protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus. In another embodiment, the analyte comprises a carbonyl group. In another embodiment, the analyte comprises an aldehyde or ketone group. In another embodiment
- 25 the analyte is malondialdehyde. In another embodiment the analyte is 4-hydroxynonenal.

In one embodiment the analyte described herein is a result of oxidative stress, such as from interaction with a superoxide, hydroxy, peroxy, alkoxy, hydroperoxy, hydrogenperoxide, hypochlorous acid, ozone, a singlet oxygen, or peroxynitrite. In another embodiment the analyte is indicative of activity or inactivity of superoxide dismutase, catalase, glutathione peroxidase or substrates relating thereto, including, vitamin A, E, ascorbate or glutathione. In another embodiment, the analyte is a biomarker for a disease such as alzheimers, parkinsons, atherosclerosis, multiple sclerosis, or cancer.

In a preferred embodiment the analyte is a result of a ROS (reactive oxygen species) pathway. In another embodiment the analyte is a product of lipid peroxidation. More particularly, the analyte is malonidialdehyde.

In another embodiment, the sample comprises live cells, intracellular fluids, extracellular fluids, biological fluids, biological fermentation media, environmental sample, industrial samples, proteins, peptides, buffer solutions biological fluids or chemical reactors. In another embodiment, the sample comprises blood cells, immune cells, cultured cells, muscle tissue, neurons, extracellular vesicles; vascular tissue, blood fluids, saliva, urine; water, soil, waste water, sea water; pharmaceuticals, foodstuffs or beverages. In another embodiment, the sample is immobilized on a polymeric membrane, within a polymeric gel, on a microparticle, on a microarray, on a silicon chip, on a glass slide, on a microwell plate, and on a microfluidic chip.

Illumination

The sample or medium in which the aldehyde/ketone sensitive dye is present is illuminated with a wavelength of light selected to give a detectable optical response, and observed with a means for detecting the optical response. Equipment that is useful for illuminating the present compounds and compositions of the invention includes, but is not limited to, hand-held ultraviolet lamps, mercury arc lamps, xenon lamps, lasers and laser diodes. These illumination sources are optically integrated into laser scanners, fluorescence microplate readers or standard or microfluorometers.


The dyes of the invention may, at any time after or during an assay, be illuminated with a wavelength of light that results in a detectable optical response, and observed with a means for detecting the optical response. Upon illumination, such as by an ultraviolet or visible wavelength emission lamp, an arc lamp, a laser, or even sunlight or ordinary room light, the fluorescent compounds, including those bound to the complementary

specific binding pair member, display intense visible absorption as well as fluorescence emission. Selected equipment that is useful for illuminating the fluorescent compounds of the invention includes, but is not limited to, hand-held ultraviolet lamps, mercury arc lamps, xenon lamps, argon lasers, laser diodes, and YAG lasers. These illumination sources are optionally integrated into laser scanners, fluorescence microplate readers, standard or mini fluorometers, or chromatographic detectors. This fluorescence emission is optionally detected by visual inspection, or by use of any of the following devices: CCD cameras, video cameras, photographic film, laser scanning devices, fluorometers, photodiodes, quantum counters, epifluorescence microscopes, scanning microscopes, flow cytometers, fluorescence microplate readers, or by means for amplifying the signal such as photomultiplier tubes. Where the sample is examined using a flow cytometer, a fluorescence microscope or a fluorometer, the instrument is optionally used to distinguish and discriminate between the fluorescent compounds of the invention and a second fluorophore with detectably different optical properties, typically by distinguishing the fluorescence response of the fluorescent compounds of the invention from that of the second fluorophore. Where a sample is examined using a flow cytometer, examination of the sample optionally includes isolation of particles within the sample based on the fluorescence response by using a sorting device. In another embodiment, the illumination source is used to form a covalent bond between the present dye and an analyte of interest. In this instance the dye comprises a photoactivatable reactive group, such as those discussed above.

A detailed description of the invention having been provided above, the following examples are given for the purpose of illustrating the invention and shall not be construed as being a limitation on the scope of the invention or claims.

EXAMPLES


Example 1: Compound 1



A mixture of 3,6-dichlorosulfofluoran (800 mg, 1.97 mmol) and *tert*-butyl carbazate (1.56 g, 11.82 mmol) in 15 ml of dry acetonitrile was stirred at 65 °C for 3 hours. The reaction mixture was then cooled to room temperature and the resulting orange-red solid collected by filtration to give 850 mg of Compound 1.

5


Example 2: Compound 2



To an ice-cooled solution of 2 ml of trifluoroacetic acid was added Compound 1 (200 mg, 0.335 mmol) in a small portion over a 5-minute period. After stirring the ice-cooled temperature for 30 minutes, chloroform (about 10 ml) was added to the reaction mixture. The resulting orange-red solid was collected by filtration, washed with ether and dried under vacuum to give 160 mg of Compound 2.

15


Example 3: Compound 3



A mixture of 3,6-dichlorosulfofluoran (500 mg, 1.23 mmol) and *tert*-butyl carbazate (650 mg, 4.93 mmol) in 10 ml of dry acetonitrile was stirred at room temperature for 12 hours. After the reaction mixture was concentrated *in vacuo*, the resulting crude mixture was purified by column chromatography on silica gel eluting with 5/95 V/V methanol-chloroform to give 350 mg of Compound 3 orange solid.


25

Example 4: Compound 4



- 5 To a solution of Compound 3 (200 mg, 0.40 mmol) in 5 ml of dry N,N-dimethylformamide was added 1 ml of sodium methoxide solution in methanol (0.5 M) and the mixture was stirred at approximately 65 °C under nitrogen atmosphere for 5 hours. The resulting reaction mixture was cooled to room temperature and concentrated *in vacuo*. The crude product was then purified by column chromatography on silica gel eluting with 7 %
- 10 methanol in chloroform to give 125 mg of Compound 4 as orange solid.


Example 5: Compound 5



- 15 To an ice-cold solution of 1 ml of trifluoroacetic acid was added Compound 4 (150 mg, 0.30 mmol) in a small portion over a 5-minute period. After stirring for 30 minutes, the resulting solid was collected by filtration, washed with ether and dried *in vacuo* to give 72 mg of Compound 5 as orange solid.

20

Example 6: Compound 6




25

- To a solution of *tert*-butyl carbazate (2 g, 15 mmol) in 10 ml acetonitrile was added 1,3-propanesulfone (0.67 ml, 7.5 mmol) and the mixture was stirred at room temperature for


two days. All the solvent was then removed *in vacuo* and the resulting crude product was purified by column chromatography on silica gel eluting with 25 % methanol in chloroform to give 1.8 g of Compound **6** as off-white solid.

5 Example 7: Compound **7**




10 A sample of 3,6-dichlorofluoran (100 mg, 0.25 mmol) and Compound **6** (156 mg, 0.62 mmol) were dissolved in a mixture of 10 ml of acetonitrile and 4 ml of N,N-dimethylformamide. After stirring at 65 °C for 2 hours, all the solvent was removed *in vacuo* and the resulting crude material was purified by column chromatography over LH-20 Sephadex eluting with water to give 80 mg of Compound **7** as orange-red solid.

15 Example 8: Compound **8**




20 To an ice-cold solution of 1 ml of trifluoroacetic acid was added Compound **7** (50 mg, 0.06 mmol) in a small portion over a 5-minute period. After stirring at ice-cold temperature for one hour, chloroform (about 10 ml) was added to the reaction mixture. The resulting orange-red solid was collected by filtration, washed with ether and dried under vacuum to give 26 mg of Compound **8** as orange-red solid.

Example 9: Compound **9**

5

To a solution of tert-butylcarbazate (250 mg, 1.9 mmol) in 10 ml of acetonitrile was added a suspension of 4-bromobutyltriphenylphosphonium bromide (95 mg, 0.63 mmol) and the mixture was heated under reflux for 12 hours. All the solvent was removed *in vacuo* and the resulting crude product was purified by column chromatography over silica gel to

10 give 250 mg of Compound **9** an off-white solid.


Example 10: Compound **10**

15

To a solution of 3,6-dichlorosulfoluoran (68 mg, 0.17 mmol) in 1 ml of N,N-dimethylformamide was added Compound **9** (90 mg, 0.17 mmol). After stirring at room temperature for 30 minutes, the mixture was concentrated *in vacuo* and the resulting crude product was purified by column chromatography over silica gel eluting with 20 % methanol in

20 chloroform to give 120 mg of Compound **10** as orange solid.


Example 11: Compound 11



5 To a solution of Compound **10** (110 mg, 0.14 mmol) in 3 ml of dichloromethane was added dibutylamine (30 μ l, 2 mmol) and the mixture was stirred near 60 ° C for 12 hours. After removing all the solvent under vacuum, the resulting crude product was purified by column chromatography over silica gel eluting with 10 % methanol in chloroform to give 68 mg of Compound **11** as orange-red solid.







10






Example 12: Compound 12





15 To an ice-cold solution of Compound **11** (20 mg, 0.02 mmol) in 1 ml of chloroform was added 250 μ l of trifluoroacetic acid and the mixture was stirred at room temperature for 2 hours. All the solvent was removed under vacuum and the crude product was purified by column chromatography over silica gel eluting with water/methanol/chloroform (2:30:68 v/v/v) to give 10 mg of Compound **12** as red solid.

Compound Table:

Compound No.	Structure
2	
5	
8	
13	
14	
15	

16	
17	
18	
19	
20	

21	
22	


Example 13: **Aldehyde Reactivity of Compounds.**

A sample of Compound **2** (5 mg, 0.0126 mmol) is dissolved in 10 ml of pH 6.5, 50
 5 mmol phosphate buffer solution. To this non-fluorescent solution, is added 25 ul, 1 mol
 solution of acetaldehyde in water at room temperature. Within a few minutes, the solution
 becomes highly fluorescent. The reaction is complete in 10 minutes to give a stable oxime
 analog. Its absorption max is at 548 nm and emission max is at 580 nm (see Fig. 3A and B).

Each of the above-cited references are hereby incorporated by reference as if
 set forth fully herein.

CLAIMS

1. A compound of Formula I:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;


R¹ and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, analyte substituted hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.


2. The compound of claim 1, wherein R^1 and R^{1a} are both H.
3. The compound of claim 1, wherein R^1 is alkyl or substituted alkyl; and R^{1a} is H.
4. The compound of claim 1, wherein R^3 is hydrazinyl or substituted hydrazinyl.
5. The compound of claim 1, wherein R^3 is methoxy or hydroxy.
6. The compound of claim 1, wherein R^3 is amino.
7. The compound of claim 1, wherein R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group or substituted fused heterocyclyl group of the formula:



R^{12} and R^{13} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

8. The compound of claim 1, wherein R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are all H.
9. The compound of claim 1, wherein R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are all H.
10. The compound of claim 1, wherein R^9 is $-SO_3^-$.
11. The compound of claim 1, wherein R^9 is $-CO_2^-$.
12. The compound of claim 1, wherein X is H.
13. The compound of claim 1, wherein X is an analyte.

14. The compound of claim 13, wherein the analyte is bound through an imine.
15. The compound of claim 13, wherein the analyte is selected from the group consisting of an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a drug, a lipid, and a synthetic polymer.
16. The compound of claim 13, wherein the analyte is malondialdehyde or 4-hydroxynonenal.
17. The compound of claim 13, wherein the analyte comprises an aldehyde group.
18. The compound of claim 1, wherein R^3 is analyte substituted hydrazinyl.
19. The compound of claim 18, wherein the analyte is bound through an imine.
20. The compound of claim 18, wherein the analyte is selected from the group consisting of an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a drug, a lipid, and a synthetic polymer.
21. The compound of claim 18, wherein the analyte comprises a carbonyl group.
22. The compound of claim 18, wherein the analyte comprises an aldehyde group.
23. The compound of claim 1, wherein R^3 is methoxy and R^9 is $-\text{CO}_2^-$.
24. A salt of the compound of claim 1.
25. A compound of Formula II:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;

Y is H or an analyte;

L is -NH- or a covalent bond;

R¹ is selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R¹¹ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl.

26. The compound of claim 25, wherein R⁹ is SO₃⁻.


27. The compound of claim 25, wherein R⁹ is CO₂⁻.

28. The compound of claim 25, wherein R², R⁴, R⁵, R⁶, R⁷, R⁸ and R¹⁰ are H.

29. The compound of claim 25, wherein R¹ and R¹¹ are H.

30. The compound of claim 25, wherein R¹ is alkyl or substituted alkyl and R¹¹ is alkyl or substituted alkyl.

31. The compound of claim 31 wherein, R^1 and R^{11} are $-(CH_2)_3SO_3^-$.
32. The compound of claim 25, wherein X and Y are H.
33. The compound of claim 25, wherein X is an analyte and Y is an analyte.
34. The compound of claim 25, wherein L is -NH-.
35. The compound of claim 25, wherein L is a covalent bond.
36. A compound of Formula III:



III

or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
wherein,


R^1 is selected from the group consisting of H, alkyl and substituted alkyl;

R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and


R^{11} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl.

37. The compound of claim 36, wherein R^2 , R^4 , R^5 , R^6 , R^7 , R^8 and R^{10} are hydrogen.

38. The compound of claim 36, wherein R^1 and R^{11} are H.
39. The compound of claim 36, wherein R^1 is alkyl or substituted alkyl and R^{11} is alkyl or substituted alkyl.
40. The compound of claim 39, wherein R^1 and R^{11} are both $-(CH_2)_3SO_3^-$.
41. A compound selected from the group consisting of:



42. A compound of Formula V:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;

Z¹ a fluorophore and Z² is R⁸; or

Z¹ is R⁸ and Z² is a fluorophore;


R⁵, R⁶, R⁸ and R⁹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R¹⁴ is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

X and R¹⁴ are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group.

43. The compound of claim 1, wherein one of Z¹ or Z² is selected from the group consisting of a xanthene, an indole and a borapolyazaindachine.

44. The compound of claim 1, wherein Z¹ or Z² is:



wherein,

R^2 , R^4 and R^{10} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^3 is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, substituted hydrazinyl, anlyte substituted hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R^2 and R^3 , R^3 and R^4 , or R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group;

R^{15} is selected from the group consisting of $=O$ and $=^{(+)}NR^{16}R^{17}$; and

R^{16} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, heteroaryl, and substituted heteroaryl; and

R^{17} is selected from the group consisting of alkyl, substituted alkyl, amino, and substituted amino.

45. The compound of claim 44, wherein R^{15} is $=O$.


46. The compound of claim 44, wherein R^{15} is $=^{(+)}NR^{16}R^{17}$.

47. The compound of claim 46, wherein R^{16} and R^{17} are methyl.

48. The compound of claim 44, wherein R^2 , R^4 and R^{10} are H.


49. The compound of claim 44, wherein R^3 is amino or substituted amino.

50. The compound of claim 44, wherein R^3 is hydroxy.
51. The compound of claim 44, wherein R^3 is $-N(CH_3)_2$.
52. The compound of claim 44, wherein R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group or substituted fused heterocyclyl group of the following structure:



wherein, R^{12} and R^{13} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

53. The compound of claim 42, wherein Z^1 or Z^2 is:




R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, $-SO_3H$, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

54. The compound of claim 53, wherein R^{19} and R^{22} are H.
55. The compound of claim 53, wherein R^{18} , R^{20} , R^{21} and R^{23} are methyl.

56. The compound of claim 53, wherein R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} are H.

57. The compound of claim 42, wherein Z^1 or Z^2 is:



R^{24} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^{25} , R^{26} , R^{27} , R^{28} and R^{29} are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, $-SO_3H$, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

58. The compound of claim 57, wherein R^{24} , R^{25} , R^{27} , R^{28} and R^{29} are H.

59. The compound of claim 57, wherein R^{26} is a carboxyl ester.

60. The compound of claim 57, wherein R^{26} is $-CO_2CH_3$.

61. The compound of claim 42, wherein R^5 , R^6 , R^8 and R^{14} are all H.


62. The compound of claim 42, wherein R^9 is $-SO_3^-$.

63. The compound of claim 42, wherein R^9 is $-CO_2^-$.

64. The compound of claim 42, wherein X is H.

65. The compound of claim 42, wherein X is an analyte.

66. The compound of claim 65, wherein the analyte is bound through an imine.
67. The compound of claim 65, wherein the analyte is selected from the group consisting of an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a drug, a lipid, and a synthetic polymer.
68. The compound of claim 65, wherein the analyte comprises a carbonyl group.
69. The compound of claim 65, wherein the analyte comprises an aldehyde group.
70. The compound of claim 42, wherein X and R¹⁴ are taken together to form a fused heteroaryl group.
71. The compound of claim 42, wherein X and R¹⁴ are taken together to form a pyrazole group.
72. A composition comprising:
 (a) an analyte; and
 (b) a compound of Formula I:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H;

R¹ and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy,

amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^3 is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or


R^2 and R^3 , R^3 and R^4 , or R^2 , R^3 and R^4 are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group.

73. The composition of claim 72, further comprising a buffer solution.

74. The composition of claim 72, wherein the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus.

75. A method for determining the presence of an analyte of interest in a sample, wherein the method comprises:

contacting the sample with a compound having Formula I:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H;

R¹ and R^{1a} are independently selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group;

incubating the sample and the compound for a sufficient amount of time to form a complex between the compound and analyte;

illuminating the complex with an appropriate wavelength to form an illuminated complex; and

detecting emissions from the illuminated complex.

76. The method of claim 75, wherein the analyte and the compound are connected by a covalent bond.

77. The method of claim 75, wherein the compound becomes fluorescent after formation of the complex.

78. The method of claim 76, wherein the analyte is bound to the compound through an imine.

79. The method of claim 75, wherein the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus.

80. The method of claim 75, wherein the analyte is malondialdehyde or 4-hydroxynonenal.


81. The method of claim 75, wherein the analyte comprises an aldehyde group.

82. The method of claim 75, wherein the sample comprises live cells, intracellular fluids, extracellular fluids, biological fluids, biological fermentation media, environmental sample, industrial samples, proteins, peptides, buffer solutions biological fluids or chemical reactors.

83. The method of claim 75, wherein the sample comprises blood cells, immune cells, cultured cells, muscle tissue, neurons, extracellular vesicles; vascular tissue, blood fluids, saliva, urine; water, soil, waste water, sea water; pharmaceuticals, foodstuffs or beverages.

84. The method of claim 75, wherein the sample is immobilized on a polymeric membrane, within a polymeric gel, on a microparticle, on a microarray, on a silicon chip, on a glass slide, on a microwell plate, and on a microfluidic chip.

85. A kit for detecting an analyte in a sample, wherein the kit comprises:
a compound of Formula I that binds the analyte:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H;

R¹ is selected from the group consisting of H, alkyl and substituted alkyl;

R², R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R³ is selected from the group consisting of alkoxy, substituted alkoxy, amino, substituted amino, hydrazinyl, alkyl, substituted alkyl, acyl, acylamino, acyloxy, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; or

R² and R³, R³ and R⁴, or R², R³ and R⁴ are taken together to form a fused heterocyclyl group, a fused substituted heterocyclyl group, a fused aryl group, a fused substituted aryl group, a fused heteroaryl group or a fused substituted heteroaryl group; and instructions for detecting the analyte.

86. The kit of claim 85, further comprising instructions for covalently bonding the

compound to the analyte.

87. The kit of claim 85, further comprising one or more of the following: a buffering agent, a purification medium, a vial comprising the analyte, or an organic solvent.

88. A compound comprising a (9-phenyl-3H-xanthen-3-ylidene)hydrazine, wherein the hydrazine group is unsubstituted or substituted with an analyte.

89. The compound of claim 88, wherein the (9-phenyl-3H-xanthen-3-ylidene)hydrazine is substituted with methoxy, amino, sulfonyl, carboxyl, sulfonylalkyl, alkyl, or a second hydrazinyl group.


90. The compound of claim 88, wherein the (9-phenyl-3H-xanthen-3-ylidene)hydrazine is (E)-(9-phenyl-3H-xanthene-6-yl-3-ylidene)bis(hydrazine).

91. The compound of claim 88, wherein the analyte comprises an aldehyde group.

92. A composition comprising:

(a) an analyte; and

(b) a compound of Formula V:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;

Z is a fluorophore;

R⁵, R⁶, R⁸ and R⁹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl

ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^{14} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or


X and R^{14} are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group.

93. The composition of claim 92, further comprising a buffer solution.

94. The composition of claim 92, wherein the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus.

95. A method for determining the presence of an analyte of interest in a sample, wherein the method comprises:

contacting the sample with compound of Formula V:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H or an analyte;

Z is a fluorophore;

R^5 , R^6 , R^8 and R^9 are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl

ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

R^{14} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

X and R^{14} are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group;

incubating the sample and the compound for a sufficient amount of time to form a complex between the compound and analyte;

illuminating the complex with an appropriate wavelength to form an illuminated complex; and

detecting emissions from the illuminated complex.

96. The method of claim 95, wherein the analyte and the compound are connected by a covalent bond.

97. The method of claim 95, wherein the compound becomes fluorescent after formation of the complex.

98. The method of claim 96, wherein the analyte is bound to the compound through an imine.

99. The method of claim 95, wherein the analyte is selected from the group consisting of: an amino acid, a peptide, a protein, a carbohydrate, a polysaccharide, a nucleoside, a nucleotide, an oligonucleotide, a nucleic acid polymer, a hapten, a psoralen, a drug, a hormone, a lipid, a lipid assembly, a synthetic polymer, a biotin-binding protein, a polymeric microparticle, a biological cell and a virus.

100. The method of claim 95, wherein the analyte is malondialdehyde or 4-hydroxynonenal.


101. The method of claim 95, wherein the analyte comprises an aldehyde group.

102. The method of claim 95, wherein the sample comprises live cells, intracellular fluids, extracellular fluids, biological fluids, biological fermentation media, environmental sample, industrial samples, proteins, peptides, buffer solutions biological fluids or chemical reactors.

103. The method of claim 95, wherein the sample comprises blood cells, immune cells, cultured cells, muscle tissue, neurons, extracellular vesicles; vascular tissue, blood fluids, saliva, urine; water, soil, waste water, sea water; pharmaceuticals, foodstuffs or beverages.

104. The method of claim 95, wherein the sample is immobilized on a polymeric membrane, within a polymeric gel, on a microparticle, on a microarray, on a silicon chip, on a glass slide, on a microwell plate, and on a microfluidic chip.

105. A kit for detecting an analyte in a sample, wherein the kit comprises:
a compound of Formula V that binds the analyte:



or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;

wherein,

X is H;

Z is a fluorophore;

R^5 , R^6 , R^8 and R^9 are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO_3^- , sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and


R^{14} is selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl; or

X and R¹⁴ are taken together to form a fused heterocyclyl group, substituted fused heterocyclyl group, fused heteroaryl group, or substituted fused heteroaryl group; and instructions for detecting the analyte.

106. The kit of claim 105, further comprising instructions for covalently bonding the compound to the analyte.

107. The kit of claim 105, further comprising one or more of the following: a buffering agent, a purification medium, a vial comprising the analyte, or an organic solvent.

108. A method of synthesizing a compound of Formula III, a tautomer thereof, a stereoisomer thereof, or a salt thereof:




or a stereoisomer, tautomer, hydrate, solvate, or salt thereof;
wherein,

R¹ and R¹¹ are each independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, substituted heterocyclyl, cycloalkyl, and substituted cycloalkyl;


R², R⁴, R⁵, R⁶, R⁷, R⁸ and R¹⁰ are each independently selected from the group consisting of H, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, halo, hydroxy, nitro, SO₃⁻, sulfonyl, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, substituted alkylthio, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

the method comprising:

contacting a compound of formula IV:



with R^1 -NH-NH- R^{12} and R^{11} -NH-NH- R^{12} , wherein R^{12} is a protecting group, thereby obtaining a compound of Formula III(a):




contacting R^{12} with a deprotecting reagent, thereby obtaining a compound of Formula III.

109. The method of claim 108, wherein the protecting group is Boc.
110. The method of claim 109, wherein the deprotecting reagent comprises an acid.
111. The method of claim 110, wherein the acid is trifluoroacetic acid (TFA).
112. The method of claim 108, further comprising a purifying step after obtaining the compound of Formula III(a).
113. The method of claim 108, further comprising a purifying step after obtaining the compound of Formula III.
114. The method of claim 112, wherein the purifying step comprises at least one of: column chromatography, trituration, recrystallization, filtration, or aqueous separation.

115. The method of claim 108, further comprising synthesizing a compound of Formula IV, comprising:

(c) contacting a compound of Formula IV(a):



IV(a)

with a chlorinating agent, thereby obtaining the compound of Formula IV.


116. The method of claim 115, wherein the chlorinating agent comprises PCl₅ or POCl₃.

117. The method of claim 115, further comprising a purifying step after obtaining the compound of Formula IV.

118. The method of claim 117, wherein the purifying step comprises at least one of: column chromatography, trituration, recrystallization, filtration, or aqueous separation.


1/3

FIG. 1



2/3

FIG. 2



3/3

FIG. 3A





FIG. 3B



INTERNATIONAL SEARCH REPORT

International application No

PCT/US2009/031830

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D311/82 C07D471/06 C07F9/28 C07F5/02 C09B11/24
G01N33/533

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D C07F C09B G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/64988 A (MOLECULAR PROBES INC [US]) 2 November 2000 (2000-11-02) the whole document	1-41, 72-91, 108-118
A	US 2005/170363 A1 (REDDINGTON MARK [US] ET AL) 4 August 2005 (2005-08-04) the whole document	1-41, 72-91, 108-118
A	US 5 433 896 A (KANG HEE C [US] ET AL) 18 July 1995 (1995-07-18) cf. esp. col. 15, 2nd paragraph the whole document	42,43, 53-56, 92-107



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

13 March 2009

Date of mailing of the international search report

19/03/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Fritz, Martin

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2009/031830

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 44-52, 57-71
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers allsearchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search reportcovers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.2

Claims Nos.: 44-52,57-71

Present claims 42-71, 88 and 92-107 relate to an extremely large number of possible products.

Support and disclosure in the sense of Article 6 and 5 PCT is to be found however for only a very small proportion of the products claimed, as there are only 22 compounds disclosed in the description which are within the subject-matter of claim 88, and the compound 22 is the only representative of the compounds of formula V.

The non-compliance with the substantive provisions is to such an extent, that the search was performed taking into consideration the non-compliance in determining the extent of the search of the above-listed claims (PCT Guidelines 9.19 and 9.23).

Consequently no search was carried out on claims 44-52, 57-71, and the search of claims 42-43, 53-56, 88 and 92-107 restricted to those claimed products which appear to be supported and a generalisation of their structural formulae, namely

those compounds V of claim 42, wherein Z1 is a fluorophore as defined in claim 53 (cf. structural formula) and Z2 is R8.

Furthermore the search of claim 88 has been restricted on the compounds claimed therein which are within the scope of the compounds I as defined in claim 1.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.2), should the problems which led to the Article 17(2)PCT declaration be overcome.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2009/031830

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0064988	A	02-11-2000	AT 233795 T	15-03-2003
			AU 751168 B2	08-08-2002
			AU 4478100 A	10-11-2000
			CA 2335359 A1	02-11-2000
			DE 60001531 D1	10-04-2003
			DE 60001531 T2	02-10-2003
			EP 1090073 A1	11-04-2001
US 2005170363	A1	04-08-2005	US 2008214811 A1	04-09-2008
US 5433896	A	18-07-1995	NONE	